

US010465217B2

(12) **United States Patent**
Kim et al.

(10) **Patent No.: US 10,465,217 B2**
(45) **Date of Patent: Nov. 5, 2019**

(54) **MICROORGANISM PRODUCING O-ACETYL HOMOSERINE AND THE METHOD OF PRODUCING O-ACETYL HOMOSERINE USING THE SAME**

(71) Applicant: **CJ CHEILJEDANG CORPORATION**, Seoul (KR)

(72) Inventors: **Hyun Ah Kim**, Seoul (KR); **Ju Hee Seo**, Gyeonggi-do (KR); **Yong Uk Shin**, Gyeonggi-do (KR); **So Young Kim**, Gyeonggi-do (KR); **Sang Kyoum Kim**, Gyeonggi-do (KR); **Kwang Ho Na**, Seoul (KR); **Jee Yeon Bae**, Seoul (KR); **Sung Kwang Son**, Seoul (KR); **Hye Ryun Yoo**, Seoul (KR); **Jin Geun Choi**, Seoul (KR)

(73) Assignee: **CJ CHEILJEDANG CORPORATION**, Seoul (KR)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **14/901,532**

(22) PCT Filed: **Jun. 22, 2015**

(86) PCT No.: **PCT/KR2015/006307**

§ 371 (c)(1),

(2) Date: **Dec. 28, 2015**

(87) PCT Pub. No.: **WO2015/199396**

PCT Pub. Date: **Dec. 30, 2015**

(65) **Prior Publication Data**

US 2017/0101655 A1 Apr. 13, 2017

(30) **Foreign Application Priority Data**

Jun. 23, 2014 (KR) 10-2014-0076779

(51) **Int. Cl.**

C12N 1/21 (2006.01)
C12N 15/54 (2006.01)
C12N 15/53 (2006.01)
C12N 15/60 (2006.01)
C12N 9/10 (2006.01)
C12P 13/06 (2006.01)
C12N 9/12 (2006.01)
C12N 9/88 (2006.01)
C12N 15/52 (2006.01)

(52) **U.S. Cl.**

CPC **C12P 13/06** (2013.01); **C12N 9/1029** (2013.01); **C12N 9/1205** (2013.01); **C12N 9/88** (2013.01); **C12N 15/52** (2013.01); **C12Y 203/01031** (2013.01); **C12Y 203/03001** (2013.01); **C12Y 207/01039** (2013.01); **C12Y 401/01031** (2013.01)

(58) **Field of Classification Search**

None

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

8,465,952 B2 6/2013 Kim et al.
2008/0293101 A1* 11/2008 Peters C12N 9/0036
435/69.1
2011/0053253 A1* 3/2011 Kim C12N 9/0006
435/252.33
2012/0252077 A1 10/2012 Figge et al.
2012/0329105 A1* 12/2012 Kuvaveva C12N 9/0016
435/109
2013/0273615 A1 10/2013 Kim et al.

FOREIGN PATENT DOCUMENTS

CN 102002473 A 4/2011
CN 103797027 A 5/2014
EP 2 290 051 A1 3/2011
EP 2 657 250 A2 10/2013
EP 2 657 345 A1 10/2013
JP 58-170488 A 10/1983
JP 2011-45360 A 3/2011
KR 10-0905381 B1 6/2009
KR 10-0951766 B1 4/2010
KR 10-2011-0023703 A 3/2011
KR 10-1117012 B1 3/2012
KR 10-2012-0070531 A 6/2012
WO 2008/013432 A1 1/2008
WO 2010/038905 A1 4/2010
WO 2011/073738 A1 6/2011
WO 2012/087039 A1 6/2012

OTHER PUBLICATIONS

Viola, The Central Enzymes of the Aspartate Family of Amino Acid Biosynthesis, Acc. Chem. Res., 2001, 34, 339-49.*
Zubieta et al., A Single Amino Acid Change Is Responsible for Evolution of Acyltransferase Specificity in Bacterial Methionine Biosynthesis, J. Biol. Chem., 2008, 283, 7561-67.*
Soma et al., Metabolic flux redirection from a central metabolic pathway toward a synthetic pathway using a metabolic toggle switch, Metabolic Eng., Feb. 2014, 23, 175-84.*
Datsenko et al., "One-step inactivation of chromosomal genes in *Escherichia coli* K-12 using PCR products," PNAS 97(12): 6640-6645, Jun. 6, 2000.
Nakashima et al., "Multiple-Gene Silencing Using Antisense RNAs in *Escherichia coli*," Michael Kaufmann and Claudia Klinger (eds.), Functional Genomics: Methods and Protocols, Methods in Molecular Biology 815: 307-319, 2012.
Soma et al., "Metabolic flux redirection from a central metabolic pathway toward a synthetic pathway using a metabolic toggle switch," Metabolic Engineering 23: 175-184, May 2014.

(Continued)

Primary Examiner — Robert B Mondesi

Assistant Examiner — Todd M Epstein

(74) Attorney, Agent, or Firm — Cooley LLP

(57) **ABSTRACT**

Disclosed is a microorganism of *Escherichia* sp. producing O-acetyl homoserine, and a method of producing O-acetyl homoserine in high yield using the microorganism.

6 Claims, 2 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

OTHER PUBLICATIONS

Stokell et al., "Probing the Roles of Key Residues in the Unique Regulatory NADH Binding Site of Type II Citrate Synthase of *Escherichia coli*," *The Journal of Biological Chemistry* 278 (37): 35435-35443, 2003.

Vandedrinck et al., "Metabolic engineering of *Escherichia coli*: construction and characterization of a *gltA* (citrate synthase) knockout mutant," *Meded Rijksuniv Gent Fak Landbouwkd Toegep Biol Wet.* 66(3a):333-6, 2001 (PubMed abstract only).


Harford et al., "Evidence for Isosteric and Allosteric Nucleotide Inhibition of Citrate Synthase from Multiple-Inhibition Studies", *Biochem. J.* (1975) 151, 455-458.

Pereira et al., "Tee Active Site Mutants of *Escherichia coli* Citrate Synthase Effects of Mutations on Catalytic and Allosteric Properties* (Received for publication, May 17, 1993)", *Jan. 7, 1994* (Jan. 7, 1994), pp. 412-417, Retrieved from the Internet: URL: <http://www.jbc.org/content/269/1/412.full.pdf>.


Kim et al., "Proteomic Response Analysis of a Threonine-Overproducing Mutant of *Escherichia coli*," *Biochem J*, 381(3):823-829 (2004).

* cited by examiner

[FIG. 1]



[FIG. 2]



1

**MICROORGANISM PRODUCING O-ACETYL
HOMOSERINE AND THE METHOD OF
PRODUCING O-ACETYL HOMOSERINE
USING THE SAME**

CROSS-REFERENCE TO RELATED
APPLICATIONS

This application is a U.S. national phase application of International PCT Patent Application No. PCT/KR2015/006307, which was filed on Jun. 22, 2015, which claims priority to Korean Patent Application Nos. 10-2014-0076779, filed Jun. 23, 2014. These applications are incorporated herein by reference in their entireties.

STATEMENT REGARDING SEQUENCE
LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is HANO_040_01US_SeqList_ST25.txt. The text file is 87 KB, was created on Dec. 21, 2015, and is being submitted electronically via EFS-Web.

TECHNICAL FIELD

The present invention relates to a microorganism of *Escherichia* sp. producing O-acetyl homoserine, and a method of producing O-acetyl homoserine in high yield using the microorganism.

BACKGROUND ART

O-Acetyl homoserine acts as a precursor of methionine, which is one of the essential amino acids in the body. Methionine has been widely used a component of medical infusion solutions and raw materials for medicinal products as well as an animal feed and food additive.

Methionine can be biologically or chemically synthesized. Recently, a two-step process, in which an L-methionine precursor produced by fermentation is converted to L-methionine by an enzyme reaction, was disclosed (International Publication No. WO 2008/013432). In the above two-step process, O-succinyl homoserine and O-acetyl homoserine may be used as the methionine precursor, and it is important that O-acetyl homoserine be produced in high yield for large-scale cost-effective production of methionine.

DISCLOSURE

Technical Problem

The present inventors, while endeavoring to improve the production of O-acetyl homoserine, discovered that the reduction of the expression or activity of citrate synthase protein can significantly increase the production capability of O-acetyl homoserine, thereby completing the present invention.

Technical Solution

An object of the present invention is to provide an O-acetyl homoserine-producing microorganism with improved O-acetyl homoserine production capability.

2

Another object of the present invention is to provide a method for producing O-acetyl homoserine using the microorganism.

Advantageous Effects

The use of the microorganism with O-acetyl homoserine production capability according to the present invention can produce O-acetyl homoserine in a higher yield and in a more environmentally friendly manner than chemical synthesis. Additionally, the thus produced O-acetyl homoserine can be used as a precursor for the synthesis of methionine and acetic acid by O-acetyl homoserine sulfhydrylase, thereby enabling bioconversion of L-methionine, and the thus converted L-methionine can be widely used in the production of foods or food additives for humans as well as animal feeds or animal feed additives.

DESCRIPTION OF DRAWINGS

FIG. 1 is an expression cassette design for the construction of a microorganism with an attenuated activity of citrate synthase.

FIG. 2 is a restriction map of pBAD24-citrate synthase antisense RNA (asRNA) vector.

BEST MODE

In an aspect, the present invention provides a microorganism of *Escherichia* sp. producing O-acetyl homoserine, in which the activity of endogenous citrate synthase protein is attenuated or inactivated.

As used herein, the term "O-acetyl homoserine", being a specific intermediate material in a methionine biosynthesis pathway of a microorganism, refers to an acetyl-derivative of L-homoserine. O-acetyl homoserine can be produced by an enzyme activity of transferring an acetyl group from acetyl-CoA to homoserine using homoserine and acetyl-CoA as substrates.


As used herein, the term "a microorganism producing O-acetyl homoserine" includes a microorganism, which, being a eukaryotic or prokaryotic microorganism producing O-acetyl homoserine within a living organism, is provided with O-acetyl homoserine-producing capability to its parent microorganism without O-acetyl homoserine-producing capability, or a microorganism which is endogenously provided with the O-acetyl homoserine-producing capability.

O-Acetyl homoserine-producing capability may be provided or promoted by improvement of species. The microorganisms having the O-acetyl homoserine-producing capability may include microorganism belonging to *Escherichia* sp., *Erwinia* sp., *Serratia* sp., *Providencia* sp., *Corynebacteria* sp., *Pseudomonas* sp., *Leptospira* sp., *Salmonella* sp., *Brevibacteria* sp., *Hypomononas* sp., *Chromobacterium* sp., and *Norcardia* sp., or fungi, or yeasts; specifically, microorganism belonging to *Escherichia* sp., *Corynebacteria* sp., *Leptospira* sp., and yeasts; and more specifically, microorganism belonging to *Escherichia* sp., as a specific example, *Escherichia coli*. The microorganisms having the O-acetyl homoserine-producing capability may be microorganisms producing L-lysine, L-threonine, L-isoleucine, or L-methionine, or derivatives thereof, but are not limited thereto.

As used herein, the term "citrate synthase (E.C. 2.3.3.1)" refers to an enzyme in the first step of the TCA cycle that mediates the reaction between oxaloacetate and acetyl-CoA. Specifically, citrate synthase mediates the condensation reaction between an acetate residue having two carbon

3

atoms, which is in acetyl-CoA, and oxaloacetate having four carbon atoms, thereby generating a citrate having six carbon atoms. In *Escherichia coli*, citrate synthase is designated GltA, and citrate synthase and GltA are interchangeably used in the present invention.



Specifically, citrate synthase may be one derived from *Escherichia* sp., and more specifically, GltA derived from *Escherichia coli*. Citrate synthase may be a protein comprising an amino acid sequence represented by SEQ ID NO: 4 or those having a homology of 70% or higher with amino acid sequence of SEQ ID NO: 4, specifically 80% or higher, or more specifically, 90% or higher. Additionally, as a sequence having a homology, if the amino acid sequence is one having the same or corresponding activity of citrate synthase with that of SEQ ID NO: 4, it is obvious that amino acid sequences with a deletion, a modification, a substitution, or an addition, in part of the sequences should also be included in the scope of the present invention. Additionally, based on the genetic code degeneracy, polynucleotide sequences encoding the same amino acid sequence and variants thereof should also be included in the scope of the present invention.

As used herein, the term “endogenous” activity refers to a natural state of a protein in a microorganism or an activity state of the corresponding protein provided in the microorganism before modification.

The “the attenuation or inactivation of a protein activity compared to its endogenous activity” refers to a reduction or elimination of the protein activity when compared with that possessed in its natural state. The attenuation is a concept referring to a case when the activity of a protein is reduced compared with that originally possessed by the microorganism due to a modification in the protein-encoding gene, a case when the level of overall protein expression is lower than that of the natural type strain of the microorganism, or a combination thereof, but is not limited thereto. The inactivation includes a case when the gene encoding the protein is not expressed at all compared to that of the natural type strain, and a case when the gene is expressed but exhibits no activity.

The attenuation or inactivation of a protein activity may be achieved by various methods well known in the art. Examples of the methods may include a method of replacing the gene encoding the protein on the chromosome with a gene modified so that the enzyme activity can be reduced including the case when the protein activity is removed; a method of introducing a modification on the expression-regulating sequence of the gene encoding the protein on the chromosome; a method of replacing the expression-regulating sequence of the gene encoding the protein with a sequence having a weak activity or no activity; a method of deleting a part of or the entire gene encoding the protein on the chromosome; a method of introducing an antisense oligonucleotide (e.g., antisense RNA), which inhibits the translation from the mRNA into a protein via a complementary binding to the transcript of the gene on the chromosome; a method of making the attachment of ribosome impossible by forming a secondary structure by artificially adding a Shine-Dalgarno (SD) sequence and its complementary sequence on the front end of the SD sequence of the gene encoding the protein; a method of reverse transcription engineering (RTE), which adds a promoter so as to be reversely transcribed on the 3' terminus of the open reading frame (ORF) of the corresponding sequence, etc., and also include a combination thereof, but are not limited thereto.

4

Specifically, the method of deleting a part of or the entire gene encoding a protein may be performed by replacing the polynucleotide, encoding the endogenous target protein within the chromosome via a vector for inserting chromosome into a microorganism, with a polynucleotide or a marker where part of the polynucleotide sequence is deleted. For example, a method of gene deletion via homologous recombination may be used, but is not limited thereto. Additionally, as used herein the term “part”, although it may vary depending on the kinds of polynucleotide, may specifically refer to 1 nucleotide to 300 nucleotides, more specifically 1 nucleotide to 100 nucleotides, and even more specifically 1 nucleotide to 50 nucleotides, but is not limited thereto.

Additionally, the method of modifying the expression regulation sequence may be performed by inducing a variation on the expression regulation sequence of the polynucleotide sequence via deletion, insertion, conservative substitution, non-conservative substitution, or a combination thereof so as to further attenuate the activity of the expression regulation sequence; or by replacing the polynucleotide sequence with a polynucleotide sequence with a weaker activity. The polynucleotide sequence may include a promoter, an operator sequence, a sequence encoding a ribosome-binding domain, and a sequence for regulating termination of transcription and translation, but is not limited thereto.

Additionally, the method of modifying the gene sequence on the chromosome may be performed by inducing a variation in the sequence via deletion, insertion, conservative substitution, non-conservative substitution, or a combination thereof so as to further attenuate the activity of the expression regulation sequence; or by replacing the sequence with a gene sequence improved to have a weaker activity or a gene sequence improved to have no activity, but is not limited thereto.

Specifically, for the attenuation of the activity of citrate synthase protein, part of an amino acid(s) in the amino acid sequence of the citrate synthase protein may be substituted with other amino acid(s). More specifically, a citrate synthase having an amino acid sequence, in which the 145th amino acid or the 167th amino acid in the amino acid sequence of the citrate synthase protein is substituted from tyrosine (Y) or lysine (K) to other amino acid(s) may be included. Even more specifically, the citrate synthase may be one having a gene sequence encoding a modified polypeptide, in which the 145th amino acid in the amino acid sequence of the citrate synthase protein is substituted from tyrosine (Y) to alanine (A), and the 167th amino acid is substituted from lysine (K) to alanine (A). In particular, the amino acid residue number was determined in sequential order after setting the amino acid positioned next to the methionine, which is encoded by the initiation codon, as the 1st amino acid. The polypeptide may respectively have an amino acid sequence represented by SEQ ID NO: 1 or 2. Additionally, if the activity of the citrate synthase is weaker than that of a wild-type, the citrate synthase may include amino acid sequences having a homology of 80% or higher with the amino acid sequence of SEQ ID NO: 1 or 2, specifically 90% or higher, more specifically 95% or higher, and even more specifically 97% or higher. As a sequence having a homology, if the amino acid sequence is one which has substantially the same or corresponding biological activity of a protein of SEQ ID NO: 1 or 2, it is obvious that amino acid sequences with a deletion, a modification, a substitution, or an addition in part of the sequences should also be included in the scope of the present invention.


5

As used herein, the term “homology” refers to a percentage in identity between two polynucleotides or polypeptide moieties. The homology between sequences from a moiety to another moiety may be determined by a technology known in the art. For example, the homology may be determined by directly arranging the sequence information between two different polynucleotide molecules or two different polypeptides using a computer program arranging and easily obtaining the sequence information. The computer program may include BLAST (NCBI), CLC Main Workbench (CLC bio), MegAlign™ (DNASTAR Inc), etc. Additionally, the homology between polynucleotides may be determined by hybridizing the polynucleotides under the condition of forming a stable double strand between homologous regions, decomposing with a single-strand-specific nuclease, and determining the decomposed fragments.

As used herein, the term “homology” refers to a relationship between proteins having “common evolutionary origin” including homologous proteins derived from superfamily proteins in all grammatical forms or with spelling variations, and those derived from different species. These proteins (and the genes encoding the same) have sequence homologies reflected by high levels of sequence similarities. However, the term “homology”, for its general use and the use in the present invention, would refer to a sequence similarity when modified by an adjective such as “very high”, instead of referring to common evolutionary origin.

In an exemplary embodiment of the present invention, the microorganism may be one in which the activity of cystathionine gamma synthase (EC 2.5.1.48), homoserine kinase (EC 2.7.1.39), or the activities of both are weaker than their endogenous activities, or inactivated.


As used herein, the term “cystathionine gamma synthase” refers to an enzyme which can synthesize cystathionine by a chemical reaction described below, using O-succinyl homoserine and L-cysteine as substrates. In the present invention, the cystathionine gamma synthase from *E. coli*, is designated as “MetB”.



Specifically, the cystathionine gamma synthase from *E. coli*, although not particularly limited thereto, may be a protein comprising an amino acid sequence represented by SEQ ID NO: 9 or those having a homology of 70% or higher with the amino acid sequence of SEQ ID NO: 9, specifically 80% or higher, and more specifically 90% or higher. Additionally, as a sequence having a homology, if the amino acid sequence is one having the same or corresponding activity of homoserine kinase with the amino acid sequence of SEQ ID NO: 9, it is obvious that amino acid sequences with a deletion, a modification, a substitution, or an addition in part of the sequences should also be included in the scope of the present invention. Additionally, based on the genetic code degeneracy, polynucleotide sequences encoding the same amino acid sequence and variants thereof should also be included in the scope of the present invention.

The method for attenuation and inactivation of the cystathionine gamma synthase activity may be performed according to the method described above.

As used herein, the term “homoserine kinase” refers to an enzyme causing the phosphorylation of homoserine, which performs the chemical reaction described below. In the present invention, the homoserine kinase from *E. coli*, is designated as “ThrB”.



6

Specifically, homoserine kinase from *Escherichia* sp., although not particularly limited thereto, may be a protein comprising an amino acid sequence represented by SEQ ID NO: 11 or those having a homology of 70% or higher with the amino sequence of SEQ ID NO: 11, specifically 80% or higher, or more specifically, 90% or higher. Additionally, as a sequence having a homology, if the amino acid sequence is one having the same or corresponding activity of homoserine kinase with the amino sequence of SEQ ID NO: 11, it is obvious that amino acid sequences with a deletion, a modification, a substitution, or an addition in part of the sequences should also be included in the scope of the present invention. Additionally, based on the genetic code degeneracy, polynucleotide sequences encoding the same amino acid sequence and variants thereof should also be included in the scope of the present invention.

The method for attenuation and inactivation of the homoserine kinase activity may be performed according to the method described above.

In a specific aspect of the present invention, the microorganism may be one, in which the activity of homoserine O-acetyltransferase is further introduced or enhanced, or the endogenous homoserine O-succinyltransferase is further modified to have the activity of homoserine O-acetyltransferase.

As used herein, the term “homoserine O-acetyltransferase (EC 2.3.1.31)” refers to an enzyme having an activity of transferring an acetyl group from acetyl-CoA to homoserine.

Specifically, the microorganism according to the present invention may be introduced with the activity of homoserine O-acetyltransferase. The homoserine O-acetyltransferase may be derived from various microorganism species, for example, from a microorganism selected from *Corynebacteria* sp., *Leptospira* sp., *Deinococcus* sp., *Deinococcus* sp., *Pseudomonas* sp., and *Mycobacterium* sp. Specifically, the homoserine O-acetyltransferase may be those which include the amino acid sequences represented by SEQ ID NO: 13 (*Leptospira meyeri*), SEQ ID NO: 14 (*Corynebacterium glutamicum*), or SEQ ID NO: 15 (*Deinococcus radiodurans*), but is not limited thereto. Additionally, the homoserine O-acetyltransferase may be a protein comprising the above amino acid sequences or those having a homology of 70% or higher with the above amino acid sequences, specifically 80% or higher, or more specifically, 90% or higher. Additionally, based on the genetic code degeneracy, polynucleotide sequences encoding the same amino acid sequence and variants thereof should also be included in the scope of the present invention.

Examples of the homoserine O-acetyltransferase to be used in the present invention are disclosed in Korean Patent Application Publication No. 10-2011-0023703 and European Patent Application Publication No. EP 2290051, and the entire specifications of these patent documents may be included as references in the present invention.

Additionally, the protein, in which the endogenous homoserine O-succinyltransferase (EC 2.3.1.46) is modified to have the activity of homoserine O-acetyltransferase, refers to a polypeptide, in which the substrate specificity of the polypeptide having the homoserine O-succinyltransferase activity is changed from succinyl-CoA to acetyl-CoA. Additionally, the modified protein, although not particularly limited thereto, may be a polypeptide having homoserine O-acetyltransferase activity, unlike its wild-type, by replacing part of the amino acid sequence of the polypeptide having homoserine O-succinyltransferase activity.

Examples of the homoserine O-succinyltransferase may be a polypeptide from *Enterobacteria* sp., *Salmonella* sp.,

Pseudomonas sp., *Bacillus* sp., or *Escherichia* sp., specifically, a polypeptide from *Escherichia* sp. having the homoserine O-succinyltransferase activity, for example, a polypeptide having homoserine O-succinyltransferase activity from *E. coli*. More specifically, the homoserine O-succinyltransferase from *E. coli* may have the amino acid sequence represented SEQ ID NO: 16, but is not limited thereto. The homoserine O-succinyltransferase from *E. coli* is designated as "MetA".

The modified homoserine O-succinyltransferase may be a variant polypeptide, in which the 111th amino acid of the polypeptide represented by SEQ ID NO: 16 or polypeptides having a homology of 95% or above with the polynucleotide sequence of SEQ ID NO: 16 is substituted with glutamic acid, and additionally, the 112nd amino acid is substituted with threonine or histidine. Specifically, the variant polypeptide may be a polypeptide having the amino acid sequence of any of SEQ ID NOS: 17 to 19. Additionally, the variant polypeptide may be a protein comprising an amino acid sequence having a homology of 70% or higher with the above amino acid sequence, specifically 80% or higher, or more specifically 90% or higher. Additionally, based on the genetic code degeneracy, polynucleotide sequences encoding the same amino acid sequence and variants thereof should also be included in the scope of the present invention. The information on the modified homoserine O-succinyltransferase can be obtained from Korean Patent Application Publication No. 10-2012-0070531 or International Publication No. WO2012/087039, and the entire specifications of these patent documents are included as references to the present invention.

As used herein, the term "introduction or enhancement of activity" refers to providing the activity of a particular protein to a microorganism which does not possess the protein; or enhancement of the intracellular activity of the protein in the microorganism which possesses the protein, and the like, and refers to the increase of the intracellular activity of the protein compared to the endogenous activity of the protein.

As used herein, the term "introduction or enhancement of protein activity" refers to not only the drawing of a higher effect than the original function due to the increase in the activity of the protein itself, but also the increase in the activity of the protein due to the increase in endogenous gene activity, endogenous gene amplification by the internal or external factors, the increase in copy number, gene introduction from outside, increase in enzyme activity due to the substitution, modification, or mutation, etc., but is not limited thereto.

In the above, the increase in gene copy number, although not particularly limited thereto, may be performed in a state operably connected to a vector, or by being inserted into the chromosome within a host cell. Specifically, the method may be executed by introducing a vector, which a polynucleotide encoding the protein of the present invention is operably connected to and can be replicated and function irrespective of a host, into a cell of the host; or introducing a vector, to which the polynucleotide is operably connected, inserting the polynucleotide into the chromosome of the host cell, into the host cell, thereby increasing the number of gene copies within the chromosome of the host cell.

The vector is a DNA construct including the polynucleotide sequence of the polynucleotide encoding the target protein, which is operably connected to a suitable regulation sequence so that the target protein can be expressed in an appropriate host, wherein the regulation sequence includes a promoter initiating transcription, a random operator

sequence for regulation of the transcription, a sequence encoding a suitable mRNA ribosome-binding domain, and a sequence for regulation of transcription and translation. The vector, after being transformed into a suitable host cell, may be replicated or function irrespective of the host genome, or may be integrated into the host genome itself.

The vector used in the present invention may not be specifically limited as long as the vector is replicable in the host cell, and any vector known in the art may be used. Examples of the vector may include natural or recombinant plasmids, cosmids, viruses, and bacteriophages. For example, as a phage vector or cosmid vector, pWE15, M13, λ MBL3, λ MBL4, λ IXII, λ ASHII, λ APII, λ t10, λ t11, Charon4A, Charon21A, etc., may be used; and as a plasmid vector, pBR-based, pUC-based, pBluescriptII-based, pGEM-based, pTZ-based, pCL-based, pET-based, etc., may be used. Specifically, pDZ, pACYC177, pACYC184, pCL, pECCG117, pUC19, pBR322, pMW118, pCC1BAC vectors, etc., may be used.

Additionally, a polynucleotide encoding an endogenous target protein may be substituted with a modified polynucleotide using a vector for inserting into the chromosome of a microorganism. The insertion of the polynucleotide into the chromosome may be performed using a known method in the art, for example, by homologous recombination. Since the vector of the present invention can be inserted into the chromosome via homologous recombination, a selection marker for confirmation of the insertion into the chromosome may be additionally included. The selection marker is used for selection of a transformed cell, i.e., in order to confirm whether the target polynucleotide has been inserted, and markers providing selectable phenotypes such as drug resistance, nutrient requirement, resistance to cytotoxic agents, and expression of surface proteins may be used, but are not limited thereto. Under the circumstances where selective agents are treated, only the cells expressing the selection markers can survive or express other phenotypic traits, and thus the transformed cells can be easily selected.

As used herein, the term "transformation" refers to a process of introducing a vector including a polynucleotide encoding a target protein into a host cell thereby enabling the expression of the polynucleotide encoded by the protein in the host cell. For the transformed polynucleotide, it does not matter whether it is inserted into the chromosome of a host cell and located therein and located outside the chromosome, as long as it can be expressed in the host cell. Additionally, the polynucleotide includes DNA and RNA which encode the target protein. The polynucleotide may be inserted in any form insofar as it can be introduced into a host cell and expressed therein. For example, the polynucleotide may be introduced into a host cell in the form of an expression cassette, which is a genetic construct including all essential elements required for self-expression. The expression cassette may conventionally include a promoter operably connected to the polynucleotide, a transcription termination signal, a ribosome-binding domain, and a translation termination signal, and may be in the form of an expression vector capable of self-replication. Additionally, the polynucleotide may be introduced into a host cell as it is, and operably connected to a sequence essential for its expression in the host cell. Additionally, as used herein, the term "operably connected" refers to a functional connection between a promoter sequence, which initiates and mediates the transcription of the polynucleotide encoding the target protein, and the gene sequence.

Then, the modification of the expression regulation sequence for increasing the expression of the polynucle-

otide, although not particularly limited thereto, may be performed by inducing a variation in the polynucleotide sequence via deletion, insertion, conservative substitution, non-conservative substitution, or a combination thereof so as to further enhance the activity of the expression regulation sequence; or by replacing the polynucleotide sequence with a polynucleotide sequence with a stronger activity. The expression regulation sequence, although not particularly limited thereto, may include a promoter, an operator sequence, a sequence encoding a ribosome-binding domain, and a sequence for regulating termination of transcription and translation, etc. Additionally, a strong exogenous promoter, instead of the original promoter, may be connected to the upper end of the expression unit of the polynucleotide.

Furthermore, the modification of the polynucleotide sequence on the chromosome, although not particularly limited thereto, may be performed by inducing a variation on the expression regulation sequence of the polynucleotide sequence via deletion, insertion, conservative substitution, non-conservative substitution, or a combination thereof so as to further enhance the activity of the polynucleotide sequence; or by replacing the polynucleotide sequence with an enhanced polynucleotide sequence with a stronger activity.

Generally, the introduction and enhancement of the protein activity may increase the activity or concentration of the corresponding protein relative to the activity or concentration of a wild-type protein or in a microorganism from at least 1%, 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400%, or 500%, to a maximum of 1000% or 2000%.

Additionally, the microorganism may be one in which the activity of the endogenous homoserine O-succinyltransferase was attenuated or inactivated compared to that of the endogenous activity, in order to enhance the biosynthesis pathway of O-acetyl homoserine by blocking the pathway of biosynthesizing O-succinyl homoserine from homoserine.

The attenuation and inactivation of the homoserine O-succinyltransferase activity may be performed according to the method explained above.

In an exemplary embodiment of the present invention, the O-acetyl homoserine-producing microorganism may be one in which the activity of an enzyme involved in the biosynthesis pathway from phosphoenolpyruvate to homoserine is additionally introduced or enhanced, in order to further increase the amount of homoserine, a substrate for the biosynthesis of O-acetyl homoserine.

Specifically, the above microorganism may be one, in which the activity of at least one protein selected from the group consisting of phosphoenolpyruvate carboxylase (ppc, EC 4.1.1.31), aspartate aminotransferase (aspC, EC 2.6.1.1), and aspartate semialdehyde dehydrogenase (asd, EC 1.2.1.11) is further introduced or enhanced.

For example, ppc gene encoding phosphoenolpyruvate carboxylate including an amino acid sequence represented by SEQ ID NO: 20, aspC gene encoding aspartate aminotransferase including an amino acid sequence represented by SEQ ID NO: 21, and asd gene encoding aspartate semialdehyde dehydrogenase including an amino acid represented by SEQ ID NO: 22 may be introduced into a microorganism. For example, the activities of the three different enzymes may be introduced and enhanced by making all the genes encoding the three different enzymes present in the chromosome of a host cell with a copy number of at least 2, but is not limited thereto. The introduction and enhancement of the activities may be performed according to the method described above.

In an exemplary embodiment of the present invention, the activity of citrate synthase protein was attenuated or inactivated by various methods, which includes deleting the citrate synthase gene in a microorganism of *E. coli* producing O-acetyl homoserine; introducing the gene encoding the modified citrate synthase protein, whose activity was attenuated compared to that of a wild-type, into the position of the citrate synthase gene; and introducing an expression vector for citrate synthase gene antisense RNA. As a result, the thus-constructed O-acetyl homoserine-producing microorganism, in which the activity of citrate synthase protein was attenuated or inactivated, showed an improved O-acetyl homoserine production capability, compared to that of the parent microorganism (Examples 1 to 4).

In another aspect, the present invention provides a method for producing O-acetyl homoserine using an O-acetyl homoserine-producing microorganism with an improved production capability of O-acetyl homoserine. Specifically, the present invention provides a method for producing O-acetyl homoserine including (a) culturing the microorganism; and (b) recovering O-acetyl homoserine produced during the cultivation of the microorganism.

The method of cultivation of the *E. coli* having O-acetyl homoserine production capability according to the present invention may be performed according to the suitable media and culture conditions known in the art. The cultivation process may be easily adjusted by a skilled person in the art depending on the microorganism to be selected. In particular, since the microorganism of the present invention is a microorganism where the activity of the citrate synthase, which is an enzyme mediating the first step of the TCA cycle, is attenuated or inactivated, the cultivation medium may include glutamate, but is not specifically limited thereto.

Examples of the culture methods may include a batch culture, a continuous culture, and a fed-batch culture, but are not limited thereto. These various methods are, for example, disclosed in "Biochemical Engineering" by James M. Lee, Prentice-Hall International Editions, pp 138-176.

The medium used in the cultivation may appropriately meet the requirement of a specific microorganism. Specifically, examples of the microorganism culture media are disclosed in "Manual of Methods for General Bacteriology" by the American Society for Bacteriology, Washington, D.C., 1981. The culture media may be those which include an appropriate carbon source, phosphorous source, inorganic compound, amino acid, and/or vitamins, etc, and cultivation may be performed in aerobic conditions while adjusting temperature, pH, etc.

Examples of the carbon source may include carbohydrates such as glucose, lactose, sucrose, lactic acid, fructose, maltose, starch, and cellulose; fats such as soybean oil, sunflower oil, castor oil, berber oil, and coconut oil; fatty acids such as palmitic acid, stearic acid, and linoleic acid; alcohols such as glycerol and ethanol; and organic acids such as acetic acid. These carbon sources may be used alone or in combination, but are not limited thereto.

Examples of the nitrogen source may include organic nitrogen sources such as peptone, yeast extract, gravy, malt extract, corn steep liquor (CSL), and bean flour; and inorganic nitrogen sources such as urea, ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate, and ammonium nitrate. These nitrogen sources may be used alone or in combination, but are not limited thereto. The culture media may further include, as a phosphorous source, potassium dihydrogen phosphate, dipotassium hydrogen phosphate, and corresponding sodium-con-

11

taining salts. The culture media may include metals such as magnesium sulfate and iron sulfate. Additionally, amino acids, vitamins and appropriate precursors may be included. These culture media or precursors may be added to the culture in the form of a batch culture or continuous culture, but are not limited thereto.

Additionally, the pH of the culture may be adjusted by adding a compound such as ammonium hydroxide, potassium hydroxide, ammonia, phosphoric acid, and sulfuric acid during cultivation in an appropriate manner. Additionally, bubble formation may be prevented during the cultivation using an antifoaming agent such as fatty acid polyglycol ester. Additionally, in order to maintain aerobic conditions in a culture liquid, an oxygen gas or a gas (e.g., air) containing an oxygen gas may be added to a culture. The culture temperature may be from 20° C. to 45° C., and specifically from 25° C. to 40° C., but is not limited thereto. The cultivation may be continued until the production of O-acetyl homoserine reaches the intended level, and specifically for 10 hours to 160 hours, but is not limited thereto.

The method of producing O-acetyl homoserine of the present invention may further include recovering O-acetyl homoserine from the cultured microorganism or a cultured product thereof. The recovery of the intended O-acetyl homoserine may be performed by a microorganism cultivation method according to the present invention, for example, an appropriate method known in the art such as a batch culture, a continuous culture, and a fed-batch culture.

The recovery may include a purification step.


The thus-recovered O-acetyl homoserine can produce methionine by a two-step process (Korean Patent No. 10-0905381).

The two-step process includes a process of producing L-methionine and an organic acid by an enzyme reaction using an enzyme having the O-acetyl homoserine sulfhydrylase activity or a microorganism possessing the enzyme, while using the O-acetyl homoserine, which was produced by the L-methionine precursor-producing microorganism, and methyl mercaptan as substrates.

More specifically, the present invention provides a method for producing L-methionine by an enzyme reaction of O-acetyl homoserine sulfhydrylase, etc., using O-acetyl homoserine, which was accumulated by the above method, as a substrate.

In the second step, when O-acetyl homoserine is used as an L-methionine precursor, O-acetyl homoserine sulfhydrylase derived from a microorganism, specifically belonging to *Leptospira* sp., *Chromobacterium* sp., and *Hyphomonas* sp., and more specifically belonging to *Leptospira meyeri*, *Pseudomonas aurogenosa*, *Hyphomonas Neptunium*, and *Chromobacterium Violaceum* may be used.

The reaction is the same as shown below.



The additional process for producing methionine is disclosed in Korean Patent No. 10-0905381, and the entire specification of the patent may be included as a reference in the present invention.

MODE FOR INVENTION

Hereinafter, the present invention will be described in more detail with reference to the following Examples. However, these Examples are for illustrative purposes only, and the invention is not intended to be limited by these Examples.

12

REFERENCE EXAMPLE

Construction of O-Acetyl Homoserine-Producing Microorganism

<1-1> Deletion of metB Gene Derived from Wild-Type *E. coli* (International Publication No. WO 2008/013432)

An O-acetyl homoserine-producing microorganism was constructed using *E. coli*, a representative microorganism among *Escherichia* sp. To this end, wild-type *E. coli* K12 W3110 (ATCC27325) obtained from American Type Culture Collection (ATCC) was used. First, in order to block the synthesis pathway from O-succinyl-L-homoserine to cystathionine, cystathionine synthase-encoding metB gene (SEQ ID NO: 10) was deleted. Specifically, the cystathionine synthase-encoding metB gene was deleted via an FRT-one-step PCR deletion method (PNAS (2000) vol 97: P6640-6645).

Specifically, the metB deletion cassette was constructed via a PCR reaction based on the pKD3 vector (PNAS (2000) vol 97: P6640-6645) as a template using primers of SEQ ID NOS: 30 and 31 as follows: 30 cycles of denaturation at 94° C. for 30 seconds, annealing at 55° C. for 30 seconds, and extension at 72° C. for 1 minute. The resulting PCR product was electrophoresed on a 1.0% agarose gel, and a 1.2 kb DNA band obtained therefrom was purified. The recovered DNA fragment was electroporated into *E. coli* (K12) W3110, which was already transformed with the pKD46 vector (PNAS (2000) vol 97: P6640-6645). For electroporation, the W3110 strain transformed with the pKD46 was cultured in an LB medium containing 100 µg/L ampicillin and 5 mM arabinose (L-arabinose) at 30° C. until OD₆₀₀=0.6, and used after washing twice with sterile distilled water and once with 10% glycerol. Electroporation was performed at 2500 V. The recovered strain was streaked on an LB plate containing 25 µg/L chloramphenicol, cultured at 37° C. overnight, and the strain showing resistance was selected. The selected strain was subjected to a PCR reaction under the same conditions based on the strain as a template using the same primers, and the deletion of metB gene was confirmed by observing the gene size of 1.2 kb on a 1.0% agarose gel. The thus-confirmed strain was cultured in an LB medium after transforming again with the pCP20 vector (PNAS (2000) vol 97: P6640-6645), and the final metB gene-deleted strain, where the gene size was reduced to 150 bp on a 1.0% agarose gel through PCR performed under the same conditions, was constructed and the removal of the chloramphenicol marker was confirmed. The thus-constructed strain was designated as "W3-B".

<1-2> Deletion of thrB Gene (International Publication No. WO 2008/013432)

In an effort to increase the amount of O-succinylhomoserine synthesis from homoserine, thrB gene, which is a homoserine kinase-encoding gene, was deleted. For deletion of the thrB gene from the W3-B strain constructed in Example 1, the FRT one step PCR deletion method used in the deletion of metB gene was used.

A thrB deletion cassette was constructed via PCR based on the pKD4 vector (PNAS (2000) vol 97: P6640-6645) as a template using primers of SEQ ID NOS: 32 and 33 as follows: 30 cycles of denaturation at 94° C. for 30 seconds, annealing at 55° C. for 30 seconds, and extension at 72° C. for 1 minute.

The resulting PCR product was electrophoresed on a 1.0% agarose gel, and a 1.6 kb DNA band obtained therefrom was purified. The recovered DNA fragment was electroporated into W3-B strain, which was already transformed

with the pKD46 vector. The recovered strain was streaked on an LB plate containing 50 µg/L kanamycin, cultured at 37° C. overnight, and the strain showing resistance was selected.

The selected strain was subjected to a PCR reaction under the same conditions directly based on the strain as a template using the same primers of SEQ ID NOS: 32 and 33, and confirmed the deletion of thrB gene by selecting the strain having the gene size of 1.6 kb on a 1.0% agarose gel. The thus-confirmed strain was cultured in an LB medium after transforming again with the pCP20 vector, and the final thrB gene-deleted strain, where the gene size was reduced to 150 bp on a 1.0% agarose gel through PCR performed under the same conditions, was constructed and the removal of the kanamycin marker was confirmed. The thus-constructed strain was designated as “W3-BT”.

<1-3> A Variant metA with Homoserine Acetyltransferase Activity (International Publication No. WO 2012/087039)

In order to strengthen the homoserine acetyltransferase activity in the strain obtained in Reference Example <1-2>, it was intended to introduce mutant type metA gene (SEQ ID NOS: 24 and 26) encoding homoserine acetyltransferase.

First, in order to construct the variant of metA gene with a strengthened activity, a PCR reaction was performed based on the chromosome of a wild-type strain W3110 as a template using the primers of SEQ ID NOS: 34 and 35, and the metA gene encoding homoserine O-succinyltransferase was amplified.

The primers used in the PCR reaction were prepared based on the polynucleotide sequence of the *E. coli* chromosome, NC_000913, registered in the NIH Gene Bank, and the primers of SEQ ID NOS: 34 and 35 have the EcoRV and HindIII restriction sites, respectively. The thus-obtained PCR product and the pCL1920 plasmid including Pcj1 were respectively treated with EcoRV and HindIII, and the PCR product was cloned into the pCL1920 plasmid. *E. coli* DH5α was transformed using the cloned plasmid, and the transformed *E. coli* DH5α was selected on LB plates containing 50 µg/mL spectinomycin, and the plasmid was obtained therefrom. The thus-obtained plasmid was designated as “pCL_Pcj1_metA”.

Then, the 111th amino acid, glycine (Gly), of O-succinyltransferase was substituted with glutamic acid (Glu) (G111E) based on the above-constructed pCL_Pcj1_metA plasmid as a template using a site directed mutagenesis kit (Stratagene, USA). The thus-constructed plasmid including the variant of G111E metA gene was designated as “pCL_Pcj1_metA (EL)”.

Additionally, in order to substitute the 111th amino acid of O-succinyltransferase from glycine to glutamic acid, and the 112nd amino acid from leucine to histidine, primers of SEQ ID NOS: 38 and 39 were used. The plasmid including metA gene, in which the 111th amino acid was substituted from glycine to glutamic acid, and the 112nd amino acid was substituted from leucine to histidine was designated as “pCL_Pcj1_metA (EH)”.

Then, a replacement cassette, for the replacement with metA (EH) into a strain, was constructed via PCR using pKD3 vector as a template along with primers of SEQ ID NOS: 40 and 41 as follows: 30 cycles of denaturation at 94° C. for 30 seconds, annealing at 55° C. for 30 seconds, and extension at 72° C. for 2 minutes. The respective PCR product was obtained using pCL-Pcj1-metA (EH) as a template for the metA (EH) part of the replacement cassette along with primers of SEQ ID NOS: 42 and 43, and primers of SEQ ID NOS: 42 and 45 for the wild-type metA part. metA (EH) replacement cassettes including the chloram-

phenicol marker part were constructed using the three different PCR products along with the primers of SEQ ID NOS: 42 and 45, and electroporated into W3-BT strain, which was already transformed with the pKD46 vector, constructed in Reference Example <1-2>. The thus-confirmed strain was cultured in an LB medium after transforming again with the pCP20 vector, and the strain, where the chloramphenicol marker was removed, and metA gene was substituted with metA (EH) was designated as “W3-BTA”.

<1-4> Construction of a Strain with 2 Copies of ppc, aspC, and asd Genes (European Patent Application Publication No. EP 2290051)

In order to increase the O-acetyl homoserine-producing capability of W3-BTA strain constructed in Reference Example <1-3>, the biosynthetic pathway was enhanced by citing the prior filed patent EP 2290051. In the same manner as in the above EP patent, a strain having 2 amplified copies each of the genes, i.e., the ppc gene encoding phosphoenolpyruvate carboxylase using primers of SEQ ID NOS: 46, 47, 48, and 49; the aspC gene encoding aspartate aminotransferase using primers of SEQ ID NOS: 50 and 51; and the asd gene encoding aspartate semialdehyde dehydrogenase using primers of SEQ ID NOS: 52, 53, 54, and 55, was constructed. In particular, the above strain with an enhanced biosynthetic pathway while producing O-acetyl homoserine was designated as “W3-BTA2PCD” (also called “WCJM”).

<1-5> Flask-Culture Experiments

The amount of O-acetyl homoserine production by the strain constructed in Reference Examples <1-3> and <1-4> was tested by an Erlenmeyer flask-culture.

Specifically, W3110, W3-BTA, and WCJM strains were inoculated into LB media, and cultured at 33° C. overnight. Then, a single colony thereof was inoculated into 3 mL of LB medium, cultured at 33° C. for 5 hours, diluted 200-fold in a 250 mL Erlenmeyer flask containing 25 mL of an O-acetyl homoserine-producing medium, cultured at 33° C. at a rate of 200 rpm for 30 hours, and the amount of O-acetyl homoserine production was examined via HPLC analysis. The media compositions used are shown in Table 1 below, and the amount of O-acetyl homoserine production examined is shown in Table 2 below.

TABLE 1

Composition of O-acetyl homoserine-producing flask-culture	
Composition	Conc. (per liter)
glucose	40 g
(NH ₄) ₂ SO ₄	17 g
KH ₂ PO ₄	1.0 g
MgSO ₄ •7H ₂ O	0.5 g
FeSO ₄ •7H ₂ O	5 mg
MnSO ₄ •8H ₂ O	5 mg
ZnSO ₄	5 mg
CaCO ₃	30 g
yeast extract	2 g
methionine	0.15 g
threonine	0.15 g

TABLE 2

	OD (562 nm)	Glucose consumption (g/L)	O-Acetyl homoserine (g/L)
W3110	14.2	40	0
W3-BTA	8.4	36	0.9
WCJM	9.6	35	1.2

15

The result revealed that the wild-type W3110 did not produce O-acetyl homoserine at all, whereas the W3-BTA strain produced 0.9 g/L of O-acetyl homoserine and the WCJM strain, which was strengthened with the biosynthesis pathway, produced 1.2 g/L of O-acetyl homoserine.

Example 1

Deletion of Citrate Synthase Activity

<1-1> Construction of a Citrate Synthase Gene-Deleted Microorganism in an O-Acetyl Homoserine-Producing Microorganism

Citrate synthase (GltA) is the enzyme in the first step of the TCA cycle, and starts with the reaction between oxaloacetate and acetyl-CoA. The growth inhibition by decrease in the TCA cycle is well known (Meded Rijksuniv Gent Fak Landbouwkd Toegep Biol Wet. 2001; 66(3a): 333-6). However, in order to increase the amount of acetyl-CoA used as a substrate for O-acetyl homoserine, W3-BTA and WCJM strains where citrate synthase activity is deleted were to be produced.

Specifically, the citrate synthase gene in W3-BTA and WCJM strains was deleted via PCR based on pKD4 vector as a template using primers of SEQ ID NOS: 56 and 57 as follows: 30 cycles of denaturation at 94° C. for 30 seconds, annealing at 55° C. for 30 seconds, and extension at 72° C. for 2 minutes. The resulting PCR product was electrophoresed on a 1.0% agarose gel, and the size of the gene was confirmed to be 1.6 kb, and its DNA was purified. The recovered DNA fragments were electroporated into W3-BTA and WCJM strains, which were already transformed with the pKD46 vector. For electroporation, W3-BTA and WCJM strains, transformed with the pKD46 vector, were cultured in an LB medium containing 100 µg/L ampicillin and 5 mM arabinose at 30° C. until OD₆₀₀=0.6, and washed twice with distilled water and once with 10% glycerol to be used. Electroporation was performed at 2500 V. The thus-recovered strains were streaked on LB plates containing 50 µg/L kanamycin, cultured at 37° C., and the strains showing resistance were selected.

The selected strains were subjected to PCR using the primers of SEQ ID NOS: 58 and 59 under the same conditions, electrophoresed on a 1.0% agarose gel, and the size of the gene was observed to be 2.5 kb, thereby confirming that a deletion cassette was inserted into the citrate synthase gene portion on the chromosome. The thus-confirmed strains were transformed again with the pCP20 vector, cultured in LB media, and strains having a deletion of the citrate synthase gene, whose size was reduced to 1.1 kb on a 1.0% agarose gel, were constructed by PCR, and it was confirmed that kanamycin markers were removed. The thus-constructed strains were designated as “W3-BTA-AD” and “WCJM-AD”, respectively.

<1-2> Evaluation of a Citrate Synthase Gene-Deleted Microorganism in an O-Acetyl Homoserine-Producing Microorganism

W3-BTA-AD and WCJM-AD strains can grow in an LB medium, but due to the deletion of the citrate synthase gene, they could not grow in an O-acetyl homoserine-containing medium. In order to test the amount of O-acetyl homoserine production, an Erlenmeyer flask-culture was performed under the condition (Table 3—a composition adding glutamate in the medium) of adding 3 g/L of glutamate into the existing composition of the culture medium.

Specifically, W3-BTA-AD and WCJM-AD strains were inoculated into LB media and cultured at 33° C. overnight.

16

Then, a single colony thereof was inoculated into 3 mL of LB medium, cultured at 33° C. for 5 hours, diluted 200-fold in a 250 mL Erlenmeyer flask containing 25 mL of an O-acetyl homoserine-producing medium (with glutamate added), cultured at 33° C. at a rate of 200 rpm for 30 hours, and the amount of O-acetyl homoserine production was examined via HPLC analysis. The amount of O-acetyl homoserine production examined is shown in Table 4 below.

TABLE 3

Composition of a medium with glutamate added to a basal medium	
Composition	Conc. (per liter)
glucose	40 g
(NH ₄) ₂ SO ₄	17 g
KH ₂ PO ₄	1.0 g
MgSO ₄ •7H ₂ O	0.5 g
FeSO ₄ •7H ₂ O	5 mg
MnSO ₄ •8H ₂ O	5 mg
ZnSO ₄	5 mg
CaCO ₃	30 g
yeast extract	2 g
methionine	0.15 g
threonine	0.15 g
glutamate	3 g

TABLE 4

Production of O-acetyl homoserine via flask-culture				
	OD (562 nm)	Glucose consumption (g/L)	O-acetyl homoserine (g/L)	Glutamate (g/L)
W3-BTA	9.9	38	0.9	3.2
W3-BTA-AD	6.1	34	1.4	2.3
WCJM	9.2	37	1.3	3.5
WCJM-AD	5.6	33	2.1	1.7

The result of the O-acetyl homoserine production via flask-culture revealed that the W3-BTA strain produced 0.9 g/L of O-acetyl homoserine, and W3-BTA-AD produced 1.4 g/L of O-acetyl homoserine, which is a 55.6% increase, although it showed a decrease in its glucose consumption. The WCJM strain produced 1.3 g/L of O-acetyl homoserine while the WCJM-AD strain produced 2.1 g/L of O-acetyl homoserine, thus confirming that the O-acetyl homoserine production capability was improved by 61.5% due to the deletion of the citrate synthase gene.

Example 2

Attenuation of the Activity of Citrate Synthase Protein

<2-1> Kinds of Citrate Synthase Gene Modifications

The WCJM-AD strain constructed in Example <1-1> showed a low culture rate, and three different types of variants, which showed an attenuated activity and a reduced binding ability to acetyl-CoA according to various modifications of citrate synthase known in numerous references (The Journal of Biological Chemistry, 2003, 278, 35435-35443), were selected. The information on the three different types of variants is shown in Table 5, which shows modified genes in which the 145th amino acid, tyrosine (Y), was substituted with alanine (A), and the 167th amino acid, lysine (K), was substituted with alanine (A), and the 204th amino acid, threonine (T), was substituted with alanine (A).

TABLE 5

Evaluation on citrate synthase (gltA) variants		
	KM VALUE [mM]	
	Acetyl-CoA	OAA
WT	0.12	0.026
Y145A	0.23	0.051
K167A	0.22	0.037
T204A	0.21	0.004

<2-2> Construction of a Microorganism with Attenuated Citrate Synthase Protein Activity in an O-Acetyl Homoserine-Producing Microorganism

The present inventors intended to increase the production capability by introducing the variants, in which the activity of citrate synthase protein was attenuated as explained in Example <2-1>, into the O-acetyl homoserine-producing microorganism.

In order to introduce the three different types of citrate synthase gene variants into the WCJM-AD strain, a modification replacement cassette was designed as shown in FIG. 1. Each variant was synthesized by substituting a primer with a nucleotide, and each cassette was constructed through 3 PCR products. For the citrate synthase gene portion, the W3110 strain was used as a template, and the modification on the 145th amino acid produced PCR reactions were performed using the primers of SEQ ID NOS: 60 and 63 and SEQ ID NOS: 62 and 61, respectively, and obtained PCR products with a size of 514 bp and 1,112 bp.

Likewise, the modification on the 167th amino acid produced PCR products with a size of 580 bp and 1,046 bp using the primers of SEQ ID NOS: 60 and 65, and the primers of SEQ ID NOS: 64 and 61, and the modification on the 204th amino acid produced PCR products with a size of 688 bp and 936 bp using the primers of SEQ ID NOS: 60 and 67 and SEQ ID NOS: 66 and 61. For the common kanamycin portion, PCR reactions were performed based on the pKD4 vector as a template using the primers of SEQ ID NOS: 68 and 69. In particular, for the insertion into the position of the citrate synthase gene, the cassette was constructed so as to include the polynucleotide sequence downstream of the citrate synthase gene in the SEQ ID NO: 69, and a PCR product with a size of 1,571 bp was obtained via electrophoresis. A sewing PCR reaction was performed based on the kanamycin DNA fragment, which is the common part with each of the two DNA fragments collected according to the modifications, respectively, using the primers of SEQ ID NOS: 60 and 69, as follows: 30 cycles of denaturation at 94° C. for 30 seconds, annealing at 55° C. for 30 seconds, and extension at 72° C. for 4 minutes. Each of the final PCR products was confirmed on a 1.0% agarose gel, and the DNA fragments with a size of 3,115 bp for the three different kinds of cassettes of citrate synthase gene modifications. The collected DNA fragments were electroporated into the WCJM-AD strain, which was already transformed with the pKD46 vector. For electroporation, the W3110 strain transformed with the pKD46 was cultured in an LB medium containing 100 µg/L ampicillin and 5 mM arabinose at 30° C. until OD₆₀₀=0.6, and used after washing twice with sterile distilled water and once with 10% glycerol. Electroporation was performed at 2500 V. The recovered strain was streaked on an LB plate containing 25 µg/L kanamycin, cultured at 37° C. overnight, and the strain showing resistance was selected. The selected strain was subjected to a PCR reaction under the same conditions based on the strain

as a template using the same primers of SEQ ID NOS: 58 and 59, and confirmed the deletion of metB gene by observing the gene size of 3.7 kb on a 1.0% agarose gel, thereby confirming that a modification cassette, in which the amino acid of the citrate synthase gene was substituted, was inserted. The thus-confirmed strain was cultured in an LB medium after transforming again with the pCP20 vector, and the three variant strains regarding the citrate synthase activity, where the gene size was reduced to 2.5 kb on a 1.0% agarose gel through PCR performed under the same conditions, were constructed and the removal of the kanamycin marker was confirmed. The thus-constructed strains were designated as "WCJM-A145", "WCJM-A167", and "WCJM-A204", and the sequence information of the citrate synthase gene introduced with modifications are shown in SEQ ID NOS: 1 to 3 (amino acid sequences) and SEQ ID NOS: 5 to 7 (nucleotide sequences), respectively.

<2-3> Evaluation of Microorganisms with Attenuated Citrate Synthase Activity in O-Acetyl Homoserine-Producing Microorganisms

An Erlenmeyer flask-culture was performed in order to examine the amount of O-acetyl homoserine production by three different strains of WCJM-A145, WCJM-A167, and WCJM-A204, in which the activity of the citrate synthase gene was attenuated. Four kinds of strains, i.e., WCJM-A145, WCJM-A167, and WCJM-A204 strains including the WCJM strain, were inoculated into LB media, and cultured at 33° C. overnight. Then, a single colony thereof was inoculated into 3 mL of LB medium, cultured at 33° C. for 5 hours, diluted 200-fold in a 250 mL Erlenmeyer flask containing 25 mL of an O-acetyl homoserine-producing medium, cultured at 33° C. at a rate of 200 rpm for 30 hours, and the amount of O-acetyl homoserine production was examined via HPLC analysis. The results are shown in Table 6 below.

TABLE 6

Production of O-acetyl homoserine via flask-culture				
	OD (562 nm)	Glucose consumption (g/L)	O-acetyl homoserine (g/L)	Glutamate (g/L)
WCJM	8.9	35	1.3	1.3
WCJM-A145	7.4	35	2.0	0
WCJM-A167	6.3	29	1.9	0
WCJM-A204	9.1	40	1.1	1.8

The result of O-acetyl homoserine production via flask-culture revealed that the WCJM strain produced 1.3 g/L of O-acetyl homoserine, and the two strains, WCJM-A145 and WCJM-A167, produced 2.0 g/L and 1.9 g/L of O-acetyl homoserine, respectively, while the amount of their glucose consumption decreased along with the decrease in their absorbance (OD). Considering the specific decrease of glutamate from 1.3 g/L to 0 g/L, it was confirmed that the result is due to the decrease in TCA cycle flow caused by the attenuation in the citrate synthase activity. However, the WCJM-A204 strain showed an increase in glutamate while showing a decrease in the amount of O-acetyl homoserine production to 0.2 g/L, thus confirming the modification is one with a strengthened activity.

Example 3

Attenuation in Expression of Citrate Synthase Protein

<3-1> Construction of Expression Vector for Citrate Synthase Gene Antisense RNA (asRNA)

The present inventors made an effort to apply an antisense RNA (asRNA) technology in order to attenuate the expres-

sion of citrate synthase protein. The antisense RNA technology is a method for reducing protein expression by neutralizing the binding between citrate synthase mRNA and ribosome, via overexpression of the complementary binding portion to the citrate synthase mRNA of the target gene. This method has the advantage in that it can regulate the level of inhibition by controlling the binding force with the mRNA of the citrate synthase gene, and this method is also useful for the construction of a recombinant microorganism because this method can effectively construct and reduce gene expression via antisense RNA controlling gene expression, not necessitating the conventional process of gene deletion.

The vector construction was performed referring to a reference (Methods Mol Biol. 2012; 815:307-19. doi: 10.1007/978-1-61779-424-7_23.), and for overexpression, the antisense RNA region of the synthase gene was to be introduced into the pBAD24 plasmid capable of induction. The pBAD24-citrate synthase asRNA vector map is shown in FIG. 2. The region where the antisense RNA of the citrate synthase gene was expressed has a size of 100 bp including the 52 bp region of the promoter region and the 48 bp region from the initiation codon of the citrate synthase, and a 38 bp paired termini (PT) structure, which reduces the instability of the antisense RNA (asRNA), is connected to both flanking regions. The antisense RNA region of the citrate synthase gene was obtained using the primers of SEQ ID NOS: 70 and 71, and the NcoI and HindIII restriction sites were included to be cloned into a vector.

The thus-obtained PCR product had a size of 194 bp, and the PCR product was cloned into the pBAD24 plasmid after treating them with EcoRV and HindIII, respectively. The thus-cloned plasmid was used to transform *E. coli* DH5 α , and the transformed *E. coli* DH5 α was selected from the LB plates containing 100 μ g/mL of ampicillin, and obtained the plasmid therefrom. The thus-obtained plasmid was designated as "pBAD24-gltA asRNA".

<3-2> Introduction of an Expression Vector of Antisense RNA of Citrate Synthase Gene into an O-Acetyl Homoserine-Producing Microorganism and Evaluation Thereof

The pBAD24-gltA-asRNA, an expression vector of antisense RNA of citrate synthase gene, was transformed into the WCJM strain, which is an O-acetyl homoserine-producing microorganism. Here, the transformed strain was designated as "WCJM/A-asRNA". In particular, it was attempted to control the expression amount of the citrate synthase protein by controlling the expression amount of the antisense RNA of the citrate synthase, and here, the expression amount of the antisense RNA can be controlled according to the concentration of arabinose.

As a result, it was confirmed that the amount of the O-acetyl homoserine production increased when the activity of the citrate synthase was attenuated as in Example 2.

Additionally, an Erlenmeyer flask-culture was performed to examine whether the amount of O-acetyl homoserine production increases as the expression amount of the citrate synthase decreases.

Specifically, WCJM and WCJM/A-asRNA strains were inoculated into LB media, and cultured at 33° C. overnight. Then, a single colony thereof was inoculated into 3 mL of LB medium, cultured at 33° C. for 5 hours, diluted 200-fold in a 250 mL Erlenmeyer flask containing 25 mL of an O-acetyl homoserine-producing medium. In particular, in order to control the expression amount of antisense RNA of the citrate synthase, arabinose was added at concentrations

of 0 mM, 2 mM, and 5 mM, and cultured at 33° C. at a rate of 200 rpm for 15 hours and 30 hours. The amount of O-acetyl homoserine production was examined via HPLC analysis, and the results are shown in Tables 7 and 8 below.

TABLE 7

15 Hours	Arabinose	OD (562 nm)	Glucose consumption (g/L)	O-acetyl homoserine (g/L)
WCJM	0 mM	4.2	9.7	0.5
WCJM	2 mM	4.5	8.9	0.6
WCJM	5 mM	4.7	8.9	0.5
WCJM/A-asRNA	0 mM	4.5	10.1	0.6
WCJM/A-asRNA	2 mM	4.2	8.8	0.6
WCJM/A-asRNA	5 mM	3.4	6.9	0.5

TABLE 8

30 Hours	Arabinose	OD (562 nm)	Glucose consumption (g/L)	O-acetyl homoserine (g/L)
WCJM	0 mM	8.9	32	1.4
WCJM	2 mM	9.1	34	1.3
WCJM	5 mM	8.9	33	1.3
WCJM/A-asRNA	0 mM	9.2	33	1.3
WCJM/A-asRNA	2 mM	8.8	32	1.6
WCJM/A-asRNA	5 mM	7.1	29	1.7

As a result, it was confirmed that, when cultured for 15 hours, the WCJM/A-asRNA strain showed a decrease in OD by about 1 according to the concentration of arabinose, whereas the concentration of O-acetyl homoserine was similar. However, when cultured for 30 hours, the WCJM strain, which is a control strain, showed the same OD and O-acetyl homoserine concentration even when the concentration of arabinose increased, whereas the WCJM/A-asRNA strain, which is a strain introduced with the expression vector for citrate synthase RNA antisense, showed a marked difference as the concentration of arabinose increased. The OD was 9.2 when the arabinose concentration was 0 mM, whereas the OD was 7.1 at 5 mM of the arabinose concentration, a decrease of 5.1, and the amount of O-acetyl homoserine increased by 30.8% although the glucose consumption was small. From these results, it was confirmed that not only the attenuation in citrate synthase activity but also in the attenuation in protein expression exhibit the same results.

Example 4

Attenuation and Inactivation of Citrate Synthase Activity in a Microorganism with High Production Yield of O-Acetyl Homoserine

<4-1> Construction of a Microorganism with High O-Acetyl Homoserine Production Yield with Inactivated Citrate Synthase Activity and Evaluation Thereof

International Publication No. WO 2012/087039 discloses in detail a method for constructing an O-acetyl homoserine-producing microorganism from a threonine-producing microorganism derived from a wild-type W3110 strain, due

21

to NTG mutation. In particular, the constructed strain producing O-acetyl homoserine with high yield was deposited at Korean Culture Center of Microorganisms (KCCM) under the Accession No of KCCM 11146P.

The KCCM11146P strain can consume 40 g/L of glucose during a flask-culture and produces about 15 g/L to 16 g/L of O-acetyl homoserine and is thus regarded as having high O-acetyl homoserine production capability. Accordingly, in order to examine whether the strain produces a higher yield of O-acetyl homoserine when the citrate synthase activity is deleted, the same was applied to the KCCM11146P strain. The construction method was the same as in Example <1-1>, and by this method, the KCCM11146P strain, where the citrate synthase activity was deleted, was constructed and designated as "KCCM11146P-AD".

The amount of O-acetyl homoserine production by the KCCM11146P strain, where the citrate synthase activity was deleted, was tested by an Erlenmeyer flask-culture. The KCCM11146P or KCCM11146P-AD strain was inoculated into an LB medium and cultured at 33° C. overnight. Then, a single colony thereof was inoculated into 3 mL of LB medium, cultured at 33° C. for 5 hours, diluted 200-fold in a 250 mL Erlenmeyer flask containing 25 mL of an O-acetyl homoserine-producing medium (with glutamate added), and cultured at 33° C. at a rate of 200 rpm for 30 hours. The amount of O-acetyl homoserine production was examined via HPLC analysis, and the results are shown in Table 9 below.

TABLE 9

Production of O-acetyl homoserine via flask-culture				
	OD (562 nm)	Glucose consumption (g/L)	O-acetyl homoserine (g/L)	Glutamate (g/L)
KCCM11146P	18.3	60	14.2	4.6
KCCM11146P-AD	14.6	60	16.7	1.8

The result of O-acetyl homoserine production via flask-culture revealed that the KCCM11146P strain produced 14.2 g/L of O-acetyl homoserine, and the KCCM11146P-AD strain produced 16.7 g/L of O-acetyl homoserine, a 17.6% increase, although it showed a decrease in absorbance (OD). <4-2> Construction of a Microorganism with High O-Acetyl Homoserine Production Yield with Attenuated Citrate Synthase Activity and Evaluation Thereof

In order to examine whether the KCCM11146P strain, which is a strain with high production yield of O-acetyl homoserine, produces a higher yield of O-acetyl homoserine even when the citrate synthase activity is attenuated, the modification on the 145th amino acid (from tyrosine (Y) to alanine (A)) and the modification on the 167th amino acid (from lysine (K) to alanine (A)), which showed the highest O-acetyl homoserine-producing capabilities among the three variant types attenuating the protein activities explained in Example <2-1>, were applied to the KCCM11146P strain.

The construction method was the same as in Example <2-2>, and by the method, two KCCM11146P strains, where the citrate synthase activity was attenuated, were constructed and designated as "KCCM11146P-A145" and "KCCM11146P-A167", respectively.

The amount of O-acetyl homoserine production by the two strains of KCCM11146P-A145 and KCCM11146P-A167, where the citrate synthase activity was attenuated, was tested by an Erlenmeyer flask-culture. The three strains, i.e., KCCM11146P-A145 and KCCM11146P-A167 strains

22

and the KCCM11146P strain, were inoculated into LB media, and cultured at 33° C. overnight. Then, a single colony thereof was inoculated into 3 mL of LB medium, cultured at 33° C. for 5 hours, diluted 200-fold in a 250 mL Erlenmeyer flask containing 25 mL of an O-acetyl homoserine-producing medium, and cultured at 33° C. at a rate of 200 rpm for 30 hours. The amount of O-acetyl homoserine production was examined via HPLC analysis, and the results are shown in Table 10 below.

TABLE 10

Production of O-acetyl homoserine via flask-culture				
	OD (562 nm)	Glucose consumption (g/L)	O-acetyl homoserine (g/L)	Glutamate (g/L)
KCCM11146P	16.3	60	15.0	1.6
KCCM11146P-A145	14.6	60	17.5	0
KCCM11146P-A167	14.2	60	17.3	0

The result of O-acetyl homoserine production via flask-culture revealed that the KCCM11146P strain produced 15.0 g/L of O-acetyl homoserine, and the two strains of KCCM11146P-A145 and KCCM11146P-A167 showed similar results as in Example <2-3>. The two strains respectively produced 17.5 g/L and 17.3 g/L of O-acetyl homoserine, an increase of about 16.7%, although they both showed a decrease in absorbance (OD).

The strain with high production yield of O-acetyl homoserine also showed a decrease in glutamate from 1.6 g/L to 0 g/L, according to the decrease in TCA cycle flow caused by the attenuation in the citrate synthase activity.

These results demonstrate that citrate synthase activity enables production of O-acetyl homoserine by applying the attenuated modification. Additionally, they also indicate that when a conversion reaction is conducted based on the O-acetyl homoserine, which was produced according to the International Publication No. WO2008/013432, as a template, and using a conversion enzyme, which additionally has the activities of cystathionine gamma synthase, O-succinylhomoserine sulfhydrylase, and O-acetyl homoserine sulfhydrylase, it is possible to simultaneously synthesize L-methionine and acetate.

The present inventors confirmed that the KCCM11146P strain, the variant on the 167th amino acid of the citrate synthase, has an improved production of O-acetyl homoserine, designated the KCCM11146P-A167 strain as "CA05-4007", and deposited it at the Korean Culture Center of Microorganisms (KCCM), an international depositary authority under the Budapest Treaty, on Nov. 22, 2013 (Accession No: KCCM 11483P).

From the foregoing, a skilled person in the art to which the present invention pertains will be able to understand that the present invention may be embodied in other specific forms without modifying the technical concepts or essential characteristics of the present invention. In this regard, the exemplary embodiments disclosed herein are only for illustrative purposes and should not be construed as limiting the scope of the present invention. On the contrary, the present invention is intended to cover not only the exemplary embodiments but also various alternatives, modifications, equivalents and other embodiments that may be included within the spirit and scope of the present invention as defined by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 71

<210> SEQ ID NO 1

<211> LENGTH: 427

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: GltA A145

<400> SEQUENCE: 1

```

Met Ala Asp Thr Lys Ala Lys Leu Thr Leu Asn Gly Asp Thr Ala Val
1          5          10          15

Glu Leu Asp Val Leu Lys Gly Thr Leu Gly Gln Asp Val Ile Asp Ile
20          25          30

Arg Thr Leu Gly Ser Lys Gly Val Phe Thr Phe Asp Pro Gly Phe Thr
35          40          45

Ser Thr Ala Ser Cys Glu Ser Lys Ile Thr Phe Ile Asp Gly Asp Glu
50          55          60

Gly Ile Leu Leu His Arg Gly Phe Pro Ile Asp Gln Leu Ala Thr Asp
65          70          75          80

Ser Asn Tyr Leu Glu Val Cys Tyr Ile Leu Leu Asn Gly Glu Lys Pro
85          90          95

Thr Gln Glu Gln Tyr Asp Glu Phe Lys Thr Thr Val Thr Arg His Thr
100         105         110

Met Ile His Glu Gln Ile Thr Arg Leu Phe His Ala Phe Arg Arg Asp
115         120         125

Ser His Pro Met Ala Val Met Cys Gly Ile Thr Gly Ala Leu Ala Ala
130         135         140

Phe Ala His Asp Ser Leu Asp Val Asn Asn Pro Arg His Arg Glu Ile
145         150         155         160

Ala Ala Phe Arg Leu Leu Ser Lys Met Pro Thr Met Ala Ala Met Cys
165         170         175

Tyr Lys Tyr Ser Ile Gly Gln Pro Phe Val Tyr Pro Arg Asn Asp Leu
180         185         190

Ser Tyr Ala Gly Asn Phe Leu Asn Met Met Phe Ser Thr Pro Cys Glu
195         200         205

Pro Tyr Glu Val Asn Pro Ile Leu Glu Arg Ala Met Asp Arg Ile Leu
210         215         220

Ile Leu His Ala Asp His Glu Gln Asn Ala Ser Thr Ser Thr Val Arg
225         230         235         240

Thr Ala Gly Ser Ser Gly Ala Asn Pro Phe Ala Cys Ile Ala Ala Gly
245         250         255

Ile Ala Ser Leu Trp Gly Pro Ala His Gly Gly Ala Asn Glu Ala Ala
260         265         270

Leu Lys Met Leu Glu Glu Ile Ser Ser Val Lys His Ile Pro Glu Phe
275         280         285

Val Arg Arg Ala Lys Asp Lys Asn Asp Ser Phe Arg Leu Met Gly Phe
290         295         300

Gly His Arg Val Tyr Lys Asn Tyr Asp Pro Arg Ala Thr Val Met Arg
305         310         315         320

Glu Thr Cys His Glu Val Leu Lys Glu Leu Gly Thr Lys Asp Asp Leu
325         330         335

Leu Glu Val Ala Met Glu Leu Glu Asn Ile Ala Leu Asn Asp Pro Tyr
340         345         350

Phe Ile Glu Lys Lys Leu Tyr Pro Asn Val Asp Phe Tyr Ser Gly Ile

```


-continued

355					360					365					
Ile	Leu	Lys	Ala	Met	Gly	Ile	Pro	Ser	Ser	Met	Phe	Thr	Val	Ile	Phe
370						375					380				
Ala	Met	Ala	Arg	Thr	Val	Gly	Trp	Ile	Ala	His	Trp	Ser	Glu	Met	His
385					390					395					400
Ser	Asp	Gly	Met	Lys	Ile	Ala	Arg	Pro	Arg	Gln	Leu	Tyr	Thr	Gly	Tyr
				405					410					415	
Glu	Lys	Arg	Asp	Phe	Lys	Ser	Asp	Ile	Lys	Arg					
			420					425							

<210> SEQ ID NO 2
 <211> LENGTH: 427
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: GltA A167

<400> SEQUENCE: 2

Met	Ala	Asp	Thr	Lys	Ala	Lys	Leu	Thr	Leu	Asn	Gly	Asp	Thr	Ala	Val
1				5					10					15	
Glu	Leu	Asp	Val	Leu	Lys	Gly	Thr	Leu	Gly	Gln	Asp	Val	Ile	Asp	Ile
			20					25					30		
Arg	Thr	Leu	Gly	Ser	Lys	Gly	Val	Phe	Thr	Phe	Asp	Pro	Gly	Phe	Thr
		35					40					45			
Ser	Thr	Ala	Ser	Cys	Glu	Ser	Lys	Ile	Thr	Phe	Ile	Asp	Gly	Asp	Glu
		50				55					60				
Gly	Ile	Leu	Leu	His	Arg	Gly	Phe	Pro	Ile	Asp	Gln	Leu	Ala	Thr	Asp
65				70						75					80
Ser	Asn	Tyr	Leu	Glu	Val	Cys	Tyr	Ile	Leu	Leu	Asn	Gly	Glu	Lys	Pro
				85					90					95	
Thr	Gln	Glu	Gln	Tyr	Asp	Glu	Phe	Lys	Thr	Thr	Val	Thr	Arg	His	Thr
			100					105					110		
Met	Ile	His	Glu	Gln	Ile	Thr	Arg	Leu	Phe	His	Ala	Phe	Arg	Arg	Asp
		115					120					125			
Ser	His	Pro	Met	Ala	Val	Met	Cys	Gly	Ile	Thr	Gly	Ala	Leu	Ala	Ala
		130				135					140				
Phe	Tyr	His	Asp	Ser	Leu	Asp	Val	Asn	Asn	Pro	Arg	His	Arg	Glu	Ile
145					150					155					160
Ala	Ala	Phe	Arg	Leu	Leu	Ser	Ala	Met	Pro	Thr	Met	Ala	Ala	Met	Cys
				165					170					175	
Tyr	Lys	Tyr	Ser	Ile	Gly	Gln	Pro	Phe	Val	Tyr	Pro	Arg	Asn	Asp	Leu
			180					185					190		
Ser	Tyr	Ala	Gly	Asn	Phe	Leu	Asn	Met	Met	Phe	Ser	Thr	Pro	Cys	Glu
		195					200					205			
Pro	Tyr	Glu	Val	Asn	Pro	Ile	Leu	Glu	Arg	Ala	Met	Asp	Arg	Ile	Leu
		210				215					220				
Ile	Leu	His	Ala	Asp	His	Glu	Gln	Asn	Ala	Ser	Thr	Ser	Thr	Val	Arg
225					230					235					240
Thr	Ala	Gly	Ser	Ser	Gly	Ala	Asn	Pro	Phe	Ala	Cys	Ile	Ala	Ala	Gly
				245					250					255	
Ile	Ala	Ser	Leu	Trp	Gly	Pro	Ala	His	Gly	Gly	Ala	Asn	Glu	Ala	Ala
			260					265					270		
Leu	Lys	Met	Leu	Glu	Glu	Ile	Ser	Ser	Val	Lys	His	Ile	Pro	Glu	Phe
		275					280					285			
Val	Arg	Arg	Ala	Lys	Asp	Lys	Asn	Asp	Ser	Phe	Arg	Leu	Met	Gly	Phe

-continued

290	295	300
Gly His Arg Val Tyr Lys Asn Tyr Asp Pro Arg Ala Thr Val Met Arg 305	310	315
Glu Thr Cys His Glu Val Leu Lys Glu Leu Gly Thr Lys Asp Asp Leu 325		330
Leu Glu Val Ala Met Glu Leu Glu Asn Ile Ala Leu Asn Asp Pro Tyr 340		345
Phe Ile Glu Lys Lys Leu Tyr Pro Asn Val Asp Phe Tyr Ser Gly Ile 355		360
Ile Leu Lys Ala Met Gly Ile Pro Ser Ser Met Phe Thr Val Ile Phe 370		375
Ala Met Ala Arg Thr Val Gly Trp Ile Ala His Trp Ser Glu Met His 385		390
Ser Asp Gly Met Lys Ile Ala Arg Pro Arg Gln Leu Tyr Thr Gly Tyr 405		410
Glu Lys Arg Asp Phe Lys Ser Asp Ile Lys Arg 420		425

<210> SEQ ID NO 3
 <211> LENGTH: 427
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: GltA A204

<400> SEQUENCE: 3

Met Ala Asp Thr Lys Ala Lys Leu Thr Leu Asn Gly Asp Thr Ala Val 1	5	10	15
Glu Leu Asp Val Leu Lys Gly Thr Leu Gly Gln Asp Val Ile Asp Ile 20		25	30
Arg Thr Leu Gly Ser Lys Gly Val Phe Thr Phe Asp Pro Gly Phe Thr 35		40	45
Ser Thr Ala Ser Cys Glu Ser Lys Ile Thr Phe Ile Asp Gly Asp Glu 50		55	60
Gly Ile Leu Leu His Arg Gly Phe Pro Ile Asp Gln Leu Ala Thr Asp 65		70	75
Ser Asn Tyr Leu Glu Val Cys Tyr Ile Leu Leu Asn Gly Glu Lys Pro 85		90	95
Thr Gln Glu Gln Tyr Asp Glu Phe Lys Thr Thr Val Thr Arg His Thr 100		105	110
Met Ile His Glu Gln Ile Thr Arg Leu Phe His Ala Phe Arg Arg Asp 115		120	125
Ser His Pro Met Ala Val Met Cys Gly Ile Thr Gly Ala Leu Ala Ala 130		135	140
Phe Tyr His Asp Ser Leu Asp Val Asn Asn Pro Arg His Arg Glu Ile 145		150	155
Ala Ala Phe Arg Leu Leu Ser Lys Met Pro Thr Met Ala Ala Met Cys 165		170	175
Tyr Lys Tyr Ser Ile Gly Gln Pro Phe Val Tyr Pro Arg Asn Asp Leu 180		185	190
Ser Tyr Ala Gly Asn Phe Leu Asn Met Met Phe Ser Ala Pro Cys Glu 195		200	205
Pro Tyr Glu Val Asn Pro Ile Leu Glu Arg Ala Met Asp Arg Ile Leu 210		215	220
Ile Leu His Ala Asp His Glu Gln Asn Ala Ser Thr Ser Thr Val Arg			

-continued

225		230		235		240
Thr Ala Gly Ser Ser Gly Ala Asn Pro Phe Ala Cys Ile Ala Ala Gly						
		245		250		255
Ile Ala Ser Leu Trp Gly Pro Ala His Gly Gly Ala Asn Glu Ala Ala						
		260		265		270
Leu Lys Met Leu Glu Glu Ile Ser Ser Val Lys His Ile Pro Glu Phe						
		275		280		285
Val Arg Arg Ala Lys Asp Lys Asn Asp Ser Phe Arg Leu Met Gly Phe						
		290		295		300
Gly His Arg Val Tyr Lys Asn Tyr Asp Pro Arg Ala Thr Val Met Arg						
305		310		315		320
Glu Thr Cys His Glu Val Leu Lys Glu Leu Gly Thr Lys Asp Asp Leu						
		325		330		335
Leu Glu Val Ala Met Glu Leu Glu Asn Ile Ala Leu Asn Asp Pro Tyr						
		340		345		350
Phe Ile Glu Lys Lys Leu Tyr Pro Asn Val Asp Phe Tyr Ser Gly Ile						
		355		360		365
Ile Leu Lys Ala Met Gly Ile Pro Ser Ser Met Phe Thr Val Ile Phe						
		370		375		380
Ala Met Ala Arg Thr Val Gly Trp Ile Ala His Trp Ser Glu Met His						
385		390		395		400
Ser Asp Gly Met Lys Ile Ala Arg Pro Arg Gln Leu Tyr Thr Gly Tyr						
		405		410		415
Glu Lys Arg Asp Phe Lys Ser Asp Ile Lys Arg						
		420		425		

<210> SEQ ID NO 4
 <211> LENGTH: 427
 <212> TYPE: PRT
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(427)
 <223> OTHER INFORMATION: GltA

<400> SEQUENCE: 4

Met Ala Asp Thr Lys Ala Lys Leu Thr Leu Asn Gly Asp Thr Ala Val						
1		5		10		15
Glu Leu Asp Val Leu Lys Gly Thr Leu Gly Gln Asp Val Ile Asp Ile						
		20		25		30
Arg Thr Leu Gly Ser Lys Gly Val Phe Thr Phe Asp Pro Gly Phe Thr						
		35		40		45
Ser Thr Ala Ser Cys Glu Ser Lys Ile Thr Phe Ile Asp Gly Asp Glu						
50		55		60		
Gly Ile Leu Leu His Arg Gly Phe Pro Ile Asp Gln Leu Ala Thr Asp						
65		70		75		80
Ser Asn Tyr Leu Glu Val Cys Tyr Ile Leu Leu Asn Gly Glu Lys Pro						
		85		90		95
Thr Gln Glu Gln Tyr Asp Glu Phe Lys Thr Thr Val Thr Arg His Thr						
		100		105		110
Met Ile His Glu Gln Ile Thr Arg Leu Phe His Ala Phe Arg Arg Asp						
		115		120		125
Ser His Pro Met Ala Val Met Cys Gly Ile Thr Gly Ala Leu Ala Ala						
		130		135		140
Phe Tyr His Asp Ser Leu Asp Val Asn Asn Pro Arg His Arg Glu Ile						
145		150		155		160

-continued

Ala Ala Phe Arg Leu Leu Ser Lys Met Pro Thr Met Ala Ala Met Cys
165 170 175

Tyr Lys Tyr Ser Ile Gly Gln Pro Phe Val Tyr Pro Arg Asn Asp Leu
180 185 190

Ser Tyr Ala Gly Asn Phe Leu Asn Met Met Phe Ser Thr Pro Cys Glu
195 200 205

Pro Tyr Glu Val Asn Pro Ile Leu Glu Arg Ala Met Asp Arg Ile Leu
210 215 220

Ile Leu His Ala Asp His Glu Gln Asn Ala Ser Thr Ser Thr Val Arg
225 230 235 240

Thr Ala Gly Ser Ser Gly Ala Asn Pro Phe Ala Cys Ile Ala Ala Gly
245 250 255

Ile Ala Ser Leu Trp Gly Pro Ala His Gly Gly Ala Asn Glu Ala Ala
260 265 270

Leu Lys Met Leu Glu Glu Ile Ser Ser Val Lys His Ile Pro Glu Phe
275 280 285

Val Arg Arg Ala Lys Asp Lys Asn Asp Ser Phe Arg Leu Met Gly Phe
290 295 300

Gly His Arg Val Tyr Lys Asn Tyr Asp Pro Arg Ala Thr Val Met Arg
305 310 315 320

Glu Thr Cys His Glu Val Leu Lys Glu Leu Gly Thr Lys Asp Asp Leu
325 330 335

Leu Glu Val Ala Met Glu Leu Glu Asn Ile Ala Leu Asn Asp Pro Tyr
340 345 350

Phe Ile Glu Lys Lys Leu Tyr Pro Asn Val Asp Phe Tyr Ser Gly Ile
355 360 365

Ile Leu Lys Ala Met Gly Ile Pro Ser Ser Met Phe Thr Val Ile Phe
370 375 380

Ala Met Ala Arg Thr Val Gly Trp Ile Ala His Trp Ser Glu Met His
385 390 395 400

Ser Asp Gly Met Lys Ile Ala Arg Pro Arg Gln Leu Tyr Thr Gly Tyr
405 410 415

Glu Lys Arg Asp Phe Lys Ser Asp Ile Lys Arg
420 425

<210> SEQ ID NO 5

<211> LENGTH: 1284

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: GltA A145

<400> SEQUENCE: 5

```

atggctgata caaaagcaaa actcaccctc aacggggata cagctgttga actggatgtg    60
ctgaaaggca cgctgggtca agatgttatt gatatccgta ctctcgggtc aaaagggtgtg    120
ttcacctttg acccaggett cacttcaacc gcatcctgcg aatctaaaat tacttttatt    180
gatggtgatg aaggtathtt gctgcaccgc ggtttcccga tcgatcagct ggcgaccgat    240
tctaactacc tggaagtttg ttacatcctg ctgaatggtg aaaaaccgac tcaggaacag    300
tatgacgaat ttaaaactac ggtgaccctg cataccatga tccacgagca gattaccctg    360
ctgttccatg ctttccgtcg cgactcgcac ccaatggcag tcatgtgtgg tattaccggc    420
gcgctggcgg cgttcgcgca cgactcgcctg gatgttaaca atcctcgtca ccgtgaaatt    480
gccgcgttcc gcctgctgtc gaaaatgccg accatggccg cgatgtgtta caagtattcc    540

```


-continued

```

attggtcagc catttgttta cccgcgcaac gatctctcct acgccggtaa cttcctgaat 600
atgatgttct ccacgccgtg cgaaccgtat gaagttaatc cgattctgga acgtgctatg 660
gaccgtattc tgatcctgca cgctgaccat gaacagaacg cctctacctc caccgtgctg 720
accgctggct cttegggtgc gaaccggtt gctgtatcg cagcaggtat tgcttactg 780
tggggacctg cgcacggcgg tgctaacgaa gcggcgtga aaatgctgga agaaatcagc 840
tccgttaaac acattccgga atttgttcgt cgtgcgaaag acaaaaatga ttctttccgc 900
ctgatgggct tcggtcaccg cgtgtacaaa aattacgacc cgcgcgccac cgtaatgcgt 960
gaaacctgcc atgaagtgc gaaagagctg ggcacgaagg atgacctgct ggaagtggct 1020
atggagctgg aaaacatcgc gctgaacgac ccgtacttta tcgagaagaa actgtaccgc 1080
aacgtcgatt tctactctgg tatcatcctg aaagcgatgg gtattccgct tccatgttc 1140
accgtcattt tcgcaatggc acgtaccgtt ggctggatcg cccactggag cgaaatgcac 1200
agtgacggta tgaagattgc ccgtccgctg cagctgtata caggatatga aaaacgcgac 1260
tttaaaagcg atatcaagcg ttaa 1284

```

```

<210> SEQ ID NO 6
<211> LENGTH: 1284
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: GltA A167

```

```

<400> SEQUENCE: 6

```

```

atggctgata caaaagcaaa actcacctc aacggggata cagctgttga actggatgtg 60
ctgaaaggca cgctgggtca agatgttatt gatatccgta ctctcggttc aaaaggtgtg 120
ttcacctttg acccaggctt cacttcaacc gcacccctcg aatctaaaat tacttttatt 180
gatggatgat aaggtatatt gctgcaccgc ggtttccoga tcgatcagct ggcgaccgat 240
tctaactacc tggaagtgtg ttacatcctg ctgaatggtg aaaaaccgac tcaggaacag 300
tatgacgaat ttaaaactac ggtgaccctg cataccatga tccacgagca gattaccctg 360
ctgttccatg ctttccgctg cgactcgcac ccaatggcag tcatgtgtgg tattaccggc 420
gcgctggcgg cgttctatca cgactcgtg gatgttaaca atcctcgtca ccgtgaaatt 480
gccgcgttcc gcctgctgtc ggcgatgccg accatggcgg cgatgtgtta caagtattcc 540
attggtcagc catttgttta cccgcgcaac gatctctcct acgccggtaa cttcctgaat 600
atgatgttct ccacgccgtg cgaaccgtat gaagttaatc cgattctgga acgtgctatg 660
gaccgtattc tgatcctgca cgctgaccat gaacagaacg cctctacctc caccgtgctg 720
accgctggct cttegggtgc gaaccggtt gctgtatcg cagcaggtat tgcttactg 780
tggggacctg cgcacggcgg tgctaacgaa gcggcgtga aaatgctgga agaaatcagc 840
tccgttaaac acattccgga atttgttcgt cgtgcgaaag acaaaaatga ttctttccgc 900
ctgatgggct tcggtcaccg cgtgtacaaa aattacgacc cgcgcgccac cgtaatgcgt 960
gaaacctgcc atgaagtgc gaaagagctg ggcacgaagg atgacctgct ggaagtggct 1020
atggagctgg aaaacatcgc gctgaacgac ccgtacttta tcgagaagaa actgtaccgc 1080
aacgtcgatt tctactctgg tatcatcctg aaagcgatgg gtattccgct tccatgttc 1140
accgtcattt tcgcaatggc acgtaccgtt ggctggatcg cccactggag cgaaatgcac 1200
agtgacggta tgaagattgc ccgtccgctg cagctgtata caggatatga aaaacgcgac 1260

```


-continued

 tttaaagcg atatcaagcg ttaa 1284

<210> SEQ ID NO 7
 <211> LENGTH: 1284
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: GltA A204

<400> SEQUENCE: 7

atggctgata caaaagcaaa actcaccctc aacggggata cagctgttga actggatgtg 60
 ctgaaaggca cgctgggtca agatgttatt gatatccgta ctctcggttc aaaagggtgtg 120
 ttcacctttg acccaggctt cacttcaacc gcacccctgc aatctaaaat tacttttatt 180
 gatggatgat aaggatattt gctgcaccgc gggttcccga tcgatcagct ggccgaccgat 240
 tctaactacc tggaagtttg ttacatcctg ctgaatgggtg aaaaaccgac tcaggaacag 300
 tatgacgaat ttaaaactac ggtgaccctg cataccatga tccacgagca gattaccctg 360
 ctgttccatg ctttccgtcg cgactcgcac ccaatggcag tcatgtgtgg tattaccggc 420
 gcgctggcgg cgttctatca cgactcgtg gatgttaaca atcctcgtca ccgtgaaatt 480
 gccgcgttcc gcctgctgtc gaaaatgccg accatggccg cgatgtgtta caagtattcc 540
 attggtcagc catttgttta cccgcgcaac gatctctcct acgcccgtta cttcctgaat 600
 atgatgttct ccgcgcctgt cgaaccgat gaagttaatc cgattctgga acgtgctatg 660
 gaccgtattc tgatcctgca cgctgaccat gaacagaacg cctctacctc caccgtgcgt 720
 accgctggct cttcgggtgc gaaccggtt gcctgtatcg cagcaggtat tgcttcactg 780
 tggggacctg cgcacggcgg tgctaacgaa gcggcgtgaa aaatgctgga agaaatcagc 840
 tccgttaaac acattccgga atttgttctg cgtgcgaaag acaaaaatga ttctttccgc 900
 ctgatgggct tcggtcaccg cgtgtacaaa aattacgacc cgcgcgccac cgtaatgcgt 960
 gaaacctgcc atgaagtgtc gaaagagctg ggcacgaagg atgacctgct ggaagtggct 1020
 atggagctgg aaaacatcgc gctgaacgac ccgtacttta tcgagaagaa actgtaccgg 1080
 aacgtcattt tctactctgg tatcatcctg aaagcgatgg gtattccgtc ttccatgttc 1140
 accgtcattt tcgcaatggc acgtaccgtt ggctggatcg cccactggag cgaaatgcac 1200
 agtgacggta tgaagattgc ccgtccgctg cagctgtata caggatatga aaaacgagc 1260
 tttaaagcg atatcaagcg ttaa 1284

<210> SEQ ID NO 8
 <211> LENGTH: 1284
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: gene
 <222> LOCATION: (1)..(1284)
 <223> OTHER INFORMATION: GltA

<400> SEQUENCE: 8

atggctgata caaaagcaaa actcaccctc aacggggata cagctgttga actggatgtg 60
 ctgaaaggca cgctgggtca agatgttatt gatatccgta ctctcggttc aaaagggtgtg 120
 ttcacctttg acccaggctt cacttcaacc gcacccctgc aatctaaaat tacttttatt 180
 gatggatgat aaggatattt gctgcaccgc gggttcccga tcgatcagct ggccgaccgat 240
 tctaactacc tggaagtttg ttacatcctg ctgaatgggtg aaaaaccgac tcaggaacag 300
 tatgacgaat ttaaaactac ggtgaccctg cataccatga tccacgagca gattaccctg 360

-continued

```

ctgttccatg ctttccgtcg cgactcgcat ccaatggcag tcatgtgtgg tattaccggc 420
gcgctggcgg cgttctatca cgactcgctg gatgttaaca atcctegtca ccgtgaaatt 480
gccgcgttcc gcctgctgtc gaaaatgccg accatggccg cgatgtgtta caagtattcc 540
attggtcagc catttgttta cccgcgcaac gatctctcct acgcggtaa cttcctgaat 600
atgatgttct ccacgccgtg cgaaccgat gaagttaatc cgattctgga acgtgctatg 660
gaccgtattc tgatcctgca cgctgacat gaacagaacg cctctacctc caccgtgcgt 720
accgctggct cttcgggtgc gaaccggtt gcctgtatcg cagcaggtat tgettactg 780
tggggacctg cgcacggcgg tgctaacgaa gcggcgctga aaatgctgga agaaatcagc 840
tccgttaaac acattccgga atttgttcgt cgtgcgaaag acaaaaatga ttctttccgc 900
ctgatgggct tcggtcaccg cgtgtacaaa aattacgacc cgcgcgccac cgtaatgcgt 960
gaaacctgcc atgaagtgtc gaaagagctg ggacgaagg atgacctgct ggaagtggct 1020
atggagctgg aaaacatcgc gctgaacgac ccgtacttta tcgagaagaa actgtaccgg 1080
aacgtcgatt tctactctgg tatcactctg aaagcgatgg gtattccgtc ttccatgttc 1140
accgtcattt tcgcaatggc acgtaccgtt ggctggatcg cccactggag cgaaatgcac 1200
agtgcaggta tgaagattgc ccgtccgct cagctgtata caggatatga aaaacgcgac 1260
tttaaagcg atatcaagcg ttaa 1284

```

```

<210> SEQ ID NO 9
<211> LENGTH: 386
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(386)
<223> OTHER INFORMATION: metB

```

```

<400> SEQUENCE: 9

```

```

Met Thr Arg Lys Gln Ala Thr Ile Ala Val Arg Ser Gly Leu Asn Asp
1          5          10          15
Asp Glu Gln Tyr Gly Cys Val Val Pro Pro Ile His Leu Ser Ser Thr
          20          25          30
Tyr Asn Phe Thr Gly Phe Asn Glu Pro Arg Ala His Asp Tyr Ser Arg
          35          40          45
Arg Gly Asn Pro Thr Arg Asp Val Val Gln Arg Ala Leu Ala Glu Leu
          50          55          60
Glu Gly Gly Ala Gly Ala Val Leu Thr Asn Thr Gly Met Ser Ala Ile
65          70          75          80
His Leu Val Thr Thr Val Phe Leu Lys Pro Gly Asp Leu Leu Val Ala
          85          90          95
Pro His Asp Cys Tyr Gly Gly Ser Tyr Arg Leu Phe Asp Ser Leu Ala
          100         105         110
Lys Arg Gly Cys Tyr Arg Val Leu Phe Val Asp Gln Gly Asp Glu Gln
          115         120         125
Ala Leu Arg Ala Ala Leu Ala Glu Lys Pro Lys Leu Val Leu Val Glu
          130         135         140
Ser Pro Ser Asn Pro Leu Leu Arg Val Val Asp Ile Ala Lys Ile Cys
145         150         155         160
His Leu Ala Arg Glu Val Gly Ala Val Ser Val Val Asp Asn Thr Phe
          165         170         175
Leu Ser Pro Ala Leu Gln Asn Pro Leu Ala Leu Gly Ala Asp Leu Val

```


-continued

180				185				190							
Leu	His	Ser	Cys	Thr	Lys	Tyr	Leu	Asn	Gly	His	Ser	Asp	Val	Val	Ala
		195					200					205			
Gly	Val	Val	Ile	Ala	Lys	Asp	Pro	Asp	Val	Val	Thr	Glu	Leu	Ala	Trp
	210					215					220				
Trp	Ala	Asn	Asn	Ile	Gly	Val	Thr	Gly	Gly	Ala	Phe	Asp	Ser	Tyr	Leu
225					230					235					240
Leu	Leu	Arg	Gly	Leu	Arg	Thr	Leu	Val	Pro	Arg	Met	Glu	Leu	Ala	Gln
			245						250					255	
Arg	Asn	Ala	Gln	Ala	Ile	Val	Lys	Tyr	Leu	Gln	Thr	Gln	Pro	Leu	Val
		260						265					270		
Lys	Lys	Leu	Tyr	His	Pro	Ser	Leu	Pro	Glu	Asn	Gln	Gly	His	Glu	Ile
		275					280					285			
Ala	Ala	Arg	Gln	Gln	Lys	Gly	Phe	Gly	Ala	Met	Leu	Ser	Phe	Glu	Leu
	290					295					300				
Asp	Gly	Asp	Glu	Gln	Thr	Leu	Arg	Arg	Phe	Leu	Gly	Gly	Leu	Ser	Leu
305					310					315					320
Phe	Thr	Leu	Ala	Glu	Ser	Leu	Gly	Gly	Val	Glu	Ser	Leu	Ile	Ser	His
			325						330					335	
Ala	Ala	Thr	Met	Thr	His	Ala	Gly	Met	Ala	Pro	Glu	Ala	Arg	Ala	Ala
		340						345					350		
Ala	Gly	Ile	Ser	Glu	Thr	Leu	Leu	Arg	Ile	Ser	Thr	Gly	Ile	Glu	Asp
	355						360					365			
Gly	Glu	Asp	Leu	Ile	Ala	Asp	Leu	Glu	Asn	Gly	Phe	Arg	Ala	Ala	Asn
	370					375					380				
Lys	Gly														
385															

<210> SEQ ID NO 10
 <211> LENGTH: 1161
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: gene
 <222> LOCATION: (1)..(1161)
 <223> OTHER INFORMATION: metB

<400> SEQUENCE: 10

```

atgacgcgta aacaggccac catcgcagtg cgtagcgggt taaatgacga cgaacagtat    60
ggttgcggtg tcccaccgat ccatctttcc agcacctata actttaccgg atttaatgaa    120
ccgcgcgcgc atgattactc gcgtcgcggc aaccaaacgc gcgatgtggt tcagcgtgcg    180
ctggcagaac tggaaggtgg tgctggtgca gtacttacta ataccggcat gtccgcgatt    240
cacctggtaa cgaccgtctt tttgaaacct ggcgatctgc tggttgcgcc gcacgactgc    300
tacggcggta gctatcgctt gttcgacagt ctggcgaaac gcggttgcta tcgcgtggtg    360
tttgttgatc aaggcgatga acaggcatta cgggcagcgc tggcagaaaa acccaactg    420
gtactggtag aaagcccaag taatccattg ttacgcgctg tggatattgc gaaaatctgc    480
catctggcaa gggaagtccg ggcggtgagc gtggtggata acaccttctt aagcccggca    540
ttacaaaatc cgctggcatt aggtgccgat ctggtggtgc attcatgcac gaaatatctg    600
aacggtcact cagacgtagt ggccggcgtg gtgattgcta aagaccgga cgttgtcact    660
gaactggcct ggtgggcaaa caatattggc gtgacgggcy gcgcgtttga cagctatctg    720
ctgctacgtg ggttgccaac gctggtgccg cgtatggagc tggcgcagcg caacgcgcag    780
  
```

-continued

```

gcgattgtga aatacctgca aaccagccg ttggtgaaaa aactgtatca cccgtcgttg 840
ccggaaaatc aggggcatga aattgccgcg cgccagcaaa aaggctttgg cgcaatggtg 900
agttttgaac tggatggcga tgagcagacg ctgctgctgtt tcttggggcg gctgtcgttg 960
tttacgctgg cggaatcatt agggggagtg gaaagttaa tctctcacgc cgcaacctg 1020
acacatgcag gcatggcacc agaagcgcgt gctgccgccc ggatctccga gacgctgctg 1080
cgtatctcca ccggtattga agatggcgaa gatttaattg ccgacctgga aaatggcttc 1140
cgggctgcaa acaaggggta a 1161

```

```

<210> SEQ ID NO 11
<211> LENGTH: 310
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(310)
<223> OTHER INFORMATION: thrB

```

```

<400> SEQUENCE: 11

```

```

Met Val Lys Val Tyr Ala Pro Ala Ser Ser Ala Asn Met Ser Val Gly
1          5          10          15
Phe Asp Val Leu Gly Ala Ala Val Thr Pro Val Asp Gly Ala Leu Leu
20          25          30
Gly Asp Val Val Thr Val Glu Ala Ala Glu Thr Phe Ser Leu Asn Asn
35          40          45
Leu Gly Arg Phe Ala Asp Lys Leu Pro Ser Glu Pro Arg Glu Asn Ile
50          55          60
Val Tyr Gln Cys Trp Glu Arg Phe Cys Gln Glu Leu Gly Lys Gln Ile
65          70          75          80
Pro Val Ala Met Thr Leu Glu Lys Asn Met Pro Ile Gly Ser Gly Leu
85          90          95
Gly Ser Ser Ala Cys Ser Val Val Ala Ala Leu Met Ala Met Asn Glu
100         105         110
His Cys Gly Lys Pro Leu Asn Asp Thr Arg Leu Leu Ala Leu Met Gly
115        120        125
Glu Leu Glu Gly Arg Ile Ser Gly Ser Ile His Tyr Asp Asn Val Ala
130        135        140
Pro Cys Phe Leu Gly Gly Met Gln Leu Met Ile Glu Glu Asn Asp Ile
145        150        155        160
Ile Ser Gln Gln Val Pro Gly Phe Asp Glu Trp Leu Trp Val Leu Ala
165        170        175
Tyr Pro Gly Ile Lys Val Ser Thr Ala Glu Ala Arg Ala Ile Leu Pro
180        185        190
Ala Gln Tyr Arg Arg Gln Asp Cys Ile Ala His Gly Arg His Leu Ala
195        200        205
Gly Phe Ile His Ala Cys Tyr Ser Arg Gln Pro Glu Leu Ala Ala Lys
210        215        220
Leu Met Lys Asp Val Ile Ala Glu Pro Tyr Arg Glu Arg Leu Leu Pro
225        230        235        240
Gly Phe Arg Gln Ala Arg Gln Ala Val Ala Glu Ile Gly Ala Val Ala
245        250        255
Ser Gly Ile Ser Gly Ser Gly Pro Thr Leu Phe Ala Leu Cys Asp Lys
260        265        270
Pro Glu Thr Ala Gln Arg Val Ala Asp Trp Leu Gly Lys Asn Tyr Leu
275        280        285

```


-continued

Gln Asn Gln Glu Gly Phe Val His Ile Cys Arg Leu Asp Thr Ala Gly
 290 295 300

Ala Arg Val Leu Glu Asn
 305 310

<210> SEQ ID NO 12
 <211> LENGTH: 933
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: gene
 <222> LOCATION: (1)..(933)
 <223> OTHER INFORMATION: thrB

<400> SEQUENCE: 12

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 60
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 120
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 180
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 240
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 300
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 360
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 420
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 480
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 540
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 600
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 660
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 720
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 780
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 840
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 900
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaa 933

<210> SEQ ID NO 13
 <211> LENGTH: 379
 <212> TYPE: PRT
 <213> ORGANISM: Leptospira meyeri
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(379)
 <223> OTHER INFORMATION: homoserine O-acetyltransferase

<400> SEQUENCE: 13

Met Pro Thr Ser Glu Gln Asn Glu Phe Ser His Gly Ser Val Gly Val
 1 5 10 15

Val Tyr Thr Gln Ser Ile Arg Phe Glu Ser Leu Thr Leu Glu Gly Gly
 20 25 30

Glu Thr Ile Thr Pro Leu Glu Ile Ala Tyr Glu Thr Tyr Gly Thr Leu
 35 40 45

Asn Glu Lys Lys Asp Asn Ala Ile Leu Val Cys His Ala Leu Ser Gly
 50 55 60

Asp Ala His Ala Ala Gly Phe His Glu Gly Asp Lys Arg Pro Gly Trp
 65 70 75 80

Trp Asp Tyr Tyr Ile Gly Pro Gly Lys Ser Phe Asp Thr Asn Arg Tyr

-continued

85				90				95							
Phe	Ile	Ile	Ser	Ser	Asn	Val	Ile	Gly	Gly	Cys	Lys	Gly	Ser	Ser	Gly
			100					105					110		
Pro	Leu	Thr	Ile	Asn	Gly	Lys	Asn	Gly	Lys	Pro	Phe	Gln	Ser	Thr	Phe
		115					120					125			
Pro	Phe	Val	Ser	Ile	Gly	Asp	Met	Val	Asn	Ala	Gln	Glu	Lys	Leu	Ile
	130					135					140				
Ser	His	Phe	Gly	Ile	His	Lys	Leu	Phe	Ala	Val	Ala	Gly	Gly	Ser	Met
145					150					155					160
Gly	Gly	Met	Gln	Ala	Leu	Gln	Trp	Ser	Val	Ala	Tyr	Pro	Asp	Arg	Leu
			165						170					175	
Lys	Asn	Cys	Ile	Val	Met	Ala	Ser	Ser	Ser	Glu	His	Ser	Ala	Gln	Gln
		180						185					190		
Ile	Ala	Phe	Asn	Glu	Val	Gly	Arg	Gln	Ala	Ile	Leu	Ser	Asp	Pro	Asn
		195					200						205		
Trp	Asn	Gln	Gly	Leu	Tyr	Thr	Gln	Glu	Asn	Arg	Pro	Ser	Lys	Gly	Leu
	210					215					220				
Ala	Leu	Ala	Arg	Met	Met	Gly	His	Ile	Thr	Tyr	Leu	Ser	Asp	Glu	Met
225					230					235					240
Met	Arg	Glu	Lys	Phe	Gly	Arg	Lys	Pro	Pro	Lys	Gly	Asn	Ile	Gln	Ser
			245						250					255	
Thr	Asp	Phe	Ala	Val	Gly	Ser	Tyr	Leu	Ile	Tyr	Gln	Gly	Glu	Ser	Phe
		260						265					270		
Val	Asp	Arg	Phe	Asp	Ala	Asn	Ser	Tyr	Ile	Tyr	Val	Thr	Lys	Ala	Leu
		275					280					285			
Asp	His	Phe	Ser	Leu	Gly	Thr	Gly	Lys	Glu	Leu	Thr	Lys	Val	Leu	Ala
	290					295					300				
Lys	Val	Arg	Cys	Arg	Phe	Leu	Val	Val	Ala	Tyr	Thr	Ser	Asp	Trp	Leu
305					310					315					320
Tyr	Pro	Pro	Tyr	Gln	Ser	Glu	Glu	Ile	Val	Lys	Ser	Leu	Glu	Val	Asn
			325						330					335	
Ala	Val	Pro	Val	Ser	Phe	Val	Glu	Leu	Asn	Asn	Pro	Ala	Gly	Arg	His
		340						345					350		
Asp	Ser	Phe	Leu	Leu	Pro	Ser	Glu	Gln	Gln	Asp	Ser	Ile	Leu	Arg	Asp
		355					360					365			
Phe	Leu	Ser	Ser	Thr	Asp	Glu	Gly	Val	Phe	Leu					
	370					375									

<210> SEQ ID NO 14
 <211> LENGTH: 379
 <212> TYPE: PRT
 <213> ORGANISM: Corynebacterium glutamicum
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(379)
 <223> OTHER INFORMATION: homoserine O-acetyltransferase

<400> SEQUENCE: 14

Met	Pro	Thr	Leu	Ala	Pro	Ser	Gly	Gln	Leu	Glu	Ile	Gln	Ala	Ile	Gly
1				5					10					15	
Asp	Val	Ser	Thr	Glu	Ala	Gly	Ala	Ile	Ile	Thr	Asn	Ala	Glu	Ile	Ala
			20					25					30		
Tyr	His	Arg	Trp	Gly	Glu	Tyr	Arg	Val	Asp	Lys	Glu	Gly	Arg	Ser	Asn
		35					40					45			
Val	Val	Leu	Ile	Glu	His	Ala	Leu	Thr	Gly	Asp	Ser	Asn	Ala	Ala	Asp
		50				55					60				

-continued

Trp Trp Ala Asp Leu Leu Gly Pro Gly Lys Ala Ile Asn Thr Asp Ile
 65 70 75 80
 Tyr Cys Val Ile Cys Thr Asn Val Ile Gly Gly Cys Asn Gly Ser Thr
 85 90 95
 Gly Pro Gly Ser Met His Pro Asp Gly Asn Phe Trp Gly Asn Arg Phe
 100 105 110
 Pro Ala Thr Ser Ile Arg Asp Gln Val Asn Ala Glu Lys Gln Phe Leu
 115 120 125
 Asp Ala Leu Gly Ile Thr Thr Val Ala Ala Val Val Leu Leu Gly Gly
 130 135 140
 Ser Met Gly Gly Ala Arg Thr Leu Glu Trp Ala Ala Met Tyr Pro Glu
 145 150 155 160
 Thr Val Gly Ala Ala Ala Val Leu Ala Val Ser Ala Arg Ala Ser Ala
 165 170 175
 Trp Gln Ile Gly Ile Gln Ser Ala Gln Ile Lys Ala Ile Glu Asn Asp
 180 185 190
 His His Trp His Glu Gly Asn Tyr Tyr Glu Ser Gly Cys Asn Pro Ala
 195 200 205
 Thr Gly Leu Gly Ala Ala Arg Arg Ile Ala His Leu Thr Tyr Arg Gly
 210 215 220
 Glu Leu Glu Ile Asp Glu Arg Phe Gly Thr Lys Ala Gln Lys Asn Glu
 225 230 235 240
 Asn Pro Leu Gly Pro Tyr Arg Lys Pro Asp Gln Arg Phe Ala Val Glu
 245 250 255
 Ser Tyr Leu Asp Tyr Gln Ala Asp Lys Leu Val Gln Arg Phe Asp Ala
 260 265 270
 Gly Ser Tyr Val Leu Leu Thr Asp Ala Leu Asn Arg His Asp Ile Gly
 275 280 285
 Arg Asp Arg Gly Gly Leu Asn Lys Ala Leu Glu Ser Ile Lys Val Pro
 290 295 300
 Val Leu Val Ala Gly Val Asp Thr Asp Ile Leu Tyr Pro Tyr His Gln
 305 310 315 320
 Gln Glu His Leu Ser Arg Asn Leu Gly Asn Leu Leu Ala Met Ala Lys
 325 330 335
 Ile Val Ser Pro Val Gly His Asp Ala Phe Leu Thr Glu Ser Arg Gln
 340 345 350
 Met Asp Arg Ile Val Arg Asn Phe Phe Ser Leu Ile Ser Pro Asp Glu
 355 360 365
 Asp Asn Pro Ser Thr Tyr Ile Glu Phe Tyr Ile
 370 375

<210> SEQ ID NO 15
 <211> LENGTH: 334
 <212> TYPE: PRT
 <213> ORGANISM: Deinococcus radiodurans
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(334)
 <223> OTHER INFORMATION: homoserine O-acetyltransferase

<400> SEQUENCE: 15

Met Thr Ala Val Leu Ala Gly His Ala Ser Ala Leu Leu Leu Thr Glu
 1 5 10 15
 Glu Pro Asp Cys Ser Gly Pro Gln Thr Val Val Leu Phe Arg Arg Glu
 20 25 30

-continued

Pro Leu Leu Leu Asp Cys Gly Arg Ala Leu Ser Asp Val Arg Val Ala
 35 40 45

 Phe His Thr Tyr Gly Thr Pro Arg Ala Asp Ala Thr Leu Val Leu His
 50 55 60

 Ala Leu Thr Gly Asp Ser Ala Val His Glu Trp Trp Pro Asp Phe Leu
 65 70 75 80

 Gly Ala Gly Arg Pro Leu Asp Pro Ala Asp Asp Tyr Val Val Cys Ala
 85 90 95

 Asn Val Leu Gly Gly Cys Ala Gly Thr Thr Ser Ala Ala Glu Leu Ala
 100 105 110

 Ala Thr Cys Ser Gly Pro Val Pro Leu Ser Leu Arg Asp Met Ala Arg
 115 120 125

 Val Gly Arg Ala Leu Leu Asp Ser Leu Gly Val Arg Arg Val Arg Val
 130 135 140

 Ile Gly Ala Ser Met Gly Gly Met Leu Ala Tyr Ala Trp Leu Leu Glu
 145 150 155 160

 Cys Pro Asp Leu Val Glu Lys Ala Val Ile Ile Gly Ala Pro Ala Arg
 165 170 175

 His Ser Pro Trp Ala Ile Gly Leu Asn Thr Ala Ala Arg Ser Ala Ile
 180 185 190

 Ala Leu Ala Pro Gly Gly Glu Gly Leu Lys Val Ala Arg Gln Ile Ala
 195 200 205

 Met Leu Ser Tyr Arg Ser Pro Glu Ser Leu Ser Arg Thr Gln Ala Gly
 210 215 220

 Gln Arg Val Pro Gly Val Pro Ala Val Thr Ser Tyr Leu His Tyr Gln
 225 230 235 240

 Gly Glu Lys Leu Ala Ala Arg Phe Asp Glu Gln Thr Tyr Cys Ala Leu
 245 250 255

 Thr Trp Ala Met Asp Ala Phe Gln Pro Ser Ser Ala Asp Leu Lys Ala
 260 265 270

 Val Arg Ala Pro Val Leu Val Val Gly Ile Ser Ser Asp Leu Leu Tyr
 275 280 285

 Pro Ala Ala Glu Val Arg Ala Cys Ala Ala Glu Leu Pro His Ala Asp
 290 295 300

 Tyr Trp Glu Leu Gly Ser Ile His Gly His Asp Ala Phe Leu Met Asp
 305 310 315 320

 Pro Gln Asp Leu Pro Glu Arg Val Gly Ala Phe Leu Arg Ser
 325 330

<210> SEQ ID NO 16
 <211> LENGTH: 309
 <212> TYPE: PRT
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(309)
 <223> OTHER INFORMATION: homoserine O-succinyl transferase

 <400> SEQUENCE: 16

Met Pro Ile Arg Val Pro Asp Glu Leu Pro Ala Val Asn Phe Leu Arg
 1 5 10 15

 Glu Glu Asn Val Phe Val Met Thr Thr Ser Arg Ala Ser Gly Gln Glu
 20 25 30

 Ile Arg Pro Leu Lys Val Leu Ile Leu Asn Leu Met Pro Lys Lys Ile
 35 40 45

 Glu Thr Glu Asn Gln Phe Leu Arg Leu Leu Ser Asn Ser Pro Leu Gln

-continued

50	55	60
Val Asp Ile Gln Leu 65	Leu Arg Ile Asp Ser 70	Arg Glu Ser Arg Asn Thr 75 80
Pro Ala Glu His Leu 85	Asn Asn Phe Tyr Cys 90	Asn Phe Glu Asp Ile Gln 95
Asp Gln Asn Phe Asp 100	Gly Leu Ile Val Thr 105	Gly Ala Pro Leu Gly Leu 110
Val Glu Phe Asn Asp 115	Val Ala Tyr Trp Pro 120	Gln Ile Lys Gln Val Leu 125
Glu Trp Ser Lys Asp 130	His Val Thr Ser Thr 135	Leu Phe Val Cys Trp Ala 140
Val Gln Ala Ala Leu 145	Asn Ile Leu Tyr Gly 150	Ile Pro Lys Gln Thr Arg 155 160
Thr Glu Lys Leu Ser 165	Gly Val Tyr Glu His 170	His Ile Leu His Pro His 175
Ala Leu Leu Thr Arg 180	Gly Phe Asp Asp Ser 185	Phe Leu Ala Pro His Ser 190
Arg Tyr Ala Asp Phe 195	Pro Ala Ala Leu Ile 200	Arg Asp Tyr Thr Asp Leu 205
Glu Ile Leu Ala Glu 210	Thr Glu Glu Gly Asp 215	Ala Tyr Leu Phe Ala Ser 220
Lys Asp Lys Arg Ile 225	Ala Phe Val Thr Gly 230	His Pro Glu Tyr Asp Ala 235 240
Gln Thr Leu Ala Gln 245	Glu Phe Phe Arg Asp 250	Val Glu Ala Gly Leu Asp 255
Pro Asp Val Pro Tyr 260	Asn Tyr Phe Pro His 265	Asn Asp Pro Gln Asn Thr 270
Pro Arg Ala Ser Trp 275	Arg Ser His Gly Asn 280	Leu Leu Phe Thr Asn Trp 285
Leu Asn Tyr Tyr Val 290	Tyr Gln Ile Thr Pro 295	Tyr Asp Leu Arg His Met 300
Asn Pro Thr Leu Asp 305		

<210> SEQ ID NO 17

<211> LENGTH: 309

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Variant polypeptide having homoserine o-acetyl transferase activity, metA EL

<400> SEQUENCE: 17

Met Pro Ile Arg Val 1 5	Pro Asp Glu Leu Pro 10	Ala Val Asn Phe Leu Arg 15
Glu Glu Asn Val Phe 20	Val Met Thr Thr Ser 25	Arg Ala Ser Gly Gln Glu 30
Ile Arg Pro Leu Lys 35	Val Leu Ile Leu Asn 40	Leu Met Pro Lys Lys Ile 45
Glu Thr Glu Asn Gln 50	Phe Leu Arg Leu Leu 55	Ser Asn Ser Pro Leu Gln 60
Val Asp Ile Gln Leu 65	Leu Arg Ile Asp Ser 70	Arg Glu Ser Arg Asn Thr 75 80
Pro Ala Glu His Leu 85	Asn Asn Phe Tyr Cys 90	Asn Phe Glu Asp Ile Gln 95

-continued

Asp Gln Asn Phe Asp Gly Leu Ile Val Thr Gly Ala Pro Leu Glu Leu
 100 105 110
 Val Glu Phe Asn Asp Val Ala Tyr Trp Pro Gln Ile Lys Gln Val Leu
 115 120 125
 Glu Trp Ser Lys Asp His Val Thr Ser Thr Leu Phe Val Cys Trp Ala
 130 135 140
 Val Gln Ala Ala Leu Asn Ile Leu Tyr Gly Ile Pro Lys Gln Thr Arg
 145 150 155 160
 Thr Glu Lys Leu Ser Gly Val Tyr Glu His His Ile Leu His Pro His
 165 170 175
 Ala Leu Leu Thr Arg Gly Phe Asp Asp Ser Phe Leu Ala Pro His Ser
 180 185 190
 Arg Tyr Ala Asp Phe Pro Ala Ala Leu Ile Arg Asp Tyr Thr Asp Leu
 195 200 205
 Glu Ile Leu Ala Glu Thr Glu Glu Gly Asp Ala Tyr Leu Phe Ala Ser
 210 215 220
 Lys Asp Lys Arg Ile Ala Phe Val Thr Gly His Pro Glu Tyr Asp Ala
 225 230 235 240
 Gln Thr Leu Ala Gln Glu Phe Phe Arg Asp Val Glu Ala Gly Leu Asp
 245 250 255
 Pro Asp Val Pro Tyr Asn Tyr Phe Pro His Asn Asp Pro Gln Asn Thr
 260 265 270
 Pro Arg Ala Ser Trp Arg Ser His Gly Asn Leu Leu Phe Thr Asn Trp
 275 280 285
 Leu Asn Tyr Tyr Val Tyr Gln Ile Thr Pro Tyr Asp Leu Arg His Met
 290 295 300
 Asn Pro Thr Leu Asp
 305

<210> SEQ ID NO 18

<211> LENGTH: 309

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Variant polypeptide having homoserine O-acetyl transferase activity, meta ET

<400> SEQUENCE: 18

Met Pro Ile Arg Val Pro Asp Glu Leu Pro Ala Val Asn Phe Leu Arg
 1 5 10 15
 Glu Glu Asn Val Phe Val Met Thr Thr Ser Arg Ala Ser Gly Gln Glu
 20 25 30
 Ile Arg Pro Leu Lys Val Leu Ile Leu Asn Leu Met Pro Lys Lys Ile
 35 40 45
 Glu Thr Glu Asn Gln Phe Leu Arg Leu Leu Ser Asn Ser Pro Leu Gln
 50 55 60
 Val Asp Ile Gln Leu Leu Arg Ile Asp Ser Arg Glu Ser Arg Asn Thr
 65 70 75 80
 Pro Ala Glu His Leu Asn Asn Phe Tyr Cys Asn Phe Glu Asp Ile Gln
 85 90 95
 Asp Gln Asn Phe Asp Gly Leu Ile Val Thr Gly Ala Pro Leu Glu Thr
 100 105 110
 Val Glu Phe Asn Asp Val Ala Tyr Trp Pro Gln Ile Lys Gln Val Leu
 115 120 125
 Glu Trp Ser Lys Asp His Val Thr Ser Thr Leu Phe Val Cys Trp Ala
 130 135 140

-continued

Val Gln Ala Ala Leu Asn Ile Leu Tyr Gly Ile Pro Lys Gln Thr Arg
 145 150 155 160

Thr Glu Lys Leu Ser Gly Val Tyr Glu His His Ile Leu His Pro His
 165 170 175

Ala Leu Leu Thr Arg Gly Phe Asp Asp Ser Phe Leu Ala Pro His Ser
 180 185 190

Arg Tyr Ala Asp Phe Pro Ala Ala Leu Ile Arg Asp Tyr Thr Asp Leu
 195 200 205

Glu Ile Leu Ala Glu Thr Glu Glu Gly Asp Ala Tyr Leu Phe Ala Ser
 210 215 220

Lys Asp Lys Arg Ile Ala Phe Val Thr Gly His Pro Glu Tyr Asp Ala
 225 230 235 240

Gln Thr Leu Ala Gln Glu Phe Phe Arg Asp Val Glu Ala Gly Leu Asp
 245 250 255

Pro Asp Val Pro Tyr Asn Tyr Phe Pro His Asn Asp Pro Gln Asn Thr
 260 265 270

Pro Arg Ala Ser Trp Arg Ser His Gly Asn Leu Leu Phe Thr Asn Trp
 275 280 285

Leu Asn Tyr Tyr Val Tyr Gln Ile Thr Pro Tyr Asp Leu Arg His Met
 290 295 300

Asn Pro Thr Leu Asp
 305

<210> SEQ ID NO 19

<211> LENGTH: 309

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Variant polypeptide having homoserine O-acetyl transferase activity, meta EH

<400> SEQUENCE: 19

Met Pro Ile Arg Val Pro Asp Glu Leu Pro Ala Val Asn Phe Leu Arg
 1 5 10 15

Glu Glu Asn Val Phe Val Met Thr Thr Ser Arg Ala Ser Gly Gln Glu
 20 25 30

Ile Arg Pro Leu Lys Val Leu Ile Leu Asn Leu Met Pro Lys Lys Ile
 35 40 45

Glu Thr Glu Asn Gln Phe Leu Arg Leu Leu Ser Asn Ser Pro Leu Gln
 50 55 60

Val Asp Ile Gln Leu Leu Arg Ile Asp Ser Arg Glu Ser Arg Asn Thr
 65 70 75 80

Pro Ala Glu His Leu Asn Asn Phe Tyr Cys Asn Phe Glu Asp Ile Gln
 85 90 95

Asp Gln Asn Phe Asp Gly Leu Ile Val Thr Gly Ala Pro Leu Glu His
 100 105 110

Val Glu Phe Asn Asp Val Ala Tyr Trp Pro Gln Ile Lys Gln Val Leu
 115 120 125

Glu Trp Ser Lys Asp His Val Thr Ser Thr Leu Phe Val Cys Trp Ala
 130 135 140

Val Gln Ala Ala Leu Asn Ile Leu Tyr Gly Ile Pro Lys Gln Thr Arg
 145 150 155 160

Thr Glu Lys Leu Ser Gly Val Tyr Glu His His Ile Leu His Pro His
 165 170 175

Ala Leu Leu Thr Arg Gly Phe Asp Asp Ser Phe Leu Ala Pro His Ser

-continued

180	185	190
Arg Tyr Ala Asp Phe Pro Ala Ala Leu Ile Arg Asp Tyr Thr Asp Leu 195 200 205		
Glu Ile Leu Ala Glu Thr Glu Glu Gly Asp Ala Tyr Leu Phe Ala Ser 210 215 220		
Lys Asp Lys Arg Ile Ala Phe Val Thr Gly His Pro Glu Tyr Asp Ala 225 230 235 240		
Gln Thr Leu Ala Gln Glu Phe Phe Arg Asp Val Glu Ala Gly Leu Asp 245 250 255		
Pro Asp Val Pro Tyr Asn Tyr Phe Pro His Asn Asp Pro Gln Asn Thr 260 265 270		
Pro Arg Ala Ser Trp Arg Ser His Gly Asn Leu Leu Phe Thr Asn Trp 275 280 285		
Leu Asn Tyr Tyr Val Tyr Gln Ile Thr Pro Tyr Asp Leu Arg His Met 290 295 300		
Asn Pro Thr Leu Asp 305		

<210> SEQ ID NO 20
 <211> LENGTH: 883
 <212> TYPE: PRT
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(883)
 <223> OTHER INFORMATION: phosphoenolpyruvate carboxylase

<400> SEQUENCE: 20

Met Asn Glu Gln Tyr Ser Ala Leu Arg Ser Asn Val Ser Met Leu Gly 1 5 10 15
Lys Val Leu Gly Glu Thr Ile Lys Asp Ala Leu Gly Glu His Ile Leu 20 25 30
Glu Arg Val Glu Thr Ile Arg Lys Leu Ser Lys Ser Ser Arg Ala Gly 35 40 45
Asn Asp Ala Asn Arg Gln Glu Leu Leu Thr Thr Leu Gln Asn Leu Ser 50 55 60
Asn Asp Glu Leu Leu Pro Val Ala Arg Ala Phe Ser Gln Phe Leu Asn 65 70 75 80
Leu Ala Asn Thr Ala Glu Gln Tyr His Ser Ile Ser Pro Lys Gly Glu 85 90 95
Ala Ala Ser Asn Pro Glu Val Ile Ala Arg Thr Leu Arg Lys Leu Lys 100 105 110
Asn Gln Pro Glu Leu Ser Glu Asp Thr Ile Lys Lys Ala Val Glu Ser 115 120 125
Leu Ser Leu Glu Leu Val Leu Thr Ala His Pro Thr Glu Ile Thr Arg 130 135 140
Arg Thr Leu Ile His Lys Met Val Glu Val Asn Ala Cys Leu Lys Gln 145 150 155 160
Leu Asp Asn Lys Asp Ile Ala Asp Tyr Glu His Asn Gln Leu Met Arg 165 170 175
Arg Leu Arg Gln Leu Ile Ala Gln Ser Trp His Thr Asp Glu Ile Arg 180 185 190
Lys Leu Arg Pro Ser Pro Val Asp Glu Ala Lys Trp Gly Phe Ala Val 195 200 205
Val Glu Asn Ser Leu Trp Gln Gly Val Pro Asn Tyr Leu Arg Glu Leu 210 215 220

-continued

Asn Glu Gln Leu Glu Glu Asn Leu Gly Tyr Lys Leu Pro Val Glu Phe
 225 230 235 240
 Val Pro Val Arg Phe Thr Ser Trp Met Gly Gly Asp Arg Asp Gly Asn
 245 250 255
 Pro Asn Val Thr Ala Asp Ile Thr Arg His Val Leu Leu Leu Ser Arg
 260 265 270
 Trp Lys Ala Thr Asp Leu Phe Leu Lys Asp Ile Gln Val Leu Val Ser
 275 280 285
 Glu Leu Ser Met Val Glu Ala Thr Pro Glu Leu Leu Ala Leu Val Gly
 290 295 300
 Glu Glu Gly Ala Ala Glu Pro Tyr Arg Tyr Leu Met Lys Asn Leu Arg
 305 310 315 320
 Ser Arg Leu Met Ala Thr Gln Ala Trp Leu Glu Ala Arg Leu Lys Gly
 325 330 335
 Glu Glu Leu Pro Lys Pro Glu Gly Leu Leu Thr Gln Asn Glu Glu Leu
 340 345 350
 Trp Glu Pro Leu Tyr Ala Cys Tyr Gln Ser Leu Gln Ala Cys Gly Met
 355 360 365
 Gly Ile Ile Ala Asn Gly Asp Leu Leu Asp Thr Leu Arg Arg Val Lys
 370 375 380
 Cys Phe Gly Val Pro Leu Val Arg Ile Asp Ile Arg Gln Glu Ser Thr
 385 390 395 400
 Arg His Thr Glu Ala Leu Gly Glu Leu Thr Arg Tyr Leu Gly Ile Gly
 405 410 415
 Asp Tyr Glu Ser Trp Ser Glu Ala Asp Lys Gln Ala Phe Leu Ile Arg
 420 425 430
 Glu Leu Asn Ser Lys Arg Pro Leu Leu Pro Arg Asn Trp Gln Pro Ser
 435 440 445
 Ala Glu Thr Arg Glu Val Leu Asp Thr Cys Gln Val Ile Ala Glu Ala
 450 455 460
 Pro Gln Gly Ser Ile Ala Ala Tyr Val Ile Ser Met Ala Lys Thr Pro
 465 470 475 480
 Ser Asp Val Leu Ala Val His Leu Leu Leu Lys Glu Ala Gly Ile Gly
 485 490 495
 Phe Ala Met Pro Val Ala Pro Leu Phe Glu Thr Leu Asp Asp Leu Asn
 500 505 510
 Asn Ala Asn Asp Val Met Thr Gln Leu Leu Asn Ile Asp Trp Tyr Arg
 515 520 525
 Gly Leu Ile Gln Gly Lys Gln Met Val Met Ile Gly Tyr Ser Asp Ser
 530 535 540
 Ala Lys Asp Ala Gly Val Met Ala Ala Ser Trp Ala Gln Tyr Gln Ala
 545 550 555 560
 Gln Asp Ala Leu Ile Lys Thr Cys Glu Lys Ala Gly Ile Glu Leu Thr
 565 570 575
 Leu Phe His Gly Arg Gly Gly Ser Ile Gly Arg Gly Gly Ala Pro Ala
 580 585 590
 His Ala Ala Leu Leu Ser Gln Pro Pro Gly Ser Leu Lys Gly Gly Leu
 595 600 605
 Arg Val Thr Glu Gln Gly Glu Met Ile Arg Phe Lys Tyr Gly Leu Pro
 610 615 620
 Glu Ile Thr Val Ser Ser Leu Ser Leu Tyr Thr Gly Ala Ile Leu Glu
 625 630 635 640

-continued

Ala Asn Leu Leu Pro Pro Pro Glu Pro Lys Glu Ser Trp Arg Arg Ile
645 650 655

Met Asp Glu Leu Ser Val Ile Ser Cys Asp Val Tyr Arg Gly Tyr Val
660 665 670

Arg Glu Asn Lys Asp Phe Val Pro Tyr Phe Arg Ser Ala Thr Pro Glu
675 680 685

Gln Glu Leu Gly Lys Leu Pro Leu Gly Ser Arg Pro Ala Lys Arg Arg
690 695 700

Pro Thr Gly Gly Val Glu Ser Leu Arg Ala Ile Pro Trp Ile Phe Ala
705 710 715 720

Trp Thr Gln Asn Arg Leu Met Leu Pro Ala Trp Leu Gly Ala Gly Thr
725 730 735

Ala Leu Gln Lys Val Val Glu Asp Gly Lys Gln Ser Glu Leu Glu Ala
740 745 750

Met Cys Arg Asp Trp Pro Phe Phe Ser Thr Arg Leu Gly Met Leu Glu
755 760 765

Met Val Phe Ala Lys Ala Asp Leu Trp Leu Ala Glu Tyr Tyr Asp Gln
770 775 780

Arg Leu Val Asp Lys Ala Leu Trp Pro Leu Gly Lys Glu Leu Arg Asn
785 790 795 800

Leu Gln Glu Glu Asp Ile Lys Val Val Leu Ala Ile Ala Asn Asp Ser
805 810 815

His Leu Met Ala Asp Leu Pro Trp Ile Ala Glu Ser Ile Gln Leu Arg
820 825 830

Asn Ile Tyr Thr Asp Pro Leu Asn Val Leu Gln Ala Glu Leu Leu His
835 840 845

Arg Ser Arg Gln Ala Glu Lys Glu Gly Gln Glu Pro Asp Pro Arg Val
850 855 860

Glu Gln Ala Leu Met Val Thr Ile Ala Gly Ile Ala Ala Gly Met Arg
865 870 875 880

Asn Thr Gly

<210> SEQ ID NO 21
<211> LENGTH: 396
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: PEPTIDE
<222> LOCATION: (1)..(396)
<223> OTHER INFORMATION: aspartate aminotransferase

<400> SEQUENCE: 21

Met Phe Glu Asn Ile Thr Ala Ala Pro Ala Asp Pro Ile Leu Gly Leu
1 5 10 15

Ala Asp Leu Phe Arg Ala Asp Glu Arg Pro Gly Lys Ile Asn Leu Gly
20 25 30

Ile Gly Val Tyr Lys Asp Glu Thr Gly Lys Thr Pro Val Leu Thr Ser
35 40 45

Val Lys Lys Ala Glu Gln Tyr Leu Leu Glu Asn Glu Thr Thr Lys Asn
50 55 60

Tyr Leu Gly Ile Asp Gly Ile Pro Glu Phe Gly Arg Cys Thr Gln Glu
65 70 75 80

Leu Leu Phe Gly Lys Gly Ser Ala Leu Ile Asn Asp Lys Arg Ala Arg
85 90 95

Thr Ala Gln Thr Pro Gly Gly Thr Gly Ala Leu Arg Val Ala Ala Asp
100 105 110

-continued

Phe Leu Ala Lys Asn Thr Ser Val Lys Arg Val Trp Val Ser Asn Pro
 115 120 125
 Ser Trp Pro Asn His Lys Ser Val Phe Asn Ser Ala Gly Leu Glu Val
 130 135 140
 Arg Glu Tyr Ala Tyr Tyr Asp Ala Glu Asn His Thr Leu Asp Phe Asp
 145 150 155 160
 Ala Leu Ile Asn Ser Leu Asn Glu Ala Gln Ala Gly Asp Val Val Leu
 165 170 175
 Phe His Gly Cys Cys His Asn Pro Thr Gly Ile Asp Pro Thr Leu Glu
 180 185 190
 Gln Trp Gln Thr Leu Ala Gln Leu Ser Val Glu Lys Gly Trp Leu Pro
 195 200 205
 Leu Phe Asp Phe Ala Tyr Gln Gly Phe Ala Arg Gly Leu Glu Glu Asp
 210 215 220
 Ala Glu Gly Leu Arg Ala Phe Ala Ala Met His Lys Glu Leu Ile Val
 225 230 235 240
 Ala Ser Ser Tyr Ser Lys Asn Phe Gly Leu Tyr Asn Glu Arg Val Gly
 245 250 255
 Ala Cys Thr Leu Val Ala Ala Asp Ser Glu Thr Val Asp Arg Ala Phe
 260 265 270
 Ser Gln Met Lys Ala Ala Ile Arg Ala Asn Tyr Ser Asn Pro Pro Ala
 275 280 285
 His Gly Ala Ser Val Val Ala Thr Ile Leu Ser Asn Asp Ala Leu Arg
 290 295 300
 Ala Ile Trp Glu Gln Glu Leu Thr Asp Met Arg Gln Arg Ile Gln Arg
 305 310 315 320
 Met Arg Gln Leu Phe Val Asn Thr Leu Gln Glu Lys Gly Ala Asn Arg
 325 330 335
 Asp Phe Ser Phe Ile Ile Lys Gln Asn Gly Met Phe Ser Phe Ser Gly
 340 345 350
 Leu Thr Lys Glu Gln Val Leu Arg Leu Arg Glu Glu Phe Gly Val Tyr
 355 360 365
 Ala Val Ala Ser Gly Arg Val Asn Val Ala Gly Met Thr Pro Asp Asn
 370 375 380
 Met Ala Pro Leu Cys Glu Ala Ile Val Ala Val Leu
 385 390 395

<210> SEQ ID NO 22
 <211> LENGTH: 367
 <212> TYPE: PRT
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: PEPTIDE
 <222> LOCATION: (1)..(367)
 <223> OTHER INFORMATION: aspartate-semialdehyde dehydrogenase

<400> SEQUENCE: 22

Met Lys Asn Val Gly Phe Ile Gly Trp Arg Gly Met Val Gly Ser Val
 1 5 10 15
 Leu Met Gln Arg Met Val Glu Glu Arg Asp Phe Asp Ala Ile Arg Pro
 20 25 30
 Val Phe Phe Ser Thr Ser Gln Leu Gly Gln Ala Ala Pro Ser Phe Gly
 35 40 45
 Gly Thr Thr Gly Thr Leu Gln Asp Ala Phe Asp Leu Glu Ala Leu Lys
 50 55 60

-continued

Ala	Leu	Asp	Ile	Ile	Val	Thr	Cys	Gln	Gly	Gly	Asp	Tyr	Thr	Asn	Glu
65					70					75					80
Ile	Tyr	Pro	Lys	Leu	Arg	Glu	Ser	Gly	Trp	Gln	Gly	Tyr	Trp	Ile	Asp
			85						90					95	
Ala	Ala	Ser	Ser	Leu	Arg	Met	Lys	Asp	Asp	Ala	Ile	Ile	Ile	Leu	Asp
			100					105						110	
Pro	Val	Asn	Gln	Asp	Val	Ile	Thr	Asp	Gly	Leu	Asn	Asn	Gly	Ile	Arg
		115					120					125			
Thr	Phe	Val	Gly	Gly	Asn	Cys	Thr	Val	Ser	Leu	Met	Leu	Met	Ser	Leu
	130					135					140				
Gly	Gly	Leu	Phe	Ala	Asn	Asp	Leu	Val	Asp	Trp	Val	Ser	Val	Ala	Thr
145					150					155					160
Tyr	Gln	Ala	Ala	Ser	Gly	Gly	Gly	Ala	Arg	His	Met	Arg	Glu	Leu	Leu
				165					170					175	
Thr	Gln	Met	Gly	His	Leu	Tyr	Gly	His	Val	Ala	Asp	Glu	Leu	Ala	Thr
			180					185					190		
Pro	Ser	Ser	Ala	Ile	Leu	Asp	Ile	Glu	Arg	Lys	Val	Thr	Thr	Leu	Thr
		195					200					205			
Arg	Ser	Gly	Glu	Leu	Pro	Val	Asp	Asn	Phe	Gly	Val	Pro	Leu	Ala	Gly
	210					215					220				
Ser	Leu	Ile	Pro	Trp	Ile	Asp	Lys	Gln	Leu	Asp	Asn	Gly	Gln	Ser	Arg
225					230					235					240
Glu	Glu	Trp	Lys	Gly	Gln	Ala	Glu	Thr	Asn	Lys	Ile	Leu	Asn	Thr	Ser
				245					250					255	
Ser	Val	Ile	Pro	Val	Asp	Gly	Leu	Cys	Val	Arg	Val	Gly	Ala	Leu	Arg
			260					265					270		
Cys	His	Ser	Gln	Ala	Phe	Thr	Ile	Lys	Leu	Lys	Lys	Asp	Val	Ser	Ile
		275					280					285			
Pro	Thr	Val	Glu	Glu	Leu	Leu	Ala	Ala	His	Asn	Pro	Trp	Ala	Lys	Val
	290					295					300				
Val	Pro	Asn	Asp	Arg	Glu	Ile	Thr	Met	Arg	Glu	Leu	Thr	Pro	Ala	Ala
305					310					315					320
Val	Thr	Gly	Thr	Leu	Thr	Thr	Pro	Val	Gly	Arg	Leu	Arg	Lys	Leu	Asn
				325					330					335	
Met	Gly	Pro	Glu	Phe	Leu	Ser	Ala	Phe	Thr	Val	Gly	Asp	Gln	Leu	Leu
		340						345					350		
Trp	Gly	Ala	Ala	Glu	Pro	Leu	Arg	Arg	Met	Leu	Arg	Gln	Leu	Ala	
		355					360					365			

<210> SEQ ID NO 23
 <211> LENGTH: 933
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: gene
 <222> LOCATION: (1)..(933)
 <223> OTHER INFORMATION: homoserine O-succinyl transferase
 <400> SEQUENCE: 23

atggttaaag tttatgcccc ggcttcagc gccaatatga gcgctcgggtt tgatgtgctc	60
ggggcggcgg tgacacctgt tgatggtgca ttgctcggag atgtagtcac gggtgaggcg	120
gcagagacat tcagtctcaa caacctcgga cgctttgccc ataagctgcc gtcagaacca	180
cgggaaaata tcgtttatca gtgctgggag cgcttttgcc aggaactggg taagcaaatt	240
ccagtggcga tgacctgga aaagaatatg ccgatcgggt cgggcttagg ctccagtgcc	300

-continued

```

tggtcgggtgg tcgcgccgct gatggcgatg aatgaacact gcggcaagcc gcttaatgac 360
actcgtttgc tggtttgat gggcgagctg gaaggccgta tctccggcag cattcattac 420
gacaacgtgg caccgtgttt tctcgggtgg atgcagttga tgatcgaaga aaacgacatc 480
atcagccagc aagtgccagg gtttgatgag tggctgtggg tgctggcgtg tccggggatt 540
aaagtctcga cggcagaagc cagggctatt ttaccggcgc agtatcgccg ccaggattgc 600
attgcgcacg ggcgacatct ggcaggcttc attcacgcct gctattcccg tcagcctgag 660
cttgcccgga agctgatgaa agatgttatc gctgaaccct accgtgaacg gttactgcca 720
ggcttccggc aggcgcggca ggcggtcgcg gaaatcgccg cggtagcgag cggtatctcc 780
ggctccggcc cgacctgtt cgctctgtgt gacaagccgg aaaccgccc ggcggttgc 840
gactggttgg gtaagaacta cctgcaaaat caggaagggt ttgttcatat ttgccggctg 900
gatacggcgg gcgcacgagt actggaaaac taa 933

```

<210> SEQ ID NO 24

<211> LENGTH: 930

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Variant polypeptide having homoserine O-acetyl transferase activity, meta EL

<400> SEQUENCE: 24

```

atgccgattc gtgtgccgga cgagctaccc gccgtcaatt tcttgctga agaaaacgtc 60
tttgtgatga caacttctcg tgcgtctggt caggaaattc gtccacttaa ggttctgatc 120
cttaacctga tgccgaagaa gattgaaact gaaaatcagt ttctgcgctt gctttcaaac 180
tcacctttgc aggtcgatat tcagctgttg cgcacgatt cccgtgaatc gcgcaacacg 240
cccgcagagc atctgaacaa cttctactgt aactttgaag atattcagga tcagaacttt 300
gacggtttga ttgtaactgg tgcgcccgtg gaactgggtg agtttaatga tgctgcttac 360
tggccgcaga tcaaacaggt gctggagtgg tcgaaagatc acgtcacctc gacgctgttt 420
gtctgctggg cggtaacagg cgcgctcaat atcctctacg gcattcctaa gcaaaactcg 480
accgaaaaac tctctggcgt ttacgagcat catattctcc atcctcatgc gcttctgacg 540
cgtggctttg atgattcatt cctggcaccg cattcgcgct atgctgactt tccggcagcg 600
ttgattcgtg attacaccga tctggaaatt ctggcagaga cggaagaagg ggatgcatat 660
ctgtttgcca gtaaagataa gcgcattgcc tttgtgacgg gccatcccga atatgatgcg 720
caaacgctgg cgcaggaatt tttccgcat gtggaagccg gactagaccc ggatgtaccg 780
tataactatt tcccgcacaa tgatccgcaa aatacacgcg gagcgagctg gcgtagtcac 840
ggtaatttac tgtttaccaa ctggctcaac tattacgtct accagatcac gccatagat 900
ctacggcaca tgaatccaac gctggattaa 930

```

<210> SEQ ID NO 25

<211> LENGTH: 930

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Variant polypeptide having homoserine O-acetyl transferase activity, meta ET

<400> SEQUENCE: 25

```

atgccgattc gtgtgccgga cgagctaccc gccgtcaatt tcttgctga agaaaacgtc 60
tttgtgatga caacttctcg tgcgtctggt caggaaattc gtccacttaa ggttctgatc 120

```

-continued

```

cttaacctga tgccgaagaa gattgaaact gaaaatcagt ttctgcgct gctttcaaac 180
tcacctttgc aggtcgatat tcagctggtg cgcacgatt cccgtgaatc gcgcaacacg 240
cccgagagc atctgaacaa cttctactgt aactttgaag atattcagga tcagaacttt 300
gacggtttga ttgtaactgg tgcgcccgtg gaaaccgtgg agtttaatga tgctgcttac 360
tggccgcaga tcaaacaggt gctggagtgg tcgaaagatc acgtcacctc gacgctgttt 420
gtctgctggg cggtagagc cgcgctcaat atcctctacg gcattcctaa gcaaactcgc 480
accgaaaaac tctctggcgt ttacgagcat catattctcc atcctcatgc gcttctgacg 540
cgtggctttg atgattcatt cctggcaccg cattcgcgct atgctgactt tccggcagcg 600
ttgattcgtg attacaccga tctggaaatt ctggcagaga cggaagaagg ggatgcatat 660
ctgtttgcca gtaaagataa ggcattgcc tttgtgacgg gccatcccga atatgatgcg 720
caaacgctgg cgcaggaatt tttccgcat gtggaagccg gactagaccg ggatgtaccg 780
tataactatt tcccgcacaa tgatccgcaa aatacaccgc gagcgagctg gcgtagtcac 840
ggtaatttac tgtttaccaa ctggctcaac tattacgtct accagatcac gccatagat 900
ctacggcaca tgaatccaac gctggattaa 930

```

<210> SEQ ID NO 26

<211> LENGTH: 930

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Variant polypeptide having homoserine O-acetyl transferase activity, meta EH

<400> SEQUENCE: 26

```

atgccgattc gtgtgccgga cgagctaccc gccgtcaatt tcttgcgta agaaaacgtc 60
tttgtgatga caacttctcg tgcgctcgtt caggaaatc gtccacttaa ggttctgatc 120
cttaacctga tgccgaagaa gattgaaact gaaaatcagt ttctgcgct gctttcaaac 180
tcacctttgc aggtcgatat tcagctggtg cgcacgatt cccgtgaatc gcgcaacacg 240
cccgagagc atctgaacaa cttctactgt aactttgaag atattcagga tcagaacttt 300
gacggtttga ttgtaactgg tgcgcccgtg gaacatgtgg agtttaatga tgctgcttac 360
tggccgcaga tcaaacaggt gctggagtgg tcgaaagatc acgtcacctc gacgctgttt 420
gtctgctggg cggtagagc cgcgctcaat atcctctacg gcattcctaa gcaaactcgc 480
accgaaaaac tctctggcgt ttacgagcat catattctcc atcctcatgc gcttctgacg 540
cgtggctttg atgattcatt cctggcaccg cattcgcgct atgctgactt tccggcagcg 600
ttgattcgtg attacaccga tctggaaatt ctggcagaga cggaagaagg ggatgcatat 660
ctgtttgcca gtaaagataa ggcattgcc tttgtgacgg gccatcccga atatgatgcg 720
caaacgctgg cgcaggaatt tttccgcat gtggaagccg gactagaccg ggatgtaccg 780
tataactatt tcccgcacaa tgatccgcaa aatacaccgc gagcgagctg gcgtagtcac 840
ggtaatttac tgtttaccaa ctggctcaac tattacgtct accagatcac gccatagat 900
ctacggcaca tgaatccaac gctggattaa 930

```

<210> SEQ ID NO 27

<211> LENGTH: 2652

<212> TYPE: DNA

<213> ORGANISM: Escherichia coli

<220> FEATURE:

<221> NAME/KEY: gene

-continued

<222> LOCATION: (1)..(2652)

<223> OTHER INFORMATION: phosphoenolpyruvate carboxylase

<400> SEQUENCE: 27

```

atgaacgaac aatattccgc attgcgtagt aatgtcagta tgctcggcaa agtgctggga      60
gaaaccatca aggatgcggt gggagaacac attcttgaac gcgtagaaac tatccgtaag     120
ttgtcgaaat cttcacgcgc tggcaatgat gctaaccgcc aggagttgct caccacctta     180
caaaatttgt cgaacgacga gctgctgccc gttgcgcgtg cgtttagtca gttcctgaac     240
ctggccaaca ccgccgagca ataccacagc atttcgccga aaggcgaagc tgccagcaac     300
ccggaagtga tcgcccgcac cctgcgtaaa ctgaaaaacc agccggaact gagcgaagac     360
accatcaaaa aagcagtgga atcgctgtcg ctggaactgg tcttcacggc tcaccaacc     420
gaaattaccg gtcgtacact gatccacaaa atggtggaag tgaacgcctg tttaaaacag     480
ctcgataaca aagatatcgc tgactacgaa cacaaccagc tgatgcgtcg cctgcgccag     540
ttgatcgccc agtcatggca taccgatgaa atccgtaagc tgcgtccaag cccggtagat     600
gaagccaaat ggggctttgc cgtagtggaa aacagcctgt ggcaaggcgt accaaattac     660
ctgcgcgaac tgaacgaaca actggaagag aacctcggct acaaactgcc cgtcgaattt     720
gttcgcgtcc gttttacttc gtggatgggc ggcgaccgcg acggcaaccg gaacgtcact     780
gccgatatca cccgccacgt cctgctactc agccgctgga aagccaccga tttgttctctg     840
aaagatattc aggtgctggt ttctgaactg tcgatggttg aagcgacccc tgaactgctg     900
gcgctggttg gcgaagaagg tgccgcagaa ccgatcgcct atctgatgaa aaacctgcgt     960
tctcgcctga tggcgacaca ggcattggctg gaagcgcgcc tgaaaggcga agaactgcca    1020
aaaccagaag gcctgctgac acaaaacgaa gaactgtggg aaccgctcta cgcttgctac    1080
cagtcacttc aggcgtgtgg catgggtatt atcgccaacg gcgatctgct cgacaccctg    1140
cgccgcgtga aatgtttcgg cgtaccgctg gtccgtattg atatcctca ggagagcacg    1200
cgtcataccg aagcgcgtgg cgagctgacc cgctacctcg gtatcggcga ctacgaaagc    1260
tggtcagagg ccgacaaaca ggcgttctctg atccgcgaac tgaactcca acgtccgctt    1320
ctgccgcgca actggcaacc aagcgcgcaa acgcgcgaag tgctcgatac ctgccaggtg    1380
attgccgaag caccgcaagg ctccattgcc gcctacgtga tctcgatggc gaaaacgccc    1440
tccgacgtac tggctgtcca cctgctgctg aaagaagcgg gtatcggggt tgcgatgccg    1500
gttgctccgc tgtttgaaac cctcgatgat ctgaacaacg ccaacgatgt catgaccag    1560
ctgctcaata ttgactggtg tcgtggcctg attcagggca aacagatggt gatgattggc    1620
tattccgact cagcaaaaga tgccgggagt atggcagctt cctgggcgca atatcaggca    1680
caggatgcat taatcaaac ctgcgaaaaa gcgggtattg agctgacgtt gttccacggc    1740
cgcgccgggt ccattggtcg cggcggcgca cctgctcatg cggcgcgtgt gtcacaaccg    1800
ccaggaagcc tgaaaggcgg cctgcgcgta accgaacagg gcgagatgat ccgctttaa    1860
tatggtctgc cagaaatcac cgtcagcagc ctgtcgcttt ataccggggc gattctggaa    1920
gccaacctgc tgccaccgcc ggagccgaaa gagagctggc gtcgcattat ggatgaactg    1980
tcagtcactc cctgcgatgt ctaccgcggc tacgtacgtg aaaacaaaga ttttgtgcct    2040
tacttccgct ccgctacgcc ggaacaagaa ctgggcaaac tgccggtggg ttcacgtccg    2100
gcgaaaacgc gcccaaccgg cggcgtcgag tcaactacgc ccattccgtg gatcttcgcc    2160
tggacgcaaa accgtctgat gctccccgcc tggctgggtg caggtacggc gctgcaaaaa    2220

```

-continued

```

gtggtcgaag acggcaaaca gagcgagctg gaggctatgt gccgcgattg gccattcttc 2280
tcgacgcgctc tcggcatgct ggagatggtc ttcgccaaag cagacctgtg gctggcgaa 2340
tactatgacc aacgcctggg agacaaagca ctgtggccgt taggtaaaga gttacgcaac 2400
ctgcaagaag aagacatcaa agtgggtgctg gcgattgcca acgattccca tctgatggcc 2460
gatctgccgt ggattgcaga gtctattcag ctacggaata tttacaccga cccgctgaac 2520
gtattgcagg ccgagttgct gcaccgctcc cgccaggcag aaaaagaagg ccaggaaccg 2580
gatcctcgcg tcgaacaagc gttaatggtc actattgccc ggattgcggc aggtatgcgt 2640
aataccggct aa 2652

```

```

<210> SEQ ID NO 28
<211> LENGTH: 1191
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: gene
<222> LOCATION: (1)..(1191)
<223> OTHER INFORMATION: aspartate aminotransferase

```

```

<400> SEQUENCE: 28

```

```

atgtttgaga acattaccgc cgctcctgcc gacccgattc tgggcctggc cgatctgttt 60
cgtgccgatg aacgtcccgg caaaattaac ctcgggattg gtgtctataa agatgagacg 120
ggcaaaacc cggactgac cagcgtgaaa aaggctgaac agtatctgct cgaaaatgaa 180
accacaaaa attacctcgg cattgacggc atccctgaat ttggtegetg cactcaggaa 240
ctgctgtttg gtaaaggtag cgccctgatc aatgacaaac gtgctcgcac ggcacagact 300
ccggggggca ctggcgact acgcgtggct gccgatttcc tggcaaaaa taccagcggt 360
aagcgtgtgt ggggtgagcaa cccaagctgg ccgaaccata agagcgtctt taactctgca 420
ggtctggaag ttcgtgaata cgcttattat gatgcccgaat atcacactct tgacttcgat 480
gcactgatta acagcctgaa tgaagctcag gctggcgacg tagtgctgtt ccatggctgc 540
tgccataacc caaccggtat cgaccctacg ctggaacaat ggcaaacact ggcacaactc 600
tccgttgaga aaggctggtt accgctgttt gacttcgctt accaggggtt tgcccgtggt 660
ctggaagaag atgctgaagg actgcgcgct ttcgcccgtc tcgataaaga gctgattggt 720
gccagttcct actctaaaaa ctttggcctg tacaacgagc gtggtggcgc ttgtactctg 780
gttgctgccg acagtgaaac cgttgatcgc gcattcagcc aatgaaagc ggcgattcgc 840
gctaaactact ctaaccacc agcacacggc gcttctgttg ttgccaccat cctgagcaac 900
gatgcgttac gtgcgatttg ggaacaagag ctgactgata tgcgccagcg tattcagcgt 960
atgcgtcagt tgttcgtcaa tacgctgcag gaaaaaggcg caaacgcga cttcagcttt 1020
atcatcaaac agaacggcat gttctccttc agtggcctga caaaagaaca agtgctgcgt 1080
ctgcgcgaag agtttggcgt atatgcccgt gcttctggtc gcgtaaagt ggccgggatg 1140
acaccagata acatggtccc gctgtgcaa gcgattgtgg cagtgtgta a 1191

```

```

<210> SEQ ID NO 29
<211> LENGTH: 1104
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: gene
<222> LOCATION: (1)..(1104)
<223> OTHER INFORMATION: aspartate-semialdehyde dehydrogenase

```

```

<400> SEQUENCE: 29

```


-continued

```

atgaaaaatg ttggttttat cggctggcgc ggtatggctg gctccgttct catgcaacgc      60
atggttgaag agcgcgactt cgacgccatt cgccctgtct tcttttctac ttctcagctt     120
ggccaggctg cgccgtcttt tggcggaacc actggcacac ttcaggatgc ctttgatctg     180
gaggcgctaa aggccctcga tatcattgtg acctgtcagg gcggcgatta taccaacgaa     240
atctatccaa agcttcgtga aagcggatgg caaggttact ggattgacgc agcatcgtct     300
ctgcgcgatga aagatgacgc catcatcatt cttgaccccg tcaatcagga cgtcattacc     360
gacggattaa ataatggcat caggactttt gttggcggtg actgtaccgt aagcctgatg     420
ttgatgtcgt tgggtggttt attcgccaat gatcttggtg attgggtgtc cgttgcaacc     480
taccaggccg cttccggcgg tggcgcgga catatgcgtg agttattaac ccagatgggc     540
catctgtatg gccatgtggc agatgaactc gcgaccccg cctctgctat tctcgatata     600
gaacgcaaag tcacaacctt aaccgtagc ggtgagctgc cggtgataa ctttggcgtg     660
ccgctggcgg gtagcctgat tccgtggatc gacaaacagc tcgataacgg tcagagccgc     720
gaagagtggg aagggcaggc ggaaaccaac aagatcctca acacatcttc cgtaattccg     780
gtagatggtt tatgtgtgcg tgtcggggca ttgcgctgcc acagccaggc attcactatt     840
aaattgaaaa aagatgtgtc tattccgacc gtggaagaac tgctggctgc gcacaatccg     900
tgggcgaaaag tcgttccgaa cgatcgggaa atcactatgc gtgagctaac cccagctgcc     960
gttaccggca cgctgaccac gccggtaggc cgctgcgta agctgaatat gggaccagag    1020
ttcctgtcag cctttaccgt gggcgaccag ctgctgtggg gggccgcgga gccgctgcgt    1080
cggatgcttc gtcaactggc gtaa                                           1104

```

```

<210> SEQ ID NO 30
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

```

```

<400> SEQUENCE: 30

```

```

ttactctggt gcctgacatt tcaccgacaa agcccaggga acttcatcac gtgtaggctg      60
gagctgcttc                                           70

```

```

<210> SEQ ID NO 31
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

```

```

<400> SEQUENCE: 31

```

```

ttacccttg tttgcagccc ggaagccatt ttccaggctg gcaattaaat catatgaata      60
tcctccttag                                           70

```

```

<210> SEQ ID NO 32
<211> LENGTH: 70
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

```

```

<400> SEQUENCE: 32

```

```

aaagaatatg ccgatcgtt cgggcttagg ctccagtgcc tggtcgtgg gtgtaggctg      60

```

-continued

gagctgcttc 70

<210> SEQ ID NO 33
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 33

agacaaccga catcgctttc aacattggcg accggagccg ggaaggcaaa catatgaata 60

tcctccttag 70

<210> SEQ ID NO 34
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 34

aattgatatc atgccgattc gtgtgccgg 29

<210> SEQ ID NO 35
 <211> LENGTH: 34
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 35

aattaagctt ttaatccagc gttggattca tgtg 34

<210> SEQ ID NO 36
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 36

ttgtaactgg tgcgccgctg gaactggtgg ggtttaatga tgtc 44

<210> SEQ ID NO 37
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 37

gacatcatta aacccacca gttccagcgg cgcaccagtt acaa 44

<210> SEQ ID NO 38
 <211> LENGTH: 44
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 38

tgtaactggt gcgccgctgg aacatgtggg gtttaatgat gtcg 44

<210> SEQ ID NO 39
 <211> LENGTH: 44

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 39

 cgacatcatt aaacccaca tgttccagcg ggcaccagt taca 44

<210> SEQ ID NO 40
 <211> LENGTH: 42
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 40

 tttccgaaac gtacctcagc aggtgtaggc tggagctgct tc 42

<210> SEQ ID NO 41
 <211> LENGTH: 43
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 41

 gaataaaatt tattcacctg ctgcatatga atatacctcct tag 43

<210> SEQ ID NO 42
 <211> LENGTH: 29
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 42

 aattgatatc atgccgattc gtgtgcccgg 29

<210> SEQ ID NO 43
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 43

 aattaagcct gctgaggtac gtttcgg 27

<210> SEQ ID NO 44
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 44

 cagcaggtga ataaatttta ttc 23

<210> SEQ ID NO 45
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 45

-continued

 cgcgaaatgga agctgtttcc 20

<210> SEQ ID NO 46
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 46

gccggaattc tgtcggatgc gatacttgcg c 31

<210> SEQ ID NO 47
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 47

gaaggagctc agaaaaccct cgcgcaaaag 30

<210> SEQ ID NO 48
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 48

aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa a 31

<210> SEQ ID NO 49
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 49

gaagggtacc agaaaaccct cgcgcaaaag 30

<210> SEQ ID NO 50
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 50

tccgagctca taagcgtagc gcatcaggca 30

<210> SEQ ID NO 51
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

<400> SEQUENCE: 51

tccgagctcg tccacctatg ttgactacat 30

<210> SEQ ID NO 52
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 52

 ccggaattcc caggagagca ataagca 27

 <210> SEQ ID NO 53
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 53

 ctagtctaga tgctctatatt aactcccg 28

 <210> SEQ ID NO 54
 <211> LENGTH: 28
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 54

 ctagtctaga ccaggagagc aataagca 28

 <210> SEQ ID NO 55
 <211> LENGTH: 27
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 55

 ccggaattct gctctattta actcccg 27

 <210> SEQ ID NO 56
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 56

 aggcgctaag gagaccttaa atggctgata caaaagcaaa actcacctc tgtaggctgg 60
 agctgcttcg 70

 <210> SEQ ID NO 57
 <211> LENGTH: 70
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 57

 gggttaaat attacaact tagcaatcaa ccattaacgc ttgatatcgc atgggaatta 60
 gccatggtcc 70

 <210> SEQ ID NO 58
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

-continued

<400> SEQUENCE: 58
cctttctata actgcgcgctc at 22

<210> SEQ ID NO 59
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 59
aggggtatag ctacgccaga a 21

<210> SEQ ID NO 60
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 60
gaaggcaaat ttaagttccg 20

<210> SEQ ID NO 61
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 61
cgaagcagct ccagcctaca ggtatagata gacgtcattt 40

<210> SEQ ID NO 62
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 62
cggcgcgctg gcggcgctcg cgcacgactc gctggatggt aa 42

<210> SEQ ID NO 63
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 63
ttaacatcca gcgagtcgtg cgccaacgcc gccagcgcgc cg 42

<210> SEQ ID NO 64
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 64
cgcggtccgc ctgctgtcgg cgatgccgac catggccgcg at 42

<210> SEQ ID NO 65
<211> LENGTH: 42

-continued

<212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 65

 atcgcggcca tggtcggcat cgccgacagc aggcggaacg cg 42

 <210> SEQ ID NO 66
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 66

 tcttgaatat gatgttctcc gcgccgtgcg aaccgtatga 40

 <210> SEQ ID NO 67
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 67

 tcatacgggtt cgcacggcgc ggagaacatc atattcagga 40

 <210> SEQ ID NO 68
 <211> LENGTH: 40
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 68

 aaatgacgtc tatctatacc tgtaggctgg agctgcttcg 40

 <210> SEQ ID NO 69
 <211> LENGTH: 76
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 69

 gcaaaactaa attattggta tcatgaattt gttgtatgat gaataaaata taggggatgg 60
 gaattagcca tgggcc 76

 <210> SEQ ID NO 70
 <211> LENGTH: 67
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Primer

 <400> SEQUENCE: 70

 ccatggagga ggaattaacc atgcagtggg ggtgggtggg gtgcgataac agctgtatcc 60
 ccgttga 67

 <210> SEQ ID NO 71
 <211> LENGTH: 67
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 71

aagcttagga ggaattaacc atgcagtggg ggtgggtggg gtgcgatgca aatttaagtt 60

ccggcag 67

The invention claimed is:

1. A microorganism of *Escherichia* sp. producing O-acetyl homoserine, wherein the endogenous activity of citrate synthase is attenuated or inactivated, and a homoserine O-acetyltransferase is further introduced or enhanced, or an endogenous homoserine O-succinyltransferase is further modified to a polypeptide having 95% identity or above with SEQ ID NO: 16 and further having mutations at the G111E position and L112T or L112H positions to have the activity of homoserine O-acetyltransferase, wherein the microorganism has an improved O-acetyl homoserine production capability compared to a corresponding unmodified *Escherichia* sp. microorganism.

2. The microorganism of *Escherichia* sp. according to claim 1, wherein the microorganism with the attenuated endogenous activity of citrate synthase has an amino acid sequence of SEQ ID NO: 1 or SEQ ID NO: 2.

10

3. The microorganism of *Escherichia* sp. according to claim 1, wherein the activity of cystathionine gamma synthase, homoserine kinase, or both is further attenuated or inactivated compared to their endogenous activities.

15

4. The microorganism of *Escherichia* sp. according to claim 1, wherein the activity of at least one protein selected from the group consisting of phosphoenolpyruvate carboxylase, aspartate aminotransferase, and aspartate semialdehyde dehydrogenase is further introduced or enhanced.

20

5. The microorganism of *Escherichia* sp. according to claim 1, wherein the microorganism is *Escherichia coli*.

6. A method for producing O-acetyl homoserine, comprising:

25

- (a) culturing the microorganism of any of claims 1 to 3, 4 and 5; and
- (b) recovering O-acetyl homoserine produced during the cultivation of the microorganism.

* * * * *