



US011052063B2

(12) **United States Patent**  
**Osterloh et al.**

(10) **Patent No.:** **US 11,052,063 B2**  
(45) **Date of Patent:** **Jul. 6, 2021**

- (54) **METHODS OF REDUCING RLP-C**
- (71) Applicant: **Amarin Pharmaceuticals Ireland Limited**, Dublin (IE)
- (72) Inventors: **Ian Osterloh**, Kent (GB); **Pierre Wicker**, Mystic, CT (US); **Rene Braeckman**, Richboro, PA (US); **Paresh Soni**, Mystic, CT (US); **Mehar Manku**, Birmingham (GB)
- (73) Assignee: **Amarin Pharmaceuticals Ireland Limited**, Dublin (IE)
- (\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.
- (21) Appl. No.: **16/713,461**
- (22) Filed: **Dec. 13, 2019**
- (65) **Prior Publication Data**  
US 2020/0121630 A1 Apr. 23, 2020

5,760,081 A 6/1998 Leaf et al.  
5,763,496 A 6/1998 Holland  
5,776,978 A 7/1998 Bruzzese  
5,792,795 A 8/1998 Buser et al.  
5,837,731 A 11/1998 Vaddadi  
5,840,944 A 11/1998 Furihata et al.  
5,886,037 A 3/1999 Klor et al.  
5,888,541 A 3/1999 Horrobin et al.  
5,948,818 A 9/1999 Buser et al.  
6,025,008 A 2/2000 Akahoshi  
6,069,168 A 5/2000 Horrobin et al.  
6,193,999 B1 2/2001 Gennadios  
6,207,699 B1 3/2001 Rothman  
6,284,268 B1 9/2001 Mishra et al.  
6,313,330 B1 11/2001 Kiyohara et al.  
6,326,031 B1 12/2001 Hsia et al.  
6,326,355 B1 12/2001 Abbruzzese et al.  
6,331,568 B1 12/2001 Horrobin  
6,362,236 B1 3/2002 Aviram  
6,368,621 B1 4/2002 Engel et al.  
6,383,482 B1 5/2002 Gorsek  
6,384,077 B1 5/2002 Peet  
6,479,544 B1 11/2002 Horrobin  
6,482,421 B2 11/2002 Weidner  
6,531,150 B1 3/2003 Sunohara et al.  
6,555,700 B1 4/2003 Horrobin et al.  
6,596,766 B1 7/2003 Igarashi et al.  
6,620,821 B2 9/2003 Robl

(Continued)

**Related U.S. Application Data**

- (63) Continuation of application No. 14/737,115, filed on Jun. 11, 2015, now Pat. No. 10,561,631.
- (60) Provisional application No. 62/077,478, filed on Nov. 10, 2014, provisional application No. 62/010,847, filed on Jun. 11, 2014.
- (51) **Int. Cl.**  
*A61K 31/232* (2006.01)  
*A61K 31/00* (2006.01)
- (52) **U.S. Cl.**  
CPC ..... *A61K 31/232* (2013.01); *A61K 31/00* (2013.01)
- (58) **Field of Classification Search**  
None  
See application file for complete search history.

**References Cited**

## U.S. PATENT DOCUMENTS

4,377,526 A 3/1983 Fujita et al.  
4,526,902 A 7/1985 Rubin  
4,920,098 A 4/1990 Cotter et al.  
4,935,243 A 6/1990 Borkan et al.  
5,013,443 A 5/1991 Higashidate et al.  
5,116,871 A 5/1992 Horrobin et al.  
5,178,873 A 1/1993 Horrobin et al.  
5,198,468 A 3/1993 Horrobin  
5,215,630 A 6/1993 Hata et al.  
5,252,333 A 10/1993 Horrobin  
5,343,389 A 8/1994 Otvos  
5,385,929 A 1/1995 Bjorge et al.  
5,457,130 A 10/1995 Tisdale et al.  
5,502,077 A 3/1996 Breivik et al.  
5,567,730 A 10/1996 Miyashita et al.  
5,589,508 A 12/1996 Schlotzer et al.  
5,603,959 A 2/1997 Horrobin et al.  
5,618,558 A 4/1997 Horrobin et al.  
5,656,667 A 8/1997 Breivik et al.  
5,698,594 A 12/1997 Breivik et al.

## FOREIGN PATENT DOCUMENTS

CA 2628305 5/2007  
CA 2653787 12/2007

(Continued)

## OTHER PUBLICATIONS

A study of AMR101 to evaluate its ability to reduce cardiovascular events in high risk patients with hypertriglyceridemia and on statin (REDUCE-IT). Available at: <http://clinicaltrials.gov/show/NCT01492361>. (3 pages).

Aarsetoey H, Gurndt H, Nygaard O. The Role of Long-Chained Marine N-3 Polyunsaturated Fatty Acids in Cardiovascular Disease. *Cardiol Res Pract.* 2012. Epub Dec. 13, 2012.

Aarsland, et al., "On the Effect of Peroxisomal beta-Oxidation and Carnitine Palmitoyltransferase Activity by Eicosapentaenoic Acid in Live and Heart of Rats." *Lipids*, 25:546-548, (Sep. 1990).

Aas, V., et al., "Eicosapentaenoic acid (20:5 n-3) increases fatty acid and glucose uptake in cultured human skeletal muscle cells." *Journal of Lipid Research*, 47:366-374 (Feb. 2006).

Abbey, M., et al., "Effect of fish oil on lipoproteins, lecithin:cholesterol acyltransferase, and lipidtransfer protein activity in humans." *Arterioscler. Thromb. Vasc. Biol.* 10:85-94 (Jan./Feb. 1990).

(Continued)

*Primary Examiner* — Savitha M Rao*Assistant Examiner* — Gregg Polansky(74) *Attorney, Agent, or Firm* — Perkins Coie LP(57) **ABSTRACT**

In various embodiments, the present invention provides methods of treating and/or preventing cardiovascular-related disease and, in particular, a method of blood lipid therapy comprising administering to a subject in need thereof a pharmaceutical composition comprising eicosapentaenoic acid or a derivative thereof.

**10 Claims, 3 Drawing Sheets**

(56)

## References Cited

## U.S. PATENT DOCUMENTS

6,689,812 B2	2/2004	Peet	2002/0055539 A1	5/2002	Bockow et al.
6,846,942 B2	1/2005	Rubin	2002/0077361 A1	6/2002	Peet et al.
7,022,713 B2	4/2006	Aoki et al.	2002/0169209 A1	11/2002	Horrobin
7,112,609 B2	9/2006	Hermelin et al.	2002/0183389 A1	12/2002	Peet
7,119,118 B2	10/2006	Peet	2002/0193439 A1	12/2002	Peet et al.
7,179,491 B1	2/2007	Mag	2002/0198177 A1	12/2002	Horrobin et al.
7,205,329 B2	4/2007	Chien et al.	2003/0100610 A1	5/2003	Shibuya
7,405,302 B2	7/2008	Hutchinson et al.	2003/0104048 A1	6/2003	Patel et al.
7,498,359 B2	3/2009	Yokoyama et al.	2003/0161918 A1	8/2003	Kendrick et al.
7,511,131 B2	3/2009	Crooke et al.	2003/0166614 A1	9/2003	Harrison
7,598,227 B2	10/2009	Crooke et al.	2003/0232385 A1	12/2003	Breit et al.
7,642,287 B2	1/2010	Guzman	2004/0009208 A1	1/2004	Edson
7,776,881 B2	8/2010	Aoki et al.	2004/0018248 A1	1/2004	Bendich
8,188,146 B2	5/2012	Peet et al.	2004/0048919 A1	3/2004	Dreon et al.
8,293,727 B2	10/2012	Manku et al.	2004/0062847 A1	4/2004	Koiki et al.
8,293,728 B2	10/2012	Manku et al.	2004/0077723 A1	4/2004	Granata
8,298,554 B2	10/2012	Manku	2004/0106591 A1	6/2004	Pacioretty et al.
8,314,086 B2	11/2012	Manku et al.	2004/0121000 A1	6/2004	Bowe et al.
8,318,715 B2	11/2012	Manku et al.	2004/0162348 A1	8/2004	Peet et al.
8,324,195 B2	12/2012	Manku et al.	2004/0204356 A1	10/2004	Guenzler-Pukall
8,357,677 B1	1/2013	Manku et al.	2004/0258645 A1	12/2004	Trejo et al.
8,367,652 B2	2/2013	Manku et al.	2005/0042214 A1	2/2005	Gershwin et al.
8,377,920 B2	2/2013	Manku et al.	2005/0137253 A1	6/2005	Phinney et al.
8,410,086 B2	4/2013	Osterloh et al.	2005/0147665 A1	7/2005	Horrobin et al.
8,431,560 B1	4/2013	Manku et al.	2005/0187292 A1	8/2005	Aoki et al.
8,440,650 B1	5/2013	Manku et al.	2005/0244367 A1	11/2005	Hui et al.
8,455,472 B2	6/2013	Osterloh et al.	2005/0272095 A1	12/2005	Wang
8,518,929 B2	8/2013	Manku et al.	2006/0034815 A1	2/2006	Guzman et al.
8,524,698 B2	9/2013	Manku et al.	2006/0051418 A1	3/2006	Cowen et al.
8,546,372 B2	10/2013	Manku et al.	2006/0088502 A1	4/2006	Sata et al.
8,551,521 B2	10/2013	Manku et al.	2006/0111437 A1	5/2006	Aoki et al.
8,563,608 B2	10/2013	Manku et al.	2006/0134178 A1	6/2006	Doisaki et al.
8,617,593 B2	12/2013	Manku et al.	2006/0134206 A1	6/2006	Iyer et al.
8,617,594 B2	12/2013	Manku et al.	2006/0135607 A1	6/2006	Kobayashi et al.
8,618,168 B2	12/2013	Fujii et al.	2006/0135610 A1	6/2006	Bortz et al.
8,623,406 B2	1/2014	Manku et al.	2006/0141022 A1	6/2006	Kawamura et al.
8,642,077 B2	2/2014	Manku et al.	2006/0142390 A1	6/2006	Manku et al.
8,669,245 B2	3/2014	Osterloh et al.	2006/0172012 A1	8/2006	Finley et al.
8,680,144 B2	3/2014	Osterloh et al.	2006/0189682 A1	8/2006	Payne et al.
8,691,871 B2	4/2014	Osterloh et al.	2006/0211749 A1	9/2006	Bobotas et al.
8,703,185 B2	4/2014	Manku et al.	2006/0211761 A1	9/2006	Kumar et al.
8,709,475 B2	4/2014	Manku et al.	2006/0211762 A1	9/2006	Rongen
8,802,718 B2	8/2014	Yokoyama et al.	2006/0211763 A1	9/2006	Fawzy et al.
8,853,256 B2	10/2014	Yokoyama et al.	2006/0217356 A1	9/2006	Wright et al.
8,906,964 B2	12/2014	Bobotas et al.	2006/0223838 A1	10/2006	Jiang et al.
9,006,285 B2	4/2015	Ohnishi	2006/0252833 A1	11/2006	Peet et al.
9,060,981 B2	6/2015	Sato et al.	2007/0021504 A1	1/2007	Yokoyama et al.
9,138,415 B2	9/2015	Manku et al.	2007/0060532 A1	3/2007	Junien et al.
9,452,121 B2	9/2016	Manku et al.	2007/0098787 A1	5/2007	Kakiuchi
9,452,150 B2	9/2016	Ueshima et al.	2007/0104779 A1	5/2007	Rongen et al.
9,603,826 B2	3/2017	Soni	2007/0105793 A1	5/2007	Hendrix
9,610,272 B2	4/2017	Soni	2007/0105954 A1	5/2007	Puri
9,623,001 B2	4/2017	Soni	2007/0141138 A1	6/2007	Feuerstein et al.
9,693,984 B2	7/2017	Soni	2007/0167520 A1	7/2007	Bruzzese
9,693,985 B2	7/2017	Soni	2007/0185198 A1	8/2007	Yokoyama et al.
9,693,986 B2	7/2017	Soni	2007/0191467 A1	8/2007	Rongen et al.
9,700,537 B2	7/2017	Yokoyama et al.	2007/0202159 A1	8/2007	Mathur et al.
9,855,237 B2	1/2018	Osterloh et al.	2007/0212411 A1	9/2007	Fawzy et al.
9,918,954 B2	3/2018	Soni	2007/0219271 A1	9/2007	Mittmann et al.
10,058,521 B2	8/2018	Bobotas et al.	2007/0265340 A1	11/2007	Shalwitz et al.
10,166,209 B2	1/2019	Manku et al.	2007/0269507 A1	11/2007	Sachetto et al.
10,220,013 B2	3/2019	Osterloh et al.	2007/0292501 A1	12/2007	Udell
10,265,290 B2	4/2019	Manku et al.	2008/0020018 A1	1/2008	Moodley et al.
10,278,935 B2	5/2019	Soni	2008/0057115 A1	3/2008	Okamoto
10,555,925 B1	2/2020	Soni	2008/0085911 A1	4/2008	Rongen et al.
10,557,856 B2	2/2020	Singbartl et al.	2008/0089876 A1	4/2008	Cavazza
10,561,631 B2 *	2/2020	Osterloh	2008/0113046 A1	5/2008	Gardette
10,568,861 B1	2/2020	Soni	2008/0125490 A1	5/2008	Svensson et al.
10,576,054 B1	3/2020	Soni	2008/0139604 A1	6/2008	Fitzpatrick et al.
2001/0035125 A1	11/2001	Talieh et al.	2008/0185198 A1	8/2008	Jones
2002/0016312 A1	2/2002	Seed et al.	2008/0200453 A1	8/2008	Cincotta
2002/0025983 A1	2/2002	Horrobin	2008/0200547 A1	8/2008	Peet et al.
2002/0035125 A1	3/2002	Shear	2008/0200707 A1	8/2008	Shimano et al.
2002/0054871 A1	5/2002	Huang	2008/0214531 A1	9/2008	Saxena et al.
2002/0055529 A1	5/2002	Bisgaier et al.	2008/0299187 A1	12/2008	Opheim et al.
			2008/0306154 A1	12/2008	Svensson et al.
			2008/0319077 A1	12/2008	Suzuki et al.
			2009/0012167 A1	1/2009	Rongen et al.
			2009/0018125 A1	1/2009	Mittmann et al.

(56)

## References Cited

## U.S. PATENT DOCUMENTS

2009/0042979	A1	2/2009	Guzman et al.	2013/0017256	A1	1/2013	Manku et al.
2009/0054329	A1	2/2009	Willemssen et al.	2013/0065956	A1	3/2013	Yokoyama
2009/0105340	A1	4/2009	Yokoyama	2013/0079409	A1	3/2013	Manku et al.
2009/0148543	A1	6/2009	Theoharides	2013/0090383	A1	4/2013	Manku et al.
2009/0156675	A1	6/2009	Yokoyama et al.	2013/0095178	A1	4/2013	Manku
2009/0182049	A1	7/2009	Opheim	2013/0095179	A1	4/2013	Davidson et al.
2009/0227602	A1	9/2009	Griffin et al.	2013/0096197	A1	4/2013	Manku
2009/0233843	A1	9/2009	Marin	2013/0102674	A1	4/2013	Manku
2009/0239927	A1	9/2009	Bobotas et al.	2013/0115284	A1	5/2013	Fujii
2009/0304784	A1	12/2009	Mane et al.	2013/0131170	A1	5/2013	Manku
2009/0311322	A1	12/2009	Dlugatch et al.	2013/0156852	A1	6/2013	Manku et al.
2010/0021555	A1	1/2010	Geiringer et al.	2013/0158120	A1	6/2013	Manku et al.
2010/0063018	A1	3/2010	Pellicciari et al.	2013/0164375	A1	6/2013	Manku et al.
2010/0069492	A1	3/2010	Geiringer et al.	2013/0165513	A1	6/2013	Manku et al.
2010/0113506	A1	5/2010	Kawano et al.	2013/0171249	A1	7/2013	Manku et al.
2010/0113811	A1	5/2010	Yadav et al.	2013/0171250	A1	7/2013	Manku et al.
2010/0119598	A1	5/2010	Yoshinari et al.	2013/0171251	A1	7/2013	Manku et al.
2010/0130608	A1	5/2010	Ryan et al.	2013/0172413	A1	7/2013	Manku
2010/0160261	A1	6/2010	Fortin	2013/0189355	A1	7/2013	Manku et al.
2010/0233280	A1	9/2010	Driscoll	2013/0195972	A1	8/2013	Manku et al.
2010/0254951	A1	10/2010	Shido et al.	2013/0252989	A1	9/2013	Manku et al.
2010/0278879	A1	11/2010	Manku	2013/0252990	A1	9/2013	Manku et al.
2010/0285121	A1	11/2010	Uchiyama et al.	2013/0253030	A1	9/2013	Osterloh et al.
2010/0298379	A1	11/2010	Jacobsen	2013/0253031	A1	9/2013	Osterloh et al.
2010/0305205	A1	12/2010	Yokoyama et al.	2013/0260403	A1	10/2013	Button et al.
2010/0311834	A1	12/2010	Manku et al.	2013/0261180	A1	10/2013	Gillies et al.
2011/0034555	A1	2/2011	Osterloh et al.	2013/0281534	A1	10/2013	Osterloh et al.
2011/0065793	A1	3/2011	Peet et al.	2013/0295173	A1	11/2013	Machielse et al.
2011/0071176	A1	3/2011	Rowe	2013/0303614	A1	11/2013	Kanehiro et al.
2011/0082119	A1	4/2011	Yano	2013/0324607	A1	12/2013	Mason
2011/0092592	A1	4/2011	Yano	2013/0331447	A1	12/2013	Manku et al.
2011/0105510	A1	5/2011	Ishikawa	2014/0004183	A1	1/2014	Soni et al.
2011/0130458	A1	6/2011	Breivik et al.	2014/0005264	A1	1/2014	Soni et al.
2011/0178105	A1	7/2011	Gillies et al.	2014/0005265	A1	1/2014	Soni et al.
2011/0195061	A1	8/2011	Minatelli	2014/0017306	A1	1/2014	Manku
2011/0218243	A1	9/2011	Rowe	2014/0057981	A1	2/2014	Fujii
2011/0223158	A1	9/2011	Sacks et al.	2014/0073692	A1	3/2014	Peet
2011/0236476	A1	9/2011	Manku	2014/0080850	A1	3/2014	Mason
2011/0268811	A1	11/2011	Minatelli et al.	2014/0080909	A1	3/2014	Manku
2011/0288171	A1	11/2011	Manku et al.	2014/0088194	A1	3/2014	Manku
2012/0035105	A1	2/2012	Geho et al.	2014/0094520	A1	4/2014	Bobotas et al.
2012/0035262	A1	2/2012	Osterloh et al.	2014/0107199	A1	4/2014	Fawzy et al.
2012/0039997	A1	2/2012	Manku et al.	2014/0127289	A1	5/2014	Osterloh et al.
2012/0046251	A1	2/2012	Schaefer et al.	2014/0128453	A1	5/2014	Mullick et al.
2012/0093922	A1	4/2012	Manku et al.	2014/0128464	A1	5/2014	Rowe
2012/0093924	A1	4/2012	Manku et al.	2014/0154310	A1	6/2014	Osterloh et al.
2012/0100208	A1	4/2012	Manku	2014/0155455	A1	6/2014	Osterloh et al.
2012/0108659	A1	5/2012	Manku et al.	2014/0155481	A1	6/2014	Osterloh et al.
2012/0108660	A1	5/2012	Manku et al.	2014/0186438	A1	7/2014	Manku et al.
2012/0108663	A1	5/2012	Manku et al.	2014/0187633	A1	7/2014	Manku et al.
2012/0121698	A1	5/2012	Manku et al.	2014/0213648	A1	7/2014	Manku et al.
2012/0156285	A1	6/2012	Manku et al.	2014/0221358	A1	8/2014	Zakrzewski
2012/0157530	A1	6/2012	Manku et al.	2014/0221452	A1	8/2014	Zakrzewski
2012/0157531	A1	6/2012	Osterloh et al.	2014/0221486	A1	8/2014	Manku et al.
2012/0172432	A1	7/2012	Manku et al.	2014/0221676	A1	8/2014	Braeckman et al.
2012/0184595	A1	7/2012	Macdonald et al.	2014/0234410	A1	8/2014	Moodley et al.
2012/0195963	A1	8/2012	Peet et al.	2014/0235716	A1	8/2014	Manku et al.
2012/0207800	A1	8/2012	Abu-Baker	2014/0243389	A1	8/2014	Zakrzewski
2012/0214771	A1	8/2012	Sampalis	2014/0249200	A1	9/2014	Braeckman et al.
2012/0225120	A1	9/2012	Manku et al.	2014/0249214	A1	9/2014	Braeckman et al.
2012/0232145	A1	9/2012	Osterloh et al.	2014/0249220	A1	9/2014	Braeckman et al.
2012/0237594	A1	9/2012	Manku et al.	2014/0249225	A1	9/2014	Mason
2012/0264824	A1	10/2012	Mizuguchi et al.	2014/0256809	A1	9/2014	Zakrzewski
2012/0295976	A1	11/2012	Yokoyama	2014/0271841	A1	9/2014	Grandolfi
2012/0302589	A1	11/2012	Manku et al.	2014/0271907	A1	9/2014	Zakrzewski
2012/0329852	A1	12/2012	Yokoyama	2014/0275252	A1	9/2014	Zakrzewski
2013/0004566	A1	1/2013	Manku et al.	2014/0275253	A1	9/2014	Zakrzewski
2013/0004567	A1	1/2013	Manku et al.	2014/0322314	A1	10/2014	Fawzy et al.
2013/0004568	A1	1/2013	Manku et al.	2014/0357717	A1	12/2014	Braeckman et al.
2013/0004572	A1	1/2013	Manku et al.	2014/0364459	A1	12/2014	Zakrzewski
2013/0005757	A1	1/2013	Osterloh et al.	2015/0045431	A1	2/2015	Zakrzewski
2013/0005809	A1	1/2013	Manku et al.	2015/0051143	A1	2/2015	Harada et al.
2013/0011471	A1	1/2013	Manku et al.	2015/0051282	A1	2/2015	Zakrzewski
2013/0011472	A1	1/2013	Manku et al.	2015/0065572	A1	3/2015	Zakrzewski
2013/0012580	A1	1/2013	Osterloh et al.	2015/0073050	A1	3/2015	Zakrzewski
				2015/0141510	A1	5/2015	Kiyohara et al.
				2015/0157592	A1	6/2015	Soni
				2015/0157593	A1	6/2015	Braeckman et al.
				2015/0164850	A1	6/2015	Osterloh et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2015/0190361 A1 7/2015 Osterloh et al.  
 2015/0216831 A1 8/2015 Manku et al.  
 2015/0250754 A1 9/2015 Ohta  
 2015/0250756 A1 9/2015 Mason  
 2015/0250757 A1 9/2015 Soni  
 2015/0258051 A1 9/2015 Manku et al.  
 2015/0265566 A1 9/2015 Osterloh et al.  
 2015/0265574 A1 9/2015 Rowe  
 2015/0272917 A1 10/2015 Manku et al.  
 2015/0283074 A1 10/2015 Fujii  
 2015/0290154 A1 10/2015 Roberts et al.  
 2015/0335607 A1 11/2015 Rowe  
 2015/0359775 A1 12/2015 Osterloh et al.  
 2016/0058729 A1 3/2016 Manku et al.  
 2016/0120837 A1 5/2016 Manku et al.  
 2016/0143875 A1 5/2016 Zakrzewski  
 2016/0151319 A1 6/2016 Kimura  
 2016/0158184 A1 6/2016 Ito  
 2016/0213636 A1 7/2016 Manku et al.  
 2016/0213639 A1 7/2016 Suzuki et al.  
 2016/0220522 A1 8/2016 Osterloh et al.  
 2016/0287546 A1 10/2016 Osterloh et al.  
 2017/0014366 A1 1/2017 Osterloh et al.  
 2017/0035722 A1 2/2017 Soni  
 2017/0056361 A1 3/2017 Soni  
 2017/0079946 A1 3/2017 Ohta  
 2017/0087111 A1 3/2017 Mason  
 2017/0100363 A9 4/2017 Zakrzewski  
 2017/0119721 A1 5/2017 Zakrzewski  
 2017/0119722 A1 5/2017 Manku et al.  
 2017/0119723 A1 5/2017 Soni  
 2017/0119724 A1 5/2017 Fujii  
 2017/0128402 A1 5/2017 Manku et al.  
 2017/0128405 A1 5/2017 Osterloh et al.  
 2017/0128406 A1 5/2017 Rowe  
 2017/0136055 A1 5/2017 Zakrzewski  
 2017/0143656 A1 5/2017 Soni  
 2017/0143657 A1 5/2017 Braeckman et al.  
 2017/0143658 A1 5/2017 Soni  
 2017/0151202 A1 6/2017 Mason  
 2017/0151206 A1 6/2017 Yokoyama  
 2017/0258753 A1 9/2017 Soni  
 2017/0258754 A1 9/2017 Soni  
 2017/0258755 A1 9/2017 Soni  
 2017/0273928 A1 9/2017 Yokoyama  
 2017/0304249 A1 10/2017 Abu-Baker  
 2017/0333377 A1 11/2017 Mason  
 2017/0348268 A1 12/2017 Kimura  
 2017/0348273 A1 12/2017 Ito  
 2017/0368184 A1 12/2017 Ito  
 2018/0015038 A1 1/2018 Ito  
 2018/0015071 A1 1/2018 Braeckman et al.  
 2018/0028480 A1 2/2018 Mason  
 2018/0028505 A1 2/2018 Oshima  
 2018/0042880 A1 2/2018 Osterloh et al.  
 2018/0042883 A1 2/2018 Manku et al.  
 2018/0064676 A1 3/2018 Zakrzewski  
 2018/0085334 A1 3/2018 Soni  
 2018/0153846 A1 6/2018 Soni  
 2018/0185320 A1 7/2018 Manku et al.  
 2018/0280334 A1 10/2018 Manku  
 2018/0289657 A1 10/2018 Soni  
 2018/0289658 A1 10/2018 Soni  
 2018/0289659 A1 10/2018 Soni  
 2018/0333383 A1 11/2018 Philip  
 2019/0038590 A1 2/2019 Manku  
 2019/0054054 A1 2/2019 Mason  
 2019/0054058 A1 2/2019 Thero  
 2019/0060308 A1 2/2019 Mason  
 2019/0070141 A1 3/2019 Osterloh  
 2019/0076388 A1 3/2019 Soni  
 2019/0076389 A1 3/2019 Soni  
 2019/0076390 A1 3/2019 Manku  
 2019/0083444 A1 3/2019 Manku  
 2019/0083445 A1 3/2019 Soni

2019/0099422 A1 4/2019 Grandolfi  
 2019/0175535 A1 6/2019 Mason  
 2019/0175537 A1 6/2019 Osterloh  
 2019/0175538 A1 6/2019 Osterloh  
 2019/0183829 A1 6/2019 Osterloh  
 2019/0183831 A1 6/2019 Osterloh  
 2019/0183840 A1 6/2019 Braeckman  
 2019/0192472 A1 6/2019 Soni  
 2019/0201364 A1 7/2019 Manku  
 2019/0209506 A1 7/2019 Mason  
 2019/0240182 A1 8/2019 Osterloh  
 2019/0240183 A1 8/2019 Manku  
 2019/0269642 A1 9/2019 Philip  
 2019/0274991 A1 9/2019 Osterloh  
 2019/0275057 A1 9/2019 Philip  
 2019/0282533 A1 9/2019 Osterloh  
 2019/0316122 A1 10/2019 Zakrzewski  
 2019/0321323 A1 10/2019 Soni  
 2019/0343788 A1 11/2019 Soni  
 2019/0358185 A1 11/2019 Mason  
 2020/0000759 A1 1/2020 Manku  
 2020/0061011 A1 2/2020 Mason  
 2020/0061012 A1 2/2020 Manku et al.  
 2020/0069632 A1 3/2020 Soni  
 2020/0078329 A1 3/2020 Soni  
 2020/0093777 A1 3/2020 Soni  
 2020/0093778 A1 3/2020 Soni  
 2020/0093790 A1 3/2020 Rowe  
 2020/0108041 A1 4/2020 Braeckman et al.  
 2020/0113862 A1 4/2020 Manku et al.  
 2020/0113864 A1 4/2020 Soni  
 2020/0121628 A1 4/2020 Osterloh et al.  
 2020/0121630 A1 4/2020 Osterloh et al.  
 2020/0138768 A1 5/2020 Soni  
 2020/0163925 A1 5/2020 Soni  
 2020/0188343 A1 6/2020 Osterloh  
 2020/0188344 A1 6/2020 Osterloh  
 2020/0197350 A1 6/2020 Manku  
 2020/0215017 A1 7/2020 Manku  
 2020/0237699 A1 7/2020 Mason  
 2020/0237700 A1 7/2020 Mason  
 2020/0246300 A1 8/2020 Manku et al.  
 2020/0261391 A1 8/2020 Soni  
 2020/0268702 A1 8/2020 Braeckman et al.

FOREIGN PATENT DOCUMENTS

CA 2675836 7/2008  
 CA 2724983 11/2009  
 CA 2772378 12/2010  
 CN 101252837 8/2008  
 EP 273708 7/1988  
 EP 277747 8/1988  
 EP 0302482 2/1989  
 EP 347509 12/1989  
 EP 0460917 12/1991  
 EP 606012 7/1994  
 EP 0610506 8/1994  
 EP 0641562 A1 3/1995  
 EP 0843972 5/1998  
 EP 1125914 8/2001  
 EP 1157692 11/2001  
 EP 1296670 4/2003  
 EP 1549299 12/2003  
 EP 1743644 1/2007  
 EP 1 790 339 A1 5/2007  
 EP 1 834 639 A1 9/2007  
 EP 1946755 7/2008  
 EP 1 982 710 A1 10/2008  
 EP 2022495 2/2009  
 EP 2395991 8/2010  
 EP 2308493 A1 4/2011  
 EP 2343066 A1 7/2011  
 EP 2433630 3/2012  
 EP 2719382 A1 4/2014  
 EP 2792746 10/2014  
 FR 2635263 2/1990  
 GB 2148713 6/1985  
 GB 2221843 2/1990

(56)

## References Cited

## FOREIGN PATENT DOCUMENTS

GB	2229363	9/1990
GB	9901809.5	1/1999
GB	2480146	11/2011
IL	55227	12/1982
JP	61035356	2/1986
JP	04182426	6/1992
JP	H0692847	4/1994
JP	09059206	3/1997
JP	2001139981	5/2001
JP	2003306690	10/2003
JP	07 238598	9/2007
JP	08 050367	3/2008
KR	10-2006-0109988	10/2006
KR	10-2007-0058460	6/2007
RU	2290185	12/2006
RU	2302248 C2	7/2007
RU	2402326 C1	10/2010
WO	WO 1990/004391	5/1990
WO	WO 1992/021335	12/1992
WO	WO 1994/028891	12/1994
WO	WO 1995/024459	9/1995
WO	WO 1996/036329	11/1996
WO	WO 1997/039759	10/1997
WO	WO 1998/016216	4/1998
WO	WO 1999/26583	6/1999
WO	WO 1999/029316	6/1999
WO	WO 2000/044361	8/2000
WO	WO 2000/051573	9/2000
WO	WO 2001/015552	3/2001
WO	WO 2002/002105	1/2002
WO	WO 2002/058793	8/2002
WO	WO 2002/089787	11/2002
WO	WO 2002/096408	12/2002
WO	WO 2003/068216	8/2003
WO	WO 2003/092673	11/2003
WO	WO 2004/050913	6/2004
WO	WO 2004/064716	8/2004
WO	WO 2004/078166	9/2004
WO	WO 2004/082402	9/2004
WO	WO 2005/060954	7/2005
WO	WO2005/065652	7/2005
WO	WO 2005/079797	9/2005
WO	WO 2005/079853	9/2005
WO	WO 2005/102301	11/2005
WO	WO 2005/123060	12/2005
WO	WO 2005/123061	12/2005
WO	WO 2006/017627	2/2006
WO	WO 2006/029577	3/2006
WO	WO 2006/062748	6/2006
WO	WO 2006/096806	9/2006
WO	WO 2007/011886	1/2007
WO	WO 2007/016256	2/2007
WO	WO 2007/017240	2/2007
WO	WO 2007/073176	6/2007
WO	WO 2007/075841	7/2007
WO	WO 2007/091338	8/2007
WO	WO2007/103557	9/2007
WO	WO 2007/128801	11/2007
WO	WO 2007/142118	12/2007
WO	WO 2008/004900	1/2008
WO	WO 2008/045465	4/2008
WO	WO 2008/088415	7/2008
WO	WO 2008/106787	9/2008
WO	WO 2008/115529	9/2008
WO	WO 2008/145170	12/2008
WO	WO 2009/004999	1/2009
WO	WO 2009/085386	7/2009
WO	WO 2009/085388	7/2009
WO	WO 2010/028067	3/2010
WO	WO 2010/093634	8/2010
WO	WO 2010/127099	11/2010
WO	WO 2010/127103	11/2010
WO	WO2010/134614	11/2010
WO	WO 2010/147994	12/2010
WO	WO 2011/028689	3/2011

WO	WO 2011/038122	3/2011
WO	WO 2011/085211	7/2011
WO	WO 2011/109724	9/2011
WO	WO2012/032414	3/2012
WO	WO 2012/074930	6/2012
WO	WO 2012/128587	9/2012
WO	WO 2013/070735	5/2013
WO	WO 2013/103958	7/2013
WO	WO 2013/148136	10/2013
WO	WO2014/004861	1/2014
WO	WO2014/004993	1/2014
WO	WO2014/005013	1/2014
WO	WO 2014/057522	4/2014
WO	WO2014/074552	5/2014
WO	WO2014/130200	8/2014
WO	WO2014/134466	9/2014
WO	WO2014/142364	9/2014
WO	WO2014/143469	9/2014
WO	WO2014/143523	9/2014
WO	WO2015/021141	2/2015
WO	WO2015/066512	5/2015
WO	WO2015/195662	12/2015
WO	WO2016/140949	9/2016
WO	WO2018/213663	11/2018
WO	WO2020/037153	2/2020
WO	WO2020/065402	4/2020
WO	WO2020/068163	4/2020
WO	WO2020/168251	8/2020

## OTHER PUBLICATIONS

Abela GS, Aziz K. "Cholesterol crystals cause mechanical damage to biological membranes: a proposed mechanism of plaque rupture and erosion leading to arterial thrombosis." *Clin. Cardiol.* (Sep. 2005);28(9):413-420.

Abelo A, Andersson TB, Antonsson M, et al. "Stereoselective metabolism of omeprazole by human cytochrome P450 enzymes." *Drug Metab. Dispos.* Aug. 28, 2000 (8): 966-72.

Ackman et al., "The 'Basic' Fatty Acid Composition of Atlantic Fish Oils: Potential Similarities Useful for Enrichment of Polyunsaturated Fatty Acids by Urea Complexation," *JAOCS*, vol. 65, 1:136-138 (Jan. 1988).

Adan, Y, et al., "Effects of docosahexaenoic and eicosapentaenoic acid on lipid metabolism, eicosanoid production, platelet aggregation and atherosclerosis." *Biosci. Biotechnol. Biochem.* 63(1), 111-119 (Jan. 1999).

Adan, Y., et al., "Concentration of serum lipids and aortic lesion size in female and male apo E-deficient mice fed docosahexaenoic acid." *Biosci. Biotechnol. Biochem.* 63(2):309-313 (Feb. 1999).

Adorini et al., "Farnesoid X receptor targeting to treat nonalcoholic steatohepatitis," *Drug Discover Today*, 14(17-18):988-997 (Sep. 2012)(available online May 28, 2012).

Agren JJ, Vaisanen S, Hanninen O, et al. "Hemostatic factors and platelet aggregation after a fish-enriched diet or fish oil or docosahexaenoic acid supplementation." *Prostaglandins Leukot Essent Fatty Acids* (Oct. 1997) 57 (4-5): 419-21.

Agren, J.J. et al., "Fatty acid composition of erythrocyte, platelet, and serum lipids in strict vegans." *Lipids* 30:365-369 (Apr. 1995).

Agren, J.J., et al., "Fish diet, fish oil and docosahexaenoic acid rich oil lower fasting and postprandial plasma lipid levels." *Eur J Clin Nutr.*, 50:765-771. (Nov. 1996).

Aguilar-Salinas et al., "High Prevalence of Low HDL Cholesterol Concentrations and Mixed Hyperlipidemia in a Mexican Nationwide Survey," *J Lipid Res.*, (Aug. 2001), 42:1298-1307.

Ai M, Otokozawa S, Asztalos BF, Ito Y, Nakajima K, White CC, Cupples LA, Wilson PW, Schaefer EJ. "Small dense LDL cholesterol and coronary heart disease: results from the Framingham Offspring Study." *Clin. Chem.* (Jun. 2010);56(6):967-976.

Ait-Said, et al., "Inhibition by eicosapentaenoic acid of IL-1 $\beta$ -induced PGHS-2 expression in human microvascular endothelial cells: involvement of lipoxygenase-derived metabolites and p38 MAPK pathway." *Biochimica et Biophysica Acta*, 1631:66-85 (Feb. 2003).

(56)

## References Cited

## OTHER PUBLICATIONS

Albert CM, Campos H, Stampfer MJ, et al. Blood Levels of Long-Chain n-3 Fatty Acids and the Risk of Sudden Death. *N Engl J Med* 346(15):1113-1138, Apr. 2002.

Alberti K, et al. Harmonizing the Metabolic Syndrome: A Joint Interim Statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation*. 120:1640-1645; Oct. 20, 2009.

Alderman, J.D., et al., "Effect of a modified, well-tolerated niacin regimen on serum total cholesterol, high density lipoprotein cholesterol and the cholesterol to high density lipoprotein ratio," *Am. J. Cardio*, 64: 725-729.A (Oct. 1989).

Alessandri, J-M., et al., "Estradiol favors the formation of eicosapentaenoic acid (20:5n-3) and n-3 docosapentaenoic acid (22:5n-3) from alpha-linolenic acid (18:3n-3) in SH-SY5Y neuroblastoma cells." *Lipids* 43:19-28 (Jan. 2008).

Allard et al. "Nutritional assessment and hepatic fatty acid composition in non-alcoholic fatty liver disease (NAFLD): a cross-sectional study." *J Hepatol*. Feb. 2008;48(2):300-7.

Allred, C., et al., "PPAR $\gamma$ 1 as a molecular target of eicosapentaenoic acid in human colon cancer (HT-29) cells." *J. Nutr.* 138:250-256 (Feb. 2008).

Almeida et al., "Effect of nebicapone on the pharmacokinetics and pharmacodynamics of warfarin in healthy subjects." *Eur J Clin Pharmacol*. (Oct. 2008);64(10):961-6.

Amarin Appoints Medpace as CRO for Two Phase 3 Cardiovascular Trials, published Oct. 19, 2009 (2 pages).

Amarin Corporation Announces First Patients Enrolled in Two Phase 3 Clinical Trials Assessing AMR101 for the Treatment of Cardiovascular Disease [online], Amarin Corporation, Jan. 11, 2010 [retrieved Apr. 27, 2011], Retrieved from the Internet: <<http://investor.amarincorp.com/releasedetail.cfm?ReleaseID=504380>> (2 pages).

Amarin Corporation, Annual Report, Jun. 24, 2010 (245 pages total)(submitted in three parts; Part I: Cover and pp. 1-39 (81 pages); Part II: pp. 40 through F-10 (81 pages); Part III: pp. F11-F51 (83 pages)).

Amarin Corporation, Executive Informational Overview, "Neurological Disease-Focused Biopharmaceutical Opportunity," SEC filing dated Oct. 11, 2005 (99 pages).

Amarin Corporation, Globe Newsire press release, "Reduce-It™ Cardiovascular Outcomes Study of Vascepa® (icosapent ethyl) Capsules Met Primary Endpoint," Sep. 24, 2018 (4 pages).

Amarin Corporation, press release (Jan. 18, 2008)(1 page).

Amarin Presentation "Next Generation Lipid Modification in Cardiovascular Disease," (Aug. 2011)(27 pages).

Amarin Presentation "Next Generation Lipid Modification in Cardiovascular Disease," (Mar. 2010)(25 pages).

Amarin Proceeding to Phase 3 with AMR101 for Hypertriglyceridemia, published Jul. 23, 2008 (1 page).

Amarin, Next Generation Lipid Modification in Cardiovascular Disease, Investor Meetings, Nov. 2010, ([http://files.shareholder.com/downloads/AMRN/0x0x417754/AA72705F-1D67-4E1D-A989-5805E5CF0244/Investor\\_Presentation\\_2010\\_Nov\\_10.pdf](http://files.shareholder.com/downloads/AMRN/0x0x417754/AA72705F-1D67-4E1D-A989-5805E5CF0244/Investor_Presentation_2010_Nov_10.pdf)), accessed Jan. 6, 2015.

Amarin's Vascepa® Briefing Document for the Endocrinologic and Metabolic Drugs Advisory Committee Meeting dated Oct. 16, 2013, (117 pages).

American Heart Association. Heart Disease and Stroke Statistics—2010 Update. Dallas, Texas: American Heart Association; 2010.

Anand RG, Alkadri M, Lavie CJ, Milani RV. The Role of Fish Oil in Arrhythmia Prevention. *J Cardioplin Rehabil Preven.*, Mar./Apr. 2008; 28:92-98.

Anber V, Griffin BA, McConnell M, Packard CJ, Shepherd J. Influence of plasma lipid and LDL-subfraction profile on the interaction between low density lipoprotein with human arterial wall proteoglycans. *Atherosclerosis*. Aug. 1996;124(2):261-271.

Anderson JL, Adams CD, Antman EM, et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction—executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non-ST-Elevation Myocardial Infarction) developed in Collaboration with the American College of Emergency Physicians, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons Endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation and the Society for Academic Emergency Medicine. *J Am Coll Cardiol* 50:652-726, Aug. 14, 2007.

Anderson TJ, Gregoire J, Hegele RA, et al. 2012 update of the Canadian Cardiovascular Society guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease in the adult. *Can. J. Cardiol*. Feb. 2013;29:151-167.

Anderson TJ, Meredith IT, Yeung AC, Frei B, Selwyn AP, Ganz P. The effect of cholesterol-lowering and antioxidant therapy on endothelium-dependent coronary vasomotion. *N. Engl. J. Med*. Feb. 1995;332:488-493.

Anderson, "Lipoprotein-Associated Phospholipase A2: An Independent Predictor of Coronary Artery Disease Events in Primary and Secondary Prevention," 101 *Am. J. Cardiology* 23F-33F (Jun. 2008).

Ando, M., et al., "Eicosapentanoic acid reduces plasma levels of remnant lipoproteins and prevents in vivo peroxidation of LDL in dialysis patients." *J. Am. Soc. Nephrol.*, 10:2177-2184 (Oct. 1999).

Ando, Y., et al., "Positional distribution of highly unsaturated fatty acids in triacyl-sn-glycerols of *Artemia Nauplii* enriched with docosahexaenoic acid ethyl ester." *Lipids* 36:733-740 (Jul. 2001).

Andrade, SE. et al., "Discontinuation of antihyperlipidaemic drugs—do rates reported in clinical trials reflect rates in primary care settings?" *New Eng. J. Med*. 332: 1125-1131. (Apr. 1995).

Andrews HE, Bruckdorfer KR, Dunn RC, Jacobs M. Low-density lipoproteins inhibit endotheliumdependent relaxation in rabbit aorta. *Nature*. May 1987;327:237-239.

Angerer et al., "n-3 Polyunsaturated Fatty Acids and the Cardiovascular System", *Current Opinion in Lipidology*, 11(1):57-63, (Feb. 2000).

Anil, Eliz, "The Impact of EPA and DHA on Blood Lipids and Lipoprotein Metabolism: Influence of ApoE Genotype", *Proceedings of the Nutrition Society*, 66:60-68, (Feb. 2007).

Annex to Rule 161 Response dated Apr. 16, 2012 (4 pages).

Antman E, Anbe D, Armstrong P, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction—executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to revise the 1999 guidelines for the management of patients with acute myocardial infarction). *J Am Coll Cardiol* 44:671-719, Aug. 4, 2004.

Aoki T et al. "Experience of the use of ethyl eicosapentaenoic acid preparation (Epadel) in patients with arteriosclerosis obliterans complicated with diabetes mellitus. A study of the long-term effects on glycemic control and blood lipids," *Rinsho to Kenkyu*; 70:625-631. (1993) (with English translation).

Appendix A to Defendants' Invalidity Contentions, 3:14-CV-02550-MLC-DEA (D.N.J.), 478 pages (Dec. 5, 2014).

Appleton, Katherine M., et al., "Effects of n-3 long-chain polyunsaturated fatty acids on depressed mood: systematic review of published trials", *Am. J. Clin. Nutr.*, 84(6):1308-1316, (Dec. 2006).

Arca et al., "Treating statin-intolerant patients," *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 4:155-156 (Apr. 28, 2011).

Armaganijan L, Lopes RD, Healey JS, Piccini JP, Nair GN, Morillo CA. Do Omega-3 fatty acids prevent atrial fibrillation after open heart surgery? A meta-analysis of randomized controlled trials. *Clinics*. 2011(accepted for publication Jul. 19, 2011); 66(11):1923-1928.

Arrol, S. et al., "The effects of fatty acids on apolipoprotein B secretion by human hepatoma cells (HEP G2)," *Atherosclerosis* 150:255-264. (Jun. 2000).

(56)

## References Cited

## OTHER PUBLICATIONS

- Arshad, A. et al., "Sudden cardiac death and the role of medical therapy." *Progress in Cardiovascular Diseases*, vol. 50, No. 6, 420-438, (May/Jun. 2008).
- Arterburn, L., et al., "Distribution, interconversion, and dose response of n-3 fatty acids in humans." *Am J Clin Nutr.*, 83:1467S-76S (Jun. 2006).
- Asahara, EPA Products What is the Clinical Significance of EpaDel? *Obesity and Diabetes* 10(6):903-905 (2011) (with English translation).
- Asano, M., et al., "Eicosapentaenoic acid inhibits vasopressin-activated Ca<sup>2+</sup> influx and cell proliferation in rat aortic smooth muscle cell lines." *European Journal of Pharmacology* 379:199-209 (Aug. 1999).
- Asano, M., et al., "Inhibitory effects of  $\omega$ -3 polyunsaturated fatty acids on receptor-mediated non-selective cation currents in rat A7r5 vascular smooth muscle cells." *British Journal of Pharmacology* 120:1367-1375, (Apr. 1997).
- ASCEND Study Collaborative Group. Effects of n-3 fatty acid supplements in diabetes mellitus. *N Engl J Med*, 379(16):1540-1550 (publication date Oct. 18, 2018; epublication date Aug. 26, 2018).
- Ascenta Health "Fish Oil as Triglycerides vs. Ethyl Esters: Why this Matters." (2015)(14 pages).
- Astarita et al., "Targeted lipidomics strategies for oxygenated metabolites of polyunsaturated fatty acids," *Biochim Biophys Acta*, 1851(4):456-468 (Apr. 2015).
- Atorvastatin Package Leaflet, Reg. No. LSR-005205/08, Sep. 30, 2016 [retrieved Sep. 30, 2016] retrieved from the internet: [academ-clinic.ru/drugs/atorvastatin](http://academ-clinic.ru/drugs/atorvastatin) (6 pages).
- ATP III guidelines, NIH publication No. 01-3305 (2001).(6 pages).
- Attie AD, et al., "Relationship between stearoyl-CoA desaturase activity and plasma triglycerides in human and mouse hypertriglyceridemia," *J. Lipid Res.* 2002;43:1899-907.
- Ault, "Prescription omega-3 fatty acid formulation approved," *ob.gyn.news*, (Jan. 15, 2005).
- Aung T, Halsey J, Kromhout D, et al. Associations of omega-3 fatty acid supplement use with cardiovascular disease risks: Meta-analysis of 10 trials involving 77917 individuals. *JAMA Cardiol* 3:225-34 (publication date Mar. 1, 2018; epublication date Jan. 31, 2018).
- Avandia [package insert]. Research Triangle Park, NC: GlaxoSmithKline; 2011.(45 pages).
- Avery et al., "Upper Gastrointestinal System," *Integrating Therapeutic and Complementary Nutrition*, Edited by Mary Marian, CRC Press (2006)(14 pages).
- Aviram M, Rosenblat M, Bisgaier CL, Newton RS. Atorvastatin and gemfibrozil metabolites, but not the parent drugs, are potent anti-oxidants against lipoprotein oxidation. *Atherosclerosis*. Jun. 1998; 138(2):271-280.
- Ayton, et al., "A pilot open case series of Ethyl-EPA supplementation in the treatment of anorexia nervosa," *Prostaglandins, Leukotrienes and Essential Fatty Acids* 71, pp. 205-209. (Oct. 2004).
- Ayton, et al., "Rapid improvement of severe anorexia nervosa during treatment with ethyl-eicosapentaenoate and micronutrients," *European Psychiatry* 19, pp. 317-319. (Aug. 2004).
- Baigent, C., et al., "Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins." *Lancet*; 366:1267-1278. (Oct. 2005).
- Baldwin RM, Ohlsson S, Pedersen RS, et al. Increased omeprazole metabolism in carriers of the CYP2C19\*7 allele; a pharmacokinetic study in healthy volunteers. *Br. J. Clin. Pharmacol.* May 2008 65 (5): 767-74.
- Baldwin SJ, Clarke SE, Chenery RJ. Characterization of the cytochrome P450 enzymes involved in the in vitro metabolism of rosiglitazone. *Br. J. Clin. Pharmacol.* Sep. 1999;48:424-432.
- Balfour et al., "Rosiglitazone," *Drugs*, 57(6):921-930 (Jun. 1999).
- Balk, E.M., et al., "Effects of omega-3 fatty acids on serum markers of cardiovascular disease risk: a systematic review. *Atherosclerosis*." 189:19-30. (Nov. 2006).
- Ballantyne CM, Bays HE, Kastelein JJ, et al. Efficacy and safety of eicosapentaenoic acid ethyl ester (AMR 101) therapy in statin-treated patients with persistent high triglycerides (from the ANCHOR study). *Am J Cardiol* Oct. 2012 110 (7): 984-92.
- Ballantyne et al., "Abstract 15071: AMR101 Lowers Triglycerides, Atherogenic Lipoprotein, Phospholipase A<sub>2</sub>, and High-sensitivity C-reactive Protein Levels in Patients with High Triglycerides and on Background Statin Therapy (the ANCHOR Study)," *Circulation*, Lippincott Williams and Wilkins, vol. 124, No. 21, Suppl., Nov. 22, 2011.
- Ballantyne et al., "Effects of icosapent ethyl on lipoprotein particle concentration and the fatty acid desaturation index in statin-treated patients with persistent high triglycerides (the ANCHOR study)." *Journ. Clin. Lipidology*, 2013, 7(3):270-271.
- Ballantyne et al., Influence of low-high density lipoprotein cholesterol and elevated triglyceride on coronary heart disease events and response to simvastatin therapy in 4S, *Circulation*, 104:3046-3051. (Dec. 2001).
- Bang HO, Dyerberg J. "Plasma lipids and Lipoproteins in Greenlandic west coast Eskimos" *Acta Med Scand*, 192:85-94. (Jul./Aug. 1972).
- Banga, A., et al., "Adiponectin translation is increased by the PPAR $\gamma$  agonists pioglitazone and  $\omega$ -3 fatty acids." *Am J Physiol Endocrinol Metab* 296:480-489 (Mar. 2009).
- Bangham et al., "Diffusion of univalent ions across the lamellae of swollen phospholipids." *J. Mol. Biol.* (Aug. 1965) 13(1):238-252
- Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM, "Fasting Compared With Nonfasting Triglycerides and Risk of Cardiovascular Events in Women," *JAMA*, 298:309-316 (Jul. 2007).
- Barter et al., "Effectiveness of Combined Statin Plus Omega-3 Fatty Acid Therapy for Mixed Dyslipidemia." *Am. J. Cardiol.* 102(8):1040-1045 (Oct. 15, 2008).
- Basu, A. et al., "Dietary Factors That Promote or Retard Inflammation." *Arterioscler. Thromb. Vasc. Biol.* 26:995-1001 (May 2006).
- Baynes JW. Role of oxidative stress in development of complications in diabetes. *Diabetes*. Apr. 1991;40(4):405-412.
- Bays HE et al. "Prescription omega 3 fatty acids and their lipid effects: physiologic mechanisms of action and clinical implications," *Expert Rev Cardiovasc Ther.*, 6:391-409. (Mar. 2008).
- Bays HE, Ballantyne CM, Braeckman RA, Stirlen WG, Soni PN. Icosapent ethyl, a pure ethyl ester of eicosapentaenoic acid: effects on circulating markers of inflammation from the MARINE and ANCHOR studies. *Am. J. Cardiovasc. Drugs*. Feb. 2013;13(1):37-46.
- Bays HE, Braeckman RA, Ballantyne CM, et al. Icosapent ethyl, a pure EPA omega-3 fatty acid: Effects on lipoprotein particle concentration and size in patients with very high triglyceride levels (the MARINE study). *J. Clin. Lipidol.* Nov./Dec. 2012;6:565-572.
- Bays HE, Safety considerations with omega-3 fatty acid therapy. *Am. J. Cardiol.* Mar. 2007 99 (6A): 35C-43C.
- Bays, H., Clinical Overview of Omacor: A Concentrated Formulation of Omega-3 Polyunsaturated Fatty Acids, *Am J Cardiol.*; 98[suppl]:71i-76i (Aug. 2006).
- Bays, H., "Rationale for Prescription Omega-3-Acid Ethyl Ester Therapy for Hypertriglyceridemia: A Primer for Clinicians," *Drugs of Today*, 44(3); 205-246. (Mar. 2008).
- Bays, H.E., Eicosapentaenoic Acid Ethyl Ester (AMR101) Therapy in Patients With Very High Triglyceride Levels (from the Multi-center, placebo-controlled, Randomized, double-blinded, 12-week study with an open-label Extension [MARINE] Trial) *Am J Cardiol*;108:682-690. (Sep. 2011).
- Bays, H.E., et al., "Long-term up to 24-month efficacy and safety of concomitant prescription omega-3-acid ethyl esters and simvastatin in hypertriglyceridemic patients." *Curr Med Res Opin.*; 26:907-915. (Apr. 2010).
- Beal, M.F., *Annals of Neurology*, vol. 38, No. 3, "Aging, Energy, and Oxidative Stress in Neurodegenerative Diseases", pp. 357-366, (Sep. 1995).
- Beaumont et al., Design of Ester Prodrugs to Enhance Oral Absorption of Poorly Permeable Compounds: Challenges to the Discovery Scientist, *Current Drug and Metabolism*. (Dec. 2003) 4:461-485.
- Becker LB, Aufderheide TP, Geocadin RG, Callaway CW, Lazar RM, Donnino MW, Nadkarni VM, Abella BS, Adrie C, Berg RA,

(56)

## References Cited

## OTHER PUBLICATIONS

Merchant RM, O'Connor RE, Meltzer DO, Holm MB, Longstreth WT, Halperin HR. AHA Consensus Statement: Primary Outcomes for Resuscitation Science Studies: A Consensus Statement From the American Heart Association. *Circulation* 2011; CIR. 0b013e3182340239 published online before print Oct. 3 2011, doi:10.1161/CIR.0b013e3182340239.

Belarbi et al., "A process for high yield and scaleable recovery of high purity eicosapentaenoic acid esters from microalgae and fish oil," *Enzyme and Microbial Technology* 26:516-529 (Apr. 2000).

Belger et al., "Assessment of prefrontal activation by infrequent visual targets and non-target noval stimuli in schizophrenia: a function MRI study," Presented at the 9th Biennial winter workshop on schizophrenia, Davos, Switzerland, Feb. 7-13, 1998, Abstract in *Schizophrenia Research*. vol. 29. No. 1/02, Jan. 1998.

Belikov, *Pharmaceutical Chemistry in Two Parts, 1/General Pharmaceutical Chemistry* 43-47 (1993) (with English translation)(9 pages).

Belmaker et al., "Addition of Omega-3 Fatty Acid to Maintenance Medication Treatment for Recurrent Unipolar Depressive Disorder," *Am. J. Psychiatry*, 159:477-479 (Mar. 2002).

Belmaker, et al., "Omega-3 Eicosapentaenoic Acid in Bipolar Depression: Report of a Small Open-Label Study," *J Clin Psychiatry*; 66:726-729. (Jun. 2005).

Bender NK, Kraynak MA, Chiquette E, et al. Effects of marine fish oils on the anticoagulation status of patients receiving chronic warfarin therapy. *J. Thromb. Thrombolysis* Jul. 5, 1998 (3): 257-61.

Bénistant, C., et al., "Docosapentaenoic acid (22:5, n-3): metabolism and effect on prostacyclin production in endothelial cells." *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 55(4):287-292, (Oct. 1996).

Benn et al., Improving Prediction of Ischemic Cardiovascular Disease in the General Population Using Apolipoprotein B: The Copenhagen City Heart Study, 27 *Arteriosclerosis, Thrombosis, & Arterial Biology* 661 (Mar. 2007).

Bennett et al., "Treatment of IgA nephropathy with eicosapentaenoic acid (EPA): a two-year prospective trial [Abstract Only]." *Clin. Nephrol.* 31(3):128-131 (Mar. 1989).

Berge, R.K., et al., "In contrast with docosahexaenoic acid, eicosapentaenoic acid and hypolipidaemic derivatives decrease hepatic synthesis and secretion of triacylglycerol by decreased diacylglycerol acyltransferase activity and stimulation of fatty acid oxidation." *Biochem J.*; 343(Pt 1):191-197. (Oct. 1999).

Berglund L, Brunzell JD, Goldberg AC, et al. Evaluation and treatment of hypertriglyceridemia: an endocrine society clinical practice guideline. *J. Clin. Endocrinol. Metab.* Sep. 2012 97 (9): 2969-89.

Berliner JA, Watson AD. A role for oxidized phospholipids in atherosclerosis. *N. Engl. J. Med.* Jul. 2005;353(1):9-11.

Bertelsen M, Anggard EE, Carrier MJ. Oxidative stress impairs insulin internalization in endothelial cells in vitro. *Diabetologia*. May 2001;44(5):605-613.

Betteridge, D.J., "Diabetic dyslipidaemia: past, present and future." *Practical Diabetes Int*, 21(2): 78-85. (Mar. 2004).

Bhatt DL, Eagle KA, Ohman EM, et al. Comparative determinants of 4-year cardiovascular event rates in stable outpatients at risk of or with atherothrombosis. *JAMA* 304(12):1350-7 (publication date Sep. 22, 2010; epublication date Aug. 30, 2010).

Bhatt DL, Fox KAA, Hacke W, et al; Charisma Investigators. Clopidogrel and aspirin versus aspirin alone for the prevention of atherothrombotic events. *N Engl J Med.* 354(16):1706-1717 (publication date Apr. 20, 2006; epublication date Mar. 12, 2006).

Bhatt DL, Hulot JS, Moliterno DJ, Harrington RA. Antiplatelet and anticoagulation therapy for acute coronary syndromes. *Circ Res* 114(12):1929-43 (publication date Jun. 6, 2014).

Bhatt DL, Steg PG, Brinton EA, et al. Rationale and design of REDUCE-IT: Reduction of Cardiovascular Events with Icosapent Ethyl-Intervention Trial. *Clin Cardiol* 40:138-48 (publication date Mar. 2017; epublication date Mar. 15, 2017).

Bhatt DL, Steg PG, Ohman EM, et al; REACH Registry Investigators. International prevalence, recognition and treatment of cardiovascular risk factors in outpatients with atherothrombosis. *JAMA*. 295(2):180-189 (publication date Jan. 11, 2006).

Bhatt et al., "Cardiovascular Risk Reduction with Icosapent Ethyl for Hypertiglyceridemia," *N. Eng. J. Med.*, Nov. 10, 2018 (epub ahead of print)(12 pages)(downloaded from [nejm.org](https://www.nejm.org) on Nov. 13, 2018 at <https://www.nejm.org/doi/full/10.1056/NEJMoa1812792>).

Bild et al., "Multi-Ethnic Study of Atherosclerosis: objectives and design," *Am J Epidemiol* 156(9):871-81 (Nov. 1, 2002).

Black et al., "Effect of intravenous eicosapentaenoic acid on cerebral blood flow, edema, and brain prostaglandins in ischemic gerbils", *Prostaglandins*, 28(4), pp. 545-546. cited by other . (Oct. 1984).

Blankenhorn D.H. et al., "Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts." *JAMA* 257: 3233-3240. (Jun. 1987).

Block, R. C., et al., "EPA and DHA in blood cell membranes from acute coronary syndrome patients and controls." *Atherosclerosis*, 197(2):821-828 (Apr. 2008).

Blumenthal, *Advanced Studies in Medicine*, 2:148-157 (2002).

Boden WE, Probstfield JL, Anderson T, Chaitman BR, Desvignes-Nickens P, Koprowicz K, IJ McBride R, Teo K, Weintraub W. and the Aim-High Investigators, "Niacin in patients with low hdl cholesterol levels receiving intensive statin therapy," *N. Engl. J. Med.* Dec. 2011;365:2255-2267.

Bonaa, KH et al., Docosahexaenoic and Eicosapentaenoic acids in plasma phospholipids are divergently associated with high density lipoprotein in humans, *Arterioscler. Thromb. Vasc. Biol.*;12:675-681 (Jun. 1992).

Bonnet et al., "Comparative Effects of 10-mg Versus 80-mg Atorvastatin on High-Sensitivity C-Reactive Protein in Patients with Stable Coronary Artery Disease: Results of the CAP(Comparative Atorvastatin Pleiotropic Effects) Study," *Clinical Therapeutics*. 30(12):2298-2313 (Dec. 2008).

Borchman D, Lamba OP, Salmassi S, Lou M, Yapped MC. The dual effect of oxidation on lipid bilayer structure. *Lipids*. Apr. 1992;27(4):261-265.

Bordin et al., "Effects of fish oil supplementation on apolipoprotein B100 production and lipoprotein metabolism in normolipidaemic males," *Eur. J. Clin. Nutr.* 52: 104-9 (Feb. 1998).

Borow et al., "Biologic plausibility, cellular effects, and molecular mechanisms of eicosapentaenoic acid (EPA) in atherosclerosis," *Atherosclerosis*, 242(1):357-66 (Sep. 2015).

Borthwick et al., "The effects of an omega-3 ethyl ester concentrate on blood lipid concentrations in patients with hyperlipidemia," *Clin. Drug Investig.* (1998) 15(5): 397-404.

Bossaller C, Habib GB, Yamamoto H, Williams C, Wells S, Henry PD. Impaired muscarinic endothelium-dependent relaxation and cyclic guanosine 5'-monophosphate formation in atherosclerotic human coronary artery and rabbit aorta. *J. Clin. Invest.* Jan. 1987;79:170-174.

Bousserouel, S., et al., "Different effects of n-6 and n-3 polyunsaturated fatty acids on the activation of rat smooth muscle cells by interleukin-1beta." *J. Lipid Res.* 44:601-611 (Mar. 2003).

Bousserouel, S., et al., "Modulation of cyclin D1 and early growth response factor-1 gene expression in interleukin-1 beta-treated rat smooth muscle cells by n-6 and n-3 polyunsaturated fatty acids." *Eur. J. Biochem.* 271:4462-4473 (Nov. 2004).

Brady, L., et al., Increased n-6 polyunsaturated fatty acids do not attenuate the effects of long-chain n-3 polyunsaturated fatty acids on insulin sensitivity or triacylglycerol reduction in Indian Asians. *Am J Clin Nutr* 79:983-91(Jun. 2004).

Braeckman et al., "Abstract 18549: Effects of AMR101, a Pure Eicosapentaenoic Omega-3 Fatty Acid, on the Fatty Acid Profile in Plasma and Red Blood Cells in Statin-Treated Patients with Persistent High Triglycerides—Results from the ANCHOR study," *Circulation* 126(21S):A15071 (Nov. 20, 2012)(2 pages).

Braeckman et al., "Effect of Concomitant Icosapent Ethyl (Eicosapentaenoic Acid Ethyl Ester) on Pharmacokinetics of Atorvastatin," *Clinical Drug Investigation*. (Jan. 2015) (3)45-51.

Braeckman RA, Manku MS, Bays HE, Stirtan WG, Soni PN. Icosapent ethyl, a pure EPA omega-3 fatty acid: effects on plasma



(56)

## References Cited

## OTHER PUBLICATIONS

and red blood cell fatty acids in patients with very high triglyceride levels (results from the MARINE study). *Prostaglandins Leukot Essent Fatty Acids*. Sep. 2013;89(4):195-201.

Braeckman RA, Stirtan WG, Soni PN. Pharmacokinetics of eicosapentaenoic acid in plasma and red blood cells after multiple oral dosing with AMR101 (ethyleicosapentaenoic acid) in healthy subjects [abstract]. Presented at: Congress of the International Society for the Study of Fatty Acids and Lipids, Vancouver, Canada, May 26-30, 2012.

Braeckman RA, Stirtan WG, Soni PN. Pharmacokinetics of eicosapentaenoic acid in plasma and red blood cells after multiple oral dosing with icosapent ethyl in healthy subjects. *Clin. Pharmacol. Drug Dev.* Mar. 2014 (epub Oct. 22, 2013); 3:101-108.

Braunersreuther V, Steffens S, Arnaud C, Pelli G, Burger F, Proudfoot A, Mach F. A novel rantes antagonist prevents progression of established atherosclerotic lesions in mice. *Arterioscler. Thromb. Vasc. Biol.* Jun. 2008;28:1090-1096.

Breslow, J., "n-3 Fatty acids and cardiovascular disease." *Am J Clin Nutr.*, 83:1477S-82S (Jun. 2006).

Brinton EA, Ballantyne CM, Bays HE, Kastelein JJ, Braeckman RA, Soni PN. Effects of AMR101 on lipid and inflammatory parameters in patients with diabetes mellitus-2 and residual elevated triglycerides (200-500 mg/dl) on statin therapy at LDL-C goal: the ANCHOR study.[abstract 629-P] *Diabetes*. 2012;61(suppl 1):A159-A160.

Brinton et al., "Effects of icosapent ethyl on lipid and inflammatory parameters in patients with diabetes mellitus-2, residual elevated triglycerides (200-500 mg/dL), and on statin therapy at LDL-C goal: the ANCHOR study," *Cardiovasc. Diabetol.* Jul. 9, 2013;12:100. doi: 10.1186/1475-2840-12-100.

Brossard, N., et al., "Retroconversion and metabolism of [13C]22:6n-3 in humans and rats after intake of a single dose of [13C]22:6n-3-3-triacylglycerols." *Am. J. Clin. Nutr.* 64:577-86 (Oct. 1996).

Brouwer, I.A., et al., "Effect of fish oil on ventricular tachyarrhythmia and death in patients with implantable cardioverter defibrillators." *JAMA*. 295(22):2613-2619 (Jun. 2006).

Brovkovych V, Dobrucki LW, Brovkovych S, Dobrucki I, Do Nascimento CA, Burewicz A, Malinski T. Nitric oxide release from normal and dysfunctional endothelium. *J. Physiol. Pharmacol.* Dec. 1999;50:575-586.

Brown et al., Simvastatin and Niacin, Antioxidant Vitamins, or the Combination for the Prevention of Coronary Disease, *N Engl J Med*, vol. 345, No. 22, 1583-1592 (Nov. 29, 2001).

Brown, A. J., et al., "Administration of n-3 Fatty Acids in the Diets of Rats or Directly to Hepatocyte Cultures Results in Different Effects on Hepatocellular ApoB Metabolism and Secretion." *Arterioscler. Thromb. Vasc. Biol.* 19:106-114 (Jan. 1999).

Brown, A. J., et al., "Persistent changes in the fatty acid composition of erythrocyte membranes after moderate intake of n-3 polyunsaturated fatty acids: study design and implications." *Am. J. Clin. Nutr.* 54:668-73 (Oct. 1991).

Brown, G., et al., "Regression of coronary artery-disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein," *B., N. Engl. J. Med.* 323: 1289-1298. (Nov. 1990).

Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature*. Dec. 2001; 414(6865):813-820.

Bryhn, M., et al., "The bioavailability and pharmacodynamics of different concentrations of omega-3 acid ethyl esters." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 75:19-24 (Jul. 2006).

Budavari, S., Editor, "The Merck Index", Merck & Co., Inc., p. 725 item 4511 and p. 279 and item 2417 (1989).

Budoff, "Triglycerides and Triglyceride-Rich Lipoproteins in the Causal Pathway of Cardiovascular Disease," *Am. J. Cardiol.*, 118(1):138-45 (Jul. 1, 2016).

Bunting et al. "Depression in Parkinson's Disease". *J Neurosci Nurs.*; 23(3):158-164. (Abstract Only) (Jun. 1991).

Burdge, G.C., et al., "Eicosapentaenoic and docosapentaenoic acids are the principal products of  $\alpha$ -linolenic acid metabolism in young men." *British Journal of Nutrition* 88:355-363 (Oct. 2002).

Burdge, G.C., et al., "Lack of effect of meal fatty acid composition on postprandial lipid, glucose and insulin responses in men and women aged 50-65 years consuming their habitual diets." *British Journal of Nutrition*, 96:489-500 (Sep. 2006).

Burdge, G.C., et al., "The effect of altering the 20:5n-3 and 22:6n-3 content of a meal on the postprandial incorporation of n-3 polyunsaturated fatty acids into plasma triacylglycerol and non-esterified fatty acids in humans." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 77:59-65 (Jul. 2007).

Burr ML, Sweetham PM, Fehily AM. Diet and reinfarction. *Eur Heart J* 15:1152-1153, Aug. 1994.

Burr, M. L., et al., "Effects of changes in fat, fish and fibre intakes on death and myocardial reinfarction: Diet and reinfarction trial." *The Lancet*, 2(8666):757-61 (Sep. 1989).

Buse JB, Ginsberg HN, Bakris GL, et al. Primary prevention of cardiovascular diseases in people with diabetes mellitus: a scientific statement from the American Heart Association and the American Diabetes Association. *Diabetes Care*. 2007;30:162-172.

Calabresi, L., et al., "Omacor in familial combined hyperlipidemia: effects on lipids and low density lipoprotein subclasses." *Atherosclerosis* 148:387-396 (Feb. 2000).

Calder PC. Omega-3 Fatty Acids and Inflammatory Processes. *Nutrients* 2(3):355-374, Mar. 2010 (epub Mar. 18, 2010).

Calder PC. The role of marine omega-3 (n-3) fatty acids in inflammatory processes, atherosclerosis and plaque stability. *Mol. Nutr. Food Res.* Jul. 2012;56(7):1073-1080.

Campos, H., et al., "Lowdensity lipoprotein size, pravastatin treatment, and coronary events." *JAMA*, 286:1468-1474 (Sep. 2001).

Canner P.L. et al., "Fifteen year mortality in Coronary Drug Project patients: long-term benefit with niacin," *J. Am. Coll. Cardiol.* 8. 1245-1255. (Dec. 1986).

Cannon CP, Blazing MA, Giugliano RP, et al; IMPROVE-IT Investigators. "Ezetimibe added to statin therapy after acute coronary syndromes." *N Engl J Med.* 372:2387-2397. (Jun. 18, 2015/epub Jun. 3, 2015)

Cannon CP, Braunwald E, McCabe CH, et al. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 350(15):1495-1504 (publication date Apr. 8, 2004; epublication date Mar. 8, 2004).

Cao H, Wang X, Huang H, Ying SZ, Guy W, Wang T, Huang CX. Omega-3 Fatty Acids in the Prevention of Atrial Fibrillation Recurrences after Cardioversion: A Meta-analysis of Randomized Controlled Trials. *Intern. Med.* 2012 (epub Sep. 15, 2012); 51:2503-2508.

Cao, et al., "Cloning, Expression, and Chromosomal Localization . . .", *Genomics*, 49:327-331, (Apr. 15, 1998).

Cao, J., et al., "Incorporation and Clearance of Omega-3 Fatty Acids in Erythrocyte Membranes and Plasma Phospholipids." *Clinical Chemistry* 52(12):2265-2272 (Dec. 2006).

Capuzzi, DM et al., "Efficacy and safety of an extended-release niacin (Niaspan): a long-term study." *Am. J. Cardiol.* 82: 74U-81U. (Dec. 17, 1998).

Carlson, L.A. & Rosenhamer G., "Reduction of mortality in the Stockholm Ischaemic Heart Disease Secondary Prevention Study by combined treatment with clofibrate and nicotinic acid." *Acta Med. Scand.* 223, 405-418 (1988).

Carlson, L.A., "Nicotinic acid: the broad spectrum lipid drug. A 50th Anniversary review", *J. Int. Med.*, 258:94-114, (Aug. 2005).

Carrero et al., "Intake of Fish Oil, Oleic Acid, Folic Acid, and Vitamins B-6 and E for 1 Year Decreases Plasma C-Reactive Protein and Reduces Coronary Heart Disease Risk Factors in Male Patients in a Cardiac Rehabilitation Program", pp. 384-390 (Feb. 2007).

Carrero, J.J. et al. "Efectos cardiovasculares de los acidos grasos omega-3 y alternativas para incrementar su ingesta," *Nutricion Hospitalaria.* (2005) (1) 63-69 [with English abstract].

Carroll, D. N., et al., "Evidence for the Cardioprotective Effects of Omega-3 Fatty Acids." *Ann Pharmacother.*, 36:1950-6 (Dec. 2002).

Carulli et al., "Chenodeoxycholic acid and ursodeoxycholic acid effects in endogenous hypertriglyceridemias. A controlled double-blind trial." *J. Clin. Pharmacol.*, 21(10):436-42 (Oct. 1981).

Caughey GE, Mantzioris E, Gibson RA, Cleland LG, James MJ. The effect on human tumor necrosis factor  $\alpha$  and interleukin  $1\beta$

(56)

## References Cited

## OTHER PUBLICATIONS

production of diets enriched in n-3 fatty acids from vegetable oil or fish oil. *Am J Clin Nutr.* Jan. 1996;63:116-122.

Cavender MA, Steg PG, Smith SC, et al; REACH Registry Investigators. Impact of diabetes mellitus on hospitalization for heart failure, cardiovascular events, and death: outcomes at 4 years from the reduction of atherothrombosis for continued health (REACH) registry. *Circulation.* 132(10):923-931 (publication date Sep. 8, 2015; epublication date Jul. 7, 2015).

Cawood AL, Ding R, Napper FL, et al. Eicosapentaenoic acid (EPA) from highly concentrated n-3 fatty acid ethyl esters is incorporated into advanced atherosclerotic plaques and higher plaque EPA is associated with decreased plaque inflammation and increased stability. *Atherosclerosis.* Sep. 2010 (epub May 20, 2010); 212:252-259.

Cazzola, R., et al., "Age- and dose-dependent effects of an eicosapentaenoic acid-rich oil on cardiovascular risk factors in healthy male subjects." *Atherosclerosis* 193:159-167 (Jul. 2007).

Ceci et al., "The effects of oral 5-hydroxytryptophan administration on feeding behavior in obese adult female subjects," *J Neural. Transm* (1989) 76(2):109-117.

Cefali, E.A., et al., "Aspirin reduces cutaneous flushing after administration of an optimised extended-release niacin formulation", *Int. J. Clin. Pharmacol. & Ther.*, 45(2):78-88, (Feb. 2007).

Center for Drug Evaluation and Research. Application No. 21-853, 21654s016, (Omacor). Statistical Review and Evaluation: Clinical Studies, Omacor (omega-3 acid ethyl ester) Capsules, 4 grams/day; 2007. Available at: [http://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2007/0218535000;%20021654s016\\_StatR.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/nda/2007/0218535000;%20021654s016_StatR.pdf). (Accessed Jan. 26, 2012) (156 pages).

Center for Drug Evaluation and Research. Approval Package for Application No. 202057Orig1s000. Review—Vascepa (formerly AMR101), 373 pages (Jul. 26, 2012)(in two parts).

Center for Drug Evaluation and Research. Approval Package for: 21-654 (Omacor/Lovaza). Statistical Review; 2004. Available at: [http://www.accessdata.fda.gov/drugsatfda\\_docs/nda/2004/21-654\\_Omacor\\_AdminCorres\\_PI.pdf](http://www.accessdata.fda.gov/drugsatfda_docs/nda/2004/21-654_Omacor_AdminCorres_PI.pdf). Accessed Jan. 26, 2012. (54 pages).

Ceriello A, Motz E. Is oxidative stress the pathogenic mechanism underlying insulin resistance, diabetes, and cardiovascular disease? The common soil hypothesis revisited. *Arterioscler. Thromb. Vasc. Biol.* (May 2004);24(5):816-823.

Chait A, Brazg RL, Tribble DL, Krauss RM. Susceptibility of small, dense, low-density lipoproteins to oxidative modification in subjects with the atherogenic lipoprotein phenotype, pattern B. *Am. J. Med.* (Apr. 1993);94(4):350-356.

Chan et al., "Effect of Atorvastatin and Fish Oil on Plasma High-Sensitivity C-Reactive Protein Concentrations in Individuals with Visceral Obesity", *Clin. Chem.*, vol. 48, pp. 877-883 (2002).

Chan et al., Factorial Study of the Effects of Atorvastatin and Fish Oil on Dyslipidaemia in Visceral Obesity, 32 *Euro. J. Clinical Investigation.* 32(6):429-36 (Jun. 2002).

Chan, D.C., et al., "Randomized controlled trial of the effect of n-3 fatty acid supplementation on the metabolism of apolipoprotein B-100 and chylomicron remnants in men with visceral obesity." *Am J Clin Nutr* 77:300-7 (2003).

Chang CL, Seo T, Du CB, Accili D, Deckelbaum RJ. n-3 Fatty Acids Decrease Arterial Low-Density Lipoprotein Cholesterol Delivery and Lipoprotein Lipase Levels in Insulin-Resistant Mice. *Arterioscler Thromb Vasc Biol.* Dec. 2010 (epub Oct. 7, 2010); 30(12):2510-2517.

Chapman, M.J., et al., "Cholesteryl ester transfer protein: at the heart of the action of lipid-modulating therapy with statins, fibrates, niacin, and cholesteryl ester transfer protein inhibitors." *Eur Heart J.*, 31(2):149-164 (Jan. 2010).

Chatterjee SN, Agarwal S. Liposomes as membrane model for study of lipid peroxidation. *Free Radic. Biol. Med.* 1988;4(1):51-72.

Chemical Book, Eicosapentaenoic acid ethyl ester, copyright 2010, printed Jun. 16, 2011 from [www.chemicalbook.com](http://www.chemicalbook.com). (2010).

Chen, H., et al., "Eicosapentaenoic acid inhibits hypoxia-reoxygenation-induced injury by attenuating upregulation of MMP-1 in adult rat myocytes." *Cardiovascular Research* 59:7-13 (Jul. 2003).

Chen, H., et al., "EPA and DHA attenuate ox-LDL-induced expression of adhesion molecules in human coronary artery endothelial cells via protein kinase B pathway." *Journal of Molecular and Cellular Cardiology* 35:769-775 (Jul. 2003).

Chen, I.S. et al., "In vitro clearance of chylomicron triglycerides containing (omega-3) eicosapentaenoate." *Atherosclerosis*, 65:193-198 (1987).

Cheng et al., "Antagonism of the prostaglandin D2 receptor 1 suppresses nicotinic acid-induced vasodilation in mice and humans," *PNAS* 103(17):6682-7 (Apr. 25, 2006).

Childs, M.T., et al., "Divergent lipoprotein Responses to Fish Oils With Various Ratios of Eicosapentaenoic Acid and Docosahexaenoic Acid", *American Society for Clinical Nutrition*, 52:632-9, (Oct. 1990).

Christensen, J. H., et al., "Effect of fish oil on heart rate variability in survivors of myocardial infarction: a double blind randomised controlled trial." *BMJ*, 312:677-678 (Mar. 16, 1996).

Christensen, M.S., et al., "Intestinal absorption and lymphatic transport of eicosapentaenoic (EPA), docosahexaenoic (DHA), and decanoic acids: dependence on intramolecular triacylglycerol structure." *Am J Clin Nutr* 61:56-61 (Jan. 1995).

Citizen Petition, Pronova BioPharma Norge AS, Docket No. FDA-2009-P-0398-0001 (Aug. 4, 2009), at ii (Appendix), available at [www.regulations.gov](http://www.regulations.gov).

Classification of Hyperlipidaemias and Hyperlipoproteinaemias, *Bulletin of the World Health Organization*, 43(6): 891-915 (1970).

Cleland, L.G., et al., "A Biomarker of n-3 compliance in patients taking fish oil for rheumatoid arthritis." *Lipids* 38:419-424 (Apr. 2003).

Clinical Trial NCT01047501, Effect of AMR101 (Ethyl Icosapentate) on Triglyceride (Tg) Levels in Patients on Statins With High Tg Levels (>200 and <500 mg/dL) (ANCHOR), *ClinicalTrials.gov* [database online], U.S. National Institute of Health, Jan. 2010 [retrieved Apr. 27, 2011], Retrieved from the Internet: <<http://clinicaltrials.gov/ct2/show/NCT01047501>> (3 pages).

Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in insulin signaling and diabetes. *American journal of physiology. Endocrinology and metabolism.* (Dec. 2003);285(6):E1151-1160.

Cohen, J.D., et al., "30-year trends in serum lipids among United States adults: results from the National Health and Nutrition Examination Surveys II, III, and 1999-2006." *Am J Cardiol.*, 106:969-975. (Dec. 15, 2010).

Cole et al., "Challenges and opportunities in the encapsulation of liquid and semi-solid formulations into capsules for oral administration," *Advanced Drug Delivery Reviews*, vol. 60, No. 6, pp. 747-756. (Mar. 17, 2007).

Colhoun, H. M., et al., "Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial." *Lancet* 364: 685-9 (Aug. 21-24, 2004).

Collins, N., et al., "Differences between Dietary Supplement and Prescription Drug Omega-3 Fatty Acid Formulations: A Legislative and Regulatory Perspective." *Journal of the American College of Nutrition*, 27 (6):659-666 (Dec. 2008).

Committee Roster for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages. (2013).

Conklin, S. M., et al., "Serum omega-3 fatty acids are associated with variation in mood, personality and behavior in hypercholesterolemic community volunteers." *Psychiatry Research* 152: 1-10 (Jul. 30, 2007).

Connor et al., "Seminars in thrombosis and hemostasis," 14:271-284. (1988).

Connor, W.E., "Importance of n-3 Fatty Acids in Health and Disease", *Am. J. Clin. Nutr.*, 71(1(S)):171S-175S, (Jan. 2000).

Conquer, J.A., et al., "Effect of supplementation with different doses of DHA on the levels of circulating DHA as non-esterified fatty acid in subjects of Asian Indian background. *J Lipid Res.*" 39:286-292. (Feb. 1998).

(56)

## References Cited

## OTHER PUBLICATIONS

- Conquer, J.A., et al., "Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects." *J Nutr.*, 126: 3032-3039. (Dec. 1996).
- Contacos et al. Effect of pravastatin and omega-3 fatty acids on plasma lipids and lipoproteins in patients with combined hyperlipidemia, pp. 1755-1762 (Dec. 1993).
- Coronary Artery Bypass Grafting, NIH, published online Feb. 23, 2012 (12 pages).
- Costanzo S, di Niro V, Castelnovo AD, et al. Prevention of postoperative atrial fibrillation in open heart surgery patients by preoperative supplementation of n-3 polyunsaturated fatty acids: An updated meta-analysis. *Periop Manga.* 2013; Apr. 12 epub.
- Coumadin [package insert], Princeton, NJ: Bristol-Myers Squibb; 2011. (10 pages).
- Cox PJ, Ryan DA, Hollis FJ, et al. Absorption, disposition, and metabolism of rosiglitazone, a potent thiazolidinedione insulin sensitizer, in humans. *Drug Metab. Dispos.* Jul. 2000;28:772-780.
- Creager MA, Gallagher SJ, Gierd XJ, Coleman SM, Dzau VJ, Cooke JP. L-arginine improves endothelium-dependent vasodilation in hypercholesterolemic humans. *J. Clin. Invest.* Oct. 1992;90:1248-1253.
- Crevel et al., "Allergenicity of Refined Vegetable Oils," *Food and Chemical Toxicology*, 38, pp. 385-393 (Apr. 2000).
- Criqui, M., "Triglycerides and Coronary Heart Disease Revisited (Again)," vol. 147 No. 6, pp. 425-427 (Sep. 18, 2007).
- Cromwell et al., "LDL particle number and risk of future cardiovascular disease in the Framingham Offspring Study—Implications for LDL Management," *Journal of Lipidology.* (Dec. 2007) 1, 583-592.
- Crowe, F. L., et al., "Serum phospholipid n-3 long-chain polyunsaturated fatty acids and physical and mental health in a population-based survey of New Zealand adolescents and adults." *Am J Clin Nutr* 86:1278-85 (Nov. 2007).
- Cruz et al., "The metabolic syndrome in children and adolescents," *Curr. Diab. Rep.*, vol. 4(1):53-62 (Feb. 2004).
- Culhane et al., "Rosuvastatin for the treatment of hypercholesterolemia," *Pharmacotherapy*, 25(7):990-1000 (Jul. 2005).
- Daggy, B., et al., Dietary fish oil decreases VLDL production rates. *Biochimica et Biophysica Acta* 920: 293-300 (Aug. 15, 1987).
- Dall et al., "Clinical utility of low-density lipoprotein particle measurement in management of cardiovascular disease: a case report," *Research Reports in Clin. Cardiol.*, vol. 2, pp. 57-62 (2011).
- Daniel et al., "The Effect of Elevated Triglycerides on the Onset and Progression of Coronary Artery Disease: A Retrospective Chart Review," *Cholesterol*, vol. 2015, Article ID 292935, 5 pages (epub Nov. 4, 2015).
- Das, U.N., Essential fatty acids as possible mediators of the actions of statins. *Prostaglandins, Leukotrienes and Essential Fatty Acids* 65(1):37-40, (Jul. 2001).
- Davidson MH, Ballantyne CM, Jacobson TA, et al. Clinical utility of inflammatory markers and advanced lipoprotein testing: advice from an expert panel of lipid specialists. *J. Clin. Lipidol.* Sep./Oct. 2011;5:338-367.
- Davidson MH, et al., Effects of prescription omega-3-acid ethyl esters on lipo protein particle concentrations, apolipoproteins AI and CIII, and lipoprotein-associated phospholipase A<sub>2</sub> mass in statin-treated subjects with hypertriglyceridemia, *J.Clin. Lipid.*, vol. 3(5), pp. 332-340 (Oct. 2009).
- Davidson MH, Rosenson RS, Maki KC, Nicholls SJ, Ballantyne CM, Mazzone T, Carlson DM, Williams LA, Kelly MT, Camp HS, Lele A, Stolzenbach JC. Effects of fenofibric acid on carotid intima-media thickness in patients with mixed dyslipidemia on atorvastatin therapy: Randomized, placebo-controlled study (first). *Arterioscler. Thromb. Vasc. Biol.* Jun. 2014;34:1298-1306.
- Davidson MH, Stein EA, Bays HE et al. "Efficacy and tolerability of adding prescription omega-3 fatty acids 4 g/d to simvastatin 40 mg/d in hypertriglyceridemic patients: an 8-week, randomized, double-blind, placebo-controlled study," *Clin Ther.*, 29:1354-1367. (Jul. 2007).
- Davidson MH., "Mechanisms for the hypotriglyceridemic effect of marine omega 3 fatty acids." *Am J Cardiol* 98(4A):27i-33i. (Aug. 21, 2006).
- Davidson, M.H., et al., "Effects of docosahexaenoic acid on serum lipoproteins in patients with combined hyperlipidemia: a randomized, doubleblind, placebo-controlled trial." *J Am Coll Nutr.*, 16:236-243. (Jun. 1997).
- Davies et al., "Rapid separation of Ldl subclasses by iodixanol gradient ultracentrifugation," *Clin. Chem.*, 49(11):1865-72. (Nov. 2003).
- Davies-Tuck et al., "Total cholesterol and triglycerides are associated with development of new bone marrow lesions in asymptomatic middle-aged women—a prospective cohort study," *Arthritis Research & Therapy.* (published online Dec. 4, 2009) pp. 1-7.
- De Caterina, R, et al., "Control of Endothelial Leukocyte Adhesion Molecules by Fatty Acids." *Lipids*, vol. 31:S57-S63 (1996).
- De Caterina, R., et al., "The Omega-3 fatty acid docosahexaenoate reduces cytokine-induced expression of proatherogenic and proinflammatory proteins in human endothelial cells." *Arterioscler. Thromb. Vasc. Biol.* 14:1829-1836 (1994).
- De Graaf J, Hak-Lemmers HL, Hectors MP, Demacker PN, Hendriks JC, Stalenhoef AF. Enhanced V susceptibility to in vitro oxidation of the dense low density lipoprotein subfraction in healthy subjects. *Arterioscler. Thromb.* 1991;11(2):298-306.
- De Moraes et al., "Evaluation of lipid extraction and fatty acid composition of human plasma," *Rev. Bras. Hematol. Hemoter.* 32(6):439-443 (2010).
- Deckelbaum R. J., et al., "Conclusions and recommendations from the symposium, Beyond Cholesterol: Prevention and Treatment of Coronary Heart Disease with n-3 Fatty Acids." *Am J Clin Nutr* 87:2010S-12S (2008).
- Defendants' Invalidity Contentions, 3:14-CV-02550-MLC-DEA (D.N. J.), 520 pages (Dec. 5, 2014).
- Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 901 pages (Dec. 5, 2014).
- DeMets DL, Lan KK. Interim Analysis: the Alpha Spending Function Approach. *Stat Med.*, Jul. 15-30, 1994; 13(13-14):1341-52.
- Dewailly, E., et al., "n-3 Fatty acids and cardiovascular disease risk factors among the Inuit of Nunavik." *Am J Clin Nutr* 74:464-73 (2001).
- Dewey FE, Gusarova V, O'Dushlaine C, et al. Supplement to: Inactivating variants in ANGPTL4 and risk of coronary artery disease. *N Engl J Med.* DOI: 10.1056/NEJMoa1510926; Mar. 24, 2016 (epub Mar. 2, 2016).
- Di Spirito, M., Morelli, G., Doyle, R.T., Johnson, J. & McKenney, J. Effect of omega-3-acid ethyl esters on steady-state plasma pharmacokinetics of atorvastatin in healthy adults. *Expert Opin. Pharmacother.* 9, 2939-2945 (2008).
- Diagnostic and Statistical Manual of Mental Disorders, 4.Ed. Text revision, published by the American Psychiatric Assoc., pp. 154-163 and 369-381 (2000).
- Diagnostic and Statistical Manual of Mental Disorders, 4.sup.th Ed., published by the American Psychiatric Assoc., pp. 285-286, (1994).
- Dijan, P., et al., *Proc. Natl. Acad. Sci.*, vol. 93, "Codon repeats in genes associated with human diseases: Fewer repeats in the genes of nonhuman primates and nucleotide substitutions concentrated at the sites of reiteration," pp. 417-421, (Jan. 9, 1996).
- Dijk, J. M., et al., "Carotid intima-media thickness and the risk of new vascular events in patients with manifest atherosclerotic disease: the SMART study." *European Heart Journal* 27:1971-1978 (2006).
- Din et al., "Omega 3 fatty acids and cardiovascular disease—fishing for a natural treatment," *BMJ*, vol. 327, No. 7430, pp. 30-35 (Jan. 3, 2004).
- Djousse L, Akinkuolie AO, Wu JHY, Ding EL, Gaziano JM. Fish consumption, omega-3 fatty acids and risk of heart failure: a meta-analysis. *Clin Nutr.* Dec. 2012 (epub Jun. 6, 2012); 31:846-853.

(56)

## References Cited

## OTHER PUBLICATIONS

- Do R, Stitzel NO, Won HH, et al. Exome sequencing identifies rare LDLR and APOA5 alleles conferring risk for myocardial infarction. *Nature*. Feb. 5, 2015 (epub Dec. 10, 2014); 518(7537):102-106.
- Do R, Willer CJ, Schmidt EM, et al. Common variants associated with plasma triglycerides and risk for coronary artery disease. *Nat Genet* Nov. 2013 (Oct. 6, 2013); 45(11):1345-52.
- Dodin, S., et al., "Flaxseed on cardiovascular disease markers in healthy menopausal women: a randomized, double-blind, placebo-controlled trial." *Nutrition* 24:23-30 (2008).
- Doi M, Nosaka K, Miyoshi T, et al. Early eicosapentaenoic acid treatment after percutaneous coronary intervention reduced acute inflammatory responses and ventricular arrhythmias in patients with acute myocardial infarction: A randomized controlled study. *Int J Cardiol.*, 176(3):577-82 (publication date Oct. 20, 2014; epublication date Aug. 19, 2014).
- Dolecek, "Epidemiological Evidence of Relationships Between Dietary Polyunsaturated Fatty Acids and Mortality in the Multiple Risk Factor Intervention Trial", *Society of Experimental Biology and Medicine*, 200(2):177-182, (1991).
- Draft Agenda for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Draft Meeting Roster for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Draft Questions for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 1 page.
- Drexler H, Zeiher AM, Meinzer K, Just H. Correction of endothelial dysfunction in coronary microcirculation of hypercholesterolaemic patients by L-arginine. *Lancet*. 1991;338:1546-1550.
- Dullenmeijer, C., et al., "n-3 Fatty acid proportions in plasma and cognitive performance in older adults." *Am J Clin Nutr* 86:1479-85 (2007).
- Duncan, R. E., et al., "Regulation of HMG-CoA reductase in MCF-7 cells by genistein, EPA, and DHA, alone and in combination with mevastatin." *Cancer Letters* 224:221-228 (2005).
- Durrington PN et al. "An omega 3 poly unsaturated fatty acid concentrate administered for one year decreased triglycerides in simvastatin treated patients with coronary heart disease and persistent Hypertriglyceridemia," *Heart*, 85:544-48 (2001).
- Dwyer, J. H., et al., "Arachidonate 5-Lipoxygenase Promoter Genotype, Dietary Arachidonic Acid, and Atherosclerosis." *N. Engl. J. Med.*, 350:1 (2004).
- Dyerberg, J., et al., "Marine Oils and Thrombogenesis." *Prog. Lipid Res.* 21:255-269 (1982).
- Egert, S., et al., "Dietary alpha-linolenic acid, EPA, and DHA have differential effects on LDL fatty acid composition but similar effects on serum lipid profiles in normolipidemic humans." *J Nutr.*, 139:861-868 (2009).
- Ehara S, Ueda M, Naruko T, Haze K, Itoh A, Otsuka M, Komatsu R, Matsuo T, Itabe H, Takano T, Tsukamoto Y, Yoshiyama M, Takeuchi K, Yoshikawa J, Becker AE. Elevated levels of oxidized low density lipoprotein show a positive relationship with the severity of acute coronary syndromes. *Circulation*. 2001;103(15):1955-1960.
- Eilat-Adar et al. "Association of Intentional Changes in Body Weight with Coronary Heart Disease Event Rates in Overweight Subjects who have an Additional Coronary Risk Factor," *Amer. Journ. Epidemiol.* 161(4)pp. 352-358 (Sep. 9, 2004).
- Eisenberg S, Bilheimer DW, Levy RI, Lindgren FT. "On the metabolic conversion of human plasma very low density lipoprotein to low density lipoprotein," *Biochim Biophys Acta*, 326:361-77 (1973).
- Eisenberg S, Rachmilewitz D. "Metabolism of rat plasma very low density lipoprotein. I. Fate in circulation of the whole lipoprotein," *Biochim Biophys Acta*, 326:378-90 (1973).
- El-Serag HB, Graham DY, Satia JA, et al. Obesity is an independent risk factor for Gerd symptoms and erosive esophagitis. *Am. J. Gastroenterol.* Jun. 2005 100 (6):1243-50.
- Elam, M.B., et al., "Effect of niacin on lipid and lipoprotein levels and glycemic control in patients with diabetes and peripheral arterial disease study: a randomized trial", *The ADMIT [Arterial Disease Multiple Intervention Trial] JAMA*, 284:1263-1270, (2000).
- El-Saadani M, Esterbauer H, El-Sayed M, Gober M, Nassar AY, Jurgens G. A spectrophotometric assay for lipid peroxides in serum lipoproteins using commercially available reagent. *J. Lipid Res.* 1989;30:627-630.
- El-Soheby, A., et al., "Regulation of Mevalonate Synthesis in Low Density Lipoprotein Receptor Knockout Mice Fed n-3 or n-6 Polyunsaturated Fatty Acids." *Lipids*, 34 (10):1037-43 (1999).
- Emsley et al., "Randomized, Placebo-Controlled Study of Ethyl-Eicosapentaenoic Acid as Supplemental Treatment in Schizophrenia," *Am. J. Psychiatry*, 159:1596-1598 (2002).
- Endo et al., "The Effects of Dietary Fatty Acids on Serum Lipids and Plasma Prostaglandin Levels in the Treatment of Obesity," *Japanese Journal of Pediatric Gastroenterology and Nutrition* 7(1):67-72 (Apr. 15, 1993) (with English translation)(22 pages).
- ENews, "Cholesterol Crystals Induce Atherosclerosis-Associated Inflammation in Mice," 1-4 (Jun. 14, 2010)(4 pages).
- Engler, et al., "Docosahexaenoic acid restores endothelial function in children with hyperlipidemia: results from the EARLY Study." *International Journal of Clinical Pharmacology and Therapeutics*, vol. 42—No. 12-2004 (672-679). (2004).
- Engler, M.B., et al., "Mechanisms of vasorelaxation induced by eicosapentaenoic acid (20:5n-3) in WKY rat aorta." *British Journal of Pharmacology* 131:1793-1799 (2000).
- Engler, M.M., et al., "The effects of a diet rich in docosahexaenoic acid on organ and vascular fatty acid composition in spontaneously hypertensive rats." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 61(5):289-295 (1999).
- Ennis JL, Cromwell WC. Clinical utility of low-density lipoprotein particles and apolipoprotein B patients with cardiovascular risk. *J. Fam. Pract.* 2013;62:1-8.
- Epadel—PubChem CID 9831415, Retrieved on Apr. 9, 2014 [Retrieved from the Internet] <URL:<http://pubchem.ncbi.nlm.nih.gov/compound/9831415>> (19 pages).
- Epadel 1990 and JELIS Study (4 pages).
- Epadel Capsules 300, *Japan Pharmaceutical Reference* 369-371 (2nd ed.) (1991). (5 pages).
- Epadel drug information brochure (2000), certified English translation(36 pages).
- Epadel Package Insert 2007 (with Translation)(6 pages).
- Epadel Summary of Product Characteristics (SPC), Mochida Pharmaceutical Co., Ltd. Tokyo, Japan, Oct. 2013.
- Epadel® [Complete prescribing information]. Update (Version 5). Tokyo, Japan: Mochida Pharmaceutical; Jan. 2007 (9 pages).
- EPANOVA® (omega-3-carboxylic acids) capsules, for oral use, Prescribing information, 5 pgs., AstraZeneca Pharmaceuticals LP, (Revised: Mar. 2017)(5 pages).
- Eritsland J, Arnesen H, Gronseth K, et al. Effect of dietary supplementation with n-3 fatty acids on coronary artery bypass graft patency. *Am. J. Cardiol.* Jan. 1996 77 (1):31-6.
- Eritsland J, Arnesen H, Seljeflot I, et al. Long-term effects of n-3 polyunsaturated fatty acids on haemostatic variables and bleeding episodes in patients with coronary artery disease. *Blood Coagul. Fibrinolysis* Feb. 6, 1995 (1): 17-22.
- Errata to the FDA Briefing Document Endocrinologic and Metabolic Drug Advisory Committee Meeting Oct. 16, 2013, 1 page.
- Esposito, "Effect of a Mediterranean-Style Diet on Endothelial Dysfunction and Markers of Vascular Inflammation in the Metabolic Syndrome: A Randomized Trial", *Journal of the American Medical Association*, 2004, 292(12), 1440-1446.
- Essentialis Inc. press release, "Essentialis Meets Primary Endpoint in Phase 2b Trial of DCCR for Treatment of Hypertriglyceridemia and is Granted Extensive Patent Coverage in the US," PR Newswire (May 17, 2009)( 2 pages).
- Exhibit A to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 48 pages (Dec. 5, 2014).
- Exhibit B to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 6 pages (Dec. 5, 2014).
- Exhibit C to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 14 pages (Dec. 5, 2014).
- Exhibit D to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 19 pages (Dec. 5, 2014).

(56)

## References Cited

## OTHER PUBLICATIONS

- Exhibit E to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 6 pages (Dec. 5, 2014).
- Exhibit F to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 10 pages (Dec. 5, 2014).
- Exhibit G to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 21 pages (Dec. 5, 2014).
- Exhibit H to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 10 pages (Dec. 5, 2014).
- Exhibit I to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 14 pages (Dec. 5, 2014).
- Exhibit J to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 13 pages (Dec. 5, 2014).
- Exhibit K to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 5 pages (Dec. 5, 2014).
- Exhibit L to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 5 pages (Dec. 5, 2014).
- Exhibit M to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 7 pages (Dec. 5, 2014).
- Exhibit N to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 15 pages (Dec. 5, 2014).
- Exhibit O to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 6 pages (Dec. 5, 2014).
- Exhibit P To Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 17 pages (Dec. 5, 2014).
- Exhibit Q to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 64 pages (Dec. 5, 2014).
- Faggini, E., et al., "Fish Oil Supplementation Prevents Neointima Formation in Nonhypercholesterolemic Balloon-Injured Rabbit Carotid Artery by Reducing Medial and Adventitial Cell Activation." *Arterioscler. Thromb. Vasc. Biol.*, 20:152-163 (2000).
- FDA Briefing Document, Endocrinologic and Metabolic Drugs Advisory Committee Meeting, dated Oct. 16, 2013, available publicly at least as of Oct. 16, 2013, 115 pages.
- FDA News Release, "FDA approves new orphan drug Kynamro to treat inherited cholesterol disorder," U.S. Food and Drug Administration, Protecting and Promoting Your Health (Jan. 29, 2013)(2 pages).
- Fer, M., et al., "Metabolism of eicosapentaenoic and docosahexaenoic acids by recombinant human cytochromes P450." *Archives of Biochemistry and Biophysics* 471:116-125 (2008).
- Ferns, G., et al., "Investigation and management of hypertriglyceridaemia." *J. Clin. Pathol.* 61:1174-1183 (2008).
- Feron O, Dessy C, Desager JP, Balligand JL. Hydroxy-methylglutaryl-coenzyme a reductase inhibition promotes endothelial nitric oxide synthase activation through a decrease in caveolin abundance. *Circulation.* 2001;103:113-118.
- Final Agenda for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Final Meeting Roster for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Final Questions for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 1 page.
- Finnen et al., "Purification and characterisation of phospholipase A2 from human epidermis," *Biochemical Society Trans*,19(2):91S, 1991.
- Fischer, R., et al., "Dietary n-3 polyunsaturated fatty acids and direct renin inhibition improve electrical remodeling in a model of high human renin hypertension." *Hypertension* 51:540-546 (2008).
- Fisher et al., *Journal of Biological Chemistry* (2001) 276(3) 27855-27863.
- Flaten, H., et al., "Fish-oil concentrate: effects on variables related to cardiovascular disease." *Am. J. Clin. Nutr.* 52:300-306 (1990).
- Food and Drug Administration (FDA), (2005) *NIASPAN niacin extended release tablets*.
- Food and Drug Administration (FDA), (2005) *Tablets ZOCOR® (SIMVASTATIN)*.
- Ford, E.S. et al., "Hypertriglyceridemia and Its Pharmacologic Treatment Among US Adults." *Arch. Intern. Med.*, 169(6): 572-78 (2009).
- Fraker TD, Fihn SD. Writing on behalf of the 2002 Chronic Stable Angina Writing Committee. 2007 chronic angina focused update of the ACC/AHA guidelines for the management of patients with chronic stable angina. A Report of the ACC/AHA Task Force on Practice Guidelines. *Circulation* 50:2264-2274, Dec. 4, 2007.
- Frangou et al., "Efficacy of ethyl-eicosapentaenoic acid in bipolar depression: randomised double-blind placebo-controlled study," *British Journ. Psychiatry*, 188, 46-50 (2006).
- Frey R, Muck W, Kirschbaum N, et al. Riociguat (BAY 63-2521) and warfarin: a pharmacodynamic and pharmacokinetic interaction study. *J. Clin. Pharmacol.* Jul. 2011 51 (7): 1051-60.
- Frick, MH, et al., "Helsinki Heart Study. Primary prevention trial with gemfibrozil in middle-aged men with dyslipidaemia. Safety of treatment, changes in risk factors and incidence of coronary heart disease", *N. Eng. J. Med.*, 317:1237-1245, (1987).
- Friedewald, W.T., et al., "Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge." *Clin Chem.*,18:499-502 (1972).
- Friedman, A. N., et al., "Fish Consumption and Omega-3 Fatty Acid Status and Determinants in Long-Term Hemodialysis." *Amer. J. Kidney Diseases*, 47(6)1064-1071 (2006).
- Frøyland et al., "Chronic administration of eicosapentaenoic acid and docosahexaenoic acid as ethyl esters reduced plasma cholesterol and changed the fatty acid composition in rat blood and organs." *Lipids* 31(2):169-78 (Feb. 1996).
- Frøyland, L., et al., "Hypotriacylglycerolemic component of fish oil." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 57 (4 & 5):387-388 (1997).
- Furuta T, Shirai N, Sugimoto M, et al. Influence of CYP2C19 pharmacogenetic polymorphism on proton pump inhibitor-based therapies. *Drug Metab. Pharmacokinet* Jun. 20, 2005 (3): 153-67.
- Futata et al., "Effect of Eicosapentaenoic Acid (EPA) Formulation on Glucose Metabolism in Non-Insulin Dependent Diabetic Patients," *Journal of Clinical and Experimental Medicine* 169(8):889-890 (May 21, 1994)(English translation, 4 pages).
- Galan P, Kesse-Guyot E, Czernichow S, et al. Effects of B vitamins and omega 3 fatty acids on cardiovascular diseases: a randomised placebo controlled trial. *Br Med J.* Nov. 29, 2010;341:c6273.
- Galeano NF, Al-Haideri M, Keyserman F, Rumsey SC, Deckelbaum RJ. Small dense low density lipoprotein has increased affinity for LDL receptor-independent cell surface binding sites: a potential mechanism for increased atherogenicity. *J. Lipid Res.* 1998;39(6):1263-1273.
- Gallagher et al., "Germline BRCA Mutations Denote a Clinicopathologic Subset of Prostate Cancer," *Amer. Assoc. Cancer Res. Clin Cancer Res.*, 16(7):2115-21 (Apr. 1, 2010).
- Ganda OP, Bhatt DL, Mason RP, Miller M, Boden WE. Unmet need for adjunctive dyslipidemia therapy in hypertriglyceridemia management. *J Am Coll Cardiol* 72(3):330-43 (publication date Jul. 17, 2018).
- Garber AJ, Abrahamson MJ, Barzilay JI, et al. American Association of Clinical Endocrinologists' comprehensive diabetes management algorithm 2013 consensus statement. *Endocr. Pract.* 2013;19(suppl 2):1-48.
- Gardner CD, Fortmann SP, Krauss RM. Association of small low-density lipoprotein particles with the incidence of coronary artery disease in men and women. *JAMA.* 1996;276(11):875-881.
- Garg, R., et al., "Niacin treatment increases plasma homocyst(e)ine levels", *Am. Heart. J.*, 138:1082-1087, (1999).
- Garnett, "Interactions with Hydroxymethylglutaryl-coenzyme a reductase inhibitors," *Am J Health-Sys Pharm* vol. 52, 1639-1645, (Aug. 1, 1995).
- Geleijnse JM, Giltay EJ, Grobbee DE, Donders ART, Kok FJ. Blood pressure response to fish oil supplementation: metaregression analysis of randomized trials. *J Hypertens.* Aug. 2002;20(8):1493-1499.
- Genest, JJ, et al., "Familial lipoprotein disorders in patients with premature coronary artery disease", 85:2025-2033, (1992).
- Geppert, et al. "Microalgal docosahexaenoic acid decreases plasma triacylglycerol in normolipidaemic vegetarians: a randomized trial." *British Journal of Nutrition*, 95, 779-786. (2006).
- Gillet L, Roger S, Bougnoux P, Le Guennec JY, Besson P. Beneficial effects of omega-3 long-chain fatty acids in breast cancer and

(56)

## References Cited

## OTHER PUBLICATIONS

cardiovascular diseases: voltage-gated sodium channels as a common feature? *Biochim. Jan.* 2011 (epub Feb. 16, 2010); 93:4-6.

Gillies, et al. "Effect of a Novel Eicosapentaenoic Acid-Rich Oil on Serum Cholesterol in Man," DuPont 2010.

Ginsberg HN, Elam MB, Lovato LC, Crouse Jr, 3rd, Leiter LA, Linz P, Friedewald WT, Buse JB, Gerstein HC, Probstfield J, Grimm RH, Ismail-Beigi F, Bigger JT, Goff DC, Jr., Cushman WC, Simons-Morton DG, Byington RP. Effects of combination lipid therapy in type 2 diabetes mellitus. *N. Engl. J. Med.* Apr. 29, 2010;362:1563-1574.

Ginsberg HN, Elam MB, Lovato LC, et al, for the ACCORD Study Group. Effects of combination lipid therapy in Type 2 diabetes mellitus. *N Engl J Med* 362:1563-1574, 2010.

Ginsberg HN. "Hypertriglyceridemia: new insights and new approaches to pharmacologic therapy," *Am J Cardiol*, 87:1174-1180 (2001).

Girotti A W. Lipid hydroperoxide generation, turnover, and effector action in biological systems. *J. Lipid Res.* 1998;39(8):1529-1542.

GISSI-HF Investigators. Effect of n-3 polyunsaturated fatty acids in patients with chronic heart failure (the GISSI-HF trial): a randomised, double-blind, placebo-controlled trial. *Lancet.* Oct. 4, 2008 (epub Aug. 29, 2008); 372(9645):1223-1230.

GISSI-Prevenzione Investigators, "Dietary Supplementation with n-3 Polyunsaturated Fatty Acids and Vitamin E after Myocardial Infarction: Results of the GISSI-Prevenzione Trial", *The Lancet*, 354:447-455, (Aug. 7, 1999).

Glod, "Recent Advances in the Pharmacotherapy of Major Depression", *Arch. Psychiatr. Nurs.*, 10(6):355-364 (Dec. 1996).

Goff DC, Lloyd-Jones DM, Bennett G, et al. ACC/AHA Prevention Guideline: 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation.* Jun. 24, 2014 (epub Nov. 12, 2013); 129:S74-S75.

Goldberg, A C: "Combination therapy of dyslipidemia," *Current Treatment Options in Cardiovascular Medicine* 200708 GB, vol. 9, No. 4, pp. 249-258 (2007).

Goodman & Gilman (Robert W. Mahley & Thomas P. Bersot) *Drug Therapy for Hypercholesterolemia and Dyslipidemia*, in Goodman & Gilman's *the Pharmacological Basis fo Therapeutics* 971 (Hardman et al., eds 10th ed. 2001)(32 pages).

Gordon, DJ. et al., High density lipoprotein cholesterol and cardiovascular disease: four prospective American studies. *Circulation.* 79: 8-15. (1989).

Gorritz JL et al., "Rhabdomyolysis and Acute Renal Failure Associated with Gemfibrozil Therapy," *Nephron* 74(2): 437-438 (1996).

Gorritz, JL, "Rhabdomyolysis and Acute Renal Failure Associated with Bezafibrate Treatment," *Nephrol Dial Transplant* 10(12):2371-2372 (1995).

Gosai, P. et al. Effect of omega-3-acid ethyl esters on the steady-state plasma pharmacokinetics of rosuvastatin in healthy adults. *Expert Opin. Pharmacother.* 9, 2947-2953 (2008).

Goto, Y. et al., "Clinical Pharmacological Trial of Ethyl Icosapentate (MND-21)-Dose Finding Study." *Journal of Clinical Therapeutic & Medicines* 8:1293-309 (1992).

Gould, A.L., et al., "Cholesterol reduction yields clinical benefit: impact of statin trials." *Circulation*, 97:946-952 (1998).

Greenblatt DJ, von Moltke LL. Interaction of warfarin with drugs, natural substances, and foods. *J. Clin. Pharmacol.* Feb. 2005 45 (2): 127-32.

Grenyer, Brin F.S., et al., "Fish Oil Supplementation in the Treatment of Major Depression: A Randomised Double-Blind Placebo-Controlled Trial", *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 31:1393-1396, (2007).

Griffin, M.D., et al., "Effects of altering the ratio of dietary n-6 to n-3 fatty acids on insulin sensitivity, lipoprotein size, and postprandial lipemia in men and postmenopausal women aged 45-70 y: the OPTILIP Study." *Am J Clin Nutr* 84:1290-8 (2006).

Grimsgaard et al., "Effects of Highly Purified Eicosapentaenoic Acid and Docosahexaenoic Acid on Hemodynamics in Humans" *American Society for Clinical Nutrition*, 68:52-9, (1998).

Grimsgaard, Kaare H. Bonna, John-Bjarne Hansen, and Arne Nordoy, "Highly purified eicosapentaenoic acid and docosahexaenoic acid in humans have similar triacylglycerol-lowering effects but divergent effects on serum fatty acids" *Am J Clin Nutr*, 66:649-659, (1997).

Grundy S.M et al., Efficacy, safety, and tolerability of once-daily niacin for the treatment of dyslipidemia associated with type 2 diabetes: results of the Assessment of Diabetes Control and Evaluation of the Efficacy of Niaspan Trial. *Arch. Intern. Med.* 162: 1568-1576 (2002).

Grundy SM, et al. Implications of Recent Clinical Trials for the National Cholesterol Education Program Adult Treatment Panel III Guidelines. *Circulation.* 2004; 110:227-39.

Grundy, Scott M., "Low-Density Lipoprotein, Non-High-Density Lipoprotein, and Apolipoprotein B as Targets of Lipid-Lowering Therapy" *Circulation.* 106:2526-2529 (2002).

Guallar, E., et al., "Omega-3 fatty acids in adipose tissue and risk of myocardial infarction—The EURAMIC study." *Arterioscler. Thromb. Vasc. Biol.*, 19:1111-1118 (1999).

Guillot, et al., "Increasing intakes of the long-chain omega-3 docosahexaenoic acid: effects on platelet functions and redox status in healthy men," *The FASEV Journal*, vol. 23, pp. 2909-2916 (2009).

Guizy, M., et al., "ω-3 and ω-6 Polyunsaturated fatty acids block HERG channels." *Am J Physiol Cell Physiol* 289:C1251-C1260 (2005).

Gyarmathy, M., "Selection from the industrial manufacturing. 5th part: Gelatine capsules. 5/2 part: Soft gelatine capsules," *Gyogyszereszet*, vol. 38, No. 2, pp. 105-109 (1994) (with English summary).

Hakonarson, H., et al., "Effects of a 5-lipoxygenase—activating protein inhibitor on biomarkers associated with risk of myocardial infarction—a randomized trial." *JAMA*, 293(8):2245-56 (May 11, 2005).

Hall, W. L., et al., "A high-fat meal enriched with eicosapentaenoic acid reduces postprandial arterial stiffness measured by digital volume pulse analysis in healthy men." *J. Nutr.* 138: 287-291 (Feb. 2008).

Hamazaki et al., "Docosahexaenoic Acid-Rich Fish Oil Does Not Affect Serum Lipid Concentrations of Normolipidemic Young Adults", *American Institute of Nutrition*, 126(11):2784-2789, Nov. (1996).

Hamazaki et al., "Effects of fish oil rich in eicosapentaenoic acid on serum lipid in hyperlipidemic hemodialysis patients," *Kidney Int'l.*, 26:81-84 (Jul. 1984).

Hamazaki et al., "Effects of Orally Administered Ethyl Ester of Eicosapentaenoic Acid (EPA: C20:5, omega-3) On PG12-Like Substance Production by Rat Aorta" *Prostaglandins*, vol. 23 No. 4, pp. 557-567 (Apr. 1982).

Hamazaki T. et al., "Reduction of microalbuminuria in diabetics by Eicosapentaenoic acid ethyl ester" *Lipids.* 25 (9):542-5 (Sep. 1990).

Hampel H, Abraham NS, El-Se rag HB. Meta-analysis: obesity and the risk for gastroesophageal reflux disease and its complications. *Ann. Intern. Med.* Aug. 2005 143 (3): 199-211.

Han, J. J., et al., "Enhancement of both reaction yield and rate of synthesis of structured triacylglycerol containing eicosapentaenoic acid under vacuum with water activity control." *Ligids* 34:989-995 (Sep. 1999).

Hanasaki, K., et al., "Potent modification of low density lipoprotein by group X secretory phospholipase A2 is linked to macrophage foam cell formation." *J. Biol. Chem.* 277(32):29116-24 (Aug. 9, 2002).

Haney, E.M., et al., "Screening for lipid disorders in children and adolescents; Systematic evidence review for the U.S. Preventive Services Task Force (evidence synthesis)." No. 47. Rockville, MD: Agency for Healthcare Research and Quality, US Department of Health and Human Services; AHRQ Publication No. 07-0598-EF-1; Jul. 2007. Available at: <http://www.uspreventiveservicestaskforce.org/uspstf07/chlipid/chlipidsyn.pdf>. (Accessed Mar. 23, 2011)(573 pages).

Hannah, J., et al., "Effect of dietary fatty acids on LDL binding." *Ann N Y Acad Sci.*, 683:178-182 (Jun. 14, 1993).

Hansen et al., "Comparative effects of prolonged intake of highly purified fish oils as ethyl ester or triglyceride on lipids, haemostasis and platelet function in normolipaemic men." *Eur. J. Clin. Nutr.* 47(7):497-507 (Jul. 1993).

(56)

## References Cited

## OTHER PUBLICATIONS

- Hansen, J.B., et al., "Effects of highly purified eicosapentaenoic acid and docosahexaenoic acid on fatty acid absorption, incorporation into serum phospholipids and postprandial triglyceridemia." *Lipids* 33:131-38 (Feb. 1998).
- Harada-Shiba et al., *Journal of Clinical and Experimental Medicine*, Jun. 30, 2007, vol. 221, No. 13, pp. 1068-1073 (with English translation).
- Harris WS. International recommendations for consumption of long-chain omega-3 fatty acids. *J Cardiovasc Med (Hagerstown)* 8(suppl 1):S50-S52, Sep. 2007.
- Harris, "n-3 Fatty acids and lipoproteins: a comparison of results from human and animal studies," *Lipids* 31, 243-252 (Mar. 1996).
- Harris, W. S. et al. "Safety and efficacy of Omacor in severe hypertriglyceridemia," *Journal of Cardiovascular Risk*, 4:385-391 (Oct.-Dec. 1997).
- Harris, W. W S., "Fish oils and plasma lipid and lipoprotein metabolism in humans: a critical review." *J Lipid Res.* 30:785-807 (Jun. 1989).
- Harris, W. W S., "The omega-3 index as a risk factor for coronary heart disease." *Am J Clin Nutr* 87:1997S-2002S (Jun. 2008).
- Harris, W. S., et al., "n-3 Fatty acids and urinary excretion of nitric oxide metabolites in humans." *Am. J. Clin. Nutr.*, 65:459-64 (Feb. 1997).
- Harris, W. S., et al., "Influence of n-3 fatty acid supplementation on the endogenous activities of plasma lipases." *Am. J. Clin. Nutr.* 66:254-60 (Aug. 1997).
- Harris, W.S., "Expert opinion: omega-3 fatty acids and bleeding—cause for concern?" *The American Journal of Cardiology* 99(6A): 45C-46C (Mar. 19, 2007).
- Harris, W.S., "n-3 Fatty acids and human lipoprotein metabolism: an update." *Lipids* 34:S257-S258 (1999).
- Harris, W.S., "n-3 Fatty acids and serum lipoproteins: human studies." *Am J Clin Nutr* 65:164S-54S (1997).
- Harris, W.S., "Omega-3 fatty acids in cardiac biopsies from heart transplantation patients." *Circulation* 110:1645-1649 (Sep. 21, 2004).
- Harris, W.S., et al., "Comparison of the effects of fish and fish-oil capsules on the n-3 fatty acid content of blood cells and plasma phospholipids." *Am J Clin Nutr* 86:1621-5 (Dec. 2007).
- Harris, W.S., et al., "Omega-3 fatty acids and coronary heart disease risk: Clinical and mechanistic perspectives." *Atherosclerosis* 197:12-24 (Mar. 2008)(epub Dec. 26, 2007).
- Harris, W.S., et al., "Stearidonic acid increases the red blood cell and heart eicosapentaenoic acid content in dogs." *Lipids* 42:325-333 (Apr. 2007)(epub Mar. 9, 2007).
- Harris, W.S., et al., "Tissue n-3 and n-6 fatty acids and risk for coronary heart disease events." *Atherosclerosis* 193:1-10 (Jul. 2007)(epub May 15, 2007).
- Hartweg, J., et al., "Potential impact of omega-3 treatment on cardiovascular disease in type 2 diabetes." *Curr Opin Lipidol.*, 20:30-38 (Feb. 2009).
- Hata et al, *Geriatric Medicine*, 30 (5), 799-852, 1992 (with English introduction).
- Hawthorne, et al., "High dose eicosapentaenoic acid ethyl ester: effects on lipids and neutrophil leukotriene production in normal volunteers." *Br. J. Clin. Pharmacol.*, vol. 30, 187-194 (Aug. 1990).
- Hayashi et al., *Decreases in Plasma Lipid Content and Thrombotic Activity by Ethyl Icosapentate Purified from Fish Oiles*, *Current Therapeutic Research*, vol. 56, No. 1, pp. 24-31 (1995).
- Herbette L, Marquardt J, Scarpa A, Blasie JK. A direct analysis of lamellar x-ray diffraction from hydrated oriented multilayers of fully functional sarcoplasmic reticulum. *Biophys. J.* Nov. 1977;20(2):245-272.
- Hibbeln, J. R., et al., "Healthy intakes of n-3 and n-6 fatty acids: estimations considering worldwide diversity." *Am J Clin Nutr.* 83:1483S-93S (Jun. 2006).
- Higashihara et al. "Effects of Eicosapentaenoic Acid on Biochemical Failure after Radical Prostatectomy for Prostate Cancer," *in vivo* 24:561-566 (Jul./Aug. 2010).
- Hilpert, K.F., et al., "Postprandial effect of n-3 polyunsaturated fatty acids on apolipoprotein B—containing lipoproteins and vascular reactivity in type 2 diabetes." *Am J Clin Nutr* 85:369-76 (Feb. 2007).
- Hirafuji, M., et al., "Docosahexaenoic acid potentiates interleukin-1 beta induction of nitric oxide synthase through mechanism involving p44/42 MAPK activation in rat vascular smooth muscle cells." *British Journal of Pharmacology* 136:613-619 (Jun. 2002).
- Hirai, A., et al., "The effects of the oral administration of fish oil concentrate on the release and the metabolism of [14C] arachidonic acid and [14C] eicosapentaenoic acid by human platelets", *Thromb. Res.*, 28:285-298, (Nov. 1, 1982).
- Hirano T, Ito Y, Koba S, Toyoda M, Ikejiri A, Saegusa H, Yamazaki J, Yoshino G. Clinical significance of small dense low-density lipoprotein cholesterol levels determined by the simple precipitation method. *Arterioscler. Thromb. Vase. Biol.* Mar. 2004;24(3):558-563.(epub Jan. 15, 2004).
- Hirano, R., et al., "Regulation by long-chain fatty acids of the expression of cholesteryl ester transfer protein in HepG2 cells." *Lipids*, 36:401-406 (Apr. 2001).
- Hofacer R, et al., Omega-3 fatty acid deficiency increases stearoyl-CoA desaturase expression and activity indices in rat liver: Positive association with non-fasting plasma triglyceride levels, Prostaglandins Leukot. Essent. Fatty Acids. *Jan./Feb. 2012;86:71-7.* (epub Nov. 1, 2011).
- Hoffman, "Atherosclerosis: Prevention through the Ages," WebMD, <https://www.webmed.com/heart/features/atherosclerosis-prevention-through-ages#1>, (Dec. 4, 2007).
- Hohenester, "Primary Biliary Cirrhosis," *Semin Immunopathol.* 31L:283-307, 285 (Sep. 2009)(epub Jul. 15, 2009).
- Holmeide, A. K., et al., "Oxidative degradation of eicosapentaenoic acid into polyunsaturated aldehydes." *Tetrahedron* 59:7157-7162 (2003).
- Holub, B.J., PhD, "Fish Oils and Cardiovascular Disease", *Canadian Medical Association Journal*, 141(10)1063 (Nov. 15, 1989).
- Holvoet P, Kritchevsky SB, Tracy RP, Mertens A, Rubin SM, Butler J, Goodpaster B, Harris TB. The metabolic syndrome, circulating oxidized LDL, and risk of myocardial infarction in wellfunctioning elderly people in the health, aging, and body composition cohort. *Diabetes.* Apr. 2004;53(4):1068-1073.
- Hom et al., "Soft Gelatin Capsules II: Oxygen Permeability Study of Capsule Shells," *J Pharm Sci.* (May 1975) 64(5):851-857.
- Hombeck, M., et al., "Biosynthesis of the algal pheromone fucoseratene by the freshwater diatom *Asterionella formosa* (Bacillariophyceae)." *Tetrahedron* 54:11033-11042 (1998).
- Hong KN, Fuster V, Rosenson RS, Rosendorff C, Bhatt DL. How low to go with glucose, cholesterol, and blood pressure in primary prevention of CVD. *J Am Coll Cardiol* 70(17):2171-85 (publication date Oct. 24, 2017; epublication date Oct. 16, 2017).
- Hoogeveen EK, Geleijnse JM, Kromhout D, et al. No effect of n-3 fatty acids supplementation on NT-proBNP after myocardial infarction: the Alpha Omega Trial. *Eur J Prev Cardiol.* May 2015;22:648-55.
- Horrobin, D.F. The Phospholipid Concept of Psychiatric Disorders and its Relationship to the Neurodevelopmental Concept of Schizophrenia. In M. Peet (ed.) *Phospholipid Spectrum Disorder in Psychiatry* pp. 1-19 (1999).
- Hoskins et al., "Combination use of statins and omega-3 fatty acids: an emerging therapy for combined hyperlipidemia," *Abstract*, 1(5): 579-591(13) (2006).
- Howe, P.R.C., et al., "Equal antithrombotic and triglyceride-lowering effectiveness of eicosapentaenoic acid-rich and docosahexaenoic acid-rich fish oil supplements." *Lipids* 34:S307-S308 (1999).
- HPS2-thrive Collaborative Group, "randomized placebo-controlled trial in 25 673 high-risk patients of er niacin/laroprant: Trial design, pre-specified muscle and liver outcomes, and reasons for stopping study treatment." *Eur. Heart J.* May 2013;34:1279-1291.
- HPS2-Thrive Collaborative Group, Landray MJ, Haynes R, et al. Effects of extended-release niacin with laropiprant in high-risk patients. *N Engl J Med.* Jul. 17, 2014; 371 (3):203-12.
- Hruska MW, Amico JA, Langae TY, Ferrell RE, Fitzgerald SM, Frye RF. The effect of trimethoprim on CYP2C8 mediated rosiglitazone

(56)

## References Cited

## OTHER PUBLICATIONS

metabolism in human liver microsomes and healthy subjects. *Br. J. Clin. Pharmacol.* Jan. 2005;59:70-79.

Hughes et al., "Fish oil produces an atherogenic lipid profile in hypertensive men," *Atherosclerosis*, 84, pp. 229-237 (Oct. 1990).

Hulthe J, Hulten LM, Fagerberg B. Low adipocyte-derived plasma protein adiponectin CJ concentrations are associated with the metabolic syndrome and small dense low-density lipoprotein particles: atherosclerosis and insulin resistance study. *Metab. Clin. Exp. Dec.* 2003;52(12):1612-1614.

Huntington's Disease Drug Works—The DHA Dilemma [http://hddrugworks.org/index2.php?option=com\\_content&task=view&id=185&pop=1&pa...](http://hddrugworks.org/index2.php?option=com_content&task=view&id=185&pop=1&pa...) Printed on Aug. 22, 2008.(2 pages).

Ignarro LJ, Buga GM, Wood KS, Byrnes RE, Chaudhuri G. Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proc. Natl. Acad. Sci. USA.* Dec. 1987;84:9265-9269.

Illingworth, DR, et al., "Comparative effects of lovastatin and niacin in primary hypercholesterolemia: A prospective trial", *Arch. Int. Med.*, 154:1586-1595, (Jul. 25, 1994).

Inoue, I., et al., "Expression of peroxisome proliferator-activated receptor  $\alpha$  (PPAR $\alpha$ ) in primary cultures of human vascular endothelial cells." *Biochem. Biophys. Res. Comm.*, 246, 370-374 (May 19, 1998).

Inzucchi et al., "Diagnosis of Diabetes," *New Engl. Journ of Med.*, 367(6):541-550 (Aug. 9, 2012).

Ishida, Y., et al., " $\alpha$ -Lipoic Acid and Insulin Autoimmune Syndrome." *Diabeters Care*, 30(9): 2240-41 (Sep. 2007).

Isley, et al., "Pilot study of combined therapy with  $\omega$ -3 fatty acids and niacin in atherogenic dyslipidemia," *Journal of Clinical Lipidology*, 1, 211-217 (Jul. 2007).

Itoh et al., "Increased adiponectin secretion by highly purified eicosapentaenoic acid in rodent models of obesity and human obese subjects," *Arterioscler. Thromb. Vasc. Biol.*, pp. 1918-1925 (together with online Supplements 1-15) (Sep. 2007)(epub Jun. 14, 2007).

Jacob RF, Mason RP. Lipid peroxidation induces cholesterol domain formation in model membranes. *J. Biol. Chem.* Nov. 25, 2005;280(47):39380-39387.(epub Sep. 28, 2005).

Jacob RF, Walter MF, Self-Medlin Y, Mason RP. Atorvastatin active metabolite inhibits oxidative modification of small dense low-density lipoprotein. *J. Cardiovasc. Pharmacol.* Aug. 2013;62(2):160-166.

Jacobson et al. "Hypertriglyceridemia and Cardiovascular Risk Reduction", *Clinical Therapeutics*, vol. 29 pp. 763-777 (May 2007).

Jacobson TA. Opening a new lipid "apo-theary": incorporating apolipoproteins as potential risk factors and treatment targets to reduce cardiovascular risk. *Mayo Clin. Proc.* Aug. 2011;86:762-780.

Jacobson, T. Secondary Prevention of Coronary Artery Disease with Omega-3 Fatty Acids. *Am J Cardiol*; 98 [suppl]: 61i-70i (Aug. 21, 2006).

Jacobson, T.A., "Role of n-3 fatty acids in the treatment of hypertriglyceridemia and cardiovascular disease." *Am J Clin Nutr* 87:1981S-90S (Jun. 2008).

Jacobson, T.A., et al., "Effects of eicosapentaenoic acid and docosahexaenoic acid on low-density lipoprotein cholesterol and other lipids: A review." *J. Clin. Lipidology*, vol. 6, pp. 5-18 (Jan./Feb. 2012).

Jakus V, Rietbrock N. Advanced glycation end-products and the progress of diabetic vascular complications. *Physiol. Res.* 2004;53(2): 131-142.

Jenner, "Presymptomatic Detection of Parkinson's Disease". *J Neural Transm Suppl.*, 40:23-36. (Abstract only) (1993).

Jialal I, Devaraj S. Antioxidants and atherosclerosis: Don't throw out the baby with the bath water. *Circulation.* Feb. 25, 2003;107:926-928.

Jialal, I. "Editorial Remnant lipoproteins: measurement and clinical significance." *Clinical Chemistry* 48(2):217-219 (Feb. 2002).

Jinno Y, Nakakuki M, Kawano H, Notsu T, Mizuguchi K, Imada K. Eicosapentaenoic acid administration attenuates the pro-

inflammatory properties of VLDL by decreasing its susceptibility to lipoprotein lipase in macrophages. *Atheroscler. Dec.* 2011;219:566-572.(epub Oct. 4, 2011).

Jong et al., "Role of ApoCs in Lipoprotein Metabolism: Function Differences Between ApoC1, ApoC2, and ApoC3," *Arteriosclerosis, Thrombosis and Vascular Biology.* (Mar. 1999) 19(3):472-484.

Jørgensen AB, Frikke-Schmidt R, Nordestgaard BG, Tybjaerg-Hansen A. Loss-of-function mutations in APOC3 and risk of ischemic vascular disease. *N Engl J Med.* Jul. 3, 2014; 371(1):32-41 (epub Jun. 18, 2014).

*Journal of Practical Pharmacy*, "Hyperlipidemia Drug," 58(4)1 303-1324 (2007) (with English abstract).

*Journal of the Japan Diabetes Society*, "The Relationship Between Postprandial ApoB48 Increase and Insulin Resistance in Type-2 Diabetes," 55(Suppl. 1):5310 (Apr. 2012) (with English Translation)(2 pages).

*Journal of the Japanese Diabetes Society*, "A Case of Familial Combined Hyperlipidemia Associated with Obesity, Type 2 Diabetes Mellitus and Severe Hypertriglyceridemia," 51(3), pp. 233-237 (Mar. 30, 2008) (with English abstract).

Jun M, Foote C, Lv J, et al. Effects of fibrates on cardiovascular outcomes: a systematic review and meta-analysis. *Lancet* 375 (9729):1875-1884, May 29, 2010 (epub May 10, 2010).

Jung, U.J., et al., "n-3 Fatty acids and cardiovascular disease: mechanisms underlying beneficial effects." *Am J Clin Nutr* 87: 2003S-9S (Jun. 2008).

Kamanna et al., "Mechanism of Action of Niacin," *The American Journal of Cardiology* (Apr. 17, 2008), 101(8), S20-S26.

Kamido et al., Lipid Composition of Platelets from Patients with Atherosclerosis:Effect of Purified Eicosapentaenoic Acid Ethyl Ester Administration, Oct. 1988, *Lipids*, 23, pp. 917-923 [Abstract only] (7 pages).

Kaminski WE, Jendraschak E, Kiefl R, et al. Dietary omega-3 fatty acids lower levels of platelet-derived growth factor mRNA in human mononuclear cells. *Blood* Apr. 1993, 81(7): 1871-9.

Kanayasu, T., et al., "Eicosapentaenoic acid inhibits tube formation of vascular endothelial cells in vitro." *Lipids* 26:271-276 (Apr. 1991).

Kastelein et al., Omega-3 Free Fatty Acids for the Treatment of Severe Hypertriglyceridemia: The EpanoVa for Lowering Very High Triglycerides (EVOLVE) Trial, *J. Clin. Lipidol. (JACL)* 597 Jan./Feb. 2014 (epub Oct. 14, 2013).

Katan, M. B., et al., "Kinetics of the incorporation of dietary fatty acids into serum cholesteryl esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study." *J. Lipid Res.* 38: 2012-2022 (Oct. 1997).

Katayama et al., Effect of long-term administration of ethyl eicosapentate (EPA-E) on local cerebral blood flow and glucose utilization in stroke-prone spontaneously hypertensive rats (SHRSP), *Brain Research*, vol. 761, pp. 300-305 (Dec. 31, 1997).

Katayama et al., "Efficacy and Safety of Ethyl Icosapentate (Epano) Given for a Long Term Against Hyperlipidemia," *Prog. Med.*, 21:457-467 (2001) (with English translation).

Kato, T., et al., "Palmitate impairs and eicosapentaenoate restores insulin secretion through regulation of SREBP-1c in pancreatic islets." *Diabetes*, 57(9):2382-2392 (2008) (published online May 5, 2008.).

Kawamura et al., "Effects of 4 weeks' intake of polyunsaturated fatty acid ethylester rich in eicosapentaenoic acid (ethylester) on plasma lipids, plasma and platelet phospholipid fatty acid composition and platelet aggregation; a double blind study," *Nihon Naika Gakkai Zasshi*, 72(1):18-24 (Jan. 10, 1983).

Kawano, H., et al., "Changes in aspects such as the collagenous fiber density and foam cell size of atherosclerotic lesions composed of foam cells, smooth muscle cells and fibrous components in rabbits caused by all-cis 5, 8, 11, 14, 17-icosapentaenoic acid," *J. Atheroscler. Thromb.*, 9:170-177, (2002).

Kawashima, H., et al., "Oral Administration of Dihomo- $\gamma$ -Linolenic Acid Prevents Development of Atopic Dermatitis in NC/Nga Mice." *Lipids* 43:37-43 (Jan. 2008)(epub Nov. 6, 2007).

Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesaniemi Y A, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso



(56)

## References Cited

## OTHER PUBLICATIONS

- M. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): Randomised controlled trial. *Lancet*. Nov. 26, 2005;366:1849-1861.
- Kelley, D. S., et al., "Docosahexaenoic Acid Supplementation Decreases Remnant-Like Particle-Cholesterol and Increases the (n-3) Index in Hypertriglyceridemic Men." *J. Nutr.* 138: 30-35 (Jan. 2008).
- Kelley, et al., "Docosahexaenoic acid supplementation improves fasting and postprandial lip profiles in hypertriglyceridemic men." *The American Journal of Clinical Nutrition*, 86: 324-333 (Aug. 2007).
- Kellner-Weibel G, Yancey PG, Jerome WG, Walser T, Mason RP, Phillips MC, Rothblat GH. Crystallization of free cholesterol in model macrophage foam cells. *Arterioscler. Thromb. Vasc. Biol.* Aug. 1999;19(8):1891-1898.
- Kendall BJ, Macdonald GA, Hayward NK, et al. The risk of Barrett's esophagus associated with abdominal obesity in males and females. *Int. J. Cancer* May 2013 132 (9): 2192-9.
- Kerr, S., Brosnan MJ, Mcintyre M, Reid JL, Dominiczak AF, Hamilton CA. Superoxide anion production is increased in a model of genetic hypertension role of the endothelium. *Hypertension*. Jun. 1999;33:1353-1358.
- Kew, S., et al., "Effects of oils rich in eicosapentaenoic and docosahexaenoic acids on immune cell composition and function in healthy humans." *Am J Clin Nutr* 79:674-81 (Apr. 2004).
- Kholodov et al., "Clinical Pharmacokinetics," *M. Medicine*. (1985) pp. 89-98, 134-138, 160, 378-380 [with English Summary](27 pages).
- Khoueiry G, Rafeh NA, Sullivan E, et al. Do omega-3 polyunsaturated fatty acids reduce risk of sudden cardiac death and ventricular arrhythmias? A meta-analysis of randomized trials. *Heart and Lung*. Jul./Aug. 2013;42:251-256. (epub May 25, 2013).
- Kim F, Tysseling KA, Rice J, Gallis B, Haji L, Giachelli CM, Raines EW, Corson MA, Schwartz MW. Activation of IKKbeta by glucose is necessary and sufficient to impair insulin signaling and nitric oxide production in endothelial cells. *J. Mol. Cell. Cardiol.* Aug. 2005;39(2):327-334.
- Kim KA, Park PW, Kim HK, Ha JM, Park JY. Effect of quercetin on the pharmacokinetics of rosiglitazone, a CYP2C8 substrate, in healthy subjects. *J. Clin. Pharmacol.* Aug. 2005;45:941-946.
- Kimura, F., et al., "Long-term supplementation of docosahexaenoic acid-rich, eicosapentaenoic acid-free microalgal oil in n-3 fatty acid-deficient rat pups." *Biosci. Biotechnol. Biochem.*, 72(2):608-610 (Feb. 2008).
- Kinoshita, "Anti-hyperlipidemic agents," *Nihon Rinsho*, 60(5):968-74 (May 2002) (with English Abstract)(11 pages).
- Kinsella, J.E., et al., "Dietary n-3 polyunsaturated fatty acids and amelioration of cardiovascular disease: possible mechanisms." *Am J Clin Nutr* 52:1-28 (Jul. 1990).
- Kitada, 9th Diabetes Drug and Drug Related Seminar Diabetes Q&A, Kanazawa Medical University, Diabetes and Endocrine Internal Medicine (<http://plaza.umin.ac.jp/iby/etcdata/yakuyaku110410.pdf>)(Apr. 10, 2011) (with English translation)(105 pages).
- Klempfner R, Erez A, Sagit BZ, et al. Elevated triglyceride level is independently associated with increased all-cause mortality in patients with established coronary heart disease: Twenty-two-year follow-up of the Bezafibrate Infarction Prevention Study and Registry. *Circ Cardiovasc Qual Outcomes* 9(2):100-8 (publication date Mar. 8, 2016).
- Knapp HR. Dietary fatty acids in human thrombosis and hemostasis. *Am. J. Clin. Nutr.* May 1997 65 (5 Suppl): 1687S-98S.
- Knopp, R.H., et al., "Contrasting effects of unmodified and time-release forms of niacin on lipoproteins in hyperlipidemic subjects: clues to mechanism of action of niacin", *Metabolism*, 34:642-650, (Jul. 1985).
- Koba S, Hirano T, Ito Y, Tsunoda F, Yokota Y, Ban Y, Iso Y, Suzuki H, Katagiri T. Significance of small dense low-density lipoprotein-cholesterol concentrations in relation to the severity of coronary heart diseases. *Atherosclerosis*. Nov. 2006;189(1):206-214. (epub Jan. 18, 2006).
- Kohno, M., et al., "Inhibition by Eicosapentaenoic Acid of Oxidized-LDL- and Lysophosphatidylcholine-Induced Human Coronary Artery Smooth Muscle Cell Production of Endothelin." *J. Vasc. Res.* 38:379-388 (Jul./Aug. 2001).
- Kojda G, Harrison DG. Interactions between no and reactive oxygen species: Pathophysiological importance in atherosclerosis, hypertension, diabetes and heart failure. *Cardiovasc. Res.* Aug. 15, 1999;43:562-571.
- Kojima, T, et al., "Long-term administration of highly purified eicosapentaenoic acid provides improvement of psoriasis." *Dermatologica*, 182:225-230 (1991).
- Koroshetz, W.J. Huntington's Disease. In Samuels, M. (ed.) *Office Practice of Neurology*, pp. 654-661 (1996).
- Kosonen, O., et al., "Inhibition by nitric oxide-releasing compounds of E-selectin expression in and neutrophil adhesion to human endothelial cells." *European Journal of Pharmacology* 394:149-156 (Apr. 7, 2000).
- Koyama et al., Plaque Reduction and Stabilization Observed in Borderline Diabetes Using Coronary CT Angiogram During Administration of Purified Eicosapentaenoic Acid (EPA) *Ther. Res.* 31(2):219-225 (Feb. 2010) (with English translation)(20 pages).
- Krauss RM. Heterogeneity of plasma low-density lipoproteins and atherosclerosis risk. *Curr. Opin. Lipidol.* Oct. 1994;5(5):339-349.
- Kris-Etherton, et al., "Fish Consumption, Fish Oil, Omega-3 Fatty Acids, and Cardiovascular Disease" *Circulation*, 106:2747-2757 (Nov. 19, 2002)(epub Jan. 28, 2003).
- Kris-Etherton, P. M., et al., "Omega-3 Fatty Acids and Cardiovascular Disease—New Recommendations From the American Heart Association." *Arterioscler Thromb Vasc Biol.* 23:151-152 (Feb. 1, 2003).
- Krzynowek et al., "Purification of Omega-3 Fatty Acids from Fish Oils Using HPLC: An Overview," *National Marine Fisheries—Proceedings of the first joint conference of the Tropical and Subtropical Fisheries Technological Society of the Americas with the Atlantic Fisheries Technological Society*, pp. 74-77 (1988).
- Ku, K., et al., "Beneficial Effects of to-3 Fatty Acid Treatment on the Recovery of Cardiac Function After Cold Storage of Hyperlipidemic Rats." *Metabolism*, 48(10):123-1209 (Oct. 1999).
- Kunimoto M, Inoue K, Nojima S. Effect of ferrous ion and ascorbate-induced lipid peroxidation on liposomal membranes. *Biochem. Biophys. Acta.* Aug. 6, 1981;646(1):169-178.
- Kurabayashi, T., et al., "Eicosapentaenoic acid effect on hyperlipidemia in menopausal Japanese women. The Niigata Epadel Study Group" *Obstet Gynecol* 96:521-8 (Oct. 2000).
- Labor Diagnostik Karlsruhe, "Target Values of Lipid Metabolism [Recommendation for lipid plasma levels in Germany]," (exact publication date unknown; circa 2006) (with English abstract)(4 pages).
- Lada et al., "Associations of Low Density Lipoprotein Particle Compositions with Atherogenicity," *Curr. Opin. Lipidol.* (Feb. 2004) 15(1):19-24.
- Lai, E., et al., "Suppression of niacin-induced vasodilation with an antagonist to prostaglandin D2 receptor subtype 1", *Clin. Pharm. & Ther.*, 81 :849-857, (Jun. 2007/epub Mar. 28, 2007).
- Laidlaw, M., et al., "Effects of supplementation with fish oil-derived n-3 fatty acids and  $\gamma$ -linolenic acid on circulating plasma lipids and fatty acid profiles in women." *Am J Clin Nutr* 77:37-42 (Jan. 2003).
- Laird et al., "Relationship of early hyperglycemia to mortality in trauma patients," *J. Trauma*, 56(5):1058-1062 (May 2004).
- Lamb RE, Goldstein BJ. Modulating an Oxidative-Inflammatory Cascade: Potential New Treatment Strategy for Improving Glucose Metabolism, Insulin Resistance, and Vascular Function. *Int. J. Clin. Pract.* Jul. 2008(epub May 16, 2008); 62(7): 1087-1095.
- Lamharzi N, Renard CB, Kramer F, Pennathur S, Heinecke JW, Chait A, Bomfeldt KE. Hyperlipidemia in concert with hyperglycemia stimulates the proliferation of macrophages in atherosclerotic lesions: potential role of glucose-oxidized LDL. *Diabetes*. Dec. 2004;53(12):3217-3225.
- Landmesser U, Dikalov S, Price SR, McCann L, Fukai T, Holland SM, Mitch WE, Harrison DG. Oxidation of tetrahydrobiopterin

(56)

## References Cited

## OTHER PUBLICATIONS

leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J. Clin. Invest.*, Apr. 2003;111:1201-1209.

LaRosa JC. Understanding risk in hypercholesterolemia. *Clin Cardiol* 26(Suppl 1):3-6, Jan. 2003.

Larsen, L.N., et al., "Heneicosapentaenoate (21:5n-3): Its incorporation into lipids and its effects on arachidonic acid and eicosanoid Synthesis." *Lipids* 32:707-714 (Jul. 1997).

Laufs et al., "Upregulation of endothelial nitric oxide synthase by hmg coa reductase inhibitors," *Circulation* (Mar. 31, 1998) 97:1129-1135.

Law TK, Yan AT, Gupta A, et al. Primary prevention of cardiovascular disease: global cardiovascular risk assessment and management in clinical practice. *Eur Heart J Qual Care Clin Outcomes*. 1(1):31-36 (publication date Jul. 2, 2015; epublication date Jul. 1, 2015).

Law, M.R., et al., "Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis." *Br Med J.*, 326:1423-1427 (Jun. 28, 2003).

Lawson et al., "Human absorption of fish oil fatty acids as triacylglycerols, free acids or ethyl esters," *Biochemical and Biophysical Research Communications* 152(1):328-335 (Apr. 15, 1988).

Leaf A, Albert CM, Josephson M, et al. for the Fatty Acid Antiarrhythmia Trial Investigators. Prevention of Fatal Arrhythmias in High-Risk Subjects by Fish Oil n-3 Fatty Acid Intake. *Circ.* Nov. 1, 2005;112:2762-2768.

Leaf A, Kang JX. Prevention of cardiac sudden death by N-3 fatty acids: a review of the evidence. *J Intern Med* 240:5-12, Jul. 1996.

Leaf, "Hypertriglyceridemia: A Guide to Assessment and Treatment," *Hospital Physician* 17-23 (Sep. 2008).

Leaf, A., "Historical overview of n3 fatty acids and coronary heart disease." *Am J Clin Nutr* 87:1978S-80S. (Jun. 2008).

Lee and G.Y.H. Lip, "The Role of Omega-3 Fatty Acids in the Secondary Prevention of Cardiovascular Disease", *Q J Med*, 96:465-480, (Jul. 2003).

Lee C, Sigari F, Segrado T, Horkko S, Hama S, Subbaiah PV, Miwa M, Navab M, Witztum JL, Reaven PD. All ApoB-containing lipoproteins induce monocyte chemotaxis and adhesion when minimally modified. Modulation of lipoprotein bioactivity by platelet-activating factor acetylhydrolase. *Arterioscler. Thromb. Vase. Biol.*, Jun. 1999; 19(6): 1437-1446.

Lee J.H., et al., "Omega-3 fatty acids for cardioprotection." *Mayo Clin Proc.*, 83(3):324-332 (Mar. 2008).

Leigh-Firbank et al., "Eicosapentaenoic acid and docosahexanoic acid from fish oils: differential associations with lipid responses," *Br. J. Nutr.* 87:435-445 (May 2002).

Lemaitre, R.N., et al., "n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study." *Am J Clin Nutr* 77:319-25 (Feb. 2003).

Leonard, Brian E., "Neurological Aspects", *Fundamentals of Psychopharmacology*, 186-187, (1997).

Leucht, S., et al., *Schizophrenia Research*, vol. 35, "Efficacy and extrapyramidal side-effects of the new antipsychotics olanzapine, quetiapine, risperidone, and sertindole compared to conventional antipsychotics and placebo. A meta-analysis of randomized controlled trials", pp. 51-68, (Jan. 4, 1999).

Levey A, et al. A New Equation to Estimate Glomerular Filtration Rate. *Ann Intern Med*. 150:604-612; May 5, 2009.

Li, D., et al., "Effect of dietary  $\alpha$ -linolenic acid on thrombotic risk factors in vegetarian men." *Am J Clin Nutr* 69:872-82 (May 1999).

Li, H., et al., "EPA and DHA reduce LPS-induced inflammation responses in HK-2 cells: Evidence for a PPAR- $\gamma$ -dependent mechanism." *Kidney Int'l.* 67:867-74 (Mar. 2005).

Libby P. Triglycerides on the rise: should we swap seats on the seesaw? *Eur Heart J.* 36(13):774-776 (publication date Apr. 1, 2015; epublication date Dec. 29, 2014).

Libby, "Inflammation and atherosclerosis," *Nature* (Dec. 2002) 420(6917):868-874.

Lichtman et al., "Depression and Coronary Heart Disease, Recommendations for Screening, Referral and Treatment," *AHA Science Advisory, Circulation* 118:1768-1775 (Sep. 29, 2008).

Lien, E.L., "Toxicology and safety of DHA." *Prostaglandins Leukot Essent Fatty Acids.*, 81:125-132 (2009).

Lin, Pao-Yen, M.D., et al., "A Meta-Analytic Review of Double-Blind, Placebo-Controlled Trials of Antidepressant Efficacy of Omega-3 Fatty Acids", *Psychiatry*, 1056-1061 (Jul. 2007).

Lin, Y., et al., "Differential effects of eicosapentaenoic acid on glycerolipid and apolipoprotein B metabolism in primary human hepatocytes compared to HepG2 cells and primary rat hepatocytes." *Biochimica et Biophysica Acta* 1256:88-96 (Apr. 28, 1995).

Lindsey, S., et al., "Low density lipoprotein from humans supplemented with n-3 fatty acids depresses both LDL receptor activity and LDLr mRNA abundance in HepG2 cells." *J Lipid Res.*, 33:647-658 (Mar. 1992).

Lipitor [package insert]. New York, NY: Parke-Davis (2012). (22 pages).

Lipitor [product information] Dublin, Ireland: Pfizer Inc. (2007). (18 pages).

Liu et al., "Effects of stable fish oil and simvastatin on plasma lipoproteins in patients with hyperlipidemia," *Nutrition Res.*, vol. 23, pp. 1027-1034 (Aug. 2003).

Liu X, et al., Stearoyl CoA Desaturase 1: Role in Cellular Inflammation and Stress, *Adv. Nutr.* Jan. 2011 (Jan. 10, 2011); 2:15-22.

Lohmussaar, E., et al., "ALOX5AP Gene and the PDE4D Gene in a Central European Population of Stroke Patients." *Stroke*, 36:731-736 (Apr. 2005)(epub Feb. 24, 2005).

Lovaza (omega-3-acid ethyl esters) Capsules, Prescribing information, GlaxoSmithKline (Nov. 2008).(9 pages).

Lovaza [package insert]. Research Triangle Park, NC: GlaxoSmithKline (2012). (14 pages).

Lovaza Side Effects, web archived webpage, archived from Drugs.com website on (Jul. 31, 2010), Retrieved from URL <<https://web.archive.org/web/20100731021902/https://www.drugs.com/sfx/lovaza-side-effects.html>> (4 pages)(Jul. 2010).

Lovaza TM (omega-3-acid ethyl esters) Capsules, Aug. 2007 (Aug. 1, 2007)m oaget 1-2, XP055589332.

Lovaza United States Prescribing Information, GlaxoSmithKline. Research Triangle Park, USA, May 2014.

Lovaza, (omega-3-acid ethyl esters) Capsules, Prescribing information Smith Kline Beechum (Jul. 2009).(17 pages).

Lovaza, GlaxoSmithKline, Lovaza Prescribing Information, Jun. 2008 [retrieved from the internet Jun. 6, 2012 <[https://web.archive.org/web/20090206170311/http://us.gsk.com/products/assets/us\\_lovaza.pdf](https://web.archive.org/web/20090206170311/http://us.gsk.com/products/assets/us_lovaza.pdf)>]; Table 3, p. 1, section entitled 'Description,' p. 3, section entitled 'Very High Triglycerides: Monotherapy,' p. 4 section entitled 'Indications and Usage' and 'Information for Patients.' (12 pages).

Lovaza® (omega-3-acid ethyl esters) Capsules, Prescribing information, GlaxoSmithKline, (Dec. 2010)(12 pages).

Lovaza®, Physicians' Desk Reference 2699-2701 (62d ed., 2008). (4 pages).

Lovegrove et al., "Moderate fish-oil supplementation reverses low-platelet, long chain n-3 polyunsaturated fatty acid status and reduced plasma triacylglycerol concentrations in British Indo-Asians," *Am. J. Clin. Nutr.*, 79:974-982 (Jun. 2004).

Lu, G., et al., "Omega-3 fatty acids alter lipoprotein subfraction distributions and the in vitro conversion of very low density lipoproteins to lowdensity lipoproteins." *J Nutr Biochem.*, 10:151-158 (Mar. 1999).

Lucas, M., et al., "Ethyl-eicosapentaenoic acid for the treatment of psychological distress and depressive symptoms in middle-aged women: a double-blind, placebo-controlled, randomized clinical trial." *Am J Clin Nutr* 89:641-51 (Feb. 2009)(epub Dec. 30, 2008).

Luria, MH, "Effect of low-dose niacin on high-density lipoprotein cholesterol and total cholesterol/high density lipoprotein cholesterol ratio", *Arch. Int. Med.*, 148:2493-2495, (Nov. 1998).

Lvovich V, Scheeline A. Amperometric sensors for simultaneous superoxide and hydrogen peroxide detection. *Anal. Chem.* Feb. 1, 1997;69:454-462.

(56)

## References Cited

## OTHER PUBLICATIONS

- Madhavi et al., "Effect of n-6 and n-3 fatty acids on the survival of vincristine sensitive and resistant human cervical carcinoma cells in vitro", *Cancer Letters*, vol. 84. No. 1, pp. 31-41 (Aug. 29, 1994).
- Madsen, L., et al., "Eicosapentaenoic and Docosahexaenoic Acid Affect Mitochondrial and Peroxisomal Fatty Acid Oxidation in Relation to Substrate Preference." *Lipids* 34:951-963 (Sep. 1999).
- Mak IT, Weglicki WB. Antioxidant properties of calcium channel blocking drugs. *Methods Enzymol.* 1994;234:620-630.
- Maki et al., "Effects of Adding Prescription Omega-3 Acid Ethyl Esters to Simvastatin (20 mg/day) on Lipids and Lipoprotein Particles in Men and Women with Mixed Dyslipidemia," *Am. J. Cardiol.*, 102:429-433 (Aug. 15, 2008)(Epub May 22, 2008).
- Maki, K.C., et al., "Baseline lipoprotein lipids and low-density lipoprotein cholesterol response to prescription omega-3 acid ethyl ester added to simvastatin therapy." *Am J Cardiol.*, 105:1409-1412 (May 15, 2010)(epub Mar. 30, 2010).
- Maki, PhD, et al., "Lipid Responses to a Dietary Docosahexaenoic Acid Supplement in Men and Women with Below Average Levels of High Density Lipoprotein Cholesterol." *Journal of the American College of Nutrition*, vol. 24, No. 3, 189-199 (Jun. 2005).
- Malinowski et al., "Elevation of Low-Density Lipoprotein Cholesterol Concentration with Over-the-Counter Fish Oil Supplementation." *Annals of Pharmacotherapy* 41:1296-1300 (Jul./Aug. 2007).
- Malinski T, Taha Z. Nitric oxide release from a single cell measured in situ by a porphyrinic-based microsensor. *Nature.* Aug. 20, 1992;358:676-678.
- Mallat, Z., et al., "Apoptosis in the vasculature: mechanisms and functional importance." *British Journal of Pharmacology* 130:947-962 (Jul. 2000).
- Mallat, Z., et al., "Protective role of interleukin-10 in atherosclerosis." *Circ. Res.* 85:e17-e24 (Oct. 15, 1999).
- Manninen V, Tenkanen L, Koskinen P, et al. Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary heart disease risk in the Helsinki Heart Study. Implications for treatment. *Circulation* 85:37-45, Jan. 1992.
- Marangell, Lauren B., M.D., et al., "A Double-Blind, Placebo-Controlled Study of the Omega-3 Fatty Acid Docosahexaenoic Acid in the Treatment of Major Depression", *Am. J. Psychiatry*, 160(5):996-998, (May 2003).
- Marchioli R, Barzi F, Bomba E, et al, GISSI-Prevenzione Investigators. Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: time-course analysis of the results of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation.* 105(16):1897-1903, Apr. 23, 2002.
- Marckmann, P., "Fishing for heart protection." *Am J Clin Nutr*, 78:1-2 (Jul. 2003).
- Marcoux et al., "Plasma remnant-like particle lipid and apolipoprotein levels in normolipidemic and hyperlipidemic subjects," *Atherosclerosis*, vol. 139, pp. 161-171 (Jul. 1998).
- Marder, "An Approach to Treatment Resistance in Schizophrenia," *British Journ. Psychiatry*, 37:19-22 (1999).
- Margolis, Simeon "What is Hyperlipidemia?" (<http://www.healthcommunities.com/highcholesterol/whathyperlipidemia.shtml>, accessed Oct. 20, 2015, published Aug. 25, 2011)(4 pages).
- Martin SS, Blaha MJ, Elshazly MB, et al. Comparison of a novel method vs the Friedewald equation for estimating low-density lipoprotein cholesterol levels from the standard lipid profile. *JAMA.* Nov. 20, 2013;310:2061-8.
- Martinez-Gonzalez J, Raposo B, Rodriguez C, Badimon L. 3-hydroxy-3-methylglutaryl coenzyme a reductase inhibition prevents endothelial no synthase downregulation by atherogenic levels of native ldl: Balance between transcriptional and posttranscriptional regulation. *Arterioscler. Thromb. Vasc. Biol.* May 2001;21:804-809.
- Martinez-Gonzalez, Jose et al., "Estatinas y acidos grasos omega-3. Disminucion de la mortalidad cardiovascular dependiente e independiente de la reduccion de la colesterolemia," (2006) *Rev Esp Cardiol Suppl.*, 6(D):20D-30D [with English abstract].
- Martin-Jadraque, R. et al., Effectiveness of low dose crystalline nicotinic acid in men with low density lipoprotein cholesterol levels. *Arch. Int. Med.* 156: 1081-1088. (May 27, 1996).
- Martz, "Moving Upstream in Huntington's," *Science-Business eXchange*, 2 pgs., Oct. 2008.
- Mason et al., "Comparative lipid antioxidant effects of omega-3 fatty acids in combination with HMG-CoA reductase inhibitors," *Journ. Clin. Lipidology* (May/Jun. 2011) 5(3):201.
- Mason et al., "Direct evidence for cholesterol crystalline domains in biological membranes: role in human pathobiology," *Biochimica et Biophysica Acta* 198-207 (Mar. 10, 2003).
- Mason et al., "Eicosapentaenoic Acid (EPA) inhibits the formation of membrane cholesterol crystalline domains by a potent antioxidant mechanism," *Journ. Clin. Lipid.*, 7(3):272-273 (May/Jun. 2013) [Abstract only].
- Mason et al., "Eicosapentaenoic acid inhibits glucose-induced membrane cholesterol crystalline domain formation through a potent antioxidant mechanism," *Biochim. Biophys. Acta.*, 1848(2):502-9, (Feb. 2015).
- Mason et al., "Eicosapentaenoic Acid Inhibits Oxidation of ApoB-containing Lipoprotein Particles of Different Size In Vitro When Administered Alone or in Combination With Atorvastatin Active Metabolite Compared With Other Triglyceride-lowering Agents," *J. Cardiovasc. Pharmacol.*, 68(1):33-40 (Jul. 2016).
- Mason et al., "Eicosapentaenoic acid reduces membrane fluidity, inhibits cholesterol domain formation, and normalizes bilayer width in atherosclerotic-like model membranes," *Biochim. Biophys. Acta.*, 1858(12):3131-3140 (Dec. 2016).
- Mason RP, Gonye GE, Chester DW, Herbette LG. Partitioning and location of Bay K 8644, 1,4-dihydropyridine calcium channel agonist, in model and biological membranes. *Biophys. J.* Apr. 1989;55(4):769-778.
- Mason RP, Jacob RF, Kubant R, Walter MF, Bellamine A, Jacoby A, Mizuno Y, Malinski T. Effect of enhanced glycemic control with saxagliptin on endothelial nitric oxide release and CD40 levels in obese rats. *J. Atheroscler. Thromb.* Epub Jun. 13, 2011;18:774-783.
- Mason RP, Jacob RF. Membrane microdomains and vascular biology: Emerging role in atherogenesis. *Circulation.* May 6, 2003; 107:2270-2273.
- Mason RP, Kalinowski L, Jacob RF, Jacoby AM, Malinski T. Nebivolol reduces nitrooxidative stress and restores nitric oxide bioavailability in endothelium of black americans. *Circulation.* Dec. 13, 2005 (epub Dec. 5, 2005); 112:3795-3801.
- Mason RP, Kubant R, Heeba G, Jacob RF, Day CA, Medlin YS, Funovics P, Malinski T. Synergistic effect of amlodipine and atorvastatin in reversing ldl-induced endothelial dysfunction. *Pharm. Res.* Aug. 2008 (epub 2007 Dec. 2018); 25:1798-1806.
- Mason RP, Walter MF, Day CA, Jacob RF. Active metabolite of atorvastatin inhibits membrane cholesterol domain formation by an antioxidant mechanism. *J. Biol. Chem.* Apr. 7, 2006 (epub Feb. 7, 2006) ;281(14):9337-9345.
- Mason RP, Walter MF, Day CA, Jacob RF. Intermolecular differences for HMG-CoA reductase inhibitors contribute to distinct pharmacologic and pleiotropic actions. *Am. J Cardiol.* Sep. 5, 2005;96(5A):11F-23F.
- Mason RP, Walter MF, Jacob RF. Effects of hmg-coa reductase inhibitors on endothelial function: Role of microdomains and oxidative stress. *Circulation.* Jun. 1, 2004;109:II34-II41.
- Mason RP, Walter MF, Mason PE. Effect of oxidative stress on membrane structure: Small angle x-ray diffraction analysis. *Free Radic. Biol. Med.* 1997;23(3):419-425.
- Mason RP. Molecular basis of differences among statins and a comparison with antioxidant vitamins. *Am. J. Cardiol.* Dec. 4, 2006 (epub Oct. 10, 2006); 98:34P-41P.
- Mataki et al., "Effect of Eicosapentaenoic Acid in Combination with HMG-CoA Reductase Inhibitor on Lipid Metabolism," *Int. Med. J.* 5(1):35-36 (Mar. 1998).
- Mater, M.K., et al., "Arachidonic acid inhibits lipogenic gene expression in 3T3-L1 adipocytes through a prostanoid pathway." *J. Lipid Res.* 39:1327-1334 (Jul. 1998).

(56)

## References Cited

## OTHER PUBLICATIONS

- Matsumoto, M., et al., "Orally administered eicosapentaenoic acid reduces and stabilizes atherosclerotic lesions in ApoE-deficient mice." *Atherosclerosis*, 197(2):524-533 (Apr. 2008/epub Sep. 4, 2007).
- Matsuzaki et al., "Incremental Effects of Eicosapentaenoic Acid on Cardiovascular Events in Statin-Treated Patients with Coronary Artery Disease," *Circ. J.* 73:1283-1290 (Jul. 2009/epub May 8, 2009).
- Matsuzawa, Y., et al., "Effect of Long-Term Administration of Ethyl Icosapentate (MND-21) in Hyperlipaemic Patients," *J. Clin Therapeutic & Medicines*, 7: 1801-16 (1991).
- Mattson MP. Modification of ion homeostasis by lipid peroxidation: roles in neuronal degeneration and adaptive plasticity. *Trends Neurosci.* Feb. 1998;21(2):53-57.
- Mayatepek, E., et al., *The Lancet*, vol. 352, Leukotriene C4-synthesis deficiency: a new inborn error of metabolism linked to a fatal developmental syndrome, pp. 1514-1517 (Nov. 7, 1998).
- Mayo Clinic at <http://www.mayoclinic.org/diseases-conditions/high-blood-cholesterol/in-depth/cholesterol> (2014)(5 pages).
- Mayo Clinic, *Diabetes Diagnosis and Treatment*, 1998, <http://www.mayoclinic.org/diseases-conditions/diabetes/diagnosis-treatment/drc-20371451> (1998-2018).
- McElroy, S.L., et al., "Clozapine in the Treatment of Psychotic Mood Disorders, Schizoaffective Disorder, and Schizophrenia", *Journal of Clinical Psychiatry*, vol. 52, No. 10, pp. 411-414 (Oct. 1991).
- McIntyre M, Hamilton CA, Rees DD, Reid JL, Dominiczak AF. Sex differences in the abundance of endothelial nitric oxide in a model of genetic hypertension. *Hypertension*. Dec. 1997;30:1517-1524.
- McKenney et al., "Prescription omega-3 fatty acids for the treatment of hypertriglyceridemia," *Am. J. Health Syst. Pharm.*, 64(6):595-605 (Mar. 15, 2007).
- McKenney et al., CMRO, "Comparison of the efficacy of rosuvastatin versus atorvastatin, simvastatin and pravastatin in achieving lipid goals: results from the STELLAR trial", 689-98 (2003).
- McKenney, J., "Niacin for dyslipidemia: considerations in product selection", *Am. J. Health Syst. Pharm.*, 60:995-1005, (May 15, 2003).
- McKenney, J.M. et al. Study of the pharmacokinetic interaction between simvastatin and prescription omega-3-acid ethyl esters. *J. Clin. Pharmacol.* 46, 785-791 (Jul. 2006).
- McKenney, James et al., "Role of prescription omega-3 fatty acids in the treatment of Hypertriglyceridemia," *Pharmacotherapy*, LNKD—Pubmed: 17461707, vol. 27, No. 5, pp. 715-728 (May 2007).
- McKeone et al., "Alterations in serum phosphatidylcholine fatty acyl species by eicosapentaenoic and docosahexaenoic ethyl esters in patients with severe hypertriglyceridemia." *J. Lipid Res.* 38:429-436 (Mar. 1997).
- McMurchie, E.J., et al., "Incorporation and effects of dietary eicosapentaenoate (20 : 5(n-3)) on plasma and erythrocyte lipids of the marmoset following dietary supplementation with differing levels of linoleic acid." *Biochimica et Biophysica Acta*, 1045:164-173 (Jul. 16, 1990).
- McNamara JR, et al., Remnant-like particle (RLP) Cholesterol is an independent cardiovascular disease risk factor in women: results from the Framingham Heart Study, *Atherosclerosis*, vol. 154(1), pp. 229-236 (Jan. 2001).
- MedlinePlus. "Coronary heart disease," Available at: <https://medlineplus.gov/ency/article/007115.htm> (review date Jul. 14, 2015)(accessed Sep. 2, 2016)(5 pages).
- Menuet, R. et al., "Importance and management of dyslipidemia in the metabolic syndrome," *American Journal of the Medical Sciences* 200512 US, vol. 33, No. 6, pp. 295-302 (2005).
- Merched, A.J., et al., "Atherosclerosis: evidence for impairment of resolution of vascular inflammation governed by specific lipid mediators." *FASEB J.* 22:3595-3606 (Oct. 2008/epub Jun. 17, 2008).
- Merkel et al., "Antisense Oligonucleotide Directed to Human Apolipoprotein B-100 Reduces Lipoprotein(a) Levels and Oxidized Phospholipids on Human Apolipoprotein B-100 Particles in Lipoprotein(a) Transgenic Mice," *Circulation*, vol. 118, pp. 743-753 (epub Jul. 28, 2008).
- Mesa, M., "Effects of oils rich in Eicosapentaenoic and docosahexaenoic acids on the oxidizability and thrombogenicity of low-density lipoprotein," *Arteriosclerosis* 175, pp. 333-343 (Aug. 2004).
- Metcalf, R.G. et al., "Effect of dietary n-3 polyunsaturated fatty acids on the inducibility of ventricular tachycardia in patients with ischemic cardiomyopathy." *Am J Cardiol* 101:758-761 (Mar. 15, 2008/epub Jan. 14, 2008).
- Metcalf, R.G., et al., "Effects of fish-oil supplementation on myocardial fatty acids in humans." *Am J Clin Nutr* 85:1222-28 (May 2007).
- Meyer et al., "Comparison of Seal Oil to Tuna Oil on Plasma Lipid Levels and Blood Pressure in Hypertiglyceridaemic Subjects," *Lipids*, 44:827-835 (Sep. 2009).
- Meyer, et al., "Dose-Dependent Effects of Docosahexaenoic Acid Supplementation on Blood Lipids in Statin-Treated Hyperlipidaemic Subjects." *Lipids*, 42:109-115 (Mar. 2007/epub Feb. 8, 2007).
- Meyers, et al., "Nicotinic acid induces secretion of prostaglandin D2 in human macrophages: An in vitro model of the niacin-flush", *Atherosclerosis*, 192:253-258, (Jun. 2007/epub Sep. 1, 2006).
- Micheletta F, Natoli S, Misuraca M, Sbarigia E, Diczfalusy U, Iuliano L. Vitamin E supplementation in patients with carotid atherosclerosis: Reversal of altered oxidative stress in plasma but not in plaque. *Arterioscler. Thromb. Vasc. Biol.* Jan. 2004 (epub Dec. 16, 2006); 24:136-140.
- Michos et al., "Niacin and Statin Combination Therapy for Atherosclerosis Regression and Prevention of Cardiovascular Disease Events," *Journ. Amer. Coll. Cardiol.*, vol. 59, No. 23:2058-2064 (Jun. 5, 2012)(epub Apr. 18, 2012).
- Mii, S., et al., "Perioperative use of eicosapentaenoic acid and patency of infrainguinal vein bypass: a retrospective chart review." *Curr Ther Res Clin Exp.* 68:161-174 (May 2007).
- Miles, et al., "Effect of orlistat in overweight and obese patients with type 2 diabetes treated with metformin," *Diabetes Care*, 25(7):1123-1128 (2002).
- Miller AK, DiCicco RA, Freed MI. The effect of ranitidine on the pharmacokinetics of rosiglitazone in healthy adult male volunteers. *Clin. Ther.* Jul. 2002;24:1062-1071.
- Miller AK, Inglis AM, Culkin KT, Jorkasky DK, Freed MI. The effect of acarbose on the pharmacokinetics of rosiglitazone. *Eur. J. Clin. Pharmacol.* May 2001 ;57:105-109.
- Miller M, Cannon CP, Murphy SA, et al. Impact of triglyceride levels beyond low-density lipoprotein cholesterol after acute coronary syndrome in the PROVE IT-TIMI 22 trial. *J Am Coll Cardiol* 51:724-730, Feb. 19, 2008.
- Miller M, Stone NJ, Ballantyne C, et al. Triglycerides and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation*. May 24, 2011 (epub Apr. 18, 2011); 123:2292-2333.
- Miller M. Current perspectives on the management of hypertriglyceridemia. *Am Heart J* 140:232-40, 2000.
- Miller, M., et al., "Impact of lowering triglycerides on raising HDL-C in hypertriglyceridemic and non-hypertriglyceridemic subjects." *International Journal of Cardiology* 119:192-195 (Jul. 10, 2007)(epub Oct. 18, 2006).
- Minihane, A.M., et al., "ApoE polymorphism and fish oil supplementation in subjects with an atherogenic lipoprotein phenotype." *Arterioscler. Thromb. Vasc. Biol.* 20:1990-1997 (Aug. 2000).
- Mishra, A., et al., "Oxidized omega-3 fatty acids inhibit NF-κB activation via a PPARα-Dependent Pathway." *Arterioscler Thromb Vasc Biol.* 24:1621-1627 (Sep. 2004)(epub Jul. 1, 2004).
- Missouri DURReport, *Statin Therapy (Oct./Nov. 2003) Drug Use Review Newsletter* 8(6):1-9.
- Mita, T. et al., Eicosapentaenoic acid reduces the progression of carotid intima-media thickness in patients with type 2 diabetes, *Atherosclerosis* 191:162-167 (Mar. 2007)(epub Apr. 17, 2006).
- Mizota M, Katsuki Y, Mizuguchi K, Endo S, Miyata H, Kojima M, Kanehiro H et al. "Pharmacological studies of eicosapentaenoic acid ethylester (EPA E) on high cholesterol diet-fed rabbits," *Nippon Yakurigaku Zasshi*, 91:255-66 (Apr. 1988) (with English abstract).

(56)

## References Cited

## OTHER PUBLICATIONS

- Mizota M, Katsuki Y, Mizuguchi K, Endo S, Miyata H, Kojima M, Kanehiro H et al. "The effects of eicosapentaenoic acid ethylester (EPA E) on arterial thrombosis in rabbits and vascular lesions in rats," *Nippon Yakurigaku Zasshi*, 91:81-9 (Feb. 1988)(with English abstract).
- Mizuguchi K, Yano T, Kojima M, Tanaka Y, Ishibashi M, Masada A, Sato M et al. "Hypolipidemic effect of ethyl all-cis-5,8,11,14,17-eicosapentaenoate (EPA-E) in rats," *Jpn J Pharmacol.*, 59(3):307-12 (Jul. 1992).
- Mizuguchi, K., et al., "Ethyl all-cis-5,8,11,14,17-icosapentaenoate modifies the biochemical properties of rat very low-density lipoprotein." *European Journal of Pharmacology*, 231:221-227 (Apr. 28, 1993).
- Mizuguchi, K., et al., "Mechanism of the lipid-lowering effect of ethyl all-cis-5,8,11,14,17-icosapentaenoate." *European Journal of Pharmacology*, 231:121-127 (Jan. 1993).
- Mochida Press Release, Pharmaceutical Col., Ltd.: Conclusion of Distributorship Agreement Concerning Switch-OTC Drug for Hyperlipidemia Treatment, Epadel, (Apr. 30, 2009)(1 page).
- Mochida, Announcement, "Mochida Announces Completion of "JELIS" Major Clinical Trial for Epadel," Mar. 22, 2005 (2 pages).
- Mochida's Epadel Reduces Risk of Stroke Recurrence—New Results of JELIS Major Clinical Trial, JCNNetwork Newswire Nov. 13, 2006 (2 pages).
- Mora, S., et al., "LDL particle subclasses, LDL particle size, and carotid atherosclerosis in the Multi-Ethnic Study of Atherosclerosis (MESA)." *Atherosclerosis.*;192:211-217 (May 2007).
- Mori et al., "Differential Effects of Eicosapentaenoic Acid and Docosahexaenoic Acid on Vascular Reactivity of the Forearm Microcirculation in Hyperlipidemic, Overweight Men," *Circulation*, 102:1264-1269 (Sep. 12, 2000).
- Mori TA, Woodman RJ. "The independent effects of eicosapentaenoic acid and docosahexaenoic acid on cardiovascular risk factors in humans," *Curr Opin Clin Nutr Metab Care* 2006; 9:95-104 (Mar. 2006).
- Mori TA. Omega-3 fatty acids and blood pressure. *Cell Mol Biol*. Feb. 25, 2010;56(1):83-92.
- Mori, et al., "Purified Eicosapentaenoic and docosahexaenoic acids have differential effects on serum lipids and lipoproteins, LDL particle size, glucose, and insulin in mildly hyperlipidemic men," *Am J Clin Nutr* 71:1085-1094 (May 2000).
- Mori, T. et al., Effect of Eicosapentaenoic acid and docosahexaenoic acid on oxidative stress and inflammatory markers in treated-hypertensive type 2 diabetic subjects, *Free Radical Biology & Medicine*, vol. 35, No. 7, pp. 772-781 (Oct. 1, 2003).
- Mori, Trevor A., et al., "Docosahexaenoic Acid but Not Eicosapentaenoic Acid Lowers Ambulatory Blood Pressure and Heart Rate in Humans", *Hypertension*, 34(2):253-60 (Aug. 1999).
- Morita, I., et al., "Effects of purified eicosapentaenoic acid on arachidonic acid metabolism in cultured murine aortic smooth muscle cells, vessel walls and platelets." *Lipids* 18:42-490 (Jan. 1983).
- Morris M, Sacks F, Rosner B. Does fish oil lower blood pressure? A meta-analysis of controlled trials. *Circ.*, Aug. 1993;88:523-533.
- Morrow, JD, "Release of markedly increased quantities of prostaglandin D2 in vivo in humans following the administration of nicotinic acid", *Prostaglandins*, 38:263-274, (Aug. 1989).
- Morton, R.E., "Specificity of lipid transfer protein for molecular species of cholesteryl ester." *J Lipid Res.*, 27:523-529 (May 1986).
- Mosher LR et al., "Nicotinic Acid Side Effects and Toxicity: A review," *Am J Psychiat.*, 126: 1290-1296 (Mar. 1970).
- Mostad et al., "Effects of Marine N-3 Fatty Acid Supplementation on Lipoprotein Subclasses Measured by Nuclear Magnetic Resonance in Subjects with Type II Diabetes," *European Journ. Clin. Nutr.*, 62(3):419-429 (Mar. 2008/epub Feb. 27, 2007).
- Mostad, I.L., et al., "Effects of n-3 fatty acids in subjects with type 2 diabetes: reduction of insulin sensitivity and time-dependent alteration from carbohydrate to fat oxidation." *Am J Clin Nutr* 84:540-50 (Sep. 2006).
- Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, et al.; on behalf of the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2016 update: a report from the American Heart Association [published online ahead of print Dec. 16, 2015]. *Circulation*. doi: 10.1161/CIR.0000000000000350.
- Mozaffarian D, Geelen A, Brouwer I, Geleijnse J, Zock P, Katan M. Effect of Fish Oil on Heart Rate in Humans a Meta-Analysis of Randomized Controlled Trials. *Circ.Sep. 27, 2005/ epub Sep. 19, 2005; 112:1945-1952.*
- Mozaffarian D, Marchioli R, Macchia A, et al. Fish Oil and Postoperative Atrial Fibrillation the Omega-3 Fatty Acids for Prevention of Post-operative Atrial Fibrillation (OPERA) Randomized Trial. *JAMA*. Nov. 21, 2012;308(19):2001-11.
- Mozaffarian D, Psaty B, Rimm E, Lemaitre R, Burke G, Lyles M, Lefkowitz D, Siscovick D. Fish Intake and Risk of Incident Atrial Fibrillation. *Circ.*, Jul. 27, 2004/epub Jul. 19, 2004; 110:368-373.
- Mozaffarian et al., "Omega-3 fatty acids and cardiovascular disease: effects on risk factors, molecular pathways and clinical events," *J. Am. Coll. Cardiol.* (Nov. 8, 2011) 58(2):2047-2067.
- Mozaffarian, "JELIS, fish oil, and cardiac events," *www.thelancet.com* vol. 369, pp. 1062-1063 (Mar. 31, 2007).
- Mozaffarian, D., "Fish and n-3 fatty acids for the prevention of fatal coronary heart disease and sudden cardiac death." *Am J Clin Nutr*, 87:1991S-6S (Jun. 2008).
- Mozaffarian, D., et al., "Dietary fish and w-3 fatty acid consumption and heart rate variability in US adults." *Circulation*, 117:1130-1137 (Mar. 4, 2008/ epub Feb. 19, 2008).
- Murck et al., "Ethyl-EPA in Huntington disease—Potentially relevant mechanism of action," *Brain Research Bulletin*, 72:159-164 (2007) (available online Nov. 15, 2006).
- Murphy SA, Cannon CP, Blazing MA, et al. Reduction in total cardiovascular events with ezetimibe/simvastatin post-acute coronary syndrome. *J Am Coll Cardiol*. 67(4):353-361 (publication date Feb. 2, 2016; epublication date Jan. 25, 2016).
- Naba, H., et al., "Improving effect of ethyl eicosapentanoate on statin-induced rhabdomyolysis in Eisai hyperbilirubinemic rats." *Biochemical and Biophysical Research Communications*, 340:215-220 (Feb. 2006/epub Dec. 9, 2005).
- Nagakawa et al., Effect of [EPA] on the Platelet Aggregation and Composition of Fatty Acid in Man: A Double Blind Study, *Atherosclerosis* 47(1):71-75 (Apr. 1983).
- Naik H, Wu JT, Palmer R, McLean L. The effects of februxostat on the pharmacokinetic parameters of rosiglitazone, a CYP2C8 substrate. *Br. J. Clin. Pharmacol.* Jan. 13, 2012;74:327-335.
- Nakamura et al., Remnant lipoproteinemia is a risk factor for endothelial vasomotor dysfunction and coronary artery disease in metabolic syndrome, *Atherosclerosis*, vol. 181(2), pp. 321-327 (Aug. 2005/epub Feb. 16, 2005).
- Nakamura, et al., "Effects of Eicosapentaenoic Acids on Remnant-like Particles, Cholesterol Concentrations and Plasma Fatty Acid Composition in Patients with Diabetes Mellitus." *in vivo* 12: 311-314 (May/Jun. 1998).
- Nakamura, H., et al., "Evaluation of ethyl icosapentate in the treatment of hypercholesterolemia in kidney transplant recipients." *Transplantation Proceedings*, 30:3047-3048 (Nov. 1998).
- Nakamura, N. et al., "Joint effects of HMG-CoA reductase inhibitors and eicosapentaenoic acids on serum lipid profile and plasma fatty acid concentrations in patients with hyperlipidemia," *International Journal of Clinical and Laboratory Research*, Springer, Berlin, DE LNKD-DOI: 10.1007/S005990050057, vol. 29, No. 1, pp. 22-25 (1999).
- Nambi V, Bhatt DL. Primary prevention of atherosclerosis: Time to take a selfie? *J Am Coll Cardiol* 2017;70(24):2992-4 (publication date Dec. 19, 2017; epublication date Dec. 11, 2017).
- Nambi, V. et al., "Combination therapy with statins and omega-3 fatty acids." *Am J Cardiol* 98:34i-38i (Aug. 21, 2006/epub May 30, 2006).
- Nasa, et al., "Long-Term Supplementation With Eicosapentaenoic Acid Salvages Cardiomyocytes From Hypoxia/Reoxygenation-Induced Injury in Rats Fed With Fish-Oil-Deprived Diet," *Jpn. J. Pharmacol.* 77, 137-146 (Jun. 1998).

(56)

## References Cited

## OTHER PUBLICATIONS

National Kidney Foundation, "Glomerular Filtration Rate (GFR)," Jan. 30, 2017 (Jan. 30, 2017), retrieved on Jul. 30, 2018 from <https://web.archive.org/web/20170130183218/https://www.kidney.org/atoz/content/gfr>; entire document, especially p. 1 paragraph 1 and p. 3, paragraph 2.

National Kidney Foundation, "The Heart and Kidney Connection," Apr. 17, 2017 (Apr. 17, 2017), retrieved on Jul. 30, 2018 from <https://web.archive.org/web/2017041700416/https://www.kidney.org/atoz/content/heart-and-kidney-connection>; entire document, especially p. 2, paragraph 1.

Natsuno et al., "Clinical Effects of Eicosapentaenoic Acid on Type-2 Diabetes Effects on Serum Lipids, Pulse Wave Speed, and Ankle-Brachial Blood Pressure Index," *Diagnosis and Treatment* 93(12):133-137 (2005)(16 pages).

Nattel, S. et al., "Atrial remodeling and atrial fibrillation: Mechanisms and implications." *Circ Arrhythmia Electrophysiol*, 1:62-73 (Apr. 2008).

Needleman P, Raz A, Minkes MS, Ferrendelli JA, Sprecher H. Triene prostaglandins: prostacyclin and thromboxane biosynthesis and unique biological properties. *Proc Natl Acad Sci USA*. Feb. 1979;76:944-948.

Negre-Salvayre, A., et al., "Advanced lipid peroxidation end products in oxidative damage to proteins. Potential role in diseases and therapeutic prospects for the inhibitors." *British Journal of Pharmacology* 153:6-20 (Jan. 2008/epub Jul. 23, 2007).

Nelson et al. "Icosapent Ethyl for Treatment of Elevated Triglyceride Levels," *Annals of Pharmacotherapy*, 47(11):1517-1523 (Nov. 2013/epub Nov. 5, 2013).

Nelson JR, Wani O, May HT, Budoff M. Potential benefits of eicosapentaenoic acid on atherosclerotic plaques. *Vascul Pharmacol*. 91:1-9 (publication date Apr. 2017; epublication date Mar. 2, 2017).

Nelson, G.J., et al., "The Effect of Dietary Docosahexaenoic Acid on Plasma Lipoproteins and Tissue Fatty Acid Composition in Humans", *Lipids*, 32(11):1137-1146, (Nov. 1997).

Nemets, Boris, M.D., "Addition of Omega-3 Fatty Acid to Maintenance Medication Treatment for Recurrent Unipolar Depressive Disorder", *Am. J. Psychiatry*, 159(3):477-479, (Mar. 2002).

Nemoto et al., "Ethyl-eicosapentaenoic Acid Reduces Liver Lipids and Lowers Plasma Levels of Lipids in Mice Fed a High-Fat Diet, in vivo," 23:685-690 (Sep./Oct. 2009).

Nenseter, MS et al., "Effect of dietary supplementation with n-3 polyunsaturated fatty acids on physical properties and metabolism of low density lipoprotein in humans," *Arterioscler. Thromb. Vasc. Biol.*, 12:369-379 (Mar. 1992).

Nestel, et al., "The n-3 fatty acids eicosapentaenoic acid and docosahexaenoic acid increase systemic arterial compliance in humans," *Am J Clin Nutr.*, 76:326-30 (Aug. 2002).

Nestel, P.J., "Effects of N-3 fatty acids on lipid metabolism." *Ann Rev Nutr.*, 10:149-167 (1990).

Nichols GA, Philip S, Reynolds K, Granowitz CB, Fazio S. Increased cardiovascular risk in hypertriglyceridemic patients with statin-controlled LDL cholesterol. *J Clin Endocrinol Metab* 103(8):3019-27 (publication date Aug. 1, 2018; epublication date May 29, 2018).

Nichols GA, Philip S, Reynolds K, Granowitz CB, Fazio S. Increased residual cardiovascular risk in patients with diabetes and high vs. normal triglycerides despite statin-controlled LDL Cholesterol. *Diabetes Obes Metab* (publication date Sep. 17, 2018; epublication date Sep. 17, 2018).

Niemi M, Backman JT, Grantors M, Laitila J, Neuvonen M, Neuvonen PJ. Gemfibrozil considerably increases the plasma concentrations of rosiglitazone. *Diabetologia*. Oct. 2003/ Jul. 29, 2003); 46: 1319-1323.

Niemi M, Backman JT, Neuvonen PJ. Effects of trimethoprim and rifampin on the pharmacokinetics of the cytochrome P450 208 substrate rosiglitazone. *Clin. Pharmacol. Ther.*, Sep. 2004;76:239-249.

Nigon F, Lesnik P, Rouis M, Chapman MJ. Discrete subspecies of human low density lipoproteins are heterogeneous in their interaction with the cellular LDL receptor. *J. Lipid Res.*, Nov. 1991;32(11):1741-1753.

Nippon Rinsho, *Metabolic Syndrome 2nd Edition—Basics and New Clinical Findings*, Jan. 20, 2011, Special Issue 1 (vol. 69), pp. 503-506 (with English translation).

Nishikawa M. et al., "Effects of Eicosapentaenoic acid (EPA) on prostacyclin production in diabetics. GC/MS analysis of PG12 and PG13 levels" *Methods Find Exp Clin Pharmacol*. 19(6):429-33 (Jul./Aug. 1997).

Nobukata, H., et al., "Age-related changes in coagulation, fibrinolysis, and platelet aggregation in male WBN/Kob rats." *Thrombosis Research* 98: 507-516 (Jun. 15, 2000).

Nobukata, H., et al., "Long-term administration of highly purified eicosapentaenoic acid ethyl ester improves the dysfunction of vascular endothelial and smooth muscle cells in male WBN/Kob rats." *Metabolism*, 49(12): 1588-1591 (Dec. 2000).

Nobukata, H., et al., "Long-term administration of highly purified eicosapentaenoic acid ethyl ester prevents diabetes and abnormalities of blood coagulation in male WBN/Kob rats." *Metabolism*, 49(12): 912-919 (Jul. 2000).

Noguchi et al., "Chemoprevention of DMBA-induced mammary carcinogenesis in rats by low-dose EPA and DHA." *Br. J. Cancer* 75(3): 348-353 (1997).

Nomura et al., "The effects of pitavastatin, eicosapentaenoic acid and combined therapy on platelet-derived microparticles and adiponectin in hyperlipidemic, diabetic patients." *Platelets*, 20(1):16-22 (Feb. 2009).

Nomura S, Shouzu A, Omoto S, et al. Effects of eicosapentaenoic acid on endothelial cell-derived microparticles, angiopoietins and adiponectin in patients with type 2 diabetes. *J Atheroscler Thromb.*, Apr. 2009;16:83-90.

Nourooz-Zadeh, J., et al., "Urinary 8-epi-PGF2 $\alpha$  and its endogenous  $\beta$ -oxidation products (2,3-dinor and 2,3-dinor-5,6-dihydro) as biomarkers of total body oxidative stress." *Biochemical and Biophysical Research Communications* 330:731-736 (May 13, 2005).

Nozaki S. et al., "Effects of purified Eicosapentaenoic acid ethyl ester on plasma lipoproteins in primary hypercholesterolemia" *Int J Vitam Nutr Res*. 62(3):256-260 (1992).

Obata, et al., "Eicosapentaenoic acid inhibits prostaglandin D2 generation by inhibiting cyclo-oxygenase in cultured human mast cells", *Clin. & Experimental Allergy*, 29:1129-1135, (Aug. 1999).

O'Donnell, C.J., et al., "Leukocyte telomere length and carotid artery intimal medial thickness—the Framingham heart study." *Arteriosclerosis, Thrombosis, and Vascular Biology*.28:1165-1171 ( Jun. 2008/epub Apr. 3, 2008).

Oemar BS, Tschudi MR, Godoy N, Brovkovich V, Malinski T, Luscher TF. Reduced endothelial nitric oxide synthase expression and production in human atherosclerosis. *Circulation.*, Jun. 30, 1998;97:2494-2498.

Oh, Robert C et al., *Management of Hypertriglyceridemia*, American Family Physician, LNKD-PUBMED: 17508532, vol. 75, No. 9, pp. 1365-1371 (May 1, 2007).

Ohara Y, Peterson TE, Harrison DG. Hypercholesterolemia increases endothelial superoxide anion production. *J. Clin. Invest*. Jun. 1993;91:2546-2551.

Ohashi, *Journal of Clinical and Experimental Medicine*, Feb. 14, 2009, vol. 228, No. 7, pp. 795-805 (with English translation).

Okuda, Y. et al., Eicosapentaenoic acid enhances nitric oxide production by cultured human endothelial cells. *Biochem. Biophys. Res. Commun.* 232: 487-491 (Mar. 17, 1997).

Okuda, Y., et al., "Long-term effects of eicosapentaenoic acid on diabetic peripheral neuropathy and serum lipids in patients with type II diabetes mellitus." *Journal of Diabetes and Its Complications* 10:280-287 (Sep./Oct. 1996).

Okumura, T., et al., "Eicosapentaenoic acid improves endothelial function in hypertriglyceridemic subjects despite increased lipid oxidizability." *Am J Med Sci* 324(5):247-253 (Nov. 2002).

Oliw, E.H., et al., "Biosynthesis of prostaglandins from 17(18)epoxy-eicosatetraenoic acid, a cytochrome P-450 metabolite of eicosapentaenoic acid." *Biochimica et Biophysica Acta*, 1126, 261-268 (Jun. 26, 1992).

(56)

## References Cited

## OTHER PUBLICATIONS

- Olofsson et al., "Apolipoprotein B: a clinically important apolipoprotein which assembles atherogenic lipoproteins and promotes the development of atherosclerosis" *Journal of Internal Medicine*, 258: 395-410 (Nov. 2005).
- Omacor Summary of Product Characteristics, Pronova BioPharma Norge AS, Lysaker, Norway, Mar. 2015.
- Omacor® Prescribing Information (Omega-3-acid ethyl esters, capsules) (2004). (9 pages).
- Omacor®, Physicians' Desk Reference 2735 (60th ed. 2006)(3 pages).
- Ona, V.O., et al., *Nature*, vol. 399, Inhibition of caspase-1 slows disease progression in a mouse model of Huntington's disease, pp. 263-267 (May 20, 1999).
- Ooi EM, "Apolipoprotein C-III: Understanding an emerging cardiovascular risk factor", *Clin.Sci. (London)*, vol. 114, pp. 611-624 (May 2008).
- Opalinska et al., "Increasing Level of Prostate-Specific Antigen and Prostate Cancer Risk Factors Among 193 Men Examined in Screening Procedure," *Ann. Univ. Curie Sklowoska Med.*, 58(2):57-63 (Abstract Only)(2003)(2 pages).
- Origin Trial Investigators (The). n-3 fatty acids and cardiovascular outcomes in patients with dysglycemia. *N Engl J Med* Jul. 6, 2012/epub Jun. 11, 2012; 367:309-318.
- O'Riordan, "DHA and EPA have differential effects on LDL-cholesterol," May 24, 2011 [online][Retrieved on Aug. 21, 2015] Retrieved from website: <http://www.medscape.com/viewarticle/743305> (2 pages).
- Osher et al., "Omega-3 Eicosapentaenoic Acid in Bipolar Depression: Report of a Small Open-Label Study," *J. Clin. Psych.* 66:726-729 (Jun. 2005).
- Otvos et al., "Clinical Implications of Discordance Between LDL Cholesterol and LDL Particle Number," *J. Clin. Lipidol.* 5(2):105-113 (Mar.-Apr. 2011)(available online Mar. 1, 2011).
- Ou Z, Ou J, Ackerman AW, Oldham KT, Pritchard KA, Jr. L-4f, an apolipoprotein a-1 mimetic, restores nitric oxide and superoxide anion balance in low-density lipoprotein-treated endothelial cells. *Circulation*. Mar. 25, 2003;107:1520-1524.
- Ozaki M, Kawashima S, Yamashita T, Hirase T, Namiki M, Inoue N, Hirata K, Yasui H, Sakurai H, Yoshida Y, Masada M, Yokoyama M. Overexpression of endothelial nitric oxide synthase accelerates atherosclerotic lesion formation in apoe-deficient mice. *J. Clin. Invest.* Aug. 2002; 110:331-340.
- Ozawa, Akio, Nakamura E, Jinbo H, Fujita T, Hirai A, Terano T, Hamazaki T et al., "Determination of higher fatty acids in various lipid fractions of human plasma, platelets, and erythrocyte membrane using thin layer chromatography and gas chromatography," *Bunseki Kagaku*, 32:174-8 (1982) (with English abstract).
- Padgett et al., "Phylogenetic and immunological definition of four lipoylated proteins from *Novosphingobium aromaticivorans*, implications for primary biliary cirrhosis," *Journ. Autoimmunity* 24:209-219 (May 2005).
- Park JH, Park DI, Kim HJ, et al. Metabolic syndrome is associated with erosive esophagitis. *World J. Gastroenterol.* Sep. 14, 2008 (35): 5442-7.
- Park JY, Kim KA, Kang MH, Kim SL, Shin JG. Effect of rifampin on the pharmacokinetics of rosiglitazone in healthy subjects. *Clin. Pharmacol. Ther.*, Mar. 2004;75:157-162.
- Park, Y., et al., "Omega-3 fatty acid supplementation accelerates chylomicron triglyceride clearance." *J. Lipid Res.* 44:455-463 (Mar. 2003).
- Pase M, Grima N, Sarris J. Do long-chain n-3 fatty acids reduce arterial stiffness? A meta-analysis of randomized controlled trials. *Br J Nutr.*, Oct. 2011; 106:974-980.
- Patel et al., "Rosiglitazone monotherapy improves glycaemic control in patients with type 2 diabetes: a twelve-week, randomized, placebo-controlled study," *Diabetes, Obesity and Metabolism*, vol. 1, pp. 165-172 (May 1999).
- Paton, CM, Ntambi, JM., Biochemical and physiological function of stearoyl-CoA desaturase, *Am. J. Physiol. Endocrinol. Metab.* Jul. 2009/epub Dec. 9, 2008; 297:E28-E37.
- PCT/GB00/00164 International Search Report dated Oct. 20, 2000 (8 pages).
- PCT/US2011/062247 International Search Report and Written Opinion dated Jun. 14, 2012 (12 pages).
- PCT/US2013/020526 International Search Report dated Mar. 29, 2013 (2 pages).
- PCT/US2013/048241 International Search Report dated Dec. 13, 2013 (3 pages).
- PCT/US2013/048516 International Search Report dated Dec. 20, 2013 (3 pages).
- PCT/US2013/048559 International Search Report dated Dec. 13, 2013 (3 pages).
- PCT/US2013/068647 International Search Report and Written Opinion dated May 13, 2014 (18 pages).
- PCT/US2014/019454 International Search Report and Written Opinion dated Jun. 3, 2014 (12 pages).
- Pedersen RS, Damkier P, Brosen K. The effects of human CYP2C8 genotype and fluvoxamine on the pharmacokinetics of rosiglitazone in healthy subjects. *Br. J. Clin. Pharmacol.* Dec. 2006/epub Jul. 12, 2006; 62:682-689.
- Pedersen, T., et al., "Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S)", *The Lancet*, No. 19, vol. 344, 8934, p. 1383-1389 (Nov. 19, 1994).
- Peet et al., "A Dose-Ranging Study of the Effects of Ethyl-Eicosapentaenoate in Patients with Ongoing Depression Despite Apparently Adequate Treatment with Standard Drugs", *Arch. Gen. Psychiatry*, 59:913-919, (Oct. 2002).
- Peet, M., et al., Phospholipid Spectrum Disorder in Psychiatry pp. 1-19, (1999).
- Pejic et al., "Hypertriglyceridemia," *Journ. Amer. Board Fam. Med.*, vol. 19(3):310-316 (May/June. 2006).
- Pennathur S, Heinecke JW. Mechanisms for oxidative stress in diabetic cardiovascular disease. *Antioxid. Redox Signal.* Jul. 2007;9(7):955-969.
- Piccini, Monica, et al., *Genomics*, vol. 47, "FACL4, a new gene encoding long-chain acyl-CoA synthetase 4, is deleted in a family with Alport syndrome, elliptocytosis, and mental retardation," pp. 350-358 (Feb. 1998).
- Piche, "Tumor Necrosis Factor-Alpha, and Fibrinogen to Abdominal Adipose Tissue, Blood Pressure, and Cholesterol and Triglyceride Levels in Healthy Postmenopausal Women", *American Journal of Cardiology*, 2005, 96(1), 92-97.
- Pike, NB, "Flushing out the role of GPR109A (HM74V) in the clinical efficacy of nicotinic acid", *J. Clin. Invest.*, 115:3400-3403, (Dec. 2005).
- Plusepa® Product brochure "Super Critically Different from Other Omega-3 Fish Oil Supplements for Depression and ADHD," by Minami Nutrition (Apr. 2009, pp. 1-6).
- Pollin TI, Damcott CM, Shen H, et al. A null mutation in human APOC3 confers a favorable plasma lipid profile and apparent cardioprotection. *Science*. Dec. 12, 2008;322(5908):1702-1705.
- Pownall, H.J., et al., "Correlation of serum triglyceride and its reduction by ω-3 fatty acids with lipid transfer activity and the neutral lipid compositions of high-density and low-density lipoproteins." *Atherosclerosis* 143:285-297 (Apr. 1999).
- Press Release: Amarin Corporation Says Huntington's Disease Drug Failed in Trials, <http://www.fiercebiotech.com/node/6607/print> (Apr. 24, 2007) (Printed on Aug. 22, 2008)(2 pages).
- Pritchard KA, Ackerman AW, Ou J, Curtis M, Smalley DM, Fontana JT, Stemerman MB, Sessa WC. Native low-density lipoprotein induces endothelial nitric oxide synthase dysfunction: Role of heat shock protein 90 and caveolin-1. *Free Radic. Biol. Med.* Jul. 2002;33:52-62.
- Pritchard KA, Jr., Groszek L, Smalley DM, Sessa WC, Wu M, Villalon P, Wolin MS, Stemerman MB. Native low-density lipoprotein increases endothelial cell nitric oxide synthase generation of superoxide anion. *Circ. Res.* 1995;77:510-518.

(56)

## References Cited

## OTHER PUBLICATIONS

- Puri et al., "Reduction in Cerebral Atrophy Associated with Ethyl-eicosapentaenoic Acid Treatment in Patients with Huntington's Disease," *Journ. Int'l. Med. Research*, 36:896-905 (Oct. 1, 2008).
- Puri, B., et al., "Eicosapentaenoic Acid in Treatment-Resistant Depression Associated with Symptom Remission, Structural Brain Changes and Reduced Neuronal Phospholipid Turnover," *Int J Clinical Practice*, 55:560-563 (Oct. 2001).
- Puri, B., et al., *Archives of General Psychiatry*, No. 55, "Sustained remission of positive and negative symptoms of schizophrenia following treatment with eicosapentaenoic acid," pp. 188-189, (Feb. 1998).
- Puri, B.K., et al., "Ethyl-EPA in Huntington Disease: A Double-Blind, Randomized, Placebo-Controlled Trial", *Neurology*, 65:286-292, (Jul. 26, 2005).
- Qi, K., et al., "Omega-3 fatty acid containing diets decrease plasma triglyceride concentrations in mice by reducing endogenous triglyceride synthesis and enhancing the blood clearance of triglyceride-rich particles." *Clinical Nutrition* 27(8):424-430 (Jun. 2008/epub Mar. 24, 2008).
- Rader, *Lipid Disorders*, in Eric J. Topol (ed.) *Textbook of Cardiovascular Medicine* pp. 55-75 (2007).
- Rahimy M, Hallen B, Narang P. Effect of tolterodine on the anticoagulant actions and pharmacokinetics of single-dose warfarin in healthy volunteers. *Arzneimittelforschung* 2002 52 (12): 890-5.
- Raitt, M.H., et al., "Fish oil supplementation and risk of ventricular tachycardia and ventricular fibrillation in patients with implantable defibrillators—a randomized controlled trial." *JAMA*. 293(23):2884-2891 (Jun. 15, 2005).
- Rambjor, Gro S., et al., "Elcosapentaenoic Acid is Primarily Responsible for Hypotriglyceridemic Effect of Fish Oil in Humans", *Fatty Acids and Lipids from Cell Biology to Human Disease: Proceedings of the 2nd international Congress of the ISSFAL (International Society for the Study of Fatty Acids and Lipids, AOCS Press, 31:S-45-S-49, (Mar. 1, 1996).*
- Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease. The Scandinavian Simvastatin Survival Study, *Lancet*. 344: 1383-1389 (1994).
- Rao MN, Mullangi R, Katneni K, et al. Lack of effect of sucralfate on the absorption and pharmacokinetics of rosiglitazone. *J. Clin. Pharmacol.* Jun. 2002;42:670-675.
- Rauch B, Rudolf R, Schneider S, et al. OMEGA, a randomized, placebo-controlled trial to test the effect of highly purified omega-3 fatty acids on top of modern guideline-adjusted therapy after myocardial infarction. *Circulation*. Nov. 23, 2010 (epub Nov. 8, 2010); 122:2152-2159.
- Rees DD, Palmer RM, Moncada S. The role of endothelium-derived nitric oxide in the regulation of blood pressure. *Proc. Natl. Acad. Sci. USA*. May 1989;86:3375-3378.
- Reich, "Formulation and physical properties of soft capsules," *Pharmaceutical capsules*. (2004) Chapter 11:201-212.
- Reiffel, J.A., et al., "Antiarrhythmic effects of omega-3 fatty acids." *Am J Cardiol* 98:50i-60i (Aug. 21, 2006/epub May 26, 2006).
- Reiner Z, Catapano AL, De BG, et al. ESC/EAS Guidelines for the management of dyslipidaemias: the Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Eur. Heart J.* Jul. 2011(epub Jun. 28, 2011); 32:1769-1818.
- Richter, Werner O., "Hypertriglyceridämie: Ein klinischer Leitfaden," *Wissenschaftliche Verlagsgesellschaft mbH Stuttgart*, front page to p. V, pp. 2 to 55, 64 to 85, 90 to 97 (2008) (with English Summary).
- Ridker PM, Everett BM, Thuren T, et al. Antiinflammatory Therapy with canakinumab for atherosclerotic disease. *N Engl J Med* 377(12):1119-31 (publication date Sep. 21, 2017; epublication date Aug. 27, 2017).
- Ridker, "C-Reactive Protein : A Simple Test to Help Predict Risk of Heart Attack and Stroke", *Circulation: Journal of the American Heart Association*, Sep. 23, 2003, 108, e81-e85.
- Riediger, N.D., et al., "A systemic review of the roles of n-3 fatty acids in health and disease." *J Am Diet Assoc.* 109:668-679. (Apr. 2009).
- Rifai, "High-Sensitivity C-Reactive Protein: A Novel and Promising Marker of Coronary Heart Disease", *Clinical Chemistry*, Mar. 2001, 47(3), 403-411.
- Risé, P., et al., "Effects of simvastatin on the metabolism of polyunsaturated fatty acids and on glycerolipid, cholesterol, and de novo lipid synthesis in THP-1 cells." *J. Lipid Res.* 38:1299-1307 (Jul. 1997).
- Risk and Prevention Study Collaborative Group, Roncaglioni MC, Tombesi M, et al. n-3 fatty acids in patients with multiple cardiovascular risk factors. *N Engl J Med.*, May 9, 2013;368(19):1800-8.
- Rissanen et al., "Fish Oil-Derived Fatty Acids, Docosahexaenoic Acid and Docosapentaenoic Acid, and the Risk of Acute Coronary Events the Kuopio Ischaemic Heart Disease Risk Factor Study," *Circulation*. (Nov. 28, 2000)(102):2677-2679 doi:10.1161/01.CIR.102.22.2677.
- Rizzo M, Bemeis K. Low-density lipoprotein size and cardiovascular risk assessment. *Q. J. Med.* Jan. 2006; 99(1): 1-14.
- Roach, P.D., et al., "The effects of dietary fish oil on hepatic high density and low density lipoprotein receptor activities in the rat." *FEBS Lett.*, 222: 159-162 (Sep. 28, 1987).
- Robinson, J.G., et al., "Meta-analysis of the relationship between non-high-density lipoprotein cholesterol reduction and coronary heart risk." *J Am Coll Cardiol.*, 53: 316-322 (Jan. 27, 2009).
- Roche, H.M., et al., "Effect of long-chain n-3 polyunsaturated fatty acids on fasting and postprandial triacylglycerol metabolism." *Am J Clin Nutr* 71:232S-7S (Jan. 2000).
- Roche, H.M., et al., "Long-chain n-3 polyunsaturated fatty acids and triacylglycerol metabolism in the postprandial state." *Lipids* 34: S259-S265 (1999).
- Rodriguez, Y., et al., "Long-chain ω6 polyunsaturated fatty acids in erythrocyte phospholipids are associated with insulin resistance in non-obese type 2 diabetics." *Clinica Chimica Acta* 354:195-199 (Apr. 2005).
- Roe MT, Armstrong PW, Fox KAA, et at; TRILOGY ACS investigators. Prasugrel versus clopidogrel for acute coronary syndromes without revascularization. *N Engl J Med.* 367(14):1297-1309 (publication date Oct. 4, 2012; epublication Aug. 25, 2012).
- Rogers, P. J., "No effect of n-3 long-chain polyunsaturated fatty acid (EPA and DHA) supplementation on depressed mood and cognitive function: a randomised controlled trial" *British Journal of Nutrition*, 99:421-431, (Feb. 2008/epub Oct. 24, 2007).
- Rost KL, Roots I. Nonlinear kinetics after high-dose omeprazole caused by saturation of genetically variable CYP2C19. *Hepatology* Jun. 23, 1996 (6): 1491-7.
- Rubins, HB, et al., "Distribution of lipids in 8,500 men with coronary artery disease: Department of Veterans Affairs HDL Intervention Trial Study Group," *Am. J. Cardiol*, 75:1196-1201, (Jun. 15, 1995).
- Rubins, HB, et al., "Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol: Veterans Affairs HDL-C Intervention Trial Study Group", *N. Eng. J. Med.*, 341:410-418, (Aug. 5, 1999).
- Ruiz-Narváez, E.A., et al., "Abdominal obesity and hyperglycemia mask the effect of a common APOC3 haplotype on the risk of myocardial infarction." *Am J Clin Nutr* 87:1932-8 (Jun. 2008).
- Ruocco MJ, Shipley GG. Interaction of cholesterol with galactocerebroside and galactocerebroside phosphatidylcholine bilayer membranes. *Biophys. J.* Dec. 1984; 46:695-707.
- Rupp, "Omega-3-Fettsäuren in der Sekundärprävention nach Myokardinfarkt," *Clin. Res. Cardiol.*, vol. 95:Suppl. 6, Vi/12/-V1-16 (2006)(with English summary).
- Rustan, A.C., et al., "Eicosapentaenoic acid inhibits cholesterol esterification in cultured parenchymal cells and isolated microsomes from rat liver." *J. Bio. Chem.* 263(17):8126-32 (Jun. 15, 1988).
- Rustan, A.C., et al., "Eicosapentaenoic acid reduces hepatic synthesis and secretion of triacylglycerol by decreasing the activity of acyl-coenzyme A:1,2-diacylglycerol acyltransferase." *J. Lipid Res.* 29:1417-1426 (Nov. 1988).
- Rustan, A.C., et al., "Postprandial decrease in plasma unesterified fatty acids during n-3 fatty acid feeding is not caused by accumu-



(56)

## References Cited

## OTHER PUBLICATIONS

- lation of fatty acids in adipose tissue." *Biochimica et Biophysica Acta* 1390:245-25 (Feb. 23, 1998).
- Ryan, A.M., et al., "Enteral nutrition enriched with eicosapentaenoic acid (EPA) preserves lean body mass following esophageal cancer surgery: results of a double-blinded randomized controlled trial." *Ann Surg* 249:355-363 (Mar. 2009).
- Ryan, A.S., et al., "Clinical overview of algal-docosahexaenoic acid: effects on triglyceride levels and other cardiovascular risk factors." *Am J Ther.*, 16:183-192 Mar./Apr. 2009
- Sacks, Frank M., "The apolipoprotein story," *Atherosclerosis Supplements*, 23-27 (Aug. 2006/epub Jul. 5, 2006).
- Saito et al., "Effects of EPA on coronary artery disease in hypercholesterolemic patients with multiple risk factors: Sub-analysis of primary prevention cases from the Japan EPA Lipid Intervention Study (JELIS)," *Atherosclerosis*, 200:135-140 (Sep. 2008/epub Jun. 19, 2008).
- Saito et al., "Results of Clinical Usage of Improved Formulation (MND-21S) Epadel Capsule 300 with Respect to Hyperlipidemia," 26(12) *Jpn. Pharmacol. Ther.* 2047-62 (1998) (with English abstract).
- Saito, J., et al., "Mechanisms of enhanced production of PGI<sub>2</sub> in cultured rat vascular smooth muscle cells enriched with eicosapentaenoic acid." *Atherosclerosis* 131: 219-228 (Jun. 1997).
- Sampath H, Ntambi JM., Role of stearoyl-CoA desaturase in human metabolic disease, *Future Lipidol.* 2008;3:2,163-73.
- Sampath H, Ntambi JM., The Role of stearoyl-CoA desaturase in obesity, insulin resistance, and inflammation, *Ann. NY. Acad. Sci.* Dec. 2011; 1243:4 7-53.
- Samuels, Martin A., M. D., et al., "Huntington's Disease", *Office Practice of Neurology*, (122):654-655, (1996).
- Sanders, A. Hinds and C.C. Pereira, "Influence of n-3 fatty acids on blood lipids in normal subjects" *Journal of Internal Medicine.* 225:99-104,(1989).
- Sanders, et al., "Influence of an algal triacylglycerol containing docosahexaenoic acid (22:6n-3) and docosapentaenoic acid (22:5n-6) on cardiovascular risk factors in healthy men and women," *British Journal of Nutrition*, 95, 525-531 (Mar. 2006).
- Sanders, T.A., et al., "Effect of varying the ratio of n-6 to n-3 fatty acids by increasing the dietary intake of  $\alpha$ -linolenic acid, eicosapentaenoic and docosahexaenoic acid, or both on fibrinogen and clotting factors VII and XII in persons aged 45-70 y: the OPTILIP Study." *Am J Clin Nutr* 84:513-22 (Sep. 2006).
- Sanders, T.A., et al., "Triglyceride-lowering effect of marine polyunsaturates in patients with hypertriglyceridemia." *Arterioscler. Thromb. Vasc. Biol.* 5:459-465 (Sep./Oct. 1985).
- Sarwar N, Danesh J, Eiriksdottir G, et al. Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation* 115:450-458, Jan. 30, 2007/epub Dec. 26, 2006.
- Sasaki J, Miwa T, Odawara M. Administration of highly purified eicosapentaenoic acid to stain-treated diabetic patients further improves vascular function. *Endocrine J.* Jan. 27, 2012; 59(4):297-304.
- Sasaki J, Yokoyama M, Matsuzaki M, et al. Relationship between coronary artery disease and non-HDL-C, and effect of highly purified EPA on the risk of coronary artery disease in hypercholesterolemic patients treated with statins: sub-analysis of the Japan EPA Lipid Intervention Study (JELIS). *J. Atheroscler. Thromb.* Dec. 17, 2012;19:194-204.
- Sasaki, Y.F., et al., "Bio-antitumor effects of unsaturated fatty acids included in fish oil—docosahexaenoic acid, docosapentaenoic acid, and eicosapentaenoic acid—in cultured Chinese hamster cells." *Mutation Research*, 320: 9-22 (Jan. 1994).
- Sato et al., "General Pharmacological Studies on 5 8 11 14 17 Eicosapentaenoic Acid Ethyl Ester EPA-E", *Folia Pharmacol JPN*, 94 (1), 35-47. cited by other (Jul. 1989) (with English abstract).
- Sato, "Effects of Highly Purified Ethyl All-cis-5,8,11,14,17-icosapentaenoate (EPA-E) on Rabbit Platelets," *Biol. Pharm. Bull.*, 16(4)362-367 (Apr. 1993).
- Satoh et al., "Highly purified eicosapentaenoic acid reduces cardiovascular index in association with decreased serum amyloid A-LDL in metabolic syndrome," *Hypertension Research* (Nov. 2009/epub Sep. 18, 2009) (32)1004-1008.
- Satoh, N., et al., "Purified eicosapentaenoic acid reduces small dense LDL, remnant lipoprotein particles, and C-reactive protein in metabolic syndrome." *Diabetes Care*, 30(1): 144-146 (Jan. 2007).
- Satoh-Asahara N, Shimatsu A, Sasaki Y, Nakaoka H, Himeno A, Tochiya M, Kono S, Takaya T, Ono K, Wada H, Suganami T, Hasegawa K, Ogawa Y., "Highly purified eicosapentaenoic acid increases interleukin-10 levels of peripheral blood monocytes in obese patients with dyslipidemia." *Diabetes Care.* Dec. 2012/epub Aug. 21, 2012; 35(12):2631-2639.
- Schaefer, E.J., et al., "Effects of eicosapentaenoic acid, docosahexaenoic acid, and olive oil on cardiovascular disease risk factors [abstract 20007]." *Circulation*, 122:A20007 (2010) (Abstract only).
- Schectman, G. & Hiatt, J., "Drug therapy for hypercholesterolemia in patients with cardiovascular disease: factors limiting achievement of lipid goals", *Am. J. Med.*, 100:197-204, (Feb. 1996).
- Schectman, G., et al., "Dietary fish oil decreases low-density-lipoprotein clearance in nonhuman primates." *Am J Clin Nutr.*, 64:215-221 (Aug. 1996).
- Schectman, G., et al., "Heterogeneity of Low Density Lipoprotein Responses to Fish-Oil Supplementation in Hypertriglyceridemic Subjects." *Arterioscler. Thromb. Vasc. Biol.* 9:345-354 (May/Jun. 1989).
- Schmidt, E.B., et al., "Lipoprotein-associated phospholipase A2 concentrations in plasma are associated with the extent of coronary artery disease and correlate to adipose tissue levels of marine n-3 fatty acids." *Atherosclerosis* 196: 420-424 (Jan. 2008).
- Schmitz PG, McCloud LK, Reikes ST, et al. Prophylaxis of hemodialysis graft thrombosis with fish oil: double-blind, randomized, prospective trial. *J. Am. Soc. Nephrol.* Jan. 13, 2002 (1): 184-90.
- Schmitz, G., et al., "The opposing effects of n-3 and n-6 fatty acids." *Progress in Lipid Research*, 47:147-155 (Mar. 2008/epub Dec. 27, 2007).
- Schreiner et al., "Lipoprotein[a] as a Risk Factor for Preclinical Atherosclerosis," *13 Atherosclerosis, Thrombosis & Vascular Biology* 6: 826-833 (1993).
- Schuirman, D.J. A comparison of the two one-sided tests procedure and the power approach for assessing the equivalence of average bioavailability. *J. Pharmacokinetic. Biopharm.* 15(6), 657-680 (Dec. 1987).
- Schunkert H, König IR, Kathiresan S, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet.* Mar. 6, 2011;43(4):333-8.
- Schwartz GG, Bessac L, Berdan LG, et al. Effect of alirocumab, a monoclonal antibody to PCSK9, on long-term cardiovascular outcomes following acute coronary syndromes: rationale and design of the ODYSSEY outcomes trial. *Am Heart J* 168(5):682-9 (publication date Nov. 2014, epublication date Aug. 7, 2017).
- Schwarz, S., et al., "Lycopene inhibits disease progression in patients with benign prostate hyperplasia." *J. Nutr.* 138: 49-53 (Jan. 2008).
- Schwellenbach et al., "The Triglyceride-Lowering Effects of a Modest Dose of Docosahexaenoic Acid Alone Versus in Combination with Low Dose Eicosapentaenoic Acid in Patients with Coronary Artery Disease and Elevated Triglycerides." *J. Am. Coll. Nutr.* 25(6):480-485 (Dec. 2006).
- Segrest et al., Structure of Apolipoprotein B-100 in Low Density Lipoproteins, *J. Lipid Res.* 42(9):1346-1367 (Sep. 2001).
- Self-Medlin Y, Byun J, Jacob RF, Mizuno Y, Mason RP. Glucose promotes membrane cholesterol crystalline domain formation by lipid peroxidation. *Biochim. Biophys. Acta.* Jun. 2009/epub Apr. 17, 2009; 1788(6): 1398-1403.
- Serhan C, Chiang N, Van Dyke T. Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol.* May 2008; 8:3449-361.
- Serhan, C.N., et al., "Resolvins: A family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals." *J. Exp. Med.* 196:1025-1037 (Oct. 21, 2002).
- Sevanian A, F. Ursini Lipid peroxidation in membranes and low-density lipoproteins: similarities and differences. *Free Radic. Biol. Med.*, Aug. 2000;29(3-4):306-311.

(56)

## References Cited

## OTHER PUBLICATIONS

- Shah, S. et al., "Eicosapentaenoic Acid (EPA) as an Adjunct in the Treatment of Schizophrenia", *Schizophrenia Research*, vol. 29, No. 1/02 (1998).
- Shan, Z., et al., "A combination study of spin-trapping, LC/ESR and LC/MS on carbon-centred radicals formed from lipoxygenase-catalysed peroxidation of eicosapentaenoic acid." *Free Radical Research*, 43(1):13-27 (Jan. 2009).
- Shearer et al., "Red Blood Cell Fatty Acid Patterns and Acute Coronary Syndrome," *PLoS ONE* 4(5): e5444, publ. May 6, 2009 (doi:10.1371/journal.pone.0005444).
- Shimizu et al., "Effects of Highly Purified Eicosapentaenoic Acid on Erythrocyte Fatty Acid Composition and Leukocyte and Colonic Mucosa Leukotriene B4 Production in Children with Ulcerative Colitis," *J. Pediatr. Gastroenterol. Nutr.*, vol. 37, No. 5, pp. 581-585 (Nov. 2003).
- Shimizu, H., et al., "Long-term effect of eicosapentaenoic acid ethyl (EPA-E) on albuminuria of non-insulin dependent diabetic patients." *Diabetes Research and Clinical Practice* 28: 35-40 (Apr. 1995).
- Shimokawa H, Flavahan NA, Vanhoutte PM. Loss of endothelial pertussis toxin-sensitive G protein function in atherosclerotic porcine coronary arteries. *Circulation*. Feb. 1991;83:652-660.
- Shinozaki K. et al., "The long-term effect of Eicosapentaenoic acid on serum levels of lipoprotein (a) and lipids in patients with vascular disease" *J Atheroscler Thromb.* 2(2):207-9 (1996).
- Shishehbor MH, Brenna ML, Aviles RJ, Fu X, Penn MS, Sprecher DL, Hazen SL. Statins promote potent systemic antioxidant effects through specific inflammatory pathways. *Circulation*. Jul. 29, 2003;108(4):426-431.
- Sicherer et al., "Prevalence of seafood allergy in the United States determined by a random telephone survey," *J. Allergy Clin. Immunol.*, 114(1):159-165 (Jul. 2004).
- Sierra, S., et al., "Dietary eicosapentaenoic acid and docosahexaenoic acid equally incorporate as decosahexaenoic acid but differ in inflammatory effects." *Nutrition* 24:245-254 (Mar. 2008).
- Silvers, Karen M., et al., "Randomised double-blind placebo-controlled trial of fish oil in the treatment of depression", *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 72:211-218, (Mar. 2005).
- Simoens, C.M., et al., "Inclusion of 10% fish oil in mixed medium-chain triacylglycerol-long chain triacylglycerol emulsions increases plasma triacylglycerol clearance and induces rapid eicosapentaenoic acid (20:5n-3) incorporation into blood cell phospholipids." *Am J Clin Nutr* 88: 282-8 (Aug. 2008).
- Simon, Joel A., et al., "Serum Fatty Acids and the Risk of Coronary Heart Disease", *American Journal of Epidemiology*, 142(5):469-476, (Sep. 1, 1995).
- Simopoulos, The Importance of the Omega-6/Omega-3 Fatty Acid Ratio in Cardiovascular Disease and Other Chronic Diseases, *Exp. Biol. Med.*, 233:674-688 (Jun. 1, 2008)(available online Jun. 1, 2008).
- Simopoulos, "Omega-3 fatty acids in health and disease and in growth and development," *Am. J. Clin. Nutr.* 54:438-63 (Sep. 1991).
- Singer, Peter, "Fluvastatin plus fish oil are more effective on cardiovascular risk factors than fluvastatin alone," Letter to the Editor, *Prostaglandins, Leukotrienes and Essential Fatty Acids*, vol. 72, pp. 379-380 (May 2005).
- Singh, R.B., et al., "Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival—4." *Cardiovascular Drugs and Therapy* 11:485-491 (Jul. 1997).
- Sirtori, C.R., et al., "One-year treatment with ethyl esters of n-3 fatty acids in patients with hypertriglyceridemia and glucose intolerance—Reduced triglyceridemia, total cholesterol and increased HDL-C." *Atherosclerosis* 137: 419-427 (Apr. 1998).
- Skinner JS, Cooper A, & Feder GS and on behalf of the Guideline Development Group. "Secondary prevention for patients following a myocardial infarction; summary of NICE guidance," *Heart*, 93:862-864 (Jul. 2007).
- Slides for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, (158 pages).
- Smith et al., Pharmacokinetics and Pharmacodynamics of Epoetin Delta in Two Studies in Health Volunteers and Two Studies in Patients with Chronic Kidney Disease, *Clinical Therapeutics*/vol. 29, pp. 1368-1380 (Jul. 2007).
- Sniderman A, Kwiterovich PO. Update on the detection and treatment of atherogenic low-density lipoproteins. *Curr. Opin. Endocrinol. Diabetes Obes.* Apr. 20, 2013;20:140-147.
- Sohma, R., et al., "Protective effect of n-3 polyunsaturated fatty acid on primary culture of rat hepatocytes without glycemic alterations." *Journal of Gastroenterology and Hepatology* 22: 1965-1970 (Nov. 2007).
- Spector, A.A. "Arachidonic acid cytochrome P450 epoxygenase pathway." *Journal of Lipid Research*, 50: S52-S56 (2009) (published online on Oct. 23, 2008.).
- Spector, A.A., et al., "Epoxyeicosatrienoic acids (EETs): metabolism and biochemical function." *Progress in Lipid Research* 43: 55-90 (Jan. 2004).
- Springer, T.A., "Traffic signals for lymphocyte recirculation and leukocyte emigration: The multistep paradigm." *Cell*, 76: 301-314 (Jan. 28, 1994).
- Squires, RW, et al., "Low-dose, time release nicotinic acid: effects in selected patients with low concentrations of high density lipoprotein cholesterol", *Mayo Clinic Proc.*, 67:855-860, (Sep. 1992).
- Srinivas, et al., "Controlled release of lysozyme from succinylated gelatin microspheres," *J. Biomater. Sci., Polymer Ed.*, vol. 12(2):137-148 (2001).
- Stalenhoef, A.F.H., et al., "The effect of concentrated n-3 fatty acids versus gemfibrozil on plasma lipoproteins, low density lipoprotein heterogeneity and oxidizability in patients with hypertriglyceridemia." *Atherosclerosis* 153: 129-138 (Nov. 2000).
- Stampfer MJ, Krauss RM, Ma J, et al. A prospective study of triglyceride level, low density lipoprotein particle diameter, and risk of myocardial infarction. *JAMA*. Sep. 1996;276:882-888.
- Stancu et al., "Statins: Mechanism of Action and Effects," *Journal of Cellular and Molecular Medicine* (Oct.-Dec. 2001), 5(4), 378-387.
- Stark, K.D. & Holub, B.J., Differential eicosapentaenoic acid elevations and altered cardiovascular disease risk factor responses after supplementation with docosahexaenoic acid in postmenopausal women receiving and not receiving hormone replacement therapy, *Am. J. Clin. Nutr.*, vol. 79, pp. 765-773 (May 2004).
- Stark, K.D., "The percentage of n-3 highly unsaturated fatty acids in total HUFA as a biomarker for omega-3 fatty acid status in tissues." *Lipids* 43:45-53 (Jan. 2008/epub Nov. 6, 2007).
- Stark, K.D., et al., "Effect of a fish-oil concentrate on serum lipids in postmenopausal women receiving and not receiving hormone replacement therapy in a placebo-controlled, double-blind trial." *Am J Clin Nutr* 72:389-94 (Aug. 2000).
- Steg PG, Bhatt DL, Wilson PWF, et al; REACH Registry Investigators. One-year cardiovascular event rates in outpatients with atherothrombosis. *JAMA*. 297(11):1197-1206 (publication date May 21, 2007).
- Stein et al., "Effect of Statin Therapy on Remnant Lipoprotein Cholesterol Levels in Patients with Combined Hyperlipidemia," *Arteriosclerosis, Thrombosis and Vascular Biology*, vol. 21, pp. 2026-2031 (Dec. 1, 2001).
- Steinberg D, Witztum JL. Is the oxidative modification hypothesis relevant to human atherosclerosis? Do the antioxidant trials conducted to date refute the hypothesis? *Circulation*. Apr. 30, 2002;105:2107-2111.
- Steinberg D. Lewis A. Conner Memorial Lecture: Oxidative modification of LDL and atherogenesis. *Circulation*. Feb. 18, 1997;95(4):1062-1071.
- Stepp DW, Ou J, Ackerman AW, Welak S, Klick D, Pritchard KA, Jr. Native ldl and minimally oxidized ldl differentially regulate superoxide anion in vascular endothelium in situ. *Am. J. Physiol.* Aug. 2002;283:H750-H759.
- Sternbach "The Glasgow Coma Scale." *The Journal of Emergency Medicine*, 19(1):67-71 (Feb. 8, 2000).

(56)

## References Cited

## OTHER PUBLICATIONS

- Stielow et al., "Novel Nox Inhibitor of oxLDL-Induced Reactive Oxygen Species Formation in Human Endothelial Cells," *Biochem. Biophys. Res. Comm.*, 344:200-205 (May 26, 2006/epub Mar. 26, 2006).
- Stiles, FDA approves EPA-only omega-3 PUFA capsule for high TG, Jul. 26, 2012, <http://www.medscape.com/viewarticle/791268>, accessed Dec. 17, 2014 (1 page).
- Stitzel N, Stirrups K, Masca N, et al. Supplement to: Coding variation in ANGPTL4, LPL, and SVEP1 and the risk of coronary disease. *N Engl J Med.* DOI: 10.1056/NEJMoa1507652; Mar. 24, 2016/epub Mar. 2, 2016.
- Stojancevic et al., "The impact of farnesoid X receptor activation on intestinal permeability in inflammatory bowel disease," *Can. J Gastroenterol.* 26(9):631-637 (Sep. 2012).
- Stoll, Andrew L. et al., "Omega 3 Fatty Acids in Bipolar Disorder", *Arch. Gen. Psychiatry*, 56:407-412, (May 1999).
- Stone NJ, Robinson J, Lichtenstein AH, et al. ACC/AHA Prevention Guideline: 2013 ACC/AHA Guideline on the Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. erratum in *Circulation.* Jun. 24, 2014;129:S46-S48.
- Su, Kuan-Pin, et al., "Omega-3 Fatty Acids in Major Depressive Disorder a Preliminary Double-Blind, Placebo-Controlled Trial", *European Neuropsychopharmacology*, 13:267-271, (Aug. 2003).
- Sugiyama et al., "A Comparison of the Hypotensive Effects of Eicosapentaenoic Acid Ethyl (EPA) on Three Diseases (Occluded Arteriosclerosis, Hyperlipidemia, and These Two Diseases Combined) P2-504 Abstract," Annual Meeting of the Japanese Society of Pharmaceutical Health Care and Sciences 20:473 (Nov. 2010) (with English translation)(3 pages).
- Sugiyama, E., et al., "Eicosapentaenoic acid lowers plasma and liver cholesterol levels in the presence of peroxisome proliferators-activated receptor alpha." *Life Sciences*, 83:19-28 (Jul. 4, 2008/epub May 1, 2008).
- Superko et al., "Lipid Management to Reduce Cardiovascular Risk: A New Strategy is Required," *Circulation*, 117:560-568 (Jan. 29, 2008).
- Surette, M.E., et al., "Dependence on dietary cholesterol for n-3 polyunsaturated fatty acid induced changes in plasma cholesterol in the Syrian hamster." *J Lipid Res.*, 33:263-271 (Feb. 1992).
- Surette, M.E., et al., "Evidence for mechanisms of the hypotriglyceridemic effect of n-3 polyunsaturated fatty acids." *Biochimica et Biophysica Acta*, 1126: 199-205 (Jun. 22, 1992).
- Tagawa H, Shimokawa H, Tagawa T, et al. Long-term treatment with eicosapentaenoic acid augments both nitric oxide-mediated and non-nitric oxide-mediated endothelium-dependent forearm vasodilatation in patients with coronary artery disease. *J Cardiovasc Pharmacol* 33(4):633-40, Apr. 1999.
- Takaki A, Umemoto S, Ono K, Seki K, Ryoke T, Fujii A, Itagaki T, Harada M, Tanaka M, Yonezawa T, Ogawa H, Matsuzaki M. Add-on therapy of epa reduces oxidative stress and inhibits the progression of aortic stiffness in patients with coronary artery disease and statin therapy: A randomized controlled study. *J. Atheroscler. Thromb.* Jun. 23, 2011;18:857-866.
- Takaku et al., Study on the Efficacy and Safety of Ethyl Icosapentate (MND-21) in Treatment of Hyperlipidemia Based on a Long-Term Administration Test, 7 *J. Clin. Ther. Med.* 191 (1991) (with English Translation)(27 pages).
- Talayero BG, Sacks FM. The role of triglycerides in atherosclerosis. *Curr. Cardiol. Rep.* 2011;13:544-552.
- Tamura, et al., "Study of the Clinical Usefulness of Ethyl Icosapentate (MND-21) in Long-Term Treatment of Hyperlipaemic Patients." *J Clin Thera & Medicines*, 7:1817-1834 (1991).
- Tanaka et al., "Genome-Wide Association Study of Plasma Polyunsaturated Fatty Acids in the InCHIANTI Study." *PLoS Genetics* 5(1):1-8 (Jan. 2009).
- Tanaka et al., "Suppression of prostaglandin synthesis by arachidonic acid or eicosapentaenoic acid in a macrophage-like cell line, RAW 264.7, treated with LPS," *Biol. Pharm. Bull.*, 22(10):1057-7 (Oct. 1999).
- Tanaka et al., "Administration of high dose eicosapentaenoic acid enhances anti-inflammatory properties of high-density lipoprotein in Japanese patients with dyslipidemia," *Atherosclerosis*, 237(2):577-83 (Dec. 2014).
- Tanaka et al., "Eicosapentaenoic Acid-Enriched High-Density Lipoproteins Exhibit Anti-Atherogenic Properties," *Circ. J.*, doi: 10.1253/circj.CJ-17/0294. [Epub ahead of print] (Jun. 23, 2017)(6 pages).
- Tanaka, K.T., et al., "Reduction in the recurrence of stroke by eicosapentaenoic acid for hypercholesterolemic patients—Subanalysis of the JELIS trial." *Stroke*, 39(7):2052-8 (Jul. 2008/epub May 1, 2008).
- Tatarczyk, et al., "Analysis of long-chain  $\omega$ -3 fatty acid content in fish-oil supplements," *Wien Klin Wochenschr*, 119/13-14: 417-422 (2007).
- Tatsuno et al., Efficacy and safety of TAK-085 compared with eicosapentaenoic acid in Japanese subjects with hypertriglyceridemia undergoing lifestyle modification: The omega-3 fatty acids randomized double-blind (ORL) study, *J. Clin. Lipid*; vol. 7(6), pp. 615-625 (Sep. 12, 2013).
- Taylor et al., "Fish allergy: fish and products thereof," *Journal Food Science* (2004) 69.8 R175-R180.
- Taylor, A.J., et al., "Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol (ARBITER) 2: a double-blind, placebo-controlled study of extended-release niacin on atherosclerosis progression in secondary prevention patients treated with statins", *Circulation*, 110:3512-3517, (Dec. 7, 2004/epub Nov. 10, 2004).
- Tedgui, A., et al., "Anti-inflammatory mechanisms in the vascular wall." *Circ. Res.* 88:877-887 (May 11, 2001).
- Teissier E, Nohara A, Chinetti G, Paumelle R, Cariou B, Fruchart JC, Brandes RP, Shah A, Staels B. Peroxisome proliferator-activated receptor alpha induces NADPH oxidase activity in macrophages, leading to the generation of LDL with PPAR-alpha activation properties. *Circ. Res.* Dec. 10, 2004/epub Nov. 11, 2004;95(12):1174-1182.
- Teramoto T, Sasaki J, Ishibashi S, et al. Diagnosis of atherosclerosis. Executive Summary of the Japan Atherosclerosis Society (JAS) Guidelines for the Diagnosis and Prevention of Atherosclerotic Cardiovascular Diseases in Japan—2012 Version. *J Atheroscler Thromb.* 2014;21(4):296-8. Electronic publication Dec. 10, 2013.
- Terano, et al., "Effect of Oral Administration of Highly Purified Eicosapentaenoic Acid on Platelet Function, Blood Viscosity and Red Cell Deformability in Healthy Human Subjects," *Atherosclerosis*, 46, 321-331 (Mar. 1983).
- The TG and HDL Working Group of the Exome Sequencing Project, National Heart, Lung, and Blood Institute. Loss-of-function mutations in APOC3, triglycerides, and coronary disease. *N Engl J Med.* Jul. 3, 2014/epub Jun. 18, 2014; 371(1):22-31.
- Theilla, M., et al., "A diet enriched in eicosapentaenoic acid, gamma-linolenic acid and antioxidants in the prevention of new pressure ulcer formation in critically ill patients with acute lung injury: A randomized, prospective, controlled study." *Clinical Nutrition* 26: 752-757 (Dec. 2007/epub Oct. 22, 2007).
- Theobald et al., "LDL Cholesterol-Raising Effect of Low-Dose Docosahexaenoic Acid in Middle-Aged Men and Women," *Am. J. Clin. Nutr.* 79:558-63 (Apr. 2004).
- Thies, F., et al., "Association of n-3 polyunsaturated fatty acids with stability of atherosclerotic plaques: a randomised controlled trial." *Lancet* 361: 477-85 (Feb. 8, 2003).
- Thies, F., et al., "Dietary supplementation with eicosapentaenoic acid, but not with other long-chain n-3 or n-6 polyunsaturated fatty acids, decreases natural killer cell activity in healthy subjects aged >55 y." *Am J Clin Nutr* 73:539-48 (Mar. 2001).
- Third Report of the NCEP Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report, NIH Publication No. 02-5215 Sep. 2002 (220 pages in three parts).
- Thomas et al., "Renal Failure—Measuring the Glomerular Filtration Rate," *Dtsch Arztebl Int.*, Dec. 18, 2009, 106(51-52); 849-54.

(56)

## References Cited

## OTHER PUBLICATIONS

- Thorwest M, Balling E, Kristensen SD, et al. Dietary fish oil reduces microvascular thrombosis in a porcine experimental model. *Thromb. Res.* Jul. 2000, 99(2): 203-8.
- Thygesen K., Alpert J., Jaffe A., et al. Third Universal Definition of Myocardial Infarction. *J Am Coll Cardiol.*, Oct. 16, 2012/epub Sep. 5, 2012; 60(16):1581-1598.
- Tilg H, Moschen AR. Inflammatory Mechanisms in the Regulation of Insulin Resistance. *Mol. Med.*, Mar./Apr. 2008;14(3-4):222-231.
- Tirosh et al., "Changes in Triglyceride Levels and Risk for Coronary Heart Disease in Young Men," American College of Physicians, pp. 377-385 (Sep. 18, 2007).
- Torrejon, C. et al., "n-3 Fatty acids and cardiovascular disease: Actions and molecular mechanisms," *Prostaglandins Leukotrienes & Essent. Fatty Acids*, 77(5-6):319-26 Nov./Dec. 2007/epub Dec. 3, 2007. doi:10.1016/j.plefa.2007.10.014 (2007).
- Toth PP, Granowitz C, Hull M, Liassou D, Anderson A, Philip S. High Triglycerides are associated with increased cardiovascular events, medical costs, and resource use: A real-world administrative claims analysis of statin-treated patients with high residual cardiovascular risk. *Journal of the American Heart Association*, 7(15):e008740 (publication date Jul. 25, 2018; epublication Aug. 7, 2018).
- Transcript from Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 76 pages.
- TREND-HD Investigators, Randomized controlled trial of ethyl-eicosapentaenoic acid in Huntington disease: the TREND-HD study, *Arch Neurol.*, vol. 65(12): 1582-9 (Dec. 2008).
- Tribble DL, Holl LG, Wood PD, Krauss RM. Variations in oxidative susceptibility among six low density lipoprotein subfractions of differing density and particle size. *Atherosclerosis*. Apr. 1992;93(3):189-199.
- Tribble DL, Rizzo M, Chait A, Lewis DM, Blanche PJ, Krauss RM. Enhanced oxidative susceptibility and reduced antioxidant content of metabolic precursors of small, dense low-density lipoproteins. *Am. J. Med.* Feb. 1, 2001;110(2):103-110.
- Trilipix Package Insert (Sep. 2010)(10 pages).
- Tsimikas S, Witztum JL, Miller ER, Sasiela WJ, Szarek M, Olsson AG, Schwartz GG. High-dose atorvastatin reduces total plasma levels of oxidized phospholipids and immune complexes present on apolipoprotein B-100 in patients with acute coronary syndromes in the MIRACL trial. *Circulation.*, Sep. 14, 2004/epub Sep. 7, 2004; 110(11):1406-1412.
- Tsuruta K., et al., "Effects of purified eicosapentaenoate ethyl ester on fibrinolytic capacity in patients with stable coronary artery disease and lower extremity ischaemia" *Coron Artery Dis.* 7(11):837-42 (Nov. 1996).
- Tulenko TN, Chen M, Mason PE, Mason RP. Physical effects of cholesterol on arterial smooth muscle membranes: Evidence of immiscible cholesterol domains and alterations in bilayer width C during atherogenesis. *J. Lipid Res.* May 1998;39:947-956.
- Tungsiripat, et al., "Dyslipidemia in HIV patients," *Cleveland Clinic Journal of Medicine*, v. 72, No. 12 (Dec. 2005).
- Turini et al., "Short-term fish oil supplementation improved innate immunity, but increased ex vivo oxidation of LDL in man—a pilot study." *Eur. J. Nutr.* 40:56-65 (Apr. 2001).
- U.S. Appl. No. 14/245,499, filed Apr. 4, 2014 (now abandoned)(43 pages).
- Ullian, M.E., "Fatty acid inhibition of angiotensin II-stimulated inositol phosphates in smooth muscle cells." *Am J Physiol Heart Circ Physiol.*, 264 (2 Pt 2):H595-603 (Feb. 1993).
- Urakaze, Masaharu, et al., "Infusion of emulsified triicosapentaenoylglycerol into rabbits. The effects on platelet aggregation, polymorphonuclear leukocyte adhesion, and fatty acid composition in plasma and platelet phospholipids", *Thromb. Res.*, 44(5):673-682 (Dec. 1986).
- Urquhart et al., "Profile of eicosanoids produced by human saphenous vein endothelial cells and the effect of dietary fatty acids," *Prostaglandins Leukot. Essent. Fatty Acid*, 65(1):15-22 (Jul. 2001).
- US Food and Drug Administration and Dept of Health and Human Services. Substances affirmed as generally recognized as safe: Menhaden Oil. *Fed Register*, 62:30751-30757 (Jun. 5, 1997).
- Vaagenes et al., "The Hypolipidaemic Effect of EPA is Potentiated by 2- and 3-Methylation." In P. Quant & S. Eaton (eds.) *Current Views of Fatty Acid Oxidation and Ketogenesis from Organelles to Point Mutations; Advances in Experimental Medicine and Biology*, vol. 466, pp. 221-226 (1999).
- Vaddadi, K.S., et al., "A Randomised, Placebo-Controlled, Double-Blind Study of Treatment of Huntington's Disease with Unsaturated Fatty Acids", *Clinical Neuroscience and Neuropathology*, 13(1):29-33, (Jan. 2002).
- Vaduganathan M, Venkataramani AS, Bhatt DL. Moving toward global primordial prevention in cardiovascular disease: The heart of the matter. *J Am Coll Cardiol* Oct. 6, 2015;66(14):1535-7.
- Van der Steeg, W.A., et al., "High-density lipoprotein cholesterol, high-density lipoprotein particle size, and apolipoprotein A-I: Significance for cardiovascular risk—the IDEAL and EPIC-Norfolk studies." *J. Am. Coll. Cardiol.* 51:634-642 (Feb. 12, 2008).
- Van Do et al., "Allergy to fish parvalbumins: Studies on the cross-reactivity of allergens from 9 commonly consumed fish," *Journ. Allergy & Clin. Immunol.*, 16(6):1314-1320 (Dec. 1, 2005).
- Van Wijk et al. Rosiglitazone improves postprandial triglyceride and free fatty acid metabolism in type 2 diabetes. *Diabetes Care*, vol. 28, No. 4, (Apr. 2005) pp. 844-849.
- Varbo A, Benn M, Tybjuerg-Hansen A, Nordestgaard BG. Reply to letters regarding article, "Elevated remnant cholesterol causes both low-grade inflammation and ischemic heart disease, whereas elevated low-density lipoprotein cholesterol causes ischemic heart disease without inflammation". *Circulation*. Jun. 17, 2014; 129(24):e656.
- Varbo et al., Remnant Cholesterol as a Causal Risk Factor for Ischemic Heart Disease, *J. Am. Coll. Cardiol.*, vol. 61(4), pp. 427-436 (Jan. 29, 2013/epub Dec. 19, 2012).
- Varbo et al., Remnant cholesterol as a cause of ischemic heart disease: Evidence, definition, measurement, atherogenicity, high risk patients, and present and future treatment, *Pharmacol. Ther.*, vol. 141(3), pp. 358-367 (Mar. 2014/epub Nov. 26, 2013).
- Vascepa [package insert], Bedminster, NJ: Amarin Pharma Inc.; Jul. 2012. (12 pages).
- Vascepa [package insert]. Bedminster, NJ: Amarin Pharma Inc.; Nov. 2013. (11 pages).
- Vasudevan et al., "Effective Use of Combination Lipid Therapy", *Curr. Atheroscl. Rep.*, vol. 8, pp. 76-84 (Jan. 2006).
- Vedin, I., et al., "Effects of docosahexaenoic acid-rich n-3 fatty acid supplementation on cytokine release from blood mononuclear leukocytes: the OmegAD study." *Am J Clin Nutr* 87:1616-22 (Jun. 2008).
- Velliquette et al., "Regulation of human stearoyl-CoA desaturase by omega-3 and omega-6 fatty acids: Implications for the dietary management of elevated serum triglycerides," *Journal of Clinical Lipidology*. (Aug. 2009/epub Jun. 21, 2009) 3:281-288.
- Vergnani L, Hatric S, Ricci F, Passaro A, Manzoli N, Zuliani G, Brovkovich V, Fellin R, Malinski T. Effect of native and oxidized low-density lipoprotein on endothelial nitric oxide and superoxide production: Key role of L-arginine availability. *Circulation*. Mar. 21, 2000; 101:1261-1266.
- Verma S, Leiter LA, Bhatt DL. CANTOS ushers in a new calculus of inflammasome targeting for vascular protection—and maybe more. *Cell Metab* 26(5):703-5 (publication date Nov. 7, 2017; epublication date Oct. 19, 2017).
- Vidal F, Colome C, Martinez-Gonzalez J, Badimon L. Atherogenic concentrations of native low density lipoproteins down-regulate nitric-oxide-synthase mRNA and protein levels in endothelial cells. *Eur. J. Biochem.* Mar. 15, 1998; 252:378-384.
- Vidgren, H.M., et al., "Incorporation of n-3 fatty acids into plasma lipid fractions, and erythrocyte membranes and platelets during dietary supplementation with fish, fish oil, and docosahexaenoic acid-rich oil among healthy young men." *Lipids* 32: 697-705 (Jul. 1997).
- Virani et al., "The Role of Lipoprotein-associated Phospholipase A2 as a marker for atherosclerosis" *Curr. Atheroscler. Rep.* 9[2]: 97-103 (Aug. 2007).

(56)

## References Cited

## OTHER PUBLICATIONS

- Volcik, K.A., et al., "Peroxisome proliferator—activated receptor  $\alpha$  genetic variation interacts with n-6 and long-chain n-3 fatty acid intake to affect total cholesterol and LDL-cholesterol concentrations in the Atherosclerosis Risk in Communities Study." *Am J Clin Nutr* 87:1926-31 (Jun. 2008).
- Von Schacky C, Baumann K, Angerer P. The effect of n-3 fatty acids on coronary atherosclerosis: results from SCIMO, an angiographic study, background and implications. *Lipids* 2001 36 Suppl: S99-102.
- Von Schacky, C., "A review of omega-3 ethyl esters for cardiovascular prevention and treatment of increased blood triglyceride levels." *Vascular Health and Risk Management* 2(3): 251-262 (2006).
- Von Schacky, C., et al., "The Effect of Dietary  $\omega$ -3 Fatty Acids on Coronary Atherosclerosis: A Randomized, Double-Blind, Placebo-Controlled Trial", *American College of Physicians-American Society of Internal Medicine*, 130(7):554-562, (Apr. 6, 1999).
- Wada, M., et al., "Enzymes and receptors of prostaglandin pathways with arachidonic acid-derived versus eicosapentaenoic acid-derived substrates and products." *J. Biol. Chem.* 282(31): 22254-22266 (Aug. 3, 2007/epub May 22, 2007).
- Wagner AH, Kohler T, Ruckschloss U, Just I, Hecker M. Improvement of nitric oxide-dependent vasodilation by hmg-coa reductase inhibitors through attenuation of endothelial superoxide anion formation. *Arterioscler. Thromb. Vasc. Biol.*, Jan. 2000;20:61-69.
- Walker G, Mandagere A, Dufton C, et al. The pharmacokinetics and pharmacodynamics of warfarin in combination with ambrisentan in healthy volunteers. *Br. J. Clin. Pharmacol.* May 2009/epub Feb. 4, 2009; 67 (5): 527-34.
- Wall R, Ross RP, Fitzgerald G, Stanton C. Fatty acids from fish: the anti-inflammatory potential of long-chain omega-3 fatty acids. *Nutr Rev.* May 2010; 68:280-289.
- Walldius, G., et al., "Editorial: Rationale for using apolipoprotein B and apolipoprotein A-I as indicators of cardiac risk and as targets for lipid-lowering therapy." *European Heart Journal* 26, 210-212 (Feb. 2005/epub Dec. 15, 2004).
- Walter MF, Jacob RF, Bjork RE, Jeffers B, Buch J, Mizuno Y, Mason RP. Circulating lipid hydroperoxides predict cardiovascular events in patients with stable coronary artery disease: the PREVENT study. *J. Am. Coll. Cardiol.*, Mar. 25, 2008;51(12):1196-1202.
- Walter MF, Jacob RF, Jeffers B, Ghadanfar MM, Preston GM, Buch J, Mason RP. Serum levels of thiobarbituric acid reactive substances predict cardiovascular events in patients with stable coronary artery disease: A longitudinal analysis of the PREVENT study. *J. Am. Coll. Cardiol.* Nov. 16, 2004; 44(10):1996-2002.
- Wander, R.C., et al., "Influence of long chain polyunsaturated fatty acids on oxidation of low density lipoprotein." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 59(2):143-151 (Aug. 1998).
- Wang Q, Liang X, Wang L, Lu X, Huang J, Cao J, Li H, Gu D. Effect of omega-3 fatty acids supplementation on endothelial function: A meta-analysis of randomized controlled trials. *Atheroscler.* Apr. 2012/epub Jan. 20, 2012; 221:563-543.
- Wang, C., et al., "n-3 Fatty acids from fish or fish-oil supplements, but not  $\alpha$ -linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary-prevention studies: a systematic review." *Am J Clin Nutr* 84:5-17 (Jan. 2006).
- Wang, L., et al., "Triglyceride-rich lipoprotein lipolysis releases neutral and oxidized FFAs that induce endothelial cell inflammation." *J. Lipid Res.* 50:204-213 (Feb. 2009/epub Sep. 23, 2008).
- Warren, Stephen T., "The Expanding World of Trinucleotide Repeats", *Science*, 271:1374-1375, (Mar. 8, 1996).
- Wassmann S, Laufs U, Muller K, Konkol C, Ahlbory K, Baumer AT, Linz W, Bohm M, Nickenig G. Cellular antioxidant effects of atorvastatin in vitro and in vivo. *Arterioscler. Thromb. Vasc. Biol.* Feb. 1, 2002; 22:300-305.
- Watanabe et al., "Bile acids lower triglyceride levels via a pathway involving FXR, SHP, and SREBP-1c," *J Clin Invest.* 113(10): 1408-1418 (May 2004).
- Watanabe T, Ando K, Daidoji H, et al. A randomized controlled trial of eicosapentaenoic acid in patients with coronary heart disease on statins. *J Cardiol* 70(6):537-44 (publication date Dec. 2017; epublication date Aug. 31, 2017).
- Watanabe, Ikuyoshi, et al., "Usefulness of EPA-E (eicosapentaenoic acid ethyl ester) in preventing neointimal formation after vascular injury", *Kokyu to Junkan*, 42(7):673-677 (1994) (with English summary).
- Weaver, K.L., et al., "Effect of Dietary Fatty Acids on Inflammatory Gene Expression in Healthy Humans." *J. Biol. Chem.*, 284(23): 15400-15407 (2009) (published online Apr. 9, 2009).
- Webcast Information for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, (1 page.).
- Weber, P., "Triglyceride-lowering effect of n-3 long chain polyunsaturated fatty acid: eicosapentaenoic acid vs. docosahexaenoic acid." *Lipids* 34: S269 (1999).
- Wei et al., Effects of [EPA] Versus [DHA] on Serum Lipids: A Systematic Review and Meta-Analysis, *13 Current Atherosclerosis Rep.* 13(6):474-483 (Dec. 2011).
- Wei LJ, Lin DY, Weissfeld L. Regression analysis of multivariate incomplete failure time data by modeling marginal distributions. *J Am Stat Assoc.* 84(408):1065-1073 (publication date Dec. 1989).
- Westerveld H.T. et al., "Effects of low-dose EPA-Eon glycemic control, lipid profile, lipoprotein(a), platelet aggregation, viscosity, and platelet and vessel wall interaction in NIDDM" *Diabetes Care* 16(5):683-8 (May 1993).
- Westphal, S., et al., "Postprandial chylomicrons and VLDLs in severe hypertriglycerolemia are lowered more effectively than are chylomicron remnants after treatment with n3 fatty acids." *Am J Clin Nutr* 71:914-20 (Apr. 2000).
- Whelan, J., et al., "Evidence that dietary arachidonic acid increases circulating triglycerides." *Lipids* 30, 425-429 (May 1995).
- Wierzbicki, A.S., "Editorial: Newer, lower, better? Lipid drugs and cardiovascular disease—the continuing story." *Int J Clin Pract*, 61(7):1064-1067 (Jul. 2007).
- Wierzbicki, A.S., "Editorial: Raising HDL-C: back to the future?" *Int J Clin Pract*, 61(7): 1069-1071 (Jul. 2007).
- Wikipedia, "Diabetes mellitus," Dec. 12, 2016 (Dec. 12, 2016), retrieved on Jul. 30, 2018 from [https://en.wikipedia.org/w/index.php?title=Diabetes\\_mellitus&oldid=754431573](https://en.wikipedia.org/w/index.php?title=Diabetes_mellitus&oldid=754431573); entire document, especially p. 1, paragraph 1.
- Wikipedia, "Ethyl eicosapentaenoic acid," Apr. 1, 2016 (Apr. 1, 2016); retrieved on Jul. 27, 2018 from [https://en.wikipedia.org/w/index.php?title=Ethyl\\_eicosapentaenoic\\_acid&oldid=713086755](https://en.wikipedia.org/w/index.php?title=Ethyl_eicosapentaenoic_acid&oldid=713086755); entire document, especially p. 1, col. 2 and p. 3, paragraph 2;.
- Williams et al., "NADPH Oxidase Inhibitors New Antihypertensive Agents?" *J. Cardiovasc Pharmacol* 50(1):9-16 (Jul. 1, 2007).
- Willumsen, N. et al., *Biochimica et Biophysica Acta*. vol. 1369, "On the effect of 2-deuterium- and 2-methyl-eicosapentaenoic acid derivatives on triglycerides, peroxisomal beta-oxidation and platelet aggregation in rats," pp. 193-203, (Mar. 2, 1998).
- Willumsen, N., et al., "Eicosapentaenoic acid, but not docosahexaenoic acid, increased mitochondrial fatty acid oxidation and upregulates 2,3-dienoyl-CoA reductase gene expression in rats." *Lipids*, 31:579-592 (Jun. 1996).
- Wilson Omega 3 fish oil: EPA versus DHA (Dietivity.com, 1-16) (2006).
- Wilt, VM & Gumm, JG, "Isolated low high-density lipoprotein cholesterol", *Ann. Pharmacol.*, 31:89-97, (Jan. 1997).
- Wink, J., et al., "Effect of very-low-dose niacin on high-density lipoprotein in patients undergoing long-term statin therapy", *Am. Heart J.*, 143:514-518, (Mar. 2002).
- Wittrup HH, Tybjærg-Hansen A, Nordestgaard BG. Lipoprotein lipase mutations, plasma lipids and lipoproteins, and risk of ischemic heart disease: a meta-analysis. *Circulation.*, Jun. 8, 1999;99:2901-2907.
- Witztum JL. The oxidation hypothesis of atherosclerosis. *Lancet*, Sep. 17, 1994;344(8925):793-795.
- Wojczynski et al., "High-fat meal effect on LDL, HDL and VLDL particle size and .No. In the Genetics of Lipid-Lowering Drugs and Diet Network (GOLDN): an interventional study," *Lipids in Health and Disease* 10:181, pp. 1-11 (Oct. 18, 2011).

(56)

## References Cited

## OTHER PUBLICATIONS

- Wojenski, C.M., et al., "Eicosapentaenoic acid ethyl ester as an antithrombotic agent: comparison to an extract of fish oil." *Biochimica et Biophysica Acta*. 1081:33-38 (Jan. 4, 1991).
- Wong, S.H., et al., "Effects of eicosapentaenoic and docosahexaenoic acids on Apoprotein B mRNA and secretion of very low density lipoprotein in HepG2 cells." *Arterioscler. Thromb. Vasc. Biol.* 9:836-841 (Nov./Dec. 1989).
- Wood et al., "Carbohydrate Restriction Alters Lipoprotein Metabolism by Modifying VLDL, LDL and HDL Subfraction Distribution and Size in Overweight Men," *Journ. of Nutrition*, 136(2):384-9 (Feb. 2006).
- Woodman et al., "Effects of Purified Eicosapentaenoic and Docosahexaenoic Acids on Glycemic Control, Blood Pressure, and Serum Lipids in Type 2 Diabetic Patients with Treated Hypertension", *The American Journal of Clinical Nutrition: Official Journal of the American Society for Clinical Nutrition, Inc.*, 76(5):1007-1015 (Nov. 1, 2002).
- Woodman, R.J., et al., "Effects of purified eicosapentaenoic acid and docosahexaenoic acid on platelet, fibrinolytic and vascular function in hypertensive type 2 diabetic patients." *Atherosclerosis* 166: 85-93 (Jan. 2003).
- Wu et al., "Diabetic dyslipidemia," *Metabolism Clinical and Experimental*, 63:1469-1479 (Dec. 2014)(available online Aug. 29, 2014).
- Wu, W.H., et al., "Effects of docosahexaenoic acid supplementation on blood lipids, estrogen metabolism, and in vivo oxidative stress in postmenopausal vegetarian women." *Eur J Clin Nutr.*, 60:386-392 (Mar. 2006).
- Xiao, Y.F., et al., "Inhibitory effect of n-3 fish oil fatty acids on cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchange currents in HEK293t cells." *Biochemical and Biophysical Research Communications* 321: 116-123 (Aug. 13, 2004).
- Xiao, Y-F., et al., "Blocking effects of polyunsaturated fatty acids on Na<sup>+</sup> channels of neonatal rat ventricular myocytes." *Proc. Natl. Acad. Sci.* 92: 11000-11004 (Nov. 21, 1995).
- Xiao, Y-F., et al., "Fatty acids suppress voltage-gated Na<sup>+</sup> currents in HEK293t cells transfected with the  $\alpha$ -subunit of the human cardiac Na<sup>+</sup> channel." *Proc. Natl. Acad. Sci.* 95: 2680-2685 (Mar. 3, 1998).
- Xydakis, AM et al., "Combination therapy for combined dyslipidemia," *American Journal of Cardiology*, Nov. 20, 2002 US, vol. 90, No. 10 Suppl. 2, p. 21 K-29K (Nov. 20, 2002).
- Yacyshyn BR, Thomson AB. The clinical importance of proton pump inhibitor pharmacokinetics. *Digestion* 2002 66 (2): 67-78.
- Yadav D, Pitchumoni CS. Issues in Hyperlipidemic Pancreatitis. *J Clin Gastroenterol* 236(1):54-62, Jan. 2003.
- Yagi K. Assay for blood plasma or serum. *Methods Enzymol.* 1984;105:328-331.
- Yamagishi K, Nettleton J, Folsom A. Plasma fatty acid composition and incident heart failure in middle-aged adults: The Atherosclerosis Risk in Communities (ARIC) Study. *Am Heart J.*, Nov. 2008/epub Aug. 29, 2008; 156:965-974.
- Yamakawa K, Shimabukuro M, Higa N, Asahi T, Ohba K, Arasaki O, Higa M, Oshiro Y, Yoshida H, Higa T, Saito T, Ueda S, Masuzaki H, Sata M. Eicosapentaenoic Acid Supplementation Changes Fatty Acid Composition and Corrects Endothelial Dysfunction in Hyperlipidemic Patients. *Cardiol Res Practice*. Dec. 26, 2012; epub Article ID 754181.
- Yamamoto, H. et al., Improvement of coronary vasomotion with Eicosapentaenoic acid does not inhibit acetylcholine-induced coronary vasospasm in patients with variant angina. *Jpn Cir J.* 59(9):608-16 (Sep. 1995).
- Yamamoto, K., et al., "4-Hydroxydocosahexaenoic acid, a potent Peroxisome Proliferator-Activated Receptor C agonist alleviates the symptoms of DSS-induced colitis." *Biochemical and Biophysical Research Communications* 367: 566-572 (Mar. 14, 2008/epub Jan. 10, 2008).
- Yamano T, Kubo T, Shiono Y, et al. Impact of eicosapentaenoic acid treatment on the fibrous cap thickness in patients with coronary atherosclerotic plaque: an optical coherence tomography study. *J Atheroscler Thromb.* 2015/epub Aug. 15, 2014;22:52-61.
- Yamashita et al., *J. Biochem.*, vol. 122, No. 1, "Acyl-transferases and Transacylases Involved in Fatty Acid Remodeling of Phospholipids and Metabolism of Bioactive Lipids in Mammalian Cells", pp. 1-16 (Jul. 1997).
- Yamashita, N., et al., "Inhibition of natural killer cell activity of human lymphocytes by eicosapentaenoic acid." *Biochem. Biophys. Res. Comm.* 138(3): 1058-1067 (Aug. 25, 1986).
- Yamazaki et al., "Changes in fatty acid composition in rat blood and organs after infusion of eicosapentaenoic acid ethyl ester", *Biochim. Biophys. ACTA*, 1128(1):35-43, (Sep. 22, 1992).
- Yamazaki, et. al., "Dissolution tests by RDC method for soft gelatin capsules containing ethyl icosapentate," *Pharm. Tech. Japan*, vol. 15, No. 4, pp. 595-603 Abstract (Apr. 1999) (with English abstract).
- Yang, S.P., et al., "Eicosapentaenoic acid attenuates vascular endothelial growth factor-induced proliferation via inhibiting Flk-1 receptor expression in bovine carotid artery endothelial cells." *J. Cell. Physio.* 176:342-349 (Aug. 1998).
- Yano T, Mizuguchi K, Takasugi K, Tanaka Y, Sato M. "Effects of ethyl all-cis-5,8,11,14,17-icosapentaenoate on low density lipoprotein in rabbits," *Yakugaku Zasshi*, 115:843-51 (Oct. 1995).
- Yano, T., et al., "Effects of ethyl-all-cis-5,8,11,14,17-icosapentaenoate (EPA-E), pravastatin and their combination on serum lipids and intimal thickening of cuff-sheathed carotid artery in rabbits." *Life Sciences*, 61(20):2007-2015 (1997).
- Yao et al., "Oxidized high density lipoprotein induces macrophage apoptosis via toll-like receptor 4-dependent CHOIP pathway," *Journ. Lipid Res.*, 58:164-177 (Jan. 2017)(First published Nov. 28, 2016).
- Yates RA, Wong J, Seiberling M, et al. The effect of anastrozole on the single-dose pharmacokinetics and anticoagulant activity of warfarin in healthy volunteers. *Br. J. Clin. Pharmacol.* May 2001 51(5): 429-35.
- Yerram, N.R., et al., "Eicosapentaenoic acid metabolism in brain microvessel endothelium: effect on prostaglandin formation." *J. Lipid Res.*30:1747-1757 (Nov. 1989).
- Yokoyama et al., "Effects of eicosapentaenoic acid on cardiovascular events in Japanese patients with hypercholesterolemia: Rationale, design, and baseline characteristics of the Japan EPA Lipid Intervention Study (JELIS)," *Amer. Heart Journal* 146(4):613-620 (Oct. 2003).
- Yokoyama et al., "Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomized open-label, blinded endpoint analysis", *Lancet*, vol. 369, pp. 1090-1098 (Mar. 31, 2007).
- Yorioka, N, "Lipid-lowering therapy and coagulation/fibrinolysis parameters in patients on peritoneal dialysis," *The International Journal of Artificial Organs*, vol. 23(1):27-32 (Jan. 2000).
- Yoshimura et al., "Effects of highly purified eicosapentaenoic acid on plasma beta thromboglobulin level and vascular reactivity to angiotensin II", *Artery*, 14(5):295-303 (1987).
- Zaima, N., et al., "Trans geometric isomers of EPA decrease LXRA-induced cellular triacylglycerol via suppression of SREBP-1c and PGC-1 $\beta$ ," *J. Lipid Res.* 47: 2712-2717 (Dec. 2006).
- Zalewski et al., Role of Lipoprotein-Associated Phospholipase A2 in Atherosclerosis: Biology, Epidemiology, and Possible Therapeutic Target, *Arteriosclerosis, Thrombosis, & Vascular Biology* 25(5):923-931 (May 2005/epub Feb. 24, 2005)
- Zanarini, et al., "Omega-3 Fatty Acid Treatment of Women with Borderline Personality Disorder: A Double-Blind, Placebo-Controlled Pilot Study," *Am J Psychiatry*, 160:167-169 (Jan. 2003).
- Zhan, S. et. al. "Meta-analysis of the effects of soy protein containing isoflavones on the lipid profile," *Am. J. Clin. Nutr.* (Feb. 2005), 81, p. 397-408.
- Zhang, M., et al., "Effects of eicosapentaenoic acid on the early stage of type 2 diabetic nephropathy in KKAY/Ta mice: involvement of anti-inflammation and antioxidative stress." *Metabolism Clinical and Experimental* 55:1590-1598 (Dec. 2006).
- Zhang, Y.W., et al., "Inhibitory effects of eicosapentaenoic acid (EPA) on the hypoxia/reoxygenation-induced tyrosine kinase activation in cultured human umbilical vein endothelial cells." *Prostaglandins, Leukotrienes and Essential FattyAcids* 67(4):253-261 (Oct. 2002).

(56)

## References Cited

## OTHER PUBLICATIONS

- Zhang, Y.W., et al., "Pretreatment with eicosapentaenoic acid prevented hypoxia/reoxygenation-induced abnormality in endothelial gap junctional intercellular communication through inhibiting the tyrosine kinase activity." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 61(1): 33-40 (Jul. 1999).
- Zhao et al., "Polyunsaturated Fatty Acids are FXR Ligands and Differentially Regulate Expression of FXR Targets," *DNA and Cell Biology*, 23(8):519-526 (Aug. 25, 2004).
- Zhao, G. et al., "Dietary  $\alpha$ -linolenic acid inhibits proinflammatory cytokine production by peripheral blood mononuclear cells in hypercholesterolemic subjects." *Am J Clin Nutr* 85:385-91 (Feb. 2007).
- Zhao, G., et al., "Dietary  $\alpha$ -linolenic acid reduces inflammatory and lipid cardiovascular risk factors in hypercholesterolemic men and women." *J. Nutr.* 134: 2991-2997 (Nov. 2004).
- Zheng et al., "Function of  $\omega$ -3 long chain unsaturated fatty acid in metabolic syndrome," *Chinese Journal of Endocrinology and Metabolism*, vol. 27, No. 9, pp. 787-790 (Sep. 30, 2011)(with English translation).
- Ziegler, D., et al., "Treatment of symptomatic diabetic polyneuropathy with the antioxidant  $\alpha$ -lipoic acid: A 7-month multicenter randomized controlled trial (ALADIN III Study)." *Diabetes Care* 22:1296-1301 (Aug. 1999).
- Zimmerman JJ, Raible DG, Harper DM, et al. Evaluation of a potential tigecycline-warfarin drug interaction. *Pharmacotherapy* Jul. 28, 2008 (7): 895-905.
- Zuijdgheest-van Leeuwen, et al., "N-3 Fatty Acids Administered as Triacylglycerols or as Ethyl Esters Have Different Effects on Serum Lipid Concentrations in Healthy Subjects," *N-3 Fatty Acids, Lipid Metabolism and Cancer*, pp. 89-100 (2000).
- Zuijdgheest-van Leeuwen, S.D., et al., "Incorporation and washout of orally administered n-3 fatty acid ethyl esters in different plasma lipid fractions." *British Journal of Nutrition* 82:481-488 (1999).
- Zuijdgheest-van Leeuwen, SD, et al., "Eicosapentaenoic acid inhibits lipolysis in weight-losing cancer patients as well as in healthy volunteers," *Eur J Gastroenterol & Hepatol.*, 10(12):A67 (1998).
- Zvyaga T, Chang SY, Chen C, et al. Evaluation of six proton pump inhibitors as inhibitors of various human cytochromes P450: focus on cytochrome P450 2C19. *Drug Metab. Dispos.* Sep. 2012 40(9): 1698-711.
- Third Report of the National Cholesterol Education Program (NCEPP) Expert Panel on Detection, Evaluation, and Treatment of High blood Cholesterol in Adults (Adult Treatment Panel III), NIH Pub. No. 02-5215, Sep. 2002, 220 pgs.
- Zimmer et al., "Danger signaling in Atherosclerosis," *Circ. Res.*, 2015; 116:323-340.
- Bays HE et al., "AMR101, a Pure Ethyl Eicosapentaenoic Acid Omega-3 Fatty Acid: Effects on Inflammation-Associated End Points from the Marine and Anchor Studies," *Journ. Clin. Lipid.*, vol. 6 No. 3, p. 279 (May 30, 2012), Abstract #150.
- Bays HE et al., Effects of prescription omega-3-acid ethyl esters on non-high-density lipoprotein cholesterol when coadministered with escalating doses of atorvastatin; *Mayo Clinic Proc.* 85(2):122-128 (Feb. 2010).
- Billman et al., "Effects of dietary omega-3 fatty acids on ventricular function in dogs with healed myocardial infarctions: in vivo and in vitro studies." *Am. J. Physiol Heart. Circ. Physiol.*, 298:H1219-H1228 (Jan. 22, 2010).
- Connor et al, "Are Fish Oils Beneficial in the Prevention and Treatment of Coronary Artery Disease?" *Am J Clin Nutr* vol. 66, No. 4, Jan. 1, 1997, pp. 1020S-1031S, XP002502041.
- Gromova, O.A et al, published Jan. 2009, [found online Dec. 11, 2019] (found from Internet: t-patient.ru/articles/6417/) with English

Machine Translation, **И ДР. СИСТЕМАТИЧЕСКИЙ АНАЛИЗ** биохимических эффектов эйкозапентаеновой

**И докозагексаеновой омега 3 ПНЖК на физиологию беременности И развитие плода. Трудный пациент.**

Январь 2009.

- Ivanova et al., "Small Dense Low-Density Lipoprotein as Biomarker for Atherosclerotic Diseases," May 9, 2017, *Oxidative Medicine and Cellular Longevity* (2017), 10 pp.
- Li, X., et al., "Protection against fine particle-induced pulmonary and systemic inflammation by omega-3 polyunsaturated fatty acids." vol. 1861, No. 3, pp. 577-584 (Dec. 21, 2016).
- Lin, Z., et al., "Cardiovascular Benefits of Fish-Oil Supplementation Against Fine Particulate Air Pollution in China." Apr. 30, 2019; *73(16):2076-2085*.
- McCabe, John B. "Literature of Resuscitation", *Resuscitation*, Elsevier, IE, vol. 19, No. 3 (Jun. 1, 1990), vol. 19, pp. 303-319, DOI: 10.1016/0300-9572(90)90109-R.
- Pepys, M.B. et al, "C-reactive protein: a critical update", *Journal of Clinical Investigation*, e-pub Jun. 15, 2003; Jul. 2003, vol. 111(12), pp. 1805-1812.
- Poirier, "Obesity and Cardiovascular Disease: Pathophysiology, Evaluation, and Effect of Weight Loss", *Circulation*, Feb. 14, 2006;113(6):898-918. Epub Dec. 27, 2005.
- Shen, W., et al., "Influence of Omega-3 Fatty Acids Intake on Human Responsiveness to Ambient Air Pollution Exposure", Apr. 1, 2017, *The FASEB Journal*; retrieved from the Internet: URL://[https://www.fasebj.org/doi/abs/10.1096/fasebj.31.1\\_supplement.971.2](https://www.fasebj.org/doi/abs/10.1096/fasebj.31.1_supplement.971.2); [retrieved on Jan. 7, 2020].
- Sherratt SCR, Mason RP. Eicosapentaenoic acid and docosahexaenoic acid have distinct membrane locations and lipid interactions as determined by X-ray diffraction. *Chem Phys Lipids* 212:73-9 (publication date May 2018, epublication date Jan. 31, 2018).
- Signori, S., et al., "Administration of omega-3 fatty acids and Raloxifene to women at high risk of breast cancer: interim feasibility and biomarkers analysis from a clinical trial," *European Journal of Clin. Nutr.*, 66, 878-884 (published online Jun. 6, 2012).
- Siscovick et al., "Dietary Intake and Cell Membrane levels of Long-chain N-3 Polyunsaturated Fatty Acids and the Risk of Primary Cardiac Arrest", *JAMA*, vol. 274, No. 17, Nov. 1, 1995, pp. 1363-1367, XP008041164.
- Third Report of the National Cholesterol Education Program (NCEPP) Expert Panel on Detection, Evaluation, and Treatment of High blood Cholesterol in Adults (Adult Treatment Panel III) May 2001, National Institutes of Health, Publication No. 01-3670.
- Thomas II et al., "Prostate Cancer Risk in Men with Baseline History of Coronary Artery Disease: Results from the REDUCE Study," *Cancer Epidemiology, Biomarkers and Prevention*, 21(4) published online Feb. 7, 2012.
- Tong, H., et al., "Omega-3 fatty acid supplementation appears to attenuate particulate air pollution-induced cardiac effects and lipid changes in healthy middle-aged adults." *Environ. Health Perspect.*, Jul. 2012, epub Apr. 19, 2012; 120(7):952-7.
- U.S. District Court of Nevada, judgment dated Mar. 30, 2020 in *Amarin Pharma, Inc. et al. v. Hikma Pharmaceuticals USA Inc. et al.*, Case 2:16-cv-02525-MMC-NJK Document 381 Filed Mar. 30, 2020 (70 pages).
- Förstermann et al., "Roles of Vascular Oxidative Stress and Nitric Oxide in the Pathogenesis of Atherosclerosis," *Circ. Res.* Feb. 17, 2017; 120(4):713-735 (Accepted Dec. 26, 2016).
- Li, Q, et al., Eicosapentaenoic acid modifies lipid composition in caveolae and induces translocation of endothelial nitric oxid synthase, *Biochimie* 89 (2007) pp. 169-177 (publ online Nov. 7, 2006).
- Martinez et al., "Serum level changes of long chain-polyunsaturated fatty acids in patients undergoing periodontal therapy combined with one year of omega-3 supplementation: a pilot randomized clinical trial," *Journal of Periodontal & Implant Science*, Aug. 28, 2014, vol. 44, pp. 169-177.
- Morin et al., "Anti-proliferative effects of a new docosapentaenoic acid monoacylglyceride in colorectal carcinoma cells;" *Prostaglandins, Leukotrienes and Essential Fatty Acids*, Aug. 7, 2013, vol. 89, pp. 203-213.
- National Kidney Foundation, *Clinical Practice Guidelines for Chronic Kidney Disease: Evaluation, Classification and Stratification*; Copyright 2002, 356 pages.

(56)

**References Cited**

OTHER PUBLICATIONS


NCBI , Table 6-1 "Factors for Converting International Units of Vitamin E<sup>a</sup> to  $\alpha$ -Tocopherol<sup>b</sup> (mg) to Meet Recommended Intake," taken from Dietary Reference Intakes for Vitamin C, Vitamin E, Selenium and Carotenoids, Institute of Medicine (US) Panel on Dietary Antioxidants and Related Compounds, Washington DC, National Academies Press (US) 2000.

Saito et al., "Effects of Ethyl Eicosapentaenoate (EPA-E), Clopidogrel, and Their Combination on Platelet Aggregation and Bleeding Time;" Japanese Pharmacology & Therapeutics, Feb. 20, 2007 (epub Jan. 2007), vol. 35, No. 2, pp. 179-185 (English abstract only).

Vega et al., "Hypercholesterolemia with Cholesterol-Enriched LDL and Normal Levels of LDL-Apolipoprotein B," Arteriosclerosis, Thrombosis and Vascular Biology, Apr. 1996; 16(4):517-22 (Accepted Jan. 3, 1996).

\* cited by examiner






\*P<0.05; \*\*P<0.01; NS=not significant.

†Median differences in percent changes vs placebo are Hodges-Lehmann medians.


FIG. 1



\*P<0.05; NS=not significant.

†Median differences in percent changes vs placebo are Hodges-Lehmann medians.

FIG. 2



\*P<0.001; \*\*P<0.05; NS=not significant. Statin intensity definitions are in the Definitions and Abbreviations section.

†Median differences in percent changes vs placebo are Hodges-Lehmann medians.

FIG. 3

## 1

## METHODS OF REDUCING RLP-C

CROSS-REFERENCE TO RELATED  
APPLICATIONS

This application is a continuation of U.S. patent application Ser. No. 14/737,115 filed Jun. 11, 2015, which claims priority to U.S. Patent Application No. 62/077,478 filed Nov. 10, 2014 and U.S. Patent Application No. 62/010,847 filed Jun. 11, 2014, the contents of which are hereby incorporated in their entirety.

## BACKGROUND

Cardiovascular disease is one of the leading causes of death in the United States and most European countries. It is estimated that over 70 million people in the United States alone suffer from a cardiovascular disease or disorder including but not limited to high blood pressure, coronary heart disease, dyslipidemia, congestive heart failure and stroke. A need exists for improved treatments for cardiovascular diseases and disorders.

Remnant-like particle cholesterol (“RLP-C”) represents the cholesterol carried by partially catabolized triglyceride (“TG”)-rich lipoproteins such as very-low-density lipoproteins (“VLDL”) in the fasted state and chylomicron remnants in the post-prandial state. Increased RLP-C levels are generally considered to be atherogenic, and may increase the risk of atherosclerotic cardiovascular disease.

## SUMMARY

In various embodiments, the present invention provides methods of treating and/or preventing cardiovascular-related diseases and, in particular, a method of blood lipid therapy comprising administering to a subject in need thereof a pharmaceutical composition comprising eicosapentaenoic acid or a derivative thereof. In one embodiment, the method comprises administering to a subject about 2 g to about 4 g per day (e.g., about 3.8 g to about 4 g per day) of eicosapentaenoic acid or derivative thereof.

In one embodiment, the composition contains not more than 10%, by weight, docosahexaenoic acid or derivative thereof, substantially no docosahexaenoic acid or derivative thereof, or no docosahexaenoic acid or derivative thereof. In another embodiment, eicosapentaenoic acid ethyl ester comprises at least 90%, by weight, of all fatty acids present in the composition. In one embodiment, the present invention provides methods of reducing the level of RLP-C in a subject compared to baseline or to placebo control.

In one embodiment, a pharmaceutical composition useful in accordance with the invention comprises, consists of or consists essentially of at least 95% by weight ethyl eicosapentaenoate (EPA-E), about 0.2% to about 0.5% by weight ethyl octadecatetraenoate (ODTA-E), about 0.05% to about 0.25% by weight ethyl nonadecapentaenoate (NDPA-E), about 0.2% to about 0.45% by weight ethyl arachidonate (AA-E), about 0.3% to about 0.5% by weight ethyl eicosatetraenoate (ETA-E), and about 0.05% to about 0.32% ethyl heneicosapentaenoate (HPA-E). In another embodiment, the composition is present in a capsule shell. In another embodiment, the composition contains substantially no or no amount of docosahexaenoic acid (DHA) or derivative thereof such as ethyl-DHA (DHA-E).

In another embodiment, the invention provides a method of treating moderate to severe hypertriglyceridemia com-

## 2

prising administering a composition as described herein to a subject in need thereof one to about four times per day.

These and other embodiments of the present invention will be disclosed in further detail herein below.

## BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows median percent change in RLP-C levels from baseline to week 12 compared to placebo in subjects having very high triglycerides (“MARINE”) or high triglycerides and on statin therapy (“ANCHOR”).

FIG. 2 shows median percent change in RLP-C levels from baseline to week 12 compared to placebo in subjects having high triglycerides by statin use.

FIG. 3 shows median percent change in RLP-C levels from baseline to week 12 compared to placebo in subjects having very high triglycerides and on statin therapy, according to intensity of the statin therapy.

## DETAILED DESCRIPTION

While the present invention is capable of being embodied in various forms, the description below of several embodiments is made with the understanding that the present disclosure is to be considered as an exemplification of the invention, and is not intended to limit the invention to the specific embodiments illustrated. Headings are provided for convenience only and are not to be construed to limit the invention in any manner. Embodiments illustrated under any heading may be combined with embodiments illustrated under any other heading.

The use of numerical values in the various quantitative values specified in this application, unless expressly indicated otherwise, are stated as approximations as though the minimum and maximum values within the stated ranges were both preceded by the word “about.” Also, the disclosure of ranges is intended as a continuous range including every value between the minimum and maximum values recited as well as any ranges that can be formed by such values. Also disclosed herein are any and all ratios (and ranges of any such ratios) that can be formed by dividing a disclosed numeric value into any other disclosed numeric value. Accordingly, the skilled person will appreciate that many such ratios, ranges, and ranges of ratios can be unambiguously derived from the numerical values presented herein and in all instances such ratios, ranges, and ranges of ratios represent various embodiments of the present invention.

In one embodiment, the invention provides a method for treatment and/or prevention of a cardiovascular-related disease. The term “cardiovascular-related disease” herein refers to any disease or disorder of the heart or blood vessels (i.e. arteries and veins) or any symptom thereof. Non-limiting examples of cardiovascular-related disease and disorders include hypertriglyceridemia, hypercholesterolemia, mixed dyslipidemia, coronary heart disease, vascular disease, stroke, atherosclerosis, arrhythmia, hypertension, myocardial infarction, and other cardiovascular events.

The term “treatment” in relation a given disease or disorder, includes, but is not limited to, inhibiting the disease or disorder, for example, arresting the development of the disease or disorder; relieving the disease or disorder, for example, causing regression of the disease or disorder; or relieving a condition caused by or resulting from the disease or disorder, for example, relieving, preventing or treating symptoms of the disease or disorder. The term “prevention” in relation to a given disease or disorder means: preventing

the onset of disease development if none had occurred, preventing the disease or disorder from occurring in a subject that may be predisposed to the disorder or disease but has not yet been diagnosed as having the disorder or disease, and/or preventing further disease/disorder development if already present.

In one embodiment, the present invention provides a method of blood lipid therapy comprising administering to a subject or subject group in need thereof a pharmaceutical composition as described herein. In another embodiment, the subject or subject group has hypertriglyceridemia, hypercholesterolemia, mixed dyslipidemia and/or very high triglycerides.

In another embodiment, the subject or subject group being treated has a baseline triglyceride level (or mean or median baseline triglyceride level in the case of a subject group), fed or fasting, of about 200 mg/dl to 499 mg/dl. In another embodiment, the subject or subject group has a baseline LDL-C level (or mean or median baseline LDL-C level), despite stable statin therapy, of about 40 mg/dl to about 115 mg/dl or about 40 to about 100 mg/dl.

In one embodiment, the subject or subject group being treated in accordance with methods of the invention is on concomitant statin therapy, for example atorvastatin, rosuvastatin or simvastatin therapy (with or without ezetimibe). In another embodiment, the subject is on concomitant stable statin therapy at time of initiation of EPA therapy.

In another embodiment, the subject or subject group being treated in accordance with methods of the invention has a body mass index (BMI or mean BMI) of not more than about 45 kg/m<sup>2</sup>.

In one embodiment, the invention provides a method of lowering triglycerides in a subject on stable statin therapy having baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl, the method comprising administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of EPA or a derivative thereof, wherein upon administering the composition to the subject daily for a period of 1 to about 12 weeks the subject exhibits at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, or at least 75% lower fasting triglycerides than a control subject maintained on stable statin therapy (and optionally placebo matching the EPA) without concomitant EPA for a period of 1 to about 12 weeks, wherein the control subject also has baseline fasting triglycerides of about 200 mg/dl to about 500 mg/dl. The term "stable statin therapy" herein means that the subject, subject group, control subject or control subject group in question has been taking a stable daily dose of a statin (e.g. atorvastatin, rosuvastatin or simvastatin) for at least 4 weeks prior to the baseline fasting triglyceride measurement (the "qualifying period"). For example, a subject or control subject on stable statin therapy would receive a constant daily (i.e. the same dose each day) statin dose for at least 4 weeks immediately prior to baseline fasting triglyceride measurement. In one embodiment, the subject's and control subject's LDL-C is maintained between about 40 mg/dl and about 115 mg/dl or about 40 mg/dl to about 100 mg/dl during the qualifying period. The subject and control subject are then continued on their stable statin dose for the 12 week period post baseline.

In one embodiment, the statin is administered to the subject and the control subject in an amount of about 1 mg to about 500 mg, about 5 mg to about 200 mg, or about 10 mg to about 100 mg, for example about 1 mg, about 2 mg, about 3 mg, about 4 mg, about 5 mg, about 6 mg, about 7

mg, about 8 mg, about 9 mg, or about 10 mg; about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 80 mg, about 90 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, or about 500 mg. In another embodiment, the subject (and optionally the control subject) has a baseline LDL-C level, despite stable statin therapy, of about 40 mg/dl to about 115 mg/dl or about 40 mg/dl to about 100 mg/dl. In another embodiment, the subject and/or control subject has a body mass index (BMI; or mean BMI) of not more than about 45 kg/m<sup>2</sup>.

In another embodiment, the invention provides a method of lowering triglycerides in a subject group on stable statin therapy having mean baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl, the method comprising administering to members of the subject group a pharmaceutical composition comprising about 1 g to about 4 g of EPA per day, wherein upon administering the composition to the members of the subject group daily for a period of 1 to about 12 weeks the subject group exhibits at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75% lower mean fasting triglycerides than a control subject group maintained on stable statin therapy without concomitant EPA (optionally with matching placebo) for a period of 1 to about 12 weeks, wherein the control subject group also has mean baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl. In a related embodiment, the stable statin therapy will be sufficient such that the subject group has a mean LDL-C level about at least about 40 mg/dl and not more than about 100 mg/dl or about 40 mg/dl to about 100 mg/dl for the 4 weeks immediately prior to the baseline fasting triglyceride measurement.

In another embodiment, the invention provides a method of lowering triglycerides in subject group on stable statin therapy and having a mean baseline fasting triglyceride level of about 200 mg/dl to 499 mg/dl, the method comprising administering to members of the subject group a pharmaceutical composition comprising about 1 g to about 4 g of EPA, wherein upon administering the composition to members of the subject group daily for a period of 1 to about 12 weeks the subject group exhibits: (a) at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75% lower mean fasting triglycerides by comparison with a control subject group maintained on stable statin therapy without concomitant EPA (optionally with matching placebo) for a period of 1 to about 12 weeks, and (b) no serum LDL-C increase, no statistically significant serum LDL-C increase, a serum LDL-C decrease, or the subject is statistically non-inferior to the control subjects (statin plus optional placebo) in regard to serum LDL-C elevation) no increase in mean serum LDL-C levels compared to baseline, wherein the control subject also has mean baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl.

In another embodiment, the invention provides a method of lowering triglycerides in subject on stable statin therapy and having mean baseline fasting triglyceride level of about 200 mg/dl to 499 mg/dl, the method comprising administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of EPA, wherein upon adminis-

5

tering the composition to the subject daily for a period of 1 to about 12 weeks the subject exhibits (a) at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, or at least 75% lower fasting triglycerides by comparison with a control subject maintained on stable statin therapy without concomitant EPA for a period of 1 to about 12 weeks and (b) no increase in serum LDL-C levels compared to baseline, wherein the control subject also has baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl.

In another embodiment, the invention provides a method of lowering triglycerides in subject group on stable statin therapy and having mean baseline fasting triglyceride level of about 200 mg/dl to 499 mg/dl, the method comprising administering to members of the subject group a pharmaceutical composition comprising about 1 g to about 4 g of EPA, wherein upon administering the composition to the members of the subject group daily for a period of 1 to about 12 weeks the subject group exhibits: (a) at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75% lower mean fasting triglycerides and (b) at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45% or at least 50% lower mean serum LDL-C levels by comparison with a control subject group maintained on stable statin therapy without concomitant EPA (optionally with matching placebo) for a period of 1 to about 12 weeks, no serum LDL-C increase, no statistically significant serum LDL-C increase, no statistically significant serum LDL-C decrease, or the subject group is statistically non-inferior to the control subject group (statin plus optional placebo) in regard to serum LDL-C elevation), wherein the control subject group also has mean baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl.

In another embodiment, the invention provides a method of lowering triglycerides in subject group on stable statin therapy and having mean baseline fasting triglyceride level of about 200 mg/dl to 499 mg/dl, the method comprising administering to members of the subject group a pharmaceutical composition comprising about 1 g to about 4 g of EPA, wherein upon administering the composition to the members of the subject group daily for a period of 1 to about 12 weeks the subject group exhibits (a) at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, at least 75% lower mean fasting triglycerides and (b) at least 5%, at least 10%, at least 15%, at least 20%, at least 25%, at least 30%, at least 35%, at least 40%, at least 45% or at least 50% lower mean serum LDL-C levels by comparison with a control subject group maintained on stable statin therapy without concomitant EPA (optionally with matching placebo) for a period of 1 to about 12 weeks, no serum LDL-C increase, no statistically significant serum LDL-C increase, no statistically significant serum LDL-C decrease, or the subject group is statistically non-inferior to the control subject group (statin plus optional placebo) in regard to serum LDL-C elevation), wherein the control subject group also has mean baseline fasting triglycerides of about 200 mg/dl to about 500 mg/dl.

In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline absolute plasma level of free total fatty acid (or mean thereof) not greater than about 300 nmol/ml,

6

not greater than about 250 nmol/ml, not greater than about 200 nmol/ml, not greater than about 150 nmol/ml, not greater than about 100 nmol/ml, or not greater than about 50 nmol/ml.

In another embodiment, the subject or subject group being treated has a baseline triglyceride level (or median baseline triglyceride level in the case of a subject group), fed or fasting, of at least about 300 mg/dl, at least about 400 mg/dl, at least about 500 mg/dl, at least about 600 mg/dl, at least about 700 mg/dl, at least about 800 mg/dl, at least about 900 mg/dl, at least about 1000 mg/dl, at least about 1100 mg/dl, at least about 1200 mg/dl, at least about 1300 mg/dl, at least about 1400 mg/dl, or at least about 1500 mg/dl, for example about 200 mg/dl to 499 mg/dl, about 400 mg/dl to about 2500 mg/dl, about 450 mg/dl to about 2000 mg/dl or about 500 mg/dl to about 1500 mg/dl.

In one embodiment, the subject or subject group being treated in accordance with methods of the invention has previously been treated with Lovaza® and has experienced an increase in, or no decrease in, LDL-C levels and/or non-HDL-C levels. In one such embodiment, Lovaza® therapy is discontinued and replaced by a method of the present invention.

In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline absolute plasma level of free EPA (or mean thereof in the case of a subject group) not greater than about 0.70 nmol/ml, not greater than about 0.65 nmol/ml, not greater than about 0.60 nmol/ml, not greater than about 0.55 nmol/ml, not greater than about 0.50 nmol/ml, not greater than about 0.45 nmol/ml, or not greater than about 0.40 nmol/ml. In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a baseline fasting plasma level (or mean thereof) of free EPA, expressed as a percentage of total free fatty acid, of not more than about 3%, not more than about 2.5%, not more than about 2%, not more than about 1.5%, not more than about 1%, not more than about 0.75%, not more than about 0.5%, not more than about 0.25%, not more than about 0.2% or not more than about 0.15%. In one such embodiment, free plasma EPA and/or total fatty acid levels are determined prior to initiating therapy.

In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline absolute plasma level of total fatty acid (or mean thereof) not greater than about 250 nmol/ml, not greater than about 200 nmol/ml, not greater than about 150 nmol/ml, not greater than about 100 nmol/ml, or not greater than about 50 nmol/ml.

In another embodiment, the subject or subject group being treated in accordance with methods of the invention exhibits a fasting baseline plasma, serum or red blood cell membrane EPA level not greater than about 70 µg/ml, not greater than about 60 µg/ml, not greater than about 50 µg/ml, not greater than about 40 µg/ml, not greater than about 30 µg/ml, or not greater than about 25 µg/ml.

In another embodiment, methods of the present invention comprise a step of measuring the subject's (or subject group's mean) baseline lipid profile prior to initiating therapy. In another embodiment, methods of the invention comprise the step of identifying a subject or subject group having one or more of the following: baseline non-HDL-C value of about 200 mg/dl to about 400 mg/dl, for example at least about 210 mg/dl, at least about 220 mg/dl, at least about 230 mg/dl, at least about 240 mg/dl, at least about 250 mg/dl, at least about 260 mg/dl, at least about 270 mg/dl, at least about 280 mg/dl, at least about 290 mg/dl, or at least

about 300 mg/dl; baseline total cholesterol value of about 250 mg/dl to about 400 mg/dl, for example at least about 260 mg/dl, at least about 270 mg/dl, at least about 280 mg/dl or at least about 290 mg/dl; baseline VLDL-C value of about 140 mg/dl to about 200 mg/dl, for example at least about 150 mg/dl, at least about 160 mg/dl, at least about 170 mg/dl, at least about 180 mg/dl or at least about 190 mg/dl; baseline HDL-C value of about 10 to about 60 mg/dl, for example not more than about 40 mg/dl, not more than about 35 mg/dl, not more than about 30 mg/dl, not more than about 25 mg/dl, not more than about 20 mg/dl, or not more than about 15 mg/dl; and/or baseline LDL-C value of about 50 to about 300 mg/dl, for example not less than about 100 mg/dl, not less than about 90 mg/dl, not less than about 80 mg/dl, not less than about 70 mg/dl, not less than about 60 mg/dl or not less than about 50 mg/dl.

In a related embodiment, upon treatment in accordance with the present invention, for example over a period of about 1 to about 200 weeks, about 1 to about 100 weeks, about 1 to about 80 weeks, about 1 to about 50 weeks, about 1 to about 40 weeks, about 1 to about 20 weeks, about 1 to about 15 weeks, about 1 to about 12 weeks, about 1 to about 10 weeks, about 1 to about 5 weeks, about 1 to about 2 weeks or about 1 week, the subject or subject group exhibits one or more of the following outcomes:

- (a) reduced triglyceride levels compared to baseline or placebo control;
- (b) reduced Apo B levels compared to baseline or placebo control;
- (c) increased HDL-C levels compared to baseline or placebo control;
- (d) no increase in LDL-C levels compared to baseline or placebo control;
- (e) a reduction in LDL-C levels compared to baseline or placebo control;
- (f) a reduction in non-HDL-C levels compared to baseline or placebo control;
- (g) a reduction in VLDL levels compared to baseline or placebo control;
- (h) an increase in apo A-I levels compared to baseline or placebo control;
- (i) an increase in apo A-I/apo B ratio compared to baseline or placebo control;
- (j) a reduction in lipoprotein A levels compared to baseline or placebo control;
- (k) a reduction in LDL particle number compared to baseline or placebo control;
- (l) an increase in LDL size compared to baseline or placebo control;
- (m) a reduction in remnant-like particle cholesterol compared to baseline or placebo control;
- (n) a reduction in oxidized LDL compared to baseline or placebo control;
- (o) no change or a reduction in fasting plasma glucose (FPG) compared to baseline or placebo control;
- (p) a reduction in hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) compared to baseline or placebo control;
- (q) a reduction in homeostasis model insulin resistance compared to baseline or placebo control;
- (r) a reduction in lipoprotein associated phospholipase A2 compared to baseline or placebo control;
- (s) a reduction in intracellular adhesion molecule-1 compared to baseline or placebo control;
- (t) a reduction in interleukin-6 compared to baseline or placebo control;
- (u) a reduction in plasminogen activator inhibitor-1 compared to baseline or placebo control;

(v) a reduction in high sensitivity C-reactive protein (hsCRP) compared to baseline or placebo control;

(w) an increase in serum or plasma EPA compared to baseline;

(x) an increase in red blood cell (RBC) membrane EPA compared to baseline or placebo control;

(y) a reduction or increase in one or more of serum phospholipid and/or red blood cell content of docosahexaenoic acid (DHA), docosapentaenoic acid (DPA), arachidonic acid (AA), palmitic acid (PA), stearidonic acid (SA) or oleic acid (OA) compared to baseline or placebo control;

(z) a reduction in one or more protein components of VLDL such as apolipoprotein C-III (hereinafter "APOC3"; also referred to as APOCIII or HALP2) compared to baseline or placebo control;

(aa) no significant increase in, or a reduction in HDL-APOC3, compared to baseline or placebo control;

(bb) no significant increase in, or a reduction in non-HDL-APOC3, compared to baseline or placebo control; and/or

(cc) no substantial increase in, or a reduction or a significant reduction in RLP-C, compared to baseline or placebo control.

In one embodiment, upon administering a composition of the invention to a subject, the subject exhibits a decrease in triglyceride levels, an increase in the concentrations of EPA and DPA (n-3) in red blood cells, and an increase of the ratio of EPA:arachidonic acid in red blood cells. In a related embodiment the subject exhibits substantially no or no increase in RBC DHA.

In one embodiment, methods of the present invention comprise measuring baseline levels of one or more markers set forth in (a)-(cc) above prior to dosing the subject or subject group. In another embodiment, the methods comprise administering a composition as disclosed herein to the subject after baseline levels of one or more markers set forth in (a)-(cc) are determined, and subsequently taking an additional measurement of said one or more markers.

In another embodiment, upon treatment with a composition of the present invention, for example over a period of about 1 to about 200 weeks, about 1 to about 100 weeks, about 1 to about 80 weeks, about 1 to about 50 weeks, about 1 to about 40 weeks, about 1 to about 20 weeks, about 1 to about 15 weeks, about 1 to about 12 weeks, about 1 to about 10 weeks, about 1 to about 5 weeks, about 1 to about 2 weeks or about 1 week, the subject or subject group exhibits any 2 or more of, any 3 or more of, any 4 or more of, any 5 or more of, any 6 or more of, any 7 or more of, any 8 or more of, any 9 or more of, any 10 or more of, any 11 or more of, any 12 or more of, any 13 or more of, any 14 or more of, any 15 or more of, any 16 or more of, any 17 or more of, any 18 or more of, any 19 or more of, any 20 or more of, any 21 or more of, any 22 or more of, any 23 or more of, any 24 or more of, any 25 or more of, any 26 or more of, any 27 or more of, any 28 or more of, or all 29 of outcomes (a)-(cc) described immediately above.

In another embodiment, upon treatment with a composition of the present invention, the subject or subject group exhibits one or more of the following outcomes:

- (a) a reduction in triglyceride level of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or placebo control;

(b) a less than 30% increase, less than 20% increase, less than 10% increase, less than 5% increase or no increase in non-HDL-C levels or a reduction in non-HDL-C levels of at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or placebo control;

(c) substantially no change in HDL-C levels, no change in HDL-C levels, or an increase in HDL-C levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or placebo control;

(d) a less than 60% increase, a less than 50% increase, a less than 40% increase, a less than 30% increase, less than 20% increase, less than 10% increase, less than 5% increase or no increase in LDL-C levels or a reduction in LDL-C levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or placebo control;

(e) a decrease in Apo B levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) as compared to baseline or placebo control;

(f) a reduction in VLDL levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(g) an increase in apo A-I levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(h) an increase in apo A-I/apo B ratio of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(i) a reduction in lipoprotein (a) levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline;

(j) a reduction in mean LDL particle number of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(k) an increase in mean LDL particle size of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about

50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(l) a reduction in remnant-like particle cholesterol of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(m) a reduction in oxidized LDL of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(n) substantially no change, no significant change, or a reduction (e.g. in the case of a diabetic subject) in fasting plasma glucose (FPG) of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(o) substantially no change, no significant change or a reduction in hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>) of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% (actual % change or median % change) compared to baseline or placebo control;

(p) a reduction in homeostasis model index insulin resistance of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(q) a reduction in lipoprotein associated phospholipase A2 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(r) a reduction in intracellular adhesion molecule-1 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(s) a reduction in interleukin-6 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(t) a reduction in plasminogen activator inhibitor-1 of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(u) a reduction in high sensitivity C-reactive protein (hsCRP) of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least



about 45%, at least about 50%, or at least about 100% (actual % change or median % change) compared to baseline or placebo control;

(v) an increase in serum, plasma and/or RBC EPA of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 100%, at least about 200% or at least about 400% (actual % change or median % change) compared to baseline or placebo control;

(w) an increase in serum phospholipid and/or red blood cell membrane EPA of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 100%, at least about 200%, or at least about 400% (actual % change or median % change) compared to baseline or placebo control;

(x) a reduction or increase in one or more of serum phospholipid and/or red blood cell DHA, DPA, AA, PA and/or OA of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) compared to baseline or placebo control;

(y) a reduction in total cholesterol of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55% or at least about 75% (actual % change or median % change) compared to baseline or placebo control;

(z) no significant increase in, or a reduction in one or more protein components of VLDL such as APOC3, of at least about 0.2%, at least about 0.3%, at least about 0.5%, at least about 0.6%, at least about 1%, at least about 5%, at least about 8%, at least about 8.5%, at least about 9%, at least about 9.4%, at least about 10%, at least about 10.1%, at least about 14.3%, at least about 15%, at least about 19%, at least about 19.2%, at least about 20%, at least about 25%, at least about 25.1%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 100%, at least about 105%, at least about 110%, at least about 115%, at least about 120%, at least about 125%, at least about 130%, at least about 135%, at least about 140%, at least about 145%, at least about 150%, at least about 155%, at least about 160%, at least about 165%, at least about 170%, at least about 175%, at least about 180%, at least about 185%, at least about 190%, at least about 195%, or at least about 200% (actual % change or median % change) compared to baseline or placebo control;

(aa) no significant increase in, or a reduction in HDL-APOC3, of at least about 0.2%, at least about 0.3%, at least about 0.5%, at least about 0.6%, at least about 1%, at least about 5%, at least about 5.9%, at least about 8%, at least about 9%, at least about 10%, at least about 15%, at least about 20%, at least about 21%, at least about 21.4%, at least about 23%, at least about 23.3%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 100%, at least about 105%, at least about 110%, at least about 115%, at least about 120%, at

least about 125%, at least about 130%, at least about 135%, at least about 140%, at least about 145%, at least about 150%, at least about 155%, at least about 160%, at least about 165%, at least about 170%, at least about 175%, at least about 180%, at least about 185%, at least about 190%, at least about 195%, or at least about 200% (actual % change or median % change) compared to baseline or placebo control; and/or

(bb) no significant increase in, or a reduction in non-HDL-APOC3, of at least about 0.2%, at least about 0.3%, at least about 0.5%, at least about 0.6%, at least about 1%, at least about 1.4%, at least about 5%, at least about 5.4%, at least about 8%, at least about 9%, at least about 10%, at least about 15%, at least about 20%, at least about 22%, at least about 22.5%, at least about 25%, at least about 30%, at least about 31%, at least about 31.8%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, at least about 55%, at least about 65%, at least about 70%, at least about 75%, at least about 80%, at least about 85%, at least about 90%, at least about 95%, at least about 100%, at least about 105%, at least about 110%, at least about 115%, at least about 120%, at least about 125%, at least about 130%, at least about 135%, at least about 140%, at least about 145%, at least about 150%, at least about 155%, at least about 160%, at least about 165%, at least about 170%, at least about 175%, at least about 180%, at least about 185%, at least about 190%, at least about 195%, or at least about 200% (actual % change or median % change) compared to baseline or placebo control; and/or

(cc) no significant increase in, or a reduction in RLP-C of at least about 5%, at least about 10%, at least about 15%, at least about 16%, about 16%, 16.1%, at least about 20%, at least about 24%, about 24%, 24.0%, at least about 25%, 25.8%, about 26%, 29.8%, about 30%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% (actual % change or median % change) compared to baseline or placebo control.

In one embodiment, methods of the present invention comprise measuring baseline levels of one or more markers set forth in (a)-(bb) prior to dosing the subject or subject group. In another embodiment, the methods comprise administering a composition as disclosed herein to the subject after baseline levels of one or more markers set forth in (a)-(bb) are determined, and subsequently taking a second measurement of the one or more markers as measured at baseline for comparison thereto.

In another embodiment, upon treatment with a composition of the present invention, for example over a period of about 1 to about 200 weeks, about 1 to about 100 weeks, about 1 to about 80 weeks, about 1 to about 50 weeks, about 1 to about 40 weeks, about 1 to about 20 weeks, about 1 to about 15 weeks, about 1 to about 12 weeks, about 1 to about 10 weeks, about 1 to about 5 weeks, about 1 to about 2 weeks or about 1 week, the subject or subject group exhibits any 2 or more of, any 3 or more of, any 4 or more of, any 5 or more of, any 6 or more of, any 7 or more of, any 8 or more of, any 9 or more of, any 10 or more of, any 11 or more of, any 12 or more of, any 13 or more of, any 14 or more of, any 15 or more of, any 16 or more of, any 17 or more of, any 18 or more of, any 19 or more of, any 20 or more of, any 21 or more of, any 22 or more of, any 23 or more of, any 24 or more of, any 25 or more of, any 26 or more of, any 27 or more of, any 28 or more of, or all 29 of outcomes (a)-(cc) described immediately above.

Parameters (a)-(cc) can be measured in accordance with any clinically acceptable methodology. For example, triglycerides, total cholesterol, HDL-C and fasting blood sugar

can be sample from serum and analyzed using standard photometry techniques. VLDL-TG, LDL-C and VLDL-C can be calculated or determined using serum lipoprotein fractionation by preparative ultracentrifugation and subsequent quantitative analysis by refractometry or by analytic ultracentrifugal methodology. Apo A1, Apo B and hsCRP can be determined from serum using standard nephelometry techniques. Lipoprotein (a) can be determined from serum using standard turbidimetric immunoassay techniques. LDL particle number and particle size can be determined using nuclear magnetic resonance (NMR) spectrometry. Remnants lipoproteins and LDL-phospholipase A2 can be determined from EDTA plasma or serum and serum, respectively, using enzymatic immunoseparation techniques. Oxidized LDL, intercellular adhesion molecule-1 and interleukin-6 levels can be determined from serum using standard enzyme immunoassay techniques. APOC3 levels can be determined by known quantitative methods including, for example, antibody-based assays such as ELISA. These techniques are described in detail in standard textbooks, for example Tietz Fundamentals of Clinical Chemistry, 6<sup>th</sup> Ed. (Burtis, Ashwood and Bortler Eds.), WB Saunders Company.

In one embodiment, the present invention provides a method of reducing an RLP-C level of a subject having baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl and on statin therapy (e.g., stable statin therapy), the method comprising administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of EPA per day, wherein upon administering the composition to the subject daily for a period of 1 to about 12 weeks, the subject exhibits a decrease in RLP-C of at least about 5% compared to a control subject on stable statin therapy without EPA, wherein the control subject also has baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl. In some embodiments, the decrease in RLP-C level is at least about 5%, at least about 10%, at least about 15%, at least about 16%, about 16%, 16.1%, at least about 20%, at least about 24%, about 24%, 24.0%, at least about 25%, 25.8%, about 26%, 29.8%, about 30%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% compared to baseline or placebo. In some embodiments, the subject is administered about 2 g per day of EPA. In some embodiments, the subject is administered about 3.8 g to about 4 g per day of EPA. In any of the methods described herein, the EPA can be ethyl eicosapentaenoate.

In one embodiment, the present invention provides a method of reducing an RLP-C level of a subject having baseline fasting triglycerides of about 500 mg/dl to about 1500 mg/dl, the method comprising administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of EPA per day, wherein upon administering the composition to the subject daily for a period of 1 to about 12 weeks, the subject exhibits a decrease in RLP-C in plasma and/or RBCs of at least about 5% compared to a control subject who has not received about 1 g to about 4 g of EPA per day, wherein the control subject also has baseline fasting triglycerides of about 500 mg/dl to about 1500 mg/dl. In some embodiments, the decrease in RLP-C level is at least about 5%, at least about 10%, at least about 15%, at least about 16%, about 16%, 16.1%, at least about 20%, at least about 24%, about 24%, 24.0%, at least about 25%, 25.8%, about 26%, 29.8%, about 30%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% compared to baseline or placebo. In some embodiments, the subject is administered about 2 g per day of EPA. In some embodiments, the subject is administered

about 3.8 g to about 4 g per day of EPA. In any of the methods described herein, the EPA can be ethyl eicosapentaenoate.

In one embodiment, subjects fast for up to 12 hours prior to blood sample collection, for example about 10 hours.

In another embodiment, the present invention provides a method of treating or preventing primary hypercholesterolemia and/or mixed dyslipidemia (Fredrickson Types IIa and IIb) in a patient in need thereof, comprising administering to the patient one or more compositions as disclosed herein. In a related embodiment, the present invention provides a method of reducing triglyceride levels in a subject or subjects when treatment with a statin or niacin extended-release monotherapy is considered inadequate (Frederickson type IV hyperlipidemia).

In another embodiment, the present invention provides a method of treating or preventing risk of recurrent nonfatal myocardial infarction in a patient with a history of myocardial infarction, comprising administering to the patient one or more compositions as disclosed herein.

In another embodiment, the present invention provides a method of slowing progression of or promoting regression of atherosclerotic disease in a patient in need thereof, comprising administering to a subject in need thereof one or more compositions as disclosed herein.

In another embodiment, the present invention provides a method of treating or preventing very high serum triglyceride levels (e.g. Types IV and V hyperlipidemia) in a patient in need thereof, comprising administering to the patient one or more compositions as disclosed herein.

In another embodiment, the present invention provides a method of treating subjects having very high serum triglyceride levels (e.g. greater than 1000 mg/dl or greater than 2000 mg/dl) and that are at risk of developing pancreatitis, comprising administering to the patient one or more compositions as disclosed herein.

In one embodiment, a composition of the invention is administered to a subject in an amount sufficient to provide a daily dose of eicosapentaenoic acid of about 1 mg to about 10,000 mg, 25 about 5000 mg, about 50 to about 3000 mg, about 75 mg to about 2500 mg, about 2 g to about 4 g, about 3.8 g to about 4 g, or about 100 mg to about 1000 mg, for example about 1 mg, about 2 mg, about 3 mg, about 4 mg, about 5 mg, about 6 mg, about 7 mg, about 8 mg, about 9 mg, about 10 mg, about 15 mg, about 20 mg, about 25 mg, about 30 mg, about 35 mg, about 40 mg, about 45 mg, about 50 mg, about 55 mg, about 60 mg, about 65 mg, about 70 mg, about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1100 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg,

about 1975 mg, about 2000 mg, about 2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225 mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about 2400 mg, about 2425 mg, about 2450 mg, about 2475 mg, about 2500 mg, about 2750 mg, about 3000 mg, about 3250 mg, about 3500 mg, about 3750 mg, about 3800 mg, about 4000 mg, about 4250 mg, about 4500 mg, about 4750 mg, about 5000 mg, about 5500 mg, about 6000 mg, about 6500 mg, about 7000 mg, about 7500 mg, about 8000 mg, about 8500 mg, about 9000 mg, about 9500 mg, or about 10,000 mg.

In another embodiment, any of the methods disclosed herein are used in treatment or prevention of a subject or subjects that consume a traditional Western diet. In one embodiment, the methods of the invention include a step of identifying a subject as a Western diet consumer or prudent diet consumer and then treating the subject if the subject is deemed a Western diet consumer. The term "Western diet" herein refers generally to a typical diet consisting of, by percentage of total calories, about 45% to about 50% carbohydrate, about 35% to about 40% fat, and about 10% to about 15% protein. A Western diet may alternately or additionally be characterized by relatively high intakes of red and processed meats, sweets, refined grains, and deserts, for example more than 50%, more than 60% or more or 70% of total calories come from these sources.

In one embodiment, a composition for use in methods of the invention comprises eicosapentaenoic acid, or a pharmaceutically acceptable ester, derivative, conjugate or salt thereof, or mixtures of any of the foregoing, collectively referred to herein as "EPA." The term "pharmaceutically acceptable" in the present context means that the substance in question does not produce unacceptable toxicity to the subject or interaction with other components of the composition.

In one embodiment, the EPA comprises all-cis eicosa-5, 8,11,14,17-pentaenoic acid. In another embodiment, the EPA comprises an eicosapentaenoic acid ester. In another embodiment, the EPA comprises a C<sub>1</sub>-C<sub>5</sub> alkyl ester of eicosapentaenoic acid. In another embodiment, the EPA comprises eicosapentaenoic acid ethyl ester, eicosapentaenoic acid methyl ester, eicosapentaenoic acid propyl ester, or eicosapentaenoic acid butyl ester. In another embodiment, the EPA comprises In one embodiment, the EPA comprises all-cis eicosa-5,8,11,14,17-pentaenoic acid ethyl ester.

In another embodiment, the EPA is in the form of ethyl-EPA, lithium EPA, mono-, di- or triglyceride EPA or any other ester or salt of EPA, or the free acid form of EPA. The EPA may also be in the form of a 2-substituted derivative or other derivative which slows down its rate of oxidation but does not otherwise change its biological action to any substantial degree.

In another embodiment, EPA is present in a composition useful in accordance with methods of the invention in an amount of about 50 mg to about 5000 mg, about 75 mg to about 2500 mg, or about 100 mg to about 1000 mg, for example about 50 mg, about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about

1025 mg, about 1050 mg, about 1075 mg, about 1100 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg, about 1975 mg, about 2000 mg, about 2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225 mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about 2400 mg, about 2425 mg, about 2450 mg, about 2475 mg, about 2500 mg, 2525 mg, about 2550 mg, about 2575 mg, about 2600 mg, about 2625 mg, about 2650 mg, about 2675 mg, about 2700 mg, about 2725 mg, about 2750 mg, about 2775 mg, about 2800 mg, about 2825 mg, about 2850 mg, about 2875 mg, about 2900 mg, about 2925 mg, about 2950 mg, about 2975 mg, about 3000 mg, about 3025 mg, about 3050 mg, about 3075 mg, about 3100 mg, about 3125 mg, about 3150 mg, about 3175 mg, about 3200 mg, about 3225 mg, about 3250 mg, about 3275 mg, about 3300 mg, about 3325 mg, about 3350 mg, about 3375 mg, about 3400 mg, about 3425 mg, about 3450 mg, about 3475 mg, about 3500 mg, about 3525 mg, about 3550 mg, about 3575 mg, about 3600 mg, about 3625 mg, about 3650 mg, about 3675 mg, about 3700 mg, about 3725 mg, about 3750 mg, about 3775 mg, about 3800 mg, about 3825 mg, about 3850 mg, about 3875 mg, about 3900 mg, about 3925 mg, about 3950 mg, about 3975 mg, about 4000 mg, about 4025 mg, about 4050 mg, about 4075 mg, about 4100 mg, about 4125 mg, about 4150 mg, about 4175 mg, about 4200 mg, about 4225 mg, about 4250 mg, about 4275 mg, about 4300 mg, about 4325 mg, about 4350 mg, about 4375 mg, about 4400 mg, about 4425 mg, about 4450 mg, about 4475 mg, about 4500 mg, about 4525 mg, about 4550 mg, about 4575 mg, about 4600 mg, about 4625 mg, about 4650 mg, about 4675 mg, about 4700 mg, about 4725 mg, about 4750 mg, about 4775 mg, about 4800 mg, about 4825 mg, about 4850 mg, about 4875 mg, about 4900 mg, about 4925 mg, about 4950 mg, about 4975 mg, or about 5000 mg.

In another embodiment, a composition useful in accordance with the invention contains not more than about 10%, not more than about 9%, not more than about 8%, not more than about 7%, not more than about 6%, not more than about 5%, not more than about 4%, not more than about 3%, not more than about 2%, not more than about 1%, or not more than about 0.5%, by weight, docosahexaenoic acid (DHA), if any. In another embodiment, a composition of the invention contains substantially no docosahexaenoic acid. In still another embodiment, a composition useful in the present invention contains no docosahexaenoic acid and/or derivative thereof.

In another embodiment, EPA comprises at least 70%, at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, by weight, of all fatty acids present in a composition that is useful in methods of the present invention.

In one embodiment, a composition of the invention comprises ultra-pure EPA. The term "ultra-pure" as used herein with respect to EPA refers to a composition comprising at least 95% by weight EPA (as the term "EPA" is defined and exemplified herein). Ultra-pure EPA comprises at least 96%

by weight EPA, at least 97% by weight EPA, or at least 98% by weight EPA, wherein the EPA is any form of EPA as set forth herein.

In another embodiment, a composition useful in accordance with methods of the invention contains less than 10%, less than 9%, less than 8%, less than 7%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, less than 1%, less than 0.5% or less than 0.25%, by weight of the total composition or by weight of the total fatty acid content, of any fatty acid other than EPA. Illustrative examples of a "fatty acid other than EPA" include linolenic acid (LA), arachidonic acid (AA), docosahexaenoic acid (DHA), alpha-linolenic acid (ALA), stearadonic acid (STA), eicosatrienoic acid (ETA) and/or docosapentaenoic acid (DPA). In another embodiment, a composition useful in accordance with methods of the invention contains about 0.1% to about 4%, about 0.5% to about 3%, or about 1% to about 2%, by weight, of total fatty acids other than EPA and/or DHA.

In another embodiment, a composition useful in accordance with the invention has one or more of the following features: (a) eicosapentaenoic acid ethyl ester represents at least about 96%, at least about 97%, or at least about 98%, by weight, of all fatty acids present in the composition; (b) the composition contains not more than about 4%, not more than about 3%, or not more than about 2%, by weight, of total fatty acids other than eicosapentaenoic acid ethyl ester; (c) the composition contains not more than about 0.6%, not more than about 0.5%, or not more than about 0.4% of any individual fatty acid other than eicosapentaenoic acid ethyl ester; (d) the composition has a refractive index (20° C.) of about 1 to about 2, about 1.2 to about 1.8 or about 1.4 to about 1.5; (e) the composition has a specific gravity (20° C.) of about 0.8 to about 1.0, about 0.85 to about 0.95 or about 0.9 to about 0.92; (e) the composition contains not more than about 20 ppm, not more than about 15 ppm or not more than about 10 ppm heavy metals, (f) the composition contains not more than about 5 ppm, not more than about 4 ppm, not more than about 3 ppm, or not more than about 2 ppm arsenic, and/or (g) the composition has a peroxide value of not more than about 5 meq/kg, not more than about 4 meq/kg, not more than about 3 meq/kg, or not more than about 2 meq/kg.

In another embodiment, a composition useful in accordance with the invention comprises, consists of or consists essentially of at least 95% by weight ethyl eicosapentaenoate (EPA-E), about 0.2% to about 0.5% by weight ethyl octadecatetraenoate (ODTA-E), about 0.05% to about 0.25% by weight ethyl nonadecapentaenoate (NDPA-E), about 0.2% to about 0.45% by weight ethyl arachidonate (AA-E), about 0.3% to about 0.5% by weight ethyl eicosatetraenoate (ETA-E), and about 0.05% to about 0.32% ethyl heneicosapentaenoate (HPA-E). In another embodiment, the composition is present in a capsule shell.

In another embodiment, compositions useful in accordance with the invention comprise, consist essential of, or consist of at least 95%, 96% or 97%, by weight, ethyl eicosapentaenoate, about 0.2% to about 0.5% by weight ethyl octadecatetraenoate, about 0.05% to about 0.25% by weight ethyl nonadecapentaenoate, about 0.2% to about 0.45% by weight ethyl arachidonate, about 0.3% to about 0.5% by weight ethyl eicosatetraenoate, and about 0.05% to about 0.32% ethyl heneicosapentaenoate. Optionally, the composition contains not more than about 0.06%, about 0.05%, or about 0.04%, by weight, DHA or derivative thereof such as ethyl-DHA. In one embodiment the composition contains substantially no or no amount of DHA or derivative thereof such as ethyl-DHA. The composition

further optionally comprises one or more antioxidants (e.g. tocopherol) or other impurities in an amount of not more than about 0.5% or not more than 0.05%. In another embodiment, the composition comprises about 0.05% to about 0.4%, for example about 0.2% by weight tocopherol. In another embodiment, about 500 mg to about 1 g of the composition is provided in a capsule shell.

In another embodiment, compositions useful in accordance with the invention comprise, consist essential of, or consist of at least 96% by weight ethyl eicosapentaenoate, about 0.22% to about 0.4% by weight ethyl octadecatetraenoate, about 0.075% to about 0.20% by weight ethyl nonadecapentaenoate, about 0.25% to about 0.40% by weight ethyl arachidonate, about 0.3% to about 0.4% by weight ethyl eicosatetraenoate and about 0.075% to about 0.25% ethyl heneicosapentaenoate. Optionally, the composition contains not more than about 0.06%, about 0.05%, or about 0.04%, by weight, DHA or derivative thereof such as ethyl-DHA. In one embodiment the composition contains substantially no or no amount of DHA or derivative thereof such as ethyl-DHA. The composition further optionally comprises one or more antioxidants (e.g. tocopherol) or other impurities in an amount of not more than about 0.5% or not more than 0.05%. In another embodiment, the composition comprises about 0.05% to about 0.4%, for example about 0.2% by weight tocopherol. In another embodiment, the invention provides a dosage form comprising about 500 mg to about 1 g of the foregoing composition in a capsule shell. In one embodiment, the dosage form is a gel or liquid capsule and is packaged in blister packages of about 1 to about 20 capsules per sheet.

In another embodiment, compositions useful in accordance with the invention comprise, consist essential of, or consist of at least 96%, 97% or 98%, by weight, ethyl eicosapentaenoate, about 0.25% to about 0.38% by weight ethyl octadecatetraenoate, about 0.10% to about 0.15% by weight ethyl nonadecapentaenoate, about 0.25% to about 0.35% by weight ethyl arachidonate, about 0.31% to about 0.38% by weight ethyl eicosatetraenoate, and about 0.08% to about 0.20% ethyl heneicosapentaenoate. Optionally, the composition contains not more than about 0.06%, about 0.05%, or about 0.04%, by weight, DHA or derivative thereof such as ethyl-DHA. In one embodiment the composition contains substantially no or no amount of DHA or derivative thereof such as ethyl-DHA. The composition further optionally comprises one or more antioxidants (e.g. tocopherol) or other impurities in an amount of not more than about 0.5% or not more than 0.05%. In another embodiment, the composition comprises about 0.05% to about 0.4%, for example about 0.2% by weight tocopherol. In another embodiment, the invention provides a dosage form comprising about 500 mg to about 1 g of the foregoing composition in a capsule shell.

In another embodiment, a composition as described herein is administered to a subject once or twice per day. In another embodiment, 1, 2, 3 or 4 capsules, each containing about 1 g of a composition as described herein, are administered to a subject daily. In another embodiment, 1 or 2 capsules, each containing about 1 g of a composition as described herein, are administered to the subject in the morning, for example between about 5 am and about 11 am, and 1 or 2 capsules, each containing about 1 g of a composition as described herein, are administered to the subject in the evening, for example between about 5 pm and about 11 pm.

In one embodiment, a subject being treated in accordance with methods of the invention is not otherwise on lipid-altering therapy, for example statin, fibrate, niacin and/or ezetimibe therapy.

In another embodiment, compositions useful in accordance with methods of the invention are orally deliverable. The terms “orally deliverable” or “oral administration” herein include any form of delivery of a therapeutic agent or a composition thereof to a subject wherein the agent or composition is placed in the mouth of the subject, whether or not the agent or composition is swallowed. Thus “oral administration” includes buccal and sublingual as well as esophageal administration. In one embodiment, the composition is present in a capsule, for example a soft gelatin capsule.

A composition for use in accordance with the invention can be formulated as one or more dosage units. The terms “dose unit” and “dosage unit” herein refer to a portion of a pharmaceutical composition that contains an amount of a therapeutic agent suitable for a single administration to provide a therapeutic effect. Such dosage units may be administered one to a plurality (i.e. 1 to about 10, 1 to 8, 1 to 6, 1 to 4 or 1 to 2) of times per day, or as many times as needed to elicit a therapeutic response.

In another embodiment, the invention provides use of any composition described herein for treating moderate to severe hypertriglyceridemia in a subject in need thereof, comprising: providing a subject having a fasting baseline triglyceride level of about 500 mg/dl to about 1500 mg/dl and administering to the subject a pharmaceutical composition as described herein. In one embodiment, the composition comprises about 1 g to about 4 g of eicosapentaenoic acid ethyl ester, wherein the composition contains substantially no docosahexaenoic acid.

In one embodiment, compositions of the invention, upon storage in a closed container maintained at room temperature, refrigerated (e.g. about 5 to about 5-10° C.) temperature, or frozen for a period of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months, exhibit at least about 90%, at least about 95%, at least about 97.5%, or at least about 99% of the active ingredient(s) originally present therein.

In one embodiment, the invention provides use of a composition as described herein in manufacture of a medicament for treatment of any of a cardiovascular-related disease. In another embodiment, the subject is diabetic.

In one embodiment, a composition as set forth herein is packaged together with instructions for using the composition to treat a cardiovascular disorder.

#### EXAMPLES

##### Example 1: Reduction of RLP-C Levels in Subjects with Baseline Fasting Triglyceride Levels of $\geq 500$ Mg/Dl

A multi-center, placebo-controlled randomized, double-blind, 12-week study with an open-label extension was performed to evaluate the efficacy and safety of AMR101 in patients with fasting triglyceride levels  $\geq 500$  mg/dL. The primary objective of the study was to determine the efficacy of  $>96\%$ , by weight of all fatty acids present, ethyl-EPA (“AMR101”) 2 g daily and 4 g daily, compared to placebo, in lowering fasting TG levels in patients with fasting TG levels  $\geq 500$  mg/dL and  $\leq 1500$  mg/dL ( $\geq 5.65$  mmol/L and  $\leq 16.94$  mmol/L).

The secondary objectives of this study were the following:

1. To determine the safety and tolerability of AMR101 2 g daily and 4 g daily;
2. To determine the effect of AMR101 on lipid and apolipoprotein profiles;
3. To determine the effect of AMR101 on low-density lipoprotein (LDL) particle number and size;
4. To determine the effect of AMR101 on oxidized LDL;
5. To determine the effect of AMR101 on fasting plasma glucose (FPG) and hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>);
6. To determine the effect of AMR101 on insulin resistance;
7. To determine the effect of AMR101 on high-sensitivity C-reactive protein (hsCRP);
8. To determine the effects of AMR101 2 g daily and 4 g daily on the incorporation of fatty acids into red blood cell membranes and into plasma phospholipids;
9. To explore the relationship between baseline fasting TG levels and the reduction in fasting TG levels; and
10. To explore the relationship between an increase in red blood cell membrane eicosapentaenoic acid (EPA) concentrations and the reduction in fasting TG levels.

The population for this study was men and women (women of childbearing potential needed to be on contraception or practice abstinence)  $>18$  years of age with a body mass index  $\leq 45$  kg/m<sup>2</sup> who were not on lipid-altering therapy or were not currently on lipid-altering therapy. Patients currently on statin therapy (with or without ezetimibe) were evaluated by the investigator as to whether this therapy could be safely discontinued at screening, or if it should have been continued. If statin therapy (with or without ezetimibe) was to be continued, dose(s) must have been stable for  $\geq 4$  weeks prior to randomization. Patients taking non-statin, lipid-altering medications (niacin  $>200$  mg/day, fibrates, fish oil, other products containing omega-3 fatty acids, or other herbal products or dietary supplements with potential lipid-altering effects), either alone or in combination with statin therapy (with or without ezetimibe), must have been able to safely discontinue non-statin, lipid-altering therapy at screening.

Approximately 240 patients were randomized at approximately 50 centers in North America, South America, Central America, Europe, India, and South Africa. The study was a 58- to 60-week, Phase 3, multi-center study consisting of 3 study periods: (1) a 6- to 8-week screening period that included a diet and lifestyle stabilization and washout period and a TG qualifying period; (2) a 12-week, double-blind, randomized, placebo-controlled treatment period; and (3) a 40-week, open-label, extension period.

During the screening period and double-blind treatment period, all visits were within  $\pm 3$  days of the scheduled time. During the open-label extension period, all visits were within  $\pm 7$  days of the scheduled time. The screening period included a 4- or 6-week diet and lifestyle stabilization period and washout period followed by a 2-week TG qualifying period.

The screening visit (Visit 1) occurred for all patients at either 6 weeks (for patients not on lipid-altering therapy at screening or for patients who did not need to discontinue their current lipid-altering therapy) or 8 weeks (for patients who required washout of their current lipid-altering therapy at screening) before randomization, as follows:

Patients who did not require a washout: The screening visit will occur at Visit 1 (Week -6). Eligible patients entered a 4-week diet and lifestyle stabilization period. At the screening visit, all patients received counseling regarding the importance of the National Cholesterol Education Program (NCEP) Therapeutic Lifestyle Changes (TLC) diet

and received instructions on how to follow this diet. Patients who required a washout: The screening visit occurred at Visit 1 (Week -8). Eligible patients began a 6-week washout period at the screening visit. Patients received counseling regarding the NCEP TLC diet and received instructions on how to follow this diet. Site personnel contacted patients who did not qualify for participation based on screening laboratory test results to instruct them to resume their prior lipid-altering medications.

At the end of the 4-week diet and lifestyle stabilization period or the 6-week diet and stabilization and washout period, eligible patients entered the 2-week TG qualifying period and had their fasting TG level measured at Visit 2 (Week -2) and Visit 3 (Week -1). Eligible patients must have had an average fasting TG level  $\geq 500$  mg/dL and  $\leq 1500$  mg/dL ( $\geq 5.65$  mmol/L and  $\leq 16.94$  mmol/L) to enter the 12-week double-blind treatment period. The TG level for qualification was based on the average (arithmetic mean) of the Visit 2 (Week -2) and Visit 3 (Week -1) values. If a patient's average TG level from Visit 2 and Visit 3 fell outside the required range for entry into the study, an additional sample for fasting TG measurement was collected 1 week later at Visit 3.1. If a third sample was collected at Visit 3.1, entry into the study was based on the average (arithmetic mean) of the values from Visit 3 and Visit 3.1.

After confirmation of qualifying fasting TG values, eligible patients entered a 12-week, randomized, double-blind treatment period. At Visit 4 (Week 0), patients were randomly assigned to one of the following treatment groups:

- AMR101 2 g daily,
- AMR101 4 g daily, or
- Placebo.

During the double-blind treatment period, patients returned to the site at Visit 5 (Week 4), Visit 6 (Week 11), and Visit 7 (Week 12) for efficacy and safety evaluations.

Patients who completed the 12-week double-blind treatment period were eligible to enter a 40-week, open-label, extension period at Visit 7 (Week 12). All patients received open-label AMR101 4 g daily. From Visit 8 (Week 16) until the end of the study, changes to the lipid-altering regimen were permitted (e.g., initiating or raising the dose of statin or adding non-statin, lipid-altering medications to the regimen), as guided by standard practice and prescribing information. After Visit 8 (Week 16), patients returned to the site every 12 weeks until the last visit at Visit 11 (Week 52).

Eligible patients were randomly assigned at Visit 4 (Week 0) to orally receive AMR101 2 g daily, AMR101 4 g daily, or placebo for the 12-week double-blind treatment period. AMR101 was provided in 1 g liquid-filled, oblong, gelatin capsules. The matching placebo capsule was filled with light liquid paraffin and contained 0 g of AMR101. During the double-blind treatment period, patients took 2 capsules (AMR101 or matching placebo) in the morning and 2 in the evening for a total of 4 capsules per day. Patients in the AMR101 2 g/day treatment group received 1 AMR101 1 g capsule and 1 matching placebo capsule in the morning and in the evening. Patients in the AMR101 4 g/day treatment group received 2 AMR101 1 g capsules in the morning and evening.

Patients in the placebo group received 2 matching placebo capsules in the morning and evening. During the extension period, patients received open-label AMR101 4 g daily. Patients took 2 AMR101 1 g capsules in the morning and 2 in the evening.

The primary efficacy variable for the double-blind treatment period was percent change in TG from baseline to

Week 12 endpoint. The secondary efficacy variables for the double-blind treatment period included the following:

Percent changes in total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), calculated low-density lipoprotein cholesterol (LDL-C), calculated non-high-density lipoprotein cholesterol (non-HDL-C), and very low-density lipoprotein cholesterol (VLDL-C) from baseline to Week 12 endpoint;

Percent change in very low-density lipoprotein TG from baseline to Week 12;

Percent changes in apolipoprotein A-I (apo A-I), apolipoprotein B (apo B), and apo A-I/apo B ratio from baseline to Week 12;

Percent changes in lipoprotein (a) from baseline to Week 12 (selected sites only);

Percent changes in LDL particle number and size, measured by nuclear magnetic resonance, from baseline to Week 12 (selected sites only);

Percent change in remnant-like particle cholesterol from baseline to Week 12 (selected sites only);

Percent change in oxidized LDL from baseline to Week 12 (selected sites only);

Changes in FPG and HbA<sub>1c</sub> from baseline to Week 12;

Change in insulin resistance, as assessed by the homeostasis model index insulin resistance, from baseline to Week 12;

Percent change in lipoprotein associated phospholipase A2 from baseline to Week 12 (selected sites only);

Change in intracellular adhesion molecule-1 from baseline to Week 12 (selected sites only);

Change in interleukin-6 from baseline to Week 12 (selected sites only);

Change in plasminogen activator inhibitor-1 from baseline to Week 12 (selected sites only);

Change in hsCRP from baseline to Week 12 (selected sites only);

Change in serum phospholipid EPA content from baseline to Week 12;

Change in red blood cell membrane EPA content from baseline to Week 12; and

Change in serum phospholipid and red blood cell membrane content in the following fatty acids from baseline to Week 12: docosapentaenoic acid, docosahexaenoic acid, arachidonic acid, palmitic acid, stearic acid, and oleic acid.

The efficacy variable for the open-label extension period was percent change in fasting TG from extension baseline to end of treatment. Safety assessments included adverse events, clinical laboratory measurements (chemistry, hematology, and urinalysis), 12-lead electrocardiograms (ECGs), vital signs, and physical examinations

For TG, TC, HDL-C, calculated LDL-C, calculated non-HDL-C, and VLDL-C, baseline was defined as the average of Visit 4 (Week 0) and the preceding lipid qualifying visit (either Visit 3 [Week -1] or if it occurs, Visit 3.1) measurements. Baseline for all other efficacy parameters was the Visit 4 (Week 0) measurement.

For TC, HDL-C, calculated LDL-C, calculated non-HDL-C, and VLDL-C, Week 12 endpoint was defined as the average of Visit 6 (Week 11) and Visit 7 (Week 12) measurements. Week 12 endpoint for all other efficacy parameters was the Visit 7 (Week 12) measurement.

The primary efficacy analysis was performed using a 2-way analysis of covariance (ANCOVA) model with treatment as a factor and baseline TG value as a covariate. The least-squares mean, standard error, and 2-tailed 95% confidence interval for each treatment group and for each comparison was estimated. The same 2-way ANCOVA model was used for the analysis of secondary efficacy variables.

The primary analysis was repeated for the per-protocol population to confirm the robustness of the results for the intent-to-treat population.

The primary efficacy variable was the percent change in fasting TG levels from baseline to Week 12. A sample size of 69 completed patients per treatment group was expected to provide 90% power to detect a difference of 30% between AMR101 and placebo in percent change from baseline in fasting TG levels, assuming a standard deviation of 45% in TG measurements and a significance level of  $p < 0.01$ . To accommodate a 15% drop-out rate from randomization to completion of the double-blind treatment period, a total of 240 randomized patients was planned (80 patients per treatment group).

Serum RLP-C was measured with an immunoseparation assay (Polymedco, Inc.). Subjects receiving 2 g/day treatment group experienced reduced RLP-C compared to placebo (data not shown). As shown in Table 1, subjects in the 4 g/day treatment group exhibited reductions in RLP-C of 16.1% compared to baseline, which corresponds to a statistically significant median placebo-adjusted reductions in RLP-C of 29.8% ( $p = 0.0041$ ). See also FIG. 1 ("MARINE").

RLP-C changes vs. placebo for subjects not receiving statin therapy are compared to RLP-C changes vs. placebo for subjects also on statin therapy in FIG. 2.

#### Example 2: Reduction of RLP-C Levels in Subjects with Baseline Fasting Triglyceride Levels of 200 mg/dl to $< 500$ mg/dl

A multi-center, placebo-controlled, randomized, double-blind, 12-week study was performed to evaluate the efficacy and safety of  $>96\%$  E-EPA in patients with fasting triglyceride levels  $\geq 200$  mg/dl and  $< 500$  mg/dl despite statin therapy (the mean of two qualifying entry values needed to be  $\geq 185$  mg/dl and at least one of the values needed to be  $\geq 200$  mg/dl). The primary objective of the study was to determine the efficacy of  $>96\%$ , by weight of all fatty acids present, ethyl-EPA ("AMR101") 2 g daily and 4 g daily, compared to placebo, in lowering fasting TG levels in patients with high risk for cardiovascular disease and with fasting TG levels  $\geq 200$  mg/dl and  $< 500$  mg/dl, despite treatment to LDL-C goal on statin therapy.

The secondary objectives of this study were the following:

1. To determine the safety and tolerability of  $>96\%$  E-EPA 2 g daily and 4 g daily;
2. To determine the effect of  $>96\%$  E-EPA on lipid and apolipoprotein profiles including total cholesterol (TC), non-high-density lipoprotein cholesterol (non-HDL-C), low density lipoprotein cholesterol (LDL-C), high density lipoprotein cholesterol (HDL-C), and very high density lipoprotein cholesterol (VHDL-C);

TABLE 1

Reductions in RLP-C.											
RLP-C Concentration										Median Change from Baseline in RLP-C	
AMR101 4 g/day			AMR101 2 g/day			Placebo			vs. Placebo		
Baseline, mg/dl (IQR)	End of Treatment, mg/dl (IQR)	Change, % (IQR)	Baseline, mg/dl (IQR)	End of Treatment, mg/dl (IQR)	Change, % (IQR)	Baseline, mg/dl (IQR)	End of Treatment, mg/dl (IQR)	Change, % (IQR)	AMR101 4 g/day % (P)	AMR101 2 g/day % (P)	
n	76		73			75					
(all subjects)											
ITT	n = 75		n = 70			n = 73					
	45.0 (53.0)	38.0 (45.0)	-16.1 (86.5)	43.0 (37.0)	44.5 (38.0)	5.8 (85.5)	47.0 (58.0)	58.0 (90.0)	14.2 (105.4)	-29.8 (0.0041)	-14.9 (0.1528)
No Statin Use	n = 56		n = 52			n = 56					
	45.0 (55.5)	39.5 (48.5)	-16.6 (85.9)	45.0 (34.5)	44.5 (36.0)	3.0 (67.6)	50.5 (61.0)	54.5 (77.5)	7.0 (107.0)	-21.4 (0.0456)	-8.5 (0.4421)
With Statin Use	n = 19		n = 18			n = 17					
	37.0 (45.0)	33.0 (27.0)	-16.1 (136.4)	31.5 (47.0)	46.5 (50.0)	29.2 (108.0)	40.0 (38.0)	65.0 (124.0)	54.3 (116.9)	-56.8 (0.0198)	-41.6 (0.1511)
Baseline TG $\leq 750$ mg/dl	n = 47		n = 42			n = 41					
	34.0 (30.0)	29.0 (28.0)	-8.8 (100.0)	27.5 (23.0)	32.5 (23.0)	18.2 (85.8)	30.0 (21.0)	33.0 (31.0)	19.0 (95.5)	-26.1 (0.0570)	-7.3 (0.6554)
Baseline TG $> 750$ mg/dl	n = 28		n = 28			n = 32					
	89.5 (109.0)	68.5 (73.0)	-25.4 (59.7)	66.0 (51.5)	65.0 (35.5)	-10.7 (72.5)	90.5 (75.5)	100.5 (89.5)	5.7 (111.1)	-37.5 (0.0196)	-25.4 (0.1016)

Values are presented in Table 1 are medians (IQR or p value). Changes vs placebo are Hodges-Lehmann estimates of median treatment differences, and p values are from the Wilcoxon rank-sum test. IQR=interquartile range. The ITT population includes subjects with RLP-C measurements.

3. To determine the effect of  $>96\%$  E-EPA on lipoprotein associated phospholipase A2 (Lp-PLA<sub>2</sub>) from baseline to week 12;
4. To determine the effect of  $>96\%$  E-EPA on low-density lipoprotein (LDL) particle number and size;

5. To determine the effect of >96% E-EPA on oxidized LDL;
6. To determine the effect of >96% E-EPA on fasting plasma glucose (FPG) and hemoglobin A<sub>1c</sub> (HbA<sub>1c</sub>);
7. To determine the effect of >96% E-EPA on insulin resistance;
8. To determine the effect of >96% E-EPA on high-sensitivity C-reactive protein (hsCRP);
9. To determine the effects of >96% E-EPA 2 g daily and 4 g daily on the incorporation of fatty acids into red blood cell membranes and into plasma phospholipids;
10. To explore the relationship between baseline fasting TG levels and the reduction in fasting TG levels; and
11. To explore the relationship between changes of fatty acid concentrations in plasma and red blood cell membranes, and the reduction in fasting TG levels.

The population for this study was men and women >18 years of age with a body mass index  $\leq 45$  kg/m<sup>2</sup> with fasting TG levels greater than or equal to 200 mg/dl and less than 500 mg/dl and on a stable dose of statin therapy (with or without ezetimibe). The statin was atorvastatin, rosuvastatin or simvastatin. The dose of statin must have been stable for  $\geq 4$  weeks prior to the LDL-C/TG baseline qualifying measurement for randomization. The statin dose was optimized such that the patients are at their LDL-C goal at the LDL-C/TG baseline qualifying measurements. The same statin at the same dose was continued until the study ended.

Patients taking any additional non-statin, lipid-altering medications (niacin >200 mg/day, fibrates, fish oil, other products containing omega-3 fatty acids, or other herbal products or dietary supplements with potential lipid-altering effects), either alone or in combination with statin therapy (with or without ezetimibe), must have been able to safely discontinue non-statin, lipid-altering therapy at screening.

Patients at high risk for CVD, i.e., patients with clinical coronary heart disease (CHD) or clinical CHD risk equivalents (10-year risk >20%) as defined in the National Cholesterol Education Program (NCEP) Adult Treatment Panel III (ATP III) Guidelines were eligible to participate in this study. Those included patients with any of the following criteria: (1) Known CVD, either clinical coronary heart disease (CHD), symptomatic carotid artery disease (CAD), peripheral artery disease (PAD) or abdominal aortic aneurism; or (2) Diabetes Mellitus (Type 1 or 2).

Approximately 702 patients were randomized at approximately 80 centers in the U.S. The study was a 18- to 20-week, Phase 3, multi-center study consisting of 2 study periods: (1) A 6- to 8-week screening period that included a diet and lifestyle stabilization, a non-statin lipid-altering treatment washout, and an LDL-C and TG qualifying period and (2) A 12-week, double-blind, randomized, placebo-controlled treatment period.

During the screening period and double-blind treatment period, all visits were within  $\pm 3$  days of the scheduled time. All patients continued to take the statin product (with or without ezetimibe) at the same dose they were taking at screening throughout their participation in the study.

The 6- to 8-week screening period included a diet and lifestyle stabilization, a non-statin lipid-altering treatment washout, and an LDL-C and TG qualifying period. The screening visit (Visit 1) occurred for all patients at either 6 weeks (for patients on stable statin therapy [with or without ezetimibe] at screening) or 8 weeks (for patients who will require washout of their current non-statin lipid-altering therapy at screening) before randomization, as follows:

Patients who did not require a washout: The screening visit occurred at Visit 1 (Week -6). Eligible patients

entered a 4-week diet and lifestyle stabilization period. At the screening visit, all patients received counseling regarding the importance of the National Cholesterol Education Program (NCEP) Therapeutic Lifestyle Changes (TLC) diet and received basic instructions on how to follow this diet.

Patients who required a washout: The screening visit occurred at Visit 1 (Week -8). Eligible patients began a 6-week washout period at the screening visit (i.e. 6 weeks washout before the first LDL-C/TG qualifying visit). Patients received counseling regarding the NCEP TLC diet and received basic instructions on how to follow this diet. Site personnel contacted patients who did not qualify for participation based on screening laboratory test results to instruct them to resume their prior lipid-altering medications.

At the end of the 4-week diet and lifestyle stabilization period or the 6-week diet and stabilization and washout period, eligible patients entered the 2-week LDL-C and TG qualifying period and had their fasting LDL-C and TG levels measured at Visit 2 (Week -2) and Visit 3 (Week -1). Eligible patients must have had an average fasting LDL-C level  $\geq 40$  mg/dL and <100 mg/dL and an average fasting TG level  $\geq 200$  mg/dL and <500 mg/dL to enter the 12-week double-blind treatment period. The LDL-C and TG levels for qualification were based on the average (arithmetic mean) of the Visit 2 (Week -2) and Visit 3 (Week -1) values. If a patient's average LDL-C and/or TG levels from Visit 2 and Visit 3 fell outside the required range for entry into the study, an additional fasting lipid profile was collected 1 week later at Visit 3.1. If a third sample was collected at Visit 3.1, entry into the study was based on the average (arithmetic mean) of the values from Visit 3 and Visit 3.1.

After confirmation of qualifying fasting LDL-C and TG values, eligible patients entered a 12-week, randomized, double-blind treatment period. At Visit 4 (Week 0), patients were randomly assigned to 1 of the following treatment groups:

- >96% E-EPA 2 g daily,
- >96% E-EPA 4 g daily, or
- Placebo.

226 to 234 patients per treatment group were randomized in this study. Stratification was by type of statin (atorvastatin, rosuvastatin or simvastatin), the presence of diabetes, and gender.

During the double-blind treatment period, patients returned to the site at Visit 5 (Week 4), Visit 6 (Week 11), and Visit 7 (Week 12) for efficacy and safety evaluations.

Eligible patients were randomly assigned at Visit 4 (Week 0) to receive orally >96% E-EPA 2 g daily, >96% E-EPA 4 g daily, or placebo.

>96% E-EPA was provided in 1 g liquid-filled, oblong, gelatin capsules. The matching placebo capsule was filled with light liquid paraffin and contained 0 g of >96% E-EPA. >96% E-EPA capsules were to be taken with food (i.e. with or at the end of a meal).

During the double-blind treatment period, patients were to take 2 capsules (>96% E-EPA or matching placebo) in the morning and 2 capsules in the evening for a total of 4 capsules per day.

Patients in the >96% E-EPA 2 g/day treatment group received 1 >96% E-EPA 1 g capsule and 1 matching placebo capsule in the morning and in the evening.

Patients in the >96% E-EPA 4 g/day treatment group received 2 >96% E-EPA 1 g capsules in the morning and evening.



Patients in the placebo group received 2 matching placebo capsules in the morning and evening.

The primary efficacy variable for the double-blind treatment period was percent change in TG from baseline to Week 12 endpoint. The secondary efficacy variables for the double-blind treatment period included the following:

Percent changes in total cholesterol (TC), high-density lipoprotein cholesterol (HDL-C), LDL-C, calculated non-HDL-C, and very low-density lipoprotein cholesterol (VLDL-C) from baseline to Week 12 endpoint;

Percent change in very low-density lipoprotein TG from baseline to Week 12;

Percent changes in apolipoprotein A-I (apo A-I), apolipoprotein B (apo B), and apo A-I/apo B ratio from baseline to Week 12;

Percent changes in lipoprotein (a) from baseline to Week 12;

Percent changes in LDL particle number and size, measured by nuclear magnetic resonance, from baseline to Week 12;

Percent change in remnant-like particle cholesterol from baseline to Week 12;

Percent change in oxidized LDL from baseline to Week 12;

Changes in FPG and HbA<sub>1c</sub> from baseline to Week 12;

Change in insulin resistance, as assessed by the homeostasis model index insulin resistance, from baseline to Week 12;

Percent change in lipoprotein associated phospholipase A<sub>2</sub> (Lp-PLA<sub>2</sub>) from baseline to Week 12;

Change in intracellular adhesion molecule-1 from baseline to Week 12;

Change in interleukin-2 from baseline to Week 12;

Change in plasminogen activator inhibitor-1 from baseline to Week 12. Note: this parameter will only be collected at sites with proper storage conditions;

Change in hsCRP from baseline to Week 12; and

Change in plasma concentration and red blood cell membrane content of fatty acid from baseline to Week 12 including EPA, docosapentaenoic acid (DPA), docosahexaenoic acid (DHA), arachidonic acid (AA), dihomo- $\gamma$ -linolenic acid (DGLA), the ratio of EPA/AA, ratio of oleic acid/stearic acid (OA/SA), and the ratio of total omega-3 acids over total omega-6 acids.

Safety assessments included adverse events, clinical laboratory measurements (chemistry, hematology, and urinalysis), 12-lead electrocardiograms (ECGs), vital signs, and physical examinations.

For TG, TC, HDL-C, LDL-C, calculated non-HDL-C, and VLDL-C, baseline was defined as the average of Visit 4 (Week 0) and the preceding lipid qualifying visit (either Visit 3 [Week -1] or if it occurs, Visit 3.1) measurements. Baseline for all other efficacy parameters was the Visit 4 (Week 0) measurement.

For TG, TC, HDL-C, LDL-C, calculated non-HDL-C, and VLDL-C, Week 12 endpoint was defined as the average of Visit 6 (Week 11) and Visit 7 (Week 12) measurements.

Week 12 endpoint for all other efficacy parameters were the Visit 7 (Week 12) measurement.

The primary efficacy analysis was performed using a 2-way analysis of covariance (ANCOVA) model with treatment as a factor and baseline TG value as a covariate. The least-squares mean, standard error, and 2-tailed 95% confidence interval for each treatment group and for each comparison were estimated. The same 2-way ANCOVA model was used for the analysis of secondary efficacy variables.

The primary analysis was repeated for the per-protocol population to confirm the robustness of the results for the intent-to-treat population.

Non-inferiority tests for percent change from baseline in LDL-C were performed between >96% E-EPA doses and placebo using a non-inferiority margin of 6% and a significant level at 0.05.

For the following key secondary efficacy parameters, treatment groups were compared using Dunnett's test to control the Type 1 error rate: TC, LDL-C, HDL-C, non-HDL-C, VLDL-C, Lp-PLA<sub>2</sub>, and apo B. For the remaining secondary efficacy parameters, Dunnett's test was used and the ANCOVA output were considered descriptive.

The evaluation of safety was based primarily on the frequency of adverse events, clinical laboratory assessments, vital signs, and 12-lead ECGs. The primary efficacy variable is the percent change in fasting TG levels from baseline to Week 12. A sample size of 194 completed patients per treatment group provided 90.6% power to detect a difference of 15% between >96% E-EPA and placebo in percent change from baseline in fasting TG levels, assuming a standard deviation of 45% in TG measurements and a significance level of  $p < 0.05$ .

Previous data on fasting LDL-C show a difference in percent change from baseline of 2.2%, with a standard deviation of 15%, between study drug and placebo. A sample size of 194 completed patients per treatment group provided 80% power to demonstrate non-inferiority ( $p < 0.05$ , one-sided) of the LDL-C response between >96% E-EPA 4 g daily and placebo, within a 6% margin. To accommodate a 10% drop-out rate from randomization to completion of the double-blind treatment period, a total of 648 randomized patients was planned (216 patients per treatment group); 702 subjects were randomized, as further described below.

## Results

Of the 702 randomized subjects, 687 were in the intent-to-treat ("ITT") population as follows:

Ultra-pure EPA, 4 g/day: 226 subjects

Ultra-pure EPA, 2 g/day: 234 subjects

Placebo: 227 subjects

Lipids were extracted from plasma and red blood cell ("RBC") suspensions and converted into fatty acid methyl esters for analysis using a standard validated gas chromatography/flame ionization detection method. Fatty acid parameters were compared between EPA treatment groups and placebo using an ANCOVA model with treatment, gender, type of statin therapy, and presence of diabetes as factors, and the baseline parameter value as a covariate. LSMs, SEs, and 2-tailed 95% confidence intervals for each treatment group and for each comparison were determined.

Baseline characteristics of the three ITT groups were comparable, with 61.4% of the ITT subjects being male, 96.3% being white, having a mean age of 61.4 years, a weight of 95.7 kg and a BMI of 32.9 kg/m<sup>2</sup>. ITT subjects with incomplete fatty acid data at baseline and/or at 12 weeks were excluded from the analyses.

Serum RLP-C was measured with an immunoseparation assay (Polymedco, Inc.). Subjects receiving 2 g/day treatment group experienced reduced RLP-C compared to placebo (data not shown). As shown in Table 2, subjects in the 4 g/day treatment group exhibited reductions in RLP-C (24.0%) compared to baseline, which corresponds to a statistically significant median placebo-adjusted reduction in RLP-C of 25.8% ( $p = 0.0001$ ). Study subjects receiving moderate- to high-intensity statin therapy exhibited greater RLP-C reductions compared to subjects receiving lower-intensity statin therapy. See also FIG. 1 ("ANCHOR").

TABLE 2

Reductions in RLP-C											
	RLP-C Concentration									Median Change from Baseline in RLP-C	
	AMR101 4 g/day			AMR101 2 g/day			Placebo			vs. Placebo	
	Baseline, mg/dl (IQR)	End of Treatment, mg/dl (IQR)	Change, % (IQR)	Baseline, mg/dl (IQR)	End of Treatment, mg/dl (IQR)	Change, % (IQR)	Baseline, mg/dl (IQR)	End of Treatment, mg/dl (IQR)	Change, % (IQR)	AMR101 4 g/day % (P)	AMR101 2 g/day % (P)
n (all subjects)	226			234			227				
ITT	n = 82	n = 84	n = 86								
	13.5 (6.0)	10.0 (6.0)	-24.0 (45.5)	15.0 (7.0)	11.0 (7.0)	-11.1 (40.0)	14.0 (7.0)	13.0 (9.0)	8.0 (66.9)	-25.8 (0.0001)	-16.7 (0.0153)
Lower Intensity	n = 5	n = 7	n = 6								
Statin	16.0 (3.0)	15.0 (5.0)	-29.2 (33.3)	15.0 (6.0)	11.0 (9.0)	-11.1 (21.1)	15.0 (6.0)	11.0 (24.0)	-19.4 (105.1)	-11.9 (0.5228)	2.7 (0.8303)
Medium and Higher Intensity	n = 77	n = 77	n = 80								
Statin	13.0 (6.0)	10.0 (6.0)	-23.1 (45.5)	15.0 (7.0)	11.0 (7.0)	-11.1 (42.3)	14.0 (7.0)	13.0 (8.5)	8.7 (66.5)	-26.7 (0.0002)	-17.3 (0.0135)
Baseline TG <259 mg/dl	n = 42	n = 40	n = 37								
Baseline	11.0 (4.0)	9.0 (5.0)	-16.0 (47.6)	11.5 (5.0)	11.0 (3.5)	-8.1 (37.4)	11.0 (5.0)	12.0 (7.0)	9.1 (58.2)	-22.2 (0.0263)	-16.4 (0.1092)
Baseline TG ≥259 mg/dl	n = 40	n = 44	n = 49								
Baseline	17.0 (5.5)	11.0 (6.5)	-31.3 (37.6)	18.0 (8.0)	13.5 (7.0)	-17.4 (37.2)	16.0 (4.0)	16.0 (11.0)	7.7 (64.1)	-30.6 (0.0010)	-18.0 (0.0658)

Values are presented in Table 2 are medians (IQR or p value). Changes vs placebo are Hodges-Lehmann estimates of median treatment differences, and p values are from the Wilcoxon rank-sum test. IQR=interquartile range. Lower-intensity statin regimens include: simvastatin 5 to 10 mg; medium-intensity statin regimens include: rosuvastatin 5 to 10 mg, atorvastatin 10 to 20 mg, simvastatin 20 to 40 mg, simvastatin 10 to 20 mg plus ezetimibe 5 to 10 mg; higher-intensity statin regimens include: rosuvastatin 20 to 40 mg, atorvastatin 40 to 80 mg, simvastatin 80 mg, simvastatin 40 to 80 mg plus ezetimibe 5 to 10 mg. Study-wide median represents the median of subjects assessed in the 4 g/day, 2 g/day and placebo groups.

Placebo-adjusted RLP-C changes for subjects receiving medium- and higher-intensity statin therapy are compared to placebo-adjusted RLP-C changes for subjects receiving low-intensity statin therapy in FIG. 3.

What is claimed is:

1. A method of reducing an RLP-C level of a subject on statin therapy, having a baseline RLP-C level of about 10 mg/dl to about 20 mg/dl, and having baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl, the method comprising administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of ethyl eicosapentaenoate per day.

2. The method of claim 1, wherein the RLP-C level is reduced by at least about 15%.

3. The method of claim 1, wherein the RLP-C level is reduced by at least about 20%.

4. The method of claim 1, wherein about 2 g per day of ethyl eicosapentaenoate is administered to the subject.

5. The method of claim 1, wherein about 3.8 g to about 4 g per day of ethyl eicosapentaenoate is administered to the subject.

6. The method of claim 1, wherein the pharmaceutical composition is administered to the subject daily for a period of 1 to about 12 weeks.

7. The method of claim 1, wherein the reduction in RLP-C level is in comparison to control subjects on statin therapy, having baseline RLP-C levels of about 10 mg/dl to about 20 mg/dl, and having baseline fasting triglycerides of about 200 mg/dl to 499 mg/dl who have received the statin but not the pharmaceutical composition.

8. The method of claim 1, wherein the pharmaceutical composition comprises at least about 95%, by weight of all fatty acids present, ethyl eicosapentaenoate.

9. The method of claim 1, wherein the pharmaceutical composition comprises no more than about 10%, by weight of all fatty acids present, docosahexaenoic acid or a derivative thereof, if any.

10. The method of claim 1, wherein the pharmaceutical composition comprises at least about 95%, by weight of all fatty acids present, ethyl eicosapentaenoate and no more than about 2%, by weight of all fatty acids present, docosahexaenoic acid or a derivative thereof, if any.

\* \* \* \* \*