

US011109596B2

(12) **United States Patent**
Leonard et al.

(10) **Patent No.: US 11,109,596 B2**
(45) **Date of Patent: *Sep. 7, 2021**

(54) **MICROORGANISMS FOR THE PRODUCTION OF INSECT PHEROMONES AND RELATED COMPOUNDS**

(71) Applicant: **Provivi, Inc.**, Santa Monica, CA (US)

(72) Inventors: **Effendi Leonard**, Santa Monica, CA (US); **Peter Meinhold**, Topanga, CA (US); **Keith Wampler**, Santa Monica, CA (US); **Pedro Coelho**, Santa Monica, CA (US); **Micah Sheppard**, Santa Monica, CA (US); **Thomas Heel**, Los Angeles, CA (US)

(73) Assignee: **Provivi, Inc.**, Santa Monica, CA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/420,566**

(22) Filed: **May 23, 2019**

(65) **Prior Publication Data**

US 2019/0338317 A1 Nov. 7, 2019

Related U.S. Application Data

(63) Continuation of application No. 15/983,706, filed on May 18, 2018, now Pat. No. 10,308,962, which is a continuation of application No. PCT/US2016/062852, filed on Nov. 18, 2016.

(60) Provisional application No. 62/351,605, filed on Jun. 17, 2016, provisional application No. 62/257,054, filed on Nov. 18, 2015.

(51) **Int. Cl.**

A01N 63/50 (2020.01)
C12P 7/64 (2006.01)
A01N 31/02 (2006.01)
C12N 9/02 (2006.01)
C12P 7/04 (2006.01)

(52) **U.S. Cl.**

CPC **A01N 63/50** (2020.01); **A01N 31/02** (2013.01); **C12N 9/0008** (2013.01); **C12N 9/0071** (2013.01); **C12P 7/04** (2013.01); **C12P 7/6436** (2013.01); **C12P 7/6472** (2013.01); **C12Y 102/0105** (2013.01); **C12Y 114/19001** (2013.01); **C12Y 114/19005** (2013.01)

(58) **Field of Classification Search**

CPC **C12Y 114/19005**; **C12Y 102/0105**; **C12N 9/0008**; **C12N 9/0071**; **C12P 7/04**; **C12P 7/6436**; **C12P 7/6472**; **A01N 31/02**; **A01N 63/10**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,231,947 A	11/1980	Schrock
4,245,131 A	1/1981	Schrock
4,427,595 A	1/1984	Schrock
4,681,956 A	7/1987	Schrock
4,727,215 A	2/1988	Schrock
5,087,710 A	2/1992	Schrock et al.
5,124,491 A	6/1992	Fleckenstein et al.
5,142,073 A	8/1992	Schrock et al.
5,146,033 A	9/1992	Schrock et al.
6,121,473 A	9/2000	Schrock et al.
6,291,742 B1	9/2001	Somerville et al.
6,346,652 B1	2/2002	Schrock et al.
7,169,959 B2	1/2007	Heck et al.
7,700,833 B2	4/2010	Renz et al.
8,110,093 B2	2/2012	Friedman et al.
8,110,670 B2	2/2012	Hu et al.
8,268,599 B2	9/2012	Schirmer et al.
8,283,143 B2	10/2012	Hu et al.
8,535,916 B2	9/2013	Del Cardayre et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CN	1656226 A	8/2005
CN	101490241 A	7/2009

(Continued)

OTHER PUBLICATIONS

Adrio, J.L., "Oleaginous Yeasts: Promising Platforms for the Production of Oleochemicals and Biofuels". Biotechnol Bioeng. (Sep. 2017); 114(9): 1915-1920. Epub May 29, 2017.

Shi and Zhao, "Metabolic Engineering of Oleaginous Yeasts for Production of Fuels and Chemicals". Front Microbiol. (Nov. 8, 2017); 8: 2185. Published online Nov. 8, 2017.

[Author Unknown] "NP 001037017: (11Z)-hexadec-11-enoyl-CoA conjugase [Bombyx mori]." NCBI Protein, Jul. 5, 2004 (Jul. 5, 2004), pp. 1-4. Retrieved from the Internet: <<https://www.ncbi.nlm.nih.gov/protein/162809332>> on Jan. 18, 2017 (Jan. 18, 2017). Entire document.

(Continued)

Primary Examiner — Ganapathirama Raghu

(74) *Attorney, Agent, or Firm* — Cooley LLP

(57) **ABSTRACT**

The present application relates to recombinant microorganisms useful in the biosynthesis of unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates which may be useful as insect pheromones, fragrances, flavors, and polymer intermediates. The C₆-C₂₄ fatty alcohols, aldehydes, and acetates described herein may be used as substrates for metathesis reactions to expand the repertoire of target compounds and pheromones. The application further relates to recombinant microorganisms co-expressing a pheromone pathway and a pathway for the production of a toxic protein, peptide, oligonucleotide, or small molecule suitable for use in an attract-and-kill pest control approach. Also provided are methods of producing unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates using the recombinant microorganisms, as well as compositions comprising the recombinant microorganisms and/or optionally one or more of the product alcohols, aldehydes, or acetates.

12 Claims, 41 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

8,658,404	B2	2/2014	Schirmer et al.
8,987,531	B2	3/2015	Grubbs et al.
9,017,984	B2	4/2015	Hu et al.
9,068,201	B2	6/2015	Hu et al.
9,200,299	B2	12/2015	Friedman et al.
9,598,706	B2	3/2017	Keasling et al.
9,776,179	B2	10/2017	Wampler et al.
10,017,455	B2	7/2018	Hu et al.
10,308,962	B1 *	6/2019	Leonard C12Y 114/19001
2006/0078973	A1	4/2006	Renz et al.
2007/0282148	A1	12/2007	Berlin et al.
2008/0009598	A1	1/2008	Herrmann et al.
2008/0119678	A1	5/2008	Hock et al.
2008/0207911	A1	8/2008	Herrmann et al.
2008/0221345	A1	9/2008	Winde et al.
2008/0275247	A1	11/2008	Kadyrov et al.
2010/0087644	A1	4/2010	Mauduit et al.
2010/0113795	A1	5/2010	Arlt et al.
2010/0170826	A1	7/2010	Friedman et al.
2010/0174068	A1	7/2010	Grela et al.
2010/0199548	A1	8/2010	Del Cardayre et al.
2010/0235934	A1	9/2010	Friedman et al.
2010/0242345	A1	9/2010	Keasling et al.
2010/0251601	A1	10/2010	Hu et al.
2011/0015430	A1	1/2011	Schrock et al.
2011/0282068	A1	1/2011	Herrmann et al.
2011/0040099	A1	2/2011	Kadyrov et al.
2011/0065915	A1	3/2011	Malcolmson et al.
2011/0077421	A1	3/2011	Schrock
2011/0097769	A1	4/2011	Del Cardayre et al.
2011/0237815	A1	9/2011	Hock et al.
2011/0256599	A1	10/2011	Hu et al.
2012/0123133	A1	5/2012	Berlin et al.
2012/0142979	A1	6/2012	Keasling et al.
2012/0264983	A1	10/2012	Hu et al.
2012/0302710	A1	11/2012	Hoveyda et al.
2012/0323000	A1	12/2012	Hoveyda et al.
2013/0079515	A1	3/2013	Grela et al.
2013/0116434	A1	5/2013	Schrock et al.
2013/0144060	A1	6/2013	Mauduit et al.
2013/0211096	A1	8/2013	Arlt et al.
2013/0245339	A1	9/2013	Keasling et al.
2013/0261312	A1	10/2013	Allen et al.
2013/0274482	A1	10/2013	Schrock et al.
2013/0281688	A1	10/2013	Di Biase et al.
2013/0281706	A1	10/2013	Hock et al.
2013/0296511	A1	11/2013	Ung et al.
2014/0171607	A1	6/2014	Grela et al.
2014/0330018	A1	11/2014	Czirok et al.
2014/0378637	A1	12/2014	Schrock et al.
2015/0018557	A1	1/2015	Nolan et al.
2015/0038723	A1	2/2015	Herrmann et al.
2015/0045558	A1	2/2015	Plenio et al.
2015/0125933	A1	5/2015	Groban et al.
2015/0166465	A1	6/2015	Chen et al.
2015/0275188	A1	10/2015	Hu et al.
2016/0046914	A1	2/2016	Hom et al.
2016/0076058	A1	3/2016	Friedman et al.
2016/0108436	A1	4/2016	Coelho et al.
2016/0222419	A1	8/2016	Stuart
2016/0304913	A1	10/2016	Gatter et al.
2017/0275651	A1	9/2017	Keasling et al.
2017/0327799	A1	11/2017	Hu et al.
2018/0162916	A1	6/2018	Borodina et al.
2018/0371510	A1	12/2018	Gatter et al.
2019/0031594	A1	1/2019	Hu et al.
2019/0136272	A1	5/2019	Otte et al.
2019/0376094	A1	12/2019	Friedman et al.
2020/0017890	A1	1/2020	Sanchez-Riera et al.
2020/0140902	A1	5/2020	Sheppard et al.

FOREIGN PATENT DOCUMENTS

CN	102264910	A	11/2011
CN	102807470	A	12/2012

CN	103842502	A	6/2014
CN	104370701	A	2/2015
CN	104781411	A	7/2015
CN	104968672	A	10/2015
CN	108138202	A	6/2018
JP	H10-506783	A	7/1998
WO	WO 1991/009825	A1	7/1991
WO	WO 1992/019631	A1	11/1992
WO	WO 2007/075427	A1	7/2007
WO	WO 2007/136762	A2	11/2007
WO	WO 2007/140954	A1	12/2007
WO	WO 2008/066754	A1	6/2008
WO	WO 2008/100251	A1	8/2008
WO	WO 2008/113041	A2	9/2008
WO	WO 2008/119082	A2	10/2008
WO	WO 2008/147781	A2	12/2008
WO	WO 2009/094201	A3	7/2009
WO	WO 2009/126831	A1	10/2009
WO	WO 2010/037550	A1	4/2010
WO	WO 2010/144296	A2	12/2010
WO	WO 2011/040963	A1	4/2011
WO	WO 2011/069134	A3	6/2011
WO	WO 2011/091980	A1	8/2011
WO	WO 2011/097642	A1	8/2011
WO	WO 2012/087964	A1	6/2012
WO	WO 2012/167171	A3	12/2012
WO	WO 2012/168183	A1	12/2012
WO	WO 2013/135776	A1	3/2013
WO	WO 2013/070725	A1	5/2013
WO	WO 2014/001291	A1	1/2014
WO	WO 2014/008054	A2	1/2014
WO	WO 2014/067767	A1	5/2014
WO	WO 2014/134333	A1	9/2014
WO	WO 2014/139679	A3	9/2014
WO	WO 2014/155185	A1	10/2014
WO	WO 2014/169014	A1	10/2014
WO	WO 2014/172534	A1	10/2014
WO	WO 2015/003814	A1	1/2015
WO	WO 2015/003815	A1	1/2015
WO	WO 2015/042306	A1	3/2015
WO	WO 2015/077752	A1	5/2015
WO	WO 2015/086684	A1	6/2015
WO	WO 2015/171057	A1	11/2015
WO	WO 2016/099568	A1	6/2016
WO	WO 2016/159869	A1	10/2016
WO	WO 2016/207339	A1	12/2016
WO	WO 2017/087846	A1	5/2017
WO	WO 2017/214133	A2	12/2017
WO	WO 2018/213554	A1	11/2018

OTHER PUBLICATIONS

Ayciriex, et al., "YPR139c/LOA1 encodes a novel lysophosphatidic acid acyltransferase associated with lipid droplets and involved in TAG homeostasis." *Mol Biol Cell* (2012); 23 (2): 233-246.

Baba, et al., "Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection." *Molecular Systems Biology* (2006); 2 (1): 1-11.

Beisson, et al., "The acyltransferase GPAT5 is required for the synthesis of suberin in seed coat and root of *Arabidopsis*." *Plant Cell* (2007); 19 (1): 351-368.

Benghezal, et al., "SLC1 and SLC4 Encode Partially Redundant Acyl-Coenzyme A 1-Acylglycerol-3-phosphate O-Acyltransferases of Budding Yeast." *The Journal of Biological Chemistry* (2007); 282 (42): 30845-30855.

Blom, et al., "Sequence and structure-based prediction of eukaryotic protein phosphorylation sites." *J. Mol. Biol.* (1999); 294 (5): 1351-1362.

Bredeweg, et al., "A molecular genetic toolbox for *Yarrowia lipolytica*." *Biotechnol Biofuels* (2017); 10: 2, pp. 1-22, ePub Jan. 3, 2017.

Broun, et al., "Catalytic plasticity of fatty acid modification enzymes underlying chemical diversity of plant lipids." *Science* (1998); 282 (5392): 1315-1317.

Brown, et al., "Limnanthes douglasii lysophosphatidic acid acyltransferases: immunological quantification, acyl selectivity and functional replacement of the *Escherichia coli* plsC gene." *Biochemical Journal* (2002); 364 (3): 795-805.

(56)

References Cited

OTHER PUBLICATIONS

Chen, et al., "The yeast acylglycerol acyltransferase LCA1 is a key component of Lands cycle for phosphatidylcholine turnover." *FEBS Letters* (2007); 581 (28): 5511-5516.

Choi, et al., "Regulatory elements that control transcription activation and unsaturated fatty acid-mediated repression of the *Saccharomyces cerevisiae* OLE1 gene." *J Biol. Chem.* (1996); 271 (7): 3581-3589.

Colby, et al., "Calculating synergistic and antagonistic responses of herbicide combinations." *Weeds*, (1967); 15 (1): 20-22.

Dahlqvist, et al., "Phospholipid:diacylglycerol acyltransferase: An enzyme that catalyzes the acyl-CoA-independent formation of triacylglycerol in yeast and plants." *Proc Natl Acad Sci USA* (2000); 97 (12): 6487-6492.

Devos and Valencia. "Practical limits of function prediction." *Proteins: Structure, Function, and Genetics* (2000); 41 (1): 98-107.

Ding, et al., "Analysis of the *Agrotis segetum* pheromone gland transcriptome in the light of sex pheromone biosynthesis" *BMC Genomics* (2015); 16 (711): 1-21.

EBI Accession EAY76846. *Oryza sativa* triacylglycerol lipase, Dec. 29, 2008 [online]. [Retrieved Sep. 21, 2008]. 3 Pages, Retrieved from the internet: <URL: <https://www.ebi.ac.uk/ena/data/view/EAY76846&display=text>>.

Endo, et al., "Chelated ruthenium catalysts for Z-selective olefin metathesis" *J Am Chem Soc.* (2011); 133 (22): 8525-8527.

Extended European Search Report for Application No. EP 16867255.8 dated Feb. 5, 2019, 12 pages.

Flook, et al., "Z-Selective and Syndioselective Ring-Opening Metathesis Polymerization (ROMP) Initiated by MonoaryloxidePyrrolide (MAP) Catalysts" *Macromolecules* (2010) 43(18):7515-7522.

Gatter, et al., "A newly identified fatty alcohol oxidase gene is mainly responsible for the oxidation of long-chain ω -hydroxy fatty acids in *Yarrowia lipolytica*." *FEMS Yeast Res.* (Sep. 2014);14(6): 858-872. Epub Jul. 2, 2014.

GenBank Accession AAL49962.1. Diacylglycerol acyltransferase 1 [Bos Taurus], Feb. 11, 2002 [online]. [Retrieved Sep. 21, 2002]. 2 pages, Retrieved from the internet:< URL: <https://www.ncbi.nlm.nih.gov/protein/AAL49962.1/>>.

GenBank Accession KTA99184.1 Alcohol O-acetyltransferase 2 [Candida] glabrata]. Feb. 9, 2016 [online]. [Retrieved Sep. 21, 2018]. 1 page, Retrieved from the internet:< URL: <https://www.ncbi.nlm.nih.gov/protein/KTA99184.1/>>.

GenBank Accession AKD01723.1 Alcohol dehydrogenase 12 [Helicoverpa armigera], Apr. 25, 2015 [online]. [retrieved Sep. 21, 2018]. 1 page, Retrieved from the internet:< URL: <https://www.ncbi.nlm.nih.gov/protein/AKD01723.1/>>.

Goelz and Cronan Jr., "The positional distribution of fatty acids in *Escherichia coli* phospholipids is not regulated by sn-glycerol 3-phosphate levels." *J Bacteriol* (1980); 144 (1): 462-464.

Gonzalez, et al., "Fatty acid-responsive control of mRNA stability. Unsaturated fatty acid-induced degradation of the *Saccharomyces* OLE1 transcript" *J. Biol. Chem.* (1996); 271 (42): 25801-25809.

Greenway and Silbert, "Altered acyltransferase activity in *Escherichia coli* associated with mutations in acyl coenzyme A synthetase." *The Journal of Biological Chemistry* (1983); 258 (21): 13034-13042.

Groot, et al., "The Genetic Basis of Pheromone Evolution in Moths." *Annu Rev Entomol.* (2016); 61: 99117. Epub Nov. 4, 2015.

Hagström, et al., "A moth pheromone brewery: production of (Z)-11-hexadecenol by heterologous co-expression of two biosynthetic genes from a noctuid moth in a yeast cell factory" *Microb. Cell Fact.* (2013); 12: 125, pp. 1-11.

Hagstrom, et al., "Semi-selective fatty acyl reductases from four heliothine moths influence the specific pheromone composition" *PLoS One* (2012); 7 (5): e37230: 1-11.

Halford, B. "Olefin Metathesis for Macrocycles—Organic Synthesis: Tungsten catalysts make macrocyclic olefins with Z-selectivity" *Chem. Eng. News* (2011); 89 (45): 11.

Hartung, et al., "Highly Z-selective and enantioselective ring-opening/cross-metathesis catalyzed by a resolved stereogenic-at-Ru complex" *J Am Chem Soc.* (2013); 135 (28): 10183-10185.

Heier, et al., "Identification of Yju3p as functional orthologue of mammalian monoglyceride lipase in the yeast *Saccharomyces cerevisiae*." *Biochimica et Biophysica Acta* (2010); 1801 (9): 1063-1071.

Herbert, et al., "Concise syntheses of insect pheromones using Z-selective cross metathesis" *Angew Chem Int Ed Engl.* (2013); 52 (1): 310-314.

Hobbs, et al., "Cloning of a cDNA encoding diacylglycerol acyltransferase from *Arabidopsis thaliana* and its functional expression." *FEBS Lett* (1999); 452 (3): 145-149.

Ingrell, et al., "NetPhosYeast: prediction of protein phosphorylation sites in yeast." *Bioinformatics* (2007); 23 (7): 895-897.

Iwama, et al., "Alcohol dehydrogenases and an alcohol oxidase involved in the assimilation of exogenous fatty alcohols in *Yarrowia lipolytica*." *FEMS Yeast Research* (May 2015); 15(3): fov014, pp. 1-12.

Jain, et al., "Identification of a Novel Lysophospholipid Acyltransferase in *Saccharomyces cerevisiae*." *The Journal of Biological Chemistry* (2007); 282 (42): 30562-30569.

Jako, et al., "Seed-Specific Over-Expression of an *Arabidopsis* cDNA Encoding a Diacylglycerol Acyltransferase Enhances Seed Oil Content and Seed Weight." *Plant Physiol* (2001); 126 (2): 861-874.

Jurenka and Rafaeli, "Regulatory Role of PBAN in Sex Pheromone Biosynthesis of Heliothine Moths." *Front. Endocrinol.* (2011); 2 (46): 1-8.

Kajiwar, et al., "Molecular cloning and characterization of the $\Delta 9$ fatty acid desaturase gene and its promoter from *Saccharomyces kluyveri*" *FEMS Yeast. Res.* (2002); 2: 333-339.

Kalscheuer and Steinbüchel, "A Novel Bifunctional Wax Ester Synthase/Acyl-CoA:Diacylglycerol Acyltransferase Mediates Wax Ester and Triacylglycerol Biosynthesis in *Acinetobacter calcoaceticus* ADP1." *The Journal of Biological Chemistry* (2002); 278 (10): 8075-8082.

Keitz, et al., "Cis-selective ring-opening metathesis polymerization with ruthenium catalysts" *J Am Chem Soc.* (2012); 134 (4): 2040-2043.

Keitz, et al., "Improved ruthenium catalysts for Z-selective olefin metathesis" *J Am Chem Soc.* (2012); 134 (1): 693-699.

Kito, et al., "Inhibition of L-Glycerol 3-Phosphate Acyltransferase from *Escherichia coli* by cis-9, 10-Methylenehexadecanoic Acid." *The Journal of Biochemistry* (1972); 71 (1): 99-105.

Lardizabal, et al., "DGAT2 is a new diacylglycerol acyltransferase gene family: purification, cloning, and expression in insect cells of two polypeptides from *Mortierella ramanniana* with diacylglycerol acyltransferase activity." *The Journal of Biological Chemistry* (2001); 276 (42): 38862-38869.

Lassner, et al., "Lysophosphatidic Acid Acyltransferase from Meadowfoam Mediates Insertion of Erucic Acid at the sn-2 Position of Triacylglycerol in Transgenic Rapeseed Oil." *Plant Physiol* (1995); 109 (4): 1389-1394.

Lee, D. "Organic chemistry: Overcoming catalytic bias" *Nature* (2011) 471 (7339): 452-453.

Liénard, et al., "Sex pheromone biosynthetic pathways are conserved between moths and the butterfly *Bicyclus anynana*." *Nature Communications* (2014); 5: 3957, pp. 1-12.

Lewin, et al., "Analysis of Amino Acid Motifs Diagnostic for the sn-Glycerol-3-phosphate Acyltransferase Reaction." *Biochemistry* (1999); 38 (18): 5764-5771.

Li, et al., "Identification of acyltransferases required for cutin biosynthesis and production of cutin with suberin-like monomers." *Proc Natl Acad Sci USA* (2007); 104 (46): 18339-18344.

Liu, et al., "Functional and Topological Analysis of Yeast Acyl-CoA:Diacylglycerol Acyltransferase 2, an Endoplasmic Reticulum Enzyme Essential for Triacylglycerol Biosynthesis." *The Journal of Biological Chemistry* (2011); 286 (15): 13115-13126.

Lu, et al., "Acyl-phosphates initiate membrane phospholipid synthesis in Gram-positive pathogens." *Mol Cell* (2006); 23 (5): 765-772.

Maniatis, et al., "Regulation of inducible and tissue-specific gene expression" *Science* (1987); 236 (4806): 1237-1245.

(56)

References Cited

OTHER PUBLICATIONS

- Marx, et al., "Stereoselective access to Z and E macrocycles by ruthenium-catalyzed Z-selective ring-closing metathesis and ethenolysis" J Am Chem Soc. (2013); 135 (1): 94-97.
- Mauersberger, et al., "Insertional Mutagenesis in the n-Alkane-Assimilating Yeast *Yarrowia lipolytica*: Generation of Tagged Mutations in Genes Involved in Hydrophobic Substrate Utilization." J. Bacteriology (2001); 183 (17): 5102-5109.
- Meek, et al. "Z-selective catalytic olefin cross-metathesis for natural product synthesis" Nature (2011); 471 (7339): 461-466.
- Miller, W. T., "Tyrosine kinase signaling and the emergence of multicellularity" Biochimica et Biophysica Acta (BBA)—Molecular Cell Research (2012); 1823 (6): 1053-1057.
- Moss, et al. "Determination of cellular fatty acid compositions of various yeasts by gas-liquid chromatography" J Clin Microbiol. (1982); 16 (6): 1073-1079.
- Moto, et al., "Involvement of a bifunctional fatty-acyl desaturase in the biosynthesis of the silkworm, *Bombyx mori*, sex pheromone." PNAS (2004); 101 (23): 8631-8636.
- Murata, et al., "Modes of Fatty-Acid Desaturation in Cyanobacteria." Plant Cell Physiol (1992); 33(7): 933-941.
- Nagiec, et al., "A suppressor gene that enables *Saccharomyces cerevisiae* to grow without making sphingolipids encodes a protein that resembles an *Escherichia coli* fatty acyltransferase." The Journal of Biological Chemistry (1993); 268 (29): 22156-22163.
- Nishida, et al., "The gene and the RNA for the precursor to the plastid-located glycerol-3-phosphate acyltransferase of *Arabidopsis thaliana*." Plant Mol Biol. (1993); 21 (2): 267-277.
- Oelkers, et al., "The DGA1 gene determines a second triglyceride synthetic pathway in yeast." The Journal of Biological Chemistry (2002); 277 (11): 8877-8881.
- Okuyama and Wakil, "Positional Specificities of Acyl Coenzyme A:Glycerophosphate and Acyl Coenzyme A: Monoacylglycerophosphate Acyltransferases in *Escherichia coli*." The Journal of Biological Chemistry (1973); 248 (14): 5197-5205.
- Ondi, et al. "High activity, stabilized formulations, efficient synthesis and industrial use of Mo-and W-based metathesis catalysts" XiMo Technology Updates, 2015: http://www.ximoinc.com/files/ximo/uploads/download/Summary_3.11.15.pdf.
- PCT/US2016/062852, International Preliminary Report on Patentability, dated May 22, 2018, 10 pages.
- PCT/US2016/062852, International Search Report and Written Opinion, dated Feb. 7, 2017, 13 pages.
- PCT/US2016/062852, Third Party Observation filed by Danmarks Tekniske Universitet on Oct. 5, 2017 with WIPO, 7 pages.
- PCT/US2017/036136, Invitation to Pay Additional Fees, mailed Sep. 20, 2017, 3 pages.
- PCT/US2017/036136, International Search Report and Written Opinion, dated Nov. 17, 2017, 15 pages.
- PCT/US2017/036136, International Preliminary Report on Patentability, dated Dec. 11, 2018, 10 pages.
- PCT/US2018/033151, Invitation to Pay Additional Fees, mailed Aug. 14, 2018, 6 pages.
- PCT/US2018/033151, International Search Report and Written Opinion, dated Oct. 15, 2018, 22 pages.
- Peryshkov, et al. "B(C₆F₅)₃ Activation of Oxo Tungsten complexes that are relevant to olefin metathesis" Organometallics (2013); 32 (19): 5256-5259.
- Peryshkov, et al., "Z-Selective olefin metathesis reactions promoted by tungsten oxo alkylidene complexes" J Am Chem Soc. (2011); 133 (51): 20754-20757.
- Riekhof, et al., "Identification and Characterization of the Major Lysophosphatidylethanolamine Acyltransferase in *Saccharomyces cerevisiae*." The Journal of Biological Chemistry (2007); 282 (39): 28344-28352.
- Rock, et al., "Phospholipid synthesis in *Escherichia coli*. Characteristics of fatty acid transfer from acyl-acyl carrier protein to sn-glycerol 3-phosphate." The Journal of Biological Chemistry (1981); 256 (2): 736-742.
- Rosenfeld, et al., "Structural and functional conservation and divergence among acyl-CoA desaturases of two noctuid species, the corn earworm, *Helicoverpa zea*, and the cabbage looper, *Trichoplusia ni*." Insect. Biochem. Mol. Biol. (2001); 31 (10): 949-964.
- Sandager, et al., "Storage lipid synthesis is non-essential in yeast." J Biol Chem (2002); 277 (8): 6478-6482.
- Schrock, et al. "Z-Selective and syndioselective ring-opening metathesis polymerization (ROMP) Initiated by monoaryloxidepyrrolide (MAP) catalysts" Macromolecules (2010); 43 (18): 7515-7522.
- Seffernick, et al., "Melamine deaminase and Atrazine chlorohydrolase: 98 percent identical but functionally different." J. Bacteriol. (2001); 183 (8): 2405-2410.
- Shi, et al., "Improving Production of Malonyl Coenzyme A-Derived Metabolites by Abolishing Snf1-Dependent Regulation of Acc1." mBio (2014); 5 (3): e01130-14: 1-8.
- Sorger and Daum, "Synthesis of Triacylglycerols by the Acyl-Coenzyme A:Diacyl-Glycerol Acyltransferase Dga1p in Lipid Particles of the Yeast *Saccharomyces cerevisiae*." J Bacteriol (2002); 184 (2):519-524.
- Ståhl, et al., "Cloning and Functional Characterization of a Phospholipid:Diacylglycerol Acyltransferase from *Arabidopsis*." Plant Physiology (2004); 135 (3):1324-1335.
- Stelinski, et al., "Sprayable microencapsulated sex pheromone formulations for mating disruption of four tortricid species: effects of application height, rate, frequency, and sticker adjuvant" J Econ. Entomol. (2007); 100(4): 1360-1369.
- Stöveken, et al., "The Wax Ester Synthase/Acyl Coenzyme A:Diacylglycerol Acyltransferase from *Acinetobacter* sp. Strain ADP1: Characterization of a Novel Type of Acyltransferase." J Bacteriol (2005); 187 (4): 1369-1376.
- Takai et al. "Construction and characterization of a *Yarrowia lipolytica* mutant lacking genes encoding cytochromes P450 subfamily 52." Fungal Genet Biol. (2012); 49 (1): 58-64. Epub Nov. 17, 2011.
- Townsend, et al. "Z-selective metathesis homocoupling of 1,3-dienes by molybdenum and tungsten monoaryloxide pyrrolide (MAP) complexes" J Am Chem Soc. (2012); 134 (28): 11334-11337.
- UniProtKB-074934 (ACOX1_YARLI): 9 (nine) pages downloaded on Oct. 23, 2018 from <https://www.uniprot.org/uniprot/074934>.
- Uniprot Accession A0A178WDE4. Acyl-coenzyme A oxidase, Apr. 12, 2017 [online]. [Retrieved on Aug. 10, 2018]. 1 page, Retrieved from the internet: <URL: <https://www.uniprot.org/uniprot/A0A178WDE4.txt?version=7>>.
- Uniprot Accession R8XW24. *Acinetobacter calcoaceticus*—Fatty acyl-CoA reductase, Apr. 13, 2013 [online]. [Retrieved Sep. 21, 2018]. 1 page, Retrieved from the internet: <URL: <https://www.uniprot.org/uniprot/R8XW24.txt?version=14>>.
- Uthoff, et al., "Thio Wax Ester Biosynthesis Utilizing the Unspecific Bifunctional Wax Ester Synthase/Acyl Coenzyme A:Diacylglycerol Acyltransferase of *Acinetobacter* sp. Strain ADP1." Appl. Environ. Microbiol. (2005); 71 (2): 790-796.
- Wahl, et al., "Antagonistic regulation of dgkA and plsB genes of phospholipid synthesis by multiple stress responses in *Escherichia coli*." Molecular Microbiology (2011); 80 (5): 1260-1275.
- Wang, et al. "Efficient and selective formation of macrocyclic disubstituted Z alkenes by ring-closing metathesis (RCM) reactions catalyzed by Mo- or W-based monoaryloxide pyrrolide (MAP) complexes: applications to total syntheses of epilachnene, yuzu lactone, ambrettolide, epothilone C, and nakadomarin A" Chemistry (2013); 19 (8): 2726-2740.
- Wang, et al., "Exploring fatty alcohol-producing capability of *Yarrowia lipolytica*." Biotechnology for Biofuels (2016); 9: 107, pp. 1-10.
- Wang, et al., "Mo-Based Complexes with Two Aryloxides and a Pentafluoroimido Ligand: Catalysts for Efficient Z-Selective Synthesis of a Macrocyclic Trisubstituted Alkene by Ring-Closing Metathesis." Angew Chem Int Ed Engl., (2013); 52 (7): 1939-1943.
- Whisstock and Lesk, "Prediction of protein function from protein sequence and structure." Q. Rev. Biophysics. (2003); 36 (3): 307-340.
- Witkowski, et al., "Conversion of a β -ketoacyl synthase to a malonyl decarboxylase by replacement of the active-site cysteine with glutamine." Biochemistry (1999); 38 (36): 11643-11650.

(56)

References Cited

OTHER PUBLICATIONS

Xia, et al., "Large Number of putative chemoreception and pheromone biosynthesis genes revealed by analyzing transcriptome from ovipositor-pheromone glands of *Chilo suppressalis*." Scientific Reports (Jan. 2015); 5: 7888. Epub Jan. 20, 2015.

Yoshimura, et al., "Involvement of the YneS/YgiH and PlsX proteins in phospholipid biosynthesis in both *Bacillus subtilis* and *Escherichia coli*." BMC Microbiology (2007); 7: 69, 13 pages.

Yousuf, et al., "Microbial conversion of olive oil mill wastewaters into lipids suitable for biodiesel production." J Agric. Food Chem. (2010); 58 (15): 8630-8635.

Yu, et al., "Enol Ethers as Substrates for Efficient Z- and Enantioselective Ring-Opening/Cross-Metathesis Reactions Promoted by Stereogenic-at-Mo Complexes: Utility in Chemical Synthesis and Mechanistic Attributes." J Am. Chem. Soc. (2012); 134(5): 2788-2799.

Yu, et al., "Synthesis of macrocyclic natural products by catalyst-controlled stereoselective ring-closing metathesis" Nature, (2011); 479 (7371): 88-93.

Zhao, et al. "Endo-selective enyne ring-closing metathesis promoted by stereogenic-at-W mono-pyrrolide complexes" Org Lett. (2011); 13 (4): 784-787.

Zheng and Zou, "The initial step of the glycerolipid pathway: identification of glycerol 3-phosphate/dihydroxyacetone phosphate dual substrate acyltransferases in *Saccharomyces cerevisiae*." The Journal of Biological Chemistry (2001); 276 (45): 417104-417116.

Zou, et al., "The *Arabidopsis thaliana* TAG1 mutant has a mutation in a diacylglycerol acyltransferase gene." The Plant Journal (1999); 19 (6): 645-653.

Chandrasekhar, et al., "One pot conversion of carboxylic acids to aldehydes with DIBAL-H". Tetrahedron Letters (Feb. 19, 1998); 39(8): 909-910.

Extended European Search Report for Application No. EP 17810850.2 dated Feb. 18, 2020, 9 pages.

Kuettel and Chapman, "The 9-hexadecenoic and 11-octadecenoic acid content of natural fats and oils". LIPIDS (1968); 3(4): 313-316.

Vickery, J.R., "The fatty acid composition of the seed oils of proteaceae: A chemotaxonomic study". Phytochemistry (Jan. 1971); 10(1): 123-130.

Wang, et al., "Comparative study of sex pheromone composition and biosynthesis in *Helicoverpa armigera*, *H. assulta* and their hybrid". Insect Biochemistry and Molecular Biology (Jun. 2005); 35(6): 575-583. Epub Mar. 16, 2005.

Zhu and Jackson, "Metabolic engineering of *Yarrowia lipolytica* for industrial applications". Current Opinion in Biotechnology (Dec. 2015); 36: 65-72.

U.S. Appl. No. 15/983,706, filed May 18, 2018, U.S. Pat. No. 10,308,962, Patented.

U.S. Appl. No. 16/307,282, filed Dec. 5, 2018, Pending.

U.S. Appl. No. 16/614,144, filed Nov. 15, 2019, Pending.

Ando, et. Al., "Lepidopteran Sex Pheromones". Topics in Current Chemistry (2004); 239: 51-96.

GenBank entry AF272342.1, *Helicoverpa zea* acyl-CoA delta-11 desaturase (PGDs1) mRNA, complete cds, Aug. 30, 2001, 2 pages, (<https://www.ncbi.nlm.nih.gov/nuccore/AF272342.1>), retrieved on Jan. 6, 2020.

Heath, et al., "Periodicity of Female Sex Pheromone Titer and Release in *Heliothis subflexa* and *H. virescens* (Lepidoptera: Noctuidae)". Annals of the Entomological Society of America (Mar. 1, 1991); vol. 84, Issue 2, pp. 182-189.

PCT/US2018/033151, International Preliminary Report on Patentability, dated Nov. 19, 2019, 14 pages.

Rosenfield, et al., "Structural and functional conservation and divergence among acyl-CoA desaturases of two noctuid species, the corn earworm, *Helicoverpa zea*, and the cabbage looper, *Trichoplusia ni*." Insect. Biochem. Mol. Biol. (2001); 31 (10): 949-964.

Sheng and Feng, "Metabolic engineering of yeast to produce fatty acid-derived biofuels: bottlenecks and solutions". Frontiers in Microbiology (2015) vol. 6, 11 pages.

Third Party Observations filed Jan. 8, 2020, in European Application No. 16867255.8, 19 pages.

European Patent Office Communication dated Jan. 13, 2020 transmitting Third Party Observations filed Jan. 8, 2020, in European Application No. 16867255.8, 1 page.

Extended European Search Report for Application No. EP 18803040.7 dated Feb. 26, 2021, 7 pages.

Flook, et al., "Synthesis of cis,syndiotactic ROMP Polymers Containing Alternating Enantiomers", J. Am. Chem. Soc. (Jan. 25, 2011); 133(6): 1784-1786.

Gerber and Schrock, "Synthesis of methyldiene complexes that contain a 2,6-dimesitylphenylimido ligand and ethenolysis of 2,3-dicarbomethoxynorbornadiene" Organometallics (2013); 32 (19): 5573-5580.

Los and Murata, "Structure and expression of fatty acid desaturases", Biochimica et Biophysica Acta (Oct. 2, 1998); 1394(1): 3-15.

Marinescu, et al. "Synthesis of variations of Stereogenic-at-Metal Imido Alkylidene Complexes of Molybdenum" Organometallics (2012); 31: 6336-6343.

Mauersberger, Stephan, "Cytochromes P450 of the Alkane-Utilising Yeast *Yarrowia lipolytica*", pp. 227-262 in G. Barth (ed.), *Yarrowia lipolytica*, Microbiology Monographs vol. 25, Springer-Verlag Berlin Heidelberg (2013); 273 pages.

Shanklin, et al., "Eight Histidine Residues Are Catalytically Essential in a Membrane-Associated Iron Enzyme, Stearoyl-CoA Desaturase, and Are Conserved in Alkane Hydroxylase and Xylene Monooxygenase", Biochemistry (Jan. 1, 1994), 33 (43): 12787-12794.

GenBank Accession No. AAD03775.1, "acyl-CoA delta11 desaturase [*Trichoplusia ni*]", Jan. 11, 1999.

GenBank Accession No. AAF81787.1, "acyl-CoA delta-11 desaturase [*Helicoverpa zea*]", Aug. 30, 2001, 1 page.

GenBank Accession No. AAF81790.2, "acyl-CoA delta-9 desaturase [*Helicoverpa zea*]", Aug. 30, 2001, 2 pages.

GenBank Accession No. AAM28480.2, "acyl-CoA desaturase HassGATD [*Helicoverpa assulta*]", May 22, 2003, 2 pages.

GenBank Accession No. AAM28481.2, "acyl-CoA desaturase HassKPSE [*Helicoverpa assulta*]", May 22, 2003, 1 page.

GenBank Accession No. AAM28483.2, "acyl-CoA desaturase HassLPAQ [*Helicoverpa assulta*]", May 22, 2003, 2 pages.

GenBank Accession No. AAM28484.2, "acyl-CoA desaturase HassNPVE [*Helicoverpa assulta*]", May 22, 2003, 1 page.

GenBank Accession No. AF416738.1, "Argyrotaenia velutinana acyl-CoA delta-11 desaturase mRNA, complete cds", Oct. 17, 2001, 2 pages.

GenBank Accession No. AF482906.2, "Helicoverpa assulta acyl-CoA desaturase HassKPSE mRNA, complete cds", May 22, 2003, 1 page.

GenBank Accession No. AF545481.1, "Choristoneura rosaceana acyl-CoA Z/E11 desaturase mRNA, complete cds", Mar. 8, 2005, 2 pages.

GenBank Accession No. AKU76402.1, "acyl-CoA desaturase 3 [*Helicoverpa armigera*]", Dec. 31, 2015, 2 pages.

GenBank Accession No. AKU76404.1, "acyl-CoA desaturase 5 [*Helicoverpa armigera*]", Dec. 31, 2015, 2 pages.

GenBank Accession No. AKU76405.1, "acyl-CoA desaturase 6 [*Helicoverpa armigera*]", Dec. 31, 2015, 1 page.

GenBank Accession No. AKU76408.1, "acyl-CoA desaturase 2 [*Helicoverpa assulta*]", Dec. 31, 2015, 1 page.

GenBank Accession No. AKU76409.1, "acyl-CoA desaturase 3 [*Helicoverpa assulta*]", Dec. 31, 2015, 1 page.

GenBank Accession No. AKU76410.1, "acyl-CoA desaturase 4 [*Helicoverpa assulta*]", Dec. 31, 2015, 1 page.

GenBank Accession No. AKU76411.1, "acyl-CoA desaturase 5 [*Helicoverpa assulta*]", Dec. 31, 2015, 1 page.

GenBank Accession No. AKU76412.1, "acyl-CoA desaturase 6 [*Helicoverpa assulta*]", Dec. 31, 2015, 2 pages.

GenBank Accession No. ATJ44449.1, "desaturase MPVE [*Helicoverpa armigera*]", Oct. 18, 2017, 1 page.

GenBank Accession No. ATJ44454.1, "desaturase LPAQ [*Helicoverpa armigera*]", Oct. 18, 2017, 2 pages.

GenBank Accession No. ATJ44456.1, "desaturase PDSN [*Helicoverpa armigera*]", Oct. 18, 2017, 1 page.

(56)

References Cited

OTHER PUBLICATIONS

GenBank Accession No. ATJ44457.1, “desaturase IPAE [*Helicoverpa armigera*]”, Oct. 18, 2017, 2 pages.

GenBank Accession No. ATJ44509.1, “desaturase KSVE [*Helicoverpa assulta*]”, Oct. 18, 2017, 1 page.

GenBank Accession No. ATJ44511.1, “desaturase MPVE [*Helicoverpa assulta*]”, Oct. 18, 2017, 1 page.

GenBank Accession No. ATJ44514.1, “desaturase PDSN [*Helicoverpa assulta*]”, Oct. 18, 2017, 1 page.

GenBank Accession No. AY493438.1, “*Thalassiosira pseudonana* delta-11 fatty acid desaturase (desN) gene, complete cds”, Apr. 6, 2004, 2 pages.

GenBank Accession No. EU152335.1, “*Lampronia capitella* acyl-CoA-delta11-desaturase mRNA, complete cds”, Oct. 12, 2008, 2 pages.

GenBank Accession No. JX679209.1, “*Agrotis segetum* fatty acyl delta-11 desaturase (D11) mRNA, complete cds”, Aug. 3, 2014, 1 page.

GenBank Accession No. JX964774.1, “*Amyelois transitella* delta 11 desaturase mRNA, complete cds”, Sep. 3, 2014, 2 pages.

GenBank Accession No. PZC81408.1, “hypothetical protein B5X24_HaOG212613 [*Helicoverpa armigera*]”, Jun. 15, 2018, 2 pages.

GenBank Accession No. PZC82682.1, “hypothetical protein B5X24_HaOG209780 [*Helicoverpa armigera*]”, Jun. 15, 2018, 2 pages.

GenBank Accession No. PZC86045.1, “hypothetical protein B5X24_HaOG213003 [*Helicoverpa armigera*]”, Jun. 15, 2018, 2 pages.

GenBank Accession No. PZC86046.1, “hypothetical protein B5X24_HaOG213004 [*Helicoverpa armigera*]”, Jun. 15, 2018, 2 pages.

GenBank Accession No. XP_021183629.1, “acyl-CoA Delta(11) desaturase [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. XP_021185492.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

GenBank Accession No. XP_021187735.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. XP_021182948.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. XP_021183600.1, “acyl-CoA Delta(11) desaturase-like isoform X1 [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

GenBank Accession No. XP_021183601.1, “acyl-CoA Delta(11) desaturase-like isoform X2 [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

GenBank Accession No. XP_021183624.1, “stearoyl-CoA desaturase 5 [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. XP_021183628.1, “acyl-CoA Delta(11) desaturase [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. XP_021183660.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

GenBank Accession No. XP_021183696.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

GenBank Accession No. XP_021190176.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

GenBank Accession No. XP_021195328.1, “stearoyl-CoA desaturase 5-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. Xp_021195974.1, “Low Quality Protein: acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 2 pages.

GenBank Accession No. XP_021200693.1, “acyl-CoA Delta(11) desaturase-like [*Helicoverpa armigera*]”, Jun. 1, 2017, 1 page.

* cited by examiner




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5

FIG. 6A




FIG. 6B




FIG. 7A





FIG. 7B






FIG. 8




FIG. 9




FIG. 10

FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D





FIG. 11E






FIG. 12A




FIG. 12B




FIG. 13A




FIG. 13B

FIG. 14A





FIG. 14B






FIG. 15




FIG. 16




FIG. 17




FIG. 18

FIG. 19A




FIG. 19B





FIG. 19C






FIG. 20

pPV0137 (mCherry_Ct)




FIG. 21B




FIG. 21D

Negative control (water)




FIG. 21A




FIG. 21C





FIG. 22






FIG. 24A




FIG. 24B

FIG. 24C






FIG. 24D

FIG. 24E






FIG. 26




FIG. 27




FIG. 28




FIG. 29




FIG. 30








FIG. 31








FIG. 34




FIG. 35




FIG. 36

FIG. 37A





FIG. 37B






FIG. 38

FIG. 39A





FIG. 39B






FIG. 40

MICROORGANISMS FOR THE PRODUCTION OF INSECT PHEROMONES AND RELATED COMPOUNDS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application is a Continuation of U.S. patent application Ser. No. 15/983,706 filed May 18, 2018 (issued as U.S. Pat. No. 10,308,962 on Jun. 4, 2019), which is a Continuation of International Application No. PCT/US2016/062852, filed on Nov. 18, 2016, which claims the benefit of priority to U.S. Provisional Application No. 62/257,054, filed Nov. 18, 2015, and U.S. Provisional Application No. 62/351,605, filed Jun. 17, 2016; each of the aforementioned applications is herein expressly incorporated by reference.

STATEMENT REGARDING THE SEQUENCE LISTING

The Sequence Listing associated with this application is provided in text format in lieu of a paper copy, and is hereby incorporated by reference into the specification. The name of the text file containing the Sequence Listing is PRVI_007_02WO_SeqList_ST25.txt. The text file is about 54 KB, was created on Nov. 15, 2016, and is being submitted electronically via EFS-Web.

TECHNICAL FIELD

This application relates to recombinant microorganisms useful in the biosynthesis of unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates which may be useful as insect pheromones, fragrances, flavors, and polymer intermediates. The application further relates to methods of producing unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates using the recombinant microorganisms, as well as compositions comprising one or more of these compounds and/or the recombinant microorganisms.

BACKGROUND

As the global demand for food grows, there is an increasing need for effective pest control. Conventional insecticides are among the most popular chemical control agents because they are readily available, rapid acting, and highly reliable. However, the overuse, misuse, and abuse of these chemicals have led to resistant pests, alteration of the natural ecology, and in some cases, environmental damage.

The use of insect pheromones to control pest populations has gained increasing popularity as a viable, safe, and environmentally-friendly alternative to conventional insecticides. Since their discovery in the late 1950s, these molecules have shown efficacy in reducing insect populations through a variety of methods, including mass trappings, attract and kill, and mating disruption. The latter method in particular represents a non-toxic means of pest control and utilizes the ability of synthetic pheromones to mask naturally occurring pheromones, thereby causing confusion and mating disruption.

Although pheromones have significant potential in agricultural insect control, the cost of synthesizing pheromones using currently available techniques is very high, which prohibits widespread use of this sustainable technology beyond high-value crops. Thus, there is an existing need to develop novel technologies for the cost-efficient production of insect pheromones and related fragrances, flavors, and

polymer intermediates. The present inventors address this need with the development of recombinant microorganisms capable of producing a wide-range of unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates including synthetic insect pheromones from low-cost feedstocks.

SUMMARY OF THE DISCLOSURE

The present application relates to recombinant microorganisms having a biosynthesis pathway for the production of one or more compounds selected from unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates. The recombinant microorganisms described herein may be used for the production of at least one compound, such as an insect pheromone, a fragrance, or a flavoring agent, selected from unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates.

In one embodiment, the recombinant microorganism comprises a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol. Accordingly, in a first aspect, the application relates to a recombinant microorganism capable of producing an unsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated C₆-C₂₄ fatty acyl-CoA, wherein the recombinant microorganism expresses (a): at least one exogenous nucleic acid molecule encoding a fatty-acyl desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-CoA to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA; and (b): at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA from (a) into the corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is an insect pheromone. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is a fragrance or flavoring agent. In some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an alcohol oxidase or an alcohol dehydrogenase, wherein the alcohol oxidase or alcohol dehydrogenase is capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from (b) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde. In some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an acetyl transferase capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from (b) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acetate.

In some embodiments, the fatty-acyl desaturase is a desaturase capable of utilizing a fatty acyl-CoA as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms.

In some embodiments, the fatty-acyl desaturase is capable of generating a double bond at position C5, C6, C7, C8, C9, C10, C11, C12, or C13 in the fatty acid or its derivatives, such as, for example, fatty acid CoA esters.

In one exemplary embodiment, the fatty-acyl desaturase is a Z11 desaturase. In various embodiments described herein, the Z11 desaturase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Agrotis segetum*, *Amyelois transitella*, *Argyrotaenia velutana*, *Choristoneura rosaceana*, *Lampronia capitella*, *Trichoplusia ni*, *Helicoverpa zea*, or *Thalassiosira pseudonana*. Further Z11-desaturases, or the nucleic acid sequences encoding them, can be isolated from *Bombyx mori*, *Manduca sexta*, *Diatraea grandiosella*, *Earias insulana*, *Earias*

3

vittella, *Plutella xylostella*, *Bombyx mori* or *Diaphania nitidalis*. In exemplary embodiments, the Z11 desaturase comprises a sequence selected from GenBank Accession Nos. JX679209, JX964774, AF416738, AF545481, EU152335, AAD03775, AAF81787, and AY493438. In some embodiments, a nucleic acid sequence encoding a Z11 desaturase from organisms of the species *Agrotis segetum*, *Amyelois transitella*, *Argyrotaenia velutiana*, *Choristoneura rosaceana*, *Lampronia capitella*, *Trichoplusia ni*, *Helicoverpa zea*, or *Thalassiosira pseudonana* is codon optimized. In some embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 9, 18, 24 and 26 from *Trichoplusia ni*. In other embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 10 and 16 from *Agrotis segetum*. In some embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 11 and 23 from *Thalassiosira pseudonana*. In certain embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 12, 17 and 30 from *Amyelois transitella*. In further embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 13, 19, 25, 27 and 31 from *Helicoverpa zea*. In some embodiments, the Z11 desaturase comprises a chimeric polypeptide. In some embodiments, a complete or partial Z11 desaturase is fused to another polypeptide. In certain embodiments, the N-terminal native leader sequence of a Z11 desaturase is replaced by an oleosin leader sequence from another species. In certain embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 15, 28 and 29.

In certain embodiments, the Z11 desaturase catalyzes the conversion of a fatty acyl-CoA into a mono- or polyunsaturated product selected from Z11-13:Acyl-CoA, E11-13:Acyl-CoA, (Z,Z)-7,11-13:Acyl-CoA, Z11-14:Acyl-CoA, E11-14:Acyl-CoA, (E,E)-9,11-14:Acyl-CoA, (E,Z)-9,11-14:Acyl-CoA, (Z,E)-9,11-14:Acyl-CoA, (Z,Z)-9,11-14:Acyl-CoA, (E,Z)-9,11-15:Acyl-CoA, (Z,Z)-9,11-15:Acyl-CoA, Z11-16:Acyl-CoA, E11-16:Acyl-CoA, (E,Z)-6,11-16:Acyl-CoA, (E,Z)-7,11-16:Acyl-CoA, (E,Z)-8,11-16:Acyl-CoA, (E,E)-9,11-16:Acyl-CoA, (E,Z)-9,11-16:Acyl-CoA, (Z,E)-9,11-16:Acyl-CoA, (Z,Z)-9,11-16:Acyl-CoA, (E,E)-11,13-16:Acyl-CoA, (E,Z)-11,13-16:Acyl-CoA, (Z,E)-11,13-16:Acyl-CoA, (Z,Z)-11,13-16:Acyl-CoA, (Z,E)-11,14-16:Acyl-CoA, (E,E,Z)-4,6,11-16:Acyl-CoA, (Z,Z,E)-7,11,13-16:Acyl-CoA, (E,E,Z,Z)-4,6,11,13-16:Acyl-CoA, Z1-17:Acyl-CoA, (Z,Z)-8,11-17:Acyl-CoA, Z11-18:Acyl-CoA, E11-18:Acyl-CoA, (Z,Z)-11,13-18:Acyl-CoA, (E,E)-11,14-18:Acyl-CoA, or combinations thereof.

In another exemplary embodiment, the fatty-acyl desaturase is a Z9 desaturase. In various embodiments described herein, the Z9 desaturase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Ostrinia furnacalis*, *Ostrinia nubilalis*, *Choristoneura rosaceana*, *Lampronia capitella*, *Helicoverpa assulta*, or *Helicoverpa zea*. In exemplary embodiments, the Z9 desaturase comprises a sequence selected from GenBank Accession Nos. AY057862, AF243047, AF518017, EU152332, AF482906, and AAF81788. In some embodiments, a nucleic acid sequence encoding a Z9 desaturase is codon optimized. In some embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 20 from *Ostrinia furnacalis*. In other embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 21 from *Lampronia capitella*. In some embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 22 from *Helicoverpa zea*.

In certain embodiments, the Z9 desaturase catalyzes the conversion of a fatty acyl-CoA into a monounsaturated or

4

polyunsaturated product selected from Z9-11:Acyl-CoA, Z9-12:Acyl-CoA, E9-12:Acyl-CoA, (E,E)-7,9-12:Acyl-CoA, (E,Z)-7,9-12:Acyl-CoA, (Z,E)-7,9-12:Acyl-CoA, (Z,Z)-7,9-12:Acyl-CoA, Z9-13:Acyl-CoA, E9-13:Acyl-CoA, (E,Z)-5,9-13:Acyl-CoA, (Z,E)-5,9-13:Acyl-CoA, (Z,Z)-5,9-13:Acyl-CoA, Z9-14:Acyl-CoA, E9-14:Acyl-CoA, (E,Z)-4,9-14:Acyl-CoA, (E,E)-9,11-14:Acyl-CoA, (E,Z)-9,11-14:Acyl-CoA, (Z,E)-9,11-14:Acyl-CoA, (Z,Z)-9,11-14:Acyl-CoA, (E,E)-9,12-14:Acyl-CoA, (Z,E)-9,12-14:Acyl-CoA, (Z,Z)-9,12-14:Acyl-CoA, Z9-15:Acyl-CoA, E9-15:Acyl-CoA, (Z,Z)-6,9-15:Acyl-CoA, Z9-16:Acyl-CoA, E9-16:Acyl-CoA, (E,E)-9,11-16:Acyl-CoA, (E,Z)-9,11-16:Acyl-CoA, (Z,E)-9,11-16:Acyl-CoA, (Z,Z)-9,11-16:Acyl-CoA, Z9-17:Acyl-CoA, E9-18:Acyl-CoA, Z9-18:Acyl-CoA, (E,E)-5,9-18:Acyl-CoA, (E,E)-9,12-18:Acyl-CoA, (Z,Z)-9,12-18:Acyl-CoA, (Z,Z,Z)-3,6,9-18:Acyl-CoA, (E,E,E)-9,12,15-18:Acyl-CoA, (Z,Z,Z)-9,12,15-18:Acyl-CoA, or combinations thereof.

In some embodiments, the recombinant microorganism may express a bifunctional desaturase capable of catalyzing the subsequent desaturation of two double bonds.

In some embodiments, the recombinant microorganism may express more than one exogenous nucleic acid molecule encoding a fatty-acyl desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-CoA to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA. For instance, the recombinant microorganism may express an exogenous nucleic acid molecule encoding a Z11 desaturase and another exogenous nucleic acid molecule encoding a Z9 desaturase.

In some embodiments, the recombinant microorganism may express a fatty-acyl conjugase that acts independently or together with a fatty-acyl desaturase to catalyze the conversion of a saturated or monounsaturated fatty acyl-CoA to a conjugated polyunsaturated fatty acyl-CoA.

In one embodiment, the disclosure provides a recombinant microorganism capable of producing a polyunsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA, wherein the recombinant microorganism expresses: (a) at least one exogenous nucleic acid molecule encoding a fatty acyl conjugase that catalyzes the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA; and (b) at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the polyunsaturated C₆-C₂₄ fatty acyl-CoA from (a) into the corresponding polyunsaturated C₆-C₂₄ fatty alcohol.

In another embodiment, the recombinant microorganism expresses at least two exogenous nucleic acid molecules encoding fatty-acyl conjugases that catalyze the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA.

In a further embodiment, the disclosure provides a recombinant microorganism capable of producing a polyunsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA, wherein the recombinant microorganism expresses: (a) at least one exogenous nucleic acid molecule encoding a fatty-acyl desaturase and at least one exogenous nucleic acid molecule encoding a fatty acyl conjugase that catalyze the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA; and (b) at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the polyunsatu-

5

rated C₆-C₂₄ fatty acyl-CoA from (a) into the corresponding polyunsaturated C₆-C₂₄ fatty alcohol.

In another embodiment, the recombinant microorganism expresses at least two exogenous nucleic acid molecules encoding fatty-acyl desaturases and at least two exogenous nucleic acid molecules encoding fatty-acyl conjugases that catalyze the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA.

In yet a further embodiment, the fatty-acyl conjugase is a conjugase capable of utilizing a fatty acyl-CoA as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms.

In certain embodiments, the conjugase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Cydia pomonella*, *Cydia nigricana*, *Lobesia botrana*, *Myelois cribrella*, *Plodia interpunctella*, *Dendrolimus punctatus*, *Lampronia capitella*, *Spodoptera litura*, *Amyelois transitella*, *Manduca sexta*, *Bombyx mori*, *Calendula officinalis*, *Trichosanthes kirilowii*, *Punica granatum*, *Momordica charantia*, *Impatiens balsamina*, and *Epiphyas postvittana*. In exemplary embodiments, the conjugase comprises a sequence selected from GenBank Accession No. or Uniprot database: A0A059TBF5, A0A0M3L9E8, A0A0M3L9S4, A0A0M3LAH8, A0A0M3LAS8, A0A0M3LAH8, B6CBS4, XP_013183656.1, XP_004923568.2, ALA65425.1, NP_001296494.1, NP_001274330.1, Q4A181, Q75PL7, Q9FPP8, AY178444, AY178446, AF182521, AF182520, Q95UJ3.

In various embodiments described herein, the fatty alcohol forming acyl-CoA reductase, i.e., fatty alcohol forming fatty-acyl reductase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Agrotis segetum*, *Spodoptera littoralis*, or *Helicoverpa armigera*. In exemplary embodiments, the reductase comprises a sequence selected from GenBank Accession Nos. JX679210 and HG423128, and UniProt Accession No. 13PN86. In some embodiments, a nucleic acid sequence encoding a fatty-acyl reductase from organisms of the species *Agrotis segetum*, *Spodoptera littoralis*, or *Helicoverpa armigera* is codon optimized. In some embodiments, the reductase comprises a sequence set forth in SEQ ID NO: 1 from *Agrotis segetum*. In other embodiments, the reductase comprises a sequence set forth in SEQ ID NO: 2 from *Spodoptera littoralis*. In some embodiments, the reductase comprises a sequence selected from SEQ ID NOs: 3 and 32 from *Helicoverpa armigera*.

In certain embodiments, the fatty alcohol forming fatty-acyl reductase catalyzes the conversion of a mono- or poly-unsaturated fatty acyl-CoA into a fatty alcohol product selected from (Z)-3-hexenol, (Z)-3-nonenol, (Z)-5-decenol, (E)-5-decenol, (Z)-7-dodecenol, (E)-8-dodecenol, (Z)-8-dodecenol, (Z)-9-dodecenol, (Z)-9-tetradecenol, (Z)-9-hexadecenol, (Z)-11-tetradecenol, (Z)-7-hexadecenol, (Z)-11-hexadecenol, (E)-11-tetradecenol, or (Z,Z)-11,13-hexadecadienol, (11Z,13E)-hexadecadienol, (E,E)-8,10-dodecadienol, (E,Z)-7,9-dodecadienol, (Z)-13-octadecenol, or combinations thereof.

In some embodiments, the recombinant microorganism may express more than one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. Such recombinant microorganisms may be advantageously used to produce blends of various insect pheromones.

6

In addition to the biosynthetic pathway described in the first aspect above, the present application provides an additional biosynthetic pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol utilizing a saturated C₆-C₂₄ fatty acyl-ACP intermediate derived from a C₆-C₂₄ fatty acid. Accordingly, in a second aspect, the application relates to a recombinant microorganism capable of producing an unsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of C₆-C₂₄ fatty acid, wherein the recombinant microorganism expresses (a): at least one exogenous nucleic acid molecule encoding an acyl-ACP synthetase that catalyzes the conversion of a C₆-C₂₄ fatty acid to a corresponding saturated C₆-C₂₄ fatty acyl-ACP; (b) at least one exogenous nucleic acid molecule encoding a fatty-acyl-ACP desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-ACP to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP; (c) one or more endogenous or exogenous nucleic acid molecules encoding a fatty acid synthase complex that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP from (b) to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP with a two carbon elongation relative to the product of (b); (d): at least one exogenous nucleic acid molecule encoding a fatty aldehyde forming fatty-acyl reductase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP from (c) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde; and (e) at least one endogenous or exogenous nucleic acid molecule encoding a dehydrogenase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde C₆-C₂₄ from (d) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is an insect pheromone. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is a fragrance or flavoring agent. In some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an alcohol oxidase or an alcohol dehydrogenase, wherein the alcohol oxidase or alcohol dehydrogenase is capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from (e) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde. In some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an acetyl transferase capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from (e) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acetate.

In some embodiments, acyl-ACP synthetase is a synthetase capable of utilizing a fatty acid as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms.

In various embodiments described herein, the acyl-ACP synthetase, or the nucleic acid that encodes it, can be isolated from organisms of the species *Vibrio harveyi*, *Rhodotorula glutinis*, or *Yarrowia lipolytica*.

In some embodiments, the fatty-acyl-ACP desaturase is a soluble desaturase. In various embodiments described herein, the fatty-acyl-ACP desaturase, or the nucleic acid that encodes it, can be isolated from organisms of the species *Pelargonium hortorum*, *Asclepias syriaca*, or *Uncaria tomentosa*.

In some embodiments, the recombinant microorganism may express more than one exogenous nucleic acid molecule encoding a fatty-acyl desaturase that catalyzes the

conversion of a saturated C₆-C₂₄ fatty acyl-ACP to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP.

As described above, fatty acid elongation enzymes, i.e., a fatty acid synthase complex, can be utilized to extend the chain length of a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP by two additional carbons at the alpha carbon. In some embodiments, the two additional carbons are derived from endogenous malonyl-CoA. In one embodiment, the one or more nucleic acid molecules encoding a fatty acid synthase complex are endogenous nucleic acid molecules, i.e., the nucleic acid molecule(s) is/are native to the recombinant microorganism. In another embodiment, the one or more nucleic acid molecules encoding a fatty acid synthase complex are exogenous nucleic acid molecules.

In various embodiments described herein, the fatty aldehyde forming acyl-ACP reductase, i.e., fatty aldehyde forming fatty-acyl reductase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species can be isolated from organisms of the species *Pelargonium hortorum*, *Asclepias syriaca*, and *Uncaria tomentosa*.

In some embodiments, the recombinant microorganism may express more than one exogenous nucleic acid molecule encoding a fatty aldehyde forming fatty-acyl reductase that catalyzes the conversion of a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde. Such recombinant microorganisms may be advantageously used to produce blends of various insect pheromones. An exemplary blend according to the instant invention comprises of (Z)-11-hexadecenal (Z11-16:Ald) and (Z)-9-hexadecenal (Z9-16:Ald). In one embodiment, the ratio of the blend is 90:10, 91:9, 92:8, 93:7, 94:6, 95:5, 96:4, 97:3, 98:2, or 99:1 ratio of (Z)-11-hexadecenal (Z11-16:Ald) to (Z)-9-hexadecenal (Z9-16:Ald). In an exemplary embodiment, the blend is a 97:3 ratio of (Z)-11-hexadecenal (Z11-16:Ald) to (Z)-9-hexadecenal (Z9-16:Ald), corresponding to key components of the *Helicoverpa* female virgin.

As noted above, the recombinant microorganism according to the second aspect comprises at least one endogenous or exogenous nucleic acid molecule encoding a dehydrogenase capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde from (d) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. In one embodiment, the dehydrogenase is encoded by an endogenous nucleic acid molecule. In another embodiment, the dehydrogenase is encoded by an exogenous nucleic acid molecule. In exemplary embodiments, the endogenous or exogenous nucleic acid molecule encoding a dehydrogenase is isolated from organisms of the species *Saccharomyces cerevisiae*, *Escherichia coli*, *Yarrowia lipolytica*, or *Candida tropicalis*.

In addition to the biosynthetic pathway described in the first and second aspects above, the present application provides an additional biosynthetic pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol utilizing a saturated C₆-C₂₄ fatty acyl-ACP intermediate derived from a C₆-C₂₄ fatty acid. Accordingly, in a third aspect, the application relates to a recombinant microorganism capable of producing an unsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of C₆-C₂₄ fatty acid, wherein the recombinant microorganism expresses (a): at least one exogenous nucleic acid molecule encoding an acyl-ACP synthetase that catalyzes the conversion of a C₆-C₂₄ fatty acid to a corresponding saturated C₆-C₂₄ fatty acyl-ACP; (b) at least one exogenous nucleic acid molecule encoding a fatty-acyl-ACP desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-ACP to a corre-

sponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP; (c) at least one exogenous fatty acyl-ACP thioesterase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP from (b) to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acid; (d) one or more endogenous or exogenous nucleic acid molecules encoding an elongase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA derived from CoA activation of the mono- or poly-unsaturated C₆-C₂₄ fatty acid from (c) to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA with a two carbon or greater elongation relative to the product of (c); and (e): at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA from (d) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is an insect pheromone. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is a fragrance or flavoring agent. In some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an alcohol oxidase or an alcohol dehydrogenase, wherein the alcohol oxidase or alcohol dehydrogenase is capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from (e) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde. In some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an acetyl transferase capable of catalyzing the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from (e) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acetate.

In some embodiments according to this third aspect, a fatty acyl-ACP thioesterase can be utilized to convert a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-ACP into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acid. In a particular embodiment, soluble fatty acyl-ACP thioesterases can be used to release free fatty acids for reactivation to a CoA thioester. Fatty acyl-ACP thioesterases including Q41635, Q39473, P05521.2, AEM72519, AEM72520, AEM72521, AEM72523, AAC49784, CAB60830, EER87824, EER96252, ABN54268, AAO77182, CAH09236, ACL08376, and homologs thereof may be used. In some embodiments, the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA may serve as a substrate for an elongase, which can be utilized to extend the chain length of a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA by two additional carbons at the alpha carbon. In some embodiments, the two additional carbons are derived from endogenous malonyl-CoA.

As described above, in some embodiments, the recombinant microorganism according to the first, second, or third aspect further comprises at least one endogenous or exogenous nucleic acid molecule encoding an alcohol oxidase capable of catalyzing the conversion of a mono- or poly-unsaturated C₆-C₂₄ fatty alcohol into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde. In certain embodiments, the alcohol oxidase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Candida boidinii*, *Komagataella pastoris*, *Tanacetum vulgare*, *Simmondsia chinensis*, *Arabidopsis thaliana*, *Lotus japonicas*, or *Candida tropicalis*. In exemplary embodiments, the alcohol oxidase comprises a sequence selected from GenBank Accession Nos. Q00922, F2QY27, Q6QIR6, Q8LDP0, and L7VFFV2.

In alternative embodiments, the fatty alcohol may be converted into a fatty aldehyde using chemical methods, including but not limited to, the use of TEMPO-bleach, TEMPO-copper-air, TEMPO-PhI(OAc)₂, Swern oxidation, or noble metal-air.

As described above, in some embodiments, the recombinant microorganism according to the first or second aspect further comprises at least one endogenous or exogenous nucleic acid molecule encoding an acetyl transferase capable of catalyzing the conversion of a C₆-C₂₄ fatty alcohol into a corresponding C₆-C₂₄ fatty acetate. In certain embodiments, the acetyl transferase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Saccharomyces cerevisiae*, *Danaus plexippus*, *Heliotis virescens*, *Bombyx mori*, *Agrotis ipsilon*, *Agrotis segetum*, *Euonymus alatus*. In exemplary embodiments, the acetyl transferase comprises a sequence selected from GenBank Accession Nos. AY242066, AY242065, AY242064, AY242063, AY242062, EHJ65205, ACX53812, NP_001182381, EHJ65977, EHJ68573, KJ579226, GU594061.

In alternative embodiments, the fatty alcohol may be converted into a fatty acetate using chemical methods, e.g., via chemical catalysis utilizing a chemical agent such as acetyl chloride, acetic anhydride, butyryl chloride, butyric anhydride, propanoyl chloride and propionic anhydride.

In some embodiments, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may further be engineered to express one or more nucleic acids encoding protein or polypeptide which, when expressed, is toxic to an insect. Exemplary toxicant producing genes suitable for the present disclosure can be obtained from entomopathogenic organism, such as *Bacillus thuringiensis*, *Pseudomonas aeruginosa*, *Serratia marcescens*, and members of the genus *Streptomyces*. In an exemplary embodiment, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may further be engineered to express a nucleic acid encoding a *Bacillus thuringiensis* ("Bt") toxin. In additional or alternative embodiments, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may further be engineered to express a nucleic acid encoding other toxic proteins such as spider venom.

In some embodiments, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may further be engineered to express an RNAi molecule which, when expressed, produces an oligonucleotide that is toxic to an insect.

In some embodiments, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may further be engineered to express a metabolic pathway which, when expressed, produces a small molecule that is toxic to an insect. Non-limiting examples of toxic small molecules include azadirachtin, spinosad, avermectin, pyrethrins, and various terpenoids.

In various embodiments described herein, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may be a eukaryotic microorganism, such as a yeast, a filamentous fungi, or an algae, or alternatively, a prokaryotic microorganism, such as a bacterium. For instance, suitable host cells can include cells of a genus

selected from the group consisting of *Yarrowia*, *Candida*, *Saccharomyces*, *Pichia*, *Hansenula*, *Clostridium*, *Zymomonas*, *Escherichia*, *Salmonella*, *Rhodococcus*, *Pseudomonas*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Alcaligenes*, *Klebsiella*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, *Brevibacterium*, and *Streptomyces*.

In some embodiments, the recombinant microorganism comprising a biosynthesis pathway for the production of an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate is a yeast. Examples of suitable yeasts include yeasts of a genus selected from the group consisting of *Yarrowia*, *Candida*, *Saccharomyces*, *Pichia*, *Hansenula*, *Kluyveromyces*, *Issatchenkia*, *Zygosaccharomyces*, *Debaryomyces*, *Schizosaccharomyces*, *Pachysolen*, *Cryptococcus*, *Trichosporon*, *Rhodotorula*, or *Myxozyma*. In certain embodiments, the yeast is an oleaginous yeast. Exemplary oleaginous yeasts suitable for use in the present disclosure include members of the genera *Yarrowia*, *Candida*, *Rhodotorula*, *Rhodospiridium*, *Cryptococcus*, *Trichosporon*, and *Lipomyces*, including, but not limited to the species of *Yarrowia lipolytica*, *Candida tropicalis*, *Rhodospiridium toruloides*, *Lipomyces starkey*, *L. lipoferus*, *C. revkaufi*, *C. pulcherrima*, *C. utilis*, *Rhodotorula minuta*, *Trichosporon pullans*, *T. cutaneum*, *Cryptococcus curvatus*, *R. glutinis*, and *R. graminis*.

As will be understood in the art, endogenous enzymes can convert critical substrates and/or intermediates upstream of or within the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate biosynthesis pathway into unwanted by-products. Accordingly, in some embodiments, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that competes with the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate biosynthesis pathway.

In one embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes the conversion of a fatty acid into a ω -hydroxyfatty acid. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more enzyme selected from XP_504406, XP_504857, XP_504311, XP_500855, XP_500856, XP_500402, XP_500097, XP_501748, XP_500560, XP_501148, XP_501667, XP_500273, BAA02041, CAA39366, CAA39367, BAA02210, BAA02211, BAA02212, BAA02213, BAA02214, AAO73952, AAO73953, AAO73954, AAO73955, AAO73956, AAO73958, AAO73959, AAO73960, AAO73961, AAO73957, XP_002546278, or homologs thereof. In the context of a recombinant bacterial microorganism, the recombinant bacterial microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more enzyme selected from BAM49649, AAB80867, AAB17462, ADL27534, AAU24352, AAA87602, CAA34612, ABM17701, AAA25760, CAB51047, AAC82967, WP_011027348, or homologs thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes the conversion of a fatty acyl-CoA into α,β -enoyl-CoA. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more enzyme selected from CAA04659, CAA04660, CAA04661, CAA04662, CAA04663, CAG79214, AAA34322,

11

AAA34361, AAA34363, CAA29901, BAA04761, AAA34891, or homologs thereof. In the context of a recombinant bacterial microorganism, the recombinant bacterial microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more enzyme selected from AAB08643, CAB15271, BAN55749, CAC44516, ADK16968, AEI37634, WP_000973047, WP_025433422, WP_035184107, WP_026484842, CEL80920, WP_026818657, WP_005293707, WP_005883960, or homologs thereof.

In embodiments where the recombinant microorganism is a yeast microorganism, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more enzyme involved in peroxisome assembly and/or peroxisome enzyme import. The recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more enzyme selected from XP_505754, XP_501986, XP_501311, XP_504845, XP_503326, XP_504029, XP_002549868, XP_002547156, XP_002545227, XP_002547350, XP_002546990, EIW11539, EIW08094, EIW11472, EIW09743, EIW08286, or homologs thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous reductase or desaturase enzymes that interferes with the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate, i.e., catalyzes the conversion of a pathway substrate or product into an unwanted by-product.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous alcohol oxidase or alcohol dehydrogenase enzymes that catalyzes the unwanted conversion of the desired product, e.g., unsaturated C₆-C₂₄ fatty alcohol into a corresponding unsaturated C₆-C₂₄ fatty aldehyde.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that competes with the biosynthesis pathway for one or more unsaturated fatty acyl-CoA intermediates. In one embodiment, the one or more endogenous enzymes comprise one or more diacylglycerol acyltransferases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more diacylglycerol acyltransferases selected from the group consisting of YALIOE32769g, YALIOD07986g and CTRG_06209, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more glycerolphospholipid acyltransferases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more glycerolphospholipid acyltransferases selected from the group consisting of YALIOE16797g and CTG_04390, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more acyl-CoA/sterol acyltransferases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more acyl-CoA/sterol acyltransferases selected from the group consisting of YALIOF06578g, CTRG_01764 and CTRG_01765, or homolog thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes

12

a reaction in a pathway that oxidizes fatty aldehyde intermediates. In one embodiment, the one or more endogenous enzymes comprise one or more fatty aldehyde dehydrogenases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more fatty aldehyde dehydrogenases selected from the group consisting of YALIOA17875g, YALIOE15400g, YALIOB01298g, YALIOF23793g, CTRG_05010 and CTRG_04471, or homolog thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that consumes fatty acetate products. In one embodiment, the one or more endogenous enzymes comprise one or more sterol esterases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more sterol esterases selected from the group consisting of YALIOE32035g, YALIOE00528g, CTRG_01138, CTRG_01683 and CTRG_04630, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more triacylglycerol lipases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more triacylglycerol lipases selected from the group consisting of YALIOD17534g, YALIOF10010g, CTRG_00057 and CTRG_06185, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more monoacylglycerol lipases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more monoacylglycerol lipases selected from the group consisting of YALIOC14520g, CTRG_03360 and CTRG_05049, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more extracellular lipases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more extracellular lipases selected from the group consisting of YALIOA20350g, YALIOD19184g, YALIOB09361g, CTRG_05930, CTRG_04188, CTRG_02799, CTRG_03052 and CTRG_03885, or homolog thereof.

In embodiments where the recombinant microorganism is a yeast microorganism, one or more of the exogenous unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate pathway genes encodes an enzyme that is localized to a yeast compartment selected from the group consisting of the cytosol, the mitochondria, or the endoplasmic reticulum. In an exemplary embodiment, one or more of the exogenous pathway genes encodes an enzyme that is localized to the endoplasmic reticulum. In another embodiment, at least two exogenous pathway genes encode an enzyme that is localized to the endoplasmic reticulum. In yet another embodiment, all exogenous pathway genes encodes an enzyme that is localized to the endoplasmic reticulum.

In a fourth aspect, the present application provides methods of producing an unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate using a recombinant microorganism as described herein. In one embodiment, the method includes cultivating the recombinant microorganism in a culture medium containing a feedstock providing a carbon source

until the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate is produced and optionally, recovering the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate. Once produced, the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate may be isolated from the fermentation medium using various methods known in the art including, but not limited to, distillation, membrane-based separation gas stripping, solvent extraction, and expanded bed adsorption.

In some embodiments, the recombinant microorganism, e.g., a yeast, may be recovered and produced in dry particulate form. In embodiments involving yeast, the yeast may be dried to produce powdered yeast. In some embodiments, the process for producing powdered yeast comprises spray drying a liquid yeast composition in air, optionally followed by further drying. In some embodiments, the recombinant microorganism composition will comprise the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate when dried.

As described herein, preferred recombinant microorganisms of the disclosure will have the ability to utilize alkanes and fatty acids as carbon sources. However, as will be understood in the art, a variety of carbon sources may be utilized, including but not limited to, various sugars (e.g., glucose, fructose, or sucrose), glycerol, alcohols (e.g., ethanol), organic acids, lignocellulose, proteins, carbon dioxide, carbon monoxide, as well as the aforementioned alkanes and fatty acids. In an exemplary embodiment, the recombinant microorganism will convert the carbon source to the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate under aerobic conditions.

As highlighted above, the present application provides methods of producing one or more unsaturated C₆-C₂₄ fatty alcohols, aldehydes, or acetates using a recombinant microorganism as described herein. In some embodiments, the product is an insect pheromone. As will be appreciated by the skilled artisan equipped with the instant disclosure, a variety of different exogenous and endogenous enzymes can be expressed in a recombinant host microorganism to produce a desired insect pheromone. Exemplary insect pheromones in the form of fatty alcohols, fatty aldehydes, or fatty acetates capable of being generated using the recombinant microorganisms and methods described herein include, but are not limited to, (Z)-11-hexadecenal, (Z)-11-hexadecenyl acetate, (Z)-9-tetradecenyl acetate, (Z,Z)-11,13-hexadecadienal, (9Z,11E)-hexadecadienal, (E,E)-8,10-dodecadien-1-ol, (7E,9Z)-dodecadienyl acetate, (Z)-3-nonen-1-ol, (Z)-5-decen-1-ol, (Z)-5-decenyl acetate, (E)-5-decen-1-ol, (E)-5-decenyl acetate, (Z)-7-dodecen-1-ol, (Z)-7-dodecenyl acetate, (E)-8-dodecen-1-ol, (E)-8-dodecenyl acetate, (Z)-8-dodecen-1-ol, (Z)-8-dodecenyl acetate, (Z)-9-dodecen-1-ol, (Z)-9-dodecenyl acetate, (Z)-9-tetradecen-1-ol, (Z)-11-tetradecen-1-ol, (Z)-11-tetradecenyl acetate, (E)-11-tetradecen-1-ol, (E)-11-tetradecenyl acetate, (Z)-7-hexadecen-1-ol, (Z)-7-hexadecenyl acetate, (Z)-9-hexadecen-1-ol, (Z)-9-hexadecenyl acetate, (Z)-11-hexadecen-1-ol, (Z)-13-octadecen-1-ol, and (Z)-13-octadecenal.

In a fifth aspect, the present application provides compositions comprising one or more of the insect pheromone-producing recombinant microorganisms described herein. In certain embodiments, the composition may further comprise one or more insect pheromones produced by the recombinant microorganism. In further embodiments, the may additionally comprise one or more toxic proteins or polypeptides produced by the recombinant microorganism.

BRIEF DESCRIPTION OF DRAWINGS

Illustrative embodiments of the disclosure are illustrated in the drawings, in which:

FIG. 1 illustrates the conversion of a saturated fatty acyl-CoA to an unsaturated fatty alcohol.

FIG. 2 illustrates the conversion of a saturated fatty acid to a mono- or poly-unsaturated fatty aldehyde, alcohol, or acetate.

FIG. 3 illustrates an additional pathway for the conversion of a saturated fatty acid to a mono- or poly-unsaturated fatty aldehyde, alcohol, or acetate.

FIG. 4 illustrates a pathway for the conversion of a saturated fatty acid to various trienes, dienes, epoxides, and odd-numbered pheromones.

FIG. 5 shows Z11-hexadecenol production from W303A and BY4742 ΔPOX1. Strain expressing empty vector (EV), *S. littoralis* reductase (FAR-SL), *H. armigera* reductase (FAR-HA), *A. segetum* reductase (FAR-AS). Error bars represent standard deviation derived from N=2 biologically-independent samples.

FIG. 6A-FIG. 6B shows sample chromatograms of biotransformation product of Z11-hexadecenoic acid using *S. cerevisiae* expressing either an empty vector (FIG. 6A), or *Helicoverpa armigera* alcohol-forming reductase (FIG. 6B). Black lines: no substrate added. Purple line: Z11-hexadecenoic acid was added as substrate.

FIG. 7A-FIG. 7B shows a comparison of GC-MS fragmentation pattern of Z11-hexadecenol authentic compound (FIG. 7A), and Z11-hexadecenol biologically derived (FIG. 7B).

FIG. 8 shows biomass at the time of harvesting for product analysis of W303A (wild type) and BY4742 ΔPOX1 (beta-oxidation deletion mutant). Strain expressing empty vector (EV), *S. littoralis* reductase (FAR-SL), *H. armigera* reductase (FAR-HA), *A. segetum* reductase (FAR-AS). Error bars represent standard deviation derived from N=2 biologically-independent samples.

FIG. 9 shows a Z11-hexadecenol calibration curve constructed using an authentic standard. The samples were generated with the extraction and analysis method described in Materials and Methods of Example 3. Error bars represent standard deviation derived from N=3 samples.

FIG. 10 shows a pOLE1 cassette comprising an extended OLE1 promoter sequence (light yellow), OLE1 promoter (orange), OLE1 leader sequence (dark grey), a synthon such as an insect desaturase sequence (light grey), and the VSP13 terminator sequence (blue).

FIG. 11A-FIG. 11E shows validation of the pOLE1 cassette, and complementation assay. FIG. 11A: YPD+palmitoleic acid; FIG. 11B: YPD-palmitoleic acid; FIG. 11C: CM-Ura glucose+palmitoleic acid; FIG. 11D: CM-Ura glucose-palmitoleic acid; FIG. 11E: Map of strains in FIG. 11A-FIG. 11D. Dasher=GFP synthon.

FIG. 12A shows complementation of ΔOLE1 growth without UFA on YPD.

FIG. 12B shows complementation of ΔOLE1 growth without UFA on CM-Ura glucose.

FIG. 13A shows the full fatty acid spectrum of a ΔOLE1 strain expressing: *S. cerevisiae* OLE1 desaturase (blue), chimeric *T. ni* desaturase (red).

FIG. 13B shows a focused fatty acid spectrum within 5.5-min-8-min retention time of *S. cerevisiae* ΔOLE1 strain expressing *S. cerevisiae* OLE1 desaturase (red) and chimeric *T. ni* desaturase (blue).

FIG. 14A-FIG. 14B shows a comparison of GC-MS fragmentation pattern of (Z)-11-hexadecenoic acid from an authentic compound (FIG. 14A) and biologically derived (FIG. 14B).

FIG. 15 shows C16 fatty alcohol production from ΔOLE1 expressing various fatty alcohol pathway variants in culture

15

supplemented with palmitic and palmitoleic acid. Error bars represent 5% uncertainty of metabolite quantification accuracy.

FIG. 16 shows representative chromatograms of biotransformation product C16 fatty acids using *S. cerevisiae* expressing fatty alcohol pathways TN_desat-HA_reduc when fed with palmitic acid (black) and when fed with palmitic and palmitoleic acids (orange). Profile of a negative control strain (harboring an empty vector) fed with palmitic acid (purple).

FIG. 17 shows that (Z)-11-hexadecenoic acid was detected in the cell pellets of *S. cerevisiae* expressing fatty alcohol pathways TN_desat-SL_reduc (blue), SC_desat-HA_reduc (red), TN_desat-HA_reduc (green), SC_desat-SL_reduc (pink).

FIG. 18 shows C16 fatty alcohol production from ΔOLE1 expressing various fatty alcohol pathway variants in culture supplemented with palmitic acid only. Error bars represent 5% uncertainty of metabolite quantification accuracy.

FIG. 19A-FIG. 19C shows detection of (Z)-11-hexadecanol. FIG. 19A: Fragmentation pattern of an authentic standard. The m/z 297.3 was used in follow up experiments to selectively detect the alcohol. To also detect the internal standard, the masses 208 and 387.3 were included too. FIG. 19B: In addition to the detection of the specific mass fragment, the retention time was used as second stage confirmation. The retention time is 6.22. FIG. 19C: Comparison of the two different regioisomers 9Z- and 11Z-hexadecanol when detected in SIM mode (297.3) with the same method.

FIG. 20 shows pXICL expression cassette architecture. The *C. albicans* OLE1 leader-*A. segetum* desaturase fusion is also shown.

FIG. 21A-FIG. 21D shows mCherry control integration. FIG. 21A: Negative (water-only) control transformation plate. FIG. 21B: pPV0137 mCherry transformation plate. FIG. 21C: Patch plates from negative control clones. FIG. 21D: Patch plates from pPV0137 clones.

FIG. 22 shows integration efficiency as a function of total observed colonies. A control plate with no DNA added to the transformation was observed to have 350 colonies (indicated by orange line). The fraction of clones confirmed to be positive integrants is positively correlated with total colony count. A sharp increase is observed above 6,000 total colonies. The data suggests that the presence of positive integrants increases the observed background growth. For some transformations the efficiency was high enough that the background population was small relative to the positive integrant population.

FIG. 23 shows a chromatogram overlay of *Candida tropicalis* SPV053 strains. Compared to the mCherry (red) control experiment a clear peak at 6.22 min is observable for the *A. transitella* (blue) and *H. zea* (green) desaturase. Therefore, the formation of Z-11-hexadecenoic acid is only observable in strains expressing an active Z11-desaturase.

FIG. 24A-FIG. 24E shows confirmation of the 11Z-regioisomer. FIG. 24A: The specific peak with an ion fragment of 245 m/z was only observed in *C. tropicalis* SPV053 expressing either the Z11-desaturase from *A. transitella* or *H. zea*. FIG. 24B: The fragmentation patterns of the authentic standard. FIG. 24D: The fragmentation patterns of the newly formed compound in samples with expressed desaturase from *H. zea* match those of the standard. FIG. 24E: The fragmentation patterns of the newly formed compound in samples with expressed desaturase from *A. transitella* match those of the standard. FIG. 24C: The fragmen-

16

tation patterns of the mCherry control significantly differ from those of FIG. 24B, FIG. 24D and FIG. 24E.

FIG. 25A-FIG. 25B shows a GC-FID chromatogram of different *C. tropicalis* SPV053 strains incubated with methyl tetradecanoate. FIG. 25A: Overall spectrum. The occurrence of the Z11-C16:1 peak is observable for the strains expressing the Z11-desaturases from *A. transitella* and *H. zea*. FIG. 25B: Zoom of the C14 to C18 area. A new peak is visible at 4.8 min, which could correspond to Z11-C14:1. Another peak near Z9-C18:1 is also visible, which could correspond to Z11-C18:1.

FIG. 26 shows only codon optimized *H. zea* desaturase variants produce detectable Z11-hexadecenoic acid in SPV140 screen. control=pPV101 integrants of SPV140, *T. ni* native=*T. ni* Z11 desaturase with native codon usage (pPV195), *T. ni* HS opt=*T. ni* Z11 desaturase with *Homo sapiens* codon optimization (pPV196), *T. ni* HS opt Y1 leader=*T. ni* Z11 desaturase with *Homo sapiens* codon optimization and swapped *Y. lipolytica* OLE1 leader sequence (pPV197), *H. zea* native=*H. zea* Z11 desaturase with native codon usage (pPV198), *H. zea* HS opt=*H. zea* Z11 desaturase with *Homo sapiens* codon optimization (pPV199), *H. zea* HS opt Y1 leader=*H. zea* Z11 desaturase with *Homo sapiens* codon optimization and swapped *Y. lipolytica* OLE1 leader sequence (pPV200), *A. transitella* native=*A. transitella* Z11 desaturase with native codon usage (pPV201). All data average of 3 biological replicates. Error bars represent standard deviation.

FIG. 27 shows only codon optimized *H. zea* desaturase variants produce detectable Z11-hexadecenoic acid in SPV300 screen. Labels indicate parent strain and plasmid of desaturase expression cassette. pPV101=hrGFP control, pPV198=*H. zea* Z11 desaturase with native codon usage, pPV199=*H. zea* Z11 desaturase with *Homo sapiens* codon optimization, pPV200=*H. zea* Z11 desaturase with *Homo sapiens* codon optimization and swapped *Y. lipolytica* OLE1 leader sequence, pPV201=*A. transitella* Z11 desaturase with native codon usage.

FIG. 28 shows final cell densities for desaturase screen in SPV140 and SPV300 backgrounds. SPV300 strains with integrated desaturase cassettes grew to higher cell densities.

FIG. 29 shows individual isolate Z11-hexadecenoic acid titers for SPV140 and SPV300 strains expressing *H. zea* Z11 desaturase with *H. sapiens* codon optimization.

FIG. 30 shows a chromatogram overlay of extracted metabolites for Z11-16OH producing strain (SPV0490) versus control strain (SPV0488) of *Candida viswanathii* (*tropicalis*).

FIG. 31 illustrates pathways that can be deleted or disrupted to reduce or eliminate competition with the biosynthesis pathway for the production of a mono- or polyunsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate.

FIG. 32A-FIG. 32B shows Z9-16OH and Z11-16OH titers in YPD (FIG. 32A) and Semi-Defined C:N=80 (FIG. 32B) media for pEXP clones. Ten isolates expressing the *H. zea* desaturase under the TEF promoter and *H. armigera* reductase under the EXP promoter from two independent competent cell preparations (Comp. Cell Preparation 1, Comp. Cell Preparation 2) were compared to a parental negative control (SPV300) and a desaturase only negative control (SPV459 Hz_desat only). Error bars represent the SEM (standard error of the mean) measured from technical replicates for each strain and condition (N=2). *One replicate from Clone 5 and Clone 18 under the Semi-Defined C:N=80 condition was lost during sample work-up so the

titers for that condition are from a single data point (N=1, Comp. Cell Preparation 1 Clone 18 and Comp. Cell Preparation 2 Clone 5).

FIG. 33A-FIG. 33B shows profiles of 16-carbon fatty acid species in YPD (FIG. 33A) and Semi-Defined C:N=80 (FIG. 33B) media for pEXP1 clones. The 16-carbon lipid profiles of 5 select clones expressing the *H. zea* desaturase under the TEF promoter and *H. armigera* reductase under the EXP promoter are compared to a parental negative control (SPV300) and a desaturase only negative control (SPV459 Hz_desat only). Error bars represent the SEM (standard error of the mean) measured from technical replicates for each strain and condition (N=2).

FIG. 34 shows Z9-16OH and Z11-16OH titers in Semi-Defined C:N=80 media for pTAL1 clones. Nine isolates expressing the *H. zea* desaturase under the TEF promoter and *H. armigera* reductase under the TAL promoter were compared to a parental negative control (SPV300) and positive Bdr pathway controls using the EXP promoter to drive *H. armigera* FAR expression (SPV575, SPV578). Error bars represent the SEM (standard error of the mean) measured from technical replicates for each strain and condition (N=2).

FIG. 35 shows profiles of 16-carbon fatty acid species in Semi-Defined C:N=80 medium for pTAL1 clones. The 16-carbon lipid profiles of 5 select clones expressing the *H. zea* desaturase under the TEF promoter and *H. armigera* reductase under the EXP promoter are compared to a parental negative control (SPV300) and positive Bdr pathway controls using the EXP promoter to drive *H. armigera* FAR expression (SPV575, SPV578). Error bars represent the SEM (standard error of the mean) measured from technical replicates for each strain and condition (N=2). * indicates clones for which one of the replicates was lost during sample processing, N=1.

FIG. 36 shows full Bdr pathway pTAL1 screen (strains expressing *H. zea* Z11 desaturase (pTEF) and *H. armigera* FAR) full lipid profiles in Semi-Defined C:N=80 medium after 48 hours of bioconversion. Error bars represent the SEM (standard error of the mean) measured from technical replicates for each strain and condition (N=2). * indicates clones for which one of the replicates was lost during sample processing, N=1.

FIG. 37A-FIG. 37B shows SPV471 (H222 Δ P Δ A Δ F expressing native *Y. lipolytica* OLE1 and *H. armigera* FAR) Z9-16OH (FIG. 37A) and fatty acid (FIG. 37B) titers in Semi-Defined C:N=80 medium after 24 hours of bioconversion. Error bars represent the SEM (standard error of the mean) measured from technical replicates for each strain and condition (N=2).

FIG. 38 shows SPV471 (H222 Δ P Δ A Δ F expressing native *Y. lipolytica* OLE1 and *H. armigera* FAR) full lipid profiles in Semi-Defined C:N=80 medium after 24 hours of bioconversion.

FIG. 39A-FIG. 39B shows SPV471 (H222 Δ P Δ A Δ F expressing native *Y. lipolytica* OLE1 and *H. armigera* FAR) Z9-16OH (FIG. 39A) and Z9-16Acid (FIG. 39B) titer time courses. Bioconversion of 16Acid was conducted in Semi-Defined C:N=80 medium using a methyl palmitate (16Acid) substrate.

FIG. 40 shows a scheme illustrating exemplary means by which the recombinant microorganisms disclosed herein can be used to create synthetic blends of insect pheromones.

A sequence listing for SEQ ID NO: 1-SEQ ID NO: 38 is part of this application and is incorporated by reference herein. The sequence listing is provided at the end of this document.

DETAILED DESCRIPTION

Definitions

The following definitions and abbreviations are to be used for the interpretation of the disclosure.

As used herein and in the appended claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to “a pheromone” includes a plurality of such pheromones and reference to “the microorganism” includes reference to one or more microorganisms, and so forth.

As used herein, the terms “comprises,” “comprising,” “includes,” “including,” “has,” “having,” “contains,” “containing,” or any other variation thereof, are intended to cover a non-exclusive inclusion. A composition, mixture, process, method, article, or apparatus that comprises a list of elements is not necessarily limited to only those elements but may include other elements not expressly listed or inherent to such composition, mixture, process, method, article, or apparatus. Further, unless expressly stated to the contrary, “or” refers to an inclusive “or” and not to an exclusive “or.”

The terms “about” and “around,” as used herein to modify a numerical value, indicate a close range surrounding that explicit value. If “X” were the value, “about X” or “around X” would indicate a value from 0.9X to 1.1X, or, in some embodiments, a value from 0.95X to 1.05X. Any reference to “about X” or “around X” specifically indicates at least the values X, 0.95X, 0.96X, 0.97X, 0.98X, 0.99X, 1.01X, 1.02X, 1.03X, 1.04X, and 1.05X. Thus, “about X” and “around X” are intended to teach and provide written description support for a claim limitation of, e.g., “0.98X.”

As used herein, the terms “microbial,” “microbial organism,” and “microorganism” include any organism that exists as a microscopic cell that is included within the domains of archaea, bacteria or eukarya, the latter including yeast and filamentous fungi, protozoa, algae, or higher Protista. Therefore, the term is intended to encompass prokaryotic or eukaryotic cells or organisms having a microscopic size and includes bacteria, archaea, and eubacteria of all species as well as eukaryotic microorganisms such as yeast and fungi. Also included are cell cultures of any species that can be cultured for the production of a chemical.

As described herein, in some embodiments, the recombinant microorganisms are prokaryotic microorganism. In some embodiments, the prokaryotic microorganisms are bacteria. “Bacteria”, or “eubacteria”, refers to a domain of prokaryotic organisms. Bacteria include at least eleven distinct groups as follows: (1) Gram-positive (gram+) bacteria, of which there are two major subdivisions: (1) high G+C group (*Actinomycetes*, *Mycobacteria*, *Micrococcus*, others) (2) low G+C group (*Bacillus*, *Clostridia*, *Lactobacillus*, *Staphylococci*, *Streptococci*, *Mycoplasmas*); (2) Proteobacteria, e.g., Purple photosynthetic+non-photosynthetic Gram-negative bacteria (includes most “common” Gram-negative bacteria); (3) Cyanobacteria, e.g., oxygenic phototrophs; (4) Spirochetes and related species; (5) Planctomyces; (6) *Bacteroides*, *Flavobacteria*; (7) *Chlamydia*; (8) Green sulfur bacteria; (9) Green non-sulfur bacteria (also anaerobic phototrophs); (10) Radioresistant micrococci and relatives; (11) *Thermotoga* and *Thermosipho* thermophiles.

“Gram-negative bacteria” include cocci, nonenteric rods, and enteric rods. The genera of Gram-negative bacteria include, for example, *Neisseria*, *Spirillum*, *Pasteurella*, *Brucella*, *Yersinia*, *Francisella*, *Haemophilus*, *Bordetella*, *Escherichia*, *Salmonella*, *Shigella*, *Klebsiella*, *Proteus*, *Vibrio*, *Pseudomonas*, *Bacteroides*, *Acetobacter*, *Aerobacter*, *Agrobacterium*, *Azotobacter*, *Spirilla*, *Serratia*, *Vibrio*, *Rhizobium*, *Chlamydia*, *Rickettsia*, *Treponema*, and *Fusobacterium*.

“Gram positive bacteria” include cocci, nonsporulating rods, and sporulating rods. The genera of gram positive bacteria include, for example, *Actinomyces*, *Bacillus*, *Clostridium*, *Corynebacterium*, *Erysipelothrix*, *Lactobacillus*, *Listeria*, *Mycobacterium*, *Myxococcus*, *Nocardia*, *Staphylococcus*, *Streptococcus*, and *Streptomyces*.

The term “recombinant microorganism” and “recombinant host cell” are used interchangeably herein and refer to microorganisms that have been genetically modified to express or to overexpress endogenous enzymes, to express heterologous enzymes, such as those included in a vector, in an integration construct, or which have an alteration in expression of an endogenous gene. By “alteration” it is meant that the expression of the gene, or level of a RNA molecule or equivalent RNA molecules encoding one or more polypeptides or polypeptide subunits, or activity of one or more polypeptides or polypeptide subunits is up regulated or down regulated, such that expression, level, or activity is greater than or less than that observed in the absence of the alteration. For example, the term “alter” can mean “inhibit,” but the use of the word “alter” is not limited to this definition. It is understood that the terms “recombinant microorganism” and “recombinant host cell” refer not only to the particular recombinant microorganism but to the progeny or potential progeny of such a microorganism. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

The term “expression” with respect to a gene sequence refers to transcription of the gene and, as appropriate, translation of the resulting mRNA transcript to a protein. Thus, as will be clear from the context, expression of a protein results from transcription and translation of the open reading frame sequence. The level of expression of a desired product in a host cell may be determined on the basis of either the amount of corresponding mRNA that is present in the cell, or the amount of the desired product encoded by the selected sequence. For example, mRNA transcribed from a selected sequence can be quantitated by qRT-PCR or by Northern hybridization (see Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press (1989)). Protein encoded by a selected sequence can be quantitated by various methods, e.g., by ELISA, by assaying for the biological activity of the protein, or by employing assays that are independent of such activity, such as western blotting or radioimmunoassay, using antibodies that recognize and bind the protein. See Sambrook et al., 1989, *supra*.

The term “polynucleotide” is used herein interchangeably with the term “nucleic acid” and refers to an organic polymer composed of two or more monomers including nucleotides, nucleosides or analogs thereof, including but not limited to single stranded or double stranded, sense or antisense deoxyribonucleic acid (DNA) of any length and, where appropriate, single stranded or double stranded, sense or antisense ribonucleic acid (RNA) of any length, including

siRNA. The term “nucleotide” refers to any of several compounds that consist of a ribose or deoxyribose sugar joined to a purine or a pyrimidine base and to a phosphate group, and that are the basic structural units of nucleic acids.

The term “nucleoside” refers to a compound (as guanosine or adenosine) that consists of a purine or pyrimidine base combined with deoxyribose or ribose and is found especially in nucleic acids. The term “nucleotide analog” or “nucleoside analog” refers, respectively, to a nucleotide or nucleoside in which one or more individual atoms have been replaced with a different atom or with a different functional group. Accordingly, the term polynucleotide includes nucleic acids of any length, DNA, RNA, analogs and fragments thereof. A polynucleotide of three or more nucleotides is also called nucleotidic oligomer or oligonucleotide.

It is understood that the polynucleotides described herein include “genes” and that the nucleic acid molecules described herein include “vectors” or “plasmids.” Accordingly, the term “gene”, also called a “structural gene” refers to a polynucleotide that codes for a particular sequence of amino acids, which comprise all or part of one or more proteins or enzymes, and may include regulatory (non-transcribed) DNA sequences, such as promoter sequences, which determine for example the conditions under which the gene is expressed. The transcribed region of the gene may include untranslated regions, including introns, 5'-untranslated region (UTR), and 3'-UTR, as well as the coding sequence.

The term “enzyme” as used herein refers to any substance that catalyzes or promotes one or more chemical or biochemical reactions, which usually includes enzymes totally or partially composed of a polypeptide or polypeptides, but can include enzymes composed of a different molecule including polynucleotides.

As used herein, the term “non-naturally occurring,” when used in reference to a microorganism organism or enzyme activity of the disclosure, is intended to mean that the microorganism organism or enzyme has at least one genetic alteration not normally found in a naturally occurring strain of the referenced species, including wild-type strains of the referenced species. Genetic alterations include, for example, modifications introducing expressible nucleic acids encoding metabolic polypeptides, other nucleic acid additions, nucleic acid deletions and/or other functional disruption of the microorganism’s genetic material. Such modifications include, for example, coding regions and functional fragments thereof, for heterologous, homologous, or both heterologous and homologous polypeptides for the referenced species. Additional modifications include, for example, non-coding regulatory regions in which the modifications alter expression of a gene or operon. Exemplary non-naturally occurring microorganism or enzyme activity includes the hydroxylation activity described above.

The term “exogenous” as used herein with reference to various molecules, e.g., polynucleotides, polypeptides, enzymes, etc., refers to molecules that are not normally or naturally found in and/or produced by a given yeast, bacterium, organism, microorganism, or cell in nature.

On the other hand, the term “endogenous” or “native” as used herein with reference to various molecules, e.g., polynucleotides, polypeptides, enzymes, etc., refers to molecules that are normally or naturally found in and/or produced by a given yeast, bacterium, organism, microorganism, or cell in nature.

The term “heterologous” as used herein in the context of a modified host cell refers to various molecules, e.g., polynucleotides, polypeptides, enzymes, etc., wherein at least

one of the following is true: (a) the molecule(s) is/are foreign (“exogenous”) to (i.e., not naturally found in) the host cell; (b) the molecule(s) is/are naturally found in (e.g., is “endogenous to”) a given host microorganism or host cell but is either produced in an unnatural location or in an unnatural amount in the cell; and/or (c) the molecule(s) differ(s) in nucleotide or amino acid sequence from the endogenous nucleotide or amino acid sequence(s) such that the molecule differing in nucleotide or amino acid sequence from the endogenous nucleotide or amino acid as found endogenously is produced in an unnatural (e.g., greater than naturally found) amount in the cell.

The term “homolog,” as used herein with respect to an original enzyme or gene of a first family or species, refers to distinct enzymes or genes of a second family or species which are determined by functional, structural, or genomic analyses to be an enzyme or gene of the second family or species which corresponds to the original enzyme or gene of the first family or species. Homologs most often have functional, structural, or genomic similarities. Techniques are known by which homologs of an enzyme or gene can readily be cloned using genetic probes and PCR. Identity of cloned sequences as homologs can be confirmed using functional assays and/or by genomic mapping of the genes.

A protein has “homology” or is “homologous” to a second protein if the amino acid sequence encoded by a gene has a similar amino acid sequence to that of the second gene. Alternatively, a protein has homology to a second protein if the two proteins have “similar” amino acid sequences. Thus, the term “homologous proteins” is intended to mean that the two proteins have similar amino acid sequences. In certain instances, the homology between two proteins is indicative of its shared ancestry, related by evolution.

The term “fatty acid” as used herein refers to a compound of structure R—COOH, wherein R is a C₆ to C₂₄ saturated, unsaturated, linear, branched or cyclic hydrocarbon and the carboxyl group is at position 1. In a particular embodiment, R is a C₆ to C₂₄ saturated or unsaturated linear hydrocarbon and the carboxyl group is at position 1.

The term “fatty alcohol” as used herein refers to an aliphatic alcohol having the formula R—OH, wherein R is a C₆ to C₂₄ saturated, unsaturated, linear, branched or cyclic hydrocarbon. In a particular embodiment, R is a C₆ to C₂₄ saturated or unsaturated linear hydrocarbon.

The term “fatty acyl-CoA” refers to a compound having the structure R—(CO)—S—R₁, wherein R₁ is Coenzyme A, and the term “fatty acyl-ACP” refers to a compound having the structure R—(CO)—S—R₁, wherein R₁ is an acyl carrier protein ACP.

Introduction

The present disclosure addresses the need for novel technologies for the cost-efficient production of valuable products from low-cost feedstocks. Specifically, the present inventors have addressed this need with the development of recombinant microorganisms capable of producing a wide-range of unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates including synthetic insect pheromones, fragrances, flavors, and polymer intermediates from low-cost feedstocks. Thus, aspects of the disclosure are based on the inventors’ discovery that recombinant microorganisms can be engineered in order to produce valuable products from low-cost feedstocks, which circumvents conventional synthetic methodologies to produce valuable products.

As discussed above, recombinant microorganisms can be engineered to synthesize mono- or poly-unsaturated C₆-C₂₄ fatty alcohols. Mono- or poly-unsaturated C₆-C₂₄ fatty alcohols synthesized as described herein can be further converted into the corresponding aldehydes or acetates. Thus, various embodiments of the present disclosure can be used to synthesize a variety of insect pheromones selected from fatty alcohols, aldehydes, and acetates. Additionally, embodiments described herein can also be used for the synthesis of fragrances, flavors, and polymer intermediates.

Pheromones

As described above, embodiments of the disclosure provide for the synthesis of one or more insect pheromones using a recombinant microorganism. A pheromone is a volatile chemical compound that is secreted by a particular insect for the function of chemical communication within the species. That is, a pheromone is secreted or excreted chemical factor that triggers a social response in members of the same species. There are, inter alia, alarm pheromones, food trail pheromones, sex pheromones, aggregation pheromones, epideictic pheromones, releaser pheromones, primer pheromones, and territorial pheromones, that affect behavior or physiology.

Non-limiting examples of insect pheromones which can be synthesized using the recombinant microorganisms and methods disclosed herein include linear alcohols, aldehydes, and acetates listed in Table 1.

TABLE 1

C ₆ -C ₂₀ Linear Pheromones	
Name	Name
(E)-2-Decen-1-ol	(E,E)-10,12-Tetradecadien-1-ol
(E)-2-Decenyl acetate	(E,E)-10,12-Tetradecadienyl acetate
(E)-2-Decenal	(E,E)-10,12-Tetradecadienal
(Z)-2-Decen-1-ol	(E,Z)-10,12-Tetradecadienyl acetate
(Z)-2-Decenyl acetate	(Z,E)-10,12-Tetradecadienyl acetate
(Z)-2-Decenal	(Z,Z)-10,12-Tetradecadien-1-ol
(E)-3-Decen-1-ol	(Z,Z)-10,12-Tetradecadienyl acetate
(Z)-3-Decenyl acetate	(E,Z,Z)-3,8,11-Tetradecatrienyl acetate
(Z)-3-Decen-1-ol	(E)-8-Pentadecen-1-ol
(Z)-4-Decen-1-ol	(E)-8-Pentadecenyl acetate
(E)-4-Decenyl acetate	(Z)-8-Pentadecen-1-ol
(Z)-4-Decenyl acetate	(Z)-8-Pentadecenyl acetate
(Z)-4-Decenal	(Z)-9-Pentadecenyl acetate
(E)-5-Decen-1-ol	(E)-9-Pentadecenyl acetate
(E)-5-Decenyl acetate	(Z)-10-Pentadecenyl acetate
(Z)-5-Decen-1-ol	(Z)-10-Pentadecenal
(Z)-5-Decenyl acetate	(E)-12-Pentadecenyl acetate
(Z)-5-Decenal	(Z)-12-Pentadecenyl acetate
(E)-7-Decenyl acetate	(Z,Z)-6,9-Pentadecadien-1-ol

TABLE 1-continued

C ₆ -C ₂₀ Linear Pheromones	
Name	Name
(Z)-7-Decenyl acetate	(Z,Z)-6,9-Pentadecadienyl acetate
(E)-8-Decen-1-ol	(Z,Z)-6,9-Pentadecadienal
(E,E)-2,4-Decadienal	(E,E)-8,10-Pentadecadienyl acetate
(E,Z)-2,4-Decadienal	(E,Z)-8,10-Pentadecadien-1-ol
(Z,Z)-2,4-Decadienal	(E,Z)-8,10-Pentadecadienyl acetate
(E,E)-3,5-Decadienyl acetate	(Z,E)-8,10-Pentadecadienyl acetate
(Z,E)-3,5-Decadienyl acetate	(Z,Z)-8,10-Pentadecadienyl acetate
(Z,Z)-4,7-Decadien-1-ol	(E,Z)-9,11-Pentadecadienal
(Z,Z)-4,7-Decadienyl acetate	(Z,Z)-9,11-Pentadecadienal
(E)-2-Undecenyl acetate	(Z)-3-Hexadecenyl acetate
(E)-2-Undecenal	(E)-5-Hexadecen-1-ol
(Z)-5-Undecenyl acetate	(E)-5-Hexadecenyl acetate
(Z)-7-Undecenyl acetate	(Z)-5-Hexadecen-1-ol
(Z)-8-Undecenyl acetate	(Z)-5-Hexadecenyl acetate
(Z)-9-Undecenyl acetate	(E)-6-Hexadecenyl acetate
(E)-2-Dodecenal	(E)-7-Hexadecen-1-ol
(Z)-3-Dodecen-1-ol	(E)-7-Hexadecenyl acetate
(E)-3-Dodecenyl acetate	(E)-7-Hexadecenal
(Z)-3-Dodecenyl acetate	(Z)-7-Hexadecen-1-ol
(E)-4-Dodecenyl acetate	(Z)-7-Hexadecenyl acetate
(E)-5-Dodecen-1-ol	(Z)-7-Hexadecenal
(E)-5-Dodecenyl acetate	(E)-8-Hexadecenyl acetate
(Z)-5-Dodecen-1-ol	(E)-9-Hexadecen-1-ol
(Z)-5-Dodecenyl acetate	(E)-9-Hexadecenyl acetate
(Z)-5-Dodecenal	(E)-9-Hexadecenal
(E)-6-Dodecen-1-ol	(Z)-9-Hexadecen-1-ol
(Z)-6-Dodecenyl acetate	(Z)-9-Hexadecenyl acetate
(E)-6-Dodecenal	(Z)-9-Hexadecenal
(E)-7-Dodecen-1-ol	(E)-10-Hexadecen-1-ol
(E)-7-Dodecenyl acetate	(E)-10-Hexadecenal
(E)-7-Dodecenal	(Z)-10-Hexadecenyl acetate
(Z)-7-Dodecen-1-ol	(Z)-10-Hexadecenal
(Z)-7-Dodecenyl acetate	(E)-11-Hexadecen-1-ol
(Z)-7-Dodecenal	(E)-11-Hexadecenyl acetate
(E)-8-Dodecen-1-ol	(E)-11-Hexadecenal
(E)-8-Dodecenyl acetate	(Z)-11-Hexadecen-1-ol
(E)-8-Dodecenal	(Z)-11-Hexadecenyl acetate
(Z)-8-Dodecen-1-ol	(Z)-11-Hexadecenal
(Z)-8-Dodecenyl acetate	(Z)-12-Hexadecenyl acetate
(E)-9-Dodecen-1-ol	(Z)-12-Hexadecenal
(E)-9-Dodecenyl acetate	(E)-14-Hexadecenal
(E)-9-Dodecenal	(Z)-14-Hexadecenyl acetate
(Z)-9-Dodecen-1-ol	(E,E)-1,3-Hexadecadien-1-ol
(Z)-9-Dodecenyl acetate	(E,Z)-4,6-Hexadecadien-1-ol
(Z)-9-Dodecenal	(E,Z)-4,6-Hexadecadienyl acetate
(E)-10-Dodecen-1-ol	(E,Z)-4,6-Hexadecadienal
(E)-10-Dodecenyl acetate	(E,Z)-6,11-Hexadecadienyl acetate
(E)-10-Dodecenal	(E,Z)-6,11-Hexadecadienal
(Z)-10-Dodecen-1-ol	(Z,Z)-7,10-Hexadecadien-1-ol
(Z)-10-Dodecenyl acetate	(Z,Z)-7,10-Hexadecadienyl acetate
(E,Z)-3,5-Dodecadienyl acetate	(Z,E)-7,11-Hexadecadien-1-ol
(Z,E)-3,5-Dodecadienyl acetate	(Z,E)-7,11-Hexadecadienyl acetate
(Z,Z)-3,6-Dodecadien-1-ol	(Z,E)-7,11-Hexadecadienal
(E,E)-4,10-Dodecadienyl acetate	(Z,Z)-7,11-Hexadecadien-1-ol
(E,E)-5,7-Dodecadien-1-ol	(Z,Z)-7,11-Hexadecadienyl acetate
(E,E)-5,7-Dodecadienyl acetate	(Z,Z)-7,11-Hexadecadienal
(E,Z)-5,7-Dodecadien-1-ol	(Z,Z)-8,10-Hexadecadienyl acetate
(E,Z)-5,7-Dodecadienyl acetate	(E,Z)-8,11-Hexadecadienal
(E,Z)-5,7-Dodecadienal	(E,E)-9,11-Hexadecadienal
(Z,E)-5,7-Dodecadien-1-ol	(E,Z)-9,11-Hexadecadienyl acetate
(Z,E)-5,7-Dodecadienyl acetate	(E,Z)-9,11-Hexadecadienal
(Z,E)-5,7-Dodecadienal	(Z,E)-9,11-Hexadecadienal
(Z,Z)-5,7-Dodecadienyl acetate	(Z,Z)-9,11-Hexadecadienal
(Z,Z)-5,7-Dodecadienal	(E,E)-10,12-Hexadecadien-1-ol
(E,E)-7,9-Dodecadienyl acetate	(E,E)-10,12-Hexadecadienyl acetate
(E,Z)-7,9-Dodecadien-1-ol	(E,E)-10,12-Hexadecadienal
(E,Z)-7,9-Dodecadienyl acetate	(E,Z)-10,12-Hexadecadien-1-ol
(E,Z)-7,9-Dodecadienal	(E,Z)-10,12-Hexadecadienyl acetate
(Z,E)-7,9-Dodecadien-1-ol	(E,Z)-10,12-Hexadecadienal
(Z,E)-7,9-Dodecadienyl acetate	(Z,E)-10,12-Hexadecadienyl acetate
(Z,Z)-7,9-Dodecadien-1-ol	(Z,E)-10,12-Hexadecadienal
(Z,Z)-7,9-Dodecadienyl acetate	(Z,Z)-10,12-Hexadecadienal
(E,E)-8,10-Dodecadien-1-ol	(E,E)-11,13-Hexadecadien-1-ol
(E,E)-8,10-Dodecadienyl acetate	(E,E)-11,13-Hexadecadienyl acetate
(E,E)-8,10-Dodecadienal	(E,E)-11,13-Hexadecadienal
(E,Z)-8,10-Dodecadien-1-ol	(E,Z)-11,13-Hexadecadien-1-ol

TABLE 1-continued

C ₆ -C ₂₀ Linear Pheromones	
Name	Name
(E,Z)-8,10-Dodecadienyl acetate	(E,Z)-11,13-Hexadecadienyl acetate
(E,Z)-8,10-Dodecadienal	(E,Z)-11,13-Hexadecadienal
(Z,E)-8,10-Dodecadien-1-ol	(Z,E)-11,13-Hexadecadien-1-ol
(Z,E)-8,10-Dodecadienyl acetate	(Z,E)-11,13-Hexadecadienyl acetate
(Z,E)-8,10-Dodecadienal	(Z,E)-11,13-Hexadecadienal
(Z,Z)-8,10-Dodecadien-1-ol	(Z,Z)-11,13-Hexadecadien-1-ol
(Z,Z)-8,10-Dodecadienyl acetate	(Z,Z)-11,13-Hexadecadienyl acetate
(Z,E,E)-3,6,8-Dodecatrien-1-ol	(Z,Z)-11,13-Hexadecadienal
(Z,Z,E)-3,6,8-Dodecatrien-1-ol	(E,E)-10,14-Hexadecadienal
(E)-2-Tridecenyl acetate	(Z,E)-11,14-Hexadecadienyl acetate
(Z)-2-Tridecenyl acetate	(E,E,Z)-4,6,10-Hexadecatrien-1-ol
(E)-3-Tridecenyl acetate	(E,E,Z)-4,6,10-Hexadecatrienyl acetate
(E)-4-Tridecenyl acetate	(E,Z,Z)-4,6,10-Hexadecatrien-1-ol
(Z)-4-Tridecenyl acetate	(E,Z,Z)-4,6,10-Hexadecatrienyl acetate
(Z)-4-Tridecenal	(E,E,Z)-4,6,11-Hexadecatrienyl acetate
(E)-6-Tridecenyl acetate	(E,E,Z)-4,6,11-Hexadecatrienal
(Z)-7-Tridecenyl acetate	(Z,Z,E)-7,11,13-Hexadecatrienal
(E)-8-Tridecenyl acetate	(E,E,E)-10,12,14-Hexadecatrienyl acetate
(Z)-8-Tridecenyl acetate	(E,E,E)-10,12,14-Hexadecatrienal
(E)-9-Tridecenyl acetate	(E,E,Z)-10,12,14-Hexadecatrienyl acetate
(Z)-9-Tridecenyl acetate	(E,E,Z)-10,12,14-Hexadecatrienal
(Z)-10-Tridecenyl acetate	(E,E,Z,Z)-4,6,11,13-Hexadecatetraenal
(E)-11-Tridecenyl acetate	(E)-2-Heptadecenal
(Z)-11-Tridecenyl acetate	(Z)-2-Heptadecenal
(E,Z)-4,7-Tridecadienyl acetate	(E)-8-Heptadecen-1-ol
(Z,Z)-4,7-Tridecadien-1-ol	(E)-8-Heptadecenyl acetate
(Z,Z)-4,7-Tridecadienyl acetate	(Z)-8-Heptadecen-1-ol
(E,Z)-5,9-Tridecadienyl acetate	(Z)-9-Heptadecenal
(Z,E)-5,9-Tridecadienyl acetate	(E)-10-Heptadecenyl acetate
(Z,Z)-5,9-Tridecadienyl acetate	(Z)-11-Heptadecen-1-ol
(Z,Z)-7,11-Tridecadienyl acetate	(Z)-11-Heptadecenyl acetate
(E,Z,Z)-4,7,10-Tridecatrienyl acetate	(E,E)-4,8-Heptadecadienyl acetate
(E)-3-Tetradecen-1-ol	(Z,Z)-8,10-Heptadecadien-1-ol
(E)-3-Tetradecenyl acetate	(Z,Z)-8,11-Heptadecadienyl acetate
(Z)-3-Tetradecen-1-ol	(E)-2-Octadecenyl acetate
(Z)-3-Tetradecenyl acetate	(E)-2-Octadecenal
(E)-5-Tetradecen-1-ol	(Z)-2-Octadecenyl acetate
(E)-5-Tetradecenyl acetate	(Z)-2-Octadecenal
(E)-5-Tetradecenal	(E)-9-Octadecen-1-ol
(Z)-5-Tetradecen-1-ol	(E)-9-Octadecenyl acetate
(Z)-5-Tetradecenyl acetate	(E)-9-Octadecenal
(Z)-5-Tetradecenal	(Z)-9-Octadecen-1-ol
(E)-6-Tetradecenyl acetate	(Z)-9-Octadecenyl acetate
(Z)-6-Tetradecenyl acetate	(Z)-9-Octadecenal
(E)-7-Tetradecen-1-ol	(E)-11-Octadecen-1-ol
(E)-7-Tetradecenyl acetate	(E)-11-Octadecenal
(Z)-7-Tetradecen-1-ol	(Z)-11-Octadecen-1-ol
(Z)-7-Tetradecenyl acetate	(Z)-11-Octadecenyl acetate
(Z)-7-Tetradecenal	(Z)-11-Octadecenal
(E)-8-Tetradecenyl acetate	(E)-13-Octadecenyl acetate
(Z)-8-Tetradecen-1-ol	(E)-13-Octadecenal
(Z)-8-Tetradecenyl acetate	(Z)-13-Octadecen-1-ol
(Z)-8-Tetradecenal	(Z)-13-Octadecenyl acetate
(E)-9-Tetradecen-1-ol	(Z)-13-Octadecenal
(E)-9-Tetradecenyl acetate	(E)-14-Octadecenal
(Z)-9-Tetradecen-1-ol	(E,Z)-2,13-Octadecadien-1-ol
(Z)-9-Tetradecenyl acetate	(E,Z)-2,13-Octadecadienyl acetate
(Z)-9-Tetradecenal	(E,Z)-2,13-Octadecadienal
(E)-10-Tetradecenyl acetate	(Z,E)-2,13-Octadecadienyl acetate
(Z)-10-Tetradecenyl acetate	(Z,Z)-2,13-Octadecadien-1-ol
(E)-11-Tetradecen-1-ol	(Z,Z)-2,13-Octadecadienyl acetate
(E)-11-Tetradecenyl acetate	(E,E)-3,13-Octadecadienyl acetate
(E)-11-Tetradecenal	(E,Z)-3,13-Octadecadienyl acetate
(Z)-11-Tetradecen-1-ol	(E,Z)-3,13-Octadecadienal
(Z)-11-Tetradecenyl acetate	(Z,E)-3,13-Octadecadienyl acetate
(Z)-11-Tetradecenal	(Z,Z)-3,13-Octadecadienyl acetate
(E)-12-Tetradecenyl acetate	(Z,Z)-3,13-Octadecadienal
(Z)-12-Tetradecenyl acetate	(E,E)-5,9-Octadecadien-1-ol
(E,E)-2,4-Tetradecadienal	(E,E)-5,9-Octadecadienyl acetate
(E,E)-3,5-Tetradecadienyl acetate	(E,E)-9,12-Octadecadien-1-ol
(E,Z)-3,5-Tetradecadienyl acetate	(Z,Z)-9,12-Octadecadienyl acetate
(Z,E)-3,5-Tetradecadienyl acetate	(Z,Z)-9,12-Octadecadienal
(E,Z)-3,7-Tetradecadienyl acetate	(Z,Z)-11,13-Octadecadienal
(E,Z)-3,8-Tetradecadienyl acetate	(E,E)-11,14-Octadecadienal
(E,Z)-4,9-Tetradecadienyl acetate	(Z,Z)-13,15-Octadecadienal
(E,Z)-4,9-Tetradecadienal	(Z,Z,Z)-3,6,9-Octadecatrienyl acetate

TABLE 1-continued

C ₆ -C ₂₀ Linear Pheromones	
Name	Name
(E,Z)-4,10-Tetradecadienyl acetate	(E,E,E)-9,12,15-Octadecatrien-1-ol
(E,E)-5,8-Tetradecadienal	(Z,Z,Z)-9,12,15-Octadecatrienyl acetate
(Z,Z)-5,8-Tetradecadien-1-ol	(Z,Z,Z)-9,12,15-Octadecatrienal
(Z,Z)-5,8-Tetradecadienyl acetate	
(Z,Z)-5,8-Tetradecadienal	
(E,E)-8,10-Tetradecadien-1-ol	
(E,E)-8,10-Tetradecadienyl acetate	
(E,E)-8,10-Tetradecadienal	
(E,Z)-8,10-Tetradecadienyl acetate	
(E,Z)-8,10-Tetradecadienal	
(Z,E)-8,10-Tetradecadien-1-ol	
(Z,E)-8,10-Tetradecadienyl acetate	
(Z,Z)-8,10-Tetradecadienal	
(E,E)-9,11-Tetradecadienyl acetate	
(E,Z)-9,11-Tetradecadienyl acetate	
(Z,E)-9,11-Tetradecadien-1-ol	
(Z,E)-9,11-Tetradecadienyl acetate	
(Z,E)-9,11-Tetradecadienal	
(Z,Z)-9,11-Tetradecadien-1-ol	
(Z,Z)-9,11-Tetradecadienyl acetate	
(Z,Z)-9,11-Tetradecadienal	
(E,E)-9,12-Tetradecadienyl acetate	
(Z,E)-9,12-Tetradecadien-1-ol	
(Z,E)-9,12-Tetradecadienyl acetate	
(Z,E)-9,12-Tetradecadienal	
(Z,Z)-9,12-Tetradecadien-1-ol	
(Z,Z)-9,12-Tetradecadienyl acetate	

In some aspects, the pheromones synthesized as taught in this disclosure include at least one pheromone listed in Table 2a to modulate the behavior of an insect listed in Table 2a. In other aspects, non-limiting examples of insect pheromones which can be synthesized using the recombinant microorganisms and methods disclosed herein include alcohols, aldehydes, and acetates listed in Table 2a. However, the

microorganisms described herein are not limited to the synthesis of C₆-C₂₀ pheromones listed in Table 1 and Table 2a. Rather, the disclosed microorganisms can also be utilized in the synthesis of various C₆-C₂₄ mono- or poly-unsaturated fatty alcohols, aldehydes, and acetates, including fragrances, flavors, and polymer intermediates.

TABLE 2a










Exemplary pheromones that can be synthesized according to methods described in the present disclosure.		
Name	Structure	Example of Biological importance
(Z)-3-hexen-1-ol		See, Sugimoto et al. (2014)
(Z)-3-nonen-1-ol		West Indian Fruity Fly male sex pheromone
(Z)-5-decan-1-ol		
(Z)-5-decenyl acetate		<i>Agrotis segetum</i> sex pheromone component
(E)-5-decen-1-ol		<i>Anarsia lineatella</i> sex pheromone component
(E)-5-decenyl acetate		<i>Anarsia lineatella</i> sex pheromone component
(Z)-7-dodecen-1-ol		
(Z)-7-dodecenyl acetate		<i>Pseudoplusia includens</i> sex pheromone <i>Agrotis segetum</i> sex pheromone component
(E)-8-dodecen-1-ol		Citrus Fruit Moth sex pheromone

TABLE 2a-continued

Exemplary pheromones that can be synthesized according to methods described in the present disclosure.		
Name	Structure	Example of Biological importance
(E)-8-dodecenyl acetate		<i>Grapholitha molesta</i> , <i>Ecdytolopha aurantiana</i> sex pheromone component
(Z)-8-dodecen-1-ol		<i>Grapholitha molesta</i> , <i>Ecdytolopha aurantiana</i> sex pheromone component
(Z)-8-dodecenyl acetate		<i>Grapholitha molesta</i> sex pheromone component
(Z)-9-dodecen-1-ol		
(Z)-9-dodecenyl acetate		<i>Eupoecilia ambiguella</i> sex pheromone
(E,E)-8,10-dodecadien-1-ol		<i>Cydia pomonella</i>
(7E,9Z)-dodecadienyl acetate		<i>Lobesia botrana</i>
(Z)-9-tetradecen-1-ol		
(Z)-9-tetradecenyl acetate		<i>Pandemis pyrusana</i> , <i>Naranga aenescens</i> , <i>Agrotis segetum</i> sex pheromone component
(Z)-11-tetradecen-1-ol		
(Z)-11-tetradecenyl acetate		<i>Pandemis pyrusana</i> , <i>Choristoneura roseceana</i> sex pheromone component
(E)-11-tetradecen-1-ol		
(E)-11-tetradecenyl acetate		<i>Choristoneura roseceana</i> , <i>Crocidolomia pavonana</i> sex pheromone component
(Z)-7-hexadecen-1-ol		
(Z)-7-hexadecenal		<i>Diatraea considerata</i> sex pheromone component
(Z)-9-hexadecen-1-ol		
(Z)-9-hexadecenal		<i>Helicoverpa zea</i> , <i>Helicoverpa armigera</i> , <i>Heliothis virescens</i> sex pheromone component
(Z)-9-hexadecenyl acetate		<i>Naranga aenescens</i> sex pheromone component
(Z)-11-hexadecen-1-ol		
(Z)-11-hexadecenal		<i>Platyptila carduidactyla</i> , <i>Heliothis virescens</i> sex pheromone <i>Helicoverpa zea</i> , <i>Helicoverpa armigera</i> , <i>Plutella xylostella</i> , <i>Diatraea considerate</i> , <i>Diatraea grandiosella</i> , <i>Diatraea saccharalis</i> , <i>Acrolepiopsis assectella</i> sex pheromone component

TABLE 2a-continued

Exemplary pheromones that can be synthesized according to methods described in the present disclosure.		
Name	Structure	Example of Biological importance
(Z)-11-hexadecenyl acetate		<i>Discestra trifolii</i> sex pheromone <i>Heliothis virescens</i> , <i>Plutella xylostella</i> , <i>Acrolepiopsis assectella</i> , <i>Crocidolomia pavonana</i> , <i>Naranga aenescens</i> sex pheromone component
(Z,Z)-11,13-hexadecadienal		<i>Amyelosis transitella</i>
(Z,Z)-11,13-hexadecadien-1-ol		<i>Amyelosis transitella</i>
(11Z,13E)-hexadecadien-1-ol		<i>Amyelosis transitella</i>
(9Z,11E)-hexadecadienal		
(Z)-13-octadecen-1-ol		
(Z)-13-octadecenal		<i>Diatraea considerata</i> , <i>Diatraea grandiosella</i> sex pheromone component
(Z,Z,Z,Z,Z)-3,6,9,12,15-tricosapentaene		<i>Amyelosis transitella</i>

Most pheromones comprise a hydrocarbon skeleton with the terminal hydrogen substituted by a functional group (Ryan M F (2002). Insect Chemoreception. Fundamental and Applied. Kluwer Academic Publishers). Table 2b shows some common functional groups, along with their formulas, prefixes and suffixes. The presence of one or more double bonds, generated by the loss of hydrogens from adjacent carbons, determines the degree of unsaturation of the molecule and alters the designation of a hydrocarbon from -ane (no multiple bonds) to -ene. The presence of two and three double bonds is indicated by ending the name with -diene and -triene, respectively. The position of each double bond is represented by a numeral corresponding to that of the carbon from which it begins, with each carbon numbered from that attached to the functional group. The carbon to which the functional group is attached is designated -1-. Pheromones may have, but are not limited to, hydrocarbon chain lengths numbering 10 (deca-), 12 (dodeca-), 14 (tetradeca-), 16 (hexadeca-), or 18 (octadeca-) carbons long. The presence of a double bond has another effect. It precludes rotation of the molecule by fixing it in one of two possible configurations, each representing geometric isomers that are different molecules. These are designated either E (from the German word Entgegen, opposite) or Z (Zusammen, together), when the carbon chains are connected on the opposite (trans) or same (cis) side, respectively, of the double bond.

TABLE 2b

Prefixes and suffixes for common functional groups			
Functional group	Formula	Prefix	Suffix
Alcohol	—OH	Hydroxy-	-ol
Aldehyde	—CH=O	Formyl-	-al
Amine	—NH ₂	Amino-	-amine
Carboxylic acid	—COOH	Carboxy-	-oic acid
Ester	—COOR	R-oxycarbonyl-	-R-oate
Ketone	>C=O	Oxo-	-one

From Howse, PE, Stevens, IDR and Jones, OT (1998). Insect pheromes and their use in pest management. London: Chapman and Hall.

Pheromones described herein can be referred to using IUPAC nomenclature or various abbreviations or variations known to one skilled in the art. For example, (11Z)-hexadecen-1-al, can also be written as Z-11-hexadecen-1-al, Z-11-hexadecenal, or Z-x-y:Ald, wherein x represents the position of the double bond and y represents the number of carbons in the hydrocarbon skeleton. Abbreviations used herein and known to those skilled in the art to identify functional groups on the hydrocarbon skeleton include “Ald,” indicating an aldehyde, “OH,” indicating an alcohol, and “Ac,” indicating an acetyl. Also, the number of carbons in the chain can be indicated using numerals rather than using the written name. Thus, as used herein, an unsaturated carbon chain comprised of sixteen carbons can be written as hexadecene or 16.

Similar abbreviation and derivations are used herein to describe pheromone precursors. For example, the fatty acyl-

CoA precursors of (11Z)-hexadecen-1-al can be identified as (11Z)-hexadecenyl-CoA or Z-11-16:Acyl-CoA.

The present disclosure relates to the synthesis of mono- or poly-unsaturated C₆-C₂₄ fatty alcohols, aldehydes, and acetates using a recombinant microorganism comprised of one or more heterologous enzymes, which catalyze substrate to product conversions for one or more steps in the synthesis process.

Desaturase

The present disclosure describes enzymes that desaturate fatty acyl substrates to corresponding unsaturated fatty acyl substrates.

In some embodiments, a desaturase is used to catalyze the conversion of a fatty acyl-CoA or acyl-ACP to a corresponding unsaturated fatty acyl-CoA or acyl-ACP. A desaturase is an enzyme that catalyzes the formation of a carbon-carbon double bond in a saturated fatty acid or fatty acid derivative, e.g., fatty acyl-CoA or fatty acyl-ACP (collectively referred to herein as "fatty acyl"), by removing at least two hydrogen atoms to produce a corresponding unsaturated fatty acid/acyl. Desaturases are classified with respect to the ability of the enzyme to selectively catalyze double bond formation at a subterminal carbon relative to the methyl end of the fatty acid/acyl or a subterminal carbon relative to the carbonyl end of the fatty acid/acyl. Omega (ω) desaturases catalyze the formation of a carbon-carbon double bond at a fixed subterminal carbon relative to the methyl end of a fatty acid/acyl. For example, an ω^3 desaturase catalyzes the formation of a double bond between the third and fourth carbon relative the methyl end of a fatty acid/acyl. Delta (Δ) desaturases catalyze the formation of a carbon-carbon double bond at a specific position relative to the carboxyl group of a fatty acid or the carbonyl group of a fatty acyl CoA. For example, a Δ^9 desaturase catalyzes the formation of a double bond between the C₉ and C₁₀ carbons with respect to the carboxyl end of the fatty acid or the carbonyl group of a fatty acyl CoA.

As used herein, a desaturase can be described with reference to the location in which the desaturase catalyzes the formation of a double bond and the resultant geometric configuration (i.e., E/Z) of the unsaturated hydrocarbon. Accordingly, as used herein, a Z9 desaturase refers to a Δ desaturase that catalyzes the formation of a double bond between the C₉ and C₁₀ carbons with respect to the carbonyl end of a fatty acid/acyl, thereby orienting two hydrocarbons on opposing sides of the carbon-carbon double bonds in the cis or Z configuration. Similarly, as used herein, a Z11 desaturase refers to a Δ desaturase that catalyzes the formation of a double bond between the C₁₁ and C₁₂ carbons with respect to the carbonyl end of a fatty acid/acyl.

Desaturases have a conserved structural motif. This sequence motif of transmembrane desaturases is characterized by [HX3-4HX7-41(3 non-His)HX2-3(1 nonHis)HHX61-189(40 non-His)HX2-3(1 non-His)HH]. The sequence motif of soluble desaturases is characterized by two occurrences of [D/EEXXH].

In some embodiments, the desaturase is a fatty acyl-CoA desaturase that catalyzes the formation of a double bond in a fatty acyl-CoA. In some such embodiments, the fatty acyl-CoA desaturase described herein is capable of utilizing a fatty acyl-CoA as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms. Thus, the desaturase used in the recombinant microorganism can be selected based on the chain length of the substrate.

In some embodiments, the fatty acyl desaturase described herein is capable of catalyzing the formation of a double

bond at a desired carbon relative to the terminal CoA on the unsaturated fatty acyl-CoA. Thus, in some embodiments, a desaturase can be selected for use in the recombinant microorganism which catalyzes double bond insertion at the 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 position with respect to the carbonyl group on a fatty acyl-CoA.

In some embodiments, the fatty acyl desaturase described herein is capable of catalyzing the formation of a double bond in a saturated fatty acyl-CoA such that the resultant unsaturated fatty acyl-CoA has a cis or trans (i.e., Z or E) geometric configuration.

In some embodiments, the desaturase is a fatty acyl-ACP desaturase that catalyzes the formation of a double bond in a fatty acyl-ACP. In some embodiments, the fatty acyl-ACP desaturase described herein is capable of utilizing a fatty acyl-CoA as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms. Thus, the desaturase used in the recombinant microorganism can be selected based on the chain length of the substrate.

In some embodiments, the fatty acyl-ACP desaturase described herein is capable of catalyzing the formation of a double bond at a desired carbon relative to the terminal carbonyl on the unsaturated fatty acyl-ACP. Thus, in some embodiments, a desaturase can be selected for use in the recombinant microorganism which catalyzes double bond insertion at the 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, or 13 position with respect to the carbonyl group on a fatty acyl-ACP.

In some embodiments, the fatty acyl desaturase described herein is capable of catalyzing the formation of a double bond in a saturated fatty acyl-CoA such that the resultant unsaturated fatty acyl-ACP has a cis or trans (i.e., Z or E) geometric configuration.

In one embodiment, the fatty acyl desaturase is a Z11 desaturase. In some embodiments, a nucleic acid sequence encoding a Z11 desaturase from organisms of the species *Agrotis segetum*, *Amyelois transitella*, *Argyrotaenia velutina*, *Choristoneura rosaceana*, *Lampronia capitella*, *Trichoplusia ni*, *Helicoverpa zea*, or *Thalassiosira pseudonana* is codon optimized. In some embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 9, 18, 24 and 26 from *Trichoplusia ni*. In other embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 10 and 16 from *Agrotis segetum*. In some embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 11 and 23 from *Thalassiosira pseudonana*. In certain embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 12, 17 and 30 from *Amyelois transitella*. In further embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 13, 19, 25, 27 and 31 from *Helicoverpa zea*. In some embodiments, the Z11 desaturase comprises a chimeric polypeptide. In some embodiments, a complete or partial Z11 desaturase is fused to another polypeptide. In certain embodiments, the N-terminal native leader sequence of a Z11 desaturase is replaced by an oleosin leader sequence from another species. In certain embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 15, 28 and 29.

In one embodiment, the fatty acyl desaturase is a Z9 desaturase. In some embodiments, a nucleic acid sequence encoding a Z9 desaturase is codon optimized. In some embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 20 from *Ostrinia furnacalis*. In other embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 21 from *Lampronia capitella*. In some

embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 22 from *Helicoverpa zea*.

Fatty Acyl Reductase

The present disclosure describes enzymes that reduce fatty acyl substrates to corresponding fatty alcohols or aldehydes.

In some embodiments, a fatty alcohol forming fatty acyl-reductase is used to catalyze the conversion of a fatty acyl-CoA to a corresponding fatty alcohol. In some embodiments, a fatty aldehyde forming fatty acyl-reductase is used to catalyze the conversion of a fatty acyl-ACP to a corresponding fatty aldehyde. A fatty acyl reductase is an enzyme that catalyzes the reduction of a fatty acyl-CoA to a corresponding fatty alcohol or the reduction of a fatty acyl-ACP to a corresponding fatty aldehyde. A fatty acyl-CoA and fatty acyl-ACP has a structure of $R-(CO)-S-R_1$, wherein R is a C_6 to C_{24} saturated, unsaturated, linear, branched or cyclic hydrocarbon, and R_1 represents CoA or ACP. In a particular embodiment, R is a C_6 to C_{24} saturated or unsaturated linear hydrocarbon. "CoA" is a non-protein acyl carrier group involved in the synthesis and oxidation of fatty acids. "ACP" is an acyl carrier protein, i.e., a polypeptide or protein subunit, of fatty acid synthase used in the synthesis of fatty acids.

Thus, in some embodiments, the disclosure provides for a fatty alcohol forming fatty acyl-reductase which catalyzes the reduction of a fatty acyl-CoA to the corresponding fatty alcohol. For example, $R-(CO)-S-CoA$ is converted to $R-CH_2OH$ and $CoA-SH$ when two molecules of $NAD(P)H$ are oxidized to $NAD(P)^+$. Accordingly, in some such embodiments, a recombinant microorganism described herein can include a heterologous fatty alcohol forming fatty acyl-reductase, which catalyzes the reduction a fatty acyl-CoA to the corresponding fatty alcohol. In an exemplary embodiment, a recombinant microorganism disclosed herein includes at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase which catalyzes the conversion of a mono- or poly-unsaturated C_6-C_{24} fatty acyl-CoA into the corresponding mono- or poly-unsaturated C_6-C_{24} fatty alcohol.

In other embodiments, the disclosure provides for a fatty aldehyde forming fatty acyl-reductase which catalyzes the reduction of a fatty acyl-ACP to the corresponding fatty aldehyde. For example, $R-(CO)-S-ACP$ is converted to $R-(CO)-H$ and $ACP-SH$ when one molecule of $NAD(P)H$ is oxidized to $NAD(P)^+$. In some such embodiments, a recombinant microorganism described herein can include a heterologous fatty aldehyde forming fatty acyl-reductase, which catalyzes the reduction a fatty acyl-ACP to the corresponding fatty aldehyde. In an exemplary embodiment, a recombinant microorganism disclosed herein includes at least one exogenous nucleic acid molecule encoding a fatty aldehyde forming fatty-acyl reductase which catalyzes the conversion of a mono- or poly-unsaturated C_6-C_{24} fatty acyl-ACP into the corresponding mono- or poly-unsaturated C_6-C_{24} fatty aldehyde.

In some embodiments, a nucleic acid sequence encoding a fatty-acyl reductase from organisms of the species *Agrotis segetum*, *Spodoptera littoralis*, or *Helicoverpa armigera* is codon optimized. In some embodiments, the fatty acyl reductase comprises a sequence set forth in SEQ ID NO: 1 from *Agrotis segetum*. In other embodiments, the fatty acyl reductase comprises a sequence set forth in SEQ ID NO: 2 from *Spodoptera littoralis*. In some embodiments, the fatty acyl reductase comprises a sequence selected from SEQ ID NOs: 3 and 32 from *Helicoverpa armigera*.

Acyl-ACP Synthetase

The present disclosure describes enzymes that ligate a fatty acid to the corresponding fatty acyl-ACP.

In some embodiments, an acyl-ACP synthetase is used to catalyze the conversion of a fatty acid to a corresponding fatty acyl-ACP. An acyl-ACP synthetase is an enzyme capable of ligating a fatty acid to ACP to produce a fatty acid acyl-ACP. In some embodiments, an acyl-ACP synthetase can be used to catalyze the conversion of a fatty acid to a corresponding fatty acyl-ACP. In some embodiments, the acyl-ACP synthetase is a synthetase capable of utilizing a fatty acid as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms. In one such embodiment, a recombinant microorganism described herein can include a heterologous acyl-ACP synthetase, which catalyzes the conversion of a fatty acid to a corresponding fatty acyl-ACP. In an exemplary embodiment, a recombinant microorganism disclosed herein includes at least one exogenous nucleic acid molecule which encodes an acyl-ACP synthetase that catalyzes the conversion of a saturated C_6-C_{24} fatty acid to a corresponding saturated C_6-C_{24} fatty acyl-ACP.

Fatty Acid Synthase Complex

The present disclosure describes enzymes that catalyze the elongation of a carbon chain in fatty acid.

In some embodiments, a fatty acid synthase complex is used to catalyze initiation and elongation of a carbon chain in a fatty acid. A "fatty acid synthase complex" refers to a group of enzymes that catalyzes the initiation and elongation of a carbon chain on a fatty acid. The ACP along with the enzymes in the fatty acid synthase (FAS) pathway control the length, degree of saturation, and branching of the fatty acids produced. The steps in this pathway are catalyzed by enzymes of the fatty acid biosynthesis (fab) and acetyl-CoA carboxylase (acc) gene families. Depending upon the desired product, one or more of these genes can be attenuated, expressed or over-expressed. In exemplary embodiments, one or more of these genes is over-expressed.

There are two principal classes of fatty acid synthases. Type I (FAS I) systems utilize a single large, multifunctional polypeptide and are common to both mammals and fungi (although the structural arrangement of fungal and mammalian synthases differ). The Type I FAS system is also found in the CMN group of bacteria (corynebacteria, mycobacteria, and *nocardia*). The Type II FAS (FAS II) is characterized by the use of discrete, monofunctional enzymes for fatty acid synthesis, and is found in archaea and bacteria.

The mechanism of FAS I and FAS II elongation and reduction is the substantially similar, as the domains of the FAS I multienzyme polypeptides and FAS II enzymes are largely conserved.

Fatty acids are synthesized by a series of decarboxylative Claisen condensation reactions from acetyl-CoA and malonyl-CoA. The steps in this pathway are catalyzed by enzymes of the fatty acid biosynthesis (fab) and acetyl-CoA carboxylase (acc) gene families. For a description of this pathway, see, e.g., Heath et al., *Prog. Lipid Res.* 40:467, 2001, which is herein incorporated by reference in its entirety. Without being limited by theory, in bacteria, acetyl-CoA is carboxylated by acetyl-CoA carboxylase (Acc, a multi-subunit enzyme encoded by four separate genes, accABCD), to form malonyl-CoA. In yeast, acetyl-CoA is carboxylated by the yeast equivalents of the acetyl-CoA carboxylase, encoded by ACC1 and ACC2. In bacteria, the malonate group is transferred to ACP by malonyl-CoA:ACP transacylase (FabD) to form malonyl-ACP. In yeast, a malonyl-palmitoyl transferase domain adds malonyl from malonyl-CoA to the ACP domain of the FAS complex. A condensa-

tion reaction then occurs, where malonyl-ACP merges with acyl-CoA, resulting in β -ketoacyl-ACP. In this manner, the hydrocarbon substrate is elongated by 2 carbons.

Following elongation, the β -keto group is reduced to the fully saturated carbon chain by the sequential action of a keto-reductase (KR), dehydratase (DH), and enol reductase (ER). The elongated fatty acid chain is carried between these active sites while attached covalently to the phosphopantetheine prosthetic group of ACP. First, the β -ketoacyl-ACP is reduced by NADPH to form β -hydroxyacyl-ACP. In bacteria, this step is catalyzed by β -ketoacyl-ACP reductase (FabG). The equivalent yeast reaction is catalyzed by the ketoreductase (KR) domain of FAS. β -hydroxyacyl-ACP is then dehydrated to form trans-2-enoyl-ACP, which is catalyzed by either β -hydroxyacyl-ACP dehydratase/isomerase (FabA) or β -hydroxyacyl-ACP dehydratase (FabZ) in bacteria or the dehydratase (DH) domain of FAS in yeast. NADPH-dependent trans-2-enoyl-ACP reductase I, II, or III (FabI, FabK, and FabL, respectively) in bacteria and the enol reductase (ER) domain of FAS in yeast reduces trans-2-enoyl-ACP to form acyl-ACP. Subsequent cycles are started by the condensation of malonyl-ACP with acyl-ACP by β -ketoacyl-ACP synthase I or β -ketoacyl-ACP synthase II (FabB and FabF, respectively, in bacteria or the beta-ketoacyl synthase (KS) domain in yeast).

In some embodiments, a fatty acid synthase complex can be used to catalyze elongation of a fatty acyl-ACP to a corresponding fatty acyl-ACP with a two carbon elongation relative to the substrate.

Dehydrogenase

The present disclosure describes enzymes that catalyze the conversion of a fatty aldehyde to a fatty alcohol. In some embodiments, an alcohol dehydrogenase (ADH, Table 3) is

used to catalyze the conversion of a fatty aldehyde to a fatty alcohol. A number of ADHs identified from alkanotrophic organisms, *Pseudomonas fluorescens* NRRL B-1244 (Hou et al. 1983), *Pseudomonas butanovora* ATCC 43655 (Vangnai and Arp 2001), and *Acinetobacter* sp. strain M-1 (Tani et al. 2000), have shown to be active on short to medium-chain alkyl alcohols (C_2 to C_{14}). Additionally, commercially available ADHs from Sigma, Horse liver ADH and Baker's yeast ADH have detectable activity for substrates with length C_{10} and greater. The reported activities for the longer fatty alcohols may be impacted by the difficulties in solubilizing the substrates. For the yeast ADH from Sigma, little to no activity is observed for C_{12} to C_{14} aldehydes by (Tani et al. 2000), however, activity for C_{12} and C_{16} hydroxy-o-fatty acids has been observed (Lu et al. 2010). Recently, two ADHs were characterized from *Geobacillus thermodenitrificans* NG80-2, an organism that degrades C_{15} to C_{36} alkanes using the LadA hydroxylase. Activity was detected from methanol to 1-triacontanol (C_{30}) for both ADHs, with 1-octanol being the preferred substrate for ADH2 and ethanol for ADH1 (Liu et al. 2009).

The use of ADHs in whole-cell bioconversions has been mostly focused on the production of chiral alcohols from ketones (Ernst et al. 2005) (Schroer et al. 2007). Using the ADH from *Lactobacillus brevis* and coupled cofactor regeneration with isopropanol, Schroer et al. reported the production of 797 g of (R)-methyl-3 hydroxybutanoate from methyl acetoacetate, with a space time yield of 29 g/L/h (Schroer et al. 2007). Examples of aliphatic alcohol oxidation in whole-cell transformations have been reported with commercially obtained *S. cerevisiae* for the conversion of hexanol to hexanal (Presecki et al. 2012) and 2-heptanol to 2-heptanone (Cappaert and Larroche 2004).

TABLE 3

Exemplary alcohol dehydrogenase enzymes.		
Organism	Gene Name	Accession No.
<i>Bactrocera oleae</i> (Olive fruit fly) (<i>Dacus oleae</i>)	ADH	Q9NAR7
<i>Cupriavidus necator</i> (<i>Alcaligenes eutrophus</i>) (<i>Ralstonia eutropha</i>)	adh	P14940
<i>Drosophila adriastola</i> (Fruit fly) (<i>Idiomyia adriastola</i>)	Adh	Q00669
<i>Drosophila affinisdisjuncta</i> (Fruit fly) (<i>Idiomyia affinisdisjuncta</i>)	Adh	P21518
<i>Drosophila ambigua</i> (Fruit fly)	Adh	P25139
<i>Drosophila borealis</i> (Fruit fly)	Adh	P48584
<i>Drosophila differens</i> (Fruit fly)	Adh	P22245
<i>Drosophila equinoxialis</i> (Fruit fly)	Adh	Q9NG42
<i>Drosophila flavomontana</i> (Fruit fly)	Adh	P48585
<i>Drosophila guanche</i> (Fruit fly)	Adh	Q09009
<i>Drosophila hawaiiensis</i> (Fruit fly)	Adh	P51549
<i>Drosophila heteroneura</i> (Fruit fly)	Adh	P21898
<i>Drosophila immigrans</i> (Fruit fly)	Adh	Q07588
<i>Drosophila insularis</i> (Fruit fly)	Adh	Q9NG40
<i>Drosophila lebanonensis</i> (Fruit fly) (<i>Scaptodrosophila lebanonensis</i>)	Adh	P10807
<i>Drosophila mauritiana</i> (Fruit fly)	Adh	P07162
<i>Drosophila madeirensis</i> (Fruit fly)	Adh	Q09010
<i>Drosophila mimica</i> (Fruit fly) (<i>Idiomyia mimica</i>)	Adh	Q00671
<i>Drosophila nigra</i> (Fruit fly) (<i>Idiomyia nigra</i>)	Adh	Q00672
<i>Drosophila orena</i> (Fruit fly)	Adh	P07159
<i>Drosophila pseudoobscura bogotana</i> (Fruit fly)	Adh	P84328
<i>Drosophila picticornis</i> (Fruit fly) (<i>Idiomyia picticornis</i>)	Adh	P23361
<i>Drosophila planitibia</i> (Fruit fly)	Adh	P23277
<i>Drosophila paulistorum</i> (Fruit fly)	Adh	Q9U8S9
<i>Drosophila silvestris</i> (Fruit fly)	Adh	P23278
<i>Drosophila subobscura</i> (Fruit fly)	Adh	Q03384
<i>Drosophila teissieri</i> (Fruit fly)	Adh	P28484
<i>Drosophila tsacasi</i> (Fruit fly)	Adh	P51550
<i>Fragaria ananassa</i> (Strawberry)	ADH	P17648

TABLE 3-continued

Exemplary alcohol dehydrogenase enzymes.		
Organism	Gene Name	Accession No.
<i>Malus domestica</i> (Apple) (<i>Pyrus malus</i>)	ADH	P48977
<i>Scaptomyza albobittata</i> (Fruit fly)	Adh	P25988
<i>Scaptomyza crassifemur</i> (Fruit fly) (<i>Drosophila crassifemur</i>)	Adh	Q00670
<i>Sulfolobus</i> sp. (strain RC3)	adh	P50381
<i>Zaprionus tuberculatus</i> (Vinegar fly)	Adh	P51552
<i>Geobacillus stearothermophilus</i> (<i>Bacillus stearothermophilus</i>)	adh	P42327
<i>Drosophila mayaguana</i> (Fruit fly)	Adh, Adh2	P25721
<i>Drosophila melanogaster</i> (Fruit fly)	Adh, CG3481	P00334
<i>Drosophila pseudoobscura pseudoobscura</i> (Fruit fly)	Adh, GA17214	Q6LCE4
<i>Drosophila simulans</i> (Fruit fly)	Adh, GD23968	Q24641
<i>Drosophila yakuba</i> (Fruit fly)	Adh, GE19037	P26719
<i>Drosophila ananassae</i> (Fruit fly)	Adh, GF14888	Q50L96
<i>Drosophila erecta</i> (Fruit fly)	Adh, GG25120	P28483
<i>Drosophila grimshawi</i> (Fruit fly) (<i>Idiomyia grimshawi</i>)	Adh, GH13025	P51551
<i>Drosophila willistoni</i> (Fruit fly)	Adh, GK18290	Q05114
<i>Drosophila persimilis</i> (Fruit fly)	Adh, GL25993	P37473
<i>Drosophila sechellia</i> (Fruit fly)	Adh, GM15656	Q9GN94
<i>Cupriavidus necator</i> (strain ATCC 17699/H16/DSM 428/Stanier 337) (<i>Ralstonia eutropha</i>)	adh, H16_A0757	Q0KDL6
<i>Mycobacterium tuberculosis</i> (strain CDC 1551/Oshkosh)	adh, MT1581	P9WQC2
<i>Staphylococcus aureus</i> (strain MW2)	adh, MW0568	Q8NXU1
<i>Mycobacterium tuberculosis</i> (strain ATCC 25618/H37Rv)	adh, Rv1530	P9WQC3
<i>Staphylococcus aureus</i> (strain N315)	adh, SA0562	Q7A742
<i>Staphylococcus aureus</i> (strain bovine RF122/ET3-1)	adh, SAB0557	Q2YSX0
<i>Sulfolobus acidocaldarius</i> (strain ATCC 33909/DSM 639/JCM 8929/NBRC 15157/NCIMB 11770)	adh, Saci_2057	Q4J781
<i>Staphylococcus aureus</i> (strain COL)	adh, SACOL0660	Q5HI63
<i>Staphylococcus aureus</i> (strain NCTC 8325)	adh, SAOUHSC_00608	Q2G0G1
<i>Staphylococcus aureus</i> (strain MRSA252)	adh, SAR0613	Q6GJ63
<i>Staphylococcus aureus</i> (strain MSSA476)	adh, SAS0573	Q6GBM4
<i>Staphylococcus aureus</i> (strain USA300)	adh, SAUSA300_0594	Q2FJ31
<i>Staphylococcus aureus</i> (strain Mu50/ATCC 700699)	adh, SAV0605	Q99W07
<i>Staphylococcus epidermidis</i> (strain ATCC 12228)	adh, SE_0375	Q8CQ56
<i>Staphylococcus epidermidis</i> (strain ATCC 35984/RP62A)	adh, SERP0257	Q5HRD6
<i>Sulfolobus solfataricus</i> (strain ATCC 35092/DSM 1617/JCM 11322/P2)	adh, SSO2536	P39462
<i>Sulfolobus tokodaii</i> (strain DSM 16993/JCM 10545/NBRC 100140/7)	adh, STK_25770	Q96XE0
<i>Anas platyrhynchos</i> (Domestic duck) (<i>Anas boschas</i>)	ADH1	P30350
<i>Apteryx australis</i> (Brown kiwi)	ADH1	P49645
<i>Ceratitis capitata</i> (Mediterranean fruit fly) (<i>Tephritis capitata</i>)	ADH1	P48814
<i>Ceratitis cosyra</i> (Mango fruit fly) (<i>Trypeta cosyra</i>)	ADH1	Q70UN9
<i>Gallus gallus</i> (Chicken)	ADH1	P23991
<i>Columba livia</i> (Domestic pigeon)	ADH1	P86883
<i>Coturnix coturnix japonica</i> (Japanese quail) (<i>Coturnix japonica</i>)	ADH1	P19631
<i>Drosophila hydei</i> (Fruit fly)	Adh1	P23236
<i>Drosophila montana</i> (Fruit fly)	Adh1	P48586
<i>Drosophila mettleri</i> (Fruit fly)	Adh1	P22246
<i>Drosophila mulleri</i> (Fruit fly)	Adh1	P07161
<i>Drosophila navojoa</i> (Fruit fly)	Adh1	P12854
<i>Geomys attwateri</i> (Attwater's pocket gopher) (<i>Geomys bursarius attwateri</i>)	ADH1	Q9Z2M2
<i>Geomys bursarius</i> (Plains pocket gopher)	ADH1	Q64413
<i>Geomys knoxjonesi</i> (Knox Jones's pocket gopher)	ADH1	Q64415
<i>Hordeum vulgare</i> (Barley)	ADH1	P05336
<i>Kluyveromyces marxianus</i> (Yeast) (<i>Candida kefir</i>)	ADH1	Q07288
<i>Zea mays</i> (Maize)	ADH1	P00333
<i>Mesocricetus auratus</i> (Golden hamster)	ADH1	P86885
<i>Pennisetum americanum</i> (Pearl millet) (<i>Pennisetum glaucum</i>)	ADH1	P14219
<i>Petunia hybrida</i> (<i>Petunia</i>)	ADH1	P25141
<i>Oryctolagus cuniculus</i> (Rabbit)	ADH1	Q03505
<i>Solanum tuberosum</i> (Potato)	ADH1	P14673

TABLE 3-continued

Exemplary alcohol dehydrogenase enzymes.		
Organism	Gene Name	Accession No.
<i>Struthio camelus</i> (Ostrich)	ADH1	P80338
<i>Trifolium repens</i> (Creeping white clover)	ADH1	P13603
<i>Zea luxurians</i> (Guatemalan teosinte) (<i>Euchlaena luxurians</i>)	ADH1	Q07264
<i>Saccharomyces cerevisiae</i> (strain ATCC 204508/S288c) (Baker's yeast)	ADH1, ADC1, YOL086C, O0947	P00330
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	ADH1, ADH, At1g77120, F22K20.19	P06525
<i>Schizosaccharomyces pombe</i> (strain 972/ATCC 24843) (Fission yeast)	adh1, adh, SPCC13B11.01	P00332
<i>Drosophila lacicola</i> (Fruit fly)	Adh1, Adh-1	Q27404
<i>Mus musculus</i> (Mouse)	Adh1, Adh-1	P00329
<i>Peromyscus maniculatus</i> (North American deer mouse)	ADH1, ADH-1	P41680
<i>Rattus norvegicus</i> (Rat)	Adh1, Adh-1	P06757
<i>Drosophila virilis</i> (Fruit fly)	Adh1, Adh-1, GJ18208	B4M8Y0
<i>Scheffersomyces stipitis</i> (strain ATCC 58785/CBS 6054/NBRC 10063/NRRL Y-11545) (Yeast) (<i>Pichia stipitis</i>)	ADH1, ADH2, PICST_68558	O00097
<i>Aspergillus flavus</i> (strain ATCC 200026/FGSC A1120/NRRL 3357/JCM 12722/SRRC 167)	adh1, AFLA_048690	P41747
<i>Neurospora crassa</i> (strain ATCC 24698/74-OR23-1A/CBS 708.71/DSM 1257/FGSC 987)	adh-1, B17C10.210, NCU01754	Q9P6C8
<i>Candida albicans</i> (Yeast)	ADH1, CAD	P43067
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	ADH1, DUPR11.3, Os11g0210300, LOC_Os11g10480, OsJ_032001	Q2R8Z5
<i>Drosophila mojavensis</i> (Fruit fly)	Adh1, GI17644	P09370
<i>Kluyveromyces lactis</i> (strain ATCC 8585/CBS 2359/DSM 70799/NBRC 1267/NRRL Y-1140/WM37) (Yeast) (<i>Candida sphaerica</i>)	ADH1, KLLA0F21010g	P20369
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	ADH1, OsL_034290	Q75ZX4
<i>Pongo abelii</i> (Sumatran orangutan) (<i>Pongo pygmaeus abelii</i>)	ADH1A	Q5RBP7
<i>Homo sapiens</i> (Human)	ADH1A, ADH1	P07327
<i>Macaca mulatta</i> (Rhesus macaque)	ADH1A, ADH1	P28469
<i>Pan troglodytes</i> (Chimpanzee)	ADH1B	Q5R1W2
<i>Papio hamadryas</i> (<i>Hamadryas</i> baboon)	ADH1B	P14139
<i>Homo sapiens</i> (Human)	ADH1B, ADH2	P00325
<i>Homo sapiens</i> (Human)	ADH1C, ADH3	P00326
<i>Papio hamadryas</i> (<i>Hamadryas</i> baboon)	ADH1C, ADH3	O97959
<i>Ceratitis capitata</i> (Mediterranean fruit fly) (<i>Tephritis capitata</i>)	ADH2	P48815
<i>Ceratitis cosyra</i> (Mango fruit fly) (<i>Trypeta cosyra</i>)	ADH2	Q70UP5
<i>Ceratitis rosa</i> (Natal fruit fly) (<i>Pterandrus rosa</i>)	ADH2	Q70UP6
<i>Drosophila arizonae</i> (Fruit fly)	Adh2	P27581
<i>Drosophila buzzatii</i> (Fruit fly)	Adh2	P25720
<i>Drosophila hydei</i> (Fruit fly)	Adh2	P23237
<i>Drosophila montana</i> (Fruit fly)	Adh2	P48587
<i>Drosophila mulleri</i> (Fruit fly)	Adh2	P07160
<i>Drosophila wheeleri</i> (Fruit fly)	Adh2	P24267
<i>Entamoeba histolytica</i>	ADH2	Q24803
<i>Hordeum vulgare</i> (Barley)	ADH2	P10847
<i>Kluyveromyces marxianus</i> (Yeast) (<i>Candida kefir</i>)	ADH2	Q9P4C2
<i>Zea mays</i> (Maize)	ADH2	P04707
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	ADH2	Q4R1E8
<i>Solanum lycopersicum</i> (Tomato) (<i>Lycopersicon esculentum</i>)	ADH2	P28032
<i>Solanum tuberosum</i> (Potato)	ADH2	P14674
<i>Scheffersomyces stipitis</i> (strain ATCC 58785/CBS 6054/NBRC 10063/NRRL Y-11545) (Yeast) (<i>Pichia stipitis</i>)	ADH2, ADH1, PICST_27980	O13309
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	ADH2, ADHIII, FDH1, At5g43940, MRH10.4	Q96533
<i>Saccharomyces cerevisiae</i> (strain ATCC 204508/S288c) (Baker's yeast)	ADH2, ADR2, YMR303C, YM9952.05C	P00331

TABLE 3-continued

Exemplary alcohol dehydrogenase enzymes.		
Organism	Gene Name	Accession No.
<i>Candida albicans</i> (strain SC5314/ATCC MYA-2876) (Yeast)	ADH2, Ca41C10.04, CaO19.12579, CaO19.5113	O94038
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	ADH2, DUPR11.1, Os11g0210500, LOC_Os11g10510	Q0ITW7
<i>Drosophila mojavensis</i> (Fruit fly)	Adh2, GI17643	P09369
<i>Kluyveromyces lactis</i> (strain ATCC 8585/CBS 2359/DSM 70799/NBRC 1267/NRRL Y-1140/WM37) (Yeast) (<i>Candida sphaerica</i>)	ADH2, KLLA0F18260g	P49383
<i>Oryctolagus cuniculus</i> (Rabbit)	ADH2-1	O46649
<i>Oryctolagus cuniculus</i> (Rabbit)	ADH2-2	O46650
<i>Hordeum vulgare</i> (Barley)	ADH3	P10848
<i>Solanum tuberosum</i> (Potato)	ADH3	P14675
<i>Kluyveromyces lactis</i> (strain ATCC 8585/CBS 2359/DSM 70799/NBRC 1267/NRRL Y-1140/WM37) (Yeast) (<i>Candida sphaerica</i>)	ADH3, KLLA0B09064g	P49384
<i>Saccharomyces cerevisiae</i> (strain ATCC 204508/S288c) (Baker's yeast)	ADH3, YMR083W, YM9582.08	P07246
<i>Homo sapiens</i> (Human)	ADH4	P08319
<i>Mus musculus</i> (Mouse)	Adh4	Q9QYY9
<i>Rattus norvegicus</i> (Rat)	Adh4	Q64563
<i>Struthio camelus</i> (Ostrich)	ADH4	P80468
<i>Kluyveromyces lactis</i> (strain ATCC 8585/ CBS 2359/DSM 70799/NBRC 1267/NRRL Y-1140/WM37) (Yeast) (<i>Candida sphaerica</i>)	ADH4, KLLA0F13530g	P49385
<i>Schizosaccharomyces pombe</i> (strain 972/ATCC 24843) (Fission yeast)	adh4, SPAC5H10.06c	Q09669
<i>Saccharomyces cerevisiae</i> (strain YJM789) (Baker's yeast)	ADH4, ZRG5, SCY_1818	A6ZTT5
<i>Saccharomyces cerevisiae</i> (strain ATCC 204508/S288c) (Baker's yeast)	ADH4, ZRG5, YGL256W, NRC465	P10127
<i>Saccharomyces pastorianus</i> (Lager yeast) (<i>Saccharomyces cerevisiae</i> × <i>Saccharomyces eubayanus</i>)	ADH5	Q6XQ67
<i>Bos taurus</i> (Bovine)	ADH5	Q3ZC42
<i>Equus caballus</i> (Horse)	ADH5	P19854
<i>Mus musculus</i> (Mouse)	Adh5, Adh-2, Adh2	P28474
<i>Rattus norvegicus</i> (Rat)	Adh5, Adh-2, Adh2	P12711
<i>Oryctolagus cuniculus</i> (Rabbit)	ADH5, ADH3	O19053
<i>Homo sapiens</i> (Human)	ADH5, ADHX, FDH	P11766
<i>Dictyostelium discoideum</i> (Slime mold)	adh5, DDB_G0281865	Q54TC2
<i>Saccharomyces cerevisiae</i> (strain ATCC 204508/S288c) (Baker's yeast)	ADH5, YBR145W, YBR1122	P38113
<i>Homo sapiens</i> (Human)	ADH6	P28332
<i>Peromyscus maniculatus</i> (North American deer mouse)	ADH6	P41681
<i>Pongo abelii</i> (Sumatran orangutan) (<i>Pongo pygmaeus abelii</i>)	ADH6	Q5R7Z8
<i>Rattus norvegicus</i> (Rat)	Adh6	Q5XI95
<i>Homo sapiens</i> (Human)	ADH7	P40394
<i>Rattus norvegicus</i> (Rat)	Adh7	P41682
<i>Mus musculus</i> (Mouse)	Adh7, Adh-3, Adh3	Q64437
<i>Mycobacterium tuberculosis</i> (strain CDC 1551/Oshkosh)	adhA, MT1911	P9WQC0
<i>Rhizobium meliloti</i> (strain 1021) (<i>Ensifer meliloti</i>) (<i>Sinorhizobium meliloti</i>)	adhA, RA0704, SMa1296	O31186
<i>Mycobacterium tuberculosis</i> (strain ATCC 25618/H37Rv)	adhA, Rv1862	P9WQC1
<i>Zymomonas mobilis</i> subsp. <i>mobilis</i> (strain ATCC 31821/ZM4/CP4)	adhA, ZMO1236	P20368
<i>Mycobacterium bovis</i> (strain ATCC BAA-935/ AF2122/97)	adhB, Mb0784c	Q7U1B9
<i>Mycobacterium tuberculosis</i> (strain CDC 1551/Oshkosh)	adhB, MT0786	P9WQC6
<i>Mycobacterium tuberculosis</i> (strain ATCC 25618/H37Rv)	adhB, Rv0761c, MTCY369.06c	P9WQC7
<i>Zymomonas mobilis</i> subsp. <i>mobilis</i> (strain ATCC 31821/ZM4/CP4)	adhB, ZMO1596	P0DJA2
<i>Zymomonas mobilis</i> subsp. <i>mobilis</i> (strain ATCC 10988/DSM 424/LMG 404/NCIMB 8938/NRRL B-806/ZM1)	adhB, Zmob_1541	F8DVL8

TABLE 3-continued

Exemplary alcohol dehydrogenase enzymes.		
Organism	Gene Name	Accession No.
<i>Mycobacterium tuberculosis</i> (strain CDC 1551/Oshkosh)	adhD, MT3171	P9WQB8
<i>Mycobacterium tuberculosis</i> (strain ATCC 25618/H37Rv)	adhD, Rv3086	P9WQB9
<i>Clostridium acetobutylicum</i> (strain ATCC 824/DSM 792/JCM 1419/LMG 5710/VKM B-1787)	adhE, aad, CA_P0162	P33744
<i>Escherichia coli</i> (strain K12)	adhE, ana, b1241, JW1228	P0A9Q7
<i>Escherichia coli</i> O157:H7	adhE, Z2016, ECs1741	P0A9Q8
<i>Rhodobacter sphaeroides</i> (strain ATCC 17023/2.4.1/NCIB 8253/DSM 158)	adhI, RHOS4_11650, RSP_2576	P72324
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	ADHIII, OsI_009236	A2XAZ3
<i>Escherichia coli</i> (strain K12)	adhP, yddN, b1478, JW1474	P39451
<i>Geobacillus stearothermophilus</i> (<i>Bacillus stearothermophilus</i>)	adhT	P12311
<i>Emericella nidulans</i> (strain FGSC A4/ATCC 38163/CBS 112.46/NRRL 194/M139)	alcA, AN8979	P08843
(<i>Aspergillus nidulans</i>)		
<i>Emericella nidulans</i> (strain FGSC A4/ATCC 38163/CBS 112.46/NRRL 194/M139)	alc, AN3741	P54202
(<i>Aspergillus nidulans</i>)		
<i>Emericella nidulans</i> (strain FGSC A4/ATCC 38163/CBS 112.46/NRRL 194/M139)	alcC, adh3, AN2286	P07754
(<i>Aspergillus nidulans</i>)		
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At1g22430, F12K8.22	Q9SK86
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At1g22440, F12K8.21	Q9SK87
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At1g32780, F6N18.16	A1L4Y2
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At1g64710, F13O11.3	Q8VZ49
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At4g22110, F1N20.210	Q0V7W6
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At5g24760, T4C12_30	Q8LEB2
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	At5g42250, K5J14.5	Q9FH04
<i>Zea mays</i> (Maize)	FDH	P93629
<i>Drosophila melanogaster</i> (Fruit fly)	Fdh, gfd, ODH, CG6598	P46415
<i>Bacillus subtilis</i> (strain 168)	gbsB, BSU31050	P71017
<i>Caenorhabditis elegans</i>	H24K24.3	Q17335
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	Os02g0815500, LOC_Os02g57040, OsJ_008550, P0643F09.4	Q0DWH1
<i>Mycobacterium tuberculosis</i> (strain ATCC 25618/H37Rv)	Rv1895	O07737
<i>Caenorhabditis elegans</i>	sodh-1, K12G11.3	Q17334
<i>Caenorhabditis elegans</i>	sodh-2, K12G11.4	O45687
<i>Pseudomonas</i> sp.	terPD	P33010
<i>Escherichia coli</i> (strain K12)	yiaY, b3589, JW5648	P37686
<i>Moraxella</i> sp. (strain TAE123)		P81786
<i>Alligator mississippiensis</i> (American alligator)		P80222
<i>Catharanthus roseus</i> (Madagascar periwinkle) (<i>Vinca rosea</i>)		P85440
<i>Gadus morhua</i> subsp. <i>callarias</i> (Baltic cod) (<i>Gadus callarias</i>)		P26325
<i>Naja naja</i> (Indian cobra)		P80512
<i>Pisum sativum</i> (Garden pea)		P12886
<i>Pelophylax perezii</i> (Perez's frog) (<i>Rana perezii</i>)		P22797
<i>Saara hardwickii</i> (Indian spiny-tailed lizard) (<i>Uromastyx hardwickii</i>)		P25405
<i>Saara hardwickii</i> (Indian spiny-tailed lizard) (<i>Uromastyx hardwickii</i>)		P25406

TABLE 3-continued

Exemplary alcohol dehydrogenase enzymes.		
Organism	Gene Name	Accession No.
<i>Equus caballus</i> (Horse)		P00327
<i>Equus caballus</i> (Horse)		P00328
<i>Geobacillus stearothermophilus</i> (<i>Bacillus stearothermophilus</i>)		P42328
<i>Gadus morhua</i> (Atlantic cod)		P81600
<i>Gadus morhua</i> (Atlantic cod)		P81601
<i>Myxine glutinosa</i> (Atlantic hagfish)		P80360
<i>Octopus vulgaris</i> (Common octopus)		P81431
<i>Pisum sativum</i> (Garden pea)		P80572
<i>Saara hardwickii</i> (Indian spiny-tailed lizard)		P80467
(<i>Uromastyx hardwickii</i>)		
<i>Scyliorhinus canicula</i> (Small-spotted catshark) (<i>Squalus canicula</i>)		P86884
<i>Sparus aurata</i> (Gilthead sea bream)		P79896

Alcohol Oxidase

The present disclosure describes enzymes that oxidize fatty alcohols to fatty aldehydes.

In some embodiments, an alcohol oxidase (AOX) is used to catalyze the conversion of a fatty alcohol to a fatty aldehyde. Alcohol oxidases catalyze the conversion of alcohols into corresponding aldehydes (or ketones) with electron transfer via the use of molecular oxygen to form hydrogen peroxide as a by-product. AOX enzymes utilize flavin adenine dinucleotide (FAD) as an essential cofactor and regenerate with the help of oxygen in the reaction medium. Catalase enzymes may be coupled with the AOX to avoid accumulation of the hydrogen peroxide via catalytic conversion into water and oxygen.

Based on the substrate specificities, AOXs may be categorized into four groups: (a) short chain alcohol oxidase, (b) long chain alcohol oxidase, (c) aromatic alcohol oxidase, and (d) secondary alcohol oxidase (Goswami et al. 2013). Depending on the chain length of the desired substrate, some members of these four groups are better suited than others as candidates for evaluation.

Short chain alcohol oxidases (including but not limited to those currently classified as EC 1.1.3.13, Table 4) catalyze the oxidation of lower chain length alcohol substrates in the range of C1-C8 carbons (van der Klei et al. 1991) (Ozimek et al. 2005). Aliphatic alcohol oxidases from methylotrophic yeasts such as *Candida boidinii* and *Komagataella pastoris* (formerly *Pichia pastoris*) catalyze the oxidation of primary alkanols to the corresponding aldehydes with a preference for unbranched short-chain aliphatic alcohols. The most broad substrate specificity is found for alcohol oxidase from the *Pichia pastoris* including propargyl alcohol, 2-chloroethanol, 2-cyanoethanol (Dienys et al. 2003). The major challenge encountered in alcohol oxidation is the high reactivity of the aldehyde product. Utilization of a two liquid phase system (water/solvent) can provide in-situ removal of the aldehyde product from the reaction phase before it is further converted to the acid. For example, hexanal production from hexanol using *Pichia pastoris* alcohol oxidase coupled with bovine liver catalase was achieved in a biphasic system by taking advantage of the presence of a stable alcohol oxidase in aqueous phase (Karra-Chaabouni et al. 2003). For example, alcohol oxidase from *Pichia pastoris* was able to oxidize aliphatic alcohols of C6 to C11 when used biphasic organic reaction system (Murray and Duff 1990). Methods for using alcohol oxidases in a biphasic

system according to (Karra-Chaabouni et al. 2003) and (Murray and Duff 1990) are incorporated by reference in their entirety.

Long chain alcohol oxidases (including but not limited to those currently classified as EC 1.1.3.20; Table 5) include fatty alcohol oxidases, long chain fatty acid oxidases, and long chain fatty alcohol oxidases that oxidize alcohol substrates with carbon chain length of greater than six (Goswami et al. 2013). Banthorpe et al. reported a long chain alcohol oxidase purified from the leaves of *Tanacetum vulgare* that was able to oxidize saturated and unsaturated long chain alcohol substrates including hex-trans-2-en-1-ol and octan-1-ol (Banthorpe 1976) (Cardemil 1978). Other plant species, including *Simmondsia chinensis* (Moreau, R. A., Huang 1979), *Arabidopsis thaliana* (Cheng et al. 2004), and *Lotus japonicas* (Zhao et al. 2008) have also been reported as sources of long chain alcohol oxidases. Fatty alcohol oxidases are mostly reported from yeast species (Hommel and Ratledge 1990) (Vanhanen et al. 2000) (Hommel et al. 1994) (Kemp et al. 1990) and these enzymes play an important role in long chain fatty acid metabolism (Cheng et al. 2005). Fatty alcohol oxidases from yeast species that degrade and grow on long chain alkanes and fatty acid catalyze the oxidation of fatty alcohols. Fatty alcohol oxidase from *Candida tropicalis* has been isolated as microsomal cell fractions and characterized for a range of substrates (Eirich et al. 2004) (Kemp et al. 1988) (Kemp et al. 1991) (Mauersberger et al. 1992). Significant activity is observed for primary alcohols of length C₈ to C₁₆ with reported K_M in the 10-50 M range (Eirich et al. 2004). Alcohol oxidases described may be used for the conversion of medium chain aliphatic alcohols to aldehydes as described, for example, for whole-cells *Candida boidinii* (Gabelman and Luzio 1997), and *Pichia pastoris* (Duff and Murray 1988) (Murray and Duff 1990). Long chain alcohol oxidases from filamentous fungi were produced during growth on hydrocarbon substrates (Kumar and Goswami 2006) (Savitha and Ratledge 1991). The long chain fatty alcohol oxidase (LjFAO1) from *Lotus japonicas* has been heterologously expressed in *E. coli* and exhibited broad substrate specificity for alcohol oxidation including 1-dodecanol and 1-hexadecanol (Zhao et al. 2008).

TABLE 4

Alcohol oxidase enzymes capable of oxidizing short chain alcohols (EC 1.1.3.13).		
Organism	Gene names	Accession No.
<i>Komagataella pastoris</i> (strain ATCC 76273/CBS 7435/CECT 11047/NRRL Y-11430/Wegner 21-1) (Yeast) (<i>Pichia pastoris</i>)	AOX1 PP7435_Ch4-0130	F2QY27
<i>Komagataella pastoris</i> (strain GS115/ATCC 20864) (Yeast) (<i>Pichia pastoris</i>)	AOX1 PAS_chr4_0821	P04842
<i>Komagataella pastoris</i> (strain ATCC 76273/CBS 7435/CECT 11047/NRRL Y-11430/Wegner 21-1) (Yeast) (<i>Pichia pastoris</i>)	AOX2 PP7435_Ch4-0863	F2R038
<i>Komagataella pastoris</i> (strain GS115/ATCC 20864) (Yeast) (<i>Pichia pastoris</i>)	AOX2 PAS_chr4_0152	C4R702
<i>Candida boidinii</i> (Yeast)	AOD1	Q00922
<i>Pichia angusta</i> (Yeast) (<i>Hansenula polymorpha</i>)	MOX	P04841
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_10802	M5CC52
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	MOX BN14_12214	M5CF32
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_10691	M5CAV1
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_09479	M5C7F4
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_10803	M5CB66
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_09900	M5C9N9
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_08302	M5C2L8
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	MOX BN14_09408	M5C784
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	MOX BN14_09478	M5C8F8
<i>Thanatephorus cucumeris</i> (strain AG1-IB/isolate 7/3/14) (Lettuce bottom rot fungus) (<i>Rhizoctonia solani</i>)	AOD1 BN14_11356	M5CH40
<i>Ogataea henricii</i>	AOD1	A5LGF0
<i>Candida methanosorbosa</i>	AOD1	A5LGE5
<i>Candida methanolovescens</i>	AOD1	A5LGE4
<i>Candida succiphila</i>	AOD1	A5LGE6
<i>Aspergillus niger</i> (strain CBS 513.88/FGSC A1513)	An15g02200	A2R501
<i>Aspergillus niger</i> (strain CBS 513.88/FGSC A1513)	An18g05480	A2RB46
<i>Moniliophthora perniciosa</i> (Witches'-broom disease fungus) (<i>Marasmius pernicius</i>)		I7CMK2
<i>Candida cariosilignicola</i>	AOD1	A5LGE3
<i>Candida pignaliae</i>	AOD1	A5LGE1
<i>Candida pignaliae</i>	AOD2	A5LGE2
<i>Candida sonorensis</i>	AOD1	A5LGD9
<i>Candida sonorensis</i>	AOD2	A5LGE0
<i>Pichia naganishii</i>	AOD1	A5LGF2
<i>Ogataea minuta</i>	AOD1	A5LGF1
<i>Ogataea philodendri</i>	AOD1	A5LGF3
<i>Ogataea wickerhamii</i>	AOD1	A5LGE8
<i>Kuraishia capsulata</i>	AOD1	A5LGE7
<i>Talaromyces stipitatus</i> (strain ATCC 10500/CBS 375.48/QM 6759/NRRL 1006) (<i>Penicillium stipitatum</i>)	TSTA_021940	B8MHF8
<i>Talaromyces stipitatus</i> (strain ATCC 10500/CBS 375.48/QM 6759/NRRL 1006) (<i>Penicillium stipitatum</i>)	TSTA_065150	B8LTH7
<i>Talaromyces stipitatus</i> (strain ATCC 10500/CBS 375.48/QM 6759/NRRL 1006) (<i>Penicillium stipitatum</i>)	TSTA_065150	B8LTH8
<i>Talaromyces stipitatus</i> (strain ATCC 10500/CBS 375.48/QM 6759/NRRL 1006) (<i>Penicillium stipitatum</i>)	TSTA_000410	B8MSB1
<i>Ogataea glucozyma</i>	AOD1	A5LGE9
<i>Ogataea parapolyomorpha</i> (strain DL-1/ATCC 26012/NRRL Y-7560) (Yeast) (<i>Hansenula polymorpha</i>)	HPODL_03886	W1QCJ3
<i>Gloeophyllum trabeum</i> (Brown rot fungus)	AOX	A8DPS4
<i>Pichia angusta</i> (Yeast) (<i>Hansenula polymorpha</i>)	mox1	A6PZG8
<i>Pichia trehalophila</i>	AOD1	A5LGF4
<i>Pichia angusta</i> (Yeast) (<i>Hansenula polymorpha</i>)	mox1	A6PZG9
<i>Pichia angusta</i> (Yeast) (<i>Hansenula polymorpha</i>)	mox1	A6PZG7
<i>Ixodes scapularis</i> (Black-legged tick) (Deer tick)	IscW_ISCW017898	B7PIZ7

TABLE 5

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Lotus japonicus</i> (<i>Lotus corniculatus</i> var. <i>japonicus</i>)	FAO1	B5WWZ8
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	FAO1 At1g03990 F21M11.7	Q9ZWB9
<i>Lotus japonicus</i> (<i>Lotus corniculatus</i> var. <i>japonicus</i>)	FAO2	B5WWZ9
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	FAO3 At3g23410 MLM24.14 MLM24.23	Q9LW56
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	FAO4A At4g19380 T5K18.160	O65709
<i>Arabidopsis thaliana</i> (Mouse-ear cress)	FAO4B At4g28570 T5F17.20	Q94BP3
<i>Microbotryum violaceum</i> (strain p1A1 Lamole)	MVLG_06864	U5HIL4
(Anther smut fungus) (<i>Ustilago violacea</i>)		
<i>Ajellomyces dermatitidis</i> ATCC 26199	BDFG_03507	T5BNQ0
<i>Gibberella zeae</i> (strain PH-1/ATCC MYA-4620/FGSC 9075/NRRL 31084) (Wheat head blight fungus) (<i>Fusarium graminearum</i>)	FG06918.1 FGSG_06918	I1RS14
<i>Pichia sorbitophila</i> (strain ATCC MYA-4447/BCRC 22081/CBS 7064/NBRC 10061/NRRL Y-12695) (Hybrid yeast)	Piso0_004410 GNLVRS01_PISO0K16268g GNLVRS01_PISO0L16269g	G8Y5E1
<i>Emericella nidulans</i> (strain FGSC A4/ATCC 38163/CBS 112.46/NRRL 194/M139) (<i>Aspergillus nidulans</i>)	AN0623.2 ANIA_00623	Q5BFQ7
<i>Pyrenophora tritici-repentis</i> (strain Pt-1C-BFP) (Wheat tan spot fungus) (<i>Drechslera tritici-repentis</i>)	PTRG_10154	B2WJW5
<i>Paracoccidioides lutzii</i> (strain ATCC MYA-826/Pb01) (<i>Paracoccidioides brasiliensis</i>)	PAAG_09117	C1HEC6
<i>Candida parapsilosis</i> (strain CDC 317/ATCC MYA-4646) (Yeast) (<i>Monilia parapsilosis</i>)	CPAR2_204420	G8BG15
<i>Pseudozyma brasiliensis</i> (strain GHG001) (Yeast)	PSEUBRA_SCAF2g03010	V5GPS6
<i>Candida parapsilosis</i> (strain CDC 317/ATCC MYA-4646) (Yeast) (<i>Monilia parapsilosis</i>)	CPAR2_204430	G8BG16
<i>Sclerotinia borealis</i> F-4157	SBOR_5750	W9CDE2
<i>Sordaria macrospora</i> (strain ATCC MYA-333/DSM 997/K(L3346)/K-hell)	SMAC_06361	F7W6K4
<i>Sordaria macrospora</i> (strain ATCC MYA-333/DSM 997/K(L3346)/K-hell)	SMAC_01933	F7VSA1
<i>Meyerozyma guilliermondii</i> (strain ATCC 6260/CBS 566/DSM 6381/JCM 1539/NBRC 10279/NRRL Y-324) (Yeast) (<i>Candida guilliermondii</i>)	PGUG_03467	A5DJL6
<i>Trichophyton rubrum</i> CBS 202.88	H107_00669	A0A023ATC5
<i>Arthrotrys oligospora</i> (strain ATCC 24927/CBS 115.81/DSM 1491) (Nematode-trapping fungus) (<i>Didymozoopphaga oligospora</i>)	AOL_s00097g516	G1XJ19
<i>Scheffersomyces stipitis</i> (strain ATCC 58785/CBS 6054/NBRC 10063/NRRL Y-11545) (Yeast) (<i>Pichia stipitis</i>)	FAO1 PICST_90828	A3LYX9
<i>Scheffersomyces stipitis</i> (strain ATCC 58785/CBS 6054/NBRC 10063/NRRL Y-11545) (Yeast) (<i>Pichia stipitis</i>)	FAO2 PICST_32359	A3LW61
<i>Aspergillus oryzae</i> (strain 3.042) (Yellow koji mold)	Ao3042_09114	I8TL25
<i>Fusarium oxysporum</i> (strain Fo5176) (<i>Fusarium</i> vascular wilt)	FOXB_17532	F9GFU8
<i>Rhizopus delemar</i> (strain RA 99-880/ATCC MYA-4621/FGSC 9543/NRRL 43880) (Mucormycosis agent) (<i>Rhizopus arrhizus</i> var. <i>delemar</i>)	RO3G_08271	I1C536
<i>Rhizopus delemar</i> (strain RA 99-880/ATCC MYA-4621/FGSC 9543/NRRL 43880) (Mucormycosis agent) (<i>Rhizopus arrhizus</i> var. <i>delemar</i>)	RO3G_00154	I1BGX0
<i>Fusarium oxysporum</i> (strain Fo5176) (<i>Fusarium</i> vascular wilt)	FOXB_07532	F9FMA2
<i>Penicillium roqueforti</i>	PROQFM164_S02g001772	W6QPY1
<i>Aspergillus clavatus</i> (strain ATCC 1007/CBS 513.65/DSM 816/NCTC 3887/NRRL 1)	ACLA_018400	A1CNB5
<i>Arthroderma otae</i> (strain ATCC MYA-4605/CBS 113480) (<i>Microsporium canis</i>)	MCYG_08732	C5G1B0
<i>Trichophyton tonsurans</i> (strain CBS 112818) (Scalp ringworm fungus)	TESG_07214	F2S8I2
<i>Colletotrichum higginsianum</i> (strain IMI 349063) (Crucifer anthracnose fungus)	CH063_13441	H1VUE7
<i>Ajellomyces capsulatus</i> (strain H143) (Darling's disease fungus) (<i>Histoplasma capsulatum</i>)	HCDG_07658	C6HN77
<i>Trichophyton rubrum</i> (strain ATCC MYA-4607/CBS 118892) (Athlete's foot fungus)	TERG_08235	F2T096
<i>Cochliobolus heterostrophus</i> (strain C5/ATCC 48332/race O) (Southern corn leaf blight fungus) (<i>Bipolaris maydis</i>)	COCHEDRAFT_1201414	M2UMT9
<i>Candida orthopsilosis</i> (strain 90-125) (Yeast)	CORT_0D04510	H8X643
<i>Candida orthopsilosis</i> (strain 90-125) (Yeast)	CORT_0D04520	H8X644

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Candida orthopsilosis</i> (strain 90-125) (Yeast)	CORT_0D04530	H8X645
<i>Pseudozyma aphidis</i> DSM 70725	PaG_03027	W3VP49
<i>Coccidioides posadasii</i> (strain C735) (Valley fever fungus)	CPC735_000380	C5P005
<i>Magnaporthe oryzae</i> (strain P131) (Rice blast fungus) (<i>Pyricularia oryzae</i>)	OOW_P131scaffold01214g15	L7IZ92
<i>Neurospora tetrasperma</i> (strain FGSC 2508/ATCC MYA-4615/P0657)	NEUTE1DRAFT_82541	F8MKD1
<i>Hypocrea virens</i> (strain Gv29-8/FGSC 10586) (<i>Gliocladium virens</i>) (<i>Trichoderma virens</i>)	TRIVIDRAFT_54537	G9MMY7
<i>Hypocrea virens</i> (strain Gv29-8/FGSC 10586) (<i>Gliocladium virens</i>) (<i>Trichoderma virens</i>)	TRIVIDRAFT_53801	G9MT89
<i>Aspergillus niger</i> (strain CBS 513.88/FGSC A1513)	An01g09620	A2Q9Z3
<i>Verticillium dahliae</i> (strain VdLs.17/ATCC MYA-4575/FGSC 10137) (<i>Verticillium</i> wilt)	VDAG_05780	G2X6J8
<i>Ustilago maydis</i> (strain 521/FGSC 9021) (Corn smut fungus)	UM02023.1	Q4PCZ0
<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> MN25	FOWG_13006	W9LNI9
<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> MN25	FOWG_02542	W9N9Z1
<i>Candida tropicalis</i> (Yeast)	FAO1	Q6QIR6
<i>Magnaporthe oryzae</i> (strain 70-15/ATCC MYA-4617/FGSC 8958) (Rice blast fungus) (<i>Pyricularia oryzae</i>)	MGG_11317	G4MVK1
<i>Candida tropicalis</i> (Yeast)	faot	Q9P8D9
<i>Candida tropicalis</i> (Yeast)	FAO2a	Q6QIR5
<i>Phaeosphaeria nodorum</i> (strain SN15/ATCC MYA-4574/FGSC 10173) (Glume blotch fungus) (<i>Septoria nodorum</i>)	SNOG_02371	Q0V0U3
<i>Candida tropicalis</i> (Yeast)	FAO2b	Q6QIR4
<i>Pestalotiopsis fici</i> W106-1	PFICI_11209	W3WU04
<i>Magnaporthe oryzae</i> (strain Y34) (Rice blast fungus) (<i>Pyricularia oryzae</i>)	OOU_Y34scaffold00240g57	L7IFT5
<i>Pseudogymnoascus destructans</i> (strain ATCC MYA-4855/20631-21) (Bat white-nose syndrome fungus) (<i>Geomyces destructans</i>)	GMDG_01756	L8G0G6
<i>Pseudogymnoascus destructans</i> (strain ATCC MYA-4855/20631-21) (Bat white-nose syndrome fungus) (<i>Geomyces destructans</i>)	GMDG_04950	L8GCY2
<i>Mycosphaerella fijiensis</i> (strain CIRAD86) (Black leaf streak disease fungus) (<i>Pseudocercospora fijiensis</i>)	MYCFIDRAFT_52380	M2Z831
<i>Bipolaris oryzae</i> ATCC 44560	COCMIDRAFT_84580	W7A0I8
<i>Cladophialophora psammophila</i> CBS 110553	A1O5_08147	W9WTM9
<i>Fusarium oxysporum</i> f. sp. <i>melonis</i> 26406	FOMG_05173	X0AEE6
<i>Fusarium oxysporum</i> f. sp. <i>melonis</i> 26406	FOMG_17829	W9ZBB7
<i>Cyphellophora europaea</i> CBS 101466	HMPREF1541_02174	W2S2S5
<i>Aspergillus kawachii</i> (strain NBRC 4308) (White koji mold) (<i>Aspergillus awamori</i> var. <i>kawachi</i>)	AKAW_00147	G7X626
<i>Aspergillus terreus</i> (strain NIH 2624/FGSC A1156)	ATEG_05086	Q0CMJ8
<i>Coccidioides immitis</i> (strain RS) (Valley fever fungus)	CIMG_02987	J3KAI8
<i>Ajellomyces dermatitidis</i> (strain ER-3/ATCC MYA-2586) (<i>Blastomyces dermatitidis</i>)	BDCG_04701	C5GLS5
<i>Fusarium oxysporum</i> f. sp. <i>cubense</i> (strain race 1) (Panama disease fungus)	FOC1_g10013865	N4U732
<i>Rhodotorula glutinis</i> (strain ATCC 204091/IIP 30/MTCC 1151) (Yeast)	RTG_00643	G0SVU8
<i>Aspergillus niger</i> (strain ATCC 1015/CBS 113.46/FGSC A1144/LSHB Ac4/NCTC 3858a/NRRL 328/USDA 3528.7)	ASPNIDRAFT_35778	G3XTM6
<i>Candida cloacae</i>	fao1	Q9P8D8
<i>Candida cloacae</i>	fao2	Q9P8D7
<i>Fusarium oxysporum</i> f. sp. <i>cubense</i> (strain race 1) (Panama disease fungus)	FOC1_g10006358	N4TUH3
<i>Candida albicans</i> (strain SC5314/ATCC MYA-2876) (Yeast)	FAO1 CaO19.13562	Q59RS8
<i>Candida albicans</i> (strain SC5314/ATCC MYA-2876) (Yeast)	orf19.13562	
<i>Chaetomium thermophilum</i> (strain DSM 1495/CBS 144.50/IMI 039719)	FAO1 CaO19.6143	Q59RP0
<i>Mucor circinelloides</i> f. <i>circinelloides</i> (strain 1006PhL) (Mucormycosis agent) (<i>Calyptromyces circinelloides</i>)	orf19.6143	
	CTHT_0018560	G0S2U9
	HMPREF1544_05296	S2JDN0

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Mucor circinelloides</i> f. <i>circinelloides</i> (strain 1006PhL) (Mucormycosis agent) (<i>Calyptromyces circinelloides</i>)	HMPREF1544_05295	S2JYP5
<i>Mucor circinelloides</i> f. <i>circinelloides</i> (strain 1006PhL) (Mucormycosis agent) (<i>Calyptromyces circinelloides</i>)	HMPREF1544_06348	S2JVK9
<i>Botryotinia fuckeliana</i> (strain BcDW1) (Noble rot fungus) (<i>Botrytis cinerea</i>)	BcDW1_6807	M7UD26
<i>Podospora anserina</i> (strain S/ATCC MYA-4624/DSM 980/FGSC 10383) (<i>Pleurage anserina</i>)	PODANS_5_13040	B2AFD8
<i>Neosartorya fumigata</i> (strain ATCC MYA-4609/ Af293/CBS 101355/FGSC A1100) (<i>Aspergillus fumigatus</i>)	AFUA_1G17110	Q4WR91
<i>Fusarium oxysporum</i> f. sp. <i>vasinfectum</i> 25433	FOTG_00686	X0MEE6
<i>Fusarium oxysporum</i> f. sp. <i>vasinfectum</i> 25433	FOTG_12485	X0LE98
<i>Trichophyton interdigitale</i> H6	H101_06625	A0A022U717
<i>Beauveria bassiana</i> (strain ARSEF 2860) (White muscardine disease fungus) (<i>Tritirachium shioteae</i>)	BBA_04100	J4UNY3
<i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> 26381	FOCG_00843	X0GQ62
<i>Fusarium oxysporum</i> f. sp. <i>radicis-lycopersici</i> 26381	FOCG_15170	X0F4T1
<i>Neurospora tetrasperma</i> (strain FGSC 2509/P0656)	NEUTE2DRAFT_88670	G4UNN6
<i>Pseudozyma hubeiensis</i> (strain SY62) (Yeast)	PHSY_000086	R9NVU1
<i>Lodderomyces elongisporus</i> (strain ATCC 11503/ CBS 2605/JCM 1781/NBRC 1676/NRRL YB-4239) (Yeast) (<i>Saccharomyces elongisporus</i>)	LELG_03289	A5E102
<i>Malassezia globosa</i> (strain ATCC MYA-4612/CBS 7966) (Dandruff-associated fungus)	MGL_3855	A8QAY8
<i>Byssosclamyces spectabilis</i> (strain No. 5/NBRC 109023) (<i>Paecilomyces variotii</i>)	PVAR5_7014	V5GBL6
<i>Ajellomyces capsulatus</i> (strain H88) (Darling's disease fungus) (<i>Histoplasma capsulatum</i>)	HCEG_03274	F0UF47
<i>Trichosporon asahii</i> var. <i>asahii</i> (strain ATCC 90039/ CBS 2479/JCM 2466/KCTC 7840/NCYC 2677/ UAMH 7654) (Yeast)	A1Q1_03669	J6FBP4
<i>Penicillium oxalicum</i> (strain 114-2/CGMCC 5302) (<i>Penicillium decumbens</i>)	PDE_00027	S7Z8U8
<i>Fusarium oxysporum</i> f. sp. <i>conglutinans</i> race 2 54008	FOPG_02304	X0IBE3
<i>Fusarium oxysporum</i> f. sp. <i>conglutinans</i> race 2 54008	FOPG_13066	X0H540
<i>Fusarium oxysporum</i> f. sp. <i>raphani</i> 54005	FOQG_00704	X0D1G8
<i>Fusarium oxysporum</i> f. sp. <i>raphani</i> 54005	FOQG_10402	X0C482
<i>Metarhizium acridum</i> (strain CQMa 102)	MAC_03115	E9DZR7
<i>Arthroderma benhamiae</i> (strain ATCC MYA-4681/ CBS 112371) (<i>Trichophyton mentagrophytes</i>)	ARB_02250	D4B1C1
<i>Fusarium oxysporum</i> f. sp. <i>cubense</i> tropical race 4 54006	FOIG_12161	X0JFI6
<i>Fusarium oxysporum</i> f. sp. <i>cubense</i> tropical race 4 54006	FOIG_12751	X0JDU5
<i>Cochliobolus heterostrophus</i> (strain C4/ATCC 48331/race T) (Southern corn leaf blight fungus) (<i>Bipolaris maydis</i>)	COCC4DRAFT_52836	N4WZZ0
<i>Trichosporon asahii</i> var. <i>asahii</i> (strain CBS 8904) (Yeast)	A1Q2_00631	K1VZW1
<i>Mycosphaerella graminicola</i> (strain CBS 115943/ IPO323) (Speckled leaf blotch fungus) (<i>Septoria tritici</i>)	MYCGRDRAFT_37086	F9X375
<i>Botryotinia fuckeliana</i> (strain T4) (Noble rot fungus) (<i>Botrytis cinerea</i>)	BofuT4_P072020.1	G2XQ18
<i>Metarhizium anisopliae</i> (strain ARSEF 23/ATCC MYA-3075)	MAA_05783	E9F0I4
<i>Cladophialophora carrionii</i> CBS 160.54	G647_05801	V9DAR1
<i>Coccidioides posadasii</i> (strain RMSCC 757/ Silveira) (Valley fever fungus)	CPSG_09174	E9DH75
<i>Rhodospiridium toruloides</i> (strain NP11) (Yeast) (<i>Rhodotorula gracilis</i>)	RHTO_06879	M7X159
<i>Puccinia graminis</i> f. sp. <i>tritici</i> (strain CRL 75-36-700-3/race SCCL) (Black stem rust fungus)	PGTG_10521	E3KIL8
<i>Trichophyton rubrum</i> CBS 288.86	H103_00624	A0A022WG28
<i>Colletotrichum fioriniae</i> PJ7	CFIO01_08202	A0A010RKZ4
<i>Trichophyton rubrum</i> CBS 289.86	H104_00611	A0A022XB46

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Cladophialophora yegresii</i> CBS 114405	A1O7_02579	W9WC55
<i>Colletotrichum orbiculare</i> (strain 104-T/ATCC 96160/CBS 514.97/LARS 414/MAFF 240422) (Cucumber anthracnose fungus) (<i>Colletotrichum lagenarium</i>)	Cob_10151	N4VFP3
<i>Drechslerella stenobrocha</i> 248	DRE_03459	W7IDL6
<i>Neosartorya fumigata</i> (strain CEA10/CBS 144.89/FGSC A1163) (<i>Aspergillus fumigatus</i>)	AFUB_016500	B0XP90
<i>Thielavia terrestris</i> (strain ATCC 38088/NRRL 8126) (<i>Acremonium alabamense</i>)	THITE_2117674	G2R8H9
<i>Gibberella fujikuroi</i> (strain CBS 195.34/IMI 58289/NRRL A-6831) (Bakanae and foot rot disease fungus) (<i>Fusarium fujikuroi</i>)	FFUJ_02948	S0DZP7
<i>Gibberella fujikuroi</i> (strain CBS 195.34/IMI 58289/NRRL A-6831) (Bakanae and foot rot disease fungus) (<i>Fusarium fujikuroi</i>)	FFUJ_12030	S0EMC6
<i>Aspergillus flavus</i> (strain ATCC 200026/FGSC A1120/NRRL 3357/JCM 12722/SRRC 167)	AFLA_109870	B8N941
<i>Togninia minima</i> (strain UCR-PA7) (Esca disease fungus) (<i>Phaeoacremonium aleophilum</i>)	UCRPA7_1719	R8BTZ6
<i>Ajellomyces dermatitidis</i> (strain ATCC 18188/CBS 674.68) (<i>Blastomyces dermatitidis</i>)	BDDG_09783	F2TUC0
<i>Macrophomina phaseolina</i> (strain MS6) (Charcoal rot fungus)	MPH_10582	K2RHA5
<i>Neurospora crassa</i> (strain ATCC 24698/74-OR23-1A/CBS 708.71/DSM 1257/FGSC 987)	NCU08977	Q7S2Z2
<i>Neosartorya fischeri</i> (strain ATCC 1020/DSM 3700/FGSC A1164/NRRL 181) (<i>Aspergillus fischerianus</i>)	NFIA_008260	A1D156
<i>Fusarium pseudograminearum</i> (strain CS3096) (Wheat and barley crown-rot fungus)	FPSE_11742	K3U9J5
<i>Spathaspora passalidarum</i> (strain NRRL Y-27907/11-Y1)	SPAPADRAFT_54193	G3AJP0
<i>Spathaspora passalidarum</i> (strain NRRL Y-27907/11-Y1)	SPAPADRAFT_67198	G3ANX7
<i>Trichophyton verrucosum</i> (strain HKI 0517)	TRV_07960	D4DL86
<i>Arthroderma gypseum</i> (strain ATCC MYA-4604/CBS 118893) (<i>Microsporum gypseum</i>)	MGYG_07264	E4V2J0
<i>Hypocrea jecorina</i> (strain QM6a) (<i>Trichoderma reesei</i>)	TRIREDRAFT_43893	G0R7P8
<i>Trichophyton rubrum</i> MR1448	H110_00629	A0A022Z1G4
<i>Aspergillus ruber</i> CBS 135680	EURHEDRAFT_512125	A0A017SPR0
<i>Glarea lozoyensis</i> (strain ATCC 20868/MF5171)	GLAREA_04397	S3D6C1
<i>Setosphaeria turcica</i> (strain 28A) (Northern leaf blight fungus) (<i>Exserohilum turcicum</i>)	SETTUDRAFT_20639	R0K6H8
<i>Paracoccidioides brasiliensis</i> (strain Pb18)	PADG_06552	C1GH16
<i>Fusarium oxysporum</i> Fo47	FOZG_13577	W9JPG9
<i>Fusarium oxysporum</i> Fo47	FOZG_05344	W9KPH3
<i>Trichophyton rubrum</i> MR1459	H113_00628	A0A022ZY09
<i>Penicillium marneffei</i> (strain ATCC 18224/CBS 334.59/QM 7333)	PMAA_075740	B6QBY3
<i>Sphaerulina musiva</i> (strain SO2202) (Poplar stem canker fungus) (<i>Septoria musiva</i>)	SEPMUDRAFT_154026	M3DAK6
<i>Gibberella moniliformis</i> (strain M3125/FGSC 7600) (Maize ear and stalk rot fungus) (<i>Fusarium verticillioides</i>)	FVEG_10526	W7N4P8
<i>Gibberella moniliformis</i> (strain M3125/FGSC 7600) (Maize ear and stalk rot fungus) (<i>Fusarium verticillioides</i>)	FVEG_08281	W7MVR9
<i>Pseudozyma antarctica</i> (strain T-34) (Yeast) (<i>Candida antarctica</i>)	PANT_22d00298	M9MGF2
<i>Paracoccidioides brasiliensis</i> (strain Pb03)	PABG_07795	C0SJD4
<i>Rhizophagus irregularis</i> (strain DAOM 181602/DAOM 197198/MUCL 43194) (Arbuscular mycorrhizal fungus) (<i>Glomus intraradices</i>)	GLOINDRAFT_82554	U9TF61
<i>Penicillium chrysogenum</i> (strain ATCC 28089/DSM 1075/Wisconsin 54-1255) (<i>Penicillium notatum</i>)	Pc21g23700 PCH_Pc21g23700	B6HJ58
<i>Baudoinia compniacensis</i> (strain UAMH 10762) (Angels' share fungus)	BAUCODRAFT_274597	M2M6Z5
<i>Hypocrea atroviridis</i> (strain ATCC 20476/IMI 206040) (<i>Trichoderma atroviride</i>)	TRIATDRAFT_280929	G9NJ32
<i>Colletotrichum gloeosporioides</i> (strain Cg-14) (Anthracnose fungus) (<i>Glomerella cingulata</i>)	CGLO_06642	T0LPH0

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Cordyceps militaris</i> (strain CM01) (Caterpillar fungus)	CCM_02665	G3JB34
<i>Pyronema omphalodes</i> (strain CBS 100304) (<i>Pyronema confluens</i>)	PCON_13062	U4LKE9
<i>Colletotrichum graminicola</i> (strain M1.001/M2/FGSC 10212) (Maize anthracnose fungus) (<i>Glomerella graminicola</i>)	GLRG_08499	E3QR67
<i>Glarea lozoyensis</i> (strain ATCC 74030/MF5533)	M7L_2117	H0EHX4
<i>Fusarium oxysporum</i> f. sp. <i>cubense</i> (strain race 4) (Panama disease fungus)	FOC4_g10002493	N1S969
<i>Fusarium oxysporum</i> f. sp. <i>cubense</i> (strain race 4) (Panama disease fungus)	FOC4_g10011461	N1RT80
<i>Cochliobolus sativus</i> (strain ND90Pr/ATCC 201652) (Common root rot and spot blotch fungus) (<i>Bipolaris sorokiniana</i>)	COCSADRAFT_295770	M2TBE4
<i>Mixia osmundae</i> (strain CBS 9802/IAM 14324/JCM 22182/KY 12970)	Mo05571 E5Q_05571	G7E7S3
<i>Mycosphaerella pini</i> (strain NZE10/CBS 128990) (Red band needle blight fungus) (<i>Dothistroma septosporum</i>)	DOTSEDRAFT_69651	N1PXR0
<i>Grosmannia clavigera</i> (strain kw 1407/UAMH 11150) (Blue stain fungus) (<i>Graphiocladiella clavigera</i>)	CMQ_1113	F0XC64
<i>Fusarium oxysporum</i> FOSC 3-a	FOYG_03004	W9IUE5
<i>Fusarium oxysporum</i> FOSC 3-a	FOYG_16040	W9HNP0
<i>Fusarium oxysporum</i> FOSC 3-a	FOYG_17058	W9HB31
<i>Nectria haematococca</i> (strain 77-13-4/ATCC MYA-4622/FGSC 9596/MPVI) (<i>Fusarium solani</i> subsp. <i>pisi</i>)	NECHADRAFT_37686	C7YQL1
<i>Nectria haematococca</i> (strain 77-13-4/ATCC MYA-4622/FGSC 9596/MPVI) (<i>Fusarium solani</i> subsp. <i>pisi</i>)	NECHADRAFT_77262	C7ZJI0
<i>Tuber melanosporum</i> (strain Mel28) (Perigord black truffle)	GSTUM_00010376001	D5GLS0
<i>Ajellomyces dermatitidis</i> (strain SLH14081) (<i>Blastomyces dermatitidis</i>)	BDBG_07633	C5JYI9
<i>Chaetomium globosum</i> (strain ATCC 6205/CBS 148.51/DSM 1962/NBRC 6347/NRRL 1970) (Soil fungus)	CHGG_09885	Q2GQ69
<i>Candida tenuis</i> (strain ATCC 10573/BCRC 21748/CBS 615/JCM 9827/NBRC 10315/NRRL Y-1498/VKM Y-70) (Yeast)	CANTEDRAFT_108652	G3B9Z1
<i>Trichophyton rubrum</i> CBS 100081	H102_00622	A0A022VKY4
<i>Pyrenophora teres</i> f. <i>teres</i> (strain 0-1) (Barley net blotch fungus) (<i>Drechslera teres</i> f. <i>teres</i>)	PTT_09421	E3RLZ3
<i>Colletotrichum gloeosporioides</i> (strain Nara gc5) (Anthracnose fungus) (<i>Glomerella cingulata</i>)	CGGC5_4608	L2GB29
<i>Gibberella zeae</i> (Wheat head blight fungus) (<i>Fusarium graminearum</i>)	FG05_06918	A0A016PCS4
<i>Trichophyton soudanense</i> CBS 452.61	H105_00612	A0A022Y6A6
<i>Sclerotinia sclerotiorum</i> (strain ATCC 18683/1980/Ss-1) (White mold) (<i>Whetzelinia sclerotiorum</i>)	SS1G_07437	A7EQ37
<i>Fusarium oxysporum</i> f. sp. <i>pisi</i> HDV247	FOVG_14401	W9NWU8
<i>Fusarium oxysporum</i> f. sp. <i>pisi</i> HDV247	FOVG_02874	W9Q5V3
<i>Ustilago hordei</i> (strain Uh4875-4) (Barley covered smut fungus)	UHOR_03009	I2G1Z4
<i>Sporisorium reilianum</i> (strain SRZ2) (Maize head smut fungus)	sr12985	E6ZYF7
<i>Bipolaris zeicola</i> 26-R-13	COCCADRAFT_81154	W6YIP8
<i>Melampsora larici-populina</i> (strain 98AG31/pathotype 3-4-7) (Poplar leaf rust fungus)	MELLADRAFT_78490	F4RUZ8
<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> (strain 4287/CBS 123668/FGSC 9935/NRRL 34936) (<i>Fusarium</i> vascular wilt of tomato)	FOXG_01901	J9MG95
<i>Fusarium oxysporum</i> f. sp. <i>lycopersici</i> (strain 4287/CBS 123668/FGSC 9935/NRRL 34936) (<i>Fusarium</i> vascular wilt of tomato)	FOXG_11941	J9N9S4
<i>Bipolaris victoriae</i> FI3	COCVIDRAFT_39053	W7EMJ8
<i>Debaryomyces hansenii</i> (strain ATCC 36239/CBS 767/JCM 1990/NBRC 0083/IGC 2968) (Yeast) (<i>Torulaspora hansenii</i>)	DEHA2E04268g	Q6BQL4
<i>Clavispora lusitaniae</i> (strain ATCC 42720) (Yeast) (<i>Candida lusitaniae</i>)	CLUG_01505	C4XZX3
<i>Candida albicans</i> (strain WO-1) (Yeast)	CAWG_02023	C4YME4
<i>Trichophyton rabrum</i> MR850	H100_00625	A0A022U0Q2

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Candida dubliniensis</i> (strain CD36/ATCC MYA-646/CBS 7987/NCPF 3949/NRRL Y-17841) (Yeast)	CD36_32890	B9WMC7
<i>Starmerella bombicola</i>	AOX1	A0A024FB95
<i>Thielavia heterothallica</i> (strain ATCC 42464/BCRC 31852/DSM 1799) (<i>Myceliophthora thermophila</i>)	MYCTH_103590	G2QJL7
<i>Claviceps purpurea</i> (strain 20.1) (Ergot fungus) (<i>Sphacelia segetum</i>)	CPUR_07614	M1WFI4
<i>Aspergillus oryzae</i> (strain ATCC 42149/RIB 40) (Yellow koji mold)	AO090023000571	Q2UH61
<i>Dictyostelium discoideum</i> (Slime mold)	DDB_0184181 DDB_G0292042	Q54DT6
<i>Triticum urartu</i> (Red wild einkorn) (<i>Crithodium urartu</i>)	TRIUR3_22733	M7YME5
<i>Solanum tuberosum</i> (Potato)	PGSC0003DMG400017211	M1BG07
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OSJNBb0044B19.5 LOC_Os10g33540	Q8W5P8
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OJ1234_B11.20 Os02g0621800	Q6K9N5
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OSJNBa0001K12.5 LOC_Os10g33520	Q8W5P3
<i>Zea mays</i> (Maize)	ZEAMMB73_809149	C0P3J6
<i>Citrus clementina</i>	CICLE_v10011111mg	V4S9P4
<i>Citrus clementina</i>	CICLE_v10018992mg	V4U4C9
<i>Citrus clementina</i>	CICLE_v10004405mg	V4S9D3
<i>Citrus clementina</i>	CICLE_v10004403mg	V4RZZ6
<i>Morus notabilis</i>	L484_011703	W9RIK0
<i>Morus notabilis</i>	L484_005930	W9RET7
<i>Medicago truncatula</i> (Barrel medic) (<i>Medicago tribuloides</i>)	MTR_1g075650	G7I4U3
<i>Arabidopsis thaliana</i> (Mouse-ear cress)		Q8LDP0
<i>Medicago truncatula</i> (Barrel medic) (<i>Medicago tribuloides</i>)	MTR_4g081080	G7JF07
<i>Simmondsia chinensis</i> (Jojoba) (<i>Buxus chinensis</i>)		L7VVFV2
<i>Prunus persica</i> (Peach) (<i>Amygdalus persica</i>)	PRUPE_ppa018458mg	M5VXL1
<i>Aphanomyces astaci</i>	H257_07411	W4GI89
<i>Aphanomyces astaci</i>	H257_07412	W4GI44
<i>Aphanomyces astaci</i>	H257_07411	W4GKE3
<i>Aphanomyces astaci</i>	H257_07411	W4GK29
<i>Aphanomyces astaci</i>	H257_07411	W4GJ79
<i>Aphanomyces astaci</i>	H257_07411	W4GI38
<i>Phaeodactylum tricornutum</i> (strain CCAP 1055/1)	PHATRDRAFT_48204	B7G6C1
<i>Hordeum vulgare</i> var. <i>distichum</i> (Two-rowed barley)		F2E4R4
<i>Hordeum vulgare</i> var. <i>distichum</i> (Two-rowed barley)		F2DZG1
<i>Hordeum vulgare</i> var. <i>distichum</i> (Two-rowed barley)		M0YPG7
<i>Hordeum vulgare</i> var. <i>distichum</i> (Two-rowed barley)		M0YPG6
<i>Hordeum vulgare</i> var. <i>distichum</i> (Two-rowed barley)		F2CUY4
<i>Ricinus communis</i> (Castor bean)	RCOM_0867830	B9S1S3
<i>Brassica rapa</i> subsp. <i>pekinensis</i> (Chinese cabbage) (<i>Brassica pekinensis</i>)	BRA014947	M4DEM5
<i>Ricinus communis</i> (Castor bean)	RCOM_0258730	B9SV13
<i>Brassica rapa</i> subsp. <i>pekinensis</i> (Chinese cabbage) (<i>Brassica pekinensis</i>)	BRA001912	M4CCI2
<i>Brassica rapa</i> subsp. <i>pekinensis</i> (Chinese cabbage) (<i>Brassica pekinensis</i>)	BRA012548	M4D7T8
<i>Brassica rapa</i> subsp. <i>pekinensis</i> (Chinese cabbage) (<i>Brassica pekinensis</i>)	BRA024190	M4E5Y6
<i>Brassica rapa</i> subsp. <i>pekinensis</i> (Chinese cabbage) (<i>Brassica pekinensis</i>)	BRA015283	M4DFL0
<i>Ricinus communis</i> (Castor bean)	RCOM_1168730	B9SS54
<i>Zea mays</i> (Maize)		C4J691
<i>Oryza glaberrima</i> (African rice)		I1P2B7
<i>Zea mays</i> (Maize)		B6SXM3
<i>Zea mays</i> (Maize)		C0HFU4
<i>Aegilops tauschii</i> (Tausch's goatgrass) (<i>Aegilops squarrosa</i>)	F775_19577	R7W4J3
<i>Solanum habrochaites</i> (Wild tomato) (<i>Lycopersicon hirsutum</i>)		R9R6T0
<i>Physcomitrella patens</i> subsp. <i>patens</i> (Moss)	PHYPADRAFT_124285	A9S535

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Physcomitrella patens</i> subsp. <i>patens</i> (Moss)	PHYPADRAFT_113581	A9RG13
<i>Physcomitrella patens</i> subsp. <i>patens</i> (Moss)	PHYPADRAFT_182504	A9S9A5
<i>Solanum pennellii</i> (Tomato) (<i>Lycopersicon pennellii</i>)		R9R6Q1
<i>Vitis vinifera</i> (Grape)	VIT_02s0087g00630	F6HJ27
<i>Vitis vinifera</i> (Grape)	VIT_07s0005g03780	F6HZM3
<i>Vitis vinifera</i> (Grape)	VIT_05s0049g01400	F6H8T4
<i>Vitis vinifera</i> (Grape)	VITISV_019349	A5AH38
<i>Capsella rubella</i>	CARUB_v10013046mg	R0HIT3
<i>Capsella rubella</i>	CARUB_v10004212mg	R0GUX4
<i>Capsella rubella</i>	CARUB_v10004208mg	R0F3X6
<i>Capsella rubella</i>	CARUB_v10012453mg	R0ILD0
<i>Capsella rubella</i>	CARUB_v10004208mg	R0GUX1
<i>Eutrema salsugineum</i> (Saltwater cress) (<i>Sisymbrium salsugineum</i>)	EUTSA_v10024496mg	V4MD54
<i>Eutrema salsugineum</i> (Saltwater cress) (<i>Sisymbrium salsugineum</i>)	EUTSA_v10020141mg	V4NM59
<i>Eutrema salsugineum</i> (Saltwater cress) (<i>Sisymbrium salsugineum</i>)	EUTSA_v10024496mg	V4LUR9
<i>Eutrema salsugineum</i> (Saltwater cress) (<i>Sisymbrium salsugineum</i>)	EUTSA_v10024528mg	V4P767
<i>Eutrema salsugineum</i> (Saltwater cress) (<i>Sisymbrium salsugineum</i>)	EUTSA_v10006882mg	V4L2P6
<i>Selaginella moellendorffii</i> (Spikemoss)	SELMODRAFT_87684	D8R6Z6
<i>Selaginella moellendorffii</i> (Spikemoss)	SELMODRAFT_87621	D8R6Z5
<i>Selaginella moellendorffii</i> (Spikemoss)	SELMODRAFT_74601	D8QN81
<i>Selaginella moellendorffii</i> (Spikemoss)	SELMODRAFT_73531	D8QN82
<i>Sorghum bicolor</i> (<i>Sorghum</i>) (<i>Sorghum vulgare</i>)	Sb04g026390	C5XXS4
	SORBIDRAFT_04g026390	
<i>Sorghum bicolor</i> (<i>Sorghum</i>) (<i>Sorghum vulgare</i>)	Sb04g026370	C5XXS1
	SORBIDRAFT_04g026370	
<i>Sorghum bicolor</i> (<i>Sorghum</i>) (<i>Sorghum vulgare</i>)	Sb01g019470	C5WYH6
	SORBIDRAFT_01g019470	
<i>Sorghum bicolor</i> (<i>Sorghum</i>) (<i>Sorghum vulgare</i>)	Sb01g019480	C5WYH7
	SORBIDRAFT_01g019480	
<i>Sorghum bicolor</i> (<i>Sorghum</i>) (<i>Sorghum vulgare</i>)	Sb01g019460	C5WYH5
	SORBIDRAFT_01g019460	
<i>Solanum pimpinellifolium</i> (Currant tomato) (<i>Lycopersicon pimpinellifolium</i>)		R9R6J2
<i>Phaseolus vulgaris</i> (Kidney bean) (French bean)	PHAVU_007G124200g	V7BGM7
<i>Phaseolus vulgaris</i> (Kidney bean) (French bean)	PHAVU_011G136600g	V7AI35
<i>Phaseolus vulgaris</i> (Kidney bean) (French bean)	PHAVU_001G162800g	V7D063
<i>Solanum tuberosum</i> (Potato)	PGSC0003DMG400024294	M1C923
<i>Solanum tuberosum</i> (Potato)	PGSC0003DMG400018458	M1BKV4
<i>Solanum tuberosum</i> (Potato)	PGSC0003DMG400018458	M1BKV3
<i>Glycine max</i> (Soybean) (<i>Glycine hispida</i>)		K7LK61
<i>Glycine max</i> (Soybean) (<i>Glycine hispida</i>)		K7KXQ9
<i>Populus trichocarpa</i> (Western balsam poplar) (<i>Populus balsamifera</i> subsp. <i>trichocarpa</i>)	POPTR_0008s16920g	B9HKS3
<i>Picea sitchensis</i> (Sitka spruce) (<i>Pinus sitchensis</i>)		B8LQ84
<i>Populus trichocarpa</i> (Western balsam poplar) (<i>Populus balsamifera</i> subsp. <i>trichocarpa</i>)	POPTR_0004s24310g	U5GKQ5
<i>Populus trichocarpa</i> (Western balsam poplar) (<i>Populus balsamifera</i> subsp. <i>trichocarpa</i>)	POPTR_0010s07980g	B9HSG9
<i>Glycine max</i> (Soybean) (<i>Glycine hispida</i>)		I1N9S7
<i>Glycine max</i> (Soybean) (<i>Glycine hispida</i>)		I1LSK5
<i>Setaria italica</i> (Foxtail millet) (<i>Panicum italicum</i>)	Si034362m.g	K4A658
<i>Solanum lycopersicum</i> (Tomato) (<i>Lycopersicon esculentum</i>)	Solyc09g072610.2	K4CUT7
<i>Setaria italica</i> (Foxtail millet) (<i>Panicum italicum</i>)	Si016380m.g	K3YQ38
<i>Solanum lycopersicum</i> (Tomato) (<i>Lycopersicon esculentum</i>)		R9R6I9
<i>Solanum lycopersicum</i> (Tomato) (<i>Lycopersicon esculentum</i>)	Solyc09g090350.2	K4CW61
<i>Solanum lycopersicum</i> (Tomato) (<i>Lycopersicon esculentum</i>)	Solyc08g005630.2	K4CI54
<i>Solanum lycopersicum</i> (Tomato) (<i>Lycopersicon esculentum</i>)	Solyc08g075240.2	K4CMP1
<i>Setaria italica</i> (Foxtail millet) (<i>Panicum italicum</i>)	Si034359m.g	K4A655
<i>Setaria italica</i> (Foxtail millet) (<i>Panicum italicum</i>)	Si034354m.g	K4A650
<i>Mimulus guttatus</i> (Spotted monkey flower) (Yellow monkey flower)	MIMGU_mgv1a001896mg	A0A022PU07
<i>Mimulus guttatus</i> (Spotted monkey flower) (Yellow monkey flower)	MIMGU_mgv1a022390mg	A0A022RAV4

TABLE 5-continued

Alcohol oxidase enzymes capable of oxidizing long chain alcohols including fatty alcohols (EC 1.1.3.20).		
Organism	Gene names	Accession No.
<i>Mimulus guttatus</i> (Spotted monkey flower) (Yellow monkey flower)	MIMGU_mgv1a001868mg	A0A022S2E6
<i>Mimulus guttatus</i> (Spotted monkey flower) (Yellow monkey flower)	MIMGU_mgv1a001883mg	A0A022S275
<i>Mimulus guttatus</i> (Spotted monkey flower) (Yellow monkey flower)	MIMGU_mgv1a001761mg	A0A022QNF0
<i>Musa acuminata</i> subsp. <i>malaccensis</i> (Wild banana) (<i>Musa malaccensis</i>)		M0SNA8
<i>Musa acuminata</i> subsp. <i>malaccensis</i> (Wild banana) (<i>Musa malaccensis</i>)		M0RUT7
<i>Musa acuminata</i> subsp. <i>malaccensis</i> (Wild banana) (<i>Musa malaccensis</i>)		M0RUK3
<i>Saprolegnia diclina</i> VS20	SDRG_10901	T0RG89
<i>Brachypodium distachyon</i> (Purple false brome) (<i>Trachynia distachya</i>)	BRADI3G49085	I1IBP7
<i>Brachypodium distachyon</i> (Purple false brome) (<i>Trachynia distachya</i>)	BRADI3G28677	I1I4N2
<i>Brachypodium distachyon</i> (Purple false brome) (<i>Trachynia distachya</i>)	BRADI3G28657	I1I4N0
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	OsI_34012	B8BHG0
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	OsI_08118	B8AFT8
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	OsI_34008	A2Z8H1
<i>Oryza sativa</i> subsp. <i>indica</i> (Rice)	OsI_34014	B8BHG1
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	LOC_Os10g33460	Q7XDG3
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	Os10g0474800	Q0IX12
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	Os10g0474966	C7J7R1
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OSJNBa0001K12.13	Q8W5N7
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OsJ_31873	B9G683
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OsJ_31875	B9G684
<i>Oryza sativa</i> subsp. <i>japonica</i> (Rice)	OSJNBa0001K12.3	Q8W5P5
<i>Arabidopsis lyrata</i> subsp. <i>lyrata</i> (Lyre-leaved rock-cress)	ARALYDRAFT_470376	D7KDA3
<i>Arabidopsis lyrata</i> subsp. <i>lyrata</i> (Lyre-leaved rock-cress)	ARALYDRAFT_479855	D7L3B6
<i>Arabidopsis lyrata</i> subsp. <i>lyrata</i> (Lyre-leaved rock-cress)	ARALYDRAFT_491906	D7MDA9
<i>Arabidopsis lyrata</i> subsp. <i>lyrata</i> (Lyre-leaved rock-cress)	ARALYDRAFT_914728	D7MGS9

Acetyl Transferase

The present disclosure describes enzymes that convert alcohols to fatty acetates.

In some embodiments, an acetyl transferase is used to catalyze the conversion of a fatty alcohol to a fatty acetate. An acetyl transferase is an enzyme that has the ability to produce an acetate ester by transferring the acetyl group from acetyl-CoA to an alcohol. In some embodiments, the acetyl transferase may have an EC number of 2.3.1.84.

The acetyl transferase, or the nucleic acid sequence that encodes it, can be isolated from various organisms, including but not limited to, organisms of the species *Saccharomyces cerevisiae*, *Danaus plexippus*, *Heliotis virescens*, *Bombyx mori*, *Agrotis ipsilon*, *Agrotis segetum*, *Euonymus alatus*. In exemplary embodiments, the acetyl transferase comprises a sequence selected from GenBank Accession Nos. AY242066, AY242065, AY242064, AY242063, AY242062, EHJ65205, ACX53812, NP_001182381, EHJ65977, EHJ68573, KJ579226, GU594061. Additional exemplary acetyl transferase peptides may be found in US2010/0199548, which is herein incorporated by reference.

Fatty Acyl-ACP Thioesterase

Acyl-ACP thioesterase releases free fatty acids from Acyl-ACPs, synthesized from de novo fatty acid biosynthesis. The reaction terminates fatty acid biosynthesis. In plants, fatty acid biosynthesis occurs in the plastid and thus requires plastid-localized acyl-ACP thioesterases. The main products

of acyl-ACP thioesterase are oleate (C18:0) and to a lesser extent palmitate (C16:0) in the vegetative tissues of all plants. The released free fatty acids are re-esterified to coenzyme A in the plastid envelope and exported out of plastid.

There are two isoforms of acyl-ACP thioesterase, FatA and FatB. Substrate specificity of these isoforms determines the chain length and level of saturated fatty acids in plants. The highest activity of FatA is with C18:1-ACP. FatA has very low activities towards other acyl-ACPs when compared with C18:1-ACP. FatB has highest activity with C16:0-ACP. It also has significant high activity with C18:1-ACP, followed by C18:0-ACP and C16:1-ACP. Kinetics studies of FatA and FatB indicate that their substrate specificities with different acyl-ACPs came from the Kcat values, rather than from Km. Km values of the two isoforms with different substrates are similar, in the micromolar order. Domain swapping of FatA and FatB indicates the N-terminus of the isoforms determines their substrate specificities (Salas J J and Ohlrogge J B (2002) Characterization of substrate specificity of plant FatA and FatB acyl-ACP thioesterases. Arch Biochem Biophys 403(1): 25-34). For those plants which predominantly accumulate medium-chain length saturated fatty acids in seeds, they evolved with specialized FatB and/or FatA thioesterases (Voelker T and Kinney A J (2001) Variations in the biosynthesis of seed-storage lipids. Annu Rev Plant Physiol Plant Mol Biol 52: 335-361). For example, laurate (12:0) is the predominant seed oil in

coconut. Correspondingly, the medium-chain specific acyl-ACP thioesterase activity was detected in coconut seeds.

In one embodiment, one or more fatty acyl-ACP thioesterases are selected from the group consisting of Q41635, Q39473, P05521.2, AEM72519, AEM72520, AEM72521, AEM72523, AAC49784, CAB60830, EER87824, EER96252, ABN54268, AAO77182, CAH09236, ACL08376, and homologs thereof.

Expression of Toxic Proteins or Polypeptides

The present disclosure describes a toxic protein, peptide, or small molecule that can be encoded by a recombinant microorganism. In some embodiments, the toxic protein, peptide, or small molecule is biosynthetically produced along with an insect pheromone.

In some embodiments, the recombinant microorganism expresses one or more nucleic acid molecules encoding a protein or polypeptide which is toxic to an insect. In some embodiments, the toxic protein or polypeptide is from an entomopathogenic organism. In some embodiments, the entomopathogenic organism is selected from *Bacillus thuringiensis*, *Pseudomonas aeruginosa*, and *Serratia marcescens*. In a particular embodiment, the nucleic acid molecule encodes a *Bacillus thuringiensis* toxin.

In some embodiments, a recombinant microorganism is engineered to express a metabolic pathway which, when expressed, produces a small molecule that is toxic to an insect.

In exemplary embodiments, an insect pheromone produced by a recombinant microorganism described herein may be used to attract a pest insect, and subsequently, the pest insect is eradicated with a toxic substance, such as a toxic protein, peptide, or small molecule, which has been co-produced by a recombinant microorganism described herein.

Biosynthesis of Pheromones Using a Recombinant Microorganism

As discussed above, in a first aspect, the present disclosure relates to a recombinant microorganism capable of producing a mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated C₆-C₂₄ fatty acyl-CoA. An illustrative embodiment of the first aspect is shown in FIG. 1. The blue lines designate biochemical pathways used to produce a saturated acyl-CoA, which acts as a substrate for unsaturated fatty-acyl CoA conversion. The substrate to unsaturated fatty acyl-CoA conversion can be performed by endogenous or exogenous enzymes in a host. Green lines indicate conversions catalyzed by an exogenous nucleic acid molecule encoding for an enzyme. Accordingly, in some embodiments, the conversion of a saturated fatty acyl-CoA to a mono- or poly-unsaturated fatty acyl-CoA is catalyzed by at least one desaturase, which is encoded by an exogenous nucleic acid molecule. In further embodiments, the conversion of the mono- or poly-unsaturated fatty acyl-CoA to a mono- or poly-unsaturated fatty alcohol is catalyzed by at least one reductase, which is encoded by an exogenous nucleic acid molecule. The dashed grey lines indicate downstream steps for the synthesis of pheromones, fragrances, flavors, and polymer intermediates, such as using an alcohol oxidase or oxidant to produce a mono- or poly-unsaturated fatty aldehyde, and an acetyl transferase or a chemical such as acetylchloride to produce a mono- or poly-unsaturated fatty acetate. The red crosses indicate deleted or down regulated pathways native to the host, which increase flux towards the engineered pathway.

Accordingly, in one embodiment, the recombinant microorganism expresses: (a) at least one exogenous nucleic acid

molecule encoding a fatty acyl desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-CoA to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA; and (b) at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA from (a) into the corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. In some embodiments, the saturated C₆-C₂₄ fatty acyl-CoA can be produced using endogenous enzymes in the host microorganism. In other embodiments, the saturated C₆-C₂₄ fatty acyl-CoA can be produced using one or more exogenous enzymes in the host microorganism.

As described above, a fatty acyl desaturase catalyzes the desaturation of the hydrocarbon chain on, e.g., a saturated fatty acyl-CoA molecule to generate a corresponding unsaturated fatty acyl CoA molecule. In some embodiments, an exogenous fatty acyl desaturase can be selected and expressed in a recombinant microorganism to catalyze the formation of at least one double bond in fatty acyl-CoA molecule having from 6 to 24 carbons in the hydrocarbon chain. Accordingly, in some embodiments, the fatty-acyl desaturase is a desaturase capable of utilizing a fatty acyl-CoA as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms.

An exogenous fatty acyl desaturase described herein can be selected to catalyze the desaturation at a desired position on the hydrocarbon chain. Accordingly, in some embodiments, the fatty-acyl desaturase is capable of generating a double bond at position C5, C6, C7, C8, C9, C10, C11, C12, or C13, in the fatty acid or its derivatives, such as, for example, fatty acid CoA esters.

One or more than one fatty acyl-CoA desaturase can be expressed in the host to catalyze desaturation at multiple positions on the hydrocarbon chain. In some embodiments, the fatty acyl-CoA desaturase is heterologous to the host microorganism. Accordingly, various embodiments provide for recombinant microorganism comprised of at least one exogenous nucleic acid molecule, which encodes a fatty acyl desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-CoA to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA.

In one exemplary embodiment, the fatty-acyl desaturase is a Z11 desaturase. The Z11 fatty-acyl desaturase catalyze double bond formation between the 11th and 12th carbons in the substrate relative to the carbonyl group. In various embodiments described herein, the Z11 desaturase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Agrotis segetum*, *Amyelois transitella*, *Argyrotaenia velutiana*, *Choristoneura rosaceana*, *Lampronia capitella*, *Trichoplusia ni*, *Helicoverpa zea*, or *Thalassiosira pseudonana*. Further Z11-desaturases, or the nucleic acid sequences encoding them, can be isolated from *Bombyx mori*, *Manduca sexta*, *Diatraea grandiosella*, *Earias insulana*, *Earias vittella*, *Plutella xylostella*, *Bombyx mori* or *Diaphania nitidalis*. In exemplary embodiments, the Z11 desaturase comprises a sequence selected from GenBank Accession Nos. JX679209, JX964774, AF416738, AF545481, EU152335, AAD03775, AAF81787, and AY493438. In some embodiments, a nucleic acid sequence encoding a Z11 desaturase from organisms of the species *Agrotis segetum*, *Amyelois transitella*, *Argyrotaenia velutiana*, *Choristoneura rosaceana*, *Lampronia capitella*, *Trichoplusia ni*, *Helicoverpa zea*, or *Thalassiosira pseudonana* is codon optimized. In some embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID

NOs: 9, 18, 24 and 26 from *Trichoplusia ni*. In other embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 10 and 16 from *Agrotis segetum*. In some embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 11 and 23 from *Thalassiosira pseudonana*. In certain embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 12, 17 and 30 from *Amyelois transitella*. In further embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 13, 19, 25, 27 and 31 from *Helicoverpa zea*. In some embodiments, the Z11 desaturase comprises a chimeric polypeptide. In some embodiments, a complete or partial Z11 desaturase is fused to another polypeptide. In certain embodiments, the N-terminal native leader sequence of a Z11 desaturase is replaced by an oleosin leader sequence from another species. In certain embodiments, the Z11 desaturase comprises a sequence selected from SEQ ID NOs: 15, 28 and 29.

In certain embodiments, the Z11 desaturase catalyzes the conversion of a fatty acyl-CoA into a mono- or polyunsaturated product selected from Z1-13:Acyl-CoA, E11-13:Acyl-CoA, (Z,Z)-7,11-13:Acyl-CoA, Z11-14:Acyl-CoA, E11-14:Acyl-CoA, (E,E)-9,11-14:Acyl-CoA, (E,Z)-9,11-14:Acyl-CoA, (Z,E)-9,11-14:Acyl-CoA, (Z,Z)-9,11-14:Acyl-CoA, (E,Z)-9,11-15:Acyl-CoA, (Z,Z)-9,11-15:Acyl-CoA, Z11-16:Acyl-CoA, E11-16:Acyl-CoA, (E,Z)-6,11-16:Acyl-CoA, (E,Z)-7,11-16:Acyl-CoA, (E,Z)-8,11-16:Acyl-CoA, (E,E)-9,11-16:Acyl-CoA, (E,Z)-9,11-16:Acyl-CoA, (Z,E)-9,11-16:Acyl-CoA, (Z,Z)-9,11-16:Acyl-CoA, (E,E)-11,13-16:Acyl-CoA, (E,Z)-11,13-16:Acyl-CoA, (Z,E)-11,13-16:Acyl-CoA, (Z,Z)-11,13-16:Acyl-CoA, (Z,E)-11,14-16:Acyl-CoA, (E,E,Z)-4,6,11-16:Acyl-CoA, (Z,Z,E)-7,11,13-16:Acyl-CoA, (E,E,Z,Z)-4,6,11,13-16:Acyl-CoA, Z1-17:Acyl-CoA, (Z,Z)-8,11-17:Acyl-CoA, Z1-18:Acyl-CoA, E11-18:Acyl-CoA, (Z,Z)-11,13-18:Acyl-CoA, (E,E)-11,14-18:Acyl-CoA, or combinations thereof.

In another exemplary embodiment, the fatty-acyl desaturase is a Z9 desaturase. The Z9 fatty-acyl desaturase catalyzes double bond formation between the 9th and 10th carbons in the substrate relative to the carbonyl group. In various embodiments described herein, the Z9 desaturase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Ostrinia furnacalis*, *Ostrinia nobilalis*, *Choristoneura rosaceana*, *Lampronia capitella*, *Helicoverpa assulta*, or *Helicoverpa zea*. In exemplary embodiments, the Z9 desaturase comprises a sequence selected from GenBank Accession Nos. AY057862, AF243047, AF518017, EU152332, AF482906, and AAF81788. In some embodiments, a nucleic acid sequence encoding a Z9 desaturase is codon optimized. In some embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 20 from *Ostrinia furnacalis*. In other embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 21 from *Lampronia capitella*. In some embodiments, the Z9 desaturase comprises a sequence set forth in SEQ ID NO: 22 from *Helicoverpa zea*.

In certain embodiments, the Z9 desaturase catalyzes the conversion of a fatty acyl-CoA into a monounsaturated or polyunsaturated product selected from Z9-11:Acyl-CoA, Z9-12:Acyl-CoA, E9-12:Acyl-CoA, (E,E)-7,9-12:Acyl-CoA, (E,Z)-7,9-12:Acyl-CoA, (Z,E)-7,9-12:Acyl-CoA, (Z,Z)-7,9-12:Acyl-CoA, Z9-13:Acyl-CoA, E9-13:Acyl-CoA, (E,Z)-5,9-13:Acyl-CoA, (Z,E)-5,9-13:Acyl-CoA, (Z,Z)-5,9-13:Acyl-CoA, Z9-14:Acyl-CoA, E9-14:Acyl-CoA, (E,Z)-4,9-14:Acyl-CoA, (E,E)-9,11-14:Acyl-CoA, (E,Z)-9,11-14:Acyl-CoA, (Z,E)-9,11-14:Acyl-CoA, (Z,Z)-9,11-14:Acyl-CoA, (E,E)-9,12-14:Acyl-CoA, (Z,E)-9,12-

14:Acyl-CoA, (Z,Z)-9,12-14:Acyl-CoA, Z9-15:Acyl-CoA, E9-15:Acyl-CoA, (Z,Z)-6,9-15:Acyl-CoA, Z9-16:Acyl-CoA, E9-16:Acyl-CoA, (E,E)-9,11-16:Acyl-CoA, (E,Z)-9,11-16:Acyl-CoA, (Z,E)-9,11-16:Acyl-CoA, (Z,Z)-9,11-16:Acyl-CoA, Z9-17:Acyl-CoA, E9-18:Acyl-CoA, Z9-18:Acyl-CoA, (E,E)-5,9-18:Acyl-CoA, (E,E)-9,12-18:Acyl-CoA, (Z,Z)-9,12-18:Acyl-CoA, (Z,Z,Z)-3,6,9-18:Acyl-CoA, (E,E,E)-9,12,15-18:Acyl-CoA, (Z,Z,Z)-9,12,15-18:Acyl-CoA, or combinations thereof.

Desaturation of a saturated C₆-C₂₄ fatty acyl-CoA can proceed through a plurality of reactions to produce a polyunsaturated C₆-C₂₄ fatty acyl-CoA. In some embodiments, the recombinant microorganism may express a bifunctional desaturase capable of catalyzing the formation at least two double bonds. In some embodiments, the recombinant microorganism may express more than one exogenous nucleic acid molecule encoding more than one fatty-acyl desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA. For example, the recombinant microorganism may express an exogenous nucleic acid molecule encoding a Z11 desaturase and another exogenous nucleic acid molecule encoding a Z9 desaturase. Thus, the resultant polyunsaturated fatty acyl-CoA would have a double bond between the 9th and 10th carbon and another double bond between the 11th and 12th carbon.

In some embodiments, the recombinant microorganism may express a fatty-acyl conjugase that acts independently or together with a fatty-acyl desaturase to catalyze the conversion of a saturated or monounsaturated fatty acyl-CoA to a conjugated polyunsaturated fatty acyl-CoA.

In one embodiment, the disclosure provides a recombinant microorganism capable of producing a polyunsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA, wherein the recombinant microorganism expresses: (a) at least one exogenous nucleic acid molecule encoding a fatty acyl conjugase that catalyzes the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA; and (b) at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the polyunsaturated C₆-C₂₄ fatty acyl-CoA from (a) into the corresponding polyunsaturated C₆-C₂₄ fatty alcohol.

In another embodiment, the recombinant microorganism expresses at least two exogenous nucleic acid molecules encoding fatty-acyl conjugases that catalyze the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA.

In a further embodiment, the disclosure provides a recombinant microorganism capable of producing a polyunsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA, wherein the recombinant microorganism expresses: (a) at least one exogenous nucleic acid molecule encoding a fatty-acyl desaturase and at least one exogenous nucleic acid molecule encoding a fatty acyl conjugase that catalyze the conversion of a saturated or monounsaturated C₆-C₂₄ fatty acyl-CoA to a corresponding polyunsaturated C₆-C₂₄ fatty acyl-CoA; and (b) at least one exogenous nucleic acid molecule encoding a fatty alcohol forming fatty-acyl reductase that catalyzes the conversion of the polyunsaturated C₆-C₂₄ fatty acyl-CoA from (a) into the corresponding polyunsaturated C₆-C₂₄ fatty alcohol.

In another embodiment, the recombinant microorganism expresses at least two exogenous nucleic acid molecules

encoding fatty-acyl desaturases and at least two exogenous nucleic acid molecules encoding fatty-acyl conjugases that catalyze the conversion of a saturated or monounsaturated C_6 - C_{24} fatty acyl-CoA to a corresponding polyunsaturated C_6 - C_{24} fatty acyl-CoA.

In yet a further embodiment, the fatty-acyl conjugase is a conjugase capable of utilizing a fatty acyl-CoA as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms.

In certain embodiments, the conjugase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Cydia pomonella*, *Cydia nigricana*, *Lobesia botrana*, *Myelois cribrrella*, *Plodia interpunctella*, *Dendrolimus punctatus*, *Lampronia capitella*, *Spodoptera litura*, *Amyelois transitella*, *Manduca sexta*, *Bombyx mori*, *Calendula officinalis*, *Trichosanthes kirilowii*, *Punica granatum*, *Momordica charantia*, *Impatiens balsamina*, and *Epiphyas postvittana*. In exemplary embodiments, the conjugase comprises a sequence selected from GenBank Accession No. or Uniprot database: A0A059TBF5, A0A0M3L9E8, A0A0M3L9S4, A0A0M3LAH8, A0A0M3LAS8, A0A0M3LAH8, B6CBS4, XP_013183656.1, XP_004923568.2, ALA65425.1, NP_001296494.1, NP_001274330.1, Q4A181, Q75PL7, Q9FPP8, AY178444, AY178446, AF182521, AF182520, Q95UJ3.

As described above, a fatty acyl reductase catalyzes the reduction of a carbonyl group, e.g., on an unsaturated fatty acyl-CoA molecule to generate a corresponding unsaturated fatty acid molecule. In some embodiments, the fatty alcohol forming fatty acyl CoA reductase is heterologous to the microorganism. Accordingly, various embodiments provide for recombinant microorganism comprised of at least one exogenous nucleic acid molecule, which encodes a fatty alcohol forming fatty acyl reductase that catalyzes the reduction of a carbonyl group on an unsaturated fatty acyl-CoA molecule to generate a corresponding unsaturated fatty acid molecule.

In some embodiments, the fatty acyl reductase is from an organism of the species *Agrotis segetum*, *Spodoptera littoralis*, or *Helicoverpa amigera*. In some embodiments, a nucleic acid sequence encoding a fatty-acyl reductase is codon optimized. In some embodiments, the fatty acyl reductase comprises a sequence set forth in SEQ ID NO: 1 from *Agrotis segetum*. In other embodiments, the fatty acyl reductase comprises a sequence set forth in SEQ ID NO: 2 from *Spodoptera littoralis*. In some embodiments, the fatty acyl reductase comprises a sequence selected from SEQ ID NOs: 3 and 32 from *Helicoverpa armigera*.

In exemplary embodiments, the fatty-acyl reductase catalyzes the conversion of a mono- or poly-unsaturated fatty acyl-CoA into a fatty alcohol product selected from (Z)-3-hexenol, (Z)-3-nonenol, (Z)-5-decenol, (E)-5-decenol, (Z)-7-dodecenol, (E)-8-dodecenol, (Z)-8-dodecenol, (Z)-9-dodecenol, (Z)-9-tetradecenol, (Z)-9-hexadecenol, (Z)-11-tetradecenol, (Z)-7-hexadecenol, (Z)-11-hexadecenol, (E)-11-tetradecenol, or (Z,Z)-11,13-hexadecadienol, (11Z,13E)-hexadecadienol, (E,E)-8,10-dodecadienol, (E,Z)-7,9-dodecadienol, (Z)-13-octadecenol, or combinations thereof.

In some embodiments, a recombinant microorganism described herein can include a plurality of fatty acyl reductases. Accordingly, in such embodiments, the recombinant microorganism expresses at least two exogenous nucleic acid molecules, which encode fatty-acyl reductases that catalyze the conversion of the mono- or poly-unsaturated C_6 - C_{24} fatty acyl-CoA into the corresponding mono- or poly-unsaturated C_6 - C_{24} fatty alcohol.

As discussed above, in a second aspect, the application relates to a recombinant microorganism capable of producing an unsaturated C_6 - C_{24} fatty alcohol from an endogenous or exogenous source of C_6 - C_{24} fatty acid. An illustrative embodiment of the second aspect is shown in FIG. 2. The blue lines designate biochemical pathways endogenous to the host, e.g., pathways for converting an n-alkane, fatty alcohol, or fatty aldehyde to a fatty acid, or the conversion of a fatty acid to fatty-acyl-CoA, acetyl-CoA, or dicarboxylic acid. The substrate to unsaturated fatty acid conversion can be performed by endogenous or exogenous enzymes in a host. Yellow lines indicate conversions catalyzed by an exogenous nucleic acid molecule encoding for an enzyme. Accordingly, in some embodiments, the conversion of a saturated fatty acid to a saturated fatty acyl-ACP can be catalyzed by at least one saturated fatty acyl-ACP synthetase, wherein the fatty acyl-ACP synthetase is encoded by an exogenous nucleic acid molecule. In further embodiments, the conversion of the saturated fatty acyl-ACP to a mono- or poly-unsaturated fatty acyl-ACP can be catalyzed by at least one fatty acyl-ACP desaturase, wherein the fatty acyl-ACP desaturase is encoded by an exogenous nucleic acid molecule. In still further embodiments, the mono- or poly-unsaturated fatty acyl-ACP can be elongated by at least 2 carbons relative using a fatty acid synthase complex and a carbon source, e.g., malonyl-ACP. In one such embodiment, the conversion of the mono- or poly-unsaturated fatty acyl-ACP to a corresponding two carbon elongated mono- or poly-unsaturated fatty acyl-ACP can be catalyzed by at least one fatty acid synthase complex, wherein the fatty acid synthase complex is encoded by one or more exogenous nucleic acid molecules. In yet further embodiments, the conversion of the elongated mono- or poly-unsaturated fatty acyl-ACP to a mono- or poly-unsaturated fatty aldehyde can be catalyzed by a fatty aldehyde forming fatty acyl reductase, wherein the fatty aldehyde forming fatty acyl reductase is encoded by an exogenous nucleic acid molecule. In some embodiments, the mono- or poly-unsaturated fatty aldehyde can be converted to a corresponding mono- or poly-unsaturated fatty alcohol, wherein the substrate to product conversion is catalyzed by a dehydrogenase, wherein the dehydrogenase is encoded by an endogenous or exogenous nucleic acid molecule. The dashed lines indicate downstream steps of the disclosure, such as utilizing an acetyl transferase or metathesis, or subsequent chemical transformations to produce functionalized pheromones. The red crosses indicate deleted or down regulated pathways native to the host, which increase flux towards the engineered pathway.

In one embodiment, the recombinant microorganism expresses (a): at least one exogenous nucleic acid molecule encoding an acyl-ACP synthetase that catalyzes the conversion of a C_6 - C_{24} fatty acid to a corresponding saturated C_6 - C_{24} fatty acyl-ACP; (b) at least one exogenous nucleic acid molecule encoding a fatty-acyl-ACP desaturase that catalyzes the conversion of a saturated C_6 - C_{24} fatty acyl-ACP to a corresponding mono- or poly-unsaturated C_6 - C_{24} fatty acyl-ACP; (c) one or more endogenous or exogenous nucleic acid molecules encoding a fatty acid synthase complex that catalyzes the conversion of the mono- or poly-unsaturated C_6 - C_{24} fatty acyl-ACP from (b) to a corresponding mono- or poly-unsaturated C_6 - C_{24} fatty acyl-ACP with a two carbon elongation relative to the product of (b); (d): at least one exogenous nucleic acid molecule encoding a fatty aldehyde forming fatty-acyl reductase that catalyzes the conversion of the mono- or poly-unsaturated C_6 - C_{24} fatty acyl-ACP from (c) into a corresponding mono- or poly-

unsaturated C₆-C₂₄ fatty aldehyde; and (e) at least one endogenous or exogenous nucleic acid molecule encoding a dehydrogenase that catalyzes the conversion of the mono- or poly-unsaturated C₆-C₂₄ fatty aldehyde C₆-C₂₄ from (d) into a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol. In some embodiments, the C₆-C₂₄ fatty acid can be produced using endogenous enzymes in the host microorganism. In other embodiments, the saturated C₆-C₂₄ fatty acid can be produced by one or more exogenous enzymes in the host microorganism.

In some embodiments, the recombinant microorganism disclosed herein includes an acyl-ACP synthetase to catalyze the conversion of a C₆-C₂₄ fatty acid to a corresponding saturated C₆-C₂₄ fatty acyl-ACP. In some embodiments the acyl-ACP synthetase is a synthetase capable of utilizing a fatty acid as a substrate that has a chain length of 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, or 24 carbon atoms. In exemplary embodiments, the recombinant microorganism can include a heterologous the acyl-ACP synthetase from an organism of the species *Vibrio harveyi*, *Rhodotorula glutinis*, or *Yarrowia lipolytica*.

In some embodiments, the recombinant microorganism includes a fatty acyl-ACP desaturase. In some embodiments, the fatty acyl-ACP desaturase is a soluble desaturase. In other embodiments, the fatty-acyl-ACP desaturase is from an organism of the species *Pelargonium hortorum*, *Asclepias syriaca*, or *Uncaria tomentosa*.

In some embodiments, the recombinant microorganism includes a fatty acid synthase complex. In some embodiments, the one or more nucleic acid molecules encoding the fatty acid synthase complex are endogenous nucleic acid molecules. In other embodiments, the one or more nucleic acid molecules encoding a fatty acid synthase complex are exogenous nucleic acid molecules.

In some embodiments, the recombinant microorganism disclosed herein includes a fatty aldehyde forming fatty-acyl reductase which catalyzes the conversion of a C₆-C₂₄ fatty acyl-ACP to the corresponding C₆-C₂₄ fatty aldehyde. In exemplary embodiments, the fatty aldehyde forming fatty-acyl reductase is from an organism of the species *Pelargonium hortorum*, *Asclepias syriaca*, and *Uncaria tomentosa*. In some embodiments, the recombinant microorganism includes a dehydrogenase to convert the unsaturated fatty aldehyde to a corresponding unsaturated fatty alcohol. In some embodiments, the nucleic acid molecule encoding the dehydrogenase is endogenous to the recombinant microorganism. In other embodiments, the nucleic acid molecule encoding a dehydrogenase is exogenous to the recombinant microorganism. In exemplary embodiments, the endogenous or exogenous nucleic acid molecule encoding a dehydrogenase is isolated from organisms of the species *Saccharomyces cerevisiae*, *Escherichia coli*, *Yarrowia lipolytica*, or *Candida tropicalis*.

As discussed above, in a third aspect, the application relates to a recombinant microorganism capable of producing an unsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of C₆-C₂₄ fatty acid. An illustrative embodiment of the second aspect is shown in FIG. 3. The blue lines designate biochemical pathways endogenous to the host, e.g., pathways for converting an n-alkane, fatty alcohol, or fatty aldehyde to a fatty acid, or the conversion of a fatty acid to fatty-acyl-CoA, acetyl-CoA, or dicarboxylic acid. The substrate to unsaturated fatty acid conversion can be performed by endogenous or exogenous enzymes in a host. Yellow lines indicate conversions catalyzed by an exogenous nucleic acid molecule encoding for an enzyme. Accordingly, in some embodiments, the conversion of a

saturated fatty acid to a saturated fatty acyl-ACP can be catalyzed by at least one saturated fatty acyl-ACP synthetase, wherein the fatty acyl-ACP synthetase is encoded by an exogenous nucleic acid molecule. The non-native saturated fatty acyl-ACP thioesters create a substrate suitable for desaturation and distinct from CoA-thioesters used for beta-oxidation or fatty acid elongation. In further embodiments, the conversion of the saturated fatty acyl-ACP to a mono- or poly-unsaturated fatty acyl-ACP can be catalyzed by at least one fatty acyl-ACP desaturase, wherein the fatty acyl-ACP desaturase is encoded by an exogenous nucleic acid molecule. In still further embodiments, the mono- or poly-unsaturated fatty acyl-ACP can be converted to a corresponding mono- or poly-unsaturated fatty acid by a fatty-acyl-ACP thioesterase. In a particular embodiment, soluble fatty acyl-ACP thioesterases can be used to release free fatty acids for reactivation to a CoA thioester. Fatty acyl-ACP thioesterases including Q41635, Q39473, P05521.2, AEM72519, AEM72520, AEM72521, AEM72523, AAC49784, CAB60830, EER87824, EER96252, ABN54268, AAO77182, CAH09236, ACL08376, and homologs thereof may be used. In an additional embodiment, the mono- or poly-unsaturated fatty acyl-CoA can be elongated by at least 2 carbons relative using an elongase and a carbon source, e.g., malonyl-ACP. In yet further embodiments, the conversion of the elongated mono- or poly-unsaturated fatty acyl-CoA to a mono- or poly-unsaturated fatty alcohol can be catalyzed by a fatty alcohol forming fatty acyl reductase, wherein the fatty alcohol forming fatty acyl reductase is encoded by an exogenous nucleic acid molecule. The dashed lines indicate downstream steps of the disclosure, such as utilizing an acetyl transferase or metathesis, or subsequent chemical transformations to produce functionalized pheromones. The red crosses indicate deleted or down regulated pathways native to the host, which increase flux towards the engineered pathway.

The fatty alcohols produced as taught herein can be further converted to produce downstream products such as insect pheromones, fragrances, flavors, and polymer intermediates, which utilize aldehydes or acetate functional groups. Thus, in some embodiments, the recombinant microorganism further comprises at least one endogenous or exogenous nucleic acid molecule encoding an alcohol oxidase or an alcohol dehydrogenase, wherein the alcohol oxidase or alcohol dehydrogenase is capable of catalyzing the conversion of a C₆-C₂₄ fatty alcohol into a corresponding C₆-C₂₄ fatty aldehyde. In other embodiments, the recombinant microorganism can further comprise at least one endogenous or exogenous nucleic acid molecule encoding an acetyl transferase capable of catalyzing the conversion of a C₆-C₂₄ fatty alcohol into a corresponding C₆-C₂₄ fatty acetate. In certain embodiments, the acetyl transferase, or the nucleic acid sequence that encodes it, can be isolated from organisms of the species *Saccharomyces cerevisiae*, *Danaus plexippus*, *Heliotis virescens*, *Bombyx mori*, *Agrotis ipsilon*, *Agrotis segetum*, *Euonymus alatus*. In exemplary embodiments, the acetyl transferase comprises a sequence selected from GenBank Accession Nos. AY242066, AY242065, AY242064, AY242063, AY242062, EHJ65205, ACX53812, NP_001182381, EHJ65977, EHJ68573, KJ579226, GU594061.

Recombinant Microorganism

The disclosure provides microorganisms that can be engineered to express various exogenous enzymes.

In various embodiments described herein, the recombinant microorganism is a eukaryotic microorganism. In some

embodiments, the eukaryotic microorganism is a yeast. In exemplary embodiments, the yeast is a member of a genus selected from the group consisting of *Yarrowia*, *Candida*, *Saccharomyces*, *Pichia*, *Hansenula*, *Kluyveromyces*, *Issatchenkia*, *Zygosaccharomyces*, *Debaryomyces*, *Schizosaccharomyces*, *Pachysolen*, *Cryptococcus*, *Trichosporon*, *Rhodotorula*, and *Myxozyma*.

The present inventors have discovered that oleaginous yeast, such as *Candida* and *Yarrowia*, have a surprisingly high tolerance to the C₆-C₂₄ fatty alcohol substrates and products. Accordingly, in one such exemplary embodiment, the recombinant microorganism of the invention is an oleaginous yeast. In further embodiments, the oleaginous yeast is a member of a genus selected from the group consisting of *Yarrowia*, *Candida*, *Rhodotorula*, *Rhodospiridium*, *Cryptococcus*, *Trichosporon*, and *Lipomyces*. In even further embodiments, the oleaginous yeast is a member of a species selected from *Yarrowia lipolytica*, *Candida tropicalis*, *Rhodospiridium toruloides*, *Lipomyces starkey*, *L. lipoferus*, *C. revkaufi*, *C. pulcherrima*, *C. utilis*, *Rhodotorula minuta*, *Trichosporon pullans*, *T. cutaneum*, *Cryptococcus curvatus*, *R. glutinis*, and *R. graminis*.

In some embodiments, the recombinant microorganism is a prokaryotic microorganism. In exemplary embodiments, the prokaryotic microorganism is a member of a genus selected from the group consisting of *Escherichia*, *Clostridium*, *Zymomonas*, *Salmonella*, *Rhodococcus*, *Pseudomonas*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Alcaligenes*, *Klebsiella*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, and *Brevibacterium*.

In some embodiments, the recombinant microorganism is used to produce a mono- or poly-unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate disclosed herein.

Accordingly, in another aspect, the present inventions provide a method of producing a mono- or poly-unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate using a recombinant microorganism described herein. In one embodiment, the method comprises cultivating the recombinant microorganism in a culture medium containing a feedstock providing a carbon source until the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate is produced. In a further embodiment, the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate is recovered. Recovery can be by methods known in the art, such as distillation, membrane-based separation gas stripping, solvent extraction, and expanded bed adsorption.

In some embodiments, the feedstock comprises a carbon source. In various embodiments described herein, the carbon source may be selected from sugars, glycerol, alcohols, organic acids, alkanes, fatty acids, lignocellulose, proteins, carbon dioxide, and carbon monoxide. In a further embodiment, the sugar is selected from the group consisting of glucose, fructose, and sucrose.

Enzyme Engineering

The enzymes in the recombinant microorganism can be engineered to improve one or more aspects of the substrate to product conversion. Non-limiting examples of enzymes that can be further engineered for use in methods of the disclosure include a desaturase (e.g., a fatty acyl-CoA desaturase or fatty acyl-ACP desaturase), an acyl-ACP synthetase, a fatty acid synthetase, a fatty acid synthase complex, an acetyl transferase, dehydrogenase, and an alcohol oxidase, and combinations thereof. These enzymes can be engineered for improved catalytic activity, improved selectivity, improved stability, improved tolerance to various fermentations conditions (temperature, pH, etc.), or

improved tolerance to various metabolic substrates, products, by-products, intermediates, etc.

Desaturase enzymes can be engineered for improved catalytic activity in the desaturation of an unsaturated substrate, for improved hydrocarbon selectivity, for improved selectivity of a Z product over an E product, or an E product over a Z product. For example, the Z9 fatty-acyl desaturase can be engineered to improve the yield in the substrate to product conversion of a saturated fatty acyl-CoA to the corresponding unsaturated fatty acyl-CoA, and, in addition or in the alternative, to improve selectivity of the desaturation at the 9 position to produce a corresponding Z-9 fatty acyl-CoA. In further non-limiting examples, the fatty acyl-ACP synthetase can be engineered for improved ACP ligation activity; a fatty acid synthase complex enzyme can be engineered for improved catalytic activity of elongation of a fatty acid substrate; a fatty alcohol forming fatty acyl-reductase can be engineered for improved catalytic activity in the reduction of a fatty acyl-CoA to a corresponding fatty alcohol; a fatty aldehyde forming fatty acyl-reductase can be engineered for improved catalytic activity in the reduction of a fatty acyl-ACP to a corresponding fatty aldehyde; a dehydrogenase can be engineered for improved catalytic activity in the conversion of a fatty acyl-ACP to a corresponding fatty alcohol; an alcohol oxidase can be engineered for improved catalytic activity in the conversion of a fatty alcohol into a corresponding fatty aldehyde; and an acetyl transferase can be engineered for improved catalytic activity in the conversion of a fatty alcohol into a corresponding fatty acetate.

The term “improved catalytic activity” as used herein with respect to a particular enzymatic activity refers to a higher level of enzymatic activity than that measured relative to a comparable non-engineered enzyme, such as a non-engineered desaturase (e.g. fatty acyl-CoA desaturase or fatty acyl-ACP desaturase), fatty alcohol or aldehyde forming fatty-acyl reductase, acyl-ACP synthetase, fatty acid synthetase, fatty acid synthase complex, acyl transferase, dehydrogenase, or an alcohol oxidase enzyme. For example, overexpression of a specific enzyme can lead to an increased level of activity in the cells for that enzyme. Mutations can be introduced into a desaturase (e.g. fatty acyl-CoA desaturase or fatty acyl-ACP desaturase), a fatty alcohol or aldehyde forming fatty-acyl reductase, a acyl-ACP synthetase, a fatty acid synthetase, a fatty acid synthase complex, a acyl transferase, a dehydrogenase, or an alcohol oxidase enzyme resulting in engineered enzymes with improved catalytic activity. Methods to increase enzymatic activity are known to those skilled in the art. Such techniques can include increasing the expression of the enzyme by increasing plasmid copy number and/or use of a stronger promoter and/or use of activating riboswitches, introduction of mutations to relieve negative regulation of the enzyme, introduction of specific mutations to increase specific activity and/or decrease the K_M for the substrate, or by directed evolution. See, e.g., *Methods in Molecular Biology* (vol. 231), ed. Arnold and Georgiou, Humana Press (2003). Metabolic Engineering—Enzyme Overexpression and Gene Deletion/Downregulation for Increased Pathway Flux

In various embodiments described herein, the exogenous and endogenous enzymes in the recombinant microorganism participating in the biosynthesis pathways described herein may be overexpressed.

The terms “overexpressed” or “overexpression” refers to an elevated level (e.g., aberrant level) of mRNAs encoding for a protein(s), and/or to elevated levels of protein(s) in cells as compared to similar corresponding unmodified cells

expressing basal levels of mRNAs or having basal levels of proteins. In particular embodiments, mRNA(s) or protein(s) may be overexpressed by at least 2-fold, 3-fold, 4-fold, 5-fold, 6-fold, 8-fold, 10-fold, 12-fold, 15-fold or more in microorganisms engineered to exhibit increased gene mRNA, protein, and/or activity.

In some embodiments, a recombinant microorganism of the disclosure is generated from a host that contains the enzymatic capability to synthesize a substrate fatty acid. In this specific embodiment it can be useful to increase the synthesis or accumulation of a fatty acid to, for example, increase the amount of fatty acid available to an engineered fatty alcohol production pathway.

In some embodiments, it may be useful to increase the expression of endogenous or exogenous enzymes involved in the fatty alcohol, aldehyde, or acetate production pathway to increase flux from the fatty acid to the fatty alcohol, aldehyde, or acetate, thereby resulting in increased synthesis or accumulation of the fatty alcohol, aldehyde, or acetate.

In some embodiments, it may be useful to increase the expression of endogenous or exogenous enzymes to increase intracellular levels of a coenzyme. In one embodiment, the coenzyme is NADH. In another embodiment, the coenzyme is NADPH. In one embodiment, the expression of proteins in the pentose phosphate pathway is increased to increase the intracellular levels of NADPH. The pentose phosphate pathway is an important catabolic pathway for supplying reduction equivalents and an important anabolic pathway for biosynthesis reactions. In one embodiment, a glucose-6-phosphate dehydrogenase that converts glucose-6-phosphate to 6-phospho D-glucono-1,5-lactone is overexpressed. In some embodiments, the glucose-6-phosphate dehydrogenase is ZWF1 from yeast. In another embodiment, the glucose-6-phosphate dehydrogenase is ZWF1 (YNL241C) from *Saccharomyces cerevisiae*. In one embodiment, a glucose-6-phosphate-1-dehydrogenase that converts D-glucopyranose-6-phosphate to 6-phospho D-glucono-1,5-lactone is overexpressed. In another embodiment, the glucose-6-phosphate-1-dehydrogenase is zwf from bacteria. In certain embodiments, the glucose-6-phosphate-1-dehydrogenase is zwf (NP_416366) from *E. coli*. In one embodiment, a 6-phosphogluconolactonase that converts 6-phospho D-glucono-1,5-lactone to D-gluconate 6-phosphate is overexpressed. In some embodiments, the 6-phosphogluconolactonase is SOL3 of yeast. In certain embodiments, the 6-phosphogluconolactonase is SOL3 (NP_012033) of *Saccharomyces cerevisiae*. In some embodiments, the 6-phosphogluconolactonase is SOL4 of yeast. In certain embodiments, the 6-phosphogluconolactonase is SOL4 (NP_011764) of *Saccharomyces cerevisiae*. In some embodiments, the 6-phosphogluconolactonase is pgl of bacteria. In certain embodiments, the 6-phosphogluconolactonase is pgl (NP_415288) of *E. coli*. In one embodiment, a 6-phosphogluconate dehydrogenase that converts D-gluconate 6-phosphate to D-ribulose 5-phosphate is overexpressed. In some embodiments, the 6-phosphogluconate dehydrogenase is GND1 from yeast. In certain embodiments, the 6-phosphogluconate dehydrogenase is GND1 (YHR183W) from *Saccharomyces cerevisiae*. In some embodiments, the 6-phosphogluconate dehydrogenase is GND2 from yeast. In certain embodiments, the 6-phosphogluconate dehydrogenase is GND2 (YGR256W) from *Saccharomyces cerevisiae*. In some embodiments, the 6-phosphogluconate dehydrogenase is gnd from bacteria. In certain embodiments, the 6-phosphogluconate dehydrogenase is gnd (NP_416533) from *E. coli*. In one embodiment, a transaldolase that interconverts D-glyceraldehyde 3-phosphate and D-sedo-

heptulose 7-phosphate to β -D-fructofuranose 6-phosphate and D-erythrose 4-phosphate is overexpressed. In some embodiments, the transaldolase is TAL1 of yeast. In certain embodiments, the transaldolase is TAL1 (NP_013458) of *Saccharomyces cerevisiae*. In some embodiments, the transaldolase is NQM1 of yeast. In certain embodiments, the transaldolase is NQM1 (NP_011557) of *Saccharomyces cerevisiae*. In some embodiments, the transaldolase is tal of bacteria. In certain embodiments, the transaldolase is talB (NP_414549) of *E. coli*. In certain embodiments, the transaldolase is talA (NP_416959) of *E. coli*. In one embodiment, a transketolase that interconverts D-erythrose 4-phosphate and D-xylulose 5-phosphate to β -D-fructofuranose 6-phosphate and D-glyceraldehyde 3-phosphate and/or interconverts D-sedoheptulose 7-phosphate and D-glyceraldehyde 3-phosphate to D-ribose 5-phosphate and D-xylulose 5-phosphate is overexpressed. In some embodiments, the transketolase is TKL1 of yeast. In certain embodiments, the transketolase is TKL1 (NP_015399) of *Saccharomyces cerevisiae*. In some embodiments, the transketolase is TKL2 of yeast. In some embodiments, the transketolase is TKL2 (NP_009675) of *Saccharomyces cerevisiae*. In some embodiments, the transketolase is tkt of bacteria. In certain embodiments, the transketolase is tktA (YP_026188) of *E. coli*. In certain embodiments, the transketolase is tktB (NP_416960) of *E. coli*. In one embodiment, a ribose-5-phosphate ketol-isomerase that interconverts D-ribose 5-phosphate and D-ribulose 5-phosphate is overexpressed. In some embodiments, the ribose-5-phosphate ketol-isomerase is RKI1 of yeast. In certain embodiments, the ribose-5-phosphate ketol-isomerase is RKI1 (NP_014738) of *Saccharomyces cerevisiae*. In some embodiments, the ribose-5-phosphate isomerase is rpi of bacteria. In certain embodiments, the ribose-5-phosphate isomerase is rpiA (NP_417389) of *E. coli*. In certain embodiments, the ribose-5-phosphate isomerase is rpiB (NP_418514) of *E. coli*. In one embodiment, a D-ribulose-5-phosphate 3-epimerase that interconverts D-ribulose 5-phosphate and D-xylulose 5-phosphate is overexpressed. In some embodiments, the D-ribulose-5-phosphate 3-epimerase is RPE1 of yeast. In certain embodiments, the D-ribulose-5-phosphate 3-epimerase is RPE1 (NP_012414) of *Saccharomyces cerevisiae*. In some embodiments, the D-ribulose-5-phosphate 3-epimerase is rpe of bacteria. In certain embodiments, the D-ribulose-5-phosphate 3-epimerase is rpe (NP_417845) of *E. coli*.

In one embodiment, the expression of an NADP+-dependent isocitrate dehydrogenase is increased to increase intracellular levels of a coenzyme. In one embodiment, an NADP+ dependent isocitrate dehydrogenase oxidizes D-threo-isocitrate to 2-oxoglutarate with concomitant generation of NADPH. In another embodiment, an NADP+ dependent isocitrate dehydrogenase oxidizes D-threo-isocitrate to 2-oxalosuccinate with concomitant generation of NADPH. In some embodiments, the NADP+-dependent isocitrate dehydrogenase is IDP from yeast. In certain embodiments, the NADP+-dependent isocitrate dehydrogenase is IDP2 (YLR174W) from *Saccharomyces cerevisiae*. In some embodiments, the NADP+-dependent isocitrate dehydrogenase is icd from bacteria. In certain embodiments, the NADP+-dependent isocitrate dehydrogenase is icd (NP_415654) from *E. coli*.

In some embodiments, the expression of a malic enzyme that decarboxylates malate to pyruvate with concomitant generation of NADH or NADPH is increased to increase intracellular levels of a coenzyme. In one embodiment, the malic enzyme is NAD+ dependent. In another embodiment,

the malic enzyme is NADP+ dependent. In one embodiment, the malic enzyme is an NAD+ dependent malate dehydrogenase from bacteria. In some embodiments, the NAD+ dependent malate dehydrogenase is maeA (NP_415996) from *E. coli*. In some embodiments, the NAD+ dependent malate dehydrogenase is maeE (CAQ68119) from *Lactobacillus casei*. In another embodiment, the malic enzyme is a mitochondrial NAD+ dependent malate dehydrogenase from yeast. In some embodiments, the NAD+ dependent malate dehydrogenase is MAE1 (YKL029C) from *S. cerevisiae*. In another embodiment, the malic enzyme is a mitochondrial NAD+ dependent malate dehydrogenase from a parasitic nematode. In some embodiments, the NAD+ dependent malate dehydrogenase is M81055 from *Ascaris suum*. In one embodiment, the malic enzyme is an NADP+ dependent malate dehydrogenase from bacteria. In some embodiments, the NADP+ dependent malate dehydrogenase is maeB (NP_416958) from *E. coli*. In one embodiment, the malic enzyme is an NADP+ dependent malate dehydrogenase from corn. In some embodiments, the NADP+ dependent malate dehydrogenase is me1 from *Zea mays*.

In some embodiments, the expression of an aldehyde dehydrogenase that oxidizes an aldehyde to a carboxylic acid with concomitant generation of NADH or NADPH is increased to increase intracellular levels of a coenzyme. In one embodiment, the aldehyde dehydrogenase is NAD+ dependent. In another embodiment, the aldehyde dehydrogenase is NADP+ dependent. In one embodiment, the aldehyde dehydrogenase is an NAD+ dependent aldehyde dehydrogenase from bacteria. In some embodiments, the NAD+ dependent aldehyde dehydrogenase is aldA (NP_415933) from *E. coli*. In another embodiment, the aldehyde dehydrogenase is a cytosolic NADP+ dependent aldehyde dehydrogenase from yeast. In some embodiments, the NADP+ dependent aldehyde dehydrogenase is ALD6 (YPL061W) from *S. cerevisiae*. In another embodiment, the aldehyde dehydrogenase is a cytosolic NADP+ dependent aldehyde dehydrogenase from bacteria. In some embodiments, the NADP+ dependent aldehyde dehydrogenase is aldB (NP_418045) from *E. coli*.

In one embodiment, overexpression of an enzyme to increase intracellular levels of a coenzyme comprises coupling supplementation of a co-substrate and overexpression of the enzyme. In one embodiment, the overexpression of an enzyme coupled with supplementation of a co-substrate of that enzyme increase flux through a biochemical pathway. In one embodiment, an NAD+ or NADP+ dependent alcohol dehydrogenase is expressed with a co-substrate. In certain embodiments, an alcohol dehydrogenase is expressed with an isopropanol co-substrate. In one embodiment, an NAD+ or NADP+ dependent glucose dehydrogenase is expressed with a co-substrate. In certain embodiments, a glucose dehydrogenase is expressed with a glucose co-substrate.

In one embodiment, the expression of a transhydrogenase is increased to interconvert NADH and NADPH. In some embodiments, the transhydrogenase is a pyridine nucleotide transhydrogenase. In some embodiments, the pyridine nucleotide transhydrogenase is from bacteria. In certain embodiments, the pyridine nucleotide transhydrogenase is pntAB (beta subunit: NP_416119; alpha subunit: NP_416120) from *E. coli*. In some embodiments, the pyridine nucleotide transhydrogenase is from human. In certain embodiments, the pyridine nucleotide transhydrogenase is NNT (NP_036475) from *Homo sapiens*. In certain embodiments, the pyridine nucleotide transhydrogenase is from

Solanum tuberosum. In certain embodiments, the pyridine nucleotide transhydrogenase is from *Spinacea oleracea*.

In some embodiments, it may be useful to increase the expression of endogenous or exogenous proteins to induce endoplasmic reticulum (ER) membrane proliferation. In some embodiments, the induction of endoplasmic reticulum membrane proliferation can improve production of fatty alcohols, aldehydes, or acetates. In one embodiment, the expression of an inactivated HMG-CoA reductase (hydroxymethylglutaryl-CoA reductase) containing one or more ER facing loops is increased. In certain embodiments, the one or more loops is between transmembrane domains 6 and 7 of an inactivated HMG-CoA reductase. In some embodiments, the inactivated HMG-CoA reductase comprises an inactivated protein or chimera which codes for the first 500 amino acids or a subsequence of the first 500 amino acids of *Yarrowia lipolytica* YALI0E04807p. In other embodiments, the inactivated HMG-CoA reductase comprises an inactivated protein or chimera which codes for the first 522 amino acids or a subsequence of the first 522 amino acids of HMG1 from *Saccharomyces cerevisiae* (NP_013636.1). In other embodiments, the inactivated HMG-CoA reductase comprises an inactivated protein or chimera which codes for the first 522 amino acids or a subsequence of the first 522 amino acids of HMG2 from *Saccharomyces cerevisiae* (NP_013555.1). In some embodiments, the expression of one or more regulatory proteins is increased to improve production of fatty alcohols, aldehydes, or acetates. In certain embodiments, the regulatory protein comprises HAC1 transcription factor from *Saccharomyces cerevisiae* (NP_116622.1). In certain embodiments, the regulatory protein comprises HAC1 transcription factor from *Yarrowia lipolytica* (YALI0B12716p).

Increased synthesis or accumulation can be accomplished by, for example, overexpression of nucleic acids encoding one or more of the above-described a fatty alcohol pathway enzymes. Overexpression of a fatty alcohol pathway enzyme or enzymes can occur, for example, through increased expression of an endogenous gene or genes, or through the expression, or increased expression, of an exogenous gene or genes. Therefore, naturally occurring organisms can be readily modified to generate non-natural, fatty alcohol producing microorganisms through overexpression of one or more nucleic acid molecules encoding a fatty alcohol biosynthetic pathway enzyme. In addition, a non-naturally occurring organism can be generated by mutagenesis of an endogenous gene that results in an increase in activity of an enzyme in the fatty alcohol biosynthetic pathways.

Equipped with the present disclosure, the skilled artisan will be able to readily construct the recombinant microorganisms described herein, as the recombinant microorganisms of the disclosure can be constructed using methods well known in the art as exemplified above to exogenously express at least one nucleic acid encoding a fatty alcohol pathway enzyme in sufficient amounts to produce a fatty alcohol.

Methods for constructing and testing the expression levels of a non-naturally occurring fatty alcohol-producing host can be performed, for example, by recombinant and detection methods well known in the art. Such methods can be found described in, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Third Ed., Cold Spring Harbor Laboratory, New York (2001); Ausubo et al., *Current Protocols in Molecular Biology*, John Wiley and Sons, Baltimore, Md. (1999).

A variety of mechanisms known in the art can be used to express, or overexpress, exogenous or endogenous genes.

81

For example, an expression vector or vectors can be constructed to harbor one or more fatty alcohol biosynthetic pathway enzyme encoding nucleic acids as exemplified herein operably linked to expression control sequences functional in the host organism. Expression vectors applicable for use in the microbial host organisms of the invention include, for example, plasmids, phage vectors, viral vectors, episomes and artificial chromosomes, including vectors and selection sequences or markers operable for stable integration into a host chromosome. Selectable marker genes also can be included that, for example, provide resistance to antibiotics or toxins, complement auxotrophic deficiencies, or supply critical nutrients not in the culture media. Expression control sequences can include constitutive and inducible promoters, transcription enhancers, transcription terminators, and the like which are well known in the art. When two or more exogenous encoding nucleic acids are to be co-expressed, both nucleic acids can be inserted, for example, into a single expression vector or in separate expression vectors. For single vector expression, the encoding nucleic acids can be operationally linked to one common expression control sequence or linked to different expression control sequences, such as one inducible promoter and one constitutive promoter. The transformation of exogenous nucleic acid sequences involved in a metabolic or synthetic pathway can be confirmed using methods well known in the art.

Expression control sequences are known in the art and include, for example, promoters, enhancers, polyadenylation signals, transcription terminators, internal ribosome entry sites (IRES), and the like, that provide for the expression of the polynucleotide sequence in a host cell. Expression control sequences interact specifically with cellular proteins involved in transcription (Maniatis et al., *Science*, 236: 1237-1245 (1987)). Exemplary expression control sequences are described in, for example, Goeddel, *Gene Expression Technology: Methods in Enzymology*, Vol. 185, Academic Press, San Diego, Calif. (1990).

In various embodiments, an expression control sequence may be operably linked to a polynucleotide sequence. By “operably linked” is meant that a polynucleotide sequence and an expression control sequence(s) are connected in such a way as to permit gene expression when the appropriate molecules (e.g., transcriptional activator proteins) are bound to the expression control sequence(s). Operably linked promoters are located upstream of the selected polynucleotide sequence in terms of the direction of transcription and translation. Operably linked enhancers can be located upstream, within, or downstream of the selected polynucleotide.

In some embodiments, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that competes with the biosynthesis pathway for the production of a mono- or poly-unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate.

In some embodiments, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes the conversion of a fatty acid into a ω -hydroxyfatty acid. In some such embodiments, the enzymes that catalyze the conversion of a fatty acid into a ω -hydroxyfatty acid are selected from the group consisting of XP_504406, XP_504857, XP_504311, XP_500855, XP_500856, XP_500402, XP_500097, XP_501748, XP_500560, XP_501148, XP_501667, XP_500273, BAA02041, CAA39366, CAA39367, BAA02210, BAA02211,

82

BAA02212, BAA02213, BAA02214, AAO73952, AAO73953, AAO73954, AAO73955, AAO73956, AAO73958, AAO73959, AAO73960, AAO73961, AAO73957, XP_002546278, BAM49649, AAB80867, AAB17462, ADL27534, AAU24352, AAA87602, CAA34612, ABM17701, AAA25760, CAB51047, AAC82967, WP_011027348, or homologs thereof.

In other embodiments, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes the conversion of a fatty acyl-CoA into α,β -enoyl-CoA. In some such embodiments, the enzymes that catalyze the conversion of a fatty acyl-CoA into α,β -enoyl-CoA are selected from the group consisting of CAA04659, CAA04660, CAA04661, CAA04662, CAA04663, CAG79214, AAA34322, AAA34361, AAA34363, CAA29901, BAA04761, AAA34891, AAB08643, CAB15271, BAN55749, CAC44516, ADK16968, AE137634, WP_000973047, WP_025433422, WP_035184107, WP_026484842, CEL80920, WP_026818657, WP_005293707, WP_005883960, or homologs thereof.

In some embodiments, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more proteins involved in peroxisome biogenesis. In such embodiments, the one or more proteins involved in peroxisome biogenesis are selected from the group consisting of XP_505754, XP_501986, XP_501311, XP_504845, XP_503326, XP_504029, XP_002549868, XP_002547156, XP_002545227, XP_002547350, XP_002546990, EIW11539, EIW08094, EIW11472, EIW09743, EIW0828, or homologs thereof.

In some embodiments, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that competes with the biosynthesis pathway for one or more unsaturated fatty acyl-CoA intermediates. In one embodiment, the one or more endogenous enzymes comprise one or more diacylglycerol acyltransferases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more diacylglycerol acyltransferases selected from the group consisting of YALI0E32769g, YALI0D07986g and CTRG_06209, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more glycerolphospholipid acyltransferases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more glycerolphospholipid acyltransferases selected from the group consisting of YALI0E16797g and CTG_04390, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more acyl-CoA/sterol acyltransferases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more acyl-CoA/sterol acyltransferases selected from the group consisting of YALI0F06578g, CTRG_01764 and CTRG_01765, or homolog thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that oxidizes fatty aldehyde intermediates. In one embodiment, the one or more endogenous enzymes comprise one or more fatty aldehyde dehydrogenases. In the context of a recombinant yeast microorganism,

the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more fatty aldehyde dehydrogenases selected from the group consisting of YALI0A17875g, YALI0E15400g, YALI0B01298g, YALI0F23793g, CTRG_05010 and CTRG_04471, or homolog thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous enzymes that catalyzes a reaction in a pathway that consumes fatty acetate products. In one embodiment, the one or more endogenous enzymes comprise one or more sterol esterases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more sterol esterases selected from the group consisting of YALI0E32035g, YALI0E00528g, CTRG_01138, CTRG_01683 and CTRG_04630, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more triacylglycerol lipases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more triacylglycerol lipases selected from the group consisting of YALI0D17534g, YALI0F10010g, CTRG_00057 and CTRG_06185, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more monoacylglycerol lipases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more monoacylglycerol lipases selected from the group consisting of YALI0C14520g, CTRG_03360 and CTRG_05049, or homolog thereof. In another embodiment, the one or more endogenous enzymes comprise one or more extracellular lipases. In the context of a recombinant yeast microorganism, the recombinant yeast microorganism is engineered to delete, disrupt, mutate, and/or reduce the activity of one or more extracellular lipases selected from the group consisting of YALI0A20350g, YALI0D19184g, YALI0B09361g, CTRG_05930, CTRG_04188, CTRG_02799, CTRG_03052 and CTRG_03885, or homolog thereof.

In another embodiment, the recombinant microorganism is manipulated to delete, disrupt, mutate, and/or reduce the activity of one or more endogenous reductase or desaturase enzymes that interferes with the unsaturated C₆-C₂₄ fatty alcohol, aldehyde, or acetate, i.e., catalyzes the conversion of a pathway substrate or product into an unwanted by-product.

Chemical Conversion of Product from Microorganism Synthesis

The present disclosure describes chemical conversions that can be used to convert a product synthesized by recombinant microorganism into a down-stream product.

In some embodiments, an unsaturated fatty alcohol, aldehyde, acetate, or carboxylic acid produced by a microorganism can undergo subsequent chemical conversion to produce a pheromone, fragrance, flavor, polymer, or polymer intermediate. Non-limiting examples of chemical transformations include esterification, metathesis, and polymerization.




Unsaturated fatty carboxylic acids can be esterified by methods known in the art. For example, Fischer esterification can be used to convert a fatty carboxylic acid to a corresponding fatty ester. See, e.g., Komura, K. et al., *Synthesis* 2008. 3407-3410.

Elongation of the carbon chain can be performed by 65 known methods to covert an unsaturated fatty alcohol into an elongated derivative thereof. Olefin metastasis catalysts

can be performed to increase the number of carbons on the fatty carbon chain and impart Z or E stereochemistry on the corresponding unsaturated product.


In general, any metathesis catalyst stable under the reaction conditions and nonreactive with functional groups on the fatty substrate (e.g., alcohol, ester, carboxylic acid, aldehyde, or acetate) can be used with the present disclosure. Such catalysts are, for example, those described by Grubbs (Grubbs, R. H., "Synthesis of large and small molecules using olefin metathesis catalysts." *PMSE Prepr.*, 2012), herein incorporated by reference in its entirety. Depending on the desired isomer of the olefin, as cis-selective metathesis catalyst may be used, for example one of those described by Shahane et al. (Shahane, S., et al. *Chem-CatChem*, 2013. 5(12): p. 3436-3459), herein incorporated by reference in its entirety. Specific catalysts 1-5 exhibiting cis-selectivity are shown below (Scheme 1) and have been described previously (Khan, R. K., et al. *J. Am. Chem. Soc.*, 2013. 135(28): p. 10258-61; Hartung, J. et al. *J. Am. Chem. Soc.*, 2013. 135(28): p. 10183-5.; Rosebrugh, L. E., et al. *J. Am. Chem. Soc.*, 2013. 135(4): p. 1276-9.; Marx, V. M., et al. *J. Am. Chem. Soc.*, 2013. 135(1): p. 94-7.; Herbert, M. B., et al. *Angew. Chem. Int. Ed. Engl.*, 2013. 52(1): p. 310-4; Keitz, B. K., et al. *J. Am. Chem. Soc.*, 2012. 134(4): p. 2040-3.; Keitz, B. K., et al. *J. Am. Chem. Soc.*, 2012. 134(1): p. 693-9.; Endo, K. et al. *J. Am. Chem. Soc.*, 2011. 133(22): p. 8525-7).

Scheme 1



85

-continued



Additional Z-selective catalysts are described in (Cannon and Grubbs 2013; Bronner et al. 2014; Hartung et al. 2014; Pribisko et al. 2014; Quigley and Grubbs 2014) and are herein incorporated by reference in their entirety. Due to their excellent stability and functional group tolerance, in some embodiments metathesis catalysts include, but are not limited to, neutral ruthenium or osmium metal carbene complexes that possess metal centers that are formally in the +2 oxidation state, have an electron count of 16, are penta-coordinated, and are of the general formula $LL'AA'M=CRbRc$ or $LL'AA'M=(C=n)CRbRc$ (Pederson and Grubbs 2002); wherein

M is ruthenium or osmium;

L and L' are each independently any neutral electron donor ligand and selected from phosphine, sulfonated phosphine, phosphite, phosphinite, phosphonite, arsine, stibnite, ether, amine, amide, imine, sulfoxide, carboxyl, nitrosyl, pyridine, thioether, or heterocyclic carbenes; and

A and A' are anionic ligands independently selected from halogen, hydrogen, C_1 - C_{20} alkyl, aryl, C_1 - C_{20} alkoxide, aryloxy, C_2 - C_{20} alkoxycarbonyl, arylcarboxylate, C_1 - C_{20} carboxylate, arylsulfonyl, C_1 - C_{20} alkylsulfonyl, C_1 - C_{20} alkylsulfinyl; each ligand optionally being substituted with C_1 - C_5 alkyl, halogen, C_1 - C_5 alkoxy; or with a phenyl group that is optionally substituted with halogen, C_1 - C_5 alkyl, or C_1 - C_5 alkoxy; and A and A' together may optionally comprise a bidentate ligand; and

R_b and R_c are independently selected from hydrogen, C_1 - C_{20} alkyl, aryl, C_1 - C_{20} carboxylate, C_1 - C_{20} alkoxy, aryloxy, C_1 - C_{20} alkoxycarbonyl, C_1 - C_{20} alkylthio, C_1 - C_{20} alkylsulfonyl and C_1 - C_{20} alkylsulfinyl, each of R_b and R_c optionally substituted with C_1 - C_5 alkyl, halogen, C_1 - C_5 alkoxy or with a phenyl group that is optionally substituted with halogen, C_1 - C_5 alkyl, or C_1 - C_5 alkoxy.

Other metathesis catalysts such as "well defined catalysts" can also be used. Such catalysts include, but are not limited

86

to, Schrock's molybdenum metathesis catalyst, 2,6-diisopropylphenylimido neophylidenemolybdenum (VI) bis (hexafluoro-t-butoxide), described by Grubbs et al. (*Tetrahedron* 1998, 54: 4413-4450) and Basset's tungsten metathesis catalyst described by Couturier, J. L. et al. (*Angew. Chem. Int. Ed. Engl.* 1992, 31: 628).





Catalysts useful in the methods of the disclosure also include those described by Peryshkov, et al. *J. Am. Chem. Soc.* 2011, 133: 20754-20757; Wang, et al. *Angewandte Chemie*, 2013, 52: 1939-1943; Yu, et al. *J. Am. Chem. Soc.*, 2012, 134: 2788-2799; Halford. *Chem. Eng. News*, 2011, 89 (45): 11; Yu, et al. *Nature*, 2011, 479: 88-93; Lee. *Nature*, 2011, 471: 452-453; Meek, et al. *Nature*, 2011: 471, 461-466; Flook, et al. *J. Am. Chem. Soc.* 2011, 133: 1784-1786; Zhao, et al. *Org. Lett.*, 2011, 13(4): 784-787; Ondi, et al. "High activity, stabilized formulations, efficient synthesis and industrial use of Mo- and W-based metathesis catalysts" *XiMo Technology Updates*, 2015: http://www.ximo-inc.com/files/ximo/uploads/download/Summary_3.11.15.pdf; Schrock, et al. *Macromolecules*, 2010: 43, 7515-7522; Peryshkov, et al. *Organometallics* 2013: 32, 5256-5259; Gerber, et al. *Organometallics* 2013: 32, 5573-5580; Marinescu, et al. *Organometallics* 2012: 31, 6336-6343; Wang, et al. *Angew. Chem. Int. Ed.* 2013: 52, 1939-1943; Wang, et al. *Chem. Eur. J.* 2013: 19, 2726-2740; and Townsend et al. *J. Am. Chem. Soc.* 2012: 134, 11334-11337.

Catalysts useful in the methods of the disclosure also include those described in International Pub. No. WO 2014/155185; International Pub. No. WO 2014/172534; U.S. Pat. Appl. Pub. No. 2014/0330018; International Pub. No. WO 2015/003815; and International Pub. No. WO 2015/003814.





Catalysts useful in the methods of the disclosure also include those described in U.S. Pat. Nos. 4,231,947; 4,245,131; 4,427,595; 4,681,956; 4,727,215; International Pub. No. WO 1991/009825; U.S. Pat. Nos. 5,087,710; 5,142,073; 5,146,033; International Pub. No. WO 1992/019631; U.S. Pat. Nos. 6,121,473; 6,346,652; 8,987,531; U.S. Pat. Appl. Pub. No. 2008/0119678; International Pub. No. WO 2008/066754; International Pub. No. WO 2009/094201; U.S. Pat. Appl. Pub. No. 2011/0015430; U.S. Pat. Appl. Pub. No. 2011/0065915; U.S. Pat. Appl. Pub. No. 2011/0077421; International Pub. No. WO 2011/040963; International Pub. No. WO 2011/097642; U.S. Pat. Appl. Pub. No. 2011/0237815; U.S. Pat. Appl. Pub. No. 2012/0302710; International Pub. No. WO 2012/167171; U.S. Pat. Appl. Pub. No. 2012/0323000; U.S. Pat. Appl. Pub. No. 2013/0116434; International Pub. No. WO 2013/070725; U.S. Pat. Appl. Pub. No. 2013/0274482; U.S. Pat. Appl. Pub. No. 2013/0281706; International Pub. No. WO 2014/139679; International Pub. No. WO 2014/169014; U.S. Pat. Appl. Pub. No. 2014/0330018; and U.S. Pat. Appl. Pub. No. 2014/0378637.

Catalysts useful in the methods of the disclosure also include those described in International Pub. No. WO 2007/075427; U.S. Pat. Appl. Pub. No. 2007/0282148; International Pub. No. WO 2009/126831; International Pub. No. WO 2011/069134; U.S. Pat. Appl. Pub. No. 2012/0123133; U.S. Pat. Appl. Pub. No. 2013/0261312; U.S. Pat. Appl. Pub. No. 2013/0296511; International Pub. No. WO 2014/134333; and U.S. Pat. Appl. Pub. No. 2015/0018557.





Catalysts useful in the methods of the disclosure also include those set forth in the following table:

Structure	Name
	dichloro[1,3-bis(2,6-isopropylphenyl)-2-imidazolidinylidene](benzylidene)(tricyclohexylphosphine)ruthenium(II)
	dichloro[1,3-bis(2,6-isopropylphenyl)-2-imidazolidinylidene](2-isopropoxyphenylmethylene)ruthenium(II)
	dichloro[1,3-bis(2-methylphenyl)-2-imidazolidinylidene](benzylidene)(tricyclohexylphosphine)ruthenium(II)
	dichloro[1,3-bis(2-methylphenyl)-2-imidazolidinylidene](2-isopropoxyphenylmethylene)ruthenium(II)

-continued

Structure	Name
	dichloro[1,3-bis(2,4,6-trimethylphenyl)-2-imidazolidinylidene](benzylidene)bis(3-bromopyridine)ruthenium(II)
	dichloro[1,3-bis(2,4,6-trimethylphenyl)-2-imidazolidinylidene](3-methyl-2-butenylidene)(tricyclohexylphosphine)ruthenium(II)
	dichloro[1,3-bis(2,4,6-trimethylphenyl)-2-imidazolidinylidene][3-(2-pyridinyl)propylidene]ruthenium(II)
	dichloro[1,3-bis(2,4,6-trimethylphenyl)-2-imidazolidinylidene] [(tricyclohexylphosphoranyl) methylidene]ruthenium(II) tetrafluoroborate

-continued

Structure	Name
	dichloro(3-methyl-2-butenylidene) bis(tricyclohexylphosphine)ruthenium(II)
	dichloro(3-methyl-2-butenylidene) bis(tricyclopentylphosphine)ruthenium(II)
	dichloro(tricyclohexylphosphine) [(tricyclohexylphosphoranyl) methylidene]ruthenium(II) tetrafluoroborate
	bis(tricyclohexylphosphine)benzylidene ruthenium(IV)dichloride

-continued

Structure	Name
	[1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene]dichloro(phenylmethylene)(tricyclohexylphosphine)ruthenium
	(1,3-bis-(2,4,6-trimethylphenyl)-2-imidazolidinylidene)dichloro(o-isopropoxyphenylmethylene)ruthenium
	dichloro(o-isopropoxyphenylmethylene)(tricyclohexylphosphine)ruthenium(II)
	[2-(1-methylethoxy-O)phenylmethyl-C](nitrate-O,O'){rel-(2R,5R,7R)-adamantane-2,1-diyl}[3-(2,4,6-trimethylphenyl)-1-imidazolidinyl-2-ylidene]ruthenium

Catalysts useful in the methods of the disclosure also include those described in U.S. Pat. Appl. Pub. No. 2008/0009598; U.S. Pat. Appl. Pub. No. 2008/0207911; U.S. Pat. Appl. Pub. No. 2008/0275247; U.S. Pat. Appl. Pub. No. 2011/0040099; U.S. Pat. Appl. Pub. No. 2011/0282068; and U.S. Pat. Appl. Pub. No. 2015/0038723.

Catalysts useful in the methods of the disclosure include those described in International Pub. No. WO 2007/140954; U.S. Pat. Appl. Pub. No. 2008/0221345; International Pub. No. WO 2010/037550; U.S. Pat. Appl. Pub. No. 2010/0087644; U.S. Pat. Appl. Pub. No. 2010/0113795; U.S. Pat. Appl. Pub. No. 2010/0174068; International Pub. No. WO 2011/091980; International Pub. No. WO 2012/168183;

U.S. Pat. Appl. Pub. No. 2013/0079515; U.S. Pat. Appl. Pub. No. 2013/0144060; U.S. Pat. Appl. Pub. No. 2013/0211096; International Pub. No. WO 2013/135776; International Pub. No. WO 2014/001291; International Pub. No. WO 2014/067767; U.S. Pat. Appl. Pub. No. 2014/0171607; and U.S. Pat. Appl. Pub. No. 2015/0045558.

The catalyst is typically provided in the reaction mixture in a sub-stoichiometric amount (e.g., catalytic amount). In certain embodiments, that amount is in the range of about 0.001 to about 50 mol % with respect to the limiting reagent of the chemical reaction, depending upon which reagent is in stoichiometric excess. In some embodiments, the catalyst is present in less than or equal to about 40 mol % relative to the limiting reagent. In some embodiments, the catalyst is present in less than or equal to about 30 mol % relative to the limiting reagent. In some embodiments, the catalyst is present in less than about 20 mol %, less than about 10 mol %, less than about 5 mol %, less than about 2.5 mol %, less than about 1 mol %, less than about 0.5 mol %, less than about 0.1 mol %, less than about 0.015 mol %, less than about 0.01 mol %, less than about 0.0015 mol %, or less, relative to the limiting reagent. In some embodiments, the catalyst is present in the range of about 2.5 mol % to about 5 mol %, relative to the limiting reagent. In some embodiments, the reaction mixture contains about 0.5 mol % catalyst. In the case where the molecular formula of the catalyst complex includes more than one metal, the amount of the catalyst complex used in the reaction may be adjusted accordingly.

In some cases, the methods described herein can be performed in the absence of solvent (e.g., neat). In some cases, the methods can include the use of one or more solvents. Examples of solvents that may be suitable for use in the disclosure include, but are not limited to, benzene, p-cresol, toluene, xylene, diethyl ether, glycol, diethyl ether, petroleum ether, hexane, cyclohexane, pentane, methylene chloride, chloroform, carbon tetrachloride, dioxane, tetrahydrofuran (THF), dimethyl sulfoxide, dimethylformamide, hexamethyl-phosphoric triamide, ethyl acetate, pyridine, triethylamine, picoline, and the like, as well as mixtures thereof. In some embodiments, the solvent is selected from benzene, toluene, pentane, methylene chloride, and THF. In certain embodiments, the solvent is benzene.

In some embodiments, the method is performed under reduced pressure. This may be advantageous in cases where a volatile byproduct, such as ethylene, may be produced during the course of the metathesis reaction. For example, removal of the ethylene byproduct from the reaction vessel may advantageously shift the equilibrium of the metathesis reaction towards formation of the desired product. In some embodiments, the method is performed at a pressure of about less than 760 torr. In some embodiments, the method is performed at a pressure of about less than 700 torr. In some embodiments, the method is performed at a pressure of about less than 650 torr. In some embodiments, the method is performed at a pressure of about less than 600 torr. In some embodiments, the method is performed at a pressure of about less than 550 torr. In some embodiments, the method is performed at a pressure of about less than 500 torr. In some embodiments, the method is performed at a pressure of about less than 450 torr. In some embodiments, the method is performed at a pressure of about less than 400 torr. In some embodiments, the method is performed at a pressure of about less than 350 torr. In some embodiments, the method is performed at a pressure of about less than 300 torr. In some embodiments, the method is performed at a pressure of about less than 250 torr. In some embodiments, the method

is performed at a pressure of about less than 200 torr. In some embodiments, the method is performed at a pressure of about less than 150 torr. In some embodiments, the method is performed at a pressure of about less than 100 torr. In some embodiments, the method is performed at a pressure of about less than 90 torr. In some embodiments, the method is performed at a pressure of about less than 80 torr. In some embodiments, the method is performed at a pressure of about less than 70 torr. In some embodiments, the method is performed at a pressure of about less than 60 torr. In some embodiments, the method is performed at a pressure of about less than 50 torr. In some embodiments, the method is performed at a pressure of about less than 40 torr. In some embodiments, the method is performed at a pressure of about less than 30 torr. In some embodiments, the method is performed at a pressure of about less than 20 torr. In some embodiments, the method is performed at a pressure of about 20 torr.

In some embodiments, the method is performed at a pressure of about 19 torr. In some embodiments, the method is performed at a pressure of about 18 torr. In some embodiments, the method is performed at a pressure of about 17 torr. In some embodiments, the method is performed at a pressure of about 16 torr. In some embodiments, the method is performed at a pressure of about 15 torr. In some embodiments, the method is performed at a pressure of about 14 torr. In some embodiments, the method is performed at a pressure of about 13 torr. In some embodiments, the method is performed at a pressure of about 12 torr. In some embodiments, the method is performed at a pressure of about 11 torr. In some embodiments, the method is performed at a pressure of about 10 torr. In some embodiments, the method is performed at a pressure of about 10 torr. In some embodiments, the method is performed at a pressure of about 9 torr. In some embodiments, the method is performed at a pressure of about 8 torr. In some embodiments, the method is performed at a pressure of about 7 torr. In some embodiments, the method is performed at a pressure of about 6 torr. In some embodiments, the method is performed at a pressure of about 5 torr. In some embodiments, the method is performed at a pressure of about 4 torr. In some embodiments, the method is performed at a pressure of about 3 torr. In some embodiments, the method is performed at a pressure of about 2 torr. In some embodiments, the method is performed at a pressure of about 1 torr. In some embodiments, the method is performed at a pressure of less than about 1 torr.

In some embodiments, the two metathesis reactants are present in equimolar amounts. In some embodiments, the two metathesis reactants are not present in equimolar amounts. In certain embodiments, the two reactants are present in a molar ratio of about 20:1, 19:1, 18:1, 17:1, 16:1, 15:1, 14:1, 13:1, 12:1, 11:1, 10:1, 9:1, 8:1, 7:1, 6:1, 5:1, 4:1, 3:1, 2:1, 1:1, 1:2, 1:3, 1:4, 1:5, 1:6, 1:7, 1:8, 1:9, 1:10, 1:11, 1:12, 1:13, 1:14, 1:15, 1:16, 1:17, 1:18, 1:19, or 1:20. In certain embodiments, the two reactants are present in a molar ratio of about 10:1. In certain embodiments, the two reactants are present in a molar ratio of about 7:1. In certain embodiments, the two reactants are present in a molar ratio of about 5:1. In certain embodiments, the two reactants are present in a molar ratio of about 2:1. In certain embodiments, the two reactants are present in a molar ratio of about 1:10. In certain embodiments, the two reactants are present in a molar ratio of about 1:7. In certain embodiments, the two reactants are present in a molar ratio of about 1:5. In certain embodiments, the two reactants are present in a molar ratio of about 1:2.

In general, the reactions with many of the metathesis catalysts disclosed herein provide yields better than 15%, better than 50%, better than 75%, or better than 90%. In addition, the reactants and products are chosen to provide at least a 5° C. difference, a greater than 20° C. difference, or a greater than 40° C. difference in boiling points. Additionally, the use of metathesis catalysts allows for much faster product formation than byproduct, it is desirable to run these reactions as quickly as practical. In particular, the reactions are performed in less than about 24 hours, less than 12 hours, less than 8 hours, or less than 4 hours.

One of skill in the art will appreciate that the time, temperature and solvent can depend on each other, and that changing one can require changing the others to prepare the pyrethroid products and intermediates in the methods of the disclosure. The metathesis steps can proceed at a variety of temperatures and times. In general, reactions in the methods of the disclosure are conducted using reaction times of several minutes to several days. For example, reaction times of from about 12 hours to about 7 days can be used. In some embodiments, reaction times of 1-5 days can be used. In some embodiments, reaction times of from about 10 minutes to about 10 hours can be used. In general, reactions in the methods of the disclosure are conducted at a temperature of from about 0° C. to about 200° C. For example, reactions can be conducted at 15-100° C. In some embodiments, reaction can be conducted at 20-80° C. In some embodiments, reactions can be conducted at 100-150° C.

Unsaturated fatty esters can be reduced using a suitable reducing agent which selectively reduces the ester to the corresponding aldehyde or alcohol but does not reduce the double bond. An unsaturated fatty ester can be reduced to the corresponding unsaturated fatty aldehyde using di-isobutyl aluminum halide (DIBAL) or Vitride®. The unsaturated fatty aldehyde can be reduced to the corresponding fatty alcohol with, e.g., DIBAL or Vitride®. In some embodiments, the unsaturated fatty ester can be reduced to the corresponding fatty alcohol using AlH₃ or 9-Borabicyclo (3.3.1)nonane (9-BBN). (See Galatis, P. *Encyclopedia of Reagents for Organic Synthesis*. 2001. New York: John Wiley & Sons; and Carey & Sunderburg. *Organic Chemistry, Part B: Reactions and Synthesis*, 5th edition. 2007. New York. Springer Sciences.)

Pheromone Compositions and Uses Thereof

As described above, products made via the methods described herein are pheromones. Pheromones prepared according to the methods of the invention can be formulated for use as insect control compositions. The pheromone compositions can include a carrier, and/or be contained in a dispenser. The carrier can be, but is not limited to, an inert liquid or solid.

Examples of solid carriers include but are not limited to fillers such as kaolin, bentonite, dolomite, calcium carbonate, talc, powdered magnesia, Fuller's earth, wax, gypsum, diatomaceous earth, rubber, plastic, China clay, mineral earths such as silicas, silica gels, silicates, attaclay, limestone, chalk, loess, clay, dolomite, calcium sulfate, magnesium sulfate, magnesium oxide, ground synthetic materials, fertilizers such as ammonium sulfate, ammonium phosphate, ammonium nitrate, thiourea and urea, products of vegetable origin such as cereal meals, tree bark meal, wood meal and nutshell meal, cellulose powders, attapulgites, montmorillonites, mica, vermiculites, synthetic silicas and synthetic calcium silicates, or compositions of these.

Examples of liquid carriers include, but are not limited to, water; alcohols, such as ethanol, butanol or glycol, as well as their ethers or esters, such as methylglycol acetate;

ketones, such as acetone, cyclohexanone, methylethyl ketone, methylisobutylketone, or isophorone; alkanes such as hexane, pentane, or heptanes; aromatic hydrocarbons, such as xylenes or alkyl naphthalenes; mineral or vegetable oils; aliphatic chlorinated hydrocarbons, such as trichloroethane or methylene chloride; aromatic chlorinated hydrocarbons, such as chlorobenzenes; water-soluble or strongly polar solvents such as dimethylformamide, dimethyl sulfoxide, or N-methylpyrrolidone; liquefied gases; waxes, such as beeswax, lanolin, shellac wax, carnauba wax, fruit wax (such as bayberry or sugar cane wax) candelilla wax, other waxes such as microcrystalline, ozocerite, ceresin, or montan; salts such as monoethanolamine salt, sodium sulfate, potassium sulfate, sodium chloride, potassium chloride, sodium acetate, ammonium hydrogen sulfate, ammonium chloride, ammonium acetate, ammonium formate, ammonium oxalate, ammonium carbonate, ammonium hydrogen carbonate, ammonium thiosulfate, ammonium hydrogen diphosphate, ammonium dihydrogen monophosphate, ammonium sodium hydrogen phosphate, ammonium thiocyanate, ammonium sulfamate or ammonium carbamate and mixtures thereof. Baits or feeding stimulants can also be added to the carrier.

Synergist

In some embodiments, the pheromone composition is combined with an active chemical agent such that a synergistic effect results. The synergistic effect obtained by the taught methods can be quantified according to Colby's formula (i.e. $(E) = X + Y - (X * Y / 100)$). See Colby, R. S., "Calculating Synergistic and Antagonistic Responses of Herbicide Combinations", 1967 Weeds, vol. 15, pp. 20-22, incorporated herein by reference in its entirety. Thus, by "synergistic" is intended a component which, by virtue of its presence, increases the desired effect by more than an additive amount. The pheromone compositions and adjuvants of the present methods can synergistically increase the effectiveness of agricultural active compounds and also agricultural auxiliary compounds.

Thus, in some embodiments, a pheromone composition can be formulated with a synergist. The term, "synergist," as used herein, refers to a substance that can be used with a pheromone for reducing the amount of the pheromone dose or enhancing the effectiveness of the pheromone for attracting at least one species of insect. The synergist may or may not be an independent attractant of an insect in the absence of a pheromone.

In some embodiments, the synergist is a volatile phytochemical that attracts at least one species of Lepidoptera. The term, "phytochemical," as used herein, means a compound occurring naturally in a plant species. In a particular embodiment, the synergist is selected from the group comprising β -caryophyllene, iso-caryophyllene, α -humulene, inalool, Z3-hexenol/yl acetate, β -farnesene, benzaldehyde, phenylacetaldehyde, and combinations thereof.

The pheromone composition can contain the pheromone and the synergist in a mixed or otherwise combined form, or it may contain the pheromone and the synergist independently in a non-mixed form.

Insecticide

The pheromone composition can include one or more insecticides. In one embodiment, the insecticides are chemical insecticides known to one skilled in the art. Examples of the chemical insecticides include one or more of pyrethroid or organophosphorus insecticides, including but are not limited to, cyfluthrin, permethrin, cypermethrin, bifenthrin, fenvalerate, flucythrinate, azinphosmethyl, methyl parathion, buprofezin, pyriproxyfen, flonicamid, acetamiprid,

dinotefuran, clothianidin, acephate, malathion, quinolphos, chloropyrifos, profenophos, bendiocarb, bifenthrin, chlorpyrifos, cyfluthrin, diazinon, pyrethrum, fenpropathrin, kinoprene, insecticidal soap or oil, neonicotinoids, diamides, avermectin and derivatives, spinosad and derivatives, azadirachtin, pyridalyl, and mixtures thereof.

In another embodiment, the insecticides are one or more biological insecticides known to one skilled in the art. Examples of the biological insecticides include, but are not limited to, azadirachtin (neem oil), toxins from natural pyrethrins, *Bacillus thuringiensis* and *Beauveria bassiana*, viruses (e.g., CYD-X™, CYD-X HP™, Germstar™, Madex HP™ and Spod-X™), peptides (Spear-T™, Spear-P™, and Spear-C™)

In another embodiment, the insecticides are insecticides that target the nerve and muscle. Examples include acetylcholinesterase (AChE) inhibitors, such as carbamates (e.g., methomyl and thiodicarb) and organophosphates (e.g., chlorpyrifos) GABA-gated chloride channel antagonists, such as cyclodiene organochlorines (e.g., endosulfan) and phenylpyrazoles (e.g., fipronil), sodium channel modulators, such as pyrethrins and pyrethroids (e.g., cypermethrin and λ -cyhalothrin), nicotinic acetylcholine receptor (nAChR) agonists, such as neonicotinoids (e.g., acetamiprid, tiacloprid, thiamethoxam), nicotinic acetylcholine receptor (nAChR) allosteric modulators, such as spinosyns (e.g., spinose and spinetoram), chloride channel activators, such as avermectins and milbemycins (e.g., abamectin, emamectin benzoate), Nicotinic acetylcholine receptor (nAChR) blockers, such as bensultap and cartap, voltage dependent sodium channel blockers, such as indoxacarb and metaflumizone, ryanodine receptor modulator, such as diamides (e.g. dhlorantraniliprole and flubendiamide). In another embodiment, the insecticides are insecticides that target respiration. Examples include chemicals that uncouple oxidative phosphorylation via disruption of the proton gradient, such as chlorfenapyr, and mitochondrial complex I electron transport inhibitors.

In another embodiment, the insecticides are insecticides that target midgut. Examples include microbial disruptors of insect midgut membranes, such as *Bacillus thuringiensis* and *Bacillus sphaericus*.

In another embodiment, the insecticides are insecticides that target growth and development. Examples include juvenile hormone mimics, such as juvenile hormone analogues (e.g. fenoxycarb), inhibitors of chitin biosynthesis, Type 0, such as benzoylureas (e.g., flufenoxuron, lufenuron, and novaluron), and ecdysone receptor agonists, such as diacylhydrazines (e.g., methoxyfenozide and tebufenozide)

Stabilizer

According to another embodiment of the disclosure, the pheromone composition may include one or more additives that enhance the stability of the composition. Examples of additives include, but are not limited to, fatty acids and vegetable oils, such as for example olive oil, soybean oil, corn oil, safflower oil, canola oil, and combinations thereof.

Filler

According to another embodiment of the disclosure, the pheromone composition may include one or more fillers. Examples of fillers include, but are not limited to, one or more mineral clays (e.g., attapulgit). In some embodiments, the attractant-composition may include one or more organic thickeners. Examples of such thickeners include, but are not limited to, methyl cellulose, ethyl cellulose, and any combinations thereof.

Solvent

According to another embodiment, the pheromone compositions of the present disclosure can include one or more solvents. Compositions containing solvents are desirable when a user is to employ liquid compositions which may be applied by brushing, dipping, rolling, spraying, or otherwise applying the liquid compositions to substrates on which the user wishes to provide a pheromone coating (e.g., a lure). In some embodiments, the solvent(s) to be used is/are selected so as to solubilize, or substantially solubilize, the one or more ingredients of the pheromone composition. Examples of solvents include, but are not limited to, water, aqueous solvent (e.g., mixture of water and ethanol), ethanol, methanol, chlorinated hydrocarbons, petroleum solvents, turpentine, xylene, and any combinations thereof.

In some embodiments, the pheromone compositions of the present disclosure comprise organic solvents. Organic solvents are used mainly in the formulation of emulsifiable concentrates, ULV formulations, and to a lesser extent granular formulations. Sometimes mixtures of solvents are used. In some embodiments, the present disclosure teaches the use of solvents including aliphatic paraffinic oils such as kerosene or refined paraffins. In other embodiments, the present disclosure teaches the use of aromatic solvents such as xylene and higher molecular weight fractions of C9 and C10 aromatic solvents. In some embodiments, chlorinated hydrocarbons are useful as co-solvents to prevent crystallization when the formulation is emulsified into water. Alcohols are sometimes used as co-solvents to increase solvent power.

Solubilizing Agent

In some embodiments, the pheromone compositions of the present disclosure comprise solubilizing agents. A solubilizing agent is a surfactant, which will form micelles in water at concentrations above the critical micelle concentration. The micelles are then able to dissolve or solubilize water-insoluble materials inside the hydrophobic part of the micelle. The types of surfactants usually used for solubilization are non-ionics: sorbitan monooleates; sorbitan monooleate ethoxylates; and methyl oleate esters.

Binder

According to another embodiment of the disclosure, the pheromone composition may include one or more binders. Binders can be used to promote association of the pheromone composition with the surface of the material on which said composition is coated. In some embodiments, the binder can be used to promote association of another additive (e.g., insecticide, insect growth regulators, and the like) to the pheromone composition and/or the surface of a material. For example, a binder can include a synthetic or natural resin typically used in paints and coatings. These may be modified to cause the coated surface to be friable enough to allow insects to bite off and ingest the components of the composition (e.g., insecticide, insect growth regulators, and the like), while still maintaining the structural integrity of the coating.

Non-limiting examples of binders include polyvinylpyrrolidone, polyvinyl alcohol, partially hydrolyzed polyvinyl acetate, carboxymethylcellulose, starch, vinylpyrrolidone/vinyl acetate copolymers and polyvinyl acetate, or compositions of these; lubricants such as magnesium stearate, sodium stearate, talc or polyethylene glycol, or compositions of these; antifoams such as silicone emulsions, long-chain alcohols, phosphoric esters, acetylene diols, fatty acids or organofluorine compounds, and complexing agents such as: salts of ethylenediaminetetraacetic acid (EDTA), salts of trinitrilotriacetic acid or salts of polyphosphoric acids, or compositions of these.

In some embodiments, the binder also acts a filler and/or a thickener. Examples of such binders include, but are not limited to, one or more of shellac, acrylics, epoxies, alkyds, polyurethanes, linseed oil, tung oil, and any combinations thereof.

Surface-Active Agents

In some embodiments, the pheromone compositions comprise surface-active agents. In some embodiments, the surface-active agents are added to liquid agricultural compositions. In other embodiments, the surface-active agents are added to solid formulations, especially those designed to be diluted with a carrier before application. Thus, in some embodiments, the pheromone compositions comprise surfactants. Surfactants are sometimes used, either alone or with other additives, such as mineral or vegetable oils as adjuvants to spray-tank mixes to improve the biological performance of the pheromone on the target. The surface-active agents can be anionic, cationic, or nonionic in character, and can be employed as emulsifying agents, wetting agents, suspending agents, or for other purposes. In some embodiments, the surfactants are non-ionics such as: alkyl ethoxylates, linear aliphatic alcohol ethoxylates, and aliphatic amine ethoxylates. Surfactants conventionally used in the art of formulation and which may also be used in the present formulations are described, in *McCutcheon's Detergents and Emulsifiers Annual*, MC Publishing Corp., Ridge-wood, N.J., 1998, and in *Encyclopedia of Surfactants*, Vol. I-III, Chemical Publishing Co., New York, 1980-81. In some embodiments, the present disclosure teaches the use of surfactants including alkali metal, alkaline earth metal or ammonium salts of aromatic sulfonic acids, for example, ligno-, phenol-, naphthalene- and dibutyl-naphthalenesulfonic acid, and of fatty acids of arylsulfonates, of alkyl ethers, of lauryl ethers, of fatty alcohol sulfates and of fatty alcohol glycol ether sulfates, condensates of sulfonated naphthalene and its derivatives with formaldehyde, condensates of naphthalene or of the naphthalenesulfonic acids with phenol and formaldehyde, condensates of phenol or phenolsulfonic acid with formaldehyde, condensates of phenol with formaldehyde and sodium sulfite, polyoxyethylene octylphenyl ether, ethoxylated isooctyl-, octyl- or nonylphenol, tributylphenyl polyglycol ether, alkylaryl polyether alcohols, isotridecyl alcohol, ethoxylated castor oil, ethoxylated triarylphenols, salts of phosphated triarylphenolethoxylates, lauryl alcohol polyglycol ether acetate, sorbitol esters, lignin-sulfite waste liquors or methylcellulose, or compositions of these.

In some embodiments, the present disclosure teaches other suitable surface-active agents, including salts of alkyl sulfates, such as diethanolammonium lauryl sulfate; alkylarylsulfonate salts, such as calcium dodecylbenzenesulfonate; alkylphenol-alkylene oxide addition products, such as nonylphenol-C18 ethoxylate; alcohol-alkylene oxide addition products, such as tridecyl alcohol-C16 ethoxylate; soaps, such as sodium stearate; alkyl-naphthalene-sulfonate salts, such as sodium dibutyl-naphthalenesulfonate; dialkyl esters of sulfosuccinate salts, such as sodium di(2-ethyl-hexyl)sulfosuccinate; sorbitol esters, such as sorbitol oleate; quaternary amines, such as lauryl trimethylammonium chloride; polyethylene glycol esters of fatty acids, such as polyethylene glycol stearate; block copolymers of ethylene oxide and propylene oxide; salts of mono and dialkyl phosphate esters; vegetable oils such as soybean oil, rapeseed/canola oil, olive oil, castor oil, sunflower seed oil, coconut oil, corn oil, cottonseed oil, linseed oil, palm oil, peanut oil, safflower oil, sesame oil, tung oil and the like; and esters of the above vegetable oils, particularly methyl esters.

Wetting Agents

In some embodiments, the pheromone compositions comprise wetting agents. A wetting agent is a substance that when added to a liquid increases the spreading or penetration power of the liquid by reducing the interfacial tension between the liquid and the surface on which it is spreading. Wetting agents are used for two main functions in agrochemical formulations: during processing and manufacture to increase the rate of wetting of powders in water to make concentrates for soluble liquids or suspension concentrates; and during mixing of a product with water in a spray tank or other vessel to reduce the wetting time of wettable powders and to improve the penetration of water into water-dispersible granules. In some embodiments, examples of wetting agents used in the pheromone compositions of the present disclosure, including wettable powders, suspension concentrates, and water-dispersible granule formulations are: sodium lauryl sulphate; sodium dioctyl sulphosuccinate; alkyl phenol ethoxylates; and aliphatic alcohol ethoxylates.

Dispersing Agent

In some embodiments, the pheromone compositions of the present disclosure comprise dispersing agents. A dispersing agent is a substance which adsorbs onto the surface of particles and helps to preserve the state of dispersion of the particles and prevents them from reaggregating. In some embodiments, dispersing agents are added to pheromone compositions of the present disclosure to facilitate dispersion and suspension during manufacture, and to ensure the particles redispense into water in a spray tank. In some embodiments, dispersing agents are used in wettable powders, suspension concentrates, and water-dispersible granules. Surfactants that are used as dispersing agents have the ability to adsorb strongly onto a particle surface and provide a charged or steric barrier to re-aggregation of particles. In some embodiments, the most commonly used surfactants are anionic, non-ionic, or mixtures of the two types.

In some embodiments, for wettable powder formulations, the most common dispersing agents are sodium lignosulphonates. In some embodiments, suspension concentrates provide very good adsorption and stabilization using polyelectrolytes, such as sodium naphthalene sulphonate formaldehyde condensates. In some embodiments, tristyrilphenol ethoxylated phosphate esters are also used. In some embodiments, such as alkylarylethylene oxide condensates and EO-PO block copolymers are sometimes combined with anionics as dispersing agents for suspension concentrates.

Polymeric Surfactant

In some embodiments, the pheromone compositions of the present disclosure comprise polymeric surfactants. In some embodiments, the polymeric surfactants have very long hydrophobic 'backbones' and a large number of ethylene oxide chains forming the 'teeth' of a 'comb' surfactant. In some embodiments, these high molecular weight polymers can give very good long-term stability to suspension concentrates, because the hydrophobic backbones have many anchoring points onto the particle surfaces. In some embodiments, examples of dispersing agents used in pheromone compositions of the present disclosure are: sodium lignosulphonates; sodium naphthalene sulphonate formaldehyde condensates; tristyrilphenol ethoxylate phosphate esters; aliphatic alcohol ethoxylates; alky ethoxylates; EO-PO block copolymers; and graft copolymers.

Emulsifying Agent

In some embodiments, the pheromone compositions of the present disclosure comprise emulsifying agents. An emulsifying agent is a substance, which stabilizes a suspension of droplets of one liquid phase in another liquid phase.

Without the emulsifying agent the two liquids would separate into two immiscible liquid phases. In some embodiments, the most commonly used emulsifier blends include alkylphenol or aliphatic alcohol with 12 or more ethylene oxide units and the oil-soluble calcium salt of dodecylbenzene sulphonic acid. A range of hydrophile-lipophile balance (“HLB”) values from 8 to 18 will normally provide good stable emulsions. In some embodiments, emulsion stability can sometimes be improved by the addition of a small amount of an EO-PO block copolymer surfactant.

Gelling Agent

In some embodiments, the pheromone compositions comprise gelling agents. Thickeners or gelling agents are used mainly in the formulation of suspension concentrates, emulsions, and suspoemulsions to modify the rheology or flow properties of the liquid and to prevent separation and settling of the dispersed particles or droplets. Thickening, gelling, and anti-settling agents generally fall into two categories, namely water-insoluble particulates and water-soluble polymers. It is possible to produce suspension concentrate formulations using clays and silicas. In some embodiments, the pheromone compositions comprise one or more thickeners including, but not limited to: montmorillonite, e.g. bentonite; magnesium aluminum silicate; and attapulgite. In some embodiments, the present disclosure teaches the use of polysaccharides as thickening agents. The types of polysaccharides most commonly used are natural extracts of seeds and seaweeds or synthetic derivatives of cellulose. Some embodiments utilize xanthan and some embodiments utilize cellulose. In some embodiments, the present disclosure teaches the use of thickening agents including, but are not limited to: guar gum; locust bean gum; carrageenan; alginates; methyl cellulose; sodium carboxymethyl cellulose (SCMC); hydroxyethyl cellulose (HEC). In some embodiments, the present disclosure teaches the use of other types of anti-settling agents such as modified starches, polyacrylates, polyvinyl alcohol, and polyethylene oxide. Another good anti-settling agent is xanthan gum.

Anti-Foam Agent

In some embodiments, the presence of surfactants, which lower interfacial tension, can cause water-based formulations to foam during mixing operations in production and in application through a spray tank. Thus, in some embodiments, in order to reduce the tendency to foam, anti-foam agents are often added either during the production stage or before filling into bottles/spray tanks. Generally, there are two types of anti-foam agents, namely silicones and non-silicones. Silicones are usually aqueous emulsions of dimethyl polysiloxane, while the nonsilicone anti-foam agents are water-insoluble oils, such as octanol and nonanol, or silica. In both cases, the function of the anti-foam agent is to displace the surfactant from the air-water interface.

Preservative

In some embodiments, the pheromone compositions comprise a preservative.

Additional Active Agent

According to another embodiment of the disclosure, the pheromone composition may include one or more insect feeding stimulants. Examples of insect feeding stimulants include, but are not limited to, crude cottonseed oil, fatty acid esters of phytol, fatty acid esters of geranyl geraniol, fatty acid esters of other plant alcohols, plant extracts, and combinations thereof.

According to another embodiment of the disclosure, the pheromone composition may include one or more insect growth regulators (“IGRs”). IGRs may be used to alter the

growth of the insect and produce deformed insects. Examples of insect growth regulators include, for example, dimilin.

According to another embodiment of the disclosure, the attractant-composition may include one or more insect sterilants that sterilize the trapped insects or otherwise block their reproductive capacity, thereby reducing the population in the following generation. In some situations allowing the sterilized insects to survive and compete with non-trapped insects for mates is more effective than killing them outright.

Sprayable Compositions

In some embodiments, the pheromone compositions disclosed herein can be formulated as a sprayable composition (i.e., a sprayable pheromone composition). An aqueous solvent can be used in the sprayable composition, e.g., water or a mixture of water and an alcohol, glycol, ketone, or other water-miscible solvent. In some embodiments, the water content of such mixture is at least about 10%, at least about 20%, at least about 30%, at least about 40%, 50%, at least about 60%, at least about 70%, at least about 80%, or at least about 90%. In some embodiments, the sprayable composition is concentrate, i.e. a concentrated suspension of the pheromone, and other additives (e.g., a waxy substance, a stabilizer, and the like) in the aqueous solvent, and can be diluted to the final use concentration by addition of solvent (e.g., water).

In some embodiments, the a waxy substance can be used as a carrier for the pheromone and its positional isomer in the sprayable composition. The waxy substance can be, e.g., a biodegradable wax, such as bees wax, carnauba wax and the like, candelilla wax (hydrocarbon wax), montan wax, shellac and similar waxes, saturated or unsaturated fatty acids, such as lauric, palmitic, oleic or stearic acid, fatty acid amides and esters, hydroxylic fatty acid esters, such as hydroxyethyl or hydroxypropyl fatty acid esters, fatty alcohols, and low molecular weight polyesters such as polyalkylene succinates.

In some embodiments, a stabilizer can be used with the sprayable pheromone compositions. The stabilizer can be used to regulate the particle size of concentrate and/or to allow the preparation of a stable suspension of the pheromone composition. In some embodiments, the stabilizer is selected from hydroxylic and/or ethoxylated polymers. Examples include ethylene oxide and propylene oxide copolymer, polyalcohols, including starch, maltodextrin and other soluble carbohydrates or their ethers or esters, cellulose ethers, gelatin, polyacrylic acid and salts and partial esters thereof and the like. In other embodiments, the stabilizer can include polyvinyl alcohols and copolymers thereof, such as partly hydrolyzed polyvinyl acetate. The stabilizer may be used at a level sufficient to regulate particle size and/or to prepare a stable suspension, e.g., between 0.1% and 15% of the aqueous solution.

In some embodiments, a binder can be used with the sprayable pheromone compositions. In some embodiments, the binder can act to further stabilize the dispersion and/or improve the adhesion of the sprayed dispersion to the target locus (e.g., trap, lure, plant, and the like). The binder can be polysaccharide, such as an alginate, cellulose derivative (acetate, alkyl, carboxymethyl, hydroxyalkyl), starch or starch derivative, dextrin, gum (arabic, guar, locust bean, tragacanth, carrageenan, and the like), sucrose, and the like. The binder can also be a non-carbohydrate, water-soluble polymer such as polyvinyl pyrrolidone, or an acidic polymer such as polyacrylic acid or polymethacrylic acid, in acid and/or salt form, or mixtures of such polymers.

Microencapsulated Pheromones

In some embodiments, the pheromone compositions disclosed herein can be formulated as a microencapsulated pheromone, such as disclosed in Ill'ichev, A L et al., *J. Econ. Entomol.* 2006; 99(6):2048-54; and Stelinki, L L et al., *J. Econ. Entomol.* 2007; 100(4):1360-9. Microencapsulated pheromones (MECs) are small droplets of pheromone enclosed within polymer capsules. The capsules control the release rate of the pheromone into the surrounding environment, and are small enough to be applied in the same method as used to spray insecticides. The effective field longevity of the microencapsulated pheromone formulations can range from a few days to slightly more than a week, depending on inter alia climatic conditions, capsule size and chemical properties.

Slow-Release Formulation

Pheromone compositions can be formulated so as to provide slow release into the atmosphere, and/or so as to be protected from degradation following release. For example, the pheromone compositions can be included in carriers such as microcapsules, biodegradable flakes and paraffin wax-based matrices. Alternatively, the pheromone composition can be formulated as a slow release sprayable.

In certain embodiments, the pheromone composition may include one or more polymeric agents known to one skilled in the art. The polymeric agents may control the rate of release of the composition to the environment. In some embodiments, the polymeric attractant-composition is impervious to environmental conditions. The polymeric agent may also be a sustained-release agent that enables the composition to be released to the environment in a sustained manner.

Examples of polymeric agents include, but are not limited to, celluloses, proteins such as casein, fluorocarbon-based polymers, hydrogenated rosins, lignins, melamine, polyurethanes, vinyl polymers such as polyvinyl acetate (PVAC), polycarbonates, polyvinylidene dinitrile, polyamides, polyvinyl alcohol (PVA), polyamide-aldehyde, polyvinyl aldehyde, polyesters, polyvinyl chloride (PVC), polyethylenes, polystyrenes, polyvinylidene, silicones, and combinations thereof. Examples of celluloses include, but are not limited to, methylcellulose, ethyl cellulose, cellulose acetate, cellulose acetate-butyrate, cellulose acetate-propionate, cellulose propionate, and combinations thereof.

Other agents which can be used in slow-release or sustained-release formulations include fatty acid esters (such as a sebacate, laurate, palmitate, stearate or arachidate ester) or a fatty alcohols (such as undecanol, dodecanol, tridecanol, tridecenol, tetradecanol, tetradecenol, tetradecadienol, pentadecanol, pentadecenol, hexadecanol, hexadecenol, hexadecadienol, octadecanol and octadecadienol).

Pheromones prepared according to the methods of the invention, as well as compositions containing the pheromones, can be used to control the behavior and/or growth of insects in various environments. The pheromones can be used, for example, to attract or repel male or female insects to or from a particular target area. The pheromones can be used to attract insects away from vulnerable crop areas. The pheromones can also be used example to attract insects as part of a strategy for insect monitoring, mass trapping, lure/attract-and-kill or mating disruption.

Lures

The pheromone compositions of the present disclosure may be coated on or sprayed on a lure, or the lure may be otherwise impregnated with a pheromone composition.

Traps

The pheromone compositions of the disclosure may be used in traps, such as those commonly used to attract any insect species, e.g., insects of the order Lepidoptera. Such traps are well known to one skilled in the art, and are commonly used in many states and countries in insect eradication programs. In one embodiment, the trap includes one or more septa, containers, or storage receptacles for holding the pheromone composition. Thus, in some embodiments, the present disclosure provides a trap loaded with at least one pheromone composition. Thus, the pheromone compositions of the present disclosure can be used in traps for example to attract insects as part of a strategy for insect monitoring, mass trapping, mating disruption, or lure/attract and kill for example by incorporating a toxic substance into the trap to kill insects caught.

Mass trapping involves placing a high density of traps in a crop to be protected so that a high proportion of the insects are removed before the crop is damaged. Lure/attract-and-kill techniques are similar except once the insect is attracted to a lure, it is subjected to a killing agent. Where the killing agent is an insecticide, a dispenser can also contain a bait or feeding stimulant that will entice the insects to ingest an effective amount of an insecticide. The insecticide may be an insecticide known to one skilled in the art. The insecticide may be mixed with the attractant-composition or may be separately present in a trap. Mixtures may perform the dual function of attracting and killing the insect.

Such traps may take any suitable form, and killing traps need not necessarily incorporate toxic substances, the insects being optionally killed by other means, such as drowning or electrocution. Alternatively, the traps can contaminate the insect with a fungus or virus that kills the insect later. Even where the insects are not killed, the trap can serve to remove the male insects from the locale of the female insects, to prevent breeding.

It will be appreciated by a person skilled in the art that a variety of different traps are possible. Suitable examples of such traps include water traps, sticky traps, and one-way traps. Sticky traps come in many varieties. One example of a sticky trap is of cardboard construction, triangular or wedge-shaped in cross-section, where the interior surfaces are coated with a non-drying sticky substance. The insects contact the sticky surface and are caught. Water traps include pans of water and detergent that are used to trap insects. The detergent destroys the surface tension of the water, causing insects that are attracted to the pan, to drown in the water. One-way traps allow an insect to enter the trap but prevent it from exiting. The traps of the disclosure can be colored brightly, to provide additional attraction for the insects.

In some embodiments, the pheromone traps containing the composition may be combined with other kinds of trapping mechanisms. For example, in addition to the pheromone composition, the trap may include one or more fluorescent lights, one or more sticky substrates and/or one or more colored surfaces for attracting moths. In other embodiments, the pheromone trap containing the composition may not have other kinds of trapping mechanisms.

The trap may be set at any time of the year in a field. Those of skill in the art can readily determine an appropriate amount of the compositions to use in a particular trap, and can also determine an appropriate density of traps/acre of crop field to be protected.

The trap can be positioned in an area infested (or potentially infested) with insects. Generally, the trap is placed on or close to a tree or plant. The aroma of the pheromone attracts the insects to the trap. The insects can then be

caught, immobilized and/or killed within the trap, for example, by the killing agent present in the trap.

Traps may also be placed within an orchard to overwhelm the pheromones emitted by the females, so that the males simply cannot locate the females. In this respect, a trap need be nothing more than a simple apparatus, for example, a protected wickable to dispense pheromone.

The traps of the present disclosure may be provided in made-up form, where the compound of the disclosure has already been applied. In such an instance, depending on the half-life of the compound, the compound may be exposed, or may be sealed in conventional manner, such as is standard with other aromatic dispensers, the seal only being removed once the trap is in place.

Alternatively, the traps may be sold separately, and the compound of the disclosure provided in dispensable format so that an amount may be applied to trap, once the trap is in place. Thus, the present disclosure may provide the compound in a sachet or other dispenser.

Dispenser

Pheromone compositions can be used in conjunction with a dispenser for release of the composition in a particular environment. Any suitable dispenser known in the art can be used. Examples of such dispensers include but are not limited to, aerosol emitters, hand-applied dispensers, bubble caps comprising a reservoir with a permeable barrier through which pheromones are slowly released, pads, beads, tubes rods, spirals or balls composed of rubber, plastic, leather, cotton, cotton wool, wood or wood products that are impregnated with the pheromone composition. For example, polyvinyl chloride laminates, pellets, granules, ropes or spirals from which the pheromone composition evaporates, or rubber septa. One of skill in the art will be able to select suitable carriers and/or dispensers for the desired mode of application, storage, transport or handling.

In another embodiment, a device may be used that contaminates the male insects with a powder containing the pheromone substance itself. The contaminated males then fly off and provide a source of mating disruption by permeating the atmosphere with the pheromone substance, or by attracting other males to the contaminated males, rather than to real females.

Behavior Modification

Pheromone compositions prepared according to the methods disclosed herein can be used to control or modulate the behavior of insects. In some embodiments, the behavior of the target insect can be modulated in a tunable manner inter alia by varying the ratio of the pheromone to the positional isomer in the composition such that the insect is attracted to a particular locus but does not contact said locus or such the insect in fact contacts said locus. Thus, in some embodiments, the pheromones can be used to attract insects away from vulnerable crop areas. Accordingly, the disclosure also provides a method for attracting insects to a locus. The method includes administering to a locus an effective amount of the pheromone composition.

The method of mating disruption may include periodically monitoring the total number or quantity of the trapped insects. The monitoring may be performed by counting the number of insects trapped for a predetermined period of time such as, for example, daily, Weekly, bi-Weekly, monthly, once-in-three months, or any other time periods selected by the monitor. Such monitoring of the trapped insects may help estimate the population of insects for that particular period, and thereby help determine a particular type and/or dosage of pest control in an integrated pest management system. For example, a discovery of a high insect population

can necessitate the use of methods for removal of the insect. Early warning of an infestation in a new habitat can allow action to be taken before the population becomes unmanageable. Conversely, a discovery of a low insect population can lead to a decision that it is sufficient to continue monitoring the population. Insect populations can be monitored regularly so that the insects are only controlled when they reach a certain threshold. This provides cost-effective control of the insects and reduces the environmental impact of the use of insecticides.

Mating Disruption

Pheromones prepared according to the methods of the disclosure can also be used to disrupt mating. Mating disruption is a pest management technique designed to control insect pests by introducing artificial stimuli (e.g., a pheromone composition as disclosed herein) that confuses the insects and disrupts mating localization and/or courtship, thereby preventing mating and blocking the reproductive cycle.

In many insect species of interest to agriculture, such as those in the order Lepidoptera, females emit an airborne trail of a specific chemical blend constituting that species' sex pheromone. This aerial trail is referred to as a pheromone plume. Males of that species use the information contained in the pheromone plume to locate the emitting female (known as a "calling" female). Mating disruption exploits the male insects' natural response to follow the plume by introducing a synthetic pheromone into the insects' habitat, which is designed to mimic the sex pheromone produced by the female insect. Thus, in some embodiments, the synthetic pheromone utilized in mating disruption is a synthetically derived pheromone composition comprising a pheromone having a chemical structure of a sex pheromone and a positional isomer thereof which is not produced by the target insect.

The general effect of mating disruption is to confuse the male insects by masking the natural pheromone plumes, causing the males to follow "false pheromone trails" at the expense of finding mates, and affecting the males' ability to respond to "calling" females. Consequently, the male population experiences a reduced probability of successfully locating and mating with females, which leads to the eventual cessation of breeding and collapse of the insect infestation.

Strategies of mating disruption include confusion, trail-masking and false-trail following. Constant exposure of insects to a high concentration of a pheromone can prevent male insects from responding to normal levels of the pheromone released by female insects. Trail-masking uses a pheromone to destroy the trail of pheromones released by females. False-trail following is carried out by laying numerous spots of a pheromone in high concentration to present the male with many false trails to follow. When released in sufficiently high quantities, the male insects are unable to find the natural source of the sex pheromones (the female insects) so that mating cannot occur.

In some embodiments, a wick or trap may be adapted to emit a pheromone for a period at least equivalent to the breeding season(s) of the midge, thus causing mating disruption. If the midge has an extended breeding season, or repeated breeding season, the present disclosure provides a wick or trap capable of emitting pheromone for a period of time, especially about two weeks, and generally between about 1 and 4 weeks and up to 6 weeks, which may be rotated or replaced by subsequent similar traps. A plurality of traps containing the pheromone composition may be placed in a locus, e.g., adjacent to a crop field. The locations

109

of the traps, and the height of the traps from ground may be selected in accordance with methods known to one skilled in the art.

Alternatively, the pheromone composition may be dispensed from formulations such as microcapsules or twist-
ties, such as are commonly used for disruption of the mating of insect pests.

Attract and Kill

The attract and kill method utilizes an attractant, such as a sex pheromone, to lure insects of the target species to an insecticidal chemical, surface, device, etc., for mass-killing and ultimate population suppression, and can have the same effect as mass-trapping. For instance, when a synthetic female sex pheromone is used to lure male pests, e.g., moths, in an attract-and-kill strategy, a large number of male moths must be killed over extended periods of time to reduce matings and reproduction, and ultimately suppress the pest population. The attract-and-kill approach may be a favorable alternative to mass-trapping because no trap-servicing or other frequent maintenance is required. In various embodiments described herein, a recombinant microorganism can co-express (i) a pathway for production of an insect pheromone and (ii) a protein, peptide, oligonucleotide, or small molecule which is toxic to the insect. In this way, the recombinant microorganism can co-produce substances suitable for use in an attract-and-kill approach.

As will be apparent to one of skill in the art, the amount of a pheromone or pheromone composition used for a particular application can vary depending on several factors such as the type and level of infestation; the type of

110

composition used; the concentration of the active components; how the composition is provided, for example, the type of dispenser used; the type of location to be treated; the length of time the method is to be used for; and environmental factors such as temperature, wind speed and direction, rainfall and humidity. Those of skill in the art will be able to determine an effective amount of a pheromone or pheromone composition for use in a given application.


As used herein, an "effective amount" means that amount of the disclosed pheromone composition that is sufficient to affect desired results. An effective amount can be administered in one or more administrations. For example, an effective amount of the composition may refer to an amount of the pheromone composition that is sufficient to attract a given insect to a given locus. Further, an effective amount of the composition may refer to an amount of the pheromone composition that is sufficient to disrupt mating of a particular insect population of interest in a given locality.

EXAMPLES


Prophetic Example 1. Production of Pheromones Products from Enzymatically-Derived Gondoic Acid Through Metathesis and Chemical Conversion

This prophetic example illustrates that different fatty acids can be used as a starting material for the biosynthetic production of a pheromone or pheromone precursor. The product obtained from the biosynthetic process disclosed herein can be subject to further chemical conversions to generate different products.

111



112



Enzymatic two carbon elongation of oleic acid yields gondoic acid. After esterification, gondoic fatty acid methyl ester (FAME) can then be converted via Z-selective olefin metathesis into C16 and C18 FAME products containing a C11 unsaturation. Upon reduction of the ester, aldehyde and fatty alcohol pheromone materials can be produced. Acetylation of the fatty alcohol product can generate the corresponding fatty acetate pheromones. Additionally, gondoic acid can be directly converted into C20 fatty aldehyde, alcohol and acetate pheromones through application of the same chemical transformation of enzymatically modified oleic acid.

Prophetic Example 2: Tailored Synthetic Blends

This prophetic example illustrates that the recombinant microorganisms disclosed herein can be used to create synthetic blends of insect pheromones.

As shown in the scheme in FIG. 40, using tetradecyl-ACP (14:ACP), a blend of E- and Z-tetradecenyl acetate (E11-14:OAc and Z11-14:OAc) pheromones can be produced with the recombinant microorganism. This blend is produced by a variety of insects, e.g., *Choristoneura roseceana* (a moth of the Tortricidae family).

Similarly, using hexadecyl-ACP (16:ACP), a blend of Z- and E hexadecenyl acetate pheromones (E11-16:OAc and Z11-16:OAc) can be produced with the recombinant microorganism.

The microorganism can be engineered with different desaturases, or other enzymes such as reductases, etc. to produce the desired blend of pheromones. One blend of particular relevance capable of being produced using the recombinant microorganisms and methods of the instant invention is a 97:3 ratio of (Z)-11-hexadecenal (Z11-16:Ald) and (Z)-9-hexadecenal (Z9-16:Ald).

Example 3: Expression of Transmembrane Alcohol-Forming Reductases in *S. cerevisiae*

Background and Rationale

Engineering microbial production of insect fatty alcohols from fatty acids entails the functional expression of a synthetic pathway. One such pathway comprises a transmembrane desaturase, and an alcohol-forming reductase to mediate the conversion of fatty acyl-CoA into regio- and stereospecific unsaturated fatty acyl-CoA, and subsequently into fatty alcohols. A number of genes encoding these enzymes are found in some insects (as well as some microalgae in the case of fatty alcohol reductase) and can be used to construct the synthetic pathway in yeasts, which are preferred production hosts. A number of transmembrane desaturases and alcohol-forming reductase variants will be screened to identify ensembles which allow high level synthesis of a single insect fatty alcohol or a blend of fatty alcohols. Additionally, these enzymes will be screened across multiple hosts (*Saccharomyces cerevisiae*, *Candida tropicalis*, and *Yarrowia lipolytica*) to optimize the search toward finding a suitable host for optimum expression of these transmembrane proteins.

Summary of Approach

Three alcohol-forming reductases of insect origin were selected.

Nucleic acids encoding the reductases were synthesized (synthons) with codon optimization for expression in *S. cerevisiae*.

Each nucleic acid encoding a given reductase was subcloned into an episomal expression cassette under the Gall promoter.

S. cerevisiae wild-type and beta-oxidation deletion mutant were transformed with expression constructs.

Heterologous protein was induced by galactose, and functional expression of the reductases was assessed in vivo via bioconversion of Z11-hexadecenoic acid into Z11-hexadecanol.

GC-MS analysis was used to identify and quantify metabolites.

Results

Alcohol-forming reductase variants were screened for activity in *S. cerevisiae* W303 (wild type) and BY4742 ΔPOX1 (beta-oxidation deletion mutant). Z11-hexadecenoic acid was chosen as a substrate in assessing enzyme activity. The in vivo bioconversion assay showed that the expression of enzyme variants derived from *Spodoptera littoralis*, *Helicoverpa armigera*, and *Agrotis segetum* (Ding, B-J., Löfstedt, C. Analysis of the *Agrotis segetum* pheromone gland transcriptome in the light of sex pheromone biosynthesis. BMC Genomics 16:711 (2015)) in W303A conferred Z11-hexadecanol production, and reached up-to ~37 μM (8 mg/L), ~70 μM (~16 mg/L), and 11 μM (~3 mg/L), respectively, within 48 h of protein induction (FIG. 5 and FIG. 6). Biologically-produced Z11-hexadecanol matched authentic Z11-hexadecanol standard (Bedoukian) as determined via GC-MS (FIG. 7). BY4742 ΔPOX1 was also explored as an expression host since deletion in the key beta-oxidation pathway enzyme could limit the degradation of Z11-hexadecenoic acid. Expressing the reductase variants in the beta-oxidation deletion mutant, however, reduced the product titer when compared to expression in the wild-type host (FIG. 5). One contributing factor of titer reduction when using BY4742 ΔPOX1 as a host was the reduction of biomass when compared to W303 (FIG. 8).

Therefore, functional expression of at least two alcohol-forming reductases in *S. cerevisiae* conferred bioconversion of Z11-hexadecenoic acid into Z11-hexadecanol.

Conclusions

Functional expression of insect transmembrane alcohol-forming reductase in *S. cerevisiae* was demonstrated. Among the reductases tested, the variant derived from *Helicoverpa armigera* is most active toward Z11-hexadecenoic acid.

The bioconversion of other fatty acid substrates can be explored to assess enzyme plasticity.

Materials & Methods

Strain Construction and Functional Expression Assay

S. cerevisiae W303 (MATA ura3-1 trp1-1 leu2-3_112 his3-11_15 ade2-1 can1-100) and BY4742 (MATa POX1::kanMX his3Δ1 leu2Δ0 lys2Δ0 ura3Δ0) were used as expression hosts. DNA sequences which encode fatty alcohol reductase variants were redesigned to optimize expression in *S. cerevisiae* (SEQ ID NOs: 1-3). Generated synthons (GenScript) were cloned into pESC-URA vector using BamHI-XhoI sites to facilitate protein expression utilizing the Gall promoter. The resulting plasmid constructs were used to transform W303, and positive transformants were selected on CM agar medium (with 2% glucose, and lacking uracil) (Teknova). To assess functional expression, two positive transformation clones that have been patched on CM agar medium (with 2% glucose, and lacking uracil) were used to seed CM liquid medium using a 24 deep-well plate format. To induce protein expression, the overnight cultures that had been grown at 28° C. were then supplemented with galactose, raffinose, and YNB to a final concentration of 2%, 1%,

and 6.7 g/L, respectively. Post 24 h of protein induction, the bioconversion substrate Z11-hexadecenoic acid (in ethanol) or heptadecanoic acid (in ethanol) was added to a final concentration of 300 mg/L. Bioconversion assay proceeded for 48 h at 28° C. prior to GC-MS analysis.

Metabolite Extraction and GC-MS Detection

The lipids were extracted according to a modified procedure of Hagström et al. (2012) (Hagström, Å. K., Liénard, M. A., Groot, A. T., Hedenström, E. & Löfstedt, C. Semi-Selective Fatty Acyl Reductases from Four Heliothine Moths Influence the Specific Pheromone Composition. *PLoS One* 7: e37230 (2012)). 1.5 mL-cell culture was transferred to a 15 mL falcon tube. The cell suspension was acidified with 1 mL 5 N HCl. 5 µL tetradecanedioic acid (10 mM in ethanol) was added as internal standard. The mixture was extracted by adding 1.5 mL hexane, then shaken for 1 h at 37° C., 250 rpm. To facilitate phase separation, the sample was centrifuged for 10 min at 2000 g. 1 mL of the organic hexane phase was then transferred to a 1.5 mL plastic tube. The solvent was removed by heating the sample 30 min at 90° C. After the sample was evaporated to dryness, 50 µL of BSTFA (N,O-bis(trimethylsilyl) trifluoroacetamide containing 1% of trimethylchlorosilane) was added. The 1.5 mL plastic tubes were shaken vigorously two times for 10 s. Prior to the transfer into a screw cap GC glass vial containing a glass insert, the sample was centrifuged for 1 min (13000 rpm). The vials were capped and heated for 30 min at 90° C. The trimethylsilyl-esters, which were generated by this method were subsequently analyzed by GC-MS analysis. GC-MS parameters are specified in Table 6. The use of SIM mode (characteristic product and IS ions) increases detection sensitivity by reducing background noise, allowing detection of the product as low as 2.4 µM (0.6 mg/L). A further reduction in the split ratio offers the possibility to further increase the sensitivity for future applications. A Z11-hexadecenol calibration curve shown in FIG. 9 was used to quantify the Z11-hexadecenol produced from yeasts. The bioconversion of heptadecanoic acid was also tested since the easily distinguished heptadecanol product could be used to benchmark successful GC-MS runs. However, none of the reductase tested showed any activity toward heptadecanoic acid.

TABLE 6

GC-MS parameters	
System	Agilent 6890 N GC, ChemStation G1701EA E.02.01.1177
Column	Rtx-5 30 m × 320 µm × 25 µm Pressure = 11.74 psi; Flow = 7.1 mL/min
Inlet	Heater = 250° C.; Pressure = 11.74 psi; Total Flow {He} = 19.5 mL/min
Carrier	He @ 147 cm/sec, 11.74 psi
Signal	Data rate = 2 Hz/0.1 min
Oven	150° C. for 1 min Ramp 12° C./min to 220° C., hold 3 min Ramp 35° C./min to 300° C., hold 4 min
Injection	Split, 250° C. Split ratio - 20:1
Detector	HP 5973 MSD in SIM mode (m/z: 297.3 and 387.3), 100 msec Dwell, EMV mode: Gain factor 1, 3 min solvent delay, 8.33 cycles/sec
Sample	Injection volume = 1 µL

Example 4: Expression of Transmembrane Desaturases in *S. cerevisiae*

Background and Rationale

Engineering microbial production of insect fatty alcohols from fatty acids requires the functional expression of a

synthetic pathway. One such pathway comprises a trans-membrane desaturase, and an alcohol-forming reductase to mediate the conversion of fatty acyl-CoA into regio- and stereospecific unsaturated fatty acyl-CoA, and subsequently into fatty alcohols. A number of genes encoding these enzymes are found in some insects as well as some microalgae. A number of transmembrane desaturases and alcohol-forming reductase variants will be screened to identify ensembles which allow high level synthesis of a single insect fatty alcohol or a blend of fatty alcohols. Additionally, these enzymes will be screened across multiple hosts (*Saccharomyces cerevisiae*, *Candida tropicalis*, and *Yarrowia lipolytica*) to optimize the search toward finding a suitable host for optimum expression of these transmembrane proteins.

Summary of Approach

A small set of desaturases (insect origin: *Agrotis segetum*, *Trichoplusia ni*, *Amyelois transitella*, *Helicoverpa zea*, and marine diatom: *Thalassiosira pseudonana*) were selected as a test case to explore and establish functional expression assays, metabolite extraction methods, and analytical chemistry.

A synthetic cassette for expression of the desaturases in *S. cerevisiae* was constructed. The cassette consists of the OLE1 promoter region, OLE1 N-terminal leader sequence, and VSP13 terminator.

The expression cassette was tested for functionality via expression of a GFP variant. Validation of the cassette allowed its utilization for exploring expression of insect desaturase.

S. cerevisiae ΔOLE1 was transformed with expression constructs containing heterologous desaturases. Functionality of the desaturases was assessed via the ability to rescue growth of ΔOLE1 without exogenous supplementation of unsaturated fatty acid (UFA). *S. cerevisiae* desaturase (OLE1) was used as a positive control of successful complementation.

Functionality of the desaturase was validated via an in vivo bioconversion of hexadecanoic acid (palmitic acid) into (Z)-11-hexadecenoic acid (palmitivaccenic acid).

GC-MS analysis was used to identify and quantify metabolites.

Results

Transmembrane desaturase variants were screened in *S. cerevisiae*. Three variants were initially tested to explore and establish functional expression assays, metabolite extraction methods, and analytical chemistry. To allow functional expression of these desaturases in *S. cerevisiae*, an episomal synthetic expression cassette termed pOLE1 cassette (FIG. 10) was constructed, which consisted of an OLE1 promoter region, an N-terminal leader sequence encoding for the first 27 amino acids of *S. cerevisiae* OLE1, and a terminator region of VPS13 (a protein involved in the protospore membrane formation, the terminator of which has been previously characterized to increase heterologous protein expression potentially by extending mRNA half-life). The functionality of the pOLE1 cassette was validated via its ability to express a GFP (FIG. 11A-FIG. 11E). Subsequently, insect desaturase synthons, and yeast OLE1 synthon were cloned into the pOLE1 cassette, and expressed in *S. cerevisiae* ΔOLE1 strain. This strain was chosen since deletion of the OLE1 allele (which encodes for palmitoyl:CoA/stearoyl:CoA (z)-9-desaturase) allows its utilization as a tool to screen for functional insect desaturase. Specifically, an active desaturase would allow complementation of growth

without requiring exogenous supplementation of UFAs. Expression of OLE1 using pOLE1 cassette complemented growth of Δ OLE1 growth without UFA (FIG. 11A-FIG. 11E); therefore, it serves as a positive control in the complementation assays. When insect desaturases were expressed, we observed that they rescued Δ OLE1 growth without UFA at varying degree. On rich medium (YPD) agar plate, expression of *S. cerevisiae* OLE1 conferred the highest level of growth, followed by *T. ni* desaturase (FIG. 12A). The latter indicated that production of unsaturated fatty acyl:CoA by *T. ni* desaturase could act as a surrogate to the missing (Z)-9-hexadecenoyl:CoA biosynthesis in Δ OLE1. Expression of *T. pseudonana* and *A. segetum* desaturases did not appear to rescue growth on YPD very well (FIG. 12A). When patched on minimal medium (CM-Ura glucose) agar plate, only expression of *S. cerevisiae* OLE1 and *T. ni* desaturase rescued Δ OLE1 growth without exogenous UFA (FIG. 12B). Expression of *T. pseudonana* and *A. segetum* desaturases did not confer growth of Δ OLE1 on minimal medium agar, suggesting their limited activity in producing UFA (results not shown). Screening a desaturase library in *Candida tropicalis* identified functional expression of *A. transitella* and *H. zea* desaturases. When these desaturases were expressed in Δ OLE1, they conferred growth without UFA on both YPD and CM-Ura glucose media similar to expression of *T. ni* desaturase (FIG. 12B).

Functional expression of the heterologous desaturases was further characterized via in vivo bioconversion of palmitic acid into insect-specific UFA. Post ~96 h-cultivation in minimal medium containing palmitic acid, total fatty acid analysis of *S. cerevisiae* Δ OLE1 expressing *T. ni* desaturase revealed production of a new fatty acid species (Z)-11-hexadecenoic acid that is not present in the control strain which expresses native yeast OLE1 desaturase (FIG. 13A-FIG. 13B). (Z)-11-hexadecenoic acid is not detected in strains expressing *A. segetum*, or *T. pseudonana* desaturase (results not shown). In addition to (Z)-11-hexadecenoic acid, (Z)-9-hexadecenoic acid was also detected in Δ OLE1 strain expressing *T. ni* desaturase (FIG. 13A-FIG. 13B). Under the cultivation condition, C16-fatty acid in the Δ OLE1 expressing *T. ni* desaturase is composed of approximately 84.7% hexadecanoic acid, 5.6% (Z)-9-hexadecenoic acid and 9.8% (Z)-11-hexadecenoic acid. In comparison, the C16 fatty acid fraction of Δ OLE1 expressing OLE1 desaturase is composed of approximately 68.6% hexadecanoic acid and 31.4% (Z)-9-hexadecenoic acid. (Z)-11-hexadecenoic acid biosynthesis in Δ OLE1 expressing *T. ni* desaturase account for ~1.5 mg/L. The amount of total fatty acids and each fatty acid within this mixture can be quantified. The biologically produced (Z)-11-hexadecenoic acid also match the retention time and fragmentation pattern of authentic standard (Z)-11-hexadecenoic acid (Larodan) as determined by GC-MS (FIG. 14A-FIG. 14B). Therefore, the regio- and stereoisomer of the biologically produced (Z)-11-hexadecenoic acid was confirmed. In vivo characterization of *A. transitella* and *H. zea* desaturase can also be done.

In summary, at least three insect desaturases capable of rescuing growth of *S. cerevisiae* Δ OLE1 without exogenous supplementation of UFA, i.e. (Z)-9-hexadecenoic acid (palmitoleic acid), were identified.

The extent of growth on rich medium (YPD) of *S. cerevisiae* Δ OLE1 bearing the expression construct was in the following order of desaturase content: OLE1, *T. ni*, *T. pseudonana*, and *A. segetum*.

The extent of growth on minimal medium (CM Glucose w/out uracil) of *S. cerevisiae* Δ OLE1 bearing the expression construct was in the following order of desaturase content: OLE1, *T. ni*.

Complementation assays using *A. transitella* and *H. zea* desaturases were also done, demonstrating functional expression in *Candida tropicalis* shown via in vivo bioconversion assay. These desaturases also complemented *S. cerevisiae* Δ OLE1 growth on rich and minimal media at least as well as *T. ni* desaturase.

Expression of *T. pseudonana* and *A. segetum* desaturases did not confer growth of *S. cerevisiae* Δ OLE1 on minimal medium without UFAs even after an extended incubation period up to 14 days. No (Z)-11-hexadecenoic acid was observed in strains harboring *T. pseudonana* or *A. segetum* desaturase.

Conclusions

Functional expression of transmembrane desaturases of insect origin in *S. cerevisiae* has been achieved.

The activity of a given heterologous desaturase can be assessed from its ability to complement growth of *S. cerevisiae* Δ OLE1 without exogenous palmitoleic supplementation, and its ability to convert palmitic acid into insect pheromone precursors (Z)-11-hexadecenoic acid.

Functional expression and/or activity of insect desaturase in *S. cerevisiae* varies widely depending on sequence origin. Variants derived from *T. ni* exhibited the best activity compared to *A. segetum* and *T. pseudonana*, as measured by the above criteria.

Desaturases derived from *A. transitella* and *H. zea* complemented Δ OLE1 as well as *T. ni* desaturase. Bioconversion assays using these desaturases can be done.

The bioconversion of other fatty acid substrates can be explored to assess enzyme plasticity.

Materials & Methods

Strain Construction and Functional Expression Assay

S. cerevisiae Δ OLE1 (MATA OLE1::LEU2 ura3-52 his4) was used as an expression host. A synthetic expression cassette termed pOLE1 (FIG. 10, SEQ ID NO: 4) which comprises the OLE1 promoter region (SEQ ID NOs: 5 and 6), nucleotides encoding for 27 N-terminal amino acids of the OLE1 leader sequence (SEQ ID NO: 7), and a VPS13 terminator sequence (SEQ ID NO: 8) was created, and cloned into pESC-URA vector in between SacI and EcoRI sites. To test the functionality of the pOLE1 cassette, Dasher GFP synthon was inserted in between SpeI and NotI sites to create pOLE1-GFP plasmid. Competent Δ OLE1 was transformed with pOLE1-GFP, and plated on CM-Ura glucose agar plate (Teknova) containing UFA (20 mm CM-URA glucose agar plate was coated with 100 μ L CM-Ura glucose medium containing 1% tergitol, and 3 μ L palmitoleic acid). After incubation at 30° C. for 5 days, Dasher GFP expression was apparent as displayed by green coloration of Δ OLE1 transformants. This result showed that the pOLE1 cassette was capable of driving heterologous protein expression. Validation of Δ OLE1 complementation was performed by restoring OLE1 activity. Specifically, native *S. cerevisiae* OLE1 synthon was inserted into pOLE1 cassette devoid of the leader sequence to create pOLE1-OLE1 plasmid. After transformation of Δ OLE1, and selection on CM-Ura glucose agar containing UFA, single colonies were patched onto YPD and CM-Ura glucose without UFA. After incubation at 30° C. for 5 days, growth was observed (FIG. 11A-FIG. 11E). As expected, Dasher GFP expression could not complement Δ OLE1 growth without UFA (FIG. 11A-FIG. 11E). DNA sequences which encode for desaturase variants were synthesized (to include nucleotide changes which

remove restriction sites used for cloning purposes), and cloned into pOLE1 using SpeI-NotI sites (Genscript, SEQ ID NOs: 9-13). Complementation assay of Δ OLE1 with insect desaturases were performed in the same way as with OLE1 desaturase.

To assess functional expression, two positive transformation clones that had been patched on CM-Ura glucose agar medium containing UFA were inoculated in 1.5 mL CM-Ura glucose liquid medium containing palmitic acid (in ethanol) at a final concentration of 300 mg/L, and with 6.7 g/L of YNB. For (z)-11-hexadecenoic isomer confirmation, a 20 mL culture was generated. Bioconversion assay proceeded for 96 h at 28° C. prior to GC-MS analysis.

Metabolite Extraction and GC-MS Detection

Total lipid composition as well as the (Z)-11-hexadecenoic acid quantification was based on modified procedures by Moss et al. (1982) (Moss, C. W., Shinoda, T. & Samuels, J. W. Determination of cellular fatty acid compositions of various yeasts by gas-liquid chromatography. *J. Clin. Microbiol.* 16: 1073-1079 (1982)) and Yousuf et al (2010) (Yousuf, A., Sannino, F., Addorisio, V. & Pirozzi, D. Microbial Conversion of Olive Oil Mill Wastewaters into Lipids Suitable for Biodiesel Production. *J. Agric. Food Chem.* 58: 8630-8635 (2010)). The pelleted cells (in 1.5 mL plastic tubes), usually about 10 mg to 80 mg, were resuspended in methanol containing 5% (w/w) of sodium hydroxide. The alkaline cell suspension was transferred into a 1.8 mL screw-cap GC-vial. The mixture was heated for 1 h in the heat block at 90° C. Prior to acidification with 400 2.5 N HCl the vial was allowed to cool to room temperature. 500 μ L chloroform containing 1 mM heptadecanoic were added and the mixture was shaken vigorously, then both aqueous and organic phase were transferred into a 1.5 mL plastic tube. The mixture was centrifuged at 13,000 rpm, afterwards 450 μ L of the organic phase were transferred into a new 1.5 mL plastic tube. The aqueous phase was extracted a second time with 500 μ L chloroform, this time without heptadecanoic acid. The combined organic phases were evaporated at 90° C. After cooling to room temperature, residual fatty acid methyl esters and free fatty acids were dissolved and derivatized in methanol containing 0.2 M TMSH (trimethylsulfonium hydroxide).

The regioselectivity of biologically produced (Z)-11-hexadecenoic acid was determined by comparing the fragmentation patterns of the dimethyl disulfide (DMDS) derivative with the DMDS derivative of an authentic standard. A yeast culture was split into 12 aliquots (to not change any parameters in the developed procedure). The cells were pelleted, which yielded 63 mg cells (ccw) on average (755 mg from 18 mL culture). The pellets were subjected to base methanolysis as described above. However, after acidification the samples were combined in a 50 mL Falcon tube. The combined sample was extracted two times with 10 mL chloroform. The mixture was centrifuged 10 min at 3000 rpm to achieve a better phase separation. The combined organic phases, which were combined in a new 50 mL Falcon and were washed consecutively with 10 mL brine and 10 mL water. The organic phase was dried with anhydrous sodium sulfate and concentrated in vacuo. The concentrated oil was dissolved in 1.5 mL chloroform and transferred to a 1.5 mL plastic tube. The chloroform was evaporated at 90° C. The remaining sample was the dissolved in 50 μ L methyl tert-butyl ether (MTBE). The 50 μ L were split into 1, 5, 10 and 20 μ L and transferred into GC-vials without insert. To each vial 200 μ L DMDS (dimethyl disulfide) and 50 μ L MTBE (containing 60 mg/mL iodine) were added. After the mixture was heated 48 h at 50°

C., excess iodine was removed by the addition of 100 μ L saturated sodium thiosulfate solution. The samples were transferred to plastic vials and extracted to times with 500 μ L dichloromethane. The combined organic phases were transferred to a new 1.5 mL plastic vial and evaporated at 90° C. The samples were taken up in 50 μ L DCM and transferred to a GC-vial. The sample was analyzed by GC-MS (Table 7) using the method of Hagström et al. (2013) (Hagström, Å. K. et al. A moth pheromone brewery: production of (Z)-11-hexadecenol by heterologous co-expression of two biosynthetic genes from a noctuid moth in a yeast cell factory. *Microb. Cell Fact.* 12: 125 (2013)).

TABLE 7

Analytical parameters used for GC-MS analysis of DMDS-derivatives	
System	Agilent 6890 N GC, ChemStation G1701EA E.02.01.1177
Column	Rtx-5 30 m $\times$ 320 μ m $\times$ 25 μ m Pressure = 11.74 psi; Flow = 7.1 mL/min
Inlet	Heater = 250° C.; Pressure = 11.74 psi; Total Flow {He} = 19.5 mL/min
Carrier	He @ 147 cm/sec, 11.74 psi
Signal	Data rate = 2 Hz/0.1 min
Oven	80° C. for 2 min Ramp 10° C./min to 180° C. Ramp 3° C./min to 260° C. Ramp 20° C./min to 280° C., hold 10 min
Injection	Split, 250° C. Split ratio -1:1
Detector	HP 5973 MSD in SCAN mode (mass range: 41 to 550 amu) 100 msec Dwell, EMV mode: Gain factor 1, 3 min solvent delay, 8.33 cycles/sec
Sample	Injection volume = 1 μ L

Example 5: *S. cerevisiae* as a Production Platform for Insect Fatty Alcohol Synthesis

Background and Rationale

Engineering microbial production of insect fatty alcohols from fatty acids requires the functional expression of a synthetic pathway. One such pathway comprises a transmembrane desaturase, and an alcohol-forming reductase to mediate the conversion of fatty acyl-CoA into regio- and stereospecific unsaturated fatty acyl-CoA, and subsequently into fatty alcohols. A number of genes encoding these enzymes are found in some insects as well as some microalgae. A number of gene variants were screened to identify enzyme activities that allow the creation of pathways capable of high level synthesis of a single or a blend of insect fatty alcohols. Additionally, these enzymes were screened across multiple hosts (*Saccharomyces cerevisiae*, *Candida tropicalis*, and *Yarrowia lipolytica*) in order to find a suitable host for optimum expression of these transmembrane proteins.

Summary of Approach

S. cerevisiae was engineered previously to express select functional transmembrane desaturase variants to allow synthesis of (Z)-11-hexadecenoic acid from palmitic acid. This allowed the identification and rank-ordering of the variants based on their bioconversion performance (see Example 4).

S. cerevisiae was engineered previously to express select functional transmembrane reductase variants to allow synthesis of (Z)-11-hexadecenol (Z11-16OH) from (Z)-11-hexadecenoic acid. This allowed the identification and rank-ordering of the variants based on their bioconversion performance (see Example 3).

Several fatty alcohol pathways comprised of the most active variant desaturases and reductases identified in the previous screens were assembled.

S. cerevisiae W303A and Δ OLE1 were transformed with the pathway constructs. Functionality of the pathway was assessed via the ability of the recombinant yeasts to synthesize Z11-16OH from palmitic acid.

GC-MS analysis was used to identify and quantify metabolites.

Results

The goal was to engineer one or more insect fatty alcohol biosynthetic pathways in *S. cerevisiae*. Previously, the functional expression of several transmembrane desaturases of insect origin in *S. cerevisiae* was demonstrated (see Example 4). Briefly, heterologous desaturase expression was enabled by designing an expression cassette which consists of an OLE1 promoter region, an N-terminal leader sequence encoding the first 27 amino acids of *S. cerevisiae* OLE1, and a terminator region of VPS13. Screening for active desaturases was done by using two approaches. First, active desaturases were screened for their ability to rescue Δ OLE1 growth without exogenous addition of unsaturated fatty acid (UFA), and second, active desaturases were screened via an in vivo screen for bioconversion of palmitic acid into (Z)-11-hexadecenoic acid. These screening strategies allowed the identification of several active variants, and the rank ordering of their relative activity. Based on these screening results, desaturases from *Trichoplusia ni* (TN_desat) and *S. cerevisiae* (SC_desat) were selected for combinatorial expression in fatty alcohol pathways. *S. cerevisiae* desaturase is known to form palmitoleic acid and oleic acid.

The functional expression of several transmembrane alcohol forming reductases of insect origin in *S. cerevisiae* had also been previously demonstrated (see Example 3). An expression cassette comprising the GAL1 promoter and CYC terminator was used to enable the functional expression of the reductases in *S. cerevisiae*. Screening several reductases via in vivo bioconversion of (Z)-11-hexadecenoic acid into Z11-16OH allowed the identification of active variants and rank ordering of their relative activity. Based on this screen, reductases from *Helicoverpa armigera* (HA_reduc), and *Spodoptera littoralis* (SL_reduc) were chosen for assembly of the fatty alcohol pathways.

Combinatorial assembly created four fatty alcohol pathways, i.e. TN_desat-HA_reduc, TN_desat-SL_reduc, SC_desat-HA_reduc, and SC_desat-SL_reduc. Pathways with SC_desat served as negative control for insect Z11-16OH synthesis. *S. cerevisiae* Δ OLE1 and W303A were transformed with constructs harboring these pathways, and transformants that grew on CM-Ura with 2% glucose and coated with palmitoleic acid were isolated. To test for fatty alcohol production, individual clones were inoculated into CM-Ura medium containing 2% glucose, 1% raffinose, 2% galactose, 300 mg/L palmitic acid, and 360 mg/L palmitoleic acid were added as bioconversion substrates. Bioconversion using palmitic acid without palmitoleic was also tested. Post ~96h-cultivation in the presence of palmitic and palmitoleic acid, culture broth analysis revealed synthesis of Z11-9OH as a major C16 alcohol product at ~0.2 mg/L, and ~0.3 mg/L in cultivation of Δ OLE1 strains harboring SC_desat-HA_reduc, and TN_desat-HA_reduc, respectively (FIG. 15, FIG. 16). A minute amount of Z11-16OH was also detected in pathways with *T. ni* or *S. cerevisiae* desaturase, and *H. armigera* reductase. In general, it was expected that in the presence of palmitic acid and palmitoleic acid, Z9-16OH synthesis was more favorable than Z11-16OH synthesis because (Z)-11-hexadecenoic acid must be biosynthesized

from *T. ni* desaturase, whereas exogenous addition of palmitoleic acid resulted in a more readily available substrate for synthesis of Z9-16OH. Fatty acid analysis was also performed. The results showed higher accumulation of (Z)-11-hexadecenoic acid (FIG. 17) in pathways containing insect desaturase than in pathways expressing *S. cerevisiae* desaturase. Albeit at minute quantities, detection of Z11-16OH, and (Z)-11-hexadecenoic acid from pathways harboring *S. cerevisiae* desaturase (which was unexpected) opens the possibility of a minor Δ 11 desaturation activity by *S. cerevisiae* desaturase. Low level synthesis of Z11-16COOH fatty acid moieties can also be derived from elongation of Z9-14COOH fatty acyl intermediate. The data shown in FIG. 15 also showed that in comparison to pathways with *H. armigera* reductase, the inclusion of *S. littoralis* reductase resulted in the reduction of (up-to ~30 fold) in Z9-16OH titer. No Z11-16OH could be detected in pathways employing *S. littoralis* reductase. These results are consistent with the reductase screening assay, which showed superior bioconversion of (Z)-11-hexadecenoic acid using *H. armigera* reductase in comparison to *S. littoralis* reductase.

The bioconversion of palmitic acid was also tested alone (without exogenous addition of palmitoleic acid) by Δ OLE1 strains expressing TN_desat-HA_reduc and TN_desat-SL_reduc (FIG. 18). Culture broth analysis determined the synthesis of Z11-16OH as the dominant unsaturated C16 fatty acid product (FIG. 16). In this assay, up to 0.22 mg/L, and 0.05 mg/L Z11-16OH was synthesized by a pathway harboring *H. armigera* reductase and *S. littoralis* reductase, respectively. The biologically produced Z11-16OH also matched the retention time and exhibited the characteristic 297.3 m/z peak like the authentic standard Z11-16OH as determined by GC-MS (SIM). Therefore, the regio- and stereoisomer of the biologically produced Z11-16OH was confirmed (FIG. 19). Furthermore, Z9-16OH (0.01 mg/L) was also observed in the cultivation of strain co-expressing *T. ni* desaturase and *H. armigera* reductase. This suggested that *T. ni* desaturase may also possess A9 desaturation activity.

OLE1 deletion impairs growth. Therefore, pathway expression was also explored in W303A, a host with intact OLE1 allele. However, despite growth improvement, pathway expression in this host resulted in more than two-fold reduction of Z11-16OH titers. This result was likely due to the repression of OLE1 promoter (which drove heterologous desaturase expression) by endogenous unsaturated fatty acyl:CoAs, the products of OLE1. The *S. cerevisiae* OLE1 promoter has been previously characterized with structural regions found to be positively and negatively regulated by saturated and unsaturated fatty acid, respectively (Choi, J-Y. et al. Regulatory Elements That Control Transcription Activation and Unsaturated Fatty Acid-mediated Repression of the *Saccharomyces cerevisiae* OLE1 Gene. J. Biol. Chem. 271: 3581-3589 (1996)). In addition to cis-transcriptional regulation, unsaturated fatty acids also interact with OLE1 promoter elements to regulate mRNA stability (Gonzales, C. I. et al. Fatty acid-responsive control of mRNA stability. Unsaturated fatty acid-induced degradation of the *Saccharomyces* OLE1 transcript. J. Biol. Chem. 271: 25801-25809 (1996)). Due to this inherent complexity of the OLE1 promoter, the utilization of unregulated orthogonal promoters, such as the OLE1 promoter from *S. kluyveri* (Kajiwara, S. Molecular cloning and characterization of the v9 fatty acid desaturase gene and its promoter region from *Saccharomyces kluyveri*. FEMS Yeast. Res. 2: 333-339 (2002)) to drive insect desaturase expression can be explored to enhance fatty alcohol production.

In summary, functional expression of synthetic pheromone pathway variants in *S. cerevisiae* Δ OLE1 resulted in the synthesis of Z11-16OH and Z9-16OH from palm oil fatty acids (palmitic acid and palmitoleic acid) up to approximately 0.2 mg/L and 0.3 mg/L, respectively.

The engineered pathway that resulted in the highest fatty alcohols is comprised of *T. ni* desaturase and *H. armigera* reductase.

Accumulation of (Z)-11-hexadecenoic acid, an intermediate of the pathway, was also observed in strains that produced Z11-16OH.

No Z11-16OH was produced and only trace Z9-16OH was detected in the negative control strain (harboring vector only).

The regio- and stereochemistry of the biologically produced Z11-16OH were confirmed by comparing the retention time and fragmentation pattern to the authentic standard compound via GC-MS.

Conclusions

The engineering of Baker's yeast for synthesis of Z11-16OH and Z9-16OH, fatty alcohol precursors of insect pheromones, was demonstrated.

Fatty alcohol production varies depending on the selection of the desaturase and reductase variants.

Accumulation of (Z)-11-hexadecenoic acid suggested the possibility of further fatty alcohol improvement by increasing the performance of alcohol forming reductase. However, it is also possible that detection of (Z)-11-hexadecenoic acid was due to its incorporation as phospholipid into any membrane other than the endoplasmic reticulum membrane (such as mitochondrial membranes, peroxisome, nuclear envelope, etc), therefore inaccessible to alcohol forming reductase (presumably translocated into the endoplasmic reticulum) which must utilize (Z)-11-hexadecenoic acid in its CoA thioester moiety as its substrate.

Culture conditions can be explored to increase fatty alcohol titers. The *T. ni* desaturase can be replaced in the pathway by *A. transitella* desaturase, another variant that also showed high activity and rescued Δ OLE1 growth faster than *T. ni* desaturase. The synthetic pathway can be imported into *Candida tropicalis* and *Yarrowia lipolytica*, which are yeasts with high adhesion property to hydrophobic substrates such as palmitic and palmitoleic acid. By increasing substrate accessibility to the microbial production platform, it is foreseeable that product titer and yield can be improved.

Materials & Methods

Strain Construction and Functional Expression Assay

S. cerevisiae Δ OLE1 (MATA OLE1::LEU2 ura3-52 his4), and W303A (MATA ura3-1 trp1-1 leu2-3_112 his3-11_15 ade2-1 can1-100) were used as expression hosts. Modular design allows combinatorial pathway assembly utilizing BamHI and XhoI to excise reductase synthons (see Example 3) and subcloning into plasmids containing pOLE1-desaturase constructs (see Example 4). Competent yeasts were transformed with pathway constructs and plated on CM-Ura glucose agar plate (Teknova). In the case of Δ OLE1 transformation, colony plating utilized 20 mM CM-Ura glucose agar plates that were coated with 100 μ L CM-Ura glucose medium containing 1% tergitol and 3 μ L palmitoleic acid.

To assess functional expression, transformants were inoculated in 20 mL CM-Ura liquid medium containing 6.7 g/L of YNB, 2% glucose, 1% raffinose, and 2% galactose. Fatty acid substrates, i.e. palmitic acid (in ethanol), was added at a final concentration of 300 mg/L. Palmitoleic acid was added at a final concentration of 360 mg/L. Bioconversion assay proceeded for 96 h at 28° C. prior to GC-MS analysis.

Metabolite Extraction and GC-MS Detection

Fatty acid analysis was as described in Example 4, except that instead of extracting the sample two times, the sample was only extracted once with chloroform containing 1 mM methyl heptadecanoate (C17:0Me). Fatty alcohol analysis was as described in Example 3, except that instead of hexane (containing tetradecanedioic acid), chloroform (containing 1 mM methyl heptadecanoate) was used. The extraction time was reduced from 1 h to 20 s. Afterwards the samples were collected in a 1.8 mL GC vial and not in a 1.5 mL plastic tube. The mass spectrometer was used in SIM mode (m/z 208, 297.3 and 387.3).

Example 6: Expression of Transmembrane Desaturases in *Candida tropicalis*

Background and Rationale

Engineering microbial production of insect fatty alcohols from fatty acids requires the functional expression of a synthetic pathway. One such pathway comprises a transmembrane desaturase, and an alcohol-forming reductase to mediate the conversion of fatty acyl-CoA into regio- and stereospecific unsaturated fatty acyl-CoA, and subsequently into fatty alcohols. A number of genes encoding these enzymes are found in some insects as well as some microalgae. A number of gene variants were screened to identify enzyme activities that allow the creation of pathways capable of high level synthesis of a single or a blend of insect fatty alcohols. Additionally, these enzymes can be screened across multiple hosts (*Saccharomyces cerevisiae*, *Candida tropicalis*, and *Yarrowia lipolytica*) to optimize the search toward finding a suitable host for optimum expression of these transmembrane proteins.

Summary of Approach

A small set of desaturases (insect origin: *Agrotis segetum*, *Amyelois transitella*, *Helicoverpa zea*, *Trichoplusia ni*, *Ostrinia furnacalis*, and *Lampronia capitella* and marine diatom: *Thalassiosira pseudonana*) were selected as a test case to explore and establish functional expression assays, metabolite extraction methods, and analytical chemistry.

Successful integration and functional expression of mCherry control from pXICL expression cassette in SPV053 were confirmed.

A recombinant desaturase library using the same pXICL vector in SPV053 background was integrated (FIG. 20). One variant, the Z11 desaturase of *Agrotis segetum*, was also cloned to produce a protein product with the first 27 amino acids of *Candida albicans* Ole1p fused to the N-terminus of the insect desaturase (SEQ ID NO: 15).

Functionality of the desaturase was validated via an in vivo bioconversion of hexadecanoic acid (palmitic acid) into (Z)-11-hexadecenoic acid (palmitvaccenic acid).

GC-FID and GC-MS analyses were used to identify and quantify metabolites.

Results

Library Construction

This study focused on the screening for transmembrane desaturase variants in *C. tropicalis* (SPV053). Five insect desaturases with reported Z11 desaturase activity on palmitoyl-CoA (C16:0) (SEQ ID NOs: 16-19, 23) and three insect desaturases with reported Z9 desaturase activity (SEQ ID NOs: 20-22) were included in the screen. One variant, the Z11 desaturase from *A. segetum* (SEQ ID NO: 16), was also cloned with 27 amino acids of the *Candida albicans* OLE1 N-terminus fused upstream of the insect sequence (FIG. 20, SEQ ID NO: 15). At the time of construction, the *A. segetum* Z11 desaturase was believed to be a positive control and the

C. albicans OLE1 fusion was constructed to test if inclusion of a *Candida* leader sequence would improve functional expression. The construct was designed to mimic those used in *Saccharomyces cerevisiae* desaturase screening (See Example 4). Finally, a control construct expressing mCherry red fluorescent protein (SEQ ID NO: 14) was included to act as a positive control for integration and expression and a negative control for recombinant desaturase activity (FIG. 21A-FIG. 21D).

Transformation efficiencies of linearized plasmids into SPV053 varied greatly across constructs. Despite low efficiencies, at least 3 clonal isolates were identified for each variant (Tables 8 and 9). It had been hypothesized that larger colonies on transformation plates were more likely to be positive integrants because the presence of the Zeocin resistance marker should increase growth rate under Zeocin selection. Analysis of the screening results suggested that the number of large colonies is not correlated to transformation efficiency. Instead total colony (small and large) count correlated best with observed efficiency (FIG. 22). In addition, in some cases positive clones were found among the small colonies. It is possible that at lower plating density growth rate may be correlated with integration events (i.e. positive integrants grow faster). A secondary screen of repatching colonies on YPD+Zeocin proved effective in enriching for positive integrants. Fast growing patches were more likely to be positive integrants than the general population of colonies on transformation plates.

TABLE 8

Desaturase transformations in SPV053. Efficiency of transformation varied across constructs with a relatively high degree of background under Zeocin selection.					
specificity	source species	pXICL plasmid	DNA ug	large colonies (control plate)	total colonies (control plate)
control	mCherry_Ct	pPV0137	1.1	60 (30)	2,000 (600)
	<i>Argotis segetum</i> -OLE1_Ca	pPV0138	1.2	120 (78)	>10,000 (320)
	<i>Agrotis segetum</i>	pPV0139	1.3	115 (78)	8,000 (320)
	<i>Amyelois transitella</i>	pPV0140	1.1	220 (78)	5,000 (320)
	<i>Trichoplusia ni</i>	pPV0141	1.1	100 (78)	>10,000 (320)
Z9	<i>Helicoverpa zea</i>	pPV0142	1.0	350 (78)	5,000 (320)
	<i>Thalassiosira pseudonana</i>	pPV0146	1.1	140 (78)	1,500 (320)
	<i>Ostrina Furnacalis</i>	pPV0143	0.9	220 (78)	6,000 (320)
	<i>Lampronia capitella</i>	pPV0144	1.2	230 (78)	5,000 (320)
	<i>Helicoverpa zea</i>	pPV0145	1.2	72 (78)	2,000 (320)

TABLE 9

Desaturase SPV053 library construction. Five insect desaturases with putative Z11 desaturation activity and 3 insect desaturases with putative Z9 desaturation activity were integrated into the SPV053 background using the pXICL vector. In addition, a control strain expressing mCherry was constructed with the same vector.					
specificity	source species	pXICL plasmid	Total positives	Total screened	Fraction positive
control	mCherry_Ct	pPV0137	16	16	1.00
	<i>Argotis segetum</i> -OLE1_Ca	pPV0138	7	12	0.58
	<i>Agrotis segetum</i>	pPV0139	12	12	1.00
	<i>Amyelois transitella</i>	pPV0140	5	60	0.08
	<i>Trichoplusia ni</i>	pPV0141	6	12	0.50
Z9	<i>Helicoverpa zea</i>	pPV0142	5	120	0.04
	<i>Thalassiosira pseudonana</i>	pPV0146	3	96	0.03
	<i>Ostrina</i>	pPV0143	3	57	0.05
	<i>Furnacalis</i>				

TABLE 9-continued

Desaturase SPV053 library construction. Five insect desaturases with putative Z11 desaturation activity and 3 insect desaturases with putative Z9 desaturation activity were integrated into the SPV053 background using the pXICL vector. In addition, a control strain expressing mCherry was constructed with the same vector.					
speci- ficity	source species	pXICL plasmid	Total positives	Total screened	Fraction positive
	<i>Lampronia capitella</i>	pPV0144	3	58	0.05
	<i>Helicoverpa zea</i>	pPV0145	3	94	0.03

Functional Expression Assay

Functional expression of the heterologous desaturases was characterized by a series of in vivo bioconversion experiments. *C. tropicalis* SPV053 derived stains expressing insect desaturases were cultured in rich (YPD) or defined (CM glucose) media supplemented with ethanol (for induction) and saturated acid substrates (palmitic acid, methyl palmitate, methyl myristate). Small scale (2 ml) cultures were cultivated for a total of 72 hours in 24 deep well plates with substrate added after the initial 24 hours.

The first screen examined multiple bioconversion media with supplementation of a palmitic acid substrate. Two

functional palmitoyl-CoA (Z)-11 desaturases were identified by fatty acid methyl ester (FAME) analysis of the cellular lipid content. Strains expressing *A. transitella* or *H. zea* Z11 desaturases (SPV0305-SPV0310) produced a fatty acid species not observed in the mCherry control strains (SPV0302-SPV0304) which eluted with the (Z)-11-hexadecenoic acid standard (FIG. 23). No other tested strains produced non-native fatty acid species (data not shown). Approximate fatty acid composition of the C16-fraction is listed in Table 10. The native palmitoyl-CoA (Z)-9 desaturase is still present in the SPV053 background which means the (Z)-9/(Z)-11 specificity of the desaturases cannot be rigorously determined. Supplementation of palmitic acid in the media increased the (Z)-11/(Z)-9 hexadecenoic acid ratio from 0.6 to 1.4 for *H. zea* desaturase expressing strains. (Z)-11-hexadecenoic acid titers were observed to be approximately 5.62 mg/L for strains expressing *A. transitella* desaturase and 5.96 mg/L for strains expressing *H. zea* desaturase. Similar performance was observed with methyl palmitate supplementation (data not shown).

TABLE 10

Composition of the C16-fatty acid fraction in different <i>C. tropicalis</i> SPV053 expressing different desaturases.				
	C16:0 [%]	Z9-C16:1 [%]	Z11-C19:1 [%]	Z11/Z9 ratio
mCherry	72.9	27.1	0.0	0.0
Hzea-YPD_NS	50.0	30.9	19.1	0.6
Hzea-YPD	58.1	17.5	24.4	1.4
AT-YPD	55.5	14.5	30.0	2.1

*NS = no substrate (hexadecanoic acid) was added.

The bioconversion assay was scaled-up to 20 ml in shake flasks in order to generate enough biomass for additional characterization of the putative (Z)-11-hexadecenoic acid species. While the observed species eluted with the (Z)-11-hexadecenoic acid standard and independently of the (Z)-9-hexadecenoic acid standard, it was possible that a different fatty acid isomer (e.g. (E)-9-hexadecenoic acid) could have a similar retention time to (Z)-11-hexadecenoic acid. As different stereoisomers elute differently on the DB-23 the occurrence of (E)-11-hexadecenoic could be excluded. Final confirmation of (Z)-11-hexadecenoic acid production was completed by using mass spectroscopy detection of DMDS derivatized fatty acids to confirm the 11-regioselectivity. Using this derivatization technique (Z)-11 and (E)-11 isomers could in principle also be resolved. The fragmentation pattern of experimental samples could be matched to the (Z)-11-hexadecenoic acid standard (FIG. 24A-24E). Using this technique, production of the specific (Z)-11-hexadecenoic acid regio- and stereoisomer was confirmed for both *A. transitella* and *H. zea* desaturase expressing strains.

Finally, methyl myristate (C14:0) was tested as substrate for the entire desaturase library. A non-native fatty acid species which elutes between myristate (C14:0) and (Z)-9-tetradecenoic acid (Z9-C14:1) was observed in strains expressing either *A. transitella* or *H. zea* Z11 desaturases (FIG. 25A). It is hypothesized that this non-native species is (Z)-11-tetradecenoic acid, and this can be confirmed with an authentic standard. In addition, *A. segetum* Z11 desaturase, *O. furnacalis* Z9 desaturase, and *H. zea* Z9 desaturase all produced a shoulder peak which eluted just after the myristate (C14:0) peak (FIG. 25B). Other C14 derived species (e.g. tetradecanedioic acid) were observed in all strains. These results suggest that *A. transitella* and *H. zea* desaturases have some activity on myristoyl-CoA. Confirmation of unknown species and quantification is required to draw further conclusions about desaturase substrate specificity in vivo.

In summary, two desaturases from *Helicoverpa zea* (AAF81787) and from *Amyelois transitella* (JX964774), were expressed in SPV053 and conferred synthesis of (Z)-11-hexadecenoic acid from either endogenously produced or supplemented palmitic acid.

Functional expression of *H. zea* and *A. transitella* desaturases in *C. tropicalis* SPV053 was confirmed using an in vivo bioconversion assay in both rich (YPD) and defined (CM glucose) media. The active desaturases generated intracellular (Z)-11-hexadecenoic acid which was not observed in mCherry expressing control strains. C16-fatty acid composition of SPV053 expressing *H. zea* desaturase is approximately 50.0% hexadecanoic acid, 30.91% (Z)-9-hexadecenoic acid and 19.1% (Z)-11-hexadecenoic acid. With palmitic acid supplementation the composition is 58.1% hexadecanoic acid, 17.5% (Z)-9-hexadecenoic acid and 24.4% (Z)-11-hexadecenoic acid. The C16-fatty acid com-

position of SPV053 expressing *A. transitella* desaturase is 55.5% hexadecanoic acid, 14.5% (Z)-9-hexadecenoic acid and 30.0% (Z)-11-hexadecenoic acid. In comparison, SPV053 expressing mCherry produced a C16-fatty acid composition of approximately 72.9% hexadecanoic acid, 27.1% (Z)-9-hexadecenoic acid and no (Z)-11-hexadecenoic acid. (Z)-11-hexadecenoic acid was produced at approximately 5.5 mg/L in both strains expressing functional Z11 desaturases.

No (Z)-11-hexadecenoic acid was observed in strains harboring *T. ni*, *T. pseudonana*, or *A. segetum* desaturase.

No difference in fatty acid composition was observed for strains expressing Z9 insect desaturases from *H. zea*, *O. furnacalis*, or *L. capitella*.

The regio- and stereoisomer of the biologically produced (Z)-11-hexadecenoic acid were confirmed by comparing the retention time and fragmentation pattern of the authentic standard compound via GC-MS after DMDS derivatization.

Bioconversions of SPV053 expressing *A. transitella* and *H. zea* desaturases with supplementation of methyl myristate produced an unidentified metabolite not observed in the mCherry expressing negative control strain. The GC retention time of this metabolite is found between myristate (C14:0) and (Z)-9-tetradecenoic acid.

Conclusions

Functional expression of transmembrane desaturase of insect origin in *C. tropicalis* SPV053 has been achieved.

The active desaturases identified via screening in *C. tropicalis* also complemented OLE1 function when expressed in *S. cerevisiae* ΔOLE1 (See Example 4).

An in vivo assay can be used to assay desaturase activity in *C. tropicalis* for non-native fatty acid isomers (e.g. (Z)-11-hexadecenoic acid). Enhanced ratios of non-native fatty acids can be produced with supplementation of saturated acid substrates such as palmitic acid or methyl myristate.

Functional expression and/or activity of insect desaturases varies widely in *C. tropicalis* SPV053 depending on sequence origin. Similar to results observed in the *S. cerevisiae* screen (See Example 4), *A. segetum* and *T. pseudonana* variants did not produce detectable (Z)-11-hexadecenoic acid. Interestingly, *T. ni* desaturase also failed to produce detectable (Z)-11-hexadecenoic acid under assay conditions. Unlike in the *S. cerevisiae* assay, the *T. ni* expression construct did not include a chimeric OLE1 leader sequence.

The inclusion of the *C. albicans* OLE1 leader sequence on the functional *H. zea* variant and non-functional *T. ni* variant can be tested.

The functional expression of additional desaturase variants to identify C14-specific desaturases can be explored.

Expression of functional desaturase with reductase variants can be done and subsequent screen for unsaturated fatty alcohol production can be performed.

Materials & Methods

Strain Construction

A conservative approach was used for recoding of genes. Native sequences were unaltered except for replacement of CTG leucine codons with TTA. All genes were cloned into pPV0053 using NcoI and NotI restriction sites by Genscript. After transformation into *E. coli* NEB10β, plasmids were miniprep using the Zyppy Plasmid Miniprep Kit (Zymo Research, Irvine, Calif.). Plasmids were linearized by digestion with BsiWI (New England Biolabs, Ipswich, Mass.) before transformation into SPV053. After digestion, DNA was isolated using Clean and Concentrator Kit (Zymo Research, Irvine, Calif.). Approximately 1 μg of DNA was

transformed by electroporation. Instead of incubation with TE+100 mM lithium acetate+DTT, cells were incubated in only TE+100 mM lithium acetate for 2 hours. Positive integrants were found to be site-specific and genotyping was conducted by check PCR. A two-stage approach was adopted for further screening of low efficiency transformations. Approximately 60 colonies were re-patched on YPD+300 µg/ml Zeocin and grown overnight. The subset of patches which grew quickly (dense growth within 24 hours) were screened by colony PCR. The vast majority of rapid growing patches were identified as positive integrants.

Functional Expression Assay

Palmitic Acid Supplementation in YPD and CM Glucose

Positive isolates were re-patched onto YPD+300 µg/ml Zeocin and grown overnight and then stored at 4° C. Strains were inoculated from patch plates into 2 ml of YPD in 24 deep well plates (square well, pyramid bottom). Three positive clones were inoculated for each desaturase variant and the mCherry expressing control strain. Deep well plates were incubated at 30° C., 1000 rpm, and 80% humidity in the Infors HT Multitron Pro plate shaker for 24 hrs. After 24 hrs of incubation, cultures were split into equal 1 ml volumes to make two sets of identical plates. Both sets of plates were pelleted by centrifugation at 500×g. One set of plates was resuspended in 2 ml of YPD+0.3% (v/v) ethanol and the second set was resuspended in 2 ml of CM glucose+0.3% ethanol. Ethanol was added at this stage to induce recombinant enzyme expression from the ICL promoter. Cultures were incubated for another 24 hours under the same conditions before 300 mg/L palmitic acid was added to cultures from a 90 g/L stock solution in ethanol. The result was the addition of a fresh 0.3% ethanol in conjunction with the palmitic acid. A subset of strains was also cultured without palmitic acid addition. These cultures had 0.3% ethanol added instead. All cultures were incubated for an additional 24 hrs before a final addition of 0.3% ethanol. After another 24 hr period of incubation, 1.5 ml of each culture was harvested in 1.7 ml microcentrifuge tubes and pelleted. Supernatant was saved in fresh tubes and pellets were processed as described below. A subset of supernatant samples was also extracted to look for free acid in the extracellular medium.

Repeated Screening with Alternate Substrates

The mCherry control and confirmed positive variants were rescreened using both palmitic acid and methyl palmitate as substrates. The culturing was conducted as described above with equimolar (1.17 mM) amounts of substrate added from ethanol stock solutions (methyl palmitate 94 g/L stock, 313 mg/L final concentration). The same protocol was also repeated with the full panel of strains using an 84 g/L stock of methyl myristate (C14:0). The final concentration of substrate was again 1.17 mM.

Confirmation of (Z)-11-Hexadecenoic Acid Isomer

The in vivo bioconversion assay was scaled up for confirmation of (Z)-11-hexadecenoic acid synthesis. 2 ml YPD seed cultures of strains SPV0302, SPV0303, and SPV0304 (mCherry), SPV0304, SPV0305, and SPV0306 (*A. transitella* Z11 desaturase), and SPV0307, SPV0308, and SPV0309 (*H. zea* Z11 desaturase) were grown overnight at 30° C., 1000 rpm, 80% humidity in the Infors HT Multitron plate shaker. 200 µl of overnight culture from each of the three clonal isolates was pooled and inoculated into a single 125 ml baffled flask containing 20 ml YPD. The resulting three flasks were grown for 24 hrs at 30° C. and 250 rpm (Infors Flask shaker). Cultures were pelleted by centrifugation at 500×g and resuspended in 20 ml of YPD+0.3% (v/v) ethanol and returned to 125 ml baffled shake flasks. Cultures

were incubated for an additional 24 hours before addition of 300 mg/L palmitic acid in a 90 g/L stock in ethanol (221 µl per flask). After 24 hours of incubation another 0.3% (v/v) ethanol (221 µl) was added to each flask for sustained induction. Flasks were incubated for an additional 24 hours before cells were harvested for FAME analysis and DMDS derivatization.

Metabolite Extraction and GC-MS Detection

Total lipid composition as well as the (Z)-11-hexadecenoic acid quantification was based on modified procedures by Moss et al. (1982) and Yousuf et al (2010). The pelleted cells (in 1.5 mL plastic tubes), usually about 10 mg to 80 mg, were resuspended in methanol containing 5% (w/w) of sodium hydroxide. The alkaline cell suspension was transferred into a 1.8 mL screw-cap GC-vial. The mixture was heated for 1 h in the heat block at 90° C. Prior to acidification with 400 2.5 N HCl the vial was allowed to cool to room temperature. 500 µL chloroform containing 1 mM heptadecanoic were added and the mixture was shaken vigorously, then both aqueous and organic phase were transferred into a 1.5 mL plastic tube. The mixture was centrifuged at 13,000 rpm, afterwards 450 µL of the organic phase were transferred into a new 1.5 mL plastic tube. The aqueous phase was extracted a second time with 500 µL chloroform, this time without heptadecanoic acid. The combined organic phases were evaporated at 90° C. After cooling to room temperature, residual fatty acid methyl esters and free fatty acids were dissolved and derivatized in methanol containing 0.2 M TMSH (trimethylsulfonium hydroxide).

The regioselectivity of biologically produced (Z)-11-hexadecenoic acid was determined by comparing the fragmentation patterns of the dimethyl disulfide (DMDS) derivative with the DMDS derivative of an authentic standard. A yeast culture was split into 12 aliquots (to not change any parameters in the developed procedure). The cells were pelleted, which yielded 63 mg cells (ccw) on average (755 mg from 18 mL culture). The pellets were subjected to base methanolysis as described above. However, after acidification the samples were combined in a 50 mL falcon tube. The combined sample was extracted two times with 10 mL chloroform. The mixture was centrifuged 10 min at 3000 rpm to achieve a better phase separation. The combined organic phases were combined in a new 50 mL falcon and were washed consecutively with 10 mL brine and 10 mL water. The organic phase was dried with anhydrous sodium sulfate and concentrated in vacuo. The concentrated oil was dissolved in 1.5 mL chloroform and transferred to a 1.5 mL plastic tube. The chloroform was evaporated at 90° C. The remaining sample was the dissolved in 50 µL methyl tert-butyl ether (MTBE). The 50 µL were split into 1, 5, 10 and 20 µL and transferred into GC-vials without insert. To each vial 200 µL DMDS (dimethyl disulfide) and 50 µL MTBE (containing 60 mg/mL iodine) were added. After the mixture was heated 48 h at 50° C., excess iodine was removed by the addition of 100 µL saturated sodium thiosulfate solution; however, due to excessive formation of detergents from the *Candida* strain, the layer did not mix properly. The samples were therefore diluted in a 15 mL falcon tube to a final sample composition of 200 µL, 3.55 mL MTBE (containing iodine and analyte), 500 µL dichloromethane, 1.5 mL water and 1 mL ethanol. The organic phase was evaporated stepwise at 85° C. in a 1.8 mL glass vial. The samples were taken up in 500 µL dichloromethane and the sample was analyzed by GC-MS using the method of Hagström et al. (2013) as in Example 4.

Example 7: Expression of Transmembrane
Desaturases in *Yarrowia lipolytica*

Background and Rationale

Engineering microbial production of insect fatty alcohols from fatty acids requires the functional expression of a synthetic pathway. One such pathway comprises a transmembrane desaturase, and an alcohol-forming reductase to mediate the conversion of fatty acyl-CoA into regio- and stereospecific unsaturated fatty acyl-CoA, and subsequently into fatty alcohols. A number of genes encoding these enzymes are found in some insects as well as some microalgae. Alternatively, regio- and stereospecific desaturases can be used to produce a microbial oil rich in fatty acid precursors. The microbial oil can then be derivatized and reduced to active ingredients. A number of gene variants were screened to identify enzyme activities that allow the creation of pathways capable of high level synthesis of a single or a blend of insect fatty acids and alcohols. Additionally, these enzymes were screened across multiple hosts (*Saccharomyces cerevisiae*, *Candida viswanathii* (*tropicalis*), and *Yarrowia lipolytica*) to optimize the search toward finding a suitable host for optimum expression of these transmembrane proteins.

Initial screening of desaturases in *S. cerevisiae* and *C. viswanathii* (*tropicalis*) identified three active Z11-C16:1 desaturase variants from *Amyelois transitella*, *Helicoverpa zea*, and *Trichoplusia ni*. The *S. cerevisiae* screening used coding sequences with an N-terminal leader sequence of the *S. cerevisiae* Ole1p Z9 desaturase fused to the full length insect Z11 desaturase sequence. This strategy has been used previously in the scientific literature to express eukaryotic desaturases in *S. cerevisiae*. All three of the above desaturases displayed Z11 desaturase activity with the Ole1p leader fusion when expressed in a OLE1 deletion background. An analogous design with a *C. albicans* Ole1p leader sequence was used with the Z11 desaturase from *H. zea*. While active, this Ole1p-*H. zea* desaturase fusion did not significantly increase Z11-hexadecenoic acid titer. Additionally, a conservatively optimized *A. transitella* Z11 desaturase was active in both *S. cerevisiae* and *C. viswanathii*. The following study focused on testing the functional expression of the *H. zea*, *T. ni*, and *A. transitella* Z11 desaturases in two different *Y. lipolytica* strains, SPV140 and SPV300. Both native and *Homo sapiens* codon optimized sequences were used for the *H. zea* and *T. ni* desaturases while only the native sequence was used for *A. transitella*. Finally, the N-terminus of the *Y. lipolytica* Ole1p Z9 stearoyl-CoA desaturase aligns more closely with insect desaturases than the N-terminus of Ole1p from either *S. cerevisiae* or *C. albicans*. Based on this alignment two additional desaturase versions were created. A putative leader sequence was swapped from the *Y. lipolytica* Ole1p onto the *T. ni* and *H. zea* desaturases.

Summary of Approach

A focused library of Z11 desaturases (insect origin: *Amyelois transitella*, *Helicoverpa zea*, *Trichoplusia ni*), which had observed activity in either *S. cerevisiae* or *C. viswanathii* were cloned into a double crossover cassette targeting the XPR2 locus with a URA3 selection marker. Protein coding sequences use either the native insect sequence (SEQ ID NOs: 24, 25), *Homo sapiens* optimized coding sequence (SEQ ID NOs: 26, 27), or the *Homo sapiens* optimized sequence with the N-terminal 84 bases (*H. zea*, SEQ ID NO: 29) or 81 bases (*T. ni*, SEQ ID NO: 28) swapped for the N-terminal 96 bases of the *Y. lipolytica* OLE1 (YALI0C05951) gene. Unlike in the *S. cerevisiae* and *C.*

viswanathii screens, the leader sequence chimeras test a direct swap of leader sequences instead of concatenating a host leader sequence to the N-terminus of the full length desaturase coding sequence. Only the native coding sequence was used for the *A. transitella* desaturase (SEQ ID NO: 30).

Each of the 7 desaturase constructs was transformed into SPV140 (PO1f) and SPV300 (H222 ΔP ΔA ΔF ΔURA3) and site-specific integrants were confirmed.

Desaturase activity was tested via an in vivo bioconversion of hexadecanoic acid (palmitic acid) into (Z)-11-hexadecenoic acid (palmitavaccenic acid) in YPD medium.

GC-FID analyses were used to identify and quantify metabolites.

Results

Strain Construction

Desaturase variants were cloned into the pPV101 vector which contains a *Y. lipolytica* expression cassette targeting integration into the XPR2 locus.

The *T. ni* and *H. zea* desaturases were each synthesized with the native insect sequence (SEQ ID NOs: 24, 25), full length insect sequence codon optimized for *Homo sapiens* (SEQ ID NOs: 26, 27), or with the putative leader sequence replaced by the leader sequence from *Y. lipolytica* OLE1 desaturase (SEQ ID NOs: 28, 29). The *A. transitella* desaturase was also synthesized using the native insect coding sequence (SEQ ID NO: 30). All seven desaturase variants were transformed into SPV140. Based on previous activity results, only the *H. zea* and *A. transitella* desaturase variants were transformed into SPV300.

Functional Expression Assay

Functional activity was assessed by a modification of the protocol used for transmembrane desaturase expression in *C. viswanathii* SPV053 (See Example 6). Briefly, *Y. lipolytica* SPV140 and SPV300 derived strains expressing insect desaturases were cultured in rich (YPD) to generate biomass. Using the YPD generated biomass, small scale (2 ml) cultures were cultivated with palmitic acid for a total of 48 hours in 24 deep well plates (See Materials & Methods for detail).

In the initial screen of *T. ni*, *H. zea*, and *A. transitella* variants, only *H. zea* desaturase variants that were codon optimized for *Homo sapiens* produced detectable Z11-hexadecenoic acid (FIG. 26). Expression of native *H. zea* desaturase conferred production of 100±5 mg/L Z11-hexadecenoic acid and the version with a *Y. lipolytica* OLE1 leader sequence produced 83±11 mg/L. As seen in FIG. 26, the distribution of the other major fatty acid species was relatively unaffected by functional desaturase expression. In the active strains, Z11-hexadecenoic acid made up ~10% (g/g) of the fatty acid species (including palmitic acid substrate which may be adsorbed to the outer cell surface).

A follow up experiment was conducted comparing active variants in the SPV140 background to SPV300 derived desaturase strains. The parent SPV300 and SPV140 expressing hrGFP were used as negative controls. The same bioconversion assay protocol was used. As in SPV140, only *H. sapiens* optimized variants produced detectable activity (FIG. 27). SPV300 strains grew to higher final cell densities (SPV300 OD600=26-28, SPV140 OD600=19-22) (FIG. 28). The highest titers were observed for strains expressing the native *H. zea* Z11-desaturase with *H. sapiens* codon optimization (pPV199). The retested SPV140 strains produced 113±1 mg/L (5.5±0.2 mg/L/OD) Z11-hexadecenoic acid which is 13% higher than titers observed in the first experiment (FIG. 29). SPV300 strains expressing the same desaturase generated a wider range of productivity. On

average they produced 89 ± 18 mg/L (3.3 ± 1.2 mg/L/OD) Z11-hexadecenoic acid, but one clone produced 124 mg/L (4.6 mg/L/OD) Z11-hexadecenoic acid.

In summary, only the *H. zea* Z11 desaturase variants with *Homo sapiens* codon optimization produced detectable Z11-hexadecenoic acid. Under the current assay condition, marginally higher titers were observed in the SPV140 background over SPV300. Table 11 summarizes the Z11-hexadecenoic acid titers.

TABLE 11

Z11-hexadecenoic acid titers obtained from expression of exemplary desaturases in <i>Yarrowia lipolytica</i>			
Desaturase	Codon optimization	Parent Strain	Z11-hexadecenoic acid titer (mg/L)
Z11 <i>T. ni</i>	Native	SPV140	ND (no detection)
Z11 <i>T. ni</i>	<i>Homo sapiens</i>	SPV140	ND
Y1 OLE1-Z11 <i>T. ni</i>	<i>Homo sapiens</i>	SPV140	ND
Z11 <i>H. zea</i>	Native	SPV140	ND
		SPV300	ND
Z11 <i>H. zea</i>	<i>Homo sapiens</i>	SPV140	100 ± 5
		SPV300	87 ± 18
Y1 OLE1-Z11 <i>H. zea</i>	<i>Homo sapiens</i>	SPV140	83 ± 11
		SPV300	55 ± 1
Z11 <i>A. transitella</i>	Native	SPV140	ND
		SPV300	ND

In SPV300, one non-site-specific integrant of pPV200 (*Y. lipolytica* OLE1-*H. zea* Z11 desaturase with *Homo sapiens* codon optimization) was tested. This integrant did not produce detectable Z11-hexadecenoic acid, while the two site-specific integrants produced 55 ± 1 mg/L.

No major hydroxy or diacid peaks were observed from pellets of SPV140 or SPV300 derived strains, and deletion of β -oxidation/ ω -oxidation genes in SPV300 did not increase Z11-hexadecenoic acid accumulation under the current assay condition (relatively low substrate concentration, rich medium).

Conclusions

The *H. zea* Z11 desaturase is active and confers production of ~ 100 mg/L Z11-hexadecenoic acid, from ~ 500 mg/L palmitic acid substrate. The functional expression was demonstrated across three positive integrants and replicate experiments in a 24 well plate assay.

H. zea desaturase required codon optimization (*Homo sapiens* or potentially *Y. lipolytica*) for activity in *Y. lipolytica*.

The *T. ni* Z11 desaturase, while active in *S. cerevisiae*, does not produce detectable Z11-hexadecenoic acid in *Y. lipolytica*.

The reproducibility of the assay for *Y. lipolytica* strains can be confirmed starting from glycerol stock.

A. transitella desaturase can be codon optimized for expression in *Y. lipolytica*.

Since *Y. lipolytica* is a candidate production host, additional copies of active desaturases can be integrated in *Y. lipolytica*, culture conditions to improve bioconversion can be identified, and substrate conversion can be quantified.

Materials & Methods

Strain Construction

All desaturase genes were synthesized (Genscript). Either native sequences or *Homo sapiens* codon optimization was used. Synthesized genes were subcloned into pPV101. Plasmids were transformed and prepped from *E. coli* EPI400 using the Zyppy Plasmid Miniprep Kit (Zymo Research, Irvine, Calif.). Approximately ~ 1 - 2 μ g of linearized DNA was transformed using Frozen-EZ Yeast Transformation II

Kit (Zymo Research, Irvine, Calif.). The entire transformation mixture was plated on CM glucose -ura agar plates. Positive integrants were found to be site-specific and genotyping was conducted by check PCR.

Functional Expression Assay

Palmitic Acid Supplementation in YPD

Positive isolates were re-patched onto YPD, grown overnight, and then stored at 4° C. Strains were inoculated from patch plates into 2 ml of YPD in 24 deep well plates (square well, pyramid bottom). Three positive clones were inoculated for each desaturase variant. Three isolates of pPV101 in SPV140 and the parent SPV300 were used as negative controls. Deep well plates were incubated at 28° C. and 250 rpm in the Infors Multitron refrigerated flask shaker for 24 hrs. After 24 hrs of incubation, a 1 ml volume of each culture was pelleted by centrifugation at $500 \times g$. Each pellet was resuspended in 2 ml of YPD. 500 mg/L palmitic acid was added to cultures from a 90 g/L stock solution in ethanol. The result was the addition of 0.5% ethanol with the palmitic acid substrate. All cultures were incubated for 48 hours before endpoint sampling. Final cell densities were measured with the Tecan Infinite 200pro plate reader. 0.75 or 0.8 ml of each culture was harvested in 1.7 ml microcentrifuge tubes and pelleted. Supernatant was removed and pellets were processed as described below.

Metabolite Extraction and GC-FID Analysis

Total lipid composition as well as the (Z)-11-hexadecenoic acid quantification was based on modified procedures by Moss et al. (1982) and Yousuf et al (2010). The pelleted cells (in 1.5 mL plastic tubes), usually about 10 mg to 80 mg, were resuspended in methanol containing 5% (w/w) of sodium hydroxide. The alkaline cell suspension was transferred into a 1.8 mL crimp vial. The mixture was heated for 1 h in the heat block at 90° C. Prior to acidification with 400 2.5 N HCl the vial was allowed to cool to room temperature. 500 μ L chloroform containing 1 mM methyl heptadecanoate were added and the mixture was shaken vigorously, then both aqueous and organic phase were transferred into a 1.5 mL plastic tube. The mixture was centrifuged at 13,000 rpm, afterwards 450 μ L of the organic phase were transferred into a GC vial. For the analysis of lipids and the quantification of fatty acids 50 μ L of 0.2 M TMSH (trimethylsulfonium hydroxide in methanol) was added and the sample analyzed by GC-FID.

Example 8: *Candida viswanathii* (*Tropicalis*) as a Production Platform for Insect Fatty Alcohol Synthesis

Background and Rationale

Variants of insect transmembrane desaturases and reductases were previously screened and rank-ordered based on their functional expression in either *Candida viswanathii* or *Saccharomyces cerevisiae* (see Examples 3, 4 and 6). *Helicoverpa zea* desaturase and *Helicoverpa armigera* reductase were selected to assemble a synthetic insect fatty alcohol pathway in *C. viswanathii*. Simultaneous expression of codon optimized *H. zea* desaturase under *Candida* isocitrate lyase (ICL) promoter, and codon optimized *H. armigera* reductase under *Candida* transcription elongation factor (TEF) promoter was achieved via genomic integration of the full fatty alcohol pathway. Accumulation of Z11-16OH was achieved in cultures of the recombinant strain (SPV0490) using simple carbon sources and palmitic acid.

Summary of Approach

Integration plasmids were designed containing a functional *Helicoverpa zea* desaturase (See Example 6) paired

135

with a *Helicoverpa armigera* reductase driven by a putatively constitutive *C. tropicalis* promoter (pTEF).

Functionality of the full pathway was assessed via an in vivo bioconversion of hexadecanoic acid (palmitic acid) into Z11-16OH.

GC-FID and GC-MS analyses were used to identify and quantify metabolites.

Results

Accumulation of Z11-16OH was detected in cultures of *Candida* engineered to express *H. zea* desaturase under an ICL promoter and *H. armigera* reductase under a TEF promoter (Table 12 and FIG. 30).

TABLE 12

Tabulated Z11-16OH titers from <i>Candida viswanathii</i> bioconversion assay. SPV088 is <i>C. viswanathii</i> which was engineered to express mCherry (negative control). SPV0490 is <i>C. viswanathii</i> which was engineered to express the insect fatty alcohol pathway. Z11-16OH titers (mg/L)		
	SPV0488 (negative control)	SPV0490
Sample 1	0.08	1.03
Sample 2	0.07	0.93
Sample 3	0.06	0.88
Average	0.07	0.95
StDev	0.01	0.06

Materials & Methods

Strain Construction

The integration plasmid (ppV0228) was designed to contain two expression cassettes. The first cassette contains *H. zea* codon-optimized desaturase (SEQ ID NO: 31) that was driven by the *C. viswanathii* ICL promoter (SEQ ID NO: 33). The second cassette contains codon-optimized *H. armigera* reductase (SEQ ID NO: 32) driven by the *C. tropicalis* TEF promoter (SEQ ID NO: 34). Gene expression in the ICL promoter cassette is terminated by the ICL terminator sequence (SEQ ID NO: 35). Gene expression in the TEF promoter cassette is terminated by the TEF terminator sequence (SEQ ID NO: 36). A conservative approach was used for recoding of genes. Native gene sequences were unaltered except for replacement of CTG leucine codons with TTA. After transformation into *E. coli* NEB100, plasmids were miniprep using the Zyppy Plasmid Miniprep Kit (Zymo Research, Irvine, Calif.). Plasmids were linearized by digestion with BsiWI (New England Biolabs, Ipswich, Mass.) before transformation into SPV053. After digestion, DNA was isolated using Clean and Concentrator Kit (Zymo Research, Irvine, Calif.). Approximately 3-5 µg of DNA was transformed by electroporation. Positive integrants were found to be site-specific and genotyping was conducted by check PCR. A two-stage approach was adopted for further screening of low efficiency transformations. Approximately 100 colonies were re-patched on YPD+250 µg/ml Zeocin and grown overnight. The subset of patches which grew quickly (dense growth within 24 hours) were screened by colony PCR.

Functional Expression Assay

Palmitic Acid Supplementation in YPD

Positive isolates were re-patched onto YPD+300 µg/ml Zeocin, grown overnight and then stored at 4° C. Strains were inoculated from patch plates into 2 ml of YPD in 24 deep well plates (square well, pyramid bottom). Four positive clones were inoculated for each desaturase and reductase variant and three positive clones were inoculated for each desaturase and mCherry expressing control strain.

136

Deep well plates were incubated at 30° C., 1000 rpm, and 80% humidity in the Infors HT Multitron Pro plate shaker for 24 hrs. After 24 hrs of incubation, a 1 ml volume of each culture was pelleted by centrifugation at 500×g. Each pellet was resuspended in 2 ml of YPD+0.3% (v/v) ethanol. Ethanol was added at this stage to induce recombinant enzyme expression from the ICL promoter. Cultures were incubated for another 24 hours under the same conditions before 300 mg/L palmitic acid was added to cultures from a 90 g/L stock solution in ethanol. The result was the addition of a fresh 0.3% ethanol in conjunction with the palmitic acid. All cultures were incubated for an additional 24 hrs before a final addition of 0.3% ethanol. After another 24 hr period of incubation, 1.5 ml of each culture was harvested in 1.7 ml microcentrifuge tubes and pelleted. Supernatant was removed and pellets were processed as described below.

Metabolite Extraction and GC-MS Detection

The pelleted cells (in 1.5 mL plastic tubes), usually about 10 mg to 80 mg, were resuspended in methanol containing 5% (w/w) of sodium hydroxide. The alkaline cell suspension was transferred into a 1.8 mL crimp vial. The mixture was heated for 1 h in a heat block at 90° C. Prior to acidification with 400 µL 2.5 N HCl the vial was allowed to cool to room temperature. 500 µL chloroform containing 1 mM methyl heptadecanoate were added and the mixture was shaken vigorously, then both aqueous and organic phase were transferred into a 1.5 mL plastic tube. The mixture was centrifuged at 13,000 rpm, afterwards 450 µL of the organic phase were transferred into a GC vial. The organic phase was evaporated in a heat block at 90° C. for 30 min. The residue was dissolved in 50 µL N,O-Bis(trimethylsilyl) trifluoroacetamide containing 1% trimethylchlorosilane. Prior to transfer into glass inserts the mixture was heated 5 min at 90° C. The samples were analyzed by GC-MS (Table 13).

TABLE 13

Analytical parameters used for GC-MS analysis of metabolites	
System	Agilent 6890 N GC, ChemStation G1701EA E.02.01.1177
Column	DB23 30 m × 25 µm × 25 µm Pressure = 11.60 psi; Flow = 0.6 mL/min
Inlet	Heater = 250° C.; Pressure = 11.74 psi; Total Flow {He} = 111 mL/min
Carrier	He @ 29 cm/sec, 11.60 psi
Signal	Data rate = 2 Hz/0.1 min
Oven	150° C. for 1 min Ramp 12° C./min to 220° C., hold 3 min Ramp 35° C./min to 300° C., hold 4 min
Injection	Splitless, 250° C.
Detector	HP 5973 MSD in SIM mode (m/z: 208.0, 297.3 and 387.3), 100 msec Dwell, EMV mode: Gain factor 1, 2.4 min solvent delay, 3.09 cycles/sec
Sample	Injection volume = 1 µL

Example 9: Insect Fatty Alcohol Production from *Yarrowia lipolytica*

Background and Rationale

Yarrowia lipolytica was engineered as a production platform for insect fatty alcohol (Z11-16OH and Z9-16OH) synthesis from palmitic acid.

After individually confirming functional expression of a Z11 desaturase (Example 7) and fatty acyl-CoA reductase (FAR), the full Z11-16OH and Z9-16OH pathways (Bdr) were engineered in *Y. lipolytica*. For the purpose of improv-

ing fatty alcohol titers, cultivations designed for promoting growth vs. for eliciting lipid storage were also explored. A growth condition favors high biomass production, but limits fatty acyl-CoA pool size used by the engineered pathway and directs fatty acyl-CoA intermediates to membrane synthesis. Conversely, a lipid storage condition creates a strong sink for production of fatty acyl-CoAs which is desirable. However, fatty acyl-CoA transport towards lipid bodies creates a strong competition for FAR activity. Under this second scenario, even though Z11-16Acid or Z9-16Acid accumulates in the cell, most of it is inaccessible to the FAR. On the other hand, there may be a continual flux of lipid remobilization under lipid storage conditions which leads to a sustained pool of Z11-16CoA or Z9-16CoA which is available to the FAR.

Summary of Approach

Two biodesaturation-reduction (B&) pathway variants were tested in the H222 ΔPΔAΔF (SPV300) background. The first combined recombinant expression of *Helicoverpa zea* Z11 desaturase paired with a *Helicoverpa armigera* fatty acyl-CoA reductase (FAR) creating a Z11-16OH synthesis

Relatively low, detectable Z11-16OH titers (0.26±0.09 mg/L) were observed in a YPD medium supplemented with 10 g/L methyl palmitate. The Z11-16Acid precursor was measured at 220±11 mg/L (across clones 2, 4, 9, 17, 23).

Higher Z11-16OH titers were observed in a semi-defined medium with C:N ratio of ~80. Across all 10 isolates Z11-16OH was produced at 2.65±0.36 mg/L. The Z11-16Acid precursor titer was 900±30 mg/L. One isolate (SPV578) produced 3.68±0.31 mg/L Z11-16OH (Z11-16Acid 840±14 mg/L).

Nine isolates expressing the *H. zea* desaturase (pTEF) and *H. armigera* reductase (pTAL1) were screened. The in vivo bioconversion assay confirmed Z11-16OH production from all isolates.

One isolate (SPV603) produced 6.82±1.11 mg/L Z11-16OH in a semi-defined medium (Z11-16Acid 1.36 g/L).

The previously tested reductase strain, SPV471 (H222 ΔPΔAΔF expressing *H. armigera* FAR), produced 4.30±2.33 mg/L Z9-16OH and 450±80 mg/L Z9-16Acid using a semi-defined medium (C:N ratio of ~80).

TABLE 14

Summary table of Z11/Z9-16OH titers from B _{dr} pathway strains in in vivo bioconversion assay.			
Strain	Medium	Z11-16OH (mg/L)	Z9-16OH (mg/L)
pTEF- <i>H. zea</i> Z11 desaturase pEXP- <i>H. armigera</i> FAR Clone 17 (SPV578)	Semi-Defined C:N = 80	3.99 ± 0.37 (n = 4)	0.22 ± 0.03 (n = 4)
pTEF- <i>H. zea</i> Z11 desaturase pTAL- <i>H. armigera</i> FAR Clone 9 (SPV603)	Semi-Defined C:N = 80	6.82 ± 1.11 (n = 2)	0.22 ± 0.01 (n = 2)
pOLE1- <i>Y. lipolytica</i> OLE1 (native) pTEF- <i>H. armigera</i> FAR (SPV471)	Semi-Defined C:N = 80	0.22 ± 0.03 (n = 2)	4.30 ± 2.23 (n = 2)

pathway. The second combined native *Y. lipolytica* Z9 desaturase activity with *H. armigera* fatty acyl-CoA reductase (FAR) expression creating a Z9-16OH pathway.

Two integration plasmids were constructed to express the *H. zea* desaturase and the *H. armigera* FAR. The TEF promoter was used for desaturase expression and the EXP1 (export protein) or the TAL1 (transaldolase) promoter was used for reductase expression.

Successful integration of the Z11-16OH pathway cassette into the H222 ΔPΔAΔF (SPV300) background was confirmed by colony PCR.

Functionality of the full Z11-16OH pathway was assessed via an in vivo bioconversion of 16Acid (palmitic acid) into Z11-16OH (Z-11-hexadecenol).

Functionality of a full Z9-16OH pathway was assessed via an in vivo bioconversion of 16Acid (palmitic acid) using previously constructed SPV471 (H222 ΔPΔAΔF derived) which expresses the *H. armigera* FAR driven by the TEF promoter.

GC-MS analysis was used to identify and quantify Z9-16OH and Z11-16OH. GC-FID analysis was used to identify and quantify fatty acids.

SUMMARY

Ten isolates expressing the *H. zea* desaturase (pTEF) and *H. armigera* reductase (pEXP1) were screened. The in vivo bioconversion assay confirmed Z11-16OH production from all isolates.

Results

Strain Construction

Evidence in the literature suggests both insect desaturases and FARs are localized in the membrane of the endoplasmic reticulum with active sites oriented towards the cytoplasm. Of the functional variants, the Z11 desaturase from *H. zea* and the FAR from *H. armigera* were selected, one hypothesis being that using enzymes from the same genus (*Helicoverpa*) could better conserve protein-protein interactions that may occur in the ER membrane.

Two new constructs were ordered from Genscript and cloned into the previously assembled *H. zea* desaturase plasmid, pPV0199. Two FAR synthons with either the EXP1 or TAL1 promoter from *Y. lipolytica* were cloned into this expression cassette.

One dual expression plasmid (with EXP1 promoter) was transformed into the parent strain SPV300 (H222 Δpox1 Δpox2 Δpox3 Δpox4 Δpox5 Δpox6 Δadh1 Δadh2 Δadh3 Δadh4 Δadh5 Δadh6 Δadh7 Δfao1 Δura3). Two different competent cell preparations of the same parent strain were transformed to study variability in strain performance resulting from competent cell preparation. Approximately 25% of URA+ clones were confirmed to be targeted integrants at the XPR2 locus (20% for preparation 1, 33% for preparation 2). Two clones from Comp. Cell Preparation 1 and all eight targeted clones from Comp. Cell Preparation 2 were selected for screening in the functional expression assay.

The second dual expression plasmid (with TAL1 promoter) was integrated into the same parent strain (SPV300).

Twenty-three colonies were screened by check PCR and 11 were found to be targeted integrants (48%). Nine integrants were selected for screening in the functional expression assay.

The construct of SPV471 (H222 Δ P Δ A Δ F expressing *H. armigera* FAR) was described previously.

Z11-16OH Functional Expression Assay

An in vivo, 24-well plate assay was used to evaluate production of Z11-16OH. The assay was based on designs used for screening desaturase and reductase variants as well as conditions used to increase fatty acid accumulation. A rich medium (YPD) and a semi-defined medium were used with 10 g/L methyl palmitate supplemented as bioconversion substrate. The semi-defined medium had a C:N ratio of ~80 and included 5 g/L glycerol and 60 g/L glucose (See Materials & Methods for further details).

The initial screen of strains harboring the *H. zea* desaturase driven by the TEF promoter and the *H. armigera* FAR driven by the EXP1 promoter confirmed that the presence of FAR was required to produce Z11-16OH. No hexadecenol was observed from both the parent and desaturase-only control strains under any condition. Under both media conditions Z11-16OH and to a lesser extent Z9-16OH were detected from clones expressing the full desaturase-reductase pathway. When the conversion was completed in rich medium, 0.26 $\pm$ 0.09 mg/L Z11-16OH and 0.06 $\pm$ 0.01 mg/L Z9-16OH were produced (FIG. 32A). A 10-fold increase in Z11-16OH titer and 3-fold increase in Z9-16OH titer was observed when the Semi-Defined medium was used (FIG. 32B). Across all pathway clones 2.65 $\pm$ 0.29 mg/L Z11-16OH and 0.18 $\pm$ 0.02 mg/L Z9-16OH were produced. The enrichment of Z11-16OH over Z9-16OH supports the potential for engineering a regiospecific Bdr pathway. Consistency between technical replicates varied across clones under the Semi-Defined medium condition. Titers for Clones 2, 4, 6, 9, and 17 were consistent with CVs <20. Clones 1, 7, and 23 have CVs >40%. The highest consistent Z11-16OH titer was observed for Clone 17, 3.68 $\pm$ 0.31 mg/L (Table 15).

TABLE 15

Summary table of Z11/Z9-16OH titers for pEXP1 clones. A population of ten isolates expressing the <i>H. zea</i> desaturase driven by pTEF and <i>H. armigera</i> reductase driven by pEXP1, from two independent competent cell preparations, were assayed for Z11-16OH and Z9-16OH production under two different media conditions. Alcohol production across isolates and from select clones are presented.					
pTEF-Hz_desat pEXP-Ha_FAR Clone(s)	Medium	Z11-16OH (mg/L)	Z9-16OH (mg/L)	Z11-16OH fold increase (relative to YPD)	Z9-16OH fold increase (relative to YPD)
All clones	YPD	0.26 $\pm$ 0.09	0.06 $\pm$ 0.01	—	—
All clones	Semi-Defined	2.65 $\pm$ 0.29	0.18 $\pm$ 0.02	10	3
Clone 2 (SPV574)	YPD	0.18 $\pm$ 0.09	0.05 $\pm$ 0.03	—	—
Clone 2 (SPV574)	Semi-Defined	2.08 $\pm$ 0.26	0.14 $\pm$ 0.04	12	3
Clone 4 (SPV575)	YPD	0.28 $\pm$ 0.01	0.11 $\pm$ 0.01	—	—
Clone 4 (SPV575)	Semi-Defined	3.24 $\pm$ 0.28	0.21 $\pm$ 0.03	12	2
Clone 9 (SPV576)	YPD	1.03 $\pm$ 0.84	0.05 $\pm$ 0.01	—	—
Clone 9 (SPV576)	Semi-Defined	1.56 $\pm$ 0.28	0.11 $\pm$ 0.02	1.5	2
Clone 23 (SPV577)	YPD	0.16 $\pm$ 0.14	0.05 $\pm$ 0.05	—	—
Clone 23 (SPV577)	Semi-Defined	3.35 $\pm$ 1.85	0.26 $\pm$ 0.15	21	5
Clone 17 (SPV578)	YPD	0.19 $\pm$ 0.01	0.06 $\pm$ 0.01	—	—
Clone 17 (SPV578)	Semi-Defined	3.68 $\pm$ 0.31	0.26 $\pm$ 0.02	14	4

The lipid profiles of the full pathway clones were also quantified. For simplicity the 16 carbon fatty acid species are plotted for select clones in FIG. 33A-33B. In general, the full Bdr pathway clones accumulated less Z11-16Acid than the desaturase only control (0.25<0.5 g/L in YPD, 0.8-1.0<1.5 g/L in Semi-Defined). Lower Z11-16Acid titers in full Bdr pathway clones may result from reduced desaturase expression in the dual expression cassette or potentially from Z11-16Acid consumption by FAR and subsequent byproduct pathways. No trend in 16Acid titer was observed in YPD, while 16Acid titers were similar for desaturase only and full pathway strains in the Semi-Defined medium.

Strains using the second dual expression cassette (pTAL-Ha_FAR) were assayed under the same Semi-Defined medium condition used to evaluate the pEXP clones. Nine pTAL clones were assayed against SPV300 (parent), SPV575 (pEXP-Ha_FAR Clone 4), and SPV578 (pEXP-Ha_FAR Clone 17) controls. As expected, no alcohol products were observed from the negative control. Alcohol titers from pEXP positive control strains replicated results observed in the initial assay of pEXP clones (FIG. 34, Table 16). Excluding one outlier clone, Clone 9, Z11-16OH titer was equivalent from pTAL clones (4.19 $\pm$ 0.16 mg/L) and pEXP clones (4.10 $\pm$ 0.22 mg/L). Clone 9 produced Z11-16OH at 6.82 $\pm$ 1.11 mg/L. As in the first assay with pEXP clones, low, but detectable titers of Z9-16OH were observed (FIG. 34, Table 16).

TABLE 16

Summary table of Z11/Z9-16OH titers for pTAL1 clones. A population of nine isolates expressing the <i>H. zea</i> desaturase under the TEF promoter and <i>H. armigera</i> reductase under the TAL promoter were assayed for Z11-16OH and Z9-16OH production under a Semi-Defined medium condition. Clones were compared to positive controls expressing the <i>H. zea</i> desaturase under the TEF promoter and <i>H. armigera</i> reductase under the EXP promoter. Alcohol production across isolates and from select clones are presented.			
pTEF-Hz_desat pEXP-Ha_FAR Clone(s)	Medium	Z11-16OH (mg/L)	Z9-16OH (mg/L)
EXP Clone 4 (SPV575)	Semi-Defined	3.91 $\pm$ 0.44	0.15 $\pm$ 0.01

TABLE 16-continued

Summary table of Z11/Z9-16OH titers for pTAL1 clones. A population of nine isolates expressing the *H. zea* desaturase under the TEF promoter and *H. armigera* reductase under the TAL promoter were assayed for Z11-16OH and Z9-16OH production under a Semi-Defined medium condition. Clones were compared to positive controls expressing the *H. zea* desaturase under the TEF promoter and *H. armigera* reductase under the EXP promoter. Alcohol production across isolates and from select clones are presented.

pTEF-Hz_desat pEXP-Ha_FAR Clone(s)	Medium	Z11-16OH (mg/L)	Z9-16OH (mg/L)
EXP Clone 17 (SPV578)	Semi-Defined	4.30 ± 0.16	0.17 ± 0.02
pTAL clones excluding Clone 9	Semi-Defined	4.19 ± 0.16	0.18 ± 0.01
pTAL Clone 9 (SPV603)	Semi-Defined	6.82 ± 1.11	0.22 ± 0.01

The lipid profiles of all strains in the second (pTAL) full pathway screen were also quantified. For simplicity the 16 carbon fatty acid species are plotted in FIG. 35. As expected, Z11-16Acid is present only for strains expressing the desaturase. Complete lipid profiles were similar to those observed previously (FIG. 36). Z9-18Acid (oleic acid) was the second most abundant fatty acid species after Z11-16Acid.

Z9-16OH Functional Expression Assay

An in vivo, flask scale assay was used to test for Z9-16OH production. The parent control strain, H222 ΔPΔΔF (SPV300), was compared to a strain expressing *H. armigera* FAR which relied on native Z9 desaturase activity to synthesize the Z9-16CoA precursor (SPV471). Biomass was generated through a YPD seed culture, mimicking the plate assay. Bioconversion flasks were inoculated at an initial OD600=1 or OD600=4 into the same Semi-Defined C:N=80 medium used in the Z11-16OH plate assay (See Materials & Methods for details). As expected, control flasks did not produce detectable Z9-16OH while SPV471 flasks produced up to 4.30±2.23 mg/L after 24 hours of incubation (FIG. 37A-FIG. 37B). While there was large variability between replicates, all SPV471 (*H. armigera* FAR) replicates exceeded 1 mg/L titer. Increased seeding density did not increase Z9-16Acid or Z9-16OH titer. The precursor Z9-16Acid titer at 24 hours was significantly less (<0.5 g/L) than the Z11-16Acid precursor observed for dual expression cassette strains used to produce Z11-16OH. The relative abundance of other fatty acid species was similar to previously observed profiles, with Z9-18Acid as the next most abundant species (FIG. 38). Both lipid and alcohol samples were taken over the course of 48 hours to produce a time course of Z9-16OH and lipid titers. Z9-16OH titer peaked at 24 hours before decreasing over the second day (FIG. 39A). Z9-16Acid increased rapidly over the first 24 hours before stabilizing or increasing slowly over the second 24 hours (FIG. 39B). Since the employed analytical method utilizes only the cell pellet, the decrease in Z9-16OH titer supports the hypothesis of downstream consumption or secretion of the alcohol products. They may be oxidized (ω-oxidation), secreted as free alcohol, or derivatized and secreted as an ester. Analysis of supernatant samples using FID and MS SCAN detection revealed no detectable Z9-16OH or Z9-16OH derivatives supporting the hypothesis of consumption via oxidation pathways.

Conclusions

Combining expression of *Helicoverpa* Z11 desaturase and fatty acyl-CoA reductase led to production of Z11-16OH in *Y. lipolytica* H222 ΔPΔΔF (SPV300) at titers >1 mg/L.

High C:N ratio conditions improved Z11-16OH titer relative to a rich medium condition.

Under lipid accumulating conditions the combination of native Z9 desaturase and *H. armigera* FAR activities are sufficient for synthesis of >1 mg/L Z9-16OH.

Titers are increased, for example, by deleting pathways consuming fatty alcohol products and/or fatty acid precursors; identifying FAR variants which exhibit higher turnover rate than *H. armigera* FAR; and/or increasing pathway copy number.

Key undesired byproducts are identified.

The possibility that some of the fatty alcohol product is converted into fatty acetate by the activity of one or more endogenous acetyltransferases is explored.

Improved host strains are engineered to eliminate the ω-oxidation pathway and components of the lipid storage pathway.

Additional copies of desaturase and FAR are integrated into *Y. lipolytica*.

Materials & Methods

Strain Construction

All desaturase and reductase genes were ordered from Genscript. *Homo sapiens* codon optimization was used (Genscript algorithm). The newly synthesized expression cassette was subcloned into pPV199 by Genscript using the SapI restriction site. Plasmids were transformed and prepped from *E. coli* EP1400 using the Zyppy Plasmid Miniprep Kit (Zymo Research, Irvine, Calif.). Plasmids were digested with PmeI (New England Biolabs, Ipswich, Mass.) and purified by gel extraction using Zymoclean Gel DNA recovery Kit (Zymo Research, Irvine, Calif.). DNA was further concentrated using Clean and Concentrator Kit (Zymo Research, Irvine, Calif.). Approximately ~1-2 μg of DNA was transformed using Frozen-EZ Yeast Transformation II Kit (Zymo Research, Irvine, Calif.). The manufacturer's protocol was modified as follows: A 2 ml YPD seed culture was inoculated at 9 am the day before competent cell preparation. The seed was grown 8 hours (until 5 pm) before 40 ml of YPD in a 250 ml baffled shake flask (or 20 ml in a 125 ml baffled flask) was inoculated to an initial OD600 of 0.0005. The culture was incubated at 28° C. and 250 rpm ~24 hours. Cells were harvested at an OD600=0.5-1. Instead of resuspending 10 ml of culture in 1 ml of Solution 2 as in the manufacturer's instructions (OD600~10), 10 ml of SPV140 culture was resuspended in 0.5 ml (OD600~20-30). All Solution 2 aliquots were slowly frozen to -80° C. by placing the tubes in a closed Styrofoam box before putting in the -80° C. freezer. 50 μl aliquots of competent cells in 1.7 ml Eppendorf tubes were thawed on ice, DNA eluted in water was added directly to the cells, and 500 μl of Solution 3 was used to suspend the cells with gentle pipetting. Tubes were incubated at 28° C. for 3 hours with gentle vortexing every 30 minutes. The entire transformation mixture was plated on CM glucose -ura agar plates. Positive integrants were found to be site-specific and genotyping was conducted by check PCR.

Z11-16OH Functional Expression Assay

Positive isolates were repatched onto YPD, grown overnight, and then stored at 4° C. Strains were inoculated from patch plates into 2 ml of YPD in 24 deepwell plates (square well, pyramid bottom). Replicate inoculations were made from each patch. Negative control strains were struck out on YPD from glycerol stocks and individual colonies were used to inoculate. Deepwell plates were incubated at 28° C. and 250 rpm in the Infors Multitron refrigerated flask shaker for 24 hrs. After 24 hrs of incubation, a 0.85 ml volume of each culture was pelleted by centrifugation at 800xg. Each pellet

was resuspended in either 2 ml of YPD or Semi-defined medium (described in Table 17 below). 10 g/L methyl palmitate (pre-warmed to ~50° C.) was added to cultures. All cultures were incubated for 48 hours before endpoint sampling. Final cell densities were measured with the Tecan Infinite 200pro plate reader. 1.5 ml (alcohol analysis) or 500 µl (lipid analysis) was transferred to 1.7 ml microcentrifuge tubes and pelleted. Supernatant was transferred to clean tubes and samples were processed as described below.

TABLE 17

Semi-defined (C:N = 80) medium composition. Components of the semi-defined base medium used to induce lipid storage are described.		
Media Components	Conc.	Units
Yeast Extract	2	g/L
Peptone	1	g/L
Potassium phosphate buffer pH 7	0.1	M
YNB w/o aa, NH4	1.7	g/L
Glucose	60	g/L
Glycerol	5	g/l

Z9-16OH Functional Expression Assay

SPV300 (negative control) and SPV471 were struck out onto YPD agar plates, grown overnight, and then stored at 4° C. Strains were inoculated from colonies into 2 ml of YPD and incubated at 28° C. and 250 rpm in 14 ml round bottom culture tubes for 8 hours. After incubation, 2 ml of culture was used to inoculate 20 ml of YPD in a 125 ml baffled shake flask. Shake flasks were incubated 24 hrs at 28° C. and 250 rpm. After incubation, cell density in shake flasks was measured using a Tecan Infinite 200pro plate reader. An appropriate volume of culture was pelleted in order to resuspend cells in 25 ml of Semi-defined C:N=80 medium (see Table 17 above) at an initial OD600=1 (~1 gDCW/L) or 4 (~4 gDCW/L). The resuspended culture was added to 250 ml baffled shake flasks. Neat methyl palmitate was added at 10 g/L final concentration after pre-heating to 50° C. After substrate addition, flasks were incubated at 28° C. and 250 rpm for two days. At 12, 18, 24, 36, 42, and 48 hours 500 µl (lipid analysis) and 1.5 ml (alcohol analysis) samples were taken in 1.7 ml microcentrifuge tubes. Samples were pelleted and the supernatant was transferred to a clean microcentrifuge tube.

Metabolite Extraction and GC-MS Detection

Alcohol Analysis

The pelleted cells (in 1.5 mL plastic tubes), usually about 10 mg to 80 mg, were resuspended in methanol containing 5% (w/w) of sodium hydroxide. The alkaline cell suspension was transferred into a 1.8 mL crimp vial. The mixture was heated for 1 h in the heat block at 90° C. Prior to acidification with 400 µL 2.5 N HCl the vial was allowed to cool to room temperature. 500 µL chloroform containing 1 mM methyl heptadecanoate were added and the mixture was shaken vigorously, then both aqueous and organic phase were transferred into a 1.5 mL plastic tube. The mixture was centrifuged at 13,000 rpm, afterwards 450 µL of the organic phase were transferred into a GC vial. The organic phase was evaporated in a heat block at 90° C. for 30 min. The residue was dissolved in 50 µL N,O-Bis(trimethylsilyl) trifluoroacetamide containing 1% trimethylchlorosilane. Prior to transfer into glass inserts the mixture was heated 5 min at 90° C. The samples were analyzed by GC-MS (Table 18).

TABLE 18

GC-MS parameters	
System	Agilent 6390 N GC, ChemStation G1701EA E.02.01.1177
Column	DB23 30 m × 25 µm × 25 µm Pressure = 11.60 psi; Flow = 0.6 mL/min
Inlet	Heater = 250° C.; Pressure = 11.74 psi;
	Total Flow {He} = 111 mL/min
Carrier	He @ 29 cm/sec, 11.60 psi
Signal	Data rate = 2 Hz/0.1 min
Oven	150° C. for 1 min Ramp 12° C./min to 220° C., hold 3 min Ramp 35° C./min to 300° C., hold 4 min
Injection	Splitless, 250° C.
Detector	Initial strain screening and first technical triplicate: HP 5973 MSD in SIM mode (m/z: 208.0, 297.3 and 387.3), SPV488/SPV490 alcohol quantification: HP 5973 MSD in SIM mode (m/z: 284.0 and 297.3), 100 msec Dwell, EMV mode: Gain factor 1, 2.4 min solvent delay, 3.09 cycles/sec
Sample	Injection volume = 1 uL

Lipid Analysis

Total lipid composition was based on modified procedures by Moss et al. (1982) and Yousuf et al (2010). The pelleted cells (in 1.5 mL plastic tubes), usually about 10 mg to 80 mg, were resuspended in methanol containing 5% (w/w) of sodium hydroxide. The alkaline cell suspension was transferred into a 1.8 mL glass crimp GC-vial. The mixture was heated for 1 h in the heat block at 90° C. Prior to acidification with 400 µL 2.5 N HCl, the vial was allowed to cool to room temperature. 500 µL chloroform containing 1 mM methyl heptadecanoate were added and the mixture was shaken vigorously, then both aqueous and organic phase were transferred into a 1.5 mL plastic tube. The mixture was centrifuged at 13,000 rpm, afterwards 450 µL of the organic phase was transferred into a new 1.8 mL glass screw-cap GC-vial. After cooling to room temperature residual fatty acid methyl esters and free fatty acids were dissolved and derivatized in methanol containing 0.2 M TMSH (trimethylsulfonium hydroxide)(Table 19).

TABLE 19

GC-MS parameters	
System	Agilent 6890 GC, ChemStation Rev. B.03.02 (341)
Column	J&W DB-23 30 m × 25 mm × 25 µm Pressure = 16 psi; Flow = 0.9 mL/min; Run Time = 14.4 min
Inlet	Heater = 240° C.; Pressure = 16 psi; Total Flow {He} = 31.4 mL/min
Carrier	H ₂ @ 1 mL/min, 9 psi, 35 cm/sec
Signal	Data rate = 2 Hz/0.1 min
Oven	150° C. for 1 min Ramp 12° C./min to 220° C., hold 3 min Ramp 35° C./min to 240° C., hold 6 min Equilibration Time: 2 min
Injection	Split, 240° C. Split ratio -30:1; 29.1 mL/min
Detector	FID, 240° C. H ₂ @ 35.0 mL/min, Air @ 350 mL/min;; Electrometer {Lit Offset} @ 2.0 pA
Sample	Injection volume = 1 uL

SEQUENCE LISTING	
SEQ ID NO: 1	<i>Agrotis segetum</i> FAR_ <i>S. cerevisiae</i> codon opt ATGCCAGTTTTGACTTCTAGAGAAGATGAAAAGT TGTCAGTTCAGAAATTTACGCTGGTAAATCTATCTTCGTTACAG GTGGTACTGGTTTCTTGGGTAAAGTTTTTATTGAAAAGTTGTTGT ACTGTTGTCCAGATATTGATAAAATCTATATGTTAATTAGAGAAA AGAAAAATTTGTCTATTGATGAAAGAATGTCAAAGTTCTTGGATG ATCCATTATTTTCTAGATTGAAGGAAGAAAGACCTGGTGACTTGG AAAAGATTGTTTTGATTCCAGGTGACATTACAGCTCCAAATTTGG GTTTATCAGCAGAAAACGAAAGAATTTTGTTAGAAAAAGTTTCTG TTATTATTAATTCAGCTGCAACTGTTAAGTTTAATGAACCATTGC CAATCGCTTGGAAGATTAATGTTGAAGGTACAAGAATGTTGTTGG CATTGTCTAGAAGAAATGAAGAGAATCGAAGTTTTTATTCATATTT CTACTGCTTACTCAAATGCATCTTCAGATAGAATCGTTGTTGATG AAATCTTGTATCCAGCTCCAGCAGATATGGATCAAGTTTACCAAT TGGTTAAAGATGGTGTTACAGAAGAAGAACTGAAAGATTGTTGA ACGGTTTGCCAAACACTTACACTTTTACTAAGGCTTTGACAGAAC ATTTGGTTGCAGAACATCAAACATACGTTCCAACATATCATCATCA GACCATCTGTTGTTGCTTCAATTAAGATGAACCAATCAGAGGTT GGTTATGTAATTGGTTTGGTGCTACAGGTATCTCTGTTTTTACTG CAAAGGGTTTGAACAGAGTTTGTGTTGGGTAAAGCTTCAAACATCG TTGATGTTATCCCAGTTGATTACGTTGCAAATTTGGTTATTGTTG CTGGTGCAAATCTGGTGGTCAAAAATCAGATGAATTAAAGATCT ATAACTGTTGTTCTTCAGATTGTAACCCAGTTACTTTGAAGAAAA TTATTAAAGAGTTTACTGAAGATACTATTAATAAATAAGTCTCATA TTATGCCATTGCCAGGTTGGTTCGTTTTTACTAAGTACAAGTGGT TGTTGACATTGTTAACTATTATTTTTCAAATGTTACCAATGTATT TGGCTGATGTTTACAGAGTTTGTACAGGTAAAATCCAAGATACA TGAAGTTGCATCATTTGGTTATTCAAACAAGATTGGGTATCGATT TCTTTACTTCTCATTCATGGGTTATGAAGACAGATAGAGTTAGAG AATTATTCGGTTCCTTGTCATTGGCAGAAAAGCATATGTTTCCAT GTGATCCATCTTCAATCGATTGGACAGATTATTTGCAATCATACT GTTACGGTGTTAGAAGATTTTTTGAAAAGAAGAAATAA
SEQ ID NO: 2	<i>Spodoptera littoralis</i> FAR1_ <i>S. cerevisiae</i> codon opt ATGGTTGTTTTGACTTCAAAGGAAAAATCAAACA TGTCTGTTGCTGATTTCTACGCTGGTAAATCTGTTTTTATTACAG GTGGTACTGGTTTCTTGGGTAAAGTTTTTATTGAAAAGTTGTTGT ACTCATGTCCAGATATTGATAAAATCTATATGTTGATCAGAGAAA AGAAAGGTCAATCTATCAGAGAAAGATTAACTAAAATTGTTGATG ATCCATTGTTTAATAGATTGAAGGATAAGAGACCAGATGATTTGG GTAAAATCGTTTTGATCCCAGGTGACATCACAGTTCCAGGTTTGG GTATTTCTGAAGAAAACGAAACAATCTTGA CTGAAAAAGTTTCAG TTGTTATTTCATTCTGCTGCAACTGTTAAGTTTAATGAACCATTGG CTACTGCATGGAACGTTAACGTTGAAGGTACAAGAATGATCATGG CATTATCAAGAAGAATGAAGAGAATCGAAGTTTTTATTCATATTT CTACTGCTTACACTAACACAAACAGAGCAGTTATTGATGAAGTTT TGTATCCACCACCAGCTGATATCAACGATGTTCATCAACATGTTA AAAATGGTGTTACAGAAGAAGAACTGAAAAGATTTTGAACGGTA GACCAAACACTTACACTTTTACTAAGGCTTTGACTGAACATTTGG TTGCAGAAAACCAATCATACATGCCAACAAATCATTGTTAGACCAT CTATTGTTGGTGCTATTAAAGATGATCCAATTAGAGGTTGGTTGG CTAATTGGTATGGTGCAACAGGTTTGTCAGTTTTTACTGCAAAGG GTTTGAACAGAGTTATATATGGTCATTCTAACCATGTTGTTGATT TGATTCCAGTTGATTACGTTGCTAATTTGGTTATTGTTGCTGGTG CAAAGACATACCATTCAAACGAAGTTACTATCTATAACTCTTGTT CTTCATCTTGTAACCCAATCACTATGAAGAGATTGGTTGGTTTGT TTATTGATTACACAGTTAAGCATAAGTCATACGTTATGCCATTGC CAGGTTGGTATGTTTACTCTAACTACAAGTGGTTGGTTTTCTTGG TTACTGTTATTTTCCAAGTTATTCCAGCTTACTTAGGTGACATTG GTAGAAGATTGTTAGGTAAAAATCCAAGATACTACAAGTTGCAAA ATTTGGTTGCTCAAACACAAGAAGCAGTTCATTTCTTTACATCAC ATACTTGGGAAATTAAATCAAAGAGA ACTTCTGAATTGTTTTTCAT CTTTGTCTTTGACAGATCAAAGAATGTTTCCATGTGATGCTAACA GAATCGATTGGACAGATTACATCACTGATTACTGTTCTGGTGTTA GACAATTTTTTGAAAAGATTAAATAA
SEQ ID NO: 3	<i>Helicoverpa armigera</i> FAR3_ <i>S. cerevisiae</i> codon. opt ATGGTTGTTTTGACTTCAAAGGAAACAAAGCCAT CTGTTGCTGAATTTTACGCTGGTAAATCAGTTTTTATTACAGGTG GTACTGGTTTCTTGGGTAAAGTTTTTATTGAAAAGTTGTTGTACT CTTGTCAGATATTGAAAATATCTATATGTTGATCAGAGAAAAGA AAGGTTTGTCAGTTTCTGAAAGAATTAAACAATTTTTAGATGATC CATTGTTTACAAGATTGAAGGATAAGAGACCAGCTGATTTGGAAA AGATTGTTTTGATCCCAGGTGACATCACTGCACCAGATTTGGGTA TTAATTCTGAAAACGAAAAGATGTTGATTGAAAAAGTTTCAGTTA TTATTCATTCTGCTGCAACTGTTAAGTTTAAATGAACCATTACCAA CAGCTTGGAAGATTAATGTTGAAGGTACTAGAATGATGTTGGCAT TGTCAAGAAGAATGAAGAGAATCGAAGTTTTTATTCATATTTCTA CAGCTTACACTAACACAAACAGAGAAGTTGTTGATGAAATCTTGT

- continued

SEQUENCE LISTING	
	ATCCAGCTCCAGCAGATATCGATCAAGTTCATCAATACGTTAAGG ATGGTATCTCAGAAGAAGATACTGAAAAGATTTTGAACGGTAGAC CAAACACTTACACTTTTACTAAGGCTTTGACAGAACATTTGGTTG CTGAAAATCAAGCATACGTTCCAACATTATTGTTAGACCATCTG TTGTTGCTGCAATTAAAGATGAACCATTGAAAGGTTGGTTGGGTA ATTGGTTTGGTGCTACAGGTTTGACTGTTTTTACAGCAAAGGTT TGAACAGAGTTATATATATGGTCATTCTTCATACATCGTTGATTGA TCCCAGTTGATTACGTTGCTAATTTGGTTATTGCTGCAGGTGCAA AATCTTCAAAGTCAACAGAATTGAAGGTTTACAACTGTTGTTCTT CATCTTGTAACCCAGTTACTATCGGTACATTGATGTCAATGTTCTG CTGATGATGCAATTAAACAAAAATCTTACGCTATGCCATTGCCAG GTTGGTACATTTTTACAAAGTACAAGTGGTTGGTTTTGTTGTTGA CATTTTTGTTCCAAGTTATTCCAGCATACGTTACTGATTTGTCAA GACATTTGATCGGTAAATCTCCAAGATACATCAAGTTGCAATCAT TGGTTAACCAAACCTAGATCATCTATCGATTCTTTACAAACCATT CTTGGGTTATGAAAGCTGATAGAGTTAGAGAATTGTACGCTTCAT TGTCTCCAGCTGATAAGTACTTATTCCCATGTGATCCAAC TGATA TCAACTGGACACATTACATCCAAGATTACTGTTGGGGTGTTAGAC ATTTCTTGAAAAAGAAATCTTACGAATAA
SEQ ID NO: 4	pOLE1 cassette CTTGCTGAAAAGATGATGTTCTGAGGTATTCTGATCGCTAGCTTCATACGCTTTT AACAAAAGTAAGCTTTTCGTTTGCAGGTTTGTTACTTTTCTGTACGAGATGATA TCGCTAAGTTTATAGTCATCTGTGAAATTTCTCAAAAACCTCATGGTTTCTCCAT CACCCATTTTTTCATTTTCAATTTGCCGGGCGGAAAAAAAAAAGGAAAAAAAAAAAA AAAAAATAAATGACACATGGAAATAAGTCAAGGATTAGCGGATATGTAGTTCCA GTCCGGGTTATACCATCACGTGATAATAAATCCAAATGAGAATGAGGGTGTCATA TCTAATCATTATGCACGTCAAGATTCTCCGTGACTATGGCTCTTTTCTGAAGCAT TTTTTCGGGCGCCCGGTGGCCAAAACTAACTCCGAGCCCGGGCATGTCCCGGGGT TAGCGGGCCCAACAAAGGCGCTTATCTGGTGGGCTTCCGTAGAAGAAAAAAGCT GTTGAGCGAGCTATTTCGGGTATCCCAGCCTTCTCTGCAGACCGCCCCAGTTGGC TTGGCTCTGGTGCTGTTTCGTTAGCATCACATCGCCTGTGACAGGCAGAGTAATA ACGGCTTAAGGTTCTCTTCGCATAGTCGGCAGCTTTCTTTCCGACGTTGAACACT CAACAAACCTTATCTAGTGCCCAACCAGGTGTGCTTCTACGAGTCTTGCTCACTC AGACACACCTATCCCTATTGTTACGGCTATGGGGATGGCACACAAAGGTGGAAAT AATAGTAGTTAACAAATATATGCAGCAAATCATCGGCTCCTGGCTCATCGAGTCTT GCAAATCAGCATATACATATATATATGSGGGGCAGATCTTGATTCAATTTATTGTTT TATTTCCATCTTTCCTACTTCTGTTTCCGTTTATATTTTGATTACGTAGAATAG AACATCATAGTAATAGATAGTTGTGGTGATCATATTATAAACAGCACTAAAACAT TACAACAAAGAATGCCAACTTCTGGAATACTATTGAATTGATTGACGACCAATT TCCAAAGGATGACTCTGCCAGCAGTGGCATTGTCGACACTAGTGCGGCCGCTCAC ATATGAAAGTATATACCCGCTTTTGTACACTATGTAGCTATAATTCAATCGTATT ATTGTAGCTCCGCACGACCATGECTTAGAAATATCCGCAGCGCG
SEQ ID NO: 5	Extended OLE1 promoter region CTTGCTGAAAAGATGATGTTCTGAGGTATTCTGATCGCTAGCTTGATACGCTTTT AACAAAAGTAAGCTTTTCGTTTGCAGGTTTGTTACTTTTCTGTACGAGATGATA TCGCTAAGTTTATAGTCATCTGTGAAATTTCTCAAAAACCTCATGGTTTCTCCAT CACCCATTTTTTCATTTTCAATTTGCCGGGCGGAAAAAAAAAAGGAAAAAAAAAAAA AAAAAATAAATGACACATGGAAATAAGTCAAGGATTAGCGGATATGTAGTTCCA GTCCGGGTTATACCATCACGTGATAATAAATCCAAATGAGAATGAGGGTGTCATA TCTAATCATTATGCACGTCAAGATTCTCCGTGACTATGGCTCTTTTCTGAAGCAT TTTTTCGGGCGCCCGGTGGCCAAAACTAACTCCGAGCCCGGGCATGTCCCGGGGT TAGCGGGCCCAACAAAGGCGCTTATCTGGTGGGCTTCCGTAGAAGAAAAAAGCT GTTGAGCGAGCTATTTCGGGTATCCCAGCCTTCTCTGCAGACCGCCCCAGTTGGC TTGGCTCTGGTGCTGTTTCGTTAGCATCACATCGCCTGTGACAGGCAGAGTAATA ACGGCTTAAGGTTCTCTTCGCATAGTCGGCAGCTTTCTTTCCGACGTTGA
SEQ ID NO: 6	OLE1 promoter region ACACTCAACAAACCTTATCTAGTGCCCAACCAGGTGTGCTTCTACGAGTCTTGCT CACTCAGACACACCTATCCCTATTGTTACGGCTATGGGGATGGCACACAAAGGTG GAAATAATAGTAGTTAACAAATATATGCAGCAAATCATCGGCTCCTGGCTCATCGA GTCTTGCAAATCAGCATATACATATATATATGSGGGGCAGATCTTGATTCATTTAT TGTTCTATTTCCATCTTTCCTACTTCTGTTTCCGTTTATATTTTGATTACGTAG AATAGAACATCATAGTAATAGATAGTTGTGGTGATCATATTATAAACAGCACTAA AACATTACAACAAAGA
SEQ ID NO: 7	OLE1 27 aa leader ATGCCAATTCTGGAATACTATTGAATTGATTGACGACCAATTTCCAAAGGATG ACTCTGCCAGCAGTGGCATTGTTCGAC
SEQ ID NO: 8	Vsp13 terminator region TCACATATGAAAGTATATACCCGCTTTTGTACACTATGTAGCTATAATTCAATCG TATTATTGTAGCTCCGCACGACCATGCCCTTAGAAATATCCGCAGCGCG

- continued

SEQUENCE LISTING

SEQ ID NO: 9	<i>T. ni</i> desaturase ATGGCTGTGATGGCTCAAACAGTACAAGAAACGGCTACAGTGTTGGAAGAGGAAG CTCGCACAGTGACTCTTGTGGCTCCAAAGACAACGCCAAGGAAATATAAATATAT ATACACCAACTTTCTTACATTTTCATATGCGCATTTAGCTGCATTATACGGACTT TATTTGTGCTTCACCTCTGCGAAATGGGAAACATTGCTATTCTCTTTCGTACTCT TCCACATGTCAAATATAGGCATCACCGCAGGGGCTCACCGACTCTGGACTCACAA GACTTTCAAAGCCAAATTGCCTTTGGAAATTGTCCTCATGATATTCAACTCTTTA GCCTTTCAAAACACGGCTATTACATGGGCTAGAGAACATCGGCTACATCACAAAT ACAGCGATACTGATGCTGATCCCCACAATGCGTCAAGAGGGTTCTTCTACTCGCA TGTTGGCTGGCTATTAGTAAAAAAACATCCCGATGTCTGAAATATGGAAAACT ATAGACATGTCGGATGTATACAATAATCCTGTGTTAAAATTTAGAAAAAGTACG CAGTACCCTTAATTGGAACAGTTTGTTTTGCTCTGCCAACTTTGATTCCAGTCTA CTGTTGGGGCGAATCGTGGAACAACGCTTGCGACATAGCCTTATTTGATACATA TTCAATCTTAACGTGACTTTCCTAGTCAACAGTGCTGCGCATATCTGGGGGAATA AGCCTTATGATAAAAGCATCTTGCCCGCTCAAAACCTGCTGGTTTCCTTCTAGC AAGTGGAGAAGGCTTCATAATTACCATCACGTCTTTCATGGGATTACCGCACA GCAGAATTAGGGAATAACTTCCTGAATTTGACGACGCTGTTTATTGATTTTGTG CCTGGTTTGGATGGCTTATGACTTGAAGTCTGTATCAGAGGATATTATAAAACA GAGAGCTAAACGAACAGGTGACGGTTCCTCAGGGGTCAATTTGGGGATGGGACGAC AAAGACATGGACCGGATATAAAATCTAAAGCTAACATTTTTTATGCTAAAAAGG AATGA
SEQ ID NO: 10	<i>A. segetum</i> desaturase ATGGCTCAAGGTGTCCAAACAACACTACGATATTGAGGGAGGAGGAGCCGTATTGA CTTTCGTGGTACCTCAAGAACCGAGAAAGTATCAAATCGTGTACCCAAACCTTAT CACATTTGGGTACTGGCATATAGCTGGTTTATACGGGCTATATTTGTGCTTTACT TCGGCAAAATGGCAACAATTTTATTCAAGTTTCATGCTCGTTGTGTTAGCAGAGT TGGGAATAACAGCCGGCGCTCACAGGTTATGGGCCACAAAACATATAAAGCGAA GCTTCCCTTACAAATTATCCTGATGATACTGAACTCCATTGCCTTCCAAAATTCC GCCATTGATTGGGTGAGGG ACCACCGTCTCCATCATAAGTACAGTGACACTGATGCAGACCCTCACAATGCTAC TCGTGGTTTCTTCTATTCTCATGTTGGATGGTTGCTCGTAAGAAAACATCCAGAA GTCAAGAGACGTGGAAAGGAACCTGACATGTCTGATATTTACAACAATCCAGTGC TGAGATTTCAAAGAAGTATGCTATACCCCTTCATCGGGGCAATGTGCTTCGGATT ACCAACTTTTATCCCTGTTTACTTCTGGGGAGAAACCTGGAGTAATGCTTGGCAT ATCACCATGCTTCGGTACATCCTCAACCTAAACATTACTTTCTTGGTCAACAGTG CTGCTCATATCTGGGGATACAAACCTTATGACATCAAAATATTGCCTGCCCAAAA TATAGCAGTTTCCATAGTAACCGGCGGCGAAGTTTCCATAACTACCACCACGTTT TTTCCTTGGGATTATCGTGCAGCAGAATTGGGGAACAATTATCTTAATTGACGA CTAAGTTCATAGATTCTTTCGCTTGGATCGGATGGGCTTACGATCTTAAGACGGT GTCCAGTGATGTTATAAAAAAGTAAGGCGGAAAGAACTGGTGATGGGACGAATCTT TGGGGTTTAGAAGACAAAGGTGAAG AAGATTTTTTGAAAATCTGGAAAGACAATTAA
SEQ ID NO: 11	<i>T. pseudonana</i> desaturase ACTAGTATGGACTTTCTCTCCGGCGATCCTTTCCGGACAC TCGTCTTGCAGCACTTGTTGTCATCGGATTTGCTGCGGCGTGGC AATGCTTCTACCCGCCGAGCATCGTCGGCAAGCCTCGTACATTAA GCAATGGTAAACTCAATACCAGAATCCATGGCAAATTGTACGACC TCTCATCGTTTCAGCATCCAGGAGGCCCCGTGCTCTTTCTCTTG TTCAAGGTCGCGACGGAACAGCTCTATTTGAGTCACACCATCCCT TCATACCTCGAAAGAATCTACTTCAGATCCTCTCCAAGTACGAGG TTCCGTCGACTGAAGACTCTGTTTTCCTTCATCGCCACCCTAGACG AACTCAATGGTGAATCTCCGTACGATTGGAAGGACATTGAAAATG ATGATTTTCGTATCTGACCTACGAGCTCTCGTAATTGAGCACTTTT CTCCTCTCGCCAAGGAAAGGGGAGTTTCACTCGTTGAGTCGTCTGA AGGCAACACCTCAGCGGTGGATGGTGGTTCTACTGCTCCTTGCGT CGTTCTTCCCTCAGCATCCCATTATATTTGAGTGGTTGTTGAGACTT TCGTTGTGCTCACTCCCATCCTCGCTTGGCTGGCGGTTGTCAATT ACTGGCACGATGCTACTCACTTTGCATTGAGCAGCAACTGGATTT TGAATGCTGCGCTCCCATATCTCCTCCCTCTCCTATCGAGTCCGT CAATGTGGTATCATCATCACGTCATTGGACATCACGCATACACCA ACATTTCCAAAAGAGATCCAGATCTTGCTCACGCTCCACAACCTCA TGAGAGAACACAAGAGTATCAAATGGAGACCATCTCACTTAAATC AAACACAGCTTCCGCGGATTCTCTTCATCTGGTCGATTGCAGTCG GTATTGGGTTGAACTTACTGAACGACGTGAGAGCACTAACCAAGC TTTCATACAACAACGTTGTTTCGGGTGGAGAAGATGTCATCGTCGC GAACATTACTCCATTTCTTGGACGTATGTTGCACATCTTTGTGA CTACACTTTGGCCCTTTTTTGGCGTTTCCGGTGTGGAAGGCCATCG TTTGGGCGACTGTACCGAATGCCATACTGAGTTTGTGCTTCATGC TGAATACGCAAATCAATCACCTCATCAACACGTGTGCACATGCTT CCGATAACAACCTTTTACAAGCATCAAGTTGTAAGTGTCTCAGAACT TTGGCCGATCAAGTGCTTTTGTCTTCATCTCTCGGGAGGTCTCA ACTACCAAATTGAACATCATTGTGTTGCCGACGGTGAACCATTGCC

- continued

SEQUENCE LISTING	
	ATTTGCCAGCTTTTGCCCCGGGTGTAGAGCGTTTGTGTAAGAAAC ACGGGGTGACATACAACTCTGTTGAAGGATACAGAGAGGCCATCA TTGCACACTTTGCACATACCAAAGATATGTCGACGAAGCCTACTG ATTGA
SEQ ID NO: 12	<i>A. transitella</i> desaturase ATGGTCCCTAACAAAGGGTTCCAGTGACGTTTTGTCTGAACATTCTGAGCCCCAGT TCACTAAACTCATAGCTCCACAAGCAGGGCCGAGGAAATACAAGATAGTGTATCG AAATTTGCTCACATTCTGGCTATTGGCACTTATCAGCTGTTTATGGGCTCTACTTG TGCTTTACTTGTGCGAAATGGGCTACCATCTTATTTGCATTTTTCTTATACGTGA TCGCGGAAATCGGTATAACAGGTGGCGCTCATAGGCTATGGGCACATCGGACTTA TAAAGCCAAGTTGCCTTTAGAGATTTTGTACTCATAATG AACTCTATTGCCTTCCAAGACACTGCTTTCACCTGGGCTCGTGATCACCGCCTTC ATCACAAATATTTCGATACTGACGCTGATCCCCACAATGCTACCAGAGGGTTTTT CTATTCACATGTAGGCTGGCTTTTGGTGAAGAAACACCCTGAAGTCAAAGCAAGA GGAAAATACTTGTCTGTAGATGATCTTAAGAATAATCCATTGCTTAAATTCCAA AGAAATACGCTATTCTAGTTATAGGCACGTTATGCTTCCTTATGCCAACATTTGT GCCCGTATACTTCTGGGGCGAGGGCATCAGCACGGCCTGGAACATCAATCTATTG CGATACGTCATGAATCTTAACATGACTTCTTAGTTAACAGTGCAGCGCATATCT TTGGCAACAAACCATACGATAAGAGCATAGCCTCAGTCCAAAATATTTCAGTTAG CTTAGCTACTTTTGGCGAAGGATTCCATAATTACCATCACACTTACCCCTGGGAT TATCGTGCGGCAGAAATTAGGAAATAATAGGCTAAATATGACTACTGCTTTCATAG ATTTCTTCGCTTGGATCGGCTGGGCTTATGACTTGAAGTCTGTGCCACAAGAGGC CATTGCAAAAAGGTGTGCGAAAACCTGGCGATGGAACGGATATGTGGGGTCGAAAA AGATAA
SEQ ID NO: 13	<i>H. zea</i> desaturase ATGGCCCAAAGCTATCAATCAACTACGGTTTTTGAGTGAGGAGAAAGAACTAACAC TGCAACATTTGGTGCCCCAAGCATCGCCCAGGAAGTATCAAATAGTGTATCCGAA CCTCATTACGTTTGGTACTGGCACATAGCCGACTTTATGGCCTTTACTTGTGC TTCAC TTCTGCTAAATGGGCTACGATTTTATTCAGCTACATCCTCTTCGTGTTAG CAGAAATAGGAATCACGGCTGGCGCTCACAGACTCTGGGCCCACAAAAC TTACAA AGCGAAACTACCATTAGAAATACTCTTAATGGTATTCAACTCCATCGCTTTTCAA AACTCAGCCATTGACTGGGTGAGGGACCACCGACTCCACCATAAGTATAGCGATA CAGATGCTGATCCCCACAATGCCAGCCGAGGGTTC TTTTATTCCCATGTAGGATG GCTACTTGTGAGAAAACATCCTGAAGTCAAAAAGCGAGGGAAGAACTCAATATG TCCGATATTTACAACAATCCTGTCTCGGTTTCAGAAAAAATACGCCATACCCT TCATTGGGGCTGTTTGTTTCGCCTTACCTACAATGATACCTGTTTACTTCTGGGG AGAAACCTGGTCCAATGCTTGGCATATCACCATGCTTCGCTACATCATGAACCTC AATGTCACCTTTTTGGTAAACAGCGCTGCTCATATATGGGGAAACAAGCCTTATG ACGCAAAAATATTACCTGCACAAAATGTAGCTGTGTGGTTCGCCACTGGTGGAGA AGGTTTCCATAATTACCACCATGTCTTCCCTGGGATTATCGAGCAGCGGAACTC GGTAACAATAGCCTCAATCTGACGACTAAATT CATAGATT TATTCGAGCAATCG GATGGGCATATGATCTGAAGACGGTTTCGGAGGATATGATAAAACAAAGGATTAA ACGCACTGGAGATGGAACGGATCTTTGGGGACACGAACAAAAC TGTGATGAAGTG TGGGATGTAAAAGATAAATCAAGTTAA
SEQ ID NO: 14	mCherry <i>C. tropicalis</i> optimized ATGGTTTTCTAAGGGTGAAGAAGACAACATGGCAATCATCAAGGAATTTATGCGTT TTAAGGTCCATATGGAAGGCTCCGTTAACGGCCACGAGTTCGAGATCGAGGGAGA AGGTGAGGGTAGACCATACGAAGGTACTCAAACCGCCAAGTTGAAAGTTACAAAG GGTGGTCCATTGCCATTTGCTTGGGATATCTTGTCCCCACAATTTATGTACGGAT CAAAGGCATATGTCAAGCATCCTGCCGACATCCCAGATTACTTGAAGTTATCCTT TCCAGAAGGTTTTAAGTGGGAGAGAGTTATGAACTTTGAAGATGGCGGAGTTGTT ACTGTTACTCAGGACTCTTCCTTGCAAGATGGTGAATTTATCTATAAAGTGAAAT TGAGAGGTACTAACTTCCATCCGACGGTCCAGTCATGCAAAAGAAGACAATGGG TTGGGAGGCTTC TCCGAAAGAATGTACCCAGAAGACGGTGCATTGAAAGGTGAA ATCAAGCAACGTTTAAAGTTGAAGGACGGTGGTCACTACGATGCCGAGGTCAAGA CCACTTATAAGGCTAAGAAGCCAGTCCAATTGCCAGGTGCTTATAACGTTAACAT CAAGTTAGATATTACTTCACACAACGAAGACTACACAATCGTTGAACAATATGAA AGAGCCGAAGGTAGACATTCTACCGGCGGCATGGACGAGTTATATAAGTAG
SEQ ID NO: 15	CaOLE1-A. <i>segetum</i> Z11 desaturase ATGACTACAGTTGAACAACTTGAAACTGTTGATATCACTAAATTGAATGCCATTG CTGCTGGTACTAATAAGAAGGTGCCAATGGCTCAAGGTGTCAAACAACTACGAT ATTGAGGGAGGAAGAGCCGTCATTGACTTTCGTGGTACCTCAAGAACCGAGAAAG TATCAAATCGTGTACCCAAACCTTATCACATTTGGGTACTGGCATATAGCTGGTT TATACGGGCTATATTTGTGCTTTACTTCGGCAAAATGGCAAACAATTTTATTCAG TTTCATGCTCGTTGTGTTAGCAGAGTTGGGAATAACAGCCGGCGCTCACAGGTTA TGGGCCCACAAAACATATAAAGCGAAGCTTCCCTTACAAATTATCTTAATGATAT TAAACTCCATTGCCTTCCAAAATTCCGCCATTGATTGGGTGAGGGACCACCGTCT CCATCATAAGTACAGTGACACTGATGCAGACCCTCACAAATGCTACTCGTGGTTTC TTCTATTCTCATGTTGGATGGTTGCTCGTAAGAAAACATCCAGAAGTCAAGAGAC GTGGAAAGGAACTTGACATGTCTGATATTTACAACAATCCAGTGTTAAGATTTCA AAAGAAGTATGCTATACCCCTTCATCGGGGCAATGTGCTTCGGATTACCAACTTTT ATCCCTGTTTACTTCTGGGGAGAAACCTGGAGTAATGCTTGGCATATCACCATGC TTCGGTACATCCTCAACCTAAACATTACTTTCCTTAGTCAACAGTGTCTGCTCATAT

- continued

SEQUENCE LISTING	
	CTGGGGATACAAACCTTATGACATCAAAATATTGCCTGCCCCAAATATAGCAGTT TCCATAGTAACCGGCGGCGAAGTTTCCATAACTACCACCACGTTTTTTTCCTTGGG ATTATCGTGCAGCAGAATTGGGGAACAATTATCTTAATTTGACGACTAAGTTCAT AGATTTCTTCGCTTGGATCGGATGGGCTTACGATCTTAAGACGGTGTCCAGTGAT GTTATAAAAAGTAAGGCGGAAAGAACTGGTGATGGGACGAATCTTTGGGGTTTAG AAGACAAAGGTGAAGAAGATTTTTTGAAAATCTGGAAAGACAATTAA
SEQ ID NO: 16	<i>A. segetum</i> Z11 desaturase ATGGCTCAAGGTGTCCAAACAACACTACGATATTGAGGGAGGAAGAGCCGTCATTGA CTTTTCGTGGTACCTCAAGAACCGAGAAAGTATCAAATCGTGTACCCAAACCTTAT CACATTTGGGTACTGGCATATAGCTGGTTTATACGGGCTATATTTGTGCTTTACT TCGGCAAAATGGCAAACAATTTTATTTCAGTTTCATGCTCGTTGTGTAGCAGAGT TGGGAATAACAGCCGCGCTCACAGGTTATGGGCCACAAAACATATAAAGCGAA GCTTCCCTTACAAATTATCTTAATGATATTAACCTCCATTGCCTTCCAAAATTCC GCCATTGATTGGGTGAGGGACCACCGTCTCCATCATAAGTACAGTGACACT GATGCAGACCCTCACAATGCTACTCGTGGTTTCTTCTATTCTCATGTTGGATGGT TGCTCGTAAGAAAACATCCAGAAGTCAAGAGACGTGGAAGGAACCTTGACATGTC TGATATTTACAACAATCCAGTGTTAAGATTTCAAAAGAAGTATGCTATACCCCTTC ATCGGGGCAATGTGCTTCGGATTACCAACTTTTATCCCTGTTTACTTCTGGGGAG AAACCTGGAGTAATGCTTGGCATATCACCATGCTTCGGTACATCCTCAACCTAAA CATTACTTTCTTAGTCAACAGTGCTGCTCATATCTGGGGATACAAACCTTATGAC ATCAAAATATTGCC TGCCCCAAATATAGCAGTTTCCATAGTAACCGGCGGCGAAG TTTCCATAACTACCACCACGTTTTTTTCCTTGGGATTATCGTGCAGCAGAATTGGG GAACAATTATCTTAATTTGACGACTAAGTTCATAGATTTCTTCGCTTGGATCGGA TGGGCTTACGATCTTAAGACGGTGTCCAGTGATGTTATAAAAAGTAAGGCGGAAA GAACTGGTGATGGGACGAATCTTTGGGGTTTAGAAGACAAAGGTGAAGAAGATTT TTTGAAAATCTGGAAAGACAATTAA
SEQ ID NO: 17	<i>A. transitella</i> Z11 desaturase ATGGTCCCTAACAGGGTTCCAGTGACGTTTTGTCTGAACATTCTGAGCCCCAGT TCACTAAACTCATAGCTCCACAAGCAGGGCCGAGGAAATACAAGATAGTGTATCG AAATTTGCTCACATTCCGGCTATTGGCACTTATCAGCTGTTTATGGGCTCTACTTG TGCTTTACTTGTGCGAAATGGGCTACCATCTTATTTGCATTTTTCTTATACGTGA TCGCGGAAATCGGTATAACAGGTGGCGCTCATAGGCTATGGGCACATCGGACTTA TAAAGCCAAGTTGCCTTTAGAGATTTTGTTACTCATAATGAATTCTATTGCCTTC CAAGACACTGCTTTCACCTGGGCTCGAGATCACCGCCTTCATCACAAATATTCCG ATACTGACGCTGATCCCCACAATGCTACCAGAGGGTTTTTCTATTCCATGTAGG CTGGCTTTTGGTGAAGAAACACCCCTGAAGTCAAAGCAAGAGGAAAATACTTGTCTG TTAGATGATCTTAAGAATAATCCATTGCTTAAATTCCAAAAGAAATACGCTATTCT TAGTTATAGGCACGTTATGCTTCCTTATGCCAACATTTGTGCCCGTATACTTCTG GGGCGAGGGCATCAGCACGGCCTGGAACATCAATCTATTGCGATACGTCATGAAT CTTAACATGACTTTCTTAGTTAACAGTGCAGCGCATATCTTTGGCAACAAACCAT ACGATAAGAGCATAGCCTCAGTCCAAAATATTTAGTTAGCTTAGCTACTTTTGG CGAAGGATTCCATAATTACCATCACACTTACCCCTGGGATTATCGTGCGGCAGAA TTAGGAAATAATAGGCTAAATATGACTACTGCTTTCATAGATTTCTTCGCTTGGA TCGGCTGGGCTTATGACTTGAAGTCTGTGCCACAAGAGGCCATTGCAAAAAGGTG TGCGAAAAC TGCGATGGAACGGATATGTGGGGTCGAAAAAGATAA
SEQ ID NO: 18	<i>T. ni</i> Z11 desaturase ATGGCTGTGATGGCTCAAACAGTACAAGAAACGGCTACAGTGTTGGAAGAGGAAG CTCGCACAGTGACTCTTGTGGCTCCAAAGACAACGCCAAGGAAATATAAATATAT ATACACCAACTTTCTTACATTTTCATATGCGCATTTAGCTGCATTATACGGACTT TATTTGTGCTTTCACCTCTGCGAAATGGGAAACATTGCTATTCTCTTTCGTACTCT TCCACATGTCAAATATAGGCATCACCGCAGGGGCTCACCGACTCTGGACTCACAA GACTTTCAAAGCCAAATTGCCTTTGGAAATTGTCCTCATGATATTCAACTCTTTA GCCTTTCAAACACGGCTATTACATGGGCTAGAGAACATCGGCTACATCACAAAT ACAGCGATAC TGATGCTGATCCCCACAATGCGTCAAGAGGGTTCCTTCTACTCGCA TGTTGGCTGGCTATTAGTAAAAAAACATCCCGATGTCTTAAATATGGAAAACT ATAGACATGTCGGATGTATACAATAATCCTGTGTAAATTTTCAAAAAAGTACG CAGTACCCTTAATTGGAACAGTTTGTTTTGCTCTTCCAACTTTGATTCCAGTCTA CTGTTGGGGCGAATCGTGGAACAACGCTTGGCACATAGCCTTATTTGATACATA TTCAATCTTAACGTGACTTTCCTAGTCAACAGTGCTGCGCATATCTGGGGGAATA AGCCTTATGATAAAAGCATCTTGCCCGCTCAAAACTTATTAGTTTCCTTCCTAGC AAGTGGAGAAGGCTTCCATAATTACCATCACGTCTTTCATGGGATTACCGCACA GCAGAATTAGGGAATAACTTCTTAAATTTGACGACGTTATTCTATTGATTTTTGTG CCTGGTTTGGATGGGCTTATGACTTGAAGTCTGTATCAGAGGATATTATAAAACA GAGAGCTAAACGAACAGGTGACGGTTCCTCAGGGGTCAATTTGGGGATGGGACGAC AAAGACATGGACCGCGATATAAAATCTAAAGCTAACATTTTTTTATGCTAAAAAGG AATGA
SEQ ID NO: 19	<i>H. zea</i> Z11 desaturase ATGGCCCAAAGCTATCAATCAACTACGGTTTTGAGTGAGGAGAAAGAACTAACAT TACAACATTTGGTGCCCCAAGCATCGCCAGGAAGTATCAAATAGTGTATCCGAA CCTCATTACGTTTGGTTACTGGCACATAGCCGACTTTATGGCCTTTACTTGTGC TTCACTTCTGCTAAATGGGCTACGATTTTATTTCAGCTACATCCTCTTCGTGTTAG CAGAAATAGGAATCACGGCTGGCGCTCACAGACTCTGGGCCACAAAACCTTACAA AGCGAAACTACCATTAGAAATACTCTTAATGGTATTCAACTCCATCGCTTTTCAA

- continued

SEQUENCE LISTING	
	AACTCAGCCATTGACTGGGTGAGGGACCACCGACTCCACCATAAGTATAGCGATA CAGATGCTGATCCCCACAATGCCAGCCGAGGGTTCTTTTATTCCCATGTAGGATG GCTACTTGTGAGAAAACATCCTGAAGTCAAAAAGCGAGGGAAAGAACTCAATATG TCCGATATTTACAACAATCCTGTCTTACGGTTTCAGAAAAAATACGCCATACCCCT TCATTGGGGCTGTTTGTTCGCCTTACCTACAATGATACCTGTTTACTTCTGGGG AGAAACCTGGTCCAATGCTTGGCATATCACCATGCTTCGCTACATCATGAACCTC AATGTCACCTTTTTGGTAAACAGCGCTGCTCATATATGGGGAAACAAGCCTTATG ACGCAAAAATATTACCTGCACAAAATGTAGCTGTGTCGGTCGCCACTGGTGGAGA AGGTTTCCATAATTACCACCATGTCTTCCCTGGGATTATCGAGCAGCGGAACTC GGTAACAATAGCCTCAATTTAACGACTAAATTCATAGATTTATTTCGCAGCAATCG GATGGGCATATGATTTAAAGACGGTTTCGGAGGATATGATAAAACAAAGATTAA ACGCACTGGAGATGGAACGGATCTTTGGGGACACGAACAAAACGTGTATGAAGTG TGGGATGTAAAAGATAAATCAAGTTAA
SEQ ID NO: 20	<i>O. furnacalis</i> Z9 desaturase ATGGCTCCTAATATTAAGGACGGAGCTGATTGAACGGAGTTTTATTTGAAGATG ACGCTAGCACCCCGATTATGCCCTTGCCACGGCCCCAGTCCAGAAAGCAGACAA CTATCCCAGAAAAC TAGTG TGAGAAACATCATACTCTTTGCATACCTTCACCTT GCCGCTGTGTATGGAGCATACCTATTCTTATTTTCAGCGAAATGGCAGACAGATA TTTTTGCCTACATTCTTTACGTGATCTCAGGACTCGGCATCACAGCGGGAGCCCA CCGCCTTTGGGCGCACAAAGTCATACAAGGCTAAGTGGCCACTTAGACTCATTCTT ATTATCTTCAACACTGTATCATTCCAGGACTCTGCTCTCGACTGGTCACGTGACC ACCGCATGCACCACAAATACTCGGAGACCGACGCCGACCCGCACAACGCGACTCG AGGGTTCTTCTCTCATATCGGCTGGTTATTAGTCCGCAAGCACCCGGAATTA AAGAGAAAGGGCAAGGGATTAGACTTAAGCGACTTGTATGCTGATCCCATCCTCC GTTTCCAGAAGAAGTACTATTTACTATTAATGCCTCTTGGCTGCTTCATCATGCC GACGGTGGTCCCGTGTA CTCTGGGGTGAGACTTGGA CTAAACGCTTCTTCGTC GCCGCGCTCTTCCGATACACCTTCATCCTCAATGTCACCTGGTTGGTCAACTCCG CCGCGCACAAAGTGGGGCCACAAGCCCTATGACAGCAGCATCAAGCCTTCCGAGAA CCTCTCAGTCTCCTTATTCGCGTTGGGCGAAGGATTCCACAAC TACCACACACA TTCCCCTGGGACTACAAAAC TGCCGAGCTCGGCAACAACAGACTCAATTTACAAA CAAAC TTCATCAACTTCTTCGCTAAAATCGGATGGGCTTACGACTTGAAAACGGT CTCCGACGAGATTATT CAGAATAGAGTCAAGCGCACAGGAGATGGCTCCCACCAC TTATGGGGTTGGGGCGACAAGGATCAACCTAAAGAGGAGGTAAACGCAGCCATTA GAATTAATCCTAAAGACGAGTAA
SEQ ID NO: 21	<i>L. capitella</i> Z9 desaturase ATGCCGCCGAACGTGACAGAGGCGAACGGAGTGTTATTTGAGAATGACGTGCAGA CTCCTGACATGGGGCTAGAAGTGGCCCCGTGTGCAGAAGGCTGACGAGCGTAAGAT CCAGCTCGTTTTGGAGGAACATCATCGCTTTTGCATGTCCTTCATTTAGCAGCTGTG TATGGAGCTTATTTATTCTTCACCTCGGCTATATGGCAGACAGACATATTTGCAT ACATCCTTTACGTTATGTCTGGATTAGGAATCACGGCGGGAGCGCACAGATTATG GGCTCATAAGTCATACAAGGCGAAGTGGCCGTTAAGATTAACTCCTCGTCGCATT AACACTTTGGCATTCCAGGATTCCGGCAATCGACTGGGCGCGCAGCACCGCATGC ACCACAAGTACTCGGAGACGGATGCGGACCCACATAACGCCACTCGCGGCTTCTT CTTTTCGCACATTGGTTGGTTACTCTGCCGAAAAACACCCGAGCTAAAGCGCAAG GGCCAGGGCCTCGACTTAAGTGACCTCTACGCAGATCCTATTATTCGCTTCCAAA AGAAGTACTACTTATTGTTAATGCCGTTAGCCTGCTTTGTTCTTCCCACCATAAT TCCGGTCTACCTCTGGGGCGAGTCTTGGA AAAACGCGTTCTTCGTAGCTGCAATG TTCCGTTACACGTT CATCCTCAACGTAACATGGCTCGTCAACTCCGCCGCCACA AATGGGGAGGCAAGCCCTATGATAAGAACATCCAGCCCGCTCAGAACATCTCTGT AGCTATCTTCGCATTAGGCGAGGGCTTCCACAAC TACCACCACACGTTCCCCTGG GACTACAAGACCGCTGAATTAGGAAACAACAGGTTAAATTTACAAC TTCGTTTA TCAATTTCTTCGCAAGCTTCGGATGGGCCTACGACTTAAAGACCGTGTCGGACGA GATTATACAACAGCGCGTTAAGAGGACGGGAGATGGGAGCCATCACTTACGGGGC TGGGGCGACCAGGACATACCGGCCGAAGAAGCTCAAGCTGCTTTACGCATTAACC GTAAAGATGATTAG
SEQ ID NO: 22	<i>H. zea</i> Z9 desaturase ATGGCTCCAAATATATCGGAGGATGTGAACGGGGTGCTCTTCGAGAGTGATGCAG CGACGCCGGACTTAGCGTTATCCACGCCGCCTGTGCAGAAGGCTGACAACAGGCC CAAGCAATTAGTGTGGAGGAACATACTATTATTTCGCGTATCTTCACTTAGCGGCT CTTTACGGAGGTTATTTATTCTCTCTCTCAGCTAAATGGCAGACAGACATATTTG CCTACATCTTATATGTGATCTCCGGGCTTGGTATCACGGCTGGAGCACATCGCTT ATGGGCCCCACAAGTCTTACAAAGCTAAATGGCCTCTCCGAGTTATCTTAGTCATC TTTAACACAGTGGCATTCAGGATGCCGCTATGGACTGGGCGCGCGACCACCGCA TGCATCACAAAGTACTCGGAAACCGATGCTGATCCTCATAATGCGACCCGAGGATT CTTCTTCTCTCACATTGGCTGGTTACTTGT CAGGAAACATCCCGACCTTAAGGAG AAGGGCAAGGGACTCGACATGAGCGACTTACTTGCTGACCCCATCTCAGGTTCC AGAAAAAATACTACTTAATCTTAATGCCCTTGGCTTGCTTCGTGATGCCTACCGT GATTCTGTGTACTTCTGGGGTGAAACCTGGACCAACGCATTCTTTGTGGCGGCC ATGTTCCGCTACGCGTT CATCCTAAATGTGACGTGGCTCGTCAACTCTGCCGCTC ACAAGTGGGGAGACAAGCCCTACGACAAAAGCATTAAGCCTTCCGAAAAC TGT GGTCGCCATGTTTCGCTCTCGGAGAAGGATTCCACAAC TACCACCACACTTCCCT TGGGACTACAAAAC TGCTGAGTTAGGCAACAACAAAAC TCAACTTCACTACCACCT TTATTAACTTCTTCGCTAAAATTGGCTGGGCTTACGACTTAAAGACAGTGTCTGA

- continued

SEQUENCE LISTING	
	TGATATCGTCAAGAACAGGGTGAAGCGCACTGGTGACGGCTCCCACCACTTATGG GGCTGGGGAGACGAAAATCAATCCAAAGAAGAAATTGATGCCGCTATCAGAATCA ATCCTAAGGACGATTAA
SEQ ID NO: 23	T. <i>pseudonana</i> Z11 desaturase ATGGACTTTCTCTCCGGCGATCCTTTCCGGACACTCGTCCTTGACGCACTTGTG TCATCGGATTTGCTGCGGCGTGGCAATGCTTCTACCCGCCGAGCATCGTCGGCAA GCCTCGTACATTAAGCAATGGTAAACTCAATACCAGAATCCATGGCAAATTGTAC GACCTCTCATCGTTTCAGCATCCAGGAGGCCCGTGGCTCTTTCTCTGTTC AAG GTGCGGACGGAACAGCTCTATTTGAGTCACACCATCCCTTCATACCTCGAAAAGAA TCTACTTCAGATCCTCTCCAAGTACGAGGTTCCGTCGACTGAAGACTCTGTTTCC TTCATCGCCACCCTAGACGAACTCAATGGTGAATCTCCGTACGATTGGAAGGACA TTGAAAATGATGATTTCGTATCTGACCTACGAGCTCTCGTAATTGAGCACTTTTC TCCTCTCGCCAAGGAAAGGGGAGTTTCACTCGTTGAGTCGTCGAAGGCAACACCT CAGCGGTGGATGGTGGTTCTATTACTCCTTGCGTCGTTCTTCCTCAGCATCCCAT TATATTTGAGTGGTTCGTGGACTTTGTTGTCGTCACTCCCATCCTCGCTTGGTT AGCGGTTGTCAATTACTGGCACGATGCTACTCACTTTGCATTGAGCAGCAACTGG ATTTTGAATGCTGCGCTCCCATATCTCCTCCTCTCCTATCGAGTCCGTCAATGT GGTATCATCATCACGTCATTGGACATCACGCATACACCAACATTTCCAAAAGAGA TCCAGATCTTGCTCACGCTCCACAACCTCATGAGAGAACACAAGAGTATCAAATGG AGACCATCTCACTTAAATCAAACACAGCTTCCGCGGATTCTCTTCATCTGGTCGA TTGCAGTCGGTATTGGGTTGAACTTATTAAACGACGTGAGAGCACTAACCAAGCT TTCATAACAACCGTTGTTGCGGTGGAGAAGATGTCATCGTCGCGAACATTACTC CATTTCTTGGACGTATGTTGCACATCTTTGTGACTACACTTTGGCCCTTTTGG CGTTTCCGGTGTGGAAGGCCATCGTTTGGGCGACTGTACCGAATGCCATATTAAG TTTGTGCTTCATGTTAAATACGCAAATCAATCACCTCATCAACACGTGTGCACAT GCTTCCGATAACAACCTTTTACAAGCATCAAGTTGTAAGTCTCAGAACTTTGGCC GATCAAGTGCCCTTTTGCTTCATCTTCTCGGGAGGTCTCAACTACCAAATTGAACA TCATTTGTTGCCGACGGTGAACCATTGCCATTTGCCAGCTTTGGCCCCGGGTGTA GAGCGTTTGTGTAAGAAACACGGGGTGACATACAACTCTGTTGAAGGATACAGAG AGGCCATCATTGCACACTTTGCACATACCAAAGATATGTCGACGAAGCCTACTGA TTGA
SEQ ID NO: 24	Native T. <i>ni</i> Z11 desaturase ATGGCTGTGATGGCTCAAACAGTACAAGAAACGGCTACAGTGTTGGAAGAGGAAG CTCGCACAGTGACTCTTGTGGCTCCAAAGACAACGCCAAGGAAATATAAAATATAT ATACACCAACTTTCTTACATTTTCATATGCGCATTTAGCTGCATTATACGGACTT TATTTGTGCTTCACCTCTGCGAAATGGGAAACATTGCTATTCTCTTTCGTACTCT TCCACATGTCAAATATAGGCATCACCGCAGGGGCTCACCGACTCTGGACTCACAA GACTTTCAAAGCCAAATTGCCTTTGGAAATTGTCCTCATGATATTCAACTCTTTA GCCTTTCAAACACGGCTATTACATGGGCTAGAGAACATCGGCTACATCACAAAT ACAGCGATACTGATGCTGATCCCCACAATGCGTCAAGAGGGTTCTTCTACTCGCA TGTTGGCTGGCTATTAGTAAAAAAACATCCCGATGTCCTGAAATATGGAAAAACT ATAGACATGTGGATGTATACAATAATCCTGTGTTAAAATTTAGAAAAAGTACG CAGTACCCTTAATTGGAACAGTTTGTTTTGTCTTGCCAACTTTGATTCCAGTCTA CTGTTGGGGCGAATCGTGGAACAACGCTTGGCACATAGCCTTATTTGATACATA TTCAATCTTAACGTGACTTTCCTAGTCAACAGTGCTGCGCATATCTGGGGGAATA AGCCTTATGATAAAAGCATCTTGCCCGCTCAAACCTGCTGGTTTCCTTCCTAGC AAGTGGAGAAGGCTTCCATAATTACCATCACGTCTTTCATGGGATTACCGCACA GCAGAATTAGGGAATAACTTCCTGAATTTGACGACGCTGTTTATTGATTTTGTG CCTGGTTTGGATGGCTTATGACTTGAAGTCTGTATCAGAGGATATTATAAAACA GAGAGCTAAACGAACAGGTGACGGTTCTTCAGGGGTCAATTTGGGGATGGGACGAC AAAGACATGGACCGGATATAAAATCTAAAGCTAACATTTTTTATGCTAAAAAGG AATGA
SEQ ID NO: 25	H. <i>zea</i> Z11 desaturase ATGGCCCAAAGCTATCAATCAACTACGGTTTTTGAGTGAGGAGAAAGAACTAACAC TGCAACATTTGGTGCCCCAAGCATCGCCAGGAAGTATCAAATAGTGTATCCGAA CCTCATTACGTTTGGTTACTGGCACATAGCCGACTTTATGGCCTTTACTTGTGC TTCACTTCTGCTAAATGGGCTACGATTTTATTAGCTACATCCTCTTCGTGTTAG CAGAAATAGGAATCACGGCTGGCGCTCACAGACTCTGGGCCACAAAACTTACAA AGCGAAACTACCATTAGAAATACTCTTAATGGTATTCAACTCCATCGCTTTTCAA AACTCAGCCATTGACTGGGTGAGGGACCACCGACTCCACCATAAGTATAGCGATA CAGATGCTGATCCCCACAATGCCAGCCGAGGGTTCTTTTATTCCCATGTAGGATG GCTACTTGTGAGAAAACATCCTGAAGTCAAAAAGCGAGGGAAAGAACTCAATATG TCCGATATTTACAACAATCCTGTCTGCGGTTTCAGAAAAAATACGCCATACCCT TCATTGGGGCTGTTTGTTCGCTTACCTACAATGATACCTGTTTACTTCTGGGG AGAAACCTGGTCCAATGCTTGGCATATCACCATGCTTCGCTACATCATGAACCTC AATGTACCTTTTTGGTAAACAGCGCTGCTCATATATGGGGAACAAGCCTTATG ACGCAAAATATTACCTGCACAAAATGTAGCTGTGTGGTCGCCACTGGTGGAGA AGGTTTCCATAATTACCACCATGTCTTCCCCTGGGATTATCGAGCAGCGGAACTC GGTAACAATAGCCTCAATCTGACGACTAAATTCATAGATTTATTTCGCAGCAATCG GATGGGCATATGATCTGAAGACGGTTTCGGAGGATATGATAAAACAAAGGATTAA ACGCACTGGAGATGGAACGGATCTTTGGGGACACGAACAAAACCTGTGATGAAGTG TGGGATGTAAAAGATAAATCAAGTTAA

- continued

SEQUENCE LISTING

SEQ ID NO: 26	<i>T. ni</i> Z11 desaturase <i>Homo sapiens</i> optimized ATGGCCGTGATGGCCCAGACCGTGCAGGAGACCGCAACAGTGCTGGAGGAGGAGG CAAGGACCGTGACACTGGTGGCACCCAAGACCACACCTAGAAAGTACAAGTATAT CTACACCAACTTCCTGACCTTCAGCTACGCACACCTGGCCGCCCTGTATGGACTG TACCTGTGCTTTACCTCCGCCAAGTGGGAGACACTGCTGTTCTCTTTTGTGCTGT TCCACATGAGCAATATCGGAATCACCGCAGGAGCACACAGGCTGTGGACCCACAA GACATTCAAGGCCAAGCTGCCTCTGGAGATCGTGCTGATGATCTTCAACTCTCTG GCCTTTTCAGAATACCGCCATCACATGGGCCCGGGAGCACAGACTGCACCACAAGT ATAGCGACACCGATGCAGACCCACACAACGCAAGCAGGGGCTTCTTTTACTCCCA CGTGGGCTGGCTGCTGGTGAAGAAGCACCCCGACGTGCTGAAGTATGGCAAGACA ATCGACATGTCCGACGTGTACAACAATCCCGTGCTGAAGTTTCAGAAGAAGTATG CCGTGCCTCTGATCGGCACCGTGTGCTTCGCCCTGCCAACACTGATCCCGTGTA TTGTTGGGGCGAGTCTTGGAACAATGCCTGGCACATCGCCCTGTTCCGGTACATC TTTAACCTGAATGTGACCTTTCTGGTGAACCTCCGCCGCCACATCTGGGGCAATA AGCCTTACGACAAGTCTATCCTGCCAGCCCAGAACCTGCTGGTGTCTTCTCTGGC CTCTGGCGAGGGCTTTCACAATTATCACCACGTGTTCCCATGGGACTACAGGACC GCAGAGCTGGGCAACAATTTTCTGAACCTGACCACACTGTTTCATCGATTTTGTG CCTGGTTCGGCTGGCCTATGACCTGAAGTCTGTGAGCGAGGATATCATCAAGCA GAGGGCAAAGAGGACAGGCGATGGCAGCTCCGGCGTGATCTGGGGATGGGACGAT AAGGATATGGACAGAGATATCAAGAGCAAGGCCAATATCTTCTACGCCAAGAAGG AGTGA
SEQ ID NO: 27	<i>H. zea</i> Z11 desaturase <i>Homo sapiens</i> optimized ATGGCACAGTCATATCAGAGCACTACCGTCCTGAGCGAAGAGAAGGAACTGACAC TGCAGCACCTGGTCCACAGGCATCACCTAGAAAGTACCAGATCGTGTATCCAAA CCTGATCACCTTCGGCTACTGGCACATCGCCGGCCTGTACGGCCTGTATCTGTGC TTTACCTCCGCCAAGTGGGCCACAATCCTGTTCTCTTACATCCTGTTTGTGCTGG CAGAGATCGGAATCACCGCAGGAGCACACAGACTGTGGGCACACAAGACATATAA GGCCAAGCTGCCCCTGAGATCCTGCTGATGGTGTTC AACAGCATCGCCTTTCAG AATTCCGCCATCGATTGGGTGCGGGACCACAGACTGCACCACAAGTACTCCGACA CCGATGCCGACCCCCACAACGCCTCTAGGGGCTTCTTTTATAGCCACGTGGGATG GCTGCTGGTGC GGAAGCACCCCTGAGGTGAAGAAGAGAGGCAAGGAGCTGAATATG TCTGATATCTACAACAATCCTGTGCTGCGCTTCAGAAGAAGTATGCCATCCCAT TCATCGGCGCGGTGTGCTTTGCCCTGCCACCATGATCCCGTGTACTTTTGGGG CGAGACATGGAGCAACGCCTGGCACATCACAAATGCTGCGGTATATCATGAACCTG AATGTGACATTCCTGGTGAACCTCCGCCGCCCACATCTGGGGCAATAAGCCATACG ACGCCAAGATCCTGCCCGCCCAGAACGTGGCCGTGAGCGTGGCAACCCGAGGAGA GGGCTTCCACAATTACCACCACGTGTTTCCTTGGGATTATCGGGCCGCCGAGCTG GGCAACAATTCTCTGAATCTGACCACAAAGTTCATCGACCTGTTTGCCGCCATCG GCTGGGCCTATGATCTGAAGACAGTGAGCGAGGACATGATCAAGCAGAGGATCAA GCGCACCGGCGATGGCACAGACCTGTGGGGGCACGAGCAGAACTGTGATGAAGTG TGGGATGTGAAAGACAAGTCCTCCTAA
SEQ ID NO: 28	<i>Y. lipolytica</i> OLE1 leader-<i>T. ni</i> Z11 desaturase <i>Homo sapiens</i> optimized ATGGTGAAGAACGTGGACCAGGTGGATCTGTCTCAGGTGGACACCATCGCAAGCG GAAGGGATGTGAATTATAAGGTGAAGTACACATCTGGCGTGAAGACCACACCAAG AAAGTACAAGTATATCTACACCAACTTCCTGACATTTTCTTACGCCACCTGGCC GCCCTGTATGGCCTGTACCTGTGCTTTACCAGCGCCAAGTGGGAGACACTGCTGT TCTCCTTTGTGCTGTTCCACATGTCTAATATCGGAATCACCGCAGGAGCACACAG GCTGTGGACCCACAAGACATTCAAGGCCAAGCTGCCCCTGAGATCGTGCTGATG ATCTTCAACTCCCTGGCCTTTTCAGAATACCGCCATCACATGGGCCCGGGAGCACA GACTGCACCACAAGTATTCTGACACCGATGCAGACCCACACAACGCAAGCAGGGG CTTCTTTTACTCCACGTGGGCTGGCTGCTGGTGAAGAAGCACCCTGACGTGCTG AAGTATGGCAAGACAATCGACATGAGCGACGTGTACAACAATCCTGTGCTGAAGT TTCAGAAGAAGTATGCCGTGCCACTGATCGGCACCGTGTGCTTCGCCCTGCCAC ACTGATCCCCGTGTACTGTTGGGGCGAGTCTTGAACAATGCCTGGCACATCGCC CTGTTCCGGTACATCTTTAACCTGAATGTGACCTTTCTGGTGAACAGCGCCGCC ACATCTGGGGCAATAAGCCATACGACAAGTCCATCCTGCCCGCCCAGAACCTGCT GGTGTCTTCTTGGCCTCTGGCGAGGGCTTTCACAATTATCACCACGTGTTCCCT TGGGACTACAGGACCGCAGAGCTGGGCAACAATTTTCTGAACCTGACCACACTGT TCATCGATTTTGTGCTGGTTCGGCTGGGCCTATGACCTGAAGTCTGTGAGCGA GGATATCATCAAGCAGAGGGCAAAGAGGACAGGCGATGGCAGCTCCGGCGTGATC TGGGGATGGGACGATAAGGATATGGACAGAGATATCAAGTCCAAGGCCAATATCT TCTACGCCAAGAAGGAGTGA
SEQ ID NO: 29	<i>Y. lipolytica</i> OLE1 leader-<i>H. zea</i> Z11 desaturase <i>Homo sapiens</i> optimized ATGGTGAAAAACGTGGACCAAGTGGATCTCTCGCAGGTCGACACCATTGCCTCCG GCCGAGATGTCAACTACAAGGTCAAGTACACCTCCGGCGTTCGCAAGTATCAGAT CGTGTATCCTAACCTGATCACCTTCGGCTACTGGCATATCGCTGGACTGTACGGA CTGTATCTGTGCTTCACTTCCGCCAAGTGGGCCACCATCCTGTTCTCTTACATCC TGTTTGTGCTGGCAGAGATCGGAATCACCGCAGGAGCACACAGACTGTGGGCACA CAAGACATATAAGGCCAAGCTGCCACTGGAGATCCTGCTGATGGTGTTC AACAGC ATCGCCTTTTCAGAATTCGCCCATCGATTGGGTGCGGGACCACAGACTGCACCACA AGTACTCCGACACAGATGCCGACCCCCACAACGCCTCTAGGGGCTTCTTTTATAG

- continued

SEQUENCE LISTING	
	CCACGTGGGATGGCTGCTGGTGCGGAAGCACCCCTGAGGTGAAGAAGAGAGGCAAG GAGCTGAATATGTCTGATATCTACAACAATCCTGTGCTGCGCTTCCAGAAGAAGT ATGCCATCCCATTTCATCGGCGCCGTGTGCTTTGCCCTGCCCACCATGATCCCCGT GTACTTTTGGGGCGAGACATGGAGCAACGCCTGGCACATCACAATGCTGCGGTAT ATCATGAACCTGAATGTGACATTCTGCTGAACTCCGCGCGCCACATCTGGGGCA ATAAGCCATACGACGCCAAGATCCTGCCCCGCCAGAACGTGGCCGTGAGCGTGGC AACC GGAGGAGAGGGCTTCCACAATTACCACCACGTGTTTCCATGGGATTATAGG GCAGCAGAGCTGGGAAACAATTCTCTGAATCTGACCACAAAGTTCATCGACCTGT TTGCCCGCCATCGGCTGGGCCTATGATCTGAAGACAGTGAGCGAGGACATGATCAA GCAGAGGATCAAGCGCACCGGCGATGGCACAGACCTGTGGGGGCACGAGCAGAAT TGTGATGAAGTGTGGGATGTGAAGGATAAAAGCAGTTGA
SEQ ID NO: 30	Native <i>A. transitella</i> Z11 desaturase ATGGTCCCTAACAGGGTTCCAGTGACGTTTTGTCTGAACATTCTGAGCCCCAGT TCACTAAACTCATAGCTCCACAAGCAGGGCCGAGGAAATACAAGATAGTGATCG AAATTTGCTCACATTGGCTATTGGCACTTATCAGCTGTTTATGGGCTCTACTTG TGCTTTACTTGTGCGAAATGGGCTACCATCTTATTTGCATTTTTCTTATACGTGA TCGCGGAAATCGGTATAACAGGTGGCGCTCATAGGCTATGGGCACATCGGACTTA TAAAGCCAAGTTGCCTTTAGAGATTTTGTACTCATAATGAATTCTATTGCCTTC CAAGACACTGCTTTCACCTGGGCTCGAGATCACCGCCTTCATCACAAATATTCGG ATACTGACGCTGATCCCCACAATGCTACCAGAGGGTTTTTCTATTACATGTAGG CTGGCTTTTGGTGAGAAACACCCTGAAGTCAAAGCAAGAGGAAAATACTTGTCTG TTAGATGATCTTAAAGAATAATCCATTGCTTAAATTCCAAAAGAAATACGCTATT TAGTTATAGGCACGTTATGCTTCCTTATGCCAACATTTGTGCCCGTATACTTCTG GGGCGAGGGCATCAGCACGGCCTGGAACATCAATCTATTGCGATACGTCATGAAT CTTAACATGACTTTCTTAGTTAACAGTGCAGCGCATATCTTTGGCAACAAACCAT ACGATAAGAGCATAGCCTCAGTCCAAAATATTTAGTTAGCTTAGCTACTTTTGG CGAAGGATTCCATAATTACCATCACACTTACCCCTGGGATTATCGTGCGGCAGAA TTAGGAAATAATAGGCTAAATATGACTACTGCTTTCATAGATTTCTTCGCTTGGA TCGGCTGGGCTTATGACTTGAAGTCTGTGCCACAAGAGGCCATTGCAAAAAGGTG TGCGAAAAC TGCGATGGAACGGATATGTGGGGTCGAAAAAGATAA
SEQ ID NO: 31	pPV0228_- Z11 Helicoverpa zea desaturase ATGGCCCAAAGCTATCAATCAACTACGGTTTTGAGTGAGGAGAAAGAACTAACAT TACAACATTTGGTGCCCCAAGCATCGCCAGGAAGTATCAAATAGTGATCCGAA CCTCATTACGTTTGTTACTGGCACATAGCCGGACTTTATGGCCTTTACTTGTGC TTCACTTCTGCTAAATGGGCTACGATTTTATTAGCTACATCCTCTTCGTGTTAG CAGAAATAGGAATCACGGCTGGCGCTCACAGACTCTGGGCCACAAAACCTACAA AGCGAAACTACCATTAGAAATACTCTTAATGGTATTCAACTCCATCGCTTTCAA AACTCAGCCATTGACTGGGTGAGGGACCACCGACTCCACCATAAGTATAGCGATA CAGATGCTGATCCCCACAATGCCAGCCGAGGGTTCTTTTATTCCCATGTAGGATG GCTACTTGTGAGAAAAATCCTGAAGTCAAAAAGCGAGGGAAAGAACTCAATATG TCCGATATTTACAACAATCCTGTCTTACGGTTTCAGAAAAAATACGCCATACCCT TCATTGGGGCTGTTTGTTCGCTTACCTACAATGATACTGTTTACTTCTGGGG AGAAACCTGGTCCAATGCTTGGCATATCACCATGCTTCGCTACATCATGAACCTC AATGTCACCTTTTTGGTAAACAGCGCTGCTCATATATGGGGAAACAAGCCTTATG ACGCAAAAATATTACCTGCACAAAATGTAGCTGTGTCGGTCGCCACTGGTGAGA AGGTTTCCATAATTACCACCATGTCTTCCCCTGGGATTATCGAGCAGCGAACTC GGTAACAATAGCCTCAATTTAACGACTAAATTCATAGATTTATTTCGCAGCAATCG GATGGGCATATGATTTAAAGACGGTTTCGGAGGATATGATAAAACAAGGATTAA ACGCACTGGAGATGGAACGGATCTTTGGGGACACGAACAAAAC TGATGAAGTG TGGGATGTAAAAGATAAATCAAGTTAA
SEQ ID NO: 32	pPV0228_- Helicoverpa armigera reductase codon optimized ATGGTCGTTTTTAACCTCTAAAGAGACAAAACCTTCAGTAGCTGAGTTTTATGCGG GAAAATCTGTTTTTATTACGGGTGGCACTGGATTCCCTTGAAAGGTATTCATAGA GAAACTTTTATATAGCTGTCCAGATATCGAGAATATCTACATGCTCATACGAGAG AAGAAAGGTC'TTCTGTAGCGAAAGAATAAAACAGTTCTTGATGACCCGCTCT TTACCAGACTAAAAGACAAAAGACCAGCTGACTTAGAGAAGATTGTATTAATACC AGGAGATATTACTGCTCCTGACTTAGGCATTAATTCTGAAAACGAGAAGATGCTT ATAGAGAAGGTATCGGTGATTATTTCATTTCGGCTGCTACGGTGAAGTTTAATGAGC CTCTCCCTACGGCTTGGAAGATCAACGTGGAAGGAACCAGAATGATGTTAGCTTT GAGTCGAAGAATGAAGCGGATTGAGGTTTTTCATTACATATCGACAGCATACACG AACACAAACAGGGAAGTGTTGACGAGATCTTATACCCAGCTCCTGCTGATATCG ACCAAGTTCATCAGTATGTCAAAGATGGAATCTCTGAGGAAGACACTGAGAAAAT ATTAAATGGTCGTCCAAATACGTACACGTTTACGAAAGCGTAACTGAGCATTTA GTTGCTGAGAACCAAGCCTACGTACCCACTATTATCGTCAGGCCGTGCTGCTGG CAGCAATAAAAGATGAGCCATTAAAAGGTTGGTTAGGCAACTGGTTTGGAGCGAC TGGTCTCACCCTGTTTACCGCTAAGGGTCTCAACCGAGTCATCTACGGTCATTCT AGCTACATCGTAGACTTAATTCCTGTGGATTATGTGCGCTAATTTAGTGATTGCTG CTGGGGCTAAGAGTAGCAAGTCAACTGAGTTGAAGGTATACAAC TGCTGCAGCAG CTCCTGCAATCCCGTCACTATTGGCACGTTAATGAGCATGTTTGTGACGATGCC ATCAAACAGAAGTCGTATGCTATGCCGCTACCGGGGTGGTACATATTCACGAAAT ATAAGTGGTTAGTTCTTCTTTAACATTTCTCTTCCAAGTTATACCGGCGTATGT CACAGATCTCTCCAGGCACTTGATTGGGAAGAGTCCACGGTACATAAAACTCCAA TCACTAGTAAATCAAACGCGCTCTTCAATCGACTTCTTACGAATCACTCCTGGG

- continued

SEQUENCE LISTING	
	TGATGAAGGCAGACAGAGTGAGAGAGTTATATGCGTCTCTTTCCCCCGCAGACAA GTACTTATTTCCCTGTGATCCTACGGACATTAAGTGGACACATTACATACAAGAC TACTGTTGGGGAGTCCGACATTTTTTTGGAGAAAAAAGCTACGAATAA
SEQ ID NO: 33	pPV0228_- _ICL_promoter TATTAGGCGAAGAGGCATCTAGTAGTAGTGGCAGTGGTGAGAACGTGGGCGCTGC TATAGTGAACAATCTCCAGTCGATGGTTAAGAAGAAGAGTGACAAACCAGCAGTG AATGACTTGTCTGGGTCCGTGAGGAAAAGAAAGAGCCCGACACAAAGGACAGTA ACGTCAAGAAACCCCAAGAAATAGGGGGGACCTGTTTAGATGTATAGGAATAAAAA CTCCGAGATGATCTCAATGTGTAATGGAGTTGTAATATTGCAAAGGGGGAAAAATC AAGACTCAAACGTGTGTATGAGTGAGCGTACGTATATCTCCGAGAGTAGTATGAC ATAATGATGACTGTGAATCATCGTAATCTCACACAAAAACCCCATTTGTCGGCCAT ATACCACACCAAGCAACACCACATATCCCCCGAAAAAAAACGTGAAAAAAGA AACAATCAAAACTACAACCTACTCCTTGATCACACAGTCATTGATCAAGTTACAG TTCCTGCTAGGGAATGACCAAGGTACAAATCAGCACCTTAATGGTTAGCACGCTC TCTTACTCTCTCTCACAGTCTTCCGGCCCCCTATTCAAATTTCTGCACTTCCATTT GACCCAGGGTTGGGAAACAGGGCCACAAAAGAAAAACCCGACGTGAATGAAAAA ACTAAGAAAAGAAAAAAATTATCACACCAGAAATTTACCTAATTGGGTAATTCC CATCGGTGTTTTTCTCGATTGTGCGCACGCACGCATGCTGAAAAAGTGTTGAG TTTTGCTTTTGCCTCGGAGTTTACGCAAGTTTTTCGATCTCGGAACCGAGGGC GGTCGCCTTGTTGTTGTGATGTCGTGCTTTGGGTGTTCTAATGTGCTGTTATTG TGCTCTTTTTTTTTCTTCTTTTTTTGGTGATCATATGATATTGCTCGGTAGATTA CTTTCGTGTGTAGGTATTCTTTTAGACGTTTGGTATTGGGTAGATATGAGAGAG AGAGAGTGGGTGGGGGAGGAGTTGGTTGTAGGAGGGACCCCTGGGAGGAAGTGTA GTTGAGTTTTCCCTGACGAATGAAAATACGTTTTTGAGAAGATAATACAGGAAAG GTGTGTCGGTGAATTTCCATCTATCCGAGGATATGAGTGGAGGAGAGTCGTGTGC GTGTGGTTAATTTAGGATCAGTGGAACACACAAAGTAAGTAAAGACAGAGAGACAG AGAGAAAAATCTGGGAAGAGACAAAGAGTCAGAGTGTGTGAGTTATTCTGTATT GTGAAATTTTTTTGCCCAACTACATAATATTGCTGAAACTAATTTTACTTAAAAA GAAAAGCCAACAACGTCCCCAGTAAAACTTTTCTATAAATATCAGCAGTTTCCC TTTCCTCCATTCCCTCTTCTTGCTTTTTTCTTACTTTCCTTTTTTATACCTTTT CATTATCATCCTTTATAATTGTCTAACCAACAACCTATATATCTATCAA
SEQ ID NO: 34	pPV0228_- _TEF_Candida tropicalis promoter region AGGAAGACAACCAAAAGAAAGATCAAATTGACTAAATGTTGAACAGACCAAAAAA AAAGAACAACAAATAGATAAAATTACAACATATTAATCTTTTGATATGTTGTTGAA TATTCTAGTAAATCTAATGATCTCAATAGTGGTTATCATTCACTCTCTTCGTCCT CCTCTCTCCCCCTCCTCCTTGCAGTATATTAAGCAATAAAAAAAAAAAAAAAAAA AAGAAAATCTGCCAACACACACAAAAAAACTTACATAGTCGTGTACCAGTGTCA ATATTTCAACAGCGCAGAGAAAAGAAGATGAACAGAAAAATTTCTCTTTGGTTT TGTCTTTGGTTTTGTATTAATCTCATTGAAAAATTTTTCTCTCTCTCTCTCTCT CTCTCTCACTCACACACTCACTCGCATTTTCGTTTGGGTACAGCAGAAGTCAGAC AGAAAAAAAAAATCGTATATAACTCTCATCAAATGCCCTAGAGAAAAATTTTCT TCTATCCTTTTTTTTTCTTCTTCTTCTTCTTCTTCTTCTTCTTCTTTTAGAAGATC TTTTTGAATTCATCAAAGATATATATTTAATCAATC
SEQ ID NO: 35	pPV0228_- _ICL_terminator AAGAAAAAGAAAAGGTAAAGAAGTTCATTTGAGATGAACTTTTGTATATGACTT TTAGTTTCTACTTTTTTTTTTATTTATTGCTTAATTTTCTTTATTTCAATCCCCC ATAGTTTGTGTAGAATATATTTATTCAATCTGGTAACTCAAACACGTAGCAAGCT CGTTGCATCTCGCCTCGTCACGGGTACAGCTCTGGAACCAAAGACAAAAA GTTGATCCGAACCTCTCGCTATTCCCTTGCTATGCTATCCACGAGATGGGGTTTA TCAGCCCAGGCAAGTCACTAAA
SEQ ID NO: 36	pPV0228_- _TEF_terminator GCTGATTAATGAATAATTAATAAGTATTGTTTTTTTTGTTTTTAATATATATATATCT TGAAATTAGTATAAAAAAATCTTTTTTTTTCTTTTTTATTTATTTATCAATAGTT TATATATATATATATAAACTTGTAAGAGATTAGGTATATCTAACAGTGATACTACT AATAGTGCTTAATATCTTTGTAAACAAGAAAATAAAATAAAC
SEQ ID NO: 37	SapI-tLIP2-pEXP1-HA FAR-SapI (insert into pPV199 creating pPV247) GCCTGAAGAGCGCTATTTATCACTCTTTACAACCTTCTACCTCAACTATCTACT TTAATAAATGAATATCGTTTATTCTCTATGATTACTGTATATGCGTTCCTCCA TGGGAGTTTGGCGCCCGTTTTTTTCGAGCCCCACACGTTTCGGTGAGTATGAGC GGCGGCAGATTCGAGCGTTTCCGGTTTCCGCGGCTGGACGAGAGCCCATGATG GGGGCTCCCACCACCAGCAATCAGGGCCCTGATTACACACCCACCTGTAATGT CATGCTGTTTCATCGTGTTAATGCTGCTGTGTGCTGTGTGTGTGTGTTGTTG GCGCTCATTGTTGCGTTATGCAGCGTACACCACAATATTGGAAGCTTATTAGC CTTTCTATTTTTTCGTTTGCAAGGCTTAACAACATTGCTGTGGAGAGGGATGG GGATATGGAGGCCGCTGGAGGGAGTCGGAGAGGCGTTTTTGGAGCGGCTTGGCC TGGCGCCAGCTCGCGAAACGCACCTAGGACCCCTTGGCACGCCGAAATGTGC CACTTTTCAGTCTAGTAACGCCTTACCTACGTCAATCCATGCATGCATGTTTG CGCCTTTTTTCCCTTGCCCTTGATCGCCACACAGTACAGTGCACGTGTACAGTG GAGGTTTTTGGGGGGTCTTAGATGGGAGCTAAAAGCGGCCCTAGCGGTACACTA GTGGGATTGTATGGAGTGGCATGGAGCCTAGGTGGAGCCTGACAGGACGCACG ACCGGCTAGCCCGTGACAGACGATGGGTGGCTCCTGTTGTCCACCGGTACAA

- continued

SEQUENCE LISTING	
	ATGTTTGGGCCAAAGTCTTGTGAGCCTTGCTTGCGAACCTAATTCCCAATTTT GTCACTTCGCACCCCCATTGATCGAGCCCTAACCCCTGCCCATCAGGCAATCC AATTAAGCTCGCATTGTCTGCCTTGTTTAGTTTGGCTCCTGCCCGTTTCGGCG TCCACTTGCACAAACACAAACAAGCATTATATATAAGGCTCGTCTCTCCCTCC CAACCACACTCACTTTTTTGCCCGTCTTCCCTTGCTAACACAAAAGTCAAGAA CACAAACAACCACCCCAACCCCTTACACACAAGACATATCTACAGCAATGGT GGTGCTGACCAGCAAGGAGACAAAGCCTTCCGTGGCCGAGTTCTACGCCGGCA AGTCCGTGTTTATCACAGGCGGCACCGGCTTCTGGGCAAGGTGTTTATCGAG AAGCTGCTGTACTCTTGCCCGAGACATCGAGAACATCTATATGCTGATCCGGGA GAAGAAGGGCCTGAGCGTGTCGAGAGAATCAAGCAGTTTCTGGACGATCCCC TGTTTACACGGCTGAAGGACAAGAGACCTGCCGATCTGGAGAAGATCGTGCTG ATCCCAGGCGACATCACCGCACCAGATCTGGGCATCAACTCCGAGAATGAGAA GATGCTGATCGAGAAGGTGTCCGTGATCATCCACTCTGCCGCCACCGTGAAGT TCAACGAGCCCCCTGCCTACAGCCTGGAAGATCAATGTGGAGGGCACCAGGATG ATGCTGGCCCTGAGCCGGAGAATGAAGCGCATCGAGGTGTTTATCCACATCTC CACAGCCTACACCAACACAAATCGGGAGGTGGTGGACGAGATCCTGTACCCAG CCCCCGCCGACATCGATCAGGTGCACCAGTATGTGAAGGACGGCATCAGCGAG GAGGATACCGAGAAGATCCTGAACGGCCGGCCAAATACCTACACATTACCAA GGCCCTGACAGAGCACCTGGTGGCCGAGAACCAGGCCTATGTGCCTACCATCA TCGTGAGACCATCCGTGGTGGCCGCCATCAAGGATGAGCCCCCTGAAGGGATGG CTGGGAAACTGGTTCGAGCAACAGGACTGACCGTGTTTACAGCCAAGGGCCT GAATAGAGTGATCTACGGCCACAGCTCCTATATCGTGGACCTGATCCCCGTGG ATTACGTGGCAAACCTGGTCAATCGCAGCAGGAGCCAAGTCTAGCAAGTCTACC GAGCTGAAGGTGTATAACTGCTGTTCCCTCTAGCTGTAATCCTGTGACCATCGG CACACTGATGTCCATGTTTCGCCGACGATGCCATCAAGCAGAAGTCTTACGCCA TGCCTCTGCCAGGCTGGTACATCTTTACAAAGTATAAGTGGCTGGTGTGCTG CTGACCTTCCTGTTTCAGGTCAATCCAGCCTACGTGACCGATCTGTCTAGGCA CCTGATCGGCAAGAGCCCCCGCTATATCAAGCTGCAGTCTCTGGTGAACCAGA CCAGGTCTCTATCGACTTCTTTACAAATCACAGCTGGGTGATGAAGGCCGAT AGGGTGCGCGAGCTGTACGCCTCTCTGAGCCCTGCCGACAAGTATCTGTTCCT CTGCGACCCCTACCGATATCAATTGGACACACTACATCCAGGATTATTGTTGGG GCGTGCGCCACTTCCTGGAGAAGAAGTCCTATGAGTGAGCCTGAAGAGC
SEQ ID NO: 38	NcoI-pTAL-AleI (insert into pPV247 creating pPV248) CCATGGGTAAGCAGGTGGCTCCGTTTGTGTCTTTGTGTTTTTCCCTCCTTTT TGGACCATTTGTGTCAGCATGTTGCGTAGGTCTGGGTGTTTGACTGTTTCAGGTGG TGGATGACGGATGCATCATCTGACGGCAGAGTGGGTACCTGGCAGTGGCAGGC TCGCAGACGAGGTAGAGAGATTCTGAAAGGAGCCATTGACAGATGGAGAATTG GATACTCCTGGTATGTCTCCGTTTCCACTTTTGACGTTGGTGACGTGCTCTG GAACGACTTTTTTCTTTTTCTTTAAAACAAAAAAGAAAGAAAAAACA TTTACTACTACCAGTAGTACACCTCAACATTGGGTCCAGAACGTCCCAACTGC ATGAGTCACTGGAGTCATGCCGAGGTCGCTAAGGTGCTGTAAAATACAACGTC AATTGAGAGAGACACAGGCGCAGCGCGCCGAGGGAGAAACGAGGCATTATCT TCTGACCCTCCTTTTTACTCGTAATCTGTATCCCGGAACCGCGTCGCATCCAT GTTAATTAAATCAACACTTACACTTGCTTGCTTCGTATGATGAAGATTCTGA CTGGCAACCCAGTCAGCAGCAGATTGGGGCAGATGTAGTAATGAAAACACTG CAAGGTGTGACGTTTGAGACACTCCAATTGGTTAGAAAGCGACAAAGAAGACG TCGGAAAAATACCGGAAAAATCGAGTCTTTTTCTTTCTGCGTATTGGGCCCTT CTGCCTCCTTTGCCGCCCTTTCCACGCTCTTCCACACCCTCACACTCCCTGA GCACTATGATCTCATTGCGCAATAAGATATACATGCACGTGCATTTGGTGAGC ACGCAGAACCTTGTTGGGGGAAGATGCCCTAACCTAAGGGCGTTCATACGG TTCGACAGAGTAACCTTGCTGTGATTATAACGCATATATAGCCCCCCCCTTC GGACCCTCCTTCTGATTTCTGTTTCTGTATCAACATTACACACAAACACACAA TGGTG

The foregoing detailed description has been given for
clearness of understanding only and no unnecessary limita-
tions should be understood there from as modifications will
be obvious to those skilled in the art.
While the disclosure has been described in connection
with specific embodiments thereof, it will be understood that
it is capable of further modifications and this application is
intended to cover any variations, uses, or adaptations of the
disclosure following, in general, the principles of the dis-

closure and including such departures from the present
disclosure as come within known or customary practice
within the art to which the disclosure pertains and as may be
applied to the essential features hereinbefore set forth and as
follows in the scope of the appended claims.
The disclosures, including the claims, figures and/or
drawings, of each and every patent, patent application, and
publication cited herein are hereby incorporated herein by
reference in their entirety.

SEQUENCE LISTING	
<160>	NUMBER OF SEQ ID NOS: 38
<210>	SEQ ID NO 1
<211>	LENGTH: 1377

-continued

<212> TYPE: DNA

<213> ORGANISM: *Agrotis segetum*

<400> SEQUENCE: 1

```
atgccagttt tgacttctag agaagatgaa aagttgtcag ttccagaatt ttacgctggt    60
aaatctatct tcgttacagg tgggtactggt ttcttgggta aagtttttat tgaaaagttg   120
ttgtactggt gtccagatat tgataaaatc tatatgttaa ttagagaaaa gaaaaatttg   180
tctattgatg aaagaatgtc aaagttcttg gatgatccat tattttctag attgaaggaa   240
gaaagacctg gtgacttggg aaagattggt ttgattccag gtgacattac agctccaaat   300
ttgggtttat cagcagaaaa cgaaagaatt ttgttagaaa aagtttctgt tattattaat   360
tcagctgcaa ctgttaagtt taatgaacca ttgccaatcg cttggaagat taatgttgaa   420
ggtacaagaa tgttgttggc attgtctaga agaatgaaga gaatcgaagt ttttattcat   480
atttctactg cttactcaaa tgcattctca gatagaatcg ttgttgatga aatcttgtat   540
ccagctccag cagatatgga tcaagtttac caattgggta aagatgggtg tacagaagaa   600
gaaactgaaa gattgttgaa cggtttgcca aacacttaca cttttactaa ggctttgaca   660
gaacatttgg ttgcagaaca tcaaacatac gttccaacta tcatcatcag accatctggt   720
gttgcttcaa ttaaagatga accaatcaga ggttggttat gtaattgggt tggtgctaca   780
ggtatctctg tttttactgc aaagggtttg aacagagttt tgttgggtaa agcttcaaac   840
atcgttgatg ttatcccagt tgattacggt gcaaatttgg ttattgttgc tggtgcaaaa   900
tctggtgggc aaaaatcaga tgaattaaag atctataact gttgttcttc agattgtaac   960
ccagttactt tgaagaaaat tattaagag tttactgaag atactattaa aaataagtct  1020
catattatgc cattgccagg ttggttcggt tttactaagt acaagtgggt gttgacattg  1080
ttaactatta tttttcaaat gttaccaatg tatttggtcg atgtttacag agttttgaca  1140
ggtaaaatcc caagatacat gaagttgcat catttggtta ttcaaacaag attgggtatc  1200
gatttcttta cttctcattc atgggttatg aagacagata gagttagaga attattcggt  1260
tctttgtcat tggcagaaaa gcatatgttt ccatgtgac catcttcaat cgattggaca  1320
gattatttgc aatcactactg ttacggtggt agaagatttt tggaaaagaa gaaataa   1377
```

<210> SEQ ID NO 2

<211> LENGTH: 1365

<212> TYPE: DNA

<213> ORGANISM: *Spodoptera littoralis*

<400> SEQUENCE: 2

```
atggttgttt tgacttcaaa ggaaaaatca aacatgtctg ttgctgattt ctacgctggt    60
aaatctgttt ttattacagg tgggtactggt ttcttgggta aagtttttat tgaaaagttg   120
ttgtactcat gtccagatat tgataaaatc tatatgttga tcagagaaaa gaaagggtcaa   180
tctatcagag aaagattaac taaaattggt gatgatccat tgtttaatag attgaaggat   240
aagagaccag atgatttggg taaaatcggt ttgatccag gtgacatcac agttccaggt   300
ttgggtattt ctgaagaaaa cgaaacaatc ttgactgaaa aagtttcagt tgttattcat   360
tctgctgcaa ctgttaagtt taatgaacca ttggctactg catggaacgt taacgttgaa   420
ggtacaagaa tgatcatggc attatcaaga agaatgaaga gaatcgaagt ttttattcat   480
atttctactg cttactacta cacaacaga gcagttattg atgaagtttt gtatccacca   540
ccagctgata tcaacgatgt tcatcaacat gttaaaaatg gtgttacaga agaagaaact   600
```

-continued

gaaaagattt	tgaacggtag	accaaact	tacactttta	ctaaggcttt	gactgaacat	660
ttggttgcag	aaaaccaatc	atacatgcc	acaatcattg	ttagaccatc	tattgttggt	720
gctattaaag	atgatccaat	tagaggttgg	ttggctaatt	ggtatggtgc	aacaggtttg	780
tcagttttta	ctgcaaagg	tttgaacaga	gttatatatg	gtcattctaa	ccatgttggt	840
gatttgattc	cagttgatta	cgttgcta	ttggttattg	ttgctggtgc	aaagacatac	900
cattcaaacg	aagtactat	ctataactct	tgttcttcat	cttgtaaccc	aatcactatg	960
aagagattgg	ttggtttgtt	tattgattac	acagttaagc	ataagtcata	cgttatgcc	1020
ttgccagggt	ggtatgttta	ctctaactac	aagtgggttg	tttcttggt	tactgttatt	1080
ttccaagtta	ttccagctta	cttaggtgac	attggtagaa	gattgttagg	taaaaatcca	1140
agatactaca	agttgcaaaa	tttggttgct	caaacacaag	aagcagttca	tttctttaca	1200
tcacatactt	gggaaattaa	atcaaagaga	acttctgaat	tgttttcac	tttgtcttg	1260
acagatcaaa	gaatgtttcc	atgtgatgct	aacagaatcg	attggacaga	ttacatcact	1320
gattactggt	ctggtgtag	acaatttttg	gaaaagatta	aataa		1365

<210> SEQ ID NO 3

<211> LENGTH: 1368

<212> TYPE: DNA

<213> ORGANISM: Helicoverpa armigera

<400> SEQUENCE: 3

atggttggtt	tgacttcaaa	ggaaacaaag	ccatctgttg	ctgaatttta	cgttggtaaa	60
tcagttttta	ttacaggtgg	tactggtttc	ttgggtaaag	tttttattga	aaagttgttg	120
tactcttgtc	cagatattga	aaatatctat	atgttgatca	gagaaaagaa	aggtttgta	180
gtttctgaaa	gaattaaaca	attttttagat	gatccattgt	ttacaagatt	gaaggataag	240
agaccagctg	atgttgaaaa	gattgttttg	atcccagggtg	acatcactgc	accagatttg	300
ggtattaatt	ctgaaaacga	aaagatgttg	attgaaaaag	tttcagttat	tattcattct	360
gctgcaactg	ttaagtttaa	tgaaccatta	ccaacagctt	ggaagattaa	tggtgaagg	420
actagaatga	tggtggcatt	gtcaagaaga	atgaagagaa	tgaagtttt	tattcatatt	480
tctacagctt	acactaacac	aaacagagaa	gttggtgatg	aaatcttgta	tccagctcca	540
gcagatatcg	atcaagttca	tcaatacgtt	aaggatggta	tctcagaaga	agatactgaa	600
aagattttga	acggttagacc	aaacacttac	acttttacta	aggctttgac	agaacatttg	660
gttgctgaaa	atcaagcata	cgttccaact	attattgtta	gaccatctgt	tggtgctgca	720
attaaagatg	aaccattgaa	aggttgggtg	ggtaattgg	ttggtgctac	aggtttgact	780
gtttttacag	caaagggttt	gaacagagtt	atatatggtc	attcttcata	catcgttgat	840
ttgatcccag	ttgattacgt	tgctaatttg	gttattgctg	cagggtgcaa	atcttcaaag	900
tcaacagaat	tgaaggttta	caactgttgt	tcttcatctt	gtaaccag	tactatcggt	960
acattgatgt	caatgttcgc	tgatgatgca	attaaacaaa	aatcttacgc	tatgccattg	1020
ccaggttggt	acatttttac	aaagtacaag	tggttggttt	tggtgttgac	atttttgttc	1080
caagttattc	cagcatatcg	tactgatttg	tcaagacatt	tgatcggtaa	atctccaaga	1140
tacatcaagt	tgcaatcatt	ggtaaccaa	actagatcat	ctatcgattt	ctttacaaac	1200
cattcttggg	ttatgaaagc	tgatagagtt	agagaattgt	acgcttcatt	gtctccagct	1260
gataagtact	tattcccatg	tgatccaact	gatatcaact	ggacacatta	catccaagat	1320
tactgttggg	gtgtagaca	tttcttgga	aagaaatctt	acgaataa		1368

-continued

<210> SEQ ID NO 4
 <211> LENGTH: 1199
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: pOLE1 cassette

<400> SEQUENCE: 4

```

cttgctgaaa agatgatgtt ctgaggtatt cgtatcgcta gcttgatacg cttttaacaa    60
aagtaagctt ttcgtttgca ggtttgggta cttttctgta cgagatgata tcgctaagtt    120
tatagtcata tgtgaaatth ctcaaaaacc tcatgggtttc tccatcacc ctttttcatt    180
tcatttgccg ggcggaaaaa aaaaaggaaa aaaaaaaaaa aaaaaataa atgacacatg    240
gaaataagtc aaggattagc ggatatgtag ttccagtcog gggtatacca tcacgtgata    300
ataaatccaa atgagaatga ggggtgtcata tctaatacatt atgcacgtca agattctccg    360
tgactatggc tcttttctga agcatttttc gggcgcccg tgccaaaaa ctaactccga    420
gcccgggcat gtccgggggt tagcggggcc acaaaaggcg cttatctggg gggttccgt    480
agaagaaaaa aagctgttga gcgagctatt tcgggtatcc cagccttctc tgcagaccgc    540
cccagttggc ttggctctgg tgctgttcgt tagcatcaca tcgctgtga caggcagagg    600
taataacggc ttaaggttct cttcgcatag tcggcagctt tctttcggac gttgaacact    660
caacaaacct tatctagtgc ccaaccaggt gtgcttctac gagtcttgct cactcagaca    720
cacctatccc tattgttacg gctatgggga tggcacacaa aggtggaaat aatagtagtt    780
aacaatatat gcagcaaatc atcggtcctt ggctcatoga gtcttgcaaa tcagcatata    840
catatatata tgggggcaga tcttgattca tttattgttc tatttccatc tttctactt    900
ctgtttccgt ttatatthtg tattacgtag aatagaacat catagtaata gatagttgtg    960
gtgatcatat tataaacagc actaaaacat tacaacaaag aatgccaaact tctggaacta   1020
ctattgaatt gattgacgac caatttccaa aggatgactc tgccagcagt ggcattgtcg   1080
acactagtgc ggcgctcac atatgaaagt atataccgcg tttgtacac tatgtagcta   1140
taattcaatc gtattattgt agctccgcac gaccatgcct tagaaatatc cgcagcgcg   1199

```

<210> SEQ ID NO 5
 <211> LENGTH: 655
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: OLE1 promoter region

<400> SEQUENCE: 5

```

cttgctgaaa agatgatgtt ctgaggtatt cgtatcgcta gcttgatacg cttttaacaa    60
aagtaagctt ttcgtttgca ggtttgggta cttttctgta cgagatgata tcgctaagtt    120
tatagtcata tgtgaaatth ctcaaaaacc tcatgggtttc tccatcacc ctttttcatt    180
tcatttgccg ggcggaaaaa aaaaaggaaa aaaaaaaaaa aaaaaataa atgacacatg    240
gaaataagtc aaggattagc ggatatgtag ttccagtcog gggtatacca tcacgtgata    300
ataaatccaa atgagaatga ggggtgtcata tctaatacatt atgcacgtca agattctccg    360
tgactatggc tcttttctga agcatttttc gggcgcccg tgccaaaaa ctaactccga    420
gcccgggcat gtccgggggt tagcggggcc acaaaaggcg cttatctggg gggttccgt    480
agaagaaaaa aagctgttga gcgagctatt tcgggtatcc cagccttctc tgcagaccgc    540

```

-continued

cccagttggc ttggctctgg tgctgttcgt tagcatcaca tcgcctgtga caggcagagg	600
taataacggc ttaaggttct cttcgcatag tcggcagctt tctttcggac gttga	655
<210> SEQ ID NO 6 <211> LENGTH: 346 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: OLE1 promoter region	
<400> SEQUENCE: 6	
acactcaaca aaccttatct agtgcccaac caggtgtgct tctacgagtc ttgtcactc	60
agacacacct atccctattg ttacggctat ggggatggca cacaagggtg gaaataatag	120
tagttaacaa tatatgcagc aaatcatcgg ctcttggtc atcgagtctt gcaaatcagc	180
atatacatat atatatgggg gcagatcttg attcatttat tgttctatct ccatctttcc	240
tacttctggt tccgtttata ttttgtatta cgtagaatag aacatcatag taatagatag	300
ttgtggtgat catattataa acagcactaa aacattacaa caaaga	346
<210> SEQ ID NO 7 <211> LENGTH: 81 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: OLE1 27aa leader	
<400> SEQUENCE: 7	
atgccaaactt ctggaactac tattgaattg attgacgacc aatttcctaaa ggatgactct	60
gccagcagtg gcattgtcga c	81
<210> SEQ ID NO 8 <211> LENGTH: 103 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Vsp13 terminator region	
<400> SEQUENCE: 8	
tcacatatga aagtatatac ccgcttttgt acactatgta gctataattc aatcgtatta	60
ttgtagctcc gcacgaccat gccttagaaa tatccgcagc gcg	103
<210> SEQ ID NO 9 <211> LENGTH: 1050 <212> TYPE: DNA <213> ORGANISM: Trichoplusia ni	
<400> SEQUENCE: 9	
atggctgtga tggctcaaac agtacaagaa acggctacag tggttgaaga ggaagctcgc	60
acagtgactc ttgtggctcc aaagacaacg ccaaggaaat ataaatatat atacaccaac	120
tttcttacat tttcatatgc gcatttagct gcattatacg gactttatct gtgcttcacc	180
tctgcgaaat gggaaacatt gctattctct ttcgtactct tccacatgct aaatataggc	240
atcaccgcag gggctcaccg actctggact cacaagactt tcaaagccaa attgcctttg	300
gaaattgtcc tcatgatatt caactcttta gcctttcaaa acacggctat tacatgggct	360
agagaacatc ggctacatca caaatacagc gatactgatg ctgatcccca caatgcgtca	420
agagggttct tctactcgca tggtggctgg ctattagtaa aaaaacatcc cgatgtcctg	480
aaatatggaa aaactataga catgtcggat gtatacaata atcctgtgtt aaaatttcag	540

-continued

aaaaagtacg cagtaccctt aattggaaca gtttgttttg ctctgccaac tttgattcca	600
gtctactggt ggggcgaatc gtggaacaac gcttggcaca tagccttatt tcgatacata	660
ttcaatctta acgtgacttt cctagtcaac agtgctgcgc atatctgggg gaataagcct	720
tatgataaaa gcatcttgcc cgctcaaaac ctgctggttt ccttcctagc aagtggagaa	780
ggcttccata attaccatca cgtctttcca tgggattacc gcacagcaga attagggaat	840
aacttcctga atttgacgac gctgttcatt gatTTTTgtg cctggtttgg atgggcttat	900
gacttgaagt ctgtatcaga ggatattata aaacagagag ctaaacgaac aggtgacggt	960
tcttcagggg tcatttgggg atgggacgac aaagacatgg accgcgatat aaaatctaaa	1020
gctaacattt tttatgctaa aaaggaatga	1050

<210> SEQ ID NO 10
<211> LENGTH: 1011
<212> TYPE: DNA
<213> ORGANISM: *Agrotis segetum*

<400> SEQUENCE: 10

atggctcaag gtgtccaaac aactacgata ttgagggagg aggagccgtc attgactttc	60
gtggtacctc aagaaccgag aaagtatcaa atcgtgtacc caaaccttat cacatttggg	120
tactggcata tagctggttt atacgggcta tatttgtgct ttacttcggc aaaatggcaa	180
acaattttat tcagtttcat gctcgttggt ttagcagagt tgggaataac agccggcgct	240
cacaggttat gggcccacaa aacatataaa gcgaagcttc cttacaaat tatectgatg	300
atactgaact ccattgcctt ccaaaattcc gccattgatt gggtgaggga ccaccgtctc	360
catcataagt acagtgcacac tgatgcagac cctcacaatg ctactcgtgg tttcttctat	420
tctcatgttg gatggttgct cgtaagaaaa catccagaag tcaagagacg tggaaaggaa	480
cttgacatgt ctgatattta caacaatcca gtgctgagat ttcaaaagaa gtatgctata	540
cccttcacgc gggcaatgtg cttcggatta ccaactttta tccctgttta cttctgggga	600
gaaacctgga gtaatgcttg gcatatcacc atgcttcggt acatcctcaa cctaaacatt	660
actttcctgg tcaacagtgc tgctcatatc tggggataca aaccttatga catcaaaata	720
ttgcctgccc aaaatatagc agtttccata gtaaccggcg gcgaagtctc cataactacc	780
accacgtttt ttccttggga ttatcgtgca gcagaattgg ggaacaatta tcttaatttg	840
acgactaagt tcatagattt cttcgttggt atcggatggg cttacgatct taagacggtg	900
tccagtgatg ttataaaaag taaggcggaa agaactggtg atgggacgaa tctttggggt	960
ttagaagaca aaggtgaaga agattttttg aaaatctgga aagacaatta a	1011

<210> SEQ ID NO 11
<211> LENGTH: 1440
<212> TYPE: DNA
<213> ORGANISM: *Thalassiosira pseudonana*

<400> SEQUENCE: 11

actagtatgg actttctctc cggcgatcct ttccggacac tcgtccttgc agcacttgtt	60
gtcatcggat ttgctgcggc gtggcaatgc ttctaccgcg cgagcatcgt cggcaagcct	120
cgtacattaa gcaatggtaa actcaatacc agaatccatg gcaaattgta cgacctctca	180
tcgtttcagc atccaggagg ccccggtggc tttctctctt ttcaaggctc cgacggaaca	240
gctctatttg agtcacacca tcccttcata cctcgaaaga atctacttca gatcctctcc	300

-continued

```

aagtacgagg ttccgtcgac tgaagactct gtttccttca tcgccaccct agacgaactc 360
aatggtgaat ctccgtacga ttggaaggac attgaaaatg atgatttcgt atctgacctc 420
cgagctctcg taattgagca cttttctcct ctcgccaagg aaaggggagt ttcactcggt 480
gagtcgtcga aggcaacacc tcagcgggtg atggtggttc tactgctcct tgcgtcgttc 540
ttcctcagca tcccattata tttgagtggg tcgtggactt tcgttgctcg cactcccatc 600
ctcgcttggtc tggcggttgt caattactgg cacgatgcta ctactttgc attgagcagc 660
aactggattt tgaatgctgc gctcccatat ctctccctc tcctatcgag tccgtcaatg 720
tggtatcatc atcacgtcat tggacatcac gcataacca acatttcaa aagagatcca 780
gatcttgctc acgctccaca actcatgaga gaacacaaga gtatcaaatg gagaccatct 840
cacttaaadc aaacacagct tccgcggatt ctcttcatct ggtcgattgc agtcggtatt 900
gggttgaact tactgaacga cgtgagagca ctaaccaagc tttcatacaa caacgttggt 960
cgggtggaga agatgtcatc gtcgcgaaca ttactccatt tccttgagc tatgttgcac 1020
atctttgtga ctacactttg gccctttttg gcgtttccgg tgtggaaggc catcgtttgg 1080
gcgactgtac cgaatgccat actgagtttg tgcttcatgc tgaatacgca aatcaatcac 1140
ctcatcaaca cgtgtgcaca tgcttccgat aacaactttt acaagcatca agttgtaact 1200
gctcagaact ttggccgatc aagtgccttt tgcttcatct tctcgggagg tctcaactac 1260
caaattgaac atcatttggt gccgacggtg aaccattgcc atttgccagc tttggccccg 1320
ggtgtagagc gtttgtgtaa gaaacacggg gtgacataca actctgttga aggatacaga 1380
gaggccatca ttgcacactt tgcacatacc aaagatatgt cgacgaagcc tactgattga 1440

```

<210> SEQ ID NO 12

<211> LENGTH: 981

<212> TYPE: DNA

<213> ORGANISM: Amyelois transitella

<400> SEQUENCE: 12

```

atggtcccta acaagggttc cagtgcggtt ttgtctgaac attctgagcc ccagttcact 60
aaactcatag ctccacaagc agggccgagg aaatacaaga tagtgtatcg aaatttgctc 120
acattcggct attggcactt atcagctggt tatgggctct acttggtgctt tacttggtgcg 180
aaatgggcta ccatcttatt tgcatttttc ttatacgtga tcgcggaaat cgggtataaca 240
ggtggcgctc ataggctatg ggcacatcgg acttataaag ccaagttgcc tttagagatt 300
ttgttactca taatgaactc tattgccttc caagacactg ctttcacctg ggctcgtgat 360
caccgccttc atcacaaata ttcggatact gacgctgatc cccacaatgc taccagaggg 420
tttttctatt cacatgtagg ctggcttttg gtgaagaaac accctgaagt caaagcaaga 480
ggaaaatact tgtcgttaga tgatcttaag aataatccat tgcttaaatt ccaaaagaaa 540
tacgctattc tagttatagg cacgttatgc ttcttatgc caacatttgt gcccgatac 600
ttctggggcg agggcatcag cacggcctgg aacatcaatc tattgcgata cgtcatgaat 660
cttaacatga ctttcttagt taacagtgcg gcgcatatct ttggcaacaa accatacgat 720
aagagcatag cctcagtcca aaatatttca gttagcttag ctacttttgg cgaaggattc 780
cataattacc atcacactta cccctgggat tatcgtgcgg cagaattagg aaataatagg 840
ctaaatatga ctactgcttt catagatttc ttcgcttgga tcggctgggc ttatgacttg 900
aagtctgtgc cacaagaggc cattgcaaaa aggtgtgcga aaactggcga tggaacggat 960
atgtgggggc gaaaaagata a 981

```

<210> SEQ ID NO 13
<211> LENGTH: 1017
<212> TYPE: DNA
<213> ORGANISM: Helicoverpa zea

<400> SEQUENCE: 13
atggcccaaa gctatcaatc aactacggtt ttgagtgagg agaaagaact aacactgcaa 60
catttggtgc cccaagcatc gcccaggaag tatcaaatag tgtatccgaa cctcattacg 120
tttggttact ggcatatagc cggactttat ggcttttact tgtgcttcac ttctgctaaa 180
tgggctacga ttttattcag ctacatcctc ttcgtgtag cagaaatagg aatcacggct 240
ggcgctcaca gactctgggc ccacaaaact tacaaagcga aactaccatt agaaatactc 300
ttaatggtat tcaactccat cgcttttcaa aactcagcca ttgactgggt gagggaccac 360
cgactccacc ataagtatag cgatacagat gctgatcccc acaatgccag ccgaggggtc 420
ttttattccc atgtaggatg gctacttggt agaaaacatc ctgaagtcaa aaagcgaggg 480
aaagaactca atatgtccga tatttacaac aatcctgtcc tgcggtttca gaaaaatac 540
gccataccct tcattggggc tgtttgtttc gccttaccta caatgatacc tgtttacttc 600
tggggagaaa cctggtccaa tgcttggtat atcaccatgc ttcgctacat catgaacctc 660
aatgtcacct ttttggtaaa cagcgctgct catatatggg gaaacaagcc ttatgacgca 720
aaaatattac ctgcacaaaa tgtagctgtg tcggtcgcca ctggtggaga aggtttccat 780
aattaccacc atgtcttccc ctgggattat cgagcagcgg aactcggtaa caatagcctc 840
aatctgacga ctaaattcat agatttattc gcagcaatcg gatgggcata tgatctgaag 900
acggtttcgg aggatatgat aaaacaaagg attaaacgca ctggagatgg aacggatctt 960
tggggacacg aacaaaactg tgatgaagtg tgggatgtaa aagataaatc aagttaa 1017

<210> SEQ ID NO 14
<211> LENGTH: 711
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: mCherry C. tropicalis optimized

<400> SEQUENCE: 14
atggtttcta agggtgaaga agacaacatg gcaatcatca aggaatttat gcgttttaag 60
gtccatatgg aaggctccgt taacggccac gaggtcgaga tcgagggaga aggtgagggg 120
agaccatacg aaggctactc aaccgccaag ttgaaagtta caaagggtgg tccattgcc 180
tttgcttggg atatcttgtc ccacaaattt atgtacggat caaaggcata tgtcaagcat 240
cctgcgcaca tcccagatta cttgaagtta tcctttccag aaggttttta gtgggagaga 300
gttatgaact ttgaagatgg cggagttggt actgttactc aggactcttc cttgcaagat 360
ggtgaattta tctataaagt gaaattgaga ggtactaact ttccatccga cggtcagtc 420
atgcaaaaga agacaatggg ttgggagggt tcttcgaaa gaatgtaccc agaagacggt 480
gcattgaaag gtgaaatcaa gcaacgttta aagttgaagg acggtggtca ctacgatgcc 540
gaggtcaaga ccacttataa ggctaagaag ccagtccaat tgccaggtgc ttataacgtt 600
aacatcaagt tagatattac ttcacacaac gaagactaca caatcgttga acaatatgaa 660
agagccgaag gtagacattc taccggcggc atggacgagt tatataagta g 711

<210> SEQ ID NO 15

-continued

<211> LENGTH: 1092	
<212> TYPE: DNA	
<213> ORGANISM: Agrotis segetum	
<400> SEQUENCE: 15	
atgactacag ttgaacaact tgaaactggt gatataccta aattgaatgc cattgctgct	60
ggtactaata agaaggtgcc aatgggtcaa ggtgtccaaa caactacgat attgagggag	120
gaagagccgt cattgacttt cgtgggtacct caagaaccga gaaagtatca aatcgtgtac	180
ccaaaacctta tcacatttgg gtactggcat atagctggtt tatacgggct atatttgtgc	240
tttacttcgg caaaatggca aacaatttta ttcagtttca tgctcgttgt gttagcagag	300
ttgggaataa cagccggcgc tcacaggtta tgggcccaca aaacatataa agcgaagctt	360
cccttacaaa ttatcttaat gatattaaac tccattgcct tccaaaattc cgccattgat	420
tgggtgaggg accaccgtct ccatcataag tacagtgaca ctgatgcaga ccctcacaat	480
gctactcgtg gtttcttcta ttctcatggt ggatgggtgc tcgtaagaaa acatccagaa	540
gtcaagagac gtggaaagga acttgacatg tctgatattt acaacaatcc agtggttaaga	600
tttcaaaaga agtatgctat acccttcacg ggggcaatgt gcttcggatt accaactttt	660
atccctgttt acttctgggg agaaacctgg agtaatgctt ggcatatcac catgcttcgg	720
tacatcctca acctaaacat tacttttcta gtcaacagtg ctgctcatat ctgggggatac	780
aaaccttatg acatcaaaat attgcctgcc caaaatatag cagtttccat agtaaccggc	840
ggcgaagttt ccataactac caccacgttt tttccttggg attatcgtgc agcagaattg	900
gggaacaatt atcttaattt gacgactaag ttcatagatt tcttcgcttg gatcggtatg	960
gcttacgata ttaagacggt gtccagtgat gttataaaaa gtaaggcgga aagaactggt	1020
gatgggacga atctttgggg tttagaagac aaagggtgaag aagatttttt gaaaatctgg	1080
aaagacaatt aa	1092
<210> SEQ ID NO 16	
<211> LENGTH: 1011	
<212> TYPE: DNA	
<213> ORGANISM: Agrotis segetum	
<400> SEQUENCE: 16	
atgggtcaag gtgtccaaac aactacgata ttgagggagg aagagccgtc attgactttc	60
gtggtacctc aagaaccgag aaagtatcaa atcgtgtacc caaaccttat cacatttggg	120
tactggcata tagctggttt atacgggcta tatttgtgct ttacttcggc aaaatggcaa	180
acaattttat tcagtttcat gctcgttgtg ttagcagagt tgggaataac agccggcgct	240
cacaggttat gggcccacaa aacatataaa gcgaagcttc ccttacaaat tatcttaatg	300
atattaaact ccattgcctt ccaaaattcc gccattgatt gggtgaggga ccaccgtctc	360
catcataagt acagtgacac tgatgcagac cctcacaatg ctactcgtgg tttcttctat	420
ttctcatgtg gatggttgct cgtaagaaaa catccagaag tcaagagacg tggaaaggaa	480
cttgacatgt ctgatattta caacaatcca gtgttaagat ttcaaaagaa gtatgctata	540
cccttcatcg gggcaatgtg cttcggatta ccaactttta tccctgttta cttctgggga	600
gaaacctgga gtaatgcttg gcatatcacc atgcttcggg acatcctcaa cctaaacatt	660
actttcttag tcaacagtgc tgctcatatc tggggataca aaccttatga catcaaaata	720
ttgcctgccc aaaatatagc agtttccata gtaaccggcg gcgaagtttc cataactacc	780
accacgtttt ttccttggga ttatcgtgca gcagaattgg ggaacaatta tcttaatttg	840

-continued

acgactaagt tcatagattt cttcgcttgg atcggatggg cttacgatct taagacggtg	900
tccagtgatg ttataaaaag taaggcggaa agaactgggt atgggacgaa tctttggggt	960
ttagaagaca aaggtgaaga agattttttg aaaatctgga aagacaatta a	1011

<210> SEQ ID NO 17
<211> LENGTH: 981
<212> TYPE: DNA
<213> ORGANISM: Amyelois transitella

<400> SEQUENCE: 17

atggtcoccta acaagggttc cagtgcggtt ttgtctgaac attctgagcc ccagttcact	60
aaactcatag ctccacaagc agggccgagg aaatacaaga tagtgtatcg aaatttgctc	120
acattcggtt attggcactt atcagctgtt tatgggctct acttgtgctt tacttgtgctg	180
aaatgggcta ccattcttatt tgcatttttc ttatacgtga tcgcggaaat cgggtataaca	240
ggtggcgctc ataggctatg ggcacatcgg acttataaag ccaagttgcc tttagagatt	300
ttgttactca taatgaattc tattgccttc caagacactg ctttcacctg ggctcgagat	360
caccgccttc atcaciaaata ttcggatact gacgctgatc cccacaatgc taccagaggg	420
tttttctatt cacatgtagg ctggcttttg gtgaagaaac accctgaagt caaagcaaga	480
ggaaaatact tgctgtaga tgatcttaag aataatccat tgcttaaatt ccaaaagaaa	540
tacgctattc tagttatagg cacgttatgc ttccttatgc caacatttgt gcccgatac	600
ttctggggcg agggcatcag cacggcctgg aacatcaatc tattgcgata cgatcatgaat	660
cttaacatga ctttcttagt taacagtgcg gcgcatactt ttggcaacaa accatacgat	720
aagagcatag cctcagtcca aaatatttca gttagcttag ctacttttgg cgaaggattc	780
cataattacc atcacactta cccctgggat tatcgtgcgg cagaattagg aaataatagg	840
ctaaatatga ctactgcttt catagatttc ttcgcttgga tcggctgggc ttatgacttg	900
aagtctgtgc cacaagaggc cattgcaaaa aggtgtgcga aaactggcga tggaacggat	960
atgtgggggtc gaaaaagata a	981

<210> SEQ ID NO 18
<211> LENGTH: 1050
<212> TYPE: DNA
<213> ORGANISM: Trichoplusia ni

<400> SEQUENCE: 18

atggctgtga tggtcacaac agtacaagaa acggctacag tggtggaaga ggaagctcgc	60
acagtgactc ttgtggctcc aaagacaacg ccaaggaaat ataaatatat atacaccaac	120
tttcttacat ttctcatatgc gcatttagct gcattatacg gactttattt gtgcttcacc	180
tctgcgaaat gggaaacatt gctattctct ttcgtactct tccacatgtc aaatataggc	240
atcacgcgag gggctcaccg actctggact cacaagactt tcaaagccaa attgcctttg	300
gaaattgtcc tcatgatatt caactcttta gcctttcaaa acacggctat tacatgggct	360
agagaacatc ggctacatca caaatacagc gatactgatg ctgatcccca caatgcgtca	420
agaggggttct tctactcgca tggtggctgg ctattagtaa aaaaacatcc cgatgtctta	480
aaatatggaa aaactataga catgtcggat gtatacaata atcctgtgtt aaaatttcag	540
aaaaagtacg cagtaccctt aattggaaca gtttggtttg ctcttccaac tttgattcca	600
gtctactgtt ggggcgaatc gtggaacaac gcttggcaca tagccttatt tcgatacata	660

-continued

ttcaatctta acgtgacttt cctagtcaac agtgctgcgc atatctgggg gaataagcct	720
tatgataaaa gcatcttgcc cgctcaaaac ttattagttt ccttcctagc aagtggagaa	780
ggcttccata attaccatca cgtctttcca tgggattacc gcacagcaga attagggaa	840
aactttcttaa atttgacgac gttattcatt gatTTTTgtg cctggtttg atgggcttat	900
gacttgaagt ctgtatcaga ggatattata aaacagagag ctaaacgaac aggtgacggt	960
tcttcagggg tcatttgggg atgggacgac aaagacatgg accgcgatat aaaatctaaa	1020
gctaacattt tttatgctaa aaaggaatga	1050

<210> SEQ ID NO 19

<211> LENGTH: 1017

<212> TYPE: DNA

<213> ORGANISM: Helicoverpa zea

<400> SEQUENCE: 19

atggcccaaa gctatcaatc aactacggtt ttgagtgagg agaaagaact aacattacaa	60
catttggtgc cccaagcatc gcccaggaag tatcaaata tagtatccga cctcattacg	120
tttggttact ggcacatagc cggactttat ggcctttact tgtgcttcac ttctgctaaa	180
tgggctacga ttttattcag ctacatcctc ttcgtgttag cagaaatagg aatcacggct	240
ggcgctcaca gactctgggc ccacaaaact tacaaaagca aactaccatt agaaatactc	300
ttaatgggat tcaactccat cgcttttcaa aactcagcca ttgactgggt gagggaccac	360
cgactccacc ataagtatag cgatacagat gctgatcccc acaatgccag ccgagggttc	420
ttttattccc atgtaggatg gctacttggt agaaaacatc ctgaagtcaa aaagcgaggg	480
aaagaactca atatgtccga tatttacaac aatcctgtct tacggtttca gaaaaatac	540
gccataccct tcattggggc tgtttgtttc gccttaccta caatgatacc tgtttacttc	600
tggggagaaa cctgggtccaa tgcttggtat atcaccatgc ttcgctacat catgaacctc	660
aatgtcacct ttttggtaaa cagcgctgct catatatggg gaaacaagcc ttatgacgca	720
aaaatattac ctgcacaaaa tgtagctgtg tcggctcgca ctgggtggaga aggtttccat	780
aattaccacc atgtcttccc ctgggattat cgagcagcgg aactcggtaa caatagcctc	840
aatttaacga ctaaattcat agatttattc gcagcaatcg gatgggcata tgatttaaag	900
acggtttcgg aggatatgat aaaacaaagg attaaacgca ctggagatgg aacggatctt	960
tggggacacg aacaaaactg tgatgaagtg tgggatgtaa aagataaatc aagttaa	1017

<210> SEQ ID NO 20

<211> LENGTH: 1068

<212> TYPE: DNA

<213> ORGANISM: Ostrinia furnacalis

<400> SEQUENCE: 20

atggctccta atattaagga cggagctgat ttgaacggag ttttatttga agatgacgct	60
agcacccccg attatgccct tgccacggcc ccagtcaga aagcagacaa ctatcccaga	120
aaactagtgt ggagaaacat catactcttt gcataccttc accttgccgc tgtgtatgga	180
gcatacctat tcttattttc agcgaaatgg cagacagata tttttgccta cattctttac	240
gtgatctcag gactcggcat cacagcggga gccacccgcc tttgggcgca caagtcatac	300
aaggctaagt ggccacttag actcattctt attatcttca aactgtatc attccaggac	360
tctgctctcg actggtcacg tgaccaccgc atgcaccaca aatactcgga gaccgacgcc	420
gaccgcaca acgcgactcg agggttcttc ttctctcata tcggctgggt attagtccgc	480

-continued

```

aagcaccggg aattaaagag aaagggcaag ggattagact taagcgactt gtatgctgat 540
cccatcctcc gtttccagaa gaagtactat ttactattaa tgctctttgg ctgcttcac 600
atgccgacgg tgggtcccgg gtacttctgg ggtgagactt ggactaacgc tttcttcgtc 660
gccgcgctct tccgatacac cttcatcctc aatgtcacct ggttgggtcaa ctccgccggc 720
cacaagtggg gccacaagcc ctatgacagc agcatcaagc cttccgagaa cctctcagtc 780
tccttattcg cgttgggcga aggattccac aactaccacc acacattccc ctgggactac 840
aaaactgccg agctcggcaa caacagactc aatttcacaa caaacttcat caacttcttc 900
gctaaaatcg gatgggctta cgacttgaac acggtctccg acgagattat tcagaataga 960
gtcaagcgca caggagatgg ctcccaccac ttatgggggt ggggcgacaa ggatcaacct 1020
aaagaggagg taaacgcagc cattagaatt aatcctaaag acgagtaa 1068

```

```

<210> SEQ ID NO 21
<211> LENGTH: 1059
<212> TYPE: DNA
<213> ORGANISM: Lampronia capitella

```

```

<400> SEQUENCE: 21

```

```

atgccgccga acgtgacaga ggccaacgga gtgttatttg agaatgacgt gcagactcct 60
gacatggggc tagaagtggc ccctgtgcag aaggctgacg agcgtaagat ccagctcggt 120
tggaggaaca tcatcgcttt tgcatgtctt catttagcag ctgtgtatgg agcttattta 180
ttcttcacct cggctatatg gcagacagac atatttgcac acatccttta cgttatgtct 240
ggattaggaa tcacggcggg agcgcacaga ttatgggctc ataagtcata caaggcgaag 300
tggcggtaaa gattaatcct cgtcgcattc aacactttgg cattccagga ttccggcaatc 360
gactggggcg gcgaccaccg catgcaccac aagtactcgg agacggatgc ggaccacat 420
aacgccactc gcggcttctt cttttcgcac attggttggg tactctgccg aaaacacccg 480
gagctaaagc gcaagggcca gggcctcgac ttaagtgacc tctacgcaga tcctattatt 540
cgcttccaaa agaagtacta cttattgtta atgccgttag cctgctttgt tcttcccacc 600
ataattccgg tctacctctg gggcgagtcc tggaaaaacg cgttcttcgt agctgcaatg 660
ttccgttaca cgttcatcct caacgtaaca tggctcgtea actccgccgc ccacaaatgg 720
ggaggcaagc cctatgataa gaacatccag cccgctcaga acatctctgt agctatcttc 780
gcattaggcg agggcttcca caactaccac cacacgttcc cctgggacta caagaccgct 840
gaattaggaa acaacagggt aaatttcaca acttcgttta tcaatttctt cgcaagcttc 900
ggatgggcct acgacttaaa gaccgtgtcg gacgagatta tacaacagcg cgtaagagg 960
acgggagatg ggagccatca cttacggggc tggggcgacc aggacatacc ggccgaagaa 1020
gctcaagctg ctttacgcat taaccgtaaa gatgattag 1059

```

```

<210> SEQ ID NO 22
<211> LENGTH: 1062
<212> TYPE: DNA
<213> ORGANISM: Helicoverpa zea

```

```

<400> SEQUENCE: 22

```

```

atggctccaa atatatcgga ggatgtgaac ggggtgctct tcgagagtga tgcagcgacg 60
ccggacttag cgttatccac gccgcctgtg cagaaggctg acaacaggcc caagcaatta 120
gtgtggagga acatactatt attcgcgtat cttcacttag cggctcttta cggagggtat 180

```

-continued

ttattcctct	tctcagctaa	atggcagaca	gacatatttg	cctacatctt	atatgtgatc	240
tccgggcttg	gtatcacggc	tggagcacat	cgcttatggg	cccacaagtc	ctacaaagct	300
aaatggcctc	tccgagttat	cttagtcatc	tttaacacag	tggcattcca	ggatgccgct	360
atggactggg	cgcgcgacca	cgcgatgcac	cacaagtact	cggaaaccga	tgctgatcct	420
cataatgcga	cccgaggatt	cttcttctct	cacattggct	ggttacttgt	caggaaacat	480
cccgacctta	aggagaaggg	caagggactc	gacatgagcg	acttacttgc	tgaccccatc	540
ctcaggttcc	agaaaaata	ctacttaatc	ttaatgccct	tggcttgctt	cgtgatgcct	600
accgtgattc	ctgtgtactt	ctggggtgaa	acctggacca	acgcattctt	tgtggcggcc	660
atgttccgct	acgcgttcac	cctaaatgtg	acgtggctcg	tcaactctgc	cgtccacaag	720
tggggagaca	agccctacga	caaaagcatt	aagccttcgg	aaaacttgtc	ggtcgccatg	780
ttcgctctcg	gagaaggatt	ccacaactac	caccacactt	tcccttggga	ctacaaaact	840
gctgagttag	gcaacaacaa	actcaacttc	actaccacct	ttattaactt	cttcgctaaa	900
attggctggg	cttacgactt	aaagacagtg	tctgatgata	tcgtcaagaa	cagggatgaag	960
cgcactggtg	acggctccca	ccacttatgg	ggctggggag	acgaaaatca	atccaaagaa	1020
gaaattgatg	ccgctatcag	aatcaatcct	aaggacgatt	aa		1062

<210> SEQ ID NO 23

<211> LENGTH: 1434

<212> TYPE: DNA

<213> ORGANISM: *Thalassiosira pseudonana*

<400> SEQUENCE: 23

atggactttc	tctccggcga	tcctttccgg	acactcgtec	ttgcagcact	tgttgtcac	60
ggatttgctg	cggcgtggca	atgcttctac	ccgccgagca	tcgtcggcaa	gcctcgtaca	120
ttaagcaatg	gtaaactcaa	taccagaatc	catggcaa	atgtacgac	ctcatcgttt	180
cagcatccag	gaggccccgt	ggctctttct	cttggtcaag	gtcgcgacgg	aacagctcta	240
tttgagtcac	accatccctt	catacctcga	aagaatctac	ttcagatcct	ctccaagtac	300
gaggttccgt	cgactgaaga	ctctgtttcc	ttcatcgcca	ccctagacga	actcaatggt	360
gaatctccgt	acgattggaa	ggacattgaa	aatgatgatt	tcgtatctga	cctacgagct	420
ctcgtaatg	agcacttttc	tcctctcgcc	aaggaaaggg	gagtttctact	cgttgagtcg	480
tcgaaggcaa	cacctcagcg	gtggatgggt	gttctattac	tccttgcgtc	gttcttcctc	540
agcatcccat	tatatattgag	tggttcgtgg	actttcgttg	tcgtcactcc	catectcgct	600
tggttagcgg	ttgtcaatta	ctggcacgat	gctactcact	ttgcattgag	cagcaactgg	660
attttgaatg	ctgcgctccc	atatctcctc	cctctcctat	cgagtcgctc	aatgtgggat	720
catcatcacg	tcattggaca	tcacgcatac	accaacattt	ccaaaagaga	tccagatctt	780
gctcacgctc	cacaactcat	gagagaacac	aagagtatca	aatggagacc	atctcactta	840
aatcaaacac	agcttccgcg	gattctcttc	atctggtcga	ttgcagtcgg	tattggggtg	900
aacttattaa	acgacgtgag	agcactaacc	aagctttcat	acaacaacgt	tgttcgggtg	960
gagaagatgt	catcgtcgcg	aacattactc	catttccttg	gacgtatgtt	gcacatcttt	1020
gtgactacac	tttggccctt	tttggcggtt	ccggtgtgga	aggccatcgt	ttgggcgact	1080
gtaccgaatg	ccatattaag	tttgtgcttc	atgttaaata	cgcaaatcaa	tcacctcatc	1140
aacacgtgtg	cacatgcttc	cgataacaac	ttttacaagc	atcaagttgt	aactgctcag	1200
aactttggcc	gatcaagtgc	cttttgcttc	atcttctcgg	gaggtctcaa	ctaccaaat	1260

-continued

gaacatcatt tgttgccgac ggtgaaccat tgccatttgc cagctttggc cccgggtgta	1320
gagcgtttgt gtaagaaaca cggggtgaca tacaactctg ttgaaggata cagagaggcc	1380
atcattgcac actttgcaca taccaaagat atgtcgacga agcctactga ttga	1434

<210> SEQ ID NO 24
 <211> LENGTH: 1050
 <212> TYPE: DNA
 <213> ORGANISM: Trichoplusia ni
 <400> SEQUENCE: 24

atggctgtga tggctcaaac agtacaagaa acggctacag tgttggaaga ggaagctcgc	60
acagtgactc ttgtggctcc aaagacaacg ccaaggaaat ataaatatat atacaccaac	120
tttcttacat tttcatatgc gcatttagct gcattatacg gactttattt gtgcttcacc	180
tctgcgaaat gggaaacatt gctattctct ttcgtactct tccacatgtc aaatataggg	240
atcacgcgag gggctcaccg actctggact cacaagactt tcaaagccaa attgcctttg	300
gaaattgtcc tcatgatatt caactcttta gcctttcaaa acacggctat tacatgggct	360
agagaacatc ggctacatca caaatacagc gatactgatg ctgatcccca caatgcgtca	420
agagggttct tctactcgca tgttggtctg ctattagtaa aaaaacatcc cgatgtcctg	480
aaatatggaa aaactataga catgtcggat gtatacaata atcctgtgtt aaaatttcag	540
aaaaagtacg cagtaccctt aattggaaca gtttgttttg ctctgccaac tttgattcca	600
gtctactgtt ggggcgaatc gtggaacaac gcttggcaca tagccttatt tcgatacata	660
ttcaatctta acgtgacttt cctagtcaac agtgctgcgc atatctgggg gaataagcct	720
tatgataaaa gcactctgcc cgctcaaaac ctgctggttt ccttcctagc aagtggagaa	780
ggcttccata attaccatca cgtctttcca tgggattacc gcacagcaga attagggaat	840
aacttctga atttgacgac gctgttcatt gatTTTTgtg cctggtttgg atgggcttat	900
gacttgaagt ctgtatcaga ggatattata aaacagagag ctaaacgaac aggtgacggt	960
tcttcagggg tcatttgggg atgggacgac aaagacatgg accgcgatat aaaatctaaa	1020
gctaacattt tttatgctaa aaaggaatga	1050

<210> SEQ ID NO 25
 <211> LENGTH: 1017
 <212> TYPE: DNA
 <213> ORGANISM: Helicoverpa zea
 <400> SEQUENCE: 25

atggcccaaa gctatcaatc aactacgggt ttgagtgagg agaaagaact aacactgcaa	60
catttggtgc cccaagcatc gccaggaag tatcaaatag tgtatccgaa cctcattacg	120
tttggttact ggcacatagc cggactttat ggcttttact tgtgcttcac ttctgctaaa	180
tgggctacga ttttattcag ctacatcctc ttcgtgttag cagaaatagg aatcacggct	240
ggcgtcaca gactctgggc ccacaaaact tacaaaagca aactaccatt agaaatactc	300
ttaatggtat tcaactccat cgcttttcaa aactcagcca ttgactgggt gagggaccac	360
cgactccacc ataagtatag cgatacagat gctgatcccc acaatgccag ccgagggttc	420
ttttattccc atgtaggatg gctacttgtg agaaaacatc ctgaagtcaa aaagcgaggg	480
aaagaactca atatgtccga tatttacaac aatcctgtcc tgcggtttca gaaaaatac	540
gccataccct tcattggggc tgtttgtttc gccttaccta caatgatacc tgtttacttc	600

-continued

tggggagaaa cctggtccaa tgcttggcat atcaccatgc ttcgctacat catgaacctc	660
aatgtcacct ttttggtaaa cagcgctgct catatatggg gaaacaagcc ttatgacgca	720
aaaatattac ctgcacaaaa tgtagctgtg tcggtcgcca ctggtggaga aggtttccat	780
aattaccacc atgtcttccc ctgggattat cgagcagcgg aactcggtaa caatagcctc	840
aatctgacga ctaaattcat agatttattc gcagcaatcg gatgggcata tgatctgaag	900
acggtttcgg aggatatgat aaaacaaagg attaaacgca ctggagatgg aacggatctt	960
tggggacacg aacaaaactg tgatgaagtg tgggatgtaa aagataaatc aagttaa	1017

<210> SEQ ID NO 26
 <211> LENGTH: 1050
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Trichoplusia ni Z11 desaturase Homo sapiens
 optimized

<400> SEQUENCE: 26

atggccgtga tggcccagac cgtgcaggag accgcaacag tgctggagga ggaggcaagg	60
accgtgacac tgggtggcacc caagaccaca cctagaaagt acaagtatat ctacaccaac	120
ttcctgacct tcagctacgc acacctggcc gccctgtatg gactgtacct gtgctttacc	180
tccgccaaagt gggagacact gctgttctct tttgtgctgt tccacatgag caatatcgga	240
atcacccgag gagcacacag gctgtggacc cacaagacat tcaaggccaa gctgcctctg	300
gagatcgtgc tgatgatctt caactctctg gcctttcaga ataccgccat cacatgggcc	360
cgggagcaca gactgcacca caagtatagc gacaccgatg cagaccaca caacgcaagc	420
aggggcttct tttactccca cgtgggctgg ctgctggatg agaagcacc cgacgtgctg	480
aagtatggca agacaatcga catgtccgac gtgtacaaca atcccgtgct gaagtttcag	540
aagaagtatg ccgtgcctct gatcggcacc gtgtgcttcg ccctgccaac actgatcccc	600
gtgtattgtt ggggcgagtc ttggaacaat gcctggcaca tcgccctgtt ccggtacatc	660
tttaacctga atgtgacctt tctggatgaac tccgccgccc acatctgggg caataagcct	720
tacgacaagt ctatcctgcc agcccagaac ctgctgggtg ccttcctggc ctctggcgag	780
ggctttcaca attatcacca cgtgttccca tgggactaca ggaccgcaga gctgggcaac	840
aattttctga acctgaccac actgttcacg gatttttgtg cctgggtcgg ctgggcctat	900
gacctgaagt ctgtgagcga ggatatcatc aagcagaggg caaagaggac aggcgatggc	960
agctccggcg tgatctgggg atgggacgat aaggatatgg acagagatat caagagcaag	1020
gccaatatct tctacgcaa gaaggagtga	1050

<210> SEQ ID NO 27
 <211> LENGTH: 1017
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Helicoverpa zea Z11 desaturase Homo sapiens
 optimized

<400> SEQUENCE: 27

atggcacagt catatcagag cactaccgtc ctgagcgaag agaaggaaact gacactgcag	60
cacctgggcc cacaggcatc acctagaaag taccagatcg tgtatccaaa cctgatcacc	120
ttcggtact ggcacatcgc cggcctgtac ggctgtatc tgtgctttac ctccgccaa	180
tgggccacaa tcctgttctc ttacatcctg tttgtgctgg cagagatcgg aatcacccga	240

-continued

ggagcacaca gactgtgggc acacaagaca tataaggcca agctgcccct ggagatcctg	300
ctgatgggtgt tcaacagcat cgccctttcag aattccgcca tcgattgggt gcgggaccac	360
agactgcacc acaagtactc cgacaccgat gccgaccccc acaacgcctc taggggcttc	420
ttttatagcc acgtgggatg gctgctgggt cggaagcacc ctgaggtgaa gaagagaggc	480
aaggagctga atatgtctga tatctacaac aatcctgtgc tgcgcttcca gaagaagtat	540
gccatcccat tcatcgggcg cgtgtgcttt gccctgcccc ccatgatccc cgtgtacttt	600
tggggcgaga catggagcaa cgctggcac atcacaatgc tgcggtatat catgaacctg	660
aatgtgacat tcctggtgaa ctccgccgcc cacatctggg gcaataagcc atacgacgcc	720
aagatcctgc ccgccagaa cgtggccgtg agcgtggcaa ccggaggaga gggcttccac	780
aattaccacc acgtgtttcc ttgggattat cgggccgccg agctgggcaa caattctctg	840
aatctgacca caaagttcat cgacctgttt gccgccatcg gctgggccta tgatctgaag	900
acagtgagcg aggacatgat caagcagagg atcaagcgca ccggcgatgg cacagacctg	960
tgggggcacg agcagaactg tgatgaagtg tgggatgtga aagacaagtc ctctaa	1017

<210> SEQ ID NO 28

<211> LENGTH: 1065

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Yarrowia lipolytica OLE1 leader - Trichoplusia ni Z11 desaturase Homo sapiens optimized

<400> SEQUENCE: 28

atggtgaaga acgtggacca ggtggatctg tctcaggtgg acaccatcgc aagcggaagg	60
gatgtgaatt ataagtgaa gtacacatct ggcgtaaga ccacaccaag aaagtacaag	120
tatatctaca ccaacttcct gacatcttct tacgcccacc tggccgccct gtatggcctg	180
tacctgtgct ttaccagcgc caagtgggag aactgctgt tctcctttgt gctgttccac	240
atgtctaata tcggaatcac cgcaggagca cacaggctgt ggaccacaa gacattcaag	300
gccaagetgc ccctggagat cgtgctgatg atcttcaact ccctggcctt tcagaatacc	360
gccatcacat gggcccggga gcacagactg caccacaagt attctgacac cgatgcagac	420
ccacacaacg caagcagggg cttcttttac tcccacgtgg gctggctgct ggtgaagaag	480
cacctgacg tgctgaagta tggcaagaca atcgacatga gcgacgtgta caacaatcct	540
gtgctgaagt ttcagaagaa gtatgccgtg cactgatcg gcaccgtgtg cttegccctg	600
cccacactga tccccgtgta ctgttggggc gagtcttggg acaatgcctg gcacatcgcc	660
ctgttcgggt acatctttaa cctgaatgtg acctttctgg tgaacagcgc cgcccacatc	720
tggggcaata agccatacga caagtccatc ctgcccgcc agaacctgct ggtgtccttc	780
ctggcctctg gcgagggtt tcacaattat caccacgtgt tcccttggga ctacaggacc	840
gcagagctgg gcaacaattt tctgaacctg accacactgt tcatcgattt ttgtgcctgg	900
ttcggtggg cctatgacct gaagtctgtg agcgaggata tcatcaagca gagggcaaag	960
aggacaggcg atggcagctc cggcgtgatc tggggatggg acgataagga tatggacaga	1020
gatatcaagt ccaaggccaa tatcttctac gccaaagaag agtga	1065

<210> SEQ ID NO 29

<211> LENGTH: 1029

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Yarrowia lipolytica OLE1 leader - Helicoverpa
zea Z11 desaturase Homo sapiens optimized

<400> SEQUENCE: 29

```

atggtgaaaa acgtggacca agtggatctc tcgcaggctcg acaccattgc ctccggccga      60
gatgtcaact acaagggtcaa gtacacctcc ggcgttcgca agtatcagat cgtgtatcct      120
aacctgatca ccttcgggta ctggcatatc gctggactgt acggactgta tctgtgcttc      180
acttcgcca agtggggccac catectgttc tcttacatcc tgtttgtgct ggcagagatc      240
ggaatcaccg caggagcaca cagactgtgg gcacacaaga catataaggc caagctgcca      300
ctggagatcc tgctgatggg gttcaacagc atcgcccttc agaattccgc catcgattgg      360
gtgcgggacc acagactgca ccacaagtac tccgacacag atgccgaccc ccacaacgcc      420
tctaggggct tcttttatag ccacgtggga tggctgctgg tgcggaagca ccctgagggtg      480
aagaagagag gcaaggagct gaatatgtct gatatctaca acaatcctgt gctgcgcttc      540
cagaagaagt atgccatccc attcatcggc gccgtgtgct ttgccctgcc caccatgatc      600
cccgtgtact tttggggcga gacatggagc aacgcctggc acatcacaat gctgcggtat      660
atcatgaacc tgaatgtgac attcctgggtg aactccgccg cccacatctg gggcaataag      720
ccatacgacg ccaagatcct gcccggcccag aacgtggccg tgagcgtggc aaccggagga      780
gagggcttcc acaattacca ccacgtgttt ccatgggatt atagggcagc agagctggga      840
aacaattctc tgaatctgac cacaagttc atcgacctgt ttgccgccat cggetgggcc      900
tatgatctga agacagtgag cgaggacatg atcaagcaga ggatcaagcg caccggcgat      960
ggcacagacc tgtgggggca cgagcagaat tgtgatgaag tgtgggatgt gaaggataaa     1020
agcagttga                                     1029

```

<210> SEQ ID NO 30

<211> LENGTH: 981

<212> TYPE: DNA

<213> ORGANISM: Amyelois transitella

<400> SEQUENCE: 30

```

atggtcccta acaagggttc cagtgcggtt ttgtctgaac attctgagcc ccagttcact      60
aaactcatag ctccacaagc agggccgagg aaatacaaga tagtgtatcg aaatttgctc      120
acattcggct attggcactt atcagctgtt tatgggctct acttgtgctt tacttgtgctg      180
aaatgggcta ccatcttatt tgcatttttc ttatacgtga tcgcggaaat cgggtataaca      240
ggtggcgctc ataggctatg ggcacatcgg acttataaag ccaagttgcc tttagagatt      300
ttgttactca taatgaattc tattgccttc caagacactg ctttcacctg ggctcgagat      360
caccgccttc atcaciaaata ttcggatact gacgctgatc cccacaatgc taccagaggg      420
tttttctatt cacatgtagg ctggcctttg gtgaagaaac accctgaagt caaagcaaga      480
ggaaaatact tgtcgtaga tgatcttaag aataatccat tgcttaaatt ccaaaagaaa      540
tacgctattc tagttatagg cacgttatgc ttccttatgc caacatttgt gcccgatac      600
ttctggggcg agggcatcag cacggcctgg aacatcaatc tattgcgata cgtcatgaat      660
cttaacatga ctttcttagt taacagtgca gcgcatatct ttggcaacaa accatacgat      720
aagagcatag cctcagttca aaatatttca gttagcttag ctacttttgg cgaaggattc      780
cataattacc atcacactta cccctgggat tatcgtgcgg cagaattagg aaataatagg      840
ctaaatatga ctactgcttt catagatttc ttcgcttgga tcggctgggc ttatgacttg      900

```

-continued

aagtctgtgc cacaagaggc cattgcaaaa aggtgtgcga aaactggcga tggaacggat 960
atgtgggggtc gaaaaagata a 981

<210> SEQ ID NO 31
<211> LENGTH: 1017
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPV0228 - Z11 Helicoverpa zea desaturase

<400> SEQUENCE: 31

atggcccaaa gctatcaatc aactacgggt ttgagtgagg agaaagaact aacattacaa 60
catttggtgc cccaagcatc gcccaggaag tatcaaatag tgtatccgaa cctcattacg 120
tttggttact ggcacatagc cggactttat ggcttttact tgtgcttcac ttctgctaaa 180
tgggctacga ttttattcag ctacatcctc ttcgtgttag cagaaatagg aatcacggct 240
ggcgctcaca gactctgggc ccacaaaact tacaaagcga aactaccatt agaaatactc 300
ttaatggtat tcaactccat cgcttttcaa aactcagcca ttgactgggt gagggaccac 360
cgactccacc ataagtatag cgatacagat gctgatcccc acaatgccag ccgaggggtc 420
ttttattccc atgtaggatg gctacttgtg agaaaacatc ctgaagtcaa aaagcgaggg 480
aaagaactca atatgtccga tatttacaac aatcctgtct tacggtttca gaaaaatac 540
gccataccct tcattggggc tgtttggttc gccttaccta caatgatacc tgtttacttc 600
tggggagaaa cctggtccaa tgcttggtat atcaccatgc ttcgctacat catgaacctc 660
aatgtcacct ttttggtaaa cagcgctgct catatatggg gaaacaagcc ttatgacgca 720
aaaatattac ctgcacaaaa tgtagctgtg tcggtcgcca ctgggtggaga aggtttccat 780
aattaccacc atgtcttccc ctgggattat cgagcagcgg aactcggtaa caatagcctc 840
aatttaacga ctaaattcat agatttatct gcagcaatcg gatgggcata tgatttaaag 900
acggtttcgg aggatatgat aaaacaaagg attaaacgca ctggagatgg aacggatctt 960
tggggacacg aacaaaactg tgatgaagtg tgggatgtaa aagataaatc aagttaa 1017

<210> SEQ ID NO 32
<211> LENGTH: 1368
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPV0228 - Helicoverpa armigera reductase codon optimized

<400> SEQUENCE: 32

atggtcgttt taacttctaa agagacaaaa ctttcagtag ctgagtttta tgccggaaaa 60
tctgttttta ttacgggtgg cactggattc cttggaaagg tattcataga gaaactttta 120
tatagctgtc cagatatcga gaatatctac atgtccatac gagagaagaa aggtctttct 180
gttagcgaaa gaataaaaca gttccttgat gaccgcgtct ttaccagact aaaagacaaa 240
agaccagctg acttagagaa gattgtatta ataccaggag atattactgc tcctgactta 300
ggcattaatt ctgaaaacga gaagatgctt atagagaagg tatcggtgat tattcattcg 360
gctgctacgg tgaagtttaa tgagcctctc cctacggctt ggaagatcaa cgtggaagga 420
accagaatga tgtagcttt gagtcgaaga atgaagcgga ttgaggtttt cattcacata 480
tcgacagcat acacgaacac aaacagggaa gtggttgacg agatcttata cccagctcct 540
gctgatatcg accaagttca tcagtatgtc aaagatggaa tctctgagga agacactgag 600

-continued

aaaatattaa atggtcgtcc aaatacgtac acgtttacga aagcgttaac tgagcattta	660
gttgctgaga accaagccta cgtaccact attatcgtea ggccgtcagt cgtggcagca	720
ataaaagatg agccattaaa aggttgggta ggcaactggg ttggagcgac tggcttcacc	780
gtgttcaccg ctaagggctc caaccgagtc atctacggtc attctagcta catcgtagac	840
ttaattcctg tggattatgt cgctaattta gtgattgctg ctggggctaa gagtagcaag	900
tcaactgagt tgaaggtata caactgctgc agcagctcct gcaatcccgt cactattggc	960
acgttaatga gcatgtttgc tgacgatgcc atcaaacaga agtcgtatgc tatgccgcta	1020
ccgggggtggg acatattcac gaaatataag tggttagttc ttcttttaac atttctcttc	1080
caagttatac cggcgtatgt cacagatctc tccaggcact tgattgggaa gaggccacgg	1140
tacataaaac tccaatcact agtaaatcaa acgcgctctt caatcgactt cttcacgaat	1200
cactcctggg tgatgaaggc agacagagtg agagagttat atgcgtctct tcccccgca	1260
gacaagtact tatttcctg tgatcctacg gacattaact ggacacatta catacaagac	1320
tactgttggg gaggccgaca ttttttgag aaaaaagct acgaataa	1368

<210> SEQ ID NO 33

<211> LENGTH: 1478

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: pPV0228 - ICL promoter

<400> SEQUENCE: 33

tattaggcga agaggcatct agtagtagtg gcagtgggga gaacgtgggc gctgctatag	60
tgaacaatct ccagtcgatg gttaagaaga agagtgacaa accagcagtg aatgacttgt	120
ctgggtccgt gaggaagaaga aagaagcccg acacaaagga cagtaacgtc aagaaccca	180
agaaataggg gggacctgtt tagatgtata ggaataaaaa ctccgagatg atctcaatgt	240
gtaatggagt tgtaatatgt caaaggggga aaatcaagac tcaaactgtg gtatgagtga	300
gcgtacgtat atctccgaga gtagtatgac ataattgatga ctgtgaatca tcgtaatctc	360
acacaaaaac cccattgtcg gccatatacc acaccaagca acaccacata tccccggaa	420
aaaaaacgt gaaaaaaga aacaatcaaa actacaacct actccttgat cacacagtca	480
ttgatcaagt tacagttcct gctaggggaat gaccaaggta caaatcagca ccttaatggg	540
tagcacgctc tcttactctc tctcacagtc ttccggcccc tattcaaaat tctgcacttc	600
catttgacct cagggttggg aaacagggcc acaaaagaaa aacccgacgt gaatgaaaa	660
actaagaaaa gaaaaaaat tatcacacca gaaatttacc taattgggta attcccatcg	720
gtgtttttcc tggattgtcg cacgcacgca tgctgaaaaa agtggttcgag ttttgctttt	780
gcctcggagt ttcacgcaag tttttcgatc tcggaaccgg agggcggtcg ccttggtgtt	840
tgtgatgtcg tgctttgggt gttctaatgt gctgttattg tgctcttttt tttcttctt	900
tttttggtga tcatatgata ttgctcggtg gattactttc gtgtgtaggt attcttttag	960
acgtttggtt attgggtaga tatgagagag agagagtggg tgggggagga gttggttgta	1020
ggagggacct ctgggaggaa gtgtagttga gttttccctg acgaatgaaa atacgttttt	1080
gagaagataa tacaggaaag gtgtgtcggg gaatttccat ctatccgagg atatgagtgg	1140
aggagagtcg tgtgcgtgtg gttaatttag gatcagtggg acacacaaag taactaagac	1200
agagagacag agagaaaaat ctggggaaga gacaaagagt cagagtgtgt gagttattct	1260

-continued

gtattgtgaa atttttttgc ccaactacat aatattgctg aaactaattt tacttaaaaa	1320
gaaaagccaa caacgtcccc agtaaaactt ttctataaat atcagcagtt ttccctttcc	1380
tccattcctc ttcttgctt ttttcttact ttcccttttt tatacctttt cattatcatc	1440
ctttataatt gtctaaccaa caactatata tctatcaa	1478

<210> SEQ ID NO 34
<211> LENGTH: 586
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPV0228 - TEF Candida tropicalis promoter region

<400> SEQUENCE: 34

aggaagacaa ccaaaagaaa gatcaaattg actaaatggt gaacagacca aaaaaaaga	60
acaacaaata gataaattac aacatattaa tcttttgata tgttggtgaa tattctagta	120
aatctaata tctcaatagt ggttatcatt cactctcttc gtctcctct ctccccctct	180
cctcttgag tatattaaaa gcaataaaaa aaaaaaaaaa aagaaaatct gccaacacac	240
acaaaaaaaa cttacatagt cgtgtaccag tgtcaatatt tcaccagcg agagaaaaga	300
agatgaacag aaaaattttc tctttgggtt tgtctttggt tttgtattaa tctcattgaa	360
aaattttttc tctctctctc tctctctctc tcactcacac actcactcgc atttcgtttg	420
ggttacagca gaagtcagac agaaaaaaaa aatcgtatat aactctcatc aaatgcccta	480
gagaaaaatt tttcttctat cttttttttt ttcttcttct tcttcttttc cttttttctt	540
ttagaagatc tttttgaatt catcaaagat atatatttaa tcaatc	586

<210> SEQ ID NO 35
<211> LENGTH: 297
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPV0228 - ICL terminator

<400> SEQUENCE: 35

aagaaaaag aaaaggtaaa gaacttcatt tgagatgaac ttttgtatat gacttttagt	60
ttctactttt ttttttattt attgcttaat tttctttatt tcaatcccc atagtttggtg	120
tagaatatat ttattcattc tggttaactca aacacgtagc aagctcgttg catctcgcct	180
cgtcacgggt acagctctgg aaccaaagac aaaaaaaaaa gttgatccga accctctcgc	240
tattccttgc tatgctatcc acgagatggg gtttatcagc ccaggcaagt cactaaa	297

<210> SEQ ID NO 36
<211> LENGTH: 217
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: pPV0228 - TEF terminator

<400> SEQUENCE: 36

gctgattaat gaataattaa taagtattgt ttttttggtt tttaatatat atatatttg	60
aaattagtat aaaaaaatc tttttttttt cttttttatt tattttatca atagtttata	120
tatatatata tataaacttg taagagatta ggtatatcta acagtgatac tactaatagt	180
gcttaatatc tttgttaaac aagaaaataa aataaac	217

<210> SEQ ID NO 37

-continued

```

<211> LENGTH: 2487
<212> TYPE: DNA
<213> ORGANISM: artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: SapI-tLIP2-pEXP1-HA_FAR-SapI (insert into
    pPV199 creating pPV247)

<400> SEQUENCE: 37

gcctgaagag cgctatttat cactctttac aacttctacc tcaactatct actttaataa      60
atgaatatcg tttattctct atgattactg tatatgcggt cctccatggg agtttggcgc      120
ccgttttttc gagccccaca cgtttcgggt agtatgagcg gcggcagatt cgagcgtttc      180
cggtttccgc ggctggacga gagcccatga tgggggctcc caccaccagc aatcagggcc      240
ctgattacac acccacctgt aatgtcatgc tgttcacgtg ggtaaatgct gctgtgtgct      300
gtgtgtgtgt gttgtttggc gctcattgtt gcgttatgca gcgtacacca caatattgga      360
agcttattag cctttctatt ttttcgtttg caaggcttaa caacattgct gtggagaggg      420
atggggatat ggaggccgct ggagggagtc ggagaggcgt tttggagcgg cttggcctgg      480
cgcccagctc gcgaaacgca cctaggacct tttggcacgc cgaaatgtgc cacttttcag      540
tctagtaacg ccttacctac gtcattccat gcatgcatgt ttgcgccttt tttcccttgc      600
ccttgatcgc cacacagtac agtgcactgt acagtggagg ttttgggggg gtcttagatg      660
ggagctaaaa gcggcctagc ggtacactag tgggattgta tggagtggca tggagcctag      720
gtggagcctg acaggacgca cgaccggcta gcccgtagca gacgatgggt ggctcctgtt      780
gtccaccgcg tacaaatgtt tgggccaaag tcttgtcagc cttgcttgcg aacctaattc      840
ccaattttgt cacttcgcac cccattgat  cgagccctaa cccctgcca tcaggcaatc      900
caattaagct cgcattgtct gccttgttta gtttggtccc tgcccgtttc ggcgccact      960
tgcacaaaca caaacaagca ttatatataa ggctcgtctc tccctcccaa ccacactcac     1020
ttttttgccc gtcttccctt gctaacacaa aagtcaagaa cacaacaac caccacaacc     1080
cccttacaca caagacatat ctacagcaat ggtggtgctg accagcaagg agacaaagcc     1140
ttccgtggcc gagttctacg ccggcaagtc cgtgtttatc acaggcggca ccggcttcct     1200
gggcaagggtg tttatcgaga agctgctgta ctcttgcca gacatcgaga acatctatat     1260
gctgatccgg gagaagaagg gcctgagcgt gtccgagaga atcaagcagt tcctggacga     1320
tcccctgttt acacggctga aggacaagag acctgccgat ctggagaaga tcgtgctgat     1380
cccaggcgac atcaccgcac cagatctggg catcaactcc gagaatgaga agatgctgat     1440
cgagaagggtg tccgtgatca tccactctgc cgccaccgtg aagttcaacg agcccctgcc     1500
tacagcctgg aagatcaatg tggagggcac caggatgatg ctggccctga gccggagaat     1560
gaagcgcacg gaggtgttta tccacatctc cacagcctac accaacacaa atcgggaggt     1620
ggtggacgag atcctgtacc cagccccgcg cgacatcgat cagggtgcacc agtatgtgaa     1680
ggacggcatc agcgaggagg ataccgagaa gatcctgaac ggccggccaa atacctacac     1740
attcaccaag gccctgacag agcacctggg ggccgagaac caggcctatg tgccctacat     1800
catcgtgaga ccatccgtgg tggccgcat caaggatgag cccctgaagg gatggctggg     1860
aaactggttc ggagcaacag gactgaccgt gtttacagcc aagggcctga atagagtgat     1920
ctacggccac agtcctata tcgtggacct gatccccgtg gattacgtgg caaacctggg     1980
catcgcagca ggagccaagt ctagcaagtc taccgagctg aagggtgata actgctgttc     2040
ctctagctgt aatcctgtga ccatcggcac actgatgtcc atgttcgccg acgatgccat     2100

```

-continued

caagcagaag tcttacgcca tgectctgcc aggctggtac atctttacaa agtataagtg	2160
gctggtgctg ctgctgacct tectgtttca ggtcatccca gcctacgtga ccgatctgtc	2220
taggcacctg atcggcaaga gccccgcta tatcaagctg cagtctctgg tgaaccagac	2280
caggctectct atcgacttct ttacaaatca cagctggggtc atgaaggccg ataggggtgcg	2340
cgagctgtac gcctctctga gccctgccga caagtatctg ttccccctgcg accctaccga	2400
tatcaattgg acacactaca tccaggatta ttgttggggc gtgcgccact tcttgagaa	2460
gaagtcttat gagtgagcct gaagagc	2487
 <210> SEQ ID NO 38 <211> LENGTH: 1012 <212> TYPE: DNA <213> ORGANISM: artificial sequence <220> FEATURE: <223> OTHER INFORMATION: NcoI-pTAL-AleI (insert into pPV247 creating pPV248) <400> SEQUENCE: 38	
ccatgggtaa gcaggtggct ccgtttgtgt ctttgtgttt ttccccctct ttttggacca	60
tttgtcagca tgttgcgtag gtctgggtgt ttgactgttc aggtgggtga tgacggatgc	120
atcatctgac ggcagagtgg gtacctggca gtggcaggct cgcagacgag gtagagagat	180
tctgaaagga gccattgaca gatggagaat tggatactcc tggatatgtcc tccgtttcca	240
cttttgacgt tggtgacgtg ctctggaacg acttttttct ttttctttaa aacaaaaaaaa	300
agaaagaaaa aaaaaacatt tactactacc agtagtacac ctcaacattg ggtccagaac	360
gtcccaactg catgagtcac tggagtcatt ccgaggtcgc taagggtgctg taaaatacaa	420
cgtcaattga gagagacaca ggccgcagcgc gccgagggag aaacgaggca tttatcttct	480
gaccctcctt tttactcgta atctgtatcc cggaaaccgcg tcgcatccat gttaattaaa	540
tcaaacactta cacttgcttg cttcgtatga tgaagatttc tgactggcaa cccagtcagc	600
agcagattgg ggcagatgta gtaatgaaaa aactgcaag gtgtgacgtt tgagacactc	660
caattggtta gaaagcgaca aagaagacgt cggaaaaata ccgaaaaaat cgagtctttt	720
tctttctgcg tattggggcc ttctgcctcc ttgtccgcc tttccacgt ctttccacac	780
cctcacactc cctgagcact atgatctcat tgcgcaataa gatatacatg cacgtgcatt	840
tggtgagcac gcagaacctt gttgggggaa gatgccctaa ccctaagggc gttccatacg	900
gttcgacaga gtaaccttgc tgtcgattat aacgcatata tagccccccc cttcggaccc	960
tcctttctgat ttctgtttct gtatcaacat tacacacaaa cacacaatgg tg	1012

What is claimed is:

1. A recombinant *Yarrowia lipolytica* capable of producing a mono- or poly-unsaturated C₆-C₂₄ fatty alcohol from an endogenous or exogenous source of saturated C₆-C₂₄ fatty acid, comprising:

55 a heterologous nucleic acid molecule encoding and expressing a *Helicoverpa* fatty acyl desaturase that catalyzes the conversion of a saturated C₆-C₂₄ fatty acyl-CoA to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA,

60 wherein the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA is converted to the corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol, and

wherein the *Helicoverpa* fatty acyl desaturase comprises a transmembrane desaturase sequence motif of

65 HX₃₋₄HX₇₋₄₁(3 non-His)HX₂₋₃(1 non-His)HHX₆₁₋₁₈₉(40 non-His)HX₂₋₃(1 non-His)HH.

2. The recombinant *Yarrowia lipolytica* of claim 1, wherein the heterologous nucleic acid molecule encodes a *Helicoverpa* Z9 or Z11 fatty acyl desaturase.
3. The recombinant *Yarrowia lipolytica* of claim 1, wherein the heterologous nucleic acid molecule encodes a *Helicoverpa zea* fatty acyl desaturase.
4. The recombinant *Yarrowia lipolytica* of claim 1, wherein the heterologous nucleic acid molecule comprises a nucleotide sequence encoding the amino acid sequence encoded by SEQ ID NO: 27.
5. The recombinant *Yarrowia lipolytica* of claim 1, comprising a disruption of the *fao1* gene.
6. The recombinant *Yarrowia lipolytica* of claim 1, comprising a disruption of the *pox1*, *pox2*, *pox3*, *pox4*, *pox5*, and/or *pox6* genes.

209

7. The recombinant *Yarrowia lipolytica* of claim 1, comprising a disruption of the adh1, adh2, adh3, adh4, adh5, adh6, and/or adh7 genes.

8. A method of producing a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA, comprising: cultivating the recombinant *Yarrowia lipolytica* of claim 1 in a culture medium containing a carbon source feedstock, such that a saturated C₆-C₂₄ fatty acyl-CoA is converted to a mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA by the catalytic activity of the desaturase.

9. The method of claim 8, wherein the mono- or poly-unsaturated C₆-C₂₄ fatty acyl-CoA is converted to a corresponding mono- or poly-unsaturated C₆-C₂₄ fatty alcohol.

10. The method of claim 9, wherein the mono- or poly-unsaturated C₆-C₂₄ fatty alcohol is a pheromone or a precursor to a pheromone.

11. The recombinant *Yarrowia lipolytica* of claim 1, wherein the heterologous nucleic acid molecule comprises a nucleotide sequence encoding an amino acid sequence having at least 80% identity to the amino acid sequence encoded by SEQ ID NO: 27.

12. The recombinant *Yarrowia lipolytica* of claim 1, wherein the heterologous nucleic acid molecule comprises the nucleotide sequence of SEQ ID NO: 27.

* * * * *

210

25