

US011124804B2

(12) **United States Patent**
DeRosa et al.

(10) **Patent No.:** **US 11,124,804 B2**
(45) **Date of Patent:** ***Sep. 21, 2021**

(54) **MULTIMERIC CODING NUCLEIC ACID AND USES THEREOF**

(2013.01); *C12P 21/00* (2013.01); *C12Y 114/16001* (2013.01); *C12Y 201/03003* (2013.01); *C12N 2830/50* (2013.01)

(71) Applicant: **Translate Bio, Inc.**, Lexington, MA (US)

(58) **Field of Classification Search**
CPC *C12N 15/85*; *C12N 15/67*; *C07H 21/02*; *A61K 31/7115*; *A61K 31/7125*
See application file for complete search history.

(72) Inventors: **Frank DeRosa**, Lexington, MA (US); **Michael Heartlein**, Lexington, MA (US); **Daniel Crawford**, Lexington, MA (US); **Shrirang Karve**, Lexington, MA (US)

(56) **References Cited**

U.S. PATENT DOCUMENTS

(73) Assignee: **Translate Bio, Inc.**, Lexington, MA (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/540,791**

(22) Filed: **Aug. 14, 2019**

(65) **Prior Publication Data**

US 2020/0172923 A1 Jun. 4, 2020

Related U.S. Application Data

(62) Division of application No. 16/280,772, filed on Feb. 20, 2019, now Pat. No. 10,428,349, which is a division of application No. 15/482,431, filed on Apr. 7, 2017, now Pat. No. 10,266,843.

(60) Provisional application No. 62/320,073, filed on Apr. 8, 2016.

(51) **Int. Cl.**

C07H 21/02 (2006.01)
A61K 31/7115 (2006.01)
A61K 31/7125 (2006.01)
C12N 15/85 (2006.01)
C07K 14/505 (2006.01)
C12N 9/10 (2006.01)
C12N 9/02 (2006.01)
C07K 14/705 (2006.01)
A61K 48/00 (2006.01)
A61K 38/17 (2006.01)
A61K 38/44 (2006.01)
A61K 38/45 (2006.01)
A61K 38/18 (2006.01)
C12N 15/67 (2006.01)
C12P 21/00 (2006.01)
A01K 67/00 (2006.01)

(52) **U.S. Cl.**

CPC *C12N 15/85* (2013.01); *A61K 31/7115* (2013.01); *A61K 31/7125* (2013.01); *A61K 38/177* (2013.01); *A61K 38/1816* (2013.01); *A61K 38/44* (2013.01); *A61K 38/45* (2013.01); *A61K 48/005* (2013.01); *C07H 21/02* (2013.01); *C07K 14/505* (2013.01); *C07K 14/705* (2013.01); *C12N 9/0071* (2013.01); *C12N 9/1018* (2013.01); *C12N 15/67*

5,705,385 A 1/1998 Bally et al.
5,976,567 A 11/1999 Wheeler
5,981,501 A 11/1999 Wheeler et al.
6,489,464 B1 12/2002 Agrawal et al.
6,534,484 B1 3/2003 Wheeler et al.
6,815,432 B2 11/2004 Wheeler et al.
7,422,902 B1 9/2008 Wheeler et al.
7,745,651 B2 6/2010 Heyes et al.
7,799,565 B2 9/2010 MacLachlan et al.
7,803,397 B2 9/2010 Heyes et al.
7,901,708 B2 3/2011 MacLachlan et al.
8,101,741 B2 1/2012 MacLachlan et al.
8,188,263 B2 5/2012 MacLachlan et al.
8,236,943 B2 8/2012 Lee et al.
8,329,070 B2 12/2012 MacLachlan et al.
8,513,403 B2 8/2013 MacLachlan et al.
8,569,256 B2 10/2013 Heyes et al.
8,710,200 B2 4/2014 Schrum et al.
8,822,663 B2 9/2014 Schrum et al.

(Continued)

FOREIGN PATENT DOCUMENTS

CA 2807552 A1 9/2012
EP 1519714 10/2010

(Continued)

OTHER PUBLICATIONS

Chassé et al., "Human recombinant adenovirus used to correct a mouse enzyme deficiency", *Clinical Nutrition*, Feb. 1990, vol. 9, No. 1, p. VIII.

Gong et al., "mRNA-mRNA duplexes that auto-elicited Staufen1-mediated mRNA decay", *Nat Struct Mol Biol.*, Jan. 1, 2013, vol. 20, No. 10, pp. 1214-1220.

Gong et al., "Supplementary Information: mRNA-mRNA duplexes that auto-elicited Staufen1-mediated mRNA decay", *Nat Struct Mol Biol.*, Sep. 22, 2013, vol. 20, No. 10, pp. 1214-1220.

Karikó et al., "Incorporation of Pseudouridine Into mRNA Yields Superior Nonimmunogenic Vector With Increased Translational Capacity and Biological Stability", *Mol Ther.*, Nov. 2008, vol. 16, No. 11, pp. 1833-1840.

(Continued)

Primary Examiner — Quang Nguyen

(74) *Attorney, Agent, or Firm* — Proskauer Rose LLP; Fangli Chen

(57) **ABSTRACT**

The present invention provides, among other things, multimeric coding nucleic acids that exhibit superior stability for in vivo and in vitro use. In some embodiments, a multimeric coding nucleic acid (MCNA) comprises two or more encoding polynucleotides linked via 3' ends such that the multimeric coding nucleic acid compound comprises two or more 5' ends.

20 Claims, 20 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS						
8,883,202	B2	11/2014	Manoharan et al.	2013/0259924 A1	10/2013	Bancel et al.
8,980,864	B2	3/2015	Hoge et al.	2013/0266640 A1	10/2013	De Fougerolles et al.
9,051,567	B2	6/2015	Fitzgerald et al.	2014/0010861 A1	1/2014	Bancel et al.
9,061,059	B2	6/2015	Chakraborty et al.	2014/0044772 A1	2/2014	MacLachlan et al.
9,089,604	B2	7/2015	Chakraborty et al.	2014/0105964 A1	4/2014	Bancel et al.
9,095,552	B2	8/2015	Chakraborty et al.	2014/0105965 A1	4/2014	Bancel et al.
9,107,886	B2	8/2015	Chakraborty et al.	2014/0147432 A1	5/2014	Bancel et al.
9,114,113	B2	8/2015	Chakraborty et al.	2014/0147454 A1	5/2014	Chakraborty et al.
9,181,319	B2	11/2015	Schrum et al.	2014/0148502 A1	5/2014	Bancel et al.
9,186,325	B2	11/2015	Manoharan et al.	2014/0155472 A1	6/2014	Bancel et al.
9,186,372	B2	11/2015	de Fougerolles et al.	2014/0155473 A1	6/2014	Bancel et al.
9,187,748	B2	11/2015	Geisbert et al.	2014/0155474 A1	6/2014	Bancel et al.
9,192,651	B2	11/2015	Chakraborty et al.	2014/0155475 A1	6/2014	Bancel et al.
9,220,755	B2	12/2015	Chakraborty et al.	2014/0171485 A1	6/2014	Bancel et al.
9,220,792	B2	12/2015	Chakraborty et al.	2014/0179756 A1	6/2014	MacLachlan et al.
9,233,141	B2	1/2016	Chakraborty et al.	2014/0179771 A1	6/2014	Bancel et al.
9,254,311	B2	2/2016	Bancel et al.	2014/0186432 A1	7/2014	Bancel et al.
9,295,689	B2	3/2016	de Fougerolles et al.	2014/0193482 A1	7/2014	Bancel et al.
9,301,993	B2	4/2016	Chakraborty et al.	2014/0194494 A1	7/2014	Bancel et al.
9,303,079	B2	4/2016	Chakraborty et al.	2014/0199371 A1	7/2014	Bancel et al.
9,334,328	B2	5/2016	Schrum et al.	2014/0200261 A1	7/2014	Hoge et al.
9,345,780	B2	5/2016	Manoharan et al.	2014/0200262 A1	7/2014	Bancel et al.
9,352,042	B2	5/2016	Heyes et al.	2014/0200263 A1	7/2014	Bancel et al.
9,352,048	B2	5/2016	Manoharan et al.	2014/0200264 A1	7/2014	Bancel et al.
9,364,435	B2	6/2016	Yaworski et al.	2014/0206752 A1	7/2014	Afeyan et al.
9,394,234	B2	7/2016	Chen et al.	2014/0206753 A1	7/2014	Guild et al.
9,404,127	B2	8/2016	Yaworski et al.	2014/0206755 A1	7/2014	Bancel et al.
9,428,535	B2	8/2016	de Fougerolles et al.	2014/0206852 A1	7/2014	Hoge et al.
9,428,751	B2	8/2016	MacDonald et al.	2014/0221465 A1	8/2014	Bancel et al.
9,447,164	B2	9/2016	Schrum et al.	2014/0243399 A1	8/2014	Schrum et al.
9,464,124	B2	10/2016	Bancel et al.	2014/0249208 A1	9/2014	Bancel et al.
9,492,386	B2	11/2016	MacLachlan et al.	2014/0255467 A1	9/2014	Bancel et al.
9,504,651	B2	11/2016	MacLachlan et al.	2014/0255468 A1	9/2014	Bancel et al.
9,504,734	B2	11/2016	Bancel et al.	2014/0275227 A1	9/2014	Hoge et al.
9,518,272	B2	12/2016	Yaworksi et al.	2014/0275229 A1	9/2014	Bancel et al.
9,572,874	B2	2/2017	Fotin-Mleczek et al.	2014/0294937 A1	10/2014	MacLachlan et al.
9,572,896	B2	2/2017	Bancel et al.	2014/0343129 A1	11/2014	de Fougerolles et al.
9,572,897	B2	2/2017	Bancel et al.	2015/0005372 A1	1/2015	Hoge et al.
9,587,003	B2	3/2017	Bancel et al.	2015/0017211 A1	1/2015	de Fougerolles et al.
9,616,084	B2	4/2017	Mutzke	2015/0044277 A1	2/2015	Bancel et al.
9,623,095	B2	4/2017	Kallen et al.	2015/0050354 A1	2/2015	Bouchon et al.
D787,703	S	5/2017	Mayer	2015/0051268 A1	2/2015	Bancel et al.
9,636,301	B2	5/2017	Weber	2015/0056253 A1	2/2015	Bancel et al.
9,655,955	B2	5/2017	Hoerr et al.	2015/0064235 A1	3/2015	Bancel et al.
9,657,295	B2	5/2017	Schrum et al.	2015/0064236 A1	3/2015	Bancel et al.
9,669,089	B2	6/2017	Thess et al.	2015/0064242 A1	3/2015	Heyes et al.
9,670,152	B2	6/2017	Payne et al.	2015/0064725 A1	3/2015	Schrum et al.
9,675,668	B2	6/2017	Bancel et al.	2015/0086614 A1	3/2015	Bancel et al.
9,682,139	B2	6/2017	Manoharan et al.	2015/0111248 A1	4/2015	Bancel et al.
9,683,233	B2	6/2017	Thess	2015/0111945 A1	4/2015	Geisbert et al.
9,687,550	B2	6/2017	Manoharan et al.	2015/0166465 A1	6/2015	Chen et al.
9,688,729	B2	6/2017	Kramps et al.	2015/0190515 A1	7/2015	Manoharan et al.
10,266,843	B2 *	4/2019	DeRosa et al.	2015/0315541 A1	11/2015	Bancel et al.
10,428,349	B2 *	10/2019	DeRosa et al.	2015/0315584 A1	11/2015	MacDonald et al.
2002/0192651	A1	12/2002	Wheeler et al.	2015/0366997 A1	12/2015	Guild et al.
2003/0181410	A1	9/2003	Wheeler et al.	2016/0082092 A1	3/2016	Hoerr et al.
2004/0043923	A1	3/2004	Parma et al.	2016/0089424 A1	3/2016	Hoerr et al.
2004/0142025	A1	7/2004	MacLachlan et al.	2016/0095924 A1	4/2016	Hope et al.
2006/0008910	A1	1/2006	MacLachlan et al.	2016/0114011 A1	4/2016	Bancel et al.
2006/0083780	A1	4/2006	Heyes et al.	2016/0115477 A1	4/2016	MacLachlan et al.
2007/0135372	A1	6/2007	MacLachlan et al.	2016/0115483 A1	4/2016	MacLachlan et al.
2009/0270481	A1	10/2009	MacLachlan et al.	2016/0136236 A1	5/2016	Hoge et al.
2010/0041152	A1	2/2010	Wheeler et al.	2016/0151284 A1	6/2016	Heyes et al.
2011/0244026	A1	10/2011	Guild et al.	2016/0158385 A1	6/2016	Bancel et al.
2011/0311583	A1	12/2011	Manoharan et al.	2016/0193299 A1	7/2016	de Fougerolles et al.
2012/0065252	A1	3/2012	Schrum et al.	2016/0194368 A1	7/2016	Hoge et al.
2012/0128760	A1	5/2012	Manoharan et al.	2016/0194625 A1	7/2016	Hoge et al.
2012/0142756	A1	6/2012	Guild et al.	2016/0199485 A1	7/2016	Manoharan et al.
2012/0202871	A1	8/2012	Heyes et al.	2016/0213785 A1	7/2016	Manoharan et al.
2012/0237975	A1	9/2012	Schrum et al.	2016/0237108 A1	8/2016	Fraley et al.
2012/0251618	A1	10/2012	Schrum et al.	2016/0237134 A1	8/2016	Hoge et al.
2012/0328668	A1	12/2012	MacLachlan et al.	2016/0250354 A1	9/2016	Manoharan et al.
2013/0195967	A1	8/2013	Guild et al.	2016/0251681 A1	9/2016	Yaworski et al.
2013/0237594	A1	9/2013	de Fougerolles et al.	2016/0256567 A1	9/2016	Heyes et al.
2013/0259923	A1	10/2013	Bancel et al.	2016/0256568 A1	9/2016	Heyes et al.
				2016/0256573 A1	9/2016	de Fougerolles et al.
				2016/0264971 A1	9/2016	Geisbert et al.
				2016/0264975 A1	9/2016	Schrum et al.
				2016/0274089 A1	9/2016	Ciufolini et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2016/0304552 A1 10/2016 Roy et al.
 2016/0317647 A1 11/2016 Ciaramella et al.
 2016/0317676 A1 11/2016 Hope et al.
 2016/0331828 A1 11/2016 Ciaramella et al.
 2016/0348099 A1 12/2016 Roy et al.
 2016/0354490 A1 12/2016 Roy et al.
 2016/0354491 A1 12/2016 Roy et al.
 2016/0354492 A1 12/2016 Roy et al.
 2016/0354493 A1 12/2016 Roy et al.
 2016/0367687 A1 12/2016 Manoharan et al.
 2016/0367702 A1 12/2016 Hoge et al.
 2016/0375134 A1 12/2016 Bancel et al.
 2016/0375137 A9 12/2016 Manoharan et al.
 2017/0000858 A1 1/2017 Fotin-Mleczek et al.
 2017/0000870 A1 1/2017 Hoerr et al.
 2017/0000871 A1 1/2017 Probst et al.
 2017/0002060 A1 1/2017 Bolen et al.
 2017/0007702 A1 1/2017 Heyes et al.
 2017/0014496 A1 1/2017 Fotin-Mleczek et al.
 2017/0028059 A1 2/2017 Baumhoff et al.
 2017/0029847 A1 2/2017 Thess
 2017/0042814 A1 2/2017 Yaworski et al.
 2017/0056528 A1 3/2017 De Fougerolles et al.
 2017/0056529 A1 3/2017 Thess et al.
 2017/0065675 A1 3/2017 Bancel et al.
 2017/0065727 A1 3/2017 Fotin-Mleczek et al.
 2017/0114378 A1 4/2017 Wochner et al.
 2017/0128549 A1 5/2017 Fotin-Mileczek et al.
 2017/0136131 A1 5/2017 Roy et al.
 2017/0136132 A1 5/2017 Roy et al.
 2017/0143631 A1 5/2017 Chen et al.
 2017/0143796 A1 5/2017 Schrum et al.
 2017/0151333 A1 6/2017 Heyes et al.
 2017/0157268 A1 6/2017 Ansell et al.
 2017/0166905 A1 6/2017 Eberle et al.
 2017/0173128 A1 6/2017 Hoge et al.
 2017/0175129 A1 6/2017 Roy et al.
 2017/0182081 A1 6/2017 Mutzke
 2017/0182150 A1 6/2017 Kallen et al.

FOREIGN PATENT DOCUMENTS

EP 2449106 A1 5/2012
 EP 2338478 B1 6/2013
 EP 2823809 A1 1/2015
 WO WO 2005/026372 A1 3/2005
 WO WO 2005/121348 A1 12/2005
 WO WO 2009/127060 A1 10/2009
 WO WO 2009/127230 A1 10/2009
 WO WO 2010/042877 A1 4/2010
 WO WO 2011/141705 A1 11/2011
 WO WO 2012/019168 A2 2/2012
 WO WO 2012/135805 A2 10/2012
 WO WO 2013/039857 A1 3/2013
 WO WO 2013/039861 A2 3/2013
 WO WO 2013/040429 A1 3/2013
 WO WO 2013/090186 A1 6/2013
 WO WO 2013/101690 A1 7/2013
 WO WO 2013/126803 A1 8/2013
 WO WO 2013/130161 A1 9/2013
 WO WO 2013/151663 A1 10/2013
 WO WO 2013/151664 A1 10/2013
 WO WO 2013/151666 A2 10/2013
 WO WO 2013/151667 A1 10/2013
 WO WO 2013/151668 A1 10/2013

WO WO 2013/151670 A2 10/2013
 WO WO 2013/151671 A1 10/2013
 WO WO 2013/151672 A1 10/2013
 WO WO 2013/151736 A2 10/2013
 WO WO 2014/113089 A1 7/2014
 WO WO 2014/144039 A1 9/2014
 WO WO 2014/144711 A1 9/2014
 WO WO 2014/144767 A1 9/2014
 WO WO 2014/152027 A1 9/2014
 WO WO 2014/152030 A1 9/2014
 WO WO 2014/152031 A1 9/2014
 WO WO 2014/152211 A1 9/2014
 WO WO 2014/152513 A1 9/2014
 WO WO 2014/152540 A1 9/2014
 WO WO 2014/153052 A2 9/2014
 WO WO 2014/158795 A1 10/2014
 WO WO 2014/159813 A1 10/2014
 WO WO 2015/006747 A2 1/2015
 WO WO 2015/023975 A1 2/2015
 WO WO 2015/048744 A2 4/2015
 WO WO 2015/051169 A2 4/2015
 WO WO 2015/051173 A2 4/2015
 WO WO 2015/058069 A1 4/2015
 WO WO 2015/061491 A1 4/2015
 WO WO 2015/085318 A2 6/2015
 WO WO 2015/089511 A2 6/2015
 WO WO 2015/011633 A1 1/2016
 WO WO 2016/054421 A1 4/2016
 WO WO 2016/071857 A1 5/2016
 WO WO 2016/077123 A1 5/2016
 WO WO 2016/077125 A1 5/2016
 WO WO 2016/118724 A1 7/2016
 WO WO 2016/118725 A1 7/2016
 WO WO 2016/154127 A2 9/2016
 WO WO 2016/164762 A1 10/2016
 WO WO 2016/183366 A2 11/2016
 WO WO 2016/197132 A1 12/2016
 WO WO 2016/197133 A1 12/2016
 WO WO 2016/201377 A1 12/2016
 WO WO 2017/019891 A2 2/2017
 WO WO 2017/049074 A1 3/2017
 WO WO 2017/049275 A2 3/2017
 WO WO 2017/049286 A1 3/2017
 WO WO 2017/102010 A1 6/2017
 WO WO 2017/103088 A1 6/2017
 WO WO 2017/108087 A1 6/2017
 WO WO 2017/109134 A1 6/2017
 WO WO 2017/109161 A1 6/2017

OTHER PUBLICATIONS

Sumida et al., "In Vitro Selection of Fab Fragments by mRNA Display and Gene-Linking Emulsion PCR", *Journal of Nucleic Acids*, Jan. 1, 2012, vol. 249, No. 4968, pp. 505-509.
 Vorobjev et al., "Nuclease resistance and RNase H sensitivity of oligonucleotides bridged by oligomethylenediol and oligoethylene glycol linkers", *Antisense Nucleic Acid Drug Dev.*, Apr. 2001, vol. 11, No. 2, pp. 77-85.
 Chaix, et al. "3'-3'-Linked Oligonucleotides: Synthesis Oligonucleotides: Synthesis and Stability Studies", *Bioorgani Biorganic & Medicinal Chemistry Letters*, Elsevier, Amsterdam, NL, vol. 6, No. 7, Jan. 1, 1996 (Jan. 1, 1996), pp. 827-832, XP002211794.
 Masuda, et al. "Synthesis, gene-silencing activity and nuclease resistance of 3'-3'-linked double short hairpin RNA." *Bioorganic & Medicinal Chemistry*, vol. 18, No. 23, Dec. 1, 2020 (Dec. 1, 2020), pp. 8277-8283, XP027546685.

* cited by examiner




FIGURE 1




FIGURE 2




FIGURE 3




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14




FIGURE 15




FIGURE 16




FIGURE 17

Serum Phenylalanine Levels in PAH KO Mice 24 Hours After Treatment
With hPAH MCNA LNPs




FIGURE 18




FIGURE 19




FIGURE 20

MULTIMERIC CODING NUCLEIC ACID AND USES THEREOF

RELATED APPLICATIONS

This application is a divisional Application of U.S. Non-Provisional application Ser. No. 16/280,772, filed Feb. 20, 2019, which is a divisional Application of U.S. Non-Provisional application Ser. No. 15/482,431, filed Apr. 7, 2017, which claims priority to U.S. Provisional Application Ser. No. 62/320,073, filed Apr. 8, 2016, the disclosure of which is hereby incorporated by reference.

SEQUENCE LISTING

The present specification makes reference to a Sequence Listing (submitted electronically as a .txt file named "MRT-1237US3_SeqListing_ST25"). The .txt file was generated Aug. 14, 2019 and is 49,326 bytes in size. The entire contents of the Sequence Listing are herein incorporated by reference.

BACKGROUND

Nucleic acid-based technologies are increasingly important for various therapeutic applications including, but not limited to, messenger RNA therapy, gene therapy, and gene editing, to name but a few. Such therapeutic applications typically require administration of exogenous polynucleotides (e.g. DNA or RNA), which is often hampered by the limited stability of such polynucleotides. For example, following their administration to a subject, many polynucleotides may be subject to nuclease (e.g. exonuclease and/or endonuclease) degradation. Nuclease degradation may negatively influence the capability of a polynucleotide to reach a target cell or to be transcribed and/or translated, the result of which is to preclude the exogenous polynucleotide from exerting an intended therapeutic effect.

SUMMARY OF THE INVENTION

The present invention provides, among other things, multimeric coding nucleic acids that exhibit superior stability for in vivo and in vitro use. The present invention also permits increased complexity and efficiency for nucleic acid based therapeutics.

In some aspects, the present invention provides a multimeric coding nucleic acid (MCNA) comprising one or more coding polynucleotides linked to one or more non-coding polynucleotides via a 3' end linkage between two or more of the polynucleotides (coding or non-coding) such that the MCNA compound comprises two or more 5' ends. In some embodiments, one or more of the 5' ends is modified to include a 5' end cap structure. In certain embodiments, one or more of the coding polynucleotides having a 5' end comprises a 5' end cap structure to facilitate translation of the coding polynucleotides. In certain embodiments, one or more of the polynucleotides having a 5' end structure comprises a 5' end cap structure to facilitate stability of the MCNA.

In some aspects, the present invention provides a multimeric coding nucleic acid (MCNA) comprising two or more encoding polynucleotides linked via 3' ends such that the multimeric coding nucleic acid compound comprises two or more 5' ends. In some embodiments, each of the two or more encoding polynucleotides is a synthetic polyribonucleotide. In some embodiments, each of the two or more encoding

polynucleotides is a synthetic polydeoxyribonucleotide. In some embodiments, each of the two or more encoding polynucleotides is a synthetic polydeoxyribonucleotide or a polyribonucleotide. In some embodiments, each of the two or more encoding polynucleotides encodes a protein of interest. In some embodiments, each of the two or more encoding polynucleotides encodes a same protein. In some embodiments, each of the two or more encoding polynucleotides encodes a distinct protein.

In some embodiments, the MCNA compound comprises three or more encoding polynucleotides. In some embodiments, the compound comprises four or more encoding polynucleotides. In some embodiments, the compound comprises five or more encoding polynucleotides.

In some embodiments, one or more of the encoding polynucleotides comprise a 5' untranslated region (5' UTR) and/or a 3' untranslated region (3' UTR). In some embodiments, the one or more of the encoding polynucleotides comprise a 3' UTR. In some embodiments, the 3' UTR is 5-2,000 nucleotides in length. In some embodiments, the 3' UTR comprises a plurality of multi-A segments with spacers in between. In some embodiments, each of the multi-A segments comprises 8-50 consecutive adenosines. In some embodiments, the plurality of multi-A segments range from 1-100. In some embodiments, the spacers are of varying lengths ranging from 5-100. In some embodiments, the spacers comprise DNA, RNA and/or modified bases. In some embodiments, the modified bases are selected from 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (Ψ U), and 1-methyl-pseudouridine. In some embodiments, the 3' UTR comprises a pseudoknot structure. In some embodiments, the 3' UTR is not followed with a polyadenylation (poly-A) tail. In some embodiments, one or more of the encoding polynucleotides comprise a poly-A tail. In some embodiments, the poly-A tail is 25-5,000 nucleotides in length. In some embodiments, the 3' UTR binds to poly-A binding proteins (PABPs). In some embodiments, the 3' UTR comprises a "kissing loop" sequence motif.

In some embodiments, the 3' ends of the two or more encoding polynucleotides are linked via an oligonucleotide bridge comprising a 3'-3' inverted phosphodiester linkage. In some embodiments, the nucleotides comprising the oligonucleotide bridge are selected from the group consisting of 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (Ψ U), and 1-methyl-pseudouridine. In some embodiments, the oligonucleotide bridge comprises at least one covalent link to an active moiety. In some embodiments, the active moiety is a targeting group, peptide, contrast agent, small molecule, protein, DNA and/or RNA. In some embodiments, nucleotides proximal to the 3'-3' inverted linkage are functionalized with one or more tri-antennary GalNac targeting agents.

In some embodiments, the encoding polynucleotides comprise one or more modified nucleotides. In some embodiments, the modified nucleotides are selected from the group consisting of 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (Ψ U), and 1-methyl-pseudouridine. In some embodiments, the modified nucleotides substitute 1-100% of corresponding native bases. In some embodiments, the at least 25% of uridines are

replaced with 2-thiouridines. In some embodiments, 100% of cytidines are replaced with 5-methylcytidines. In some embodiments, the modified nucleotides are further modified with a 4'-thio substitution on the ribose ring. In some embodiments, the native nucleotides are modified with a 4'-thio substitution on the ribose ring.

In some embodiments, one or more encoding polynucleotides in the MCNA comprise a polynucleotide portion that encodes a therapeutic protein. In some embodiments, one or more encoding polynucleotides in the MCNA comprise a polynucleotide portion that encodes an enzyme, a receptor, a ligand, a light chain or heavy chain of an antibody, a nuclease, or a DNA-binding protein. In certain embodiments, one or more encoding polynucleotides in the MCNA comprise a polynucleotide portion that encodes a nuclease.

In some embodiments, the two or more encoding polynucleotides in the MCNA each comprise a polynucleotide portion that encodes a therapeutic protein. In some embodiments, the two or more encoding polynucleotides in the MCNA each comprise a polynucleotide portion that encodes an enzyme, a receptor, a ligand, a light chain or heavy chain of an antibody, a nuclease, and/or a DNA-binding protein. In some embodiments, the two or more encoding polynucleotides in the MCNA each comprise a polynucleotide portion that encodes a nuclease.

In some embodiments, a first encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a first protein and a second encoding polynucleotide in the MCNA comprising a polynucleotide portion that encodes a second protein that is the same protein as the first protein. In some embodiments, a first encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a first protein and a second encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a second protein that is distinct from the first protein. In certain embodiments, a first encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a first protein in a class of an enzyme, a receptor, a ligand, a light chain or heavy chain of an antibody, a nuclease, or a DNA-binding protein, and a second encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a second protein that is distinct from the first protein but in the same class as the first protein. In certain embodiments, a first encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a first protein in a class of an enzyme, a receptor, a ligand, a light chain or heavy chain of an antibody, a nuclease, or a DNA-binding protein, and a second encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a second protein that is distinct from the first protein and in a different class from the first protein. In certain embodiments, a first encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a light chain of an antibody and a second encoding polynucleotide in the MCNA comprises a polynucleotide portion that encodes a heavy chain in the antibody.

In some aspects, the present invention provides a multi-meric nucleic acid (MNA) comprising two or more polynucleotides linked via at least one 3' end linkage between two or more of the polynucleotides such that the MNA compound comprises two or more 5' ends. In some embodiments, one or more of the 5' ends is modified to facilitate stability of the MNA. In certain embodiments, the two or more polynucleotides linked via the at least one 3' end linkage each are non-coding nucleotides.

In some aspects, the present invention provides a composition comprising the MCNA as described above, encap-

sulated or complexed with a delivery vehicle. In some embodiments, the delivery vehicle is selected from the group consisting of liposomes, lipid nanoparticles, solid-lipid nanoparticles, polymers, viruses, sol-gels, and nano-gels.

In some aspects, the present invention provides methods of delivering MCNA for in vivo protein production, comprising administering the MCNA as described above to a subject in need of delivery. In some embodiments, the MCNA is administered via a route of delivery selected from the group consisting of intravenous delivery, subcutaneous delivery, oral delivery, subdermal delivery, ocular delivery, intratracheal injection pulmonary delivery (e.g. nebulization), intramuscular delivery, intrathecal delivery, or intrarticular delivery.

It is to be understood that all embodiments as described above are applicable to all aspects of the present invention.

BRIEF DESCRIPTION OF THE DRAWING

The drawings are for illustration purposes only, not for limitation.

FIG. 1 shows an exemplary MCNA comprising two RNA species linked via a 3'-3' inverted RNA nucleotide dimer.

FIG. 2 shows an exemplary MCNA comprising two RNA species linked via a 3'-3' inverted RNA nucleotide dimer wherein the MCNA is functionalized with a tri-antennary GalNac targeting agent.

FIG. 3 shows an exemplary MCNA comprising two RNA species linked via a 3'-3' inverted RNA nucleotide dimer wherein the MCNA is functionalized with two tri-antennary GalNac targeting agent.

FIG. 4 shows a general scheme for synthesis of MCNA.

FIG. 5 shows exemplary results of synthesized EPO MCNA detected via gel electrophoresis. Constructs were synthesized under the following conditions: RNA Ligase 1 (A); RNA Ligase 1+10% PEG (B); and RNA Ligase 2 (C).

FIG. 6 shows exemplary results of synthesized EPO MCNA detected via gel electrophoresis. Lane 1 show capped EPO RNA with no tail. Lane 2 shows an EPO MCNA mixture with no DNase treatment. Lane 3 shows an EPO MCNA mixture treated with DNase.

FIG. 7 shows an exemplary graph of the level of hEPO protein secreted after transfection of HEK293T cells with synthetic constructs comprising untailed EPO mRNA or MCNA comprising hEPO mRNA (1 microgram per construct).

FIG. 8 shows exemplary results of synthesized EPO MCNA detected via gel electrophoresis. Lane 1 contains an RNA Ladder, Lane 2 contains a ligation product for EPO MCNA that was not purified, Lane 3 contains purified unreacted/partially reacted product and Lane 4 contains purified EPO MCNA ligation product.

FIG. 9 shows an exemplary graph of the level of hEPO protein secreted after transfection of HEK293T cells with synthetic constructs comprising untailed EPO mRNA or purified MCNA comprising hEPO mRNA (250 nanogram per construct).

FIG. 10 shows an exemplary graph of the level of hOTC protein activity measured in cell lysate after transfection of HEK293T cells with synthetic constructs comprising untailed hOTC mRNA (hOTC monomer) or MCNA comprising hOTC mRNA.

FIG. 11 shows an exemplary graph of the level of hPAH protein produced after transfection of HEK293T cells with synthetic constructs comprising untailed hPAH mRNA (hPAH monomer) or MCNA comprising hPAH mRNA.

FIG. 12 shows an exemplary Western blot demonstrating hCFTR protein production after transfection of HEK293T cells with synthetic constructs comprising untailed hCFTR mRNA (hCFTR monomer) or MCNA comprising hCFTR mRNA.

FIG. 13 shows an exemplary graph of citrulline production measured in livers of mice after treatment with hOTC MCNA encapsulated in lipid nanoparticles.

FIG. 14 shows an exemplary Western blot demonstrating hOTC production detected in livers of mice after treatment with hOTC MCNA or hOTC monomers encapsulated in lipid nanoparticles.

FIG. 15 shows an exemplary graph of citrulline production measured in livers of mice after treatment with hOTC mRNA encapsulated in lipid nanoparticles.

FIG. 16 shows an exemplary graph comparing citrulline production 1 week after administration as a percentage of citrulline production 24 hours after administration in mice treated with hOTC mRNA or hOTC MCNA encapsulated in lipid nanoparticles.

FIG. 17 shows an exemplary graph of hPAH protein detected in livers of PAH knock-out (KO) mice 24 hours after they were administered either hPAH MCNA or hPAH monomers encapsulated in lipid nanoparticles.

FIG. 18 shows an exemplary graph of serum phenylalanine levels in PAH knock-out (KO) mice 24 hours after they were administered either hPAH MCNA or hPAH monomers encapsulated in lipid nanoparticles.

FIG. 19 shows an exemplary graph of hEPO protein detected in the serum of wild-type mice 24 hours after they were administered either hEPO MCNA or hEPO monomers encapsulated in lipid nanoparticles.

FIG. 20 shows exemplary immunohistochemical detection of human Cystic Fibrosis Transmembrane Conductance Regulator (hCFTR) protein in CFTR KO mouse lung 24 hours and 7 days after treatment with hCFTR MCNA encapsulated in lipid nanoparticles via aerosolization.

DEFINITIONS

In order for the present invention to be more readily understood, certain terms are first defined below. Additional definitions for the following terms and other terms are set forth throughout the specification. The publications and other reference materials referenced herein to describe the background of the invention and to provide additional detail regarding its practice are hereby incorporated by reference.

Amino acid: As used herein, the term “amino acid,” in its broadest sense, refers to any compound and/or substance that can be incorporated into a polypeptide chain. In some embodiments, an amino acid has the general structure $H_2N-C(H)(R)-COOH$. In some embodiments, an amino acid is a naturally occurring amino acid. In some embodiments, an amino acid is a synthetic amino acid; in some embodiments, an amino acid is a d-amino acid; in some embodiments, an amino acid is an l-amino acid. “Standard amino acid” refers to any of the twenty standard l-amino acids commonly found in naturally occurring peptides. “Nonstandard amino acid” refers to any amino acid, other than the standard amino acids, regardless of whether it is prepared synthetically or obtained from a natural source. As used herein, “synthetic amino acid” encompasses chemically modified amino acids, including but not limited to salts, amino acid derivatives (such as amides), and/or substitutions. Amino acids, including carboxy- and/or amino-terminal amino acids in peptides, can be modified by methylation, amidation, acetylation, protecting groups, and/or

substitution with other chemical groups that can change the peptide’s circulating half-life without adversely affecting their activity. Amino acids may participate in a disulfide bond. Amino acids may comprise one or posttranslational modifications, such as association with one or more chemical entities (e.g., methyl groups, acetate groups, acetyl groups, phosphate groups, formyl moieties, isoprenoid groups, sulfate groups, polyethylene glycol moieties, lipid moieties, carbohydrate moieties, biotin moieties, etc.). The term “amino acid” is used interchangeably with “amino acid residue,” and may refer to a free amino acid and/or to an amino acid residue of a peptide. It will be apparent from the context in which the term is used whether it refers to a free amino acid or a residue of a peptide.

Animal: As used herein, the term “animal” refers to any member of the animal kingdom. In some embodiments, “animal” refers to humans, at any stage of development. In some embodiments, “animal” refers to non-human animals, at any stage of development. In certain embodiments, the non-human animal is a mammal (e.g., a rodent, a mouse, a rat, a rabbit, a monkey, a dog, a cat, a sheep, cattle, a primate, and/or a pig). In some embodiments, animals include, but are not limited to, mammals, birds, reptiles, amphibians, fish, insects, and/or worms. In some embodiments, an animal may be a transgenic animal, genetically-engineered animal, and/or a clone.

Approximately or about: As used herein, the term “approximately” or “about,” as applied to one or more values of interest, refers to a value that is similar to a stated reference value. In certain embodiments, the term “approximately” or “about” refers to a range of values that fall within 25%, 20%, 19%, 18%, 17%, 16%, 15%, 14%, 13%, 12%, 11%, 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, or less in either direction (greater than or less than) of the stated reference value unless otherwise stated or otherwise evident from the context (except where such number would exceed 100% of a possible value).

Biologically active: As used herein, the term “biologically active” refers to a characteristic of any agent that has activity in a biological system, and particularly in an organism. For instance, an agent that, when administered to an organism, has a biological effect on that organism, is considered to be biologically active.

Delivery: As used herein, the term “delivery” encompasses both local and systemic delivery. For example, delivery of MCNA encompasses situations in which an MCNA is delivered to a target tissue and the encoded protein is expressed and retained within the target tissue (also referred to as “local distribution” or “local delivery”), and situations in which an MCNA is delivered to a target tissue and the encoded protein is expressed and secreted into patient’s circulation system (e.g., serum) and systematically distributed and taken up by other tissues (also referred to as “systemic distribution” or “systemic delivery”).

Expression: As used herein, “expression” of a nucleic acid sequence refers to translation of an MCNA into a polypeptide, assemble multiple polypeptides into an intact protein (e.g., enzyme) and/or post-translational modification of a polypeptide or fully assembled protein (e.g., enzyme). In this application, the terms “expression” and “production,” and grammatical equivalent, are used interchangeably.

Functional: As used herein, a “functional” biological molecule is a biological molecule in a form in which it exhibits a property and/or activity by which it is characterized.

Half-life: As used herein, the term “half-life” is the time required for a quantity such as nucleic acid or protein

concentration or activity to fall to half of its value as measured at the beginning of a time period.

Improve, increase, or reduce: As used herein, the terms “improve,” “increase” or “reduce,” or grammatical equivalents, indicate values that are relative to a baseline measurement, such as a measurement in the same individual prior to initiation of the treatment described herein, or a measurement in a control subject (or multiple control subject) in the absence of the treatment described herein. A “control subject” is a subject afflicted with the same form of disease as the subject being treated, who is about the same age as the subject being treated.

In Vitro: As used herein, the term “in vitro” refers to events that occur in an artificial environment, e.g., in a test tube or reaction vessel, in cell culture, etc., rather than within a multi-cellular organism.

In Vivo: As used herein, the term “in vivo” refers to events that occur within a multi-cellular organism, such as a human and a non-human animal. In the context of cell-based systems, the term may be used to refer to events that occur within a living cell (as opposed to, for example, in vitro systems).

Isolated: As used herein, the term “isolated” refers to a substance and/or entity that has been (1) separated from at least some of the components with which it was associated when initially produced (whether in nature and/or in an experimental setting), and/or (2) produced, prepared, and/or manufactured by the hand of man. Isolated substances and/or entities may be separated from about 10%, about 20%, about 30%, about 40%, about 50%, about 60%, about 70%, about 80%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% of the other components with which they were initially associated. In some embodiments, isolated agents are about 80%, about 85%, about 90%, about 91%, about 92%, about 93%, about 94%, about 95%, about 96%, about 97%, about 98%, about 99%, or more than about 99% pure. As used herein, a substance is “pure” if it is substantially free of other components. As used herein, calculation of percent purity of isolated substances and/or entities should not include excipients (e.g., buffer, solvent, water, etc.).

messenger RNA (mRNA): As used herein, the term “messenger RNA (mRNA)” or “mRNA” refers to a polynucleotide that encodes at least one polypeptide. mRNA as used herein encompasses both modified and unmodified RNA. mRNA may contain one or more coding and non-coding regions. mRNA can be purified from natural sources, produced using recombinant expression systems and optionally purified, chemically synthesized, etc. Where appropriate, e.g., in the case of chemically synthesized molecules, mRNA can comprise nucleoside analogs such as analogs having chemically modified bases or sugars, backbone modifications, etc. An mRNA sequence is presented in the 5' to 3' direction unless otherwise indicated. A typical mRNA molecule has a 5' end and a 3' end. In some embodiments, an mRNA is or comprises natural nucleosides (e.g., adenosine, guanosine, cytidine, uridine); nucleoside analogs (e.g., 2-aminoadenosine, 2-thiothymidine, inosine, pyrrolo-pyrimidine, 3-methyl adenosine, 5-methylcytidine, C-5 propynyl-cytidine, C-5 propynyl-uridine, 2-aminoadenosine, C5-bromouridine, C5-fluorouridine, C5-iodouridine, C5-propynyl-uridine, C5-propynyl-cytidine, C5-methylcytidine, 2-aminoadenosine, 7-deazaadenosine, 7-deazaguanosine, 8-oxoadenosine, 8-oxoguanosine, O(6)-methylguanine, and 2-thiocytidine); chemically modified bases; biologically modified bases (e.g., methylated bases); intercalated bases;

modified sugars (e.g., 2'-fluororibose, ribose, 2'-deoxyribose, arabinose, and hexose); and/or modified phosphate groups (e.g., phosphorothioates and 5'-N-phosphoramidite linkages).

5 Nucleic acid: As used herein, the term “nucleic acid,” in its broadest sense, refers to any compound and/or substance that is or can be incorporated into a polynucleotide chain. In some embodiments, a nucleic acid is a compound and/or substance that is or can be incorporated into a polynucleotide chain via a phosphodiester linkage. In some embodiments, “nucleic acid” refers to individual nucleic acid residues (e.g., nucleotides and/or nucleosides). In some
10 embodiments, “nucleic acid” refers to a polynucleotide chain comprising individual nucleic acid residues. In some
15 embodiments, “nucleic acid” encompasses RNA as well as single and/or double-stranded DNA and/or cDNA.

Patient: As used herein, the term “patient” or “subject” refers to any organism to which a provided composition may be administered, e.g., for experimental, diagnostic, prophylactic, cosmetic, and/or therapeutic purposes. Typical
20 patients include animals (e.g., mammals such as mice, rats, rabbits, non-human primates, and/or humans). In some embodiments, a patient is a human. A human includes pre- and post-natal forms.

25 Pharmaceutically acceptable: The term “pharmaceutically acceptable”, as used herein, refers to substances that, within the scope of sound medical judgment, are suitable for use in contact with the tissues of human beings and animals without excessive toxicity, irritation, allergic response, or
30 other problem or complication, commensurate with a reasonable benefit/risk ratio.

Pharmaceutically acceptable salt: Pharmaceutically acceptable salts are well known in the art. For example, S. M. Berge et al., describes pharmaceutically acceptable salts in detail in *J. Pharmaceutical Sciences* (1977) 66:1-19. Pharmaceutically acceptable salts of the compounds of this invention include those derived from suitable inorganic and organic acids and bases. Examples of pharmaceutically acceptable, nontoxic acid addition salts are salts of an amino group formed with inorganic acids such as hydrochloric acid, hydrobromic acid, phosphoric acid, sulfuric acid and perchloric acid or with organic acids such as acetic acid, oxalic acid, maleic acid, tartaric acid, citric acid, succinic acid or malonic acid or by using other methods used in the art such as ion exchange. Other pharmaceutically acceptable
35 salts include adipate, alginate, ascorbate, aspartate, benzenesulfonate, benzoate, bisulfate, borate, butyrate, camphorate, camphorsulfonate, citrate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, formate, fumarate, glucoheptonate, glycerophosphate, gluconate, hemisulfate, heptanoate, hexanoate, hydroiodide, 2-hydroxy-ethanesulfonate, lactobionate, lactate, laurate, lauryl sulfate, malate, maleate, malonate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, nitrate, oleate, oxalate,
40 palmitate, pamoate, pectinate, persulfate, 3-phenylpropionate, phosphate, picrate, pivalate, propionate, stearate, succinate, sulfate, tartrate, thiocyanate, p-toluenesulfonate, undecanoate, valerate salts, and the like. Salts derived from appropriate bases include alkali metal, alkaline earth metal, ammonium and $N^+(C_{1-4} \text{ alkyl})_4$ salts. Representative alkali or alkaline earth metal salts include sodium, lithium, potassium, calcium, magnesium, and the like. Further pharmaceutically acceptable salts include, when appropriate, nontoxic ammonium, quaternary ammonium, and amine cations
45 formed using counterions such as halide, hydroxide, carboxylate, sulfate, phosphate, nitrate, sulfonate and aryl sulfonate. Further pharmaceutically acceptable salts include
50
55
60
65

salts formed from the quarternization of an amine using an appropriate electrophile, e.g., an alkyl halide, to form a quarternized alkylated amino salt.

Systemic distribution or delivery: As used herein, the terms “systemic distribution,” “systemic delivery,” or gram-
5 matical equivalent, refer to a delivery or distribution mechanism or approach that affect the entire body or an entire organism. Typically, systemic distribution or delivery is accomplished via body’s circulation system, e.g., blood stream. Compared to the definition of “local distribution or
10 delivery.”

Subject: As used herein, the term “subject” refers to a human or any non-human animal (e.g., mouse, rat, rabbit, dog, cat, cattle, swine, sheep, horse or primate). A human includes pre- and post-natal forms. In many embodiments, a
15 subject is a human being. A subject can be a patient, which refers to a human presenting to a medical provider for diagnosis or treatment of a disease. The term “subject” is used herein interchangeably with “individual” or “patient.” A subject can be afflicted with or is susceptible to a disease or disorder but may or may not display symptoms of the disease or disorder.

Substantially: As used herein, the term “substantially” refers to the qualitative condition of exhibiting total or near-total extent or degree of a characteristic or property of
25 interest. One of ordinary skill in the biological arts will understand that biological and chemical phenomena rarely, if ever, go to completion and/or proceed to completeness or achieve or avoid an absolute result. The term “substantially” is therefore used herein to capture the potential lack of
30 completeness inherent in many biological and chemical phenomena.

Target tissues: As used herein, the term “target tissues” refers to any tissue that is affected by a disease to be treated. In some embodiments, target tissues include those tissues
35 that display disease-associated pathology, symptom, or feature.

Therapeutically effective amount: As used herein, the term “therapeutically effective amount” of a therapeutic agent means an amount that is sufficient, when administered
40 to a subject suffering from or susceptible to a disease, disorder, and/or condition, to treat, diagnose, prevent, and/or delay the onset of the symptom(s) of the disease, disorder, and/or condition. It will be appreciated by those of ordinary skill in the art that a therapeutically effective amount is typically administered via a dosing regimen comprising at
45 least one unit dose.

Treating: As used herein, the term “treat,” “treatment,” or “treating” refers to any method used to partially or completely alleviate, ameliorate, relieve, inhibit, prevent, delay
50 onset of, reduce severity of and/or reduce incidence of one or more symptoms or features of a particular disease, disorder, and/or condition. Treatment may be administered to a subject who does not exhibit signs of a disease and/or exhibits only early signs of the disease for the purpose of decreasing the risk of developing pathology associated with the disease.

DETAILED DESCRIPTION

The present invention provides, among other things, methods for synthesizing and compositions comprising mul-
65 timeric coding nucleic acids (MCNA). In particular, the present invention provides MCNA compounds comprising two or more encoding polynucleotides linked via their 3' ends such that the MCNA compound comprises two or more 5' ends and methods of synthesizing the same. In some

embodiments, each of the two or more encoding polynucleotides is a synthetic polyribonucleotide. In some embodiments, each of the two or more encoding polynucleotides is a synthetic polydeoxyribonucleotide. In some embodiments,
5 a synthetic polyribonucleotide or polydeoxyribonucleotide of the invention codes for a polypeptide, protein, enzyme, antibody, or receptor. In some embodiments, the present invention provides a multimeric nucleic acid (MNA) comprising two or more polynucleotides linked via at least one
10 3' end linkage between two or more of the polynucleotides such that the MNA compound comprises two or more 5' ends. In some embodiments, one or more of the 5' ends is modified to facilitate stability of the MNA. In certain embodiments, the two or more polynucleotides linked via
15 the at least one 3' end linkage each are non-coding nucleotides. In some embodiments, a MNA comprises a synthetic polyribonucleotide or polydeoxyribonucleotide that does not code for a polypeptide, protein, enzyme, antibody, or receptor. In some embodiments, MNA comprising a synthetic
20 polyribonucleotide or polydeoxyribonucleotide inhibits gene expression. In some embodiments, a synthetic polyribonucleotide of the invention that inhibits gene expression is a small interfering ribonucleic acid (siRNA), a microRNA (miRNA), or a short hairpin RNA (shRNA).

While the administration of exogenous polynucleotides (e.g. DNA or RNA) represents a meaningful advancement for the treatment of diseases, the administration of such exogenous polynucleotides is often hampered by the limited
25 stability of such polynucleotides, particularly following their in vivo administration. For example, following their administration to a subject, many polynucleotides may be subject to nuclease (e.g. exonuclease and/or endonuclease) degradation. Nuclease degradation may negatively influence the capability of a polynucleotide to reach a target cell or to be
30 transcribed and/or translated, the result of which is to preclude the exogenous polynucleotide from exerting an intended therapeutic effect.

In some embodiments, the MCNA of the present invention exhibit increased in vivo stability compared to a single polynucleotide not linked to another polynucleotide by its 3'
35 end (hereinafter “monomeric polynucleotide”). In some embodiments, the MCNA of the present invention, when delivered in vivo, lead to enhanced protein production compared to a monomeric polynucleotide encoding the same protein. In some embodiments, the MCNA of the present invention, when delivered to a subject, are tolerated better by the subject compared to a corresponding monomeric polynucleotide.

Multimeric Coding Nucleic Acids (MCNA)

In some embodiments, the present invention provides compositions comprising multimeric coding nucleic acids (MCNA) and methods for synthesizing the same. In particular, the present invention provides MCNA compounds comprising two or more encoding polynucleotides linked via
40 their 3' ends such that the MCNA compound comprises two or more 5' ends and methods of synthesizing the same. In some embodiments, each of the two or more encoding polynucleotides is a synthetic polyribonucleotide. In some embodiments, each of the two or more encoding polynucleotides is a synthetic polydeoxyribonucleotide or a polyribonucleotide. In some embodiments, each of the two or more encoding polynucleotides encodes a protein of interest. In
45 some embodiments, each of the two or more encoding polynucleotides encodes a same protein. In some embodiments, each of the two or more encoding polynucleotides

encodes a distinct protein. In some embodiments, each of the two or more encoding polynucleotides encoding a distinct protein are present in equal numbers. In some embodiments, each of the two or more encoding polynucleotides encoding a distinct protein are present in unequal numbers (e.g., 2 5 copies of a polynucleotide encoding protein of interest #1 and 1 copy of a polynucleotide encoding protein of interest #2). In some embodiments, a MCNA compound comprises three or more encoding polynucleotides. In some embodiments, a MCNA compound comprises four or more encoding 10 polynucleotides. In some embodiments, a MCNA compound comprises five or more encoding polynucleotides.

In some embodiments, the present invention provides a multimeric nucleic acid (MNA) comprising two or more polynucleotides linked via at least one 3' end linkage 15 between two or more of the polynucleotides such that the MNA compound comprises two or more 5' ends. In some embodiments, one or more of the 5' ends is modified to facilitate stability of the MNA. In certain embodiments, at least one of the two or more polynucleotides linked via the 20 at least one 3' end linkage is an encoding polynucleotide and at least one of the two or more polynucleotides linked via the at least one 3' end linkage is a non-coding polynucleotide, thereby constituting a multimeric coding nucleic acid (MCNA). In certain embodiments, the encoding polynucleotide 25 encodes a protein of interest and the non-coding polynucleotide inhibits gene expression (e.g., small interfering ribonucleic acid (siRNA), a microRNA (miRNA), or a short hairpin RNA (shRNA)).

In some embodiments, a MCNA compound comprising 30 two or more encoding polynucleotides encodes one or more chains of an antibody or antibody fragment. In some embodiments, the two or more encoding polynucleotides encode a heavy chain and light chain of an antibody. In some embodiments, the antibody is an intact immunoglobulin, (Fab)₂, (Fab')₂, Fab, Fab' or scFv. In some embodiments, the antibody is an IgG. In some embodiments, the antibody is selected from the group consisting of anti-CCl2, anti-lysyl 35 oxidase-like-2 (LOXL2), anti-Flt-1, anti-TNF- α , anti-Interleukin-2R α receptor (CD25), anti-TGF β , anti-B-cell activating factor, anti-alpha-4 integrin, anti-BAGE, anti- β -catenin/m, anti-Bcr-abl, anti-CS, anti-CA125, anti-CAMEL, anti-CAP-1, anti-CASP-8, anti-CD4, anti-CD19, anti-CD20, anti-CD22, anti-CD25, anti-CDC27/m, anti-CD 30, anti-CD33, anti-CD52, anti-CD56, anti-CD80, anti-CDK4/m,

anti-CEA, anti-CT, anti-CTL4, anti-Cyp-B, anti-DAM, anti-EGFR, anti-ErbB3, anti-ELF2M, anti-EMMPRIN, anti-Ep-Cam, anti-ETV6-AML1, anti-HER2, anti-G250, anti-GAGE, anti-GnT-V, anti-Gp100, anti-HAGE, anti-HER-2/ 5 neu, anti-HLA-A*0201-R1701, anti-IGF-1R, anti-IL-2R, anti-IL-S, anti-MC1R, anti-myosin/m, anti-MUC1, anti-MUM-1, -2, -3, anti-proteinase-3, anti-p190 minor bcr-abl, anti-Pml/RAR α , anti-PRAMS, anti-PSA, anti-PSM, anti-PSMA, anti-RAGE, anti-RANKL, anti-RU1 or RU2, anti-SAGE, anti-SART-1 or anti-SART-3, anti-survivin, anti-TEL/AML1, anti-TPI/m, anti-TRP-1, anti-TRP-2, anti-TRP-2/INT2, and anti-VEGF or anti-VEGF receptor.

In some embodiments, a MCNA compound comprising two or more encoding polynucleotides encodes one or more 15 nucleases. In some embodiments, each of the one or more nucleases is selected from the group comprising Cas9, zinc-finger nucleases (ZFN), TALEN, homing endonucleases, homing meganucleases, and combinations thereof. Exemplary nucleases include Afu Uracil-DNA Glycosylase (UDG), Tma Endonuclease III, Tth Endonuclease IV, Antarctic Thermolabile UDG, APE 1, Cas9 Nuclease NLS (*S. pyogenes*), Cas9 Nuclease (*S. pyogenes*), DNase I, Endonuclease IV, Endonuclease V, Endonuclease VIII, Exonuclease I, Exonuclease III (*E. coli*), Exonuclease T, Exonuclease V 20 (RecBCD), Exonuclease VII, Exonuclease VIII (truncated), Fpg, hAAG, hOGG1, hSMUG1, Lambda Exonuclease, Micrococcal Nuclease, Mung Bean Nuclease, Nuclease BAL-31, RecA γ , RecJ γ , T4 PDG (T4 Endonuclease V), T5 Exonuclease, T7 Endonuclease I, T7 Exonuclease, Thermostable FEN1, Uracil Glycosylase Inhibitor (UGI). Exemplary homing nucleases include I-AabMI, I-AniI, I-CeuI, I-CkaMI, I-CpaMI, I-CreI, I-DmoI, I-GpeMI, I-GpiI, I-GzeI, I-GzeII, I-HjeMI, I-LtrI, I-LtrWI, I-MpeMI, I-MsoI, I-OnuI, I-PanMI, I-SceI, I-SmaMI, I-Vdi141I, PI-SceI, 25 I-CreI (m), I-MsoI (m), I-OnuI (E2), I-AniI/I-OnuI, I-DmoI/I-CreI, I-GpiI/I-OnuI, I-GzeI/I-PanMI, I-LtrI/I-PanMI, I-OnuI/I-LtrI, I-AaeMIP, I-ApaMIP, I-GzeMIIP, I-NcrMIP, I-OsoMIIP, I-OsoMIP, I-PanMIIP, I-PanMIIP, I-ScuMIIP, I-ScuMIIP, and I-ScuMIVP.

In some embodiments, a MCNA compound comprises 30 two more polynucleotides that include one, two, or more encoding polynucleotides, wherein each encoding polynucleotide comprises a polynucleotide portion that is an mRNA transcript for a gene and/or for a protein selected from Table 1, Table 2, Table 3, Table 4, Table 5 or Table 6.

TABLE 1

DISEASE/DISORDERS	GENE(S)
Neoplasia	PTEN; ATM; ATR; EGFR; ERBB2; ERBB3; ERBB4; Notch1; Notch2; Notch3; Notch4; AKT; AKT2; AKT3; HIF; H1Fla; HIF3a; Met; HRG; Bcl2; PPARalpha; PPAR gamma; WT1 (Wilms Tumor); FGF Receptor Family members (5 members: 1, 2, 3, 4, 5); CDKN2a; APC; RB (retinoblastoma); MEN1; VHL; BRCA1; BRCA2; AR (Androgen Receptor); TSG101; IGF; IGF Receptor; Igf1 (4 variants); Igf2 (3 variants); Igf1 Receptor; Igf2 Receptor; Bax; Bcl2; caspases family (9 members: 1, 2, 3, 4, 6, 7, 8, 9, 12); Kras; Apc
Age-related Macular Degeneration	Aber; Ccl2; Cc2; cp (ceruloplasmin); Timp3; cathepsinD; Vldlr; Ccr2
Schizophrenia Disorders	Neuregulin1 (Nrg1); Erb4 (receptor for Neureguline); Complexin1 (Cplx1); Tph1 Tryptophan hydroxylase; Tph2 Tryptophan hydroxylase 2; Neurexin 1; GSK3; GSK3a; GSK3b; 5-HTT (Slc6a4); COMT; DRD (Drd1a); SLC6A3; DAOA; DTNBP1; Dao (Dao1)
Trinucleotide Repeat Disorders	HTT (Huntington's Dx); SBMA/SMAXI/AR (Kennedy's Dx); FXN/X25 (Friedrich's Ataxia); ATX3 (Machado-Joseph's Dx); ATXN1 and ATXN2 (spinocerebellar ataxias); DMPK (myotonic dystrophy); Atrophin-1 and

TABLE 1-continued

DISEASE/DISORDERS	GENE(S)
	Atn1(DRPLA Dx); CBP (Creb-BP-global instability); VLDLR (Alzheimer's); Atxn7; Atxn10
Fragile X Syndrome	FMR2; FXR1; FXR2; mGLUR5
Secretase Related Disorders	APH-1 (alpha and beta); Presenilin (Psen1); nicastrin (Ncstn); PEN-2
Others	Nos1; Parp1; Nat1; Nat2
Prion-related Disorders	Prp
ALS	SOD1; ALS2; STEX; FUS; TARD BP; VEGF (VEGF-a; VEGF-b; VEGF-c)
Drug Addiction	Prkce (alcohol); Drd2; Drd4; ABAT (alcohol); GRIA2; Grm5; Grin1; Htr1b; Grin2a; Drd3; Pdyn; Gria1 (alcohol)
Autism	Mecp2; BZRAP1; MDGA2; Sema5A; Neurexin 1; Fragile X (FMR2 (AFF2); FXR1; FXR2; Mglur5)
Alzheimer's Disease	E1; CHIP; UCH; UBB; Tau; LRP; PICALM; Clusterin; PS1; SORL1; CR1; Vld1r; Uba1; Uba3; CHIP28 (Aqp1, Aquaporin 1); Uchl1; Uchl3; APP
Inflammation	IL-10; IL-1 (IL-1a; IL-1b); IL-13; IL-17 (IL-17a (CTLA8); IL-17b; IL-17c; IL-17d; IL-171); 11-23; Cx3crl; ptpn22; TNFa; NOD2/CARD15 for IBD; IL-6; IL-12 (IL-12a; IL-12b); CTLA4; Cx3cell
Parkinson's Disease	x-Synuclein; DJ-1; LRRK2; Parkin; PINK1

TABLE 2

CELLULAR FUNCTION	GENES
Blood and coagulation diseases and disorders	Anemia (CRAN1, CDA1, RPS19, DBA, PKLR, PK1, NT5C3, UMPH1, PSNI, RHAG, RH50A, NRAMP2, SPTB, ALAS2, ANH1, ASB, ABCB7, ABC7, ASAT); Bare lymphocyte syndrome (TAPBP, TPSN, TAP2, ABCB3, PSF2, RING11, MHC2TA, C2TA, RFX5, RFXAP, RFX5), Bleeding disorders (TBXA2R, P2RX1, P2X1); Factor Hand factor H-like 1 (HF1, CFH, HUS); Factor V and Factor VIII (MCFD2); Factor VII deficiency (F7); Factor X deficiency (FIO); Factor XI deficiency (F11); Factor XII deficiency (F12, HAF); Factor XIIIa deficiency (F13AI, F13A); Factor XIIIb deficiency (F13B); Fanconi anemia (FANCA, FACA, FA1, FA, FAA, FAAP95, FAAP90, FLJ34064, FANCB, FANCC, FACC, BRCA2, FANCDI, FANCD2, FANCD, FACD, FAD, FANCE, FACE, FANCF, XRCC9, FANCG, BR1PI, BACH1, FANCI, PHF9, FANCL, FANCM, KIAA1596); Hemophagocytic lymphohistiocytosis disorders (PRF1, HPLH2, UNC13D, MUNC13-4, HPLH3, HLH3, FHL3); Hemophilia A (F8, FSC, HEMA); Hemophilia B (F9, HEMB), Hemorrhagic disorders (PI, ATT, F5); Leukocyte deficiencies and disorders (ITGB2, CD18, LCAMB, LAD, EIF2B1, EIF2BA, EIF2B2, EIF2B3, EIF2B5, LVWM, CACH, CLE, EIF2B4); Sickle cell anemia (HBB); Thalassemia (HBA2, HBB, HBD, LCRB, HBA1).
Cell dysregulation and oncology diseases and disorders	B-cell non-Hodgkin lymphoma (BCL7A, BCL7); Leukemia (TALI, TCL5, SCL, TAL2, FLT3, NBS1, NBS, ZNFN1AI, 1KI, LYF1, HOXD4, HOX4B, BCR, CML, PHL, ALL, ARNT, KRAS2, RASK2, GMPS, AFIO, ARHGEF12, LARG, KIAA0382, CALM, CLTH, CEBPA, CEBP, CHIC2, BTL, FLT3, KIT, PBT, LPP, NPML, NUP214, D9S46E, CAN, CAIN, RUNXI, CBFA2, AML1, WHSC1LI, NSD3, FLT3, AF1Q, NPM1, NUMA1, ZNF145, PLZF, PML, MYL, STAT5B, AF1Q, CALM, CLTH, ARL11, ARLTS1, P2RX7, P2X7, BCR, CML, PHL, ALL, GRAF, NF1, VRNF, WSS, NFNS, PTPNII, PTP2C, SHP2, NS1, BCL2, CCND1, PRAD1, BCL1, TCRA, GATA1, GF1, ERYF1, NFE1, ABLI, NQO1, DIA4, NMOR1, NUP214, D9S46E, CAN, CAIN).
Inflammation and immune related diseases and disorders	AIDS (KIR3DL1, NKAT3, NKB1, AMB11, K1R3DS1, IFNG, CXCL12, SD F1); Autoimmune lymphoproliferative syndrome (TNFRSF6, APT1, FAS, CD95, ALPS1A); Combined immunodeficiency, (IL2RG, SCIDX1, SCIDX, IMD4); HN-1 (CCL5, SCYA5, D17S136E, TCP228), HIV susceptibility or infection (IL10, CSIF, CMKBR2, CCR2, CMKBR5, CCCR5 (CCR5)); Immunodeficiencies (CD3E, CD3G, AICDA, AID, HIGM2, TNFRSF5, CD40, UNG, DGU, HIGM4, TNFSFS, CD40LG, HIGM1, IGM, FOXP3, IPEX, AIID, XPID, PIDX, TNFRSF14B, TAC1; Inflammation (IL-10, IL-1 (IL-1a, IL-1b), IL-13, IL-17 (IL-17a (CTLA8), IL-17b, IL-17c, IL-17d, IL-171), 11-23, Cx3crl, ptpn22, TNFa, NOD2/CARD15 for IBD, IL-6, IL-12 (IL-12a, IL-12b), CTLA4, Cx3cII); Severe combined immunodeficiencies (SCIDs)(JAK3, JAKL, DCLREIC, ARTEMIS, SCIDA, RAG1, RAG2, ADA, PTPRC, CD45, LCA, IL7R, CD3D, T3D, IL2RG, SCIDXI, SCIDX, IMD4).

TABLE 2-continued

CELLULAR FUNCTION	GENES
Metabolic, liver, kidney and protein diseases and disorders	Amyloid neuropathy (TTR, PALB); Amyloidosis (APOA1, APP, AAA, CVAP, AD1, GSN, FGA, LYZ, TTR, PALB); Cirrhosis (KRT18, KRT8, CIRH1A, NAIC, TEX292, KIAA1988); Cystic fibrosis (CFTR, ABCC7, CF, MRP7); Glycogen storage diseases (SLC2A2, GLUT2, G6PC, G6PT, G6PT1, GAA, LAMP2, LAMPB, AGL, GDE, GBE1, GYS2, PYGL, PFKM); Hepatic adenoma, 142330 (TCF1, HNF1A, MODY3); Hepatic failure, early onset, and neurologic disorder (SCOD1, SCO1); Hepatic lipase deficiency (LIPC); Hepatoblastoma, cancer and carcinomas (CTNNB1, PDGFRL, PDGRL, PRLTS, AX1NI, AXIN, CTNNB1, TP53, P53, LFS1, IGF2R, MPRI, MET, CASP8, MCH5; Medullary cystic kidney disease (UMOD, HNFJ, FJHN, MCKD2, ADMCKD2); Phenylketonuria (PAH, PKU1, QDPR, DHPR, PTS); Polycystic kidney and hepatic disease (FCYT, PKHD1, ARPKD, PKD1, PKD2, PKD4, PKDTS, PRKCSH, G19P1, PCLD, SEC63).
Muscular/skeletal diseases and disorders	Becker muscular dystrophy (DMD, BMD, MYF6), Duchenne Muscular Dystrophy (DMD, BMD); Emery-Dreifuss muscular dystrophy (LMNA, LMN1, EMD2, FPLD, CMDIA, HGPS, LGMDIB, LMNA, LMNI, EMD2, FPLD, CMD1A); Facioscapulohumeral muscular dystrophy (FSHMD1A, FSHD1A); Muscular dystrophy (FKRP, MDC1C, LGMD2I, LAMA2, LAMM, LARGE, KIAA0609, MDC1D, FCMD, TTID, MYOT, CAPN3, CANP3, DYSF, LGMD2B, SGCG, LGMD2C, DMDA1, SCG3, SGCA, ADL, DAG2, LGMD2D, DMDA2, SGCB, LGMD2E, SGCD, SGD, LGMD2F, CMD1L, TCAP, LGMD2G, CMD1N, TRIM32, HT2A, LGMD2H, FKRP, MDC1C, LGMD21, TTN, CMD1G, TMD, LGMD2J, POMT1, CAV3, LGMD1C, SEPN1, SELN, RSMD1, PLEC1, PLTN, EBS1); Osteopetrosis (LRP5, BMND1, LRP7, LR3, OPPG, VBCH2, CLCN7, CLC7, OPTA2, OSTMI, GL, TCIRG1, TIRC7, OC116, OPTB1); Muscular atrophy (VAPB, VAPC, ALS8, SMN1, SMA1, SMA2, SMA3, SMA4, BSCL2, SPG17, GARS, SMAD1, CMT2D, HEXB, IGHMBP2, SMUBP2, CATF1, SMARD1).
Neurological and neuronal diseases and disorders	ALS (SOD1, ALS2, STEX, FUS, TARDBP, VEGF (VEGF-a, VEGF-b, VEGF-c); Alzheimer disease (APP, AAA, CVAP, AD1, APOE, AD2, PSEN2, AD4, STM2, APBB2, FE65LI, NOS3, PLA2, URK, ACE, DCPI, ACEI, MPO, PAC1PI, PAXIPIL, PTIP, A2M, BLMH, BMH, PSEN1, AD3); Autism (MeCP2, BZRAP1, MDGA2, Sema5A, Neurexin 1, GLO1, MECP2, RTT, PPMX, MRX16, MRX79, NLGN3, NLGN4, KIAA1260, AUTSX2); Fragile X Syndrome (FMR2, FXR1, FXR2, mGLUR5); Huntington's disease and disease like disorders (HD, IT15, PRNP, PRIP, JPH3, JP3, HDL2, TBP, SCA17); Parkinson disease (NR4A2, NURR1, NOT, TINUR, SNCAIP, TBP, SCA17, SNCA, NACP, PARK1, PARK4, DJ1, PARK7, LRRK2, PARK8, PINK1, PARK6, UCHL1, PARK5, SNCA, NACP, PARK1, PARK4, PRKN, PARK2, PDJ, DBH, NDUFV2); Rett syndrome (MECP2, RTT, PPMX, MRX16, MRX79, CDKL5, STK9, MECP2, RTT, PPMX, MRX16, MRX79, x-Synuclein, DJ-1); Schizophrenia (Neuregulin1 (Nrg1), Erb4 (receptor for Neuregulin), Complexin1 (Cplx1), Tph1 Tryptophan hydroxylase, Tph2, Tryptophan hydroxylase 2, Neurexin 1, GSK3, GSK3a, GSK3b, 5-HTT (Slc6a4), CONT, DRD (Drd1a), SLC6A β , DAOA, DTNBP1, Dao (Dao1)); Secretase Related Disorders (APH-1 (alpha and beta), Presenilin (Psen1), nicastrin, (Ncstn), PEN-2, Nos1, Parp1, Nat1, Nat2); Trinucleotide Repeat Disorders (HTT (Huntington's Dx), SBMA/SMAX1/AR (Kennedy's Dx), FXN/X25 (Friedrich's Ataxia), ATX3 (Machado-Joseph's Dx), ATXN1 and ATXN2 (spinocerebellar ataxias), DMPK (myotonic dystrophy), Atrophin-1 and Atm1 (DRPLA Dx), CBP (Creb-BP-global instability), VLDLR (Alzheimer's), Atxn7, Atxn10).
Ocular diseases and disorders	Age-related macular degeneration (Aber, Ccl2, Cc2, cp (ceruloplasmin), Timp3, cathepsinD, Vldlr, Ccr2); Cataract (CRYAA, CRYA1, CRYBB2, CRYB2, PITX3, BFSP2, CP49, CP47, CRYAA, CRYAI, PAX6, AN2 MGDA, CRYBA1, CRYB1, CRYGC, CRYG3, CCL, LIM2, MP19, CRYGD, CRYG4, BFSP2, CP49, CP47, HSF4, CTM, HSF4, CTM, MIP, AQPO, CRYAB, CRYA2, CTPP2, CRYBB1, CRYGD, CRYG4, CRYBB2, CRYB2, CRYGC, CRYG3, CCL, CRYAA, CRYA1, GJA8, CX50, CAE1, GJA3, CX46, CZP3, CAE3, CCM1, CAM, KRIT1); Corneal clouding and dystrophy (APOA1, TGFBI, CSD2, CDGG1, CSD, BIGH3, CDG2, TACSTD2, TROP2, M1SI, VSX1, RINX, PPCD, PPD, KTCN, COL8A2, FECD, PPCD2, PIP5K3, CFD); Cornea plana congenital (KERA, CNA2); Glaucoma (MYOC, TIGR, GLCIA, JOAG, GPOA, OPTN, GLC1E, FIP2, HYPL, NRP, CYP1BI, GLC3A, OPA1, NTG, NPG, CYP1BI, GLC3A); Leber congenital amaurosis (CRB1, RP12, CRX, CORD2, CRD, RPGRIP1, LCA6, CORD9, RPE65, RP20, AIPL1, LCA4, GUCY2D, GUC2D, LCA1, CORD6, RDH12, LCA3); Macular dystrophy (ELOVL4, ADMD, STGD2, STGD3, RDS, RP7, PRPH2, PRPH, AVMD, AOFMD, VMD2).
Epilepsy Duchenne muscular dystrophy	NHLRC1, EPM2A, EPM2B DMD, BMD

TABLE 2-continued

CELLULAR FUNCTION	GENES
AIDS	KIR3DL1, NKAT3, NKB1, AMB11, KIR3DS1, IFNG, CDDCL12, SDF1
Alpha 1-Antitrypsin Deficiency	SERPINA1 [serpin peptidase inhibitor, cladeA (alpha-1 antiproteinase, antitrypsin), member 1]; SERPINA2 [serpin peptidase inhibitor, cladeA (alpha-1 antiproteinase, antitrypsin), member 2]; SERPINA3 [serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 3]; SERPINA5 [serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 5]; SERPINA6 [serpin peptidase inhibitor, clade A (alpha-1 antiproteinase, antitrypsin), member 6]; SERPINA7 [serpin peptidase inhibitor, Glade A (alpha-1 antiproteinase, antitrypsin), member 7]; SERPINA6 (serpin peptidase inhibitor, cladeA (alpha-1 antiproteinase, antitrypsin), member 6)

TABLE 3

CELLULAR FUNCTION	GENES
PI3K/AKT Signaling	PRKCE; ITGAM; ITGA5; IRAK1; PRKAA2; EIF2AK2; PTEN; EIF4E; PRKCZ; GRK6; MAPK1; TSC1; PLK1; AKT2; IKBKB; PIK3CA; CDK8; CDKN1B; NFKB2; BCL2; PIK3CB; PPP2R1A; MAPK8; BCL2LI; MAPK3; TSC2; ITGA1; KRAS; EIF4EBP1; RELA; PRKCD; NOS3; PRKAA1; MAPK9; CDK2; PPP2CA; PIM1; ITGB7; YWHAZ; ILK; TP53; RAF1; IKBKG; RELB; DYRK1A; CDKN1A; ITGB1; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; CHUK; PDPK1; PPP2R5C; CTNBN1; MAP2K1; NFKB1; PAK3; ITGB3; CCND1; GSK3A; FRAP1; SFN; ITGA2; TTK; CSNK1A1; BRAF; GSK3B; AKT3; FOXO1; SGK; HSP90AA1; RPS6KB1
ERK/MAPK Signaling	PRKCE; ITGAM; ITGA5; HSPB1; IRAK1; PRKAA2; EIF2AK2; RAC1; RAP1A; TLN1; EIF4E; ELK1; GRK6; MAPK1; RAC2; PLK1; AKT2; PIK3CA; CDK8; CREB1; PRKC1; PTK2; FOS; RPS6KA4; PIK3CB; PPP2R1A; PIK3C3; MAPK8; MAPK3; ITGA1; ETS1; KRAS; MYCN; EIF4EBP1; PPARC; PRKCD; PRKAA1; MAPK9; SRC; CDK2; PPP2CA; PIM1; PIK3C2A; ITGB7; YWHAZ; PPP1CC; KSR1; PXN; RAF1; FYN; DYRK1A; ITGB1; MAP2K2; PAK4; PIK3R1; STAT3; PPP2R5C; MAP2KI; PAK3; ITGB3; ESR1; ITGA2; MYC; TTK; CSNK1A1; CRKL; BRAF; ATF4; PRKCA; SRF; STAT1; SGK
Glucocorticoid Receptor Signaling	RAC1; TAF4B; EP300; SMAD2; TRAF6; PCAF; ELK1; MAPK1; SMAD3; AKT2; IKBKB; NCOR2; UBE21; PIK3CA; CREB1; FOS; HSPA5; NFKB2; BCL2; MAP3K14; STAT5B; PIK3CB; PIK3C3; MAPK8; BCL2L1; MAPK3; TSC22D3; MAPK10; NRIP1; KRAS; MAPK13; RELA; STAT5A; MAPK9; NOS2A; PBX1; NR3C1; PIK3C2A; CDKN1C; TRAF2; SERPINE1; NCOA3; MAPK14; TNF; RAF1; IKBKG; MAP3K7; CREBBP; CDKN1A; MAP2K2; JAK1; IL8; NCOA2; AKT1; JAK2; PIK3R1; CHUK; STAT3; MAP2K1; NFKB1; TGFBR1; ESR1; SMAD4; CEBPB; WN; AR; AKT3; CCL2; MMP1; STAT1; IL6; HSP90AA1
Axonal Guidance Signaling	PRKCE; ITGAM; ROCK1; ITGA5; CXCR4; ADAM12; IGF1; RAC1; RAP1A; EIF4E; PRKCZ; NRP1; NTRK2; ARHGAP7; SMO; ROCK2; MAPK1; PGF; RAC2; PTPN11; GNAS; AKT2; PIK3CA; ERBB2; PRKCI; PTK2; CFL1; GNAQ; PIK3CB; CXCL12; PIK3C3; WNT11; PRKD1; GNB2L1; ABL1; MAPK3; ITGA1; KRAS; RHOA; PRKCD; PIK3C2A; ITGB7; GLI2; PXN; VASP; RAF1; FYN; ITGB1; MAP2K2; PAK4; ADAM17; AKT1; PIK3R1; GLI1; WNT5A; ADAM10; MAP2K1; PAK3; ITGB3; CDC42; VEGFA; ITGA2; EPHA8; CRKL; RND1; GSK3B; AKT3; PRKCA
Ephrin Receptor Signaling	PRKCE; ITGAM; ROCK1; ITGA5; CXCR4; IRAK1; PRKAA2; EIF2AK2; RAC1; RAP1A; GRK6; ROCK2; MAPK1; PGF; RAC2; PTPN11; GNAS; PLK1; AKT2; DOK1; CDK8; CREB1; PTK2; CFL1; GNAQ; MAP3K14; CXCL12; MAPK8; GNB2L1; ABL1; MAPK3; ITGA1; KRAS; RHOA; PRKCD; PRKAA1; MAPK9; SRC; CDK2; PIM1; ITGB7; PXN; RAF1; FYN; DYRK1A; ITGB1;

TABLE 3-continued

CELLULAR FUNCTION	GENES
Actin Cytoskeleton Signaling	MAP2K2; PAK4, AKT1; JAK2; STAT3; ADAM10; MAP2K1; PAK3; ITGB3; CDC42; VEGFA; ITGA2; EPHA8; TTK; CSNK1A1; CRKL; BRAF; PTPN13; ATF4; AKT3; SGK ACTN4; PRKCE; ITGAM; ROCK1; ITGA5; IRAK1; PRKAA2; EIF2AK2; RAC1; INS; ARHGEF7; GRK6; ROCK2; MAPK1; RAC2; PLK1; AKT2; PIK3CA; CDK8; PTK2; CPL1; PIK3CB; MYH9; DIAPH1; PIK3C3; MAPK8; F2R; MAPK3; SLC9A1; ITGA1; KRAS; RHOA; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; ITGB7; PPP1CC; PXN; VIL2; RAF1; GSN; DYRK1A; ITGB1; MAP2K2; PAK4; PIP5K1A; PIK3R1; MAP2K1; PAK3; ITGB3; CDC42; APC; ITGA2; TTK; CSNK1A1; CRKL; BRAF; VAV3; SGK
Huntington's Disease Signaling	PRKCE; IGF1; EP300; RCOR1; PRKCZ; HDAC4; TGM2; MAPK1; CAPNS1; AKT2; EGFR; NCOR2; SP1; CAPN2; PIK3CA; HDAC5; CREB1; PRKCI; HSPA5; REST; GNAQ; PIK3CB; PIK3C3; MAPK8; IGF1R; PRKD1; GNB2L1; BCL2L1; CAPN1; MAPK3; CASP8; HDAC2; HDAC7A; PRKCD; HDAC11; MAPK9; HDAC9; PIK3C2A; HDAC3; TP53; CASP9; CREBBP; AKT1; PIK3R1; PDPK1; CASP1; APAF1; FRAP1; CASP2; JUN; BAX; ATF4; AKT3; PRKCA; CLTC; SGK; HDAC6; CASP3
Apoptosis Signaling	PRKCE; ROCK1; BID; IRAK1; PRKAA2; EIF2AK2; BAK1; BIRC4; GRK6; MAPK1; CAPNS1; PLK1; AKT2; IKBKB; CAPN2; CDK8; FAS; NFKB2; BCL2; MAP3K14; MAPK8; BCL2L1; CAPN1; MAPK3; CASP8; KRAS; RELA; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; TP53; TNF; RAF1; IKBKG; RELB; CASP9; DYRK1A; MAP2K2; CHUK; APAF1; MAP2K1; NFKB1; PAK3; LMNA; CASP2; BIRC2; TTK; CSNK1A1; BRAF; BAX; PRKCA; SGK; CASP3; BIRC3; PARP1
B Cell Receptor Signaling	RAC1; PTEN; LYN; ELK1; MAPK1; RAC2; PTPN11; AKT2; IKBKB; PIK3CA; CREB1; SYK; NFKB2; CAMK2A; MAP3K14; PIK3CB; PIK3C3; MAPK8; BCL2L1; ABL1; MAPK3; ETS1; KRAS; MAPK13; RELA; PTPN6; MAPK9; EGR1; PIK3C2A; BTK; MAPK14; RAF1; IKBKG; RELB; MAP3K7; MAP2K2; AKT1; PIK3R1; CHUK; MAP2K1; NFKB1; CDC42; GSK3A; FRAP1; BCL6; BCL10; JUN; GSK3B; ATF4; AKT3; VAV3; RPS6KB1
Leukocyte Extravasation Signaling	ACTN4; CD44; PRKCE; ITGAM; ROCK1; CXCR4; CYBA; RAC1; RAP1A; PRKCZ; ROCK2; RAC2; PTPN11; MMP14; PIK3CA; PRKCI; PTK2; PIK3CB; CXCL12; PIK3C3; MAPK8; PRKD1; ABL1; MAPK10; CYBB; MAPK13; RHOA; PRKCD; MAPK9; SRC; PIK3C2A; BTK; MAPK14; NOX1; PXN; VIL2; VASP; ITGB1; MAP2K2; CTNND1; PIK3R1; CTNND1; CLDN1; CDC42; F11R; ITK; CRKL; VAV3; CTTN; PRKCA; MMP1; MMP9
Integrin Signaling	ACTN4; ITGAM; ROCK1; ITGA5; RAC1; PTEN; RAP1A; TLN1; ARHGEF7; MAPK1; RAC2; CAPNS1; AKT2; CAPN2; PIK3CA; PTK2; PIK3CB; PIK3C3; MAPK8; CAV1; CAPN1; ABL1; MAPK3; ITGA1; KRAS; RHOA; SRC; PIK3C2A; ITGB7; PPP1CC; ILK; PXN; VASP; RAF1; FYN; ITGB1; MAP2K2; PAK4; AKT1; PIK3R1; TNK2; MAP2K1; PAK3; ITGB3; CDC42; RND3; ITGA2; CRKL; BRAF; GSK3B; AKT3
Acute Phase Response Signaling	IRAK1; SOD2; MYD88; TRAF6; ELK1; MAPK1; PTPN11; AKT2; IKBKB; PIK3CA; FOS; NFKB2; MAP3K14; PIK3CB; MAPK8; RIPK1; MAPK3; IL6ST; KRAS; MAPK13; IL6R; RELA; SOCS1; MAPK9; FTL; NR3C1; TRAF2; SERPINE1; MAPK14; TNF; RAF1; PDK1; IKBKG; RELB; MAP3K7; MAP2K2; AKT1; JAK2; PIK3R1; CHUK; STAT3; MAP2K1; NFKB1; FRAP1; CEBPB; JUN; AKT3; IL1R1; IL6
PTEN Signaling	ITGAM; ITGA5; RAC1; PTEN; PRKCZ; BCL2L11; MAPK1; RAC2; AKT2; EGFR; IKBKB; CBL; PIK3CA; CDKN1B; PTK2; NFKB2; BCL2; PIK3CB; BCL2L1; MAPK3; ITGA1; KRAS; ITGB7; ILK; PDGFRB; INSR; RAF1; IKBKG; CASP9; CDKN1A; ITGB1; MAP2K2; AKT1; PIK3R1; CHUK; PDGFRA; PDPK1; MAP2K1; NFKB1; ITGB3; CDC42; CCND1; GSK3A; ITGA2; GSK3B; AKT3; FOXO1; CASP3; RPS6KB1
p53 Signaling	PTEN; EP300; BBC3; PCAF; FASN; BRCA1; GADD45A; BIRC5; AKT2; PIK3CA; CHEK1; TP53INP1; BCL2; PIK3CB; PIK3C3; MAPK8; THBS1; ATR; BCL2L1; E2F1; PMAIP1; CHEK2; TNFRSF10B; TP73; RB1; HDAC9; CDK2; PIK3C2A; MAPK14; TP53; LRDD; CDKN1A;

TABLE 3-continued

CELLULAR FUNCTION	GENES
Aryl Hydrocarbon Receptor Signaling	HIPK2; AKT1; PIK3R1; RRM2B; APAF1; CTNNB1; SIRT1; CCND1; PRKDC; ATM; SFN; CDKN2A; JUN; SNAI2; GSK3B; BAX; AKT3 HSPB1; EP300; FASN; TGM2; RXRA; MAPK1; NQO1; NCOR2; SP1; ARNT; CDKN1B; FOS; CHEK1; SMARCA4; NEKB2; MAPK8; ALDH1A1; ATR; E2F1; MAPK3; NRIP1; CHEK2; RELA; TP73; GSTP1; RB1; SRC; CDK2; AHR; NFE2L2; NCOA3; TP53; TNF; CDKN1A; NCOA2; APAF1; NFKB1; CCND1; ATM; ESR1; CDKN2A; MYC; JUN; ESR2; BAX; IL6; CYP1B1; HSP90AA1
Xenobiotic Metabolism Signaling	PRKCE; EP300; PRKCZ; RXRA; MAPK1; NQO1; NCOR2; PIK3CA; ARNT; PRKCI; NFKB2; CAMK2A; PIK3CB; PPP2R1A; PIK3C3; MAPK8; PRKD1; ALDH1A1; MAPK3; NRIP1; KRAS; MAPK13; PRKCD; GSTP1; MAPK9; NOS2A; ABCB1; AHR; PPP2CA; FTL; NFE2L2; PIK3C2A; PPARGC1A; MAPK14; TNF; RAF1; CREBBP; MAP2K2; PIK3R1; PPP2R5C; MAP2K1; NFKB1; KEAP1; PRKCA; EIF2AK3; IL6; CYP1B1; HSP90AA1
SAPK/JNK Signaling	PRKCE; IRAK1; PRKAA2; EIF2AK2; RAC1; ELK1; GRK6; MAPK1; GADD45A; RAC2; PLK1; AKT2; PIK3CA; FADD; CDK8; PIK3CB; PIK3C3; MAPK8; RIPK1; GNB2L1; IRS1; MAPK3; MAPK10; DAXX; KRAS; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; TRAF2; TP53; LCK; MAP3K7; DYRK1A; MAP2K2; PIK3R1; MAP2K1; PAK3; CDC42; JUN; TTK; CSNK1A1; CRKL; BRAF; SGK
PPAr/RXR Signaling	PRKAA2; EP300; INS; SMAD2; TRAF6; PPARA; FASN; RXRA; MAPK1; SMAD3; GNAS; IKBKB; NCOR2; ABCA1; GNAQ; NFKB2; MAP3K14; STAT5B; MAPK8; IRS1; MAPK3; KRAS; RELA; PRKAA1; PPARGC1A; NCOA3; MAPK14; INSR; RAF1; IKBKG; RELB; MAP3K7; CREBBP; MAP2K2; JAK2; CHUK; MAP2K1; NFKB1; TGFB1; SMAD4; JUN; IL1R1; PRKCA; IL6; HSP90AA1; ADIPOQ
NF-KB Signaling	IRAK1; EIF2AK2; EP300; INS; MYD88; PRKCZ; TRAF6; TBK1; AKT2; EGFR; IKBKB; PIK3CA; BTRC; NFKB2; MAP3K14; PIK3CB; PIK3C3; MAPK8; RIPK1; HDAC2; KRAS; RELA; PIK3C2A; TRAF2; TLR4; PDGFRB; TNF; INSR; LCK; IKBKG; RELB; MAP3K7; CREBBP; AKT1; PIK3R1; CHUK; PDGFRA; NFKB1; TLR2; BCL10; GSK3B; AKT3; TNFAIP3; IL1R1
Neuregulin Signaling	ERBB4; PRKCE; ITGAM; ITGA5; PTEN; PRKCZ; ELK1; MAPK1; PTPN11; AKT2; EGFR; ERBB2; PRKCI; CDKN1B; STAT5B; PRKD1; MAPK3; ITGA1; KRAS; PRKCD; STAT5A; SRC; ITGB7; RAF1; ITGB1; MAP2K2; ADAM1 7; AKT1; PIK3RI; PDPK1; MAP2K1; ITGB3; EREG; FRAP1; PSEN1; ITGA2; MYC; NRG1; CRKL; AKT3; PRKCA; HSP90AA1; RPS6KB1
Wnt & Beta catenin Signaling	CD44; EP300; LRP6; DVL3; CSNK1E; GJA1; SMO; AKT2; PIN1; CDH1; BTRC; GNAQ; MARK2; PPP2R1A; WNT11; SRC; DKK1; PPP2CA; SOX6; SFRP2; ILK; LEF1; SOX9; TP53; MAP3K7; CREBBP; TCF7L2; AKT1; PPP2R5C; WNT5A; LRP5; CTNNB1; TGFB1; CCND1; GSK3A; DVL1; APC; CDKN2A; MYC; CSNK1A1; GSK3B; AKT3; SOX2
Insulin Receptor Signaling	PTEN; INS; EIF4E; PTPN1; PRKCZ; MAPK1; TSC1; PTPN11; AKT2; CBL; PIK3CA; PRKCI; PIK3CB; PIK3C3; MAPKS; IRS1; MAPK3; TSC2; KRAS; EIF4EBP1; SLC2A4; PIK3C2A; PPP1CC; INSR; RAF1; FYN; MAP2K2; JAK1; AKT1; JAK2; PIK3RI; PDPK1; MAP2K1; GSK3A; FRAP1; CRKL; GSK3B; AKT3; FOXO1; SGK; RPS6KB1
IL-6 Signaling	HSPB1; TRAF6; MAPKAPK2; ELK1; MAPK1; PTPN11; IKBKB; FOS; NFKB2; MAP3K14; MAPKS; MAPK3; MAPK10; IL6ST; KRAS; MAPK13; IL6R; RELA; SOCS1; MAPK9; ABCB1; TRAF2; MAPK14; TNF; RAF1; IKBKG; RELB; MAP3K7; MAP2K2; IL8; JAK2; CHUK; STAT3; MAP2KI; NFKB1; CEBPB; JUN; IL1R1; SRF; IL6
Hepatic Cholestasis	PRKCE; IRAK1; INS; MYDSS; PRKCZ; TRAF6; PPARA; RXRA; IKBKB; PRKCI; NFKB2; MAP3K14; MAPKS; PRKD1; MAPK10; RELA; PRKCD; MAPK9; ABCB1; TRAF2; TLR4; TNF; INSR; IKBKG; RELB; MAP3K7; IL8; CHUK; NR1H2; TJP2; NFKB1; ESR1; REBF1; FGFR4; JUN; IL1R1; PRKCA; IL6

TABLE 3-continued

CELLULAR FUNCTION	GENES
IGF-1 Signaling	IGF1; PRKCZ; ELK1; MAPK1; PTPN11; NEDD4; AKT2; PIK3CA; PRKCI; PTK2; FOS; PIK3CB; PIK3C3; MAPKS; IGF1R; IRS1; MAPK3; IGFBP7; KRAS; PIK3C2A; YWHAZ; PXN; RAF1; CASP9; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; IGFBP2; SFN; JUN; CYR61; AKT3; FOXO1; SRF; CTGF; RPS6KB1
NRF2-mediated Oxidative Stress Response	PRKCE; EP300; SOD2; PRKCZ; MAPK1; SQSTM1; NQO1; PIK3CA; PRKCI; FOS; PIK3CB; PIK3C3; MAPK8; PRKD1; MAPK3; KRAS; PRKCD; GSTP1; MAPK9; FTL; NFE2L2; PIK3C2A; MAPK14; RAF1; MAP3K7; CREBBP; MAP2K2; AKT1; PIK3R1; MAP2K1; PPIB; JUN; KEAP1; GSK3B; ATF4; PRKCA; EIF2AK3; HSP90AA1
Hepatic Fibrosis/Hepatic Stellate Cell Activation	EDN1; IGF1; KDR; FLT1; SMAD2; FGFR1; MET; PGF; SMAD3; EGFR; FAS; CSF1; NFKB2; BCL2; MYH9; IGF1R; IL6R; RELA; TLR4; PDGFRB; TNF; RELB; IL8; PDGFRA; NFKB1; TGFBR1; SMAD4; VEGFA; BAX; IL1R1; CCL2; HGF; MMP1; STAT1; IL6; CTGF; MMP9
PPAR Signaling	EP300; INS; TRAF6; PPARA; RXRA; MAPK1; IKBKB; NCOR2; FOS; NFKB2; MAP3K14; STAT5B; MAPK3; NRIP1; KRAS; PPARG; RELA; STAT5A; TRAF2; PPARGC1A; PDGFRB; TNF; INSR; RAF1; IKBKG; RELB; MAP3K7; CREBBP; MAP2K2; CHUK; PDGFRA; MAP2KI; NFKB1; JUN; IL1R1; HSP90AA1
Fc Epsilon R1 Signaling	PRKCE; RAC1; PRKCZ; LYN; MAPK1; RAC2; PTPN11; AKT2; PIK3CA; SYK; PRKCI; PIK3CB; PIK3C3; MAPK8; PRKD1; MAPK3; MAPK10; KRAS; MAPK13; PRKCD; MAPK9; PIK3C2A; BTK; MAPK14; TNF; RAF1; FYN; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; AKT3; VAV3; PRKCA
G-Protein Coupled Receptor Signaling	PRKCE; RAP1A; RGS16; MAPK1; GNAS; AKT2; IKBKB; PIK3CA; CREB1; GNAQ; NFKB2; CAMK2A; PIK3CB; PIK3C3; MAPK3; KRAS; RELA; SRC; PIK3C2A; RAF1; IKBKG; RELB; FYN; MAP2K2; AKT1; PIK3R1; CHUK; PDPK1; STAT3; MAP2K1; NFKB1; BRAF; ATF4; AKT3; PRKCA
Inositol Phosphate Metabolism	PRKCE; IRAK1; PRKAA2; EIF2AK2; PTEN; GRK6; MAPK1; PLK1; AKT2; PIK3CA; CDK8; PIK3CB; PIK3C3; MAPK8; MAPK3; PRKCD; PRKAA1; MAPK9; CDK2; PIM1; PIK3C2A; DYRK1A; MAP2K2; PIP5K1A; PIK3R1; MAP2K1; PAK3; ATM; TTK; CSNK1A1; BRAF; SGK
PDGF Signaling	EIF2AK2; ELK1; ABL2; MAPK1; PIK3CA; FOS; PIK3CB; PIK3C3; MAPK8; CAV1; ABL1; MAPK3; KRAS; SRC; PIK3C2A; PDGFRB; RAF1; MAP2K2; JAK1; JAK2; PIK3R1; PDGFRA; STAT3; SPHK1; MAP2K1; MYC; JUN; CRKL; PRKCA; SRF; STAT1; SPHK2
VEGF Signaling	ACTN4; ROCK1; KDR; FLT1; ROCK2; MAPK1; PGF; AKT2; PIK3CA; ARNT; PTK2; BCL2; PIK3CB; PIK3C3; BCL2L1; MAPK3; KRAS; HIF1A; NOS3; PIK3C2A; PXN; RAF1; MAP2K2; ELAVL1; AKT1; PIK3R1; MAP2K1; SFN; VEGFA; AKT3; FOXO1; PRKCA
Natural Killer Cell Signaling	PRKCE; RAC1; PRKCZ; MAPK1; RAC2; PTPN11; KIR2DL3; AKT2; PIK3CA; SYK; PRKCI; PIK3CB; PIK3C3; PRKD1; MAPK3; KRAS; PRKCD; PTPN6; PIK3C2A; LCK; RAF1; FYN; MAP2K2; PAK4; AKT1; PIK3R1; MAP2K1; PAK3; AKT3; VAV3; PRKCA
Cell Cycle: G1/S Checkpoint Regulation	HDAC4; SMAD3; SUV39H1; HDAC5; CDKN1B; BTRC; ATR; ABL1; E2F1; HDAC2; HDAC7A; RB1; HDAC11; HDAC9; CDK2; E2F2; HDAC3; TP53; CDKN1A; CCND1; E2F4; ATM; RBL2; SMAD4; CDKN2A; MYC; NRG1; GSK3B; RBL1; HDAC6
T Cell Receptor Signaling	RAC1; ELK1; MAPK1; IKBKB; CBL; PIK3CA; FOS; NFKB2; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; RELA; PIK3C2A; BTK; LCK; RAF1; IKBKG; RELB; FYN; MAP2K2; PIK3R1; CHUK; MAP2K1; NFKB1; ITK; BCL10; JUN; VAV3
Death Receptor Signaling	CRADD; HSPB1; BID; BIRC4; TBK1; IKBKB; FADD; FAS; NFKB2; BCL2; MAP3K14; MAPK8; RIPK1; CASP8; DAXX; TNFRSF10B; RELA; TRAF2; TNF; IKBKG; RELB; CASP9; CHUK; APAF1; NFKB1; CASP2; BIRC2; CASP3; BIRC3
FGF Signaling	RAC1; FGFR1; MET; MAPKAPK2; MAPK1; PTPN11; AKT2; PIK3CA; CREB1; PIK3CB; PIK3C3; MAPK8; MAPK3; MAPK13; PTPN6; PIK3C2A; MAPK14; RAF1; AKT1; PIK3R1; STAT3; MAP2K1; FGFR4; CRKL; ATF4; AKT3; PRKCA; HGF
GN-CSF Signaling	LYN; ELK1; MAPK1; PTPN11; AKT2; PIK3CA; CAMK2A; STAT5B; PIK3CB; PIK3C3; GNB2L1; BCL2L1; MAPK3;

TABLE 3-continued

CELLULAR FUNCTION	GENES
	ETS1; KRAS; RUNX1; PIM1; PIK3C2A; RAF1; MAP2K2; AKT1; JAK2; PIK3R1; STAT3; MAP2K1; CCND1; AKT3; STAT1
Amyotrophic Lateral Sclerosis Signaling	BID; IGF1; RAC1; BIRC4; PGF; CAPNS1; CAPN2; PIK3CA; BCL2; PIK3CB; PIK3C3; BCL2L1; CAPN1; PIK3C2A; TP53; CASP9; PIK3R1; RAB5A; CASP1; APAF1; VEGFA; BIRC2; BAX; AKT3; CASP3; BIRC3
JAK/Stat Signaling	PTPN1; MAPK1; PTPN11; AKT2; PIK3CA; STAT5B; PIK3CB; PIK3C3; MAPK3; KRAS; SOCS1; STAT5A; PTPN6; PIK3C2A; RAF1; CDKN1A; MAP2K2; JAK1; AKT1; JAK2; PIK3R1; STAT3; MAP2K1; FRAP1; AKT3; STAT1
Nicotinate and Nicotinamide Metabolism	PRKCE; IRAK1; PRKAA2; EIF2AK2; GRK6; MAPK1; LK1; AKT2; T2; CDK8; MAPK8; MAPK3; PRKCD; PRKAA1; PBEF1; MAPK9; CDK2; PIMI; DYRK1A; MAP2K2; MAP2K1; PAK3; NT5E; TTK; CSNK1A1; BRAF; SGK
Chemokine Signaling	CXCR4; ROCK2; MAPK1; PTK2; FOS; CFL1; GNAQ; CAMK2A; CXCL12; MAPK8; MAPK3; KRAS; MAPK13; RHOA; CCR3; SRC; PPP1CC; MAPK14; NOX1; RAF1; MAP2K2; MAP2K1; JUN; CCL2; PRKCA
IL-2 Signaling	ELK1; MAPK1; PTPN11; AKT2; PIK3CA; SYK; FOS; STAT5B; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; SOCS1; STAT5A; PIK3C2A; LCK; RAF1; MAP2K2; JAK1; AKT1; PIK3R1; MAP2K1; JUN; AKT3
Synaptic Long Term Depression	PRKCE; IGF1; PRKCZ; PRDX6; LYN; MAPK1; GNAS; PRKCI; GNAQ; PPP2R1A; IGF1R; PRKD1; MAPK3; KRAS; GRN; PRKCD; NOS3; NOS2A; PPP2CA; YWHAZ; RAF1; MAP2K2; PPP2R5C; MAP2K1; PRKCA
Estrogen Receptor Signaling	TAF4B; EP300; CARM1; PCAF; MAPK1; NCOR2; SMARCA4; MAPK3; NRIP1; KRAS; SRC; NR3C1; HDAC3; PPARGG1A; RBM9; NCOA3; RAF1; CREBBP; MAP2K2; NCOA2; MAP2K1; PRKDC; ESR1; ESR2
Protein Ubiquitination Pathway	TRAF6; SMURF1; BIRC4; BRCA1; UCHL1; NEDD4; CBL; UBE2I; BTRC; HSPA5; USP7; USP10; FBXW7; USP9X; STUB1; USP22; B2M; BIRC2; PARK2; USP8; USP1; VHL; HSP90AA1; BIRC3
IL-10 Signaling	TRAF6; CCR1; ELK1; IKBKB; SP1; FOS; NFKB2; MAP3K14; MAPK8; MAPK13; RELA; MAPK14; TNF; IKBKG; RELB; MAP3K7; JAK1; CHUK; STAT3; NFKB1; JUN; IL1R1; IL6
VDR/RXR Activation	PRKCE; EP300; PRKCZ; RXRA; GADD45A; HES1; NCOR2; SP1; PRKCI; CDKN1B; PRKD1; PRKCD; RUNX2; KLF4; YY1; NCOA3; CDKN1A; NCOA2; SPP1; LRP5; CEBPB; FOXO1; PRKCA
TGF-beta Signaling	EP300; SMAD2; SMURF1; MAPK1; SMAD3; SMAD1; FOS; MAPK8; MAPK3; KRAS; MAPK9; RUNX2; SERPINE1; RAF1; MAP3K7; CREBBP; MAP2K2; MAP2K1; TGFBR1; SMAD4; JUN; SMAD5
Toll-like Receptor Signaling	IRAK1; EIF2AK2; MYD88; TRAF6; PPARA; ELK1; IKBKB; FOS; NFKB2; MAP3K14; MAPK8; MAPK13; RELA; TLR4; MAPK14; IKBKG; RELB; MAP3K7; CHUK; NFKB1; TLR2; JUN
P38 MAPK Signaling	HSPB1; IRAK1; TRAF6; MAPKAPK2; ELK1; FADD; FAS; CREB1; DDIT3; RPS6KA4; DAXX; MAPK13; TRAF2; MAPK14; TNF; MAP3K7; TGFBR1; MYC; ATF4; IL1R1; SRF; STAT1
Neurotrophin/TRK Signaling	NTRK2; MAPK1; PTPN11; PIK3CA; CREB1; FOS; PIK3CB; PIK3C3; MAPK8; MAPK3; KRAS; PIK3C2A; RAF1; MAP2K2; AKT1; PIK3R1; PDPK1; MAP2K1; CDC42; JUN; ATF4
FXR/RXR Activation	INS; PPARA; FASN; RXRA; AKT2; SDC1; MAPK8; APOB; MAPK10; PPARG; MTPP; MAPK9; PPARGC1A; TNF; CREBBP; AKT1; SREBF1; FGFR4; AKT3; FOXO1
Synaptic Long Term Potentiation	PRKCE; RAPIA; EP300; PRKCZ; MAPK1; CREB1; PRKCI; GNAQ; CAMK2A; PRKD1; MAPK3; KRAS; PRKCD; PPP1CC; RAF1; CREBBP; MAP2K2; MAP2K1; ATF4; PRKCA
Calcium Signaling	RAP1A; EP300; HDAC4; MAPK1; HDAC5; CREB1; CAMK2A; MYH9; MAPK3; HDAC2; HDAC7A; HDAC11; HDAC9; HDAC3; CREBBP; CALR; CAMKK2; ATF4; HDAC6
EGF Signaling	ELK1; MAPK1; EGFR; PIK3CA; FOS; PIK3CB; PIK3C3; MAPK8; MAPK3; PIK3C2A; RAF1; JAK1; PIK3R1; STAT3; MAP2K1; JUN; PRKCA; SRF; STAT1
Hypoxia Signaling in the Cardiovascular System	EDN1; PTEN; EP300; NQO1; UBE2I; CREB1; ARNT; HIF1A; SLC2A4; NOS3; TP53; LDHA; AKT1; ATM; VEGFA; JUN; ATF4; VHL; HSP90AA1

TABLE 3-continued

CELLULAR FUNCTION	GENES
LPS/IL-1 Mediated Inhibition of RXR Function LXR/RXR Activation	IRAK1; MYD88; TRAF6; PPARA; RXRA; ABCA1; MAPK8; ALDH1A1; GSTP1; MAPK9; ABCB1; TRAF2; TLR4; TNF; MAP3K7; NR1H2; SREBF1; JUN; IL1R1 FASN; RXRA; NCOR2; ABCA1; NFKB2; IRF3; RELA; NOS2A; TLR4; TNF; RELB; LDLR; NR1H2; NFKB1; SREBF1; IL1R1; CCL2; IL6; MMP9
Amyloid Processing	PRKCE; CSNK1E; MAPK1; CAPNS1; AKT2; CAPN2; CAPN1; MAPK3; MAPK13; MAPT; MAPK14; AKT1; PSEN1; CSNK1A1; GSK3B; AKT3; APP
IL-4 Signaling	AKT2; PIK3CA; PIK3CB; PIK3C3; IRS1; KRAS; SOCS1; PTPN6; NR3C1; PIK3C2A; JAK1; AKT1; JAK2; PIK3R1; FRAP1; AKT3; RPS6KB1
Cell Cycle: G2/M DNA Damage Checkpoint Regulation	EP300; PCAF; BRCA1; GADD45A; PLK1; BTRC; CHEK1; ATR; CHEK2; YWHAZ; TP53; CDKN1A; PRKDC; ATM; SFN; CDKN2A
Nitric Oxide Signaling in the Cardiovascular System	KDR; FLT1; PGF; AKT2; PIK3CA; PIK3CB; PIK3C3; CAV1; PRKCD; NOS3; PIK3C2A; AKT1; PIK3R1; VEGFA; AKT3; HSP90AA1
Purine Metabolism	NME2; SMARCA4; MYH9; RRM2; ADAR; EIF2AK4; PKM2; ENTPD1; RAD51; RRM2B; TJP2; RAD51C; NT5E; POLD1; NME1
cAMP-mediated Signaling	RAP1A; MAPK1; GNAS; CREB1; CAMK2A; MAPK3; SRC; RAF1; MAP2K2; STAT3; MAP2K1; BRAF; ATF4
Mitochondrial Dysfunction	SOD2; MAPK8; CASP8; MAPK10; MAPK9; CASP9; PARK7; PSEN1; PARK2; APP; CASP3
Notch Signaling	HES1; JAG1; NUMB; NOTCH4; ADAM17; NOTCH2; PSEN1; NOTCH3; NOTCH1; DLL4
Endoplasmic Reticulum Stress Pathway	HSPA5; MAPK8; XBP1; TRAF2; ATF6; CASP9; ATF4; EIF2AK3; CASP3
Pyrimidine Metabolism	NME2; AICDA; RRM2; EIF2AK4; ENTPD1; RRM2B; NT5E; POLD1; NME1
Parkinson's Signaling	UCHL1; MAPK8; MAPK13; MAPK14; CASP9; PARK7; PARK2; CASP3
Cardiac & Beta Adrenergic Signaling	GNAS; GNAQ; PPP2R1A; GNB2L1; PPP2CA; PPP1CC; PPP2R5C
Glycolysis/ Gluconeogenesis	HK2; GCK; GPI; ALDH1A1; PKM2; LDHA; HK1
Interferon Signaling	IRF1; SOCS1; JAK1; JAK2; IFITM1; STAT1; IFIT3
Sonic Hedgehog Signaling	ARRB2; SMO; GLI2; DYRK1A; GLI1; GSK39; DYRK1B
Glycerophospholipid Metabolism	PLD1; GRN; GPAM; YWHAZ; SPHK1; SPHK2
Phospholipid Degradation	PRDX6; PLD1; GRN; YWHAZ; SPHK1; SPHK2
Tryptophan Metabolism	SIAH2; PRMT5; NEDD4; ALDH1A1; CYP1B1; SIAH1
Lysine Degradation	SUV39H1; EHMT2; NSD1; SETD7; PPP2R5C
Nucleotide Excision Repair Pathway	ERCC5; ERCC4; XPA; XPC; ERCC1
Starch and Sucrose Metabolism	UCHL1; HK2; GCK; GPI; HK1
Aminosugars Metabolism	NQO1; HK2; GCK; HK1
Arachidonic Acid Metabolism	PRDX6; GRN; YWHAZ; CYP1B1
Circadian Rhythm Signaling	CSNK1E; CREB1; ATF4; NR1D1
Coagulation System	BDKRB1; F2R; SERPINE1; F3
Dopamine Receptor Signaling	PPP2R1A; PPP2CA; PPP1CC; PPP2R5C
Glutathione Metabolism	IDH2; GSTP1; ANPEP; IDH1
Glycerolipid Metabolism	ALDH1A1; GPAM; SPHK1; SPHK2
Linoleic Acid Metabolism	PRDX6; GRN; YWHAZ; CYP1B1
Methionine Metabolism	DNMT1; DNMT3B; AHCY; DNMT3A
Pyruvate Metabolism	GLO1; ALDH1A1; PKM2; LDHA
Arginine and Proline Metabolism	ALDH1A1; NOS3; NOS2A
Eicosanoid Signaling	PRDX6; GRN; YWHAZ
Fructose and Mannose Metabolism	HK2; GCK; HK1
Galactose Metabolism	HK2; GCK; HK1
Stilbene, Coumarine and Lignin Biosynthesis	PRDX6; PRDX1; TYR
Antigen Presentation Pathway	CALR; B2M
Biosynthesis of Steroids	NQO1; DHCR7
Butanoate Metabolism	ALDH1A1; NLGN1
Citrate Cycle	IDH2; IDH1
Fatty Acid Metabolism	ALDH1A1; CYP1B1

TABLE 3-continued

CELLULAR FUNCTION	GENES
Glycerophospholipid Metabolism	PRDX6; CHKA
Histidine Metabolism	PRMT5; ALDH1A1
Inositol Metabolism	ERO1L; APEX1
Metabolism of Xenobiotics by Cytochrome p450	GSTP1; CYP1B1
Methane Metabolism	PRDX6; PRDX1
Phenylalanine Metabolism	PRDX6; PRDX1
Propanoate Metabolism	ALDH1A1; LDHA
Selenoamino Acid Metabolism	PRMT5; AHCY
Sphingolipid Metabolism	SPHK1; SPHK2
Aminophosphonate Metabolism	PRMT5
Androgen and Estrogen Metabolism	PRMT5
Ascorbate and Aldarate Metabolism	ALDH1A1
Bile Acid Biosynthesis	ALDH1A1
Cysteine Metabolism	LDHA
Fatty Acid Biosynthesis	FASN
Glutamate Receptor Signaling	GNB2L1
NRF2-mediated Oxidative Stress Response	PRDX1
Pentose Phosphate Pathway	GPI
Pentose and Glucuronate Interconversions	UCHL1
Retinol Metabolism	ALDH1A1
Riboflavin Metabolism	TYR
Tyrosine Metabolism	PRMT5, TYR
Ubiquinone Biosynthesis	PRMT5
Valine, Leucine and Isoleucine Degradation	ALDH1A1
Glycine, Serine and Threonine Metabolism	CHKA
Lysine Degradation	ALDH1A1
Pain/Taste	TRPM5; TRPA1
Pain	TRPM7; TRPC5; TRPC6; TRPC1; Cnr1; cnr2; Grk2; Trpa1; Pomc; Cgrp; Crf; Pka; Era; Nr2b; TRPM5; Prkaca; Prkacb; Prkar1a; Prkar2a
Mitochondrial Function	AIF; CytC; SMAC (Diablo); Aifm-1; Aifm-2
Developmental Neurology	BMP-4; Chordin (Chrd); Noggin (Nog); WNT (Wnt2; Wnt2b, Wnt3a, Wnt4; Wnt5a; Wnt6; Wnt7b; Wnt8b; Wnt9a; Wnt9b; Wnt10a; Wnt10b; Wnt16); beta-catenin; Dkk-1; Frizzled related proteins; Otx-2; Gbx2; FGF-8; Reelin; Dab1; unc-86 (Pou4f1 or Brn3a); Numb; Reln

TABLE 4

INDICATION(S)	THERAPEUTIC PROTEIN
Maple syrup urine disease	3-methyl-2-oxobutanoate dehydrogenase
Medium-chain acyl-CoA dehydrogenase deficiency	Acyl-CoA dehydrogenase
Alpha 1-antitrypsin deficiency	Alpha 1 protease inhibitor
Pompe disease	Alpha glucosidase
Paroxysmal nocturnal hemoglobinuria	Anti-complement factor C5 Mab
Familial dysbetalipoproteinemia	Apolipoprotein E
Argininemia	Arginase
Argininosuccinic acidemia	Argininosuccinate lyase
Citrullinemia, type I	Argininosuccinate synthase
Short-chain acyl-CoA dehydrogenase deficiency	Butyryl-CoA dehydrogenase
Hereditary angioedema	C1 esterase inhibitor
Carbamylphosphate synthetase deficiency	Carbamylphosphate synthetase
Cystic fibrosis	CFTR
Hemophilia B	Factor IX

TABLE 4-continued

INDICATION(S)	THERAPEUTIC PROTEIN
Hemophilia A, Hemophilia B	Factor VII
Hemophilia A	Factor VIII
Classical galactosemia	Galactose-1-phosphate uridylyltransferase
von Gierke's disease	Glucose-6-phosphatase
Glutaric acidemia, type I	Glutaryl-CoA dehydrogenase
Isovaleric aciduria	Isovaleric acid CoA dehydrogenase deficiency
Homozygous familial hypercholesterolemia	LDL receptor
Long-chain 3-OH acyl-CoA dehydrogenase deficiency	Long-chain-3-hydroxyacyl-CoA dehydrogenase
Very long-chain acyl-CoA dehydrogenase deficiency	Long-chain-acyl-CoA dehydrogenase
Methylmalonyl-CoA mutase deficiency	Methylmalonyl-CoA mutase
Ornithine transcarbamylase deficiency	Ornithine transcarbamylase
Phenylketonuria	Phenylalanine hydroxylase
Acute intermittent porphyria	Porphobilinogen deaminase
Propionic acidemia	Propionyl-CoA carboxylase
Hyperoxaluria, type I	Serine-pyruvate aminotransferase
Crigler-Najjar syndrome	UDP-glucuronosyltransferase
Non-Hodgkin lymphoma	Anti-CD20 mAb
Allergic asthma	Anti-IgE mAb
Psoriasis	Anti-IL-12 & IL-23 mAb
Rheumatoid arthritis	Anti-interleukin-6 (IL-6) mAb
Anemia	Erythropoietin
Rheumatoid arthritis	T-cell costimulation blocker
Rheumatoid arthritis	TNF-alpha inhibitors (including anti-TNF-alpha mAb)
Gout	Urate oxidase
Familial chylomicronemia	Lipoprotein lipase
Melanoma	Anti-CTLA4 mAb
Head and neck cancer, Metastatic colorectal cancer	Anti-EGFR mAb
HER2+ breast cancer, gastric cancer	Anti-HER2 mAb
Metastatic colorectal cancer, NSCLC, others	Anti-VEGF mAb
Blepharospasm, Cervical dystonia, Chronic migraine, more	Botulinum toxin
Female infertility	Follicle stimulating hormone
Type 2 diabetes mellitus	Glucagon-like peptide 1 (GLP-1) agonist
Growth hormone deficiency	Growth hormone 1/Growth hormone 2
Type 2 diabetes mellitus	Insulin
Hypoparathyroidism	Parathyroid hormone
Asthma	SERCA2
Asthma	FoxP3
Surfactant Deficiency	Pulmonary surfactants (SFTPA1, SFTPB, SFTPC, SFTPD)
Pulmonary Alveolar proteinosis	GM-CSF Receptor (CSF2RA, CSF2RB)
alport syndrome	Col4A5
Stargardt's Disease	ABCA4
Retinitis pigmentosa	Rhodopsins
Adrenoleukodystrophy	ABCD1
Adenosine deaminase deficiency	Adenosine deaminase
Familial adenomatous polyposis	APC
Autosomal recessive polycystic kidney disease	ARPKD
Metachromatic leukodystrophy	Arylsulfatase A
Batten disease	Battenin + others
Beta-thalassemia	Beta globin
X-linked agammaglobulinemia	Bruton's tyrosine kinase
Becker muscular dystrophy	Dystrophin
Duchenne muscular dystrophy	Dystrophin
Marfan syndrome	FBN1
Fragile X syndrome	FMRP
Krabbe disease	Galactocerebrosidase
Sickle cell disease	Hemoglobin
Sanfilippo syndrome, type A (MPS IIIA)	Heparan N-sulfatase
GM2 gangliosidosis	HEXA, HEXB
Hemachromatosis	HFE protein
Huntington disease	Huntingtin
Lesch-Nyhan syndrome	Hypoxanthine phosphoribosyltransferase 1
McArdle disease	Muscle glycogen phosphorylase
Sanfilippo syndrome, type B (MPS IIIB)	N-acetyl-alpha-D-glucosaminidase
Leber's hereditary optic neuropathy	NADH dehydrogenase
Neurofibromatosis, type 1	NF-1

TABLE 4-continued

INDICATION(S)	THERAPEUTIC PROTEIN
Niemann Pick disease, type C	NPC1
Alpers' disease	POLG
Von Hippel-Lindau disease	pVHL
Paget disease of bone	Sequestosome 1
Carnitine uptake defect	SLC22A5
Cystinuria	SLC7A9
Niemann Pick disease, type A/B	SMPD1
Spinal muscular atrophy	Survival motor neuron protein
Li-Fraumeni syndrome	TP53
Fabry disease	Alpha galactosidase
Alpha-mannosidosis	Alpha-D-mannosidase
Hurler syndrome (MPS I)	Alpha-L iduronidase
Hemolytic uremic syndrome	Anti-complement factor C5 mAb
Morquio syndrome, type B (MPS IVB)	Beta-galactosidase
Multiple carboxylase deficiency	Biotin-methylcrotonoyl-CoA-carboxylase ligase
Homocystinuria	Cystathionine beta-synthase
Cystinosis	Cystinosin
Cystic fibrosis	Deoxyribonuclease I
Erythropoietic protoporphyria	Ferrochelataase
Tyrosinemia, type I	Fumarylacetoacetase
GALK deficiency	Galactokinase
Morquio syndrome, type A (MPS IVA)	Galactose 6-sulfate sulfatase
GALE deficiency	Galactose epimerase
Gaucher disease	Glucocerebrosidase
Alkaptonuria	Homogentisate 1,2-dioxygenase
Hunter syndrome (MPS II)	Iduronate-2-sulfatase
Lysosomal acid lipase deficiency	Lysosomal acid lipase
Hypermethioninemia	Methionine adenosyltransferase
3-Methylcrotonyl-CoA carboxylase deficiency	Methylcrotonoyl-CoA carboxylase
3-Methylglutaconic aciduria	Methylglutaconyl-CoA hydratase
Maroteaux-Lamy syndrome (MPS VI)	N-acetylgalactosamine 4-sulfatase
Familial mediterranean fever	Pyrin (MEFV)
Tetrahydrobiopterin-deficient hyperphenylalaninemia	Tetrahydrobiopterin
Juvenile rheumatoid arthritis	TNF-alpha inhibitors
Psoriatic arthritis	TNF-alpha inhibitors
Hypophosphatasia	TNSALP
Gilbert syndrome	UDP-glucuronosyltransferase
Porphyria cutanea tarda	Uroporphyrinogen decarboxylase
Wilson disease	Wilson disease protein
Systemic lupus erythematosus	Anti-BAFF
Osteoporosis	Anti-RANKL mAb
Multiple sclerosis	Anti-VLA-4 mAb
Neutropenia	G-CSF
Immunoglobulin deficiency	Immunoglobulin
Primary humoral immune deficiencies (e.g., CVID)	Immunoglobulin
Infectious diseases vaccines	Infectious antigen
Hepatitis B, Hepatitis C	Interferon alpha
Multiple sclerosis	Interferon beta
Chronic immune thrombocytopenia	Thrombopoietin
Ehlers-Danlos syndrome, type 1	Proteins encoded by ADAMTS2, B3GALT6, B4GALT7, CHST14, COL1A1, COL1A2, COL3A1, COL5A1, COL5A2, DSE, FKBP14, PLOD1, PRDM5, SLC39A13, TNXB, and ZNF469
Stickler syndrome	Proteins encoded by COL11A1, COL11A2, COL2A1, COL9A1, COL9A2, and COL9A3
Hereditary hemorrhagic telangiectasia	Proteins encoded by ACVRL1, ENG, and SMAD4
Hereditary spherocytosis	Proteins encoded by ANK1, EPB42, SLC4A1, SPTA1 and SPTB
Brugada syndrome	Proteins encoded by CACNA1C, CACNA2D1, CACNB2, GPD1L, HCN4, KCND3, KCNE3, KCNE5, KCNJ8, RANGRF, SCN1B, SCN2B, SCN3B, SCN5A, SLMAP, and TRPM4
Osteopetrosis	Proteins encoded by CA2, CLCN7, IKBKG, ITGB3, OSTM1, PLEKHM1, TCIRG1, TNFRSF11A, and TNFSF11
Mitochondrial oxidative phosphorylation disorders	Proteins encoded by FBXL4, and NDUFB9

TABLE 5

INDICATION(S)	THERAPEUTIC PROTEIN	GENE
Achromatopsia type 2	Cyclic nucleotide-gated channel, $\alpha 3$ subunit	CNGA3
Achromatopsia type 3	Cyclic nucleotide-gated channel, $\beta 3$ subunit	CNGB3
Aland Island eye disease	Cav1.4: calcium channel, voltage-gated, L type, $\alpha 1F$ subunit	CACNA1F
Andersen-Tawil syndrome	Kir2.1: potassium channel, inwardly-rectifying, subfamily J, member 2	KCNJ2
Benign familial infantile epilepsy	Nav2.1: sodium channel, voltage-gated, type II, α subunit	SCN2A
	Kv7.2: potassium channel, voltage-gated, KQT-like subfamily, member 2	KCNQ2
	Kv7.3: potassium channel, voltage-gated, KQT-like subfamily, member 3	KCNQ3
Bestrophinopathy, autosomal-recessive	Bestrophin 1	BEST1
Central core disease	RyR1: ryanodine receptor 1	RYR1
Charcot-Marie-Tooth disease type 2C	Transient receptor potential cation channel, subfamily V, member 4	TRPV4
Childhood absence epilepsy	γ -aminobutyric acid A receptor, $\alpha 1$ subunit	GABRA1
	γ -aminobutyric acid A receptor, $\alpha 6$ subunit	GABRA6
	γ -aminobutyric acid A receptor, $\beta 3$ subunit	GABRB3
	γ -aminobutyric acid A receptor, $\gamma 2$ subunit	GABRG2
	Cav3.2: calcium channel, voltage-gated, T type, $\alpha 1H$ subunit	CACNA1H
Cognitive impairment with or without cerebellar ataxia	Nav1.6: sodium channel, voltage-gated, type VIII, α subunit	SCN8A
Cone-rod dystrophy, X-linked, type 3	Cav1.4: calcium channel, voltage-gated, L type, $\alpha 1F$ subunit	CACNA1F
Congenital distal spinal muscular atrophy	Transient receptor potential cation channel, subfamily V, member 4	TRPV4
Congenital indifference to pain, autosomal-recessive	Nav1.7: Sodium channel, voltage-gated, type IX, α subunit	SCN9A
Congenital myasthenic syndrome	Cholinergic receptor, muscle nicotinic, $\alpha 1$ subunit	CHRNA1
	Cholinergic receptor, muscle nicotinic, $\beta 1$ subunit	CHRNA1
	Cholinergic receptor, muscle nicotinic, δ subunit	CHRNA1
	Cholinergic receptor, muscle nicotinic, ϵ subunit	CHRNA1
	Nav1.4: sodium channel, voltage-gated, type IV, α subunit	SCN4A
Congenital stationary night blindness type 1C	Transient receptor potential cation channel, subfamily M, member 1	TRPM1
Congenital stationary night blindness type 2A	Cav1.4: calcium channel, voltage-gated, L type, $\alpha 1F$ subunit	CACNA1F
Deafness, autosomal-dominant, type 2A	Kv7.4: potassium channel, voltage-gated, KQT-like subfamily, member 4	KCNQ4
Deafness, autosomal-recessive, type 4, with enlarged vestibular aqueduct	Kir4.1: potassium channel, inwardly-rectifying, subfamily J, member 10	KCNJ10
Dravet syndrome	Nav1.1: sodium channel, voltage-gated, type I, α subunit	SCN1A
Early infantile epileptic encephalopathy type 7	γ -aminobutyric acid A receptor, $\gamma 2$ subunit	GABRG2
	Kv7.2: potassium channel, voltage-gated, KQT-like subfamily, member 2	KCNQ2
Early infantile epileptic encephalopathy type 11	Nav2.1: sodium channel, voltage-gated, type II, α subunit	SCN2A
Early infantile epileptic encephalopathy type 13	Nav1.6: sodium channel, voltage-gated, type VIII, α subunit	SCN8A
Early infantile epileptic encephalopathy type 14	KCa4.1: potassium channel, subfamily T, member 1	KCNT1
EAST/SeSAME syndrome	Kir4.1: potassium channel, inwardly-rectifying, subfamily J, member 10	KCNJ10

TABLE 5-continued

INDICATION(S)	THERAPEUTIC PROTEIN	GENE
Episodic ataxia type 1	Kv1.1: potassium channel, voltage-gated, shaker-related subfamily, member 1	KCNA1
Episodic ataxia type 2	Cav2.1: calcium channel, voltage-gated, P/Q type, α 1A subunit	CACNA1A
Episodic ataxia type 5	Cav β 4: calcium channel, voltage-gated, β 4 subunit	CACNB4
Familial episodic pain syndrome	Transient receptor potential cation channel, subfamily A, member 1	TRPA1
Familial hemiplegic migraine type 1	Cav2.1: calcium channel, voltage-gated, P/Q type, α 1A subunit	CACNA1A
Familial hemiplegic migraine type 3	Nav1.1: sodium channel, voltage-gated, type I, α subunit	SCN1A
Generalized epilepsy with febrile seizures plus (GEFS+)	Nav β 1: sodium channel, voltage-gated, type I, β subunit	SCN1B
	Nav1.1: sodium channel, voltage-gated, type I, α subunit	SCN1A
	γ -aminobutyric acid A receptor, γ 2 subunit	GABRG2
Generalized epilepsy with paroxysmal dyskinesia	KCa1.1: potassium channel, calcium-activated, large conductance, subfamily M, α 1 subunit	KCNMA1
Hereditary hyperekplexia	Glycine receptor, α 1 subunit	GLRA1
	Glycine receptor, β subunit	GLRB
Hyperkalemic periodic paralysis	Nav1.4: sodium channel, voltage-gated, type IV, α subunit	SCN4A
Hypokalemic periodic paralysis type 1	Cav1.1: calcium channel, voltage-gated, L type, α 1S subunit	CACNA1S
Hypokalemic periodic paralysis type 2	Nav1.4: sodium channel, voltage-gated, type IV, α subunit	SCN4A
Juvenile macular degeneration	Cyclic nucleotide-gated channel, β 3 subunit	CNGB3
Juvenile myoclonic epilepsy	γ -aminobutyric acid A receptor, α 1 subunit	GABRA1
	Cav β 4: calcium channel, voltage-gated, β 4 subunit	CACNB4
Malignant hyperthermia susceptibility	RyR1: ryanodine receptor 1	RYR1
	Cav1.1: calcium channel, voltage-gated, L type, α 1S subunit	CACNA1S
Mucopolidosis type IV	TRPML1/mucolipin 1	MCOLN1
Multiple pterygium syndrome, lethal type	Cholinergic receptor, muscle nicotinic, α 1 subunit	CHRNA1
Multiple pterygium syndrome, nonlethal type (Escobar variant)	Cholinergic receptor, muscle nicotinic, δ subunit	CHRND
	Cholinergic receptor, muscle nicotinic, γ subunit	CHRNG
Myotonia congenita, autosomal-dominant (Thomsen disease)	ClC-1: chloride channel 1, voltage-gated	CLCN1
Myotonia congenita, autosomal-recessive (Becker disease)	ClC-1: chloride channel 1, voltage-gated	CLCN1
Nocturnal frontal lobe epilepsy type 1	Cholinergic receptor, neuronal nicotinic, α 4 subunit	CHRNA4
Nocturnal frontal lobe epilepsy type 3	Cholinergic receptor, neuronal nicotinic, β 2 subunit	CHRNA2
Nocturnal frontal lobe epilepsy type 4	Cholinergic receptor, neuronal nicotinic, α 2 subunit	CHRNA2
Nocturnal frontal lobe epilepsy type 5	KCa4.1: potassium channel, subfamily T, member 1	KCNT1
Paramyotonia congenita	Nav1.4: sodium channel, voltage-gated, type IV, α subunit	SCN4A
Paroxysmal extreme pain disorder	Nav1.7: Sodium channel, voltage-gated, type IX, α subunit	SCN9A
Potassium-aggravated myotonia	Nav1.4: sodium channel, voltage-gated, type IV, α subunit	SCN4A
Primary erythralgia	Nav1.7: sodium channel, voltage-gated, type IX, α subunit	SCN9A
Retinitis pigmentosa type 45, autosomal-recessive	Cyclic nucleotide-gated channel, β 1 subunit	CNGB1
Retinitis pigmentosa type 49, autosomal-recessive	Cyclic nucleotide-gated channel, α 1 subunit	CNGA1
Retinitis pigmentosa type 50, autosomal-dominant	Bestrophin 1	BEST1
Scapuloperoneal spinal muscular atrophy	Transient receptor potential cation channel, subfamily V, member 4	TRPV4

TABLE 5-continued

INDICATION(S)	THERAPEUTIC PROTEIN	GENE
Small fiber neuropathy	Nav1.7: sodium channel, voltage-gated, type IX, α subunit	SCN9A
Spinocerebellar ataxia type 6	Cav2.1: calcium channel, voltage-gated, P/Q type, α 1A subunit	CACNA1A
Spinocerebellar ataxia type 13	Kv3.3: potassium channel, voltage-gated, Shaw-related subfamily, member 3	KCNC3
Vitelliform macular dystrophy	Bestrophin 1	BEST1
Vitreoretinopathy	Bestrophin 1	BEST1

TABLE 6

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
A1E959	Odontogenic ameloblast-associated protein	ODAM
A1KZ92	Peroxidasin-like protein	PXDNL
A1L453	Serine protease 38	PRSS38
A1L4H1	Soluble scavenger receptor cysteine-rich domain-containing protein SSC5D	SSC5D
A2RUU4	Colipase-like protein 1	CLPSL1
A2VDF0	Fucose mutarotase	FUOM
A2VEC9	SCO-spondin	SSPO
A3KMH1	von Willebrand factor A domain-containing protein 8	VWA8
A4D0S4	Laminin subunit beta-4	LAMB4
A4D1T9	Probable inactive serine protease 37	PRSS37
A5D8T8	C-type lectin domain family 18 member A	CLEC18A
A6NC86	phospholipase A2 inhibitor and Ly6/PLAUR domain-containing protein	PINLYP
A6NCI4	von Willebrand factor A domain-containing protein 3A	VWA3A
A6ND01	Probable folate receptor delta	FOLR4
A6NDD2	Beta-defensin 108B-like	
A6NE02	BTB/POZ domain-containing protein 17	BTBD17
A6NEF6	Growth hormone 1	GH1
A6NF02	NPIP-like protein LOC730153	
A6NFB4	HCG1749481, isoform CRA_k	CSH1
A6NFZ4	Protein FAM24A	FAM24A
A6NG13	Glycosyltransferase 54 domain-containing protein	
A6NGN9	IgLON family member 5	IGLON5
A6NHN0	Otolin-1	OTOL1
A6NHN6	Nuclear pore complex-interacting protein-like 2	NPIPL2
A6NI73	Leukocyte immunoglobulin-like receptor subfamily A member 5	LILRA5
A6NIT4	Chorionic somatomammotropin hormone 2 isoform 2	CSH2
A6NJ69	IgA-inducing protein homolog	IGIP
A6NKQ9	Choriogonadotropin subunit beta variant 1	CGB1
A6NMZ7	Collagen alpha-6(VI) chain	COL6A6
A6NNS2	Dehydrogenase/reductase SDR family member 7C	DHRS7C
A6XGL2	Insulin A chain	INS
A8K0G1	Protein Wnt	WNT7B
A8K2U0	Alpha-2-macroglobulin-like protein 1	A2ML1
A8K7I4	Calcium-activated chloride channel regulator 1	CLCA1
A8MTL9	Serpin-like protein HMSD	HMSD
A8MV23	Serpin E3	SERPINE3
A8MZH6	Oocyte-secreted protein 1 homolog	OOSP1
A8TX70	Collagen alpha-5(VI) chain	COL6A5
B0ZBE8	Natriuretic peptide	NPPA
B1A4G9	Somatotropin	GH1
B1A4H2	HCG1749481, isoform CRA_d	CSH1
B1A4H9	Chorionic somatomammotropin hormone	CSH2
B1AJZ6	Protein Wnt	WNT4
B1AKI9	Isthmin-1	ISM1
B2RNN3	Complement C1q and tumor necrosis factor-related protein 9B	C1QTNF9B
B2RUY7	von Willebrand factor C domain-containing protein 2-like	VWC2L
B3GLJ2	Prostate and testis expressed protein 3	PATE3
B4DI03	SEC11-like 3 (<i>S. cerevisiae</i>), isoform CRA_a	SEC11L3
B4DJF9	Protein Wnt	WNT4

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
B4DUL4	SEC11-like 1 (<i>S. cerevisiae</i>), isoform CRA_d	SEC11L1
B5MCC8	Protein Wnt	WNT10B
B8A595	Protein Wnt	WNT7B
B8A597	Protein Wnt	WNT7B
B8A598	Protein Wnt	WNT7B
B9A064	Immunoglobulin lambda-like polypeptide 5	IGLL5
C9J3H3	Protein Wnt	WNT10B
C9J8I8	Protein Wnt	WNT5A
C9JAF2	Insulin-like growth factor II Ala-25 Del	IGF2
C9JCI2	Protein Wnt	WNT10B
C9JL84	HERV-H LTR-associating protein 1	HHLA1
C9JNR5	Insulin A chain	INS
C9JUI2	Protein Wnt	WNT2
D6RF47	Protein Wnt	WNT8A
D6RF94	Protein Wnt	WNT8A
E2RYF7	Protein PBMUCL2	HCG22
E5RFR1	PENK(114-133)	PENK
E7EML9	Serine protease 44	PRSS44
E7EPC3	Protein Wnt	WNT9B
E7EVP0	Nociceptin	PNOC
E9PD02	Insulin-like growth factor I	IGF1
E9PH60	Protein Wnt	WNT16
E9PJL6	Protein Wnt	WNT11
F5GYM2	Protein Wnt	WNT5B
F5H034	Protein Wnt	WNT5B
F5H364	Protein Wnt	WNT5B
F5H7Q6	Protein Wnt	WNT5B
F8WCM5	Protein INS-IGF2	INS-IGF2
F8WDR1	Protein Wnt	WNT2
H0Y663	Protein Wnt	WNT4
H0YK72	Signal peptidase complex catalytic subunit SEC11A	SEC11A
H0YK83	Signal peptidase complex catalytic subunit SEC11A	SEC11A
H0YM39	Chorionic somatomammotropin hormone	CSH2
H0YMT7	Chorionic somatomammotropin hormone	CSH1
H0YN17	Chorionic somatomammotropin hormone	CSH2
H0YNA5	Signal peptidase complex catalytic subunit SEC11A	SEC11A
H0YNG3	Signal peptidase complex catalytic subunit SEC11A	SEC11A
H0YNX5	Signal peptidase complex catalytic subunit SEC11A	SEC11A
H7BZB8	Protein Wnt	WNT10A
H9KV56	Choriogonadotropin subunit beta variant 2	CGB2
I3L0L8	Protein Wnt	WNT9B
J3KNZ1	Choriogonadotropin subunit beta variant 1	CGB1
J3KP00	Choriogonadotropin subunit beta	CGB7
J3QT02	Choriogonadotropin subunit beta variant 1	CGB1
O00175	C-C motif chemokine 24	CCL24
O00182	Galectin-9	LGALS9
O00187	Mannan-binding lectin serine protease 2	MASP2
O00230	Cortistatin	CORT
O00253	Agouti-related protein	AGRP
O00270	12-(S)-hydroxy-5,8,10,14-eicosatetraenoic acid receptor	GPR31
O00292	Left-right determination factor 2	LEFTY2
O00294	Tubby-related protein 1	TULP1
O00295	Tubby-related protein 2	TULP2
O00300	Tumor necrosis factor receptor superfamily member 11B	TNFRSF11B
O00339	Matrilin-2	MATN2
O00391	Sulfhydryl oxidase 1	QSOX1
O00468	Agtrin	AGRN
O00515	Ladinin-1	LAD1
O00533	Processed neural cell adhesion molecule L1-like protein	CHL1
O00584	Ribonuclease T2	RNASET2
O00585	C-C motif chemokine 21	CCL21
O00602	Ficolin-1	FCN1
O00622	Protein CYR61	CYR61
O00626	MDC(5-69)	CCL22
O00634	Netrin-3	NTN3
O00744	Protein Wnt-10b	WNT10B
O00755	Protein Wnt-7a	WNT7A

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
O14498	Immunoglobulin superfamily containing leucine-rich repeat protein	ISLR
O14511	Pro-neuregulin-2, membrane-bound isoform	NRG2
O14594	Neurocan core protein	NCAN
O14625	C-X-C motif chemokine 11	CXCL11
O14638	Ectonucleotide pyrophosphatase/phosphodiesterase family member 3	ENPP3
O14656	Torsin-1A	TOR1A
O14657	Torsin-1B	TOR1B
O14786	Neuropilin-1	NRP1
O14788	Tumor necrosis factor ligand superfamily member 11, membrane form	TNFSF11
O14791	Apolipoprotein L1	APOL1
O14793	Growth/differentiation factor 8	MSTN
O14904	Protein Wnt-9a	WNT9A
O14905	Protein Wnt-9b	WNT9B
O14944	Proepiregulin	EREG
O14960	Leukocyte cell-derived chemotaxin-2	LECT2
O15018	Processed PDZ domain-containing protein 2	PDZD2
O15041	Semaphorin-3E	SEMA3E
O15072	A disintegrin and metalloproteinase with thrombospondin motifs 3	ADAMTS3
O15123	Angiopoietin-2	ANGPT2
O15130	Neuropeptide FF	NPFF
O15197	Ephrin type-B receptor 6	EPHB6
O15204	ADAM DEC1	ADAMDEC1
O15230	Laminin subunit alpha-5	LAMA5
O15232	Matrilin-3	MATN3
O15240	Neuroendocrine regulatory peptide-1	VGF
O15263	Beta-defensin 4A	DEFB4A
O15335	Chondroadherin	CHAD
O15393	Transmembrane protease serine 2 catalytic chain	TMPRSS2
O15444	C-C motif chemokine 25	CCL25
O15467	C-C motif chemokine 16	CCL16
O15496	Group 10 secretory phospholipase A2	PLA2G10
O15520	Fibroblast growth factor 10	FGF10
O15537	Retinoschisin	RS1
O43157	Plexin-B1	PLXNB1
O43184	Disintegrin and metalloproteinase domain-containing protein 12	ADAM12
O43240	Kallikrein-10	KLK10
O43278	Kunitz-type protease inhibitor 1	SPINT1
O43320	Fibroblast growth factor 16	FGF16
O43323	Desert hedgehog protein C-product	DHH
O43405	Cochlin	COCH
O43508	Tumor necrosis factor ligand superfamily member 12, membrane form	TNFSF12
O43555	Progonadoliberin-2	GNRH2
O43557	Tumor necrosis factor ligand superfamily member 14, soluble form	TNFSF14
O43692	Peptidase inhibitor 15	PI15
O43699	Sialic acid-binding Ig-like lectin 6	SIGLEC6
O43820	Hyaluronidase-3	HYAL3
O43827	Angiopoietin-related protein 7	ANGPTL7
O43852	Calumenin	CALU
O43854	EGF-like repeat and discoidin I-like domain-containing protein 3	EDIL3
O43866	CD5 antigen-like	CD5L
O43897	Tolloid-like protein 1	TLL1
O43915	Vascular endothelial growth factor D	FIGF
O43927	C-X-C motif chemokine 13	CXCL13
O60218	Aldo-keto reductase family 1 member B10	AKR1B10
O60235	Transmembrane protease serine 11D	TMPRSS11D
O60258	Fibroblast growth factor 17	FGF17
O60259	Kallikrein-8	KLK8
O60383	Growth/differentiation factor 9	GDF9
O60469	Down syndrome cell adhesion molecule	DSCAM
O60542	Persephin	PSPN
O60565	Gremlin-1	GREM1
O60575	Serine protease inhibitor Kazal-type 4	SPINK4
O60676	Cystatin-8	CST8
O60687	Sushi repeat-containing protein SRPX2	SRPX2
O60844	Zymogen granule membrane protein 16	ZG16
O60882	Matrix metalloproteinase-20	MMP20
O60938	Keratocan	KERA

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
O75015	Low affinity immunoglobulin gamma Fc region receptor III-B	FCGR3B
O75077	Disintegrin and metalloproteinase domain-containing protein 23	ADAM23
O75093	Slit homolog 1 protein	SLIT1
O75094	Slit homolog 3 protein	SLIT3
O75095	Multiple epidermal growth factor-like domains protein 6	MEGF6
O75173	A disintegrin and metalloproteinase with thrombospondin motifs 4	ADAMTS4
O75200	Nuclear pore complex-interacting protein-like 1	NPIPL1
O75339	Cartilage intermediate layer protein 1 C1	CILP
O75354	Ectonucleoside triphosphate diphosphohydrolase 6	ENTPD6
O75386	Tubby-related protein 3	TULP3
O75398	Deformed epidermal autoregulatory factor 1 homolog	DEAF1
O75443	Alpha-tectorin	TECTA
O75445	Usherin	USH2A
O75462	Cytokine receptor-like factor 1	CRLF1
O75487	Glypican-4	GPC4
O75493	Carbonic anhydrase-related protein 11	CA11
O75594	Peptidoglycan recognition protein 1	PGLYRP1
O75596	C-type lectin domain family 3 member A	CLEC3A
O75610	Left-right determination factor 1	LEFTY1
O75629	Protein CREG1	CREG1
O75636	Ficolin-3	FCN3
O75711	Scrapie-responsive protein 1	SCRG1
O75715	Epididymal secretory glutathione peroxidase	GPX5
O75718	Cartilage-associated protein	CRTAP
O75829	Chondrosurfactant protein	LECT1
O75830	Serpin I2	SERPINI2
O75882	Attractin	ATRN
O75888	Tumor necrosis factor ligand superfamily member 13	TNFSF13
O75900	Matrix metalloproteinase-23	MMP23A
O75951	Lysozyme-like protein 6	LYZL6
O75973	C1q-related factor	C1QL1
O76038	Secretagogin	SCGN
O76061	Stanniocalcin-2	STC2
O76076	WNT1-inducible-signaling pathway protein 2	WISP2
O76093	Fibroblast growth factor 18	FGF18
O76096	Cystatin-F	CST7
O94769	Extracellular matrix protein 2	ECM2
O94813	Slit homolog 2 protein C-product	SLIT2
O94907	Dickkopf-related protein 1	DKK1
O94919	Endonuclease domain-containing 1 protein	ENDOD1
O94964	N-terminal form	SOGA1
O95025	Semaphorin-3D	SEMA3D
O95084	Serine protease 23	PRSS23
O95150	Tumor necrosis factor ligand superfamily member 15	TNFSF15
O95156	Neurexophilin-2	NXPH2
O95157	Neurexophilin-3	NXPH3
O95158	Neurexophilin-4	NXPH4
O95388	WNT1-inducible-signaling pathway protein 1	WISP1
O95389	WNT1-inducible-signaling pathway protein 3	WISP3
O95390	Growth/differentiation factor 11	GDF11
O95393	Bone morphogenetic protein 10	BMP10
O95399	Urotensin-2	UTS2
O95407	Tumor necrosis factor receptor superfamily member 6B	TNFRSF6B
O95428	Papilin	PAPLN
O95445	Apolipoprotein M	APOM
O95450	A disintegrin and metalloproteinase with thrombospondin motifs 2	ADAMTS2
O95460	Matrilin-4	MATN4
O95467	LHAL tetrapeptide	GNAS
O95631	Netrin-1	NTN1
O95633	Follistatin-related protein 3	FSTL3
O95711	Lymphocyte antigen 86	LY86
O95715	C-X-C motif chemokine 14	CXCL14
O95750	Fibroblast growth factor 19	FGF19
O95760	Interleukin-33	IL33
O95813	Cerberus	CER1
O95841	Angiopoietin-related protein 1	ANGPTL1

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
O95897	Noelin-2	OLFM2
O95925	Eppin	EPPIN
O95965	Integrin beta-like protein 1	ITGBL1
O95967	EGF-containing fibulin-like extracellular matrix protein 2	EFEMP2
O95968	Secretoglobin family 1D member 1	SCGB1D1
O95969	Secretoglobin family 1D member 2	SCGB1D2
O95970	Leucine-rich glioma-inactivated protein 1	LGI1
O95972	Bone morphogenetic protein 15	BMP15
O95994	Anterior gradient protein 2 homolog	AGR2
O95998	Interleukin-18-binding protein	IL18BP
O96009	Napsin-A	NAPSA
O96014	Protein Wnt-11	WNT11
P00450	Ceruloplasmin	CP
P00451	Factor VIIIa light chain	F8
P00488	Coagulation factor XIII A chain	F13A1
P00533	Epidermal growth factor receptor	EGFR
P00709	Alpha-lactalbumin	LALBA
P00734	Prothrombin	F2
P00738	Haptoglobin beta chain	HP
P00739	Haptoglobin-related protein	HPR
P00740	Coagulation factor IXa heavy chain	F9
P00742	Factor X heavy chain	F10
P00746	Complement factor D	CFD
P00747	Plasmin light chain B	PLG
P00748	Coagulation factor XIIa light chain	F12
P00749	Urokinase-type plasminogen activator long chain A	PLAU
P00750	Tissue-type plasminogen activator	PLAT
P00751	Complement factor B Ba fragment	CFB
P00797	Renin	REN
P00973	2'-5'-oligoadenylate synthase 1	OAS1
P00995	Pancreatic secretory trypsin inhibitor	SPINK1
P01008	Antithrombin-III	SERPINC1
P01009	Alpha-1-antitrypsin	SERPINA1
P01011	Alpha-1-antichymotrypsin His-Pro-less	SERPINA3
P01019	Angiotensin-1	AGT
P01023	Alpha-2-macroglobulin	A2M
P01024	Acylation stimulating protein	C3
P01031	Complement C5 beta chain	C5
P01033	Metalloproteinase inhibitor 1	TIMP1
P01034	Cystatin-C	CST3
P01036	Cystatin-S	CST4
P01037	Cystatin-SN	CST1
P01042	Kininogen-1 light chain	KNG1
P01127	Platelet-derived growth factor subunit B	PDGFB
P01135	Transforming growth factor alpha	TGFA
P01137	Transforming growth factor beta-1	TGFB1
P01138	Beta-nerve growth factor	NGF
P01148	Gonadoliberin-1	GNRH1
P01160	Atrial natriuretic factor	NPPA
P01178	Oxytocin	OXT
P01185	Vasopressin-neurophysin 2-copeptin	AVP
P01189	Corticotropin	POMC
P01210	PENK(237-258)	PENK
P01213	Alpha-neoendorphin	PDYN
P01215	Glycoprotein hormones alpha chain	CGA
P01222	Thyrotropin subunit beta	TSHB
P01225	Follitropin subunit beta	FSHB
P01229	Lutropin subunit beta	LHB
P01233	Choriogonadotropin subunit beta	CGB8
P01236	Prolactin	PRL
P01241	Somatotropin	GH1
P01242	Growth hormone variant	GH2
P01243	Chorionic somatomammotropin hormone	CSH2
P01258	Katacalcin	CALCA
P01266	Thyroglobulin	TG
P01270	Parathyroid hormone	PTH
P01275	Glucagon	GCG
P01282	Intestinal peptide PHM-27	VIP
P01286	Somatoliberin	GHRH
P01298	Pancreatic prohormone	PPY
P01303	C-flanking peptide of NPY	NPY
P01308	Insulin	INS
P01344	Insulin-like growth factor II	IGF2
P01350	Big gastrin	GAST

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P01374	Lymphotoxin-alpha	LTA
P01375	C-domain 1	TNF
P01562	Interferon alpha-1/13	IFNA1
P01563	Interferon alpha-2	IFNA2
P01566	Interferon alpha-10	IFNA10
P01567	Interferon alpha-7	IFNA7
P01568	Interferon alpha-21	IFNA21
P01569	Interferon alpha-5	IFNA5
P01570	Interferon alpha-14	IFNA14
P01571	Interferon alpha-17	IFNA17
P01574	Interferon beta	IFNB1
P01579	Interferon gamma	IFNG
P01583	Interleukin-1 alpha	IL1A
P01584	Interleukin-1 beta	IL1B
P01588	Erythropoietin	EPO
P01591	Immunoglobulin J chain	IGJ
P01732	T-cell surface glycoprotein CD8 alpha chain	CD8A
P01833	Polymeric immunoglobulin receptor	PIGR
P01857	Ig gamma-1 chain C region	IGHG1
P01859	Ig gamma-2 chain C region	IGHG2
P01860	Ig gamma-3 chain C region	IGHG3
P01861	Ig gamma-4 chain C region	IGHG4
P01871	Ig mu chain C region	IGHM
P01880	Ig delta chain C region	IGHD
P02452	Collagen alpha-1(I) chain	COL1A1
P02458	Chondrocalcin	COL2A1
P02461	Collagen alpha-1(III) chain	COL3A1
P02462	Collagen alpha-1(IV) chain	COL4A1
P02647	Apolipoprotein A-I	APOA1
P02649	Apolipoprotein E	APOE
P02652	Apolipoprotein A-II	APOA2
P02654	Apolipoprotein C-I	APOC1
P02655	Apolipoprotein C-II	APOC2
P02656	Apolipoprotein C-III	APOC3
P02671	Fibrinogen alpha chain	FGA
P02675	Fibrinopeptide B	FGB
P02679	Fibrinogen gamma chain	FGG
P02741	C-reactive protein	CRP
P02743	Serum amyloid P-component(1-203)	APCS
P02745	Complement C1q subcomponent subunit A	C1QA
P02746	Complement C1q subcomponent subunit B	C1QB
P02747	Complement C1q subcomponent subunit C	C1QC
P02748	Complement component C9b	C9
P02749	Beta-2-glycoprotein 1	APOH
P02750	Leucine-rich alpha-2-glycoprotein	LRG1
P02751	Ugl-Y2	FN1
P02753	Retinol-binding protein 4	RBP4
P02760	Trypstatin	AMBP
P02763	Alpha-1-acid glycoprotein 1	ORM1
P02765	Alpha-2-HS-glycoprotein chain A	AHSG
P02766	Transthyretin	TTR
P02768	Serum albumin	ALB
P02771	Alpha-fetoprotein	AFP
P02774	Vitamin D-binding protein	GC
P02775	Connective tissue-activating peptide III	PPBP
P02776	Platelet factor 4	PF4
P02778	CXCL10(1-73)	CXCL10
P02786	Transferrin receptor protein 1	TFRC
P02787	Serotransferrin	TF
P02788	Lactoferrin-C	LTF
P02790	Hemopexin	HPX
P02808	Statherin	STATH
P02810	Salivary acidic proline-rich phosphoprotein 1/2	PRH2
P02812	Basic salivary proline-rich protein 2	PRB2
P02814	Peptide D1A	SMR3B
P02818	Osteocalcin	BGLAP
P03950	Angiogenin	ANG
P03951	Coagulation factor XIa heavy chain	F11
P03952	Plasma kallikrein	KLKB1
P03956	27 kDa interstitial collagenase	MMP1
P03971	Muellerian-inhibiting factor	AMH
P03973	Antileukoprotease	SLPI
P04003	C4b-binding protein alpha chain	C4BPA
P04004	Somatomedin-B	VTN
P04054	Phospholipase A2	PLA2G1B
P04085	Platelet-derived growth factor subunit A	PDGFA

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P04090	Relaxin A chain	RLN2
P04114	Apolipoprotein B-100	APOB
P04118	Colipase	CLPS
P04141	Granulocyte-macrophage colony-stimulating factor	CSF2
P04155	Trefoil factor 1	TFF1
P04180	Phosphatidylcholine-sterol acyltransferase	LCAT
P04196	Histidine-rich glycoprotein	HRG
P04217	Alpha-1B-glycoprotein	A1BG
P04275	von Willebrand antigen 2	VWF
P04278	Sex hormone-binding globulin	SHBG
P04279	Alpha-inhibin-31	SEMG1
P04280	Basic salivary proline-rich protein 1	PRB1
P04628	Proto-oncogene Wnt-1	WNT1
P04745	Alpha-amylase 1	AMY1A
P04746	Pancreatic alpha-amylase	AMY2A
P04808	Prorelaxin H1	RLN1
P05000	Interferon omega-1	IFNW1
P05013	Interferon alpha-6	IFNA6
P05014	Interferon alpha-4	IFNA4
P05015	Interferon alpha-16	IFNA16
P05019	Insulin-like growth factor I	IGF1
P05060	GAWK peptide	CHGB
P05090	Apolipoprotein D	APOD
P05109	Protein S100-A8	S100A8
P05111	Inhibin alpha chain	INHA
P05112	Interleukin-4	IL4
P05113	Interleukin-5	IL5
P05120	Plasminogen activator inhibitor 2	SERPINB2
P05121	Plasminogen activator inhibitor 1	SERPINE1
P05154	Plasma serine protease inhibitor	SERPINA5
P05155	Plasma protease C1 inhibitor	SERPING1
P05156	Complement factor I heavy chain	CFI
P05160	Coagulation factor XIII B chain	F13B
P05161	Ubiquitin-like protein ISG15	ISG15
P05230	Fibroblast growth factor 1	FGF1
P05231	Interleukin-6	IL6
P05305	Big endothelin-1	EDN1
P05408	C-terminal peptide	SCG5
P05451	Lithostathine-1-alpha	REG1A
P05452	Tetranectin	CLEC3B
P05543	Thyroxine-binding globulin	SERPINA7
P05814	Beta-casein	CSN2
P05997	Collagen alpha-2(V) chain	COL5A2
P06276	Cholinesterase	BCHE
P06307	Cholecystokinin-12	CCK
P06396	Gelsolin	GSN
P06681	Complement C2	C2
P06702	Protein S100-A9	S100A9
P06727	Apolipoprotein A-IV	APOA4
P06734	Low affinity immunoglobulin epsilon Fc receptor soluble form	FCER2
P06744	Glucose-6-phosphate isomerase	GPI
P06850	Corticoliberin	CRH
P06858	Lipoprotein lipase	LPL
P06881	Calcitonin gene-related peptide 1	CALCA
P07093	Glia-derived nexin	SERPINE2
P07098	Gastric triacylglycerol lipase	LIPF
P07225	Vitamin K-dependent protein S	PROS1
P07237	Protein disulfide-isomerase	P4HB
P07288	Prostate-specific antigen	KLK3
P07306	Asialoglycoprotein receptor 1	ASGR1
P07355	Annexin A2	ANXA2
P07357	Complement component C8 alpha chain	C8A
P07358	Complement component C8 beta chain	C8B
P07360	Complement component C8 gamma chain	C8G
P07477	Alpha-trypsin chain 2	PRSS1
P07478	Trypsin-2	PRSS2
P07492	Neuromedin-C	GRP
P07498	Kappa-casein	CSN3
P07585	Decorin	DCN
P07911	Uromodulin	UMOD
P07942	Laminin subunit beta-1	LAMB1
P07988	Pulmonary surfactant-associated protein B	SFTPB
P07998	Ribonuclease pancreatic	RNASE1
P08118	Beta-microseminoprotein	MSMB

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P08123	Collagen alpha-2(I) chain	COL1A2
P08185	Corticosteroid-binding globulin	SERPINA6
P08217	Chymotrypsin-like elastase family member 2A	CELA2A
P08218	Chymotrypsin-like elastase family member 2B	CELA2B
P08253	72 kDa type IV collagenase	MMP2
P08254	Stromelysin-1	MMP3
P08294	Extracellular superoxide dismutase [Cu—Zn]	SOD3
P08476	Inhibin beta A chain	INHBA
P08493	Matrix Gla protein	MGP
P08572	Collagen alpha-2(IV) chain	COL4A2
P08581	Hepatocyte growth factor receptor	MET
P08603	Complement factor H	CFH
P08620	Fibroblast growth factor 4	FGF4
P08637	Low affinity immunoglobulin gamma Fc region receptor III-A	FCGR3A
P08697	Alpha-2-antiplasmin	SERPINF2
P08700	Interleukin-3	IL3
P08709	Coagulation factor VII	F7
P08833	Insulin-like growth factor-binding protein 1	IGFBP1
P08887	Interleukin-6 receptor subunit alpha	IL6R
P08949	Neuromedin-B-32	NMB
P08F94	Fibrocystin	PKHD1
P09038	Fibroblast growth factor 2	FGF2
P09228	Cystatin-SA	CST2
P09237	Matrilysin	MMP7
P09238	Stromelysin-2	MMP10
P09341	Growth-regulated alpha protein	CXCL1
P09382	Galectin-1	LGALS1
P09466	Glycodelin	PAEP
P09486	SPARC	SPARC
P09529	Inhibin beta B chain	INHBB
P09544	Protein Wnt-2	WNT2
P09603	Processed macrophage colony-stimulating factor 1	CSF1
P09681	Gastric inhibitory polypeptide	GIP
P09683	Secretin	SCT
P09919	Granulocyte colony-stimulating factor	CSF3
P0C091	FRAS1-related extracellular matrix protein 3	FREM3
P0C0L4	C4d-A	C4A
P0C0L5	Complement C4-B alpha chain	C4B
P0C0P6	Neuropeptide S	NPS
P0C7L1	Serine protease inhibitor Kazal-type 8	SPINK8
P0C862	Complement C1q and tumor necrosis factor-related protein 9A	C1QTNF9
P0C8F1	Prostate and testis expressed protein 4	PATE4
P0CG01	Gastrokine-3	GKN3P
P0CG36	Cryptic family protein 1B	CFC1B
P0CG37	Cryptic protein	CFC1
P0CJ68	Humanin-like protein 1	MTRNR2L1
P0CJ69	Humanin-like protein 2	MTRNR2L2
P0CJ70	Humanin-like protein 3	MTRNR2L3
P0CJ71	Humanin-like protein 4	MTRNR2L4
P0CJ72	Humanin-like protein 5	MTRNR2L5
P0CJ73	Humanin-like protein 6	MTRNR2L6
P0CJ74	Humanin-like protein 7	MTRNR2L7
P0CJ75	Humanin-like protein 8	MTRNR2L8
P0CJ76	Humanin-like protein 9	MTRNR2L9
P0CJ77	Humanin-like protein 10	MTRNR2L10
P0DJD7	Pepsin A-4	PGA4
P0DJD8	Pepsin A-3	PGA3
P0DJD9	Pepsin A-5	PGA5
P0DJI8	Amyloid protein A	SAA1
P0DJI9	Serum amyloid A-2 protein	SAA2
P10082	Peptide YY(3-36)	PYY
P10092	Calcitonin gene-related peptide 2	CALCB
P10124	Seryglycin	SRGN
P10145	MDNCF-a	IL8
P10147	MIP-1-alpha(4-69)	CCL3
P10163	Peptide P-D	PRB4
P10451	Osteopontin	SPP1
P10599	Thioredoxin	TXN
P10600	Transforming growth factor beta-3	TGFB3
P10643	Complement component C7	C7
P10645	Vasostatin-2	CHGA
P10646	Tissue factor pathway inhibitor	TFPI
P10720	Platelet factor 4 variant(4-74)	PF4V1

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P10745	Retinol-binding protein 3	RBP3
P10767	Fibroblast growth factor 6	FGF6
P10909	Clusterin alpha chain	CLU
P10912	Growth hormone receptor	GHR
P10915	Hyaluronan and proteoglycan link protein 1	HAPLN1
P10966	T-cell surface glycoprotein CD8 beta chain	CD8B
P10997	Islet amyloid polypeptide	IAPP
P11047	Laminin subunit gamma-1	LAMC1
P11150	Hepatic triacylglycerol lipase	LIPC
P11226	Mannose-binding protein C	MBL2
P11464	Pregnancy-specific beta-1-glycoprotein 1	PSG1
P11465	Pregnancy-specific beta-1-glycoprotein 2	PSG2
P11487	Fibroblast growth factor 3	FGF3
P11597	Cholesteryl ester transfer protein	CETP
P11684	Uteroglobin	SCGB1A1
P11686	Pulmonary surfactant-associated protein C	SFTPC
P12034	Fibroblast growth factor 5	FGF5
P12107	Collagen alpha-1(XI) chain	COL11A1
P12109	Collagen alpha-1(VI) chain	COL6A1
P12110	Collagen alpha-2(VI) chain	COL6A2
P12111	Collagen alpha-3(VI) chain	COL6A3
P12259	Coagulation factor V	F5
P12272	PTHrP[1-36]	PTH LH
P12273	Prolactin-inducible protein	PIP
P12544	Granzyme A	GZMA
P12643	Bone morphogenetic protein 2	BMP2
P12644	Bone morphogenetic protein 4	BMP4
P12645	Bone morphogenetic protein 3	BMP3
P12724	Eosinophil cationic protein	RNASE3
P12821	Angiotensin-converting enzyme, soluble form	ACE
P12838	Neutrophil defensin 4	DEFA4
P12872	Motilin	MLN
P13232	Interleukin-7	IL7
P13236	C-C motif chemokine 4	CCL4
P13284	Gamma-interferon-inducible lysosomal thiol reductase	IFI30
P13500	C-C motif chemokine 2	CCL2
P13501	C-C motif chemokine 5	CCL5
P13521	Secretogranin-2	SCG2
P13591	Neural cell adhesion molecule 1	NCAM1
P13611	Versican core protein	VCAN
P13671	Complement component C6	C6
P13688	Carcinoembryonic antigen-related cell adhesion molecule 1	CEACAM1
P13725	Oncostatin-M	OSM
P13726	Tissue factor	F3
P13727	Eosinophil granule major basic protein	PRG2
P13942	Collagen alpha-2(XI) chain	COL11A2
P13987	CD59 glycoprotein	CD59
P14138	Endothelin-3	EDN3
P14174	Macrophage migration inhibitory factor	MIF
P14207	Folate receptor beta	FOLR2
P14222	Perforin-1	PRF1
P14543	Nidogen-1	NID1
P14555	Phospholipase A2, membrane associated	PLA2G2A
P14625	Endoplasmic reticulum chaperone	HSP90B1
P14735	Insulin-degrading enzyme	IDE
P14778	Interleukin-1 receptor type 1, soluble form	IL1R1
P14780	82 kDa matrix metalloproteinase-9	MMP9
P15018	Leukemia inhibitory factor	LIF
P15085	Carboxypeptidase A1	CPA1
P15086	Carboxypeptidase B	CPB1
P15151	Poliovirus receptor	PVR
P15169	Carboxypeptidase N catalytic chain	CPN1
P15248	Interleukin-9	IL9
P15291	N-acetyllactosamine synthase	B4GALT1
P15309	PAPf39	ACPP
P15328	Folate receptor alpha	FOLR1
P15374	Ubiquitin carboxyl-terminal hydrolase isozyme L3	UCHL3
P15502	Elastin	ELN
P15509	Granulocyte-macrophage colony-stimulating factor receptor subunit alpha	CSF2RA
P15515	Histatin-1	HTN1
P15516	His3-(31-51)-peptide	HTN3
P15692	Vascular endothelial growth factor A	VEGFA

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P15814	Immunoglobulin lambda-like polypeptide 1	IgLL1
P15907	Beta-galactoside alpha-2,6-sialyltransferase 1	ST6GAL1
P15941	Mucin-1 subunit beta	MUC1
P16035	Metalloproteinase inhibitor 2	TIMP2
P16112	Aggrecan core protein 2	ACAN
P16233	Pancreatic triacylglycerol lipase	PNLIP
P16442	Histo-blood group ABO system transferase	ABO
P16471	Prolactin receptor	PRLR
P16562	Cysteine-rich secretory protein 2	CRISP2
P16619	C-C motif chemokine 3-like 1	CCL3L1
P16860	BNP(3-29)	NPPB
P16870	Carboxypeptidase E	CPE
P16871	Interleukin-7 receptor subunit alpha	IL7R
P17213	Bactericidal permeability-increasing protein	BPI
P17538	Chymotrypsinogen B	CTRB1
P17931	Galectin-3	LGALS3
P17936	Insulin-like growth factor-binding protein 3	IGFBP3
P17948	Vascular endothelial growth factor receptor 1	FLT1
P18065	Insulin-like growth factor-binding protein 2	IGFBP2
P18075	Bone morphogenetic protein 7	BMP7
P18428	Lipopolysaccharide-binding protein	LBP
P18509	PACAP-related peptide	ADCYAP1
P18510	Interleukin-1 receptor antagonist protein	IL1RN
P18827	Syndecan-1	SDC1
P19021	Peptidylglycine alpha-hydroxylating monooxygenase	PAM
P19235	Erythropoietin receptor	EPOR
P19438	Tumor necrosis factor-binding protein 1	TNFRSF1A
P19652	Alpha-1-acid glycoprotein 2	ORM2
P19801	Amiloride-sensitive amine oxidase [copper-containing]	ABP1
P19823	Inter-alpha-trypsin inhibitor heavy chain H2	ITIH2
P19827	Inter-alpha-trypsin inhibitor heavy chain H1	ITIH1
P19835	Bile salt-activated lipase	CEL
P19875	C-X-C motif chemokine 2	CXCL2
P19876	C-X-C motif chemokine 3	CXCL3
P19883	Follistatin	FST
P19957	Elafin	PI3
P19961	Alpha-amylase 2B	AMY2B
P20061	Transcobalamin-1	TCN1
P20062	Transcobalamin-2	TCN2
P20142	Gastricsin	PGC
P20155	Serine protease inhibitor Kazal-type 2	SPINK2
P20231	Tryptase beta-2	TPSB2
P20333	Tumor necrosis factor receptor superfamily member 1B	TNFRSF1B
P20366	Substance P	TAC1
P20382	Melanin-concentrating hormone	PMCH
P20396	Thyroliberin	TRH
P20742	Pregnancy zone protein	PZP
P20774	Mimecan	OGN
P20783	Neurotrophin-3	NTF3
P20800	Endothelin-2	EDN2
P20809	Interleukin-11	IL11
P20827	Ephrin-A1	EFNA1
P20849	Collagen alpha-1(IX) chain	COL9A1
P20851	C4b-binding protein beta chain	C4BPB
P20908	Collagen alpha-1(V) chain	COL5A1
P21128	Poly(U)-specific endoribonuclease	ENDOU
P21246	Pleiotrophin	PTN
P21583	Kit ligand	KITLG
P21741	Midkine	MDK
P21754	Zona pellucida sperm-binding protein 3	ZP3
P21781	Fibroblast growth factor 7	FGF7
P21802	Fibroblast growth factor receptor 2	FGFR2
P21810	Biglycan	BGN
P21815	Bone sialoprotein 2	IBSP
P21860	Receptor tyrosine-protein kinase erbB-3	ERBB3
P21941	Cartilage matrix protein	MATN1
P22003	Bone morphogenetic protein 5	BMP5
P22004	Bone morphogenetic protein 6	BMP6
P22079	Lactoperoxidase	LPO
P22105	Tenascin-X	TNXB
P22301	Interleukin-10	IL10
P22303	Acetylcholinesterase	ACHE
P22352	Glutathione peroxidase 3	GPX3

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P22362	C-C motif chemokine 1	CCL1
P22455	Fibroblast growth factor receptor 4	FGFR4
P22466	Galanin message-associated peptide	GAL
P22692	Insulin-like growth factor-binding protein 4	IGFBP4
P22749	Granulysin	GNLY
P22792	Carboxypeptidase N subunit 2	CPN2
P22891	Vitamin K-dependent protein Z	PROZ
P22894	Neutrophil collagenase	MMP8
P23142	Fibulin-1	FBLN1
P23280	Carbonic anhydrase 6	CA6
P23352	Anosmin-1	KAL1
P23435	Cerebellin-1	CBLN1
P23560	Brain-derived neurotrophic factor	BDNF
P23582	C-type natriuretic peptide	NPPC
P23946	Chymase	CMA1
P24043	Laminin subunit alpha-2	LAMA2
P24071	Immunoglobulin alpha Fc receptor	FCAR
P24347	Stromelysin-3	MMP11
P24387	Corticotropin-releasing factor-binding protein	CRHBP
P24592	Insulin-like growth factor-binding protein 6	IGFBP6
P24593	Insulin-like growth factor-binding protein 5	IGFBP5
P24821	Tenascin	TNC
P24855	Deoxyribonuclease-1	DNASE1
P25067	Collagen alpha-2(VIII) chain	COL8A2
P25311	Zinc-alpha-2-glycoprotein	AZGP1
P25391	Laminin subunit alpha-1	LAMA1
P25445	Tumor necrosis factor receptor superfamily member 6	FAS
P25940	Collagen alpha-3(V) chain	COL5A3
P25942	Tumor necrosis factor receptor superfamily member 5	CD40
P26022	Pentraxin-related protein PTX3	PTX3
P26927	Hepatocyte growth factor-like protein beta chain	MST1
P27169	Serum paraoxonase/arylesterase 1	PON1
P27352	Gastric intrinsic factor	GIF
P27487	Dipeptidyl peptidase 4 membrane form	DPP4
P27539	Embryonic growth/differentiation factor 1	GDF1
P27658	Vastatin	COL8A1
P27797	Calreticulin	CALR
P27918	Properdin	CFP
P28039	Acyloxyacyl hydrolase	AOAH
P28300	Protein-lysine 6-oxidase	LOX
P28325	Cystatin-D	CST5
P28799	Granulin-1	GRN
P29122	Proprotein convertase subtilisin/kexin type 6	PCSK6
P29279	Connective tissue growth factor	CTGF
P29320	Ephrin type-A receptor 3	EPHA3
P29400	Collagen alpha-5(IV) chain	COL4A5
P29459	Interleukin-12 subunit alpha	IL12A
P29460	Interleukin-12 subunit beta	IL12B
P29508	Serpin B3	SERPINB3
P29622	Kallistatin	SERPINA4
P29965	CD40 ligand, soluble form	CD40LG
P30990	Neurotensin/neuromedin N	NTS
P31025	Lipocalin-1	LCN1
P31151	Protein S100-A7	S100A7
P31371	Fibroblast growth factor 9	FGF9
P31431	Syndecan-4	SDC4
P31947	14-3-3 protein sigma	SFN
P32455	Interferon-induced guanylate-binding protein 1	GBP1
P32881	Interferon alpha-8	IFNA8
P34096	Ribonuclease 4	RNASE4
P34130	Neurotrophin-4	NTF4
P34820	Bone morphogenetic protein 8B	BMP8B
P35030	Trypsin-3	PRSS3
P35052	Secreted glypican-1	GPC1
P35070	Betacellulin	BTC
P35225	Interleukin-13	IL13
P35247	Pulmonary surfactant-associated protein D	SFTPD
P35318	ADM	ADM
P35542	Serum amyloid A-4 protein	SAA4
P35555	Fibrillin-1	FBN1
P35556	Fibrillin-2	FBN2
P35625	Metalloproteinase inhibitor 3	TIMP3

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P35858	Insulin-like growth factor-binding protein complex acid labile subunit	IGFALS
P35916	Vascular endothelial growth factor receptor 3	FLT4
P35968	Vascular endothelial growth factor receptor 2	KDR
P36222	Chitinase-3-like protein 1	CHI3L1
P36952	Serpine B5	SERPINE5
P36955	Pigment epithelium-derived factor	SERPINF1
P36980	Complement factor H-related protein 2	CFHR2
P39059	Collagen alpha-1(XV) chain	COL15A1
P39060	Collagen alpha-1(XVIII) chain	COL18A1
P39877	Calcium-dependent phospholipase A2	PLA2G5
P39900	Macrophage metalloelastase	MMP12
P39905	Glial cell line-derived neurotrophic factor	GDNF
P40225	Thrombopoietin	THPO
P40967	M-alpha	PMEL
P41159	Leptin	LEP
P41221	Protein Wnt-5a	WNT5A
P41222	Prostaglandin-H2 D-isomerase	PTGDS
P41271	Neuroblastoma suppressor of tumorigenicity 1	NBL1
P41439	Folate receptor gamma	FOLR3
P42127	Agouti-signaling protein	ASIP
P42702	Leukemia inhibitory factor receptor	LIFR
P42830	ENA-78(9-78)	CXCL5
P43026	Growth/differentiation factor 5	GDF5
P43251	Biotinidase	BTD
P43652	Afamin	AFM
P45452	Collagenase 3	MMP13
P47710	Casoxin-D	CSN1S1
P47929	Galectin-7	LGALS7B
P47972	Neuronal pentraxin-2	NPTX2
P47989	Xanthine oxidase	XDH
P47992	Lymphotactin	XCL1
P48023	Tumor necrosis factor ligand superfamily member 6, membrane form	FASLG
P48052	Carboxypeptidase A2	CPA2
P48061	Stromal cell-derived factor 1	CXCL12
P48304	Lithostathine-1-beta	REG1B
P48307	Tissue factor pathway inhibitor 2	TFPI2
P48357	Leptin receptor	LEPR
P48594	Serpine B4	SERPINE4
P48645	Neuromedin-U-25	NMU
P48740	Mannan-binding lectin serine protease 1	MASP1
P48745	Protein NOV homolog	NOV
P48960	CD97 antigen subunit beta	CD97
P49223	Kunitz-type protease inhibitor 3	SPINT3
P49747	Cartilage oligomeric matrix protein	COMP
P49763	Placenta growth factor	PGF
P49765	Vascular endothelial growth factor B	VEGFB
P49767	Vascular endothelial growth factor C	VEGFC
P49771	Fms-related tyrosine kinase 3 ligand	FLT3LG
P49862	Kallikrein-7	KLK7
P49863	Granzyme K	GZMK
P49908	Selenoprotein P	SEPP1
P49913	Antibacterial protein FALL-39	CAMP
P50607	Tubby protein homolog	TUB
P51124	Granzyme M	GZMM
P51512	Matrix metalloproteinase-16	MMP16
P51654	Glypican-3	GPC3
P51671	Eotaxin	CCL11
P51884	Lumican	LUM
P51888	Prolargin	PRELP
P52798	Ephrin-A4	EFNA4
P52823	Stanniocalcin-1	STC1
P53420	Collagen alpha-4(IV) chain	COL4A4
P53621	Coatamer subunit alpha	COPA
P54108	Cysteine-rich secretory protein 3	CRISP3
P54315	Pancreatic lipase-related protein 1	PNLIPRP1
P54317	Pancreatic lipase-related protein 2	PNLIPRP2
P54793	Arylsulfatase F	ARSF
P55000	Secreted Ly-6/uPAR-related protein 1	SLURP1
P55001	Microfibrillar-associated protein 2	MFAP2
P55056	Apolipoprotein C-IV	APOC4
P55058	Phospholipid transfer protein	PLTP
P55075	Fibroblast growth factor 8	FGF8
P55081	Microfibrillar-associated protein 1	MFAP1
P55083	Microfibrillar-associated glycoprotein 4	MFAP4

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P55107	Bone morphogenetic protein 3B	GDF10
P55145	Mesencephalic astrocyte-derived neurotrophic factor	MANF
P55259	Pancreatic secretory granule membrane major glycoprotein GP2	GP2
P55268	Laminin subunit beta-2	LAMB2
P55773	CCL23(30-99)	CCL23
P55774	C-C motif chemokine 18	CCL18
P55789	FAD-linked sulfhydryl oxidase ALR	GFER
P56703	Proto-oncogene Wnt-3	WNT3
P56704	Protein Wnt-3a	WNT3A
P56705	Protein Wnt-4	WNT4
P56706	Protein Wnt-7b	WNT7B
P56730	Neurotrypsin	PRSS12
P56851	Epididymal secretory protein E3-beta	EDDM3B
P56975	Neuregulin-3	NRG3
P58062	Serine protease inhibitor Kazal-type 7	SPINK7
P58215	Lysyl oxidase homolog 3	LOXL3
P58294	Prokineticin-1	PROK1
P58335	Anthrax toxin receptor 2	ANTXR2
P58397	A disintegrin and metalloproteinase with thrombospondin motifs 12	ADAMTS12
P58417	Neurexophilin-1	NXPH1
P58499	Protein FAM3B	FAM3B
P59510	A disintegrin and metalloproteinase with thrombospondin motifs 20	ADAMTS20
P59665	Neutrophil defensin 1	DEFA1B
P59666	Neutrophil defensin 3	DEFA3
P59796	Glutathione peroxidase 6	GPX6
P59826	BPI fold-containing family B member 3	BPIFB3
P59827	BPI fold-containing family B member 4	BPIFB4
P59861	Beta-defensin 131	DEFB131
P60022	Beta-defensin 1	DEFB1
P60153	Inactive ribonuclease-like protein 9	RNASE9
P60827	Complement C1q tumor necrosis factor-related protein 8	C1QTNF8
P60852	Zona pellucida sperm-binding protein 1	ZP1
P60985	Keratinocyte differentiation-associated protein	KRTDAP
P61109	Kidney androgen-regulated protein	KAP
P61278	Somatostatin-14	SST
P61366	Osteocrin	OSTN
P61626	Lysozyme C	LYZ
P61769	Beta-2-microglobulin	B2M
P61812	Transforming growth factor beta-2	TGFB2
P61916	Epididymal secretory protein E1	NPC2
P62502	Epididymal-specific lipocalin-6	LCN6
P62937	Peptidyl-prolyl cis-trans isomerase A	PPIA
P67809	Nuclease-sensitive element-binding protein 1	YBX1
P67812	Signal peptidase complex catalytic subunit SEC11A	SEC11A
P78310	Coxsackievirus and adenovirus receptor	CXADR
P78333	Secreted glypican-5	GPC5
P78380	Oxidized low-density lipoprotein receptor 1	OLR1
P78423	Processed fractalkine	CX3CL1
P78509	Reelin	RELN
P78556	CCL20(2-70)	CCL20
P80075	MCP-2(6-76)	CCL8
P80098	C-C motif chemokine 7	CCL7
P80108	Phosphatidylinositol-glycan-specific phospholipase D	GPLD1
P80162	C-X-C motif chemokine 6	CXCL6
P80188	Neutrophil gelatinase-associated lipocalin	LCN2
P80303	Nucleobindin-2	NUCB2
P80511	Calcitermin	S100A12
P81172	Hepcidin-25	HAMP
P81277	Prolactin-releasing peptide	PRLH
P81534	Beta-defensin 103	DEFB103A
P81605	Dermcidin	DCD
P82279	Protein crumbs homolog 1	CRB1
P82987	ADAMTS-like protein 3	ADAMTSL3
P83105	Serine protease HTRA4	HTRA4
P83110	Serine protease HTRA3	HTRA3
P83859	Orexigenic neuropeptide QRFP	QRFP
P98088	Mucin-5AC	MUC5AC
P98095	Fibulin-2	FBLN2

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
P98160	Basement membrane-specific heparan sulfate proteoglycan core protein	HSPG2
P98173	Protein FAM3A	FAM3A
Q00604	Norrin	NDP
Q00796	Sorbitol dehydrogenase	SORD
Q00887	Pregnancy-specific beta-1-glycoprotein 9	PSG9
Q00888	Pregnancy-specific beta-1-glycoprotein 4	PSG4
Q00889	Pregnancy-specific beta-1-glycoprotein 6	PSG6
Q01523	HD5(56-94)	DEFA5
Q01524	Defensin-6	DEFA6
Q01955	Collagen alpha-3(IV) chain	COL4A3
Q02297	Pro-neuregulin-1, membrane-bound isoform	NRG1
Q02325	Plasminogen-like protein B	PLGLB1
Q02383	Semenogelin-2	SEMG2
Q02388	Collagen alpha-1(VII) chain	COL7A1
Q02505	Mucin-3A	MUC3A
Q02509	Otoconin-90	OC90
Q02747	Guanylin	GUCA2A
Q02763	Angiopoietin-1 receptor	TEK
Q02817	Mucin-2	MUC2
Q02985	Complement factor H-related protein 3	CFHR3
Q03167	Transforming growth factor beta receptor type 3	TGFBR3
Q03403	Trefoil factor 2	TFF2
Q03405	Urokinase plasminogen activator surface receptor	PLAUR
Q03591	Complement factor H-related protein 1	CFHR1
Q03692	Collagen alpha-1(X) chain	COL10A1
Q04118	Basic salivary proline-rich protein 3	PRB3
Q04756	Hepatocyte growth factor activator short chain	HGFAC
Q04900	Sialomucin core protein 24	CD164
Q05315	Eosinophil lysophospholipase	CLC
Q05707	Collagen alpha-1(XIV) chain	COL14A1
Q05996	Processed zona pellucida sperm-binding protein 2	ZP2
Q06033	Inter-alpha-trypsin inhibitor heavy chain H3	ITIH3
Q06141	Regenerating islet-derived protein 3-alpha	REG3A
Q06828	Fibromodulin	FMOD
Q07092	Collagen alpha-1(XVI) chain	COL16A1
Q07325	C-X-C motif chemokine 9	CXCL9
Q07507	Dermatopontin	DPT
Q075Z2	Binder of sperm protein homolog 1	BSPH1
Q07654	Trefoil factor 3	TFF3
Q07699	Sodium channel subunit beta-1	SCN1B
Q08345	Epithelial discoidin domain-containing receptor 1	DDR1
Q08380	Galectin-3-binding protein	LGALS3BP
Q08397	Lysyl oxidase homolog 1	LOXL1
Q08431	Lactadherin	MFG8
Q08629	Testican-1	SPOCK1
Q08648	Sperm-associated antigen 11B	SPAG11B
Q08830	Fibrinogen-like protein 1	FGL1
Q10471	Polypeptide N-acetylgalactosaminyltransferase 2	GALNT2
Q10472	Polypeptide N-acetylgalactosaminyltransferase 1	GALNT1
Q11201	CMP-N-acetylneuramate-beta-galactosamide-alpha-2,3-sialyltransferase 1	ST3GAL1
Q11203	CMP-N-acetylneuramate-beta-1,4-galactoside alpha-2,3-sialyltransferase	ST3GAL3
Q11206	CMP-N-acetylneuramate-beta-galactosamide-alpha-2,3-sialyltransferase 4	ST3GAL4
Q12794	Hyaluronidase-1	HYAL1
Q12805	EGF-containing fibulin-like extracellular matrix protein 1	EFEMP1
Q12836	Zona pellucida sperm-binding protein 4	ZP4
Q12841	Follistatin-related protein 1	FSTL1
Q12904	Aminoacyl tRNA synthase complex-interacting multifunctional protein 1	AIMP1
Q13018	Soluble secretory phospholipase A2 receptor	PLA2R1
Q13072	B melanoma antigen 1	BAGE
Q13093	Platelet-activating factor acetylhydrolase	PLA2G7
Q13103	Secreted phosphoprotein 24	SPP2
Q13162	Peroxiredoxin-4	PRDX4
Q13201	Platelet glycoprotein Ia*	MMRN1
Q13214	Semaphorin-3B	SEMA3B

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q13219	Pappalysin-1	PAPPA
Q13231	Chitotriosidase-1	CHIT1
Q13253	Noggin	NOG
Q13261	Interleukin-15 receptor subunit alpha	IL15RA
Q13275	Semaphorin-3F	SEMA3F
Q13291	Signaling lymphocytic activation molecule	SLAMF1
Q13316	Dentin matrix acidic phosphoprotein 1	DMP1
Q13361	Microfibrillar-associated protein 5	MFAP5
Q13410	Butyrophilin subfamily 1 member A1	BTN1A1
Q13421	Mesothelin, cleaved form	MSLN
Q13429	Insulin-like growth factor I	IGF-I
Q13443	Disintegrin and metalloproteinase domain-containing protein 9	ADAM9
Q13519	Neuropeptide 1	PNOC
Q13751	Laminin subunit beta-3	LAMB3
Q13753	Laminin subunit gamma-2	LAMC2
Q13790	Apolipoprotein F	APOF
Q13822	Ectonucleotide pyrophosphatase/phosphodiesterase family member 2	ENPP2
Q14031	Collagen alpha-6(IV) chain	COL4A6
Q14050	Collagen alpha-3(IX) chain	COL9A3
Q14055	Collagen alpha-2(IX) chain	COL9A2
Q14112	Nidogen-2	NID2
Q14114	Low-density lipoprotein receptor-related protein 8	LRP8
Q14118	Dystroglycan	DAG1
Q14314	Fibroleukin	FGL2
Q14393	Growth arrest-specific protein 6	GAS6
Q14406	Chorionic somatomammotropin hormone-like 1	CSHL1
Q14507	Epididymal secretory protein E3-alpha	EDDM3A
Q14508	WAP four-disulfide core domain protein 2	WFDC2
Q14512	Fibroblast growth factor-binding protein 1	FGFBP1
Q14515	SPARC-like protein 1	SPARCL1
Q14520	Hyaluronan-binding protein 2 27 kDa light chain	HABP2
Q14563	Semaphorin-3A	SEMA3A
Q14623	Indian hedgehog protein	IHH
Q14624	Inter-alpha-trypsin inhibitor heavy chain H4	ITIH4
Q14667	UPF0378 protein KIAA0100	KIAA0100
Q14703	Membrane-bound transcription factor site-1 protease	MBTPS1
Q14766	Latent-transforming growth factor beta-binding protein 1	LTBP1
Q14767	Latent-transforming growth factor beta-binding protein 2	LTBP2
Q14773	Intercellular adhesion molecule 4	ICAM4
Q14993	Collagen alpha-1(XIX) chain	COL19A1
Q14CN2	Calcium-activated chloride channel regulator 4, 110 kDa form	CLCA4
Q15046	Lysine--tRNA ligase	KARS
Q15063	Periostin	POSTN
Q15109	Advanced glycosylation end product-specific receptor	AGER
Q15113	Procollagen C-endopeptidase enhancer 1	PCOLCE
Q15166	Serum paraoxonase/lactonase 3	PON3
Q15195	Plasminogen-like protein A	PLGLA
Q15198	Platelet-derived growth factor receptor-like protein	PDGFRL
Q15223	Poliovirus receptor-related protein 1	PVRL1
Q15238	Pregnancy-specific beta-1-glycoprotein 5	PSG5
Q15363	Transmembrane emp24 domain-containing protein 2	TMED2
Q15375	Ephrin type-A receptor 7	EPHA7
Q15389	Angiopoietin-1	ANGPT1
Q15465	Sonic hedgehog protein	SHH
Q15485	Ficolin-2	FCN2
Q15517	Corneodesmosin	CDSN
Q15582	Transforming growth factor-beta-induced protein ig-h3	TGFBI
Q15661	Tryptase alpha/beta-1	TPSAB1
Q15726	Metastin	KISS1
Q15782	Chitinase-3-like protein 2	CHI3L2
Q15828	Cystatin-M	CST6
Q15846	Clusterin-like protein 1	CLUL1
Q15848	Adiponectin	ADIPOQ

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q16206	Protein disulfide-thiol oxidoreductase	ENOX2
Q16270	Insulin-like growth factor-binding protein 7	IGFBP7
Q16363	Laminin subunit alpha-4	LAMA4
Q16378	Proline-rich protein 4	PRR4
Q16557	Pregnancy-specific beta-1-glycoprotein 3	PSG3
Q16568	CART(42-89)	CARTPT
Q16610	Extracellular matrix protein 1	ECM1
Q16619	Cardiotrophin-1	CTF1
Q16623	Syntaxin-1A	STX1A
Q16627	HCC-1(9-74)	CCL14
Q16651	Prostasin light chain	PRSS8
Q16661	Guanylate cyclase C-activating peptide 2	GUCA2B
Q16663	CCL15(29-92)	CCL15
Q16674	Melanoma-derived growth regulatory protein	MIA
Q16769	Glutaminyl-peptide cyclotransferase	QPCT
Q16787	Laminin subunit alpha-3	LAMA3
Q16842	CMP-N-acetylneuraminate-beta-galactosamide-alpha-2,3-sialyltransferase 2	ST3GAL2
Q17RR3	Pancreatic lipase-related protein 3	PNLIPRP3
Q17RW2	Collagen alpha-1(XXIV) chain	COL24A1
Q17RY6	Lymphocyte antigen 6K	LY6K
Q1L6U9	Prostate-associated microseminoprotein	MSMP
Q1W4C9	Serine protease inhibitor Kazal-type 13	SPINK13
Q1ZYL8	Izumo sperm-egg fusion protein 4	IZUMO4
Q29960	HLA class I histocompatibility antigen, Cw-16 alpha chain	HLA-C
Q2I0M5	R-spondin-4	RSPO4
Q2L4Q9	Serine protease 53	PRSS53
Q2MKA7	R-spondin-1	RSPO1
Q2MV58	Tectonic-1	TCTN1
Q2TAL6	Brorin	VWC2
Q2UY09	Collagen alpha-1(XXVIII) chain	COL28A1
Q2VPA4	Complement component receptor 1-like protein	CR1L
Q2WEN9	Carcinoembryonic antigen-related cell adhesion molecule 16	CEACAM16
Q30KP8	Beta-defensin 136	DEFB136
Q30KP9	Beta-defensin 135	DEFB135
Q30KQ1	Beta-defensin 133	DEFB133
Q30KQ2	Beta-defensin 130	DEFB130
Q30KQ4	Beta-defensin 116	DEFB116
Q30KQ5	Beta-defensin 115	DEFB115
Q30KQ6	Beta-defensin 114	DEFB114
Q30KQ7	Beta-defensin 113	DEFB113
Q30KQ8	Beta-defensin 112	DEFB112
Q30KQ9	Beta-defensin 110	DEFB110
Q30KR1	Beta-defensin 109	DEFB109P1
Q32P28	Prolyl 3-hydroxylase 1	LEPRE1
Q3B7J2	Glucose-fructose oxidoreductase domain-containing protein 2	GFOD2
Q3SY79	Protein Wnt	WNT3A
Q3T906	N-acetylglucosamine-1-phosphotransferase subunits alpha/beta	GNPTAB
Q495T6	Membrane metallo-endopeptidase-like 1	MMEL1
Q49AH0	Cerebral dopamine neurotrophic factor	CDNF
Q4G0G5	Secretoglobin family 2B member 2	SCGB2B2
Q4G0M1	Protein FAM132B	FAM132B
Q4LDE5	Sushi, von Willebrand factor type A, EGF and pentraxin domain-containing protein 1	SVEP1
Q4QY38	Beta-defensin 134	DEFB134
Q4VAJ4	Protein Wnt	WNT10B
Q4W5P6	Protein TMEM155	TMEM155
Q4ZHG4	Fibronectin type III domain-containing protein 1	FNDC1
Q53H76	Phospholipase A1 member A	PLA1A
Q53RD9	Fibulin-7	FBLN7
Q53S33	BOLA-like protein 3	BOLA3
Q5BLP8	Neuropeptide-like protein C4orf48	C4orf48
Q5DT21	Serine protease inhibitor Kazal-type 9	SPINK9
Q5EBL8	PDZ domain-containing protein 11	PDZD11
Q5FYB0	Arylsulfatase J	ARSJ
Q5FYB1	Arylsulfatase I	ARSI
Q5GAN3	Ribonuclease-like protein 13	RNASE13
Q5GAN4	Ribonuclease-like protein 12	RNASE12
Q5GAN6	Ribonuclease-like protein 10	RNASE10

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q5GFL6	von Willebrand factor A domain-containing protein 2	VWA2
Q5H8A3	Neuromedin-S	NMS
Q5H8C1	FRAS1-related extracellular matrix protein 1	FREM1
Q5IJ48	Protein crumbs homolog 2	CRB2
Q5J5C9	Beta-defensin 121	DEFB121
Q5JS37	NHL repeat-containing protein 3	NHLRC3
Q5JTB6	Placenta-specific protein 9	PLAC9
Q5JU69	Torsin-2A	TOR2A
Q5JXM2	Methyltransferase-like protein 24	METTL24
Q5JZY3	Ephrin type-A receptor 10	EPHA10
Q5K4E3	Polyserase-2	PRSS36
Q5SRR4	Lymphocyte antigen 6 complex locus protein G5c	LY6G5C
Q5T1H1	Protein eyes shut homolog	EYS
Q5T4F7	Secreted frizzled-related protein 5	SFRP5
Q5T4W7	Artemin	ARTN
Q5T7M4	Protein FAM132A	FAM132A
Q5TEH8	Protein Wnt	WNT2B
Q5TIE3	von Willebrand factor A domain-containing protein 5B1	VWA5B1
Q5UCC4	ER membrane protein complex subunit 10	EMC10
Q5VST6	Abhydrolase domain-containing protein FAM108B1	FAM108B1
Q5VTL7	Fibronectin type III domain-containing protein 7	FNDC7
Q5VUM1	UPF0369 protein C6orf57	C6orf57
Q5VV43	Dyslexia-associated protein KIAA0319	KIAA0319
Q5VWW1	Complement C1q-like protein 3	C1QL3
Q5VXI9	Lipase member N	LIPN
Q5VXJ0	Lipase member K	LIPK
Q5VXM1	CUB domain-containing protein 2	CDCP2
Q5VYX0	Renalase	RNLS
Q5VYY2	Lipase member M	LIPM
Q5W186	Cystatin-9	CST9
Q5W5W9	Regulated endocrine-specific protein 18	RESP18
Q5XG92	Carboxylesterase 4A	CES4A
Q63HQ2	Pikachurin	EGFLAM
Q641Q3	Meteorin-like protein	METRNL
Q66K79	Carboxypeptidase Z	CPZ
Q685J3	Mucin-17	MUC17
Q68BL7	Olfactomedin-like protein 2A	OLFML2A
Q68BL8	Olfactomedin-like protein 2B	OLFML2B
Q68DV7	E3 ubiquitin-protein ligase RNF43	RNF43
Q6B9Z1	Insulin growth factor-like family member 4	IGFL4
Q6BAA4	Fc receptor-like B	FCRLB
Q6E0U4	Dermokine	DMKN
Q6EMK4	Vasorin	VASN
Q6FHJ7	Secreted frizzled-related protein 4	SFRP4
Q6GPI1	Chymotrypsin B2 chain B	CTRB2
Q6GTS8	Probable carboxypeptidase PM20D1	PM20D1
Q6H9L7	Isthmin-2	ISM2
Q6IE36	Ovostatin homolog 2	OVOS2
Q6IE37	Ovostatin homolog 1	OVOS1
Q6IE38	Serine protease inhibitor Kazal-type 14	SPINK14
Q6ISS4	Leukocyte-associated immunoglobulin-like receptor 2	LAIR2
Q6JVE5	Epididymal-specific lipocalin-12	LCN12
Q6JVE6	Epididymal-specific lipocalin-10	LCN10
Q6JVE9	Epididymal-specific lipocalin-8	LCN8
Q6KF10	Growth/differentiation factor 6	GDF6
Q6MZW2	Follistatin-related protein 4	FSTL4
Q6NSX1	Coiled-coil domain-containing protein 70	CCDC70
Q6NT32	Carboxylesterase 5A	CES5A
Q6NT52	Choriogonadotropin subunit beta variant 2	CGB2
Q6NUI6	Chondroadherin-like protein	CHADL
Q6NUJ1	Saposin A-like	PSAPL1
Q6P093	Arylacetamide deacetylase-like 2	AADA2L2
Q6P4A8	Phospholipase B-like 1	PLBD1
Q6P5S2	UPF0762 protein C6orf58	C6orf58
Q6P988	Protein notum homolog	NOTUM
Q6PCB0	von Willebrand factor A domain-containing protein 1	VWA1
Q6PDA7	Sperm-associated antigen 11A	SPAG11A
Q6PEW0	Inactive serine protease 54	PRSS54
Q6PEZ8	Podocan-like protein 1	PODNL1

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q6PKH6	Dehydrogenase/reductase SDR family member 4-like 2	DHRS4L2
Q6Q788	Apolipoprotein A-V	APOA5
Q6SPF0	Atherin	SAMD1
Q6UDR6	Kunitz-type protease inhibitor 4	SPINT4
Q6URK8	Testis, prostate and placenta-expressed protein	TEPP
Q6UW01	Cerebellin-3	CBLN3
Q6UW10	Surfactant-associated protein 2	SFTA2
Q6UW15	Regenerating islet-derived protein 3-gamma	REG3G
Q6UW32	Insulin growth factor-like family member 1	IGFL1
Q6UW78	UPF0723 protein C11orf83	C11orf83
Q6UW88	Epigen	EPGN
Q6UWE3	Colipase-like protein 2	CLPSL2
Q6UWF7	NXPE family member 4	NXPE4
Q6UWF9	Protein FAM180A	FAM180A
Q6UWM5	GLIPR1-like protein 1	GLIPR1L1
Q6UWN8	Serine protease inhibitor Kazal-type 6	SPINK6
Q6UWP2	Dehydrogenase/reductase SDR family member 11	DHRS11
Q6UWP8	Suprabasin	SBSN
Q6UWQ5	Lysozyme-like protein 1	LYZL1
Q6UWQ7	Insulin growth factor-like family member 2	IGFL2
Q6UWR7	Ectonucleotide pyrophosphatase/ phosphodiesterase family member 6 soluble form	ENPP6
Q6UWT2	Adropin	ENHO
Q6UWU2	Beta-galactosidase-1-like protein	GLB1L
Q6UWW0	Lipocalin-15	LCN15
Q6UWX4	HHIP-like protein 2	HHIPL2
Q6UWY0	Arylsulfatase K	ARSK
Q6UWY2	Serine protease 57	PRSS57
Q6UWY5	Olfactomedin-like protein 1	OLFML1
Q6UX06	Olfactomedin-4	OLFM4
Q6UX07	Dehydrogenase/reductase SDR family member 13	DHRS13
Q6UX39	Amelotin	AMTN
Q6UX46	Protein FAM150B	FAM150B
Q6UX73	UPF0764 protein C16orf89	C16orf89
Q6UXB0	Protein FAM131A	FAM131A
Q6UXB1	Insulin growth factor-like family member 3	IGFL3
Q6UXB2	VEGF co-regulated chemokine 1	CXCL17
Q6UXF7	C-type lectin domain family 18 member B	CLEC18B
Q6UXH0	Hepatocellular carcinoma-associated protein TD26	C19orf80
Q6UXH1	Cysteine-rich with EGF-like domain protein 2	CRELD2
Q6UXH8	Collagen and calcium-binding EGF domain-containing protein 1	CCBE1
Q6UXH9	Inactive serine protease PAMR1	PAMR1
Q6UXI7	Vitrin	VIT
Q6UXI9	Nephronectin	NPNT
Q6UXN2	Trem-like transcript 4 protein	TREML4
Q6UXS0	C-type lectin domain family 19 member A	CLEC19A
Q6UXT8	Protein FAM150A	FAM150A
Q6UXT9	Abhydrolase domain-containing protein 15	ABHD15
Q6UXV4	Apolipoprotein O-like	APOOL
Q6UXX5	Inter-alpha-trypsin inhibitor heavy chain H6	ITIH6
Q6UXX9	R-spondin-2	RSPO2
Q6UY14	ADAMTS-like protein 4	ADAMTSL4
Q6UY27	Prostate and testis expressed protein 2	PATE2
Q6W4X9	Mucin-6	MUC6
Q6WN34	Chordin-like protein 2	CHRD12
Q6WRI0	Immunoglobulin superfamily member 10	IGSF10
Q6X4U4	Sclerostin domain-containing protein 1	SOSTDC1
Q6X784	Zona pellucida-binding protein 2	ZPBP2
Q6XE38	Secretoglobin family 1D member 4	SCGB1D4
Q6XPR3	Repetin	RPTN
Q6XZB0	Lipase member I	LIP1
Q6ZMM2	ADAMTS-like protein 5	ADAMTSL5
Q6ZMP0	Thrombospondin type-1 domain-containing protein 4	THSD4
Q6ZNF0	Iron/zinc purple acid phosphatase-like protein	PAPL
Q6ZRI0	Otogelin	OTOG
Q6ZRP7	Sulfhydryl oxidase 2	QSOX2
Q6ZWJ8	Kielin/chordin-like protein	KCP
Q75N90	Fibrillin-3	FBN3
Q765I0	Urotensin-2B	UTS2D
Q76B58	Protein FAM5C	FAM5C

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q76LX8	A disintegrin and metalloproteinase with thrombospondin motifs 13	ADAMTS13
Q76M96	Coiled-coil domain-containing protein 80	CCDC80
Q7L1S5	Carbohydrate sulfotransferase 9	CHST9
Q7L513	Fc receptor-like A	FCRLA
Q7L8A9	Vasohibin-1	VASH1
Q7RTM1	Otopetrin-1	OTOP1
Q7RTW8	Otoancorin	OTOA
Q7RTY5	Serine protease 48	PRSS48
Q7RTY7	Ovochymase-1	OVCH1
Q7RTZ1	Ovochymase-2	OVCH2
Q7Z304	MAM domain-containing protein 2	MAMDC2
Q7Z3S9	Notch homolog 2 N-terminal-like protein	NOTCH2NL
Q7Z4H4	Intermedin-short	ADM2
Q7Z4P5	Growth/differentiation factor 7	GDF7
Q7Z4R8	UPF0669 protein C6orf120	C6orf120
Q7Z4W2	Lysozyme-like protein 2	LYZL2
Q7Z5A4	Serine protease 42	PRSS42
Q7Z5A7	Protein FAM19A5	FAM19A5
Q7Z5A8	Protein FAM19A3	FAM19A3
Q7Z5A9	Protein FAM19A1	FAM19A1
Q7Z5J1	Hydroxysteroid 11-beta-dehydrogenase 1-like protein	HSD11B1L
Q7Z5L0	Vitellogenesis membrane outer layer protein 1 homolog	VMO1
Q7Z5L3	Complement C1q-like protein 2	C1QL2
Q7Z5L7	Podocan	PODN
Q7Z5P4	17-beta-hydroxysteroid dehydrogenase 13	HSD17B13
Q7Z5P9	Mucin-19	MUC19
Q7Z5Y6	Bone morphogenetic protein 8A	BMP8A
Q7Z7B7	Beta-defensin 132	DEFB132
Q7Z7B8	Beta-defensin 128	DEFB128
Q7Z7C8	Transcription initiation factor TFIID subunit 8	TAF8
Q7Z7H5	Transmembrane emp24 domain-containing protein 4	TMED4
Q86SG7	Lysozyme g-like protein 2	LYG2
Q86SI9	Protein CE1	C5orf38
Q86TE4	Leucine zipper protein 2	LUZP2
Q86TH1	ADAMTS-like protein 2	ADAMTSL2
Q86U17	Serpin A11	SERPINA11
Q86UU9	Endokinin-A	TAC4
Q86UW8	Hyaluronan and proteoglycan link protein 4	HAPLN4
Q86UX2	Inter-alpha-trypsin inhibitor heavy chain H5	ITIH5
Q86V24	Adiponectin receptor protein 2	ADIPOR2
Q86VB7	Soluble CD163	CD163
Q86VR8	Four-jointed box protein 1	FJX1
Q86WD7	Serpin A9	SERPINA9
Q86WN2	Interferon epsilon	IFNE
Q86WS3	Placenta-specific 1-like protein	PLAC1L
Q86X52	Chondroitin sulfate synthase 1	CHSY1
Q86XP6	Gastrokin-2	GKN2
Q86XS5	Angiopoietin-related protein 5	ANGPTL5
Q86Y27	B melanoma antigen 5	BAGE5
Q86Y28	B melanoma antigen 4	BAGE4
Q86Y29	B melanoma antigen 3	BAGE3
Q86Y30	B melanoma antigen 2	BAGE2
Q86Y38	Xylosyltransferase 1	XYLT1
Q86Y78	Ly6/PLAUR domain-containing protein 6	LYPD6
Q86YD3	Transmembrane protein 25	TMEM25
Q86YJ6	Threonine synthase-like 2	THNSL2
Q86YW7	Glycoprotein hormone beta-5	GPHB5
Q86Z23	Complement C1q-like protein 4	C1QL4
Q8IU57	Interleukin-28 receptor subunit alpha	IL28RA
Q8IUA0	WAP four-disulfide core domain protein 8	WFDC8
Q8IUB2	WAP four-disulfide core domain protein 3	WFDC3
Q8IUB3	Protein WFDC10B	WFDC10B
Q8IUB5	WAP four-disulfide core domain protein 13	WFDC13
Q8IUH2	Protein CREG2	CREG2
Q8IUK5	Plexin domain-containing protein 1	PLXDC1
Q8IUL8	Cartilage intermediate layer protein 2 C2	CILP2
Q8IUX7	Adipocyte enhancer-binding protein 1	AEBP1
Q8IUX8	Epidermal growth factor-like protein 6	EGFL6
Q8IVL8	Carboxypeptidase O	CPO
Q8IVN8	Somatomedin-B and thrombospondin type-1 domain-containing protein	SBSPON
Q8IVW8	Protein spinster homolog 2	SPNS2

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q8IW75	Serpin A12	SERPINA12
Q8IW92	Beta-galactosidase-1-like protein 2	GLB1L2
Q8IWL1	Pulmonary surfactant-associated protein A2	SFTPA2
Q8IWL2	Pulmonary surfactant-associated protein A1	SFTPA1
Q8IWV2	Contactin-4	CNTN4
Q8IWY4	Signal peptide, CUB and EGF-like domain-containing protein 1	SCUBE1
Q8IX30	Signal peptide, CUB and EGF-like domain-containing protein 3	SCUBE3
Q8IXA5	Sperm acrosome membrane-associated protein 3, membrane form	SPACA3
Q8IXB1	DnaJ homolog subfamily C member 10	DNAJC10
Q8IXL6	Extracellular serine/threonine protein kinase Fam20C	FAM20C
Q8IYD9	Lung adenoma susceptibility protein 2	LAS2
Q8IYP2	Serine protease 58	PRSS58
Q8IYS5	Osteoclast-associated immunoglobulin-like receptor	OSCAR
Q8IZC6	Collagen alpha-1(XXVII) chain	COL27A1
Q8IZJ3	C3 and PZP-like alpha-2-macroglobulin domain-containing protein 8	CPAMD8
Q8IZN7	Beta-defensin 107	DEFB107B
Q8NOV4	Leucine-rich repeat LGI family member 2	LGI2
Q8N104	Beta-defensin 106	DEFB106B
Q8N119	Matrix metalloproteinase-21	MMP21
Q8N129	Protein canopy homolog 4	CNPY4
Q8N135	Leucine-rich repeat LGI family member 4	LGI4
Q8N145	Leucine-rich repeat LGI family member 3	LGI3
Q8N158	Glypican-2	GPC2
Q8N1E2	Lysozyme g-like protein 1	LYG1
Q8N2E2	von Willebrand factor D and EGF domain-containing protein	VWDE
Q8N2E6	Prosalusin	TOR2A
Q8N2S1	Latent-transforming growth factor beta-binding protein 4	LTBP4
Q8N302	Angiogenic factor with G patch and FHA domains 1	AGGF1
Q8N307	Mucin-20	MUC20
Q8N323	NXPE family member 1	NXPE1
Q8N387	Mucin-15	MUC15
Q8N3Z0	Inactive serine protease 35	PRSS35
Q8N436	Inactive carboxypeptidase-like protein X2	CPXM2
Q8N474	Secreted frizzled-related protein 1	SFRP1
Q8N475	Follistatin-related protein 5	FSTL5
Q8N4F0	BPI fold-containing family B member 2	BPIFB2
Q8N4T0	Carboxypeptidase A6	CPA6
Q8N5W8	Protein FAM24B	FAM24B
Q8N687	Beta-defensin 125	DEFB125
Q8N688	Beta-defensin 123	DEFB123
Q8N690	Beta-defensin 119	DEFB119
Q8N6C5	Immunoglobulin superfamily member 1	IGSF1
Q8N6C8	Leukocyte immunoglobulin-like receptor subfamily A member 3	LILRA3
Q8N6G6	ADAMTS-like protein 1	ADAMTSL1
Q8N6Y2	Leucine-rich repeat-containing protein 17	LRRC17
Q8N729	Neuropeptide W-23	NPW
Q8N8U9	BMP-binding endothelial regulator protein	BMPER
Q8N907	DAN domain family member 5	DAND5
Q8NAT1	Glycosyltransferase-like domain-containing protein 2	GTDC2
Q8NAU1	Fibronectin type III domain-containing protein 5	FNDC5
Q8NB37	Parkinson disease 7 domain-containing protein 1	PDDC1
Q8NBI3	Draxin	DRAXIN
Q8NBM8	Preylcysteine oxidase-like	PCYOX1L
Q8NBP7	Proprotein convertase subtilisin/kexin type 9	PCSK9
Q8NBQ5	Estradiol 17-beta-dehydrogenase 11	HSD17B11
Q8NBV8	Synaptotagmin-8	SYT8
Q8NCC3	Group XV phospholipase A2	PLA2G15
Q8NCF0	C-type lectin domain family 18 member C	CLEC18C
Q8NCW5	NAD(P)H-hydrate epimerase	APOA1BP
Q8NDA2	Hemicentin-2	HMCN2
Q8NDX9	Lymphocyte antigen 6 complex locus protein G5b	LY6G5B
Q8NDZ4	Deleted in autism protein 1	C3orf58

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q8NEB7	Acrosin-binding protein	ACRBP
Q8NES8	Beta-defensin 124	DEFB124
Q8NET1	Beta-defensin 108B	DEFB108B
Q8NEX5	Protein WFDC9	WFDC9
Q8NEX6	Protein WFDC11	WFDC11
Q8NF86	Serine protease 33	PRSS33
Q8NFM7	Interleukin-17 receptor D	IL17RD
Q8NMQ5	BPI fold-containing family B member 6	BPIFB6
Q8NMQ6	BPI fold-containing family C protein	BPIFC
Q8NFU4	Follicular dendritic cell secreted peptide	FDCSP
Q8NFW1	Collagen alpha-1(XXII) chain	COL22A1
Q8NG35	Beta-defensin 105	DEFB105B
Q8NG41	Neuropeptide B-23	NPB
Q8NHW6	Otospiralin	OTOS
Q8NI99	Angiopoietin-related protein 6	ANGPTL6
Q8TAA1	Probable ribonuclease 11	RNASE11
Q8TAG5	V-set and transmembrane domain-containing protein 2A	VSTM2A
Q8TAL6	Fin bud initiation factor homolog	FIBIN
Q8TAT2	Fibroblast growth factor-binding protein 3	FGFBP3
Q8TAX7	Mucin-7	MUC7
Q8TB22	Spermatogenesis-associated protein 20	SPATA20
Q8TB73	Protein NDNF	NDNF
Q8TB96	T-cell immunomodulatory protein	ITFG1
Q8TC92	Protein disulfide-thiol oxidoreductase	ENOX1
Q8TCV5	WAP four-disulfide core domain protein 5	WFDC5
Q8TD06	Anterior gradient protein 3 homolog	AGR3
Q8TD33	Secretoglobin family 1C member 1	SCGB1C1
Q8TD46	Cell surface glycoprotein CD200 receptor 1	CD200R1
Q8TDE3	Ribonuclease 8	RNASE8
Q8TDF5	Neuropilin and tolloid-like protein 1	NETO1
Q8TDL5	BPI fold-containing family B member 1	BPIFB1
Q8TE56	A disintegrin and metalloproteinase with thrombospondin motifs 17	ADAMTS17
Q8TE57	A disintegrin and metalloproteinase with thrombospondin motifs 16	ADAMTS16
Q8TE58	A disintegrin and metalloproteinase with thrombospondin motifs 15	ADAMTS15
Q8TE59	A disintegrin and metalloproteinase with thrombospondin motifs 19	ADAMTS19
Q8TE60	A disintegrin and metalloproteinase with thrombospondin motifs 18	ADAMTS18
Q8TE99	Acid phosphatase-like protein 2	ACPL2
Q8TER0	Sushi, nidogen and EGF-like domain-containing protein 1	SNED1
Q8TEU8	WAP, kazal, immunoglobulin, kunitz and NTR domain-containing protein 2	WFIKKN2
Q8WTQ1	Beta-defensin 104	DEFB104B
Q8WTR8	Netrin-5	NTN5
Q8WTU2	Scavenger receptor cysteine-rich domain-containing group B protein	SRCRB4D
Q8WU66	Protein TSPEAR	TSPEAR
Q8WUA8	Tsukushin	TSKU
Q8WUF8	Protein FAM172A	FAM172A
Q8WUJ1	Neuferricin	CYB5D2
Q8WUY1	UPF0670 protein THEM6	THEM6
Q8WVN6	Secreted and transmembrane protein 1	SECTM1
Q8WVQ1	Soluble calcium-activated nucleotidase 1	CANT1
Q8WWA0	Intelectin-1	ITLN1
Q8WWG1	Neuregulin-4	NRG4
Q8WWQ2	Inactive heparanase-2	HPSE2
Q8WWU7	Intelectin-2	ITLN2
Q8WWY7	WAP four-disulfide core domain protein 12	WFDC12
Q8WWY8	Lipase member H	LIPH
Q8WWZ8	Oncoprotein-induced transcript 3 protein	OIT3
Q8WX39	Epididymal-specific lipocalin-9	LCN9
Q8WXA2	Prostate and testis expressed protein 1	PATE1
Q8WXD2	Secretogranin-3	SCG3
Q8WXF3	Relaxin-3 A chain	RLN3
Q8WXI7	Mucin-16	MUC16
Q8WXQ8	Carboxypeptidase A5	CPA5
Q8WXS8	A disintegrin and metalloproteinase with thrombospondin motifs 14	ADAMTS14
Q92484	Acid sphingomyelinase-like phosphodiesterase 3a	SMPDL3A

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q92485	Acid sphingomyelinase-like phosphodiesterase 3b	SMPDL3B
Q92496	Complement factor H-related protein 4	CFHR4
Q92520	Protein FAM3C	FAM3C
Q92563	Testican-2	SPOCK2
Q92583	C-C motif chemokine 17	CCL17
Q92626	Peroxidasin homolog	PXDN
Q92743	Serine protease HTRA1	HTRA1
Q92752	Tenascin-R	TNR
Q92765	Secreted frizzled-related protein 3	FRZB
Q92819	Hyaluronan synthase 2	HAS2
Q92820	Gamma-glutamyl hydrolase	GGH
Q92824	Proprotein convertase subtilisin/kexin type 5	PCSK5
Q92832	Protein kinase C-binding protein NELL1	NELL1
Q92838	Ectodysplasin-A, membrane form	EDA
Q92874	Deoxyribonuclease-1-like 2	DNASE1L2
Q92876	Kallikrein-6	KLK6
Q92913	Fibroblast growth factor 13	FGF13
Q92954	Proteoglycan 4 C-terminal part	PRG4
Q93038	Tumor necrosis factor receptor superfamily member 25	TNFRSF25
Q93091	Ribonuclease K6	RNASE6
Q93097	Protein Wnt-2b	WNT2B
Q93098	Protein Wnt-8b	WNT8B
Q95460	Major histocompatibility complex class I-related gene protein	MR1
Q969D9	Thymic stromal lymphopoietin	TSLP
Q969E1	Liver-expressed antimicrobial peptide 2	LEAP2
Q969H8	UPF0556 protein C19orf10	C19orf10
Q969Y0	NXPE family member 3	NXPE3
Q96A54	Adiponectin receptor protein 1	ADIPOR1
Q96A83	Collagen alpha-1(XXVI) chain	EMID2
Q96A84	EMI domain-containing protein 1	EMID1
Q96A98	Tuberoinfundibular peptide of 39 residues	PTH2
Q96A99	Pentraxin-4	PTX4
Q96BH3	Epididymal sperm-binding protein 1	ELSPBP1
Q96BQ1	Protein FAM3D	FAM3D
Q96CG8	Collagen triple helix repeat-containing protein 1	CTHRC1
Q96DA0	Zymogen granule protein 16 homolog B	ZG16B
Q96DN2	von Willebrand factor C and EGF domain-containing protein	VWCE
Q96DR5	BPI fold-containing family A member 2	BPIFA2
Q96DR8	Mucin-like protein 1	MUCL1
Q96DX4	RING finger and SPRY domain-containing protein 1	RSPRY1
Q96EE4	Coiled-coil domain-containing protein 126	CCDC126
Q96GS6	Abhydrolase domain-containing protein FAM108A1	FAM108A1
Q96GW7	Brevican core protein	BCAN
Q96HF1	Secreted frizzled-related protein 2	SFRP2
Q96I82	Kazal-type serine protease inhibitor domain-containing protein 1	KAZALD1
Q96ID5	Immunoglobulin superfamily member 21	IGSF21
Q96II8	Leucine-rich repeat and calponin homology domain-containing protein 3	LRCH3
Q96IY4	Carboxypeptidase B2	CPB2
Q96JB6	Lysyl oxidase homolog 4	LOXL4
Q96JK4	HHIP-like protein 1	HHIPL1
Q96KN2	Beta-Ala-His dipeptidase	CNDP1
Q96KW9	Protein SPACA7	SPACA7
Q96KX0	Lysozyme-like protein 4	LYZL4
Q96L15	Ecto-ADP-ribosyltransferase 5	ART5
Q96LB8	Peptidoglycan recognition protein 4	PGLYRP4
Q96LB9	Peptidoglycan recognition protein 3	PGLYRP3
Q96LC7	Sialic acid-binding Ig-like lectin 10	SIGLEC10
Q96LR4	Protein FAM19A4	FAM19A4
Q96MK3	Protein FAM20A	FAM20A
Q96MS3	Glycosyltransferase 1 domain-containing protein 1	GLT1D1
Q96NY8	Processed poliovirus receptor-related protein 4	PVRL4
Q96NZ8	WAP, kazal, immunoglobulin, kunitz and NTR domain-containing protein 1	WFIKKN1
Q96NZ9	Proline-rich acidic protein 1	PRAP1
Q96P44	Collagen alpha-1(XXI) chain	COL21A1
Q96PB7	Noelin-3	OLFM3

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q96PC5	Melanoma inhibitory activity protein 2	MIA2
Q96PD5	N-acetylmuramoyl-L-alanine amidase	PGLYRP2
Q96PH6	Beta-defensin 118	DEFB118
Q96PL1	Secretoglobin family 3A member 2	SCGB3A2
Q96PL2	Beta-tectorin	TECTB
Q96QH8	Sperm acrosome-associated protein 5	SPACA5
Q96QR1	Secretoglobin family 3A member 1	SCGB3A1
Q96QU1	Protocadherin-15	PCDH15
Q96QV1	Hedgehog-interacting protein	HHIP
Q96RW7	Hemicentin-1	HMCN1
Q96S42	Nodal homolog	NODAL
Q96S86	Hyaluronan and proteoglycan link protein 3	HAPLN3
Q96SL4	Glutathione peroxidase 7	GPX7
Q96SM3	Probable carboxypeptidase X1	CPXM1
Q96T91	Glycoprotein hormone alpha-2	GPHA2
Q99062	Granulocyte colony-stimulating factor receptor	CSF3R
Q99102	Mucin-4 alpha chain	MUC4
Q99217	Amelogenin, X isoform	AMELX
Q99218	Amelogenin, Y isoform	AMELY
Q99435	Protein kinase C-binding protein NELL2	NELL2
Q99470	Stromal cell-derived factor 2	SDF2
Q99542	Matrix metalloproteinase-19	MMP19
Q99574	Neuroserpin	SERPINI1
Q99584	Protein S100-A13	S100A13
Q99616	C-C motif chemokine 13	CCL13
Q99645	Epiphycan	EPYC
Q99674	Cell growth regulator with EF hand domain protein 1	CGREF1
Q99715	Collagen alpha-1(XII) chain	COL12A1
Q99727	Metalloproteinase inhibitor 4	TIMP4
Q99731	C-C motif chemokine 19	CCL19
Q99748	Neurturin	NRTN
Q99935	Proline-rich protein 1	PROL1
Q99942	E3 ubiquitin-protein ligase RNF5	RNF5
Q99944	Epidermal growth factor-like protein 8	EGFL8
Q99954	Submaxillary gland androgen-regulated protein 3A	SMR3A
Q99969	Retinoic acid receptor responder protein 2	RARRES2
Q99972	Myocilin	MYOC
Q99983	Osteomodulin	OMD
Q99985	Semaphorin-3C	SEMA3C
Q99988	Growth/differentiation factor 15	GDF15
Q9BPW4	Apolipoprotein L4	APOL4
Q9BQ08	Resistin-like beta	RETNLB
Q9BQ16	Testican-3	SPOCK3
Q9BQ51	Programmed cell death 1 ligand 2	PDCD1LG2
Q9BQB4	Sclerostin	SOST
Q9BQI4	Coiled-coil domain-containing protein 3	CCDC3
Q9BQP9	BPI fold-containing family A member 3	BPIFA3
Q9BQR3	Serine protease 27	PRSS27
Q9BQY6	WAP four-disulfide core domain protein 6	WFDC6
Q9BRR6	ADP-dependent glucokinase	ADPGK
Q9BS86	Zona pellucida-binding protein 1	ZBPB
Q9BSG0	Protease-associated domain-containing protein 1	PRADC1
Q9BSG5	Retbindin	RTBDN
Q9BT30	Probable alpha-ketoglutarate-dependent dioxygenase ABH7	ALKBH7
Q9BT56	Spexin	C12orf39
Q9BT67	NEDD4 family-interacting protein 1	NDFIP1
Q9BTY2	Plasma alpha-L-fucosidase	FUCA2
Q9BU40	Chordin-like protein 1	CHRDL1
Q9BUD6	Spondin-2	SPON2
Q9BUN1	Protein MENT	MENT
Q9BUR5	Apolipoprotein O	APOO
Q9BV94	ER degradation-enhancing alpha-mannosidase-like 2	EDEM2
Q9BWP8	Collectin-11	COLEC11
Q9BWS9	Chitinase domain-containing protein 1	CHID1
Q9BX67	Junctional adhesion molecule C	JAM3
Q9BX93	Group XIIB secretory phospholipase A2-like protein	PLA2G12B
Q9BXI9	Complement C1q tumor necrosis factor-related protein 6	C1QTNF6
Q9BXJ0	Complement C1q tumor necrosis factor-related protein 5	C1QTNF5

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q9BXJ1	Complement C1q tumor necrosis factor-related protein 1	C1QTNF1
Q9BXJ2	Complement C1q tumor necrosis factor-related protein 7	C1QTNF7
Q9BXJ3	Complement C1q tumor necrosis factor-related protein 4	C1QTNF4
Q9BXJ4	Complement C1q tumor necrosis factor-related protein 3	C1QTNF3
Q9BXJ5	Complement C1q tumor necrosis factor-related protein 2	C1QTNF2
Q9BXN1	Asporin	ASPN
Q9BXP8	Pappalysin-2	PAPPA2
Q9BXR6	Complement factor H-related protein 5	CFHR5
Q9BXS0	Collagen alpha-1(XXV) chain	COL25A1
Q9BXX0	EMILIN-2	EMILIN2
Q9BXY4	R-spondin-3	RSPO3
Q9BY15	EGF-like module-containing mucin-like hormone receptor-like 3 subunit beta	EMR3
Q9BY50	Signal peptidase complex catalytic subunit SEC11C	SEC11C
Q9BY76	Angiopoietin-related protein 4	ANGPTL4
Q9BYF1	Processed angiotensin-converting enzyme 2	ACE2
Q9BYJ0	Fibroblast growth factor-binding protein 2	FGFBP2
Q9BYW3	Beta-defensin 126	DEFB126
Q9BYX4	Interferon-induced helicase C domain-containing protein 1	IFIH1
Q9BYZ8	Regenerating islet-derived protein 4	REG4
Q9BZ76	Contactin-associated protein-like 3	CNTNAP3
Q9BZG9	Ly-6/neurotoxin-like protein 1	LYNX1
Q9BZJ3	Tryptase delta	TPSD1
Q9BZM1	Group XIIA secretory phospholipase A2	PLA2G12A
Q9BZM2	Group IIF secretory phospholipase A2	PLA2G2F
Q9BZM5	NKG2D ligand 2	ULBP2
Q9BZP6	Acidic mammalian chitinase	CHIA
Q9BZZ2	Sialoadhesin	SIGLEC1
Q9COB6	Protein FAM5B	FAM5B
Q9GZM7	Tubulointerstitial nephritis antigen-like	TINAGL1
Q9GZN4	Brain-specific serine protease 4	PRSS22
Q9GZP0	Platelet-derived growth factor D, receptor-binding form	PDGFD
Q9GZT5	Protein Wnt-10a	WNT10A
Q9GZU5	Nyctalopin	NYX
Q9GZV7	Hyaluronan and proteoglycan link protein 2	HAPLN2
Q9GZV9	Fibroblast growth factor 23	FGF23
Q9GZX9	Twisted gastrulation protein homolog 1	TWSG1
Q9GZZ7	GDNF family receptor alpha-4	GFRA4
Q9GZZ8	Extracellular glycoprotein lacritin	LACRT
Q9H0B8	Cysteine-rich secretory protein LCCL domain-containing 2	CRISPLD2
Q9H106	Signal-regulatory protein delta	SIRPD
Q9H114	Cystatin-like 1	CSTL1
Q9H173	Nucleotide exchange factor SIL1	SIL1
Q9H1E1	Ribonuclease 7	RNASE7
Q9H1F0	WAP four-disulfide core domain protein 10A	WFDC10A
Q9H1J5	Protein Wnt-8a	WNT8A
Q9H1J7	Protein Wnt-5b	WNT5B
Q9H1M3	Beta-defensin 129	DEFB129
Q9H1M4	Beta-defensin 127	DEFB127
Q9H1Z8	Augurin	C2orf40
Q9H239	Matrix metalloproteinase-28	MMP28
Q9H2A7	C-X-C motif chemokine 16	CXCL16
Q9H2A9	Carbohydrate sulfotransferase 8	CHST8
Q9H2R5	Kallikrein-15	KLK15
Q9H2X0	Chordin	CHRD
Q9H2X3	C-type lectin domain family 4 member M	CLEC4M
Q9H306	Matrix metalloproteinase-27	MMP27
Q9H324	A disintegrin and metalloproteinase with thrombospondin motifs 10	ADAMTS10
Q9H336	Cysteine-rich secretory protein LCCL domain-containing 1	CRISPLD1
Q9H3E2	Sorting nexin-25	SNX25
Q9H3R2	Mucin-13	MUC13
Q9H3U7	SPARC-related modular calcium-binding protein 2	SMOC2
Q9H3Y0	Peptidase inhibitor R3HDML	R3HDML
Q9H4A4	Aminopeptidase B	RNPEP

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q9H4F8	SPARC-related modular calcium-binding protein 1	SMOC1
Q9H4G1	Cystatin-9-like	CST9L
Q9H5V8	CUB domain-containing protein 1	CDCP1
Q9H6B9	Epoxide hydrolase 3	EPHX3
Q9H6E4	Coiled-coil domain-containing protein 134	CCDC134
Q9H741	UPF0454 protein C12orf49	C12orf49
Q9H772	Gremlin-2	GREM2
Q9H7Y0	Deleted in autism-related protein 1	CXorf36
Q9H8L6	Multimerin-2	MMRN2
Q9H9S5	Fukutin-related protein	FKRP
Q9HAT2	Sialate O-acetyltransferase	SIAE
Q9HB40	Retinoid-inducible serine carboxypeptidase	SCPEP1
Q9HB63	Netrin-4	NTN4
Q9HBJ0	Placenta-specific protein 1	PLAC1
Q9HC23	Prokineticin-2	PROK2
Q9HC57	WAP four-disulfide core domain protein 1	WFDC1
Q9HC73	Cytokine receptor-like factor 2	CRLF2
Q9HC84	Mucin-5B	MUC5B
Q9HCB6	Spondin-1	SPON1
Q9HCQ7	Neuropeptide NPSF	NPVF
Q9HCT0	Fibroblast growth factor 22	FGF22
Q9HD89	Resistin	RETN
Q9NNX1	Tuftelin	TUFT1
Q9NNX6	CD209 antigen	CD209
Q9NP55	BPI fold-containing family A member 1	BPIFA1
Q9NP70	Ameloblastin	AMBN
Q9NP95	Fibroblast growth factor 20	FGF20
Q9NP99	Triggering receptor expressed on myeloid cells 1	TREM1
Q9NPA2	Matrix metalloproteinase-25	MMP25
Q9NPE2	Neugrin	NGRN
Q9NPH0	Lysophosphatidic acid phosphatase type 6	ACP6
Q9NPH6	Odorant-binding protein 2b	OBP2B
Q9NQ30	Endothelial cell-specific molecule 1	ESM1
Q9NQ36	Signal peptide, CUB and EGF-like domain-containing protein 2	SCUBE2
Q9NQ38	Serine protease inhibitor Kazal-type 5	SPINK5
Q9NQ76	Matrix extracellular phosphoglycoprotein	MEPE
Q9NQ79	Cartilage acidic protein 1	CRTAC1
Q9NR16	Scavenger receptor cysteine-rich type 1 protein M160	CD163L1
Q9NR23	Growth/differentiation factor 3	GDF3
Q9NR71	Neutral ceramidase	ASAH2
Q9NR99	Matrix-remodeling-associated protein 5	MXRA5
Q9NRA1	Platelet-derived growth factor C	PDGFC
Q9NRC9	Otoraplin	OTOR
Q9NRE1	Matrix metalloproteinase-26	MMP26
Q9NRJ3	C-C motif chemokine 28	CCL28
Q9NRM1	Enamelin	ENAM
Q9NRN5	Olfactomedin-like protein 3	OLFML3
Q9NRR1	Cytokine-like protein 1	CYTL1
Q9NS15	Latent-transforming growth factor beta-binding protein 3	LTBP3
Q9NS62	Thrombospondin type-1 domain-containing protein 1	THSD1
Q9NS71	Gastrokin-1	GKN1
Q9NS98	Semaphorin-3G	SEMA3G
Q9NSA1	Fibroblast growth factor 21	FGF21
Q9NT22	EMILIN-3	EMILIN3
Q9NTU7	Cerebellin-4	CBLN4
Q9NVR0	Kelch-like protein 11	KLHL11
Q9NWH7	Spermatogenesis-associated protein 6	SPATA6
Q9NXC2	Glucose-fructose oxidoreductase domain-containing protein 1	GFOD1
Q9NY56	Odorant-binding protein 2a	OBP2A
Q9NY84	Vascular non-inflammatory molecule 3	VNN3
Q9NZ20	Group 3 secretory phospholipase A2	PLA2G3
Q9NZC2	Triggering receptor expressed on myeloid cells 2	TREM2
Q9NZK5	Adenosine deaminase CECR1	CECR1
Q9NZK7	Group IIE secretory phospholipase A2	PLA2G2E
Q9NZP8	Complement C1r subcomponent-like protein	C1RL
Q9NZV1	Cysteine-rich motor neuron 1 protein	CRIM1
Q9NZW4	Dentin sialoprotein	DSPP
Q9P0G3	Kallikrein-14	KLK14

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q9P0W0	Interferon kappa	IFNK
Q9P218	Collagen alpha-1(XX) chain	COL20A1
Q9P2C4	Transmembrane protein 181	TMEM181
Q9P2K2	Thioredoxin domain-containing protein 16	TXNDC16
Q9P2N4	A disintegrin and metalloproteinase with thrombospondin motifs 9	ADAMTS9
Q9UBC7	Galanin-like peptide	GALP
Q9UBD3	Cytokine SCM-1 beta	XCL2
Q9UBD9	Cardiotrophin-like cytokine factor 1	CLCF1
Q9UBM4	Opticin	OPTC
Q9UBP4	Dickkopf-related protein 3	DKK3
Q9UBQ6	Exostosin-like 2	EXTL2
Q9UBR5	Chemokine-like factor	CKLF
Q9UBS5	Gamma-aminobutyric acid type B receptor subunit 1	GABBR1
Q9UBT3	Dickkopf-related protein 4 short form	DKK4
Q9UBU2	Dickkopf-related protein 2	DKK2
Q9UBU3	Ghrelin-28	GHRL
Q9UBV4	Protein Wnt-16	WNT16
Q9UBX5	Fibulin-5	FBLN5
Q9UBX7	Kallikrein-11	KLK11
Q9UEF7	Klotho	KL
Q9UFP1	Protein FAM198A	FAM198A
Q9UGM3	Deleted in malignant brain tumors 1 protein	DMBT1
Q9UGM5	Fetuin-B	FETUB
Q9UGP8	Translocation protein SEC63 homolog	SEC63
Q9UHF0	Neurokinin-B	TAC3
Q9UHF1	Epidermal growth factor-like protein 7	EGFL7
Q9UHG2	ProSAAS	PCSK1N
Q9UHI8	A disintegrin and metalloproteinase with thrombospondin motifs 1	ADAMTS1
Q9UHL4	Dipeptidyl peptidase 2	DPP7
Q9UI42	Carboxypeptidase A4	CPA4
Q9UIG4	Psoriasis susceptibility 1 candidate gene 2 protein	PSORS1C2
Q9UIK5	Tomoregulin-2	TMEFF2
Q9UIQ6	Leucyl-cystinyl aminopeptidase, pregnancy serum form	LNPEP
Q9UJA9	Ectonucleotide pyrophosphatase/phosphodiesterase family member 5	ENPP5
Q9UJH8	Meteorin	METR
Q9UJJ9	N-acetylglucosamine-1-phosphotransferase subunit gamma	GNPTG
Q9UJW2	Tubulointerstitial nephritis antigen	TINAG
Q9UK05	Growth/differentiation factor 2	GDF2
Q9UK55	Protein Z-dependent protease inhibitor	SERPINA10
Q9UK85	Dickkopf-like protein 1	DKKL1
Q9UKJ1	Paired immunoglobulin-like type 2 receptor alpha	PILRA
Q9UKP4	A disintegrin and metalloproteinase with thrombospondin motifs 7	ADAMTS7
Q9UKP5	A disintegrin and metalloproteinase with thrombospondin motifs 6	ADAMTS6
Q9UKQ2	Disintegrin and metalloproteinase domain-containing protein 28	ADAM28
Q9UKQ9	Kallikrein-9	KLK9
Q9UKR0	Kallikrein-12	KLK12
Q9UKR3	Kallikrein-13	KLK13
Q9UKU9	Angiopoietin-related protein 2	ANGPTL2
Q9UKZ9	Procollagen C-endopeptidase enhancer 2	PCOLCE2
Q9UL52	Transmembrane protease serine 11E non-catalytic chain	TMPRSS11E
Q9ULC0	Endomucin	EMCN
Q9ULI3	Protein HEG homolog 1	HEG1
Q9ULZ1	Apelin-13	APLN
Q9ULZ9	Matrix metalloproteinase-17	MMP17
Q9UM21	Alpha-1,3-mannosyl-glycoprotein 4-beta-N-acetylglucosaminyltransferase A soluble form	MGAT4A
Q9UM22	Mammalian ependymin-related protein 1	EPDR1
Q9UM73	ALK tyrosine kinase receptor	ALK
Q9UMD9	97 kDa linear IgA disease antigen	COL17A1
Q9UMX5	Neudesin	NENF
Q9UN73	Protocadherin alpha-6	PCDHA6
Q9UNA0	A disintegrin and metalloproteinase with thrombospondin motifs 5	ADAMTS5

TABLE 6-continued

Secreted Proteins		
Uniprot ID	Protein Name	Gene Name
Q9UNI1	Chymotrypsin-like elastase family member 1	CELA1
Q9UNK4	Group IID secretory phospholipase A2	PLA2G2D
Q9UP79	A disintegrin and metalloproteinase with thrombospondin motifs 8	ADAMTS8
Q9UPZ6	Thrombospondin type-1 domain-containing protein 7A	THSD7A
Q9UQ72	Pregnancy-specific beta-1-glycoprotein 11	PSG11
Q9UQ74	Pregnancy-specific beta-1-glycoprotein 8	PSG8
Q9UQC9	Calcium-activated chloride channel regulator 2	CLCA2
Q9UQE7	Structural maintenance of chromosomes protein 3	SMC3
Q9UQP3	Tenascin-N	TNN
Q9Y223	UDP-N-acetylglucosamine 2-epimerase	GNE
Q9Y240	C-type lectin domain family 11 member A	CLEC11A
Q9Y251	Heparanase 8 kDa subunit	HPSE
Q9Y258	C-C motif chemokine 26	CCL26
Q9Y264	Angiotensinogen-converting enzyme 1	ANGPT4
Q9Y275	Tumor necrosis factor ligand superfamily member 13b, membrane form	TNFSF13B
Q9Y287	BRI2 intracellular domain	ITM2B
Q9Y2E5	Epididymis-specific alpha-mannosidase	MAN2B2
Q9Y334	von Willebrand factor A domain-containing protein 7	VWA7
Q9Y337	Kallikrein-5	KLK5
Q9Y3B3	Transmembrane emp24 domain-containing protein 7	TMED7
Q9Y3E2	BOLA-like protein 1	BOLA1
Q9Y426	C2 domain-containing protein 2	C2CD2
Q9Y4K0	Lysyl oxidase homolog 2	LOXL2
Q9Y4X3	C-C motif chemokine 27	CCL27
Q9Y5C1	Angiotensinogen-converting enzyme 1	ANGPTL3
Q9Y5I2	Protocadherin alpha-10	PCDHA10
Q9Y5I3	Protocadherin alpha-1	PCDHA1
Q9Y5K2	Kallikrein-4	KLK4
Q9Y5L2	Hypoxia-inducible lipid droplet-associated protein	HILPDA
Q9Y5Q5	Atrial natriuretic peptide-converting enzyme	CORIN
Q9Y5R2	Matrix metalloproteinase-24	MMP24
Q9Y5U5	Tumor necrosis factor receptor superfamily member 18	TNFRSF18
Q9Y5W5	Wnt inhibitory factor 1	WIF1
Q9Y5X9	Endothelial lipase	LIPG
Q9Y625	Secreted glypican-6	GPC6
Q9Y646	Carboxypeptidase Q	CPQ
Q9Y6C2	EMILIN-1	EMILIN1
Q9Y6F9	Protein Wnt-6	WNT6
Q9Y6I9	Testis-expressed sequence 264 protein	TEX264
Q9Y6L7	Tolloid-like protein 2	TLL2
Q9Y6N3	Calcium-activated chloride channel regulator family member 3	CLCA3P
Q9Y6N6	Laminin subunit gamma-3	LAMC3
Q9Y6R7	IgGf-binding protein	FCGBP
Q9Y6Y9	Lymphocyte antigen 96	LY96
Q9Y6Z7	Collectin-10	COLEC10

In one set of embodiments, the MCNA compound comprises two encoding polynucleotides. For example, the MCNA compound may be a palindromic coding nucleic acid (PCNA) having two encoding polynucleotides each having a polynucleotide portion that codes for the same protein.

In some embodiments, a MCNA compound comprises an encoding polynucleotide that encodes Cystic Fibrosis Transmembrane Conductance Regulator (hCFTR) mRNA, linked to a non-coding polynucleotide via a 3' end linkage between the polynucleotides. In some embodiments, a MCNA compound comprises two or more encoding polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein at least one of the encoding polynucleotides encodes hCFTR. In some embodiments, a MCNA compound is a palindromic coding nucleic acid (PCNA) com-

prising two encoding polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein each encoding polynucleotide codes for hCFTR. In some embodiments, a MCNA compound comprises two or more polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein at least one polynucleotide is an encoding polynucleotide that encodes hCFTR and at least one polynucleotide acts as a protecting group.

In some embodiments, a MCNA compound comprises an encoding polynucleotide that encodes human phenylalanine hydroxylase (hPAH) mRNA, linked to a non-coding polynucleotide via a 3' end linkage between the polynucleotides. In some embodiments, a MCNA compound comprises two or more encoding polynucleotides linked via a 3' end linkage

between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein at least one of the encoding polynucleotides encodes hPAH. In some embodiments, a MCNA compound is a palindromic coding nucleic acid (PCNA) comprising two encoding polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein each encoding polynucleotide codes for hPAH. In some embodiments, a MCNA compound comprises two or more polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein at least one polynucleotide is an encoding polynucleotide that encodes hPAH and at least one polynucleotide acts as a protecting group.

In some embodiments, a MCNA compound comprises an encoding polynucleotide that encodes human Ornithine transcarbamylase (hOTC) mRNA, linked to a non-coding polynucleotide via a 3' end linkage between the polynucleotides. In some embodiments, a MCNA compound comprises two or more encoding polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein at least one of the encoding polynucleotides encodes hOTC. In some embodiments, a MCNA compound is a palindromic coding nucleic acid (PCNA) comprising two encoding polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein each polynucleotide codes for hOTC. In some embodiments, a MCNA compound comprises two or more polynucleotides linked via a 3' end linkage between the polynucleotides such that the MCNA compound comprises two or more 5' ends, wherein at least one polynucleotide is an encoding polynucleotide that encodes hOTC and at least one polynucleotide acts as a protecting group.

Bridge (w/3'-3' Linkage)

In some embodiments, a MCNA compound comprises two or more polynucleotides wherein the 3' ends of each polynucleotide are linked via an oligonucleotide bridge (also "bridging oligonucleotide" or "bridging oligo") comprising a 3'-3' inverted phosphodiester linkage. In some embodiments, the oligonucleotide bridge comprises modified nucleotides. In some embodiments, the oligonucleotide bridge comprises 2'-O-methyl RNA. In some embodiments, the oligonucleotide bridge comprises DNA. In some embodiments, the oligonucleotide bridge is between 2 and 1000 nucleotides in length. In some embodiments, the oligonucleotide bridge comprises one or more active moieties that are bound to the bridge by covalent links. In some embodiments, an active moiety is a targeting group, peptide, contrast agent, small molecule, protein, DNA and/or RNA. In some embodiments, an active moiety binds a receptor ligand for a cell surface receptor. In some embodiments, the active moiety is one or more tri-antennary GalNAc targeting agents.

MCNA Synthesis

In some embodiments, the present invention provides methods of synthesizing MCNA. In some embodiments, the synthesis of MCNA comprises ligating two or more polynucleotides such that the 3' end of each polynucleotide is ligated to the 5' end of an oligonucleotide bridge, wherein the oligonucleotide bridge comprises two 5' ends and an internal 3'-3' inverted phosphodiester linkage. In some embodiments, the method of synthesizing MCNA comprises the use of oligonucleotide splints complementary to regions of the two or more polynucleotides such that a ligase can join each polynucleotide to a 5' end of an oligonucleotide bridge. In some embodiments, oligonucleotide splints are

complementary to regions of the two or more polynucleotides such that a ligase joins perfect ends of each polynucleotide to a 5' end of an oligonucleotide bridge. In some embodiments, oligonucleotide splints are complementary to regions of the two or more polynucleotides such that a ligase joins the 3' end of each polynucleotide to a 5' end of an oligonucleotide bridge. In some embodiments, an oligonucleotide splint comprises DNA. In some embodiments, a ligase is RNA Ligase. In some embodiments, a ligase is T4 RNA Ligase 1. In some embodiments, a ligase is T4 RNA Ligase 2.

In some embodiments, the molar ratio of polynucleotide to oligonucleotide bridge to oligonucleotide splint when synthesizing MCNA is 2:1:2. In some embodiments, the molar ratio of polynucleotide to oligonucleotide bridge when synthesizing MCNA is 2:1. In some embodiments, the molar ratio of polynucleotide to oligonucleotide splint when synthesizing MCNA is 2:2. In some embodiments, synthesis of MCNA further comprises PEG.

In some embodiments, MCNA can be prepared by splint ligation of the 3' end of two copies of an RNA to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5' untranslated region (UTR) and a 3' UTR flanking an RNA coding sequence is transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This transcript is then ligated in a single step to a "bridge" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt using either (A) T4 RNA ligase 1, (B) T4 RNA ligase 1+PEG 8K, or (C) T4 RNA Ligase 2 and a DNA oligonucleotide "splint" complementary to the 3'-UTR and bridging oligo. To prepare the samples for ligation, the bridging oligo is 5'-end phosphorylated in a reaction containing 50 μ M oligo, ATP, 1 \times PNK Buffer and T4 Polynucleotide Kinase at 37 $^{\circ}$ C. for 1 hour. Phosphorylated bridging oligo is then desalted using a Sephadex G-25 desalting column and hybridized to the transcript and splint in a reaction containing capped RNA transcript, 1 \times bridging oligo and 2 \times splint oligo by heating to 75 $^{\circ}$ C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction is subsequently prepared to contain a 50% diluted hybridization reaction and (A) 1 \times RNA ligase Buffer, ATP and T4 RNA ligase 1 (NEB), (B) 1 \times RNA ligase Buffer, ATP, 10% PEG and T4 RNA ligase 1 (NEB), or (C) 1 \times T4RNA Ligase 2 Buffer and T4 RNA ligase 2 (NEB). Each is reacted for 90 minutes at 37 $^{\circ}$ C. The completed ligation reaction is then purified using an RNeasy Mini Kit (Qiagen). The purified MCNA product is subsequently treated with DNase I to remove residual bridge oligonucleotide.

In some embodiments, MCNA can be prepared by splint-independent ligation of the 3' end of two copies of an RNA to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence.

Untranslated Regions

Typically, mRNA synthesis includes the addition of a "cap" on the 5' end, and a "tail" on the 3' end. The presence of the cap is important in providing resistance to nucleases found in most eukaryotic cells. The presence of a "tail" serves to protect the mRNA from exonuclease degradation.

In some embodiments, one or more polynucleotides of the MCNA include a 5' and/or 3' untranslated region. In some embodiments, a 5' untranslated region (5' UTR) includes one or more elements that affect an mRNA's stability or trans-

lation, for example, an iron responsive element. In some embodiments, a 5' untranslated region may be between about 50 and 500 nucleotides in length.

In some embodiments, a 3' untranslated region (3' UTR) includes one or more of a polyadenylation signal, a binding site for proteins that affect MCNA's stability of location in a cell, or one or more binding sites for miRNAs. In some embodiments, a 3' untranslated region may be between 50 and 500 nucleotides in length or longer. In some embodiments, a 3' untranslated region may be between 5 and 2,000 nucleotides in length.

Exemplary 3' and/or 5' UTR sequences can be derived from nucleic acid molecules that are stable (e.g., globin, actin, GAPDH, tubulin, histone, or citric acid cycle enzymes) to increase the stability of the sense MCNA molecule. For example, a 5' UTR sequence may include a partial sequence of a CMV immediate-early 1 (IE1) gene, or a fragment thereof to improve the nuclease resistance and/or improve the half-life of the polynucleotide. Also contemplated is the inclusion of a sequence encoding human growth hormone (hGH), or a fragment thereof to the 3' end or untranslated region of the polynucleotide (e.g., MCNA) to further stabilize the polynucleotide. Generally, these modifications improve the stability and/or pharmacokinetic properties (e.g., half-life) of the polynucleotide relative to their unmodified counterparts, and include, for example modifications made to improve such polynucleotides' resistance to in vivo nuclease digestion.

3' UTR

In some embodiments, a 3' UTR comprises a plurality of multi-A segments with spacers in between. In some embodiments, spacers comprise DNA, RNA and/or modified bases. In some embodiments, each of the multi-A segments comprises 8-50 consecutive adenosines. In some embodiments, the plurality of multi-A segments range from 1-100 in number. In some embodiments, the spacers are of varying lengths ranging from 5-100. In some embodiments, a 3' UTR comprises a pseudoknot structure. A pseudoknot can be defined as an RNA structure minimally composed of two helical segments connected by single stranded regions or loops (Staple, D. W. et al., *PLoS Biology*, 2005, 3, e213). They are predominantly formed through secondary structures such as hairpin or stem loops and a distal single strand region. In some embodiments, a 3' UTR comprises a "kissing loop" sequence motif. Broadly defined, a kissing loop can be described as the structure formed when unpaired nucleotides in a stem/hairpin loop of one RNA molecule base pair with unpaired nucleotides of a second stem/hairpin loop of a separate RNA molecule. In some embodiments, a 3' UTR is not followed with a polyadenylation (poly-A) tail. In some embodiments, a 3' UTR binds to poly-A binding proteins (PABPs).

In some embodiments, MCNA include a 3' poly(A) tail structure. In some embodiments, a poly-A tail is 25-5,000 nucleotides in length. A poly-A tail on the 3' terminus of MCNA typically includes about 10 to 300 adenosine nucleotides (e.g., about 10 to 200 adenosine nucleotides, about 10 to 150 adenosine nucleotides, about 10 to 100 adenosine nucleotides, about 20 to 70 adenosine nucleotides, or about 20 to 60 adenosine nucleotides). In some embodiments, mRNAs include a 3' poly(C) tail structure. A suitable poly-C tail on the 3' terminus of MCNA typically include about 10 to 200 cytosine nucleotides (e.g., about 10 to 150 cytosine nucleotides, about 10 to 100 cytosine nucleotides, about 20 to 70 cytosine nucleotides, about 20 to 60 cytosine nucleo-

tidies, or about 10 to 40 cytosine nucleotides). The poly-C tail may be added to the poly-A tail or may substitute the poly-A tail.

Typically, the presence of a "tail" serves to protect the MCNA from exonuclease degradation. The poly A tail is thought to stabilize natural messengers and synthetic sense MCNA. Therefore, in certain embodiments a long poly A tail can be added to an MCNA molecule thus rendering the MCNA more stable. Poly A tails can be added using a variety of art-recognized techniques. For example, long poly A tails can be added to synthetic or in vitro transcribed RNA using poly A polymerase (Yokoe, et al. *Nature Biotechnology*, 1996; 14: 1252-1256). A transcription vector can also encode long poly A tails. In addition, poly A tails can be added by transcription directly from PCR products. Poly A may also be ligated to the 3' end of a sense RNA with RNA ligase (see, e.g., *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press: 1991 edition)).

In some embodiments, one or more polynucleotides of the MCNA includes a 3' poly(A) tail structure. Typically, the length of the poly-A tail can be at least about 10, 50, 100, 200, 300, 400 at least 500 nucleotides. In some embodiments, a poly-A tail on the 3' terminus of MCNA typically includes about 10 to 300 adenosine nucleotides (e.g., about 10 to 200 adenosine nucleotides, about 10 to 150 adenosine nucleotides, about 10 to 100 adenosine nucleotides, about 20 to 70 adenosine nucleotides, or about 20 to 60 adenosine nucleotides). In some embodiments, MCNA include a 3' poly-C tail structure. A suitable poly-C tail on the 3' terminus of MCNA typically include about 10 to 200 cytosine nucleotides (e.g., about 10 to 150 cytosine nucleotides, about 10 to 100 cytosine nucleotides, about 20 to 70 cytosine nucleotides, about 20 to 60 cytosine nucleotides, or about 10 to 40 cytosine nucleotides). The poly-C tail may be added to the poly-A tail or may substitute the poly-A tail.

In some embodiments, the length of the poly-A or poly-C tail is adjusted to control the stability of a modified sense MCNA molecule of the invention and, thus, the transcription of protein that is coded for by one or more of the encoding polynucleotides of the MCNA. For example, since the length of the poly-A tail can influence the half-life of a sense MCNA molecule, the length of the poly-A tail can be adjusted to modify the level of resistance of the MCNA to nucleases and thereby control the time course of polynucleotide expression and/or polypeptide production in a target cell.

5' UTR

In some embodiments, MCNA include a 5' cap structure. A 5' cap is typically added as follows: first, an RNA terminal phosphatase removes one of the terminal phosphate groups from the 5' nucleotide, leaving two terminal phosphates; guanosine triphosphate (GTP) is then added to the terminal phosphates via a guanylyl transferase, producing a 5'5'5' triphosphate linkage; and the 7-nitrogen of guanine is then methylated by a methyltransferase. Examples of cap structures include, but are not limited to, m⁷G(5')ppp(5')G(5')ppp(5')A and G(5')ppp(5')G.

Naturally occurring cap structures comprise a 7-methyl guanosine that is linked via a triphosphate bridge to the 5'-end of the first transcribed nucleotide, resulting in a dinucleotide cap of m⁷G(5')ppp(5')N, where N is any nucleoside. In vivo, the cap is added enzymatically. The cap is added in the nucleus and is catalyzed by the enzyme guanylyl transferase. The addition of the cap to the 5' terminal end of RNA occurs immediately after initiation of transcription. The terminal nucleoside is typically a guanos-

ine, and is in the reverse orientation to all the other nucleotides, i.e., G(5')ppp(5')GpNpNp.

One cap for MCNA produced by in vitro transcription is m⁷G(5')ppp(5')G, which has been used as the dinucleotide cap in transcription with T7 or SP6 RNA polymerase in vitro to obtain MCNA having a cap structure in their 5'-termini. A method for the in vitro synthesis of capped MCNA employs a pre-formed dinucleotide of the form m⁷G(5')ppp(5')G (“m⁷GpppG”) as an initiator of transcription.

To date, a usual form of a synthetic dinucleotide cap used in in vitro translation experiments is the Anti-Reverse Cap Analog (“ARCA”) or modified ARCA, which is generally a modified cap analog in which the 2' or 3' OH group is replaced with —OCH₃.

Additional cap analogs include, but are not limited to, a chemical structures selected from the group consisting of m⁷GpppG, m⁷GpppA, m⁷GpppC; unmethylated cap analogs (e.g., GpppG); dimethylated cap analog (e.g., m^{2,7}GpppG), trimethylated cap analog (e.g., m^{2,2,7}GpppG), dimethylated symmetrical cap analogs (e.g., m⁷Gpppm⁷G), or anti reverse cap analogs (e.g., ARCA; m^{7,2'}OmeGpppG, m^{7,2'd}GpppG, m^{7,3'}OmeGpppG, m^{7,3'd}m GpppG and their tetraphosphate derivatives) (see, e.g., Jemielity, J. et al., “Novel ‘anti-reverse’ cap analogs with superior translational properties”, RNA, 9: 1108-1122 (2003)).

In some embodiments, a suitable cap is a 7-methyl guanylate (“m⁷G”) linked via a triphosphate bridge to the 5'-end of the first transcribed nucleotide, resulting in m⁷G(5')ppp(5')N, where N is any nucleoside. A preferred embodiment of a m⁷G cap utilized in embodiments of the invention is m⁷G(5')ppp(5')G.

In some embodiments, the cap is a Cap0 structure. Cap0 structures lack a 2'-O-methyl residue of the ribose attached to bases 1 and 2. In some embodiments, the cap is a Cap1 structure. Cap1 structures have a 2'-O-methyl residue at base 2. In some embodiments, the cap is a Cap2 structure. Cap2 structures have a 2'-O-methyl residue attached to both bases 2 and 3.

A variety of m⁷G cap analogs are known in the art, many of which are commercially available. These include the m⁷GpppG described above, as well as the ARCA 3'-OCH₃ and 2'-OCH₃ cap analogs (Jemielity, J. et al., RNA, 9: 1108-1122 (2003)). Additional cap analogs for use in embodiments of the invention include N7-benzylated dinucleoside tetraphosphate analogs (described in Grudzien, E. et al., RNA, 10: 1479-1487 (2004)), phosphorothioate cap analogs (described in Grudzien-Nogalska, E., et al., RNA, 13: 1745-1755 (2007)), and cap analogs (including biotinylated cap analogs) described in U.S. Pat. Nos. 8,093,367 and 8,304,529, incorporated by reference herein.

Nucleotide Modifications

In some embodiments, MCNA according to the present invention may be synthesized as unmodified or modified nucleic acid. Typically, nucleic acids are modified to enhance stability. Modifications of MCNA can include, for example, modifications of the nucleotides of the MCNA. A modified MCNA according to the invention can thus include, for example, backbone modifications, sugar modifications or base modifications. In some embodiments, MCNA may be synthesized from naturally occurring nucleotides and/or nucleotide analogues (modified nucleotides) including, but not limited to, purines (adenine (A), guanine (G)) or pyrimidines (thymine (T), cytosine (C), uracil (U)), and as modified nucleotides analogues or derivatives of purines and pyrimidines, such as, e.g. 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine,

2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (ΨU), and 1-methyl-pseudouridine, 1-methyl-adenine, 2-methyl-adenine, 2-methylthio-N-6-isopentenyl-adenine, N6-methyl-adenine, N6-isopentenyl-adenine, 2-thio-cytosine, 3-methyl-cytosine, 4-acetyl-cytosine, 5-methyl-cytosine, 2,6-diaminopurine, 1-methyl-guanine, 2-methyl-guanine, 2,2-dimethyl-guanine, 7-methyl-guanine, inosine, 1-methyl-inosine, pseudouracil (5-uracil), dihydro-uracil, 2-thio-uracil, 4-thio-uracil, 5-carboxymethylaminomethyl-2-thio-uracil, 5-(carboxyhydroxymethyl)-uracil, 5-fluoro-uracil, 5-bromo-uracil, 5-carboxymethylaminomethyl-uracil, 5-methyl-2-thio-uracil, 5-methyl-uracil, N-uracil-5-oxyacetic acid methyl ester, 5-methylaminomethyl-uracil, 5-methoxyaminomethyl-2-thio-uracil, 5'-methoxycarbonyl-methyl-uracil, 5-methoxy-uracil, uracil-5-oxyacetic acid methyl ester, uracil-5-oxyacetic acid (v), 1-methyl-pseudouracil, queosine, .beta.-D-mannosyl-queosine, wybutoxosine, and phosphoramidates, phosphorothioates, peptide nucleotides, methylphosphonates, 7-deazaguanosine, 5-methylcytosine and inosine. The preparation of such analogues is known to a person skilled in the art e.g., from the U.S. Pat. Nos. 4,373,071, 4,401,796, 4,415,732, 4,458,066, 4,500,707, 4,668,777, 4,973,679, 5,047,524, 5,132,418, 5,153,319, 5,262,530 and 5,700,642, the disclosures of which are incorporated by reference in their entirety.

In some embodiments, MCNA of the of the present invention comprise encoding polynucleotides that comprise one or more modified nucleotides. In some embodiments, the one or more modified nucleotides are selected from the group consisting of 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (ΨU), and 1-methyl-pseudouridine. In some embodiments, the modified nucleotides substitute 1-100% of corresponding native bases. In some embodiments, at least 25% of uridines are replaced with 2-thiouridines. In some embodiments, 100% cytidines are replaced with 5-methylcytidines. In some embodiments, modified nucleotides are further modified with a 4'-thio substitution on the ribose ring. In some embodiments, native nucleotides are modified with a 4'-thio substitution on the ribose ring.

In some embodiments, MCNA may contain nucleic acid backbone modifications. Typically, a backbone modification is a modification in which the phosphates of the backbone of the nucleotides contained in the MCNA are modified chemically. Exemplary backbone modifications typically include, but are not limited to, modifications from the group consisting of methylphosphonates, methylphosphoramidates, phosphoramidates, phosphorothioates (e.g. cytidine 5'-O-(1-thiophosphate)), boranophosphates, positively charged guanidinium groups etc., which means by replacing the phosphodiester linkage by other anionic, cationic or neutral groups.

In some embodiments, MCNA may contain sugar modifications. A typical sugar modification is a chemical modification of the sugar of the nucleotides it contains including, but not limited to, sugar modifications chosen from the group consisting of 2'-deoxy-2'-fluoro-oligoribonucleotide (2'-fluoro-2'-deoxycytidine 5'-triphosphate, 2'-fluoro-2'-deoxyuridine 5'-triphosphate), 2'-deoxy-2'-deamine-oligoribonucleotide (2'-amino-2'-deoxycytidine 5'-triphosphate, 2'-amino-2'-deoxyuridine 5'-triphosphate), 2'-O-alkyloligoribonucleotide, 2'-deoxy-2'-C-alkyloligoribonucleotide (2'-O-methylcytidine 5'-triphosphate, 2'-methyluridine 5'-triphosphate), 2'-C-alkyloligoribonucleotide, and isomers thereof (2'-aracytidine 5'-triphosphate, 2'-arauridine 5'-tri-

phosphate), or azidotriphosphates (2'-azido-2'-deoxycytidine 5'-triphosphate, 2'-azido-2'-deoxyuridine 5'-triphosphate).

In some embodiments, MCNA may contain modifications of the bases of the nucleotides (base modifications). A modified nucleotide which contains a base modification is also called a base-modified nucleotide. Examples of such base-modified nucleotides include, but are not limited to, 2-amino-6-chloropurine riboside 5'-triphosphate, 2-amino-adenosine 5'-triphosphate, 2-thiocytidine 5'-triphosphate, 2-thiouridine 5'-triphosphate, 4-thiouridine 5'-triphosphate, 5-aminoallylcytidine 5'-triphosphate, 5-aminoallyluridine 5'-triphosphate, 5-bromocytidine 5'-triphosphate, 5-bromouridine 5'-triphosphate, 5-iodocytidine 5'-triphosphate, 5-iodouridine 5'-triphosphate, 5-methylcytidine 5'-triphosphate, 5-methyluridine 5'-triphosphate, 6-azacytidine 5'-triphosphate, 6-azauridine 5'-triphosphate, 6-chloropurine riboside 5'-triphosphate, 7-deazaadenosine 5'-triphosphate, 7-deazaguanosine 5'-triphosphate, 8-azaadenosine 5'-triphosphate, 8-azidoadenosine 5'-triphosphate, benzimidazole riboside 5'-triphosphate, N1-methyladenosine 5'-triphosphate, N1-methylguanosine 5'-triphosphate, N6-methyladenosine 5'-triphosphate, O6-methylguanosine 5'-triphosphate, pseudouridine 5'-triphosphate, puromycin 5'-triphosphate or xanthosine 5'-triphosphate. In some embodiments, MCNA comprises modified bases selected from 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (Ψ U), and 1-methyl-pseudouridine.

Delivery Vehicles

According to the present invention, MCNA as described herein may be delivered as naked polynucleotides or via delivery vehicles. As used herein, the terms "delivery vehicle", "transfer vehicle", "nanoparticle" or grammatical equivalent, are used interchangeably.

In some embodiments, MCNA may be delivered via a single delivery vehicle. In some embodiments, MCNA may be delivered via one or more delivery vehicles each of a different composition. According to various embodiments, suitable delivery vehicles include, but are not limited to polymer based carriers, such as polyethyleneimine (PEI), lipid nanoparticles and liposomes, nanoliposomes, ceramide-containing nanoliposomes, proteoliposomes, both natural and synthetically-derived exosomes, natural, synthetic and semi-synthetic lamellar bodies, nanoparticulates, calcium phosphor-silicate nanoparticulates, calcium phosphate nanoparticulates, silicon dioxide nanoparticulates, nanocrystalline particulates, semiconductor nanoparticulates, poly(D-arginine), sol-gels, nanodendrimers, starch-based delivery systems, micelles, emulsions, niosomes, multi-domain-block polymers (vinyl polymers, polypropyl acrylic acid polymers, dynamic polyconjugates), dry powder formulations, plasmids, viruses, calcium phosphate nucleotides, aptamers, peptides and other vectorial tags.

Liposomal Delivery Vehicles

In some embodiments, a suitable delivery vehicle is a liposomal delivery vehicle, e.g., a lipid nanoparticle. As used herein, liposomal delivery vehicles, e.g., lipid nanoparticles, are usually characterized as microscopic vesicles having an interior aqua space sequestered from an outer medium by a membrane of one or more bilayers. Bilayer membranes of liposomes are typically formed by amphiphilic molecules, such as lipids of synthetic or natural origin that comprise spatially separated hydrophilic and hydrophobic domains (Lasic, Trends Biotechnol., 16: 307-321, 1998). Bilayer membranes of the liposomes can also be formed by


amphiphilic polymers and surfactants (e.g., polymerosomes, niosomes, etc.). In the context of the present invention, a liposomal delivery vehicle typically serves to transport a desired MCNA to a target cell or tissue.

Cationic Lipids

In some embodiments, liposomes may comprise one or more cationic lipids. As used herein, the phrase "cationic lipid" refers to any of a number of lipid species that have a net positive charge at a selected pH, such as physiological pH. Several cationic lipids have been described in the literature, many of which are commercially available. Particularly suitable cationic lipids for use in the compositions and methods of the invention include those described in international patent publications WO 2010/053572 (and particularly, CI 2-200 described at paragraph [00225]) and WO 2012/170930, both of which are incorporated herein by reference. In certain embodiments, the compositions and methods of the invention employ a lipid nanoparticles comprising an ionizable cationic lipid described in U.S. provisional patent application 61/617,468, filed Mar. 29, 2012 (incorporated herein by reference), such as, e.g., (15Z, 18Z)—N,N-dimethyl-6-(9Z, 12Z)-octadeca-9, 12-dien-1-yl) tetracos-15,18-dien-1-amine (HGT5000), (15Z, 18Z)—N,N-dimethyl-6-((9Z, 12Z)-octadeca-9, 12-dien-1-yl)tetracos-4,15,18-trien-1-amine (HGT5001), and (15Z,18Z)—N,N-dimethyl-6-((9Z, 12Z)-octadeca-9, 12-dien-1-yl) tetracos-5, 15, 18-trien-1-amine (HGT5002).

In some embodiments, provided liposomes include a cationic lipid described in WO 2013/063468 and in U.S. provisional application entitled "Lipid Formulations for Delivery of Messenger RNA" filed concurrently with the present application on even date, both of which are incorporated by reference herein.

In some embodiments, a cationic lipid comprises a compound of formula I-c1-a:



or a pharmaceutically acceptable salt thereof, wherein:
 each R^2 independently is hydrogen or C_{1-3} alkyl;
 each q independently is 2 to 6;
 each R' independently is hydrogen or C_{1-3} alkyl;
 and each R^L independently is C_{8-12} alkyl.

In some embodiments, each R^2 independently is hydrogen, methyl or ethyl. In some embodiments, each R^2 independently is hydrogen or methyl. In some embodiments, each R^2 is hydrogen.

101

In some embodiments, each q independently is 3 to 6. In some embodiments, each q independently is 3 to 5. In some embodiments, each q is 4.

In some embodiments, each R' independently is hydrogen, methyl or ethyl. In some embodiments, each R' independently is hydrogen or methyl. In some embodiments, each R' independently is hydrogen.


In some embodiments, each R^L independently is C8-12 alkyl. In some embodiments, each R^L independently is n -C₈₋₁₂ alkyl. In some embodiments, each R^L independently is C₉₋₁₁ alkyl. In some embodiments, each R^L independently is n -C₉₋₁₁ alkyl. In some embodiments, each R^L independently is C₁₀ alkyl. In some embodiments, each R^L independently is n -C₁₀ alkyl.

In some embodiments, each R^2 independently is hydrogen or methyl; each q independently is 3 to 5; each R' independently is hydrogen or methyl; and each R^L independently is C₈₋₁₂ alkyl.

In some embodiments, each R^2 is hydrogen; each q independently is 3 to 5; each R' is hydrogen; and each R^L independently is C₈₋₁₂ alkyl.

In some embodiments, each R^2 is hydrogen; each q is 4; each R' is hydrogen; and each R^L independently is C₈₋₁₂ alkyl.


In some embodiments, a cationic lipid comprises a compound of formula I-g:




or a pharmaceutically acceptable salt thereof, wherein each R^L independently is C₈₋₁₂ alkyl. In some embodiments, each R^L independently is n -C₈₋₁₂ alkyl. In some embodiments, each R^L independently is C₉₋₁₁ alkyl. In some embodiments, each R^L independently is n -C₉₋₁₁ alkyl. In some embodiments, each R^L independently is C₁₀ alkyl. In some embodiments, each R^L is n -C₁₀ alkyl.

In particular embodiments, provided liposomes include a cationic lipid cKK-E12, or (3,6-bis(4-(bis(2-hydroxydecyl)amino)butyl)piperazine-2,5-dione). The structure of cKK-E12 is shown below:

102




Additional exemplary cationic lipids include those of formula I:




and pharmaceutically acceptable salts thereof,

wherein,


R is



R is




R is



or

R is



(see, e.g., Fenton, Owen S., et al. “Bioinspired Alkenyl Amino Alcohol Ionizable Lipid Materials for Highly Potent In Vivo mRNA Delivery.” *Advanced materials* (2016)).

In some embodiments, the one or more cationic lipids may be N-[1-(2,3-dioleoyloxy)propyl]-N,N,N-trimethylammonium chloride or “DOTMA” (Feigner et al. (Proc. Nat’l Acad. Sci. 84, 7413 (1987); U.S. Pat. No. 4,897,355). DOTMA can be formulated alone or can be combined with the neutral lipid, dioleoylphosphatidyl-ethanolamine or “DOPE” or other cationic or non-cationic lipids into a liposomal transfer vehicle or a lipid nanoparticle, and such liposomes can be used to enhance the delivery of nucleic acids into target cells. Other suitable cationic lipids include, for example, 5-carboxyspermylglycinedioctadecylamide or “DOGS,” 2,3-dioleoyloxy-N-[2(spermine-carboxamido)ethyl]-N,N-dimethyl-1-propanaminium or “DOSPA” (Behr et al. Proc. Nat.’l Acad. Sci. 86, 6982 (1989); U.S. Pat. Nos. 5,171,678; 5,334,761), 1,2-Dioleoyl-3-Dimethylammonium-Propane or “DODAP”, 1,2-Dioleoyl-3-Trimethylammonium-Propane or “DOTAP”.

Additional exemplary cationic lipids also include 1,2-distearoyloxy-N,N-dimethyl-3-aminopropane or “DSDMA”, 1,2-dioleoyloxy-N,N-dimethyl-3-aminopropane or “DODMA”, 1,2-dilinoleyloxy-N,N-dimethyl-3-aminopropane or “DLinDMA”, 1,2-dilinolenyloxy-N,N-dimethyl-3-aminopropane or “DLinDMA”, N-dioleoyl-N,N-dimethylammonium chloride or “DODAC”, N,N-distearyl-N,N-dimethylammonium bromide or “DDAB”, N-(1,2-dimyristyloxyprop-3-yl)-N,N-dimethyl-N-hydroxyethyl ammonium bromide or “DMRIE”, 3-dimethylamino-2-(cholest-5-en-3-beta-oxybutan-4-oxy)-1-(cis,cis-9,12-octadecadienoxy)propane or “CLinDMA”, 2-[5’-(cholest-5-en-3-beta-oxy)-3’-oxapentoxy]-3-dimethyl-1-(cis,cis-9’,12’-octadecadienoxy)propane or “CpLinDMA”, N,N-dimethyl-3,4-dioleoyloxybenzylamine or “DMOBA”, 1,2-N,N’-dioleoylcarbonyl-3-dimethylaminopropane or “DOcarbDAP”, 2,3-Dilinoleoyloxy-N,N-dimethylpropylamine or “DLinDAP”, 1,2-N,N’-Dilinoleylcarbonyl-3-dimethylaminopropane or “DLincarbDAP”, 1,2-Dilinoleoylcarbonyl-3-dimethylaminopropane or “DLinCDAP”, 2,2-dilinoleyl-4-dimethylaminomethyl-[1,3]-dioxolane or “DLin-DMA”, 2,2-dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane or “DLin-K-XTC2-DMA”, and 2-(2,2-di((9Z,12Z)-octadeca-9,12-dien-1-yl)-1,3-dioxolan-4-yl)-N,N-dimethylethanamine (DLin-KC2-DMA)) (See, WO 2010/042877; Semple et al., Nature Biotech. 28: 172-176 (2010)), or mixtures thereof. (Heyes, J., et al., J Controlled Release 107: 276-287 (2005); Morrissey, D V., et al., Nat. Biotechnol. 23(8): 1003-1007 (2005); PCT Publication WO2005/121348A1). In some embodiments, one or more of the cationic lipids comprise at least one of an imidazole, dialkylamino, or guanidinium moiety.

In some embodiments, the one or more cationic lipids may be chosen from XTC (2,2-Dilinoleyl-4-dimethylaminoethyl-[1,3]-dioxolane), MC3 (((6Z,9Z,28Z,31Z)-heptatriaconta-6,9,28,31-tetraen-19-yl 4-(dimethylamino)butanoate), ALNY-100 ((3aR,5s,6aS)—N,N-dimethyl-2,2-di((9Z,12Z)-octadeca-9,12-dienyl)tetrahydro-3aH-cyclopenta[d]

[1,3]dioxol-5-amine)), NC98-5 (4,7,13-tris(3-oxo-3-(undecylamino)propyl)-N1,N16-diundecyl-4,7,10,13-tetraaza-hexadecane-1,16-diamide), DODAP (1,2-dioleoyl-3-dimethylammonium propane), HGT4003 (WO 2012/170889, the teachings of which are incorporated herein by reference in their entirety), ICE (WO 2011/068810, the teachings of which are incorporated herein by reference in their entirety), HGT5000 (U.S. Provisional Patent Application No. 61/617,468, the teachings of which are incorporated herein by reference in their entirety) or HGT5001 (cis or trans) (Provisional Patent Application No. 61/617,468), aminoalcohol lipidoids such as those disclosed in WO2010/053572, DOTAP (1,2-dioleoyl-3-trimethylammonium propane), DOTMA (1,2-di-O-octadecenyl-3-trimethylammonium propane), DLinDMA (Heyes, J.; Palmer, L.; Bremner, K.; MacLachlan, I. “Cationic lipid saturation influences intracellular delivery of encapsulated nucleic acids” J. Contr. Rel. 2005, 107, 276-287), DLin-KC2-DMA (Semple, S. C. et al. “Rational Design of Cationic Lipids for siRNA Delivery” Nature Biotech. 2010, 28, 172-176), C12-200 (Love, K. T. et al. “Lipid-like materials for low-dose in vivo gene silencing” PNAS 2010, 107, 1864-1869).

In some embodiments, the percentage of cationic lipid in a liposome may be greater than 10%, greater than 20%, greater than 30%, greater than 40%, greater than 50%, greater than 60%, or greater than 70%. In some embodiments, cationic lipid(s) constitute(s) about 30-50% (e.g., about 30-45%, about 30-40%, about 35-50%, about 35-45%, or about 35-40%) of the liposome by weight. In some embodiments, the cationic lipid (e.g., cKK-E12) constitutes about 30%, about 35%, about 40%, about 45%, or about 50% of the liposome by molar ratio.

Non-Cationic/Helper Lipids

In some embodiments, provided liposomes contain one or more non-cationic (“helper”) lipids. As used herein, the phrase “non-cationic lipid” refers to any neutral, zwitterionic or anionic lipid. As used herein, the phrase “anionic lipid” refers to any of a number of lipid species that carry a net negative charge at a selected H, such as physiological pH. Non-cationic lipids include, but are not limited to, distearoylphosphatidylcholine (DSPC), dioleoylphosphatidylcholine (DOPC), dipalmitoylphosphatidylcholine (DPPC), dioleoylphosphatidylglycerol (DOPG), dipalmitoylphosphatidylglycerol (DPPG), dioleoylphosphatidylethanolamine (DOPE), palmitoyloleoylphosphatidylcholine (POPC), palmitoyloleoyl-phosphatidylethanolamine (POPE), dioleoyl-phosphatidylethanolamine 4-(N-maleimidomethyl)-cyclohexane-1-carboxylate (DOPE-mal), dipalmitoyl phosphatidyl ethanolamine (DPPE), dimyristoylphosphoethanolamine (DMPE), distearoyl-phosphatidyl-ethanolamine (DSPE), 16-O-monomethyl PE, 16-O-dimethyl PE, 18-1-trans PE, 1-stearoyl-2-oleoyl-phosphatidylethanolamine (SOPE), or a mixture thereof.

In some embodiments, such non-cationic lipids may be used alone, but are preferably used in combination with other excipients, for example, cationic lipids. In some embodiments, the non-cationic lipid may comprise a molar ratio of about 5% to about 90%, or about 10% to about 70% of the total lipid present in a liposome. In some embodiments, a non-cationic lipid is a neutral lipid, i.e., a lipid that does not carry a net charge in the conditions under which the composition is formulated and/or administered. In some embodiments, the percentage of non-cationic lipid in a liposome may be greater than 5%, greater than 10%, greater than 20%, greater than 30%, or greater than 40%.

Cholesterol-Based Lipids

In some embodiments, provided liposomes comprise one or more cholesterol-based lipids. For example, suitable cholesterol-based cationic lipids include, for example, DC-Choi (N,N-dimethyl-N-ethylcarboxamidcholesterol), 1,4-bis(3-N-oleylamino-propyl)piperazine (Gao, et al. *Biochem. Biophys. Res. Comm.* 179, 280 (1991); Wolf et al. *BioTechniques* 23, 139 (1997); U.S. Pat. No. 5,744,335), or ICE. In some embodiments, the cholesterol-based lipid may comprise a molar ration of about 2% to about 30%, or about 5% to about 20% of the total lipid present in a liposome. In some embodiments, The percentage of cholesterol-based lipid in the lipid nanoparticle may be greater than 5, %, 10%, greater than 20%, greater than 30%, or greater than 40%.

PEGylated Lipids

In some embodiments, provided liposomes comprise one or more PEGylated lipids. For example, the use of polyethylene glycol (PEG)-modified phospholipids and derivatized lipids such as derivatized ceramides (PEG-CER), including N-Octanoyl-Sphingosine-1-[Succinyl(Methoxy Polyethylene Glycol)-2000] (C8 PEG-2000 ceramide) is also contemplated by the present invention in combination with one or more of the cationic and, in some embodiments, other lipids together which comprise the liposome. Contemplated PEG-modified lipids include, but are not limited to, a polyethylene glycol chain of up to 5 kDa in length covalently attached to a lipid with alkyl chain(s) of C₆-C₂₀ length. In some embodiments, a PEG-modified or PEGylated lipid is PEGylated cholesterol or PEG-2K. The addition of such components may prevent complex aggregation and may also provide a means for increasing circulation lifetime and increasing the delivery of the lipid-nucleic acid composition to the target cell, (Klibanov et al. (1990) *FEBS Letters*, 268 (1): 235-237), or they may be selected to rapidly exchange out of the formulation in vivo (see U.S. Pat. No. 5,885,613).

In some embodiments, particularly useful exchangeable lipids are PEG-ceramides having shorter acyl chains (e.g., C₁₄ or C₁₈). The PEG-modified phospholipid and derivitized lipids of the present invention may comprise a molar ratio from about 0% to about 15%, about 0.5% to about 15%, about 1% to about 15%, about 4% to about 10%, or about 2% of the total lipid present in the liposome.

According to various embodiments, the selection of cationic lipids, non-cationic lipids and/or PEG-modified lipids which comprise the lipid nanoparticle, as well as the relative molar ratio of such lipids to each other, is based upon the characteristics of the selected lipid(s), the nature of the intended target cells, the characteristics of the MCNA to be delivered. Additional considerations include, for example, the saturation of the alkyl chain, as well as the size, charge, pH, pKa, fusogenicity and toxicity of the selected lipid(s). Thus the molar ratios may be adjusted accordingly.

Formation of Liposomes

The liposomal transfer vehicles for use in the compositions of the invention can be prepared by various techniques which are presently known in the art. The liposomes for use in provided compositions can be prepared by various techniques which are presently known in the art. For example, multilamellar vesicles (MLV) may be prepared according to conventional techniques, such as by depositing a selected lipid on the inside wall of a suitable container or vessel by dissolving the lipid in an appropriate solvent, and then evaporating the solvent to leave a thin film on the inside of the vessel or by spray drying. An aqueous phase may then added to the vessel with a vortexing motion which results in the formation of MLVs. Unilamellar vesicles (ULV) can then be formed by homogenization, sonication or extrusion

of the multilamellar vesicles. In addition, unilamellar vesicles can be formed by detergent removal techniques.

In certain embodiments, provided compositions comprise a liposome wherein the MCNA is associated on both the surface of the liposome and encapsulated within the same liposome. For example, during preparation of the compositions of the present invention, cationic liposomes may associate with the MCNA through electrostatic interactions. For example, during preparation of the compositions of the present invention, cationic liposomes may associate with the MCNA through electrostatic interactions.

In some embodiments, the compositions and methods of the invention comprise MCNA encapsulated in a liposome. In some embodiments, the one or more MCNA species may be encapsulated in the same liposome. In some embodiments, the one or more MCNA species may be encapsulated in different liposomes. In some embodiments, the MCNA is encapsulated in one or more liposomes, which differ in their lipid composition, molar ratio of lipid components, size, charge (Zeta potential), targeting ligands and/or combinations thereof. In some embodiments, the one or more liposome may have a different composition of cationic lipids, neutral lipid, PEG-modified lipid and/or combinations thereof. In some embodiments the one or more liposomes may have a different molar ratio of cationic lipid, neutral lipid, cholesterol and PEG-modified lipid used to create the liposome.

The process of incorporation of a desired MCNA into a liposome is often referred to as "loading". Exemplary methods are described in Lasic, et al., *FEBS Lett.*, 312: 255-258, 1992, which is incorporated herein by reference. The liposome-incorporated nucleic acids may be completely or partially located in the interior space of the liposome, within the bilayer membrane of the liposome, or associated with the exterior surface of the liposome membrane. The incorporation of a nucleic acid into liposomes is also referred to herein as "encapsulation" wherein the nucleic acid is entirely contained within the interior space of the liposome. The purpose of incorporating a MCNA into a transfer vehicle, such as a liposome, is often to protect the nucleic acid from an environment which may contain enzymes or chemicals that degrade nucleic acids and/or systems or receptors that cause the rapid excretion of the nucleic acids. Accordingly, in some embodiments, a suitable delivery vehicle is capable of enhancing the stability of the MCNA contained therein and/or facilitate the delivery of MCNA to the target cell or tissue.

Liposome Size

Suitable liposomes in accordance with the present invention may be made in various sizes. In some embodiments, provided liposomes may be made smaller than previously known mRNA encapsulating liposomes. In some embodiments, decreased size of liposomes is associated with more efficient delivery of MCNA. Selection of an appropriate liposome size may take into consideration the site of the target cell or tissue and to some extent the application for which the liposome is being made.

In some embodiments, an appropriate size of liposome is selected to facilitate systemic distribution of polypeptide encoded by the MCNA. In some embodiments, it may be desirable to limit transfection of the MCNA to certain cells or tissues. For example, to target hepatocytes a liposome may be sized such that its dimensions are smaller than the fenestrations of the endothelial layer lining hepatic sinusoids in the liver; in such cases the liposome could readily penetrate such endothelial fenestrations to reach the target hepatocytes.

Alternatively or additionally, a liposome may be sized such that the dimensions of the liposome are of a sufficient diameter to limit or expressly avoid distribution into certain cells or tissues. For example, a liposome may be sized such that its dimensions are larger than the fenestrations of the endothelial layer lining hepatic sinusoids to thereby limit distribution of the liposomes to hepatocytes.

In some embodiments, the size of a liposome is determined by the length of the largest diameter of the liposome particle. In some embodiments, a suitable liposome has a size no greater than about 250 nm (e.g., no greater than about 225 nm, 200 nm, 175 nm, 150 nm, 125 nm, 100 nm, 75 nm, or 50 nm). In some embodiments, a suitable liposome has a size ranging from about 10-250 nm (e.g., ranging from about 10-225 nm, 10-200 nm, 10-175 nm, 10-150 nm, 10-125 nm, 10-100 nm, 10-75 nm, or 10-50 nm). In some embodiments, a suitable liposome has a size ranging from about 100-250 nm (e.g., ranging from about 100-225 nm, 100-200 nm, 100-175 nm, 100-150 nm). In some embodiments, a suitable liposome has a size ranging from about 10-100 nm (e.g., ranging from about 10-90 nm, 10-80 nm, 10-70 nm, 10-60 nm, or 10-50 nm). In a particular embodiment, a suitable liposome has a size less than about 100 nm.

A variety of alternative methods known in the art are available for sizing of a population of liposomes. One such sizing method is described in U.S. Pat. No. 4,737,323, incorporated herein by reference. Sonicating a liposome suspension either by bath or probe sonication produces a progressive size reduction down to small ULV less than about 0.05 microns in diameter. Homogenization is another method that relies on shearing energy to fragment large liposomes into smaller ones. In a typical homogenization procedure, MLV are recirculated through a standard emulsion homogenizer until selected liposome sizes, typically between about 0.1 and 0.5 microns, are observed. The size of the liposomes may be determined by quasi-electric light scattering (QELS) as described in Bloomfield, *Ann. Rev. Biophys. Bioeng.*, 10:421-150 (1981), incorporated herein by reference. Average liposome diameter may be reduced by sonication of formed liposomes. Intermittent sonication cycles may be alternated with QELS assessment to guide efficient liposome synthesis.

Polymers

In some embodiments, a suitable delivery vehicle is formulated using a polymer as a carrier, alone or in combination with other carriers including various lipids described herein. Thus, in some embodiments, liposomal delivery vehicles, as used herein, also encompass polymer containing nanoparticles. Suitable polymers may include, for example, polyacrylates, polyalkylcyanoacrylates, polylactide, polylactide-polyglycolide copolymers, polycaprolactones, dextran, albumin, gelatin, alginate, collagen, chitosan, cyclodextrins, protamine, PEGylated protamine, PLL, PEGylated PLL and polyethylenimine (PEI). When PEI is present, it may be branched PEI of a molecular weight ranging from 10 to 40 kDa, e.g., 25 kDa branched PEI (Sigma #408727).

A suitable liposome for the present invention may include one or more of any of the cationic lipids, non-cationic lipids, cholesterol lipids, PEGylated lipids and/or polymers described herein at various ratios. As non-limiting examples, a suitable liposome formulation may include a combination selected from cKK-E12, DOPE, cholesterol and DMG-PEG2K; C12-200, DOPE, cholesterol and DMG-PEG2K; HGT4003, DOPE, cholesterol and DMG-PEG2K; or ICE, DOPE, cholesterol and DMG-PEG2K.

In various embodiments, cationic lipids (e.g., cKK-E12, C12-200, ICE, and/or HGT4003) constitute about 30-60%

(e.g., about 30-55%, about 30-50%, about 30-45%, about 30-40%, about 35-50%, about 35-45%, or about 35-40%) of the liposome by molar ratio. In some embodiments, the percentage of cationic lipids (e.g., cKK-E12, C12-200, ICE, and/or HGT4003) is or greater than about 30%, about 35%, about 40%, about 45%, about 50%, about 55%, or about 60% of the liposome by molar ratio.

In some embodiments, the ratio of cationic lipid(s) to non-cationic lipid(s) to cholesterol-based lipid(s) to PEGylated lipid(s) may be between about 30-60:25-35:20-30:1-15, respectively. In some embodiments, the ratio of cationic lipid(s) to non-cationic lipid(s) to cholesterol-based lipid(s) to PEGylated lipid(s) is approximately 40:30:20:10, respectively. In some embodiments, the ratio of cationic lipid(s) to non-cationic lipid(s) to cholesterol-based lipid(s) to PEGylated lipid(s) is approximately 40:30:25:5, respectively. In some embodiments, the ratio of cationic lipid(s) to non-cationic lipid(s) to cholesterol-based lipid(s) to PEGylated lipid(s) is approximately 40:32:25:3, respectively. In some embodiments, the ratio of cationic lipid(s) to non-cationic lipid(s) to cholesterol-based lipid(s) to PEGylated lipid(s) is approximately 50:25:20:5.

Pharmaceutical Compositions

To facilitate expression of MCNA in vivo, delivery vehicles such as liposomes can be formulated in combination with one or more additional nucleic acids, carriers, targeting ligands or stabilizing reagents, or in pharmacological compositions where it is mixed with suitable excipients. Techniques for formulation and administration of drugs may be found in "Remington's Pharmaceutical Sciences," Mack Publishing Co., Easton, Pa., latest edition.

In some embodiments, a composition comprises MCNA encapsulated or complexed with a delivery vehicle. In some embodiments, the delivery vehicle is selected from the group consisting of liposomes, lipid nanoparticles, solid-lipid nanoparticles, polymers, viruses, sol-gels, and nanogels.

Provided liposomally-encapsulated or liposomally-associated MCNA, and compositions containing the same, may be administered and dosed in accordance with current medical practice, taking into account the clinical condition of the subject, the site and method of administration, the scheduling of administration, the subject's age, sex, body weight and other factors relevant to clinicians of ordinary skill in the art. The "effective amount" for the purposes herein may be determined by such relevant considerations as are known to those of ordinary skill in experimental clinical research, pharmacological, clinical and medical arts. In some embodiments, the amount administered is effective to achieve at least some stabilization, improvement or elimination of symptoms and other indicators as are selected as appropriate measures of disease progress, regression or improvement by those of skill in the art. For example, a suitable amount and dosing regimen is one that causes at least transient protein (e.g., enzyme) production.

The present invention provides methods of delivering MCNA for in vivo protein production, comprising administering MCNA to a subject in need of delivery. In some embodiments, MCNA is administered via a route of delivery selected from the group consisting of intravenous delivery, subcutaneous delivery, oral delivery, subdermal delivery, ocular delivery, intratracheal injection pulmonary delivery (e.g. nebulization), intramuscular delivery, intrathecal delivery, or intraarticular delivery.

Suitable routes of administration include, for example, oral, rectal, vaginal, transmucosal, pulmonary including intratracheal or inhaled, or intestinal administration; paren-

teral delivery, including intradermal, transdermal (topical), intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, or intranasal. In particular embodiments, the intramuscular administration is to a muscle selected from the group consisting of skeletal muscle, smooth muscle and cardiac muscle. In some embodiments the administration results in delivery of the MCNA to a muscle cell. In some embodiments the administration results in delivery of the MCNA to a hepatocyte (i.e., liver cell). In a particular embodiment, the intramuscular administration results in delivery of the MCNA to a muscle cell.

Alternatively or additionally, liposomally-encapsulated MCNA and compositions of the invention may be administered in a local rather than systemic manner, for example, via injection of the pharmaceutical composition directly into a targeted tissue, preferably in a sustained release formulation. Local delivery can be affected in various ways, depending on the tissue to be targeted. For example, aerosols containing compositions of the present invention can be inhaled (for nasal, tracheal, or bronchial delivery); compositions of the present invention can be injected into the site of injury, disease manifestation, or pain, for example; compositions can be provided in lozenges for oral, tracheal, or esophageal application; can be supplied in liquid, tablet or capsule form for administration to the stomach or intestines, can be supplied in suppository form for rectal or vaginal application; or can even be delivered to the eye by use of creams, drops, or even injection. Formulations containing provided compositions complexed with therapeutic molecules or ligands can even be surgically administered, for example in association with a polymer or other structure or substance that can allow the compositions to diffuse from the site of implantation to surrounding cells. Alternatively, they can be applied surgically without the use of polymers or supports.

Provided methods of the present invention contemplate single as well as multiple administrations of a therapeutically effective amount of the therapeutic agents (e.g., MCNA) described herein. Therapeutic agents can be administered at regular intervals, depending on the nature, severity and extent of the subject's condition. In some embodiments, a therapeutically effective amount of the therapeutic agents (e.g., MCNA) of the present invention may be administered intrathecally periodically at regular intervals (e.g., once every year, once every six months, once every five months, once every three months, bimonthly (once every two months), monthly (once every month), biweekly (once every two weeks), twice a month, once every 30 days, once every 28 days, once every 14 days, once every 10 days, once every 7 days, weekly, twice a week, daily or continuously).

In some embodiments, provided liposomes and/or compositions are formulated such that they are suitable for extended-release of the MCNA contained therein. Such extended-release compositions may be conveniently administered to a subject at extended dosing intervals. For example, in one embodiment, the compositions of the present invention are administered to a subject twice a day, daily or every other day. In a preferred embodiment, the compositions of the present invention are administered to a subject twice a week, once a week, once every 7 days, once every 10 days, once every 14 days, once every 28 days, once every 30 days, once every two weeks, once every three weeks, or more preferably once every four weeks, once a month, twice a month, once every six weeks, once every eight weeks, once every other month, once every three months, once every four months, once every six months, once every eight

months, once every nine months or annually. Also contemplated are compositions and liposomes which are formulated for depot administration (e.g., intramuscularly, subcutaneously, intravitreally) to either deliver or release MCNA over extended periods of time. Preferably, the extended-release means employed are combined with modifications made to the MCNA to enhance stability.

As used herein, the term "therapeutically effective amount" is largely determined based on the total amount of the therapeutic agent contained in the pharmaceutical compositions of the present invention. Generally, a therapeutically effective amount is sufficient to achieve a meaningful benefit to the subject (e.g., treating, modulating, curing, preventing and/or ameliorating a disease or disorder). For example, a therapeutically effective amount may be an amount sufficient to achieve a desired therapeutic and/or prophylactic effect. Generally, the amount of a therapeutic agent (e.g., MCNA) administered to a subject in need thereof will depend upon the characteristics of the subject. Such characteristics include the condition, disease severity, general health, age, sex and body weight of the subject. One of ordinary skill in the art will be readily able to determine appropriate dosages depending on these and other related factors. In addition, both objective and subjective assays may optionally be employed to identify optimal dosage ranges.

A therapeutically effective amount is commonly administered in a dosing regimen that may comprise multiple unit doses. For any particular therapeutic protein, a therapeutically effective amount (and/or an appropriate unit dose within an effective dosing regimen) may vary, for example, depending on route of administration, on combination with other pharmaceutical agents. Also, the specific therapeutically effective amount (and/or unit dose) for any particular patient may depend upon a variety of factors including the disorder being treated and the severity of the disorder; the activity of the specific pharmaceutical agent employed; the specific composition employed; the age, body weight, general health, sex and diet of the patient; the time of administration, route of administration, and/or rate of excretion or metabolism of the specific protein employed; the duration of the treatment; and like factors as is well known in the medical arts.

In some embodiments, the therapeutically effective dose ranges from about 0.005 mg/kg body weight to 500 mg/kg body weight, e.g., from about 0.005 mg/kg body weight to 400 mg/kg body weight, from about 0.005 mg/kg body weight to 300 mg/kg body weight, from about 0.005 mg/kg body weight to 200 mg/kg body weight, from about 0.005 mg/kg body weight to 100 mg/kg body weight, from about 0.005 mg/kg body weight to 90 mg/kg body weight, from about 0.005 mg/kg body weight to 80 mg/kg body weight, from about 0.005 mg/kg body weight to 70 mg/kg body weight, from about 0.005 mg/kg body weight to 60 mg/kg body weight, from about 0.005 mg/kg body weight to 50 mg/kg body weight, from about 0.005 mg/kg body weight to 40 mg/kg body weight, from about 0.005 mg/kg body weight to 30 mg/kg body weight, from about 0.005 mg/kg body weight to 25 mg/kg body weight, from about 0.005 mg/kg body weight to 20 mg/kg body weight, from about 0.005 mg/kg body weight to 15 mg/kg body weight, from about 0.005 mg/kg body weight to 10 mg/kg body weight.

In some embodiments, the therapeutically effective dose is greater than about 0.1 mg/kg body weight, greater than about 0.5 mg/kg body weight, greater than about 1.0 mg/kg body weight, greater than about 3 mg/kg body weight, greater than about 5 mg/kg body weight, greater than about

10 mg/kg body weight, greater than about 15 mg/kg body weight, greater than about 20 mg/kg body weight, greater than about 30 mg/kg body weight, greater than about 40 mg/kg body weight, greater than about 50 mg/kg body weight, greater than about 60 mg/kg body weight, greater than about 70 mg/kg body weight, greater than about 80 mg/kg body weight, greater than about 90 mg/kg body weight, greater than about 100 mg/kg body weight, greater than about 150 mg/kg body weight, greater than about 200 mg/kg body weight, greater than about 250 mg/kg body weight, greater than about 300 mg/kg body weight, greater than about 350 mg/kg body weight, greater than about 400 mg/kg body weight, greater than about 450 mg/kg body weight, greater than about 500 mg/kg body weight. In a particular embodiment, the therapeutically effective dose is 1.0 mg/kg. In some embodiments, the therapeutically effective dose of 1.0 mg/kg is administered intramuscularly or intravenously.

Also contemplated herein are lyophilized pharmaceutical compositions comprising one or more of the liposomes disclosed herein and related methods for the use of such compositions as disclosed for example, in U.S. Provisional Application No. 61/494,882, filed Jun. 8, 2011, the teachings of which are incorporated herein by reference in their entirety. For example, lyophilized pharmaceutical compositions according to the invention may be reconstituted prior to administration or can be reconstituted in vivo. For example, a lyophilized pharmaceutical composition can be formulated in an appropriate dosage form (e.g., an intradermal dosage form such as a disk, rod or membrane) and administered such that the dosage form is rehydrated over time in vivo by the individual's bodily fluids.

Provided liposomes and compositions may be administered to any desired tissue. In some embodiments, the MCNA delivered by provided liposomes or compositions is expressed in the tissue in which the liposomes and/or compositions were administered. In some embodiments, the MCNA delivered is expressed in a tissue different from the tissue in which the liposomes and/or compositions were administered. Exemplary tissues in which delivered MCNA may be delivered and/or expressed include, but are not limited to the liver, kidney, heart, spleen, serum, brain, skeletal muscle, lymph nodes, skin, and/or cerebrospinal fluid.

In some embodiments, administering the provided composition results in an increased MCNA expression level in a biological sample from a subject as compared to a baseline expression level before treatment. Typically, the baseline level is measured immediately before treatment. Biological samples include, for example, whole blood, serum, plasma, urine and tissue samples (e.g., muscle, liver, skin fibroblasts). In some embodiments, administering the provided composition results in an increased MCNA expression level by at least about 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90%, or 95% as compared to the baseline level immediately before treatment. In some embodiments, administering the provided composition results in an increased MCNA expression level as compared to a MCNA expression level in subjects who are not treated

According to various embodiments, the timing of expression of delivered MCNA can be tuned to suit a particular medical need. In some embodiments, the expression of the protein encoded by delivered MCNA is detectable 1, 2, 3, 6, 12, 24, 48, 72, and/or 96 hours after administration of provided liposomes and/or compositions. In some embodi-

ments, the expression of the protein encoded by delivered MCNA is detectable 1 week, two weeks, and/or 1 month after administration.

EXAMPLES

While certain compounds, compositions and methods of the present invention have been described with specificity in accordance with certain embodiments, the following examples serve only to illustrate the compounds of the invention and are not intended to limit the same.

Example 1. Exemplary Synthesis of Multimeric Coding Nucleic Acid (MCNA)

This example provides exemplary schemes for synthesizing the MCNA described in this application, for effective delivery and expression of MCNA encoding therapeutic proteins in vivo.

Synthesis of MCNA was attempted by ligating a synthetic oligonucleotide containing a 3'-3' phosphodiester bond to multiple polynucleotides using a complementary DNA splint. Several different T4 RNA ligases were tested for the ability to ligate a synthetic oligonucleotide containing a 3'-3' phosphodiester bond to multiple polynucleotides using a complementary DNA splint. The first RNA ligase ("RNA Ligase 1") was a "single-strand" RNA ligase that ligated single RNA strands, double RNA strands and double RNA strands designed to implement a single strand overhang. The second RNA ligase ("RNA Ligase 2") was a "double-stranded" RNA ligase that ligated nicks in RNA bound to a complementary oligonucleotide. Both RNA Ligase 1 and RNA Ligase 2 required phosphorylated 5' ends of the oligonucleotide bridge to proceed with adenylation for the ligation reaction.

As a non-limiting example, Erythropoietin (EPO) mRNA was ligated to a bridging oligo containing a 3'-3' phosphodiester bond using a complementary DNA splint. Examples of a bridging oligonucleotide that contains a 3'-3' phosphodiester bond and DNA splints are described below. The exemplary sequence for EPO used in the examples herein are listed below.

Erythropoietin (EPO) mRNA (including 5' and 3' UTR) :

(SEQ ID NO: 1)

GGACAGAUCCGCGGAGACGCCAUCACGCUGUUUGACCUCAUAGA
 AGACACCGGGACCGAUCAGCCUCCGCGCCGGGAACGGUGCAUUGGA
 ACGCGGAUUCGCGGCAAGAGUGACUCACCGUCCUUGACACGAUGG
 GGGUGCACGAAUGUCUGCCUGGCUGUGGCUUCUCCUGUCCUGCUGU
 CGCUCCUCUGGGCCUCCAGUCCUGGGCGCCCCACCACGCCUCAUCU
 GUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAGGCCAAGGAGGCCG
 AGAAUAUCACGACGGGCGUGUGCUGAACACUGCAGCUUGAAUGAGAAUA
 UCACUGUCCAGACACCAAAGUUAUUUCUAUGCCUGGAAGAGGAUGG
 AGGUCGGGCGAGCAGGCCGUAAGAAGUCUGGCAGGGCCUGGCCUGCUGU
 CGGAAGCUGUCCUGCGGGGCCAGGCCUGUUGGUCAACUCUCCAGC
 CGUGGGAGCCCCUGCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUC
 GCAGCCUACACUCUGCUUCGGGCUUGGGAGCCCAGAAGGAAGCCA

-continued

UCUCCCCUCCAGAUGC... GACGACUUUCCGCAAACUCUUC... GAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGGGGACAGAU... GACGGGUGGCAUCCUGUGACCCCUC... GAAGUUGCCACUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGU... UGCAUCAAGCU

Erythropoietin (EPO) mRNA (including 5' and 3' UTR with 200 A poly(A) Tail):

(SEQ ID NO: 2)

GGACAGAU... AGACACCGGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGA... ACGCGGAU... GGGUGCACGAAUGUCCUGCCUGGCUGUGGCUCUCCUGUCCUGCUGU... CGCUCCCUCUGGGCCUCCAGUCCUGGGCGCCCCACCACGCCUCAUCU... GUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAGGCCAAGGAGGCCG... AGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAUGAGAAUA... UCACUGUCCAGACACCAAAGUUAAUUUCU... AGGUCGGGCAGCAGGCCGUAGAAGUCUGGCAGGGCCUGGCCUGCUGU... CGGAAGCUGUCCUGCGGGGCCAGGCCUGUUGGUCAACUCUCCAGC... CGUGGGAGCCCUGCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUC... GCAGCCUACACACUCUGCUUCGGGCUCUGGGAGCCAGAAGGAAGCCA... UCUCCCCUCAGAUCCGCGCCUCAGCUGCUCACUCCGAACAAUCACUG... CUGACACUUUCCGCAAACUCUUC... GAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGGGGACAGAU... GACGGGUGGCAUCCUGUGACCCCUC... GAAGUUGCCACUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGU... UGCAUCAAGCUAAA... AA... AA... AA... AAAAAAAAAAAAAAAAAAAAAAAAAAAAAA

Erythropoietin (EPO) mRNA (including 5' and 3' UTR with internal 65A poly(A) region in 3' UTR):

(SEQ ID NO: 3)

GGACAGAU... AGACACCGGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGA... ACGCGGAU... GGGUGCACGAAUGUCCUGCCUGGCUGUGGCUCUCCUGUCCUGCUGU... CGCUCCCUCUGGGCCUCCAGUCCUGGGCGCCCCACCACGCCUCAUCU... GUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAGGCCAAGGAGGCCG... AGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAUGAGAAUA... UCACUGUCCAGACACCAAAGUUAAUUUCU... AGGUCGGGCAGCAGGCCGUAGAAGUCUGGCAGGGCCUGGCCUGCUGU... CGGAAGCUGUCCUGCGGGGCCAGGCCUGUUGGUCAACUCUCCAGC

-continued

CGUGGGAGCCCUCGAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUC... GCAGCCUACACACUCUGCUUCGGGCUCUGGGAGCCCAGAAGGAAGCCA... 5 UCUCCCCUCCAGAUGC... CUGACACUUUCCGCAAACUCUUC... GAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGGGGACAGAU... 10 GACGGGUGGCAUCCUGUGACCCCUC... GAAGUUGCCACUCCAGUGCCCAC... AAAGCCUUGU... 15 CCUAAUAAAAUUAAGUUGCAUCAAGCU

Erythropoietin (EPO) mRNA (including 5' and 3' UTR with multiple short internal poly(A) regions in 3' UTR):

(SEQ ID NO: 4)

GGACAGAU... AGACACCGGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGA... ACGCGGAU... GGGUGCACGAAUGUCCUGCCUGGCUGUGGCUCUCCUGUCCUGCUGU... 25 CGCUCCCUCUGGGCCUCCAGUCCUGGGCGCCCCACCACGCCUCAUCU... GUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAGGCCAAGGAGGCCG... 30 AGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAUGAGAAUA... UCACUGUCCAGACACCAAAGUUAAUUUCU... AGGUCGGGCAGCAGGCCGUAGAAGUCUGGCAGGGCCUGGCCUGCUGU... 35 CGGAAGCUGUCCUGCGGGGCCAGGCCUGUUGGUCAACUCUCCAGC... CGUGGGAGCCCUCGAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUC... GCAGCCUACACACUCUGCUUCGGGCUCUGGGAGCCCAGAAGGAAGCCA... 40 UCUCCCCUCAGAUCCGCGCCUCAGCUGCUCACUCCGAACAAUCACUG... CUGACACUUUCCGCAAACUCUUC... GAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGGGGACAGAU... 45 GACGGGUGGCAAAAAAAAAAAAAAAAAAUCCUGUGACCCCUC... AAAAAAAAAAAGUCCUCUCCUGGCCUGGAAAAAAAAAAAAAAAAAGUU... GCCACUCCAGUGCCCACCAAAAAAAAAAAAAAAAAAGCCUUGUCCUAAUAA... AAUUAAGUUGCAUCAAGCU

Bridging Oligonucleotide 1:

(SEQ ID NO: 5)

5'-CGA CUC UCG G-3'-PO4-3'-G GCU CUC AGC-5'

The bases included in SEQ ID NO: 5 are 2'-O-methyl RNA and the 3'-3' bridge comprises PO4.

Bridging Oligonucleotide 2:

(SEQ ID NO: 6)

5'-AAAAAAAAA-3'-PO4-3'-AAAAAAAAA-5'

Bridging Oligonucleotide 3:

(SEQ ID NO: 7)

5'-AAA-3'-PO4-3'-AAA-5'

Bridging Oligonucleotide 4:

(SEQ ID NO: 8)

5'-A-3'-PO4-3'-A-5'

-continued

Splint Oligonucleotide 1: (SEQ ID NO: 9)
 5'-CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT
 TAG G-3'

Splint Oligonucleotide 2: (SEQ ID NO: 10)
 5'-CCG AGA GTG ATG CAA CTT AAT TTT ATT AGG-3'

Splint Oligonucleotide 3: (SEQ ID NO: 11)
 5'-TTT TTT TTT TAG CTT GAT GCA ACT TAA TTT TAT
 TAG G-3'

Splint Oligonucleotide 4: (SEQ ID NO: 12)
 5'-CCG AGA GTC GTT TTT TTT TTT TTT TTT TTT-3'

Splint Oligonucleotide 5: (SEQ ID NO: 13)
 3'-G GAT TAT TTT AAT TCA ACG TAG TTC GAG CTG AGA
 GCC-5'-PO₄-5'-CCG AGA GTC GAG CTT GAT GCA ACT
 TAA TTT TAT TAG G-3'

Splint Oligonucleotide 6: (SEQ ID NO: 14)
 3'-GGA TTA TTT TAA TTC AAC GTA GTG AGA GCC-5'-
 PO₄-5'-CCG AGA GTG ATG CAA CTT AAT TTT ATT
 AGG-3'

Splint Oligonucleotide 7: (SEQ ID NO: 15)
 3'-G GAT TAT TTT AAT TCA ACG TAG TTC GAT TTT TTT
 TTT-5'-PO₄-5'-TTT TTT TTT TAG CTT GAT GCA ACT
 TAA TTT TAT TAG G-3'

Splint Oligonucleotide 8: (SEQ ID NO: 16)
 3'-TTT TTT TTT TTT TTT TTT TTG CTG AGA
 GCC-5'-PO₄-5'-CCG AGA GTC GTT TTT TTT TTT TTT
 TTT TTT-3'

EPO MCNA #1 (No Poly A Tail)

MCNA 1 (SEQ ID NO: 17) was prepared by splint ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5' untranslated region (UTR) and a 3' UTR flanking an RNA sequence encoding hEPO was transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure, and purified. This hEPO transcript was then ligated in a single step to a 2'-hydroxymethylated RNA (OMeRNA) "bridging" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (bridging oligo 1 (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', bold bases OMeRNA) using either (A) T4 RNA ligase 1+PEG 8K, (B) T4 RNA ligase 1 or (C) T4 RNA Ligase 2 and a DNA oligonucleotide "splint" complementary to the 3'-UTR and bridging oligo 1 (splint oligo 1 (SEQ ID NO: 9); 5' CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). Alternatively, MCNA was prepared using splint oligonucleotide 5 (SEQ ID NO: 13), a palindromic sequence containing 2 copies of oligo 2 connected

with a 5'-5' phosphodiester bond. To prepare the samples for ligation, bridging oligo 1 was 5'-end phosphorylated in a reaction containing 50 μM bridging oligo 1, 1 mM ATP, 1×PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 h. Phosphorylated bridging oligo 1 was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.2 μM capped hEPO transcript, 1.5 μM bridging oligo 1 and 3 μM splint oligo 1 (or 1.5 μM splint oligo 5) by heating to 75° C. for 5 min followed by gradual cooling to room temperature over 5 min. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and (A) 1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25° C.), 1 mM ATP and 1 U/μL T4 RNA ligase 1 (NEB), (B) 1× RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/μL T4 RNA ligase 1 (NEB) or (C) 1× T4RNA Ligase 2 Buffer (NEB; 50 mM Tris-HCl, 2 mM MgCl₂, 1 mM DTT, 400 μM ATP at pH 7.5 at 25° C.) and 1 U/μL T4 RNA ligase 2 (NEB). Each was reacted for 90 minutes at 37° C. The completed ligation reaction was then purified using an RNeasy Mini Kit (Qiagen). A portion of the purified MCNA 1 product was subsequently treated with DNase I to remove residual bridge oligonucleotide to prevent potential endogenous RNase H cleavage of PCNA 1 in cells.

Alternatively, MCNA 1 (SEQ ID NO: 17) was prepared by splint ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR) and a 3' UTR flanking an RNA sequence encoding hEPO was transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hEPO transcript was then ligated in a single step to a 2'-hydroxymethylated RNA (OMeRNA) "bridging" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (bridging oligo 1 (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', bold bases OMeRNA) using either (A) T4 RNA ligase 1+PEG 8K, (B) T4 RNA ligase 1 or (C) T4 RNA Ligase 2 and a DNA oligonucleotide "splint" complementary to the 3'-UTR and bridging oligo 1 (splint oligo 1 (SEQ ID NO: 9); 5' CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). Alternatively, MCNA was prepared using splint oligonucleotide 6 (SEQ ID NO: 14), and a palindromic sequence containing 2 copies of oligo 2 connected with a 5'-5' phosphodiester bond. To prepare the samples for ligation, bridging oligo 1 was 5'-end phosphorylated in a reaction containing 50 μM bridging oligo 1, 1 mM ATP, 1×PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated bridging oligo 1 was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.2 μM capped hEPO transcript, 1.5 μM bridging oligo 1 and 3 μM splint oligo 1 (or 1.5 μM splint oligo 6) by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and (A) 1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25°

117

C.), 1 mM ATP and 1 U/ μ L T4 RNA ligase 1 (NEB), (B) 1 \times RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/ μ L T4 RNA ligase 1 (NEB) or (C) 1 \times T4RNA Ligase 2 Buffer (NEB; 50 mM Tris-HCl, 2 mM MgCl₂, 1 mM DTT, 400 μ M ATP pH 7.5 at 25° C.) and 1 U/ μ L T4 RNA ligase 2 (NEB). Each was reacted for 90 minutes 37° C. The completed ligation reaction was then purified using an RNeasy Mini Kit (Qiagen). A portion of the purified MCNA 1 product was subsequently treated with DNase I to remove residual bridge oligonucleotide to prevent potential endogenous RNase H cleavage of PCNA 1 in cells. MCNA 1 (No Poly(A) Tail Sequence):

(SEQ ID NO: 17)
 5' -GGACAGAUCCGCGGAGACGCCAUCCACGCUGUUUUGACCUCCAUAG
 AAGACACCGGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAA
 CGCGGAUUCGCGUGCCAAGAGUGACUCACCGUCCUUGACACGAUGGGGG
 UGCACGAAUGUCCUGCCUGGCUUGGCUUCUCCUGUCCUGCUGUCGUCUC
 CCUCUGGGCCUCCAGUCCUGGGCGCCCCACCAGCCUCAUCUGUGACAG
 CCGAGUCCUGGAGAGGUACCUCUUGGAGGCCAAGGAGGCCGAGAAUAUCA
 CGACGGGCGUGUGCAACACUGCAGCUUGAAUGAGAAUAUCACUGUCCCA
 GACACCAAAGUUAUUUUAUGCCUGGAAGAGGAUGGAGGUCGGGCAGCA
 GGCCGUAGAAGUCUGGCAGGGCCUGGCCUGCUGUCGGAAGCUGUCCUGC
 GGGGCCAGGCCUGUUGGUCAACUCUCCAGCCUGGGAGCCCCUGCAG
 CUGCAUGUGGAUAAAGCCGUCAGUGGCCUUCGCGAGCCUACCCACUCUGCU
 UCGGGCUCUGGGAGCCCAGAAGGAAGCAUCUCCUCCAGAUCCGGCCU
 CAGCUGCUCACUCCGAACAAUCACUGCUGACACUUUCCGAAACUCUUC
 CGAGUCUACUCCAAUUUCCUCCGGGGAAAGCUGAAGCUGUACACAGGGGA
 GGCCUGCAGGACAGGGGACAGAUGACGGGUGGCAUCCUGUGACCCUCC
 CCAGUGCCUCCUGGCCUGGAAGUUGCCACUCCAGUGCCACCAGCCU
 UGUCCUAAUAAAUAAGUUGCAUCAAGCU CGACUCUCGG-3'-PO₄-
3'-GGCUCUCAGCUCGAACUACGUUGAAUUAUUUUAUCCUGUCCGAC
 CACCCGUGACCUCACCGUUGAAGGUCCCGGUCUCUCCGUGACCCUCC
 CAGUGUCCUACGGUGGGCAGUAGACAGGGGACAGGACGUCCGGAGGGGA
 CACAUGUCGAAGUCGAAAGGGCCUCCUUUAACCUCAUCUGAGCCUUCUC
 AAACGCCUUUCACAGUCGUCACUAAAGCCUACCCUCGUCGACUCCGGC
 GUAGACUCCUCCUACCGAAGGAAGACCCGAGGGUCUCGGGCUUCGUCU
 CACCACUCCGACGCUCCGGUGACUGCCGAAUAGGUGUACGUCGACGUC
 CCCGAGGGUGCCGACCCUUCUAAACUGGUUGUCCCGACCGGGGCGUCCU

118

-continued

GUCGAAGGCGUCGUCUCCCGGUCGCGGACGGUCUGAAGAUGCCGGACGACC
 GGCUGGAGGUAGGAGAAGGUCCGUUAUCUUUAAUUGAAACCACAGACCUG
 5 UCACUAUAAGAGUAAGUUCGACGUCACAAGUCGUGUCGGGCAGCACUAUA
 AGAGCCGGAGGAACCGGAGGUUCUCAUGGAGAGGUCCUGAGCCGACAGU
 GUCUACUCCGACACCCCGCGGGUCCUGACCCUCCGGGUCUCCUCCGCU
 10 GUCGUCUCCUGUCCUUCGUGUCGGUCCGUCGUCGUAAGCACGUGGGGGU
 AGCACAGUCCUGCCACUCAGUGAGAACCUGUCCUAGGCGCAAGGUU
 ACGUGGCAAGGGCCGGCGCCUCCGACCUAGCCAGGGCCACAGAAGAUACC
 15 UCCAGUUUUGUCGCACCUACCGCAGAGGUCCGCUAGACAGG-5'

EPO MCNA #2

MCNA 2 (SEQ ID NO: 18) was prepared by splint ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR) and a 3' UTR flanking an RNA sequence encoding hEPO was transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hEPO transcript was then ligated in a single step to an RNA "bridging" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (bridging oligo 2 (SEQ ID NO: 6); 5'-AAAAAAAAAAAA-3'-3'-A AAAAAAAAAAAA-5', underlined bases RNA) using T4 RNA ligase 1+PEG 8K and a DNA oligonucleotide "splint" complementary to the 3'-UTR and bridging oligo 2 (splint oligo 3 (SEQ ID NO: 11); 5' TTT TTT TAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). Alternatively, MCNA was prepared using splint oligo 7 (SEQ ID NO: 15), a palindromic sequence containing 2 copies of splint oligo 7 connected with a 5'-5' phosphodiester bond. To prepare the samples for ligation, bridging oligo 2 was 5'-end phosphorylated in a reaction containing 50 μ M oligo 3, 1 mM ATP, 1 \times PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT, pH 7.6 at 25° C.) and 0.5 U/ μ L T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated bridging oligo 2 was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.2 μ M capped hEPO transcript, 1.5 μ M bridging oligo 2 and 3 μ M splint oligo 3 (or 1.5 μ M splint oligo 7) by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and 1 \times RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT, pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/ μ L T4 RNA ligase 1 (NEB), and was reacted for 90 min at 37° C. The completed ligation reaction was then purified using an RNeasy Mini Kit (Qiagen). EPO PCNA #2 (10A-10A Bridge):

(SEQ ID NO: 18)

5' -
 GGACAGAUCCGCGGAGACGCCAUCCACGCUGUUUUGACCUCCAUAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUCGCG
 UGCCAAGAGUGACUCACCGUCCUUGACACGAUGGGGGUGCACGAAUGUCCUGCCU

-continued

GGCUGUGGCUUCUCCUGUCUCCUGUCGUCUCCUCCUGGGCCUCCAGUCCUGGG
CGCCCCACCACGCCUCAUCUGUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAG
GCCAAGGAGGCCGAGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAU
GAGAAUAUCACUGUCCAGACACCAAAGUAAUUUCUAUGCCUGGAAGAGGAUG
GAGGUCGGGCAGCAGGCCGUGAGAAGUCUGGCAGGGCCUGGCCCUGCUGUCGGAAG
CUGUCCUGCGGGGCCAGGCCUGUUGGUCAACUCUCCAGCCUGGGAGCCCCU
GCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUCGCAGCCUCACCACUCUGCUU
CGGGCUCUGGGAGCCCAGAAGGAAGCCAUCUCCUCCAGAUCCGGCCUCAGCUG
CUCCACUCCGAACAACUACUGCUGACACUUCCGCAAACUCUCCGAGUCUACUC
CAAUUUCCUCGGGGAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGG
GGACAGAUGACGGGUGGCAUCCUGUGACCCCUCUCCAGUGCCUCUCCUGGCCCU
GGAAGUUGCCACUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGUUGCAUC
AAGCUAAAAAAAAA - 3' - PO₄ - 3' -

AAAAAAAAAUCGAACUACGUUAAUAAAAUAAUCCUGUCCGACCACCCGUGAC
CUCACCGUUGAAGGUCCCGUCCUCUCCUGACCCCUCUCCAGUGUCCUACGGU
GGCAGUAGACAGGGGACAGGACGUCCGGAGGGGACACAUGUCGAAGUCGAAAG
GGGCCUCCUUAAACCUCAUCUGAGCCUUCUCAAACGCCUUUCACAGUCGUCACUA
ACAAGCCUACCCUCGUCGACUCCGGCGUAGACCUCUCCUCCUACCGAAGGAAGACC
CGAGGGUCUCGGGCUUCGUCUACCCACUCCGACGCUUCGGUGACUGCCGAAUA
GGUGUACGUCGACGUCCCGAGGGUGCCGACCCUUCUCAAACUGGUUGUCCCGGAC
CGGGGCGUCCUGUCGAAGGCGUUCGUCUCCGGUCCGGGACGGUCUGAAGAUGCCGG
ACGACGGGCUGGAGGUAGGAGAAGGUCGUAUCUUUAAUUGAAACCACAGACCC
UGUCACUAUAAGAGUAAGUUCGACGUCACAAGUCGUGUCGGGCAGCACUAUAAG
AGCCGGAGGAACCGGAGGUUCUCCAUUGGAGAGGUCCUGAGCCGACAGUGUCUACU
CCGCACCACCCCGCGGGUCCUGACCCUCCGGGUCUCCUCCUGCUGUCGUCUCCUGUCC
UCUUCGGUGUCGGUCCGUCUGUAAGCACGUGGGGUAGCACAGUUCUGCCACU
CAGUGAGAACCUGUCCUAGGGCGAAGGUUACGUGGCAAGGGCCGGCGCCUCC
GACCUAGCCAGGGCCACAGAAGAUACCUCAGUUUUGUCGCACCUACCGCAGAGG
UCCGCUAGACAGG- 5'

EPO MCNA #3

MCNA 3 (SEQ ID NO: 19) was prepared by splint ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5' untranslated region (UTR), a 3' UTR with both UTRs flanking an RNA sequence encoding hEPO was transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. The construct was treated further to incorporate a poly(A) tail length of ~200 As using poly(A) polymerase. This hEPO transcript was then ligated in a single step to OMeRNA "bridge" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (bridging oligo 1 (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', bold bases OMeRNA) using T4 RNA ligase 1+PEG 8K and

50 a DNA oligonucleotide "splint" complementary to the 3'-UTR and bridging oligo 1 (splint oligo 4 (SEQ ID NO: 12); 5' CCG AGA GTC GTT TTT TTT TTT TTT TTT 3'; all bases DNA). Alternatively, MCNA could be prepared using splint oligo 8 (SEQ ID NO: 16), a palindromic sequence containing 2 copies of splint oligo 4 connected with a 5'-5' phosphodiester bond. To prepare the samples for ligation, bridging oligo 1 was 5'-end phosphorylated in a reaction containing 50 μM oligo 1, 1 mM ATP, 1×PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT, pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated bridging oligo 1 was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.2 μM capped hEPO transcript, 1.5 μM bridging oligo 1 and 3 μM splint oligo 4 by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An

121

RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and 1xRNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT, pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/μL T4 RNA

122

ligase 1 (NEB), and was reacted for 90 minutes at 37° C. The completed ligation reaction was then purified using an RNeasy Mini Kit (Qiagen).
EPO PCNA #3 (Includes 200A Poly(A) Tail):

(SEQ ID NO: 19)

5' -
 GGACAGAUCCGCGGAGACGCCAUCACCGCUGUUUUGACCUCCAUAAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUCCTCCG
 UGCCAAGAGUGACUCACCGUCUUGACACGAUGGGGGUGCACGAAUGUCUGCCU
 GGCUGUGGCUUCUCCUGUCCUGCUGUCGCUCUCCUGGGCCUCCAGUCCUGGG
 CGCCCCACCGCCUCAUCUGUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAG
 GCCAAGGAGGCCGAGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAU
 GAGAAUAUCACUGUCCAGACACCAAAGUAAUUUCUAUGCCUGGAAGAGGAUG
 GAGGUCGGGCAGCAGGCCGUAAGAAGUCUGGCAGGGCCUGGCCUUGCUGUCGGAAG
 CUGUCCUGCGGGCCAGGCCUGUUGGUCAACUCUCCAGCCUGGGAGCCCU
 GCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUCGCAGCCUACACUCUGCUU
 CGGGCUCUGGGAGCCAGAAGGAAGCCAUCUCCUCCAGAUCCGGCCUCAGCUG
 CUCCACUCGAAACAAUCACUGCUGACACUUUCGCAAACUCUCCGAGUCUACUC
 CAUUUCCUCCGGGGAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGG
 GGACAGAUAGCGGGUGGCAUCCUGUGACCCUCCUCCAGUGCCUUCUCCUGGCCU
 GGAAGUUGCCACUCCAGUGCCACCAGCCUUGUCUAAUAAAUAAGUUGCAUC
 AAGCUAA
 AA
 AA
 AA
 AA - CGACUCUCGG-
3' - PO₄ - 3' - GGCUCUCAGC -
 AA
 AA
 AA
 AAAUCGAAUCGUUGAAU
 UAAAAUAAUCUGUCCGACCACCCGUGACCUACCGUUGAAGGUCCCGUCCUC
 UCCGUGACCCUCCUCCAGUGUCCUACGGUGGGCAGUAGACAGGGGACAGGACGU
 CCGGAGGGGACACAUGUCGAAGUCGAAAGGGCCUCCUUAAACCUCAUCUGAGCC
 UUCUCAAAACGCCUUUCACAGUCGUCACUAAACAGCCUACCUUGUCGACUCGGC
 GUAGACUCCCCUCUACCGAAGGAAGACCCGAGGGUCUCGGGCUUCGUCUCACCA
 CUCCGACGCUUCGGUGACUGCCGAAAUAGGUGUACGUCGACGUCUCCGAGGGUG
 CCGACCUUCUCAACUGGUUGUCCCGACCGGGGCGUCUGUCGAAGGCUGUCGU
 CCCGGUCCGGGACGGUCUGAAGAUGCCGGACGACGGGCUGGAGGUAGGAGAAGG
 UCCGUUUCUUAAUUGAAACACAGACCCUGUCACUAUAAGAGUAAGUUCGACG
 UCACAAGUCGUGUCGGGCAGCACUAUAAGAGCCGGAGGAACCGGAGGUUCUCAU
 GGAGAGGUCCUGAGCCGACAGUGUCUACUCCGCACCACCCCGGGUCCUGACCC
 UCCGGGUCUCCUCCUGCUGUCGUCCUGUCCUUCGGUGUCGGUCCGUCUGUAA

-continued

AA
 AA
 UAAAAUAAUCCUGUUCGACCACCCGUGACCUCACCGUUGAAGGUCCCGUCCUC
 UCCGUGACCCUCCCGAGUGUCCUACGUGGGCAGUAGACAGGGGACAGGACGU
 CCGGAGGGGACACAUGUCGAAGUCGAAAGGGGCCUCUUUAAACCUCAUCUGAGCC
 UUCUCAAAACGCCUUUCACAGUCGUCACUAACAAGCCUACCCUGUCGACUCCGGC
 GUAGACCUCUCCUACCGAAGGAAGACCCGAGGGUUCGCGGCUUCGUCUCACCA
 CUCCGACGCUCCGGUGACUGCCGAAUAGGUGUACGUCGACGUCUCCGAGGGUG
 CCGACCUCUUCUAAUCUGGUUGUCCCGGACCGGGGCGUCUGUCGAAGGCUGUCGU
 CCCGGUCCGGGACGGUCUGAAGAUGCCGGACGACGGGCUGGAGGUAGGAGAAGG
 UCCGUAUCUUUAAUUGAAACCACAGACCCUGUCACUAUAAGAGUAAGUUCGACG
 UCACAAGUCGUGUCGGGCAGCACUAUAAGAGCCGGAGGAACCGGAGGUUCUCCA
 GGAGAGGUCCUGAGCCGACAGUGUCUACUCCGCACCACCCCGGGGUCCUGACCC
 UCCGGGUCUCCUUCGUGUCGUCUCCUGUCCUCUUCGGUGUCGGUCCGUCUGUAA
 GCACGUGGGGUAGCACAGUUCUCCGACUCAGUGAGAACCUGUCCCUUAGGCCG
 CAAGGUUACGUGGCAAGGGCCGGCGCCUCCGACCUGCCAGGGCCACAGAAGUA
 CCUCCAGUUUUGUCGCACCUACCCGACAGAGGUCCGCUAGACAGG-5'

EPO MCNA #5

MCNA 5 (SEQ ID NO: 21) was prepared by splint-independent ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single dinucleotide containing two A's linked by a 3'-3' phosphodiester bond. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR), a 3' UTR with both UTRs flanking an RNA sequence encoding hEPO was transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. The construct was treated further to incorporate a poly(A) tail length of ~200 As using poly(A) polymerase. This hEPO transcript was then ligated via two steps to an RNA "bridging" dinucleotide containing an A with a 3'-3' phosphodiester linkage to another A (bridging oligo 4 (SEQ ID NO: 8); 5'-A-3'-3'-A-5', underlined bases RNA) using T4 RNA ligase 1+PEG 8K. To prepare the samples for ligation, bridging oligo 4 was 5'-end phosphorylated in a reaction containing 50 μM bridging oligo 4, 1 mM ATP, 1×PNK

30 Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT, pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated bridging oligo 4 was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and denatured in a reaction containing 2.4 μM capped and tailed hEPO transcript and 50 μM bridging oligo 4 by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and 1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT, pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/μL T4 RNA ligase 1 (NEB), and was reacted for 90 minutes at 37° C. The partial ligation reaction was then purified using an RNeasy Mini Kit (Qiagen). The ligation reaction was repeated using a 1:1 molar ratio of the partial ligation product and additional capped and tailed hEPO transcript, and purified as previously.

EPO PCNA #5 (Includes 200A Poly(A) Tail with 1A-1A Bridge):

(SEQ ID NO: 21)

5' -
 GGACAGAUCGCCUGGAGACGCCAUCCACGCUGUUUUGACCUCUAGAAAGACACC
 GGGACCGAUCAGCCUCCGCGGGCCGGGAACGGUGCAUUGGAACGCGGAUUCCTCG
 UGCCAAGAGUGACUCACCGUCCUUGACACGAUGGGGUGCACGAAUGUCCUGCCU
 GGCUGGGCUUCUCCUGUCCUUGUCGUCUCCUUGGGCCUCCAGUCCUGGG
 CGCCCCACCACGCCUCAUCUGUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAG
 GCCAAGGAGGCCGAGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAU
 GAGAAUAUCACUGUCCAGACACCAAAGUUAAUUUCUAUGCCUGGAAGAGGAUG
 GAGGUCGGGCAGCAGGCCGUAAGAAGUCUGGCAGGGCCUGGCCUUGCUGUCGGAAG
 CUGUCCUGCGGGCCAGGCCUGUUGGUCAACUCUCCAGCCUGGGAGCCCU

-continued

GCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUCGAGCCUCACCACUCUGCUU
 CGGGCUCUGGGAGCCAGAAAGGAAGCCAUCUCCCCUCCAGAUGCGGCCUCAGCUG
 CUCCACUCCGAACAAUCACUGCUGACACUUCCGCAAACUCUUCGAGUCUACUC
 CAAUUUCCUCCGGGGAAAGCUGAAGCUGUACACAGGGGAGGCCUGCAGGACAGG
 GGACAGAUGACGGGUGGCAUCCUUGUGACCCUCCCCAGUGCCUCUCUGGCCCU
 GGAAGUUGCCACUCCAGUGCCACCAGCCUUGUCCUAAUAAAAUUAAGUUGCAUC
 AAGCUAA
 AA
 AA
 AA
 AA
 AA
 AA
 AA
 AA
 AA
 UAAAAUAAUCUGUUCGACCACCCGUGACCUCACCGUUGAAGGUCCCGUCCUC
 UCCGUGACCCUCCCCAGUGUCCUACGGUGGGCAGUAGACAGGGGACAGGACGU
 CCGGAGGGGACACAUGUCGAAAGGCGCCUCUUUAAACCUCAUCUGAGCC
 UUCUCAAAACGCCUUUCACAGUCGUCACUAACAAGCCUACCCUGUCGACUCCGGC
 GUAGACCUCCCUACCGAAGGAAGACCCGAGGGUUCGGGCUUCGUCUCACCA
 CUCCGACGCUUCGGUGACUGCCGAAAUAGGUGUACGUCGACGUCCCGAGGGUG
 CCGACCCUUCUAAUCUGGUUGUCCGGACCGGGGCGUCCUGUCGAAGGCUGUCGU
 CCCGGUCCGGGACGGUCUGAAGAUCCGGACGACGGGCGUGGAGGUAGGAGAAGG
 UCCGUAUCUUAAUUGAAACCACAGACCCUGUCACUAUAAGAGUAAGUUCGACG
 UCACAAGUCGUGUCGGGACGACUAUAAGAGCCGGAGGAACCGGAGGUUCUCCA
 GGAGAGGUCCUGAGCCGACAGUGUCUACUCCGCACCACCCCGGGGUCCUGACCC
 UCCGGGUCUCCUCGUGUCGUCUCCUGUCCUCUUCGGUGUCGGUCCGUCUGUAA
 GCACGUGGGGUAGCACAGUUCUGCCACUCAGUGAGAACCUGGCCCUUAGGCG
 CAAGGUUACGUGGCAAGGGCCGGCCUCCGACCUAGCCAGGGCCACAGAAGUA
 CCUCCAGUUUUGUCGCACCUACCCGAGAGGUCCGCUAGACAGG-5'

EPO PCNA #6

PCNA 6 (SEQ ID NO: 22) is prepared by splint ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR), a 3' UTR containing an internal section of 65 consecutive As with both UTRs flanking an RNA sequence encoding hEPO is transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hEPO transcript is then ligated in a single step to a OMeRNA "bridging" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (bridging oligo 1 (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', underlined bases OMeRNA) using T4

RNA ligase 1+PEG 8K and a DNA oligonucleotide "splint" complementary to the 3' UTR and bridging oligo 1 (splint oligo 1 (SEQ ID NO: 9); 5' CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). To prepare the samples for ligation, bridging oligo 1 is 5'-end phosphorylated in a reaction containing 50 μM bridging oligo 1, 1 mM ATP, 1×PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated bridging oligo 1 is then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.2 μM capped hEPO transcript, 1.5 μM bridging oligo 1 and 3 μM splint oligo 1 by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction is subsequently prepared to contain a 50% diluted hybridization reaction and

129

1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT, pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/μL T4 RNA ligase 1 (NEB), and is reacted for 90

130

minutes at 37° C. The completed ligation reaction is then purified using an RNeasy Mini Kit (Qiagen). EPO PCNA #6 (Includes Internal 65A Poly(A) Region):

(SEQ ID NO: 22)

5' -
 GGACAGAU CGCCUGGAGACGCCAUCACGCUGUUUGACCUC CAUAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUC CCG
 UGCCAAGAGUGACUCACCGUCCUUGACACGAUGGGGGUGCACGAAUGUC CUGCCU
 GGCUGUGGCUUCUCUGUCCUGCUGUCGCUC CUCUGGGCCUCCAGUCCUGGG
 CGCCCCACCACGCCUCAUCUGUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAG
 GCCAAGGAGGCCGAGAAUAUCACGACGGGCUGUGCUAACACUGCAGCUUGAAU
 GAGAAUAUCACUGUCCAGACACCAAAGUUAUUUCUAUGCCUGGAAGAGGAUG
 GAGGUCGGGCAGCAGGCCGUAAGAAGUCUGGCAGGGCCUGGCCUCUGUCGGAAG
 CUGUCCUGCGGGCCAGGCCUGUUGGUAACUCUCCAGCCGUGGGAGCCCU
 GCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUCGCAGCCUCACCACUCUGCUU
 CGGGCUCUGGGAGCCAGAAGGAAGCCAUCUC CCCCAGAUCCGCGCCUCAGCUG
 CUCCACUCGAAACAAUCACUGCUGACACUUUCGCAAACUCUCCGAGUCUACUC
 CAAUUUCUCCGGGGAAAGCUGAAGCUGUACACAGGGGAGGCCUCAGGACAGG
 GGACAGAUGACGGGUGGCAUCCUGUGACCCCUCC CAGUGCCUCUCUGGCCCU
 GGAAGUUGCCACUCCAGUGCCACCAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAGCCUUGUCCUAAUAAAAU
 UAAGUUGCAUCAAGCUCGACUCUCGG - 3' - PO₄ - 3' -
 GGCUCUCAGCUCGAACUACGUUGAAUUAUUUUUAUUCUGUUCGAAAAAAAAA
 AAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAAA
 ACCACCGUGACCUCACCGUUGAAGGUCCCGGUCUCUCGUGACCCUCC CAG
 UGUCCUACGGUGGGCAGUAGACAGGGACAGGACGUCCGGAGGGGACACAUGU
 CGAAGUCGAAAGGGCCUCUUUAACCUCAUCUGAGCCUUCUCAAACGCCUUUCA
 CAGUCGUCACUAACAAGCCUCACUCGUCGACUCCGGCGUAGACCUC CUCUAC
 CGAAGGAAGACCCGAGGGUCUCGGGCUUCGUCUCACCACUCCGACGCUCCGGUG
 ACUGCCGAAUAGGUGUACGUCGACGUCCCGAGGGUGCCGACCCUUCUCAACUG
 GUUGUCCCGACCGGGCGUCUGUCGAAGGCUGUCGUCCCGUCCGGGACGGUC
 UGAAGAUGCCGGACGACGGGCUUGGAGGUAGGAGAAGGUCCGUAUCUUUAAUUGA
 AACCACAGACCCUGUCACUAUAAGAGUAAGUUCGACGUCACAAGUCGUGUCGGGC
 AGCACUAUAAGAGCCGAGGAACCGGAGGUUCUCAUGGAGAGGUCCUGAGCCG
 ACAGUGUCUACUCCGCACCACCCCGGGGUCUGACCCUCGGGUCUCCUCGCU
 GUCGUCUCCUGUCUUCUGGUGUCGGUCCGUCUGUAAGCACGUGGGGUAGCAC
 AGUUCUGCCACUCAGUGAGAACCUGCCCCUAGGCGCAAGGUUACGUGGCAAG
 GGCCGGCGCCUCCGACCUAGCCAGGGCCACAGAAGAUACCUCAGUUUUGUCGCA
 CCUACCGCAGAGGUCCGCUAGACAGG - 5'

EPO PCNA #7

PCNA7 (SEQ ID NO: 23) is prepared by splint ligation of the 3' end of two copies of an RNA encoding the human Erythropoietin (hEPO) protein to the 5' ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5' untranslated region (UTR), a 3' UTR containing 3 stretches of 15 As and 1 stretch of 16 As with both UTRs flanking an RNA sequence encoding hEPO is transcribed using T7 RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hEPO transcript is then ligated in a single step to a OMeRNA "bridge" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (bridging oligo 1 (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', underlined bases OMeRNA) using T4 RNA ligase 1+PEG 8K and a DNA oligonucleotide "splint" complementary to the 3'-UTR and bridging oligo 1 (splint oligo 1 (SEQ ID NO: 9); 5' CCG AGAGTC GAG CTT GAT GCAACT TAA TTT

TAT TAG G 3'; all bases DNA). To prepare the samples for ligation, oligo 1 is 5'-end phosphorylated in a reaction containing 50 μM bridging oligo 1, 1 mM ATP, 1×PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT, pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated bridging oligo 1 is then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.2 μM capped hEPO transcript, 1.5 μM bridging oligo 1 and 3 μM splint oligo 1 by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction is subsequently prepared to contain a 50% diluted hybridization reaction and 1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT, pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 1 U/μL T4 RNA ligase 1 (NEB), and is reacted for 90 min at 37° C. The completed ligation reaction is then purified using an RNeasy Mini Kit (Qiagen). EPO PCNA #7 (Includes Multiple Short Internal Poly(A) Regions):

(SEQ ID NO: 23)

5' -
GGACAGAUCGCCUGGAGACGCCAUCCACGCUGUUUUGACCUCCAUAGAAGACACC
GGGACCGAUC CAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUC CCG
UGCCAAGAGUGACUCACCGUCCUUGACACGAUGGGGUGCACGAAUGUCCUGCCU
GGCUGUGGCUCUCUGUCCUGCUGUCGCUCCUCUGGGCCUCCAGUCCUGGG
CGCCCCACACGCCUCAUCUGUGACAGCCGAGUCCUGGAGAGGUACCUCUUGGAG
GCCAAGGAGGCCGAGAAUAUCACGACGGGCUGUGCUGAACACUGCAGCUUGAAU
GAGAAUAUCACUGUCCAGACACCAAAGUAAUUAUUGCCUGGAAGAGGAUG
GAGGUCGGGCAGCAGGCCGUAGAAGUCUGGCAGGGCCUGGCCUCUGUCGGAAG
CUGUCCUGCGGGCCAGGCCUGUUGGUCAACUCUCCAGCCGUGGGAGCCCU
GCAGCUGCAUGUGGAUAAAGCCGUCAGUGGCCUUCGCAGCCUACACUCUGCUU
CGGGCUCUGGGAGCCAGAAGGAAGCCAUCUCCCUCCAGAUGCGGCCUCAGCUG
CUCCACUCCGAACAAUCACUGCUGACACUUUCGCAAACUCUCCGAGUCUACUC
CAAUUUCCUCGGGGAAAGCUGAAGCUGUACACAGGGGAGGCCUCGAGGACAGG
GGACAGAUGACGGGUGGCAAAAAAAAAAAAAAAAAUCCUGUGACCCUCCCAAAA
AAAAAAAAAAGUGCCUCUCUGGCCUCUGGAAAAAAAAAAAAAAAAAGUUGCCAC
UCCAGUGCCACCAAAAAAAAAAAAAAAAAAGCCUUGUCCUAAUAAAAUUAAGUUGC
AUCAAGCUCGACUCUCGG-3'-PO₄-3'-
GGCUCUCAGCUCGAACUACGUUGAAUAAAAAAAAUCCUGUCCGAAAAAAAAA
AAAAACACCCGUGACCUCACCGUUGAAAAAAAAAAAAAAAAAGGUCCGGUCCUCU
CCGUGAAAAAAAAAAAAAAAAACCCUCCAGUGUCCUAAAAAAAAAAAAAAAAAC
GGUGGGCAGUAGACAGGGACAGGACGUCCGGAGGGGACACAUGUCGAAGUCGA
AAGGGCCUCCUUUAACCUCAUCUGAGCCUUCUCAACGCCUUUCACAGUCGUCA
CUAACAAGCCUCACUCGUCGACUCGGCGUAGACCUCUCCUUAACCGAAGGAAG
ACCCGAGGGUCUCGGGCUUCGUCUCACACUCCGACGCUCCGGUGACUGCCGAA
AUAGGUGUACGUCGACGUCCCGAGGGUGCCGACCCUUCUCAACUGGUUGUCCCG
GACCGGGCGUCCUGUCGAAGGCUGUCGUCCCGGUCCGGACGGUCUGAAGAUGC

-continued

CGGACGACGGGCGUGGAGGUAGGAGAAGGUCCGUAUCUUUAAUUGAAACCACAGA
 CCCUGUCACUAUAAGAGUAAGUUCGACGUCACAAGUCGUGUCGGGCAGCACUAUA
 AGAGCCGGAGGAACCGGAGGUUCUCAUGGAGAGGUCCUGAGCCGACAGUGUCU
 ACUCCGCACCACCCCGCGGGUCCUGACCCUCCGGGUCUCCUCGUGUCGUCUCCUG
 UCCUCUUCGGUGUCGGUCCGUCCUGUAAGCACGUGGGGUAGCACAGUUCUGCC
 ACUCAGUGAGAACCGUGCCCUUAGGCGCAAGGUUACGUGGCAAGGGCCGGCGCC
 UCCGACCUAGCCAGGGCCACAGAAGAUACCUCAGUUUUGUCGCACCUACCAGCAG
 AGGUCCGCUAGACAGG-5'

FIG. 5 shows the results of MCNA detected via gel electrophoresis. MCNA run in lanes 1-15 were the result of a ligation reaction comprising an EPO mRNA to bridging oligonucleotide to DNA splint (SEQ ID NO: 9) molar ratio of 2:1:2. The molar amounts of EPO mRNA and RNA ligase are included in the below table:

Lane	EPO (μ M)	Ligase (μ M)
1	1.7	2.25 RNA Ligase 1
2	1.7	0.6 RNA Ligase 1
3	0.85	0.6 RNA Ligase 1
4	0.425	0.6 RNA Ligase 1
5	0.2125	0.6 RNA Ligase 1
6	1.7	2.25 RNA Ligase 1 + 10% PEG
7	1.7	0.6 RNA Ligase 1 + 10% PEG
8	0.85	0.6 RNA Ligase 1 + 10% PEG
9	0.425	0.6 RNA Ligase 1 + 10% PEG
10	0.2125	0.6 RNA Ligase 1 + 10% PEG
11	1.7	0.3 RNA Ligase 2
12	1.7	0.6 RNA Ligase 2
13	0.85	0.6 RNA Ligase 2
14	0.425	0.6 RNA Ligase 2
15	0.2125	0.6 RNA Ligase 2

FIG. 5 demonstrates that RNA Ligase 1 was superior to RNA Ligase 2 in producing MCNA comprising EPO RNA under the conditions tested. Further, the addition of 10% PEG to the reaction conditions enhanced ligation.

FIG. 6 shows MCNA detected via gel electrophoresis. Lane 1 shows Capped EPO mRNA (no poly(A) tail). Lane 2 shows a MCNA mixture of full length MCNA ligation product mixed with unreacted/partially reacted EPO RNA product (with no DNase treatment). Lane 3 shows a MCNA mixture of full length MCNA ligation product mixed with unreacted/partially reacted EPO RNA product (with DNase treatment).

FIG. 8 shows MCNA detected via gel electrophoresis. Lane 1 shows a RNA sizing ladder. Lane 2 shows a MCNA mixture of full length MCNA ligation product mixed with unreacted/partially reacted EPO RNA product. Lane 3 shows purified unreacted/partially reacted EPO RNA product. Lane 4 shows purified EPO MCNA ligation product.

MCNA-OTC Preparation
 MCNA-OTC comprising human Ornithine Transcarbamylase (hOTC) RNA (SEQ ID NO: 24) was prepared by

15 splint ligation of the 3'-end of two copies of an RNA encoding the hOTC protein to the 5'-ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR) and a 3' UTR flanking an RNA sequence encoding hOTC was transcribed using RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hOTC transcript was then ligated in a single step to a 2'-hydroxymethylated RNA (OMeRNA) "bridge" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (oligo 1 (bridge) (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', bold bases OMeRNA) using either T4 RNA ligase 1+PEG 8K and a DNA oligonucleotide "splint" complementary to the 3'-UTR and oligo 1 (oligo 2 (splint) (SEQ ID NO: 9); 5' CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). To prepare the samples for ligation, oligo 1 was 5'-end phosphorylated in a reaction containing 50 μ M oligo 1, 1 mM ATP, 1 \times PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT pH 7.6 at 25° C.) and 0.5 U/ μ L T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated oligo 1 (bridge) was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 3.3 μ M capped hOTC transcript, 1.5 μ M oligo 1 and 3.3 μ M oligo 2 by heating to 75° C. for 5 minutes, followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and 1 \times RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 0.33 U/ μ L T4 RNA ligase 1. Each was reacted for 60 minutes at 37° C. The completed ligation reaction was then reacted with DNase I and subsequently purified using an RNeasy Maxi Kit (Qiagen). The reaction products were evaluated for ligation efficiency using TBE/agarose gel electrophoresis. The isolated MCNA-OTC product was equilibrated with Lipofectamine and transfected into adherent HEK293 cells. Unfractionated cell lysate was then assayed for citrulline production from ornithine and carbamoyl phosphate (FIG. 10).
 MCNA-OTC

(SEQ ID NO: 24)

5' -
 GGACAGAUCCGCGGAGACGCCAUCCACGCUGUUUUGACCUCCAUAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUCCTCCG
 UGCCAAGAGUGACUCACCGUCCUUGACACGAUGCUGUUCACCUUCGGAUCUUGC

- continued

UGAACAACGCUGCGUUCGGAUUGGUCACAACUUCUUGGUCCGGAACUUCAGAUG
 CGGCCAGCCGCUCCAGAACAAGGUGCAGUCUAGGGGAGGGACCUCUCACCCUG
 AAAAAUUCACCGGAGAAGAGAUCAAGUACAUGCUGUGGCUGUCAGCCGACCUCA
 AAUUCGGAUCAAGCAGAAGGGCGAAUACCUUCCUUUGCUGCAGGGAAAGUCCCU
 GGGGAUGAUCUUCGAGAAGCGCAGCACUCGCACUAGACUGUCAACUGAAACCGGC
 UUCGCGCUGCUGGGAGGACACCCUGCUUCCUGACCACCAAGAUUCCAUCUGG
 GUGUGAACGAAUCCUCACCGACACAGCGCGGGUGCUGUCGUCUUGGCAGACGC
 GGUCUCGCCC CGUGUACAAGCAGUCUGAUCUGGACACUCUGGCCAAGGAAGCC
 UCCAUUCUAUCAUUAUUGGAUUGUCCGACCUCUACCAUCCAUCCAGAUUCUGG
 CCGAUUAUCUGACUCUGCAAGAACAUACAGCUCUCCUGAAGGGGUUACCCUUUC
 GUGGAUCGGCGACGGCAACAACAUUCUGCACAGCAUUAUGAUGAGCGCUGCCAAG
 UUUGGAAUGCACCUCCAAGCAGCGACCCCGAAGGGAUACGAGCCAGACGCCUCCG
 UGACGAAGCUGGCUGAGCAGUACGCCAAGGAGAACGGCACUAAGCUGCUGCUCAC
 CAACGACCCUCUCGAAGCCGCCACGGUGGCAACGUGCUGAUCACCGAUACCGG
 AUCUCCAUGGGACAGGAGGAGAAAAGAAGAAGCGCCUGCAAGCAUUUCAGGGG
 UACCAGGUGACUAUGAAAACCGCCAAGGUCGCGCCUCGGACUGGACCUUCUUGC
 ACUGUCUGCCAGAAAGCCCGAAGAGGUGGACGACGAGGUGUUCUACAGCCCGCG
 GUCGUCUGGUCUUUCGAGGCCGAAAACAGGAAGUGGACUAUCAUGGCCGUGAU
 GGUGUCUCCUGCUGACCGAUUACUCCCGCAGCUGCAGAAACCAAAGUUCUGACGG
 GUGGCAUCCUGUGACCCUCCCGAGUGCCUCUCUGGCCUGGAAGUUGCCACU
 CCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGUUGCAUCAAGCU CGACUCUC

GG-3' - PO₄-3' -

GGUCUCAGCUCGAACUACGUUGAAUUAUUAAUUAUCCUGUUCGACCACCCGUGA
 CCUCACCGUUGAAGGUCCCGGUCUCUCGUGACCCUCCCGAGUGUCCUACGG
 UGGCAGUCUUGAAACCAAAGACGUCGACGCCCCUCAUAGCCAGUCGUCCUGU
 GGUAGUGCCGGUACUAUCAGGUGAAGGACAAAAGCCGGAGGCCUUUCUGGUCGC
 UGGCGCCGACAUCUUGUGGAGCAGCAGGUGGAGAAGCCCGAAAGACCCGUCUGU
 CACGUUCUUCAGGUCAGGCUCGCGCUGGAACCGCCAAAAGUAUCAGUGGACC
 AUGGGGACUUUACGAACGUCCGGAAGAAGAAAAGGAGGAGGACAGGGUACCUC
 UAGGUCCAUAGCCACUAGUCGUGCAACGGUGGCACCCGCCGAAGCUCUCCAGCA
 ACCACUCGUCGUCGAAUCACGGCAAGAGGAACCGCAUGACGAGUCGGUCGAAGCA
 GUGCCUCCGACAGCCGAGCAUAGGGAAGCCCAGCGACGAACCUCACGUAAGGU
 UUGAACCGUCGCGAGUAGUAUACGACACGUCUUAACAACGGCAGCGGCUAGG
 UGCUUUCCAUUCGGGAAGUCCUCGACAUUACAAGAACGUCUCAGUCUAUUAG
 CCGGUCUAGACCUACCCUACCAUCUCAGCCUGUAGGUAAUUAUAUCCUAC
 CUCCGAAGGAACCGGUCUCACAGGUCUAGUCUGACGAACAUGUGCGCCCGCUCU
 GGCGCAGACGGUACCUGCUGUCGUGGGCGGACACAGCCACUCCUAAGCAAGUG
 UGGGUCUACCUAUAAGAACCACCAGUCCUUCGUCCCCACAGGAGGGUCGUCGCGC
 UUCGGCCAAAGUCAACUGUCAGAUACGCUCACGACGCGAAGAGCUUCUAGUAGG

-continued

GGUCCCUGAAAGGGACGUCGUUUCCUCCAUAAGCGGGAAGACGAACUAGGCCUU
 AACUC CAGCCGACUGUCGGUGUCGUACAUGAACUAGAGAAGAGGCCACUUCAAA
 AAGUCCCACUCCUCCAGGGAGGGGAACUCGACGUGGAACAAGACCUCGCCGACCG
 GCGUAGACUUCAAGGCCUGGUACUUAACACUGGUAAGGCCUUGCGUCGCAACAA
 GUCGUUCUAGGCCUCCAACUUGUCGUAGCACAGUUCUGCCACUCAGUGAGAACC
 GUGCCCCUAGGCCGCAAGGUUACGUGGCAAGGGCCGGCGCCUCCGACCUAGCCAG
 GGCCACAGAAGAUACCUCAGUUUUGUCGCACCUACCGCAGAGGUCCGCUAGACA
 GG-5' (Bold base are OMeRNA)

MCNA-PAH Preparation

MCNA-PAH comprising human Phenylalanine Hydroxylase (hPAH) RNA (SEQ ID NO: 25) was prepared by splint ligation of the 3'-end of two copies of an RNA encoding the hPAH protein to the 5'-ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR) and a 3' UTR flanking an RNA sequence encoding hPAH was transcribed using RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hPAH transcript was then ligated in a single step to a 2'-hydroxymethylated RNA (OMeRNA) "bridge" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (oligo 1 (bridge) (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', bold bases OMeRNA) using either T4 RNA ligase 1+PEG 8K and a DNA oligonucleotide "splint" complementary to the 3'-UTR and oligo 1 (oligo 2 (splint) (SEQ ID NO: 9); 5' CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). To prepare the samples for ligation, oligo 1 was 5'-end phosphorylated in a reaction containing 50 μM oligo 1, 1 mM ATP, 1×PNK

15 Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated oligo 1 (bridge) was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 2.7 μM capped hPAH transcript, 1.2 μM oligo 1 and 2.7 μM oligo 2 by heating to 75° C. for 5 minutes, followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and 1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 0.33 U/μL T4 RNA ligase 1. Each was reacted for 60 minutes at 37° C. The completed ligation reaction was then reacted with DNase I and subsequently purified using an RNeasy Maxi Kit (Qiagen). The reaction products were evaluated for ligation efficiency using TBE/agarose gel electrophoresis. The isolated MCNA-PAH reaction product was equilibrated with Lipofectamine and transfected into adherent HEK293 cells. Unfractionated cell lysate was then assayed for PAH protein expression using a PAH-specific ELISA (FIG. 11).
 MCNA-PAH

(SEQ ID NO: 25)

5' -
 GGACAGAUCGCCUGGAGACGCCAUCCACGCUGUUUUGACCUCUCCAUGAAGACACC
 GGGACCGAUC CAGCCUCCGCGGGCGGGAACGGUGCAUUGGAACGCGGAUUC CCGG
 UGCCAAGAGUGACUCACCGUCCUUGACACGAUGAGCACC GCGGUCUGGAGAACC
 CCGGCCUGGGCCGCAAGCUGAGCGACUUCGGCCAGGAGACCAGCUACAUCGAGGA
 CAACUGCAACCAGAACGGCGCCAUCAGCCUGAUCUUCAGCCUGAAGGAGGAGGUG
 GGCGCCUGGCCAAGGUGCUGCGCCUGUUCGAGGAGAACGACGUGAACCUGACCC
 ACAUCGAGAGCCGCCCCAGCCGCCUGAAGAAGGACGAGUACGAGUUCUUCACCCA
 CCUGGACAAGCGCAGCCUGCCCGCCUGACCAACAUCAUCAAGAUCCUGCGCCAC
 GACAUCGGCGCCACCGUGCACGAGCUGAGCCGCGACAAGAAGAAGGACACCGUGC
 CCUGGUUCCCCCGCACCAUCCAGGAGCUGGACCGCUUCGCCAACCAGAUCCUGAG
 CUACGGCGCCGAGCUGGACGCGCACCCCGGCUUCAAGGACCCCGUGUACCGG
 GCCCGCCGCAAGCAGUUCGCGACAUCGCCUACAACUACCGCCACGGCCAGCCCA
 UCCCCGCGUGGAGUACAUGGAGGAGGAGAAGAAGACCUGGGGCACCGUGUUCA
 AGACCCUGAAGAGCCUGUACAAGACCCACGCCUGCUACGAGUACAACCACAUCUU
 CCCCUGCUGGAGAAGUACUGCGGCUUCACGAGGACAACAUC CCCCAGCUGGAG
 GACGUGAGCCAGUUCUGCAGACCUGCACCGGCUUCGCGCCUGCGCCCCGUGGCCG

- continued

GCCUCGUCGAGCAGCCGCGACUUCUGGGCGGCCUGGCCUUCGCGUGUUCACUCG
 CACCAGUACAUCGCCACGGCAGCAAGCCCAUGUACACCCCGAGCCCGACAUC
 UGCCACGAGCUGCUGGGCCACGUGCCCUGUUCAGCGACCCGAGCUUCGCCAGU
 UCAGCCAGGAGAUCGGCCUGGCCAGCCUGGGCGCCCCGACGAGUACAUCGAGAA
 GCUGGCCACCAUCUACUGGUUCACCGUGGAGUUCGGCCUGUGCAAGCAGGGCGAC
 AGCAUCAAGGCCUACGGCGCCGGCCUGCUGAGCAGCUUCGGCGAGCUGCAGUACU
 GCCUGAGCGAGAAGCCCAAGCUGCUGCCCUGGAGCUGGAGAAGACCGCCAUCCA
 GAACUACACCGUGACCGAGUUCAGCCCCUGUACUACGUGGCCGAGAGCUUCAAC
 GACGCCAAGGAGAAGGUGCGCAACUUCGCCGCCACCAUCCCCGCCCCUUCAGCG
 UGCGCUACGACCCCUACACCCAGCGCAUCGAGGUGCUGGACAACACCCAGCAGCU
 GAAGAUCCUGGCCGACAGCAUCAACAGCGAGAUCCGCAUCCUGUGCAGCGCCUG
 CAGAAGAUCAAGUAACGGGUGGCAUCCUGUGACCCUCCCCAGUGCCUCUCCUG
 GCCCUGGAAGUUGCCACUCCAGUGCCCACCAGCCUUGUCCUAAUAAAAUUAAGUU
 GCAUCAAGCU CGACUCUCGG - 3' - PO₄ - 3' -

GGUCUCAGCUCGAAUCUACGUUGAAUAAAAUAAUCUGUUCGACCACCCGUGA
 CCUCACCGUUGAAGGUCCCGGUCUCUCCGUGACCCUCCCCAGUGUCCUACGG
 UGGCAAUGAACUAGAAGACGUCCCGCGACGUGUCCUACGGCUAGAGCGACAACU
 ACGACAGCCGGUCCUAGAAGUCGACGACCCACAACAGGUCGUGGAGCUACGCGAC
 CCACAUCCCCAGCAUCGCGUGCGACUUCGCCGCCCUACCACCGCCGCUUCAACG
 CGUGGAAGAGGAACCGCAGCAACUUCGAGAGCCGGUGCAUCAUGUCCCCGACCUCU
 GAGCCAGUGCCACAUCAAGACCUACCGCCAGAAGAGGUCGAGGUCCCGUCGUCG
 AACCCGAAGAGCGAGUCCGUCUAGACGUCGAGCGGCUUCGACGAGUCGUCGGCC
 GCGGCAUCCGGAACUACGACAGCGGGACGAACGUGUCCGGCUUGAGGUGCCACUU
 GGUCAUCUACCACCGGUCGAAGAGCUACAUGAGCAGCCCCGCGGGUCCGACCGG
 UCCGGCUAGAGGACCGACUUGACCCGCUUCGACGCCAGCGACUUGUCCCCGUGCA
 CCGGGUCGUCGAGCACCUCUACAGCCCGAGCCCCACAUGUACCCGAACGACGG
 CACCGCCUACAUGACCCACGUCACCUUGUGCGCCUUCGGUCCGGCGGGUCCUUC
 AGCGCCGACGAGUCGUCGGCCGGUGCCCCGCGUCCGCCUUCGGCCACGUC CAGA
 CGUCCUUGACCAGUGCAGGAGGUCGACCCCUACAACAGGAGCACCUCGGCGU
 CAUGAAGAGGUCGUCUUUUUACACCAACAUGAGCAUCGUCCGCACCAGAAC
 AUGUCCGAGAAGUCCAGAACUUGUGCCACGGGGUCCAGAAGAAGAGGAGGAGG
 UACAUGAGGUGCGCCCCUACCCGACCGGCACCGCCAUCAACAUCGCUACAGCC
 GCUUGACGAACCGCCCGCGCAUGUGCCCCAGGAACUUCGGCCCCACGAGCCG
 CAGGUCGAGCCGCGGCAUCGAGUCCUAGACCAACCGCUUCGCCAGGUCGAGGACC
 UACCACGCCCCUUGGUCCCGUGCCACAGGAAGAAGAACAGCGCCGAGUCGAGCA
 CGUGCCACCGCGGCUACAGCACCGGUCUAGAACUACUACAACCAGUCCCGCCC
 GUCCGACCGGAACAGGUCCACCCACUUCUUGAGCAUGAGCAGGAAGAAGUCCGCC
 GACCCCGCCGAGAGCUACACCCAGUCCAAGUGCAGCAAGAGGAGCUUGUCCGCGU
 CGUGGAACCGGUCCCGCGGGUGGAGGAGGAAGUCCGACUUCUAGUCCGACUACCG

-continued

CGGCAAGACCAACGUCAACAGGAGCUACAUCGACCAGAGGACCGGCUUCAGCGAG
 UCGAACCGCCGGGUCCGGCCCCAAGAGGUCGUGCCGCCACGAGUAGCACAGUUCU
 GCCACUCAGUGAGAACCUGGCCUUAGGCGCAAGGUUACGUGGCAAGGGCCGGC
 GCCUCCGACCUAGCCAGGGCCACAGAAGAUACCUCCAGUUUUGUCGCACCUACCG
 CAGAGGUCCGCUAGACAGG-5' (Bold base are OMeRNA)

MCNA-CFTR Preparation

MCNA-CFTR comprising human Cystic Fibrosis Transmembrane Conductance Regulator (hCFTR) RNA (SEQ ID NO: 26) was prepared by splint ligation of the 3'-end of two copies of an RNA encoding the hCFTR protein to the 5'-ends of a single oligonucleotide containing two 5' ends and a linked 3'-3' phosphodiester bond within the sequence. Briefly, a 5'-capped RNA containing a 5'-untranslated region (UTR) and a 3' UTR flanking an RNA sequence encoding hCFTR was transcribed using RNA polymerase, enzymatically capped to contain a 5'-Cap 1 structure and purified. This hCFTR transcript was then ligated in a single step to a 2'-hydroxymethylated RNA (OMeRNA) "bridge" oligonucleotide containing a 20 nucleotide (nt) palindromic sequence with a 3'-3' phosphodiester linkage between the 10th and 11th nt (oligo 1 (bridge) (SEQ ID NO: 5); 5'-CGACUCUCGG-3'-3'-GGCUCUCAGC-5', bold bases OMeRNA) using either T4 RNA ligase 1+PEG 8K and a DNA oligonucleotide "splint" complementary to the 3'-UTR and oligo 1 (oligo 2 (splint) (SEQ ID NO: 9); 5' CCG AGA GTC GAG CTT GAT GCA ACT TAA TTT TAT TAG G 3'; all bases DNA). To prepare the samples for ligation, oligo 1 was 5'-end phosphorylated in a reaction containing 50 μM

oligo 1, 1 mM ATP, 1×PNK Buffer (NEB; 70 mM Tris-HCl, 10 mM MgCl₂, 5 mM DTT pH 7.6 at 25° C.) and 0.5 U/μL T4 Polynucleotide Kinase (NEB) at 37° C. for 1 hour. Phosphorylated oligo 1 (bridge) was then desalted using a Sephadex G-25 desalting column (Princeton Separations) and hybridized to the transcript and splint in a reaction containing 0.92 μM capped hCFTR transcript, 0.42 μM oligo 1 and 0.92 μM oligo 2 by heating to 75° C. for 5 minutes followed by gradual cooling to room temperature over 5 minutes. An RNA ligation reaction was subsequently prepared to contain a 50% diluted hybridization reaction and 1×RNA ligase Buffer (NEB; 50 mM Tris-HCl, 10 mM MgCl₂, 1 mM DTT pH 7.5 at 25° C.), 1 mM ATP, 10% PEG and 0.33 U/μL T4 RNA ligase 1. Each was reacted for 60 minutes at 37° C. The completed ligation reaction was then reacted with DNase I and subsequently purified using an RNeasy Maxi Kit (Qiagen). The reaction products were evaluated for ligation efficiency using TBE/agarose gel electrophoresis. The isolated MCNA-CFTR product was equilibrated with Lipofectamine and transfected into adherent HEK293 cells. Unfractionated cell lysate was then assayed for CFTR protein expression using CFTR-specific Western Blotting (FIG. 12).
 MCNA-CFTR

(SEQ ID NO: 26)

5' -
 GGACAGAU CGCCUGGAGACGCCAUCCACGCUGUUUUGACCUC CAUAGAAGACACC
 GGGACCGAUCCAGCCUCCGCGGCCGGGAACGGUGCAUUGGAACGCGGAUUC CCG
 UGCCAAGAGUGACUCACCGUCCUUGACACGAUGCAACGCUCUCUCUUGAAAAGG
 CCUCGGUGGUGUCCAAGCUCUUCUUCGUGGACUAGACCCAUC CUGAGAAAGGG
 GUACAGACAGCGCUUGGAGCUGUCCGAUAUCUAUCAAUCCUUC CUGGACUCC
 GCGGACAACCUGUCCGAGAAGCUCGAGAGAGAAUGGGACAGAGAA CUCGCCUCAA
 AGAAGAACC CGAAGCUGAUUAAUGCGCUUAGGCGGUGCUUUUUCUGGCGGUUCA
 UGUUCUACGGCAUCUUCUACCUUGGGAGAGGUCACCAAGGCCGUGCAGCCCCU
 GUUGCUGGGACGGAUUAUUGCCUCCUACGACCCCGACAACAAGGAAGAAAGAAGC
 AUCGCUAUCUACUUGGGCAUCGGUCUGUGCCUGCUUUUCAUCGUC CGGACCCUCU
 UGUUGCAUCCUGCUAUUUUCGGCCUGCAUCAUUGGCAUGCAGAU GAGAAUUG
 CCAUGUUUUC CUGAUCUACAAGAAAACUCUGAAGCUCUCGAGCCGCGUGCUUGA
 CAAGAUUUC AUCCGCGCCAGCUCGUGUCCUGCUCUCCAACAAUCUGAACAAGUUC
 GACGAGGGCCUCGCCUGGCCACUUCGUGUGGAUCGCCCCUCUGCAAGUGGCGC
 UUCUGAUGGGCCUGAUCUGGGAGCUGCUGCAAGCCUCGGCAUUCUGUGGGCUUG
 GAUUCUGAUCGUGCUGGCACUGUUC CAGGCCGACUGGGGCGGAUGAUGAUGA
 AGUACAGGGACCAGAGAGCCGGAAAGAUUCCGAACGGCUGGUGAUCACUUCGG
 AAAUGAUCGAAAACAUC CAGUCAGUGAAGGCCUACUGCUGGGAAGAGGCCAUGG

- continued

AAAAGAUGAUUGAAAACCUCCGGCAAACCGAGCUGAAGCUGACCCGCAAGGCCG
UUACGUGCGCUAUUUCAACUCGUCGCUUUUCUUCUUCUCCGGGUUCUUCGUGGUG
UUUCUCUCCGUGCUCUCCUACGCCUGAUUAAGGGAAUCAUCCUCAGGAAGAUCU
UCACCACCAUUUCCUUCUGUAUCGUGCUCGCAUGGCCGUGACCCGGCAGUUCUCC
AUGGGCCGUGCAGACUUGGUACGACUCCUGGGAGCCAUAACAAGAUCAGGAC
UUCUUCAAAGCAGGAGUACAAGACCUCGAGUACAACCUGACUACUACCGAGG
UCGUGAUGGAAAACGUCACCGCUUUUUGGGAGGAGGGAUUUGGCGAACUGUUCG
AGAAGGCCAAGCAGAACAACAACAACCGCAAGACCUCGAACGGUGACGACUCCCU
CUUCUUUCAAACUUCAGCCUGCUCGGGACGCCCGUGCUGAAGGACAUUAACUUC
AAGAUCGAAAGAGGACAGCUCUUGGCGGUGGCCGGAUCGACCGGAGCCGAAAG
ACUUCUCCUGCUGAUGGUGAUCUAGGGAGAGCUUGAACCUAGCGAGGGAAAGAUC
AAGCACUCCGGCCGAUCAGCUUCUGUAGCCAGUUUUCUGGAUCAUGCCCGGAA
CCAUUAAGGAAAACAUCUUCUUCGGCGUGUCCUACGAUGAAUACCGCUACCGGUC
CGUGAUCAAAGCCUGCCAGCUGGAAGAGGAUUAUUCAAAGUUCGCGGAGAAAGA
UAACAUCGUGCUGGGCGAAGGGGUUAUACCUUGUCGGGGGCCAGCGGGCUAG
AAUCUCGCGUGGCCAGAGCCUGUAUAAGGACGCCGACCUGUAUCUCCUGGACUCC
CCUUCGGAUACCUGGACGUCUUGACCGAAAAGGAGAUUCUGAAUCGUGCGUGU
GCAAGCUGAUGGCUAACAAGACUCGCAUCCUCGUGACCUCAAAAUGGAGCACCU
GAAGAAGGCAGACAAGAUUCUGAUUCUGCAUGAGGGGUCCUCCUACUUUUACGG
CACCUUCUCGGAGUUGCAGAACUUGCAGCCGACUUCUCAUCGAAGCUGAUGGGU
UGCGACAGCUUCGACCAGUUCUCCGCCGAAAGAAGGAACUCGAUCUGACCGGAAA
CCUUGCACCGCUUCUUCUUUGGAAGGGCAGCCCCUGUGUCAUGGACCGAGACUAA
GAAGCAGAGCUUCAAGCAGACCGGGGAAUUCGGCGAAAAGAGGAAGAACAGCAU
CUUGAACCCCAUUAACUCCAUCGCAAGUUCUCAAUCGUGCAAAGACGCCACUG
CAGAUGAACGGCAUUGAGGAGACUCCGACGAACCCUUGAGAGGCGCCUGUCCC
UGGUGCCGGACAGCGAGCAGGGAGAAGCCAUCUGCCUCGGAUUUCUGAUCUC
CACUGGUCGACGCUCAAGCCGGCGGCGCAGUCCGUGCUGAACCUGAUGACC
CACAGCGUGAACAGGGCCAAAACAUCACCGCAAGACUACCGCAUCCACCGGA
AAGUGUCCUGGCACCUCAAGCGAAUCUUAACCGAGCUCGACAUACUCCCGGAG
ACUGUCGAGGAAACCGGGUCGAAAUUCGGAAGAAUCAACGAGGAGGAUCU
GAAAGAGUGCUUCUUCGACGAUAUGGAGUCGAUACCCGCCGUGACGACUUGGAA
CACUUAUCUGCGGUACAUCACUGUGCACAAGUCAUUGAUCUUCGUGCUGAUUUG
GUGCCUGGUGAUUUUCCUGGCCGAGGUCGCGGCCUCACUGGUGGUGCUCUGGCUG
UUGGAAACACGCCUCUGCAAGACAAGGGAAACUCCACGCACUCGAGAAACAACA
GCUAUGCCGUGAUUAUCACUUCACCUCUUAUUAACGUGUUCUACUACUACGU
CGGAGUGGCGGAUACCGUCGCGAUGGGUUUCUUCAGAGGACUGCCGCGUGGUC
CACACC'UUGAUCACCGUCAGCAAGAUUCUUCACCACAAGAUUGUUGCAUAGCGUGC
UGCAGGCCCCAUGUCCACCUCAACACUCUGAAGGCCGGAGGCAUUCUGAACAG
AUUCUCCAAGGACAUCGCUAUCUUGGACGAUCUCCUGCCGCUUACCAUCUUUGAC
UUCAUCCAGCUGCUGCUGAUCGUGAUUGGAGCAAUCGCAGUGGUGGCGGUGCUG
CAGCCUUAUUAUUUCGUGGCCACUGUGCCGGUCAUUGGGCGUUAUCAUGCUGC

- continued

GGGCCUACUUCUCCAAACCAGCCAGCAGCUGAAGCAACUGGAAUCCGAGGGACG
 AUCCCCAUCUUCACUCACCUUGUGACGUCGUUGAAGGGACUGUGGACCCUCCGG
 GCUUUCGGACGGCAGCCUACUUCGAAACCCUUCUCCACAAGGCCUGAACCUCC
 ACACCGCCAAUUGGUUCCUGUACCUGUCCACCCUGCGGUGGUUCCAGAUGCAGAU
 CGAGAUGAUUUUCGUCAUCUUCUUCUACGCGGUCACAUUCAUCAGCAUCCUGACU
 ACCGGAGAGGGAGAGGGACGGGUCGGAUAUUCUGACCCUCCGCAUGAACAUU
 AUGAGCACCCUGCAGUGGGCAGUGAACAGCUCGAUCGACGUGGACAGCCUGAUGC
 GAAGCGUCAGCCGCGUGUUCAGUUCUUCGACAUCCUACUGAGGGAAAACCCAC
 UAAGUCCACUAAGCCUACAAAAUGGCCAGCUGAGCAAGGUCAUGAUCAUCGAA
 AACUCCACGUGAAGAAGGACGAUAUUUGGCCUCCGGAGGUCAAUGACCGUGA
 AGGACCUGACCGCAAAGUACACCGAGGGAGGAAACGCCAUUCUCGAAAACAUCAG
 CUUCUCAUUUCGCCGGGACAGCGGGUCGGCCUUCUCGGGCGGACCGGUUCGGG
 AAGUCAACUCUGCUGCGGCUUUCUCCGGCUGCUGAAUACCGAGGGGGAAAUC
 AAAUUGACGGCGUGUCUUGGGAUUCAUUACUCUGCAGCAGUGGCGGAAGGCCU
 UCGGGCUGAUCCCCAGAAAGGUGUUAUCUUCUGGGUACCUUCGGAAGAACC
 GGAUCCUUCAGAGCAGUGGAGCGACCAAGAAAUCUGGAAGGUCGCCGACGAGGU
 CGGCCUGCGCUCUGAUUGAACAAUUUCUGGAAAGCUGGACUUCGUGCUCGUC
 GACGGGGGAUGUGUCCUGUCGCACGGACAUAAGCAGCUCAUGUGCCUCGCACGGU
 CCGUGCUCUCAAGGCCAAGAUUCUGCUGCUGGACGAAACUUCGGCCACCUGGA
 UCCGGUACCUACCAGAUCAUCAGGAGGACCCUGAAGCAGGCCUUUGCCGAUUGC
 ACCGUGAUUCUCUGCGAGCACCGCAUCGAGGCCAUGCUGGAGUGCCAGCAGUUC
 UGGUCAUCGAGGAGAACAAGGUCGCCAAUACGACUCCAUAUCAAAGCUCUCAA
 CGAGCGGUCGUCUGUUCAGACAAGCUAUUUACCGUCCGAUAGAGUGAAGCUUUC
 CCGCAUCGGAACAGCUCAAAGUGCAAUUCGAAGCCGAGAUCCGAGCCUUGAAGG
 AAGAGACUGAGGAAGAGGUGCAGGACACCCGGCUUUAACGGGUGGCAUCCUGU
 GACCCUCCCCAGUGCCUUCUUGGCCUGGAAGUUGCCACUCCAGUGCCCACCA
 GCCUUGUCUAAUAAAAUUAAGUUGCAUCAAGCU CGACUCUCGG-3'-PO₄-3'-
GGCUCUCAGCUCGAACUACGUUGAAUAAAAUAAUCCUGUUCGACCACCCGUGA
 CCUCACCGUUGAAGGUCCCGGUCUCUCCGUGACCCUCCCCAGUGUCCUACGG
 UGGGCAAUUUCGGCCACAGGACGUGGAGAAGGAGUCAGAGAAGGAAGUCCGA
 CGCUAGACGCCGAAGCUAAACGUGAAACUCGACAAGGCUACGCCUUCUCGAAGU
 GAGAUAGCCUGCCACUUUAUCGAACAGACUUGUCGUGGCGAGCAACUCCUGAA
 AACUUACCUCAGCAUAACCGCCUGGAACAAGAGGAGCUACUGGUCCUUGACGACC
 GUGAGGUCGUACCGGAGCUACGCCACGAGCGUCUCUAGUGCCACGUUAGCCGUU
 UCCGGACGAAGUCCAGGAGGACUACUAGACCAUCCACUGGCCUAGGUCCACCCG
 GCUUCCAAGCAGGUCGUCGUCUUGAACCAGAACCUUCUGGUCUGGCACGCUCC
 GUGUACUCGACGAAUACAGGCACGUCUGUAGGGGGCAGCUGCUCGUGCU
 UCAGGUCCGAAAGGUCCUUUAACAAGUAGUGCCUCGCGUCGGCUGGAGCAGCCG
 CUGGAAGGUCUAAAGAACCAGCGAGGUGACGAGCAUUCUAGGUCCAAGAAGGC

-continued

CUUCCAUGGGCUUUCUACUUGUGGAAGACCCCUAGUGCGGCUUCCGGAAGGCG
GUGACGACGUCUCAUUAACUAGGGUUCUGUGCGGCAGUUAACCUAAAGGGGG
AGCCAUAAGUCGUCGCGCCUCCUUUCGGCUGUCGUCUCAAACUGAAGGGCCUUGGCC
AGGCGGGCUCUUCGGCUGGGCGACAGGGCCGCUUUAACUUCUUCGACUACAAAAG
CUCUUACCGCAAAGGAGGGAGCCACAUGAAACGCCAGUCCAGGAAGUGCCAGUAA
ACUGGAGGCCUCCCGUUUAUAGCAGGAAGAAGUGCACCCUCAAAGCUACUAGU
ACUGGAACGAGUCGACCGGUAAAAACAUCGAAUCACUGAAUCACCCAAAAGG
GAGUCAUCCGUACAGCUACUUGAACUUGUGCGCCGACUGCGAAGCGUAGUCCGAC
AGGUGCAGCUAGCUCGACAAGUGACGGGUGACGUCCACGAGUAUUACAAGUACC
GCUCCAGUCUAAUAAGGCUGGGCAGGGAGAGGGAGAGGCCAUCAGUCCUACGA
CUACUUACACUGGCGUACUUCUUCUACUGCUUUUAGUAGAGCUACGCGUAGACC
UUGGUGGGUCCACCUGUCCAUGUCCUUGGUUAACCGCCACACCUCCAAGUCC
GGAAACCCUUCUCCCAAAGCUUCAUCCGACGGCAGGCUUUCGGGCCUCCAGGU
GUCAGGGAAGUUGCUGCAGUGUCCACUCACUUCUACCCCUAGCAGGGAGCCUA
AGGUCAACGAAGUCGACGACCGACCAAACUCCUUCUACUCCGGGCGUCGUACUACU
UGCGGUGUACUGGCGGUGUACCGGUGCUUUUACAUCGACGUCGUGGGCGGUG
GUGACGCUAACGAGGUAGUGCUAGUCGUCGUCGACCUACUUCAGUUUCUACCAU
UCGCCGUCCUAGCAGGUCCUACGCUACAGGAACCUUAGACAAGUCUACG
GAGGCCGGAAGUCUCACAACUCCACCUGUACCCCGGACGUCGUGCGAUACGUU
GUAGAACACCACUUCUAGAACGACUGCCACUAGUUCACACCUGGUCGCCGUA
GGAGACUUCUUGGGUAGCGUCGUCUCCAUAGGCGGUGAGGCUGCAUCUACUUCU
UGUGCAUUAUUCUCCUCCACCUCACUUAUAGUGCCGUUUCGACAACAAGAGCU
CACGCACCUCAAAGGGAACAGAACGUCUCCGCACAAAGGUUGUCGGUCUCGUGG
UGGUCACUCCGGCGCUGGAGCCGGUCCUUUAGUGGUCGUGGUUAGUCGUGCU
UCUAGUUACUGAACACGUGUCACUACAUGGCGUCUUAUCAAGGUUCAGCAGU
GCCGCCCAUAGCUGAGGUUAGCAGCUUCUUCGUGAGAAAGUCUAGGAGGAGCA
ACUAAAGAAGCCUUUAAAGCUCGGGCCAAAGGACGUCGUCAGAGGCCUCAUCUA
CAGCUCGAGCAUUCUAAGCGAACUCCACGGUCCUGUGAAAGGCCACCUCACGC
CAUCAGAACGCCACUUAACAAAACCGGGACCAAGUGCGACACCCAGUAGUCCAAGU
CGUGCCUGACGGCGGGCGGCCGAACCUCCGAGCCUGGUCACCCUAGUGCCUUUA
GGCUCCGUCCUACCGAAGAGGGACGAGCGACAGGCCGUGGUCCUGUCCGCGGAG
AGUUCCTCAAGCAGCCUCAGGAGGAGUACGGCAAGUAGACGUCACCCGAGAAAA
CGUGCUAACUCUUGAACGCCUACUCAAUUACCCCAAGUUUAACGACAAGAAGGA
GAAAAGCGGCUUAAGGGGCAGACGAACUUCGAGACGAAGAAUCAGAGCCAGGU
ACUGUGUCCCGCAGCGGAAGGUUUCUUCGCCACGUUCCAAAGGCAGUCCUAG
CUCAAGGAAGAAAGCCGCCUCUUGACCAGCUUCGACAGCGUUGGGUAGUCGAAGC
UACUCUUCAGCCCAGCUUCAAGACGUUGAGGCUCUUCACGGCAUUUUAUCCU
CCUGGGGAGUACGUCUAGUCUUAAGAACAGACGGAAGAAGUCCACGAGGUAAAA
CCUCCAGUGCUCUACGCUCAAGAAACUUCGGUAGUCGAACGUGUGCGUGCUAAGC
UUCUAGAGGAAAAGCCAGUCCUGCAGGUCCAAGGCUUCCCCUCAGGUCCUUA
UGUCCAGCCGAGGAUAUUGGCCGAGACCGGUCGUCUUAAGAUCGGGCGACCGG

-continued

GGGGCUGUUC CAUUAUGGGGAAGCGGGUCGUGCACAAUAGAAAGAGGCGCUU
GAAACUUUAUAGGAGAAGGUCGACCGUCCGAAACUAGUGCCUGGCCAUCGCCAUA
AGUAGCAUCCUGUGCGGCUCUACUACAAAAGGAAUACCAAGGCCCGUACUAGG
UCCUUUUGACCGAUGUCUUCGACUACGCCGGCCUCACGAAACUAGAAAGGGAGCGA
UCCAAGUUCGAGAGGGUACUAGUGGUAGUCGUCCUUCAGAAAGGCCGAGGCCA
GCUAGGCCGGUGGCGGUCCUCGACAGGAGAAAGCUAGAACUUCAAUUACAGGAA
GUCGUGCCCGCAGGGCUCGUCCGACUUCAAACUUUUUUCUUCUCCUCAGCAGUGGC
AAGCUC CAGAACGCCAACAAACAAGACGAACCGGAAGAGCUUGUCAAGCGGUU
UAGGGAGGAGGGUUUCCGCCACUGCAAAGGUAGUGCUGGAGCCAUCAUCAGU
CCAACAUGAGCUC CAGAACAUGAGGACGAAAACUCCUUCAGGACCUAGAACAA
UUACCGAGGGUCCUCAGCAUGGUUCAGACGUGCCGGGUACCCUUGACGGCCAG
UGCCGGUACGCCUCGUGCUAUGUCUUUCUUUACCACCACUUCUAGAAGGACUCCU
ACUAAGGGAAUUAGUCCCGCAUCCUUCGUGCCUCUUCUUGUGGUGCUUCUUGGG
CCUCUUUCUUUCGCCUGCUCAACUUUAUCGCGUGCAUUCGCCGGAACGCCCCAG
UCGAAUGCGAGCCAAACGGCCUCCAAAGUUAGUAGAAAAGGUACCGGAGAAGG
GUCGUCAUCCGGAAGUGACUGACCUACAAAAGCUAGUAAAGGCUUCACUAGUGG
UCGGCAAGCCUUUAGAAAGGCCGAGAGACCAGGGACAUGAAGUAGUAGUAGGCG
GGUUCAGGCCGGACCUUGUCACGGUCGUGCUAGUCCUAGGUUCGGGUGUCUUC
GGCUC CCGAACGUCGUCGAGGGUCUAGUCCGGUAGUCUUCGCGGUGAACGUCUCC
CCGCUAGGUGUCUUCACCCGGUCCCGCUCGCGGAGCAGCUUGAACAGUCUAAC
AACCUCUCGUCCUGUGUCGACCGGCUACCUUAGAACAGUUCGUGCGCCGAGC
UCUCGAAGUCUCAAAAGAACAUCUAGUCCUUUUGUACCGUUAAGAGUAGACGU
ACGGUUACACUACGUCCGGCUUUUAUCGUCCUACGUUGUUCUCCAGGCCUGCUA
CUUUUCGUCCGUGUCUGGCUACGGGUUCAUCUAUCGCUACGAAGAAAGAAGGAA
CAACAGCCCCAGCAUCCUCGUAUUAGGCAGGGUCGUUGUCCCGACGUGCCGG
AACCACUGGAGAGGGUCCAUCUCUUCUACGGCAUCUUGUACUUGGCGGUCUUUU
UCGUGGCGGAUUCGCGUAAUAGUCGAAGCCCAAGAAGAAACUCCGCUCAAGAG
ACAGGGUAAGAGAGAGCUCGAAGAGCCUGUCCAACAGGCGCCUCAGGUGCCUUC
CUAAACUAUCUAUAGCCUGUCGAGGUUCGCGACAGACAUGGGGAAAGAGUCCUA
CCCAGAU CAGGUGUCUUCUUCUCGAACCUGUGGUGGCUC CGGAAAAGUUCUCCU
CUCGCAACGUAGCACAGUUCUGCCACUCAGUGAGAACCGUGCCCCUAGGCGCA
AGGUUACGUGGCAAGGGCCGGCGCCUCCGACCUAGCCAGGGCCACAGAAGAUACC
UCCAGUUUUGUCGCACCUACCGCAGAGGUCCGCUAGACAGG-5' (Bold base are
OMeRNA)

Example 2. Exemplary Protein Production with MCNA

This example demonstrates the production of protein encoded by mRNA linked by their 3' ends to a bridging oligonucleotide.

MCNA comprising human erythropoietin (hEPO) mRNA were synthesized as described above and used to transfect HEK293T cells (1 microgram RNA transfection per sample). FIG. 7 shows the results of an experiment com-

paring the amount of secreted hEPO protein from HEK293T cells when the cells were transfected with either a) mRNA encoding hEPO that lacked a polyA tail, b) MCNA comprising hEPO mRNA, or c) MCNA comprising hEPO mRNA that had been treated with DNase. A clear increase in protein production was achieved when the cells were transfected with either the MCNA comprising hEPO mRNA or the DNase-treated MCNA comprising hEPO mRNA compared to the untailed hEPO mRNA.

FIG. 9 shows the results of an experiment comparing the amount of secreted hEPO protein from HEK293T cells when the cells were transfected with either a) mRNA encoding hEPO that lacked a polyA tail, b) unpurified mixture of MCNA comprising hEPO mRNA with unreacted/partially reacted EPO RNA, c) purified unreacted/partially reacted EPO RNA, or d) purified EPO MCNA. All samples were transfected with a total of 250 nanograms RNA. A clear increase in protein production was achieved when the cells were transfected with purified EPO MCNA compared to the mixture or unreacted hEPO RNA. FIG. 10 shows the results of an experiment comparing the amount of human OTC protein activity (as measured by citrulline production) within HEK293T cells when the cells were transfected with either a) mRNA encoding hOTC that lacked a polyA tail (hOTC monomer), or b) MCNA comprising hOTC mRNA. Detectable protein production was achieved only when the cells were transfected with the MCNA comprising hOTC as compared to the hOTC monomer.

FIG. 11 shows the results of an experiment comparing the amount of human PAH protein produced within HEK293T cells when the cells were transfected with either a) mRNA encoding hPAH that lacked a polyA tail (hPAH monomer), or b) MCNA comprising hPAH mRNA. Significantly higher protein production was achieved when the cells were transfected with the MCNA comprising hPAH as compared to the hPAH monomer.

FIG. 12 shows the results of an experiment comparing the amount of human CFTR protein produced within HEK293T cells when the cells were transfected with either a) mRNA encoding hCFTR that lacked a polyA tail (hCFTR monomer), or b) MCNA comprising hCFTR mRNA. Detectable protein production was achieved only when the cells were transfected with the MCNA comprising hCFTR as compared to the hCFTR monomer.

Example 3. Exemplary In Vivo Protein Production with MCNA

This example demonstrates the in vivo production of protein encoded by mRNA linked by their 3' ends to a bridging oligonucleotide.

MCNA comprising human ornithine carbamoyltransferase (hOTC) mRNA were synthesized as described above. *spf^{ash}* mice were treated intravenously with hOTC MCNA encapsulated in lipid nanoparticles. Animals were sacrificed and their livers were isolated either 24 hours or 7 days post-administration. Citrulline production was measured in the liver samples and it was found that the level of hOTC protein activity 7 days post-administration was comparable to the level of hOTC protein activity 24 hours post-administration (FIG. 13). At both time points, hOTC protein activity was significantly greater than in the livers of control *spf^{ash}* mice. Further, substantial hOTC protein was detected via Western blot at both 1 day and 8 days post-administration, but for only the *spf^{ash}* mice treated with hOTC MCNA

LNPs, not the mice treated with the hOTC monomer LNPs (FIG. 14), consistent with the observed activity data. In comparison, when *spf^{ash}* mice were treated intravenously with hOTC mRNA, levels of hOTC protein activity were higher 24 hours post-administration than they were 7 days post-administration (FIG. 15). As clearly shown in FIG. 16, when hOTC protein activity 7 day post-administration was calculated as a percentage of activity levels after 24 hours, more sustained in vivo activity is observed for hOTC MCNA (109% of 24 hour activity) than for hOTC mRNA (38% of 24 hour activity).

In another study, MCNA comprising human phenylalanine hydroxylase (hPAH) were synthesized as described above. PAH knock-out (KO) mice were treated intravenously with either hPAH MCNA or an hPAH monomer (hPAH mRNA with a 5' cap but without a polyA tail) encapsulated in lipid nanoparticles. Animals were sacrificed and their livers were isolated 24 hours post-administration. More than 27 times more hPAH protein was detected in the livers of mice treated with hPAH MCNA than was detected in the livers of mice treated with the hPAH monomer (FIG. 17).

Further, a demonstration of efficacy was achieved after treatment of PAH knock-out (KO) mice with hPAH MCNA LNPs. Specifically, serum phenylalanine levels were significantly reduced 24 hours after treatment with hPAH MCNA while no reduction in serum phenylalanine was seen 24 hours after treatment with hPAH monomer LNPs (FIG. 18).

In another study, MCNA comprising human erythropoietin (hEPO) were synthesized as described above. Wild-type mice were treated intravenously with either hEPO MCNA or an hEPO monomer (hEPO mRNA with a 5' cap but without a polyA tail) encapsulated in lipid nanoparticles. Serum samples from the animals were obtained 24 hours post-administration. More than 480 times more hEPO protein was detected in the serum of mice treated with hEPO MCNA than was detected in the serum of mice treated with the hEPO monomer (FIG. 19).

In another study, MCNA comprising human cystic fibrosis transmembrane conductance regulator (hCFTR) were synthesized as described above. CFTR KO mice were treated via aerosolization of hCFTR MCNA encapsulated in lipid nanoparticles. Animals were sacrificed and their lungs were isolated either 24 hours or 7 days post-administration. As shown in FIG. 20, MCNA-derived hCFTR protein was detected in both the bronchial epithelial airways (top row) as well as alveolar regions (bottom row) both 24 hours and 7 days post-administration (brown staining).

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. The scope of the present invention is not intended to be limited to the above Description, but rather is as set forth in the following claims:

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 26

<210> SEQ ID NO 1

<211> LENGTH: 827

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

-continued

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 1

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
 cgauccagcc uccgcgccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu 120
 gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
 ugucccugcu gucgcuccu cugggccucc caguccuggg cgccccacca cgccucaucu 240
 gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aauaucacga 300
 cgggcugugc ugaacacugc agcuugaug agaauaucac ugucccagac accaaaguaa 360
 auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
 uggcccugcu gucggaagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
 cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
 cucugcuucg ggcucuggga gcccagaagg aagccaucuc cccuccagau gcggccucag 600
 cugcuccacu ccgaacaau acugcugaca cuuuccgcaa acucuuccga gucuacucca 660
 auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
 gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
 ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcu 827

<210> SEQ ID NO 2

<211> LENGTH: 1027

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 2

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
 cgauccagcc uccgcgccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu 120
 gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
 ugucccugcu gucgcuccu cugggccucc caguccuggg cgccccacca cgccucaucu 240
 gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aauaucacga 300
 cgggcugugc ugaacacugc agcuugaug agaauaucac ugucccagac accaaaguaa 360
 auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
 uggcccugcu gucggaagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
 cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
 cucugcuucg ggcucuggga gcccagaagg aagccaucuc cccuccagau gcggccucag 600
 cugcuccacu ccgaacaau acugcugaca cuuuccgcaa acucuuccga gucuacucca 660
 auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
 gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
 ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcuaaa aaaaaaaaaa 840
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 900
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 960
 aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1020
 aaaaaaa 1027

-continued

<210> SEQ ID NO 3
 <211> LENGTH: 891
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 3

```

ggacagaucg ccuggagacg ccauccacgc uuuuuugacc uccauagaag acaccgggac    60
cgauccagcc uccgcggccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu    120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc    180
ugucccugcu gucgcucccu cugggccucc caguccuggg cgccccacca cgccucaucu    240
gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aaauacacga    300
cgggcugugc ugaacacugc agcuugaug agaauaucac ugucccagac accaaaguua    360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc    420
uggcccugcu gucggagcu guccugcggg gccaggcccu guuggucaac ucuucccagc    480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca    540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc ccucccagau gcggccucag    600
cugcuccacu ccgaacaau acugcugaca cuuuccgcaa acucuuccga gucuacucca    660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau    720
gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu    780
ccagugccca ccaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa    840
aaaaaaaaa aaaaaaagcc uuguccuaau aaaaauaagu ugcaucaagc u                891
  
```

<210> SEQ ID NO 4
 <211> LENGTH: 883
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 4

```

ggacagaucg ccuggagacg ccauccacgc uuuuuugacc uccauagaag acaccgggac    60
cgauccagcc uccgcggccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu    120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc    180
ugucccugcu gucgcucccu cugggccucc caguccuggg cgccccacca cgccucaucu    240
gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aaauacacga    300
cgggcugugc ugaacacugc agcuugaug agaauaucac ugucccagac accaaaguua    360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc    420
uggcccugcu gucggagcu guccugcggg gccaggcccu guuggucaac ucuucccagc    480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca    540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc ccucccagau gcggccucag    600
cugcuccacu ccgaacaau acugcugaca cuuuccgcaa acucuuccga gucuacucca    660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau    720
gacggguggc aaaaaaaaa aaaaaucccu gugacccuc ccaaaaaaaaa aaaaaaaaaaag    780
ugccucuccu ggcccuggaa aaaaaaaaaa aaaguugcca cuccagugcc caccaaaaaa    840
aaaaaaaaaag ccuuguccua auaaaauuaa guugcauca gcu                883
  
```

-continued

<210> SEQ ID NO 5
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (10)..(11)
 <223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
 10 and 11 comprises PO4

 <400> SEQUENCE: 5

 cgacucucgg ggcucucagc 20

<210> SEQ ID NO 6
 <211> LENGTH: 20
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (10)..(11)
 <223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
 10 and 11 comprises PO4

 <400> SEQUENCE: 6

 aaaaaaaaaa aaaaaaaaaa 20

<210> SEQ ID NO 7
 <211> LENGTH: 6
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (3)..(4)
 <223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
 3 and 4 comprises PO4

 <400> SEQUENCE: 7

 aaaaaa 6

<210> SEQ ID NO 8
 <211> LENGTH: 2
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: MISC_FEATURE
 <222> LOCATION: (1)..(2)
 <223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
 1 and 2 comprises PO4

 <400> SEQUENCE: 8

 aa 2

<210> SEQ ID NO 9
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide

 <400> SEQUENCE: 9

 ccgagagtcg agcttgatgc aacttaattt tattagg 37

-continued

<210> SEQ ID NO 10
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 10

ccgagagtga tgcaacttaa ttttattagg 30

<210> SEQ ID NO 11
 <211> LENGTH: 37
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 11

tttttttttt agcttgatgc aacttaattt tattagg 37

<210> SEQ ID NO 12
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide

<400> SEQUENCE: 12

ccgagagtcg tttttttttt tttttttttt 30

<210> SEQ ID NO 13
 <211> LENGTH: 74
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (37)..(38)
 <223> OTHER INFORMATION: 5'-5' bridge between nucleotides at positions
 37 and 38 comprises P04

<400> SEQUENCE: 13

ggattatttt aattcaacgt agttcgagct gagagccccg agagtcgagc ttgatgcaac 60

ttaattttat tagg 74

<210> SEQ ID NO 14
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (30)..(31)
 <223> OTHER INFORMATION: 5'-5' bridge between nucleotides at positions
 30 and 31 comprises P04

<400> SEQUENCE: 14

ggattatttt aattcaacgt agtgagagcc ccgagagtga tgcaacttaa ttttattagg 60

<210> SEQ ID NO 15
 <211> LENGTH: 74
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (37)..(38)
 <223> OTHER INFORMATION: 5'-5' bridge between nucleotides at positions
 37 and 38 comprises P04

<400> SEQUENCE: 15

ggattat tttt aattcaacgt agttcgattt tttttttttt ttttttttagc ttgatgcaac 60
 ttaattttat tagg 74

<210> SEQ ID NO 16
 <211> LENGTH: 60
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (30)..(31)
 <223> OTHER INFORMATION: 5'-5' bridge between nucleotides at positions
 30 and 31 comprises P04

<400> SEQUENCE: 16

tttttttttt tttttttttt gctgagagcc ccgagagtcg tttttttttt tttttttttt 60

<210> SEQ ID NO 17
 <211> LENGTH: 1674
 <212> TYPE: RNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Chemically synthesized nucleotide
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (837)..(838)
 <223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
 837 and 838 comprises P04

<400> SEQUENCE: 17

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccaauagaag acaccgggac 60
 cgauccagcc uccgeggccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu 120
 gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
 ugucccugcu gucgcuccu cugggccucc caguccuggg cgccccacca cgccucaucu 240
 gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aauaucacga 300
 cgggcugugc ugaacacugc agcuugaaug agaauaucac ugucccagac accaaaguua 360
 auuucuaugc cuggaagagg auggaggucg ggcagcagc cguagaaguc uggcagggcc 420
 uggcccugcu gucggaaagc guccugcggg gccaggcccu guuggucaac ucuucccagc 480
 cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
 cucugcuucg ggcucuggga gcccagaagg aagccaucuc cccuccagau gcggccucag 600
 cugcuccacu ccgaacaau acugcugaca cuuuccgaa acucuuccga gucuacucca 660
 auuuccuccg gggaaagcug aagcuguaca caggggagc cugcaggaca ggggacagau 720
 gacggguggc aucccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
 ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcucga cucucggggc 840
 ucucagcucg aacuacguug aaauaaaaa auccuguucc gaccaccgu gaccucaccg 900
 uugaaggucc cgguccucuc cgugaccccu cccaguguc ccuacggugg gcaguagaca 960
 ggggacagga cguccggagg ggacacaugu cgaagucgaa aggggccucc uuaaaccuca 1020

-continued

```

ucugagccuu cucaaagcc uuucacaguc gucacuaaca agccucaccu cgucgacucc 1080
ggcguagacc uccccucuac cgaaggaaga cccgaggguc ucgggcuucg ucucaccacu 1140
ccgacgcuuc cggugacugc cgaaauaggu guacgucgac guccccgagg gugccgaccc 1200
uucucaacug guugucccg accggggcgu ccugucgaag gcugucgucc cgguccggga 1260
cggucugaag augccggacg acgggcugga gguaggagaa gguccguauc uuuuuugaa 1320
accacagacc cugucacuau aagaguaagu ucgacgucac aagucguguc gggcagcacu 1380
auaagagccg gaggaaccgg agguucucca uggagagguc cugagccgac agugucuacu 1440
ccgcaccacc ccgcgggucc ugaccucccg ggucuccuc gcugucgucc cuguccucu 1500
cggugucggg cgguccgua agcacguggg gguagcacag uuccugccac ucagugagaa 1560
ccgugccccc uaggcgcaag guuacguggc aaggccggc gccuccgacc uagccagggc 1620
cacagaagau accuccaguu uugucgcacc uaccgcagag guccgcuaga cagg 1674

```

<210> SEQ ID NO 18

<211> LENGTH: 1674

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (837)..(838)

<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions 837 and 838 comprises P04

<400> SEQUENCE: 18

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcgggcg ggaacggugc auuggaacgc ggauuccccc ugccaagagu 120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
ugucccugcu gucgcucccu cuggggccucc caguccuggg cgcaccacca cgccucaucu 240
gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aauaucacga 300
cgggcugugc ugaacacugc agcuugaaug agaauaucac ugucccagac accaaaguua 360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
uggcccugcu gucggaaagc guccugcggg gccaggcccu guugguacac ucuucccagc 480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc cccuccagau gcggccucag 600
cugcuccacu ccgaacaauc acugcugaca cuuuccgaa acucuuccga gucuacucca 660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcuaaa aaaaaaaaaa 840
aaaaaaaaucg aacuacguug aaauaaaaua auccguucc gaccaccgug gaccucaccg 900
uugaaggucc cgguccucuc cgugaccccu cccaguguc ccuacggugc gcaguagaca 960
ggggacagga cguccggagg ggacacaugu cgaagucgaa aggggcccuc uuuuaccuca 1020
ucugagccuu cucaaagcc uuucacaguc gucacuaaca agccucaccu cgucgacucc 1080
ggcguagacc uccccucuac cgaaggaaga cccgaggguc ucgggcuucg ucucaccacu 1140
ccgacgcuuc cggugacugc cgaaauaggu guacgucgac guccccgagg gugccgaccc 1200
uucucaacug guugucccg accggggcgu ccugucgaag gcugucgucc cgguccggga 1260

```

-continued

```

cggucugaag augccggacg acgggcugga gguaggagaa gguccguauc uuuuuugaa 1320
accacagacc cugucacuau aagaguaagu ucgacgucac aagucguguc gggcagcacu 1380
auaagagccg gaggaaccgg agguucucca uggagagguc cugagccgac agugucuacu 1440
ccgcaccacc ccgcgggucc ugaccucccg ggucuccuc gcugucgucc cuguccucu 1500
cggugucggu ccguccugua agcacguggg gguagcacag uuccugccac ucagugagaa 1560
ccgugcccu uaggcgcaag guuacguggc aaggccggc gccuccgacc uagccagggc 1620
cacagaagau accuccaguu uugucgcacc uaccgcagag guccgcuaga cagg 1674

```

<210> SEQ ID NO 19

<211> LENGTH: 2074

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1037)..(1038)

<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
1037 and 1038 comprises PO4

<400> SEQUENCE: 19

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcgcccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu 120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
uguccugcu gucgcuccu cugggcccucc caguccuggg cgcaccacca cgccucaucu 240
gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aaauacacga 300
cgggucuguc ugaacacugc agcuugaau agaaauacac ugucccagac accaaaguuu 360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
uggcccugcu gucggaagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc ccuccagau gggccucag 600
cugcuccacu ccgaacaau acugcugaca cuuuccgaa acucuuccga gucuacucca 660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcuaaa aaaaaaaaaa 840
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 900
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 960
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1020
aaaaaacga cucucggggc ucucagcaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1080
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaucg aacuacguug 1260
auuuuuuuu auccuguucc gaccaccguc gaccucaccg uugaagguc cgguccucuc 1320
cgugaccccu cccaguguc ccuacggugg gcaguagaca ggggacagga cguccggagg 1380
ggacacaugu cgaagucgaa aggggcccucc uuuaaccuca ucugagccuu cucaaaccgc 1440
uuucacaguc gucacuaaca agccucaccu cgucgacucc ggcguagacc ucccucucac 1500

```

-continued

```

cgaaggaaga cccgaggguc ucgggcuucg ucucaccacu cgcagcguuc cggugacugc 1560
cgaauuaggu guacgucgac guccccgagg gugccgaccc uucucaacug guugucccg 1620
accggggcgu ccugucgaag gcugucgucc cgguccggga cggucugaag augccggacg 1680
acgggcugga gguaggagaa gguccguauc uuuuuugaa accacagacc cugucacuau 1740
aagaguaagu ucgacgucac aagucguguc gggcagcacu auaagagccg gaggaaccgg 1800
agguucucca uggagagguc cugagccgac agugucuacu ccgcaccacc ccgcccggucc 1860
ugaccucccg ggucuccuc gcugucgucc cuguccucuu cggugucggu ccguccgua 1920
agcacguggg gguagcacag uuccugccac ucagugagaa ccgugcccu uaggcgcaag 1980
guuacguggc aaggccggc gccuccgacc uagccagggc cacagaagau accuccaguu 2040
uugucgcacc uaccgagag guccgcuaga cagg 2074

```

```

<210> SEQ ID NO 20
<211> LENGTH: 2060
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized nucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1030)..(1031)
<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
1030 and 1031 comprises P04

```

```

<400> SEQUENCE: 20

```

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcgccg ggaacggugc auugaacgc ggauucccg ugccaagagu 120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
ugucccugcu gucgcuccu cugggcccucc caguccuggg cgcaccacca cgccucaucu 240
gugacagccg aguccuggag agguaccucu uggaggcca ggaggccgag aauaucacga 300
cgggcugugc ugaacacugc agcuugaug agaauaucac ugucccagac accaaaguaa 360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
uggcccugcu gucggagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
cucugcuucg ggcucugga gcccagaagg aagccaucuc ccuccagau gcggccucag 600
cugcuccacu ccgaacaau acugcugaca cuuuccgaa acucuuccga gucuacucca 660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcuaaa aaaaaaaaaa 840
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 900
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 960
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1020
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1080
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaucgaacu acguugaau aaaauaaucc 1260
uguuccgacc acccgugacc ucaccguuga aggucccggc ccucuccgug accccucccc 1320

```

-continued

```

aguguccua cggugggag uagacagggg acaggacguc cggaggggac acaugucgaa 1380
gucgaaaggg gccuccuuua accucaucug agccuuuca aacgccuuuc acagucguca 1440
cuaacaagcc ucaccucguc gacuccggcg uagaccuccc cucuaccgaa ggaagacccg 1500
agggucucgg gcuucgucuc accacuccga cgcuuccggu gacugccgaa auagguguac 1560
gucgacgucc cggagggugc cgacccuucu caacugguug ucccggaccg gggcguccug 1620
ucgaaggcug ucgucccggu cggggacggu cugaagaugc cggacgacgg gcuggaggua 1680
ggagaagguc cguaucuuua auugaaacca cagaccucgu cacuaaaga guaaguucga 1740
cgucacaagu cgugucgggc agcacuaaaa gagccggagg aaccggaggu ucuccaugga 1800
gagguccuga gccgacagug ucuacuccgc accaccccgcc ggguccugac ccuccggguc 1860
uccucgcug ucgucccugu ccucuucggu gucgguccgu ccuguaagca cgugggggua 1920
gcacaguucc ugccacucag ugagaaccgu gccccuagg cgcaagguua cguggcaagg 1980
gccggcgccu ccgaccuagc cagggccaca gaagauaccu ccaguuuugu cgcaccuacc 2040
gcagaggucc gcuagacagg 2060

```

<210> SEQ ID NO 21

<211> LENGTH: 2056

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1028)..(1029)

<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions 1028 and 1029 comprises PO4

<400> SEQUENCE: 21

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccaugaag acaccgggac 60
cgauccagcc uccgcgcccg ggaacggugc auuggaacgc ggauuccccg ugccaagagu 120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
ugucccugcu gucgcucccu cugggcccucc caguccuggg cgcaccacca cgccucaucu 240
gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aaauacacga 300
cgggcugugc ugaacacugc agcuugaaug agaauaucac uguccagac accaaaguuu 360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
uggcccugcu gucggaaagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc ccuccagau gcggccucag 600
cugcuccacu ccgaacaauc acugcugaca cuuuccgcaa acucuuccga gucuacucca 660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
gacggguggc auccuguga cccucccca guggcucucc uggeccugga aguugccacu 780
ccagugccca ccagccuugu ccuaauaaaa uuaaguugca ucaagcuaaa aaaaaaaaaa 840
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 900
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 960
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1020
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1080
aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1140

```

-continued

```

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 1200
aaaaaaaaa aaaaaaaaaa aaaaaaaaaa cgaacuacgu ugaauuaaaa uaauccuguu 1260
ccgaccacc gugaccucac cguugaaggu cccgguccuc uccgugaccc cuccccagug 1320
ucccuacggu gggcaguaga caggggacag gacguccgga ggggacacau gucgaagucg 1380
aaaggggccu ccuuuaaccu caucugagcc uucucaaacg ccuucacag ucgucacuaa 1440
caagccucac cucgucgacu cgggcguaga ccuccccucu accgaaggaa gacccgaggg 1500
ucucgggcuu cgucucacca cuccgacgcu uccggugacu gccgaaauag guguaucgucg 1560
acguccccga gggugccgac ccuucucaac ugguuugucc ggaccggggc guccugucga 1620
aggcugucgu cccgguccgg gacggucuga agauggccga cgaccgggug gagguaggag 1680
aagguccgua ucuuuaauug aaaccacaga ccugucacu auaagagua guucgacguc 1740
acaagucgug ucgggcagca cuauaagagc cggaggaacc ggagguucuc cauggagagg 1800
uccugagccg acagugucua cuccgcacca ccccgcgguu ccugaccuc cgggucuccc 1860
ucgucugcgu cccuguccuc uccggugucg guccguccug uaagcacgug gggguagcac 1920
aguuccugcc acucagugag aaccgugccc cuuagggcga agguuacgug gcaagggccg 1980
gcgccuccga ccuagccagg gccacagaag auaccuccag uuuugucgca ccuaccgag 2040
agguccgcu gacagg 2056

```

<210> SEQ ID NO 22

<211> LENGTH: 1802

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (901)..(902)

<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
901 and 902 comprises P04

<400> SEQUENCE: 22

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcgcccg ggaacggugc auuggaacgc ggauuccccg ugccaagagu 120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
ugucccugcu gucgcuccu cugggcccucc caguccuggg cgcaccacca cgccucaucu 240
gugacagccg aguccuggag agguaccucu uggaggccaa ggaggccgag aaauacacga 300
cgggucuguc ugaacacugc agcuugaaug agaauaucac ugucccagac accaaaguua 360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
uggcccugcu gucggaagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc cccuccagau gcggccucag 600
cugcuccacu ccgaacaau acugcugaca cuuuccgcaa acucuuccga gucuaucucca 660
auuuccuccg gggaaagcug aagcuguaca caggggaggg cugcaggaca ggggacagau 720
gacggguggc auccuguga cccucccca gugccucucc uggcccugga aguugccacu 780
ccagugccca ccaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa 840
aaaaaaaaa aaaaaaagcc uuguccuaau aaaaauaagu ugcaucaagc ucgacucucg 900
gggcucucag cucgaacuac guugaauuaa aauaauccug uuccgaaaaa aaaaaaaaaa 960

```

-continued

```

aaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa aaaaaaaaaa ccacccguga 1020
ccucaccguu gaaggucccg guccucuccg ugacccuccc ccaguguccc uacggugggc 1080
aguagacagg ggacaggacg uccggagggg acacaugucg aagucgaaag gggccuccuu 1140
uaaccucauc ugagccuucu caaacgccuu ucacagucgu cacuaacaag ccucaccucg 1200
ucgacuccgg cguagaccuc cccucuaccg aaggaagacc cgagggucuc gggcuucguc 1260
ucaccacucc gacgcuuccg gugacugccg aaauaggugu acgucgacgu ccccgagggg 1320
gccgaccuuu cucaacuggu ugucccgac cggggcgucc ugucgaaggc ugucgucccg 1380
guccgggacg gucugaagau gccggacgac gggcuggagg uaggagaagg uccguaucuu 1440
uaauugaaac cacagaccu gucacuaaa gaguaaguuc gacgucaca gucgugucgg 1500
gcagcacuau aagagccgga ggaaccggag guucuccaug gagagguccu gagccgacag 1560
ugucuacucc gcaccacccc gcggguccug acccuccggg ucuccucgc ugucgucccu 1620
guccucuucg gugucggucc guccuguaag cacguggggg uagcacaguu ccugccacuc 1680
agugagaacc gugcccuua ggcgcaaggu uacguggcaa gggccggcgc cuccgaccua 1740
gccagggcca cagaagauac cuccaguuuu gucgcaccua ccgagaggu ccgcuagaca 1800
gg 1802

```

```

<210> SEQ ID NO 23
<211> LENGTH: 1786
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized nucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (893)..(894)
<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
      893 and 894 comprises PO4

```

```

<400> SEQUENCE: 23
ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcgcccg ggaacggugc auugaaacgc ggauuccccg ugccaagagu 120
gacucaccgu ccuugacacg augggggugc acgaaugucc ugccuggcug uggcuucucc 180
ugucccugcu gucgcucccu cugggccucc caguccuggg cgcaccacca cgccucaucu 240
gugacagccg aguccuggag agguaccucu uggaggcca gaggccgag aauaucacga 300
cgggcugugc ugaacacugc agcuugaaug agaauaucac ugucccagac accaaaguua 360
auuucuaugc cuggaagagg auggaggucg ggcagcaggc cguagaaguc uggcagggcc 420
uggcccugcu gucggagcu guccugcggg gccaggcccu guuggucaac ucuucccagc 480
cgugggagcc ccugcagcug cauguggaua aagccgucag uggccuucgc agccucacca 540
cucugcuucg ggcucuggga gcccagaagg aagccaucuc cccuccagau gcggccucag 600
cugcuccacu ccgaacaau acugcugaca cuuuccgaa acucuuccga gucuacucca 660
auuuccuccg gggaaagcug aagcuguaca caggggaggc cugcaggaca ggggacagau 720
gacggguggc aaaaaaaaaa aaaaauccu gugacccuc ccaaaaaaaaa aaaaaaaag 780
ugccucuccu ggcccuggaa aaaaaaaaaa aaaguugcca cuccagugcc caccaaaaaa 840
aaaaaaaaag ccuuguccua auaaaauuaa guugcauca gcucgacucu cggggcucuc 900
agcucgaacu acguugaauu aaaaauaacc uguuccgaaa aaaaaaaaaa aaccaccgu 960
gaccucaccg uugaaaaaaaa aaaaaaagg ucccguuccu cuccgugaaa aaaaaaaaaa 1020

```

-continued

```

aaacccucc ccagugucc uaaaaaaaaa aaaaaacggu gggcaguaga caggggacag 1080
gacguccgga ggggacacau gucgaagucg aaaggggccu ccuuuaaccu caucugagcc 1140
uucucaaacg ccuuucacag ucgucacuaa caagccucac cucgucgacu ccggcguaga 1200
ccuccccucu accgaaggaa gacccgaggg ucucgggcuu cgucucacca cuccgacgcu 1260
uccggugacu gccgaaauag guguacgucg acgucgccga gggugccgac ccuucucaac 1320
ugguugucc ggaccggggc guccugucga aggucgucgu cccgguccgg gacggucuga 1380
agaugccgga cgacgggucg gagguaggag aagguccgua ucuuuauug aaaccacaga 1440
cccugucacu auaagaguaa guucgacguc acaagucgug ucgggcagca cuauaagagc 1500
cggaggaacc ggagguucuc cauggagagg uccugagccg acagugucua cuccgcacca 1560
ccccgcgggu ccugaccuc cgggucucc ucgucgucgu cccuguccuc uucggugucg 1620
guccguccug uaagcacgug gggguagcac aguuccugcc acucagugag aaccgugccc 1680
cuuaggcgca agguuacgug gcaagggccg gcgccuccga ccuagccagg gccacagaag 1740
auaccuccag uuuugucgca ccuaccgagc agguccgcu gacagg 1786

```

<210> SEQ ID NO 24

<211> LENGTH: 2640

<212> TYPE: RNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Chemically synthesized nucleotide

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1320)..(1321)

<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions 1320 and 1321 comprises PO4

<400> SEQUENCE: 24

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcggccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu 120
gacucaccgu ccuugacacg augcuguuca accuucggau cuugcugaac aacgcugcgu 180
uccggaaugg ucacaacuuc augguccgga acuucagaug cggccagccg cuccagaaca 240
aggugcagcu caaggggagg gaccuccuca cccugaaaaa cuucaccgga gaagagauca 300
aguacaugcu guggcuguca gccgaccuca aaauccggau caagcagaag ggccaauacc 360
uuccuuugcu gcagggaaag ucccugggga ugaucuuuga gaagcgcagc acucgcacua 420
gacugucaac ugaaacccgc uucgcgcugc ugggaggaca cccugcuuc cugaccaccc 480
aagauaucca ucugggugug aacgaauccc ucaccgacac agcgcgggug cugucgucca 540
uggcagacgc gguccucgcc cgcguguaca agcagucuga ucuggacacu cuggccaagg 600
aagccuccau uccuaucauu aauggauugu ccgaccucua ccauccauc cagauucugg 660
ccgauuaucu gacucugcaa gaacauuaca gcucccugaa ggggcuuacc cuuucgugga 720
ucggcgacgg caacaacauu cugcacagca uuauaugag cgcugccaag uuuggaaugc 780
accuccaagc agcgaccccg aagggauacg agccagacgc cuccgugacg aagcuggcug 840
agcaguacgc caaggagaac ggcacuaagc ugcugcucac caacgaccu cucgaagccg 900
cccacggugg caacgugcug aucaccgaa cccuggaucuc caugggacag gaggaggaaa 960
agaagaagcg ccugcaagca uuucaggggu accaggugac uaugaaaacc gccaaaggucg 1020
ccgccucgga cuggaccuuc uugcacuguc ugcccagaaa gcccgaaagag guggacgacg 1080
agguguucua cagcccgcgg ucgucggucu uuccggaggc cgaaaacagg aaguggacua 1140

```

-continued

```

ucauggccgu gauggugucc cugcugaccg auuacucucc gcagcugcag aaaccaaagu 1200
ucugacgggu ggcaucccug ugacccucc ccagugccuc uccuggccu ggaaguugcc 1260
acuccagugc ccaccagccu uguccuaaua aaauuaaguu gcaucaagcu cgacucucgg 1320
ggcucucagc ucgaacuacg uugaauuaaa auaauccugu uccgaccacc cgugaccuca 1380
ccguugaagg ucccgguccu cuccgugacc ccucccagu gucccuacgg ugggcagucu 1440
ugaaaccaa gacgucgacg cccucuaaua gccagucguc ccugugguag ugccgguacu 1500
aucaggugaa ggacaaaagc cggaggccuu ucuggucgcu ggcgcccagc aucuugugga 1560
gcagcaggug gagaagcccg aaagaccgcu cugucacguu cuuccagguc aggcuccgcc 1620
gcuggaaccg caaaaguau caguggacca uggggacuuu acgaacguc gcgaagaaga 1680
aaaggaggag gacagguac cucuaggucc auagccacua gucgugcaac gguggcacc 1740
gccgaagcuc ucccagcaac cacucgucgu cgaauccagg caagaggaac cgcaugacga 1800
gucggucgaa gcagugccuc cgcagaccga gcauaggaa gcccagcga cgaaccucca 1860
cguaaaguuu gaaccgucg gaguaguau acgacacguc uuacaacaac ggcagcggcu 1920
aggugcuuuc ccuucgggg aagucccucg acauuacaag aacgucucag ucuuuagcc 1980
ggucuuagac cuaccuacc aucuccagcc uguuagguaa uuacuauccu uaccuccgaa 2040
ggaaccgguc ucacaggucu agucugacga acaugugcgc ccgucuccg cgcagacggu 2100
accugcuguc gugggcgca cacagccacu ccuaagcaa gugugggucu accuauagaa 2160
cccaccaguc cuucgucucc acaggagggu cugcgcgcuu cggccaaagu caacugucag 2220
aucacgcuca cgacgcgaag agcuucuagu aggggucucc gaaaggagc ucguuuccuu 2280
ccauaagcgg gaagacgaac uaggccuaa acuccagccg acugucggug ucguacauga 2340
acuagagaag aggccacuuc aaaaagucc acuccuccag ggagggaac ucgacgugga 2400
acaagaccuc gccgaccgca guagacuua aggcucggu cuucaacacu gguaaggccu 2460
ugcugcga caagucguuc uaggcuucca acuugucgua gcacaguucc ugccacucag 2520
ugagaaccgu gcccuuagg cgcaagguua cguggcaagg gccggcgccu ccgaccuagc 2580
cagggccaca gaagauaccu ccaguuuugu cgcaccuacc gcagaggucc gcuagacagg 2640

```

```

<210> SEQ ID NO 25
<211> LENGTH: 3228
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized nucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1614)..(1615)
<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
1614 and 1615 comprises PO4

```

```

<400> SEQUENCE: 25
ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccaugaag acaccgggac 60
cgauccagcc uccgcccggc ggaacggugc auuggaacgc ggauuccccg ugccaagagu 120
gacucaccgu ccuugacacg augagcaccg ccgucgugga gaaccccggc cugggcccga 180
agcugagcga cuucggccag gagaccagcu acaucgagga caacugcaac cagaacggcg 240
ccaucagccu gauuucagc cugaaggagg aggugggccc ccuggccaag gucgucgccc 300
uguucgagga gaacgacgug aaccugacc acaucgagag ccgcccagc cgccugaaga 360
aggacgagua cgaguucuc acccaccugg acaagcgcag ccugcccgcc cugaccaaca 420

```


-continued

ucaucaagau	ccugcgccac	gacaucggcg	ccaccgugca	cgagcugagc	cgcgacaaga	480
agaaggacac	cgugcccugg	uucggccgca	ccauccagga	gcuggaccgc	uucggccaacc	540
agauccugag	cuacggcgcc	gagcuggacg	ccgaccaccc	cgguucaag	gacggcgugu	600
accgccccg	ccgcaagcag	uucggcgaca	ucgccuacaa	cuaccgccac	ggccagccca	660
uucggccgcu	ggaguacaug	gaggaggaga	agaagaccug	gggcaccgug	uucaagacc	720
ugaagagccu	guacaagacc	cacgcccugcu	acgaguacaa	ccacaucuu	cccugcugg	780
agaaguacug	cgguuuccac	gaggacaaca	uucggcagcu	ggaggacgug	agccaguucc	840
ugcagaccug	caccggcuuc	cgcccugcgcc	ccguggccgg	ccugcugagc	agcccgagcu	900
uccuggggcg	ccuggccuuc	cgcguguuuc	acugcaccca	guacaucgc	cacggcagca	960
agcccAugua	cacggccgag	cccgacaucu	gccacgagcu	gcuggggccac	gugcccugcu	1020
ucagcgaccg	cagcuucgcc	caguucagcc	aggagaucgg	ccuggccagc	cuggggcgccc	1080
ccgacgagua	caucgagaag	cuggccacca	ucuaucgguu	caccguggag	uucggccugcu	1140
gcaagcaggg	cgacagcauc	aaggccuacg	gcgcccggccu	gcugagcagc	uucggcgagc	1200
ugcaguacug	ccugagcgag	aagcccagc	ugcugccccc	ggagcuggag	aagaccgcca	1260
uccagaacua	caccgugacc	gaguuccagc	cccuguacua	cguggccgag	agcuucaacg	1320
acgccaagga	gaaggugcgc	aacuucggcg	ccaccaucac	ccgcccuc	agcgugcgcu	1380
acgacccua	caccagcgc	aucgaggugc	uggacaacac	ccagcagcug	aagaucgug	1440
ccgacagcau	caacagcgag	aucggcaucc	ugugcagcgc	ccugcagaag	aucaaguaac	1500
ggguggcauc	ccugugacc	cucggcagug	ccucuccugg	cccuggaagu	ugccacucca	1560
gugcccacca	gccuuguccu	aaauuuuu	aguugcauca	agcucgacuc	ucggggcucu	1620
cagcucgaac	uacguugaau	uaaaauaac	cuguuccgac	caccggugac	cucaccguug	1680
aaggucggcg	uccucuccgu	gacggcucc	cagugucccu	acggugggca	augaacuaga	1740
agacgucgg	cgacgugucc	uacggcuaga	gcgacaacua	cgacagccgg	uccuagaagu	1800
cgacgaccca	caacaggucc	uggagcuacg	cgaccacau	cccagcauc	gcgugcgacu	1860
uucggccccc	cuaccaccgc	cgcuucaacg	cguggaagag	gaaccgcagc	aacuucgaga	1920
gccggugcau	caugucggcg	accuugagcc	agugccacau	caagaccuac	cgccagaaga	1980
ggucgagguc	cccugcugcg	aaccggaaga	gcgaguccgu	caugacgucg	agcgguucg	2040
acgagucguc	cgcccgcgcc	aucgggaacu	acgacagcgg	gacgaacgug	uccggcuuga	2100
ggugccacuu	ggucaucua	caccggucga	agagcuacau	gagcagcccc	cgcgggucgg	2160
accggucgg	cuagaggacc	gacuugacc	gcuucgacgc	cagcgacuug	uucggcugca	2220
ccgggucguc	gagcaccguc	uacagcccga	gccccacau	guaccggaac	gacggcaccg	2280
ccuacaugac	ccacgucacc	uugugcgccu	uccggucgg	cgguccuuc	agcgccgacg	2340
agucgucgg	ccggugcccc	gcgucggccu	ucggccacgu	ccagacgucc	uugaccgagu	2400
gcaggagguc	gacggccuac	aacaggagca	ccuucggcg	caugaagagg	ucgucggccu	2460
ucuaaccaa	caugagcauc	gucggcacc	agaacauguc	cgagaagucc	cagaacuugu	2520
gccacgggg	ccagaagaag	aggaggaggu	acaugaggug	cgccccuac	ccgaccggca	2580
ccgccaucua	cauccgcuac	agccgcuuga	cgaacggcg	ccgcccag	ugcccagga	2640
acuucggccc	caccagcgc	aggucgagcc	gcggcaucga	guccuagacc	aaccgcuucg	2700
ccaggucgag	gaccuaccac	gccccuugg	uucggugcca	caggaagaag	aacagcgccg	2760
agucgagcac	gugccaccgc	ggcuacagca	ccgcuucca	gaacuacuac	aaccagucc	2820

-continued

```

gcccguccga cgccaacagg uccacccacu ucuugagcau gagcaggaag aaguccgccg 2880
accccgccga gagcuacacc caguccaagu gcagcaagag gagcuugucc gcgucgugga 2940
accggucccg cggguggagg aggaaguccg acuuuaguc cgacuaccgc ggcaagacca 3000
acgucaacag gagcuacauc gaccagagga ccggcuucag cgagucgaac gccggguccg 3060
gccccaaagag gucgugccgc cacgaguagc acaguuccug ccacucagug agaaccgugc 3120
ccuuuaggcg caagguuacg uggcaagggc cggcgccucc gaccuagcca gggccacaga 3180
agauaccucc aguuuugucg caccuaccgc agagguccgc uagacagg 3228

```

```

<210> SEQ ID NO 26
<211> LENGTH: 9396
<212> TYPE: RNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Chemically synthesized nucleotide
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (4698)..(4699)
<223> OTHER INFORMATION: 3'-3' bridge between nucleotides at positions
4698 and 14699 comprises PO4

```

```

<400> SEQUENCE: 26

```

```

ggacagaucg ccuggagacg ccauccacgc uguuuugacc uccauagaag acaccgggac 60
cgauccagcc uccgcggccg ggaacggugc auuggaacgc ggauuucccg ugccaagagu 120
gacucaccgu ccuugacacg augcaacgcu cuccucuuga aaaggccucg guggugucca 180
agcucuucuu cucguggacu agacccaucc ugagaaaggg guacagacag cgcuuggagc 240
uguccgauau cuaucaaauc ccuuccgugg acuccgcgga caaccugucc gagaagcucg 300
agagagaaug ggacagagaa cucgccucaa agaagaacc gaagcugauu aaugcgcuua 360
ggcggugcuu uuucuggcgg uucauguucu acggcaucuu ccucuaccug ggagagguca 420
ccaaggccgu gcagccccug uugcugggac ggauuuuugc cuccuacgac cccgacaaca 480
aggaagaaag aagcaucgcu aucuacuugg gcaucggucu gugccugcuu uucaucgucc 540
ggaccucuu guugcauccu gcuauuuucg gccugcauca cauuggcaug cagaugagaa 600
uugccauguu uucccugauc uacaagaaaa cucugaagcu cucgagccgc gugcuugaca 660
agauuuccau cggccagcuc gugucccugc ucuccaaca ucugaacaag uucgacgagg 720
gccucgccc uggccacuuc guguggaucg cccucugca aguggcgcuu cugaugggccc 780
ugaucuggga gcugcugcaa gccucggcau ucugugggcu uggauuccug aucgugcugg 840
cacuguucca ggccggacug gggcggauga ugaugaagua cagggaccag agagccggaa 900
agauuuccga acggcuggug aucacuucgg aaaugaucga aaacauccag ucagugaagg 960
ccuacugcug ggaagaggcc auggaaaaga ugaauaaaa ccuccggcaa accgagcuga 1020
agcugacccg caaggccgcu uacgugcgcu auuacaacuc guccgcuuc uucuucuccg 1080
gguucucgu gguguuucuc uccgugcucc ccuacgccc gauuaaggga aucauccuca 1140
ggaagauuu caccaccauu uccuucugua ucgugcuccg cauggccgug acccggcagu 1200
ucccaugggc cgugcagacu ugguacgacu ccugggagc cauaacaag auccaggacu 1260
uccuucaaaa gcaggaguac aagaccucg aguacaaccu gacuacuacc gaggucguga 1320
uggaaaacgu caccgccuuu ugggaggagg gauuuggcga acuguucgag aaggccaagc 1380
agaacaaca caaccgcaag accucgaacg gugacgacuc ccucuucuu ucaaacuua 1440
gccugcucgg gacgcccug cugaaggaca uuaacuuaa gaucgaaaga ggacagcucc 1500

```

-continued

uggcgguggc	cggaucgacc	ggagccgga	agacuuccu	gcugauggug	aucaugggag	1560
agcuugaacc	uagcgaggga	aagaucaagc	acuccggccg	caucagcuuc	uguagccagu	1620
uuuccuggau	caugcccga	accuuuaagg	aaaacaucau	cuucggcgug	uccuacgaug	1680
aauaccgcu	cgguccgug	aucaaagccu	gccagcugga	agaggauuu	ucaaaguucg	1740
cggagaaaga	uaacaucgug	cugggccaag	ggguuuuac	cuugucgggg	ggccagcggg	1800
cuagaaucuc	gcuggccaga	gccguguaua	aggacgccga	ccuguaucuc	cuggacucucc	1860
ccuucggaua	ccuggacguc	cugaccgaaa	aggagauuu	cgaaucgugc	gugugcaagc	1920
ugauggcu	caagacucg	auccucguga	ccuccaaaau	ggagcaccug	aagaaggcag	1980
acaagauucu	gauucugcau	gagggguccu	ccuacuuuu	cggcaccuuc	ucggaguugc	2040
agaacuugca	gcccgaucuc	ucaucgaagc	ugauggguug	cgacagcuuc	gaccaguucu	2100
ccgccgaaag	aaggaacucg	auccugacgg	aaaccuugca	ccgcuucucu	uuggaaggcg	2160
acgccccugu	gucauggacc	gagacuaaga	agcagagcuu	caagcagacc	ggggaauucg	2220
gcgaaaagag	gaagaacagc	aucuugaacc	ccauuaacuc	cauccgcaag	uucucaaucg	2280
ugaaaagac	gccacugcag	augaacggca	uugaggagga	cuccgacgaa	ccccuugaga	2340
ggcgccuguc	ccuggugccg	gacagcgagc	agggagaagc	cauccugccu	cggauuuccg	2400
ugaucuccac	ugguccgacg	cuccaagccc	ggcgccgcca	guccgugcug	aaccugauga	2460
cccacagcgu	gaaccagggc	caaaacauuc	accgcaagac	uaccgcaucc	accggaaag	2520
ugucccuggc	accucaagcg	aaucuuaccg	agcucgacau	cuacucccg	agacugucgc	2580
aggaaaccgg	gcucgaaau	uccgaagaaa	ucaacgagga	ggaucugaaa	gagugcuucu	2640
ucgacgauau	ggagucgaua	cccgccguga	cgacuuggaa	cacuuaucug	cgguaacauca	2700
cugugcacia	gucauugauc	uucgugcuga	uuuggugccu	ggugauuuuc	cuggccgagg	2760
ucgcccuc	acugguggug	cucuggcugu	ugggaaacac	gccucugcaa	gacaaggga	2820
acuccacgca	cucgagaaac	aacagcuau	ccgugauuu	cacuuccacc	uccucuauu	2880
acguguucua	caucuacguc	ggaguggcgg	auaccucgcu	cgcgaugggu	uucuucagag	2940
gacugccgcu	gguccacacc	uugaucaccg	ucagcaagau	ucuucaccac	aagauguugc	3000
auagcgugcu	gcaggcccc	auguccacc	ucaacacucu	gaaggccgga	ggcauucuga	3060
acagauucuc	caaggacauc	gcuauccugg	acgaucuccu	gccgcuuacc	aucuuugacu	3120
ucauccagcu	gcugcugauc	gugauuggag	caaucgcagu	gguggcggug	cugcagccuu	3180
acauuuucgu	ggccacugug	ccggucauug	uggcguucau	caugcugcgg	gccuacuucc	3240
uccaaaccag	ccagcagcug	aagcaacugg	aauccgaggg	acgaucccc	aucuucacuc	3300
accuugugac	gucguugaag	ggacugugga	cccuccgggc	uuucggacgg	cagccuacu	3360
ucgaaaccu	cuuccacaag	gcccugaacc	uccacaccgc	caauugguuc	cuguaccugu	3420
ccaccucg	gugguuccag	augcgcaucg	agaugauuuu	cgucaucuc	uucaucgcg	3480
ucacauucau	cagcauccug	acuaccggag	aggagagggg	acgggucgga	auaaucuga	3540
cccucgccau	gaacauuau	agcaccucg	agugggcagu	gaacagcucg	aucgacgugg	3600
acagccugau	gcgaagcug	agccgcgugu	ucaaguucau	cgacaugccu	acugagggaa	3660
aaccacuaa	guccacuaag	ccuacaaaa	auggccagcu	gagcaagguc	augaucaucg	3720
aaaacucca	cgugaagaag	gacgauuuu	ggccuccgg	aggucaaaug	accgugaagg	3780
accugaccg	aaaguacacc	gagggaggaa	acgccauucu	cgaaaacauc	agcuucucca	3840

-continued

uuucgccggg	acagcggguc	ggccuucucg	ggcggaccgg	uuccgggaag	ucaacucugc	3900
ugucggcuuu	ccuccggcug	cugaauaccg	agggggaaau	ccaaaugac	ggcgugucuu	3960
gggauuccau	uacucugcag	caguggcgga	aggccuucgg	cgugaucccc	cagaaggugu	4020
ucaucuucuc	ggguaccuuc	cggaagaacc	uggauccuua	cgagcagugg	agcgaccaag	4080
aaaucuggaa	ggucgccgac	gaggucggcc	ugcgcuccgu	gauugaacaa	uuuccuggaa	4140
agcuggacuu	cgugcucguc	gacgggggau	guguccuguc	gcacggacau	aagcagcuca	4200
ugugccucgc	acgguccgug	cucuccaagg	ccaagauucu	gcugcuggac	gaaccuucgg	4260
cccaccugga	uccggucacc	uaccagauca	ucaggaggac	ccugaagcag	gccuuugccg	4320
auugcaccgu	gauucucugc	gagcaccgca	ucgaggccau	gcuggagugc	cagcaguucc	4380
uggucaucga	ggagaacaag	guccgccaa	acgacuccau	ucaaagcuc	cucaacgagc	4440
ggucgcuguu	cagacaagcu	auuucaccgu	ccgauagagu	gaagcucuuc	ccgcaucgga	4500
acagcucaaa	gugcaaacg	aagccgcaga	ucgagccuu	gaaggaagag	acugaggaag	4560
aggugcagga	caccggcuu	uaacgggugg	caucccugug	acccucccc	agugccucuc	4620
cuggcccugg	aaguugccac	uccagugccc	accagccuug	uccuaaaaa	auuaaguugc	4680
aucaagcucg	acucucgggg	cucucagcuc	gaacuacguu	gaauaaaau	aauccuguuc	4740
cgaccacccg	ugaccucacc	guugaagguc	ccgguccucu	ccgugacccc	uccccagugu	4800
cccuacggug	ggcaauuucg	gcccacagga	cguggagaag	gagucagaga	aggaaguucc	4860
gacgcuagac	gccgaagcua	aacgugaaac	ucgacaaggc	uacgcccuc	ucgaagugag	4920
auagccugcc	acuuuucga	acagacuugu	cgucggcgag	caacuccucg	aaaacuacc	4980
ucagcauac	cgccuggaac	aagaggagcu	acugguccuu	gacgaccgug	aggucguacc	5040
ggagcuacgc	cacgagcugc	ucuuagugcc	acguuagccg	uuuccggacg	aagucccagg	5100
aggacuacua	gaccauccac	uggccuaggu	ccaccggcu	uccaagcagg	ucgucgucuu	5160
agaaccggaa	ccucucgugc	cuggcacgcu	ccguguacuc	gacgaauaca	ggcacgcugu	5220
ccuguguagg	gggcagcugc	ucgugcuuca	ggucgaaagg	uccuuuaaca	aguuagugcc	5280
ucgcguccgg	cuggagcagc	cguggaagg	ucuaaagaac	cagcgaggug	acgagcauuc	5340
cuagguccaa	gaaggccuuc	caugggcucu	ucuacuugug	gaagacccc	uagugcggu	5400
uccggaaggc	ggugacgacg	ucucauuacc	uuagguucu	gugcggcagu	uaaacuaaa	5460
gggggagcca	uaagucgucg	gccuccuuc	ggcugucguc	ucaacugaag	ggccuuggcc	5520
aggcgggcuc	uuccggcugg	gcgacagggc	cgcuuuaccu	cuucgacuac	aaaagcucuu	5580
accgcaaagg	agggagccac	augaaacgcc	aguccaggaa	gugccaguaa	acuggaggcc	5640
ucccgguua	uagcaggaag	aagugcacc	ucaaaagcua	cuaguacugg	aacgagucga	5700
ccgguaaaa	caucccgau	caccugauc	accaaaagg	gagucauccg	uacagcuacu	5760
ugaacuugug	cgccgacugc	gaagcguagu	ccgacaggug	cagcuagcuc	gacaagugac	5820
gggugacguc	ccacgaguau	uacaaguacc	gcucccaguc	cuaauaaggc	ugggcagggg	5880
gagggagagg	ccaucagucc	uacgacuacu	uacacuggcg	cuacuucuc	uacugcuuuu	5940
aguagagcua	cgcuagacc	uugguggcgu	cccaccuguc	cauguccuug	guuaaccgcc	6000
acaccuccaa	gucccgaac	accuucucc	aaagcuucau	cccgacggca	ggcuuucggg	6060
ccucccaggu	gucagggag	uugcugcagu	guuccacuca	cuucuacccc	cuagcagggg	6120
gccuaagguc	aacgaagucg	acgaccgacc	aaaccuccuu	cauccgggcg	ucguacuacu	6180
ugcggugua	cuggccgugu	caccggugcu	uuuacaucc	gacgucgugg	cgguggugac	6240

- continued

gcuaacgagg	uuagugcuag	ucgucgucga	ccuacuucag	uuucuaccu	ucgccguc	6300
cuagcagguc	cuacgcucac	aggaaccucu	uagacaaguc	uuacggaggc	cggaagucuc	6360
acaacuccca	ccuguacccc	cggacgucgu	gcgauacguu	guagaacacc	acuucuuaga	6420
acgacugcca	cuaguuccac	accuggucgc	cgucaggaga	cuucuuggg	uagcgcucgu	6480
cccuaaggcg	gugaggcugc	aucuacaucu	ugugcauuau	ucuccuccac	cuucacuauu	6540
agugccguau	cgacaacaaa	gagcucacgc	accucaaagg	gaacagaacg	ucuccgcaca	6600
aaggguguc	ggucucgug	uggucacucc	ggcgcuggag	ccgguccuuu	uaguggucg	6660
ugguuuaguc	gugcuucua	uuacugaaca	cgugcacua	cauggcgucu	auucacaagg	6720
uucagcagug	ccgcccauag	cugagguaua	gcagcuucuu	cgugagaaag	ucuaggagga	6780
gcaacuaaag	aagccuuuaa	agcucgggcc	aaaggacgcu	gucagaggcc	cucaucuaca	6840
gucgcagcca	uucuaagcga	acuccacggu	cccugugaaa	ggcccaccua	cgccaucaga	6900
acgccacuua	caaaaccggg	accaagugcg	accccagua	guccaagucg	ugccugacgg	6960
cggcgccccg	aaccucgcag	ccuggucacc	ucuagugccu	uuaggcuccg	uccuaccgaa	7020
gagggacgag	cgacaggccg	uggucccugu	ccgcggagag	uuccccaagc	agccucagga	7080
ggaguucgg	caaguagacg	ucaccgcaga	aaacgugcua	acucuugaac	gccuaccuca	7140
auuaccccaa	guucuacgac	aagaaggaga	aaagcggcuu	aaggggcccag	acgaacuucg	7200
agacgaagaa	ucagagccag	guacuguguc	cccgcagcgg	aagguuucuc	uucgccacgu	7260
uccaaaggca	guccuagcuc	aaggaagaaa	gccgcucuu	gaccagcuc	gacagcguug	7320
gguagucgaa	gcuacucuc	agcccagcgu	ucaagacguu	gaggcucuc	cacggcauuu	7380
ucauccuccu	ggggaguacg	ucuuagucuu	agaacagacg	gaagaagucc	acgagguaaa	7440
accuccagug	cuccuacgcu	cagaacaauc	gguagucgaa	cgugugcgug	cuaagcuucu	7500
agaggaaaag	ccaguccugc	agguccauag	gcuucucccu	cagguccucu	auguccagcc	7560
gcaggaauau	gugccgagac	cggucgcucu	aagaucgggc	gaccgggggg	cuguuccauu	7620
augggggaag	cgggucgugc	uacaauagaa	agaggcguu	gaaacuuuau	aggagaaggu	7680
cgaccguccg	aaacuagugc	cuggccaucg	ccauaaguag	cauccugugc	ggcuucucuc	7740
acaaaaggaa	uuaccaaggc	ccguacuagg	uccuuugac	cgaugucuc	gacuacgccg	7800
gccucacgaa	cuagaaagg	agcgauccaa	guucgagagg	guacuagug	uagucguc	7860
uucagaaagg	ccgaggccag	cuaggccggu	ggcgguccuc	gacaggagaa	agcuagaacu	7920
ucauuuacag	gaagucgugc	ccgcagggcu	cguccgacuu	caaacuuiuc	uucuccuca	7980
gcaguggcaa	gcuccagaac	gccaacaaca	acaagacgaa	ccggaagagc	uugucaagcg	8040
guuuaggag	gaggguuuuc	cgccacugca	aaagguagug	cuggagccau	caucagucca	8100
acaugagcuc	ccagaacaug	aggacgaaaa	cuuccuucag	gaccuagaac	aaauaccgag	8160
gguccucag	caugguucag	acgugccggg	uaccuugac	ggcccagugc	cgguacgccu	8220
cgugcuau	cuuccuuuac	caccacuucu	agaaggacuc	cuacuaagg	aaauagucc	8280
gcauuccuc	gugccucucu	uuguggugcu	ucuuaggccu	cuucucuuu	cgccugcuc	8340
acuuuucgc	gugcauucgc	cggaacgccc	agucgaaguc	gagccaaacg	gccuccaaaa	8400
guuaguagaa	aagguaccgg	agaagggucg	ucauccggaa	gugacugacc	uacaaaagcu	8460
aguaaaaggcu	ucacuagug	ucggcaagcc	uuuagaaagg	ccgagagacc	agggacauga	8520
aguaguagua	ggcgggguca	ggccggaccu	ugucacgguc	gugcuagucc	uuagguucgg	8580

-continued

gugucuuacg gcuccgaacg ucgucgaggg ucuaguccgg guagucuucg cggugaacgu	8640
cuccecgua ggugugcuuc acccgguccc gcuccgggag cagcuugaac aagucuaaca	8700
accucucguc ccugugcucg accggcuacc uuuagaacag uucgugcgcc gagcucucga	8760
agucuaaaaa gaacaucuag uccuuuuugu accguuaaga guagacguac gguuacacua	8820
cguccggcuu uuaucguccu acguuguucu cccaggccug cuacuuuucg uccgugucug	8880
gcuacggguu caucuaucgc uacgaagaaa gaaggaacaa cagccccagc auccuccguu	8940
auuaggcagg gucguugucc ccgacgugcc ggaaccacug gagagggucc aucuccuucu	9000
acggcaucuu guacuuggcg gucuuuuucg uggcggauuc gcguaauuag ucgaagccca	9060
agaagaaacu ccgcucaaga gacaggguaa gagagagcuc gaagagccug uccaacaggc	9120
gccucaggug ccuucccuaa acuaucuaa gccugucgag guucgcgaca gacaugggga	9180
aagaguccua cccagaucag gugcucuucu ucucgaaccu gugguggcuc cggaaaaguu	9240
cuccucucgc aacguagcac aguuccugcc acucagugag aaccgugccc cuuaggcgca	9300
agguuacgug gcaagggccg gcgccuccga ccuagccagg gccacagaag auaccuccag	9360
uuuugucgca ccuaccgag agguccgcua gacagg	9396

We claim:

1. A composition comprising a multimeric coding nucleic acid (MCNA) encapsulated or complexed with a liposomal delivery vehicle,

wherein the MCNA comprises two messenger RNAs (mRNAs) linked at 3' ends, via stable linkage, such that the multimeric coding nucleic acid has two 5' ends, and wherein the stable linkage is an oligonucleotide bridge comprising an internal 3'-to- 3' inverted phosphodiester linkage.

2. The composition of claim 1, wherein the delivery vehicle comprises one or more cationic lipids, one or more non-cationic lipids, one or more cholesterol-based lipids, and one or more PEGylated lipids.

3. A method of delivering MCNA for in vivo protein production, comprising administering the composition of claim 1 to a subject in need of delivery.

4. The method of claim 3, wherein the composition is administered via a route of delivery selected from the group consisting of intravenous delivery, subcutaneous delivery, oral delivery, subdermal delivery, ocular delivery, pulmonary delivery, intramuscular delivery, intrathecal delivery, and intraarticular delivery.

5. The method of claim 3, wherein the composition comprises human Cystic Fibrosis Transmembrane Conductance Regulator (hCFTR) mRNA.

6. The method of claim 3, wherein the composition comprises human phenylalanine hydroxylase (hPAH) mRNA.

7. The composition of claim 1, wherein each of the mRNAs encodes a protein of interest.

8. The composition of claim 7, wherein each of the mRNAs encodes a same protein.

9. The composition of claim 7, wherein each of the mRNAs encodes a distinct protein.

10. The composition of claim 1, wherein the mRNAs comprise a 5' untranslated region (5' UTR) and/or a 3' untranslated region (3' UTR).

11. The composition of claim 1, wherein the mRNAs comprise a 3' UTR.

12. The composition of claim 11, wherein the 3' UTR comprises a plurality of multi-A segments with spacers in between.

13. The composition of claim 1, wherein the oligonucleotide bridge comprises nucleosides selected from the group consisting of 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (ψ U), and 1-methyl-pseudouridine.

14. The composition of claim 1, wherein the mRNAs comprise one or more modified nucleosides.

15. The composition of claim 14, wherein the modified nucleosides are selected from the group consisting of 2'-OMe-A, 2'-OMe-G, 2'-OMe-C, 2'-OMe-U, 2'-F-A, 2'-F-G, 2'-F-C, 2'-F-U, LNA-A, LNA-G, LNA-C, LNA-U, N6-methyl-adenosine, 2-thiouridine (2sU), 5-methyl-cytidine (5mC), pseudouridine (ψ U), and 1-methyl-pseudouridine.

16. The composition of claim 1, wherein each of the mRNAs encodes an enzyme, a receptor, a ligand, a light chain or heavy chain of an antibody, a nuclease, or a DNA-binding protein.

17. The composition of claim 11, wherein the 3' UTR does not include a polyA tail.

18. The composition of claim 1, wherein each of the mRNAs is unmodified.

19. The composition of claim 1, wherein the oligonucleotide bridge is unmodified.

20. The composition of claim 1, wherein each of the mRNAs has a 5' cap.

* * * * *