

US011261233B2

(12) **United States Patent**
Liu et al.

(10) **Patent No.:** **US 11,261,233 B2**
(45) **Date of Patent:** **Mar. 1, 2022**

(54) **CD80 AND CD86 BINDING PROTEIN COMPOSITIONS AND USES THEREOF**

(71) Applicants: **ONCOC4, INC.**, Rockville, MD (US);
CHILDREN'S RESEARCH INSTITUTE, CHILDREN'S NATIONAL MEDICAL CENTER, Washington, DC (US)

(72) Inventors: **Yang Liu**, Washington, DC (US); **Pan Zheng**, Washington, DC (US); **Martin Devenport**, Gaithersburg, MD (US); **Wei Wu**, Washington, DC (US); **Xuexiang Du**, Washington, DC (US); **Mingyue Liu**, Washington, DC (US); **Fei Tang**, Washington, DC (US)

(73) Assignees: **ONCOC4, INC.**, Rockville, MD (US); **CHILDREN'S NATIONAL MEDICAL CENTER**, Washington, DC (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 335 days.

(21) Appl. No.: **16/333,888**

(22) PCT Filed: **Sep. 19, 2017**

(86) PCT No.: **PCT/US2017/052264**
§ 371 (c)(1),
(2) Date: **Mar. 15, 2019**

(87) PCT Pub. No.: **WO2018/053506**
PCT Pub. Date: **Mar. 22, 2018**

(65) **Prior Publication Data**
US 2019/0263889 A1 Aug. 29, 2019

Related U.S. Application Data

(60) Provisional application No. 62/396,667, filed on Sep. 19, 2016.

(51) **Int. Cl.**
C07K 14/705 (2006.01)
C07K 16/46 (2006.01)
A61K 38/00 (2006.01)
A61P 37/06 (2006.01)
A61P 35/00 (2006.01)
C07K 16/28 (2006.01)
A61K 39/395 (2006.01)
C07K 14/00 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**
CPC .. **C07K 14/70596** (2013.01); **A61K 39/39541** (2013.01); **A61P 35/00** (2018.01); **A61P 37/06** (2018.01); **C07K 14/00** (2013.01); **C07K 16/2818** (2013.01); **C07K 16/2827** (2013.01); **A61K 38/00** (2013.01); **A61K 2039/507** (2013.01); **C07K 2317/24** (2013.01); **C07K 2317/34** (2013.01); **C07K 2317/76** (2013.01); **C07K 2319/30** (2013.01)

(58) **Field of Classification Search**
None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

10,618,960 B2 * 4/2020 Liu A61P 35/00
2011/0081354 A1 4/2011 Korman et al.
2015/0051158 A1 2/2015 Akamatsu et al.
2019/0127468 A1 * 5/2019 Liu C07K 16/2878
2020/0283526 A1 * 9/2020 Liu A61P 35/00
2021/0047410 A1 * 2/2021 Liu C07K 16/2827

OTHER PUBLICATIONS

Buchbinder, E., et al., "CTLA-4 and PD-1 Pathways Similarities, Differences, and Implications of Their Inhibition," *Am J Clin. Oncol.*, vol. 39, No. 1, pp. 98-106 (Feb. 3, 2016).
Grosso, J., et al., "CTLA-4 Blockade in Tumor Models: An Overview of Preclinical and Translational Research," *Cancer Immun.*, vol. 13, No. 5, pp. 1-14 (Jan. 22, 2013).
Lipson, E., et al., "Successful Administration of Ipilimumab to Two Kidney Transplantation Patients with Metastatic Melanoma," *J. Clin. Oncol.*, vol. 32, No. 19, pp. e69-e71 (2014).
International Search Report of PCT/US2017/52264 dated Jan. 23, 2018.
Written Opinion of PCT/US2017/52264 dated Jan. 23, 2018.

* cited by examiner

Primary Examiner — Ilia I Ouspenski
(74) *Attorney, Agent, or Firm* — Polsinelli PC; Ron Galant

(57) **ABSTRACT**

This invention relates to cytotoxic T-lymphocyte-associated antigen-4 (CTLA-4) protein compositions and their use in the mitigation of autoimmune adverse events associated with cancer immunotherapy. Specifically, the disclosure provides a CTLA-4 protein comprising a mutant extracellular domain of CTLA-4, wherein the CTLA-4 protein exhibits reduced binding to an anti-CTLA-4 antibody as compared to a wild-type extracellular domain of CTLA-4, wherein the anti-CTLA-4 antibody has anti-cancer immunotherapeutic activity.

19 Claims, 35 Drawing Sheets

Specification includes a Sequence Listing.

FIG. 1A

MACLGQRHKAQLNLARTWPCTLLFFLLFIPVFCKAMHVAQPAVVLASSRGIASFVCEYASPGKAT
EVRTVLRQADSQVTEVCAATYMMGNELTFLDSICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL
MYPPYYLGIGNGTQIYVIDPEPCPDSDFLWILAAVSSGLFFYSFLITAVSLSKMLKRRSPLTTGVYVK
MPPTPECEKQFQPYFIPIN

FIG. 1B

MHVAQPAVVLASSRGIASFVCEYASPGKATEVRTVLRQADSQVTEVCAATYMMGNELTFLDDS
ICTGTSSGNQVNLTIQGLRAMDTGLYICKVELMYPPYYLGIGNGTQIYVIDPEPCPDSDQEPKSSD
KTHTSPPAPPELLGGSSVFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAK
TKPREEQYNSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRD
ELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPPVLDSDGSGFFLYSKLTVDKSRWQQGNVFN
SCSVMHEALHNHYTQKSLSLSPGK




FIG. 2




FIG. 3

Weight of hCTLA4-KI mice (12.30.15-Day42)





FIG. 4

FIG. 5






FIG. 6 D




FIG. 6A

FIG. 6B




FIG. 6C




FIG. 8




FIG. 9




FIG. 10

FIG. 11A





FIG. 11B






FIG. 12




FIG. 13




FIG. 14

CTLA-4Fc WT

MHVAQPAVVLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDS ICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGGSSVFLFPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEOYNSTYRVVSVLTVLHODWLNQKEYCKVSNKALPAPIEK
TISKAKGQPREPOVYITLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWOOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTLA-4Fc M1

iqvtqpsVVLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDS ICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGGSSVFLFPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEOYNSTYRVVSVLTVLHODWLNQKEYCKVSNKALPAPIEK
TISKAKGQPREPOVYITLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWOOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTLA-4Fc M2

MHVAQPAVVLASSRGIAS**Fp**CEYASPGKATEVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDS ICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGGSSVFLFPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEOYNSTYRVVSVLTVLHODWLNQKEYCKVSNKALPAPIEK
TISKAKGQPREPOVYITLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWOOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTLA-4Fc M3

MHVAQPAVVLASSRGIASFVCEYAS P **shnTd**EVVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDS ICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLLGIGNGTQIYVIDPEPCPD
DSDQEPKSSDKTHTSPSPAPELLGGSSVFLFPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEOYNSTYRVVSVLTVLHODWLNQKEYCKVSNKALPAPIEK
KTISKAKGQPREPOVYITLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWOOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTLA-4Fc M4

MHVAQPAVVLASSRGIASFVCEYASPGKATEVRVTVLR **tnDqm**VTEVCAATYMMGNELTFLDSDS ICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGGSSVFLFPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEOYNSTYRVVSVLTVLHODWLNQKEYCKVSNKALPAPIEK
TISKAKGQPREPOVYITLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWOOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTLA-4Fc M5

MHVAQPAVVLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAAT **ftekntvg**FLDSDS ICTGTSSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGGSSVFLFPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNKATKPREEOYNSTYRVVSVLTVLHODWLNQKEYCKVSNKALPAPIEK
TISKAKGQPREPOVYITLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPPVLDSDGSFFLYSKLTVDKSRWOOQGNVFSCVMHEALHNHYTQKSLSLSPGK

FIG. 16A

CTLA-4Fc M6

MHVAQPAVWLASSRGIASFVCEYASPGKATEVRVTVLRROADSQVTEVCAATYMMGNELTFLDSDICTGT'SSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGSSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKENWYVDGVEVHNAKTKPREEOYNSTRVVSVLTVLHODWLNNGKEYKCKVSNKALPAPIEK
TISKAKGOPREPOVYTLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTIA-4Fc M7

MHVAQPAVWLASSRGIASFVCEYASPGKATEVRVTVLRROADSQVTEVCAATYMMGNELTFLD**Ypfcs**GTS SGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGSSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKENWYVDGVEVHNAKTKPREEOYNSTRVVSVLTVLHODWLNNGKEYKCKVSNKALPAPIEK
TISKAKGOPREPOVYTLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTLA-4Fc M8

MHVAQPAVWLASSRGIASFVCEYASPGKATEVRVTVLRROADSQVTEVCAATYMMGNELTFLDSDICTGT'**nesr**VNLTIQGLRAMDTGLYICKVEL**MYPPPY**YLGIGNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGSSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKENWYVDGVEVHNAKTKPREEOYNSTRVVSVLTVLHODWLNNGKEYKCKVSNKALPAPIEK
TISKAKGOPREPOVYTLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTIA-4Fc M9

MHVAQPAVWLASSRGIASFVCEYASPGKATEVRVTVLRROADSQVTEVCAATYMMGNELTFLDSDICTGT'SSGNQVNLTIQGLRAMVDTGLYICKVEL**MYPPPY**YLGIGNGTQIYVIDPEPCPD
DSDQEPKSSDKTHTSPSPAPELLGSSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKENWYVDGVEVHNAKTKPREEOYNSTRVVSVLTVLHODWLNNGKEYKCKVSNKALPAPIEK
TISKAKGOPREPOVYTLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOQGNVFSCVMHEALHNHYTQKSLSLSPGK





CTLA-4Fc M10

MHVAQPAVWLASSRGIASFVCEYASPGKATEVRVTVLRROADSQVTEVCAATYMMGNELTFLDSDICTGT'SSGNQVNLTIQGLRAMMDTGLY**1**CKVEL**MYPPPY**YLGIGNGTQIYVIDPEPCPD
DSDQEPKSSDKTHTSPSPAPELLGSSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKENWYVDGVEVHNAKTKPREEOYNSTRVVSVLTVLHODWLNNGKEYKCKVSNKALPAPIEK
TISKAKGOPREPOVYTLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOQGNVFSCVMHEALHNHYTQKSLSLSPGK

CTIA-4Fc M11


MHVAQPAVWLASSRGIASFVCEYASPGKATEVRVTVLRROADSQVTEVCAATYMMGNELTFLDSDICTGT'SSGNQVNLTIQGLRAMDTGLYICKVEL**MYPPPYfvGm**GNGTQIYVIDPEPCPD
SDQEPKSSDKTHTSPSPAPELLGSSVFLFPPKPKDTLMI SRTPEVTCVVVDVSHEDPEVKENWYVDGVEVHNAKTKPREEOYNSTRVVSVLTVLHODWLNNGKEYKCKVSNKALPAPIEK
TISKAKGOPREPOVYTLPPSRDELTKNOVSLTCLVKGFYPSDIAVEWESNGOPENNYKTTTPVLDSDGSFFLYSKLTVDKSRWQOQGNVFSCVMHEALHNHYTQKSLSLSPGK

FIG. 16B



Protein Concentration

FIG. 18



CTLA-4Fc M12
 iqvtqPsVVLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAATftekntvgFLDSDSICTGTfnesrVNLTIQGLRAMDTGLYICKVELMYPPPYfvGmGNGTQIYVIDPEPCPD
 SDQEPKSSDKTHTSPSPAPPELLGGSSVFLFPPKPKD~~TL~~MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYCKVSNKAL
 PAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK~~TT~~PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP
GK

CTLA-4Fc M13
 MHVAQPAVVLASSRGIASFVCEYASPGKyTEVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDSICTGTSSGNQVNLTIQGLRAMDTGLYICKVELMYPPPYfvGmGNGTQIYVI
 DPEPCDSDQEPKSSDKTHTSPSPAPPELLGGSSVFLFPPKPKD~~TL~~MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKC
 KVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK~~TT~~PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQ
KLSLSLSPGK

CTLA-4Fc M14
 iqvtqPsVVLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAATftekntvgFLDSDSICTGTSSGNQVNLTIQGLRAMDTGLYICKVELMYPPPYfvGmGNGTQIYVIDPEPCPD
 DSDQEPKSSDKTHTSPSPAPPELLGGSSVFLFPPKPKD~~TL~~MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYCKVSNKA
 LPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK~~TT~~PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSP
GK

CTLA-4Fc M15 (belatacept)
 MHVAQPAVVLASSRGIASFVCEYASPGKyTEVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDSICTGTSSGNQVNLTIQGLRAMDTGLYICKVELMYPPPYfeGIGNGTQIYVID
 PEP~~CP~~DSDQEPKSSDKTHTSPSPAPPELLGGSSVFLFPPKPKD~~TL~~MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYCK
 VSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK~~TT~~PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQK
SLSLSPGK

CTLA-4Fc M16
 iqvtqPsVVLASSRGIASFVCEYASPGKATEVRVTVLRQADSQVTEVCAATftekntvgFLDSDSICTGTfnesrVNLTIQGLRAMDTGLYICKVELMYPPPYfvGmGNGTQIYVIDPEPCPD
 SDQEPKSSDKTHTSPSPAPPELLGGSSVFLFPPKPKD~~TL~~ytRePEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYCKVSNKALP
 APIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK~~TT~~PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQKSLSLSPG
K

CTLA-4Fc M17
 MHVAQPAVVLASSRGIASFVCEYASPGKyTEVRVTVLRQADSQVTEVCAATYMMGNELTFLDSDSICTGTSSGNQVNLTIQGLRAMDTGLYICKVELMYPPPYfeGmGNGTQIYVI
 DPEPCDSDQEPKSSDKTHTSPSPAPPELLGGSSVFLFPPKPKD~~TL~~MISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVSVLTVLHQDWLNGKEYKC
 KVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYK~~TT~~PPVLDSDGSFFLYSKLTVDKSRWQQGNVFSCSVMHEALHNHYTQ
KLSLSLSPGK

FIG. 19

FIG. 20A

Binding ELISA-5.3.16
(coat 1 µg/ml hCTLA4 mutain protein+ Biotin-hB7.1)




FIG. 20B

Binding ELISA-8.25.16
(coat 1 µg/ml hCTLA4 mutain protein+ Biotin-hB7.1)




FIG. 21A

Binding ELISA-12.2.16
(coat 1 µg/ml hCTLA4 mutain protein+ Biotin-hB7.1)




FIG. 21B

Binding ELISA-12.2.16
(coat 1 µg/ml hCTLA4 mutain protein+ Biotin-hB7.2)




FIG. 22A




FIG. 22B

Binding ELISA
(coat 1 $\mu\text{g/ml}$ hCTLA4 mutain protein + biotin-Treme




FIG. 23B






FIG. 23A



mAb4 concentration (µg/ml)

FIG. 24A



mAb5 concentration (µg/ml)

FIG. 24B




FIG. 25A

Binding ELISA – 12.3.16

(coat 1 µg/ml hCTLA4 mutain protein + biotin-mAb4)





FIG. 25B

Binding ELISA – 12.3.16

(coat 1 µg/ml hCTLA4 mutain protein + biotin-mAb5)







FIG. 26E





FIG. 26F



E006-E042-CTLA4^{h/h} KI-female pups




FIG. 27

FIG. 28A




FIG. 28B




FIG. 29A

E072-irAEs-NSG-1.27.2017




FIG. 29B




FIG. 30A




FIG. 30B




FIG. 30C




FIG. 30D




FIG. 30E





FIG. 30F






FIG. 31

FIG. 32B




FIG. 32D




FIG. 32A





FIG. 32C



CD80 AND CD86 BINDING PROTEIN COMPOSITIONS AND USES THEREOF

BACKGROUND OF THE INVENTION

Treatment with anti-CTLA-4 antibodies has been shown to be a powerful tool for enhancing anti-tumor immunity in preclinical models (10). Monotherapy with an antibody against CTLA-4 promoted rejection of transplantable tumors of various origins. Based on promising preclinical tumor model studies, the clinical potential of antibodies against CTLA-4 has been explored in different human malignancies. Although anti-CTLA-4 (Ipilimumab, marketed as Yervoy, disclosed in U.S. Pat. No. 6,984,720) has demonstrated efficacy in treating melanoma, treatment and targeting of CTLA-4 is associated with autoimmune-like toxicities. In addition, anti-CTLA-4 mAbs such as Ipilimumab and Tremelimumab are used in combination therapy with anti-PD-1/PD-L1 antibodies with superior therapeutic effect. However, the improved therapeutic effect is associated even higher rates of grade 3 and grade 4 organ toxicity. Characteristic side effects from inhibition of CTLA-4 are generally called immune-related adverse events (irAEs) and the most common irAEs are skin rash, hepatitis, colitis and endocrinopathies, particularly hypopituitarism. Therefore, there is a large unmet medical need to treat irAE while preserving the cancer therapeutic effect of anti-CTLA-4 monoclonal antibody (mAb).

The inventors have demonstrated that both clinically proven therapeutic anti-human CTLA-4 mAb and two anti-mouse Ctl-4 mAbs induce tumor rejection without blocking B7-CTLA-4 interactions under physiologically relevant conditions. Therefore, such blockade was not necessary for tumor rejection even for the mAb that can potently block B7-CTLA-4 interactions. These data refute the hypothesis that anti-CTLA-4 mAb confers an immunotherapeutic effect through checkpoint blockade (108). In support of this notion, it has further been shown that the immunotherapeutic effect mediated by Ipilimumab and possibly other anti-CTLA-4 mAbs, was unaffected by blocking B7-1 and B7-2, which is critical for pathogenesis of autoimmune diseases.

Accumulating data demonstrated that the human CTLA-4 gene encodes two different isoforms of proteins through alternative splicing: one with a trans-membrane domain which is thus likely to be anchored in membrane, and another that lacks the trans-membrane domain and is predicted to be secreted (sCTLA-4) (128). Importantly, genetic studies demonstrated that a polymorphism of CTLA-4 that reduces the relative abundance of the soluble isoform strongly associates with multiple autoimmune diseases (64). The fact that subjects with autoimmune prone alleles express less sCTLA-4 mRNA suggests that sCTLA-4 may be protective. This is notion is supported by the broad therapeutic effect of abatacept (129,130), which is a form of soluble CTLA-4, and by a genetic study in which the selective ablation of the sCTLA-4 isoform accelerated the development of type I diabetes in the mice (131). Based on these genetic data, the inventors had the insight that an anti-CTLA-4 antibody that shows the poorest binding to soluble CTLA-4 should give the least irAE. Indeed, based on the impact of the antibodies on the body weight gain in mice treated with anti-CTLA-4 antibodies during the perinatal period, a strong correlation was found among four anti-CTLA-4 mAbs: Ipilimumab has the strongest binding for sCTLA-4 and is the most toxic anti-CTLA-4, whereas antibody L3D10 had the weaker binding to sCTLA-4 and was the least toxic. Furthermore, humanized L3D10 variants

that preferentially reduced binding to sCTLA-4 showed further improved the safety profile over the parent antibody.

The protective function of soluble CTLA-4 molecules further support an approach of using soluble CTLA-4 fusion proteins to mitigate, reduce or treat irAE. However, since patients with anti-CTLA-4 mAb induced irAE have circulating anti-CTLA-4 mAbs, CTLA-4 fusion proteins such as abatacept will not only reduce the therapeutic effect of anti-CTLA-4 mAbs by preventing them from binding to cell-associated CTLA-4 molecules, but also be rendered ineffective because they will be cleared from circulation after forming immune-complex with circulating anti-CTLA-4 mAbs. Accordingly, there is a need in the art for improved CTLA-4 immunotherapy compositions and methods.

SUMMARY OF THE INVENTION

This invention relates to human CTLA-4 proteins and anti-B7-1 and anti-B7-2 compositions, and their use for immunotherapy and the treatment of autoimmune disease and inflammation, and for the reduction of autoimmune side effects associated with anti-CTLA-4 immunotherapy. Specifically, the invention relates to a CTLA-4 protein that exhibits reduced or eliminated binding to anti-CTLA-4 antibodies used in cancer immunotherapy but retains the ability to bind to B7-1 and B7-2.

The CTLA-4 protein may comprise a CTLA-4 protein in which the amino acids of an anti-CTLA-4 antibody binding epitope are modified or mutated in order to reduce or abolish binding of an anti-CTLA-4 antibody to the CTLA-4 protein, as compared to a wild-type extracellular domain of CTLA-4. The CTLA-4 protein may have anti-cancer immunotherapeutic activity. The anti-CTLA-4 antibody may be Ipilimumab. The CTLA-4 protein may retain its ability to bind at least one of B7-1 and B7-2. The CTLA-4 protein may not block the anti-cancer immunotherapeutic effects of the anti-CTLA-4 antibody.

The CTLA-4 protein may comprise an extracellular domain of mature human CTLA-4, a variant thereof, or an active fragment thereof. The CTLA-4 protein may comprise an amino acid sequence selected from the group consisting of SEQ ID NOS: 34, 36, 38, 39, 40, and 46-48, a variant thereof, and an active fragment thereof. The CTLA-4 protein may be soluble. The CTLA-4 protein may comprise a CTLA-4 fusion protein wherein the extracellular domain of human CTLA-4 is attached to a human immunoglobulin heavy chain constant region. The CTLA-4 protein may comprise an amino acid sequence selected from the group consisting of SEQ ID NO: 17, 19, 21, 22, 23, and 42-44. The CTLA-4 protein may be belatacept.

In another embodiment, the invention relates to antibody compositions that bind at least one of human B7-1 and B7-2, and their use for immunotherapy and the treatment of autoimmune disease and inflammation, and for the reduction of autoimmune side effects associated with anti-CTLA-4 immunotherapy.

Also provided herein is a pharmaceutical composition comprising a therapeutically effective amount of a protein or antibody described herein. The pharmaceutical composition may comprise a physiologically acceptable carrier or excipient.

In another aspect, provided herein are methods for reducing one or more immune functions or responses in a subject, comprising administering to a subject in need thereof the CTLA-4 protein or anti-B7 antibody composition, or a pharmaceutical composition thereof. In a specific embodi-

ment, presented herein are methods for preventing, treating, or managing a disease in which it is desirable to inhibit or reduce one or more immune functions or responses. The disease may be an autoimmune disease, such as rheumatoid arthritis (RA) or Juvenile Idiopathic Arthritis (JIA).

The compositions described herein may also be used to mitigate, minimize or treat the immune related adverse effects associated with immunotherapy. In particular, the composition may comprise a molecule that blocks or reduces the function of B7-1 and B7-2 without affecting the cancer immunotherapeutic activity of an anti-CTLA-4 antibody. The molecule may be a CTLA-4 protein, an anti-B7 antibody, or a pharmaceutical composition thereof. The molecule may be an antibody that can functionally block binding of at least one of B7-1 and B7-2 to at least one of CD28 and CTLA-4, and may be anti-B7-1 or anti-B7-2. The antibody may be capable of binding both B7-1 and B7-2, and may comprise a binding site that reacts with both B7-1 and B7-2.

A composition described herein may be administered to a subject, who may have cancer, in combination with, or on a background of, anti-CTLA-4 immunotherapy. The composition may be used prophylactically to prevent irAEs before anti-CTLA-4 treatment is initiated or the before the clinical signs of irAEs emerge. In another embodiment, the composition is used therapeutically to treat irAEs after anti-CTLA-4 treatment is initiated and the clinical symptoms are diagnosed.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Human CTLA-4 sequence. FIG. 1A shows the sequence of human CTLA-4 protein (NCBI accession number NP_005205; SEQ ID NO: 1) with the signal peptide in bold, the IgV domain underlined and the transmembrane domain double underlined. FIG. 1B shows the sequence of abatacept (SEQ ID NO: 2) with the CTLA-4 sequence shown in bold with the remainder of the protein comprising the IgG1 Fc region.

FIG. 2. Therapeutic effect of chimeric L3D10 and 10D1 in the MC38 tumor model. Human CTLA-4-knock-in mice with body weight of approximately 20 grams were used for the study. 1×10^6 MC38 tumor cells were injected subcutaneously into *Ctla-4^{h/h}* mice and when the tumor reached a size of 0.5 cm in diameter, tumor bearing mice were randomized into three groups with 5 or 6 mice each. Mice were then treated (i.p.) with 100 μ g/injection of 10D1, chimeric L3D10 or control hIgGFc on days 7, 10, 13, and 16 as indicated by the arrows. The results of duplicate experiments are shown (left and right panels) and data shown are means and S.D. of tumor size (n=6 per group in the left panel, n=5 per group in the right panel). L3D10 and 10D1 have similar therapeutic effect in this model and are both able to induce complete remission of established tumors. The diameters (d) of the tumors were calculated using the following formula: $D = \sqrt[3]{(ab)}$, $V = ab^2/2$, where a is the long diameter, while b is the short diameter. Statistical analyses were performed by two-way repeated measures ANOVA (treatment \times time). For the left panel: P=10D1 vs. hIgGFc: 5.71×10^{-7} ; L3D10 vs. hIgGFc: 5.53×10^{-7} ; 10D1 vs. L3D10: P=0.869.

FIG. 3. Adverse effects of chimeric L3D10 and 10D1 in combination with anti-PD-1. Top panel depicts the experimental design. 10-day old female-only human CTLA-4-knockin mice with body weight of greater than 4 grams were used for the study. They received indicated proteins or their combinations. Arrows indicate time of treatment (100

μ g/mice/treatment). Data shown are means and S.D. of % weight gains. Chimeric L3D10 and 10D1 have comparable cancer therapeutic effect in adult mice (FIG. 2) but distinct adverse effects are seen when 10D1 is combined with the anti-PD-1 mAb.

FIG. 4. Adverse effects of chimeric L3D10 and 10D1 in combination with anti-PD-1. The graph shows the terminal body weight on Day 42 in the mice from the experiment outlined in FIG. 3 that received either control IgG, 10D1+anti-PD-1 or chimeric L3D10+anti-PD-1 (n=5 per group). A significant reduction in weight is observed with the anti-PD1+10D1 combination, which was not seen with the anti-PD-1+Chimeric L3D10 combination.

FIG. 5. Pathological effects of chimeric L3D10 and 10D1 in combination with anti-PD-1. Group 1 is hIgG, Group 2 is 10D1+anti-PD1, and Group 3 is L3D10+anti-PD1. To further examine to relative toxicity of L3D10 compared to 10D1 when administered in combination with anti-PD-1, the gross anatomy of the mice described in FIG. 4 above was investigated. The Uterus/Ovary/Bladder and thymus were noticeably smaller in mice treated with 10D1+PD-1, whereas the organs in mice treated with L3D10+anti-PD-1 was comparable to hIgG control. In contrast, the hearts dissected from mice treated with 10D1 appeared larger in size with a noticeably whiter appearance.

FIG. 6. Treatment with 10D1 in combination with anti-PD-1 results in abnormal erythropoiesis. Given the differences in the hearts observed in FIG. 5, erythropoiesis was investigated within the mice and clear differences were observed in the mice treated with 10D1+anti-PD-1 relative to the groups treated with L3D10+anti-PD-1 or control antibody (hIgG), which were fairly similar. The bone marrow from mice treated with 10D1+anti-PD-1 had a noticeably whiter color (FIG. 6A) and the isolated blood was almost completely white in color (FIG. 6B). In accordance with this, when differentiation of the red blood cells was analyzed using distribution of CD119 and CD71 markers, a statistically significant reduction in the number of cells undergoing Stage IV development was observed in the 10D1+anti-PD-1 treated mice. Representative FACS profiles are shown in FIG. 6C, while summary data are presented in FIG. 6D.

FIG. 7. Toxicity scores of mice treated with chimeric L3D10 and 10D1 in combination with anti-PD-1. This tissue data is summarized and shows the high toxicity scores of mice treated with 10D1+anti-PD-1 relative to L3D10+anti-PD-1 which has scores only marginally higher than the hIgG control mouse group.

FIG. 8. L3D10 and 10D1 display similar binding patterns for plate immobilized CTLA-4. ELISA plates were coated with 1 μ g/ml of CTLA-4-His protein (Sino Biological, China). The given concentration of biotinylated binding proteins were added and binding measured using HRP-conjugated streptavidin. 10D1-1 and -2 are two independent material lots of the same antibody. hIgG-Fc is a human Ig negative control.

FIG. 9. L3D10 displays reduced binding soluble CTLA-4. Given concentration of anti-human CTLA-4 mAbs were coated on the plate overnight, after washing and blocking with bovine serum albumin, biotinylated CTLA-4-Fc was added at 0.25 μ g/ml. After incubation and washing, the amounts of captured CTLA-4-Fc were measured using HRP-labeled streptavidin.

FIG. 10. 10D1+anti-PD-1 do not have significant toxicity in the *Ctla-4^{h/m}* mice as evidenced by normal body weight gains in mice that received antibody treatment during the perinatal period. The mice received treatments with given

5

antibody or combinations on days 10, 13, 16, 19 and 22 intraperitoneally (100 μ g/mice/injection/antibody). Mice were weighed at least once every 3 days.

FIG. 11. Anti-tumor activity of humanized L3D10 antibodies compared to 10D1. Using the MC38 mouse tumor model in human CTLA-4 knockin mice, the anti-tumor activity of humanized L3D10 antibodies was investigated compared to the chimeric L3D10 antibody and 10D1. FIG. 11A shows the treatment schedule of the in vivo experiment. Mice were given a total of 4 doses of antibody every 3 days starting on day 7 after inoculation. All humanized antibodies (n=6 per group) completely eradicated the tumors and were comparable to 10D1 (FIG. 11B).

FIG. 12. Comparison among 10D1, mAb4 and mAb5 females for their combined toxicity with anti-PD-1 mAb. Female *Ctla-4^{h/h}* mice were treated on days 10 or 11 days after birth with four injections of antibodies (100 μ g/mice/injection, once every three days) or control Fc as specified in the legends. Mice were weighted once every 3 days. Data shown are means and SEM of % weight gain over a 30 day period. All mice were sacrificed on day 43 for histological analysis. The number of mice used per group is shown in the parentheses of labels.

The p-values for comparisons between various treatments are as follows.

hIg vs. α PD1 + L3D10	P value = 0.16
hIg vs. α PD1 + hIg	P value = 0.0384*
hIg vs. α PD1 + 10D1	P value = $<2e-16^{***}$
hIg vs. α PD1 + mAb4	P value = 0.16
hIg vs. α PD1 + mAb5	P value = 0.00207**
α PD1 + L3D10 vs. α PD1 + hIg	P value = 0.00654**
α PD1 + L3D10 vs. α PD1 + 10D1	P value = $<2e-16^{***}$
α PD1 + L3D10 vs. α PD1 + mAb4	P value = 0.492
α PD1 + L3D10 vs. α PD1 + mAb5	P value = 0.000124***
α PD1 + hIg vs. α PD1 + 10D1	P value = $<2e-16^{***}$
α PD1 + hIg vs. α PD1 + mAb4	P value = 0.0579
α PD1 + hIg vs. α PD1 + mAb5	P value = 0.409
α PD1 + 10D1 vs. α PD1 + mAb4	P value = $<2e-16^{***}$
α PD1 + 10D1 vs. α PD1 + mAb5	P value = $<2e-16^{***}$
α PD1 + mAb4 vs. α PD1 + mAb5	P value = 0.000446***

FIG. 13. Humanization of L3D10 does not affect binding to immobilized CTLA-4. The capacity of the humanized L3D10 antibodies to bind immobilized CTLA-4 was determined as described in FIG. 23. X-axis indicates the concentration of anti-CTLA-4 mAbs added into solution. Humanization does not affect binding to immobilized CTLA-4 and all three humanized antibodies demonstrated similar binding to the parental chimeric L3D10 antibody and 10D1. Similar patterns were observed when CTLA-4-Ig was used instead of CTLA-4-his.

FIG. 14. Humanization further reduces L3D10 binding to soluble CTLA-4. The capacity of the humanized L3D10 antibodies to bind soluble CTLA-4 was determined as described in FIG. 23. X-axis indicates the concentration of anti-CTLA-4 mAbs coated onto ELISA plates. Humanization further reduces binding to soluble CTLA-4 relative to the parental L3D10 chimeric antibody. Similar patterns were observed when CTLA-4-Ig was used instead of CTLA-4-his.

FIG. 15. Alignment of the human, macaque and mouse CTLA-4 extracellular domains. The amino acid sequences of the human (Hm; amino acids 1-124 of SEQ ID NO: 2), macaque (Mk) and mouse (Ms) CTLA-4 protein extracellular domains are aligned and the conserved amino acids (relative to the human sequence) are shown with dashes (-). In order to help the alignment, the mouse sequence has a

6

deletion and insertion (relative to the human and monkey sequences) at the positions highlighted. The known B7-1Ig binding site is shown in bold and underlined. The sequences demonstrate that the human and monkey sequences are highly conserved, whereas the mouse sequence has a number of amino acid differences. Based on this sequence alignment, 11 mutant (M1-M11) human CTLA-4Fc proteins were designed that incorporate murine specific amino acids—the amino acids incorporated into each mutant protein are shown in blue.

FIGS. 16A and B. Amino acid sequence composition of the WT (SEQ ID NO: 2) and mutant CTLA-4Fc proteins (SEQ ID NOs: 7-17). DNA constructs encoding the WT CTLA-4Fc protein and 11 mutant proteins, M1-M11, incorporating murine *Ctla-4* amino acids were designed as shown. The amino acid sequences are for mature proteins, including the IgG1 Fc portion, but not the signal peptide. The known B7-1Ig binding site is shown in large letters and double-underlined. The replaced murine amino acid residues in the mutant are shown lower case. The IgG1 Fc portion of the proteins in underlined.

FIGS. 17A-D. Mutation in M11 (AA103-106, YLGI>fcGm) selectively abolish antibody binding to human CTLA-4. Data shown are means of duplicates, depicting the binding of B7-1Fc (FIG. 17A), L3D10 (FIG. 17B), mAb4 (FIG. 17C), and mAb5 (FIG. 17D) binding to plate-coated hCTLA-4-Fc (open circles), mCTLA-4-Fc (filled triangles), M11 (filled circles) and IgG1-Fc (open triangles).

FIG. 18. Mapping L3D10, mAb4 and mAb5 to an epitope adjacent to the B7-1 binding site in a 3-D structure of the B7-1-CTLA-4 complex. The B7-1 binding motif and antibody epitope are shaded differently. B7-1 is depicted above CTLA-4 with a space-filled ribbon, while that of CTLA-4 is depicted as an unfilled ribbon.

FIG. 19. Amino acid sequence composition of the mutant CTLA-4Fc proteins, M12-M17 (SEQ ID NOs: 18-23). DNA constructs encoding the 6 mutant CTLA-4Fc proteins, M12-M17, incorporating murine *Ctla-4* amino acids were designed as shown. The amino acid sequences are for mature proteins, including the IgG1 Fc portion, but not the signal peptide. The known B7-1Ig binding site is shown in large letters and double-underlined. The replaced murine amino acid residues in the mutant are shown lower case. The IgG1 Fc portion of the proteins is underlined.

FIGS. 20A and B. Identifying CTLA-4Fc mutant proteins that retain binding to biotinylated B7-1Fc. CTLA4-Fc fusion proteins M1-M17 were coated onto 96-well plates. Biotinylated B7-1Fc at the given concentration was incubated with the plates and detected with streptavidin-conjugated HRP. Data shown are means of duplicate experiments.

FIGS. 21A and B. Identifying CTLA-4Fc mutant proteins that retain binding to biotinylated B7-1 and B7-2 proteins. Mutant CTLA4-Fc fusion proteins (1 μ g/ml) were coated onto 96-well plates. Biotinylated B7-1Fc (FIG. 21A) and B7-2Fc (FIG. 21B) were incubated with the plates at the given concentrations and detected with streptavidin-conjugated HRP. Data shown are means of duplicate experiments.

FIGS. 22A and B. Identifying CTLA-4Fc mutant proteins that have lost binding to biotinylated Ipilimumab (FIG. 22A) and Tremelimumab (FIG. 22B). Mutant CTLA4-Fc fusion proteins (1 μ g/ml) were coated onto 96-well plates. Biotinylated antibody mAb2 was incubated with the plates at the given concentrations and detected with streptavidin-conjugated HRP. Data shown are means of duplicate experiments.

FIGS. 23A and B. Identifying CTLA-4Fc mutant proteins that have lost binding to biotinylated mAb4. CTLA4-Fc fusion proteins M1-M17 were coated onto 96-well plates.

Biotinylated antibody mAb4 was incubated with the plates at the given concentrations and detected with streptavidin-conjugated HRP. Data shown are means of duplicate experiments.

FIGS. 24A and B. Identifying CTLA-4Fc mutant proteins that have lost binding to biotinylated mAb5. CTLA4-Fc fusion proteins M1-M17 were coated onto 96-well plates. Biotinylated antibody mAb5 was incubated with the plates at the given concentrations and detected with streptavidin-conjugated HRP. Data shown are means of duplicate experiments.

FIGS. 25A and B. Identifying CTLA-4Fc mutant proteins that have lost binding to biotinylated mAb4 (FIG. 25A) and mAb5 (FIG. 25B). Mutant CTLA4-Fc fusion proteins (1 $\mu\text{g}/\text{ml}$) were coated onto 96-well plates. Biotinylated antibody mAb4 (FIG. 25A) or mAb5 (FIG. 25B) was incubated with the plates at the given concentrations and detected with streptavidin-conjugated HRP. Data shown are means of duplicate experiments.

FIGS. 26A-F. The therapeutic effect of Ipilimumab is not achieved by blocking B7-1 and/or B7-2. FIG. 26A. Confirmation of the blocking activities of anti-B7 mAbs. CHO cells expressing mouse B7-1 or B7-2 were incubated with a mixture of 20 mg/ml anti-B7-1 (upper panel) or anti-B7-2 (lower panel) and biotinylated human CTLA-4Fc (2 mg/ml) for 1 hour. After washing away unbound proteins, the cell surface CTLA-4Fc was detected by PE-conjugated streptavidin and measured by flow cytometry. Data shown are representative FACS profiles and have been repeated twice. FIG. 26B. Diagram of the experimental design: MC38 tumor-bearing *Ctla-4^{h/m}* mice received anti-B7-1 and anti-B7-2 antibodies (300 $\mu\text{g}/\text{mouse}/\text{injection}$, once every 3 days for a total of 3 injections) in combination with either control Ig or Ipilimumab. Mice that received Ipilimumab without anti-B7-1 and anti-B7-2 were used as a positive control for tumor rejection and hIg was used as a negative control. FIGS. 26C and D. Saturation of B7-1 and B7-2 by antibody treatments as diagrammed in FIG. 26B. The PBL from mice treated as shown in FIG. 26B were stained with fluoro-chrome-conjugated anti-B7-1 and anti-B7-2 mAbs at 24 hours after anti-B7 treatment. PBL from *Cd80^{-/-}Cd86^{-/-}* mice were used as negative control. FIG. 26E. Functional blockade of B7 by anti-B7-1 and anti-B7-2 mAbs based on ablation of antibody responses. Sera were collected at day 22 after tumor challenge to evaluate anti-human IgG antibody responses. FIG. 26F. Saturating blocking of B7-1 and B7-2 by anti-B7-1 and anti-B7-2 mAbs does not affect the immunotherapeutic effect of Ipilimumab. Data shown are tumor volumes over time and have been repeated twice with similar results.

FIG. 27. CTLA-4-Fc fusion proteins M11 and M15 protect mice against irAE caused by combination therapy with anti-PD-1 and anti-CTLA-4 (Ipilimumab).

FIGS. 28A and B. CTLA4-Fc mutants M15 and M17 do not interfere with the immunotherapeutic effect of Ipilimumab. The *Ctla-4^{h/m}* heterozygous mice were transplanted with MC38 tumors and treated with Ipilimumab in combination with either control IgG Fc or CTLA-4 mutant proteins at time points indicated by the arrows. FIG. 28A. Reduction in tumor sizes. FIG. 28B. Survival of tumor-bearing mice, using tumor sizes of reached 2 cm in diameter as early removal criteria.

FIGS. 29A and B. Protective function of M15 and M17-2 against Ipilimumab-induced weight-loss (FIG. 29A) and potential liver toxicity of M15 (FIG. 29B) in NSG mice reconstituted with human CD34⁺ hematopoietic stem cells. 16 female humanized mice were randomized into 4 groups

based on the weight human leukocyte reconstitution and % of human T cells in the blood. The 4 groups were treated, respectively, with either control human IgG Fc (400 $\mu\text{g}/\text{injection}$), Ipilimumab (100 $\mu\text{g}/\text{injection}$)+human IgG Fc (300 $\mu\text{g}/\text{injection}$), Ipilimumab (100 $\mu\text{g}/\text{injection}$)+M15 (300 $\mu\text{g}/\text{injection}$), or Ipilimumab (100 $\mu\text{g}/\text{injection}$)+M17-2 (300 $\mu\text{g}/\text{injection}$) on days 0, 3 and 6. On day 18, the mice received one more injection in which the dose of Ipilimumab were 200 $\mu\text{g}/\text{mice}$ +human IgG Fc (300 $\mu\text{g}/\text{injection}$), Ipilimumab (100 $\mu\text{g}/\text{injection}$)+M15 (300 $\mu\text{g}/\text{injection}$), or Ipilimumab (100 $\mu\text{g}/\text{injection}$)+M17-2 (300 $\mu\text{g}/\text{injection}$). Mice were monitored over 30 days for weight changes, using day 0 as 100%. FIG. 29B. Elevation of ALT was observed in mice that received Ipilimumab along with M15 but not M17-2.

FIGS. 30A-F. Differential effect of M15 and M17-2 in Ipilimumab-induced T cell activation based on the analysis of spleen T cells on day 31. Mice were treated as described in FIG. 29 and sacrificed on day 31 for flow cytometry. CD4 (FIGS. 30A-C) and CD8 (FIGS. 30D-F) frequency of CD4 and CD8 T cells (FIGS. 30A and D), central memory (CD44^{hi}CD62L^{hi}) (FIGS. 30B and E) and effector memory cells (CD44^{hi}CD62L^{lo}) (FIGS. 30C and F) cells.

FIG. 31. Differential effect of M15 and M17 on the expansion of NKT cells. As detailed in FIG. 30, except that NK T cells (CD3⁺CD56⁺) were analyzed. Mice were treated as described in FIG. 29 and sacrificed on day 31 for flow cytometry.

FIGS. 32A-D. Differential effect of M15 and M17-2 on the frequency of Treg (FIGS. 32A and B) and their expression of CTLA-4 protein (FIGS. 32C and D). Mice were treated as described in FIG. 29 and sacrificed on day 31 for flow cytometry.

DETAILED DESCRIPTION

The inventors have discovered soluble CTLA-4 fusion proteins containing mutations that prevent their binding by most anti-CTLA-4 mAbs while retaining their ability to bind B7-1 and B7-2 on antigen presenting cells (APC). Surprisingly, these fusion proteins prevent the autoimmune adverse effects of anti-CTLA-4 mAbs without affecting the cancer therapeutic effects of anti-CTLA-4.

1. Definitions

The terminology used herein is for the purpose of describing particular embodiments only and is not intended to be limiting. As used in the specification and the appended claims, the singular forms “a,” “an” and “the” include plural referents unless the context clearly dictates otherwise.

For recitation of numeric ranges herein, each intervening number there between with the same degree of precision is explicitly contemplated. For example, for the range of 6-9, the numbers 7 and 8 are contemplated in addition to 6 and 9, and for the range 6.0-7.0, the numbers 6.0, 6.1, 6.2, 6.3, 6.4, 6.5, 6.6, 6.7, 6.8, 6.9 and 7.0 are explicitly contemplated.

As used herein, the term “protein”, “peptide” or “polypeptide” refers to a chain of amino acids of any length, regardless of modification (e.g., phosphorylation or glycosylation). The amino acids may be natural, synthetic, or a modification or combination of natural and synthetic. A polypeptide of the present invention may be a recombinant polypeptide, a natural polypeptide or a synthetic polypeptide, preferably a recombinant polypeptide.

As used herein, the terms “portion,” “segment,” and “fragment,” when used in relation to polypeptides, refer to a continuous sequence of residues, such as amino acid

residues, which sequence forms a subset of a larger sequence. For example, if a polypeptide were subjected to treatment with any of the common endopeptidases, such as trypsin or chymotrypsin, the oligopeptides resulting from such treatment would represent portions, segments or frag- 5 ments of the starting polypeptide. A “fragment” of a polypeptide thus refers to any subset of the polypeptide that is a shorter polypeptide of the full length protein. Generally, fragments will be five or more amino acids in length.

As used herein, the term “soluble portion” of a protein 10 means that portion of the full length polypeptide that does not include any part of the transmembrane portion or segment. For example, with respect to CTLA-4, a soluble portion would include the extracellular portion (with or without the N-terminal signal sequence) but would not include any part of the transmembrane portion (or, at least, not enough to reduce solubility). Thus, the ECD of human CTLA-4 is shown as SEQ ID NO: 3 (i.e., amino acids 36-161 of the full length sequence (SEQ ID NO: 1), where amino acids 1-35 comprise the signal sequence and are not 15 included in the mature extracellular, and thus soluble, protein.

As used herein, the term “fusion protein” is defined as one or more amino acid sequences joined together using methods known in the art. The joined amino acid sequences thereby form one fusion protein. Fusion proteins known in the art include the hexa-histidine peptide, the hemagglutinin “HA” tag, which corresponds to an epitope derived from the influenza hemagglutinin protein (Wilson, I. A. et al. (1984) “The Structure Of An Antigenic Determinant In A Protein,” 20 *Cell*, 37:767-778) and the “flag” tag (Knappik, A. et al. (1994) “An Improved Affinity Tag Based On The FLAG Peptide For The Detection And Purification Of Recombinant Antibody Fragments,” *Biotechniques* 17(4):754-761).

A derivative, analog or homolog, of a polypeptide (or 25 fragment thereof) of the invention may be (i) one in which one or more of the amino acid residues are substituted with a conserved or non-conserved amino acid residue (preferably a conserved amino acid residue) and such substituted amino acid residue may or may not be one encoded by the genetic code, or (ii) one in which one or more of the amino acid residues includes a substituent group, or (iii) one in which the mature polypeptide is fused with another compound, such as a compound to increase the half-life of the polypeptide (for example, polyethylene glycol), or (iv) one 30 in which the additional amino acids are fused to the mature polypeptide, such as a leader or secretory sequence or a sequence which is employed for purification of the mature polypeptide or a proprotein sequence. Such derivatives and analogs are deemed to be within the scope of those skilled 35 in the art from the teachings herein.

As used herein, the term “antibody” is intended to denote an immunoglobulin molecule that possesses a “variable region” antigen recognition site. The term “variable region” is intended to distinguish such domain of the immunoglobulin from domains that are broadly shared by antibodies (such as an antibody Fc domain). The variable region comprises a “hypervariable region” whose residues are responsible for antigen binding. The hypervariable region comprises amino acid residues from a “Complementarity Determining Region” or “CDR” (i.e., typically at approximately residues 24-34 (L1), 50-56 (L2) and 89-97 (L3) in the light chain variable domain and at approximately residues 27-35 (H1), 50-65 (H2) and 95-102 (H3) in the heavy chain variable domain; ref. 44) and/or those residues from a “hypervariable 40 loop” (i.e., residues 26-32 (L1), 50-52 (L2) and 91-96 (L3) in the light chain variable domain and 26-32 (H1), 53-55

(H2) and 96-101 (H3) in the heavy chain variable domain; Ref. 45). “Framework Region” or “FR” residues are those variable domain residues other than the hypervariable region residues as herein defined. The term antibody includes 5 monoclonal antibodies, multi-specific antibodies, human antibodies, humanized antibodies, synthetic antibodies, chimeric antibodies, camelized antibodies, single chain antibodies, disulfide-linked Fvs (sdFv), intrabodies, and anti-idiotypic (anti-Id) antibodies (including, e.g., anti-Id and anti-anti-Id antibodies to antibodies of the invention). In particular, such antibodies include immunoglobulin molecules of any type (e.g., IgG, IgE, IgM, IgD, IgA and IgY), class (e.g., IgG₁, IgG₂, IgG₃, IgG₄, IgA₁ and IgA₂) or subclass.

As used herein, the term “active fragment” refers to a portion of a natural polypeptide or antibody, or a polypeptide with high sequence homology (for example, at least 80%, 85%, 90%, 95%, 98%, or 99% amino acid sequence identity) to a natural polypeptide or antibody and which retains 15 biological activity. Representative examples of “biological activity” include the ability to bind B7 proteins and to bind their natural receptors or to be bound by a specific antibody. For example, an active fragment of CTLA-4 would be capable of binding B7.1 or B7.2 or by binding to a ligand of CTLA-4. In preferred embodiments, such a fragment would consist of the extracellular domain (ECD) of a CTLA-4 20 protein.

As used herein the term “isolated” is meant to describe a compound of interest (e.g., either a polynucleotide or a polypeptide) that is in an environment different from that in which the compound naturally occurs e.g. separated from its natural milieu such as by concentrating a peptide to a concentration at which it is not found in nature. “Isolated” is meant to include compounds that are within samples that 25 are substantially enriched for the compound of interest and/or in which the compound of interest is partially or substantially purified.

“Substantially identical” may mean that a first and second amino acid sequence are at least 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% over a region of 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100, 110, 120, 130, 140, 150, 160, 170, 180, 190, 200, 210, 220, 230, 240, 250, 260, 270, 280, 290, or 300 amino acids.

“Treatment” or “treating,” when referring to protection of an animal from a disease, means preventing, suppressing, repressing, or completely eliminating the disease. Preventing the disease involves administering a composition of the present invention to an animal prior to onset of the disease. Suppressing the disease involves administering a composition of the present invention to an animal after induction of the disease but before its clinical appearance. Repressing the disease involves administering a composition of the present invention to an animal after clinical appearance of the disease. 50

As used herein, a “variant” polypeptide contains at least one amino acid sequence alteration as compared to the amino acid sequence of the corresponding wild-type polypeptide. As used herein, an “amino acid sequence alteration” can be, for example, a substitution, which may be conservative, a deletion, or an insertion of one or more amino acids. Variant may also mean a protein with an amino acid sequence that is substantially identical to a referenced protein with an amino acid sequence, and may retain at least one biological activity. A variant may be a derivative, analog or 65

homolog, of a polypeptide. A variant may also be a soluble portion of a polypeptide. A conservative substitution of an amino acid, i.e., replacing an amino acid with a different amino acid of similar properties (e.g., hydrophilicity, degree and distribution of charged regions) is recognized in the art as typically involving a minor change. These minor changes can be identified, in part, by considering the hydropathic index of amino acids, as understood in the art. Kyte et al., *J. Mol. Biol.* 157:105-132 (1982). The hydropathic index of an amino acid is based on a consideration of its hydrophobicity and charge. It is known in the art that amino acids of similar hydropathic indexes can be substituted and still retain protein function. In one aspect, amino acids having hydropathic indexes of ± 2 are substituted. The hydrophilicity of amino acids can also be used to reveal substitutions that would result in proteins retaining biological function. A consideration of the hydrophilicity of amino acids in the context of a peptide permits calculation of the greatest local average hydrophilicity of that peptide, a useful measure that has been reported to correlate well with antigenicity and immunogenicity. U.S. Pat. No. 4,554,101, incorporated fully herein by reference. Substitution of amino acids having similar hydrophilicity values can result in peptides retaining biological activity, for example immunogenicity, as is understood in the art. Substitutions may be performed with amino acids having hydrophilicity values within ± 2 of each other. Both the hydrophobicity index and the hydrophilicity value of amino acids are influenced by the particular side chain of that amino acid. Consistent with that observation, amino acid substitutions that are compatible with biological function are understood to depend on the relative similarity of the amino acids, and particularly the side chains of those amino acids, as revealed by the hydrophobicity, hydrophilicity, charge, size, and other properties.

2. B7-CTLA-4 Interactions and Immunotherapy

The ability of T cells to optimally mediate an immune response against an antigen requires two distinct signaling interactions. First, antigen that has been arrayed on the surface of antigen-presenting cells (APC) must be presented to antigen-specific naive T cells in the form of MHC: peptide complex (1, 2). Such presentation delivers a signal via the T cell receptor (TCR) that directs the T cell to initiate an immune response that will be specific to the presented antigen. Second, a series of co-stimulatory signals, mediated through interactions between the APC and distinct T cell surface molecules, triggers first the activation and proliferation of the T cells and ultimately their inhibition (3-5). Thus, the first signal confers specificity to the immune response whereas the second signal serves to determine the nature, magnitude and duration of the response while limiting immunity to self. Of particular importance among these second signal molecules is binding between the B7.1 (CD80) (6) and B7.2 (CD86) (7-9) ligands of the APC and the CD28 and CTLA-4 receptors (10-12) of the T-lymphocyte.

CD28 (Cluster of Differentiation 28) and Cytotoxic T-lymphocyte-associated antigen-4 (CTLA-4) are members of the immunoglobulin superfamily of proteins and are involved in the regulation of T-cell activation. CD28 is constitutively expressed on naive T cells and binds B7.1 and B7.2 providing a co-stimulatory signal required for T cell activation and survival. In contrast, upon T cell activation, CTLA-4 is upregulated on T cells and competes with CD28 for binding to B7.1 and B7.2, thereby transmitting a suppressive signal for T cell activation.

B7 family molecules have a membrane proximal IgC (constant) domain and a membrane distal IgV (variable)

domain. The CD28-like family of receptors for these ligands share a common extracellular IgV-like domain. Interactions of receptor-ligand pairs are mediated predominantly through residues in the IgV domains of the ligands and receptors (Schwartz, et al., *Nature Immunol.*, 3:427-434 (2002)). Crystallographic analysis revealed that the CTLA-4/B7 binding interface is dominated by the interaction of the CDR3-analogous loop from CTLA-4, composed of a MYPPPY motif (Schwartz, et al., *Nature*, 410:604-608 (2001); and Stamper, et al., *Nature*, 410:608-611 (2001)).

Antibodies against CTLA-4 have been shown to block the interaction between CTLA-4 and the costimulatory molecules B7.1 and B7.2 in vitro. This blockade removes the CTLA-4-mediated inhibitory signal on T-cells and thereby stimulates a natural immune response that can be used as a therapy against cancer. Treatment with anti-CTLA-4 antibodies has been shown to be a powerful tool for enhancing anti-tumor immunity in preclinical models (10). Monotherapy with an antibody against CTLA-4 promoted rejection of transplantable tumors of various origins. Based on promising preclinical tumor model studies, the clinical potential of antibodies against CTLA-4 has been explored in different human malignancies. Although anti-CTLA-4 (Ipilimumab, marketed as Yervoy, disclosed in U.S. Pat. No. 6,984,720) has demonstrated efficacy in treating melanoma, treatment and targeting of CTLA-4 is associated with autoimmune like toxicities. Characteristic side effects from inhibition of CTLA-4 are generally called immune-related adverse events (irAEs) and the most common irAEs are skin rash, hepatitis, colitis and endocrinopathies, particularly hypopituitarism. Therefore, there is a desire to improve the safety of anti-CTLA-4 antibodies by reducing the associated irAEs.

The inventors have surprisingly discovered that both clinically proven therapeutic anti-human CTLA-4 mAb and two anti-mouse Ctl4 mAbs induce tumor rejection without blocking B7-CTLA-4 interactions under physiologically relevant conditions. Therefore, blocking CTLA-4 interactions with B7.1 or B7.2 is not necessary for tumor rejection even for the mAb that can potently block these B7-CTLA-4 interactions. These data refute the hypothesis that anti-CTLA-4 mAb confers immunotherapeutic effect through checkpoint blockade (108). Furthermore, the inventors have identified an anti-CTLA-4 antibody, L3D10, with reduced immune related toxicities demonstrating that cancer immunity and irAEs can be uncoupled genetically.

3. Soluble CTLA-4

Accumulating data demonstrated that the human CTLA-4 gene encodes two different isoforms of proteins through alternative splicing: one with a trans-membrane domain which is thus likely to be anchored in the membrane, and another that lacks the trans-membrane domain and is predicted to be secreted (sCTLA-4, SEQ ID NO: 4) (128). Importantly, genetic studies demonstrated that a polymorphism of CTLA-4 that reduces the relative abundance of the soluble isoform strongly associates with multiple autoimmune diseases (64). The fact that subjects with autoimmune prone alleles express less sCTLA-4 mRNA suggests that sCTLA-4 may be protective. This notion is supported by the broad therapeutic effect of abatacept (129,130), which is a form of soluble CTLA-4, and by the genetic study in which the selective ablation of the sCTLA-4 isoform accelerated the development of type I diabetes in the mice (131).

Based on these genetic data it the inventors had the insight that the antibody that shows the poorest binding to sCTLA-4 should be associated with the fewest irAE. Indeed, the inventors identified an anti-CTLA-4 antibody, L3D10, with

reduced immune related toxicities that was shown to have reduced binding to sCTLA-4 relative to membrane bound or immobilized CTLA-4. Based on the impact of the antibodies on the body weight gain in mice treated with anti-CTLA-4 antibodies during the perinatal period, a strong correlation was found among four anti-CTLA-4 mAbs: Ipilimumab has the strongest binding for sCTLA-4 and is the most toxic anti-CTLA-4, whereas L3D10 had the weakest binding to sCTLA-4 and was the least toxic. Furthermore, humanization of the L3D10 antibody preferentially reduced binding to sCTLA-4 and appeared to further improve the safety profile over the parent antibody.

In order to map the CTLA-4 binding epitope of the L3D10 parent antibody and humanized variants, the inventors took advantage of the fact that the mouse and human CTLA-4 proteins are cross-reactive between species to B7-1, but not to the anti-CTLA-4 antibodies. Accordingly, the inventors designed a number of mutants of the human CTLA-4Fc protein in which clusters of amino acids from the human CTLA-4 protein were replaced with amino acids from the murine CtlA-4 protein (SEQ ID NO: 5). As the anti-CTLA-4 antibodies used in this study do not bind to murine CtlA-4, binding of the anti-human CTLA-4 antibodies should be abolished when key residues of the antibody binding epitope are replaced with murine amino acids. Accordingly, the inventors have shown that the L3D10 binding site of CTLA-4 maps directly adjacent to the B7-1 binding motif, MYPPPY (SEQ ID NO: 50). This correlates well with the demonstrated ability of the antibody to block B7-CTLA-4 interactions both in vitro and in vivo. By contrast, Ipilimumab, does not block binding of B7.1 or B7.2 to abatacept under physiological conditions and so it must also bind a region that does not include the MYPPPY motif. However, as it does not show reducing binding to sCTLA-4, the binding domain must be different from antibody L3D10.

Endogenous sCTLA-4 is produced by fusion of the truncated C-terminal end of the extracellular IgV domain directly to the intracellular domain, without the intervening transmembrane domain. Accordingly, there are only 12 amino acids of the CTLA-4 extracellular domain C-terminal to the MYPPPY motif in sCTLA-4, whereas the endogenous membrane bound version has 21 amino acids. Based on this, without being bound by theory, it appears that an antibody that binds to polymorphic C-terminal domain residues, such as L3D10, is more likely to lose reactivity to sCTLA-4. Furthermore, the inventors have demonstrated that it is possible to mutate these amino acids in the CTLA-4Fc protein in this region so that they no longer bind anti-CTLA-4 antibodies.

4. Recombinant CTLA-4 Proteins

CTLA-4Fc fusion protein (abatacept, marketed as Orenia) is a selective costimulation modulator comprising the extracellular domain of CTLA-4 fused to the Fc region of human IgG1 (shown in FIG. 1B; SEQ ID NO:2) that inhibits T cell (T lymphocyte) activation by binding to B7.1 and B7.2, thereby blocking interaction with CD28. In vitro, abatacept decreases T cell proliferation and inhibits the production of the cytokines TNF alpha (TNF α), interferon- γ , and interleukin-2. Activated T lymphocytes are implicated in the pathogenesis of Rheumatoid Arthritis (RA) and are found in the synovium of patients with RA. In a rat collagen-induced arthritis model, abatacept suppresses inflammation, decreases anti-collagen antibody production, and reduces antigen specific production of interferon- γ . Based on this preclinical activity, abatacept has been developed and approved for the treatment of RA and Juvenile Idiopathic Arthritis (JIA) in humans.

Although capable of reducing immune responses, CTLA-4Fc fusion proteins comprising the endogenous CTLA-4 extracellular domain, such as abatacept, are incapable of reducing or mitigating the immune related toxicities associated with anti-CTLA-4 immunotherapies as the activity of the two molecules are offsetting. The CTLA-4 fusion protein such as the abatacept will not only reduce the therapeutic effect of anti-CTLA-4 mAbs by binding directly to the antibodies to prevent them from binding to endogenous cell-associated CTLA-4 molecules, but will also be rendered ineffective themselves as they will be cleared from circulation after forming immune-complex with circulating anti-CTLA-4 mAbs.

Given that both blocking and non-blocking anti-CTLA-4 antibodies have anti-tumor effects and antibodies demonstrating reduced irAEs are associated with reduced binding to sCTLA-4, the inventors engineered a panel of soluble CTLA-4 proteins with mutations that prevent binding by most anti-CTLA-4 mAbs that were tested while retaining their ability to bind B7-1 and B7-2 on the antigen-presenting cells, which is critical for the pathogenesis of autoimmune disease and irAEs. Such proteins, by virtue of their retained B7 binding activity, can be used in the treatment of autoimmune diseases such as RA and JIA. Furthermore, such proteins can be used in combination with anti-CTLA-4 immunotherapy treatment to reduce the associated irAEs, while leaving the anti-tumor activity of the immunotherapy intact.

a. CTLA-4 Proteins

Provided herein is a CTLA-4 protein comprising (a) the extracellular domain of mature human CTLA-4 (SEQ ID NO: 3) or mouse CTLA-4 (SEQ ID NO: 6), (b) an amino acid set forth in one of SEQ ID NOS: 24-40 and 46-49, (c) a variant of the foregoing, or (d) an active fragment of the foregoing. The CTLA-4 protein may be soluble. The CTLA-4 protein may further comprise an N-terminal signal peptide. The CTLA-4 protein may retain the ability to bind human B7.1 and B7.2 but lack the ability to bind an anti-CTLA-4 antibody. In one specific embodiment, the CTLA-4 protein does not bind to Ipilimumab or Tremelimumab under physiological conditions. In particular, the CTLA-4 protein may comprise an amino acid sequence set forth in SEQ ID NO: 34, 36, 38, 39, 40, 46, 47, or 48. The CTLA-4 protein may be isolated.

b. CTLA-4 Fusion Proteins

The CTLA-4 protein may be fused at its N- or C-terminal end to another protein. The other protein may be a portion of a mammalian Ig protein, which may be human or mouse. The portion may comprise a Fc region of the Ig protein. The Fc region may comprise the hinge region and CH2 and CH3 domains of the Ig protein. The Ig protein may be human IgG1, IgG2, IgG3, IgG4, or IgA. The Ig protein may also be IgM, and the Fc portion may comprise the hinge region and CH3 and CH4 domains of IgM. In one embodiment, the Fc region is human IgG1 comprising the amino acid sequence set forth in SEQ ID NO: 41. The CTLA-4 protein may comprise an amino acid sequence selected from the group consisting of SEQ ID NOS: 7-23 and 42-45, and particularly selected from the group consisting of SEQ ID NO: 17, 19, 21, 22, 23, 42, 43, and 44.

The CTLA-4 protein may also be fused at its N- or C-terminus to a protein tag, which may comprise GST, His, or FLAG. Methods for making fusion proteins and purifying fusion proteins are well known in the art.

5. Anti-B7 Antibodies

Another embodiment of the invention relates to an anti-B7 antibody composition that binds at least one of human

B7-1 and B7-2. The anti-B7 antibody may bind to one or both of endogenous B7-1 and B7-2, and may neutralize activity without affecting the cancer immunotherapeutic activity of anti-CTLA-4 antibodies. The anti-B7 antibody may be humanized to minimize anti-drug antibodies. The anti-B7 antibody may be a monoclonal antibody, and may be cross-reactive with both B7-1 and B7-2. In yet another embodiment, a bispecific antibody may comprise antigen-binding domains of two antibodies, respectively binding to B7-1 or B7-2

6. Production

The CTLA-4 protein or anti-B7 antibody may be prepared using a eukaryotic expression system. The expression system may entail expression from a vector in mammalian cells, such as Chinese Hamster Ovary (CHO) cells. The system may also be a viral vector, such as a replication-defective retroviral vector that may be used to infect eukaryotic cells. The CTLA-4 protein may also be produced from a stable cell line that expresses CTLA-4 protein from a vector or a portion of a vector that has been integrated into the cellular genome. The stable cell line may express CTLA-4 protein from an integrated replication-defective retroviral vector. The expression system may be GPEX™.

The CTLA-4 protein or anti-B7 antibody can be purified using, for example, chromatographic methods such as affinity chromatography, ion exchange chromatography, hydrophobic interaction chromatography, DEAE ion exchange, gel filtration, and hydroxylapatite chromatography. In some embodiments, fusion proteins can be engineered to contain an additional domain containing amino acid sequence that allows the polypeptides to be captured onto an affinity matrix. For example, a CTLA-4 protein or anti-B7 antibody comprising the Fc region of an immunoglobulin domain can be isolated from cell culture supernatant or a cytoplasmic extract using a protein A column. In addition, a tag such as c-myc, hemagglutinin, polyhistidine, or Flag™ (Kodak) can be used to aid polypeptide purification. Such tags can be inserted anywhere within the polypeptide, including at either the carboxyl or amino terminus. Other fusions that can be useful include enzymes that aid in the detection of the polypeptide, such as alkaline phosphatase. Immunoaffinity chromatography also can be used to purify polypeptides. Fusion proteins can additionally be engineered to contain a secretory signal (if there is not a secretory signal already present) that causes the fusion protein to be secreted by the cells in which it is produced. The secreted fusion proteins can then conveniently be isolated from the cell media.

7. Pharmaceutical Compositions

The invention further concerns a pharmaceutical composition comprising a therapeutically effective amount of one or more of the above-described CTLA-4 proteins and anti-B7 antibodies, and a physiologically acceptable carrier or excipient. Preferably, compositions of the invention comprise a prophylactically or therapeutically effective amount of the CTLA-4 protein or anti-B7 antibody and a pharmaceutically acceptable carrier

In a specific embodiment, the term “pharmaceutically acceptable” means approved by a regulatory agency of the Federal or a state government or listed in the U.S. Pharmacopeia or other generally recognized pharmacopeia for use in animals, and more particularly in humans. The term “carrier” refers to a diluent, adjuvant (e.g., Freund’s adjuvant (complete and incomplete), excipient, or vehicle with which the therapeutic is administered. Such pharmaceutical carriers may be sterile liquids, such as water and oils, including those of petroleum, animal, vegetable or synthetic origin, such as peanut oil, soybean oil, mineral oil, sesame

oil and the like. Water is a preferred carrier when the pharmaceutical composition is administered intravenously. Saline solutions and aqueous dextrose and glycerol solutions can also be employed as liquid carriers, particularly for injectable solutions. Suitable pharmaceutical excipients include starch, glucose, lactose, sucrose, gelatin, malt, rice, flour, chalk, silica gel, sodium stearate, glycerol monostearate, talc, sodium chloride, dried skim milk, glycerol, propylene, glycol, water, ethanol and the like. The composition, if desired, may also contain minor amounts of wetting or emulsifying agents, or pH buffering agents. These compositions may take the form of solutions, suspensions, emulsion, tablets, pills, capsules, powders, sustained-release formulations and the like.

Generally, the ingredients of compositions of the invention may be supplied either separately or mixed together in unit dosage form, for example, as a dry lyophilized powder or water free concentrate in a hermetically sealed container such as an ampoule or sachette indicating the quantity of active agent. Where the composition is to be administered by infusion, it can be dispensed with an infusion bottle containing sterile pharmaceutical grade water or saline. Where the composition is administered by injection, an ampoule of sterile water for injection or saline may be provided so that the ingredients may be mixed prior to administration.

The compositions of the invention may be formulated as neutral or salt forms. Pharmaceutically acceptable salts include, but are not limited to, those formed with anions such as those derived from hydrochloric, phosphoric, acetic, oxalic, tartaric acids, etc., and those formed with cations such as those derived from sodium, potassium, ammonium, calcium, ferric hydroxides, isopropylamine, triethylamine, 2-ethylamino ethanol, histidine, procaine, etc.

8. Methods of Treatment

a. Autoimmune Disease

The CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof may be used to treat an inflammatory or autoimmune disease. Representative inflammatory or autoimmune diseases and disorders that may be treated using B7-H4 fusion polypeptides include, but are not limited to, rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), alopecia areata, anklosing spondylitis, antiphospholipid syndrome, autoimmune Addison’s disease, autoimmune hemolytic anemia, autoimmune hepatitis, autoimmune inner ear disease, autoimmune lymphoproliferative syndrome (alps), autoimmune thrombocytopenic purpura (ATP), Behcet’s disease, bullous pemphigoid, cardiomyopathy, celiac sprue-dermatitis, chronic fatigue syndrome immune deficiency, syndrome (CFIDS), chronic inflammatory demyelinating polyneuropathy, cicatricial pemphigoid, cold agglutinin disease, Crest syndrome, Crohn’s disease, Deigo’s disease, dermatomyositis, dermatomyositis—juvenile, discoid lupus, essential mixed cryoglobulinemia, fibromyalgia—fibromyositis, grave’s disease, guillain-barre, hashimoto’s thyroiditis, idiopathic pulmonary fibrosis, idiopathic thrombocytopenia purpura (ITP), Iga nephropathy, insulin dependent diabetes (Type I), juvenile arthritis, Meniere’s disease, mixed connective tissue disease, multiple sclerosis (MS), myasthenia gravis, pemphigus vulgaris, pernicious anemia, polyarteritis nodosa, polychondritis, polyglancular syndromes, polymyalgia rheumatica, polymyositis and dermatomyositis, primary agammaglobulinemia, primary biliary cirrhosis, psoriasis, Raynaud’s phenomenon, Reiter’s syndrome, rheumatic fever, sarcoidosis, scleroderma, Sjogren’s syndrome, stiff-man syndrome, Takayasu arteritis, temporal arteritis/giant cell arteritis, ulcerative colitis, uveitis, vasculitis, vitiligo, and Wegener’s

granulomatosis. In a particular embodiment, the autoimmune disease may be rheumatoid arthritis (RA) or Juvenile Idiopathic Arthritis (JIA). The CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof may be administered to a subject in need thereof. The subject may be a mammal such as a human.

The CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof may be combined with another drug, such as a disease-modifying antirheumatic drug (DMARD). The drug may be a nonsteroid anti-inflammatory drug (NSAID), which may be a propionic acid derivative, an acetic acid derivative, an enolic acid derivative, a fenamic acid derivative, or a selective Cox2 inhibitor. The drug may also be a corticosteroid or Methotrexate. The drug may be a biologic, which may be a TNF- α antagonist such as an anti-TNF- α antibody or a fusion protein that binds to TNF- α (Enbrel), an anti-CD20 mAb, an antagonist of costimulatory molecule CD80 and CD86 such as a monoclonal antibody or a fusion protein (CTLA-4Ig) that binds to the two molecules, or an antagonist for a receptor of either IL-1 or IL-6. The CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof, and the other drug may be administered together or sequentially.

b. Immunotherapy

The CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof may also be used to mitigate, reduce or treat the immune related adverse events (irAEs) associated with anti-CTLA-4 immunotherapy in cancer patients. The CTLA-4 protein may be administered prophylactically (before the emergence of irAEs) or therapeutically (after the emergence of irAEs). In particular, the CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof may be administered to a subject in combination with, or on a background of, anti-CTLA-4 immunotherapy. The subject may be cancer patient. In one embodiment, the CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof is used prophylactically to prevent irAEs before anti-CTLA-4 treatment is initiated or the before the clinical signs of irAEs emerge. In another embodiment, the CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof is used therapeutically to treat irAEs after anti-CTLA-4 treatment is initiated and the clinical symptoms are diagnosed.

c. Methods of Administration

The CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof may be administered by a method including, but not limited to, parenteral administration (e.g., intradermal, intramuscular, intraperitoneal, intravenous and subcutaneous), epidural, and mucosal (e.g., intranasal and oral routes). In a specific embodiment, the CTLA-4 protein or anti-B7 antibody, or pharmaceutical composition thereof is administered intramuscularly, intravenously, or subcutaneously. The composition may be administered by any convenient route, for example, by infusion or bolus injection, by absorption through epithelial or mucocutaneous linings (e.g., oral mucosa, rectal and intestinal mucosa, etc.) and may be administered together with other biologically active agents. Administration can be systemic or local.

EXAMPLES

Example 1

Chimeric L3D10 Antibody has Equivalent Activity as 10D1 in Causing Tumor Rejection

In the clinic, the anti-CTLA-4 antibody, Ipilimumab (mab 10D1), has been shown to improve the survival of cancer

patients but induces significant autoimmune adverse effect. Using human CTLA-4 gene knock-in mice and hu-PBL-Scid mice, it was previously demonstrated that mouse anti-human CTLA-4 antibodies reduced tumor growth, and identified L3D10 as the most effective among the panel of mAbs tested (21).

The availability of human CTLA-4 gene knockin mice (20) provided with an unprecedented opportunity to test biological activity of the chimeric L3D10 anti-human CTLA-4 antibody, comprising the parental L3D10 variable region and human IgG1 constant domain (the Fc region), with clinically used anti-CTLA-4 mAb, 10D1. In this humanized mouse model, a CTLA-4 gene encoding a product with 100% identity to human CTLA-4 protein is expressed under the control of endogenous mouse CTLA-4 locus. When the anti-tumor activity of the chimeric L3D10 and 10D1 were directly compared in the MC38 tumor model in human CTLA-4-knockin mice, it is clear that both antibodies were comparable in causing tumor rejection, whereas the tumors grew progressively in IgG control group. FIG. 2 shows the results of antibody treatment on tumor size from duplicate experiments.

Example 2

Reduction of Autoimmune Adverse Effect by Other Immunotherapeutic Antibodies

Recent clinical studies have revealed that combination therapy between anti-PD-1 and anti-CTLA-4 mAb further increase the survival of end-stage melanoma patients. However, 55% of the patients that received the combination therapy developed grades 3 and 4 immune related adverse events (irAEs). It is therefore critical to develop antibodies with less toxicity. We have developed an in vivo model that recapitulates the irAEs associated with the combination therapy of anti-CTLA-4 and anti-PD-1 mAbs observed in the clinic. In this model we treated human CTLA-4 gene knockin mice (CTLA-4^{h/h}) during the perinatal period with high doses of anti-PD-1 and anti-CTLA-4 mAbs. We found that while the young mice tolerate treatment of individual mAbs, combination therapy with anti-PD-1 and 10D1 causes severe irAE with multiple organ inflammation, anemia and, as shown in FIG. 3, severely stunted growth. In contrast, when combined with anti-PD-1, chimeric L3D10 exhibits only mild irAE as demonstrated by normal weight gain.

To further examine to relative toxicity of chimeric L3D10 compared to 10D1 when administered in combination with anti-PD-1, we looked at the pathological effects in the CTLA-4^{h/h} knockin mice at 42 days after administration. As shown in FIG. 4, terminal body weight (day 42) in mice treated with L3D10+anti-PD-1 was similar to mice treated with hIgG negative control antibody. However, by comparison, the weight of mice treated with 10D1+anti-PD-1 was much lower. Accordingly, when we looked at the gross anatomy of these mice, the Uterus/Ovary/Bladder and thymus were noticeably smaller in mice treated with 10D1+PD-1 (FIG. 5). Again, the organs in mice treated with L3D10+anti-PD-1 was comparable to hIgG control. In contrast, the hearts dissected from mice treated with 10D1 appeared slightly larger in size with a noticeably whiter appearance. As a result we decided to look at erythropoiesis within the mice and observed clear differences in the mice treated with 10D1+anti-PD-1 relative to the groups treated with L3D10+anti-PD-1 or control antibody, which were fairly similar. As shown in FIG. 6A, the bone marrow from

mice treated with 10D1+anti-PD-1 had a noticeably whiter color and the isolated blood was almost completely white in color (FIG. 6B). In accordance with this, when we took a closer look at the cells undergoing the different stages of blood development using CD71 and CD119 markers. Representative FACS profiles are shown in FIG. 6C, while summary data are presented in FIG. 6d. These data revealed a statistically significant reduction in the number of cells undergoing Stage IV development in the 10D1+anti-PD-1 treated mice (FIG. 6D).

To further determine the toxicology of L3D10 vs 10D1 in combination with anti-PD-1, we performed histological analysis of the heart, lung, salivary gland and the kidney and liver following fixation in 10% formalin for at least 24 hours. In each of the tissues studied, mice treated with 10D1+anti-PD-1 displayed a high level of T cell infiltration. The toxicity score, based on severity of inflammation, are summarized in FIG. 7, which shows the high toxicity scores of mice treated with 10D1+anti-PD-1 relative to L3D10+anti-PD-1 which has scores only marginally higher than the hIgG control mouse group.

Example 3

L3D10 has Reduced Binding for Soluble CTLA-4

L3D10 and 10D1 display similar binding patterns for plate immobilized CTLA-4 (FIG. 8). As a possible explanation for the reduced toxicity of L3D10 relative to 10D1, particularly the increased T cell infiltration/activity associated with 10D1, we decided to look at the binding to soluble CTLA-4. We chose to look at this because the association between CTLA-4 polymorphism and multiple autoimmune diseases relates to the defective production of soluble CTLA-4 (67) and genetic silencing of the sCTLA-4 isoform increased the onset of type I diabetes in mice (131). Furthermore, soluble CTLA-4 (abatacept and belatacept) is a widely used drug for immune suppression. In accordance with this idea, when we looked at the relative binding to soluble CTLA-4, we observed a marked decrease in the binding of L3D10 relative to 10D1 (FIG. 9).

We have demonstrated that anti-CTLA-4 mAb induce robust tumor injection in heterozygous *Ctla-4^{h/m}* mice in which only 50% of CTLA-4 molecules can bind to anti-human CTLA-4 mAbs. To determine if engagement of 50% of CTLA-4 is sufficient to induce irAE, we treated the *Ctla-4^{h/m}* mice with anti-PD-1+10D1. As shown in FIG. 10, anti-PD-1+10D1 failed to induce weight loss in the *Ctla-4^{h/m}* mice. Therefore, irAE and cancer immunity can be uncoupled genetically.

In vivo activity demonstrates that the L3D10 antibody retains its anti-tumor activity but displays reduced autoimmune adverse effect observed with other immunotherapeutic antibodies such as 10D1, indicating it is possible to enhance anti-tumor activity without exacerbating autoimmune adverse events. Accordingly, autoimmune side effects are not a necessary price for cancer immunity and that it is possible to uncouple these two activities. Characterization of L3D10 demonstrated that its ability to block the interaction of CTLA-4 with B7.1 and B7.2 is more effective than by 10D1. Further characterization demonstrates that L3D10 and 10D1 bind to immobilized CTLA-4 with a similar

binding profile. However, L3D10 demonstrates much lower binding affinity to soluble CTLA-4 than 10D1.

Example 4

Humanized L3D10 Anti-CTLA-4 Antibodies

Humanized L3D10 antibodies were designed by creating multiple hybrid sequences that fuse select parts of the parental antibody sequence with the human framework sequences, including grafting of the CDR sequences into acceptor frameworks. We evaluated the anti-tumor activity of three of the humanized antibodies (mAb4, mAb5 and mAb6) compared to 10D1 and the chimeric L3D10 antibody using the syngeneic MC38 mouse tumor model in human CTLA-4-knockin mice described in Example 1 above. FIG. 11A shows the treatment schedule of the in vivo experiment; mice were given a total of 4 doses of antibody every 3 days starting on day 7 after inoculation. As shown in FIG. 11B, all humanized antibodies completely eradicated the tumor and were comparable to 10D1. Similar efficacy was seen using the syngeneic MC38 mouse tumor model in heterozygous *Ctla-4^{h/m}* mice and the syngeneic B16-F1 melanoma mouse tumor model in human CTLA-4-knockin mice (data not shown).

To test if the superior safety profiles of L3D10 can be maintained after humanization, we compared mAb4 and mAb5 with 10D1 for their adverse effects when used in combination with anti-PD-1. As shown in FIG. 12, both mAb4 and mAb5 are less toxic than 10D1 when used in combination with anti-PD-1.

Example 5

Binding Characteristics of the Humanized Anti-CTLA-4 Antibodies

In order to confirm that the humanized antibodies retained their CTLA-4 binding characteristics, we looked at binding to immobilized and plate bound CTLA-4. Humanization did not affect binding to immobilized CTLA-4 and all 3 humanized antibodies demonstrated similar binding to the parental chimeric L3D10 antibody (FIG. 13). However, humanization further reduces L3D10 binding to soluble CTLA-4 (FIG. 14).

We have demonstrated that chimeric L3D10 has a 1000-fold higher B7 blocking activity than 10D1. This raised an interesting possibility that blocking B7-CTLA-4 interactions may explain its lack of irAE. However, neither mAb4 nor mAb5 block B7-CTL-A4 interactions in vitro and in vivo. The fact that mAb4 and mAb5 show diminished irAE further supported the notion that blocking B7-CTLA-4 interaction is not responsible for improved safety of L3D10.

Given the proposed role for CTLA-4 in the protection against autoimmune diseases, we proposed reduced binding to soluble CTLA-4 as an underlying mechanism for improved safety profiles. To test this hypothesis, we used the growth weight gain among the female mice that received anti-PD-1+anti-CTLA-4 mAbs during the perinatal period as the basic indicator for irAE. Severe reduction in weight gain was observed in the mice that received both 10D1 and anti-PD-1, whereas those that received mAb5+anti-PD-1

21

had the lowest irAE, followed by mAb4 and then L3D10 (data not shown). The strict inverse correlation with reduced binding to sCTLA-4 are consistent with the central hypothesis.

Example 6

Epitope Mapping of the L3D10 and Humanized Antibodies

In order to map the CTLA-4 binding epitope of the L3D10 antibody and the humanized variants, mAb4 and mAb5, we took advantage of the fact that the mouse and human CTLA-4 proteins are cross-reactive to B7-1, but not to the anti-CTLA-4 antibodies. The fact that anti-human CTLA-4 antibodies do not cross react with murine Ctla-4, presumably reflects differences in the amino acid sequence between human and mouse CTLA-4 in the extracellular domain. FIG. 15 shows the alignment of the human, macaque and mouse CTLA-4 extracellular domains and highlight the sequence conservation between human and macaque, while showing the numerous differences between the murine and primate sequences. Due to conservation of the MYPPPY binding motif (SEQ ID NO: 50), mouse and human CTLA-4 proteins are cross-reactive to B7-1 (72).

Accordingly, we designed 11 mutants of the human CTLA-4Fc protein, designated M1-M11 (SEQ ID NOS: 7-17) in which clusters of amino acids from the human CTLA-4 protein were replaced with amino acids from the murine Ctla-4 protein. The amino acids incorporated into each of the 11 mutants is shown in FIG. 15, and the amino acids sequences of the WT and mutant CTLA-4Fc proteins is shown in FIGS. 16A and B. As the anti-CTLA-4 antibodies used in this study do not bind to murine Ctla-4, binding of the anti-human CTLA-4 antibodies should be abolished when key residues of the antibody binding epitope are replaced with murine amino acids.

DNA vectors encoding 11 CTLA-4Fc mutant proteins were constructed based on the wild type human CTLA-4Fc sequence and proteins were produced by transient transfection in HEK293 at the 0.01 mL scale followed by one-step Protein A chromatography purification, and the yield is provided in Table 1. Many of the mutations appear to affect

22

protein expression as indicated by their yields relative to the WT human CTLA-4Fc protein.

TABLE 1

WT and mutant CTLA-4Fc proteins produced transiently in HEK293 cells.	
Protein name	Yield (mg)
CTLA-4Fc	0.72
WT control	
Mutant 1	1.29
Mutant 2	0.03
Mutant 3	0.21
Mutant 4	0.11
Mutant 5	1.89
Mutant 6	0.38
Mutant 7	0.25
Mutant 8	1.61
Mutant 9	0.01
Mutant 10	0.04
Mutant 11	1.70

The capacity of chimeric L3D10 and the humanized antibodies mAb4 and mAb5 to bind the immobilized CTLA-4Fc mutant constructs was then determined by ELISA in which plates were coated with the CTLA-4 mutant proteins at 1 µg/mL and biotinylated anti-CTLA-4 antibodies, or B7-1 Ig control protein, were added and binding measured using HRP-conjugated streptavidin. The results of binding assays are shown in Tables 2-5. As expected, all 4 binding proteins demonstrated nice dose-dependent binding for the WT CTLA-4Fc protein. However, mutations that were introduced into the M9 and M10 proteins appear to alter the overall structure and these mutants failed to bind B7-1Fc. Mutations introduced in M2 and M4 also partially altered CTLA-4 conformation as indicated by reduced binding relative to the WT protein. Consistent with this notion, all 4 of these mutants (M2, M4, M9 and M10) were expressed at much lower yield (Table 1). In contrast, using binding to the WT CTLA-4Fc protein and binding of the B7-1Fc proteins as references, M11 clearly stands out as a protein that is expressed well, binds B7-1Fc efficiently but failed to bind two humanized anti-CTLA-4 antibodies. Its binding to original L3D10 is also reduced by approximately 100-fold (Table 3). As expected, the mutations that affect the overall conformation also affected the binding to the anti-CTLA-4 antibodies.

TABLE 2

Integrity of CTLA-4Ig mutants as indicated by their binding to B7-1 Ig fusion protein. Binding to CTLA-4Fc proteins was performed by ELISA, with the amounts of biotinylated protein bound measured by horse-radish peroxidase (HRP)-conjugated streptavidin.												
Protein Conc.	WT	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11
0	0.193	0.196	0.202	0.184	0.182	0.182	0.184	0.185	0.18	0.172	0.175	0.174
0	0.19	0.182	0.177	0.175	0.171	0.171	0.173	0.17	0.168	0.164	0.162	0.163
10 ng/ml	0.259	0.328	0.204	0.267	0.199	0.286	0.255	0.218	0.293	0.166	0.167	0.22
10 ng/ml	0.257	0.311	0.187	0.249	0.184	0.271	0.244	0.22	0.276	0.154	0.159	0.217
100 ng/ml	1.137	1.594	0.316	1.087	0.359	1.513	1.093	0.785	1.468	0.164	0.164	0.884
100 ng/ml	1.111	1.553	0.299	1.082	0.34	1.221	1.049	0.695	1.375	0.155	0.15	1.045
1 ug/ml	2.813	3.147	1.179	3.147	1.375	2.877	3.053	2.703	3.253	0.199	0.171	3.053
1 ug/ml	2.651	3.053	0.986	2.864	1.413	3.025	2.983	2.716	2.93	0.218	0.172	3.159

Values shown are the OD450 measurements.

WT = wild type CTLA-4Fc.

M1-M11 are CTLA-4Fc mutant proteins.

TABLE 3

Epitope mapping of chimeric L3D10 antibody. Binding to CTLA-4Fc proteins was performed by ELISA, with the amounts of biotinylated protein bound measured by horse-radish peroxidase (HRP)-conjugated streptavidin.

Protein Conc.	WT	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11
0	0.202	0.196	0.2	0.187	0.184	0.189	0.192	0.198	0.187	0.179	0.179	0.183
0	0.195	0.187	0.185	0.18	0.176	0.176	0.176	0.176	0.17	0.166	0.166	0.167
10 ng/ml	1.433	2.47	0.375	0.62	0.507	1.539	1.033	0.714	1.233	0.18	0.18	0.202
10 ng/ml	1.518	2.432	0.317	0.587	0.356	1.366	0.976	0.738	1.237	0.171	0.169	0.203
100 ng/ml	3.053	3.253	1.384	2.318	2.142	2.841	2.699	2.495	2.909	0.295	0.215	0.635
100 ng/ml	3.025	3.239	1.164	2.354	1.409	2.991	2.771	2.483	2.841	0.304	0.216	0.759
1 ug/ml	3.373	3.268	2.387	3.184	2.651	3.025	3.092	3.147	3.136	0.916	0.804	2.841
1 ug/ml	3.114	2.967	2.619	3.124	2.659	3.034	3.072	2.991	3.034	0.916	0.868	2.983

Values shown are the OD450 measurements.

WT = wild type CTLA-4Fc.

M1-M11 are CTLA-4Fc mutant proteins

TABLE 4

Epitope mapping of humanized antibody mAb4. Binding to CTLA-4Fc proteins was performed by ELISA, with the amounts of biotinylated protein bound measured by horse-radish peroxidase (HRP)-conjugated streptavidin.

Protein Conc.	WT	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11
10 ng/ml	0.312	2.264	0.207	0.198	0.194	0.407	0.22	0.194	0.247	0.177	0.181	0.172
10 ng/ml	0.29	2.297	0.184	0.178	0.174	0.378	0.202	0.185	0.222	0.154	0.16	0.164
100 ng/ml	1.077	2.827	0.203	0.27	0.219	1.371	0.459	0.281	0.725	0.171	0.17	0.172
100 ng/ml	0.841	3.061	0.194	0.264	0.208	1.589	0.42	0.277	0.801	0.154	0.155	0.159
1 ug/ml	2.51	2.881	0.339	0.882	0.473	2.79	1.992	1.169	2.33	0.175	0.17	0.178
1 ug/ml	2.471	2.958	0.263	1.121	0.573	2.795	2.016	1.243	2.642	0.167	0.169	0.185

Values shown are the OD450 measurements.

WT = wild type CTLA-4Fc.

M1-M11 are CTLA-4Fc mutant proteins

TABLE 5

Epitope mapping of humanized antibody mAb5. Binding to CTLA-4Fc proteins was performed by ELISA, with the amounts of biotinylated protein bound measured by horse-radish peroxidase (HRP)-conjugated streptavidin.

Protein Conc.	WT	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11
10 ng/ml	0.597	2.307	0.195	0.544	0.189	1.239	0.603	0.19	0.5	0.373	0.169	0.157
10 ng/ml	0.535	2.244	0.162	0.195	0.435	1.188	0.516	0.535	0.47	0.148	0.15	0.152
100 ng/ml	1.947	2.632	0.182	0.389	0.248	2.601	1.296	0.521	2.001	0.15	0.15	0.152
100 ng/ml	2.229	2.186	0.175	0.364	0.221	2.425	0.875	0.405	2	0.137	0.139	0.148
1 ug/ml	2.724	2.05	0.259	1.662	0.725	2.654	2.355	1.418	2.548	0.157	0.151	0.162
1 ug/ml	2.742	2.297	0.274	1.549	0.724	2.84	2.374	1.369	2.69	0.147	0.143	0.165

Values shown are the OD450 measurements.

WT = wild type CTLA-4Fc.

M1-M11 are CTLA-4Fc mutant proteins

TABLE 6

Raw data from a repeat study showing specific loss of antigenic epitope only in M11. As in Tables 2-5, except that additional controls were included to shown specificity of the binding.

	Ab Conc.	hCTLA4-Fc	M1	M2	M3	M4	M5	M6	M7	M8	M9	M10	M11
Biotin-LD310	0	0.18	0.187	0.377	0.183	0.22	0.177	0.183	0.186	0.368	0.15	0.215	0.171
	0	0.177	0.222	0.538	0.167	0.18	0.229	0.177	0.142	0.217	0.293	0.114	0.155
	10 ng/ml	1.705	2.692	0.469	0.623	0.817	1.853	1.244	0.837	1.27	0.158	0.169	0.19
	10 ng/ml	1.799	2.779	0.333	0.593	0.563	1.802	1.881	0.884	1.454	0.213	0.157	0.194
	100 ng/ml	3.316	3.195	1.313	2.244	2.233	3.251	3.032	2.672	3.015	0.419	0.26	0.752
	100 ng/ml	3.458	3.567	1.37	2.535	2.356	3.316	3.032	2.875	3.157	0.346	0.272	0.746
	1000 ng/ml	3.567	3.509	2.833	3.333	3.08	3.413	3.282	3.299	3.352	1.124	0.945	2.888
Biotin-hB7-1	1000 ng/ml	3.672	3.509	2.755	3.299	3.149	3.537	3.316	3.352	3.435	1.181	0.941	2.914
	0	0.195	0.2	0.202	0.198	0.192	0.197	0.195	0.198	0.192	0.186	0.185	0.186
	0	0.192	0.185	0.181	0.192	0.178	0.178	0.178	0.187	0.173	0.169	0.168	0.161
	10 ng/ml	0.316	0.37	0.216	0.304	0.22	0.345	0.279	0.258	0.326	0.177	0.176	0.239
	10 ng/ml	0.31	0.356	0.21	0.414	0.26	0.331	0.279	0.253	0.297	0.159	0.167	0.236
	100 ng/ml	1.581	1.882	0.333	1.245	0.527	1.813	1.235	0.899	1.357	0.176	0.172	1.092
	100 ng/ml	1.525	1.928	0.323	1.345	0.489	1.735	1.385	0.987	1.643	0.162	0.155	1.283
Biotin-mAb4	1000 ng/ml	3.76	3.6	1.167	3.435	1.573	3.316	3.413	3.101	3.635	0.232	0.189	3.568
	1000 ng/ml	3.6	3.673	1.316	3.51	2.009	3.459	3.413	3.183	3.635	0.215	0.181	3.673
	10 ng/ml	0.451	2.812	0.207	0.202	0.194	0.626	0.23	0.207	0.327	0.197	0.205	0.181
	10 ng/ml	0.417	2.693	0.181	0.179	0.177	0.642	0.22	0.195	0.32	0.158	0.182	0.162
	100 ng/ml	1.868	3.568	0.212	0.29	0.256	2.618	0.589	0.345	1.532	0.172	0.174	0.171
	100 ng/ml	1.938	3.317	0.203	0.274	0.247	2.126	0.571	0.305	1.419	0.155	0.155	0.162
	1000 ng/ml	2.99	3.568	0.268	1.181	0.712	2.922	2.187	1.329	2.817	0.181	0.17	0.177
Biotin-mAb5	1000 ng/ml	3.033	3.51	0.268	1.184	0.759	3.071	2.358	1.475	2.809	0.144	0.171	0.187
	10 ng/ml	0.983	2.654	0.202	0.218	0.197	1.409	0.429	0.218	0.727	0.176	0.176	0.17
	10 ng/ml	0.955	2.604	0.184	0.2	0.168	1.389	0.380	0.21	0.761	0.148	0.154	0.152
	100 ng/ml	2.669	3.007	0.232	0.334	0.319	2.908	1.839	0.523	2.609	0.145	0.161	0.16
	100 ng/ml	2.741	3.188	0.203	0.354	0.374	2.895	1.741	0.478	2.604	0.145	0.148	0.157
	1000 ng/ml	3.183	3.146	0.327	1.837	1.019	2.968	2.817	1.72	3.042	0.173	0.163	0.174
	1000 ng/ml	3.209	3.316	0.321	1.867	1.015	3.196	2.857	1.766	3.031	0.143	0.163	0.187

Ab conc.	Biotin-L3D10 mCTLA4-Fc	Biotin-L3D10 hIg-Fc	Biotin-hB7-1 mCTLA4-Fc	Biotin-hB7-1 hIg-Fc	Biotin-HL12	Biotin-HL12	Biotin-HL32
0	0.19	0.198	0.202	0.191			
0	0.189	0.184	0.18	0.185			
10 ng/ml	0.201	0.201	0.338	0.181	0.179	0.188	0.185
10 ng/ml	0.18	0.182	0.318	0.164	0.165	0.162	0.17
100 ng/ml	0.303	0.315	1.635	0.176	0.171	0.177	0.185
100 ng/ml	0.314	0.326	1.668	0.165	0.162	0.168	0.185
1000 ng/ml	0.942	1.385	3.569	0.18	0.177	0.182	0.184
1000 ng/ml	0.94	1.473	3.353	0.179	0.172	0.177	0.176
	mCTLA4-Fc	hIgG	mCTLA4-Fc	hIgG	mCTLA4-Fc	hIgG	mCTLA4-Fc
	Biotin-L3D10		Biotin-hB7-1		Biotin-HL12		Biotin-HL32

Since L3D10 retained significant binding the M11, we tested if the binding is specific. We coated plate with human CTLA-4-Fc (hCTLA-4Fc), mouse CTLA-4-Fc (mCTLA-4-Fc), Control IgG1-Fc or all mutant hCTLA-4-Fc and measured their binding to B7-1Fc along with L3D10, mAb4 and mAb5. The bulk of the data are presented in Table 6. As shown in FIG. 17, biotinylated B7-1 binds hCTLA-4, mCTLA-4 and M11, equally well. The specificity of the assay is demonstrated by lack of binding to IgG1-Fc. Interesting, while L3D10-binding to M11 is stronger than those to IgG1-Fc and mCTLA-4-Fc, significant binding to IgG1-Fc suggest that the chimeric antibody binding to M11 maybe nonspecific. In contrast, none of the humanized antibodies bind to M11, mCTLA-4, and IgG1-Fc control. These data demonstrate that mutations introduced in M11 selectively ablated L3D10, mAb4 and mAb5 binding to CTLA-4.

Using known complex structure 133, we mapped the CTLA-4 epitope in a 3-D structure. As shown in FIG. 18, the epitope recognized by these mAbs localized within the area covered by B7-1. As such, L3D10, mAb4 and mAb5 binding to CTLA-4 would be mutually exclusive to that of B7-1. The poor blocking of mAb4 and mAb5 is due to lower avidity rather than distinctive binding domains.

Using a number of mutants of the human CTLA-4Fc protein in which clusters of amino acids from the human

CTLA-4 protein were replaced with amino acids from the murine CtlA-4 protein, we clearly demonstrate that when we replace 4 amino acids that immediately follow the known B7-1 binding domain of CTLA-4, MYPPPY, dose-dependent binding of the antibodies is largely abolished. The fact that the binding epitope maps directly adjacent to the B7-1 binding domain correlates well with the demonstrated ability of the L3D10 antibodies to block B7-CTLA-4 interactions both in vitro and in vivo.

Example 7

Generation of Fusion Proteins that Show Diminished Binding to One or More Cancer Immunotherapeutic Anti-CTLA-4 mAbs

We have generated a panel of 17 CTLA-4-Fc fusion proteins with various mutations, designated M1-M17 (SEQ ID NOS: 7-23), which include the 11 proteins identified in Example 10 (FIG. 16) plus 6 additional mutants (FIG. 19). Further mutations were made to M17. The new mutants are called M17-1, M17-2, M17-3 and M17-4 (SEQ ID NOS: 42-45). These mutations were generated anticipating that they would ablate binding to a broad spectrum of anti-CTLA-4 antibodies but retain binding to the CTLA-4

ligands, CD80 (B7-1) and CD86 (B7-2). As the first step, we coated a given concentration of the fusion proteins on the plate and added biotinylated B7-1Fc. As shown in FIGS. 20A and B, all but 2 fusion proteins (M9 and M10) retain binding to B7-1, although a significant reduction in B7-1 5 binding was found in M2, M4 and M7. Since these proteins are functionally less active, it is unlikely that these mutants are optimal for in vivo protection against autoimmune adverse effects. Additional data demonstrate that M15, M17, M17-1 and M17-2 retained strong binding to both CD80 and CD86 (FIG. 21).

Next, we evaluated if these mutant proteins (M1-M17, and M17-1 to M17-4) have lost binding to anti-CTLA4 mAbs that are either approved for clinical use or being developed for clinical therapy. A total of 4 mAbs were tested. For mAb1 (Ipilimumab), fusion proteins M11, M13, M15, M17, M17-1, M17-2 and M17-3 have lost binding, and thus can be used for protection against autoimmune diseases induced by this mAb (FIG. 22A). For autoimmune diseases induced by mAb2 (Tremelimumab), M15, M17, M17-2 and M17-3 will be effective (FIG. 22B). For mAb4 and mAb5, M11, M12, M14, and M16 are likely the most effective, although M3 and M17 may also work (FIGS. 23, 24 and 25).

Therefore, as summarized in Table 7, CTLA4-Fc fusion protein M17 has broad spectrum activity for use in protection against autoimmune diseases induced by anti-CTLA4 antibodies. M17-3 shows low binding to anti-CTLA4 antibodies but low binding to B7-1 and B7-2. However, M17-2 showed better binding to B7-1 and B7-2 and yet shows acceptably low cross-reactivity against a broad range of anti-human CTLA-4 mAbs.

TABLE 7

Identification of CTLA4-Fc fusion proteins that retains binding to B7-1, but not to anti-CTLA4 mAb. Relative binding is presented by "+" or "-", where "++++" is the strongest binding and "-" indicates no binding.						
CTLA4-Fc Fusion protein	hB7-1Fc	hB7-2Fc	mAb1 (Ipilimumab)	mAb2 (Tremelimumab)	mAb4	mAb5
abatacept	+++	ND	+++	+++	+++	+++
hIgGFc ctl	-	-	-	-	-	-
M1	+++	ND	+++	ND	++++	++++
M2	+	ND	+	ND	-	-
M3	+++	ND	+++	ND	+	+
M4	+	ND	+	ND	+	+
M5	+++	ND	+++	ND	+++	+++
M6	+++	ND	+++	ND	+	++
M7	++	ND	+	ND	+	+
M8	+++	ND	+++	ND	+++	+++
M9	-	ND	-	ND	-	-
M10	-	ND	-	ND	-	-
M11	+++	ND	-	+++	-	-
M12	+++	ND	+	+++	-	-
M13	+++	ND	-	+++	++	++
M14	+++	ND	+	+++	-	-
M15	+++	+++	-	-	+++	++++
M16	+++	ND	+	+++	-	-
M17	+++	++	-	-	+	+
M17-1	+++	++	-	+++	++	-
M17-2	+++	+++	-	-	+/-	-
M17-3	++	+/-	-	-	-	-
M17-4	+++	+	+	+++	++	-

ND = not determined.

Blocking B7-1 and B7-2 does not Prevent Immunotherapeutic Effect of Ipilimumab

A critical requirement for treating anti-CTLA-4-immunotherapy related adverse effects with CTLA-4-Fc mutants that bind to B7-1/B7-2 is that blocking B7-1 and B7-2 does not affect the immunotherapeutic effect. Since mice with targeted mutations of Cd80 (encoding B7-1) and Cd86 (encoding B7-2) do not have Treg (17) and thus express very little Ctla4, we tested this prediction by using a saturating dose of anti-B7-1 (1G10) and anti-B7-2 (GL-1) mAbs, which block the binding of human CTLA-4 to mB7-1 and mB7-2, respectively (FIG. 26A). As diagrammed in FIG. 26B, we treated MC38 tumor-bearing mice with Ipilimumab in conjunction with either control Ig or a combination of anti-mB7-1 and anti-mB7-2 mAbs. The anti-mB7 mAbs used completely masked all B7-1 and B7-2 in the PBL as their binding to new anti-mB7 mAb is reduced to what was observed in mice with targeted mutations of both Cd80 and Cd86 (dKO) (FIG. 26C, D). Functional blockade of B7 by anti-B7-1 and anti-B7-2 mAbs was evaluated using sera collected at day 22 after tumor challenge to evaluate anti-human IgG antibody response and confirmed by the fact that antibody response against Ipilimumab was abrogated (FIG. 26E). Importantly, saturating blockade of mB7 did not affect the Ipilimumab-induced tumor rejection as anti-mB7 and control Ig treated mice are equally responsive to Ipilimumab (FIG. 26F). These data indicate that anti-B7-1 and anti-B7-2 as well as B7-binding CTLA-4 fusion proteins that do not bind anti-CTLA-4 antibodies can be used to treat irAE

associated with anti-CTLA-4 immunotherapy, while leaving the protective immunotherapeutic activity intact.

Example 9

Use of CTLA-4 Fusion Proteins to Treat Autoimmune Adverse Effects Associated with Anti-CTLA-4 and Anti-PD-1 Combination Therapy

Combination therapy with anti-CTLA-4 and anti-PD-1 represents the most effective cancer immunotherapy. However, with greater than 50% of patients receiving immunotherapy developing irAEs, there is an urgent need to develop therapeutic approaches to treat or prevent irAE. As demonstrated in Example 8, the immunotherapeutic effect of Ipilimumab (mAb1) is unaffected by the blockade of B7-1 and B7-2. These findings prompted us to test if mutant CTLA-4Fc can be used to treat irAE associated with Ipilimumab+ anti-PD-1 combination therapy. Since Ipilimumab does not bind to CTLA-4-Fc fusion proteins M11 or M15 (belatacept), we tested to see if co-administration of belatacept or M11 or M15 can ameliorate irAE, using retarded growth as a readout of irAE (see Example 2). As shown in FIG. 27, low doses of either M11 (100 µg/injection) or M15 (200 µg/injection) prevented irAE. In addition to M11 and M15, other proteins with similar properties, including M13, M16 and M17 may also be used.

Example 10

Identification of Fusion Proteins that do not Interfere with Anti-CTLA4 Immunotherapeutic Activity

In order to determine whether CTLA-4-Fc fusion proteins interfere with anti-CTLA4 immunotherapeutic effects, we injected MC38-tumor bearing mice with either control IgG Fc (200 µg/injection) or Ipilimumab (100 µg/injection) in combination with control IgGFc (100 µg/injection) or the CTLA4 fusion proteins, Abetacept, M15 or M17 (100 µg/injection) on days 7, 10, 13 and 17. The tumor growth rate were shown in FIG. 28A, while the mouse survival to early removal endpoint are shown in FIG. 28B. As expected, Abatacept ablated the immunotherapeutic effect of the Ipilimumab. In contrast, M15 and M17 did not interfere with the therapeutic effect of Ipilimumab.

Example 11

Modulation of T Cell Activation and its Associated Adverse Effect in NSG Mice Reconstituted with Human Hematopoietic System

To determine the impact of CTLA-4 fusion proteins on the activation of human T cells and associated irAE in vivo, we reconstituted the 3 week old NSG mice with human hematopoietic stem cells and monitored T cell activation. As shown in FIGS. 29-32, different CTLA-4 mAbs appear to have different effects on T cell phenotype and the adverse effects of Ipilimumab. Thus, as shown in FIG. 29A, treatment with Ipilimumab prevents NSG mice from gaining weight, and this is prevented by co-administration of M15 or M17-2. Interestingly, M15 induced elevation of Alanine transaminase when used in conjunction with Ipilimumab (FIG. 29B). These data suggest that M15 may induce liver

damage when used in conjunction with Ipilimumab. Importantly, no such effect was observed when M17-2 was used in conjunction with M15.

To understand the immunological basis of the differentiation between M15 and M17, we sacrificed the NSG mice on day 31 and analyzed the composition of T cell subsets. As shown in FIG. 30, M17 induced more central memory T cells, while reducing effector T cells. The reduction of effector T cells may explain the reduced adverse effect as this is the subset that migrates into organs to cause immune destruction. Furthermore, since NKT cells are the primary mediator of hepatitis, we compared M15 and M17-2 for their impact on NKT cells, based on expression of both CD56 and CD3. As shown in FIG. 31, M17-2 reduced the number of NKT while M15 enhanced it.

The regulatory T cells suppress autoimmune diseases and cancer immunity. It is therefore desirable to reduce Treg to enhance cancer immunity. However, if the reduction is too severe, one may induce autoimmune disease. Consistent with a critical role for B7-CD28 interactions in the generation and maintenance of Treg in mice, both M15 and M17-2 significantly reduced the % of Treg (FIGS. 32A and B). However, M15 is much more effective in reducing Treg (FIGS. 32A and B). Furthermore, Treg from M15-treated mice had substantially lower CTLA-4 (FIGS. 32C and D). Given the essential role for CTLA-4 in Treg function in vivo, it is likely that Treg from the M15-treated mice are functionally impaired. Thus, a severe reduction of Treg combined with selective reduction of CTLA-4 protein levels suggest an intriguing possibility that M15 may induce autoimmune destruction when used in conjunction Ipilimumab.

All publications and patents mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference in its entirety. While the invention has been described in connection with specific embodiments thereof, it will be understood that it is capable of further modifications and this application is intended to cover any variations, uses, or adaptations of the invention following, in general, the principles of the invention and including such departures from the present disclosure as come within known or customary practice within the art to which the invention pertains and as may be applied to the essential features hereinbefore set forth.

REFERENCES CITED

10. Leach D R, Krummel M F, Allison J P. Enhancement of antitumor immunity by CTLA-4 blockade [see comments]. *Science*. 1996; 271(5256):1734-6.
- 108 Guinan, E. C. et al. Transplantation of anergic histoincompatible bone marrow allografts. *N. Engl. J. Med.* 340, 1704-1714, doi:10.1056/NEJM199906033402202 (1999).
- 128 Magistrelli, G. et al. A soluble form of CTLA-4 generated by alternative splicing is expressed by nonstimulated human T cells. *Eur. J. Immunol.* 29, 3596-3602, doi:10.1002/(SICI)1521-4141(199911)29:11<3596::AID-IMMU3596>3.0.CO;2-Y (1999).
- 64 Ueda, H. et al. Association of the T-cell regulatory gene CTLA-4 with susceptibility to autoimmune disease. *Nature* 423, 506-511 (2003).
- 129 Kremer, J. M. et al. Treatment of rheumatoid arthritis by selective inhibition of T-cell activation with fusion pro-

tein CTLA-4Ig. *N. Engl. J. Med.* 349, 1907-1915, doi:10.1056/NEJMoa035075 (2003).

130 Abrams, J. R. et al. CTLA-4Ig-mediated blockade of T-cell costimulation in patients with psoriasis vulgaris. *J. Clin. Invest.* 103, 1243-1252, doi:10.1172/JCI5857 (1999).

131 Gerold, K. D. et al. The soluble CTLA-4 splice variant protects from type 1 diabetes and potentiates regulatory T-cell function. *Diabetes* 60, 1955-1963, doi:10.2337/db11-0130 (2011).

132 Peach, R. J. et al. Complementarity determining region 1 (CDR1)- and CDR3-analogous regions in CTLA-4 and CD28 determine the binding to B7-1. *J. Exp. Med.* 180, 2049-2058 (1994).

1. Townsend A R M, Tothbard J, Gotch F M, Bahadur G, Wraith D, McMichael A J. The epitope of influenza nucleoprotein recognized by cytotoxic lymphocytes can be defined with short synthetic peptides. *Cell.* 1986; 44:959-68.

2. Zinkernagel R M, Doherty P C. Restriction of in vitro T cell-mediated cytotoxicity in lymphocytic choriomeningitis within a syngeneic or semiallogeneic system. *Nature.* 1974; 248:701-2.

3. Lafferty K J, Prowse S J, Simeonovic C J, Warren H S. Immunobiology of tissue transplantation: a return to the passenger leukocyte concept. *Annu Rev Immunol.* 1983; 1:143-73.

4. Liu Y, Linsley P S. Costimulation of T-cell growth. *Curr Opin Immunol.* 1992; 4(3):265-70.

5. Schwartz R H. Costimulation of T lymphocytes: the role of CD28, CTLA-4, and B7/BB1 in interleukin-2 production and immunotherapy. *Cell.* 1992; 71(7):1065-8.

6. Freeman G J, Freedman A S, Segil J M, Lee G, Whitman J F, Nadler L M. B7, a new member of the Ig superfamily with unique expression on activated and neoplastic B cells. *J Immunol.* 1989; 143(8):2714-22.

7. Freeman G J, Gribben J G, Boussiotis V A, Ng J W, Restivo V A, Jr., Lombard L A, et al. Cloning of B7-2: a CTLA-4 counter-receptor that costimulates human T cell proliferation [see comments]. *Science.* 1993; 262(5135):909-11.

8. Hathcock K S, Laszlo G, Dickler H B, Bradshaw J, Linsley P, Hodes R J. Identification of an alternative CTLA-4 ligand costimulatory for T cell activation [see comments]. *Science.* 1993; 262(5135):905-7.

9. Wu Y, Guo Y, Liu Y. A major costimulatory molecule on antigen-presenting cells, CTLA-4 ligand A, is distinct from B7. *J Exp Med.* 1993; 178(5):1789-93.

10. Leach D R, Krummel M F, Allison J P. Enhancement of antitumor immunity by CTLA-4 blockade [see comments]. *Science.* 1996; 271(5256):1734-6.

11. Linsley P S, Brady W, Urnes M, Grosmaire L S, Damle N K, Ledbetter J A. CTLA-4 is a second receptor for the B cell activation antigen B7. *J Exp Med.* 1991; 174(3):561-9.

12. Linsley P S, Clark E A, Ledbetter J A. T-cell antigen CD28 mediates adhesion with B cells by interacting with activation antigen B7/BB-1. *Proc Natl Acad Sci USA.* 1990; 87(13):5031-5.

21. May K F, Roychowdhury S, Bhatt D, Kocak E, Bai X F, Liu J Q, et al. Anti-human CTLA-4 monoclonal antibody promotes T cell expansion and immunity in a hu-PBL-SCID model: a new method for preclinical screening of costimulatory monoclonal antibodies. *Blood.* 2005; 105:1114-20. PubMed PMID: 15486062.

20. Lute K D, May K F, Lu P, Zhang H, Kocak E, Mosinger B, et al. Human CTLA-4-knock-in mice unravel the quantitative link between tumor immunity and autoimmunity induced by anti-CTLA-4 antibodies. *Blood.* 2005. PubMed PMID: 16037385.

49. Keler, T. et al. Activity and safety of CTLA-4 blockade combined with vaccines in cynomolgus macaques. *J. Immunol.* 171, 6251-6259 (2003).

50. Wing, K. et al. CTLA-4 control over Foxp3+ regulatory T cell function. *Science* 322, 271-275, doi:10.1126/science.1160062 (2008).

66. Qureshi, O. S. et al. Trans-endocytosis of CD80 and CD86: a molecular basis for the cell-extrinsic function of CTLA-4. *Science* 332, 600-603, doi:10.1126/science.1202947 (2011).

30. Walunas, T. L., et al., CTLA-4 can function as a negative regulator of T cell activation. *Immunity*, 1994. 1(5): p. 405-13.

31. Krummel, M. F. and J. P. Allison, CD28 and CTLA-4 have opposing effects on the response of T cells to stimulation. *J Exp Med*, 1995. 182(2): p. 459-65.

52. Simpson, T. R. et al. Fc-dependent depletion of tumor-infiltrating regulatory T cells co-defines the efficacy of anti-CTLA-4 therapy against melanoma. *J. Exp. Med.* 210, 1695-1710, doi:10.1084/jem.20130579 (2013).

55. Korman, A. J., Peggs, K. S. & Allison, J. P. Checkpoint blockade in cancer immunotherapy. *Adv. Immunol.* 90, 297-339, doi:10.1016/50065-2776(06)90008-X (2006).

56. Ribas, A. et al. Tremelimumab (CP-675,206), a cytotoxic T lymphocyte associated antigen 4 blocking monoclonal antibody in clinical development for patients with cancer. *Oncologist* 12, 873-883, doi:10.1634/theoncologist.12-7-873 (2007).

57. Ribas, A. et al. Phase III randomized clinical trial comparing tremelimumab with standard-of-care chemotherapy in patients with advanced melanoma. *J. Clin. Oncol.* 31, 616-622, doi:10.1200/JCO.2012.44.6112 (2013).

58. Lee, K. M. et al. Molecular basis of T cell inactivation by CTLA-4 [In Process Citation]. *Science* 282, 2263-2266 (1998).

59. Marengere, L. E. et al. Regulation of T cell receptor signaling by tyrosine phosphatase SYP association with CTLA-4 [published errata appear in *Science* 1996 Dec. 6; 274(5293):1597 and 1997 Apr. 4; 276(5309):21]. *Science* 272, 1170-1173 (1996).

60. Liu, Y. Is CTLA-4 a negative regulator for T-cell activation? *Immunol. Today* 18, 569-572 (1997).

61. Tivol, E. A. et al. Loss of CTLA-4 leads to massive lymphoproliferation and fatal multiorgan tissue destruction, revealing a critical negative regulatory role of CTLA-4. *Immunity* 3, 541-547 (1995).

62. Waterhouse, P. et al. Lymphoproliferative disorders with early lethality in mice deficient in CTLA-4 [see comments]. *Science* 270, 985-988 (1995).

63. Bachmann, M. F., Kohler, G., Ecabert, B., Mak, T. W. & Kopf, M. Cutting edge: lymphoproliferative disease in the absence of CTLA-4 is not T cell autonomous. *J. Immunol.* 163, 1128-1131 (1999).

64. Bachmann, M. F. et al. Normal pathogen-specific immune responses mounted by CTLA-4-deficient T cells: a paradigm reconsidered. *Eur. J. Immunol.* 31, 450-458 (2001).

65. Nguyen, T. V., Ke, Y., Zhang, E. E. & Feng, G. S. Conditional deletion of Shp2 tyrosine phosphatase in thymocytes suppresses both pre-TCR and TCR signals. *J. Immunol.* 177, 5990-5996 (2006).

72 Peach R J, Bajorath J, Brady W, Leytze G, Greene J, Naemura J, et al. Complementarity determining region 1 (CDR1)- and CDR3-analogous regions in CTLA-4 and CD28 determine the binding to B7-1. *J Exp Med.* 1994; 180(6):2049-58.

17. Proietto A II, Lahoud M H, Wu L. Distinct functional capacities of mouse thymic and splenic dendritic cell popu-

lations. Immunol Cell Biol. 2008 November-December; 86(8):700-8. doi: 10.1038/icb.2008.63. Epub 2008 Sep. 9.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 50

<210> SEQ ID NO 1

<211> LENGTH: 223

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

Met Ala Cys Leu Gly Phe Gln Arg His Lys Ala Gln Leu Asn Leu Ala
1           5           10           15
Thr Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro
20           25           30
Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
35           40           45
Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
50           55           60
Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
65           70           75           80
Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
85           90           95
Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
100          105          110
Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
115          120          125
Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly
130          135          140
Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser
145          150          155          160
Asp Phe Leu Leu Trp Ile Leu Ala Ala Val Ser Ser Gly Leu Phe Phe
165          170          175
Tyr Ser Phe Leu Leu Thr Ala Val Ser Leu Ser Lys Met Leu Lys Lys
180          185          190
Arg Ser Pro Leu Thr Thr Gly Val Tyr Val Lys Met Pro Pro Thr Glu
195          200          205
Pro Glu Cys Glu Lys Gln Phe Gln Pro Tyr Phe Ile Pro Ile Asn
210          215          220

```

<210> SEQ ID NO 2

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1           5           10           15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
20           25           30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
35           40           45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
50           55           60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65           70           75           80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu

```

-continued

85					90					95					
Met	Tyr	Pro	Pro	Pro	Tyr	Tyr	Leu	Gly	Ile	Gly	Asn	Gly	Thr	Gln	Ile
			100					105					110		
Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro	Lys
		115					120					125			
Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu	Leu
		130					135					140			
Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
145						150					155				160
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val
				165					170					175	
Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val
			180					185					190		
Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser
		195					200					205			
Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu
		210					215					220			
Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala
225						230					235				240
Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro
				245					250					255	
Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln
			260					265					270		
Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala
		275					280					285			
Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
		290				295					300				
Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
305						310					315				320
Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser
				325					330					335	
Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser
			340					345					350		
Leu	Ser	Pro	Gly	Lys											
			355												

<210> SEQ ID NO 3

<211> LENGTH: 126

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

Lys	Ala	Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg
1				5					10					15	
Gly	Ile	Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr
			20					25					30		
Glu	Val	Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu
		35					40					45			
Val	Cys	Ala	Ala	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp
		50				55					60				
Asp	Ser	Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	Gln	Val	Asn	Leu	Thr
65				70					75					80	
Ile	Gln	Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val
			85						90					95	

-continued

Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr
 100 105 110

Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120 125

<210> SEQ ID NO 4
 <211> LENGTH: 174
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Ala Cys Leu Gly Phe Gln Arg His Lys Ala Gln Leu Asn Leu Ala
 1 5 10 15

Thr Arg Thr Trp Pro Cys Thr Leu Leu Phe Phe Leu Leu Phe Ile Pro
 20 25 30

Val Phe Cys Lys Ala Met His Val Ala Gln Pro Ala Val Val Leu Ala
 35 40 45

Ser Ser Arg Gly Ile Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly
 50 55 60

Lys Ala Thr Glu Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln
 65 70 75 80

Val Thr Glu Val Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr
 85 90 95

Phe Leu Asp Asp Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val
 100 105 110

Asn Leu Thr Ile Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile
 115 120 125

Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly
 130 135 140

Asn Gly Thr Gln Ile Tyr Val Ile Ala Lys Glu Lys Lys Pro Ser Tyr
 145 150 155 160

Asn Arg Gly Leu Cys Glu Asn Ala Pro Asn Arg Ala Arg Met
 165 170

<210> SEQ ID NO 5
 <211> LENGTH: 223
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 5

Met Ala Cys Leu Gly Leu Arg Arg Tyr Lys Ala Gln Leu Gln Leu Pro
 1 5 10 15

Ser Arg Thr Trp Pro Phe Val Ala Leu Leu Thr Leu Leu Phe Ile Pro
 20 25 30

Val Phe Ser Glu Ala Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala
 35 40 45

Ser Ser His Gly Val Ala Ser Phe Pro Cys Glu Tyr Ser Pro Ser His
 50 55 60

Asn Thr Asp Glu Val Arg Val Thr Val Leu Arg Gln Thr Asn Asp Gln
 65 70 75 80

Met Thr Glu Val Cys Ala Thr Thr Phe Thr Glu Lys Asn Thr Val Gly
 85 90 95

Phe Leu Asp Tyr Pro Phe Cys Ser Gly Thr Phe Asn Glu Ser Arg Val
 100 105 110

Asn Leu Thr Ile Gln Gly Leu Arg Ala Val Asp Thr Gly Leu Tyr Leu
 115 120 125

-continued

Cys Lys Val Glu Leu Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly
 130 135 140

Asn Gly Thr Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser
 145 150 155 160

Asp Phe Leu Leu Trp Ile Leu Val Ala Val Ser Leu Gly Leu Phe Phe
 165 170 175

Tyr Ser Phe Leu Val Ser Ala Val Ser Leu Ser Lys Met Leu Lys Lys
 180 185 190

Arg Ser Pro Leu Thr Thr Gly Val Tyr Val Lys Met Pro Pro Thr Glu
 195 200 205

Pro Glu Cys Glu Lys Gln Phe Gln Pro Tyr Phe Ile Pro Ile Asn
 210 215 220

<210> SEQ ID NO 6
 <211> LENGTH: 126
 <212> TYPE: PRT
 <213> ORGANISM: Mus musculus

<400> SEQUENCE: 6

Glu Ala Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser His
 1 5 10 15

Gly Val Ala Ser Phe Pro Cys Glu Tyr Ser Pro Ser His Asn Thr Asp
 20 25 30

Glu Val Arg Val Thr Val Leu Arg Gln Thr Asn Asp Gln Met Thr Glu
 35 40 45

Val Cys Ala Thr Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp
 50 55 60

Tyr Pro Phe Cys Ser Gly Thr Phe Asn Glu Ser Arg Val Asn Leu Thr
 65 70 75 80

Ile Gln Gly Leu Arg Ala Val Asp Thr Gly Leu Tyr Leu Cys Lys Val
 85 90 95

Glu Leu Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr
 100 105 110

Gln Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120 125

<210> SEQ ID NO 7
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 7

Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
 20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80

Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95

Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 8
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 8

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser His Gly Leu 1 5 10 15
Ala Ser Phe Pro Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 9
 <211> LENGTH: 358
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 9

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Ser His Asn Thr Asp Glu 20 25 30
Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val 35 40 45
Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp 50 55 60
Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile 65 70 75 80
Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu 85 90 95
Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln

-continued

100					105					110					
Ile	Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro
		115					120					125			
Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu
	130					135					140				
Leu	Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp
145					150					155					160
Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp
				165						170					175
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly
			180					185						190	
Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn
		195					200					205			
Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp
	210					215					220				
Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro
225					230					235					240
Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu
				245					250					255	
Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn
			260					265						270	
Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile
		275					280					285			
Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr
		290				295					300				
Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys
305					310					315					320
Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys
				325					330					335	
Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu
			340					345					350		
Ser	Leu	Ser	Pro	Gly	Lys										
		355													

<210> SEQ ID NO 10

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 10

Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
1				5					10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr	Glu	Val
			20					25					30		
Arg	Val	Thr	Val	Leu	Arg	Thr	Asn	Asp	Gln	Met	Val	Thr	Glu	Val	Cys
		35					40					45			
Ala	Ala	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp	Asp	Ser
		50				55					60				
Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	Gln	Val	Asn	Leu	Thr	Ile	Gln
65					70				75					80	
Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val	Glu	Leu
				85					90					95	
Met	Tyr	Pro	Pro	Pro	Tyr	Tyr	Leu	Gly	Ile	Gly	Asn	Gly	Thr	Gln	Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 11
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 11

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile

-continued

100				105				110							
Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro	Lys
		115					120					125			
Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu	Leu
	130					135					140				
Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
145					150					155					160
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val
				165					170					175	
Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val
			180					185					190		
Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser
		195					200					205			
Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu
	210					215					220				
Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala
225					230					235					240
Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro
				245					250					255	
Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln
			260					265					270		
Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala
		275					280					285			
Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
	290					295					300				
Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
305					310					315					320
Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser
				325					330					335	
Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser
			340				345						350		
Leu	Ser	Pro	Gly	Lys											
		355													

<210> SEQ ID NO 12

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 12

Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
1			5						10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr	Glu	Val
		20						25					30		
Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu	Val	Cys
		35				40						45			
Ala	Thr	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp	Asp	Ser
	50					55					60				
Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	Gln	Val	Asn	Leu	Thr	Ile	Gln
65					70				75					80	
Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val	Glu	Leu
			85					90						95	
Met	Tyr	Pro	Pro	Pro	Tyr	Tyr	Leu	Gly	Ile	Gly	Asn	Gly	Thr	Gln	Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 13
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 13

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Tyr Pro 50 55 60
Phe Cys Ser Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile

-continued

100				105				110							
Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro	Lys
		115					120					125			
Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu	Leu
	130					135					140				
Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
145					150					155					160
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val
				165					170					175	
Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val
			180					185					190		
Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser
		195					200					205			
Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu
	210					215					220				
Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala
225					230					235					240
Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro
				245					250					255	
Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln
			260					265					270		
Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala
		275					280					285			
Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
	290					295					300				
Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
305					310					315					320
Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser
				325					330					335	
Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser
			340				345						350		
Leu	Ser	Pro	Gly	Lys											
		355													

<210> SEQ ID NO 14

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 14

Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
1			5						10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr	Glu	Val
		20						25					30		
Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu	Val	Cys
		35				40						45			
Ala	Ala	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp	Asp	Ser
		50				55					60				
Ile	Cys	Thr	Gly	Thr	Phe	Asn	Glu	Ser	Arg	Val	Asn	Leu	Thr	Ile	Gln
65					70					75					80
Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val	Glu	Leu
				85					90					95	
Met	Tyr	Pro	Pro	Pro	Tyr	Tyr	Leu	Gly	Ile	Gly	Asn	Gly	Thr	Gln	Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 15
 <211> LENGTH: 358
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 15

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Val Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu 85 90 95
Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln

-continued

100				105				110							
Ile	Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro
		115					120					125			
Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu
	130					135					140				
Leu	Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp
145					150					155					160
Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp
				165					170					175	
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly
			180					185					190		
Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn
	195						200					205			
Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp
	210					215					220				
Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro
225					230					235					240
Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu
				245					250					255	
Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn
			260					265					270		
Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile
		275					280					285			
Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr
	290					295					300				
Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys
305					310					315					320
Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys
				325					330				335		
Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu
			340					345					350		
Ser	Leu	Ser	Pro	Gly	Lys										
	355														

<210> SEQ ID NO 16
 <211> LENGTH: 358
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 16

Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
1				5					10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr	Glu	Val
			20					25					30		
Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu	Val	Cys
		35				40						45			
Ala	Ala	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp	Asp	Ser
		50				55					60				
Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	Gln	Val	Asn	Leu	Thr	Ile	Gln
65					70				75					80	
Gly	Leu	Arg	Ala	Met	Met	Asp	Thr	Gly	Leu	Tyr	Leu	Cys	Lys	Val	Glu
				85					90					95	
Leu	Met	Tyr	Pro	Pro	Pro	Tyr	Tyr	Leu	Gly	Ile	Gly	Asn	Gly	Thr	Gln

-continued

100				105				110							
Ile	Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro
		115					120					125			
Lys	Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu
	130					135					140				
Leu	Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp
145					150					155					160
Thr	Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp
				165					170					175	
Val	Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly
			180					185					190		
Val	Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn
	195					200						205			
Ser	Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp
	210					215					220				
Leu	Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro
225					230					235					240
Ala	Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu
				245					250					255	
Pro	Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn
			260					265					270		
Gln	Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile
		275				280						285			
Ala	Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr
	290					295					300				
Thr	Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys
305					310					315					320
Leu	Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys
				325					330				335		
Ser	Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu
			340					345					350		
Ser	Leu	Ser	Pro	Gly	Lys										
	355														

<210> SEQ ID NO 17
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 17

Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
1				5					10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr	Glu	Val
			20					25					30		
Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu	Val	Cys
		35				40						45			
Ala	Ala	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp	Asp	Ser
		50				55					60				
Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	Gln	Val	Asn	Leu	Thr	Ile	Gln
65					70				75					80	
Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val	Glu	Leu
				85				90						95	
Met	Tyr	Pro	Pro	Pro	Tyr	Phe	Val	Gly	Met	Gly	Asn	Gly	Thr	Gln	Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 18
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 18

Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Phe Asn Glu Ser Arg Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 19
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 19

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 20
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 20

Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 21
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 21

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Tyr Glu Gly Ile Gly Asn Gly Thr Gln Ile

-continued

100					105					110					
Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro	Lys
		115					120					125			
Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu	Leu
		130					135					140			
Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
							150					155			160
Leu	Met	Ile	Ser	Arg	Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val
				165					170					175	
Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val
			180					185					190		
Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser
			195				200					205			
Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu
							215					220			
Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala
							230					235			240
Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro
				245					250					255	
Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln
			260					265					270		
Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala
			275				280					285			
Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
							295					300			
Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
				310					315					320	
Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser
				325					330					335	
Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser
			340				345						350		
Leu	Ser	Pro	Gly	Lys											
			355												

<210> SEQ ID NO 22

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 22

Ile	Gln	Val	Thr	Gln	Pro	Ser	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
1				5					10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Ala	Thr	Glu	Val
			20					25					30		
Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu	Val	Cys
			35				40						45		
Ala	Ala	Thr	Phe	Thr	Glu	Lys	Asn	Thr	Val	Gly	Phe	Leu	Asp	Asp	Ser
			50				55					60			
Ile	Cys	Thr	Gly	Thr	Phe	Asn	Glu	Ser	Arg	Val	Asn	Leu	Thr	Ile	Gln
65						70						75			80
Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val	Glu	Leu
				85					90					95	
Met	Tyr	Pro	Pro	Pro	Tyr	Phe	Val	Gly	Met	Gly	Asn	Gly	Thr	Gln	Ile

-continued

100					105					110					
Tyr	Val	Ile	Asp	Pro	Glu	Pro	Cys	Pro	Asp	Ser	Asp	Gln	Glu	Pro	Lys
		115					120					125			
Ser	Ser	Asp	Lys	Thr	His	Thr	Ser	Pro	Pro	Ser	Pro	Ala	Pro	Glu	Leu
		130					135					140			
Leu	Gly	Gly	Ser	Ser	Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr
		145					150					155			160
Leu	Tyr	Ile	Thr	Arg	Glu	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val
				165					170					175	
Ser	His	Glu	Asp	Pro	Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val
			180					185					190		
Glu	Val	His	Asn	Ala	Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser
		195					200					205			
Thr	Tyr	Arg	Val	Val	Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu
		210					215					220			
Asn	Gly	Lys	Glu	Tyr	Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala
		225					230					235			240
Pro	Ile	Glu	Lys	Thr	Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro
			245						250					255	
Gln	Val	Tyr	Thr	Leu	Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln
			260					265					270		
Val	Ser	Leu	Thr	Cys	Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala
		275					280					285			
Val	Glu	Trp	Glu	Ser	Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr
		290					295					300			
Pro	Pro	Val	Leu	Asp	Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu
		305					310					315			320
Thr	Val	Asp	Lys	Ser	Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser
			325						330					335	
Val	Met	His	Glu	Ala	Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser
			340				345						350		
Leu	Ser	Pro	Gly	Lys											
		355													

<210> SEQ ID NO 23

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 23

Met	His	Val	Ala	Gln	Pro	Ala	Val	Val	Leu	Ala	Ser	Ser	Arg	Gly	Ile
			5						10					15	
Ala	Ser	Phe	Val	Cys	Glu	Tyr	Ala	Ser	Pro	Gly	Lys	Tyr	Thr	Glu	Val
			20					25					30		
Arg	Val	Thr	Val	Leu	Arg	Gln	Ala	Asp	Ser	Gln	Val	Thr	Glu	Val	Cys
			35				40						45		
Ala	Ala	Thr	Tyr	Met	Met	Gly	Asn	Glu	Leu	Thr	Phe	Leu	Asp	Asp	Ser
			50				55					60			
Ile	Cys	Thr	Gly	Thr	Ser	Ser	Gly	Asn	Gln	Val	Asn	Leu	Thr	Ile	Gln
			65				70					75			80
Gly	Leu	Arg	Ala	Met	Asp	Thr	Gly	Leu	Tyr	Ile	Cys	Lys	Val	Glu	Leu
			85						90					95	
Met	Tyr	Pro	Pro	Pro	Tyr	Phe	Glu	Gly	Met	Gly	Asn	Gly	Thr	Gln	Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys 115 120 125		
Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu 130 135 140		
Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr 145 150 155 160		
Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val 165 170 175		
Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val 180 185 190		
Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser 195 200 205		
Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu 210 215 220		
Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala 225 230 235 240		
Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro 245 250 255		
Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln 260 265 270		
Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala 275 280 285		
Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr 290 295 300		
Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu 305 310 315 320		
Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser 325 330 335		
Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser 340 345 350		
Leu Ser Pro Gly Lys 355		

<210> SEQ ID NO 24
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 24

Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser Arg Gly Ile 1 5 10 15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val 20 25 30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys 35 40 45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser 50 55 60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln 65 70 75 80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu 85 90 95
Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile

-continued

100	105	110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp		
115	120	

<210> SEQ ID NO 25
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 25

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser His Gly Leu															
1			5					10						15	

Ala Ser Phe Pro Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val															
		20					25						30		

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys															
	35					40				45					

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser															
	50					55				60					

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln															
65				70				75						80	

Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu															
			85					90						95	

Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile															
		100					105					110			

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp															
	115						120								

<210> SEQ ID NO 26
 <211> LENGTH: 125
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 26

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile															
1			5					10						15	

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Ser His Asn Thr Asp Glu															
		20					25						30		

Val Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val															
	35					40				45					

Cys Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp															
	50					55				60					

Ser Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile															
65				70				75						80	

Gln Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu															
			85					90						95	

Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln															
		100					105					110			

Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp															
	115						120						125		

<210> SEQ ID NO 27
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 27

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1           5           10           15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
          20           25           30
Arg Val Thr Val Leu Arg Thr Asn Asp Gln Met Val Thr Glu Val Cys
          35           40           45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
          50           55           60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65           70           75           80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
          85           90           95
Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile
          100          105          110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
          115          120

```

<210> SEQ ID NO 28

<211> LENGTH: 124

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 28

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1           5           10           15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
          20           25           30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
          35           40           45
Ala Ala Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp Asp Ser
          50           55           60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65           70           75           80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
          85           90           95
Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile
          100          105          110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
          115          120

```

<210> SEQ ID NO 29

<211> LENGTH: 124

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 29

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1           5           10           15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
          20           25           30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys

```


-continued

Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln Ile
 100 105 110

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120

<210> SEQ ID NO 32
 <211> LENGTH: 125
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 32

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
 20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80

Gly Leu Arg Ala Met Val Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu
 85 90 95

Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln
 100 105 110

Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120 125

<210> SEQ ID NO 33
 <211> LENGTH: 125
 <212> TYPE: PRT
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 33

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
 20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80

Gly Leu Arg Ala Met Met Asp Thr Gly Leu Tyr Leu Cys Lys Val Glu
 85 90 95

Leu Met Tyr Pro Pro Pro Tyr Tyr Leu Gly Ile Gly Asn Gly Thr Gln
 100 105 110

Ile Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120 125

<210> SEQ ID NO 34
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 34

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1          5          10          15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
20          25          30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
35          40          45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
50          55          60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65          70          75          80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
85          90          95
Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile
100         105         110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
115         120

```

<210> SEQ ID NO 35

<211> LENGTH: 124

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 35

```

Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser Arg Gly Ile
1          5          10          15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
20          25          30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
35          40          45
Ala Ala Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp Asp Ser
50          55          60
Ile Cys Thr Gly Thr Phe Asn Glu Ser Arg Val Asn Leu Thr Ile Gln
65          70          75          80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
85          90          95
Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile
100         105         110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
115         120

```

<210> SEQ ID NO 36

<211> LENGTH: 124

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 36

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1          5          10          15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
20          25          30

```


-continued

Met Tyr Pro Pro Pro Tyr Tyr Glu Gly Ile Gly Asn Gly Thr Gln Ile
 100 105 110

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120

<210> SEQ ID NO 39
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 39

Ile Gln Val Thr Gln Pro Ser Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Ala Thr Glu Val
 20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45

Ala Ala Thr Phe Thr Glu Lys Asn Thr Val Gly Phe Leu Asp Asp Ser
 50 55 60

Ile Cys Thr Gly Thr Phe Asn Glu Ser Arg Val Asn Leu Thr Ile Gln
 65 70 75 80

Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95

Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile
 100 105 110

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120

<210> SEQ ID NO 40
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 40

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
 20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80

Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95

Met Tyr Pro Pro Pro Tyr Phe Glu Gly Met Gly Asn Gly Thr Gln Ile
 100 105 110

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
 115 120

<210> SEQ ID NO 41
 <211> LENGTH: 233
 <212> TYPE: PRT

-continued

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

Gln Glu Pro Lys Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro
 1 5 10 15
 Ala Pro Glu Leu Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys
 20 25 30
 Pro Lys Asp Thr Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val
 35 40 45
 Val Val Asp Val Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr
 50 55 60
 Val Asp Gly Val Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu
 65 70 75 80
 Gln Tyr Asn Ser Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His
 85 90 95
 Gln Asp Trp Leu Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys
 100 105 110
 Ala Leu Pro Ala Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln
 115 120 125
 Pro Arg Glu Pro Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu
 130 135 140
 Thr Lys Asn Gln Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro
 145 150 155 160
 Ser Asp Ile Ala Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn
 165 170 175
 Tyr Lys Thr Thr Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu
 180 185 190
 Tyr Ser Lys Leu Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val
 195 200 205
 Phe Ser Cys Ser Val Met His Glu Ala Leu His Asn His Tyr Thr Gln
 210 215 220
 Lys Ser Leu Ser Leu Ser Pro Gly Lys
 225 230

<210> SEQ ID NO 42

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 42

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15
 Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
 20 25 30
 Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45
 Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60
 Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80
 Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95
 Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile
 100 105 110

-continued

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys
 115 120 125
 Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu
 130 135 140
 Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
 145 150 155 160
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 165 170 175
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val
 180 185 190
 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser
 195 200 205
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 210 215 220
 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala
 225 230 235 240
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 245 250 255
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln
 260 265 270
 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 275 280 285
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 290 295 300
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 305 310 315 320
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 325 330 335
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 340 345 350
 Leu Ser Pro Gly Lys
 355

<210> SEQ ID NO 43
 <211> LENGTH: 357
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 43

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15
 Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
 20 25 30
 Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45
 Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60
 Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80
 Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95
 Met Tyr Pro Pro Pro Tyr Phe Glu Gly Ile Gly Asn Gly Thr Gln Ile
 100 105 110

-continued

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys
 115 120 125
 Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu
 130 135 140
 Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
 145 150 155 160
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 165 170 175
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val
 180 185 190
 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser
 195 200 205
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 210 215 220
 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala
 225 230 235 240
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 245 250 255
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln
 260 265 270
 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 275 280 285
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 290 295 300
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 305 310 315 320
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 325 330 335
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 340 345 350
 Leu Ser Pro Gly Lys
 355

<210> SEQ ID NO 44

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 44

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15
 Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
 20 25 30
 Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45
 Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60
 Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80
 Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95
 Met Tyr Pro Pro Pro Tyr Phe Asp Gly Ile Gly Asn Gly Thr Gln Ile
 100 105 110

-continued

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys
 115 120 125
 Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu
 130 135 140
 Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
 145 150 155 160
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 165 170 175
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val
 180 185 190
 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser
 195 200 205
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 210 215 220
 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala
 225 230 235 240
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 245 250 255
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln
 260 265 270
 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 275 280 285
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 290 295 300
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 305 310 315 320
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 325 330 335
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 340 345 350
 Leu Ser Pro Gly Lys
 355

<210> SEQ ID NO 45

<211> LENGTH: 357

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Fusion protein

<400> SEQUENCE: 45

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15
 Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
 20 25 30
 Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45
 Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60
 Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80
 Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95
 Met Tyr Pro Pro Pro Tyr Phe Val Gly Ile Gly Asn Gly Thr Gln Ile
 100 105 110

-continued

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp Gln Glu Pro Lys
 115 120 125
 Ser Ser Asp Lys Thr His Thr Ser Pro Pro Ser Pro Ala Pro Glu Leu
 130 135 140
 Leu Gly Gly Ser Ser Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr
 145 150 155 160
 Leu Met Ile Ser Arg Thr Pro Glu Val Thr Cys Val Val Val Asp Val
 165 170 175
 Ser His Glu Asp Pro Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val
 180 185 190
 Glu Val His Asn Ala Lys Thr Lys Pro Arg Glu Glu Gln Tyr Asn Ser
 195 200 205
 Thr Tyr Arg Val Val Ser Val Leu Thr Val Leu His Gln Asp Trp Leu
 210 215 220
 Asn Gly Lys Glu Tyr Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala
 225 230 235 240
 Pro Ile Glu Lys Thr Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro
 245 250 255
 Gln Val Tyr Thr Leu Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln
 260 265 270
 Val Ser Leu Thr Cys Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala
 275 280 285
 Val Glu Trp Glu Ser Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr
 290 295 300
 Pro Pro Val Leu Asp Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu
 305 310 315 320
 Thr Val Asp Lys Ser Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser
 325 330 335
 Val Met His Glu Ala Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser
 340 345 350
 Leu Ser Pro Gly Lys
 355

<210> SEQ ID NO 46
 <211> LENGTH: 124
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 46

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
 1 5 10 15
 Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
 20 25 30
 Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
 35 40 45
 Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
 50 55 60
 Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
 65 70 75 80
 Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
 85 90 95
 Met Tyr Pro Pro Pro Tyr Phe Val Gly Met Gly Asn Gly Thr Gln Ile
 100 105 110

-continued

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
115 120

<210> SEQ ID NO 47
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 47

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
35 40 45

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
50 55 60

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65 70 75 80

Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
85 90 95

Met Tyr Pro Pro Pro Tyr Phe Glu Gly Ile Gly Asn Gly Thr Gln Ile
100 105 110

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
115 120

<210> SEQ ID NO 48
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutant

<400> SEQUENCE: 48

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1 5 10 15

Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
20 25 30

Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
35 40 45

Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
50 55 60

Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65 70 75 80

Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
85 90 95

Met Tyr Pro Pro Pro Tyr Phe Asp Gly Ile Gly Asn Gly Thr Gln Ile
100 105 110

Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
115 120

<210> SEQ ID NO 49
<211> LENGTH: 124
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Mutant

-continued

<400> SEQUENCE: 49

```

Met His Val Ala Gln Pro Ala Val Val Leu Ala Ser Ser Arg Gly Ile
1          5          10          15
Ala Ser Phe Val Cys Glu Tyr Ala Ser Pro Gly Lys Tyr Thr Glu Val
          20          25          30
Arg Val Thr Val Leu Arg Gln Ala Asp Ser Gln Val Thr Glu Val Cys
          35          40          45
Ala Ala Thr Tyr Met Met Gly Asn Glu Leu Thr Phe Leu Asp Asp Ser
          50          55          60
Ile Cys Thr Gly Thr Ser Ser Gly Asn Gln Val Asn Leu Thr Ile Gln
65          70          75          80
Gly Leu Arg Ala Met Asp Thr Gly Leu Tyr Ile Cys Lys Val Glu Leu
          85          90          95
Met Tyr Pro Pro Pro Tyr Phe Val Gly Ile Gly Asn Gly Thr Gln Ile
          100          105          110
Tyr Val Ile Asp Pro Glu Pro Cys Pro Asp Ser Asp
          115          120

```

<210> SEQ ID NO 50

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 50

```

Met Tyr Pro Pro Pro Tyr
1          5

```

The invention claimed is:

1. A CTLA-4 protein comprising an extracellular domain of CTLA-4 comprising the sequence set forth in any one of SEQ ID NOS: 34, 36, 39, 40, and 46-48.

2. The CTLA-4 protein of claim 1, wherein the CTLA-4 protein has the ability to bind at least one of B7-1 and B7-2.

3. The CTLA-4 protein of claim 2 wherein the CTLA-4 protein does not block the anti-cancer immunotherapeutic effects of an anti-CTLA-4 antibody.

4. The CTLA-4 protein of claim 3, wherein the anti-CTLA-4 antibody is Ipilimumab.

5. The CTLA-4 protein of claim 1, wherein the extracellular domain of CTLA-4 is fused to a Fc region of a human immunoglobulin protein.

6. The CTLA-4 protein of claim 5, wherein the sequence of the CTLA-4 protein comprises the sequence set forth in any one of SEQ ID NOS: 17, 19, 22, 23, and 42-44.

7. A pharmaceutical composition comprising a therapeutically effective amount of the CTLA-4 protein of claim 1, and a physiologically acceptable carrier or excipient.

8. A method of treating immune-related adverse events associated with anti-CTLA-4 immunotherapy in a subject in need thereof, comprising administering to the subject the CTLA-4 protein of claim 1.

9. The method of claim 8, wherein the anti-CTLA-4 immunotherapy is an anti-CTLA-4 antibody.

35 10. The method of claim 9, wherein the subject has cancer.

11. The method of claim 8, wherein the extracellular domain of CTLA-4 is fused to a Fc region of a human immunoglobulin protein.

40 12. The method of claim 11, wherein the sequence of the CTLA-4 protein comprises the sequence set forth in any one of SEQ ID NOS: 17, 19, 22, 23, and 42-44.

13. The method of claim 12, where the CTLA-4 protein comprises the sequence set forth in SEQ ID NO: 17.

45 14. The method of claim 12, where the CTLA-4 protein comprises the sequence set forth in SEQ ID NO: 23.

15. A method of minimizing autoimmune adverse events associated with cancer immunotherapy in a subject in need thereof, comprising administering to the subject the CTLA-4 protein of claim 1.

50 16. The method of claim 15, wherein the extracellular domain of CTLA-4 is fused to a Fc region of a human immunoglobulin protein.

17. The method of claim 16, wherein the sequence of the CTLA-4 protein comprises the sequence set forth in any one of SEQ ID NOS: 17, 19, 22, 23, and 42-44.

55 18. The method of claim 17, wherein the CTLA-4 protein comprises the sequence set forth in SEQ ID NO: 17.

19. The method of claim 17, wherein the CTLA-4 protein comprises the sequence set forth in SEQ ID NO: 23.

* * * * *