

US011517631B2

(12) **United States Patent**
Nathwani et al.

(10) **Patent No.:** **US 11,517,631 B2**
(45) **Date of Patent:** ***Dec. 6, 2022**

(54) **FACTOR IX ENCODING NUCLEOTIDES**

(71) Applicant: **UCL Business LTD**, London (GB)

(72) Inventors: **Amit Nathwani**, Rickmansworth (GB);
Jenny McIntosh, Surrey (GB)

(73) Assignee: **UCL BUSINESS LTD**, London (GB)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 688 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/398,840**

(22) Filed: **Apr. 30, 2019**

(65) **Prior Publication Data**

US 2020/0054768 A1 Feb. 20, 2020

Related U.S. Application Data

(62) Division of application No. 16/105,583, filed on Aug. 20, 2018, now Pat. No. 10,842,885.

(51) **Int. Cl.**

A61K 48/00 (2006.01)

C07K 14/81 (2006.01)

A61P 7/00 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 48/0066** (2013.01); **A61K 48/0075** (2013.01); **A61K 48/0091** (2013.01); **A61P 7/00** (2018.01); **C07K 14/8107** (2013.01); **C12Y 304/21022** (2013.01)

(58) **Field of Classification Search**

CPC **A61K 48/0066**; **A61K 48/0075**; **A61K 48/0091**; **A61P 7/00**; **C07K 14/8107**; **C12Y 304/21022**

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

6,271,025 B1 8/2001 Negrier et al.
6,419,921 B1 7/2002 Négrier et al.
6,723,551 B2 4/2004 Kotin et al.
6,800,461 B2 10/2004 Negrier et al.
6,884,616 B1 4/2005 Negrier et al.
7,351,813 B2 4/2008 Miao
8,030,065 B2 10/2011 Gray
8,168,425 B2 5/2012 Gray
8,198,421 B2 6/2012 Samulski
9,764,045 B2 9/2017 Nathwani et al.
2002/0086427 A1 7/2002 Leiden et al.
2003/0022378 A1 1/2003 Ehrhardt et al.
2003/0130221 A1 7/2003 High et al.
2003/0148506 A1 8/2003 Kotin et al.
2003/0186291 A1 10/2003 Faust et al.
2004/0053870 A1 3/2004 Yew et al.
2007/0003521 A1 1/2007 Yew
2007/0180546 A1 8/2007 Rapp et al.
2008/0102115 A1 5/2008 Oyhenart et al.
2008/0167219 A1 7/2008 Lin et al.
2008/0269125 A1 10/2008 Ballance et al.

2008/0305991 A1 12/2008 Defrees et al.
2011/0070241 A1 3/2011 Yang
2012/0009222 A1 1/2012 Nguyen et al.
2013/0236974 A1 9/2013 de Fougerolles
2014/0271550 A1 9/2014 Rabinowitz et al.
2016/0122739 A1 5/2016 Sheehan et al.
2016/0222414 A1 8/2016 Rabinowitz et al.
2016/0361427 A1 12/2016 Defrees et al.
2016/0375110 A1 12/2016 High et al.
2017/0136104 A1 5/2017 Defrees et al.

FOREIGN PATENT DOCUMENTS

CN 106497949 A 3/2017
EP 1010762 A1 6/2000
EP 1026250 A1 8/2000
EP 1038959 A1 9/2000
EP 1048726 A2 11/2000
EP 1048736 A1 11/2000
EP 1048735 B1 9/2006
EP 2067488 A1 6/2009
EP 2149603 A1 2/2010
EP 2492347 A1 8/2012
EP 2216409 B1 12/2014
WO WO-9417810 A1 8/1994

(Continued)

OTHER PUBLICATIONS

Monahan et al entitled "Employing a Gain-of Function Factor IX Variant R338L to Advance the Efficacy and Safety of Hemophilia B Human Gene Therapy: Preclinical Evaluation Supporting an Ongoing Adeno-Associated Virus Clinical Trial" (Human Gene Therapy Feb. 2015 vol. 26: pp. 69-81). (Year: 2015).*

EP19762195.6 Examination Report dated May 3, 2021.

"Satya et al., 2003 "A Pattern Matching Algorithm for Codon Optimization and CpG Motif-Engineering in DNA Expression Vectors" Proceeding IEEE Computer Society Bioinformatic Conference, 2003 pp. 294-305 vol. 2".

"Bauer et al., 2010 "The impact of intragenic CpG content on gene expression" Nucleic Acids Research, Jul. 2010, vol. 38, pp. 3891-3908".

(Continued)

Primary Examiner — Catherine S Hibbert

(74) *Attorney, Agent, or Firm* — Wilson Sonsini Goodrich & Rosati

(57) **ABSTRACT**

The present invention relates to polynucleotides comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or fragment thereof and wherein a portion of the coding sequence is not wild type. The present invention further relates to viral particles comprising a recombinant genome comprising the polynucleotide of the invention, compositions comprising the polynucleotides or viral particles, and methods and uses of the polynucleotides, viral particles or compositions.

14 Claims, 13 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO WO-9742900 A1 11/1997
 WO WO-9841240 A1 9/1998
 WO WO-9903496 A1 1/1999
 WO WO-9949803 A1 10/1999
 WO WO-9949880 A1 10/1999
 WO WO-0014262 A2 3/2000
 WO WO-0049147 A1 8/2000
 WO WO-0054787 A1 9/2000
 WO WO-0136620 A2 5/2001
 WO WO-0166149 A2 9/2001
 WO WO-0170763 A1 9/2001
 WO WO-0175092 A2 10/2001
 WO WO-0177137 A1 10/2001
 WO WO-0179271 A1 10/2001
 WO WO-0170968 A3 12/2001
 WO WO-0198482 A2 12/2001
 WO WO-0234296 A1 5/2002
 WO WO-0240544 A2 5/2002
 WO WO-02062376 A1 8/2002
 WO WO-02062377 A2 8/2002
 WO WO-02064799 A2 8/2002
 WO WO-02071843 A1 9/2002
 WO WO-02079447 A2 10/2002
 WO WO-02099105 A2 12/2002
 WO WO-03020764 A2 3/2003
 WO WO-03025146 A2 3/2003
 WO WO-03048364 A2 6/2003
 WO WO-2004027019 A2 4/2004
 WO WO-2004080162 A2 9/2004
 WO WO-2004092351 A2 10/2004
 WO WO-2005040215 A2 5/2005
 WO WO-2005058283 A2 6/2005
 WO WO-2005084430 A1 9/2005
 WO WO-2004098532 A3 1/2006
 WO WO-2006015789 A2 2/2006
 WO WO-2006018204 A1 2/2006
 WO WO-2006026238 A2 3/2006
 WO WO-2006036502 A2 4/2006
 WO WO-2006093847 A1 9/2006
 WO WO-2006103258 A1 10/2006
 WO WO-2006127896 A2 11/2006
 WO WO-2006110689 A3 4/2007
 WO WO-2007036233 A2 4/2007
 WO WO-2007046703 A2 4/2007
 WO WO-2007047706 A2 4/2007
 WO WO-2007120533 A2 10/2007
 WO WO-2007130453 A2 11/2007
 WO WO-2007135182 A2 11/2007
 WO WO-2007148971 A2 12/2007
 WO WO-2007149406 A2 12/2007
 WO WO-2007149852 A2 12/2007
 WO WO-2008091311 A1 7/2008
 WO WO-2008092643 A2 8/2008
 WO WO-2008092644 A2 8/2008
 WO WO-2008124724 A1 10/2008
 WO WO-2008153366 A2 12/2008
 WO WO-2009014445 A2 1/2009
 WO WO-2009026393 A2 2/2009
 WO WO-2009038462 A1 3/2009
 WO WO-2009051717 A2 4/2009
 WO WO-2009059056 A2 5/2009
 WO WO-2009061369 A2 5/2009
 WO WO-2009102085 A1 8/2009
 WO WO-2009130198 A2 10/2009
 WO WO-2009137254 A2 11/2009
 WO WO-2009140015 A2 11/2009
 WO WO-2008118507 A3 12/2009
 WO WO-2010012451 A1 2/2010
 WO WO-2010029178 A1 3/2010
 WO WO-2010055413 A1 5/2010
 WO WO-2011005968 A1 1/2011
 WO WO-2011011841 A1 2/2011
 WO WO-2011014890 A1 2/2011
 WO WO-2011054994 A1 5/2011

WO WO-2011122950 A1 10/2011
 WO WO-2011154520 A1 12/2011
 WO WO-2012061654 A1 5/2012
 WO WO-2012135805 A2 10/2012
 WO WO-2013078400 A1 5/2013
 WO WO-2013090648 A1 6/2013
 WO WO-2013173512 A2 11/2013
 WO WO-2012170930 A9 1/2014
 WO WO-2014016580 A1 1/2014
 WO WO-2014063108 A1 4/2014
 WO WO-2014063753 A1 5/2014
 WO WO-2014064277 A1 5/2014
 WO WO-2014070349 A1 5/2014
 WO WO-2014081831 A1 5/2014
 WO WO-2014152940 A1 9/2014
 WO WO-2014193716 A2 12/2014
 WO WO-2015012924 A2 1/2015
 WO WO-2015013313 A2 1/2015
 WO WO-2015038625 A1 3/2015
 WO WO-2015073988 A1 5/2015
 WO WO-2014182684 A3 6/2015
 WO WO-2015085276 A1 6/2015
 WO WO-2015086406 A2 6/2015
 WO WO-2015139093 A1 9/2015
 WO WO-2015162302 A2 10/2015
 WO WO-2016004113 A1 1/2016
 WO WO-2016028872 A2 2/2016
 WO WO-2016041588 A1 3/2016
 WO WO-2016073837 A1 5/2016
 WO WO-2016075473 A2 5/2016
 WO WO-2016123200 A1 8/2016
 WO WO-2016127057 A1 8/2016
 WO WO-2016146757 A1 9/2016
 WO WO-2016168728 A3 11/2016
 WO WO-2016179644 A1 11/2016
 WO WO-2016181122 A1 11/2016
 WO WO-2016181123 A1 11/2016
 WO WO-2016210170 A1 12/2016
 WO WO-2017011519 A1 1/2017
 WO WO-2017021359 A1 2/2017
 WO WO-2017024060 A1 2/2017
 WO WO-2017070167 A1 4/2017
 WO WO-2017075619 A1 5/2017
 WO WO-2017083762 A1 5/2017
 WO WO-2017083764 A1 5/2017
 WO WO-2017093482 A1 6/2017
 WO WO-2017096039 A1 6/2017
 WO WO-2017180861 A1 10/2017
 WO WO-2017191274 A3 2/2018
 WO WO-2018160686 A1 9/2018
 WO WO-2018176027 A1 9/2018
 WO WO-2018199214 A1 11/2018
 WO WO-2018206168 A1 11/2018
 WO WO-2018213786 A1 11/2018
 WO WO-2018217731 A1 11/2018
 WO WO-2018222792 A1 12/2018
 WO WO-2018222890 A1 12/2018
 WO WO-2018226887 A1 12/2018
 WO WO-2019011893 A1 1/2019
 WO WO-2019032898 A1 2/2019
 WO WO-2018022844 A3 4/2019
 WO WO-2019067766 A1 4/2019
 WO WO-2019070888 A1 4/2019
 WO WO-2019079215 A1 4/2019
 WO WO-2019094521 A1 5/2019
 WO WO-2019153009 A1 8/2019
 WO WO-2019154939 A1 8/2019
 WO WO-2019210267 A2 10/2019
 WO WO-2019219649 A1 11/2019
 WO WO-2019222132 A1 11/2019
 WO WO-2019241486 A1 12/2019
 WO WO-2020033863 A1 2/2020
 WO WO-2020039183 A1 2/2020

OTHER PUBLICATIONS

“Binny et al., 2012 “Vector Systems for Prenatal Gene Therapy: Principles of Adeno-Associated Virus Vector Design and Produc-

(56)

References Cited

OTHER PUBLICATIONS

- tion" *Methods in Molecular Biology*, Apr. 25, 2012 pp. 109-131 vol. 891".
- Chen, Zhi-Ying, et al., "Improved Production and Purification of Minicircle DNA Vector Free of Plasmid Bacterial Sequences and Capable of Persistent Transgene Expression in Vivo", *Human Gene Therapy*, 16(1):126-131 (2005).
- "Database UniProt [Online] May 10, 2017 (May 10, 2017), "SubName:Full=coagulation factor IX isoform X2 {ECO:00003131RefSeq:XP008059975.1}"; XP002796360, retrieved from EBI accession No. UNIPROT:A0A1U7TQC."
- Faust, Susan M., et al., "CpG-Depleted Adeno-Associated Virus Vectors Evade Immune Detection" *J. Clin Invest* (2013) 123(7):2994-3001.
- Faust, Susan M., et al., "Escaping Immune Activation Through the Use of CpG-Depleted AAV Vectors" *Molecular Therapy* vol. 21, Supplement 1, May 2013.
- "Frumkin et al., 2018 "Codon usage of highly expressed genes affects proteome-wide translation efficiency" *PNAS*, May 7, 2018 pp. E4940-E4949 vol. 115 No. 21".
- Hodges BL, et al., "Long-Term Transgene Expression From Plasmid DNA Gene Therapy Vectors is Negatively Affected by CpG Dinucleotides", *Molecular Therapy*. 10(2):269-278 (2004).
- Hyde, Stephen C., et al., "CpG-Free Plasmids Confer Reduced Inflammation and Sustained Pulmonary Gene Expression", *Nature Biology* (2008) 26: 549-551.
- Inouye et al. "Protein expression of preferred human codon-optimized *Gussia luciferase* genes with an artificial open reading frame in mammalian and bacterial cells." *Protein Expression and Purification* 128 (2016) 93-100.
- "Kao C Y et al., 2013 "Incorporation of the factor IX Padua mutation into FIX-Triple improves clotting activity in vitro and in vivo" *Thrombosis and Haemostasis*, May 16, 2013 pp. 244-256 vol. 110 No. 2".
- "Krinner et al., 2014 "CpG domains downstream of TSSs promote high levels of gene expression" *Nucleic Acids Research*, Apr. 2014, vol. 42, pp. 3551-3564".
- Mauro "Codon Optimization in the Production of Recombinant Biotherapeutics: Potential Risks and Considerations" *BioDrugs* (2018) 32:69-81.
- "Nair et al., 2011 "Effect of different UCOE-promoter combinations in creation of engineered cell lines for the production of Factor VIII" *BMC Research Notes*, Jun. 10, 2011 vol. 4 Art. 178".
- "Pierce & Iorio 2018 "Past, present and future of haemophilia gene therapy: From vectors and transgenes to known and unknown outcomes" *Haemophilia*, May 21, 2018 pp. 60-67 vol. 24 Suppl. 6".
- "Terry 2016 "Shire Kills Baxalta's Hemophilia B Program; Clears Path for BioMarin, Spark Therapeutics and uniQure." *BioSpace*, Aug. 4, 2016 <https://www.biospace.com/article/shire-kills-baxaltas-hemophilia-b-program-clears-path-for-biomarin-spark-therapeutics-and-uniquire/>".
- "Wang et al., 1997 "A factor IX-deficient Mouse Model of hemophilia B gene therapy". *Proceeding of the National Academy of Sciences of the United States of America*, 1997 pp. 11563-11566 vol. 94".
- "Yew et al., 2002 "CpG-depleted Plasmid DNA Vectors With Enhanced Safety and Long-Term Gene Expression in Vivo" *Molecular Therapy*, Jun. 2002 pp. 731-738 vol. 5 No. 6".
- Allan et al. Evolutionary Duplication of a Hepatic Control Region in the Human Apolipoprotein E Gene Locus. *The Journal of Biological Chemistry*, vol. 270(44):26278-26281, (Nov. 3, 1995).
- Anson et al. The gene structure of human anti-haemophilic factor IX. *The EMBO Journal* 3(5):1053-1060 (1984).
- Arruda et al. Emerging therapies for haemophilia: controversies and unanswered questions [version 1; referees: 4 approved]. *F1000Research* 2018, 7(F1000 Faculty Rev):489 Last updated: Apr. 24, 2018.
- Arruda et al. Safety and efficacy of factor IX gene transfer to skeletal muscle in murine and canine hemophilia B models by adeno-associated viral vector serotype 1. *Blood* 103(1):85-92 (Jan. 1, 2004).
- Bantel-Schaal, et al., Human Adeno-Associated Virus Type 5 is Only Distantly Related to Other Known Primate Helper-Dependent Parvoviruses, *Journal of Virology*, (1999) vol. 73 (2), pp. 939-947.
- Berns, Parvoviridae: The Viruses and Their Replication, *Fields Virology*; Feb. 7, 2018; pp. 2173-2197.
- Cantore, et al., Hyperfunctional coagulation factor IX improves the efficacy of gene therapy in hemophilic mice, *Blood*, (2012), vol. 120, pp. 4517-4520.
- Chang et al. Changing Residue 338 in Human Factor IX from Arginine to Alanine Causes an Increase in Catalytic Activity. *The Journal of Biological Chemistry*, 273(20):12089-12094 (May 15, 1998).
- Chiorini J.A. et al. Cloning of adeno-associated virus type 4 (AAV4) and generation of recombinant AAV4 particles. *Journal of Virology*, 71:6823-6833 (1997).
- Chiorini, J.A. et al. Cloning and characterization of adeno-associated virus type 5, *Journal of Virology* 73:1309-1319 (1999).
- Dang, et al. In Vivo Footprinting Analysis of the Hepatic Control Region of the Human Apolipoprotein E/C-I/C-IV/C-II Gene Locus. *The Journal of Biological Chemistry*, 271(45):28667-28676 (Nov. 8, 1996).
- Dang et al. Structure of the Hepatic Control Region of the Human Apolipoprotein E/C-I Gene Locus. *The Journal of Biological Chemistry*, 270(38):22577-22585 (Sep. 22, 1995).
- Enjolras et al. The Three In-frame ATG, Clustered in the Translation Initiation Sequence of Human Factor IX Gene, Are Required for an Optimal Protein Production. *Thrombosis and Haemostasis*, Schattauer GMBH Germany, 82(4):1264-1269 (Oct. 1, 1999).
- Fagone, et al., Systemic Errors in Quantitative Polymerase Chain Reaction Titration of Self-Complementary Adeno-Associated Viral Vectors and Improved Alternative Methods, *Hum Gene Ther Methods*. Feb. 23, 2012 (1):1-7.
- George et al. Hemophilia B Gene Therapy with a High-Specificity Activity Factor IX Variant. *The New England Journal of Medicine*, 377(23):2215-2227 (Dec. 7, 2017).
- Giannelli et al. Haemophilia B: database of point mutations and short additions and deletions, Oxford University Press, *Nucleic Acids Research*, 18(14):4053-4059 (1990).
- Haas et al. Codon usage limitation in the expression of HIV-1 envelope glycoprotein, *Current Biology* 6(3):315-324 (1996).
- Hafenrichter et al. Quantitative Evaluation of Liver-Specific Promoters From Retroviral Vectors After in Vivo Transduction of Hepatocytes, *Blood*, 84(10):3394-3404 (Nov. 15, 1994).
- Ketterling et al. The Rates of G:C->T:A and G:C->C:G Transversions at CpG Dinucleotides in the Human Factor IX Gene, *Am. J. Hum. Genet.* 54:831-835(1994).
- Kurachi et al. Role of Intron I in Expression of the Human Factor IX Gene, *Journal of Biological Chemistry, American Society for Biochemistry and Molecular Biology, US*, 270(10):5276-5281 (Mar. 10, 1995).
- Lu et al. Gene therapy for hemophilia B mediated by recombinant adeno-associated viral vector with hFIXR338A, a high catalytic activity mutation of human coagulation factor IX, *Science in China (Series C)*, 44(6):585-592(Dec. 2001).
- Mathur et al. Protease and EGF1 Domains of Factor IXa Play Distinct Roles in Binding to Factor VIIIa, *The Journal of Biological Chemistry*, 274(26):18477-18486 (Jun. 25, 1999).
- McIntosh J. et al. Therapeutic levels of FVIII following a single peripheral vein administration of rAAV vector encoding a novel human factor VIII variant. *Blood* 121(17):3335-44 (Apr. 25, 1993).
- Miao, C.H. et al. Inclusion of the hepatic locus control region, an intron, and untranslated region increases and stabilizes hepatic factor IX gene expression in vivo but not in vitro. *Mol Ther.* 1(6):522-32(Jun. 2000).
- Nathwani, A.C. et al. Self-complementary adeno-associated virus vectors containing a novel liver-specific human factor IX expression cassette enable highly efficient transduction of murine and nonhuman primate liver. *Blood*, 107(7): 2653-2661 (Apr. 1, 2006).
- Nathwani et al. Adenovirus-Associated Virus Vector-Mediated Gene Transfer in Hemophilia B. *New England Journal of Medicine*, 365(25):2357-65 (Dec. 22, 2011).

(56)

References Cited

OTHER PUBLICATIONS

- Nathwani et al. Our Journey to Successful Gene Therapy for Hemophilia B, *Human Gene Therapy*, 25(11):923-926 (Nov. 1, 2014).
- Nathwani, et al., Long-term Safety and Efficacy Following Systemic Administration of a Self-complementary AAV Vector Encoding Human FIX Pseudotyped With Serotype 5 and 8 Capsid Proteins, *Mol Ther.* May 2011, vol. 19. (5), pp. 876-885.
- Needleman, S.B. et al. A general method applicable to the search for similarities in the amino acid sequence of two proteins. *Journal Molecular Biology* 48(3):443-53 (Mar. 1970).
- Okuyama T. et al. Liver-directed gene therapy: a retroviral vector with a complete LTR and the ApoE enhancer-alpha 1-antitrypsin promoter dramatically increases expression of human alpha 1-antitrypsin in vivo. *Human Gene Therapy* 7(5):637-45 (Mar. 20, 1996).
- Altschul, et al., Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, *Nucleic Acids Research*, (1997) vol. 25(17), pp. 3389-3402.
- Altschul, et al, Basic Local Alignment Search Tool, *J. Mol. Biol.* 215 (1990), pp. 403-410.
- Paul E. Monahan et al., "Employing a Gain-of-Function Factor IX Variant R338L to Advance the Efficacy and Safety of Hemophilia B Human Gene Therapy: Preclinical Evaluation Supporting an Ongoing Adeno-Associated Virus Clinical Trial", *Human Gene Therapy* 26:69-81 (Feb. 2015), DOI: 10.1089/hum.2014.106.
- Bryan et al., 2013, <http://www.elsevierblogs.com/currentcomments/?p=962>, Implications of protein fold switching, pp. 1-4.
- Dunbar et al., 2018, *Science*, vol. 359, ean4672, pp. 1-10.
- Lenzi et al., 2014, NCBI Bookshelf, a Service of the National Library of Medicine, National Institute of Health, Oversight and Review of Clinical Gene Transfer Protocols: Assessing the Role of the Recombinant DNA Advisory Committee. Washington (DC): National Academies Press (US), pp. 1-16.
- Shim et al., 2017, *Current Gene Therapy*, vol. 17, No. 5, pp. 1-18.
- Suwanmanee et al., 2014, *Molecular Therapy*, Integration-deficient lentiviral vectors expressing codon-optimized R338L human FIX restore normal hemostasis in Hemophilia B mice, vol. 22, No. 3, pp. 567-574.
- Rodriguez et al. Biosynthesis of FVIII in megakaryocytic cells: improved production and biochemical characterization, Blackwell Publishing Ltd, *British Journal of Haematology*, 127:568-575 (2004).
- Rutledge, E.A. et al. Infectious clones and vectors derived from adeno-associated virus (AAV) serotypes other than AAV type 2. *Journal of Virology* 72:309-319 (1998).
- Sabatino et al. Novel hemophilia B mouse models exhibiting a range of mutations in the Factor IX gene, *Blood*, 104(9): 2767-2774 (Nov. 1, 2004).
- Schuettrumpf et al. Factor IX variants improve gene therapy efficacy for hemophilia B, *Blood*, 105(6):2316-2323 (Mar. 15, 2005).
- Simioni et al. Evidence of the first X-linked thrombophilia due to a novel mutation in clotting factor IX gene resulting in hyperfunctional fix: factor IX arginine 338 leucine (factor IX padua), 2009 the Authors. *Journal Compilation. International Society on Thrombosis and Haemostasis* 7 (Suppl. 2). Abstract PL-TU-0004.
- Simioni et al. X-Linked Thrombophilia with a Mutant Factor IX (Factor IX Padua). *New England Journal of Medicine* 361:1671-5(2009).
- Kaur et al., 2009, *Current Gene Therapy*, vol. 9, pp. 434-458.
- Maqbool et al., 2015, *Biochemical Society Transactions*, vol. 43, No. 5, pp. 1011-1017.
- Plantier et al., 2001, *Thromb Haemost*, a factor VIII minigene comprising the truncated intron I of factor IX highly improves the in vitro production of factor VIII, 86(2), pp. 596-603.
- Srivastava, A. et al. Nucleotide sequence and organization of the adeno-associated virus 2 genome, *Journal of Virology* 45(2): 555-64(Feb. 1983).
- Wang et al. Sustained correction of bleeding disorder in hemophilia B mice by gene therapy. *Proc Natl Acad. Sci. USA* 96(7): 3906-3910 (Mar. 30, 1999).
- Welch et al. You're one in a googol: optimizing genes for protein expression, *J.R. Soc. Interface* 6(Suppl. 4):S467-76(Aug. 9, 2009). E-published: Mar. 11, 2009.
- Wooddell et al. Sustained liver-specific transgene expression from the albumin promoter in mice following hydrodynamic plasmid DNA delivery. *The Journal of Gene Medicine*, 10: 551-563 (2008).
- Wu, et al., Optimization of Self-complementary AAV Vectors for Liver-directed Expression Results in Sustained Correction of Hemophilia B at Low Vector Dose, *Mol Ther.* Feb. 2008; vol. 16(2), pp. 280-289.
- Wu, P. et al. Mutational analysis of the adeno-associated virus type 2 (AAV2) capsid gene and construction of AAV2 vectors with altered tropism. *Journal of Virology*, 74(18):8635-47 (Sep. 2000).
- Yan et al. Transgenic Mice Can Express Mutant Human Coagulation Factor IX with Higher Level of Clotting Activity. *Biochemical Genetics*, 44(7/8):349-360 (Aug. 2006).

* cited by examiner




FIG. 1A

ssHLP2.T1-ACNP-FIX-GoF (HTAG)




FIG. 1B

ssHLP2.T1-codop-FIX-GoF (HTFG)




FIG. 1C

ssLP1-FIXco (FIXco)

FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E






FIG. 2F










FIG. 5






FIG. 7A




FIG. 7B





FIG. 7C








FIG. 9




FIG. 10




FIG. 11A




FIG. 11B




FIG. 12





FIG. 13



1**FACTOR IX ENCODING NUCLEOTIDES****CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a divisional of U.S. application Ser. No. 16/105,583, filed Aug. 20, 2018, which is herein incorporated by reference in its entirety.

FIELD

The present invention relates to polynucleotides comprising a nucleotide sequence encoding Factor IX, viral particles comprising the polynucleotides and treatments utilising the polynucleotides.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Aug. 20, 2018, is named 52186-706.201 SL.txt and is 79,915 bytes in size.

BACKGROUND

Haemophilia B, an X-linked life threatening bleeding disorder affects 1:30,000 males. Current treatment involves frequent intravenous injections (2-3 times per week) of Factor IX (FIX) protein. This treatment is highly effective at arresting bleeding but it is not curative and is extremely expensive (£150,000/patient/year), thus making it unaffordable by the majority of haemophilia B patients in the world. Gene therapy for haemophilia B offers the potential for a cure through persistent, endogenous production of Factor IX following the transfer of a functioning copy of the Factor IX gene to an affected patient.

The present application relates to a gene therapy approach for treating haemophilia B, involving administering a vector comprising a polynucleotide encoding Factor IX. Such a gene therapy approach would avoid the need for frequent intravenous injections of Factor IX. However, it is difficult to provide an effective gene therapy vector, i.e. one that allows for a high level of Factor IX expression and of the expression of Factor IX which is highly active.

SUMMARY

The present application demonstrates that various modifications to a polynucleotide comprising a Factor IX nucleotide sequence can help to improve the expression level and the activity of the expressed Factor IX polypeptide. For example, the present application demonstrates that the following can improve the efficacy of a polynucleotide comprising a Factor IX nucleotide sequence for treatment of haemophilia B:

- using a codon optimised sequence;
- maintaining a portion of the Factor IX polypeptide that is not codon optimised;
- including an intron or a fragment of an intron;
- providing sequences flanking the intron or fragment of an intron that are not codon optimised;
- using a gain of function mutation;
- using a specific promoter; and/or
- maintaining an AAV genome, comprising the nucleotide, in single stranded form.

2

These modifications provide a Factor IX sequence which is expressed highly, and which encodes a highly active Factor IX polypeptide or fragment thereof. As demonstrated in the Examples, the polynucleotide of the invention expresses, and provides overall Factor IX activity, at higher levels than other Factor IX encoding polynucleotides, for example those disclosed in WO16/075473.

Accordingly, in a first aspect of the invention, there is provided a polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or fragment thereof and wherein a portion of the coding sequence is not wild type.

In a second aspect of the invention, there is provided a polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or a fragment thereof and the coding sequence comprises: (i) a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1; and (ii) a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15.

In a third aspect of the invention, there is provided a polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence encodes a Factor IX protein or fragment thereof and has at least 97%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 5.

In a fourth aspect of the invention, there is provided a viral particle comprising a recombinant genome comprising the polynucleotide of the invention.

In a fifth aspect of the invention, there is provided a composition comprising the polynucleotide or viral particle of the invention and a pharmaceutically acceptable excipient.

In a sixth aspect of the invention, there is provided a method of treatment comprising administering an effective amount of the polynucleotide or viral particle of the invention to a patient.

In a seventh aspect of the invention, there is provided a use of the polynucleotide, viral particle or composition of the invention in the manufacture of a medicament for use in a method of treatment.

The invention described herein also relates to the following aspects:

1. A polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or fragment thereof and wherein a portion of the coding sequence is not wild type.
2. The polynucleotide of aspect 1, wherein the portion of the coding sequence that is not wild type is codon optimised.
3. A polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or a fragment thereof and the coding sequence comprises:
 - (i) a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1; and
 - (ii) a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15.

3

4. The polynucleotide of aspect 3, wherein the sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, or at least 99.8% identical to SEQ ID NO. 1 is codon optimised.
5. A polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence encodes a Factor IX protein or fragment thereof and has at least 97%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 5.
6. The polynucleotide of aspect 5, wherein the Factor IX nucleotide sequence comprises a coding sequence and a portion of the coding sequence is codon optimised.
7. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises DNA or RNA.
8. The polynucleotide of any one of aspects 2, 4, 6 or 7, wherein the portion of the coding sequence that is codon optimised is a contiguous portion.
9. The polynucleotide of aspect 2, 4, 6, 7 or 8, wherein the portion of the coding sequence that is codon optimised is codon optimised for expression in the human liver.
10. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to a reference wild type Factor IX nucleotide sequence.
11. The polynucleotide of any one of aspects 2, 4, 6 or 7, wherein the portion of the coding sequence that is codon optimised is at least 800, at least 900, at least 1100, less than 1500, less than 1300, less than 1200, between 800 and 1500, between 900 and 1300, between 1100 and 1200, or around 1191 nucleotides in length.
12. The polynucleotide of any one of aspects 2, 4 or 6-11, wherein the portion of the coding sequence that is codon optimised comprises 1, 2, 3, 4, 5 or all of:
- exon 3 or a portion of at least 10, at least 15, at least 20, less than 25, between 10 and 25, between 15 and 25, or between 20 and 25 nucleotides of exon 3;
 - exon 4 or a portion of at least 80, at least 90, at least 100, less than 114, between 80 and 114, between 90 and 114, or between 100 and 114 nucleotides of exon 4;
 - exon 5 or a portion of at least 90, at least 100, at least 110, less than 129, between 90 and 129, between 100 and 129, or between 110 and 129 nucleotides of exon 5;
 - exon 6 or a portion of at least 150, at least 180, at least 200, less than 203, between 150 and 203, between 180 and 203, or between 200 and 203 nucleotides of exon 6;
 - exon 7 or a portion of at least 70, at least 80, at least 90, at least 100, less than 115, between 70 and 115, between 80 and 115, between 90 and 115, or between 100 and 115 nucleotides of exon 7; and/or
 - exon 8 or a portion of at least 400, at least 450, at least 500, less than 548, between 400 and 548, between 450 and 548, or between 500 and 548 nucleotides of exon 8.
13. The polynucleotide of aspect 12, wherein the portion of the coding sequence that is codon optimised comprises a), b), c), d), e) and f).
14. The polynucleotide of aspect 12 or aspect 13, wherein the portion of the coding sequence that is codon optimised comprises a portion of at least 20 nucleotides of exon 3, a portion of at least 100 nucleotides of exon 4, a portion of at least 110 nucleotides of exon 5, a portion of at least 180 nucleotides of exon 6, a portion of at least 100 nucleotides of exon 7, and a portion of at least 500 nucleotides of exon 8.

4

15. The polynucleotide of any one of aspects 12-14, wherein the portion of the coding sequence that is codon optimised comprises exon 3, exon 4, exon 5, exon 6, exon 7, and exon 8.
16. The polynucleotide of any one of aspects 2, 4 or 6-15, wherein the portion of the coding sequence that is codon optimised comprises a portion of exon 2, and the portion of exon 2 is less than 160, less than 150, less than 100, less than 75, less than 60, at least 20, at least 30, at least 40, at least 50, between 20 and 160, between 30 and 150, between 30 and 100, between 40 and 75, or around 56 nucleotides in length.
17. The polynucleotide of any one of aspects 2, 4 or 6-16, wherein the portion of the coding sequence that is codon optimised comprises a portion of exon 2 that is between 30 and 100 nucleotides in length.
18. The polynucleotide of any one of aspects 2, 4 or 6-17, wherein the portion of the coding sequence that is codon optimised comprises a reduced number of CpGs compared to a corresponding portion of a reference wild type Factor IX sequence.
19. The polynucleotide of aspect 18, wherein the portion of the coding sequence that is codon optimised comprises less than 40, less than 20, less than 18, less than 10, less than 5, or less than 1 CpG.
20. The polynucleotide of aspect 18 or 19, wherein the portion of the coding sequence that is codon optimised is CpG free.
21. The polynucleotide of any one of aspects 2, 4 or 6-20, wherein, in the portion of the coding sequence that is codon optimised, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, or at least 73% of the codons are selected from the group consisting of:
- TTC;
 - CTG;
 - ATC;
 - GTG;
 - GTC;
 - AGC;
 - CCC;
 - ACC;
 - GCC;
 - TAC;
 - CAC;
 - CAG;

5

-continued

m)
AAC;n)
AAA;o)
AAG;p)
GAC;q)
TGC;r)
AGG;s)
GGC;
andt)
GAG.

22. The polynucleotide of any one of aspects 2, 4 or 6-21, wherein, in the portion of the coding sequence that is codon optimised:

- a) at least 1, at least 2, at least 4, or at least 5 codons that encode phenylalanine is/are replaced with TTC compared to a reference wild type Factor IX sequence;
- b) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT; and/or
- d) the codons that encode phenylalanine are TTC, except where the following codon starts with a G.

23. The polynucleotide of any one of aspects 2, 4 or 6-22, wherein, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 10, at least 15, or at least 16 codons that encode leucine is/are replaced with CTG compared to a reference wild type Factor IX sequence;
- b) at least 90%, or at least 94% of the codons that encode leucine are CTG; and/or
- c) at least 90%, or at least 94% of the codons that encode leucine are CTG and the remainder are CTC.

24. The polynucleotide of any one of aspects 2, 4, 6-23, wherein, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 10, at least 11, or at least 12 codons that encode isoleucine is/are replaced with ATC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon ATC is/are replaced with ATT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC;
- d) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT; and/or
- e) the codons that encode isoleucine are ATC, except where the following codon starts with a G.

25. The polynucleotide of any one of aspects 2, 4 or 6-24, wherein, in the portion of the coding sequence that is codon optimised:

- a) at least 10, at least 15, at least 20, or at least 25 codons that encode valine is/are replaced with GTG compared to a reference wild type Factor IX sequence;

6

b) at least 1 codon that encodes valine is/are replaced with GTC compared to a reference wild type Factor IX sequence;

c) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG; and/or

d) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG and the remainder are GTC.

26. The polynucleotide of any one of aspects 2, 4 or 6-25, wherein, in the portion of the coding sequence that is codon optimised:

a) at least 5, at least 10, at least 12, or at least 13 codons that encode serine is/are replaced with AGC compared to a reference wild type Factor IX sequence;

b) at least 1, at least 2, or at least 4 codons that encode serine is/are replaced with TCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;

c) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC; and/or

d) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC and the remainder are TCT or TCC.

27. The polynucleotide of any one of aspects 2, 4 or 6-26, wherein, in the portion of the coding sequence that is codon optimised:

a) at least 1, at least 2, or at least 5 codons that encode proline is/are replaced with CCC compared to a reference wild type Factor IX sequence;

b) at least 1 codons that encode proline is/are replaced with CCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;

c) at least 50%, at least 55%, or at least 60% of the codons that encode proline are CCC;

d) at 50%, at least 55%, or at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT; and/or

e) the codons that encode proline are CCC, except where the following codon starts with a G.

28. The polynucleotide of any one of aspects 2, 4 or 6-27, wherein, in the portion of the coding sequence that is codon optimised:

a) at least 6, at least 7, at least 8, or at least 10 codons that encode threonine is/are replaced with ACC compared to a reference wild type Factor IX sequence;

b) at least 1, or at least 2, codons that encode threonine is/are replaced with ACT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;

c) at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC;

d) at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC and the remainder are ACT; and/or

e) the codons that encode threonine are ACC, except where the following codon starts with a G.

29. The polynucleotide of any one of aspects 2, 4 or 6-28, wherein, in the portion of the coding sequence that is codon optimised:

a) at least 1, at least 2, at least 3, or at least 4 codons that encode alanine is/are replaced with GCC compared to a reference wild type Factor IX sequence;

b) at least 1, at least 2, or at least 3 codons that encode alanine is/are replaced with GCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;

7

- c) at least 35%, at least 40%, or at least 43% of the codons that encode alanine are GCC;
- d) at least 35%, at least 40%, or at least 45% of the codons that encode alanine are GCC and the remainder are GCT; and/or
- e) the codons that encode alanine are GCC, except where the following codon starts with a G.
30. The polynucleotide of any one of aspects 2, 4 or 6-29, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 1, or at least 2 codons that encode tyrosine is/are replaced with TAC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon TAC is/are replaced with TAT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC;
- d) at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT; and/or
- e) the codons that encode tyrosine are TAC, except where the following codon starts with a G.
31. The polynucleotide of any one of aspects 2, 4 or 6-30, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 1 codons that encode histidine is/are replaced with CAC compared to a reference wild type Factor IX sequence;
- b) at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC;
- c) at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC and the remainder are CAT; and/or
- d) the codons that encode histidine are CAC, except where the following codon starts with a G.
32. The polynucleotide of any one of aspects 2, 4 or 6-31, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 1, at least 2, at least 4, or at least 5 codons that encode glutamine is/are replaced with CAG compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon CAG is/are replaced with CAA compared to a reference wild type Factor IX sequence;
- c) at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG; and/or
- d) at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG and the remainder are CAA.
33. The polynucleotide of any one of aspects 2, 4 or 6-32, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 1, at least 2, at least 4, or at least 5 codons that encode asparagine is/are replaced with AAC compared to a reference wild type Factor IX sequence;
- b) at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC and the remainder are AAT; and/or
- d) the codons that encode asparagine are AAC, except where the following codon starts with a G.
34. The polynucleotide of any one of aspects 2, 4 or 6-33, wherein, in the portion of the coding sequence that is codon optimised:

8

- a) at least 5, at least 7, at least 8, or at least 9 codons that encode lysine is/are replaced with AAG compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon AAG is/are replaced with AAA compared to a reference wild type Factor IX sequence;
- c) at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG; and/or
- d) at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG and the remainder are AAA.
35. The polynucleotide of any one of aspects 2, 4 or 6-34, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 1, at least 2, at least 3, or at least 4 codons that encode aspartate is/are replaced with GAC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon GAC is/are replaced with GAT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC;
- d) at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC and the remainder are GAT; and/or
- e) the codons that encode aspartate are GAC, except where the following codon starts with a G.
36. The polynucleotide of any one of aspects 2, 4 or 6-35, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 15, at least 20, at least 25, or at least 26 codons that encode glutamate is/are replaced with GAG compared to a reference wild type Factor IX sequence;
- b) at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG; and/or
- c) at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG and the remainder are GAA.
37. The polynucleotide of any one of aspects 2, 4, or 6-36, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 5, at least 6, at least 7, or at least 8 codons that encode cysteine is/are replaced with TGC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon TGC is/are replaced with TGT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC;
- d) at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC and the remainder are TGT; and/or
- e) the codons that encode cysteine are TGC, except where the following codon starts with a G.
38. The polynucleotide of any one of aspects 2, 4, or 6-37, wherein, in the portion of the coding sequence that is codon optimised the codons that encode tryptophan are TGG.
39. The polynucleotide of any one of aspects 2, 4, or 6-38, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 5, at least 8, at least 10, or at least 11 codons that encode arginine is/are replaced with AGG compared to a reference wild type Factor IX sequence;
- b) at least 1 codon that encodes arginine is/are replaced with AGA compared to a reference wild type Factor IX sequence;

- c) at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG; and/or
- d) at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG and the remainder are AGA. 5
40. The polynucleotide of any one of aspects 2, 4, or 6-39, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 5, at least 10, at least 12, or at least 13 codons that encode glycine is/are replaced with GGC compared to a reference wild type Factor IX sequence; 10
- b) at least 5, at least 6, at least 7, or at least 8 codons that encode glycine is/are replaced with GGG compared to a reference wild type Factor IX sequence, where the following codon starts with a G; 15
- c) at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC;
- d) at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC and the remainder are GGG; and/or 20
- e) the codons that encode glycine are GGC, except where the following codon starts with a G.
41. The polynucleotide of any one of aspects 2, 4, or 6-40, wherein the portion of the coding sequence that is codon optimised comprises codons that encode phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine. 25
42. The polynucleotide of any one of aspects 2, 4, or 6-41, wherein the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the codon optimised portion: 30
- a) at least 5 codons that encode phenylalanine is/are replaced with TTC compared to a reference wild type Factor IX sequence; 40
- b) at least 16 codons that encode leucine is/are replaced with CTG compared to a reference wild type Factor IX sequence;
- c) at least 12 codons that encode isoleucine is/are replaced with ATC compared to a reference wild type Factor IX sequence; 45
- d) at least 25 codons that encode valine is/are replaced with GTG compared to a reference wild type Factor IX sequence;
- e) at least 13 codons that encode serine is/are replaced with AGC compared to a reference wild type Factor IX sequence; 50
- f) at least 5 codons that encode proline is/are replaced with CCC compared to a reference wild type Factor IX sequence; 55
- g) at least 10 codons that encode threonine is/are replaced with ACC compared to a reference wild type Factor IX sequence;
- h) at least 4 codons that encode alanine is/are replaced with GCC compared to a reference wild type Factor IX sequence; 60
- i) at least 2 codons that encode tyrosine is/are replaced with TAC compared to a reference wild type Factor IX sequence;
- j) at least 1 codons that encode histidine is/are replaced with CAC compared to a reference wild type Factor IX sequence; 65

- k) at least 5 codons that encode glutamine is/are replaced with CAG compared to a reference wild type Factor IX sequence;
- l) at least 5 codons that encode asparagine is/are replaced with AAC compared to a reference wild type Factor IX sequence;
- m) at least 9 codons that encode lysine is/are replaced with AAG compared to a reference wild type Factor IX sequence;
- n) at least 4 codons that encode aspartate is/are replaced with GAC compared to a reference wild type Factor IX sequence;
- o) at least 26 codons that encode glutamate is/are replaced with GAG compared to a reference wild type Factor IX sequence;
- p) at least 8 codons that encode cysteine is/are replaced with TGC compared to a reference wild type Factor IX sequence;
- q) the codons that encode tryptophan are TGG;
- r) at least 11 codons that encode arginine is/are replaced with AGG compared to a reference wild type Factor IX sequence; and
- s) at least 13 codons that encode glycine is/are replaced with GGC compared to a reference wild type Factor IX sequence.
43. The polynucleotide of any one of aspects 2, 4, or 6-42, wherein the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the codon optimised portion: 30
- a) at least 70% of the codons that encode phenylalanine are TTC;
- b) at least 94% of the codons that encode leucine are CTG;
- c) at least 75% of the codons that encode isoleucine are ATC;
- d) at least 95% of the codons that encode valine are GTG;
- e) at least 70% of the codons that encode serine are AGC;
- f) at least 60% of the codons that encode proline are CCC;
- g) at least 55% of the codons that encode threonine are ACC;
- h) at least 43% of the codons that encode alanine are GCC;
- i) at least 48% of the codons that encode tyrosine are TAC;
- j) at least 65% of the codons that encode histidine are CAC;
- k) at least 90% of the codons that encode glutamine are CAG;
- l) at least 70% of the codons that encode asparagine are AAC;
- m) at least 95% of the codons that encode lysine are AAG;
- n) at least 60% of the codons that encode aspartate are GAC;
- o) at least 95% of the codons that encode glutamate are GAG;
- p) at least 55% of the codons that encode cysteine are TGC;
- q) the codons that encode tryptophan are TGG;
- r) at least 75% of the codons that encode arginine are AGG; and
- s) at least 60% of the codons that encode glycine are GGC.
44. The polynucleotide of any one of aspects 2, 4, or 6-43, wherein the portion of the coding sequence that is codon

11

- optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the codon optimised portion:
- a) at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT;
 - b) at least 94% of the codons that encode leucine are CTG and the remainder are CTC;
 - c) at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT;
 - d) at least 95% of the codons that encode valine are GTG;
 - e) at least 70% of the codons that encode serine are AGC;
 - f) at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT;
 - g) at least 55% of the codons that encode threonine are ACC and the remainder are ACT;
 - h) at least 43% of the codons that encode alanine are GCC and the remainder are GCT;
 - i) at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT;
 - j) at least 65% of the codons that encode histidine are CAC and the remainder are CAT;
 - k) at least 90% of the codons that encode glutamine are CAG and the remainder are CAA;
 - l) at least 70% of the codons that encode asparagine are AAC and the remainder are AAT;
 - m) at least 95% of the codons that encode lysine are AAG and the remainder are AAA;
 - n) at least 60% of the codons that encode aspartate are GAC and the remainder are GAT;
 - o) at least 95% of the codons that encode glutamate are GAG and the remainder are GAA;
 - p) at least 55% of the codons that encode cysteine are TGC and the remainder are TGT;
 - q) the codons that encode tryptophan are TGG;
 - r) at least 75% of the codons that encode arginine are AGG and the remainder are AGA; and
 - s) at least 60% of the codons that encode glycine are GGC and the remainder are GGG.
45. The polynucleotide of any one of aspects 10-44, wherein the reference wild type Factor IX sequence is SEQ ID NO. 9 or SEQ ID NO. 19.
46. The polynucleotide of any one of aspects 2, 4 or 6-45, wherein the portion of the coding sequence that is codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 800, at least 900, at least 1100, less than 1191, less than 1100, less than 1000, between 800 and 1191, between 900 and 1191, or around 1191 nucleotides of SEQ ID NO. 1.
47. The polynucleotide of aspect 46, wherein the portion of the coding sequence that is codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 1.
48. The polynucleotide of aspect 46 or 47, wherein the portion of the coding sequence that is codon optimised is at least 95% identical to a fragment of between 900 and 1191 nucleotides of SEQ ID NO. 1.
49. The polynucleotide of any one of aspects 46-48, wherein the portion of the coding sequence that is codon optimised is at least 95%, or at least 98% identical to SEQ ID NO. 1.
50. The polynucleotide of any one of the preceding aspects, wherein the coding sequence comprises a portion that is not codon optimised.

12

51. The polynucleotide of aspect 50, wherein the portion that is not codon optimised is at least 100, at least 150, at least 170, at least 190, less than 250, less than 225, less than 200, or around 195 nucleotides.
52. The polynucleotide of any one of aspects 50 or 51, wherein the portion that is not codon optimised comprises exon 1 or a portion of at least 60, at least 70, at least 80, between 60 and 88, between 70 and 88, or between 80 and 88 nucleotides of exon 1.
53. The polynucleotide of any one of aspects 50-52, wherein the portion that is not codon optimised comprises a portion of at least 50, at least 75, at least 80, at least 90, at least 100, less than 140, less than 120, between 50 and 140, between 75 and 120, or around 107 nucleotides of exon 2.
54. The polynucleotide of any one of aspects 50-53, wherein the portion that is not codon optimised comprises CpGs.
55. The polynucleotide of aspect 54, wherein the portion that is not codon optimised comprises at least 1 or at least 2 CpGs per 100 nucleotides.
56. The polynucleotide of any one of aspects 50-55, wherein the portion that is not codon optimised comprises less than 50%, less than 45%, less than 40%, or less than 35% codons selected from the group consisting of:

a)
TTC;

b)
CTG;

c)
ATC;

d)
GTG;

e)
GTC;

f)
AGC;

g)
CCC;

h)
ACC;

i)
GCC;

j)
TAC;

k)
CAC;

l)
CAG;

m)
AAC;

n)
AAA;

o)
AAG;

p)
GAC;

q)
TGC;

-continued

r)
AGG;s)
GGC;
andt)
GAG.

57. The polynucleotide of any one of aspects 50-56, wherein the portion that is not codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 150, at least 175, less than 195, less than 190, or less than 180 nucleotides of SEQ ID NO. 15.

58. The polynucleotide of aspect 57, wherein the portion that is not codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15.

59. The polynucleotide of any one of aspects 50-58, wherein the portion that is not codon optimised is wild type.

60. The polynucleotide of any one of aspects 50-59, wherein the portion that is not codon optimised is at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO: 15.

61. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide further comprises an intron or a fragment of an intron that interrupts the coding sequence.

62. The polynucleotide of aspect 61, wherein the intron or the fragment of an intron is a portion of a wild type Factor IX intron.

63. The polynucleotide of aspect 61 or 62, wherein the fragment of an intron is less than 500, less than 400, less than 350, less than 300, at least 100, at least 200, at least 250, at least 290, between 100 and 500, between 200 and 400, between 250 and 350, or around 299 nucleotides.

64. The polynucleotide of any one of aspects 61-63, wherein the fragment of an intron is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 200, at least 250, or at least 290 nucleotides of SEQ ID NO. 3.

65. The polynucleotide of any one of aspects 61-64, wherein the intron or the fragment of an intron is at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.3.

66. The polynucleotide of aspect 65, wherein the intron or the fragment of an intron is at least 95%, or at least 98% identical to SEQ ID NO.3.

67. The polynucleotide of any one of aspects 61-66, wherein the intron or the fragment of an intron interrupts the portion that is not codon optimised.

68. The polynucleotide of aspect 67, wherein the intron or the fragment of an intron is flanked by at least 60, at least 70, at least 80, at least 90, or at least 100 nucleotides that are not codon optimised.

69. The polynucleotide of aspect 68, wherein the intron or the fragment of an intron is flanked by between 110 and 120 nucleotides that are not codon optimised at the 5' end and between 100 and 110 nucleotides that are not codon optimised at the 3' end.

70. The polynucleotide of any one of aspects 61-69, wherein the intron or the fragment of an intron is positioned between exon 1 and exon 2.

71. The polynucleotide of any one of aspects 61-70, wherein the intron or the fragment of the intron is a fragment of native intron 1 (intron 1a).

72. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide further comprises a transcription regulatory element.

73. The polynucleotide of aspect 72, wherein the transcription regulatory element comprises a liver-specific promoter.

74. The polynucleotide of aspect 72 or aspect 73, wherein the transcription regulatory element comprises an A1AT promoter or a fragment of an A1AT promoter.

75. The polynucleotide of aspect 74, wherein the fragment of an A1AT promoter is at least 100, at least 120, at least 150, at least 180, less than 255, between 100 and 255, between 150 and 225, between 150 and 300, or between 180 and 255 nucleotides in length.

76. The polynucleotides of aspect 75, wherein the fragment of an A1AT promoter is between 150 and 300 nucleotides in length.

77. The polynucleotides, of any one of aspects 72-76, wherein the transcription regulatory element comprises an enhancer.

78. The polynucleotide of aspect 77, wherein the enhancer is an HCR enhancer or a fragment of an HCR enhancer.

79. The polynucleotide of aspect 78, wherein the fragment of an HCR enhancer is a fragment of at least 80, at least 90, at least 100, less than 192, between 80 and 192, between 90 and 192, between 100 and 250, or between 117 and 192 nucleotides in length.

80. The polynucleotide of aspect 79, wherein the fragment of an HCR enhancer is between 100 and 250 nucleotides in length.

81. The polynucleotide of any one of aspects 72-80, wherein the transcription regulatory element is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 6.

82. The polynucleotide of aspect 81, wherein the transcription regulatory element has a sequence of SEQ ID NO. 6.

83. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises an enhancer that is at least 80%, at least 85%, at least 90%, at least 95% at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13.

84. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises an enhancer that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13.

85. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises an enhancer of SEQ ID NO. 13.

86. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a promoter that is at least 80%, at least 85%, at least 90%, at least 95% at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14.

87. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a promoter that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14.

88. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a promoter of SEQ ID NO. 14.

89. The polynucleotide of any one of the preceding aspects, wherein the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to codon 384 of wild type factor IX, and wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes alanine or leucine.
90. The polynucleotide of aspect 89, wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX is CTX, wherein X is any nucleotide.
91. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a Factor IX nucleotide sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment at least 1200, at least 1350, or at least 1650 nucleotides of SEQ ID NO. 5.
92. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a Factor IX nucleotide sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.5.
93. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1; and
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.
94. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a coding sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and
 - (iii) the polynucleotide comprises a promoter element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14 and/or an enhancer element that is at least 98%, at least 99%, at least 99.5%, at least 99.8% or 100% identical to SEQ ID NO. 13.
95. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and

- (iii) the polynucleotide comprise a transcription regulatory element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 6.
96. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
 - (ii) the Factor IX nucleotide sequence comprises a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a corresponding portion of SEQ ID NO: 2; and
 - (iii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.
97. The polynucleotide of any one of aspects 95 or 96, wherein the Factor IX nucleotide sequence comprises an intron or a fragment of an intron, and the fragment of an intron is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 3.
98. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a coding sequence and a portion of the coding sequence is not codon optimised; and
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.
99. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 12 and a transcription regulatory element of SEQ ID NO. 7.
100. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 6.
101. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at a level at least 2, or at least 3 times greater than a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 12 or SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 7 or SEQ ID NO. 6.
102. A viral particle comprising a recombinant genome comprising the polynucleotide of any one of the preceding aspects.
103. The viral particle of aspect 102, which is an AAV, adenoviral, or lentiviral viral particle.
104. The viral particle of aspect 103, which is an AAV viral particle.

105. The viral particle of any one of aspects 102-104, wherein the recombinant genome further comprises:
- AAV2 ITRs;
 - a poly A sequence;
 - an origin of replication; and/or
 - two resolvable ITRs.
106. The viral particle of aspect 105, wherein the recombinant genome is single-stranded and/or comprises two resolvable ITRs.
107. The viral particle of any one of aspects 102-106, wherein the viral particle comprises a capsid selected from the group consisting of:
- a capsid having at least 96%, at least 98%, at least 99%, at least 99.5%, at least 99.8% identity or 100% identity to SEQ ID NO.10;
 - a capsid having at least 96%, at least 98%, at least 99%, at 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 17;
 - AAVMutC; and
 - AAV5.
108. The viral particle of any one of aspects 102-107, wherein on transduction into Huh7 cells, the viral particle expresses Factor IX protein or a fragment thereof having a Factor IX activity greater than the activity of Factor IX expressed from a viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO: 12 and a transcription regulatory element of SEQ ID NO. 7 and/or a viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 6.
109. The viral particle of aspect 108, wherein the activity is measured using a chromogenic substrate which is specific for Factor Xa.
110. The polynucleotide or viral particle of any one of the preceding aspects, wherein the Factor IX protein fragment is at least 200, at least 250, at least 300, between 200 and 415, between 250 and 415, or between 300 and 415 amino acids in length.
111. The polynucleotide or viral particle of any one of the preceding aspects, wherein the Factor IX protein or fragment thereof comprises a sequence:
- at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 8; or
 - at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of SEQ ID NO. 8 at least 200, at least 250, at least 300, between 200 and 415, between 250 and 415, or between 300 and 415 amino acids in length.
112. A composition comprising the polynucleotide or viral particle of any one of the preceding aspects and a pharmaceutically acceptable excipient.
113. The polynucleotide, viral particle or composition of any one of the preceding aspects for use in a method of treatment.
114. The polynucleotide, viral particle or composition for use of aspect 113, wherein the method of treatment comprises administering an effective amount of the polynucleotide or viral particle of any one of aspects 1-111 to a patient.
115. A method of treatment comprising administering an effective amount of the polynucleotide or viral particle of any one of aspects 1-111 to a patient.
116. Use of the polynucleotide, viral particle or composition of any one of aspects 1-111 in the manufacture of a medicament for use in a method of treatment.

117. The use of aspect 116, wherein the method of treatment comprises administering an effective amount of the polynucleotide or viral particle of any one of aspects 1-111 to a patient.
118. The polynucleotide, viral particle, composition, use or method of any one of aspects 112-117, wherein the method of treatment is a method of treating haemophilia.
119. The polynucleotide, viral particle, composition, use or method of aspect 118, wherein the haemophilia is haemophilia B.
120. The polynucleotide, viral particle, composition, use or method of aspect 119, wherein the patient has antibodies or inhibitors to Factor IX.

BRIEF DESCRIPTION OF THE DRAWINGS

- FIG. 1A, FIG. 1B, and FIG. 1C—Schematic of FIX transgene cassettes ssLP1.FIXco (FIG. 1C), ssHLP2.TI-codop-FIX-GoF (HTFG) (FIG. 1B) and ssHLP2.TI-ACNP-FIX-GoF (HTAG) (FIG. 1A). ITR=Inverted Terminal Repeat; HLP2 and LP1 are transcription regulatory elements of SEQ ID NOs: 6 and 7, respectively; E=Exon; T1=Truncated Intron 1A; WT=Wild Type; CO=Codon Optimised; ACNP=a codon optimised sequence of the invention.
- FIG. 2A, FIG. 2B, FIG. 2C, FIG. 2D, FIG. 2E, and FIG. 2F—Results from HUH7 transduction with AAV2/Mut C vectors—Experiment 1; FIG. 2A, FIG. 2B, and FIG. 2C show the level of FIX antigen in supernatant. FIG. 2A shows level of FIX antigen after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^3 , FIG. 2B shows level of FIX antigen after HUH7 transduction with AAV2/MutC vectors at MOI of 5×10^3 , FIG. 2C shows level of FIX antigen after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^4 ; FIG. 2D, FIG. 2E, and FIG. 2F show the level of FIX antigen after normalisation using the number of vector genomes present in cell lysate. FIG. 2D shows the level of FIX antigen after normalisation, for transduction with AAV2/MutC vectors at MOI of 1×10^3 , FIG. 2E shows the level of FIX antigen after normalisation, for transduction with AAV2/MutC vectors at MOI of 5×10^3 , FIG. 2F shows the level of FIX antigen after normalisation, for transduction with AAV2/MutC vectors at MOI of 1×10^4 ; Error bars represent mean \pm SD of n=2. $1e3=1 \times 10^3$; $5e3=5 \times 10^3$; $1e4=1 \times 10^4$; MOI=multiplicity of infection.
- FIG. 3A, FIG. 3B, and FIG. 3C—Results from HUH7 transduction with AAV2/Mut C vectors—Experiment 2, showing the level of FIX antigen in supernatant. Error bars represent mean \pm SD of n=3. $1e3=1 \times 10^3$; $5e3=5 \times 10^3$; $1e4=1 \times 10^4$, MOI=multiplicity of infection. FIG. 3A shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^3 . FIG. 3B shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 5×10^3 . FIG. 3C shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^4 .
- FIG. 4A, FIG. 4B, FIG. 4C, FIG. 4D, and FIG. 4E—Results from HUH7 transduction with AAV2/Mut C vectors—Experiment 2; FIG. 4A shows the level of FIX antigen in supernatant. FIG. 4A shows level of FIX antigen in the supernatant after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^3 , FIG. 4B shows level of FIX antigen in the supernatant after HUH7 transduction with AAV2/MutC vectors at MOI of 5×10^3 , FIG. 4C shows level of FIX antigen in the supernatant after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^4 ; FIG. 4D and FIG. 4E show the level of FIX antigen after normalisation using the number of vector genomes present in cell lysate. FIG. 4D shows the level of FIX antigen after normalisation, for

transduction with AAV2/MutC vectors at MOI of 5×10^3 , FIG. 4E shows the level of FIX antigen after normalisation, for transduction with AAV2/MutC vectors at MOI of 1×10^4 ; Error bars represent mean \pm SD of $n=3$. $1e3=1 \times 10^3$; $5e3=5 \times 10^3$; $1e4=1 \times 10^4$; MOI=multiplicity of infection.

FIG. 5—Combined data (from Experiments 1 and 2) for AAV2/Mut C transduction of HUH7 cells at MOI 5×10^3 . Error bars represent mean \pm SD of $n=12$. $P=0.001$ by Student's T-test. $5e3=5 \times 10^3$; MOI=multiplicity of infection.

FIG. 6A, FIG. 6B, and FIG. 6C—The activity of FIX for MOI 1×10^3 , 5×10^3 and 1×10^4 is shown after HUH7 transduction with AAV2/Mut C vectors (Experiment 1). Error bars represent mean \pm SD of $n=2$ duplicate wells. $1e3=1 \times 10^3$; $5e3=5 \times 10^3$; $1e4=1 \times 10^4$ MOI=multiplicity of infection. FIG. 6A shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^3 . FIG. 6B shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 5×10^3 . FIG. 6C shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^4 .

FIG. 7A, FIG. 7B, and FIG. 7C—The activity of FIX is shown after HUH7 transduction with AAV2/Mut C vectors (Experiment 2). Error bars represent mean \pm SD of $n=3$. $1e3=1 \times 10^3$; $5e3=5 \times 10^3$; $1e4=1 \times 10^4$, MOI=multiplicity of infection. FIG. 7A shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^3 . FIG. 7B shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 5×10^3 . FIG. 7C shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^4 .

FIG. 8A, FIG. 8B, and FIG. 8C—The activity of FIX is shown after HUH7 transduction with AAV2/Mut C vectors (Experiment 2). Error bars represent mean \pm SD of $n=3$. $1e3=1 \times 10^3$; $5e3=5 \times 10^3$; $1e4=1 \times 10^4$, MOI=multiplicity of infection. FIG. 8A shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^3 . FIG. 8B shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 5×10^3 . FIG. 8C shows activity of FIX after HUH7 transduction with AAV2/MutC vectors at MOI of 1×10^4 .

FIG. 9—Combined data (from Experiments 1 and 2; FIG. 6B and FIG. 7B) showing activity of FIX for MOI 5×10^3 shown after HUH7 transduction with AAV2/Mut C vectors. Error bars represent mean \pm SD of $n=12$. $5e3=5 \times 10^3$; MOI=multiplicity of infection. Statistical significance determined using a Student's T-test ($p=0.0195$).

FIG. 10—Normalised level of human FIX in murine plasma after administration of AAV2/8.LP1.FIXco and AAV2/8.HLP2.HTFG (Experiment 3). FIX:Ag levels were normalised to vector copies/cell. Error bars represent mean \pm SD of $n=4$ mice. P -value <0.05 (Student's T-test)

FIG. 11A and FIG. 11B—Comparison of alternate codon optimisation of FIX in C57BL/6 mice (Experiment 4). Mice were injected with AAV2/8 vectors containing ssHLP2.HTFG, ssHLP2.HTAG or scLP1.FIXco (control). FIG. 11A shows level of FIX antigen, wherein the level of FIX antigen was assessed 3 weeks post-injection ($P=0.0007$ between ssHLP2.HTAG and sc.LP1.FIXco and $p=0.0198$ between ssHLP2.HTAG and ssHLP2.HTFG). FIG. 11B shows level of FIX antigen, wherein antigen levels were normalised to vector genome ($p=0.0009$ between ssHLP2.HTAG and sc.LP1.FIXco and $p=0.0039$ between ssHLP2.HTAG and ssHLP2.HTFG). $n=4$ mice. P -values were determined using one-way ANOVA (multiple comparison).

FIG. 12—Comparison of alternate codon optimisation of FIX in C57BL/6 mice (Experiment 4). Mice were injected

with AAV2/8 vectors containing ssHLP2.HTFG, ssHLP2.HTAG or scLP1.FIXco (control). The level of FIX activity was assessed 3 weeks post-injection. $n=4$ mice. $P=0.0008$ between ssHLP2.HTAG and scLP1.FIXco and $p=0.01$ between ssHLP2.HTAG and ssHLP2.HTFG; p values were determined using one-way ANOVA (multiple comparison).

FIG. 13—Schematic of Factor IX structure. The numbers above the schematic represent amino acid positions in the complete Factor IX polypeptide including the signal peptide and the pro-peptide region (encoded by SEQ ID NO. 9). The numbers below the schematic represent equivalent amino acid positions in mature Factor IX (which corresponds to the portion of coding sequence in SEQ ID NO.19).

DETAILED DESCRIPTION

General Definitions

Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by a person skilled in the art to which this invention belongs.

In general, the term “comprising” is intended to mean including but not limited to. For example, the phrase “a polynucleotide comprising a Factor IX nucleotide sequence” should be interpreted to mean that the polynucleotide has a Factor IX nucleotide sequence, but the polynucleotide may contain additional nucleotides.

In some embodiments of the invention, the word “comprising” is replaced with the phrase “consisting of”. The term “consisting of” is intended to be limiting. For example, the phrase “a polynucleotide consisting of a Factor IX nucleotide sequence” should be understood to mean that the polynucleotide has a Factor IX nucleotide sequence and no additional nucleotides.

The terms “protein” and “polypeptide” are used interchangeably herein, and are intended to refer to a polymeric chain of amino acids of any length.

For the purpose of this invention, in order to determine the percent identity of two sequences (such as two polynucleotide or two polypeptide sequences), the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in a first sequence for optimal alignment with a second sequence). The nucleotide residues at nucleotide positions are then compared. When a position in the first sequence is occupied by the same nucleotide residue as the corresponding position in the second sequence, then the nucleotides are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (i.e., % identity=number of identical positions/total number of positions in the reference sequence $\times 100$).

Typically the sequence comparison is carried out over the length of the reference sequence. For example, if the user wished to determine whether a given (“test”) sequence is 95% identical to SEQ ID NO. 5, SEQ ID NO. 5 would be the reference sequence. For example, to assess whether a sequence is at least 80% identical to SEQ ID NO. 5 (an example of a reference sequence), the skilled person would carry out an alignment over the length of SEQ ID NO. 5, and identify how many positions in the test sequence were identical to those of SEQ ID NO. 5. If at least 80% of the positions are identical, the test sequence is at least 80% identical to SEQ ID NO. 5. If the sequence is shorter than SEQ ID NO. 5, the gaps or missing positions should be considered to be non-identical positions.

The skilled person is aware of different computer programs that are available to determine the homology or identity between two sequences. For instance, a comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In an embodiment, the percent identity between two amino acid or nucleic acid sequences is determined using the Needleman and Wunsch (1970) algorithm which has been incorporated into the GAP program in the Accelrys GCG software package (available at <http://www.accelrys.com/products/gcg/>), using either a Blosum 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6.

For the purposes of the present invention, the term “fragment” refers to a contiguous portion of a sequence. For example, a fragment of SEQ ID NO. 5 of 50 nucleotides refers to 50 contiguous nucleotides of SEQ ID NO. 5.

A Polynucleotide

In one aspect, the present invention provides a polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or fragment thereof and wherein a portion of the Factor IX nucleotide sequence is not wild type.

The polynucleotide may further comprise one or more of the following features. The polynucleotide may comprise a portion that is not codon optimised. The polynucleotide may comprise an intron or a fragment of an intron. The polynucleotide may comprise a mutation in a codon corresponding to codon 384 of wild type Factor IX.

The term “polynucleotide” refers to a polymeric form of nucleotides of any length, deoxyribonucleotides, ribonucleotides, or analogs thereof. For example, the polynucleotide may comprise DNA (deoxyribonucleotides) or RNA (ribonucleotides). The polynucleotide may consist of DNA. The polynucleotide may be mRNA. Since the polynucleotide

may comprise RNA or DNA, all references to T (thymine) nucleotides may be replaced with U (uracil).

A Factor IX Nucleotide Sequence

The polynucleotide comprises a Factor IX nucleotide sequence. The Factor IX nucleotide sequence comprises a coding sequence that encodes the Factor IX protein or fragment thereof.

A “coding sequence” is a sequence that encodes a polynucleotide, and excludes non coding regions such as introns. A coding sequence may be interrupted by non-coding nucleotides (e.g. an intron), but only nucleotides that encode the polypeptide should be considered to be part of the coding sequence. For example, a coding sequence that encodes a Factor IX protein will comprise any codons that encode an amino acid forming part of the Factor IX protein that is expressed from that coding sequence, irrespective of whether those codons are contiguous in sequence or separated by one or more non-coding nucleotides. In other words, a polynucleotide which contains stretches of coding nucleotides interrupted by a stretch of non-coding nucleotides will be considered to comprise a “coding sequence” consisting of the non-contiguous coding stretches immediately juxtaposed (i.e. minus the non-coding stretch). However, herein, the stop codon will be considered to be part of the full length coding sequence.

The term “sequence that encodes” refers to a nucleotide sequence comprising codons that encode the encoded polypeptide. For example, a nucleotide sequence that encodes a Factor IX protein or fragment thereof comprises codons that encode the amino acid sequence of the Factor IX protein or fragment thereof. A suitable nucleotide sequence is provided in SEQ ID NO. 5.

The following Table describes codons that encode each amino acid:

Amino Acid	Codon	Amino Acid	Codon	Amino Acid	Codon
Phenylalanine	TTC	Proline	CCT	Asparagine	AAT
	TTT		CCC		AAC
			CCA		
			CCG		
Leucine	TTA	Threonine	ACT	Lysine	AAA
	TTG		ACC		AAG
	CTT		ACA		
	CTC		ACG		
	CTA				
	CTG				
Isoleucine	ATT	Alanine	GCT	Aspartic Acid	GAT
	ATC		GCC		GAC
	ATA		GCA		
			GCG		
Methionine	ATG	Tyrosine	TAT	Glutamic Acid	GAA
			TAC		GAG
Valine	GTT	Histidine	CAT	Cysteine	TGT
	GTC		CAC		TGC
	GTA				
	GTG				
Serine	TCT	Glutamine	CAA	Tryptophan	TGG
	TCC		CAG		
	TCA				
	TCG				
	AGT				
	AGC				

Amino Acid	Codon	Amino Acid	Codon	Amino Acid	Codon
Arginine	CGT	Glycine	GGT		
	CGC		GGC		
	CGA		GGA		
	CGG		GGG		
	AGA				
	AGG				

The corresponding RNA codons will contain Us in place of the Ts in the Table above.

One aspect of the present invention provides a polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence encodes a Factor IX protein or fragment thereof and has at least 97%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 5. Optionally, the Factor IX nucleotide sequence comprises a coding sequence and a portion of the coding sequence is codon optimised.

In general, the Factor IX nucleotide sequence may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 1200, at least 1350, or at least 1650 nucleotides of SEQ ID NO. 5. The Factor IX nucleotide sequence may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a contiguous fragment of at least 1200, at least 1350, or at least 1650 nucleotides of SEQ ID NO. 5. The Factor IX nucleotide sequence may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.5. For example, the Factor IX nucleotide sequence may be at least 98% identical to SEQ ID NO.5.

Factor IX Protein or Fragment Thereof

The polynucleotide comprises a Factor IX nucleotide sequence comprising a coding sequence that encodes a Factor IX protein or fragment thereof.

Wild type Factor IX is a serine protease, which forms part of the coagulation cascade. Lack of or mutated Factor IX can lead to reduced blood clotting and the disease haemophilia B. A typical wild type Factor IX polypeptide is encoded by SEQ ID NO. 9 (sometimes referred to as Factor IX Malmö B) or SEQ ID NO. 19. An alternative wild type Factor IX polypeptide differs from that encoded by SEQ ID NO. 9 at codon 194, for example codon 194 may encode threonine ("Malmö A") instead of alanine.

Factor IX (e.g. a Factor IX of SEQ ID NO. 16) as initially expressed as a precursor "immature" form, comprising a hydrophobic signal peptide (amino acids 1-28 of SEQ ID NO. 16), a pro-peptide region (amino acids 29-46 of SEQ ID NO. 16) and a mature polypeptide region, as set out in FIG. 13. The mature (zymogen) form of Factor IX lacks the hydrophobic signal peptide and the pro-peptide region. The term "mature Factor IX" refers to a Factor IX polypeptide that does not comprise the hydrophobic signal peptide or the pro-peptide region, such as SEQ ID NO. 8.

During clotting the single-chain zymogen form is cleaved by Factor XIa or Factor VIIa to produce an active two-chain form (Factor IXa), with the two chains linked by a disulphide bridge. The activated form can catalyse the hydrolysis of an arginine-isoleucine bond in Factor X to form Factor Xa. Wild type Factor IX is inhibited by thrombin. The wild type Factor IX protein has four protein domains, a Gla

domain, two tandem copies of the EGF domain and a C-terminal trypsin-like peptidase domain which is responsible for catalytic cleavage.

The term "Factor IX protein" refers to the single-chain zymogen form of Factor IX, the activated two-chain form and variants thereof, and may refer to the mature Factor IX polypeptide or a Factor IX polypeptide comprising the pro-peptide region and/or the signal peptide region.

Preferably the Factor IX fragment is at least 200, at least 250, at least 300, between 200 and 461, between 250 and 461, or between 300 and 461 amino acids in length. In an embodiment, the Factor IX protein or fragment thereof comprises a sequence:

- a) at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 8; or
- b) at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of SEQ ID NO. 8 at least 200, at least 250, at least 300, between 200 and 415, between 250 and 415, or between 300 and 415 amino acids in length.

In an embodiment, the Factor IX protein or fragment thereof is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.16; or at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of SEQ ID NO.16 at least 200, at least 250, at least 300, between 200 and 461, between 250 and 461, or between 300 and 461 amino acids in length. In an embodiment, the Factor IX protein or fragment thereof is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.16; or at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of SEQ ID NO.16 at least 300, or between 300 and 461 amino acids in length. The Factor IX protein or fragment thereof may have a sequence of SEQ ID NO: 16 or SEQ ID NO: 8.

Preferably the Factor IX protein or fragment thereof is functional. A functional Factor IX protein or fragment is one which carries out hydrolysis of an arginine-isoleucine bond in Factor X to form Factor Xa.

It is within the abilities of the person skilled in the art to determine whether a Factor IX protein or fragment encoded by a Factor IX nucleotide sequence is functional. The skilled person merely needs to express the Factor IX nucleotide sequence, and test whether the expressed protein is active. For example, the skilled person could prepare a viral particle of the invention comprising the Factor IX nucleotide sequence linked to an operable promoter, and transduce cells with the viral particle under conditions suitable for expression of the Factor IX protein or fragment thereof. The activity of the expressed Factor IX protein or fragment thereof can be analysed using a chromogenic assay, such as the activity assay described in Example 3.

For example, a suitable chromogenic assay is as follows. Factor IX is mixed with thrombin, phospholipids, calcium, thrombin activated Factor VIII and Factor XIa. Under these

conditions, the Factor XIa activates the Factor IX to form Factor IXa, and the activity of the Factor IXa can catalyse cleavage of a chromogenic substrate (SXA-11) to produce pNA. The level of pNA generated can be measured by determining absorbance at 405 nm, and this is proportional to the activity of the Factor IX in the sample.

The activity can be normalised to compensate for different concentrations of Factor IX in the sample, by measuring the concentration of Factor IX in the sample using a standard ELISA assay, such as the assay described in Example 4, and dividing the activity by the Factor IX concentration. For example, an antibody that binds to Factor IX could be bound to a plate. The sample, comprising the Factor IX at unknown concentration, could be passed over the plate. A second detection antibody that binds to Factor IX could be applied to the plate, and any excess washed off. The detection antibody that remains (i.e. is not washed off) will be bound to Factor IX. The detection antibody could be linked to an enzyme such as horse radish peroxidase. The level of detection antibody that binds to the Factor IX on the plate could be measured by measuring the amount of the detection antibody. For example, if the detection antibody is linked to horse radish peroxidase, the horse radish peroxidase can catalyse the production of a blue reaction product from a substrate such as TMB (3,3',5,5'-tetramethylbenzidine), and the level of the blue product can be detected by absorbance at 450 nm. The level of the blue product is proportional to the amount of detection antibody that remained after the washing step, which is proportional to the amount of Factor IX in the sample.

Optionally, the Factor IX protein or fragment thereof has an activity greater than that of the Factor IX polypeptide encoded by SEQ ID NO. 9, SEQ ID NO. 19, or SEQ ID NO. 12. Optionally, the activity is measured using a chromogenic substrate which is specific for Factor IX, i.e. a substrate which may be altered by Factor IXa to provide a chromogenic signal. A suitable chromogenic substrate is SXa-11.

In an embodiment, the Factor IX protein or fragment thereof comprises a mutation at a position corresponding to position 384 of wild type Factor IX. For example, position 384 (numbering from the start of the signal peptide, i.e. a position corresponding to amino acid 384 of SEQ ID NO. 16) of wild type Factor IX is an arginine residue (R384), but this can be replaced by a different residue. In an embodiment, R384 is replaced with a small, hydrophobic amino acid. For example, the small, hydrophobic amino acid could be alanine, isoleucine, leucine, valine or glycine. Preferably, the Factor IX protein or fragment thereof comprises a leucine at a position corresponding to position 384 in wild type Factor IX, as shown in SEQ ID NO. 16.

A mutation at a position corresponding to position 384 of the wild type sequence may cause a gain-of-function (GoF) mutation, resulting in Factor IX that is hyperfunctional. The advantage of expressing a Factor IX protein containing a mutation at position 384 is that a relatively small increase in protein amount produces a larger increase in overall protein activity.

It is within the abilities of the person skilled in the art to determine whether a given polypeptide has a mutation at a position corresponding to position 384. The person skilled in the art merely needs to align the sequence of the polypeptide sequence with that of a wild type (precursor, immature) Factor IX polypeptide, and determine whether the residue of the former that aligns with the 384th residue of the latter is an arginine. If not, the polypeptide has a mutation at a position corresponding to position 384 in wild type Factor

IX. The alignment may be performed using any suitable algorithm such as that of Needleman and Wunsch described above.

A Portion of the Coding Sequence is not Wild Type

5 A portion of the coding sequence may not be wild type. The wild type Factor IX-encoding nucleotide sequence is represented by SEQ ID NO. 9, and a coding sequence that comprises a portion differing from that of SEQ ID NO. 9 comprises a portion that is not wild type (providing such 10 portion also differs from other Factor IX coding sequences which are regarded also as wild type, for example the Malmo A variant mentioned previously).

In an embodiment, the portion of the coding sequence that is not wild type is codon optimised. To identify whether a 15 coding sequence comprises a portion that is codon optimised, one can align the coding sequence with SEQ ID NO. 9. If any portions of the sequence are not identical to SEQ ID NO. 9, the user should then determine whether they are codon optimised, i.e., whether they comprise at least one 20 codon that has been replaced with a favoured codon, i.e., one of TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG. If the portion that is not wild type comprises at least one codon that has been replaced with a 25 favoured codon, then it is codon optimised. Preferably, a contiguous portion of the coding sequence is codon optimised. However, in some embodiments, the portion of the coding sequence which is codon optimised could be split over 2, 3, 4 or 5 regions of the coding sequence. Optionally, 30 the portion of the coding sequence which is not codon optimised is split over less than 3 or less than 2 regions of the coding sequence. A nucleotide sequence can be codon optimised by replacing codons with other codons that are favoured (i.e. reflective of codon bias) in a particular organ or a particular organism (so-called favoured codons). Such 35 a codon optimisation improves expression of the nucleotide sequence in the particular organ or particular organism. For example, if a nucleotide sequence is codon optimised for the human liver, the nucleotide sequence is modified to increase the number of codons that are favoured in the human liver. The skilled person would appreciate that codon-optimising a sequence may not entail changing every codon as at some 40 positions a "favoured codon" may already be present.

Such codon optimisation may be subject to other factors. 45 For example, it can be seen that the presence of CpGs has an adverse effect on expression and so the user may decide not to use favoured codons if their use at certain positions introduces CpGs into the sequence; this will still be considered to be codon optimisation. In an embodiment, a favoured 50 codon that ends with a C nucleotide will not be included in the portion of the coding sequence that is codon optimised, where the next codon in the sequence begins with a G. For example, codon CTC encodes leucine. CTC should not be used for encoding leucine where the next codon in the 55 sequence begins with a G, such as codon GTT.

The present application discloses that certain codons are favoured for expression in the human liver and that reducing the CpG content of a coding sequence, whilst maintaining a high proportion of those favoured codons, improves expression of the coding sequence. The favoured codons are TTC, 60 CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG.

In one embodiment, the portion of the coding sequence that is codon optimised is codon optimised for expression in the liver, optionally the human liver. A portion of the coding sequence that is codon optimised for expression in the liver

may comprise a higher proportion of codons that are favoured in the liver, such as favoured codons TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG.

In an embodiment, the following codons are collectively overrepresented in the portion of the coding sequence that is not wild type or is codon optimised: TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG. By “collectively overrepresented”, is meant that the total number of favoured codons in the portion of the coding sequence which is codon optimised or not wild type is higher than the total number of the favoured codons in the corresponding portion of a wild type Factor IX nucleotide sequence (such as that as SEQ ID NO. 9 or SEQ ID NO. 19).

In a preferred embodiment, in the portion of the coding sequence that is codon optimised there is a greater frequency of the following codons compared to the corresponding portion of a wild type Factor IX nucleotide sequence (such as that of SEQ ID NO.9): TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG. Optionally, the following codons are collectively overrepresented in the portion of the coding sequence that is codon optimised, except where their presence results in a CpG: TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG.

Optionally, the portion of the coding sequence that is codon optimised comprises at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, or at least 65%, at least 70% or at least 73% of codons selected from the group consisting of: TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG.

The codon usage in a codon optimised portion of a polynucleotide of the invention (HLP2.T1-ACNP-FIX-GoF) is compared with the codon usage in a corresponding stretch of wild type Factor IX nucleotide sequence (SEQ ID NO.9) in the following table.

TABLE 1

Amino Acid	Codon	HTAG	%age of codons	Wild type
Phe	TTT	5	26	12
	TTC	14	74	9
Leu	CTT	0		9
	CTC	1	5	5
	CTA	0		2
	CTG	19	95	3
	TTA	0		6
	TTG	0		3
Ile	ATT	5	24	17
	ATC	16	76	7
	ATA	0		1
Met	TTG	0		0
	ATG	2	100	6
Val	GTT	0		22
	GTC	1	3	3
	GTA	0		5
	GTG	33	97	7
Ser	TCT	6	25	6
	TCC	1	4	5

TABLE 1-continued

Amino Acid	Codon	HTAG	%age of codons	Wild type	
5	TCA	0		6	
	TCG	0		0	
	AGT	0		7	
	AGC	17	71	3	
10	Pro	CCT	3	23	4
	CCC	8	62	3	
	CCA	2	15	8	
	CCG	0		0	
15	Thr	ACT	11	39	13
	ACC	17	61	7	
	ACA	0		10	
20	Ala	ACG	0	1	
	GCT	11	55	9	
	GCC	9	45	5	
25	GCA	0		8	
	GCG	0		0	
	Tyr	TAT	7	50	11
30	TAC	7	50	5	
	His	CAT	3	33	6
35	CAC	6	67	4	
	Gln	CAA	1	8	7
40	CAG	11	92	7	
	Asn	AAT	7	27	15
45	AAC	19	73	17	
	Lys	AAA	1	4	12
50	AAG	24	96	16	
	Asp	GAT	7	39	12
55	GAC	11	61	7	
	Glu	GAA	1	3	33
60	GAG	35	97	10	
	Cys	TGT	9	43	19
65	TGC	12	57	5	
	Trp	TGG	7	88	7
70	TGA	1	13	0	
	Arg	CGT	0		1
75	CGC	0		1	
	CGA	0		6	
80	CGG	0		3	
	AGA	3	20	8	
85	AGG	12	80	1	
	Gly	GGT	0		8
90	GGC	21	66	9	
	GGA	0		15	
95	GGG	11	34	4	

The total number of favoured codons in SEQ ID NO. 9 in this region is 120 (30% of the sequence). On the other hand, the total number of favoured codons in the codon optimised portion of HTAG is 293 (73% of the codons).

It is straightforward to determine whether a given portion of a polynucleotide comprises favoured codons. In order to determine the frequency of each codon used in a portion of a nucleotide sequence, the skilled person merely needs to enter the sequence of that portion into one of the readily available algorithms that looks at codon usage and review the results. Alternatively, the user could simply count them.

The codons that are replaced in the codon optimised portion of HTAG compared to the corresponding region of SEQ ID NO.9 are set out in the following table.

TABLE 2

Amino Acid	Codon replacements	Frequency
Pro	CCA to CCC	2
	CCA to CCT	2
	CCT to CCC	3
Leu	TTA to CTG	5
	CTC to CTG	3
	CTT to CTG	6
	TTG to CTG	2
	CTA to CTG	1
	TTA to TTG	0
Gly	GGC to GGG	1
	GGA to GGC	9
	GGT to GGG	4
	GGT to GGC	3
	GGG to GGC	2
	GGA to GGG	5
Ile	ATT to ATC	12
	ATC to ATT	1
	ATA to ATC	1
Val	GTA to GTG	4
	GTC to GTG	3
	GTG to GTA	
	GTT to GTG	20
	GTA to GTC	1
Lys	AAA to AAG	10
	AAC to AAG	
	AAG to AAA	1
Tyr	TAT to TAC	3
	TAC to TAT	1
Gln	CAA to CAG	6
	CAG to CAA	1
His	CAT to CAC	2
Glu	GAA to GAG	27
Cys	TGT to TGC	9
	TGC to TGT	1
Ser	AGT to AGC	3
	TCC to AGC	4
	AGT to TCT	3
	TCA to AGC	3
	TCT to AGC	4
	TCA to TCT	1
Ala	GCA to GCC	3
	GCA to GCT	4
	GCT to GCC	2
Arg	CGA to AGG	5
	AGA to AGG	5
	CGT to AGG	1
	CGG to AGG	1
	CGG to AGA	1
Thr	ACA to ACC	6
	ACT to ACC	4
	ACA to ACT	3
	ACG to ACC	1
Phe	TTT to TTC	5
Asp	GAT to GAC	5
	GAC to GAT	1

TABLE 2-continued

Amino Acid	Codon replacements	Frequency
Asn	AAT to AAC	6
	stop	1
	GoF mutation	1

In an embodiment, in the portion of the coding sequence that is codon optimised:

a) at least 1, at least 2, at least 4, or at least 5 codons that encode phenylalanine is/are replaced with TTC compared to a reference wild type Factor IX sequence;

b) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC;

c) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT; and/or

d) the codons that encode phenylalanine are TTC, except where the following codon starts with a G.

For example, when we say at least 1 of codon A is replaced with at least 1 of codon B, this refers to replacement of codon A with codon B in at least 1 position compared to a wild type sequence, such as SEQ ID NO. 9. To determine whether such a replacement has taken place, one merely needs to align the test sequence to a wild type Factor IX sequence and see which codons are different. If at least 1 codon in the test sequence corresponding to codon A of wild type Factor IX is codon B in the test sequence, then at least 1 of codon A has been replaced by codon B. For example, if the first codon is TTT in the test sequence and TTC in the wild type Factor IX sequence, the test sequence comprises at least 1 of codon TTC replaced with TTT.

In an embodiment, in the portion of the coding sequence that is codon optimised:

a) at least 5, at least 10, at least 15, or at least 16 codons that encode leucine is/are replaced with CTG compared to a reference wild type Factor IX sequence;

b) at least 90%, or at least 94% of the codons that encode leucine are CTG; and/or

c) at least 90%, or at least 95% of the codons that encode leucine are CTG and the remainder are CTC.

In an embodiment, in the portion of the coding sequence that is codon optimised:

a) at least 5, at least 10, at least 11, or at least 12 codons that encode isoleucine is/are replaced with ATC compared to a reference wild type Factor IX sequence;

b) at least 1 of codon ATC is/are replaced with ATT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;

c) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC;

d) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT; and/or

e) the codons that encode isoleucine are ATC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

a) at least 10, at least 15, at least 20, or at least 25 codons that encode valine is/are replaced with GTG compared to a reference wild type Factor IX sequence;

31

- b) at least 1 codon that encodes valine is/are replaced with GTC compared to a reference wild type Factor IX sequence;
- c) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG; and/or
- d) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG and the remainder are GTC.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 10, at least 12, or at least 13 codons that encode serine is/are replaced with AGC compared to a reference wild type Factor IX sequence;
- b) at least 1, at least 2, or at least 4 codons that encode serine is/are replaced with TCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC; and/or
- d) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC and the remainder are TCT or TCC.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1, at least 2, or at least 5 codons that encode proline is/are replaced with CCC compared to a reference wild type Factor IX sequence;
- b) at least 1 codons that encode proline is/are replaced with CCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 50%, at least 55%, or at least 60% of the codons that encode proline are CCC;
- d) at 50%, at least 55%, or at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT; and/or
- e) the codons that encode proline are CCC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 6, at least 7, at least 8, or at least 10 codons that encode threonine is/are replaced with ACC compared to a reference wild type Factor IX sequence;
- b) at least 1, or at least 2, codons that encode threonine is/are replaced with ACT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC;
- d) at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC and the remainder are ACT; and/or
- e) the codons that encode threonine are ACC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1, at least 2, at least 3, or at least 4 codons that encode alanine is/are replaced with GCC compared to a reference wild type Factor IX sequence;
- b) at least 1, at least 2, or at least 3 codons that encode alanine is/are replaced with GCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 35%, at least 40%, or at least 43% of the codons that encode alanine are GCC;
- d) at least 35%, at least 40%, or at least 45% of the codons that encode alanine are GCC and the remainder are GCT; and/or

32

- e) the codons that encode alanine are GCC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1, or at least 2 codons that encode tyrosine is/are replaced with TAC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon TAC is/are replaced with TAT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC;
- d) at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT; and/or
- e) the codons that encode tyrosine are TAC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1 codons that encode histidine is/are replaced with CAC compared to a reference wild type Factor IX sequence;
- b) at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC;
- c) at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC and the remainder are CAT; and/or
- d) the codons that encode histidine are CAC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1, at least 2, at least 4, or at least 5 codons that encode glutamine is/are replaced with CAG compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon CAG is/are replaced with CAA compared to a reference wild type Factor IX sequence;
- c) at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG; and/or
- d) at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG and the remainder are CAA.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1, at least 2, at least 4, or at least 5 codons that encode asparagine is/are replaced with AAC compared to a reference wild type Factor IX sequence;
- b) at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC and the remainder are AAT; and/or
- d) the codons that encode asparagine are AAC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 7, at least 8, or at least 9 codons that encode lysine is/are replaced with AAG compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon AAG is/are replaced with AAA compared to a reference wild type Factor IX sequence;
- c) at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG; and/or
- d) at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG and the remainder are AAA.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 1, at least 2, at least 3, or at least 4 codons that encode aspartate is/are replaced with GAC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon GAC is/are replaced with GAT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC;
- d) at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC and the remainder are GAT; and/or
- e) the codons that encode aspartate are GAC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 15, at least 20, at least 25, or at least 26 codons that encode glutamate is/are replaced with GAG compared to a reference wild type Factor IX sequence;
- b) at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG; and/or
- c) at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG and the remainder are GAA.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 6, at least 7, or at least 8 codons that encode cysteine is/are replaced with TGC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon TGC is/are replaced with TGT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC;
- d) at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC and the remainder are TGT; and/or
- e) the codons that encode cysteine are TGC, except where the following codon starts with a G.

In an embodiment, in the portion of the coding sequence that is codon optimised in the portion of the coding sequence that is codon optimised the codons that encode tryptophan are TGG.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 8, at least 10, or at least 11 codons that encode arginine is/are replaced with AGG compared to a reference wild type Factor IX sequence;
- b) at least 1 codon that encodes arginine is/are replaced with AGA compared to a reference wild type Factor IX sequence;
- c) at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG; and/or
- d) at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG and the remainder are AGA.

Preferably at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG.

In an embodiment, in the portion of the coding sequence that is codon optimised:

- a) at least 5, at least 10, at least 12, or at least 13 codons that encode glycine is/are replaced with GGC compared to a reference wild type Factor IX sequence;
- b) at least 5, at least 6, at least 7, or at least 8 codons that encode glycine is/are replaced with GGG compared to

a reference wild type Factor IX sequence, where the following codon starts with a G;

- c) at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC;
- d) at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC and the remainder are GGG; and/or
- e) the codons that encode glycine are GGC, except where the following codon starts with a G.

In an embodiment, the portion of the coding sequence that is codon optimised comprises codons that encode phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine.

In an embodiment, the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine and glycine, and in the codon optimised portion:

- a) at least 5 codons that encode phenylalanine is/are replaced with TTC compared to a reference wild type Factor IX sequence;
- b) at least 16 codons that encode leucine is/are replaced with CTG compared to a reference wild type Factor IX sequence;
- c) at least 12 codons that encode isoleucine is/are replaced with ATC compared to a reference wild type Factor IX sequence;
- d) at least 25 codons that encode valine is/are replaced with GTG compared to a reference wild type Factor IX sequence;
- e) at least 13 codons that encode serine is/are replaced with AGC compared to a reference wild type Factor IX sequence;
- f) at least 5 codons that encode proline is/are replaced with CCC compared to a reference wild type Factor IX sequence;
- g) at least 10 codons that encode threonine is/are replaced with ACC compared to a reference wild type Factor IX sequence;
- h) at least 4 codons that encode alanine is/are replaced with GCC compared to a reference wild type Factor IX sequence;
- i) at least 2 codons that encode tyrosine is/are replaced with TAC compared to a reference wild type Factor IX sequence;
- j) at least 1 codons that encode histidine is/are replaced with CAC compared to a reference wild type Factor IX sequence;
- k) at least 5 codons that encode glutamine is/are replaced with CAG compared to a reference wild type Factor IX sequence;
- l) at least 5 codons that encode asparagine is/are replaced with AAC compared to a reference wild type Factor IX sequence;
- m) at least 9 codons that encode lysine is/are replaced with AAG compared to a reference wild type Factor IX sequence;
- n) at least 4 codons that encode aspartate is/are replaced with GAC compared to a reference wild type Factor IX sequence;
- o) at least 26 codons that encode glutamate is/are replaced with GAG compared to a reference wild type Factor IX sequence;

35

- p) at least 8 codons that encode cysteine is/are replaced with TGC compared to a reference wild type Factor IX sequence;
- q) the codons that encode tryptophan are TGG;
- r) at least 11 codons that encode arginine is/are replaced with AGG compared to a reference wild type Factor IX sequence; and
- s) at least 13 codons that encode glycine is/are replaced with GGC compared to a reference wild type Factor IX sequence.

In an embodiment, the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine and glycine, and in the codon optimised portion:

- a) at least 70% of the codons that encode phenylalanine are TTC;
- b) at least 94% of the codons that encode leucine are CTG;
- c) at least 75% of the codons that encode isoleucine are ATC;
- d) at least 95% of the codons that encode valine are GTG;
- e) at least 70% of the codons that encode serine are AGC;
- f) at least 60% of the codons that encode proline are CCC;
- g) at least 55% of the codons that encode threonine are ACC;
- h) at least 43% of the codons that encode alanine are GCC;
- i) at least 48% of the codons that encode tyrosine are TAC;
- at least 65% of the codons that encode histidine are CAC;
- k) at least 90% of the codons that encode glutamine are CAG;
- l) at least 70% of the codons that encode asparagine are AAC;
- m) at least 95% of the codons that encode lysine are AAG;
- n) at least 60% of the codons that encode aspartate are GAC;
- o) at least 95% of the codons that encode glutamate are GAG;
- p) at least 55% of the codons that encode cysteine are TGC;
- q) the codons that encode tryptophan are TGG;
- r) at least 75% of the codons that encode arginine are AGG; and
- s) at least 60% of the codons that encode glycine are GGC.

In an embodiment, the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine and glycine, and in the codon optimised portion:

- a) at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT;
- b) at least 94% of the codons that encode leucine are CTG and the remainder are CTC;
- c) at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT;
- d) at least 95% of the codons that encode valine are GTG;
- e) at least 70% of the codons that encode serine are AGC;
- f) at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT;
- g) at least 55% of the codons that encode threonine are ACC and the remainder are ACT;

36

- h) at least 43% of the codons that encode alanine are GCC and the remainder are GCT;
- i) at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT;
- j) at least 65% of the codons that encode histidine are CAC and the remainder are CAT;
- k) at least 90% of the codons that encode glutamine are CAG and the remainder are CAA;
- l) at least 70% of the codons that encode asparagine are AAC and the remainder are AAT;
- m) at least 95% of the codons that encode lysine are AAG and the remainder are AAA;
- n) at least 60% of the codons that encode aspartate are GAC and the remainder are GAT;
- o) at least 95% of the codons that encode glutamate are GAG and the remainder are GAA;
- p) at least 55% of the codons that encode cysteine are TGC and the remainder are TGT;
- q) the codons that encode tryptophan are TGG;
- r) at least 75% of the codons that encode arginine are AGG and the remainder are AGA; and
- s) at least 60% of the codons that encode glycine are GGC and the remainder are GGG.

The reference wild type Factor IX sequence may be SEQ ID NO. 9 or SEQ ID NO. 19.

The portion that is codon optimised can correspond to a sequence encoding part of or an entire Factor IX protein. For example, the Factor IX protein could be a full length coding sequence (such as a sequence encoding SEQ ID NO. 8 or SEQ ID NO. 16) or a variant thereof, and the entire coding sequence could be codon optimised. Hence, reference herein to "a portion of the coding sequence is codon optimised" should be understood to mean "at least a portion of the coding sequence is codon optimised". In some embodiments, however, a portion of the coding sequence is not codon optimised, for example a portion of the coding sequence is not codon optimised for expression in the liver. In some embodiments, the portion of the coding sequence that is codon optimised is at least 800, at least 900, at least 1100, less than 1500, less than 1300, less than 1200, between 800 and 1500, between 900 and 1300, between 1100 and 1200, or around 1191 nucleotides in length.

In an embodiment, the portion of the coding sequence that is codon optimised comprises exon 3 or a portion of at least 10, at least 15, at least 20, less than 25, between 10 and 25, between 15 and 25, or between 20 and 25 nucleotides of exon 3. In a further embodiment, the portion of the coding sequence that is codon optimised comprises exon 4 or a portion of at least 80, at least 90, at least 100, less than 114, between 80 and 114, between 90 and 114, or between 100 and 114 nucleotides of exon 4. In a further embodiment, the portion of the coding sequence that is codon optimised comprises exon 5 or a portion of at least 90, at least 100, at least 110, less than 129, between 90 and 129, between 100 and 129, or between 110 and 129 nucleotides of exon 5. In a further embodiment, the portion of the coding sequence that is codon optimised comprises exon 6 or a portion of at least 150, at least 180, at least 200, less than 203, between 150 and 203, between 180 and 203, or between 200 and 203 nucleotides of exon 6. In a further embodiment, the portion of the coding sequence that is codon optimised comprises exon 7 or a portion of at least 70, at least 80, at least 90, at least 100, less than 115, between 70 and 115, between 80 and 115, between 90 and 115, or between 100 and 115 nucleotides of exon 7. In a further embodiment, the portion of the coding sequence that is codon optimised comprises exon 8 or a portion of at least 400, at least 450, at least 500, less than

548, between 400 and 548, between 450 and 548, or between 500 and 548 nucleotides of exon 8.

Exon 3 comprises nucleotides 253-277 of wild type Factor IX (such as a Factor IX of SEQ ID NO. 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. Exon 4 comprises nucleotides 278-391 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. Exon 5 comprises nucleotides 392-520 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. Exon 6 comprises nucleotides 521-723 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. Exon 7 comprises nucleotides 724-838 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. Exon 8 comprises nucleotides 839-1386 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence.

Preferably a portion of at least 20 nucleotides of exon 3, a portion of at least 100 nucleotides of exon 4, a portion of at least 110 nucleotides of exon 5, a portion of at least 180 nucleotides of exon 6, a portion of at least 100 nucleotides of exon 7, and a portion of at least 500 nucleotides of exon 8 are codon optimised. The portion of the coding sequence that is codon optimised may comprise exon 3, exon 4, exon 5, exon 6, exon 7 and exon 8. In an embodiment, the portion of the coding sequence that is codon optimised comprises exon 3, exon 4, exon 5, exon 6, exon 7 and exon 8.

In an embodiment, the portion of the coding sequence that is codon optimised comprises a portion of exon 2, and the portion of exon 2 is less than 160, less than 150, less than 100, less than 75, less than 60, at least 20, at least 30, at least 40, at least 50, between 20 and 160, between 30 and 150, between 30 and 100, between 40 and 75, or around 56 nucleotides in length. Exon 2 comprises nucleotides 89-252 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. In a preferred embodiment, the portion of the coding sequence that is codon optimised comprises a portion of exon 2 that is between 30 and 100 nucleotides in length.

It is within the capabilities of the person skilled in the art to determine whether a portion of a sequence encoding a Factor IX protein or fragment thereof corresponds, for example, to exon 8 of wild type Factor IX. The person skilled in the art merely needs to perform a sequence alignment of the sequence encoding the Factor IX protein or fragment thereof with exon 8 using a suitable alignment algorithm such as that of Needleman and Wunsch described above, and determine whether at least part of the nucleotide sequence has greater than 90%, greater than 95%, or greater than 98% identity to exon 8 of SEQ ID NO. 9 (as described above, exon 8 of SEQ ID NO. 9 consists of nucleotides 839-1386 of SEQ ID NO.9).

As discussed above, providing a polynucleotide sequence comprising a coding sequence that is partially or wholly codon optimised can ensure that the encoded polypeptide is expressed at a high level. In one embodiment, a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to the reference wild type Factor IX sequence. The reference wild type Factor IX sequence may be SEQ ID NO: 9. In an embodiment, a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at

higher levels compared to a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO: 12 and a transcription regulatory element of SEQ ID NO: 7. In an embodiment, a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO: 18 and a transcription regulatory element of SEQ ID NO: 6.

In an embodiment, a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at a level at least 1.5, at least 2, at least 2.5, or at least 3 times greater than a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 12 or SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 7 or SEQ ID NO. 6. Optionally, a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at a level at least 1.5, at least 2, at least 2.5, or at least 3 times greater than a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 12 and a transcription regulatory element of SEQ ID NO. 7. Optionally, a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at a level at least 1.5, at least 2, at least 2.5, or at least 3 times greater than a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 6.

The skilled person may determine whether the Factor IX nucleotide sequence is expressed at higher levels compared to a reference sequence by transducing host cells with a viral particle comprising the Factor IX nucleotide sequence, and some cells with a vector comprising the reference sequence. The cells may be cultured under conditions suitable for expressing the Factor IX protein or fragment thereof encoded by the Factor IX nucleotide sequence, and the level of expressed Factor IX protein can be compared. The level of expressed Factor IX protein can be assessed using an ELISA such as that described in the section entitled "Factor IX protein or fragment thereof". Suitable host cells include cultured human liver cells, such as Huh 7 cells.

As discussed above, the presence of CpGs (i.e. CG dinucleotides) may reduce expression efficiency. This is because CpGs may be methylated, and their methylation may lead to gene silencing thereby reducing expression. For this reason, it is preferred that the portion of the coding sequence that is codon optimised comprises a reduced number of CpGs compared to a corresponding portion of a reference wild type Factor IX sequence. In a preferred embodiment, the portion of the coding sequence that is codon optimised comprises than 40, less than 20, less than 10, less than 5, or less than 1 CpG. Preferably, the portion of the coding sequence that is codon optimised is CpG free, i.e. contains no (0) CG dinucleotides.

In an embodiment, the portion of the coding sequence that is codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 800, at least 900, at least 1100, less than 1191, less than 1100, less than 1000, between 800 and 1191, between 900 and 1191, or around 1191 nucleotides of SEQ ID NO. 1. In an embodiment, the portion of the coding sequence that is codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 1. In an embodiment, the portion of the coding sequence that is codon optimised is at least 95% identical to a fragment of

between 900 and 1191 nucleotides of SEQ ID NO. 1. In an embodiment, the portion of the coding sequence that is codon optimised is at least 95%, or at least 98% identical to SEQ ID NO. 1.

The present invention provides a polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or a fragment thereof and the coding sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1 and a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15. Optionally, the sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, or at least 99.8% identical to SEQ ID NO. 1 is codon optimised.

Portion of the Coding Sequence that is not Codon Optimised

In an embodiment, the Factor IX nucleotide sequence comprises a portion that is not codon optimised. The portion that is not codon optimised may be a contiguous portion. Including a portion that is not codon optimised may improve expression of the coding sequence, as the portion that is not codon optimised may interact beneficially with other portions of the coding sequence such as an intron or a fragment of an intron. For example, the Factor IX nucleotide sequence may comprise an intron, or a fragment of an intron, and in such cases flanking the intron or the fragment of an intron with wild type Factor IX sequence may help to ensure correct splicing.

The portion that is not codon optimised is not modified to include a greater number of favoured codons compared to the wild type sequence. For example, the portion that is not codon optimised may comprise a similar number of favoured codons to a wild type sequence. The portion that is not codon optimised may comprise less than 50% of codons TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG. Optionally, the portion that is not codon optimised comprises less than 50%, less than 45%, or less than 40% codons selected from the group consisting of: TTC, CTG, ATC, GTG, GTC, AGC, CCC, ACC, GCC, TAC, CAC, CAG, AAC, AAA, AAG, GAC, TGC, AGG, GGC, and GAG.

Optionally, the portion that is not codon optimised is not codon optimised for expression in human liver cells. In an embodiment, the portion that is not codon optimised comprises substantially the same number of favoured codons as a corresponding portion of SEQ ID NO. 9. For example, the portion that is not codon optimised may comprise at least 90% of the number of favoured codons as a corresponding portion of SEQ ID NO. 9.

Optionally, the portion that is not codon optimised is at least 100, at least 150, at least 170, at least 190, less than 250, less than 225, less than 200, or around 195 nucleotides in length.

As discussed in more detail below, the Factor IX nucleotide sequence may comprise an intron or a fragment of an intron. In such cases, the intron or the fragment of an intron may be flanked by the portion that is not codon optimised, i.e. some of the portion that is not codon optimised may be adjacent to the 3' end of the intron or the fragment of an intron and some of the portion that is not codon optimised may be adjacent to the 5' end of the intron or the fragment of the intron. The intron or the fragment of an intron may be between exon 1 and exon 2. In such cases, it is advantageous

to include a portion that is not codon optimised which portion comprises a portion of exon 1 and a portion of exon 2.

Optionally, the portion that is not codon optimised comprises exon 1 or a portion of at least 60, at least 70, at least 80, between 60 and 88, between 70 and 88, or between 80 and 88 contiguous nucleotides of exon 1. Exon 1 comprises nucleotides 1-88 of wild type Factor IX (such as a Factor IX of SEQ ID NO: 9), or a corresponding sequence in a non-wild-type Factor IX nucleotide sequence. Part of exon 1 may encode the signal peptide region and the pro-peptide region. Optionally, the portion that is not codon optimised comprises or does not comprise the signal peptide and/or pro-peptide regions. Exon 1 may also comprise an additional non-coding stretch of 29 nucleotide at the 5' end. If the Factor IX nucleotide sequence comprises an intron or a fragment of an intron, it is preferable that the portion that is not codon optimised comprises a portion of exon 1 that is adjacent to the intron or the fragment of an intron. For example, if the intron or the fragment of an intron is between exon 1 and exon 2, it is preferable that the portion that is not codon optimised comprises a portion of exon 1 that corresponds to nucleotides 80-88, 70-88, 60-88, 40-88, or 20-88 of SEQ ID NO.9.

Optionally, the portion that is not codon optimised comprises a portion of at least 50, at least 75, at least 80, at least 90, at least 100, less than 140, less than 120, between 50 and 140, between 75 and 120, or around 107 nucleotides of exon 2. For example, if the intron or the fragment of an intron is between exon 1 and exon 2, it is preferable that the portion that is not codon optimised comprises a portion of exon 2 that corresponds to nucleotides 89-100, 89-120, 89-140, 89-160, 89-180, or 89-196 of SEQ ID NO.9.

The portion that is not codon optimised may comprise CpGs. For example, the portion that is not codon optimised may comprise the same number of CpGs as a corresponding portion of SEQ ID NO. 9. The portion that is not codon optimised may comprise at least 1, at least 1.5, or at least 2 CpGs per 100 nucleotides. The portion that is not codon optimised may comprise at least 1, at least 2, at least 3, between 1 and 5, between 2 and 5, or around 5 CpGs.

The portion that is not codon optimised may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 150, at least 175, less than 195, less than 190, or less than 180 of SEQ ID NO. 15 or SEQ ID NO: 2. The portion that is not codon optimised may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15 or SEQ ID NO: 2. For example, the portion that is not codon optimised may be at least 98% identical to SEQ ID NO. 15 or SEQ ID NO. 2.

The portion that is not codon optimised may be wild type. SEQ ID NO. 9 is an example of a wild type Factor IX nucleotide coding sequence. Thus, the portion that is not codon optimised may be at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a corresponding portion of SEQ ID NO: 9. The Factor IX Nucleotide Sequence May Comprise an Intron or a Fragment of an Intron

The Factor IX nucleotide sequence may comprise an intron or a fragment of an intron that interrupts the coding sequence. An intron is a sequence of nucleotides that is excised during the process of expression, and does not form part of the coding sequence.

A genomic wild type Factor IX nucleotide sequence comprises introns, that interrupt the Factor IX coding sequence. The presence of an intron may assist in maintaining a high level of expression of wild type Factor IX. Thus, it may be advantageous to include an intron, or at least a fragment of an intron, in a Factor IX nucleotide sequence of the invention. For example, the Factor IX nucleotide sequence may comprise an intron or a fragment of an intron that corresponds to intron 1 in wild type Factor IX. Suitably, the intron is a fragment of intron 1A of wild type Factor IX, such as SEQ ID NO: 3. It has been found that truncating the sequence of intron 1 causes expression of the Factor IX nucleotide sequence to be increased. It is thought that the truncation of intron 1 to form intron 1A may delete a repressor element in the intron. Truncation of the intron 1 sequence also results in the Factor IX nucleotide sequence being shorter which allows more efficient packaging of the Factor IX nucleotide sequence into a viral delivery system in gene therapy embodiments.

The fragment of an intron may be less than 500, less than 400, less than 350, less than 300, at least 100, at least 200, at least 250, at least 290, between 100 and 500, between 200 and 400, between 250 and 350, or around 299 nucleotides. The fragment of an intron may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 200, at least 250, or at least 290 nucleotides of SEQ ID NO. 3. The intron or fragment of an intron may be at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.3. For example, the intron or the fragment of an intron may be at least 95%, or at least 98% identical to SEQ ID NO.3.

Preferably, the intron or the fragment of an intron interrupts the portion that is not codon optimised i.e. the intron is 5' to a portion that is not codon optimised and 3' to a portion that is not codon optimised in the Factor IX nucleotide sequence. An intron is "flanked by" a sequence that is not codon optimised if the nucleotides immediately 3' and 5' of the intron or close to the 3' and 5' sections of the intron are not codon optimised. "Close to the intron" refers to within 1, within 2, within 3, within 4, within 5, within 6, within 7, within 8, within 8 or within 10 nucleotides of the intron. As discussed above, flanking the intron or the fragment of an intron with a nucleotide sequence that is not codon optimised may help to ensure correct splicing. Optionally, the intron or the fragment of an intron is flanked by at least 60, at least 70, at least 80, at least 90, or at least 100 nucleotides that are not codon optimised. For example, an intron is flanked by 60 nucleotides that are not codon optimised if 40 nucleotides that are immediately 3' of the intron and 20 nucleotides that are immediately 5' of the intron are not codon optimised, or if 30 nucleotides that are immediately 3' of the intron and 30 nucleotides that are immediately 5' of the intron are not codon optimised. Optionally, the intron or the fragment of an intron is flanked by between 110 and 120 nucleotides that are not codon optimised at the 5' end (e.g. immediately 5' of the intron) and between 100 and 110 nucleotides that are not codon optimised at the 3' end (e.g. immediately 3' of the intron).

The intron or the fragment of an intron may be positioned between portions of the coding sequence corresponding to exon 1 and exon 2 of a Factor IX nucleotide sequence. If the intron or the fragment of an intron corresponds to a fragment of intron 1 in wild type Factor IX, it is preferable that the intron or the fragment of an intron is between portions of the

coding sequence corresponding to exon 1 and exon 2 of a Factor IX nucleotide sequence.

The Polynucleotide May Further Comprise a Transcription Regulatory Element

5 The polynucleotide may comprise a transcription regulatory element.

In one embodiment, the transcription regulatory element is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 6. In an embodiment, the polynucleotide comprises a transcription regulatory element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO: 6. Optionally, the polynucleotide comprises a transcription regulatory element of SEQ ID NO: 6.

15 Any appropriate transcription regulatory element may be used, such as HLP2, HLP1, LP1, HCR-hAAT, ApoE-hAAT, and LSP, which are all liver specific transcription regulatory elements. These transcription regulatory elements are described in more detail in the following references: HLP1: McIntosh J. et al., Blood 2013 Apr. 25, 121(17):3335-44; LP1: Nathwani et al., Blood. 2006 Apr. 1, 107(7): 2653-2661; HCR-hAAT: Miao et al., Mol Ther. 2000; 1: 522-532; ApoE-hAAT: Okuyama et al., Human Gene Therapy, 7, 637-645 (1996); and LSP: Wang et al., Proc Natl Acad Sci USA. 1999 Mar. 30, 96(7): 3906-3910. The HLP2 transcription regulatory element has a sequence of SEQ ID NO: 6.

The transcription regulatory element may comprise a promoter and/or an enhancer, such as the promoter element and/or enhancer element from HLP2, HLP1, LP1, HCR-hAAT, ApoE-hAAT, and LSP. Each of these transcription regulatory elements comprises a promoter, an enhancer, and optionally other nucleotides.

In an embodiment, the transcription regulatory element comprises an enhancer which is the human apolipoprotein E (ApoE) hepatic locus control region (HCR; Miao et al (2000), Molecular Therapy 1(6):522), or a fragment thereof. In an embodiment, the transcription regulatory element comprises a fragment of the HCR enhancer which is a fragment of at least 80, at least 90, at least 100, less than 192, between 80 and 192, between 90 and 192, between 100 and 250, or between 117 and 192 nucleotides in length. Optionally, the fragment of the HCR enhancer is between 100 and 250 nucleotides in length.

45 A suitable HCR enhancer element fragment is described in SEQ ID NO. 13. Optionally, the transcription regulatory element comprises an enhancer that is at least 80, at least 90, at least 100, less than 192, between 80 and 192, between 90 and 192, between 100 and 250, or between 117 and 192 nucleotides in length and the enhancer comprises a polynucleotide sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical SEQ ID NO. 13. Optionally, the transcription regulatory element comprises an enhancer that is between 117 and 192 nucleotides in length and the enhancer comprises a polynucleotide sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical SEQ ID NO. 13. Optionally, the transcription regulatory element comprises an enhancer that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 90, at least 100, or at least 110 nucleotides of SEQ ID NO. 13. Optionally, the polynucleotide comprises an enhancer that is at least 80%, at least 85%, at least 90%, at least 95% at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13. Optionally, the polynucleotide

comprises an enhancer that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13. Optionally, the polynucleotide comprises an enhancer of SEQ ID NO. 13.

In an embodiment, the transcription regulatory element comprises a promoter which is a human alpha-1 anti-trypsin promoter (A1AT; Miao et al (2000), *Molecular Therapy* 1(6):522), or a fragment thereof. Optionally, a fragment of an A1AT promoter which is at least 100, at least 120, at least 150, at least 180, less than 255, between 100 and 255, between 150 and 225, between 150 and 300, or between 180 and 255 nucleotides in length. Optionally, the fragment of an A1AT promoter is between 150 and 300 nucleotides in length.

A suitable A1AT promoter fragment is described in SEQ ID NO. 14. Optionally, the transcription regulatory element comprises a promoter that is at least 100, at least 120, at least 150, at least 180, less than 255, between 100 and 255, between 150 and 300, or between 180 and 255 nucleotides in length and the promoter comprises a polynucleotide sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14. Optionally, the transcription regulatory element comprises a promoter that is between 180 and 255 nucleotides in length and the promoter comprises a polynucleotide sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14. Optionally, the polynucleotide comprises a promoter that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 120, or at least 150 nucleotides of SEQ ID NO. 14. Optionally, the polynucleotide comprises a promoter that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14. Optionally, the polynucleotide comprises a promoter that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14. Optionally, the polynucleotide comprises a promoter of SEQ ID NO. 14.

If the polynucleotide is intended for expression in the liver, the promoter may be a liver-specific promoter. Optionally, the promoter is a human liver-specific promoter.

A "liver-specific promoter" is a promoter that provides a higher level of expression in liver cells compared to other cells in general. For example, the skilled person can determine whether a promoter is a liver-specific promoter by comparing expression of the polynucleotide in liver cells (such as Huh 7 cells) with expression of the polynucleotide in cells from other tissues. If the level of expression is higher in the liver cells, compared to the cells from other tissues, the promoter is a liver-specific promoter.

Gain of Function Mutation

The Factor IX protein or fragment thereof may comprise a gain of function mutation. A gain of function mutation is a mutation that increases the activity of the Factor IX protein or fragment thereof. For example, the gain of function mutation may result in a Factor IX protein or fragment thereof that has an activity at least 1.5-fold, at least 2-fold, at least 2.5-fold, at least 3-fold, at least 4-fold, at least 5-fold, at least 6-fold, at least 6.5-fold, at least 7-fold, at least 7.5-fold, or at least 8-fold or more greater than wild type Factor IX (such as the Factor IX encoded by SEQ ID NO. 9 or SEQ ID NO. 19).

The Factor IX protein or fragment thereof may comprise a mutation at a position corresponding to position 384 of wild type Factor IX (corresponding to codon 384 of SEQ ID

NO. 9 or amino acid 384 of the immature polypeptide encoded by SEQ ID NO. 9). A mutation at a position corresponding to position 384 of wild type Factor IX may be a gain of function mutation. For example, replacement of arginine 384 with leucine can lead to a substantial increase in activity.

Whether or not a Factor IX protein comprises a mutation at a position corresponding to position 384 in Factor IX can be determined by aligning the Factor IX protein with SEQ ID NO. 16 using a suitable algorithm such as that of Needleman and Wunsch described above, and determining whether the amino acid that aligns to amino acid 384 (which is leucine in SEQ ID NO. 16) is an arginine residue. If the amino acid that aligns to amino acid 384 of SEQ ID NO. 16 is not an arginine residue then the Factor IX protein has a mutation at a position corresponding to position 384 of wild type Factor IX.

Whether or not a mutation is a gain of function mutation can be determined by comparing the activity of a Factor IX protein comprising the mutation with the activity of a reference Factor IX protein that is identical except for the putative gain of function mutation. The relative activities of these two proteins can be determined using a chromogenic assay such as that discussed under the heading "Factor IX protein or fragment thereof". If the activity of the Factor IX protein comprising the mutation is higher than the activity of the reference protein, the mutation is a gain of function mutation.

Accordingly, the Factor IX nucleotide sequence may comprise a codon that encodes a mutation at a position corresponding to position 384 in Factor IX. For example, the Factor IX nucleotide sequence may comprise a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX that is a small, hydrophobic amino acid. The small, hydrophobic amino acid may be alanine, leucine, isoleucine, glycine, or valine. For example, the small, hydrophobic amino acid may be alanine or leucine. Preferably the small, hydrophobic amino acid is leucine.

The codon that encodes a mutation at a position corresponding to position 384 in wild type Factor IX can be a codon that encodes leucine such as CTX, where X is any nucleotide. Preferably, X is C or G. The codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX may be CTC, such as in SEQ ID NO. 4. In alternative embodiments, the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX is TTG or CTG, such as in SEQ ID NO. 11 or SEQ ID NO. 26. For example, reference to SEQ ID NO: 1 herein may be replaced by reference to the corresponding portions of SEQ ID NOs: 26 or 11. In other words, SEQ ID NO:1 may be substituted at nucleotide 957 (C) with G, or at nucleotides 955 (C) and 957 (C) with T and G respectively.

The polynucleotide may comprise a Factor IX sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 1200, at least 1350, or at least 1650 nucleotides of SEQ ID NO. 5. For example, the Factor IX nucleotide sequence may be at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 5.

Suitably,

- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1; and
- (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.

Suitably,

- (i) the Factor IX nucleotide sequence comprises a coding sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
- (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and
- (iii) the polynucleotide comprises an enhancer element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13.

Suitably,

- (i) the Factor IX nucleotide sequence comprises a coding sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
- (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and
- (iii) the polynucleotide comprises a promoter element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14.

Suitably,

- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
- (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and
- (iii) the polynucleotide comprises a transcription regulatory element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 6.

Suitably,

- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
- (ii) the Factor IX nucleotide sequence comprises a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO: 2; and
- (iii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the

codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.

Suitably, the Factor IX nucleotide sequence comprises an intron or a fragment of an intron, and the fragment of an intron is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 3.

A Viral Particle Comprising the Polynucleotide

The invention further provides a viral particle comprising a recombinant genome comprising polynucleotides of the invention. For the purposes of the present invention, the term "viral particle" refers to all or part of a virion. For example, the viral particle comprises a recombinant genome and may further comprise a capsid. The viral particle may be a gene therapy vector. Herein, the terms "viral particle" and "vector" are used interchangeably. For the purpose of the present application, a "gene therapy" vector is a viral particle that can be used in gene therapy, i.e. a viral particle that comprises all the required functional elements to express a transgene, such as a Factor IX nucleotide sequence, in a host cell after administration.

Suitable viral particles include a parvovirus, a retrovirus, a lentivirus or a herpes simplex virus. The parvovirus may be an adeno-associated virus (AAV). The viral particle is preferably a recombinant adeno-associated viral (AAV) vector or a lentiviral vector. More preferably, the viral particle is an AAV viral particle. The terms AAV and rAAV are used interchangeably herein.

The genomic organization of all known AAV serotypes is very similar. The genome of AAV is a linear, single-stranded DNA molecule that is less than about 5,000 nucleotides in length. Inverted terminal repeats (ITRs) flank the unique coding nucleotide sequences for the non-structural replication (Rep) proteins and the structural (VP) proteins. The VP proteins (VP1, -2 and -3) form the capsid. The terminal 145 nt are self-complementary and are organized so that an energetically stable intramolecular duplex forming a T-shaped hairpin may be formed. These hairpin structures function as an origin for viral DNA replication, serving as primers for the cellular DNA polymerase complex. Following wild type (wt) AAV infection in mammalian cells the Rep genes (i.e. encoding Rep78 and Rep52 proteins) are expressed from the P5 promoter and the P19 promoter, respectively, and both Rep proteins have a function in the replication of the viral genome. A splicing event in the Rep ORF results in the expression of actually four Rep proteins (i.e. Rep78, Rep68, Rep52 and Rep40). However, it has been shown that the unspliced mRNA, encoding Rep78 and Rep52 proteins, in mammalian cells are sufficient for AAV vector production. Also in insect cells the Rep78 and Rep52 proteins suffice for AAV vector production.

The recombinant viral genome of the invention may comprise ITRs. It is possible for an AAV vector of the invention to function with only one ITR. Thus, the viral genome comprises at least one ITR, but, more typically, two ITRs (generally with one either end of the viral genome, i.e. one at the 5' end and one at the 3' end). There may be intervening sequences between the polynucleotide and one or more of the ITRs. The polynucleotide of the invention may be incorporated into a viral particle located between two regular ITRs or located on either side of an ITR engineered with two D regions.

AAV sequences that may be used in the present invention for the production of AAV vectors can be derived from the genome of any AAV serotype. Generally, the AAV serotypes have genomic sequences of significant homology at the amino acid and the nucleic acid levels, provide an identical

set of genetic functions, produce virions which are essentially physically and functionally equivalent, and replicate and assemble by practically identical mechanisms. For the genomic sequence of the various AAV serotypes and an overview of the genomic similarities see e.g. GenBank Accession number U89790; GenBank Accession number J01901; GenBank Accession number AF043303; GenBank Accession number AF085716; Chiorini et al, 1997; Srivastava et al, 1983; Chiorini et al, 1999; Rutledge et al, 1998; and Wu et al, 2000. AAV serotype 1, 2, 3, 3B, 4, 5, 6, 7, 8, 9, 10, 11 or 12 may be used in the present invention. The sequences from the AAV serotypes may be mutated or engineered when being used in the production of gene therapy vectors.

Optionally, an AAV vector comprises ITR sequences which are derived from AAV1, AAV2, AAV4 and/or AAV6. Preferably the ITR sequences are AAV2 ITR sequences. Herein, the term AAVx/y refers to a viral particle that comprises some components from AAVx (wherein x is a AAV serotype number) and some components from AAVy (wherein y is the number of the same or different serotype). For example, an AAV2/8 vector may comprise a portion of a viral genome, including the ITRs, from an AAV2 strain, and a capsid derived from an AAV8 strain.

In an embodiment, the viral particle is an AAV viral particle comprising a capsid. AAV capsids are generally formed from three proteins, VP1, VP2 and VP3. The amino acid sequence of VP1 comprises the sequence of VP2. The portion of VP1 which does not form part of VP2 is referred to as VP1 unique or VP1U. The amino acid sequence of VP2 comprises the sequence of VP3. The portion of VP2 which does not form part of VP3 is referred to as VP2 unique or VP2U. Preferably the capsid is an AAV5 capsid or a Mut C capsid. The Mut C capsid may have at least 96%, at least 98%, at least 99%, at least 99.5%, at least 99.8% identity or 100% identity to SEQ ID NO.10. The AAV capsid may have at least 96%, at least 98%, at least 99%, at 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 17. In an alternative embodiment, the capsid has a VP2U and/or VP3 of SEQ ID NO. 17 and a VP1U sequence having at least 96%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identity to SEQ ID NO: 20, SEQ ID NO: 21, SEQ ID NO: 22, SEQ ID NO: 23, SEQ ID NO: 24 or SEQ ID NO: 25.

A viral particle of the invention may be a "hybrid" particle in which the viral ITRs and viral capsid are from different parvoviruses, such as different AAV serotypes. Preferably, the viral ITRs and capsid are from different serotypes of AAV, in which case such viral particles are known as transcapsidated or pseudotyped. Likewise, the parvovirus may have a "chimeric" capsid (e. g., containing sequences from different parvoviruses, preferably different AAV serotypes) or a "targeted" capsid (e. g., a directed tropism).

In some embodiments, the recombinant AAV genome comprises intact ITRs, comprising functional terminal resolution sites (TRS). Such an AAV genome may contain one or two resolvable ITRs, i.e. ITRs containing a functional TRS at which site-specific nicking can take place to create a free 3' hydroxyl group which can serve as a substrate for DNA polymerase to unwind and copy the ITR. Preferably, the recombinant genome is single-stranded (i.e., it is packaged into the viral particle in a single-stranded form). Optionally, the recombinant genome is not packaged in self-complementary configuration, i.e. the genome does not comprise a single covalently-linked polynucleotide strand with substantial self-complementary portions that anneal in the viral particle. Alternatively, the recombinant genome may be packaged in "monomeric duplex" form. "Mono-

meric duplexes" are described in WO 2011/122950. The genome may be packaged as two substantially complementary but non-covalently linked polynucleotides which anneal in the viral particle.

The viral particle may further comprise a poly A sequence. The poly A sequence may be positioned downstream of the nucleotide sequence encoding a functional Factor IX protein. The poly A sequence may be a bovine growth hormone poly A sequence (bGHpA). The poly A sequence may be between 250 and 270 nucleotides in length.

The viral particle of the invention optionally expresses highly in host cells. For example, on transduction in Huh7 cells, the viral particle expresses Factor IX protein or a fragment thereof having a Factor IX activity greater than the activity of Factor IX protein expressed from a viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO: 12 and a transcription regulatory element of SEQ ID NO. 7 and/or a viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 6. Optionally, after transduction into a population of Huh7 cells, the viral particle expresses Factor IX protein, or a fragment thereof, having a Factor IX activity greater than the activity of Factor IX expressed from a comparable viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO: 12 and a transcription regulatory element of SEQ ID NO. 7 transduced into a comparable population of Huh7 cells. Optionally, after transduction into a population of Huh7 cells, the viral particle expresses Factor IX protein, or a fragment thereof, having a Factor IX activity greater than the activity of Factor IX expressed from a comparable viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO: 18 and a transcription regulatory element of SEQ ID NO. 6 transduced into a comparable population of Huh7 cells. In such embodiments, the term "comparable viral particle" refers to a viral particle that is the same as an AAV viral particle of the invention, except the comparable viral particle comprises a different Factor IX nucleotide sequence and a different transcription regulatory element (those of SEQ ID NO: 12 and SEQ ID NO: 7 or SEQ ID NO: 18 and SEQ ID NO: 6). Optionally, the activity is assessed using a chromogenic assay such as the chromogenic assay discussed above. In this case, however, the activity is not normalised for the Factor IX concentration, so the activity is a function of the level of expression as well as the inherent activity of the Factor IX protein.

Compositions, Methods and Uses

In a further aspect of the invention, there is provided a composition comprising the polynucleotide or vector/viral particle of the invention and a pharmaceutically acceptable excipient.

The pharmaceutically acceptable excipients may comprise carriers, diluents and/or other medicinal agents, pharmaceutical agents or adjuvants, etc. Optionally, the pharmaceutically acceptable excipients comprise saline solution. Optionally, the pharmaceutically acceptable excipients comprise human serum albumin.

The invention further provides a polynucleotide, vector/viral particle or composition of the invention for use in a method of treatment. Optionally the method of treatment comprises administering an effective amount of the polynucleotide or vector/viral particle of the invention to a patient.

The invention further provides a method of treatment comprising administering an effective amount of the polynucleotide or vector/viral particle of the invention to a patient.

The invention further provides use of the polynucleotide, vector/viral particle or composition of the invention in the manufacture of a medicament for use in a method of treatment. Optionally the method of treatment comprises administering an effective amount of the polynucleotide or vector/viral particle of the invention to a patient.

Optionally the method of treatment is a gene therapy. A “gene therapy” involves administering a vector/viral particle of the invention that is capable of expressing a transgene (such as a Factor IX nucleotide sequence) in the host to which it is administered.

Optionally, the method of treatment is a method of treating a coagulopathy such as haemophilia (for example haemophilia A or B) or Van Willebrands’ disease. Preferably, the coagulopathy is characterised by increased bleeding and/or reduced clotting. Optionally, the method of treatment is a method of treating haemophilia, for example haemophilia B. In some embodiments, the patient is a patient suffering from haemophilia B. Optionally the patient has antibodies or inhibitors to Factor IX. Optionally, the polynucleotide and/or vector/viral particle is administered intravenously. Optionally, the polynucleotide and/or vector/viral particle is for administration only once (i.e. a single dose) to a patient.

When haemophilia B is “treated” in the above method, this means that one or more symptoms of haemophilia are ameliorated. It does not mean that the symptoms of haemo-

philia are completely remedied so that they are no longer present in the patient, although in some methods, this may be the case. The method of treatment may result in one or more of the symptoms of haemophilia B being less severe than before treatment. Optionally, relative to the situation pre-administration, the method of treatment results in an increase in the amount/concentration of circulating Factor IX in the blood of the patient, and/or the overall level of Factor IX activity detectable within a given volume of blood of the patient, and/or the specific activity (activity per amount of Factor IX protein) of the Factor IX in the blood of the patient.

A “therapeutically effective amount” refers to an amount effective, at dosages and for periods of time necessary, to achieve the desired therapeutic result, such as raising the level of functional factor IX in a subject (so as to lead to functional factor IX production at a level sufficient to ameliorate the symptoms of haemophilia B).

Optionally, the vector/viral particle is administered at a dose of less than 1×10^{11} , less than 1×10^{12} , less than 5×10^{12} , less than 2×10^{12} , less than 1.5×10^{12} , less than 3×10^{12} , less than 1×10^{13} , less than 2×10^{13} , or less than 3×10^{13} vector genomes per kg of weight of patient. Optionally, the dose of vector/viral particle that is administered is selected such that the subject expresses Factor IX at an activity of 10%-90%, 20%-80%, 30%-70%, 25%-50%, 20%-150%, 30%-140%, 40%-130%, 50%-120%, 60%-110% or 70%-100% of the Factor IX activity of a non-haemophilic healthy subject.

TABLE 3

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
1	Codon optimised portion of TI-ACNP-FIX-GoF coding sequence	GAGGAGAAGTGCAGCTTTGAGGAGGCCAGGGAGGTGTTTGAGAACACT GAGAGGACCCTGAGTTCTGGAAGCAGTATGTGGATGGGGACCAGTGT GAGAGCAACCCCTGCCTGAATGGGGGCGAGCTGCAAGGATGACATCAAC AGCTATGAGTGTGTTGCCCTTTGGCTTTGAGGGCAAGAACGTGAG CTGGATGTGACCTGCAACATCAAGAATGGCAGATGTGAGCAGTTCTGC AAGAACTCTGCTGACAACAAGGTGGTGTGAGCTGCACTGAGGGCTAC AGGCTGGCTGAGAACCAGAAGAGCTGTGAGCCTGCTGTGCCATTCCCA TGTGGCAGAGTGTCTGTGAGCCAGACCAGCAAGCTGACCAGGGCTGAG GCTGTGTTCCCTGATGTGGACTATGTGAACAGCACTGAGGCTGAAACC ATCCTGGACAACATCACCCAGAGCACCCAGAGCTTCAATGACTTCAACC AGGGTGGTGGGGGGGAGGATGCCAAGCCTGGCCAGTTCCCTGGCAA GTGGTGC TGAATGGCAAGGTGGATGCCCTTCTGTGGGGCAGCATTGTG AATGAGAAGTGGATTGTGACTGCTGCCACTGTGTGGAGACTGGGGTG AAGATCACTGTGGTGGCTGGGGAGCACACATTGAGGAGACTGAGCAC ACTGAGCAGAAGAGGAATGTGATCAGGATCATCCCCACCACAACACTAC AATGCTGCCATCAACAAGTACAACCATGACATTGCCCTGTGGAGCTG GATGAGCCCCCTGGTGTGAAACAGCTATGTGACCCCATCTGCATTGCT GACAAGGAGTACACCAACATCTTCTGAAAGTTTGGCTCTGGCTATGTG TCTGGCTGGGGCAGGGTGTCCACAAGGGCAGGTCTGCCCTGGTGTG CAGTACCTGAGGGTGGCCCTGGTGGACAGGGCCACCTGCCTGCTCAGC ACCAAGTTCACCATCTACAACAACATGTTCTGTGCTGGCTTCCATGAG GGGGCAGGGACAGCTGCCAGGGGGACTCTGGGGGCCCCCATGTGACT GAGGTGGAGGGCACCAGCTTCTGACTGGCATCATCAGCTGGGGGGAG GAGTGTGCCATGAAGGGCAAGTATGGCATCTACACCAAGTCTCCAGA TATGTGAAC TGGATCAAGGAGAAGACCAAGCTGACCTGA
2	Wild type portion of TI-ACNP-FIX-GoF coding sequence, including intron	ATGCAGCGCGTGAACATGATCATGGCAGAATCACAGGCCTCATCACC ATCTGCCTTTTAGGATATCTACTCAGTGTGAATGTACAGGTTTGT CCTTTTTTAAAATACATTGAGTATGCTTGCTTTTAGATATAGAAATA TCTGATGCTGTCTTCTTCACTAAATTTGATTACATGATTTGACAGCA ATATTGAAGAGTCTAACAGCCAGCACGAGGTTGGTAAGTACTGTGGG AACATCACAGATTTTGGCTCCATGCCCTAAAGAGAAATTGGCTTTTCCAG ATTATTTGGATTAAAAACAAAGACTTTCTTAAGAGATGTAATAATTTT ATGATGTTTTCTTTTGTCTAAACTAAAGAATTATTCTTTTACATTT CAGTTTTTCTTGATCATGAAAACGCCAACAAAATTCTGAATCGGCCAA AGAGGTATAATTCAGGTAAATTGGAAGAGTTTGTTCAGGGGAACCTTG AGAGAGAATGTATG

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
3	Truncated FIX intron 1A	GTTTGTTCCTTTTTTAAATACATTGAGTATGCTTGCCTTTAGATA TAGAAATATCTGATGCTGTCTTCTTCACTAAATTTGATTACATGATT TGACAGCAATATTGAAGAGTCTAACAGCCAGCACGCAGGTTGGTAAGT ACTGTGGGAACATCACAGATTTTGGCTCCATGCCCTAAAGAGAAATTG GCTTTCAGATTATTTGGATTAAAAACAAAGACTTCTTAAAGAGATGTA AAATTTTCATGATGTTTTCTTTTTGCTAAAACAAAGAATTATTCTT TTACATTCAG
4	Coding sequence of TI-ACNP-FIX-GoF	ATGCAGCGCGTGAACATGATCATGGCAGAATCACCAGGCCTCATCACC ATCTGCCTTTTAGGATATCTACTCAGTGCTGAATGTACAGTTTTCTT GATCATGAAAACGCCAACAAAATCTGAATCGGCCAAAGAGGTATAAT TCAGGTAAATTGGAAGAGTTTGTTCAGGGAACCTTGAGAGAGAATGT ATGGAGGAGAAGTGCAGCTTTGAGGAGGCCAGGGAGGTGTTTGAGAAC ACTGAGAGGACCACTGAGTTCTGGAAGCAGTATGTGGATGGGGACCAG TGTGAGAGCAACCCCTGCCTGAATGGGGCAGCTGCAAGGATGACATC AACAGCTATGAGTGCTGGTGCCTTTGGCTTTGAGGGCAAGAAGTGT GAGCTGGATGTGACCTGCAACATCAAGAATGGCAGATGTGAGCAGTTC TGCAAGAACTCTGCTGACAACAAGGTGGTGTGCAGCTGCACTGAGGGC TACAGGCTGGCTGAGAACCAGAAGAGCTGTGAGCCTGCTGTGCCATT CCATGTGGCAGAGTGTCTGTGAGCCAGACCAGCAAGCTGACCAGGGCT GAGGCTGTGTTCCCTGATGTGGACTATGTGAACAGCACTGAGGCTGAA ACCATCCTGGACAACATCACCCAGAGCACCCAGAGCTTCAATGACTTC ACCAGGGTGGTGGGGGGGAGGATGCCAAGCCTGGCCAGTTCCTTGG CAAGTGGTGTGAATGGCAAGGTGGATGCCCTTCTGTGGGGCAGCATT GTGAATGAGAAGTGGATTGTGACTGCTGCCACTGTGTGAGACTGGG GTGAAGATCACTGTGGTGGCTGGGGAGCACAACATTGAGGAGACTGAG CACACTGAGCAGAAGAGGAATGTGATCAGGATCATCCCCACCACAAC TACAATGCTGCCATCAACAAGTACAACCATGACATTGCCCTGCTGGAG CTGGATGAGCCCTGGTGTGACAGCTATGTGACCCCATCTGCATT GCTGACAAGGAGTACCCAACATCTTCTGAAGTTTGGCTCTGGCTAT GTGTCTGGCTGGGGCAGGGTGTTCACAAGGGCAGGTCTGCCCTGGTG CTGCAGTACCTGAGGGTGCCTTGGTGGACAGGGCCACCTGCCTGCTC AGCACAAGTTCACCATCTACAACAACATGTTCTGTGCTGGCTTCCAT GAGGGGGCAGGGACAGCTGCCAGGGGGACTCTGGGGGCCCCCATGTG ACTGAGGTGGAGGGCACCAGCTTCTGACTGGCATCATCAGCTGGGGG GAGGAGTGTGCCATGAAGGGCAAGTATGGCATCTACACCAAAGTCTCC AGATATGTGAACCTGGATCAAGGAGAAGACCAAGCTGACCTGA
5	Coding sequence of TI-ACNP-FIX-GoF Factor IX sequence, including intron	ATGCAGCGCGTGAACATGATCATGGCAGAATCACCAGGCCTCATCACC ATCTGCCTTTTAGGATATCTACTCAGTGCTGAATGTACAGTTTTGTTT CCTTTTTTAAATACATTGAGTATGCTTGCCTTTTAGATATAGAAATA TCTGATGCTGTCTTCTTCACTAAATTTTATTACATGATTGACAGCA ATATTGAAGAGTCTAACAGCCAGCACGCAGGTTGGTAAGTACTGTGGG AACATCACAGATTTTGGCTCCATGCCCTAAAGAGAAATTGGCTTTTCCAG ATTATTTGGATTAAAAACAAAGACTTCTTAAAGAGATGTAATTTTTC ATGATGTTTTCTTTTTTGTAAAACAAAGAATTATTCTTTTACATTT CAGTTTTTCTTGATCATGAAAACGCCAACAAAATCTGAATCGGCCAA AGAGGTATAATTCAGGTAAATTGGAAGAGTTTGTTCAGGGGAACCTTG AGAGAGAATGTATGGAGGAGAAGTGCAGCTTTGAGGAGGCCAGGGAGG TGTTTGAGAACACTGAGAGGACCACTGAGTTCTGGAAGCAGTATGTGG ATGGGGACCAGTGTGAGAGCAACCCCTGCCTGAATGGGGCAGCTGCA AGGATGACATCAACAGCTATGAGTGCTGGTGCCTTTTGGCTTTGAGG GCAAGAACTGTGAGCTGGATGTGACCTGCAACATCAAGAATGGCAGAT GTGAGCAGTTCTGCAAGAACTCTGCTGACAACAAGGTGGTGTGAGCT GCACTGAGGGCTACAGGCTGGCTGAGAACCAGAAGAGCTGTGAGCCTG CTGTGCCATTCCATGTGGCAGAGTGTCTGTGAGCCAGACCAGCAAGC TGACCAGGGCTGAGGCTGTGTTCCCTGATGTGGACTATGTGAACAGCA CTGAGGCTGAAACCATCTGGACAACATCACCCAGAGCACCCAGAGCT TCAATGACTTCACCAGGGTGGTGGGGGGGAGGATGCCAAGCCTGGCC AGTTCCTTGGCAAGTGGTGTGAATGGCAAGGTGGATGCCCTTCTGTG GGGGCAGCATTGTGAATGAGAAGTGGATTGTGACTGCTGCCACTGTG TGGAGACTGGGGTGAAGATCACTGTGGTGGCTGGGGAGCACAACATTG AGGAGACTGAGCACACTGAGCAGAAGAGGAATGTGATCAGGATCATCC CCCACCACAACATAATGCTGCCATCAACAAGTACAACCATGACATTG CCCTGCTGGAGCTGGATGAGCCCTGGTGTGAACAGCTATGTGACCC CCATCTGCATTGCTGACAAGGAGTACCCAACATCTTCTGAAGTTTGG GCTCTGGCTATGTGTCTGGCTGGGGCAGGGTGTTCACAAGGGCAGGT CTGCCCTGGTGTGCTGAGTACCTGAGGGTGCCTTGGTGGACAGGGCCA CCTGCCCTGCTCAGCACCAGTTCACCATCTACAACAACATGTTCTGTG CTGGCTTCCATGAGGGGGCAGGGACAGCTGCCAGGGGGACTCTGGGG GCCCCATGTGACTGAGGTGGAGGGCACCAGCTTCTGACTGGCATCA TCAGCTGGGGGGAGGAGTGTGCCATGAAGGGCAAGTATGGCATCTACA CAAAGTCTCCAGATATGTGAACCTGGATCAAGGAGAAGACCAAGCTGA CCTGA

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
6	HLP2 transcription regulatory element sequence	ccCTAAATGGGCAAACATTGCAAGCAGCAAACAGCAAACACACAGCC CTCCCTGCCTGCTGACCTTGGAGCTGGGGCAGAGGTCAGACACCTCTC TGGGCCATGCCACCTCCAACCTGGACACAGGACGCTGTGGTTTCTGAG CCAGGGGGCGACTCAGATCCCAGCCAGTGGACTTAGCCCTGTGCT CCTCCGATAACTGGGGTGACCTTGGTTAATATTACCAGCAGCCTCCC CCGTGCCCCCTCTGGATCCACTGCTTAAATACGGACGAGGACAGGGCC CTGTCTCCTCAGCTTACGGACACCACCTGACCTGGGACAGTGAAT
7	LP1 transcription regulatory element sequence	CCCTAAATGGGCAAACATTGCAAGCAGCAAACAGCAAACACACAGCC CTCCCTGCCTGCTGACCTTGGAGCTGGGGCAGAGGTCAGACACCTCTC TGGGCCATGCCACCTCCAACATCCACTCGACCCCTTGAATTTTCGGT GGAGAGGAGCAGAGGTTGCTCTGGCGTGGTTTAGGT AGTGTGAGAG GGGAATGACT CCTTTCGGTA AGTGCAGTGG AAGCTGTACA CTGCCAGGC AAAGCGTCCG GGCAGCGTAG GCGGGCGACT CAGATCCCAG CCAGTGGACT TAGCCCTGT TTGCTCCTCC GATAACTGGG GTGACCTTGG TTAATATTCA CCAGCAGCCT CCCCCGTTGC CCCTCTGGAT CCACTGCTTA AATACGGACG AGGACAGGGC CCTGTCTCCT CAGCTTCAGG CACCACCACT GACCTGGGAC AGTGAAT
8	"Mature" Factor IX amino acid sequence encoded by SEQ ID NO. 4	YNSGKLEEFVQGNLERECMEEKCSFEEAREVFENTERTTEFWKQYVDG DQCESNPCLNGGSKDDINSYECWCPFGFEGKNCLELDVTCNINKGRCE QFCKNSADNKVVCSCTEGYRLAENQKSCEPAVFPFCGRVSVSQTSLT RAEAVFPDVDYVNSTEAETILDNI TQSTQSENDFTRVVGGEDAKPGQF PWQVVLNGKVDAFCCGSIVNEKWI VTAACHVETGVKI TVVAGEHNI EE TEHTEQKRNVI RII PHHNYNAAINKYNHDTALLELDEPLVLSYVTP CIADKEYTNI FLKEGSGYVSGWGRV FHKGRSALVLYLRVPLVDRATC LLSTKFTIYNMECAGEHEGGRDSCQDSSGGPHVTEVEGTSFLTGI IS WGEECAMKGYGIYTKVSRVYVNWIKKTKLT
9	Wild type Factor IX (Malmo B variant) coding sequence	ATGCAGCGCTGAACATGATCATGGCAGAATCACCAGGCCTCATCACC ATCTGCCTTTTAGGATATCTACTCAGTGTGAATGTACAGTTTTTCTT GATCATGAAAACGCCAACAAATTTCTGAATCGGCCAAAGAGGTATAAT TCAGGTAATTTGGAAGAGTTTGTTC AAGGGAACCTTGAGAGAGAATGT ATGGAAGAAAAGTGTAGTTTTGAAGAAGCAGCAGAGAAGTTTTGAAAAC ACTGAAAGAACAACGAATTTTGAAGCAGTATGTTGATGGAGATCAG TGTGAGTCCAATCCATGTTTAAATGGCGCAGTTGCAAGGATGACATT AATTCCTATGAATGTTGGTGTCCCTTTGGATTTGAAGGAAAGAACTGT GAATTAGATGTAACATGTAACATTAAGAATGGCAGATGCGAGCAGTTT TGTAATAATAGTGTGATAACAAGGTGGTTTGTCTCCTGTACTGAGGGA TATCGACTTGCAGAAAACCAGAAGTCTGTGAACCAGCAGTGCCATTT CCATGTGGAAGAGTTTCTGTTTCAAAAACCTTCTAAGCTCACCCGTGCT GAGGCTGTTTTTCTGATGTGGACTATGTAATTTCTACTGAAGCTGAA ACCATTTTGGATAACATCACTCAAAGCACCAATCATTTAATGACTTC ACTCGGTTGTTGGTGGAGAAGATGCCAAAACAGGTCAATTCCTTGG CAGGTGTTTTGAATGGTAAAGTTGATGATTCTGTGGAGGCTCTATC GTTAATGAAAATGGATTGTAAGTGTGCTGCCACTGTGTTGAACTGGT GTTAAAATTACAGTTGTCGCAGGTGAACATAATATTGAGGAGACAGAA CATAACAGACAAAAGCGAAATGTGATTGCAATTATTCCTCACCACAAC TACAATGCAGCTATTAATAAGTACAACCATGACATTGCCCTTCTGGAA CTGGACGAACCCCTAGTGCTAACAGCTACGTTACACCTATTTGCATT GCTGACAAGGAATACACGAACATCTTCTCAAATTTGGATCTGGCTAT GTAAGTGGCTGGGGAAGAGTCTTCCAAAAGGGAGATCAGCTTTAGTT CTTCAGTACCTTAGAGTCCACTTGTGACCGAGCCACATGCTTTCGA TCTACAAAGTTCAACATCTATAACAACATGTTCTGTGCTGGCTTCCAT GAAGGAGGTAGAGATTGATGCAAGGAGATAGTGGGGACCCCATGTT ACTGAAGTGAAGGGACAGTTTCTTAACTGGAATTATTAGCTGGGGT GAAGAGTGTGCAATGAAAGGCAATATGGAATATATACCAAGGTATCC CGGTATGTCAACTGGATTAAGGAAAAACAAAGCTCACTTAA
10	Mut C capsid polypeptide sequence	MAADGYLPDWLEDNLSEGIREWALKPGAPKPKANQQKQDDGRGLVLP GYKYLGFNGLDKGEPVNAADAAALEHDKAYDQQLQAGDNPYLRYNHA DAEFQERLQEDTSFGGNLGRAVFQAKKRVLEPLGLVEEGAATAPGKKR PVDQSPQEPDSSSGVGSKGQPARKRLNFGQTGDSVDPDQPLGEP AAPTSLGSNTMASGGGAPMADNNEGADGVGNSGNWHCDSQWLGRVI TTSTRWALPTYNNHLYKQISSQSGASNDNHYFGYSTPWGYEDFNRFH CHFSPRDWQRLINNNWGRFPKLSFKLENIQVKEVTQNDGTTTANL TSTVQVFTDSEYQLPYVLSAHQCLPPFPADVFMVPQYGYLTLNNGS QAVGRSSFYCLEYFPSQMLRTGNFQFSYTFEDVPFHSSYAHQS LDR LMNPLIDQYLYLNRQTGTTSGTTNQSRLLFQAGPQSMQLQARNWLP GPCYRQRLSKTANDNNNSNFPWTAASKYHLNDRSLVNP GPAMASHK DDEEKFPFMHGNLIFGKEGTTASNAELDNVMTDEEIRTTNPVATEQ YGTVANLQSSNTAPTRTVNDQ GALPGMVWQDRDVY LQGF IWAKI PH

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
		TDGHFHPSPLMGGEGLEKHPQPIMIKNTPVPANPPTTESPAKFASFIT QYSTGQVSVEIEWELQKENSKRWNPEIQYTSNYNKSXNVDFTVDTNGV YSEPRPIGTRYLTRL
11	FIXco coding sequence with TTG G oF codon	ATGCAGAGGGTGAACATGATCATGGCTGAGAGCCCTGGCCTGATCACC ATCTGCCTGCTGGGCTACCTGCTGTCTGCTGAGTGCACTGTGTTCTTG GACCATGAGAATGCCAACAAGATCCTGAACAGGCCCAAGAGATAACAAC TCTGGCAAGCTGGAGGAGTTTGTGCAGGGCAACCTGGAGAGGGAGTGC ATGGAGGAGAAGTGCAGCTTTGAGGAGGCCAGGGAGGTGTTTGAGAAC ACTGAGAGGACCACTGAGTTCTGGAAGCAGTATGTGGATGGGGACCAG TGTGAGAGCAACCCCTGCCTGAATGGGGCAGCTGCAAGGATGACATC AACAGCTATGAGTGCTGGTGCCCTTTGGCTTTGAGGGCAAGAAGTGT GAGCTGGATGTGACCTGCAACATCAAGAATGGCAGATGTGAGCAGTTC TGCAAGAAGTCTGCTGACAACAAGGTGGTGTGCAGCTGCACTGAGGGC TACAGGCTGGCTGAGAACCAGAAGAGCTGTGAGCCTGCTGTGCCATTC CCATGTGGCAGAGTGCTGTGAGCCAGACCAGCAAGCTGACCAGGGCT GAGGCTGTGTTCCCTGATGTGGACTATGTGAACAGCACTGAGGCTGAA ACCATCCTGGACAACATCACCCAGAGCACCCAGAGCTTCAATGACTTC ACCAGGGTGGTGGGGGGGAGGATGCCAAGCCTGGCCAGTTCCTCTGG CAAGTGGTGTGATGAATGGCAAGGTGGATGCCCTTCTGTGGGGCAGCATT GTGAATGAGAAGTGGATTGTGACTGCTGCCACTGTGTGGAGACTGGG GTGAAGATCACTGTGGTGGCTGGGGAGCACAACATTGAGGAGACTGAG CACACTGAGCAGAAGAGGAATGTGATCAGGATCATCCCCACCACAAC TACAATGCTGCCATCAACAAGTACAACCATGACATTGCCCTGCTGGAG CTGGATGAGCCCTGGTGTGACAGCTATGTGACCCCATCTGCATT GCTGACAAGGAGTACACCAACATCTTCTGAAGTTTGGCTCTGGCTAT GTGTCTGGCTGGGGCAGGGTGTTCACAAGGGCAGGTCTGCCCTGGTG CTGCAGTACCTGAGGGTGGCCCTGGTGGACAGGGCCACCTGCCTGTTG AGCACCAGTTTACCATCTACAACAACATGTTCTGTGCTGGCTTCCAT GAGGGGGCAGGGACAGCTGCCAGGGGACTCTGGGGGCCCATGTG ACTGAGGTGGAGGGCACCAGCTTCTGACTGGCATCATCAGCTGGGGG GAGGAGTGTGCCATGAAGGGCAAGTATGGCATCTACACCAAGTCTCC AGATATGTGAAGTGGATCAAGGAGAAGACCAAGCTGACCTGA
12	FIXco coding sequence	ATGCAGAGGGTGAACATGATCATGGCTGAGAGCCCTGGCCTGATCACC ATCTGCCTGCTGGGCTACCTGCTGTCTGCTGAGTGCACTGTGTTCTTG GACCATGAGAATGCCAACAAGATCCTGAACAGGCCCAAGAGATAACAAC TCTGGCAAGCTGGAGGAGTTTGTGCAGGGCAACCTGGAGAGGGAGTGC ATGGAGGAGAAGTGCAGCTTTGAGGAGGCCAGGGAGGTGTTTGAGAAC ACTGAGAGGACCACTGAGTTCTGGAAGCAGTATGTGGATGGGGACCAG TGTGAGAGCAACCCCTGCCTGAATGGGGCAGCTGCAAGGATGACATC AACAGCTATGAGTGCTGGTGCCCTTTGGCTTTGAGGGCAAGAAGTGT GAGCTGGATGTGACCTGCAACATCAAGAATGGCAGATGTGAGCAGTTC TGCAAGAAGTCTGCTGACAACAAGGTGGTGTGCAGCTGCACTGAGGGC TACAGGCTGGCTGAGAACCAGAAGAGCTGTGAGCCTGCTGTGCCATTC CCATGTGGCAGAGTGCTGTGAGCCAGACCAGCAAGCTGACCAGGGCT GAGGCTGTGTTCCCTGATGTGGACTATGTGAACAGCACTGAGGCTGAA ACCATCCTGGACAACATCACCCAGAGCACCCAGAGCTTCAATGACTTC ACCAGGGTGGTGGGGGGGAGGATGCCAAGCCTGGCCAGTTCCTCTGG CAAGTGGTGTGATGAATGGCAAGGTGGATGCCCTTCTGTGGGGCAGCATT GTGAATGAGAAGTGGATTGTGACTGCTGCCACTGTGTGGAGACTGGG GTGAAGATCACTGTGGTGGCTGGGGAGCACAACATTGAGGAGACTGAG CACACTGAGCAGAAGAGGAATGTGATCAGGATCATCCCCACCACAAC TACAATGCTGCCATCAACAAGTACAACCATGACATTGCCCTGCTGGAG CTGGATGAGCCCTGGTGTGACAGCTATGTGACCCCATCTGCATT GCTGACAAGGAGTACACCAACATCTTCTGAAGTTTGGCTCTGGCTAT GTGTCTGGCTGGGGCAGGGTGTTCACAAGGGCAGGTCTGCCCTGGTG CTGCAGTACCTGAGGGTGGCCCTGGTGGACAGGGCCACCTGCCTGAGG AGCACCAGTTTACCATCTACAACAACATGTTCTGTGCTGGCTTCCAT GAGGGGGCAGGGACAGCTGCCAGGGGACTCTGGGGGCCCATGTG ACTGAGGTGGAGGGCACCAGCTTCTGACTGGCATCATCAGCTGGGGG GAGGAGTGTGCCATGAAGGGCAAGTATGGCATCTACACCAAGTCTCC AGATATGTGAAGTGGATCAAGGAGAAGACCAAGCTGACCTGA
13	Enhancer element from HLP2	CCCTAAAATGGGCAACATTGCAAGCAGCAAACAGCAAACACACAGCC CTCCCTGCCTGCTGACCTTGGAGCTGGGGCAGAGGTGAGACACCTCTC TGGGCCATGCCACCTCCAAC
14	Promoter element from HLP2	GGGCGACTCAGATCCCAGCCAGTGGACTTAGCCCCGTGTTGCTCCTCC GATAACTGGGGTGCCTTGGTTAATATTACCAGCAGCCTCCCCGTT GCCCCCTGGATCCACTGCTTAAATACGGACGAGGACAGGGCCCTGTC TCCTCAGCTTACGGCACCACTGACCTGGGACAGTGAAT

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
15	Wild type portion of TI-ACNP-FIX-GoF coding sequence, excluding the intron	ATGCAGCGCGTGAACATGATCATGGCAGAATCACCAGGCCTCATCACC ATCTGCCTTTTAGGATATCTACTCAGTGCTGAATGTACAGTTTTTCTT GATCATGAAAACGCCAACAAAATTCTGAATCGGCCAAAGAGGTATAAT TCAGGTAAATTGGAAGAGTTTGTTC AAGGGAACCTTGAGAGAGAATGT ATG
16	"Immature" Factor IX amino acid sequence encoded by SEQ ID NO. 4	MQRVNMIMAESPLITICLLGYLLSAECTVFLDHENANKILNRPKRYN SGKLEEFVQGNLERECMEEEKCSFEEAREVFENTERTEFWKQYVDGDQ CESNPCLNGGSKDDINSYECWCPFGFEGKNCELDVTCNIKNGRCEQF CKNSADNKVVCSCTEGYRLAENQKSCPEAVPFPCGRVSVSQTSLKTRA EAVFPDQVYVNSTEAETILDNI TQSTQSENDFTRVVGGEDAKPGQFPW QVVLNGKVDAFCCGSIVNEKWI VTAACHVETGVKI TVVAGEHNIETE HTEQKRNVIIRIIPHHNYNAAINKYNHDIALLELDEPLVLSYVTPICI ADKEYTNIIFLKEGSGYVSGWGRVEHKGRSALVLYLRVPLVDRATCLL STKFTIYMNMECAGEHEGGRDSCQGDSSGGPHVTEVEGTSFLTGIISWG EECAMKGYGIYTKVSRVYVNIKEKTKLT
17	AAV5 capsid polypeptide sequence	MSFVDHPPDWLEEVGEGLEFLGLEAGPKPKPNQQHQDQARGLVLP YNYLGPNGLDRGEPVNRADDEVAREHDI SYNEQLEAGDNPYLKYNHAD AEFQEKLADDT SFGGNL GKAVFQA KKRVL E PFG LVEE GAKTAPT GKRI DDHFPKRKKARTEEDSKPSTSSDAEAGPSGSQQLQIPAPASSLGADT MSAGGGPLGDNNQGADGVGNASGDWHCDS TWMGDRVVTKSTRTWVLP SYNNHQYREIKSGSVDGSNANAYFGYSTPWGYFDNFREHSHWSPRDWQ RLINNYWGFRRSLRVKIFNIQVKEVTVQDSTTTIANLSTTVQVFTD DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS FFCLEYFPSKMLRTGNNEFFTYNFEEVPHSSFAPSQNLFKLANPLVD QYLYRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGR TQGNLGS VNRASVSAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTMI ENSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQSS TTAPATGTYNLQEI VPGSVWMERDVYLGQPIWAKI PETGAHFHPSAM GGFGLKHPPMMLIKNT PVPGNIT SFSDVPVSSFI TQYSTGQVTVEME WELKKENSKRWNP EIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRYL TRPL
18	Coding sequence of TI-codop-FIX-GoF Factor IX sequence, including intron	ATGCAGCGCGTGAACATGATCATGGCAGAATCACCAGGCCTCATCACC ATCTGCCTTTTAGGATATCTACTCAGTGCTGAATGTACAGTTTTGTTT CCTTTTAAATAACATGAGTATGCTTGCCTTTTAGATATAGAAATA TCTGATGCTGTCTTCTTCACTAAATTTGATTACATGATTTGACAGCA ATATTGAAGAGTCTAACAGCCAGCACGCAGGTTGGTAACTACTGTGGG AACATCACAGATTTTGGCTCCATGCCCTAAAGAGAAATTTGGCTTTCAG ATTATTTGGATTA AAAACAAGACTTTCTTAAGAGATGTA AAAATTTT ATGATGTTTTCTTTTGTCTAAACTAAAGAATTATTTCTTTACATTT CAGTTTTCTTGATCATGAAAACGCCAACAAAATCTGAATCGGCCAA AGAGGTATAATTCAGGTAATTTGGAAGAGTTTGTTC AAGGGAACCTTG AGAGAGAATGTATGGAGGAGAAGTGTCTTTTCGAGGAGCGAGAGAGG TTTTCGAGAATACTGAGCGAACCAACGAATCTGGAAACAATATGTGG ATGGCGACCAATGTGAATCTAATCCCTGCCTCAACGGTGGCTCATGCA AAGACGATATCAACAGCTACGAGTGTGGTGCCCTTTGGTTTCGAGG GAAAGAATTGCGAGCTTGATGTAACCTGTAACATTAAGAATGGGCGCT GCGAACAGTTTTGCAAGAACAGCGCCGACAATAAGGTCGTCGAGTT GTACCGAAGGCTATAGGCTTGACAGAAATCAGAAGAGTTGCGAGCCTG CTGTGCCGTTCCCATGTGGCAGAGTCAGTGTGTCCAACTAGCAAGC TGACAAGAGCAGAAGCCGTTTTCCCGATGTGGACTACGTGAATTCCA CTGAAGCCGAAACGATCCTGGACAATATCACACAGAGCACTCAGTCTT TCAACGACTTCACACGGGTTGTGGGAGGAGAGGACGCCAAAACCGGCC AGTTTCCTTGGCAAGTCGTTCTAACGGCAAGGTCGACGCTTTTGTG GAGGGAGTATTGTGAACGAGAAATGGATTGTCACCGCTGCTCATTGTG TTGAACTGGGGTAAAATCACTGTTGTGCGAGGAGAGCAATATCG AAGAGACAGAACACACCGAGCAGAAACGCAACGTTATTCGGATCATTC CACATCACAACTACAATGCTGCCATCAACAAGTACAACCACGACATTG CGCTGTGGAGTTGGATGAACCTCTCGTGCTCAACTCCTATGTGACCC CAATCTGCATAGCAGATAAGGAGTATACCAACATCTTCTGAAGTTG GGTCAGGTTATGTGTGAGGCTGGGGACGAGTGTTCATAAAGGGAGAT CAGCACTGGTGTGAGTATCTGCGCGTACCCTGGTGGATCGGGCTA CTTGCTGCTAAGCACAAAATCACCATCTACAACAACATGTTTTGTG CCGTTTTACGAAGGCGGACGGGACAGCTGTCAGGGAGATTCCGGAG GGCTCATGTACAGAGGTCGAGGGCACCTCTTTCTCACTGGGATTA TAAGCTGGGAGAGAATGCGCCATGAAAGGGAAGTACGGCATATACA CGAAAGTGTCTAGATACGTGAATTTGGATTAAGGAAAAGACCAAACCTGA CGTGA

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
19	Wild type Factor IX (Malmo B variant) coding sequence corresponding to mature FIX polypeptide	TATAATTCAGGTAAATTGGAAGAGTTTGTTC AAGGGAACCTTGAGAGA GAATGTATGGAAGAAAAGTGTAGTTTTGAAGAAGCACGAGAAGTTTTT GAAAACACTGAAAGAACAAC TGAATTTTGAAGCAGTATGTTGATGGA GATCAGTGTGAGTCCAATCCATGTTTAAATGGCGGCAGTTGCAAGGAT GACATTAATTCCTATGAATGTTGGTGTCCCTTTGGATTTGAAGGAAAG AACTGTGAATTAGATGTAACATGTAACATTAAGAATGGCAGATGCGAG CAGTTTTGTAAAAATAGTGTGATAACAAGGTGGTTTGTCTCTGTACT GAGGGATATCGACTTGCAGAAAACCAGAAGTCCCTGTGAACCAGCAGTG CCATTTCCATGTGGAAGAGTTTCTGTTTCCAAAACCTTAAGCTCACC CGTGCTGAGGCTGTTTTTCCATGATGTGGACTATGTAATTTCTACTGAA GCTGAAACCATTTTGGATAACATCACTCAAAGCACCCAAATCATTTAAT GACTTCACTCGGGTTGTTGGTGGAGAAGATGCCAAACCAGGTCAATTC CCTTGGCAGGTTGTTTTGAATGGTAAAGTTGATGCATTTCTGTGGAGGC TCTATCGTTAATGAAAAATGGATTGTAAGTCTGCCCACTGTGTTGAA ACTGGTGTAAAATTACAGTTGTCGCAGGTGAACATAATATGAGGAG ACAGAACATACAGAGCAAAAGCGAAATGTGATTGCAATTAATTCCTCAC CACAACTACAATGCAGCTATTAATAAGTACAACCATGACATTGCCCTT CTGGAACCTGGACGAACCCTTAGTGCTAAACAGCTACGTTACACCTATT TGCAATGCTGACAAGGAATACACGAACATCTTCCCAAATTTGGATCT GGCTATGTAAGTGGCTGGGGAAGAGTCTTCCACAAAGGGAGATCAGCT TTAGTTCCTCAGTACCTTAGAGTTCCACTTGTGACCGAGCCACATGT CTTCGATCTACAAAGTTCACCATCTATAACAACATGTTCTGTGCTGGC TTCCATGAAGGAGGTAGAGATTCAATGTCAGGAGATAGTGGGGACCC CATGTTACTGAAGTGAAGGGACCAGTTTCTTAAC TGAAT TATTAGC TGGGGTGAAGAGTGTGCAATGAAAGGCAAATATGGAATATATACCAAG GTATCCCGGTATGTCAACTGGATTAAGGAAAAACAAAGCTCACTTAA
20	AAV2-5 hybrid VP1u variant 1	MAADGYLPDWLEEVGEGLEFLGLEAGPPKPKPNQQHQDQARGLVLP YNYLGPNGLDRGEPVNRADDEVAREHDI SYNEQLEAGDNPYLKYNHAD AEFQEKLADDT SFGGNL GKAVFQAKKRVLEPFGLVEEGAKTAPTGRKI DDHFPKRKKARTEEDSKPSTSSDAEAGPSGSQQQLQI PAQPASSLGADT MSAGGGGPLGDNNQGADGVGNASGDWHCDSTWMDRVTVKSTRTWVLP SYNNHQYREIKSGSVDGSNANAYFGYSTPWGYFDENREHSHWSPRDWQ RLINNYWGFRRSLRVKIFNIQVKEVTVDSTTTIANNLSTVQVFTD DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS SFECLYFPSKMLRTGNNFEFTYNFEEVPHSSFAPSQNLFLKLANPLVD QYLYRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGRTOGWNLGS VNRASVSFAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTMI FNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQS TTAPATGTYNLQEI VPGSVWMERDVYLQGP IWAKI PETGAHFHPSPA GGFGLKHPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEME WELKKENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRYL TRPL
21	AAV2-5 hybrid VP1u variant 2	MAADGYLPDWLEDTLSEGIRQWKLKPGPPPPKPAERHKDDSRGLVLP GYNYLGPNGLDRGEPVNRADDEVAREHDI SYNEQLEAGDNPYLKYNHA DAEFQEKLADDT SFGGNL GKAVFQAKKRVLEPFGLVEEGAKTAPTGRKI IDHFPKRKKARTEEDSKPSTSSDAEAGPSGSQQQLQI PAQPASSLGAD TMSAGGGGPLGDNNQGADGVGNASGDWHCDSTWMDRVTVKSTRTWVLP PSYNNHQYREIKSGSVDGSNANAYEGYSTPWGYFDENREHSHWSPRDW QRLINNYWGFRRSLRVKIFNIQVKEVTVDSTTTIANNLSTVQVFTD DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS SFECLYFPSKMLRTGNNFEFTYNFEEVPHSSFAPSQNLFLKLANPLV DQYLYRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGRTOGWNLGS GVNRASVSFAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTMI IFNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQS STTAPATGTYNLQEI VPGSVWMERDVYLQGP IWAKI PETGAHFHPSPA MGFGLKHPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEME EWELKKENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRYL LTRPL
22	AAV2-5 hybrid VP1u variant 3	MAADGYLPDWLEDTLSEGIRQWKLKPGPPPPKPAERHKDDSRGLVLP GYKYLGPNGLDKGEVNRADDEVAREHDI SYNEQLEAGDNPYLKYNHA DAEFQEKLADDT SFGGNL GKAVFQAKKRVLEPFGLVEEGAKTAPTGRKI IDHFPKRKKARTEEDSKPSTSSDAEAGPSGSQQQLQI PAQPASSLGAD TMSAGGGGPLGDNNQGADGVGNASGDWHCDSTWMDRVTVKSTRTWVLP PSYNNHQYREIKSGSVDGSNANAYEGYSTPWGYFDENREHSHWSPRDW QRLINNYWGFRRSLRVKIFNIQVKEVTVDSTTTIANNLSTVQVFTD DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS SFECLYFPSKMLRTGNNFEFTYNFEEVPHSSFAPSQNLFLKLANPLV DQYLYRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGRTOGWNLGS GVNRASVSFAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTMI IFNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQS STTAPATGTYNLQEI VPGSVWMERDVYLQGP IWAKI PETGAHFHPSPA

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
		MGGFGLKHPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEM EWELKKENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRY LTRPL
23	AAV2-5 hybrid VP1u variant 4	MAADGYLPDWLEDTLSEGIQWKLKPGPPPKPAERHKDDSRGLVLP GYKYLGPFNGLDKGEPVNEADAAALEHDKAYDRQLDSGDNPYLKYNHA DAEFQEKLADDTSGGNGLKVAFQAKKRVLEPFGLVEEGAKTAPTGKR IDHFPKRKKARTEEDSKPSTSDAEAGPSGSQQLQIPAPASSLGAD TMSAGGGGGLGDNNGADGVGNASGDWHCDSTWMDRVTKSTRTWVL PSYNNHQYREIKSGSVDGSNANAYEGYSTPWGYFDENREHSHWSPRDW QRLINNYWGFRRSLRVKIFNIQVKEVTVQDSTTTIANNLSTVQVFT DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS SFECLEYFPKMLRTGNNFEFTYNFEEVPHSSFAPSQNLFKLANPLV DQYLRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGRTQGWNLGS GVNRASVSAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTM IFNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQS STTAPATGTYNLQEI VPGSVWMERDVYLQGPWAKIPEPGAHFHPSPA MGGFGLKHPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEM EWELKKENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRY LTRPL
24	AAV2-5 hybrid VP1u variant 5	MAADGYLPDWLEDTLSEGIQWKLKPGPPPKPAERHKDDSRGLVLP GYKYLGPFNGLDKGEPVNEADAAALEHDKAYDRQLDSGDNPYLKYNHA DAEFQERLKEDTSFGGNGLGRAVFQAKKRVLEPFGLVEEGAKTAPTGKR IDHFPKRKKARTEEDSKPSTSDAEAGPSGSQQLQIPAPASSLGAD TMSAGGGGGLGDNNGADGVGNASGDWHCDSTWMDRVTKSTRTWVL PSYNNHQYREIKSGSVDGSNANAYFGYSTPWGYFDNRFHSHWSPRDW QRLINNYWGFRRSLRVKIFNIQVKEVTVQDSTTTIANNLSTVQVFT DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS SFFCLEYFPKMLRTGNNFEFTYNFEEVPHSSFAPSQNLFKLANPLV DQYLRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGRTQGWNLGS GVNRASVSAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTM IFNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQS STTAPATGTYNLQEI VPGSVWMERDVYLQGPWAKIPEPGAHFHPSPA MGGFGLKHPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEM EWELKKENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRY LTRPL
25	AAV2-5 hybrid VP1u variant 6	MAADGYLPDWLEDTLSEGIQWKLKPGPPPKPAERHKDDSRGLVLP GYKYLGPFNGLDKGEPVNEADAAALEHDKAYDRQLDSGDNPYLKYNHA DAEFQERLKEDTSFGGNGLGRAVFQAKKRVLEPLGLVEEPVKTAPTGKR IDHFPKRKKARTEEDSKPSTSDAEAGPSGSQQLQIPAPASSLGAD TMSAGGGGGLGDNNGADGVGNASGDWHCDSTWMDRVTKSTRTWVL PSYNNHQYREIKSGSVDGSNANAYFGYSTPWGYFDNRFHSHWSPRDW QRLINNYWGFRRSLRVKIFNIQVKEVTVQDSTTTIANNLSTVQVFT DDYQLPYVVGNGTEGCLPAFPQVFTLPQYGYATLNRDNTENPTERS SFFCLEYFPKMLRTGNNFEFTYNFEEVPHSSFAPSQNLFKLANPLV DQYLRFVSTNNTGGVQFNKNLAGRYANTYKNWFPGPMGRTQGWNLGS GVNRASVSAFATTNRMELEGASYQVPPQPNGMTNNLQGSNTYALENTM IFNSQPANPGTTATYLEGNMLITSESETQPVNRVAYNVGGQMATNNQS STTAPATGTYNLQEI VPGSVWMERDVYLQGPWAKIPEPGAHFHPSPA MGGFGLKHPPMMLIKNTPVPGNITSFSDVPVSSFITQYSTGQVTVEM EWELKKENSKRWNPEIQYTNNYNDPQFVDFAPDSTGEYRTRPIGTRY LTRPL
26	FIXco coding sequence with CTG G oF codon	ATGCAGAGGGTGAACATGATCATGGCTGAGAGCCCTGGCCTGATCACC ATCTGCCTGCTGGGCTACCTGCTGCTGCTGAGTGCCTGTGTTCCCTG GACCATGAGAATGCCAACAAGATCCTGAACAGGCCCAAGAGATACAAC TCTGGCAAGCTGGAGGAGTTTGTGAGGGCAACCTGGAGAGGGAGTGC ATGGAGGAGAAGTGCAGCTTTGAGGAGGCCAGGGAGGTGTTTGAGAAC ACTGAGAGGACCACTGAGTTCTGGAAGCAGTATGTGGATGGGGACCAG TGTGAGAGCAACCCCTGCCTGAATGGGGCAGCTGCAAGGATGACATC AACAGCTATGAGTGTGTTGCCCTTTGGCTTTGAGGGCAAGAACTGT GAGCTGGATGTGACCTGCAACATCAAGAATGGCAGATGTGAGCAGTTC TGCAAGAATCTGCTGACAACAAGGTGGTGTGAGCAGCTGACTGAGGGC TACAGGCTGGCTGAGAACCAGAAGAGCTGTGAGCCTGCTGTGCCATTC CCATGTGGCAGAGTGTCTGTGAGCCAGACCAGCAAGCTGACCAGGGCT GAGGCTGTGTTCCCTGATGTGACTATGTGAACAGCACTGAGGCTGAA ACCATCCTGGACAACATCACCCAGAGCACCCAGAGCTTCAATGACTTC ACCAGGGTGGTGGGGGGGAGGATGCCAAGCCTGGCCAGTTCCTCTGG CAAGTGGTGTGAATGGCAAGGTGGATGCCTTCTGTGGGGGAGCATT GTGAAGATCACTGTGGTGGCTGGGGAGCACAACTTGAAGAGACTGAG CACACTGAGCAGAAGAGGAATGTGATCAGGATCATCCCCACCACAAC

TABLE 3-continued

Sequence Listing		
Sequence identity number	Sequence description	Nucleotide or Amino Acid Sequence
		TACAATGCTGCCATCAACAAGTACAACCATGACATTGCCCTGCTGGAG CTGGATGAGCCCCTGGTGCTGAACAGCTATGTGACCCCCATCTGCATT GCTGACAAGGAGTACACCAACATCTTCTGAAGTTTGGCTCTGGCTAT GTGTCTGGCTGGGGCAGGGTGTCCACAAGGGCAGGTCTGCCCTGGTG CTGCAGTACCTGAGGGTGGCCCTGGTGGACAGGGCCACCTGCCTGCTG AGCACCAAGTTCACCATCTACAACAACATGTTCTGTGCTGGCTTCCAT GAGGGGGCAGGGACAGCTGCCAGGGGGACTCTGGGGGCCCCCATGTG ACTGAGGTGGAGGGCACCAGCTTCTGACTGGCATCATCAGCTGGGGG GAGGAGTGTGCCATGAAGGGCAAGTATGGCATCTACACCAAAGTCTCC AGATATGTGAAGTGGATCAAGGAGAAGACCAAGCTGACCTGA

EXAMPLES

In the following examples, experiments were performed with recombinant AAV carrying the FIX transgene cassettes ssLP1.FIXco (FIXco herein), ssHLP2.TI-codop-FIX-GoF (HTFG herein) and ssHLP2.TI-ACNP-FIX-GoF (HTAG herein). SsHLP2.TI-codop-FIX-GoF is a version of the ssHLP2.TI-codop-FIX construct disclosed in WO2016/075473 modified to encode leucine (L) instead of arginine (R) at position 384 of the encoded FIX polypeptide. These cassettes are shown in FIG. 1A, FIG. 1B, and FIG. 1C. ssLP1.FIXco contains a fully codon-optimised FIX coding sequence (SEQ ID NO: 12) preceded 5' by an SV40 intron, with expression driven by the LP1 promoter (SEQ ID NO: 7). ssHLP2.TI-codop-FIX-GoF and ssHLP2.TI-ACNP-FIX-GoF share the structure of having the shorter HLP2 transcription regulatory element (SEQ ID NO. 6) 5' to a FIX coding sequence which is interrupted by a truncated version of the native intron 1A and in which the exon 1 and part of the exon 2 nucleotide sequence is wild type (non-codon-optimised), with the remainder of the coding sequence codon-optimised (SEQ ID NO. 18 for HTFG and SEQ ID NO. 5 for HTAG). The nucleotide sequence of the codon-optimised portions is what differs between the respective two constructs. Unlike the wild type FIX protein encoded by ssLP1.FIXco, ssHLP2.TI-codop-FIX-GoF and ssHLP2.TI-ACNP-FIX-GoF encode a hyper-active FIX having an arginine (R) to leucine (L) substitution at position 384 of the FIX polypeptide.

Example 1—Methods

AAV Vector Production and Quantification

1. AAV vector stocks were prepared by standard triple plasmid transfection of human embryonic kidney (HEK293) cells with a combination of plasmids consisting of a vector genome plasmid, an adenoviral helper plasmid, and a packaging plasmid containing AAV Rep and Cap (AAV8 or AAVMut C) functions. As the recombinant AAV particles contained a genome based on AAV2, and a capsid from serotype 8 or a synthetic capsid comprising portions from two serotypes (Mut C'; SEQ ID NO: 10), they are referred to as 'pseudotyped'.

2. Vectors were purified by density gradient centrifugation with iodixanol.

3. Vector genomes were titred by qPCR with primers directed to the promoter region.

In Vitro Transduction and Detection of FIX Expression

1. HUH7 cells were plated at 5×10^5 cells per well in 12-well plates.

2. Cells were then stimulated with mitomycin C for 1 hour before transduction with AAV particles carrying a FIX-encoding transgene cassette.

3. Five days after transduction, supernatant was collected and analysed for the level of FIX using a commercially available ELISA kit (Stago Asserachrom IX:Ag kit Ref #00943). Activity of FIX was analysed using the commercially available chromogenic kit from Quadrachem (Biophen FIX (6) kit Ref #221806).

4. Vector genome DNA was extracted using the Qiagen DNeasy Blood and Tissue kit (Ref #69506) according to the manufacturer's instructions, and quantified by qPCR.

30 Detection of FIX in vivo

1. Adult C57BL/6 mice were injected with 5×10^{10} vector genomes (vg) of AAV particles carrying a FIX-encoding transgene cassette via the tail vein (n=4 per group).

2. Two weeks after injection mice were anaesthetised and blood collected via cardiac puncture, added to sodium citrate (1/10 dilution), and centrifuged at $3000 \times g$ for 15 minutes at 4°C . to collect the plasma, which was frozen at -80°C . for analysis.

3. Liver was harvested and snap frozen in liquid nitrogen before storage at -80°C . for DNA extraction (Qiagen DNeasy Blood and Tissue kit, Ref #69506) for vector genome analysis.

4. The level of FIX present in murine plasma was determined using a FIX ELISA kit (Stago Asserachrom IX:Ag kit REF 00943). The activity of human FIX was determined using the commercially available chromogenic kit from Quadrachem (Biophen FIX (6) kit Ref #221806).

Example 2—Analysis of In Vitro FIX Trans Gene Expression Using ELISA

HUH7 (human hepatocyte) cells were cultured in standard cell culture conditions. FIX transgenes were expressed by treating HUH7 cells with mitomycin C for 1 hour then subsequently transducing with pseudotyped ssAAV2/Mut C. AAV particles were first generated by transfection of HEK293 cells with recombinant vector genome plasmid, in addition to AAV helper and packaging plasmids, and culturing for a further 48 hours. ssAAV2/Mut C vectors were purified from the HEK293 cells by density gradient centrifugation and iodixanol. Vector genomes were titred by qPCR utilizing primers directed towards the promoter region of the transgene expression cassette. FIX expression cassettes were compared to determine their relative ability to express a FIX transgene in vitro by measuring FIX levels 5 days post-transduction. Vectors being evaluated were ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF and ssAAV2/Mut

C.HLP2.TI-ACNP-FIX-GoF. FIX levels in the culture supernatant were analysed through the use of a commercially available ELISA kit (Stago Asserachrom IX:Ag kit Ref #00943). In two separate experiments (see FIG. 2D, FIG. 2E, and FIG. 2F and FIG. 4D and FIG. 4E) FIX expression levels—as derived from ELISA assays utilizing the supernatant of HUH7 cultured cells—were normalised against copies of the vector genome per cell following the harvesting of HUH7 cell DNA using the Qiagen DNeasy Blood and Tissue kit (Ref #69506).

5 days post-transduction with ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF and ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF at a MOI of 1×10^3 vector genomes (vg), FIX levels were greater in HUH7 cells transduced with ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF than in HUH7 cells transduced with ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF ($n=2$; FIG. 2A, FIG. 2B, and FIG. 2C, FIG. 3A, FIG. 3B, and FIG. 3C, and FIG. 4A, FIG. 4B, and FIG. 4C). Similarly, when identical FIX expression assays were performed in HUH7 cells with increased MOI (5×10^3 and 1×10^4), FIX levels were greater in cells transduced with ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF than in cells transduced with ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF ($n=2$; FIG. 2A, FIG. 2B, FIG. 2C, FIG. 3A, FIG. 3B, FIG. 3C and FIG. 4A, FIG. 4B, and FIG. 4C). When FIX expression levels were normalised against viral vector genome copies per cell, at each of the three MOIs tested (1×10^3 , 5×10^3 and 1×10^4) FIX levels were consistently higher in HUH7 cells transduced with ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF relative to ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF ($n=2$; FIG. 2D, FIG. 2E, and FIG. 2F and FIG. 4D and FIG. 4E). When the 5×10^3 transduction data from the experiments FIG. 2A, FIG. 2B, FIG. 2C, FIG. 2D, FIG. 2E, and FIG. 2F, FIG. 3A, FIG. 3B, and FIG. 3C, and FIG. 4A, FIG. 4B, FIG. 4C, FIG. 4D, and FIG. 4E is combined, it shows significantly superior expression from ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF relative to ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF (FIG. 5).

Example 3—Analysis of In Vitro FIX Trans Gene Activity

FIX activity was assessed by harvesting HUH7 cell supernatant and using the BIOPHEN Factor IX kit (Quadrachem #221806, #222101, #223201). Partially codon-optimised FIX transgenes (HLP2.TI-codop-FIX-GoF and HLP2.TI-ACNP-FIX-GoF) were compared in vitro to determine relative FIX activity following ssAAV2/Mut C transduction (at a MOI of 1×10^3 , 5×10^3 and 1×10^4) of HUH7 cells. Supernatant was isolated from the HUH7 cells 5 days post-transduction, and FIX activity was determined using the BIOPHEN Factor IX kit. Regardless of the MOI, greater mean FIX activity was observed in supernatant derived from cells transduced with ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF (FIG. 6A, FIG. 6B, and FIG. 6C, FIG. 7A, FIG. 7B, and FIG. 7C, and FIG. 8A, FIG. 8B, and FIG. 8C). When the 5×10^3 transduction data from the experiments of FIG. 6A, FIG. 6B, and FIG. 6C and FIG. 7A, FIG. 7B, and FIG. 7C is combined, it shows significantly superior expression from ssAAV2/Mut C.HLP2.TI-ACNP-FIX-GoF relative to ssAAV2/Mut C.HLP2.TI-codop-FIX-GoF (FIG. 9).

Example 4—Analysis of In Vivo FIX Transgene Expression Using ELISA

FIX transgenes were expressed in C57Bl/6 mice following transduction by tail-vein injection with 5×10^{10} vector

genomes (vg) of pseudotyped ssAAV2/8 vectors. AAV particles were first generated by transfection of HEK293 cells with recombinant vector genome plasmid, in addition to AAV helper and packaging plasmids, and culturing for a further 48 hours. ssAAV2/8 vectors were purified from the HEK293 cells by density gradient centrifugation and iodixanol. Vector genomes were titred by qPCR utilizing primers directed towards the promoter region of the transgene expression cassette. FIX expression cassettes LP1.FIXco, HLP2.TI-codop-FIX-GoF and HLP2.TI-ACNP-FIX-GoF were compared to determine their relative ability to express a FIX transgene.

In a first experiment involving ssAAV2/8.HLP2.TI-codop-FIX-GoF and ssAAV2/8.LP1.FIXco, 2 weeks post-dosing blood was collected from anaesthetised mice via cardiac puncture. Subsequently, plasma was isolated via the addition of sodium citrate (1/10 dilution) and centrifugation at $3000 \times g$ for 15 minutes at $4^\circ C$. Circulating levels of FIX were determined using a FIX ELISA kit (Stago Asserachrom IX:Ag kit Ref #00943).

FIX expression levels were normalised against copies of vector genome per cell following the harvesting of mouse liver. Normalised FIX expression levels were determined as being significantly higher ($p < 0.05$) after transduction with ssAAV2/8.HLP2.TI-codop-FIX-GoF relative to ssAAV2/8.LP1.FIXco ($n=4$ mice; FIG. 10).

In a further experiment the partially codon-optimised FIX transgenes (HLP2.TI-codop-FIX-GoF and HLP2.TI-ACNP-FIX-GoF) were compared in vivo to determine relative FIX expression following C57Bl/6 mouse transduction with ssAAV2/8. Concurrently, FIX expression was determined following transduction of C57Bl/6 mice with the scAAV2/8.LP1.FIXco vector. Plasma was isolated from the mice 3 weeks post-dosing, and FIX antigen levels were determined using the ELISA assay. Mean FIX expression levels were lowest in mice transduced with scAAV2/8.LP1.FIXco, whilst levels were greater in mice transduced with ssAAV2/8.HLP2.TI-codop-FIX-GoF ($n=4$ mice; FIG. 11A). Mice transduced with ssAAV2/8.HLP2.TI-ACNP-FIX-GoF had significantly greater FIX expression than mice transduced with ssAAV2/8.HLP2.TI-codop-FIX-GoF ($n=4$ mice; FIG. 11A). When FIX expression levels were normalised against viral vector genome copies per cell the trend in FIX expression was maintained, whereby ssAAV2/8.HLP2.TI-ACNP-FIX-GoF produces significantly more FIX than both ssAAV2/8.HLP2.TI-codop-FIX-GoF and scAAV2/8.LP1.FIXco ($n=4$; FIG. 11B). Furthermore, ssAAV2/8.HLP2.TI-codop-FIX-GoF exhibited greater FIX expression than scAAV2/8.LP1.FIXco ($n=4$ mice; FIG. 11B).

Example 5—Analysis of In Vivo FIX Transgene Activity

BIOPHEN Factor IX kit (Quadrachem #221806, #222101, #223201) is a chromogenic assay for measuring Factor IX activity in human citrated plasma or in Factor IX concentrates, using a manual chromogenic method.

In the presence of thrombin, phospholipids and calcium, first Factor XIa, supplied in the assay at a constant concentration and in excess, activates FIX, present in the tested sample, into FIXa, which forms an enzymatic complex with thrombin activated factor VIII:C, also supplied in the assay at a constant concentration and in excess, phospholipids (PLPs) and Calcium, that activates Factor X, present in the assay system, into Factor Xa. This activity is directly related to the amount of Factor IX, which is the limiting factor. Generated Factor Xa is then exactly measured by its specific

activity on Factor Xa chromogenic substrate (SXa-11). Factor Xa cleaves the substrate and releases pNA. The amount of pNA generated is directly proportional to the Factor IXa activity. Finally, there is a direct relationship between the amount of Factor IX in the assayed sample and the Factor Xa activity generated, measured by the amount of pNA released, determined by colour development at 405 nm.

The partially codon-optimised FIX transgenes (HLP2.TI-codop-FIX-GoF and HLP2.TI-ACNP-FIX-GoF) were compared in vivo to determine relative FIX activity following C57Bl/6 mouse transduction with ssAAV2/8. Concurrently, FIX activity was determined following transduction of C57Bl/6 mice with scAAV2/8.LP1.FIXco. Plasma was isolated from the mice 3 weeks post-dosing, and FIX activity was determined using the BIOPHEN Factor IX kit. Mean FIX activity was lowest in mice transduced with scAAV2/8.LP1.FIXco, whilst activity was greater in mice transduced with ssAAV2/8.HLP2.TI-codop-FIX-GoF (n=4 mice; FIG. 12). Mice transduced with ssAAV2/8.HLP2.TI-ACNP-FIX-GoF had significantly greater FIX activity than mice transduced with ssAAV2/8.HLP2.TI-codop-FIX-GoF (n=4 mice; FIG. 12).

Embodiments

The invention described herein also relates to the following aspects:

1. A polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or fragment thereof and wherein a portion of the coding sequence is not wild type.
2. The polynucleotide of aspect 1, wherein the portion of the coding sequence that is not wild type is codon optimised.
3. A polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or a fragment thereof and the coding sequence comprises:
 - (i) a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1; and
 - (ii) a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15.
4. The polynucleotide of aspect 3, wherein the sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, or at least 99.8% identical to SEQ ID NO. 1 is codon optimised.
5. A polynucleotide comprising a Factor IX nucleotide sequence, wherein the Factor IX nucleotide sequence encodes a Factor IX protein or fragment thereof and has at least 97%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 5.
6. The polynucleotide of aspect 5, wherein the Factor IX nucleotide sequence comprises a coding sequence and a portion of the coding sequence is codon optimised.
7. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises DNA or RNA.
8. The polynucleotide of any one of aspects 2, 4, 6 or 7, wherein the portion of the coding sequence that is codon optimised is a contiguous portion.
9. The polynucleotide of aspect 2, 4, 6, 7 or 8, wherein the portion of the coding sequence that is codon optimised is codon optimised for expression in the human liver.
10. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide

sequence is expressed in human liver cells at higher levels compared to a reference wild type Factor IX nucleotide sequence.

11. The polynucleotide of any one of aspects 2, 4, 6 or 7, wherein the portion of the coding sequence that is codon optimised is at least 800, at least 900, at least 1100, less than 1500, less than 1300, less than 1200, between 800 and 1500, between 900 and 1300, between 1100 and 1200, or around 1191 nucleotides in length.
12. The polynucleotide of any one of aspects 2, 4 or 6-11, wherein the portion of the coding sequence that is codon optimised comprises 1, 2, 3, 4, 5 or all of:
 - a) exon 3 or a portion of at least 10, at least 15, at least 20, less than 25, between 10 and 25, between 15 and 25, or between 20 and 25 nucleotides of exon 3;
 - b) exon 4 or a portion of at least 80, at least 90, at least 100, less than 114, between 80 and 114, between 90 and 114, or between 100 and 114 nucleotides of exon 4;
 - c) exon 5 or a portion of at least 90, at least 100, at least 110, less than 129, between 90 and 129, between 100 and 129, or between 110 and 129 nucleotides of exon 5;
 - d) exon 6 or a portion of at least 150, at least 180, at least 200, less than 203, between 150 and 203, between 180 and 203, or between 200 and 203 nucleotides of exon 6;
 - e) exon 7 or a portion of at least 70, at least 80, at least 90, at least 100, less than 115, between 70 and 115, between 80 and 115, between 90 and 115, or between 100 and 115 nucleotides of exon 7; and/or
 - f) exon 8 or a portion of at least 400, at least 450, at least 500, less than 548, between 400 and 548, between 450 and 548, or between 500 and 548 nucleotides of exon 8.
13. The polynucleotide of aspect 12, wherein the portion of the coding sequence that is codon optimised comprises a), b), c), d), e) and f).
14. The polynucleotide of aspect 12 or aspect 13, wherein the portion of the coding sequence that is codon optimised comprises a portion of at least 20 nucleotides of exon 3, a portion of at least 100 nucleotides of exon 4, a portion of at least 110 nucleotides of exon 5, a portion of at least 180 nucleotides of exon 6, a portion of at least 100 nucleotides of exon 7, and a portion of at least 500 nucleotides of exon 8.
15. The polynucleotide of any one of aspects 12-14, wherein the portion of the coding sequence that is codon optimised comprises exon 3, exon 4, exon 5, exon 6, exon 7, and exon 8.
16. The polynucleotide of any one of aspects 2, 4 or 6-15, wherein the portion of the coding sequence that is codon optimised comprises a portion of exon 2, and the portion of exon 2 is less than 160, less than 150, less than 100, less than 75, less than 60, at least 20, at least 30, at least 40, at least 50, between 20 and 160, between 30 and 150, between 30 and 100, between 40 and 75, or around 56 nucleotides in length.
17. The polynucleotide of any one of aspects 2, 4 or 6-16, wherein the portion of the coding sequence that is codon optimised comprises a portion of exon 2 that is between 30 and 100 nucleotides in length.
18. The polynucleotide of any one of aspects 2, 4 or 6-17, wherein the portion of the coding sequence that is codon optimised comprises a reduced number of CpGs compared to a corresponding portion of a reference wild type Factor IX sequence.

69

19. The polynucleotide of aspect 18, wherein the portion of the coding sequence that is codon optimised comprises less than less than 40, less than 20, less than 18, less than 10, less than 5, or less than 1 CpG.
20. The polynucleotide of aspect 18 or 19, wherein the portion of the coding sequence that is codon optimised is CpG free.
21. The polynucleotide of any one of aspects 2, 4 or 6-20, wherein, in the portion of the coding sequence that is codon optimised, at least 40%, at least 45%, at least 50%, at least 55%, at least 60%, at least 65%, at least 70%, or at least 73% of the codons are selected from the group consisting of:

a)
TTC;

b)
CTG;

c)
ATC;

d)
GTG;

e)
GTC;

f)
AGC;

g)
CCC;

h)
ACC;

i)
GCC;

j)
TAC;

k)
CAC;

l)
CAG;

m)
AAC;

n)
AAA;

o)
AAG;

p)
GAC;

q)
TGC;

r)
AGG;

s)
GGC;
and

t)
GAG.

22. The polynucleotide of any one of aspects 2, 4 or 6-21, wherein, in the portion of the coding sequence that is codon optimised:

70

- a) at least 1, at least 2, at least 4, or at least 5 codons that encode phenylalanine is/are replaced with TTC compared to a reference wild type Factor IX sequence;
- b) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT; and/or
- d) the codons that encode phenylalanine are TTC, except where the following codon starts with a G.
23. The polynucleotide of any one of aspects 2, 4 or 6-22, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 5, at least 10, at least 15, or at least 16 codons that encode leucine is/are replaced with CTG compared to a reference wild type Factor IX sequence;
- b) at least 90%, or at least 94% of the codons that encode leucine are CTG; and/or
- c) at least 90%, or at least 94% of the codons that encode leucine are CTG and the remainder are CTC.
24. The polynucleotide of any one of aspects 2, 4, 6-23, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 5, at least 10, at least 11, or at least 12 codons that encode isoleucine is/are replaced with ATC compared to a reference wild type Factor IX sequence;
- b) at least 1 of codon ATC is/are replaced with ATT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC;
- d) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT; and/or
- e) the codons that encode isoleucine are ATC, except where the following codon starts with a G.
25. The polynucleotide of any one of aspects 2, 4 or 6-24, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 10, at least 15, at least 20, or at least 25 codons that encode valine is/are replaced with GTG compared to a reference wild type Factor IX sequence;
- b) at least 1 codon that encodes valine is/are replaced with GTC compared to a reference wild type Factor IX sequence;
- c) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG; and/or
- d) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG and the remainder are GTC.
26. The polynucleotide of any one of aspects 2, 4 or 6-25, wherein, in the portion of the coding sequence that is codon optimised:
- a) at least 5, at least 10, at least 12, or at least 13 codons that encode serine is/are replaced with AGC compared to a reference wild type Factor IX sequence;
- b) at least 1, at least 2, or at least 4 codons that encode serine is/are replaced with TCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC; and/or
- d) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC and the remainder are TCT or TCC.

27. The polynucleotide of any one of aspects 2, 4 or 6-26, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1, at least 2, or at least 5 codons that encode proline is/are replaced with CCC compared to a reference wild type Factor IX sequence;
 - at least 1 codons that encode proline is/are replaced with CCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 50%, at least 55%, or at least 60% of the codons that encode proline are CCC;
 - at 50%, at least 55%, or at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT; and/or
 - the codons that encode proline are CCC, except where the following codon starts with a G.
28. The polynucleotide of any one of aspects 2, 4 or 6-27, wherein, in the portion of the coding sequence that is codon optimised:
- at least 6, at least 7, at least 8, or at least 10 codons that encode threonine is/are replaced with ACC compared to a reference wild type Factor IX sequence;
 - at least 1, or at least 2, codons that encode threonine is/are replaced with ACT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC;
 - at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC and the remainder are ACT; and/or
 - the codons that encode threonine are ACC, except where the following codon starts with a G.
29. The polynucleotide of any one of aspects 2, 4 or 6-28, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1, at least 2, at least 3, or at least 4 codons that encode alanine is/are replaced with GCC compared to a reference wild type Factor IX sequence;
 - at least 1, at least 2, or at least 3 codons that encode alanine is/are replaced with GCT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 35%, at least 40%, or at least 43% of the codons that encode alanine are GCC;
 - at least 35%, at least 40%, or at least 45% of the codons that encode alanine are GCC and the remainder are GCT; and/or
 - the codons that encode alanine are GCC, except where the following codon starts with a G.
30. The polynucleotide of any one of aspects 2, 4 or 6-29, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1, or at least 2 codons that encode tyrosine is/are replaced with TAC compared to a reference wild type Factor IX sequence;
 - at least 1 of codon TAC is/are replaced with TAT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC;
 - at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT; and/or
 - the codons that encode tyrosine are TAC, except where the following codon starts with a G.

31. The polynucleotide of any one of aspects 2, 4 or 6-30, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1 codons that encode histidine is/are replaced with CAC compared to a reference wild type Factor IX sequence;
 - at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC;
 - at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC and the remainder are CAT; and/or
 - the codons that encode histidine are CAC, except where the following codon starts with a G.
32. The polynucleotide of any one of aspects 2, 4 or 6-31, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1, at least 2, at least 4, or at least 5 codons that encode glutamine is/are replaced with CAG compared to a reference wild type Factor IX sequence;
 - at least 1 of codon CAG is/are replaced with CAA compared to a reference wild type Factor IX sequence;
 - at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG; and/or
 - at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG and the remainder are CAA.
33. The polynucleotide of any one of aspects 2, 4 or 6-32, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1, at least 2, at least 4, or at least 5 codons that encode asparagine is/are replaced with AAC compared to a reference wild type Factor IX sequence;
 - at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC;
 - at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC and the remainder are AAT; and/or
 - the codons that encode asparagine are AAC, except where the following codon starts with a G.
34. The polynucleotide of any one of aspects 2, 4 or 6-33, wherein, in the portion of the coding sequence that is codon optimised:
- at least 5, at least 7, at least 8, or at least 9 codons that encode lysine is/are replaced with AAG compared to a reference wild type Factor IX sequence;
 - at least 1 of codon AAG is/are replaced with AAA compared to a reference wild type Factor IX sequence;
 - at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG; and/or
 - at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG and the remainder are AAA.
35. The polynucleotide of any one of aspects 2, 4 or 6-34, wherein, in the portion of the coding sequence that is codon optimised:
- at least 1, at least 2, at least 3, or at least 4 codons that encode aspartate is/are replaced with GAC compared to a reference wild type Factor IX sequence;
 - at least 1 of codon GAC is/are replaced with GAT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC;
 - at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC and the remainder are GAT; and/or

- e) the codons that encode aspartate are GAC, except where the following codon starts with a G.
36. The polynucleotide of any one of aspects 2, 4 or 6-35, wherein, in the portion of the coding sequence that is codon optimised:
- at least 15, at least 20, at least 25, or at least 26 codons that encode glutamate is/are replaced with GAG compared to a reference wild type Factor IX sequence;
 - at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG; and/or
 - at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG and the remainder are GAA.
37. The polynucleotide of any one of aspects 2, 4, or 6-36, wherein, in the portion of the coding sequence that is codon optimised:
- at least 5, at least 6, at least 7, or at least 8 codons that encode cysteine is/are replaced with TGC compared to a reference wild type Factor IX sequence;
 - at least 1 of codon TGC is/are replaced with TGT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC;
 - at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC and the remainder are TGT; and/or
 - the codons that encode cysteine are TGC, except where the following codon starts with a G.
38. The polynucleotide of any one of aspects 2, 4, or 6-37, wherein, in the portion of the coding sequence that is codon optimised the codons that encode tryptophan are TGG.
39. The polynucleotide of any one of aspects 2, 4, or 6-38, wherein, in the portion of the coding sequence that is codon optimised:
- at least 5, at least 8, at least 10, or at least 11 codons that encode arginine is/are replaced with AGG compared to a reference wild type Factor IX sequence;
 - at least 1 codon that encodes arginine is/are replaced with AGA compared to a reference wild type Factor IX sequence;
 - at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG; and/or
 - at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG and the remainder are AGA.
40. The polynucleotide of any one of aspects 2, 4, or 6-39, wherein, in the portion of the coding sequence that is codon optimised:
- at least 5, at least 10, at least 12, or at least 13 codons that encode glycine is/are replaced with GGC compared to a reference wild type Factor IX sequence;
 - at least 5, at least 6, at least 7, or at least 8 codons that encode glycine is/are replaced with GGG compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC;
 - at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC and the remainder are GGG; and/or
 - the codons that encode glycine are GGC, except where the following codon starts with a G.
41. The polynucleotide of any one of aspects 2, 4, or 6-40,

- leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine.
42. The polynucleotide of any one of aspects 2, 4, or 6-41, wherein the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the codon optimised portion:
- at least 5 codons that encode phenylalanine is/are replaced with TTC compared to a reference wild type Factor IX sequence;
 - at least 16 codons that encode leucine is/are replaced with CTG compared to a reference wild type Factor IX sequence;
 - at least 12 codons that encode isoleucine is/are replaced with ATC compared to a reference wild type Factor IX sequence;
 - at least 25 codons that encode valine is/are replaced with GTG compared to a reference wild type Factor IX sequence;
 - at least 13 codons that encode serine is/are replaced with AGC compared to a reference wild type Factor IX sequence;
 - at least 5 codons that encode proline is/are replaced with CCC compared to a reference wild type Factor IX sequence;
 - at least 10 codons that encode threonine is/are replaced with ACC compared to a reference wild type Factor IX sequence;
 - at least 4 codons that encode alanine is/are replaced with GCC compared to a reference wild type Factor IX sequence;
 - at least 2 codons that encode tyrosine is/are replaced with TAC compared to a reference wild type Factor IX sequence;
 - at least 1 codons that encode histidine is/are replaced with CAC compared to a reference wild type Factor IX sequence;
 - at least 5 codons that encode glutamine is/are replaced with CAG compared to a reference wild type Factor IX sequence;
 - at least 5 codons that encode asparagine is/are replaced with AAC compared to a reference wild type Factor IX sequence;
 - at least 9 codons that encode lysine is/are replaced with AAG compared to a reference wild type Factor IX sequence;
 - at least 4 codons that encode aspartate is/are replaced with GAC compared to a reference wild type Factor IX sequence;
 - at least 26 codons that encode glutamate is/are replaced with GAG compared to a reference wild type Factor IX sequence;
 - at least 8 codons that encode cysteine is/are replaced with TGC compared to a reference wild type Factor IX sequence;
 - the codons that encode tryptophan are TGG;
 - at least 11 codons that encode arginine is/are replaced with AGG compared to a reference wild type Factor IX sequence; and
 - at least 13 codons that encode glycine is/are replaced with GGC compared to a reference wild type Factor IX sequence.

43. The polynucleotide of any one of aspects 2, 4, or 6-42, wherein the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the codon optimised portion:
- at least 70% of the codons that encode phenylalanine are TTC;
 - at least 94% of the codons that encode leucine are CTG;
 - at least 75% of the codons that encode isoleucine are ATC;
 - at least 95% of the codons that encode valine are GTG;
 - at least 70% of the codons that encode serine are AGC;
 - at least 60% of the codons that encode proline are CCC;
 - at least 55% of the codons that encode threonine are ACC;
 - at least 43% of the codons that encode alanine are GCC;
 - at least 48% of the codons that encode tyrosine are TAC;
 - at least 65% of the codons that encode histidine are CAC;
 - at least 90% of the codons that encode glutamine are CAG;
 - at least 70% of the codons that encode asparagine are AAC;
 - at least 95% of the codons that encode lysine are AAG;
 - at least 60% of the codons that encode aspartate are GAC;
 - at least 95% of the codons that encode glutamate are GAG;
 - at least 55% of the codons that encode cysteine are TGC;
 - the codons that encode tryptophan are TGG;
 - at least 75% of the codons that encode arginine are AGG; and
 - at least 60% of the codons that encode glycine are GGC.
44. The polynucleotide of any one of aspects 2, 4, or 6-43, wherein the portion of the coding sequence that is codon optimised comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the codon optimised portion:
- at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT;
 - at least 94% of the codons that encode leucine are CTG and the remainder are CTC;
 - at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT;
 - at least 95% of the codons that encode valine are GTG;
 - at least 70% of the codons that encode serine are AGC;
 - at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT;
 - at least 55% of the codons that encode threonine are ACC and the remainder are ACT;
 - at least 43% of the codons that encode alanine are GCC and the remainder are GCT;
 - at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT;
 - at least 65% of the codons that encode histidine are CAC and the remainder are CAT;
 - at least 90% of the codons that encode glutamine are CAG and the remainder are CAA;

- at least 70% of the codons that encode asparagine are AAC and the remainder are AAT;
 - at least 95% of the codons that encode lysine are AAG and the remainder are AAA;
 - at least 60% of the codons that encode aspartate are GAC and the remainder are GAT;
 - at least 95% of the codons that encode glutamate are GAG and the remainder are GAA;
 - at least 55% of the codons that encode cysteine are TGC and the remainder are TGT;
 - the codons that encode tryptophan are TGG;
 - at least 75% of the codons that encode arginine are AGG and the remainder are AGA; and
 - at least 60% of the codons that encode glycine are GGC and the remainder are GGG.
45. The polynucleotide of any one of aspects 10-44, wherein the reference wild type Factor IX sequence is SEQ ID NO. 9 or SEQ ID NO. 19.
46. The polynucleotide of any one of aspects 2, 4 or 6-45, wherein the portion of the coding sequence that is codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 800, at least 900, at least 1100, less than 1191, less than 1100, less than 1000, between 800 and 1191, between 900 and 1191, or around 1191 nucleotides of SEQ ID NO. 1.
47. The polynucleotide of aspect 46, wherein the portion of the coding sequence that is codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 1.
48. The polynucleotide of aspect 46 or 47, wherein the portion of the coding sequence that is codon optimised is at least 95% identical to a fragment of between 900 and 1191 nucleotides of SEQ ID NO. 1.
49. The polynucleotide of any one of aspects 46-48, wherein the portion of the coding sequence that is codon optimised is at least 95%, or at least 98% identical to SEQ ID NO. 1.
50. The polynucleotide of any one of the preceding aspects, wherein the coding sequence comprises a portion that is not codon optimised.
51. The polynucleotide of aspect 50, wherein the portion that is not codon optimised is at least 100, at least 150, at least 170, at least 190, less than 250, less than 225, less than 200, or around 195 nucleotides.
52. The polynucleotide of any one of aspects 50 or 51, wherein the portion that is not codon optimised comprises exon 1 or a portion of at least 60, at least 70, at least 80, between 60 and 88, between 70 and 88, or between 80 and 88 nucleotides of exon 1.
53. The polynucleotide of any one of aspects 50-52, wherein the portion that is not codon optimised comprises a portion of at least 50, at least 75, at least 80, at least 90, at least 100, less than 140, less than 120, between 50 and 140, between 75 and 120, or around 107 nucleotides of exon 2.
54. The polynucleotide of any one of aspects 50-53, wherein the portion that is not codon optimised comprises CpGs.
55. The polynucleotide of aspect 54, wherein the portion that is not codon optimised comprises at least 1 or at least 2 CpGs per 100 nucleotides.

56. The polynucleotide of any one of aspects 50-55, wherein the portion that is not codon optimised comprises less than 50%, less than 45%, less than 40%, or less than 35% codons selected from the group consisting of:

- a)
TTC;
- b)
CTG;
- c)
ATC;
- d)
GTG;
- e)
GTC;
- f)
AGC;
- g)
CCC;
- h)
ACC;
- i)
GCC;
- j)
TAC;
- k)
CAC;
- l)
CAG;
- m)
AAC;
- n)
AAA;
- o)
AAG;
- p)
GAC;
- q)
TGC;
- r)
AGG;
- s)
GGC;
and
- t)
GAG.

57. The polynucleotide of any one of aspects 50-56, wherein the portion that is not codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 150, at least 175, less than 195, less than 190, or less than 180 nucleotides of SEQ ID NO. 15.

58. The polynucleotide of aspect 57, wherein the portion that is not codon optimised is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 15.

59. The polynucleotide of any one of aspects 50-58, wherein the portion that is not codon optimised is wild type.

60. The polynucleotide of any one of aspects 50-59, wherein the portion that is not codon optimised is at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO: 15.

61. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide further comprises an intron or a fragment of an intron that interrupts the coding sequence.

62. The polynucleotide of aspect 61, wherein the intron or the fragment of an intron is a portion of a wild type Factor IX intron.

63. The polynucleotide of aspect 61 or 62, wherein the fragment of an intron is less than 500, less than 400, less than 350, less than 300, at least 100, at least 200, at least 250, at least 290, between 100 and 500, between 200 and 400, between 250 and 350, or around 299 nucleotides.

64. The polynucleotide of any one of aspects 61-63, wherein the fragment of an intron is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of at least 100, at least 200, at least 250, or at least 290 nucleotides of SEQ ID NO. 3.

65. The polynucleotide of any one of aspects 61-64, wherein the intron or the fragment of an intron is at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.3.

66. The polynucleotide of aspect 65, wherein the intron or the fragment of an intron is at least 95%, or at least 98% identical to SEQ ID NO.3.

67. The polynucleotide of any one of aspects 61-66, wherein the intron or the fragment of an intron interrupts the portion that is not codon optimised.

68. The polynucleotide of aspect 67, wherein the intron or the fragment of an intron is flanked by at least 60, at least 70, at least 80, at least 90, or at least 100 nucleotides that are not codon optimised.

69. The polynucleotide of aspect 68, wherein the intron or the fragment of an intron is flanked by between 110 and 120 nucleotides that are not codon optimised at the 5' end and between 100 and 110 nucleotides that are not codon optimised at the 3' end.

70. The polynucleotide of any one of aspects 61-69, wherein the intron or the fragment of an intron is positioned between exon 1 and exon 2.

71. The polynucleotide of any one of aspects 61-70, wherein the intron or the fragment of the intron is a fragment of native intron 1 (intron 1a).

72. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide further comprises a transcription regulatory element.

73. The polynucleotide of aspect 72, wherein the transcription regulatory element comprises a liver-specific promoter.

74. The polynucleotide of aspect 72 or aspect 73, wherein the transcription regulatory element comprises an A1AT promoter or a fragment of an A1AT promoter.

75. The polynucleotide of aspect 74, wherein the fragment of an A1AT promoter is at least 100, at least 120, at least 150, at least 180, less than 255, between 100 and 255, between 150 and 225, between 150 and 300, or between 180 and 255 nucleotides in length.

76. The polynucleotides of aspect 75, wherein the fragment of an A1AT promoter is between 150 and 300 nucleotides in length.

77. The polynucleotides, of any one of aspects 72-76, wherein the transcription regulatory element comprises an enhancer.
78. The polynucleotide of aspect 77, wherein the enhancer is an HCR enhancer or a fragment of an HCR enhancer.
79. The polynucleotide of aspect 78, wherein the fragment of an HCR enhancer is a fragment of at least 80, at least 90, at least 100, less than 192, between 80 and 192, between 90 and 192, between 100 and 250, or between 117 and 192 nucleotides in length.
80. The polynucleotide of aspect 79, wherein the fragment of an HCR enhancer is between 100 and 250 nucleotides in length.
81. The polynucleotide of any one of aspects 72-80, wherein the transcription regulatory element is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 6.
82. The polynucleotide of aspect 81, wherein the transcription regulatory element has a sequence of SEQ ID NO. 6.
83. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises an enhancer that is at least 80%, at least 85%, at least 90%, at least 95% at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13.
84. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises an enhancer that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 13.
85. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises an enhancer of SEQ ID NO. 13.
86. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a promoter that is at least 80%, at least 85%, at least 90%, at least 95% at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14.
87. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a promoter that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14.
88. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a promoter of SEQ ID NO. 14.
89. The polynucleotide of any one of the preceding aspects, wherein the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to codon 384 of wild type factor IX, and wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes alanine or leucine.
90. The polynucleotide of aspect 89, wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX is CTX, wherein X is any nucleotide.
91. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a Factor IX nucleotide sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment at least 1200, at least 1350, or at least 1650 nucleotides of SEQ ID NO. 5.

92. The polynucleotide of any one of the preceding aspects, wherein the polynucleotide comprises a Factor IX nucleotide sequence that is at least 80%, at least 85%, at least 90%, at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.5.
93. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1; and
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.
94. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a coding sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and
 - (iii) the polynucleotide comprises a promoter element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 14 and/or an enhancer element that is at least 98%, at least 99%, at least 99.5%, at least 99.8% or 100% identical to SEQ ID NO. 13.
95. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine; and
 - (iii) the polynucleotide comprise a transcription regulatory element that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 6.
96. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a sequence that is at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO.1;
 - (ii) the Factor IX nucleotide sequence comprises a sequence that is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a corresponding portion of SEQ ID NO: 2; and
 - (iii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.

97. The polynucleotide of any one of aspects 95 or 96, wherein the Factor IX nucleotide sequence comprises an intron or a fragment of an intron, and the fragment of an intron is at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 3.
98. The polynucleotide of any one of the preceding aspects, wherein:
- (i) the Factor IX nucleotide sequence comprises a coding sequence and a portion of the coding sequence is not codon optimised; and
 - (ii) the Factor IX nucleotide sequence comprises a codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX wherein the codon that encodes an amino acid at a position corresponding to position 384 of wild type Factor IX encodes leucine.
99. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 12 and a transcription regulatory element of SEQ ID NO. 7.
100. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 6.
101. The polynucleotide of any one of the preceding aspects, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at a level at least 2, or at least 3 times greater than a polypeptide encoded by a nucleotide sequence comprising a Factor IX nucleotide sequence of SEQ ID NO. 12 or SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 7 or SEQ ID NO. 6.
102. A viral particle comprising a recombinant genome comprising the polynucleotide of any one of the preceding aspects.
103. The viral particle of aspect 102, which is an AAV, adenoviral, or lentiviral viral particle.
104. The viral particle of aspect 103, which is an AAV viral particle.
105. The viral particle of any one of aspects 102-104, wherein the recombinant genome further comprises:
- a) AAV2 ITRs;
 - b) a poly A sequence;
 - c) an origin of replication; and/or
 - d) two resolvable ITRs.
106. The viral particle of aspect 105, wherein the recombinant genome is single-stranded and/or comprises two resolvable ITRs.
107. The viral particle of any one of aspects 102-106, wherein the viral particle comprises a capsid selected from the group consisting of:
- (i) a capsid having at least 96%, at least 98%, at least 99%, at least 99.5%, at least 99.8% identity or 100% identity to SEQ ID NO.10;
 - (ii) a capsid having at least 96%, at least 98%, at least 99%, at 99.5%, at least 99.8%, or 100% identity to SEQ ID NO. 17;
 - (iii) AAVMutC; and
 - (iv) AAV5.

108. The viral particle of any one of aspects 102-107, wherein on transduction into Huh7 cells, the viral particle expresses Factor IX protein or a fragment thereof having a Factor IX activity greater than the activity of Factor IX expressed from a viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO: 12 and a transcription regulatory element of SEQ ID NO. 7 and/or a viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO. 18 and a transcription regulatory element of SEQ ID NO. 6.
109. The viral particle of aspect 108, wherein the activity is measured using a chromogenic substrate which is specific for Factor Xa.
110. The polynucleotide or viral particle of any one of the preceding aspects, wherein the Factor IX protein fragment is at least 200, at least 250, at least 300, between 200 and 415, between 250 and 415, or between 300 and 415 amino acids in length.
111. The polynucleotide or viral particle of any one of the preceding aspects, wherein the Factor IX protein or fragment thereof comprises a sequence:
- a) at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to SEQ ID NO. 8; or
 - b) at least 95%, at least 98%, at least 99%, at least 99.5%, at least 99.8%, or 100% identical to a fragment of SEQ ID NO. 8 at least 200, at least 250, at least 300, between 200 and 415, between 250 and 415, or between 300 and 415 amino acids in length.
112. A composition comprising the polynucleotide or viral particle of any one of the preceding aspects and a pharmaceutically acceptable excipient.
113. The polynucleotide, viral particle or composition of any one of the preceding aspects for use in a method of treatment.
114. The polynucleotide, viral particle or composition for use of aspect 113, wherein the method of treatment comprises administering an effective amount of the polynucleotide or viral particle of any one of aspects 1-111 to a patient.
115. A method of treatment comprising administering an effective amount of the polynucleotide or viral particle of any one of aspects 1-111 to a patient.
116. Use of the polynucleotide, viral particle or composition of any one of aspects 1-111 in the manufacture of a medicament for use in a method of treatment.
117. The use of aspect 116, wherein the method of treatment comprises administering an effective amount of the polynucleotide or viral particle of any one of aspects 1-111 to a patient.
118. The polynucleotide, viral particle, composition, use or method of any one of aspects 112-117, wherein the method of treatment is a method of treating haemophilia.
119. The polynucleotide, viral particle, composition, use or method of aspect 118, wherein the haemophilia is haemophilia B.
120. The polynucleotide, viral particle, composition, use or method of aspect 119, wherein the patient has antibodies or inhibitors to Factor IX.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 26

<210> SEQ ID NO 1
 <211> LENGTH: 1191
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 1

```

gaggagaagt gcagccttga ggaggccagg gaggtgtttg agaacactga gaggaccact    60
gagttctgga agcagtatgt ggatggggac cagtgtgaga gcaaccctg cctgaatggg    120
ggcagctgca aggatgacat caacagctat gagtgtcggg gcccttttg ctttgagggc    180
aagaactgtg agctggatgt gacctgcaac atcaagaatg gcagatgtga gcagttctgc    240
aagaactctg ctgacaacaa ggtgggtgtg agctgcactg agggctacag gctggctgag    300
aaccagaaga gctgtgagcc tgctgtgcca ttcccatgtg gcagagtgtc tgtgagccag    360
accagcaagc tgaccagggc tgaggctgtg ttccctgatg tggactatgt gaacagcact    420
gaggctgaaa ccatcctgga caacatcacc cagagcacc agagcttcaa tgacttcacc    480
agggtggtgg ggggggagga tgccaagcct ggccagttcc cctggcaagt ggtgctgaat    540
ggcaagggtg atgccttctg tgggggcagc attgtgaatg agaagtggat tgtgactgct    600
gccactgtg tggagactgg ggtgaagatc actgtggtgg ctggggagca caacattgag    660
gagactgagc aactgagca gaagaggaat gtgatcagga tcatcccca ccacaactac    720
aatgctgcca tcaacaagta caaccatgac attgccttgc tggagctgga tgagccctg    780
gtgctgaaca gctatgtgac ccccatctgc attgctgaca aggagtacac caacatcttc    840
ctgaagtttg gctctggcta tgtgtctggc tggggcaggg tgttccaca gggcaggtct    900
gccctggtgc tgcagtacct gaggggtgcc ctggtggaca gggccacctg cctgctcagc    960
accaagttca ccatctaca caacatgttc tgtgctggct tccatgaggg gggcagggac   1020
agctgccagg gggactctgg gggcccccat gtgactgagg tggagggcac cagcttctg    1080
actggcatca tcagctgggg ggaggagtgt gccatgaagg gcaagtatgg catctacacc   1140
aaagtctcca gatatgtgaa ctggatcaag gagaagacca agctgacctg a           1191

```

<210> SEQ ID NO 2
 <211> LENGTH: 494
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 2

```

atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcaccat ctgcctttta    60
ggatatctac tcagtgtgta atgtacaggt ttgtttcctt ttttaaata cattgagtat    120
gcttgccctt tagatataga aatatctgat gctgtcttct tactaaatt ttgattacat    180
gatttgacag caatattgaa gagtctaaca gccagcacgc aggttggtaa gtactgtggg    240
aacatcacag attttggtc catgccctaa agagaaattg gctttcagat tatttgatt    300
aaaaacaaag actttcttaa gagatgtaaa attttcatga tgttttctt tttgctaaaa    360
ctaaagaatt attcttttac atttcagttt ttcttgatca tgaaaacgcc aacaaaattc    420
tgaatcggcc aaagaggtat aattcaggta aattggaaga gtttgttcaa gggaaccttg    480
agagagaatg tatg                                     494

```

-continued

<210> SEQ ID NO 3
 <211> LENGTH: 299
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

 <400> SEQUENCE: 3

 gtttgtttcc ttttttaaaa tacattgagt atgcttgcc tttagatata gaaatatctg 60
 atgctgtctt cttcactaaa ttttgattac atgatttgac agcaatattg aagagtctaa 120
 cagccagcac gcaggttgg aagtactgtg ggaacatcac agattttggc tccatgccct 180
 aaagagaaat tggctttcag attatttggga ttaaaaaaaa agactttctt aagagatgta 240
 aaattttcat gatgttttct tttttgctaa aactaaagaa ttattctttt acatttcag 299

<210> SEQ ID NO 4
 <211> LENGTH: 1386
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

 <400> SEQUENCE: 4

 atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcacat ctgcctttta 60
 ggatatctac tcagtgtgta atgtacagtt tttcttgatc atgaaaacgc caacaaaatt 120
 ctgaatcggc caaagaggta taattcaggt aaattggaag agtttgttca agggaacctt 180
 gagagagaat gtatggagga gaagtgcagc tttgaggagg ccaggagggt gtttgagaac 240
 actgagagga cactgagtt ctggaagcag tatgtggatg gggaccagtg tgagagcaac 300
 cctgcctga atgggggcag ctgcaaggat gacatcaaca gctatgagt ctggtgcccc 360
 tttggctttg agggcaagaa ctgtgagctg gatgtgacct gcaacatcaa gaatggcaga 420
 tgtgagcagt tctgcaagaa ctctgctgac aacaaggtgg tgtgcagctg cactgagggc 480
 tacaggctgg ctgagaacca gaagagctgt ggcctgctg tgccattccc atgtggcaga 540
 gtgtctgtga gccagaccag caagctgacc agggctgagg ctgtgttccc tgatgtggac 600
 tatgtgaaca gactgaggc tgaaccatc ctggacaaca tcaccagag caccagagc 660
 ttcaatgact tcaccagggt ggtggggggg gaggatgcca agcctggcca gttcccctgg 720
 caagtgggtg tgaatggcaa ggtggatgcc ttctgtgggg gcagcattgt gaatgagaag 780
 tggattgtga ctgctgcccc ctgtgtggag actgggggtga agatcactgt ggtggctggg 840
 gagcacaaca ttgaggagac tgagcacact gagcagaaga ggaatgtgat caggatcatc 900
 ccccaccaca actacaatgc tgccatcaac aagtacaacc atgacattgc cctgctggag 960
 ctggatgagc ccctggtgct gaacagctat gtgaccccc tctgcattgc tgacaaggag 1020
 tacaccaaca tcttctgaa gtttggctct ggctatgtgt ctggctgggg caggggttct 1080
 cacaagggca ggtctgccct ggtgctgcag tacctgaggg tgccccctgt ggacagggcc 1140
 acctgcctgc tcagaccaa gttcaccatc tacaacaaca tgttctgtgc tggcttccat 1200
 gaggggggca gggacagctg ccagggggac tctgggggcc cccatgtgac tgaggtggag 1260
 ggcaccagct tctgactgg catcatcagc tggggggagg agtgtgccc gaagggcaag 1320
 tatggcatct acaccaaagt ctccagatat gtgaactgga tcaaggagaa gaccaagctg 1380
 acctga 1386

-continued

```

<210> SEQ ID NO 5
<211> LENGTH: 1685
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 5
atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcacat ctgcctttta    60
ggatatctac tcagtgetga atgtacaggt ttgtttcctt ttttaaaata cattgagtat    120
gcttgccctt tagatataga aatatctgat gctgtcttct tcaactaaatt ttgattacat    180
gatttgacag caatattgaa gagtctaaca gccagcacgc aggttggtaa gtactgtggg    240
aacatcacag attttggtc catgccctaa agagaaattg gctttcagat tatttgatt    300
aaaaacaaag actttcttaa gagatgtaaa attttcatga tgttttctt tttgctaaaa    360
ctaaagaatt attcttttac atttcagttt ttcttgatca tgaaaacgcc aacaaaattc    420
tgaatcggcc aaagaggtat aattcaggta aattggaaga gtttgttcaa gggaaccttg    480
agagagaatg tatggaggag aagtgcagct ttgaggaggc caggagggtg tttgagaaca    540
ctgagaggac cactgagttc tgggaagcagt atgtggatgg ggaccagtgt gagagcaacc    600
cctgcctgaa tggggggcagc tgcaaggatg acatcaacag ctatgagtgc tggtgcccct    660
ttggctttga gggcaagaac tgtgagctgg atgtgacctg caacatcaag aatggcagat    720
gtgagcagtt ctgcaagaac tctgctgaca acaaggtggt gtgcagctgc actgagggct    780
acaggctggc tgagaaccag aagagctgtg agcctgctgt gccattcca tgtggcagag    840
tgtctgtgag ccagaccagc aagctgacca gggctgaggc tgtgttcctt gatgtggact    900
atgtgaacag cactgaggct gaaaccatcc tggacaacat caccagagc acccagagct    960
tcaatgactt caccaggtg gtgggggggg aggatgcaa gcctggccag tcccctggc    1020
aagtgggtgct gaatggcaag gtggatgcct tctgtggggg cagcattgtg aatgagaagt    1080
ggattgtgac tgctgccac tgtgtggaga ctggggtgaa gatcactgtg gtggctgggg    1140
agcacaacat tgaggagact gagcacactg agcagaagag gaatgtgatc aggatcatcc    1200
cccaccacaa ctacaatgct gccatcaaca agtacaacca tgacattgcc ctgctggagc    1260
tggatgagcc cctggtgctg aacagctatg tgacccccat ctgcattgct gacaaggagt    1320
acaccaacat cttcctgaag tttggctctg gctatgtgtc tggctggggc aggggtgttc    1380
acaagggcag gtctgcctg gtgctgcagt acctgagggt gcccctggtg gacagggcca    1440
cctgcctgct cagcaccaag ttcaccatct acaacaacat gttctgtgct ggcttccatg    1500
aggggggcag ggacagctgc cagggggact ctgggggccc ccatgtgact gaggtggagg    1560
gcaccagctt cctgactggc atcatcagct ggggggagga gtgtgccatg aagggaagt    1620
atggcatcta caccaaagtc tccagatatg tgaactggat caaggagaag accaagctga    1680
cctga    1685

```

```

<210> SEQ ID NO 6
<211> LENGTH: 334
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic regulatory element

<400> SEQUENCE: 6
ccctaaaatg ggcaaacatt gcaagcagca aacagcaaac acacagccct cctgcctgc    60

```

-continued

```

tgacctgga gctggggcag aggtcagaca cctctctggg cccatgccac ctccaactgg 120
acacaggacg ctgtggtttc tgagccaggg ggcgactcag atcccagcca gtggacttag 180
ccctgtttg ctctccgat aactggggtg accttggtta atattacca gcagcctccc 240
ccgttgcccc tctggatcca ctgcttaaat acggacgagg acagggcct gtctcctcag 300
cttcaggcac caccactgac ctgggacagt gaat 334

```

```

<210> SEQ ID NO 7
<211> LENGTH: 447
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic regulatory element

```

```

<400> SEQUENCE: 7

```

```

ccctaaaatg ggcaaacatt gcaagcagca aacagcaaac acacagccct cctgcttgc 60
tgacctgga gctggggcag aggtcagaga cctctctggg cccatgccac ctccaacatc 120
cactcgaccc cttggaattt cgggtgagag gagcagaggt tgtcctggcg tggtttaggt 180
agtgtgagag gggaatgact cctttcggta agtgcagtgg aagctgtaca ctgcccaggc 240
aaagcgtccg ggcagcgtag gcgggcgact cagatcccag ccagtggact tagcccctgt 300
ttgctcctcc gataactggg gtgaccttgg ttaatattca ccagcagcct cccccgttgc 360
ccctctggat ccactgctta aatacggacg aggacagggc cctgtctcct cagcttcagg 420
caccaccact gacctgggac agtgaat 447

```

```

<210> SEQ ID NO 8
<211> LENGTH: 415
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 8

```

```

Tyr Asn Ser Gly Lys Leu Glu Glu Phe Val Gln Gly Asn Leu Glu Arg
1          5          10          15
Glu Cys Met Glu Glu Lys Cys Ser Phe Glu Glu Ala Arg Glu Val Phe
20          25          30
Glu Asn Thr Glu Arg Thr Thr Glu Phe Trp Lys Gln Tyr Val Asp Gly
35          40          45
Asp Gln Cys Glu Ser Asn Pro Cys Leu Asn Gly Gly Ser Cys Lys Asp
50          55          60
Asp Ile Asn Ser Tyr Glu Cys Trp Cys Pro Phe Gly Phe Glu Gly Lys
65          70          75          80
Asn Cys Glu Leu Asp Val Thr Cys Asn Ile Lys Asn Gly Arg Cys Glu
85          90          95
Gln Phe Cys Lys Asn Ser Ala Asp Asn Lys Val Val Cys Ser Cys Thr
100         105         110
Glu Gly Tyr Arg Leu Ala Glu Asn Gln Lys Ser Cys Glu Pro Ala Val
115         120         125
Pro Phe Pro Cys Gly Arg Val Ser Val Ser Gln Thr Ser Lys Leu Thr
130         135         140
Arg Ala Glu Ala Val Phe Pro Asp Val Asp Tyr Val Asn Ser Thr Glu
145         150         155         160
Ala Glu Thr Ile Leu Asp Asn Ile Thr Gln Ser Thr Gln Ser Phe Asn
165         170         175
Asp Phe Thr Arg Val Val Gly Gly Glu Asp Ala Lys Pro Gly Gln Phe

```

-continued

180				185				190							
Pro	Trp	Gln	Val	Val	Leu	Asn	Gly	Lys	Val	Asp	Ala	Phe	Cys	Gly	Gly
	195						200					205			
Ser	Ile	Val	Asn	Glu	Lys	Trp	Ile	Val	Thr	Ala	Ala	His	Cys	Val	Glu
	210					215				220					
Thr	Gly	Val	Lys	Ile	Thr	Val	Val	Ala	Gly	Glu	His	Asn	Ile	Glu	Glu
	225				230					235					240
Thr	Glu	His	Thr	Glu	Gln	Lys	Arg	Asn	Val	Ile	Arg	Ile	Ile	Pro	His
			245					250						255	
His	Asn	Tyr	Asn	Ala	Ala	Ile	Asn	Lys	Tyr	Asn	His	Asp	Ile	Ala	Leu
		260						265					270		
Leu	Glu	Leu	Asp	Glu	Pro	Leu	Val	Leu	Asn	Ser	Tyr	Val	Thr	Pro	Ile
	275						280					285			
Cys	Ile	Ala	Asp	Lys	Glu	Tyr	Thr	Asn	Ile	Phe	Leu	Lys	Phe	Gly	Ser
	290					295					300				
Gly	Tyr	Val	Ser	Gly	Trp	Gly	Arg	Val	Phe	His	Lys	Gly	Arg	Ser	Ala
	305				310					315					320
Leu	Val	Leu	Gln	Tyr	Leu	Arg	Val	Pro	Leu	Val	Asp	Arg	Ala	Thr	Cys
			325						330					335	
Leu	Leu	Ser	Thr	Lys	Phe	Thr	Ile	Tyr	Asn	Asn	Met	Phe	Cys	Ala	Gly
			340					345					350		
Phe	His	Glu	Gly	Gly	Arg	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro
		355					360					365			
His	Val	Thr	Glu	Val	Glu	Gly	Thr	Ser	Phe	Leu	Thr	Gly	Ile	Ile	Ser
	370					375					380				
Trp	Gly	Glu	Glu	Cys	Ala	Met	Lys	Gly	Lys	Tyr	Gly	Ile	Tyr	Thr	Lys
	385				390					395					400
Val	Ser	Arg	Tyr	Val	Asn	Trp	Ile	Lys	Glu	Lys	Thr	Lys	Leu	Thr	
				405					410					415	

<210> SEQ ID NO 9

<211> LENGTH: 1386

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```

atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcacat ctgcctttta    60
ggatatctac tcagtgctga atgtacagtt tttcttgatc atgaaaacgc caacaaaatt    120
ctgaatcggc caaagaggta taattcaggt aaattggaag agtttgttca agggaacctt    180
gagagagaat gtatggaaga aaagtgtagt tttgaagaag cacgagaagt ttttgaaaac    240
actgaaagaa caactgaatt ttggaagcag tatgttgatg gagatcagtg tgagtccaat    300
ccatgtttaa atggcggcag ttgcaaggat gacattaatt cctatgaatg ttggtgtccc    360
tttgatttg aaggaaagaa ctgtgaatta gatgtaacat gtaacattaa gaatggcaga    420
tgcgagcagt tttgtaaaaa tagtgctgat aacaagggtg tttgctctg tactgagggg    480
tatcgacttg cagaaaacca gaagtcctgt gaaccagcag tgccatttcc atgtggaaga    540
gtttctgttt cacaaacttc taagctcacc cgtgctgagg ctgtttttcc tgatgtggac    600
tatgtaaatt ctactgaagc tgaaccatt ttggataaca tcaactcaag cacccaatca    660
tttaatgact tcactcgggt tgttggtgga gaagatgcca aaccaggtea attcccttgg    720
caggttgttt tgaatggtaa agttgatgca ttctgtggag gctctatcgt taatgaaaaa    780
tggattgtaa ctgctgccca ctgtgttgaa actggtgta aaattacagt tgtcgcaggt    840

```

-continued

```

gaacataata ttgaggagac agaacataca gagcaaaagc gaaatgtgat tcgaattatt    900
cctcaccaca actacaatgc agctattaat aagtacaacc atgacattgc ccttctggaa    960
ctggacgaac ccttagtgct aacagctac gttacaccta tttgcattgc tgacaaggaa   1020
tacacgaaca tcttctcaa atttggatct ggctatgtaa gtggctgggg aagagtcttc   1080
caciaaggga gatcagcttt agttcttcag taccttagag ttccacttgt tgaccgagcc   1140
acatgtcttc gatctacaaa gttcaccatc tataacaaca tgttctgtgc tggcttccat   1200
gaaggaggta gagattcatg tcaaggagat agtggggggac cccatgttac tgaagtggaa   1260
gggaccagtt tcttaactgg aattattagc tggggggaag agtgtgcaat gaaaggcaaa   1320
tatggaatat ataccaaggt atcccgggat gtcaactgga ttaaggaaaa aacaaagctc   1380
acttaa                                           1386

```

```

<210> SEQ ID NO 10
<211> LENGTH: 736
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 10

```

```

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Asn Leu Ser
 1           5           10           15
Glu Gly Ile Arg Glu Trp Trp Ala Leu Lys Pro Gly Ala Pro Lys Pro
          20           25           30
Lys Ala Asn Gln Gln Lys Gln Asp Asp Gly Arg Gly Leu Val Leu Pro
          35           40           45
Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
          50           55           60
Val Asn Ala Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
 65           70           75           80
Gln Gln Leu Gln Ala Gly Asp Asn Pro Tyr Leu Arg Tyr Asn His Ala
          85           90           95
Asp Ala Glu Phe Gln Glu Arg Leu Gln Glu Asp Thr Ser Phe Gly Gly
          100          105          110
Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
          115          120          125
Leu Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Gly Lys Lys Arg
          130          135          140
Pro Val Asp Gln Ser Pro Gln Glu Pro Asp Ser Ser Ser Gly Val Gly
 145          150          155          160
Lys Ser Gly Lys Gln Pro Ala Arg Lys Arg Leu Asn Phe Gly Gln Thr
          165          170          175
Gly Asp Ser Glu Ser Val Pro Asp Pro Gln Pro Leu Gly Glu Pro Pro
          180          185          190
Ala Ala Pro Thr Ser Leu Gly Ser Asn Thr Met Ala Ser Gly Gly Gly
          195          200          205
Ala Pro Met Ala Asp Asn Asn Glu Gly Ala Asp Gly Val Gly Asn Ser
          210          215          220
Ser Gly Asn Trp His Cys Asp Ser Gln Trp Leu Gly Asp Arg Val Ile
 225          230          235          240
Thr Thr Ser Thr Arg Thr Trp Ala Leu Pro Thr Tyr Asn Asn His Leu
          245          250          255

```


-continued

Tyr Lys Gln Ile Ser Ser Gln Ser Gly Ala Ser Asn Asp Asn His Tyr
 260 265 270
 Phe Gly Tyr Ser Thr Pro Trp Gly Tyr Phe Asp Phe Asn Arg Phe His
 275 280 285
 Cys His Phe Ser Pro Arg Asp Trp Gln Arg Leu Ile Asn Asn Asn Trp
 290 295 300
 Gly Phe Arg Pro Lys Lys Leu Ser Phe Lys Leu Phe Asn Ile Gln Val
 305 310 315 320
 Lys Glu Val Thr Gln Asn Asp Gly Thr Thr Thr Ile Ala Asn Asn Leu
 325 330 335
 Thr Ser Thr Val Gln Val Phe Thr Asp Ser Glu Tyr Gln Leu Pro Tyr
 340 345 350
 Val Leu Gly Ser Ala His Gln Gly Cys Leu Pro Pro Phe Pro Ala Asp
 355 360 365
 Val Phe Met Val Pro Gln Tyr Gly Tyr Leu Thr Leu Asn Asn Gly Ser
 370 375 380
 Gln Ala Val Gly Arg Ser Ser Phe Tyr Cys Leu Glu Tyr Phe Pro Ser
 385 390 395 400
 Gln Met Leu Arg Thr Gly Asn Asn Phe Gln Phe Ser Tyr Thr Phe Glu
 405 410 415
 Asp Val Pro Phe His Ser Ser Tyr Ala His Ser Gln Ser Leu Asp Arg
 420 425 430
 Leu Met Asn Pro Leu Ile Asp Gln Tyr Leu Tyr Tyr Leu Asn Arg Thr
 435 440 445
 Gln Gly Thr Thr Ser Gly Thr Thr Asn Gln Ser Arg Leu Leu Phe Ser
 450 455 460
 Gln Ala Gly Pro Gln Ser Met Ser Leu Gln Ala Arg Asn Trp Leu Pro
 465 470 475 480
 Gly Pro Cys Tyr Arg Gln Gln Arg Leu Ser Lys Thr Ala Asn Asp Asn
 485 490 495
 Asn Asn Ser Asn Phe Pro Trp Thr Ala Ala Ser Lys Tyr His Leu Asn
 500 505 510
 Gly Arg Asp Ser Leu Val Asn Pro Gly Pro Ala Met Ala Ser His Lys
 515 520 525
 Asp Asp Glu Glu Lys Phe Phe Pro Met His Gly Asn Leu Ile Phe Gly
 530 535 540
 Lys Glu Gly Thr Thr Ala Ser Asn Ala Glu Leu Asp Asn Val Met Ile
 545 550 555 560
 Thr Asp Glu Glu Glu Ile Arg Thr Thr Asn Pro Val Ala Thr Glu Gln
 565 570 575
 Tyr Gly Thr Val Ala Asn Asn Leu Gln Ser Ser Asn Thr Ala Pro Thr
 580 585 590
 Thr Arg Thr Val Asn Asp Gln Gly Ala Leu Pro Gly Met Val Trp Gln
 595 600 605
 Asp Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp Ala Lys Ile Pro His
 610 615 620
 Thr Asp Gly His Phe His Pro Ser Pro Leu Met Gly Gly Phe Gly Leu
 625 630 635 640
 Lys His Pro Pro Pro Gln Ile Met Ile Lys Asn Thr Pro Val Pro Ala
 645 650 655
 Asn Pro Pro Thr Thr Phe Ser Pro Ala Lys Phe Ala Ser Phe Ile Thr
 660 665 670
 Gln Tyr Ser Thr Gly Gln Val Ser Val Glu Ile Glu Trp Glu Leu Gln

-continued

675	680	685
Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln Tyr Thr Ser Asn		
690	695	700
Tyr Asn Lys Ser Val Asn Val Asp Phe Thr Val Asp Thr Asn Gly Val		
705	710	715
Tyr Ser Glu Pro Arg Pro Ile Gly Thr Arg Tyr Leu Thr Arg Asn Leu		
	725	730
		735

<210> SEQ ID NO 11
 <211> LENGTH: 1386
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 11

```

atgcagaggg tgaacatgat catggctgag agccctggcc tgatcacat ctgctgctg      60
ggctacctgc tgtctgctga gtgcactgtg ttctggacc atgagaatgc caacaagatc     120
ctgaacaggc ccaagagata caactctggc aagctggagg agtttgtgca gggcaacctg     180
gagagggagt gcatggagga gaagtgcagc tttgaggagg ccagggaggt gtttgagaac     240
actgagagga cactgagtt ctggaagcag tatgtggatg gggaccagtg tgagagcaac     300
ccctgctga atgggggag ctgcaaggat gacatcaaca gctatgagtg ctggtgcccc     360
tttgctttg agggcaagaa ctgtgagctg gatgtgacct gcaacatcaa gaatggcaga     420
tgtgagcagt tctgcaagaa ctctgctgac aacaaggtgg tgtgcagctg cactgagggc     480
tacaggctgg ctgagaacca gaagagctgt gagcctgctg tgccattccc atgtggcaga     540
gtgtctgtga gccagaccag caagctgacc agggctgagg ctgtgttccc tgatgtggac     600
tatgtgaaca gcactgaggc tgaaacatc ctggacaaca tcaccagag caccagagc     660
ttcaatgact tcaccagggt ggtggggggg gaggatgcca agcctggcca gttcccctgg     720
caagtgggag tgaatggcaa ggtggatgcc ttctgtgggg gcagcattgt gaatgagaag     780
tggattgtga ctgctgcca ctgtgtggag actgggggta agatcactgt ggtggctggg     840
gagcacaaca ttgaggagac tgagcacact gagcagaaga ggaatgtgat caggatcatc     900
ccccaccaca actacaatgc tgccatcaac aagtacaacc atgacattgc cctgctggag     960
ctggatgagc ccctggtgct gaacagctat gtgaccccc tctgcattgc tgacaaggag    1020
tacaccaaca tcttctgaa gtttggtctt ggctatgtgt ctggctgggg caggggtgtc    1080
cacaagggca ggtctgccct ggtgctgcag tacctgaggg tgccccctgt ggacagggcc    1140
acctgcctgt tgagcaccaa gttcaccatc tacaacaaca tgttctgtgc tggttccat    1200
gaggggggca gggacagctg ccagggggac tctgggggccc cccatgtgac tgaggtggag    1260
ggcaccagct tcctgactgg catcatcagc tggggggagg agtgtgcat gaagggcaag    1320
tatggcatct acaccaaagt ctccagatat gtgaactgga tcaaggagaa gaccaagctg    1380
acctga                                           1386
  
```

<210> SEQ ID NO 12
 <211> LENGTH: 1386
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 12

-continued

```

atgcagaggg tgaacatgat catggctgag agccctggcc tgatcacat ctgectgctg 60
ggctacctgc tgtctgctga gtgactgtg ttctggacc atgagaatgc caacaagatc 120
ctgaacaggg ccaagagata caactctggc aagctggagg agtttgtgca gggcaacctg 180
gagagggagt gcatggagga gaagtgcagc tttgaggagg ccaggagggt gtttgagaac 240
actgagagga cactgagtt ctggaagcag tatgtggatg gggaccagtg tgagagcaac 300
ccctgcctga atgggggag ctgcaaggat gacatcaaca gctatgagtg ctggtgcccc 360
tttgctttg agggcaagaa ctgtgagctg gatgtgacct gcaacatcaa gaatggcaga 420
tgtgagcagt tctgcaagaa ctctgctgac aacaaggagg tgtgcagctg cactgagggc 480
tacaggctgg ctgagaacca gaagagctgt gagcctgctg tgccattccc atgtggcaga 540
gtgtctgtga gccagaccag caagctgacc agggctgagg ctgtgttccc tgatgtggac 600
tatgtgaaca gactgagggc tgaaacatc ctggacaaca tccccagag cccccagagc 660
ttcaatgact tcaccagggt ggtggggggg gaggatgcca agcctggcca gttcccctgg 720
caagtggggt tgaatggcaa ggtggatgcc ttctgtgggg gcagcattgt gaatgagaag 780
tggattgtga ctgctgcccc ctgtgtggag actgggggtga agatcactgt ggtggctggg 840
gagcacaaca ttgaggagac tgagcacact gagcagaaga ggaatgtgat caggatcatc 900
ccccaccaca actacaatgc tgccatcaac aagtacaacc atgacattgc cctgctggag 960
ctggatgagc ccctggtgct gaacagctat gtgaccccc tctgcattgc tgacaaggag 1020
tacaccaaca tcttctgaa gtttggtctt ggctatgtgt ctggctgggg cagggtgttc 1080
cacaagggca ggtctgcctt ggtgctgcag tacctgaggg tgcccctggt ggacagggcc 1140
acctgcctga ggagaccaa gttcaccatc tacaacaaca tgttctgtgc tggcttccat 1200
gaggggggca gggacagctg ccagggggac tctgggggccc cccatgtgac tgaggtggag 1260
ggcaccagct tcctgactgg catcatcagc tggggggagg agtgtgccc gaagggcaag 1320
tatggcatct acaccaaagt ctccagatat gtgaactgga tcaaggagaa gaccaagctg 1380
acctga 1386

```

```

<210> SEQ ID NO 13
<211> LENGTH: 117
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic enhancer element

```

```

<400> SEQUENCE: 13

```

```

ccctaaaatg ggcaaacatt gcaagcagca aacagcaaac acacagccct cctgcctgc 60
tgaccttggg gctggggcag aggtcagaca cctctctggg cccatgccac ctccaac 117

```

```

<210> SEQ ID NO 14
<211> LENGTH: 185
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic promoter element

```

```

<400> SEQUENCE: 14

```

```

gggcgactca gatcccagcc agtggactta gccctggtt gctcctccga taactgggggt 60
gaccttgggt aatattcacc agcagcctcc cccgttgcce ctctggatcc actgcttaa 120
tacggacgag gacagggccc tgtctcctca gcttcaggca ccaccactga cctgggacag 180
tgaat 185

```

-continued

<210> SEQ ID NO 15
 <211> LENGTH: 195
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

 <400> SEQUENCE: 15

 atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcacat ctgcctttta 60
 ggatatctac tcagtgtga atgtacagtt tttcttgatc atgaaaacgc caacaaaatt 120
 ctgaatcggc caaagaggta taattcaggt aaattggaag agtttgttca agggaacctt 180
 gagagagaat gtagt 195

<210> SEQ ID NO 16
 <211> LENGTH: 461
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

 <400> SEQUENCE: 16

 Met Gln Arg Val Asn Met Ile Met Ala Glu Ser Pro Gly Leu Ile Thr
 1 5 10 15
 Ile Cys Leu Leu Gly Tyr Leu Leu Ser Ala Glu Cys Thr Val Phe Leu
 20 25 30
 Asp His Glu Asn Ala Asn Lys Ile Leu Asn Arg Pro Lys Arg Tyr Asn
 35 40 45
 Ser Gly Lys Leu Glu Glu Phe Val Gln Gly Asn Leu Glu Arg Glu Cys
 50 55 60
 Met Glu Glu Lys Cys Ser Phe Glu Glu Ala Arg Glu Val Phe Glu Asn
 65 70 75 80
 Thr Glu Arg Thr Thr Glu Phe Trp Lys Gln Tyr Val Asp Gly Asp Gln
 85 90 95
 Cys Glu Ser Asn Pro Cys Leu Asn Gly Gly Ser Cys Lys Asp Asp Ile
 100 105 110
 Asn Ser Tyr Glu Cys Trp Cys Pro Phe Gly Phe Glu Gly Lys Asn Cys
 115 120 125
 Glu Leu Asp Val Thr Cys Asn Ile Lys Asn Gly Arg Cys Glu Gln Phe
 130 135 140
 Cys Lys Asn Ser Ala Asp Asn Lys Val Val Cys Ser Cys Thr Glu Gly
 145 150 155 160
 Tyr Arg Leu Ala Glu Asn Gln Lys Ser Cys Glu Pro Ala Val Pro Phe
 165 170 175
 Pro Cys Gly Arg Val Ser Val Ser Gln Thr Ser Lys Leu Thr Arg Ala
 180 185 190
 Glu Ala Val Phe Pro Asp Val Asp Tyr Val Asn Ser Thr Glu Ala Glu
 195 200 205
 Thr Ile Leu Asp Asn Ile Thr Gln Ser Thr Gln Ser Phe Asn Asp Phe
 210 215 220
 Thr Arg Val Val Gly Gly Glu Asp Ala Lys Pro Gly Gln Phe Pro Trp
 225 230 235 240
 Gln Val Val Leu Asn Gly Lys Val Asp Ala Phe Cys Gly Gly Ser Ile
 245 250 255
 Val Asn Glu Lys Trp Ile Val Thr Ala Ala His Cys Val Glu Thr Gly
 260 265 270
 Val Lys Ile Thr Val Val Ala Gly Glu His Asn Ile Glu Glu Thr Glu

-continued

275				280				285							
His	Thr	Glu	Gln	Lys	Arg	Asn	Val	Ile	Arg	Ile	Ile	Pro	His	His	Asn
	290					295					300				
Tyr	Asn	Ala	Ala	Ile	Asn	Lys	Tyr	Asn	His	Asp	Ile	Ala	Leu	Leu	Glu
305					310					315					320
Leu	Asp	Glu	Pro	Leu	Val	Leu	Asn	Ser	Tyr	Val	Thr	Pro	Ile	Cys	Ile
				325					330					335	
Ala	Asp	Lys	Glu	Tyr	Thr	Asn	Ile	Phe	Leu	Lys	Phe	Gly	Ser	Gly	Tyr
			340					345					350		
Val	Ser	Gly	Trp	Gly	Arg	Val	Phe	His	Lys	Gly	Arg	Ser	Ala	Leu	Val
		355					360					365			
Leu	Gln	Tyr	Leu	Arg	Val	Pro	Leu	Val	Asp	Arg	Ala	Thr	Cys	Leu	Leu
	370					375					380				
Ser	Thr	Lys	Phe	Thr	Ile	Tyr	Asn	Asn	Met	Phe	Cys	Ala	Gly	Phe	His
385					390					395					400
Glu	Gly	Gly	Arg	Asp	Ser	Cys	Gln	Gly	Asp	Ser	Gly	Gly	Pro	His	Val
				405					410					415	
Thr	Glu	Val	Glu	Gly	Thr	Ser	Phe	Leu	Thr	Gly	Ile	Ile	Ser	Trp	Gly
			420					425					430		
Glu	Glu	Cys	Ala	Met	Lys	Gly	Lys	Tyr	Gly	Ile	Tyr	Thr	Lys	Val	Ser
		435					440					445			
Arg	Tyr	Val	Asn	Trp	Ile	Lys	Glu	Lys	Thr	Lys	Leu	Thr			
	450					455					460				

<210> SEQ ID NO 17

<211> LENGTH: 724

<212> TYPE: PRT

<213> ORGANISM: adeno-associated virus 5

<400> SEQUENCE: 17

Met	Ser	Phe	Val	Asp	His	Pro	Pro	Asp	Trp	Leu	Glu	Glu	Val	Gly	Glu
1				5					10					15	
Gly	Leu	Arg	Glu	Phe	Leu	Gly	Leu	Glu	Ala	Gly	Pro	Pro	Lys	Pro	Lys
			20					25					30		
Pro	Asn	Gln	Gln	His	Gln	Asp	Gln	Ala	Arg	Gly	Leu	Val	Leu	Pro	Gly
		35					40					45			
Tyr	Asn	Tyr	Leu	Gly	Pro	Gly	Asn	Gly	Leu	Asp	Arg	Gly	Glu	Pro	Val
	50					55					60				
Asn	Arg	Ala	Asp	Glu	Val	Ala	Arg	Glu	His	Asp	Ile	Ser	Tyr	Asn	Glu
65					70					75					80
Gln	Leu	Glu	Ala	Gly	Asp	Asn	Pro	Tyr	Leu	Lys	Tyr	Asn	His	Ala	Asp
				85					90					95	
Ala	Glu	Phe	Gln	Glu	Lys	Leu	Ala	Asp	Asp	Thr	Ser	Phe	Gly	Gly	Asn
			100					105					110		
Leu	Gly	Lys	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Val	Leu	Glu	Pro	Phe
		115					120					125			
Gly	Leu	Val	Glu	Glu	Gly	Ala	Lys	Thr	Ala	Pro	Thr	Gly	Lys	Arg	Ile
	130					135					140				
Asp	Asp	His	Phe	Pro	Lys	Arg	Lys	Lys	Ala	Arg	Thr	Glu	Glu	Asp	Ser
145					150					155					160
Lys	Pro	Ser	Thr	Ser	Ser	Asp	Ala	Glu	Ala	Gly	Pro	Ser	Gly	Ser	Gln
				165					170					175	
Gln	Leu	Gln	Ile	Pro	Ala	Gln	Pro	Ala	Ser	Ser	Leu	Gly	Ala	Asp	Thr
			180					185						190	

-continued

Met	Ser	Ala	Gly	Gly	Gly	Gly	Pro	Leu	Gly	Asp	Asn	Asn	Gln	Gly	Ala
		195					200					205			
Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asp	Trp	His	Cys	Asp	Ser	Thr	Trp
	210					215					220				
Met	Gly	Asp	Arg	Val	Val	Thr	Lys	Ser	Thr	Arg	Thr	Trp	Val	Leu	Pro
225					230					235					240
Ser	Tyr	Asn	Asn	His	Gln	Tyr	Arg	Glu	Ile	Lys	Ser	Gly	Ser	Val	Asp
				245					250					255	
Gly	Ser	Asn	Ala	Asn	Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	Tyr
			260					265					270		
Phe	Asp	Phe	Asn	Arg	Phe	His	Ser	His	Trp	Ser	Pro	Arg	Asp	Trp	Gln
	275						280					285			
Arg	Leu	Ile	Asn	Asn	Tyr	Trp	Gly	Phe	Arg	Pro	Arg	Ser	Leu	Arg	Val
	290					295				300					
Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Val	Gln	Asp	Ser	Thr
305					310					315					320
Thr	Thr	Ile	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr	Asp
				325					330					335	
Asp	Asp	Tyr	Gln	Leu	Pro	Tyr	Val	Val	Gly	Asn	Gly	Thr	Glu	Gly	Cys
			340					345					350		
Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val	Phe	Thr	Leu	Pro	Gln	Tyr	Gly	Tyr
		355					360					365			
Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr	Glu	Asn	Pro	Thr	Glu	Arg	Ser	Ser
	370					375					380				
Phe	Phe	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Lys	Met	Leu	Arg	Thr	Gly	Asn
385					390					395					400
Asn	Phe	Glu	Phe	Thr	Tyr	Asn	Phe	Glu	Glu	Val	Pro	Phe	His	Ser	Ser
				405					410					415	
Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe	Lys	Leu	Ala	Asn	Pro	Leu	Val	Asp
		420						425					430		
Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser	Thr	Asn	Asn	Thr	Gly	Gly	Val	Gln
		435					440					445			
Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg	Tyr	Ala	Asn	Thr	Tyr	Lys	Asn	Trp
	450					455					460				
Phe	Pro	Gly	Pro	Met	Gly	Arg	Thr	Gln	Gly	Trp	Asn	Leu	Gly	Ser	Gly
465					470					475					480
Val	Asn	Arg	Ala	Ser	Val	Ser	Ala	Phe	Ala	Thr	Thr	Asn	Arg	Met	Glu
				485					490					495	
Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val	Pro	Pro	Gln	Pro	Asn	Gly	Met	Thr
			500					505					510		
Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr	Tyr	Ala	Leu	Glu	Asn	Thr	Met	Ile
		515					520					525			
Phe	Asn	Ser	Gln	Pro	Ala	Asn	Pro	Gly	Thr	Thr	Ala	Thr	Tyr	Leu	Glu
	530					535					540				
Gly	Asn	Met	Leu	Ile	Thr	Ser	Glu	Ser	Glu	Thr	Gln	Pro	Val	Asn	Arg
545					550					555					560
Val	Ala	Tyr	Asn	Val	Gly	Gly	Gln	Met	Ala	Thr	Asn	Asn	Gln	Ser	Ser
				565					570					575	
Thr	Thr	Ala	Pro	Ala	Thr	Gly	Thr	Tyr	Asn	Leu	Gln	Glu	Ile	Val	Pro
		580						585					590		
Gly	Ser	Val	Trp	Met	Glu	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile	Trp
		595					600					605			
Ala	Lys	Ile	Pro	Glu	Thr	Gly	Ala	His	Phe	His	Pro	Ser	Pro	Ala	Met

-continued

610	615	620
Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys Asn 625	630	635
Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val Ser 645	650	655
Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met Glu 660	665	670
Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln 675	680	685
Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro Asp 690	695	700
Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr Leu 705	710	715
720		
Thr Arg Pro Leu		

<210> SEQ ID NO 18
 <211> LENGTH: 1685
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 18

```

atgcagcgcg tgaacatgat catggcagaa tcaccaggcc tcatcaccat ctgcctttta    60
ggatatctac tcagtgetga atgtacaggt ttgtttcctt ttttaaata cattgagtat    120
gcttgccctt tagatataga aatatctgat gctgtcttct tcactaaatt ttgattacat    180
gatttgacag caatattgaa gagtctaaca gccagcacgc aggttggtaa gtactgtggg    240
aacatcacag attttggtc catgccttaa agagaaattg gctttcagat tatttgatt    300
aaaaacaaag actttcttaa gagatgtaaa attttcatga tgttttctt tttgctaaaa    360
ctaaagaatt attcttttac atttcagttt ttcttgatca tgaaaacgcc aaaaaattc    420
tgaatcggcc aaagaggtat aattcaggta aattggaaga gtttggtcaa gggaaccttg    480
agagagaatg tatggaggag aagtgttctt tcgaggaggc gagagaggtt ttcgagaata    540
ctgagcgaac aaccgaattc tggaaacaat atgtggatgg cgaccaatgt gaatctaate    600
cctgcctcaa cgggtggtca tgcaaagacg atatcaacag ctacgagtgt tggtgccct    660
ttggtttcga gggaaagaat tgcgagcttg atgtaacctg taacattaag aatgggctg    720
gcgaacagtt ttgcaagaac agcgccgaca ataaggctgt ctgcagttgt accgaaggct    780
ataggcttgc agagaatcag aagagttgag agcctgctgt gccgttcca tgtggcagag    840
tcagtgtgtc ccaaactagc aagctgacaa gagcagaagc cgttttccc gatgtggact    900
acgtgaattc cactgaagcc gaaacgatcc tggacaatat cacacagagc actcagtctt    960
tcaacgactt cacacgggtt gtgggaggag aggacgcaa acccgccag tttccttggc    1020
aagtcgttct taacggcaag gtcgagcct tttgtggagg gagtattgtg aacgagaaat    1080
ggattgtcac cgctgctcat tgtgttgaat ctggggtgaa aatcactgtt gtcgaggag    1140
agcacaatat cgaagagaca gaacacaccg agcagaaacg caacgttatt cggatcattc    1200
cacatcacia ctacaatgct gccatcaaca agtacaacca cgacattgag ctgctggagt    1260
tggatgaacc tctcgtgctc aactcctatg tgacccaat ctgcatagca gataaggagt    1320
ataccaacat cttcctgaag tttgggtcag gttatgtgct aggctgggga cgagtgttcc    1380
ataaaggag atcagcactg gtgttgagc atctgcgctg accactggtg gatcgggcta    1440

```

-continued

```

cttgectgct aagcacaaaa ttcaccatct acaacaacat gttttgtgcc ggttttcacg 1500
aaggcggcag ggacagctgt caggagatt ccggagggcc tcatgtcaca gaggtcgagg 1560
gcacctcctt tctcactggg attataagct ggggagaaga atgcgccatg aaaggaagt 1620
acggcatata cacgaaagtg tctagatacg tgaattggat taaggaaaag accaaactga 1680
cgtga 1685

```

```

<210> SEQ ID NO 19
<211> LENGTH: 1248
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 19

```

```

tataattcag gtaaattgga agagtttggt caaggaacc ttgagagaga atgtatggaa 60
gaaaagtgta gttttgaaga agcacgagaa gttttgaaa aactgaaag aacaactgaa 120
ttttggaagc agtatgttga tggagatcag tgtgagtcca atccatgttt aaatggcggc 180
agttgcaagg atgacattaa ttccatgaa tgttggtgtc ctttggatt tgaaggaaag 240
aactgtgaat tagatgtaac atgtaacatt aagaatggca gatgagca gttttgtaa 300
aatagtgtg ataacaagggt ggtttgctcc tgtactgagg gatatcgact tgcagaaaac 360
cagaagtct gtgaaccagc agtgccattt ccatgtgaa gagtttctgt ttcacaaact 420
tctaagctca cccgtgctga ggctgtttt cctgatgtgg actatgtaa ttctactgaa 480
gctgaaacca ttttgataa catcactcaa agcaccaat catttaatga cttcactcgg 540
gttgttggtg gagaagatgc caaaccagg caattccctt ggcagggtgt tttgaatggt 600
aaagttgatg cattctgtgg aggtctatc gttaatgaaa aatggattgt aactgctgcc 660
cactgtgttg aaactggtgt taaaattaca gttgtcgcag gtgaacataa tattgaggag 720
acagaacata cagagcaaaa gcgaaatgtg attcgaatta ttccacca caactacaat 780
gcagctatta ataagtacaa ccatgacatt gcccttctgg aactggacga acccttagtg 840
ctaaacagct acgttacacc tatttgcat gctgacaagg aatacacgaa catcttctc 900
aaatttgat ctggctatgt aagtggctgg ggaagagtct tccacaaagg gagatcagct 960
ttagttcttc agtaccttag agttccactt gttgaccgag ccacatgtct tcatctaca 1020
aagttcacca tctataacaa catgttctgt gctggcttcc atgaaggagg tagagattca 1080
tgcaaggag atagtggggg accccatggt actgaagtgg aaggaccag tttcttaact 1140
ggaattatta gctgggtga agagtgtgca atgaaaggca aatatggaat atataccaag 1200
gtatccoggt atgtcaactg gattaaggaa aaaacaaagc tcacttaa 1248

```

```

<210> SEQ ID NO 20
<211> LENGTH: 724
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic polypeptide

```

```

<400> SEQUENCE: 20

```

```

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Glu Val Gly Glu
1           5           10           15
Gly Leu Arg Glu Phe Leu Gly Leu Glu Ala Gly Pro Pro Lys Pro Lys
20          25          30
Pro Asn Gln Gln His Gln Asp Gln Ala Arg Gly Leu Val Leu Pro Gly
35          40          45

```


-continued

Tyr Asn Tyr Leu Gly Pro Gly Asn Gly Leu Asp Arg Gly Glu Pro Val
 50 55 60

Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp Ile Ser Tyr Asn Glu
 65 70 75 80

Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala Asp
 85 90 95

Ala Glu Phe Gln Glu Lys Leu Ala Asp Asp Thr Ser Phe Gly Gly Asn
 100 105 110

Leu Gly Lys Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro Phe
 115 120 125

Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Thr Gly Lys Arg Ile
 130 135 140

Asp Asp His Phe Pro Lys Arg Lys Lys Ala Arg Thr Glu Glu Asp Ser
 145 150 155 160

Lys Pro Ser Thr Ser Ser Asp Ala Glu Ala Gly Pro Ser Gly Ser Gln
 165 170 175

Gln Leu Gln Ile Pro Ala Gln Pro Ala Ser Ser Leu Gly Ala Asp Thr
 180 185 190

Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly Ala
 195 200 205

Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr Trp
 210 215 220

Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu Pro
 225 230 235 240

Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val Asp
 245 250 255

Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly Tyr
 260 265 270

Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp Gln
 275 280 285

Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg Val
 290 295 300

Lys Ile Phe Asn Ile Gln Val Lys Glu Val Thr Val Gln Asp Ser Thr
 305 310 315 320

Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Thr Asp
 325 330 335

Asp Asp Tyr Gln Leu Pro Tyr Val Val Gly Asn Gly Thr Glu Gly Cys
 340 345 350

Leu Pro Ala Phe Pro Pro Gln Val Phe Thr Leu Pro Gln Tyr Gly Tyr
 355 360 365

Ala Thr Leu Asn Arg Asp Asn Thr Glu Asn Pro Thr Glu Arg Ser Ser
 370 375 380

Phe Phe Cys Leu Glu Tyr Phe Pro Ser Lys Met Leu Arg Thr Gly Asn
 385 390 395 400

Asn Phe Glu Phe Thr Tyr Asn Phe Glu Glu Val Pro Phe His Ser Ser
 405 410 415

Phe Ala Pro Ser Gln Asn Leu Phe Lys Leu Ala Asn Pro Leu Val Asp
 420 425 430

Gln Tyr Leu Tyr Arg Phe Val Ser Thr Asn Asn Thr Gly Gly Val Gln
 435 440 445

Phe Asn Lys Asn Leu Ala Gly Arg Tyr Ala Asn Thr Tyr Lys Asn Trp
 450 455 460

-continued

Phe Pro Gly Pro Met Gly Arg Thr Gln Gly Trp Asn Leu Gly Ser Gly
 465 470 475 480
 Val Asn Arg Ala Ser Val Ser Ala Phe Ala Thr Thr Asn Arg Met Glu
 485 490 495
 Leu Glu Gly Ala Ser Tyr Gln Val Pro Pro Gln Pro Asn Gly Met Thr
 500 505 510
 Asn Asn Leu Gln Gly Ser Asn Thr Tyr Ala Leu Glu Asn Thr Met Ile
 515 520 525
 Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu Glu
 530 535 540
 Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn Arg
 545 550 555 560
 Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser Ser
 565 570 575
 Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val Pro
 580 585 590
 Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile Trp
 595 600 605
 Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala Met
 610 615 620
 Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys Asn
 625 630 635 640
 Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val Ser
 645 650 655
 Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met Glu
 660 665 670
 Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile Gln
 675 680 685
 Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro Asp
 690 695 700
 Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr Leu
 705 710 715 720
 Thr Arg Pro Leu

<210> SEQ ID NO 21
 <211> LENGTH: 725
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 21

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
 20 25 30
 Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Asn Tyr Leu Gly Pro Gly Asn Gly Leu Asp Arg Gly Glu Pro
 50 55 60
 Val Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp Ile Ser Tyr Asn
 65 70 75 80
 Glu Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Lys Leu Ala Asp Asp Thr Ser Phe Gly Gly

-continued

100					105					110					
Asn	Leu	Gly	Lys	Ala	Val	Phe	Gln	Ala	Lys	Lys	Arg	Val	Leu	Glu	Pro
		115					120					125			
Phe	Gly	Leu	Val	Glu	Glu	Gly	Ala	Lys	Thr	Ala	Pro	Thr	Gly	Lys	Arg
	130					135					140				
Ile	Asp	Asp	His	Phe	Pro	Lys	Arg	Lys	Lys	Ala	Arg	Thr	Glu	Glu	Asp
145					150					155					160
Ser	Lys	Pro	Ser	Thr	Ser	Ser	Asp	Ala	Glu	Ala	Gly	Pro	Ser	Gly	Ser
				165					170					175	
Gln	Gln	Leu	Gln	Ile	Pro	Ala	Gln	Pro	Ala	Ser	Ser	Leu	Gly	Ala	Asp
			180					185					190		
Thr	Met	Ser	Ala	Gly	Gly	Gly	Gly	Pro	Leu	Gly	Asp	Asn	Asn	Gln	Gly
	195						200					205			
Ala	Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asp	Trp	His	Cys	Asp	Ser	Thr
	210					215					220				
Trp	Met	Gly	Asp	Arg	Val	Val	Thr	Lys	Ser	Thr	Arg	Thr	Trp	Val	Leu
225					230					235					240
Pro	Ser	Tyr	Asn	Asn	His	Gln	Tyr	Arg	Glu	Ile	Lys	Ser	Gly	Ser	Val
				245					250					255	
Asp	Gly	Ser	Asn	Ala	Asn	Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly
			260					265					270		
Tyr	Phe	Asp	Phe	Asn	Arg	Phe	His	Ser	His	Trp	Ser	Pro	Arg	Asp	Trp
	275						280					285			
Gln	Arg	Leu	Ile	Asn	Asn	Tyr	Trp	Gly	Phe	Arg	Pro	Arg	Ser	Leu	Arg
	290					295					300				
Val	Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Val	Gln	Asp	Ser
305					310					315					320
Thr	Thr	Thr	Ile	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr
				325					330					335	
Asp	Asp	Asp	Tyr	Gln	Leu	Pro	Tyr	Val	Val	Gly	Asn	Gly	Thr	Glu	Gly
			340					345					350		
Cys	Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val	Phe	Thr	Leu	Pro	Gln	Tyr	Gly
	355						360						365		
Tyr	Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr	Glu	Asn	Pro	Thr	Glu	Arg	Ser
	370					375					380				
Ser	Phe	Phe	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Lys	Met	Leu	Arg	Thr	Gly
385					390					395					400
Asn	Asn	Phe	Glu	Phe	Thr	Tyr	Asn	Phe	Glu	Glu	Val	Pro	Phe	His	Ser
				405					410					415	
Ser	Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe	Lys	Leu	Ala	Asn	Pro	Leu	Val
		420						425					430		
Asp	Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser	Thr	Asn	Asn	Thr	Gly	Gly	Val
		435					440						445		
Gln	Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg	Tyr	Ala	Asn	Thr	Tyr	Lys	Asn
	450					455					460				
Trp	Phe	Pro	Gly	Pro	Met	Gly	Arg	Thr	Gln	Gly	Trp	Asn	Leu	Gly	Ser
465					470					475					480
Gly	Val	Asn	Arg	Ala	Ser	Val	Ser	Ala	Phe	Ala	Thr	Thr	Asn	Arg	Met
				485					490					495	
Glu	Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val	Pro	Pro	Gln	Pro	Asn	Gly	Met
			500					505					510		
Thr	Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr	Tyr	Ala	Leu	Glu	Asn	Thr	Met
			515				520					525			

-continued

Ile Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu
 530 535 540
 Glu Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn
 545 550 555 560
 Arg Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser
 565 570 575
 Ser Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val
 580 585 590
 Pro Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile
 595 600 605
 Trp Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala
 610 615 620
 Met Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys
 625 630 635 640
 Asn Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val
 645 650 655
 Ser Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met
 660 665 670
 Glu Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile
 675 680 685
 Gln Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro
 690 695 700
 Asp Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr
 705 710 715 720
 Leu Thr Arg Pro Leu
 725

<210> SEQ ID NO 22
 <211> LENGTH: 725
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide
 <400> SEQUENCE: 22

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
 1 5 10 15
 Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
 20 25 30
 Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
 35 40 45
 Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
 50 55 60
 Val Asn Arg Ala Asp Glu Val Ala Arg Glu His Asp Ile Ser Tyr Asn
 65 70 75 80
 Glu Gln Leu Glu Ala Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
 85 90 95
 Asp Ala Glu Phe Gln Glu Lys Leu Ala Asp Asp Thr Ser Phe Gly Gly
 100 105 110
 Asn Leu Gly Lys Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
 115 120 125
 Phe Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Thr Gly Lys Arg
 130 135 140
 Ile Asp Asp His Phe Pro Lys Arg Lys Lys Ala Arg Thr Glu Glu Asp
 145 150 155 160

-continued

Ser Lys Pro Ser Thr Ser Ser Asp Ala Glu Ala Gly Pro Ser Gly Ser
 165 170 175

Gln Gln Leu Gln Ile Pro Ala Gln Pro Ala Ser Ser Leu Gly Ala Asp
 180 185 190

Thr Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly
 195 200 205

Ala Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr
 210 215 220

Trp Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu
 225 230 235 240

Pro Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val
 245 250 255

Asp Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
 260 265 270

Tyr Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp
 275 280 285

Gln Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg
 290 295 300

Val Lys Ile Phe Asn Ile Gln Val Lys Glu Val Thr Val Gln Asp Ser
 305 310 315 320

Thr Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Thr
 325 330 335

Asp Asp Asp Tyr Gln Leu Pro Tyr Val Val Gly Asn Gly Thr Glu Gly
 340 345 350

Cys Leu Pro Ala Phe Pro Pro Gln Val Phe Thr Leu Pro Gln Tyr Gly
 355 360 365

Tyr Ala Thr Leu Asn Arg Asp Asn Thr Glu Asn Pro Thr Glu Arg Ser
 370 375 380

Ser Phe Phe Cys Leu Glu Tyr Phe Pro Ser Lys Met Leu Arg Thr Gly
 385 390 395 400

Asn Asn Phe Glu Phe Thr Tyr Asn Phe Glu Glu Val Pro Phe His Ser
 405 410 415

Ser Phe Ala Pro Ser Gln Asn Leu Phe Lys Leu Ala Asn Pro Leu Val
 420 425 430

Asp Gln Tyr Leu Tyr Arg Phe Val Ser Thr Asn Asn Thr Gly Gly Val
 435 440 445

Gln Phe Asn Lys Asn Leu Ala Gly Arg Tyr Ala Asn Thr Tyr Lys Asn
 450 455 460

Trp Phe Pro Gly Pro Met Gly Arg Thr Gln Gly Trp Asn Leu Gly Ser
 465 470 475 480

Gly Val Asn Arg Ala Ser Val Ser Ala Phe Ala Thr Thr Asn Arg Met
 485 490 495

Glu Leu Glu Gly Ala Ser Tyr Gln Val Pro Pro Gln Pro Asn Gly Met
 500 505 510

Thr Asn Asn Leu Gln Gly Ser Asn Thr Tyr Ala Leu Glu Asn Thr Met
 515 520 525

Ile Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu
 530 535 540

Glu Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn
 545 550 555 560

Arg Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser
 565 570 575

-continued

Ser Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val
580 585 590

Pro Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile
595 600 605

Trp Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala
610 615 620

Met Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys
625 630 635 640

Asn Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val
645 650 655

Ser Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met
660 665 670

Glu Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile
675 680 685

Gln Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro
690 695 700

Asp Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr
705 710 715 720

Leu Thr Arg Pro Leu
725

<210> SEQ ID NO 23
 <211> LENGTH: 725
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 23

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
1 5 10 15

Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
20 25 30

Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Glu Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Lys Leu Ala Asp Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Lys Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
115 120 125

Phe Gly Leu Val Glu Glu Gly Ala Lys Thr Ala Pro Thr Gly Lys Arg
130 135 140

Ile Asp Asp His Phe Pro Lys Arg Lys Lys Ala Arg Thr Glu Glu Asp
145 150 155 160

Ser Lys Pro Ser Thr Ser Ser Asp Ala Glu Ala Gly Pro Ser Gly Ser
165 170 175

Gln Gln Leu Gln Ile Pro Ala Gln Pro Ala Ser Ser Leu Gly Ala Asp
180 185 190

Thr Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly
195 200 205

-continued

Ala	Asp	Gly	Val	Gly	Asn	Ala	Ser	Gly	Asp	Trp	His	Cys	Asp	Ser	Thr	210	215	220	
Trp	Met	Gly	Asp	Arg	Val	Val	Thr	Lys	Ser	Thr	Arg	Thr	Trp	Val	Leu	225	230	235	240
Pro	Ser	Tyr	Asn	Asn	His	Gln	Tyr	Arg	Glu	Ile	Lys	Ser	Gly	Ser	Val	245	250	255	
Asp	Gly	Ser	Asn	Ala	Asn	Ala	Tyr	Phe	Gly	Tyr	Ser	Thr	Pro	Trp	Gly	260	265	270	
Tyr	Phe	Asp	Phe	Asn	Arg	Phe	His	Ser	His	Trp	Ser	Pro	Arg	Asp	Trp	275	280	285	
Gln	Arg	Leu	Ile	Asn	Asn	Tyr	Trp	Gly	Phe	Arg	Pro	Arg	Ser	Leu	Arg	290	295	300	
Val	Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Val	Gln	Asp	Ser	305	310	315	320
Thr	Thr	Thr	Ile	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr	325	330	335	
Asp	Asp	Asp	Tyr	Gln	Leu	Pro	Tyr	Val	Val	Gly	Asn	Gly	Thr	Glu	Gly	340	345	350	
Cys	Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val	Phe	Thr	Leu	Pro	Gln	Tyr	Gly	355	360	365	
Tyr	Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr	Glu	Asn	Pro	Thr	Glu	Arg	Ser	370	375	380	
Ser	Phe	Phe	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Lys	Met	Leu	Arg	Thr	Gly	385	390	395	400
Asn	Asn	Phe	Glu	Phe	Thr	Tyr	Asn	Phe	Glu	Glu	Val	Pro	Phe	His	Ser	405	410	415	
Ser	Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe	Lys	Leu	Ala	Asn	Pro	Leu	Val	420	425	430	
Asp	Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser	Thr	Asn	Asn	Thr	Gly	Gly	Val	435	440	445	
Gln	Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg	Tyr	Ala	Asn	Thr	Tyr	Lys	Asn	450	455	460	
Trp	Phe	Pro	Gly	Pro	Met	Gly	Arg	Thr	Gln	Gly	Trp	Asn	Leu	Gly	Ser	465	470	475	480
Gly	Val	Asn	Arg	Ala	Ser	Val	Ser	Ala	Phe	Ala	Thr	Thr	Asn	Arg	Met	485	490	495	
Glu	Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val	Pro	Pro	Gln	Pro	Asn	Gly	Met	500	505	510	
Thr	Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr	Tyr	Ala	Leu	Glu	Asn	Thr	Met	515	520	525	
Ile	Phe	Asn	Ser	Gln	Pro	Ala	Asn	Pro	Gly	Thr	Thr	Ala	Thr	Tyr	Leu	530	535	540	
Glu	Gly	Asn	Met	Leu	Ile	Thr	Ser	Glu	Ser	Glu	Thr	Gln	Pro	Val	Asn	545	550	555	560
Arg	Val	Ala	Tyr	Asn	Val	Gly	Gly	Gln	Met	Ala	Thr	Asn	Asn	Gln	Ser	565	570	575	
Ser	Thr	Thr	Ala	Pro	Ala	Thr	Gly	Thr	Tyr	Asn	Leu	Gln	Glu	Ile	Val	580	585	590	
Pro	Gly	Ser	Val	Trp	Met	Glu	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile	595	600	605	
Trp	Ala	Lys	Ile	Pro	Glu	Thr	Gly	Ala	His	Phe	His	Pro	Ser	Pro	Ala	610	615	620	
Met	Gly	Gly	Phe	Gly	Leu	Lys	His	Pro	Pro	Pro	Met	Met	Leu	Ile	Lys				

-continued

260				265				270							
Tyr	Phe	Asp	Phe	Asn	Arg	Phe	His	Ser	His	Trp	Ser	Pro	Arg	Asp	Trp
		275					280					285			
Gln	Arg	Leu	Ile	Asn	Asn	Tyr	Trp	Gly	Phe	Arg	Pro	Arg	Ser	Leu	Arg
		290				295					300				
Val	Lys	Ile	Phe	Asn	Ile	Gln	Val	Lys	Glu	Val	Thr	Val	Gln	Asp	Ser
		305			310					315				320	
Thr	Thr	Thr	Ile	Ala	Asn	Asn	Leu	Thr	Ser	Thr	Val	Gln	Val	Phe	Thr
				325					330					335	
Asp	Asp	Asp	Tyr	Gln	Leu	Pro	Tyr	Val	Val	Gly	Asn	Gly	Thr	Glu	Gly
			340					345					350		
Cys	Leu	Pro	Ala	Phe	Pro	Pro	Gln	Val	Phe	Thr	Leu	Pro	Gln	Tyr	Gly
		355					360					365			
Tyr	Ala	Thr	Leu	Asn	Arg	Asp	Asn	Thr	Glu	Asn	Pro	Thr	Glu	Arg	Ser
		370				375					380				
Ser	Phe	Phe	Cys	Leu	Glu	Tyr	Phe	Pro	Ser	Lys	Met	Leu	Arg	Thr	Gly
				385		390				395				400	
Asn	Asn	Phe	Glu	Phe	Thr	Tyr	Asn	Phe	Glu	Glu	Val	Pro	Phe	His	Ser
				405					410					415	
Ser	Phe	Ala	Pro	Ser	Gln	Asn	Leu	Phe	Lys	Leu	Ala	Asn	Pro	Leu	Val
			420					425				430			
Asp	Gln	Tyr	Leu	Tyr	Arg	Phe	Val	Ser	Thr	Asn	Asn	Thr	Gly	Gly	Val
		435					440					445			
Gln	Phe	Asn	Lys	Asn	Leu	Ala	Gly	Arg	Tyr	Ala	Asn	Thr	Tyr	Lys	Asn
		450				455					460				
Trp	Phe	Pro	Gly	Pro	Met	Gly	Arg	Thr	Gln	Gly	Trp	Asn	Leu	Gly	Ser
				465		470				475				480	
Gly	Val	Asn	Arg	Ala	Ser	Val	Ser	Ala	Phe	Ala	Thr	Thr	Asn	Arg	Met
				485					490					495	
Glu	Leu	Glu	Gly	Ala	Ser	Tyr	Gln	Val	Pro	Pro	Gln	Pro	Asn	Gly	Met
			500					505				510			
Thr	Asn	Asn	Leu	Gln	Gly	Ser	Asn	Thr	Tyr	Ala	Leu	Glu	Asn	Thr	Met
			515				520					525			
Ile	Phe	Asn	Ser	Gln	Pro	Ala	Asn	Pro	Gly	Thr	Thr	Ala	Thr	Tyr	Leu
				530		535					540				
Glu	Gly	Asn	Met	Leu	Ile	Thr	Ser	Glu	Ser	Glu	Thr	Gln	Pro	Val	Asn
				545		550				555				560	
Arg	Val	Ala	Tyr	Asn	Val	Gly	Gly	Gln	Met	Ala	Thr	Asn	Asn	Gln	Ser
				565					570					575	
Ser	Thr	Thr	Ala	Pro	Ala	Thr	Gly	Thr	Tyr	Asn	Leu	Gln	Glu	Ile	Val
			580					585				590			
Pro	Gly	Ser	Val	Trp	Met	Glu	Arg	Asp	Val	Tyr	Leu	Gln	Gly	Pro	Ile
			595			600						605			
Trp	Ala	Lys	Ile	Pro	Glu	Thr	Gly	Ala	His	Phe	His	Pro	Ser	Pro	Ala
				610		615					620				
Met	Gly	Gly	Phe	Gly	Leu	Lys	His	Pro	Pro	Pro	Met	Met	Leu	Ile	Lys
				625		630				635				640	
Asn	Thr	Pro	Val	Pro	Gly	Asn	Ile	Thr	Ser	Phe	Ser	Asp	Val	Pro	Val
				645				650				655			
Ser	Ser	Phe	Ile	Thr	Gln	Tyr	Ser	Thr	Gly	Gln	Val	Thr	Val	Glu	Met
			660					665				670			
Glu	Trp	Glu	Leu	Lys	Lys	Glu	Asn	Ser	Lys	Arg	Trp	Asn	Pro	Glu	Ile
			675			680						685			

-continued

Gln Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro
690 695 700

Asp Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr
705 710 715 720

Leu Thr Arg Pro Leu
725

<210> SEQ ID NO 25
 <211> LENGTH: 725
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic polypeptide

<400> SEQUENCE: 25

Met Ala Ala Asp Gly Tyr Leu Pro Asp Trp Leu Glu Asp Thr Leu Ser
1 5 10 15

Glu Gly Ile Arg Gln Trp Trp Lys Leu Lys Pro Gly Pro Pro Pro Pro
20 25 30

Lys Pro Ala Glu Arg His Lys Asp Asp Ser Arg Gly Leu Val Leu Pro
35 40 45

Gly Tyr Lys Tyr Leu Gly Pro Phe Asn Gly Leu Asp Lys Gly Glu Pro
50 55 60

Val Asn Glu Ala Asp Ala Ala Ala Leu Glu His Asp Lys Ala Tyr Asp
65 70 75 80

Arg Gln Leu Asp Ser Gly Asp Asn Pro Tyr Leu Lys Tyr Asn His Ala
85 90 95

Asp Ala Glu Phe Gln Glu Arg Leu Lys Glu Asp Thr Ser Phe Gly Gly
100 105 110

Asn Leu Gly Arg Ala Val Phe Gln Ala Lys Lys Arg Val Leu Glu Pro
115 120 125

Leu Gly Leu Val Glu Glu Pro Val Lys Thr Ala Pro Thr Gly Lys Arg
130 135 140

Ile Asp Asp His Phe Pro Lys Arg Lys Lys Ala Arg Thr Glu Glu Asp
145 150 155 160

Ser Lys Pro Ser Thr Ser Ser Asp Ala Glu Ala Gly Pro Ser Gly Ser
165 170 175

Gln Gln Leu Gln Ile Pro Ala Gln Pro Ala Ser Ser Leu Gly Ala Asp
180 185 190

Thr Met Ser Ala Gly Gly Gly Gly Pro Leu Gly Asp Asn Asn Gln Gly
195 200 205

Ala Asp Gly Val Gly Asn Ala Ser Gly Asp Trp His Cys Asp Ser Thr
210 215 220

Trp Met Gly Asp Arg Val Val Thr Lys Ser Thr Arg Thr Trp Val Leu
225 230 235 240

Pro Ser Tyr Asn Asn His Gln Tyr Arg Glu Ile Lys Ser Gly Ser Val
245 250 255

Asp Gly Ser Asn Ala Asn Ala Tyr Phe Gly Tyr Ser Thr Pro Trp Gly
260 265 270

Tyr Phe Asp Phe Asn Arg Phe His Ser His Trp Ser Pro Arg Asp Trp
275 280 285

Gln Arg Leu Ile Asn Asn Tyr Trp Gly Phe Arg Pro Arg Ser Leu Arg
290 295 300

Val Lys Ile Phe Asn Ile Gln Val Lys Glu Val Thr Val Gln Asp Ser
305 310 315 320

-continued

Thr Thr Thr Ile Ala Asn Asn Leu Thr Ser Thr Val Gln Val Phe Thr
 325 330 335
 Asp Asp Asp Tyr Gln Leu Pro Tyr Val Val Gly Asn Gly Thr Glu Gly
 340 345 350
 Cys Leu Pro Ala Phe Pro Pro Gln Val Phe Thr Leu Pro Gln Tyr Gly
 355 360 365
 Tyr Ala Thr Leu Asn Arg Asp Asn Thr Glu Asn Pro Thr Glu Arg Ser
 370 375 380
 Ser Phe Phe Cys Leu Glu Tyr Phe Pro Ser Lys Met Leu Arg Thr Gly
 385 390 395 400
 Asn Asn Phe Glu Phe Thr Tyr Asn Phe Glu Glu Val Pro Phe His Ser
 405 410 415
 Ser Phe Ala Pro Ser Gln Asn Leu Phe Lys Leu Ala Asn Pro Leu Val
 420 425 430
 Asp Gln Tyr Leu Tyr Arg Phe Val Ser Thr Asn Asn Thr Gly Gly Val
 435 440 445
 Gln Phe Asn Lys Asn Leu Ala Gly Arg Tyr Ala Asn Thr Tyr Lys Asn
 450 455 460
 Trp Phe Pro Gly Pro Met Gly Arg Thr Gln Gly Trp Asn Leu Gly Ser
 465 470 475 480
 Gly Val Asn Arg Ala Ser Val Ser Ala Phe Ala Thr Thr Asn Arg Met
 485 490 495
 Glu Leu Glu Gly Ala Ser Tyr Gln Val Pro Pro Gln Pro Asn Gly Met
 500 505 510
 Thr Asn Asn Leu Gln Gly Ser Asn Thr Tyr Ala Leu Glu Asn Thr Met
 515 520 525
 Ile Phe Asn Ser Gln Pro Ala Asn Pro Gly Thr Thr Ala Thr Tyr Leu
 530 535 540
 Glu Gly Asn Met Leu Ile Thr Ser Glu Ser Glu Thr Gln Pro Val Asn
 545 550 555 560
 Arg Val Ala Tyr Asn Val Gly Gly Gln Met Ala Thr Asn Asn Gln Ser
 565 570 575
 Ser Thr Thr Ala Pro Ala Thr Gly Thr Tyr Asn Leu Gln Glu Ile Val
 580 585 590
 Pro Gly Ser Val Trp Met Glu Arg Asp Val Tyr Leu Gln Gly Pro Ile
 595 600 605
 Trp Ala Lys Ile Pro Glu Thr Gly Ala His Phe His Pro Ser Pro Ala
 610 615 620
 Met Gly Gly Phe Gly Leu Lys His Pro Pro Pro Met Met Leu Ile Lys
 625 630 635 640
 Asn Thr Pro Val Pro Gly Asn Ile Thr Ser Phe Ser Asp Val Pro Val
 645 650 655
 Ser Ser Phe Ile Thr Gln Tyr Ser Thr Gly Gln Val Thr Val Glu Met
 660 665 670
 Glu Trp Glu Leu Lys Lys Glu Asn Ser Lys Arg Trp Asn Pro Glu Ile
 675 680 685
 Gln Tyr Thr Asn Asn Tyr Asn Asp Pro Gln Phe Val Asp Phe Ala Pro
 690 695 700
 Asp Ser Thr Gly Glu Tyr Arg Thr Thr Arg Pro Ile Gly Thr Arg Tyr
 705 710 715 720
 Leu Thr Arg Pro Leu
 725

-continued

<210> SEQ ID NO 26
 <211> LENGTH: 1386
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic construct

 <400> SEQUENCE: 26

 atgcagaggg tgaacatgat catggctgag agccctggcc tgatcacat ctgectgctg 60
 ggctacctgc tgtctgetga gtgcactgtg ttcttgacc atgagaatgc caacaagatc 120
 ctgaacaggc ccaagagata caactctggc aagctggagg agtttgtgca gggcaacctg 180
 gagagggagt gcatggagga gaagtgcagc tttgaggagg ccaggagggt gtttgagaac 240
 actgagagga cactgagtt ctggaagcag tatgtggatg gggaccagtg tgagagcaac 300
 cctgcctga atgggggag ctgcaaggat gacatcaaca gctatgagt ctggtgcccc 360
 tttggctttg agggcaagaa ctgtgagctg gatgtgacct gcaacatcaa gaatggcaga 420
 tgtgagcagt tctgcaagaa ctctgctgac aacaaggtgg tgtgcagctg cactgagggc 480
 tacaggctgg ctgagaacca gaagagctgt gagcctgctg tgccattccc atgtggcaga 540
 gtgtctgtga gccagaccag caagctgacc agggctgagg ctgtgttccc tgatgtggac 600
 tatgtgaaca gactgaggc tgaaccatc ctggacaaca taccagag caccagagc 660
 ttcaatgact tcaccagggt ggtggggggg gaggatgcca agcctggcca gttcccctgg 720
 caagtgggct tgaatggcaa ggtggatgcc ttctgtgggg gcagcattgt gaatgagaag 780
 tggattgtga ctgctgcca ctgtgtggag actgggggta agatcactgt ggtggctggg 840
 gagcacaaca ttgaggagac tgagcact gagcagaaga ggaatgtgat caggatcacc 900
 cccaccaca actacaatgc tgccatcaac aagtacaacc atgacattgc cctgctggag 960
 ctggatgagc cctggtgct gaacagctat gtgaccccc tctgcattgc tgacaaggag 1020
 tacaccaaca tcttctgaa gtttggctct ggctatgtgt ctggctgggg cagggtgttc 1080
 cacaagggca ggtctgcct ggtgctgcag tacctgaggg tgcccctggt ggacagggcc 1140
 acctgctgc tgagcaccaa gttcaccatc tacaacaaca tgttctgtgc tggcttccat 1200
 gaggggggca gggacagctg ccagggggac tctggggggc cccatgtgac tgaggtggag 1260
 ggcaccagct tctgactgg catcatcagc tggggggagg agtgtgcat gaagggcaag 1320
 tatggcatct acaccaaagt ctccagatat gtgaactgga tcaaggagaa gaccaagctg 1380
 acctga 1386

The invention claimed is:

1. A polynucleotide comprising a Factor IX nucleotide sequence corresponding to a reference wild-type Factor IX sequence of SEQ ID NO: 9, wherein:

- (i) the Factor IX nucleotide sequence comprises a coding sequence that encodes a Factor IX protein or fragment thereof;
- (ii) a portion of the coding sequence is codon optimized compared to the reference wild-type Factor IX sequence of SEQ ID NO: 9;
- (iii) the portion of the coding sequence that is codon optimized is at least 1100 nucleotides in length;
- (iv) in the portion that is codon optimized at least 73% of codons are selected from the group consisting of:
 TTC encoding phenylalanine,
 CTG encoding leucine,
 ATC encoding isoleucine,

50

55

60

65

- GTG or GTC encoding valine,
 AGC encoding serine,
 CCC encoding proline,
 ACC encoding threonine,
 GCC encoding alanine,
 TAC encoding tyrosine,
 CAC encoding histidine,
 CAG encoding glutamine,
 AAC encoding asparagine,
 AAA or AAG encoding lysine,
 GAC encoding aspartate,
 TGC encoding cysteine,
 AGG encoding arginine,
 GGC encoding glycine, and
 GAG encoding glutamate;
- (v) the portion of the coding sequence that is codon optimized is CpG free;

135

- (vi) the polynucleotide further comprises a transcription regulatory element comprising:
- (a) an A1AT promoter or a fragment of an A1AT promoter between 150 and 300 nucleotides in length; and/or
 - (b) an HCR enhancer or a fragment of an HCR enhancer between 100 and 250 nucleotides in length; and
- (vii) the Factor IX nucleotide sequence comprises a codon that encodes a leucine at a position corresponding to position 384 of the reference wild type Factor IX sequence of SEQ ID NO: 9.
2. The polynucleotide of claim 1, wherein the polynucleotide comprises a promoter selected from the group consisting of:
- (i) a promoter comprising at least 80% identity to SEQ ID NO. 14;
 - (ii) a promoter comprising at least 98% identity to SEQ ID NO. 14;
 - (iii) a promoter comprising the sequence of SEQ ID NO. 14; and
 - (iv) a liver-specific promoter.
3. The polynucleotide of claim 1, wherein the portion of the coding sequence that is codon optimized comprises at least a portion of exon 3, at least a portion of exon 4, at least a portion of exon 5, at least a portion of exon 6, at least a portion of exon 7, and at least a portion of exon 8.
4. The polynucleotide of claim 1, wherein, in the portion of the coding sequence that is codon optimized, the codon optimization is selected from the group consisting of:
- (i) a) at least 1, at least 2, at least 4, or at least 5 codons that encode phenylalanine is/are replaced with TTC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
 - b) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC;
 - c) at least 60%, at least 65%, or at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT; or
 - d) the codons that encode phenylalanine are TTC, except where the following codon starts with a G;
 - (ii) a) at least 5, at least 10, at least 15, or at least 16 codons that encode leucine is/are replaced with CTG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
 - b) at least 90%, or at least 94% of the codons that encode leucine are CTG; or
 - c) at least 90%, or at least 94% of the codons that encode leucine are CTG and the remainder are CTC;
 - (iii) a) at least 5, at least 10, at least 11, or at least 12 codons that encode isoleucine is/are replaced with ATC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
 - b) at least 1 of codon ATC is/are replaced with ATT compared to a reference wild type Factor IX sequence, where the following codon starts with a G;
 - c) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC;
 - d) at least 60%, at least 70%, or at least 75% of the codons that encode isoleucine are ATC and the remainder are ATT; or
 - e) the codons that encode isoleucine are ATC, except where the following codon starts with a G;
 - (iv) a) at least 10, at least 15, at least 20, or at least 25 codons that encode valine is/are replaced with GTG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;

136

- b) at least 1 codon that encodes valine is/are replaced with GTC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- c) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG; or
- d) at least 80%, at least 90%, or at least 95% of the codons that encode valine are GTG and the remainder are GTC;
- (v) a) at least 5, at least 10, at least 12, or at least 13 codons that encode serine is/are replaced with AGC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1, at least 2, or at least 4 codons that encode serine is/are replaced with TCT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC; or
- d) at least 60%, at least 65%, or at least 70% of the codons that encode serine are AGC and the remainder are TCT or TCC;
- (vi) a) at least 1, at least 2, or at least 5 codons that encode proline is/are replaced with CCC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 codon that encodes proline is/are replaced with CCT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 50%, at least 55%, or at least 60% of the codons that encode proline are CCC;
- d) at least 50%, at least 55%, or at least 60% of the codons that encode proline are CCC and the remainder are CCA or CCT; or
- e) the codons that encode proline are CCC, except where the following codon starts with a G;
- (vii) a) at least 6, at least 7, at least 8, or at least 10 codons that encode threonine is/are replaced with ACC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1, or at least 2 codons that encode threonine is/are replaced with ACT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC;
- d) at least 45%, at least 50%, or at least 55% of the codons that encode threonine are ACC and the remainder are ACT; or
- e) the codons that encode threonine are ACC, except where the following codon starts with a G;
- (viii) a) at least 1, at least 2, or at least 3 codons that encode alanine is/are replaced with GCC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1, at least 2, at least 3, or at least 4 codons that encode alanine is/are replaced with GCT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 35%, at least 40%, or at least 43% of the codons that encode alanine are GCC;
- d) at least 35%, at least 40%, or at least 43% of the codons that encode alanine are GCC and the remainder are GCT; or
- e) the codons that encode alanine are GCC, except where the following codon starts with a G;

137

- (ix) a) at least 1, or at least 2 codons that encode tyrosine is/are replaced with TAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 of codon TAC is/are replaced with TAT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC;
- d) at least 40%, at least 45%, or at least 48% of the codons that encode tyrosine are TAC and the remainder are TAT; or
- e) the codons that encode tyrosine are TAC, except where the following codon starts with a G;
- (x) a) at least 1 codon that encodes histidine is/are replaced with CAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 50%, at least 60%, or at least 65% of codons that encode histidine are CAC;
- c) at least 50%, at least 60%, or at least 65% of the codons that encode histidine are CAC and the remainder are CAT; or
- d) the codons that encode histidine are CAC, except where the following codon starts with a G;
- (xi) a) at least 1, at least 2, at least 4, or at least 5 codons that encode glutamine is/are replaced with CAG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 of codon CAG is/are replaced with CAA compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- c) at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG; or
- d) at least 80%, at least 85%, or at least 90% of the codons that encode glutamine are CAG and the remainder are CAA;
- (xii) a) at least 1, at least 2, at least 4, or at least 5 codons that encode asparagine is/are replaced with AAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC;
- c) at least 60%, at least 65%, or at least 70% of the codons that encode asparagine are AAC and the remainder are AAT; or
- d) the codons that encode asparagine are AAC, except where the following codon starts with a G;
- (xiii) a) at least 5, at least 7, at least 8, or at least 9 codons that encode lysine is/are replaced with AAG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 of codon AAG is/are replaced with AAA compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- c) at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG; or
- d) at least 80%, at least 90%, or at least 95% of the codons that encode lysine are AAG and the remainder are AAA;
- (xiv) a) at least 1, at least 2, at least 3, or at least 4 codons that encode aspartate is/are replaced with GAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 of codon GAC is/are replaced with GAT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;

138

- c) at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC;
- d) at least 45%, at least 50%, or at least 60% of the codons that encode aspartate are GAC and the remainder are GAT; or
- e) the codons that encode aspartate are GAC, except where the following codon starts with a G;
- (xv) a) at least 15, at least 20, at least 25, or at least 26 codons that encode glutamate is/are replaced with GAG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG; or
- c) at least 80%, at least 90%, or at least 95% of the codons that encode glutamate are GAG and the remainder are GAA;
- (xvi) a) at least 5, at least 6, at least 7, or at least 8 codons that encode cysteine is/are replaced with TGC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 of codon TGC is/are replaced with TGT compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC;
- d) at least 40%, at least 50%, or at least 55% of the codons that encode cysteine are TGC and the remainder are TGT; or
- e) the codons that encode cysteine are TGC, except where the following codon starts with a G;
- (xvii) wherein, codons that encode tryptophan are TGG;
- (xviii) a) at least 5, at least 8, at least 10, or at least 11 codons that encode arginine is/are replaced with AGG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 1 codon that encodes arginine is/are replaced with AGA compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- c) at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG; or
- d) at least 60%, at least 70%, or at least 75% of the codons that encode arginine are AGG and the remainder are AGA; and
- (xix) a) at least 5, at least 10, at least 12, or at least 13 codons that encode glycine is/are replaced with GGC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 5, at least 6, at least 7, or at least 8 codons that encode glycine is/are replaced with GGG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9, where the following codon starts with a G;
- c) at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC;
- d) at least 50%, at least 55%, or at least 60% of the codons that encode glycine are GGC and the remainder are GGG; or
- e) the codons that encode glycine are GGC, except where the following codon starts with a G.
5. The polynucleotide of claim 1, wherein the portion of the coding sequence that is codon optimized comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the portion of the coding sequence that is codon optimized:

- a) at least 5 codons that encode phenylalanine is/are replaced with TTC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- b) at least 16 codons that encode leucine is/are replaced with CTG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- c) at least 12 codons that encode isoleucine is/are replaced with ATC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- d) at least 25 codons that encode valine is/are replaced with GTG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- e) at least 13 codons that encode serine is/are replaced with AGC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- f) at least 5 codons that encode proline is/are replaced with CCC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- g) at least 10 codons that encode threonine is/are replaced with ACC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- h) at least 4 codons that encode alanine is/are replaced with GCC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- i) at least 2 codons that encode tyrosine is/are replaced with TAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- j) at least 1 codon that encodes histidine is/are replaced with CAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- k) at least 5 codons that encode glutamine is/are replaced with CAG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- l) at least 5 codons that encode asparagine is/are replaced with AAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- m) at least 9 codons that encode lysine is/are replaced with AAG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- n) at least 4 codons that encode aspartate is/are replaced with GAC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- o) at least 26 codons that encode glutamate is/are replaced with GAG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- p) at least 8 codons that encode cysteine is/are replaced with TGC compared to the reference wild type Factor IX sequence of SEQ ID NO: 9;
- q) codons that encode tryptophan are TGG;
- r) at least 11 codons that encode arginine is/are replaced with AGG compared to the reference wild type Factor IX sequence of SEQ ID NO: 9; and
- s) at least 13 codons that encode glycine is/are replaced with GGC compared the reference wild type Factor IX sequence of SEQ ID NO: 9.
6. The polynucleotide of claim 1, wherein the portion of the coding sequence that is codon optimized comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the portion of the coding sequence that is codon optimized:
- a) at least 70% of codons that encode phenylalanine are TTC;
- b) at least 94% of codons that encode leucine are CTG;
- c) at least 75% of codons that encode isoleucine are ATC;
- d) at least 95% of codons that encode valine are GTG;

- e) at least 70% of codons that encode serine are AGC;
- f) at least 60% of codons that encode proline are CCC;
- g) at least 55% of codons that encode threonine are ACC;
- h) at least 43% of codons that encode alanine are GCC;
- i) at least 48% of codons that encode tyrosine are TAC;
- j) at least 65% of codons that encode histidine are CAC;
- k) at least 90% of codons that encode glutamine are CAG;
- l) at least 70% of codons that encode asparagine are AAC;
- m) at least 95% of codons that encode lysine are AAG;
- n) at least 60% of codons that encode aspartate are GAC;
- o) at least 95% of codons that encode glutamate are GAG;
- p) at least 55% of codons that encode cysteine are TGC;
- q) codons that encode tryptophan are TGG;
- r) at least 75% of codons that encode arginine are AGG; and
- s) at least 60% of codons that encode glycine are GGC.
7. The polynucleotide of claim 1, wherein the portion of the coding sequence that is codon optimized comprises codons encoding phenylalanine, leucine, isoleucine, valine, serine, proline, threonine, alanine, tyrosine, histidine, glutamine, asparagine, lysine, aspartate, glutamate, cysteine, tryptophan, arginine, and glycine, and in the portion of the coding sequence that is codon optimized:
- a) at least 70% of the codons that encode phenylalanine are TTC and the remainder are TTT;
- b) at least 94% of codons that encode leucine are CTG and the remainder are CTC;
- c) at least 75% of codons that encode isoleucine are ATC and the remainder are ATT;
- d) at least 95% of codons that encode valine are GTG;
- e) at least 70% of codons that encode serine are AGC;
- f) at least 60% of codons that encode proline are CCC and the remainder are CCA or CCT;
- g) at least 55% of codons that encode threonine are ACC and the remainder are ACT;
- h) at least 43% of codons that encode alanine are GCC and the remainder are GCT;
- i) at least 48% of codons that encode tyrosine are TAC and the remainder are TAT;
- j) at least 65% of codons that encode histidine are CAC and the remainder are CAT;
- k) at least 90% of codons that encode glutamine are CAG and the remainder are CAA;
- l) at least 70% of codons that encode asparagine are AAC and the remainder are AAT;
- m) at least 95% of codons that encode lysine are AAG and the remainder are AAA;
- n) at least 60% of codons that encode aspartate are GAC and the remainder are GAT;
- o) at least 95% of codons that encode glutamate are GAG and the remainder are GAA;
- p) at least 55% of codons that encode cysteine are TGC and the remainder are TGT;
- q) codons that encode tryptophan are TGG;
- r) at least 75% of codons that encode arginine are AGG and the remainder are AGA; and
- s) at least 60% of codons that encode glycine are GGC and the remainder are GGG.
8. The polynucleotide of claim 1, wherein the polynucleotide comprises an enhancer selected from the group consisting of:
- (i) an enhancer comprising at least 80% identity to SEQ ID NO. 13;
- (ii) an enhancer comprising at least 98% identity to SEQ ID NO. 13; and
- (iii) an enhancer comprising the sequence of SEQ ID NO. 13.

9. The polynucleotide of claim 2, wherein a polypeptide encoded by the Factor IX nucleotide sequence is expressed in human liver cells at higher levels compared to the reference wild type Factor IX sequence of SEQ ID NO: 9.

10. The polynucleotide of claim 1, wherein the codon that encodes an amino acid at the position corresponding to position 384 of the reference wild type Factor IX sequence of SEQ ID NO: 9 is CTC. 5

11. The polynucleotide of claim 1, wherein the codon that encodes an amino acid at the position corresponding to position 384 of the reference wild type Factor IX sequence of SEQ ID NO: 9 is CTG. 10

12. An AAV viral particle comprising a recombinant genome comprising the polynucleotide of claim 1.

13. The AAV viral particle of claim 12, wherein the recombinant genome comprises two resolvable ITRs. 15

14. The AAV viral particle of claim 12, wherein after transduction into a population of Huh7 cells, the AAV viral particle expresses the Factor IX protein or a fragment thereof, wherein the Factor IX protein or the fragment thereof has an activity greater than an activity of a Factor IX protein or a fragment thereof expressed from a comparable AAV viral particle comprising a Factor IX nucleotide sequence of SEQ ID NO: 12 and a transcription regulatory element of SEQ ID NO: 7 transduced into a comparable population of Huh7 cells. 20 25

* * * * *