



US011708412B2

(12) **United States Patent**
Johnson et al.

(10) **Patent No.:** **US 11,708,412 B2**
(45) **Date of Patent:** **Jul. 25, 2023**

(54) **METHODS FOR TREATING HEMATOLOGIC CANCERS**

(71) Applicants: **Novartis AG**, Basel (CH); **The Medical College of Wisconsin, Inc.**, Milwaukee, WI (US)

(72) Inventors: **Bryon Duane Johnson**, Jackson, WI (US); **Robert Millman**, Boston, MA (US)

(73) Assignees: **Novartis AG**, Basel (CH); **The Medical College of Wisconsin, Inc.**, Milwaukee, WI (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 206 days.

(21) Appl. No.: **16/724,759**

(22) Filed: **Dec. 23, 2019**

(65) **Prior Publication Data**

US 2020/0377600 A1 Dec. 3, 2020

Related U.S. Application Data

(62) Division of application No. 15/024,396, filed as application No. PCT/US2014/057491 on Sep. 25, 2014, now Pat. No. 10,570,204.

(60) Provisional application No. 62/017,192, filed on Jun. 25, 2014, provisional application No. 61/882,702, filed on Sep. 26, 2013.

(51) **Int. Cl.**

C07K 16/28 (2006.01)
A61K 39/395 (2006.01)
A61P 35/00 (2006.01)
A61K 47/68 (2017.01)
A61K 51/10 (2006.01)
A61N 5/10 (2006.01)
A61K 39/00 (2006.01)

(52) **U.S. Cl.**

CPC **C07K 16/2827** (2013.01); **A61K 39/39558** (2013.01); **A61K 47/6849** (2017.08); **A61K 51/1096** (2013.01); **A61N 5/10** (2013.01); **C07K 16/2803** (2013.01); **C07K 16/2818** (2013.01); **A61K 2039/505** (2013.01); **A61K 2039/507** (2013.01); **C07K 2317/24** (2013.01); **C07K 2317/31** (2013.01); **C07K 2317/54** (2013.01); **C07K 2317/55** (2013.01); **C07K 2317/622** (2013.01); **C07K 2317/626** (2013.01); **C07K 2317/76** (2013.01)

(58) **Field of Classification Search**

None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,417,984 A 3/1947 Kingsley
2,581,113 A 1/1952 Chester et al.

5,324,514 A 6/1994 Sipos
5,434,131 A 7/1995 Linsley et al.
5,629,204 A 5/1997 Honjo et al.
5,698,520 A 12/1997 Honjo et al.
5,811,097 A 9/1998 Allison et al.
5,897,862 A 4/1999 Hardy et al.
5,968,511 A 10/1999 Akita et al.
6,051,227 A 4/2000 Allison et al.
6,084,083 A 7/2000 Levinson
6,204,371 B1 3/2001 Levinson
6,288,218 B1 9/2001 Levinson
6,414,117 B1 7/2002 Levinson
6,562,343 B1 5/2003 Levinson
6,632,927 B2 10/2003 Adair et al.
6,632,976 B1 10/2003 Tomizuka et al.
6,803,192 B1 10/2004 Chen
6,808,710 B1 10/2004 Wood et al.
6,936,704 B1 8/2005 Freeman et al.
7,029,674 B2 4/2006 Carreno et al.
7,038,013 B2 5/2006 Freeman et al.
7,041,474 B2 5/2006 Kingsbury
7,101,550 B2 9/2006 Wood et al.
7,105,328 B2 9/2006 Wood et al.
7,122,372 B2 10/2006 Hardy et al.
7,169,791 B2 1/2007 Breitenstein et al.
7,172,750 B2 2/2007 Levinson
7,329,639 B2 2/2008 Hardy et al.
7,332,582 B2 2/2008 Hardy et al.
7,414,171 B2 8/2008 Honjo et al.
7,423,128 B2 9/2008 Gazit-Bomstein et al.
7,449,300 B2 11/2008 Chen et al.
7,470,428 B2 12/2008 Kuchroo et al.
7,488,802 B2 2/2009 Collins et al.
7,521,051 B2 4/2009 Collins et al.
7,524,498 B2 4/2009 Hardy et al.
7,553,939 B2 6/2009 McIntire et al.
7,563,869 B2 7/2009 Honjo et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 774391 B2 6/2004
CN 102492038 A 6/2012

(Continued)

OTHER PUBLICATIONS

Restifo et al. (2012) Adoptive immunotherapy for cancer: harnessing the T cell response. *Nat Rev Immunol.* 12(4): 269-281.*
Ding et al. (2012) Polyfunctional CD4+ T cells are essential for eradicating advanced B-cell lymphoma after chemotherapy. *Blood* 120(11): 2229-2239.*
Dudley et al. (2005) Adoptive cell transfer therapy following non-myeloablative but lymphodepleting chemotherapy for the treatment of patients with refractory metastatic melanoma. *J Clin Oncol.* 23(10): 2346-2357.*
Kearl et al. (2013) Programmed Death Receptor-1/Programmed Death Receptor Ligand-1 Blockade after Transient Lymphodepletion To Treat Myeloma. *J Immunol.* 190: 5620-5628 (prepublished online Apr. 24, 2013).*

(Continued)

Primary Examiner — Ilia I Ouspenski
(74) *Attorney, Agent, or Firm* — K&L Gates LLP

(57) **ABSTRACT**

The present invention relates to methods of treating hematologic cancers using a combination of inhibitors of PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA4.

12 Claims, 20 Drawing Sheets

Specification includes a Sequence Listing.

(56)

References Cited

U.S. PATENT DOCUMENTS

7,595,048 B2	9/2009	Honjo et al.	9,884,913 B2	2/2018	Sabatos-Peyton et al.
7,598,257 B2	10/2009	Rodgers et al.	9,908,936 B2	3/2018	Triebel et al.
7,635,757 B2	12/2009	Freeman et al.	9,944,645 B2	4/2018	Zhuo et al.
7,695,715 B2	4/2010	Hardy et al.	9,988,452 B2	6/2018	Freeman et al.
7,722,868 B2	5/2010	Freeman et al.	10,344,089 B2 *	7/2019	Thudium A61P 31/12
7,767,675 B2	8/2010	Zhuo et al.	10,251,893 B2	9/2019	Cheng et al.
7,794,710 B2	9/2010	Chen et al.	10,472,419 B2	11/2019	Sabatos-Peyton et al.
7,838,220 B2	11/2010	McIntire et al.	10,570,204 B2 *	2/2020	Johnson A61K 47/6849
7,858,746 B2	12/2010	Honjo et al.	10,711,060 B2	7/2020	Triebel et al.
7,943,743 B2	5/2011	Korman et al.	10,752,687 B2	8/2020	Freeman et al.
8,008,449 B2	8/2011	Korman et al.	10,851,165 B2	12/2020	Freeman et al.
8,039,273 B2	10/2011	Jeffrey	10,981,990 B2	4/2021	Sabatos-Peyton et al.
8,039,479 B2	10/2011	Michellys et al.	11,155,620 B2	10/2021	Sabatos-Peyton et al.
8,088,905 B2	1/2012	Collins et al.	11,344,620 B2	5/2022	Lebwohl et al.
8,101,176 B2	1/2012	Kuchroo et al.	2002/0164660 A1	11/2002	Spaulding et al.
8,168,179 B2	5/2012	Honjo et al.	2003/0039653 A1	2/2003	Chen et al.
8,168,757 B2	5/2012	Finnefrock et al.	2003/0232323 A1	12/2003	Freeman et al.
8,217,149 B2	7/2012	Irving et al.	2004/0073957 A1	4/2004	Tomizuka et al.
8,287,856 B2	10/2012	Li et al.	2004/0241745 A1	12/2004	Honjo et al.
8,329,660 B2	12/2012	Kuchroo et al.	2005/0191721 A1	9/2005	Kuchroo et al.
8,354,509 B2	1/2013	Carven et al.	2005/0276756 A1	12/2005	Hoo et al.
8,361,736 B2	1/2013	Majeti et al.	2006/0110383 A1	5/2006	Honjo et al.
8,415,355 B2	4/2013	Brain et al.	2006/0210567 A1	9/2006	Collins et al.
8,460,886 B2	6/2013	Shibayama et al.	2007/0041982 A1	2/2007	Ponath et al.
8,460,927 B2	6/2013	Chen	2007/0065427 A1	3/2007	Freeman et al.
8,461,330 B2	6/2013	Zhuo et al.	2007/0071675 A1	3/2007	Wu et al.
8,501,758 B2	8/2013	Huang et al.	2007/0122378 A1	5/2007	Freeman et al.
8,546,336 B2	10/2013	Chen et al.	2007/0202100 A1	8/2007	Wood et al.
8,546,396 B2	10/2013	Cheng et al.	2008/0025979 A1	1/2008	Honjo et al.
8,552,002 B2	10/2013	Ding et al.	2008/0167287 A1	7/2008	Zhuo et al.
8,552,154 B2	10/2013	Freeman et al.	2008/0311117 A1	12/2008	Collins et al.
8,552,156 B2	10/2013	Takayanagi et al.	2009/0055944 A1	2/2009	Korman et al.
8,568,728 B2	10/2013	Jeffrey	2009/0076250 A1	3/2009	Honjo et al.
8,580,247 B2	11/2013	Li et al.	2009/0155275 A1	6/2009	Wu et al.
8,609,089 B2	12/2013	Langermann et al.	2009/0304693 A1	12/2009	Ghayur et al.
8,617,546 B2	12/2013	Kang et al.	2009/0311253 A1	12/2009	Ghayur et al.
8,647,623 B2	2/2014	Takayanagi et al.	2010/0028330 A1	2/2010	Collins et al.
8,685,980 B2	4/2014	Besong et al.	2010/0040614 A1 *	2/2010	Ahmed A61P 1/16 424/173.1
8,697,069 B2	4/2014	Kuchroo et al.	2010/0056576 A1	3/2010	Burger et al.
8,709,412 B2	4/2014	Jones et al.	2010/0061992 A1	3/2010	Anderson et al.
8,709,416 B2	4/2014	Langermann et al.	2010/0074900 A1	3/2010	Ghayur et al.
8,709,429 B2	4/2014	Majeti et al.	2010/0076178 A1	3/2010	Ghayur et al.
8,715,619 B2	5/2014	Karsunky	2010/0105667 A1	4/2010	Furet et al.
8,728,474 B2	5/2014	Honjo et al.	2010/0136009 A1	6/2010	Papkoff et al.
8,735,551 B2	5/2014	Garner et al.	2010/0203056 A1	8/2010	Irving et al.
8,735,553 B1	5/2014	Li et al.	2010/0233079 A1	9/2010	Jakob et al.
8,779,105 B2	7/2014	Korman et al.	2010/0233183 A1	9/2010	Triebel et al.
8,779,108 B2	7/2014	Queva et al.	2010/0260668 A1	10/2010	Ghayur et al.
8,841,418 B2	9/2014	Karsunky et al.	2010/0266531 A1	10/2010	Hsieh et al.
8,900,587 B2	12/2014	Carven et al.	2010/0266617 A1	10/2010	Carven et al.
8,927,697 B2	1/2015	Davis et al.	2011/0008766 A1	1/2011	Ghayur et al.
8,952,136 B2	2/2015	Carven et al.	2011/0044894 A1	2/2011	Karsunky
9,045,545 B1	6/2015	Clube	2011/0044980 A1	2/2011	Ghayur et al.
9,067,999 B1	6/2015	Honjo et al.	2011/0054151 A1	3/2011	Lazar et al.
9,073,994 B2	7/2015	Honjo et al.	2011/0059106 A1	3/2011	Kuchroo et al.
9,084,776 B2	7/2015	Korman et al.	2011/0081341 A1	4/2011	Honjo et al.
9,102,727 B2	8/2015	Freeman et al.	2011/0091372 A1	4/2011	Ghayur et al.
9,103,832 B2	8/2015	Takayanagi et al.	2011/0091463 A1	4/2011	Ghayur et al.
9,109,034 B1	8/2015	Clube	2011/0097337 A1 *	4/2011	Preville A61P 37/02 424/153.1
9,132,281 B2	9/2015	Zeng et al.	2011/0110852 A1	5/2011	Miller et al.
9,175,082 B2	11/2015	Zhou et al.	2011/0123550 A1	5/2011	Shibayama et al.
9,238,646 B2	1/2016	Cheng et al.	2011/0136781 A1	6/2011	Zhuo et al.
9,333,256 B2	5/2016	Kuchroo et al.	2011/0150892 A1 *	6/2011	Thudium C07K 16/2803 435/69.6
9,358,289 B2	6/2016	Korman et al.	2011/0171220 A1	7/2011	Davis
9,387,247 B2	7/2016	Korman et al.	2011/0177088 A1	7/2011	Olive et al.
9,409,970 B2	8/2016	Mikesell et al.	2011/0195068 A1	8/2011	Langermann et al.
9,457,080 B2	10/2016	Freeman et al.	2011/0209230 A1	8/2011	Korman et al.
9,492,539 B2	11/2016	Korman et al.	2011/0212094 A1	9/2011	Ghayur et al.
9,492,540 B2	11/2016	Korman et al.	2011/0229461 A1	9/2011	Tyson
9,505,839 B2	11/2016	Lonberg et al.	2011/0236375 A1	9/2011	Lazar et al.
9,605,070 B2	3/2017	Sabatos-Peyton et al.	2011/0237573 A1	9/2011	Cheng et al.
9,683,048 B2	6/2017	Freeman et al.	2011/0263827 A1	10/2011	Ghayur et al.
9,815,898 B2	11/2017	Freeman et al.	2011/0271358 A1	11/2011	Freeman et al.
9,815,901 B2	11/2017	Brogdon et al.	2011/0280800 A1	11/2011	Wu et al.
9,834,605 B2	12/2017	Carven et al.	2011/0280877 A1	11/2011	Tamada
			2011/0287009 A1	11/2011	Scheer et al.

(56)

References Cited

U.S. PATENT DOCUMENTS

2011/0300183 A1* 12/2011 Sitkovsky A61P 31/00
435/375

2012/0014957 A1 1/2012 Ghayur et al.
2012/0034160 A1 2/2012 Ghayur et al.
2012/0039870 A9 2/2012 Dolk et al.
2012/0039906 A1 2/2012 Olive
2012/0070450 A1 3/2012 Ishikawa et al.
2012/0076805 A1 3/2012 Sharpe et al.
2012/0087858 A1 4/2012 Ghayur et al.
2012/0100131 A1 4/2012 Takayanagi et al.
2012/0107234 A1 5/2012 Pedersen et al.
2012/0114649 A1 5/2012 Langermann et al.
2012/0141501 A1 6/2012 Yoshida et al.
2012/0195900 A1 8/2012 Ghayur et al.
2012/0201746 A1 8/2012 Liu et al.
2012/0201824 A1 8/2012 Wasik
2012/0258108 A1 10/2012 Ghayur et al.
2012/0263722 A1 10/2012 Ghayur et al.
2013/0005216 A1 1/2013 Rittberger
2013/0017199 A1 1/2013 Langermann
2013/0022623 A1 1/2013 Karsunky et al.
2013/0095098 A1 4/2013 Tyson
2013/0109843 A1 5/2013 Carven et al.
2013/0133091 A1 5/2013 Korman et al.
2013/0156774 A1 6/2013 Kuchroo et al.
2013/0183688 A1 7/2013 Kuchroo et al.
2013/0202623 A1 8/2013 Chomont et al.
2013/0230514 A1 9/2013 Langermann et al.
2013/0310375 A1 11/2013 Cheng et al.
2013/0324515 A1 12/2013 Zhuo et al.
2014/0044728 A1 2/2014 Takayanagi et al.
2014/0044738 A1 2/2014 Langermann et al.
2014/0065142 A1 3/2014 Roschke et al.
2014/0093511 A1* 4/2014 Lonberg A61K 45/06
435/69.6

2014/0155678 A1 6/2014 Zeng et al.
2014/0178370 A1 6/2014 Freeman et al.
2014/0212422 A1 7/2014 Korman et al.
2014/0234320 A1 8/2014 Croft et al.
2014/0242094 A1 8/2014 Kuchroo et al.
2014/0274788 A1 9/2014 Ishikawa et al.
2014/0294852 A1 10/2014 Korman et al.
2014/0314714 A1 10/2014 Honjo et al.
2014/0328833 A1 11/2014 Korman et al.
2014/0341902 A1 11/2014 Maecker et al.
2014/0341978 A1 11/2014 Kim et al.
2014/0348743 A1 11/2014 Korman et al.
2015/0017185 A1 1/2015 Akbar et al.
2015/0023986 A1 1/2015 Jones et al.
2015/0079109 A1 3/2015 Li et al.
2015/0086574 A1 3/2015 Karsunky et al.
2015/0093380 A1 4/2015 Honjo et al.
2015/0125386 A1 5/2015 Hansen et al.
2015/0210769 A1 7/2015 Freeman et al.
2015/0218274 A1 8/2015 Sabatos-Peyton et al.
2015/0232555 A1 8/2015 Carven et al.
2015/0259420 A1 9/2015 Triebel et al.
2015/0290316 A1 10/2015 Graziano et al.
2015/0307609 A1 10/2015 Lonberg et al.
2016/0002334 A1 1/2016 Kuchroo et al.
2016/0068601 A1 3/2016 Brogdon et al.
2016/0082014 A1 3/2016 Cheng et al.
2016/0108123 A1 4/2016 Freeman et al.
2016/0122707 A1* 5/2016 Swee A61K 47/68
435/372

2016/0159905 A1 6/2016 Abdiche et al.
2016/0222116 A1* 8/2016 Korman C07K 16/2818
2016/0222121 A1 8/2016 Johnson et al.
2016/0326178 A1 11/2016 Zhuo et al.
2017/0044259 A1 2/2017 Tipton et al.
2017/0088615 A1 3/2017 Korman et al.
2017/0137514 A1 5/2017 Lonberg et al.
2017/0190777 A1 7/2017 Sabatos-Peyton et al.
2017/0198041 A1 7/2017 Sabatos-Peyton et al.
2017/0209574 A1 7/2017 Cao et al.

2017/0210804 A1 7/2017 Friebel et al.
2017/0247456 A1 8/2017 Freeman et al.
2017/0281624 A1 10/2017 Peters et al.
2017/0296659 A1 10/2017 Lebwohl et al.
2017/0304443 A1 10/2017 Lebwohl et al.
2017/0340733 A1 11/2017 Cao
2018/0000832 A1 1/2018 Cheng et al.
2018/0066054 A1 3/2018 Thudium et al.
2018/0086830 A1 3/2018 Triebel et al.
2018/0155427 A1 6/2018 Freeman et al.
2018/0186882 A1 7/2018 Freeman et al.
2018/0207273 A1 7/2018 Dranoff et al.
2018/0222982 A1 8/2018 Dranoff et al.
2018/0282340 A1 10/2018 Zhuo et al.
2018/0340025 A1 11/2018 Dranoff et al.
2018/0371093 A1 12/2018 Bilic et al.
2019/0365746 A1 12/2019 Dobson et al.
2020/0030442 A1 1/2020 Cao
2020/0172617 A1 6/2020 Stein et al.
2020/0223917 A1 7/2020 Sabatos-Peyton et al.
2020/0223924 A1 7/2020 Stein et al.
2020/0277378 A1 9/2020 Sabatos-Peyton et al.
2020/0308277 A1 10/2020 Sabatos-Peyton et al.
2020/0339689 A1 10/2020 Freeman et al.
2020/0369762 A1 11/2020 Bruederle et al.
2020/0377600 A1 12/2020 Johnson et al.
2021/0000951 A1 1/2021 Cao et al.
2021/0009687 A1 1/2021 Triebel et al.
2021/0284737 A1 9/2021 Freeman et al.
2022/0133889 A1 5/2022 Dranoff et al.
2022/0153835 A1 5/2022 Dranoff et al.
2022/0185883 A1 6/2022 Sabatos-Peyton et al.
2022/0306737 A1 9/2022 Eliasson et al.
2023/0013364 A1 1/2023 Dranoff et al.

FOREIGN PATENT DOCUMENTS

CN 103079644 A 5/2013
CN 103242448 A 8/2013
EP 0670369 A2 9/1995
EP 0742795 A1 11/1996
EP 1165616 A1 1/2002
EP 1210424 A1 6/2002
EP 1334659 A1 8/2003
EP 1385533 A1 2/2004
EP 1445264 A1 8/2004
EP 1537878 A1 6/2005
EP 1576014 A1 9/2005
EP 1591527 A1 11/2005
EP 1870399 A1 12/2007
EP 1896582 A1 3/2008
EP 2161336 A1 3/2010
EP 2195347 A1 6/2010
EP 2206517 A1 7/2010
EP 2243493 A1 10/2010
EP 2270051 A2 1/2011
EP 2307050 A1 4/2011
EP 2328920 A2 6/2011
EP 2342228 A1 7/2011
EP 2342229 A1 7/2011
EP 2360254 A1 8/2011
EP 2370593 A2 10/2011
EP 2397155 A1 12/2011
EP 2397156 A1 12/2011
EP 2412825 A1 2/2012
EP 2417984 A1 2/2012
EP 2418278 A2 2/2012
EP 2439272 A2 4/2012
EP 2439273 A2 4/2012
EP 2482849 A2 8/2012
EP 2504364 A1 10/2012
EP 2099447 B1 11/2012
EP 2535354 A1 12/2012
EP 2545076 A1 1/2013
EP 2545078 A1 1/2013
EP 2051990 B1 2/2013
EP 2581113 A1 4/2013
EP 2170959 B1 10/2013
EP 2691112 A1 2/2014

(56)

References Cited

FOREIGN PATENT DOCUMENTS

EP	2699264	A1	2/2014	WO	2008060617	A2	5/2008
EP	2723381	A2	4/2014	WO	2008064157	A1	5/2008
EP	2320940	B1	3/2015	WO	2008071447	A2	6/2008
EP	2905030	A1	8/2015	WO	2008073687	A2	6/2008
EP	2344474	B1	9/2015	WO	2008083174	A2	7/2008
EP	2927240	A1	10/2015	WO	2008085562	A2	7/2008
EP	2474545	B1	11/2016	WO	2008156712	A1	12/2008
EP	3222634	A1	9/2017	WO	2009014708	A2	1/2009
JP	H07291996	A	11/1995	WO	2009024531	A1	2/2009
JP	2002194491	A	7/2002	WO	2009029342	A2	3/2009
JP	2003029846	A	1/2003	WO	2009091547	A1	7/2009
JP	2004512005	A	4/2004	WO	2009097394	A2	8/2009
JP	2006521783	A	9/2006	WO	2009101611	A1	8/2009
JP	2006340714	A	12/2006	WO	2009114335	A2	9/2009
JP	2010514791	A	6/2010	WO	2009120905	A2	10/2009
JP	2012503984	A	2/2012	WO	2009141386	A1	11/2009
RU	2494107	C2	9/2013	WO	2010001617	A1	1/2010
WO	8808135	A1	10/1988	WO	2010019570	A2	2/2010
WO	9520605	A1	8/1995	WO	2010026124	A1	3/2010
WO	1996027603	A1	9/1996	WO	2010027423	A2	3/2010
WO	9707671	A1	3/1997	WO	2010027827	A2	3/2010
WO	0032231	A1	6/2000	WO	2010027828	A2	3/2010
WO	0058363	A1	10/2000	WO	2010029082	A1	3/2010
WO	0071078	A2	11/2000	WO	2010029434	A1	3/2010
WO	0073498	A1	12/2000	WO	2010029435	A1	3/2010
WO	0114424	A2	3/2001	WO	2010036959	A2	4/2010
WO	0114556	A1	3/2001	WO	2010051502	A2	5/2010
WO	0114557	A1	3/2001	WO	2010063011	A2	6/2010
WO	0139722	A2	6/2001	WO	2010077634	A1	7/2010
WO	01077342	A1	10/2001	WO	2010084999	A1	7/2010
WO	01083750	A2	11/2001	WO	2010089411	A2	8/2010
WO	200194413	A2	12/2001	WO	2010098788	A2	9/2010
WO	0200692	A2	1/2002	WO	2010101849	A1	9/2010
WO	0200730	A2	1/2002	WO	2010102278	A1	9/2010
WO	2002/22577	A2	3/2002	WO	2010110346	A1	9/2010
WO	0224891	A2	3/2002	WO	2010117057	A1	10/2010
WO	0232378	A2	4/2002	WO	2011005481	A1	1/2011
WO	0234205	A2	5/2002	WO	2011011027	A1	1/2011
WO	0239813	A1	5/2002	WO	2011025927	A1	3/2011
WO	02078731	A1	10/2002	WO	2011034605	A2	3/2011
WO	02079499	A1	10/2002	WO	2011041613	A2	4/2011
WO	02086083	A2	10/2002	WO	2011066342	A2	6/2011
WO	03000066	A1	1/2003	WO	2011066389	A1	6/2011
WO	03002722	A2	1/2003	WO	2011069104	A2	6/2011
WO	03011911	A1	2/2003	WO	2011076786	A1	6/2011
WO	03033644	A2	4/2003	WO	2011100841	A1	8/2011
WO	03042402	A2	5/2003	WO	201110604	A1	9/2011
WO	03063792	A2	8/2003	WO	201110621	A1	9/2011
WO	03077914	A1	9/2003	WO	2011131472	A1	10/2011
WO	03099196	A2	12/2003	WO	2011155607	A1	12/2011
WO	2004004771	A1	1/2004	WO	2011159877	A2	12/2011
WO	2004005281	A1	1/2004	WO	2012018538	A2	2/2012
WO	2004007679	A2	1/2004	WO	2012022814	A1	2/2012
WO	2004056875	A1	7/2004	WO	2012064733	A2	5/2012
WO	2004072286	A1	8/2004	WO	2012079000	A1	6/2012
WO	2004078928	A2	9/2004	WO	2012106587	A1	8/2012
WO	2005027854	A2	3/2005	WO	2012135408	A1	10/2012
WO	2005033144	A2	4/2005	WO	2012145493	A1	10/2012
WO	2005097211	A2	10/2005	WO	2012177624	A2	12/2012
WO	2006004988	A2	1/2006	WO	2012177788	A1	12/2012
WO	2006021955	A2	3/2006	WO	2013006490	A2	1/2013
WO	2006042237	A2	4/2006	WO	2013006727	A1	1/2013
WO	2006121168	A1	11/2006	WO	2013019906	A1	2/2013
WO	2006124269	A2	11/2006	WO	2013043647	A1	3/2013
WO	2006133396	A2	12/2006	WO	2013066761	A1	5/2013
WO	2007005874	A2	1/2007	WO	2013079174	A1	6/2013
WO	2007011968	A2	1/2007	WO	2013079945	A1	6/2013
WO	2007024705	A2	3/2007	WO	2013169693	A1	11/2013
WO	2007024715	A2	3/2007	WO	2013173223	A1	11/2013
WO	2007070514	A1	6/2007	WO	2013181452	A1	12/2013
WO	2007082154	A2	7/2007	WO	2013181634	A2	12/2013
WO	2007084786	A1	7/2007	WO	2013184757	A1	12/2013
WO	2007113648	A2	10/2007	WO	2014008218	A1	1/2014
WO	2007146968	A2	12/2007	WO	2014018632	A1	1/2014
WO	2008016893	A1	2/2008	WO	2014022138	A2	2/2014
				WO	2014022332	A1	2/2014
				WO	2014022758	A1	2/2014
				WO	2014047350	A1	3/2014
				WO	2014055648	A1	4/2014

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO 2014055897 A2 4/2014
 WO 2014072493 A1 5/2014
 WO 2014085318 A1 6/2014
 WO 2014100079 A1 6/2014
 WO 2014165082 A2 10/2014
 WO 2014165422 A1 10/2014
 WO 2014179664 A2 11/2014
 WO 2014189805 A1 11/2014
 WO 2014195852 A1 12/2014
 WO 2014204791 A1 12/2014
 WO 2015009856 A2 1/2015
 WO 2015026634 A1 2/2015
 WO 2015026684 A1 2/2015
 WO 2015035606 A1 3/2015
 WO 2015036394 A1 3/2015
 WO 2015036499 A1 3/2015
 WO 2015036511 A1 3/2015
 WO 2015042246 A1 3/2015
 WO 2015048312 A1 4/2015
 WO 2015048520 A1 4/2015
 WO 2015061668 A1 4/2015
 WO 2015070060 A1 5/2015
 WO 2015075725 A1 5/2015
 WO 2015081158 A1 6/2015
 WO 2015085847 A1 6/2015
 WO 2015088847 A1 6/2015
 WO 2015095423 A2 6/2015
 WO 2015095811 A2 6/2015
 WO 2015103602 A1 7/2015
 WO 2015109124 A2 7/2015
 WO 2015109391 A1 7/2015
 WO 2015112800 A1 7/2015
 WO 2015112805 A1 7/2015
 WO 2015112900 A1 7/2015
 WO 2015117002 A1 8/2015
 WO 2015118175 A2 8/2015
 WO 2015119944 A1 8/2015
 WO 2015120198 A1 8/2015
 WO 2015134605 A1 9/2015
 WO 2015138920 A1 9/2015
 WO 2015176033 A1 11/2015
 WO 2015181342 A1 12/2015
 WO 2015195163 A1 12/2015
 WO 2016000619 A1 1/2016
 WO 2016028672 A1 2/2016
 WO 2016040880 A1 3/2016
 WO 2016040882 A1 3/2016
 WO 2016040892 A1 3/2016
 WO 2016054555 A2 4/2016
 WO 2016061142 A1 4/2016
 WO 2016069727 A1 5/2016
 WO 2016079049 A1 5/2016
 WO 2016100882 A1 6/2016
 WO 2016161239 A1 10/2016
 WO 2016161270 A1 10/2016
 WO 2016168716 A1 10/2016
 WO 2017017623 A1 2/2017
 WO 2017017624 A1 2/2017
 WO 2017019894 A1 2/2017
 WO 2017019896 A1 2/2017
 WO 2017019897 A1 2/2017
 WO 2017034916 A1 3/2017
 WO 2017097407 A1 6/2017
 WO 2017106656 A1 6/2017
 WO 2017189433 A1 11/2017
 WO 2018150312 A1 8/2018
 WO 2019006007 A1 1/2019
 WO 2019018640 A1 1/2019
 WO 2019018730 A1 1/2019
 WO 2019099838 A1 5/2019
 WO 2019180576 A1 9/2019
 WO 2019200229 A1 10/2019
 WO 2020128636 A1 6/2020
 WO 2021053490 A1 3/2021
 WO 2021079188 A1 4/2021

WO 2021079195 A1 4/2021
 WO 2021123902 A1 6/2021
 WO 2021144657 A1 7/2021
 WO 2022195551 A1 9/2022

OTHER PUBLICATIONS

Rosenberg et al. (2008) Adoptive cell transfer: a clinical path to effective cancer immunotherapy. *Nat Rev Cancer* 8(4): 299-308.*
 Till et al. (2012) CD20-specific adoptive immunotherapy for lymphoma using a chimeric antigen receptor with both CD28 and 4-1BB domains: pilot clinical trial results. *Blood* 119(17): 3940-3950.*
 Woo et al. (2011) Immune inhibitory molecules LAG-3 and PD-1 synergistically regulate T-cell function to promote tumoral immune escape. *Cancer Res* 72: 917-927.*
 Zhou et al. (2009) Depletion of Endogenous Tumor-Associated Regulatory T Cells Improves the Efficacy of Adoptive Cytotoxic T-Cell Immunotherapy in Murine Acute Myeloid Leukemia. *Blood* 114: 3793-3802.*
 Fourcade et al., "Upregulation of Tim-3 and PD-1 expression is associated with tumor antigen-specific CD8+ T cell dysfunction in melanoma patients" *The Journal of Experimental Medicine* (2010) vol. 207 No 10 pp. 2175-2186.
 Franklin, M.C., et al., "Insights into ErbB signaling from the structure of the ErbB2-pertuzumab complex," *Cancer Cell* 5 (4):317-328, Cell Press, United States (Apr. 2004).
 Freeman et al., "Cloning of B7-2: a CTLA-4 counter-receptor that costimulates human T cell proliferation," *Science* (1999) vol. 262, pp. 909-911.
 Freeman et al., "Engagement of the PD-1 immunoinhibitory receptor by a novel B7 family member leads to negative regulation of lymphocyte activation", *J Exp Med.* (2000) vol. 192, No. 7, pp. 1027-1034.
 Freeman et al., "TIM genes: a family of cell surface phosphatidylserine receptors that regulate innate and adaptive immunity" *Immunol Rev* (2010) vol. 235 No. 1 pp. 172-189.
 Fujimori et al. "A Modeling Analysis of Monoclonal Antibody Percolation Through Tumors: A Binding-Site Barrier" *J Mucl Med* (1990) vol. 31, pp. 1191-1198.
 Fukushima et al., "Antibodies to T-cell Ig and mucin domain-containing proteins (Tim)-1 and -3 suppress the induction and progression of murine allergic conjunctivitis" *Biochemical and Biophysical Research Communications* (2006) vol. 353 No. 1 p. 211-16.
 Fulda, "Molecular Pathways: Targeting Inhibitor of Apoptosis Proteins in Cancer—From Molecular Mechanism to Therapeutic Application," *Clin Cancer Res* (2013) vol. 20, No. 2, pp. 289-295.
 Gao et al., "TIM-3 Expression Characterizes Regulatory T Cells in Tumor Tissues and Is Associated with Lung Cancer Progression" *PLoS ONE* (2012) vol. 7 No. 2 e30676.
 Garcia et al: "The Pan PIM Kinase Inhibitor LGH447 Shows Activity In PIM2 Dependent Multiple Myeloma and In AML Models", *Blood* (2013) Retrieved from the Internet: URL:<http://www.bloodjournal.org/content/122/21/1666> [retrieved on Apr. 14, 2016].
 Garrison K et al: "The small molecule TGF-[beta] signaling inhibitor SM16 synergizes with agonistic OX40 antibody to suppress established mammary tumors and reduce spontaneous metastasis" *Cancer Immunology, Immunotherapy* (2012) vol. 61 No. 4 pp. 511-521.
 Ge, X., et al., "CD134 Allodepletion Allows Selective Elimination of Alloreactive Human T Cells without Loss of Virus-Specific and Leukemia-Specific Effectors," *Biology of Blood and Marrow Transplantation* 14(5):518-530, American Society for Blood and Marrow Transplantation, United States (2008).
 Geng et al., "Soluble Form of T Cell Ig Mucin 3 Is an Inhibitory Molecule in T Cell-Mediated Immune Response" *The Journal of Immunology* (2006) vol. 176 pp. 1411-1420.
 Gettinger et al. "Safety and Response 1-98 With Nivolumab (Anti-PD-1; BMS-936558, ONO-1538) Plus Erlotinib in Patients (Pts) With Epidermal Growth Factor Receptor Mutant (EGFR MT) Advanced Non-Small Cell Lung Cancer (NSCLC) Metastatic Non-

(56)

References Cited

OTHER PUBLICATIONS

Small Cell Lung Cancer” *International Journal of Radiation: Oncology Biology Physics* (2014) vol. 90, No. 5, pp. S34-S35.

Ghebeh et al., “Doxorubicin downregulates cell surface B7-H1 expression and upregulates its nuclear expression in breast cancer cells: role of B7-H1 as an anti-apoptotic molecule,” *Breast Cancer Research* (2010) vol. 12, No. 4, Article R48, 12 pages.

Golden-Mason et al., “Negative Immune Regulator Tim-3 Is Overexpressed on T Cells in Hepatitis C Virus Infection and Its Blockade Rescues Dysfunctional CD4+ and CD8+ T Cells,” *J Virol* (2009) vol. 83, No. 18, pp. 9122-9130.

Greenspan, N.S., “Epitopes, paratopes and other topes: do immunologists know what they are talking about?” *Bulletin de l’Institut Pasteur* 90(4):267-279, Elsevier, France (1992).

Grygielewicz Paulina et al: “Epithelial-mesenchymal transition confers resistance to selective FGFR inhibitors in SNU-16 gastric cancer cells”. *Gastric Cancer*. Springer Japan. Tokyo, vol. 19. No. 1., Nov. 19, 2014 (Nov. 19, 2014). pp. 53-62.

Haitov, “Immunology: Structure and Functions of the Immune System,” Geotar-Media Publishing Group, Moscow (2013) p. 110. Russian.

Hallett et al., “Immunosuppressive Effects of Multiple Myeloma Are Overcome by PD-L1 Blockade” *Biol Blood Marrow Transplant* (2011) vol. 17, No. 8, pp. 1133-1145.

Hamid et al., “Safety and Tumor Responses with Lembroizumab (Anti-PD-1) in Melanoma” *New England Journal of Medicine* (2013) vol. 369 No. 2 pp. 134-144.

Hansen, J.A., et al., “Monoclonal Antibodies Identifying a Novel T-Cell Antigen and Ia Antigens of Human Lymphocytes,” *Immunogenetics* 10:247-260, Springer-Verlag (1980).

Harlow, E. and Lane, D., “Using Antibodies: A Laboratory Manual,” Cold Spring Harbor Laboratory Press, Cold Spring Harbor, United States various pages (1999).

Hastings et al., “TIM-3 is Expressed on Activated Human CD4+ T Cells and Regulates Th1 and Th17 Cytokines” *Eur J Immunol* (2009) vol. 39 No. 9 pp. 2492-2501.

He, Y-F., et al., “Blocking programmed death-1 ligand-PD-1 interactions by local gene therapy results in enhancement of antitumor effect of secondary lymphoid tissue chemokine,” *The Journal of Immunology* 173(8):4919-4928, The American Association of Immunologists, United States (2004).

Henry et al., “Structure and evolution of the extended B7 family,” *Immunol Today* (1999) vol. 20, No. 6, pp. 285-288.

Herbst et al., “A study of MPDL3280A, an engineered PD-L1 antibody in patients with locally advanced or metastatic tumors,” *J Clin Oncol* (2013) vol. 31, No. 15 (Supp), Abstract 3000.

Hirano, F., et al., “Blockade of B7-H1 and PD-1 by monoclonal antibodies potentiates cancer therapeutic immunity,” *Cancer Research* 65(3):1089-1096, American Association for Cancer Research, United States (2005).

Hofstra et al., “Prevention of Th2-like cell responses by coadministration of IL-12 and IL-18 is associated with inhibition of antigen-induced airway hyperresponsiveness, eosinophilia, and serum IgE levels.” *Journal of Immunology* (1998) vol. 161 No. 9 pp. 5054-5060.

Hogenesch et al. “Challenges in pre-clinical testing of anti-cancer drugs in cell culture and in animal models” *Journal of Controlled Release* (2012) vol. 164, No. 2, pp. 183-186.

Hu Yi et al: “Essential role of AKT in tumor cells addicted to FGFR.” *Anti-Cancer Drugs*, vol. 25, No. 2, Feb. 2014 (Feb. 2014), pp. 183-188.

Huang et al., “Lymphoma endothelium preferentially expresses Tim-3 and facilitates the progression of lymphoma by mediating immune evasion” *The Journal of Experimental Medicine* (2010) vol. 207 No. 3 pp. 505-520.

Huang et al., “Role of LAG-3 in Regulatory T Cells,” *Immunity* (2004) vol. 21, pp. 503-513.

Huang, Z., “Structural chemistry and therapeutic intervention of protein-protein interactions in immune response, human immuno-

deficiency virus entry, and apoptosis,” *Pharmacology & Therapeutics* 86(3):201-215, Pergamon Press, England (2000). cited by applicant.

Huard et al., “Lymphocyte-activation gene 3/major histocompatibility complex class II interaction modulates the antigenic response of CD4+ T lymphocytes,” *Eur J Immunol* (1994) vol. 24, pp. 3216-3221.

Huard et al., “T cell major histocompatibility complex class II molecules down-regulate CD4+ T cell clone responses following LAG-3 binding,” *Eur J Immunol* (1996) vol. 26, pp. 1180-1186.

Hutloff, A., et al., “ICOS is an inducible T-cell co-stimulator structurally and functionally related to CD28,” *Nature* 397 (6716):263-266, Nature Publishing Group, England (1999).

International Search Report and Written Opinion for International Application No. PCT/US2015/049826 dated Dec. 16, 2015.

International Search Report and Written Opinion for International Application No. PCT/US2015/055390, dated Dec. 17, 2015.

International Search Report and Written Opinion for International Application No. PCT/US2015/066812 dated Mar. 23, 2016.

International Search Report and Written Opinion for International Application No. PCT/US2016/044545 dated Oct. 28, 2016.

International Search Report and Written Opinion for International Application No. PCT/US2016/044547 dated Oct. 18, 2016.

International Search Report and Written Opinion for International Application No. PCT/US2016/044549 dated Oct. 14, 2016.

International Search Report and Written Opinion for PCT/US2014/057491 dated Jan. 7, 2015.

International Search Report and Written Opinion for PCT/US2015/012754 dated May 20, 2015.

International Search Report and Written Opinion for PCT/US2015/013913 dated May 4, 2015.

International Search Report and Written Opinion for PCT/US2015/049826 dated Dec. 16, 2015.

International Search Report and Written Opinion for PCT/US2015/053799 dated May 17, 2016.

International Search Report and Written Opinion issued in International Application No. PCT/US2016/067200, dated Apr. 10, 2017.

International Search Report and Written Opinion issued in International Application No. PCT/US2018/039825, dated Oct. 4, 2018.

Sakuishi et al., “TIM3+FOXP3+ regulatory T cells are tissue-specific promoters of T-cell dysfunction in cancer” *Oncotarget* (2013) vol. 2 No. 4 pp. e23849-1-e23849-9.

Salama, A.D., et al., “Critical role of the programmed death-1 (PD-1) pathway in regulation of experimental autoimmune encephalomyelitis,” *The Journal of Experimental Medicine* 198(1):71-78, The Rockefeller University Press, United States (2003).

Sanmamed et al. “Agonists of Co-stimulation in Cancer Immunotherapy Directed Against CD137, OX40, GITR, CD27, CD28, and ICOS” *Seminars in Oncology* (2015) vol. 42, No. 4, pp. 640-655.

Santiago et al., “Structures of T Cell Immunoglobulin Mucin Receptors 1 and 2 Reveal Mechanisms for Regulation of Immune Responses by the TIM Receptor Family” *Immunity* (2007) vol. 26 pp. 299-310.

Schroll, A. et al., “Tim3 Is Upregulated and Protective in Nephrotoxic Serum Nephritis”, *The American Journal of Pathology*, vol. 176, No. 4, pp. 1716-1742, Apr. 2010.

Search Report and Written Opinion issued in Singapore Application No. 11201605627T dated Aug. 15, 2017.

Search Report and Written Opinion issued in Singapore Application No. 11201702401R, completed Mar. 29, 2018.

Shakhov et al., “SMUCKLER/TIM4 is a distinct member of TIM family expressed by stromal cells of secondary lymphoid tissues and associated with lymphotoxin signaling” *Eur. J Immunol* (2004) vol. 34 pp. 494-503.

Sher et al. “Regulation of Immunity to Parasites by T cells and T Cell-derived Cytokines” *Annual Review Immunol* (1992) vol. 10, pp. 385-409.

Shinohara, T., et al., “Structure and chromosomal localization of the human PD-1 gene (PDCD1),” *Genomics* 23 (3):704-706, Academic Press, United States (1994).

(56)

References Cited

OTHER PUBLICATIONS

- Sho, M., et al., "Possibility of Clinical Applications for Novel Cancer Immunotherapy via Inhibition of T-cell Negative Pathway," *Magazine of the Japan Society of Clinical Oncology* 40(2):590 (PS22-5) (2005).
- Simmons et al., "Tim-3+ T-bet+ Tumor-Specific Th1 Cells Colocalize with and Inhibit Development and Growth of Murine Neoplasms" *The Journal of Immunology* (2005) vol. 174 pp. 1405-1415.
- Soares et al. "Recombinant Human Tumor Antigen MUC1 Expressed in Insect Cells: Structure and Immunogenicity" *Protein Expression and Purification* (2001) vol. 22, pp. 92-100.
- Soh, E.Y., et al., "Neutralizing vascular endothelial growth factor activity inhibits thyroid cancer growth in vivo," *Surgery* 128(6):1059-1066, Mosby, United States (2000).
- Song et al.: "3681 Phenotypic and Functional Effects of Novel HDAC Inhibitor LBH589 On Human Lymphocyte Populations", 51st ASH Annual Meeting and Exposition (2009) Retrieved from the Internet: URL: <https://jash.confex.com/jash/2889/webprogram/Paper22684.html> [retrieved on Apr. 14, 2016].
- Song W et al: "HDAC inhibition by LBH589 affects the phenotype and function of human myeloid dendritic cells.", *LEUKEMIA* JAN 2811, vol. 25, No. 1, Jan. 2011 (Jan. 2011), pp. 161-168.
- Stewart et al., "MEDI4736: Delivering effective blockade of immunosuppression to enhance tumour rejection: Monoclonal antibody discovery and practical development," *Cancer Res* (2011) vol. 71, No. 8 (Supp), Abstract LB-158.
- Strome et al., "A Mechanistic Perspective of Monoclonal Antibodies in Cancer Therapy Beyond Target-Related Effects," *The Oncologist* (2007) vol. 12, pp. 1084-1095.
- Supplementary Partial European Search Report for European Application No. EP 1484888, dated Mar. 1, 2017. 10 pages.
- Swallow et al., "B7h, a Novel Costimulatory Homolog of B7.1 and B7.2, Is Induced by TNF α ," *Immunity* (1999) vol. 11, pp. 423-432.
- Sánchez-Fueyo et al., "Tim-3 inhibits T helper type 1-mediated auto- and alloimmune responses and promotes immunological tolerance," *Nat Immunol* (2003) vol. 4, No. 11, pp. 1093-1101.
- Takamura et al., "Premature Terminal Exhaustion of Friend Virus-Specific Effector CD8+ T Cells by Rapid Induction of Multiple Inhibitory Receptors" *J Immunol* (2010) vol. 184 pp. 4696-4707.
- Talmadge et al. "Murine Models to Evaluate Novel and Conventional Therapeutic Strategies for Cancer" *The American Journal of Pathology* (2007) vol. 170, No. 3, pp. 793-804.
- Tamura, H., et al., "B7-H1 costimulation preferentially enhances CD28-independent T-helper cell function," *Blood* 97 (6):1809-1816, The American Society of Hematology, United States (2001).
- Tang et al. "Immunotherapy and tumor microenvironment" *Cancer Letters* (2016) vol. 370, pp. 85-90.
- Teeling, J.L., et al., "Characterization of new human CD20 monoclonal antibodies with potent cytolytic activity against non-Hodgkin Lymphomas," *Blood* 104(6):1793-1800, American Society of Hematology, United States (2004).
- Thangamathesvaran et al., "Immune checkpoint inhibitors and radiotherapy—concept and review of current literature," *Ann Transl Med* (2018) vol. 6, No. 8, Article 155, 11 pages.
- Thomas et al. "Combined Effects of RU486 and Tamoxifen on the Growth and Cell Cycle Phases of the MCF-7 Cell Line" *Journal of Clinical Endocrinology and Metabolism* (1992) vol. 75, Issue 3, pp. 865-870.
- Thomas et al. "Effects of Gossypol on the Cell Cycle Phases in T-47D Human Breast Cancer Cells" *Anticancer Research* (1991) vol. 11, No. 4, pp. 1469-1476.
- Thomas, M.L., "Of ITAMs and ITIMs: turning on and off the B cell antigen receptor," *The Journal of Experimental Medicine* 181(6):1953-1956, The Rockefeller University Press, United States (1995).
- Thompson et al., "Tumor B7-H1 Is Associated with Poor Prognosis in Renal Cell Carcinoma Patients with Long-term Follow-up" *Cancer Res.* (2006) vol. 66, pp. 3381-3385.
- Thompson, R. Houston, et al.; "PD-1 is expressed by tumor-infiltrating immune cells and is associated with poor outcome for patients with renal cell carcinoma"; *Clin. Cancer Res* 13(6):1757-1761 (2007).
- Thurber et al. "Antibody tumor penetration: Transport opposed by systemic and antigen-mediated clearance" *Advanced Drug Delivery Reviews* (2008) vol. 60, pp. 1421-1434.
- Tomlinson, I.M., et al., "The repertoire of human germline VH sequences reveals about fifty groups of VH segments with different hypervariable loops," *Journal of Molecular Biology* 227(3):776-798, Elsevier, Netherlands (1992).
- Topalian et al., "Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer," *N Engl J Med* (2012) vol. 366, No. 26, pp. 2443-2454.
- Topalian, S., "Q&A: Suzanne Topalian on Immune Therapies", *Cancer Discovery* 3(7):712, American Association for Cancer Research, United States, published online Jun. 27, 2013.
- Topalian, S., et al., "Nivolumab (anti-PD-1; BMS-936558; ONO-4538) in patients with advanced solid tumors: Survival and long-term safety in a phase I trial," accessed at <http://meetinglibrary.asco.org/content/113543-132>, accessed on Jun. 1, 2016, 2pages.
- Topalian, S.L., et al., "Safety, Activity, and Immune Correlates of Anti-PD-1 Antibody in Cancer," *The New England Journal of Medicine* 366(26):2443-2454, Massachusetts Medical Society, United States (2012).
- Trautmann, Lydie, et al.; "Upregulation of PD-1 expression on HIV-specific CD8+ T cells leads to reversible immune dysfunction"; *Nat. Med.*; 12(10):1198-1202 (2006).
- Triebel et al., "LAG-3, A Novel Lymphocyte Activation Gene Closely Related to CD4," *J Exp Med* (1990) vol. 171, pp. 1393-1405.
- Triebel, "LAG-3: a regulator of T-cell and DC responses and its use in therapeutic vaccination," *Trends Immunol* (2003) vol. 24, No. 12, pp. 619-622.
- Tsai, C.-J., et al., "Protein Allosteric, signal transmission and dynamics: a classification scheme of allosteric mechanisms," *Molecular BioSystems* 5(3):207-216, Royal Society of Chemistry, England (2009).
- Tsushima, Fumihiko, et al.; "Predominant expression of B7-H1 and its immunoregulatory roles in oral squamous cell carcinoma"; *Oral Oncology*; 42:268-274 (2006).
- Tuskan et al., "Real-time PCR analysis of candidate imprinted genes on mouse chromosome 11 shows balanced expression from the maternal and paternal chromosomes and strain-specific variation in expression levels" *Epigenetics* (2008) vol. 3 No. 1 pp. 43-50.
- U.S. Appl. No. 15/536,718, filed Jun. 16, 2017.
- UniProtKB/Swiss-Prot Database entry, PDCD1.sub.-HUMAN, accessed at <http://www.uniprot.org/uniprot/Q15116.txt>, accessed on Jun. 1, 2016, 5 pages.
- Van De Weyer et al. "A highly conserved tyrosine of Tim-3 is phosphorylated upon stimulation by its ligand galectin-9" *Biochemical and Biophysical Research Communications* (2006) vol. 351, pp. 571-576.
- Van Regenmortel, M.H.V., "The recognition of Proteins and Peptides by Antibodies," *Journal of Immunoassay* 21 (2-3):85-108, Taylor & Francis, England (2000).
- Vanneman et al: "Combining immunotherapy and targeted therapies in cancer treatment" *Nature Reviews Cancer* (2012) vol. 12 No. 4 pp. 237-251.
- Verbrugge et al: "The curative outcome of radioimmunotherapy in a mouse breast cancer model relies on mTOR signaling", *Radiation Research. Radiation Research Society, GB*, (2014) vol. 182 No 2 pp. 219-229.
- Nishimura, H., et al., "Autoimmune dilated cardiomyopathy Science in PD-1 receptor-deficient mice," *Science* 291 (5502):1319-322, American Association for the Advancement of Science, United States (2001).
- Nishimura, H., et al., "Development of lupus-like autoimmune diseases by disruption of the PD-Igene encoding an ITIM motif-carrying immunoreceptor," *Immunity* 11(2):141-151, Cell Press, United States (1999).
- Nishimura, H., et al., "Immunological studies on PD-1 deficient mice: implication of PD-1 as a negative regulator for B cell

(56)

References Cited

OTHER PUBLICATIONS

responses," *International Immunology* 10(10):1563-1572, Oxford University Press, England (1998).

Nomi, T., et al., "Clinical significance and therapeutic potential of the programmed death-1 ligand/programmed death-1 pathway in human pancreatic cancer," *Clinical Cancer Research* 13(7):2151-2157, The Association, United States (2007). cited by applicant.

Ohigashi et al, "Clinical Significance of Programmed Death-1 Ligand-1 and Programmed Death-1 Ligand-2 Expression in Human Esophageal Cancer" *Clin Cancer Research* (2005) vol. 11, pp. 2947-2953.

Ohno et al., "Antigen-binding specificities of antibodies are primarily determined by seven residues of VH," *PNAS* (1985) vol. 82, pp. 2945-2949.

Okamoto et al., "T-Helper Type 1/T-Helper Type 2 Balance in Malignant Pleural Effusions Compared to Tuberculous Pleural Effusions" *Chest* (2005) vol. 128 pp. 4030-4035.

Okazaki et al., "PD-1 and PD-1 ligands: from discovery to clinical application" *Intern. Immun.* (2007) vol. 19, pp. 813-824.

Okazaki, T., et al., "New regulatory co-receptors: inducible co-stimulator and PD-1," *Current Opinion in Immunology* 14 (6):779-782, Elsevier, England (2002).

Okazaki, T., et al., "PD-1 immunoreceptor inhibits B cell receptor-mediated signaling by recruiting src homology 2-domain-containing tyrosine phosphatase 2 to phosphotyrosine," *Proceeding of the National Academy of Sciences* 98 (24):13866-13871, National Academy of Sciences, United States (2001).

Oki Y et al: "Immune regulatory effects of panobinostat in patients with Hodgkin lymphoma through modulation of serum cytokine levels and T-cell PD1 expression," *Blood Cancer Journal*, vol. 4, E236, 2014, pp. 1-4.

Okudaira et al., "A modified version of galectin-9 suppresses cell growth and induces apoptosis of human T-cell leukemia virus type 1-infected T-cell lines" *Int. J. Cancer* (2007) vol. 120 pp. 2251-2261.

OPDIVO (nivolumab) Dmg Label, Initial U.S. Approval: 2014, Revised May 2016.

OPDIVO (nivolumab) Drug Label, Initial U.S. Approval: 2014, Revised Nov. 2016.

Opposition filed by ASILFA AG in corresponding Chilean Application No. 2017-00888 on Oct. 24, 2017, assigned litigation file number by the National Institute of Industrial Property of Chile (INAPI) on Feb. 27, 2018, notified by INAPI to agent on Mar. 2, 2018.

Opposition filed by Laboratorios Legrand S.A. in corresponding Colombian Application No. NC2017/0003490 on Dec. 7, 2017, admitted Dec. 19, 2017, published Dec. 20, 2017.

Opposition filed in Colombian Application No. NC2016/0001001, filed Feb. 15, 2017.

Ozaki, S., et al., "Immunotherapy of Multiple Myeloma with a Monoclonal Antibody Directed Against a Plasma Cell-specific Antigen, HM1.24," *Blood* 90(8):3179-3186, American Society of Hematology, United States (1997).

Ozkaynak, E., et al., "Programmed death-1 targeting can promote allograft survival," *The Journal of Immunology* 169 (11):6546-6553, The American Association of Immunologists, United States (2002).

Pal et al., "Programmed Death-1 Inhibition in Renal Cell Carcinoma: Clinical Insights and Future Directions," *Clinical Adv Hem One* (2014) vol. 12, Issue 2, pp. 90-99.

Panka, et al.; "Variable region framework differences result in decreased or increased affinity of variant anti-digoxin antibodies"; *Proc. Natl. Acad. Sci. USA*; 85:3080-3084 (1988).

Pardoll et al. "The blockade of immune checkpoints in cancer immunotherapy" *Nature Reviews Cancer* (2012) vol. 12, pp. 252-264.

Park, J.W. and Smolen, J., "Monoclonal antibody therapy," *Advances in Protein Chemistry* 56:369-421, Academic Press, United States (2001).

Parry, Richard V., et al.; "CTLA-4 and PD-1 Receptors Inhibit T-Cell Activation by Distinct Mechanisms"; *Molecular and Cellular Biology*; 25(21):9543-9553 (2005).

Patel et al., "Taming dendritic cells with TIM-3: another immunosuppressive strategy used by tumors" *Immunotherapy* (2012) vol. 4 No. 12 pp. 1795-1798.

Paterson et al., "The Programmed Death-1 Ligand 1:B7-1 Pathway Restrains Diabetogenic Effector T Cells In Vivo," *J Immunol* (2011) vol. 187, pp. 1097-1105.

Patsoukis, N., et al., "PD-1 Increases PTEN Phosphatase Activity While Decreasing PTEN Protein Stability by Inhibiting Casein Kinase 2," *Molecular and Cellular Biology* 33(16):3091-3098, American Society for Microbiology, United States (Aug. 2013).

Patsoukis, N., et al., "PD-1 inhibits T cell proliferation by upregulating p. 27 and p. 15 and suppressing Cdc25A," *Cell Cycle* 11(23):1-5, Landes Bioscience, United States (Dec. 2012).

Patsoukis, N., et al., "Selective Effects of PD-1 on Akt and Ras Pathways Regulate Molecular Components of the Cell Cycle and Inhibit T Cell Proliferation," *Science Signaling* 5(230): ra46, pp. 1-14, American Association for the Advancement of Science, United States (Jun. 2012).

Perez-Gracia et al., "Orchestrating immune check-point blockade for cancer immunotherapy in combinations", *Current Opinion in Immunology* (2014) vol. 27 pp. 89-97.

Phong et al., "Tim-3 enhances Fc[epsilon]RI-proximal signaling to modulate mast cell activation," *J Experimental Medicine* (2015) vol. 212, No. 13, pp. 2289-2304.

Pinzon-Ortiz et al: "S710: The combination of JAK inhibitor, ruxolitinib, pan-PIM inhibitor, LGH447, and CDK4/6 inhibitor, LEE011, in a preclinical mouse model of myeloproliferative neoplasia", *Haematologica, The Hematology Journal: Official Organ of the European Hematology Association*, vol. 99. No. Supp 1 (2014) p. 252.

Polyak, M.J. and Deans, J.P., "Alanine-170 and proline-172 are critical determinants for extracellular CD20 epitopes; heterogeneity in the fine specificity of CD20 monoclonal antibodies is defined by additional requirements imposed by both amino acid sequence and quaternary structure," *Blood* 99(9):3256-3262, American Society of Hematology, United States (2002).

Post Grant Opposition filed in Colombian Patent Application No. NC2016/0001001, dated Jul. 31, 2018.

Postel-Vinay et al. "Challenges of phase 1 clinical trials evaluating immune checkpoint-targeted antibodies" *Annals of Oncology* (2016) vol. 27, pp. 214-224.

Powles et al., "MPDL3280A (anti-PD-L1) treatment leads to clinical activity in metastatic bladder cancer," *Nature* (2014) vol. 515, pp. 558-562.

Prigent et al., "Lymphocyte activation gene-3 induces tumor regression and antitumor immune responses," *Eur J Immunol* (1999) vol. 29, pp. 3867-3876.

Prokunina, L. and Alarcon-Riquelme, M., "The genetic basis of systemic lupus erythematosus-knowledge of today and thoughts for tomorrow," *Human Molecular Genetics* 13(1):R143-R148, Oxford University Press, England (2004).

Quintarelli et al.: "Selective strong synergism of Ruxolitinib and second generation tyrosine kinase inhibitors to overcome bone marrow stroma related drug resistance in chronic myelogenous leukemia", *Leukemia Research*, New York, NY, US, vol. 38, No. 2, Nov. 15, 2013 (Nov. 15, 2013), pp. 236-242.

Raziorrouh et al. "The Immunoregulatory Role of CD244 in Chronic Hepatitis B Infection and its Inhibitory Potential on Virus-Specific CD8+ T-cell Function" *Hepatology* (2010) vol. 52 pp. 1934-1947.

Richter et al., "On the role of the inhibitory receptor LAG-3 in acute and chronic LCMV infection," *Int Immunol* (2009) vol. 22, No. 1, pp. 13-23.

Riley, J.L. and June, C.H., "The CD28 family: a T-cell rheostat for therapeutic control of T-cell activation," *Blood* 105 (1):13-21, American Society of Hematology, United States (Jan. 2005).

Rothe et al. "Enhancing dendritic cell-induced T-cell responses by immunomodulating molecules" 13th CIMT Annual Meeting (2015) p. 74.

(56)

References Cited

OTHER PUBLICATIONS

- Rowe et al., "Innate IFN-gamma is essential for programmed death ligand-1-mediated T cell stimulation following *Listeria monocytogenes* infection," *J Immunol* (2012) vol. 189, No. 2, pp. 876-884.
- Rudikoff, et al.; "Single Amino Acid Substitution Altering Antigen-binding Specificity"; *Proc. Natl. Acad. Sci. USA*; 79:1979-1983 (1982).
- Rudnick et al. "Affinity and Avidity in Antibody-Based Tumor Targeting" *Cancer Biotherapy and Radiopharmaceuticals* (2009) vol. 24, No. 2, pp. 155-160.
- Sabatos et al. "Interaction of Tim-3 and Tim-3 ligand regulates T helper type 1 responses and induction of peripheral tolerance" *Nature Immunology* (2003) vol. 4, pp. 1102-1110.
- Sabatos-Peyton et al., "Blockade of Tim-3 binding to phosphatidylserine and CEACAM1 is a shared feature of anti-Tim-3 antibodies that have functional efficacy," *Oncoimmunology* (2018) vol. 7, No. 2, Article e1385690, 9 pages.
- Sakuishi et al., "Emerging Tim-3 functions in anti-microbial and tumor immunity" *Trends Immunol* (2011) vol. 32 No. 8 pp. 345-349.
- Sakuishi et al., "Targeting Tim-3 and PD-1 pathways to reverse T cell exhaustion and restore anti-tumor immunity," *J Experimental Medicine* (2010) vol. 207, No. 10, pp. 2187-2194.
- Carreno, B.M, and Collins, M., "The B7 family of ligands and its receptors: new pathways for costimulation and inhibition of immune responses," *Annual Review of Immunology* 20:29-53, Annual Reviews, United States (2002).
- Carreno, B.M , "BTLA: a new inhibitory receptor with a B7-like ligand," *TRENDS in Immunology* 24(10):524-527, Elsevier, England (2003).
- Carter, L.L. and Carreno, B.M., et al., "Cytotoxic T-lymphocyte antigen-4 and programmed death-1 function as negative regulators of lymphocyte activation," *Immunologic Research* 28(1):49-59, Humana Press, United States (2003).
- Carter, L.L., et al., "PD-1:PD-L inhibitory pathway affects both CD4(+) and CD8(+) T cells and is overcome by IL-2," *European Journal of Immunology* 32(3):634-643, WILEY-VCH Verlag GmbH, Germany (2002).
- Carvalho et al., "Doxorubicin: the good, the bad and the ugly effect," *Current Medicinal Chemistry* (2009) vol. 16, No. 25, pp. 3267-3285.
- Casati et al., "Soluble Human LAG-3 Molecule Amplifies the In vitro Generation of Type 1 Tumor-Specific Immunity," *Cancer Res* (2006) vol. 66, No. 8, pp. 4450-4460.
- Catherine Sabatos-Peyton, MBG453: A high affinity, ligand-blocking anti-TIM-3 monoclonal Ab. American Association for Cancer Research (AACR) Annual Meeting, Apr. 17, 2016, New Orleans, Louisiana.
- Ceeraz et al., "B7 family checkpoint regulators in immune regulation and disease," *Trends Immunol* (2013) vol. 34, No. 11, pp. 556-563.
- Cespedes "Mouse models in oncogenesis and cancer therapy" *Clin. Tranl. Oncol.* (2006) vol. 8, No. 5, pp. 318-329.
- Chan et al. "Therapeutic antibodies for autoimmunity and inflammation" *Nature Reviews Immunology* (2010) vol. 10, pp. 301-316.
- Chen, L., "Co-inhibitory molecules of the B7-CD28 family in the control of T-cell immunity," *Nature Reviews Immunology* 4(5):336-347, Nature Publishing Group, England (2004).
- Chen, Y., et al., "Selection and Analysis of an Optimized Anti-VEGF Antibody: Crystal Structure of an Affinity-matured Fab in Complex with Antigen," *Journal of Molecular Biology* 293(4):865-881, Academic Press, England (1999).
- Chervontseva A M et al: "Effect of cytarabine on expression of cell adhesion molecules and on endothelium-eukocyte interaction in vitro", *Terapevticheskii ARKHIV*2006, vol. 78, No. 7, 2006, pp. 67-72.
- Chesi et al., "IAP antagonists induce anti-tumor immunity in multiple myeloma," *Nat Med* (2016) vol. 22, No. 12, pp. 1411-1420.
- Christiansen et al: "Eradication of solid tumors using histone deacetylase inhibitors combined with immune-stimulating antibodies", *Proceedings of the National Academy of Sciences*, vol. 108 No 10, Feb. 22, 2011 (Feb. 22, 2011), pp. 4141-4146.
- Christiansson Lisa et al: "The tyrosine kinase inhibitors imatinib and dasatinib reduce myeloid suppressor cells and release effector lymphocyte responses.", *Molecular Cancer Therapeutics*, vol. 14, No. 5, May 2015 (May 2015), pp. 1181-1191.
- ClinicalTrials.gov Identifier: NCT01988896 "A Phase 1b Study of MPDL3280A (an Engineered Anti-PDL1 Antibody) in Combination With Cobimetinib in Patients With Locally Advanced or Metastatic Solid Tumors" *Clinicaltrials.gov*, last updated Dec. 1, 2014.
- ClinicalTrials.gov Identifier: NCT02323126 "Study of Efficacy and Safety of Nivolumab in Combination With EGF816 and of Nivolumab in Combination With INC280 in Patients With Previously Treated Non-small Cell Lung Cancer (EGF816)," *Clinicaltrials.gov*, last updated Jun. 6, 2018.
- ClinicalTrials.gov Identifier: NCT02040064 "Tolerability and Efficacy of Tremelimumab in Combination With Gefitinib in NSCLC Patients", *ClinicalTrials.gov*; last updated Jan. 17, 2014.
- ClinicalTrials.gov Identifier: NCT02263508 "A Phase 1 b/3, Multicenter, Open-label Trial of Tafimogene Laherparepvec in Combination With Pembrolizumab (MK-3475) for Treatment of Unresected, Stage IIIB to IVM1c Melanoma (MASTERKEY-265)", *ClinicalTrials.gov*; last updated Jun. 22, 2015.
- ClinicalTrials.gov Identifier: NCT02339571 "Randomized Phase II/III Study of Nivolumab Plus Ipilimumab Plus Sargramostim Versus Nivolumab Plus Ipilimumab in Patients With Unresectable Stage III or Stage IV Melanoma", *ClinicalTrials.gov*; last updated Apr. 9, 2015.
- ClinicalTrials.gov Identifier: NCT02608268, Safety and Efficacy of MBG453 as Single Agent and in Combination With PDR001 in Patients With Advanced Malignancies, Information provided by Novartis (Novartis Pharmaceuticals), last updated Oct. 13, 2016.
- ClinicalTrials.gov Identifier: NCT02817633, A Phase 1 Study of TSR-022, an Anti-TIM-3 Monoclonal Antibody, in Patients With Advanced Solid Tumors, Information provided by Tesaro, Inc., last updated Aug. 26, 2016.
- Cloekaert, A., et al., "O-Polysaccharide epitopic heterogeneity at the surface of *Brucella* spp. studied by enzyme-linked immunosorbent assay and flow cytometry," *Clinical and Diagnostic Laboratory Immunology* 5(6):862-870, American Society for Microbiology, United States (1998).
- Cohen et al., "Image Cytometry of Estrogen Receptors in Breast Carcinomas" *Cytometry* (1988) vol. 9 pp. 579-587.
- Collins et al., "The B7 family of immune-regulatory ligands" *Genome Biology* (2005) vol. 6 No. 223.
- Cragg, M.S. et al., "Complement-mediated lysis by anti-CD20 mAb correlated with segregation into lipid rafts," *Blood* 101(3):1045-1052, American Society of Hematology, United States (2003).
- Creelan, B.C., "Update on Immune Checkpoint Inhibitors in Lung Cancer," *Journal of the Moffitt Cancer Center* 21 (1):80-89, H. Lee Moffitt Cancer Center and Research Institute, United States (2014).
- Cruse, J.M. and Lewis, R.E., "Antigens and Immunogens," in *Atlas of Immunology*, 2nd ed., pp. 105-126, CRC Press, United States (Dec. 29, 2003).
- Davies, D.R. and Cohen, G. H., "Interactions of protein antigens with antibodies," *Proceedings of the National Academy of Sciences USA* 93(1):7-12, National Academy of Sciences, United States (1996).
- Davies, Julian, et al.; "Affinity improvement of single antibody VH domains: residues in all three hypervariable regions affect antigen binding"; *J Mol Biol*; 2:169-179 (1996).
- Dekruyff et al., "T Cell/Transmembrane, Ig, and Mucin-3 Allelic Variants Differentially Recognize Phosphatidylserine and Mediate Phagocytosis of Apoptotic Cells" *The Journal of Immunology* (2010) vol. 184 pp. 1918-1930.
- Del-Rio, Maria-Luisa, et al.; "Antibody-mediated signaling through PD-1 costimulates T cells and enhances CD28-dependent proliferation"; *Eur. J Immunol.*; 35(12):3545-3560 (2005).
- Demaria et al., "Immune-mediated inhibition of Metastases after Treatment with Local Radiation and CTLA-4 Blockade in a Mouse Model of Breast Cancer," *Clinical Cancer Research* (2005) vol. 11, pp. 728-734.
- Dennis "Off by a whisker" *Nature* (2006) vol. 442, pp. 739-741.

(56)

References Cited

OTHER PUBLICATIONS

- Dey et al: "Nutl in-3 inhibits the NF[kappa]B Pathway in a p53 Dependent Manner: Implications in Lung Cancer Therapy". *Cell Cycle*, vol. 6, No. 17, Sep. 1, 2007 (Sep. 1, 2007), pp. 2178-2185.
- Dong et al., "Tumor-associated B7-H1 promotes T-cell apoptosis: A potential mechanism of immune evasion", *Nat Med* (2002) vol. 8 pp. 793-800.
- Dong, H. and Chen, L., "B7-H1 pathway and its role in the evasion of tumor immunity," *Journal of Molecular Medicine* 81(5):281-287, Springer, Germany (2003).
- Dong, H., et al., "Tumor-associated B7-H1 promotes T-cell apoptosis: a potential mechanism of immune evasion," *Nature Medicine* 8(8):793-800, Nature Publishing Company, United States (2002).
- Dorfman et al., "The phosphatidylserine receptors, T cell immunoglobulin mucin proteins 3 and 4, are markers of histiocytic sarcoma and other histiocytic and decdritic cell neoplasms" *Hum Pathol* (2010) vol. 41 No. 10 pp. 1486-1494.
- Dougan et al., "Regulation of innate and adaptive antitumor immunity by IAP antagonists," *Immunotherapy* (2018) vol. 10, No. 9, pp. 787-796.
- Dougan, D.A., et al., "Effects of substitutions in the binding surface of an antibody on antigen affinity," *Protein Engineering* 11(1):165-74, Oxford University Press, England (1998).
- Drake et al., "Blocking the regulatory T cell molecule LAG-3 augments in vivo anti-tumor immunity in an autochthonous model of prostate cancer," *J Clin Oncol* (2006) vol. 24, No. 18S, Abstract 2573.
- Du Manoir et al., "Ki-67 Labeling in Postmitotic Cells Defines Different Ki-67 Pathways Within the 2c Compartment" *Cytometry* (1991) vol. 12 pp. 455-463.
- El Mir et al., "A Soluble Lymphocyte Activation Gene-3 Molecule Used as a Vaccine Adjuvant Elicits Greater Humoral and Cellular Immune Responses to Both Particulate and Soluble Antigens," *J Immunol* (2000) vol. 164, pp. 5583-5589.
- Entzminger et al., "De novo design of antibody complementarity determining regions binding a FLAG tetrapeptide," *Sci Rep* (2017), retrived from www.nature.com/articles/s41598-017-10737-9.pdf?origin=ppub, last accessed Jan. 12, 2018.
- Extended European Search Report for European Application No. EP 1484888, dated May 31, 2017.
- Finger, L.R., et al., "The human PD-1 gene: complete cDNA, genomic organization, and developmentally regulated expression in B cell progenitors," *Gene* 197(1-2):177-187, Elsevier, United States (1997).
- Fivash, M., et al., "BIAcore for macromolecular interaction," *Current Opinion in Biotechnology* 9(1):97-101, Current Biology, England (1998).
- Fleischer, Bernhard, et al.; "T cell stimulation by *Staphylococcal enterotoxins*"; *J. Exp. Medicine*; 167(5):1697-1707 (1988).
- International Search Report and Written Opinion issued in PCT/US2018/061534, dated Apr. 1, 2019, 9 pages.
- Koya et al., "BRAF Inhibitor Vemurafenib Improves the Antitumor Activity of Adoptive Cell Immunotherapy," *Cancer Res* (2012) vol. 72, No. 16, pp. 3928-3937.
- Liu et al., "The BRAF and MEK Inhibitors Dabrafenib and Trametinib: Effects on Immune Function and in Combination with Immunomodulatory Antibodies Targeting PD-1, PD-L1, and CTLA-4," *Clin Cancer Res* (2015) vol. 21, No. 7, pp. 1639-1651.
- Mach et al., "Phase (Ph) II Study of MBG453 + Spartalizumab in Patients (PTS) with Non-Small Cell Lung Cancer (NSCLC) and Melanoma Pretreated with Anti-PD-1/L1 Therapy," Poster Display Session 3 from ESMO Congress, Sep. 30, 2019, 3 pages.
- Najjar et al., "Myeloid-Derived Suppressor Cell Subset Accumulation in Renal Cell Carcinoma Parenchyma Is Associated with Intratumoral Expression of IL1beta, IL8, CXCL5, and Mip-1alpha," *Clinical Cancer Research* (2016) vol. 23, No. 9, pp. 2346-2355.
- Neubert et al., "T cell-induced CSF1 promotes melanoma resistance to PD1 blockade," *Science Translational Medicine* (2018) vol. 10, No. 436, Article eaan3311, 14 pages.
- Sachdev et al., "Phase 1/2a study of double immune suppression blockade by combining a CSF1R inhibitor (pexidartinib/PLX3397) with an anti-PD-1 antibody (pembrolizumab) to treat advanced melanoma and other solid tumors," *Gynecologic Oncology* (2016) vol. 141, pp. 147-148.
- Simeone et al., "Immunomodulating antibodies in the treatment of metastatic melanoma: The experience with anti-CTLA-4, anti-CD137, and anti-PD1," *Journal of Immunotoxicology* (2012) vol. 9, No. 3, pp. 241-247.
- Steele et al., "CXCR2 Inhibition Profoundly Suppresses Metastases and Augments Immunotherapy in Pancreatic Ductal Adenocarcinoma," *Cancel Cell* (2016) vol. 29, pp. 832-845.
- Stromnes et al., "Pancreatic Cancer: Planning Ahead for Metastatic Spread," *Cancel Cell* (2016) vol. 29, No. 6, pp. 774-776.
- Villalobos, "Primer organismo con ADN ailincial logra reproducirse," retrieved from www.fayerwayer.com/2014/05/primer-organismo-con-adn-artificial-logra-reproducirse (2014) last accessed Oct. 18, 2019.
- Zhang, "P2.01-097—Phase 3 Study of Pembrolizumab vs Platinum-Based Chemotherapy for PD-L1," (NSCLC ID 2182) Abstract from Poster Presentation from International Association for the Study of Lung Cancer, Aug. 9, 2015, retrieved from library.iaslc.org/search-speaker?search_speaker=30076.
- Zhu et al., "CSF1/CSF1R Blockade Reprograms Tumor-Infiltrating Macrophages and Improves Response to T-cell Checkpoint Immunotherapy in Pancreatic Cancer Models," *Cancer Research* (2014) vol. 74, No. 18, pp. 5057-5069.
- Miko et al., "Involvement of Galectin-9/TIM-3 Pathway in the Systemic Inflammatory Response in Early-Onset Preeclampsia," *PLOS One* (2013) vol. 8, No. 8, Article e71811, 9 pages.
- [No Author Listed] Press Release: "Bristol-Myers Squibb Announces Collaboration to Evaluate Opdivo (nivolumab) in Combination with Targeted Therapies from Novartis to Treat Non-Small Cell Lung Cancer (NSCLC)," dated Oct. 6, 2014.
- Abbas et al. "Functional diversity of helper T lymphocytes" *Nature* (1996) vol. 383, pp. 787-793.
- Abbas, A.K., et al., *Cellular and Molecular Immunobiology*, 2nd ed., pp. 8, 47-50, W.B. Saunders Company, United States (1991).
- Acquaviva et al.: "FGFR3 Translocations in Bladder Cancer: Differential Sensitivity to HSP90 Inhibition Based on Drug Metabolism". *Molecular Cancer Research*, vol. 12. no. 7. Jul. 1, 2014 (Jul. 1, 2014). pp. 1042-1054.
- Adams, G.P. and Weiner, L.M., "Monoclonal antibody therapy of cancer," *Nature Biotechnology* 23(9): 1147-1157, Nature Publishing Group, United States (2005).
- Agata, Y., et al., "Expression of the PD-1 antigen on the surface of stimulated mouse T and B lymphocytes," *International Immunology* 8(5):765-772, Oxford University Press, England (1996).
- Allard et al. "Targeting CD73 Enhances the Antitumor Activity of Anti-FD-1 and Anti-CTLA-4 mAbs" *Clinical Cancer Research* (2013) vol. 19, No. 20, pp. 5626-5635.
- Allison, J.P. and Krummel, M.F., "The Yin and Yang of T Cell Costimulation," *Science* 270(5238):932-933, American Association for the Advancement of Science, United States (1995).
- Almagro et al. "Humanization of Antibodies" *Frontiers in Bioscience* (2008) vol. 13, pp. 1619-1633.
- Amin et al: "Nivolumab (anti-PD-1; BMS-936558, ONO-4538) in combination with sunitinib or pazopanib in patients (pts) with metastatic renal cell carcinoma (mRCC)" *Journal of Clinical Oncology* (2014) vol. 32, No. 15 suppl, Abstract 5010.
- Anderson "TIM-3 as a therapeutic target in human inflammatory diseases" *Expert Opinion on Therapeutic Targets* (2007) vol. 11, issue 8, pp. 1005-1009.
- Anderson et al. "Promotion of Tissue Inflammation by the Immune Receptor Tim-3 Expressed on Innate Immune Cells" *Science* (2007) vol. 318, pp. 1141-1143.
- Anderson et al. "Tim-3, a negative regulator of anti-tumor immunity" *Current Opinion in Immunology* (2012) vol. 24, No. 2, pp. 213-216.
- Anderson, "Tim-3: An Emerging Target in the Cancer Immunotherapy Landscape" *Cancer Immunology Research* (2014) vol. 2 No. 5 pp. 393-398.

(56)

References Cited

OTHER PUBLICATIONS

- Anderson et al. "Lag-3, Tim-3, and TIGIT: Co-inhibitory Receptors with Specialized Functions in Immune Regulation" *Immunity* (2016) vol. 44, pp. 989-1004.
- Andre, E., et al., "Precise Characterization of the Epitope Recognized by a Monoclonal Antibody Against *Escherichia coli* RNA Polymerase," *Hybridoma* 24(1):1-5, Mary Ann Liebert, Inc., United States (Feb. 2005).
- Ansari, M.J., et al., "The programmed death-1 (PD-1) pathway regulates autoimmune diabetes in nonobese diabetic (NOD) mice," *The Journal of Experimental Medicine* 198(1):63-69, The Rockefeller University Press, United States (2003). cited by applicant.
- Ansell, S.M., et al., "PD-1 Blockade with Nivolumab in Relapsed or Refractory Hodgkin's Lymphoma," *The New England Journal of Medicine* 372(4):311-319, Massachusetts Medical Society, United States (Jan. 22, 2015).
- Armand, P., et al., "289 Nivolumab in Patients with relapsed or Refractory Hodgkin Lymphoma-Preliminary Safety, Efficacy and Biomarker Results of a Phase I Study," 56th ASH Annual Meeting and Exposition, Abstracts & Program, San Francisco, CA, Dec. 6-9, 2014.
- Ascierto et al. "Future perspectives in melanoma research" meeting report from the "Melanoma Bridge", Napoli, Dec. 5-8, 2013 *Journal of Translational Medicine* (2014) vol. 12, No. 277, pp. 1-29.
- Ashworth et al. "Management of a Patient With Advanced BRAF-Mutant Melanoma" *Journal of the National Comprehensive Cancer Network* (2014) vol. 12, No. 3, pp. 315-319.
- Aspeshlagh et al. "Rationale for anti-OX40 cancer immunotherapy" *European Journal of Cancer* (2016) vol. 52, pp. 50-66.
- Baixeras et al., "Characterization of the Lymphocyte Activation Gene 3-Encoded Protein. A New Ligand for Human Leukocyte Antigen Class II Antigens," *J Exp Med* (1992) vol. 176, pp. 327-337.
- Barber, Daniel L., et al.; "Restoring function in exhausted CD8 T cells during chronic viral infection"; *Nature* 439:682-687 (2006).
- Batus et al. "Optimal Management of Metastatic Melanoma: Current Strategies and Future Directions" *Am. J. Clin. Dermatol.* (2013) vol. 14, No. 3, pp. 179-194.
- Beckman et al. "Antibody Constructs in Cancer Therapy" *Cancer* (2007) vol. 109, No. 2, pp. 170-179.
- Bellucci et al.: "JAK1 and JAK2 Modulate Tumor Cell Susceptibility To Natural Killer (NK) Cells Through Regulation of PDL1 Expression", *Blood* (Nov. 15, 2013), Retrieved from the Internet: URL:<http://www.bloodjournal.org/content/122/21/3472.full.pdf> [retrieved on Apr. 14, 2016].
- Bennett, F., et al., "Program death-1 engagement upon TCR activation has distinct effects on costimulation and cytokine-driven proliferation: attenuation of ICOS, IL-4, and IL-21, but not CD28, IL-7, and IL-15 responses," *The Journal of Immunology* 170(2):711-718, The American Association of Immunologists, United States (2003).
- Benson et al. "The PD-1/PD-L1 axis modulates the natural killer cell versus multiple myeloma effect: a therapeutic target for CT-011, a novel monoclonal anti-PD-1 antibody" *Blood* (2010) vol. 116, No. 13, pp. 2286-2294.
- Berg, J.M., et al., "The Immune System," in *Biochemistry* 5th ed., pp. 921-950, W.H. Freeman and Company, United States (2002).
- Berger, R., et al., "Phase I safety and pharmacokinetic study of CT-011, a humanized antibody interacting with PD-1, in patients with advanced hematologic malignancies," *Clinical Cancer Research* 14(10):3044-3051, American Association for Cancer Research, United States (2008).
- Berrien-Elliott et al., "Durable Adoptive Immunotherapy for Leukemia Produced by Manipulation of Multiple Regulatory Pathways of CD8+ T-Cell Tolerance," *Cancer Research* (2012) vol. 73, pp. 605-616.
- Bigras, et al. "Spatial distribution of DNA ploidy in colorectal carcinoma" *Analytic Cellular Pathology* (1994) vol. 7, pp. 289-300.
- Blank et al "Combination of targeted therapy and immunotherapy in melanoma" *Cancer Immunol Immunother* (2011) vol. 60, pp. 1359-1371.
- Blank, C., et al., "Interaction of PD-L1 on tumor cells with PD-1 on tumor-specific T cells as a mechanism of immune evasion: implications for tumor immunotherapy," *Cancer Immunology Immunotherapy* 54(4):307-314, Springer-Verlag, Germany (2005).
- Blank, C., et al., "PD-L1/B7H-1 inhibits the effector phase of tumor rejection by T cell receptor (TCR) transgenic CD8+ T cells," *Cancer Research* 64(3):1140-1145, American Association for Cancer Research, United States (2004).
- Blank, Christian, et al.; "Contribution of the PD-L1/PD-1 pathway to T-cell exhaustion: an update on implications for chronic infections and tumor evasion"; *Cancer Immunol. Immunotherapy*; 56(5):739-745 (2007).
- Blazar, B.R., et al., "Infusion of anti-B7.1 (CD80) and anti-B7.2 (CD86) monoclonal antibodies inhibits murine graft-versus-host disease lethality in part via direct effects on CD4+ and CD8+ T cells," *The Journal of Immunology* 157 (8):3250-3259, The American Association of Immunologists, United States (1996).
- Brahmer et al., "Safety and Activity of Anti-PD-L1 Antibody in Patients with Advanced Cancer," *N Engl J Med* (2012) vol. 366, pp. 2455-2465.
- Brahmer, J.R., et al., "Safety and activity of MDX-1106 (ONO-4538) anti-PD-1 monoclonal antibody in patients with selected refractory or relapsed malignancies," *Journal of Clinical Oncology* 26:Abstract No. 3006, American Society of Clinical Oncology, United States (2008).
- Brahmer, J.R., et al., "Phase I study of single-agent anti-programmed death-1 (MDX-1106) in refractory solid tumors Safety, clinical activity, pharmacodynamics, and immunologic correlates," *Journal of Clinical Oncology* 28 (19):3167-3175, American Society of Clinical Oncology, United States (2005).
- Brand et al., "Prospect for Anti-HER2 Receptor Therapy in Breast Cancer," *Anticancer Research* (2006) vol. 26, pp. 463-470.
- Brown et al., "Blockade of Programmed Death-1 Ligands on Dendritic Cells Enhances T Cell Activation and Cytokine Production", *J Immunol.* (2003) vol. 170, pp. 1257-1266.
- Brown et al., "Tolerance to Single, but Not Multiple, Amino Acid Replacements in Antibody VH CDR2: A Means of Minimizing B Cell Wastage from Somatic Hypermutation?" *J Immunol* (1996) vol. 156, pp. 3285-3291.
- Brown, J.A., et al., "Expression and functional consequences PD-1 ligands on natural APCs and tumors," *The FASEB Journal* 15(4):A345 (abstract No. 275.23), Federation of American Societies for Experimental Biology, United States (2001). cited by applicant.
- Brunet et al., "A new member of the immunoglobulin superfamily CTLA-4," *Nature* (1987) vol. 328, pp. 267-270.
- Butte et al., "Interaction of human PD-L1 and B7-1", *Mol Immunol* (2008) vol. 45, pp. 3567-3572.
- Butte et al., "Programmed Death-1 Ligand 1 Interacts Specifically with the B7-1 Costimulatory Molecule to Inhibit T Cell Responses," *Immunity* (2007) vol. 27, pp. 111-122.
- Campbell, A.M., "Characterisation of monoclonal antibodies," in *Laboratory Techniques in Biochemistry and Molecular Biology, Monoclonal Antibody Technology: The Production and Characterization of Rodent and Human Hybridomas*, vol. 13, pp. 186-215, Elsevier, the Netherlands (1984).
- Cao et al. "Genetic variations and haplotypes in TIM-3 gene and the risk of gastric cancer" *Cancer Immunol Immunother* (2010) vol. 59 pp. 1851-1857.
- Extended European Search Report issued in European Patent Application No. 20174880.3, dated Nov. 6, 2020, 15 pages.
- Extended European Search Report issued in European Application No. 19206634.8, dated Sep. 10, 2020, 26 pages.
- Extended European Search Report issued in European Application No. 21166347.1 dated Nov. 9, 2021.
- Edwards et al., "The Remarkable Flexibility of the Human Antibody Repertoire; Isolation of Over One Thousand Different Antibodies to a Single Protein, BLYS," *J Mol Biol* (2003) vol. 334, pp. 103-118.
- Felip et al., "Efficacy and safety of capmatinib plus nivolumab in pretreated patients with EGFR wild-type non-small cell lung cancer," IASLC 2020 World Conference on Lung Cancer, Abstract No. 817, downloaded from https://cpaper.ctimeetingtech.com/wclc2020/submission/preview/print?publication_id=817.

(56)

References Cited

OTHER PUBLICATIONS

- Janku et al. "Abstract CT034: Phase I study of WNT974 + spartalizumab in patients (pts) with advanced solid tumors" *Cancer Research* (2020) vol. 80, No. 16, pp. 1-4.
- Liu et al., "Targeting Wnt-driven cancer through the inhibition of Porcupine by LGK974," *PNAS* (2013) vol. 110, No. 50, pp. 20224-20229.
- Marchalonis et al., "The antibody repertoire in evolution: Chance, selection, and continuity," *Developmental and Comparative Immunology* (2006) vol. 30, pp. 223-247.
- Spigel et al., "Phase 1/2 Study of the Safety and Tolerability of Nivolumab Plus Crizotinib for the First-Line Treatment of Anaplastic Lymphoma Kinase Translocation—Positive Advanced Non-Small Cell Lung Cancer (CheckMate 370)," *Journal of Thoracic Oncology* (2018) vol. 13, No. 5, pp. 682-688.
- International Search Report and Written Opinion issued in PCT/US2007/085100, dated Apr. 28, 2008, 11 pages.
- Iouzalen et al., "LAP, a lymphocyte activation gene-3 (LAG-3)-associated protein that binds to a repeated EP motif in the intracellular region of LAG-3, may participate in the down-regulation of the CD3/TCR activation pathway," *Eur J Immunol* (2001) vol. 31, pp. 2885-2891.
- Ishida et al., "Induced expression of PD-1, a novel member of the immunoglobulin gene superfamily, upon programmed cell death," *The EMBO Journal* (1992) vol. 11, No. 11, pp. 3887-3895.
- Ishima, R. and Torchia, D.A., "Protein Dynamics from NMR," *Nature Structural Biology* 7(9):740-743, Nature Publishing Company, United States (2000).
- Iwai et al., "Involvement of PD-L1 on tumor cells in the escape from host immune system and tumor immunotherapy by PD-L1 blockade" *PNAS* (2002) vol. 99, pp. 12293-12297.
- Iwai et al., "PD-1 blockade inhibits hematogenous spread of poorly immunogenic tumor cells by enhanced recruitment of effector T cells," *International Immunology* (2005) vol. 17, No. 2, pp. 133-144.
- Iwai, Y., et al., "Microanatomical localization of PD-1 in human tonsils," *Immunology Letters* 83(3):215-220, Elsevier, Netherlands (2002).
- Iwai, Y., et al., "PD-1 inhibits antiviral immunity at the effector phase in the liver," *The Journal of Experimental Medicine* 198(1):39-50, The Rockefeller University Press, United States (2003).
- Jan et al., "Prospective separation of normal and leukemic stem cells based on differential expression of TIM3, a human acute myeloid leukemia stem cell marker" *PNAS* (2011) vol. 108 No 12 pp. 5009-5014.
- Jason-Moller, L., et al., "Overview of Biacore Systems and Their Applications," *Current Protocols in Protein Science* S45:19.13.1-19.13.14, John Wiley & Sons, Inc., United States (2006).
- Jiang et al., "mTOR Kinase Inhibitor AZD8855 Enhances the Immunotherapeutic Activity of an Agonist CD40 Antibody in Cancer Treatment" *Cancer Research* (2011) vol. 71 No 12, pp. 4074-4084.
- Jiang X et al: "The activation of MAPK in melanoma cells resistant to BRAF inhibition promotes PD-L1 expression that is reversible by MEK and PI3K inhibition", *Clinical Cancer Research*, the American Association for Cancer Research, US, vol. 19, No. 3, Feb. 1, 2013 (Feb. 1, 2013). pp. 598-609.
- Jin et al. "Cooperation of Tim-3 and PD-1 in CD8 T-cell exhaustion during chronic viral infection" *PNAS* (2010) vol. 107, Issue 33, p. 14733-14738.
- Jing et al., "Combined immune checkpoint protein blockade and low dose whole body irradiation as immunotherapy for melanoma," *Journal for Immunotherapy of Cancer* (2015) vol. 3, No. 2, 15 pages.
- Johne, B., "Protocol: Epitope Mapping by Surface Plasmon Resonance in the BIAcore," *Molecular Biotechnology* 9 (1):65-71, Humana Press, United States (1998).
- Jones et al., "Tim-3 expression defines a novel population of dysfunctional T cells with highly elevated frequencies in progressive HIV-1 infection," *J Exp Med* (2008) vol. 205, No. 12, pp. 2763-2779.
- Ju et al., "T cell immunoglobulin-and mucin-domain-containing molecule-3 (Tim-3) mediates natural killer cell suppression in chronic hepatitis B" *Journal of Hepatology* (2010) vol. 52 No. 3 pp. 322-329.
- Kanai, T., et al., "Blockade of B7-H1 Suppresses the Development of Chronic Intestinal Inflammation," *The Journal of Immunology* 171(8):4156-4163, American Association of Immunologists, Inc., United States (2003).
- Karu et al., "Recombinant Antibody Technology," *ILAR Journal*, vol. 37, No. 3, pp. 132-141 (1995).
- Kasagi, S., et al., "Anti-programmed cell death 1 antibody reduces CD4+PD-1+ T cells and relieves the lupus-like Tephritis of NZB/W FI mice," *The Journal Immunology* 184(5):2337-2347, The American Association of Immunologists, United States (2010).
- Kaveri, S., "Epitope and idiotope mapping using monoclonal antibodies," *Methods in Molecular Biology* 51:171-181, Humana Press, United States (1995).
- Kearl et al., "PD-1/PD-L1 Blockade after Transient Lymphodepletion to Treat Myeloma," Presentation from Society for Immunotherapy of Cancer Conference, Oct. 27, 2012, North Bethesda, Maryland, 19 pages.
- Kearl et al., "Programmed Death Receptor-1/Programmed Death Receptor Ligand-1 Blockage after transient Lymphodepletion To Treat Myeloma," *J Immunol* (2013) vol. 190, pp. 5620-5628.
- Kearley et al., "Th-2 driven, allergen-induced airway inflammation is reduced after treatment with anti-Tim-3 antibody in vivo" *The Journal of Experimental Medicine* (2007) vol. 204 No. 6 pp. 1289-1294.
- KEYTRUDA (pembrolizumab) Drug Label, Initial U.S. Approval: 2014, Revised Aug. 2016.
- KEYTRUDA (pembrolizumab) Drug Label, Initial U.S. Approval: 2014, Revised Oct. 2016.
- Khaitov, *Immunologia*, Moscow, (2011) "GEOTAR-Media", p. 103.
- Khalil et al. "The New Era of Cancer Immunotherapy: Manipulating T-Cell Activity to Overcome Malignancy" *Immunotherapy of Cancer In: Advances in Cancer Research* (2015) vol. 128, pp. 1-68.
- Kier et al., "PD-1 and its ligands in tolerance and immunity" *Annu. Rev. Immunol.* (2008) vol. 26 pp. 677-704.
- Kikushige et al. "TIM-3 as a therapeutic target for malignant stem cells in acute myelogenous leukemia" *New York Academy of Sciences* (2012) vol. 1266, pp. 118-123.
- Kikushige et al., "TIM-3 Is a Promising Target to Selectively Kill Acute Myeloid Leukemia Stem Cells" *Cell Stem Cell* (2010) vol. 7 pp. 708-717.
- Kim et al: "Eradication of metastatic mouse cancers resistant to immune checkpoint blockade by suppression of myeloid-derived cells. (Includes Supporting Information)", *Proceedings of the National Academy of Sciences of the United States of America*, vol. 111, No. 32, Aug. 12, 2014 (Aug. 12, 2014), pp. 11774-11777.
- Kirkwood et al., "Immunotherapy of cancer in 2012" *CA: A Cancer Journal for Clinicians* (2012) vol. 62 No 5 pp. 309-335.
- Klein Jan M et al: "The histone deacetylase inhibitor LBH589 (panobinostat) modulates the crosstalk of lymphocytes with Hodgkin lymphoma cell lines", *PLOS ONE* (2013) vol. 8, No. 11, E79502, 2813, pp. 1-6.
- Klibi et al. "Blood diffusion and Th1-suppressive effects of galectin-9-containing exosomes released by Epstein-Barr virus-infected nasopharyngeal carcinoma cells" *Blood* (2009) vol. 113 No. 9 pp. 1957-1966.
- Knight et al. "Host immunity contributes to the antimelanoma activity of BRAF inhibitors" *The Journal of Clinical Investigation* (2013) vol. 123, No. 3, pp. 1371-1381.
- Knights et al., "Inhibitor of apoptosis protein (IAP) antagonists demonstrate divergent immunomodulatory properties in human immune subsets with implications for combination therapy" *Cancer Immunology and Immunotherapy* (2013) vol. 62 No. 2 pp. 321-335.
- Koga et al., "Blockade of the Interaction Between PD-1 and PD-L1 Accelerates Graft Arterial Disease in Cardiac Allografts," *Arterioscler Thromb Vase Biol* (2004) vol. 24, pp. 2057-2062.
- Konishi, J., et al., "B7-H1 expression on non-small cell lung cancer cells and its relationship with tumor-infiltrating lymphocytes and

(56)

References Cited

OTHER PUBLICATIONS

their PD-1 expression," *Clinical Cancer Research* 10(15):5094-5100, American Association for Cancer Research, United States (2004).

Korman et al, "Checkpoint Blockade in Cancer Immunotherapy" *Adv Immunol* (2006) vol. 90 pp. 297-339.

Kuchroo et al. "The TIM Gene Family: Emerging Roles in Immunity and Disease" *Nature Reviews Immunology* (2003) vol. 3, pp. 454-462.

Kuchroo et al., "B7-1 and B7-2 Costimulatory Molecules Activate Differentially the Th1/Th2 Developmental Pathways: Application to Autoimmune Disease Therapy" *Cell* (1995) vol. 80 No. 707-18.

Kwong et al., "Molecular Analysis of Tumor-Promoting CD8+ Cells in Two-Stage Cutaneous Chemical Carcinogenesis" *J Invest Dermatol* (2010) vol. 130 No. 6 pp. 1726-1736.

Lack et al. "Nebulized but not parenteral IFN-gamma decreases IgE production and normalizes airways function in a murine model of allergen sensitization" *Journal of Immunology* (1994) vol. 152, pp. 2546-2554.

Ladner, R.C., "Mapping the epitopes of Antibodies," *Biotechnology and Genetic Engineering Reviews* 24(1):1-30, Taylor & Francis, England (2007).

Laricchia, Robbio, L., et al., "Mapping of Monoclonal Antibody-and Receptor-Binding Domains on Human Granulocyte-Macrophage Colony-Stimulation Factor (rhGM-CSF) Using a Surface Plasmon Resonance-Based Biosensor," *Hybridoma* 15(5):343-350, Mary Ann Liebert, Inc. United States (1996).

Latchman et al., "PD-L2 is a second ligand for PD-1 and inhibits T cell activation," *Nature Immunol* (2001) vol. 2, No. 3, pp. 261-268.

Leach, D.R., et al., "Enhancement of antitumor immunity by CTLA-4 blockade," *Science* 271(5256):1734-1736, American Association for the Advancement of Science, United States (1996).

Lee et al. "The inhibition of the T-cell immunoglobulin and mucin domain 3 (Tim3) pathway enhances the efficacy of tumor vaccine" *Biochemical and Biophysical Research Communications* (2010) vol. 402, pp. 88-93.

Lehmann et al, "Identification of human triple-negative breast cancer subtypes and preclinical models for selection of targeted therapies", *J Clin Invest.*, 121(7): 2750-2767.

Vietta et al. "Considering Therapeutic Antibodies" *Science* (2006) vol. 313, pp. 308-309.

Vivier et al., "Immunoreceptor tyrosine-based inhibition motifs," *Immunol Today* (1997) vol. 18, No. 6, pp. 286-291.

Voskoglou-Nomikos et al. "Clinical Predictive Value of the in Vitro Cell Line, Human Xenograft, and Mouse Allograft Preclinical Cancer Models" *Clinical Cancer Research* (2003) vol. 9, pp. 4227-4239.

Waldmann, Thomas A.; "Effective cancer therapy through immunomodulation"; *Annual Rev.*; 57(1):65-81 (2006).

Walunas et al., "CTLA-4 can function as a negative regulator of T cell activation," *Immunity* (1994) vol. 1, No. 5, pp. 405-413.

Wang et al. "In Vitro Characterization of the Anti-PD-1 Antibody Nivolumab, BMS-936558, and In Vivo Toxicology in Non-Human Primates" *Cancer Immunology Research* (2014) vol. 2, No. 9, pp. 846-856.

Wang et al., "The Mdm2 inhibitor, NVP-CGM097, in combination with the BRAF inhibitor NVP-LGX818 elicits synergistic antitumor effects in melanoma" *Cancer Research* (2014), Abstract 5466, Retrieved from the Internet: URL: http://cancerres.aacrjournals.org/content/74/19_Supplement/5466 [retrieved on Apr. 14, 2016].

Wang et al., "VISTA, a novel mouse Ig superfamily ligand that negatively regulates T cell responses" *The Journal of Experimental Medicine* (2011) vol. 208 No. 3 pp. 577-592.

Wang et al.: "Abstract 2929: The Mdm2 inhibitor NVP-CGM097 enhances the anti-tumor activity of NVP-LDK378 in ALK mutant neuroblastoma models", *Cancer Research* (2014) Retrieved from the Internet: URL: http://cancerres.aacrjournals.org/content/74/19_Supplement/2929 [retrieved on Apr. 14, 2016].

Wang, "Interaction of TIM4-TIM1 Modulates the Function of CD4+CD25+Treg in Food Allergic Mice," Master's Thesis submitted at Zhengzhou University (2010) 65 pages, Chinese with English Abstract.

Watanabe et al., "Current approaches for the treatment of multiple myeloma," *Int J Hematol* (2013) vol. 97, pp. 333-344.

Weber, J.S., et al., "Safety, Efficacy, and Biomarkers of Nivolumab with Vaccine in Ipilimumab-Refractory or -Naive Melanoma," *Journal of Clinical Oncology* 31 (34):4311-4318, American Society of Clinical Oncology, United States (2013). cited by applicant.

Weisberg et al., "Smac mimetics: implications for enhancement of targeted therapies in leukemia: Treating leukemia with Smac mimetics," *Leukemia* (2010) vol. 24, No. 12, pp. 2100-2109.

Wiener et al., "TIM-3 Is Expressed in Melanoma Cells and Is Upregulated in TGF-Beta Stimulated Mast Cells" *Journal of Investigative Dermatology* (2007) vol. 127 pp. 906-914.

Wilson, I.A. and Stanfield, R.L., "Antibody-antigen interactions," *Current Opinion in Structural Biology* 3:113-118, Current Biology, United States (1993).

Winslow, R., "New Cancer Drugs Harness Power of Immune System", *The Wall Street Journal*, May 15, 2013, accessed at <http://www.wsj.com/articles/SB10001424127887323398204578485401089823868>, accessed on Jun. 1, 2016, 4 pages.

Wolchok et al., "Nivolumab plus Ipilimumab in Advanced Melanoma," *N Engl J Med* (2013) vol. 369, pp. 122-133.

Wolchok, J.D., et al., "Nivolumab plus Ipilimumab in Advanced Melanoma," *The New England Journal of Medicine* 369 (2):122-133, Massachusetts Medical Society, United States (2013).

Wong, R.M., et al., "Programmed death-1 blockade enhances expansion and functional capacity of human melanoma antigen-specific CTLs," *International Immunology* 19(10):1223-1234, Oxford University Press, England (2007).

Woo et al. "Immune Inhibitory Molecules LAG-3 and PD-1 Synergistically Regulate T-cell Function to Promote Tumoral Immune Escape" *Cancer Research* (2011) vol. 72, No. 4, pp. 917-927.

Woods David M et al: "HDAC Inhibition Upregulates PD-1 Ligands in Melanoma and Augments Immunotherapy with PD-1 Blockade," *Cancer Immunology Research*, vol. 3, No. 12, Dec. 2015 (Dec. 2015), pp. 1375-1385.

Woods David M et al: "The antimelanoma activity of the histone deacetylase inhibitor panobinostat (LBH589) is mediated by direct tumor cytotoxicity and increased tumor immunogenicity", *Melanoma Research*, vol. 23, No. 5, Oct. 2013 (Oct. 2013), pp. 341-348.

Woods et al: "Abstract 4090: Inhibition of class I histone deacetylases promotes robust and durable enhancement of PDL1 expression in melanoma: Rationale for combination therapy", *Cancer Research* (2014) Retrieved from the Internet: URL: http://cancerres.aacrjournals.org/content/74/19_Supplement/4090.short [retrieved on Apr. 14, 2016].

Workman et al., "Negative Regulation of T Cell Homeostasis by Lymphocyte Activation Gene-3 (CD223)," *J Immunol* (2005) vol. 174, pp. 688-695.

Wu et al., "Endothelial cell-expressed Tim-3 facilitates metastasis of melanoma cells by activating the NF-kB pathway" *Oncology Reports* (2010) vol. 24 pp. 693-699.

Wu et al., "Immunotherapies: The Blockade of Inhibitory Signals," *Int J Biol Sci* (2012) vol. 8, No. 10, pp. 1420-1430.

Wu, K-P, et al., "Structural Basis of a Flavivirus Recognized by Its Neutralizing Antibody: Solution Structure of the Domain III of the Japanese Encephalitis Virus Envelope Protein," *The Journal of Biological Chemistry* 278 (46):46007-46013, American Society for Biochemistry and Molecular Biology, Inc., United States (Nov. 2003).

Yamazaki, T., et al., "Expression of programmed death 1 ligands by murine T cells and APC," *The Journal of Immunology* 169(10):5538-5545, The American Association of Immunologists, United States (2002).

Yan et al., "Tim-3 Expression Defines Regulatory T Cells in Human Tumors" *PLoS ONE* (2013) vol. 8 No. 3 e58006.

Yang et al., "Lack of TIM-3 Immunoregulation in Multiple Sclerosis" *The Journal of Immunology* (2008) vol. 180 No. 7 pp. 4409-4414.

(56)

References Cited

OTHER PUBLICATIONS

- Yervoy (ipilimumab) Drug Label, Initial U.S. Approval: 2011, Revised Oct. 2015.
- Yi, J., et al., "Mapping the Epitope of an Inhibitory Monoclonal Antibody to the C-terminal DNA-binding Domain of HIV-1 Integrase," *The Journal of Biological Chemistry* 277(14):12164-12174, American Society for Biochemistry and Molecular Biology, Inc., United States (2002).
- Youngnak, Pompan, et al.; "Differential binding properties of B7-H1 and B7-DC to programmed death-1"; *Biochem. Biophys Res Commun.*; 307:672-677 (2003).
- Yuan Z et al., "Blockade of inhibitors of apoptosis (IAPs) in combination with tumor-targeted delivery of tumor necrosis factor-[alpha] leads to synergistic antitumor activity" *Cancer Gene Therapy* (2013) vol. 20 No. 1 pp. 46-56.
- Zamarin et al. "Immune checkpoint modulation: Rational design of combination strategies" *Pharmacology & Therapeutics* (2015) vol. 150, pp. 23-32.
- Zehavi-Willner, Tova, et al.; "The mitogenic activity of *Staphylococcal enterotoxin B* (Seb): a monovalent T cell mitogen that stimulates cytolytic T lymphocytes but cannot mediate their lytic interaction"; *J. Immunol.*; 137(8):2682-2687 (1986).
- Zhang et al., "Tim-3 regulates pro-and anti-inflammatory cytokine expression in human CD14+ monocytes" *Journal of Leucocyte Biology* (2012) vol. 91 pp. 189-196.
- Zhang, "Oral 31—PD1 Axis Inhibition," Abstract from Poster Presentation from International Association for the Study of Lung Cancer, Aug. 9, 2015, retrieved from library.iaslc.org/search-speaker?search_speaker=30076.
- Zhang, Xuewu, et al.; "Structural and Functional Analysis of the Costimulatory Receptor Programmed Death-1" *Immunity*; 20:337-347 (2004).
- Zhou et al., "Coexpression of Tim-3 and PD-1 identifies a CD8+ T-cell exhaustion phenotype in mice with disseminated acute myelogenous leukemia" *Blood* (2011) vol. 117 No. 17 pp. 4501-4510.
- Zhuang J et al: "Selective IAP inhibition results in sensitization of unstimulated but not CD40-stimulated chronic lymphocytic leukaemia cells to TRAIL-induced apoptosis" *Pharmacology Research & Perspectives* (2014) vol. 2 Issue 6, Article E00081, 14 pages.
- Zou, W. and Chen, L., "Inhibitory B7-family Molecules in the Tumour Microenvironment," *Nature Reviews Immunology* 8(6):467-477, Nature Publishing Group, England (2008).
- Zuberek, K., "The role of in vivo PD-1/PD-L1 interactions in syngeneic and allogeneic antitumor responses in murine tumor models," *Blood* 98(11):42B (2001).
- Zuberek, Krystyna, et al.; "In vitro and in vivo expression regulation of PD-1 and PD-L1 in murine tumor models" *Blood*; 98(11 Part 1):25a (2001).
- [No Author Listed] "Crean un ADN sintético capaz de evolucionar," retrieved from www.nationalgeographic.es/ciencia/crean-un-adn-sintetico-capaz-de-evolucionar, Apr. 25, 2012, last accessed Oct. 18, 2019.
- Borate et al., "Phase Ib Study of the Anti-TIM-3 Antibody MBG453 in Combination with Decitabine in Patients with High-Risk Myelodysplastic Syndrome (MDS) and Acute Myeloid Leukemia (AML)," *Blood* (2019) vol. 134, Supp. 1, Abstract 637, 5 pages. [ClinicalTrials.gov](https://clinicaltrials.gov) "History of Changes for Study NCT01454102," [ClinicalTrials.gov](https://clinicaltrials.gov), latest version submitted on Nov. 3, 2017.
- Curigliano et al., "Abstract CT183: Phase (Ph) I/II study of MBG453 +/- spartalizumab (PDR001) in patients (pts) with advanced malignancies," AACR Annual Meeting 2019, Cancer Research, vol. 79, Issue 13, Supplement, 2 pages.
- Extended European Search Report issued in European Application No. 19199437.5, dated Mar. 12, 2020, 8 pages.
- Highfill et al., "Disruption of CXCR2-Mediated MDSC Tumor Trafficking Enhances Anti-PD1 Efficacy," *Science Translational Medicine* (2014) vol. 6, Issue 237, Article 237ra67, 15 pages.
- Lenschow et al., "Expression and functional significance of an additional ligand for CTLA-2," *PNAS* (1993) vol. 90, pp. 11054-11058.
- Lenschow et al., "Long-term survival of xenogeneic pancreatic islet grafts induced by CTLA4lg," *Science* (1992) vol. 257, Issue 5071, pp. 789-792.
- Lesokhin, A.M., et al., "291 Preliminary Results of a Phase I Study of Nivolumab (BMS-936558) in Patients with Relapsed of Refractory Lymphoid Malignancies," 56th ASH Annual Meeting and Exposition, Abstracts & Program, San Francisco, CA, United States, Dec. 6-9, 2014.
- Lewis, D.E., et al., "Tumor Necrosis Factor-alpha, and CD80 Modulate CD28 Expression through a Similar Mechanism of T-cell Receptor-Independent of Transcription," *The Journal of Biological Chemistry* 279 (28):29130-29138, The American Society for Biochemistry and Molecular Biology, Inc., United States (2004).
- Li et al., "Contribution of PD-L1 to oncogenesis of lymphoma and its RNAi-based targeting therapy" *Leukemia & Lymphoma* (2012) vol. 53, No. 10, pp. 2015-2023.
- Li, L., et al., "A pathway regulated by cell cycle inhibitor p27Kip1 and checkpoint inhibitor Smad3 is involved in the induction of T cell tolerance," *Nature Immunology* 7(11):1157-1165, Nature Publishing, United States (2006).
- Li, L., et al., "CD4+CD25+ regulatory T-cell lines from human cord blood have functional and molecular properties and T-cell anergy," *Blood* 106(9):3068-3073, American Society of Hematology, United States (Nov. 2005).
- Li, L., et al., "IL-1 beta-Mediated Signals Preferentially Drive Conversion of Regulatory T Cells but Not Conventional T Cells into IL-17-Producing Cells," *The Journal of Immunology* 185(7):4148-4153, American Association of Immunologists, Inc., United States (2010).
- Li, L., et al., "Rap1-GTP is a Negative Regulator of Th Cell Function and Promotes the Generation of CD4+CD103+ Regulatory T Cells In Vivo," *The Journal of Immunology* 175(5):3133-3139, American Association of Immunologists, Inc., United States (Sep. 2005).
- Li, L., et al., "The cyclin dependent kinase inhibitor (R)-roscovitine prevents alloreactive T cell clonal expansion and protects against acute GvHD," *Cell Cycle* 8(11):1794-1802, Landes Bioscience, United States (2009).
- Liblau et al. "Th1 and Th2 CD4+ T cells in the pathogenesis of organ-specific autoimmune diseases" *Immunology Today* (1995) vol. 16, Issue 1, pp. 34-38.
- Lin, David Yin-Wei, et al.; "The PD-1/PD-L1 complex resembles the antigen-binding Fv domains of antibodies and T cell receptors"; *Proc. Natl. Acad. Sci. USA* 105(8):3011-3016 (2008).
- Linsley et al., "CTLA-4 Is a Second Receptor for the B Cell Activation Antigen B7," *J Exp Med* (1991) vol. 174, pp. 561-569.
- Linsley et al., "Immunosuppression in Vivo by a Soluble Form of the CTLA-4 T Cell Activation Molecule," *Science* (1992) vol. 257, pp. 792-795.
- Linsley et al., "Intracellular Trafficking of CTLA-4 and Focal Localization Towards Sites of TCR Engagement" *Immunity* (1996) vol. 4 pp. 535-543.
- List of clinical trials identified in [ClinicalTrials.gov](https://clinicaltrials.gov) relating to PDR001 as of Dec. 22, 2016.
- Liu et al., "A Novel Kinase Inhibitor, INC28060, Blocks c-MET-Dependent Signaling, Neoplastic Activities, and Cross-Talk with EGFR and HER-3," *Clin Cancer Res* (2011) vol. 17, No. 22, pp. 7127-7138.
- Lonberg, N., et al., "Antigen-specific human antibodies from mice comprising four distinct genetic modifications," *Nature* 368(6474):856-859, Nature Publishing Group, United States (1994).
- Loser et al., "IL-10 Controls Ultraviolet-Induced Carcinogenesis in Mice" *The Journal of Immunology* (2007) vol. 179 pp. 365-371.
- Lute, K.D., et al., "Human CTLA4 knock-in mice unravel the quantitative link between tumor immunity and autoimmunity induced by anti-CTLA-4 antibodies," *Blood* 106(9):3127-3133, American Society of Hematology, United States (2005).
- Mahoney et al. "The Next Immune-Checkpoint Inhibitors: PD-1/PD-L1 Blockade in Melanoma" *Clinical Therapeutics* (2015) vol. 37, No. 4, pp. 764-782.

(56)

References Cited

OTHER PUBLICATIONS

Maier et al., "PD-1: PD-L1 Interactions Contribute to the Functional Suppression of Virus-Specific CD8+ T Lymphocytes in the Liver," *J Immunol* (2007) vol. 178, pp. 2714-2720.

Makishi et al. "Retracted: A modified version of galectin-9 induces cell cycle arrest and apoptosis of Burkitt and Hodgkin lymphoma cells" *British Journal of Hematology* (2008) vol. 142 pp. 583-594.

Manning et al., "A model of multiple myeloma: culture of 5T33 murine myeloma cells and evaluation of tumorigenicity in the C57BL/KaLwRij mouse", *Br J Cancer.*, 66(6): 1088-1093 (1992).

Marri et al., "Human Biochemistry" Mir Publishing, Moscow (1993) vol. 1, p. 34. Russian.

Masters et al., "Abstract 5016: Antitumor activity of anti -PD-1 in combination with tyrosine kinase inhibitors in a preclinical renal cell carcinoma model" *AACR Annual Meeting* (2014) vol. 74, No. 5016.

May, K.F., Jr., et al., "Anti-human CTLA-4 monoclonal antibody promotes T-cell expansion and immunity in a hu-PBL-SCID model: a new method for preclinical screening of costimulatory monoclonal antibodies," *Blood* 105(3):1114-1120, American Society of Hematology, United States (2005).

Maçon-Lemaitre et al., "The negative regulatory function of the lymphocyte-activation gene-3 co-receptor (CD223) on human T cells," *Immunology* (2005) vol. 115, pp. 170-178.

Menzies et al. "Systemic treatment for BRAF-mutant melanoma: where do we go next?" *Lancet Oncology* (2014) vol. 15, pp. e371-e381.

Menzies et al., "Recent advances in melanoma systemic therapy. BRAF inhibitors, CTLA antibodies and beyond" *European Journal of Cancer* (2013) vol. 49 No. 15 pp. 3229-3241.

Miska et al., "Autoimmunity-mediated antitumor immunity: Tumor as an immunoprivileged self," *Eur J Immunol* (2012) vol. 42, pp. 2584-2596.

Mittendorf Elizabeth A et al: "PD-L1 expression in triple-negative breast cancer." *Cancer Immunology Research*, vol. 2. No. 4. Apr. 2014 (Apr. 2014). pp. 361-370.

Mocellin et al., "CTLA-4 blockade and the renaissance of cancer immunotherapy," *Biochim Biophys Acta* (2013) vol. 1836, pp. 187-196.

Mokyr et al., "Realization of the Therapeutic Potential of CTLA-4 Blockade in Low-Dose Chemotherapy-treated Tumor-bearing Mice", *Cancer Research*, 58:5301-5304 (1998).

Monney et al. "Th1-specific cell surface protein Tim-3 regulates macrophage activation and severity of an autoimmune disease" *Nature* (2002) vol. 415, pp. 536-541.

Moreira Da Silva, "Nivolumab Anti-PD-1 monoclonal antibody cancer immunotherapy" *Drugs of the Future* (2014) vol. 39 No. 1 pp. 15-24.

Mosmann et al. "The expanding universe of T-cell subsets: Th1, Th2 and more" *Immunology Today* (1996) vol. 17, Issue 3, pp. 138-146.

Mossman et al. "Two Types of Murine Helper T Cell Clone" *Journal of Immunology* (1986) vol. 136, Issue 7, pp. 2348-2357).

Mou et al., "Association Between TIM-1 Gene Polymorphisms and Allergic Rhinitis in a Han Chinese Population" *J Investig Allergol Clin Immunol* (2010) vol. 20 No. 1 pp. 3-8.

Murrey et al., *Biokhimiya cheloveka*, Mir (1993) vol. 1, p. 34.

Nagahara et al., "Galectin-9 Increases Tim-3+ Dendritic Cells and CF+ T Cells and Enhances Antitumor Immunity via Galectin-9-Tim-3 Interactions" *The Journal of Immunology* (2008) vol. 181 pp. 7660-7669.

Naing et al. "A first-in-human phase I study of the anti-PD-1 antibody PDR001 in patients with advanced solid tumors" 2016 ASCO Annual Meeting, *J Clin Oncol* 34, 2016 (suppl; abstr 3060).

Nakae et al., "Mast cells enhance T cell activation: importance of mast cell costimulatory molecules and secreted TNF" *The Journal of Immunology* (2006) vol. 176 No. 4 pp. 2238-2248.

Nellore, A., et al., "The cyclin dependent kinase inhibitor (R)-roscovitine mediates selective suppression of alloreactive human T cells but preserves pathogen-specific and leukemia-specific effectors," *Clinical Immunology* 152(1-2):48-57 (May-Jun. 2014; Epub Mar. 12, 2014).

Ngiow et al. "Prospects for TIM3-Targeted Antitumor Immunotherapy" *Cancer Research* (2011) vol. 71, Issue 21, pp. 6567-6571.

Ngiow et al., "Anti-TIM3 Antibody Promotes T Cell IFN- γ -Mediated Antitumor Immunity and Suppresses Established Tumors" *Cancer Research* (2011) vol. 71 No. 10 pp. 3540-3551.

Nicholson et al., "An Altered Peptide Ligand Mediates Immune Deviation and Prevents Autoimmune Encephalomyelitis" *Immunity* (1995) vol. 3 pp. 397-405.

Nielsen, C., et al., "A putative regulatory polymorphism in PD-1 is associated with nephropathy in a population-based cohort of systemic lupus erythematosus patients," *Lupus* 13(7):510-516, SAGE, England (2004).

Nirschl et al., "Molecular Pathways: Coexpression of Immune Checkpoint Molecules: Signaling Pathways and Implications for Cancer Immunotherapy," *Cancer Res* (2013) vol. 19, No. 18, pp. 4917-4924.

Nishimura et al., "Developmentally regulated expression of the PD-1 protein on the surface of double-negative (CD4-CD8-) thymocytes," *Int Immunol* (1996) vol. 8, No. 5, pp. 773-780.

Christensen et al., "c-Met as a target for human cancer and characterization of inhibitors for therapeutic intervention," *Cancer Letters* (2005) vol. 225, pp. 1-26.

Extended European Search Report issued in European Patent Application No. 18211373.8, dated Jun. 25, 2019.

Fessas et al., "A molecular and preclinical comparison of the PD-1-targeted T-cell checkpoint inhibitors nivolumab and pembrolizumab," *Seminars in Oncology* (2017) vol. 44, pp. 136-140.

Lu et al., "Everolimus enhances the cytotoxicity of bendamustine in multiple myeloma cells through a network of pro-apoptotic and cell-cycle-progression regulatory proteins," *Acta Biochim Biophys Sin* (2013) vol. 45, pp. 683-691.

OPDIVO (nivolumab) Drug Label, Initial U.S. Approval: 2014, Revised Nov. 2016, 58 pages.

Partial European Search Report issued in European Patent Application No. 19206634.8, dated Apr. 30, 2020.

* cited by examiner




Figure 1A




Figure 1B




Figure 1B
Continued




Figure 1C




Figure 2




Figure 3A




Figure 3B




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

Experimental Design




Figure 9A




Figure 9B




Figure 9C




Figure 9D




Figure 9E




Figure 10A




Figure 10B




Figure 11A




Figure 11B




Figure 12




Figure 13




Figure 14

METHODS FOR TREATING HEMATOLOGIC CANCERS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a divisional of U.S. application Ser. No. 15/024,396, filed Mar. 24, 2016, now U.S. Pat. No. 10,570,204, which is a U.S. National Stage Application under 35 U.S.C. § 371 of International Application No. PCT/US2014/057491, filed Sep. 25, 2014, which claims the benefit of U.S. Provisional Application No. 61/882,702, filed Sep. 26, 2013 and U.S. Provisional Application No. 62/017,192, filed Jun. 25, 2014. The contents of the aforesaid applications are hereby incorporated by reference in their entirety.

STATEMENT OF RIGHTS

This invention was made with government support under Grant NCATS 8UL1TR000055 awarded by the National Institutes of Health. The U.S. government has certain rights in the invention. This statement is included solely to comply with 37 C.F.R. § 401.14(a)(f)(4) and should not be taken as an assertion or admission that the application discloses and/or claims only one invention.

SEQUENCE LISTING

The instant application contains a Sequence Listing which has been submitted electronically in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Oct. 22, 2014, is named C2160-7004WO_SL.txt and is 46,210 bytes in size.

BACKGROUND OF THE INVENTION

Hematologic malignancies encompass cancers affecting blood, bone marrow, and lymph nodes and are particularly affected by modulation of the immune system since immune system cells are derived from hematologic lineages. Although hematologic cancers have traditionally been treated with conventional drug therapies, such as alkylating and other DNA damaging compounds, it is increasingly becoming recognized that immune checkpoint regulators play critical roles in determining whether hematologic cancer cells are tolerated or attacked by the immune system (Wu et al. (2012) *Int. J. Biol. Sci.* 8:1420-1430; Nirschl and Drake (2013) *Clin. Cancer Res.*, electronically published July 18; Ceeraz et al. (2013) *Trends Immunol.*, electronically published August 13). However, immune checkpoint regulators, such as CTLA-4, PD-1, VISTA, B7-H2, B7-H3, PD-L1, B7-H4, B7-H6, 2B4, ICOS, HVEM, PD-L2, CD160, gp49B, PIR-B, KIR family receptors, TIM-1, TIM-3, TIM-4, LAG-3, BTLA, SIRPalpha (CD47), CD48, 2B4 (CD244), B7.1, B7.2, ILT-2, ILT-4, TIGIT, LAG-3, BTLA, A2aR and many more, negatively regulate immune response progression based on complex and combinatorial interactions between numerous inputs. While some progress has been made to determine which interventions at which particular nodes of the immune checkpoint regulatory system can be targeted for benefiting the treatment of hematologic cancers (Kearl et al. (2013) *J. Immunol.* 190:5620-5628; Hallett et al. (2011) *Biol. Blood Marrow Transplant.* 17:1133-1145; Pardoll et al. (2012) *Nat. Rev. Cancer* 12:252-264; Brahmer et al. (2012) *N. Engl. J. Med.* 366:2455-2465; Mocellin et al. (2013) *Biochim. Biophys. Acta*

1836:187-196; Topalian et al. (2012) *N. Engl. J. Med.* 366:2443-2454; and Wolchok et al. (2013) *N. Engl. J. Med.* 369:122-133), it is not currently possible to identify specific interactions having synergistic anti-cancer therapeutic efficacy. Accordingly, there is a great need in the art to define specific and synergistic combinations of immune checkpoint regulators useful for treating hematologic cancers.

SUMMARY OF THE INVENTION

The present invention is based, at least in part, on the discovery that inhibiting or blocking Programmed Death 1 (PD-1) or Programmed Death-1 Ligand (PD-L1) in combination with inhibition of an immune checkpoint inhibitor (e.g., one or more of TIM-3, LAG-3 or CTLA4) results in a synergistic therapeutic benefit for treating a hematologic cancer, e.g., a myeloma. This finding is unexpected given the lack of such benefit observed for inhibiting or blocking other combinations of immune checkpoint regulators.

Accordingly, in one aspect, the invention features a method of treating a subject afflicted with a hematologic cancer comprising administering to the subject an inhibitor of PD-1 or PD-L1, and an inhibitor of an immune checkpoint regulator (e.g., an inhibitor of one or more of TIM-3, LAG-3 or CTLA4). In one embodiment, an inhibitor of PD-1 or PD-L1 is administered in combination with an inhibitor of TIM-3. In another embodiment, an inhibitor of PD-1 or PD-L1 is administered in combination with an inhibitor of LAG-3. In yet another embodiment, an inhibitor of PD-1 or PD-L1 is administered in combination with an inhibitor of CTLA-4. Inhibition as described herein can be performed by inhibition at the DNA, RNA or protein level. In embodiments, an inhibitory nucleic acid (e.g., a dsRNA, siRNA or shRNA), can be used to inhibit expression of an inhibitory molecule. In other embodiments, the inhibitor is a polypeptide, e.g., a soluble ligand, or an antibody or antigen-binding fragment thereof, that binds to PD-1 or PD-L1, or other immune checkpoint regulator. Examples of each of the aforesaid inhibitors are provided in more detail below. The term "inhibition" or "inhibitor" includes a reduction in a certain parameter, e.g., an activity, of a given molecule, e.g., an immune checkpoint regulator. For example, inhibition of an activity, e.g., a PD-1 activity, of at least 5%, 10%, 20%, 30%, 40% or more is included by this term. Thus, inhibition need not be 100%. Activities for the immune checkpoint regulators can be determined as described herein or assays known in the art.

In one embodiment, the inhibitor is a bispecific or multispecific antibody selective for PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA4. In another embodiment, a combination of inhibitors comprising a first inhibitor that selectively inhibits or blocks PD-1 or PD-L1 and a second inhibitor that selectively inhibits or blocks TIM-3, LAG-3 or CTLA4 is provided. In one embodiment, the inhibitor is a soluble ligand, e.g., a soluble ligand of PD-1, PD-L1, TIM-3, LAG-3 or CTLA-4 (e.g., a CTLA-4-Ig). In still another embodiment, the first inhibitor and/or second inhibitor is an antibody or an antigen binding fragment thereof, which specifically binds to PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4. In yet another embodiment, the antibody, or antigen binding fragment thereof, is murine, chimeric, humanized, composite, or human. In another embodiment, the antibody, or antigen fragment thereof, is detectably labeled, comprises an effector domain, comprises an Fc domain, and/or is selected from the group consisting of Fv, Fav, F(ab')₂, Fab', dsFv, scFv, sc(Fv)₂, and diabodies fragments. In still another embodiment, the antibody, or

antigen binding fragment thereof, is conjugated to a cytotoxic agent (e.g., a chemotherapeutic agent, a biologic agent, a toxin, a radioactive isotope, and the like).

For example, an anti-PD-1 or PD-L1 antibody, or antigen binding fragment thereof, can be administered in combination with an anti-LAG-3 antibody or an antigen-binding fragment thereof. In another embodiment, an anti-PD-1 or PD-L1 antibody, or antigen binding fragment thereof, is administered in combination with an anti-TIM-3 antibody or antigen-binding fragment thereof. In yet other embodiments, an anti-PD-1 or PD-L1 antibody, or antigen binding fragment thereof, is administered in combination with an anti-LAG-3 antibody and an anti-TIM-3 antibody, or antigen-binding fragments thereof. In yet other embodiments, an anti-PD-1 or PD-L1 antibody, or antigen binding fragment thereof, is administered in combination with an anti-CTLA-4 antibody, or antigen binding fragment thereof (e.g., ipilimumab). Any combination of the aforesaid antibodies can be used in the methods described herein. The combination of antibodies recited herein can be administered separately, e.g., as separate antibodies, or linked, e.g., as a bispecific or trispecific antibody molecule. In one embodiment, a bispecific antibody that includes an anti-PD-1 or PD-L1 antibody, or antigen binding fragment thereof, and an anti-TIM-3, anti-LAG-3 antibody or anti-CTLA4 antibody, or antigen-binding fragment thereof, is administered.

In addition to, or in place of antibodies and antigen binding fragments thereof, numerous other agents are contemplated. For example, in one embodiment, the agent comprises an RNA interfering agent which inhibits expression of PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4 (e.g., a small interfering RNA (siRNA), small hairpin RNA (shRNA), or a microRNA (miRNA)). In another embodiment, the agent comprises an antisense oligonucleotide complementary to PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4. In still another embodiment, the agent comprises a peptide or peptidomimetic that inhibits or blocks PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4. In yet another embodiment, the agent comprises a small molecule that inhibits or blocks PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4 (e.g., a small molecule that inhibits a protein-protein interaction between PD-L1 and a PD-L1 receptor and/or TIM-3 and a TIM-3 receptor). In another embodiment, the agent comprises an aptamer that inhibits or blocks PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4.

Numerous adaptations to the methods described herein are contemplated. For example, in one embodiment, that at least one agent is administered in a pharmaceutically acceptable formulation. In another embodiment, the method further comprises administering to the subject a therapeutic agent for treating the hematologic cancer. In still another embodiment, the method further comprises a step of transient or complete lymphodepletion (e.g., sublethal whole body irradiation used for transient lymphodepletion or lethal whole body irradiation used for complete lymphodepletion). In yet another embodiment, the step of lymphodepletion occurs before, concurrently with, or after the step of agent administration. In another embodiment, the hematologic cancer is selected from the group consisting of multiple myeloma, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, small lymphocytic lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, mantle cell lymphoma, follicular lymphoma, Waldenstrom's macroglobulinemia, B-cell lymphoma and diffuse large B-cell lymphoma, precursor B-lymphoblastic leukemia/lymphoma, B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma, B-cell prolymphocytic leu-

kemia, lymphoplasmacytic lymphoma, splenic marginal zone B-cell lymphoma (with or without villous lymphocytes), hairy cell leukemia, plasma cell myeloma/plasmacytoma, extranodal marginal zone B-cell lymphoma of the MALT type, nodal marginal zone B-cell lymphoma (with or without monocytoid B cells), Burkitt's lymphoma; precursor T-lymphoblastic lymphoma/leukemia, T-cell prolymphocytic leukemia, T-cell granular lymphocytic leukemia, aggressive NK cell leukemia, adult T-cell lymphoma/leukemia (HTLV 1-positive), nasal-type extranodal NK/T-cell lymphoma, enteropathy-type T-cell lymphoma, hepatosplenic γ - δ T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, mycosis fungoides/Sézary syndrome, anaplastic large cell lymphoma (T/null cell, primary cutaneous type), anaplastic large cell lymphoma (T-/null-cell, primary systemic type), peripheral T-cell lymphoma not otherwise characterized, angioimmunoblastic T-cell lymphoma, polycythemia vera (PV), myelodysplastic syndrome (MDS), indolent Non-Hodgkin's Lymphoma (iNHL) and aggressive Non-Hodgkin's Lymphoma (aNHL). In some embodiments, the hematologic cancer is selected from the group consisting of B-cell lymphoma, myeloid leukemia and multiple myeloma or can be multiple myeloma alone. In another embodiment, the subject is a mammal, optionally wherein the mammal is a human.

In another aspect, a kit for treating a subject afflicted with a hematologic cancer comprising an inhibitor of PD-1 or PD-L1, and an inhibitor of an immune checkpoint regulator (e.g., an inhibitor of one or more of TIM-3, LAG-3 or CTLA4), is provided. In one embodiment, the inhibitor is a bispecific or multispecific antibody, or antigen binding fragment thereof, selective for both PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA4. Similarly, a kit for treating a subject afflicted with a hematologic cancer comprising a first agent that selectively inhibits or blocks PD-1 or PD-L1 and a second agent that selectively inhibits or blocks TIM-3, LAG-3 or CTLA4, is provided. In one embodiment, the first agent and/or second agent is an antibody, or an antigen binding fragment thereof, which specifically binds to PD-1 or PD-L1 protein and/or TIM-3, LAG-3 or CTLA4 protein. Any antibody, or antigen binding fragment thereof, provided in a kit can be murine, chimeric, humanized, composite, or human. In another embodiment, the antibody, or antigen binding fragment thereof, is detectably labeled, comprises an effector domain, comprises an Fc domain, and/or is selected from the group consisting of Fv, Fav, F(ab')₂, Fab', dsFv, scFv, sc(Fv)₂, and diabodies fragments. In still another embodiment, the antibody, or antigen binding fragment thereof, is conjugated to a cytotoxic agent (e.g., a chemotherapeutic agent, a biologic agent, a toxin, and a radioactive isotope). In yet another embodiment, the agent is selected from the group consisting of a) an RNA interfering agent which inhibits expression of PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4, optionally wherein said RNA interfering agent is an small interfering RNA (siRNA), small hairpin RNA (shRNA), or a microRNA (miRNA); b) an antisense oligonucleotide complementary to PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4; c) a peptide or peptidomimetics that inhibits or blocks PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4; d) a small molecule that inhibits or blocks PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4, optionally wherein said small molecule inhibits a protein-protein interaction between PD-L1 and a PD-L1 receptor and/or other immune checkpoint regulator (e.g., TIM-3 and a TIM-3

receptor); and e) an aptamer that inhibits or blocks PD-1 or PD-L1 and/or TIM-3, LAG-3 or CTLA4.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1A shows representative flow cytometry dot plots depicting accumulation of tumor cells in myeloma bearing mice over time.

FIG. 1B shows percentages of CD4⁺ and CD8⁺ T cells expressing indicated immune checkpoint proteins in myeloma bearing mice over time. Data shown are representative of more than four independent analyses. *p<0.05, **p<0.01, ***p<0.001 as compared to T cells from naïve non-myeloma bearing mice.

FIG. 1C shows expression of Tim-3 and PD-1, Lag-3 and PD-1, or 2B4 and PD-1 on gated CD8⁺ or CD4⁺ T cells from bone marrow of moribund mice.

FIG. 2 shows representative flow cytometry dot plots depicting expression of indicated immune checkpoint proteins on CD4⁺ Treg cells in myeloma bearing mice and control mice.

FIG. 3A shows expression of Tim-3 and PD-1, Lag-3 and PD-1, or 2B4 and PD-1 on gated CD8⁺ T cells from myeloma bearing mice treated with sublethal whole body irradiation and anti-PD-L1 antibody. Data shown are representative of more than four independent analyses.

FIG. 3B shows frequency of CD8⁺Tim-3⁺, CD8⁺Lag-3⁺ and CD8⁺2B4⁺ cells in spleens of anti-PD-L1 antibody treated myeloma bearing mice compared with spleens of control antibody treated. ***p<0.001.

FIG. 4 shows a schematic diagram of the experimental treatment protocol used to establish and treat myelomas in mice in Example 1.

FIG. 5 shows survival data of myeloma bearing mice treated with inter alia a blocking anti-PD-L1 antibody, a blocking anti-TIM-3 antibody, or combinations thereof.

FIG. 6 shows survival data of myeloma bearing mice treated with inter alia a blocking anti-PD-L1 antibody, a blocking anti-LAG-3 antibody, or combinations thereof.

FIG. 7 shows survival data of myeloma bearing mice treated with inter alia a blocking anti-TIM-3 antibody, a blocking anti-LAG-3 antibody, or combinations thereof.

FIG. 8 shows survival data of myeloma bearing mice treated with inter alia a blocking anti-PD-L1 antibody and a blocking anti-PD-1 antibody.

FIG. 9A shows a schematic diagram of the experimental treatment protocol used to establish and treat myelomas in mice in Example 2.

FIG. 9B shows survival curves of myeloma bearing mice treated with a blocking anti-Tim-3 antibody only, a blocking anti-Lag-3 antibody only, or in combination with a blocking anti-PD-L1 antibody. Survival was compared with control antibody treated mice or mice treated with anti-PD-L1 antibody only. A combination of anti-Lag-3 and anti-Tim-3 antibodies was also tested. Survival curves represent combined data from three independent experiments; n=10-15 mice per experimental group.

FIG. 9C shows survival curves of myeloma bearing mice treated with a blocking anti-CTLA4 antibody only, or in combination with a blocking anti-PD-L1 antibody. Survival was compared with control antibody treated mice or mice treated with anti-PD-L1 antibody only. Survival curves represent combined data from three independent experiments; n=10-15 mice per experimental group.

FIG. 9D shows survival curves of myeloma bearing mice treated with a blocking anti-CD48 antibody only, or in combination with a blocking anti-PD-L1 antibody. Survival

was compared with control antibody treated mice or mice treated with anti-PD-L1 antibody only. Survival curves represent combined data from two independent experiments; n=10-15 mice per experimental group.

FIG. 9E shows survival curves of some of the survivors from FIGS. 9B-9C re-challenged with 5T33 myeloma cells on day 110. P values were determined by the log-rank test.

FIG. 10A shows frequencies of tumor-reactive CD8⁺ and CD4⁺ T cells in the spleens and bone marrow of mice treated with combinations of immune checkpoint protein blockade. The graphs are representative of three independent experiments in which the CD8⁺ or CD4⁺ T cells for each group were pooled from 5-7 individual mice.

FIG. 10B shows levels of cytokine production by CD8⁺ T cells purified from the spleens of mice treated with combinations of immune checkpoint protein blockade. The graphs are representative of two independent experiments in which the CD8⁺ T cells for each group were pooled from 5 individual mice. *p<0.05, **p<0.01 as compared with T cells from mice treated with anti-PD-L1 alone.

FIG. 11A shows expression of PD-1 on gated CD8⁺ T cells from spleen and bone marrow of mice treated with different blocking antibodies or control IgG.

FIG. 11B shows tumor-reactive IFN- γ -secreting cell frequencies in the presence of anti-PD-L1 or control IgG (10 μ g/ml). The graphs are representative of two independent experiments in which the CD8⁺ T cells for each group were pooled from five to seven individual mice. P values were determined by the Student t test. *p<0.05, **p<0.01, ***p<0.001.

FIG. 12 shows levels of cytokine production by CD4⁺ T cells isolated from spleens of myeloma bearing mice treated with combinations of immune checkpoint protein blockade. The graphs are representative of two independent experiments in which the CD4⁺ T cells for each group were pooled from 5 individual mice. *p<0.05, **p<0.01 as compared with T cells from mice treated with anti-PD-L1 alone.

FIG. 13 shows expression of immune checkpoint proteins PD-1, Tim-3, Lag-3 and 2B4 on CD4⁺ and CD8⁺ T cells in mice with other hematologic cancers.

FIG. 14 shows a working model of combined checkpoint blockade and lymphodepleting whole body irradiation.

DETAILED DESCRIPTION OF THE INVENTION

Methods are provided for treating a hematologic cancer, for example, in vitro, ex vivo, or in vivo in a subject, comprising contacting a cancerous cell or administering to a subject a therapeutically effective amount of at least one agent that selectively inhibits or blocks PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA4. In some embodiments, the methods involve a combination of an inhibitor of PD-1 or PD-L1 and an inhibitor of an immune checkpoint inhibitor (e.g., an inhibitor of one or more of TIM-3, LAG-3 or CTLA4). In one embodiment, an inhibitor of PD-1 or PD-L1 is administered in combination with an inhibitor of TIM-3. In another embodiment, an inhibitor of PD-1 or PD-L1 is administered in combination with an inhibitor of LAG-3. In yet another embodiment, an inhibitor of PD-1 or PD-L1 is administered in combination with an inhibitor of CTLA-4. Exemplary inhibitors include a bispecific antibody, multispecific antibody, or combination of individual antibodies that inhibit or block an activity of PD-1 or PD-L1 and another immune checkpoint inhibitor. Such combinations can provide therapeutic benefit for treating hematologic cancers. Such discoveries are especially surprising and unexpected given

reports regarding the lack of activities reported for immune checkpoint regulators, such as TIM-3, in human T cell activation (see, for example, Leitner et al. (2013) *PLoS Pathog.* 9:e1003253).

It will be appreciated that the methods and compositions described herein may be combined with other treatment regimens and/or other predictive biomarkers and methods of using same. It will also be appreciated that the present invention is not limited to the particular embodiments described herein, but can be carried out in variations well known to the skilled artisan.

I. Definitions

The articles “a” and “an” are used herein to refer to one or to more than one (i.e. to at least one) of the grammatical object of the article. By way of example, “an element” means one element or more than one element.

The “amount” of a marker, e.g., expression or copy number of a marker, or protein level of a marker, in a subject is “significantly” higher or lower than the normal amount of a marker, if the amount of the marker is greater or less, respectively, than the normal level by an amount greater than the standard error of the assay employed to assess amount, and preferably at least twice, and more preferably three, four, five, ten or more times that amount. Alternately, the amount of the marker in the subject can be considered “significantly” higher or lower than the normal amount if the amount is at least about two, and preferably at least about three, four, or five times, higher or lower, respectively, than the normal amount of the marker.

The term “altered level of expression” of a marker refers to an expression level or copy number of a marker in a test sample e.g., a sample derived from a subject suffering from cancer, that is greater or less than the standard error of the assay employed to assess expression or copy number, and is preferably at least twice, and more preferably three, four, five or ten or more times the expression level or copy number of the marker or chromosomal region in a control sample (e.g., sample from a healthy subject not having the associated disease) and preferably, the average expression level or copy number of the marker or chromosomal region in several control samples. The altered level of expression is greater or less than the standard error of the assay employed to assess expression or copy number, and is preferably at least twice, and more preferably three, four, five or ten or more times the expression level or copy number of the marker in a control sample (e.g., sample from a healthy subject not having the associated disease) and preferably, the average expression level or copy number of the marker in several control samples.

The term “altered activity” of a marker refers to an activity of a marker which is increased or decreased in a disease state, e.g., in a cancer sample, as compared to the activity of the marker in a normal, control sample. Altered activity of a marker may be the result of, for example, altered expression of the marker, altered protein level of the marker, altered structure of the marker, or, e.g., an altered interaction with other proteins involved in the same or different pathway as the marker, or altered interaction with transcriptional activators or inhibitors.

Unless otherwise specified herein, the terms “antibody” and “antibodies” broadly encompass naturally-occurring forms of antibodies (e.g. IgG, IgA, IgM, IgE) and recombinant antibodies such as single-chain antibodies, chimeric and humanized antibodies and multi-specific antibodies, as well as fragments and derivatives of all of the foregoing,

which fragments and derivatives have at least an antigenic binding site. Antibody derivatives may comprise a protein or chemical moiety conjugated to an antibody.

The term “antibody” as used herein also includes an “antigen-binding portion” of an antibody (or simply “antibody portion”). The term “antigen-binding portion”, as used herein, refers to one or more fragments of an antibody that retain the ability to specifically bind to an antigen (e.g., polypeptide or fragment thereof of PD-1, PD-L1, LAG-3, CTLA-4 and/or TIM-3). It has been shown that the antigen-binding function of an antibody can be performed by fragments of a full-length antibody. Examples of binding fragments encompassed within the term “antigen-binding portion” of an antibody include (i) a Fab fragment, a monovalent fragment consisting of the VL, VH, CL and CH1 domains; (ii) a F(ab')₂ fragment, a bivalent fragment comprising two Fab fragments linked by a disulfide bridge at the hinge region; (iii) a Fd fragment consisting of the VH and CH1 domains; (iv) a Fv fragment consisting of the VL and VH domains of a single arm of an antibody, (v) a dAb fragment (Ward et al., (1989) *Nature* 341:544-546), which consists of a VH domain; and (vi) an isolated complementarity determining region (CDR). Furthermore, although the two domains of the Fv fragment, VL and VH, are coded for by separate genes, they can be joined, using recombinant methods, by a synthetic linker that enables them to be made as a single protein chain in which the VL and VH regions pair to form monovalent polypeptides (known as single chain Fv (scFv); see e.g., Bird et al. (1988) *Science* 242:423-426; and Huston et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:5879-5883; and Osbourn et al. 1998, *Nature Biotechnology* 16: 778). Such single chain antibodies are also intended to be encompassed within the term “antigen-binding portion” of an antibody. Any VH and VL sequences of specific scFv can be linked to human immunoglobulin constant region cDNA or genomic sequences, in order to generate expression vectors encoding complete IgG polypeptides or other isotypes. VH and VL can also be used in the generation of Fab, Fv or other fragments of immunoglobulins using either protein chemistry or recombinant DNA technology. Other forms of single chain antibodies, such as diabodies are also encompassed. Diabodies are bivalent, bispecific antibodies in which VH and VL domains are expressed on a single polypeptide chain, but using a linker that is too short to allow for pairing between the two domains on the same chain, thereby forcing the domains to pair with complementary domains of another chain and creating two antigen binding sites (see e.g., Holliger, P., et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6444-6448; Poljak, R. J., et al. (1994) *Structure* 2:1121-1123).

Still further, an antibody or antigen-binding portion thereof may be part of larger immunoadhesion polypeptides, formed by covalent or noncovalent association of the antibody or antibody portion with one or more other proteins or peptides. Examples of such immunoadhesion polypeptides include use of the streptavidin core region to make a tetrameric scFv polypeptide (Kipriyanov, S. M., et al. (1995) *Human Antibodies and Hybridomas* 6:93-101) and use of a cysteine residue, a marker peptide and a C-terminal polyhistidine tag to make bivalent and biotinylated scFv polypeptides (Kipriyanov, S. M., et al. (1994) *Mol. Immunol.* 31:1047-1058). Antibody portions, such as Fab and F(ab')₂ fragments, can be prepared from whole antibodies using conventional techniques, such as papain or pepsin digestion, respectively, of whole antibodies. Moreover, antibodies,

antibody portions and immunoadhesion polypeptides can be obtained using standard recombinant DNA techniques, as described herein.

Antibodies may be polyclonal or monoclonal; xenogeneic, allogeneic, or syngeneic; or modified forms thereof (e.g., humanized, chimeric, etc.). Antibodies may also be fully human. Preferably, antibodies of the invention bind specifically or substantially specifically to PD-1, PD-L1, LAG-3, CTLA-4 and/or TIM-3 polypeptides or fragments thereof. They may also be selective for such antigens such that they can distinguish such antigens from closely related antigens, such as other B7 family members. The terms “monoclonal antibodies” and “monoclonal antibody composition”, as used herein, refer to a population of antibody polypeptides that contain only one species of an antigen binding site capable of immunoreacting with a particular epitope of an antigen, whereas the term “polyclonal antibodies” and “polyclonal antibody composition” refer to a population of antibody polypeptides that contain multiple species of antigen binding sites capable of interacting with a particular antigen. A monoclonal antibody composition typically displays a single binding affinity for a particular antigen with which it immunoreacts.

As used herein, a “blocking” antibody or an antibody “antagonist” or “inhibitor” is one which inhibits or reduces at least one biological activity of the antigen(s) it binds. For example, an anti-PD-L1 or anti-TIM-3 antibody binds PD-L1 or TIM-3, respectively, and inhibits the ability of PD-L1 to, for example, bind PD-1, and inhibits the ability of TIM-3 to, for example, bind galectin-9 or phosphatidylserine. In certain embodiments, the blocking antibodies or antagonist antibodies or fragments thereof described herein substantially or completely inhibit a given biological activity of the antigen(s).

The term “body fluid” refers to fluids that are excreted or secreted from the body as well as fluid that are normally not (e.g. amniotic fluid, aqueous humor, bile, blood and blood plasma, cerebrospinal fluid, cerumen and earwax, cowper’s fluid or pre-ejaculatory fluid, chyle, chyme, stool, female ejaculate, interstitial fluid, intracellular fluid, lymph, menses, breast milk, mucus, pleural fluid, pus, saliva, sebum, semen, serum, sweat, synovial fluid, tears, urine, vaginal lubrication, vitreous humor, vomit).

The term “bispecific antibody” or “multispecific antibody” refers to an antibody that recognized more than one epitope. Such antibodies are useful for targeting different proteins using the same agent. Methods of making such antibodies are well known in art (see, at least U.S. Pat. Nos. 5,798,229; 5,989,830; and Holliger et al. (2005) *Nat. Biotech.* 23:1126-1136).

The terms “cancer” or “tumor” or “hyperproliferative disorder” refer to the presence of cells possessing characteristics typical of cancer-causing cells, such as uncontrolled proliferation, immortality, metastatic potential, rapid growth and proliferation rate, and certain characteristic morphological features. Cancer cells are often in the form of a tumor, but such cells may exist alone within an animal, or may be a non-tumorigenic cancer cell, such as a leukemia cell. Cancers include, but are not limited to, B cell cancer, e.g., multiple myeloma, Waldenstrom’s macroglobulinemia, the heavy chain diseases, such as, for example, alpha chain disease, gamma chain disease, and mu chain disease, benign monoclonal gammopathy, and immunocytic amyloidosis, melanomas, breast cancer, lung cancer, bronchus cancer, colorectal cancer, prostate cancer, pancreatic cancer, stomach cancer, ovarian cancer, urinary bladder cancer, brain or central nervous system cancer, peripheral nervous system

cancer, esophageal cancer, cervical cancer, uterine or endometrial cancer, cancer of the oral cavity or pharynx, liver cancer, kidney cancer, testicular cancer, biliary tract cancer, small bowel or appendix cancer, salivary gland cancer, thyroid gland cancer, adrenal gland cancer, osteosarcoma, chondrosarcoma, cancer of hematologic tissues, and the like. Other non-limiting examples of types of cancers applicable to the methods encompassed by the present invention include human sarcomas and carcinomas, e.g., fibrosarcoma, myxosarcoma, liposarcoma, chondrosarcoma, osteogenic sarcoma, chordoma, angiosarcoma, endotheliosarcoma, lymphangiosarcoma, lymphangioendotheliosarcoma, synovialoma, mesothelioma, Ewing’s tumor, leiomyosarcoma, rhabdomyosarcoma, colon carcinoma, colorectal cancer, pancreatic cancer, breast cancer, ovarian cancer, prostate cancer, squamous cell carcinoma, basal cell carcinoma, adenocarcinoma, sweat gland carcinoma, sebaceous gland carcinoma, papillary carcinoma, papillary adenocarcinomas, cystadenocarcinoma, medullary carcinoma, bronchogenic carcinoma, renal cell carcinoma, hepatoma, bile duct carcinoma, liver cancer, choriocarcinoma, seminoma, embryonal carcinoma, Wilms’ tumor, cervical cancer, bone cancer, brain tumor, testicular cancer, lung carcinoma, small cell lung carcinoma, bladder carcinoma, epithelial carcinoma, glioma, astrocytoma, medulloblastoma, craniopharyngioma, ependymoma, pinealoma, hemangioblastoma, acoustic neuroma, oligodendroglioma, meningioma, melanoma, neuroblastoma, retinoblastoma; leukemias, e.g., acute lymphocytic leukemia and acute myelocytic leukemia (myeloblastic, promyelocytic, myelomonocytic, monocytic and erythroleukemia); chronic leukemia (chronic myelocytic (granulocytic) leukemia and chronic lymphocytic leukemia); and polycythemia vera, lymphoma (Hodgkin’s disease and non-Hodgkin’s disease), multiple myeloma, Waldenstrom’s macroglobulinemia, and heavy chain disease. In some embodiments, cancers are epithelial in nature and include but are not limited to, bladder cancer, breast cancer, cervical cancer, colon cancer, gynecologic cancers, renal cancer, laryngeal cancer, lung cancer, oral cancer, head and neck cancer, ovarian cancer, pancreatic cancer, prostate cancer, or skin cancer. In other embodiments, the cancer is breast cancer, prostate cancer, lung cancer, or colon cancer. In still other embodiments, the epithelial cancer is non-small-cell lung cancer, nonpapillary renal cell carcinoma, cervical carcinoma, ovarian carcinoma (e.g., serous ovarian carcinoma), or breast carcinoma. The epithelial cancers may be characterized in various other ways including, but not limited to, serous, endometrioid, mucinous, clear cell, Brenner, or undifferentiated.

The term “control” refers to any reference standard suitable to provide a comparison to the expression products in the test sample. In one embodiment, the control comprises obtaining a “control sample” from which expression product levels are detected and compared to the expression product levels from the test sample. Such a control sample may comprise any suitable sample, including but not limited to a sample from a control cancer patient (can be stored sample or previous sample measurement) with a known outcome; normal tissue or cells isolated from a subject, such as a normal patient or the cancer patient, cultured primary cells/tissues isolated from a subject such as a normal subject or the cancer patient, adjacent normal cells/tissues obtained from the same organ or body location of the cancer patient, a tissue or cell sample isolated from a normal subject, or a primary cells/tissues obtained from a depository. In another preferred embodiment, the control may comprise a reference standard expression product level from any suitable source,

including but not limited to housekeeping genes, an expression product level range from normal tissue (or other previously analyzed control sample), a previously determined expression product level range within a test sample from a group of patients, or a set of patients with a certain outcome (for example, survival for one, two, three, four years, etc.) or receiving a certain treatment (for example, standard of care cancer therapy). It will be understood by those of skill in the art that such control samples and reference standard expression product levels can be used in combination as controls in the methods of the present invention. In one embodiment, the control may comprise normal or non-cancerous cell/tissue sample. In another preferred embodiment, the control may comprise an expression level for a set of patients, such as a set of cancer patients, or for a set of cancer patients receiving a certain treatment, or for a set of patients with one outcome versus another outcome. In the former case, the specific expression product level of each patient can be assigned to a percentile level of expression, or expressed as either higher or lower than the mean or average of the reference standard expression level. In another preferred embodiment, the control may comprise normal cells, cells from patients treated with combination chemotherapy, and cells from patients having benign cancer. In another embodiment, the control may also comprise a measured value for example, average level of expression of a particular gene in a population compared to the level of expression of a housekeeping gene in the same population. Such a population may comprise normal subjects, cancer patients who have not undergone any treatment (i.e., treatment naive), cancer patients undergoing standard of care therapy, or patients having benign cancer. In another preferred embodiment, the control comprises a ratio transformation of expression product levels, including but not limited to determining a ratio of expression product levels of two genes in the test sample and comparing it to any suitable ratio of the same two genes in a reference standard; determining expression product levels of the two or more genes in the test sample and determining a difference in expression product levels in any suitable control; and determining expression product levels of the two or more genes in the test sample, normalizing their expression to expression of housekeeping genes in the test sample, and comparing to any suitable control. In particularly preferred embodiments, the control comprises a control sample which is of the same lineage and/or type as the test sample. In another embodiment, the control may comprise expression product levels grouped as percentiles within or based on a set of patient samples, such as all patients with cancer. In one embodiment a control expression product level is established wherein higher or lower levels of expression product relative to, for instance, a particular percentile, are used as the basis for predicting outcome. In another preferred embodiment, a control expression product level is established using expression product levels from cancer control patients with a known outcome, and the expression product levels from the test sample are compared to the control expression product level as the basis for predicting outcome. As demonstrated by the data below, the methods of the invention are not limited to use of a specific cut-point in comparing the level of expression product in the test sample to the control.

The term "hematologic cancer" refers to cancers of cells derived from the blood. In some embodiments, the hematologic cancer is selected from the group consisting of acute lymphocytic leukemia, myeloid leukemia including acute myeloid leukemia and chronic myelogenous leukemia, chronic lymphocytic leukemia, small lymphocytic lymphoma,

multiple myeloma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, mantle cell lymphoma, follicular lymphoma, Waldenstrom's macroglobulinemia, B-cell lymphoma and including diffuse large B-cell lymphoma (including primary mediastinal B-cell lymphoma and intravascular large B-cell lymphoma), follicular lymphoma, chronic lymphocytic leukemia, small lymphocytic lymphoma, mantle cell lymphoma, mucosa-associated lymphoid tissue (MALT) lymphomas 9 e.g., extranodal marginal zone B-cell lymphoma of the MALT type, nodal marginal zone B-cell lymphoma (with or without monocytoid B cells), marginal zone B-cell lymphomas (e.g., nodal marginal zone B-cell lymphoma and splenic marginal zone B-cell lymphoma (with or without villous lymphocytes)), Burkitt lymphoma, lymphoplasmacytic lymphoma (Waldenstrom macroglobulinemia), mediastinal large B cell lymphoma, precursor B-lymphoblastic leukemia/lymphoma and, B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma, B-cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, splenic marginal zone B-cell lymphoma (with or without villous lymphocytes), hairy cell leukemia, plasma cell myeloma/plasmacytoma, extranodal marginal zone B-cell lymphoma of the MALT type, nodal marginal zone B-cell lymphoma (with or without monocytoid B cells), Burkitt's lymphoma; precursor T-lymphoblastic lymphoma/leukemia, T-cell prolymphocytic leukemia, T-cell granular lymphocytic leukemia, aggressive NK cell leukemia, adult T-cell lymphoma/leukemia (HTLV 1-positive), nasal-type extranodal NK/T-cell lymphoma, enteropathy-type T-cell lymphoma, hepatosplenic γ - δ T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, mycosis fungoides/Sézary syndrome, anaplastic large cell lymphoma (T/null cell, primary cutaneous type), anaplastic large cell lymphoma (T-/null-cell, primary systemic type), peripheral T-cell lymphoma not otherwise characterized, angioimmunoblastic T-cell lymphoma, polycythemia vera (PV), myelodysplastic syndrome (MDS). NHL may include indolent Non-Hodgkin's Lymphoma (iNHL) or aggressive Non-Hodgkin's Lymphoma (aNHL).

As used herein, the term "coding region" refers to regions of a nucleotide sequence comprising codons which are translated into amino acid residues, whereas the term "non-coding region" refers to regions of a nucleotide sequence that are not translated into amino acids (e.g., 5' and 3' untranslated regions).

As used herein, the term "complementary" refers to the broad concept of sequence complementarity between regions of two nucleic acid strands or between two regions of the same nucleic acid strand. It is known that an adenine residue of a first nucleic acid region is capable of forming specific hydrogen bonds ("base pairing") with a residue of a second nucleic acid region which is antiparallel to the first region if the residue is thymine or uracil. Similarly, it is known that a cytosine residue of a first nucleic acid strand is capable of base pairing with a residue of a second nucleic acid strand which is antiparallel to the first strand if the residue is guanine. A first region of a nucleic acid is complementary to a second region of the same or a different nucleic acid if, when the two regions are arranged in an antiparallel fashion, at least one nucleotide residue of the first region is capable of base pairing with a residue of the second region. Preferably, the first region comprises a first portion and the second region comprises a second portion, whereby, when the first and second portions are arranged in an antiparallel fashion, at least about 50%, and preferably at least about 75%, at least about 90%, or at least about 95% of the nucleotide residues of the first portion are capable of

base pairing with nucleotide residues in the second portion. More preferably, all nucleotide residues of the first portion are capable of base pairing with nucleotide residues in the second portion.

As used herein, the term “determining a suitable treatment regimen for the subject” is taken to mean the determination of a treatment regimen (i.e., a single therapy or a combination of different therapies that are used for the prevention and/or treatment of the cancer in the subject) for a subject that is started, modified and/or ended based or essentially based or at least partially based on the results of the analysis according to the present invention. One example is starting an adjuvant therapy after surgery whose purpose is to decrease the risk of recurrence, another would be to modify the dosage of a particular chemotherapy. The determination can, in addition to the results of the analysis according to the present invention, be based on personal characteristics of the subject to be treated. In most cases, the actual determination of the suitable treatment regimen for the subject will be performed by the attending physician or doctor.

“Homologous” as used herein, refers to nucleotide sequence similarity between two regions of the same nucleic acid strand or between regions of two different nucleic acid strands. When a nucleotide residue position in both regions is occupied by the same nucleotide residue, then the regions are homologous at that position. A first region is homologous to a second region if at least one nucleotide residue position of each region is occupied by the same residue. Homology between two regions is expressed in terms of the proportion of nucleotide residue positions of the two regions that are occupied by the same nucleotide residue. By way of example, a region having the nucleotide sequence 5'-AT-TGCC-3' and a region having the nucleotide sequence 5'-TATGGC-3' share 50% homology. Preferably, the first region comprises a first portion and the second region comprises a second portion, whereby, at least about 50%, and preferably at least about 75%, at least about 90%, or at least about 95% of the nucleotide residue positions of each of the portions are occupied by the same nucleotide residue. More preferably, all nucleotide residue positions of each of the portions are occupied by the same nucleotide residue.

As used herein, the term “host cell” is intended to refer to a cell into which a nucleic acid of the invention, such as a recombinant expression vector of the invention, has been introduced. The terms “host cell” and “recombinant host cell” are used interchangeably herein. It should be understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

The term “humanized antibody,” as used herein, is intended to include antibodies made by a non-human cell having variable and constant regions which have been altered to more closely resemble antibodies that would be made by a human cell. For example, by altering the non-human antibody amino acid sequence to incorporate amino acids found in human germline immunoglobulin sequences. Humanized antibodies may include amino acid residues not encoded by human germline immunoglobulin sequences (e.g., mutations introduced by random or site-specific mutagenesis in vitro or by somatic mutation in vivo), for example in the CDRs. The term “humanized antibody”, as used herein, also includes antibodies in which CDR sequences

derived from the germline of another mammalian species, such as a mouse, have been grafted onto human framework sequences.

As used herein, the term “immune cell” refers to cells that play a role in the immune response. Immune cells are of hematopoietic origin, and include lymphocytes, such as B cells and T cells; natural killer cells; myeloid cells, such as monocytes, macrophages, eosinophils, mast cells, basophils, and granulocytes.

As used herein, the term “immune checkpoints” or “immune checkpoint regulators” means a group of molecules on the cell surface of CD4⁺ and CD8⁺ T cells. These molecules fine-tune immune responses by down-modulating or inhibiting an immune response, e.g., an anti-tumor immune response. Immune checkpoint proteins are known in the art and include, without limitation, CTLA-4, PD-1, VISTA, B7-H2, B7-H3, PD-L1, B7-H4, B7-H6, 2B4, ICOS, HVEM, PD-L2, CD160, gp49B, PIR-B, KIR family receptors, TIM-1, TIM-3, TIM-4, LAG-3, BTLA, SIRPalpha (CD47), CD48, 2B4 (CD244), B7.1, B7.2, ILT-2, ILT-4, TIGIT, LAG-3, BTLA, and A2aR (see, for example, WO 2012/177624). Immunotherapeutic agents that can act as immune checkpoint inhibitors useful in the methods of the present invention, include, but are not limited to, inhibitors of PD-1, PD-L1, TIM-3, LAG-3 and CTLA-4 (e.g., soluble peptide inhibitors or antibodies, e.g., anti-PD-1 antibodies, anti-PD-L1 antibodies, anti-CTLA-4 antibodies, anti-TIM-3 antibodies, and anti-LAG-3 antibodies).

As used herein, the term “immune response” includes T cell mediated and/or B cell mediated immune responses. Exemplary immune responses include T cell responses, e.g., cytokine production and cellular cytotoxicity. In addition, the term immune response includes immune responses that are indirectly effected by T cell activation, e.g., antibody production (humoral responses) and activation of cytokine responsive cells, e.g., macrophages.

As used herein, the term “immunotherapeutic agent” can include any molecule, peptide, antibody or other agent which can stimulate a host immune system to generate an immune response to a tumor or cancer in the subject. Various immunotherapeutic agents are useful in the compositions and methods described herein.

An “inducible” promoter is a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a living human cell substantially only when an inducer which corresponds to the promoter is present in the cell.

As used herein, the term “inhibit” refers to any decrease in, for example a particular action, function, or interaction. For example, cancer is “inhibited” if at least one symptom of the cancer is reduced, slowed, or delayed. As used herein, cancer is also “inhibited” if recurrence or metastasis of the cancer is reduced, slowed, or delayed.

As used herein, the term “interaction,” when referring to an interaction between two molecules, refers to the physical contact (e.g., binding) of the molecules with one another. Generally, such an interaction results in an activity (which produces a biological effect) of one or both of said molecules.

An “isolated antibody” is intended to refer to an antibody that is substantially free of other antibodies having different antigenic specificities (e.g., an isolated antibody that specifically binds PD-L1 polypeptide or a fragment thereof, or TIM-3 polypeptide or a fragment thereof, is substantially free of antibodies that specifically bind antigens other than

said polypeptide or a fragment thereof). Moreover, an isolated antibody may be substantially free of other cellular material and/or chemicals.

As used herein, an “isolated protein” refers to a protein that is substantially free of other proteins, cellular material, separation medium, and culture medium when isolated from cells or produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. An “isolated” or “purified” protein or biologically active portion thereof is substantially free of cellular material or other contaminating proteins from the cell or tissue source from which the antibody, polypeptide, peptide or fusion protein is derived, or substantially free from chemical precursors or other chemicals when chemically synthesized. The language “substantially free of cellular material” includes preparations, in which compositions of the invention are separated from cellular components of the cells from which they are isolated or recombinantly produced. In one embodiment, the language “substantially free of cellular material” includes preparations of having less than about 30%, 20%, 10%, or 5% (by dry weight) of cellular material. When an antibody, polypeptide, peptide or fusion protein or fragment thereof, e.g., a biologically active fragment thereof, is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation.

A “kit” is any manufacture (e.g. a package or container) comprising at least one reagent, e.g. a probe or small molecule, for specifically detecting and/or affecting the expression of a marker of the invention. The kit may be promoted, distributed, or sold as a unit for performing the methods of the present invention. The kit may comprise one or more reagents necessary to express a composition useful in the methods of the present invention. In certain embodiments, the kit may further comprise a reference standard, e.g., a nucleic acid encoding a protein that does not affect or regulate signaling pathways controlling cell growth, division, migration, survival or apoptosis. One skilled in the art can envision many such control proteins, including, but not limited to, common molecular tags (e.g., green fluorescent protein and beta-galactosidase), proteins not classified in any of pathway encompassing cell growth, division, migration, survival or apoptosis by GeneOntology reference, or ubiquitous housekeeping proteins. Reagents in the kit may be provided in individual containers or as mixtures of two or more reagents in a single container. In addition, instructional materials which describe the use of the compositions within the kit can be included.

A “marker” is a gene whose altered level of expression in a tissue or cell from its expression level in normal or healthy tissue or cell is associated with a disease state, such as cancer. A “marker nucleic acid” is a nucleic acid (e.g., mRNA, cDNA) encoded by or corresponding to a marker of the invention. Such marker nucleic acids include DNA (e.g., cDNA) comprising the entire or a partial sequence of any of the nucleic acid sequences set forth in the Sequence Listing or the complement of such a sequence. The marker nucleic acids also include RNA comprising the entire or a partial sequence of any of the nucleic acid sequences set forth in the Sequence Listing or the complement of such a sequence, wherein all thymidine residues are replaced with uridine residues. A “marker protein” is a protein encoded by or corresponding to a marker of the invention. A marker protein comprises the entire or a partial sequence of any of the

sequences set forth in the Sequence Listing. The terms “protein” and “polypeptide” are used interchangeably.

The “normal” level of expression of a marker is the level of expression of the marker in cells of a subject, e.g., a human patient, not afflicted with a cancer. An “over-expression” or “significantly higher level of expression” of a marker refers to an expression level in a test sample that is greater than the standard error of the assay employed to assess expression, and is preferably at least twice, and more preferably 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 times or more higher than the expression activity or level of the marker in a control sample (e.g., sample from a healthy subject not having the marker associated disease) and preferably, the average expression level of the marker in several control samples. A “significantly lower level of expression” of a marker refers to an expression level in a test sample that is at least twice, and more preferably 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 times or more lower than the expression level of the marker in a control sample (e.g., sample from a healthy subject not having the marker associated disease) and preferably, the average expression level of the marker in several control samples.

An “over-expression” or “significantly higher level of expression” of a marker refers to an expression level in a test sample that is greater than the standard error of the assay employed to assess expression, and is preferably at least twice, and more preferably 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 times or more higher than the expression activity or level of the marker in a control sample (e.g., sample from a healthy subject not having the marker associated disease) and preferably, the average expression level of the marker in several control samples. A “significantly lower level of expression” of a marker refers to an expression level in a test sample that is at least twice, and more preferably 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3, 3.5, 4, 4.5, 5, 5.5, 6, 6.5, 7, 7.5, 8, 8.5, 9, 9.5, 10, 10.5, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 times or more lower than the expression level of the marker in a control sample (e.g., sample from a healthy subject not having the marker associated disease) and preferably, the average expression level of the marker in several control samples.

The term “peripheral blood cell subtypes” refers to cell types normally found in the peripheral blood including, but is not limited to, eosinophils, neutrophils, T cells, monocytes, NK cells, granulocytes, and B cells.

The term “probe” refers to any molecule which is capable of selectively binding to a specifically intended target molecule, for example, a nucleotide transcript or protein encoded by or corresponding to a marker. Probes can be either synthesized by one skilled in the art, or derived from appropriate biological preparations. For purposes of detection of the target molecule, probes may be specifically designed to be labeled, as described herein. Examples of molecules that can be utilized as probes include, but are not limited to, RNA, DNA, proteins, antibodies, and organic molecules.

The term “prognosis” includes a prediction of the probable course and outcome of cancer or the likelihood of recovery from the disease. In some embodiments, the use of statistical algorithms provides a prognosis of cancer in an individual. For example, the prognosis can be surgery, development of a clinical subtype of cancer (e.g., hematologic cancers, such as multiple myeloma), development of

one or more clinical factors, development of intestinal cancer, or recovery from the disease.

The term “response to cancer therapy” or “outcome of cancer therapy” relates to any response of the hyperproliferative disorder (e.g., cancer) to a cancer therapy, preferably to a change in tumor mass and/or volume after initiation of neoadjuvant or adjuvant chemotherapy. Hyperproliferative disorder response may be assessed, for example for efficacy or in a neoadjuvant or adjuvant situation, where the size of a tumor after systemic intervention can be compared to the initial size and dimensions as measured by CT, PET, mammogram, ultrasound or palpation. Response may also be assessed by caliper measurement or pathological examination of the tumor after biopsy or surgical resection for solid cancers. Responses may be recorded in a quantitative fashion like percentage change in tumor volume or in a qualitative fashion like “pathological complete response” (pCR), “clinical complete remission” (cCR), “clinical partial remission” (cPR), “clinical stable disease” (cSD), “clinical progressive disease” (cPD) or other qualitative criteria. Assessment of hyperproliferative disorder response may be done early after the onset of neoadjuvant or adjuvant therapy, e.g., after a few hours, days, weeks or preferably after a few months. A typical endpoint for response assessment is upon termination of neoadjuvant chemotherapy or upon surgical removal of residual tumor cells and/or the tumor bed. This is typically three months after initiation of neoadjuvant therapy. In some embodiments, clinical efficacy of the therapeutic treatments described herein may be determined by measuring the clinical benefit rate (CBR). The clinical benefit rate is measured by determining the sum of the percentage of patients who are in complete remission (CR), the number of patients who are in partial remission (PR) and the number of patients having stable disease (SD) at a time point at least 6 months out from the end of therapy. The shorthand for this formula is $CBR = CR + PR + SD$ over 6 months. In some embodiments, the CBR for a particular cancer therapeutic regimen is at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, or more. In other embodiments, the percentage of patients who are in either CR, PR, and/or SD in any combination at least 30 days, 2 months, 3 months, 4 months, 5 months, 6 months, 7 months, 8 months, 9 months, 10 months, 11 months, 12 months, 18 months, 24 months, 30 months, 36 months, 60 months, or longer is at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, or more. In some embodiments, the percentage is 100% over such a time period. Additional criteria for evaluating the response to cancer therapies are related to “survival,” which includes all of the following: survival until mortality, also known as overall survival (wherein said mortality may be either irrespective of cause or tumor related); “recurrence-free survival” (wherein the term recurrence shall include both localized and distant recurrence); metastasis free survival; disease free survival (wherein the term disease shall include cancer and diseases associated therewith). The length of said survival may be calculated by reference to a defined start point (e.g., time of diagnosis or start of treatment) and end point (e.g., death, recurrence or metastasis). In addition, criteria for efficacy of treatment can be expanded to include response to chemotherapy, probability of survival, probability of metastasis within a given time period, and probability of tumor recurrence. For example, in order to determine appropriate threshold values, a particular cancer therapeutic regimen can be administered to a population of subjects and the outcome can be correlated to copy number, level of expression, level of activity, etc. of a marker determined

prior to administration of any cancer therapy. The outcome measurement may be pathologic response to therapy given in the neoadjuvant setting. Alternatively, outcome measures, such as overall survival and disease-free survival can be monitored over a period of time for subjects following cancer therapy for whom the measurement values are known. In certain embodiments, the same doses of cancer therapeutic agents are administered to each subject. In related embodiments, the doses administered are standard doses known in the art for cancer therapeutic agents. The period of time for which subjects are monitored can vary. For example, subjects may be monitored for at least 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 25, 30, 35, 40, 45, 50, 55, or 60 months. Biomarker threshold values that correlate to outcome of a cancer therapy can be determined using methods such as those described in the Examples section. Outcomes can also be measured in terms of a “hazard ratio” (the ratio of death rates for one patient group to another; provides likelihood of death at a certain time point), “overall survival” (OS), and/or “progression free survival.” In certain embodiments, the prognosis comprises likelihood of overall survival rate at 1 year, 2 years, 3 years, 4 years, or any other suitable time point. The significance associated with the prognosis of poor outcome in all aspects of the present invention is measured by techniques known in the art. For example, significance may be measured with calculation of odds ratio. In a further embodiment, the significance is measured by a percentage. In one embodiment, a significant risk of poor outcome is measured as odds ratio of 0.8 or less or at least about 1.2, including by not limited to: 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.5, 3.0, 4.0, 5.0, 10.0, 15.0, 20.0, 25.0, 30.0 and 40.0. In a further embodiment, a significant increase or reduction in risk is at least about 20%, including but not limited to about 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% and 98%. In a further embodiment, a significant increase in risk is at least about 50%. Thus, the present invention further provides methods for making a treatment decision for a cancer patient, comprising carrying out the methods for prognosing a cancer patient according to the different aspects and embodiments of the present invention, and then weighing the results in light of other known clinical and pathological risk factors, in determining a course of treatment for the cancer patient. For example, a cancer patient that is shown by the methods of the invention to have an increased risk of poor outcome by combination chemotherapy treatment can be treated with more aggressive therapies, including but not limited to radiation therapy, peripheral blood stem cell transplant, bone marrow transplant, or novel or experimental therapies under clinical investigation.

The term “resistance” refers to an acquired or natural resistance of a cancer sample or a mammal to a cancer therapy (i.e., being nonresponsive to or having reduced or limited response to the therapeutic treatment), such as having a reduced response to a therapeutic treatment by 25% or more, for example, 30%, 40%, 50%, 60%, 70%, 80%, or more, to 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 15-fold, 20-fold or more. The reduction in response can be measured by comparing with the same cancer sample or mammal before the resistance is acquired, or by comparing with a different cancer sample or a mammal who is known to have no resistance to the therapeutic treatment. A typical acquired resistance to chemotherapy is called “multidrug resistance.” The multidrug resistance can be mediated by P-glycoprotein or can be mediated by other mechanisms, or it can occur when a mammal is infected with a multi drug-resistant

microorganism or a combination of microorganisms. The determination of resistance to a therapeutic treatment is routine in the art and within the skill of an ordinarily skilled clinician, for example, can be measured by cell proliferative assays and cell death assays as described herein as “sensitizing.” In some embodiments, the term “reverses resistance” means that the use of a second agent in combination with a primary cancer therapy (e.g., chemotherapeutic or radiation therapy) is able to produce a significant decrease in tumor volume at a level of statistical significance (e.g., $p < 0.05$) when compared to tumor volume of untreated tumor in the circumstance where the primary cancer therapy (e.g., chemotherapeutic or radiation therapy) alone is unable to produce a statistically significant decrease in tumor volume compared to tumor volume of untreated tumor. This generally applies to tumor volume measurements made at a time when the untreated tumor is growing log rhythmically.

An “RNA interfering agent” as used herein, is defined as any agent which interferes with or inhibits expression of a target gene, e.g., a marker of the invention, by RNA interference (RNAi). Such RNA interfering agents include, but are not limited to, nucleic acid molecules including RNA molecules which are homologous to the target gene, e.g., a marker of the invention, or a fragment thereof, short interfering RNA (siRNA), and small molecules which interfere with or inhibit expression of a target gene by RNA interference (RNAi).

“RNA interference (RNAi)” is an evolutionally conserved process whereby the expression or introduction of RNA of a sequence that is identical or highly similar to a target gene results in the sequence specific degradation or specific post-transcriptional gene silencing (PTGS) of messenger RNA (mRNA) transcribed from that targeted gene (see Coburn and Cullen (2002) *J. Virol.* 76(18):9225), thereby inhibiting expression of the target gene. In one embodiment, the RNA is double stranded RNA (dsRNA). This process has been described in plants, invertebrates, and mammalian cells. In nature, RNAi is initiated by the dsRNA-specific endonuclease Dicer, which promotes processive cleavage of long dsRNA into double-stranded fragments termed siRNAs. siRNAs are incorporated into a protein complex that recognizes and cleaves target mRNAs. RNAi can also be initiated by introducing nucleic acid molecules, e.g., synthetic siRNAs or RNA interfering agents, to inhibit or silence the expression of target genes. As used herein, “inhibition of target gene expression” or “inhibition of marker gene expression” includes any decrease in expression or protein activity or level of the target gene (e.g., a marker gene of the invention) or protein encoded by the target gene, e.g., a marker protein of the invention. The decrease may be of at least 30%, 40%, 50%, 60%, 70%, 80%, 90%, 95% or 99% or more as compared to the expression of a target gene or the activity or level of the protein encoded by a target gene which has not been targeted by an RNA interfering agent.

The term “sample” used for detecting or determining the presence or level of at least one biomarker is typically whole blood, plasma, serum, saliva, urine, stool (e.g., feces), tears, and any other bodily fluid (e.g., as described above under the definition of “body fluids”), or a tissue sample (e.g., biopsy) such as a small intestine, colon sample, or surgical resection tissue. In certain instances, the method of the present invention further comprises obtaining the sample from the individual prior to detecting or determining the presence or level of at least one marker in the sample.

The term “sensitize” means to alter cancer cells or tumor cells in a way that allows for more effective treatment of the

associated cancer with a cancer therapy (e.g., chemotherapeutic or radiation therapy). In some embodiments, normal cells are not affected to an extent that causes the normal cells to be unduly injured by the cancer therapy (e.g., chemotherapy or radiation therapy). An increased sensitivity or a reduced sensitivity to a therapeutic treatment is measured according to a known method in the art for the particular treatment and methods described herein below, including, but not limited to, cell proliferative assays (Tanigawa N, Kern D H, Kikasa Y, Morton D L, *Cancer Res* 1982; 42: 2159-2164), cell death assays (Weisenthal L M, Shoemaker R H, Marsden J A, Dill P L, Baker J A, Moran E M, *Cancer Res* 1984; 94: 161-173; Weisenthal L M, Lippman M E, *Cancer Treat Rep* 1985; 69: 615-632; Weisenthal L M, In: Kaspers G J L, Pieters R, Twentymann P R, Weisenthal L M, Veerman A J P, eds. *Drug Resistance in Leukemia and Lymphoma*. Langhorne, P A: Harwood Academic Publishers, 1993: 415-432; Weisenthal L M, *Contrib Gynecol Obstet* 1994; 19: 82-90). The sensitivity or resistance may also be measured in animal by measuring the tumor size reduction over a period of time, for example, 6 month for human and 4-6 weeks for mouse. A composition or a method sensitizes response to a therapeutic treatment if the increase in treatment sensitivity or the reduction in resistance is 25% or more, for example, 30%, 40%, 50%, 60%, 70%, 80%, or more, to 2-fold, 3-fold, 4-fold, 5-fold, 10-fold, 15-fold, 20-fold or more, compared to treatment sensitivity or resistance in the absence of such composition or method. The determination of sensitivity or resistance to a therapeutic treatment is routine in the art and within the skill of an ordinarily skilled clinician. It is to be understood that any method described herein for enhancing the efficacy of a cancer therapy can be equally applied to methods for sensitizing hyperproliferative or otherwise cancerous cells (e.g., resistant cells) to the cancer therapy.

“Short interfering RNA” (siRNA), also referred to herein as “small interfering RNA” is defined as an agent which functions to inhibit expression of a target gene, e.g., by RNAi. An siRNA may be chemically synthesized, may be produced by in vitro transcription, or may be produced within a host cell. In one embodiment, siRNA is a double stranded RNA (dsRNA) molecule of about 15 to about 40 nucleotides in length, preferably about 15 to about 28 nucleotides, more preferably about 19 to about 25 nucleotides in length, and more preferably about 19, 20, 21, or 22 nucleotides in length, and may contain a 3' and/or 5' overhang on each strand having a length of about 0, 1, 2, 3, 4, or 5 nucleotides. The length of the overhang is independent between the two strands, i.e., the length of the overhang on one strand is not dependent on the length of the overhang on the second strand. Preferably the siRNA is capable of promoting RNA interference through degradation or specific post-transcriptional gene silencing (PTGS) of the target messenger RNA (mRNA). In another embodiment, an siRNA is a small hairpin (also called stem loop) RNA (shRNA). In one embodiment, these shRNAs are composed of a short (e.g., 19-25 nucleotide) antisense strand, followed by a 5-9 nucleotide loop, and the analogous sense strand. Alternatively, the sense strand may precede the nucleotide loop structure and the antisense strand may follow. These shRNAs may be contained in plasmids, retroviruses, and lentiviruses and expressed from, for example, the pol III U6 promoter, or another promoter (see, e.g., Stewart, et al. (2003) *RNA* April; 9(4):493-501 incorporated by reference herein). RNA interfering agents, e.g., siRNA molecules, may be administered to a subject having or at risk for having cancer, to inhibit expression of a marker gene of the inven-

tion, e.g., a marker gene which is overexpressed in cancer (such as the markers listed in Table 3) and thereby treat, prevent, or inhibit cancer in the subject.

As used herein, “subject” refers to any healthy animal, mammal or human, or any animal, mammal or human afflicted with a cancer, e.g., lung, ovarian, pancreatic, liver, breast, prostate, and colon carcinomas, as well as melanoma and multiple myeloma. The term “subject” is interchangeable with “subject”.

The language “substantially free of chemical precursors or other chemicals” includes preparations of antibody, polypeptide, peptide or fusion protein in which the protein is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. In one embodiment, the language “substantially free of chemical precursors or other chemicals” includes preparations of antibody, polypeptide, peptide or fusion protein having less than about 30% (by dry weight) of chemical precursors or non-antibody, polypeptide, peptide or fusion protein chemicals, more preferably less than about 20% chemical precursors or non-antibody, polypeptide, peptide or fusion protein chemicals, still more preferably less than about 10% chemical precursors or non-antibody, polypeptide, peptide or fusion protein chemicals, and most preferably less than about 5% chemical precursors or non-antibody, polypeptide, peptide or fusion protein chemicals.

As used herein, the term “survival” includes all of the following: survival until mortality, also known as overall survival (wherein said mortality may be either irrespective of cause or tumor related); “recurrence-free survival” (wherein the term recurrence shall include both localized and distant recurrence); metastasis free survival; disease free survival (wherein the term disease shall include cancer and diseases associated therewith). The length of said survival may be calculated by reference to a defined start point (e.g. time of diagnosis or start of treatment) and end point (e.g. death, recurrence or metastasis). In addition, criteria for efficacy of treatment can be expanded to include response to chemotherapy, probability of survival, probability of metastasis within a given time period, and probability of tumor recurrence.

The term “synergistic effect” refers to the combined effect of two or more anticancer agents can be greater than the sum of the separate effects of the anticancer agents or alone. In some embodiments, in can provide for similar efficacy of monotherapy but with other unexpected improvements relative to monotherapy, such as reducing unwanted side effects.

A “tissue-specific” promoter is a nucleotide sequence which, when operably linked with a polynucleotide which encodes or specifies a gene product, causes the gene product to be produced in a living human cell substantially only if the cell is a cell of the tissue type corresponding to the promoter.

A “transcribed polynucleotide” or “nucleotide transcript” is a polynucleotide (e.g. an mRNA, hnRNA, a cDNA, or an analog of such RNA or cDNA) which is complementary to or homologous with all or a portion of a mature mRNA made by transcription of a marker of the invention and normal post-transcriptional processing (e.g. splicing), if any, of the RNA transcript, and reverse transcription of the RNA transcript.

An “underexpression” or “significantly lower level of expression or copy number” of a marker refers to an expression level or copy number in a test sample that is greater than the standard error of the assay employed to assess expression or copy number, but is preferably at least twice, and more preferably three, four, five or ten or more

times less than the expression level or copy number of the marker in a control sample (e.g., sample from a healthy subject not afflicted with cancer) and preferably, the average expression level or copy number of the marker in several control samples.

As used herein, the term “vector” refers to a nucleic acid capable of transporting another nucleic acid to which it has been linked. One type of vector is a “plasmid”, which refers to a circular double stranded DNA loop into which additional DNA segments may be ligated. Another type of vector is a viral vector, wherein additional DNA segments may be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as “recombinant expression vectors” or simply “expression vectors”. In general, expression vectors of utility in recombinant DNA techniques are often in the form of plasmids. In the present specification, “plasmid” and “vector” may be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

There is a known and definite correspondence between the amino acid sequence of a particular protein and the nucleotide sequences that can code for the protein, as defined by the genetic code (shown below) Likewise, there is a known and definite correspondence between the nucleotide sequence of a particular nucleic acid and the amino acid sequence encoded by that nucleic acid, as defined by the genetic code.

GENETIC CODE

Alanine (Ala, A)	GCA, GCC, GCG, GCT
Arginine (Arg, R)	AGA, ACG, CGA, CGC, CGG, CGT
Asparagine (Asn, N)	AAC, AAT
Aspartic acid (Asp, D)	GAC, GAT
Cysteine (Cys, C)	TGC, TGT
Glutamic acid (Glu, E)	GAA, GAG
Glutamine (Gln, Q)	CAA, CAG
Glycine (Gly, G)	GGA, GGC, GGG, GGT
Histidine (His, H)	CAC, CAT
Isoleucine (Ile, I)	ATA, ATC, ATT
Leucine (Leu, L)	CTA, CTC, CTG, CTT, TTA, TTG
Lysine (Lys, K)	AAA, AAG
Methionine (Met, M)	ATG
Phenylalanine (Phe, F)	TTC, TTT
Proline (Pro, P)	CCA, CCC, CCG, CCT
Serine (Ser, S)	AGC, AGT, TCA, TCC, TCG, TCT
Threonine (Thr, T)	ACA, ACC, ACG, ACT
Tryptophan (Trp, W)	TGG
Tyrosine (Tyr, Y)	TAC, TAT
Valine (Val, V)	GTA, GTC, GTG, GTT
Termination signal (end)	TAA, TAG, TGA

An important and well known feature of the genetic code is its redundancy, whereby, for most of the amino acids used to make proteins, more than one coding nucleotide triplet may be employed (illustrated above). Therefore, a number of different nucleotide sequences may code for a given amino acid sequence. Such nucleotide sequences are considered functionally equivalent since they result in the

production of the same amino acid sequence in all organisms (although certain organisms may translate some sequences more efficiently than they do others). Moreover, occasionally, a methylated variant of a purine or pyrimidine may be found in a given nucleotide sequence. Such methylations do not affect the coding relationship between the trinucleotide codon and the corresponding amino acid.

In view of the foregoing, the nucleotide sequence of a DNA or RNA coding for a fusion protein or polypeptide of the invention (or any portion thereof) can be used to derive the fusion protein or polypeptide amino acid sequence, using the genetic code to translate the DNA or RNA into an amino acid sequence. Likewise, for fusion protein or polypeptide amino acid sequence, corresponding nucleotide sequences that can encode the fusion protein or polypeptide can be deduced from the genetic code (which, because of its redundancy, will produce multiple nucleic acid sequences for any given amino acid sequence). Thus, description and/or disclosure herein of a nucleotide sequence which encodes a fusion protein or polypeptide should be considered to also include description and/or disclosure of the amino acid sequence encoded by the nucleotide sequence. Similarly, description and/or disclosure of a fusion protein or polypeptide amino acid sequence herein should be considered to also include description and/or disclosure of all possible nucleotide sequences that can encode the amino acid sequence.

Finally, nucleic acid and amino acid sequence information for the loci and biomarkers of the present invention (e.g., biomarkers listed in Table 1) are well known in the art and readily available on publicly available databases, such as the National Center for Biotechnology Information (NCBI). For example, exemplary nucleic acid and amino acid sequences derived from publicly available sequence databases are provided below.

For example, the term "PD-1" refers to a member of the immunoglobulin gene superfamily that functions as a coinhibitory receptor having PD-L1 and PD-L2 as known ligands. PD-1 was previously identified using a subtraction cloning based approach to select for proteins involved in apoptotic cell death. PD-1 is a member of the CD28/CTLA-4 family of molecules based on its ability to bind to PD-L1. Like CTLA-4, PD-1 is rapidly induced on the surface of T-cells in response to anti-CD3 (Agata et al. (1996) *Int. Immunol.* 8:765). In contrast to CTLA-4, however, PD-1 is also induced on the surface of B-cells (in response to anti-IgM). PD-1 is also expressed on a subset of thymocytes and myeloid cells (Agata et al. (1996) *supra*; Nishimura et al. (1996) *Int. Immunol.* 8:773).

The nucleic acid and amino acid sequences of a representative human PD-1 biomarker is available to the public at the GenBank database under NM 005018.2 and NP 005009.2 and is shown in Table 1 (see also Ishida et al. (1992) 20 *EMBO J* 11:3887; Shinohara et al. (1994) *Genomics* 23:704; U.S. Pat. No. 5,698,520). PD-1 has an extracellular region containing immunoglobulin superfamily domain, a transmembrane domain, and an intracellular region including an immunoreceptor tyrosine-based inhibitory motif (ITIM) (Ishida et al. (1992) *EMBO J.* 11:3887; Shinohara et al. (1994) *Genomics* 23:704; and U.S. Pat. No. 5,698,520). These features also define a larger family of polypeptides, called the immunoinhibitory receptors, which also includes gp49B, PIR-B, and the killer inhibitory receptors (KIRs) (Vivier and Daron (1997) *Immunol. Today* 18:286). It is often assumed that the tyrosyl phosphorylated ITIM motif of these receptors interacts with SH2-domain containing phosphatases, which leads to inhibitory signals.

A subset of these immunoinhibitory receptors bind to MHC polypeptides, for example the KIRs, and CTLA4 binds to B7-1 and B7-2. It has been proposed that there is a phylogenetic relationship between the MHC and B7 genes (Henry et al. (1999) *Immunol. Today* 20(6):285-8). Nucleic acid and polypeptide sequences of PD-1 orthologs in organisms other than humans are well known and include, for example, mouse PD-1 (NM_008798.2 and NP_032824.1), rat PD-1 (NM_001106927.1 and NP_001100397.1), dog PD-1 (XM_543338.3 and XP_543338.3), cow PD-1 (NM_001083506.1 and NP_001076975.1), and chicken PD-1 (XM_422723.3 and XP_422723.2).

PD-1 polypeptides are inhibitory receptors capable of transmitting an inhibitory signal to an immune cell to thereby inhibit immune cell effector function, or are capable of promoting costimulation (e.g., by competitive inhibition) of immune cells, e.g., when present in soluble, monomeric form. Preferred PD-1 family members share sequence identity with PD-1 and bind to one or more B7 family members, e.g., B7-1, B7-2, PD-1 ligand, and/or other polypeptides on antigen presenting cells.

The term "PD-1 activity," includes the ability of a PD-1 polypeptide to modulate an inhibitory signal in an activated immune cell, e.g., by engaging a natural PD-1 ligand on an antigen presenting cell. PD-1 transmits an inhibitory signal to an immune cell in a manner similar to CTLA4. Modulation of an inhibitory signal in an immune cell results in modulation of proliferation of, and/or cytokine secretion by, an immune cell. Thus, the term "PD-1 activity" includes the ability of a PD-1 polypeptide to bind its natural ligand(s), the ability to modulate immune cell costimulatory or inhibitory signals, and the ability to modulate the immune response.

The term "PD-1 ligand" refers to binding partners of the PD-1 receptor and includes both PD-L1 (Freeman et al. (2000) *J. Exp. Med.* 192:1027) and PD-L2 (Latchman et al. (2001) *Nat. Immunol.* 2:261). At least two types of human PD-1 ligand polypeptides exist. PD-1 ligand proteins comprise a signal sequence, and an IgV domain, an IgC domain, a transmembrane domain, and a short cytoplasmic tail. Both PD-L1 (See Freeman et al. (2000) *J. Exp. Med.* 192:1027 for sequence data) and PD-L2 (See Latchman et al. (2001) *Nat. Immunol.* 2:261 for sequence data) are members of the B7 family of polypeptides. Both PD-L1 and PD-L2 are expressed in placenta, spleen, lymph nodes, thymus, and heart. Only PD-L2 is expressed in pancreas, lung and liver, while only PD-L1 is expressed in fetal liver. Both PD-1 ligands are upregulated on activated monocytes and dendritic cells, although PD-L1 expression is broader. For example, PD-L1 is known to be constitutively expressed and upregulated to higher levels on murine hematopoietic cells (e.g., T cells, B cells, macrophages, dendritic cells (DCs), and bone marrow-derived mast cells) and non-hematopoietic cells (e.g., endothelial, epithelial, and muscle cells), whereas PD-L2 is inducibly expressed on DCs, macrophages, and bone marrow-derived mast cells (see, Butte et al. (2007) *Immunity* 27:111).

PD-1 ligands comprise a family of polypeptides having certain conserved structural and functional features. The term "family" when used to refer to proteins or nucleic acid molecules, is intended to mean two or more proteins or nucleic acid molecules having a common structural domain or motif and having sufficient amino acid or nucleotide sequence homology, as defined herein. Such family members can be naturally or non-naturally occurring and can be from either the same or different species. For example, a family can contain a first protein of human origin, as well as other, distinct proteins of human origin or alternatively, can

contain homologues of non-human origin. Members of a family may also have common functional characteristics. PD-1 ligands are members of the B7 family of polypeptides. The term “B7 family” or “B7 polypeptides” as used herein includes costimulatory polypeptides that share sequence homology with B7 polypeptides, e.g., with B7-1 (CD80), B7-2 (CD86), inducible costimulatory ligand (ICOS-L), B7-H3, B7-H4, VISTA, B7-H6, B7h (Swallow et al. (1999) *Immunity* 11:423), and/or PD-1 ligands (e.g., PD-L1 or PD-L2). For example, human B7-1 and B7-2 share approximately 26% amino acid sequence identity when compared using the BLAST program at NCBI with the default parameters (Blosum62 matrix with gap penalties set at existence 11 and extension 1 (see the NCBI website)). The term B7 family also includes variants of these polypeptides which are capable of modulating immune cell function. The B7 family of molecules share a number of conserved regions, including signal domains, IgV domains and the IgC domains. IgV domains and the IgC domains are art-recognized Ig superfamily member domains. These domains correspond to structural units that have distinct folding patterns called Ig folds. Ig folds are comprised of a sandwich of two β sheets, each consisting of anti-parallel β strands of 5-10 amino acids with a conserved disulfide bond between the two sheets in most, but not all, IgC domains of Ig, TCR, and MHC molecules share the same types of sequence patterns and are called the C1-set within the Ig superfamily. Other IgC domains fall within other sets. IgV domains also share sequence patterns and are called V set domains. IgV domains are longer than IgC domains and contain an additional pair of β strands.

The term “PD-L1” refers to a specific PD-1 ligand. Two forms of human PD-L1 molecules have been identified. One form is a naturally occurring PD-L1 soluble polypeptide, i.e., having a short hydrophilic domain at the COOH-terminal end and no transmembrane domain, and is referred to herein as PD-L1S (shown in Table 1 as SEQ ID NO: 4). The second form is a cell-associated polypeptide, i.e., having a transmembrane and cytoplasmic domain, referred to herein as PD-L1M (shown in SEQ ID NO: 6). The nucleic acid and amino acid sequences of representative human PD-L1 biomarkers regarding PD-L1M are also available to the public at the GenBank database under NM_014143.3 and NP_054862.1. PD-L1 proteins comprise a signal sequence, and an IgV domain and an IgC domain. The signal sequence of SEQ ID NO: 4 is shown from about amino acid 1 to about amino acid 18. The signal sequence of SEQ ID NO: 6 is shown: from about amino acid 1 to about amino acid 18. The IgV domain of SEQ ID NO: 4 is shown from about amino acid 19 to about amino acid 134 and the IgV domain of SEQ ID NO: 6 is shown from about amino acid 19 to about amino acid 134. The IgC domain of SEQ ID NO: 4 is shown from about amino acid 135 to about amino acid 227 and the IgC domain of SEQ ID NO: 6 is shown from about amino acid 135 to about amino acid 227. The hydrophilic tail of the PD-L1 exemplified in SEQ ID NO: 4 comprises a hydrophilic tail shown from about amino acid 228 to about amino acid 245. The PD-L1 polypeptide exemplified in SEQ ID NO: 6 comprises a transmembrane domain shown from about amino acids 239 to about amino acid 259 of SEQ ID NO: 6 and a cytoplasmic domain shown of about 30 amino acids from 260 to about amino acid 290 of SEQ ID NO: 6. In addition, nucleic acid and polypeptide sequences of PD-L1 orthologs in organisms other than humans are well known and include, for example, mouse PD-L1 (NM_021893.3 and NP_068693.1), rat PD-L1 (NM_001191954.1 and NP_001178883.1), dog PD-L1

(XM_541302.3 and XP_541302.3), cow PD-L1 (NM_001163412.1 and NP_001156884.1), and chicken PD-L1 (XM_424811.3 and XP_424811.3).

The term “TIM-3” refers to a type I cell-surface glycoprotein that comprises an N-terminal immunoglobulin (Ig)-like domain, a mucin domain with O-linked glycosylations and with N-linked glycosylations close to the membrane, a single transmembrane domain, and a cytoplasmic region with tyrosine phosphorylation motif(s) (see, for example, U.S. Pat. Publ. 2013/0156774). TIM-3 is a member of the T cell/transmembrane, immunoglobulin, and mucin (TIM) gene family. Nucleic acid and polypeptide sequences of human TIM-3 are well known in the art and are publicly available, for example, as described in NM_032782.4 and NP_116171.3. The term, as described above for useful markers such as PD-L1 and PD-1, encompasses any naturally occurring allelic, splice variants, and processed forms thereof. Typically, TIM-3 refers to human TIM-3 and can include truncated forms or fragments of the TIM-3 polypeptide. In addition, nucleic acid and polypeptide sequences of TIM-3 orthologs in organisms other than humans are well known and include, for example, mouse TIM-3 (NM_134250.2 and NP_599011.2), chimpanzee TIM-3 (XM_518059.4 and XP_518059.3), dog TIM-3 (NM_001254715.1 and NP_001241644.1), cow TIM-3 (NM_001077105.2 and NP_001070573.1), and rat TIM-3 (NM_001100762.1 and NP_001094232.1). In addition, neutralizing anti-TIM-3 antibodies are well known in the art (see, at least U.S. Pat. Publ. 2013/0183688, Ngiow et al. (2011) *Cancer Res.* 71:3540-3551; and antibody 344823 from R&D Biosystems, as well as clones 2C23, 5D12, 2E2, 4A4, and IG5, which are all published and thus publicly available).

TIM-3 was originally identified as a mouse Th1-specific cell surface protein that was expressed after several rounds of in vitro Th1 differentiation, and was later shown to also be expressed on Th17 cells. In humans, TIM-3 is expressed on a subset of activated CD4+ T cells, on differentiated Th1 cells, on some CD8+ T cells, and at lower levels on Th17 cells (Hastings et al. (2009) *Eur. J. Immunol.* 39:2492-2501). TIM-3 is also expressed on cells of the innate immune system including mouse mast cells, subpopulations of macrophages and dendritic cells (DCs), NK and NKT cells, human monocytes, human dendritic cells, and on murine primary bronchial epithelial cell lines. TIM-3 expression is regulated by the transcription factor T-bet. TIM-3 can generate an inhibitory signal resulting in apoptosis of Th1 and Tc1 cells, and can mediate phagocytosis of apoptotic cells and cross-presentation of antigen. Polymorphisms in TIM-1 and TIM-3 can reciprocally regulate the direction of T-cell responses (Freeman et al. (2010) *Immunol. Rev.* 235:172-89).

TIM-3 has several known ligands, including galectin-9, phosphatidylserine, and HMGB1. For example, galectin-9 is an S-type lectin with two distinct carbohydrate recognition domains joined by a long flexible linker, and has an enhanced affinity for larger poly-N-acetyllactosamine-containing structures. Galectin-9 does not have a signal sequence and is localized in the cytoplasm. However, it can be secreted and exerts its function by binding to glycoproteins on the target cell surface via their carbohydrate chains (Freeman et al. (2010) *Immunol. Rev.* 235:172-89). Engagement of TIM-3 by galectin-9 leads to Th1 cell death and a consequent decline in IFN-gamma production. When given in vivo, galectin-9 had beneficial effects in several murine disease models, including an EAE model, a mouse model of arthritis, in cardiac and skin allograft transplant models, and

contact hypersensitivity and psoriatic models (Freeman et al. (2010) *Immunol. Rev.* 235:172-89). Residues important for TIM-3 binding to galectin-9 include TIM-3(44), TIM-3(74), and TIM-3(100), which undergo N- and/or O-glycosylation. It is also known that TIM-3 mediates T-cell dysfunction associated with chronic viral infections (Golden-Mason et al. (2009) *J. Virol.* 83:9122-9130; Jones et al. (2008) *J. Exp. Med.* 205:2763-2779) and increases HIV-1-specific T cell responses when blocked ex vivo (Golden-Mason et al. (2009) *J. Virol.* 83:9122-9130). In addition, in chronic HCV infection, TIM-3 expression was increased on CD4⁺ and CD8⁺ T cells, specifically HCV-specific CD8⁺ cytotoxic T cells (CTLs) in chronic HCV infection and treatment with a blocking monoclonal antibody to TIM-3 reversed HCV-specific T cell exhaustion (Jones et al. (2008) *J. Exp. Med.* 205:2763-2779).

The term "LAG-3," also known as CD223, refers to a member of the immunoglobulin supergene family and is structurally and genetically related to CD4 (see, U.S. Pat. Publ. 2011/0150892). LAG-3 is generally known as a membrane protein encoded by a gene located on the distal part of the short arm of chromosome 12, near the CD4 gene, suggesting that the LAG-3 gene may have evolved through gene duplication (Triebel et al. (1990) *J. Exp. Med.* 171:1393-1405). However, secreted forms of the protein are known (e.g., for human and mouse TIM-3). Nucleic acid and polypeptide sequences of human LAG-3 are well known in the art and are publicly available, for example, as described in NM_002286.5 and NP_002277.4.

The term encompasses any naturally occurring allelic, splice variants, and processed forms thereof. Typically, LAG-3 refers to human LAG-3 and can include truncated forms or fragments of the LAG-3 polypeptide. In addition, nucleic acid and polypeptide sequences of LAG-3 orthologs in organisms other than humans are well known and include, for example, mouse LAG-3 (NM_008479.2 and NP_032505.1), chimpanzee LAG-3 (XM_508966.4 and XP_508966.2), monkey LAG-3 (XM_001108923.2 and XP_001108923.1), cow LAG-3 (NM_00124949.1 and NP_001232878.1), rat LAG-3 (NM_212513.2 and NP_997678.2), and chicken LAG-3 (XM_416510.3, XP_416510.2, XM_004938117.1, and XP_004938174.1). In addition, neutralizing anti-LAG-3 antibodies are well known in the art (see, at least U.S. Pat. Publs. 2011/0150892 and 2010/0233183; Macon-Lemaitre and Triebel (2005) *Immunology* 115:170-178; Drake et al. (2006) *J. Clin. Oncol.* 24:2573; Richter et al. (2010) *Int. Immunol.* 22:13-23).

LAG-3 is not expressed on resting peripheral blood lymphocytes but is expressed on activated T cells and NK cells and has a number of functions (see, U.S. Pat. Publ. 2011/0150892). Similar to CD4, LAG-3 has been demonstrated to interact with MHC Class II molecules but, unlike CD4, LAG-3 does not interact with the human immunodeficiency virus gp120 protein (Baixeras et al. (1992) *J. Exp. Med.* 176:327-337). Studies using a soluble LAG-3 immunoglobulin fusion protein (sLAG-3Ig) demonstrated direct and specific binding of LAG-3 to MHC class II on the cell

surface (Huard et al. (1996) *Eur. J. Immunol.* 26:1180-1186). In in vitro studies of antigen-specific T cell responses, the addition of anti-LAG-3 antibodies led to increased T cell proliferation and higher expression of activation antigens such as CD25, supporting a role for the LAG-/MHC class II interaction in down-regulating antigen-dependent stimulation of CD4⁺T lymphocytes (Huard et al. (1994) *Eur. J. Immunol.* 24:3216-3221). The intra-cytoplasmic region of LAG-3 has been demonstrated to interact with a protein termed LAP, which is thought to be a signal transduction molecule involved in the downregulation of the CD3/TCR activation pathway (Iouzalén et al. (2001) *Eur. J. Immunol.* 31:2885-2891). Furthermore, CD4⁺CD25⁺ regulatory T cells (T_{reg}) have been shown to express LAG-3 upon activation and antibodies to LAG-3 inhibit suppression by induced regulatory T cells, both in vitro and in vivo, suggesting that LAG-3 contributes to the suppressor activity of regulatory T cells (Huang et al. (2004) *Immunity* 21:503-513). Still further, LAG-3 has been shown to negatively regulate T cell homeostasis by regulatory T cells in both T cell-dependent and independent mechanisms (Workman and Vignali (2005) *J. Immunol.* 174:688-695).

In certain circumstances, LAG-3 also has been shown to have immunostimulatory effects. For example, LAG-3 transfected tumor cells transplanted into syngeneic mice showed marked growth reduction or complete regression as compared to untransfected tumor cells, suggesting that LAG-3 expression on the tumor cells stimulated an anti-tumor response by triggering antigen presenting cells via MHC class II molecules (Prigent et al. (1999) *Eur. J. Immunol.* 29:3867-3876). Additionally, soluble LAG-3 Ig fusion protein has been shown to stimulate both humoral and cellular immune responses when administered to mice together with an antigen, indicating that soluble LAG-3Ig can function as a vaccine adjuvant (El Mir and Triebel (2000) *J. Immunol.* 164:5583-5589). Furthermore, soluble human LAG-3Ig has been shown to amplify the in vitro generation of type I tumor-specific immunity (Casati et al. (2006) *Cancer Res.* 66:4450-4460). The functional activity of LAG-3 is reviewed further in Triebel (2003) *Trends Immunol.* 24:619-622.

CTLA-4 is a T cell surface molecule that was originally identified by differential screening of a murine cytolytic T cell cDNA library, Brunet et al. (1987) *Nature* 328:267-270. The role of CTLA-4 as a second receptor for B7 is discussed in Linsley et al. (1991) *J. Exp. Med.* 174:561-569. Freeman et al. (1993) *Science* 262:907-909 discusses CTLA-4 in B7 deficient mice. Ligands for CTLA-4 are described in Lenschow et al. (1993) P.N.A.S. 90:11054-11058. Linsley et al. (1992) *Science* 257:792-795 describes immunosuppression in vivo by a soluble form of CTLA-4. Lenschow et al. (1992) *Science* 257:789-792 discusses long term survival of pancreatic islet grafts induced by CTLA-4Ig. It is suggested in Walunas et al. (1994) *Immunity* 1:405-413, that CTLA-4 can function as a negative regulator of T cell activation. The amino acid and nucleotide sequence of CTLA-4 (e.g., human CTLA-4) are known in the art (e.g., as described in U.S. Pat. Nos. 5,811,097 and 5,434,131, incorporated herein by reference).

TABLE 1

SEQ ID NO: 1 Human PD-1 cDNA Sequence

cactctgggtg gggctgctcc aggc atg cag atc cca cag gcg ccc tgg cca 51
 Met Gln Ile Pro Gln Ala Pro Trp Pro
 1 5

TABLE 1-continued

gtc gtc tgg gcg gtg cta caa ctg ggc tgg cgg cca gga tgg ttc tta	99
Val Val Trp Ala Val Leu Gln Leu Gly Trp Arg Pro Gly Trp Phe Leu	
10 15 20 25	
gac tcc cca gac agg ccc tgg aac ccc ccc acc ttc tcc cca gcc ctg	147
Asp Ser Pro Asp Arg Pro Trp Asn Pro Pro Thr Phe Ser Pro Ala Leu	
30 35 40	
ctc gtg gtg acc gaa ggg gac aac gcc acc ttc acc tgc agc ttc tcc	195
Leu Val Val Thr Glu Gly Asp Asn Ala Thr Phe Thr Cys Ser Phe Ser	
45 50 55	
aac aca tcg gag agc ttc gtg cta aac tgg tac cgc atg agc ccc agc	243
Asn Thr Ser Glu Ser Phe Val Leu Asn Trp Tyr Arg Met Ser Pro Ser	
60 65 70	
aac cag acg gac aag ctg gcc gcc ttc ccc gag gac cgc agc cag ccc	291
Asn Gln Thr Asp Lys Leu Ala Ala Phe Pro Glu Asp Arg Ser Gln Pro	
75 80 85	
ggc cag gac tgc cgc ttc cgt gtc aca caa ctg ccc aac ggg cgt gac	339
Gly Gln Asp Cys Arg Phe Arg Val Thr Gln Leu Pro Asn Gly Arg Asp	
90 95 100 105	
ttc cac atg agc gtg gtc agg gcc cgg cgc aat gac agc ggc acc tac	387
Phe His Met Ser Val Val Arg Ala Arg Asn Asp Ser Gly Thr Tyr	
110 115 120	
ctc tgt ggg gcc atc tcc ctg gcc ccc aag gcg cag atc aaa gag agc	435
Leu Cys Gly Ala Ile Ser Leu Ala Pro Lys Ala Gln Ile Lys Glu Ser	
125 130 135	
ctg cgg gca gag ctc agg gtg aca gag aga agg gca gaa gtg ccc aca	483
Leu Arg Ala Glu Leu Arg Val Thr Glu Arg Arg Ala Glu Val Pro Thr	
140 145 150	
gcc cac ccc agc ccc tca ccc agg tca gcc ggc cag ttc caa acc ctg	531
Ala His Pro Ser Pro Ser Pro Arg Ser Ala Gly Gln Phe Gln Thr Leu	
155 160 165	
gtg gtt ggt gtc gtg ggc ggc ctg ctg ggc agc ctg gtg ctg cta gtc	579
Val Val Gly Val Val Gly Gly Leu Leu Gly Ser Leu Val Leu Leu Val	
170 175 180 185	
tgg gtc ctg gcc gtc atc tgc tcc cgg gcc gca cga ggg aca ata gga	627
Trp Val Leu Ala Val Ile Cys Ser Arg Ala Ala Arg Gly Thr Ile Gly	
190 195 200	
gcc agg cgc acc ggc cag ccc ctg aag gag gac ccc tca gcc gtg cct	675
Ala Arg Arg Thr Gly Gln Pro Leu Lys Glu Asp Pro Ser Ala Val Pro	
205 210 215	
gtg ttc tct gtg gac tat ggg gag ctg gat ttc cag tgg cga gag aag	723
Val Phe Ser Val Asp Tyr Gly Glu Leu Asp Phe Gln Trp Arg Glu Lys	
220 225 230	
acc ccg gag ccc ccc gtg ccc tgt gtc cct gag cag acg gag tat gcc	771
Thr Pro Glu Pro Pro Val Pro Cys Val Pro Glu Gln Thr Glu Tyr Ala	
235 240 245	
acc att gtc ttt cct agc gga atg ggc acc tca tcc ccc gcc cgc agg	819
Thr Ile Val Phe Pro Ser Gly Met Gly Thr Ser Ser Pro Ala Arg Arg	
250 255 260 265	
ggc tca gct gac ggc cct cgg agt gcc cag cca ctg agg cct gag gat	867
Gly Ser Ala Asp Gly Pro Arg Ser Ala Gln Pro Leu Arg Pro Glu Asp	
270 275 280	
gga cac tgc tct tgg ccc ctc tgaccggcctt ccttggccac cagtgtctg cag	921
Gly His Cys Ser Trp Pro Leu	
285	

SEQ ID NO: 2 Human PD-1 Amino Acid Sequence

Met Gln Ile Pro Gln Ala Pro Trp Pro Val Val Trp Ala Val Leu Gln	
1 5 10 15	
Leu Gly Trp Arg Pro Gly Trp Phe Leu Asp Ser Pro Asp Arg Pro Trp	
20 25 30	

TABLE 1-continued

Asn	Pro	Pro	Thr	Phe	Ser	Pro	Ala	Leu	Leu	Val	Val	Thr	Glu	Gly	Asp
		35					40					45			
Asn	Ala	Thr	Phe	Thr	Cys	Ser	Phe	Ser	Asn	Thr	Ser	Glu	Ser	Phe	Val
	50					55					60				
Leu	Asn	Trp	Tyr	Arg	Met	Ser	Pro	Ser	Asn	Gln	Thr	Asp	Lys	Leu	Ala
	65				70					75					80
Ala	Phe	Pro	Glu	Asp	Arg	Ser	Gln	Pro	Gly	Gln	Asp	Cys	Arg	Phe	Arg
				85					90					95	
Val	Thr	Gln	Leu	Pro	Asn	Gly	Arg	Asp	Phe	His	Met	Ser	Val	Val	Arg
			100					105					110		
Ala	Arg	Arg	Asn	Asp	Ser	Gly	Thr	Tyr	Leu	Cys	Gly	Ala	Ile	Ser	Leu
		115					120					125			
Ala	Pro	Lys	Ala	Gln	Ile	Lys	Glu	Ser	Leu	Arg	Ala	Glu	Leu	Arg	Val
	130					135					140				
Thr	Glu	Arg	Arg	Ala	Glu	Val	Pro	Thr	Ala	His	Pro	Ser	Pro	Ser	Pro
	145				150					155					160
Arg	Ser	Ala	Gly	Gln	Phe	Gln	Thr	Leu	Val	Val	Gly	Val	Val	Gly	Gly
			165						170					175	
Leu	Leu	Gly	Ser	Leu	Val	Leu	Leu	Val	Trp	Val	Leu	Ala	Val	Ile	Cys
			180					185					190		
Ser	Arg	Ala	Ala	Arg	Gly	Thr	Ile	Gly	Ala	Arg	Arg	Thr	Gly	Gln	Pro
		195					200					205			
Leu	Lys	Glu	Asp	Pro	Ser	Ala	Val	Pro	Val	Phe	Ser	Val	Asp	Tyr	Gly
	210					215					220				
Glu	Leu	Asp	Phe	Gln	Trp	Arg	Glu	Lys	Thr	Pro	Glu	Pro	Pro	Val	Pro
	225				230					235					240
Cys	Val	Pro	Glu	Gln	Thr	Glu	Tyr	Ala	Thr	Ile	Val	Phe	Pro	Ser	Gly
			245						250					255	
Met	Gly	Thr	Ser	Ser	Pro	Ala	Arg	Arg	Gly	Ser	Ala	Asp	Gly	Pro	Arg
		260					265						270		
Ser	Ala	Gln	Pro	Leu	Arg	Pro	Glu	Asp	Gly	His	Cys	Ser	Trp	Pro	Leu
		275					280					285			

SEQ ID NO: 3 Human PD-L1S cDNA Acid Sequence

gcttcccagag gctccgcacc agccgcgctt ctgtccgcct gcagggcatt ccagaaag	58
atg agg ata ttt gct gtc ttt ata ttc atg acc tac tgg cat ttg ctg	106
Met Arg Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu	
1 5 10 15	
aac gca ttt act gtc acg gtt ccc aag gac cta tat gtg gta gag tat	154
Asn Ala Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr	
20 25 30	
ggt agc aat atg aca att gaa tgc aaa ttc cca gta gaa aaa caa tta	202
Gly Ser Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu	
35 40 45	
gac ctg gct gca cta att gtc tat tgg gaa atg gag gat aag aac att	250
Asp Leu Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile	
50 55 60	
att caa ttt gtg cat gga gag gaa gac ctg aag gtt cag cat agt agc	298
Ile Gln Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser	
65 70 75 80	
tac aga cag agg gcc cgg ctg ttg aag gac cag ctc tcc ctg gga aat	346
Tyr Arg Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn	
85 90 95	
gct gca ctt cag atc aca gat gtg aaa ttg cag gat gca ggg gtg tac	394
Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr	
100 105 110	

TABLE 1-continued

cgc tgc atg atc agc tat ggt ggt gcc gac tac aag cga att act gtg	442
Arg Cys Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val	
115 120 125	
aaa gtc aat gcc cca tac aac aaa atc aac caa aga att ttg gtt gtg	490
Lys Val Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val	
130 135 140	
gat cca gtc acc tct gaa cat gaa ctg aca tgt cag gct gag ggc tac	538
Asp Pro Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr	
145 150 155 160	
ccc aag gcc gaa gtc atc tgg aca agc agt gac cat caa gtc ctg agt	586
Pro Lys Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser	
165 170 175	
ggt aag acc acc acc acc aat tcc aag aga gag gag aag ctt ttc aat	634
Gly Lys Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn	
180 185 190	
gtg acc agc aca ctg aga atc aac aca aca act aat gag att ttc tac	682
Val Thr Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr	
195 200 205	
tgc act ttt agg aga tta gat cct gag gaa aac cat aca gct gaa ttg	730
Cys Thr Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu	
210 215 220	
gtc atc cca ggt aat att ctg aat gtg tcc att aaa ata tgt cta aca	778
Val Ile Pro Gly Asn Ile Leu Asn Val Ser Ile Lys Ile Cys Leu Thr	
225 230 235 240	
ctg tcc cct agc acc tagcatgatg tctgcctatc atagtcattc agtgattggt	833
Leu Ser Pro Ser Thr	
245	
gaataaatga atgaatgaat aacactatgt ttacaaaata tatectaatt cctcacctcc	893
atcattccaa accatattgt tacttaataa acattcagca gatatttatg gaataaaaaa	953
aaaaaaaaa aaaaa	968

SEQ ID NO: 4 Human PD-L1S Amino Acid Sequence

Met Arg Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu	
1 5 10 15	
Asn Ala Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr	
20 25 30	
Gly Ser Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu	
35 40 45	
Asp Leu Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile	
50 55 60	
Ile Gln Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser	
65 70 75 80	
Tyr Arg Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn	
85 90 95	
Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr	
100 105 110	
Arg Cys Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val	
115 120 125	
Lys Val Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val	
130 135 140	
Asp Pro Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr	
145 150 155 160	
Pro Lys Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser	
165 170 175	
Gly Lys Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn	
180 185 190	

TABLE 1-continued

Val	Thr	Ser	Thr	Leu	Arg	Ile	Asn	Thr	Thr	Thr	Asn	Glu	Ile	Phe	Tyr
		195					200					205			
Cys	Thr	Phe	Arg	Arg	Leu	Asp	Pro	Glu	Glu	Asn	His	Thr	Ala	Glu	Leu
	210					215					220				
Val	Ile	Pro	Gly	Asn	Ile	Leu	Asn	Val	Ser	Ile	Lys	Ile	Cys	Leu	Thr
225					230					235					240
Leu	Ser	Pro	Ser	Thr											
				245											

SEQ ID NO: 5 Human PD-L1M cDNA Acid Sequence

cgaggctccg caccagccgc gcttctgtcc gctgcaggg cattccagaa agatgagg	58
	Met Arg
	1
ata ttt gct gtc ttt ata ttc atg acc tac tgg cat ttg ctg aac gca	106
Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu Asn Ala	
	5 10 15
ttt act gtc acg gtt ccc aag gac cta tat gtg gta gag tat ggt agc	154
Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr Gly Ser	
	20 25 30
aat atg aca att gaa tgc aaa ttc cca gta gaa aaa caa tta gac ctg	202
Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu Asp Leu	
	35 40 45 50
gct gca cta att gtc tat tgg gaa atg gag gat aag aac att att caa	250
Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile Ile Gln	
	55 60 65
ttt gtg cat gga gag gaa gac ctg aag gtt cag cat agt agc tac aga	298
Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser Tyr Arg	
	70 75 80
cag agg gcc cgg ctg ttg aag gac cag ctc tcc ctg gga aat gct gca	346
Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn Ala Ala	
	85 90 95
ctt cag atc aca gat gtg aaa ttg cag gat gca ggg gtg tac cgc tgc	394
Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr Arg Cys	
	100 105 110
atg atc agc tat ggt ggt gcc gac tac aag cga att act gtg aaa gtc	442
Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val Lys Val	
	115 120 125 130
aat gcc cca tac aac aaa atc aac caa aga att ttg gtt gtg gat cca	490
Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val Asp Pro	
	135 140 145
gtc acc tct gaa cat gaa ctg aca tgt cag gct gag ggc tac ccc aag	538
Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr Pro Lys	
	150 155 160
gcc gaa gtc atc tgg aca agc agt gac cat caa gtc ctg agt ggt aag	586
Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser Gly Lys	
	165 170 175
acc acc acc acc aat tcc aag aga gag gag aag ctt ttc aat gtg acc	634
Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn Val Thr	
	180 185 190
agc aca ctg aga atc aac aca aca act aat gag att ttc tac tgc act	682
Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr Cys Thr	
	195 200 205 210
ttt agg aga tta gat cct gag gaa aac cat aca gct gaa ttg gtc atc	730
Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu Val Ile	
	215 220 225
cca gaa cta cct ctg gca cat cct cca aat gaa agg act cac ttg gta	778
Pro Glu Leu Pro Leu Ala His Pro Pro Asn Glu Arg Thr His Leu Val	
	230 235 240
att ctg gga gcc atc tta tta tgc ctt ggt gta gca ctg aca ttc atc	826
Ile Leu Gly Ala Ile Leu Leu Cys Leu Gly Val Ala Leu Thr Phe Ile	

TABLE 1-continued

245	250	255	
ttc cgt tta aga aaa ggg aga atg atg gat gtg aaa aaa tgt ggc atc			874
Phe Arg Leu Arg Lys Gly Arg Met Met Asp Val Lys Lys Cys Gly Ile			
260	265	270	
caa gat aca aac tca aag aag caa agt gat aca cat ttg gag gag acg			922
Gln Asp Thr Asn Ser Lys Lys Gln Ser Asp Thr His Leu Glu Glu Thr			
275	280	285	290
taatccagca ttggaacttc tgatcttcaa gcagggatc tcaacctgtg gtttaggggt			982
tcacgggggc tgagcgtgac aagaggaagg aatgggcccg tgggatgcag gcaatgtggg			1042
acttaaaagg cccaagcact gaaaatggaa cctggcgaaa gcagaggagg agaatgaaga			1102
aagatggagt caaacagga gcctggaggg agaccttgat actttcaaat gcctgagggg			1162
ctcatcgacg cctgtgacag ggagaaagga tacttctgaa caaggagcct ccaagcaaat			1222
catccattgc tcaccttagg aagacggggt gagaatccct aatttgaggg tcagttcctg			1282
cagaagtgcc ctttgccctcc actcaatgcc tcaatttggt ttctgcatga ctgagagtct			1342
cagtgttga acgggacagt atttatgtat gaggttttcc tatttatattt gaggctgtga			1402
ggtcttcttg tcagtgtagt gtggttgtga atgatttctt ttgaagatat attgtagtag			1462
atgttacaat tttgtcgcca aactaaactt gctgcttaat gatttgctca catctagtaa			1522
aacatggagt atttgtaaaa aaaaaaaaaa a			1553

SEQ ID NO: 6 Human PD-L1M Amino Acid Sequence

Met Arg Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu	1	5	10	15	
Asn Ala Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr		20	25	30	
Gly Ser Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu		35	40	45	
Asp Leu Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile		50	55	60	
Ile Gln Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser		65	70	75	80
Tyr Arg Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn		85	90	95	
Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr		100	105	110	
Arg Cys Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val		115	120	125	
Lys Val Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val		130	135	140	
Asp Pro Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr		145	150	155	160
Pro Lys Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser		165	170	175	
Gly Lys Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn		180	185	190	
Val Thr Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr		195	200	205	
Cys Thr Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu		210	215	220	
Val Ile Pro Glu Leu Pro Leu Ala His Pro Pro Asn Glu Arg Thr His		225	230	235	240
Leu Val Ile Leu Gly Ala Ile Leu Leu Cys Leu Gly Val Ala Leu Thr					

TABLE 1-continued

245			250			255									
Phe	Ile	Phe	Arg	Leu	Arg	Lys	Gly	Arg	Met	Met	Asp	Val	Lys	Lys	Cys
			260					265				270			
Gly	Ile	Gln	Asp	Thr	Asn	Ser	Lys	Lys	Gln	Ser	Asp	Thr	His	Leu	Glu
			275					280				285			
Glu	Thr														
	290														

SEQ ID NO: 7 Mouse PD-L1 cDNA Sequence

1 atgaggatat ttgctggcat tatattcaca gcctgctgctc acttgctacg ggcgtttact
 61 atcacggctc caaaggactt gtacgtgggtg gagtatggca gcaacgtcac gatggagtgc
 121 agattccctg tagaacggga gctggacctg cttgcgcttag tgggtactg ggaaaaggaa
 181 gatgagcaag tgattcagtt tgtggcagga gaggaggacc ttaagcctca gcacagcaac
 241 ttcaggggga gagcctcgct gccaaaggac cagcttttga agggaaatgc tgcccttcag
 301 atcacagacg tcaagctgca ggacgcaggc gtttactgct gcataatcag ctacggtggt
 361 gcgagctaca agcgaatcac gctgaaagtc aatgccccat accgcaaat caaccagaga
 421 atttccgtgg atccagccac ttctgagcat gaactaatat gtcaggccga gggttatcca
 481 gaagctgagg taatctggac aaacagtgc caccaacccg tgagtgggaa gagaagtgtc
 541 accacttccc ggacagaggg gatgcttctc aatgtgacca gcagtctgag ggtcaacgcc
 601 acagcgaatg atgttttcta ctgtacgttt tggagatcac agccagggca aaaccacaca
 661 gcgagctga tcatcccaga actgcctgca acacatcctc cacagaacag gactcactgg
 721 gtgcttctgg gatccatcct gttgttctc attgtagtgt ccacggtcct cctcttcttg
 781 agaaaacaag tgagaatgct agatgtggag aaatgtggcg ttgaagatac aagctcaaaa
 841 aaccgaaatg atacacaatt cgaggagacg taa

SEQ ID NO: 8 Mouse PD-L1 Amino Acid Sequence

1 mrifagiift acchllraft itapkdlyvv eygsnvtmec rfpvereldl lalvvyweke
 61 deqvivfvag eedlqpqhsn frgraslpkd qlkgnaalq itdvklqdag vycciiisygg
 121 adykritlkv napyrkinqr isvdpatseh elicqaegyp eaeviwtnsd hqpvsgkrsv
 181 ttsrtegmll nvtsslrva tandvfycf wsrqpgqnh taeliipelpa thppqnrthw
 241 vllgsillfl ivvstvlfl rkqvrml dve kcgvedtssk nrndtqfeet

SEQ ID NO: 9 Human TIM-3 cDNA Sequence

1 atgttttcac atcttccctt tgactgtgctc ctgctgctgc tgctgctact acttacaagg
 61 tccctcagaag tggaatacag agcggaggtc ggtcagaatg cctatctgcc ctgcttctac
 121 accccagccg cccagggaa cctcgtgcc gtctgctggg gcaaaggagc ctgtcctgtg
 181 tttgaatgtg gcaacgtggt gctcaggact gatgaaagg atgtgaatta ttggacatcc
 241 agatactggc taaatgggga tttccgcaa ggagatgtgt cctgaccat agagaatgtg
 301 actctagcag acagtgggat ctactgctgc cggatccaaa tcccaggcat aatgaatgat
 361 gaaaaattta acctgaagtt ggtcatcaaa ccagccaagg tcaccctgc accgactcgg
 421 cagagagact tctctgcagc ctttccaagg atgcttacca ccaggggaca tggcccagca
 481 gagacacaga cactggggag cctccctgat ataaatctaa cacaaatc cacattggcc
 541 aatgagttac gggactctag attggccaat gacttacggg actctggagc aaccatcaga
 601 ataggcatct acatcggagc agggatctgt gctgggctgg ctctggctct tatcttcggc
 661 gctttaattt tcaaatggta ttctcatagc aaagagaaga tacagaattt aagcctcatc

TABLE 1-continued

721 tctttggcca acctccctcc ctcaggattg gcaaatgcag tagcagaggg aattcgctca
 781 gaagaaaaca tctataccat tgaagagaac gtatatgaag tggaggagcc caatgagtat
 841 tattgctatg tcagcagcag gcagcaaccc tcacaacctt tgggttgctg ctttgcaatg
 901 ccatag

SEQ ID NO: 10 Human TIM-3 Amino Acid Sequence

1 mfshlpfdcv lllllllltr sseveyraev gqnaylpcfy tpaapgnlvp vewgkgacpv
 61 fecgnvvlrt derdvnywts rywlngdfrk gdvshtienv tladsgiycc riqipgimnd
 121 ekfnlklvik pakvtpaptr qrdftaafpr mltrrhgpa etqtlgslpd initqistla
 181 nelrdsrlan dlrdsgatir igiyigagic aglallalifg alifkwysys kekiqnslsi
 241 slanlppsgl anavaegirs eeniytieen vyeveepney ycyvssrqqp sqplgcrfam
 301 p

SEQ ID NO: 11 Mouse TIM-3 cDNA Sequence

1 atgttttcag gtcttacct caactgtgct ctgctgctgc tgcaactact acttgcaagg
 61 tcattggaaa atgcttatgt gtttgaggtt ggtaagaatg cctatctgcc ctgcagttac
 121 actctatcta cacctggggc acttgctgct atgtgctggg gcaagggatt ctgtccttgg
 181 tcacagtgta ccaacgagtt gctcagaact gatgaaagaa atgtgacata tcagaaatcc
 241 agcagatacc agctaaaggc cgatctcaac aaaggagacg tgtctctgat cataaagaat
 301 gtgactctgg atgaccatgg gacctactgc tgcaggatac agttccctgg tcttatgaat
 361 gataaaaaat tagaactgaa attagacatc aaagcagcca aggtcactcc agctcagact
 421 gcccatgggg actctactac agcttctcca agaaccctaa ccacggagag aatgggttca
 481 gagacacaga cactggtgac cctccataat aacaatggaa caaaaatttc cacatgggct
 541 gatgaaatta aggactctgg agaaacgatc agaactgcta tccacattgg agtgggagtc
 601 tctgctgggt tgacctggc acttatcatt ggtgtcttaa tctttaaag gtattcctgt
 661 aagaaaaaga agttatcgag tttgagcctt attacactgg ccaacttgcc tccaggaggg
 721 ttggcaaatg caggagcagt caggattcgc tctgaggaaa atatctacac catcgaggag
 781 aacgtatatg aagtggagaa ttcaaatgag tactactgct acgtcaacag ccagcagcca
 841 tctga

SEQ ID NO: 12 Mouse TIM-3 Amino Acid Sequence

1 mfsqtlncv llllqlllar slenayvfev gknaylpcsy tlstpgalvp mcwgkgfcpw
 61 sqctnellrt dernvtyqks sryqlkgdln kgdvsliikn vtlddhgtyc criqfpglmn
 121 dkklelklldi kaakvtpaqt ahgdsttasp rtltternngs etqtiavlhn nngtkistwa
 181 deikdsgeti rtaihigvgv sagltlalii gvlilkwysc kkkklssls1 itlanlppgg
 241 lanagavrir seeniytiee nvyevensne ycyvnsqqp s

SEQ ID NO: 13 Human LAG-3 cDNA Sequence

1 atgtgggagg ctcagttcct gggcttgctg tttctgcagc cgctttgggt ggctccagtg
 61 aagcctctcc agccaggggc tgaggctccc gtgggtgtggg cccaggaggg ggctcctgcc
 121 cagctcccct gcagccccac aatccccctc caggatctca gccttctgag aagagcaggg
 181 gtcacttggc agcatcagcc agacagtggc ccgcccgtg ccgccccgg ccaccccctg
 241 gccccgggc ctcacccggc ggcgcctcc tctggggggc ccaggccccg ccgctacacg
 301 gtgctgagcg tgggtcccgg aggctgcgc agcgggaggc tgcccctgca gccccgcgtc

TABLE 1-continued

361 cagctggatg agcgcggccg gcagcggcgg gacttctcgc tatggctgcg cccagcccgg
 421 cgcgccgacg ccggcgagta ccgcgcccg gtgcacctca gggaccgcgc cctctcctgc
 481 cgctccgctc tgcgctggg ccaggcctcg atgactgcca gccccccagg atctctcaga
 541 gcctccgact gggtcatttt gaactgctcc ttcagccgcc ctgaccgccc agcctctgtg
 601 cattgggtcc ggaaccgggg ccagggccga gtccctgtcc gggagtcccc ccatcaccac
 661 ttagcggaaa gcttctctt cctgccccaa gtcagcccca tggactctgg gccctggggc
 721 tgcacacctc cctacagaga tggcttcaac gtctccatca tgtataacct cactgttctg
 781 ggtctggagc ccccaactcc cttgacagtg tacgctggag caggttccag ggtggggctg
 841 ccctgccgcc tgctgctgg tgtggggacc cggctcttcc tcaactgcaa gtggactcct
 901 cctgggggag gccctgacct cctggtgact ggagacaatg gcgactttac ccttcgacta
 961 gaggatgtga gccaggccca ggctgggacc tacacctgcc atatccatct gcaggaacag
 1021 cagctcaatg ccaactgtcac attggcaatc atcacagtga ctcccaaate ctttgggtca
 1081 cctggatccc tggggaagct gctttgtgag gtgactccag tatctggaca agaacgcttt
 1141 gtgtggagct ctctggacac cccatcccag aggagtctt caggaccttg gctggaggca
 1201 caggaggccc agctccttcc ccagccttgg caatgccagc tgtaccaggg ggagaggctt
 1261 cttggagcag cagtgtactt cacagagctg tctagcccag gtgcccacg ctctgggaga
 1321 gccccagggtg ccctcccagc aggccacctc ctgctgttcc tcatccttgg tgcctttct
 1381 ctgctccttt tggtagctgg agcctttggc tttcaccttt ggagaagaca gtggcgacca
 1441 agacgatttt ctgccttaga gcaagggatt caccctccgc aggctcagag caagatagag
 1501 gagctggagc aagaaccgga gccggagccg gagccggaac cggagcccga gcccagccc
 1561 gagccggagc agctctga

SEQ ID NO: 14 Human LAG-3 Amino Acid Sequence

1 mweaqflgll flqplwvapv kplqpgaevp vvwagegapa qlpcspitpl qdlsllrrag
 61 vtwhqhpdsq ppaaapghpl apgphpaaps swgprprryt vlsvpggplr sgrlplqpry
 121 qldergrqrg dfsllwrpar radageyraa vhlrdralsc rlrlrlgqas mtasppgslr
 181 asdwilncs fsrpdrrpasv hwfrnrgqgr vpvresphhh laesflflpq vspmdsgpwg
 241 ciltyrdgfn vsimynitvl glepptpltv yagagsrvgl perlpagvgt rsfltakwtp
 301 pgggpdllvt gdnqdftrl edvsqaqagt ytchihlqeq qlnatvtlai itvtpksfgs
 361 pgslgkllce vtpvsgqerf vwssldtspq rsfsgpwlea qeaqlsqpw qcqlyggerl
 421 lgaavyftel sspgaqrsgr apgalpaghl llflilgvls llllvtgafg fhlwrrqwrp
 481 rrfaleqgi hppqaqskie eleqepepep epepepepep epeql

SEQ ID NO: 15 Mouse LAG-3 cDNA Sequence

1 atgagggagg acctgctcct tggctttttg cttctgggac tgctttggga agctccagtt
 61 gtgtcttcag gcctgggaa agagctcccc gtgggtgtggg cccaggaggg agctcccgtc
 121 catcttcctc gcagcctcaa atcccccaac ctggatccta actttctacg aagaggaggg
 181 gttatctggc aacatcaacc agacagtggc caaccactc ccatcccggc ccttgacctt
 241 caccagggga tgcctcggc tagacaacct gcaccgggtc gctacacggt gctgagcgtg
 301 gctccaggag gctgcgagc cgggaggcag cccctgcatc cccacgtgca gctggaggag
 361 cgcgccctcc agcgcgggga cttctctctg tggttgcgcc cagctctgag caccgatgag
 421 ggcgagtacc acgccaccgt gcgctcccg aaccgcgccc tctcctgag tctccgctg

TABLE 1-continued

```

481 cgcgctggcc aggcctcgat gattgctagt ccctcaggag tectcaagct gtctgattgg
541 gtccttttga actgctcctt cagccgtcct gaccgcccag tctctgtgca ctggttccag
601 ggccagaacc gaggcctgt ctacaactca ccgcgtcatt ttttagctga aactttcctg
661 ttactgcccc aagtcagccc cctggactct gggacctggg gctgtgtcct cacctacaga
721 gatggcttca atgtctccat cacgtacaac ctcaaggttc tgggtctgga gcccgtagcc
781 cctctgacag tgtacgtgc tgaaggttct aggggtggagc tgcctgtca tttgccccca
841 ggagtgggga ccccttcttt gctcattgcc aagtggactc ctctggagg aggtcctgag
901 ctccccgtgg ctggaaagag tggcaatttt acccttcacc ttgaggctgt gggctctggca
961 caggctggga cctacacctg tagcatccat ctgcaggac agcagctcaa tgccactgtc
1021 acgttggcgg tcatcacagt gactcccaaa tccttcgggt tacctggctc ccggggggaag
1081 ctggtgtgtg agtaacccc ggcattctga aaggaaagat ttgtgtggcg tcccctgaac
1141 aatctgtcca ggagttgccc gggcctgtg ctggagattc aggaggccag gctccttget
1201 gagcgatggc agtgtcagct gtacgagggc cagaggcttc ttggagcgac agtgtacgcc
1261 gcagagtcta gtcaggcgc ccacagtgtc aggagaatct caggtgacct taaaggaggc
1321 catctcgttc tcgttctcat ccttgggtgc ctctcctgt tccttttggg ggcgggggccc
1381 tttggctttc actggtggag aaaacagttg ctactgagaa gattttctgc cttagaacat
1441 gggattcagc catttccggc tcagaggaag atagaggagc tggagcgaga actggagacg
1501 gagatgggac aggagccgga gcccagccg gagccacagc tggagccaga gcccaggcag
1561 ctctga

```

SEQ ID NO: 16 Mouse LAG-3 Amino Acid Sequence

```

1 mredlllgfl llgllweapv vssgpgkelp vvwagegapv hlpcslkspn ldpnflrrgg
61 viwqhqpdsq qptpipaldl hqgmpsprqp apgrytvlsv apgglrsgrq plhphvqlee
121 rglqrgdfsl wlrpalrtda geyhatvrlp nralscslrl rvgqasmias psgvlklsdw
181 vllncsfserp drpvsvhwfq gqnrpvvyns prhflaetfl llpqvsplds gtwgcvltyr
241 dgfnvsityn lkvlglepva pltvyaags rvelpchlpp gvgtpslia kwtppgggpe
301 lpvagksqnf tlhleavgl qagtytcsih lggqqlnatv tlavitvtpk sfglpgsrgk
361 llcevtpasg kerfvwrpln nlsrscpgpv leiqearlla erwqcqlyeg qrllgatvya
421 aesssgahsa rrisgdlkqg hlvlvlilga lsflllvaga fgfhwvrkql llrrfsaleh
481 giqpfpaqrk ieelerelet emgqepepep epqlepeprq l

```

Inhibitors of immune checkpoint modulators are known in the art, and can be used in any of the methods described herein. For example, anti-PD-1 antibodies or soluble polypeptide inhibitors can be used. In some embodiments, the anti-PD-1 antibody is chosen from MDX-1106, Merck 3475 or CT-011. MDX-1106, also known as MDX-1106-04, ONO-4538, BMS-936558 or Nivolumab. In some embodiments, the anti-PD-1 antibody is Nivolumab (CAS Registry Number: 946414-94-4). Nivolumab (also referred to as BMS-936558 or MDX1106; Bristol-Myers Squibb) is a fully human IgG4 monoclonal antibody which specifically blocks PD1. Nivolumab (clone 5C4) and other human monoclonal antibodies that specifically bind to PD1 are disclosed in U.S. Pat. No. 8,008,449 and WO2006/121168. Lambrolizumab (also referred to as MK03475; Merck) is a humanized IgG4 monoclonal antibody that binds to PD1.

Lambrolizumab and other humanized anti-PD1 antibodies are disclosed in U.S. Pat. No. 8,354,509 and WO2009/114335. Pidilizumab (CT-011; Cure Tech) is a humanized IgG1k monoclonal antibody that binds to PD1. Pidilizumab and other humanized anti-PD1 monoclonal antibodies are disclosed in WO2009/101611. Other anti-PD1 antibodies include AMP 514 (Amplimmune), among others, e.g., anti-PD1 antibodies disclosed in U.S. Pat. No. 8,609,089, US 2010028330, and/or US 20120114649. In some embodiments, the PD-1 inhibitor is an immunoadhesin (e.g., an immunoadhesin comprising an extracellular or PD-1 binding portion of PD-L1 or PD-L2 fused to a constant region (e.g., an Fc region of an immunoglobulin sequence). In some embodiments, the PD-1 inhibitor is AMP-224. AMP-224 (B7-DCIg; Amplimmune; e.g., disclosed in WO2010/

027827 and WO2011/066342), is a PD-L2 Fc fusion soluble receptor that blocks the interaction between PD1 and B7-H1.

In other embodiments, anti-PD-L1 antibodies or soluble polypeptide inhibitors can be used. In some embodiments, the PD-L1 inhibitor is anti-PD-L1 antibody. In other embodiments, the anti-PD-L1 binding antagonist is chosen from YW243.55.570, MPDL3280A or MDX-1105. MDX-1105, also known as BMS-936559, is an anti-PD-L1 antibody described in WO2007/005874. MDPL3280A (Genentech/Roche) is a human Fc optimized IgG1 monoclonal antibody that binds to PD-L1. MDPL3280A and other human monoclonal antibodies to PD-L1 are disclosed in U.S. Pat. No. 7,943,743 and U.S. Publication No.: 20120039906. Antibody YW243.55.570 (heavy and light chain variable region sequences shown in SEQ ID Nos. 20 and 21, respectively) is an anti-PD-L1 described in WO 2010/077634.

Exemplary anti-CTLA4 antibodies that can be used in the methods disclosed herein include, but are not limited to, Tremelimumab (IgG2 monoclonal antibody available from Pfizer, formerly known as ticilimumab, CP-675,206); and Ipilimumab (CTLA-4 antibody, also known as MDX-010, CAS No. 477202-00-9). Antibodies to T cell costimulatory molecules such as CTLA-4 (e.g., U.S. Pat. No. 5,811,097), CTLA-4 inhibitor (e.g., CP-675,206, ipilimumab)

II. Methods of Treating Hematologic Cancers

a. Agents Useful for Treating Hematologic Cancers

It is demonstrated herein that inhibiting or blocking a function of PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA-4 synergistically and significantly blocks the establishment and progression of malignancies (e.g., hematologic cancers such as multiple myeloma) in all subjects analyzed. Inhibition or blockade of PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA-4 function can block the establishment and progression of similar malignancies. Thus, the agents of the present invention described herein that modulate the interaction between, for example, PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA-4, whether directly or indirectly, can upregulate or downregulate the immune system and, thereby, upregulate or downregulate an immune response.

PD-1, PD-L1 and TIM-3, LAG-3 and CTLA-4 are immune checkpoint regulators that deliver co-inhibitory immune signals. Thus, in one embodiment, agents that neutralize an activity of PD-1 or PD-L1 and TIM-3, LAG-3 or CTLA-4 can prevent inhibitory signaling and upregulate an immune response. In another embodiment, agents which directly block the interaction between PD-1 or PD-L1 and its natural receptor(s), and TIM-3, LAG-3 or CTLA-4 and its natural receptor(s) (e.g., blocking antibodies) can prevent inhibitory signaling and upregulate an immune response. Alternatively, agents that indirectly block the interaction between PD-1 or PD-L1 and its natural receptor(s), and TIM-3, LAG-3 or CTLA-4 and its natural receptor(s) can prevent inhibitory signaling and upregulate an immune response. For example, soluble B7-1 or soluble PD-1, by binding to a PD-L1 polypeptide indirectly reduces the effective concentration of PD-L1 polypeptide available to bind to the immunoinhibitor receptor, PD-1. Exemplary agents for upregulating an immune response include antibodies against PD-1, PD-L1, LAG-3, CTLA-4 and/or TIM-3 that block the interaction between the immune checkpoint regulator and its natural receptor(s); a non-activating form of PD-1, PD-L1, LAG-3, CTLA-4 and/or TIM-3 (e.g., a dominant negative polypeptide), small molecules or peptides that

block the interaction between the immune checkpoint regulator and its natural receptor(s); fusion proteins (e.g. the extracellular portion of -1, PD-L1, LAG-3, CTLA-4 or TIM-3 fused to the Fc portion of an antibody or immunoglobulin) that bind to their natural receptor(s); nucleic acid molecules that block PD-1, PD-L1, LAG-3, CTLA-4 and/or TIM-3 gene expression, e.g., transcription or translation; and the like.

Additional agents useful in the methods of the present invention include antibodies, small molecules, peptides, peptidomimetics, natural ligands, and derivatives of natural ligands, that can either bind and/or inactivate or inhibit protein biomarkers of the invention, including the biomarkers listed in Table 1, or fragments thereof; as well as RNA interference, antisense, nucleic acid aptamers, etc. that can downregulate the expression and/or activity of the biomarkers of the invention, including the biomarkers listed in Table 1, or fragments thereof.

In one embodiment, isolated nucleic acid molecules that specifically hybridize with or encode one or more biomarkers of the invention, listed in Table 1 for example, or biologically active portions thereof. As used herein, the term "nucleic acid molecule" is intended to include DNA molecules (i.e., cDNA or genomic DNA) and RNA molecules (i.e., mRNA) and analogs of the DNA or RNA generated using nucleotide analogs. The nucleic acid molecule can be single-stranded or double-stranded, but preferably is double-stranded DNA. An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the isolated nucleic acid molecules corresponding to the one or more biomarkers listed in Table 1 or described herein can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb or 0.1 kb of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived (i.e., a lymphoma cell). Moreover, an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized.

A nucleic acid molecule of the present invention, e.g., a nucleic acid molecule having the nucleotide sequence of one or more biomarkers listed in Table 1 or a nucleotide sequence which is at least about 50%, preferably at least about 60%, more preferably at least about 70%, yet more preferably at least about 80%, still more preferably at least about 90%, and most preferably at least about 95% or more (e.g., about 98%) homologous to the nucleotide sequence of one or more biomarkers listed in Table 1 or a portion thereof (i.e., 100, 200, 300, 400, 450, 500, or more nucleotides), can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, a human cDNA can be isolated from a human cell line using all or portion of the nucleic acid molecule, or fragment thereof, as a hybridization probe and standard hybridization techniques (i.e., as described in Sambrook, J., Fritsh, E. F., and Maniatis, T. *Molecular Cloning: A Laboratory Manual*. 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989). Moreover, a nucleic acid molecule encompassing all or a portion of the nucleotide sequence of one or more biomark-

ers listed in Table 1 or a nucleotide sequence which is at least about 50%, preferably at least about 60%, more preferably at least about 70%, yet more preferably at least about 80%, still more preferably at least about 90%, and most preferably at least about 95% or more homologous to the nucleotide sequence, or fragment thereof, can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon the sequence of the one or more biomarkers listed in Table 1, or fragment thereof, or the homologous nucleotide sequence. For example, mRNA can be isolated from muscle cells (i.e., by the guanidinium-thiocyanate extraction procedure of Chirgwin et al. (1979) *Biochemistry* 18: 5294-5299) and cDNA can be prepared using reverse transcriptase (i.e., Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, Md.; or AMV reverse transcriptase, available from Seikagaku America, Inc., St. Petersburg, Fla.). Synthetic oligonucleotide primers for PCR amplification can be designed according to well-known methods in the art. A nucleic acid of the invention can be amplified using cDNA or, alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplification techniques. The nucleic acid so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to the nucleotide sequence of one or more biomarkers listed in Table 1 can be prepared by standard synthetic techniques, i.e., using an automated DNA synthesizer.

Probes based on the nucleotide sequences of one or more biomarkers listed in Table 1 can be used to detect or confirm the desired transcripts or genomic sequences encoding the same or homologous proteins. In preferred embodiments, the probe further comprises a label group attached thereto, i.e., the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a diagnostic test kit for identifying cells or tissue which express one or more biomarkers listed in Table 1, such as by measuring a level of one or more biomarkers nucleic acid in a sample of cells from a subject, i.e., detecting mRNA levels of one or more biomarkers listed in Table 1.

Nucleic acid molecules encoding proteins corresponding to one or more biomarkers listed in Table 1, or portions thereof, from different species are also contemplated. For example, rat or monkey cDNA can be identified based on the nucleotide sequence of a human and/or mouse sequence and such sequences are well known in the art. In one embodiment, the nucleic acid molecule(s) of the invention encodes a protein or portion thereof which includes an amino acid sequence which is sufficiently homologous to an amino acid sequence of one or more biomarkers listed in Table 1, such that the protein or portion thereof modulates (e.g., enhance), one or more of the following biological activities: a) binding to the biomarker; b) modulating the copy number of the biomarker; c) modulating the expression level of the biomarker; and d) modulating the activity level of the biomarker.

As used herein, the language "sufficiently homologous" refers to proteins or portions thereof which have amino acid sequences which include a minimum number of identical or equivalent (e.g., an amino acid residue which has a similar side chain as an amino acid residue in one or more biomarkers listed in Table 1, or fragment thereof) amino acid residues to an amino acid sequence of the biomarker, or fragment thereof, such that the protein or portion thereof modulates (e.g., enhance) one or more of the following biological activities: a) binding to the biomarker; b) modu-

lating the copy number of the biomarker; c) modulating the expression level of the biomarker; and d) modulating the activity level of the biomarker.

In another embodiment, the protein is at least about 50%, preferably at least about 60%, more preferably at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homologous to the entire amino acid sequence of the biomarker, or a fragment thereof.

Portions of proteins encoded by nucleic acid molecules of the one or more biomarkers listed in Table 1 are preferably biologically active portions of the protein. As used herein, the term "biologically active portion" of one or more biomarkers listed in Table 1 is intended to include a portion, e.g., a domain/motif, that has one or more of the biological activities of the full-length protein.

Standard binding assays, e.g., immunoprecipitations and yeast two-hybrid assays, as described herein, or functional assays, e.g., RNAi or overexpression experiments, can be performed to determine the ability of the protein or a biologically active fragment thereof to maintain a biological activity of the full-length protein.

The invention further encompasses nucleic acid molecules that differ from the nucleotide sequence of the one or more biomarkers listed in Table 1, or fragment thereof due to degeneracy of the genetic code and thus encode the same protein as that encoded by the nucleotide sequence, or fragment thereof. In another embodiment, an isolated nucleic acid molecule of the invention has a nucleotide sequence encoding a protein having an amino acid sequence of one or more biomarkers listed in Table 1, or fragment thereof, or a protein having an amino acid sequence which is at least about 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more homologous to the amino acid sequence of the one or more biomarkers listed in Table 1, or fragment thereof. In another embodiment, a nucleic acid encoding a polypeptide consists of nucleic acid sequence encoding a portion of a full-length fragment of interest that is less than 195, 190, 185, 180, 175, 170, 165, 160, 155, 150, 145, 140, 135, 130, 125, 120, 115, 110, 105, 100, 95, 90, 85, 80, 75, or 70 amino acids in length.

It will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences of the one or more biomarkers listed in Table 1 may exist within a population (e.g., a mammalian and/or human population). Such genetic polymorphisms may exist among individuals within a population due to natural allelic variation. As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding one or more biomarkers listed in Table 1, preferably a mammalian, e.g., human, protein. Such natural allelic variations can typically result in 1-5% variance in the nucleotide sequence of the one or more biomarkers listed in Table 1. Any and all such nucleotide variations and resulting amino acid polymorphisms in the one or more biomarkers listed in Table 1 that are the result of natural allelic variation and that do not alter the functional activity of the one or more biomarkers listed in Table 1 are intended to be within the scope of the invention. Moreover, nucleic acid molecules encoding one or more biomarkers listed in Table 1 from other species.

In addition to naturally-occurring allelic variants of the one or more biomarkers listed in Table 1 that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into the nucleotide sequence, or fragment thereof, thereby leading to changes in the amino acid sequence of the encoded one or more biomarkers listed in Table 1, without altering the functional

ability of the one or more biomarkers listed in Table 1. For example, nucleotide substitutions leading to amino acid substitutions at “non-essential” amino acid residues can be made in the sequence, or fragment thereof. A “non-essential” amino acid residue is a residue that can be altered from the wild-type sequence of the one or more biomarkers listed in Table 1 without altering the activity of the one or more biomarkers listed in Table 1, whereas an “essential” amino acid residue is required for the activity of the one or more biomarkers listed in Table 1. Other amino acid residues, however, (e.g., those that are not conserved or only semi-conserved between mouse and human) may not be essential for activity and thus are likely to be amenable to alteration without altering the activity of the one or more biomarkers listed in Table 1.

The comparison of sequences and determination of percent homology between two sequences can be accomplished using a mathematical algorithm. Preferably, the alignment can be performed using the Clustal Method. Multiple alignment parameters include GAP Penalty=10, Gap Length Penalty=10. For DNA alignments, the pairwise alignment parameters can be Htuple=2, Gap penalty=5, Window=4, and Diagonal saved=4. For protein alignments, the pairwise alignment parameters can be Ktuple=1, Gap penalty=3, Window=5, and Diagonals Saved=5.

In a preferred embodiment, the percent identity between two amino acid sequences is determined using the Needleman and Wunsch (*J. Mol. Biol.* (48):444-453 (1970)) algorithm which has been incorporated into the GAP program in the GCG software package (available online), using either a Blossom 62 matrix or a PAM250 matrix, and a gap weight of 16, 14, 12, 10, 8, 6, or 4 and a length weight of 1, 2, 3, 4, 5, or 6. In yet another preferred embodiment, the percent identity between two nucleotide sequences is determined using the GAP program in the GCG software package (available online), using a NWSgapdna.CMP matrix and a gap weight of 40, 50, 60, 70, or 80 and a length weight of 1, 2, 3, 4, 5, or 6. In another embodiment, the percent identity between two amino acid or nucleotide sequences is determined using the algorithm of E. Meyers and W. Miller (*CABIOS*, 4:11-17 (1989)) which has been incorporated into the ALIGN program (version 2.0) (available online), using a PAM120 weight residue table, a gap length penalty of 12 and a gap penalty of 4.

An isolated nucleic acid molecule encoding a protein homologous to one or more biomarkers listed in Table 1, or fragment thereof, can be created by introducing one or more nucleotide substitutions, additions or deletions into the nucleotide sequence, or fragment thereof, or a homologous nucleotide sequence such that one or more amino acid substitutions, additions or deletions are introduced into the encoded protein. Mutations can be introduced by standard techniques, such as site-directed mutagenesis and PCR-mediated mutagenesis. Preferably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A “conservative amino acid substitution” is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), nonpolar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), branched side chains (e.g., threonine, valine, isoleucine) and aromatic side

chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in one or more biomarkers listed in Table 1 is preferably replaced with another amino acid residue from the same side chain family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of the coding sequence of the one or more biomarkers listed in Table 1, such as by saturation mutagenesis, and the resultant mutants can be screened for an activity described herein to identify mutants that retain desired activity. Following mutagenesis, the encoded protein can be expressed recombinantly according to well-known methods in the art and the activity of the protein can be determined using, for example, assays described herein.

The levels of one or more biomarkers listed in Table 1 may be assessed by any of a wide variety of well-known methods for detecting expression of a transcribed molecule or protein. Non-limiting examples of such methods include immunological methods for detection of proteins, protein purification methods, protein function or activity assays, nucleic acid hybridization methods, nucleic acid reverse transcription methods, and nucleic acid amplification methods.

In preferred embodiments, the levels of one or more biomarkers listed in Table 1 are ascertained by measuring gene transcript (e.g., mRNA), by a measure of the quantity of translated protein, or by a measure of gene product activity. Expression levels can be monitored in a variety of ways, including by detecting mRNA levels, protein levels, or protein activity, any of which can be measured using standard techniques. Detection can involve quantification of the level of gene expression (e.g., genomic DNA, cDNA, mRNA, protein, or enzyme activity), or, alternatively, can be a qualitative assessment of the level of gene expression, in particular in comparison with a control level. The type of level being detected will be clear from the context.

In a particular embodiment, the mRNA expression level can be determined both by in situ and by in vitro formats in a biological sample using methods known in the art. The term “biological sample” is intended to include tissues, cells, biological fluids and isolates thereof, isolated from a subject, as well as tissues, cells and fluids present within a subject. Many expression detection methods use isolated RNA. For in vitro methods, any RNA isolation technique that does not select against the isolation of mRNA can be utilized for the purification of RNA from cells (see, e.g., Ausubel et al., ed., *Current Protocols in Molecular Biology*, John Wiley & Sons, New York 1987-1999). Additionally, large numbers of tissue samples can readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski (1989, U.S. Pat. No. 4,843,155).

The isolated mRNA can be used in hybridization or amplification assays that include, but are not limited to, Southern or Northern analyses, polymerase chain reaction analyses and probe arrays. One preferred diagnostic method for the detection of mRNA levels involves contacting the isolated mRNA with a nucleic acid molecule (probe) that can hybridize to the mRNA encoded by the gene being detected. The nucleic acid probe can be, for example, a full-length cDNA, or a portion thereof, such as an oligonucleotide of at least 7, 15, 30, 50, 100, 250 or 500 nucleotides in length and sufficient to specifically hybridize under stringent conditions to a mRNA or genomic DNA encoding one or more biomarkers listed in Table 1. Other suitable probes for use in the diagnostic assays of the invention are described herein. Hybridization of an mRNA

with the probe indicates that one or more biomarkers listed in Table 1 is being expressed.

In one format, the mRNA is immobilized on a solid surface and contacted with a probe, for example by running the isolated mRNA on an agarose gel and transferring the mRNA from the gel to a membrane, such as nitrocellulose. In an alternative format, the probe(s) are immobilized on a solid surface and the mRNA is contacted with the probe(s), for example, in a gene chip array, e.g., an Affymetrix™ gene chip array. A skilled artisan can readily adapt known mRNA detection methods for use in detecting the level of the one or more biomarkers listed in Table 1.

An alternative method for determining mRNA expression level in a sample involves the process of nucleic acid amplification, e.g., by RT-PCR (the experimental embodiment set forth in Mullis, 1987, U.S. Pat. No. 4,683,202), ligase chain reaction (Barany, 1991, *Proc. Natl. Acad. Sci. USA*, 88:189-193), self-sustained sequence replication (Guatelli et al., 1990, *Proc. Natl. Acad. Sci. USA* 87:1874-1878), transcriptional amplification system (Kwoh et al., 1989, *Proc. Natl. Acad. Sci. USA* 86:1173-1177), Q-Beta Replicase (Lizardi et al., 1988, *Bio/Technology* 6:1197), rolling circle replication (Lizardi et al., U.S. Pat. No. 5,854,033) or any other nucleic acid amplification method, followed by the detection of the amplified molecules using techniques well-known to those of skill in the art. These detection schemes are especially useful for the detection of nucleic acid molecules if such molecules are present in very low numbers. As used herein, amplification primers are defined as being a pair of nucleic acid molecules that can anneal to 5' or 3' regions of a gene (plus and minus strands, respectively, or vice-versa) and contain a short region in between. In general, amplification primers are from about 10 to 30 nucleotides in length and flank a region from about 50 to 200 nucleotides in length. Under appropriate conditions and with appropriate reagents, such primers permit the amplification of a nucleic acid molecule comprising the nucleotide sequence flanked by the primers.

For in situ methods, mRNA does not need to be isolated from the cells prior to detection. In such methods, a cell or tissue sample is prepared/processed using known histological methods. The sample is then immobilized on a support, typically a glass slide, and then contacted with a probe that can hybridize to the one or more biomarkers listed in Table 1.

As an alternative to making determinations based on the absolute expression level, determinations may be based on the normalized expression level of one or more biomarkers listed in Table 1. Expression levels are normalized by correcting the absolute expression level by comparing its expression to the expression of a non-biomarker gene, e.g., a housekeeping gene that is constitutively expressed. Suitable genes for normalization include housekeeping genes such as the actin gene, or epithelial cell-specific genes. This normalization allows the comparison of the expression level in one sample, e.g., a subject sample, to another sample, e.g., a normal sample, or between samples from different sources.

The level or activity of a protein corresponding to one or more biomarkers listed in Table 1 can also be detected and/or quantified by detecting or quantifying the expressed polypeptide. The polypeptide can be detected and quantified by any of a number of means well known to those of skill in the art. These may include analytic biochemical methods such as electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), hyperdiffusion chromatography, and the like, or various immunological methods such as fluid or gel

precipitin reactions, immunodiffusion (single or double), immunoelectrophoresis, radioimmunoassay (RIA), enzyme-linked immunosorbent assays (ELISAs), immunofluorescent assays, Western blotting, and the like. A skilled artisan can readily adapt known protein/antibody detection methods for use in determining whether cells express the biomarker of interest.

The present invention further provides soluble, purified and/or isolated polypeptide forms of one or more biomarkers listed in Table 1, or fragments thereof. In addition, it is to be understood that any and all attributes of the polypeptides described herein, such as percentage identities, polypeptide lengths, polypeptide fragments, biological activities, antibodies, etc. can be combined in any order or combination with respect to any biomarker listed in Table 1 and combinations thereof.

In one aspect, a polypeptide may comprise a full-length amino acid sequence corresponding to one or more biomarkers listed in Table 1 or a full-length amino acid sequence with 1 to about 20 conservative amino acid substitutions. An amino acid sequence of any described herein can also be at least 50, 55, 60, 65, 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 99.5% identical to the full-length sequence of one or more biomarkers listed in Table 1, which is either described herein, well known in the art, or a fragment thereof. In another aspect, the present invention contemplates a composition comprising an isolated polypeptide corresponding to one or more biomarkers listed in Table 1 and less than about 25%, or alternatively 15%, or alternatively 5%, contaminating biological macromolecules or polypeptides.

The present invention further provides compositions related to producing, detecting, or characterizing such polypeptides, or fragment thereof, such as nucleic acids, vectors, host cells, and the like. Such compositions may serve as compounds that modulate the expression and/or activity of one or more biomarkers described herein or, for example, listed in Table 1.

An isolated polypeptide or a fragment thereof (or a nucleic acid encoding such a polypeptide) corresponding to one or more biomarkers of the invention, including the biomarkers listed in Table 1 or fragments thereof, can be used as an immunogen to generate antibodies that bind to said immunogen, using standard techniques for polyclonal and monoclonal antibody preparation according to well-known methods in the art. An antigenic peptide comprises at least 8 amino acid residues and encompasses an epitope present in the respective full length molecule such that an antibody raised against the peptide forms a specific immune complex with the respective full length molecule. Preferably, the antigenic peptide comprises at least 10 amino acid residues. In one embodiment such epitopes can be specific for a given polypeptide molecule from one species, such as mouse or human (i.e., an antigenic peptide that spans a region of the polypeptide molecule that is not conserved across species is used as immunogen; such non conserved residues can be determined using an alignment such as that provided herein).

In one embodiment, an antibody binds substantially specifically to PD-L1 and inhibits or blocks its immunoinhibitory function, such as by interrupting its interaction with an inhibitory receptor like PD-1. In another embodiment, an antibody binds substantially specifically to TIM-3 and inhibits or blocks its immunoinhibitory function, such as by interrupting its interaction with galectin-9 or phosphatidylserine.

For example, a polypeptide immunogen typically is used to prepare antibodies by immunizing a suitable subject (e.g., rabbit, goat, mouse or other mammal) with the immunogen. An appropriate immunogenic preparation can contain, for example, a recombinantly expressed or chemically synthesized molecule or fragment thereof to which the immune response is to be generated. The preparation can further include an adjuvant, such as Freund's complete or incomplete adjuvant, or similar immunostimulatory agent. Immunization of a suitable subject with an immunogenic preparation induces a polyclonal antibody response to the antigenic peptide contained therein.

Polyclonal antibodies can be prepared as described above by immunizing a suitable subject with a polypeptide immunogen. The polypeptide antibody titer in the immunized subject can be monitored over time by standard techniques, such as with an enzyme linked immunosorbent assay (ELISA) using immobilized polypeptide. If desired, the antibody directed against the antigen can be isolated from the mammal (e.g., from the blood) and further purified by well-known techniques, such as protein A chromatography, to obtain the IgG fraction. At an appropriate time after immunization, e.g., when the antibody titers are highest, antibody-producing cells can be obtained from the subject and used to prepare monoclonal antibodies by standard techniques, such as the hybridoma technique (originally described by Kohler and Milstein (1975) *Nature* 256:495-497) (see also Brown et al. (1981) *J. Immunol.* 127:539-46; Brown et al. (1980) *J. Biol. Chem.* 255:4980-83; Yeh et al. (1976) *Proc. Natl. Acad. Sci.* 76:2927-31; Yeh et al. (1982) *Int. J. Cancer* 29:269-75), the more recent human B cell hybridoma technique (Kozbor et al. (1983) *Immunol. Today* 4:72), the EBV-hybridoma technique (Cole et al. (1985) *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc., pp. 77-96) or trioma techniques. The technology for producing monoclonal antibody hybridomas is well known (see generally Kenneth, R. H. in *Monoclonal Antibodies: A New Dimension In Biological Analyses*, Plenum Publishing Corp., New York, N.Y. (1980); Lerner, E. A. (1981) *Yale J. Biol. Med.* 54:387-402; Gefter, M. L. et al. (1977) *Somatic Cell Genet.* 3:231-36). Briefly, an immortal cell line (typically a myeloma) is fused to lymphocytes (typically splenocytes) from a mammal immunized with an immunogen as described above, and the culture supernatants of the resulting hybridoma cells are screened to identify a hybridoma producing a monoclonal antibody that binds to the polypeptide antigen, preferably specifically.

Any of the many well-known protocols used for fusing lymphocytes and immortalized cell lines can be applied for the purpose of generating a monoclonal antibody against one or more biomarkers of the invention, including the biomarkers listed in Table 1, or a fragment thereof (see, e.g., Galfre, G. et al. (1977) *Nature* 266:55052; Gefter et al. (1977) *supra*; Lerner (1981) *supra*; Kenneth (1980) *supra*). Moreover, the ordinary skilled worker will appreciate that there are many variations of such methods which also would be useful. Typically, the immortal cell line (e.g., a myeloma cell line) is derived from the same mammalian species as the lymphocytes. For example, murine hybridomas can be made by fusing lymphocytes from a mouse immunized with an immunogenic preparation of the present invention with an immortalized mouse cell line. Preferred immortal cell lines are mouse myeloma cell lines that are sensitive to culture medium containing hypoxanthine, aminopterin and thymidine ("HAT medium"). Any of a number of myeloma cell lines can be used as a fusion partner according to standard techniques, e.g., the P3-NS1/1-Ag4-1, P3-x63-Ag8.653 or

Sp2/O—Ag14 myeloma lines. These myeloma lines are available from the American Type Culture Collection (ATCC), Rockville, Md. Typically, HAT-sensitive mouse myeloma cells are fused to mouse splenocytes using polyethylene glycol ("PEG"). Hybridoma cells resulting from the fusion are then selected using HAT medium, which kills unfused and unproductively fused myeloma cells (unfused splenocytes die after several days because they are not transformed). Hybridoma cells producing a monoclonal antibody of the invention are detected by screening the hybridoma culture supernatants for antibodies that bind a given polypeptide, e.g., using a standard ELISA assay.

As an alternative to preparing monoclonal antibody-secreting hybridomas, a monoclonal specific for one of the above described polypeptides can be identified and isolated by screening a recombinant combinatorial immunoglobulin library (e.g., an antibody phage display library) with the appropriate polypeptide to thereby isolate immunoglobulin library members that bind the polypeptide. Kits for generating and screening phage display libraries are commercially available (e.g., the Pharmacia *Recombinant Phage Antibody System*, Catalog No. 27-9400-01; and the Stratagene *SurfZAP™ Phage Display Kit*, Catalog No. 240612). Additionally, examples of methods and reagents particularly amenable for use in generating and screening an antibody display library can be found in, for example, Ladner et al. U.S. Pat. No. 5,223,409; Kang et al. International Publication No. WO 92/18619; Dower et al. International Publication No. WO 91/17271; Winter et al. International Publication WO 92/20791; Markland et al. International Publication No. WO 92/15679; Breitling et al. International Publication WO 93/01288; McCafferty et al. International Publication No. WO 92/01047; Garrard et al. International Publication No. WO 92/09690; Ladner et al. International Publication No. WO 90/02809; Fuchs et al. (1991) *Biotechnology* (NY) 9:1369-1372; Hay et al. (1992) *Hum. Antibod. Hybridomas* 3:81-85; Huse et al. (1989) *Science* 246:1275-1281; Griffiths et al. (1993) *EMBO J.* 12:725-734; Hawkins et al. (1992) *J. Mol. Biol.* 226:889-896; Clarkson et al. (1991) *Nature* 352:624-628; Gram et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:3576-3580; Garrard et al. (1991) *Biotechnology* (NY) 9:1373-1377; Hoogenboom et al. (1991) *Nucleic Acids Res.* 19:4133-4137; Barbas et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:7978-7982; and McCafferty et al. (1990) *Nature* 348:552-554.

Since it is well known in the art that antibody heavy and light chain CDR3 domains play a particularly important role in the binding specificity/affinity of an antibody for an antigen, the recombinant monoclonal antibodies of the present invention prepared as set forth above preferably comprise the heavy and light chain CDR3s of variable regions of the antibodies described herein and well known in the art. Similarly, the antibodies can further comprise the CDR2s of variable regions of said antibodies. The antibodies can further comprise the CDR1s of variable regions of said antibodies. In other embodiments, the antibodies can comprise any combinations of the CDRs.

The CDR1, 2, and/or 3 regions of the engineered antibodies described above can comprise the exact amino acid sequence(s) as those of variable regions of the present invention disclosed herein. However, the ordinarily skilled artisan will appreciate that some deviation from the exact CDR sequences may be possible while still retaining the ability of the antibody to bind a desired target, such as PD-L1, TIM-3, or LAG-3 effectively (e.g., conservative sequence modifications). Accordingly, in another embodiment, the engineered antibody may be composed of one or

more CDRs that are, for example, 50%, 60%, 70%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or 99.5% identical to one or more CDRs of the present invention described herein or otherwise publicly available.

The structural features of non-human or human antibodies (e.g., a rat anti-mouse/anti-human PD-L1 antibody) can be used to create structurally related human antibodies that retain at least one functional property of the antibodies of the present invention, such as binding to PD-L1, TIM-3, or LAG-3. Another functional property includes inhibiting binding of the original known, non-human or human antibodies in a competition ELISA assay.

In some embodiments, monoclonal antibodies capable of binding and inhibiting/blocking PD-L1, TIM-3, and/or LAG-3 are provided, comprising a heavy chain wherein the variable domain comprises at least a CDR having a sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical from the group of heavy chain variable domain CDRs presented herein or otherwise publicly available.

Similarly, monoclonal antibodies binding and inhibiting/blocking PD-L1, TIM-3, and/or LAG-3, comprising a light chain wherein the variable domain comprises at least a CDR having a sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical from the group of light chain variable domain CDRs presented herein or otherwise publicly available, are also provided.

Monoclonal antibodies capable of binding and inhibiting/blocking PD-L1, TIM-3, and/or LAG-3, comprising a heavy chain wherein the variable domain comprises at least a CDR having a sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical from the group of heavy chain variable domain CDRs presented herein or otherwise publicly available; and comprising a light chain wherein the variable domain comprises at least a CDR having a sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical from the group of light chain variable domain CDRs presented herein or otherwise publicly available, are also provided.

A skilled artisan will note that such percentage homology is equivalent to and can be achieved by introducing 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, or more conservative amino acid substitutions within a given CDR.

The monoclonal antibodies of the present invention can comprise a heavy chain, wherein the variable domain comprises at least a CDR having a sequence selected from the group consisting of the heavy chain variable domain CDRs presented herein or otherwise publicly available and a light chain, wherein the variable domain comprises at least a CDR having a sequence selected from the group consisting of the light chain variable domain CDRs presented herein or otherwise publicly available.

Such monoclonal antibodies can comprise a light chain, wherein the variable domain comprises at least a CDR having a sequence selected from the group consisting of CDR-L1, CDR-L2, and CDR-L3, as described herein; and/or a heavy chain, wherein the variable domain comprises at least a CDR having a sequence selected from the group consisting of CDR-H1, CDR-H2, and CDR-H3, as described herein. In some embodiments, the monoclonal antibodies capable of binding human Gall comprises or consists of CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3, as described herein.

The heavy chain variable domain of the monoclonal antibodies of the present invention can comprise or consist

of the vH amino acid sequence set forth herein or otherwise publicly available and/or the light chain variable domain of the monoclonal antibodies of the present invention can comprise or consist of the vk amino acid sequence set forth herein or otherwise publicly available.

The present invention further provides fragments of said monoclonal antibodies which include, but are not limited to, Fv, Fab, F(ab')₂, Fab', dsFv, scFv, sc(Fv)₂ and diabodies; and multispecific antibodies formed from antibody fragments.

Other fragments of the monoclonal antibodies of the present invention are also contemplated. For example, individual immunoglobulin heavy and/or light chains are provided, wherein the variable domains thereof comprise at least a CDR presented herein or otherwise publicly available. In one embodiment, the immunoglobulin heavy chain comprises at least a CDR having a sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical from the group of heavy chain or light chain variable domain CDRs presented herein or otherwise publicly available. In another embodiment, an immunoglobulin light chain comprises at least a CDR having a sequence that is at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, 99.5% or 100% identical from the group of light chain or heavy chain variable domain CDRs presented herein or otherwise publicly available, are also provided.

In some embodiments, the immunoglobulin heavy and/or light chain comprises a variable domain comprising at least one of CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, or CDR-H3 described herein. Such immunoglobulin heavy chains can comprise or consist of at least one of CDR-H1, CDR-H2, and CDR-H3. Such immunoglobulin light chains can comprise or consist of at least one of CDR-L1, CDR-L2, and CDR-L3.

In other embodiments, an immunoglobulin heavy and/or light chain according to the present invention comprises or consists of a vH or vk variable domain sequence, respectively, provided herein or otherwise publicly available.

The present invention further provides polypeptides which have a sequence selected from the group consisting of vH variable domain, vk variable domain, CDR-L1, CDR-L2, CDR-L3, CDR-H1, CDR-H2, and CDR-H3 sequences described herein.

Antibodies, immunoglobulins, and polypeptides of the invention can be used in an isolated (e.g., purified) form or contained in a vector, such as a membrane or lipid vesicle (e.g. a liposome).

Amino acid sequence modification(s) of the antibodies described herein are contemplated. For example, it may be desirable to improve the binding affinity and/or other biological properties of the antibody. It is known that when a humanized antibody is produced by simply grafting only CDRs in VH and VL of an antibody derived from a non-human animal in FRs of the VH and VL of a human antibody, the antigen binding activity is reduced in comparison with that of the original antibody derived from a non-human animal. It is considered that several amino acid residues of the VH and VL of the non-human antibody, not only in CDRs but also in FRs, are directly or indirectly associated with the antigen binding activity. Hence, substitution of these amino acid residues with different amino acid residues derived from FRs of the VH and VL of the human antibody would reduce binding activity and can be corrected by replacing the amino acids with amino acid residues of the original antibody derived from a non-human animal.

Modifications and changes may be made in the structure of the antibodies described herein, and in the DNA sequences encoding them, and still obtain a functional molecule that encodes an antibody and polypeptide with desirable characteristics. For example, certain amino acids may be substituted by other amino acids in a protein structure without appreciable loss of activity. Since the interactive capacity and nature of a protein define the protein's biological functional activity, certain amino acid substitutions can be made in a protein sequence, and, of course, in its DNA encoding sequence, while nevertheless obtaining a protein with like properties. It is thus contemplated that various changes may be made in the antibodies sequences of the invention, or corresponding DNA sequences which encode said polypeptides, without appreciable loss of their biological activity.

In making the changes in the amino sequences of polypeptide, the hydrophobic index of amino acids may be considered. The importance of the hydrophobic amino acid index in conferring interactive biologic function on a protein is generally understood in the art. It is accepted that the relative hydrophobic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like. Each amino acid has been assigned a hydrophobic index on the basis of their hydrophobicity and charge characteristics these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (<RTI 3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydrophobic index or score and still result in a protein with similar biological activity, i.e. still obtain a biological functionally equivalent protein.

As outlined above, amino acid substitutions are generally therefore based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions which take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

Another type of amino acid modification of the antibody of the invention may be useful for altering the original glycosylation pattern of the antibody to, for example, increase stability. By "altering" is meant deleting one or more carbohydrate moieties found in the antibody, and/or adding one or more glycosylation sites that are not present in the antibody. Glycosylation of antibodies is typically N-linked. "N-linked" refers to the attachment of the carbohydrate moiety to the side chain of an asparagine residue. The tripeptide sequences asparagine-X-serine and asparagine-X-threonine, where X is any amino acid except proline, are the recognition sequences for enzymatic attachment of the carbohydrate moiety to the asparagine side chain. Thus, the presence of either of these tripeptide sequences in a polypeptide creates a potential glycosylation site. Addition of glycosylation sites to the antibody is conveniently accomplished by altering the amino acid sequence such that it contains one or more of the above-described tripeptide sequences (for N-linked glycosylation sites). Another type

of covalent modification involves chemically or enzymatically coupling glycosides to the antibody. These procedures are advantageous in that they do not require production of the antibody in a host cell that has glycosylation capabilities for N- or O-linked glycosylation. Depending on the coupling mode used, the sugar(s) may be attached to (a) arginine and histidine, (b) free carboxyl groups, (c) free sulfhydryl groups such as those of cysteine, (d) free hydroxyl groups such as those of serine, threonine, or hydroxyproline, (e) aromatic residues such as those of phenylalanine, tyrosine, or tryptophan, or (f) the amide group of glutamine. For example, such methods are described in WO87/05330.

Similarly, removal of any carbohydrate moieties present on the antibody may be accomplished chemically or enzymatically. Chemical deglycosylation requires exposure of the antibody to the compound trifluoromethanesulfonic acid, or an equivalent compound. This treatment results in the cleavage of most or all sugars except the linking sugar (N-acetylglucosamine or N-acetylgalactosamine), while leaving the antibody intact. Chemical deglycosylation is described by Sojahn et al. (1987) and by Edge et al. (1981). Enzymatic cleavage of carbohydrate moieties on antibodies can be achieved by the use of a variety of endo- and exo-glycosidases as described by Thotakura et al. (1987).

Other modifications can involve the formation of immunoconjugates. For example, in one type of covalent modification, antibodies or proteins are covalently linked to one of a variety of non proteinaceous polymers, e.g., polyethylene glycol, polypropylene glycol, or polyoxyalkylenes, in the manner set forth in U.S. Pat. Nos. 4,640,835; 4,496,689; 4,301,144; 4,670,417; 4,791,192 or 4,179,337.

Conjugation of antibodies or other proteins of the present invention with heterologous agents can be made using a variety of bifunctional protein coupling agents including but not limited to N-succinimidyl (2-pyridyldithio) propionate (SPDP), succinimidyl (N-maleimidomethyl)cyclohexane-1-carboxylate, iminothiolane (IT), bifunctional derivatives of imidoesters (such as dimethyl adipimidate HCL), active esters (such as disuccinimidyl suberate), aldehydes (such as glutaraldehyde), bis-azido compounds (such as bis (p-azidobenzoyl) hexanediamine), bis-diazonium derivatives (such as bis-(p-diazoniumbenzoyl)-ethylenediamine), diisocyanates (such as toluene 2,6diisocyanate), and bis-active fluorine compounds (such as 1,5-difluoro-2,4-dinitrobenzene). For example, carbon labeled 1-isothiocyanatobenzyl methyl-diethylene triamine-pentaacetic acid (MX-DTPA) is an exemplary chelating agent for conjugation of radionucleotide to the antibody (WO 94/11026).

In another aspect, the present invention features antibodies conjugated to a therapeutic moiety, such as a cytotoxin, a drug, and/or a radioisotope. When conjugated to a cytotoxin, these antibody conjugates are referred to as "immunotoxins." A cytotoxin or cytotoxic agent includes any agent that is detrimental to (e.g., kills) cells. Examples include taxol, cytochalasin B, gramicidin D, ethidium bromide, emetine, mitomycin, etoposide, teniposide, vincristine, vinblastine, colchicin, doxorubicin, daunorubicin, dihydroxy anthracin dione, mitoxantrone, mithramycin, actinomycin D, 1-dehydrotestosterone, glucocorticoids, procaine, tetracaine, lidocaine, propranolol, and puromycin and analogs or homologs thereof. Therapeutic agents include, but are not limited to, antimetabolites (e.g., methotrexate, 6-mercaptopurine, 6-thioguanine, cytarabine, 5-fluorouracil decarbazine), alkylating agents (e.g., mechlorethamine, thioepa chlorambucil, melphalan, carmustine (BSNU) and lomustine (CCNU), cyclophosphamide, busulfan, dibromomannitol, streptozotocin, mitomycin C, and cis-dichlorodiamine

platinum (II) (DDP) cisplatin), anthracyclines (e.g., daunorubicin (formerly daunomycin) and doxorubicin), antibiotics (e.g., dactinomycin (formerly actinomycin), bleomycin, mithramycin, and anthramycin (AMC)), and anti-mitotic agents (e.g., vincristine and vinblastine). An antibody of the present invention can be conjugated to a radioisotope, e.g., radioactive iodine, to generate cytotoxic radiopharmaceuticals for treating a related disorder, such as a cancer.

Conjugated antibodies, in addition to therapeutic utility, can be useful for diagnostically or prognostically to monitor polypeptide levels in tissue as part of a clinical testing procedure, e.g., to determine the efficacy of a given treatment regimen. Detection can be facilitated by coupling (i.e., physically linking) the antibody to a detectable substance. Examples of detectable substances include various enzymes, prosthetic groups, fluorescent materials, luminescent materials, bioluminescent materials, and radioactive materials. Examples of suitable enzymes include horseradish peroxidase, alkaline phosphatase, β -galactosidase, or acetylcholinesterase; examples of suitable prosthetic group complexes include streptavidin/biotin and avidin/biotin; examples of suitable fluorescent materials include umbelliferone, fluorescein, fluorescein isothiocyanate (FITC), rhodamine, dichlorotriazinylamine fluorescein, dansyl chloride or phycoerythrin (PE); an example of a luminescent material includes luminol; examples of bioluminescent materials include luciferase, luciferin, and aequorin, and examples of suitable radioactive material include ^{125}I , ^{131}I , ^{35}S , or ^3H . [0134] As used herein, the term "labeled", with regard to the antibody, is intended to encompass direct labeling of the antibody by coupling (i.e., physically linking) a detectable substance, such as a radioactive agent or a fluorophore (e.g. fluorescein isothiocyanate (FITC) or phycoerythrin (PE) or Indocyanine (Cy5)) to the antibody, as well as indirect labeling of the antibody by reactivity with a detectable substance.

The antibody conjugates of the present invention can be used to modify a given biological response. The therapeutic moiety is not to be construed as limited to classical chemical therapeutic agents. For example, the drug moiety may be a protein or polypeptide possessing a desired biological activity. Such proteins may include, for example, an enzymatically active toxin, or active fragment thereof, such as abrin, ricin A, *Pseudomonas* exotoxin, or diphtheria toxin; a protein such as tumor necrosis factor or interferon- γ ; or biological response modifiers such as, for example, lymphokines, interleukin-1 ("IL-1"), interleukin-2 ("IL-2"), interleukin-6 ("IL-6"), granulocyte macrophage colony stimulating factor ("GM-CSF"), granulocyte colony stimulating factor ("G-CSF"), or other cytokines or growth factors.

Techniques for conjugating such therapeutic moiety to antibodies are well known, see, e.g., Arnon et al., "Monoclonal Antibodies For Immunotargeting Of Drugs In Cancer Therapy", in *Monoclonal Antibodies And Cancer Therapy*, Reisfeld et al. (eds.), pp. 243-56 (Alan R. Liss, Inc. 1985); Hellstrom et al., "Antibodies For Drug Delivery", in *Controlled Drug Delivery* (2nd Ed.), Robinson et al. (eds.), pp. 623-53 (Marcel Dekker, Inc. 1987); Thorpe, "Antibody Carriers Of Cytotoxic Agents In Cancer Therapy: A Review", in *Monoclonal Antibodies '84: Biological And Clinical Applications*, Pinchera et al. (eds.), pp. 475-506 (1985); "Analysis, Results, And Future Prospective Of The Therapeutic Use Of Radiolabeled Antibody In Cancer Therapy", in *Monoclonal Antibodies For Cancer Detection And Therapy*, Baldwin et al. (eds.), pp. 303-16 (Academic Press 1985), and Thorpe et al., "The Preparation And

Cytotoxic Properties Of Antibody-Toxin Conjugates", *Immunol. Rev.*, 62:119-58 (1982).

In some embodiments, conjugations can be made using a "cleavable linker" facilitating release of the cytotoxic agent or growth inhibitory agent in a cell. For example, an acid-labile linker, peptidase-sensitive linker, photolabile linker, dimethyl linker or disulfide-containing linker (See e.g. U.S. Pat. No. 5,208,020) may be used. Alternatively, a fusion protein comprising the antibody and cytotoxic agent or growth inhibitory agent may be made, by recombinant techniques or peptide synthesis. The length of DNA may comprise respective regions encoding the two portions of the conjugate either adjacent one another or separated by a region encoding a linker peptide which does not destroy the desired properties of the conjugate.

Additionally, recombinant polypeptide antibodies, such as chimeric and humanized monoclonal antibodies, comprising both human and non-human portions, which can be made using standard recombinant DNA techniques, are within the scope of the invention. Such chimeric and humanized monoclonal antibodies can be produced by recombinant DNA techniques known in the art, for example using methods described in Robinson et al. International Patent Publication PCT/US86/02269; Akira et al. European Patent Application 184,187; Taniguchi, M. European Patent Application 171,496; Morrison et al. European Patent Application 173,494; Neuberger et al. PCT Application WO 86/01533; Cabilly et al. U.S. Pat. No. 4,816,567; Cabilly et al. European Patent Application 125,023; Better et al. (1988) *Science* 240:1041-1043; Liu et al. (1987) *Proc. Natl. Acad. Sci. USA* 84:3439-3443; Liu et al. (1987) *J. Immunol.* 139:3521-3526; Sun et al. (1987) *Proc. Natl. Acad. Sci.* 84:214-218; Nishimura et al. (1987) *Cancer Res.* 47:999-1005; Wood et al. (1985) *Nature* 314:446-449; Shaw et al. (1988) *J. Natl. Cancer Inst.* 80:1553-1559; Morrison, S. L. (1985) *Science* 229:1202-1207; Oi et al. (1986) *Biotechniques* 4:214; Winter U.S. Pat. No. 5,225,539; Jones et al. (1986) *Nature* 321:552-525; Verhoeyan et al. (1988) *Science* 239:1534; and Beidler et al. (1988) *J. Immunol.* 141:4053-4060.

In addition, humanized antibodies can be made according to standard protocols such as those disclosed in U.S. Pat. No. 5,565,332. In another embodiment, antibody chains or specific binding pair members can be produced by recombination between vectors comprising nucleic acid molecules encoding a fusion of a polypeptide chain of a specific binding pair member and a component of a replicable generic display package and vectors containing nucleic acid molecules encoding a second polypeptide chain of a single binding pair member using techniques known in the art, e.g., as described in U.S. Pat. Nos. 5,565,332, 5,871,907, or 5,733,743. The use of intracellular antibodies to inhibit protein function in a cell is also known in the art (see e.g., Carlson, J. R. (1988) *Mol. Cell. Biol.* 8:2638-2646; Biocca, S. et al. (1990) *EMBO J.* 9:101-108; Werge, T. M. et al. (1990) *FEBS Lett.* 274:193-198; Carlson, J. R. (1993) *Proc. Natl. Acad. Sci. USA* 90:7427-7428; Marasco, W. A. et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:7889-7893; Biocca, S. et al. (1994) *Biotechnology* (NY) 12:396-399; Chen, S-Y. et al. (1994) *Hum. Gene Ther.* 5:595-601; Duan, L et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:5075-5079; Chen, S-Y. et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:5932-5936; Beerli, R. R. et al. (1994) *J. Biol. Chem.* 269:23931-23936; Beerli, R. R. et al. (1994) *Biochem. Biophys. Res. Commun.* 204:666-672; Mhashilkar, A. M. et al. (1995) *EMBO J.* 14:1542-1551; Richardson, J. H. et al. (1995) *Proc. Natl. Acad. Sci.*

USA 92:3137-3141; PCT Publication No. WO 94/02610 by Marasco et al.; and PCT Publication No. WO 95/03832 by Duan et al.).

Additionally, fully human antibodies could be made against biomarkers of the invention, including the biomarkers listed in Table 1, or fragments thereof. Fully human antibodies can be made in mice that are transgenic for human immunoglobulin genes, e.g. according to Hogan, et al., "Manipulating the Mouse Embryo: A Laboratory Manual," Cold Spring Harbor Laboratory. Briefly, transgenic mice are immunized with purified immunogen. Spleen cells are harvested and fused to myeloma cells to produce hybridomas. Hybridomas are selected based on their ability to produce antibodies which bind to the immunogen. Fully human antibodies would reduce the immunogenicity of such antibodies in a human.

In one embodiment, an antibody for use in the instant invention is a bispecific or multispecific antibody. A bispecific antibody has binding sites for two different antigens within a single antibody polypeptide. Antigen binding may be simultaneous or sequential. Triomas and hybrid hybridomas are two examples of cell lines that can secrete bispecific antibodies. Examples of bispecific antibodies produced by a hybrid hybridoma or a trioma are disclosed in U.S. Pat. No. 4,474,893. Bispecific antibodies have been constructed by chemical means (Staerz et al. (1985) *Nature* 314:628, and Perez et al. (1985) *Nature* 316:354) and hybridoma technology (Staerz and Bevan (1986) *Proc. Natl. Acad. Sci. USA*, 83:1453, and Staerz and Bevan (1986) *Immunol. Today* 7:241). Bispecific antibodies are also described in U.S. Pat. No. 5,959,084. Fragments of bispecific antibodies are described in U.S. Pat. No. 5,798,229.

Bispecific agents can also be generated by making heterohybridomas by fusing hybridomas or other cells making different antibodies, followed by identification of clones producing and co-assembling both antibodies. They can also be generated by chemical or genetic conjugation of complete immunoglobulin chains or portions thereof such as Fab and Fv sequences. The antibody component can bind to a polypeptide or a fragment thereof of one or more biomarkers of the invention, including one or more biomarkers listed in Table 1, or a fragment thereof. In one embodiment, the bispecific antibody could specifically bind to both a polypeptide or a fragment thereof and its natural binding partner(s) or a fragment(s) thereof.

In another aspect of this invention, peptides or peptide mimetics can be used to antagonize the activity of one or more biomarkers of the invention, including one or more biomarkers listed in Table 1, or a fragment(s) thereof. In one embodiment, variants of one or more biomarkers listed in Table 1 which function as a modulating agent for the respective full length protein, can be identified by screening combinatorial libraries of mutants, e.g., truncation mutants, for antagonist activity. In one embodiment, a variegated library of variants is generated by combinatorial mutagenesis at the nucleic acid level and is encoded by a variegated gene library. A variegated library of variants can be produced, for instance, by enzymatically ligating a mixture of synthetic oligonucleotides into gene sequences such that a degenerate set of potential polypeptide sequences is expressible as individual polypeptides containing the set of polypeptide sequences therein. There are a variety of methods which can be used to produce libraries of polypeptide variants from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be performed in an automatic DNA synthesizer, and the synthetic gene then ligated into an appropriate expression

vector. Use of a degenerate set of genes allows for the provision, in one mixture, of all of the sequences encoding the desired set of potential polypeptide sequences. Methods for synthesizing degenerate oligonucleotides are known in the art (see, e.g., Narang, S. A. (1983) *Tetrahedron* 39:3; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477).

In addition, libraries of fragments of a polypeptide coding sequence can be used to generate a variegated population of polypeptide fragments for screening and subsequent selection of variants of a given polypeptide. In one embodiment, a library of coding sequence fragments can be generated by treating a double stranded PCR fragment of a polypeptide coding sequence with a nuclease under conditions wherein nicking occurs only about once per polypeptide, denaturing the double stranded DNA, renaturing the DNA to form double stranded DNA which can include sense/antisense pairs from different nicked products, removing single stranded portions from reformed duplexes by treatment with S1 nuclease, and ligating the resulting fragment library into an expression vector. By this method, an expression library can be derived which encodes N-terminal, C-terminal and internal fragments of various sizes of the polypeptide.

Several techniques are known in the art for screening gene products of combinatorial libraries made by point mutations or truncation, and for screening cDNA libraries for gene products having a selected property. Such techniques are adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of polypeptides. The most widely used techniques, which are amenable to high through-put analysis, for screening large gene libraries typically include cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates isolation of the vector encoding the gene whose product was detected. Recursive ensemble mutagenesis (REM), a technique which enhances the frequency of functional mutants in the libraries, can be used in combination with the screening assays to identify variants of interest (Arkin and Youvan (1992) *Proc. Natl. Acad. Sci. USA* 89:7811-7815; Delagrave et al. (1993) *Protein Eng.* 6(3): 327-331). In one embodiment, cell based assays can be exploited to analyze a variegated polypeptide library. For example, a library of expression vectors can be transfected into a cell line which ordinarily synthesizes one or more biomarkers of the invention, including one or more biomarkers listed in Table 1, or a fragment thereof. The transfected cells are then cultured such that the full length polypeptide and a particular mutant polypeptide are produced and the effect of expression of the mutant on the full length polypeptide activity in cell supernatants can be detected, e.g., by any of a number of functional assays. Plasmid DNA can then be recovered from the cells which score for inhibition, or alternatively, potentiation of full length polypeptide activity, and the individual clones further characterized.

Systematic substitution of one or more amino acids of a polypeptide amino acid sequence with a D-amino acid of the same type (e.g., D-lysine in place of L-lysine) can be used to generate more stable peptides. In addition, constrained peptides comprising a polypeptide amino acid sequence of interest or a substantially identical sequence variation can be generated by methods known in the art (Rizo and Gierasch (1992) *Annu. Rev. Biochem.* 61:387, incorporated herein by

reference); for example, by adding internal cysteine residues capable of forming intramolecular disulfide bridges which cyclize the peptide.

The amino acid sequences disclosed herein will enable those of skill in the art to produce polypeptides corresponding peptide sequences and sequence variants thereof. Such polypeptides can be produced in prokaryotic or eukaryotic host cells by expression of polynucleotides encoding the peptide sequence, frequently as part of a larger polypeptide. Alternatively, such peptides can be synthesized by chemical methods. Methods for expression of heterologous proteins in recombinant hosts, chemical synthesis of polypeptides, and in vitro translation are well known in the art and are described further in Maniatis et al. *Molecular Cloning: A Laboratory Manual* (1989), 2nd Ed., Cold Spring Harbor, N.Y.; Berger and Kimmel, *Methods in Enzymology*, Volume 152, *Guide to Molecular Cloning Techniques* (1987), Academic Press, Inc., San Diego, Calif.; Merrifield, J. (1969) *J. Am. Chem. Soc.* 91:501; Chaiken I. M. (1981) *CRC Crit. Rev. Biochem.* 11: 255; Kaiser et al. (1989) *Science* 243:187; Merrifield, B. (1986) *Science* 232:342; Kent, S. B. H. (1988) *Annu. Rev. Biochem.* 57:957; and Offord, R. E. (1980) *Semisynthetic Proteins*, Wiley Publishing, which are incorporated herein by reference).

Peptides can be produced, typically by direct chemical synthesis. Peptides can be produced as modified peptides, with nonpeptide moieties attached by covalent linkage to the N-terminus and/or C-terminus. In certain preferred embodiments, either the carboxy-terminus or the amino-terminus, or both, are chemically modified. The most common modifications of the terminal amino and carboxyl groups are acetylation and amidation, respectively. Amino-terminal modifications such as acylation (e.g., acetylation) or alkylation (e.g., methylation) and carboxy-terminal-modifications such as amidation, as well as other terminal modifications, including cyclization, can be incorporated into various embodiments of the invention. Certain amino-terminal and/or carboxy-terminal modifications and/or peptide extensions to the core sequence can provide advantageous physical, chemical, biochemical, and pharmacological properties, such as: enhanced stability, increased potency and/or efficacy, resistance to serum proteases, desirable pharmacokinetic properties, and others. Peptides disclosed herein can be used therapeutically to treat disease, e.g., by altering costimulation in a patient.

Peptidomimetics (Fauchere (1986) *Adv. Drug Res.* 15:29; Veber and Freidinger (1985) *TINS* p. 392; and Evans et al. (1987) *J. Med. Chem.* 30:1229, which are incorporated herein by reference) are usually developed with the aid of computerized molecular modeling. Peptide mimetics that are structurally similar to therapeutically useful peptides can be used to produce an equivalent therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biological or pharmacological activity), but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of: —CH₂NH—, —CH₂S—, —CH₂-CH₂-, —CH=CH— (cis and trans), —COCH₂-, —CH(OH)CH₂-, and —CH₂SO—, by methods known in the art and further described in the following references: Spatola, A. F. in “*Chemistry and Biochemistry of Amino Acids, Peptides, and Proteins*” Weinstein, B., ed., Marcel Dekker, New York, p. 267 (1983); Spatola, A. F., Vega Data (March 1983), Vol. 1, Issue 3, “Peptide Backbone Modifications” (general review); Morley, J. S. (1980) *Trends Pharm. Sci.* pp. 463-468 (general review); Hudson, D. et al. (1979) *Int. J. Pept. Prot. Res.* 14:177-185 (—CH₂NH—,

CH₂CH₂-); Spatola, A. F. et al. (1986) *Life Sci.* 38:1243-1249 (—CH₂-S); Hann, M. M. (1982) *J. Chem. Soc. Perkin Trans. I.* 307-314 (—CH—CH—, cis and trans); Almquist, R. G. et al. (190) *J. Med. Chem.* 23:1392-1398 (—COCH₂-); Jennings-White, C. et al. (1982) *Tetrahedron Lett.* 23:2533 (—COCH₂-); Szelke, M. et al. *European Appln.* EP 45665 (1982) CA: 97:39405 (1982)(—CH(OH)CH₂-); Holladay, M. W. et al. (1983) *Tetrahedron Lett.* (1983) 24:4401-4404 (—C(OH)CH₂-); and Hruby, V. J. (1982) *Life Sci.* (1982) 31:189-199 (—CH₂-S—); each of which is incorporated herein by reference. A particularly preferred non-peptide linkage is —CH₂NH—. Such peptide mimetics may have significant advantages over polypeptide embodiments, including, for example: more economical production, greater chemical stability, enhanced pharmacological properties (half-life, absorption, potency, efficacy, etc.), altered specificity (e.g., a broad-spectrum of biological activities), reduced antigenicity, and others. Labeling of peptidomimetics usually involves covalent attachment of one or more labels, directly or through a spacer (e.g., an amide group), to non-interfering position(s) on the peptidomimetic that are predicted by quantitative structure-activity data and/or molecular modeling. Such non-interfering positions generally are positions that do not form direct contacts with the macropolypeptides(s) to which the peptidomimetic binds to produce the therapeutic effect. Derivatization (e.g., labeling) of peptidomimetics should not substantially interfere with the desired biological or pharmacological activity of the peptidomimetic.

Also encompassed by the present invention are small molecules which can modulate (either enhance or inhibit) interactions, e.g., between biomarkers described herein or listed in Table 1 and their natural binding partners. The small molecules of the present invention can be obtained using any of the numerous approaches in combinatorial library methods known in the art, including: spatially addressable parallel solid phase or solution phase libraries; synthetic library methods requiring deconvolution; the ‘one-bead one-compound’ library method; and synthetic library methods using affinity chromatography selection. (Lam, K. S. (1997) *Anti-cancer Drug Des.* 12:145).

Examples of methods for the synthesis of molecular libraries can be found in the art, for example in: DeWitt et al. (1993) *Proc. Natl. Acad. Sci. USA* 90:6909; Erb et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:11422; Zuckermann et al. (1994) *J. Med. Chem.* 37:2678; Cho et al. (1993) *Science* 261:1303; Carrell et al. (1994) *Angew. Chem. Int. Ed. Engl.* 33:2059; Carell et al. (1994) *Angew. Chem. Int. Ed. Engl.* 33:2061; and in Gallop et al. (1994) *J. Med. Chem.* 37:1233.

Libraries of compounds can be presented in solution (e.g., Houghten (1992) *Biotechniques* 13:412-421), or on beads (Lam (1991) *Nature* 354:82-84), chips (Fodor (1993) *Nature* 364:555-556), bacteria (Ladner U.S. Pat. No. 5,223,409), spores (Ladner USP ’409), plasmids (Cull et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:1865-1869) or on phage (Scott and Smith (1990) *Science* 249:386-390); (Devlin (1990) *Science* 249:404-406); (Cwirla et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6378-6382); (Felici (1991) *J. Mol. Biol.* 222:301-310); (Ladner supra.). Compounds can be screened in cell based or non-cell based assays. Compounds can be screened in pools (e.g. multiple compounds in each testing sample) or as individual compounds.

The invention also relates to chimeric or fusion proteins of the biomarkers of the invention, including the biomarkers listed in Table 1, or fragments thereof. As used herein, a “chimeric protein” or “fusion protein” comprises one or more biomarkers of the invention, including one or more

biomarkers listed in Table 1, or a fragment thereof, operatively linked to another polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the respective biomarker. In a preferred embodiment, the fusion protein comprises at least one biologically active portion of one or more biomarkers of the invention, including one or more biomarkers listed in Table 1, or fragments thereof. Within the fusion protein, the term “operatively linked” is intended to indicate that the biomarker sequences and the non-biomarker sequences are fused in-frame to each other in such a way as to preserve functions exhibited when expressed independently of the fusion. The “another” sequences can be fused to the N-terminus or C-terminus of the biomarker sequences, respectively.

Such a fusion protein can be produced by recombinant expression of a nucleotide sequence encoding the first peptide and a nucleotide sequence encoding the second peptide. The second peptide may optionally correspond to a moiety that alters the solubility, affinity, stability or valency of the first peptide, for example, an immunoglobulin constant region. In another preferred embodiment, the first peptide consists of a portion of a biologically active molecule (e.g. the extracellular portion of the polypeptide or the ligand binding portion). The second peptide can include an immunoglobulin constant region, for example, a human C γ 1 domain or C γ 4 domain (e.g., the hinge, CH2 and CH3 regions of human IgC γ 1, or human IgC γ 4, see e.g., Capon et al. U.S. Pat. Nos. 5,116,964; 5,580,756; 5,844,095 and the like, incorporated herein by reference). Such constant regions may retain regions which mediate effector function (e.g. Fc receptor binding) or may be altered to reduce effector function. A resulting fusion protein may have altered solubility, binding affinity, stability and/or valency (i.e., the number of binding sites available per polypeptide) as compared to the independently expressed first peptide, and may increase the efficiency of protein purification. Fusion proteins and peptides produced by recombinant techniques can be secreted and isolated from a mixture of cells and medium containing the protein or peptide. Alternatively, the protein or peptide can be retained cytoplasmically and the cells harvested, lysed and the protein isolated. A cell culture typically includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. Protein and peptides can be isolated from cell culture media, host cells, or both using techniques known in the art for purifying proteins and peptides. Techniques for transfecting host cells and purifying proteins and peptides are known in the art.

Preferably, a fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992).

Particularly preferred Ig fusion proteins include the extracellular domain portion or variable region-like domain of PD-L1, TIM-3, LAG-3, or other biomarker listed in Table 1, coupled to an immunoglobulin constant region (e.g., the Fc region). The immunoglobulin constant region may contain genetic modifications which reduce or eliminate effector activity inherent in the immunoglobulin structure. For example, DNA encoding the extracellular portion of a polypeptide of interest can be joined to DNA encoding the hinge, CH2 and CH3 regions of human IgG γ 1 and/or IgG γ 4 modified by site directed mutagenesis, e.g., as taught in WO 97/28267.

In another embodiment, the fusion protein contains a heterologous signal sequence at its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of a polypeptide can be increased through use of a heterologous signal sequence.

The fusion proteins of the invention can be used as immunogens to produce antibodies in a subject. Such antibodies may be used to purify the respective natural polypeptides from which the fusion proteins were generated, or in screening assays to identify polypeptides which inhibit the interactions between one or more biomarkers polypeptide or a fragment thereof and its natural binding partner(s) or a fragment(s) thereof.

Also provided herein are compositions comprising one or more nucleic acids comprising or capable of expressing at least 1, 2, 3, 4, 5, 10, 20 or more small nucleic acids or antisense oligonucleotides or derivatives thereof, wherein said small nucleic acids or antisense oligonucleotides or derivatives thereof in a cell specifically hybridize (e.g., bind) under cellular conditions, with cellular nucleic acids (e.g., small non-coding RNAs such as miRNAs, pre-miRNAs, pri-miRNAs, miRNA*, anti-miRNA, a miRNA binding site, a variant and/or functional variant thereof, cellular mRNAs or a fragments thereof). In one embodiment, expression of the small nucleic acids or antisense oligonucleotides or derivatives thereof in a cell can inhibit expression or biological activity of cellular nucleic acids and/or proteins, e.g., by inhibiting transcription, translation and/or small nucleic acid processing of, for example, one or more biomarkers of the invention, including one or more biomarkers listed in Table 1, or fragment(s) thereof. In one embodiment, the small nucleic acids or antisense oligonucleotides or derivatives thereof are small RNAs (e.g., microRNAs) or complements of small RNAs. In another embodiment, the small nucleic acids or antisense oligonucleotides or derivatives thereof can be single or double stranded and are at least six nucleotides in length and are less than about 1000, 900, 800, 700, 600, 500, 400, 300, 200, 100, 50, 40, 30, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, or 10 nucleotides in length. In another embodiment, a composition may comprise a library of nucleic acids comprising or capable of expressing small nucleic acids or antisense oligonucleotides or derivatives thereof, or pools of said small nucleic acids or antisense oligonucleotides or derivatives thereof. A pool of nucleic acids may comprise about 2-5, 5-10, 10-20, 10-30 or more nucleic acids comprising or capable of expressing small nucleic acids or antisense oligonucleotides or derivatives thereof.

In one embodiment, binding may be by conventional base pair complementarity, or, for example, in the case of binding to DNA duplexes, through specific interactions in the major groove of the double helix. In general, “antisense” refers to the range of techniques generally employed in the art, and includes any process that relies on specific binding to oligonucleotide sequences.

It is well known in the art that modifications can be made to the sequence of a miRNA or a pre-miRNA without disrupting miRNA activity. As used herein, the term “functional variant” of a miRNA sequence refers to an oligo-nucleotide sequence that varies from the natural miRNA sequence, but retains one or more functional characteristics of the miRNA (e.g. cancer cell proliferation inhibition, induction of cancer cell apoptosis, enhancement of cancer cell susceptibility to chemotherapeutic agents, specific miRNA target inhibition). In some embodiments, a functional variant of a miRNA sequence retains all of the functional characteristics of the miRNA. In certain embodiments, a functional variant of a miRNA has a nucleobase sequence that is at least about 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99% identical to the miRNA or precursor thereof over a region of about 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 100 or more nucleobases, or that the functional variant hybridizes to the complement of the miRNA or precursor thereof under stringent hybridization conditions. Accordingly, in certain embodiments the nucleobase sequence of a functional variant is capable of hybridizing to one or more target sequences of the miRNA.

miRNAs and their corresponding stem-loop sequences described herein may be found in miRBase, an online searchable database of miRNA sequences and annotation, found on the world wide web at microrna.sanger.ac.uk. Entries in the miRBase Sequence database represent a predicted hairpin portion of a miRNA transcript (the stem-loop), with information on the location and sequence of the mature miRNA sequence. The miRNA stem-loop sequences in the database are not strictly precursor miRNAs (pre-miRNAs), and may in some instances include the pre-miRNA and some flanking sequence from the presumed primary transcript. The miRNA nucleobase sequences described herein encompass any version of the miRNA, including the sequences described in Release 10.0 of the miRBase sequence database and sequences described in any earlier Release of the miRBase sequence database. A sequence database release may result in the re-naming of certain miRNAs. A sequence database release may result in a variation of a mature miRNA sequence.

In some embodiments, miRNA sequences of the invention may be associated with a second RNA sequence that may be located on the same RNA molecule or on a separate RNA molecule as the miRNA sequence. In such cases, the miRNA sequence may be referred to as the active strand, while the second RNA sequence, which is at least partially complementary to the miRNA sequence, may be referred to as the complementary strand. The active and complementary strands are hybridized to create a double-stranded RNA that is similar to a naturally occurring miRNA precursor. The activity of a miRNA may be optimized by maximizing uptake of the active strand and minimizing uptake of the complementary strand by the miRNA protein complex that regulates gene translation. This can be done through modification and/or design of the complementary strand.

In some embodiments, the complementary strand is modified so that a chemical group other than a phosphate or hydroxyl at its 5' terminus. The presence of the 5' modification apparently eliminates uptake of the complementary strand and subsequently favors uptake of the active strand by the miRNA protein complex. The 5' modification can be any of a variety of molecules known in the art, including NH₂, NHCOCH₃, and biotin.

In another embodiment, the uptake of the complementary strand by the miRNA pathway is reduced by incorporating nucleotides with sugar modifications in the first 2-6 nucleotides of the complementary strand. It should be noted that such sugar modifications can be combined with the 5' terminal modifications described above to further enhance miRNA activities.

In some embodiments, the complementary strand is designed so that nucleotides in the 3' end of the complementary strand are not complementary to the active strand. This results in double-strand hybrid RNAs that are stable at the 3' end of the active strand but relatively unstable at the 5' end of the active strand. This difference in stability enhances the uptake of the active strand by the miRNA pathway, while reducing uptake of the complementary strand, thereby enhancing miRNA activity.

Small nucleic acid and/or antisense constructs of the methods and compositions presented herein can be delivered, for example, as an expression plasmid which, when transcribed in the cell, produces RNA which is complementary to at least a unique portion of cellular nucleic acids (e.g., small RNAs, mRNA, and/or genomic DNA). Alternatively, the small nucleic acid molecules can produce RNA which encodes mRNA, miRNA, pre-miRNA, pri-miRNA, miRNA*, anti-miRNA, or a miRNA binding site, or a variant thereof. For example, selection of plasmids suitable for expressing the miRNAs, methods for inserting nucleic acid sequences into the plasmid, and methods of delivering the recombinant plasmid to the cells of interest are within the skill in the art. See, for example, Zeng et al. (2002) *Mol. Cell* 9:1327-1333; Tuschl (2002), *Nat. Biotechnol.* 20:446-448; Brummelkamp et al. (2002) *Science* 296:550-553; Miyagishi et al. (2002) *Nat. Biotechnol.* 20:497-500; Paddison et al. (2002) *Genes Dev.* 16:948-958; Lee et al. (2002) *Nat. Biotechnol.* 20:500-505; and Paul et al. (2002) *Nat. Biotechnol.* 20:505-508, the entire disclosures of which are herein incorporated by reference.

Alternatively, small nucleic acids and/or antisense constructs are oligonucleotide probes that are generated ex vivo and which, when introduced into the cell, results in hybridization with cellular nucleic acids. Such oligonucleotide probes are preferably modified oligonucleotides that are resistant to endogenous nucleases, e.g., exonucleases and/or endonucleases, and are therefore stable in vivo. Exemplary nucleic acid molecules for use as small nucleic acids and/or antisense oligonucleotides are phosphoramidate, phosphothioate and methylphosphonate analogs of DNA (see also U.S. Pat. Nos. 5,176,996; 5,264,564; and 5,256,775). Additionally, general approaches to constructing oligomers useful in antisense therapy have been reviewed, for example, by Van der Krol et al. (1988) *BioTechniques* 6:958-976; and Stein et al. (1988) *Cancer Res* 48:2659-2668.

Antisense approaches may involve the design of oligonucleotides (either DNA or RNA) that are complementary to cellular nucleic acids (e.g., complementary to biomarkers listed in Table 1). Absolute complementarity is not required. In the case of double-stranded antisense nucleic acids, a single strand of the duplex DNA may thus be tested, or triplex formation may be assayed. The ability to hybridize will depend on both the degree of complementarity and the length of the antisense nucleic acid. Generally, the longer the hybridizing nucleic acid, the more base mismatches with a nucleic acid (e.g., RNA) it may contain and still form a stable duplex (or triplex, as the case may be). One skilled in the art can ascertain a tolerable degree of mismatch by use of standard procedures to determine the melting point of the hybridized complex.

Oligonucleotides that are complementary to the 5' end of the mRNA, e.g., the 5' untranslated sequence up to and including the AUG initiation codon, should work most efficiently at inhibiting translation. However, sequences complementary to the 3' untranslated sequences of mRNAs have recently been shown to be effective at inhibiting translation of mRNAs as well (Wagner (1994) *Nature* 372: 333). Therefore, oligonucleotides complementary to either the 5' or 3' untranslated, non-coding regions of genes could be used in an antisense approach to inhibit translation of endogenous mRNAs. Oligonucleotides complementary to the 5' untranslated region of the mRNA may include the complement of the AUG start codon. Antisense oligonucleotides complementary to mRNA coding regions are less efficient inhibitors of translation but could also be used in accordance with the methods and compositions presented herein. Whether designed to hybridize to the 5', 3' or coding region of cellular mRNAs, small nucleic acids and/or antisense nucleic acids should be at least six nucleotides in length, and can be less than about 1000, 900, 800, 700, 600, 500, 400, 300, 200, 100, 50, 40, 30, 25, 24, 23, 22, 21, 20, 19, 18, 17, 16, 15, or 10 nucleotides in length.

Regardless of the choice of target sequence, it is preferred that in vitro studies are first performed to quantitate the ability of the antisense oligonucleotide to inhibit gene expression. In one embodiment these studies utilize controls that distinguish between antisense gene inhibition and non-specific biological effects of oligonucleotides. In another embodiment these studies compare levels of the target nucleic acid or protein with that of an internal control nucleic acid or protein. Additionally, it is envisioned that results obtained using the antisense oligonucleotide are compared with those obtained using a control oligonucleotide. It is preferred that the control oligonucleotide is of approximately the same length as the test oligonucleotide and that the nucleotide sequence of the oligonucleotide differs from the antisense sequence no more than is necessary to prevent specific hybridization to the target sequence.

Small nucleic acids and/or antisense oligonucleotides can be DNA or RNA or chimeric mixtures or derivatives or modified versions thereof, single-stranded or double-stranded. Small nucleic acids and/or antisense oligonucleotides can be modified at the base moiety, sugar moiety, or phosphate backbone, for example, to improve stability of the molecule, hybridization, etc., and may include other appended groups such as peptides (e.g., for targeting host cell receptors), or agents facilitating transport across the cell membrane (see, e.g., Letsinger et al. (1989) *Proc. Natl. Acad. Sci. U.S.A.* 86:6553-6556; Lemaitre et al. (1987) *Proc. Natl. Acad. Sci. U.S.A.* 84:648-652; PCT Publication No. WO88/09810) or the blood-brain barrier (see, e.g., PCT Publication No. WO89/10134), hybridization-triggered cleavage agents. (See, e.g., Krol et al. (1988) *BioTech.* 6:958-976) or intercalating agents. (See, e.g., Zon (1988) *Pharm. Res.* 5:539-549). To this end, small nucleic acids and/or antisense oligonucleotides may be conjugated to another molecule, e.g., a peptide, hybridization triggered cross-linking agent, transport agent, hybridization-triggered cleavage agent, etc.

Small nucleic acids and/or antisense oligonucleotides may comprise at least one modified base moiety which is selected from the group including but not limited to 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xantine, 4-acetylcytosine, 5-(carboxyhydroxyethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine,

1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methyl ester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Small nucleic acids and/or antisense oligonucleotides may also comprise at least one modified sugar moiety selected from the group including but not limited to arabinose, 2-fluoroarabinose, xylulose, and hexose.

In certain embodiments, a compound comprises an oligonucleotide (e.g., a miRNA or miRNA encoding oligonucleotide) conjugated to one or more moieties which enhance the activity, cellular distribution or cellular uptake of the resulting oligonucleotide. In certain such embodiments, the moiety is a cholesterol moiety (e.g., antagomirs) or a lipid moiety or liposome conjugate. Additional moieties for conjugation include carbohydrates, phospholipids, biotin, phenazine, folate, phenanthridine, anthraquinone, acridine, fluoresceins, rhodamines, coumarins, and dyes. In certain embodiments, a conjugate group is attached directly to the oligonucleotide. In certain embodiments, a conjugate group is attached to the oligonucleotide by a linking moiety selected from amino, hydroxyl, carboxylic acid, thiol, unsaturations (e.g., double or triple bonds), 8-amino-3,6-dioxaoctanoic acid (ADO), succinimidyl 4-(N-maleimidomethyl) cyclohexane-1-carboxylate (SMCC), 6-amino-hexanoic acid (AHEX or AHA), substituted C1-C10 alkyl, substituted or unsubstituted C2-C10 alkenyl, and substituted or unsubstituted C2-C10 alkynyl. In certain such embodiments, a substituent group is selected from hydroxyl, amino, alkoxy, carboxy, benzyl, phenyl, nitro, thiol, thioalkoxy, halogen, alkyl, aryl, alkenyl and alkynyl.

In certain such embodiments, the compound comprises the oligonucleotide having one or more stabilizing groups that are attached to one or both termini of the oligonucleotide to enhance properties such as, for example, nuclease stability. Included in stabilizing groups are cap structures. These terminal modifications protect the oligonucleotide from exonuclease degradation, and can help in delivery and/or localization within a cell. The cap can be present at the 5'-terminus (5'-cap), or at the 3'-terminus (3'-cap), or can be present on both termini. Cap structures include, for example, inverted deoxy abasic caps.

Suitable cap structures include a 4',5'-methylene nucleotide, a 1-(beta-D-erythrofuransyl) nucleotide, a 4'-thio nucleotide, a carbocyclic nucleotide, a 1,5-anhydrohexitol nucleotide, an L-nucleotide, an alpha-nucleotide, a modified base nucleotide, a phosphorodithioate linkage, a threopentofuransyl nucleotide, an acyclic 3',4'-seco nucleotide, an acyclic 3,4-dihydroxybutyl nucleotide, an acyclic 3,5-dihydroxypentyl nucleotide, a 3'-3'-inverted nucleotide moiety, a 3'-3'-inverted abasic moiety, a 3'-2'-inverted nucleotide moiety, a 3'-2'-inverted abasic moiety, a 1,4-butanediol phosphate, a 3'-phosphoramidate, a hexylphosphate, an aminohexyl phosphate, a 3'-phosphate, a 3'-phosphorothioate, a phosphorodithioate, a bridging methylphosphonate moiety, and a non-bridging methylphosphonate moiety 5'-aminoalkyl phosphate, a 1,3-diamino-2-propyl phosphate, 3-aminopropyl phosphate, a 6-aminohexyl phosphate, a 1,2-aminododecyl phosphate, a hydroxypropyl phosphate, a 5'-5'-inverted nucleotide moiety, a 5'-5'-inverted abasic moiety, a

5'-phosphoramidate, a 5'-phosphorothioate, a 5'-amino, a bridging and/or non-bridging 5'-phosphoramidate, a phosphorothioate, and a 5'-mercapto moiety.

Small nucleic acids and/or antisense oligonucleotides can also contain a neutral peptide-like backbone. Such molecules are termed peptide nucleic acid (PNA)-oligomers and are described, e.g., in Perry-O'Keefe et al. (1996) Proc. Natl. Acad. Sci. U.S.A. 93:14670 and in Eglom et al. (1993) Nature 365:566. One advantage of PNA oligomers is their capability to bind to complementary DNA essentially independently from the ionic strength of the medium due to the neutral backbone of the DNA. In yet another embodiment, small nucleic acids and/or antisense oligonucleotides comprises at least one modified phosphate backbone selected from the group consisting of a phosphorothioate, a phosphorodithioate, a phosphoramidothioate, a phosphoramidate, a phosphordiamidate, a methylphosphonate, an alkyl phosphotriester, and a formacetal or analog thereof.

In a further embodiment, small nucleic acids and/or antisense oligonucleotides are α -anomeric oligonucleotides. An α -anomeric oligonucleotide forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual β -units, the strands run parallel to each other (Gautier et al. (1987) Nucl. Acids Res. 15:6625-6641). The oligonucleotide is a 2'-O-methylribonucleotide (Inoue et al. (1987) Nucl. Acids Res. 15:6131-6148), or a chimeric RNA-DNA analogue (Inoue et al. (1987) FEBS Lett. 215:327-330).

Small nucleic acids and/or antisense oligonucleotides of the methods and compositions presented herein may be synthesized by standard methods known in the art, e.g., by use of an automated DNA synthesizer (such as are commercially available from Biosearch, Applied Biosystems, etc.). As examples, phosphorothioate oligonucleotides may be synthesized by the method of Stein et al. (1988) Nucl. Acids Res. 16:3209, methylphosphonate oligonucleotides can be prepared by use of controlled pore glass polymer supports (Sarin et al. (1988) Proc. Natl. Acad. Sci. U.S.A. 85:7448-7451), etc. For example, an isolated miRNA can be chemically synthesized or recombinantly produced using methods known in the art. In some instances, miRNA are chemically synthesized using appropriately protected ribonucleoside phosphoramidites and a conventional DNA/RNA synthesizer. Commercial suppliers of synthetic RNA molecules or synthesis reagents include, e.g., Prologo (Hamburg, Germany), Dharmacon Research (Lafayette, Colo., USA), Pierce Chemical (part of Perbio Science, Rockford, Ill., USA), Glen Research (Sterling, Va., USA), ChemGenes (Ashland, Mass., USA), Cruachem (Glasgow, UK), and Exiqon (Vedbaek, Denmark).

Small nucleic acids and/or antisense oligonucleotides can be delivered to cells in vivo. A number of methods have been developed for delivering small nucleic acids and/or antisense oligonucleotides DNA or RNA to cells; e.g., antisense molecules can be injected directly into the tissue site, or modified antisense molecules, designed to target the desired cells (e.g., antisense linked to peptides or antibodies that specifically bind receptors or antigens expressed on the target cell surface) can be administered systematically.

In one embodiment, small nucleic acids and/or antisense oligonucleotides may comprise or be generated from double stranded small interfering RNAs (siRNAs), in which sequences fully complementary to cellular nucleic acids (e.g. mRNAs) sequences mediate degradation or in which sequences incompletely complementary to cellular nucleic acids (e.g., mRNAs) mediate translational repression when expressed within cells. In another embodiment, double

stranded siRNAs can be processed into single stranded antisense RNAs that bind single stranded cellular RNAs (e.g., microRNAs) and inhibit their expression. RNA interference (RNAi) is the process of sequence-specific, post-transcriptional gene silencing in animals and plants, initiated by double-stranded RNA (dsRNA) that is homologous in sequence to the silenced gene. *in vivo*, long dsRNA is cleaved by ribonuclease III to generate 21- and 22-nucleotide siRNAs. It has been shown that 21-nucleotide siRNA duplexes specifically suppress expression of endogenous and heterologous genes in different mammalian cell lines, including human embryonic kidney (293) and HeLa cells (Elbashir et al. (2001) Nature 411:494-498). Accordingly, translation of a gene in a cell can be inhibited by contacting the cell with short double stranded RNAs having a length of about 15 to 30 nucleotides or of about 18 to 21 nucleotides or of about 19 to 21 nucleotides. Alternatively, a vector encoding for such siRNAs or short hairpin RNAs (shRNAs) that are metabolized into siRNAs can be introduced into a target cell (see, e.g., McManus et al. (2002) RNA 8:842; Xia et al. (2002) Nature Biotechnology 20:1006; and Brummelkamp et al. (2002) Science 296:550). Vectors that can be used are commercially available, e.g., from OligoEngine under the name pSuper RNAi System™.

Ribozyme molecules designed to catalytically cleave cellular mRNA transcripts can also be used to prevent translation of cellular mRNAs and expression of cellular polypeptides, or both (See, e.g., PCT International Publication WO90/11364, published Oct. 4, 1990; Sarver et al. (1990) Science 247:1222-1225 and U.S. Pat. No. 5,093,246). While ribozymes that cleave mRNA at site specific recognition sequences can be used to destroy cellular mRNAs, the use of hammerhead ribozymes is preferred. Hammerhead ribozymes cleave mRNAs at locations dictated by flanking regions that form complementary base pairs with the target mRNA. The sole requirement is that the target mRNA have the following sequence of two bases: 5'-UG-3'. The construction and production of hammerhead ribozymes is well known in the art and is described more fully in Haseloff and Gerlach (1988) Nature 334:585-591. The ribozyme may be engineered so that the cleavage recognition site is located near the 5' end of cellular mRNAs; i.e., to increase efficiency and minimize the intracellular accumulation of non-functional mRNA transcripts.

The ribozymes of the methods presented herein also include RNA endoribonucleases (hereinafter "Cech-type ribozymes") such as the one which occurs naturally in *Tetrahymena thermophila* (known as the IVS, or L-19 IVS RNA) and which has been extensively described by Thomas Cech and collaborators (Zaug et al. (1984) Science 224:574-578; Zaug et al. (1986) Science 231:470-475; Zaug et al. (1986) Nature 324:429-433; WO 88/04300; and Been et al. (1986) Cell 47:207-216). The Cech-type ribozymes have an eight base pair active site which hybridizes to a target RNA sequence whereafter cleavage of the target RNA takes place. The methods and compositions presented herein encompasses those Cech-type ribozymes which target eight base-pair active site sequences that are present in cellular genes.

As in the antisense approach, the ribozymes can be composed of modified oligonucleotides (e.g., for improved stability, targeting, etc.). A preferred method of delivery involves using a DNA construct "encoding" the ribozyme under the control of a strong constitutive pol III or pol II promoter, so that transfected cells will produce sufficient quantities of the ribozyme to destroy endogenous cellular messages and inhibit translation. Because ribozymes unlike

antisense molecules, are catalytic, a lower intracellular concentration is required for efficiency.

Nucleic acid molecules to be used in triple helix formation for the inhibition of transcription of cellular genes are preferably single stranded and composed of deoxyribonucleotides. The base composition of these oligonucleotides should promote triple helix formation via Hoogsteen base pairing rules, which generally require sizable stretches of either purines or pyrimidines to be present on one strand of a duplex. Nucleotide sequences may be pyrimidine-based, which will result in TAT and CGC triplets across the three associated strands of the resulting triple helix. The pyrimidine-rich molecules provide base complementarity to a purine-rich region of a single strand of the duplex in a parallel orientation to that strand. In addition, nucleic acid molecules may be chosen that are purine-rich, for example, containing a stretch of G residues. These molecules will form a triple helix with a DNA duplex that is rich in GC pairs, in which the majority of the purine residues are located on a single strand of the targeted duplex, resulting in CGC triplets across the three strands in the triplex.

Alternatively, the potential sequences that can be targeted for triple helix formation may be increased by creating a so called "switchback" nucleic acid molecule. Switchback molecules are synthesized in an alternating 5'-3', 3'-5' manner, such that they base pair with first one strand of a duplex and then the other, eliminating the necessity for a sizable stretch of either purines or pyrimidines to be present on one strand of a duplex.

Small nucleic acids (e.g., miRNAs, pre-miRNAs, pri-miRNAs, miRNA*, anti-miRNA, or a miRNA binding site, or a variant thereof), antisense oligonucleotides, ribozymes, and triple helix molecules of the methods and compositions presented herein may be prepared by any method known in the art for the synthesis of DNA and RNA molecules. These include techniques for chemically synthesizing oligodeoxyribonucleotides and oligoribonucleotides well known in the art such as for example solid phase phosphoramidite chemical synthesis. Alternatively, RNA molecules may be generated by *in vitro* and *in vivo* transcription of DNA sequences encoding the antisense RNA molecule. Such DNA sequences may be incorporated into a wide variety of vectors which incorporate suitable RNA polymerase promoters such as the T7 or SP6 polymerase promoters. Alternatively, antisense cDNA constructs that synthesize antisense RNA constitutively or inducibly, depending on the promoter used, can be introduced stably into cell lines.

Moreover, various well-known modifications to nucleic acid molecules may be introduced as a means of increasing intracellular stability and half-life. Possible modifications include but are not limited to the addition of flanking sequences of ribonucleotides or deoxyribonucleotides to the 5' and/or 3' ends of the molecule or the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages within the oligodeoxyribonucleotide backbone. One of skill in the art will readily understand that polypeptides, small nucleic acids, and antisense oligonucleotides can be further linked to another peptide or polypeptide (e.g., a heterologous peptide), e.g., that serves as a means of protein detection. Non-limiting examples of label peptide or polypeptide moieties useful for detection in the invention include, without limitation, suitable enzymes such as horseradish peroxidase, alkaline phosphatase, beta-galactosidase, or acetylcholinesterase; epitope tags, such as FLAG, MYC, HA, or HIS tags; fluorophores such as green fluorescent protein; dyes; radioisotopes; digoxigenin; biotin; antibodies; polymers; as well as others known in the art, for

example, in *Principles of Fluorescence Spectroscopy*, Joseph R. Lakowicz (Editor), Plenum Pub Corp, 2nd edition (July 1999).

The modulatory agents described herein (e.g., antibodies, small molecules, peptides, fusion proteins, or small nucleic acids) can be incorporated into pharmaceutical compositions and administered to a subject *in vivo*. The compositions may contain a single such molecule or agent or any combination of agents described herein. "Single active agents" described herein can be combined with other pharmacologically active compounds ("second active agents") known in the art according to the methods and compositions provided herein. It is believed that certain combinations work synergistically in the treatment of conditions that would benefit from the modulation of immune responses. Second active agents can be large molecules (e.g., proteins) or small molecules (e.g., synthetic inorganic, organometallic, or organic molecules). For example, anti-PD-L1 and anti-TIM-3 antibodies can be further combined with anti-LAG-3, anti-PD-1, anti-PD-L2, anti-CTLA4, etc. antibodies or combinations thereof.

Examples of large molecule active agents include, but are not limited to, hematopoietic growth factors, cytokines, and monoclonal and polyclonal antibodies. Typical large molecule active agents are biological molecules, such as naturally occurring or artificially made proteins. Proteins that are particularly useful in this invention include proteins that stimulate the survival and/or proliferation of hematopoietic precursor cells and immunologically active poietic cells *in vitro* or *in vivo*. Others stimulate the division and differentiation of committed erythroid progenitors in cells *in vitro* or *in vivo*. Particular proteins include, but are not limited to: interleukins, such as IL-2 (including recombinant IL-2 ("rIL2") and canarypox IL-2), IL-10, IL-12, and IL-18; interferons, such as interferon alfa-2a, interferon alfa-2b, interferon alpha-n1, interferon alpha-n3, interferon beta-1a, and interferon gamma-1b; GM-CSF and GM-CSF; and EPO.

Particular proteins that can be used in the methods and compositions provided herein include, but are not limited to: filgrastim, which is sold in the United States under the trade name Neupogen® (Amgen, Thousand Oaks, Calif.); sargramostim, which is sold in the United States under the trade name Leukine® (Immunex, Seattle, Wash.); and recombinant EPO, which is sold in the United States under the trade name Epogen® (Amgen, Thousand Oaks, Calif.). Recombinant and mutated forms of GM-CSF can be prepared as described in U.S. Pat. Nos. 5,391,485; 5,393,870; and 5,229,496; all of which are incorporated herein by reference. Recombinant and mutated forms of G-CSF can be prepared as described in U.S. Pat. Nos. 4,810,643; 4,999,291; 5,528,823; and 5,580,755; all of which are incorporated herein by reference.

Similarly, chemotherapeutic agents are well known in the art. For example, chemotherapeutic agents include alkylating agents such as thiotepa and cyclophosphamide (Cytosan™); alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzodopa, carboquone, meturedopa, and uredopa; emylerumines and memylamelamines including alfretamine, triemylenemelumine, triethylenephosphoramidate, triethylenethiophosphoramidate, and trimethylolmelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including synthetic analogue topotecan); bryostatin; callistatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CBI-TMI); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mus-

tards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosoureas such as carmustine, chlorozotocin, foremustine, lomustine, nimustine, ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammall and calicheamicin phili); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromomorphores), aclacinomysins, actinomycin, anthracycline, azaserine, bleomycins, cactinomycin, carabycin, carminomycin, carzinophilin, chromomycins, dactinomycin, daunorubicin, detorubicin, 6-diazo-5-oxo-L-norleucine, doxorubicin (Adramycin™) (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil (5-FU); folic acid analogues such as demopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogues such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitio stanol, testolactone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replinisher such as frolic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; hestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformthine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidamine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidamol; nitracrine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK™; razoxane; rhizoxin; sizofiran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-tricyclopropylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethane; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiopeta; taxoids, e.g., paclitaxel (Taxol™, Bristol Meyers Squibb Oncology, Princeton, N.J.) and docetaxel (Taxotere™, Rhone-Poulenc Rorer, Antony, France); chlorambucil; gemcitabine (Gemzar™); 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vancristine; vinorelbine (Navelbine™); novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeoloda; ibandronate; CPT-11; topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; and pharmaceutically acceptable salts, acids or derivatives of any of the above. Also included in the definition of "chemotherapeutic agent" are anti-hormonal agents that act to regulate or inhibit hormone action on tumors such as anti-estrogens and selective estrogen receptor modulators (SERMs), including, for example, tamoxifen (including Nolvadex™), raloxifene, droloxifene, 4-hydroxytamoxifen, trioxifene, keoxifene, LY117018, onapristone, and toremifene (Fareston™); inhibitors of the enzyme aromatase, which regulates estrogen production in the adrenal glands, such as, for example,

4(5)-imidazoles, aminoglutethimide, megestrol acetate (Megace™), exemestane, formestane, fadrozole, vorozole (Rivisor™), letrozole (Femara™), and anastrozole (Arimidex™); and anti-androgens such as flutamide, nilutamide, bicalutamide, leuprorelin, and goserelin; and pharmaceutically acceptable salts, acids or derivatives of any of the above. In some embodiments, the inhibitor downregulates Rac 1 output. Additional examples of chemotherapeutic and other anti-cancer agents are described in US Pat. Pubs. 2013/0239239 and 2009/0053224.

b. Pharmaceutical Compositions

Agents that modulate (e.g., inhibit or block) the function of PD-1 or PD-L1 and LAG-3, CTLA-4 or TIM-3, including, e.g., blocking antibodies, peptides, fusion proteins, or small molecules, can be incorporated into pharmaceutical compositions suitable for administration to a subject. Such compositions typically comprise the antibody, peptide, fusion protein or small molecule and a pharmaceutically acceptable carrier. As used herein, "pharmaceutically acceptable carrier" is intended to include any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions. A pharmaceutical composition of the invention is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, e.g., intravenous, intradermal, subcutaneous, oral (e.g., inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerin, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, N.J.) or phosphate buffered saline (PBS). In all cases, the composition should be sterile and should be fluid to the extent that easy syringeability exists. It must be stable under the conditions of manufacture and storage and should be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Pre-

vention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. They can be enclosed in gelatin capsules or compressed into tablets. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash, wherein the compound in the fluid carrier is applied orally and swished and expectorated or swallowed. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the compounds are delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, e.g., a gas such as carbon dioxide, or a nebulizer.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The compounds can also be prepared in the form of suppositories (e.g., with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

In one embodiment, modulatory agents are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be

used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Methods for preparation of such formulations should be apparent to those skilled in the art. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Pat. No. 4,522,811.

It is especially advantageous to formulate oral or parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. Dosage unit form as used herein refers to physically discrete units suited as unitary dosages for the subject to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by, and directly dependent on, the unique characteristics of the active compound, the particular therapeutic effect to be achieved, and the limitations inherent in the art of compounding such an active compound for the treatment of individuals.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD50 (the dose lethal to 50% of the population) and the ED50 (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD50/ED50. Compounds which exhibit large therapeutic indices are preferred. While compounds that exhibit toxic side effects can be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED50 with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC50 (i.e., the concentration of the test compound which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma can be measured, for example, by high performance liquid chromatography.

The above described modulating agents may be administered in the form of expressible nucleic acids which encode said agents. Such nucleic acids and compositions in which they are contained, are also encompassed by the present invention. For instance, the nucleic acid molecules of the invention can be inserted into vectors and used as gene therapy vectors. Gene therapy vectors can be delivered to a subject by, for example, intravenous injection, local administration (see U.S. Pat. No. 5,328,470) or by stereotactic injection (see e.g., Chen et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:3054-3057). The pharmaceutical preparation of the gene therapy vector can include the gene therapy vector in

an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery vector can be produced intact from recombinant cells, e.g., retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The pharmaceutical compositions can be included in a container, pack, or dispenser together with instructions for administration.

c. Prophylactic Methods

In one aspect, the present invention provides a method for preventing in a subject, a cancer, such as a hematologic cancer like multiple myeloma, associated with a less than desirable immune response. Subjects at risk for such a disease can be identified, for example, by any or a combination of diagnostic or prognostic assays known in the art. Administration of a prophylactic agent(s) can occur prior to the manifestation of symptoms associated with an unwanted or less than desirable immune response. The appropriate agent(s) used for treatment (e.g. antibodies, peptides, fusion proteins or small molecules) can be determined based on clinical indications and can be identified using diagnostic assays well known in the art, as well as those described herein.

d. Therapeutic Methods

Another aspect of the invention pertains to therapeutic methods of modulating an immune response, e.g., by inhibiting or blocking the function of PD-1 or PD-L1 and LAG-3, CTLA-4 or TIM-3.

Modulatory methods of the present invention involve contacting a cell with an agent that inhibits or blocks the function of PD-1 or PD-L1 and LAG-3, CTLA-4 or TIM-3. Exemplary agents useful in such methods are described above. Such agents can be administered in vitro or ex vivo (e.g., by contacting the cell with the agent) or, alternatively, in vivo (e.g., by administering the agent to a subject). As such, the present invention provides methods useful for treating an individual afflicted with a cancer, such as a hematologic cancer like multiple myeloma.

Agents that upregulate immune responses can be in the form of enhancing an existing immune response or eliciting an initial immune response. Thus, enhancing an immune response using the subject compositions and methods is useful for treating cancer, but can also be useful for treating an infectious disease (e.g., bacteria, viruses, or parasites), a parasitic infection, and an immunosuppressive disease.

Immune responses can also be enhanced in an infected patient through an ex vivo approach, for instance, by removing immune cells from the patient, contacting immune cells in vitro with an agent described herein and reintroducing the in vitro stimulated immune cells into the patient.

In certain instances, it may be desirable to further administer other agents that upregulate immune responses, for example, forms of other B7 family members that transduce signals via costimulatory receptors, in order to further augment the immune response.

Agents that upregulate an immune response can be used prophylactically in vaccines against various polypeptides (e.g., polypeptides derived from pathogens). Immunity against a pathogen (e.g., a virus) can be induced by vaccinating with a viral protein along with an agent that upregulates an immune response, in an appropriate adjuvant.

In another embodiment, upregulation or enhancement of an immune response function, as described herein, is useful in the induction of tumor immunity.

In another embodiment, the immune response can be stimulated by the methods described herein, such that pre-

existing tolerance, clonal deletion, and/or exhaustion (e.g., T cell exhaustion) is overcome. For example, immune responses against antigens to which a subject cannot mount a significant immune response, e.g., to an autologous antigen, such as a tumor specific antigens can be induced by administering appropriate agents described herein that upregulate the immune response. In one embodiment, an autologous antigen, such as a tumor-specific antigen, can be coadministered. In another embodiment, the subject agents can be used as adjuvants to boost responses to foreign antigens in the process of active immunization.

In one embodiment, immune cells are obtained from a subject and cultured ex vivo in the presence of an agent as described herein, to expand the population of immune cells and/or to enhance immune cell activation. In a further embodiment the immune cells are then administered to a subject. Immune cells can be stimulated in vitro by, for example, providing to the immune cells a primary activation signal and a costimulatory signal, as is known in the art. Various agents can also be used to costimulate proliferation of immune cells. In one embodiment immune cells are cultured ex vivo according to the method described in PCT Application No. WO 94/29436. The costimulatory polypeptide can be soluble, attached to a cell membrane, or attached to a solid surface, such as a bead.

In still another embodiment, agents described herein useful for upregulating immune responses can further be linked, or operatively attached, to toxins using techniques that are known in the art, e.g., crosslinking or via recombinant DNA techniques. Such agents can result in cellular destruction of desired cells. In one embodiment, a toxin can be conjugated to an antibody, such as a bispecific antibody. Such antibodies are useful for targeting a specific cell population, e.g., using a marker found only on a certain type of cell. The preparation of immunotoxins is, in general, well known in the art (see, e.g., U.S. Pat. No. 4,340,535, and EP 44167). Numerous types of disulfide-bond containing linkers are known which can successfully be employed to conjugate the toxin moiety with a polypeptide. In one embodiment, linkers that contain a disulfide bond that is sterically "hindered" are preferred, due to their greater stability in vivo, thus preventing release of the toxin moiety prior to binding at the site of action. A wide variety of toxins are known that may be conjugated to polypeptides or antibodies of the invention. Examples include: numerous useful plant-, fungus- or even bacteria-derived toxins, which, by way of example, include various A chain toxins, particularly ricin A chain, ribosome inactivating proteins such as saporin or gelonin, a-sarcin, aspergillin, restrictocin, ribonucleases, such as placental ribonuclease, angiogenic, diphtheria toxin, and *Pseudomonas* exotoxin, etc. A preferred toxin moiety for use in connection with the invention is toxin A chain which has been treated to modify or remove carbohydrate residues, deglycosylated A chain. (U.S. Pat. No. 5,776,427). Infusion of one or a combination of such cytotoxic agents, (e.g., ricin fusions) into a patient may result in the death of immune cells.

In yet another embodiment, the efficacy of the treatment methods described herein can be enhanced by incorporating a step of lymphodepletion prior to, concurrently with, or after the administration of agents the inhibit or block PD-1, PD-L1, CTLA-4, TIM-3, and/or LAG-3 function. For example, therapeutic benefits of administering the described anti-cancer agents can be synergistically enhanced by performing such administration after or in conjunction with lymphodepletion. Methods for achieving lymphodepletion in various forms and at various levels are well known in the

art (see, for example, U.S. Pat. No. 7,138,144). For example, the term “transient lymphodepletion” refers to destruction of lymphocytes and T cells, usually prior to immunotherapy. This can be accomplished in a number of ways, including “sublethal irradiation,” which refers to administration of one or more doses of radiation that is generally non-lethal to all members of a population of subjects to which the administration is applied. Transient lymphodepletion is generally not myeloablative, as would be the case in complete lymphodepletion, such that the subjects hematopoietic or immunological capacity remains sufficiently intact to regenerate the destroyed lymphocyte and T cell populations. By contrast, “lethal irradiation” occurs when the administration is generally lethal to some but not all members of the population of subjects and “supralethal irradiation” occurs when the administration is generally lethal to all members of the population of subjects.

Depending on the application and purpose, transient lymphodepletion or complete lymphodepletion may be effected, for example, by any combination of irradiation, treatment with a myeloablative agent, and/or treatment with an immunosuppressive agent, according to standard protocols. For example, biological methods include, for example, administration of immunity-suppressing cells or by administration of biological molecules capable of inhibiting immunoreactivity, such as, for example, Fas-ligand and CTLA4-Ig. Examples of myeloablative agents include busulfan, dimethyl mileran, melphalan and thiotepa. Examples of immunosuppressive agents include prednisone, methyl prednisolone, azathioprine, cyclosporine A, cyclophosphamide, fludarabin, CTLA4-Ig, anti-T cell antibodies, etc.

Regarding irradiation, a sublethal dose of irradiation is generally within the range of 1 to 7.5 Gy whole body irradiation, a lethal dose is generally within the range of 7.5 to 9.5 Gy whole body irradiation, and a supralethal dose is within the range of 9.5 to 16.5 Gy whole body irradiation.

Depending on the purpose and application, the dose of irradiation may be administered as a single dose or as a fractionated dose. Similarly, administering one or more doses of irradiation can be accomplished essentially exclusively to the body part or to a portion thereof, so as to induce myeloreduction or myeloablation essentially exclusively in the body part or the portion thereof. As is widely recognized in the art, a subject can tolerate as sublethal conditioning ultra-high levels of selective irradiation to a body part such as a limb, which levels constituting lethal or supralethal conditioning when used for whole body irradiation (see, for example, Breitz (2002) *Cancer Biother Radiopharm.* 17:119; Limit (1997) *J. Nucl. Med.* 38:1374; and Dritschilo and Sherman (1981) *Environ. Health Perspect.* 39:59). Such selective irradiation of the body part, or portion thereof, can be advantageously used to target particular blood compartments, such as specific lymph nodes, in treating hematopoietic cancers.

e. Administration of Agents

The immune modulating agents of the invention are administered to subjects in a biologically compatible form suitable for pharmaceutical administration in vivo, to enhance immune cell mediated immune responses. By “biologically compatible form suitable for administration in vivo” is meant a form to be administered in which any toxic effects are outweighed by the therapeutic effects. The term “subject” is intended to include living organisms in which an immune response can be elicited, e.g., mammals. Examples of subjects include humans, dogs, cats, mice, rats, and transgenic species thereof. Administration of an agent as described herein can be in any pharmacological form includ-

ing a therapeutically active amount of an agent alone or in combination with a pharmaceutically acceptable carrier.

Administration of a therapeutically active amount of the therapeutic composition of the present invention is defined as an amount effective, at dosages and for periods of time necessary, to achieve the desired result. For example, a therapeutically active amount of an agent may vary according to factors such as the disease state, age, sex, and weight of the individual, and the ability of peptide to elicit a desired response in the individual. Dosage regimens can be adjusted to provide the optimum therapeutic response. For example, several divided doses can be administered daily or the dose can be proportionally reduced as indicated by the exigencies of the therapeutic situation.

Inhibiting or blocking the function of PD-1 or PD-L1 and LAG-3, CTLA-4 or TIM-3, or in some embodiments, inhibiting or blocking a combination of these agents, can be accomplished by combination therapy with the modulatory agents described herein. Combination therapy describes a therapy in which PD-1 or PD-L1 and LAG-3, CTLA-4 or TIM-3, are inhibited or blocked simultaneously. Simultaneous inhibition or blockade may be achieved by administration of the modulatory agents described herein simultaneously (e.g., in a combination dosage form or by simultaneous administration of single agents) or by administration of single agents according to a schedule that results in effective amounts of each modulatory agent present in the patient at the same time.

The therapeutic agents described herein can be administered in a convenient manner such as by injection (subcutaneous, intravenous, etc.), oral administration, inhalation, transdermal application, or rectal administration. Depending on the route of administration, the active compound can be coated in a material to protect the compound from the action of enzymes, acids and other natural conditions which may inactivate the compound. For example, for administration of agents, by other than parenteral administration, it may be desirable to coat the agent with, or co-administer the agent with, a material to prevent its inactivation.

An agent can be administered to an individual in an appropriate carrier, diluent or adjuvant, co-administered with enzyme inhibitors or in an appropriate carrier such as liposomes. Pharmaceutically acceptable diluents include saline and aqueous buffer solutions. Adjuvant is used in its broadest sense and includes any immune stimulating compound such as interferon. Adjuvants contemplated herein include resorcinols, non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecyl polyethylene ether. Enzyme inhibitors include pancreatic trypsin inhibitor, diisopropylfluorophosphate (DEEP) and trasylol. Liposomes include water-in-oil-in-water emulsions as well as conventional liposomes (Sterna et al. (1984) *J. Neuroimmunol.* 7:27).

The agent may also be administered parenterally or intraperitoneally. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof, and in oils. Under ordinary conditions of storage and use, these preparations may contain a preservative to prevent the growth of microorganisms.

Pharmaceutical compositions of agents suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. In all cases the composition will preferably be sterile and must be fluid to the extent that easy syringeability exists. It will preferably be stable under the conditions of manufacture and storage and preserved against the contami-

nating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it is preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating an agent of the invention (e.g., an antibody, peptide, fusion protein or small molecule) in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the agent plus any additional desired ingredient from a previously sterile-filtered solution thereof.

When the agent is suitably protected, as described above, the protein can be orally administered, for example, with an inert diluent or an assimilable edible carrier. As used herein "pharmaceutically acceptable carrier" includes any and all solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like. The use of such media and agents for pharmaceutically active substances is well known in the art. Except insofar as any conventional media or agent is incompatible with the active compound, use thereof in the therapeutic compositions is contemplated. Supplementary active compounds can also be incorporated into the compositions.

It is especially advantageous to formulate parenteral compositions in dosage unit form for ease of administration and uniformity of dosage. "Dosage unit form", as used herein, refers to physically discrete units suited as unitary dosages for the mammalian subjects to be treated; each unit containing a predetermined quantity of active compound calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. The specification for the dosage unit forms of the invention are dictated by, and directly dependent on, (a) the unique characteristics of the active compound and the particular therapeutic effect to be achieved, and (b) the limitations inherent in the art of compounding such an active compound for the treatment of sensitivity in individuals.

In one embodiment, an agent of the invention is an antibody. As defined herein, a therapeutically effective amount of antibody (i.e., an effective dosage) ranges from about 0.001 to 30 mg/kg body weight, preferably about 0.01 to 25 mg/kg body weight, more preferably about 0.1 to 20 mg/kg body weight, and even more preferably about 1 to 10 mg/kg, 2 to 9 mg/kg, 3 to 8 mg/kg, 4 to 7 mg/kg, or 5 to 6 mg/kg body weight. The skilled artisan will appreciate that certain factors may influence the dosage required to effec-

tively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of an antibody can include a single treatment or, preferably, can include a series of treatments. In a preferred example, a subject is treated with antibody in the range of between about 0.1 to 20 mg/kg body weight, one time per week for between about 1 to 10 weeks, preferably between 2 to 8 weeks, more preferably between about 3 to 7 weeks, and even more preferably for about 4, 5, or 6 weeks. It will also be appreciated that the effective dosage of antibody used for treatment may increase or decrease over the course of a particular treatment. Changes in dosage may result from the results of diagnostic assays.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application, as well as the Figures, are incorporated herein by reference.

III. Kits

The present invention also encompasses kits for treating cancers, such as hematologic cancers like multiple myeloma, using agents that inhibit or block PD-1 or PD-L1 and LAG-3,

CTLA-4 or TIM-3 function. For example, the kit can comprise an antibody as described herein, e.g., an antibody against PD-1, PD-L1, LAG-3, CTLA-4 and/or TIM-3, packaged in a suitable container and can further comprise instructions for using such antibodies to treat cancers in a patient in need thereof. The kit may also contain other components, such as administration tools like packaged in a separate container.

EXAMPLES

Example 1: Combined PD-L1 and TIM-3 Blockade as Immunotherapy for Hematologic Cancers

a. Materials and Methods

Mice

All mice were housed in the Medical College of Wisconsin Biomedical Resource Center, an AAALAC-accredited facility. C47BL6/KaLwRij mice were used in the experiments. All animal work was reviewed and approved by the Medical College of Wisconsin Institutional Animal Care and Use Committee.

Tumor Cells

The 5T33 murine multiple myeloma (MM) cell line was derived from myeloma that spontaneously arose in a C57B1/KaLwRij mouse (Radl et al. (1988) *Am. J. Pathol.* 132:593-597; and Manning et al. (1992) *Br. J. Cancer* 66:1088-1093). For experiments, 5T33 cells were thawed from a large frozen stock and cultured in RPMI 1640+10% fetal bovine serum for no longer than 2 weeks prior to inoculation of mice. Mice were inoculated with tumor as follows: 2×10^6 5T33 cells intravenously (i.v.). 5T33-bearing mice were considered moribund and euthanized when they developed paraparesis or paraplegia. Occasionally, 5T33-inoculated mice developed tumor masses or lesions and were euthanized when the size of the mass or lesion exceeded 250 mm²; other symptoms of advanced 5T33 included splenomegaly, hepatomegaly, or neurologic impairment.

Antibodies

The clone designations for the antibodies used are as follows: anti-PD-L1 (clone 10F.9G2; Paterson et al. (2011) *J. Immunol.* 187:1097-1105; Maier et al. (2007) *J. Immunol.* 178:2714-2720), anti-PD-1 (clone 332.8H3), anti-Lag-3 (clone C9B7W; available from eBioscience as catalog number eBioC9B7W and other manufacturers), anti-Tim-3 (clone 5D12; available from EMD Millipore as catalog number MABF73). Isotype control antibodies included Armenian hamster IgG and rat IgG2a kappa.

Irradiation

In general, myeloma-bearing recipient mice were given total body irradiation as a single sublethal (500 cGy) dose seven days after myeloma inoculation. Radiation was administered by a Shepherd Mark I Cesium Irradiator in accordance with established guidelines.

Statistics

Survival curves were compared using the log rank (Mantel Cox) test based on n=5-6 mice per group. P-values of <0.05 were considered significant. Statistical analysis was done using Prism version 5.0a software (GraphPad Software, La Jolla, Calif.).

Other materials and methods are described in the "Results" section below.

b. Results

Multiple myeloma is characterized by the presence of transformed neoplastic plasma cells in the bone marrow and is generally considered to be an incurable disease. Successful treatments will likely require multi-faceted approaches incorporating conventional drug therapies, immunotherapy and other novel treatments. It has previously been determined that a combination of transient lymphodepletion (sublethal whole body irradiation) and PD-1 blockade generated anti-myeloma T cell reactivity capable of eliminating established disease (Kearl et al. (2013) *J. Immunol.* 190: 5620-5628; Hallett et al. (2011) *Biol. Blood Marrow Transplant.* 17:1133-1145). Besides expression of the immune checkpoint protein PD-1, T cells within tumor environment may develop a dysfunctional phenotype accompanied by the increased expression of other checkpoint proteins.

It was hypothesized that the anti-myeloma effect of transient lymphodepletion and PD-1 blockade would be increased by blocking other immune checkpoint protein interactions.

Expression of Immune Checkpoint Proteins on T Cells in Bone Marrow of Myeloma Bearing Mice Over Time

Accordingly, an extensive phenotypic analysis (flow cytometry) of bone marrow and splenic tissues from myeloma-bearing mice was performed to temporally examine T cells for expression of immune checkpoint proteins and assess the tissues for presence of immune regulatory T (Treg) cells. KaLwRij mice were inoculated with 2×10^5 T33-GFP cells intravenously. Myeloma bearing mice were euthanized between days 7 and 28 after inoculation or when moribund (day 29-40), and femoral bone marrow cells were harvested. Tumor cell accumulation was monitored by flow cytometry (GFP+ tumor cells) (FIG. 1A), and CD4⁺ (top) and CD8⁺ (bottom) T cells were analyzed by flow cytometry for expression of various immune checkpoint proteins over time (FIG. 1B). Naïve non-myeloma bearing mice were analyzed as controls. Immune checkpoint protein percentages were based on isotype controls.

As shown in FIG. 1B, PD-1, 2B4, LAG3, and TIM-3 were the most prominent immune checkpoint proteins present on T cells in myeloma bearing mice. As shown in FIG. 1C, a relatively large percentage of PD-1⁺ T cells co-expressed other inhibitory checkpoint proteins such as Tim-3, Lag-3

and 2B4. It was also determined that Treg cells in the tumor microenvironment also had increased expression of PD-1 and other inhibitory receptors (such as Tim-3, Lag-3 and 2B4 shown in FIG. 2) compared to Tregs cells from non-myeloma bearing mice, which is believed to be related with enhanced suppressive function by these cells.

Increased Expression of Immune Checkpoint Proteins on T Cells in Mice Treated with Sublethal Whole Body Irradiation and Anti-PD-L1 Antibody

Myeloma bearing KaLwRij mice were treated with 500 cGy whole body irradiation 7 days after tumor cell inoculation. Treatment with anti-PD-L1 antibody or control IgG (200µg i.p.) was initiated 5 days later and specifically given 12, 14, and 19 days after tumor inoculation. Mice were euthanized at day 21, splenocytes were harvested, and the CD8 T cells were analyzed by flow cytometry for immune checkpoint protein expression. As shown in FIGS. 3A-3B, the frequencies of CD8⁺Tim-3⁺, CD8⁺Lag-3⁺ and CD8⁺2B4⁺ cells in spleens of anti-PD-L1 antibody treated mice were higher compared with spleens of control antibody treated.

Blocking PD-L1 in Combination with Tim-3 after Lymphodepleting Whole Body Irradiation Synergistically Improved Survival

It was then examined whether blocking various immune checkpoint proteins could provide additive or synergistic anti-myeloma effects when combined with PD-L1 blockade (FIG. 4). In this Example, the combined blockade of PD-1 and TIM-3 was most effective and proved to be synergistic, as myeloma was surprisingly rejected in 100% of these mice (FIG. 5). Inhibition of certain other immune checkpoint proteins, either alone or in combinations, did not produce such robust therapeutic benefits (FIGS. 6-8).

Thus, the data indicate that dual blockade of PD-L1 and TIM-3 represents a surprisingly and unexpectedly potent immunotherapeutic intervention for treating hematologic cancers, such as multiple myeloma.

Example 2: Combined Immune Checkpoint Protein Blockade and Lymphodepletion as Immunotherapy for Hematologic Cancers

a. Materials and methods are essentially the same as described in Example 1 unless specifically indicated below in the "Results" section below.

b. Results

As shown in this Example, combined immune checkpoint protein blockade and lymphodepletion provide an effective immunotherapy for hematologic cancers such as myeloma. Blocking of PD-L1 in Combination with Tim-3, Lag-3 or CTLA-4 after Lymphodepleting Whole Body Irradiation Synergistically Improved Survival

FIG. 9A depicts the experimental design. KaLwRij mice received 500 cGy irradiation 7 days after tumor cell inoculation. The treatment with blocking antibody or control IgG (200 µg i.p.) was initiated 5 days later and specifically given 12, 14, 19, 21, 26, and 28 days after tumor inoculation.

As shown in FIGS. 9B-9D, blocking PD-L1 in combination with Tim-3 (FIG. 9B), Lag-3 (FIG. 9B), or CTLA-4 (FIG. 9C) after lymphodepleting whole body irradiation synergistically improved survival of myeloma bearing mice, whereas blocking PD-L1 in combination with CD48 (FIG. 9D) did not have synergistic effect on survival. As shown in FIG. 9E, 100% of re-challenged mice that had received anti-PD-L1 antibody alone, or the combination of anti-PD-

L1 antibody with anti-Tim-3 antibody, anti-Lag-3 antibody, or anti-CTLA-4 antibody, survived to day 110, compared to control mice.

Combined Checkpoint Blockade after Lymphodepleting Whole Body Irradiation Increased Frequencies of Tumor-Reactive T Cells

The experimental design shown in FIG. 9A was used. CD4⁺ or CD8⁺ T cells were isolated from spleens and bone marrow 21 days after tumor cell inoculation (i.e., 14 days after irradiation) in mice treated with control IgG, anti-PD-L1 antibody only, or the combination of anti-PD-L1 antibody with anti-Tim-3 antibody, anti-Lag-3 antibody, or anti-CTLA-4 antibody. The CD8⁺ or CD4⁺ T cells were tested in IFN- γ ELISPOT assays using 5T33 or MHC class II+ 5T33 tumor cells as stimulators, respectively, to determine tumor-reactive IFN- γ -secreting cell frequencies.

As shown in FIG. 10A, the frequencies of tumor-reactive CD8⁺ and CD4⁺ T cells were increased in the spleens (top row) and bone marrow (bottom row) of mice treated with combinations of immune checkpoint protein blockade. Combined checkpoint blockade, such as the combination of anti-PD-L1 antibody with anti-Tim-3 antibody, anti-Lag-3 antibody, or anti-CTLA-4 antibody, increased frequencies of tumor-reactive T cells.

Combined Checkpoint Blockade after Lymphodepleting Whole Body Irradiation Increased Cytokine Production by CD8⁺ T Cells

CD8⁺ T cells purified from the spleens of myeloma bearing mice treated with anti-PD-L1 antibody only, or the combination of anti-PD-L1 antibody with anti-Tim-3 antibody, anti-Lag-3 antibody or anti-CTLA4 antibody, were stimulated with 5T33 for 48 hours. Supernatants were collected and cytokine levels from were determined using a multiplex cytokine assay. As shown in FIG. 10B, combined checkpoint blockade, such as the combination of anti-PD-L1 antibody with anti-Tim-3 antibody, anti-Lag-3 antibody, or anti-CTLA-4 antibody, increased the production of cytokines (e.g., IL-2, IFN- γ and GM-CSF) by CD8⁺ T cells.

Combined Blockade of Immune Checkpoint Proteins Increased PD-1 Expression on CD8⁺ T Cells as Well as Increased Frequency of Tumor Specific Cytotoxic T Lymphocytes

The experimental design in FIG. 9A was used. CD8⁺ T cells were isolated from spleens and bone marrow 21 days after tumor cell inoculation in mice treated with control IgG, anti-PD-L1 antibody only, or the combination of anti-PD-L1 antibody with anti-Tim-3 antibody, anti-Lag-3 antibody, or anti-CTLA4 antibody.

FIG. 11A shows increased expression of PD-1 on gated CD8⁺ T cells from spleens and bone marrow (BM) of mice treated with different blocking antibodies or control IgG.

The CD8⁺ T cells were assayed in IFN- γ ELISPOT assays with tumor cell stimulators to determine tumor-reactive IFN- γ -secreting cell frequencies in the presence of anti-PD-L1 antibody or control IgG (10 μ g/ml). As shown in FIG. 11B, in the presence of blocking anti-PD-L1 antibody in vitro, all combinations of blocking antibodies in vivo resulted in significantly increased cytotoxic T lymphocyte (CTL) frequencies versus blockade of PD-L1 only (p<0.001).

Combined Blockade of Immune Checkpoint Proteins Enhanced Th1 and Th2 Cytokine Secretion

The experimental design shown in FIG. 9A was used. CD4⁺ T cells were isolated from spleens 21 days after tumor cell inoculation (i.e., 14 days after irradiation) in myeloma bearing mice treated with control anti-PD-L1, or the combination of anti-PD-L1 antibody with anti-Tim-3 antibody,

anti-Lag-3 antibody, or anti-CTLA4 antibody. CD4⁺ T cells purified from the spleen were stimulated with MHC class II negative 5T33-WT, or MHC class II positive 5T33-CIITA, or no stimulation for 48 hours. Supernatants were collected and cytokine (IFN- γ , IL-4 and IL-5) levels from were determined using a multiplex cytokine assay.

As shown in FIG. 12, combined blockade of immune checkpoint proteins such as anti-PD-L1 and anti-CTLA-4 antibodies enhanced Th1 and Th2 cytokine secretion.

Expression of Immune Checkpoints on T Cells in Mice with Other Hematologic Cancers

Murine hematologic cancers other than myeloma express PD-L1 and respond to whole body irradiation plus PD-L1 blockade (Kearl et al. (2013) *J. Immunol.* 190:5620-5628).

To examine the expression of immune checkpoint proteins on T cells in mice with other hematologic cancers, mice were injected i.v. with A20 B cell lymphoma cells, C1498 acute myeloid leukemia cells, or EL4 lymphoma cells (x-axis in FIG. 13). Bone marrow was collected from moribund animals. CD4⁺ (top row in FIG. 13) and CD8⁺ (bottom row in FIG. 13) T cells in the bone marrow were analyzed for expression of checkpoint proteins PD-1, Tim-3, Lag-3 and 2B4 by flow cytometry (n=4-5 mice for each). Naïve non-cancer-bearing mice were used as controls. FIG. 13 shows expression of PD-1, Tim-3, Lag-3 and 2B4 on T cells in mice bearing the indicated hematologic cancer cells. A correlation can exist between the T cell expression profile of these checkpoint proteins and increased anti-tumor response after co-blockade of the respective pathway(s) in hematologic malignancy models other than myeloma.

A Model of Combined Immune Checkpoint Blockade and Lymphodepleting Whole Body Irradiation for Treating Hematologic Cancers

A working model of combined immune checkpoint blockade and lymphodepleting whole body irradiation is illustrated in FIG. 14. In hematologic cancers, dysfunctional antigen-activated T cells (e.g., CD4⁺ and CD8⁺ T cells) are unable to kill cancer cells. Lymphopenia-induced T cell proliferation allows for reactivation of those T cells. For reactivated T cells to remain functional and kill cancer cells, immune checkpoint proteins must be blocked. Lymphopenic environment can be achieved by low-dose whole body irradiation (WBI). Lymphodepleting chemotherapy or low doses of T cell-depleting antibodies can also be used instead of whole body irradiation.

INCORPORATION BY REFERENCE

All publications, patents, and patent applications mentioned herein are hereby incorporated by reference in their entirety as if each individual publication, patent or patent application was specifically and individually indicated to be incorporated by reference. In case of conflict, the present application, including any definitions herein, will control.

Also incorporated by reference in their entirety are any polynucleotide and polypeptide sequences which reference an accession number correlating to an entry in a public database, such as those maintained by The Institute for Genomic Research (TIGR) on the world wide web at tigr.org and/or the National Center for Biotechnology Information (NCBI) on the World Wide Web at ncbi.nlm.nih.gov.

EQUIVALENTS

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1

<211> LENGTH: 921

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (25)..(888)

<400> SEQUENCE: 1

```

cactctgggtg gggctgctcc aggc atg cag atc cca cag gcg ccc tgg cca      51
      Met Gln Ile Pro Gln Ala Pro Trp Pro
      1                5

gtc gtc tgg gcg gtg cta caa ctg ggc tgg cgg cca gga tgg ttc tta      99
Val Val Trp Ala Val Leu Gln Leu Gly Trp Arg Pro Gly Trp Phe Leu
10                15                20                25

gac tcc cca gac agg ccc tgg aac ccc ccc acc ttc tcc cca gcc ctg      147
Asp Ser Pro Asp Arg Pro Trp Asn Pro Pro Thr Phe Ser Pro Ala Leu
      30                35                40

ctc gtg gtg acc gaa ggg gac aac gcc acc ttc acc tgc agc ttc tcc      195
Leu Val Val Thr Glu Gly Asp Asn Ala Thr Phe Thr Cys Ser Phe Ser
      45                50                55

aac aca tcg gag agc ttc gtg cta aac tgg tac cgc atg agc ccc agc      243
Asn Thr Ser Glu Ser Phe Val Leu Asn Trp Tyr Arg Met Ser Pro Ser
      60                65                70

aac cag acg gac aag ctg gcc gcc ttc ccc gag gac cgc agc cag ccc      291
Asn Gln Thr Asp Lys Leu Ala Ala Phe Pro Glu Asp Arg Ser Gln Pro
      75                80                85

ggc cag gac tgc cgc ttc cgt gtc aca caa ctg ccc aac ggg cgt gac      339
Gly Gln Asp Cys Arg Phe Arg Val Thr Gln Leu Pro Asn Gly Arg Asp
90                95                100                105

ttc cac atg agc gtg gtc agg gcc cgg cgc aat gac agc ggc acc tac      387
Phe His Met Ser Val Val Arg Ala Arg Asn Asp Ser Gly Thr Tyr
      110                115                120

ctc tgt ggg gcc atc tcc ctg gcc ccc aag gcg cag atc aaa gag agc      435
Leu Cys Gly Ala Ile Ser Leu Ala Pro Lys Ala Gln Ile Lys Glu Ser
      125                130                135

ctg cgg gca gag ctc agg gtg aca gag aga agg gca gaa gtg ccc aca      483
Leu Arg Ala Glu Leu Arg Val Thr Glu Arg Arg Ala Glu Val Pro Thr
      140                145                150

gcc cac ccc agc ccc tca ccc agg tca gcc ggc cag ttc caa acc ctg      531
Ala His Pro Ser Pro Ser Pro Arg Ser Ala Gly Gln Phe Gln Thr Leu
      155                160                165

gtg gtt ggt gtc gtg ggc ggc ctg ctg ggc agc ctg gtg ctg cta gtc      579
Val Val Gly Val Val Gly Gly Leu Leu Gly Ser Leu Val Leu Leu Val
170                175                180                185

tgg gtc ctg gcc gtc atc tgc tcc cgg gcc gca cga ggg aca ata gga      627
Trp Val Leu Ala Val Ile Cys Ser Arg Ala Ala Arg Gly Thr Ile Gly
      190                195                200

gcc agg cgc acc ggc cag ccc ctg aag gag gac ccc tca gcc gtg cct      675
Ala Arg Arg Thr Gly Gln Pro Leu Lys Glu Asp Pro Ser Ala Val Pro
      205                210                215

gtg ttc tct gtg gac tat ggg gag ctg gat ttc cag tgg cga gag aag      723
Val Phe Ser Val Asp Tyr Gly Glu Leu Asp Phe Gln Trp Arg Glu Lys
      220                225                230

acc ccg gag ccc ccc gtg ccc tgt gtc cct gag cag acg gag tat gcc      771
Thr Pro Glu Pro Pro Val Pro Cys Val Pro Glu Gln Thr Glu Tyr Ala
      235                240                245

acc att gtc ttt cct agc gga atg ggc acc tca tcc ccc gcc cgc agg      819
Thr Ile Val Phe Pro Ser Gly Met Gly Thr Ser Ser Pro Ala Arg Arg

```


-continued

250	255	260	265	
ggc tca gct gac	ggc cct cgg agt gcc cag cca ctg agg cct gag gat			867
Gly Ser Ala Asp	Gly Pro Arg Ser Ala Gln Pro Leu Arg Pro Glu Asp			
	270	275	280	
gga cac tgc tct tgg ccc ctc	tgaccggctt ccttggccac cagtgttctg cag			921
Gly His Cys Ser Trp Pro Leu				
	285			

<210> SEQ ID NO 2
 <211> LENGTH: 288
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Gln Ile Pro Gln Ala Pro Trp Pro Val Val Trp Ala Val Leu Gln	
1	5 10 15
Leu Gly Trp Arg Pro Gly Trp Phe Leu Asp Ser Pro Asp Arg Pro Trp	
	20 25 30
Asn Pro Pro Thr Phe Ser Pro Ala Leu Leu Val Val Thr Glu Gly Asp	
	35 40 45
Asn Ala Thr Phe Thr Cys Ser Phe Ser Asn Thr Ser Glu Ser Phe Val	
	50 55 60
Leu Asn Trp Tyr Arg Met Ser Pro Ser Asn Gln Thr Asp Lys Leu Ala	
65	70 75 80
Ala Phe Pro Glu Asp Arg Ser Gln Pro Gly Gln Asp Cys Arg Phe Arg	
	85 90 95
Val Thr Gln Leu Pro Asn Gly Arg Asp Phe His Met Ser Val Val Arg	
	100 105 110
Ala Arg Arg Asn Asp Ser Gly Thr Tyr Leu Cys Gly Ala Ile Ser Leu	
	115 120 125
Ala Pro Lys Ala Gln Ile Lys Glu Ser Leu Arg Ala Glu Leu Arg Val	
	130 135 140
Thr Glu Arg Arg Ala Glu Val Pro Thr Ala His Pro Ser Pro Ser Pro	
145	150 155 160
Arg Ser Ala Gly Gln Phe Gln Thr Leu Val Val Gly Val Val Gly Gly	
	165 170 175
Leu Leu Gly Ser Leu Val Leu Leu Val Trp Val Leu Ala Val Ile Cys	
	180 185 190
Ser Arg Ala Ala Arg Gly Thr Ile Gly Ala Arg Arg Thr Gly Gln Pro	
	195 200 205
Leu Lys Glu Asp Pro Ser Ala Val Pro Val Phe Ser Val Asp Tyr Gly	
	210 215 220
Glu Leu Asp Phe Gln Trp Arg Glu Lys Thr Pro Glu Pro Pro Val Pro	
225	230 235 240
Cys Val Pro Glu Gln Thr Glu Tyr Ala Thr Ile Val Phe Pro Ser Gly	
	245 250 255
Met Gly Thr Ser Ser Pro Ala Arg Arg Gly Ser Ala Asp Gly Pro Arg	
	260 265 270
Ser Ala Gln Pro Leu Arg Pro Glu Asp Gly His Cys Ser Trp Pro Leu	
	275 280 285

<210> SEQ ID NO 3
 <211> LENGTH: 968
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS

-continued

<222> LOCATION: (59)..(793)

<400> SEQUENCE: 3

```

gcttcccagag gctccgcacc agccgcgctt ctgtccgctt gcagggcatt ccagaaag      58

atg agg ata ttt gct gtc ttt ata ttc atg acc tac tgg cat ttg ctg      106
Met Arg Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu
1           5           10           15

aac gca ttt act gtc acg gtt ccc aag gac cta tat gtg gta gag tat      154
Asn Ala Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr
           20           25           30

ggg agc aat atg aca att gaa tgc aaa ttc cca gta gaa aaa caa tta      202
Gly Ser Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu
           35           40           45

gac ctg gct gca cta att gtc tat tgg gaa atg gag gat aag aac att      250
Asp Leu Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile
50           55           60

att caa ttt gtg cat gga gag gaa gac ctg aag gtt cag cat agt agc      298
Ile Gln Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser
65           70           75           80

tac aga cag agg gcc cgg ctg ttg aag gac cag ctc tcc ctg gga aat      346
Tyr Arg Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn
           85           90           95

gct gca ctt cag atc aca gat gtg aaa ttg cag gat gca ggg gtg tac      394
Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr
100           105           110

cgc tgc atg atc agc tat ggt ggt gcc gac tac aag cga att act gtg      442
Arg Cys Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val
115           120           125

aaa gtc aat gcc cca tac aac aaa atc aac caa aga att ttg gtt gtg      490
Lys Val Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val
130           135           140

gat cca gtc acc tct gaa cat gaa ctg aca tgt cag gct gag ggc tac      538
Asp Pro Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr
145           150           155           160

ccc aag gcc gaa gtc atc tgg aca agc agt gac cat caa gtc ctg agt      586
Pro Lys Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser
165           170           175

ggg aag acc acc acc acc aat tcc aag aga gag gag aag ctt ttc aat      634
Gly Lys Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn
180           185           190

gtg acc agc aca ctg aga atc aac aca aca act aat gag att ttc tac      682
Val Thr Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr
195           200           205

tgc act ttt agg aga tta gat cct gag gaa aac cat aca gct gaa ttg      730
Cys Thr Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu
210           215           220

gtc atc cca ggt aat att ctg aat gtg tcc att aaa ata tgt cta aca      778
Val Ile Pro Gly Asn Ile Leu Asn Val Ser Ile Lys Ile Cys Leu Thr
225           230           235           240

ctg tcc cct agc acc tagcatgatg tctgcctatc atagtcattc agtgattggt      833
Leu Ser Pro Ser Thr
245

gaataaatga atgaatgaat aacactatgt ttacaaaata taccctaatt cctcacctcc      893

attcatocaa accatattgt tacttaataa acattcagca gatatttatg gaataaaaaa      953

aaaaaaaaaa aaaaaa      968

```

<210> SEQ ID NO 4

<211> LENGTH: 245

-continued

```

<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Arg Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu
1           5           10           15

Asn Ala Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr
20           25           30

Gly Ser Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu
35           40           45

Asp Leu Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile
50           55           60

Ile Gln Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser
65           70           75           80

Tyr Arg Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn
85           90           95

Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr
100          105          110

Arg Cys Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val
115          120          125

Lys Val Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val
130          135          140

Asp Pro Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr
145          150          155          160

Pro Lys Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser
165          170          175

Gly Lys Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn
180          185          190

Val Thr Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr
195          200          205

Cys Thr Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu
210          215          220

Val Ile Pro Gly Asn Ile Leu Asn Val Ser Ile Lys Ile Cys Leu Thr
225          230          235          240

Leu Ser Pro Ser Thr
245

```

```

<210> SEQ ID NO 5
<211> LENGTH: 1553
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (53)..(922)

```

```

<400> SEQUENCE: 5

cgaggctccg caccagccgc gcttctgtcc gctgcaggg cattccagaa ag atg agg      58
                                         Met Arg
                                         1

ata ttt gct gtc ttt ata ttc atg acc tac tgg cat ttg ctg aac gca      106
Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu Asn Ala
5           10           15

ttt act gtc acg gtt ccc aag gac cta tat gtg gta gag tat ggt agc      154
Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr Gly Ser
20           25           30

aat atg aca att gaa tgc aaa ttc cca gta gaa aaa caa tta gac ctg      202
Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu Asp Leu
35           40           45           50

```


-continued

gct gca cta att gtc tat tgg gaa atg gag gat aag aac att att caa	250
Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile Ile Gln	
55 60 65	
ttt gtg cat gga gag gaa gac ctg aag gtt cag cat agt agc tac aga	298
Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser Tyr Arg	
70 75 80	
cag agg gcc cgg ctg ttg aag gac cag ctc tcc ctg gga aat gct gca	346
Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn Ala Ala	
85 90 95	
ctt cag atc aca gat gtg aaa ttg cag gat gca ggg gtg tac cgc tgc	394
Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr Arg Cys	
100 105 110	
atg atc agc tat ggt ggt gcc gac tac aag cga att act gtg aaa gtc	442
Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val Lys Val	
115 120 125 130	
aat gcc cca tac aac aaa atc aac caa aga att ttg gtt gtg gat cca	490
Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val Asp Pro	
135 140 145	
gtc acc tct gaa cat gaa ctg aca tgt cag gct gag ggc tac ccc aag	538
Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr Pro Lys	
150 155 160	
gcc gaa gtc atc tgg aca agc agt gac cat caa gtc ctg agt ggt aag	586
Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser Gly Lys	
165 170 175	
acc acc acc acc aat tcc aag aga gag gag aag ctt ttc aat gtg acc	634
Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn Val Thr	
180 185 190	
agc aca ctg aga atc aac aca aca act aat gag att ttc tac tgc act	682
Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr Cys Thr	
195 200 205 210	
ttt agg aga tta gat cct gag gaa aac cat aca gct gaa ttg gtc atc	730
Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu Val Ile	
215 220 225	
cca gaa cta cct ctg gca cat cct cca aat gaa agg act cac ttg gta	778
Pro Glu Leu Pro Leu Ala His Pro Pro Asn Glu Arg Thr His Leu Val	
230 235 240	
att ctg gga gcc atc tta tta tgc ctt ggt gta gca ctg aca ttc atc	826
Ile Leu Gly Ala Ile Leu Leu Cys Leu Gly Val Ala Leu Thr Phe Ile	
245 250 255	
ttc cgt tta aga aaa ggg aga atg atg gat gtg aaa aaa tgt ggc atc	874
Phe Arg Leu Arg Lys Gly Arg Met Met Asp Val Lys Lys Cys Gly Ile	
260 265 270	
caa gat aca aac tca aag aag caa agt gat aca cat ttg gag gag acg	922
Gln Asp Thr Asn Ser Lys Lys Gln Ser Asp Thr His Leu Glu Glu Thr	
275 280 285 290	
taatccagca ttggaacttc tgatcttcaa gcagggatcc tcaacctgtg gtttaggggt	982
tcatcggggc tgagcgtgac aagaggaagg aatgggcccg tgggatgcag gcaatgtggg	1042
acttaaaagg cccaagcact gaaaatggaa cctggcgaaa gcagaggagg agaatgaaga	1102
aagatggagt caaacaggga gcctggaggg agaccttgat actttcaaat gcctgagggg	1162
ctcatcgacg cctgtgacag ggagaaagga tactttctgaa caaggagcct ccaagcaaat	1222
catccattgc tcatcctagg aagacggggt gagaatccct aatttgaggg tcagttcctg	1282
cagaagtgcc ctttgctcc actcaatgcc tcaatttggt ttctgcatga ctgagagtct	1342
cagtgttga acgggacagt atttatgtat gagtttttcc tattttattt gagtctgtga	1402
ggtcttcttg tcatgtgagt gtggttgtga atgatttctt ttgaagatat attgtagtag	1462

-continued

 atgttacaat tttgtcgcca aactaaactt gctgcttaat gatttgctca catctagtaa 1522

aacatggagt atttgtaaaa aaaaaaaaaa a 1553

<210> SEQ ID NO 6

<211> LENGTH: 290

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

 Met Arg Ile Phe Ala Val Phe Ile Phe Met Thr Tyr Trp His Leu Leu
 1 5 10 15

 Asn Ala Phe Thr Val Thr Val Pro Lys Asp Leu Tyr Val Val Glu Tyr
 20 25 30

 Gly Ser Asn Met Thr Ile Glu Cys Lys Phe Pro Val Glu Lys Gln Leu
 35 40 45

 Asp Leu Ala Ala Leu Ile Val Tyr Trp Glu Met Glu Asp Lys Asn Ile
 50 55 60

 Ile Gln Phe Val His Gly Glu Glu Asp Leu Lys Val Gln His Ser Ser
 65 70 75 80

 Tyr Arg Gln Arg Ala Arg Leu Leu Lys Asp Gln Leu Ser Leu Gly Asn
 85 90 95

 Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr
 100 105 110

 Arg Cys Met Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Val
 115 120 125

 Lys Val Asn Ala Pro Tyr Asn Lys Ile Asn Gln Arg Ile Leu Val Val
 130 135 140

 Asp Pro Val Thr Ser Glu His Glu Leu Thr Cys Gln Ala Glu Gly Tyr
 145 150 155 160

 Pro Lys Ala Glu Val Ile Trp Thr Ser Ser Asp His Gln Val Leu Ser
 165 170 175

 Gly Lys Thr Thr Thr Thr Asn Ser Lys Arg Glu Glu Lys Leu Phe Asn
 180 185 190

 Val Thr Ser Thr Leu Arg Ile Asn Thr Thr Thr Asn Glu Ile Phe Tyr
 195 200 205

 Cys Thr Phe Arg Arg Leu Asp Pro Glu Glu Asn His Thr Ala Glu Leu
 210 215 220

 Val Ile Pro Glu Leu Pro Leu Ala His Pro Pro Asn Glu Arg Thr His
 225 230 235 240

 Leu Val Ile Leu Gly Ala Ile Leu Leu Cys Leu Gly Val Ala Leu Thr
 245 250 255

 Phe Ile Phe Arg Leu Arg Lys Gly Arg Met Met Asp Val Lys Lys Cys
 260 265 270

 Gly Ile Gln Asp Thr Asn Ser Lys Lys Gln Ser Asp Thr His Leu Glu
 275 280 285

 Glu Thr
 290

<210> SEQ ID NO 7

<211> LENGTH: 873

<212> TYPE: DNA

<213> ORGANISM: Mus sp.

<400> SEQUENCE: 7

atgaggatat ttgctggcat tatattcaca gcctgctgctc acttgctacg ggcgtttact 60

-continued

```

atcacggctc caaaggactt gtacgtggtg gagtatggca gcaacgtcac gatggagtgc 120
agattccctg tagaacggga gctggacctg cttgcgtagg tgggtgtactg ggaaaaggaa 180
gatgagcaag tgattcagtt tgtggcagga gaggaggacc ttaagcctca gcacagcaac 240
ttcaggggga gagcctcgct gccaaaggac cagcttttga agggaaatgc tgcccttcag 300
atcacagacg tcaagctgca ggacgcaggc gtttactgct gcataatcag ctacggtggt 360
gcgactaca agcgaatcac gctgaaagtc aatgccccat accgcaaat caaccagaga 420
atttccgtgg atccagccac ttctgagcat gaactaatat gtcaggccga gggttatcca 480
gaagctgagg taatctggac aaacagtgc caccaaccgg tgagtgggaa gagaagtgtc 540
accacttccc ggacagaggg gatgcttctc aatgtgacca gcagtctgag ggtcaacgcc 600
acagcgaatg atgttttcta ctgtacgttt tggagatcac agccagggca aaaccacaca 660
gcgagctga tcatcccaga actgcctgca acacatcctc cacagaacag gactcactgg 720
gtgcttctgg gatccatcct gttgttctc attgtagtgt ccacggctct cctcttcttg 780
agaaaacaag tgagaatgct agatgtggag aaatgtggcg ttgaagatac aagctcaaaa 840
aaccgaaatg atacacaatt cgaggagacg taa 873

```

```

<210> SEQ ID NO 8
<211> LENGTH: 290
<212> TYPE: PRT
<213> ORGANISM: Mus sp.

```

```

<400> SEQUENCE: 8

```

```

Met Arg Ile Phe Ala Gly Ile Ile Phe Thr Ala Cys Cys His Leu Leu
1           5           10           15
Arg Ala Phe Thr Ile Thr Ala Pro Lys Asp Leu Tyr Val Val Glu Tyr
20           25           30
Gly Ser Asn Val Thr Met Glu Cys Arg Phe Pro Val Glu Arg Glu Leu
35           40           45
Asp Leu Leu Ala Leu Val Val Tyr Trp Glu Lys Glu Asp Glu Gln Val
50           55           60
Ile Gln Phe Val Ala Gly Glu Glu Asp Leu Lys Pro Gln His Ser Asn
65           70           75           80
Phe Arg Gly Arg Ala Ser Leu Pro Lys Asp Gln Leu Leu Lys Gly Asn
85           90           95
Ala Ala Leu Gln Ile Thr Asp Val Lys Leu Gln Asp Ala Gly Val Tyr
100          105          110
Cys Cys Ile Ile Ser Tyr Gly Gly Ala Asp Tyr Lys Arg Ile Thr Leu
115          120          125
Lys Val Asn Ala Pro Tyr Arg Lys Ile Asn Gln Arg Ile Ser Val Asp
130          135          140
Pro Ala Thr Ser Glu His Glu Leu Ile Cys Gln Ala Glu Gly Tyr Pro
145          150          155          160
Glu Ala Glu Val Ile Trp Thr Asn Ser Asp His Gln Pro Val Ser Gly
165          170          175
Lys Arg Ser Val Thr Thr Ser Arg Thr Glu Gly Met Leu Leu Asn Val
180          185          190
Thr Ser Ser Leu Arg Val Asn Ala Thr Ala Asn Asp Val Phe Tyr Cys
195          200          205
Thr Phe Trp Arg Ser Gln Pro Gly Gln Asn His Thr Ala Glu Leu Ile
210          215          220
Ile Pro Glu Leu Pro Ala Thr His Pro Pro Gln Asn Arg Thr His Trp

```


-continued

225	230	235	240
Val Leu Leu Gly Ser Ile Leu Leu Phe Leu Ile Val Val Ser Thr Val			
	245	250	255
Leu Leu Phe Leu Arg Lys Gln Val Arg Met Leu Asp Val Glu Lys Cys			
	260	265	270
Gly Val Glu Asp Thr Ser Ser Lys Asn Arg Asn Asp Thr Gln Phe Glu			
	275	280	285
Glu Thr			
290			

<210> SEQ ID NO 9
 <211> LENGTH: 906
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

```

atgttttcac atcttcctt tgactgtgtc ctgctgctgc tgctgctact acttacaagg      60
tcctcagaag tggaatacag agcggaggtc ggtcagaatg cctatctgcc ctgcttctac     120
accccagccg ccccagggaa cctcgtgccc gtctgctggg gcaaaggagc ctgtcctgtg     180
tttgaatgtg gcaacgtggt gctcaggact gatgaaaggg atgtgaatta ttggacatcc     240
agatactggc taaatgggga tttccgcaaa ggagatgtgt ccctgacat agagaatgtg     300
actctagcag acagtgggat ctactgctgc cggatccaaa tcccaggcat aatgaatgat     360
gaaaaattta acctgaagtt ggatcatcaa ccagccaagg tcaccctgc accgactcgg     420
cagagagact tcaactgcagc ctttccaagg atgcttacca ccaggggaca tggcccagca     480
gagacacaga cactggggag cctccctgat ataaatctaa cacaaatc caccattggcc     540
aatgagttac gggactctag attggccaat gacttacggg actctggagc aaccatcaga     600
ataggcatct acatcggagc agggatctgt gctgggctgg ctctggctct tatcttcggc     660
gctttaattt tcaaattgta ttctcatagc aaagagaaga tacagaattt aagcctcatc     720
tctttggcca acctccctcc ctcaggattg gcaaatgcag tagcagaggg aattcgctca     780
gaagaaaaca tctataccat tgaagagaac gtatatgaag tggaggagcc caatgagtat     840
tattgctatg tcagcagcag gcagcaacct tcacaacctt tgggttgctg ctttgcaatg     900
ccatag                                             906
  
```

<210> SEQ ID NO 10
 <211> LENGTH: 301
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

Met Phe Ser His Leu Pro Phe Asp Cys Val Leu Leu Leu Leu Leu Leu															
1			5				10						15		
Leu Leu Thr Arg Ser Ser Glu Val Glu Tyr Arg Ala Glu Val Gly Gln															
			20				25						30		
Asn Ala Tyr Leu Pro Cys Phe Tyr Thr Pro Ala Ala Pro Gly Asn Leu															
			35				40						45		
Val Pro Val Cys Trp Gly Lys Gly Ala Cys Pro Val Phe Glu Cys Gly															
			50				55						60		
Asn Val Val Leu Arg Thr Asp Glu Arg Asp Val Asn Tyr Trp Thr Ser															
65							70						75		80
Arg Tyr Trp Leu Asn Gly Asp Phe Arg Lys Gly Asp Val Ser Leu Thr															
			85										90		95

-continued

Ile Glu Asn Val Thr Leu Ala Asp Ser Gly Ile Tyr Cys Cys Arg Ile
 100 105 110

Gln Ile Pro Gly Ile Met Asn Asp Glu Lys Phe Asn Leu Lys Leu Val
 115 120 125

Ile Lys Pro Ala Lys Val Thr Pro Ala Pro Thr Arg Gln Arg Asp Phe
 130 135 140

Thr Ala Ala Phe Pro Arg Met Leu Thr Thr Arg Gly His Gly Pro Ala
 145 150 155 160

Glu Thr Gln Thr Leu Gly Ser Leu Pro Asp Ile Asn Leu Thr Gln Ile
 165 170 175

Ser Thr Leu Ala Asn Glu Leu Arg Asp Ser Arg Leu Ala Asn Asp Leu
 180 185 190

Arg Asp Ser Gly Ala Thr Ile Arg Ile Gly Ile Tyr Ile Gly Ala Gly
 195 200 205

Ile Cys Ala Gly Leu Ala Leu Ala Leu Ile Phe Gly Ala Leu Ile Phe
 210 215 220

Lys Trp Tyr Ser His Ser Lys Glu Lys Ile Gln Asn Leu Ser Leu Ile
 225 230 235 240

Ser Leu Ala Asn Leu Pro Pro Ser Gly Leu Ala Asn Ala Val Ala Glu
 245 250 255

Gly Ile Arg Ser Glu Glu Asn Ile Tyr Thr Ile Glu Glu Asn Val Tyr
 260 265 270

Glu Val Glu Glu Pro Asn Glu Tyr Tyr Cys Tyr Val Ser Ser Arg Gln
 275 280 285

Gln Pro Ser Gln Pro Leu Gly Cys Arg Phe Ala Met Pro
 290 295 300

<210> SEQ ID NO 11
 <211> LENGTH: 846
 <212> TYPE: DNA
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 11

atgttttcag gtcttaccct caactgtgtc ctgctgctgc tgcaactact acttgcaagg 60
 tcattggaaa atgcttatgt gtttgagggt ggtaagaatg cctatctgcc ctgcagttac 120
 actctatcta cacctggggc acttgtgcct atgtgctggg gcaagggatt ctgtccttgg 180
 tcacagtgta ccaacgagtt gctcagaact gatgaaagaa atgtgacata tcagaaatcc 240
 agcagatacc agctaaaggg cgatctcaac aaaggagacg tgtctctgat cataaagaat 300
 gtgactctgg atgacctgg gacctactgc tgcaggatac agttccctgg tcttatgaat 360
 gataaaaaat tagaactgaa attagacatc aaagcagcca aggtcactcc agctcagact 420
 gcccatgggg actctactac agcttctcca agaaccctaa ccacggagag aaatggttca 480
 gagacacaga cactggtgac cctccataat aacaatggaa caaaaatttc cacatgggct 540
 gatgaaatta aggactctgg agaaacgatc agaactgcta tccacattgg agtgggagtc 600
 tctgctgggt tgaccctggc acttatcatt ggtgtcttaa tccttaaagtg gtattcctgt 660
 aagaaaaaga agttatcgag tttgagcctt attacactgg ccaacttggc tccaggaggg 720
 ttggcaaatg caggagcagt caggattcgc tctgaggaaa atatctacac catcgaggag 780
 aacgtatatg aagtggagaa ttcaaatgag tactactgct acgtcaacag ccagcagcca 840
 tcctga 846

-continued

<210> SEQ ID NO 12
 <211> LENGTH: 281
 <212> TYPE: PRT
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 12

Met Phe Ser Gly Leu Thr Leu Asn Cys Val Leu Leu Leu Leu Gln Leu
 1 5 10 15
 Leu Leu Ala Arg Ser Leu Glu Asn Ala Tyr Val Phe Glu Val Gly Lys
 20 25 30
 Asn Ala Tyr Leu Pro Cys Ser Tyr Thr Leu Ser Thr Pro Gly Ala Leu
 35 40 45
 Val Pro Met Cys Trp Gly Lys Gly Phe Cys Pro Trp Ser Gln Cys Thr
 50 55 60
 Asn Glu Leu Leu Arg Thr Asp Glu Arg Asn Val Thr Tyr Gln Lys Ser
 65 70 75 80
 Ser Arg Tyr Gln Leu Lys Gly Asp Leu Asn Lys Gly Asp Val Ser Leu
 85 90 95
 Ile Ile Lys Asn Val Thr Leu Asp Asp His Gly Thr Tyr Cys Cys Arg
 100 105 110
 Ile Gln Phe Pro Gly Leu Met Asn Asp Lys Lys Leu Glu Leu Lys Leu
 115 120 125
 Asp Ile Lys Ala Ala Lys Val Thr Pro Ala Gln Thr Ala His Gly Asp
 130 135 140
 Ser Thr Thr Ala Ser Pro Arg Thr Leu Thr Thr Glu Arg Asn Gly Ser
 145 150 155 160
 Glu Thr Gln Thr Leu Val Thr Leu His Asn Asn Asn Gly Thr Lys Ile
 165 170 175
 Ser Thr Trp Ala Asp Glu Ile Lys Asp Ser Gly Glu Thr Ile Arg Thr
 180 185 190
 Ala Ile His Ile Gly Val Gly Val Ser Ala Gly Leu Thr Leu Ala Leu
 195 200 205
 Ile Ile Gly Val Leu Ile Leu Lys Trp Tyr Ser Cys Lys Lys Lys Lys
 210 215 220
 Leu Ser Ser Leu Ser Leu Ile Thr Leu Ala Asn Leu Pro Pro Gly Gly
 225 230 235 240
 Leu Ala Asn Ala Gly Ala Val Arg Ile Arg Ser Glu Glu Asn Ile Tyr
 245 250 255
 Thr Ile Glu Glu Asn Val Tyr Glu Val Glu Asn Ser Asn Glu Tyr Tyr
 260 265 270
 Cys Tyr Val Asn Ser Gln Gln Pro Ser
 275 280

<210> SEQ ID NO 13
 <211> LENGTH: 1578
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

atgtgggagg ctcagttcct gggcttgctg tttctgcagc cgctttgggt ggctccagtg 60
 aagcctctcc agccaggggc tgagggtccc gtggtgtggg cccaggaggg ggctcctgcc 120
 cagctcccct gcagccccc aatccccctc caggatctca gccttctgcg aagagcaggg 180
 gtcacttggc agcatcagcc agacagtggc ccgcccgtg ccgccccgg ccataccctg 240
 gccccggcc ctcaccggc ggcgcctcc tcttgggggc ccaggcccc ccgetacag 300

-continued

```

gtgctgagcg tgggtcccgg aggcctgcgc agcgggaggg tgcccctgca gccccgcgtc 360
cagctggatg agcgcggccg gcagcgcggg gacttctcgc tatggctgcg cccagcccgg 420
cgcgcgagcg cgggcgagta ccgcgcccgc gtgcacctca gggaccgcgc cctctcctgc 480
cgctccgctc tgcgcctggg ccaggcctcg atgactgcca gccccccagg atctctcaga 540
gcctccgact gggtcatttt gaactgctcc ttcagccgcc ctgaccgccc agcctctgtg 600
cattggttcc ggaaccgggg ccagggccga gtccctgtcc gggagtcccc ccatcaccac 660
ttagcggaaa gcttctctt cctgccccaa gtcagcccca tggactctgg gcctggggc 720
tgcacctca cctacagaga tggcttcaac gtctccatca tgtataacct cactgttctg 780
ggtctggagc ccccaactcc cttgacagtg tacgctggag caggttccag ggtggggctg 840
ccctgcgcgc tgctgctgg tgtggggacc cggcttttcc tcaactgcaa gtggactcct 900
cctgggggag gccctgacct cctggtgact ggagacaatg gcgactttac ccttcgacta 960
gaggatgtga gccaggccca ggctgggacc tacacctgcc atatccatct gcaggaacag 1020
cagctcaatg ccactgtcac attggcaatc atcacagtga ctcccaaatc ctttgggtca 1080
cctggatccc tggggaagct gctttgtgag gtgactccag tatctggaca agaacgcttt 1140
gtgtggagct ctctggacac cccatcccag aggagtttct caggaccttg gctggaggca 1200
caggaggccc agctcctttc ccagccttgg caatgccagc tgtaccaggg ggagaggctt 1260
cttgagcag cagtgtactt cacagagctg tctagcccag gtgcccaacg ctctgggaga 1320
gccccaggty ccctcccagc aggccacctc ctgetgtttc tcatecttgg tgcctttct 1380
ctgctccttt tgggtgactgg agcctttggc tttcaccttt ggagaagaca gtggcgacca 1440
agacgatttt ctgccttaga gcaagggatt caccctccgc aggetcagag caagatagag 1500
gagctggagc aagaaccgga gccggagccg gagccggaac cggagcccga gcccgagccc 1560
gagccggagc agctctga 1578

```

<210> SEQ ID NO 14

<211> LENGTH: 525

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

```

Met Trp Glu Ala Gln Phe Leu Gly Leu Leu Phe Leu Gln Pro Leu Trp
1           5           10          15
Val Ala Pro Val Lys Pro Leu Gln Pro Gly Ala Glu Val Pro Val Val
20          25          30
Trp Ala Gln Glu Gly Ala Pro Ala Gln Leu Pro Cys Ser Pro Thr Ile
35          40          45
Pro Leu Gln Asp Leu Ser Leu Leu Arg Arg Ala Gly Val Thr Trp Gln
50          55          60
His Gln Pro Asp Ser Gly Pro Pro Ala Ala Ala Pro Gly His Pro Leu
65          70          75          80
Ala Pro Gly Pro His Pro Ala Ala Pro Ser Ser Trp Gly Pro Arg Pro
85          90          95
Arg Arg Tyr Thr Val Leu Ser Val Gly Pro Gly Gly Leu Arg Ser Gly
100         105         110
Arg Leu Pro Leu Gln Pro Arg Val Gln Leu Asp Glu Arg Gly Arg Gln
115         120         125
Arg Gly Asp Phe Ser Leu Trp Leu Arg Pro Ala Arg Arg Ala Asp Ala
130         135         140

```


-continued

Gly Glu Tyr Arg Ala Ala Val His Leu Arg Asp Arg Ala Leu Ser Cys
 145 150 155 160
 Arg Leu Arg Leu Arg Leu Gly Gln Ala Ser Met Thr Ala Ser Pro Pro
 165 170 175
 Gly Ser Leu Arg Ala Ser Asp Trp Val Ile Leu Asn Cys Ser Phe Ser
 180 185 190
 Arg Pro Asp Arg Pro Ala Ser Val His Trp Phe Arg Asn Arg Gly Gln
 195 200 205
 Gly Arg Val Pro Val Arg Glu Ser Pro His His His Leu Ala Glu Ser
 210 215 220
 Phe Leu Phe Leu Pro Gln Val Ser Pro Met Asp Ser Gly Pro Trp Gly
 225 230 235 240
 Cys Ile Leu Thr Tyr Arg Asp Gly Phe Asn Val Ser Ile Met Tyr Asn
 245 250 255
 Leu Thr Val Leu Gly Leu Glu Pro Pro Thr Pro Leu Thr Val Tyr Ala
 260 265 270
 Gly Ala Gly Ser Arg Val Gly Leu Pro Cys Arg Leu Pro Ala Gly Val
 275 280 285
 Gly Thr Arg Ser Phe Leu Thr Ala Lys Trp Thr Pro Pro Gly Gly Gly
 290 295 300
 Pro Asp Leu Leu Val Thr Gly Asp Asn Gly Asp Phe Thr Leu Arg Leu
 305 310 315 320
 Glu Asp Val Ser Gln Ala Gln Ala Gly Thr Tyr Thr Cys His Ile His
 325 330 335
 Leu Gln Glu Gln Gln Leu Asn Ala Thr Val Thr Leu Ala Ile Ile Thr
 340 345 350
 Val Thr Pro Lys Ser Phe Gly Ser Pro Gly Ser Leu Gly Lys Leu Leu
 355 360 365
 Cys Glu Val Thr Pro Val Ser Gly Gln Glu Arg Phe Val Trp Ser Ser
 370 375 380
 Leu Asp Thr Pro Ser Gln Arg Ser Phe Ser Gly Pro Trp Leu Glu Ala
 385 390 395 400
 Gln Glu Ala Gln Leu Leu Ser Gln Pro Trp Gln Cys Gln Leu Tyr Gln
 405 410 415
 Gly Glu Arg Leu Leu Gly Ala Ala Val Tyr Phe Thr Glu Leu Ser Ser
 420 425 430
 Pro Gly Ala Gln Arg Ser Gly Arg Ala Pro Gly Ala Leu Pro Ala Gly
 435 440 445
 His Leu Leu Leu Phe Leu Ile Leu Gly Val Leu Ser Leu Leu Leu Leu
 450 455 460
 Val Thr Gly Ala Phe Gly Phe His Leu Trp Arg Arg Gln Trp Arg Pro
 465 470 475 480
 Arg Arg Phe Ser Ala Leu Glu Gln Gly Ile His Pro Pro Gln Ala Gln
 485 490 495
 Ser Lys Ile Glu Glu Leu Glu Gln Glu Pro Glu Pro Glu Pro Glu Pro
 500 505 510
 Glu Pro Glu Pro Glu Pro Glu Pro Glu Pro Glu Gln Leu
 515 520 525

<210> SEQ ID NO 15
 <211> LENGTH: 1566
 <212> TYPE: DNA
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 15

-continued

```

atgagggagg acctgctcct tggctttttg cttctgggac tgctttggga agctccagtt    60
gtgtcttcag ggcttgggaa agagctcccc gtggtgtggg cccaggaggg agctcccgtc    120
catcttcctt gcagcctcaa atcccccaac ctggatccta actttctacg aagaggaggg    180
gttatctggc aacatcaacc agacagtggc caaccactc ccatcccggc ccttgacctt    240
caccagggga tgcctcgcct tagacaacct gcaccgggtc gctacacggt gctgagcgtg    300
gtcccaggag gcctgcgcag cgggaggcag ccctgcac cccacgtgca gctggaggag    360
cgcgccctcc agcgcgggga cttctctctg tggttgccgc cagctctgcg caccgatgcg    420
ggcgagtacc acgccaccgt ggcctccccg aaccgcgccc tctcctgcag tctccgctg    480
cgcgctggcc aggcctcgat gattgctagt ccctcaggag tctcaagct gtctgattgg    540
gtccttttga actgctcctt cagccgtcct gaccgcccag tctctgtgca ctggttccag    600
ggccagaacc gagtgctgtt ctacaactca ccgcgtcatt ttttagctga aactttcctg    660
ttactgcccc aagtcagccc cctggactct gggacctggg gctgtgtcct cacctacaga    720
gatggcttca atgtctccat cacgtacaac ctcaaggttc tgggtctgga gcccgtagcc    780
cctctgacag tgtacgtgc tgaaggttct agggaggagc tgcctgtca tttgccccca    840
ggagtgggga ccccttcttt gctcattgcc aagtggactc ctctggagg aggtcctgag    900
ctccccgtgg ctggaagag tggcaatfff acccttcacc ttgaggctgt gggctctggca    960
caggctggga cctacacctg tagcatccat ctgcagggac agcagctcaa tgccactgtc   1020
acgttggcgg tcatcacagt gactcccaaa tccttcgggt tacctggctc ccgggggaag   1080
ctgttggtg aggtaacccc ggcatctgga aaggaaagat ttgtgtggcg tcccctgaac   1140
aatctgtcca ggagttgccc gggccctgtg ctggagattc aggaggccag gctccttgct   1200
gagcagatggc agtgtcagct gtacgagggc cagaggcttc ttggagcgc agtgtacgcc   1260
gcagagteta gctcaggcgc ccacagtgct aggagaatct cagggtgacct taaaggaggc   1320
catctcgttc tcgtttcat ccttgggtgcc ctctccctgt tccttttggg ggccggggcc   1380
tttggtttc actggtggag aaaacagttg ctactgagaa gattttctgc cttagaacat   1440
gggattcagc catttccggc tcagaggaag atagaggagc tggagcgaga actggagacg   1500
gagatgggac aggagccgga gcccagaccg gagccacagc tggagccaga gcccaggcag   1560
ctctga                                         1566

```

```

<210> SEQ ID NO 16
<211> LENGTH: 521
<212> TYPE: PRT
<213> ORGANISM: Mus sp.

```

```

<400> SEQUENCE: 16

```

```

Met Arg Glu Asp Leu Leu Leu Gly Phe Leu Leu Leu Gly Leu Leu Trp
1          5          10          15
Glu Ala Pro Val Val Ser Ser Gly Pro Gly Lys Glu Leu Pro Val Val
20          25          30
Trp Ala Gln Glu Gly Ala Pro Val His Leu Pro Cys Ser Leu Lys Ser
35          40          45
Pro Asn Leu Asp Pro Asn Phe Leu Arg Arg Gly Gly Val Ile Trp Gln
50          55          60
His Gln Pro Asp Ser Gly Gln Pro Thr Pro Ile Pro Ala Leu Asp Leu
65          70          75          80
His Gln Gly Met Pro Ser Pro Arg Gln Pro Ala Pro Gly Arg Tyr Thr

```


-continued

85			90			95									
Val	Leu	Ser	Val	Ala	Pro	Gly	Gly	Leu	Arg	Ser	Gly	Arg	Gln	Pro	Leu
			100					105					110		
His	Pro	His	Val	Gln	Leu	Glu	Glu	Arg	Gly	Leu	Gln	Arg	Gly	Asp	Phe
			115					120					125		
Ser	Leu	Trp	Leu	Arg	Pro	Ala	Leu	Arg	Thr	Asp	Ala	Gly	Glu	Tyr	His
						135					140				
Ala	Thr	Val	Arg	Leu	Pro	Asn	Arg	Ala	Leu	Ser	Cys	Ser	Leu	Arg	Leu
					150					155					160
Arg	Val	Gly	Gln	Ala	Ser	Met	Ile	Ala	Ser	Pro	Ser	Gly	Val	Leu	Lys
				165						170				175	
Leu	Ser	Asp	Trp	Val	Leu	Leu	Asn	Cys	Ser	Phe	Ser	Arg	Pro	Asp	Arg
			180					185					190		
Pro	Val	Ser	Val	His	Trp	Phe	Gln	Gly	Gln	Asn	Arg	Val	Pro	Val	Tyr
			195					200				205			
Asn	Ser	Pro	Arg	His	Phe	Leu	Ala	Glu	Thr	Phe	Leu	Leu	Leu	Pro	Gln
			210				215				220				
Val	Ser	Pro	Leu	Asp	Ser	Gly	Thr	Trp	Gly	Cys	Val	Leu	Thr	Tyr	Arg
					230					235					240
Asp	Gly	Phe	Asn	Val	Ser	Ile	Thr	Tyr	Asn	Leu	Lys	Val	Leu	Gly	Leu
				245					250					255	
Glu	Pro	Val	Ala	Pro	Leu	Thr	Val	Tyr	Ala	Ala	Glu	Gly	Ser	Arg	Val
			260					265					270		
Glu	Leu	Pro	Cys	His	Leu	Pro	Pro	Gly	Val	Gly	Thr	Pro	Ser	Leu	Leu
			275				280					285			
Ile	Ala	Lys	Trp	Thr	Pro	Pro	Gly	Gly	Gly	Pro	Glu	Leu	Pro	Val	Ala
						295					300				
Gly	Lys	Ser	Gly	Asn	Phe	Thr	Leu	His	Leu	Glu	Ala	Val	Gly	Leu	Ala
					310					315					320
Gln	Ala	Gly	Thr	Tyr	Thr	Cys	Ser	Ile	His	Leu	Gln	Gly	Gln	Gln	Leu
				325					330					335	
Asn	Ala	Thr	Val	Thr	Leu	Ala	Val	Ile	Thr	Val	Thr	Pro	Lys	Ser	Phe
			340					345					350		
Gly	Leu	Pro	Gly	Ser	Arg	Gly	Lys	Leu	Leu	Cys	Glu	Val	Thr	Pro	Ala
			355				360					365			
Ser	Gly	Lys	Glu	Arg	Phe	Val	Trp	Arg	Pro	Leu	Asn	Asn	Leu	Ser	Arg
			370				375				380				
Ser	Cys	Pro	Gly	Pro	Val	Leu	Glu	Ile	Gln	Glu	Ala	Arg	Leu	Leu	Ala
					390					395					400
Glu	Arg	Trp	Gln	Cys	Gln	Leu	Tyr	Glu	Gly	Gln	Arg	Leu	Leu	Gly	Ala
				405					410					415	
Thr	Val	Tyr	Ala	Ala	Glu	Ser	Ser	Ser	Gly	Ala	His	Ser	Ala	Arg	Arg
			420					425					430		
Ile	Ser	Gly	Asp	Leu	Lys	Gly	Gly	His	Leu	Val	Leu	Val	Leu	Ile	Leu
			435				440					445			
Gly	Ala	Leu	Ser	Leu	Phe	Leu	Leu	Val	Ala	Gly	Ala	Phe	Gly	Phe	His
					455						460				
Trp	Trp	Arg	Lys	Gln	Leu	Leu	Leu	Arg	Arg	Phe	Ser	Ala	Leu	Glu	His
					470					475					480
Gly	Ile	Gln	Pro	Phe	Pro	Ala	Gln	Arg	Lys	Ile	Glu	Glu	Leu	Glu	Arg
				485					490					495	
Glu	Leu	Glu	Thr	Glu	Met	Gly	Gln	Glu	Pro	Glu	Pro	Glu	Pro	Glu	Pro
			500					505						510	

-continued

Gln Leu Glu Pro Glu Pro Arg Gln Leu
515 520

What is claimed is:

1. A method of treating a subject afflicted with a hematologic cancer comprising administering to the subject a therapeutically effective amount of: (i) a first inhibitor that is an inhibitor of PD-L1 or PD-1 and (ii) a second inhibitor that is an inhibitor of LAG-3; and wherein the method further comprises a step of transient lymphodepletion.
2. The method of claim 1 wherein an inhibitor of PD-L1 is administered in combination with an inhibitor of LAG-3.
3. The method of claim 1 wherein an inhibitor of PD-1 is administered in combination with an inhibitor of LAG-3.
4. The method of claim 1, wherein the PD-1, PD-L1, or LAG-3 inhibitor:
 - (a) is an antibody or antigen-binding fragment thereof, that binds to one or more targets selected from the group consisting of PD-1, PD-L1, and LAG-3; or
 - (b) is a bispecific or multispecific antibody, or antigen binding fragment thereof, selective for (i) PD-1 or PD-L1 and (ii) LAG-3.
5. The method of claim 1, wherein a combination of inhibitors comprising a first inhibitor that selectively inhibits or blocks PD-1 or PD-L1 and a second inhibitor that selectively inhibits or blocks LAG-3 is administered.
6. The method of claim 5, wherein the first inhibitor, second inhibitor, or both inhibitors, is an antibody or an antigen binding fragment thereof, which specifically binds to (i) PD-1 or PD-L1 and/or (ii) LAG-3.
7. The method of claim 6, wherein:
 - (a) said antibody, or antigen binding fragment thereof, is murine, chimeric, humanized, composite, or human; or
 - (b) said antibody, or antigen binding fragment thereof, is detectably labeled, comprises an effector domain, comprises an Fc domain, and/or is selected from the group consisting of Fv, Fav, F(ab')₂, Fab', dsFv, scFv, sc(Fv)₂, and diabodies fragments.
8. The method of claim 1, wherein said first and/or second inhibitor is administered in a pharmaceutically acceptable formulation.
9. The method of claim 1, further comprising administering to the subject a therapeutic agent for treating the hematologic cancer.
10. The method of claim 1, wherein:
 - (a) sublethal whole body irradiation is used for transient lymphodepletion; or
 - (b) the step of lymphodepletion occurs before, concurrently with, or after the step of first and/or second inhibitor administration.
11. The method of claim 1, wherein the hematologic cancer is selected from the group consisting of:
 - (a) multiple myeloma, acute lymphocytic leukemia, acute myeloid leukemia, chronic lymphocytic leukemia, small lymphocytic lymphoma, non-Hodgkin's lymphoma, Hodgkin's lymphoma, mantle cell lymphoma, follicular lymphoma, Waldenstrom's macroglobulinemia, B-cell lymphoma and diffuse large B-cell lymphoma, precursor B-lymphoblastic leukemia/lymphoma, B-cell chronic lymphocytic leukemia/small lymphocytic lymphoma, B-cell prolymphocytic leukemia, lymphoplasmacytic lymphoma, splenic marginal zone B-cell lymphoma (with or without villous lymphocytes), hairy cell leukemia, plasma cell myeloma/plasmacytoma, extranodal marginal zone B-cell lymphoma of the MALT type, nodal marginal zone B-cell lymphoma (with or without monocytoid B cells), Burkitt's lymphoma; precursor T-lymphoblastic lymphoma/leukemia, T-cell prolymphocytic leukemia, T-cell granular lymphocytic leukemia, aggressive NK cell leukemia, adult T-cell lymphoma/leukemia (HTLV 1-positive), nasal-type extranodal NK/T-cell lymphoma, enteropathy-type T-cell lymphoma, hepatosplenic γ - δ T-cell lymphoma, subcutaneous panniculitis-like T-cell lymphoma, mycosis fungoides/Sezary syndrome, anaplastic large cell lymphoma (T/null cell, primary cutaneous type), anaplastic large cell lymphoma (T-/null-cell, primary systemic type), peripheral T-cell lymphoma not otherwise characterized, angioimmunoblastic T-cell lymphoma, polycythemia vera (PV), myelodysplastic syndrome (MDS), indolent Non-Hodgkin's Lymphoma (iNHL), and aggressive Non-Hodgkin's Lymphoma (aNHL); or
 - (b) B-cell lymphoma, myeloid leukemia, and multiple myeloma.
12. The method of claim 1, wherein the subject is a human.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 11,708,412 B2
APPLICATION NO. : 16/724759
DATED : July 25, 2023
INVENTOR(S) : Bryon Duane Johnson et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:


In the Claims

At Column 119, Claim number 2, Line number 15, delete “The method of claim 1 wherein” and insert --The method of claim 1, wherein--.

At Column 119, Claim number 3, Line number 17, delete “The method of claim 1 wherein” and insert --The method of claim 1, wherein--.

At Column 120, Claim number 11, Line number 36, delete “mycosis fungoides/Sezary syndrome” and insert --mycosis fungoides/Sézary syndrome--.

Signed and Sealed this
Twenty-sixth Day of December, 2023



Katherine Kelly Vidal
Director of the United States Patent and Trademark Office