



US011773063B1

(12) **United States Patent**  
**Gray et al.**(10) **Patent No.: US 11,773,063 B1**  
(45) **Date of Patent: Oct. 3, 2023**(54) **PHARMACEUTICALLY ACCEPTABLE  
SALTS AND COMPOSITIONS THEREOF**(71) Applicant: **BECKLEY PSYTECH LIMITED,**  
Oxford (GB)(72) Inventors: **Jason Gray**, Oxford (GB); **Susana Del  
Rio Gancedo**, Oxford (GB); **Dita  
Davis**, Oxford (GB); **David James  
Pearson**, Oxford (GB); **Daniel Rixson**,  
Oxford (GB)(73) Assignee: **Beckley Psytech Limited**, Oxford (GB)(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 0 days.(21) Appl. No.: **17/941,410**(22) Filed: **Sep. 9, 2022**(30) **Foreign Application Priority Data**

Aug. 19, 2022 (GB) ..... 2212116

(51) **Int. Cl.**  
**C07D 209/16** (2006.01)(52) **U.S. Cl.**  
CPC ..... **C07D 209/16** (2013.01); **C07B 2200/13**  
(2013.01)(58) **Field of Classification Search**  
CPC ..... **C07D 209/16**  
See application file for complete search history.(56) **References Cited**

## U.S. PATENT DOCUMENTS

2,774,763	A	12/1956	Garbrecht
2,997,470	A	8/1961	Pioch
3,078,214	A	2/1963	Hofmann et al.
3,224,945	A	12/1965	Tyler, Jr.
5,811,436	A	9/1998	Leonard et al.
11,518,742	B2	12/2022	Feilding-Mellen et al.
11,518,743	B2	12/2022	Feilding-Mellen et al.
2005/0019411	A1	1/2005	Colombo et al.
2012/0108510	A1	5/2012	Young et al.
2017/0348303	A1	12/2017	Bosse et al.
2018/0021326	A1	1/2018	Stamets
2018/0147142	A1	5/2018	Knight
2020/0179349	A1	6/2020	Yun et al.
2020/0187777	A1	6/2020	Luderer et al.
2021/0058956	A1	2/2021	Chatterjee et al.
2021/0069170	A1	3/2021	Stamets
2021/0085671	A1	3/2021	Chadeayne
2021/0322743	A1	10/2021	Rinti et al.
2022/0062238	A1*	3/2022	Layzell ..... A61K 31/4045
2022/0396552	A1	12/2022	Feilding-Mellen et al.
2023/0031944	A1	2/2023	Feilding-Mellen et al.

## FOREIGN PATENT DOCUMENTS

CH	578565	A5	8/1976
CN	103816150	A	5/2014
CN	113288883	A	8/2021
DE	2617738	A1	11/1976
EP	0026899	A1	4/1981
EP	3868364	A1	8/2021

GB	981192	A	1/1965
GB	1410349	A	10/1975
GB	1584464	A	2/1981
GB	2596884	A	1/2022
WO	0115677	A2	3/2001
WO	0115677	A3	3/2001
WO	0238142	A2	5/2002
WO	2004000849	A2	12/2003
WO	2008003028	A2	1/2008
WO	2010054202	A2	5/2010
WO	2012/173701	A1	12/2012
WO	2013063492	A1	5/2013
WO	2018195455	A1	10/2018
WO	2019073379	A1	4/2019
WO	2019081764	A1	5/2019
WO	2019173797	A1	9/2019
WO	2019246532	A1	12/2019
WO	2020169850	A1	8/2020
WO	2020169851	A1	8/2020
WO	2020176597	A1	9/2020
WO	2020181194	A1	9/2020
WO	2020212951	A1	10/2020
WO	2021003467	A1	1/2021
WO	2021041407	A1	3/2021
WO	2021089872	A1	5/2021
WO	2021209815	A1	10/2021
WO	2021222885	A1	11/2021
WO	2021225796	A1	11/2021
WO	2021250435	A1	12/2021
WO	WO 2022/246572	*	12/2022
WO	2023002005	A1	1/2023

## OTHER PUBLICATIONS

Berge et al., "Pharmaceutical Salts" *Journal of Pharmaceutical Sciences*; 1977; vol. 66, No. 1; pp. 1-19.\*  
Registry No. 2761182-82-3, File Registry on STN, entered STN: Mar. 3, 2022.\*  
International Search Report in International Application No. PCT/GB2021/051475, dated Sep. 16, 2021 (3 pages).  
International Search Report in International Application No. PCT/GB2021/051476, dated Sep. 15, 2021 (5 pages).  
Uthaug, M.V et al., "Prospective examination of synthetic 5-methoxy-N,N-dimethyltryptamine inhalation: effects on IL-6, cortisol levels, affect, and non-judgment," *Psychopharmacology*, vol. 237, pp. 773-785 (2020).  
Database Registry. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1H-Indole-3-ethanamine, 5-methoxy-N,N dimethyl-, benzoate (1:1); RN 282103-25-7; ED Aug. 1, 2000 (1 page).  
Benington, F. et al., "Synthesis of O- and N-Methylated Derivatives of 5-Hydroxytryptamine," *The Journal of Organic Chemistry*, vol. 23, pp. 1977-1979 (1958).  
Falkenberg, G. et al., "The Crystal and Molecular Structure of 5-Methoxy-(N,N)-dimethyltryptamine Hydrochloride," *Acta Crystallographica Section B*, vol. 27, pp. 411-418 (1971).

(Continued)

*Primary Examiner* — Rebecca L Anderson  
(74) *Attorney, Agent, or Firm* — Bookoff McAndrews, PLLC

(57) **ABSTRACT**

Pharmaceutically acceptable salts of 5-methoxy-N,N-dimethyltryptamine are described, as well as compositions/formulations and uses thereof as a medicament.

**10 Claims, 112 Drawing Sheets**

(56)

## References Cited

## OTHER PUBLICATIONS

- Roseman, L. et al., "Increased amygdala responses to emotional faces after psilocybin treatment-resistant depression," *Neuropharmacology*, vol. 142, pp. 263-269 (2018).
- Griffiths, R.R. et al., "Psilocybin produces substantial and sustained decreases in depression and anxiety in patients with life-threatening cancer: a randomized double-blind trial," *Journal of Psychopharmacology*, vol. 30, pp. 1181-1197 (2016).
- Carhart-Harris, R.L. et al., "Psilocybin with psychological support for treatment-resistant depression: six-month follow-up," *Psychopharmacology*, vol. 235, pp. 399-408 (2018).
- Monte, A.P. et al., "Stereoselective LSD-like Activity in a Series of d-Lysergic Acid Amides of (R) and (S)-2-Aminoalkanes," *Journal of Medicinal Chemistry*, vol. 38, pp. 958-966 (1995).
- Ishii, H. et al., "Studies on Lysergic Acid Diethylamide and Related Compounds. Part 8. Structural Identification of New Metabolites of Lysergic Acid Diethylamide obtained by Microbial Transformation using *Streptomyces roseochromogenes*," *Journal of the Chemical Society, Perkin Transactions 1: Organic & Bio-organic Chemistry*, vol. 4, pp. 902-905 (1980).
- Nakahara, Y. et al., "Studies on Lysergic Acid Diethylamide and Related Compounds. III. Improvement of Amidation of Lysergic Acid," *Yakugaku Zasshi*, vol. 94, pp. 407-412 (1974).
- Huang, X. et al., "Drug Discrimination and Receptor Binding Studies of N-Isopropyl Lysergamide Derivatives," *Pharmacology Biochemistry and Behavior*, vol. 47, pp. 667-673 (1994).
- Ishii, H. et al., "Studies on Lysergic Acid Diethylamide and Related Compounds. IX. Microbial Transformation of Amides Related to Lysergic Acid Diethylamide by *Streptomyces roseochromogenes*," *Chemical & Pharmaceutical Bulletin*, vol. 27, pp. 3029-3038 (1979).
- Johnson, F.N. et al., "Emetic Activity of Reduced Lysergamides," *Journal of Medicinal Chemistry*, vol. 16, pp. 532-537 (1973).
- Vangveravong, S. et al., "Synthesis and Serotonin Receptor Affinities of a Series of trans-2-(Indol-3-yl) cyclopropylamine Derivatives," *Journal of Medicinal Chemistry*, vol. 41, pp. 4995-5001 (1998).
- Schneller, S.W. et al., "Synthesis of 4-Amino-1H-pyrrolo[2,3-b]pyridine (1,7-Dideazaadenine) and 1H-Pyrrolo[2,3-b]pyridin-4-ol (1,7-Dideazahypoxanthine)," *The Journal of Organic Chemistry A*, vol. 45, pp. 4045-4048 (1980).
- Singh, S.K. et al., "An ab Initio Study of the Effect of Substituents on the  $n \rightarrow \pi^*$  Interactions between 7-Azaindole and 2,6-Difluorosubstituted Pyridines," *The Journal of Physical Chemistry A*, vol. 120, pp. 6258-6269 (2016).
- Monson, C.M. et al., "MDMA-facilitated cognitive-behavioural conjoint therapy for posttraumatic stress disorder: an uncontrolled trial," *European Journal of Psychotraumatology*, vol. 11, pp. 1-7 (2020).
- Wolfson, P.E. et al., "MDMA-assisted psychotherapy for treatment of anxiety and other psychological distress related to life-threatening illnesses: a randomized pilot study," *Scientific Reports*, vol. 10, pp. 1-15 (2020).
- Yazar-Klosinski, B.B. et al., "Potential Psychiatric Uses for MDMA," *Developments*, vol. 101, pp. 194-196 (2017).
- PharmaTher Holdings Ltd. Dec. 14, 2021. PharmaTher Announces Positive Research Results for LSD Microneedle Patch. Press Release. <URL: <https://psychedelicinvest.com/pharmather-announces-positive-research-results-for-isd-microneedle-patch/>>.
- Szabo, A. et al., "Psychedelic N,N-Dimethyltryptamine and 5-Methoxy-N,N-Dimethyltryptamine Modulate Innate and Adaptive Inflammatory Responses through the Sigma-1 Receptor of Human Monocyte-Derived Dendritic Cells," *PLoS One*, vol. 9, pp. 1-12 (2014).
- Sherwood, A.M. et al., "Synthesis and Characterization of 5-MeO-DMT Succinate for Clinical Use," *ACS Omega*, vol. 5, pp. 32067-32075 (2020).
- Galeffi, C. et al., "N,N-Dimethyl-5-Methoxytryptamine, a Component of a Dart Poison of the Yanoáma Indians," *Journal of Natural Products*, vol. 46, pp. 586-587 (1983).
- CAPLUS. Chemical Abstracts Service: Columbus. CAPLUS Accession No. 2017:1595854. Title: Preparation of tetrahydropyrroloindolylcycloalkylacrylic acid derivatives and analogs for us as estrogen receptor modulators. Inventor: Huang, P.Q. et al. (4 pages).
- Sohlberg, E. et al., "The impact of the site of blood sampling on pharmacokinetic parameters following sublingual dosing to dogs," *Journal of Pharmacological and Toxicological Methods*, vol. 67, pp. 1-4 (2013).
- Illum, L. et al., "The Effect of Blood Sampling Site and Physicochemical Characteristics of Drugs on Bioavailability after Nasal Administration in the Sheep Model," *Pharmaceutical Research*, vol. 20, pp. 1474-1484 (2003).
- Gupta, S.P., "QSAR Studies on Drugs Acting at the Central Nervous System," *Chemical Reviews*, vol. 89, pp. 1765-1800 (1989).
- Stoll, A. et al., "49. Amide der stereoisomeren Lysergsäuren und Dihydro-lysergsäuren," *Helvetica Chimica Acta*, vol. 38, pp. 421-433 (1955).
- Halberstadt, A.L. et al., "Pharmacological characterization of the LSD analog N-ethyl-N-cyclopropyl lysergamide (ECPLA)," *Psychopharmacology*, vol. 236, pp. 799-808 (2019).
- McKenna, D.J. et al., "Differential interactions of indolealkylamines with 5-hydroxytryptamine receptor subtypes," *Neuropharmacology*, vol. 29, pp. 193-198 (1990).
- Glennon, R.A. et al., "Serotonin Receptor Binding Affinities of Tryptamine Analogues," *Journal of Medicinal Chemistry*, vol. 22, pp. 428-432 (1979).
- Klein, A.K. et al., "Investigation of the Structure—Activity Relationships of Psilocybin Analogues," *ACS Pharmacology & Translational Science*, vol. 4, pp. 533-542 (2021).
- Sard, H. et al., "SAR of psilocybin analogs: Discovery of a selective 5-HT<sub>2C</sub> agonist," *Bioorganic & Medicinal Chemistry Letters*, vol. 15, pp. 4555-4559 (2005).
- Medline Plus, "Cancer," National Institute of Health (2007) <URL: [www.nlm.nih.gov/medlineplus.cancer.html](http://www.nlm.nih.gov/medlineplus.cancer.html)>.
- Lala, P.K. et al., "Role of nitric oxide in tumor progression: Lessons from experimental tumors," *Cancer and Metastasis Reviews*, vol. 17, pp. 91-106 (1998).
- Golub, T.R. et al., *Molecular Classification of Cancer: Class Discovery and Class Prediction by Gene Expression Monitoring*, vol. 286, pp. 531-537 (1999).
- Glässer, A., "Some Pharmacological Actions of D-Lysergic Acid Methyl Carbinolamide," *Nature*, vol. 189, pp. 313-314 (1961).
- Lyon, R.A. et al., "Indolealkylamine analogs share 5-HT<sub>2</sub> binding characteristics with phenylalkylamine hallucinogens," *European Journal of Pharmacology*, vol. 145, pp. 291-297 (1988).
- U.S. Appl. No. 18/065,030, filed Dec. 13, 2022 (55 pages).
- U.S. Appl. No. 18/162,976, filed Feb. 1, 2023 (176 pages).
- Shen et al., "Psychedelic 5-Methoxy-N,N-dimethyltryptamine: Metabolism, Pharmacokinetics, Drug Interactions, and Pharmacological Actions," *Curr Drug Metab.*, Oct. 2010, 11(8), pp. 659-666.
- DMT-Nexus Wiki. Feb. 10, 2023. Psychedelic Compounds Chemical and Physical Properties. <URL: [http://wiki.dmt-nexus.me/Psychedelic\\_Compounds\\_Chemical\\_and\\_Physical\\_Properties#DMT\\_Benzoate](http://wiki.dmt-nexus.me/Psychedelic_Compounds_Chemical_and_Physical_Properties#DMT_Benzoate)>.
- Shulgin, A. et al. TiHKAL: The Continuation. #38. 5-MEO-DMT. Tryptamine, 5-Methoxy-N,N-Dimethyl; Indole, 5-Methoxy-3-[2-(Dimethylamino)]; 5-Methoxy-N,N-Dimethyltryptamine; 5-Methoxy-3-[2-(Dimethylamino)Ethyl]indole; N,N,O-Trimethylserotonin; N,N,O-Tms; Bufotenine Methyl Ether; O-Methylbufotenine; OMB., Feb. 21, 2015. <URL: [https://erowid.org/library/books\\_online/tihkal/tihkal38.shtml](https://erowid.org/library/books_online/tihkal/tihkal38.shtml)>.
- CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 4H-Pyrrolo[2,3-b]pyridine-4-one, 1,7-dihydro-. CHEMCATS Accession No. 1756550559. Catalog Name: Sagechem Limited Product List. Order No. Catalog: S243355. CAS Registry No. 1076197-59-5, May 28, 2020 (1 page).
- CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1-methyl-1H-pyrrolo[2,3-b]pyridin-4-ol. CHEMCATS Accession No. 1545199867. Catalog Name: Azepine Product List. Order No. Catalog: AZ04819515. CAS Registry No. 1781876-60-5, Mar. 1, 2019 (1 page).
- CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1-methylpyrrolo[2,3-b]pyridin-4-ol. CHEMCATS Acces-

(56)

**References Cited**

## OTHER PUBLICATIONS

sion No. 1442516433. Catalog Name: Aurora Building Blocks 2. Order No. Catalog: 115.267.167. CAS Registry No. 1781876-60-5, Apr. 19, 2021 (1 page).

CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1-methyl-1H-pyrrolo[2,3-b]pyridin-4-ol. CHEMCATS Accession No. 0002254898. Catalog Name: FCH Group Reagents for Synthesis. Order No. Catalog: FCH1635008. CAS Registry No. 1781876-60-5, May 5, 2021 (1 page).

CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1H-Pyrrolo[2,3-B]Pyridin-4(7H)-One. CHEMCATS Accession No. 2022337458. Catalog Name: Chemieliva Pharmaceutical Product List. Order No. Catalog: CE0957308. CAS Registry No. 1076197-59-5, Jan. 28, 2021 (1 page).

CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1H-Pyrrolo[2,3-b]pyridin-4(7H)-one. CHEMCATS Accession No. 1621739382. Catalog Name: Ambeed, Inc. Product List. Order No. Catalog: A763560. CAS Registry No. 1076197-59-5, May 7, 2021 (1 page).

CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1H-Pyrrolo[2,3-b]pyridin-4-ol hydrate. CHEMCATS Accession No. 1773869211. Catalog Name: Aurora Building Blocks 3. Order No. Catalog: 129.194.895. CAS Registry No. 2031269-35-Apr. 7, 19, 2021 (1 page).

CHEMCATS. Chemical Abstracts Service: Columbus, OH. Chemical Name: 1H-pyrrolo[2,3-b]pyridin-4-ol hydrate. CHEMCATS Accession No. 0968477988. Catalog Name: ASW MedChem Product List. Order No. Catalog: TH-45275. CAS Registry No. 2031269-35-7, Jun. 5, 2020 (1 page).

\* cited by examiner




FIG. 1

DXD2202-003-03-TGA




FIG. 2




FIG. 3




FIG. 4





FIG. 5



2Theta (Coupled TwoTheta/Theta) WL-1.54060

FIG. 6






FIG. 7




FIG. 8




FIG. 9




FIG. 10

Sample: DXD2203-004-35  
Size: 5.1640 mg  
Method: DJP Double cycle

File: C:\TA\Data\TGA\DXD2203-004-35  
Operator: DJP  
Run Date: 04-Mar-2022 17:38  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 11

Sample: DXD2203-004-35  
Size: 5.1640 mg  
Method: DJP Double cycle

File: H:\03-Mar\DXD2203-004-35.001  
Operator: DJP  
Run Date: 04-Mar-2022 17:38  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 12







FIG. 13



2Theta (Coupled Two Theta/Theta) WL-1.54060

FIG. 14






FIG. 17




FIG. 18




FIG. 19






FIG. 20




FIG. 21




FIG. 22

DXD2203-007-08




FIG. 23

DXD2203-007-08




FIG. 24

DXD2203-007-08.421.fid




FIG. 25

Sample: DXD2203-007-07  
Size: 13.5480 mg  
Method: DJP Double cycle

File: C:\TA\Data\TGA\DXD2203-007-07  
Operator: DJP  
Run Date: 05-Mar-2022 15:51  
Instrument: TGA Q5000 V3.17 Build 265

TGA




FIG. 26





FIG. 27



2Theta (Coupled TwoTheta/Theta) WL-1.54060

FIG. 28




FIG. 29






FIG. 30








FIG. 33




FIG. 34




FIG. 35




FIG. 36




FIG. 37

DXD2203-009-09




FIG. 38




FIG. 39




FIG. 40




FIG. 41




FIG. 42






FIG. 43




FIG. 44




FIG. 45

DXD2203-009-20




FIG. 46




FIG. 47




FIG. 48




FIG. 49




FIG. 50




FIG. 51




FIG. 52

DXD2203-009-21




FIG. 53




FIG. 54




FIG. 55






FIG. 56




FIG. 57




FIG. 58




FIG. 59




FIG. 60




FIG. 61




FIG. 62




FIG. 63




FIG. 64




FIG. 65

DXD2203-010-03




FIG. 66






FIG. 67




FIG. 68




FIG. 69




FIG. 70




FIG. 71




FIG. 72




FIG. 73

Sample: DXD2203-010-04  
Size: 2.8470 mg  
Method: DJP Double cycle  
Comment: oxoglutarate

TGA

File: DXD2203-010-04  
Operator: DJP  
Run Date: 26-Feb-2022 20:19  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 74




FIG. 75




FIG. 76




FIG. 77




FIG. 78






FIG. 79

DXD2203-010-05.420.fid




FIG. 80




FIG. 81




FIG. 82




FIG. 83




FIG. 84




FIG. 85





FIG. 86



2Theta (Coupled TwoTheta/Theta) WL-1.54060

FIG. 87




FIG. 88







FIG. 89



2Theta (Coupled TwoTheta/Theta) WL-1.54060

FIG. 90






FIG. 91



DXD2203-010-08\_3d-40\_75  
DXD2203-010-08

FIG. 92



DXD2203-004-08\_53  
DXD2203-015-03

FIG. 93




FIG. 94




FIG. 95




FIG. 96






FIG. 97




FIG. 98




FIG. 99




FIG. 100




FIG. 101

Sample: DXD2203-013-03  
Size: 17.1710 mg  
Method: DJP Double cycle

TGA

File: DXD2203-013-03  
Operator: DJP  
Run Date: 20-Jul-2022 10:53  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 102




FIG. 103




FIG. 104




FIG. 105

Sample: DXD2203-014-01  
Size: 23.4050 mg  
Method: DJP Double cycle

TGA

File: DXD2203-014-01  
Operator: DJP  
Run Date: 21-Jul-2022 09:06  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 106

Sample: DXD2203-014-01  
Size: 23.4050 mg  
Method: DJP Double cycle

TGA

File: Q:\...107-Jul\DXD2203-014-01  
Operator: DJP  
Run Date: 21-Jul-2022 09:06  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 107





FIG. 108





2Theta (Coupled TwoTheta/Theta) WL-1.54060

FIG. 109

Sample: DXD2203-015-03  
Size: 25.1200 mg  
Method: DJP Double cycle

TGA

File: DXD2203-015-03  
Operator: DJP  
Run Date: 22-Jul-2022 07:04  
Instrument: TGA Q5000 V3.17 Build 265




FIG. 110

Universal V4.5A TA Instruments

Sample: DXD2203-015-03  
Size: 25.1200 mg  
Method: DJP Double cycle

File: Q:\...107-Jul\DXD2203-015-03.001  
Operator: DJP  
Run Date: 22-Jul-2022 07:04  
Instrument: TGA Q5000 V3.17 Build 265





FIG. 111



2Theta (Coupled TwoTheta/Theta) WL-1.54060

FIG. 112




FIG. 113




FIG. 114




FIG. 115

DR-2186-45-01




FIG. 116




FIG. 117




FIG. 118






FIG. 119




FIG. 120




FIG. 121




FIG. 122




FIG. 123




FIG. 124




FIG. 125




FIG. 126




FIG. 127




FIG. 128




FIG. 129




FIG. 130




FIG. 131




FIG. 132




FIG. 133




FIG. 134






FIG. 135




FIG. 136




FIG. 137




FIG. 138




FIG. 139




FIG. 140




FIG. 141




FIG. 142




FIG. 143




FIG. 144




FIG. 145




FIG. 146




FIG. 147




FIG. 148




FIG. 149




FIG. 150






FIG. 151




FIG. 152




FIG. 153




FIG. 154




FIG. 155




FIG. 156




FIG. 157




FIG. 158




FIG. 159




FIG. 160




FIG. 161

DJP2202-003-01




FIG. 162




FIG. 163

DJP2202006-01-TGA




FIG. 164




FIG. 165




FIG. 166






FIG. 167

DJP2202-006-03-TGA




FIG. 168




FIG. 169




FIG. 170




FIG. 171

Concentration vs Peak area




FIG. 172

1

**PHARMACEUTICALLY ACCEPTABLE  
SALTS AND COMPOSITIONS THEREOF**

CROSS-REFERENCE TO RELATED  
APPLICATIONS


This application claims priority to GB Application No. 2212116.4, filed on Aug. 19, 2022, which is incorporated by reference herein in its entirety.

FIELD OF THE INVENTION

This invention relates to pharmaceutically acceptable salts of 5-methoxy-N,N-dimethyltryptamine. In particular, though not exclusively, the invention relates to compositions/formulations and uses of the same as a medicament.

BACKGROUND OF THE INVENTION

5-methoxy-N,N-dimethyltryptamine (5-MeO-DMT) is a pharmacologically active compound of the tryptamine class and has the chemical formula:



5-MeO-DMT is a psychoactive/psychedelic substance found in nature and is believed to act mainly through serotonin receptors. It is also believed to have a high affinity for the 5-HT2 and 5-HT1A subtypes, and/or inhibits monoamine reuptake.

However, 5-MeO-DMT is not well understood and uses of this compound have not been well explored. Further, 5-MeO-DMT is not easy to handle, and there are challenges in formulating it for effective delivery in pharmaceutically useful compositions/formulations.

There remains a need in the art for improved compositions/formulations and uses of 5-MeO-DMT.

SUMMARY OF THE INVENTION

Herein disclosed is a non-hygroscopic salt of 5-MeO-DMT.

In an embodiment, the non-hygroscopic salt is 5-MeO-DMT hydrobromide.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT hydrobromide.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT hydrobromide, characterised by peaks in an XRPD diffractogram at 14.6, 16.8, 20.8, 24.3, 24.9 and 27.5° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT hydrobromide, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 109 or FIG. 171.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT hydrobromide, characterised by peaks in an

2

XRPD diffractogram at 14.6, 21.6 and 24.3° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT hydrobromide, characterised by peaks in an XRPD diffractogram at 18.6, 19.7 and 24.8° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT hydrobromide, characterised by peaks in an XRPD diffractogram at 14.6, 20.8, 21.6, 24.3 and 25.4° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT hydrobromide. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising crystalline 5-MeO-DMT hydrobromide. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT phosphate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT phosphate, characterised by peaks in an XRPD diffractogram at 12.9, 20.4 and 23.1° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT phosphate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 6.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT phosphate.

In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT phosphate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT fumarate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT fumarate, characterised by peaks in an XRPD diffractogram at 13.0, 16.3 and 22.1° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT fumarate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 14.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT fumarate.

In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT fumarate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT oxalate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT oxalate, characterised by peaks in an XRPD diffractogram at 13.0, 19.9 and 26.0° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT oxalate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 19 or FIG. 28.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT oxalate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT oxalate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT tartrate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT tartrate, characterised by peaks in an XRPD diffractogram at 18.3, 18.6, and 20.7°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT tartrate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 34 or FIG. 41.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT tartrate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT tartrate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT benzenesulfonate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT benzenesulfonate, characterised by peaks in an XRPD diffractogram at 9.5, 21.2, and 23.6°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT benzenesulfonate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 42.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT benzenesulfonate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT benzenesulfonate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT tosylate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT tosylate, characterised by peaks in an XRPD diffractogram at 19.3, 23.6 and 24.1°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT tosylate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 49 or 56.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT tosylate.

In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT tosylate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT glycolate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT glycolate, characterised by peaks in an XRPD diffractogram at 20.2, 21.1 and 23.4°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT glycolate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 62.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT glycolate.

In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT glycolate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT ketoglutarate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT ketoglutarate, characterised by peaks in an XRPD diffractogram at 14.4, 18.2 and 20.9°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT ketoglutarate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 69.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT ketoglutarate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT ketoglutarate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT malate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT malate, characterised by peaks in an XRPD diffractogram at 18.3, 18.7 and 18.9°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT malate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 76.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT malate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT malate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided 5-MeO-DMT saccharinate.

In an embodiment, there is provided a crystalline form of 5-MeO-DMT saccharinate, characterised by peaks in an XRPD diffractogram at 8.7, 15.2 and 20.9°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

## 5

In an embodiment, there is provided a crystalline form of 5-MeO-DMT saccharinate, characterised by peaks in an XRPD diffractogram as shown in, or substantially as shown in, FIG. 81.

In an embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT saccharinate. In an embodiment, the pharmaceutical composition is for use as a medicament.

In an embodiment, there is provided a pharmaceutical composition comprising a crystalline form of 5-MeO-DMT saccharinate. In an embodiment, the pharmaceutical composition is for use as a medicament.

Herein disclosed, there is provided a composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable salt of 5-methoxy-N,N-dimethyltryptamine (5-MeO-DMT).

In a first aspect of the invention, there is provided a composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable hydrobromide salt of 5-methoxy-N,N-dimethyltryptamine (5-MeO-DMT).

The invention provides for improved formulations and uses of 5-MeO-DMT salts.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.05 mg to 100 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.1 mg to 50 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.5 mg to 25 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.5 mg to 10 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 1 mg to 10 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 1 mg to 8 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 3 mg to 15 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.005 mg to 100 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.001 mg to 100 mg.

In an embodiment the composition comprises a dosage amount of 5-MeO-DMT in the range of 0.0005 mg to 100 mg.

The level of the active agent can be adjusted as required by need for example to suit a certain patient group (e.g. the elderly) or the conditions being treated.

In an embodiment the composition is formulated in a dosage form selected from: oral, transdermal, inhalable, intravenous, subcutaneous or rectal dosage form.

It is advantageous to be able to deliver the active agent in different forms, for example to suit a certain patient group (e.g. the elderly) or the conditions being treated.

In an embodiment the composition is formulated in a dosage form selected from: tablet, capsule, granules, powder, free-flowing powder, inhalable powder, aerosol, nebulised, vaping, buccal, sublingual, sublabial, injectable, or suppository dosage form.

In an embodiment the powder is suitable for administration by inhalation via a medicament dispenser selected from a reservoir dry powder inhaler, a unit-dose dry powder inhaler, a pre-metered multi-dose dry powder inhaler, a nasal inhaler or a pressurized metered dose inhaler.

In an embodiment the powder comprises particles, the particles having a median diameter of less than 2000  $\mu\text{m}$ , 1000  $\mu\text{m}$ , 500  $\mu\text{m}$ , 250  $\mu\text{m}$ , 100  $\mu\text{m}$ , 50  $\mu\text{m}$ , or 1  $\mu\text{m}$ .

## 6

In an embodiment the powder comprises particles, the particles having a median diameter of greater than 500  $\mu\text{m}$ , 250  $\mu\text{m}$ , 100  $\mu\text{m}$ , 50  $\mu\text{m}$ , 1  $\mu\text{m}$  or 0.5  $\mu\text{m}$ .

In an embodiment the powder comprises particles, and wherein the powder has a particle size distribution of  $d_{10}=20-60 \mu\text{m}$ , and/or  $d_{50}=80-120 \mu\text{m}$ , and/or  $d_{90}=130-300 \mu\text{m}$ .

The nature of the powder can be adjusted to suit need. For example, if being made for nasal inhalation, then the particles may be adjusted to be much finer than if the powder is going to be formulated into a gelatine capsule, or differently again if it is going to be compacted into a tablet.

In an embodiment the 5-MeO-DMT salt is amorphous or crystalline. In an embodiment, the 5-MeO-DMT salt is in a polymorphic crystalline form.

For the salt, the dosage amount is the equivalent amount of the free base delivered when the salt is taken. So 100 mg dosage amount of 5MeODMT corresponds to 117 mg of the hydrochloride salt (i.e. both providing the same molar amount of the active substance). The greater mass of the salt needed is due to the larger formula weight of the hydrogen chloride salt (i.e. 218.3 g/mol for the free base as compared to 254.8 g/mol for the salt). Similarly, for the deuterated or trituted version of 5MeODMT (also considered within the scope of the invention), a slight increase in mass can be expected due to the increased formula weight of these isotopic compounds.

Amorphous and crystalline substances often show different chemical/physical properties, e.g. improved rate of dissolution in a solvent, or improved thermal stability. Similarly, different polymorphs may also show different and useful chemical/physical properties.

In an embodiment the composition comprises one or more pharmaceutically acceptable carriers or excipients.

In an embodiment the composition comprises one or more of: mucoadhesive enhancer, penetrating enhancer, cationic polymers, cyclodextrins, Tight Junction Modulators, enzyme inhibitors, surfactants, chelators, and polysaccharides.

In an embodiment the composition comprises one or more of: chitosan, chitosan derivatives (such as N,N,N-trimethyl chitosan (TMC), n-propyl-(QuatPropyl), n-butyl-(QuatButyl) and n-hexyl (QuatHexyl)-N,N-dimethyl chitosan, chitosan chloride),  $\beta$ -cyclodextrin, *Clostridium perfringens* enterotoxin, zonula occludens toxin (ZOT), human neutrophil elastase inhibitor (ER143), sodium taurocholate, sodium deoxycholate sodium, sodium lauryl sulphate, glycodeoxycholat, palmitic acid, palmitoleic acid, stearic acid, oleyl acid, oleyl alcohol, capric acid sodium salt, DHA, EPA, dipalmitoyl phophatidyl choline, soybean lecithin, lysophosphatidylcholine, dodecyl maltoside, tetradecyl maltoside, EDTA, lactose, cellulose, and citric acid.

In an embodiment the composition disclosed herein is for use as a medicament. In an embodiment the composition disclosed herein is for use in a method of treatment of a human or animal subject by therapy.

In an embodiment the method of treatment is a method of treatment of:

- conditions caused by dysfunctions of the central nervous system,
- conditions caused by dysfunctions of the peripheral nervous system,
- conditions benefiting from sleep regulation (such as insomnia),
- conditions benefiting from analgesics (such as chronic pain),
- migraines,

trigeminal autonomic cephalgias (such as short-lasting unilateral neuralgiform headache with conjunctival injection and tearing (SUNCT), and short-lasting neuralgiform headaches with cranial autonomic symptoms (SUNA)),

conditions benefiting from neurogenesis (such as stroke, traumatic brain injury, Parkinson's dementia),

conditions benefiting from anti-inflammatory treatment, depression,

treatment resistant depression

anxiety,

substance use disorder,

addictive disorder,

gambling disorder,

eating disorders,

obsessive-compulsive disorders, or

body dysmorphic disorders,

optionally the condition is SUNCT and/or SUNA.

Treatment of the above conditions may be beneficially improved by taking the invention.

In an embodiment, the method of treatment is a method of treatment of alcohol-related diseases and disorders, eating disorders, impulse control disorders, nicotine-related disorders, tobacco-related disorders, methamphetamine-related disorders, amphetamine-related disorders, cannabis-related disorders, cocaine-related disorders, hallucinogen use disorders, inhalant-related disorders, benzodiazepine abuse or dependence related disorders, and/or opioid-related disorders.

In an embodiment, the method of treatment is a method of treatment of tobacco addiction. In an embodiment, the method is a method of reducing tobacco use. In an embodiment, the method of treatment is a method of treatment of nicotine addiction. In an embodiment, the method is a method of reducing nicotine use.

In an embodiment, the method of treatment is a method of treating alcohol abuse and/or addiction. In an embodiment, the method of treatment is a method of reducing alcohol use.

In an embodiment, the method of treatment is a method of treating or preventing heavy drug use.

In an embodiment, the method of treatment is a method of treating or preventing heavy drug use, including, but not limited to, alcohol, tobacco, nicotine, cocaine, methamphetamine, other stimulants, phencyclidine, other hallucinogens, marijuana, sedatives, tranquilizers, hypnotics, and opiates. It will be appreciated by one of ordinary skill in the art that heavy use or abuse of a substance does not necessarily mean the subject is dependent on the substance.

In an embodiment the method of treatment is a method of treatment of more than one of the above conditions, for example, the method of treatment may be a method of treatment of depression and anxiety.

In an embodiment the composition is administered one or more times a year.

In an embodiment the composition is administered one or more times a month.

In an embodiment the composition is administered one or more times a week.

In an embodiment the composition is administered one or more times a day.

In an embodiment the composition is administered at such a frequency as to avoid tachyphylaxis.

In an embodiment the composition is administered together with a complementary treatment and/or with a further active agent.

In an embodiment the further active agent is a psychedelic compound, optionally a tryptamine.

In an embodiment the further active agent is lysergic acid diethylamide (LSD), psilocybin, psilocin or a prodrug thereof.

In an embodiment the further active agent is an antidepressant compound.

In an embodiment the further active agent is selected from an SSRI, SNRI, TCA or other antidepressant compounds.

In an embodiment the further active agent is selected from Citalopram (Celexa, Cipramil), Escitalopram (Lexapro, Cipralex), Fluoxetine (Prozac, Sarafem), Fluvoxamine (Luvox, Faverin), Paroxetine (Paxil, Seroxat), Sertraline (Zoloft, Lustral), Desvenlafaxine (Pristiq), Duloxetine (Cymbalta), Levomilnacipran (Fetzima), Milnacipran (Ixel, Savella), Venlafaxine (Effexor), Vilazodone (Viibryd), Vortioxetine (Trintellix), Nefazodone (Dutonin, Nefadar, Serzone), Trazodone (Desyrel), Reboxetine (Edronax), Teniloxazine (Luceclan, Metatone), Viloxazine (Vivalan), Bupropion (Wellbutrin), Amitriptyline (Elavil, Endep), Amitriptylinoxide (Amioxid, Ambivalon, Equilibrin), Clomipramine (Anafranil), Desipramine (Norpramin, Pertofrane), Dibenzepin (Noveril, Victoril), Dimetacrine (Istonil), Dosulepin (Prothiaden), Doxepin (Adapin, Sinequan), Imipramine (Tofranil), Lofepramine (Lomont, Gamanil), Melitracen (Dixeran, Melixeran, Trausabun), Nitroxazepine (Sintamil), Nortriptyline (Pamelor, Aventyl), Noxiptiline (Agedal, Elronon, Nogedal), Opipramol (Insidon), Pipofezine (Azafen/Azaphen), Protriptyline (Vivactil), Trimipramine (Surmontil), Amoxapine (Asendin), Maprotiline (Ludiomil), Mianserin (Tolvon), Mirtazapine (Remeron), Setiptiline (Tecipul), Isocarboxazid (Marplan), Phenelzine (Nardil), Tranylcypromine (Parnate), Selegiline (Eldepryl, Zelapar, Emsam), Caroxazone (Surodil, Timostenil), Metralindole (Inkazan), Moclobemide (Aurorix, Manerix), Pirlindole (Pirazidol), Toloxatone (Humoryl), Agomelatine (Valdoxan), Esketamine (Spravato), Ketamine (Ketalar), Tandospiro (Sediell), Tianeptine (Stablon, Coaxil), Amisulpride (Solian), Aripiprazole (Abilify), Brexpiprazole (Rexulti), Lurasidone (Latuda), Olanzapine (Zyprexa), Quetiapine (Seroquel), Risperidone (Risperdal), Trifluoperazine (Stelazine), Buspirone (Buspar), Lithium (Eskalith, Lithobid), Modafinil (Provigil), Thyroxine (T4), Triiodothyronine (T3).

In an embodiment the further active agent is selected from Celexa (citalopram), Cymbalta (duloxetine), Effexor (venlafaxine), Lexapro (escitalopram), Luvox (fluvoxamine), Paxil (paroxetine), Prozac (fluoxetine), Remeron (mirtazapine), Savella (milnacipran), Trintellix (vortioxetine), Vestra (reboxetine), Viibryd (vilazodone), Wellbutrin (bupropion), Zoloft (sertraline).

In an embodiment the complementary treatment is psychotherapy.

In an embodiment, there is provided a composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable salt of 5MeODMT for use in a method of treatment of treatment resistant depression.

In an embodiment, there is provided a composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable salt of 5MeODMT for use in a method of treatment of depression.

In an embodiment, there is provided a composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable salt of 5MeODMT for use in a method of treatment of PTSD.

In an embodiment, there is provided a composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable salt of 5MeODMT for use in a method of treatment of addiction/substance misuse disorders.

In an embodiment, there is provided a nasal inhalation composition comprising a pharmaceutically effective amount of a pharmaceutically acceptable salt of 5MeODMT for use in a method of treatment of treatment resistant depression.

Treatment of the above conditions may be beneficially improved by taking the invention together with some complementary treatments; also these treatments may occur much less regularly than some other treatments that require daily treatments or even multiple treatments a day.

For the avoidance of doubt, the person skilled in the art will appreciate that numerical values relating to measurements are subject to instrument setup and measurement errors which can result in small discrepancies in the measurement values obtained. As such, it will be readily understood that where herein, for example, an XRPD peak is given a value of  $XX.Y^{\circ}2\theta$ , included with the scope of the disclosure of this application are values  $XX.Y^{\circ}2\theta \pm 0.1^{\circ}2\theta$ ,  $XX.Y^{\circ}2\theta \pm 0.2^{\circ}2\theta$ ,  $XX.Y^{\circ}2\theta \pm 0.3^{\circ}2\theta$ ,  $XX.Y^{\circ}2\theta \pm 0.4^{\circ}2\theta$  and  $XX.Y^{\circ}2\theta \pm 0.5^{\circ}2\theta$ . The skilled person will doubtless understand that the same applies for numerical values given for temperatures and enthalpies (joules). Again, solely as an example, a temperature value of  $XX.Y^{\circ}C.$  (or  $XX^{\circ}C.$ ) will be understood to encompass values of  $XX.Y^{\circ}C. \pm 0.1^{\circ}C.$ ,  $XX.Y^{\circ}C. \pm 0.2^{\circ}C.$ ,  $XX.Y^{\circ}C. \pm 0.3^{\circ}C.$ ,  $XX.Y^{\circ}C. \pm 0.4^{\circ}C.$  and  $XX.Y^{\circ}C. \pm 0.5^{\circ}C.$

Similarly, it should be understood that values measured herein may be rounded down and that these rounded values are within the scope of the original disclosure. For example, values measured at 2 decimal places herein may be expressed at 1 decimal place (with the appropriate rounding) and so are still within the original disclosure.

Terms such as “a”, “an,” and “the” are not intended to refer to only a singular entity but include the general class of which a specific example may be used for illustration.

As used herein, the terms “about” and/or “around” refer to a value that is within 10% above or below the value being described.

The different polymorphic forms of the various salts described herein have been labelled sequentially as Pattern 1, Pattern 2 etc., principally numbering these patterns in the order in which they appear in the application. For the sake of completeness, we add that no inference should be taken from the ordering of the polymorphs using this numbering system.

#### BRIEF DESCRIPTION OF THE FIGURES

FIG. 1. XRPD Diffractogram of free base, Batch: DXD2203-003-01.

FIG. 2. TGA Thermogram of free base, Batch: DXD2203-003-01.

FIG. 3. DSC Thermogram (1st heat) of free base, Batch: DXD2203-003-01.

FIG. 4. DSC Thermograms of free base, Batch: DXD2203-003-01, cooling (blue trace), 2nd heat (green trace).

FIG. 5. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of free base, Batch: DXD2203-003-01.

FIG. 6. XRPD Diffractograms of Phosphate salt isolated from IPA (blue trace, top), THF (black trace, middle) and Ethyl Acetate (red trace, bottom).

FIG. 7. TGA Thermogram of Phosphate salt, Batch: DXD2203-04-35.

FIG. 8. Heat-cool-reheat DSC thermogram of Phosphate salt, 1st heating (blue trace), cooling ramp (green trace) and 2nd heating (red trace), Batch: DXD2203-004-35.

FIG. 9. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Phosphate salt, Batch: DXD2203-004-35

FIG. 10. <sup>31</sup>P NMR (d<sub>6</sub>-DMSO) Spectrum of Phosphate salt, Batch: DXD2203-004-35

FIG. 11. DVS Isotherm plot of Phosphate salt, Batch: DXD2203-004-35.

FIG. 12. DVS Kinetic plot of Phosphate salt, Batch: DXD2203-004-35.

FIG. 13. XRPD Diffractograms of Phosphate salt, Batch: DXD2203-004-35 (red trace, top) and post-DVS (black trace, bottom).

FIG. 14. XRPD Diffractograms of Fumarate salt isolated from Ethyl Acetate (red trace, top), Acetone (black trace, middle) and IPA (blue trace, bottom).

FIG. 15. TGA Thermogram of Fumarate salt, Batch: DXD2203-004-53.

FIG. 16. DSC Thermogram (1st heating) of Fumarate salt, Batch: DXD2203-004-53.

FIG. 17. DSC Thermograms of Fumarate salt, cooling (green trace) and 2nd heating (blue trace).

FIG. 18. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Fumarate salt, Batch: DXD2203-004-53.

FIG. 19. XRPD Diffractograms of Oxalate salt isolated from (from top to bottom of the XRPD) acetone (pink trace), ethyl acetate (brown trace), acetonitrile (black trace), THF (red trace), IPA (blue trace) and 5% water: ethanol (green trace).

FIG. 20. TGA Thermogram of Oxalate salt, Batch: DXD2203-007-08.

FIG. 21. DSC Thermogram (1st heating) of Oxalate salt, Batch: DXD2203-007-08.

FIG. 22. DSC Thermograms of Oxalate salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 23. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Oxalate salt, Batch: DXD2203-007-08.

FIG. 24. <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) Spectrum of Oxalate salt, Batch: DXD2203-007-08.

FIG. 25. Quantitative <sup>13</sup>C NMR (d<sub>6</sub>-DMSO) Spectrum of Oxalate salt, Batch: DXD2203-007-08.

FIG. 26. DVS Isotherm plot of Oxalate salt, Batch: DXD2203-007-07.

FIG. 27. Sorption kinetic plot of Oxalate salt, Batch: DXD2203-007-07.

FIG. 28. XRPD Diffractograms of Oxalate salt, Batch: DXD2203-007-07 (red trace, bottom) and post-DVS (black trace, top).

FIG. 29. XRPD Diffractograms of Batch: DXD2203-007-44 isolated from ethyl acetate (black trace, top), free base (blue trace, middle) and adipic acid (red trace, bottom).

FIG. 30. TGA Thermogram of Batch: DXD2203-007-44.

FIG. 31. DSC Thermogram (1st heating) of Batch: DXD2203-007-44.

FIG. 32. DSC Thermograms of Batch: DXD2203-007-44, cooling (blue trace) and 2nd heating (green trace).

FIG. 33. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Batch: DXD2203-007-44.

FIG. 34. XRPD Diffractograms (from top to bottom) of Tartrate salt isolated from IPA/hexane (green trace), THF/hexane (blue trace), THF at 40° C. (black trace) and ethanol at 40° C. (red trace).

FIG. 35. TGA Thermogram of Tartrate salt, Batch: DXD2203-009-09.

FIG. 36. DSC Thermogram (1st heating) of Tartrate salt, Batch: DXD2203-009-09.

FIG. 37. DSC Thermograms of Tartrate salt, cooling (blue trace) and 2nd heating (green trace).



## 11

FIG. 38. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Tartrate salt, Batch: DXD2203-009-09.

FIG. 39 shows the DVS isotherm plot for Tartrate salt.

FIG. 40. Sorption kinetic plot of Tartrate salt, Batch: DXD2203-009-08.

FIG. 41. XRPD Diffractograms of Tartrate salt, Batch: DXD2203-009-08 (red trace) post DVS (black trace).

FIG. 42. XRPD Diffractograms of Benzenesulfonate salt isolated from IPA/hexane (red trace), Benzenesulfonic acid (black trace) and Free Base (blue trace).

FIG. 43. TGA Thermogram of Benzenesulfonate salt, Batch: DXD2203-009-20.

FIG. 44. DSC Thermogram (1st heating) of Benzenesulfonate salt, Batch: DXD2203-009-20.

FIG. 45. DSC Thermograms of Benzenesulfonate salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 46. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Benzenesulfonate salt, Batch: DXD2203-009-20.

FIG. 47. DVS Isotherm plot of Benzenesulfonate salt, Batch: DXD2203-009-20.

FIG. 48. Sorption kinetic plot of Benzenesulfonate salt, Batch: DXD2203-009-20.

FIG. 49. XRPD Diffractograms of Tosylate salt isolated from IPA/Hexane (red trace, top), p-toluene sulfonic acid (black trace, middle) and Free Base (blue trace, bottom).

FIG. 50. TGA Thermogram of Tosylate salt, Batch: DXD2203-009-21.

FIG. 51. DSC Thermogram (1st heating) of Tosylate salt, Batch: DXD2203-009-21.

FIG. 52. DSC Thermograms of Tosylate salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 53. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Tosylate salt, Batch: DXD2203-009-21.

FIG. 54. DVS Isotherm plot of Tosylate salt, Batch: DXD2203-009-21.

FIG. 55. Sorption kinetic plot of Tosylate salt, Batch: DXD2203-009-21.

FIG. 56. XRPD Diffractograms of Tosylate salt, Batch: DXD2203-009-21 (red trace) and post DVS (black trace).

FIG. 57. XRPD Diffractograms (from top to bottom) of Hydrobromide salt isolated from acetonitrile/toluene (black trace), DMF/toluene (red trace), methanol/MTBE (green trace) and Free Base (blue trace).

FIG. 58. TGA Thermogram of Hydrobromide salt, Batch: DXD2203-010-02.

FIG. 59. DSC Thermogram (1st heating) of Hydrobromide salt, Batch: DXD2203-010-02.

FIG. 60. DSC Thermograms of Hydrobromide salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 61. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Hydrobromide salt, Batch: DXD2203-010-02.

FIG. 62. XRPD Diffractograms of glycolate salt isolated from IPAC-40° C. (red trace, top), Glycolic acid (black trace, middle) and Free Base (blue trace, bottom).

FIG. 63. TGA Thermogram of glycolate salt, Batch: DXD2203-010-03.

FIG. 64. DSC Thermogram (1st heating) of glycolate salt, Batch: DXD2203-010-03.

FIG. 65. DSC Thermograms of glycolate salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 66. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of glycolate salt, Batch: DXD2203-010-03.

FIG. 67. DVS Isotherm plot of glycolate salt, Batch: DXD2203-010-03.

FIG. 68. DVS Kinetic plot of glycolate salt, Batch: DXD2203-010-03.

## 12

FIG. 69. XRPD Diffractograms (from top to bottom) of Ketoglutarate salt isolated from methanol/MTBE (red trace), ethanol/MTBE (green trace), Ketoglutaric acid (black trace) and Free Base (blue trace).

FIG. 70. TGA Thermogram of Ketoglutarate salt, Batch: DXD2203-010-04.

FIG. 71. DSC Thermogram (1st heating) of Ketoglutarate salt, Batch: DXD2203-010-04.

FIG. 72. DSC Thermograms of Ketoglutarate salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 73. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Ketoglutarate salt, Batch: DXD2203-010-04.

FIG. 74. DVS Isotherm plot of Ketoglutarate salt, Batch: DXD2203-010-04.

FIG. 75. Sorption kinetic plot of Ketoglutarate salt, Batch: DXD2203-010-04.

FIG. 76. XRPD Diffractograms (from top to bottom) of Malate salt isolated from ethanol/MTBE (red trace), IPAC/MTBE (green trace), L-Malic acid (black trace) and Free Base (blue trace).

FIG. 77. TGA Thermogram of Malate salt, Batch: DXD2203-010-05.

FIG. 78. DSC Thermogram (1st heating) of Malate salt, Batch: DXD2203-010-05.

FIG. 79. DSC Thermograms of Malate salt, cooling (blue trace) and 2nd heating (green trace).

FIG. 80. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Malate salt, Batch: DXD2203-015-05.

FIG. 81. XRPD Diffractograms of Saccharinate salt (red trace, top), Saccharine (black trace, middle) and Free Base (blue trace, bottom).

FIG. 82. TGA Thermogram of Saccharinate salt, Batch: DXD2203-010-01.

FIG. 83. DSC Thermogram (1st heating) of Saccharinate salt, Batch: DXD2203-010-01.

FIG. 84. DSC Thermograms of Saccharinate salt, cooling (green trace) and 2nd heating (blue trace).

FIG. 85. <sup>1</sup>H NMR (d<sub>6</sub>-DMSO) Spectrum of Saccharinate salt, Batch: DXD2203-010-01.

FIG. 86. XRPD Diffractograms of Phosphate salt, initial (red trace, bottom) and after 3 days at 40° C./75% RH (black trace, top).

FIG. 87. XRPD Diffractograms of Fumarate salt, initial (red trace, bottom) and after 3 days at 40° C./75% RH (black trace, top).

FIG. 88. XRPD Diffractograms of Fumarate salt, initial (red trace), after 3 days at 40° C./75% RH (black trace), Fumaric acid (green trace) and free base (blue trace).

FIG. 89. XRPD Diffractograms of Tartrate salt, initial (red trace) and after 3 days at 40° C./75% RH (black trace).

FIG. 90. XRPD Diffractograms of Tosylate salt, initial (red trace) and after 3 days at 40° C./75% RH (black trace).

FIG. 91. XRPD Diffractograms of Saccharinate salt, initial (red trace) and after 3 days at 40° C./75% RH (black trace).

FIG. 92. XRPD Diffractograms of Hydrobromide salt, initial (red trace) and after 3 days at 40° C./75% RH (black trace).

FIG. 93. XRPD Diffractograms of Fumarate salt produced during scale-up versus that initially analysed during the salt screen.

FIG. 94. XRPD Diffractograms of Oxalate salt produced during scale-up versus that initially analysed during the salt screen.

FIG. 95. XRPD Diffractograms of Hydrobromide salt produced during scale-up versus that initially analysed during the salt screen.

## 13

FIG. 96. TGA Thermogram of Hydrobromide salt produced during scale-up.

FIG. 97. DSC Thermogram (1st heating) of Hydrobromide salt produced during scale-up.

FIG. 98. DSC Thermogram (cooling) of Hydrobromide salt produced during scale-up.

FIG. 99. DSC Thermograms of Hydrobromide salt produced during scale-up (top) versus that initially analysed during the salt screen (bottom).

FIG. 100.  $^1\text{H}$  NMR spectrum of Hydrobromide salt produced during scale-up.

FIG. 101.  $^{13}\text{C}$  NMR spectrum of Oxalate salt produced during scale-up.

FIG. 102. DVS Isotherm plot of Oxalate salt produced during scale-up.

FIG. 103. Sorption kinetic plot of Oxalate salt produced during scale-up.

FIG. 104. XRPD Diffractograms of Oxalate salt produced during scale-up, (black trace, top) and post-DVS (red trace, bottom).

FIG. 105. XRPD Diffractograms of Oxalate salt produced during scale-up, (black trace, top), post-DVS (red trace, middle) and post storage (bottom) at  $40^\circ\text{C}/75\%$  RH for 1 week.

FIG. 106. DVS Isotherm plot of Hydrobromide salt produced during scale-up.

FIG. 107. Sorption kinetic plot of Hydrobromide salt produced during scale-up.

FIG. 108. XRPD Diffractograms of Hydrobromide salt produced during scale-up, (black trace, top) and post-DVS (red trace, bottom).

FIG. 109. XRPD Diffractograms of Hydrobromide salt produced during scale-up, (blue trace, top), post-DVS (red trace, middle) and post storage (bottom) at  $40^\circ\text{C}/75\%$  RH for 1 week.

FIG. 110. DVS Isotherm plot of Fumarate salt produced during scale-up.

FIG. 111. Sorption kinetic plot of Fumarate salt produced during scale-up.

FIG. 112. XRPD Diffractograms of Fumarate salt produced during scale-up, (black trace, top) and post-DVS (red trace, bottom).

FIG. 113. TGA Thermogram of Fumarate salt produced during scale-up post-DVS.

FIG. 114. DSC Thermogram of Fumarate salt produced during scale-up post-DVS.

FIG. 115. XRPD Diffractograms of Fumarate salt produced during scale-up, (blue trace, top), post-DVS (red trace, middle) and post storage (bottom) at  $40^\circ\text{C}/75\%$  RH for 1 week.

FIG. 116. XRPD Diffractograms of Phosphate salt.

FIG. 117. XRPD Diffractograms of Phosphate salt.

FIG. 118. XRPD Diffractograms of various samples of Phosphate salt.

FIG. 119. XRPD Diffractograms of various samples of Tartrate salt.

FIG. 120. XRPD Diffractograms of various samples of Tartrate salt (low intensity samples).

FIG. 121.  $^1\text{H}$ -NMR Spectrum of 5-MeO-DMT Phosphate salt Pattern 1.

FIG. 122. XRPD Diffractogram of 5-MeO-DMT Phosphate Pattern 1.

FIG. 123. Thermal analysis (TGA and DSC) of 5-MeO-DMT Phosphate Pattern 1.

FIG. 124. DVS Isotherm plot of 5-MeO-DMT Phosphate Pattern 1.

## 14

FIG. 125. Sorption kinetic plot of 5-MeO-DMT Phosphate Pattern 1.

FIG. 126. XRPD Diffractogram comparison of 5-MeO-DMT Phosphate pre- and post-DVS showing the form remains Pattern 1.

FIG. 127. HPLC chromatogram and purity analysis of 5-MeO-DMT Phosphate Pattern 1.

FIG. 128. XRPD diffractogram overlay of 5-MeO-DMT Phosphate Pattern 1 before and after storage at  $25^\circ\text{C}/97\%$  RH and  $40^\circ\text{C}/75\%$  RH for 7 days.

FIG. 129. HPLC chromatogram and purity analysis of 5-MeO-DMT Phosphate Pattern 1 before and after storage at  $40^\circ\text{C}/75\%$  RH for 7 days.

FIG. 130. HPLC chromatogram and purity analysis of 5-MeO-DMT Phosphate Pattern 1 before and after storage at  $25^\circ\text{C}/97\%$  RH for 7 days.

FIG. 131. XRPD diffractogram of 5-MeO-DMT Tartrate Pattern 1.

FIG. 132. XRPD diffractogram overlay of 5-MeO-DMT Tartrate Pattern 1.

FIG. 133.  $^1\text{H}$ -NMR Spectrum of 5-MeO-DMT Tartrate salt Pattern 1.

FIG. 134. Thermal analysis (TGA and DSC) of 5-MeO-DMT Tartrate Pattern 1.

FIG. 135. DVS Isotherm plot of 5-MeO-DMT Tartrate Pattern 1.

FIG. 136. Sorption kinetic plot of 5-MeO-DMT Tartrate Pattern 1.

FIG. 137. XRPD diffractogram overlay of 5-MeO-DMT Tartrate Pattern 1 pre- and post-DVS showing the form remains Pattern 1.

FIG. 138. HPLC chromatogram and purity analysis of 5-MeO-DMT Tartrate Pattern 1.

FIG. 139. XRPD diffractogram overlay of 5-MeO-DMT Tartrate Pattern 1 before and after storage at  $25^\circ\text{C}/97\%$  RH and  $40^\circ\text{C}/75\%$  RH for 7 days.

FIG. 140. HPLC chromatogram and purity analysis of 5-MeO-DMT Tartrate Pattern 1 before and after storage at  $40^\circ\text{C}/75\%$  RH for 7 days.

FIG. 141. HPLC chromatogram and purity analysis of 5-MeO-DMT Tartrate Pattern 1 before and after storage at  $25^\circ\text{C}/97\%$  RH for 7 days.

FIG. 142. XRPD Diffractogram of hydrochloride salt lot RPI-014-022.

FIG. 143. TGA Thermogram of hydrochloride salt lot RPI-014-022.

FIG. 144. DSC Thermogram of first heat cycle of hydrochloride salt lot RPI-014-022.

FIG. 145. DSC Thermogram of cool and reheat cycles of hydrochloride salt lot RPI-014-022.

FIG. 146. DSC Thermograms for reheating of hydrochloride lot RPI-014-022 at different heating rates.

FIG. 147.  $^1\text{H}$  NMR Spectrum of hydrochloride salt lot RPI-014-022.

FIG. 148.  $^1\text{H}$ - $^{13}\text{C}$  HSQC Spectrum of hydrochloride salt lot RPI-014-022.

FIG. 149. XRPD Diffractogram of lyophilised hydrochloride salt (red) vs supplied pattern 1 (black).

FIG. 150. XRPD Diffractogram of pattern 2 lot DJP2202-007-01 from dioxane (red) compared to pattern 1.

FIG. 151. TGA Thermogram of Hydrochloride pattern 2 lot DJP2202-007-01.

FIG. 152. DSC Thermogram of Hydrochloride pattern 2 lot DJP2202-007-01.

FIG. 153.  $^1\text{H}$  NMR Spectrum of Hydrochloride pattern 2 lot DJP2202-007-01.

## 15

FIG. 154. XRPD Diffractogram of Hydrochloride pattern 3 (blue) and pattern 2 (red) and pattern 1 (black).

FIG. 155. TGA Thermogram of Hydrochloride pattern 3 lot DJP2202-007-14.

FIG. 156. DSC Thermogram of Hydrochloride pattern 3 lot DJP2202-007-14.

FIG. 157. <sup>1</sup>H NMR Spectrum of hydrochloride pattern 3 lot DJP2202-007-14.

FIG. 158. XRPD Diffractogram of pattern 1 resulting from evaporation of 5-MeO-DMT HCl/solvent combinations: HCl/MEK (green, top), HCl/IPA (blue, immediately below green), HCl/EtOH (red, immediately below blue) and HCl/1-PrOH (black, bottom).

FIG. 159. XRPD Diffractogram of benzoate salt pattern 2 (red) vs supplied pattern 1 (black).

FIG. 160. TGA Thermogram of benzoate salt pattern 2 lot DJP2202-003-01.

FIG. 161. DSC Thermogram of benzoate salt pattern 2 lot DJP2202-003-01.

FIG. 162. <sup>1</sup>H NMR Spectrum of benzoate salt pattern 2 lot DJP2202-003-01.

FIG. 163. XRPD Diffractogram of benzoate pattern 3 (blue) compared to pattern 2 (red) and pattern 1 (black).

FIG. 164. TGA Thermogram of benzoate salt pattern 3 lot DJP2202-006-01.

FIG. 165. DSC Thermogram of benzoate salt pattern 3 lot DJP2202-006-01.

FIG. 166. <sup>1</sup>H NMR Spectrum of benzoate salt pattern 3 lot DJP2202-006-01.

FIG. 167. XRPD Diffractogram of benzoate salt pattern 4 (green) compared to pattern 3 (blue), pattern 2 (red) and pattern 1 (black).

FIG. 168. TGA Thermogram of benzoate salt pattern 4 lot DJP2202-006-03.

FIG. 169. DSC Thermogram of benzoate salt pattern 4 lot DJP2202-006-03.

FIG. 170. <sup>1</sup>H NMR Spectrum of benzoate salt pattern 4 lot DJP2202-006-03.

FIG. 171. XRPD Diffractogram of hydrobromide salt pattern 2.

FIG. 172. Calibration curve of free base.

## DESCRIPTION OF THE INVENTION

An object of the present invention is to provide 5-MeO-DMT salts. Moreover, another object of the present invention is to provide 5-MeO-DMT salts which neither easily convert into hydrates, even when a pharmaceutical composition comprising a 5-MeO-DMT salt is stored for a long period of time. Hygroscopicity is the phenomenon of attracting and holding water molecules via either adsorption or absorption from the surrounding environment. Pharmaceuticals that pick up less than 0.2% moisture at 80% RH are considered non hygroscopic. Pharmaceuticals that pick up between 0.2% and 2.0% moisture at 80% RH are considered slightly hygroscopic. Pharmaceuticals that pick up between 2.0% and 15.0% moisture at 80% RH are considered moderately hygroscopic. Pharmaceuticals that pick up more than 15.0% moisture at 80% RH are considered very hygroscopic. Hygroscopic substances are difficult to handle and costly and burdensome measures must be taken in order to ensure they are not exposed to moisture during process and formulation. Exposed to moisture, hygroscopic substances can take on water and convert to a hydrous form. This presents several disadvantages. First, the hydrous forms may have the disadvantage of being less bioavailable and less dissoluble than the anhydrous forms. Second, the variation

## 16

in the amount of hydrous versus anhydrous substance from batch to batch could fail to meet specifications set by drug regulatory agencies. Third, processes like milling may cause the drug substance to adhere to manufacturing equipment which may further result in processing delay, increased operator involvement, increased cost, increased maintenance and lower production yield. Fourth, in addition to problems caused by introduction of moisture during the processing of these hygroscopic substances, the potential for absorbance of moisture during storage and handling would adversely affect the dissolubility of the drug substance. Thus shelf-life of the product could be significantly decreased and/or packaging costs could be significantly increased.

The inventors have surprisingly discovered that 5-MeO-DMT hydrobromide is a non-hygroscopic salt of 5-MeO-DMT. The tartrate salt of 5-MeO-DMT is moderately hygroscopic, the tosylate salt and the phosphate salt are both slightly hygroscopic.

The inventors have further surprisingly discovered that 5-MeO-DMT hydrobromide, whilst being non-hygroscopic, has high solubility compared to other moderately hygroscopic salts of 5-MeO-DMT for example the benzoate or oxalate salts. The non-hygroscopic, highly soluble HBr salt of 5-MeO-DMT therefore affords the advantage of removing the need for costly and burdensome processing measures, for example the need for low humidity manufacturing environment. The high solubility of the HBr salt of 5-MeO-DMT also facilitates the use of simplified solid formulations without the need for costly solubility enhancement techniques.

The inventors have further surprisingly discovered multiple polymorphic forms of crystalline 5-MeO-DMT hydrobromide, including a form referred to as form/pattern 2 with desirable qualities. The XRPD for this crystalline form can be seen in FIG. 171 and the peaks are tabulated in Tables 21, 21a and 21b.

## Example 1: Salt Screen

5-methoxy-N,N-dimethyltryptamine (5-MeO-DMT) was supplied as a HCl salt. In order to preform salt screening experiments, the HCl salt was converted into free base. The crystalline nature of isolated free base was confirmed by XRPD and further analysed by TGA, DSC, <sup>1</sup>H NMR analyses.

Salt screening studies with 24 selected counter-ions were performed to determine if 5-MeO-DMT free base is amenable to salt formation. On completion of this study, 11 crystalline salts were successfully generated and are displayed in Table 1. Salts were examined by XRPD, TGA, DSC, <sup>1</sup>H NMR and some by DVS analyses. Short term physical stability was examined by storage of salts at 40° C./75% RH for three days.

TABLE 1

Produced crystalline salts
Salt
Phosphate
Fumarate
Oxalate
Tartrate
Benzenesulfonate
Tosylate
Hydrobromide
Glycolate
Ketoglutarate

17

TABLE 1-continued

Produced crystalline salts Salt
Saccharinate Malate

## Instruments

## X-Ray Powder Diffraction (XRPD)

XRPD diffractograms were acquired using Bruker D2 Phaser diffractometer equipped with LYNXEYE detector. Samples were prepared using a zero-background sample holder. The samples were scanned from 5 to 32° (2θ) using a step size of 0.02° and a time per step of 0.13 second whilst spinning the sample. Diffractograms were plotted using the EVA program from Bruker.

## Thermo—Gravimetric Analysis (TGA)

TGA thermograms were obtained with a TA Instrument Discovery 550 in Al pans. The heating rate used was 10° C./min linear ramp from 25 to 400° C. with a nitrogen purging at a rate of 60 ml/min. TGA thermograms were analysed using TRIOS software.

## Differential Scanning Calorimetry (DSC)

DSC analyses were performed on a TA Instrument DSC250 with a Tzero cell purged at constant flow rate of 50 ml min<sup>-1</sup> with dry nitrogen and a refrigerated cooling system RCS90. The instrument was calibrated using Indium as a standard. A small quantity of the samples was weighed into TA Tzero Aluminium pan with pierced lid. Samples were heating at 10° C./min in heat-cool-reheat method. TRIOS software was used to analyse DSC scans.

## Nuclear Magnetic Resonance Spectroscopy (NMR)

The 1H NMR spectra were measured on Bruker NEO spectrometer operating at 400.13 MHz for protons. Samples were dissolved in d6-DMSO. Data were processed using MestReNovax64 software.

## Dynamic Vapor Sorption (DVS)

DVS analyses were performed on TA Instrument DVS Q5000. Samples were added to a pre-tared metallised quartz crucible and run at 25° C. from 40% RH to 90% RH, down to zero and back to 40%. This cycle was repeated in increments of 10% RH.

## Preparation and Initial Characterisation of Free Base

5-MeO-DMT Hydrochloride salt (2×5 g) was used for preparation of free base.

Free base was isolated from NaHCO<sub>3</sub>-Ethyl acetate extraction (5.7 g, 67% yield).

5-MeO-DMT free base was characterised by XRPD, TGA, DSC and <sup>1</sup>H NMR.

## X-Ray Powder Diffraction (XRPD)

XRPD diffractogram in FIG. 1 displayed crystalline peaks confirming the crystallinity of the free base. This was nominated as free base pattern 1. The XRPD peak data is shown in Table 2.

TABLE 2

XRPD Peak data for free base pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	5.464°	16.160	0.010
2	7.588°	11.641	0.006
3	9.294°	9.507	0.008
4	10.450°	8.459	0.037
5	10.867°	8.135	0.201

18

TABLE 2-continued

XRPD Peak data for free base pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
6	11.867°	7.451	0.002
7	12.779°	6.922	0.009
8	13.217°	6.693	0.083
9	15.062°	5.878	0.011
10	15.833°	5.593	0.019
11	16.251°	5.450	0.041
12	16.644°	5.322	0.069
13	17.192°	5.154	0.070
14	17.933°	4.942	0.027
15	18.478°	4.798	0.080
16	19.480°	4.553	0.017
17	19.802°	4.480	0.020
18	20.402°	4.349	0.036
19	20.814°	4.264	0.051
20	21.106°	4.206	0.071
21	21.652°	4.101	1.000
22	21.971°	4.042	0.141
23	22.976°	3.868	0.019
24	23.686°	3.753	0.008
25	24.958°	3.565	0.030
26	25.340°	3.512	0.015
27	25.705°	3.463	0.013
28	26.111°	3.410	0.101
29	26.764°	3.328	0.058
30	27.082°	3.290	0.258
31	28.120°	3.171	0.006
32	28.439°	3.136	0.005
33	29.176°	3.058	0.005
34	29.811°	2.995	0.010
35	30.329°	2.945	0.005
36	31.067°	2.876	0.005

TABLE 2a

XRPD Peak data for free base pattern 1 (2 d.p.).			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	5.46°	16.16	0.01
2	7.59°	11.64	0.01
3	9.29°	9.51	0.01
4	10.45°	8.46	0.04
5	10.87°	8.14	0.20
6	11.87°	7.45	0.00
7	12.78°	6.92	0.01
8	13.22°	6.69	0.08
9	15.06°	5.88	0.01
10	15.83°	5.59	0.02
11	16.25°	5.45	0.04
12	16.64°	5.32	0.07
13	17.19°	5.15	0.07
14	17.93°	4.94	0.03
15	18.48°	4.80	0.08
16	19.48°	4.55	0.02
17	19.8°	4.48	0.02
18	20.4°	4.35	0.04
19	20.81°	4.26	0.05
20	21.11°	4.21	0.07
21	21.65°	4.10	1.00
22	21.97°	4.04	0.14
23	22.98°	3.87	0.02
24	23.69°	3.75	0.01
25	24.96°	3.57	0.03
26	25.34°	3.51	0.02
27	25.71°	3.46	0.01
28	26.11°	3.41	0.10
29	26.76°	3.33	0.06
30	27.08°	3.29	0.26
31	28.12°	3.17	0.01

19

TABLE 2a-continued

XRPD Peak data for free base pattern 1 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
32	28.44°	3.14	0.01
33	29.18°	3.06	0.01
34	29.81°	3.00	0.01
35	30.33°	2.95	0.01
36	31.07°	2.88	0.01

TABLE 2b

XRPD Peak data for free base pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	5.5°	16.2	0.0
2	7.6°	11.6	0.0
3	9.3°	9.5	0.0
4	10.5°	8.5	0.0
5	10.9°	8.1	0.2
6	11.9°	7.5	0.0
7	12.8°	6.9	0.0
8	13.2°	6.7	0.1
9	15.1°	5.9	0.0
10	15.8°	5.6	0.0
11	16.3°	5.5	0.0
12	16.6°	5.3	0.1
13	17.2°	5.2	0.1
14	17.9°	4.9	0.0
15	18.5°	4.8	0.1
16	19.5°	4.6	0.0
17	19.8°	4.5	0.0
18	20.4°	4.3	0.0
19	20.8°	4.3	0.1
20	21.1°	4.2	0.1
21	21.7°	4.1	1.0
22	22.0°	4.0	0.1
23	23.0°	3.9	0.0
24	23.7°	3.8	0.0
25	25.0°	3.6	0.0
26	25.3°	3.5	0.0
27	25.7°	3.5	0.0
28	26.1°	3.4	0.1
29	26.8°	3.3	0.1
30	27.1°	3.3	0.3
31	28.1°	3.2	0.0
32	28.4°	3.1	0.0
33	29.2°	3.1	0.0
34	29.8°	3.0	0.0
35	30.3°	2.9	0.0
36	31.1°	2.9	0.0

## Thermo—Gravimetric Analysis (TGA)

TGA thermogram of free base in FIG. 2 showed no weight loss between 25 to 150° C. and good thermal stability up to 150° C. followed by a rapid weight loss due to the thermal degradation of the API.

## Differential Scanning Calorimetry (DSC)

The first heating ramp displayed a sharp endothermic event with  $T_{onset}$  at 66.4° C. and heat of fusion 96.4 J/g, which corresponds to melting of the free base as shown in, or substantially as shown in, FIG. 3.

The cooling ramp of 10° C./min from 205° C. to -90° C. displayed a vitrification at around -15.3° C. The 2nd heating ramp showed an endothermic shift in the baseline around -11.9° C. ( $T_g$ ), followed by recrystallisation exotherm with the onset temperature of 36° C. The sharp endotherm with onset temperature of 65.4° C. corresponds to melting of pattern 1 as demonstrated in FIG. 4.

20

## Nuclear Magnetic Resonance Spectroscopy (NMR)

The  $^1\text{H}$  NMR spectrum of free base in  $d_6$ -DMSO shown in FIG. 5 provided structure confirmation of the material. There is no obvious evidence of residual solvents present in the sample.

## Salt Screen Studies

Salt screens experiments were consisted of combining solutions of the API and 1.05 stoichiometric amounts of counter ions.

Selected counter ions for salt studies are tabulated in Table 3.

TABLE 3

Selected counter ions	
Counter ions - Set 1	
1	Sulphuric acid 98%
2	p-Toluene sulphonic acid
3	Methane sulphonic acid
4	Benzene sulphonic acid
5	Maleic acid
6	Phosphoric acid
7	Ethane sulphonic acid 70 wt % in water
8	L-Tartaric acid
9	Fumaric acid
10	(2S)-5-oxopyrrolidine-2- carboxylic acid
11	L-Lactic acid
12	Citric acid
Counter ions - Set 2	
13	Hydrobromic acid
14	Oxalic acid
15	2-Hydroxy ethanesulphonic acid
16	L-Glutamic acid
17	Ketoglutaric acid
18	L-Malic acid
19	Glycolic acid
20	Adipic acid
21	Acetic acid
22	Propionic acid
23	Hippuric acid
24	Saccharin

Free base (2.16 g) was dissolved in 1,4-dioxane (72 ml) at room temperature. Approximately 1 ml of this the stock solution was dispensed to 72 (2 ml) HPLC vials. This preparation was carried out twice for each set of 12 counter ions.

## Set 1:

Free Base dioxane solutions were then frozen at -20° C. for 5 hours. After this time frozen samples were lyophilised for approximately 60 hours.

## Set 2:

Free Base dioxane solutions were then frozen at -20° C. for 5 hours. After this time frozen samples were lyophilised for 12 hours.

To freeze dried samples approximately 0.5 ml of solvent was added. Solvents used in this study are tabulated in Table 4.

TABLE 4

Solvents used in salt screening experiments	
Acetone	
EtOAc	
MeCN	
THF	
IPA	
5% water:EtOH	

Acid stock solutions in 1.05 eq. ratio were added to free base samples. Solvents used for preparation of acid stock solutions are summarised in Table 5. Due to poor solubility of L-Glutamic acid in examined solvents, L-Glutamic acid was added as a solid in 1.05 eq. to free base samples.

TABLE 5

Acid stock solutions			
Counter ions-Set 1	Stock Solvent	Counter ions - Set 2	Stock Solvent
Sulphuric acid 98%	Water	Hydrobromic acid	Water
p-Toluene sulphonic acid	Water	Oxalic acid	Water
Methane sulphonic acid	Water	2-Hydroxy ethanesulphonic acid	Water
Benzene sulphonic acid	Water	L-Glutamic acid	N/A
Maleic acid	Water	Ketoglutaric acid	Water
Phosphoric acid	Water	L-Malic acid	Water
Ethane sulphonic acid 70 wt % in water	Water	Glycolic acid	Water
L-Tartaric acid	Water	Adipic acid	THF
Fumaric acid	5% Water: EtOH	Acetic acid	Water
(2S)-5-oxopyrrolidine-2-carboxylic acid	EtOH	Propionic acid	Water

TABLE 5-continued

Acid stock solutions			
Counter ions-Set 1	Stock Solvent	Counter ions - Set 2	Stock Solvent
L-Lactic acid	Water	Hippuric acid	THF:MeOH (1:1; v/v)
Citric acid	Water	Saccharin	THF

No formation of solid phases was observed after mixing acids with API solutions.

Thermal cycling experiment was performed on samples between ambient and 40° C. Temperature was held for 4 hours at each condition. Thermal cycling—Set-1 for 20 hours. Thermal cycling—Set-2 for 24 hours.

Where solids were observed after thermal cycling, these were isolated by centrifuge filtration using Nylon 0.2 micrometre centrifuge filter tubes and analysed by XRPD.

Any new crystalline forms were also analysed by TGA, DSC, <sup>1</sup>H NMR and DVS analyses.

Remaining solutions were first cooled to 4° C. for 2 hours to promote precipitation. As no precipitation occurred solutions were allowed to evaporate under ambient conditions.

The outcome of experiments is summarised in Table 6 and Table 7, respectively.

TABLE 6

Outcome summary of experiments, Set-1				
Acid	Solvent	Batch	After thermal cycling	After evaporation to dryness
Sulphuric acid	Acetone	DXD2203-004-001	Yellow solution	Yellow glass
	EtOAc	DXD2203-004-002	Dark pink solution	Dark pink glass
	MeCN	DXD2203-004-003	Pink solution	Pink glass
	THF	DXD2203-004-004	Pink solution	Pink glass
	IPA	DXD2203-004-005	Pink solution	Pink glass
	5% water:EtOH	DXD2203-004-006	Pink solution	Pink glass
p-toluene sulphonic acid	Acetone	DXD2203-004-007	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-008	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-009	Light yellow solution	Light yellow glass
	THF	DXD2203-004-010	Pink solution	Pink glass
	IPA	DXD2203-004-011	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-004-012	Pink solution	Pink glass
Methane sulphonic acid	Acetone	DXD2203-004-013	Yellow solution	Yellow glass
	EtOAc	DXD2203-004-014	Pink solution	Pink glass
	MeCN	DXD2203-004-015	Yellow solution	Yellow glass
	THF	DXD2203-004-016	Yellow solution	Yellow glass
	IPA	DXD2203-004-017	Pink solution	Pink glass
	5% water:EtOH	DXD2203-004-018	Pink solution	Pink glass
Benzene sulphonic acid	Acetone	DXD2203-004-019	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-020	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-021	Light yellow solution	Light yellow glass
	THF	DXD2203-004-022	Light yellow solution	Light yellow glass
	IPA	DXD2203-004-023	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-004-024	Light yellow solution	Light yellow glass
Maleic acid	Acetone	DXD2203-004-025	Yellow solution	Yellow glass
	EtOAc	DXD2203-004-026	Yellow solution	Yellow glass
	MeCN	DXD2203-004-027	Yellow solution	Yellow glass
	THF	DXD2203-004-028	Yellow solution	Yellow glass
	IPA	DXD2203-004-029	Yellow solution	Yellow glass
	5% water:EtOH	DXD2203-004-030	Yellow solution	Yellow glass
Phosphoric acid	Acetone	DXD2203-004-031	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-032	White precipitate	N/A
	MeCN	DXD2203-004-033	Light yellow solution	Light yellow glass
	THF	DXD2203-004-034	White precipitate	N/A
	IPA	DXD2203-004-035	White precipitate	N/A
	5% water:EtOH	DXD2203-004-036	Light yellow solution	Light yellow glass
Ethane sulphonic acid	Acetone	DXD2203-004-037	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-038	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-039	Light yellow solution	Light yellow glass
	THF	DXD2203-004-040	Light yellow solution	Light yellow glass
	IPA	DXD2203-004-041	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-004-042	Light yellow solution	Light yellow glass

TABLE 6-continued

Outcome summary of experiments, Set-1				
Acid	Solvent	Batch	After thermal cycling	After evaporation to dryness
L-tartaric acid	Acetone	DXD2203-004-043	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-044	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-045	Light yellow solution	Light yellow glass
	THF	DXD2203-004-046	Light yellow solution	Light yellow glass
	IPA	DXD2203-004-047	Light yellow solution	Light yellow glass
Fumaric acid	5% water:EtOH	DXD2203-004-048	Light yellow solution	Light yellow glass
	Acetone	DXD2203-004-049	White precipitate	N/A
	EtOAc	DXD2203-004-050	White precipitate	N/A
	MeCN	DXD2203-004-051	Light yellow solution	Light yellow glass
	THF	DXD2203-004-052	Light pink solution	Light pink glass
	IPA	DXD2203-004-053	White precipitate	N/A
	5% water:EtOH	DXD2203-004-054	Light yellow solution	Light yellow glass
(2S)-5oxopyrrolidine-2carboxylic acid	Acetone	DXD2203-004-055	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-056	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-057	Light yellow solution	Light yellow glass
	THF	DXD2203-004-058	Light yellow solution	Light yellow glass
	IPA	DXD2203-004-059	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-004-060	Light yellow solution	Light yellow glass
L-lactic acid	Acetone	DXD2203-004-061	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-062	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-063	Light yellow solution	Light yellow glass
	THF	DXD2203-004-064	Light yellow solution	Light yellow glass
	IPA	DXD2203-004-065	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-004-066	Light yellow solution	Light yellow glass
Citric acid	Acetone	DXD2203-004-067	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-004-068	Light yellow solution	Light yellow glass
	MeCN	DXD2203-004-069	Light yellow solution	Light yellow glass
	THF	DXD2203-004-070	Light yellow solution	Light yellow glass
	IPA	DXD2203-004-071	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-004-072	Light yellow solution	Light yellow glass

TABLE 7

Outcome summary of experiments, Set-2				
Acid	Solvent	Batch	After thermal cycling	After evaporation to dryness
Hydrobromic acid	Acetone	DXD2203-007-01	Yellow solution	Yellow glass
	EtOAc	DXD2203-007-02	Yellow solution	Yellow glass
	MeCN	DXD2203-007-03	Light pink solution	Light pink glass
	THF	DXD2203-007-04	Light pink solution	Light pink glass
	IPA	DXD2203-007-05	Light pink solution	Light pink glass
	5% water:EtOH	DXD2203-007-06	Light pink solution	Light pink glass
Oxalic acid	Acetone	DXD2203-007-07	Off white precipitate	N/A
	EtOAc	DXD2203-007-08	White precipitate	N/A
	MeCN	DXD2203-007-09	Off white precipitate	N/A
	THF	DXD2203-007-10	Off white precipitate	N/A
	IPA	DXD2203-007-11	Off white precipitate	N/A
	5% water:EtOH	DXD2203-007-12	Off white precipitate	N/A
2-hydroxy ethanesulfonic acid	Acetone	DXD2203-007-13	Light yellow solution	Yellow glass
	EtOAc	DXD2203-007-14	Light yellow solution	Yellow glass
	MeCN	DXD2203-007-15	Light pink solution	Light pink glass
	THF	DXD2203-007-16	Light pink solution	Light pink glass
	IPA	DXD2203-007-17	Light pink solution	Light pink glass
L-glutamic acid	5% water:EtOH	DXD2203-007-18	Light pink solution	Light pink glass
	Acetone	DXD2203-007-19	Light yellow solution + undissolved acid	N/A
	EtOAc	DXD2203-007-20	Light yellow-solution + undissolved acid	N/A
	MeCN	DXD2203-007-21	Light yellow-solution + undissolved acid	N/A


TABLE 7-continued

Outcome summary of experiments. Set-2				
Acid	Solvent	Batch	After thermal cycling	After evaporation to dryness
	THF	DXD2203-007-22	Light yellow solution + undissolved acid	N/A
	IPA	DXD2203-007-23	Light yellow solution + undissolved acid	N/A
	5% water:EtOH	DXD2203-007-24	Light yellow solution + undissolved acid	N/A
Ketoglutaric acid	Acetone	DXD2203-007-25	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-007-26	Light yellow solution	Light yellow glass
	MeCN	DXD2203-007-27	Light yellow-solution	Light yellow glass
	THF	DXD2203-007-28	Light yellow-solution	Light yellow glass
	IPA	DXD2203-007-29	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-007-30	Light yellow solution	Light yellow glass
L-malic acid	Acetone	DXD2203-007-31	Yellow solution	Yellow glass
	EtOAc	DXD2203-007-32	Yellow solution	Yellow glass
	MeCN	DXD2203-007-33	Yellow solution	Yellow glass
	THF	DXD2203-007-34	Yellow solution	Yellow glass
	IPA	DXD2203-007-35	Yellow solution	Yellow glass
	5% water:EtOH	DXD2203-007-36	Yellow solution	Yellow glass
Glycolic acid	Acetone	DXD2203-007-37	Yellow solution	Yellow glass
	EtOAc	DXD2203-007-38	Yellow solution	Yellow glass
	MeCN	DXD2203-007-39	Yellow solution	Yellow glass
	THF	DXD2203-007-40	Yellow solution	Yellow glass
	IPA	DXD2203-007-41	Yellow solution	Yellow glass
	5% water:EtOH	DXD2203-007-42	Yellow solution	Yellow glass
Adipic acid	Acetone	DXD2203-007-43	Yellow solution	Yellow glass
	EtOAc	DXD2203-007-44	Off white precipitate	N/A
	MeCN	DXD2203-007-45	Yellow solution	Yellow glass
	THF	DXD2203-007-46	Yellow solution	Yellow glass
Acetic acid	IPA	DXD2203-007-47	Yellow solution	Yellow glass
	5% water:EtOH	DXD2203-007-48	Yellow solution	Yellow glass
	Acetone	DXD2203-007-49	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-007-50	Yellow solution	Yellow glass
	MeCN	DXD2203-007-51	Yellow solution	Yellow glass
	THF	DXD2203-007-52	Yellow solution	Yellow glass
Propionic acid	IPA	DXD2203-007-53	Yellow solution	Yellow glass
	5% water:EtOH	DXD2203-007-54	Yellow solution	Yellow glass
	Acetone	DXD22 03-007-55	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-007-56	Yellow solution	Yellow glass
	MeCN	DXD2203-007-57	Yellow solution	Yellow solution
	THF	DXD2203-007-58	Yellow solution	Yellow solution
Hippuric acid	IPA	DXD2203-007-59	Yellow solution	Yellow solution
	5% water:EtOH	DXD2203-007-60	Yellow solution	Yellow solution
	Acetone	DXD2203-007-61	Light yellow solution	Light yellow glass
	EtOAc	DXD2203-007-62	Light yellow solution	Light yellow glass
	MeCN	DXD2203-007-63	Light yellow solution	Light yellow glass
	THF	DXD2203-007-64	Light yellow solution	Light yellow glass
Saccharin	IPA	DXD2203-007-65	Light yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-007-66	Light yellow solution	Light yellow glass
	Acetone	DXD2203-007-67	Yellow solution	Light yellow glass
	EtOAc	DXD2203-007-68	Yellow solution	Light yellow glass
	MeCN	DXD2203-007-69	Yellow solution	Light yellow glass
	THF	DXD2203-007-70	Yellow solution	Light yellow glass
	IPA	DXD2203-007-71	Yellow solution	Light yellow glass
	5% water:EtOH	DXD2203-007-72	Yellow solution	Light yellow glass



27

Phosphate Salt



The Phosphate salt showed the same crystalline XRPD pattern for all three solids isolated from different solvents as displayed in FIG. 6. This crystalline form was nominated as pattern 1 and XRPD peak data are tabulated in Table 8, Table 8a or Table 8b.

TABLE 8

XRPD Peak data for Phosphate pattern 1.			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	6.012°	14.690	0.296
2	6.529°	13.527	0.961
3	9.045°	9.769	0.142
4	11.855°	7.459	0.061
5	12.888°	6.863	1.000
6	14.357°	6.165	0.404
7	16.929°	5.233	0.042
8	18.004°	4.923	0.253
9	18.809°	4.714	0.087
10	19.319°	4.591	0.748
11	20.353°	4.360	0.922
12	23.104°	3.847	0.795
13	24.607°	3.615	0.012
14	25.343°	3.512	0.068
15	26.625°	3.345	0.036
16	27.128°	3.284	0.025
17	27.982°	3.136	0.018
18	28.784°	3.099	0.039
19	30.815°	2.899	0.032
20	31.505°	2.837	0.016

TABLE 8a

XRPD Peak data for Phosphate pattern 1 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	6.01°	14.69	0.30
2	6.53°	13.53	0.96
3	9.05°	9.77	0.14
4	11.86°	7.46	0.06
5	12.89°	6.86	1.00
6	14.36°	6.17	0.40
7	16.93°	5.23	0.04
8	18.00°	4.92	0.25
9	18.81°	4.71	0.09
10	19.32°	4.59	0.75
11	20.35°	4.36	0.92
12	23.1°	3.85	0.80
13	24.61°	3.62	0.01
14	25.34°	3.51	0.07
15	26.63°	3.35	0.04
16	27.13°	3.28	0.03
17	27.98°	3.19	0.02
18	28.78°	3.10	0.04
19	30.82°	2.90	0.03
20	31.51°	2.34	0.02

28

TABLE 8b

XRPD Peak data for Phosphate pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	6.0°	14.7	0.3
2	6.5°	13.5	1.0
3	9.0°	9.8	0.1
4	11.9°	7.5	0.1
5	12.9°	6.9	1.0
6	14.4°	6.2	0.4
7	16.9°	5.2	0.0
8	18.0°	4.9	0.3
9	18.3°	4.7	0.1
10	19.3°	4.6	0.7
11	20.4°	4.4	0.9
12	23.1°	3.8	0.8
13	24.6°	3.6	0.0
14	25.3°	3.5	0.1
15	26.6°	3.3	0.0
16	27.1°	3.3	0.0
17	28.0°	3.2	0.0
18	28.3°	3.1	0.0
19	30.8°	2.9	0.0
20	31.5°	2.8	0.0

The TGA thermogram of Phosphate salt presented in FIG. 7 displayed a two-step weight loss before the thermal decomposition. From ambient temperature to 60° C. the weight loss of 2.8%, which corresponds to loss of IPA (~0.15 eq) from surface of particles. The second weight loss of 4.9% between 60 to 130° C. is due to dehydration of ~0.9 eq of water.

The 1st heating DSC thermogram in FIG. 8 displayed a broad endotherm corresponding to desolvation/dehydration process. The melting endothermic event of the Phosphate salt with  $T_{onset}$  around 90.1° C. and heat of fusion 163.6 J/g is followed by the thermal degradation of the material. The cooling ramp of 10° C./min from 215° C. to -90° C. displayed a vitrification around 71.7° C. and the 2nd heat cycle the glass transition around 75.1° C.

The <sup>1</sup>H NMR spectrum of Phosphate salt in d<sub>6</sub>-DMSO solvent is shown in FIG. 9. Proton chemical shift changes when compared to <sup>1</sup>H NMR spectrum of free base indicate salt formation. Approximately 0.16 eq of IPA solvent was observed.

The <sup>31</sup>P NMR spectrum of Phosphate salt in d<sub>6</sub>-DMSO shows a singlet peak at around 0 ppm, confirming the presence of phosphoric acid as displayed in FIG. 10.

DVS analysis was performed using a small sample mass due to material constraints. The sample shows no evidence of form change and only shows evidence of the material drying out. It would be wise to repeat this experiment if more material becomes available. FIG. 11 displays DVS Isotherm plot of Phosphate salt.

The DVS kinetic plot of Phosphate salt DXD220-004-35 is shown in FIG. 12. It can be seen from the XRPD diffractogram in FIG. 13 that the post DVS Phosphate salt is missing peaks at around 5.9, 11.9 and 18.8 2 $\theta$  when compared to the input material as indicated by arrows. Also, the post DVS sample displayed a new shoulder at around 12.6 and 19.9 2 $\theta$ , respectively as indicated by asterisks, which are not characteristics of free base.

In one embodiment, there is provided 5-MeO-DMT phosphate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT phosphate. In one embodiment, there is provided crystalline 5-MeO-DMT phosphate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT phosphate, as characterised by one or more of:

29

An XRPD pattern as shown in, or substantially as shown in, FIG. 6;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, or twenty peaks in an XRPD diffractogram as detailed in Table 8, Table 8a or Table 8b;

One or more, two or more, three or more, four or more, five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 8, Table 8a or Table 8b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 7;

A weight loss of 2.8% between ambient temperature and 60° C., as measured by TGA thermogram;

A weight loss of between 1.5 to 3.5% between ambient temperature and 60° C., as measured by TGA thermogram;

A weight loss of 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9, 3.0, 3.1, 3.2, 3.3, 3.4 or 3.5% between ambient temperature and 60° C., as measured by TGA thermogram;

A weight loss of 4.9% between 60 to 130° C., as measured by TGA thermogram;

A weight loss of between 3.5 to 6.5% between 60 to 130° C., as measured by TGA thermogram;

A weight loss of 3.5, 3.6, 3.7, 3.8, 3.9, 4.0, 4.1, 4.2, 4.3, 4.4, 4.5, 4.6, 4.7, 4.8, 4.9, 5.0, 5.1, 5.2, 5.3, 5.4, 5.5, 5.6, 5.7, 5.8, 5.9, 6.0, 6.1, 6.2, 6.3, 6.4 or 6.5% between 60 and 130° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 8;

A melting endothermic event with an onset of around 90.1° C. and a heat of fusion of 163.6 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 85 to 95° C. and a heat of fusion of around 155 to 170 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 85, 86, 87, 88, 89, 90, 91, 92, 93, 94 or 95° C. and a heat of fusion of around 155, 156, 157, 158, 159, 160, 161, 162, 163, 164, 165, 166, 167, 168, 169 or 170 J/g, as measured in a DSC thermogram;

A vitrification around 71.7° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 65-75° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 65, 66, 67, 68, 69, 70, 71, 72, 73, 74 or 75° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 75.1° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 70-80° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 or 80° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 9;

30


A <sup>31</sup>P NMR spectrum as shown in, or substantially as shown in, FIG. 10;

A DVS isotherm as shown in, or substantially as shown in, FIG. 11; and/or

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 12.

In one embodiment, there is provided crystalline 5-MeO-DMT phosphate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT phosphate, as characterised by an XRPD pattern as shown in, or substantially as shown in, FIG. 13.

Fumarate Salt



The XRPD results shown in FIG. 14 confirmed the crystallinity of Fumarate salt. All three isolated samples from different solvents displayed the same crystalline pattern. This was nominated as pattern 1 with XRPD data peak presented in Table 9, Table 9a or Table 9b.

TABLE 9

XRPD data peak for Fumarate pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. intensity
1	5.911°	14.940	0.015
2	6.459°	13.673	0.073
3	10.694°	8.266	0.012
4	12.107°	7.305	0.104
5	12.981°	6.815	1.000
6	14.203°	6.231	0.021
7	14.975°	5.911	0.034
8	16.304°	5.432	0.673
9	17.532°	5.055	0.233
10	18.166°	4.879	0.498
11	19.220°	4.614	0.604
12	19.443°	4.561	0.345
13	20.396°	4.351	0.598
14	20.783°	4.271	0.073
15	21.487°	4.132	0.037
16	22.052°	4.028	0.859
17	22.834°	3.891	0.523
18	24.255°	3.667	0.015
19	25.064°	3.550	0.243
20	25.410°	3.502	0.310
21	26.116°	3.409	0.200
22	27.542°	3.236	0.135
23	28.051°	3.178	0.174
24	28.709°	3.107	0.035
25	30.166°	2.960	0.071
26	30.482°	2.930	0.024
27	30.764°	2.904	0.031
28	31.022°	2.880	0.034
29	31.555°	2.833	0.008

TABLE 9a

XRPD data peak for Fumarate pattern 1 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	5.91°	14.94	0.02
2	6.46°	13.67	0.07
3	10.69°	8.27	0.01
4	12.11°	7.31	0.10
5	12.98°	6.82	1.00
6	14.20°	6.23	0.02
7	14.98°	5.91	0.03
8	16.30°	5.43	0.67
9	17.53°	5.06	0.23
10	18.17°	4.88	0.50
11	19.22°	4.61	0.60
12	19.45°	4.56	0.35
13	20.40°	4.35	0.60
14	20.78°	4.27	0.07
15	21.49°	4.13	0.04
16	22.05°	4.03	0.86
17	22.83°	3.89	0.52
18	24.26°	3.67	0.02
19	25.06°	3.55	0.24
20	25.41°	3.50	0.31
21	26.12°	3.41	0.20
22	27.54°	3.24	0.14
23	28.05°	3.18	0.17
24	28.71°	3.11	0.04
25	30.17°	2.96	0.07
26	30.48°	2.93	0.02
27	30.76°	2.90	0.03
28	31.02°	2.88	0.03
29	31.56°	2.83	0.01

TABLE 9b

XRPD data peak for Fumarate pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	5.9°	14.9	0.0
2	6.5°	13.7	0.1
3	10.7°	8.3	0.0
4	12.1°	7.3	0.1
5	13.0°	6.8	1.0
6	14.2°	6.2	0.0
7	15.0°	5.9	0.0
8	16.3°	5.4	0.7
9	17.5°	5.1	0.2
10	18.2°	4.9	0.5
11	19.2°	4.6	0.6
12	19.4°	4.6	0.3
13	20.4°	4.4	0.6
14	20.8°	4.3	0.1
15	21.5°	4.1	0.0
16	22.1°	4.0	0.9
17	22.8°	3.9	0.5
18	24.3°	3.7	0.0
19	25.1°	3.6	0.2
20	25.4°	3.5	0.3
21	26.1°	3.4	0.2
22	27.5°	3.2	0.1
23	28.1°	3.2	0.2
24	28.7°	3.1	0.0
25	30.2°	3.0	0.1
26	30.5°	2.9	0.0
27	30.8°	2.9	0.0
28	31.0°	2.9	0.0
29	31.6°	2.8	0.0

The TGA thermogram of Fumarate salt presented in FIG. 15, displayed no presence of residual solvents and good thermal stability up to 150° C.

The 1st heating DSC data showed two small endothermic events around 93.7° C. and 134.0° C. respectively, corre-

sponding to solid-state transformations. The sharp endotherm with  $T_{onset}$  around 176.5° C. and heat of fusion 92.3 J/g corresponds to melting, followed by the thermal degradation of the Fumarate salt as demonstrated in FIG. 16.

The 10° C./min cooling ramp from 215° C. to -90° C. displayed a vitrification around 45.1° C. and the 2nd heating cycle exhibited a glass transition around 51.1° C. as shown in, or substantially as shown in, FIG. 17.

The <sup>1</sup>H NMR spectrum of Fumarate salt in d<sub>6</sub>-DMSO shown in FIG. 18 displayed around 1.4 eq of fumaric acid present as well as traces of IPA solvent.

In one embodiment, there is provided 5-MeO-DMT fumarate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT fumarate. In one embodiment, there is provided crystalline 5-MeO-DMT fumarate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT fumarate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 14;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, or twenty nine peaks in an XRPD diffractogram as detailed in Table 9, Table 9a or Table 9b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 9, Table 9a or Table 9b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 15;

A DSC thermogram as shown in, or substantially as shown in, FIG. 16;

Two small endothermic events around 93.7° C. and 134.0° C. respectively as measured in a DSC thermogram;

Two small endothermic events around 85-100° C. and 130-140° C. respectively as measured in a DSC thermogram;

Two small endothermic events around 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100° C. and 130, 131, 132, 133, 134, 135, 136, 137, 138, 139 or 140° C. respectively as measured in a DSC thermogram;

A sharp endotherm with an onset of around 176.5° C. and heat of fusion 92.3 J/g as measured in a DSC thermogram;

A sharp endotherm with an onset of around 165-185° C. and heat of fusion about 88-100 J/g as measured in a DSC thermogram;

A sharp endotherm with an onset of around 165, 166, 167, 168, 169, 170, 171, 172, 173, 174, 175, 176, 177, 178, 179, 180, 181, 182, 183, 184, or 185° C. and heat of fusion about 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99, or 100 J/g as measured in a DSC thermogram;

A vitrification around 45.1° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 40-50° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

## 33

A vitrification around 40, 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;


A glass transition around 51.1° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 45-55° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 or 55° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.; and/or

A 1H NMR spectrum as shown in, or substantially as shown in, FIG. 18.

Oxalate Salt



XRPD diffractograms of Oxalate salt displayed the same crystalline solid form for all samples as shown in, or substantially as shown in, FIG. 19. This was nominated as pattern 1 with XRPD data peak presented in Table 10, 10a and 10b.

TABLE 10

XRPD Peak data for Oxalate pattern 1.			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	7.923°	11.150	0.062
2	10.999°	8.037	0.012
3	11.688°	7.565	0.002
4	12.963°	6.824	1.000
c	13.697°	6.460	0.095
6	14.779°	5.989	0.005
7	16.600°	5.336	0.001
8	17.217°	5.146	0.002
9	17.777°	4.985	0.004
10	18.284°	4.843	0.001
11	18.570°	4.774	0.010
12	18.861°	4.701	0.003
13	20.147°	4.404	0.005
14	19.928°	4.452	0.214
15	21.360°	4.157	0.004
16	21.967°	4.043	0.100
17	23.560°	3.773	0.077
18	23.933°	3.715	0.003
19	24.348°	3.653	0.003
20	25.159°	3.537	0.002
21	25.573°	3.481	0.010
22	25.961°	3.429	0.377
23	26.348°	3.380	0.015
24	26.605°	3.348	0.001
25	27.458°	3.246	0.080
26	27.960°	3.189	0.001
27	29.719°	3.004	0.011
28	30.525°	2.926	0.005
29	31.562°	2.832	0.015

## 34

TABLE 10a

XRPD Peak data for Oxalate pattern 1 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	7.92°	11.15	0.06
2	11.00°	8.04	0.01
3	11.69°	7.57	0.00
4	12.96°	6.82	1.00
5	13.7°	6.46	0.10
6	14.78°	5.99	0.01
7	16.60°	5.34	0.00
8	17.22°	5.15	0.00
9	17.78°	4.99	0.00
10	18.28°	4.85	0.00
11	18.57°	4.77	0.01
12	18.86°	4.70	0.00
13	20.15°	4.40	0.01
14	19.93°	4.45	0.21
15	21.36°	4.16	0.00
16	21.97°	4.04	0.10
17	23.56°	3.77	0.08
18	23.93°	3.72	0.00
19	24.35°	3.65	0.00
20	25.16°	3.54	0.00
21	25.57°	3.48	0.01
22	25.96°	3.43	0.38
23	26.35°	3.38	0.02
24	26.61°	3.35	0.00
25	27.46°	3.25	0.08
26	27.96°	3.19	0.00
27	29.72°	3.00	0.01
28	30.53°	2.93	0.01
29	31.56°	2.83	0.02

TABLE 10b

XRPD Peak data for Oxalate pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
1	7.9°	11.2	0.1
2	11.0°	8.0	0.0
3	11.7°	7.6	0.0
4	13.0°	6.8	1.0
5	13.7°	6.5	0.1
6	14.8°	6.0	0.0
7	16.6°	5.3	0.0
8	17.2°	5.1	0.0
9	17.8°	5.0	0.0
10	18.3°	4.8	0.0
11	18.6°	4.8	0.0
12	18.9°	4.7	0.0
13	20.1°	4.4	0.0
14	19.9°	4.5	0.2
15	21.4°	4.2	0.0
16	22.0°	4.0	0.1
17	23.6°	3.3	0.1
18	23.9°	3.7	0.0
19	24.3°	3.7	0.0
20	25.2°	3.5	0.0
21	25.6°	3.5	0.0
22	26.0°	3.4	0.4
23	26.3°	3.4	0.0
24	26.6°	3.3	0.0
25	27.5°	3.2	0.1
26	28.0°	3.2	0.0
27	29.7°	3.0	0.0
28	30.5°	2.9	0.0
29	31.6°	2.8	0.0

TGA analysis of the Oxalate salt shows 0.6% weight loss between 25-180° C. (~0.02 moles EtOAc) followed by a single step thermal degradation as shown in, or substantially as shown in, FIG. 20. The 1st heating DSC thermogram displayed a single melting endotherm with  $T_{onst}$  around 176.1° C., followed by the decomposition of the material.

35

The enthalpy associated with the endothermic peak is 157.5 J/g as shown in, or substantially as shown in, FIG. 21. DSC thermograms of Oxalate salt showed a vitrification around 50.7° C. upon cooling and glass transition at 58.0° C. during the 2nd heating cycle as shown in, or substantially as shown in, FIG. 22.

Proton chemical shift changes in NMR ( $d_6$ -DMSO) spectrum indicate Oxalate salt formation. Traces of EtOAc were also detected in the spectrum as shown in, or substantially as shown in, FIG. 23.

$^{13}\text{C}$  NMR spectrum of Oxalate salt in  $d_6$ -DMSO presented in FIG. 24, showed a signal for carbon at 164 ppm, confirming the presence of the oxalic acid. The quantitative  $^{13}\text{C}$  NMR spectrum of Oxalate salt in  $d_6$ -DMSO is presented in FIG. 25. The signal for carbon at 164 ppm, confirming the presence of ~1 eq of the oxalic acid. DVS Analysis of the Oxalate salt lot DXD2203-007-07 was performed and the isotherm plot is shown in FIG. 26. The DVS kinetic plot of the Oxalate salt lot DXD2203-007-07 is displayed in FIG. 27 and showed no evidence of a form change. XRPD analysis performed on post DVS Oxalate salt showed that no change in the crystalline form occurred during the DVS experiment as demonstrated in FIG. 28.

In one embodiment, there is provided 5-MeO-DMT oxalate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT oxalate. In one embodiment, there is provided crystalline 5-MeO-DMT oxalate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT oxalate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 19;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, or twenty nine peaks in an XRPD diffractogram as detailed in Table 10, Table 10a or Table 10b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 10, Table 10a or Table 10b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 20;

A weight loss of around 0.6% between 25-180° C., as measured by TGA thermogram;

A weight loss of around 0.1-1.0% between 25-180° C., as measured by TGA thermogram;

A weight loss of around 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0% between 25-180° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 21;

A melting endothermic event with an onset of around 176.1° C. and an enthalpy of 157.5 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 170-180° C. and an enthalpy of around 152-162 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 170, 171, 172, 173, 174, 175, 176, 177, 178, 179 or 180° C.

36

and an enthalpy of around 152, 153, 154, 155, 156, 157, 158, 159, 160, 161 or 162 J/g, as measured in a DSC thermogram;

A vitrification around 50.7° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 45-55° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 or 55° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 58.0° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 53-63° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 53, 54, 55, 56, 57, 58, 59, 60, 61, 62 or 63° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A  $^1\text{H}$  NMR spectrum as shown in, or substantially as shown in, FIG. 23;

A  $^{13}\text{C}$  NMR spectrum as shown in, or substantially as shown in, FIG. 24;


A  $^{13}\text{C}$  NMR spectrum as shown in, or substantially as shown in, FIG. 25;

A DVS isotherm as shown in, or substantially as shown in, FIG. 26;

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 27; and/or

An XRPD pattern as shown in, or substantially as shown in, FIG. 28.

Adipate Salt



XRPD diffractogram of isolated solid from ethyl acetate displayed a crystalline pattern which does not correspond to free base pattern 1 or adipic acid as shown in, or substantially as shown in, FIG. 29. Later analysis indicated this to be a free form and as such nominated as pattern 2.

TABLE 11

XRPD Peak data for DXD2203-007-44, pattern 2.

Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.368°	9.433	1.000
2	14.136°	6.260	0.493
3	14.535°	6.089	0.039
4	17.694°	5.009	0.611
5	18.017°	4.919	0.448
6	18.692°	4.743	0.052
7	19.490°	4.551	0.414

TABLE 11-continued

XRPD Peak data for DXD2203-007-44, pattern 2.			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
8	20.846°	4.258	0.309
9	21.557°	4.119	0.338
10	21.831°	4.068	0.433
11	23.433°	3.793	0.368
12	24.948°	3.566	0.027
13	25.633°	3.473	0.117
14	25.831°	3.446	0.308
15	26.875°	3.315	0.157
16	28.213°	3.161	0.214
17	28.503°	3.129	0.058
18	30.224°	2.955	0.079
19	30.499°	2.929	0.052
20	31.203°	2.864	0.054
21	31.867°	2.806	0.033

TABLE 11a

XRPD Peak data for DXD2203-007-44, pattern 2 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.37°	9.43	1.00
2	14.14°	6.26	0.49
3	14.54°	6.09	0.04
4	17.69°	5.01	0.61
5	18.02°	4.92	0.45
6	18.69°	4.74	0.05
7	19.49°	4.55	0.41
8	20.85°	4.26	0.31
9	21.56°	4.12	0.34
10	21.83°	4.07	0.43
11	23.43°	3.79	0.37
12	24.95°	3.57	0.03
13	25.63°	3.47	0.12
14	25.83°	3.45	0.31
15	26.88°	3.32	0.16
16	28.21°	3.16	0.21
17	28.50°	3.13	0.06
18	30.22°	2.96	0.08
19	30.50°	2.93	0.05
20	31.20°	2.86	0.05
21	31.87°	2.81	0.03

TABLE 11b

XRPD Peak data for DXD2203-007-44, pattern 2 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.4°	9.4	1.0
2	14.1°	6.3	0.5
3	14.5°	6.1	0.0
4	17.7°	5.0	0.6
5	18.0°	4.9	0.4
6	18.7°	4.7	0.1
7	19.5°	4.6	0.4
8	20.8°	4.3	0.3
9	21.6°	4.1	0.3
10	21.8°	4.1	0.4
11	23.4°	3.8	0.4
12	24.9°	3.6	0.0
13	25.6°	3.5	0.1
14	25.8°	3.4	0.3
15	26.9°	3.3	0.2
16	28.2°	3.2	0.2

TABLE 11b-continued

XRPD Peak data for DXD2203-007-44, pattern 2 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
17	28.5°	3.1	0.1
18	30.2°	3.0	0.1
19	30.5°	2.9	0.1
20	31.2°	2.9	0.1
21	31.9°	2.8	0.0

The TGA thermograph displayed in FIG. 30 showed a weight loss of 0.5% between 25-170° C., (~0.03 moles EtOAc) followed by the thermal degradation of the material.

The 1<sup>st</sup> heating DSC thermogram shown in FIG. 31, displayed a broad endotherm with onset temperature of 73.9° C. and heat of fusion 90.9 J/g corresponding to the melt of the material.

As shown in, or substantially as shown in, FIG. 32, the cooling ramp from 215° C. to -90° C. at 10° C./min displayed a vitrification at around 3.1° C. The glass transition at around 7.4° C. was observed during the 2<sup>nd</sup> heating ramp.

<sup>1</sup>H NMR spectrum (d<sub>6</sub>-DMSO) of DXD2203-007-44 solid displayed in FIG. 33 showed traces of ethyl acetate and no presence of adipic acid. This suggests that isolated material is a free base of different crystalline form as the XRPD pattern of DXD2203-007-44 does not match with the input (DXD2203003-01). This was nominated as pattern 2 of free base.

However, it is interesting to note the significant difference in glass transitions. As mentioned previously, free base pattern 1 displayed Tg around -11.9° C. during the second heating cycle, whereas free base pattern 2 showed Tg around 7.4° C.

In one embodiment, there is provided 5-MeO-DMT adipate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT adipate. In one embodiment, there is provided crystalline 5-MeO-DMT adipate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT adipate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 29;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, or twenty one peaks in an XRPD diffractogram as detailed in Table 11, Table 11a or Table 11b;

One or more, two or more, three or more, four or more, or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 11, Table 11a or Table 11b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 30;

A weight loss of 0.5% between 25-170° C., as measured by TGA thermogram;

A weight loss of around 0.1-1.0% between 25-170° C., as measured by TGA thermogram;

A weight loss of around 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0% between 25-170° C., as measured by TGA thermogram;

39

A DSC thermogram as shown in, or substantially as shown in, FIG. 31 or 32;

A melting endothermic event with an onset of around 73.9° C. and an enthalpy of 90.9 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 68-80° C. and an enthalpy of around 85-95 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 or 80° C. and an enthalpy of around 85, 86, 87, 88, 89, 90, 91, 92, 93, 94 or 95 J/g, as measured in a DSC thermogram;

A vitrification around 3.1° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 0-10° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A vitrification around 0, 1, 2, 3, 4, 5, 6, 7, 8, 9 or 10° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 7.4° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 2-12° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A glass transition around 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 215° C. to -90° C.;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 33.

#### Solvent—Anti-Solvent Experiments

Amorphous salts produced by a slow evaporation to dryness at RT (DXD2203-004 and DXD2203-007) were used for solvent/anti-solvent experiments.

Approximately 1 ml of solvent was added to amorphous salt and samples were placed to 40° C. chamber for one hour. After this time clear solutions were allowed to cool down to room temperature and antisolvent (approximately 2 ml) was added dropwise. Solvent/anti-solvent systems used are tabulated in Table 12.

In some cases, precipitates were formed during cooling solutions to the room temperature. Saccharinate amorphous salt did not dissolve after one hour at 40° C. and the IPA solvent was allowed to evaporate.

Produced solids were isolated by centrifuge filtration using Nylon 0.2 micrometre centrifuge filter tubes and analysed by XRPD.

Any new crystalline forms were also analysed by TGA, DSC, <sup>1</sup>H NMR and DVS analyses.

TABLE 12


Solvent/Anti-solvent experiments.			
Salt	Batch	Solvent	Anti-solvent
Tartrate	DXD2203-009-08	IPA	hexane
	DXD2203-009-09	THF	hexane
	DXD2203-009-12	EtOH (40° C.)	N/A
	DXD2203-009-13	THF (40° C.)	N/A
	DXD2203-009-10	THF	hexane
Fumarate	DXD2203-009-11	THF (40° C.)	N/A
	DXD2203-009-20	IPA	hexane
Tosylate	DXD2203-009-21	IPA	Hexane
Saccharinate	DXD2203-010-01	IPA	N/A

40

TABLE 12-continued

Solvent/Anti-solvent experiments.			
Salt	Batch	Solvent	Anti-solvent
Hydrobromide	DXD2203-010-02	MeOH	MTBE
	DXD2203-010-08	DMF	toluene
	DXD2203-010-09	MeCN	toluene
Glycolate	DXD2203-010-03	IPAC (40° C.)	N/A
	DXD2203-010-04	EtOH	MTBE
Ketoglutarate	DXD2203-010-06	MeOH	MTBE
	DXD2203-010-05	EtOH	MTBE
Malate	DXD2203-010-07	IPAC	MTBE

#### Tartrate Salt



XRPD diffractograms of Tartrate salt showed that same crystalline solid form was isolated from all solvent systems as demonstrated in FIG. 34. This was nominated as pattern 1 with XRPD peak data displayed in Table 13, Table 13a or Table 13b.

TABLE 13

XRPD Peak data for Tartrate pattern 1.			
Peak No.	Angle 2θ	d Value	Rel. Intensity
1	10.147°	8.711	0.045
2	10.987°	8.046	0.012
3	11.794°	7.497	0.050
4	13.080°	6.763	0.246
5	14.786°	5.986	0.015
6	15.214°	5.819	0.331
7	15.464°	5.725	0.093
8	16.534°	5.357	0.138
9	17.793°	4.981	0.374
10	18.267°	4.853	0.692
11	18.579°	4.772	0.687
12	18.785°	4.720	0.684
13	19.360°	4.581	0.044
14	19.630°	4.519	0.187
15	20.311°	4.369	0.683
16	20.730°	4.281	1.000
17	21.467°	4.136	0.161
18	22.125°	4.015	0.461
19	22.639°	3.924	0.179
20	23.041°	3.857	0.114
21	23.679°	3.754	0.258
22	24.856°	3.579	0.079
23	25.431°	3.500	0.135
24	26.034°	3.420	0.749
25	26.368°	3.377	0.042
26	27.029°	3.296	0.068
27	27.299°	3.264	0.130
28	27.870°	3.199	0.064

41

TABLE 13-continued

XRPD Peak data for Tartrate pattern 1.			
Peak No.	Angle $2\theta$	d Value	Rel. Intensity
29	28.937°	3.083	0.024
30	29.311°	3.045	0.088
31	30.009°	2.975	0.121
32	31.153°	2.869	0.067
33	31.707°	2.820	0.047

TABLE 13a

XRPD Peak data for Tartrate pattern 1 (2 d.p.).			
Peak No.	Angle $2\theta$	d Value	Rel. Intensity
1	10.15°	8.71	0.05
2	10.99°	8.05	0.01
3	11.79°	7.50	0.05
4	13.08°	6.76	0.25
5	14.79°	5.99	0.02
6	15.21°	5.82	0.33
7	15.46°	5.73	0.09
8	16.53°	5.36	0.14
9	17.79°	4.98	0.37
10	18.27°	4.85	0.69
11	18.58°	4.77	0.69
12	18.79°	4.72	0.68
13	19.36°	4.58	0.04
14	19.63°	4.52	0.19
15	20.31°	4.37	0.68
16	20.73°	4.28	1.00
17	21.47°	4.14	0.16
18	22.13°	4.02	0.46
19	22.64°	3.92	0.18
20	23.04°	3.86	0.11
21	23.68°	3.75	0.26
22	24.86°	3.58	0.08
23	25.43°	3.50	0.14
24	26.03°	3.42	0.75
25	26.37°	3.38	0.04
26	27.03°	3.30	0.07
27	27.30°	3.26	0.13
28	27.87°	3.20	0.06
29	28.94°	3.08	0.02
30	29.31°	3.05	0.09
31	30.01°	2.98	0.12
32	31.15°	2.87	0.07
33	31.71°	2.82	0.05

TABLE 13b

XRPD Peak data for Tartrate pattern 1 (1 d.p.).			
Peak No.	Angle $2\theta$	d Value	Rel. Intensity
1	10.1°	8.7	0.0
2	11.0°	8.0	0.0
3	11.8°	7.5	0.1
4	13.1°	6.8	0.2
5	14.8°	6.0	0.0
6	15.2°	5.8	0.3
7	15.5°	5.7	0.1
8	16.5°	5.4	0.1
9	17.8°	5.0	0.4
10	18.3°	4.9	0.7
11	18.6°	4.8	0.7
12	18.8°	4.7	0.7
13	19.4°	4.6	0.0

42

TABLE 13b-continued

XRPD Peak data for Tartrate pattern 1 (1 d.p.).			
Peak No.	Angle $2\theta$	d Value	Rel. Intensity
14	19.6°	4.5	0.2
15	20.3°	4.4	0.7
16	20.7°	4.3	1.0
17	21.5°	4.1	0.2
18	22.1°	4.0	0.5
19	22.6°	3.9	0.2
20	23.0°	3.9	0.1
21	23.7°	3.8	0.3
22	24.9°	3.6	0.1
23	25.4°	3.5	0.1
24	26.0°	3.4	0.7
25	26.4°	3.4	0.0
26	27.0°	3.3	0.1
27	27.3°	3.3	0.1
28	27.9°	3.2	0.1
29	28.9°	3.1	0.0
30	29.3°	3.0	0.1
31	30.0°	3.0	0.1
32	31.2°	2.9	0.1
33	31.7°	2.8	0.0

25 The TGA thermogram of Tartrate salt showed 1% of weight loss between 25-170° C. (~0.05 moles THF) and good thermal stability up to around 170° C. as displayed in FIG. 35.

30 The 1<sup>st</sup> heating cycle DSC thermogram of Tartrate salt displayed a single endothermic event with onset temperature around 138.9° C. and heat of fusion 97.0 J/g as shown in, or substantially as shown in, FIG. 36 which correspond to the melting of the Tartrate salt.

35 The cooling ramp from 200° C. to -90° C. at 10° C./min showed a vitrification at around 49.0° C. and the 2<sup>nd</sup> heating cycle displayed a glass transition of 54.2° C. as shown in, or substantially as shown in, FIG. 37.

40 <sup>1</sup>H NMR spectrum of Tartrate salt in d<sub>6</sub>-DMSO presented in FIG. 38, displayed around 0.9 eq of tartaric acid present and traces of THF. The <sup>1</sup>H NMR result confirmed the formation of Tartrate salt. The total water uptake between 0% RH and 90% RH at 25° C. was observed to be approximately 3.3% w/w (moderately hygroscopic). Adsorption/ 45 desorption profiles are reversible and overlap, which indicates that sorption of moisture at higher humidity does not affect the internal structure of Tartrate salt. The XRPD post DVS analyses showed that Tartrate salt did not undergo any 50 solid form transformation when exposed to moisture and remained the same crystalline solid form as demonstrated in FIG. 41.

In one embodiment, there is provided 5-MeO-DMT tartrate. In one embodiment, there is provided a pharmaceutical 55 composition comprising 5-MeO-DMT tartrate. In one embodiment, there is provided crystalline 5-MeO-DMT tartrate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT tartrate, as characterised by one or 60 more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 34;

One or more, two or more, three or more, four or more, 65 five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen



43

or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, twenty nine or more, thirty or more, thirty one or more, thirty two or more, or thirty three peaks in an XRPD diffractogram as detailed in Table 13, Table 13a or Table 13b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 13, Table 13a or Table 13b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 35;

A weight loss of 1% between 25-170° C., as measured by TGA thermogram;

A weight loss of around 0.1-1.0% between 25-170° C., as measured by TGA thermogram;

A weight loss of around 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0% between 25-170° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 36 or FIG. 37;

A melting endothermic event with an onset of around 138.9° C. and an enthalpy of 97.0 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 130-145° C. and an enthalpy of around 92-102 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 130, 131, 132, 133, 134, 135, 136, 137, 138, 139, 140, 141, 142, 143, 144 or 145° C. and an enthalpy of around 92, 93, 94, 95, 96, 97, 98, 99, 100, 101 or 102 J/g, as measured in a DSC thermogram;

A vitrification around 49° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A vitrification around 45-55° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A vitrification around 45, 46, 47, 48, 49, 50, 51, 52, 53, 54 or 55° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A glass transition around 54.2° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A glass transition around 50-60° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A glass transition around 50, 51, 52, 53, 54, 55, 56, 57, 58, 59 or 60° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;


A total water uptake between 0% RH and 90% RH at 25° C. of approximately 3.3% w/w;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 38; and/or

An XRPD pattern as shown in, or substantially as shown in, FIG. 41.

44

Benzenesulfonate Salt



The XRPD pattern of Benzenesulfonate salt is shown in FIG. 42. It displayed peaks at different 2 theta values when compared to Benzenesulfonic acid and Free Base, thus confirming the salt formation. This was nominated as pattern 1 with XRPD peak data shown in Table 14, Table 14a or Table 14b.

TABLE 14

XRPD Peak data for Benzenesulfonate pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	8.579°	10.298	0.006
2	9.490°	9.312	1.000
3	9.738°	9.076	0.051
4	12.189°	7.255	0.006
5	13.449°	6.578	0.064
6	14.164°	6.248	0.260
7	14.703°	6.020	0.225
8	14.997°	5.903	0.315
9	16.791°	5.276	0.105
10	17.770°	4.987	0.213
11	18.035°	4.915	0.393
12	18.506°	4.791	0.025
13	18.872°	4.698	0.266
14	20.035°	4.428	0.024
15	20.332°	4.364	0.092
16	21.215°	4.185	0.535
17	21.954°	4.045	0.014
18	22.526°	3.944	0.025
19	22.964°	3.870	0.023
20	23.338°	3.809	0.175
21	23.571°	3.771	0.704
22	23.911°	3.719	0.039
23	24.435°	3.640	0.389
24	24.655°	3.608	0.331
25	25.050°	3.552	0.188
26	26.152°	3.405	0.104
27	26.536°	3.356	0.083
28	26.962°	3.304	0.015
29	27.316°	3.262	0.017
30	28.333°	3.147	0.295
31	28.900°	3.087	0.010
32	29.249°	3.051	0.076
33	29.621°	3.013	0.007
34	30.064°	2.970	0.013
35	30.564°	2.923	0.011
36	31.017°	2.881	0.010
37	31.901°	2.803	0.054

45

TABLE 14a

XRPD Peak data for Benzenesulfonate pattern 1 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.58°	10.30	0.01
2	9.49°	9.31	1.00
3	9.74°	9.08	0.05
4	12.19°	7.26	0.01
5	13.45°	6.58	0.06
6	14.16°	6.25	0.26
7	14.70°	6.02	0.23
8	15.00°	5.90	0.32
9	16.79°	5.28	0.11
10	17.77°	4.99	0.21
11	18.04°	4.92	0.39
12	18.51°	4.79	0.03
13	18.87°	4.70	0.27
14	20.04°	4.43	0.02
15	20.33°	4.36	0.09
16	21.22°	4.19	0.54
17	21.95°	4.05	0.01
18	22.53°	3.94	0.03
19	22.96°	3.87	0.02
20	23.34°	3.81	0.18
21	23.57°	3.77	0.70
22	23.91°	3.72	0.04
23	24.44°	3.64	0.39
24	24.66°	3.61	0.33
25	25.05°	3.55	0.19
26	26.15°	3.41	0.10
27	26.54°	3.36	0.08
28	26.96°	3.30	0.02
29	27.32°	3.26	0.02
30	28.33°	3.15	0.30
31	28.90°	3.09	0.01
32	29.25°	3.05	0.08
33	29.62°	3.01	0.01
34	30.06°	2.97	0.01
35	30.56°	2.92	0.01
36	31.02°	2.88	0.01
37	31.90°	2.80	0.05

TABLE 14b

XRPD Peak data for Benzenesulfonate pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.6°	10.3	0.0
2	9.5°	9.3	1.0
3	9.7°	9.1	0.1
4	12.2°	7.3	0.0
5	13.4°	6.6	0.1
6	14.2°	6.2	0.3
7	14.7°	6.0	0.2
8	15.0°	5.9	0.3
9	16.8°	5.3	0.1
10	17.8°	5.0	0.2
11	18.0°	4.9	0.4
12	18.5°	4.8	0.0
13	18.9°	4.7	0.3
14	20.0°	4.4	0.0
15	20.3°	4.4	0.1
16	21.2°	4.2	0.5
17	22.0°	4.0	0.0
18	22.5°	3.9	0.0
19	23.0°	3.9	0.0
20	23.3°	3.8	0.2
21	23.6°	3.8	0.7
22	23.9°	3.7	0.0
23	24.4°	3.6	0.4
24	24.7°	3.6	0.3
25	25.1°	3.6	0.2

46

TABLE 14b-continued

XRPD Peak data for Benzenesulfonate pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
26	26.2°	3.4	0.1
27	26.5°	3.4	0.1
28	27.0°	3.3	0.0
29	27.3°	3.3	0.0
30	28.3°	3.1	0.3
31	28.9°	3.1	0.0
32	29.2°	3.1	0.1
33	29.6°	3.0	0.0
34	30.1°	3.0	0.0
35	30.6°	2.9	0.0
36	31.0°	2.9	0.0
37	31.9°	2.8	0.1

The TGA thermogram of Benzenesulfonate salt is shown in FIG. 43. It showed a good thermal stability up to 250° C. The observed weight loss of 1.5% between 25-250° C. corresponding to around 0.1 moles of IPA.

The DSC analysis of the Benzenesulfonate salt was performed. The 1<sup>st</sup> heating thermogram in FIG. 44 displayed a broad endothermic event with  $T_{onset}$  around 76.2° C. and heat of fusion 66.5 J/g due to the melting of the Benzenesulfonate salt. A weak exothermic event around 140° C. was also observed.

The cooling ramp and the 2<sup>nd</sup> heating DSC thermograms in FIG. 45 displayed a vitrification around 17.7° C. and glass transition around 23.3° C., respectively.

<sup>1</sup>H NMR spectrum of Benzenesulfonate salt in d<sub>6</sub>-DMSO shown in FIG. 46 confirmed presence of 1.0 eq of benzene sulfonic acid. Residual traces of IPAC were also observed in the spectrum.

The FIG. 47 shows the DVS isotherm plot for Benzenesulfonate salt. The first sorption isotherm showed hysteresis between 40-80% RH. Firstly, a slight water uptake up to 60% RH (1.2% w/w) was observed. When the salt was exposed to 70% RH and 80% RH it exhibited moisture absorption of 7% and 14% w/w, respectively and then the salt begins to deliquesce. The total moisture uptake between 0% RH and 90% RH at 25° C. was observed to be approximately 21% w/w in both cycles.

The DVS kinetic plot of Benzenesulfonate salt is presented in FIG. 48. Due to deliquescence of Benzenesulfonate salt during the DVS experiment, XRPD analyses were not performed on post DVS sample.

In one embodiment, there is provided 5-MeO-DMT benzenesulfonate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT benzenesulfonate. In one embodiment, there is provided crystalline 5-MeO-DMT benzenesulfonate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT benzenesulfonate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 42;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, twenty nine or more, thirty or more, thirty one or more,

47

thirty two or more, thirty three or more, thirty four or more, thirty five or more, thirty six or more, or thirty seven peaks in an XRPD diffractogram as detailed in Table 14, Table 14a or Table 14b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 14, Table 14a or Table 14b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 43;

A weight loss of 1.5% between 25-250° C., as measured by TGA thermogram;

A weight loss of around 1.0-2.0% between 25-250° C., as measured by TGA thermogram;

A weight loss of around 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8, 1.9 or 2.0% between 25-250° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 44 or FIG. 45;

A melting endothermic event with an onset of around 76.2° C. and an enthalpy of 66.5 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 70-80° C. and an enthalpy of around 60-70 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 or 80° C. and an enthalpy of around 60, 61, 62, 63, 64, 65, 66, 67, 68, 69 or 70 J/g, as measured in a DSC thermogram;

A weak exothermic event at around 140° C. as measured in a DSC thermogram;

A weak exothermic event at around 135-145° C. as measured in a DSC thermogram;

A weak exothermic event at around 135, 136, 137, 138, 139, 140, 141, 142, 143, 144 or 145° C. as measured in a DSC thermogram;

A vitrification around 17.7° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A vitrification around 12-22° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A vitrification around 12, 13, 14, 15, 16, 17, 18, 19, 20, 21 or 22° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A glass transition around 23.3° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A glass transition around 18-28° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A glass transition around 18, 19, 20, 21, 22, 23, 24, 25, 26, 27 or 28° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;


A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 46;

A DVS isotherm plot as shown in, or substantially as shown in, FIG. 47; and/or

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 48.

48

Tosylate Salt



As shown in, or substantially as shown in, FIG. 49, the crystalline form of Tosylate salt has a distinctively different XRPD pattern when compared to the free base and p-toluenesulfonic acid confirming the salt formation. This was nominated as pattern 1 with XRPD peak data presented in Table 15, Table 15a or Table 15b.

TABLE 15

XRPD Peak data for Tosylate pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	9.609°	9.197	0.296
2	12.001°	7.369	0.041
3	12.934°	6.839	0.007
4	13.798°	6.413	0.386
5	14.391°	6.150	0.315
6	16.105°	5.499	0.034
7	16.625°	5.328	0.138
8	17.489°	5.067	0.157
9	18.114°	4.893	0.142
10	19.342°	4.585	0.622
11	19.945°	4.448	0.119
12	22.076°	4.023	0.034
13	23.161°	3.837	0.057
14	23.571°	3.771	0.472
15	24.054°	3.697	1.000
16	25.037°	3.554	0.130
17	25.979°	3.427	0.023
18	26.435°	3.369	0.107
19	27.302°	3.264	0.420
20	27.712°	3.216	0.097
21	28.880°	3.089	0.123
22	29.371°	3.039	0.013
23	29.769°	2.999	0.026
24	30.161°	2.961	0.046
25	30.981°	2.884	0.032
26	31.738°	2.817	0.048

TABLE 15a

XRPD Peak data for Tosylate pattern 1 (2 d.p.).			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	9.61°	9.20	0.30
2	12.00°	7.37	0.04
3	12.93°	6.84	0.01
4	13.80°	6.41	0.39
5	14.39°	6.15	0.32
6	16.11°	5.50	0.03
7	16.63°	5.33	0.14
8	17.49°	5.07	0.16
9	18.11°	4.89	0.14
10	19.34°	4.59	0.62
11	19.95°	4.45	0.12

TABLE 15a-continued

XRPD Peak data for Tosylate pattern 1 (2 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
12	22.08°	4.02	0.03
13	23.16°	3.84	0.06
14	23.57°	3.77	0.47
15	24.05°	3.70	1.00
16	25.04°	3.55	0.13
17	25.98°	3.43	0.02
18	26.44°	3.37	0.11
19	27.30°	3.26	0.42
20	27.71°	3.22	0.10
21	28.88°	3.09	0.12
22	29.37°	3.04	0.01
23	29.77°	3.00	0.03
24	30.16°	2.96	0.05
25	30.98°	2.88	0.03
26	31.74°	2.82	0.05

TABLE 15b

XRPD Peak data for Tosylate pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.6°	9.2	0.3
2	12.0°	7.4	0.0
3	12.9°	6.8	0.0
4	13.8°	6.4	0.4
5	14.4°	6.2	0.3
6	16.1°	5.5	0.0
7	16.6°	5.3	0.1
8	17.5°	5.1	0.2
9	18.1°	4.9	0.1
10	19.3°	4.6	0.6
11	19.9°	4.4	0.1
12	22.1°	4.0	0.0
13	23.2°	3.8	0.1
14	23.6°	3.8	0.5
15	24.1°	3.7	1.0
16	25.0°	3.6	0.1
17	26.0°	3.4	0.0
18	26.4°	3.4	0.1
19	27.3°	3.3	0.4
20	27.7°	3.2	0.1
21	28.9°	3.1	0.1
22	29.4°	3.0	0.0
23	29.8°	3.0	0.0
24	30.2°	3.0	0.0
25	31.0°	2.9	0.0
26	31.7°	2.8	0.0

The TGA thermogram of Tosylate salt displayed a weight loss of 1.0% between 25-230° C. (0.1 moles IPA) followed by the thermal degradation as shown in, or substantially as shown in, FIG. 50.

The 1<sup>st</sup> heating DSC thermogram of Tosylate salt presented in FIG. 51, exhibited a single endothermic event with the onset temperature of 109.7° C. and heat of fusion of 89.3 J/g corresponding to the melting.

As presented in FIG. 52, the DSC thermograms of Tosylate salt upon cooling and 2<sup>nd</sup> heating displayed a vitrification and a glass transition around 24.3 and 30.2° C., respectively.

<sup>1</sup>H NMR spectrum in d<sub>6</sub>-DMSO of Tosylate salt presented in FIG. 53 displayed the proton signals corresponding to the p-toluene sulfonic acid which were integrated as 1.0 eq. Traces of IPA were detected in the spectrum.

Evaluation of the DVS results obtained for Tosylate salt shows that the material is slightly hygroscopic with a water

uptake of approximately 1.9% w/w between 0% RH and 90% RH at 25° C. The material takes up moisture reversibly without hysteresis which indicates that sorption at higher humidity does not affect the internal structure of the Tartrate salt as shown in, or substantially as shown in, FIG. 54.

The DVS kinetic plot of Tosylate salt is displayed in FIG. 55.

It can be seen from the XRPD diffractogram in FIG. 56 that the Tosylate salt did not undergo any solid-state changes during the DVS experiment as the XRPD pattern of post DVS sample remained unchanged.

In one embodiment, there is provided 5-MeO-DMT tosylate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT tosylate. In one embodiment, there is provided crystalline 5-MeO-DMT tosylate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT tosylate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 49 or 56;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, or twenty six peaks in an XRPD diffractogram as detailed in Table 15, Table 15a or Table 15b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 15, Table 15a or Table 15b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 50;

A weight loss of 1.0% between 25-230° C., as measured by TGA thermogram;

A weight loss of around 0.5-1.5% between 25-230° C., as measured by TGA thermogram;

A weight loss of around 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4 or 1.5% between 25-230° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 52;

A melting endothermic event with an onset of around 109.7° C. and an enthalpy of 89.3 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 105-115° C. and an enthalpy of around 85-95 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 105, 106, 107, 108, 109, 110, 111, 112, 113, 114 or 115° C. and an enthalpy of around 85, 86, 87, 88, 89, 90, 91, 92, 93, 94 or 95 J/g, as measured in a DSC thermogram;

A vitrification around 24.3° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A vitrification around 20-30° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A vitrification around 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

## 51

A glass transition around 30.2° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A glass transition around 25-35° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A glass transition around 25, 26, 27, 28, 29, 30, 31, 32, 33, 34 or 35° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 230° C. to -90° C.;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 53;

A total water uptake between 0% RH and 90% RH at 25° C. of approximately 1.9% w/w;


A total water uptake between 0% RH and 90% RH at 25° C. of approximately 1.5-2.5% w/w;

A total water uptake between 0% RH and 90% RH at 25° C. of approximately 1.5, 1.6, 1.7, 1.8, 1.9, 2.0, 2.1, 2.2, 2.3, 2.4 or 2.5% w/w;

A DVS isotherm plot as shown in, or substantially as shown in, FIG. 54; and/or

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 55.

Hydrobromide Salt



XRPD diffractograms of Hydrobromide salt displayed same crystalline patterns for all three solvent/anti-solvent systems used as demonstrated in FIG. 57. The XRPD diffractograms of Hydrobromide salt showed a distinct diffraction profile when compared to free base, confirming the salt formation. This was nominated as pattern 1 with XRPD peak data shown in Tables 16, 16a or 16b.

TABLE 16

XRPD Peak data for Hydrobromide pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	6.202°	14.238	0.187
2	9.099°	9.711	0.022
3	12.115°	7.299	0.004
4	12.359°	7.156	0.192
5	13.322°	6.641	0.002
6	14.046°	6.300	0.229
7	14.667°	6.035	0.104
8	15.219°	5.817	0.007
9	16.722°	5.297	0.002
10	18.566°	4.775	1.000
11	18.780°	4.721	0.030
12	19.740°	4.494	0.361
13	20.937°	4.239	0.026
14	22.900°	3.880	0.026
15	23.540°	3.776	0.106
16	24.169°	3.679	0.016
17	24.824°	3.584	0.299
18	25.714°	3.462	0.005
19	26.109°	3.410	0.014

## 52

TABLE 16-continued

XRPD Peak data for Hydrobromide pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
20	26.746°	3.331	0.007
21	27.427°	3.249	0.042
22	28.245°	3.157	0.036
23	28.771°	3.101	0.002
24	30.020°	2.974	0.004
25	30.556°	2.923	0.006
26	31.132°	2.871	0.020

TABLE 16a

XRPD Peak data for Hydrobromide pattern 1. (2 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	6.20°	14.24	0.19
2	9.10°	9.71	0.02
3	12.12°	7.30	0.00
4	12.36°	7.16	0.19
5	13.32°	6.64	0.00
6	14.05°	6.30	0.23
7	14.67°	6.04	0.10
8	15.22°	5.82	0.01
9	16.72°	5.30	0.00
10	18.57°	4.78	1.00
11	18.78°	4.72	0.03
12	19.74°	4.49	0.36
13	20.94°	4.24	0.03
14	22.90°	3.88	0.03
15	23.54°	3.78	0.11
16	24.17°	3.68	0.02
17	24.82°	3.58	0.30
18	25.71°	3.46	0.01
19	26.11°	3.41	0.01
20	26.75°	3.33	0.01
21	27.43°	3.25	0.04
22	28.25°	3.16	0.04
23	28.77°	3.10	0.00
24	30.02°	2.97	0.00
25	30.56°	2.92	0.01
26	31.13°	2.87	0.02

TABLE 16b

XRPD Peak data for Hydrobromide pattern 1 (1 d.p.).			
Peak No.	Angle 2 θ	d Value	Rel. intensity
1	6.2°	14.2	0.2
2	9.1°	9.7	0.0
3	12.1°	7.3	0.0
4	12.4°	7.2	0.2
5	13.3°	6.6	0.0
6	14.0°	6.3	0.2
7	14.7°	6.0	0.1
8	15.2°	5.8	0.0
9	16.7°	5.3	0.0
10	18.6°	4.8	1.0
11	18.8°	4.7	0.0
12	19.7°	4.5	0.4
13	20.9°	4.2	0.0
14	22.9°	3.9	0.0
15	23.5°	3.8	0.1
16	24.2°	3.7	0.0
17	24.8°	3.6	0.3
18	25.7°	3.5	0.0
19	26.1°	3.4	0.0

53

TABLE 16b-continued

XRPD Peak data for Hydrobromide pattern 1 (1 d.p.).			
Peak No.	Angle 2 $\theta$	d Value	Rel. intensity
20	26.7°	3.3	0.0
21	27.4°	3.7	0.0
22	28.2°	3.2	0.0
23	28.8°	3.1	0.0
24	30.0°	3.0	0.0
25	30.6°	2.9	0.0
26	31.1°	2.9	0.0

The TGA thermogram of Hydrobromide salt in FIG. 58 showed that the material is thermally stable up to 220° C., then the thermal degradation occurs. 0.45% weight loss between 25-220° C. is due to release of volatiles.

The DSC analysis of the Hydrobromide salt was performed and thermograms are presented in FIG. 59. The 1st heating DSC thermogram shows a single endothermic event with the onset temperature of 148.7° C. and heat of fusion of 97.3 J/g, which corresponds to the melting of the Hydrobromide salt.

The cooling ramp of 10° C./min from 220° C. to -90° C. displayed a vitrification at around 34.8° C. and the 2<sup>nd</sup> heating cycle a glass transition at around 42.2° C. as shown in, or substantially as shown in, FIG. 60.

<sup>1</sup>H NMR spectrum of Hydrobromide salt in d<sub>6</sub>-DMSO is shown in FIG. 61. It displayed traces of MTBE. The observed proton chemical shift changes in the NMR spectrum are indicative of the salt formation.

In one embodiment, there is provided 5-MeO-DMT hydrobromide. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT hydrobromide. In one embodiment, there is provided crystalline 5-MeO-DMT hydrobromide, or a pharmaceutical composition comprising crystalline 5-MeO-DMT hydrobromide, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 57;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, or twenty six peaks in an XRPD diffractogram as detailed in Table 16, Table 16a or Table 16b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 16, Table 16a or Table 16b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 58;

A weight loss of about 0.45% between 25-220° C., as measured by TGA thermogram;

A weight loss of about 0.35-0.55% between 25-220° C., as measured by TGA thermogram;

A weight loss of about 0.35, 0.36, 0.37, 0.38, 0.39, 0.40, 0.41, 0.42, 0.43, 0.44, 0.45, 0.46, 0.47, 0.48, 0.49, 0.50, 0.51, 0.52, 0.53, 0.54 or 0.55% between 25-220° C., as measured by TGA thermogram;

A weight loss of about 0.1-1.0% between 25-220° C., as measured by TGA thermogram;

54

A weight loss of about 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9 or 1.0% between 25-220° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 59 or FIG. 60;

A melting endothermic event with an onset of around 148.7° C. and an enthalpy of 97.3 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 143-153° C. and an enthalpy of around 92-102 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 143, 144, 145, 146, 147, 148, 149, 150, 151, 152 or 153 and an enthalpy of around 92, 93, 94, 95, 96, 97, 98, 99, 100, 101, or 102 J/g, as measured in a DSC thermogram;

A vitrification around 34.8° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A vitrification around 30-40° C. as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A vitrification around 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, or 40° C. as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;


A glass transition around 42.2° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A glass transition around 37-47° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A glass transition around 37, 38, 39, 40, 41, 42, 43, 44, 45, 46 or 47° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.; and/or

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 61.

Glycolate Salt



The XRPD diffractogram of the glycolate salt exhibited a crystalline form which has a different XRPD pattern when compared to the free base and glycolic acid as shown in, or substantially as shown in, FIG. 62. This form was nominated as pattern 1 with XRPD peak data presented in Table 17, Table 17a or Table 17b.

TABLE 17

XRPD Peak data for glycolate pattern 1.			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.837°	9.999	0.004
2	10.114°	8.739	0.551

55

TABLE 17-continued

XRPD Peak data for glycolate pattern 1.			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
3	11.674°	7.574	0.255
4	15.436°	5.736	0.063
5	16.557°	5.350	0.011
6	17.187°	5.155	0.107
7	17.923°	4.945	0.047
8	19.656°	4.513	0.025
9	20.235°	4.385	1.000
10	21.085°	4.210	0.624
11	21.908°	4.054	0.034
12	22.401°	3.966	0.088
13	22.979°	3.867	0.073
14	23.408°	3.797	0.923
15	24.340°	3.654	0.364
16	24.944°	3.567	0.011
17	25.495°	3.491	0.081
18	26.873°	3.315	0.079
19	27.449°	3.247	0.169
20	27.781°	3.209	0.012
21	28.878°	3.089	0.015
22	29.704°	3.005	0.005
23	30.565°	2.922	0.267
24	31.124°	2.871	0.113
25	31.793°	2.812	0.008

TABLE 17a

XRPD Peak data for glycolate pattern 1. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.84°	10.00	0.00
2	10.11°	8.74	0.55
3	11.67°	7.57	0.26
4	15.44°	5.74	0.06
5	16.56°	5.35	0.01
6	17.19°	5.16	0.11
7	17.92°	4.95	0.05
8	19.66°	4.51	0.03
9	20.24°	4.39	1.00
10	21.09°	4.21	0.62
11	21.91°	4.05	0.03
12	22.40°	3.97	0.09
13	22.98°	3.87	0.07
14	23.41°	3.80	0.92
15	24.34°	3.65	0.36
16	24.94°	3.57	0.01
17	25.50°	3.49	0.08
18	26.87°	3.32	0.08
19	27.45°	3.25	0.17
20	27.78°	3.21	0.01
21	28.88°	3.09	0.02
22	29.70°	3.01	0.01
23	30.57°	2.92	0.27
24	31.12°	2.87	0.11
25	31.79°	2.81	0.01

TABLE 17b

XRPD Peak data for glycolate pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.8°	10.0	0.0
2	10.1°	8.7	0.6
3	11.7°	7.6	0.3
4	15.4°	5.7	0.1
5	16.6°	5.4	0.0
6	17.2°	5.2	0.1

56

TABLE 17b-continued

XRPD Peak data for glycolate pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
7	17.9°	4.9	0.0
8	19.7°	4.5	0.0
9	20.2°	4.4	1.0
10	21.1°	4.2	0.6
11	21.9°	4.1	0.0
12	22.4°	4.0	0.1
13	23.0°	3.9	0.1
14	23.4°	3.8	0.9
15	24.3°	3.7	0.4
16	24.9°	3.6	0.0
17	25.5°	3.5	0.1
18	26.9°	3.3	0.1
19	27.4°	3.2	0.2
20	27.8°	3.2	0.0
21	28.9°	3.1	0.0
22	29.7°	3.0	0.0
23	30.6°	2.9	0.3
24	31.1°	2.9	0.1
25	31.8°	2.8	0.0

The TGA thermogram of glycolate salt showed a weight loss of 1.4% between 25-155° C. (~0.07 moles IPAC). The material is thermally stable up to around 155° C. as demonstrated in FIG. 63. The DSC analysis of the glycolate salt was performed. The 1<sup>st</sup> heating DSC thermogram displayed a melting endotherm with T<sub>onset</sub> around 95.2° C. and heat of fusion of 100.5 J/g, followed by the thermal degradation at higher temperature as shown in, or substantially as shown in, FIG. 64.

The cooling ramp from 200° C. to -90° C. displayed a vitrification around 7.5° C. and the 2<sup>nd</sup> heating cycle showed a glass transition around 14.5° C. as presented in FIG. 65.

<sup>1</sup>H NMR spectrum in d<sub>6</sub>-DMSO of glycolate salt shown in FIG. 66, confirmed the presence of 1.0 eq glycolic acid. Traces of isopropyl acetate were also observed in the spectrum.

The FIG. 67 shows the DVS isotherm plot for glycolate salt. The first sorption isotherm showed hysteresis between 40-80% RH. Firstly, a slight water uptake up to 60% RH (2.3% w/w) was observed.

When the salt was exposed to 70% RH and 80% RH it exhibited moisture absorption 11% and 33% w/w respectively and the glycolate salt begins to deliquesce. The total moisture uptake between 0% RH and 90% RH at 25° C. was observed to be approximately 62% w/w for 1<sup>st</sup> cycle and 59% w/w for 2<sup>nd</sup> cycle.

As the glycolate salt underwent deliquescence during the DVS experiment, XRPD analyses were not carry out on post DVS sample.

In one embodiment, there is provided 5-MeO-DMT glycolate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT glycolate. In one embodiment, there is provided crystalline 5-MeO-DMT glycolate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT glycolate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 62;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or

57

more, twenty two or more, twenty three or more, twenty four or more, or twenty five peaks in an XRPD diffractogram as detailed in Table 17, Table 17a or Table 17b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 17, Table 17a or Table 17b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 63;

A weight loss of about 1.4% between 25-155° C., as measured by TGA thermogram;

A weight loss of about 0.9-1.9% between 25-155° C., as measured by TGA thermogram;

A weight loss of about 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6, 1.7, 1.8 or 1.9% between 25-155° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 64 or FIG. 65;

A melting endothermic event with an onset of around 95.2° C. and an enthalpy of 100.5 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 90-100° C. and an enthalpy of around 95-105 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100° C. and an enthalpy of around 95, 96, 97, 98, 99, 100, 101, 102, 103, 104 or 105 J/g, as measured in a DSC thermogram;

A vitrification around 7.5° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A vitrification around 2-12° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A vitrification around 2, 3, 4, 5, 6, 7, 8, 9, 10, 11 or 12° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A glass transition around 14.5° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A glass transition around 10-20° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;


A glass transition around 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 200° C. to -90° C.;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 66;

A DVS isotherm plot as shown in, or substantially as shown in, FIG. 67; and/or

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 68.

Ketoglutarate Salt



58

Both samples isolated from different solvent/anti-solvent systems displayed the same XRPD crystalline pattern which is distinctively different when compared to free base and ketoglutaric acid confirming the salt formation as shown in, or substantially as shown in, FIG. 69. This was nominated as pattern 1 with XRPD peak data displayed in Table 18, Table 18a or Table 18b.

TABLE 18

XRPD Peak data for Ketoglutarate pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	9.964°	8.870	0.098
2	11.503°	7.687	0.056
3	14.386°	6.152	0.348
4	15.796°	5.606	0.020
5	17.613°	5.031	0.116
6	18.157°	4.882	1.000
7	18.922°	4.686	0.185
8	19.940°	4.449	0.053
9	20.460°	4.337	0.039
10	20.914°	4.244	0.399
11	21.420°	4.145	0.057
12	22.517°	3.946	0.263
13	23.165°	3.836	0.208
14	23.861°	3.726	0.202
15	24.534°	3.625	0.010
16	25.616°	3.475	0.337
17	26.071°	3.415	0.083
18	26.670°	3.340	0.130
19	27.361°	3.257	0.091
20	27.700°	3.218	0.115
21	28.957°	3.081	0.016
22	29.518°	3.024	0.026
23	30.025°	2.974	0.042
24	30.580°	2.921	0.035

TABLE 18a

XRPD Peak data for Ketoglutarate pattern 1. (2 d p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	9.96°	8.87	0.10
2	11.50°	7.69	0.06
3	14.39°	6.15	0.35
4	15.80°	5.61	0.02
5	17.61°	5.03	0.12
6	18.16°	4.83	1.00
7	18.92°	4.69	0.19
8	19.94°	4.45	0.05
9	20.46°	4.34	0.04
10	20.91°	4.24	0.40
11	21.42°	4.15	0.06
12	22.52°	3.95	0.26
13	23.17°	3.84	0.21
14	23.86°	3.73	0.20
15	24.53°	3.63	0.01
16	25.62°	3.48	0.34
17	26.07°	3.42	0.08
18	26.67°	3.34	0.13
19	27.36°	3.26	0.09
20	27.70°	3.22	0.12
21	28.96°	3.08	0.02
22	29.52°	3.02	0.03
23	30.03°	2.97	0.04
24	30.58°	2.92	0.04



TABLE 18b

XRPD Peak data for Ketoglutarate pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	10.0°	8.9	0.1
2	11.5°	7.7	0.1
3	14.4°	6.2	0.3
4	15.8°	5.6	0.0
5	17.6°	5.0	0.1
6	18.2°	4.9	1.0
7	13.9°	4.7	0.2
8	19.9°	4.4	0.1
9	20.5°	4.3	0.0
10	20.9°	4.2	0.4
11	21.4°	4.1	0.1
12	22.5°	3.9	0.3
13	23.2°	3.8	0.2
14	23.9°	3.7	0.2
15	24.5°	3.6	0.0
16	25.6°	3.5	0.3
17	26.1°	3.4	0.1
18	26.7°	3.3	0.1
19	27.4°	3.3	0.1
20	27.7°	3.2	0.1
21	29.0°	3.1	0.0
22	29.5°	3.0	0.0
23	30.0°	3.0	0.0
24	30.6°	2.9	0.0

The TGA thermogram of Ketoglutarate salt displayed a weight loss of 1.2% between 25-150° C., which corresponds to ~0.1 moles EtOH and 0.05 moles MTBE. The Ketoglutaric salt is thermally stable up to 150° C. as shown in, or substantially as shown in, FIG. 70.

The DSC analysis of the ketoglutaric salt was performed and the results displayed in FIG. 71. The 1<sup>st</sup> heating DSC thermogram of Ketoglutarate salt exhibited a single endothermic event with the onset temperature of 85.5° C. and heat of fusion of 92.4 J/g, which corresponds to the melting of the material.

During cooling from 150° C. to -90° C./10 min a vitrification around 21.9° C. was observed and the 2<sup>nd</sup> heating cycle displayed a glass transition of 28.5° C. as shown in, or substantially as shown in, FIG. 72.

<sup>1</sup>H NMR spectrum of Ketoglutarate salt in d<sub>6</sub>-DMSO presented in FIG. 73, showed all associated peaks with approximately 1.0 eq of ketoglutaric acid being present. The spectrum also displayed 0.13 eq EtOH and 0.1 eq MTBE in the sample.

The isotherm plot of Ketoglutarate salt is presented in FIG. 74. The first sorption isotherm displayed a gradual water uptake between 40% RH (0.13% w/w) to 80% RH (2.55% w/w). A rapid increase to 28.14% w/w was observed at 90% RH. This indicates that Ketoglutarate salt deliquesce at high RH. The sorption kinetic plot of Ketoglutarate salt is presented in FIG. 75.

Due to deliquescence of the Ketoglutarate salt during DVS analyses, the post DVS sample was not analysed by XRPD.

In one embodiment, there is provided 5-MeO-DMT ketoglutarate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT ketoglutarate. In one embodiment, there is provided crystalline 5-MeO-DMT ketoglutarate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT ketoglutarate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 69;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, or twenty four peaks in an XRPD diffractogram as detailed in Table 18, Table 18a or Table 18b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 18, Table 18a or Table 18b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 70;

A weight loss of about 1.2% between 25-150° C., as measured by TGA thermogram;

A weight loss of about 0.7-1.7% between 25-150° C., as measured by TGA thermogram;

A weight loss of about 0.7, 0.8, 0.9, 1.0, 1.1, 1.2, 1.3, 1.4, 1.5, 1.6 or 1.7% between 25-150° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 71 or FIG. 72;

A melting endothermic event with an onset of around 85.5° C. and an enthalpy of 92.4 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 80-90° C. and an enthalpy of around 87-97 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 80, 81, 82, 83, 84, 85, 86, 87, 88, 89 or 90 C and an enthalpy of around 87, 88, 89, 90, 91, 92, 93, 94, 95, 96 or 97 J/g, as measured in a DSC thermogram;

A vitrification around 21.9° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 150° C. to -90° C.;

A vitrification around 16-26° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 150° C. to -90° C.;

A vitrification around 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or 26° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 150° C. to -90° C.;

A glass transition around 28.5° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 150° C. to -90° C.;

A glass transition around 23-33° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 150° C. to -90° C.;

A glass transition around 23, 24, 25, 26, 27, 28, 29, 30, 31, 32 or 33° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 150° C. to -90° C.;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 73;

A gradual water uptake between 40% RH (0.13% w/w) to 80% RH (2.55% w/w), optionally with a rapid increase to 28.14% w/w at 90% RH;


A gradual water uptake between 40% RH (0.05-0.2% w/w) to 80% RH (1.50-3.5% w/w), optionally with a rapid increase to 20-40% w/w at 90% RH;

A DVS isotherm plot as shown in, or substantially as shown in, FIG. 74; and/or

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 75.

61

Malate Salt



XRPD diffractograms showed that same crystalline forms were produced from ethanol/MTBE and IPAC/MTBE solvent/anti-solvent systems, which does not correspond to free base and/or L-malic acid as demonstrated in FIG. 76. This was nominated as pattern 1 with XRPD peak data presented in Table 19, 19a or 19b.

TABLE 19

XRPD Peak data for Malate pattern 1.			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	11.823°	7.479	0.136
2	14.733°	6.008	0.392
3	15.588°	5.680	0.058
4	18.254°	4.856	1.000
5	18.666°	4.750	0.746
6	18.889°	4.694	0.963
7	20.330°	4.365	0.035
8	20.782°	4.271	0.025
9	21.080°	4.211	0.052
10	21.608°	4.109	0.475
11	21.963°	4.044	0.335
12	23.014°	3.861	0.068
13	23.662°	3.757	0.456
14	24.119°	3.687	0.474
15	26.084°	3.413	0.505
16	26.608°	3.347	0.168
17	27.179°	3.278	0.131
18	27.839°	3.202	0.210
19	29.242°	3.052	0.011
20	29.649°	3.011	0.022
21	30.113°	2.965	0.113
22	30.700°	2.910	0.046
23	31.219°	2.863	0.024
24	31.821°	2.810	0.056

TABLE 19a

XRPD Peak data for Malate pattern 1. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	11.82°	7.48	0.14
2	14.73°	6.01	0.39
3	15.59°	5.68	0.06
4	18.25°	4.86	1.00
5	18.67°	4.75	0.75
6	18.89°	4.69	0.97
7	20.33°	4.37	0.04
8	20.78°	4.27	0.03
9	21.08°	4.21	0.05
10	21.61°	4.11	0.48
11	21.96°	4.04	0.34
12	23.01°	3.86	0.07
13	23.66°	3.76	0.46
14	24.12°	3.69	0.47
15	26.08°	3.41	0.51

62

TABLE 19a-continued

XRPD Peak data for Malate pattern 1. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
16	26.61°	3.35	0.17
17	27.18°	3.28	0.13
18	27.84°	3.20	0.21
19	29.24°	3.05	0.01
20	29.65°	3.01	0.02
21	30.11°	2.97	0.11
22	30.70°	2.91	0.05
23	31.22°	2.86	0.02
24	31.82°	2.81	0.06

TABLE 19b

XRPD Peak data for Malate pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	11.8°	7.5	0.1
2	14.7°	6.0	0.4
3	15.6°	5.7	0.1
4	13.3°	4.9	1.0
5	13.7°	4.8	0.7
6	18.9°	4.7	1.0
7	20.3°	4.4	0.0
8	20.8°	4.3	0.0
9	21.1°	4.2	0.1
10	21.6°	4.1	0.5
11	22.0°	4.0	0.3
12	23.0°	3.9	0.1
13	23.7°	3.8	0.5
14	24.1°	3.7	0.5
15	26.1°	3.4	0.5
16	26.6°	3.3	0.2
17	27.2°	3.3	0.1
18	27.8°	3.2	0.2
19	29.2°	3.1	0.0
20	29.6°	3.0	0.0
21	30.1°	3.0	0.1
22	30.7°	2.9	0.0
23	31.2°	2.9	0.0
24	31.8°	2.8	0.1

The TGA thermogram of Malate salt displayed a weight loss of 2.6% between ambient temperature and 170° C., due to loss of moisture from surface of particles. The material is thermally stable up to around 170° C. as shown in, or substantially as shown in, FIG. 77.

The DSC analysis of the Malate salt was performed. The 1<sup>st</sup> heating DSC thermogram of Malate salt exhibited a broad endothermic event with onset temperature of 80.9° C. and heat of fusion of 87.0 J/g, which corresponds to the melting of the salt as shown in, or substantially as shown in, FIG. 78.

The cooling ramp 10° C./min from 170 to -90° C. displayed vitrification around 15.1° C. and the 2<sup>nd</sup> heating cycle showed a glass transition around 22.0° C. as demonstrated in FIG. 79.

<sup>1</sup>H NMR spectrum of Malate salt in d<sub>6</sub>-DMSO is shown in FIG. 80. It displayed traces of ethanol (~0.1 eq) and MTBE (~0.08 eq). Signals corresponding to L-Malic acid were integrated as approximately 0.7 eq.

In one embodiment, there is provided 5-MeO-DMT malate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT malate. In one embodiment, there is provided crystalline 5-MeO-DMT malate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT malate, as characterised by one or more of:

63

An XRPD pattern as shown in, or substantially as shown in, FIG. 76;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, or twenty four peaks in an XRPD diffractogram as detailed in Table 19, Table 19a or Table 19b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 19, Table 19a or Table 19b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 77;

A weight loss of about 2.6% between ambient temperature and 170° C., as measured by TGA thermogram;

A weight loss of about 2.0-3.0% between ambient temperature and 170° C., as measured by TGA thermogram;

A weight loss of about 2.0, 2.1, 2.2, 2.3, 2.4, 2.5, 2.6, 2.7, 2.8, 2.9 or 3.0% between ambient temperature and 170° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 78 or FIG. 79;

A melting endothermic event with an onset of around 80.9° C. and an enthalpy of 87.0 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 75-85° C. and an enthalpy of around 82-92 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 75, 76, 77, 78, 79, 80, 81, 82, 83, 84 or 85° C. and an enthalpy of around 82, 83, 84, 85, 86, 87, 88, 89, 90, 91 or 92 J/g, as measured in a DSC thermogram;

A vitrification around 15.1° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 170° C. to -90° C.;

A vitrification around 10-20° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 170° C. to -90° C.;

A vitrification around 10, 11, 12, 13, 14, 15, 16, 17, 18, 19 or 20° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 170° C. to -90° C.;

A glass transition around 22° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 170° C. to -90° C.;


A glass transition around 17-27° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 170° C. to -90° C.;

A glass transition around 17, 18, 19, 20, 21, 22, 23, 24, 25, 26 or 27° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 170° C. to -90° C.; and/or

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 80.

64

Saccharinate Salt



As shown in, or substantially as shown in, FIG. 81 the crystalline form of Saccharinate salt exhibited a different XRPD pattern when compared to free base and saccharin, which suggests the salt formation. This was nominated as pattern 1 with XRPD peak data tabulated in Table 20, 20a or 20b.

TABLE 20

XRPD Peak data for Saccharinate pattern 1.			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	5.196°	16.993	0.314
2	9.572°	9.232	0.019
3	10.180°	8.682	0.928
4	11.395°	7.759	0.138
5	12.951°	6.830	0.051
6	13.423°	6.591	0.159
7	14.998°	5.902	0.362
8	15.182°	5.331	0.366
9	16.169°	5.477	0.228
10	16.806°	5.271	0.206
11	17.542°	5.052	0.295
12	17.845°	4.966	0.075
13	18.421°	4.813	0.027
14	18.736°	4.732	0.056
15	19.167°	4.627	0.037
16	19.672°	4.509	0.092
17	20.213°	4.390	0.034
18	20.868°	4.253	1.000
19	21.701°	4.092	0.113
20	22.138°	4.012	0.036
21	22.613°	3.929	0.069
22	22.987°	3.866	0.293
23	23.949°	3.713	0.025
24	24.396°	3.646	0.337
25	25.231°	3.527	0.076
26	25.463°	3.495	0.175
27	25.948°	3.431	0.084
28	26.616°	3.346	0.165
29	27.555°	3.234	0.247
30	23.286°	3.153	0.137
31	28.735°	3.104	0.039
32	29.676°	3.008	0.032
33	29.961°	2.980	0.037
34	30.413°	2.937	0.048
35	31.037°	2.879	0.040
36	31.781°	2.813	0.025

TABLE 20a

XRPD Peak data for Saccharinate pattern 1. (2 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	5.20°	16.99	0.31
2	9.57°	9.23	0.02
3	10.18°	8.68	0.93

65

TABLE 20a-continued

XRPD Peak data for Saccharinate pattern 1. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
4	11.40°	7.76	0.14
5	12.95°	6.83	0.05
6	13.42°	6.59	0.16
7	15.00°	5.90	0.36
8	15.13°	5.83	0.37
9	16.17°	5.48	0.23
10	16.81°	5.27	0.21
11	17.54°	5.05	0.30
12	17.85°	4.97	0.08
13	18.42°	4.81	0.03
14	18.74°	4.73	0.06
15	19.17°	4.63	0.04
16	19.67°	4.51	0.09
17	20.21°	4.39	0.08
18	20.87°	4.25	1.00
19	21.70°	4.09	0.11
20	22.14°	4.01	0.04
21	22.61°	3.93	0.07
22	22.99°	3.87	0.29
23	23.95°	3.71	0.03
24	24.40°	3.65	0.34
25	25.23°	3.53	0.08
26	25.46°	3.50	0.18
27	25.95°	3.43	0.08
28	26.62°	3.35	0.17
29	27.56°	3.23	0.25
30	28.29°	3.15	0.19
31	28.74°	3.10	0.04
32	29.68°	3.01	0.03
33	29.96°	2.98	0.04
34	30.41°	2.94	0.05
35	31.04°	2.88	0.04
36	31.78°	2.81	0.03

TABLE 20b

XRPD Peak data for Saccharinate pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	5.2°	17.0	0.3
2	9.6°	9.2	0.0
3	10.2°	8.7	0.9
4	11.4°	7.8	0.1
5	13.0°	6.8	0.1
6	13.4°	6.6	0.2
7	15.0°	5.9	0.4
8	15.2°	5.8	0.4
9	16.2°	5.5	0.2
10	16.8°	5.3	0.2
11	17.5°	5.1	0.3
12	17.8°	5.0	0.1
13	18.4°	4.8	0.0
14	18.7°	4.7	0.1
15	19.2°	4.6	0.0
16	15.7°	4.5	0.1
17	20.2°	4.4	0.1
18	20.9°	4.3	1.0
19	21.7°	4.1	0.1
20	22.1°	4.0	0.0
21	22.6°	3.9	0.1
22	23.0°	3.9	0.3
23	23.9°	3.7	0.0
24	24.4°	3.6	0.3
25	25.2°	3.5	0.1
26	25.5°	3.5	0.2
27	25.9°	3.4	0.1
28	26.6°	3.3	0.2
29	27.6°	3.2	0.2
30	28.3°	3.2	0.2
31	28.7°	3.1	0.0

66

TABLE 20b-continued

XRPD Peak data for Saccharinate pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
32	29.7°	3.0	0.0
33	30.0°	3.0	0.0
34	30.4°	2.9	0.0
35	31.0°	2.9	0.0
36	31.8°	2.8	0.0

The TGA thermogram of Saccharinate salt presented in FIG. 82, showed that the material is thermally stable up to 200° C., then a single step thermal degradation follows. 0.7% of weight loss was observed between 25-200° C., due to desolation of process solvents.

The DSC thermogram of Saccharinate salt presented in FIG. 83 showed a broad endothermic melting peak with onset temperature at 100.0° C. and heat of fusion of 76.6 J/g.

The cooling ramp at 10° C./min from 220° C. to -90° C. showed a vitrification around 27.1° C. and the 2<sup>nd</sup> heat cycle displayed a glass transition around 33.8° C. as demonstrated in FIG. 84.

<sup>1</sup>H NMR spectrum in d<sub>6</sub>-DMSO for Saccharinate salt is displayed in FIG. 85. The signals in the spectrum confirmed a presence of 1.0 eq saccharin. Traces of residual process solvents were also observed in the spectrum.

In one embodiment, there is provided 5-MeO-DMT saccharinate. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT saccharinate. In one embodiment, there is provided crystalline 5-MeO-DMT saccharinate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT saccharinate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 81;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, twenty nine or more, thirty or more, thirty one or more, thirty two or more, thirty three or more, thirty four or more, thirty five or more, or thirty six peaks in an XRPD diffractogram as detailed in Table 20, Table 20a or Table 20b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 20, Table 20a or Table 20b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 82;

A weight loss of about 0.7% between 25-200° C., as measured by TGA thermogram;

A weight loss of about 0.2-1.2% between 25-200° C., as measured by TGA thermogram;

A weight loss of about 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8, 0.9, 1.0, 1.1 or 1.2% between 25-200° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 83 or FIG. 84;

A melting endothermic event with an onset of around 100.0° C. and an enthalpy of 76.6 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 95-105° C. and an enthalpy of around 70-80 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 95, 96, 97, 98, 99, 100, 101, 102, 103, 104 or 105° C. and an enthalpy of around 70, 71, 72, 73, 74, 75, 76, 77, 78, 79 or 80 J/g, as measured in a DSC thermogram;

A vitrification around 27.1° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A vitrification around 22-32° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A vitrification around 22, 23, 24, 25, 26, 27, 28, 29, 30, 31 or 32° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A glass transition around 33.8° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.; and/or

A glass transition around 28-38° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A glass transition around 28, 29, 30, 31, 32, 33, 34, 35, 36, 37 or 38° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 85.

#### Physical Stability of Salts

Physical stability stress tests were conducted on generated crystalline salts at controlled temperature and relative humidity. The powder samples were stored at 40° C./75% RH for three days. After this time samples were analysed by XRPD to see whether the physical state of salts changed during the storage.

#### Phosphate Salt

As shown in, or substantially as shown in, FIG. 86, the crystalline Phosphate salt did not undergo any transformation. The XRPD diffractogram of Phosphate salt after three days of storage at 40° C./75% RH agrees with the input material.

#### Fumarate Salt

XRPD of Fumarate salt after three days of storage at 40° C./75% RH showed the same XRPD crystalline pattern when compared to XRPD diffractogram of the input material. However, some additional peaks were also observed as indicated by arrows in FIG. 87.

To understand what phase transformation might happened during the storage, observed additional peaks were compared with XRPD pattern of free base and fumaric acid. XRPD diffractogram showed that positions of additional peaks are not characteristic of the free base and/or fumaric acid as displayed in FIG. 88. More experiments would need to be carried out to see if a new polymorph or hydrate form of Fumarate salt was generated during the storage for three days at 40° C./75% RH.

#### Tartrate Salt

No change in the crystalline form was observed for Tartrate salt after three days of storage at 40° C./75% RH as shown in, or substantially as shown in, FIG. 89.

#### Malate Salt

The crystalline Malate salt underwent conversion to a "gum" material after 3 days of storage at 40° C./75% RH, therefore XRPD analysis were not performed.

#### Tosylate Salt

Crystalline Tosylate salt after three days storage at 40° C./75% RH exhibited the identical XRPD pattern as the input material as demonstrated in FIG. 90.

#### Saccharinate Salt

Saccharinate salt after the storage at 40° C./75% RH for three days displayed the same XRPD pattern as the input material (FIG. 91).

#### Hydrobromide Salt

XRPD diffractogram of Hydrobromide salt after three days of storage at 40° C./75% RH remained the same. No change in crystalline form was observed as shown in, or substantially as shown in, FIG. 92.

### Example 2: Salt Scale-Up

#### Fumarate Salt

5-MeO-DMT Free Base (12.43 g) was dissolved in acetone (60 ml). 1.05 eqv. of Fumaric acid (0.5M in 5% water:EtOH (v/v)) was added. No precipitation was observed upon addition. The volume was reduced by rotary evaporation to form a "Honey" like liquid. This was then thermally cycled between ambient and 40° C. overnight—the temperature was held for 4 hrs at each condition. No precipitation was observed. To this THF (100 ml) was added. The sample was stirred for 30 mins at RT. The formation of an off-white (light brown) solid was observed. This solid was collected by filtration and dried under vacuum at 80° C. for ~20 hrs. A total of 12.64 g was produced (64.65% yield).

An XRPD diffractogram of the produced material versus that analysed in the screening can be seen in FIG. 93. The diffractograms are considered to represent the same crystalline form.

#### Oxalate Salt

5-MeO-DMT Free Base (2x5 g) was dissolved in acetone (50 ml). 1.05 eqv. of Oxalic acid (1.0 M in water) was added. No precipitation observed upon addition. The volume was reduced by rotary evaporation to form an off-white solid. The solid was collected by filtration and dried under vacuum at 80° C. overnight. A total of 10.39 g was produced (73.48% yield).

An XRPD diffractogram of the produced material versus that analysed in the screening can be seen in FIG. 94. The diffractograms are considered to represent the same crystalline form.

<sup>13</sup>C NMR of Oxalate salt produced during scale-up can be seen in FIG. 101 (pp=ZGIG, D1=60 seconds, >35 hr acquisition time, mono salt).

#### Hydrobromide Salt

5-MeO-DMT Free Base (17.18 g) was dissolved in MeOH (80 ml). 1.05 eqv. of HBr acid (1.0 M in MeOH) was added. No precipitation observed upon addition. Volume reduced by rotary evaporation. No precipitation. Methyl tert-butyl ether (50 ml) was added. Formation of off-white (light brown) solid during addition. Solid collected by filtration and dried under vacuum at 80° C. for ~18 hrs. A total of 16.06 g was produced (68.20% yield).

An XRPD diffractogram of the produced material versus that analysed in the screening can be seen in FIG. 95. The diffractograms are considered to represent two different crystalline forms, pattern or form 1 from the original screening and pattern or form 2 from the scale-up.

This was nominated as pattern 2 with XRPD peak data tabulated in Table 21, 21a and 21b. The XRPD can be seen in FIG. 171.

69

TABLE 21

XRPD Peak data for Hydrobromide pattern 2.		
2θ	d Value	Rel. Intensity
12.693°	6.968	0.053
14.633°	6.049	0.384
16.792°	5.275	0.075
17.175°	5.159	0.160
20.750°	4.277	0.209
21.561°	4.118	1.000
22.574°	3.936	0.087
22.740°	3.907	0.111
24.334°	3.655	0.722
24.871°	3.577	0.195
25.441°	3.498	0.366
26.421°	3.371	0.106
27.484°	3.243	0.174
28.558°	3.123	0.074
29.072°	3.069	0.066
29.388°	3.037	0.126
31.314°	2.854	0.057
31.652°	2.825	0.071

TABLE 21a

XRPD Peak data for Hydrobromide pattern 2. (2 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	12.69°	6.97	0.05
2	14.63°	6.05	0.33
3	16.79°	5.28	0.08
4	17.18°	5.16	0.16
5	20.75°	4.28	0.21
6	21.56°	4.12	1.00
7	22.57°	3.94	0.09
8	22.74°	3.91	0.11
9	24.33°	3.66	0.72
10	24.87°	3.58	0.20
11	25.44°	3.50	0.37
12	26.42°	3.37	0.11
13	27.48°	3.24	0.17
14	28.56°	3.12	0.07
15	29.07°	3.07	0.07
16	29.39°	3.04	0.13
17	31.31°	2.85	0.06
18	31.65°	2.83	0.07

TABLE 21b

XRPD Peak data for Hydrobromide pattern 2. (2 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	12.7°	7.0	0.1
2	14.6°	6.1	0.4
3	16.8°	5.3	0.1
4	17.2°	5.2	0.2
5	20.8°	4.3	0.2
6	21.6°	4.1	1.0
7	22.6°	3.9	0.1
8	22.7°	3.9	0.1
9	24.3°	3.7	0.7
10	24.9°	3.6	0.2
11	25.4°	3.5	0.4
12	26.4°	3.4	0.1
13	27.5°	3.2	0.2
14	28.6°	3.1	0.1
15	29.1°	3.1	0.1
16	29.4°	3.0	0.1
17	31.3°	2.9	0.1
18	31.7°	2.8	0.1

70

No weight loss for this crystalline form of the salt due to moisture content was observed during TGA (Ramp 10° C./min to 400° C.) with 95% of weight remaining at 269° C., as shown in, or substantially as shown in, FIG. 96.

5 The DSC thermogram of Hydrobromide salt presented in FIG. 97 showed an onset temperature at 166.10° C. and heat of fusion of 104.04 J/g.

The cooling ramp showed a vitrification around 42.7° C. and the 2<sup>nd</sup> heat cycle displayed a glass transition around 10 47.0° C. as demonstrated in FIG. 98.

The DSC Thermograms of Hydrobromide salt produced during scale-up (top) versus that initially analysed during the salt screen (bottom) can be seen in FIG. 99.

15 <sup>1</sup>H NMR spectrum for the Hydrobromide salt produced during scale-up is displayed in FIG. 100.

In one embodiment, there is provided 5-MeO-DMT Hydrobromide. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT Hydrobromide. In one embodiment, there is provided crystalline 20 5-MeO-DMT Hydrobromide, or a pharmaceutical composition comprising crystalline 5-MeO-DMT Hydrobromide, as characterised by one or more of:

25 An XRPD pattern as shown in, or substantially as shown in, FIG. 109 or FIG. 171;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more peaks in an XRPD diffractogram as detailed in Tables 21, 21a or 21b;

30 One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Tables 21, 21a or 21b;

A DVS isotherm as shown in, or substantially as shown in, FIG. 106;

A DVS kinetic plot as shown in, or substantially as shown in, FIG. 107;

40 A TGA thermogram as shown in, or substantially as shown in, FIG. 96;

A DSC thermogram as shown in, or substantially as shown in, FIG. 97 or FIG. 98;

45 A melting endothermic event with an onset of around 166.10° C. and an enthalpy of 104.04 J/g, as measured in a DSC thermogram;

A melting endothermic event with an onset of around 160-170° C. and an enthalpy of 100-110 J/g, as measured in a DSC thermogram;

50 A melting endothermic event with an onset of around 160, 161, 162, 163, 164, 165, 166, 167, 168, 169 or 170° C. and an enthalpy of 100, 101, 102, 103, 104, 105, 106, 107, 108, 109 or 110 J/g, as measured in a DSC thermogram;

55 A vitrification around 42.7° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A vitrification around 35-45°C, as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

60 A vitrification around 35, 36, 37, 38, 39, 40, 41, 42, 43, 44 or 45° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

65 A glass transition around 478° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

## 71

A glass transition around 42-52° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.;

A glass transition around 42, 43, 44, 45, 46, 47, 48, 49, 51, or 52° C., as measured in a DSC thermogram with a cooling ramp of 10° C./min from 220° C. to -90° C.; and/or

A <sup>1</sup>H NMR spectrum as shown in, or substantially as shown in, FIG. 100.

## HPLC

The HPLC method was as detailed in the Table below:

Column	XSelect CSH C18, 2.5 µm, 4.6 × 30 mm	
Mobile Phase	A	10 mM Ammonium Acetate
	B	Acetonitrile
Autosampler Temperature	Ambient (28° C.)	
Column Temperature	40° C.	
Injection volume	5 µL	
Wavelength	224 nm	
Flow Rate	2.0 mL/min	
Gradient Time	% MPA	% MPB
0.00	95	5
1.00	95	5
4.00	5	95
4.01	0	100
4.50	0	100
4.51	95	5
6.00	95	5
Run Time	6 minutes	
Sample Concentration	1 mg/mL	
Typical RT	2.08 minutes	

Oxalate Salt HPLC Purity, DVS, Stability at 40° C./75% RH (XRPD and HPLC), PSD/Morphology Assessment

Purity of Oxalate salt produced during scale-up:

Sample ID	Description	Purity (Area %)
DXD2203-11-01	Free base	97.9
DXD2203-13-03	Oxalate Salt	98.3

DVS analysis of the Oxalate salt produced during scale-up was performed and the isotherm plot is shown in FIG. 102. The DVS kinetic plot of the salt is displayed in FIG. 103. XRPD analysis was performed on post DVS Oxalate salt and showed no change in crystalline form occurred during the DVS experiment as demonstrated in FIG. 104. XRPD analysis was performed post storage at 40° C./75% RH for 1 week and no change in crystalline form occurred as demonstrated in FIG. 105. No change in the purity, as analysed by HPLC, was seen after 1 week:

Sample ID	Description	Purity (Area %)	
		T = 0	T = 7 days
DXD2203-13-03	Oxalate Salt	98.3	98.3

Hydrobromide Salt HPLC Purity, DVS, Stability at 40° C./75% RH (XRPD and HPLC), PSD/Morphology Assessment

## 72

Purity of Hydrobromide salt produced during scale-up:

Sample ID	Description	Purity (Area %)
DXD2203-11-01	Free base	97.9
DXD2203-14-01	Hydrogen Bromide Salt	99.7

DVS analysis of the Hydrobromide salt produced during scale-up was performed and the isotherm plot is shown in FIG. 106. The DVS kinetic plot of the salt is displayed in FIG. 107. XRPD analysis was performed on post DVS Hydrobromide salt and showed no change in crystalline form occurred during the DVS experiment as demonstrated in FIG. 108. XRPD analysis was performed post storage at 40° C./75% RH for 1 week and no change in crystalline form occurred as demonstrated in FIG. 109. No change in the purity, as analysed by HPLC, was seen after 1 week:

Sample ID	Description	Purity (Area %)	
		T = 0	T = 7 days
DXD2203-14-01	Hydrogen Bromide Salt	99.7	99.7

Fumarate Salt HPLC Purity, DVS, Stability at 40° C./75% RH (XRPD and HPLC), PSD/Morphology Assessment

Purity of Fumarate salt produced during scale-up:

Sample ID	Description	Purity (Area %)
DXD2203-11-01	Free base	97.9
DXD2203-15-03	Fumarate Salt	99.0

DVS analysis of the Fumarate salt produced during scale-up was performed and the isotherm plot is shown in FIG. 110. ~2.84% mass difference between 40% RH and 0% RH, ~0.45 eq water for hemi fumarate salt.

The DVS kinetic plot of the salt is displayed in FIG. 111. XRPD analysis was performed on post DVS Fumarate salt and showed a change in crystalline form occurred during the DVS experiment as demonstrated in FIG. 112. The new form has been nominated as form/pattern 2. TGA analysis was performed post DVS and the 2.52% loss correlates well with the DVS data and can be seen in FIG. 113. DSC analysis was also performed post-DVS and the results of this can be seen in FIG. 114.

XRPD analysis was performed post storage at 40° C./75% RH for 1 week and the material is a mixture of form/pattern 1 and form/pattern 2 as demonstrated in FIG. 109. No change in the purity, as analysed by HPLC, was seen after 1 week:

Sample ID	Description	Purity (Area %)	
		T = 0	T = 7 days
DXD2203-15-03	Fumarate Salt	99.0	99.0

The XRPD Peak Data for Form/Pattern 2 can be Seen Tabulated in Table 28, 28a and 28b.

Table 28			Table 28a			Table 28b		
Angle 2θ	d Value	Rel. Intensity	Angle 2θ	d Value	Rel. Intensity	Angle 2θ	d Value	Rel. Intensity
1	6.355	13.896	6.36	13.90	0.17	6.4	13.9	0.2
2	10.830	8.162	10.83	8.16	0.05	10.8	8.2	0.0
3	12.111	7.302	12.11	7.30	0.28	12.1	7.3	0.3
4	12.683	6.974	12.68	6.97	0.66	12.7	7.0	0.7
5	13.607	6.502	13.61	6.50	0.07	13.6	6.5	0.1
6	14.097	6.277	14.10	6.28	0.07	14.1	6.3	0.1
7	14.651	6.041	14.65	6.04	0.15	14.7	6.0	0.1
8	15.372	5.759	15.37	5.76	0.83	15.4	5.8	0.8
9	16.332	5.423	16.33	5.42	0.14	16.3	5.4	0.1
10	16.823	5.266	16.82	5.27	0.73	16.8	5.3	0.7
11	17.098	5.182	17.10	5.18	0.21	17.1	5.2	0.2
12	17.522	5.057	17.52	5.06	0.48	17.5	5.1	0.5
13	18.119	4.892	18.12	4.89	0.22	18.1	4.9	0.2
14	18.348	4.831	18.35	4.83	1.00	18.3	4.8	1.0
15	18.882	4.696	18.88	4.70	0.45	18.9	4.7	0.4
16	19.210	4.617	19.21	4.62	0.32	19.2	4.6	0.3
17	19.545	4.538	19.55	4.54	0.10	19.5	4.5	0.1
18	20.328	4.365	20.33	4.37	0.18	20.3	4.4	0.2
19	20.799	4.267	20.80	4.27	0.05	20.8	4.3	0.1
20	20.961	4.235	20.96	4.23	0.09	21.0	4.2	0.1
21	21.593	4.112	21.59	4.11	0.55	21.6	4.1	0.6
22	21.770	4.079	21.77	4.08	0.38	21.8	4.1	0.4
23	22.063	4.026	22.06	4.03	0.24	22.1	4.0	0.2
24	22.432	3.960	22.43	3.96	0.10	22.4	4.0	0.1
25	22.781	3.900	22.78	3.90	0.23	22.8	3.9	0.2
26	23.148	3.839	23.15	3.84	0.23	23.1	3.8	0.2
27	23.732	3.746	23.73	3.75	0.13	23.7	3.7	0.1
28	24.242	3.669	24.24	3.67	0.06	24.2	3.7	0.1
29	24.385	3.647	24.39	3.65	0.15	24.4	3.6	0.1
30	24.877	3.576	24.88	3.58	0.30	24.9	3.6	0.3
31	25.396	3.504	25.40	3.50	0.32	25.4	3.5	0.3
32	25.719	3.461	25.72	3.46	0.16	25.7	3.5	0.2
33	26.035	3.420	26.04	3.42	0.11	26.0	3.4	0.1
34	26.197	3.399	26.20	3.40	0.19	26.2	3.4	0.2
35	26.921	3.309	26.92	3.31	0.15	26.9	3.3	0.1
36	27.621	3.227	27.62	3.23	0.05	27.6	3.2	0.1
37	28.715	3.106	28.71	3.11	0.03	28.7	3.1	0.0
38	29.125	3.064	29.13	3.06	0.03	29.1	3.1	0.0
39	30.132	2.964	30.13	2.96	0.03	30.1	3.0	0.0
40	30.429	2.935	30.43	2.94	0.04	30.4	2.9	0.0
41	30.764	2.904	30.76	2.90	0.04	30.8	2.9	0.0

In one embodiment, there is provided crystalline 5-MeO-DMT fumarate form/pattern 2, or a pharmaceutical composition comprising crystalline 5-MeO-DMT fumarate form/pattern 2, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 112;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, twenty nine or more, thirty or more, thirty one or more, thirty two or more, thirty three or more, thirty four or more, thirty five or more, thirty six or more, thirty seven or more, thirty eight or more, thirty nine or more, forty or more or forty one peaks in an XRPD diffractogram as detailed in Tables 28, 28a or 28b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Tables 28, 28a or 28b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 113; and/or

A DSC thermogram as shown in, or substantially as shown in, FIG. 114.

#### Example 3: Further Salt Characterisation

##### Salt Cracking

5-MeO-DMT HCl (J11635, HCl Pattern 1, 4.94 g) was dissolved in 10 volumes (50 ml) of water giving a clear brown solution. To this was added 1 equivalent of NaOH dropwise as a 4M aqueous solution (4.85 mL) giving a tan suspension which after stirring for 10 minutes formed a brown oil. The oil was extracted with 3×10 vol (50 mL) of 2-MeTHF. The organic phases were combined and washed with brine before being concentrated using a rotary evaporator giving a thick brown oil. The oil was dissolved in 10 vol (50 mL) of 2-MeTHF and concentrated again to a brown oil with a small amount of solid material was present. The sample was dried further in a vacuum oven at RT overnight giving a tan solid. Sample ID: DR-2186-43-01. (Yield=3.856 g).

DR-2186-43-01 (3.865 g) was suspended in 5 volumes (19.3 mL) of water in a round bottom flask and stirred overnight at room temperature. The suspension was filtered through a Buchner funnel using a Whatman grade 1 filter paper and vacuum. The cake was dried under suction for 30 minutes before being transferred to a vacuum oven at RT for 2 hours. Yield=4.233 g (81.1%).



75

Sample ID	DR-2186-43-02
XRPD	Free Form Pattern 1
<sup>1</sup> H-NMR	Consistent with structure, trace residual solvent
IC	No ions detected
HPLC Purity (Pharmorphix Generic Method)	96.5%

#### Phosphate Formation

In a 20 mL scintillation vial 5-MeO-DMT (1.00 g, Free Form Pattern 1, DR-2186-46-02) was dissolved in 5 volumes (5 mL) of IPA:water 9:1 at 50° C. on a Polar Bear heat/cool block with magnetic bottom stirring (500 RPM), giving a brown solution. 1.1 mol eq. (5.05 mL) of phosphoric acid was added at a 1M solution in THF dropwise over 2 minutes. Initially this gave a white precipitate, on further addition a light brown gummy solid formed on the base of the vial which became more solid over 5 minutes of stirring. The encrusted solid was agitated using a spatula and after a further 5 minutes a light tan suspension was obtained. The crystallisation was then cooled to 5° C. at 0.1° C./min and held there overnight.

An aliquot (ca. 0.3 mL) of suspension was filtered using a cartridge and frit along with positive pressure. The solid was dried briefly under a stream of N<sub>2</sub> before collecting an XRPD (DR-2186-45-01\_A1) shown in FIG. 116. This showed that PHO Pattern 1+extra peaks had been formed (matched DR-2186-26-10). The suspension was stirred at 5° C. for a further 24 hours and a second aliquot taken and XRPD collected (DR-2186-45-01\_A2). This showed no change.

The bulk sample was isolated by vacuum filtration using a Buchner funnel and 55 mm Whatman grade 1 filter paper. The vial and cake were washed with 1 mL of cold IPA:water 9:1. The material was dried under suction for 30 minute.

#### Phosphate Formation—Re-Crystallisation

DR-2186-45-01 (1.00 g, PHO Pattern 1+extra peaks) was weighed into a 20 ml scintillation vial and a stirrer bar and 5 volumes (5 mL) of MeOH added. On a Polar Bear heating

76

block the suspension was heated to 50° C., 500 RPM. Sequential aliquots of hot MeOH were added to the sample according to the Table below, noting observations, looking for sample dissolution.

Volumes of MeOH	50° C.					60° C.	
	5	8	12	16	18	18	20
Observation	X	X	X	X	X	X	X

At 20 vols (20 mL) dissolution had still not been achieved so the suspension was transferred to a 50 mL Easymax vessel fitted with overhead stirring.

Volumes of MeOH	60° C., 150 RPM			60° C., 250 RPM
	20	22	24	25
Observation	X	X	✓/X	✓

✓ = solution, ✓/X = turbid solution, X = suspension

Dissolution was achieved at 25 volumes of MeOH at 60° C. giving a clear yellow/brown solution. This was cooled at 0.5° C./min to 5° C. At 50° C. seeding was attempted with ca. 5 mg of PHO Pattern 1 (DR-2186-34-10), however, no visual change in solution turbidity was noted. Turbidity was observed to be starting to increase at 34° C. The crystallisation was held at 5° C. for 1 hour and was a thick off white suspension. The solid was isolated by vacuum filtration through a Whatman grade 1 filter paper and Buchner funnel. The cake was dried under suction for 30 mins. (Yield=296 mg, 29.7%). There was some fouling/encrustation on the vessel which a sample of was collected separately (ID: DR-2186-48-01\_crust). An XRPD analysis of the bulk sample, the encrusted material and the Phosphate pattern 1 reference can be seen in FIG. 117.

A summary of the characteristics of the 5-MeO-DMT phosphate salt can be seen in the Table below:

Sample ID	DR-2186-48-01
XRPD	PHO Pattern 1
<sup>1</sup> H-NMR	Consistent with structure, 0.1 mol eq. of residual MeOH in sample
IC	1.43 mol eq. of phosphate
SEM	Long, thin lath shaped particles with some larger particles that are more plat like, up to 200 µm long
PLM	Needle/long thin lath morphology with some larger plates ca. 200 µm in length. Crystals exhibit birefringence.
HSM (Hot Stage Microscopy)	Melting of sample occurs from 155-164° C.
TGA	0.4 wt. % (0.04 mol. eq. MeOH) mass loss from 40-125° C. A further 0.95% (0.1 mol eq. MeOH) was lost during the melt, 130-170° C. Decomposition onset from 180° C.
DSC	Small endotherm onset 68.6° C. (3 J/g) associated with first mass loss. Large sharp endotherm onset at 161.7° C. (77 J/g), assigned as the melt based on the HSM data. This is immediately followed by an exotherm and charge in baseline which may indicate decomposition beginning.
GVS	Small total reversible mass change of 0.34 wt. % (0.06 mol eq. water) from 0-90% RH with a small hysteresis observed on the second desorption cycle. The material is classed as slightly hygroscopic (based on European Pharmacopeia definitions). The solid form of the material was largely unchanged by XRPD but very small peaks can be seen that may be consistent with the additional peaks previously observed.
HPLC Purity (Pharmorphix Generic Method 30 mins)	98.0%

-continued

Sample ID	DR-2186-48-01
Static Storage 7 days - 40° C./75% RH	XRPD - PHO Pattern 1 HPLC - 98.1%
Static Storage 7 days - 25° C./97% RH	XRPD - PHO Pattern 1 HPLC - 98.0%

Assessment of Polymorphic Behaviour of 5-MeO-DMT Phosphate <sup>10</sup>

DR-2186-45-01 (20 mg, PHO Pattern 1+Extra Peaks) was weighed into 10 HPLC vials. To this was added 10 volumes (200 µL) of solvent, a stirrer bar and the samples heated to 50° C. on a Polar Bear heat/cool block (400 RPM). After holding at 50° C. for 30 minutes and observation was made and the samples cooled to 5° C. where a further observation was made. All samples were then subjected to heat/cool cycles between 5 and 50° C. holding for 4 hours at each temperature for 24 hours.

-continued

No.	Pos. [°2θ]	Rel. Int. [%]
7	15.6	6
8	17.4	22
9	17.6	24.4
10	17.8	13.5
11	18.5	3.3
12	19.1	5.1

Sample	Solvent	Observation at 50° C.	Observation at 5° C.	Observation at Isolation	XPRD
DR-2186-52-01	MeOH	X	X	White Suspension	PHO Pattern 1
DR-2186-52-02	Acetone	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-03	CAN	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-04	IPA:Water 9:1	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-05	2-MeTHF	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-06	EtOAc	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-07	Ethanol	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-08	Toluene	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-09	MEK	X	X	White Suspension	PHO Pattern 1 + Extra Peaks
DR-2186-52-10	TBME	X	X	White Suspension	PHO Pattern 1 + Extra Peaks

The results of the XRPD analysis of the samples can be seen in FIG. 118.

Non of the samples completely dissolved during the experiment. PHO Pattern 1+Extra Peaks (which was the input material) was obtained from all solvents other than methanol which yielded pure PHO Pattern 1.

5-MeO-DMT Phosphate Pattern 1

5-MeO-DMT Phosphate Pattern 1 (XRPD analysis shown in FIG. 122, peaks in the Table below) was crystallised as a phase pure form by recrystallization of a mixture of Pattern 1 and another form from methanol. The resulting solid has a HPLC purity of 98.0% (input material 96.5%) and the IC showed that there was 1.43 mol. eq. of phosphate present in the sample (see FIG. 127).

-continued

No.	Pos. [°2θ]	Rel. Int. [%]
13	19.5	45.3
14	20.4	100
15	21.3	27
16	21.8	19.7
17	22.4	8
18	23.4	54.7
19	23.9	19.7
20	24.3	43.2
21	25.3	27.3
22	25.8	45.1
23	26.7	3.7
24	27.1	6.6
25	27.9	6
26	28.4	13.9
27	28.9	8.8
28	29.2	4
29	30.6	6.1
30	31.5	4

No.	Pos. [°2θ]	Rel. Int. [%]
1	5.5	24
2	7.2	6.5
3	10.6	10.2
4	13.5	26.2
5	14.5	11.4
6	15.3	18.7

This higher stoichiometry maybe the driving force behind the formation of a mixture of forms in the initial salt formation step as only 1.1 equivalents of phosphoric acid were added. The solid crystallised as long thin colourless

lath crystals with some of the larger ones reaching 200  $\mu\text{m}$  in length and becoming more plate like.

The  $^1\text{H-NMR}$  spectrum is (FIG. 121) consistent with the structure and the sample found to contain 0.1 mol. eq. of MeOH. The thermal data suggests a small amount of residual solvent, 0.4 wt. % lost between 40-125° C. and a further 0.95 wt. % (both losses equate to 0.14 mol. eq. of MeOH) of solvent lost from 130-170° C. The DSC contains a large sharp endotherm with an onset at 161.7° C. which is assigned as the melt, corroborated by the melt observed in the HSM. The TGA and DSC Thermograms can be seen in FIG. 123.

The sample is classed as slightly hygroscopic with a total reversible mass change of 0.3 wt. % between 0-90% RH. The DVS isotherm and DVS kinetic plot can be seen in FIGS. 124 and 125, respectively. The solid form was unchanged after a double-cycle experiment, the XRPD analysis confirming this can be seen in FIG. 126. Phosphate Pattern 1 was also found to be stable to storage at both 40° C./75% RH and 25° C./97% RH, with no change observed by XRPD or HPLC, see FIGS. 128-130.

A polymorph assessment was carried out in 10 different solvents using PHO Pattern 1+extra peaks as the input material, maturing a slurry of the salt between 5° C. and 50° C. for 24 hours. The results of the XRPD analysis of the samples can be seen in FIG. 118. The isolated solid was the same as the input material in all cases other than MeOH.

In one embodiment, there is provided crystalline 5-MeO-DMT phosphate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT phosphate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 122;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, twenty nine or more or thirty peaks in an XRPD diffractogram as detailed the Table above;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in the Table above;

A TGA thermogram as shown in, or substantially as shown in, FIG. 123;

A weight loss of about 0.4% between 40 and 125° C. and a weight loss of about 0.95% between 130 and 170° C., as measured by TGA thermogram;

A weight loss of about 0.2-0.6% between 40 and 125° C. and a weight loss of about 0.85-1.05% between 130 and 170° C., as measured by TGA thermogram;

A weight loss of about 0.2, 0.3, 0.4, 0.5 or 0.6% between 40 and 125° C. and a weight loss of about 0.85, 0.9, 0.95, 1.0 or 1.05% between 130 and 170° C., as measured by TGA thermogram;

A DSC thermogram as shown in, or substantially as shown in, FIG. 123;

A melting endothermic event with an onset of around 161.7° C., as measured in a DSC thermogram;

A melting endothermic event with an onset of around 155-165° C., as measured in a DSC thermogram; and/or

A melting endothermic event with an onset of around 155, 156, 157, 158, 159, 160, 161, 162, 163, 164 or 165° C., as measured in a DSC thermogram.

#### L-Tartrate Formation

In a 20 mL scintillation vial 5-MeO-DMT (1.00 g, Free Form Pattern 1, DR-2186-46-02) was dissolved in 5 volumes (5 mL) of IPA:water 9:1 at 50° C. on a Polar Bear heat/cool block with magnetic bottom stirring (500 RPM), giving a brown solution. 1.1 mol eq. (5.05 mL) of L-tartaric acid was added at a 1M solution in THF dropwise over 2 minutes. This formed a brown oil, which formed a thick light tan suspension after stirring at 50° C. for 5 minutes. The crystallisation was then cooled to 5° C. at 0.1° C./min and held there overnight.

An aliquot (ca. 0.3 mL) of suspension was filtered using a cartridge and frit along with positive pressure. The solid was dried briefly under a stream of N<sub>2</sub> before collecting an XRPD. This showed that TAR Pattern 1 had been formed. (XRPD analysis shown in FIGS. 131 and 132, peaks in the Table below).

No.	Pos. [°2 $\theta$ ]	Rel. Int. [%]
1	5.0	100
2	10.1	2.6
3	11.7	2.8
4	13.0	10.7
5	15.1	12.1
6	15.5	6.1
7	16.5	5.2
8	17.7	21.9
9	18.2	40.1
10	18.7	38.3
11	19.6	9.2
12	20.2	19.5
13	20.7	34.7
14	21.4	8
15	22.1	23.4
16	22.6	8.7
17	23.0	5
18	23.6	13.2
19	24.8	3.7
20	25.3	4.8
21	26.0	37.1
22	26.3	1.9
23	27.0	3.5
24	27.2	6.6
25	27.8	3
26	29.3	3.8
27	29.9	6.2
28	31.1	4.6
29	31.7	3.7

The bulk sample was isolated by vacuum filtration using a Buchner funnel and 55 mm Whatman grade 1 filter paper. The vial and cake were washed with 1 mL of cold IPA:water 9:1. The material was dried under suction for 15 minutes then transferred to a vacuum oven at RT for 4.5 hours. Yield=1.413 g (83.7%).

A summary of the characteristics of the 5-MeO-DMT tartrate salt can be seen in the Table below:

Sample ID	DR-2186-46-01
XRPD	TAR Pattern 1
$^1\text{H-NMR}$	Consistent with structure, 0.97 mol eq. tartrate, 0.02 mol eq. of residual THF and IPA
SEM	Very small plate particles <1-20 $\mu\text{m}$ in size
PLM	Very small crystals exhibiting birefringence

-continued

Sample ID	DR-2186-46-01
HSM (Hot Stage Microscopy)	Melting of sample observed from 135° C. and complete by 143° C.
TGA	No mass loss observed in sample before decomposition.
DSC	Large sharp endotherm with onset at 145.0° C. (116 J/g), this has been assigned as a melt due to good agreement with observations made during the hot stage microscopy
GVS	Total mass change of 0.58 wt. % from 0-90% RH (0.12 mol eq. of water) and is reversible with no hysteresis. The material is classed as slightly hygroscopic (based on European Pharmacopeia definitions). The solid form of the material was unchanged by XRPD.
HPLC Purity (Pharmorphix Generic Method 30 mins)	98.0%
Static Storage 7 days-40° C./ 75% RH	KRPD-TAR Pattern 1 HPLC-97.8%

DR-2186-46-01 (20 mg, TAR Pattern 1) was weighed into 10 HPLC vials. To this was added 10 volumes (200  $\mu$ L) of solvent, a stirrer bar and the samples heated to 50° C. on a Polar Bear heat/cool block (400 RPM). After holding at 50° C. for 30 minutes and observation was made and the samples cooled to 5° C. where a further observation was made. All samples were then subjected to heat/cool cycles between 5 and 50° C. holding for 4 hours at each temperature for 24 hours.

Sample	Solvent	Observation at 50° C.	Observation at 5° C.	Observation at Isolation	XRPD
DR-2186-52-01	MeOH	X	X	White Suspension	TAR Pattern 1
DR-2186-52-03	Acetone	X	X	White Suspension	TAR Pattern 1
DR-2186-52-03	ACN	X	X	White Suspension	TAR Pattern 1
DR-2186-52-04	IPA:Water 9:1	X	X	White Suspension	TAR Pattern 1
DR-2186-52-05	2-MeTHF	X	X	White Suspension	TAR Pattern 1
DR-2186-52-06	EtOAc	X	X	White Suspension	TAR Pattern 1
DR-2186-52-07	Ethanol	X	X	White Suspension	TAR Pattern 1
DR-2186-52-08	Toluene	X	X	White Suspension	TAR Pattern 1
DR-2186-52-09	MEK	X	X	White Suspension	TAR Pattern 1
DR-2186-52-10	TBME	X	X	White Suspension	TAR Pattern 1

The results of the XRPD analysis of the samples can be seen in FIG. 119. FIG. 120 shows the XRPD analysis results for low intensity samples (solvents: EtOH, ACN).

No dissolution was observed during the experiment and only TAR Pattern 1 was obtained after the 24 hours of maturation.

5-MeO-DMT Tartrate Pattern 1

5-MeO-DMT Tartrate Pattern 1 (FIG. 131) was crystallised from an THF:IPA:Water (ca. 10:9:1) solvent system. The resulting solid has a HPLC purity of 98.0% (input material 96.5%) and the <sup>1</sup>H-NMR showed that there was 1.0 mol. eq. of tartrate present in the sample with very low amounts of residual IPA or THF (FIG. 133). The sample is comprised of very small plate like crystals 1-20  $\mu$ m in size.

The TGA (FIG. 134) showed no mass loss until the onset of decomposition at 175° C. The DSC contains a large sharp endotherm with an onset at 145.0° C. which is assigned as the melt, corroborated by the melt observed in the HSM.

The sample is classed as slightly hygroscopic with a total reversible mass change of 0.6 wt. % between 0-90% RH

(FIG. 135). The solid-form was unchanged after the double-cycle experiment (FIG. 136). Tartrate Pattern 1 was also found to be stable to storage at both 40° C./75% RH and 25° C./97% RH, with no change observed by XRPD or HPLC (FIGS. 137-141).

A polymorph assessment was carried out in 10 different solvents, maturing a slurry of the salt between 5° C. and 50° C. for 24 hours.

In one embodiment, there is provided crystalline 5-MeO-DMT tartrate, or a pharmaceutical composition comprising crystalline 5-MeO-DMT tartrate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 131 or FIG. 132;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more or twenty nine peaks in an XRPD diffractogram as detailed the peak Table above;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in the peak Table above;

A TGA thermogram as shown in, or substantially as shown in, FIG. 134;

A DSC thermogram as shown in, or substantially as shown in, FIG. 134;

A melting endothermic event with an onset of around 145.0° C., as measured in a DSC thermogram;

A melting endothermic event with an onset of around 140-150° C., as measured in a DSC thermogram;

A melting endothermic event with an onset of around 140, 141, 142, 143, 144, 145, 146, 147, 148, 149, 150° C., as measured in a DSC thermogram;

A total reversible mass change of around 0.6% between 0-90% RH;

A total reversible mass change of around 0.4-0.8% between 0-90% RH;

A total reversible mass change of around 0.4, 0.5, 0.6, 0.7 or 0.8% between 0-90% RH; and/or

A DVS Isotherm as shown in, or substantially as shown in, FIG. 135.

Example 4: Further Characterisation of  
5-MeO-DMT Hydrochloride

## Instruments

## X-Ray Powder Diffraction (XRPD)

XRPD diffractograms were acquired using Bruker D2 Phaser diffractometer. A copper x-ray source at 300 W was used in conjunction with a Lynxeye detector. Samples were prepared using a zero-background sample holder. The samples were scanned from 5 to 32° (2 $\theta$ ) using a step size of 0.02° and a time per step of 0.13 second whilst spinning the sample. Diffractograms were plotted using the EVA program from Bruker.

## Thermo—Gravimetric Analysis (TGA)

TGA thermograms were obtained with a TA Instrument Discovery 550 in Al pans. The heating rate used was 10° C./min linear ramp from 25 to 400° C. with a nitrogen purging at a rate of 60 ml/min. TGA thermograms were analysed using TRIOS software.

## Differential Scanning Calorimetry (DSC)

DSC analyses were performed on a TA Instrument DSC250 with a Tzero cell purged at constant flow rate of 50 ml min<sup>-1</sup> with dry nitrogen and a refrigerated cooling system RCS90. The instrument was calibrated using Indium as a standard. A small quantity of the samples was weighed into TA Tzero Aluminium pan with pierced lid. Samples were heating at 10° C./min in heat-cool-reheat method. TRIOS software was used to analyse DSC scans.

## Nuclear Magnetic Resonance Spectroscopy (NMR)

The NMR spectra were measured on Bruker NEO spectrometer operating at 400.13 MHz for protons. Samples were dissolved in d<sub>6</sub>-DMSO. Data were processed using MestReNovax64 software.

## Initial Characterisation of Hydrochloride Salt

Baseline analysis of the hydrochloride salt was performed to compare to data generated in later studies. The hydrochloride was analysed by XRPD and the diffractogram is shown in FIG. 142. The material is crystalline and this was designated as pattern 1. The XRPD peak data is shown in Table 22, 22a or 22b:

TABLE 22

XRPD Peak data for hydrochloride pattern 1			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.191°	9.614	0.59
2	12.275°	7.205	0.24
3	13.601°	6.505	0.16
4	14.030°	6.307	0.34
5	14.925°	5.931	1.00
6	15.513°	5.708	0.01
7	18.403°	4.817	0.59
8	18.396°	4.693	0.06
9	19.613°	4.523	0.58
10	21.305°	4.167	0.13
11	22.899°	3.881	0.02
12	23.133°	3.842	0.06
13	23.436°	3.793	0.03
14	23.826°	3.732	0.53
15	24.565°	3.621	0.08
16	25.048°	3.552	0.23
17	25.716°	3.461	0.04
18	25.974°	3.428	0.14
19	26.226°	3.395	0.03
20	26.783°	3.326	0.13
21	27.271°	3.268	0.04
22	27.547°	3.235	0.06
23	28.110°	3.172	0.16
24	23.955°	3.081	0.05

TABLE 22-continued

XRPD Peak data for hydrochloride pattern 1			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
25	30.045°	2.972	0.01
26	30.670°	2.913	0.06
27	31.009°	2.882	0.03
28	31.431°	2.844	0.02

TABLE 22a

XRPD Peak data for hydrochloride pattern 1. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.19°	9.61	0.59
2	12.28°	7.21	0.24
3	13.60°	6.51	0.16
4	14.03°	6.31	0.34
5	14.93°	5.93	1.00
6	15.51°	5.71	0.01
7	18.40°	4.82	0.59
8	18.90°	4.69	0.06
9	19.61°	4.52	0.58
10	21.31°	4.17	0.13
11	22.90°	3.88	0.02
12	23.13°	3.84	0.06
13	23.44°	3.79	0.03
14	23.83°	3.73	0.53
15	24.57°	3.62	0.08
16	25.05°	3.55	0.23
17	25.72°	3.46	0.04
18	25.97°	3.43	0.14
19	26.23°	3.40	0.03
20	26.78°	3.33	0.13
21	27.27°	3.27	0.04
22	27.55°	3.24	0.06
23	28.11°	3.17	0.16
24	28.96°	3.08	0.05
25	30.05°	2.97	0.01
26	30.67°	2.91	0.06
27	31.01°	2.88	0.03
28	31.43°	2.84	0.02

TABLE 22b

XRPD Peak data for hydrochloride pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	9.2°	9.6	0.6
2	12.3°	7.2	0.2
3	13.6°	6.5	0.2
4	14.0°	6.3	0.3
5	14.9°	5.9	1.0
6	15.5°	5.7	0.0
7	18.4°	4.8	0.6
8	18.9°	4.7	0.1
9	19.6°	4.5	0.6
10	21.3°	4.2	0.1
11	22.9°	3.9	0.0
12	23.1°	3.8	0.1
13	23.4°	3.8	0.0
14	23.8°	3.7	0.5
15	24.6°	3.6	0.1
16	25.1°	3.6	0.2
17	25.7°	3.5	0.0
18	26.0°	3.4	0.1
19	26.2°	3.4	0.0
20	26.8°	3.3	0.1
21	27.3°	3.3	0.0
22	27.6°	3.2	0.1

TABLE 22b-continued

XRPD Peak data for hydrochloride pattern 1. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
23	28.1°	3.2	0.2
24	29.0°	3.1	0.1
25	30.1°	3.0	0.0
26	30.7°	2.9	0.1
27	31.0°	2.9	0.0
28	31.4°	2.8	0.0

Thermal analysis was performed and the TGA thermogram is shown in FIG. 143. This shows the material is an anhydrous form with good thermal stability up to  $\sim 235^\circ\text{C}$ . where gross decomposition is taking place. The DSC thermogram for the first heat cycle is shown in FIG. 144. This shows a single sharp endothermic event, melting, with an onset of  $146.5^\circ\text{C}$ . and an enthalpy of  $121.7\text{ J/g}$ . The sample was then cooled in the DSC to  $\sim -90^\circ\text{C}$  and reheated. The thermograms are shown in FIG. 145 and show vitrification at  $40.2^\circ\text{C}$ . in the cool cycle and a  $T_g$  at  $44.8^\circ\text{C}$ . in the second heat cycle. There was no evidence of recrystallization during this experiment when performed at a heating or cooling rate of  $10^\circ\text{C. min}^{-1}$ .

To look for new crystal forms via recrystallisation from the melt, the DSC experiment was repeated but different cool and reheat rates were used in the second cycle. Three separate samples were all heated to  $200^\circ\text{C}$ . at  $10^\circ\text{C. min}^{-1}$  and then cooled to  $-90^\circ\text{C}$ . and reheated to  $200^\circ\text{C}$ . at three different rates of  $5^\circ\text{C. min}^{-1}$ ,  $2^\circ\text{C. min}^{-1}$  and  $1^\circ\text{C. min}^{-1}$ . The cool cycles only showed vitrification (data not shown) but the reheat cycles showed a recrystallisation event followed by a melt, but the melt is consistent with the recrystallised material being pattern 1 and as such, no new crystalline forms were observed in these experiments. The three thermograms are shown in FIG. 146.

The hydrochloride was analysed by  $^1\text{H}$  and  $^1\text{H}$ - $^{13}\text{C}$  HSQC NMR. The  $^1\text{H}$  NMR spectrum is shown in FIG. 147 and is consistent with the supplied structure. There are no obvious evidence of residual process solvents and the material shows a high chemical purity. The  $^1\text{H}$ - $^{13}\text{C}$  HSQC spectrum is shown in FIG. 148 and is consistent with the supplied structure and shows the expected DEPT editing.

In one embodiment, there is provided 5-MeO-DMT HCl. In one embodiment, there is provided a pharmaceutical composition comprising 5-MeO-DMT HCl. In one embodiment, there is provided crystalline 5-MeO-DMT HCl, or a pharmaceutical composition comprising crystalline 5-MeO-DMT HCl, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 142;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more or twenty eight peaks in an XRPD diffractogram as detailed in Table 22, Table 22a or Table 22b;

One or more, two or more, three or more, four or more, five or more peaks in an XRPD diffractogram with a

relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 22, Table 22a or Table 22b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 143;

A DSC thermogram as shown in, or substantially as shown in, FIG. 144;

A melting endothermic event with an onset of around  $146.5^\circ\text{C}$ . and an enthalpy of  $121.7\text{ J/g}$ , as measured in a DSC thermogram;

A melting endothermic event with an onset of around  $140$  to  $150^\circ\text{C}$ . and an enthalpy of around  $115$  to  $125\text{ J/g}$ , as measured in a DSC thermogram;

A melting endothermic event with an onset of around  $140$ ,  $141$ ,  $142$ ,  $143$ ,  $144$ ,  $145$ ,  $146$ ,  $147$ ,  $148$ ,  $149$ ,  $150^\circ\text{C}$ . and an enthalpy of around  $155$ ,  $156$ ,  $157$ ,  $158$ ,  $159$ ,  $160$ ,  $161$ ,  $162$ ,  $163$ ,  $164$ ,  $165$ ,  $166$ ,  $167$ ,  $168$ ,  $169$  or  $170\text{ J/g}$ , as measured in a DSC thermogram;

A vitrification around  $40.2^\circ\text{C}$ ., as measured in a DSC thermogram with a cooling ramp of  $10^\circ\text{C./min}$  from  $230^\circ\text{C}$ . to  $-90^\circ\text{C}$ .;

A vitrification around  $35$ - $45^\circ\text{C}$ ., as measured in a DSC thermogram with a cooling ramp of  $10^\circ\text{C./min}$  from  $230^\circ\text{C}$ . to  $-90^\circ\text{C}$ .;

A vitrification around  $35$ ,  $36$ ,  $37$ ,  $38$ ,  $39$ ,  $40$ ,  $41$ ,  $42$ ,  $43$ ,  $44$  or  $45^\circ\text{C}$ ., as measured in a DSC thermogram with a cooling ramp of  $10^\circ\text{C./min}$  from  $230^\circ\text{C}$ . to  $-90^\circ\text{C}$ .;

A glass transition around  $44.8^\circ\text{C}$ ., as measured in a DSC thermogram with a cooling ramp of  $10^\circ\text{C./min}$  from  $230^\circ\text{C}$ . to  $-90^\circ\text{C}$ .;

A glass transition around  $40$ - $50^\circ\text{C}$ ., as measured in a DSC thermogram with a cooling ramp of  $10^\circ\text{C./min}$  from  $230^\circ\text{C}$ . to  $-90^\circ\text{C}$ .;

A glass transition around  $41$ ,  $42$ ,  $43$ ,  $44$ ,  $45$ ,  $46$ ,  $47$ ,  $48$ ,  $49$  or  $50^\circ\text{C}$ ., as measured in a DSC thermogram with a cooling ramp of  $10^\circ\text{C./min}$  from  $230^\circ\text{C}$ . to  $-90^\circ\text{C}$ .;

A  $^1\text{H}$  NMR spectrum as shown in, or substantially as shown in, FIG. 147; and/or

A  $^1\text{H}$ - $^{13}\text{C}$  HSQC NMR spectrum as shown in, or substantially as shown in, FIG. 148.

Polymorphism Screen

To render the material amorphous and thus remove seeds of the pattern 1 the material was rendered amorphous. To  $2.605\text{ g}$  of the hydrochloride lot RPI-014-022 dioxane ( $55\text{ ml}$ ) and water ( $5\text{ ml}$ ) was added.

The mixture was agitated gently and warmed to aid dissolution. The clear solution was then divided equally between  $60\text{ HPLC}$  vials ( $\sim 40\text{ mg}$  salt in each vial) and the vials were then frozen at  $\sim -18^\circ\text{C}$ . for  $6\text{ hours}$  and then dried by lyophilisation overnight. One sample was analysed by XRPD and the diffractogram shown in FIG. 149 demonstrates that although not  $100\%$  amorphous, the crystallinity is reduced, albeit still pattern 1 in nature.

Thermal Cycling

Twenty five of the lyophilised samples were treated with solvent and thermally cycled between ambient and  $40^\circ\text{C}$ . with four hours spent under each condition. After three days and solids were isolated by centrifuge filtration and analysed by XRPD. Any solutions were allowed to evaporate at RT but this did not yield any new solids. The solvents, observations, isolation and XRPD results are summarised in the Table below:

Summary of thermal cycle of hydrochloride salt

Experiment	Solvent	Volume added (ml)	Initial Observations	3-day Observations	Isolation	XRPD Result
DJP2202-007-01	1,4-Dioxane	200	no change	solid mass	After thermal cycle	Pattern 2
DJP2202-007-02	1-Propanol	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-03	2-Butanol	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-04	2-Ethoxyethanol	200	dissolved	clear solution	Evaporated but no solid	n/a
DJP2202-007-05	2-Methyltetrahydrofuran	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-06	2-Propanol	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-07	Acetone	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-08	Acetonitrile	200	no change	lump of material	After thermal cycle	Pattern 1
DJP2202-007-09	Anisole	200	no change	lump of material	After thermal cycle	Pattern 1
DJP2202-007-10	Chlorobenzene	200	no change	white solid	After thermal cycle	Pattern 1
DJP2202-007-11	Ethanol	150	no change	white solid	After thermal cycle	Pattern 1
DJP2202-007-12	Ethyl acetate	200	no change	white solid	After thermal cycle	Pattern 1
DJP2202-007-13	Isopropyl acetate	200	no change	white solid	After thermal cycle	Pattern 1
DJP2202-007-14	Methanol	200	dissolved	clear solution	After thermal cycle	Pattern 3
DJP2202-007-15	Propyl acetate	200	no change	white solid	After thermal cycle	Pattern 1
DJP2202-007-16	Methylethyl ketone	200	no change	white crystals	After thermal cycle	Pattern 2
DJP2202-007-17	Formamide	150	dissolved	few crystals	Evaporated but no solid	n/a
DJP2202-007-18	N,N-Dimethylacetamide	200	dissolved	clear solution	Evaporated but no solid	n/a
DJP2202-007-19	N-Methylpyrrolidone	200	no change	clear solution	Evaporated but no solid	n/a
DJP2202-007-20	iso-Propyl acetate	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-21	2-Me-1-PrOH	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-22	Tetrahydrofuran	200	no change	few crystals	After thermal cycle	Pattern 1
DJP2202-007-23	MeOH/water 953/47 Calc Aw 0.214	200	dissolved	clear solution	Evaporated to oil	n/a
DJP2202-007-24	MeOH/water 693/307 Calc Aw 0.599	200	dissolved	clear solution	Evaporated to oil	n/a
DJP2202-007-25	MeOH/water 360/640 Calc Aw 0.821	200	dissolved	clear solution	Evaporated but no solid	n/a

The XRPD for the new pattern 2 is shown in FIG. 150 compared to the supplied pattern 1. The XRPD peak data is shown in Table 23, 23a or 23b:

TABLE 23

XRPD Peak data for hydrochloride pattern 2			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.454°	10.450	0.01
2	11.786°	7.502	0.17
3	12.650°	6.992	0.01
4	13.042°	6.783	0.25
5	13.789°	6.417	0.02
6	14.600°	6.062	0.01
7	14.968°	5.914	0.49

TABLE 23-continued

XRPD Peak data for hydrochloride pattern 2			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
8	16.120°	5.494	0.02
9	17.283°	5.127	0.14
10	17.529°	5.055	0.29
11	17.791°	4.982	1.00
12	18.363°	4.828	0.05
13	18.530°	4.784	0.90
14	19.571°	4.532	0.37
15	19.858°	4.467	0.05
16	20.354°	4.360	0.18
17	20.883°	4.250	0.06
18	21.344°	4.160	0.08

89

TABLE 23-continued

XRPD Peak data for hydrochloride pattern 2			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
19	22.244°	3.993	0.17
20	22.715°	3.912	0.38
21	23.321°	3.811	0.09
22	23.645°	3.760	0.66
23	24.751°	3.594	0.21
24	25.620°	3.474	0.15
25	26.224°	3.396	0.12
26	26.593°	3.349	0.51
27	26.947°	3.306	0.21
28	27.103°	3.287	0.30
29	27.442°	3.248	0.04
30	27.706°	3.217	0.02
31	28.051°	3.178	0.16
32	28.541°	3.125	0.11
33	28.813°	3.096	0.05
34	29.442°	3.031	0.05
35	30.057°	2.971	0.15
36	30.648°	2.915	0.10
37	31.078°	2.875	0.08
38	31.474°	2.840	0.08

TABLE 23a

XRPD Peak data for hydrochloride pattern 2. (2. d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.45°	10.45	0.01
2	11.79°	7.50	0.17
3	12.65°	6.99	0.01
4	13.04°	6.78	0.25
5	13.79°	6.42	0.02
6	14.60°	6.06	0.01
7	14.97°	5.91	0.49
8	16.12°	5.49	0.02
9	17.28°	5.13	0.14
10	17.53°	5.06	0.29
11	17.79°	4.98	1.00
12	18.36°	4.83	0.05
13	18.53°	4.78	0.90
14	19.57°	4.53	0.37
15	19.86°	4.47	0.05
16	20.35°	4.36	0.18
17	20.88°	4.25	0.06
18	21.34°	4.16	0.08
19	22.24°	3.99	0.17
20	22.72°	3.91	0.38
21	23.32°	3.81	0.09
22	23.65°	3.76	0.66
23	24.75°	3.59	0.21
24	25.62°	3.47	0.15
25	26.22°	3.40	0.12
26	26.59°	3.35	0.51
27	26.95°	3.31	0.21
28	27.10°	3.29	0.30
29	27.44°	3.25	0.04
30	27.71°	3.22	0.02
31	28.05°	3.18	0.16
32	28.54°	3.13	0.11
33	28.81°	3.10	0.05
34	29.44°	3.03	0.05
35	30.06°	2.97	0.15
36	30.65°	2.92	0.10
37	31.08°	2.88	0.08
38	31.47°	2.84	0.08

90

TABLE 23b

XRPD Peak data for hydrochloride pattern 2. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.5°	10.5	0.0
2	11.8°	7.5	0.2
3	12.7°	7.0	0.0
4	13.0°	6.8	0.3
5	13.8°	6.4	0.0
6	14.6°	6.1	0.0
7	15.0°	5.9	0.5
8	16.1°	5.5	0.0
9	17.3°	5.1	0.1
10	17.5°	5.1	0.3
11	17.8°	5.0	1.0
12	18.4°	4.8	0.1
13	18.5°	4.8	0.9
14	19.6°	4.5	0.4
15	19.9°	4.5	0.1
16	20.4°	4.4	0.2
17	20.9°	4.3	0.1
18	21.3°	4.2	0.1
19	22.2°	4.0	0.2
20	22.7°	3.9	0.4
21	23.3°	3.8	0.1
22	23.7°	3.8	0.7
23	24.8°	3.6	0.2
24	25.6°	3.5	0.2
25	26.2°	3.4	0.1
26	26.6°	3.4	0.5
27	27.0°	3.3	0.2
28	27.1°	3.3	0.3
29	27.4°	3.3	0.0
30	27.7°	3.2	0.0
31	28.1°	3.2	0.2
32	28.5°	3.1	0.1
33	28.8°	3.1	0.1
34	29.4°	3.0	0.1
35	30.1°	3.0	0.2
36	30.7°	2.9	0.1
37	31.1°	2.9	0.1
38	31.5°	2.8	0.1

The pattern 2 material was analysed further by TGA and DSC. The TGA thermogram is shown in FIG. 151 and demonstrates that the material appears to be a non-solvated/hydrated form with gross decomposition starting at around 256° C. The DSC thermogram in FIG. 152 shows a small endothermic event with an onset of 136.8° C. followed by an exothermic event with an onset of 139.1° C. followed by a final large endothermic event with an onset of 146.1° C. It would appear that the pattern 2 material melts, recrystallizes to pattern 1 and then melts again. The pattern 2 material was analysed by <sup>1</sup>H NMR and this showed that the material was still a salt, and only contained ~0.018 eq. of dioxane confirming it is not a solvated form. The <sup>1</sup>H NMR spectrum is shown in FIG. 153.

In one embodiment, there is provided crystalline 5-MeO-DMT hydrochloride or a pharmaceutical composition comprising crystalline 5-MeO-DMT hydrochloride, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 150;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more,



twenty nine or more, thirty or more, thirty one or more, thirty two or more, thirty three or more, thirty four or more, thirty five or more, thirty six or more, thirty seven or more or thirty eight peaks in an XRPD diffractogram as detailed in Table 23, 23a or 23b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 23, 23a or 23b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 151;

A DSC thermogram as shown in, or substantially as shown in, FIG. 152;

A small endothermic event with an onset of around 136.8°

C. followed by an exothermic event with an onset of around 139.1° C. followed by a final large endothermic event with an onset of around 146.1° C.;

A small endothermic event with an onset of 134-138° C. followed by an exothermic event with an onset of 137-141° C. followed by a final large endothermic event with an onset of 144-148° C.;

A small endothermic event with an onset of 134, 135, 136, 137 or 138° C. followed by an exothermic event with an onset of 137, 138, 139, 140 or 141° C. followed by a final large endothermic event with an onset of 144, 145, 146, 147 or 148° C.; and/or

An <sup>1</sup>H NMR spectrum as shown, or substantially as shown, in FIG. 153.

The XRPD for the new pattern 3 is shown in FIG. 154 compared to the pattern 1 and pattern 2. The XRPD peak data is shown in the table below:

TABLE 24

XRPD Peak data for hydrochloride pattern 3			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	6.509°	13.568	0.06
2	12.341°	7.166	0.04
3	12.979°	6.815	1.00
4	14.420°	6.137	0.03
5	16.706°	5.303	0.21
6	17.643°	5.023	0.10
7	17.805°	4.978	0.16
8	19.519°	4.544	0.25
9	20.226°	4.387	0.09
10	21.056°	4.216	0.08
11	21.882°	4.059	0.03
12	22.175°	4.006	0.44
13	23.525°	3.779	0.34
14	24.239°	3.669	0.11
15	25.454°	3.496	0.17
16	26.101°	3.411	0.19
17	27.201°	3.276	0.06
18	28.161°	3.166	0.14
19	28.408°	3.139	0.05
20	29.220°	3.054	0.12
21	29.841°	2.992	0.02
22	30.361°	2.942	0.03
23	30.876°	2.894	0.02

TABLE 24a

XRPD Peak data for hydrochloride pattern 3 (2 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	6.51°	13.57	0.06
2	12.34°	7.17	0.04
3	12.98°	6.82	1.00

TABLE 24a-continued

XRPD Peak data for hydrochloride pattern 3 (2 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
4	14.42°	6.14	0.03
5	16.71°	5.30	0.21
6	17.64°	5.02	0.10
7	17.81°	4.98	0.16
8	19.52°	4.54	0.25
9	20.23°	4.39	0.09
10	21.06°	4.22	0.08
11	21.88°	4.06	0.03
12	22.18°	4.01	0.44
13	23.53°	3.78	0.34
14	24.24°	3.67	0.11
15	25.45°	3.50	0.17
16	26.10°	3.41	0.19
17	27.20°	3.28	0.06
18	28.16°	3.17	0.14
19	23.41°	3.14	0.05
20	29.22°	3.05	0.12
21	29.84°	2.99	0.02
22	30.36°	2.94	0.03
23	30.88°	2.89	0.02

TABLE 24b

XRPD Peak data for hydrochloride pattern 3. (1 d.p.)			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	6.5°	13.6	0.1
2	12.3°	7.2	0.0
3	13.0°	6.8	1.0
4	14.4°	6.1	0.0
5	16.7°	5.3	0.2
6	17.6°	5.0	0.1
7	17.8°	5.0	0.2
8	19.5°	4.5	0.3
9	20.2°	4.4	0.1
10	21.1°	4.2	0.1
11	21.9°	4.1	0.0
12	22.2°	4.0	0.4
13	23.5°	3.8	0.3
14	24.2°	3.7	0.1
15	25.5°	3.5	0.2
16	26.1°	3.4	0.2
17	27.2°	3.3	0.1
18	28.2°	3.2	0.1
19	28.4°	3.1	0.1
20	29.2°	3.1	0.1
21	29.8°	3.0	0.0
22	30.4°	2.9	0.0
23	30.9°	2.9	0.0

The pattern 3 material was analysed further by TGA and DSC. The TGA thermogram is shown in FIG. 155 and demonstrates that the material appears to be a solvated/hydrated form with gross decomposition starting at around 260° C. The DSC thermogram shown in FIG. 156 shows a large endothermic event with an onset of 49.0° C. which corresponds with the weight loss observed in the TGA.

There are no other significant thermal events observed apart from degradation.

The pattern 3 material was analysed by <sup>1</sup>H NMR and this showed that the material was still a salt, and only contained a trace of dioxane and no methanol confirming it is not a solvated form. The <sup>1</sup>H NMR spectrum is shown in FIG. 157. The weight loss in the TGA and lack of solvent in the <sup>1</sup>H NMR suggest this material is a hydrated form.

93

In one embodiment, there is provided crystalline 5-MeO-DMT hydrochloride or a pharmaceutical composition comprising crystalline 5-MeO-DMT hydrochloride, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 154;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more or twenty three peaks in an XRPD diffractogram as detailed in Table 24, 24a or 24b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 24, 24a or 24b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 155;

A DSC thermogram as shown in, or substantially as shown in, FIG. 156;

A large endothermic event with an onset of 49.0° C., as measured in a DSC thermogram;

A large endothermic event with an onset of 46-52° C., as measured in a DSC thermogram;

A large endothermic event with an onset of 46, 47, 48, 49, 50, 51 or 52° C., as measured in a DSC thermogram; and/or

An <sup>1</sup>H NMR spectrum as shown, or substantially as shown, in FIG. 157.

#### Evaporations of Hydrochloride Salt

Twelve samples of the lyophilised material were treated with a minimal volume (ca 0.5 ml) of warm solvent, syringe filtered through a 0.22 m filter and added to a clean HPLC vial and allowed to evaporate at RT. The solvents used and XRPD results following evaporation are detailed in the Table below:


#### Summary of Results from Evaporations of Hydrochloride Salt

Experiment	Solvent	XRPD Results
DJP2202-011-01	Ethylene glycol	Failed to evaporate fully
DJP2202-011-02	1-PrOH	Pattern 1
DJP2202-011-03	Water	Failed to evaporate fully
DJP2202-011-04	MIBK	Insufficient material
DJP2202-011-05	EtOH	Pattern 1
DJP2202-011-06	IPA	Pattern 1
DJP2202-011-07	THF	Insufficient material
DJP2202-011-08	Dioxane	Insufficient material
DJP2202-011-09	Chlorobenzene	Insufficient material
DJP2202-011-10	IPAc	Insufficient material
DJP2202-011-11	MEK	Pattern 1
DJP2202-011-12	MeCN	Oil

The low solubility in many of the solvents meant that there was often insufficient solid for XRPD analysis. The XRPD diffractograms for the solids obtained can be seen in FIG. 158.

94

#### Example 5: Further Characterisation of 5-MeO-DMT Benzoate



To render the amorphous and thus remove seeds of the supplied pattern 1 the supplied material was lyophilised in an attempt to render it amorphous. To 2.396 g of the supplied benzoate salt lot 800674000 dioxane (55 ml) and water (5 ml) was added. The mixture was agitated gently and warmed to aid dissolution. The clear solution was then divided equally between 60 HPLC vials (~40 mg salt in each vial) and the vials were then frozen at ~-18° C. overnight and then dried by lyophilisation overnight. One sample was analysed by XRPD and the diffractogram shown in FIG. 159 and shows this to be a new crystalline form, designated as pattern 2. The XRPD peak data is shown in the Tables 25, 25a or 25b below:

TABLE 25

XRPD Peak data for benzoate pattern 2			
Peak No.	Angle 2 θ	d Value	Rel. Intensity
1	6.451°	13.689	0.12
2	9.114°	9.696	0.13
3	9.395°	9.406	0.01
4	12.880°	6.868	0.45
5	13.109°	6.748	0.13
5	14.615°	6.056	0.16
7	15.976°	5.543	0.04
8	18.405°	4.817	0.99
9	13.894°	4.693	0.09
10	19.512°	4.546	1.00
11	19.971°	4.442	0.06
12	20.481°	4.333	0.10
13	21.545°	4.121	0.11
14	22.506°	3.947	0.09
15	22.907°	3.879	0.41
16	23.429°	3.794	0.09
17	24.254°	3.667	0.23
18	25.531°	3.486	0.11
19	25.911°	3.436	0.03
20	26.360°	3.378	0.02
21	26.812°	3.322	0.07
22	27.149°	3.282	0.06
23	27.599°	3.229	0.08
24	29.065°	3.070	0.11
25	29.420°	3.034	0.03
26	30.607°	2.919	0.01
27	31.288°	2.857	0.06

TABLE 25a

XRPD Peak data for benzoate pattern 2 (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	6.45°	13.69	0.12
2	9.11°	9.70	0.13
3	9.40°	9.41	0.01
4	12.88°	6.87	0.45
5	13.11°	6.75	0.13
6	14.62°	6.06	0.16
7	15.98°	5.54	0.04
8	18.41°	4.82	0.99
9	18.89°	4.69	0.09
10	19.51°	4.55	1.00
11	19.97°	4.44	0.06
12	20.48°	4.33	0.10
13	21.55°	4.12	0.11
14	22.51°	3.95	0.09
15	22.91°	3.88	0.41
16	23.43°	3.79	0.09
17	24.25°	3.67	0.23
18	25.53°	3.49	0.11
19	25.91°	3.44	0.03
20	26.36°	3.38	0.02
21	26.81°	3.32	0.07
22	27.15°	3.28	0.06
23	27.60°	3.23	0.08
24	29.07°	3.07	0.11
25	29.42°	3.03	0.03
26	30.61°	2.92	0.01
27	31.29°	2.86	0.06

TABLE 25b

XRPD Peak data for benzoate pattern 2 (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	6.5°	13.7	0.1
2	9.1°	9.7	0.1
3	9.4°	9.4	0.0
4	12.9°	6.9	0.5
5	13.1°	6.8	0.1
6	14.6°	6.1	0.2
7	16.0°	5.5	0.0
8	18.4°	4.8	1.0
9	18.9°	4.7	0.1
10	19.5°	4.6	1.0
11	20.0°	4.4	0.1
12	20.5°	4.3	0.1
13	21.6°	4.1	0.1
14	22.5°	4.0	0.1
15	22.9°	3.9	0.4
16	23.4°	3.8	0.1
17	24.3°	3.7	0.2
18	25.5°	3.5	0.1
19	25.9°	3.4	0.0
20	26.4°	3.4	0.0
21	26.8°	3.3	0.1
22	27.2°	3.3	0.1
23	27.6°	3.2	0.1
24	29.1°	3.1	0.1
25	29.4°	3.0	0.0
26	30.6°	2.9	0.0
27	31.3°	2.9	0.1

The samples were used in further experiments.

The pattern 2 material was analysed further by TGA and DSC and the TGA thermogram is shown in FIG. 160. This shows the material is non-hydrated non-solvated form and shows good thermal stability to ~170° C. The DSC thermogram is shown in FIG. 161 and shows two endothermic

events very close to each other. The first has an onset of 119.6° C. and an enthalpy of 32.6 J/g and the second has an onset of 123.2° C. and an enthalpy of 34.5 J/g. It should be noted that the enthalpy values are probably under reported due to the integration algorithm, and that both of these endothermic events are at higher temperature compared to the pattern 1 material.

The material was analysed by <sup>1</sup>H NMR and the spectrum (FIG. 162) confirmed that the material is non solvated and still the benzoate salt and not a free form. Only traces of dioxane, ~0.04 eq. were observed.

In one embodiment, there is provided crystalline 5-MeO-DMT benzoate or a pharmaceutical composition comprising crystalline 5-MeO-DMT benzoate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 159;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more or twenty seven peaks in an XRPD diffractogram as detailed in Table 25, 25a or 25b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 25, 25a or 25b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 160;

A DSC thermogram as shown in, or substantially as shown in, FIG. 161;

Two endothermic events very close to each other, the first having an onset of around 119.6° C. and an enthalpy of around 32.6 J/g and the second having an onset of around 123.2° C. and an enthalpy of around 34.5 J/g, as measured in a DSC thermogram;

Two endothermic events very close to each other, the first having an onset of around 117-122° C. and an enthalpy of around 30-33 J/g and the second having an onset of around 122-125° C. and an enthalpy of around 33-36 J/g, as measured in a DSC thermogram;

Two endothermic events very close to each other, the first having an onset of around 117, 118, 119, 120, 121 or 122° C. and an enthalpy of around 30, 31, 32 or 33 J/g and the second having an onset of around 122, 123, 124 or 125° C. and an enthalpy of around 33, 34, 35 or 36 J/g, as measured in a DSC thermogram; and/or

An <sup>1</sup>H NMR spectrum as shown, or substantially as shown, in FIG. 162.

Thermal Cycling of Benzoate Salt

Twenty five of the lyophilised samples were treated with solvent and thermally cycled between ambient and 40° C. with four hours spent under each condition. After three days and solids were isolated by centrifuge filtration and analysed by XRPD. Any solutions were allowed to evaporate at RT but this did not yield any new solids. The solvents, observations, isolation and XRPD results are summarised in the Table below:

Summary of thermal cycle of benzoate salt

Experiment	Solvent	Volume added $\mu$ l	Initial Observations	3-day Observations	Isolation	XRPD Result
DJP2202-006-01	1,4-Dioxane	200	no change	solid mass	After thermal cycle	Pattern 3
DJP2202-006-02	1-Propanol	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-03	2-Butanol	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-04	2-Ethoxyethanol	200	dissolved	clear solution	Evaporated to oil	n/a
DJP2202-006-05	2-Methyltetrahydrofuran	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-06	2-Propanol	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-07	Acetone	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-08	Acetonitrile	200	no change	lump of material	After thermal cycle	Pattern 4
DJP2202-006-09	Anisole	200	no change	lump of material	After thermal cycle	Pattern 4
DJP2202-006-10	Chlorobenzene	200	no change	white solid	After thermal cycle	Pattern 4 + extra peaks
DJP2202-006-11	Ethanol	150	no change	white solid	After thermal cycle	Pattern 4
DJP2202-006-12	Ethyl acetate	200	no change	white solid	After thermal cycle	Pattern 4
DJP2202-006-13	Isopropyl acetate	200	no change	white solid	After thermal cycle	Pattern 4
DJP2202-006-14	Methanol	200	dissolved	clear solution	Evaporated to solid	Pattern 1
DJP2202-006-15	Propyl acetate	200	no change	white solid	After thermal cycle	Pattern 4
DJP2202-006-16	Methylethyl ketone	200	no change	white crystals	After thermal cycle	Pattern 4
DJP2202-006-17	Formamide	150	dissolved	few crystals	After thermal cycle	Pattern 4
DJP2202-006-18	N,N-Dimethylacetamide	200	dissolved	clear solution	Evaporated-no solid	n/a
DJP2202-006-19	N-Methylpyrrolidone	200	no change	clear solution	Evaporated-no solid	n/a
DJP2202-006-20	iso-Propyl acetate	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-21	2-Me-1-PrOH	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-22	Tetrahydrofuran	200	no change	few crystals	After thermal cycle	Pattern 4
DJP2202-006-23	MeOH/water 953/47 Calc Aw 0.214	200	dissolved	clear solution	Evaporated to oil	n/a
DJP2202-006-24	MeOH/water 693/307 Calc Aw 0.599	200	dissolved	clear solution	Evaporated to solid	Pattern 1
DJP2202-006-25	MeOH/water 360/640 Calc Aw 0.821	200	dissolved	clear solution	Evaporated to solid	Pattern 1

The XRPD for the new pattern 3 is shown in FIG. 164 compared to the supplied pattern 1 and pattern 2. The XRPD peak data is shown in the Tables 26, 26a or 26b below:

TABLE 26

XRPD Peak pick of Benzoate pattern 3			
Peak No.	Angle $2\theta$	d Value	Rel. Intensity
1	5.193°	17.002	0.29
2	6.122°	14.426	0.33
3	9.563°	9.241	0.01
4	10.220°	8.649	0.03
5	11.373°	7.774	0.03
6	14.018°	6.313	0.01
7	14.806°	5.979	0.02
8	15.320°	5.779	0.16

TABLE 26-continued

XRPD Peak pick of Benzoate pattern 3				
	Peak No.	Angle $2\theta$	d Value	Rel. Intensity
55	9	18.254°	4.856	0.93
	10	19.260°	4.605	0.12
	11	20.450°	4.339	1.00
	12	22.663°	3.920	0.02
60	13	23.666°	3.756	0.04
	14	24.400°	3.645	0.18
	15	25.593°	3.478	0.08
	16	26.232°	3.395	0.01
	17	27.407°	3.252	0.01
	18	29.826°	2.993	0.01
65	19	30.812°	2.900	0.01

TABLE 26a

XRPD Peak pick of Benzoate pattern 3. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	5.19°	17.00	0.29
2	6.12°	14.43	0.33
3	9.56°	9.24	0.01
4	10.22°	8.65	0.03
5	11.37°	7.77	0.03
6	14.02°	6.31	0.01
7	14.81°	5.98	0.02
8	15.32°	5.78	0.16
9	18.25°	4.86	0.93
10	19.26°	4.61	0.12
11	20.45°	4.34	1.00
12	22.66°	3.92	0.02
13	23.67°	3.76	0.04
14	24.40°	3.65	0.18
15	25.59°	3.48	0.08
16	26.23°	3.40	0.01
17	27.41°	3.25	0.01
18	29.83°	2.99	0.01
19	30.81°	2.90	0.01

TABLE 26b

XRPD Peak pick of Benzoate pattern 3. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	5.2°	17.0	0.3
2	6.1°	14.4	0.3
3	9.6°	9.2	0.0
4	10.2°	8.7	0.0
5	11.4°	7.8	0.0
6	14.0°	6.3	0.0
7	14.8°	6.0	0.0
8	15.3°	5.8	0.2
9	18.3°	4.9	0.9
10	19.3°	4.6	0.1
11	20.5°	4.3	1.0
12	22.7°	3.9	0.0
13	23.7°	3.3	0.0
14	24.4°	3.7	0.2
15	25.6°	3.5	0.1
16	26.2°	3.4	0.0
17	27.4°	3.3	0.0
18	29.8°	3.0	0.0
19	30.8°	2.9	0.0

The new pattern 3 material was analysed further by TGA and DSC. The TGA thermogram is shown in FIG. 164 and shows an initial loss of 9.77% which would equate to 0.42 eq of dioxane. The material then undergoes complete mass loss starting at approximately 190° C. The DSC thermogram is shown in FIG. 165. This shows a single endothermic event with an onset of 123.2° C., which is similar to the second thermal event observed in the pattern 2 material.

The material was analysed further by <sup>1</sup>H NMR and the spectrum is shown in FIG. 166. This shows the material is still the benzoate salt, but also shows ~1.22 eq of dioxane indicating this is probably a solvated form of the material.

In one embodiment, there is provided crystalline 5-MeO-DMT benzoate or a pharmaceutical composition comprising crystalline 5-MeO-DMT benzoate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 163;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or

more, sixteen or more, seventeen or more, eighteen or more or nineteen peaks in an XRPD diffractogram as detailed in Table 26, 26a or 26b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 26, 26a or 26b;

A TGA thermogram as shown in, or substantially as shown in, FIG. 164;

A DSC thermogram as shown in, or substantially as shown in, FIG. 165;

A single endothermic event with an onset of around 123.2° C., as measured in a DSC thermogram;

A single endothermic event with an onset of around 120-125° C., as measured in a DSC thermogram;

A single endothermic event with an onset of around 120, 121, 122, 123, 124 or 125° C., as measured in a DSC thermogram; and/or

An <sup>1</sup>H NMR spectrum as shown, or substantially as shown, in FIG. 166.

The XRPD for the new pattern 4 is shown in FIG. 167 compared to the supplied pattern 1, pattern 2 and pattern 3. The XRPD peak data is shown in the Tables below:

TABLE 27

XRPD Peak pick of benzoate pattern 4			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.761°	10.085	0.06
2	9.053°	9.760	1.00
3	10.763°	8.213	0.04
4	11.411°	7.748	0.10
5	12.275°	7.205	0.08
6	14.401°	6.146	0.02
7	16.154°	5.482	0.15
8	16.409°	5.398	0.14
9	17.439°	5.081	0.39
10	17.745°	4.994	0.11
11	18.029°	4.916	0.03
12	18.387°	4.821	0.22
13	20.033°	4.429	0.04
14	20.514°	4.326	0.01
15	20.823°	4.262	0.29
16	21.519°	4.126	0.02
17	21.783°	4.077	0.02
18	22.116°	4.016	0.01
19	22.738°	3.908	0.20
20	23.039°	3.857	0.03
21	24.222°	3.672	0.01
22	24.640°	3.610	0.18
23	25.215°	3.529	0.21
24	26.134°	3.407	0.01
25	26.371°	3.377	0.04
26	27.153°	3.281	0.10
27	28.203°	3.162	0.03
28	28.730°	3.105	0.02
29	30.215°	2.955	0.06
30	30.822°	2.899	0.02
31	31.411°	2.846	0.01
32	31.695°	2.821	0.01

TABLE 27a

XRPD Peak pick of benzoate pattern 4. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.76°	10.09	0.06
2	9.05°	9.76	1.00
3	10.76°	8.21	0.04
4	11.41°	7.75	0.10
5	12.28°	7.21	0.08

TABLE 27a-continued

XRPD Peak pick of benzoate pattern 4. (2 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
6	14.40°	6.15	0.02
7	16.15°	5.48	0.15
8	16.41°	5.40	0.14
9	17.44°	5.08	0.39
10	17.75°	4.99	0.11
11	18.03°	4.92	0.03
12	18.39°	4.82	0.22
13	20.03°	4.43	0.04
14	20.51°	4.33	0.01
15	20.82°	4.26	0.29
16	21.52°	4.13	0.02
17	21.78°	4.08	0.02
18	22.12°	4.02	0.01
19	22.74°	3.91	0.20
20	23.04°	3.86	0.03
21	24.22°	3.67	0.01
22	24.64°	3.61	0.18
23	25.22°	3.53	0.21
24	26.13°	3.41	0.01
25	26.37°	3.38	0.04
26	27.15°	3.28	0.10
27	28.20°	3.16	0.03
28	28.73°	3.11	0.02
29	30.22°	2.96	0.06
30	30.82°	2.90	0.02
31	31.41°	2.85	0.01
32	31.70°	2.82	0.01

TABLE 27b

XRPD Peak pick of benzoate pattern 4. (1 d.p.)			
Peak No.	Angle 2 $\theta$	d Value	Rel. Intensity
1	8.8°	10.1	0.1
2	9.1°	9.8	1.0
3	10.8°	8.2	0.0
4	11.4°	7.8	0.1
5	12.3°	7.2	0.1
6	14.4°	6.2	0.0
7	16.2°	5.5	0.2
8	16.4°	5.4	0.1
9	17.4°	5.1	0.4
10	17.8°	5.0	0.1
11	18.0°	4.9	0.0
12	18.4°	4.8	0.2
13	20.0°	4.4	0.0
14	20.5°	4.3	0.0
15	20.8°	4.3	0.3
16	21.5°	4.1	0.0
17	21.8°	4.1	0.0
18	22.1°	4.0	0.0
19	22.7°	3.9	0.2
20	23.0°	3.9	0.0
21	24.2°	3.7	0.0
22	24.6°	3.6	0.2
23	25.2°	3.5	0.2
24	26.1°	3.4	0.0
25	26.4°	3.4	0.0
26	27.2°	3.3	0.1
27	28.2°	3.2	0.0
28	28.7°	3.1	0.0
29	30.2°	3.0	0.1
30	30.8°	2.9	0.0
31	31.4°	2.9	0.0
32	31.7°	2.8	0.0

The new pattern 4 material was analysed further by TGA and DSC. The TGA thermogram is shown in FIG. 168 and shows the material to be an anhydrous non solvated form with good thermal stability to ~190°. The material then undergoes complete mass loss starting at approximately 190° C. The DSC thermogram is shown in FIG. 169. This

shows a single endothermic event with an onset of 123.4° C., which is similar to the second thermal event observed in the pattern 2 material.

The material was analysed further by <sup>1</sup>H NMR and the spectrum is shown in FIG. 170. This shows the material is still the benzoate salt and only shows trace amounts of the process solvent (2-BuOH).

In one embodiment, there is provided crystalline 5-MeO-DMT benzoate or a pharmaceutical composition comprising crystalline 5-MeO-DMT benzoate, as characterised by one or more of:

An XRPD pattern as shown in, or substantially as shown in, FIG. 167;

One or more, two or more, three or more, four or more, five or more, six or more, seven or more, eight or more, nine or more, ten or more, eleven or more, twelve or more, thirteen or more, fourteen or more, fifteen or more, sixteen or more, seventeen or more, eighteen or more, nineteen or more, twenty or more, twenty one or more, twenty two or more, twenty three or more, twenty four or more, twenty five or more, twenty six or more, twenty seven or more, twenty eight or more, twenty nine or more, thirty or more, thirty one or more or thirty two peaks in an XRPD diffractogram as detailed in Table 27, 27a or 27b;

One or more, two or more, three or more, four or more or five or more peaks in an XRPD diffractogram with a relative intensity of over 0.1, 0.2, 0.3, 0.4, 0.5, 0.6, 0.7, 0.8 or 0.9 as detailed in Table 27, 27a or 267;

A TGA thermogram as shown in, or substantially as shown in, FIG. 168;

A DSC thermogram as shown in, or substantially as shown in, FIG. 169;

A single endothermic event with an onset of around 123.4° C., as measured in a DSC thermogram;

A single endothermic event with an onset of around 120-125° C., as measured in a DSC thermogram;

A single endothermic event with an onset of around 120, 121, 122, 123, 124 or 125° C., as measured in a DSC thermogram; and/or

An <sup>1</sup>H NMR spectrum as shown, or substantially as shown, in FIG. 170.

#### Evaporations of Benzoate Salt

Twelve samples of the lyophilised material were treated with a minimal volume (ca 0.5 ml) of warm solvent, syringe filtered through a 0.22 micrometre filter and added to a clean HPLC vial and allowed to evaporate at RT. The solvents used and XRPD results following evaporation are detailed in the Table below:

#### Summary of Results from Evaporations of Benzoate Salt

Experiment	Solvent	XRPD Result
DJP2202-010-01	Ethylene glycol	Failed to evaporate fully
DJP2202-010-02	1-PrOH	Oil
DJP2202-010-03	Water	Failed to evaporate fully
DJP2202-010-04	MIBK	Oil
DJP2202-010-05	EtOH	Oil
DJP2202-010-06	IPA	Pattern 4
DJP2202-010-07	THF	Pattern 4
DJP2202-010-08	dioxane	Pattern 4

103

-continued

Experiment	Solvent	XRPD Result
DJP2202-010-09	chlorobenzene	Pattern 4
DJP2202-010-10	IPAc	Pattern 2
DJP2202-010-11	MEK	Pattern 4
DJP2202-010-12	MeCN	Pattern 4

#### Antisolvent Additions of Benzoate Salt

For antisolvent additions thirteen samples of the lyophilised material were dissolved in a small amount of hot solvent (~0.5 ml). Extra supplied material, ca 30 mg was added to each and the clear solution was syringe filtered through a 0.22 µm filter into a 20 ml scintillation vial. A large excess of antisolvent was then added, the vials sealed. After storage at RT overnight, and samples where no solid had been produced were first cooled to 4° C. for ~24 hrs, and again if no solid was produced, they were cooled to -18° C. Any solids were isolated by centrifuge filtration and analysed by XRPD. The solvents, antisolvents, isolation conditions and XRPD results are summarised in the Table below:

#### Summary of Results from Antisolvent Additions of Benzoate Salt

Experiment	Solvent	Anti Solvent	Observations	XRPD Results
DJP2202-012-01	MeOH/water Aw 0.2	MeOH	Cooled -18 C.	No solid
DJP2202-012-02	MeOH	TBME	Cooled -18 C.	No solid
DJP2202-012-03	EtOH	TBME	Cooled -18 C.	No solid
DJP2202-012-04	EtOH	Heptane	Some solid at RT	Pattern 1
DJP2202-012-05	THF	TBME	Cooled -18 C.	No solid
DJP2202-012-06	dioxane	TBME	Cooled -18 C.	No solid
DJP2202-012-07	DMF	PhMe	Cooled -18 C.	No solid
DJP2202-012-08	IPAc	TBME	Cooled -18 C.	No solid
DJP2202-012-09	MEK	TBME	Cooled -18 C.	No solid
DJP2202-012-10	MEK	Heptane	Some solid at RT	Pattern 2
DJP2202-012-11	MeCN	TBME	Cooled -18 C.	No solid
DJP2202-012-12	MeCN	PhMe	Some solid at RT	Pattern 4

#### Cooling of Benzoate Salt

For the cooling experiments twelve samples of the lyophilised material were dissolved in a small amount of hot solvent (~0.5 ml) and syringe filtered through a 0.22 µm filter into a clean HPLC vial. Samples were allowed to cool to RT, and if no solid was produced to 4° C. and then to -18° C. Solids were isolated by centrifuge filtration and analysed by XRPD. The solvents, isolation conditions and XRPD results are summarised in the Table below:

#### Summary of Results from Cooling Experiments of Benzoate Salt

Experiment	Solvent	Observations	XRPD Results
DJP2202-013-01	MeOH/water Aw 0.2	Cooled -18 C.	Pattern 4
DJP2202-013-02	MeOH/water Aw 0.6	Cooled -18 C.	no solids
DJP2202-013-03	Water	Cooled -18 C.	Pattern 4
DJP2202-013-04	MeOH	Cooled -18 C.	no solids
DJP2202-013-05	EtOH	Solid at RT	Pattern 4
DJP2202-013-06	IPA	Solid at RT	Pattern 4

104

-continued

Experiment	Solvent	Observations	XRPD Results
5 DJP2202-013-07	THF	Solid at RT	Pattern 4
DJP2202-013-08	dioxane	Solid at RT	Pattern 3
DJP2202-013-09	chlorobenzene	Solid at RT	Looks like mix #4 #2
DJP2202-013-10	IPAc	Solid at RT	Pattern 4
DJP2202-013-11	MEK	Solid at RT	Pattern 2
10 DJP2202-013-12	MeCN	Solid at RT	Pattern 4

#### Example 6: Solubility Experiments

Solubility assessments of 5-MeO-DMT oxalate, hydrobromide fumarate and benzoate were performed in four different media:

Media	100 mM buffer
SNF	Sodium chloride/Calcium chloride/Potassium chloride
pH 1.2 buffer	Potassium chloride/Hydrochloric acid
pH 4.5 buffer	Sodium acetate/Acetic acid
25 pH 6.8 buffer	Sodium phosphate dibasic/Potassium dihydrogen phosphate

#### Oxalate Salt (DXD2203-013-02) 250 mg per vial results

Media	pH after addition of 1.5 ml media at 37° C.	Observation
SNF	2.13	Suspension
pH 1.2	1.87	Suspension
pH 4.5	3.05	Suspension
35 pH 6.8	3.45	Suspension

#### Hydrobromide Salt (DXD2203-014-01) 250 mg per vial results

Media	pH after addition of 1.0 ml media at 37° C.	Observation
SNF	8.13	Clear solution
pH 1.2	8.09	Clear solution
45 pH 4.5	7.65	Clear solution
pH 6.8	7.82	Clear solution

The hydrobromide salt has a solubility of, at least, 250 mg/ml.

#### 50 Fumarate Salt (DXD2203-015-03) 250 mg per vial results

Media	pH after addition of 1.0 ml media at 37° C.	Observation
SNF	5.14	Clear solution
55 pH 1.2	5.24	Clear solution
pH 4.5	5.31	Clear solution
pH 6.8	5.79	Clear solution

60 The fumarate salt has a solubility of, at least, 250 mg/ml.  
Benzoate Salt (21/32/68/FP1) 50 mg per vial results

Media	pH after addition of 0.5 ml media at 37° C.	Observation
65 SNF	6.77	Clear solution
pH 1.2	5.28	Clear solution

-continued

Media	pH after addition of 0.5 ml media at 37° C.	Observation
pH 4.5	5.22	Clear solution
pH 6.8	6.82	Clear solution

The benzoate salt has a solubility of, at least, 100 mg/ml. The calibration curve of the free base was prepared between 0.031-0.500 mg/ml and can be seen in FIG. 172.

#### Oxalate Salt Solubility by HPLC

Mother liquors were adjusted to the desired pH before HPLC quantification.

Media	Solubility (mg/ml) of free base
SNF	20.22
pH 12.	19.02
pH 4.5	35.53
pH 6.8	26.85

Isolated solids were analyzed by XRPD and no change in form was observed.

#### CLAUSES

1. A non-hygroscopic salt of 5-MeO-DMT wherein the non-hygroscopic salt is 5-MeO-DMT hydrobromide.
2. A crystalline form of the non-hygroscopic salt of clause 1.
3. A crystalline form of the non-hygroscopic salt of clause 1, characterised by peaks in an XRPD diffractogram at 14.6, 16.8, 20.8, 24.3, 24.9 and 27.5° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
4. A crystalline form of the non-hygroscopic salt of clause 1, characterised by peaks in an XRPD diffractogram at 14.6, 21.6 and 24.3° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
5. A crystalline form of the non-hygroscopic salt of clause 1, characterised by peaks in an XRPD diffractogram at 18.6, 19.7 and 24.8° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
6. The crystalline form of the non-hygroscopic salt of clause 4, characterised by peaks in an XRPD diffractogram at 14.6, 20.8, 21.6, 24.3 and 25.4° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
7. A pharmaceutical composition comprising the non-hygroscopic salt of 5-MeO-DMT of clause 1.
8. A pharmaceutical composition comprising the crystalline form of the non-hygroscopic salt of 5-MeO-DMT of any one of clauses 3 to 6.
9. The pharmaceutical composition of clause 7 for use as a medicament.
10. The pharmaceutical composition of clause 8 for use as a medicament.
11. 5-MeO-DMT phosphate.
12. A crystalline form of the 5-MeO-DMT phosphate of clause 11.
13. A crystalline form of the 5-MeO-DMT phosphate of clause 11, characterised by peaks in an XRPD diffractogram at 12.9, 20.4 and 23.1° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

14. The crystalline form of the 5-MeO-DMT phosphate of clause 13, characterised by peaks in an XRPD diffractogram at 12.9, 14.4, 19.3, 20.4 and 23.1° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
15. A pharmaceutical composition comprising the 5-MeO-DMT phosphate of clause 11.
16. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT phosphate of clause 13 or clause 14.
17. The pharmaceutical composition of clause 15 for use as a medicament.
18. The pharmaceutical composition of clause 16 for use as a medicament.
19. 5-MeO-DMT fumarate.
20. A crystalline form of the 5-MeO-DMT fumarate of clause 19.
21. A crystalline form of the 5-MeO-DMT fumarate of clause 19, characterised by peaks in an XRPD diffractogram at 13.0, 16.3 and 22.1° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
22. A crystalline form of the 5-MeO-DMT fumarate of clause 21, characterised by peaks in an XRPD diffractogram at 13.0, 16.3, 19.2, 20.4 and 22.1° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
23. A pharmaceutical composition comprising the 5-MeO-DMT fumarate of clause 19.
24. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT fumarate of clause 21 or clause 22.
25. The pharmaceutical composition of clause 23 for use as a medicament.
26. The pharmaceutical composition of clause 24 for use as a medicament.
27. 5-MeO-DMT oxalate.
28. A crystalline form of the 5-MeO-DMT oxalate of clause 27.
29. A crystalline form of the 5-MeO-DMT oxalate of clause 28, characterised by peaks in an XRPD diffractogram at 13.0, 19.9 and 26.0° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
30. A crystalline form of the 5-MeO-DMT oxalate of clause 29, characterised by peaks in an XRPD diffractogram at 13.0, 14.0, 19.9, 22.0 and 26.0° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
31. A pharmaceutical composition comprising the 5-MeO-DMT oxalate of clause 27.
32. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT oxalate of clause 29 or clause 30.
33. The pharmaceutical composition of clause 31 for use as a medicament.
34. The pharmaceutical composition of clause 32 for use as a medicament.
35. 5-MeO-DMT tartrate.
36. A crystalline form of the 5-MeO-DMT tartrate of clause 35.
37. A crystalline form of the 5-MeO-DMT tartrate of clause 35, characterised by peaks in an XRPD diffractogram at 18.3, 18.6, and 20.7° 2θ±0.1° 2θ as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.



38. A crystalline form of the 5-MeO-DMT tartrate of clause 35, characterised by peaks in an XRPD diffractogram at 18.3, 18.6, 18.8, 20.3 and 20.7°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å. 5
39. A pharmaceutical composition comprising the 5-MeO-DMT tartrate of clause 35.
40. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT tartrate of clause 37 or clause 38. 10
41. The pharmaceutical composition of clause 39 for use as a medicament.
42. The pharmaceutical composition of clause 40 for use as a medicament. 15
43. 5-MeO-DMT benzenesulfonate.
44. A crystalline form of the 5-MeO-DMT benzenesulfonate of clause 43.
45. A crystalline form of the 5-MeO-DMT benzenesulfonate of clause 44, characterised by peaks in an XRPD diffractogram at 9.5, 21.2, and 23.6°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å. 20
46. A crystalline form of the 5-MeO-DMT benzenesulfonate of clause 45, characterised by peaks in an XRPD diffractogram at 9.5, 18.0, 21.2, 23.6 and 24.4°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å. 25
47. A pharmaceutical composition comprising the 5-MeO-DMT benzenesulfonate of clause 43. 30
48. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT benzenesulfonate of clause 45 or clause 46.
49. The pharmaceutical composition of clause 47 for use as a medicament. 35
50. The pharmaceutical composition of clause 48 for use as a medicament.
51. 5-MeO-DMT tosylate.
52. A crystalline form of the 5-MeO-DMT tosylate of clause 51. 40
53. A crystalline form of the 5-MeO-DMT tosylate of clause 51, characterised by peaks in an XRPD diffractogram at 19.3, 23.6 and 24.1°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å. 45
54. A crystalline form of the 5-MeO-DMT tosylate of clause 51, characterised by peaks in an XRPD diffractogram at 13.8, 19.3, 23.6, 24.1 and 27.3°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å. 50
55. A pharmaceutical composition comprising the 5-MeO-DMT tosylate of clause 51.
56. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT tosylate of clause 53 or clause 54. 55
57. The pharmaceutical composition of clause 55 for use as a medicament.
58. The pharmaceutical composition of clause 56 for use as a medicament.
59. 5-MeO-DMT glycolate. 60
60. A crystalline form of the 5-MeO-DMT glycolate of clause 59.
61. A crystalline form of the 5-MeO-DMT glycolate of clause 59, characterised by peaks in an XRPD diffractogram at 20.2, 21.1 and 23.4°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å. 65

62. A crystalline form of the 5-MeO-DMT glycolate of clause 61, characterised by peaks in an XRPD diffractogram at 10.1, 20.2, 21.1, 23.4 and 24.3°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
63. A pharmaceutical composition comprising the 5-MeO-DMT glycolate of clause 59.
64. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT glycolate of clause 61 or clause 62.
65. The pharmaceutical composition of clause 63 for use as a medicament.
66. The pharmaceutical composition of clause 64 for use as a medicament.
67. 5-MeO-DMT ketoglutarate.
68. A crystalline form of the 5-MeO-DMT ketoglutarate of clause 67.
69. A crystalline form of the 5-MeO-DMT ketoglutarate of clause 67, characterised by peaks in an XRPD diffractogram at 14.4, 18.2 and 20.9°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
70. A crystalline form of the 5-MeO-DMT ketoglutarate of clause 69, characterised by peaks in an XRPD diffractogram at 14.4, 18.2, 20.9, 22.5 and 25.6°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
71. A pharmaceutical composition comprising the 5-MeO-DMT ketoglutarate of clause 67.
72. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT ketoglutarate of clause 69 or clause 70.
73. The pharmaceutical composition of clause 71 for use as a medicament.
74. The pharmaceutical composition of clause 72 for use as a medicament.
75. 5-MeO-DMT malate.
76. A crystalline form of the 5-MeO-DMT malate of clause 75.
77. A crystalline form of the 5-MeO-DMT malate of clause 75, characterised by peaks in an XRPD diffractogram at 18.3, 18.7 and 18.9°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
78. A crystalline form of the 5-MeO-DMT malate of clause 77, characterised by peaks in an XRPD diffractogram at 18.3, 18.7, 18.9, 21.6 and 26.1°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
79. A pharmaceutical composition comprising the 5-MeO-DMT malate of clause 75.
80. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT malate of clause 77 or clause 78.
81. The pharmaceutical composition of clause 79 for use as a medicament.
82. The pharmaceutical composition of clause 80 for use as a medicament.
83. 5-MeO-DMT saccharinate.
84. A crystalline form of the 5-MeO-DMT saccharinate of clause 83.
85. A crystalline form of the 5-MeO-DMT saccharinate of clause 83, characterised by peaks in an XRPD diffractogram at 8.7, 15.2 and 20.9°  $2\theta \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.

## 109

86. A crystalline form of the 5-MeO-DMT saccharinate of clause 85, characterised by peaks in an XRPD diffractogram at 5.2, 8.7, 15.0, 15.2 and  $20.9 \pm 0.1^\circ$   $2\theta$  as measured by x-ray powder diffraction using an x-ray wavelength of 1.5406 Å.
87. A pharmaceutical composition comprising the 5-MeO-DMT saccharinate of clause 85.
88. A pharmaceutical composition comprising the crystalline form of 5-MeO-DMT saccharinate of clause 85 or clause 86.
89. The pharmaceutical composition of clause 87 for use as a medicament.
90. The pharmaceutical composition of clause 88 for use as a medicament.

The invention claimed is:

1. A non-hygroscopic salt of 5-methoxy-N,N-dimethyltryptamine (5-MeO-DMT), wherein the non-hygroscopic salt is 5-MeO-DMT hydrobromide and the non-hygroscopic salt is in crystalline form.

2. The non-hygroscopic salt of claim 1, characterised by peaks in an x-ray powder diffraction (XRPD) diffractogram at  $2\theta$  values of  $14.6 \pm 0.1^\circ$ ,  $16.8 \pm 0.1^\circ$ ,  $20.8 \pm 0.1^\circ$ ,  $24.3 \pm 0.1^\circ$ ,  $24.9 \pm 0.1^\circ$ , and  $27.5 \pm 0.1^\circ$  as measured using an x-ray wavelength of 1.5406 Å.

## 110

3. The non-hygroscopic salt of claim 1, characterised by peaks in an x-ray powder diffraction (XRPD) diffractogram at  $2\theta$  values of  $14.6 \pm 0.1^\circ$ ,  $21.6 \pm 0.1^\circ$ , and  $24.3 \pm 0.1^\circ$  as measured using an x-ray wavelength of 1.5406 Å.
4. The non-hygroscopic salt of claim 1, characterised by peaks in an x-ray powder diffraction (XRPD) diffractogram at  $2\theta$  values of  $18.6 \pm 0.1^\circ$ ,  $19.7 \pm 0.1^\circ$ , and  $24.8 \pm 0.1^\circ$  as measured using an x-ray wavelength of 1.5406 Å.
5. The non-hygroscopic salt of claim 3, characterised by peaks in an x-ray powder diffraction (XRPD) diffractogram at  $2\theta$  values of  $14.6 \pm 0.1^\circ$ ,  $20.8 \pm 0.1^\circ$ ,  $21.6 \pm 0.1^\circ$ ,  $24.3 \pm 0.1^\circ$ , and  $25.4 \pm 0.1^\circ$  as measured using an x-ray wavelength of 1.5406 Å.
6. A pharmaceutical composition comprising the non-hygroscopic salt of claim 1.
7. A pharmaceutical composition comprising the non-hygroscopic salt of claim 2.
8. A pharmaceutical composition comprising the non-hygroscopic salt of claim 3.
9. A pharmaceutical composition comprising the non-hygroscopic salt of claim 4.
10. A pharmaceutical composition comprising the non-hygroscopic salt of claim 5.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**


PATENT NO. : 11,773,063 B1  
APPLICATION NO. : 17/941410  
DATED : October 3, 2023  
INVENTOR(S) : Jason Gray et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

In the Claims

In Claim 4, Column 110, Line 7, delete "20" and insert -- 20 --.

Signed and Sealed this  
Twenty-first Day of November, 2023  


Katherine Kelly Vidal  
*Director of the United States Patent and Trademark Office*