



US011970500B1

(12) **United States Patent**
Wang et al.

(10) **Patent No.:** **US 11,970,500 B1**
(45) **Date of Patent:** ***Apr. 30, 2024**

(54) **CRYSTALLINE FORM OF
(S)-7-(1-ACRYLOYLPYPERIDIN-4-YL)-
2-(4-PHENOXYPHENYL)-4,5,6,7-TETRA-
HYDOPYRAZOLO[1,5-A]PYRIMIDINE-
3-CARBOXAMIDE, PREPARATION, AND
USES THEREOF**

(52) **U.S. Cl.**
CPC *C07D 487/04* (2013.01); *A61P 35/02*
(2018.01); *A61P 35/04* (2018.01); *C07B*
2200/13 (2013.01)

(58) **Field of Classification Search**
CPC *A61P 35/02*; *A61P 35/04*; *A61P 35/00*;
C07B 2200/13; *C07D 487/04*;
(Continued)

(71) Applicant: **BEIGENE SWITZERLAND GMBH,**
Basel (CH)

(56) **References Cited**

(72) Inventors: **Zhiwei Wang,** Beijing (CN); **Yunhang
Guo,** Beijing (CN); **Gongyin Shi,**
Beijing (CN); **Lai Wang,** Beijing (CN)

U.S. PATENT DOCUMENTS

(73) Assignee: **BeiGene Switzerland GmbH,** Basel
(CH)

7,393,848 B2 7/2008 Currie et al.
7,514,444 B2 4/2009 Honigberg et al.
(Continued)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

FOREIGN PATENT DOCUMENTS

This patent is subject to a terminal dis-
claimer.

CN 1771231 A 5/2006
CN 102656173 A 9/2012
(Continued)

(21) Appl. No.: **18/526,535**

OTHER PUBLICATIONS

(22) Filed: **Dec. 1, 2023**

Armour, K. et al., "Recombinant human IgG molecules lacking Fcγ receptor I binding and monocyte triggering activities," *Eur. J. Immunol.*, Aug. 1999, vol. 29, No. 8, pp. 2613-2624.
(Continued)

Related U.S. Application Data

(63) Continuation of application No. 17/858,826, filed on
Jul. 6, 2022, now Pat. No. 11,884,674, which is a
(Continued)

Primary Examiner — Savitha M Rao

(74) *Attorney, Agent, or Firm* — Ice Miller LLP

(30) **Foreign Application Priority Data**


Aug. 16, 2016 (WO) PCT/CN2016/09510

(57) **ABSTRACT**

(51) **Int. Cl.**
C07D 487/04 (2006.01)
A61P 35/02 (2006.01)
A61P 35/04 (2006.01)

The present invention relates to a crystalline form of (S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydro-pyrazolo[1,5-a]pyrimidine-3-carboxamide for inhibiting Btk, methods of preparation thereof and pharmaceutical compositions, and use of the crystalline form above in the treatment of a disease, or in the manufacturing of a medicament for the treatment of a disease.

28 Claims, 8 Drawing Sheets



Related U.S. Application Data

continuation of application No. 17/146,855, filed on Jan. 12, 2021, which is a continuation of application No. 16/325,447, filed as application No. PCT/IB2017/054955 on Aug. 15, 2017, now Pat. No. 10,927,117.

(58) **Field of Classification Search**

CPC C07D 471/04; A61K 2300/00; A61K 31/519; A61K 2039/505; A61K 39/395; A61K 39/3955; A61K 39/39558; A61K 45/06; A61K 9/0019; Y02P 20/55; C07K 16/2818; C07K 16/2887

See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

7,718,662	B1	5/2010	Chen et al.
8,084,620	B2	12/2011	Liu et al.
8,735,553	B1	5/2014	Li et al.
9,447,106	B2	9/2016	Wang et al.
9,556,188	B2	1/2017	Wang et al.
10,005,782	B2	6/2018	Wang et al.
10,570,139	B2	2/2020	Wang et al.
10,709,699	B2	7/2020	Ding et al.
10,786,507	B2	9/2020	Lu et al.
10,927,117	B2	2/2021	Wang et al.
11,142,528	B2	10/2021	Wang et al.
11,512,132	B2	11/2022	Li et al.
11,555,038	B2	1/2023	Guo et al.
11,591,340	B2	2/2023	Wang et al.
11,597,768	B2	3/2023	Wang et al.
11,673,951	B2	6/2023	Li et al.
11,786,529	B2	10/2023	Hilger et al.
11,814,389	B2	11/2023	Wang et al.
2002/0094989	A1	7/2002	Hale et al.
2006/0178367	A1	8/2006	Currie et al.
2006/0183746	A1	8/2006	Currie et al.
2008/0076921	A1	3/2008	Honigberg et al.
2008/0139582	A1	6/2008	Honigberg et al.
2009/0105209	A1	4/2009	Dewdney et al.
2009/0318441	A1	12/2009	Brain et al.
2010/0004231	A1	1/2010	Dewdney et al.
2010/0016296	A1	1/2010	Singh et al.
2010/0016301	A1	1/2010	Dewdney et al.
2010/0029610	A1	2/2010	Singh et al.
2010/0035841	A1	2/2010	Jankowski et al.
2010/0087464	A1	4/2010	Mi et al.
2010/0105676	A1	4/2010	Liu et al.
2010/0144705	A1	6/2010	Miller
2010/0160292	A1	6/2010	Whitney et al.
2010/0160303	A1	6/2010	Liu et al.
2010/0222325	A1	9/2010	Berthel et al.
2010/0249092	A1	9/2010	Singh et al.
2010/0254905	A1	10/2010	Honigberg et al.
2011/0118233	A1	5/2011	Blomgren et al.
2011/0124640	A1	5/2011	Liu et al.
2011/0224235	A1	9/2011	Honigberg et al.
2011/0301145	A1	12/2011	Barbosa, Jr. et al.
2012/0028981	A1	2/2012	Miller
2012/0040961	A1	2/2012	Gray et al.
2012/0053189	A1	3/2012	Loury
2012/0058996	A1	3/2012	Liu et al.
2012/0077832	A1	3/2012	Witowski et al.
2012/0082702	A1	4/2012	Delucca et al.
2012/0129852	A1	5/2012	Duan et al.
2012/0157442	A1	6/2012	Bui et al.
2012/0157443	A1	6/2012	Bui et al.
2012/0232054	A1	9/2012	Moriarty et al.
2013/0079327	A1	3/2013	Yamamoto et al.
2013/0096118	A1	4/2013	Liu et al.
2013/0116213	A1	5/2013	Cha et al.
2013/0261103	A1	10/2013	Currie et al.
2013/0281432	A1	10/2013	Currie et al.

2014/0045833	A1	2/2014	Laurent et al.
2014/0094459	A1	4/2014	Goldstein et al.
2014/0107151	A1	4/2014	Goldstein et al.
2014/0162983	A1	6/2014	Hodous et al.
2014/0221398	A1	8/2014	Goldstein et al.
2014/0243306	A1	8/2014	Heng et al.
2015/0079109	A1	3/2015	Li et al.
2015/0118222	A1	4/2015	Levy et al.
2015/0259354	A1	9/2015	Wang et al.
2015/0315274	A1	11/2015	Li et al.
2016/0083392	A1	3/2016	Wang et al.
2017/0073349	A1	3/2017	Wang et al.
2018/0251466	A1	9/2018	Wang et al.
2019/0169201	A1	6/2019	Wang et al.
2020/0148690	A1	5/2020	Wang et al.
2020/0181150	A1	6/2020	Wang et al.
2020/0368237	A1	11/2020	Hilger et al.
2021/0130363	A1	5/2021	Wang et al.
2021/0275530	A1	9/2021	Hu et al.
2021/0332049	A1	10/2021	Guo et al.
2022/0241285	A1	8/2022	Wang et al.
2022/0249491	A1	8/2022	Qiu et al.
2022/0274994	A1	9/2022	Wang et al.
2022/0281876	A1	9/2022	Wang et al.
2022/0281881	A1	9/2022	Wang et al.
2022/0298163	A1	9/2022	Wang et al.
2022/0340583	A1	10/2022	Wang et al.
2022/0340584	A1	10/2022	Wang et al.
2022/0387404	A1	12/2022	Jiang et al.
2023/0011916	A1	1/2023	Jiang et al.
2023/0013862	A1	1/2023	Wang et al.
2023/0057716	A1	2/2023	Wang et al.
2023/0089557	A1	3/2023	Yu et al.
2023/0149411	A1	5/2023	Wang et al.
2023/0242542	A1	8/2023	Wang
2023/0303689	A1	9/2023	Li et al.
2023/0364096	A1	11/2023	Hu et al.

FOREIGN PATENT DOCUMENTS

CN	104884458	A	9/2015
CN	107530348	A	1/2018
CN	108778281	B	9/2021
GB	1412017	A	10/1975
JP	H07278148	A	10/1995
JP	2006510582	A	3/2006
JP	2010504324	A	2/2010
WO	WO-0116138	A1	3/2001
WO	WO-0119829	A2	3/2001
WO	WO-2002020740		3/2002
WO	WO-0250071	A1	6/2002
WO	WO-02072576	A1	9/2002
WO	WO-03004497	A1	1/2003
WO	WO-2004017908	A2	3/2004
WO	WO-2004099249	A2	11/2004
WO	WO-2005005429	A1	1/2005
WO	WO-2005011597	A2	2/2005
WO	WO-2005014599	A1	2/2005
WO	WO-2005047290	A2	5/2005
WO	WO-2006053121	A2	5/2006
WO	WO-2006065946	A1	6/2006
WO	WO-2006099075	A2	9/2006
WO	WO-2007026720	A1	3/2007
WO	WO-2007026950	A1	3/2007
WO	WO-2007027594	A1	3/2007
WO	WO-2007027729	A1	3/2007
WO	WO-2007087068	A2	8/2007
WO	WO-2007136790	A2	11/2007
WO	WO-2008033834	A1	3/2008
WO	WO-2008033854	A1	3/2008
WO	WO-2008033857	A2	3/2008
WO	WO-2008039218	A2	4/2008
WO	WO-2008054827	A2	5/2008
WO	WO-2008144253	A1	11/2008
WO	WO-2009039397	A2	3/2009
WO	WO-2009051822	A1	4/2009
WO	WO-2009077334	A1	6/2009
WO	WO-2009098144	A1	8/2009
WO	WO-2009158571	A1	12/2009

(56)

References Cited

FOREIGN PATENT DOCUMENTS

WO	WO-2010000633	A1	1/2010
WO	WO-2010006947	A1	1/2010
WO	WO-2010006970	A1	1/2010
WO	WO-2010028236	A1	3/2010
WO	WO-2010051549	A1	5/2010
WO	WO-2010065898	A2	6/2010
WO	WO-2010068788	A1	6/2010
WO	WO-2010068806	A1	6/2010
WO	WO-2010068810	A2	6/2010
WO	WO-2010122038	A1	10/2010
WO	WO-2011006074	A1	1/2011
WO	WO-2011140488	A1	11/2011
WO	WO-2011153514	A2	12/2011
WO	WO-2012020008	A1	2/2012
WO	WO-2012135801	A1	10/2012
WO	WO-2012143522	A1	10/2012
WO	WO-2012156334	A1	11/2012
WO	WO-2012158795	A1	11/2012
WO	WO-2014173289	A1	10/2014
WO	WO-2015035606	A1	3/2015
WO	WO-2015061752	A1	4/2015
WO	WO-2016008411	A1	1/2016
WO	WO-2016024228	A1	2/2016
WO	WO-2016025720	A1	2/2016
WO	WO-2016087994	A1	6/2016
WO	WO-2016100914	A1	6/2016
WO	WO-2016105582	A1	6/2016
WO	WO-2017046746	A1	3/2017
WO	WO-2017059224	A2	4/2017
WO	WO-2017218844	A2	12/2017
WO	WO-2018033135	A1	2/2018
WO	WO-2018033853	A2	2/2018
WO	WO-2018137681	A1	8/2018
WO	WO-2018193105	A1	10/2018
WO	WO-2019034009	A1	2/2019
WO	WO-2019108795	A1	6/2019
WO	WO-2019183226	A1	9/2019
WO	WO-2020249001	A1	12/2020
WO	WO-2020249002	A1	12/2020
WO	WO-2021047623	A1	3/2021
WO	WO-2021170045	A1	9/2021

OTHER PUBLICATIONS

Atkinson, B. T., "Tec regulates platelet activation by GPVI in the absence of Btk," *Blood*, Nov. 15, 2003, vol. 102, Issue 19, pp. 3592-3599.

Balar, A. et al., "Pembrolizumab (pembro) as first-line therapy for advanced/unresectable or metastatic urothelial cancer: Preliminary results from the phase 2 KEYNOTE-052 study," *Annals of Oncology*, 2016, vol. 27 (Supplement 6), pp. vi552-vi587.

Balbach, S., et al., "Pharmaceutical evaluation of early development candidates: 'The 100 mg-approach'," *International Journal of Pharmaceutics*, 2004, vol. 275 pp. 1-12.

Beigene Co., Ltd., "Phase I Clinical Research to Evaluate Safety, Tolerability and Pharmacokinetics/Pharmacodynamics Characteristics of BTK Inhibitor, BGB-3111, in Treating Chinese B Lymphocyte Tumor Patients," [Online], May 2016, Retrieved from the Internet: <http://www.chinadrugtrials.org.cn/clinicaltrials.searchlistdetail.dhtml>, Retrieved on: Mar. 21, 2022, 17 pages (with Machine Translation).

Beigene, "Efficacy and Safety of Zanubrutinib in Relapsed or Refractory Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma," [Online], Jul. 2017, Retrieved from the Internet: <https://clinicaltrials.gov/ct2/show/NCT03206918>, Retrieved on: Apr. 4, 2022, 7 pages.

Bellmunt, J. et al., "Pembrolizumab as Second-Line Therapy for Advanced Urothelial Carcinoma," *N. Engl. J. Med.*, Mar. 2017, vol. 376, No. 11, pp. 1015-1026.

Bellmunt, J. et al., "Randomized Phase III Study Comparing Paclitaxel/Cisplatin/Gemcitabine and Gemcitabine/Cisplatin in Patients

with Locally Advanced or Metastatic Urothelial Cancer Without Prior Systemic Therapy: EORTC Intergroup Study 30987," *J Clin Oncol.*, Apr. 1, 2012, vol. 30, No. 10, pp. 1107-1113.

Benson, D. M., Jr. et al., "The PD-1/PD-L1 axis modulates the natural killer cell versus multiple myeloma effect: a therapeutic target for CT-011, a novel monoclonal anti-PD-1 antibody," *Blood*, Sep. 2010, vol. 116, No. 13, pp. 2286-2294.

Berger, R. et al., "Phase I safety and pharmacokinetic study of CT-011, a humanized antibody interacting with PD-1, in patients with advanced hematologic malignancies," *Clinical Cancer Research*, May 2008, vol. 14, No. 10, pp. 3044-3051.

Bowman, B.J. et al., "Hard gelatin capsules. Drug Release from HGC," Chapter 15 in *Protein-Based Films and Coatings*. Aristippos Gennadios (Ed.) CRC Press, 2002; p. 385.

Boyd et al., "Deep sequencing and human antibody repertoire analysis," *Current Opinion in Immunology*, Jun. 1, 2016, vol. 40, pp. 103-109.

Bradshaw, J. M., "The Src, Syk, and Tec family kinases: distinct types of molecular switches," *Cell Signal.*, Aug. 2010, vol. 22, No. 8, pp. 1175-1184.

Brinckerhoff, C., "A Fresh Look at the Lead Compound Analysis," *PharmaPatents*, Foley & Lardner LLP, Jan. 22, 2019, 4 pages.

Buske, C. et al., "A Head-to-Head Phase 3 Study Comparing BGB-3111 and Ibrutinib in Patients With Waldenstrom Macroglobulinemia," *Hematological Oncology*, Jun. 7, 2017, vol. 35, pp. 422-423.

Byrd, J. C. et al., "Acalabrutinib (ACP-196) in Relapsed Chronic Lymphocytic Leukemia," *N. Engl. J. Med.*, Jan. 28, 2016, vol. 374, No. 4, pp. 323-332.

Byrd, J. C. et al., "Targeting BTK with Ibrutinib in Relapsed Chronic Lymphocytic Leukemia," *N. Engl. J. Med.*, Jul. 4, 2013, vol. 369, No. 1, pp. 32-42.

Byrd, J. C. et al., "Three-year follow-up of treatment-naïve and previously treated patients with CLL and SLL receiving single-agent ibrutinib," *Blood*, Apr. 15, 2015, vol. 125, Issue 16, pp. 2497-2506.

Caira, M. R., "Crystalline Polymorphism of Organic Compounds", *Topics in Current Chemistry*, Jan. 1, 1998, vol. 198, pp. 163-208.

Cartigny, D. et al., "General Asymmetric Hydrogenation of 2-Alkyl- and 2-Aryl-Substituted Quinoxaline Derivatives Catalyzed by Iridium-Difluorophos: Unusual Halide Effect and Synthetic Application," *J. Org. Chem.*, Apr. 2012, vol. 77, No. 10, pp. 4544-4556.

Casset, F., et al., "A Peptide Mimetic of an Anti-cd4 Monoclonal Antibody by Rational Design," *Biochemical and Biophysical Research Communications*, Jul. 2003, vol. 307, No. 1, pp. 198-205.

Cermakova, K et al., "Next-Generation Drugs and Probes for Chromatin Biology: From Targeted Protein Degradation to Phase Separation," *Molecules*, Aug. 2018, vol. 23, No. 1958, 26 pages.

Chan., et al., "Excipients: powders and solid dosage forms," *Encyclopedia of pharmaceutical technology*, 3rd ed., Informa Healthcare USA, Inc, New York 3, 2007, pp. 1646-1655.

Chen, J. et al., "Impact of Bruton's Tyrosine Kinase Inhibitors on Collagen-Induced Platelet Aggregation: a Pharmacokinetic/Pharmacodynamic Perspective," *EHA Learning Center, Meeting Abstract 2016*, 4 pages.

ClinicalTrials.gov Identifier: NCT03053440, A Study Comparing BGB-3111 and Ibrutinib in Subjects With Waldenström's Macroglobulinemia (WM) (Aug. 8, 2017), 4 pages.

ClinicalTrials.gov Identifier: NCT03145064, Study of BTK Inhibitor BGB-3111 in Subjects With Relapsed/Refractory Non-GCB Type Diffuse Large B Cell Lymphoma (May 5, 2017), 5 pages.

ClinicalTrials.gov Identifier: NCT03189524, A Study to Investigate Zanubrutinib in Chinese Participants With B-cell Lymphoma (Jun. 15, 2017), 5 pages.

ClinicalTrials.gov Identifier: NCT03206970, Study to Evaluate Efficacy and Safety of BGB-3111 in Participants With Relapsed or Refractory Mantle Cell Lymphoma (MCL) (Jul. 3, 2017), 6 pages.

ClinicalTrials.gov Identifier: NCT04470908, The Effect of Moderate CYP3A Inducer Rifabutin on the Pharmacokinetics of Zanubrutinib in Healthy Males, Sep. 21, 2021, 6 pages.

Conley, M. E. et al., "Primary B Cell Immunodeficiencies: Comparisons and Contrasts," *Annu. Rev. Immunol.*, Apr. 23, 2009, vol. 27, pp. 199-227.

(56)

References Cited

OTHER PUBLICATIONS

- Davis, R. E. et al., "Chronic active B-cell-receptor signaling in diffuse large B-cell lymphoma," *Nature*, Jan. 7, 2019, vol. 463, No. 7277, pp. 88-92.
- Di Paolo et al., "Specific Btk inhibition suppresses B cell- and myeloid cell-mediated arthritis," *Nat. Chem. Biol.*, Jan. 2011, vol. 7, No. 1, pp. 41-50.
- Eng, M., "United States: Selecting a Lead Compound: A Balancing Act in the Chemical Obviousness Inquiry," Monday, Mar. 8, 2018, 2 pages, retrieved from the Internet on Nov. 28, 2022, <https://www.mondaq.com/unitedstates/patent/680778/selecting-a-lead-compound-a-balancing-act-in-the-chemical-obviousness-inquiry>.
- Extended European Search Report for European Application No. 14787642.9, dated Jan. 26, 2016, 5 pages.
- Extended European Search Report for European Application No. 17841107.0, dated Feb. 21, 2020, 12 pages.
- Extended European Search Report for European Application No. 17841172.4, dated Mar. 5, 2020, 6 pages.
- Extended European Search Report for European Application No. 18744173.8, dated Oct. 21, 2020, 12 pages.
- Extended European Search Report for European Application No. 20822606.8, dated Jun. 9, 2023, 9 pages.
- Gurcan, H. M. et al., "A review of the current use of rituximab in autoimmune diseases," *Int. Immunopharmacol.*, Jan. 2009, vol. 9, No. 1, pp. 10-25.
- Hackam et al., "Translation of research evidence from animals to humans," *JAMA*, Oct. 11, 2006, vol. 296, No. 14, pp. 1727-1732.
- He, X., "Part III. Design, Development, and Scale-Up of Formulation and Process. 18. Integration of Physical, Chemical, Mechanical, and Biopharmaceutical Properties in Solid Oral Dosage Form Development," In *Developing Solid Oral Dosage Forms Pharmaceutical Theory and Practice*, Elsevier Science, 2009, pp. 414-415.
- Heda et al., "8 Dry-Fill Formulation," *Pharmaceutical Dosage Forms: Capsules*, Oct. 30, 2017, pp. 211-236.
- Hirayama, Y., "Handbook for organic compound crystal—Principle and know-how," 2008, 28 pages.
- Hu, Nan et al., "BTK inhibitor BGB-3111 demonstrates anti-tumor activity in solid tumor models," *Cancer Research*, Jul. 1, 2017, vol. 77, Supplement 13, 4 pages.
- Humphries, L. A. et al., "Tec kinases mediate sustained calcium influx via site-specific tyrosine phosphorylation of the phospholipase Cgamma Src homology 2-Src homology 3 linker," *J. Biol.Chem.*, Sep. 2004, vol. 279, No. 36, pp. 37651-37661.
- "International Nonproprietary Names for Pharmaceutical Substances (INN)," *WHO Drug Information*, 2017, vol. 31, No. 2, pp. 241-242, 355.
- International Search Report and Written Opinion for International Application No. PCT/CN2014/075943, dated Jul. 18, 2014, 10 pages.
- International Search Report and Written Opinion for International Application No. PCT/CN2017/098023, dated Nov. 16, 2017, 10 pages.
- International Search Report and Written Opinion for International Application No. PCT/CN2018/074108, dated Apr. 23, 2018, 12 pages.
- International Search Report and Written Opinion for International Application No. PCT/CN2018/100145, dated Nov. 14, 2018, 8 pages.
- International Search Report and Written Opinion for International Application No. PCT/CN2020/095352, dated Sep. 16, 2020, 23 pages.
- International Search Report and Written Opinion for International Application No. PCT/CN2020/095353, dated Sep. 15, 2020, 23 pages.
- International Search Report and Written Opinion for International Application No. PCT/IB2017/054955, dated Sep. 10, 2018, 16 pages.
- International Search Report and Written Opinion for International Application No. PCT/US2018/063068, dated Feb. 27, 2019, 7 pages.
- Jenkins, S. M. et al., "Substituent variation in azabicyclic triazole- and tetrazole-based muscarinic receptor ligands," *J. Med. Chem.*, Jun. 1992, vol. 35, No. 13, pp. 2392-2406.
- Jiangsu Chia Tai Qingjiang Pharmaceutical Co, Ltd., Clinical Trial Reg. No. CTR20160513, Oct. 10, 2016, retrieved online at <http://www.chinadrugtrials.org.cn/clinicaltrials.searchlistdetail.dhtml> on May 22, 2023, 5 pages. (ENG Translation).
- Jie, L., "Deuterated Drugs Progress," *Chemical Engineering Design Communication Medicine and Chemical Industry*, 2016, vol. 42, No. 4, p. 199.
- Jordan "Tamoxifen: a most unlikely pioneering medicine," *Nature Reviews Drug Discovery*, Mar. 2003, vol. 2, No. 3, pp. 205-213.
- Kaur, V. et al., "Ibrutinib in CLL: a focus on adverse events, resistance, and novel approaches beyond ibrutinib," *Ann. Hematol.*, Jul. 2017, vol. 96, No. 7, pp. 1175-1184.
- Kersseboom, R. et al., "Constitutive activation of Bruton's tyrosine kinase induces the formation of autoreactive IgM plasma cells," *Eur. J. Immunol.*, Sep. 2010, vol. 40, pp. 2643-2654.
- Khan, W. N., "Regulation of B lymphocyte development and activation by Bruton's tyrosine kinase," *Immunol. Res.*, Apr. 2001, vol. 23, pp. 147-156.
- Kim, K.-H. et al., "Imidazo[1,5-a]quinoxalines as irreversible BTK inhibitors for the treatment of rheumatoid arthritis," *Bioorg. Med. Chem. Lett.*, Nov. 1, 2011, vol. 21, No. 21, pp. 6258-6263.
- Kourtzelis, I., et al., "The Dual Role of Complement in Cancer and its Implication in Anti-Tumor Therapy," *Annals of Translational Medicine*, Jul. 2016, vol. 4, No. 14, 14 pages.
- Li et al., "Skin toxicities associated with epidermal growth factor receptor inhibitors," *Targeted Oncol.*, Jun. 2009, vol. 4, pp. 107-119.
- Li, N. et al., "BGB-3111 is a novel and highly selective Bruton's tyrosine kinase (BTK) inhibitor," *Cancer Center, 106th Annual Meeting of the American Association for Cancer Research, AACR 2015, Philadelphia, PA, United States, Aug. 2015, vol. 75, No. 15, Supp. 1, Abstract No. 2597, 2 pages.*
- Lou, Y. et al., "Bruton's tyrosine kinase inhibitors: Approaches to potent and selective inhibition, preclinical and clinical evaluation for inflammatory diseases and B cell malignancies," *J. Med. Chem.*, May 24, 2023, vol. 55, No. 10, pp. 4539-4550.
- Luo, J. et al., "Modern Physical Pharmaceutics Theory and Practice," *Shang Hai Science and Technology Literature Publishing House*, Apr. 2005, pp. 293-295.
- Medchemexpress, "Zanubrutinib," Product Data Sheet, Retrieved from the Internet: www.medchemexpress.com, Retrieved Aug. 17, 2021, 2 pages.
- Melosky et al., "Supportive Care Treatments for Toxicities of Anti-EGFR and Other Targeted Agents," *Curr. Oncol.*, Jun. 2012, vol. 19, Suppl. 1, pp. 59-63.
- Miller, C., "Prior Art Chemical Structures Must be More Than a 'Code Name'," *Jones Day*, Apr. 18, 2018, retrieved online Nov. 28, 2022, 2 pages, retrieved online at <https://www.ptablitigationblog.com/prior-art-chemical-structures-must-be-more-than-a-code-name/>.
- Mohamed, A. J. et al., "Bruton's tyrosine kinase (Btk): function, regulation, and transformation with special emphasis on the PH domain," *Immunol. Rev.*, Mar. 2009, vol. 228, pp. 58-73.
- Neklesa, T.K. et al., "Targeted protein degradation by PROTACs," *Pharmacology & Therapeutics*, Jun. 1, 2017, vol. 174, pp. 138-144.
- Noonan, K., "*Novartis Pharmaceuticals Corp. v. West-Ward Pharmaceuticals Int'l* (Fed. Cir. 2019)," *Patent Docs*, May 16, 2019, 3 pages, retrieved from the internet on Nov. 28, 2022, <https://www.patentdocs.org/2019/05/novartis-pharmaceuticals-corp-v-west-ward-pharmaceuticals-intl-fed-cir-2019.html>.
- Office Action for Japanese Application No. 2019-508889, dated Jun. 22, 2021, 7 pages.
- Ou, Y. et al., "Evaluation of drug interaction potential of zanubrutinib with cocktail probes representative of CYP34A, CYP2C9, CYP2C19, P-gp and BCRP," *British Journal of Clinical Pharmacology*, 2021, vol. 87, pp. 2926-2936.
- Pacoud, O. et al., "A Woman with Relapsed Chronic Lymphocytic Leukemia and Upper Lobe Consolidation," *American Thoracic Society*, Nov. 2021, vol. 18, No. 11, pp. 1901-1906.
- Pan, Z. "Bruton's tyrosine kinase as a drug discovery target," *Drug News Perspect*, Sep. 1, 2008, vol. 21, No. 7, pp. 357-362.

(56)

References Cited

OTHER PUBLICATIONS

- Pan, Z. et al., "Discovery of Selective Irreversible Inhibitors for Bruton's Tyrosine Kinase," *ChemMedChem*, Jan. 2007, vol. 2, No. 1, pp. 58-61.
- Park, B.K. et al., "The Role of Cytochrome P450 Enzymes in Hepatic and Extrahepatic Human Drug Toxicity," *Pharmac. Ther.*, Jan. 1, 1995, vol. 68, No. 3, pp. 385-424.
- Pirmohamed, M. et al., "The Role of Active Metabolites in Drug Toxicity," *Drug Safety*, Aug. 1994, vol. 11, No. 2, pp. 114-144.
- Raedler, L. A., "Farydak (Panobinostat): First HDAC Inhibitor Approved for Patients with Relapsed Multiple Myeloma," *Am. Health Drug Benefits*, Mar. 2016, vol. 9, pp. 84-87.
- REVLIMID Prescribing Information, Initial US Approval 2005 (Celgene Corp), Revised Sep. 2014 (aka Revlimid 2014), 35 pages.
- Rokosz, L. L. et al., "Kinase inhibitors as drugs for chronic inflammatory and immunological diseases: progress and challenges," *Expert Opin. Ther. Targets*, Jul. 1, 2008, vol. 12, No. 7, pp. 883-903.
- Seymour, J. F., et al., "High Overall Response Rate With the BTK Inhibitor BGB-3111 in Patients With Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma: An Update on Safety and Activity," *Hematological Oncology*, 2017, vol. 35, pp. 234-235.
- Shioji, Y., "Production Technology of Solid Preparations," Tokyo, CMC Publication, Jan. 27, 2003, Popular Edition, pp. 9 and 12-13.
- Singhal, D., et al., "Drug Polymorphism and Dosage Form Design: A practical perspective," *Advanced Drug Delivery Reviews*, 2004, vol. 56, pp. 335-347.
- Smith, C. I. et al., "Expression of Bruton's agammaglobulinemia tyrosine kinase gene, BTK, is selectively down-regulated in T lymphocytes and plasma cells," *J. Immunol.*, Jan. 1994, vol. 152, No. 2, pp. 557-565.
- Takada, N., "Bulk Drug Form Screening and Selection at Drug Discovery Phase," *Pharm Stage*, Jan. 2007, vol. 6, No. 10, pp. 20-25.
- Takayama, T. et al., "Effects of the novel and potent lymphocyte-specific protein tyrosine kinase inhibitor TKM0150 on mixed lymphocyte reaction and contact hypersensitivity in mice," *Arzneimittelforschung*, 2010, vol. 60, No. 5, pp. 282-285.
- Takayama, T. et al., "Ring-fused pyrazole derivatives as potent inhibitors of lymphocyte-specific kinase (Lck): Structure, synthesis, and SAR," *Bioorganic & Medicinal Chemistry Letters*, Jan. 2010, vol. 20, No. 1, pp. 112-116.
- Tam, C. et al., "The BTK Inhibitor, Bgb-3111, is Safe, Tolerable, and Highly Active in Patients with Relapsed/ Refractory B-Cell Malignancies: Initial Report of a Phase 1 First-in-Human Trial," *Blood*, Dec. 3, 2015, vol. 126, No. 23, 8 pages.
- Tam, C. S. et al., "A head-to-head Phase III study comparing zanubrutinib versus ibrutinib in patients with Waldenstrom macroglobulinemia," *Future Oncology*, Sep. 2018, vol. 14, No. 22, pp. 2229-2237.
- Tam, C. S. et al., "Phase 1 study of the selective BTK inhibitor zanubrutinib in B-cell malignancies and safety and efficacy evaluation in CLL," *The American Society of Hematology*, Sep. 12, 2019, vol. 134, No. 11, pp. 851-859.
- Tam, C.S. et al., "Clinical pharmacology and PK/PD translation of the second-generation Bruton's tyrosine kinase inhibitor, zanubrutinib," *Expert Review of Clinical Pharmacology*, Nov. 2, 2021, vol. 14, No. 11, pp. 1329-1344.
- Tam, C.S. et al., "High Major Response Rate, Including Very Good Partial Responses (VGPR), in Patients (pts) with Waldenstrom Macroglobulinemia (WM) Treated with the Highly Specific BTK Inhibitor Bgb-3111: Expansion Phase Results from an Ongoing Phase I Study," *Blood*, Dec. 2, 2016, vol. 128, No. 22, 8 pages.
- Tam, C.S. et al., "Twice Daily Dosing with the Highly Specific BTK Inhibitor, Bgb-3111, Achieves Complete and Continuous BTK Occupancy in Lymph Nodes, and is Associated with Durable Responses in Patients (pts) with Chronic Lymphocytic Leukemia (CLL)/Small Lymphocytic Lymphoma (SLL)," *Blood*, Dec. 2, 2016, vol. 128, No. 22, 10 pages.
- Tuloup, V. et al., "Model-Based Comparative Analysis of Rifampicin and Rifabutin Drug-Drug Interaction Profile," *Antimicrobial Agents and Chemotherapy*, Sep. 2021, vol. 65, No. 9, e01043-21, 7 pages.
- Uckun, F. M. et al., "Bruton's tyrosine kinase as a new therapeutic target," *Anti-Cancer Agents in Medicinal Chemistry*, Nov. 1, 2007, vol. 7, No. 6, pp. 624-632.
- Vetrie, D. et al., "The gene involved in X-linked agammaglobulinaemia is a member of the src family of protein-tyrosine kinases," *Nature*, Jan. 21, 1993, vol. 361, No. 6409, pp. 226-233.
- Wang, M. L. et al., "Targeting BTK with Ibrutinib in Relapsed or Refractory Mantle-Cell Lymphoma," *N. Engl. J. Med.*, Aug. 8, 2013, vol. 369, pp. 507-516.
- Wenzel, S-S et al., "MCL1 is deregulated in subgroups of diffuse large B-cell lymphoma," *Leukemia*, Jun. 2013, vol. 27, No. 6, pp. 1381-1390.
- Wilson, W. H. et al., "686—The Bruton's Tyrosine Kinase (Btk) Inhibitor, Ibrutinib (PCI-32765), Has Preferential Activity in the ABC Subtype of Relapsed/Refractory De Novo Diffuse Large B-Cell Lymphoma (DLBCL): Interim Results of a Multicenter, Open-Label, Phase 2 Study," *Poster #686, 54th American Society of Hematology (ASH) Annual Meeting Abstract*, Dec. 10, 2012, 3 pages.
- Woof, J.M., "Fc receptors and their interaction with antibodies," *Biochemical Society Transactions*, 1990, vol. 18, pp. 217-218.
- Wu, J. et al., "Second-generation inhibitors of Bruton tyrosine kinase," *Journal of Hematology & Oncology*, Sep. 2, 2016, vol. 9, No. 1, pp. 80.




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7A




FIG. 7B




FIG. 8




FIG. 9




FIG. 10




FIG. 11

1

**CRYSTALLINE FORM OF
(S)-7-(1-ACRYLOYLPYPERIDIN-4-YL)-
2-(4-PHENOXYPHENYL)-4,5,6,7-TETRA-
HYDROPIRAZOLO[1,5-A]PYRIMIDINE-
3-CARBOXAMIDE, PREPARATION, AND
USES THEREOF**

**CROSS-REFERENCE TO RELATED
APPLICATIONS**

This application is a continuation of U.S. application Ser. No. 17/146,855, filed on Jan. 12, 2021, which is a continuation of U.S. application Ser. No. 16/325,447, now U.S. Pat. No. 10,927,117, filed on Feb. 14, 2019, which is a U.S. National Stage Application under 35 U.S.C. § 371 of International Application No. PCT/IB2017/054955, filed on Aug. 15, 2017, which claims priority to Application No. PCT/CN2016/095510 (CN), filed Aug. 16, 2016.

FIELD OF THE INVENTION

The present invention relates to a crystalline form of (S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetra-hydropyrazolo[1,5-a]pyrimidine-3-carboxamide. The present invention also relates to methods of preparing the crystalline form and methods of using the crystalline form as a Btk inhibitor.

BACKGROUND OF THE INVENTION

Bruton's tyrosine kinase (Btk) belongs to the Tec tyrosine kinase family (Vetrie et al., *Nature* 361: 226-233, 1993; Bradshaw, *Cell Signal.* 22: 1175-84, 2010). Btk is primarily expressed in most hematopoietic cells such as B cells, mast cells and macrophages (Smith et al., *J. Immunol.* 152: 557-565, 1994) and is localized in bone marrow, spleen and lymph node tissue. Btk plays important roles in B-cell receptor (BCR) and FcR signaling pathways, which involve in B-cell development, differentiation (Khan, *Immunol. Res.* 23: 147, 2001). Btk is activated by upstream Src-family kinases. Once activated, Btk in turn phosphorylates PLC gamma, leading to effects on B-cell function and survival (Humphries et al., *J. Biol. Chem.* 279: 37651, 2004).

These signaling pathways must be precisely regulated. Mutations in the gene encoding Btk cause an inherited B-cell specific immunodeficiency disease in humans, known as X-linked agammaglobulinemia (XLA) (Conley et al., *Annu. Rev. Immunol.* 27: 199-227, 2009). Aberrant BCR-mediated signaling may result in dysregulated B-cell activation leading to a number of autoimmune and inflammatory diseases. Preclinical studies show that Btk deficient mice are resistant to developing collagen-induced arthritis. Moreover, clinical studies of Rituxan, a CD20 antibody to deplete mature B-cells, reveal the key role of B-cells in a number of inflammatory diseases such as rheumatoid arthritis, systemic lupus erythematosus and multiple sclerosis (Gurcan et al., *Int. Immunopharmacol.* 9: 10-25, 2009). Therefore, Btk inhibitors can be used to treat autoimmune and/or inflammatory diseases.


In addition, aberrant activation of Btk plays an important role in pathogenesis of B-cell lymphomas indicating that inhibition of Btk is useful in the treatment of hematological malignancies (Davis et al., *Nature* 463: 88-92, 2010). Preliminary clinical trial results showed that the Btk inhibitor PCI-32765 was effective in treatment of several types of B-cell lymphoma (for example, 54th American Society of Hematology (ASH) annual meeting abstract, December

2

2012: 686 The Bruton's Tyrosine Kinase (Btk) Inhibitor, Ibrutinib (PCI-32765), Has Preferential Activity in the ABC Subtype of Relapsed/Refractory De Novo Diffuse Large B-Cell Lymphoma (DLBCL): Interim Results of a Multi-center, Open-Label, Phase I Study). Because Btk plays a central role as a mediator in multiple signal transduction pathways, inhibitors of Btk are of great interest as anti-inflammatory and/or anti-cancer agents (Mohamed et al., *Immunol. Rev.* 228: 58-73, 2009; Pan, *Drug News perspect* 21: 357-362, 2008; Rokosz et al., *Expert Opin. Ther. Targets* 12: 883-903, 2008; Uckun et al., *Anti-cancer Agents Med. Chem.* 7: 624-632, 2007; Lou et al, *J. Med. Chem.* 55(10): 4539-4550, 2012).

International application WO2014173289A disclosed a series of fused heterocyclic compounds as Btk inhibitors. In particular, WO2014173289A disclosed (S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetra-hydropyrazolo[1,5-a]pyrimidine-3-carboxamide (hereinafter Compound 1)

Compound 1



Compound 1 is a potent, specific and irreversible BTK kinase inhibitor. The data generated in preclinical studies using biochemical, cell based and animal studies suggested that Compound 1 could offer significant benefit in inhibiting tumor growth in B-cell malignancies. As Compound 1 was shown to be more selective than ibrutinib for inhibition of BTK vs. EGFR, FGR, FRK, HER2, HER4, ITK, JAK3, LCK, and TEC, it is expected to give rise to less side-effects than ibrutinib in clinic. In addition, Compound 1 showed significantly less inhibition of rituximab-induced antigen-dependent cell-mediated cytotoxicity (ADCC) than ibrutinib due to weaker ITK inhibition, and therefore may provide better efficacy when combined with rituximab or other ADCC-dependent antibody in treating B-cell malignancies.

Preclinical safety evaluation has demonstrated that Compound 1 was safer than ibrutinib in terms of the overall tolerance and severe toxicities in both rat and dog single and repeat dose toxicity studies up to 28 days. Additionally, Compound 1 had better bioavailability without accumulation issues observed for ibrutinib. These unique characteristics warrant further evaluation of Compound 1 in clinical studies.

However, Compound 1 was found to be an amorphous form according to the preparation method for Compound 27 in WO 2014173289A, which was further confirmed by the X-Ray Powder Diffraction pattern of FIG. 7A. The amorphous form was shown to have a low glass transition temperature as shown in FIG. 7B, indicating some difficulties in the drug formulation with the amorphous form, such


3

as low stability and hard to purify. Therefore, it's necessary to develop a new form of Compound 1 which possesses characteristics such as high melting point and better stability, suitable for drug formulation.

SUMMARY OF THE INVENTION

The inventors have unexpectedly found a crystalline form of Compound 1, which possesses a high melting point and shows an extremely stable profile even when stored at 25° C./60% RH for up to 24 months or stored at 40° C./75% RH condition for up to 6 months.


In a first aspect, disclosed herein is a crystalline form of Compound 1,



Compound 1

In some embodiments, the crystalline form of Compound 1 is a crystalline anhydrate (herein referred to as "Crystalline Form A").

In a second aspect, disclosed herein is a crystalline form of Compound BG-13, which has an X-ray powder diffraction pattern substantially in accordance with FIG. 11.




Compound BG-13

In a third aspect, disclosed herein is a method of preparing Compound 1.


Also disclosed herein is an intermediate compound of Formula Ie or a salt thereof, or Formula If or a salt thereof used to prepare Compound 1,

4

Formula Ie



Formula If



In a fourth aspect, disclosed herein is a method of preparing Crystalline Form A disclosed herein.

In a fifth aspect, disclosed herein is a pharmaceutical composition comprising a therapeutically effective amount of Crystalline Form A disclosed herein.

In a sixth aspect, disclosed herein is a method of treating a disease associated with undesirable Btk activity in a subject by administering to a subject Crystalline Form A disclosed herein.

In a seventh aspect, disclosed herein is a method of treating a disease selected from an allergic disease, an autoimmune disease, an inflammatory disease, a cancer, or a combination of two or more thereof, in a subject by administering to the subject Crystalline Form A disclosed herein.

In an eighth aspect, disclosed herein is a method of treating a B-cell proliferative disease, selected from B-cell malignancies, or relapsed/refractory B-cell malignancies, in a subject by administering to the subject Crystalline Form A disclosed herein. In some embodiment of this aspect, disclosed herein is a method of treating a B-cell proliferative disease, selected from chronic lymphocytic, non-Hodgkin's lymphoma, diffuse large B cell lymphoma, mantle cell lymphoma, follicular lymphoma, chronic lymphocytic leukemia, small lymphocytic lymphoma, waldenstrom macroglobulinemia, marginal zone lymphoma, Hairy cell leukemia, Burkitt's-like leukemia or a combination of two or more thereof, in a subject by administering to the subject Crystalline Form A disclosed herein.

In a ninth aspect, disclosed herein is a use of Crystalline Form A disclosed herein in manufacturing a medicament for treatment of at least one disease associated with undesirable Btk activity, in a subject.

5

In a tenth aspect, disclosed herein is a use of Crystalline Form A disclosed herein in manufacturing a medicament for treatment of a disease selected from an allergic disease, an autoimmune disease, an inflammatory disease, a cancer, or a combination of two or more thereof, in a subject.

In an eleventh aspect, disclosed herein is a use of Crystalline Form A disclosed herein in manufacturing a medicament for treatment of a B-cell proliferative disease selected from B-cell malignancies, or relapsed/refractory B-cell malignancies, in a subject. In some embodiment of this aspect, disclosed herein is a use of Crystalline Form A disclosed herein in manufacturing a medicament for treatment of a B-cell proliferative disease selected from chronic lymphocytic, non-Hodgkin's lymphoma, diffuse large B cell lymphoma, mantle cell lymphoma, follicular lymphoma, chronic lymphocytic leukemia, small lymphocytic lymphoma, waldenstrom macroglobulinemia, marginal zone lymphoma, Hairy cell leukemia, Burkitt's-like leukemia, or a combination of two or more thereof, in a subject.

In a twelfth aspect, disclosed herein is a process for preparing a crystalline form A of Compound 1, comprising mixing amorphous form of compound 1 with the following solvent system to form a clear solution; keeping the solution at room temperature or heat with or without stirring for a certain period of time to precipitate the crystalline form A, wherein the solvent system is:

ethyl acetate: hexane=1:0.6-0.7 by volume ratio;
ethyl acetate: heptane=1:0.6-0.7 by volume ratio;
ethyl acetate: cyclohexane=1:0.6-1.2 by volume ratio;
methyl acetate: hexane=1:0.6-1.2 by volume ratio;
toluene: hexane=1.0:0.2-0.4 by volume ratio;
toluene: cyclohexane=1.0:0.1-0.2 by volume ratio;
methyl acetate: cyclohexane=0.6-0.8:1.0 by volume ratio;
IPAC: cyclohexane=1.0:0.2-1.0 by volume ratio; or
Isobutyl acetate: cyclohexane=1.0:0.2-1.0 by volume ratio.

In one embodiment, the amorphous form of compound 1 has an ee value more than 90%. In other embodiment, the amorphous form of compound 1 has an ee value of 97%.

BRIEF DESCRIPTIONS OF THE DRAWINGS

FIG. 1 shows the XRPD pattern of Crystalline Form A.
FIG. 2 shows the DSC curve of Crystalline Form A.
FIG. 3 shows the TGA curve of Crystalline Form A.
FIG. 4 shows the ¹H-NMR of Crystalline Form A.
FIG. 5 shows the ¹³C-NMR of Crystalline Form A.
FIG. 6 shows DVS plot of Crystalline Form A.
FIG. 7A shows the XRPD pattern of the amorphous form of Compound 1.
FIG. 7B shows the mDSC curve of the amorphous form of Compound 1, showing the glass transition temperature of the amorphous form is 79.7° C. (mid-point temperature).
FIG. 8 shows the absolute structure of single crystal of BG-13.

6

FIG. 9 illustrates hydrogen bonds of single crystal of BG-13.

FIG. 10 shows a crystal packing of single crystal of BG-13.

FIG. 11 shows the XRPD pattern of single crystal of BG-13.

DETAILED DESCRIPTION OF THE INVENTION

The inventors have unexpectedly found that Compound 1 in a crystalline form, named as Crystalline Form A, can only be obtained at a particular conditions, depending on the ee value of the starting materials, and the ratio of the co-solvents and so on. A polymorph study was also performed through methods of slow evaporation, anti-solvent addition, slow cooling, vapor diffusion and polymer-induced crystallization. Most of experiments failed to get crystalline form, which indicates the obtaining of Crystalline Form A is not straight forward.

Further characterization results have revealed that Crystalline Form A is an anhydrate with a melting point of 139.4±2° C. (onset temperature). To evaluate stability, the sample of Crystalline Form A was stored at 80° C. for 2 days, 25° C./60% RH for up to 24 months or 40° C./75% RH condition for up to 6 months, and characterized by XRPD before, during and after the stability test. Results showed no crystal form change was observed for all the above periods, indicating good physical stability of Crystalline Form A at 80° C. or stored at 25° C./60% RH for up to 24 months and at 40° C./75% RH condition for up to 6 months.

In some embodiments, Crystalline Form A has an X-ray powder diffraction pattern comprising diffraction peaks having 2θ angle values independently selected from: approximately 14.8±0.2°, 16.4±0.2° and 21.4±0.2°.

In some embodiments, Crystalline Form A has an X-ray powder diffraction pattern comprising diffraction peaks having 2θ angle values independently selected from: approximately 14.8±0.2°, 15.6±0.2°, 16.4±0.2° and 21.4±0.2°.

In some embodiments, Crystalline Form A has an X-ray powder diffraction pattern comprising diffraction peaks having 2θ angle values independently selected from: approximately 12.2±0.2°, 12.9±0.2°, 14.8±0.2°, 15.6±0.2°, 16.4±0.2° and 21.4±0.2°.

In some embodiments, Crystalline Form A has an X-ray powder diffraction pattern comprising diffraction peaks having 2θ angle values independently selected from: approximately 12.2±0.2°, 12.9±0.2°, 14.8±0.2°, 15.6±0.2°, 16.4±0.2°, 17.7±0.2°, 18.5±0.2°, 20.7±0.2° and 21.4±0.2°.

In some embodiments, Crystalline Form A has an X-ray powder diffraction pattern substantially in accordance with FIG. 1.

In some embodiments, Crystalline Form A has an X-ray powder diffraction pattern summarized in Table 1.

TABLE 1

X-ray Diffraction Pattern of Crystalline Form A			
Peak#	Diffraction angle (2-theta)	Spacing	Relative intensity
1	5.432	16.26908	7.37
2	10.799	8.19295	2.40
3	12.188	7.26191	13.19
4	12.942	6.84040	13.51
5	14.820	5.97780	28.09
6	15.587	5.68534	19.63
7	16.350	5.42177	29.30
8	17.662	5.02158	13.62
9	18.452	4.80853	11.39
10	18.689	4.74791	8.26
11	20.729	4.28515	11.07
12	21.420	4.14847	100.00
13	22.035	4.03409	7.59
14	22.864	3.88958	6.70
15	23.684	3.75673	5.24
16	25.111	3.54646	2.43
17	26.525	3.36044	5.13
18	26.906	3.31381	6.41
19	27.126	3.28741	6.92
20	29.641	3.01393	4.61
21	30.755	2.90724	2.58
22	36.421	2.46692	1.29

In some preferred embodiments, Crystalline Form A has a melting point of $139 \pm 2^\circ \text{C}$. (onset temperature).

In some preferred embodiments, Crystalline Form A has a DSC substantially in accordance with FIG. 2.


In some preferred embodiments, Crystalline Form A has a TGA substantially in accordance with FIG. 3.

In some embodiments, the crystalline Form A is slightly hygroscopic. In some embodiments, the crystalline Form A is unsolvated.

In some embodiments, the crystalline Form A has substantially the same X-ray powder diffraction (XRPD) pattern post storage at 40°C . and 75% RH for up to 6 months. In some embodiments, the crystalline Form A has substantially the same X-ray powder diffraction (XRPD) pattern post storage at 25°C . and 60% RH for up to 24 months.

Also disclosed herein is a crystalline form of Compound BG-13, which has an X-ray powder diffraction pattern substantially in accordance with FIG. 11,

Compound BG-13




In some of embodiments, the crystalline form of BG-13 is a single crystal, which has a unit cell dimensions comprising $a=16.7939(4)\text{\AA}$, $b=7.9871(2)\text{\AA}$, $c=23.5438(5)\text{\AA}$, $\alpha=90.00\text{deg.}$, $\beta=108.0460(10)\text{deg.}$, $\gamma=90.00\text{deg.}$


The inventors have deduced the absolute configurations of Compound 1 to be S from the single crystal X-ray structural analysis of intermediate BG-13.

Also disclosed herein is a method for preparing Compound 1 and deuterium-labeled Compound 1, such as the procedures depicted in Scheme 1. The new synthetic methods and the crystallization/recrystallization procedures of Compound 1 via crystalline Form A disclosed herein overcome many issues associated with the processes reported previously, such as preparation of the key chiral intermediate with $>98\%$ optical purity, improve the purity of Compound 1 to reach the acceptance criteria in the specification, control the impurities in Compound 1 and provide many advantages over the existing processes. Notably, the methods disclosed herein are especially suitable for reproducible, commercial-scale manufacture of Compound 1 in high quality and good yields. In an alternative process, BG-9 or its analogs in Scheme 1 could be asymmetrically reduced with low to excellent enantioselectivities (5% ee. to 95% ee). The process of other steps are similar to those listed in Scheme 1.

Scheme 1: Preparation of Compound 1 and deuterium-labeled Compound 1




-continued




11

12


-continued




13



-continued



Also disclosed herein is a method for preparing the compound of Formula Ia, comprising asymmetrically reducing the compound of Formula I in the presence of the catalyst and/or reductant to produce the compound of Formula Ia,



wherein R¹ is hydrogen or an amino protecting group.

In some embodiments, the amino protecting group includes, but not limit to, acetyl, propionyl, butyryl, phenylacetyl, benzoyl, toluyl, Phenoxyacetyl (POA), methoxycarbonyl, ethoxycarbonyl, 2,2,2-trichloroethoxycarbonyl, tert-

14



butyloxycarbonyl (BOC), 2-iodoethoxycarbonyl, carbobenzoxy (CBZ), 4-methoxybenzyloxycarbonyl, (Fluoren-9-ylmethoxy)carbonyl (Fmoc), 4-methoxy-2,3,6-trimethylbenzenesulfonyl (Mtr), benzyl, methyl or 4-methoxybenzyl.

In some embodiments, wherein the catalyst is a neutral catalyst system or a cationic catalyst system. In some preferred embodiments, the catalyst is an iridium catalyst system including, but not limited to, [Ir(COD)Cl]₂/(R or S)-MeO-Biphep, [Ir(COD)Cl]₂/(R or S)-Binap, [Ir(COD)Cl]₂/(R or S)-Tol-Binap, [Ir(COD)Cl]₂/(R or S)-xyl-Binap, [Ir(COD)Cl]₂/(S,S or R,R)-Diop, [Ir(COD)Cl]₂/(R or S)-P-Phos, [Ir(COD)Cl]₂/(R or S)-Tol-P-Phos, [Ir(COD)Cl]₂/(R or S)-Xyl-P-Phos, [Ir(COD)Cl]₂/(R,R or S,S)-Me-DuPhos, [Ir(COD)Cl]₂/(R or S)-SegPhos, [Ir(μ-Cl)(cod)]₂/(R or S)-Ship, [Ir(μ-Cl)(cod)]₂/(R or S)-Siphos, [Ir(μ-Cl)(cod)]₂/(R or S)-Siphos-PE, [Ir(μ-Cl)(cod)]₂/(R or S)-MonoPhos, [Ir(μ-Cl)(cod)]₂/(R or S)-tol-SDP, [Ir(μ-Cl)(cod)]₂/(S,S or R,R)-Diop, [Ir(μ-Cl)(cod)]₂/(S,R or R,S)-Josiphos, [Ir(μ-Cl)(cod)]₂/(R or S)-Binap, [Ir(μ-Cl)(cod)]₂/(R or S)-MeO-Biphep, [Ir(μ-Cl)(cod)]₂/(R or S)-Synphos, or [Ir(μ-Cl)(cod)]₂/(R or S)-Difluorphosor [Ir(cod)₂]⁺X⁻ (X: e.g. BF₄, NO₃, OTf, PF₆, SbF₆ and BarF) plus related ligands as described above (Wen Bo et al., *J. AM. CHEM SOC.* 125, 10536-10537 2003. Damien et al., *J. Org. Chem.* 77, 4544-4556, 2012. Milos et al., *Org. Process Res. Dev.* 16, 1293-1300, 2012.); a rhodium catalyst system including, but not limited to, [Rh(COD)₂]BF₄ plus ligands described above (Xiang-Ping et al., *Top Organomet Chem* 36, 313-354, 2011); or, a ruthenium catalyst system including, but not limited to, RuCl₂ (R or S)-BINAP/(R or S)-DAIPEN, RuCl₂ (R or S)-BINAP/(R,R or S,S)-DPEN, RuCl₂(S or R)-BINAP (S,S or R,R)-DACH, RuCl₂[(R or S)-Tol-BINAP] [(S,S or R,R)-DPEN], RuCl₂ (R,R or S,S)-Me-DuPHOS/(R,R or S,S)-DPEN, RuCl₂ (R,R or S,S)-Et-DuPHOS/(R,R or S,S)-DPEN, RuCl₂ (R,R or S,S)-Et-DuPHOS/(R,R or S,S)-DACH, RuCl₂(S,S or R,R)-i-Pr-DuPHOS/(R,R or S,S)-DPEN, RuCl₂ (R or S)-HexaPHEMP/(R,R or S,S)-DPEN, RuCl₂ (R or S)-MeO-BIPHEP/(R,R or S,S)-DPEN (Christopher et al., *Adv. Synth. Catal.* 345, 195-201, 2003. Julian et al., *Adv. Synth. Catal.* 345, 300-307, 2003.).

The above method was found to produce excellent enantioselectivities up to 95% ee by using the above catalyst, especially the neutral or cationic iridium catalyst system.

15

Also disclosed herein is a method for resolving the compound of Formula IIa to produce the compound of Formula IIb, or improving the chiral purity of the compound of Formula IIb, comprising treating the racemic compound of Formula IIa with a chiral acid,





wherein R^1 is hydrogen, methyl, benzyl, 4-methoxybenzyl or the other conventional amino protecting groups as mentioned above.

In some embodiments, the chiral acid includes, but not limited to, L-malic acid, D-malic acid, L-Mandelic acid, D-Mandelic acid, L-camphorsulfonic acid, D-camphorsulfonic acid, L-tartaric acid, D-tartaric acid, L-DBTA, D-DBTA, L-DTTA, or D-DTTA.

Also disclosed herein is a method for resolving a compound of Formula Ic to produce a compound of Formula Id or improving the chiral purity of formula Id, comprising treating the racemic compound of Formula Ic with a chiral acid,


16



wherein R^1 is hydrogen, methyl, benzyl, 4-methoxybenzyl or the other conventional amino protecting groups as mentioned above.


In some embodiments, the chiral acid includes, but not limited to, L-malic acid, D-malic acid, L-Mandelic acid, D-Mandelic acid, L-camphorsulfonic acid, D-camphorsulfonic acid, L-tartaric acid, D-tartaric acid, L-DBTA, D-DBTA, L-DTTA, or D-DTTA.

Also disclosed herein is a compound of Formula Ie or a salt thereof, or Formula If or a salt thereof used to prepare Compound 1,



17

-continued



Further, the present also provides methods of preparing Crystalline Form A. The crystalline form disclosed herein can be prepared by crystallizing the compound disclosed herein from a suitable solvent system comprising at least one solvent, which can be achieved by methods of spontaneous precipitation (evaporation), cooling, and/or adding anti-solvent (in which the compound disclosed herein has relatively lower solubility), in order to achieve oversaturation in a solvent system. Crystallization can also be achieved by using or not using crystal seeds which is suitable for crystallizing the crystalline forms disclosed herein.

In some embodiments, the method of preparing Crystalline Form A comprises the steps of dissolving (S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (Compound 1) in DCM, swapping to solvent EA, recrystallizing from EA/MTBE, to obtain the target crystalline form.

In some embodiments, the method of preparing Crystalline Form A comprises the steps of dissolving Compound 1 in EA, adding hexane, to obtain the target crystalline form.

In some embodiments, the method of preparing Crystalline Form A is achieved by adding an anti-solvent into the solution of the solid Compound 1 or crude Form A in a solvent for dissolving the solid, wherein the anti-solvent including, but not limited to, H₂O and n-heptane, and the solvent for dissolving the solid including, but not limited to, acetone, DMAc, EtOAc, DCM, Toluene, and 2-MeTHF.

In some embodiments, the method of preparing Crystalline Form A is achieved by adding the solution of the solid Compound 1 or crude Form A in a solvent into an anti-solvent, and allow sufficient time for organic vapor to interact with the solution in a sealed reactor, wherein the solvent including, but not limited to, acetone, and EtOAc, and the anti-solvent including, but not limited to, n-heptane.

Also disclosed herein is a pharmaceutical composition comprises a therapeutically effective amount of Crystalline Form A, and a pharmaceutically acceptable excipient. In some embodiments, the pharmaceutical composition is used in an oral administration. In some preferred embodiments, the pharmaceutical composition comprises 1 wt % to 99 wt % of Crystalline Form A. In some more preferred embodiments, the pharmaceutical composition comprises 1 wt % to 70 wt % of Crystalline Form A. In some most embodiments, the pharmaceutical composition comprises 10 wt % to 30 wt % of Crystalline Form A.

The present invention also provide a method of treating or preventing a disease associated with undesirable Btk activity in a subject by administering to a subject Crystalline Form A.

18

The present invention also provide a method of treating or preventing a disease selected from an allergic disease, an autoimmune disease, an inflammatory disease, a cancer, or a combination of two or more thereof in a subject by administering to the subject Crystalline Form A.

The present invention also provide a method of treating or preventing a B-cell proliferative disease in a subject by administering Crystalline Form A to the subject.

In some embodiments, the B-cell proliferative disease is B-cell malignancies including but not limited to, lymphoma, non-Hodgkin's lymphoma (NHL), diffuse large B cell lymphoma (DLBCL), mantle cell lymphoma (MCL), follicular lymphoma (FL), chronic lymphocytic leukemia (CLL), small lymphocytic lymphoma (SLL), Waldenstrom macroglobulinemia (WM), marginal zone lymphoma (MZL), Hairy cell leukemia (HCL), Burkitt's-like leukemia (BL).

In some embodiments, the B-cell proliferative disease is relapsed/refractory (R/R) B-cell malignancies including, but limited to, R/R MCL, R/R CLL, R/R SLL, R/R WM.

The Crystalline Form A disclosed herein can be used in manufacturing a medicament for treatment of at least one disease associated with undesirable Btk activity, in a subject.

The Crystalline Form A disclosed herein can be used in manufacturing a medicament for the treatment of a disease selected from an allergic disease, an autoimmune disease, an inflammatory disease, a cancer, or a combination of two or more thereof, in a subject.

The Crystalline Form A disclosed herein can be used in manufacturing a medicament for the treatment of a B-cell proliferative disease selected from B-cell malignancies, or relapsed/refractory B-cell malignancies, in a subject.

The updated clinical trials continue to demonstrate that Compound 1 is well tolerated in treatment naïve (TN) and relapsed/refractory (R/R) B-cell malignancies, eg., in WM, with a very good partial response (VGPR) rate of over 40% in an evaluable population of 42 patients and with an overall response rate (ORR) of 90% in 42 efficacy-evaluable patients with a median follow-up time of 12.3 months, and in CLL/SLL, with a high overall response rate (94%) and a very low treatment discontinuation rate (3%) at a median follow-up of 10.5 months for efficacy evaluation.

Definitions

Unless specifically defined elsewhere in this document, all other technical and scientific terms used herein have the meaning commonly understood by one of ordinary skill in the art to which this invention belongs.

As used herein, including the appended claims, the singular forms of words such as "a", "an", and "the", include their corresponding plural references unless the context clearly dictates otherwise. Thus, for example, reference to "a crystalline form" includes one or more of such different crystalline forms and reference to "the method" includes reference to equivalent steps and methods known to those of ordinary skill in the art that could be modified or substituted for the methods described herein.

As disclosed herein, the crystalline form is an approximately pure crystalline. The term "approximately pure" as herein used refers to at least 85 wt %, preferably at least 95 wt %, more preferably at least 99 wt % of Crystalline Form A disclosed herein.

For crystalline forms disclosed herein, only the main peaks (i.e., the most characteristic, significant, unique and/or reproducible peaks) are summarized; additional peaks may be obtained from the diffraction spectra by conventional methods. The main peaks described above can be repro-

duced within the margin of error (± 2 at the last given decimal place, or ± 0.2 at the stated value).

As disclosed herein, “an X-ray powder diffraction pattern substantially in accordance with FIG. 1” refers to the X-ray powder diffraction pattern that show major peaks as in FIG. 1, wherein major peaks refer to those with the relative intensity greater than 10%, preferably greater than 20%, relative to the highest peak (with its relative intensity designated to be 100%) in FIG. 1.

Throughout this specification and the claims which follow, unless the context requires otherwise, the word “comprise”, and variations such as “comprises” and “comprising”, will be understood to imply the inclusion of a stated integer or step or group of integers or steps but not the exclusion of any other integer or step or group of integer or step. When used herein the term “comprising” can be substituted with the term “containing” or sometimes when used herein with the term “having”.

The term “therapeutically effective amount” as herein used, refers to the amount of a compound that, when administered to a subject for treating a disease, or at least one of the clinical symptoms of a disease or disorder, is sufficient to affect such treatment for the disease, disorder, or symptom. The “therapeutically effective amount” can vary with the compound, the disease, disorder, and/or symptoms of the disease or disorder, severity of the disease, disorder, and/or symptoms of the disease or disorder, the age of the subject to be treated, and/or the weight of the subject to be treated. An appropriate amount in any given instance can be apparent to those skilled in the art or can be determined by routine experiments. In the case of combination therapy, the “therapeutically effective amount” refers to the total amount of the combination objects for the effective treatment of a disease, a disorder or a condition.

The pharmaceutical composition comprising the compound disclosed herein can be administered via oral, inhalation, rectal, parenteral or topical administration to a subject in need thereof. For oral administration, the pharmaceutical composition may be a regular solid formulation such as tablets, powder, granule, capsules and the like, a liquid formulation such as water or oil suspension or other liquid formulation such as syrup, solution, suspension or the like; for parenteral administration, the pharmaceutical composition may be solution, water solution, oil suspension concentrate, lyophilized powder or the like. Preferably, the formulation of the pharmaceutical composition is selected from tablet, coated tablet, capsule, suppository, nasal spray or injection, more preferably tablet or capsule. The pharmaceutical composition can be a single unit administration with an accurate dosage. In addition, the pharmaceutical composition may further comprise additional active ingredients.

All formulations of the pharmaceutical composition disclosed herein can be produced by the conventional methods in the pharmaceutical field. For example, the active ingredient can be mixed with one or more excipients, then to make the desired formulation. The “pharmaceutically acceptable excipient” refers to conventional pharmaceutical carriers suitable for the desired pharmaceutical formulation, for example: a diluent, a vehicle such as water, various organic solvents, etc, a filler such as starch, sucrose, etc a binder such as cellulose derivatives, alginates, gelatin and polyvinylpyrrolidone (PVP); a wetting agent such as glycerol; a disintegrating agent such as agar, calcium carbonate and sodium bicarbonate; an absorption enhancer such as quaternary ammonium compound; a surfactant such as hexadecanol; an absorption carrier such as Kaolin and soap clay; a lubricant such as talc, calcium stearate, magnesium stear-

ate, polyethylene glycol, etc. In addition, the pharmaceutical composition further comprises other pharmaceutically acceptable excipients such as a decentralized agent, a stabilizer, a thickener, a complexing agent, a buffering agent, a permeation enhancer, a polymer, aromatics, a sweetener, and a dye.

The term “disease” refers to any disease, discomfort, illness, symptoms or indications, and can be interchangeable with the term “disorder” or “condition”.

Abbreviations:

AcOH	Acetic acid
AEs	Adverse events
BID	Twice a day
CLL	Chronic lymphocytic leukemia
Con.	Concentrated
D-DBTA	(2S, 3S)-Dibenzoyl tartaric acid
DDQ	2,3-Dichloro-5,6-dicyano-1,4-benzoquinone
DCM	Dichloromethane
DIEA	N,N-diisopropylethylamine
DLBCL	Diffuse large B cell lymphoma
DMAc	N,N-dimethylacetamide
DMF	N,N-dimethylformamide
DMF-DMA	N,N-dimethylformamide dimethyl acetal
DMSO	Dimethylsulfoxide
DSC	Differential Scanning Calorimetry
DVS	Dynamic Vapor Sorption
EA	Ethyl Acetate, EtOAc
EDCI	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
EtOH	Ethanol
FL	Follicular lymphoma
GC	Gas Chromatograph
GCMS	Gas Chromatography-Mass Spectrometry
HOAc	Acetic Acid
HOBt	Hydroxybenzotriazole
HPLC	High Performance Liquid Chromatography
IPA	Isopropyl alcohol
IPAc	Isopropyl acetate
IPC	In Process Control
KF	Karl-Fischer
L-DBTA	(2R, 3R)-Dibenzoyl tartaric acid
LOQ	Limit of Quantification
MCL	Mantle cell lymphoma
MeCN or ACN	Acetonitrile
MeMgBr	Methyl Magnesium Bromide
MeOH	Methanol
2-MeTHF	2-Methyltetrahydrofuran
MIBE	4-methyl-2-pentanone
MsOH	Methanesulfonic Acid
MTBE	Methyl tertiary butyl ether
NHL	non-Hodgkin's lymphoma
NLT	not less than
NMP	1-Methyl-2-pyrrolidone
NMR	Nuclear Magnetic Resonance
NMT	Not more than
ORR	Overall response rate
Pd	Palladium
pH	Hydrogen ion concentration
POA	Phenoxyacetyl
QD	Once a day
RH	Relative Humidity
SLL	Small lymphocytic lymphoma
RT	Room Temperature
TEA	Triethylamine
TGA	Thermo-gravimetric Analysis
THF	Tetrahydrofuran
TN	Treatment naïve
VGPR	very good partial response
XRPD	X-ray Powder Diffraction
WM	Waldenstrom macroglobulinemia


Example

The present invention is further exemplified, but not limited, by the following examples that illustrate the invention.

21


Example 1 Preparation of (S)-7-(1-Acryloylpiperidin-4-yl)-2-(4-Phenoxyphenyl)-4,5,6,7-Tetrahydropyrazolo[1,5-a]Pyrimidine-3-Carboxamide (Compound 1) and Crystalline Form a Thereof

Step 1: Synthesis of BG-2



Under nitrogen atmosphere, to a solution of EA (5 v), HOBT (1.2 eq.), EDCI (1.2 eq.), 4-phenoxybenzoic acid (BG-1, 80 Kg, 1.0 eq.) and malononitrile (1.2 eq.) was added TEA (2.4 eq.) at 10° C. The mixture was then stirred at RT until the reaction was completed. The mixture was then centrifuged and the cake was washed with EA. The filtrate was washed with aqueous NaHCO₃ twice and NH₄Cl. The organic phase was washed with 1.5 N H₂SO₄ twice and stirred. Concentrated, precipitated from methanol and purified water. The solid was collected by centrifugation and dried under vacuum. This gave 79.9 Kg of BG-2. ¹H NMR (DMSO-d₆) δ 7.62 (d, J=8.6 Hz, 2H), 7.46-7.38 (m, 2H), 7.18 (t, J=7.4 Hz, 1H), 7.06 (d, J=8.0 Hz, 2H), 6.94 (d, J=8.6 Hz, 2H).

Step 2: Synthesis of BG-3



Under nitrogen atmosphere, a solution of BG-2 (79.9 kg, 1.0 eq.) in MeCN (5.0 v) was added into trimethoxymethane (12.0 v) at 85° C. The resultant mixture was stirred until the reaction was completed. Sampled for HPLC analysis. Concentrated under vacuum. The residue was precipitated from i-PrOH and hexane. The mixture was centrifuged, and the cake was washed with hexane and dried under vacuum. This gave 71.7 Kg of product. ¹H NMR (400 MHz, DMSO-d₆) δ

22

7.70 (d, J=8.4 Hz, 2H), 7.52-7.45 (m, 2H), 7.28 (t, J=7.6 Hz, 1H), 7.22-7.06 (m, 4H), 3.93 (s, 3H).

Step 3: Synthesis of BG-4

5

BG-4

BG-2

10

15

20

25

30

Steps 4 to 6: Synthesis of BG-8

35


40

BG-3

45

50

55




To a mixture of DCM (8.0 v), BG-5 (80.0 Kg, 1.0 eq.), N,O-dimethylhydroxylamine hydrochloride (1.2 eq.), HOBT (1.2 eq.) and EDCI (1.2 eq.), TEA (2.6 eq.) was charged dropwise below 15° C. the mixture was stirred at RT until the reaction was completed, centrifuged and the cake was washed with DCM (1.0 v) twice. The filtrate was washed with 20% aqueous NH₄Cl (3×4.0 v). The filtrate was concentrated under vacuum to give the crude product BG-6,

23

which was used in the next step without further purification. The residue was dissolved in toluene (5.0 v) and THF (1.0 v), cooled to 10° C., charged dropwise MeMgBr (1.4 eq.) at 10° C. and then stirred at RT until the reaction was completed. The solution was cooled below 10° C. Saturated aqueous NH₄Cl was charged dropwise below 10° C. The mixture was centrifuged, separated, filtrated, and the organic phase was washed with aqueous NaCl twice. The organic phase was concentrated to give the crude product, which was used in the next step without further purification. The residue in DMF (2.5 v) and DMF-DMA (2.5 v) was stirred at 110° C. until the reaction was completed. The reaction mixture was cooled, concentrated and then DCM was added. The final mixture was washed with saturated aqueous NH₄Cl. The organic layer was concentrated and precipitated by charging hexane. The mixture was centrifuged and the cake was collected. The cake was dried under vacuum. This gave 82.2 Kg of the desired product. ¹H NMR (DMSO-d₆) δ 7.49 (d, J=12.6 Hz, 1H), 5.01 (d, J=12.6 Hz, 1H), 3.99-3.82 (m, 2H), 3.14-2.94 (m, 2H), 2.89-2.61 (m, 6H), 2.49-2.37 (m, 1H), 1.66-1.56 (m, 2H), 1.39 (s, 9H), 1.39-1.20 (m, 2H).

Step 7: Synthesis of BG-9




Under nitrogen atmosphere, a mixture of toluene (8.0 v), AcOH (0.5 v), BG-8 (1.2 eq.) and BG-4 (66.9 Kg 1.0 eq.) was heated to 95° C. and stirred until the reaction was completed. The mixture was cooled, concentrated and precipitated from methanol. The mixture was centrifuged and the cake was washed with methanol. The cake was dried under vacuum. This gave 107.8 Kg of product. ¹H NMR (DMSO-d₆) δ 8.78 (d, J=4.6 Hz, 1H), 8.15-8.07 (m, 2H), 7.51-7.41 (m, 2H), 7.34 (d, J=4.6 Hz, 1H), 7.27-7.19 (m, 3H), 7.17-7.10 (m, 2H), 4.24-4.02 (m, 2H), 3.81-3.69 (m, 1H), 3.12-3.82 (m, 2H), 2.15-2.04 (m, 2H), 1.76-1.60 (m, 2H), 1.43 (s, 9H).

24

Step 8: Synthesis of BG-10


BG-10



To a mixture of THF (10.0 v), BG-9 (13.0 Kg, 1.0 eq.) and D-DBTA (1.0 eq) under N₂ was charged Pd/C (10% w/w), hydrogen gas was introduced into the reactor and the hydrogen pressure was maintained to 1.8 MPa. The reactor was heated to 40° C. slowly and stirred until the reaction was completed. The mixture was then cooled, filtered, and the cake was washed with THF. The filtrate was collected, and concentrated under vacuum. DCM was added. The residue was washed with aq. NaHCO₃, concentrated and precipitated from MTBE and hexane, then centrifuged. The cake was collected and dried under vacuum to give the desired compound (yield:94.8% and purity:98.5%). ¹H-NMR (DMSO-d₆) δ 7.82-7.76 (m, 2H), 7.56-7.51 (m, 1H), 7.45-7.37 (m, 2H), 7.21-7.14 (m, 1H), 7.12-7.03 (m, 4H), 4.09-3.91 (m, 3H), 3.30-3.22 (m, 2H), 2.82-2.55 (m, 2H), 2.18-1.99 (m, 2H), 1.98-1.86 (m, 1H), 1.69-1.58 (m, 1H), 1.56-1.45 (m, 1H), 1.38 (s, 9H), 1.32-1.13 (m, 2H).

Step 9: Synthesis of BG-11


BG-11



To a solution of BG-10 (100.0 Kg 1.0 eq.) in DCM (6.0 v) was added dropwise HCl in EtOH (20.9% w/w, 2.0 v) under nitrogen atmosphere. The mixture is stirred until the reaction was completed. MTBE (4.0 v) was added to the solution, cooled. The cakes was collected by centrifugation and washed with hexane (2.0 V), then the cake was slurried in hexane (5 v), and centrifuged again. The cake was washed with hexane (2.0 V) and dried under vacuum. This gave 85.2


25

Kg product. $^1\text{H-NMR}$ (DMSO-d_6) δ 9.25-8.85 (m, 2H), 7.84-7.70 (m, 2H), 7.47-7.37 (m, 2H), 7.18 (t, $J=7.4$ Hz, 1H), 7.12-7.03 (m, 4H), 5.73 (br s, 2H), 4.12-4.03 (m, 1H), 3.25-3.19 (m, 4H), 2.90-2.73 (m, 2H), 2.28-2.12 (m, 1H), 2.10-2.00 (m, 1H), 1.99-1.86 (m, 1H), 1.84-1.52 (m, 4H).
Step 10: Synthesis of BG-11A



A mixture of BG-11 (85.0 Kg, 1.0 eq) in water (6.0 v) and NaOH (3.0 eq) was stirred until the reaction was completed at RT. The cake was collected and slurried in MTBE (6.0 v). The mixture was then centrifuged to collect the cake. The cake was dried under vacuum. This gave 71.3 Kg product.
 $^1\text{H-NMR}$ (DMSO-d_6) δ 7.82-7.74 (m, 2H), 7.54-7.49 (m, 1H), 7.45-7.38 (m, 2H), 7.21-7.14 (m, 1H), 7.12-7.04 (m, 4H), 4.03-3.95 (m, 1H), 3.29-3.21 (m, 2H), 3.00-2.87 (m, 2H), 2.46-2.31 (m, 2H), 2.11-1.83 (m, 3H), 1.58-1.12 (m, 4H).

Step 11: Synthesis of BG-11B



A mixture of ethanol/water/acetic acid (7:3:1, 46 v) and BG-11A (30 kg, 1.0 eq.) in a reactor was heated to $70\pm 5^\circ\text{C}$. under nitrogen atmosphere, then a solution of D-DBTA (1.20 eq.) in ethanol/water/acetic acid (7:3:1, 4 v) was added dropwise with the temperature not less than 65°C . The resulting solution was stirred for 16 hrs at $60-65^\circ\text{C}$., then cooled to RT. The solid was collected by centrifugation and washed with ethanol (2.0 v). The cake was slurried in the mixed solvent of ethanol/water/AcOH (7:3:1, 20 v) for 16 hrs at 55°C . and cooled to RT. The solid was collected by centrifugation, washed with ethanol (2.0 v). The cake was

26

dried under vacuum (Yield: 37.9%). $^1\text{H-NMR}$ (DMSO-d_6) δ 8.76 (br s, 2H), 7.99-7.89 (m, 4H), 7.83-7.75 (m, 2H), 7.66-7.57 (m, 3H), 7.52-7.45 (m, 4H), 7.45-7.39 (m, 2H), 7.21-7.14 (m, 1H), 7.13-7.03 (m, 4H), 5.64 (s, 2H), 4.08-4.00 (m, 1H), 3.29-3.19 (m, 4H), 2.85-2.72 (m, 2H), 2.21-1.40 (m, 7H).
Step 12: Synthesis of BG-11C

BG-11A


10

15

20

25

BG-11C



30

35

40

45

50

55

60


65

To a mixture of dichloromethane (15.0 v) and 20.0% aqueous KOH (3.0 v) was added bachwise BG-11B (48.0 kg, 1.0 eq.) under nitrogen atmosphere at RT. After the reaction was completed, the organic layer was collected and the water layer was extracted with dichloromethane (5.0 v). The organic layers were combined. Con. HCl (0.36 v) was added to the above organic layers at RT. The resulting mixture was stirred until the reaction was completed. The solid was collected by centrifugation and washed with dichloromethane (1.0 v). The collected solid was slurried with MTBE (6.0 v). The solid was collected by centrifugation and washed with MTBE (1.0 v), then was dried under vacuum. This gave 31.5 Kg product (Yield:100%).

Step 12: Synthesis of BG-11D (Alternative Intermediate)

ACN (5.0 v), soft water (10.0 v), KOH (5.0 eq) was charged to a reactor and stirred for at least 15 min. BG-11B (1.0 eq) was charge to the reactor in portion-wise. The mixture was stirred until the reaction was completed. The cake was collected by centrifugation, slurried in ACN (1.0 v) and soft water (5.0 v), and dried under vacuum to give the product.

Step 13: Synthesis of BG-12




BG-12

27

A solution of BG-11C (15.0 Kg 1.0 eq.) in MsOH (2.5 v) was stirred at 85° C. under nitrogen atmosphere until the reaction was completed. After cooling to 5° C. purified water (4.0 v) was added dropwise to the system and kept the temperature not more than 35° C. (temperature increased obviously). The resulting solution was stirred for 16 hrs at 30° C., and then washed with DCM (2×3.0 v). The aqueous phase was collected. DCM (6.0 v) was added to the aqueous phase, the mixture was cooled to 5° C. The pH value was adjusted to 11.2 with 20% aqueous NaOH (temperature increased obviously) with stirring with the temperature not more than 30° C. The organic phase was separated and collected. The aqueous was extracted with DCM (3.0 v). The organic layers were combined and concentrated. MTBE (4.0 v) was added to the residue. The mixture was then concentrated and precipitated from n-heptane. The solid was collected by centrifugation and dried in a vacuum oven. This gave 12.55 Kg product (Yield: 94.9%). ¹H-NMR (DMSO-d₆) δ 7.52-7.46 (m, 2H), 7.45-7.38 (m, 2H), 7.21-7.13 (m, 1H), 7.12-7.03 (m, 4H), 6.64 (s, 1H), 3.99-3.90 (m, 1H), 3.29-3.22 (m, 2H), 3.03-2.90 (m, 2H), 2.48-2.36 (m, 2H), 2.03 (dd, J=13.9, 5.6 Hz, 2H), 2.14-1.99 (m, 1H), 1.97-1.85 (m, 1H), 1.65-1.15 (m, 3H).


Step 14: Synthesis of BG-13



A mixture of MeOH (13.5 v), purified water (4.5 v) and BG-12 (8.5 Kg, 1.0 eq.) in a reactor was heated to 50° C. under N₂ atmosphere. To the mixture was charged dropwise a solution of L-DBTA (0.7 eq) in MeOH/purified water (1.5 v/0.5 v) while keeping the temperature at 50° C. After addition, the mixture was stirred for at least 2 hrs at 50° C., and then cooled to RT and stirred for at least 16 hrs at RT. The cake was collected by Centrifugation and was washed with MeOH (2.0 v). The cake was dried in a vacuum oven. This gave 9.08 Kg product (Yield: 74.8%, ee value >98%).

28

Step 15: Synthesis of (S)-7-(1-Acryloylpiperidin-4-yl)-2-(4-Phenoxyphenyl)-4,5,6,7-Tetrahydropyrazolo[1,5-a]Pyrimidine-3-Carboxamide (Compound 1)




Under N₂ atmosphere, ACN (12.0 v), water (12.5 v), BG-13 (8.0 Kg, 1.0 eq), and NaHCO₃ (2.5 eq.) were added to a reactor. The mixture was then cooled to -5-0° C. To the mixture, the solution of acryloyl chloride (1.1 eq.) in MeCN (0.5 v) was added dropwise and stirred until the reaction was completed. EA (6.0 v) was then added to the reactor, and stirred. The organic phase was collected. The aqueous layer was further extracted with EA (3.0 v). The organic phases were combined and washed with brine. The organic layer was collected and concentrated.

BG-13

The residue was purified by silica gel (2 wt) column, eluted with 3% w/w methanol in DCM (21.0 v). The Compound 1 solution was collected and concentrated under vacuum. The residue was precipitated from EA/MTBE (2.0 v). The cake was collected by centrifugation as the product.

Step 15: Synthesis of (S)-7-(1-Acryloylpiperidin-4-yl)-2-(4-Phenoxyphenyl)-4,5,6,7-Tetrahydropyrazolo[1,5-a]Pyrimidine-3-Carboxamide (Compound 1, Alternative Method)



A mixture of CH₃CN (10.0 v), purified water (5.0 v), NaOH (1.5 eq.) and BG-13 (1.0 eq.) was stirred to get a clear solution. EtOAc (6.0 v) was then charged to the reaction and separated. The organic phase was collected and washed with 15% brine (3.0 v) twice. The organic phase prepared above was concentrated and the solvent was swapped to CH₃CN (residue volume: NMT 5.0 v). CH₃CN (7.5 v) and purified

water (12.5 v) were charged and cooled to 15-20° C. L-(+)-tartaric acid (0.5 eq) and NaHCO₃(2.5 eq.) were charged to the reaction mixture. A solution of acryloyl chloride (1.1 eq.) in CH₃CN (0.5 v) was charged drop-wise to the reaction mixture. After the reaction was completed, EtOAc (6.0 v) was charged to the reaction mixture and organic layer was collected. Aqueous phase was further extracted with EA (3.0 v). The organic layers were combined, washed with 15% brine (5.0 v) and concentrated. The solvent was swapped to DCM (volume of residue: 1.5-2.0 v) and purified by silica gel column (silica gel: 100-200 mesh, 2.0 w/w; eluent: 3% w/w MeOH in DCM (about 50 v). The collected solution was concentrated and swapped to EtOAc (4.0 v). MTBE (6.4 v) was charged drop-wise to residue at 50° C. The mixture was then cooled to 5° C. and the cake was collected centrifugation.

Step 16: Preparation of Crystalline Form a of Compound 1

The above cake of Compound 1 was dissolved in 7.0 volumes of DCM, and then swapped to solvent EA. After recrystallization from EA/MTBE, the cakes was collected by centrifugation, and was dried under vacuum. This gave 4.44 Kg product (Yield: 70.2%).

The product was then characterized by X-ray powder diffraction (XRPD) pattern method, which was generated on a PANalytical Empyrean X-ray powder diffractometer with the XRPD parameters as follows: X-Ray wavelength (Cu, $\text{K}\alpha$, $\text{K}\alpha_1$ (Å): 1.540598, $\text{K}\alpha_2$ (Å): 1.544426; $\text{K}\alpha_2/\text{K}\alpha_1$ intensity ratio: 0.50); X-Ray tube setting (45 Kv, 40 mA); divergence slit (automatic); scan mode (Continuous); scan range (° 2TH) (3°-40); step size (° 2TH) (0.0131); scan speed (°/min) (about 10). The XRPD result found the resultant product as a crystalline shown in FIG. 1.

The differential scanning calorimetry (DSC) curves shown as in FIG. 2 was generated on a TA Q2000 DSC from TA Instruments. The DSC parameters used includes: temperature (25° C.-desired temperature); heating rate (10° C./min); method (ramp); sample pan (aluminum, crimped); purge gas (N₂). DSC result showed a sharp melting point at 139.4° C. (onset temperature).

The thermo-gravimetric analysis (TGA) curves shown as in FIG. 3 was generated on a TA Q5000 TGA from TA Instruments. The TGA parameters used includes: temperature (RT-desired temperature); heating rate (10° C./min); method (ramp); sample pan (platinum, open); purge gas (N₂). TGA result showed is anhydrous with no weight loss even up to 110° C.

The proton nuclear magnetic resonance (¹H-NMR) shown as in FIG. 4 was collected on a Bruker 400M NMR Spectrometer in DMSO-d₆. ¹H-NMR (DMSO-d₆) δ 7.50 (d, J=8.6 Hz, 2H), 7.46-7.38 (m, 2H), 7.17 (t, J=7.6 Hz, 1H), 7.08 (d, J=7.6 Hz, 2H), 7.05 (d, J=8.8 Hz, 2H), 6.85-6.72 (m, 1H), 6.67 (s, 1H), 6.07 (dd, J=16.8, 2.2 Hz, 1H), 5.64 (dd, J=10.4 Hz, 2.2 Hz, 1H), 4.55-4.38 (m, 1H), 4.17-3.94 (m, 2H), 3.33-3.22 (m, 2H), 3.08-2.88 (m, 1H), 2.67-2.51 (m, 1H), 2.36-2.15 (m, 1H), 2.12-1.82 (m, 2H), 1.79-1.65 (m, 1H), 1.63-1.49 (m, 1H), 1.38-1.08 (m, 2H).

The carbon nuclear magnetic resonance (¹³C-NMR) shown as in FIG. 5 was collected on a Bruker 400M NMR Spectrometer in DMSO-d₆. ¹³C-NMR spectra for Crystalline Form A of Compound 1.

Example 2 Preparation of Crystalline Form A of Compound 1

(S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (Compound 1) was prepared by the method disclosed in

WO2014173289A, and further lyophilized to obtain amorphous form of Compound 1. A solution of Compound 1 (200 mg, ee value >97%) in EA (8 mL) was heated to 50° C., to the above solution was added dropwise hexane (8 mL) at 50° C. The mixture was cooled to RT and stirred for 16 hr then was filtered to give 110 mg as a white solid. The solid obtained were characterized by XRPD to be Form A.

Example 3 Preparation of Crystalline Form A of Compound 1 (Anti-solvent Addition)

About 15 mg of sample (Crystalline Form A) was weighed into a 20-mL glass vial, followed by the addition of 0.4-1.2 mL corresponding solvent (see Table 2) to dissolve all the solid. The mixture was then magnetically stirred at the speed of 800 rpm to get a clear solution at RT. Subsequently, the relative anti-solvent (see Table 2) was added to the solution to induce precipitation or until the total amount of anti-solvent reached 15.0 mL. If no precipitation occurs, the solution was then transferred to slow evaporation at RT. The solids obtained were characterized by XRPD to be Form A.

TABLE 2

Anti-Solvent Addition Experiments		
Experiment ID	Solvent	Anti-solvent
1	Acetone	H ₂ O
2	DMAc	H ₂ O
3	EtOAc	n-heptane
4	DCM	n-heptane
5	Toluene	n-heptane
6	2-MeTHF	n-heptane

Example 4 Preparation of Crystalline Form A of Compound 1 (Solution Vapor Diffusion)

About 15 mg of sample (Crystalline Form A) was dissolved in 0.5-1.5 mL of the corresponding solvent (acetone or EtOAc) to obtain a clear solution in a 3-mL vial. Subsequently, the solution was placed into a 20-mL vial with 3 mL of relative anti-solvent (n-heptane). The 20-mL vial was sealed with a cap and kept at RT, allowing sufficient time for organic vapor to interact with the solution. At the end of 11 days, clear solutions were transferred to evaporation at RT. The solid obtained were characterized by XRPD to be Form A.

Example 5 Stability Test of Crystalline Form A of Compound 1 and Purity of Compound 1

(1) Physical Stability Test

The Crystalline Form A of Compound 1 was stored at 80° C. for two days as a thermo-stability test, and the XRPD patterns before and after the test showed no crystal form change.

The long term stability studies of Crystalline Form A of Compound 1 showed there was no significant chemical purity change occurred when stored at 25° C./60% RH for up to 24 months (% area: T0=99.2% and T12=99.2%) and at 40° C./75% RH condition for up to 6 months (% area: T0=99.1% and T6=99.4%). In addition, no crystal form and optical purity changes were observed when stored at 25° C./60% RH for up to 24 months and at 40° C./75% RH condition for up to 6 months.

(2) Hygroscopic Test

The dynamic vapor sorption (DVS) plots shown as in FIG. 6 was collected a SMS (Surface Measurement Systems) DVS Intrinsic. The DVS parameters used includes: temperature (25° C.); dm/dt (0.002%/min); Min. dm/dt stability duration (10 min); Max. equilibrium time (180 min); RH range (0% RH to 95% RH); RH step size (10% RH from 0% RH to 90% RH, 5% RH from 90% RH to 95% RH). As shown in FIG. 6, there is a very slight increase of mass at 80% RH, which was about 0.8% for Crystalline Form A of Compound 1.


(3) Crystallization/Recrystallization Via Form a to Improve the Purity of Compound 1

Crystallization/Recrystallization via Form A is an efficient way to improve the purity of Compound 1 and control the impurities in Compound 1 to reach the acceptance criteria in the specification. See an example as shown in Table 3.

TABLE 3

Purity Changing after Crystallization/Recrystallization via Form A	
Conditions	Purity of Compound 1
After Silica Gel Chromatography Purification	98.5% area
After First Recrystallization	99.3% area
After Second Recrystallization	99.5% area

Example 6 Preparation of Deuterium-Labeled (S)-7-(1-Acryloylpiperidin-4-yl)-2-(4-Phenoxyphenyl)-4,5,6,7-Tetrahydropyrazolo[1,5-a]Pyrimidine-3-Carboxamide (Deuterium-Labeled Compound 1)



To a solution of acrylic-2,3,3-d3 acid (50 mg, 0.67 mmol) and DMF (one drop) in DCM (20 mL) was added dropwise oxalyl chloride (1.6 N, 40.9 mL, 65.5 mmol) at 05° C. then stirred for 2 hours at RT. The mixture was concentrated under reduced pressure to give the crude acryloyl-d₃ chloride.

To a solution of (S)-2-(4-phenoxyphenyl)-7-(piperidin-4-yl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (dissociated from BG-13, see step 15, Compound 1, alternative method; 278 mg, 0.67 mmol) in DCM (20 mL) and aqu. NaHCO₃ (10 mL) was added dropwise a solution of the above acryloyl-d₃ chloride in DCM (5 mL) at 0-5° C. and stirred for 2 hours at RT. The organic combined layers were dried over anhydrous sodium sulfate, filtered, concen-

trated under reduced pressure and purified by prep-TLC to afford 55 mg (17.5%) of (S)-7-(1-(acryloyl-d₃)piperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide as an off-white solid. ¹H NMR (400 MHz, DMSO) δ 7.50-7.44 (m, 2H), 7.42-7.35 (m, 2H), 7.17-7.10 (m, 1H), 7.09-6.99 (m, 4H), 6.64 (s, 1H), 4.52-4.40 (m, 1H), 4.10-3.95 (m, 2H), 2.29-3.25 (m, 2H), 3.04-2.86 (m, 1H), 2.63-2.50 (m, 1H), 2.32-2.13 (m, 1H), 2.06-1.81 (m, 2H), 1.75-1.45 (m, 2H), 1.35-1.08 (m, 2H). MS (ESI, m/e) [M+1]⁺ 475.2.

Example 7 Polymorph Study of Compound 1

(1) Polymorph Study from Amorphous Form—Preparation of Form a from an Amorphous Form of Compound 1

(S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide (Compound 1) was prepared by the method disclosed in WO2014173289A, and further lyophilized to obtain amorphous form of Compound 1.

For each experiment in Tables 4a to 4k, Tables 5a to 5e and Table 6, about 20 mg of Compound 1 as amorphous form was weighed into a glass vial, followed by the addition of corresponding solvent. The mixture was heated to give a clear solution if needed. Then the mixture was kept at RT without stirring for 1-2 days to see any solid generated from the clear solution. The solid was monitored by Polarized light microscopy.

Table 4 Compound 1 (ee value=90%) as Starting Material

TABLE 4a

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (mL)	Hexane (mL)			
1-1	0.5	0.1	Y	—	No solid
1-2	0.5	0.2	Y	—	Little solid
1-3	0.5	0.3	N	Y	Oil
1-4	0.5	0.4	N	Y	Oil
1-5	1	0.2	Y	—	No solid
1-6	1	0.3	Y	—	No solid
1-7	1	0.4	Y	—	No solid
1-8	1	0.5	Y	—	No solid
1-9	1	0.6	Y	—	Little solid
1-10	1	0.7	Y	—	Little solid
1-11	1	0.8	N	Y	Oil
1-12	1	0.9	N	Y	Oil

TABLE 4b

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (mL)	Heptane (mL)			
2-1	0.5	0.1	Y	—	No solid
2-2	0.5	0.2	Y	—	Little solid
2-3	0.5	0.3	N	Y	Oil
2-4	0.5	0.4	N	N	Oil
2-5	1	0.2	Y	—	No solid
2-6	1	0.3	Y	—	No solid
2-7	1	0.4	Y	—	No solid
2-8	1	0.5	Y	—	No solid
2-9	1	0.6	Y	—	Little solid
2-10	1	0.7	Y	—	Little solid
2-11	1	0.8	Y	—	Oil
2-12	1	0.9	N	Y	Oil

33

TABLE 4c

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (mL)	Cyclohexane (mL)			
3-1	0.5	0.2	Y	—	No solid
3-2	0.5	0.3	Y	—	Little solid
3-3	0.5	0.4	Y	—	Oil
3-4	0.5	0.5	Y	—	Oil
3-5	0.5	0.6	N	Y	Oil
3-6	1	0.6	Y	—	Little solid
3-7	1	0.8	Y	—	Little solid
3-8	1	1.0	Y	—	Little solid
3-9	1	1.2	Y	—	Little solid
3-10	1	1.4	Y	—	Oil

TABLE 4d

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	DCM (mL)	Hexane (mL)			
4-1	0.5	0.4	Y	—	No solid
4-2	0.5	0.6	Y	—	No solid
4-3	0.5	0.8	Y	—	No solid
4-4	0.5	1.0	N	Y	Oil
4-5	1.0	1.4	Y	—	No solid
4-6	1.0	1.6	Y	—	No solid
4-7	1.0	1.8	Y	—	No solid
4-8	1.0	2.0	N	Y	Oil

TABLE 4e

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	1,2-Dichloroethane (mL)	Hexane (mL)			
5-1	0.5	0.6	Y	—	No solid
5-2	0.5	0.8	Y	—	No solid
5-3	0.5	1.0	Y	—	No solid
5-4	0.5	1.1	N	Y	Oil
5-5	1.0	1.4	Y	—	No solid
5-6	1.0	1.6	Y	—	No solid
5-7	1.0	1.8	Y	—	No solid
5-8	1.0	2.0	Y	—	No solid
5-9	1.0	2.2	N	Y	Oil

TABLE 4f

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	MeOAc (mL)	Hexane (mL)			
6-1	0.5	0.3	Y	—	Little solid
6-2	0.5	0.4	Y	—	Oil
6-3	0.5	0.5	Y	—	Oil
6-4	0.5	0.6	N	Y	Oil
6-5	1.0	0.6	Y	—	Little solid
6-6	1.0	0.8	Y	—	Little solid
6-7	1.0	1.0	Y	—	Little solid
6-8	1.0	1.2	Y	—	Little solid
6-9	1.0	1.4	N	Y	Oil

TABLE 4g

Experiment ID	Solvent		RT	heat	Result (1-2 d)
	Toluene (mL)	Hexane (mL)			
7-1	1.0	0.2	Y	—	Little solid
7-2	1.0	0.3	Y	—	Little solid

34

TABLE 4g-continued

Experiment ID	Solvent		RT	heat	Result (1-2 d)
	Toluene (mL)	Hexane (mL)			
7-3	1.0	0.4	Y	—	Little solid
7-4	1.0	0.5	N	Y	Oil
7-5	1.0	0.6	N	Y	Oil

TABLE 4h

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	Toluene (mL)	Cyclohexane (mL)			
8-1	1.0	0.1	Y	—	Little solid
8-2	1.0	0.2	Y	—	Little solid
8-3	1.0	0.3	Y	—	Oil
8-4	1.0	0.4	N	Y	Oil
8-5	1.0	0.5	N	Y	Little solid
8-6	1.5	0.4	Y	—	Little solid
8-7	1.5	0.5	Y	—	Little solid

TABLE 4i

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	MeOAc (mL)	Cyclohexane (mL)			
9-1	0.4	1.0	Y	—	Little solid
9-2	0.5	1.0	Y	—	Little solid
9-3	0.6	1.0	Y	—	Little solid
9-4	0.8	1.0	Y	—	Little solid
9-5	1.0	1.0	Y	—	Little solid

TABLE 4j

Experiment ID	Solvent		RT	Heat	Result (2-3 d)
	IPAC (ml)	Cyclohexane (ml)			
10-1	1.0	0.2	Y	—	Little solid
10-2	1.0	0.4	Y	—	Little solid
10-3	1.0	0.6	Y	—	Little solid
10-4	1.0	0.8	Y	—	Little solid
10-5	1.0	1.0	Y	—	Little solid
10-6	1.0	1.2	N	Y	Oil

TABLE 4k

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	Isobutyl acetate (mL)	Cyclohexane (mL)			
11-1	1.0	0.2	Y	—	Little solid
11-2	1.0	0.4	Y	—	Little solid
11-3	1.0	0.6	Y	—	Little solid

35

TABLE 4k-continued

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	Isobutyl acetate (mL)	Cyclohexane (mL)			
11-4	1.0	0.8	Y	—	Little solid
11-5	1.0	1.0	Y	—	Little solid
11-6	1.0	1.2	N	Y	Oil

Y = Yes, and N = No.

Table 5 Compound 1 (ee value=97%) as Starting Material

TABLE 5a

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (mL)	Hexane (mL)			
12-1	1	0.2	Y	—	No solid
12-2	1	0.3	Y	—	No solid
12-3	1	0.4	Y	—	No solid
12-4	1	0.5	Y	—	No solid
12-5	1	0.6	Y	—	Solid
12-6	1	0.7	Y	—	Solid
12-7	1	0.8	N	Y	Oil
12-8	1	0.9	N	Y	Oil

TABLE 5b

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (mL)	Heptane (mL)			
13-1	1	0.2	Y	—	No solid
13-2	1	0.3	Y	—	No solid
13-3	1	0.4	Y	—	No solid
13-4	1	0.5	Y	—	No solid
13-5	1	0.6	Y	—	Solid
13-6	1	0.7	Y	—	Solid
13-7	1	0.8	Y	—	Oil
13-8	1	0.9	N	Y	Oil

TABLE 5c

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	MeOAc (ml)	Cyclohexane (ml)			
14-1	0.4	1.0	Y	—	No solid
14-2	0.5	1.0	Y	—	No solid
14-3	0.6	1.0	Y	—	Solid
14-4	0.8	1.0	Y	—	Solid
14-5	1.0	1.0	Y	—	No solid
14-6	1.0	1.5	Y	—	Solid
14-7	1.0	2.0	Y	—	Solid
14-8	1.0	2.2	N	Y	Oil

TABLE 5d

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (ml)	Cyclohexane (ml)			
15-1	0.5	0.2	Y	—	No solid
15-2	0.5	0.3	Y	—	solid
15-3	0.5	0.4	Y	—	Oil
15-4	0.5	0.5	Y	—	Oil

36

TABLE 5d-continued

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	EA (ml)	Cyclohexane (ml)			
5					
10					
15					
15-5	0.5	0.6	N	Y	Oil
15-6	1	0.6	Y	—	solid
15-7	1	0.8	Y	—	solid
15-8	1	1.0	Y	—	solid
15-9	1	1.2	Y	—	solid
15-10	1	1.4	Y	—	Oil

TABLE 5e

Experiment ID	Solvent		RT	Heat	Result (1-2 d)
	MeOAc (ml)	Hexane (ml)			
20					
25					
30					
35					
40					
45					
50					
55					
16-1	0.5	0.3	Y	—	solid
16-2	0.5	0.4	Y	—	No solid
16-3	0.5	0.5	Y	—	No solid
16-4	0.5	0.6	N	Y	No solid
16-5	1.0	0.6	Y	—	solid
16-6	1.0	0.8	Y	—	solid
16-7	1.0	1.0	Y	—	solid
16-8	1.0	1.2	Y	—	solid
16-9	1.0	1.4	N	Y	Oil

Y = Yes, and N = No.

Experiments in Tables 4a to 4k and Tables 5a to 5e were conducted on the same scale (i.e., the amount of the starting material—amorphous Compound 1 is about 20 mg). However, the ee value of the starting material appeared to have significant influence in the amount of the solid to be formed in each experiment. As shown in the experiments in Tables 4a to 4k starting from 90% ee of amorphous Compound 1, the solids thus formed are of little amount. The experiments in Tables 5a to 5e starting from 97% ee of amorphous Compound 1 resulted in noticeable amount of solid. Also, the obtained solid in the experiments in Tables 4a to 4k showed low ee value when crystallization from starting material with 90% ee. One solid sample from Tables 4a to 4k using EA/Hexane as crystallization system only showed 45% ee value.

The results of Table 5a were further confirmed by a scale-up experiment similar to those in Example 2, which confirmed that the resultant solid was in the desired crystalline form (Form A).

As shown in the above Tables 4a to 4k and Tables 5a to 5e, the formation of the crystalline solid may vary depending on the specific solvents, the ratio of the solvents, and so on.

The results in Table 6 further confirms that the formation of the crystalline solid depends on the specific ratio of the solvent.

TABLE 6

Solvent-1 (0.5 mL)	Solvent-2											
	Hexane		MTBE		Heptane		Cyclohexane		H ₂ O		Ether	
	V/mL	Results	V/mL	Results	V/mL	Results	V/mL	Results	V/mL	Results	V/mL	Results
THF	0.4	Little solid	2.0	Little solid	0.4	Little solid	0.4	Little solid	—	—	2.0	No solid
Me—THF	0.4	Little solid	1.5	No solid	0.4	No solid	—	—	—	—	—	—
DME	0.6	Little solid	1.5	No solid	0.6	Little solid	0.8	Little solid	—	—	1.5	No solid
EtOH	2.0	No solid	5.0	No solid	2.5	oil	2.0	No solid	5.0	No solid	5.0	No solid
i-PrOH	1.5	No solid	2.5	No solid	—	—	2.0	No solid	2.0	Little solid	—	—
pyridine	1.0	No solid	4.0	No solid	—	—	1.0	No solid	2.0	No solid	—	—
Propyl acetate	0.3	No solid	0.8	No solid	—	—	—	—	—	—	—	—
Isobutyl acetate	0.2	No solid	2.0	No solid	—	—	—	—	—	—	—	—
n-BuOH	0.5	No solid	2.0	No solid	—	—	—	—	0.5	No solid	—	—
1,2-Dichloroethane	1.0	No solid	5.0	No solid	—	—	1.2	No solid	—	—	—	—
DCM	1.0	oil	5.0	No solid	—	—	1.2	No solid	—	—	—	—
Toluene	0.2	Oil and Little solid	0.5	No solid	0.2	No solid	0.2	Oil and Little solid	—	—	0.5	No solid

(2) Polymorph Study from Crystalline Form—Preparation of Form a from Crystalline Form Slow Evaporation ²⁵

About 15 mg of sample (Crystalline Form A) was weighed into a 3-mL glass vial, followed by the addition of corresponding solvent or solvent mixture (see Table 7) to get a clear solution. Subsequently, the vial was covered with parafilm with 3-4 pinholes, and kept at RT to allow the solution to evaporate slowly. The solids were isolated for XRPD analysis. However, no crystal form was produced, as summarized in Table 7.

TABLE 7

Slow Evaporation Experiments		
Experiment ID	Solvent (v/v)	Solid Form
1	MeOH	Amorphous
2	EtOH	Amorphous
3	IPA	Amorphous
4	ACN	Amorphous
5	Acetone	Oil
6	EtOAc	Oil
7	THF	Oil
8	DCM	Amorphous
9	Toluene	Oil
10	Acetic acid	Oil
11	EtOH/H ₂ O (4:1)	Amorphous
12	Acetone/H ₂ O (4:1)	Amorphous
13	THF/H ₂ O (4:1)	Amorphous
14	DCM/n-heptane (4:1)	Amorphous
15	EtOH/n-heptane (4:1)	Amorphous
16	EtOAc/n-heptane (6.5:1)	Oil
17	ACN/MTBE (4:1)	Oil

Anti-Solvent Addition

About 15 mg of sample (Crystalline Form A) was weighed into a 20-mL glass vial, followed by the addition of 0.4-1.2 mL corresponding solvent (see Table 8). The mixture was then magnetically stirred at the speed of 800 rpm to get a clear solution at RT. Subsequently, the relative anti-solvent (see Table 8) was added to the solution to induce precipitation or until the total amount of anti-solvent reached 15.0 mL. If no precipitation occurs, the solution was then transferred to slow evaporation at RT. Results summarized in Table 8.

TABLE 8

Anti-solvent Addition Experiments			
Experiment ID	Anti-Solvent	Solvent	Solid Form
1	H ₂ O	EtOH	Oil
2	H ₂ O	THF	Oil
3	H ₂ O	Acetic acid	Oil
4	n-heptane	1,4-dioxane	Oil
5	MTBE	ACN	Oil
6	MTBE	NMP	N/A
7	MTBE	EtOH	Oil
8	MTBE	DCM	Oil

N/A: no solid was obtained.

Slow Cooling

About 20 mg of sample (Crystalline Form A) was suspended in 1.0 mL of corresponding solvent (see Table 9) in a 3-mL glass vial at RT. The suspension was transferred to slurry at 50° C. using magnetic stirring with the speed of 800 rpm. The sample was equilibrated at 50° C. for 2 hrs and filtered using a 0.45 μm Nylon membrane. Subsequently, the filtrate was slowly cooled down from 50° C. to 5° C. at a rate of 0.1° C./min. The obtained solids were kept isothermal at 5° C. before isolated for XRPD analysis. No crystal form was obtained, as summarized in Table 9.

TABLE 9

Slow Cooling Experiments		
Experiment ID	Solvent (v/v)	Solid Form
1	IPA	N/A
2	MIBK	N/A
3	IPAc	N/A
4	Toluene	N/A
5	EtOH/H ₂ O (1:2)	Gel
6	Acetone/H ₂ O (1:2)	Gel
7	EtOAc/n-heptane (1:2)	Amorphous
8	CHCl ₃ /n-heptane (1:2)	N/A
9	THF/n-heptane (1:2)	Oil*
10	ACN/MTBE (1:2)	Oil*

N/A: no solid was obtained.

*clear solution was transferred to evaporate at RT.

Solution Vapor Diffusion

About 15 mg of sample (Crystalline Form A) was dissolved in 0.5-1.5 mL of corresponding solvent (see Table 10) to obtain a clear solution in a 3-mL vial. Subsequently, the solution was placed into a 20-mL vial with 3 mL of relative anti-solvent. The 20-mL vial was sealed with a cap and kept at RT, allowing sufficient time for organic vapor to interact with the solution. At the end of 11 days, clear solutions were transferred to evaporation at RT. The solids obtained were characterized by XRPD. The results summarized in Table 10.

TABLE 10

Solution Vapor Diffusion Experiments			
Experiment ID	Solvent	Anti-solvent	Solid Form
1	EtOH	H ₂ O	Amorphous*
2	ACN	H ₂ O	N/A
3	Acetone	H ₂ O	Amorphous*
4	THF	H ₂ O	Amorphous*
5	Acetic acid	H ₂ O	Oil
6	EtOH	n-heptane	N/A
7	THF	n-heptane	Amorphous*
8	DCM	n-heptane	Amorphous*
9	DMAc	MTBE	N/A
10	IPA	MTBE	N/A
11	1,4-Dioxane	MTBE	N/A
12	Toluene	MTBE	N/A
12	NMP	MTBE	N/A

N/A: no solid was obtained.

*solid was generated via slow evaporation.

Polymer-Induced Crystallization Experiments

About 15 mg of sample (Crystalline Form A) was dissolved in 1.0 mL of corresponding solvent (see Table 11) to obtain a clear solution in a 3-mL vial. The solution was then

filtered using 0.45 μ m Nylon membrane. About 2 mg of polymer mixture was added into the filtrate. The mixture was stirred at RT to induce precipitation. The solids were isolated for XRPD analysis. No crystal form was obtained, as summarized in Table 11.

TABLE 11

Polymer-induced crystallization Experiments			
Experiment ID	Solvent (v/v)	Polymer Mixture	Solid Form
1	EtOH	A	Amorphous
2	ACN	A	Amorphous
3	Acetone	A	Amorphous
4	THF	A	Amorphous
5	DCM	A	Amorphous
6	EtOAc	A	Amorphous
7	EtOH	B	Amorphous
8	ACN	B	Amorphous
9	Acetone	B	Amorphous
10	THF	B	Amorphous
11	DCM	B	Amorphous
12	EtOAc	B	Amorphous

Polymer mixture A: polyvinyl pyrrolidone (PVP), polyvinyl alcohol (PVA), polyvinyl chloride (PVC), polyvinyl acetate (PVAc), hypromellose (HPMC), methyl cellulose (MC) (mass ratio of 1:1:1:1:1:1). Polymer mixture B: polycaprolactone (PCL), polyethylene glycol (PEG), poly(methyl methacrylate) (PMMA) sodium alginate (SA), and hydroxyethyl cellulose (HEC) (mass ratio of 1:1:1:1:1).

Example 8 Determination of Absolute Configuration of Compound 1

Preparation of BG-13 Single Crystal

Six single crystal growth experiments (see Table 12) were performed via slow cooling. Suitable single crystals of BG-13 were obtained by slow cooling in MeOH/H₂O (1:1, v/v). The crystal data and structure refinement are listed in Table 13.

TABLE 12

single Crystal Growth Experiments						
Experiment ID	Weight (mg)	Solvent (1 mL, v/v)	Method	Temperature (° C.)	Dissolved (Y/N*)	Observation
1	5.6	IPA/H ₂ O (3/1)	cooling	60	N	Block-like crystal
2	5.5	IPA/H ₂ O (3/1)	cooling	60	N	Block-like crystal
3	5.4	IPA/H ₂ O (3/1)	cooling	60	N	Block-like crystal
4	5.5	IPA/H ₂ O (3/1)	cooling	60	N	Block-like crystal
5	4.7	MeOH/H ₂ O (2/1)	cooling	60	N	Crystal
6	5.5	MeOH/H ₂ O (1/1)	cooling	60	N	Crystal

TABLE 13

Single Crystal Data and Structure Refinement of BG-13
The data of single crystal were generated on a Bruker APEX DUO single-crystal diffractometer with CCD detector (Cu K α , λ = 1.54178 Å, 173.15K).

Empirical formula	C ₃₃ H ₃₄ N ₅ O ₆	—
Formula weight	596.65	—
Temperature	173.15	—
Wavelength	1.54178 Å	—
Crystal system, space group	monoclinic	C2
Unit cell dimensions	a = 16.7939(4) Å b = 7.9871(2) Å c = 23.5438(5) Å	alpha = 90.00 deg. beta = 108.0460(10) deg. gamma = 90.00 deg.
Volume	3002.69(12) Å ³	—
Z, Calculated density	4	1.320 mg/mm ³
Absorption coefficient	0.756 mm ⁻¹	—
F(000)	1260.0	—
Crystal size	0.3 × 0.21 × 0.08 mm ³	—
Theta range for data collection	1.97 to 64.96 deg.	—

TABLE 13-continued


Single Crystal Data and Structure Refinement of BG-13		
The data of single crystal were generated on a Bruker APEX DUO single-crystal diffractometer with CCD detector (Cu K α , λ = 1.54178 Å, 173.15K).		
Limiting indices	-19 \leq h \leq 17, -7 \leq k \leq 9, -27 \leq l \leq 24	—
Reflections collected/unique	5073/3756[R(int) = 0.1062]	—
Completeness	92.8%	—
Refinement method	Full matrix least squares on F ²	—
Data/restraints/parameters	3756/1/398	—
Goodness-of-fit on F ²	1.192	—
Final R indices [I > 2 σ (I)]	R ₁ = 0.0819	wR ₂ = 0.2294
Absolute structure Flack	0.0(3)	—
Largest diff. peak and hole	0.50 and -0.57 e · Å ⁻³	—

BG-13 was confirmed to be a (2R, 3R)-dibenzoyl tartaric acid (L-DBTA) salt and the molar ratio of freebase to L-DBTA is 2:1. Configuration of both carbons (C32 and C32') in L-DBTA was confirmed to be R. Configuration of C6 in freebase was determined to be S, as shown in FIG. 8 to FIG. 10. A powder X-ray diffraction pattern method was also used to characterize the structure of the single crystals, as shown in FIG. 11.


Absolute Configuration of Compound 1

The absolute configurations of Compound 1 was deduced to be S from the single crystal X-ray structural analysis of intermediate BG-13.

Example 9 Chiral Resolution of BG-11A



-continued



General procedure: To a solution of compound BG-11A in a prepared solvent system was added a chiral acid at elevated temperature. After stirring at this temperature, it was cooled to RT and stirred overnight at RT. The solid was filtered and washed with the prepared solvent system. The ee value was tested by chiral HPLC directly from the related salt or its Boc-derivative (see Table 14). Other chiral acids or solvent system gave no ee value, low ee value or not desired chiral compound.


TABLE 14

Chiral Resolution of BG-11A					
Solvent System	Amount of 11A	Chiral acid	temperature	ee value	Yield
EtOH/H ₂ O/AcOH 7/3/1 (1.9 L)	40.0 g	D-DBTA (0.5 eq.)	70° C. to RT	>85% ee	42.2%
i-PrOH/H ₂ O/AcOH 7/3/1 (25 mL)	500 mg	D-DBTA (1.0 eq.)	70° C. to RT	77% ee	38.5%
i-PrOH/H ₂ O/AcOH 7/3/1 (25 mL)	500 mg	D-DBTA (0.5 eq.)	70° C. to RT	85% ee	38.9%
EtOH/H ₂ O/AcOH 7/3/1 (25 mL)	500 mg	D-DBTA (0.5 eq.)	70° C. to RT	86% ee	39.8%
MeOH/H ₂ O/AcOH 7/3/1 (25 mL)	500 mg	D-DBTA (0.5 eq.)	70° C. to RT	82% ee	42.2%
AcOH/H ₂ O 3/1 (40 mL)	1 g	D-DBTA (0.55 eq.)	60° C. to RT	83% ee	27.6%

TABLE 14-continued

Chiral Resolution of BG-11A					
Solvent System	Amount of 11A	Chiral acid	temperature	ee value	Yield
1,4-dioxane/H ₂ O 1/1 (2.5 mL)	25 mg	D-DBTA (2.0 eq.)	60° C. to RT	No Solid	No Solid
MeOH/H ₂ O 1/1 (2.5 mL)	25 mg	D-DBTA (2.0 eq.)	60° C. to RT	36% ee	Not weigh
CH ₃ CN/H ₂ O 9/1 (2.5 mL)	25 mg	D-DBTA (2.0 eq.)	60° C. to RT	14% ee	Not weigh
CH ₃ CN/H ₂ O 6/1 (2.5 mL)	25 mg	D-DBTA (2.0 eq.)	60° C. to RT	89% ee	Not weigh
i-PrOH/H ₂ O 1/1 (2.5 mL)	25 mg	D-DBTA (2.0 eq.)	60° C. to RT	79% ee	Not weigh
CH ₃ CN/H ₂ O 4/1 (1 mL)	50 mg	D-DBTA (1.0 eq.)	60° C. to RT	24% ee	46%
CH ₃ CN/H ₂ O 4/1 (1.5 mL)	50 mg	D-DBTA (1.0 eq.)	60° C. to RT	91% ee	33.7%

Example 10 Chiral Resolution of BG-12A and
Improve the Chiral Purity



Chiral acid
→

20


25

30

35

40

-continued



General procedure: To a solution of compound BG-12A in a prepared solvent system was added chiral acid in at elevated temperature. After stirring at this temperature, it was cooled to RT and stirred overnight at RT. The solid was filtered and washed with the prepared solvent system. The chiral purity was tested by chiral HPLC directly from the related salt or free base (see Table 15). Other chiral acids or solvent system gave no ee value, low ee value or not desired chiral compound.

TABLE 15

Chiral Resolution of BG-12A					
Solvent System	Amount of BG-12A	Chiral acid	Temperature	ee value	Yield
MeOH/H ₂ O 3/1 (1500 mL)	50 g	L-DBTA (0.35 eq.)	50° C. to RT	85.6% ee	43.1%
EtOH/H ₂ O 6/1 (250 mL)	14.4 g	L-DBTA (0.55 eq.)	78° C. to RT	79.1% ee	41.8%
n-BuOH/H ₂ O 6/1 (20 mL)	1 g	L-DBTA (0.8 eq.)	80° C. to RT	95% ee	20%
MeOH (4 mL)	500 mg	L-DBTA (1.1 eq.)	Reflux to RT	No solid	
EtOH (17 mL)	1.0 g	L-DBTA (1.1 eq.)	Reflux to RT	40% ee	Not weigh
EtOH (30 mL)	500 mg	L-DBTA (2.2 eq.)	Reflux to RT	No ee	Not weigh

The obtained L-DBTA salt (31 g, 85.6% ee) was added to THF/H₂O (1/1, 1034 mL), the suspension was warmed to 70° C. and stirred until all solid dissolved. Then 517 mL of water was added. The solution was then slowly cooled to 40° C. and added seed crystal (10 mg). After stirring for about 2 hrs, the solution was slowly cooled to ambient temperature and stirred for 2 days. Filtered, the solid was washed with THF/H₂O=1/1 (20 mL) and dried over under reduced pressure to give the product as a white solid (22.5 g, 72% yield, >98.5 ee value).

The obtained free base (6.02 g, 79.1% ee) was dissolved in (1 g/15 mL) EtOH/H₂O (6/1, 90 mL), stirred at 78° C. allowing all the starting material to be dissolved. Then, a solution of L-DBTA (2.84 g, 7.9 mmol, 0.55 eq) in EtOH/H₂O (6/1, 7 mL) was added. The solid quickly formed, the mixture was stirred at this temperature for 1 h before removing the heating system. The mixture was allowed to cool to RT. Filtered, the solid was washed with EtOH/H₂O (6/1, 10 mL). The collected solid was converted to free base using in NaOH aqueous solution and DCM to get the product (4.7 g, yield: 32.6%, 93% ee) as a white foam.

A suspension of the obtained free base (70.0 g, 90.5% ee) in CH₃CN/H₂O (1/1, 700 mL) was heated to 60° C. to give a clear solution. To the above solution was then added L-DBTA (33 g, 0.55 eq). After stirring at 60° C. for about 2 hr, the mixture was slowly cooled to RT and stirred for overnight. Filtered, the solid was washed with CH₃CN/H₂O (1/1, 50 mL), dried over under reduced pressure to give the product as a off-white solid (80 g, yield: 80% ee value >98%).

Example 11 Efficacy Tests

(S)-7-(1-acryloylpiperidin-4-yl)-2-(4-phenoxyphenyl)-4,5,6,7-tetrahydropyrazolo[1,5-a]pyrimidine-3-carboxamide was tested hereinafter by using its Crystalline Form A.

Test 1: Inhibition and Selectivity of the Kinases
Methods:

(1) BTK Kinase Enzymatic Assays

Crystalline Form A of Compound 1 was tested for inhibition of BTK kinase (aa2-659, Carna Biosciences) in assays based on the time-resolved fluorescence-resonance energy transfer (TR-FRET) methodology. The assays were carried out in 384-well low volume black plates in a reaction mixture containing BTK kinase, 5 μM ATP, 2 μM peptide substrate and 0-10 μM compound in buffer containing 50 mM Tris pH7.4, 10 mM MgCl₂, 2 mM MnCl₂, 0.1 mM EDTA, 1 mM DTT, 0.005% Tween-20, 20 nM SEB and 0.01% BSA. The kinase was incubated with compound for 60 minutes at room temperature and the reaction was initiated by the addition of ATP and peptide substrate. After reaction at room temperature for 60 minutes, an equal volume of stop/detection solution was added according to the manufacture's instruction (CisBio Bioassays). The stop/detection solution contained Eu³⁺cryptate-conjugated mouse monoclonal antibody (PT66) anti-phosphotyrosine and XL665-conjugated streptavidin in buffer containing 50 mM HEPES pH7.0, 800 mM KF, 20 mM EDTA, and 0.1% BSA. Plates were sealed and incubated at room temperature for 1 hour, and the TR-FRET signals (ratio of fluorescence emission at 665 nm over emission at 620 nm with excitation at 337 nm wavelength) were recorded on a PHERAstar FS plate reader (BMG Labtech). Phosphorylation of peptide substrate led to the binding of anti-phosphotyrosine antibody to the biotinylated peptide substrate, which places fluorescent donor (Eu³⁺cryptate) in close proximity to the acceptor (Streptavidin-XL665), thus resulting in a high

degree of fluorescence resonance energy transfer from the donor fluorophore (at 620 nm) to the acceptor fluorophore (at 665 nm). Inhibition of BTK kinase activity resulted in decrease of the TR-FRET signal. The IC₅₀ for Compound 1 was derived from fitting the data to the four-parameter logistic equation by Graphpad Prism software.

(2) Biochemical Kinase Selectivity

Selectivity of Crystalline Form A was profiled against a panel of 342 kinases at 1 μM at Reaction Biology Corp. Crystalline Form A displayed less than 70% inhibition against 329 kinases, and greater than 70% inhibition against 13 kinases including BTK. IC₅₀ of Crystalline Form A (see Table 13), including ITK, TEC, JAK3 and EGFR assays carried out in-house at BeiGene by using a TR-FRET assay and corresponding peptides as the substrate.

IC₅₀ determination of ITK: The protocol of ITK assay is similar to BTK assay except for the following modification: 3 μM ATP and 2 μM TK substrate were used in the kinase reaction.

IC₅₀ determination of TEC: The protocol of Tec assay is similar to BTK assay except for the following modifications: 1) 280 μM ATP and 2 nM Poly-GT substrate were used in the kinase reaction; 2) the reaction buffer doesn't contain SEB.

IC₅₀ determination of JAK3: The protocol of JAK3 assay is similar to BTK assay except for the following modifications: 1) 3.4 μM ATP and 3 μM peptide substrate (B-EE-15, Biotin-EQEDEPEGDYFEWLE) were used in the kinase reaction; 2) the reaction buffer contains 50 mM Tris pH7.8, 10 mM MgCl₂, 5 mM DTT, 0.01% Triton X-100 and 0.01% BSA.

IC₅₀ determination of EGFR: The protocol of EGFR assay is similar to BTK assay except for the following modifications: 1) 20 μM ATP, 1.44 μM TK substrate-biotin (one universal substrate for tyrosine kinases) and 0-1000 nM compound (the final concentration of 1% DMSO) were used in the kinase reaction; 2) the reaction buffer contains 50 mM HEPES pH7.5, 10 mM MgCl₂, 1 mM EGTA, 0.01% Brij-35, 2.5 mM DTT and 0.1% BSA; 3) the stop/detection solution buffer contains 25 mM HEPES pH7.5, 400 mM KF, 50 mM EDTA, 0.01% Triton-X100 and 0.1% BSA.

Results

IC₅₀ of Crystalline Form A for BTK kinase was 0.27 nM. Crystalline Form A was shown to be a potent, specific and irreversible BTK kinase inhibitor. In terms of its selectivity, Crystalline Form A inhibited only 13 other kinases more than 70% when profiled against a panel of 342 human kinases at 1 μM.

TABLE 16

Enzymatic Inhibition Activities of Crystalline Form A	
Enzyme	IC ₅₀ (nM)
BTK	0.27
ITK	53
TEC	1.9
JAK3	600
EGFR	3.6
BLK	1.13
BMX/ETK	0.62
BRK	33
ERBB4/HER4	1.58
FGR	155

TABLE 16-continued

Enzymatic Inhibition Activities of Crystalline Form A	
Enzyme	IC ₅₀ (nM)
FRK/PTK5	379
LCK	187
TXK	2.95

Note:

BTK, EGFR, ITK, TEC and JAK3 assays were carried out by using a TR-FRET assay and corresponding peptides as substrate. IC₅₀s of Crystalline Form A were measured at K_M of ATP for the five kinases and with 1-hour pre-incubation. HER4, BMX, TXK, BLK, FGR, LCK, FRK/PTK5 assays were carried out at Reaction Biology Corp. using ³³P-ATP and filter-binding assay. IC₅₀s of Crystalline Form A were measured at 1 μM ATP and with 1-hour pre-incubation.

Test 2: BTKpY223 Cellular Assay by Crystalline Form a Methods:

BTKpY223 cellular assay is a HTRF based assay intended to quantitatively determine the endogenous levels of phosphorylation at BTK Tyr223. Phosphorylated Tyr223 is necessary for full activation of BTK. The assay was performed in Ramos cells (CRL-1596, ATCC) with a BTKpY223 assay kit (63IDC000, Cisbio).

Briefly, Ramos cells were serum starved in 0.5% FBS-containing RPMI1640 for 2 hours. Following starvation, the cells were incubated with Crystalline Form A to be detected at various concentrations in a CO₂ incubator for 1 hour. After incubation, cells were stimulated with 1 mM pervanadate (PV) or Na₃VO₄ (OV) for 20 min. Then, the cells were spun down and lysed with 1× lysis buffer at RT for 10 min (4× lysis buffer supplied in the kit). During incubation, 1× antibody mix was prepared by diluting anti-BTK-d2 and anti-pBTK-K in detection buffer (supplied in the kit). 2 ul/well of 1× antibody mixture was dispensed into the Opti-Plate-384 assay plate (6005620, Perkin Elmer). After that, 18 μL of cell lysate was transferred to the assay plate pre-loaded with antibody solution. After mixing gently and spinning briefly, the plate was sealed up and kept in dark at RT for 18 hours. The fluorescence emission was measured at two different wavelengths (665 nm and 620 nm) on a compatible HTRF reader (PHERAstar FS, BMG). The potency of Compound 1 was calculated basing on the inhibition of ratio between signal intensities at 665 nm and 620 nm. IC₅₀ values were calculated with GraphPad Prism software using the sigmoidal doseresponse function.

Results:

Crystalline Form A inhibited the phosphorylation of BTK in the B cell lymphoma cell line, Ramos, at concentration as low as 1.8±0.2 nM (n=3).

Test 3: Effects of Crystalline Form a on Tumor Cell Proliferation in Haematological Cancer Lines (Rec-1, Mino, JEKO-1 and TMD-8)

Methods:

3 MCL cell lines (Rec-1, Mino and JEKO-1) and an ABC type diffuse large B-cell lymphoma cell line (TMD8) were used in this study. Cell lines were maintained in RPMI-1640 supplemented with 10% fetal bovine serum/FBS (Thermo Scientific); 100 units/ml penicillin (Gibco) and 0.1 mg/ml streptomycin (Gibco) and kept at 37° C. in a humidified atmosphere of 5% CO₂ in air. Cell lines were reinstated from frozen stocks that were laid down within 30 passages from the original cells purchased.

The growth-inhibitory activity of compounds in Rec-1, Mino, JEKO-1 and TMD-8 cells was determined using CellTiter-Glo luminescent cell viability assay (Promega). The number of cells seeded per well of a 96-well plate was optimized for each cell line to ensure logarithmic growth

over 6 days treatment period. Cells were treated in triplicate with a 10-point dilution series. Following a 6-day exposure to the compound, a volume of CellTiter-Glo reagent equal to the volume of cell culture medium present in each well was added. Mixture was mixed on an orbital shaker for 2 minutes to allow cell lysis, followed by 10 minutes incubation at room temperature to allow development and stabilization of luminescent signal, which corresponded to quantity of ATP and thus the quantity of metabolically active cells. Luminescent signal was measured using PHERAstar FS reader (BMG Labtech). IC₅₀ values for cell viability were determined with GraphPad Prism software and were the mean of 3 independent assays.

Results:

Crystalline Form A of Compound 1 exhibited specific and potent inhibitory effect on cellular proliferation in 3 MCL cell lines and an ABC type diffuse large B-cell lymphoma cell line (TMD8) (Table 17).

TABLE 17

Inhibition of Crystalline Form A on hematic tumor cell proliferation			
Cell line	Cell Type	Potency IC ₅₀ (nM)	Standard deviation(nM)
Rec-1	MCL	0.36	0.03
Mino	MCL	3.8	1.8
JEKO-1	MCL	20.0	NA
TMD-8	DLBCL(ABC)	0.54	0.3

Test 4: Pharmacokinetics Study of Crystalline Form a in Mouse

Methods:

For time course study, mice were randomly assigned into 7 groups with 4 mice per group. Mice were treated with single dose of Crystalline Form A of Compound 1 and euthanized using carbon dioxide at different time points (30 minutes, 1, 2, 4, 12, 24 hrs) after dosing. For dose dependency study, mice were randomly assigned into 9 groups with 4 mice per group. Mice were treated with different dose levels of Crystalline Form A of Compound 1 and euthanized using carbon dioxide at 4 hrs after dosing. Treatments were administered by oral gavage (p.o.) in a volume of 10 ml/kg body weight. Body weight was assessed immediately before dosing and volume dosed was adjusted accordingly.

PK SAMPLE PREPARATION: For time course study, blood samples (50 μL per mouse) were collected from the retro-orbital sinus under isoflurane/oxygen anesthesia at 15 min after dosing (this group of mice were also used for 24 hr time point) or heart puncture after euthanization for the other time points. For dose dependency study, blood samples were collected from the retro-orbital sinus under isoflurane/oxygen anesthesia at 30 minutes after dosing. Plasma was collected by centrifugation at 3,000 g for 10 minutes and was kept frozen in -80° C. until analysis.

PK Analysis: maximum plasma concentration (C_{max}) and time to reach C_{max} (T_{max}) were taken directly from the plasma concentration versus time profiles.

Results:

Crystalline Form A was quickly absorbed and eliminated in ICR mice.

Test 5: Efficacy Study of Crystalline Form a for in TMD-8 Xenograft Model

Tumor Implantation Methods:

Animals were pre-treated with cyclophosphamide (prepared in saline, 150 mg/kg i.p.) and disulfiram (prepared in

0.8% Tween 80 in saline, 125 mg/kg p.o., one hour after each dose of cyclophosphamide) once daily for two days. Animals were then inoculated with TMD-8 cells 24 hours after the second dose of cyclophosphamide. On the day of implantation, cell culture medium was replaced with fresh medium. Four hours later, media was removed and cells were collected as described above. Cells were re-suspended in cold (4° C.) PBS and same volume of matrigel (BD, Cat #356237) was added to give a final concentration of 2.5×10^7 cells/ml. Resuspended cells were placed on ice prior to inoculation. The right axilla region of each mouse was cleaned with 75% ethanol prior to cell inoculation. Each animal was injected subcutaneously with 5×10^6 cells in 200 μ l of cell suspension in the right front flank via a 26-gauge needle.

For in vivo efficacy studies, starting from day 3 after cell inoculation, animals were randomly assigned into desired number of groups with 10 mice per group. Mice were treated twice daily (BID) with vehicle (0.5% carboxymethylcellulose (CMC)+0.2% Tween 80), and different dose levels of Crystalline Form A of Compound 1 for 39 days. Treatments were administered by oral gavage (p.o.) in a volume of 10 ml/kg body weight. Body weight was assessed immediately before dosing and volume dosed was adjusted accordingly. Tumor volume was measured twice weekly in two dimensions using a calliper (measurable from day 11 post inoculation in this study). Tumor volume was calculated using the formula: $V=0.5 \times (a \times b^2)$ where a and b are the long and short diameters of the tumor, respectively. Statistical analysis was conducted using the student T-test. $P < 0.05$ was considered statistically significant. One individual was responsible for tumor measurement for the entire duration of the study. Body weights were also recorded twice weekly. Mice were also being monitored daily for clinical signs of toxicity for the duration of the study.

Results:

In vivo efficacy of Crystalline Form A was examined in TMD-8 DLBCL xenografts grown subcutaneously in NOD/SCID mice. Following daily oral administration at well tolerated at different dose levels twice daily (BID), Crystalline Form A of Compound 1 induced dose-dependent anti-tumor effects. Crystalline Form A of Compound 1 at lowest dose tested already showed strong anti-tumor activity. All treatment groups had no significant impact on animal body weight throughout the study.

Test 6: Efficacy Study of Crystalline Form a in Systemic REC-1 Xenograft Model

Tumor Implantation Methods:

Animals were pre-treated with cyclophosphamide (prepared in saline, 150 mpk i.p.) and disulfiram (prepared in 0.8% TW-80 in saline, 125 mpk p.o., one hour after each dose of cyclophosphamide) once daily for two days. Animals were then inoculated with REC-1 cells 24 hours after the second dose of cyclophosphamide. On the day of implantation, cell culture medium was replaced with fresh medium. Four hours later, media was removed and cells were collected as described above. Cells were re-suspended in cold (4° C.) PBS to give a final concentration of 1×10^8 cells/ml. Resuspended cells were placed on ice prior to implantation. Each animal was injected intravenously via tail vein with 1×10^7 cells in 100 μ l of cell suspension.

For in vivo efficacy studies, starting from day 8 after cell inoculation, animals were randomly assigned into desired number of groups with 10 mice per group. Mice were treated either twice daily (BID) with vehicle (0.5% carboxymethylcellulose (CMC)+0.2% Tween 80), different dose levels of Crystalline Form A of Compound 1 for 71 days. All dosing

was stopped on day 78 after inoculation. Treatments were administered by oral gavage (p.o.) in a volume of 10 ml/kg body weight. Body weight was assessed immediately before dosing and volume dosed was adjusted accordingly. Body weight was recorded twice weekly (changed to three times per week from day 33). Mice were also watched daily for clinical signs of sickness for the duration of the study. The endpoint of the study is overall survival. In the case of severe toxic effect, such as loss of movement, mice were euthanized and scored as death.

For Data Analysis: Survival analysis was performed by Kaplan-Meier method. The survival time was defined as the time from the day of tumor cell inoculation to the date of animal death or being euthanized. For each group, median survival time (MST), range of survival time (RST) with 95% confidence interval and increase in life-span (ILS) were calculated. Median survival is defined as the time when 50% of mice have died. ILS was calculated using the following formula:

$$\% ILS = (MST - MST_{(vehicle)}) / MST_{(vehicle)} \times 100$$

Statistical Analysis was Conducted Between Each Group Using Gehan-Breslow-Wilcoxon Test. $P < 0.05$ was Considered as Statistically Significant.

Results:

Crystalline Form A of Compound 1 demonstrated dose-dependent anti-tumor activity against systemic REC-1 MCL engrafts in NOD/SCID mice. Crystalline Form A of Compound 1 was significantly effective in this xenograft model.

Test 7: Toxicology of Crystalline Form a

A comprehensive nonclinical toxicity study program, including 28-day GLP studies in rats and dogs and several investigational studies, was conducted for the evaluation of the preclinical safety of Crystalline Form A of Compound 1 at different doses. These studies took account the available regulatory guidance for preclinical development of anticancer drugs. In these studies, Compound 1 demonstrated a favorable toxicology and safety pharmacology profile. No test article-related mortality occurred at any dose levels throughout the study. No toxicologically significant changes in clinical chemistry or coagulation were noted throughout the study. None of these changes were noted after the recovery phase.

Test 8: Pharmacokinetics of Crystalline Form a

The fully-validated LC-MS/MS method was well used for the pharmacokinetic (PK) studies of Crystalline Form A of Compound 1 in Sprague-Dawley rats and beagle dogs following single- and multiple-dose administrations.

Crystalline Form A of Compound 1 has good oral bioavailability in rats. It was quickly absorbed and exhibited high plasma clearance (CL) in rats. The kinetics was linear over the dose range in female rats. The linearity in male rats was not as good. There was no statistically significant accumulation of Compound 1 following multiple oral dosing in both male and female rats. Crystalline Form A of Compound 1 exhibited moderate clearance (CL), reasonably good bioavailability (F %), linear PK over the dose range and no accumulation of Compound 1 following multiple oral dosing in dogs.

Test 9: ADME of Crystalline Form a

Compound 1 was widely distributed to various tissues, but was low in brain tissue, indicating the drug does not easily cross the blood-brain barrier.

IC_{50} values of Crystalline Form A of Compound 1 for seven major drug metabolizing CYP isozymes (CYP1A2, CYP2B6, CYP2C8, CYP2C9, CYP2C19, CYP2D6, and CYP3A) were determined in human liver microsomes and

the time-dependent inhibition potential on major CYP isozymes of Compound 1 was also evaluated. It showed weak inhibition on CYP2C8 ($IC_{50}=4.03 \mu M$), CYP2C9 ($IC_{50}=5.69 \mu M$) and CYP2C19 ($IC_{50}=7.58 \mu M$), but lower inhibition on other CYP isozymes. Compound 1 is unlikely to be time dependent CYP inhibitors on these 7 major human CYPs. CYP3A is the major CYP isoform responsible for the metabolism in human liver microsomes.

Example 12 Clinical Trail Study

(1) Ongoing Clinical Trial Phase 1 Result on Compound 1 in Patients with Advanced B Cell Malignancies

The first-in-human multi-center, open-label phase 1 trial of Compound 1 is being conducted in Australia and New Zealand and is comprised of two parts—a dose-escalation phase involving 25 patients and a dose-expansion phase, in which we plan to enroll a total of 100 patients. A total of 39 patients, including all 25 patients from the initial dose-escalation component and 14 patients from the ongoing dose-expansion component were enrolled. Based on the pharmacokinetics, pharmacodynamics, safety and efficacy of Compound 1 in the dose-escalation phase, 320 mg once daily (QD) and 160 mg twice daily (BID) are being further explored in the ongoing dose-expansion trial.

As of Oct. 19, 2015, the cutoff date for data analysis, 29 objective responses have been observed, including 3 complete responses (CRs), 1 very good partial response (VGPR), and 25 partial responses (PRs). Responses by histology are summarized in Table 18. 31 of the 39 patients remain on study treatment, free of progression, including all patients to date who have achieved an objective response.

TABLE 18

Responses by histology of Patients						
	Follow-up Days Median (Range)	Best Response				ORR (CR + PR)
		CR	PR	SD	PD	
Chronic Lymphocytic Leukemia	220 (83-329)	0/14 (0%)	13/14 ¹ (93%)	1/14 (7%)	0 (0%)	13/14 (93%)
Mantle Cell Lymphoma	148 (84-392)	2/10 (20%)	6/10 (60%)	1/10 (10%)	1/10 (10%)	8/10 (80%)
Waldenström's Macroglobulinemia	271 (11-398)	0/7 (0%)	6/7 ² (86%)	0/7 (0%)	1/7 (14%)	6/7 (86%)
DLBCL	29 (20-236)	1/4 (25%)	0/4 (0%)	0/4 (0%)	3/4 (75%)	1/4 (25%)
Indolent NHL	233 (215-250)	0/2 (0%)	0/2 (0%)	2/2 (100%)	0/2 (0%)	0/2 (0%)
Hairy Cell Leukemia	362	0/1 (0%)	1/1 (100%)	0/1 (0%)	0/1 (0%)	1/1 (100%)
Burkitt's-like Lymphoma	84	0/1 (0%)	0/1 (0%)	0/1 (0%)	1/1 (100%)	0/1 (0%)

¹Includes five patients with lymphocytosis at latest assessment;

²Includes one patient with VGPR

Note:

CR = complete response;

PR = partial response;

SD = stable disease;

PD = progressive disease;

ORR = objective response rate

8 patients discontinued Compound 1, including 6 due to disease progression and 2 due to adverse events related to their underlying malignancy. 3 patients died during study as a result of disease progression or complications of disease

progression. There were no drug-related serious adverse events (SAEs). The vast majority of adverse events, regardless of relationship to treatment, were Grade 1 or 2 in severity and not treatment-limiting. Of the 19 >Grade 3 AEs, 4 were assessed by investigators as possibly drug-related—all were self-limited neutropenia, not requiring treatment discontinuation. There was one case of major hemorrhage, defined as a bleeding event grade 3 or higher or an intracranial bleeding event of any grade: GI hemorrhage in a mantle cell lymphoma patient with lymphomatous involvement of the GI tract; this bleeding event occurred during drug hold, and resolved rapidly with re-initiation of Compound 1 treatment, and therefore is not considered to be drug-related. 6 patients had a baseline history of atrial fibrillation/flutter (AF), and no exacerbation or new event of AF was reported.

(2) Ongoing Clinical Trial Phase 1 Result on Compound 1 in Patients with Waldenström's Macroglobulinemia (WM)

The multi-center, open-label Phase 1 trial of Compound 1 in B-cell malignancies is being conducted in Australia, New Zealand, South Korea, and the United States and consists of a dose-escalation phase and a dose-expansion phase in disease-specific cohorts, which include treatment naïve and relapsed/refractory waldenström's macroglobulinemia (R/R WM). The dose-escalation component of the trial tested total daily doses ranging from 40 mg to 320 mg, and the ongoing dose-expansion phase is testing doses of 160 mg twice a day (BID) or 320 mg once a day (QD). As of Mar. 31, 2017, 48 patients with WM were enrolled in the study. Responses were determined according to the modified Sixth International Workshop on WM (IWWM) criteria.

Compound 1 was shown to be well tolerated with no discontinuation for Compound 1-related toxicity to date. Adverse events (AEs) were generally mild in severity and self-limited. The most frequent AEs (>10%) of any attribution among 48 patients evaluable for safety were petechiae/purpura/contusion (35%), upper respiratory tract infection (31%), constipation (25%), diarrhea (19%), epistaxis (19%), nausea (17%), cough (15%), anemia (15%), headache (15%), neutropenia (13%), and rash (13%), all of which were grade 1 or 2 in severity except for grade 3 or 4 anemia and neutropenia (8% each) as well as grade 3 or 4 diarrhea and headache (2% each). Five serious AEs were assessed to be possibly related to Compound 1; these included one case each of hemothorax, atrial fibrillation, colitis, febrile neutropenia, and headache. Among AEs of special interest, there were a total of three cases of atrial fibrillation (all grade 1 or 2), and one case of serious hemorrhage (hemothorax), defined as grade 3 or higher hemorrhage or central nervous system hemorrhage of any grade. Three events led to treatment discontinuation: one case each of bronchiectasis, prostate adenocarcinoma, and adenocarcinoma of pylorus.

At the time of the data cutoff, 42 patients were evaluable for response. Patients not evaluable for efficacy included two patients with less than 12 weeks of follow-up, three patients with IgM <500 mg/dl at baseline, and one patient with inaccurate baseline IgM due to cryoprotein. At a median follow-up of 12.3 months (4.4-30.5 months), the ORR was 90% (38/42 patients) and the major response rate was 76% (32/42 patients), with VGPRs in 43% (18/42) of patients and partial responses in 33% (14/42) of patients.

(3) Ongoing Clinical Trial Phase 1 Result on Compound 1 in Patients with Chronic Lymphocytic Leukemia and Small Lymphocytic Lymphoma (CLL/SLL)

The multi-center, open-label Phase 1 trial of Compound 1 in patients with B-cell malignancies is being conducted in Australia, New Zealand, South Korea, and the United States


and consists of a dose-escalation phase and a dose-expansion phase in disease-specific cohorts, which include treatment naïve (TN) and relapsed/refractory (R/R) CLL/SLL. The dose-escalation component of the trial tested total daily doses between 40 mg and 320 mg, and the ongoing dose-expansion component is testing doses of 160 mg twice a day (5 BID) or 320 mg once a day (QD). As of Mar. 31, 2017, 69 patients with CLL or SLL (18 TN, 51 R/R) were enrolled in the study.

Compound 1 was shown to be well tolerated in CLL/SLL. The most frequent adverse events (AEs) (>10%) of any attribution were petechiae/purpura/contusion (46%), fatigue (29%), upper respiratory tract infection (28%), cough (23%), diarrhea (22%), headache (19%), hematuria (15%), nausea (13%), rash (13%), arthralgia (12%), muscle spasms (12%), and urinary tract infection (12%); all of these events were grade 1 or 2 except for one case of grade 3 purpura (subcutaneous hemorrhage), which was the only major bleeding event. Additional adverse events of interest included one case of each grade 2 diarrhea and grade 2 atrial fibrillation. A total of 18 serious AEs (SAES) occurred in 13 patients, with no SAE occurring in more than one patient. Only one patient discontinued treatment due to an AE, a grade 2 pleural effusion.

At the time of the data cutoff, 66 patients (16 TN and 50 R/R) had more than 12 weeks of follow-up and were evaluable for efficacy, and three other patients had less than 12 weeks of follow-up. After a median follow-up of 10.5 months (2.2-26.8 months), the overall response rate (ORR) was 94% (62/66) with complete responses (CRs) in 3% (2/66), partial responses (PRs) in 82% (54/66), and PRs with lymphocytosis (PR-Ls) in 9% (6/66) of patients. Stable disease (SD) was observed in 5% (3/66) of patients. The patient with pleural effusion discontinued treatment prior to week 12 and was not evaluable for response. There was one instance of Hodgkin's transformation. In TN CLL/SLL, at a median follow-up time of 7.6 months (3.7-11.6 months), the ORR was 100% (16/16) with CRs in 6% (1/16), PRs in 81% (13/16) and PR-Ls in 13% (2/16) of patients. In R/R CLL/SLL, at a median follow-up time of 14.0 months (2.2-26.8 months), the ORR was 92% (46/50) with CRs in 2% (1/50), PRs in 82% (41/50), and PR-Ls in 8% (4/50) of patients. Stable disease was observed in 6% (3/50) patients.

The invention claimed is:

1. A method for treating a B-cell proliferative disease in a subject, comprising administering to the subject in need thereof Compound 1,



wherein the B-cell proliferative disease is selected from a group consisting of chronic lymphocytic leukemia, small lymphocytic lymphoma, mantle cell lymphoma, Waldenström's macroglobulinemia, marginal zone lymphoma, and follicular lymphoma; and Compound 1 is administered at a dose of 320 mg once a day (QD).

2. The method of claim 1, wherein Compound 1 has a purity of at least 99.3%.

3. The method of claim 1, wherein Compound 1 has a purity of at least 99.5%.

4. The method of claim 3, wherein the B-cell proliferative disease is mantle cell lymphoma.

5. The method of claim 4, wherein the subject has received at least one prior therapy.

6. The method of claim 3, wherein the B-cell proliferative disease is Waldenström's macroglobulinemia.

7. The method of claim 3, wherein the B-cell proliferative disease is marginal zone lymphoma.

8. The method of claim 7, wherein the subject has received at least one prior therapy.

9. The method of claim 8, wherein the marginal zone lymphoma is relapsed or refractory marginal zone lymphoma.

10. The method of claim 3, wherein the B-cell proliferative disease is chronic lymphocytic leukemia.

11. The method of claim 3, wherein the B-cell proliferative disease is small lymphocytic lymphoma.

12. The method of claim 3, wherein the B-cell proliferative disease is follicular lymphoma.

13. The method of claim 3, wherein an amorphous form or a crystalline form of Compound 1 is administered.

14. The method of claim 13, wherein a crystalline form of Compound 1 is administered.

15. The method of claim 14, wherein the crystalline form exhibits an X-ray powder diffraction pattern comprising diffraction peaks having 2θ angle values at 14.8±0.2°, 16.4±0.2° and 21.4±0.2°.

16. The method of claim 15, wherein the crystalline form exhibits an X-ray powder diffraction pattern comprising diffraction peaks having 2θ angle values at 14.8±0.2°, 15.6±0.2°, 16.4±0.2° and 21.4±0.2°.

17. The method of claim 16, wherein the B-cell proliferative disease is mantle cell lymphoma, wherein the subject has received at least one prior therapy.

18. The method of claim 16, wherein the B-cell proliferative disease is Waldenström's macroglobulinemia.

19. The method of claim 16, wherein the B-cell proliferative disease is relapsed or refractory marginal zone lymphoma, wherein the subject has received at least one prior therapy.

20. The method of claim 16, wherein the B-cell proliferative disease is chronic lymphocytic leukemia or small lymphocytic lymphoma.

21. The method of claim 13, wherein an amorphous form of Compound 1 is administered.

22. The method of claim 21, wherein the amorphous form of Compound 1 has a mid-point temperature of a glass transition temperature at 79.7° C.

23. The method of claim 22, wherein the amorphous form has an enantiomeric excess value of at least 97%.

24. The method of claim 23, wherein the B-cell proliferative disease is mantle cell lymphoma, wherein the subject has received at least one prior therapy.

25. The method of claim 23, wherein the B-cell proliferative disease is Waldenström's macroglobulinemia.

26. The method of claim 23, wherein the B-cell proliferative disease is relapsed or refractory marginal zone lymphoma, wherein the subject has received at least one prior therapy.

27. The method of claim 23, wherein the B-cell proliferative disease is chronic lymphocytic leukemia or small lymphocytic lymphoma. 5

28. The method of claim 3, wherein Compound 1 is administered orally.

* * * * *