



US 20030165946A1

(19) **United States**

(12) **Patent Application Publication**
Evans

(10) **Pub. No.: US 2003/0165946 A1**

(43) **Pub. Date: Sep. 4, 2003**

(54) **METHOD FOR THE COMPLETE
CHEMICAL SYNTHESIS AND ASSEMBLY OF
GENES AND GENOMES**

(60) Provisional application No. 60/059,017, filed on Sep. 16, 1997.

Publication Classification

(75) Inventor: **Glen A. Evans**, San Marcos, CA (US)

(51) **Int. Cl.⁷ C12Q 1/68; C12P 19/34**

(52) **U.S. Cl. 435/6; 435/91.2**

Correspondence Address:
CAMPBELL & FLORES LLP
4370 LA JOLLA VILLAGE DRIVE
7TH FLOOR
SAN DIEGO, CA 92122 (US)

(57) **ABSTRACT**

(73) Assignee: **Egea Biosciences, Inc.**

The present invention relates generally to the fields of oligonucleotide synthesis. More particularly, it concerns the assembly of genes and genomes of completely synthetic artificial organisms. Thus, the present invention outlines a novel approach to utilizing the results of genomic sequence information by computer directed gene synthesis based on computing on the human genome database. Specifically, the present invention contemplates and describes the chemical synthesis and resynthesis of genes defined by the genome sequence in a host vector and transfer and expression of these sequences into suitable hosts.

(21) Appl. No.: **10/322,360**

(22) Filed: **Dec. 17, 2002**

Related U.S. Application Data

(63) Continuation of application No. 09/554,929, filed on May 12, 2000, now Pat. No. 6,521,427, filed as 371 of international application No. PCT/US98/19312, filed on Sep. 16, 1998.

Figure 1. Jurassic Park Paradigm For The Construction Of Synthetic Organism

DNA SEQUENCE INFORMATION

- Step 1. Determine/design DNA sequence of the genome
- Step 2. Synthesize and assemble the genomic DNA
- Step 3. Introduce the DNA into an enucleated pleuripotent host cell.
- Step 4. Introduce the host cell into a foster mother animal

SYNTHETIC ORGANISM

Figure 2. Strategy for synthetic genetics and assembly of organisms

1. Design genome, containing prokaryotic origin of replication and drug selection vector.
2. SynGen 2.0, breaks down genome into component overlapping oligonucleotides, programs oligonucleotide synthesizer
3. Chemical synthesis of component oligonucleotide using MERMADE high throughput synthesizer
3. Combinatoric assembly of component oligonucleotides using robotic processing
4. Transformation into component bacteria

Figure 3
Strategy for Synthetic
Genetics and Combinatoric
Assembly




Figure 4A
Synlux/4 synthetic plasmid, luciferase, Knr 4800 bp (SEQ ID NO:1)

```

10      20      30      40      50      60      70      80      90     100
aagcttacctcgatttgaggacgttacaaagtattactgttaaggagcgttagattaaaaaattgaaaattgaaaatttagaattggccttaataaac
agaatcaccaaaaaaggaaatagagatgaaatttggaataatctgtttttcgtatcaaccaccagggtgaaactcataagctaaagtaattggatcgcctctgtt
luxA --- >

```

```

cggttgggtatcgccctcagaagagtaggggttttgatcacatattggaccttagaacatcattttacagaggtttggcttaccgggaaaatttatttggctgctgc
ggctaacctgtttaggagaactaaaacattaaattgtttggcactatgggggtttgttatcccgacagcacaccagttccgacagtttgaagaagcttttattt
ttagatcaaatgtcgaagggtcgtttttaaatttggaaaccttcgagggctataccataaagattttcggatatttgggtgttgatatggaagagtcctcgag
caattactcaaaaatttctaccagatgataatggaaagctttacagacaggaaccattagctctgatatgattacattcaatttccctaaaggttgatgtata
tcccaaaagtgtactcaaaaaattgtaccaaacctgtatgactgtgagtcgcgaagtacgacagaatggctagcaatacaaaagggttacccaattggctttagl
tggatttattggactaataatgaaaaaaagcacagatggaaactctataatgaaattggacagaaatattggctcattgatatcttaaaatagatcattgtatga
cttataatttgttctgttgatgatgatgacaaaagggcaagatgtttgctgggagtttctgaaaaattgggtatgactcattatgtaaatggaccataat
ctttaatgatagcaatcaaaactcgtgttatgattatcataaaaggtcaatggcgtgatttttacaaggacatatacaaacaccatcggacgtgttgat
tatagcaattgggtattaaacctgttaggcactcctcagcagtgatttgaatcattcaacgtgatttggatgcaacgggtattacaacattacatggcggat
ttgaaagcctaatggaaactgaagatgaaataattgcttccatggacgctttatgacacaaagtcgctccttctttaaagaacccaataaaattacttattt
galactagagataataaaggaaacaagttatgaaatttggattatttttctaaacttccagaaagatggaaataacatctgaaagaaacgtttggataaatggg

```

luxB --- >

```

taaaagctgtacgttaattgattccaactaaatcatttttaatactgcccctttgttaatgaacatcactttttcaaaaaattggtattgttggagacacctat
taccgacgctgggtttttttattagggttaacaaaataaaattacatattgggttcaataaacagtaattaccacccaaccctgtactgttagcagaagaa
ggcagtttattagatccaaatgtcagagggacgcttcttgggttttagtgactcggaaagtgatttcgaaaaggaaatttttttagacgtcatalctcat
caaggcaacaacaatttgaagcactgtatgaaataatlaatgacgcatlaactacaggttatgtcattccccaaaaacgacttttattgattttcccaaaaggt
ttcaattaatcccacactgttacagtgagaatggaccctaaagcaatattgtatccgtacatcaaaaagaaagtcgtcatgtggcagcgaaaaaaggcactgcct
tbaacatttaagtgggagataaatttagaaaacaaaagacgctatgcaattctataataataaaacagcacaacaaactatgggtattgatatctcggatgttg
atcatcaatlaactglaattgogaacttaaatgctgatagaagtcggctcaagaagaagtgagagaataacttaaaagactatatacactgaaacttacc
tcaaatggacagagatgaaaaaaattaaactgcaattatgaaagagaatgcagttgggtctcattgatgactattatgaatcgacaaaaattagcagtggaaaaa
acagggctcaaaaaatatttattatccttbbgaatcaatgtccgatatataaagatgtaaaaagctattatgatattgttgaaacccaaaaaatcgaaatgaaatt
taccataataaaaaattaaaggcaatttctatatatbagattgccccttttggggatccctctagaaaatttttattctgatlaataagatgagaattcacttggccg

```

pUC 19 --- >

```

tcgttttacaaactcgtgactgggaaaaacccctggcgtttacccaacttaatcgcccttgacgacatcccccttttggccagctggcgttaatagcgaagagggc
ccgcaaccgatcgcccttcccacagcttggcagcctgaaaggcgaattggcgcctgatgggtattttctccttacgcactctgtgcggtatttccacaccgc
atattggcactctcagtaaatctgctctgatgcccgaatagttaaagccagcccgcaccccgaaccccgcctgacgcgcctgacggccttgcctgtctgc
tcccggcatccgcttacagacaagctgtgacccgtctccgggagctgcatgtgtcagaggttttccaccgtcaccacgaacccgcgcgagacgaaaggccct
cgtgatcgcctatttttataaggttaattgcatgataataatgggttcttagacgtcaggtggcactttcggggaaatgtgcgcggaaccccctatttgt
tatttttctaaaaagcttcaagctcagcagcactcagggcgaagggctgctaaagggaagcgaacacgttagaaaagccagtcocgagaacgggtgct
gaccccggatgaatgtcagctactgggctatctggacaaggggaaaaacgcaagcgaagagaaagcaggttagcttgcagttgggttacaatggcgatagct
agactgggcccgttttattggacagcgaagcgaacgggaattgcccagctgggcccctctggttaaggtttgggaagccctgcaaaagtaaaactggatggccttcc
ttggcccacaaggatctgatggcaggggatcaagatctgatcaagagacaggtaggatcgtttcgcgatgtgaacaagatggatggcagcaggtt

```

kan/neo phosphotransferase ---

Fig. 5A List of component oligonucleotides derived from the sequence of Synlux4 in Figure 4 (SEQ ID No: 2 through to SEQ ID No: 193)

oligoF01, AAGCTTACCTCGATTTGAGGACGTTACAAGTATTACTGTTAAGGAGCGTA
oligoF02, GATTAAAAAATGAAATTGAAAATGAATTATTAGAATTGGCTTAAATAAAC
oligoF03, AGAATCACCAAAAAGGAATAGAGTATGAAGTTTGGAAATATTTGTTTTTC
oligoF04, GTATCAACCACCAGGTGAAACTCATAAGCTAAGTAATGGATCGCTTTGTT
oligoF05, CGGCTTGGTATCGCCTCAGAAGAGTAGGGTTTGATACATATTGGACCTTA
oligoF06, GAACATCATTTTACAGAGTTTGGTCTTACGGGAAATTTATTTGTTGCTGC
oligoF07, GGCTAACCTGTTAGGAAGAATAAAACATTAAATGTTGGCACTATGGGGG
oligoF08, TTGTTATTCCGACAGCACACCCAGTTCGACAGTTAGAAGACGTTTTATTA
oligoF09, TTAGATCAAATGTCGAAAGGTCGTTTTAATTTTGGAAACCGTTCGAGGGCT
oligoF10, ATACCATAAAGATTTTCGAGTATTTGGTGTGATATGGAAGAGTCTCGAG
oligoF11, CAATTACTCAAATTTCTACCAGATGATAATGGAAAGCTTACAGACAGGA
oligoF12, ACCATTAGCTCTGATAGTGATTACATTCAATTTCTAAGGTTGATGTATA
oligoF13, TCCCAAAGTGTACTCAAAAAATGTACCAACCTGTATGACTGCTGAGTCCG
oligoF14, CAAGTACGACAGAATGGCTAGCAATACAAGGGCTACCAATGGTTCTTAGT
oligoF15, TGGATTATTGGTACTAATGAAAAAAAAGCACAGATGGAACCTATAATGA
oligoF16, AATTGCGACAGAATATGGTCATGATATATCTAAAATAGATCATTGTATGA
oligoF17, CTTATATTTGTTCTGTTGATGATGATGCACAAAAGGCGCAAGATGTTTGT
oligoF18, CGGGAGTTTCTGAAAAATTGGTATGACTCATATGTAAATGCGACCAATAT
oligoF19, CTTTAATGATAGCAATCAAACCTCGTGGTTATGATTATCATAAAGGTCAAT
oligoF20, GCGTGATTTTGTTTTACAAGGACATACAAACACCAATCGACGTGTTGAT
oligoF21, TATAGCAATGGTATTAACCCTGTAGGCACTCCTGAGCAGTGTATTGAAAT
oligoF22, CATTCAACGTGATATTGATGCAACGGGTATTACAAACATTACATGCGGAT
oligoF23, TTGAAGCTAATGGAACCTGAAGATGAAATAATTGCTTCCATGCGACGCTTT
oligoF24, ATGACACAAGTCGCTCCTTTCTTAAAAGAACCTAAATAAATTACTTATTT
oligoF25, GATACTAGAGATAATAAGGAACAAGTTATGAAATTTGGATTATTTTTTTCT
oligoF26, AAACCTTCAGAAAGATGGAATAACATCTGAAGAAACGTTGGATAATATGG
oligoF27, TAAAGACTGTCACGTTAATTGATTCAACTAAATATCATTTTAATACTGCC
oligoF28, TTTGTTAATGAACATCACTTTTCAAAAAATGGTATTGTTGGAGCACCTAT
oligoF29, TACCGCAGCTGGTTTTTTATTAGGGTTAACAAATAAATTACATATTGGTT
oligoF30, CATTAAATCAAGTAATTACCACCCATCACCCCTGTACGTGTAGCAGAAGAA
oligoF31, GCCAGTTTATTAGATCAAATGTCAGAGGGACGCTTCATTCTTGGTTTTAG
oligoF32, TGACTGCGAAAGTGATTTTCGAAATGGAATTTTTTTAGACGTCATATCTCATV
oligoF33, CAAGGCAACAACAATTTGAAGCATGCTATGAAATAATTAATGACGCATTA
oligoF34, ACTACAGGTTATTGTCATCCCCAAAACGACTTTTATGATTTTCCAAAGGT
oligoF35, TTCAATTAATCCACACTGTTACAGTGAGAATGGACCTAAGCAATATGTAT
oligoF36, CCGCTACATCAAAGAAGTCGTCATGTGGGCAGCGAAAAAGGCACTGCCT
oligoF37, TTAACATTTAAGTGGGAGGATAATTTAGAAACCAAAGAACGCTATGCAAT
oligoF38, TCTATATAATAAAACAGCACACAATATGGTATTGATATTTCCGGATGTTG
oligoF39, ATCATCAATTAAGTGAATTTGCGAACTTAAATGCTGATAGAAGTACGGCT
oligoF40, CAAGAAGAAGTGAGAGAATACTTAAAAGACTATATCACTGAAACTTACCC
oligoF41, TCAAATGGACAGAGATGAAAAAATTAAGTGCATTATTGAAGAGAATGCAG
oligoF42, TTGGGTCTCATGATGACTATTATGAATCGACAAAATTAGCAGTGGAAAAA
oligoF43, ACAGGGTCTAAAAATATTTTATTATCCTTTGAATCAATGTCCGATATTAA
oligoF44, AGATGTAAAAGATAFTATTGATATGTTGAACCAAAAAATCGAAATGAATT
oligoF45, TACCATAATAAAATTAAGGCAATTTCTATATTAGATTGCCTTTTTGGGG
oligoF46, ATCCTCTAGAAATATTTTATCTGATTAATAAGATGAGAATTCAGTGGCCC
oligoF47, TCGTTTTACAACGTCGTGACTGGGAAAACCTGGCGTTACCCAACCTAAT
oligoF48, CGCCTTGACAGCACATCCCCCTTTCGCCAGCTGGCGTAATAGCGAAGAGGC
oligoF49, CCGCACCGATCGCCCTTCCCAACAGTTGCGCAGCCTGAATGGCGAATGGC

Figure 5B

oligoF50, GCCTGATGCGGTATTTTCTCCTTACGCATCTGTGCGGTATTTACACCCGC
oligoF51, ATATGGTGCACCTCTCAGTACAATCTGCTCTGATGCCGCATAGTTAAGCCA
oligoF52, GCCCCGACACCCGCCAACACCCGCTGACGCGCCCTGACGGGCTTGTCTGC
oligoF53, TCCCGGCATCCGCTTACAGACAAGCTGTGACCGTCTCCGGGAGCTGCATG
oligoF54, TGTCAGAGGTTTTTCACCGTCATCACCGAAACGCGCGAGACGAAAGGGCCT
oligoF55, CGTGATACGCCTATTTTTATAGGTTAATGTCATGATAATAATGGTTTCTT
oligoF56, AGACGTCAGGTGGCACTTTTCGGGGAAATGTGCGCGGAACCCCTATTTGT
oligoF57, TTATTTTTCTAAAAAGCTTTCACGCTGCCGCAAGCACTCAGGGCGCAAGGG
oligoF58, CTGCTAAAGGAAGCGGAACACGTAGAAAGCCAGTCCGCAGAAACGGTGCT
oligoF59, GACCCCGGATGAATGTCAGCTACTGGGCTATCTGGACAAGGGAAAACGCA
oligoF60, AGCGCAAAGAGAAAGCAGGTAGCTTGCAGTGGGCTTACATGGCGATAGCT
oligoF61, AGACTGGGCGGTTTTTATGGACAGCAAGCGAACCAGGAATTGCCAGCTGGGG
oligoF62, CGCCCTCTGGTAAGGTTGGGAAGCCCTGCAAAGTAAACTGGATGGCTTTC
oligoF63, TTGCCGCCAAGGATCTGATGGCGCAGGGGATCAAGATCTGATCAAGAGAC
oligoF64, AGGATGAGGATCGTTTCGCATGATTGAACAAGATGGATTGCACGCAGGTT
oligoF65, CTCCGGCCGCTTGGGTGGAGAGGCTATTCGGCTATGACTGGGCACAACAG
oligoF66, ACAATCGGCTGCTCTGATGCCGCCGTGTTCCGGCTGTCAGCGCAGGGGCG
oligoF67, CCCGGTTCTTTTTGTCAAGACCGACCTGTCCGGTGCCCTGAATGAACTGC
oligoF68, AGGACGAGGCAGCGCGGCTATCGTGGCTGGCCACGACGGGCGTTCCTTGC
oligoF69, GCAGCTGTGCTCGACGTTGTCACCTGAAGCGGGAAGGGACTGGCTGCTATT
oligoF70, GGGCGAAGTGCCGGGGCAGGATCTCCTGTCATCTCACCTTGCTCCTGCCG
oligoF71, AGAAAGTATCCATCATGGCTGATGCAATGCGGCGGCTGCATACGCTTGAT
oligoF72, CCGGCTACCTGCCATTTCGACCACCAAGCGAAACATCGCATCGAGCGAGC
oligoF73, ACGTACTCGGATGGAAGCCGGTCTTGTTCGATCAGGATGATCTGGACGAAG
oligoF74, AGCATCAGGGGCTCGCGCCAGCCGAACTGTTCCGCCAGGCTCAAGGCGCGC
oligoF75, ATGCCCGACGGCGAGGATCTCGTTCGTGACCCATGGCGATGCCTGCTTGCC
oligoF76, GAATATCATGGTGGAAAATGGCCGCTTTTTCTGGATTTCATCGACTGTGGCC
oligoF77, GGCTGGGTGTGGCGGACCGCTATCAGGACATAGCGTTGGCTACCCGTGAT
oligoF78, ATTGCTGAAGAGCTTGGCGGCGAATGGGCTGACCGCTTCCTCGTGCTTTA
oligoF79, CGGTATCGCCGCTCCCGATTTCGACGCGCATCGCCTTCTATCGCCTTCTTG
oligoF80, ACGAGTTCTTCTGAGCGGGACTCTGGGGTTTCGAAATGACCGACCAAGCGA
oligoF81, CGCCCAACCTGCCATCACGAGATTTTCGATTCCACCGCCGCCTTCTATGAA
oligoF82, AGGTTGGGCTTCGGAATCGTTTTCCGGGACGCGGGCTGGATGATCCTCCA
oligoF83, GCGCGGGGATCTCATGCTGGAGTTCTTCGCCCACCCCGGGCATGACCAAA
oligoF84, ATCCCTTAACGTGAGTTTTTCGTTCCACTGAGCGTCAGACCCCGTAGAAAA
oligoF85, GATCAAAGGATCTTCTTGAGATCCTTTTTTTCTGCGCGTAATCTGCTGCT
oligoF86, TGCAAACAAAAAAACCACCGCTACCAGCGGTGGTTTGTTTGGCCGGATCAA
oligoF87, GAGCTACCAACTCTTTTTCCGAAGGTAACCTGGCTTCAGCAGAGCGCAGAT
oligoF88, ACCAAATACTGTCCTTCTAGTGTAGCCGTAGTTAGGCCACCACTTCAAGA
oligoF89, ACTCTGTAGCACCGCCTACATACTCGCTCTGCTAATCCTGTTACCAGTG
oligoF90, GCTGCTGCCAGTGGCGATAAGTTCGTGCTTACCAGGGTTGGACTCAAGACG
oligoF91, ATAGTTACCGGATAAGGCGCAGCGGTCCGGGCTGAACGGGGGGTTTCGTGCA
oligoF92, CACAGCCCAGCTTGGAGCGAACGACCTACACCGAACTGAGATACCTACAG
oligoF93, CGTGAGCTATGAGAAAGCGCCACGCTTCCCAGAGGGAGAAAGGCGGACAG
oligoF94, GTATCCGGTAAGCGGCAGGGTCCGGAACAGGAGAGCGCACGAGGGAGCTTC
oligoF95, CAGGGGGAAACGCCTGGTATCTTTATAGTCTGTGCGGGTTTCGCCACCTC
oligoF96, TGACTTGAGCGTCGATTTTTGTGATGCTCGTCAGGGGGGCGGAGCCTATG
oligoR01, CATCACAAAAATCGACGCTCAAGTCAGAGGTGGCGAAACCCGACAGGACT
oligoR02, ATAAAGATAACCAGGCGTTTTCCCCCTGGAAGCTCCCTCGTGCGCTCTCCTG
oligoR03, TTCCGACCCTGCCGCTTACCAGGATACCTGTCCGCCTTTCTCCCTTCGGGA
oligoR04, AGCGTGGCGCTTTCTCATAGCTCACGCTGTAGGTATCTCAGTTCGGTGTA
oligoR05, GGTCGTTTCGCTCCAAGCTGGGCTGTGTGCACGAACCCCCCGTTCAGCCCG
oligoR06, ACCGCTGCGCCTTATCCGGTAACTATCGTCTTGAGTCCAACCCGGTAAGA

Figure 5C

oligoR07, CACGACTTATCGCCACTGGCAGCAGCCACTGGTAACAGGATTAGCAGAGC
oligoR08, GAGGTATGTAGGCGGTGCTACAGAGTTCTTGAAGTGGTGGCCTAACTACG
oligoR09, GCTACACTAGAAGGACAGTATTTGGTATCTGCGCTCTGCTGAAGCCAGTT
oligoR10, ACCTTCGGAAAAAGAGTTGGTAGCTCTTGATCCGGCAAACAAACCACCGC
oligoR11, TGGTAGCGGTGGTTTTTTTTGTTTTGCAAGCAGCAGATTACGCGCAGAAAA
oligoR12, AAGGATCTCAAGAAGATCCTTTGATCTTTTCTACGGGGTCTGACGCTCAG
oligoR13, TGGAACGAAAACCTCACGTTAAGGGATTTTGGTTCATGCCCGGGGTGGGCGA
oligoR14, AGAACTCCAGCATGAGATCCCCGCGCTGGAGGATCATCCAGCCGGCGTCC
oligoR15, CGGAAAACGATTCCGAAGCCCAACCTTTCATAGAAGGCGGCGGTGGAATC
oligoR16, GAAATCTCGTGATGGCAGGTTGGGCGTTCGCTTGGTTCGGTCATTTTGAACC
oligoR17, CCAGAGTCCCGCTCAGAAGAAGTCTGCAAGAAGGCGATAGAAGGCGATGC
oligoR18, GCTGCGAATCGGGAGCGGCGATAACCGTAAAGCACGAGGAAGCGGTCAGCC
oligoR19, CATTGCGCCCAAGCTCTTCAGCAATATCACGGGTAGCCAACGCTATGTC
oligoR20, CTGATAGCGGTCCGCCACACCCAGCCGGCCACAGTCGATGAATCCAGAAA
oligoR21, AGCGGCCATTTTCCACCATGATATTCGGCAAGCAGGCATCGCCATGGGTC
oligoR22, ACGACGAGATCCTCGCCGTCGGGCATGCGCGCCTTGAGCCTGGCGAACAG
oligoR23, TTCGGCTGGCGCGAGCCCCTGATGCTCTTCGTCCAGATCATCCTGATCGA
oligoR24, CAAGACCGGCTTCCATCCGAGTACGTGCTCGCTCGATGCGATGTTTTGCT
oligoR25, TGGTGGTGAATGGGCAGGTAGCCGGATCAAGCGTATGCAGCCGCCGCAT
oligoR26, TGCATCAGCCATGATGGATACTTTCTCGGCAGGAGCAAGGTGAGATGACA
oligoR27, GGAGATCCTGCCCCGGCACTTCCGCCAATAGCAGCCAGTCCCTTCCCGCT
oligoR28, TCAGTGACAACGTGAGCACAGCTGCGCAAGGAACGCCCGTTCGTGGCCAG
oligoR29, CCACGATAGCCGCGCTGCCTCGTCTGAGTTCATTCAGGGCACCGGACA
oligoR30, GGTCCGTCTTGACAAAAAGAACCAGGGCGCCCCCTGCGCTGACAGCCGGAAC
oligoR31, ACGGCGGCATCAGAGCAGCCGATGTCTGTGTGCCCCAGTCATAGCCGAA
oligoR32, TAGCCTCTCCACCCAAGCGGCCGGAGAACCCTGCGTGCAATCCATCTTGTT
oligoR33, CAATCATGCGAAACGATCCTCATCCTGTCTCTTGATCAGATCTTGATCCC
oligoR34, CTGCGCCATCAGATCCTTGGCGGCAAGAAAGCCATCCAGTTTACTTTGCA
oligoR35, GGGCTTCCCAACCTTACCAGAGGGCGCCCCAGCTGGCAATTCCGGTTCGC
oligoR36, TTGCTGTCCATAAAACCGCCAGTCTAGCTATCGCCATGTAAGCCCCTG
oligoR37, CAAGCTACCTGCTTTCTCTTTGCGCTTTCGTTTTCCCTTGTCCAGATAGC
oligoR38, CCAGTAGCTGACATTCATCCGGGGTTCAGCACCCGTTTCTGCGGACTGGCTT
oligoR39, TCTACGTGTTCCGCTTCTTTAGCAGCCCTTTCGCGCCCTGAGTGCTTTCGG
oligoR40, CAGCGTGAAGCTTTTTAGAAAAATAAAACAAATAGGGGTTCGCGCACATT
oligoR41, TCCCCGAAAAGTGCCACCTGACGTCTAAGAAACCATTATTATCATGACAT
oligoR42, TAACCTATAAAAATAGGCGTATCACGAGGCCCTTTCGTCTCGCGCGTTTC
oligoR43, GGTGATGACGGTGAAAACCTCTGACACATGCAGCTCCCGGAGACGGTCAC
oligoR44, AGCTTGTCTGTAAGCGGATGCCGGGAGCAGACAAGCCCGTCAGGGCGCGT
oligoR45, CAGCGGGTGTGGCGGGTGTGCGGGCTGGCTTAACTATGCGGCATCAGAG
oligoR46, CAGATTGTAAGGAGTGCACCATATGCGGTGTGAAATACCGCACAGATG
oligoR47, CGTAAGGAGAAAATACCGCATCAGGCGCCATTTCGCCATTTCAGGCTGCGCA
oligoR48, ACTGTTGGGAAGGGCGATCGGTGCGGGCCTCTTCGCTATTACGCCAGCTG
oligoR49, GCGAAAGGGGGATGTGCTGCAAGGCGATTAAGTTGGGTAAACGCCAGGGTT
oligoR50, TTCCCAGTCACGACGTTGTAACAAACGACGGCCAGTGAATTCTCATCTTATT
oligoR51, AATCAGATAAAAATATTTCTAGAGGATCCCCAAAAGGCAATCTAATATAG
oligoR52, AAATTGCCTTTAATTTTATTATGGTAAATTCATTTTCGATTTTTTGGTTCA
oligoR53, ACATATCAATAATATCTTTTACATCTTTAATATCGGACATTGATTCAAAG
oligoR54, GATAATAAAAATATTTTAGACCCTGTTTTTTCCACTGCTAATTTTGTGCA
oligoR55, TTCATAATAGTCATCATGAGACCCAACCTGCATTCTCTTCAATAATGCAGT
oligoR56, TAATTTTTTTCATCTCTGTCCATTTGAGGGTAAGTTTCAGTGATATAGTCT
oligoR57, TTTAAGTATTCTCTCACTTCTTCTTGAGCCGTAATTCTATCAGCATTTAA
oligoR58, GTTCGCAATTACAGTTAATTGATGATCAACATCCGAAATATCAATACCAT
oligoR59, ATTGTTGTGCTGTTTTATTATATAGAATTGCATAGCGTTCTTTGGTTTCT
oligoR60, AAATTATCCTCCACTTAAATGTTAAAGGCAGTGCCTTTTTTCGCTGCCCA

Figure 5D

oligoR61, CATGACGACTTCTTTTGATGTAGCGGATACATATTGCTTAGGTCCATTCT
oligoR62, CACTGTAACAGTGTGGATTAATTGAAACCTTTGGAAAATCATAAAAGTCG
oligoR63, TTTTGGGGATGACAATAACCTGTAGTTAATGCGTCATTAATTATTTTCATA
oligoR64, GCATGCTTCAAATTGTTGTTGCCTTGATGAGATATGACGTCTAAAAAATT
oligoR65, CCATTCGAAATCACTTTCGCAGTCACTAAAACCAAGAATGAAGCGTCCC
oligoR66, TCTGACATTTGATCTAATAAACTGGCTTCTTCTGCTACACGTACAGGGTG
oligoR67, ATGGGTGGTAATTACTTGATTTAATGAACCAATATGTAATTTATTTGTTA
oligoR68, ACCCTAATAAAAAACCAGCTGCGGTAATAGGTGCTCCAACAATACCATTT
oligoR69, TTTGAAAAGTGATGTTCAATTAACAAAGGCAGTATTAAAATGATATTTAGT
oligoR70, TGAATCAATTAACGTGACAGTCTTTACCATATTATCCAACGTTTCTTCAG
oligoR71, ATGTTATTCCATCTTTCTGAAAGTTTAGAAAAATAATCCAAATTTTCATA
oligoR72, ACTTGTTCCCTTATTATCTCTAGTATCAAATAAGTAATTTATTTAGGTTCT
oligoR73, TTTAAGAAAGGAGCGACTTGTGTCATAAAGCGTCGCATGGAAGCAATTAT
oligoR74, TTCATCTTCAGTTCATTAGCTTCAAATCCGCATGTAATGTTTGTAATAC
oligoR75, CCGTTGCATCAATATCACGTTGAATGATTTCAATACACTGCTCAGGAGTG
oligoR76, CCTACAGGGTTAATAACCATTGCTATAATCAACACGTCGATTGGTGTGTTGT
oligoR77, ATGTCCTTGTA AAAACAAAATCACGCCATTGACCTTTATGATAATCATAAC
oligoR78, CACGAGTTTGATTGCTATCATTAAAGATATTGGTCGCATTTACATATGAG
oligoR79, TCATACCAATTTTTCAGAACTCCCGACAAACATCTTGCGCCTTTTGTGC
oligoR80, ATCATCATCAACAGAACAAATATAAGTCATACAATGATCTATTTTAGATA
oligoR81, TATCATGACCATATTCTGTGCGCAATTTCAATTATAGAGTTCATCTGTGCT
oligoR82, TTTTTTTCATTAGTACCAATAATCCAACCTAAGAACCATTGGTAGCCCTTG
oligoR83, TATTGCTAGCCATTCTGTGCTACTTGCGGACTCAGCAGTCATACAGGTTG
oligoR84, GTACATTTTTTGAGTACACTTTGGGATATACATCAACCTTAGGAAATTGA
oligoR85, ATGTAATCACTATCAGAGCTAATGGTTCCTGTCTGTAAGCTTTCCATTAT
oligoR86, CATCTGGTAGAAATTTTGAGTAATTGCTCGAGACTCTTCCATATCAACAC
oligoR87, CAAATACTCGAAAATCTTTATGGTATAGCCCTCGAACGGTTCCAAATA
oligoR88, AAACGACCTTTCGACATTTGATCTAATAATAAAACGTCTTCTAACTGTCG
oligoR89, AACTGGGTGTGCTGTGCGGAATAACAACCCCATAGTGCCAACATTTAATG
oligoR90, TTTTAGTTCTTCCTAACAGGTTAGCCGCAGCAACAAATAAATTTCCCGTA
oligoR91, AGACCAAACCTCTGTA AAAATGATGTTCTAAGGTCCAATATGTATCAAACCC
oligoR92, TACTCTTCTGAGGCGATACCAAGCCGAACAAAGCGATCCATTACTTAGCT
oligoR93, TATGAGTTTACCTGGTGGTTGATACGAAAAACAAATATTTCCAAACTTC
oligoR94, ATACTCTATTCCTTTTTGGTGATTCTGTTTATTTAAGCCAATTCTAATAA
oligoR95, TTCATTTTCAATTTCAATTTTTTAATCTACGCTCCTTAACAGTAATACTTG
oligoR96, TAACGTCCTCAAATCGAGGTAAGCTTCATAGGCTCCGCCCCCTGACGAG

FIG. 6 Combinatoric assembly

Instruction set for 192 oligos (96 pairs).

- | | | | |
|----|--|----|---------------------------------|
| 1. | -F A1 → -C A1
-F A2 → -C A2
-F A3 → -C A3
-F A4 → -C A4
repeat with all wells to H12
-R A1 → -C A1
-R A2 → -C A2
-R A3 → -C A3
-R A4 → -C A4
repeat with all wells to H12 | 6. | B4 → C8
D12 → F4
G8 → H12 |
| | | 7. | C8 → H12
F4 → H12 |

All remaining operations on -C plate

- | | |
|----|--|
| 2. | A1 → A2
A3 → A4
A5 → A6
A7 → A8
A9 → A10
A11 → A12
repeat with each letter |
| 3. | A2 → A4
A6 → A8
A10 → A12
repeat with each letter |
| 4. | A4 → A8
A12 → B4
B8 → B12
C4 → C8
C12 → D4
D8 → D12
E4 → E8
E12 → F4
F8 → F12
G4 → G8
G12 → H4
H8 → H12 |
| 5. | A8 → B4
B12 → C8
D4 → D12
E8 → F4
F12 → G8
H4 → H12 |

SynGene Formatter v.4

Figure 7A

```

program Syn_Gene_Formatter (input, output, f, g, h);

{Synthetic Gene Formatting Program}
{This is a draft experimental program designed to break down a designer gene or genome }
{into oligonucleotides for synthesis. The program is for complete synthetic designer gene}
{construction. The program is based upon a an original program for formatting DNA sequences}
{written in 1988 by G . Evans for DNA sequence analysis and formatting}
{}
{This program is copyright (c) 1997 Glen A. Evans. All rights reserved}

const
  maxlength = 5000; {maximum length of sequence}
  searchlength = 10; {maximum length of search string}

var
  f: text; {inputfile of sequence}
  g: text; {output file of sequence}
  h: text; {output file of sequence}

{arrays for sequence formatting}

  dna: array[1..maxlength] of char;
  rdna: array[1..maxlength] of char;
  oligo: array[1..100] of char;

  l, k, seqlength: integer;
  nucin: char;
  oligolength, offset: integer;

  infile, outfile: string;

procedure initialize;

{This procedure initializes the program and opens the input file}

var
  s: string;

begin
  repeat
    write('>');
    readln(s);
  until length(s) = 0;
  writeln('Welcome to Syn_Gene_Formatter Version 1.0 - copyright (c) Glen A. Evans 1997');
  write('Enter the input file name: ');
  readln(infile);
  write('Enter the outfile name: ');
  readln(outfile);
  write('Enter the length of oligos you wish to use: ');

```

SynGene Formatter v.4

Figure 7B

```
readln(oligolength);
write('Enter the reverse oligo offset value: ');
readln(offset);
writeln('Thank you. ');
write('The program will now format the sequence into oligonucleotide fragments of length ');
writeln(oligolength);
writeln;
writeln;

end; {initialize}

procedure readinseq;
var
  j: integer;
  nuc: char;

begin
  writeln('reading input file');
  seqlength := 1;
  while not eof(f) do
    begin
      read(f, nuc);
      if nuc <> '' then
        begin
          if nuc = 'G' then
            dna[seqlength] := nuc;
          if nuc = 'A' then
            dna[seqlength] := nuc;
          if nuc = 'T' then
            dna[seqlength] := nuc;
          if nuc = 'C' then
            dna[seqlength] := nuc;
          if nuc = 'X' then
            dna[seqlength] := nuc;
          if nuc = 'N' then
            dna[seqlength] := nuc;
          seqlength := seqlength + 1;
        end;
      end;

      seqlength := seqlength - 1;
    end; {readinseq}

procedure readinfile;

begin
```

SynGene Formatter v.4

Figure 7C

```
reset(f, infile);
readinseq;
close(f);

end; {readinfile}

procedure writeforseq;

var

i, h, b, on: integer;

begin
write('fragmenting sequence into forward oligos');
b := 1;
on := 1;
rewrite(g, outfile);
writeln(g, infile);

while b < seqlength + 1 do
begin
write('.');
write(g, 'Oligo No.', on, ' ');
begin
for h := 1 to oligolength do
begin
write(g, dna[b]);
b := b + 1;
end;
on := on + 1;
writeln(g);
end;
end;
writeln;

end; {writeforseq}

procedure reverseseq;

{This procedure generates the reverse complement of the sequence}

var

i, h, b, a, on: integer;

begin
```

SynGene v.4

Figure 7D

```

write('generating the reverse complement');
b := seqlength;
for a := 1 to seqlength do
  begin
    if dna[b] = 'G' then
      rdna[a] := 'C';
    if dna[b] = 'A' then
      rdna[a] := 'T';
    if dna[b] = 'T' then
      rdna[a] := 'A';
    if dna[b] = 'C' then
      rdna[a] := 'G';
    b := b - 1;
    write('.');
  end;
writeln;

```

```
end; {reverseseq}
```

```
procedure writerevseq;
```

{This procedure fragments the reverse complement sequence starting at the offset value}

```
var
```

```
  i, h, b, on: integer;
```

```
begin
```

```
  write('fragmenting sequence into reverse oligos');
```

```
  on := 1;
```

```
  b := offset;
```

```
  while b < seqlength do
```

```
    begin
```

```
      writeln(g);
```

```
      write(g, 'Oligo No.', on, '. ');
```

```
      begin
```

```
        for h := 1 to oligolength do
```

```
          begin
```

```
            write(g, rdna[b]);
```

```
            b := b + 1;
```

```
          end;
```

```
        on := on + 1;
```

```
        write('.');
```

```
      end;
```

SynGene Formatter v.4

Figure 7E

```
end; {writerevseq}

procedure finaloligo;

var
  b, a: integer;

begin
  writeln;
  writeln('generating the last portion of the final oligo...');
  for a := 1 to offset do
    begin
      write(g, rdna[a])

    end;
  writeln(g);
  close(g);

end; {finaloligo}

begin {main}

  initialize;
  readinfile;
  writeforseq;
  reverseseq;
  writerevseq;
  finaloligo;
  writeln('processing completed');
  writeln('Have a nice day . ');

end. {main}
```


METHOD FOR THE COMPLETE CHEMICAL SYNTHESIS AND ASSEMBLY OF GENES AND GENOMES

BACKGROUND OF THE INVENTION

[0001] 1. Field of the Invention

[0002] The present invention relates generally to the fields of oligonucleotide synthesis. More particularly, it concerns the assembly of genes and genomes of completely synthetic artificial organisms.

[0003] 2. Description of Related Art

[0004] Present research and commercial applications in molecular biology are based upon recombinant DNA developed in the 1970's. A critical facet of recombinant DNA is molecular cloning in plasmids, covered under seminal patent of Cohen and Boyer (U.S. Pat. No. 4,740,470 "Biologically functional molecular chimeras"). This patent teaches a method for the "cutting and splicing" of DNA molecules based upon restriction endonucleases, the introduction of these "recombinant" molecules into host cells, and their replication in the bacterial hosts. This technique is the basis of all molecular cloning for research and commercial purposes carried out for the past 20 years and the basis of the field of molecular biology and genetics.

[0005] Recombinant DNA technology is a powerful technology, but is limited in utility to modifications of existing DNA sequences which are modified through 1) restriction enzyme cleavage sites, 2) PAC primers' for amplification, 3) site-specific mutagenesis, and other techniques. The creation of an entirely new molecule, or the substantial modification of existing molecules, is extremely time consuming, expensive, requires complex and multiple steps, and in some cases is impossible. Recombinant DNA technology does not permit the creation of entirely artificial molecules, genes, genomes or organisms, but only modifications of naturally-occurring organisms.

[0006] Current biotechnology for industrial production, for drug design and development, for potential applications of vaccine development and genetic therapy, and for agricultural and environmental use of recombinant DNA, depends on naturally-occurring organisms and DNA molecules. To create or engineer new or novel functions, or to modify organisms for specialized use (such as producing a human hormone), requires substantially complex, time consuming and difficult manipulations of naturally-occurring DNA molecules. In some cases, changes to naturally-occurring DNA are so complex that they are not possible in practice. Thus, there is a need for technology that allows the creation of novel DNA molecules in a single step without requiring the use of any existing recombinant or naturally-occurring DNA.

SUMMARY OF THE INVENTION

[0007] The present invention addresses the limitations in present recombinant nucleic acid manipulations by providing a fast, efficient means for generating practically any nucleic acid sequence, including entire genes, chromosomal segments, chromosomes and genomes. Because this approach is based on an completely synthetic approach, there are no limitations, such as the availability of existing nucleic acids, to hinder the construction of even very large segments of nucleic acid.

[0008] Thus, in a first embodiment, there is provided a method for the construction of a double-stranded DNA segment comprising the steps of (i) providing two sets of single-stranded oligonucleotides, wherein (a) the first set comprises the entire plus strand of said DNA segment, (b) the second set comprises the entire minus strand of said DNA segment, and (c) each of said first set of oligonucleotides being complementary to two oligonucleotides of said second set of oligonucleotides, (ii) annealing said first and said second set of oligonucleotides, and (iii) treating said annealed oligonucleotides with a ligating enzyme. Optional steps provide for the synthesis of the oligonucleotide sets and the transformation of host cells with the resulting DNA segment.

[0009] In particular embodiments, the DNA segment is 100, 200, 300, 400, 800, 100, 1500, 200, 4000, 8000, 10000, 12000, 18,000, 20000, 40,000, 80,000; 10^6 , 10^7 , 10^8 , 10^9 or more base pairs in length. Indeed, it is contemplated that the methods of the present invention will be able to create entire artificial genomes of lengths comparable to known bacterial, yeast, viral, mammalian, amphibian, reptilian, avian genomes. In more particular embodiments, the DNA segment is a gene encoding a protein of interest. The DNA segment further may include non-coding elements such as origins of replication, telomeres, promoters, enhancers, transcription and translation start and stop signals, introns, exon splice sites, chromatin scaffold components and other regulatory sequences. The DNA segment may comprise multiple genes, chromosomal segments, chromosomes and even entire genomes. The DNA segments may be derived from prokaryotic or eukaryotic sequences including bacterial, yeast, viral, mammalian, amphibian, reptilian, avian, plants, archbacteria and other DNA containing living organisms.

[0010] The oligonucleotide sets preferably are comprised oligonucleotides of between about 15 and 100 bases and more preferably between about 20 and 50 bases. Specific lengths include, but are not limited to 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 and 100. Depending on the size, the overlap between the oligonucleotides of the two sets may be designed to be between 5 and 75 bases per oligonucleotide pair.

[0011] The oligonucleotides preferably are treated with polynucleotide kinase, for example, T4 polynucleotide kinase. The kinasing can be performed prior to mixing of the oligonucleotides set or after, but before annealing. After annealing, the oligonucleotides are treated with an enzyme having a ligating function. For example, a DNA ligase typically will be employed for this function. However, topoisomerase, which does not require 5' phosphorylation, is rapid and operates at room temperature, and may be used instead of ligase.

[0012] In a second embodiment, there is provided a method for construction of a double-stranded DNA segment comprising the steps of (i) providing two sets of single-stranded oligonucleotides, wherein (a) the first set comprises the entire plus strand of said DNA segment, (b) the second set comprises the entire minus strand of said DNA segment,

and (c) each of said first set of oligonucleotides being complementary to two oligonucleotides of said second set of oligonucleotides, (ii) annealing pairs of complementary oligonucleotides to produce a set of first annealed products, wherein each pair comprises an oligonucleotide from each of said first and said second sets of oligonucleotides, (iii) annealing pairs of first annealed products having complementary sequences to produce a set of second annealed products, (iv) repeating the process until all annealed products have been annealed into a single DNA segment, and (v) treating said annealed products with ligating enzyme.

[0013] In a third embodiment, there is provided a method for the construction of a double-stranded DNA segment comprising the steps of (i) providing two sets of single-stranded oligonucleotides, wherein (a) the first set comprises the entire plus strand of said DNA segment, (b) the second set comprises the entire minus strand of said DNA segment, and (c) each of said first set of oligonucleotides being complementary to two oligonucleotides of said second set of oligonucleotides, (ii) annealing said the 5' terminal oligonucleotide of said first set of oligonucleotide with the 3' terminal oligonucleotide of said second set of oligonucleotides, (iii) annealing the next most 5' terminal oligonucleotide of said first set of oligonucleotides with the product of step (ii), (iv) annealing the next most 3' terminal oligonucleotide of said second set of oligonucleotides with the product of step (iii), (v) repeating the process until all oligonucleotides of said first and said second sets have been annealed, and (vi) treating said annealed oligonucleotides with ligating enzyme. Optional steps provide for the synthesis of the oligonucleotide sets and the transformation of host cells with the resulting DNA segment. In a preferred embodiment, the 5' terminal oligonucleotide of the first set is attached to a support, which process may include the additional step of removing the DNA segment from the support. The support may be any support known in the art, for example, a microtiter plate, a filter, polystyrene beads, polystyrene tray, magnetic beads, agarose and the like.

[0014] Annealing conditions may be adjusted based on the particular strategy used for annealing, the size and composition of the oligonucleotides, and the extent of overlap between the oligonucleotides of the first and second sets. For example, where all the oligonucleotides are mixed together prior to annealing, heating the mixture to 80° C., followed by slow annealing for between 1 to 12 h is conducted. Thus, annealing may be conducted for about 2, about 3, about 4, about 5, about 6, about 7, about 8, about 9, or about 10 h. However, in other embodiments, the annealing time may be as long as 24 h.

[0015] With the aid of a computer, the inventor is able to direct synthesis of a vector/gene combination using a high throughput oligonucleotide synthesizer as a set of overlapping component oligonucleotides. The oligonucleotides are assembled using a robotic combinatoric assembly strategy and the assembly ligated using DNA ligase or topoisomerase, followed by transformation into a suitable host strain. In a particular embodiment, this invention generates a set of bacterial strains containing a viable expression vector for all genes in a defined region of the genome. In other embodiments, a yeast or baculovirus expression vector system is also contemplated to allow expression of each gene in a chromosomal region in a eukaryotic host. In yet another embodiment, it the present invention allows one of

skill in the art to devise a "designer gene" strategy wherein a gene or genomes or virtually any structure may be readily designed, synthesized and expressed. Thus, eventually the technology described herein may be employed to create entire genomes for introduction into host cells for the creation of entirely artificial designer living organisms.

[0016] In specific embodiments, the present invention provides a method for the synthesis of a replication-competent, double-stranded polynucleotide, wherein the polynucleotide comprises an origin of replication, a first coding region and a first regulatory element directing the expression of the first coding region.

[0017] Additionally the method may further comprise the step of amplifying the double-stranded polynucleotide. In specific embodiments, the double-stranded polynucleotide comprises 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 5000, 10×10^3 , 20×10^3 , 30×10^3 , 40×10^3 , 50×10^3 , 60×10^3 , 70×10^3 , 80×10^3 , 90×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 or 1×10^{10} base pairs in length. The first regulatory element may be a promoter. In certain embodiments, the double-stranded polynucleotide further comprises a second regulatory element, the second regulatory element being a polyadenylation signal. In yet further embodiments, the double-stranded polynucleotide comprises a plurality of coding regions and a plurality of regulatory elements. Specifically, it is contemplated that the coding regions encode products that comprise a biochemical pathway. In particular embodiments the biochemical pathway is glycolysis. More particularly, it is contemplated that the coding regions encode enzymes selected from the group consisting of hexokinase, phosphohexose isomerase, phosphofructokinase-1, aldolase, triose-phosphate isomerase, glyceraldehyde-3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase and pyruvate kinase enzymes of the glycolytic pathway.

[0018] In other embodiments, the biochemical pathway is lipid synthesis, cofactor synthesis. Particularly contemplated are synthesis of lipoic acid, riboflavin synthesis nucleotide synthesis the nucleotide may be a purine or a pyrimidine.

[0019] In certain other embodiments it is contemplated that the coding regions encode enzymes involved in a cellular process selected from the group consisting of cell division, chaperone, detoxification, peptide secretion, energy metabolism, regulatory function, DNA replication, transcription, RNA processing and tRNA modification. In preferred embodiments, the energy metabolism is oxidative phosphorylation.

[0020] It is contemplated that the double-stranded polynucleotide is a DNA or an RNA. In preferred embodiments, the double-stranded polynucleotide may be a chromosome. The double-stranded polynucleotide may be an expression construct. Specifically, the expression construct may be a bacterial expression construct, a mammalian expression construct or a viral expression construct. In particular embodiments, the double-stranded polynucleotide comprises a genome selected from the group consisting of bacterial genome, yeast genome, viral genome, mammalian genome, amphibian genome and avian genome.

[0021] In those embodiments in which the genome is a viral genome, the viral genome may be selected from the group consisting of retrovirus, adenovirus, vaccinia virus, herpesvirus and adeno-associated virus.

[0022] The present invention further provides a method of producing a viral particle.

[0023] Another embodiment provides a method of producing an artificial genome, wherein the chromosome comprises all coding regions and regulatory elements found in a corresponding natural chromosome. In specific embodiments, the corresponding natural chromosome is a human mitochondrial genome. In other embodiments, the corresponding natural chromosome is a chloroplast genome.

[0024] Also provided is a method of producing an artificial genetic system, wherein the system comprises all coding regions and regulatory elements found in a corresponding natural biochemical pathway. Such a biochemical pathway will likely possess a group of enzymes that serially metabolize a compound. In particularly preferred embodiments, the biochemical pathway comprises the activities required for glycolysis. In other embodiments, the biochemical pathway comprises the enzymes required for electron transport. In still further embodiments, the biochemical pathway comprises the enzyme activities required for photosynthesis.

[0025] Other objects, features and advantages of the present invention will become apparent from the following detailed description. It should be understood, however, that the detailed description and the specific examples, while indicating preferred embodiments of the invention, are given by way of illustration only, since various changes and modifications within the spirit and scope of the invention will become apparent to those skilled in the art from this detailed description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0026] The following drawings form part of the present specification and are included to further demonstrate certain aspects of the present invention. The invention may be better understood by reference to one or more of these drawings in combination with the detailed description of specific embodiments presented herein.

[0027] **FIG. 1.** Flow diagram of the Jurassic Park paradigm for the reassembly of living organisms.

[0028] **FIG. 2.** Flow diagram of the strategy of synthetic genetics.

[0029] **FIG. 3.** Flow diagram of the strategy for combinatoric assembly of oligonucleotides into complete genes or genomes.

[0030] **FIG. 4.** Design of plasmid synlux4. The sequence of 4800 is annotated with the locations of lux A+B genes, neomycin/kanamycin phosphotransferase and pUC 19 sequences.

[0031] **FIG. 5.** List of component oligonucleotides derived from the sequence of Synlux4 in **FIG. 4.**

[0032] **FIG. 6.** Schema for the combinatoric assembly of synthetic plasmids from component oligonucleotides.

[0033] **FIG. 7.** SynGene program for generating overlapping oligonucleotides sufficient to reassemble the gene or plasmid.

DESCRIPTION OF ILLUSTRATIVE EMBODIMENTS

[0034] The complete sequence of complex genomes, including the human genome, make large scale functional

approaches to genetics possible. The present invention outlines a novel approach to utilizing the results of genomic sequence information by computer directed gene synthesis based on computing on the human genome database. Specifically, the invention describes chemical synthesis and resynthesis of genes for transfer of these genes into a suitable host cells.

[0035] The present invention provides methods that can be used to synthesize de novo, DNA segments that encode sets of genes, either naturally occurring genes expressed from natural or artificial promoter constructs or artificial genes derived from synthetic DNA sequences, which encodes elements of biological systems that perform a specified function or attribution of an artificial organism as well as entire genomes. In producing such systems and genomes, the present invention provides the synthesis of a replication-competent, double-stranded polynucleotide, wherein the polynucleotide has an origin of replication, a first coding region and a first regulatory element directing the expression of the first coding region. By replication competent, it is meant that the polynucleotide is capable of directing its own replication. Thus, it is envisioned that the polynucleotide will possess all the cis-acting signals required to facilitate its own synthesis. In this respect, the polynucleotide will be similar to a plasmid or a virus, such that once placed within a cell, it is able to be replicated by a combination of the polynucleotide's and cellular functions.

[0036] Thus, using the techniques of the present invention, one of skill in the art can create an artificial genome that is capable of encoding all the activities required for sustaining its own existence. Also contemplated are artificial genetic systems that are capable of encoding enzymes and activities of a particular biochemical pathway. In such a system, it will be desirable to have all the activities present such that the whole biochemical pathway will operate. The co-expression of a set of enzymes required for a particular pathway constitutes a complete genetic or biological system. For example, the co-expression of the enzymes involved in glycolysis constitutes a complete genetic system for the production of energy in the form of ATP from glucose. Such systems for energy production may include groups of enzymes which naturally or artificially serially metabolize a set of compounds.

[0037] The types of biochemical pathways would include but are not limited to those for the biosynthesis of cofactors prosthetic groups and carriers (lipoate synthesis, riboflavin synthesis pyridine nucleotide synthesis); the biosynthesis of the cell envelopes (membranes, lipoproteins, porins, surface polysaccharides, lipopolysaccharides, antigens and surface structures); cellular processes including cell division, chaperones, detoxification, protein secretion, central intermediary metabolism (energy production vi phosphorus compounds and other); energy metabolism including aerobic, anaerobic, ATP proton motive force interconversions, electron transport, glycolysis triose phosphate pathway, pyruvate dehydrogenase, sugar metabolism; purine, pyrimidine nucleotide synthesis, including 2'deoxyribonucleotide synthesis, nucleotide and nucleoside interconversion, salvage of nucleoside and nucleotides, sugar-nucleotide biosynthesis and conversion; regulatory functions including transcriptional and translational controls, DNA replication including degradation of DNA, DNA replication, restriction modification, recombination and repair; transcription including

degradation of DNA, DNA-dependent RNA polymerase and transcription factors; RNA processing; translation including amino acyl tRNA synthetases, degradation of peptides and glycopeptides, protein modification, ribosome synthesis and modification, tRNA modification; translation factors transport and binding proteins including amino acid, peptide, amine carbohydrate, organic alcohol, organic acid and cation transport; and other systems for the adaptation, specific function or survival of an artificial organism.

[0038] A. Definitions

[0039] DNA segment—a linear piece of DNA having a double-stranded region and both 5'- and 3'-ends; the segment may be of any length sufficiently long to be created by the hybridization of at least two oligonucleotides having complementary regions.

[0040] Oligonucleotides—small DNA segments, single-stranded or double-stranded, comprised of the nucleotide bases A, T, G and C linked through phosphate bonds; oligonucleotides typically range from about 10 to 100 base pairs.

[0041] Plus strand—by convention, the single-strand of a double-stranded DNA that starts with the 5' end to the left as one reads the sequence.

[0042] Minus strand—by convention, the single-strand of a double-stranded DNA that starts with the 3' end to the left as one reads the sequence.

[0043] Complementary—where two nucleic acids have at least a portion of their sequences, when read in opposite (5'→3'; 3'→5') direction, that pair sequential nucleotides in the following fashion: A-T, G-C, T-A, G-C.

[0044] Oligonucleotide sets—a plurality of oligonucleotides that, taken together, comprise the sequence of a plus or minus strand of a DNA segment.

[0045] Annealed products—two or more oligonucleotides having complementary regions, where they are permitted, under proper conditions, to base pair, thereby producing double stranded regions.

[0046] B. The Present Invention

[0047] The present invention describes methods for enabling the creation of DNA molecules, genomes and entire artificial living organisms based upon information only, without the requirement for existing genes, DNA molecules or genomes.

[0048] The methods of the present invention are diagrammed in **FIG. 1** and **FIG. 2** and generally involve the following steps. Generally, using simple computer software, comprising sets of gene parts and functional elements it is possible to construct a virtual polynucleotide in the computer. This polynucleotide consists of a string of DNA bases, G, A, T or C, comprising for example an entire artificial genome in a linear string. For transfer of the synthetic gene into for example, bacterial cells the polynucleotide should contain the sequence for a bacterial (such as pBR322) origin of replication. For transfer into eukaryotic cells, it should contain the origin of replication of a mammalian virus, chromosome or subcellular component such as mitochondria.

[0049] Following construction, simple computer software is then used to break down the genome sequence into a set of overlapping oligonucleotides of specified length. This results in a set of shorter DNA sequences which overlap to cover the entire genome in overlapping sets. Typically, a gene of 1000 bases pairs would be broken down into 20 100-mers where 10 of these comprise one strand and 10 of these comprise the other strand. They would be selected to overlap on each strand by 25 to 50 base pairs.

[0050] This step is followed by direction of chemical synthesis of each of the overlapping set of oligonucleotides using an array type synthesizer and phosphoramidite chemistry resulting in an array of synthesized oligomers. The next step is to balance concentration of each oligomer and pool the oligomers so that a single mixture contains equal concentrations of each. The mixed oligonucleotides are treated with T4 polynucleotide kinase to 5' phosphorylate the oligonucleotides. The next step is to carry out a “slow” annealing step to co-anneal all of the oligomers into the sequence of the predicted gene or genome. This is done by heating the mixture to 80° C., then allowing it to cool slowly to room temperature over several hours. The mixture of oligonucleotides is then treated with T4 DNA ligase (or alternatively topoisomerase) to join the oligonucleotides. The oligonucleotides are then transferred into competent host cells.

[0051] The above technique represents a “combinatorial” assembly strategy where all oligonucleotides are jointly co-annealed by temperature-based slow annealing. A variation on this strategy, which may be more suitable for very long genes or genomes, such as greater than 5,000 base pairs final length, is as follows. Using simple computer software, comprising sets of gene parts and functional elements, a virtual gene or genome is constructed in the computer. This gene or genome would consist of a string of DNA bases, G, A, T or C, comprising the entire genome in a linear string. For transfer of the synthetic gene into bacterial cells, it should contain the sequence for a bacterial (such as pBR322) origin of replication.

[0052] The next step is to carry out a ligation chain reaction using a new oligonucleotide addition each step. With this procedure, the first oligonucleotide in the chain is attached to a solid support (such as an agarose bead). The second is added along with DNA ligase, and annealing and ligation reaction carried out, and the beads are washed. The second, overlapping oligonucleotide from the opposite strand is added, annealed and ligation carried out. The third oligonucleotide is added and ligation carried out. This procedure is replicated until all oligonucleotides are added and ligated. This procedure is best carried out for long sequences using an automated device. The DNA sequence is removed from the solid support, a final ligation (is circular) is carried out, and the molecule transferred into host cells.

[0053] Alternatively, it is contemplated that if the ligation kinetics allow all the oligonucleotides may be placed in a mixture and ligation be allowed to proceed. In yet another embodiment, a series of smaller polynucleotides may be made by ligating 2, 3, 4, 5, 6, or 7 oligonucleotides into one sequence and adding this to another sequence comprising a similar number of oligonucleotides parts.

[0054] The ligase chain reaction (“LCR”), disclosed in EPO No. 320 308, is incorporated herein by reference in its

entirety. In LCR, two complementary probe pairs are prepared, and in the presence of the target sequence, each pair will bind to opposite complementary strands of the target such that they abut. In the presence of a ligase, the two probe pairs will link to form a single unit. By temperature cycling, as in PCR™, bound ligated units dissociate from the target and then serve as “target sequences” for ligation of excess probe pairs. U.S. Pat. No. 4,883,750 describes a method similar to LCR for binding probe pairs to a target sequence. The following sections describe these methods in further detail.

[0055] C. Nucleic Acids

[0056] The present invention discloses the artificial synthesis of genes. In one embodiment of the present invention, the artificial genes can be transferred into cells to confer a particular function either as discrete units or as part of artificial chromosomes or genome. One will generally prefer to design oligonucleotides having stretches of 15 to 100 nucleotides, 25 to 200 nucleotides or even longer where desired. Such fragments may be readily prepared by, directly synthesizing the fragment by chemical means as described below.

[0057] Accordingly, the nucleotide sequences of the invention may be used for their ability to selectively form duplex molecules with complementary stretches of genes or RNAs or to provide primers for amplification of DNA or RNA from tissues. Depending on the application envisioned, one will desire to employ varying conditions of hybridization to achieve varying degrees of hybridization selectivity. Typically high selectivity is favored.

[0058] For applications requiring high selectivity, one typically will desire to employ relatively stringent conditions to form the hybrids, e.g., one will select relatively low salt and/or high temperature conditions, such as provided by about 0.02 M to about 0.10 M NaCl at temperatures of about 50° C. to about 70° C. Such high stringency conditions tolerate little, if any, mismatch between the oligonucleotide and the template or target strand. It generally is appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide.

[0059] For certain applications, for example, by analogy to, substitution of nucleotides by site-directed mutagenesis, it is appreciated that lower stringency conditions may be used. Under these conditions, hybridization may occur even though the sequences of probe and target strand are not perfectly complementary, but are mismatched at one or more positions. Conditions may be rendered less stringent by increasing salt concentration and decreasing temperature. For example, a medium stringency condition could be provided by about 0.1 to 0.25 M NaCl at temperatures of about 37° C. to about 55° C., while a low stringency condition could be provided by about 0.15 M to about 0.9 M salt, at temperatures ranging from about 20° C. to about 55° C. Thus, hybridization conditions can be readily manipulated depending on the desired results.

[0060] In certain embodiments, it will be advantageous to determine the hybridization of oligonucleotides by employing as a label. A wide variety of appropriate indicator means are known in the art, including fluorescent, radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of being detected. In preferred embodiments, one

may desire to employ a fluorescent label or an enzyme tag such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other environmentally undesirable reagents. In the case of enzyme tags, colorimetric indicator substrates are known that can be employed to provide a detection means visible to the human eye or spectrophotometrically, to identify whether specific hybridization with complementary oligonucleotide has occurred.

[0061] In embodiments involving a solid phase, for example the first oligonucleotide is adsorbed or otherwise affixed to a selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to hybridization with the complementary oligonucleotides under desired conditions. The selected conditions will also depend on the particular circumstances based on the particular criteria required (depending, for example, on the G+C content, type of target nucleic acid, source of nucleic acid, size of hybridization probe, etc.). Following washing of the hybridized surface to remove non-specifically bound oligonucleotides, the hybridization may be detected, or even quantified, by means of the label.

[0062] For applications in which the nucleic acid segments of the present invention are incorporated into vectors, such as plasmids, cosmids or viruses, these segments may be combined with other DNA sequences, such as promoters, polyadenylation signals, restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol.

[0063] DNA segments encoding a specific gene may be introduced into recombinant host cells and employed for expressing a specific structural or regulatory protein. Alternatively, through the application of genetic engineering techniques, subportions or derivatives of selected genes may be employed. Upstream regions containing regulatory regions such as promoter regions may be isolated and subsequently employed for expression of the selected gene.

[0064] The nucleic acids employed may encode antisense constructs that hybridize, under intracellular conditions, to a nucleic acid of interest. The term “antisense construct” is intended to refer to nucleic acids, preferably oligonucleotides, that are complementary to the base sequences of a target DNA. Antisense oligonucleotides, when introduced into a target cell, specifically bind to their target nucleic acid and interfere with transcription, RNA processing, transport, translation and/or stability. Antisense constructs may be designed to bind to the promoter and other control regions, exons, introns or even exon-intron boundaries of a gene.

[0065] Other sequences with lower degrees of homology also are contemplated. For example, an antisense construct which has limited regions of high homology, but also contains a non-homologous region (e.g., a ribozyme) could be designed. These molecules, though having less than 50% homology, would bind to target sequences under appropriate conditions.

[0066] In certain embodiments, one may wish to employ antisense constructs which include other elements, for example, those which include C-5 propyne pyrimidines.

Oligonucleotides which contain C-5 propyne analogues of uridine and cytidine have been shown to bind RNA with high affinity and to be potent antisense inhibitors of gene expression (Wagner et al., 1993).

[0067] According to the present invention, DNA segments of a variety of sizes will be produced. These DNA segments will, by definition, be linear molecules. As such, they typically will be modified before further use. These modifications include, in one embodiment, the restriction of the segments to produce one or more “sticky ends” compatible with complementary ends of other molecules, including those in vectors capable of supporting the replication of the DNA segment. This manipulation facilitates “cloning” of the segments.

[0068] Typically, cloning involves the use of restriction endonucleases, which cleave at particular sites within DNA strands, to prepare a DNA segment for transfer into a cloning vehicle. Ligation of the compatible ends (which include blunt ends) using a DNA ligase completes the reaction. Depending on the situation, the cloning vehicle may comprise a relatively small portion of DNA, compared to the insert. Alternatively, the cloning vehicle may be extremely complex and include a variety of features that will affect the replication and function of the DNA segment. In certain embodiments, a rare cutter site may be introduced into the end of the polynucleotide sequence.

[0069] Cloning vehicles include plasmids such as the pUC series, Bluescript™ vectors and a variety of other vehicles with multipurpose cloning sites, selectable markers and origins of replication. Because of the nature of the present invention, the cloning vehicles may include such complex molecules as phagemids and cosmids, which hold relatively large pieces of DNA. In addition, the generation of artificial chromosomes, and even genomes.

[0070] Following cloning into a suitable vector, the construct then is transferred into a compatible host cell. A variety of different gene transfer techniques are described elsewhere in this document. Culture of the host cells for the intended purpose (amplification, expression, subcloning) follows.

[0071] Throughout this application, the term “expression construct” is meant to include a particular kind of cloning vehicle containing a nucleic acid coding for a gene product in which part or all of the nucleic acid encoding sequence is capable of being transcribed. The transcript may be translated into a protein, but it need not be. Thus, in certain embodiments, expression includes both transcription of a gene and translation of a RNA into a gene product. In other embodiments, expression only includes transcription of the nucleic acid, for example, to generate antisense constructs.

[0072] In preferred embodiments, the nucleic acid is under transcriptional control of a promoter. A “promoter” refers to a DNA sequence recognized by the synthetic machinery of the cell, or introduced synthetic machinery, required to initiate the specific transcription of a gene. The phrase “under transcriptional control” means that the promoter is in the correct location and orientation in relation to the nucleic acid to control RNA polymerase initiation and expression of the gene.

[0073] The term promoter will be used here to refer to a group of transcriptional control modules that are clustered

around the initiation site for RNA polymerase II. Much of the thinking about how promoters are organized derives from analyses of several viral promoters, including those for the HSV thymidine kinase (tk) and SV40 early transcription units. These studies, augmented by more recent work, have shown that promoters are composed of discrete functional modules, each consisting of approximately 7-20 bp of DNA, and containing one or more recognition sites for transcriptional activator or repressor proteins.

[0074] At least one module in each promoter functions to position the start site for RNA synthesis. The best known example of this is the TATA box, but in some promoters lacking a TATA box, such as the promoter for the mammalian terminal deoxynucleotidyl transferase gene and the promoter for the SV40 late genes, a discrete element overlying the start site itself helps to fix the place of initiation.

[0075] Additional promoter elements regulate the frequency of transcriptional-initiation.

[0076] Typically, these are located in the region 30-110 bp upstream of the start site, although a number of promoters have recently been shown to contain functional elements downstream of the start site as well. The spacing between promoter elements frequently is flexible, so that promoter function is preserved when elements are inverted or moved relative to one another. In the tk promoter, the spacing between promoter elements can be increased to 50 bp apart before activity begins to decline. Depending on the promoter, it appears that individual elements can function either co-operatively or independently to activate transcription.

[0077] The particular promoter that is employed to control the expression of a nucleic acid is not believed to be critical, so long as it is capable of expressing the nucleic acid in the targeted cell. Thus, where a human cell is targeted, it is preferable to position the nucleic acid coding region adjacent to and under the control of a promoter that is capable of being expressed in a human cell. Generally speaking, such a promoter might include either a human or viral promoter. Preferred promoters include those derived from HSV. Another preferred embodiment is the tetracycline controlled promoter.

[0078] In various other embodiments, the human cytomegalovirus (CMV) immediate early gene promoter, the SV40 early promoter and the Rous sarcoma virus long terminal repeat can be used to obtain high-level expression of transgenes. The use of other viral or mammalian cellular or bacterial phage promoters which are well-known in the art to achieve expression of a transgene is contemplated as well, provided that the levels of expression are sufficient for a given purpose. It is envisioned that any elements/promoters may be employed in the context of the present invention. Below is a list of viral promoters, cellular promoters/enhancers and inducible promoters/enhancers that could be used in combination with the nucleic acid encoding a gene of interest in an expression construct. Enhancer/promoter elements contemplated for use with the present invention include but are not limited to Immunoglobulin Heavy Chain, Immunoglobulin Light, Chain T-Cell Receptor, HLA DQ α and DQ β , β -Interferon, Interleukin-2, Interleukin-2 Receptor, MHC Class II 5, MHC Class II HLA-DR α , -Actin, Muscle Creatine Kinase, Prealbumin (Transthyretin), Elastase I, Metallothionein, Collagenase, Albumin Gene, α -Fetoprotein, τ -Globin, β -Globin, e-fos, c-HA-ras, Insulin,

Neural Cell Adhesion Molecule (NCAM), α 1-Antitrypsin, H₂B (TH2B) Histone, Mouse or Type I Collagen, Glucose-Regulated Proteins (GRP94 and GRP78), Rat Growth Hormone, Human Serum Amyloid A (SAA), Troponin I (TN I), Platelet-Derived Growth Factor, Duchenne Muscular Dystrophy, SV40, Polyoma, Retroviruses, Papilloma Virus, Hepatitis B Virus, Human Immunodeficiency Virus, Cytomegalovirus, Gibbon Ape Leukemia Virus. Inducible promoter elements and their associated inducers are listed in Table 2 below. This list is not intended to be exhaustive of all the possible elements involved in the promotion of transgene expression but, merely, to be exemplary thereof. Additionally, any promoter/enhancer combination (as per the Eukaryotic Promoter Data Base EPDB) could also be used to drive expression of the gene. Eukaryotic cells can support cytoplasmic transcription from certain bacterial promoters if the appropriate bacterial polymerase is provided, either as part of the delivery complex or as an additional genetic expression construct.

[0079] Enhancers were originally detected as genetic elements that increased transcription from a promoter located at a distant position on the same molecule of DNA. This ability to act over a large distance had little precedent in classic studies of prokaryotic transcriptional regulation. Subsequent work showed that regions of DNA with enhancer activity are organized much like promoters. That is, they are composed of many individual elements, each of which binds to one or more transcriptional proteins.

[0080] The basic distinction between enhancers and promoters is operational. An enhancer region as a whole must be able to stimulate transcription at a distance; this need not be true of a promoter region or its component elements. On the other hand, a promoter must have one or more elements that direct initiation of RNA synthesis at a particular site and in a particular orientation, whereas enhancers lack these specificities. Promoters and enhancers are often overlapping and contiguous, often seeming to have a very similar modular organization.

TABLE 2

Element	Inducer
MT II	Phorbol Ester (TPA) Heavy metals
MMTV (mouse mammary tumor virus)	Glucocorticoids
β -Interferon	poly(rI)X poly(rc)
Adenovirus 5 E2	Ela
c-jun	Phorbol Ester (TPA), H ₂ O ₂
Collagenase	Phorbol Ester (TPA)
Stromelysin	Phorbol Ester (TPA), IL-1
SV40	Phorbol Ester (TPA)
Murine MX Gene	Interferon, Newcastle Disease Virus
GRP78 Gene	A23187
α -2-Macroglobulin	IL-6
Vimentin	Serum
MHC Class I Gene H-2kB	Interferon
HSP70	Ela, SV40 Large T Antigen
Proliferin	Phorbol Ester-TPA
Tumor Necrosis Factor	FMA
Thyroid Stimulating Hormone α Gene	Thyroid Hormone

[0081] Use of the baculovirus system will involve high level expression from the powerful polyhedron promoter.

One will typically include a polyadenylation signal to effect proper polyadenylation of the transcript. The nature of the polyadenylation signal is not believed to be crucial to the successful practice of the invention, and any such sequence may be employed. Preferred embodiments include the SV40 polyadenylation signal and the bovine growth hormone polyadenylation signal, convenient and known to function well in various target cells. Also contemplated as an element of the expression cassette is a terminator. These elements can serve to enhance message levels and to minimize read through from the cassette into other sequences.

[0082] A specific initiation signal also may be required for efficient translation of coding sequences. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may need to be provided. One of ordinary skill in the art would readily be capable of determining this and providing the necessary signals. It is well known that the initiation codon must be "in-frame" with the reading frame of the desired coding sequence to ensure translation of the entire insert. The exogenous translational control signals and initiation codons can be either natural or synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements (Bittner et al., 1987).

[0083] In certain embodiments, it may be desirable to include specialized regions known as telomeres at the end of a genome sequence. Telomeres are repeated sequences found at chromosome ends and it has long been known that chromosomes with truncated ends are unstable, tend to fuse with other chromosomes and are otherwise lost during cell division. Some data suggest that telomeres interaction the nucleoprotein complex and the nuclear matrix. One putative role for telomeres includes stabilizing chromosomes and shielding the ends from degradative enzyme.

[0084] Another possible role for telomeres is in replication. According to present doctrine, replication of DNA requires starts from short RNA primers annealed to the 3'-end of the template. The result of this mechanism is an "end replication problem" in which the region corresponding to the RNA primer is not replicated. Over many cell divisions, this will result in the progressive truncation of the chromosome. It is thought that telomeres may provide a buffer against this effect, at least until they are themselves eliminated by this effect. A further structure to be included in DNA segments is a centromere.

[0085] In certain embodiments of the invention, the delivery of a nucleic acid in a cell may be identified in vitro or in vivo by including a marker in the expression construct. The marker would result in an identifiable change to the transfected cell permitting easy identification of expression.

[0086] A number of selection systems may be used, including, but not limited, to the herpes simplex virus thymidine kinase (Wigler et al., 1977), hypoxanthine-guanine phosphoribosyltransferase (Szybalska et al., 1962) and adenine phosphoribosyltransferase genes (Lowy et al., 1980), in tk⁻, hgprt⁻ or aprt⁻ cells, respectively. Also, antimetabolite resistance can be used as the basis of selection for dhfr, which confers resistance to methotrexate (Wigler et al., 1980; O'Hare et al., 1981); gpt, which confers resistance to mycophenolic acid (Mulligan et al., 1981); neo,

which confers resistance to the aminoglycoside G-418 (Colberre-Garapin et al., 1981); and hygromycin, which confers resistance to hygromycin.

[0087] Usually the inclusion of a drug selection marker aids in cloning and in the selection of transformants, for example, neomycin, puromycin, hygromycin, DHFR, GPT, zeocin and histidinol. Alternatively, enzymes such as herpes simplex virus thymidine kinase (tk) (eukaryotic) or chloramphenicol acetyltransferase (CAT) (prokaryotic) may be employed. Immunologic markers also can be employed. The selectable marker employed is not believed to be important, so long as it is capable of being expressed simultaneously with the nucleic acid encoding a gene product. Further examples of selectable markers are well known to one of skill in the art.

[0088] In certain embodiments of the invention, the use of internal ribosome binding sites (IRES) elements are used to create multigene, or polycistronic, messages. IRES elements are able to bypass the ribosome scanning model of 5' methylated Cap dependent translation and begin translation at internal sites (Pelletier and Sonenberg, 1988). IRES elements from two members of the picornavirus family (polio and encephalomyocarditis) have been described (Pelletier and Sonenberg, 1988), as well an IRES from a mammalian message (Macejak and Sarnow, 1991). IRES elements can be linked to heterologous open reading frames. Multiple open reading frames can be transcribed together, each separated by an IRES, creating polycistronic messages. By virtue of the IRES element, each open reading frame is accessible to ribosomes for efficient translation. Multiple genes can be efficiently expressed using a single promoter/enhancer to transcribe a single message.

[0089] Any heterologous open reading frame can be linked to IRES elements. This includes genes for secreted proteins, multi-subunit proteins, encoded by independent genes, intracellular or membrane-bound proteins and selectable markers. In this way, expression of several proteins can be simultaneously engineered into a cell with a single construct and a single selectable marker.

[0090] D. Encoded Proteins

[0091] In this application, the inventors use genetic information for creative or synthetic purposes. The complete genome sequence will give a catalog of all genes necessary for the survival, reproduction, evolution and speciation of an organism and, given suitable high tech tools, the genomic information may be modified or even created from "scratch" in order to synthesize life. Thus it is contemplated that a combination of suitable energy generation genes, regulatory genes, and other functional genes could be constructed which would be sufficient to render an artificial organism with the basic functionalities to enable independent survival.

[0092] To meet this goal, the present invention utilizes known cDNA sequences for any given gene to express proteins in an artificial organism. Any protein so expressed in this invention may be modified for particular purposes according to methods well known to those of skill in the art. For example, particular peptide residues may be derivatized or chemically modified in order to alter the immune response or to permit coupling of the peptide to other agents. It also is possible to change particular amino acids within the peptides without disturbing the overall structure or antigen-

nicity of the peptide. Such changes are therefore termed "conservative" changes and tend to rely on the hydrophilicity or polarity of the residue. The size and/or charge of the side chains also are relevant factors in determining which substitutions are conservative.

[0093] Once the entire coding sequence of a gene has been determined, the gene can be inserted into an appropriate expression system. The gene can be expressed in any number of different recombinant DNA expression systems to generate large amounts of the polypeptide product, which can then be purified and used to vaccinate animals to generate antisera with which further studies may be conducted.

[0094] Examples of expression systems known to the skilled practitioner in the art include bacteria such as *E. coli*, yeast such as *Saccharomyces cerevisia* and *Pichia pastoris*, baculovirus, and mammalian expression systems such as in COS or CHO cells. In one embodiment, polypeptides are expressed in *E. coli* and in baculovirus expression systems. A complete gene can be expressed or, alternatively, fragments of the gene encoding portions of polypeptide can be produced.

[0095] In one embodiment, the gene sequence encoding the polypeptide is analyzed to detect putative transmembrane sequences. Such sequences are typically very hydrophobic and are readily detected by the use of standard sequence analysis software, such as MacVector (IBI, New Haven, Conn.). The presence of transmembrane sequences is often deleterious when a recombinant protein is synthesized in many expression systems, especially *E. coli*, as it leads to the production of insoluble aggregates that are difficult to renature into the native conformation of the protein. Deletion of transmembrane sequences typically does not significantly alter the conformation of the remaining protein structure.

[0096] Moreover, transmembrane sequences, being by definition embedded within a membrane, are inaccessible. Therefore, antibodies to these sequences will not prove useful for in vivo or in situ studies. Deletion of transmembrane-encoding sequences from the genes used for expression can be achieved by standard techniques. For example, fortuitously-placed restriction enzyme sites can be used to excise the desired gene fragment, or PCRTM-type amplification can be used to amplify only the desired part of the gene. The skilled practitioner will realize that such changes must be designed so as not to change the translational reading frame for downstream portions of the protein-encoding sequence.

[0097] In one embodiment, computer sequence analysis is used to determine the location of the predicted major antigenic determinant epitopes of the polypeptide. Software capable of carrying out this analysis is readily available commercially, for example MacVector (IBI, New Haven, Conn.). The software typically uses standard algorithms such as the Kyte/Doolittle or Hopp/Woods methods for locating hydrophilic sequences which are characteristically found on the surface of proteins and are, therefore, likely to act as antigenic determinants.

[0098] Once this analysis is made, polypeptides can be prepared that contain at least the essential features of the antigenic determinant and that can be employed in the

generation of antisera against the polypeptide. Minigenes or gene fusions encoding these determinants can be constructed and inserted into expression vectors by standard methods, for example, using PCR™ methodology.

[0099] The gene or gene fragment encoding a polypeptide can be inserted into an expression vector by standard subcloning techniques. In one embodiment, an *E. coli* expression vector is used that produces the recombinant polypeptide as a fusion protein, allowing rapid affinity purification of the protein. Examples of such fusion protein expression systems are the glutathione S-transferase system (Pharmacia, Piscataway, N.J.), the maltose binding protein system (NEB, Beverly, Mass.), the FLAG system (IBI, New Haven, Conn.), and the 6xHis system (Qiagen, Chatsworth, Calif.).

[0100] Some of these systems produce recombinant polypeptides bearing only a small number of additional amino acids, which are unlikely to affect the antigenic ability of the recombinant polypeptide. For example, both the FLAG system and the 6xHis system add only short sequences, both of that are known to be poorly antigenic and which do not adversely affect folding of the polypeptide to its native conformation. Other fusion systems produce polypeptide where it is desirable to excise the fusion partner from the desired polypeptide. In one embodiment, the fusion partner is linked to the recombinant polypeptide by a peptide sequence containing a specific recognition sequence for a protease. Examples of suitable sequences are those recognized by the Tobacco Etch Virus protease (Life Technologies, Gaithersburg, Md.) or Factor Xa (New England Biolabs, Beverly, Mass.).

[0101] Recombinant bacterial cells, for example *E. coli*, are grown in any of a number of suitable media, for example LB, and the expression of the recombinant polypeptide induced by adding IPTG to the media or switching incubation to a higher temperature. After culturing the bacteria for a further period of between 2 and 24 h, the cells are collected by centrifugation and washed to remove residual media. The bacterial cells are then lysed, for example, by disruption in a cell homogenizer and centrifuged to separate the dense inclusion bodies and cell membranes from the soluble cell components. This centrifugation can be performed under conditions whereby the dense inclusion bodies are selectively enriched by incorporation of sugars such as sucrose into the buffer and centrifugation at a selective speed.

[0102] In another embodiment, the expression system used is one driven by the baculovirus polyhedron promoter. The gene encoding the polypeptide can be manipulated by standard techniques in order to facilitate cloning into the baculovirus vector. One baculovirus vector is the pBlueBac vector (Invitrogen, Sorrento, Calif.). The vector carrying the gene for the polypeptide is transfected into *Spodoptera frugiperda* (Sf9) cells by standard protocols, and the cells are cultured and processed to produce the recombinant antigen. See Summers et al., A MANUAL OF METHODS FOR BACULOVIRUS VECTORS AND INSECT CELL CULTURE PROCEDURES, Texas Agricultural Experimental Station.

[0103] In designing a gene that encodes a particular polypeptide, the hydrophatic index of amino acids may be considered. Table 3 provides a codon table showing the nucleic acids that encode a particular amino acid. The

importance of the hydrophatic amino acid index in conferring interactive biologic function on a protein is generally understood in the art (Kyte & Doolittle, 1982). The following is a brief discussion of the the hydrophatic amino acid index for use in the present invention.

TABLE 3

Amino Acids		Codons				
Alanine	Ala	A	GCA	GCC	GCG	GCU
Cysteine	Gys	C	UGC	UGU		
Aspartic acid	Asp	D	GAC	GAU		
Glutamic acid	Glu	E	GAA	GAG		
Phenylalanine	Phe	F	UUC	UUU		
Glycine	Gly	G	GGA	GGC	GGG	GGU
Histidine	His	H	CAC	CAU		
Isoleucine	Ile	I	AUA	AUG	AUU	
Lysine	Lys	K	AAA	AAG		
Leucine	Leu	L	UUA	UUG	CUA	CUC
Methionine	Met	M	AUG			
Asparagine	Asn	N	AAG	AAU		
Proline	Pro	P	CCA	CCC	CCG	CCU
Glutamine	Gln	Q	CAA	CAG		
Arginine	Arg	R	AGA	AGG	CGA	CGC
Serine	Ser	S	AGC	AGU	UGA	UCC
Threonine	Thr	T	ACA	ACC	ACG	ACU
Valine	Val	V	GUA	GUC	GUG	GUU
Tryptophan	Trp	W	UGG			
Tyrosine	Tyr	Y	UAC	UAU		

[0104] It is accepted that the relative hydrophatic character of the amino acid contributes to the secondary structure of the resultant protein, which in turn defines the interaction of the protein with other molecules, for example, enzymes, substrates, receptors, DNA, antibodies, antigens, and the like.

[0105] Each amino acid has been assigned a hydrophatic index on the basis of their hydrophobicity and charge characteristics (Kyte & Doolittle, 1982), these are: Isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5);

[0106] asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

[0107] It is known in the art that certain amino acids may be substituted by other amino acids having a similar hydrophatic index or score and still result in a protein with similar biological activity, i.e., still obtain a biological functionally equivalent protein. In making such changes, the substitution of amino acids whose hydrophatic indices are within ± 2 is preferred, those which are within +1 are particularly preferred, and those within +0.5 are even more particularly preferred.

[0108] It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Pat. No. 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with a biological property of the protein.

[0109] As detailed in U.S. Pat. No. 4,554,101, the following hydrophilicity values have been assigned to amino acid

residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0±1); glutamate (+3.0±1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5±1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

[0110] It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent and immunologically equivalent protein. In such changes, the substitution of amino acids whose hydrophilicity values are within ±2 is preferred, those that are within ±1 are particularly preferred, and those within ±0.5 are even more particularly preferred.

[0111] As outlined above, amino acid substitutions are generally based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. Exemplary substitutions that take various of the foregoing characteristics into consideration are well known to those of skill in the art and include: arginine and lysine; glutamate and aspartate; serine and threonine; glutamine and asparagine; and valine, leucine and isoleucine.

[0112] E. Expression of and Delivery of Genes

[0113] I. Expression

[0114] Once the designer gene, genome or biological system has been made according the methods described herein, the polynucleotides can be expressed as encoded peptides or proteins of the gene, genome or biological system. The engineering of the polynucleotides for expression in a prokaryotic or eukaryotic system may be performed by techniques generally known to those of skill in recombinant expression. Therefore, promoters and other elements specific to a bacterial mammalian or other system may be included in the polynucleotide sequence. It is believed that virtually any expression system may be employed in the expression of the claimed nucleic acid sequences.

[0115] The artificially generated polynucleotide sequences are suitable for eukaryotic expression, as the host cell will generally process the genomic transcripts to yield functional mRNA for translation into protein. It is believed that the use of a designer gene version will provide advantages in that the size of the gene will generally be much smaller and more readily employed to transfect the targeted cell than will a genomic gene, which will typically be up to an order of magnitude larger than the designer gene. However, the inventor does not exclude the possibility of employing a genomic version of a particular gene where desired.

[0116] As used herein, the terms “engineered” and “recombinant” cells are intended to refer to a cell into which an exogenous polynucleotide described herein has been introduced. Therefore, engineered cells are distinguishable from naturally-occurring cells which do not contain a recombinantly introduced exogenous polynucleotide. Engineered cells are thus cells having a gene or genes introduced through the hand of man. Recombinant cells include those having an introduced polynucleotides, and also include polynucleotides positioned adjacent to a promoter not naturally associated with the particular introduced gene.

[0117] To express a recombinant encoded protein or peptide, whether mutant or wild-type, in accordance with the

present invention one would prepare an expression vector that comprises one of the claimed isolated nucleic acids under the control of one or more promoters. To bring a coding sequence “under the control of” a promoter, one positions the 5' end of the translational initiation site of the reading frame generally between about 1 and 50 nucleotides “downstream” of (i.e., 3' of) the chosen promoter. The “upstream” promoter stimulates transcription of the inserted DNA and promotes expression of the encoded recombinant protein. This is the meaning of “recombinant expression” in the context used here.

[0118] Many standard techniques are available to construct expression vectors containing the appropriate nucleic acids and transcriptional/translational control sequences in order to achieve protein or peptide expression in a variety of host-expression systems. Cell types available for expression include, but are not limited to, bacteria, such as *E. coli* and *B. subtilis* transformed with recombinant phage DNA, plasmid DNA or cosmid DNA expression vectors.

[0119] Certain examples of prokaryotic hosts are *E. coli* strain RRI, *E. coli* LE392, *E. coli* B, *E. coli* χ 1776(ATCC No. 31537) as well as *E. coli* W3110 (F—, lambda-, prototrophic, ATCC No. 273325); bacilli such as *Bacillus subtilis*; and other enterobacteriaceae such as *Salmonella typhimurium*, *Serratia marcescens*, and various *Pseudomonas* species.

[0120] In general, plasmid vectors containing replicon and control sequences that are derived from species compatible with the host cell are used in connection with these hosts. The vector ordinarily carries a replication site, as well as marking sequences that are capable of providing phenotypic selection in transformed cells. For example, *E. coli* is often transformed using pBR322, a plasmid derived from an *E. coli* species. Plasmid pBR322 contains genes for ampicillin and tetracycline resistance and thus provides easy means for identifying transformed cells. The pBR322 plasmid, or other microbial plasmid or phage must also contain, or be modified to contain, promoters that can be used by the microbial organism for expression of its own proteins.

[0121] In addition, phage vectors containing replicon and control sequences that are compatible with the host microorganism can be used as transforming vectors in connection with these hosts. For example, the phage lambda GEM™-11 may be utilized in making a recombinant phage vector that can be used to transform host cells, such as *E. coli* LE392.

[0122] Further useful vectors include pIN vectors (Inouye et al., 1985); and pGEX vectors, for use in generating glutathione S-transferase (GST) soluble fusion proteins for later purification and separation or cleavage. Other suitable fusion proteins are those with β -galactosidase, ubiquitin, or the like.

[0123] Promoters that are most commonly used in recombinant DNA construction include the lactamase (penicillinase), lactose and tryptophan (trp) promoter systems. While these are the most commonly used, other microbial promoters have been discovered and utilized, and details concerning their nucleotide sequences have been published, enabling those of skill in the art to ligate them functionally with plasmid vectors.

[0124] For expression in *Saccharomyces*, the plasmid YRp7, for example, is commonly used (Stinchcomb et al.,

1979; Kingsman et al., 1979; Tschemper et al., 1980). This plasmid contains the *trp1* gene, which provides a selection marker for a mutant strain of yeast lacking the ability to grow in tryptophan, for example ATCC No. 44076 or PEP4-1 (Jones, 1977). The presence of the *trp1* lesion as a characteristic of the yeast host cell genome then provides an effective environment for detecting transformation by growth in the absence of tryptophan.

[0125] Suitable promoting sequences in yeast vectors include the promoters for 3-phosphoglycerate kinase (Hitzman et al., 1980) or other glycolytic enzymes (Hess et al., 1968; Holland et al., 1978), such as enolase, glyceraldehyde-3-phosphate dehydrogenase, hexokinase, pyruvate decarboxylase, phosphofructokinase, glucose-6-phosphate isomerase, 3-phosphoglycerate mutase, pyruvate kinase, triosephosphate isomerase, phosphoglucose isomerase, and glucokinase. In constructing suitable expression plasmids, the termination sequences associated with these genes are also ligated into the expression vector 3' of the sequence desired to be expressed to provide polyadenylation of the mRNA and termination.

[0126] Other suitable promoters, which have the additional advantage of transcription controlled by growth conditions, include the promoter region for alcohol dehydrogenase 2, isocytochrome C, acid phosphatase, degradative enzymes associated with nitrogen metabolism, and the aforementioned glyceraldehyde-3-phosphate dehydrogenase, and enzymes responsible for maltose and galactose utilization.

[0127] In addition to micro-organisms, cultures of cells derived from multicellular organisms may also be used as hosts. In principle, any such cell culture is workable, whether from vertebrate or invertebrate culture. In addition to mammalian cells, these include insect cell systems infected with recombinant virus expression vectors (e.g., baculovirus); and plant cell systems infected with recombinant virus expression vectors (e.g., cauliflower mosaic virus, CAMV; tobacco mosaic virus, TMV) or transformed with recombinant plasmid expression vectors (e.g., Ti plasmid) containing one or more coding sequences.

[0128] In a useful insect system, *Autographa californica* nuclear polyhydrosis virus (AcNPV) is used as a vector to express foreign genes. The virus grows in *Spodoptera frugiperda* cells. The isolated nucleic acid coding sequences are cloned into non-essential regions (for example the polyhedron gene) of the virus and placed under control of an AcNPV promoter (for example, the polyhedron promoter). Successful insertion of the coding sequences results in the inactivation of the polyhedron gene and production of non-occluded recombinant virus (i.e., virus lacking the proteinaceous coat coded for by the polyhedron gene). These recombinant viruses are then used to infect *Spodoptera frugiperda* cells in which the inserted gene is expressed (e.g., U.S. Pat. No. 4,215,051).

[0129] Examples of useful mammalian host cell lines are VERO and HeLa cells, Chinese hamster ovary (CHO) cell lines, WI38, BHK, COS-7, 293, HepG2, NIH3T3, RIN and MDCK cell lines. In addition, a host cell may be chosen that modulates the expression of the inserted sequences, or modifies and processes the gene product in the specific fashion desired. Such modifications (e.g., glycosylation) and processing (e.g., cleavage) of protein products may be important for the function of the encoded protein.

[0130] Different host cells have characteristic and specific mechanisms for the post-translational processing and modification of proteins. Appropriate cell lines or host systems can be chosen to ensure the correct modification and processing of the foreign protein expressed. Expression vectors for use in mammalian cells ordinarily include an origin of replication (as necessary), a promoter located in front of the gene to be expressed, along with any necessary ribosome binding sites, RNA splice sites, polyadenylation site, and transcriptional terminator sequences. The origin of replication may be provided either by construction of the vector to include an exogenous origin, such as may be derived from SV40 or other viral (e.g., Polyoma, Adeno, VSV, BPV) source, or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter is often sufficient.

[0131] The promoters may be derived from the genome of mammalian cells (e.g., metallothionein promoter) or from mammalian viruses (e.g., the adenovirus late promoter; the vaccinia virus 7.5K promoter). Further, it is also possible, and may be desirable, to utilize promoter or control sequences normally associated with the desired gene sequence, provided such control sequences are compatible with the host cell systems.

[0132] Specific initiation signals may also be required for efficient translation of the claimed isolated nucleic acid coding sequences. These signals include the ATG initiation codon and adjacent sequences. Exogenous translational control signals, including the ATG initiation codon, may additionally need to be provided. One of ordinary skill in the art would readily be capable of determining this need and providing the necessary signals. It is well known that the initiation codon must be in-frame (or in-phase) with the reading frame of the desired coding sequence to ensure translation of the entire insert. These exogenous translational control signals and initiation codons can be of a variety of origins, both natural and synthetic. The efficiency of expression may be enhanced by the inclusion of appropriate transcription enhancer elements or transcription terminators (Bittner et al., 1987).

[0133] In eukaryotic expression, one will also typically desire to incorporate into the transcriptional unit an appropriate polyadenylation site (e.g., 5'-AATAAA-3') if one was not contained within the original cloned segment. Typically, the poly A addition site is placed about 30 to 2000 nucleotides "downstream" of the termination site of the protein at a position prior to transcription termination.

[0134] For long-term, high-yield production of recombinant proteins, stable expression is preferred. For example, cell lines that stably express constructs encoding proteins may be engineered. Rather than using expression vectors that contain viral origins of replication, host cells can be transformed with vectors controlled by appropriate expression control elements (e.g., promoter, enhancer, sequences, transcription terminators, polyadenylation sites, etc.), and a selectable marker. Following the introduction of foreign DNA, engineered cells may be allowed to grow for 1-2 days in an enriched medium, and then are switched to a selective medium. The selectable marker in the recombinant plasmid confers resistance to the selection and allows cells to stably integrate the plasmid into their chromosomes and grow to form foci, which in turn can be cloned and expanded into cell lines.

[0135] It is contemplated that the nucleic acids of the invention may be “overexpressed”, i.e., expressed in increased levels relative to its natural expression in human cells, or even relative to the expression of other proteins in the recombinant host cell. Such overexpression may be assessed by a variety of methods, including radio-labeling and/or protein purification. However, simple and direct methods are preferred, for example, those involving SDS/PAGE and protein staining or western blotting, followed by quantitative analyses, such as densitometric scanning of the resultant gel or blot. A specific increase in the level of the recombinant protein or peptide in comparison to the level in natural human cells is indicative of overexpression, as is a relative abundance of the specific protein in relation to the other proteins produced by the host cell and, e.g., visible on a gel.

[0136] II. Delivery In various embodiments of the invention, the expression construct may comprise a virus or engineered construct derived from a viral genome. The ability of certain viruses to enter cells via receptor-mediated endocytosis and to integrate into the host cell genome and express viral genes stably and efficiently have made them attractive candidates for the transfer of foreign genes into mammalian cells (Ridgeway, 1988; Nicolas and Rubenstein, 1988; Baichwal and Sugden, 1986; Temin, 1986). The first viruses used as vectors were DNA viruses including the papovaviruses (simian virus 40, bovine papilloma virus, and polyoma) (Ridgeway, 1988; Baichwal and Sugden, 1986) and adenoviruses (Ridgeway, 1988; Baichwal and Sugden, 1986) and adeno-associated viruses. Retroviruses also are attractive gene transfer vehicles (Nicolas and Rubenstein, 1988; Temin, 1986) as are vaccinia virus (Ridgeway, 1988) and adeno-associated virus (Ridgeway, 1988). Such vectors may be used to (i) transform cell lines *in vitro* for the purpose of expressing proteins of interest or (ii) to transform cells *in vitro* or *in vivo* to provide therapeutic polypeptides in a gene therapy scenario. Herpes simplex virus (HSV) is another attractive candidate, especially where neurotropism is desired. HSV also is relatively easy to manipulate and can be grown to high titers. Thus, delivery is less of a problem, both in terms of volumes needed to attain sufficient MOI and in a lessened need for repeat dosings.

[0137] With the recent recognition of defective hepatitis B viruses, new insight was gained into the structure-function relationship of different viral sequences. *In vitro* studies showed that the virus could retain the ability for helper-dependent packaging and reverse transcription despite the deletion of up to 80% of its genome (Horwich et al., 1990). This suggested that large portions of the genome could be replaced with foreign genetic material. The hepatotropism and persistence (integration) were particularly attractive properties for liver-directed gene transfer. Chang et al., recently introduced the chloramphenicol acetyltransferase (CAT) gene into duck hepatitis B virus genome in the place of the polymerase, surface, and pre-surface coding sequences. It was co-transfected with wild-type virus into an avian hepatoma cell line. Culture media containing high titers of the recombinant virus were used to infect primary duckling hepatocytes. Stable CAT gene expression was detected for at least 24 days after transfection (Chang et al., 1991).

[0138] Several non-viral methods for the transfer of expression constructs into cultured mammalian cells also are

contemplated by the present invention. These include calcium phosphate precipitation (Graham and Van Der Eb, 1973; Chen and Okayama, 1987; Rippe et al., 1990) DEAE-dextran (Gopal, 1985), electroporation (Tur-Kaspa et al., 1986; Potter et al., 1984), direct microinjection (Harland and Weintraub, 1985), DNA-loaded liposomes (Nicolau and Sene, 1982; Fraley et al., 1979) and lipofectamine-DNA complexes, cell sonication (Fechheimer et al., 1987), gene bombardment using high velocity microprojectiles (Yang et al., 1990), and receptor-mediated transfection (Wu and Wu, 1987; Wu and Wu, 1988). Some of these techniques may be successfully adapted for *in vivo* or *ex vivo* use.

[0139] Once the expression construct has been delivered into the cell the nucleic acid encoding the gene of interest may be positioned and expressed at different sites. In certain embodiments, the nucleic acid encoding the gene may be stably integrated into the genome of the cell. This integration may be in the cognate location and orientation via homologous recombination (gene replacement) or it may be integrated in a random, non-specific location (gene augmentation). In yet further embodiments, the nucleic acid may be stably maintained in the cell as a separate, episomal segment of DNA. Such nucleic acid segments or “episomes” encode sequences sufficient to permit maintenance and replication independent of or in synchronization with the host cell cycle. How the expression construct is delivered to a cell and where in the cell the nucleic acid remains is dependent on the type of expression construct employed.

[0140] In one embodiment, the expression construct may simply consist of naked recombinant DNA or plasmids. Transfer of the construct may be performed by any of the methods mentioned above which physically or chemically permeabilize the cell membrane. This is particularly applicable for transfer *in vitro* but it may be applied to *in vivo* use as well. Dubensky et al., (1984) successfully injected polyomavirus DNA in the form of calcium phosphate precipitates into liver and spleen of adult and newborn mice demonstrating active viral replication and acute infection. Benvenisty and Neshif (1986) also demonstrated that direct intraperitoneal injection of calcium phosphate-precipitated plasmids results in expression of the transfected genes. It is envisioned that DNA encoding a gene of interest may also be transferred in a similar manner *in vivo* and express the gene product.

[0141] Another embodiment of the invention for transferring a naked DNA expression construct or DNA segment into cells may involve particle bombardment. This method depends on the ability to accelerate DNA-coated microprojectiles to a high velocity allowing them to pierce cell membranes and enter cells without killing them (Klein et al., 1987). Several devices for accelerating small particles have been developed. One such device relies on a high voltage discharge to generate an electrical current, which in turn provides the motive force (Yang et al., 1990). The microprojectiles used have consisted of biologically inert substances such as tungsten or gold beads.

[0142] Selected organs including the liver, skin, and muscle tissue of rats and mice have been bombarded *in vivo* (Yang et al., 1990; Zelenin et al., 1991). This may require surgical exposure of the tissue or cells, to eliminate any intervening tissue between the gun and the target organ, i.e.,

ex vivo treatment. Again, DNA encoding a particular gene may be delivered via this method and still be incorporated by the present invention.

[0143] In a further embodiment of the invention, the DNA segment or expression construct may be entrapped in a liposome. Liposomes are vesicular structures characterized by a phospholipid bilayer membrane and an inner aqueous medium. Multilamellar liposomes have multiple lipid layers separated by aqueous medium. They form spontaneously when phospholipids are suspended in an excess of aqueous solution. The lipid components undergo self-rearrangement before the formation of closed structures and entrap water and dissolved solutes between the lipid bilayers (Ghosh and Bachhawat, 1991). Also contemplated are lipofectamine-DNA complexes.

[0144] Liposome-mediated nucleic acid delivery and expression of DNA in vitro has been very successful. Wong et al., (1980) demonstrated the feasibility of liposome-mediated delivery and expression of foreign DNA in cultured chick embryo, HeLa and hepatoma cells. Nicolau et al., (1987) accomplished successful liposome-mediated gene transfer in rats after intravenous injection.

[0145] In certain embodiments, the liposome may be complexed with a hemagglutinating virus (HVJ). This has been shown to facilitate fusion with the cell membrane and promote cell entry of liposome-encapsulated DNA (Kaneda et al., 1989). In other embodiments, the liposome may be complexed or employed in conjunction with nuclear non-histone chromosomal proteins (HMG-1) (Kato et al., 1991). In yet further embodiments, the liposome may be complexed or employed in conjunction with both HVJ and HMG-1. In that such expression constructs have been successfully employed in transfer and expression of nucleic acid in vitro and in vivo, then they are applicable for the present invention. Where a bacterial promoter is employed in the DNA construct, it also will be desirable to include within the liposome an appropriate bacterial polymerase.

[0146] Other expression constructs which can be employed to deliver a nucleic acid encoding a particular gene into cells are receptor-mediated delivery vehicles. These take advantage of the selective uptake of macromolecules by receptor-mediated endocytosis in almost all eukaryotic cells. Because of the cell type-specific distribution of various receptors, the delivery can be highly specific (Wu and Wu, 1993).

[0147] Receptor-mediated gene targeting vehicles generally consist of two components: a cell receptor-specific ligand and a DNA-binding agent. Several ligands have been used for receptor-mediated gene transfer. The most extensively characterized ligands are asialoorosomuroid (ASOR) (Wu and Wu, 1987) and transferrin (Wagner et al., 1990). Recently, a synthetic neoglycoprotein, which recognizes the same receptor as ASOR, has been used as a gene delivery vehicle (Ferkol et al., 1993; Perales et al., 1994) and epidermal growth factor (EGF) has also been used to deliver genes to squamous carcinoma cells (Myers, EPO 0273085).

[0148] In other embodiments, the delivery vehicle may comprise a ligand and a liposome. For example, Nicolau et al., (1987) employed lactosyl-ceramide, a galactose-terminal asialganglioside, incorporated into liposomes and observed an increase in the uptake of the insulin gene by

hepatocytes. Thus, it is feasible that a nucleic acid encoding a particular gene also may be specifically delivered into a cell type such as lung, epithelial or tumor cells, by any number of receptor-ligand systems with or without liposomes.

[0149] In certain embodiments, gene transfer may more easily be performed under ex vivo conditions. Ex vivo gene therapy refers to the isolation of cells from an organism, the delivery of a nucleic acid into the cells in vitro, and then the return of the modified cells back into an organism. This may involve the surgical removal of tissue/organs from an animal or the primary culture of cells and tissues. Anderson et al., U.S. Pat. No. 5,399,346, and incorporated herein in its entirety, disclose ex vivo therapeutic methods. F. Oligonucleotide Synthesis Oligonucleotide synthesis is well known to those of skill in the art. Various different mechanisms of oligonucleotide synthesis have been disclosed in for example, U.S. Pat. Nos. 4,659,774, 4,816,571, 5,141,813, 5,264,566, 4,959,463, 5,428,148, 5,554,744, 5,574,146, 5,602,244, each of which is incorporated herein by reference.

[0150] Phosphoramidite chemistry (Beaucage, and Lye, 1992) has become by far the most widely used coupling chemistry for the synthesis of oligonucleotides. As is well known to those skilled in the art, phosphoramidite synthesis of oligonucleotides involves activation of nucleoside phosphoramidite monomer precursors by reaction with an activating agent to form activated intermediates, followed by sequential addition of the activated intermediates to the growing oligonucleotide chain (generally anchored at one end to a suitable solid support) to form the oligonucleotide product.

[0151] Tetrazole is commonly used for the activation of the nucleoside phosphoramidite monomers. Tetrazole has an acidic proton which presumably protonates the basic nitrogen of the diisopropylamino phosphine group, thus making the diisopropylamino group a leaving group. The negatively charged tetrazolium ion then makes an attack on the trivalent phosphorous, forming a transient phosphorous tetrazolide species. The 5'-OH group of the solid support bound nucleoside then attacks the active trivalent phosphorous species, resulting in the formation of the internucleotide linkage. The trivalent phosphorous is finally oxidized to the pentavalent phosphorous. The US patents listed above describe other activators and solid supports for oligonucleotide synthesis.

[0152] High throughput oligonucleotide synthesis can be achieved using a synthesizer. The Genome Science and Technology Center, as one aspect of the automation development effort, recently developed a high throughput large scale oligonucleotide synthesizer. This instrument, denoted the MERMADE, is based on a 96-well plate format and uses robotic control to carry out parallel synthesis on 192 samples (296-well plates). This device has been variously described in the literature and in presentations, is generally available in the public domain (licensed from the University of Texas and available on contract from Avantec). The device has gone through various generations with differing operating parameters. The device may be used to synthesize 192 oligonucleotides simultaneously with 99% success. It has virtually 100% success for oligomers less than 60 bp; operates at 20 mM synthesis levels, and gives a product yield of >99% complete synthesis. Using these systems the

inventor has synthesized over 10,000 oligomers used for sequencing, PCRTM amplification and recombinant DNA applications. For most uses, including cloning, synthesis success is sufficient such that post synthesis purification is not required.

[0153] Once the genome has been synthesized using the methods of the present invention it may be necessary to screen the sequences for analysis of function. Specifically contemplated by the present inventor are chip-based DNA technologies such as those described by Hacia et al. (1996) and Shoemaker et al. (1996). Briefly, these techniques involve quantitative methods for analyzing large numbers of genes rapidly and accurately. By tagging genes with oligonucleotides or using fixed probe arrays, one can employ chip technology to segregate target molecules as high density arrays and screen these molecules on the basis of hybridization. See also Pease et al. (1994); Fodor et al. (1991).

[0154] The use of combinatorial synthesis and high throughput screening assays are well known to those of skill in the art, e.g. U.S. Pat. Nos. 5,807,754; 5,807,683; 5,804,563; 5,789,162; 5,783,384; 5,770,358; 5,759,779; 5,747,334; 5,686,242; 5,198,346; 5,738,996; 5,733,743; 5,714,320; 5,663,046 (each specifically incorporated herein by reference). These patents teach various aspects of the methods and compositions involved in the assembly and activity analyses of high density arrays of different polysubunits (polynucleotides or polypeptides). As such it is contemplated that the methods and compositions described in the patents listed above may be useful in assay the activity profiles of the compositions of the present invention.

[0155] The present invention produces a replication competent polynucleotide. Viruses are naturally occurring replication competent pieces of DNA, to the extent that disclosure regarding viruses may be useful in the context of the present invention, the following is a disclosure of viruses. Researchers note that viruses have evolved to be able to deliver their DNA to various host tissues despite the human body's various defensive mechanisms. For this reason, numerous viral vectors have been designed by researchers seeking to create vehicles for therapeutic gene delivery. Some of the types of viruses that have been engineered are listed below.

[0156] II. Adenovirus

[0157] Adenovirus is a 36 kB, linear, double-strained DNA virus that allows substitution of large pieces of adenoviral DNA with foreign sequences up to 7 kB (Grunhaus and Horwitz, 1992). Adenovirus DNA does not integrate into the host cell chromosomal because adenoviral DNA can replicate in an episomal manner. Also, adenoviruses are structurally stable, and no genome rearrangement has been detected after extensive amplification. Adenovirus can infect virtually all epithelial cells regardless of their cell cycle stage. This means that adenovirus can infect non-dividing cells. So far, adenoviral infection appears to be linked only to mild disease such as acute respiratory disease in humans. This group of viruses can be obtained in high titers, e.g., 10^9 - 10^{11} plaque-forming units per ml, and they are highly infective.

[0158] Both ends of the viral genome contain 100-200 base pair inverted repeats (ITRs), which are cis elements necessary for viral DNA replication and packaging. The

early (E) and late (L) regions of the genome contain different transcription units that are divided by the onset of viral DNA replication. The E1 region (E1A and E1B) encodes proteins responsible for the regulation of transcription of the viral genome and a few cellular genes. The expression of the E2 region (E2A and E2B) results in the synthesis of the proteins for viral DNA replication. These proteins are involved in DNA replication, late gene expression and host cell shut-off (Renan, 1990). The products of the late genes, including the majority of the viral capsid proteins, are expressed only after significant processing of a single primary transcript issued by the major late promoter (MLP). The MLP, (located at 16.8 m.u.) is particularly efficient during the late phase of infection, and all the mRNA's issued from this promoter possess a 5'-tripartite leader (TPL) sequence which makes them preferred mRNA's for translation.

[0159] The E3 region encodes proteins that appears to be necessary for efficient lysis of Ad infected cells as well as preventing TNF-mediated cytolysis and CTL mediated lysis of infected cells. In general, the E4 region encodes is believed to encode seven proteins, some of which activate the E2 promoter. It has been shown to block host mRNA transport and enhance transport of viral RNA to cytoplasm. Further the E4 product is in part responsible for the decrease in early gene expression seen late in infection. E4 also inhibits E1A and E4 (but not E1B), expression during lytic growth. Some E4 proteins are necessary for efficient DNA replication however the mechanism for this involvement is unknown. E4 is also involved in post-transcriptional events in viral late gene expression; i.e., alternative splicing of the tripartite leader in lytic growth. Nevertheless, E4 functions are not absolutely required for DNA replication but their lack will delay replication. Other functions include negative regulation of viral DNA synthesis, induction of sub-nuclear reorganization normally seen during adenovirus infection, and other functions that are necessary for viral replication, late viral mRNA accumulation, and host cell transcriptional shut off.

[0160] II. Retroviruses

[0161] The retroviruses are a group of single-stranded RNA viruses characterized by an ability to convert their RNA to double-stranded DNA to infected cells by a process of reverse-transcription (Coffin, 1990). The resulting DNA then stably integrates into cellular chromosomes as a provirus and directs synthesis of viral proteins. The integration results in the retention of the viral gene sequences in the recipient cell and its descendants. The retroviral genome contains three genes, gag, pol, and env that code for capsid proteins, polymerase enzyme, and envelope components, respectively. A sequence found upstream from the gag gene, termed ψ components is constructed (Mann et al., 1983). When a recombinant plasmid containing a human cDNA, together with the retroviral LTR and ψ sequences is introduced into this cell line (by calcium phosphate precipitation for example), the ψ sequence allows the RNA transcript of the recombinant plasmid to be packaged into viral particles, which are then secreted into the culture media (Nicolas and Rubenstein, 1988; Temin, 1986; Mann et al., 1983). The media containing the recombinant retroviruses is then collected, optionally concentrated, and used for gene transfer. Retroviral vectors are able to infect a broad variety of cell types. However, integration requires the division of host cells (Paskind et al., 1975).

[0162] The retrovirus family includes the subfamilies of the oncoviruses, the lentiviruses and the spumaviruses. Two oncoviruses are Moloney murine leukemia virus (MMLV) and feline leukemia virus (FeLV). The lentiviruses include human immunodeficiency virus (HIV), simian immunodeficiency virus (SIV) and feline immunodeficiency virus (FIV). Among the murine viruses such as MMLV there is a further classification. Murine viruses may be ecotropic, xenotropic, polytropic or amphotropic. Each class of viruses target different cell surface receptors in order to initiate infection.

[0163] Further advances in retroviral vector design and concentration methods have allowed production of amphotropic and xenotropic viruses with titers of 10^8 to 10^9 cfu/ml (Bowles et al., 1996; Irwin et al., 1994; Jolly, 1994; Kitten et al., 1997).

[0164] Replication defective recombinant retroviruses are not acute pathogens in primates (Chowdhury et al., 1991). They have been successfully applied in cell culture systems to transfer the CFTR gene and generate cAMP-activated Cl^- secretion in a variety of cell types including human airway epithelia (Drumm et al., 1990, Olsen et al., 1992; Anderson et al., 1991; Olsen et al., 1993). While there is evidence of immune responses to the viral gag and env proteins, this does not prevent successful readministration of vector (McCormack et al., 1997). Further, since recombinant retroviruses have no expressed gene products other than the transgene, the risk of a host inflammatory response due to viral protein expression is limited (McCormack et al., 1997). As for the concern about insertional mutagenesis, to date there are no examples of insertional mutagenesis arising from any human trial with recombinant retroviral vectors.

[0165] More recently, hybrid lentivirus vectors have been described combining elements of human immunodeficiency virus (HIV) (Naldini et al., 1996) or feline immunodeficiency virus (FIV) (Poeschla et al., 1998) and MMLV. These vectors transduce nondividing cells in the CNS (Naldini et al., 1996; Blomer et al., 1997), liver (Kafri et al., 1997), muscle (Kafri et al., 1997) and retina (Miyoshi et al., 1997). However, a recent report in xenograft models of human airway epithelia suggests that in well-differentiated epithelia, gene transfer with VSV-G pseudotyped HIV-based lentivirus is inefficient (Goldman et al., 1997).

[0166] III. Adeno-Associated Virus

[0167] In addition, AAV possesses several unique features that make it more desirable than the other vectors. Unlike retroviruses, AAV can infect non-dividing cells; wild-type AAV has been characterized by integration, in a site-specific manner, into chromosome 19 of human cells (Kotin and Berns, 1989; Kotin et al., 1990; Kotin et al., 1991; Samulski et al., 1991); and AAV also possesses anti-oncogenic properties (Ostrove et al., 1981; Berns and Giraud, 1996). Recombinant AAV genomes are constructed by molecularly cloning DNA sequences of interest between the AAV ITRs, eliminating the entire coding sequences of the wild-type AAV genome. The AAV vectors thus produced lack any of the coding sequences of wild-type AAV, yet retain the property of stable chromosomal integration and expression of the recombinant genes upon transduction both in vitro and in vivo (Berns, 1990; Berns and Bohensky, 1987; Bertran et al., 1996; Kearns et al., 1996; Ponnazhagan et al., 1997a). Until recently, AAV was believed to infect almost all cell types, and even cross species barriers. However, it now has

been determined that AAV infection is receptor-mediated (Ponnazhagan et al., 1996; Mizukami et al., 1996).

[0168] AAV utilizes a linear, single-stranded DNA of about 4700 base pairs. Inverted terminal repeats flank the genome. Two genes are present within the genome, giving rise to a number of distinct gene products. The first, the cap gene, produces three different virion proteins (VP), designated VP-1, VP-2 and VP-3. The second, the rep gene, encodes four non-structural proteins (NS). One or more of these rep gene products is responsible for transactivating AAV transcription. The sequence of AAV is provided by Srivastava et al. (1983), and in U.S. Pat. No. 5,252,479 (entire text of which is specifically incorporated herein by reference).

[0169] The three promoters in AAV are designated by their location, in map units, in the genome. These are, from left to right, p5, p19 and p40. Transcription gives rise to six transcripts, two initiated at each of three promoters, with one of each pair being spliced. The splice site, derived from map units 42-46, is the same for each transcript. The four non-structural proteins apparently are derived from the longer of the transcripts, and three virion proteins all arise from the smallest transcript.

[0170] AAV is not associated with any pathologic state in humans. Interestingly, for efficient replication, AAV requires "helping" functions from viruses such as herpes simplex virus I and II, cytomegalovirus, pseudorabies virus and, of course, adenovirus. The best characterized of the helpers is adenovirus, and many "early" functions for this virus have been shown to assist with AAV replication. Low level expression of AAV rep proteins is believed to hold AAV structural expression in check, and helper virus infection is thought to remove this block.

[0171] IV. Vaccinia Virus

[0172] Vaccinia viruses are a genus of the poxyvirus family. Vaccinia virus vectors have been used extensively because of the ease of their construction, relatively high levels of expression obtained, wide host range and large capacity for carrying DNA. Vaccinia contains a linear, double-stranded DNA genome of about 186 kB that exhibits a marked "A-T" preference. Inverted terminal repeats of about 10.5 kB flank the genome. The majority of essential genes appear to map within the central region, which is most highly conserved among poxyviruses. Estimated open reading frames in vaccinia virus number from 150 to 200. Although both strands are coding, extensive overlap of reading frames is not common. U.S. Pat. No. 5,656,465 (specifically incorporated by reference) describes in vivo gene delivery using pox viruses.

[0173] V. Papovavirus

[0174] The papovavirus family includes the papillomaviruses and the polyomaviruses. The polyomaviruses include Simian Virus 40 (SV40), polyoma virus and the human polyomaviruses BKV and JCV. Papillomaviruses include the bovine and human papillomaviruses. The genomes of polyomaviruses are circular DNAs of a little more than 5000 bases. The predominant gene products are three virion proteins (VP1-3) and Large T and Small T antigens. Some have an additional structural protein, the agnoprotein, and others have a Middle T antigen. Papillomaviruses are somewhat larger, approaching 8 kB Little is known about the

cellular receptors for polyomaviruses, but polyoma infection can be blocked by treating with sialidase. SV40 will still infect sialidase-treated cells, but JCV cannot hemagglutinate cells treated with sialidase. Because interaction of polyoma V1 with the cell surface activates c-myc and c-fos, it has been hypothesized that the virus receptor may have some properties of a growth factor receptor. Papillomaviruses are specifically tropic for squamous epithelia, though the specific receptor has not been identified.

[0175] VI. Paramyxovirus

[0176] The paramyxovirus family is divided into three genera: paramyxovirus, morbillivirus and pneumovirus. The paramyxovirus genus includes the mumps virus and Sendai virus, among others, while the morbilliviruses include the measles virus and the pneumoviruses include respiratory syncytial virus (RSV). Paramyxovirus genomes are RNA based and contain a set of six or more genes, covalently linked in tandem. The genome is something over 15 kB in length. The viral particle is 150-250 nm in diameter, with "fuzzy" projections or spikes protruding therefrom. These are viral glycoproteins that help mediate attachment and entry of the virus into host cells.

[0177] A specialized series of proteins are involved in the binding and entry of paramyxoviruses. Attachment in Paramyxoviruses and Morbilliviruses is mediated by glycoproteins that bind to sialic acid-containing receptors. Other proteins anchor the virus by embedding hydrophobic regions in the lipid bilayer of the cell's surface, and exhibit hemagglutinating and neuraminidase activities. In Pneumoviruses, the glycoprotein is heavily glycosylated with O-glycosidic bonds. This molecule lacks the exhibit hemagglutinating and neuraminidase activities of its relatives.

[0178] VII. Herpesvirus.

[0179] Because herpes simplex virus (HSV) is neurotropic, it has generated considerable interest in treating nervous system disorders. Moreover, the ability of HSV to establish latent infections in non-dividing neuronal cells without integrating in to the host cell chromosome or otherwise altering the host cell's metabolism, along with the existence of a promoter that is active during latency makes HSV an attractive vector. And though much attention has focused on the neurotropic applications of HSV, this vector also can be exploited for other tissues given its wide host range.

[0180] Another factor that makes HSV an attractive vector is the size and organization of the genome. Because HSV is large, incorporation of multiple genes or expression cassettes is less problematic than in other smaller viral systems. In addition, the availability of different viral control sequences with varying performance (temporal, strength, etc.) makes it possible to control expression to a greater extent than in other systems. It also is an advantage that the virus has relatively few spliced messages, further easing genetic manipulations.

[0181] HSV also is relatively easy to manipulate and can be grown to high titers. Thus, delivery is less of a problem, both in terms of volumes needed to attain sufficient MOI and in a lessened need for repeat dosings. For a review of HSV as a gene therapy vector, see Glorioso et al. (1995).

[0182] HSV, designated with subtypes 1 and 2, are enveloped viruses that are among the most common infectious

agents encountered by humans, infecting millions of human subjects worldwide. The large, complex, double-stranded DNA genome encodes for dozens of different gene products, some of which derive from spliced transcripts. In addition to virion and envelope structural components, the virus encodes numerous other proteins including a protease, a ribonucleotides reductase, a DNA polymerase, a ssDNA binding protein, a helicase/primase, a DNA dependent ATPase, a dUTPase and others.

[0183] HSV genes form several groups whose expression is coordinately regulated and sequentially ordered in a cascade fashion (Honess and Roizman, 1974; Honess and Roizman 1975; Roizman and Sears, 0.1995). The expression of α genes, the first set of genes to be expressed after infection, is enhanced by the virion protein number 16, or α -transducing factor (Post et al., 1981; Batterson and Roizman, 1983; Campbell et al., 1983). The expression of β genes requires functional α gene products, most notably ICP4, which is encoded by the α 4 gene (DeLuca et al., 1985). γ genes, a heterogeneous group of genes encoding largely virion structural proteins, require the onset of viral DNA synthesis for optimal expression (Holland et al., 1980).

[0184] In line with the complexity of the genome, the life cycle of HSV is quite involved. In addition to the lytic cycle, which results in synthesis of virus particles and, eventually, cell death, the virus has the capability to enter a latent state in which the genome is maintained in neural ganglia until some as of yet undefined signal triggers a recurrence of the lytic cycle. Avirulent variants of HSV have been developed and are readily available for use in gene therapy contexts (U.S. Pat. No. 5,672,344).

G. EXAMPLES

[0185] The following examples are included to demonstrate preferred embodiments of the invention. It should be appreciated by those of skill in the art that the techniques disclosed in the examples which follow represent techniques discovered by the inventor to function well in the practice of the invention, and thus can be considered to constitute preferred modes for its practice. However, those of skill in the art should, in light of the present disclosure, appreciate that many changes can be made in the specific embodiments which are disclosed and still obtain a like or similar result without departing from the spirit and scope of the invention.

Example 1

Combinatoric gene assembly

[0186] The inventor has developed a strategy of oligomer assembly into larger DNA molecules denoted combinatoric assembly. The procedure is carried out as follows: one may design a plasmid using one of a number of commercial or public domain computer programs to contain the genes, promoters, drug selection, origin of replication, etc. required. SynGene v.2.0 is a program that generates a list of overlapping oligonucleotides sufficient to reassemble the gene or plasmid (see FIG. 7). For instance, for a 5000 bp gene, SynGene 2.0 can generate two lists of 100 component 50 mers from one strand and 100 component 50 mers from the complementary strand such that each pair of oligomers will overlap by 25 base pairs. The program checks the sequence for repeats and produces a MERMADE input file

which directly programs the oligonucleotide synthesizer. The synthesizer produces two sets of 96-well plates containing the complementary oligonucleotides. A SynGene program is depicted in **FIG. 7**. This program is designed to break down a designer gene or genome into oligonucleotides for synthesis. The program is for the complete synthetic designer gene and is based upon an original program for formatting DNA sequences written by Dr. Glen Evans.

[0187] Combinatoric assembly is best carried out using a programmable robotic workstation such as a Beckman Biomek 2000. In short, pairs of oligomers which overlap are mixed and annealed. Following annealing, a smaller set of duplex oligomers is generated. These are again paired and annealed, forming a smaller set of larger oligomers. Sequentially, overlapping oligomers are allowed to anneal until the entire reassembly is completed. Annealing may be carried out in the absence of ligase, or each step may be followed by ligation. In one configuration, oligomers are annealed in the presence of topoisomerase 2, which does not require 5' phosphorylation of the oligomer, occurs at room temperature, and is a rapid (5 minute) reaction as opposed to 12 h ligation at 12°. Following the complete assembly, the resulting DNA molecule can be used for its designed purpose, usually transformation into a bacterial host for replication. The steps in this cycle are outlined in **FIG. 3**.

[0188] This approach has a major advantage over traditional recombinant DNA based cloning. While it is technically feasible to make virtually any modification or mutation in existing DNA molecules, the effort required, as well as the high technical skill, make some constructions difficult or tedious. This method, while having been used for many years, is not applicable to automated gene cloning or large scale creation of entirely novel DNA sequences.

Example 2

Production of Artificial Genes

[0189] In one example, the present invention will produce a known gene of about 1000 base pairs in length by the following method. A set of oligonucleotides, each of 50 bases, is generated such that the entire plus strand of the gene is represented. A second set of oligonucleotides, also comprised of 50-mers, is generated for the minus strand. This set is designed, however, such that complementary pairing with the first and second sets results in overlap of "paired" sequences, i.e., each oligonucleotide of the first set is complementary with regions from two oligonucleotides of the second set (with the possible exception of the terminal oligonucleotides). The region of overlap is set at 30 bases, leaving a 20 base pair overhang for each pair. The first and said second set of oligonucleotides is annealed in a single mixture and treated with a ligating enzyme.

[0190] In another example, the gene to be synthesized is about 5000 base pairs. Each set of oligonucleotides is made up of fifty 100-mers with overlapping regions, of complementary oligonucleotides, of 75 bases, leaving 25 base "sticky ends." In this embodiment, the 5' terminal oligonucleotide of the first oligonucleotide set is annealed with the 3' terminal oligonucleotide of the second set to form a first annealed product, then the next most 5' terminal oligonucleotide of the first set is annealed with the first annealed product to form a second annealed product, and the process

is repeated until all oligonucleotides of said first and said second sets have been annealed. Ligation of the products may occur between steps or at the conclusion of all hybridizations.

[0191] In a third example, a gene of 100,000 bp is synthesized from one thousand 100-mers. Again, the overlap between "pairs" of plus and minus oligonucleotides is 75 bases, leaving a 25 base pair overhang. In this method, a combinatorial approach is used where corresponding pairs of partially complementary oligonucleotides are hybridized in first step. A second round of hybridization then is undertaken with appropriately complementary pairs of products from the first round. This process is repeated a total of 10 times, each round of hybridization reducing the number of products by half. Ligation of the products then is performed.

Example 3

Large Scale Expression of Human Gene Products

[0192] Once the human genome has been characterized, functional analysis of the human genome, based upon the complete sequence, will require a variety of approaches to structural, functional and network biology. The approach proposed herein for producing a series of expression constructs representing all potential human gene products and the assembly of sets of bacterial and/or yeast expressing these products will provide an important avenue into the beginnings of functional analysis.

[0193] Secondly, the approach described here, when developed to its theoretical optima, will allow the large scale transfer of genes to cell lines or organisms for functional analysis. The long term goal of this concept is the creation of living organisms entirely based on bioinformatics and information processing. Obviously, the knowledge of the complete sequence is not sufficient to appreciate the myriad of biological concepts inherent in life.

Example 4

Construction of a Synthetic Plasmid

[0194] A DNA molecule was designed using synthetic parts of previously known plasmids. As a demonstration of this technique, plasmid synlux⁴ was designed. Synlux4 consists of 4800 base pairs of DNA. Within this sequence are included the sequence of lux A and lux B, the A and B components of the luciferase protein from *Vibrio Fisherii*, portions of plasmid. pUC19 including the origin of replication and replication stability sequences, the promoter and coding sequence for tn9 kanamycin/neomycin phosphotransferase. The sequence was designed on a computer using Microsoft Word and Vector NTI (InforMax, Inc.). The sequence is listed in **FIG. 4**.

[0195] Following design, a computer program SynGene 2.0 was used to break the sequence down into components consisting of overlapping 50-mer oligonucleotides. From the 4800 base pair sequence, 192 50-mers were designed. The component oligonucleotides are listed in **FIG. 5**. These component oligonucleotides were synthesized using a custom 96-well oligonucleotide synthesizer (Rayner, et al.) Genome Research, 8, 741-747 (1998). The component oligonucleotides were produced in two 96-well microtitre plates, each plate holding one set of component oligonucle-

otides. Thus, plate one held the forward strand oligos and plate 2 held the reverse strand oligos.

[0196] The oligonucleotides were assembled and ligations carried out using a Biomek 1000 robotic workstation (Beckman). Sequential transfers of oligonucleotides were done by pipetting from one well to a second well of the plate and a ligation reaction carried out using T4 ligase. The pattern of assembly is delineated in FIG. 6.

[0197] Following assembly, the resulting ligation mix was used to transform competent *E. coli* strain DH5a. The transformation mix was plated on LB plates containing 25 µg/ml kanamycin sulfate and recombinant colonies obtained. The resulting recombinant clones were isolated, cloned, and DNA prepared. The DNA was analyzed on 1% agarose gels in order to detect recombinant molecules. Clones were shown to contain the expected 4800 base pair plasmid containing lux A and B genes.

[0198] All of the compositions and/or methods disclosed and claimed herein can be made and executed without undue experimentation in light of the present disclosure. While the compositions and methods of this invention have been described in terms of preferred embodiments, it will be apparent to those of skill in the art that variations may be applied to the compositions and/or methods and in the steps or in the sequence of steps of the method described herein without departing from the concept, spirit and scope of the invention. More specifically, it will be apparent that certain agents which are both chemically and physiologically related may be substituted for the agents described herein while the same or similar results would be achieved. All such similar substitutes and modifications apparent to those skilled in the art are deemed to be within the spirit, scope and concept of the invention as defined by the appended claims.

REFERENCES

[0199] The following references, to the extent that they provide exemplary procedural or other details supplementary to those set forth herein, are specifically incorporated herein by reference.

- [0200] Anderson et al. U.S. Pat. No. 5,399,346, 1995.
- [0201] Anderson et al., *Science*, 253:202-205, 1991. Baichwal and Sugden, "Vectors for gene transfer derived from animal DNA viruses: Transient and stable expression of transferred genes," In: *Gene Transfer*, Kucherlapati (ed.), New York, Plenum Press, pp. 117-148, 1-986.
- [0202] Batterson and Roizman, *J. Virol.*, 46:371-377, 1983.
- [0203] Beaucage, and Lyer, *Tetrahedron*, 48:2223-2311, 1992
- [0204] Benvenisty and Neshif, "Direction introduction of genes into rats and expression of the genes," *Proc. Nat. Acad. Sci. USA*, 83:9551-9555, 1986.
- [0205] Berns and Bohenzky, *Adv. Virus Res.*, 32:243-307, 1987.
- [0206] Bertran, et al., *J Virol.*, 70(10):6759-6766, 1996.
- [0207] Bittner et al., *Methods in Enzymol*, 153:516-544, 1987.
- [0208] Blomer et al., Highly efficient and sustained gene transfer in adult neurons with a lentivirus vector. *J. Virol.* 71:6641-6649, 1997
- [0209] Bowles et al., *Hum. Gene Ther.*, 7:1735-1742, 1996.
- [0210] Chang et al., "Foreign gene delivery and expression in hepatocytes using a hepatitis B virus vector," *Hepatology*, 14:124A, 1991.
- [0211] Chen and Okayama, "High-efficiency transfection of mammalian cells by plasmid DNA," *Mol. Cell Biol.*, 7:2745-2752, 1987.
- [0212] Coffin, "Retroviridae and their replication," In: *Virology*, Fields, Knipe (ed.), New York: Raven Press, pp. 1437-1500, 1990.
- [0213] Coffin, In: *Virology*, ed., New York: Raven Press, pp. 1437-1500, 1990.
- [0214] Colberre-Garapin et al., *J. Mol. Biol.*, 150:1, 1981.
- [0215] Couch et al, "Immunization with types 4 and 7 adenovirus by selective infection of the intestinal tract," *Am. Rev. Resp. Dis.*, 88:394-403, 1963.
- [0216] Coupar et al., "A general method for the construction of recombinant vaccinia virus expressing multiple foreign genes," *Gene*, 68:1-10, 1988.
- [0217] Davey et al., *EPO No.* 329 822.
- [0218] DeLuca et al., *J. Virol.*, 56:558-570, 1985.
- [0219] Dnummetal., *Cell*, 62:1227-1233, 1990.
- [0220] Dubensky et al., "Direct transfection of viral and plasmid DNA into the liver or spleen of mice," *Proc. Nat. Acad. Sci. USA*, 81:7529-7533, 1984.
- [0221] Fechheimer et al., "Transfection of mammalian cells with plasmid DNA by scrape loading and sonication loading," *Proc. Natl. Acad. Sci USA*, 84:8463-8467, 1987.
- [0222] Ferkol et al., "Regulation of the phosphoenolpyruvate carboxykinase/human factor IX gene introduced into the livers of adult rats by receptor-mediated gene transfer," *FASEB J.*, 7:1081-1091, 1993.
- [0223] Fraley et al., "Entrapment of a bacterial plasmid in phospholipid vesicles: Potential for gene transfer," *Proc. Natl. Acad. Sci. USA*, 76:3348-3352, 1979.
- [0224] Freifelder, *Physical Biochemistry Applications to Biochemistry and Molecular Biology*, 2nd ed. Wm. Freeman and Co., New York, N.Y., 1982.
- [0225] Friedmann, "Progress toward human gene therapy," *Science*, 244:1275-1281, 1989.
- [0226] Frohman, In: *PCR™ Protocols: A Guide To Methods And Applications*, Academic Press, N.Y., 1990.
- [0227] Ghosh-Choudhury et al, *EMBO J*, 6:1733-1739, 1987.
- [0228] Ghosh and Bachhawat, "Targeting of liposomes to hepatocytes," In: *Liver diseases, targeted diagnosis*

- and therapy using specific receptors and ligands, Wu, Wu (ed.), New York:, Marcel Dekker, pp. 87-104, 1991.
- [0229] Gingeras et al., PCT Application WO 88/10315.
- [0230] Glorioso et al., *Ann. Rev. Microbiol.*, 49:675-710, 1995.
- [0231] Goldman et al., *Hum Gene Ther.* 8(18): 2261-2268, 1997.
- [0232] Gomez-Foix et al., "Adenovirus-mediated transfer of the muscle glycogen phosphorylase gene into hepatocytes confers altered regulation of glycogen," *J. Biol. Chem.*, 267:25129-25134, 1992.
- [0233] Gopal, "Gene transfer method for transient gene expression, stable transfection, and cotransfection of suspension cell cultures," *Mol Cell Biol.*, 5:1188-1190, 1985.
- [0234] Graham and Prevec, "Adenovirus-based expression vectors and recombinant vaccines," *Biotech.*, 20:363-390, 1992.
- [0235] Graham and Prevec, "Manipulation of adenovirus vector," In: *Methods in Molecular Biology: Gene Transfer and Expression Protocol*, Clifton and Murray (ed.), NJ, Humana Press, 7:109-128, 1991.
- [0236] Graham and Van Der Eb, "A new technique for the assay of infectivity of human adenovirus 5 DNA," *Virology*, 52:456-467, 1973.
- [0237] Graham et al., "Characteristics of a human cell line transformed by DNA from human adenovirus type 5," *J. Gen. Virol.*, 36:59-72, 1977.
- [0238] Grunhaus and Horwitz, "Adenovirus as cloning vector," *Seminar in Virology*, 3:237-252, 1992.
- [0239] Harland and Weintraub, "Translation of mammalian mRNA injected into *Xenopus* oocytes is specifically inhibited by antisense RNA," *J. Cell Biol.*, 101:1094-1099, 1985.
- [0240] Hermonat and Muzycska, "Use of adenoassociated virus as a mammalian DNA cloning vector: Transduction of neomycin resistance into mammalian tissue culture cells," *Proc. Nat. Acad. Sci. USA*, 81:6466-6470, 1984.
- [0241] Hersdorffer et al., "Efficient gene transfer in live mice using a unique retroviral packaging line," *DNA Cell Biol.*, 9:713-723, 1990.
- [0242] Herz and Gerard, "Adenovirus-mediated transfer of low density lipoprotein receptor gene acutely accelerates cholesterol clearance in normal mice," *Proc. Natl. Acad. Sci. USA* 90:2812-2816, 1993.
- [0243] Hess et al, *J. Adv. Enzyme Reg.*, 7:149, 1968.
- [0244] Hitzeman et al., *J. Biol. Chem.*, 255:2073, 1980.
- [0245] Holland et al., *Biochemistry*, 17:4900, 1978.
- [0246] Holland et al., *Virology*, 101:10-18, 1980.
- [0247] Honess and Roizman, *J Virol.*, 14:8-19, 1974.
- [0248] Honess and Roizman, *J Virol.*, 16:1308-1326, 1975
- [0249] Horwich et al., "Synthesis of hepadenavirus particles that contain replication-defective duck hepatitis B virus genomes in cultured HuH7 cells," *J. Virol.*, 64:642-650, 1990.
- [0250] Innis et al., *PCR™ Protocols*, Academic Press, Inc., San Diego Calif., 1990.
- [0251] Inouye et al., *Nucleic Acids Res.*, 13: 3101-3109, 1985.
- [0252] Irwin et al., *J. Virol.*, 68:5036-5044, 1994.
- [0253] Johnson et al., "Peptide Turn Mimetics" IN: *Biotechnology And Pharmacy*, Pezzuto et al., (eds.), Chapman and Hall, New York, 1993.
- [0254] Jolly, "Viral vector systems for gene therapy," *Can. Gene Ther.*, 1:51-64, 1994.
- [0255] Jones and Shenk, "Isolation of deletion and substitution mutants of adenovirus type 5," *Cell*, 13:181-188, 1978.
- [0256] Jones, *Genetics*, 85: 12, 1977.
- [0257] Kafri et al., Sustained expression of genes delivered directly into liver and muscle by lentiviral vectors. *Nat. Genet.* 17:314-317, 1997.
- [0258] Kaneda et al., "Increased expression of DNA cointroduced with nuclear protein in adult rat liver," *Science*, 243:375-378, 1989.
- [0259] Karlsson et al, *EMBO J*, 5:2377-2385, 1986.
- [0260] Kato et al, "Expression of hepatitis β virus surface antigen in adult rat liver," *J Biol. Chem.*, 266:3361-3364, 1991.
- [0261] Kearns et al., *Gene Ther.*, 3:748-755, 1996.
- [0262] Kingsman et al., *Gene*, 7: 141, 1979.
- [0263] Kitten et al. *Hum. Gene Ther.*, 8:1491-1494, 1997.
- [0264] Klein et al., "High-velocity microprojectiles for delivering nucleic acids into living cells," *Nature*, 327:70-73, 1987.
- [0265] Kotin and Berns, *Virol.*, 170:460-467, 1989.
- [0266] Kotin et al., *Genomics*, 10:831-834, 1991.
- [0267] Kotin et al, *Proc. Natl. Acad. Sci. USA*, 87:2211-2215, 1990.
- [0268] Kwoh et al., *Proc. Nat. Acad. Sci. USA*, 86:1173, 1989.
- [0269] Kyte and Doolittle, "A simple method for displaying the hydrophobic character of a protein," *J. Mol. Biol.*, 157(1):105-132, 1982.
- [0270] Le Gal La Salle et al., "An adenovirus vector for gene transfer into neurons and glia in the brain," *Science*, 259:988-990, 1993.
- [0271] Levrero et al, *Gene*, 101:195-202, 1991.
- [0272] Lishanski et al., *Proc. Nat'l. Acad. Sci USA.*, 91:2674-2678, (1994) Lowy et al., *Cell*, 22: 817, 1980.
- [0273] Macejak and Sarnow, *Nature*, 353:90-94, 1991.

- [0274] Mann et al., "Construction of a retrovirus packaging mutant and its use to produce helper-free defective retrovirus," *Cell*, 33:153-159, 1983.
- [0275] Mann et al., *Cell*, 33:153-159, 1983.
- [0276] Markowitz et al., "A safe packaging line for gene transfer: Separating viral genes on two different plasmids," *J. Virol.*, 62:1120-1124, 1988.
- [0277] McCormack et al. *Hum. Gene Ther.* 8: 1263-1273, 1997.
- [0278] Miller et al., PCT Application WO 89/06700
- [0279] Miyoshi et al., *Proc. Natl. Acad. Sci. USA* 94:10319-10323.
- [0280] Mizukami et al., *Virology*, 217:124-130, 1996.
- [0281] Mulligan et al, *Proc. Nat'l Acad. Sci. USA*, 78: 2072, 1981.
- [0282] Mulligan, "The basic science of gene therapy," *Science*, 260:926-932, 1993.
- [0283] Myers, EP 0273085
- [0284] Naldini et al. *Science* 272:263-267.
- [0285] Nicolas and Rubenstein, "Retroviral vectors," In: *Vectors: A survey of molecular cloning vectors and their uses*, Rodriguez and Denhardt (eds.), Stoneham: Butterworth, pp. 493-513, 1988.
- [0286] Nicolas and Rubenstein, In: *Vectors: A survey of molecular cloning vectors and their uses*, Rodriguez and Denhardt (eds.), Stoneham: Butterworth, pp. 493-513, 1988.
- [0287] Nicolau and Sene, "Liposome-mediated DNA transfer in eukaryotic cells," *Biochim. Biophys. Acta*, 721:185-190, 1982.
- [0288] Nicolau et al., "Liposomes as carriers for in vivo gene transfer and expression," *Methods Enzymol.*, 149:157-176, 1987.
- [0289] O'Hare et al., *Proc. Nat'l Acad. Sci. USA*, 78: 1527, 1981.
- [0290] Ohara et al., *Proc. Nat'l Acad. Sci. USA*, 86: 5673-5677, 1989.
- [0291] Olsen, J. C., L. G. Johnson, J. M. Stutts, B. Sarkadi, J. R. Yankaskas, R. Swanstrom, and R. C. Boucher. 1992. Correction of the apical membrane chloride permeability defect in polarized cystic fibrosis airway epithelia following retroviral-mediated gene transfer. *Hum. Gene Ther.* 3:253-266.
- [0292] Olsen, Johnson, Wong-Sun, Moore, Swanstrom, and Boucher, "Retrovirus-mediated gene transfer to cystic fibrosis airway epithelial cells: effect of selectable marker sequences on long-term expression," *Nucleic Acids Res.*, 21(3):663-669, 1993.
- [0293] Ostrove et al., *Virology*, 113:532-533, 1981.
- [0294] Paskind et al., "Dependence of moloney murine leukemia virus production on cell growth," *Virology*, 67:242-248, 1975.
- [0295] Paskind et al., *Virology*, 67:242-248, 1975.
- [0296] Pelletier and Sonenberg, *Nature*, 334:320-325, 1988.
- [0297] Perales et al., "Gene transfer in vivo: Sustained expression and regulation of genes introduced into the liver by receptor-targeted uptake," *Proc. Natl. Acad. Sci.* 91:4086-4090, 1994.
- [0298] Pignon et al., *Hum. Mutat.*, 3: 126-132; 1994.
- [0299] Poeschla, E. M., F. W-Staal, and D. L. Looney. 1998. Efficient transduction of non-dividing human cells by feline immunodeficiency virus lentiviral vectors. *Nature Med.* 4:354-357.
- [0300] Ponnazhagan et al., *Hum. Gene Ther.*, 8:275-284, 1997a.
- [0301] Ponnazhagan et al., *J. Gen. Virol.*, 77:1111-1122, 1996.
- [0302] Post et al., *Cell*, 24:555-565, 1981.
- [0303] Potter et al., "Enhancer-dependent expression of human k immunoglobulin genes introduced into mouse pre-B lymphocytes by electroporation," *Proc. Nat. Acad. Sci. USA*, 81:7161-7165, 1984.
- [0304] Racher et al., *Biotechnology Techniques*, 9:169-174, 1995.
- [0305] Ragot et al.; "Efficient adenovirus-mediated transfer of a human minidystrophin-gene to skeletal muscle of mdx mice," *Nature*, 361:647-650, 1993.
- [0306] Renan, "Cancer genes: Current status, future prospects and applications in radiotherapy/oncology," *Radiother. Oncol*, 19:197-218, 1990.
- [0307] Renan, *Radiother. Oncol.*, 19:197-218, 1990.
- [0308] Rich et al., "Development and analysis of recombinant adenoviruses for gene therapy of cystic fibrosis," *Hum. Gene Ther.*, 4:461-476, 1993.
- [0309] Ridgeway, "Mammalian expression vectors," In: Rodriguez R L, Denhardt D T, ed. *Vectors: A survey of molecular cloning vectors and their uses*. Stoneham: Butterworth, pp. 467-492, 1988.
- [0310] Rippe et al., "DNA-mediated gene transfer into adult rat hepatocytes in primary culture," *Mol Cell Biol.*, 10:689-695, 1990.
- [0311] Roizman and Sears, In *Fields' Virology*, 3rd Edition, eds. Fields et al. (Raven Press, New York, N.Y.), pp.2231-2295, 1995.
- [0312] Rosenfeld et al., "In vivo transfer of the human cystic fibrosis transmembrane conductance regulator gene to the airway epithelium," *Cell*, 68:143-155, 1992.
- [0313] Roux et al., "A versatile and potentially general approach to the targeting of specific cell types by retroviruses: Application to the infection of human cells by means of major histocompatibility complex class I and class II antigens by mouse ecotropic murine leukemia virus-derived viruses," *Proc. Natl Acad. Sci. USA*, 86:9079-9083, 1989.
- [0314] Sambrook et al., *Molecular cloning: A laboratory manual*, 2d Ed., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1989.

- [0315] Samulski et al, *EMBO J.*, 10:3941-3950, 1991.
- [0316] Srivastava et al., *J. Virol.*, 45:555-564, 1983.
- [0317] Stinchcomb et al., *Nature*, 282: 39, 1979.
- [0318] Stratford-Perricaudet and Perricaudet, "Gene transfer into animals: the promise of adenovirus," p. 51-61, In: *HUMAN GENE TRANSFER*, Cohen-Haguenaer and Boiron (eds.), Editions John Libbey Eurotext, France, 1991.
- [0319] Stratford-Perricaudet et al., "Evaluation of the transfer and expression in mice of an enzyme-encoding gene using a human adenovirus vector," *Hum. Gene Ther.*, 1:241-256, 1990.
- [0320] Summers et al. "A manual of methods for baculovirus vectors and insect cell culture procedures," Texas Agriculture Experimental Station.
- [0321] Szybalska et al., *Proc. Nat'l Acad. Sci. USA*, 48:2026, 1962.
- [0322] Temin, "Retrovirus vectors for gene transfer: Efficient integration into and expression of exogenous DNA in vertebrate cell genome," In: *Gene Transfer*, Kucherlapati (ed.), New York, Plenum Press, pp. 149-188, 1986.
- [0323] Temin, In: *Gene Transfer*, Kucherlapati (ed.), New York: Plenum Press, pp. 149-188, 1986.
- [0324] Top et al., "Immunization with live types 7 and 4 adenovirus vaccines. II. Antibody response and protective effect against acute respiratory disease due to adenovirus type 7," *J Infect. Dis.*, 124:155-160, 1971.
- [0325] Tschemper et al., *Gene*, 10: 157, 1980.
- [0326] Tur-Kaspa et al., "Use of electroporation to introduce biologically active foreign genes into primary rat hepatocytes," *Mol. Cell Biol.*, 6:716-718, 1986.
- [0327] U.S. Pat. No. 4,683,195
- [0328] U.S. Pat. No. 4,683,202
- [0329] U.S. Pat. No. 4,800,159
- [0330] Varmus et al., "Retroviruses as mutagens: Insertion and excision of a nontransforming provirus alter the expression of a resident transforming provirus," *Cell*, 25:23-36, 1981.
- [0331] Wagner et al., *Proc. Natl. Acad. Sci.* 87(9):3410-3414, 1990.
- [0332] Wagner et al., *Science*, 260:1510-1513, 1993.
- [0333] Walker et al., *Proc. Nat'l Acad. Sci. USA* 89:392-396 1992.
- [0334] Wigler et al., *Cell*, 11: 223, 1977.
- [0335] Wigler et al., *Proc. Nat'l Acad. Sci. USA*, 77: 3567, 1980.
- [0336] Wong et al., "Appearance of P-lactamase activity in animal cells upon liposome mediated gene transfer," *Gene*, 10:87-94, 1980.
- [0337] Wu and Wu, "Evidence for targeted gene delivery to HepG2 hepatoma cells in vitro," *Biochem.*, 27:887-892, 1988.
- [0338] Wu and Wu, "Receptor-mediated in vitro gene transfections by a soluble DNA carrier system," *J. Biol. Chem.*, 262:4429-4432, 1987.
- [0339] Wu and Wu, *Adv. Drug Delivery Rev.*, 12:159-167, 1993.
- [0340] Wu et al., *Genomics*, 4:560, 1989.
- [0341] Yang et al., "In vivo and in vitro gene transfer to mammalian somatic cells by particle bombardment," *Proc. Natl. Acad. Sci USA*, 87:9568-9572, 1990.
- [0342] Zelenin et al., "High-velocity mechanical DNA transfer of the chloramphenicol acetyltransferase gene into rodent liver, kidney and mammary gland cells in organ explants and in vivo," *FEBS Lett.*, 280:94-96, 1991.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 193

<210> SEQ ID NO 1

<211> LENGTH: 4800

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic plasmid

<400> SEQUENCE: 1

```

aagcttacct cgatttgagg acgttacaag tattactggt aaggagcgta gattaaaaa      60
tgaaattgaa aatgaattat tagaattggc ttaaataaac agaatcacca aaaaggaata      120
gagtatgaag tttggaata tttgtttttc gtatcaacca ccaggtgaaa ctcataagct      180
aagtaatgga tcgctttggt cggcttggtc tcgcctcaga agagtagggt ttgatacata      240
ttggacctta gaacatcatt ttacagagtt tggctctacg ggaaatttat ttgttgctgc      300

```

-continued

ggctaacctg	ttaggaagaa	ctaaaacatt	aatggttggc	actatggggg	ttgttattcc	360
gacagcacac	ccagttcgac	agttagaaga	cgttttatta	ttagatcaaa	tgtcgaaagg	420
tcgttttaat	tttgaaccg	ttcgagggt	ataccataaa	gattttcgag	tatttgggtg	480
tgatattgaa	gagtctcgag	caattactca	aaatttctac	cagatgataa	tggaaagctt	540
acagacagga	accattagct	ctgatagtga	ttacattcaa	tttcctaagg	ttgatgtata	600
tcccaaagtg	tactcaaaaa	atgtaccaac	ctgtatgact	gctgagtccg	caagtacgac	660
agaatggcta	gcaatacaag	ggctaccaat	ggttcttagt	tggattattg	gtactaatga	720
aaaaaaagca	cagatggaac	tctataatga	aattgcgaca	gaatatggtc	atgatataatc	780
taaaatagat	cattgtatga	cttatatttg	ttctgttgat	gatgatgcac	aaaaggcgca	840
agatgtttgt	cgggagtctt	tgaaaaattg	gtatgactca	tatgtaaag	cgaccaatat	900
ctttaatgat	agcaatcaaa	ctcgtgggta	tgattatcat	aaaggccaat	ggcgtgattt	960
tgttttacaa	ggacatacaa	acaccaatcg	acgtgttgat	tatagcaatg	gtattaacct	1020
tgtaggcact	cctgagcagt	gtattgaaat	cattcaacgt	gatattgatg	caacgggtat	1080
tacaaacatt	acatgcgat	ttgaagctaa	tggaaactgaa	gatgaaataa	ttgcttccat	1140
gcgacgcttt	atgacacaag	tcgctccttt	cttaaaagaa	cctaaataaa	ttacttattt	1200
gatactagag	ataataagga	acaagttatg	aaatttggat	tattttttct	aaactttcag	1260
aaagatggaa	taacatctga	agaaacggtg	gataaatatg	taaagactgt	cacgttaatt	1320
gattcaacta	aatatcattt	taatactgcc	tttggttaatg	aacatcactt	ttcaaaaaat	1380
ggatttgggt	gagcacctat	taccgcagct	ggttttttat	taggggtaac	aaataaatta	1440
catattgggt	cattaaatca	agtaattacc	acctatcacc	ctgtacgtgt	agcagaagaa	1500
gccagtttat	tagatcaaat	gtcagagggg	cgcttcattc	ttggtttttag	tgactgcgaa	1560
agtgatttcg	aatggaatt	ttttagacgt	catatctcat	caaggcaaca	acaatttgaa	1620
gcatgctatg	aaataattaa	tgacgcatta	actacagggt	attgtcatcc	ccaaaacgac	1680
ttttatgatt	ttcaaagggt	ttcaattaat	ccacactggt	acagtgagaa	tggacctaa	1740
caatatgtat	ccgctacatc	aaaagaagtc	gtcatgtggg	cagcgaaaaa	ggcactgcct	1800
ttaacattta	agtgggagga	taatttagaa	accaagaagc	gctatgcaat	tctatataat	1860
aaaacagcac	aacaatatgg	tattgatatt	tcggatgttg	atcatcaatt	aactgtaatt	1920
gcgaacttaa	atgctgatag	aagtacggct	caagaagaag	tgagagaata	cttaaaagac	1980
tatatcactg	aaacttacc	tcaaatggac	agagatgaaa	aaattaactg	cattattgaa	2040
gagaatgcag	ttgggtctca	tgatgactat	tatgaatcga	caaaattagc	agtggaaaaa	2100
acaggttcta	aaaatatttt	attatccttt	gaatcaatgt	ccgatattaa	agatgtaaaa	2160
gatattattg	atatgttgaa	ccaaaaaatc	gaaatgaatt	taccataata	aaattaaagg	2220
caatttctat	attagattgc	ctttttgggg	atcctctaga	aatattttat	ctgattaata	2280
agatgagaat	tcactggccg	tcgttttaca	acgtcgtgac	tgggaaaacc	ctggcgttac	2340
ccaacttaat	cgcttgcag	cacatcccc	tttccgacg	tggcgtaata	gcgaagaggc	2400
ccgcaccgat	cgcccttccc	aacagttgcg	cagcctgaat	ggcgaatggc	gcctgatgcg	2460
gtattttctc	cttacgcatc	tgtgcggtat	ttcacaccgc	atatgggtgca	ctctcagtac	2520
aatctgctct	gatgccgat	agttaagcca	gccccgacac	ccgccaacac	ccgctgacgc	2580

-continued

gccctgacgg	gcttgtctgc	tcccggcatc	cgcttacaga	caagctgtga	ccgtctccgg	2640
gagctgcatg	tgtcagaggt	tttcaccgtc	atcaccgaaa	cgcgcgagac	gaaagggcct	2700
cgtgatacgc	ctatTTTTat	aggTTaatgt	catgataata	atggTTTTctt	agacgtcagg	2760
tggcactttt	cggggaaatg	tgcgcggaac	ccctatTTTgt	ttatTTTTtct	aaaaagcttc	2820
acgctgccgc	aagcactcag	ggcgcaaggg	ctgctaaagg	aagcggaaca	cgtagaaagc	2880
cagtccgcag	aaacggTgct	gaccccggat	gaatgtcagc	tactgggcta	tctggacaag	2940
ggaaaacgca	agcgcaaaga	gaaagcaggT	agctTgcagt	gggctTacat	ggcgatagct	3000
agactgggcg	gttttatgga	cagcaagcga	accggaattg	ccagctgggg	cgccctctgg	3060
taaggTtggg	aagccctgca	aagTaaactg	gatggctTtc	ttgccgcaa	ggatctgatg	3120
gcgcagggga	tcaagatctg	atcaagagac	aggatgagga	tcgTttcgca	tgattgaaca	3180
agatggattg	cacgcaggTt	ctccggccgc	ttgggtggag	aggctattcg	gctatgactg	3240
ggcacaacag	acaatcggtc	gctctgatgc	cgccgtgttc	cggtgtcag	cgcaggggcg	3300
cccgtTtctt	tttgtcaaga	ccgacctgtc	cggtgccttg	aatgaactgc	aggacgaggc	3360
agcgcggcta	tcgtggctgg	ccacgacggg	cgTtccttgc	gcagctgtgc	tcgacgtTgt	3420
cactgaagcg	ggaagggact	ggctgctatt	gggogaagTg	ccggggcagg	atctcctgtc	3480
atctcacctt	gctcctgccg	agaaagtatc	catcatggct	gatgcaatgc	ggcggctgca	3540
tacgcttgat	ccggctacct	gcccattcga	ccaccaagcg	aaacatcgca	tcgagcgagc	3600
acgtactcgg	atggaagccg	gtcttgtcga	tcaggatgat	ctggacgaag	agcatcaggg	3660
gctcgcgcca	gccgaactgt	tcgccaggct	caaggcgcgc	atgcccgacg	gcgaggatct	3720
cgTcgtgacc	catggcgatg	cctgcttgcc	gaatatcatg	gtggaaaatg	gccgcttttc	3780
tggattcatc	gactgtggcc	ggctgggtgt	ggcggaccgc	tatcaggaca	tagcgttggc	3840
taccctgat	attgctgaag	agctTggcgg	cgaatgggct	gaccgcttcc	tcgtgcttta	3900
cggTatcgcc	gctcccgatt	cgcagcgcac	cgcttctat	cgcttcttg	acgagTtctt	3960
ctgagcggga	ctctggggTt	cgaaatgacc	gaccaagcga	cgcccaacct	gcatcacga	4020
gatttcgatt	ccaccgccgc	cttctatgaa	aggTtgggct	tcggaatcgt	tttccgggac	4080
gccggctgga	tgatcctcca	gcgcggggat	ctcatgctgg	agTtcttcgc	ccaccccggg	4140
catgacaaa	atcccttaac	gtgagTtttc	gttccactga	gcgtcagacc	ccgtagaaaa	4200
gatcaaagga	tcttcttgag	atcctTTTTt	tctgcgcgta	atctgctgct	tgcaaacaaa	4260
aaaaccaccg	ctaccagcgg	tggTttgttt	gccggatcaa	gagctaccaa	ctctttttcc	4320
gaaggtaact	ggcttcagca	gagcgcagat	accaaatact	gtccttctag	tgtagccgta	4380
gttaggccac	cacttcaaga	actctgtagc	accgcctaca	tacctcgctc	tgctaactct	4440
gttaccagtg	gctgctgcca	gtggcgataa	gtcgtgtctt	accgggttgg	actcaagacg	4500
atagttaccg	gataagggcg	agcggTcggg	ctgaacgggg	ggTtctgTca	cacagcccag	4560
cttgagcga	acgacctaca	ccgaactgag	atacctacag	cgtgagctat	gagaaagcgc	4620
cacgcttccc	gaagggagaa	aggcggacag	gtatccggta	agcggcaggg	tcggaacagg	4680
agagcgcacg	agggagcttc	cagggggaaa	cgctggTat	ctttatagtc	ctgtcgggtt	4740
tcgccacctc	tgacttgagc	gtcgatTTTT	gtgatgctcg	tcaggggggc	ggagcctatg	4800

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 2

aagcttacct cgatttgagg acgttacaag tattactggt aaggagcgta 50

<210> SEQ ID NO 3
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 3

gattaaaaa tgaaattgaa aatgaattat tagaattggc ttaaataaac 50

<210> SEQ ID NO 4
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 4

agaatcacca aaaaggaata gagtatgaag tttggaaata tttgtttttc 50

<210> SEQ ID NO 5
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 5

gtatcaacca ccagtgaaa ctcataagct aagtaatgga tcgctttggt 50

<210> SEQ ID NO 6
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 6

cggcttggtgta tcgcctcaga agagtaggggt ttgatacata ttggacctta 50

<210> SEQ ID NO 7
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 7

gaacatcatt ttacagagtt tggctttacg ggaaatttat ttgttgctgc 50

<210> SEQ ID NO 8
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 8

ggctaacctg ttaggaagaa ctaaaacatt aaatgttggc actatggggg 50

<210> SEQ ID NO 9

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 9

ttgttattcc gacagcacac ccagttcgac agttagaaga cgttttatta 50

<210> SEQ ID NO 10

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 10

ttagatcaaa tgtcgaaagg tcgttttaaat tttggaaccg ttcgagggct 50

<210> SEQ ID NO 11

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 11

ataccataaa gattttcgag tatttgggtgt tgatatggaa gagtctcgag 50

<210> SEQ ID NO 12

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 12

caattactca aaatttctac cagatgataa tggaaagctt acagacagga 50

<210> SEQ ID NO 13

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 13

accattagct ctgatagtgga ttacattcaa tttcctaagg ttgatgtata 50

<210> SEQ ID NO 14

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 14

-continued

tcccaaagtg tactcaaaaa atgtaccaac ctgtatgact gctgagtccg 50

<210> SEQ ID NO 15
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 15

caagtacgac agaatggcta gcaatacaag ggctaccaat ggttcttagt 50

<210> SEQ ID NO 16
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 16

tggattattg gtactaatga aaaaaagca cagatggaac tctataatga 50

<210> SEQ ID NO 17
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 17

aattgcgaca gaatatggtc atgatatatc taaaatagat cattgtatga 50

<210> SEQ ID NO 18
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 18

cttatatttg ttctgttgat gatgatgcac aaaaggcgca agatgtttgt 50

<210> SEQ ID NO 19
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 19

cgggagtttc tgaaaaattg gtatgactca tatgtaaagtg cgaccaatat 50

<210> SEQ ID NO 20
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 20

ctttaatgat agcaatcaaa ctcgtgggta tgattatcat aaaggtcaat 50

<210> SEQ ID NO 21

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 21

ggcgtgattt tgttttacaa ggacatacaa acaccaatcg acgtgttgat 50

<210> SEQ ID NO 22
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 22

tatagcaatg gtattaacct tgtaggcact cctgagcagt gtattgaaat 50

<210> SEQ ID NO 23
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 23

cattcaacgt gatattgatg caacgggtat tacaacatt acatgcggat 50

<210> SEQ ID NO 24
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 24

ttgaagctaa tggaactgaa gatgaaataa ttgcttccat gcgacgcttt 50

<210> SEQ ID NO 25
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 25

atgacacaag tcgctccttt cttaaagaa cctaaataaa ttacttattt 50

<210> SEQ ID NO 26
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 26

gatactagag ataataagga acaagttatg aaatttgat tattttttct 50

<210> SEQ ID NO 27
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 27

aaactttcag aaagatggaa taacatctga agaaacgttg gataatatgg 50

<210> SEQ ID NO 28
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 28

taaagactgt cacgtaatt gattcaacta aatatcattt taatactgcc 50

<210> SEQ ID NO 29
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 29

tttgtaatg aacatcactt ttcaaaaaat ggtattggtg gagcacctat 50

<210> SEQ ID NO 30
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 30

taccgcagct ggtttttat tagggtaac aaataaatta catattggtt 50

<210> SEQ ID NO 31
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 31

cattaaatca agtaattacc acccatcacc ctgtacgtgt agcagaagaa 50

<210> SEQ ID NO 32
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 32

gccagtttat tagatcaaat gtcagagga cgcttcattc ttggttttag 50

<210> SEQ ID NO 33
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 33

-continued

tgactgcgaa agtgatttcg aatggaatt ttttagacgt catatctcat 50

<210> SEQ ID NO 34
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 34

caaggcaaca acaatttgaa gcatgctatg aaataattaa tgacgcatta 50

<210> SEQ ID NO 35
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 35

actacaggtt attgtcatcc caaaacgac ttttatgatt ttccaaggt 50

<210> SEQ ID NO 36
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 36

ttcaattaat ccacactggt acagtgagaa tggacctaa caatatgtat 50

<210> SEQ ID NO 37
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 37

ccgctacatc aaaagaagtc gtcattgtgg cagcgaaaaa ggcactgcct 50

<210> SEQ ID NO 38
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 38

ttaacattta agtgggagga taatttagaa accaaagaac gctatgcaat 50

<210> SEQ ID NO 39
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 39

tctatataat aaaacagcac aacaatatgg tattgatatt tcggatggtg 50

<210> SEQ ID NO 40

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 40

atcatcaatt aactgtaatt gcgaacttaa atgctgatag aagtacggct 50

<210> SEQ ID NO 41
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 41

caagaagaag tgagagaata cttaaagac tatatcactg aaacttacc 50

<210> SEQ ID NO 42
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 42

tcaaatggac agagatgaaa aaattaactg cattattgaa gagaatgcag 50

<210> SEQ ID NO 43
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 43

ttgggtctca tgatgactat tatgaatcga caaaattagc agtggaaaa 50

<210> SEQ ID NO 44
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 44

acaggtctta aaaatatttt attatccttt gaatcaatgt ccgatattaa 50

<210> SEQ ID NO 45
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 45

agatgtaaaa gatattattg atatgttgaa ccaaaaaatc gaaatgaatt 50

<210> SEQ ID NO 46
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 46
taccataata aaattaaagg caatttctat attagattgc ctttttgggg 50

<210> SEQ ID NO 47
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 47
atcctctaga aatattttat ctgattaata agatgagaat tcaactggccg 50

<210> SEQ ID NO 48
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 48
tcgttttaca acgtcgtgac tgggaaaacc ctggcggttac ccaacttaat 50

<210> SEQ ID NO 49
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 49
cgccttgacg cacatcccc tttcgccagc tggcgtaata gcgaagaggg 50

<210> SEQ ID NO 50
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 50
ccgcaccgat cgccttccc aacagttgag cagcctgaat ggccaatggc 50

<210> SEQ ID NO 51
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 51
gcctgatgag gtattttctc cttacgcacg tgtgcgggat ttcacaccgc 50

<210> SEQ ID NO 52
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 52

-continued

atatggtgca ctctcagtac aatctgctct gatgccgcat agttaagcca 50

<210> SEQ ID NO 53
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 53

gccccgacac ccgccaacac ccgctgacgc gccctgacgg gcttgtctgc 50

<210> SEQ ID NO 54
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 54

tcccggcatc cgcttacaga caagctgtga ccgtctccgg gagctgcatg 50

<210> SEQ ID NO 55
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 55

tgtcagaggt tttcaccgtc atcaccgaaa cgcgcgagac gaaagggcct 50

<210> SEQ ID NO 56
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 56

cgtgatagc ctatttttat aggttaatgt catgataata atggtttctt 50

<210> SEQ ID NO 57
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 57

agacgtcagg tggcactttt cgggaaatg tgcgcggaac ccctatttgt 50

<210> SEQ ID NO 58
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 58

ttatttttct aaaaagcttc acgctgccgc aagcactcag ggcgcaaggg 50

<210> SEQ ID NO 59

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 59

ctgctaaagg aagcgaaca cgtagaaagc cagtccgcag aaacggtgct 50

<210> SEQ ID NO 60
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 60

gaccccgat gaatgtcagc tactgggcta tctggacaag ggaaaacgca 50

<210> SEQ ID NO 61
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 61

agcgcaaaga gaaagcaggt agcttgcaagt gggcttacat ggcgatagct 50

<210> SEQ ID NO 62
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 62

agactgggcg gttttatgga cagcaagcga accggaattg ccagctgggg 50

<210> SEQ ID NO 63
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 63

cgccctctgg taaggttggg aagccctgca aagtaaactg gatggctttc 50

<210> SEQ ID NO 64
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 64

ttgccgcaa ggatctgatg ggcagggga tcaagatctg atcaagagac 50

<210> SEQ ID NO 65
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 65

aggatgagga tcgtttcgca tgattgaaca agatggattg cacgcaggtt 50

<210> SEQ ID NO 66

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 66

ctccggccgc ttgggtggag aggctattcg gctatgactg ggcacaacag 50

<210> SEQ ID NO 67

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 67

acaatcggct gctctgatgc cgccgtgttc cggctgtcag cgcaggggcg 50

<210> SEQ ID NO 68

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 68

cccgttctt tttgtcaaga ccgacctgtc cggtgccctg aatgaactgc 50

<210> SEQ ID NO 69

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 69

aggacgaggc agcgcggcta tcgtggctgg ccacgacggg cgttccttgc 50

<210> SEQ ID NO 70

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 70

gggcgaagtg ccggggcagg atctcctgtc atctcacctt gctcctgccg 50

<210> SEQ ID NO 71

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 71

-continued

gggcgaagtg ccggggcagg atctcctgtc atctcacctt gctcctgccg 50

<210> SEQ ID NO 72
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 72

agaaagtatc catcatggct gatgcaatgc ggcggctgca tacgcttgat 50

<210> SEQ ID NO 73
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 73

ccggctacct gccattcga ccaccaagcg aaacatcgca tcgagcgagc 50

<210> SEQ ID NO 74
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 74

acgtactcgg atggaagccg gtcttgctga tcaggatgat ctggacgaag 50

<210> SEQ ID NO 75
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 75

agcatcaggg gctcgcgcca gccgaactgt tcgccaggct caaggcgcgc 50

<210> SEQ ID NO 76
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 76

atgcccgacg gcgaggatct cgtcgtgacc catggcgatg cctgcttgcc 50

<210> SEQ ID NO 77
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 77

gaatatcatg gtggaaaatg gccgcttttc tggattcatc gactgtggcc 50

<210> SEQ ID NO 78

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 78

ggctgggtgt ggcggaccgc tatcaggaca tagcgttggc taccctgat 50

<210> SEQ ID NO 79
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 79

attgctgaag agcttggcgg cgaatgggct gaccgcttcc tcgtgcttta 50

<210> SEQ ID NO 80
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 80

cggatcggc gctcccgatt cgcagcgcac cgccttctat cgccttcttg 50

<210> SEQ ID NO 81
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 81

acgagttctt ctgagcggga ctctgggggtt cgaaatgacc gaccaagcga 50

<210> SEQ ID NO 82
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 82

cgcccaacct gccatcacga gatttcgatt ccaccgccgc cttctatgaa 50

<210> SEQ ID NO 83
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 83

aggttgggct tcggaatcgt tttccgggac gccggctgga tgatcctcca 50

<210> SEQ ID NO 84
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 84

gcgcgggat ctcatgctgg agttcttcgc ccaccccggg catgaccaa 50

<210> SEQ ID NO 85

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 85

atcccttaac gtgagtttc gttccactga gcgtcagacc ccgtagaaaa 50

<210> SEQ ID NO 86

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 86

gatcaaagga tcttcttgag atcctttttt totgdcgta atctgctgct 50

<210> SEQ ID NO 87

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 87

tgcaaacaaa aaaaccaccg ctaccagcgg tggtttggtt gccggatcaa 50

<210> SEQ ID NO 88

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 88

gagctaccaa ctctttttcc gaaggtaact ggcttcagca gagcgcagat 50

<210> SEQ ID NO 89

<211> LENGTH: 51

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 89

accaaatact gtccttctag tgtagccgta gttaggccac cacttcaatg a 51

<210> SEQ ID NO 90

<211> LENGTH: 50

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 90

-continued

actctgtagc accgcctaca tacctcgctc tgctaatacct gttaccagtg 50

<210> SEQ ID NO 91
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 91

gctgctgccca gtggcgataa gtcgtgtcctt accgggttgg actcaagacg 50

<210> SEQ ID NO 92
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 92

atagttaccg gataaggcgc agcggtcggg ctgaacgggg ggttcgtgca 50

<210> SEQ ID NO 93
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 93

cacagcccag cttggagcga acgacctaca ccgaactgag atacctacag 50

<210> SEQ ID NO 94
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 94

cgtgagctat gagaaagcgc cacgcttccc gaagggagaa aggcggacag 50

<210> SEQ ID NO 95
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 95

gtatccgta agcggcaggg tcggaacagg agagcgcacg agggagcttc 50

<210> SEQ ID NO 96
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 96

cagggggaaa cgcctggtat ctttatagtc ctgtcggggtt tcgccacctc 50

<210> SEQ ID NO 97

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 97

tgacttgagc gtcgattttt gtgatgctcg tcaggggggc ggagcctatg 50

<210> SEQ ID NO 98
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 98

catcaca aaa atcgacgctc aagtcagagg tggcgaaacc cgacaggact 50

<210> SEQ ID NO 99
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 99

ataaagatac caggcgtttc cccctggaag ctccctcgtg cgctctcctg 50

<210> SEQ ID NO 100
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 100

ttccgaccct gccgcttacc ggatacctgt cgcctttct cccttcggga 50

<210> SEQ ID NO 101
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 101

agcgtggcgc tttctcatag ctcacgctgt aggtatctca gttcgggtga 50

<210> SEQ ID NO 102
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 102

ggtcgttcgc tccaagctgg gctgtgtgca cgaaccccc gttcagcccg 50

<210> SEQ ID NO 103
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 103
accgctgcg cttatccggt aactatcgtc ttgagtccaa cccggtaaga 50

<210> SEQ ID NO 104
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 104
cacgacttat cgccactggc agcagccact ggtaacagga ttagcagagc 50

<210> SEQ ID NO 105
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 105
gaggtatgta ggcggtgcta cagagttcct gaagtgggtg cctaactacg 50

<210> SEQ ID NO 106
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 106
gctacactag aaggacagta tttggtatct gcgctctgct gaagccagtt 50

<210> SEQ ID NO 107
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 107
accttcggaa aaagagttgg tagctcttga tccggcaaac aaaccaccgc 50

<210> SEQ ID NO 108
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 108
tggtagcggg ggttttttg tttgcaagca gcagattacg cgcagaaaaa 50

<210> SEQ ID NO 109
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 109

-continued

aaggatctca agaagatcct ttgatctttt ctacggggtc tgacgctcag 50

<210> SEQ ID NO 110
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 110

tggaacgaaa actcacgta agggattttg gtcatgcccg gggtagggcga 50

<210> SEQ ID NO 111
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 111

agaactccag catgagatcc ccgcgctgga ggatcatcca gccggcgtcc 50

<210> SEQ ID NO 112
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 112

cggaaaacga ttccgaagcc caacctttca tagaaggcgg cggtagaatc 50

<210> SEQ ID NO 113
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 113

gaaatctcgt gatggcaggt tgggcgtcgc ttggtcggtc atttcgaacc 50

<210> SEQ ID NO 114
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 114

ccagagtccc gctcagaaga actcgtcaag aaggcgatag aaggcgatgc 50

<210> SEQ ID NO 115
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 115

gctgcgaatc gggagcggcg ataccgtaaa gcacgaggaa gcggtcagcc 50

<210> SEQ ID NO 116

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 116

cattcgccgc caagctcttc agcaatatca cgggtagcca acgctatgtc 50

<210> SEQ ID NO 117
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 117

ctgatagcgg tccgccacac ccagccggcc acagtcgatg aatccagaaa 50

<210> SEQ ID NO 118
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 118

agcggccatt ttccacatg atattcggca agcagggcatc gccatgggtc 50

<210> SEQ ID NO 119
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 119

acgacgagat cctcgccgtc gggcatgcgc gccttgagcc tggcgaacag 50

<210> SEQ ID NO 120
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 120

ttcggctggc gcgagcccct gatgctcttc gtccagatca tcctgatcga 50

<210> SEQ ID NO 121
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 121

caagaccggc ttccatccga gtacgtgctc gctcogatgc atgtttcgtc 50

<210> SEQ ID NO 122
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 122
tgggtggtcga atgggcaggt agccggatca agcgtatgca gccgccgcat 50

<210> SEQ ID NO 123
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 123
tgcacagcc atgatggata ctttctcggc aggagcaagg tgagatgaca 50

<210> SEQ ID NO 124
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 124
ggagatcctg ccccggaact tcgcccaata gcagccagtc cttcccgt 50

<210> SEQ ID NO 125
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 125
tcagtgaaa cgtcgagcac agctgcgcaa ggaacgcccg tcgtggccag 50

<210> SEQ ID NO 126
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 126
ccacgatagc cgcgctgcct cgtcctgcag ttcattcagg gcaccggaca 50

<210> SEQ ID NO 127
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 127
ggtcggcttt gacaaaaaga accgggccc cctgcgctga cagccggaac 50

<210> SEQ ID NO 128
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 128

-continued

acggcgcat cagagcagcc gattgtctgt tgtgcccagt catagccgaa 50

<210> SEQ ID NO 129
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 129

tagcctctcc acccaagcgg ccggagaacc tgcgtgcaat ccatcttgtt 50

<210> SEQ ID NO 130
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 130

caatcatgcg aaacgatcct catcctgtct cttgatcaga tcttgatccc 50

<210> SEQ ID NO 131
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 131

ctgcgccatc agatccttgg cggcaagaaa gccatccagt ttactttgca 50

<210> SEQ ID NO 132
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 132

gggcttccca acctaccag agggcgcccc agctggcaat tccggttcgc 50

<210> SEQ ID NO 133
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 133

ttgctgtcca taaaaccgcc cagtctagct atcgccatgt aagcccactg 50

<210> SEQ ID NO 134
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 134

caagctacct gctttctctt tgcgcttgcg ttttccttg tccagatagc 50

<210> SEQ ID NO 135

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 135

ccagtagctg acattcatcc ggggtcagca ccgtttctgc ggactggctt 50

<210> SEQ ID NO 136
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 136

tctacgtggt ccgcttcctt tagcagccct tgcgccctga gtgcttgagg 50

<210> SEQ ID NO 137
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 137

cagcgtgaag ctttttagaa aaataaacia ataggggttc cgcgcacatt 50

<210> SEQ ID NO 138
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 138

tccccgaaaa gtgccacctg acgtctaaga aaccattatt atcatgacat 50

<210> SEQ ID NO 139
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 139

taacctataa aataggcgt atcacgaggc cctttcgtct cgcgcgtttc 50

<210> SEQ ID NO 140
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 140

ggtgatgacg gtgaaaacct ctgacacatg cagctcccgg agacggtcac 50

<210> SEQ ID NO 141
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 141
agcttgctctg taagcggatg ccgggagcag acaagcccgt cagggcgcgt 50

<210> SEQ ID NO 142
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 142
cagcgggtgt tggcgggtgt cggggctggc ttaactatgc ggcatcagag 50

<210> SEQ ID NO 143
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 143
cagattgtac tgagagtgc ccatatgcgg tgtgaaatac cgcacagatg 50

<210> SEQ ID NO 144
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 144
cgtaaggaga aaataccgca tcaggcgcca ttogccattc aggctgcgca 50

<210> SEQ ID NO 145
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 145
actgttggga agggcgatcg gtgcgggcct cttoctatt acgccagctg 50

<210> SEQ ID NO 146
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 146
gcgaaagggg gatgtgctgc aaggcgatta agttgggtaa cgccagggtt 50

<210> SEQ ID NO 147
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 147

-continued

ttcccagtca cgacgttgta aaacgacggc cagtgaattc tcatcttatt 50

<210> SEQ ID NO 148
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 148

aatcagataa aatatttcta gaggatcccc aaaaaggcaa tctaataatag 50

<210> SEQ ID NO 149
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 149

aaattgcctt taattttatt atggtaaatt catttcgatt ttttggttca 50

<210> SEQ ID NO 150
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 150

acatatcaat aatatctttt acatctttaa tatcggacat tgattcaaag 50

<210> SEQ ID NO 151
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 151

gataataaaa tatttttaga ccctgttttt tccactgcta attttgtcga 50

<210> SEQ ID NO 152
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 152

ttcataatag tcatcatgag acccaactgc atttctttca ataatgcagt 50

<210> SEQ ID NO 153
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 153

taattttttc atctctgtcc atttgagggt aagtttcagt gatatagtct 50

<210> SEQ ID NO 154

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 154

tttaagtatt ctctcacttc ttcttgagcc gtacttctat cagcatttaa 50

<210> SEQ ID NO 155
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 155

gttcgcaatt acagttaatt gatgatcaac atccgaaata tcaataccat 50

<210> SEQ ID NO 156
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 156

attggtgtgc tgttttatta tatagaattg catagcgttc tttggtttct 50

<210> SEQ ID NO 157
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 157

aaattatcct cccacttaaa tgtaaaggc agtgcctttt tcgctgcca 50

<210> SEQ ID NO 158
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 158

catgacgact tcttttgatg tagcggatac atattgctta ggtccattct 50

<210> SEQ ID NO 159
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 159

cactgtaaca gtgtggatta attgaaacct ttggaaaatc ataaaagtcg 50

<210> SEQ ID NO 160
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 160
ttttgggat gacaataacc tgtagttaat gcgtcattaa ttatttcata 50

<210> SEQ ID NO 161
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 161
gcatgcttca aattgttgtt gccttgatga gatatgacgt ctaaaaaatt 50

<210> SEQ ID NO 162
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 162
ccatttcgaa atcactttcg cagtcactaa aaccaagaat gaagcgtccc 50

<210> SEQ ID NO 163
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 163
tctgacattt gatctaataa actggcttct tctgctacac gtacagggtg 50

<210> SEQ ID NO 164
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 164
atgggtggta attacttgat ttaatgaacc aatatgtaat ttatttgta 50

<210> SEQ ID NO 165
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 165
accctaataa aaaaccagct gcggtaatag gtgctccaac aataccattt 50

<210> SEQ ID NO 166
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide
<400> SEQUENCE: 166

-continued

tttgaaaagt gatgttcatt aacaaaggca gtattaaaat gatatttagt 50

<210> SEQ ID NO 167
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 167

tgaatcaatt aacgtgacag tctttacat attatccaac gtttcttcag 50

<210> SEQ ID NO 168
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 168

atgttattcc atctttctga aagtttagaa aaaataatcc aaatttcata 50

<210> SEQ ID NO 169
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 169

acttgttcct tattatctct agtatcaaat aagtaattta ttaggttct 50

<210> SEQ ID NO 170
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 170

tttaagaaag gagcgacttg tgtcataaag cgtcgcacatg aagcaattat 50

<210> SEQ ID NO 171
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 171

ttcatcttca gttccattag cttcaaatcc gcatgtaatg tttgtaatac 50

<210> SEQ ID NO 172
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 172

ccggttgcac aatatcacgt tgaatgattt caatacactg ctcaggagtg 50

<210> SEQ ID NO 173

-continued

<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 173

cctacagggt taataccatt gctataatca acacgtcgat tgggtgttgt 50

<210> SEQ ID NO 174
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 174

atgtccttgt aaaacaaaat cacgccattg acctttatga taatcataac 50

<210> SEQ ID NO 175
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 175

cacgagtttg attgctatca ttaaagatat tggtcgcatt tacatatgag 50

<210> SEQ ID NO 176
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 176

tcataccaat ttttcagaaa ctcccgacaa acatcttgcg ctttttgtgc 50

<210> SEQ ID NO 177
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 177

atcatcatca acagaacaaa tataagtcac acaatgatct attttagata 50

<210> SEQ ID NO 178
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 178

tatcatgacc atattctgtc gcaatttcat tatagagttc catctgtgct 50

<210> SEQ ID NO 179
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 179

ttttttcat tagtaccaat aatccaacta agaaccattg gtagcccttg 50

<210> SEQ ID NO 180
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 180

tattgctagc cattctgtcg tacttgcgga ctcagcagtc atacaggttg 50

<210> SEQ ID NO 181
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 181

gtacattttt tgagtacact ttgggatata catcaacctt aggaaattga 50

<210> SEQ ID NO 182
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 182

atgtaatcac tatcagagct aatggttcct gtotgtaagc tttccattat 50

<210> SEQ ID NO 183
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 183

catctggtag aaattttgag taattgctcg agactottcc atatcaacac 50

<210> SEQ ID NO 184
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 184

caaatactcg aaaatcttta tggatatagcc ctogaacggt tccaaaatta 50

<210> SEQ ID NO 185
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 185

-continued

aaacgacctt tcgacatttg atctaataat aaaacgtctt ctaactgtcg 50

<210> SEQ ID NO 186
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 186

aactgggtgt gctgtcggaa taacaacccc catagtgccca acatttaatg 50

<210> SEQ ID NO 187
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 187

ttttagttct tcctaacagg ttagccgcag caacaaataa atttcccgta 50

<210> SEQ ID NO 188
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 188

agaccaaact ctgtaaaatg atgttctaag gtccaatatg tatcaaacc 50

<210> SEQ ID NO 189
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 189

tactcttctg aggcgatacc aagccgaaca aagcgatcca ttacttagct 50

<210> SEQ ID NO 190
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 190

tatgagtttc acctggtggt tgatacgaaa aacaaatatt tccaaacttc 50

<210> SEQ ID NO 191
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 191

atactctatt cctttttggt gattctgttt atttaagcca attctaataa 50

<210> SEQ ID NO 192

-continued

<211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 192

ttcattttca atttcatttt ttaatctacg ctcttaaca gtaatacttg 50

<210> SEQ ID NO 193
 <211> LENGTH: 50
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic Oligonucleotide

<400> SEQUENCE: 193

taacgtcctc aaatcgaggt aagcttcata ggctccgcc cctgacgag 50

1. A method for the synthesis of a replication-competent, double-stranded polynucleotide, wherein said polynucleotide comprises an origin of replication, a first coding region and a first regulatory element directing the expression of said first coding region, comprising the steps of:

- (a) generating a first set of oligonucleotides corresponding to the entire plus strand of said double-stranded polynucleotide;
- (b) generating a second set of oligonucleotides corresponding to the entire minus strand of said double-stranded polynucleotide; and
- (c) annealing said first and said second set of oligonucleotides;

wherein each of said oligonucleotides of said second set of oligonucleotides overlaps with and hybridizes to two complementary oligonucleotides of said first set of oligonucleotides, except that two oligonucleotides at a 5' or 3' end of said double-stranded polynucleotide will hybridize with only one complementary oligonucleotide.

2. The method of claim 1, further comprising the step of treating said annealed oligonucleotides with a ligating enzyme to generate continuous strands of said double-stranded polynucleotide.

3. The method of claim 1, further comprising the step of amplifying said double-stranded polynucleotide.

4. The method of claim 1, wherein said double-stranded polynucleotide comprises 100, 200, 300, 400, 500, 600, 700, 800, 900, 1000, 5000, 10×10^3 , 20×10^3 , 30×10^3 , 40×10^3 , 50×10^3 , 60×10^3 , 70×10^3 , 80×10^3 , 90×10^3 , 1×10^4 , 1×10^5 , 1×10^6 , 1×10^7 , 1×10^8 , 1×10^9 or 1×10^{10} base pairs in length.

5. The method of claim 1, wherein said first regulatory element is a promoter.

6. The method of claim 5, wherein said double-stranded polynucleotide comprises a second regulatory element, said second regulatory element being a polyadenylation signal.

7. The method of claim 1, wherein said double-stranded polynucleotide comprises a plurality of coding regions and a plurality of regulatory elements.

8. The method of claim 7, wherein said coding regions encode products that comprise a biochemical pathway.

9. The method of claim 8, wherein said biochemical pathway is glycolysis.

10. The method of claim 9, wherein said coding regions encode enzymes selected from the group consisting of hexokinase, phosphohexose isomerase, phosphofructokinase-1, aldolase, triose-phosphate isomerase, glyceraldehyde-3-phosphate dehydrogenase, phosphoglycerate kinase, phosphoglycerate mutase, enolase and pyruvate kinase.

11. The method of claim 8, wherein said biochemical pathway is lipid synthesis.

12. The method claim 7, wherein said biochemical pathway is cofactor synthesis.

13. The method of claim 13, wherein said pathway involves lipoic acid.

14. The method of claim 13, wherein said biochemical pathway is riboflavin synthesis.

15. The method of claim 7, wherein said biochemical pathway is nucleotide synthesis.

16. The method of claim 15, wherein said nucleotide is a purine.

17. The method of claim 15, wherein said nucleotide is a pyrimidine.

18. The method of claim 7, wherein said coding regions encode enzymes involved in a cellular process selected from the group consisting of cell division, chaperone, detoxification, peptide secretion, energy metabolism, regulatory function, DNA replication, transcription, RNA processing and tRNA modification.

19. The method of claim 18, wherein said energy metabolism is oxidative phosphorylation.

20. The method of claim 1, wherein said double-stranded polynucleotide is a DNA.

21. The method of claim 1, wherein said double-stranded polynucleotide is an RNA.

22. The method of claim 1, wherein said double-stranded polynucleotide is an expression construct.

23. The method of claim 22, wherein said expression construct is a bacterial expression construct.

24. The method of claim 22, wherein said expression construct is a mammalian expression construct.

25. The method of claim 17, wherein said expression construct is a viral expression construct.

26. The method of claim 1, wherein said double-stranded polynucleotide comprises a genome selected from the group consisting of bacterial genome, yeast genome, viral genome, mammalian genome, amphibian genome and avian genome.

27. The method of claim 1, wherein said overlap between the oligonucleotides of said first and said second set of oligonucleotides is between about 5 base pairs and about 75 base pairs.

28. The method of claim 1, wherein said overlap is about 10 base pairs, about 15 base pairs, about 20 base pairs, about 25 base pairs, about 30 base pairs, about 35 base pairs, about 40 base pairs, about 45 base pairs, about 50 base pairs, about 55 base pairs, about 60 base pairs, about 65 base pairs, or about 70 base pairs.

29. The method of claim 5, wherein said promoter is selected from the group consisting of CMV IE, SV40 IE, RSV, β -actin, tetracycline regulatable and ecdysone regulatable.

30. The method of claim 26, wherein said genome is a viral genome.

31. The method of claim 30, wherein said viral genome is selected from the group consisting of retrovirus, adenovirus, vaccinia virus, herpesvirus and adeno-associated virus.

32. The method of claim 1, wherein said double-stranded polynucleotide is a chromosome.

33. A method of producing a viral particle comprising the steps of:

- (a) providing a host cell;
- (b) transforming said host cell with an artificial viral genome prepared by:
 - (i) generating a first set of oligonucleotides corresponding to the entire plus strand of said viral genome;
 - (ii) generating a second set of oligonucleotides corresponding to the entire minus strand of said viral genome; and
 - (iii) annealing said first and said second set of oligonucleotides;

wherein each of said oligonucleotides of said second set of oligonucleotides overlaps with and hybridizes to two complementary oligonucleotides of said first set of oligonucleotides, except that two oligonucleotides at a 5' or 3' end of said viral genome will hybridize with only one complementary oligonucleotide; and

- (c) culturing said transformed host cell under conditions such that said viral particle is expressed.

34. The method of claim 33, wherein said viral genome is selected from the group consisting of retrovirus, adenovirus, vaccinia virus, herpesvirus and adeno-associated virus.

35. A method of producing an artificial genome, wherein said chromosome comprises all coding regions and regulatory elements found in a corresponding natural chromosome, comprising the steps of:

- (a) generating a first set of oligonucleotides corresponding to the entire plus strand of said chromosome;
- (b) generating a second set of oligonucleotides corresponding to the entire minus strand of said chromosome; and
- (c) annealing said first and said second set of oligonucleotides;

wherein each of said oligonucleotides of said second set of oligonucleotides overlaps with and hybridizes to two complementary oligonucleotides of said first set of oligonucleotides, except that two oligonucleotides at a 5' or 3' end of said chromosome will hybridize with only one complementary oligonucleotide.

36. The method of claim 35, wherein said corresponding natural chromosome is a human mitochondrial genome.

37. The method of claim 35, wherein said corresponding natural chromosome is a chloroplast genome.

38. A method of producing an artificial genetic system, wherein said system comprises all coding regions and regulatory elements found in a corresponding natural biochemical pathway, comprising the steps of:

- (a) generating a first set of oligonucleotides corresponding to the entire plus strand of said chromosome;
- (b) generating a second set of oligonucleotides corresponding to the entire minus strand of said chromosome; and
- (c) annealing said first and said second set of oligonucleotides;

wherein each of said oligonucleotides of said second set of oligonucleotides overlaps with and hybridizes to two complementary oligonucleotides of said first set of oligonucleotides, except that two oligonucleotides at a 5' or 3' end of said chromosome will hybridize with only one complementary oligonucleotide

wherein expression of said biochemical pathway coding regions results in the expression of a group of enzymes that serially metabolize a compound.

39. The method of claim 38, wherein said biochemical pathway comprises the activities required for glycolysis.

40. The method of claim 38, wherein said biochemical pathway comprises the enzymes required for electron transport.

41. The method of claim 38, wherein said biochemical pathway comprises the enzyme activities required for photosynthesis.

* * * * *