

US 20060084111A1

(19) **United States**

(12) **Patent Application Publication**

Ruan et al.

(10) **Pub. No.: US 2006/0084111 A1**

(43) **Pub. Date: Apr. 20, 2006**

(54) **METHOD FOR GENE IDENTIFICATION
SIGNATURE (GIS) ANALYSIS**

(75) Inventors: **Yijun Ruan**, Singapore (SG); **Patrick
Ng**, Singapore (SG); **Chialin Wei**,
Singapore (SG)

Correspondence Address:
**ROTHWELL, FIGG, ERNST & MANBECK,
P.C.
1425 K STREET, N.W.
SUITE 800
WASHINGTON, DC 20005 (US)**

(73) Assignee: **Agency For Science, Technology and
Research**, Singapore (ES)

(21) Appl. No.: **11/304,538**
(22) Filed: **Dec. 16, 2005**

Related U.S. Application Data

(62) Division of application No. 10/664,234, filed on Sep.
17, 2003.

Publication Classification

(51) **Int. Cl.**
C40B 40/08 (2006.01)
C12Q 1/68 (2006.01)
C07H 21/02 (2006.01)
(52) **U.S. Cl.** **435/6; 536/23.1**

(57) **ABSTRACT**

An isolated oligonucleotide comprising at least one ditag,
wherein the ditag comprises two joined first and second
sequence tags, wherein the first tag comprises the 5'-termi-
nus sequence and the second tag comprises the 3'-terminus
sequence of a nucleic acid molecule or a fragment thereof.
The ditag analysis is useful for gene discovery and genome
mapping.

Figure 1




Figure 2

GIS analysis (PCR approach)




Figure 3




Figure 4




Figure 5




Figure 6




Figure 7




Figure 8

pZErO-1




Figure 9




FIGURE 10

pGISI sequence

NotI

XhoI ----- MmeI

EcoRI ----- BamHI

1 GGGCGAATTC TCGAGCGGCC GCGGATCCGA CGAGAGCGCC TCGGTACGSC TCGCCGCGGT GGCTGGCGCT ACTTCGGAGG AGCCCGACGC GCGCGGGTCG
CCCGCTTAAG AGCTCGCCGG CGCCTAGGCT GCTCTCGGG ACGCATGCCG ACGGGCGCCA CCGACCGCGA TGAAGCCTCC TCGGGCTGCG CCGCGCCAGC

101 TTTTTATACA TTCCCGCGCG GAGGCAACGG AAGGGCGGGG CGCCTCGTGA TTAGGCGCGG GAGGTCACAG GCTCTGTTGT CATGAAGGTG AAAATTAAAT
AAAAATATGT AAGGGCGCGC CTCCGTTGCC TTCCCGCCCC GCGGAGCACT AATCCGGCGC CTCCAGTGTC CGAGACAACA GTACTTCCAC TTTTAATTTA
MmeI

201 GTTGGAAATGG TGTGGCCACT TGGCTCTGGG TAGCCAATGA TGAGAACTGC GGCATCTGCA GGATGGCGTT TAATGGCTGC TGTCCAGACT GTAAGGTGCC
CAACCTTACC ACACCGGTGA ACCGAGACCC ATCGGTTACT ACTCTTGACG CCGTAGACGT CCTACCGCAA ATTACCGACG ACAGGTCTGA CATTCCACGG

301 TGGTGATGAC TCGCCCTCG TGTGGGACA GTGCTCCCAC TGCTTCCACA TGCATGTCAT CCTCAAGTGG CTGAATGCGC AGCAGGTGCA GCAGCACTGC
ACCACTACTG ACGGGGAGC ACACCCCTGT CACGAGGGTG ACGAAGGTGT ACGTGACGTA GGAGTTCACC GACTTACGCG TCGTCCACGT CGTCGTGACG

401 CCCATGTGTC GCCAGGAGTG GAAGTTCAA GAGTGAAGCC CGTGCCGTGC CACTTCCCTC TCCTGTGCTG TGCCAGGCTC AGCCCTTCC CTCCCTCCCC
GGGTACACAG CGGTCTCAC CTTCAGTTT CTCACTTCGG GCACGGCAGC GTGAAGGGAG AGGACACGAC ACGGTCCGAG TCGGGGAAGG GAGGGAGGGG

501 TCCCCCAGAT ACAGCACCCC AAGTCCCCTC CACACAGCAC AGTGGTGCCC AGAGATCTCG GTCTGTGCCG GGGACAAGGA TGCTTTCTGT TTGGCTGGGA
AGGGGTCTA TGTCTGCGG TTCAGGGGAG GTGTGTCTGT TCACCACGGG TCTCTAGAGC CAGACACGGC CCCTGTTCTT ACGAAAGACA AACCGACCTT
MmeI

601 CAAGGTTGAA AGGAGCTTTG CTGACTGTTT TGTTTTCCCA TCACATTGAC ACTTTATTCA ATAAGTAAAA CTCATTACAG TTCCAAGTCG GATCCTGGGT
GTTCCAACCT TCCTCGAAAC GACTGACAAA ACAAAGGGT AGTGTAACGT TGAAATAAGT TATTCATTTT GAGTAATGTC AAGGTTACGC CTAGGACCCA
SalI

701 CGACCTGCAG GCATGCAAGC TTGAGTATTC TATAGTGTC CTTAAATAGC TTGGCGTAAT CATGGTCATA GCTGTTTCCT GTGTGAAATT GTTATCCGCT
GCTGACGTC CGTACGTTCC AACTCATAAG ATATCACAGT GGATTTATCG AACCGCATTA GTACCAGTAT CGACAAAGGA CACACTTTAA CAATAGGCGA

801 CACAATTCCA CACAACATAC GAGCCGGAAG CATAAAGTGT AAAGCCTGGG GTGCCATAAT AGTGAGCTAA CTCACATTAA TTGCGTTGCG CTCACTGCC
GTGTTAAGGT GTGTTGTATG CTCGGCCTTC GTATTTCACA TTTCGGACCC CACGGATTAC TCACCTGATT GAGTGTAATT AACGCAACGC GAGTGACGGG

901 GCTTTCAGT CGGGAACCT GTCGTGCCAG CTGCATTAAT GAATCGGCCA ACGCGCGGGG AGAGGCGGTT TCGGTATTGG GCGCTCTTCC GCTTCTTCGC
CGAAAGGTCA GCCCTTTGGA CAGCACGGTC GACGTAATTA CTTAGCCGGT TGCGCGCCCC TCTCCGCCAA ACGCATAACC CGCGAGAAGG CGAAGGAGCG

1001 TCACTGACTC GCTGCGCTCG GTCGTTCCGG TCGCGCGAGC GGTATCAGCT CACTCAAAGG CGGTAATACG GTTATCCACA GAATCAGGGG ATAACGCAAG
AGTGACTGAG CGACGCGAGC CAGCAAGCCG ACGCCGCTCG CCATAGTCGA GTGAGTTTCC GCCATTATGC CAATAGGTGT CTTAGTCCCC TATTGCGTCC

1101 AAAGAACATG TGAGCAAAAG GCCAGCAAAA GGCCAGGAAC CGTAAAAAGG CCGCGTTGCT GGCGTTTTTC GATAGGCTCC GCCCCCTGA CGAGCATCAC
TTTCTGTAC ACTCGTTTTC CGGTGTTTTT CCGGTCTTTC GCATTTTTC GCGCAACGA CCGCAAAAAG CTATCCGAGG CGGGGGGACT GCTCGTAGTG

1201 AAAAATCGAC GCTCAAGTCA GAGGTGGCGA AACCAGACAG GACTATAAAG ATACCAGGCG TTTCCCTCTG GAAGCTCCCT CGTGCGCTCT CCTGTACCGA
TTTTTAGCTG CGAGTTCAGT CTCCACCGCT TTGGGCTGTC CTGATATTTT TATGGTCCGC AAAGGGGGAC CTTGAGGGGA GCACGCGAGA GGACATGGCT

1301 CCCTGCCGCT TACCGGATAC CTGTCCGCTT TTCTCCCTTC GGAAGCGGTG GCGCTTTCTC ATAGCTCACG CTGTAGGTAT CTCAGTTCGG TGTAGGTGCT
GGGACGGCGA ATGGCCTATG GACAGGCGGA AAGAGGGAAG CCCTTCGCAC CCGCAAGAG TATCGAGTGC GACATCCATA GAGTCAAGCC ACATCCAGCA

1401 TCGCTCCAAG CTGGGCTGTG TGCACGAACC CCCCCTTCAG CCCGACCGCT GCGCCTTATC CGGTAACCTAT CGTCTTGAGA CCAACCCGGT AAGACACGAC
AGCGAGGTTT GACCCGACAC ACGTGCTTGG GGGGCAAGTC GGCTGCGGCA CGCGGAATAG GCCATTGATA GCAGAACTCT GGTGGGGCCA TTCTGTGCTG

1501 TTATCGCCAC TGGCAGCAGC CACTGGTAAC AGGATTAGCA GAGCGAGGTA TGTAGGCGGT GCTACAGAGT TCTTGAAGTG GTGGCCTAAC TACGGCTACA
AATAGCGGTG ACCGTCGTCG GTGACCAATG TCCTAATCCT CTCGCTCCAT ACATCCGCCA CGATGCTCA AGAAGCTTAC CACCGGATTG ATGCCGATGT

1601 CTAGAAGGAC AGTATTTGGT ATCTGCGCTC TGCTGAAGCC AGTTACCTTC GGAAAAAGAG TTGGTAGCTC TTGATCCGGC AAACAACCA CCGCTGGTAG
GATCTTCTCG TCATAAACCA TAGACGCGAG ACGACTTCGG TCAATGGAAG CCTTTTCTC AACCATCGAG AACTAGGCCG TTTGTTTGGT GGCGACCATC

1701 CGGTGGTTTT TTTGTTTGCA AGCAGCAGAT TACGCGCAGA AAAAAGGAT CTCAAGAAGA TCCTTTGATC TTTTCTACGG GGTCTGACGC TCAGTGGAAC
GCCACCAAAA AAACAACGT TCGTCGTCTA ATGCGCGTCT TTTTTCCTA GAGTCTTCT AGGAACTAG AAAAGATGCC CCAGACTGCG AGTCACCTTG

1801 GAAAACCTAC GTTAAGGGAT TTTGGTCATG AGATTATCAA AAAGGATCTT CACCTAGATC CTTTTAAAT AAAAATGAAG TTTTAAATCA ATCTAAAGTA
CTTTGAGTG CAATTCCTTA AAACAGTAC TCTAATAGTT TTTCCTAGAA GTGGATCTAG GAAAATTTAA TTTTACTTC AAAATTTAGT TAGATTTTAT

1901 TATATGAGTA AACTTGGTCT GACAGTTACC AATGCTTAAT CAGTGAGGCA CCTATCTCAG CGATCTGTCT ATTTGCTTCA TCCATAGTTG CCTGACTCCC
ATATACTCAT TTGAACCAGA CTGTCAATGG TTACGAATTA GTCACTCCGT GGATAGAGTC GCTAGACAGA TAAAGCAAGT AGGTATCAAC GGACTGAGGG

2001 CGTCGTGTAG ATAACACGA TACGGGAGGG CTTACCATCT GGGCCAGTG CTGCAATGAT ACCGCGAGAC CCACGCTCAC CGGCTCCAGA TTTATCAGCA
GCAGCACATC TATTGATGCT ATGCCCTCCC GAATGGTAGA CCGGGGTCAC GACGTTACTA TGGCGCTCTG GGTGCGAGTG GCCGAGGTCT AAATAGTCGT

2101 ATAAACCAGC CAGCCGGAAG GGCCGAGCGC AGAAGTGGTC CTGCAACTTT ATCCGCTTCC ATCCAGTCTA TTAATTGTTG CCGGGAAGCT AGAGTAAGTA
TATTTGGTCG GTCGGCCTTC CCGGCTCGCG TCTTACCAG GACGTTGAAA TAGGCGGAGG TAGGTGAGT AATTAACAAC GGCCCTTCGA TCTCATTCAT

2201 GTTCGCCAGT TAATAGTTTG CGCAACGTTG TTGGCATTGC TACAGGCATC GTGGTGTCAC GCTCGTCGTT TGGTATGGCT TCATTGAGCT CCGGTTCCCA
CAAGCGGTCA ATTATCAAC GCGTTGCAAC AACCGTAACG ATGTCCGTAG CACCACAGTG CGAGCAGCAA ACCATACCGA AGTAAGTCGA GGCCAAGGGT

2301 ACGATCAAGG CGAGTTACAT GATCCCCCAT GTGTGCAAAA AAAGCGGTTA GCTCCTTCGG TCCTCCGATC GTTGTGAGAA GTAAGTTGGC CGCAGTGTTA
TGCTAGTTCC GCTCAATGTA CTAGGGGGTA CAACACGTTT TTTCGCCAAT CGAGGAAGCC AGGAGGCTAG CAACAGTCTT CATTCAACCG GCGTCACAA

2401 TCACTCATGG TTATGGCAGC ACTGCATAAT TCTCTTACTG TCATGCCATC CGTAAGATGC TTTTCTGTGA CTGGTGAGTA CTCAACCAAG TCATTCTGAG
AGTGAGTACC AATACCGTCG TGACGTATTA AGAGAATGAC AGTACGGTAG GCATTCTACG AAAAGACACT GACCACTCAT GAGTTGGTTC AGTAAGACTC

2501 AATAGTGTAT GCGGCGACCG AGTTGCTCTT GCCCGGCGTC AATACGGGAT AATACCGCGC CACATAGCAG AACTTTAAAA GTGCTCATCA TTGGAAAACG
TTATACATA CGCCGCTGGC TCAACGAGAA CCGGCGCGAG TTATGCCCTA TTATGGCGCG GTGTATCGTC TTGAAATTTT TCGAGTAGT AACCTTTTGC

2601 TTCTTCGGGG CGAAAACCT CAAGGATCTT ACCGCTGTTG AGATCCAGTT CGATGTAACC CACTCGTGCA CCCAACTGAT CTTGAGCATC TTTTACTTTT
AAGAAGCCCC GCTTTTGAGA GTTCCTAGAA TGGCGACAAC TCTAGGTCAA GCTACATTGG GTGAGCACGT GGGTTGACTA GAAGTCGTAG AAAATGAAAG

FIGURE 10 (CONTINUED)

2701	ACCAGCGTTT CTGGGTGAGC AAAAACAGGA AGGCAAAATG CCGCAAAAAA GGGAATAAGG GCGACACGGA AATGTTGAAT ACTCATACTC TTCCTTTTTC	TGGTCGCAAA GACCCACTCG TTTTGTCTCT TCCGTTTAC GCGGTTTTTT CCCTTATTCC CGCTGTGCCT TTACAACCTA TGAGTATGAG AAGGAAAAAG
2801	AATATTATTG AAGCATTTAT CAGGGTTATT GTCTCATGAG CCGATACATA TTTGAATGTA TTTAGAAAAA TAAACAAATA GGGGTTCCGC GCACATTTCC	TTATAATAAC TTCGTAAATA GTCCCAATAA CAGAGTACTC GCCTATGTAT AAACCTACAT AAATCTTTTT ATTTGTTTAT CCCCAGGCG CGTGTAAGG
2901	CCGAAAAGTG CCACCTGACG TCTAAGAAAC CATTATTATC ATGACATTAA CCTATAAAAA TAGGCGTATC ACGAGGCCCT TTCGTCTCGC GCGTTTCGGT	GGCTTTTCAC GGTGGACTGC AGATTCTTTG GTAATAATAG TACTGTAATT GGATATTTTT ATCCGCATAG TGCTCCGGGA AAGCAGAGCG CGCAAAGCCA
3001	GATGACGGTG AAAACCTCTG ACACATGCAG CTCCCGGAGA CGGTCACAGC TTGTCTGTAA GCGGATGCCG GGAGCAGACA AGCCCGTCAG GCGCGCTCAG	CTACTGCCAC TTTTGAGAGC TGTGTACGTC GAGGGCCTCT GCCAGTGTCT AACAGACATT CGCCTACGGC CCTCGTCTGT TCGGGCAGTC CCGCGCAGTC
3101	CGGGTGTGG CGGGTGTGCG GGCTGGCTTA ACTATGCGGC ATCAGAGCAG ATTGTACTGA GAGTGCACCA TATGCGGTGT GAAATACCGC ACAGATGCGT	GCCCACAACC GCCCACAGCC CCGACCGAAT TGATACGCCG TAGTCTCGTC TAACATGACT CTCACGTGGT ATACGCCACA CTTTATGGCG TGTCTACGCA
3201	AAGGAGAAAA TACCGCATCA GCGCCATTC GCCATTCAGG CTGCGCAACT GTTGGGAAGG GCGATCGGTG CGGGCCTCTT CGCTATTACG CCAGCTGGCG	TTCTCTTTT ATGGCGTAGT CCGCGGTAAG CGGTAAGTCC GACGCGTTGA CAACCTTCC CGCTAGCCAC GCCCGAGAA GCGATAATGC GGTGACCGC
3301	AAAGGGGGAT GTGCTGCAAG GCGATTAAGT TGGGTAACGC CAGGGTTTTT CCAGTCACGA CGTTGTAAAA CGACGGCCAG TGAATTGTAA TACGACTCAC	TTTCCCCCTA CACGACGTTT CGCTAATTCA ACCCATGCG GTCCCAAAG GGTCAGTGCT GCAACATTTT GCTGCCGGTC ACTTAACATT ATGCTGAGTG
3401	TATA	ATAT

METHOD FOR GENE IDENTIFICATION SIGNATURE (GIS) ANALYSIS

[0001] This application is a divisional of U.S. Ser. No. 10/664,234, filed Sep. 17, 2003.

FIELD OF THE INVENTION

[0002] The present invention relates generally to the field of gene and transcript expression and specifically to a method for the serial analysis of a large number of transcripts by identification of a gene signature (GIS) corresponding to defined regions within a transcript.

BACKGROUND OF THE INVENTION

[0003] One of the most important goals of the human genome project is to provide complete lists of genes for the genomes of human and model organisms. Complete genome annotation of genes relies on comprehensive transcriptome analysis by experimental and computational approaches. Ab initio predictions of genes must be validated by experimental data. An ideal solution is to clone all full-length transcripts and completely sequence them. This approach has gained recognition recently (Strausberg, R. L., et al., 1999, *Science*, 286: 455-457) and progress has been made (Jongeneel C. V., et al., 2003, *Proc Natl Acad Sci U S A*, 100, 4702-4705). However, due to the complexity and immense volume of transcripts expressed in the various developmental stages of an organism's life cycle, complete sequencing analysis of all different transcriptomes still remains unrealistic.

[0004] To get around such a dilemma, a cDNA tagging strategy that obtains partial sequences that represent full transcripts has been developed and widely applied in determining genes and characterizing transcriptomes in the past decade.

[0005] In the expressed sequence tag (EST) approach, cDNA clones are sequenced from 5' and/or 3' ends (Adams, M., et al., 1991, *Science*, 252, 1651-1656). Each EST sequence read would generate on average a 500 bp tag per transcript. The number of same or overlapping ESTs would manifest the relative level of gene expression activity. Though EST is effective in identifying genes, it is prohibitively expensive to tag every transcript in a transcriptome. In practice, sequencing usually ceases after 10,000 or less ESTs are obtained from a cDNA library where millions of transcripts might be cloned.

[0006] To increase the efficiency in sequencing and counting large numbers of transcripts, Serial Analysis of Gene Expression (SAGE) (Velculescu, V. E., et al., 1995, *Science*, 270, 484-487; Saha S., et al., 2002, *Nature Biotechnology*, 20, 508-12; U.S. Pat. No. 6,498,013; U.S. Pat. No. 6,383,743) and the recent Massively Parallel Signature Sequencing (MPSS) technique (Mao C., et al., 2000, *Proc Natl Acad Sci USA*, 97, 1665-1670; Brenner S., et al., 2000, *Nature Biotechnology*, 18, 630-634) were developed based on the fact that a short signature sequence (14-20 bp) of a transcript can be sufficiently specific to represent that gene.

[0007] Experimentally, short tags can be extracted from cDNA one tag per transcript. Such short tags can be efficiently sequenced either by a concatenation tactic (as for SAGE) or by a hybridization-based methodology for MPSS. For example, in SAGE, multiple tags are concatenated into

long DNA fragments and cloned for sequencing. Each SAGE sequence readout can usually reveal 20-30 SAGE tags. A modest SAGE sequencing effort of less than 10,000 reads will have significant coverage of a transcriptome. Transcript abundance is measured by simply counting the numerical frequency of the SAGE tags.

[0008] With the availability of many assembled genome sequences in public databases, the use of a short tag strategy for transcriptome characterization is becoming popular (Jongeneel et al., 2003, *Proc. Natl. Acad. Sci. USA* 100: 4702-4705). In theory, short DNA tags of about 20 bp can be specifically mapped to a single location within a complex mammalian genome and uniquely represent a transcript in the content of whole transcriptome. However, in reality, there still exist a large number of "ambiguous" SAGE tags (14-21 bp) and MPSS tags (17 bp) that have multiple locations in a genome, and may be shared by many genes. Limited by the availability of type II restriction enzymes that can cut longer than 21 bp, the SAGE method currently can not generate any longer tags to improve specificity.

[0009] Further, SAGE and MPSS methods only produce a single signature per transcript in the middle of the gene. In view of the "internal" nature of the tag in a transcript, these methods provide only limited tag information.

[0010] Therefore, despite their usefulness in sequencing efficiency, the utility of methods such as SAGE or MPSS is severely undermined by their lack of specificity and consequent inconclusiveness.

[0011] There is a need in the art for more efficient methods which retain the sequencing efficiency and at the same time improve the use of the tagging strategy for transcriptome characterization and facilitate the annotation of genomes.

SUMMARY OF THE INVENTION

[0012] The present invention solves the problems mentioned above by providing two tags (a ditag) per nucleic acid molecule, therefore increasing the specificity of the tags to represent a nucleic acid molecule (for example a gene). The two tags are extracted from the 5' and 3' ends of the same nucleic acid molecule, and therefore ditags are more informative to reflect the structure of the nucleic acid molecules. Critically, the invention provides a method to link the 5' and 3' tags of the same nucleic acid molecule into a single ditag unit. Therefore, the pairs of 5' and 3' tags that represent the nucleic acid molecule can be easily recognized by simple sequencing analysis. The invention can be used for the identification of new genes, for the measure of transcript abundance in transcriptomes, for the annotation of genome sequences and at the same time enhancing sequencing efficiency.

[0013] In particular, the invention provides an isolated oligonucleotide comprising at least one ditag, wherein the ditag comprises two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a nucleic acid molecule.

[0014] The oligonucleotide of the invention, further comprises at least two adapters flanking the ditag, wherein each adapter comprises at least one restriction site. In particular, each adapter comprises at least: a first restriction site proximal to the tag which is an asymmetric recognition site (for

example, a homing endonuclease recognition site, or a type II recognition site) and at least a second restriction site. The second or further restriction site may be any restriction site known in the art may be used. For example, BamHI. Also, any asymmetric restriction site different from the first restriction site may be used. The recognition site for this enzyme however must be absent from the vector backbone after insertion of the ditag.

[0015] The nucleic acid molecule may be the full-length sequence of a gene or a fragment thereof. For example, RNA, mRNA, genomic DNA, full-length cDNA or cDNA.

[0016] The ditag may vary in nucleotide number. According to one embodiment, it is obtained by splicing the 5' terminus and the 3' terminus of a nucleic acid molecule in presence of at least one restriction enzyme and the size of the sequence tags is determined by the restriction enzyme used. Accordingly, the number of nucleotides of the ditag can vary according to the restriction enzyme used.

[0017] When MmeI is used, this enzyme recognizes a sequence inside each of the two adapters that flank the nucleic acid molecule which one intends to reduce, but cuts inside the nucleic acid molecule forming a tag comprising 19-21 nucleotides. Two such tags may be additionally processed by blunting and ligation to form a ditag comprising 34-38 nucleotides. The ditag is hence obtained by splicing together the 5' terminus and the 3' terminus of the same nucleic acid molecule.

[0018] The ditag of the invention can be of any size, preferably 12-60 bp.

[0019] The oligonucleotide may comprise a concatemer of ditags, for example 1 to 1000 ditags.

[0020] The invention also provides a vector comprising the oligonucleotide of the invention. In particular, the vector comprises at least a nucleic acid molecule and at least two adapters flanking the nucleic acid molecule, wherein each adapter comprises at least: a first restriction site which is a asymmetric restriction site (asymmetric restriction site is, for example, a homing endonuclease recognition site, or a type II recognition site) and at least a second restriction site (for example Bam HI), and the backbone of the vector does not comprise the asymmetric restriction site and the second or further restriction site. A preferable, asymmetric restriction site is the type II restriction site MmeI.

[0021] The invention also provides a vector having the sequence indicated in SEQ ID NO:18.

[0022] The invention further provides a cDNA library, wherein every cDNA clone comprises the at least one oligonucleotide of the invention.

[0023] According to another aspect, the invention also provides a method for preparing at least one oligonucleotide comprising at least one ditag comprising:

[0024] producing at least one nucleic acid molecule;

[0025] isolating the 5' terminus and the 3' terminus of the nucleic acid molecule or fragment thereof, and

[0026] linking the 5' terminus and 3' terminus to create the at least one ditag.

[0027] In particular, it is provided a method for preparing at least one oligonucleotide comprising at least one ditag comprising:

[0028] producing at least one nucleic acid molecule flanked by two adapters;

[0029] isolating the 5' terminus and the 3' terminus of the nucleic acid molecule; and

[0030] linking the 5' terminus and 3' terminus to create the at least one oligonucleotide comprising at least one ditag flanked by the two adapters.

[0031] The nucleic acid molecule desired to be reduced in form of a ditag may be a full nucleic acid molecule or a portion inside the nucleic acid molecule.

[0032] The nucleic acid molecule may correspond to the full-length of a gene or fragment thereof.

[0033] The method may further comprise the step of determining the nucleotide sequence of the at least one ditag to detect gene expression.

[0034] According to a further aspect, the method of the invention may further comprise the steps of:

[0035] determining the sequence of the at least one ditag; and

[0036] comparing the ditag nucleotide sequence to a database comprising genomic sequences whereby matching 5' and 3' termini sequences are identified.

[0037] According to a particular embodiment, the invention provides a method comprising:

[0038] producing at least one nucleic acid molecule, preferably a full-length cDNA, flanked by two adapters, wherein each adapter comprises at least one restriction site; splicing the 5' terminus and the 3' terminus of the nucleic acid molecule to produce at least one ditag by adding at least one restriction enzyme recognizing the recognition sites.

[0039] Preferably, each adapter comprises at least: a first restriction site which is an asymmetric restriction site and a second restriction site.

[0040] As restriction enzyme, any useful enzyme can be used. For example, a restriction enzyme recognizing two asymmetric recognition sites.

[0041] Asymmetric recognition site can be: i) homing endonuclease asymmetric recognition site sequences or ii) restriction endonuclease asymmetric cleavage sites sequences recognizable by type II restriction enzymes.

[0042] According to a particular embodiment, the splicing step is carried out by using MmeI (together with T4 DNA polymerase and T4 DNA ligase) and the ditag of 34-38 nucleotides, flanked by two adapters, is produced.

[0043] According to a further aspect, the ditag of any embodiment of the invention can be linked to other ditags to produce concatemers of ditag. For example, 1 to 1000 ditags.

[0044] According to another further aspect, it is provided a method for genome mapping, comprising:

[0045] preparing at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a nucleic acid

molecule, the nucleic acid molecule corresponding to the full-length of a gene or fragment thereof;

[0046] mapping each of the two tags of the at least one ditag on the genome; and

[0047] defining the structural region of the corresponding gene on the genome map.

[0048] According to a still another aspect, the invention provides a method of gene discovery comprising:

[0049] preparing at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a nucleic acid molecule, the nucleic acid molecule corresponding to the full-length of a gene or fragment thereof;

[0050] comparing the obtained at least one ditag with a genome map and/or a gene database; if the 5' and 3' termini tags of a ditag are matched to the genome sequence but not in known gene databases, then the detected ditags may represent new genes in the given genomes.

[0051] Such ditags can directly guide the process of recovering the full-length nucleic acid molecule corresponding to the newly identified genes.

[0052] It is also an aspect of the invention a method for recovering the full-length cDNA of new and/or other interesting genes comprising:

[0053] preparing, from a full-length cDNA library, at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a full-length cDNA library;

[0054] sequencing the obtained oligonucleotide ditag, preferably a large number of the obtained ditags;

[0055] determining the ditag of interest (for example, based on biological aspects); and

[0056] recovering the full-length cDNA corresponding to the ditag of interest from the parental full-length cDNA library.

[0057] Further, the invention also provides a method for quantifying the transcriptional activity of a gene comprising:

[0058] preparing, from a full-length cDNA library, at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a full-length cDNA;

[0059] sequencing the obtained oligonucleotide ditag, preferably a large number of the obtained ditags;

[0060] determining the frequency of the sequenced ditag which corresponds to the transcriptional activity of the gene.

BRIEF DESCRIPTION OF THE FIGURES

[0061] FIG. 1 shows the GIS analysis experimental workflow (bacterial transformation approach). In the figure, the

letters N, B, M, S either in capital or small letters denotes the recognition sites for the restriction enzymes Not I, Bam HI, Mme I and Sal I, respectively. The text "Me" represents methylation of the newly-synthesized first-strand cDNA.

[0062] FIG. 2 shows the GIS analysis experimental workflow (PCR-based approach). In the figure, the letters N, B, M, S either in capital or small letters denotes the recognition sites for the restriction enzymes Not I, Bam HI, Mme I and Sal I, respectively. The text "Me" represents methylation of the newly-synthesized first-strand cDNA.

[0063] FIG. 3 shows the GIS application of mapping transcriptome to genome.

[0064] FIG. 4 is an electrophoresis gel showing MmeI digestion of a mix of original full-length cDNA clones. Lane 1: original supercoiled plasmid preparation. Lane 2: 1 kb DNA ladder. Lane 3: MmeI digestion products. The arrow-head shows the position of all the linearized tagged-plasmids.

[0065] FIG. 5 is an electrophoresis gel related to the preparation of GIS ditags. The plasmid DNA of GIS ditag library is digested with BamHI. The 50 bp ditag fragments are separated and purified from the vector using a 10% polyacrylamide gel. Lane 1: DNA size markers. Lane 2-8: formation of 50 bp GIS ditags.

[0066] FIG. 6 is an electrophoresis gel related to the preparation of GIS ditags by PCR. The ditag-containing PCR fragments generated from the GIS full-length cDNA library are digested by BamHI. The 50 bp ditag fragments are separated and purified from adaptor arms in 10% polyacrylamide gel. Lane 1: DNA size markers. Lane 2-15: large scale preparation of 50 bp GIS ditags.

[0067] FIG. 7 shows the pGIS1 vector construct.

[0068] FIG. 8 shows the commercial pZerO-1 vector construct (Invitrogen) The positions of the various sequencing/PCR primer binding sites (PMR003, PMR004, PMR011 and PMR012) are shown.

[0069] FIG. 9 shows a typical example of the QC (quality check) performed on multiple clones from the GIS library using PCR. Lane 1: pZerO-1 vector as negative control. M: 1kb+DNA ladder. Lanes 2-25: randomly-picked clones.

[0070] FIG. 10 shows the double strand nucleotide sequence of pGIS1. The region between the restriction sites Not I and Sal I is the stuffer fragment that is removed during cloning. It is highlighted in bold and italic type. The single strand nucleotide sequence is also reported as SEQ ID NO:18. The region representing the stuffer fragment is between nucleotide 15 to 704 (both nucleotides included).

DETAILED DESCRIPTION OF THE INVENTION

[0071] The present invention provides a Gene Identification Signatures (GIS) and a GIS analysis method: useful, for example, for the rapid analysis of numerous transcripts in order to identify the overall pattern of transcript expression (transcriptome), for the selection and/or construction of cDNA and full-length cDNAs, tag sequencing, gene discovery, genome mapping and annotation. In general, the GIS and GIS analysis method according to the invention greatly facilitates the collection of gene information by experimental approach.

[0072] For the purpose of the present application, GIS means a ditag (also indicated as GIS ditag) or an oligonucleotide comprising at least one ditag, wherein the ditag comprises the 5' terminus (or end region) and the 3' terminus (or end region) of a nucleic acid molecule, which it is desired to reduce, "shrink" or represent.

[0073] The ditag is shorter than the original nucleic acid molecule from which it originates or which it represents. Preferably, the ditag must be much shorter than the original nucleic acid molecule. As consequence of the "shrinking", the ditag essentially comprises the 5' end region (also indicated as 5' tag) and 3' end region (also indicated as 3' tag) of the original nucleic acid molecule. Hence, the portion of the original nucleic acid molecule which is between or inside the 5' tag and 3' tag is not included in the ditag. The ditag according to the invention retains the most informative features of the original nucleic acid molecule, viz. the start and the end signatures of the nucleic acid. It is thereby also more specific and accurate than SAGE or MPSS methods in characterizing transcriptomes and defining gene structure by mapping the GIS tags to genome sequences.

[0074] Accordingly, the invention provides an isolated oligonucleotide comprising at least one ditag, wherein the ditag comprises two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a nucleic acid molecule or fragment thereof.

[0075] The oligonucleotide of the invention may further comprise two adapters flanking the ditag, wherein each adapter comprises at least one restriction site (see **FIG. 1** and **FIG. 2**). In particular, each adapter comprises at least: a first restriction site which is an asymmetric restriction site and at least a second adjacent restriction site. Therefore, the number of restriction sites present in each adapter may be two or more. Examples of asymmetric restriction sites are homing endonuclease asymmetric recognition sites, and type II (or class II) recognition sites. A list of possible asymmetric restriction sites and corresponding restriction enzymes recognizing such asymmetric sites is reported below. Example of second and further restriction sites may be for example BamHI. This second restriction site is for the purpose of subsequent isolation of a pool of ditags that can then be ligated together to form concatemers.

[0076] The original nucleic acid molecule that one intends to reduce (to shrink) may be any natural, any modified or any synthetic nucleic acid molecule. It can also be of any size. The nucleic acid molecule can be a gene (the full-length of a gene) or a fragment thereof. The nucleic acid may be RNA, mRNA, genomic DNA, full-length cDNA, or cDNA or a fragment thereof.

[0077] The ditag can also be fully chemically synthesized by comprising the 5' end and 3' end of a nucleic acid molecule which the ditag intends to represent.

[0078] The molecule that one intends to reduce may also be a portion or fragment inside a nucleic acid molecule. Accordingly, it is possible to use restriction enzymes recognizing restrictions sites flanking the region which is intended to be reduced. The desired restriction sites may be placed into the appropriate position during the preparation of the nucleic acid molecule, for example a cDNA or full-length cDNA.

[0079] According to a particular aspect, the nucleic acid desired to be reduced is a full-length cDNA. Full-length cDNA can be prepared according to any method known in the art. See for example, the cap-trapper approach, for example Caminci et al., 1996, *Genomics*, Vol. 37, 327-336; U.S. Pat. No. 6,143,528; Edery et al., 1995, *Mol. Cell. Biol.*, Vol. 15, No. 6, 3363-3371.

[0080] Those of skill in the art will know other capture systems, for example, those based on biotin/streptavidin, digoxigenin/anti-digoxigenin for isolation of the full-length cDNAs can be used.

[0081] The ditag can be prepared according to any technique known in the art. For example, the original nucleic acid molecule may be cut through any chemical reaction and the obtained 5' and 3' termini ligated to create the ditag.

[0082] The nucleic acid molecule which is intended to be reduced, which is preferably prepared comprising two adapters flanking the molecule, may be inserted into a vector. In a particular realisation, each adapter comprises at least one restriction site, preferably comprises at least a first restriction site comprising an asymmetric restriction site and a second restriction site. Accordingly, in the vector used, it is important that the backbone of the vector does not comprise the restriction site or sites present in the adapters.

[0083] Accordingly, a library of nucleic acid molecule (for example, a library of full-length cDNAs) is prepared.

[0084] Preferably, the nucleic acid molecule is spliced into a ditag or oligonucleotide comprising a ditag by using restriction enzymes which recognize restriction sites flanking the nucleic acid molecule to be reduced. Accordingly, the recognition sites are placed upstream of the 5' terminus and downstream of the 3' terminus of the nucleic acid molecule or fragment thereof desired to be reduced (preferably into the adapters). Accordingly, the oligonucleotide obtained by splicing comprises two adapters flanking the ditag. Each adapter comprising at least one restriction site. Preferably, comprising at least one first restriction site which is an asymmetric site (for example a type II restriction site, like Mmel) and at least a second restriction site (any known restriction site may be used, for example BamHI).

[0085] The 5' tag and 3' tag forming the ditag may have the same or different size. Preferably, they have the same number of nucleotides.

[0086] The ditag can be of any size, but needs to be meaningful and advantageous over the size of the parental sequence from which it is derived. The preferred size of a tag or ditag is determined by genome complexity. For a bacterial genome a tag from about 8 bp to about 16 bp may be sufficient whereas for a complex genome like the human genome, a 16-20 bp tag (or in other words a 32-40 bp ditag) may be considered. In general, the size of the ditag is from about 12-60 bp.

[0087] For the purpose of the present application, the terms 5'-terminus, 5'-end and 5'-tag are equivalent to each other and can be used interchangeably. In the same way, the terms 3'-terminus, 3'-end and 3'-tag are equivalent to each other and can be used interchangeably. In an original nucleic acid molecule or portion inside a nucleic acid molecule that one intends to reduce or represent, each of the 5'-end and

3'-end represents a region or portion most closer to the extremity and most far from the middle region of the molecule.

[0088] According to one aspect, the 5'-tag and 3'-tag comprised in the ditag are the regions of the molecule cleaved by a restriction enzyme most closer to the 5'-end and 3'-end, respectively, of the nucleic acid molecule or portion thereof which is intended to be reduced or represented. Accordingly, the size of the ditag can be determined by the restriction enzyme or enzymes used. The invention, therefore, relates to an oligonucleotide comprising at least one ditag, wherein the ditag is obtained by splicing the 5' terminus and the 3' terminus of the nucleic acid molecule in the presence of at least one restriction enzyme, which recognizes the restriction sites flanking the nucleic acid molecule. Accordingly, the size of the sequence tags is determined by the restriction enzyme used.

[0089] When preparing the nucleic acid molecule, for example a full-length cDNA, desired restriction sites flanking the 5'-end and 3'-end of the region which is intended to be reduced or represented are inserted. An example of construction of a full-length cDNA by insertion of desired restriction sites flanking the 5'-end and 3'-end is shown in **FIG. 1** and **FIG. 2**. A full-length cDNA library is then prepared, following which a GIS ditag library is subsequently prepared.

[0090] As an example, a restriction enzyme recognizing an asymmetric restriction site can be used for the purpose of the preparation of the ditag according to the invention. In particular a type II enzyme, for example MmeI.

[0091] As an example, asymmetric sites can be introduced. Asymmetric site sequences useful for the purpose of the present invention are: i) two homing endonuclease asymmetric recognition site sequences or ii) restriction endonuclease asymmetric cleavage sites sequences recognizable by type II restriction enzymes.

[0092] Homing endonucleases are sold and described by New England Biolabs, Inc.; a description of the asymmetric site sequences is also available in the New England Biolabs Catalog. These homing endonuclease asymmetric recognition site sequences are from 18 to 39 bp. However, in the present invention the recognition site sequences are not limited to those sequences nor to these sizes. Preferably, the restriction homing endonucleases capable of cutting the asymmetric site sequences are selected from the group consisting of: I-CeuI, PI-SceI, PI-PspI and I-SceI. The list mentioned above however is not exhaustive. Other homing endonucleases known in the art and those which may be later discovered are included in the scope of the present invention.

[0093] Examples of type II restriction enzymes include: AarI, AceIII, AclI, BaeI, Bbr71, BbvI, BbvI, BccI, Bce83I, BceAI, BceFI, BcgI, BciVI, BfiI, BinI, BpII, BsaXI, BscAI, BseMII, BseRI, BsgI, BsmI, BsmAI, BsmFI, Bsp24I, BspCNI, BspMI, BsrI, BsrDI, BstF5I, BtgZI, BtsI, CjeI, CjePI, EciI, Eco31T, Eco57I, Eco57MI, Esp3I, Fall, FauI, FokI, GsuI, HaeIV, HgaI, Hin4I, HphI, HpyAV, Ksp632I, MboII, MlyI, MmeI, MnlI, PleI, PpiI, PstI, RleAI, SapI, SfaNI, SspD5I, Sth132I, StsI, TaqII, TspDTI, TspGWI, TspRI and Tth11II (the list in the web site of Rebase Enzymes®: <http://rebase.neb.com/cgi-bin/outsidelist>; see

also Szybalski, W., 1985, *Gene*, 40:169). The list mentioned above however is not exhaustive. Other type II enzymes known in the art and those which may be later discovered are included in the scope of the present invention.

[0094] Examples of recognition sites and cleavage sites of several class II restriction enzymes are (into parenthesis are the recognition site and the cleavage site): BbvI (GCAGC 8/12), HgaI (GACGC 5/10), BsmFI (GGGAC 10/14) SfaNI (GCATC 5/9), and Bsp I (ACCTGC 4/8).

[0095] The ditag of the invention can conveniently be ligated or joined in order to form concatemers of ditag. Accordingly, the invention relates to an oligonucleotide comprising 1 to 1000 ditags, in particular 1 to 200, more in particular 8 to 20 ditags. When ditags are concatemerized, a higher yield of information is achieved because the oligonucleotide, vector or clone comprises more ditags. Hence, the concatenation of ditags allows an efficient analysis of the nucleic acid molecules, like full-length cDNAs, in a serial manner by sequencing multiple ditags within a single vector or clone.

[0096] The oligonucleotide, ditag or concatemers of ditags can be inserted into a vector either before or after concatemerization.

[0097] According to one aspect, the oligonucleotide comprising the ditag is amplified. For example, by using PCR or any other known amplification methods. Accordingly, suitable PCR primers corresponding to specific regions inside the vector are used. Such regions flank the oligonucleotide comprising the ditag and adapters. PCR can be performed directly on the ligation (self-circularization) reaction to obtain short (for example 200 bp) PCR products (see the PCR approach in **FIG. 2**). These PCR products that contain the required GIS ditags will then be cut with an enzyme recognizing the at least second restriction site (inside the adapters) to generate the required short cohesive ditags. As restriction enzyme recognizing the second or further restriction site, BamHI can for example be used, and cohesive ditags of 50 bp are generated. The advantage of this amplification step is that of generating GIS ditags circumventing the need to produce a GIS ditag library amplification, which can be avoided by not transforming the self-circularized tagged plasmids. The amplified oligonucleotide can then subsequently be excised from the vector (in this example, by digestion with BamHI) and concatenated in long stretches of DNA or RNA for subsequent cloning and sequencing analysis (see **FIG. 1** and **FIG. 2**).

[0098] As a particular aspect, the invention discloses a cDNA library wherein the oligonucleotide comprises at least one ditag, and wherein the ditag comprises 34-38 nucleotides and is obtained by splicing nucleotides from the 5' terminus and nucleotides from the 3' terminus of a full-length cDNA or fragment thereof.

[0099] The ditag library according to the invention is representative of the library comprising the original nucleic acid molecules. For example, when the library comprising the nucleic acid molecules is a full-length cDNA library, the ditag library is representative of the full-length ditag library. Each ditag clone comprises sufficient information characterizing the specific full-length clone. More important, the ditag of the invention comprises the 5'-end and 3'-end of the original full-length cDNA. Hence, the ditag is representative of the structure of the full-length cDNA.

[0100] Accordingly, it is sufficient to sequence and analyze the ditag clones of the ditag library. In case a ditag of interest is found, the corresponding full-length cDNA can be selected and prepared from the full-length cDNA library, for example by PCR or directly from target RNA samples by RT-PCR.

[0101] The invention provides a method for the preparation of at least one oligonucleotide comprising at least one ditag comprising:

[0102] producing at least one nucleic acid molecule;

[0103] isolating the 5' terminus and the 3' terminus of the nucleic acid molecule or fragment thereof;

[0104] linking the 5' terminus and 3' terminus to create the at least one ditag.

[0105] In particular, the invention provides a method for preparing at least one oligonucleotide comprising at least one ditag comprising:

[0106] producing at least one nucleic acid molecule flanked by two adapters;

[0107] isolating the 5' terminus and the 3' terminus of the nucleic acid molecule; and

[0108] linking the 5' terminus and 3' terminus to create the at least one oligonucleotide comprising at least one ditag flanked by the two adapters.

[0109] The method further comprising including the oligonucleotide comprising the at least one ditag flanked by the adapters into a vector.

[0110] The nucleic acid molecule which is intended to shrink or represent may be RNA, mRNA, genomic DNA, full-length cDNA, or cDNA.

[0111] The nucleic acid molecule may be the full-length sequence of a gene or a fragment thereof.

[0112] The method of the invention may further comprise the step of determining the nucleotide sequence of the at least one ditag to detect gene expression. The method may further comprise the steps of: determining the sequence of the at least one ditag; and comparing the ditag nucleotide sequence to a database comprising genomic sequences whereby matching 5' and 3' termini sequences are identified.

[0113] More in particular, the invention relates to a method comprising:

[0114] producing at least one nucleic acid molecule, for example a full-length cDNA, flanked by two adapters, wherein each adapter comprises at least one restriction; splicing the 5' terminus and the 3' terminus of the nucleic acid molecule or fragment thereof to produce at least one ditag by adding at least one restriction enzyme recognizing the recognition sites.

[0115] Any recognition site known in the art may be used. Restriction enzyme recognizing at least one recognition site within the nucleic acid molecule and which can be used will be evident to those skilled in the art (see for example, Current Protocols in Molecular Biology, Vol. 2, 1995, Ed. Ausubel, et al., Greene Publish. Assoc. & Wiley Interscience, Unit 3.1.15; New England Biolabs Catalog, 1995).

[0116] For example, the two recognition sites may be asymmetric recognition sites:

[0117] The asymmetric recognition site are: i) homing endonuclease asymmetric recognition site sequences or ii) restriction endonuclease asymmetric cleavage sites sequences recognizable by type II restriction enzymes.

[0118] The type II restriction enzyme is selected from the group Consisting of AarI, AceIII, AclI, BaeI, Bbr7I, BbvI, BbvII, BccI, Bce83I, BceAI, BceII, BcgI, BciVI, BfiI, BinI, BpII, BsaXI, BscAI, BseMII, BseRI, BsgI, BsmI, BsmAI, BsmFI, Bsp24I, BspCNI, BspMI, BsrI, BsrDI, BstF5I, BtgZI, BtsI, CjeI, CjePI, EciI, Eco31I, Eco57I, Eco57MI, Esp3I, Fall, FauI, FokI, GsuI, HaeIV, HgaI, Hin4I, HphI, HpyAV, Ksp632I, MboII, MlyI, MmeI, MnlI, PleI, PpiI, PsrT, RleAI, SapI, SfaNI, SspD5I, Sth132I, StsI, TaqII, TspDTI, TspGWI, TspRI and Tth111I (see the list in the web site of Rebase Enzymes®: <http://rebase.neb.com/cgi-bin/outsidelist>; see also Szybalski, W., 1985, Gene, 40:169; and). The list mentioned above however is not exhaustive. Other type II enzymes known in the art and those which may be later discovered are included in the scope of the present invention.

[0119] The enzyme recognizing the homing endonuclease asymmetric restriction site is selected from the group consisting of: I-CeuI, PI-SceI, PI-PspI and I-SceI. The list mentioned above however is not exhaustive. Other homing endonucleases known in the art and those which may be later discovered are included in the scope of the present invention.

[0120] A particularly preferred tagging enzyme, according to the invention, is an enzyme which cleaves 20/18 nucleotides 3' of its recognition site forming 3' overhanging ends, such as MmeI

[0121] Artificial restriction endonucleases can also be used. These endonucleases may be prepared by protein engineering. For example, the endonuclease FokI has been engineered by insertions so that it cleaves one nucleotide further away from its recognition site on both strands of the DNA substrates. See Li and Chandrasegaran, Proc. Nat. Acad. Sciences USA 90:2764-8, 1993. Such techniques can be applied to prepare restriction endonucleases with desirable recognition sequences and desirable distances from recognition site to cleavage site.

[0122] The method further comprises producing concatemers of ditag. The concatemers may be generally about 1 to 1000 ditags, in particular 1 to 200 ditags, more in particular 8 to 20 ditags. While these are preferred concatemers, it will be apparent that the number of ditags which can be concatenated depends on the length of the individual tags and can be readily determined by those of skilled in the art without undue experimentation. After formation of concatemers, multiple tags may be cloned into a vector for sequence analysis, or ditags or concatemers can be directly sequenced without cloning by methods known to those of skill in the art.

[0123] The ditags present in a particular clone can be sequenced by standard methods (see for example, Current Protocols in Molecular Biology, supra, Unit 7) either manually or using automated methods.

[0124] As described above, the method comprises introducing the oligonucleotide comprising the at least one ditag in a vector.

[0125] With the term vector or recombinant vector it is intended a plasmid, virus or other vehicle known in the art that has been manipulated by insertion or incorporation of the ditag genetic sequences. Such vectors contain a promoter sequence which facilitates the efficient transcription. The vector typically contains an origin of replication, a promoter, as well as specific genes which allow phenotypic selection of the transformed cells. Vectors suitable for use in the present invention include for example, pBlueScript (Stratagene, La Jolla, Calif.); pBC, pZerO-1 (Invitrogen, Carlsbad, Calif.) (see **FIG. 8**) and pGEM3z (Promega, Madison, Wis.) or modified vectors thereof as well as other similar vectors known to those of skill in the art. As a particular realisation, the pGEM3z vector has been modified, and will be referred to as pGIS1 (see also **FIGS. 7 and 10**). pGEM vectors have also been disclosed in U.S. Pat. No. 4,766,072, herein incorporated by reference.

[0126] For the production of the parental nucleic acid molecule, for example full-length libraries and the GIS ditag libraries, suitable vectors are used. Accordingly, suitable vectors, which are within the scope of the present invention, are those wherein the backbone of the vector does not comprise the same restriction site comprised in the adapters flanking the parental nucleic acid molecule or the ditag, after insertion of the parental nucleic acid molecule. Preferably, the invention provides a vector wherein the vector backbone (other than within the stuffer region that is removed during insertion of the parental nucleic acid molecule) does not comprise the asymmetric restriction site and the second or further restriction site which are comprised into the adapters. In particular, the vector does not comprise the at least asymmetric II restriction site (for example type II restriction site) and the at least second or further restriction site comprised in the adapters. More preferably, the vector backbone (other than within the stuffer region that is removed during insertion of the parental nucleic acid molecule) does not comprise MmeI and BamHI.

[0127] An example of such a vector not comprising MmeI in any region outside of the stuffer is the vector pGIS1 shown in **FIG. 7** and **FIG. 10**. In pGIS1 the MmeI recognition sites were deleted by mutagenesis. The sequence is shown in **FIG. 10** and in SEQ ID NO:18. In **FIG. 10**, the stuffer region between the sites Not I and Sal I has been highlighted. The invention also related to the pGIS vector comprising the oligonucleotide according to any embodiment of the invention.

[0128] The oligonucleotide(s), ditag(s) or concatemer(s) of the invention may also be ligated into a vector for sequencing purposes.

[0129] Vectors in which the ditags are cloned can be transferred into a suitable host cell. Host cells are cells in which a vector can be propagated and its DNA expressed. The term also includes any progeny of the subject host cell. It is understood that all progeny may not be identical to the parental cell since there may be mutations that occur during replication. However, such progeny are included when the term host cell is used. Methods of stable transfer, meaning that the foreign DNA is continuously maintained in the host, are known in the art.

[0130] Transformation of a host cell with a vector containing ditag(s) may be carried out by conventional techniques as are well known to those skilled in the art. Where

the host is prokaryotic, such as *E. coli*, competent cells which are capable of DNA uptake can be prepared from cells harvested after exponential growth phase and subsequently treated by the CaCl_2 method using procedures well known in the art. Alternatively, MgCl_2 or RbCl can be used. Transformation can also be performed by electroporation or other commonly used methods in the art.

[0131] An embodiment of this is shown in **FIG. 1** and **FIG. 2**. According to this embodiment, the method of the invention comprises:

[0132] producing at least one nucleic acid molecule comprising a full-length cDNA molecule flanked by two adapters; each adapter comprising MmeI recognition sites and another recognition site, which may be BamHI, flanking the 5' terminus and 3' terminus of the full-length cDNA;

[0133] splicing the 5' terminus and the 3' terminus of the full-length cDNA to produce at least one ditag, comprising cleaving the full-length cDNA with MmeI which forms 3' overhanging tag ends, and ligating the two 5' and 3' termini tags to produce the ditag.

[0134] As shown in **FIG. 1** and **FIG. 2**, the use of restriction enzymes may leave 5' and 3' double stranded end comprising a short overhanging end (also referred to as sticky end or cohesive end) consisting of few nucleotides. In particular, by using MmeI, the produced 5' and 3' ends consist each of a 20 bp double strand and two nucleotides as 3' overhanging ends. The two tags may be followed by blunt-ending and intra-molecular self ligation to produce tagged plasmids that contain 18 bp signature sequence as 5' end and another 18 bp signature sequence as 3' end of the parental transcript. However, the number of nucleotides cut by MmeI is variable. Accordingly, the ditag obtained by using MmeI may be of 34-38 bp.

[0135] The vector which has been used for the preparation of full-length cDNA library is pGIS1. As mentioned above, pGIS1 does not contain in its backbone MmeI restriction sites, other than within the stuffer region between Not I and Sal I, this stuffer region being subsequently removed during production of the libraries.

[0136] The oligonucleotide comprising the ditag flanked by the adapters is cut out from the GIS ditag library and linked to other oligonucleotides comprising ditag and adapters to form concatemers of ditags. The concatemers of ditag are then cloned into a vector for sequencing analysis.

[0137] Before cutting the oligonucleotide out from the GIS ditag library, it can be amplified directly from the ligation (self-circularization) reaction mix, for example by PCR using suitable primers. The recovered amplified oligonucleotide comprising ditag and adapters is then linked to other oligonucleotides comprising ditag and adapters to form concatemers of ditags. The concatemers of ditag are then cloned into a vector for sequencing analysis.

[0138] The method may further comprise the steps of:

[0139] determining the nucleotide sequence of the ditag;

[0140] detecting the gene expression;

[0141] and/or comparing the determined nucleotide sequence to a database comprising genomic sequences whereby matching 5' and 3' termini sequences are identified.

[0142] In particular, the at least one ditag comprises 36 nucleotides and the first and second sequence tags comprise each 18 nucleotides.

[0143] As mentioned above, the ditag according to the invention includes the "signature" (consisting of the 5' and 3' ends) of the nucleic acid molecule which is intended to be reduced or represented. Such ditags, preferably cDNA ditags, of a library may be concatenated and sequenced. The paired 5' and 3' signature sequences (tags) of a transcript in a ditag delineate the starting and ending points of transcripts. The ditag can be split up in the two tags during data analysis and mapped head-to-head in a specific region within a reasonable distance on a chromosome of an assembled genome sequence. The genomic DNA sequence in between these two tags is the full structural content of the prospective gene, including exons and introns.

[0144] A general description of genome mapping using the ditag of the invention is shown in **FIG. 3**.

[0145] A modest sequencing run can generate sufficient data to characterize a transcriptome not only by determining the level of transcript abundance but also by defining the structure of transcripts using the revealed 5' and 3' regions. This results in about over 20-fold more efficient than EST sequencing.

[0146] Because the tags of the ditag can be matched to any genome, for example to human genomic sequences, PCR and RT-PCR primers can then be designed based on the matching genomic sequence.

[0147] Accordingly, a further aspect of the invention relates to a method for genome mapping, comprising:

[0148] preparing at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a nucleic acid molecule, the nucleic acid molecule corresponding to the full-length of a gene or fragment thereof;

[0149] mapping each of the two tags of the at least one ditag on the genome; and

[0150] defining the structural region of the corresponding gene on the genome map.

[0151] Further, it is also an aspect of the invention to provide a method of gene discovery comprising:

[0152] preparing at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a nucleic acid molecule, the nucleic acid molecule corresponding to

the full-length of a gene or fragment thereof, comparing the obtained at least one ditag with a genome map and/or a gene database;

[0153] detecting matching of the 5' and 3' termini tags on the genome map but detecting no match on one or more of the known gene database;

The method further comprises the step of recovering the full-length nucleic acid molecule corresponding to the newly discovered gene.

[0154] The invention also provides a method for recovering full-length cDNA comprising:

[0155] preparing, from a full-length cDNA library, at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a full-length cDNA;

[0156] sequencing the obtained oligonucleotide ditag;

[0157] determining the ditag of interest; and

[0158] recovering the full-length cDNA corresponding to the ditag of interest from the full-length cDNA library.

[0159] The invention also provides a method for quantifying the transcriptional activity of a gene comprising:

[0160] preparing, from a full-length cDNA library, at least one oligonucleotide comprising at least one ditag, the ditag comprising two joined first and second sequence tags, wherein the first tag comprises the 5'-terminus sequence and the second tag comprises the 3'-terminus sequence of a full-length cDNA;

[0161] sequencing the obtained oligonucleotide ditag;

[0162] determining the frequency of the sequenced ditag which corresponds to the transcriptional activity of the gene.

[0163] Having now generally described the invention, the same will be more readily understood through reference to the following examples which are provided by way of illustration, and are not intended to be limiting of the present invention.

EXAMPLES

[0164] GIS oligonucleotides for cDNA synthesis, the structure of a generic 50 bp cohesive ditag, primers used for the construction of vector pGIS1, and ds-DNA adapter which are used in the examples are listed below.

GIS Analysis Oligos for cDNA Synthesis

[0165] GsuI-Oligo dT Primer:

GsuI-oligo dT primer:
5'-GAGCTCCTTCTGGAGTTTTTTTTTTTTTTTTVN-3' (SEQ ID NO:1)

NotI/BamHI/MmeI(N)6 primer linker (top):
5'-AATTCGGCGCGCTTGGATCCGACNNNNNN (SEQ ID NO:2)

NotI/BamHI/MmeI(N) primer linker (bottom):
5'-p-GTCGGATCCAAGCGCCGCG-3' (SEQ ID NO:3)

-continued

NotI/BamHI/MmeI(N)5 primer linker (top):
 5'-AATTCGCGGCCGCTTGATCCGACGNNNNN (SEQ ID NO:4)

MmeI/BamHI/SalI adapter (top):
 5'-TCGACCCAGGATCCAACTT-3' (SEQ ID NO:5)

MmeI/BamHI/SalI adapter (bottom):
 5'-p-GTTGGATCCTGGG-3' (SEQ ID NO:6)

PMR003:
 5'-gtaaaacgacggccagt-3' (SEQ ID NO:7)

PMR004:
 5'-ggaaacagctatgacatg-3' (SEQ ID NO:8)

PMR006:
 5'-taatacgactcactatagg-3' (SEQ ID NO:9)

PMR011:
 5'-GATGTGCTGCAAGGCGATTAAG-3' (SEQ ID NO:10)

PMR012:
 5'-agcggataacaatttcacacagg-3'. (SEQ ID NO:11)

[0166] Structure of a Generic 50 bp Cohesive Ditag

5'-gatccgacxxxxxxxxxxxxxxxxnnnnnnnnnnnnnnnaagttg (SEQ ID NO:12)

GCTGXXXXXXXXXXXXXXXXNNNNNNNNNNNNNTTCAACCTAG-5' (SEQ ID NO:13)

Wherein X and N may be any of A, C, G or T.

[0167] Primers Used for the Construction of Vector pGIS
1

[0173] (3) The GIS library for clones of concatenated GIS ditags;

[0174] (4) GIS sequencing analysis.

Mme_mut1:
 5'-p-CGCTCTCCTgTACCgACCCCTgCCgCTTAC-3' (SEQ ID NO:14)

Mme_mut2:
 5'-p-AACTATCgTCTTgAgACCAACCCgGTAAG-3' (SEQ ID NO:15)

[0168] ds-DNA Adapter

5'-AATTCTCgAgCgGCCgGATATCg-3' (SEQ ID NO:16)

3'-GAGCTCGCCGGCGCTATAGCTTAA-p-5' (SEQ ID NO:17)

pGIS1 Sequence

[0169] The sequence of pGIS1 (SEQ ID NO:18) is shown in **FIG. 10**.

Example 1**The Method**

[0170] The experimental procedure of GIS ditag analysis has been carried out according to the following modules of cDNA library construction and analysis:

[0171] (1) The full-length cDNA library which introduces the MmeI sites flanking both ends of each cDNA insert;

[0172] (2) The GIS ditag library in which each clone contains a 5' 18 bp signature and a 3' 18 bp signature of a transcriptional unit;

1. GIS Full-Length cDNA Library with Addition of MmeI sites for Each cDNA Inserts

[0175] The outline of procedure of this section was as follows: starting from high quality mRNA, the first cDNA was synthesized with a GsuI-oligo dT primer (SEQ ID NO:1).

[0176] The first strand cDNA/RNA hybrids was subjected to a full-length enrichment procedure by the biotinylation-based cap-trapper approach. Any cap-trapper approach known in the art can be used, for example Carninci et al., 1996, Genomics, Vol. 37, 327-336; U.S. Pat. No. 6,143,528; Edery et al., 1995, Mol. Cell. Biol., Vol. 15, No. 6, 3363-3371).

[0177] The enriched full-length first strand cDNA was the template for second cDNA synthesis primed with adapter-primer (NotI/BamHI/MmeI-(N)5 and —(N)6, (SEQ ID NOS: 2-4) that contain a MmeI, a BamHI, and a NotI site.

[0178] After the double strand cDNA was made, the cDNA poly-A/T tail was cleaved off by GsuI restriction enzyme. GsuI is another Type-II endonuclease that cleaves

DNA 16 bp from its recognition site. At the *GsuI* cleavage end, an adapter containing a *MmeI*, a *BamHI* site, and a *Sall* cohesive end was ligated to the cDNA (SEQ ID NOS: 5-7).

[0179] Following a *NotI* digestion, the full-length cDNA was inserted into the vector pGIS1, between the *NotI* and *Sall* sites in the polylinker. The vector pGIS1 (see FIGS. 7 and 10) is modified from pGEM3z (Promega).

1-1 mRNA Preparation

[0180] The total mRNA has been prepared from mouse embryonic stem cell line E14 using Trizol reagent (Invitrogen). However, any standard method (as those described in Sambrook J. and Russell D. W., 2001, Molecular Cloning, Cold Spring Harbor Laboratory Press) may also be used.

[0181] mRNA (polyA RNA) was purified by oligo dT magnetic beads according to standard techniques (for example, Sambrook and Russell, 2001, as above). Alternatively, purification may be carried out by affinity column according to standard techniques (for example, Sambrook and Russell, 2001, as above).

1-2 First Strand cDNA Synthesis and Full-Length Selection

[0182] In this step, the first cDNA is synthesized with a *GsuI*-oligo dT primer. Then, the first strand cDNA/RNA hybrids are subjected to a full-length enrichment procedure by the biotinylation-based cap-trapper approach.

[0183] *GsuI*-Oligo dT Primer:

5' -GAGCTCCTTCTGGAGTTTTTTTTTTTTTTTTVN-3' (SEQ ID NO:1)

[0184] The following were mixed:

GsuI-oligo dT primer (7 µg/µl)	2 µl
PolyA RNA (20 µg)	18 µl

[0185] The obtained solution was heated to 65° C. for 10 min and 37° C. for 1 min. Then, spin tube in microfuge and the following substances were added:

2X GC-I buffer (Takara)	75 µl
RNase inhibitor Promega)	1 µl
10 mM dNTP (with methyl-dCTP)	4 µl
Saturated trehalose	10 µl
4.9M sorbitol	26 µl
Superscript II reverse transcriptase (Invitrogen)	15 ul

[0186] The obtained solution was incubated at 37° C. for 10 min, 42° C. for 30 min 50° C. for 20 min and 55° C. for 20 min. 2 µl of proteinase K (20 mg/ml) were added. The obtained solution was Incubated at 45° C. for 15 min followed by phenol/chloroform extraction and isopropanol precipitation (according to standard technique, eg. Sambrook and Russel, 2001, as above).

[0187] The RNA/cDNA heteroduplex was re-suspended into 44.5 µl of ddH₂O. 3 µl of 1.1 M NaOAc pH 4.5 and 2.5 µl of 100 mM NaIO₄ were added to oxidize the diol

structures of the mRNA. 50 µl of the reaction solution were incubated on ice in the dark for 45 min followed by adding 0.5 µl of 10% SDS, 11 µl of 5 M NaCl and 61 µL of isopropanol to precipitate the RNA/DNA. The precipitated RNA/DNA was resuspended in 50 µl of ddH₂O. 5 µL 1M NaOAc (pH6.1), 5 µL 10% (w/v) SDS and 150 µL 10 mM long-arm biotin hydrazide were added to biotinylate the RNA. The reaction was incubated at room temperature in dark overnight. The biotinylated RNA/DNA was precipitated by adding 5 µL 5M NaCl, 750 µL 1M RNase-free NaOAc (pH6.1), and 750 µL 100% EtOH or 200 µL of 100% Isopropanol. Incubate at -80° C. for 30 min by spill 14 krpm at 4° C. for 30 min.

[0188] The pellet was washed w/70% (v/v) EtOH/30%, DEPC-treated ddH₂O (DEPC is diethylpyrocarbonate, which is an RNase inhibitor), and 14 krpm spin was carried out at 4° C. for 10 min. The extra liquid was carefully removed. Then, the pellet was air-dried. and resuspended in 400 µL DEPC-ddH₂O. Then 50 µL 10×RNaseI buffer and 25 units RNaseI/µg of starting mRNA were added. The obtained solution was incubated at 37° C. for 30 min. 10 µL of 10 mg/mL Yeast tRNA (Ambioni) and 150 µL of 5M NaCl were added to stop the reaction.

[0189] While biotinylating the RNA-DNA heteroduplex, the Streptavidin Dynabeads were prepared as follows: 400 µL of M-280 Streptavidin beads (Dyna) were pipetted into an RNase-free Eppendorf tube, the beads placed on a magnet, left staying for at least 30 min, and then the

supernatant was removed. The beads were re-suspended in 400 µL 1× binding buffer (2M NaCl, 50 mM EDTA, pH 8.0). The tube was placed on a magnet, waited at least 30 min, and then the supernatant was removed. The 1× binding buffer wash was repeated for 2 more times. The beads were re-suspended in 400 µL 1× binding buffer with 100 µg of Yeast tRNA, and then incubated at 4° C. for 30 min with occasional mixing. The tube was placed on a magnet stand, waited at least 30 seconds, and the supernatant was removed. The beads were washed with 1× binding buffer for 3 times. The beads and RNA/DNA heteroduplex were mixed (the total volume now was 660 µL, and the binding condition was at 1 M NaCl). The mixture was rotated at room temperature for 30 mini.

[0190] The tube was placed on a magnet stand, waited at least 30 seconds, and the supernatant removed (the supernatant was saved as "unbound").

[0191] The beads were washed two times with 400 µL of 1× binding buffer. Washed with 400 µL of 0.4% (w/v) SDS plus 50 µg/mL Yeast tRNA. Washed with 400 µL of 1× wash buffer (10 mM Tris-HCl pH7.5, 0.2 mM EDTA, 10 mM NaCl & 20% (v/v) glycerol, 40 µg/mL Yeast tRNA). And washed w/400 µL of 50 µg/mL Yeast tRNA. For all washes the tube was placed on a magnet stand, waited for at least 30 seconds, and the supernatant was removed.

[0192] The first strand cDNA was released by alkali hydrolysis of RNA. The following was added: 50 µL 50 mM NaOH and 5 mM EDTA (pH8.0). The tube was rotated at 65° C. for 10 min. The tube was placed on a magnet stand, and the supernatant transferred to another tube containing 50 µL 1M Tris-Cl (pH7.5).

[0193] The lysis procedure was repeated for 2 more times. The final volume of supernatant was 300 µL (containing the first strand cDNA).

[0194] The cDNA was extracted by phenol/chloroform extraction and precipitate by 1 mL ethanol with glycogen. The cDNA was re-suspended in 5 µL LoTE (0.1×) LoTE is low salt Tris-EDTA buffer (3 mM Tris-HCl pH 7.5 and 0.2 mM EDTA pH7.5)).

1-3 Second Strand cDNA Synthesis

[0195] The following reagents were added to the each corresponding tube on ice.

cDNA in LoTE	5 µL
Linker (N5)	1.6 µg
Linker (N6)	0.4 µg
Soln II (Takara ligation kit)	10 µL
Soln I (Takara ligation kit)	20 µL

[0196] Linker (N6) is:

NotI/BamHI/MmeI(N)6 primer linker (top):
5'-AATTCGCGGCCGCTTGGATCCGACNNNNNN (SEQ ID NO:2)

NotI/BamHI/MmeI(N) primer linker (bottom):
5'-p-GTCGGATCCAAGCGCCGCG-3' (SEQ ID NO:3)

[0197] Linker (N5) is:

NotI/BamHI/MmeI(N)5 primer linker (top):
5'-AATTCGCGGCCGCTTGGATCCGACGNNNNNN (SEQ ID NO:4)

NotI/BamHI/MmeI(N) primer linker (bottom): is the sequence (SEQ ID NO:3) indicated above.

[0198] The cDNA and linker mixture was incubated at 16° C. overnight. And the following were added:

ddH ₂ O	20 µL
10XExTaq buffer (Takara)	8 µL
2.5 mM dNTP	8 µL
ExTaq polymerase (Takara)	4 µL

[0199] The mixture was preheated in a thermo-cycler 65° C., 5 min→68° C., 30 min→72° C., 10 min., followed by phenol/chloroform extraction and ethanol ppt with glyco-gen, and re-suspended in 85 µl ddH₂O.

1-4. Removal of polyA Tail by GsuI Digestion

[0200] The following reagents were added to the tube.

cDNA	85 µL
GsuI (Fermentas)	5 µL
10X bufferB (Fermentas)	10 µL

[0201] The mixture was incubated at 30° C. for 2 hours, followed by phenol/chloroform and ethanol precipitation. The pellet was re-suspended in 10 µl ddH₂O, and the following 3' adaptor ligation reaction was carried out.

1-5. Addition of 3' Adaptor with MmeI and BamHI and SalI Sites.

[0202] The following components were added to the tube containing 10 µl of sample. The 10 µl of sample was the double-stranded full-length cDNA which has had the poly(A) tail removed by GsuI digestion.

5X ligation buffer	10 µL
GsuI SalI adapter (0.4 µg/µL)	25 µL

[0203] [The GsuI SalI Adapter is MmeI/BamHI/SalI Adapter)

T4 DNA ligase (5 units/ul) (Invitrogen)	5 µL
---	------

[0204]

MmeI/BamHI/SalI adapter (top):
5'-TCGACCCAGGATCCAACCTT-3' (SEQ ID NO:5)

MmeI/BamHI/SalI adapter (bottom):
5'-p-GTTGGATCCTGGG-3' (SEQ ID NO:6)

[0205] The reaction was incubated at 16° C. overnight, followed by phenol/chloroform extraction and ethanol pre-cipitation, and the pellet re-suspended in 41 µl dH₂O.

1-6. NotI Digestion and cDNA Size Fractionation

[0206] The following were added on ice and in order:

NEB Buffer 3	5 µL
NotI (10 units/µl) (NEB)	4 µL

[0207] The obtained solution was incubated at 37° C. for 1-2 hours.

[0208] cDNA Size Fractionation Columns were prepared (the Invitrogen instructions were followed: uncap the column (bottom first) and allow it to drain completely; wash 5 times with 800 μ L $T_{10}E_{0.1}N_{25}$ Buffer, allowing the column to drain completely each time). The DNA sample was loaded onto the column. The flow-through was collected in an Eppendorf tube (fraction 1). 100 μ L of $T_{10}E_{0.1}N_{25}$ Buffer were added. The flow-through was collected in an Eppendorf tube (fraction 2). Another 100 μ L of $T_{10}E_{0.1}N_{25}$ Buffer was added. The flow-through collected, one drop per pre-numbered Eppendorf tube (beginning with fraction 3, each drop was about 30-40 μ L).

[0209] Whenever the column runs dry, another 100 μ L of $T_{10}E_{0.1}N_{25}$ Buffer may be added.

[0210] Up to drop 20 should be collected (according to the Invitrogen protocol). 3 μ L of each fraction were run on agarose gel to visualize the cDNA size in each fraction. Pool fractions were showing cDNA ≥ 1.0 kbp (usually up to 2-3 kbp). Pooled samples were kept neat (using a cuvette soaked at least 30' in slightly acidified 100% EtOH, rinsed 5 times with ddH₂O, and saving sample. This is what has to be ligated).

[0211] If only one fraction is to be used, precipitate it and use the half to all of it, depending on what the gel looks like.

[0212] At this point the cDNA fragments have the NotI cohesive end at 5' side and SalI cohesive end at 3' side, and are ready to be cloned in vector.

1-7. Ligation of cDNA with Linearized Plasmid pGIS1.

[0213] 1-7-1 The Cloning Vector pGIS1 was Prepared by NotI and SalI Digestion.

[0214] The vector sequence of pGIS1 is shown in **FIGS. 7 and 9**.

pGIS1 Cloning Vector Construction

(I) Site-Specific Mutagenesis of pGEM3z to Create MmeI-Minus Vector

[0215] The vector pGIS-1 was derived from pGEM3z (Promega). pGEM3z originally contained two MmeI recognition sites that were knocked-out by site-directed mutagenesis. The QuikChange Multi kit (Stratagene) was used, together with mutagenic primers:

Mme_mut1:
5'-p-CgCTCTCCTgTACCGACCCtGCCgCTTAC-3' (SEQ ID NO:14)

Mme_mut2:
5'-p-AACTATCgTCTTgAgACCAACCCgTAAg-3' (SEQ ID NO:15)

(II) Modification of Polylinker Region

[0216] The polylinker region was modified by simple insertion of a ds-DNA adapter at the existing EcoRI site. Additional recognition sites thus introduced are: XhoI, NotI and EcoRV (EcoRV is deleted upon insertion of the stuffer fragment (see below)).

[0217] ds-DNA Adapter:

5'-AATTCTCgAgCggCCgCgATATCg-3' (SEQ ID NO:16)

3'-GAGCTCGCCGGCGCTATAGCTTAA-p-5' (SEQ ID NO:17)

(III) Stuffer Fragment Insertion

[0218] An approximately 690 bp fragment was inserted between the NotI and SalI sites of the modified vector (see vector sequence in **FIG. 10**). This facilitated the production of NotI/SalI double-digested vector, as the stuffer can be clearly visualized and excised during gel-purification.

[0219] The linearized plasmid was gel purified.

[0220] 1-7-2 The cDNA was Ligated to the pGIS I Vector Overnight and the Constructs were Transferred into Electrocompetent *E. coli* TOP10 Cells by Electroporation according to Standard Techniques (see Sambrook and Russel, 2001, as above).

1-8. Library QC (QC=Quality Check)

[0221] A dilution series of 1-100 μ L of transformants was plated out onto LB agar plates with antibiotic selection. The colonies were incubated overnight and counted to determine the library titer.

[0222] Between 24 to 96 colonies (arbitrary numbers) were picked and the inserts size determined by direct colony PCR and agarose gel electrophoresis (according to standard techniques, eg. Sambrook and Russel, 2001, see above). The percentage of cDNA insert and the average insert size were estimated.

[0223] At this stage, the GIS full-length cDNA library may be stored as ligation reactions or as transformants in *E. coli* cells, according to standard methodology (Sambrook and Russel, 2001, see above).

Example 2

2. GIS Ditag Library

[0224] The cDNA clones made from steps 1-1 to 1-8 contained a MmeI site (TCCGAC) at the 5 side and another MmeI site (TCCAAC) in reverse orientation at the 3 end. Note that these two MmeI recognition sites are two isoforms that can be recognized by MmeI (TCCRAC 20/18, where R=(A/G)). The sequence difference here will be useful later for directional indication. MmeI restriction enzyme will

cleave these clones 20 bp into the cDNA fragments from their 5 and 3 ends. Consequently, despite the variable sizes of the digested cDNA, the vector plus the 20 bp cDNA signature tags on each end of all clones will be of a constant size that can be easily recognized upon agarose gel electrophoresis, and can be easily purified from the unwanted cDNA fragments.

[0225] The gel-purified vector plus tags can then be self-ligated to give a agged plasmid containing the 5 and 3 GIS signature tags.

2-1. Plasmid Preparation

[0226] The GIS full-length cDNA library was amplified once by plating an appropriate number of clones on large (22×22 cm) agar plates (Genetix). The number of colonies required was determined by the estimated transcriptome size. After an overnight 37 C incubation, the resultant bacterial colonies were harvested and pelleted by centrifugation at 3000 g for 30 min. Plasmid DNA preparation was performed using the Qiagen HiSpeed Plasmid Maxi kit. The quality of the DNA obtained was examined by agarose gel electrophoresis and restriction digestion. Approximately 300,000 colonies can be processed to yield at least 1 mg of plasmid DNA.

2-2. MmeI Digestion

[0227] Approximately 10 µg of plasmid DNA was digested using MmeI as per manufacturer's conditions

mately 20 ng/µl. Self-ligation (intramolecular recircularization) was carried out as follows:

Approx. 350 ng DNA	15.0 µL
Ligation Solution I (Takara Ligation Kit 2)	15.0 µL

[0232] Incubated at 16 C, 2 hr to overnight

[0233] 2-5. Creation of Di-Signature Tags (Ditags)

[0234] The goal of this step was to obtain the GIS di-signature tags in a form quantitatively representative of the original cDNA library from which the tagged-plasmids were derived.

[0235] Structure of a Generic 50 bp Cohesive Ditag

5' -GATCCGACXXXXXXXXXXXXXXXXXXXXNNNNNNNNNNNNNNNNNAAGTTG (SEQ ID NO:12)

GCTGXXXXXXXXXXXXXXXXXXXXNNNNNNNNNNNNNNNNNTTCAACCTAG-5' (SEQ ID NO:13)

(NEB), ensuring that the number of units of enzyme used was always less than 4-fold excess to prevent methylation-induced inhibition. Digestion proceeded at 37 C for 2-6 hrs.

[0228] An aliquot of the digestion reaction was examined on an agarose gel: a strong band of approximately 2800 bp in size corresponding to the linearized vector containing the GIS signature tags were easily observed, together with a number of fragments derived from the excision of cDNA from the original plasmids (see **FIG. 4**).

2-3. Linear Vector-GIS Ditag Purification

[0229] The digestion reaction was electrophoresed on 0.7% agarose, and the 2800 bp vector-GIS tag band was excised and purified using the Qiagen agarose gel extraction kit.

2-4. Vector-GIS Ditag Self Ligation to Create Agged-Plasmids

[0230] MmeI digestion resulted in a 2 bp overhang on both the 5 and 3 signature tags. These were removed (polished off) using T4 DNA polymerase (Promega), leaving behind 18 bp tags:

(0.5–2.0 ug) DNA	50 µL
10x Y+/TANGO buffer (Fermentas)	6.0 µL
0.1 M DTT	0.3 µL
T4 DNA polymerase	5 units/µg
10 mM dNTP	0.6 µL
ddH ₂ O	to 60.0 µL

Incubated at 37 C, for 5 min, then inactivate at 75 C for 10 min

[0231] The purified, blunted DNA was then ethanol precipitated and resuspended at a concentration of approxi-

Wherein X and N may be any of A, C, G or T.

[0236] We used two approaches to this:

[0237] (i) Bacterial transformation, tagged-plasmid purification and restriction digest to release 50 bp cohesive ditags;

[0238] (ii) Direct PCR on the ligation reaction followed by restriction digest of the PCR products to release 50 bp cohesive ditags.

2-5-1 Transformation and Propagation; Preparation of Tagged-Plasmids (See **FIG. 1**)

[0239] 1 µl of the ligation reaction (Section 2-4) were transformed per 50 µl of electrocompetent TOP10 cells (Invitrogen) by electroporation. Recovered in 1 ml SOC media at 37 C for 1 hr, then plated out several dilutions on LB agar+ampicillin for QC and titering.

[0240] QC (Quality Check): plasmid DNA was prepared from several colonies and tested by digestion with BamHI: tagged-plasmids release a 50 bp cohesive ditag.

[0241] This process was then scaled-up by plating the remaining culture on large agar plates, and performing maxipreps using Qiagen HiSpeed Plasmid Maxi kit.

[0242] As an example, approximately 5,000 colonies was processed to yield at least 40 ug of tagged-plasmid DNA.

[0243] This plasmid DNA was then BamHI-digested to generate 50 bp cohesive ditags (see **FIG. 5** as example result).

2-5-2 PCR-Based Retrieval of Cohesive Ditags (See **FIG. 2**)

[0244] PCR was performed on the ligation reaction using primers PMR003 and PMR004 that bind to vector sequences flanking the ditags.

PMR003: 5' -GTAAAACGACGGCCAGT-3' (SEQ ID NO:7)

PMR004: 5' -GGAAACAGCTATGACCATG-3' (SEQ ID NO:8)

[0245] The amount of starting material was determined empirically by doing a series of dilutions and choosing the conditions that result in a clean, specific PCR product of approximately 200 bp

[0246] (e.g. 1:200) diluted ligation reaction 5.0 μ L

10x HiFi buffer	2.0 μ L
10 mM dNTP	0.4 μ L
PMR003 (100 ng/ μ L)	1.0 μ L
PMR004 (100 ng/ μ L)	1.0 μ L
Eppendorf TripleMaster polymerase	0.2 μ L
dH ₂ O	10.4 μ L

(the HiFi buffer was the reaction buffer provided with the Eppendorf TripleMaster enzyme)

[0247] Thermo-Cycling Conditions:

[0248] Step 1: 95 C \times 2 min

[0249] Step 2: 95 C \times 30 sec

[0250] Step 3: 55 C \times 1 ml-n

[0251] Step 4: 72 C \times 30 sec

[0252] Go to step 2, repeat steps (2-4) 24 \times

[0253] Step 5: 72 C \times 4 min 16° C. forever

[0254] The PCR products were analyzed on a 1.5% agarose gel.

[0255] For negative controls, the PCR reaction was performed using (i) no template, and (ii) no ligase. To obtain sufficient 200 bp PCR product for subsequent 50 bp cohesive ditag production, the reaction was scaled-up: do 96 PCR reactions using a 96-well PCR plate; this generates approx. 50 μ g of 200 bp ditag. The individual PCR reactions are then combined and ethanol precipitated before Bam HI digest to generate 50 bp cohesive ditags (see FIG. 6 as an example result).

3. GIS Library

3-1. Tagged-Plasmid Preparation

[0256] This applies only to the bacterial transformation-based approach (see Section 2-5-1).

[0257] 3-2. BamHI Digestion and Purification of GIS Tags

[0258] 3-2-1 BamHI Digestion of Tagged-Plasmids (Section 2-5-1) Released 50 bp Cohesive Ditags:

DNA (tagged-plasmids)	40 μ g
10x unique BamHI buffer (NEB)	100 μ L
100x BSA	10 μ L
BamHI (20 U/ μ L, NEB)	10 μ L
dH ₂ O	to 1 mL

[0259] The choice of value of 40 μ g of DNA (tagged-plasmids) was arbitrary.

[0260] Aliquots were divided into 10 \times 100 μ l for more efficient digestion, and incubated at 37° C., for 4 hrs.

[0261] After digest, they were inactivated at 65 C, for 15 min, then phenol-chloroform extraction and ethanol precipitation were performed. Then, the pellet comprising 50 bp cohesive ditags and the rest of the cleavage products after the BamHI digest was resuspended in LoTE buffer for gel-purification.

[0262] 3-2-2 BamHI Digestion of or 200 bp Ditags Retrieved by PCR (Section 2-5-2) Released 50 bp Cohesive Ditags:

DNA (PCR products)	40 μ g
10x unique BamHI buffer (NEB)	100 μ L
100x BSA	10 μ L
BamHI (20 U/ μ L, NEB)	10 μ L
dH ₂ O	to 1 mL

[0263] The choice of value of 40 μ g of DNA (tagged-plasmids) was arbitrary.

[0264] Aliquots were divide into 10 \times 100 μ l for more efficient digestion, incubated at 37° C., for 4 hrs.

[0265] After digest, they were inactivated at 65 C, for 15 min, then phenol-chloroform extraction and ethanol precipitation were performed. Then, the pellet comprising 50 bp cohesive ditags and the rest of the cleavage products after the BamHI digest was resuspended in LoTE buffer for gel-purification.

3-3 Gel-Purification of 50 bp Cohesive Ditags

[0266] The BamHI-digested DNA according to both section 3-2-1 or 3-2-2 was separated on a large (Hoefer Ruby 600,15 \times 15 cm, 1.5 mm thick) 10% polyacrylamide gel. Electrophoresis proceeded at 200V for approx. 2 hrs until the Bromophenol Blue (standard tracking dye) band almost reached the bottom of the gel. The gel was stained in SYBR Green I (Molecular Probes, Inc.) for 30 min before visualisation and excision of the 50 bp cohesive ditags.

[0267] At this stage it is convenient not to load more than 5 μ g per lane, or fluorescence quenching occurs.

[0268] The 50 bp cohesive ditags were excised and collected into 0.6 ml microfuge tubes (2 gel pieces per tube) which have been pierced at the bottom with a 21 G needle. This pierced tube was placed inside a 1.7 ml microfuge tube, and centrifuged at 12 K g, 4 C for 2-5 min. The gel pieces were thus shredded and collected in the 1.7 ml tube.

[0269] 150 μ l of LoTE:NH₄OAc (125:25) were added to each tube and left overnight at 4 C to elute. The next day, the eluate was collected with the aid of microspin filter units (SpinX, Costar), and ethanol precipitation performed to retrieve the purified 50 bp ditags, which were resuspended in LoTE. Starting from 70 μ g 200 bp ditag, we expected to retrieve several hundred ng of 50 bp ditag.

3-4. Ditag Concatenation and Gel-Purification

[0270] Some optimization (ligation time, amount of starting material) may be necessary to ensure that the concatenation of the 50 bp ditags results in a smear of products ranging from approx. 300 bp to >1000 bp. The conditions below are suggested as a starting point:

50 bp cohesive ditags	150–500 ng
5x buffer (with PEG; BRL)	2.0 μ L
T4 DNA ligase (5 U/ μ L)	1.0 μ L
dH ₂ O	to 10 μ L

[0271] Incubated at 16° C. for 1 hr.

[0272] Loading buffer was added and the entire sample heated at 65 C for 15 min. The sample loaded in a single well of an 8% polyacrylamide minigel and run at 200V for about 1 hr, or until Bromophenol Blue was about 2 cm from bottom.

[0273] The smear of ligation products can be excised as 2 or more fractions, eg. 200-500 bp; 500-1000 bp; >1000 bp.

[0274] Elution of DNA from the gel pieces was performed as detailed in Section 3-3. The eluate was extracted with phenol-chloroform then ethanol precipitated. Resuspend the DNA pellet in 6 μ l LoTE.

3-5. Cloning of Concatemers

[0275] The cloning vector was prepared by digesting 2 μ g of pZErO-1 plasmid DNA (Invitrogen) (**FIG. 8**) (**FIG. 8** shows the sequencing/PCR primer binding sites) with 10 units of BamHI for 3 hours at 37 C. The digested DNA was phenol-chloroform extracted and ethanol precipitated, then resuspended in LoTE at a concentration of 33 ng/ μ l. The ligation reaction was performed as follows:

Concatemer DNA	6.0 μ L
BamHI/pZErO-1	1.0 μ L
5x ligase buffer	2.0 μ L
T4 DNA ligase (5 U/ μ L)	1.0 μ L

Incubated at 16° C. overnight.

[0276] The vector self-ligation was also performed in parallel as a control.

[0277] The ligation products were purified before electroporation. The phenol-chloroform extraction was followed by ethanol precipitation; the pellet was washed 3 times with 75% ethanol before re-suspending in 121 μ l TE (0.1X). 1 μ l of this DNA was used to transform 50 μ l of electro-competent TOP10 bacterial cells. After recovery (see also Section 2-5-1), 50 μ l were plated on a small agar plate (containing Low Salt LB agar (Lennox L) plus Zeocin (50 g/ml) and IPTG (50 μ g/ml) and incubated overnight at 37 C. As a background control, bacteria were plated out that have been similarly transformed with the vector self-ligation reaction above. (IPTG is optional when using TOP10 cells but may reduce background).

3-6. GIS Library QC (Quality Check)

[0278] The following day, 10-30 colonies were picked to check for insert size by PCR. For each reaction, a single colony was picked into a PCR tube containing:

10x HiFi buffer	2.0 μ L
10 mM dNTP	0.4 μ L
PMR003 (100 ng/ μ l)	1.0 μ L
PMR004 (100 ng/ μ l)	1.0 μ L
Eppendorf TripleMaster polymerase	0.2 μ L
dH ₂ O	11.4 μ L

[0279] Thermo-Cycling Conditions:

[0280] Step 1: 95 C×2 min

[0281] Step 2: 95 C×30 sec

[0282] Step 3: 55 C×1 min

[0283] Step 4: 72 C×3 min

[0284] Go to step 2, repeat steps (2-4) 24×

[0285] Step 5: 72 C×4 min

[0286] 16° C. forever

[0287] The PCR products were visualized on 1% agarose gel. A typical result is shown in **FIG. 9**.

[0288] The primer pair PMR003/PMR004 (SEQ ID NO:7/SEQ ID NO:8) gives a band of approx. 220 bp in the absence of any cloned insert. If the quality of the library thus produced appears good, the remaining transformation mixture can be plated out (Section 3-5) on large agar plates in preparation for DNA sequencing analysis.

[0289] The primer pair PMR003/PMR004 is also convenient for checking the quality of the library, but for the actual preparation of PCR templates for sequencing, primer pair PMR012/PMR003 (SEQ ID NO:11/SEQ ID NO:7) were preferred (see Section 4-2).

PMR012:
5' -AGCGGATAACAATTTTCACACAGG-3' . (SEQ ID NO:11)

4. Sequencing Analysis of GIS Tags

4-1. Library Plating and Colony Picking

[0290] The transformed TOP10 (Invitrogen) bacteria cells were plated out on 22×22 cm agar plates with colony density less than 3,000 per plate. Individual colonies were picked and cultured in 384-well plates with LB plus Zeocin (see above in section 3.5) at 37° C. overnight. Multiple copies of 384-well plates are replicated and stored in -80° C.

4-2. Template Preparation

[0291] Bacterial cultures in 384-well plates were inoculated in pre-mixed PCR cocktails. PCR was performed using primer pair PMR012/PMR003.

[0292] This primer pair gives a band of 245 bp in the absence of any concatemer insert. Nonetheless, this set of primers is preferred as it allows the use of sequencing primers PMR004 (M13 reverse; 68 bp from BamHI site) and

PMR006 (SEQ ID NO:9)(M13 forward; 87 bp from BamHI site).

PMR006: 5'-TAATACGACTCACTATAGGG-3' (SEQ ID NO:9)

4-3. Sequencing

[0293] PCR templates were sequenced using the sequencing primers PMR004 and PMR006 to sequence in both directions.

Example 3

[0294] The GIS analysis method according to any embodiment of the invention is a complete gene discovery platform. It combines full-length cDNA library construction, cDNA tag sequencing, genome mapping and annotation into one operation from the same starting materials. For example, to study the genes expressed in human stem cells, we start with the stem cell mRNA, construct a stem cell GIS full-length cDNA library, and then the GIS library. We will only need

to sequence 50,000 clones of the GIS library to reveal over a million transcripts. Such deep sampling will allow us to capture nearly all unique transcripts expressed in the human stem cell transcriptome. Each of the GIS ditags can be specifically mapped to the genome and therefore define the structural regions of the corresponding genes on the chromosomes. Most of the GIS ditags map to known genes on chromosomes and the counts of the GIS ditags provide the measurement of expression activity. Some of the GIS ditags may map to desert ("no1 gene") regions of the genome, which may suggest the identification of new genes that are expressed in the stem cell transcriptome. In such a way the genome annotation for genes is further refined by this whole transcriptome-to-whole genome approach. Based on the GIS ditag sequences, these putative new genes can be readily cloned from the original GIS full-length cDNA library.

[0295] We can apply this GIS gene discovery system not only to human stem cells, but also to all other biological systems, such as development of cells, tissues and organs of human and model organisms.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 29

<210> SEQ ID NO 1
 <211> LENGTH: 33
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: oligonucleotide primer with homology to
 bacterial cloning vector
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(33)
 <223> OTHER INFORMATION: n is a,c,g, or t
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(33)
 <223> OTHER INFORMATION: v is a,c,g

<400> SEQUENCE: 1

gagctccttc tggagttttt tttttttttt tvn

33

<210> SEQ ID NO 2
 <211> LENGTH: 30
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:
 <223> OTHER INFORMATION: oligonucleotide primer with homology to
 bacterial cloning vector
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(30)
 <223> OTHER INFORMATION: n is a,t,c or g

<400> SEQUENCE: 2

aattcgcggc cgcttgatc cgacnnnnnn

30

<210> SEQ ID NO 3
 <211> LENGTH: 20
 <212> TYPE: DNA
 <213> ORGANISM: Artificial
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<400> SEQUENCE: 3

gtcggatcca agcggccgcg

20

<210> SEQ ID NO 4

<211> LENGTH: 30

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(30)

<223> OTHER INFORMATION: n is a,t,c or g

<400> SEQUENCE: 4

aattcgcggc cgcttgatc cgacgnnnnn

30

<210> SEQ ID NO 5

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<400> SEQUENCE: 5

tcgacccagg atccaactt

19

<210> SEQ ID NO 6

<211> LENGTH: 13

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(1)

<223> OTHER INFORMATION: phosphorylation

<400> SEQUENCE: 6

gttggatcct ggg

13

<210> SEQ ID NO 7

<211> LENGTH: 17

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<400> SEQUENCE: 7

gtaaaacgac ggccagt

17

<210> SEQ ID NO 8

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

-continued

<400> SEQUENCE: 8

ggaaacagct atgaccatg

19

<210> SEQ ID NO 9

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<400> SEQUENCE: 9

taatacgact cactataggg

20

<210> SEQ ID NO 10

<211> LENGTH: 22

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<400> SEQUENCE: 10

gatgtgctgc aaggcgatta ag

22

<210> SEQ ID NO 11

<211> LENGTH: 23

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: oligonucleotide primer with homology to
bacterial cloning vector

<400> SEQUENCE: 11

agcggataac aatttcacac agg

23

<210> SEQ ID NO 12

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Oligionucleotide with homolgy to a
bacteria cloning vector

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(48)

<223> OTHER INFORMATION: n is a,t,c or g

<400> SEQUENCE: 12

gatccgacnn nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnaagttg

48

<210> SEQ ID NO 13

<211> LENGTH: 48

<212> TYPE: DNA

<213> ORGANISM: Artificial

<220> FEATURE:

<223> OTHER INFORMATION: Oligionucleotide with homolgy to a
bacteria cloning vector

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (1)..(48)

<223> OTHER INFORMATION: n is a,t,c or g

<400> SEQUENCE: 13

gatccaactt nnnnnnnnnn nnnnnnnnnn nnnnnnnnnn nnnngtcg

48

-continued

<210> SEQ ID NO 14
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligionucleotide primer with homolgy to a
bacteria cloning vector

<400> SEQUENCE: 14

cgctctcctg taccgaccct gccgcttac 29

<210> SEQ ID NO 15
<211> LENGTH: 29
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligionucleotide primer with homolgy to a
bacteria cloning vector

<400> SEQUENCE: 15

aactatcgtc ttgagaccaa cccggtaag 29

<210> SEQ ID NO 16
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligionucleotide adapter with homolgy to a
bacteria cloning vector

<400> SEQUENCE: 16

aattctcgag cggccgcgat atcg 24

<210> SEQ ID NO 17
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: Oligionucleotide adapter with homolgy to a
bacteria cloning vector

<400> SEQUENCE: 17

gagctcgccg gcgctatagc ttaa 24

<210> SEQ ID NO 18
<211> LENGTH: 3404
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: bacterial cloning vector

<400> SEQUENCE: 18

gggcgaattc tcgagcggcc gcggatccga cgagagcgcc tgcgtacggc tcgccgcggt 60
ggctggcgct acttcggagg agcccgacgc ggcgcggtcg tttttataca ttcccgcgcg 120
gaggcaacgg aagggcgggg cgcctcgtga ttaggccgcg gaggtcacag gctctgttgt 180
catgaagggtg aaaattaaat gttggaatgg tgtggccact tggctctggg tagccaatga 240
tgagaactgc ggcactctgca ggatggcggt taatggctgc tgtccagact gtaagggtgcc 300
tggtgatgac tgccccctcg tgtggggaca gtgctccac tgcttcaca tgcaactgcat 360
cctcaagtgg ctgaatgcgc agcaggtgca gcagcactgc cccatgtgtc gccaggagtg 420

-continued

gaagttcaaa	gagtgaagcc	cgtgccgtgc	cacttccctc	tcctgtgctg	tgccaggctc	480
agcccccttc	ctccctcccc	ttccccagat	acagcacccc	aagtccccctc	cacacagcac	540
agtggtgccc	agagatctcg	gtctgtgccg	gggacaagga	tgctttctgt	ttggctggga	600
caagggtgaa	aggagctttg	ctgactgttt	tgttttccca	tcacattgac	actttattca	660
ataagtaaaa	ctcattacag	ttccaagtcg	gatcctgggt	cgacctgcag	gcatgcaagc	720
ttgagtattc	tatagtgtca	cctaaatagc	ttggcgtaat	catggtcata	gctgtttcct	780
gtgtgaaatt	gttatccgct	cacaattcca	cacaacatac	gagccggaag	cataaagtgt	840
aaagcctggg	gtgcctaattg	agtgaagctaa	ctcacattaa	ttgcgttgcg	ctcactgccc	900
gctttccagt	cgggaaacct	gtcgtgccag	ctgcattaat	gaatcggcca	acgcgcgggg	960
agaggcggtt	tgcgatttgg	gcgctcttcc	gcttcctcgc	tcactgactc	gctgcgctcg	1020
gtcgttcggc	tgccggcgagc	ggtatcagct	cactcaaagg	cggtaatacg	gttatccaca	1080
gaatcagggg	ataacgcagg	aaagaacatg	tgagcaaaag	gccagcaaaa	ggccaggaac	1140
cgtaaaaagg	ccgcgttgct	ggcgtttttc	gataggtctc	gccccctga	cgagcatcac	1200
aaaaatcgac	gctcaagtca	gaggtggcga	aacccgacag	gactataaag	ataccaggcg	1260
tttccccctg	gaagctccct	cgtgcgctct	cctgtaccga	ccctgccgct	taccggatac	1320
ctgtccgcct	ttctcccttc	gggaagcgtg	gcgctttctc	atagctcacg	ctgtaggtat	1380
ctcagttcgg	tgtaggtcgt	tcgctccaag	ctgggctgtg	tgcacgaacc	ccccgttcag	1440
cccgaccgct	gcgccttatc	cggtaactat	cgtcttgaga	ccaacccggt	aagacacgac	1500
ttatcgccac	tggcagcagc	cactggtaac	aggattagca	gagcgaggta	tgtaggcggt	1560
gctacagagt	tcttgaagtg	gtggcctaac	tacggctaca	ctagaaggac	agtatttggt	1620
atctgcgctc	tgctgaagcc	agttaccttc	ggaaaaagag	ttggtagctc	ttgatccggc	1680
aaacaaacca	ccgctggtag	cgggtggtttt	tttgtttgca	agcagcagat	tacgcgcaga	1740
aaaaaaggat	ctcaagaaga	tcctttgatc	ttttctacgg	ggtctgacgc	tcagtggaac	1800
gaaaactcac	gttaagggat	tttgggtcatg	agattatcaa	aaaggatctt	cacctagatc	1860
cttttaaatt	aaaaatgaag	ttttaaatca	atctaaagta	tatatgagta	aacttggtct	1920
gacagttacc	aatgcttaat	cagtgaggca	cctatctcag	cgatctgtct	atttcgttca	1980
tccatagttg	cctgactccc	cgtcgtgtag	ataactacga	tacgggaggg	cttaccatct	2040
ggccccagtg	ctgcaatgat	accgcgagac	ccacgctcac	cggctccaga	tttatcagca	2100
ataaaccagc	cagccggaag	ggccgagcgc	agaagtggtc	ctgcaacttt	atccgcctcc	2160
atccagtcta	ttaattgttg	ccgggaagct	agagtaagta	gttcgccagt	taatagtttg	2220
cgcaacgttg	ttggcattgc	tacaggcatc	gtggtgtcac	gctcgtcgtt	tggtatggct	2280
tcattcagct	ccggttccca	acgatcaagg	cgagttacat	gatcccccat	gttgtgcaaa	2340
aaagcggtta	gctccttcgg	tcctccgatc	gttgtcagaa	gtaagttggc	cgcagtgtta	2400
tcactcatgg	ttatggcagc	actgcataat	tctcttactg	tcatgccatc	cgtaagatgc	2460
ttttctgtga	ctggtgagta	ctcaaccaag	tcattctgag	aatagtgtat	gcggcgaccg	2520
agttgctctt	gcccggcgctc	aatacgggat	aataccgcgc	cacatagcag	aactttaaaa	2580
gtgctcatca	ttggaaaacg	ttcttcgggg	cgaaaactct	caaggatctt	accgctgttg	2640
agatccagtt	cgatgtaacc	cactcgtgca	cccaactgat	cttcagcatc	ttttactttc	2700

-continued

accagcgttt ctgggtgagc aaaaacagga aggcaaatg ccgcaaaaaa gggaataagg 2760
gcgacacgga aatgttgaat actcatactc ttcctttttc aatattattg aagcatttat 2820
cagggttatt gtctcatgag cggatacata tttgaatgta tttagaaaaa taaacaaata 2880
ggggttccgc gcacatttcc ccgaaaagtg ccacctgacg tctaagaaac cattattatc 2940
atgacattaa cctataaaaa taggcgtatc acgaggccct ttcgtctcgc gcgtttcggg 3000
gatgacgggtg aaaacctctg acacatgcag ctcccgagga cggtcacagc ttgtctgtaa 3060
gcggatgccg ggagcagaca agcccgtcag ggcgcgtcag cgggtgttgg cgggtgtcgg 3120
ggctggctta actatgcggc atcagagcag attgtactga gagtgcacca tatgcgggtg 3180
gaaataccgc acagatgcgt aaggagaaaa taccgcatca ggcgccattc gccattcagg 3240
ctgcgcaact gttgggaagg gcgatcgggtg cgggcctctt cgctattacg ccagctggcg 3300
aaagggggat gtgctgcaag gcgattaagt tgggtaacgc cagggttttc ccagtcacga 3360
cgttgtaaaa cgacggccag tgaattgtaa tacgactcac tata 3404

<210> SEQ ID NO 19
<211> LENGTH: 10
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: mammalian p53 consensus sequence
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: r is a purine (A or G)
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: w is A or T
<220> FEATURE:
<221> NAME/KEY: misc_feature
<223> OTHER INFORMATION: y is a pyrimidine (C or T)

<400> SEQUENCE: 19

rrrcwgyyyy 10

<210> SEQ ID NO 20
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: mammalian p53 consensus sequence

<400> SEQUENCE: 20

gaacatgtcc caacatgttg 20

<210> SEQ ID NO 21
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: mammalian p53 consensus sequence

<400> SEQUENCE: 21

agacaagccc gggcaaggcc 20

<210> SEQ ID NO 22
<211> LENGTH: 2770
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:

-continued

<223> OTHER INFORMATION: Bacterial Cloning Vector

<400> SEQUENCE: 22

gggcgaattc gatatcgcg	ccgcgaggag tatggatccg actcgagtcg gatcctggct	60
cctcgtcgac ctgcaggcat gcaagcttga gtattctata gtgtcaccta aatagcttgg		120
cgtaatcatg gtcatactgt tttcctgtgt gaaattgtta tccgctcaca attccacaca		180
acatacgagc cggaagcata aagtgtaaag cctgggggtgc ctaatgagtg agctaactca		240
cattaattgc gttgcgctca ctgcccgtt tccagtcggg aaacctgtcg tgccagctgc		300
attaatgaat cggccaacgc gcggggagag gcggtttgcg tattggggcg tcttccgctt		360
cctcgtcac tgactcgtg cgctcggctg ttcggctgcg gcgagcggta tcagctcact		420
caaaggcggg aatacgggta tccacagaat caggggataa cgcaggaaag aacatgtgag		480
caaaaggcca gcaaaaggcc aggaaccgta aaaaggccgc gttgctggcg tttttcgata		540
ggctccgccc ccctgacgag catcacaaaa atcgacgctc aagtcagagg tggcgaaacc		600
cgacaggact ataaagatac caggcgtttc cccctggaag ctccctcgtg cgctctcctg		660
taccgacctt gccgcttacc ggatacctgt ccgcctttct cccttcggga agcgtggcgc		720
tttctcatag ctacgctgt aggtatctca gttcgggtga ggtcgttcgc tccaagctgg		780
gctgtgtgca cgaaccccc gttcagccc accgctgcgc cttatccggt aactatcgtc		840
ttgagaccaa cccggtaaag cacgacttat cgccactggc agcagccact ggtaacagga		900
ttagcagagc gaggtatgta ggcggtgcta cagagttctt gaagtgggtg cctaactacg		960
gctacactag aaggacagta tttggtatct gcgctctgct gaagccagtt accttcggaa		1020
aaagagttgg tagctcttga tccggcaaac aaaccaccgc tggtagcggg gggttttttg		1080
tttgcaagca gcagattacg cgcagaaaaa aaggatctca agaagatcct ttgatctttt		1140
ctacggggtc tgacgctcag tggaacgaaa actcacgta agggattttg gtcatgagat		1200
tatcaaaaag gatcttcacc tagatccttt taaattaaaa atgaagtttt aaatcaatct		1260
aaagtatata tgagtaaaact tggctctgaca gttaccaatg cttaatcagt gaggcaccta		1320
tctcagcgat ctgtctattt cgttcatcca tagttgcctg actccccgtc gtgtagataa		1380
ctacgatacg ggagggttta ccatctggcc ccagtgcctg aatgataccg cgagaccac		1440
gctcaccggc tccagattta tcagcaataa accagccagc cggaagggcc gagcgagaa		1500
gtggtcctgc aactttatcc gcctccatcc agtctattaa ttggtgcccg gaagctagag		1560
taagtagttc gccagttaat agtttgcgca acgttggtg cattgctaca ggcatcgtgg		1620
tgtcacgctc gtcgttttgt atggcttcat tcagctccgg tcccaacga tcaaggcgag		1680
ttacatgatc ccccatgttg tgcaaaaaag cggttagctc cttcggctct ccgatcgttg		1740
tcagaagtaa gttggccgca gtgttatcac tcatggttat ggcagcactg cataattctc		1800
ttactgtcat gccatccgta agatgctttt ctgtgactgg tgagtactca accaagtcac		1860
tctgagaata gtgtatgcgg cgaccgagtt gctcttgccc ggcgtcaata cgggataata		1920
ccgcgccaca tagcagaact ttaaaagtgc tcatcattgg aaaacgttct tcggggcgaa		1980
aactctcaag gatcttaccg ctgttgagat ccagttcgat gtaaccact cgtgcacca		2040
actgatcttc agcatctttt actttcacca gcgtttcttg gtgagcaaaa acaggaaggc		2100
aaaatgccgc aaaaaaggga ataaggcgca cacggaaatg ttgaatactc atactcttcc		2160

-continued

tttttcaata ttattgaagc atttatcagg gttattgtct catgagcgga tacatatttg	2220
aatgtattta gaaaaataaa caaatagggg ttccgcgcac atttccccga aaagtgccac	2280
ctgacgtcta agaaaccatt attatcatga cattaaccta taaaaatagg cgtatcacga	2340
ggccctttcg tctcgcgcgt ttcggtgatg acggtgaaaa cctctgacac atgcagctcc	2400
cggagacggg cacagcttgt ctgtaagcgg atgccgggag cagacaagcc cgtcagggcg	2460
cgtcagcggg tgttgggcgg tgtcggggct ggcttaacta tgcggcatca gagcagattg	2520
tactgagagt gcaccatatg cgggtgtgaaa taccgcacag atgcgtaagg agaaaatacc	2580
gcatcaggcg ccattcgcca ttcaggctgc gcaactgttg ggaagggcga tcggtgcggg	2640
cctcttcgct attacgccag ctggcgaaag ggggatgtgc tgcaaggcga ttaagttggg	2700
taacgccagg gttttcccag tcacgacgtt gtaaacgac ggccagtga ttgtaatacg	2760
actcactata	2770
 <210> SEQ ID NO 23 <211> LENGTH: 54 <212> TYPE: DNA <213> ORGANISM: Artificial <220> FEATURE: <223> OTHER INFORMATION: oligonucleotide with homology to bacterial cloning vector	
<400> SEQUENCE: 23	
gcggccgcga ggagtatgga tccgactcga gtcggatcct ggctcctcgt cgac	54
 <210> SEQ ID NO 24 <211> LENGTH: 26 <212> TYPE: DNA <213> ORGANISM: Artificial <220> FEATURE: <223> OTHER INFORMATION: oligonucleotide with homology to bacterial cloning vector	
<400> SEQUENCE: 24	
gcggccgcga ggagtatgga tccgac	26
 <210> SEQ ID NO 25 <211> LENGTH: 24 <212> TYPE: DNA <213> ORGANISM: Artificial <220> FEATURE: <223> OTHER INFORMATION: oligonucleotide with homology to bacterial cloning vector	
<400> SEQUENCE: 25	
gtcggatcct ggctcctcgt cgac	24
 <210> SEQ ID NO 26 <211> LENGTH: 10 <212> TYPE: DNA <213> ORGANISM: mammalian <220> FEATURE: <221> NAME/KEY: polyA_site <222> LOCATION: (1)..(10)	
<400> SEQUENCE: 26	
aaaaaaaaa	10
 <210> SEQ ID NO 27	

-continued

```

<211> LENGTH: 54
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide with homology to bacterial
        cloning vector

<400> SEQUENCE: 27

gcgcggcgct cctcatacct aggctgagct cagcctagga ccgaggagca gctg          54

<210> SEQ ID NO 28
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide with homology to bacterial
        cloning vector

<400> SEQUENCE: 28

cgccggcgct cctcatacct aggctg          26

<210> SEQ ID NO 29
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: oligonucleotide with homology to bacterial
        cloning vector

<400> SEQUENCE: 29

cagcctagga ccgaggagca gctg          24

```

What is claimed is:

1. An isolated oligonucleotide comprising at least one ditag, wherein the ditag includes two joined first and second sequence tags, and wherein the first tag includes a 5'-terminus sequence and a second tag comprises the 3'-terminus sequence of a nucleic acid molecule or fragment thereof.

2. The oligonucleotide of claim 1, further comprising two adapters flanking the ditag, wherein each adapter includes at least one restriction site.

3. The oligonucleotide of claim 2, wherein each adapter comprises at least a first restriction site which is an asymmetric restriction site and at least a second restriction site.

4. The oligonucleotide of claim 3, wherein the asymmetric restriction site comprises a type II restriction site.

5. The oligonucleotide of claim 1, wherein the nucleic acid molecule comprises the full-length sequence of a gene or a fragment thereof.

6. The oligonucleotide of claim 1, wherein the nucleic acid comprises RNA, mRNA, genomic DNA, full-length cDNA, or cDNA.

7. The oligonucleotide of claim 1, wherein the ditag is obtained by splicing the 5' terminus and the 3' terminus of the nucleic acid molecule or fragment thereof in the presence of at least one restriction enzyme and the size of the sequence tags is determined by the restriction enzyme used.

8. The oligonucleotide of claim 7, wherein the restriction enzyme is a type II restriction enzyme.

9. The oligonucleotide of claim 7, wherein the restriction enzyme is selected from the group consisting of AarI, AceIII, AclI, BaeI, Bbr7I, BbvI, BbvII, BccI, Bce83I,

BceAI, Bcefl, BcgI, BciVI, BfiI, BinI, BpII, BsaXI, BscAI, BseMII, BseRI, BsgI, BsmI, BsmAI, BsmFI, Bsp24I, BspCNI, BspMI, BsrI, BsrDI, BstF51, BtgZI, BtsI, CjeI, CjePT, EciI, Eco31I, Eco57I, Eco57MI, Esp3I, Fall, Faul, FokI, GsuI, HaeIV, HgaI, Hin4I, HphI, HpyAV, Ksp632I, MboII, MlyI, MmeI, MnlI, PleI, PpiI, PstI, RleAI, SapI, SfaNI, SspD51, Sth132I, StsI, TaqII, TspDTI, TspGWI, TspRI, Tth111II, I-CeuI, PI-SceI, PI-PspI and I-SceI.

10. The oligonucleotide of claim 1, wherein the ditag size is 12-60 bp.

11. The oligonucleotide of claim 1, wherein the ditag comprises 34-38 nucleotides and the size of each tag is determined by the use of restriction enzyme MmeI.

12. The oligonucleotide of claim 1, wherein the first and second tag have the same or a different number of nucleotides.

13. The oligonucleotide of claim 1, wherein the oligonucleotide consists of 1 to 1000 ditags.

14. A vector comprising an isolated oligonucleotide including at least one ditag, wherein the ditag includes two joined first and second sequence tags, and wherein the first tag includes a 5'-terminus sequence and a second tag comprises the 3'-terminus sequence of a nucleic acid molecule or fragment thereof.

15. The vector of claim 14 further comprising two adapters flanking the ditag, wherein each adapter includes at least one restriction site and wherein each adapter includes at least a first restriction site which is an asymmetric restriction site and at least a second restriction site.

16. The vector of claim 15, wherein the backbone of the vector does not comprise the asymmetric restriction site or the second restriction site.

17. The vector of claim 16, wherein the asymmetric restriction site is a type II restriction site.

18. A vector comprising at least a nucleic acid molecule and two adapters flanking the nucleic acid molecule, wherein each adapter comprises at least: a first restriction site which is a type II restriction site and at least a second restriction site, and wherein the backbone of the vector does not comprise the type II restriction site, or the second restriction site.

19. The vector of claim 18, wherein the type T1 restriction site is MmeI.

20. A vector comprising SEQ ID NO:18.

21. The vector of claim 20, comprising at least one ditag, wherein the ditag includes two joined first and second sequence tags, and wherein the first tag includes a 5'-terminus sequence and a second tag comprises the 3'-terminus sequence of a nucleic acid molecule or fragment thereof.

22. The vector of claim 21, comprising comprising two adapters flanking the ditag, wherein each adapter includes at least one restriction site and wherein each adapter includes at least a first restriction site which is an asymmetric restriction site and at least a second restriction site.

23. A cDNA library, wherein every cDNA clone of the library comprises at least one oligonucleotide including at least one ditag, wherein the ditag includes two joined first and second sequence tags, and wherein the first tag includes a 5'-terminus sequence and a second tag comprises the 3'-terminus sequence of a nucleic acid molecule or fragment thereof.

24. The cDNA library of claim 23, wherein the at least one oligonucleotide comprises 1-1000 ditags.

25. A method for recovering full-length cDNA comprising:

preparing, from a full-length cDNA library, at least one oligonucleotide including at least one ditag, the ditag including two joined first and second sequence tags, wherein the first tag includes the 5'-terminus sequence and the second tag includes the 3'-terminus sequence of a full-length cDNA;

sequencing the obtained oligonucleotide ditag;

determining the ditag of interest; and

recovering the full-length cDNA corresponding to the ditag of interest from the full-length cDNA library.

26. A method for quantifying the transcriptional activity of a gene comprising:

preparing, from a full-length cDNA library, at least one oligonucleotide comprising at least one ditag, the ditag including two joined first and second sequence tags, wherein the first tag includes the 5'-terminus sequence and the second tag includes the 3'-terminus sequence of a full-length cDNA;

sequencing the obtained oligonucleotide ditag;

determining the frequency of the sequenced ditag which corresponds to the transcriptional activity of the gene.

* * * * *