

US 20090042266A1

(19) **United States**

(12) **Patent Application Publication**
VEHMAANPERA et al.

(10) **Pub. No.: US 2009/0042266 A1**

(43) **Pub. Date: Feb. 12, 2009**

(54) **TREATMENT OF CELLULOSIC MATERIAL AND ENZYMES USEFUL THEREIN**

(60) Provisional application No. 60/753,258, filed on Dec. 22, 2005.

(75) Inventors: **Jari VEHMAANPERA**, Klaukkala (FI); **Marika Alapuranen**, Rajamaki (FI); **Terhi Puranen**, Nurmijarvi (FI); **Matti Siika-Aho**, Helsinki (FI); **Jarno Kallio**, Jarvenpaa (FI); **Satu Hooman**, Espoo (FI); **Sanni Voutilainen**, Virkkala (FI); **Teemu Halonen**, Espoo (FI); **Liisa Viikari**, Helsinki (FI)

(30) **Foreign Application Priority Data**

Dec. 22, 2005 (FI) 20051318

Publication Classification

(51) **Int. Cl.**
C12P 7/10 (2006.01)
C12N 9/42 (2006.01)
C07H 21/00 (2006.01)
C12N 15/64 (2006.01)
C12N 1/21 (2006.01)
C12N 1/20 (2006.01)

Correspondence Address:
BANNER & WITCOFF, LTD.
28 STATE STREET, 28th FLOOR
BOSTON, MA 02109-9601 (US)

(52) **U.S. Cl. 435/165; 435/209; 536/23.1; 435/320.1; 435/252.33; 435/252.8**

(73) Assignee: **Roal Oy**, Rajamaki (FI)

(57) **ABSTRACT**

(21) Appl. No.: **12/141,976**

The present invention relates to the production of sugar hydrolysates from cellulosic material. The method may be used e.g. for producing fermentable sugars for the production of bioethanol from lignocellulosic material. Cellulolytic enzymes and their production by recombinant technology is described, as well as uses of the enzymes and enzyme preparations.

(22) Filed: **Jun. 19, 2008**

Related U.S. Application Data

(63) Continuation of application No. PCT/FI2006/050558, filed on Dec. 15, 2006.

Fig. 1A.




Fig. 1B.




Fig. 1C.




Fig. 1D.




Fig. 1E.




Fig. 1F.




Fig. 2.




Fig. 3A.




Fig. 3B.




Fig. 4A.




Fig. 4B.




Fig. 5A.




Fig. 5B.




Fig. 6A.




Fig. 6B.




Fig. 7A.




Fig. 7B.




Fig. 8A.




Fig 8B.




Fig. 9.




Fig. 10.




Fig. 11.




Fig. 12.




Fig. 13.




Fig. 14.





Fig. 15.



TREATMENT OF CELLULOSIC MATERIAL AND ENZYMES USEFUL THEREIN

RELATED APPLICATIONS

[0001] This application is a continuation of PCT application no. PCT/FI2006/050558, designating the United States and filed Dec. 15, 2006; which claims the benefit of the filing date of Finnish application no. 20051318, filed Dec. 22, 2005; and U.S. application No. 60/753,258, filed Dec. 22, 2005; each of which is hereby incorporated herein by reference in its entirety for all purposes.

FIELD

[0002] The present invention relates to the production of sugar hydrolysates from cellulosic material. More precisely the invention relates to production of fermentable sugars from lignocellulosic material by enzymatic conversion. The fermentable sugars are useful e.g. in the production of bioethanol, or for other purposes. In particular the invention is directed to a method for treating cellulosic material with cellobiohydrolase, endoglucanase, beta-glucosidase, and optionally xylanase, and to enzyme preparations and the uses thereof. The invention is further directed to novel cellulolytic polypeptides, polynucleotides encoding them, and to vectors and host cells containing the polynucleotides. Still further the invention is directed to uses of the polypeptides and to a method of preparing them.

BACKGROUND

[0003] Sugar hydrolysates can be used for microbial production of a variety of fine chemicals or biopolymers, such as organic acids e.g. lactic acid, or ethanol or other alcohols e.g. n-butanol, 1,3-propanediol, or polyhydroxyalkanoates (PHAs). The sugar hydrolysates may also serve as raw material for other non-microbial processes, e.g., for enrichment, isolation and purification of high value sugars or various polymerization processes. One of the major uses of the sugar hydrolysates is in the production of biofuels. The production of bioethanol and/or other chemicals may take place in an integrated process in a biorefinery (Wyman 2001).

[0004] Limited resources of fossil fuels, and increasing amounts of CO₂ released from them and causing the greenhouse phenomenon have raised a need for using biomass as a renewable and clean source of energy. One promising, alternative technology is the production of biofuels i.e. ethanol from cellulosic materials. In the transportation sector biofuels are for the time being the only option, which could reduce the CO₂ emissions by an order of magnitude. The ethanol can be used in existing vehicles and distribution systems and thus it does not require expensive infrastructure investments. Sugars derived from lignocellulosic renewable raw materials can also be used as raw materials for a variety of chemical products that can replace oil-based chemicals.

[0005] Most of the carbohydrates in plants are in the form of lignocellulose, which essentially consists of cellulose, hemicellulose, pectin and lignin. In a lignocellulose-to-ethanol process the lignocellulosic material is first pretreated either chemically or physically to make the cellulose fraction more accessible to hydrolysis. The cellulose fraction is then hydrolysed to obtain sugars that can be fermented by yeast into ethanol. Lignin is obtained as a main co-product that may be used as a solid fuel.

[0006] Bioethanol production costs are high and the energy output is low, and there is continuous research for making the process more economical. Enzymatic hydrolysis is considered the most promising technology for converting cellulosic biomass into fermentable sugars. However, enzymatic hydrolysis is used only to a limited amount at industrial scale, and especially when using strongly lignified material such as wood or agricultural waste the technology is not satisfactory. The cost of the enzymatic step is one of the major economical factors of the process. Efforts have been made to improve the efficiency of the enzymatic hydrolysis of the cellulosic material (Badger 2002).

[0007] US 2002/0192774 A1 describes a continuous process for converting solid lignocellulosic biomass into combustible fuel products. After pretreatment by wet oxidation or steam explosion the biomass is partially separated into cellulose, hemicellulose and lignin, and is then subjected to partial hydrolysis using one or more carbohydrase enzymes (EC 3.2). Celluclast™, a commercial product by Novo Nordisk A/S containing cellulase and xylanase activities is given as an example.

[0008] US 2004/000 5674 A1 describes novel enzyme mixtures that can be used directly on lignocellulose substrate, whereby toxic waste products formed during pretreatment processes may be avoided, and energy may be saved. The synergistic enzyme mixture contains a cellulase and an auxiliary enzyme such as cellulase, xylanase, ligninase, amylase, protease, lipase or glucuronidase, or any combination thereof. Cellulase is considered to include endoglucanase (EG), beta-glucosidase (BG) and cellobiohydrolase (CBH). The examples illustrate the use of a mixture of *Trichoderma* xylanase and cellulase preparations.

[0009] Kurabi et al. (2005) have investigated enzymatic hydrolysis of steam-exploded and ethanol organosolv-pretreated Douglas-fir by novel and commercial fungal cellulases. They tested two commercial *Trichoderma reesei* cellulase preparations, and two novel preparations produced by mutant strains of *Trichoderma* sp. and *Penicillium* sp. The *Trichoderma* sp. preparation showed significantly better performance than the other preparations. The better performance was believed to be at least partly due to a significantly higher beta-glucosidase activity, which relieves product inhibition of cellobiohydrolase and endoglucanase.

[0010] US 2004/005 3373 A1 pertains a method of converting cellulose to glucose by treating a pretreated lignocellulosic substrate with an enzyme mixture comprising cellulase and a modified cellobiohydrolase I (CBHI). The CBHI has been modified by inactivating its cellulose binding domain (CBD). Advantages of CBHI modification are e.g. better recovery and higher hydrolysis rate with high substrate concentration. The cellulase is selected from the group consisting of EG, CBH and BG. The CBHI is preferably obtained from *Trichoderma*.

[0011] US 2005/016 4355 A1 describes a method for degrading lignocellulosic material with one or more cellulolytic enzymes in the presence of at least one surfactant. Additional enzymes such as hemicellulases, esterase, peroxidase, protease, laccase or mixture thereof may also be used. The presence of surfactant increases the degradation of lignocellulosic material compared to the absence of surfactant. The cellulolytic enzymes may be any enzyme involved in the degradation of lignocellulose including CBH, EG, and BG.

[0012] There is a huge number of publications disclosing various cellulases and hemicellulases.

[0013] Cellobiohydrolases (CBHs) are disclosed e.g. in WO 03/000 941, which relates to CBHI enzymes obtained from various fungi. No physiological properties of the enzymes are provided, nor any examples of their uses. Hong et al. (2003b) characterizes CBHI of *Thermoascus aurantiacus* produced in yeast. Applications of the enzyme are not described. Tuohy et al. (2002) describe three forms of cellobiohydrolases from *Talaromyces emersonii*.

[0014] Endoglucanases of the cel5 family (EGs fam 5) are described e.g. in WO 03/062 409, which relates to compositions comprising at least two thermostable enzymes for use in feed applications. Hong et al. (2003a) describe production of thermostable endo- β -1,4-glucanase from *T. aurantiacus* in yeast. No applications are explained. WO 01/70998 relates to β -glucanases from *Talaromyces*. They also describe β -glucanases from *Talaromyces emersonii*. Food, feed, beverage, brewing, and detergent applications are discussed. Lignocellulose hydrolysis is not mentioned. WO 98/06 858 describes beta-1,4-endoglucanase from *Aspergillus niger* and discusses feed and food applications of the enzyme. WO 97/13853 describes methods for screening DNA fragments encoding enzymes in cDNA libraries. The cDNA library is of yeast or fungal origin, preferably from *Aspergillus*. The enzyme is preferably a cellulase. Van Petegem et al. (2002) describe the 3D-structure of an endoglucanase of the cel5 family from *Thermoascus aurantiacus*. Parry et al. (2002) describe the mode of action of an endoglucanase of the cel5 family from *Thermoascus aurantiacus*.

[0015] Endoglucanases of the cel7 family (EGs fam 7) are disclosed e.g. in U.S. Pat. No. 5,912,157, which pertains *Myceliphthora* endoglucanase and its homologues and applications thereof in detergent, textile, and pulp. U.S. Pat. No. 6,071,735 describes cellulases exhibiting high endoglucanase activity in alkaline conditions. Uses as detergent, in pulp and paper, and textile applications are discussed. Bioethanol is not mentioned. U.S. Pat. No. 5,763,254 discloses enzymes degrading cellulose/hemicellulose and having conserved amino acid residues in CBD.

[0016] Endoglucanases of the cel45 family (EGs fam 45) are described e.g. in U.S. Pat. No. 6,001,639, which relates to enzymes having endoglucanase activity and having two conserved amino acid sequences. Uses in textile, detergent, and pulp and paper applications are generally discussed and treating of lignocellulosic material is mentioned but no examples are given. WO 2004/053039 is directed to detergent applications of endoglucanases. U.S. Pat. No. 5,958,082 discloses the use of endoglucanase, especially from *Thielavia terrestris* in textile application. EP 0495258 relates to detergent compositions containing *Humicola* cellulase. U.S. Pat. No. 5,948,672 describes a cellulase preparation containing endoglucanase, especially from *Humicola* and its use in textile and pulp applications. Lignocellulose hydrolysis is not mentioned.

[0017] A small amount of beta-glucosidase (BG) enhances hydrolysis of biomass to glucose by hydrolyzing cellobiose produced by cellobiohydrolases. Cellobiose conversion to glucose is usually the major rate-limiting step. Beta-glucosidases are disclosed e.g. in US 2005/021 4920, which relates to BG from *Aspergillus fumigatus*. The enzyme has been produced in *Aspergillus oryzae* and *Trichoderma reesei*. Use of the enzyme in degradation of biomass or detergent applications is generally discussed but not exemplified. WO02/095 014 describes an *Aspergillus oryzae* enzyme having cellobiase activity. Use in the production of ethanol from biomass is generally discussed but not exemplified. WO2005/074656

discloses polypeptides having cellulolytic enhancing activity derived e.g. from *T. aurantiacus*; *A. fumigatus*; *T. terrestris* and *T. aurantiacus*. WO02/26979 discloses enzymatic processing of plant material. U.S. Pat. No. 6,022,725 describes cloning and amplification of the beta-glucosidase gene of *Trichoderma reesei*, and U.S. Pat. No. 6,103,464 describes a method for detecting DNA encoding a beta-glucosidase from a filamentous fungus. No application examples are given.

[0018] Xylanases are described e.g. in FR2786784, which relates to a heat-stable xylanase, useful e.g. in treating animal feed and in bread making. The enzyme is derived from a thermophilic fungus, particularly of the genus *Thermoascus*.

[0019] U.S. Pat. No. 6,197,564 describes enzymes having xylanase activity, and obtained from *Aspergillus aculeatus*. Their application in baking is exemplified. WO 02/24926 relates to *Talaromyces xylanases*. Feed and baking examples are given. WO01/42433 discloses thermostable xylanase from *Talaromyces emersonii* for use in food and feed applications.

[0020] The best-investigated and most widely applied cellulolytic enzymes of fungal origin have been derived from *Trichoderma reesei* (the anamorph of *Hypocrea jecorina*). Consequently also most of the commercially available fungal cellulases are derived from *Trichoderma reesei*. However, the majority of cellulases from less known fungi have not been applied in processes of practical importance such as in degrading cellulosic material, including lignocellulose.

[0021] There is a continuous need for new methods of degrading cellulosic substrates, in particular lignocellulosic substrates, and for new enzymes and enzyme mixtures, which enhance the efficiency of the degradation. There is also a need for processes and enzymes, which work at high temperatures, thus enabling the use of high biomass consistency and leading to high sugar and ethanol concentrations. This approach may lead to significant saving in energy and investments costs. The high temperature also decreases the risk of contamination during hydrolysis. The present invention aims to meet at least part of these needs.

BRIEF DESCRIPTION

[0022] It has now surprisingly been found that cellulolytic enzymes, and especially cellobiohydrolases obtainable from *Thermoascus aurantiacus*, *Acremonium thermophilum*, or *Chaetomium thermophilum* are particularly useful in hydrolyzing cellulosic material. In addition to cellobiohydrolases these fungi also have endoglucanases, beta-glucosidases and xylanases that are very suitable for degrading cellulosic material. The enzymes are kinetically very effective over a broad range of temperatures, and although they have high activity at high temperatures, they are also very efficient at standard hydrolysis temperatures. This makes them extremely well suited for varying cellulosic substrate hydrolysis processes carried out both at conventional temperatures and at elevated temperatures.

[0023] The present invention provides a method for treating cellulosic material with cellobiohydrolase, endoglucanase and beta-glucosidase, whereby said cellobiohydrolase comprises an amino acid sequence having at least 80% identity to SEQ ID NO: 2, 4, 6 or 8, or to an enzymatically active fragment thereof.

[0024] The invention further provides an enzyme preparation comprising cellobiohydrolase, endoglucanase and beta-glucosidase, wherein said cellobiohydrolase comprises an

amino acid sequence having at least 80% identity to SEQ ID NO: 2, 4, 6 or 8, or to an enzymatically active fragment thereof.

[0025] The use of said enzyme preparation for degrading cellulosic material is also provided, as well as the use of said method in a process for preparing ethanol from cellulosic material.

[0026] The invention is also directed to a polypeptide comprising a fragment having cellulolytic activity and being selected from the group consisting of:

[0027] a) a polypeptide comprising an amino acid sequence having at least 66% identity to SEQ ID NO:4, 79% identity to SEQ ID NO:6, 78% identity to SEQ ID NO:12, 68% identity to SEQ ID NO:14, 72% identity to SEQ ID NO:16, 68% identity to SEQ ID NO:20, 74% identity to SEQ ID NO:22 or 24, or 78% identity to SEQ ID NO:26;

[0028] b) a variant of a) comprising a fragment having cellulolytic activity; and

[0029] c) a fragment of a) or b) having cellulolytic activity.

[0030] One further object of the invention is an isolated polynucleotide selected from the group consisting of:

[0031] a) a nucleotide sequence of SEQ ID NO: 3, 5, 11, 13, 15, 19, 21, 23 or 25, or a sequence encoding a polypeptide of claim 35;

[0032] b) a complementary strand of a)

[0033] c) a fragment of a) or b) comprising at least 20 nucleotides; and

[0034] d) a sequence that is degenerate as a result of the genetic code to any one of the sequences as defined in a), b) or c).

[0035] The invention still further provides a vector, which comprises said polynucleotide as a heterologous sequence, and a host cell comprising said vector. *Escherichia coli* strains having accession number DSM 16728, DSM 16729, DSM 17324, DSM 17323, DSM 17729, DSM 16726, DSM 16725, DSM 17325 or DSM 17667 are also included in the invention.

[0036] Other objects of the invention are enzyme preparations comprising at least one of the novel polypeptides, and the use of said polypeptide or enzyme preparation in fuel, textile, detergent, pulp and paper, food, feed or beverage industry.

[0037] Further provided is a method for preparing a polypeptide comprising a fragment having cellulolytic activity and being selected from the group consisting of:

[0038] a) a polypeptide comprising an amino acid sequence having at least 66% identity to SEQ ID NO:4, 79% identity to SEQ ID NO:6, 78% identity to SEQ ID NO:12, 68% identity to SEQ ID NO:14, 72% identity to SEQ ID NO:16, 68% identity to SEQ ID NO:20, 74% identity to SEQ ID NO:22 or 24, or 78% identity to SEQ ID NO:26;

[0039] b) a variant of a) comprising a fragment having cellulolytic activity; and

[0040] c) a fragment of a) or b) having cellulolytic activity,

[0041] said method comprising transforming a host cell with a vector encoding said polypeptide, and culturing said host cell under conditions enabling expression of said polypeptide, and optionally recovering and purifying the polypeptide produced.

[0042] Still further provided is a method of treating cellulosic material with a spent culture medium of at least one microorganism capable of producing a polypeptide as defined above, wherein the method comprises reacting the cellulosic material with the spent culture medium to obtain hydrolysed cellulosic material.

[0043] Specific embodiments of the invention are set forth in the dependent claims.

[0044] Other objects, details and advantages of the present invention will become apparent from the following drawings, detailed description and examples.

BRIEF DESCRIPTION OF THE DRAWINGS

[0045] FIG. 1. Temperature dependencies of the cellulase and beta-glucosidase activities in the supernatants of the tested six fungal strains. The incubation time in the assay was 60 min at the given temperature, the assay pH was 5.0 (MUL-activity) or 4.8 (CMCase or BGU). Activity obtained at 60° C. is set as the relative activity of 100%. A) *Thermoascus aurantiacus* ALKO4239, B) *Thermoascus aurantiacus* ALKO4242, C) *Acremonium thermophilum* ALKO4245, D) *Talaromyces thermophilus* ALKO4246, E) *Chaetomium thermophilum* ALKO4261, F) *Chaetomium thermophilum* ALKO4265.

[0046] FIG. 2. Schematic picture of the expression cassettes used in the transformation of *Trichoderma reesei* protoplasts for producing the recombinant fungal proteins. The recombinant genes were under the control of *T. reesei* cbh1 (cel7A) promoter (cbh1 prom) and the termination of the transcription was ensured by using *T. reesei* cbh1 terminator sequence (cbh1 term). The amdS gene was included as a transformation marker.

[0047] FIG. 3. A) pH optima of the recombinant CBH/Cel7 protein preparations from *Thermoascus aurantiacus* ALKO4242, *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 determined on 4-methylumbelliferyl- β -D-lactoside (MUL) at 50° C., 10 min. The results are given as mean (\pm SD) of three separate measurements. B) Thermal stability of recombinant CBH/Cel7 protein preparations from *Thermoascus aurantiacus* ALKO4242, *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 determined on 4-methylumbelliferyl- β -D-lactoside (MUL) at the optimum pH for 60 min. The results are given as mean (\pm SD) of three separate measurements. Both reactions contained BSA (100 μ g/ml) as a stabilizer.

[0048] FIG. 4. Crystalline cellulose (Avicel) hydrolysis by the purified recombinant cellobiohydrolases at 45° C. Substrate concentration 1% (w/v), pH 5.0, enzyme concentration 1.4 μ M. A) Cellobiohydrolases harboring a CBD, B) cellobiohydrolases (core) without a CBD.

[0049] FIG. 5. Crystalline cellulose (Avicel) hydrolysis by the purified recombinant cellobiohydrolases at 70° C. Substrate concentration 1% (w/v), pH 5.0, enzyme concentration 1.4 μ M. A) Cellobiohydrolases harboring a CBD, B) cellobiohydrolases (core) without a CBD.

[0050] FIG. 6. A) The pH dependency of the heterologously produced *Acremonium* EG_40/Cel45A, EG_40_like/Cel45B and *Thermoascus* EG_28/Cel5A activity was determined with CMC substrate in a 10 min reaction at 50° C. B) Temperature optimum of the *Acremonium* EG_40/Cel45A, EG_40_like/Cel45B and *Thermoascus* EG_28/Cel5A was determined at pH 5.5, 4.8, and 6.0, respectively. The reaction containing CMC as substrate was performed for 60 min, except for EG_28/Cel5A for 10 min. BSA (100 μ g/ml) was added as a stabilizer.

[0051] FIG. 7. A) The pH dependency of the heterologously produced *Acremonium* BG_101/Cel3A, *Chaetomium* BG_76/Cel3A, and *Thermoascus* BG_81/Cel3A activity was determined with 4-nitrophenyl- β -D-glucopyranoside

substrate in a 10 min reaction at 50° C. B) Temperature optimum of the *Acremonium* β G_101/Cel3A, *Chaetomium* β G_76/Cel3A, and *Thermoascus* β G_81/Cel3A was determined at pH 4.5, 5.5, and 4.5, respectively. The reaction containing 4-nitrophenyl- β -D-glucopyranosid as substrate was performed for 60 min, BSA (100 μ g/ml) was added as a stabilizer.

[0052] FIG. 8. A) The pH dependency of the heterologously produced *Thermoascus* XYN_30/Xyn10A xylanase activity was determined with birch xylan substrate in a 10 min reaction at 50° C. B) Temperature optimum of XYN_30/Xyn10A was determined at pH 5.3 in a 60 min reaction, BSA (100 μ g/ml) was added as a stabilizer.

[0053] FIG. 9. Hydrolysis of washed steam exploded spruce fibre (10 mg/ml) with a mixture of thermophilic enzymes (MIXTURE 1) and *T. reesei* enzymes at 55 and 60° C. Enzyme dosage is given by FPU/g dry matter of substrate, FPU assayed at 50° C., pH 5. Hydrolysis was carried out for 72 h at pH 5, with mixing. The results are given as mean (\pm SD) of three separate measurements.

[0054] FIG. 10. Hydrolysis of steam exploded corn stover (10 mg/ml) with a mixture of thermophilic enzymes (MIXTURE 2) and *T. reesei* enzymes at 45, 55 and 57.5° C. Enzyme dosage was for "MIXTURE 2" 5 FPU/g dry matter of substrate and for *T. reesei* enzymes 5 FPU/g dry matter Celluclast supplemented with 100 nkat/g dry matter Novozym 188 (filter paper activity was assayed at 50° C., pH 5). Hydrolysis was carried out for 72 h at pH 5, with mixing. The results are given as mean (\pm SD) of three separate measurements. The substrate contained soluble reducing sugars (ca 0.7 mg/ml). This background sugar content was subtracted from the reducing sugars formed during the hydrolysis.

[0055] FIG. 11. Hydrolysis of steam exploded corn stover (10 mg/ml) with a mixture of thermophilic enzymes containing a new thermophilic xylanase from *Thermoascus aurantiacus* (MIXTURE 3) and *T. reesei* enzymes at 45, 55 and 60° C. Enzyme dosage was for "MIXTURE 3" 5 FPU/g dry matter of substrate and for *T. reesei* enzymes 5 FPU/g dry matter Celluclast supplemented with 100 nkat/g dry matter Novozym 188 (filter paper activity was assayed at 50° C., pH 5). Hydrolysis was carried out for 72 h at pH 5, with mixing. The results are given as mean (\pm SD) of three separate measurements. The substrate contained soluble reducing sugars (ca 0.7 mg/ml). This background sugar content was subtracted from the reducing sugars formed during the hydrolysis.

[0056] FIG. 12. Hydrolysis of steam exploded spruce fibre (10 mg/ml) with a mixture of thermophilic enzymes containing a new thermophilic xylanase XYN_30/Xyn10A from *Thermoascus aurantiacus* (MIXTURE 3) and *T. reesei* enzymes at 45, 55 and 60° C. Enzyme dosage for "MIXTURE 3" was 5 FPU/g dry matter of substrate and for *T. reesei* enzymes 5 FPU/g dry matter Celluclast supplemented with 100 nkat/g dry matter Novozym 188 (filter paper activity was assayed at 50° C., pH 5). Hydrolysis was carried out for 72 h at pH 5, with mixing. The results are given as mean (\pm SD) of three separate measurements.

[0057] FIG. 13. The effect of glucose on activity of different β -glucosidase preparations. The standard assay using p-nitrophenyl- β -D-glucopyranoside as substrate was carried out in the presence of glucose in the assay mixture. The activity is presented as percentage of the activity obtained without glucose.

[0058] FIG. 14. FPU activities of the enzyme mixtures at temperatures from 50° C. to 70° C., presented as a percentage of the activity under the standard conditions (50° C., 1 h).

[0059] FIG. 15. The relative cellulase activity of two different *T. reesei* strains grown in media containing untreated Nutriose (NO) or BG_81/Cel3A pretreated Nutriose (NBG81) as a carbon source.

DETAILED DESCRIPTION

[0060] Cellulose is the major structural component of higher plants. It provides plant cells with high tensile strength helping them to resist mechanical stress and osmotic pressure. Cellulose is a β -1,4-glucan composed of linear chains of glucose residues joined by β -1,4-glycosidic linkages. Cellobiose is the smallest repeating unit of cellulose. In cell walls cellulose is packed in variously oriented sheets, which are embedded in a matrix of hemicellulose and lignin. Hemicellulose is a heterogeneous group of carbohydrate polymers containing mainly different glucans, xylans and mannans. Hemicellulose consists of a linear backbone with β -1,4-linked residues substituted with short side chains usually containing acetyl, glucuronyl, arabinosyl and galactosyl. Hemicellulose can be chemically cross-linked to lignin. Lignin is a complex cross-linked polymer of variously substituted p-hydroxyphenylpropane units that provides strength to the cell wall to withstand mechanical stress, and it also protects cellulose from enzymatic hydrolysis.

[0061] Lignocellulose is a combination of cellulose and hemicellulose and polymers of phenol propanol units and lignin. It is physically hard, dense, and inaccessible and the most abundant biochemical material in the biosphere. Lignocellulose containing materials are for example: hardwood and softwood chips, wood pulp, sawdust and forestry and wood industrial waste; agricultural biomass as cereal straws, sugar beet pulp, corn stover and cobs, sugar cane bagasse, stems, leaves, hulls, husks, and the like; waste products as municipal solid waste, newspaper and waste office paper, milling waste of e.g. grains; dedicated energy crops (e.g., willow, poplar, switchgrass or reed canarygrass, and the like). Preferred examples are corn stover, switchgrass, cereal straw, sugar-cane bagasse and wood derived materials.

[0062] "Cellulosic material" as used herein, relates to any material comprising cellulose, hemicellulose and/or lignocellulose as a significant component. "Lignocellulosic material" means any material comprising lignocellulose. Such materials are e.g. plant materials such as wood including softwood and hardwood, herbaceous crops, agricultural residues, pulp and paper residues, waste paper, wastes of food and feed industry etc. Textile fibres such as cotton, fibres derived from cotton, linen, hemp, jute and man made cellulosic fibres as modal, viscose, lyocel are specific examples of cellulosic materials.

[0063] Cellulosic material is degraded in nature by a number of various organisms including bacteria and fungi. Cellulose is typically degraded by different cellulases acting sequentially or simultaneously. The biological conversion of cellulose to glucose generally requires three types of hydrolytic enzymes: (1) Endoglucanases which cut internal β -1,4-glycosidic bonds; (2) Exocellobiohydrolases that cut the disaccharide cellobiose from the end of the cellulose polymer chain; (3) β -1,4-glucosidases which hydrolyze the cellobiose and other short cello-oligosaccharides to glucose.

In other words the three major groups of cellulases are cellobiohydrolases (CBH), endoglucanases (EG) and beta-glucosidases (BG).

[0064] Degradation of more complex cellulose containing substrates requires a broad range of various enzymes. For example lignocellulose is degraded by hemicellulases, like xylanases and mannanases. Hemicellulase is an enzyme hydrolysing hemicellulose.

[0065] “Cellulolytic enzymes” are enzymes having “cellulolytic activity,” which means that they are capable of hydrolysing cellulosic substrates or derivatives thereof into smaller saccharides. Cellulolytic enzymes thus include both cellulases and hemicellulases. Cellulases as used herein include cellobiohydrolase, endoglucanase and beta-glucosidase.

[0066] *T. reesei* has a well known and effective cellulase system containing two CBHs, two major and several minor EGs and BGs. *T. reesei* CBHI (Cel7A) cuts sugar from the reducing end of the cellulose chain, has a C-terminal cellulose binding domain (CBD) and may constitute up to 60% of the total secreted protein. *T. reesei* CBHII (Cel6A) cuts sugar from the non-reducing end of the cellulose chain, has an N-terminal cellulose binding domain and may constitute up to 20% of the total secreted protein. Endoglucanases EGI (Cel7B), and EGV (Cel45A) have a CBD in their C-terminus, EGII (Cel5A) has an N-terminal CBD and EGIII (Cel12A) does not have a cellulose binding domain at all. CBHI, CBHII, EGI and EGII are so called “major cellulases” of *Trichoderma* comprising together 80-90% of total secreted proteins. It is known to a man skilled in the art that an enzyme may be active on several substrates and enzymatic activities can be measured using different substrates, methods and conditions. Identifying different cellulolytic activities is discussed for example in van Tilbeurgh et al. 1988.

[0067] In addition to a catalytic domain/core expressing cellulolytic activity cellulolytic enzymes may comprise one or more cellulose binding domains (CBDs), also named as carbohydrate binding domains/modules (CBD/CBM), which can be located either at the N- or C-terminus of the catalytic domain. CBDs have carbohydrate-binding activity and they mediate the binding of the cellulase to crystalline cellulose but have little or no effect on cellulase hydrolytic activity of the enzyme on soluble substrates. These two domains are typically connected via a flexible and highly glycosylated linker region.

[0068] “Cellobiohydrolase” or “CBH” as used herein refers to enzymes that cleave cellulose from the end of the glucose chain and produce mainly cellobiose. They are also called 1,4-beta-D-glucan cellobiohydrolases or cellulose 1,4-beta-cellobiosidases. They hydrolyze the 1,4-beta-D-glucosidic linkages from the reducing or non-reducing ends of a polymer containing said linkages, such as cellulose, whereby cellobiose is released. Two different CBHs have been isolated from *Trichoderma reesei*, CBHI and CBHII. They have a modular structure consisting of a catalytic domain linked to a cellulose-binding domain (CBD). There are also cellobiohydrolases in nature that lack CBD.

[0069] “Endoglucanase” or “EG” refers to enzymes that cut internal glycosidic bonds of the cellulose chain. They are classified as EC 3.2.1.4. They are 1,4-beta-D-glucan 4-glucanohydrolases and catalyze endohydrolysis of 1,4-beta-D-glycosidic linkages in polymers of glucose such as cellulose and derivatives thereof. Some naturally occurring endoglucanases

have a cellulose binding domain, while others do not. Some endoglucanases have also xylanase activity (Bailey et al., 1993).

[0070] “Beta-glucosidase” or “BG” or “G” refers to enzymes that degrade small soluble oligosaccharides including cellobiose to glucose. They are classified as EC 3.2.1.21. They are beta-D-glucoside glucohydrolases, which typically catalyze the hydrolysis of terminal non-reducing beta-D-glucose residues. These enzymes recognize oligosaccharides of glucose. Typical substrates are cellobiose and cellotriose. Cellobiose is an inhibitor of cellobiohydrolases, wherefore the degradation of cellobiose is important to overcome end-product inhibition of cellobiohydrolases.

[0071] Xylanases are enzymes that are capable of recognizing and hydrolyzing hemicellulose. They include both exohydrolytic and endohydrolytic enzymes. Typically they have endo-1,4-beta-xylanase (EC 3.2.1.8) or beta-D-xylosidase (EC 3.2.1.37) activity that breaks down hemicellulose to xylose. “Xylanase” or “Xyn” in connection with the present invention refers especially to an enzyme classified as EC 3.2.1.8 hydrolyzing xylose polymers of lignocellulosic substrate or purified xylan.

[0072] In addition to this cellulases can be classified to various glycosyl hydrolase families according their primary sequence, supported by analysis of the three dimensional structure of some members of the family (Henrissat 1991, Henrissat and Bairoch 1993, 1996). Some glycosyl hydrolases are multifunctional enzymes that contain catalytic domains that belong to different glycosylhydrolase families. Family 3 consists of beta-glucosidases (EC 3.2.1.21) such as Ta BG_81, At BG_101 and Ct BG_76 described herein. Family 5 (formerly known as celA) consists mainly of endoglucanases (EC 3.2.1.4) such as Ta EG_28 described herein. Family 7 (formerly cellulase family celC) contains endoglucanases (EC 3.2.1.4) and cellobiohydrolases (EC 3.2.1.91) such as Ct EG_54, Ta CBH, At CBH_A, At CBH_C and Ct CBH described herein. Family 10 (formerly celF) consists mainly of xylanases (EC 3.2.1.8) such as Ta XYN_30 and At XYN_60 described herein. Family 45 (formerly celK) contains endoglucanases (EC 3.2.1.4) such as At EG_40 and At EG_40_like described herein.

[0073] Cellulolytic enzymes useful for hydrolyzing cellulosic material are obtainable from *Thermoascus aurantiacus*, *Acremonium thermophilum*, or *Chaetomium thermophilum*. “Obtainable from” means that they can be obtained from said species, but it does not exclude the possibility of obtaining them from other sources. In other words they may originate from any organism including plants. Preferably they originate from microorganisms e.g. bacteria or fungi. The bacteria may be for example from a genus selected from *Bacillus*, *Azospirillum* and *Streptomyces*. More preferably the enzyme originates from fungi (including filamentous fungi and yeasts), for example from a genus selected from the group consisting of *Thermoascus*, *Acremonium*, *Chaetomium*, *Achaetomium*, *Thielavia*, *Aspergillus*, *Botrytis*, *Chrysosporium*, *Collybia*, *Fomes*, *Fusarium*, *Humicola*, *Hypocrea*, *Lentinus*, *Melanocarpus*, *Myceliophthora*, *Myriococcum*, *Neurospora*, *Penicillium*, *Phanerochaete*, *Phlebia*, *Pleurotus*, *Podospora*, *Polyporus*, *Rhizoctonia*, *Scytalidium*, *Pycnoporus*, *Trametes* and *Trichoderma*.

[0074] According to a preferred embodiment of the invention the enzymes are obtainable from *Thermoascus aurantiacus* strain ALKO4242 deposited as CBS 116239, strain ALKO4245 deposited as CBS 116240 presently classified as

Acremonium thermophilum, or *Chaetomium thermophilum* strain ALKO4265 deposited as CBS 730.95.

[0075] The cellobiohydrolase preferably comprises an amino acid sequence having at least 80% identity to SEQ ID NO: 2, 4, 6 or 8, or an enzymatically active fragment thereof.

Cellobio- hydrolase	Gene	Obtainable from	CBD	nucleic acid	amino acid
				SEQ ID NO:	SEQ ID NO:
Ta CBH	Ta cel7A	<i>T. aurantiacus</i>	-	1	2
At CBH_A	At cel7B	<i>A. thermophilum</i>	-	3	4
At CBH_C	At cel7A	<i>A. thermophilum</i>	+	5	6
Ct CBH	Ct cel7A	<i>C. thermophilum</i>	+	7	8

[0076] These CBHs have an advantageous cellulose inhibition constant compared to that of *Trichoderma reesei* CBH, and they show improved hydrolysis results when testing various cellulosic substrates. SEQ ID NO: 2 and 4 do not comprise a CBD. Particularly enhanced hydrolysis results may be obtained when a cellulose binding domain (CBD) is attached to a CBH that has no CBD of its own. The CBD may be obtained e.g. from a *Trichoderma* or *Chaetomium* species, and it is preferably attached to the CBH via a linker. The resulting fusion protein containing a CBH core region attached to a CBD via a linker may comprise an amino acid sequence having at least 80% identity to SEQ ID NO: 28 or 30. Polynucleotides comprising a sequence of SEQ ID NO: 27 or 29 encode such fusion proteins.

[0077] The endoglucanase may comprise an amino acid sequence having at least 80% identity to SEQ ID NO: 10, 12, 14 or 16, or an enzymatically active fragment thereof. These endoglucanases have good thermostability.

Endo- glucanase	Gene	Obtainable from	CBD	nucl. acid	amino acid
				SEQ ID NO:	SEQ ID NO:
Ta EG_28	Ta cel5A	<i>T. aurantiacus</i>	-	9	10
At EG_40	At cel45A	<i>A. thermophilum</i>	+	11	12
At EG40_like	At cel45B	<i>A. thermophilum</i>	-	13	14
Ct EG_54	Ct cel7B	<i>C. thermophilum</i>	+	15	16

[0078] The beta-glucosidase may comprise an amino acid sequence having at least 80% identity to SEQ ID NO: 22, 24 or 26, or an enzymatically active fragment thereof. These beta-glucosidases have good resistance to glucose inhibition, which is advantageous to avoid end product inhibition during enzymatic hydrolysis of cellulosic material. The beta-glucosidases may also be used in preparing sophorose, a cellulase inducer used in cultivation of *T. reesei*.

Beta- glucosidase	Gene	Obtainable from	nucleic acid	amino acid
			SEQ ID NO:	SEQ ID NO:
Ta BG_81	Ta cel3A	<i>T. aurantiacus</i>	21	22
At BG_101	At cel3A	<i>A. thermophilum</i>	23	24
Ct BG_76	Ct cel3A	<i>C. thermophilum</i>	25	26

[0079] The xylanase may comprise an amino acid sequence having at least 80% identity to SEQ ID NO: 18 or 20, or an enzymatically active fragment thereof.

Xylanase	Gene	Obtainable from	CBD	nucleic acid	amino acid
				SEQ ID NO:	SEQ ID NO:
Xyn_30	Ta xyn10A	<i>T. aurantiacus</i>	+	17	18
Xyn_60	At xyn10A	<i>A. thermophilum</i>	-	19	20

[0080] By the term “identity” is here meant the global identity between two amino acid sequences compared to each other from the first amino acid encoded by the corresponding gene to the last amino acid. The identity of the full-length sequences is measured by using Needleman-Wunsch global alignment program at EMBOSS (European Molecular Biology Open Software Suite; Rice et al., 2000) program package, version 3.0.0, with the following parameters: EMBL-SUM62, Gap penalty 10.0, Extend penalty 0.5. The algorithm is described in Needleman and Wunsch (1970). The man skilled in the art is aware of the fact that results using Needleman-Wunsch algorithm are comparable only when aligning corresponding domains of the sequence. Consequently comparison of e.g. cellulase sequences including CBD or signal sequences with sequences lacking those elements cannot be done.

[0081] According to one embodiment of the invention, a cellulolytic polypeptide is used that has at least 80, 85, 90, 95 or 99% identity to SEQ ID NO: 2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24 or 26 or at least to its enzymatically active fragment.

[0082] By the term “enzymatically active fragment” is meant any fragment of a defined sequence that has cellulolytic activity. In other words an enzymatically active fragment may be the mature protein part of the defined sequence, or it may be only an fragment of the mature protein part, provided that it still has cellobiohydrolase, endoglucanase, beta-glucosidase or xylanase activity.

[0083] The cellulolytic enzymes are preferably recombinant enzymes, which may be produced in a generally known manner. A polynucleotide fragment comprising the enzyme gene is isolated, the gene is inserted under a strong promoter in an expression vector, the vector is transferred into suitable host cells and the host cells are cultivated under conditions provoking production of the enzyme. Methods for protein production by recombinant technology in different host systems are well known in the art (Sambrook et al., 1989; Coen, 2001; Gellissen, 2005). Preferably the enzymes are produced as extracellular enzymes that are secreted into the culture medium, from which they can easily be recovered and isolated. The spent culture medium of the production host can be used as such, or the host cells may be removed therefrom, and/or it may be concentrated, filtrated or fractionated. It may also be dried.

[0084] Isolated polypeptide in the present context may simply mean that the cells and cell debris have been removed from the culture medium containing the polypeptide. Conveniently the polypeptides are isolated e.g. by adding anionic and/or cationic polymers to the spent culture medium to enhance precipitation of cells, cell debris and some enzymes that have unwanted side activities. The medium is then fil-

trated using an inorganic filtering agent and a filter to remove the precipitants formed. After this the filtrate is further processed using a semi-permeable membrane to remove excess of salts, sugars and metabolic products.

[0085] According to one embodiment of the invention, the heterologous polynucleotide comprises a gene similar to that included in a microorganism having accession number DSM 16723, DSM 16728, DSM 16729, DSM 16727, DSM 17326, DSM 17324, DSM 17323, DSM 17729, DSM 16724, DSM 16726, DSM 16725, DSM 17325 or DSM 17667.

[0086] The production host can be any organism capable of expressing the cellulolytic enzyme. Preferably the host is a microbial cell, more preferably a fungus. Most preferably the host is a filamentous fungus. Preferably the recombinant host is modified to express and secrete cellulolytic enzymes as its main activity or one of its main activities. This can be done by deleting major homologous secreted genes e.g. the four major cellulases of *Trichoderma* and by targeting heterologous genes to a locus that has been modified to ensure high expression and production levels. Preferred hosts for producing the cellulolytic enzymes are in particular strains from the genus *Trichoderma* or *Aspergillus*.

[0087] The enzymes needed for the hydrolysis of the cellulosic material according to the invention may be added in an enzymatically effective amount either simultaneously e.g. in the form of an enzyme mixture, or sequentially, or as a part of the simultaneous saccharification and fermentation (SSF). Any combination of the cellobiohydrolases comprising an amino acid sequence having at least 80% identity to SEQ ID NO: 2, 4, 6 or 8 or to an enzymatically active fragment thereof may be used together with any combination of endoglucanases and beta-glucosidases. If the cellulosic material comprises hemicellulose, hemicellulases, preferably xylanases are additionally used for the degradation. The endoglucanases, beta-glucosidases and xylanases may be selected from those described herein, but are not limited to them. They can for example also be commercially available enzyme preparations. In addition to cellulases and optional hemicellulases one or more other enzymes may be used, for example proteases, amylases, laccases, lipases, pectinases, esterases and/or peroxidases. Another enzyme treatment may be carried out before, during or after the cellulase treatment.

[0088] The term “enzyme preparation” denotes to a composition comprising at least one of the desired enzymes. The preparation may contain the enzymes in at least partially purified and isolated form. It may even essentially consist of the desired enzyme or enzymes. Alternatively the preparation may be a spent culture medium or filtrate containing one or more cellulolytic enzymes. In addition to the cellulolytic activity, the preparation may contain additives, such as mediators, stabilizers, buffers, preservatives, surfactants and/or culture medium components. Preferred additives are such, which are commonly used in enzyme preparations intended for a particular application. The enzyme preparation may be in the form of liquid, powder or granulate. Preferably the enzyme preparation is spent culture medium. “Spent culture medium” refers to the culture medium of the host comprising the produced enzymes. Preferably the host cells are separated from the said medium after the production.

[0089] According to one embodiment of the invention the enzyme preparation comprises a mixture of CBH, EG and BG, optionally together with xylanase and/or other enzymes. The CBH comprises an amino acid sequence having at least 80% identity to SEQ ID NO: 2, 4, 6 or 8 or to an enzymatically

active fragment thereof, and it may be obtained from *Thermoascus aurantiacus*, *Acremonium thermophilum*, or *Chaetomium thermophilum*, whereas EG, BG and xylanase may be of any origin including from said organisms. Other enzymes that might be present in the preparation are e.g. proteases, amylases, laccases, lipases, pectinases, esterases and/or peroxidases.

[0090] Different enzyme mixtures and combinations may be used to suit different process conditions. For example if the degradation process is to be carried out at a high temperature, thermostable enzymes are chosen. A combination of a CBH of family 7 with an endoglucanase of family 45, optionally in combination with a BG of family 3 and/or a xylanase of family 10 had excellent hydrolysis performance both at 45° C., and at elevated temperatures.

[0091] Cellulolytic enzymes of *Trichoderma reesei* are conventionally used at temperatures in the range of about 40-50° C. in the hydrolysis, and at 30-40° C. in SSF. CBH, EG, BG and Xyn obtainable from *Thermoascus aurantiacus*, *Acremonium thermophilum*, or *Chaetomium thermophilum* are efficient at these temperatures too, but in addition most of them also function extremely well at temperatures between 50° C. and 75° C., or even up to 80° C. and 85° C., such as between 55° C. and 70° C., e.g. between 60° C. and 65° C. For short incubation times enzyme mixtures are functional up to even 85° C., for complete hydrolysis lower temperatures are normally used.

[0092] The method for treating cellulosic material with CBH, EG, BG and Xyn is especially suitable for producing fermentable sugars from lignocellulosic material. The fermentable sugars may then be fermented by yeast into ethanol, and used as fuel. They can also be used as intermediates or raw materials for the production of various chemicals or building blocks for the processes of chemical industry, e.g. in so called biorefinery. The lignocellulosic material may be pretreated before the enzymatic hydrolysis to disrupt the fiber structure of cellulosic substrates and make the cellulose fraction more accessible to the cellulolytic enzymes. Current pretreatments include mechanical, chemical or thermal processes and combinations thereof. The material may for example be pretreated by steam explosion or acid hydrolysis.

[0093] A number of novel cellulolytic polypeptides were found in *Thermoascus aurantiacus*, *Acremonium thermophilum*, and *Chaetomium thermophilum*. The novel polypeptides may comprise a fragment having cellulolytic activity and be selected from the group consisting of a polypeptide comprising an amino acid sequence having at least 66%, preferably 70% or 75%, identity to SEQ ID NO: 4, 79% identity to SEQ ID NO: 6, 78% identity to SEQ ID NO: 12, 68%, preferably 70% or 75%, identity to SEQ ID NO: 14, 72%, preferably 75%, identity to SEQ ID NO: 16, 68%, preferably 70% or 75%, identity to SEQ ID NO: 20, 74% identity to SEQ ID NO: 22 or 24, or 78% identity to SEQ ID NO: 26.

[0094] The novel polypeptides may also be variants of said polypeptides. A “variant” may be a polypeptide that occurs naturally e.g. as an allelic variant within the same strain, species or genus, or it may have been generated by mutagenesis. It may comprise amino acid substitutions, deletions or insertions, but it still functions in a substantially similar manner to the enzymes defined above i.e. it comprises a fragment having cellulolytic activity.

[0095] The cellulolytic polypeptides are usually produced in the cell as immature polypeptides comprising a signal sequence that is cleaved off during secretion of the protein.

They may also be further processed during secretion both at the N-terminal and/or C-terminal end to give a mature, enzymatically active protein. A polypeptide “comprising a fragment having cellulolytic activity” thus means that the polypeptide may be either in immature or mature form, preferably it is in mature form, i.e. the processing has taken place.

[0096] The novel polypeptides may further be a “fragment of the polypeptides or variants” mentioned above. The fragment may be the mature form of the proteins mentioned above, or it may be only an enzymatically active part of the mature protein. According to one embodiment of the invention, the polypeptide has an amino acid sequence having at least 80, 85, 90, 95, or 99% identity to SEQ ID NO: 4, 6, 12, 14, 16, 20, 22, 24 or 26, or to a cellulolytically active fragment thereof. It may also be a variant thereof, or a fragment thereof having cellobiohydrolase, endoglucanase, xylanase, or beta-glucosidase activity. According to another embodiment of the invention, the polypeptide consists essentially of a cellulolytically active fragment of a sequence of SEQ ID NO: 4, 6, 12, 14, 16, 20, 22, 24 or 26.

[0097] The novel polynucleotides may comprise a nucleotide sequence of SEQ ID NO: 3, 5, 11, 13, 15, 19, 21, 23 or 25, or a sequence encoding a novel polypeptide as defined above, including complementary strands thereof. Polynucleotide as used herein refers to both RNA and DNA, and it may be single stranded or double stranded. The polynucleotide may also be a fragment of said polynucleotides comprising at least 20 nucleotides, e.g. at least 25, 30 or 40 nucleotides. According to one embodiment of the invention it is at least 100, 200 or 300 nucleotides in length. Further the polynucleotide may be degenerate as a result of the genetic code to any one of the sequences as defined above. This means that different codons may code for the same amino acid.

[0098] According to one embodiment of the invention the polynucleotide is “comprised in” SEQ ID NO: 3, 5, 11, 13, 15, 19, 21, 23 or 25, which means that the sequence has at least part of the sequence mentioned. According to another embodiment of the invention, the polynucleotide comprises a gene similar to that included in a microorganism having accession number DSM 16728, DSM 16729, DSM 17324, DSM 17323, DSM 17729, DSM 16726, DSM 16725, DSM 17325 or DSM 17667.

[0099] The novel proteins/polypeptides may be prepared as described above. The novel polynucleotides may be inserted into a vector, which is capable of expressing the polypeptide encoded by the heterologous sequence, and the vector may be inserted into a host cell capable of expressing said polypeptide. The host cell is preferably of the genus *Trichoderma* or *Aspergillus*.

[0100] A heterologous gene encoding the novel polypeptides has been introduced on a plasmid into an *Escherichia coli* strain having accession number DSM 16728, DSM 16729, DSM 17324, DSM 17323, DSM 17729, DSM 16726, DSM 16725, DSM 17325 or DSM 17667.

[0101] The novel enzymes may be components of an enzyme preparation. The enzyme preparation may comprise one or more of the novel polypeptides, and it may be e.g. in the form of spent culture medium, powder, granules or liquid. According to one embodiment of the invention it comprises cellobiohydrolase, endoglucanase, beta-glucosidase, and optionally xylanase activity and/or other enzyme activities. It may further comprise any conventional additives.

[0102] The novel enzymes may be applied in any process involving cellulolytic enzymes, such as in fuel, textile, deter-

gent, pulp and paper, food, feed or beverage industry, and especially in hydrolysing cellulosic material for the production of biofuel comprising ethanol. In the pulp and paper industry they may be used to modify cellulosic fibre for example in treating kraft pulp, mechanical pulp, or recycled paper.

[0103] The invention is illustrated by the following non-limiting examples. It should be understood, however, that the embodiments given in the description above and in the examples are for illustrative purposes only, and that various changes and modifications are possible within the scope of the invention.

EXAMPLES

Example 1

Screening for Strains Expressing Cellulolytic Activity and their Cultivation for Purification

[0104] About 25 fungal strains from the Roal Oy culture collection were tested for cellulolytic activity including beta-glucosidases. After preliminary screening six strains were chosen for further studies. These were *Thermoascus aurantiacus* ALKO4239 and ALKO4242, *Acremonium thermophilum* ALKO4245, *Talaromyces thermophilus* ALKO4246 and *Chaetomium thermophilum* ALKO4261 and ALKO4265.

[0105] The strains ALKO4239, ALKO4242 and ALKO4246 were cultivated in shake flasks at 42° C. for 7 d in the medium 3×B, which contains g/litre: Solka Floc cellulose 18, distiller's spent grain 18, oats spelt xylan 9, CaCO₃ 2, soybean meal 4.5, (NH₄)HPO₄ 4.5, wheat bran 3.0, KH₂PO₄ 1.5, MgSO₄·H₂O 1.5, NaCl 0.5, KNO₃ 0.9, locust bean gum 9.0, trace element solution #1 0.5, trace element solution #2 0.5 and Struktol (Stow, Ohio, USA) antifoam 0.5 ml; the pH was adjusted to 6.5. Trace element solution #1 has g/litre: MnSO₄ 1.6, ZnSO₄·7H₂O 3.45 and CoCl₂·6H₂O 2.0; trace element solution #2 has g/litre: FeSO₄·7H₂O 5.0 with two drops of concentrated H₂SO₄.

[0106] The strain ALKO4261 was cultivated in shake flasks in the medium 1×B, which has one third of each of the constituents of the 3×B medium (above) except it has same concentrations for CaCO₃, NaCl and the trace elements. The strain was cultivated at 45° C. for 7 d.

[0107] The strain ALKO4265 was cultivated in shake flasks in the following medium, g/l: Solka Floc cellulose 40, Pharmamedia™ (Traders Protein, Memphis, Tenn., USA) 10, corn steep powder 5, (NH₄)₂SO₄ 5 and KH₂PO₄ 15; the pH was adjusted to 6.5. The strain was cultivated at 45° C. for 7 d.

[0108] After the cultivation the cells and other solids were collected by centrifugation down and the supernatant was recovered. For the shake flask cultivations, protease inhibitors PMSF (phenylmethyl-sulphonyl fluoride) and pepstatin A were added to 1 mM and 10 µg/ml, respectively. If not used immediately, the preparations were stored in aliquots at -20° C.

[0109] For the estimation of the thermoactivity of the enzymes, assays were performed of the shake flask cultivation preparations at 50° C., 60° C., 65° C., 70° C. and 75° C. for 1 h, in the presence of 100 µg bovine serum albumin (BSA)/ml as a stabilizer. Preliminary assays were performed at 50° C. and 65° C. at two different pH values (4.8/5.0 or 6.0) in order to clarify, which pH was more appropriate for the thermoactivity assay.

[0110] All shake flask supernatants were assayed for the following activities:

[0111] Cellobiohydrolase I—like activity ('CBHI') and the endoglucanase I—like activity ('EGI'):

[0112] These were measured in 50 mM Na-acetate buffer with 0.5 mM MUL (4-methylumbelliferyl-beta-D-lactoside) as the substrate. Glucose (100 mM) was added to inhibit any interfering beta-glucosidase activity. The liberated 4-methylumbelliferyl was measured at 370 nm. The 'CBHI' and the 'EGI' activities were distinguished by measuring the activity in the presence and absence of cellobiose (5 mM). The activity that is not inhibited by cellobiose represents the 'EGI' activity and the remaining MUL activity represents the 'CBHI' activity (van Tilbeurgh et al, 1988). The assay was performed at pH 5.0 or 6.0 (see below).

[0113] The endoglucanase (CMCase) activity:

[0114] This was assayed with 2% (w/v) carboxymethylcellulose (CMC) as the substrate in 50 mM citrate buffer essentially as described by Bailey and Nevalainen 1981; Haakana et al. 2004. Reducing sugars were measured with the DNS reagent. The assay was performed at pH 4.8 or 6.0 (see below).

[0115] Beta-glucosidase (BGU) activity:

[0116] This was assayed with 4-nitrophenyl-β-D-glucopyranoside (1 mM) in 50 mM citrate buffer as described by Bailey and Nevalainen 1981. The liberated 4-nitrophenol was measured at 400 nm. The assay was performed at pH 4.8 or 6.0 (see below).

[0117] The relative activities of the enzymes are presented in FIG. 1. The relative activities were presented by setting the activity at 60° C. as 100% (FIG. 1). All strains produced enzymes, which had high activity at high temperatures (65° C.-75° C.).

[0118] For protein purifications. ALKO4242 was also grown in a 2 litre bioreactor (Braun Biostat® B, Braun, Melsungen, Germany) in the following medium, g/litre: Solka Floc cellulose 40, soybean meal 10, NH₄NO₃ 5, KH₂PO₄ 5, MgSO₄·7H₂O 0.5, CaCl₂·2H₂O 0.05, trace element solution #1 0.5, trace element solution #2 0.5. The aeration was 1 vvm, antifoam control with Struktol, stirring 200-800 rpm and temperature at 47° C. Two batches were run, one at pH 4.7±0.2 (NH₃/H₂SO₄) and the other with initial pH of pH 4.5. The cultivation time was 7 d. After the cultivation the cells and other solids were removed by centrifugation.

[0119] The strain ALKO4245 was grown in 2 litre bioreactor (Braun Biostat® B, Braun, Melsungen, Germany) in the following medium, g/litre: Solka Floc cellulose 40, corn steep powder 15, distiller's spent grain 5, oats spelt xylan 3, locust bean gum 3, (NH₄)₂SO₄ 5 and KH₂PO₄ 5. The pH range was 5.2±0.2 (NH₃/H₂SO₄), aeration 1 vvm, stirring 300-600 rpm, antifoam control with Struktol and the temperature 42° C. The cultivation time was 4 d. After the cultivation the cells and other solids were removed by centrifugation.

[0120] For enzyme purification, ALKO4261 was grown in a 10 litre bioreactor (Braun Biostat® ED, Braun, Melsungen, Germany) in the following medium, g/litre: Solka Floc cellulose 30, distiller's spent grain 10, oats spelt xylan 5, CaCO₃ 2, soybean meal 10, wheat bran 3.0, (NH₄)₂SO₄ 5, KH₂PO₄ 5, MgSO₄·7H₂O 0.5, NaCl 0.5, KNO₃ 0.3, trace element solution #1 0.5 and trace element solution #2 0.5. The pH range was 5.2±0.2 (NH₃/H₂SO₄), aeration 1 vvm, stirring 200-600 rpm, antifoam control with Struktol and the temperature 42° C. The cultivation time was 5 d. A second batch was grown under similar conditions except that Solka Floc was added to

40 g/l and spent grain to 15 g/l. The supernatants were recovered by centrifugation and filtering through Seitz-K 150 and EK filters (Pall SeitzSchenk Filtersystems GmbH, Bad Kreuznach, Germany). The latter supernatant was concentrated about ten fold using the Pellicon mini ultrafiltration system (filter NMWL 10 kDa; Millipore, Billerica, Mass., USA).

[0121] For enzyme purification, ALKO4265 was also grown in a 10 litre bioreactor (Braun Biostat® ED, Braun, Melsungen, Germany) in the same medium as above, except KH₂PO₄ was added to 2.5 g/l. The pH range was 5.3±0.3 (NH₃/H₃PO₄), aeration 0.6 vvm, stirring 500 rpm, antifoam control with Struktol and the temperature 43° C. The cultivation time was 7 d. The supernatants were recovered by centrifugation and filtering through Seitz-K 150 and EK filters (Pall SeitzSchenk Filtersystems GmbH, Bad Kreuznach, Germany). The latter supernatant was concentrated about 20 fold using the Pellicon mini ultrafiltration system (filter NMWL 10 kDa; Millipore, Billerica, Mass., USA).

Example 2

Purification and Characterization of Cellobiohydrolases from *Acremonium thermophilum* ALKO4245 and *Chaetomium thermophilum* ALKO4265

[0122] *Acremonium thermophilum* ALKO4245 and *Chaetomium thermophilum* ALKO4265 were grown as described in Example 1. The main cellobiohydrolases were purified using p-aminobenzyl 1-thio-β-cellobioside-based affinity column, prepared as described by Tomme et al., 1988.

[0123] The culture supernatants were first buffered into 50 mM sodium acetate buffer pH 5.0, containing 1 mM δ-glucanolactone and 0.1 M glucose in order to retard ligand hydrolysis in the presence of β-glucosidases. Cellobiohydrolases were eluted with 0.1 M lactose and finally purified by gel filtration chromatography using Superdex 200 HR 10/30 columns in the ÄKTA system (Amersham Pharmacia Biotech). The buffer used in gel filtration was 50 mM sodium phosphate pH 7.0, containing 0.15 M sodium chloride.

[0124] Purified cellobiohydrolases were analysed by SDS-polyacrylamide gel electrophoresis and the molecular mass of both proteins was determined to be approximately 70 kDa evaluated on the basis of the molecular mass standards (Low molecular weight calibration kit, Amersham Biosciences). Purified *Acremonium* and *Chaetomium* cellobiohydrolases were designated as At Cel7A and Ct Cel7A, respectively, following the scheme in Henrissat et al. (1998) (Henrissat, 1991; Henrissat and Bairoch, 1993).

[0125] The specific activity of the preparations was determined using 4-methylumbelliferyl-β-D-lactoside (MUL), 4-methylumbelliferyl-β-D-cellobioside (MUG2) or 4-methylumbelliferyl-β-D-cellotrioside (MUG3) as substrate (van Tilbeurgh et al., 1988) in 0.05 M sodium citrate buffer pH 5 at 50° C. for 10 min. Endoglucanase and xylanase activities were determined by standard procedures (according to IUPAC, 1987) using carboxymethyl cellulose (CMC) and birch glucuronoxylan (Bailey et al., 1992) as substrates. Specific activity against Avicel was calculated on the basis of reducing sugars formed in a 24 h reaction at 50° C., pH 5.0, with 1% substrate and 0.25 μM enzyme dosage. The protein content of the purified enzyme preparations was measured according to Lowry et al., 1951. To characterize the end products of hydrolysis, soluble sugars liberated in 24 h hydrolysis experiment, as described above, were analysed by

HPLC (Dionex). Purified cellobiohydrolase I (CBHI/Cel7A) of *Trichoderma reesei* was used as a reference.

[0126] The specific activities of the purified enzymes and that of *T. reesei* CBHI/Cel7A are presented in Table 1. The purified At Cel7A and Ct Cel7A cellobiohydrolases possess higher specific activities against small synthetic substrates as compared to *T. reesei* CBHI/Cel7A. The specific activity against Avicel was clearly higher with the herein disclosed enzymes. Low activities of the purified enzyme preparations against xylan and CMC may either be due to the properties of the proteins themselves, or at least partially to the remaining minor amounts of contaminating enzymes. The major end product of cellulose hydrolysis by all purified enzymes was cellobiose which is typical to cellobiohydrolases.

TABLE 1

Specific activities (nkat/mg) of the purified cellobiohydrolases and the reference enzyme of <i>T. reesei</i> (50° C., pH 5.0, 24 h).			
Substrate	<i>A. thermophilum</i>	<i>C. thermophilum</i>	<i>T. reesei</i>
	ALKO4245 Cel7A	ALKO4265 Cel7A	
Xylan	11.3	6.7	1.3
CMC	26.2	5.5	1.0
MUG2	9.2	18.9	4.3
MUG3	1.3	1.5	0.9
MUL	21.5	54.0	21.9
Avicel	1.8	1.4	0.6

[0127] Thermal stability of the purified cellobiohydrolases was determined at different temperatures. The reaction was performed in the presence of 0.1% BSA at pH 5.0 for 60 min using 4-methylumbelliferyl- β -D-lactoside as substrate. *C. thermophilum* ALKO4265 CBH/Cel7A and *A. thermophilum* ALKO4245 CBH/Cel7A were stable up to 65° and 60° C., respectively. The *T. reesei* reference enzyme (CBHI/Cel7A) retained 100% of activity up to 55° C.

Example 3

Purification and Characterization of an Endoglucanase from *Acremonium thermophilum* ALKO4245

[0128] *Acremonium thermophilum* ALKO4245 was grown as described in Example 1. The culture supernatant was incubated at 70° C. for 24 hours after which it was concentrated by ultrafiltration. The pure endoglucanase was obtained by sequential purification with hydrophobic interaction and cation exchange chromatography followed by gel filtration. The endoglucanase activity of the fractions collected during purification was determined using carboxymethyl cellulose (CMC) as substrate (procedure of IUPAC 1987). Protein content was measured by BioRad Assay Kit (Bio-Rad Laboratories) using bovine serum albumine as standard.

[0129] The concentrated culture supernatant was applied to a HiPrep 16/10 Butyl FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate buffer pH 6.0, containing 1 M (NH₄)₂SO₄. Bound proteins were eluted with the linear gradient from the above buffer to 5 mM potassium phosphate, pH 6.0. Fractions were collected and the endoglucanase activity was determined as described above. The endoglucanase activity was eluted in a broad conductivity area of 120 to 15 mS/cm.

[0130] Combined fractions were applied to a HiTrap SPXL cation exchange column equilibrated with 8 mM sodium acetate, pH 4.5. Bound proteins were eluted with a linear

gradient from 0 to 0.25 M NaCl in the equilibration buffer. The protein containing endoglucanase activity was eluted at the conductivity area of 3-7 mS/cm. Cation exchange chromatography was repeated and the protein eluate was concentrated by freeze drying.

[0131] The dissolved sample was loaded onto a Superdex 75 HR10/30 gel filtration column equilibrated with 20 mM sodium phosphate buffer pH 7.0, containing 0.15 M NaCl. The main protein fraction was eluted from the column with the retention volume of 13.3 ml. The protein eluate was judged to be pure by SDS-polyacryl amide gel electrophoresis and the molecular weight was evaluated to be 40 kDa. The specific activity of the purified protein, designated as At EG-40, at 50° C. was determined to be 450 nkat/mg (procedure of IUPAC 1987, using CMC as substrate).

[0132] Thermal stability of the purified endoglucanase was determined at different temperatures. The reaction was performed in the presence of 0.1 mg/ml BSA at pH 5.0 for 60 min using carboxymethyl cellulose as substrate. *A. thermophilum* EG_40/Cel45A was stable up to 80° C. The *T. reesei* reference enzymes EGI (Cel7B) and EGII (Cel5A) retained 100% of activity up to 60° C. and 65° C., respectively.

Example 4

Purification of an Endoglucanase from *Chaetomium thermophilum* ALKO4261

[0133] *Chaetomium thermophilum* ALKO4261 was grown as described in Example 1. The pure endoglucanase was obtained by sequential purification with hydrophobic interaction and cation exchange chromatography followed by gel filtration. The endoglucanase activity of the fractions collected during purification was determined using carboxymethyl cellulose (CMC) as substrate (procedure of IUPAC 1987).

[0134] Ammonium sulfate was added to the culture supernatant to reach the same conductivity as 20 mM potassium phosphate pH 6.0, containing 1 M (NH₄)₂SO₄. The sample was applied to a HiPrep 16/10 Phenyl FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate pH 6.0, containing 1 M (NH₄)₂SO₄. Elution was carried out with a linear gradient of 20 to 0 mM potassium phosphate, pH 6.0, followed by 5 mM potassium phosphate, pH 6.0 and water. Bound proteins were eluted with a linear gradient of 0 to 6 M Urea. Fractions were collected and the endoglucanase activity was analysed as described above. The protein containing endoglucanase activity was eluted in the beginning of the urea gradient.

[0135] The fractions were combined, equilibrated to 16 mM Tris-HCl pH 7.5 (I=1.4 mS/cm) by 10DG column (Bio-Rad) and applied to a HiTrap DEAE FF anion exchange column equilibrated with 20 mM Tris-HCl, pH 7.5. Bound proteins were eluted with a linear gradient from 0 to 1 M NaCl in the equilibration buffer. Fractions were collected and analyzed for endoglucanase activity as described above. The protein was eluted in the range of 10-20 mS/cm.

[0136] The sample was equilibrated to 15 mM sodium acetate, pH 4.5 by 10DG column (Bio-Rad) and applied to a HiTrap SP XL cation exchange column equilibrated with 20 mM sodium acetate pH 4.5. Proteins were eluted with a linear gradient from 0 to 0.4 M sodium acetate, pH 4.5. Endoglucanase activity was eluted in the range of 1-10 mS/cm. The collected sample was lyophilized.

[0137] The sample was dissolved in water and applied to a Superdex 75 HR 10/30 gel filtration column equilibrated with 20 mM sodium phosphate pH 6.0, containing 0.15 M NaCl. Fractions were collected and those containing endoglucanase activity were combined. The protein eluate was judged to be pure by SDS-polyacrylamide gel electrophoresis and the molecular mass was evaluated on the basis of molecular mass standards (prestained SDS-PAGE standards, Broad Range, Bio-Rad) to be 54 kDa. The pI of the purified protein, designated as Ct EG_54 was determined with PhastSystem (Pharmacia) to be ca 5.5.

Example 5

Purification of an Endoglucanase from *Thermoascus aurantiacus* ALKO4242

[0138] *Thermoascus aurantiacus* ALKO4242 was grown as described in Example 1. The pure endoglucanase was obtained by sequential purification with hydrophobic interaction and anion exchange chromatography followed by gel filtration. The endoglucanase activity of the fractions collected during purification was determined using carboxymethyl cellulose (CMC) as substrate (procedure of IUPAC 1987). Protein content was measured by BioRad Assay Kit (Bio-Rad Laboratories) using bovine serum albumine as standard.

[0139] The culture supernatant was applied to a HiPrep 16/10 Butyl hydrophobic interaction column equilibrated with 20 mM potassium phosphate buffer pH 6.0, containing 0.7 M $(\text{NH}_4)_2\text{SO}_4$. Bound proteins were eluted with 0.2 M $(\text{NH}_4)_2\text{SO}_4$ (I=39 mS/cm). Fractions containing endoglucanase activity were combined and concentrated by ultrafiltration.

[0140] The sample was desalted in 10DG columns (Bio-Rad) and applied to a HiTrap DEAE FF anion exchange column equilibrated with 15 mM Tris-HCL, pH 7.0. Bound proteins were eluted with a linear gradient from 0 to 0.4 M NaCl in the equilibration buffer. The protein containing endoglucanase activity was eluted at the conductivity area of 15-21 mS/cm. Collected fractions were combined and concentrated as above.

[0141] The sample was applied to a Sephacryl S-100 HR 26/60 gel filtration column equilibrated with 50 mM sodium acetate buffer pH 5.0, containing 0.05 M NaCl. The protein fraction containing endoglucanase activity was eluted from the column with a retention volume corresponding to a molecular weight of 16 kDa. Collected fractions were combined, concentrated and gel filtration was repeated. The protein eluate was judged to be pure by SDS-polyacrylamide gel electrophoresis and the molecular weight was evaluated to be 28 kDa. The pI of the purified protein, designated as Ta EG_28, was determined in an IEF gel (PhastSystem, Pharmacia) to be about 3.5. The specific activity of Ta EG_28 at 50° C. was determined to be 4290 nkat/mg (procedure of IUPAC 1987, using CMC as substrate).

Example 6

Purification and Characterization of a β -Glucosidase from *Acremonium thermophilum* ALKO4245

[0142] *Acremonium thermophilum* ALKO4245 was grown as described in Example 1. The pure β -glucosidase was obtained by sequential purification with hydrophobic interaction and anion exchange chromatography followed by gel

filtration. The β -glucosidase activity of the fractions collected during purification was determined using 4-nitrophenyl- β -D-glucopyranoside as substrate (Bailey and Linko, 1990). Protein content was measured by BioRad Assay Kit (Bio-Rad Laboratories) using bovine serum albumine as standard.

[0143] The culture supernatant was applied to a HiPrep 16/10 Phenyl Sepharose FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate pH 6.0, containing 1 M $(\text{NH}_4)_2\text{SO}_4$. Bound proteins were eluted with a linear gradient from the equilibration buffer to 5 mM potassium phosphate in the conductivity area 137-16 mS/cm. Collected fractions were combined and concentrated by ultrafiltration.

[0144] The sample was desalted in 10DG columns (Bio-Rad) and applied to a HiTrap DEAE FF anion exchange column equilibrated with 10 mM potassium phosphate pH 7.0. Bound proteins were eluted with a linear gradient from the equilibration buffer to the same buffer containing 0.25 M NaCl in the conductivity area 1.5-12 mS/cm. Anion exchange chromatography was repeated as above, except that 4 mM potassium phosphate buffer pH 7.2 was used. Proteins were eluted at the conductivity area of 6-9 mS/cm. Fractions containing β -glucosidase activity were collected, combined, and concentrated.

[0145] The active material from the anion exchange chromatography was applied to a Sephacryl S-300 HR 26/60 column equilibrated with 20 mM sodium phosphate pH 6.5, containing 0.15 M NaCl. The protein with β -glucosidase activity was eluted with a retention volume corresponding to a molecular weight of 243 kDa. The protein was judged to be pure by SDS-polyacrylamide gel electrophoresis and the molecular weight was evaluated to be 101 kDa. The pI of the purified protein, designated as At β G_101, was determined in an IEF gel (PhastSystem, Pharmacia) to be in the area of 5.6-4.9. The specific activity of At β G_101 at 50° C. was determined to be 1100 nkat/mg (using 4-nitrophenyl- β -D-glucopyranoside as substrate, Bailey and Linko, 1990).

[0146] Thermal stability of the purified β -glucosidase was determined at different temperatures. The reaction was performed in the presence of 0.1 mg/ml BSA at pH 5.0 for 60 min using 4-nitrophenyl- β -D-glucopyranoside as substrate. *A. thermophilum* β G_101 was stable up to 70° C. The *Aspergillus* reference enzyme (Novozym 188) retained 100% of activity up to 600.

Example 7

Purification of a β -Glucosidase from *Chaetomium thermophilum* ALKO4261

[0147] *Chaetomium thermophilum* ALKO4261 was grown as described in Example 1. The pure β -glucosidase was obtained by sequential purification with hydrophobic interaction, anion and cation exchange chromatography followed by gel filtration. The β -glucosidase activity of the fractions collected during purification was determined using 4-nitrophenyl- β -D-glucopyranoside as substrate (Bailey and Linko, 1990).

[0148] The culture supernatant was applied to a HiPrep 16/10 Phenyl Sepharose FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate pH 6.0, containing 0.8 M $(\text{NH}_4)_2\text{SO}_4$. The elution was carried out with a linear gradient from the equilibration buffer to 3 mM potassium phosphate, pH 6.0, followed by elution with water and 6 M urea. The first fractions with β -glucosidase activity were

eluted in the conductivity area of 80-30 mS/cm. The second β -glucosidase activity was eluted with 6 M urea. The active fractions eluted by urea were pooled and desalted in 10DG columns (Bio-Rad) equilibrated with 10 mM Tris-HCl pH 7.0.

[0149] After desalting, the sample was applied to a HiTrap DEAE FF anion exchange column equilibrated with 15 mM Tris-HCl pH 7.0. The protein did not bind to the column but was eluted during the sample feed. This flow-through fraction was desalted in 1 ODG columns (Bio-Rad) equilibrated with 7 mM Na acetate, pH 4.5.

[0150] The sample from the anion exchange chromatography was applied to a HiTrap SP FF cation exchange column equilibrated with 10 mM sodium acetate pH 4.5. Bound proteins were eluted with a linear gradient from 10 mM to 400 mM sodium acetate, pH 4.5. The fractions with β -glucosidase activity eluting in conductivity area of 6.5-12 mS/cm were collected, desalted in 10DG columns (Bio-Rad) equilibrated with 7 mM sodium acetate, pH 4.5 and lyophilized.

[0151] The lyophilized sample was diluted to 100 μ l of water and applied to a Superdex 75 HF10/30 gel filtration column equilibrated with 20 mM sodium phosphate pH 4.5, containing 0.15 M NaCl. The β -glucosidase activity was eluted at a retention volume of 13.64 ml. Collected fractions were combined, lyophilized and dissolved in water. The protein was judged to be pure by SDS-polyacryl amide gel electrophoresis and the molecular weight was evaluated to be 76 kDa. The protein was designated as Ct β G₇₆.

Example 8

Purification and Characterization of a β -glucosidase from *Thermoascus aurantiacus* ALKO4242

[0152] *Thermoascus aurantiacus* ALKO4242 was grown as described in Example 1. The pure β -glucosidase was obtained by sequential purification with hydrophobic interaction, anion and cation exchange chromatography followed by gel filtration. The β -glucosidase activity of the fractions collected during purification was determined using 4-nitrophenyl- β -D-glucopyranoside as substrate (Bailey and Linko, 1990). Protein content was measured by BioRad Assay Kit (Bio-Rad Laboratories) using bovine serum albumine as standard.

[0153] The culture supernatant was applied to a HiPrep 16/10 Phenyl Sepharose FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate pH 6.0, containing 0.7 M $(\text{NH}_4)_2\text{SO}_4$. Bound proteins were eluted with a linear gradient from 0.2 M $(\text{NH}_4)_2\text{SO}_4$ to 5 mM potassium phosphate, pH 6.0. The β -glucosidase activity was eluted during the gradient in the conductivity area of 28.0-1.1 mS/cm. Fractions were combined and concentrated by ultrafiltration.

[0154] The sample was desalted in 10DG columns (Bio-Rad) and applied to a HiTrap DEAE FF anion exchange column equilibrated with 20 mM Tris-HCl pH 7.0. The enzyme was eluted with a linear gradient from 0 to 0.2 M NaCl in the equilibration buffer and with delayed elution by 20 mM Tris-HCl, containing 0.4 M NaCl. The sample eluting in the conductivity area of ca. 10-30 mS/cm was concentrated by ultrafiltration and desalted by 10DG column (Bio-Rad).

[0155] The sample was applied to a HiTrap SP XL cation exchange column equilibrated with 9 mM sodium acetate pH 4.5. The enzyme was eluted with a linear gradient from 10 mM to 400 mM NaAc and by delayed elution using 400 mM

NaAc pH 4.5. Proteins with β -glucosidase activity were eluted broadly during the linear gradient in the conductivity area of 5.0-11.3 mS/cm.

[0156] The active material from the cation exchange chromatography was applied to a Sephacryl S-300 HR 26/60 column equilibrated with 20 mM sodium phosphate pH 7.0, containing 0.15 M NaCl. The protein with β -glucosidase activity was eluted with a retention volume corresponding to a molecular weight of 294 kDa. Collected fractions were combined, lyophilized and dissolved in water. The protein was judged to be pure by SDS-polyacrylamide gel electrophoresis and the molecular weight was evaluated to be 81 kDa, representing most likely the monomeric form of the protein. Isoelectric focusing (IEF) was carried out using a 3-9 μ l gel. After silver staining, a broad area above pI 5.85 was stained in addition to a narrow band corresponding to pI 4.55. The specific activity of the purified protein, designated as Ta β G₈₁, at 50° C. was determined to be 600 nkat/mg using 4-nitrophenyl- β -D-glucopyranoside as substrate (Bailey and Linko, 1990).

[0157] Thermal stability of the purified β -glucosidase was determined at different temperatures. The reaction was performed in the presence of 0.1 mg/ml BSA at pH 5.0 for 60 min using 4-nitrophenyl- β -D-glucopyranoside as substrate. *T. aurantiacus* β G₈₁ was stable up to 75° C. The *Aspergillus* reference enzyme (Novozym 188) retained 100% of activity up to 60° C.

Example 9

Purification of a Xylanase from *Acremonium thermophilum* ALKO4245

[0158] *Acremonium thermophilum* ALKO4245 was grown as described in Example 1. The culture supernatant was incubated at 70° C. for 24 hours after which, it was concentrated by ultrafiltration. The pure xylanase was obtained by sequential purification with hydrophobic interaction and cation exchange chromatography followed by gel filtration. The xylanase activity was determined using birch xylan as substrate (procedure of IUPAC 1987). Protein was assayed by BioRad Protein Assay Kit (Bio-Rad Laboratories) using bovine serum albumin as standard.

[0159] The concentrated culture supernatant was applied to a HiPrep 16/10 Butyl FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate buffer pH 6.0, containing 1 M $(\text{NH}_4)_2\text{SO}_4$. Bound proteins were eluted with the linear gradient from the above buffer to 5 mM potassium phosphate, pH 6.0. The protein fraction was eluted in a broad conductivity area of 120 to 15 mS/cm.

[0160] The sample from the hydrophobic interaction column was applied to a HiTrap SP XL cation exchange column equilibrated with 8 mM sodium acetate, pH 4.5. The protein did not bind to this column but was eluted in the flow-through during sample feed. This eluate was concentrated by ultrafiltration. The hydrophobic chromatography was repeated as described above. The unbound proteins were collected and freeze dried.

[0161] The dissolved sample was loaded onto the Superdex 75 HR10/30 gel filtration column equilibrated with 20 mM sodium phosphate buffer pH 7.0, containing 0.15 M NaCl. The protein eluted from the column with the retention volume of 11.2 ml was judged to be pure by SDS-polyacryl amide gel electrophoresis. The molecular mass of the purified protein was evaluated on the basis of molecular mass standards

(prestained SDS-PAGE standards, Broad Range, Bio-Rad) to be 60 kDa. The specific activity of the protein, designated as At XYN_60, at 50° C. was determined to be 1800 nkat/mg (procedure of IUPAC 1987, using birch xylan as substrate). The relative activity was increased about 1.2 fold at 60° C. and 1.65 fold at 70° C. (10 min, pH 5.0) as compared to 50° C. The specific activity against MUG2 (4-methylumbelliferyl-β-D-cellobioside), MUL (4-methylumbelliferyl-beta-D-lactoside) and MUG3 (4-methylumbelliferyl-β-D-celotrioside) were 54, 33 and 78 nkat/mg (50° C. pH 5.0 10 min), respectively. This is in agreement with the fact that the family 10 xylanases also show activity against the aryl glucopyranosides (Biely et al. 1997).

Example 10

Purification of a Xylanase from *Thermoascus aurantiacus* ALKO4242

[0162] *Thermoascus aurantiacus* ALKO4242 was grown as described in Example 1. The pure xylanase was obtained by sequential purification with hydrophobic interaction, anion, and cation exchange chromatography followed by gel filtration. The xylanase activity was determined using birch xylan as substrate (procedure of IUPAC 1987). Protein was assayed by BioRad Protein Assay Kit (Bio-Rad Laboratories) using bovine serum albumin as standard.

[0163] The culture supernatant was applied to a HiPrep 16/10 Phenyl Sepharose FF hydrophobic interaction column equilibrated with 20 mM potassium phosphate buffer pH 6.0, containing 0.7 M (NH₄)₂SO₄. Bound proteins were eluted with a two-step elution protocol. The elution was carried out by dropping the salt concentration first to 0.2 M (NH₄)₂SO₄ and after that a linear gradient from 20 mM potassium phosphate pH 6.0, containing 0.2 M (NH₄)₂SO₄ to 5 mM potassium phosphate pH 6.0 was applied. The protein was eluted with 0.2 M (NH₄)₂SO₄ (I=39 mS/cm).

[0164] The sample was desalted in 10DG columns (Bio-Rad) and applied to a HiTrap DEAE FF anion exchange column equilibrated with 15 mM Tris-HCL, pH 7.0. The protein did not bind to the anion exchange column but was eluted in the flow-through. The conductivity of the sample was adjusted to correspond that of 20 mM sodium acetate, pH 4.5 by adding water and pH was adjusted to 4.5 during concentration by ultrafiltration.

[0165] The sample was applied to a HiTrap SP XL cation exchange column equilibrated with 20 mM sodium acetate, pH 4.5. Bound proteins were eluted with a linear gradient from the equilibration buffer to the same buffer containing 1 M NaCl. The enzyme was eluted at the conductivity area of 1-7 mS/cm. The sample was lyophilized and thereafter dissolved in water.

[0166] The lyophilized sample was dissolved in water and applied to a Superdex 75 HR 10/30 gel filtration column equilibrated with 20 mM sodium phosphate pH 7.0, containing 0.15 M NaCl. The protein was eluted from the column with a retention volume corresponding to a molecular weight of 26 kDa. The protein was judged to be pure by SDS-polyacrylamide gel electrophoresis. The molecular mass of the pure protein was 30 kDa as evaluated on the basis of molecular mass standards (prestained SDS-PAGE standards, Broad Range, Bio-Rad). The pI of the purified protein, designated as Ta XYN_30 was determined with PhastSystem (Pharmacia) to be ca. 6.8. The specific activity of Ta XYN_30 at 50° C.

was determined to be 4800 nkat/mg (procedure of IUPAC 1987, using birch xylan as substrate).

Example 11

Internal Amino Acid Sequencing

[0167] The internal peptides were sequenced by electrospray ionization combined to tandem mass spectrometry (ESI-MS/MS) using the Q-TOF1 (Micromass) instrument. The protein was first alkylated and digested into tryptic peptides. Generated peptides were desalted and partially separated by nano liquid chromatography (reverse-phase) before applying to the Q-TOF1 instrument. The internal peptide sequences for *Chaetomium thermophilum* and *Acremonium thermophilum* cellobiohydrolases are shown in Table 2. The peptides from *Chaetomium* CBH were obtained after the corresponding cbh gene had been cloned. The peptides determined from *Acremonium* CBH were not utilized in the cloning of the corresponding gene.

TABLE 2

Internal peptide sequences determined from <i>Chaetomium thermophilum</i> ALKO4265 CBH (1_C-4_C) and <i>Acremonium thermophilum</i> ALKO4245 CBH (1_A-4_A).	
Peptide	Sequence
Peptide 1_C	T P S T N D A N A G F G R
Peptide 2_C	V A F S N T D D F N R
Peptide 3_C	F S N T D D F N R K
Peptide 4_C	P G N S L/I T Q E Y C D A Q/K K
Peptide 1_A	V T Q F I/L T G
Peptide 2_A	M G D T S F Y G P G
Peptide 3_A	C D P D G C D F N
Peptide 4_A	S G N S L/I T T D F

I/L = leucine and isoleucine have the same molecular mass and cannot be distinguished in ESI-MS/MS analysis

Q/K = the molecular mass of glutamine and lysine differs only 0.036 Da and cannot be distinguished in ESI-MS/MS analysis

[0168] The internal peptide sequences of purified endoglucanases, β-glucosidases, and xylanases of *Acremonium thermophilum* ALKO4245, *Chaetomium thermophilum* ALKO4261 and *Thermoascus aurantiacus* ALKO4242 are listed in Table 3, Table 4 and Table 5.

TABLE 3

Internal peptide sequences determined from <i>Acremonium thermophilum</i> ALKO4245 EG_40, <i>Chaetomium thermophilum</i> ALKO4261 EG_54 and <i>Thermoascus aurantiacus</i> ALKO4242 EG_28 endoglucanases.	
Protein Peptide	Sequence ^(a)
At EG_40 Peptide 1	Q S C S S F P A P L K P G C Q W R
Peptide 2	Y A L T F N S G P V A G K
Peptide 3	V Q C P S E L T S R
Peptide 4	N Q P V F S C S A D W Q R

TABLE 3-continued

Internal peptide sequences determined from <i>Acremonium thermophilum</i> ALKO4245 EG_40, <i>Chaetomium thermophilum</i> ALKO4261 EG_54 and <i>Thermoascus aurantiacus</i> ALKO4242 EG_28 endoglucanases.		
Protein	Peptide	Sequence ^(a)
	Peptide 5	Y W D C C K P S C G W P G K
	Peptide 6	P T F T
Ct EG_54	Peptide 1	E P E P E V T Y Y V
	Peptide 2	Y Y L L D Q T E Q Y
	Peptide 3	R Y C A C M D L W E A N S R
	Peptide 4	P G N T P E V H P Q/K
	Peptide 5	S I/L A P H P C N Q/K
	Peptide 6	Q Q Y E M F R
	Peptide 7	A L N D D F C R
	Peptide 8	W G N P P P R
Ta EG_28	Peptide 1	I/L T S A T Q W L R
	Peptide 2	G C A I/L S A T C V S S T I/L G Q E R
	Peptide 3	P F M M E R
	Peptide 4	Q Y A V V D P H N Y G R

^(a)I/L = leucine and isoleucine have the same molecular mass and cannot be distinguished in ESI-MS/MS analysis, Q/K = the molecular mass of glutamine and lysine differs only 0.036 Da and cannot be distinguished in ESI-MS/MS analysis.

TABLE 4

Internal peptide sequences determined from <i>Acremonium thermophilum</i> ALKO4245 βG_101, <i>Chaetomium thermophilum</i> ALKO4261 βG_76 and <i>Thermoascus aurantiacus</i> ALKO4242 βG_81 beta- glucosidases.		
Protein	Peptide	Sequence ^(a)
At βG_101	Peptide 1	S P F T W G P T R
	Peptide 2	V V V G D D A G N P C
	Peptide 3	A F V S Q L T L L E K
	Peptide 4	G T D V L/I Y T P N N K
	Peptide 5	Q P N P A G P N A C V L/I R
Ct βG_76	Peptide 1	E G L F I D Y R
	Peptide 2	P G Q S G T A T F R
	Peptide 3	E T M S S N V D D R
	Peptide 4	I A L V G S A A V V
	Peptide 5	M W L C E N D R
	Peptide 6	Y P Q L C L Q D G P L G I R
	Peptide 7	E L N G Q N S G Y P S I
Ta βG_81	Peptide 1	T P F T W G K
	Peptide 2	L C L Q D S L P G V R
	Peptide 3	G V D V Q L G P V A G V A P R
	Peptide 4	V N L T L E
	Peptide 5	F T G V F G E D V V G
	Peptide 6	N D L P L T G Y E K

^(a)I/L = leucine and isoleucine have the same molecular mass and cannot be distinguished in ESI-MS/MS analysis

TABLE 5

Internal peptide sequences determined from <i>Acremonium thermophilum</i> ALKO4245 XYN_60 and <i>Thermoascus aurantiacus</i> ALKO4242 XYN_30 xylanases.		
Protein	Peptide	Sequence
At XYN_60	Peptide 1	Y N D Y N L E Y N Q K
	Peptide 2	F G Q V T P E N
	Peptide 3	V D G D A T Y M S Y V N N K
	Peptide 4	K P A W T S V S S V L A A K
	Peptide 5	S Q G D I V P R A K
Ta XYN_30	Peptide 1	V Y F G V A T D Q N R
	Peptide 2	N A A I I Q A D F G Q V T P E N S M K
	Peptide 3	G H T L V W H S Q L P S W V S S I T D K
	Peptide 4	N H I T T L M T R
	Peptide 5	A W D V V N E A F N E D G S L R
	Peptide 6	L Y I N D Y N L D S A S Y P K
	Peptide 7	A S T T P L L F D G N F N P K P A Y N A I V Q D L Q Q
	Peptide 8	Q T V F L N V I G E D Y I P I A F Q T A R

Example 12

Construction of Genomic Libraries for *Thermoascus aurantiacus*, *Chaetomium thermophilum* and *Acremonium thermophilum*

[0169] The genomic library of *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 were made to Lambda DASH®II vector (Stratagene, USA) according to the instructions from the supplier. The chromosomal DNAs, isolated by the method of Raeder and Broda (1985), were partially digested with Sau3A. The digested DNAs were size-fractionated and the fragments of the chosen size (≈5-23 kb) were dephosphorylated and ligated to the BamHI digested lambda vector arms. The ligation mixtures were packaged using Gigapack III Gold packaging extracts according to the manufacturer's instructions (Stratagene, USA). The titers of the *Chaetomium thermophilum* and *Acremonium thermophilum* genomic libraries were 3.6×10^6 pfu/ml and 3.7×10^5 pfu/ml and those of the amplified libraries were 6.5×10^{10} pfu/ml and 4.2×10^8 pfu/ml, respectively.

[0170] Lambda FIX® II/Xho I Partial Fill-In Vector Kit (Stratagene, USA) was used in the construction of the genomic libraries for *Thermoascus aurantiacus* ALKO4242 and *Chaetomium thermophilum* ALKO4261 according to the instructions from the supplier. The chromosomal DNAs, isolated by the method of Raeder and Broda (1985), were partially digested with Sau3A. The digested DNAs were size-fractionated and the fragments of the chosen size (6-23 kb) were filled-in and ligated to the XhoI digested Lambda FIX II vector arms. The ligation mixtures were packaged using Gigapack III Gold packaging extracts according to the manufacturer's instructions (Stratagene, USA). The titers of the *Thermoascus aurantiacus* ALKO4242 and *Chaetomium thermophilum* ALKO4261 genomic libraries were 0.2×10^6 and 0.3×10^6 pfu/ml and those of the amplified libraries were 1.8×10^9 and 3.8×10^9 pfu/ml, respectively.

Example 13

Cloning of the Cellobiohydrolase (cbh/cel7) Genes from *Thermoascus aurantiacus*, *Chaetomium thermophilum* and *Acremonium thermophilum*

[0171] Standard molecular biology methods were used in the isolation and enzyme treatments of DNA (plasmids, DNA fragments), in *E. coli* transformations, etc. The basic methods used are described in the standard molecular biology handbooks, e.g., Sambrook et al. (1989) and Sambrook and Russell (2001).

[0172] The probes for screening the genomic libraries which were constructed as described in Example 12 were amplified by PCR using the *Thermoascus aurantiacus* ALKO4242, *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 genomic DNAs as templates in the reactions. Several primers tested in PCR reactions were designed according to the published nucleotide sequence (WO 03/000941, Hong et al., 2003b). The PCR reaction mixtures contained 50 mM Tris-HCl, pH 9.0, 15 mM (NH₄)₂SO₄, 0.1% Triton X-100, 1.5 mM MgCl₂, 0.2 mM dNTPs, 5 μM each primer and 1 units of Dynazyme EXT DNA polymerase (Finnzymes, Finland) and 0.5-1 μg of the genomic DNA. The conditions for the PCR reactions were the following: 5 min initial denaturation at 95° C., followed by 30 cycles of 1 min at 95° C., either 1 min annealing at 62° C. (±8° C. gradient) for *Thermoascus* ALKO4242 and *Chaetomium* ALKO4265 templates or 1 min annealing at 58° C. (±6° C. gradient) for *Acremonium* ALKO4245 template, 2 min extension at 72° C. and a final extension at 72° C. for 10 min.

[0173] DNA products of the expected sizes (calculated from published cbh sequences) were obtained from all genomic templates used. The DNA fragments of the expected sizes were isolated from the most specific PCR reactions and they were cloned to pCR® Blunt-TOPO® vector (Invitrogen, USA). The inserts were characterized by sequencing and by performing Southern blot hybridizations to the genomic DNAs digested with several restriction enzymes. The PCR fragments, which were chosen to be used as probes for screening of the *Thermoascus aurantiacus*, *Chaetomium thermophilum* and *Acremonium thermophilum* genomic libraries are presented in Table 6.

[0174] The deduced amino acid sequences from all these probes had homology to several published CBH sequences (BLAST program, version 2.2.9 at NCBI, National Center for Biotechnology Information; Altschul et al., 1990) of glycoside hydrolase family 7 (Henrissat, 1991; Henrissat and Bairoch, 1993).

[0175] The inserts from the plasmids listed in Table 6 were labeled with digoxigenin according to the supplier's instructions (Roche, Germany), and the amplified genomic libraries (2×10⁵-3×10⁵ plaques) were screened with the labeled probe fragments. The hybridization temperature for the filters was 68° C. and the filters were washed 2×5 min at RT using 2×SSC-0.1% SDS followed by 2×15 min at 68° C. using 0.1×SSC-0.1% SDS with the homologous probes used. Several positive plaques were obtained from each of the hybridizations. In screening of the *Acremonium* ALKO4245 genomic libraries, some of the positive plaques were strongly hybridizing to the probe in question but, in addition, there was an amount of plaques hybridizing more weakly to the probes. This suggested that other cellobiohydrolase gene(s) might be present in the genome, causing cross-reaction. From four to five strongly hybridizing plaques were purified from *Thermoascus* ALKO4242 and *Chaetomium* ALKO4265 genomic library screenings. In the case of the *Acremonium thermophilum* ALKO4245, four out of six purified plaques hybridized weakly by the probe used. The phage DNAs were isolated and characterized by Southern blot hybridizations. The chosen restriction fragments hybridizing to the probe were subcloned to pBluescript II KS+vector and the relevant regions of the clones were sequenced.

[0176] In total four cbh/cel7 genes were cloned; one from *Thermoascus aurantiacus* ALKO4242, one from *Chaetomium thermophilum* ALKO4265 and two from *Acremonium thermophilum* ALKO4245 (at the early phase of the work, these had the codes At_cbh_C and At_cbh_A, and were then designated as At cel7A and At cel7B, respectively). Table 7 summarizes the information on the probes used for screening the genes, the phage clones from which the genes were isolated, the chosen restriction fragments containing the full-length genes with their promoter and terminator regions, the plasmid names, and the DSM deposit numbers for the *E. coli* strains carrying these plasmids.

TABLE 6

The primers used in the PCR reactions and probes chosen for screening of the cbh/cel7 genes from <i>Thermoascus aurantiacus</i> , <i>Chaetomium thermophilum</i> and <i>Acremonium thermophilum</i> genomic libraries. The genomic template DNA and the name of the plasmid containing the probe fragment are shown.					
Gene	Forward primer	Reverse primer	Template DNA	Fragment (kb)	Plasmid
Ta cbh	TCEL11 atgcgaactggcgcttgggtcc	TCEL12 gaatttggagctagtgtcgacg	<i>Thermoascus</i> ALKO4242	0.8 kb	pALK1633
Ct cbh	TCEL7 cgatgccaaactggcgctggac	TCEL8 ttcttggtggtgtcgacggtc	<i>Chaetomium</i> ALKO4265	0.8 kb	pALK1632
At cbh	TCEL13 agctcgaccaactgctacacg	TCEL4 accgtgaacttcttctgctggtg	<i>Acremonium</i> ALKO4245	0.7 kb	pALK1634

TABLE 7

The probes used for cloning of cbh/cel7 genes, the phage clone and the subclones chosen, the plasmid number and the number of the deposit of the corresponding <i>E. coli</i> strain.					
Gene	Probe used in screening	Phage clone	The fragment subcloned to pBluescript II	Plasmid no	<i>E. coli</i> deposit no
Ta cel7A	pALK1633	F12	3.2 kb XbaI	pALK1635	DSM 16723
Ct cel7A	pALK1632	F36	2.3 kb PvuI - HindIII	pALK1642	DSM 16727
At cel7B	pALK1634	F6	3.1 kb EcoRI	pALK1646	DSM 16728
At cel7A	pALK1634	F2	3.4 kb XhoI	pALK1861	DSM 16729

the deduced amino acid sequences of the clones containing the Ct cel7A and At cel7A genes. Thus, it could be concluded that the genes encoding the purified CBH/Cel7 proteins from *Chaetomium thermophilum* and *Acremonium thermophilum* were cloned.

TABLE 8

Summary on the cbh/cel7 genes isolated from <i>Thermoascus aurantiacus</i> ALKO4242, <i>Chaetomium thermophilum</i> ALKO4265 and <i>Acremonium thermophilum</i> ALKO4245.					
Cbh gene	Length with introns (bp) ^(a)	Coding region (bp) ^(b)	No of introns	Lengths of introns (bp)	SEQ ID NO:
Ta cel7A	1439	1371	1	65	1
Ct cel7A	1663	1596	1	64	7
At cel7B	1722	1377	3	134, 122, 87	3
At cel7A	1853	1569	4	88, 53, 54, 86	5

^(a)The STOP codon is included.

^(b)The STOP codon is not included.

TABLE 9

Summary of amino acid sequences deduced from the cbh/cel7 gene sequences from <i>Thermoascus aurantiacus</i> ALKO4242, <i>Chaetomium thermophilum</i> ALK4265 and <i>Acremonium thermophilum</i> ALKO4245.							
CBH protein	No of aas	Length of ss NN/HMM ^(a)	C-terminal CBD ^(b)	Predicted MW (Da, ss not incl) ^(c)	Predicted pl (ss not incl)	Putative N-glycosylation sites ^(d)	SEQ ID NO:
Ta Cel7A	457	17/17	NO	46 873	4.44	2	2
Ct Cel7A	532	18/18	YES, T497 to L532	54 564	5.05	3	8
At Cel7B	459	21/21	NO	47 073	4.83	2	4
At Cel7A	523	17/17	YES, Q488 to L523	53 696	4.67	4	6

ss, signal sequence.

^(a)The prediction on the signal sequence was made using the program SignalP V3.0 (Nielsen et al., 1997; Bendtsen et al., 2004); the NN value was obtained using neural networks and HMM value using hidden Markov models.

^(b)The cellulose-binding domain (CBD), the amino acids of the C-terminal CBD region are indicated (M1 (Met #1) included in numbering)

^(c)The predicted signal sequence was not included. The prediction was made using the Compute pl/MW tool at ExPASy server (Gasteiger et al., 2003).

^(d)The number of sequences N-X-S/T.

[0177] The relevant information on the genes and the deduced protein sequences (SEQ ID NO: 1-8) are summarized in Table 8 and Table 9, respectively.

[0178] The peptide sequences of the purified CBH proteins from *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 (Table 2) were found from

[0179] The deduced amino acid sequences of *Thermoascus aurantiacus* Cel7A and *Acremonium thermophilum* Cel7A (core, without the CBD) were most homologous to each other (analyzed by Needleman-Wunsch global alignment, EMBOSS 3.0.0 Needle, with Matrix EBLOSUM62, Gap Penalty 10.0 and Extend Penalty 0.5; Needleman and Wun-

sch, 1970). In addition, the deduced *Acremonium thermophilum* Cel7A had a lower identity to the deduced *Chaetomium thermophilum* Cel7A. The *Acremonium thermophilum* Cel7B was most distinct from the CBH/Cel7 sequences of the invention.

[0180] The deduced *Chaetomium* Cel7A sequence possessed the highest identities (analyzed by Needleman-Wunsch global alignment, EMBOSS Needle, see above) to polypeptides of *Chaetomium thermophilum*, *Scytalidium thermophilum* and *Thielavia australiensis* CBHI described in WO 03/000941. Similarly, the deduced *Thermoascus aurantiacus* Cel7A sequence was highly identical to the published CBHI of the *Thermoascus aurantiacus* (WO 03/000941, Hong et al., 2003b). *Acremonium thermophilum* Cel7B had significantly lower identities to the previously published sequences, being more closely related to the CBHI polypeptide from *Oryza sativa*. The highest homologies of the deduced *Acremonium thermophilum* Cel7A sequence were to *Exidia glandulosa* and *Acremonium thermophilum* CBHI polynucleotides (WO 03/000941). The alignment indicates that the cloned *Thermoascus aurantiacus* ALKO4242, *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 sequences encode the CBH proteins having high homology to the polypeptides of the glycoside hydrolase family 7, therefore these were designated as Cel7A or Cel7B (Henrissat et al. 1998).

[0181] The comparison of the deduced amino acid sequences of the cbh/cel7 genes from *Thermoascus aurantiacus* ALKO4242, *Chaetomium thermophilum* ALKO4265 and *Acremonium thermophilum* ALKO4245 *Thielavia* to each other, and further to the sequences found from the databases, are shown in Table 10.

TABLE 10

The highest homology sequences to the deduced amino acid sequences of the cbh/cel7 genes from <i>Thermoascus aurantiacus</i> ALKO4242, <i>Chaetomium thermophilum</i> ALKO4265 and <i>Acremonium thermophilum</i> ALKO4245. The alignment was made using Needleman-Wunsch global alignment (EMBLOSUM62, Gap penalty 10.0, Extend penalty 0.5).	
Organism, enzyme and accession number	Identity, (%)
* <i>Thermoascus aurantiacus</i> Cel7A	100.0
<i>Thermoascus aurantiacus</i> , AY840982	99.6
<i>Thermoascus aurantiacus</i> , AX657575	99.1
<i>Thermoascus aurantiacus</i> , AF421954	97.8
<i>Talaromyces emersonii</i> , AY081766	79.5
<i>Chaetomidium pingtungium</i> , AX657623	76.4
<i>Trichophaea saccata</i> , AX657607	73.4
* <i>Acremonium thermophilum</i> Cel7A (core)	70.6
<i>Emericella nidulans</i> , AF420020 (core)	70.4
* <i>Chaetomium thermophilum</i> Cel7A (core)	66.4
* <i>Chaetomium thermophilum</i> Cel7A	100.0
<i>Chaetomium thermophilum</i> , AY861347	91.9
<i>Chaetomium thermophilum</i> , AX657571	91.7
<i>Scytalidium thermophilum</i> , AX657627	74.7
<i>Thielavia australiensis</i> , AX657577	74.6
<i>Acremonium thermophilum</i> , AX657569	72.3
<i>Exidia glandulosa</i> , AX657613	68.0
* <i>Acremonium thermophilum</i> Cel7A	66.9
* <i>Thermoascus aurantiacus</i> Cel7A (core)	66.4
<i>Exidia glandulosa</i> , AX657615	60.8
<i>Chaetomium pingtungium</i> , AX657623	60.7
* <i>Acremonium thermophilum</i> Cel7B (core)	60.2
* <i>Acremonium thermophilum</i> Cel7B	100.0
<i>Oryza sativa</i> , AK108948	66.1

TABLE 10-continued

The highest homology sequences to the deduced amino acid sequences of the cbh/cel7 genes from <i>Thermoascus aurantiacus</i> ALKO4242, <i>Chaetomium thermophilum</i> ALKO4265 and <i>Acremonium thermophilum</i> ALKO4245. The alignment was made using Needleman-Wunsch global alignment (EMBLOSUM62, Gap penalty 10.0, Extend penalty 0.5).	
Organism, enzyme and accession number	Identity, (%)
<i>Exidia glandulosa</i> , AX657615	65.0
<i>Acremonium thermophilum</i> , AX657569 (core)	64.8
<i>Thermoascus aurantiacus</i> , AX657575	64.8
* <i>Acremonium thermophilum</i> Cel7A	64.6
* <i>Thermoascus aurantiacus</i> Cel7A	64.4
<i>Trichophaea saccata</i> , AX657607	63.6
* <i>Chaetomium thermophilum</i> Cel7A (core)	60.2
* <i>Acremonium thermophilum</i> Cel7A	100.0
<i>Exidia glandulosa</i> , AX657613	77.9
<i>Exidia glandulosa</i> , AX657615	77.9
<i>Acremonium thermophilum</i> , AX657569	77.5
<i>Thielavia australiensis</i> , AX657577	71.0
* <i>Thermoascus aurantiacus</i> Cel7A (core)	70.6
<i>Scytalidium thermophilum</i> , AX657627	67.5
<i>Chaetomium thermophilum</i> , AX657571	67.5
<i>Chaetomium pingtungium</i> , AX657623	67.3
* <i>Chaetomium thermophilum</i> Cel7A	66.9
* <i>Acremonium thermophilum</i> Cel7B (core)	64.6

*indicates an amino acid sequence derived from one of the cellobiohydrolase genes cloned in this work. 'Core' indicates alignment without the CBD.

Example 14

Production of Recombinant CBH/Cel7 Proteins in *Trichoderma reesei*

[0182] Expression plasmids were constructed for production of the recombinant CBH/Cel7 proteins from *Thermoascus aurantiacus* (Ta Cel7A), *Chaetomium thermophilum* (Ct Cel7A) and *Acremonium thermophilum* (At Cel7A, At Cel7B; at early phase of the work these proteins had the temporary codes At CBH_C and At CBH_A, respectively). The expression plasmids constructed are listed in Table 11. The recombinant cbh/cel7 genes, including their own signal sequences, were exactly fused to the *T. reesei* cbh1 (cel7A) promoter by PCR. The transcription termination was ensured by the *T. reesei* cel7A terminator and the *A. nidulans* amdS marker gene was used for selection of the transformants as described in Paloheimo et al. (2003). The linear expression cassettes (FIG. 2), were isolated from the vector backbones after EcoRI digestion and were transformed into *T. reesei* A96 and A98 protoplasts (both strains have the genes encoding the four major cellulases CBHI/Cel7A, CBHII/Cel6A, EGI/Cel7B and EGII/Cel5A deleted). The transformations were performed as in Penttilä et al. (1987) with the modifications described in Karhunen et al. (1993), selecting with acetamide as a sole nitrogen source. The transformants were purified on selection plates through single conidia prior to sporulating them on PD.

TABLE 11

The expression cassettes constructed to produce CBH/Cel7 proteins of *Thermoascus aurantiacus* ALKO4242 (Ta Cel7A), *Chaetomium thermophilum* ALKO4265 (Ct Cel7A), and *Acremonium thermophilum* ALKO4245 (At Cel7A, At Cel7B) in *Trichoderma reesei*. The overall structure of the expression cassettes was as described in FIG. 2. The cloned cbh/cel7 genes were exactly fused to the *T. reesei* cbh1/cel7A promoter.

CBH/Cel7	Expression plasmid	Size of the expr. cassette ^(a)	cel7A terminator ^(b)
Ta Cel7A	pALK1851	9.0 kb	245 bp (XbaI)
Ct Cel7A	pALK1857	9.2 kb	240 bp (HindIII)
At Cel7B	pALK1860	9.4 kb	361 bp (EcoRI)
At Cel7A	pALK1865	9.5 kb	427 bp (EcoRV)

^(a)The expression cassette for *T. reesei* transformation was isolated from the vector backbone by using EcoRI digestion.

^(b)The number of the nucleotides from the genomic cbh1/cel7A terminator region after the STOP codon. The restriction site at the 3'-end, used in excising the genomic gene fragment, is included in the parenthesis.

[0183] The CBH/Cel7 production of the transformants was analysed from the culture supernatants of the shake flask cultivations (50 ml). The transformants were grown for 7 days at 28° C. in a complex lactose-based cellulase-inducing medium (Joutsjoki et al. 1993) buffered with 5% KH₂PO₄. The cellobiohydrolase activity was assayed using 4-methylumbelliferyl-β-D-lactoside (MUL) substrate according to van Tilbeurgh et al., 1988. The genotypes of the chosen transformants were confirmed by using Southern blots in which several genomic digests were included and the respective expression cassette was used as a probe. Heterologous expression of the Ta Cel7A, Ct Cel7A, At Cel7A and At Cel7B proteins was analyzed by SDS-PAGE with subsequent Coomassive staining. The findings that no cellobiohydrolase activity or heterologous protein production in SDS-PAGE could be detected for the At Cel7B transformants containing integrated expression cassette, suggest that At Cel7B is produced below detection levels in *Trichoderma* using the described experimental design.

[0184] The recombinant CBH/Cel7 enzyme preparations were characterized in terms of pH optimum and thermal stability. The pH optimum of the recombinant CBH/Cel7 proteins from *Thermoascus aurantiacus*, *Chaetomium thermophilum*, and *Acremonium thermophilum* were determined in the universal McIlvaine buffer within a pH range of 3.0-8.0 using 4-methylumbelliferyl-β-D-lactoside (MUL) as a substrate (FIG. 3 A). The pH optimum for Ct Cel7A and At Cel7A enzymes is at 5.5, above which the activity starts to gradually drop. The pH optimum of the recombinant crude Ta Cel7A is at 5.0 (FIG. 3 A). Thermal stability of the recombinant Cel7 enzymes was determined by measuring the MUL activity in universal McIlvaine buffer at the optimum pH with reaction time of 1 h. As shown from the results Ta Cel7A and Ct Cel7A retained more than 60% of their activities at 70° C., whereas At Cel7A showed to be clearly less stable at the higher temperatures (≥65° C.) (FIG. 3 B).

[0185] The chosen CBH/Cel7 transformants were cultivated in lab bioreactors at 28° C. in the medium indicated above for 3-4 days with pH control 4.4±0.2 (NH₃/H₃PO₄) to obtain material for the application tests. The supernatants were recovered by centrifugation and filtering through Seitz-K 150 and EK filters (Pall SeitzSchenk Filtersystems GmbH, Bad Kreuznach, Germany).

Example 15

Production of the Recombinant *Thermoascus aurantiacus* Cel7A+CBD Fusion Proteins in *T. reesei*

[0186] *Thermoascus aurantiacus* Cel7A (AF478686, Hong et al., 2003b; SEQ ID. NO: 1) was fused to linker and CBD of *Trichoderma reesei* CBH1/Cel7A (AR088330, Srisodsuk et al. 1993) (=Tr CBD) followed by the production of the fusion protein (SEQ ID NO: 28 corresponding nucleic acid SEQ ID. NO: 27) in the *T. reesei* as was described in F120055205/U.S. Ser. No. 11/119,526; filed Apr. 29, 2005. In addition, *Thermoascus aurantiacus* Cel7A was fused to linker and CBD of *Chaetomium thermophilum* Cel7A (SEQ ID. NO: 7) (Ct CBD). For that purpose, the coding sequence of the linker and the CBD of *Chaetomium thermophilum* Cel7A were synthesized by PCR using following primers:

(forward sequence)
5' - TTAACATATGTTATCTACTCCAACATCAAGGTCGGACCCATCGGCT

CGACCGTCCCTGGCCTTGAC - 3'
and

(reverse sequence)
5' - TATATGCGCCGCAAGCTTTACCATCAAGTTACTCCAGCAAATCAGG

GAACTG - 3'.

[0187] The PCR reaction mixture contained 1× DyNAzyme™ EXT reaction buffer (Finnzymes, Finland), 15 mM Mg²⁺, 0.2 mM dNTPs, 2 μM of each primer, 0.6 units of DyNAzyme™ EXT DNA polymerase (Finnzymes, Finland), and approximately 75 ng/30 μl of template DNA, containing full-length cel7A gene from the *Chaetomium thermophilum*. The conditions for the PCR reaction were the following: 2 min initial denaturation at 98° C., followed by 30 cycles of 30 sec at 98° C., 30 sec annealing at 68° C. (+4° C. gradient), 30 sec extension at 72° C. and a final extension at 72° C. for 10 min. The specific DNA fragment in PCR reaction was obtained at annealing temperature range from 64° C. to 68.5° C. The synthesized CBD fragment of the *Chaetomium thermophilum* was ligated after *Thermoascus aurantiacus* cel7A gene resulting in a junction point of GPIGST between the domains. The PCR amplified fragment in the plasmid was confirmed by sequencing (SEQ ID. NO: 29). The constructed fusion cel7A gene was exactly fused to the *T. reesei* cbh1 (cel7A) promoter. The transcription termination was ensured by the *T. reesei* cel7A terminator and the *A. nidulans* amdS marker gene was used for selection of the transformants as described in Paloheimo et al. (2003).

[0188] The linear expression cassette was isolated from the vector backbone after NotI digestion and was transformed to *T. reesei* A96 protoplasts. The transformations were performed as in Penttila et al. (1987) with the modifications described in Karhunen et al. (1993), selecting with acetamide as a sole nitrogen source. The transformants were purified on selection plates through single conidia prior to sporulating them on PD.

[0189] *Thermoascus aurantiacus* Cel7A+CBD (SEQ ID. NO: 28 and 30) production of the transformants was analyzed from the culture supernatants of the shake flask cultivations (50 ml). The transformants were grown for 7 days in a complex cellulase-inducing medium (Joutsjoki et al. 1993) buffered with 5% KH₂PO₄ at pH 5.5. The cellobiohydrolase activity was assayed using 4-methylumbelliferyl-β-D-lactoside (MUL) substrate according to van Tilbeurgh et al., 1988. The

genotypes of the chosen transformants were confirmed by using Southern blots in which several genomic digests were included and the expression cassette was used as a probe. The SDS-PAGE analyses showed that the recombinant *Thermoplasma aurantiacus* Cel7A+CBD enzymes were produced as stable fusion proteins in *T. reesei*.

[0190] The chosen transformant producing the Ta Cel7A+Tr CBD fusion protein (SEQ ID. NO: 28) was also cultivated in 2 litre bioreactor at 28° C. in the medium indicated above for 3-4 days with pH control 4.4±0.2 (NH₃/H₃PO₄) to obtain material for the application tests. The supernatants were recovered by centrifugation and filtering through Seitz-K 150 and EK filters (Pall SeitzSchenk Filtersystems GmbH, Bad Kreuznach, Germany).

Example 16

Comparison of the Michaelis-Menten and Cellobiose Inhibition Constants of Purified Recombinant Cellobiohydrolases

[0191] The Michaelis-Menten and cellobiose inhibition constants were determined from the cellobiohydrolases produced heterologously in *T. reesei* (Examples 14 and 15). The enzymes were purified as described in Example 2. Protein concentrations of purified enzymes were measured by their absorption at 280 nm using a theoretical molar extinction co-efficient, which were calculated from the amino acid sequences (Gill and von Hippel, 1989).

[0192] Kinetic constants (Km and kcat values) and cellobiose inhibition constant (Ki) for Tr CBHI/Cel7A, Ta CBH/Cel7A, At CBH/Cel7A and Ct CBH/Cel7A, were measured using CNPLac (2-Chloro-4-nitrophenyl-β-D-lactoside) as substrate at ambient temperature (22° C.) in 50 mM sodium phosphate buffer, pH 5.7. For the determination of the inhibition constant (Ki), eight different substrate concentrations (31-4000 μM) in the presence of a range of five inhibitor concentrations (0-100 μM or 0-400 μM), which bracket the K_i value, were used. All experiments were performed in microtiter plates and the total reaction volume was 200 μl. The initial rates were in each case measured by continuous monitoring the release of the chloro-nitrophenolate anion (CNP, 2-Chloro-4-nitrophenolate) through measurements at 405 nm using Varioscan (Thermolabsystems) microtiter plate reader. The results were calculated from CNP standard curve (from 0 to 100 μM). Enzyme concentrations used were: Tr CBHI/Cel7A 2.46 μM, Ta CBH/Cel7A 1.58 μM, Ct CBH/Cel7A 0.79 μM and At CBH/Cel7A 3 μM. The Km and kcat constants were calculated from the fitting of the Michaelis-Menten equation using the programme of Origin. Lineweaver-Burk plots, replots (LWB slope versus [Glc₂; cellobiose]) and Hanes plots were used to distinguish between competitive and mixed type inhibition and to determine the inhibition constants (Ki).

[0193] The results from the kinetic measurements are shown in Table 12 and Table 13. As can be seen, Ct CBH/Cel7A has clearly the higher turnover number (kcat) on CNPLac and also the specificity constant (kcat/Km) is higher as compared to CBHI/Cel7A of *T. reesei*. Cellobiose (Glc₂) is a competitive inhibitor for all the measured cellulases, and the Tr CBHI/Cel7A (used as a control) has the strongest inhibition (i.e. the lowest K_i value) by cellobiose. The At CBH/Cel7A had over 7-fold higher inhibition constant as compared to that of Tr CBHI/Cel7A. These results indicate that all three novel cellobiohydrolases could work better on cellulose

hydrolysis due to decreased cellobiose inhibition as compared to *Trichoderma reesei* Cel7A cellobiohydrolase I.

TABLE 12

Comparison of the cellobiose inhibition constants of four GH family 7 cellobiohydrolases, measured on CNPLac in 50 mM sodium phosphate buffer pH 5.7, at 22° C.		
Enzyme	Ki (μM)	Type of inhibition
Ct Cel7A	39	competitive
Ta Cel7A	107	competitive
At Cel7A	141	competitive
Tr Cel7A	19	competitive

TABLE 13

Comparison of the Michaelis-Menten kinetic constants of <i>Chaetomium thermophilum</i> cellobiohydrolase Cel7A to CBHI/Cel7A of <i>T. reesei</i> , measured on CNPLac in 50 mM sodium phosphate buffer pH 5.7, at 22° C.			
Enzyme	kcat (min ⁻¹)	Km (μM)	kcat/Km (min ⁻¹ M ⁻¹)
Ct Cel7A	18.8	1960	9.5 103
Tr Cel7A	2.6	520	5.0 103

Example 17

Hydrolysis of Crystalline Cellulose (Avicel) by the Recombinant Cellobiohydrolases

[0194] The purified recombinant cellobiohydrolases Ct Cel7A, Ta Cel7A, Ta Cel7A+ TrCBD, Ta Cel7A+ CtCBD, At Cel7A as well as the core version of Ct Cel7A (see below) were tested in equimolar amounts in crystalline cellulose hydrolysis at two temperatures, 45° C. and 70° C.; the purified *T. reesei* Tr Cel7A and its core version (see below) were used as comparison. The crystalline cellulose (Ph 101, Avicel; Fluka, Buchs, Switzerland) hydrolysis assays were performed in 1.5 ml tube scale 50 mM sodium acetate, pH 5.0. Avicel was shaken at 45° C. or at 70° C., with the enzyme solution (1.4 μM), and the final volume of the reaction mixture was 325 μl. The hydrolysis was followed up to 24 hours taking samples at six different time points and stopping the reaction by adding 163 μl of stop reagent containing 9 vol of 94% ethanol and 1 vol of 1 M glycine (pH 11). The solution was filtered through a Millex GV13 0.22 μm filtration unit (Millipore, Billerica, Mass., USA). The formation of soluble reducing sugars in the supernatant was determined by parahydroxybenzoic-acidhydrazide (PAHBAH) method (Lever, 1972) using a cellobiose standard curve (50 to 1600 μM cellobiose). A freshly made 0.1 M PAHBAH (Sigma-Aldrich, St. Louis, Mo., USA) in 0.5 M NaOH (100 μl) solution was added to 150 μl of the filtered sample and boiled for 10 minutes after which the solution was cooled on ice. The absorbance of the samples at 405 nm was measured.

[0195] The core versions of the cellobiohydrolases harboring a CBD in their native form were obtained as follows: Ct Cel7A and Tr Cel7A were exposed to proteolytic digestion to remove the cellulose-binding domain. Papain (Papaya Latex, 14 U/mg, Sigma) digestion of the native cellobiohydrolases

was performed at 37° C. for 24 h in a reaction mixture composed of 10 mM L-cystein and 2 mM EDTA in 50 mM sodium acetate buffer (pH 5.0) with addition of papain (two papain concentrations were tested: of one fifth or one tenth amount of papain of the total amount of the Cel7A in the reaction mixture). The resultant core protein was purified with DEAE Sepharose FF (Pharmacia, Uppsala, Sweden) anion exchange column as described above. The product was analysed in SDS-PAGE.

[0196] The hydrolysis results at 45° C. and 70° C. are shown in FIG. 4 and FIG. 5, respectively. The results show clearly that all the cellobiohydrolases show faster and more complete hydrolysis at both temperatures as compared to the state-of-art cellobiohydrolase *T. reesei* Cel7A. At 70° C. the thermostable cellobiohydrolases from *Thermoascus aurantiacus* ALKO4242 and *Chaetomium thermophilum* ALKO4265 are superior as compared to the *T. reesei* Cel7A, also in the case where the *Thermoascus Cel7A* core is linked to the CBD of *T. reesei* Cel7A (Ta Cel7A+Tr CBD). It was surprising that the cellobiohydrolases isolated and cloned in this work are superior, when harboring a CBD, in the rate and product formation in crystalline cellulose hydrolysis also at the conventional hydrolysis temperature of 45° C. when compared to the state-of-art cellobiohydrolase *T. reesei* Cel7A (CBHI) at the same enzyme concentration. The results are also in agreement with those enzyme preparations (At Cel7A and Ct Cel7A), which were purified from the original hosts and tested in Avicel hydrolysis (50° C., 24 h) (Example 2, Table 1).

Example 18

Cloning of *Acremonium thermophilum* ALKO4245, *Chaetomium thermophilum* ALKO4261, and *Thermoascus aurantiacus* ALKO4242 Endoglucanase Genes

[0197] Standard molecular biology methods were used as described in Example 13. The construction of the *Acremonium*, *Chaetomium*, and *Thermoascus* genomic libraries has been described in Example 12.

[0198] The peptides derived from the purified *Acremonium* and *Chaetomium* endoglucanases shared homology with several endoglucanases of glycosyl hydrolase family 45 such as *Melanocarpus albomyces* Cel45A endoglucanase (AJ515703) and *Humicola insolens* endoglucanase (A35275), respectively. Peptides derived from the *Thermoascus endoglucanase* shared almost 100% identity with the published *Thermoascus aurantiacus* EG1 endoglucanase sequence (AF487830). To amplify a probe for screening of the *Acremonium* and *Chaetomium* genomic libraries, degenerate primers were designed on the basis of the peptide sequences. The order of the peptides in the protein sequence and the corresponding sense or anti-sense nature of the primers was deduced from the comparison with the homologous published endoglucanases. Primer sequences and the corresponding peptides are listed in Table 14. Due to almost 100% identity of the *Thermoascus* peptides with the published sequence, the endoglucanase gene was amplified by PCR directly from the genomic DNA.

TABLE 14

Oligonucleotides synthesized and used as PCR primers to amplify a probe for screening of <i>Acremonium thermophilum</i> cel45A (EG_40) and <i>Chaetomium thermophilum</i> cel7B (EG_54) gene from the corresponding genomic libraries.		
Protein	Peptide	Primer location ^(a) Primer sequence ^(b)
At	EG_40 Peptide 5	1-6 TAYTGGGAYTGYTGAAARCC WFQNAADN ^(c) RTTRTCNGCRTTYTGRAACCA
Ct	EG_54 Peptide 7	3-7 GCAAGCTTCGRCARAARTCRT CRTT ^(d)
	Peptide 2	5-9 GGAATTCGAYCARACNGARCA RTA ^(e)

^(a)Amino acids of the peptide used for designing the primer sequence

^(b)N = A, C, G, or T; R = A or G; Y = C or T

^(c)Peptide not derived from the purified *Acremonium* EG_40 protein, but originates from the *M. albomyces* Cel45A sequence (AJ515703) homologous to EG 40.

^(d)A Hind III restriction site was added to the 5' end of the oligonucleotide

^(e)An EcoRI restriction site was added to the 5' end of the oligonucleotide

[0199] The *Acremonium thermophilum* cel45A gene specific probe to screen the genomic library was amplified with the forward (TAYTGGGAYTGYTGAAARCC) and reverse (RTTRTCNGCRTTYTGRAACCA) primers using genomic DNA as a template. The PCR reaction mixtures contained 50 mM Tris-HCl, pH 9.0, 15 mM (NH₄)₂SO₄, 0.1% Triton X-100, 1.5 mM MgCl₂, 0.1 mM dNTPs, 0.5 µg each primer, 1 unit of Dynazyme EXT DNA polymerase (Finnzymes, Finland) and approximately 0.5 µg of *Acremonium* genomic DNA. The conditions for PCR reactions were the following: 5 min initial denaturation at 95° C., followed by 30 cycles of 1 min at 95° C., 1 min annealing at 50-60° C., 2 min extension at 72° C. and a final extension at 72° C. for 10 min. For amplification of the *Chaetomium thermophilum* cel7B gene (coding for Ct EG_54) specific probe, a forward primer (GGAATTCGAYCARACNGARCART) and a reverse primer (GCAAGCTTCGRCARAARTCRTT) were used. The PCR reaction mixtures contained 10 mM Tris-HCl, pH 8.8, 50 mM KCl, 0.1% Triton X-100, 1.5 mM MgCl₂, 0.2 mM dNTPs, 250 µmol each primer, 2 unit of Dynazyme II DNA polymerase (Finnzymes, Finland) and approximately 2 µg of *Chaetomium* genomic DNA. The conditions for PCR reaction were as described above, except that annealing was performed at 45-50° C.

[0200] Two PCR products were obtained from the *Acremonium* PCR reaction. DNA fragments of about 0.6 kb and 0.8 kb were isolated from agarose gel and were cloned into the pCR4-TOPO® TA vector (Invitrogen, USA) resulting in plasmids pALK1710 and pALK1711, respectively. The DNA products were characterized by sequencing and by performing Southern blot hybridizations to the genomic *Acremonium* DNA digested with several restriction enzymes. The hybridization patterns obtained with the two fragments in stringent washing conditions suggest that two putative endoglucanase genes could be screened from the *Acremonium* genomic library. The deduced amino acid sequences of both PCR products have homology to several published endoglucanase sequences of glycosyl hydrolase family 45 (BLAST program, National Center for Biotechnology Information; Altschul et al., 1990).

[0201] One PCR product of expected size (estimated from the homologous *Humicola insolens* endoglucanase sequence,

A35275) was obtained from the Chaetomium PCR reaction. This DNA fragment of about 0.7 kb was cloned into the pCR4-TOPO® TA vector (Invitrogen, USA) resulting in plasmid pALK2005 and analyzed as described above. The deduced amino acid sequence of the PCR product has homology to several published cellulase sequences of glycosyl hydrolase family 7 (BLAST program, version 2.2.9 at NCBI, National Center for Biotechnology Information; Altschul et al., 1990).

[0202] The insert from plasmids pALK1710, pALK1711, and pALK2005 was isolated by restriction enzyme digestion and labeled with digoxigenin according to the supplier's instructions (Roche, Germany). About $1-2 \times 10^5$ plaques from the amplified *Acremonium* or *Chaetomium* genomic library were screened. The temperature for hybridisation was 68° C. and the filters were washed 2×5 min at RT using 2×SSC-0.1% SDS followed by 2×15 min at 68° C. using 0.1×SSC-0.1% SDS. Several positive plaques were obtained, of which five to six strongly hybridizing plaques were purified from each screening. Phage DNAs were isolated and analysed by Southern blot hybridization. Restriction fragments hybridizing to the probe were subcloned into the pBluescript II KS+vector (Stratagene, USA) and the relevant parts were sequenced. In all cases the subcloned phage fragment contains the full-length gene of interest. Table 15 summarises the information of the probes used for screening of the endoglucanase genes, phage clones from which the genes were isolated, chosen restriction fragments containing the full-length genes with their promoter and terminator regions, names of plasmids containing the subcloned phage fragment, and the deposit numbers in the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH culture collection (DSM) for *E. coli* strains carrying these plasmids.

[0203] *Thermoascus aurantiacus* cel5A gene (coding for EG_28) (SEQ ID NO: 9) was amplified directly from the isolated genomic DNA by PCR reaction. The forward (AT-TAACCGCGGACTGCGCATCAT-GAAGCTCGGCTCTCTCGTGCTC) and reverse (AACT-GAGGCATAGAACTGACGTCATATT) primers that were used for amplification were designed on the basis of the published *T. aurantiacus* eg1 gene (AF487830). The PCR reaction mixtures contained 1× Phusion HF buffer, 0.3 mM dNTPs, 0.5 μM of each primer, 2 units of Phusion™ DNA polymerase (Finnzymes, Finland) and approximately 0.25 μg of *Thermoascus* genomic DNA. The conditions for PCR reactions were the following: 5 min initial denaturation at 95° C., followed by 25 cycles of 30 s at 95° C., 30 s annealing at 57-67° C., 2.5 min extension at 72° C. and a final extension at 72° C. for 5 min. The amplified 1.3 kb product containing the exact gene (from START to STOP codon) was cloned as a SacII-PstI fragment into the pBluescript II KS+vector. Two independent clones were sequenced and one clone was selected and designated as pALK1926. The deposit number of the *E. coli* strain containing pALK1926 in the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH culture collection is DSM 17326.

[0204] Relevant information of the genes and the deduced protein sequences (SEQ ID NO: 9-16) are summarized in Table 16 and Table 17, respectively. Peptide sequences of the purified *Acremonium* EG-40 (gene At cel45A), *Chaetomium* EG-54 (gene Ct cel7B), and *Thermoascus* EG_28 (gene Ta cel5A) endoglucanases were found in the corresponding deduced amino acid sequences of the cloned genes confirming that appropriate genes were cloned.

TABLE 15

Probes used for cloning of endoglucanase gene, phage clone and the subclone chosen, plasmid name and the corresponding deposit number of the <i>E. coli</i> strain.						
Gene	Genomic library	Probe used in screening	Phage clone	Subcloned fragment	Plasmid	<i>E. coli</i> deposit no.
At cel45A	<i>A. thermophilum</i> ALKO4245	pALK1710	P24	5.5 kb SmaI	pALK1908	DSM 17324
At cel45B	<i>A. thermophilum</i> ALKO4245	pALK1711	P41	6.0 kb XhoI	pALK1904	DSM 17323
Ct cel7B	<i>C. thermophilum</i> ALKO4261	pALK2005	P55	5.1 kb BamHI	pALK2010	DSM 17729

TABLE 16

Summary of the endoglucanase genes isolated from <i>Acremonium thermophilum</i> , <i>Chaetomium thermophilum</i> , and <i>Thermoascus aurantiacus</i> .					
Endoglucanase gene	Length with introns (bp) ^(a)	Coding region (bp) ^(b)	No of introns	Lengths of introns (bp)	SEQ ID NO:
At cel45A	1076	891	2	59, 123	11
At cel45B	1013	753	2	155, 102	13
Ct cel7B	1278	1275	—	—	15
Ta cel5A	1317	1005	5	55, 60, 59, 74, 61	9

^(a)The STOP codon is included.

^(b)The STOP codon is not included.

TABLE 17

Summary of the deduced endoglucanase sequences of <i>Acremonium thermophilum</i> , <i>Chaetomium thermophilum</i> , and <i>Thermoascus aurantiacus</i> .							
Endoglucanase protein	No of aas	Length of ss NN/HMM ^(a)	CBD ^(b)	Predicted MW (Da, ss not incl) ^(c)	Predicted pI (ss not incl)	Putative N-glycosylation sites ^(d)	SEQ ID NO:
At EG_40	297	21/21	Yes, K265 to L297	28625	4.79	2	12
At EG_40_like	251	20/20	No	23972	6.11	2	14
Ct EG_54	425	17/17	No	45358	5.44	1	16
Ta EG_28	335	30(e)	No	33712	4.30	1	10

ss, signal sequence.

^(a)The prediction of the signal sequence was made using the program SignalP V3.0 (Nielsen et al., 1997; Bendtsen et al., 2004); the NN value was obtained using neural networks and HMM value using hidden Markov models.

^(b)Presence of a cellulose binding domain in the protein, the amino acids of the C-terminal CBD are indicated (numbering according to the full length polypeptide)

^(c)The predicted signal sequence is not included. Prediction was made using the Compute pI/MW tool at ExPASy server (Gasteiger et al., 2003).

^(d)The putative N-glycosylation sites N-X-S/T were predicted using the program NetNGlyc 1.0 (Gupta et al., 2004).

^(e)According to Hong et al. 2003a

[0205] The deduced protein sequences of *Acremonium* EG_40 (At Cel45A) and EG_40_like (At Cel45B), *Chaetomium* EG_54 (Ct Cel7B), and *Thermoascus* EG_28 (Ta Cel5A) endoglucanases share homology with cellulases of glycosyl hydrolase family 45 (*Acremonium*), family 7 (*Chaetomium*), and family 5 (*Thermoascus*), thus identifying the isolated genes as members of these gene families. The closest homologies of the *Acremonium* endoglucanases EG_40/Cel45A and EG_40_like/Cel45B are endoglucanases of *Thielavia terrestris* (CQ827970, 77.3% identity) and *Myceliophthora thermophila* (AR094305, 66.9% identity), respectively (Table 18). The two isolated *Acremonium* family 45 endoglucanases share only an identity of 53.7% with each other. Of these enzymes only EG-40/Cel45A contains a cellulose binding domain (CBD).

[0206] The closest homology for the predicted protein sequence of *Chaetomium* EG_54/Cel7B endoglucanase is found in the *Melanocarpus albomyces* Cel7A cellulase sequence (AJ515704). The identity between these two protein sequences is 70.6%.

[0207] The protein sequence of the isolated *Thermoascus aurantiacus* endoglucanase is completely identical with that

of the published *T. aurantiacus* EGI (AF487830, Table 18). The closest homology was found in a β -glucanase sequence of *Talaromyces emersonii* (AX254752, 71.1% identity).

TABLE 18

Comparison of the deduced <i>Acremonium thermophilum</i> EG_40, EG_40_like/Cel45B, <i>Chaetomium thermophilum</i> EG_54/Cel7B, and <i>Thermoascus aurantiacus</i> EG_28/Cel5A endoglucanases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.	
Organism, enzyme, and accession number	Identity (%)
<i>Acremonium thermophilum</i> EG_40	100.0
<i>Thielavia terrestris</i> EG45, CQ827970	77.3
<i>Melanocarpus albomyces</i> Cel45A, AJ515703	75.3
<i>Neurospora crassa</i> , hypothetical XM_324477	68.9

TABLE 18-continued

Comparison of the deduced <i>Acremonium thermophilum</i> EG_40, EG_40_like/Cel45B, <i>Chaetomium thermophilum</i> EG_54/Cel7B, and <i>Thermoascus aurantiacus</i> EG_28/Cel5A endoglucanases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.	
Organism, enzyme, and accession number	Identity (%)
<i>Humicola grisea</i> var <i>thermoidea</i> , EGL3, AB003107	67.5
<i>Humicola insolens</i> EG5, A23635	67.3
<i>Myceliophthora thermophila</i> fam 45, AR094305	57.9
* <i>Acremonium thermophilum</i> EG_40_like	53.7
<i>Acremonium thermophilum</i> EG_40_like	100.0
<i>Myceliophthora thermophila</i> fam 45, AR094305	66.9
<i>Magnaporthe grisea</i> 70-15 hypothetical XM_363402	61.9
<i>Thielavia terrestris</i> EG45, CQ827970	56.8
* <i>Acremonium thermophilum</i> EG_40	56.8
<i>Melanocarpus albomyces</i> Cel45A, AJ515703	53.7
	52.8
<i>Chaetomium thermophilum</i> EG_54	100.0
<i>Melanocarpus albomyces</i> Cel7A, AJ515704	70.6
<i>Humicola grisea</i> var <i>thermoidea</i> EGI, D63516	68.8
<i>Humicola insolens</i> EGI, AR012244	67.7

TABLE 18-continued

Comparison of the deduced *Acremonium thermophilum* EG_40, EG_40_like/Cel45B, *Chaetomium thermophilum* EG_54/Cel7B, and *Thermoascus aurantiacus* EG_28/Cel5A endoglucanases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.

Organism, enzyme, and accession number	Identity (%)
<i>Myceliophthora thermophila</i> EGI, AR071934	61.7
<i>Fusarium oxysporum</i> var <i>lycopercisi</i> EGI, AF29210	53.5
<i>Fusarium oxysporum</i> EGI, AR012243	52.6
<i>Thermoascus aurantiacus</i> EG_28	100.0
<i>Thermoascus aurantiacus</i> EG, AX812161	100.0
<i>Thermoascus aurantiacus</i> EGI, AY055121	99.4
<i>Talaromyces emersonii</i> β -glucanase, AX254752	71.1
<i>Talaromyces emersonii</i> EG, AF440003	70.4
<i>Aspergillus niger</i> EG, A69663	70.1
<i>Aspergillus niger</i> EG, A62441	69.9
<i>Aspergillus niger</i> EG, AF331518	69.6
<i>Aspergillus aculeatus</i> EGV, AF054512	68.5

*indicates an endoglucanase encoded by a gene cloned in this work.

Example 19

Production of Recombinant Endoglucanases in *Trichoderma reesei*

[0208] Expression plasmids were constructed for production of the recombinant *Acremonium* EG_40/Cel45A, EG_40_like/Cel45B, and *Thermoascus* EG_28/Cel5A proteins as described in Example 14. Linear expression cassettes (Table 19) were isolated from the vector backbone by restriction enzyme digestion, transformed into *T. reesei* A96 and transformants purified as described in Example 14.

TABLE 19

The expression cassettes constructed for production of *Acremonium thermophilum* EG_40/Cel45A, EG_40_like/Cel45B, and *Thermoascus aurantiacus* EG_28/Cel5A endoglucanases in *Trichoderma reesei*. The schematic structure of the expression cassettes is described in FIG. 2.

Endoglucanase	Expression plasmid	Size of the expression cassette ^(a)	Heterologous terminator ^(b)
At EG_40	pALK1920	10.9 kb NotI	156 bp (HindIII)
At EG_40_like	pALK1921	8.6 kb EcoRI	282 bp (SspI)
Ta EG_28	pALK1930	8.6 kb NotI	none

^(a)The expression cassette for *T. reesei* transformation was isolated from the vector backbone by EcoRI or NotI digestion.

^(b)The number of nucleotides after the STOP codon of the cloned gene that are included in the expression cassette are indicated. The restriction site at the 3'-region of the gene that was used in construction of the expression cassette is indicated in parenthesis.

[0209] The endoglucanase production of the transformants was analyzed from the culture supernatants of shake flask cultivations (50 ml). Transformants were grown as in Example 14 and the enzyme activity of the recombinant protein was measured from the culture supernatant as the release of reducing sugars from carboxymethylcellulose (2% (w/v) CMC) at 50° C. in 50 mM citrate buffer pH 4.8 essentially as described by Bailey and Nevalainen 1981; Haakana et al. 2004. Production of the recombinant proteins was also detected from culture supernatants by SDS-polyacrylamide gel electrophoresis. *Acremonium* EG_40-specific polyclonal

antibodies were produced in rabbits (University of Helsinki, Finland). The expression of EG_40 was verified by Western blot analysis with anti-EG_40 antibodies using the ProtoBlot Western blot AP system (Promega). The genotypes of the chosen transformants were analysed by Southern blotting using the expression cassette as a probe.

[0210] The pH optimum of the heterologously produced endoglucanases was determined in the universal McIlvaine's buffer within a pH range of 4.0-8.0 using carboxymethylcellulose as substrate. As shown in FIG. 6 A the broadest pH range (4.5-6.0) is that of the *Acremonium* EG_40/Cel45A protein, the optimum being at pH 5.5. The pH optima for the other heterologously produced endoglucanases are pH 5.0-5.5 and 6.0 for *Acremonium* EG_40_like/Cel45B and *Thermoascus* EG_28/Cel5A, respectively. The optimal temperature for enzymatic activity of these endoglucanases was determined at the temperature range of 50-85° C. as described above. The highest activity of the enzymes was determined to be at 75° C., 60° C., and 75° C. for the *Acremonium* EG_40/Cel45A, EG_40_like/Cel45B, and *Thermoascus* EG_28/Cel5A, respectively (FIG. 6 B).

[0211] The chosen transformants were cultivated, as described in Example 14, in a 2 litre bioreactor for four days (28° C., pH 4.2) to obtain material for the application tests.

Example 20

Cloning of *Acremonium thermophilum* ALKO4245, *Chaetomium thermophilum* ALKO4261, and *Thermoascus aurantiacus* ALKO4242 Beta-Glucosidase Genes

[0212] Standard molecular biology methods were used as described in Example 13. The construction of the *Acremonium*, *Chaetomium*, and *Thermoascus* genomic libraries has been described in Example 12.

[0213] The peptides derived from the purified *Acremonium*, *Chaetomium*, and *Thermoascus* β -glucosidases shared homology with several β -glucosidases of glycosyl hydrolase family 3 such as *Acremonium cellulolyticus* (BD168028), *Trichoderma viride* (AY368687), and *Talaromyces emersonii* (AY072918) β -glucosidases, respectively. To amplify a probe for screening of the *Acremonium*, *Chaetomium*, or *Thermoascus* genomic libraries, degenerate primers were designed on the basis of the peptide sequences. The order of the peptides in the protein sequence and the corresponding sense or anti-sense nature of the primers was deduced from the comparison with the homologous published β -glucosidases. Primer sequences and the corresponding peptides are listed in Table 20.

TABLE 20

Oligonucleotides synthesized and used as PCR primers to amplify a probe for screening of *Acremonium thermophilum* cel3A (β G_101), *Chaetomium thermophilum* cel3A (β G_76), and *Thermoascus aurantiacus* cel3A (β G_81) gene from the corresponding genomic libraries.

Protein	Peptide	Primer location ^(a)	Primer Sequence ^(b)
At β G_101	EKVNL ^(c) Peptide 4 6-11		GARAARGTNAAYCTNAC YTTRCCRTTRTTSGGRGTR TA

TABLE 20-continued

Oligonucleotides synthesized and used as PCR primers to amplify a probe for screening of <i>Acremonium thermophilum</i> cel3A (β G_101), <i>Chaetomium thermophilum</i> cel3A (β G_76), and <i>Thermoascus aurantiacus</i> cel3A (β G_81) gene from the corresponding genomic libraries.			
Protein	Peptide	Primer location ^(a)	Primer Sequence ^(b)
Ct β G_76	Peptide 6	4-9	TNTGYCTNCARGAYGG
	Peptide 1	3-8	TCRAARTGSCGRTARTCRA TRAASAG
Ta β G_81	Peptide 3	1-5	AARGGYGTSGAYGTSCAR
	Peptide 1	2-7	YTTRCCCCASGTRAASGG

^(a)Amino acids of the peptide used for designing the primer sequence

^(b)To reduce degeneracy, some codons were chosen according to fungal preference. N = A, C, G, or T; R = A or G; S = C or G; Y = C or T

^(c)Peptide not derived from the purified *Acremonium* β G_101 protein, but originates from the *A. cellulolyticus* β -glucosidase sequence (BD168028) homologous to β G_101.

[0214] The probes for screening genomic libraries constructed were amplified with the listed primer combinations (Table 20) using *Acremonium*, *Chaetomium*, or *Thermoascus* genomic DNA as template. The PCR reaction mixtures contained 50 mM Tris-HCl, pH 9.0, 15 mM (NH₄)₂SO₄, 0.1% Triton X-100, 1.5 mM MgCl₂, 0.1-0.2 mM dNTPs, 0.25 μ g each primer, 1 unit of Dynazyme EXT DNA polymerase (Finnzymes, Finland) and approximately 0.5 μ g of genomic DNA. The conditions for PCR reactions were the following: 5 min initial denaturation at 95° C., followed by 30 cycles of 1 min at 95° C., 1 min annealing at 40° C. (*Acremonium* DNA as a template), at 50° C. (*Chaetomium* DNA as a template), or at 63° C. (*Thermoascus* DNA as a template), 2-3 min extension at 72° C. and a final extension at 72° C. for 5-10 min.

[0215] Specific PCR products of expected size (estimated from the homologous β -glucosidase sequences BD168028,

AY072918, and AY368687) were isolated from the agarose gel. DNA fragments of about 1.8 kb (*Acremonium*), 1.5 kb (*Chaetomium*), and 1.52 kb (*Thermoascus*) were cloned into the pCR4-TOPO® TA vector (Invitrogen, USA) resulting in plasmids pALK1924, pALK1935, and pALK1713, respectively. The DNA products were characterized by sequencing and by performing Southern blot hybridizations to the genomic DNA digested with several restriction enzymes. The hybridization patterns in stringent washing conditions suggest that one putative β -glucosidase gene could be isolated from the *Acremonium*, *Chaetomium*, and *Thermoascus* genomic library. The deduced amino acid sequences of all three PCR products have homology to several published β -glucosidase sequences of glycosyl hydrolase family 3 (BLAST program, National Center for Biotechnology Information; Altschul et al., 1990).

[0216] The insert from plasmids pALK1713, pALK1924, and pALK1935 was isolated by restriction enzyme digestion and labeled with digoxigenin according to the supplier's instructions (Roche, Germany). About 1-2 \times 10⁵ plaques from the amplified *Acremonium*, *Chaetomium*, or *Thermoascus* genomic library were screened as described in Example 18. Several positive plaques were obtained, of which five to six strongly hybridizing plaques were purified from each screening. Phage DNAs were isolated and analysed by Southern blot hybridization. Restriction fragments hybridizing to the probe were subcloned into the pBluescript II KS+vector (Stratagene, USA) and the relevant parts were sequenced. In all cases the subcloned phage fragment contains the full-length gene of interest. Table 21 summarises the information of the probes used for screening of the β -glucosidase genes, phage clones from which the genes were isolated, chosen restriction fragments containing the full-length genes with their promoter and terminator regions, names of plasmids containing the subcloned phage fragment, and the deposit numbers in the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH culture collection (DSMZ) for *E. coli* strains carrying these plasmids.

TABLE 21

Probes used for cloning of β -glucosidase gene, phage clone and the subclone chosen, plasmid name and the corresponding deposit number of the <i>E. coli</i> strain.						
Gene	Genomic library	Probe			Plasmid	<i>E. coli</i> deposit no.
		used in screening	Phage clone	Subcloned fragment		
At cel3A	<i>A. thermophilum</i> ALKO4245	pALK1924	P44	6.0 kb HindIII	pALK1925	DSM 17325
Ct cel3A	<i>C. thermophilum</i> ALKO4261	pALK1935	P51	7.0 kb XbaI	pALK2001	DSM 17667
Ta cel3A	<i>T. aurantiacus</i> ALKO4242	pALK1713	P21	5.3 kb BamHI	pALK1723	DSM 16725

[0217] Relevant information of the genes and deduced protein sequences (SEQ ID NO: 21-26) are summarized in Table 22 and Table 23, respectively. Peptide sequences of the purified *Acremonium* β G_101 (At Cel3A), *Chaetomium* β G_76 (Ct Cel3A), and *Thermoascus* β G_81 (Ta Cel3A) proteins were found in the corresponding deduced amino acid sequences of the cloned genes confirming that appropriate genes were cloned.

TABLE 22

Summary of the β -glucosidase genes isolated from <i>Acremonium thermophilum</i> , <i>Chaetomium thermophilum</i> , and <i>Thermoascus aurantiacus</i> .					
β -glucosidase gene	Length with introns (bp) ^(a)	Coding region bp ^(b)	No of introns	Lengths of introns (bp)	SEQ ID NO:
At cel3A	2821	2583	3	92, 74, 69	23
Ct cel3A	2257	2202	1	52	25
Ta cel3A	3084	2529	7	134, 67, 56, 64, 59, 110, 62	21

^(a)The STOP codon is included.

^(b)The STOP codon is not included.

TABLE 23

Summary of the deduced β -glucosidase sequences of <i>Acremonium thermophilum</i> , <i>Chaetomium thermophilum</i> , and <i>Thermoascus aurantiacus</i> .							
β -glucosidase protein	No of aas	Length of ss NN/HMM ^(a)	CBD ^(b)	Predicted MW (Da, ss not incl) ^(c)	Predicted pl ss not incl)	Putative N-glycosylation sites ^(d)	SEQ ID NO:
At β G_101	861	19/18	No	91434	5.46	8	24
Ct β G_76	734	20/20	No	76457	6.3	2	26
Ta β G_81	843	19/19	No	89924	4.95	8	22

ss, signal sequence.

^(a)The prediction of the signal sequence was made using the program SignalP V3.0 (Nielsen et al., 1997; Bendtsen et al., 2004); the NN value was obtained using neural networks and HMM value using hidden Markov models.

^(b)Presence of a cellulose binding domain in the protein.

^(c)The predicted signal sequence is not included. Prediction was made using the Compute pl/MW tool at ExPASy server (Gasteiger et al., 2003).

^(d)The putative N-glycosylation sites N-X-S/T were predicted using the program NetNGlyc 1.0 (Gupta et al., 2004).

[0218] The deduced protein sequences of *Acremonium* β G-101/Cel3A, *Chaetomium* β G_76/Cel3A, and *Thermoascus* β G_81/Cel3A β -glucosidases share homology with enzymes of glycosyl hydrolase family 3, thus identifying that the isolated genes belong to this gene family. The closest counterparts of the *Acremonium*, *Chaetomium*, and *Thermoascus* β -glucosidases are those of *Magnaporthe grisea* (β -glucosidase, AY849670), *Neurospora crassa* (hypothetical, XM_324308), and *Talaromyces emersonii* (β -glucosidase, AY072918), respectively (Table 24). The highest sequence identity (73.2%) found was that of *C. thermophilum* β G_76/Cel3A to *N. crassa* hypothetical protein indicating that novel enzymes genes were cloned.

TABLE 24

Comparison of the deduced <i>Acremonium thermophilum</i> β G_101/Cel3A, <i>Chaetomium thermophilum</i> β G_76/Cel3A, and <i>Thermoascus aurantiacus</i> β G_81/Cel3A β -glucosidases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.	
Organism, enzyme, and accession number	Identity (%)
* <i>Acremonium thermophilum</i> β G_101	100.0
<i>Magnaporthe grisea</i> β -glucosidase, AY849670	73.1
<i>Neurospora crassa</i> hypothetical, XM_330871	71.1
<i>Trichoderma reesei</i> Cel3B, AY281374	65.2
* <i>Thermoascus aurantiacus</i> β G_81	62.2
<i>Aspergillus aculeatus</i> β -glucosidase, D64088	59.5
<i>Talaromyces emersonii</i> β -glucosidase, AY072918	58.9
<i>Aspergillus oryzae</i> , AX616738	58.2
<i>Acremonium cellulolyticus</i> β -glucosidase, BD168028	57.2
* <i>Chaetomium thermophilum</i> β G_76	40.9
<i>Chaetomium thermophilum</i> β G_76	100.0
<i>Neurospora crassa</i> , hypothetical XM_324308	76.9
<i>Magnaporthe grisea</i> , hypothetical XM_364573	70.2
<i>Trichoderma viridae</i> BGI, AY368687	65.8
<i>Acremonium cellulolyticus</i> β -glucosidase, BD168028	41.2
* <i>Acremonium thermophilum</i> β G_101	40.9
<i>Trichoderma reesei</i> Cel3B, AY281374	40.0

TABLE 24-continued

Comparison of the deduced <i>Acremonium thermophilum</i> β G_101/Cel3A, <i>Chaetomium thermophilum</i> β G_76/Cel3A, and <i>Thermoascus aurantiacus</i> β G_81/Cel3A β -glucosidases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.	
Organism, enzyme, and accession number	Identity (%)
* <i>Thermoascus aurantiacus</i> β G_81	39.9
* <i>Thermoascus aurantiacus</i> β G_81	100.0
<i>Talaromyces emersonii</i> β -glucosidase, AY072918	73.2
<i>Aspergillus oryzae</i> , AX616738	69.5
<i>Aspergillus aculeatus</i> β -glucosidase, D64088	68.0
<i>Acremonium cellulolyticus</i> β -glucosidase, BD168028	65.7

TABLE 24-continued

Comparison of the deduced *Acremonium thermophilum* β G_101/Cel3A, *Chaetomium thermophilum* β G_76/Cel3A, and *Thermoascus aurantiacus* β G_81/Cel3A β -glucosidases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.

Organism, enzyme, and accession number	Identity (%)
* <i>Acremonium thermophilum</i> β G_101	62.2
<i>Trichoderma reesei</i> Cel3B, AY281374	57.9
* <i>Chaetomium thermophilum</i> β G_76	39.9

*indicates a β -glucosidase encoded by a gene cloned in this work.

Example 21

Production of Recombinant Beta-Glucosidases in
Trichoderma reesei

[0219] Expression plasmids were constructed for production of the recombinant *Acremonium* β G_101/Cel3A, *Chaetomium* β G_76/Cel3A, and *Thermoascus* β G_81/Cel3A proteins as described in Example 14. Linear expression cassettes (Table 25) were isolated from the vector backbone by restriction enzyme digestion, transformed into *T. reesei* A96 or A33 (both strains have the genes encoding the four major cellulases CBHI/Cel7A, CBHII/Cel6A, EGI/Cel7B and EGII/Cel5A deleted) and transformants purified as described in Example 14.

TABLE 25

The expression cassettes constructed for production of *Acremonium thermophilum* β G_101/Cel3A, *Chaetomium thermophilum* β G_76/Cel3A, and *Thermoascus aurantiacus* β G_81/Cel3A β -glucosidases in *Trichoderma reesei*. The schematic structure of the expression cassettes is described in FIG. 2.

β -glucosidase	Expression plasmid	Size of the expression cassette ^(a)	Heterologous terminator ^(b)
At β G_101	pALK1933	10.5 kb NotI	300 bp (HindIII)
Ct β G_76	pALK2004	10.1 kb EcoRI	528 bp (XbaI)
Ta β G_81	pALK1914	10.9 kb EcoRI	452 bp (ApoI)

^(a)The expression cassette for *T. reesei* transformation was isolated from the vector backbone by EcoRI or NotI digestion.

^(b)The number of nucleotides after the STOP codon of the cloned gene that are included in the expression cassette are indicated. The restriction site at the 3'-region of the gene that was used in construction of the expression cassette is indicated in parenthesis.

[0220] The beta-glucosidase production of the transformants was analyzed from the culture supernatants of shake flask cultivations (50 ml). Transformants were grown as in Example 14 and the enzyme activity of the recombinant protein was measured from the culture supernatant using 4-nitrophenyl- β -D-glucopyranoside substrate as described by Bailey and Nevalainen 1981. Production of the recombinant proteins was also detected from culture supernatants by SDS-polyacrylamide gel electrophoresis. In addition, the expression of *Thermoascus* β G_81 was verified by Western blot analysis with anti- β G_81 antibodies as described in Example 19. The genotypes of the chosen transformants were analysed by Southern blotting using the expression cassette as a probe.

[0221] The pH optimum of the heterologously produced β -glucosidases was determined in the universal McIlvaine's buffer within a pH range of 3.0-8.0 using 4-nitrophenyl- β -D-glucopyranoside as substrate. The pH optima for the *Acremonium* β G_101, *Chaetomium* β G_76, and *Thermoascus* β G_81 are pH 4.5, 5.5, and 4.5, respectively (FIG. 7 A). The optimal temperature for enzymatic activity of these β -glucosidases was determined at the temperature range of 50-85° C. as described above. The highest activity of the enzymes was determined to be at 70° C., 65° C., and 75° C. for the *Acremonium* β G_101/Cel3A, *Chaetomium* β G_76/Cel3A, and *Thermoascus* β G_81/Cel3A, respectively (FIG. 7 B).

[0222] The chosen transformants were cultivated, as described in Example 14, in a 2 litre bioreactor for four days (28° C., pH 4.2) to obtain material for the application tests.

Example 22

Cloning of *Acremonium thermophilum* ALKO4245
and *Thermoascus aurantiacus* ALKO4242 Xylanase
Genes

[0223] Standard molecular biology methods were used as described in Example 13. The construction of the *Acremonium* genomic library has been described in Example 12.

[0224] The peptides derived from the purified *Acremonium* xylanase shared homology with xylanases of the glycosyl hydrolase family 10 such as *Humicola grisea* XYNI (AB01030). All peptides derived from the *Thermoascus* xylanase were completely identical with the published *Thermoascus aurantiacus* XYNA sequence (AJ132635) thus identifying the purified protein as the same enzyme. Due to this the *Thermoascus* xylanase gene was amplified by PCR from the genomic DNA.

[0225] To amplify a probe for screening of the *Acremonium* xylanase gene from the genomic library, degenerate primers were designed on the basis of the peptide sequences (Example 11, Table 5). The order of the peptides in the protein sequence and the corresponding sense or antisense nature of the primers was deduced from the comparison with the homologous *Humicola insolens* XYNI sequence (AB001030). The sense primer sequence (GAYGGYGA-YGCSACYTAYATG) is based on Peptide 3 (amino acids 2-8) and anti-sense primer (YTTYTGRTCRTAYTCSAGRT-TRTA) on Peptide 1 (amino acids 4-11).

[0226] A PCR product of expected size (estimated from the homologous *Humicola insolens* XYNI sequence AB001030) was obtained from the reaction. This DNA fragment of about 0.7 kb was cloned into the pCR4-TOPO® TA vector (Invitrogen, USA) resulting in plasmid pALK1714, and was characterized by sequencing. The deduced amino acid sequence of the PCR product has homology to several published xylanase sequences of glycosyl hydrolase family 10 (BLAST program, National Center for Biotechnology Information; Altschul et al., 1990).

[0227] The insert from plasmid pALK1714 was isolated by restriction enzyme digestion and labeled with digoxigenin according to the supplier's instructions (Roche, Germany). About $1-2 \times 10^5$ plaques from the amplified *Acremonium* genomic library were screened as described in Example 18. Several positive plaques were obtained, of which five strongly hybridizing plaques were purified. Phage DNAs were isolated and analysed by Southern blot hybridization. A 3.0 kb XbaI restriction fragment hybridizing to the probe was subcloned into the pBluescript II KS+ vector (Stratagene,

USA) resulting in plasmid pALK1725. Relevant parts of pALK1725 were sequenced and found to contain the full-length *Acremonium thermophilum* xyn10A gene (SEQ ID NO: 19). The deposit number of the *E. coli* strain containing pALK1725 in the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH culture collection is DSM 16726.

[0228] *Thermoascus aurantiacus* xyn10A gene (SEQ ID NO: 17) was amplified directly from the isolated genomic DNA by PCR reaction. The forward (TTATACCGCGG-GAAGCCATGGTTCGACCAACGATCCTAC) and reverse (TTATAGGATCCACCGGTCTATACTCACT-GCTGCAGGTCCTG) primers that were used in the amplification of the gene were designed on the basis of the published *T. aurantiacus* xynA gene (AJ132635). The PCR reaction mixtures contained 50 mM Tris-HCl, pH 9.0, 15 mM

TABLE 26

Summary of the xylanase genes isolated from <i>Acremonium thermophilum</i> and <i>Thermoascus aurantiacus</i> .					
Xylanase gene	Length with introns (bp) ^(a)	Coding region (bp) ^(b)	No of introns	Lengths of introns (bp)	SEQ ID NO:
At xyn10A	1471	1248	2	135, 85	19
Ta xyn10A	1913	987	10	73, 74, 68, 103, 69, 65, 93, 66, 100, 212	17

^(a)The STOP codon is included.

^(b)The STOP codon is not included.

TABLE 27

Summary of the deduced xylanase sequences of <i>Acremonium thermophilum</i> and <i>Thermoascus aurantiacus</i>							
Xylanase protein	No of aas	Length of ss NN/HMM ^(a)	CBD ^(b)	Predicted MW (Da, ss not incl) ^(c)	Predicted pl (ss not incl)	Putative N-glycosylation sites ^(d)	SEQ ID NO:
At XYN_60	416	19/19	Yes, W385 to L416	42533	6.32	1-2	20
Ta XYN_30	329	26(e)	No	32901	5.81	0	18

ss, signal sequence.

^(a)The prediction of the signal sequence was made using the program SignalP V3.0 (Nielsen et al., 1997; Bendtsen et al., 2004); the NN value was obtained using neural networks and HMM value using hidden Markov models.

^(b)Presence of a carbohydrate binding domain CBD, the amino acids of the C-terminal CBD are indicated (numbering according to the full length polypeptide)

^(c)The predicted signal sequence is not included. Prediction was made using the Compute pl/MW tool at ExPASy server (Gasteiger et al., 2003).

^(d)The putative N-glycosylation sites N-X-S/T were predicted using the program NetNGlyc 1.0 (Gupta et al., 2004).

^(e)According to Lo Leggio et al., 1999

(NH₄)₂SO₄, 0.1% Triton X-100, 1.5 mM MgCl₂, 0.3 mM dNTPs, 1 μM each primer, 1 unit of Dynazyme EXT DNA polymerase (Finnzymes, Finland) and approximately 0.5 μg of *Thermoascus* genomic DNA. The conditions for PCR reactions were the following: 5 min initial denaturation at 95° C., followed by 30 cycles of 1 min at 95° C., 1 min annealing at 60-66° C., 3 min extension at 72° C. and a final extension at 72° C. for 10 min. The amplified 1.9 kb product containing the exact gene (from START to STOP codon) was cloned as a SacII-BamHI fragment into the pBluescript II KS+vector. Three independent clones were sequenced and one clone was selected and designated as pALK1715. The deposit number of the *E. coli* strain containing pALK1715 in the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH culture collection is DSM 16724.

[0229] Relevant information of the genes and deduced protein sequences (SEQ ID NO: 17-20) are summarized in Table 26 and Table 27, respectively. Peptide sequences of the purified *Acremonium* XYN_60 and *Thermoascus* XYN_30 proteins were found in the corresponding deduced amino acid sequences of the cloned genes (At xyn10A and Ta xyn10A, respectively) confirming that appropriate genes were cloned.

[0230] The deduced protein sequences of *Acremonium* and *Thermoascus* xylanases share homology with several enzymes of glycosyl hydrolase family 10, identifying the corresponding genes as members of family 10 xylanases. The closest counterpart for the *Acremonium* XYN-60/Xyn10A found is the *Humicola grisea* XYLI (AB001030) showing 67.1% identity with XYN_60 (Table 28). The predicted protein sequence of the isolated *Thermoascus aurantiacus* XYN_30/Xyn10A xylanase is completely identical with that of the published *T. aurantiacus* XYNA (P23360, Table 28). The closest homology was found in a xylanase sequence of *Aspergillus niger* (A62445, 69.7% identity).

TABLE 28

Comparison of the deduced <i>Acremonium thermophilum</i> XYN_60/Xyn10A and <i>Thermoascus aurantiacus</i> XYN_30/Xyn10A xylanases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.	
Organism, enzyme, and accession number	Identity (%)
* <i>Thermoascus aurantiacus</i> XYN_30	100.0
<i>Thermoascus aurantiacus</i> XynA, P23360	100.0

TABLE 28-continued

Comparison of the deduced *Acremonium thermophilum* XYN_60/Xyn10A and *Thermoascus aurantiacus* XYN_30/Xyn10A xylanases with their homologous counterparts. The alignment was performed using the Needle programme of the EMBOSS programme package.

Organism, enzyme, and accession number	Identity (%)
<i>Thermoascus aurantiacus</i> XynA, AF127529	99.4
<i>Aspergillus niger</i> xylanase, A62445	69.7
<i>Aspergillus aculeatus</i> xylanase, AR137844	69.9
<i>Aspergillus terreus</i> fam 10 xyn, DQ087436	65.0
<i>Aspergillus sojae</i> , XynXI AB040414	63.8
<i>Penicillium chrysogenum</i> xylanase, AY583585	62.5
* <i>Acremonium thermophilum</i> XYN_60	100.0
<i>Humicola grisea</i> XYL I, AB001030	67.1
<i>Magnaporthe grisea</i> 70-15, hypothetical XM_364947	63.8
<i>Aspergillus aculeatus</i> xylanase, AR149839	53.7
<i>Talaromyces emersonii</i> xylanase, AX403831	51.8
<i>Gibberella zeae</i> xylanase, AY575962	51.4
<i>Magnaporthe grisea</i> XYL5, AY144348	48.5
<i>Talaromyces emersonii</i> , AX172287	46.9

*indicates a xylanase encoded by a gene cloned in this work.

Example 23

Production of Recombinant Xylanases in *Trichoderma reesei*

[0231] Expression plasmids were constructed for production of the recombinant *Acremonium* XYN-60/Xyn10A and *Thermoascus* XYN_30/Xyn10A proteins as described in Example 14. Linear expression cassettes (Table 29) were isolated from the vector backbone by restriction enzyme digestion, transformed into *T. reesei* A96, and transformants purified as described in Example 14.

TABLE 29

The expression cassettes constructed for production of *Acremonium thermophilum* XYN_60/Xyn10A and *Thermoascus aurantiacus* XYN_30/Xyn10A xylanases in *Trichoderma reesei*. The schematic structure of the expression cassettes is described in FIG. 2.

Xylanase	Expression plasmid	Size of the expression cassette ^a	Heterologous terminator ^b
At XYN_60	pALK1912	9.0 kb	150 bp (BamHI)
Ta XYN_30	pALK1913	9.3 kb	none

^aThe expression cassette for *T. reesei* transformation was isolated from the vector backbone by EcoRI digestion.

^bThe number of nucleotides after the STOP codon of the cloned gene that are included in the expression cassette are indicated. The restriction site at the 3'-region of the gene that was used in construction of the expression cassette is indicated in parenthesis.

[0232] The xylanase production of the transformants was analyzed from the culture supernatants of shake flask cultivations (50 ml). Transformants were grown as in Example 14 and the enzyme activity of the recombinant protein was measured from the culture supernatant as the release of reducing sugars from birch xylan (1% w/v) at 50° C. in 50 mM citrate buffer pH 5.3 as described by Bailey and Poutanen 1989. Production of the recombinant protein was also analyzed from culture supernatant by SDS-polyacrylamide gel electrophoresis. In addition, the expression of both xylanases was determined by Western blot analysis with anti-XYN_30 or

anti-XYN_60 antibodies as described in Example 19. The genotypes of the chosen transformants were analysed by Southern blotting using the expression cassette as a probe.

[0233] *Thermoascus* XYN-30/Xyn10A was produced in *T. reesei* and the pH optimum of the heterologously produced protein was determined in the universal McIlvaine's buffer within a pH range of 3.0-8.0 using birch xylan as substrate (FIG. 8 A). The optimal pH was determined to be 4.5. The temperature optimum for the enzymatic activity of XYN_30 was determined to be 75° C. (FIG. 8 B).

[0234] The chosen transformants were cultivated, as described in Example 14, in a 2 litre bioreactor for four days (28° C., pH 4.2) to obtain material for the application tests.

Example 24

Performance of the Recombinant Cellobiohydrolases in the Hydrolysis

[0235] The performance of the purified recombinant cellobiohydrolases was evaluated in the hydrolysis studies with purified *T. reesei* enzymes. Hydrolysis was carried out with controlled mixtures of purified enzymes on several pre-treated substrates. Culture filtrates of *T. reesei*, containing different cloned CBH/Cel7 enzymes were obtained as described in Examples 14 and 15, and the CBH enzymes were purified by affinity chromatography as described in Example 2. In addition, pure *T. reesei* cellulases (purified as described by Suurnakki et al., 2000) were used in the enzyme mixtures. The cellobiohydrolases used in the experiment were:

[0236] *Thermoascus aurantiacus* ALKO4242 CBH (Ta Cel7A)

[0237] *Thermoascus aurantiacus* ALKO4242 CBH (Ta Cel7A) with genetically attached CBD of *Trichoderma reesei* (Ta Cel7A+Tr CBD)

[0238] *Thermoascus aurantiacus* ALKO4242 CBH (Ta Cel7A) with genetically attached CBD of *Chaetomium thermophilum* (Ta Cel7A+Ct CBD)

[0239] *Acremonium thermophilum* ALKO4245 CBH (At Cel7A)

[0240] *Chaetomium thermophilum* ALKO4265 CBH (Ct Cel7A).

[0241] Each CBH/Cel7 to be tested (dosage 14.5 mg/g dry matter of substrate) was used either together with EGII/Cel5A of *T. reesei* (3.6 mg/g) or with a mixture containing *T. reesei* EGI/Cel7B (1.8 mg/g), EGII/Cel5A (1.8 mg/g), xylanase pl 9 (Tenkanen et al. 1992) (5000 nkat/g) and acetyl xylan esterase (AXE) (Sundberg and Poutanen, 1991) (250 nkat/g). All mixtures were supplemented with additional β -glucosidase from a commercial enzyme preparation Novozym 188 (176 nkat/g d.w.). Triplicate tubes containing the enzyme mixture and 10 mg (dry matter)/ml of the substrate suspended in 0.05 M sodium acetate were incubated in mixing by magnetic stirring at 45° C. for 48 h. Reference samples with inactivated enzymes and corresponding substrates were also prepared. The release of hydrolysis products was measured as reducing sugars with DNS method using glucose as standard (Table 30).

[0242] The following substrates were used in the experiment:

[0243] Crystalline cellulose (Avicel)

[0244] Washed steam pre-treated spruce fibre (impregnation with 3% w/w SO₂ for 20 min, followed by steam pre-treatment at 215° C. for 5 min), dry matter 25.9% (SPRUCE).

[0245] Washed wet oxidized corn stover fibre (WOCS).

[0246] Washed steam pre-treated willow fibre (pre-treatment for 14 min at 210° C.), dry matter 23.0% (WILLOW).

philum Cel7A and *C. thermophilum* Cel7A. Considering also the better temperature stability of the herein produced cello-

TABLE 30

Hydrolysis products with CBH enzymes (45° C., pH 5.0). Reaction products after 48 h hydrolysis as reducing sugars (mg/ml), measured glucose as standard.					
Enzymes		Substrates			
CBH	Additional enzymes	Avicel	SPRUCE	WOCS	WILLOW
Ta Cel7A	EGII, bG	2.0	2.0	2.8	2.0
Ta Cel7A + Tr CBD	EGII, bG	5.8	4.0	4.4	4.0
Ta Cel7A + Ct CBD	EGII, bG	4.9	3.7	4.6	3.7
At Cel7A	EGII, bG	5.3	3.3	4.5	3.3
Ct Cel7A	EGII, bG	6.0	2.6	3.4	2.6
Cel7A of <i>T. reesei</i>	EGII, bG	4.7	2.9	2.9	2.9
Ta Cel7A	EGII, EGI, XYL, AXE, bG	nd	nd	4.3	2.8
Ta Cel7A + Tr CBD	EGII, EGI, XYL, AXE, bG	nd	nd	7.2	5.9
Ta Cel7A + Ct CBD	EGII, EGI, XYL, AXE, bG	nd	nd	7.2	5.6
At Cel7A	EGII, EGI, XYL, AXE, bG	nd	nd	6.4	5.4
Ct Cel7A	EGII, EGI, XYL, AXE, bG	nd	nd	5.6	4.0
Cel7A of <i>T. reesei</i>	EGII, EGI, XYL, AXE, bG	nd	nd	6.0	4.1

Abbreviations:

CBH = cellobiohydrolase;

EGI = endoglucanase I (Cel7B) of *T. reesei*,

EGII = endoglucanase II (Cel5A) of *T. reesei*;

bG = β -glucosidase (from Novozym 188);

XYL = xylanase pl 9 (XYN II) of *T. reesei*,

AXE = acetyl xylan esterase of *T. reesei*;

nd = not done.

[0247] In Table 30 the different cellobiohydrolases have been compared based on the same protein dosage in the hydrolysis. The results show that on cellulosic substrates (Avicel and spruce fibre) Cel7A of *Thermoascus aurantiacus* with genetically attached CBD showed clearly higher hydrolysis than *T. reesei* CBHI/Cel7A. Without CBD, *T. aurantiacus* Cel7A was less efficient on these substrates. The performance of *Acremonium thermophilum* and *Chaetomium thermophilum* cellobiohydrolases was also better than that of *T. reesei* CBHI/Cel7A on several substrates; in particular, *C. thermophilum* Cel7A showed high efficiency on pure cellulose (Avicel).

[0248] In the case of substrates containing notable amounts of hemicellulose (willow and corn stover) the CBH/Cel7 enzymes clearly needed additionally both hemicellulases and endoglucanases to perform efficiently. If no additional hemicellulases were present, Cel7A of *T. aurantiacus* with genetically attached CBD showed again clearly highest hydrolysis. With the most important hemicellulose-degrading enzymes (xylanase, acetyl xylan esterase and EGI) Cel7A of *T. aurantiacus* with genetically attached CBD performed again with highest efficiency. *A. thermophilum* Cel7A was more efficient than *T. reesei* enzyme and *C. thermophilum* Cel7A produced hydrolysis products on the same level than *T. reesei* CBHI/Cel7A. The cellulose binding domain of *T. reesei* seemed to give slightly better efficiency than CBD of *C. thermophilum* in the hydrolytic performance of *T. aurantiacus* Cel7A, even though the difference was rather small.

[0249] It can be concluded that when CBHI/Cel7A was replaced in the mixture of *Trichoderma* enzymes by the herein produced cellobiohydrolases, the hydrolysis efficiency as judged by this experimental arrangements was clearly improved in the case of *T. aurantiacus* Cel7A with genetically attached CBD, and also improved in the case of *A. thermo-*

biohydrolases, the results indicate that the performance of cellulase enzyme mixtures in higher temperatures than 45° C. can be clearly improved by using the herein produced cellobiohydrolases.

Example 25

Performance of the Recombinant Endoglucanases in the Hydrolysis

[0250] The preparations containing the endoglucanases were compared in hydrolysis studies mixed with the purified CBH/Cel7 and CBH/Cel6 enzymes on several pre-treated substrates. Culture filtrates of *T. reesei*, containing different cloned endoglucanase enzymes were obtained as described in Example 19. The enzymes were enriched by removing thermolabile proteins from the mixtures by a heat treatment (60° C., 2 h, pH 5) and the supernatant was used for the hydrolysis studies. In addition, pure *T. reesei* cellulases (purified as described by Suurnakki et al., 2000) were used in the enzyme mixtures. The endoglucanases used in the experiment were:

[0251] *Acremonium thermophilum* ALKO4245 endoglucanase At EG_40/Cel45A (ALKO4245 EG_40)

[0252] *Acremonium thermophilum* ALKO4245 endoglucanase At EG_40_like/Cel45B (ALKO4245 EG_40_like)

[0253] *Thermoascus aurantiacus* ALKO4242 endoglucanase Ta EG_28/Cel5A (ALKO4242 EG_28).

[0254] The following substrates were used in the experiment:

[0255] Washed steam pre-treated spruce fibre (impregnation with 3% SO₂ for 20 min, followed by steam pre-treatment at 215° C. for 5 min), dry matter 25.9% (SPRUCE).

[0256] Steam exploded corn stover fibre (steam pre-treatment at 21° C. for 5 min), dry matter 31.0% (SECS).

[0257] The endoglucanases to be studied (dosage 840 nkat/g dry matter, based on endoglucanase activity against HEC according to IUPAC, 1987) were used either with cellobiohydrolases of *T. reesei* (CBHI/Cel7A, 8.1 mg/g d.m. and CBHII/Cel6A, 2.0 mg/g d.m.) or with *Thermoascus aurantiacus* Cel7A with genetically attached CBD of *T. reesei* (10.1 mg/g d.m.). Purified (Suurnakki et al., 2000) EGI (Cel7B) and EGII (Cel5A) of *T. reesei* were also included in the experiments for comparison. All mixtures were supplemented with additional β -glucosidase from Novozym 188 (to make the total β -glucosidase dosage 560 nkat/g d.w., the relatively high dosage was used to compensate the differences in the background activities of the different EG preparations). Triplicate tubes were incubated in mixing at 45° C. for 48 h and reference samples with inactivated enzymes and corresponding substrates were prepared. The release of hydrolysis products was measured as reducing sugars with DNS method using glucose as standard (Table 31).

TABLE 31

Hydrolysis products with different endoglucanase preparations when used together with cellobiohydrolases from <i>T. reesei</i> or with <i>T. aurantiacus</i> Cel7A harbouring CBD of <i>T. reesei</i> . Reaction products after 48 h hydrolysis (45° C., pH 5.0) as reducing sugars (mg/ml), measured glucose as standard.			
Enzymes		Substrate	
Endoglucanase	CBH/Cel7	SPRUCE	SECS
no added EG	CBHI and CBHII of <i>T. reesei</i>	2.4	3.2
EGI	CBHI and CBHII of <i>T. reesei</i>	3.5	4.6
EGII	CBHI and CBHII of <i>T. reesei</i>	3.8	3.5
At EG_40	CBHI and CBHII of <i>T. reesei</i>	4.9	4.3
At EG_40like	CBHI and CBHII of <i>T. reesei</i>	4.5	4.8
Ta EG_28	CBHI and CBHII of <i>T. reesei</i>	3.0	3.9
no added EG	<i>T. aurantiacus</i> Cel7A + Tr CBD	1.8	2.1
EG I	<i>T. aurantiacus</i> Cel7A + Tr CBD	nd.	4.2
EG II	<i>T. aurantiacus</i> Cel7A + Tr CBD	3.2	nd.
At EG_40	<i>T. aurantiacus</i> Cel7A + Tr CBD	4.8	4.0
Ta EG_28	<i>T. aurantiacus</i> Cel7A + Tr CBD	1.5	nd.

Abbreviations:

CBHI = cellobiohydrolase I (Cel7A) of *T. reesei*;
 CBHII = cellobiohydrolase II (Cel6A) of *T. reesei*;
 EGI = endoglucanase I (Cel7B) of *T. reesei*;
 EGII = endoglucanase II (Cel5A) of *T. reesei*;
 bG = β -glucosidase (from Novozym 188);
 nd. = not done.

[0258] In Table 31 the different endoglucanases have been compared based on the same activity dosage in the hydrolysis. This may favour enzymes with low specific activity against the substrate (hydroxyethyl cellulose) used in the assay and underestimate the efficiency of enzymes with high specific activity against hydroxyethyl cellulose. In any case, the results show that *Acremonium thermophilum* endoglucanases perform very well in the hydrolysis when affecting together with both cellobiohydrolases used in the mixture. *A. thermophilum* endoglucanases have similar performance to *T. reesei* EGI/Cel7B which is a very efficient enzyme on hemicellulose-containing corn stover substrate due to its strong xylanase side activity. *T. aurantiacus* endoglucanase Cel5A (ALKO4242 EG_28) showed lower hydrolysis than *T. reesei* enzymes.

[0259] It can be concluded that the endoglucanases from *A. thermophilum* perform with comparable or enhanced efficiency when compared to the corresponding *Trichoderma* enzymes in the hydrolysis as judged by this experimental arrangement. Considering also the temperature stability of

the herein described endoglucanases, the results indicate that the performance of cellulase enzyme mixtures in higher temperatures than 45° C. can be improved by using the herein described endoglucanases.

Example 26

Hydrolysis of Steam Pre-Treated Spruce at High Temperatures

[0260] Washed steam exploded spruce fibre (impregnation with 3% w/w SO₂ for 20 min, followed by steam pre-treatment at 215° C. for 5 min), with dry matter of 25.9% was suspended in 5 ml of 0.05 M sodium acetate buffer in the consistency of 10 mg/ml. This substrate was hydrolysed using different enzyme mixtures in test tubes with magnetic stirring in the water bath adjusted in different temperatures for 72 h. For each sample point, a triplicate of test tubes was withdrawn from hydrolysis, boiled for 10 min in order to terminate the enzyme hydrolysis, centrifuged, and the supernatant was analysed for reaction products from hydrolysis. The blanks containing the substrate alone (only buffer added instead of enzymes) were also incubated in the corresponding conditions.

[0261] A mixture of thermophilic cellulases was prepared using the following components:

[0262] Thermophilic CBH/Cel7 preparation containing *Thermoascus aurantiacus* ALKO4242 Cel7A with genetically attached CBD of *T. reesei* CBHI/Cel7A. The protein preparation was produced as described in Example 15 and purified according to Example 2 resulting in the purified Ta Cel7A+Tr CBD preparation with protein content of 5.6 mg/ml.

[0263] Thermophilic endoglucanase preparation containing *Acremonium thermophilum* ALKO4245 endoglucanase At EG_40/Cel45A. The protein was produced in *T. reesei* as described in Example 19. In order to enrich the thermophilic components, the spent culture medium was heat treated (60° C. for 2 hours). The preparation obtained contained protein 4.9 mg/ml and endoglucanase activity (according to IUPAC, 1987) 422 nkat/ml.

[0264] Thermophilic β -glucosidase preparation prepared as described in Example 21 containing *Thermoascus aurantiacus* ALKO4242 β -glucosidase Ta β G_81/Cel3A. In order to enrich the thermophilic components, the fermentor broth was heat treated (65° C. for 2 hours). The preparation obtained contained 4.3 mg/ml protein and β -glucosidase activity of 6270 nkat/ml (according to Bailey and Linko, 1990).

[0265] These enzyme preparations were combined as follows (per 10 ml of mixture): CBH/Cel7-preparation 4.51 ml, endoglucanase preparation 5.19 ml and β -glucosidase preparation 0.29 ml. This mixture was used as "MIXTURE 1" of the thermophilic enzymes.

[0266] As a comparison and reference, a state-of art mixture of commercial *Trichoderma reesei* enzymes was constructed combining (per 10 ml): 8.05 ml Celluclast 1.5 L FG (from Novozymes A/S) and 1.95 ml Novozym 188 (from Novozymes A/S). This was designated as "*T. Reesei* ENZYMES."

[0267] Enzymes were dosed on the basis of the FPU activity of the mixtures: "MIXTURE 1" using the dosage of 5.5 FPU per 1 gram of dry matter in the spruce substrate, and "*T. Reesei* ENZYMES" using 5.8 FPU per 1 gram of dry matter in the spruce substrate.

[0268] Samples were taken from the hydrolysis after 24, 48 and 72 h and treated as described above. The hydrolysis products were quantified using the assay for reducing sugars (Bernfeld, 1955), using glucose as standard. The amount of hydrolysis products as reducing sugars is presented in FIG. 9.

[0269] The results clearly show better performance of the herein described enzymes as compared to the state-of-art *Trichoderma* enzymes in 55° C. and 60° C. on the spruce substrate. On the basis of HPLC analysis the maximum yield of sugars from the substrate would be 5.67 mg per 10 mg of dry spruce substrate. Because of the relatively low dosage of enzyme the final sugar yields were clearly lower. For thermostable enzymes the sugar yield based on reducing sugar assay was 66% and 57% of theoretical in 55° C. and 60° C., respectively. For state-of art *Trichoderma* enzymes it was only 31% and 11% in 55° C. and 60° C., respectively.

Example 27

Hydrolysis of Steam Pre-Treated Corn Stover at High Temperatures

[0270] Steam exploded corn stover fibre (treatment at 195° C. for 5 min), with dry matter of 45.3% was suspended in 5 ml of 0.05 M sodium acetate buffer in the consistency of 10 mg/ml. The treatments and measurements were performed as described in Example 26.

[0271] A mixture of herein described thermophilic cellulases was constructed using the following components:

[0272] Thermophilic CBH preparation containing *Thermoascus aurantiacus* ALKO4242 Cel7A with genetically attached CBD of *T. reesei* CBHI/Cel7A (Ta Cel7A+ Tr CBD, Example 15). The protein content of the preparation was 31 mg/ml.

[0273] Thermophilic endoglucanase preparation containing *Acremonium thermophilum* ALKO4245 endoglucanase At EG_40/Cel45A was obtained as described in Example 19. The concentrated enzyme preparation contained endoglucanase activity (according to IUPAC, 1987) of 2057 nkat/ml.

[0274] Thermophilic β -glucosidase preparation containing *Thermoascus aurantiacus* ALKO 4242 β -glucosidase Ta β G_81/Cel3A was obtained as described in Example 21 containing β -glucosidase activity (according to Bailey and Linko, 1990) of 11500 nkat/ml.

[0275] Thermophilic xylanase product containing an AM24 xylanase originating from *Nonomuraea flexuosa* DSM43186. The product was prepared by using a recombinant *Trichoderma reesei* strain that had been transformed with the expression cassette pALK1502, as described in WO2005/100557. The solid product was dissolved in water to make a 10% solution and an enzyme preparation with xylanase activity (assayed according to Bailey et al., 1992) of 208000 nkat/ml was obtained.

[0276] These enzyme preparations were combined as follows (per 10 ml of mixture): CBH/Cel7 preparation 7.79 ml, endoglucanase preparation 0.96 ml, β -glucosidase preparation 1.14 ml and xylanase preparation 0.31 ml. This mixture was used as "MIXTURE 2" of the thermophilic enzymes.

[0277] As a comparison and reference, a state-of art mixture of commercial *Trichoderma reesei* enzymes was constructed by combining (per 10 ml) 8.05 ml Celluclast 1.5 L FG (from Novozymes A/S) and 1.95 ml Novozym 188 (from Novozymes A/S). This was designated as "*T. Reesei* ENZYMES."

[0278] Samples were taken from the hydrolysis after 24, 48 and 72 h and treated as described above. The hydrolysis products were quantified using the assay for reducing sugars (Bernfeld, 1955), using glucose as standard. The results from the substrate blanks were subtracted from the samples with enzymes, and the concentration of hydrolysis products as reducing sugars is presented in FIG. 10.

[0279] The results clearly show better performance of the herein described enzymes as compared to the state-of-art *Trichoderma* enzymes. In 45° C. the mixture of thermophilic enzymes showed more efficient hydrolysis as compared to *T. reesei* enzymes: The hydrolysis was faster and higher sugar yields were also obtained. On the basis of HPLC analysis the maximum yield of sugars (including free soluble sugars in the unwashed substrate that was used) from the substrate would be 5.73 mg per 10 mg of dry substrate. Thus, the hydrolysis by the MIXTURE 2 enzymes was nearly complete within 48 hours. In 55° C. and 57.5° C. the herein described thermophilic enzymes showed also clearly better performance in the hydrolysis as compared to the state-of art *Trichoderma* enzymes.

Example 28

Hydrolysis of Pre-Treated Corn Stover at High Temperatures Using Mixture with a Thermostable Xylanase

[0280] The procedure explained in Example 27 was repeated except that the xylanase product XT 02026A3 was replaced by thermophilic xylanase preparation containing *Thermoascus aurantiacus* ALKO4242 xylanase Ta XYN_30/XynI OA produced in *T. reesei*. The fermentor broth, produced as described in Example 23 contained xylanase activity of 132 000 nkat/ml (assayed according to Bailey et al., 1992).

[0281] These enzyme preparations were combined as follows (per 10 ml of mixture): CBH/Cel7-preparation 7.64 ml, endoglucanase preparation 0.96 ml, β -glucosidase preparation 1.15 ml and xylanase preparation 0.25 ml. This mixture was used as "MIXTURE 3" of the thermophilic enzymes.

[0282] As a comparison and reference, a state-of-art mixture of commercial *Trichoderma reesei* enzymes was constructed by combining (per 10 ml) 8.05 ml Celluclast 1.5 L FG (from Novozymes A/S) and 1.95 ml Novozym 188 (from Novozymes A/S). This was designated as "*T. Reesei* ENZYMES."

[0283] Samples were taken from the hydrolysis after 24, 48 and 72 h and treated as described above. The hydrolysis products were quantified using the assay for reducing sugars (Bernfeld, 1955), using glucose as standard. The results from the substrate blanks were subtracted from the samples with enzymes, and the concentration of hydrolysis products as reducing sugars is presented in FIG. 11.

[0284] The results clearly show better performance of the mixture of the herein described enzymes as compared to the state-of-art *Trichoderma* enzymes. In 45° C. the mixture of thermophilic enzymes showed more efficient hydrolysis as compared to *T. reesei* enzymes. In 55° C. and 60° C. the herein described thermophilic enzymes showed clearly better performance in the hydrolysis as compared to the state-of art *Trichoderma* enzymes. The performance of the new enzyme

mixture at 60° C. was at the same level than the performance of state-of-art enzymes at 45° C.

Example 29

Hydrolysis of Pre-Treated Spruce at High Temperatures Using Mixture with a Thermostable Xylanase

[0285] Procedure as described in Example 28 was repeated with washed steam exploded spruce fibre (impregnation with 3% w/w SO₂ for 20 min, followed by steam pre-treatment at 215° C. for 5 min, with dry matter of 25.9%) as substrate using hydrolysis temperatures 45° C., 55° C. and 60° C. Samples were taken from the hydrolysis after 24, 48 and 72 h and treated as described above. The hydrolysis products were quantified using the assay for reducing sugars (Bernfeld, 1955), using glucose as standard. The results from the substrate blanks were subtracted from the samples with enzymes, and the concentration of hydrolysis products as reducing sugars is presented in FIG. 12.

[0286] The results clearly show better performance of the mixture of herein described enzymes as compared to the state-of-art *Trichoderma* enzymes in all the temperatures studied. At 45° C. the mixture of thermophilic enzymes showed more efficient hydrolysis as compared to *T. reesei* enzymes, evidently due to the better stability in long term hydrolysis. At 55° C. the efficiency of the mixture of herein described enzymes was still on the same level than at 45° C., whereas the state-of-art mixture was inefficient with the substrate used in this temperature. At 60° C. the herein described thermophilic enzymes showed decreased hydrolysis although the hydrolysis was nearly at the same level as the performance of the state-of-art enzymes at 45° C.

Example 30

Evaluation of Glucose Inhibition of β -Glucosidases from *Acremonium thermophilum* ALKO4245, *Chaetomium thermophilum* ALKO4261 and *Thermoascus aurantiacus* ALKO4242

[0287] The culture filtrates produced by *Acremonium thermophilum* ALKO4245, *Chaetomium thermophilum* ALKO4261 and *Thermoascus aurantiacus* ALKO4242 strains are described in Example 1. The β -glucosidase activities (measured according to Bailey and Linko, 1990) of these preparations were 21.4 nkat/ml, 5.6 nkat/ml and 18.6 nkat/ml, respectively. For comparison, commercial enzymes Celluclast 1.5 L (β -glucosidase 534 nkat/ml) and Novozym 188 (β -glucosidase 5840 nkat/ml) were also included in the experiment.

[0288] In order to evaluate the sensitivity of the different β -glucosidases towards glucose inhibition, the standard activity assay procedure was performed in the presence of different concentrations of glucose. The substrate (p-nitrophenyl- β -D-glucopyranoside) solutions for β -glucosidase activity assay were supplemented by glucose so that the glucose concentration in the assay mixture was adjusted to the values from 0 to 0.5 M. Except this glucose addition the assay was performed using the standard procedure (Bailey and Linko, 1990). The activities in the presence of varying glucose concentrations as a percentage of the activity without glucose are presented in FIG. 13.

[0289] The results show that β -glucosidases from *C. thermophilum* and *T. aurantiacus* were affected less by glucose inhibition than the β -glucosidases present in the commercial

enzymes: *Aspergillus*-derived β -glucosidase in Novozym 188 or *Trichoderma*-derived β -glucosidase in Celluclast 1.5 L. *A. thermophilum* enzyme showed behaviour comparable to *T. reesei* enzyme of Celluclast. Especially *C. thermophilum* enzyme was clearly less affected by high glucose concentration. Thus, these results indicate that considering glucose inhibition the use of the new β -glucosidases, especially from strains *Acremonium thermophilum* ALKO4242 and *Chaetomium thermophilum* ALKO4261, would give clear advantages in hydrolysis in industrial conditions with high glucose concentration.

Example 31

Filter Paper Activity of Enzyme Mixtures in High Temperatures

[0290] Filter paper activity of enzyme preparations was measured according to the method of IUPAC (1987) as described in the procedure except enzyme reaction was performed at temperatures from 50° C. to 70° C. The calculated FPU activity is based on the amount of enzyme required to hydrolyse 4% of filter paper substrate in 1 h under the experimental conditions. The FPU activity is considered to represent the total overall cellulase activity of an enzyme preparation.

[0291] The enzyme mixtures were MIXTURE 2 prepared as described in Example 27, MIXTURE 3 prepared as described in Example 28, and MIXTURE 4. MIXTURE 4 was prepared by combining enzyme preparations described in Example 27 as follows (per 10 ml of mixture): CBH/Cel7-preparation 7.84 ml, endoglucanase preparation 0.99 ml and β -glucosidase preparation 1.17 ml.

[0292] The enzyme mixtures used as reference, representing the state-of-art-mixtures, were:

[0293] "*T. Reesei* ENZYMES A" prepared as preparation "*T. Reesei* ENZYMES" described in Example 26.

[0294] "*T. Reesei* ENZYMES B" was constructed combining (per 10 ml) 8.05 ml Econase CE (a commercial *T. reesei* cellulase preparation from AB Enzymes Oy, Rajamaki, Finland) and 1.95 ml Novozym 188 (from Novozymes A/S).

[0295] The FPU activities measured for the enzyme preparations at different temperatures are presented in FIG. 14 as percentages of the activity under standard (IUPAC, 1987) conditions (at 50° C.).

[0296] Results clearly show that the mixtures of the invention show higher overall cellulase activity in elevated (60-70°) temperatures as compared to the state-of-art mixtures based on enzymes from *Trichoderma* and *Aspergillus*.

Example 32

Use of the Novel Beta-Glucosidases in Preparation of Sophorose

[0297] A high concentration starch hydrolysate mixture (Nutriose 74/968, Roquette) was treated with *Thermoascus aurantiacus* β G_81/Cel3A enriched enzyme preparation produced as described in Example 21 to produce a sugar mixture containing appreciable amounts of cellulase inducer (sophorose) to overcome the glucose repression.

[0298] The Ta β G_81/Cel3A enriched enzyme preparation was added to a 70% (w/w) Nutriose solution to a final concentration of 1 g total protein/litre. The container of the mixture was incubated in a water bath at 65° C. for 3 days with constant stirring and used as a carbon source in a shake flask

medium for two different *Trichoderma*-strains (A47 and Rut-C30). The effect of the enzyme treatment was measured as an endoglucanase activity formed during a 7 days shake flask cultivation. As a reference cultivations were performed under the same conditions with untreated Nutriose as a carbon source. More than two-fold increase in the activities was obtained in the shake flask cultivations performed on Ta β G₈₁/Cel3A pretreated Nutriose media with the strains tested. Results are shown in FIG. 15.

List of deposited organisms

Strain	Plasmid contained	Deposition authority	Deposition date	Deposition number
<i>Acremonium thermophilum</i> ALKO4245	—	CBS ⁽¹⁾	20 Sep. 2004	CBS 116240
<i>Thermoascus aurantiacus</i> ALKO4242	—	CBS ⁽¹⁾	20 Sep. 2004	CBS 116239
<i>Chaetomium thermophilum</i> ALKO4265	—	CBS ⁽²⁾	Nov. 8, 1995	CBS 730.95 ⁽⁴⁾
<i>Escherichia coli</i>	pALK1635	DSMZ ⁽³⁾	16 Sep. 2004	DSM 16723
<i>Escherichia coli</i>	pALK1642	DSMZ	16 Sep. 2004	DSM 16727
<i>Escherichia coli</i>	pALK1646	DSMZ	16 Sep. 2004	DSM 16728
<i>Escherichia coli</i>	pALK1861	DSMZ	16 Sep. 2004	DSM 16729
<i>Escherichia coli</i>	pALK1715	DSMZ	16 Sep. 2004	DSM 16724
<i>Escherichia coli</i>	pALK1723	DSMZ	16 Sep. 2004	DSM 16725
<i>Escherichia coli</i>	pALK1725	DSMZ	16 Sep. 2004	DSM 16726
<i>Escherichia coli</i>	pALK1904	DSMZ	13 May 2005	DSM 17323
<i>Escherichia coli</i>	pALK1908	DSMZ	13 May 2005	DSM 17324
<i>Escherichia coli</i>	pALK1925	DSMZ	13 May 2005	DSM 17325
<i>Escherichia coli</i>	pALK1926	DSMZ	13 May 2005	DSM 17326
<i>Escherichia coli</i>	pALK2001	DSMZ	18 Oct. 2005	DSM 17667
<i>Escherichia coli</i>	pALK2010	DSMZ	18 Nov. 2005	DSM 17729

⁽¹⁾the Centralbureau Voor Schimmelcultures at Uppsalalaan 8, 3584 CT, Utrecht, the Netherlands

⁽²⁾the Centralbureau Voor Schimmelcultures at Oosterstraat 1, 3742 SK BAARN, The Netherlands

⁽³⁾Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ), Mascheroder Weg 1 b, D-38124 Braunschweig, Germany

⁽⁴⁾[After termination of the current deposit period, samples will be stored under agreements as to make the strain available beyond the enforceable time of the patent.]

REFERENCES

- [0299] Altschul S., Gish W., Miller W., Myers E. W. and Lipman D. J., (1990) "Basic local alignment search tool," *J. Mol. Biol.* 215: 403-410.
- [0300] Badger, P. C. (2002) "Ethanol from cellulose: a general review," In: *Trends in New Crops and New Uses*, J. Janick and A. Whipkey (eds.). ASHS Press, Alexandria, Va., USA, pp. 17-21.
- [0301] Bailey M. J. and K. M. H. Nevalainen (1981) "Induction, isolation and testing of stable *Trichoderma reesei* mutants with improved production of solubilizing cellulose," *Enz. Microbiol. Technol.*, 3:153-157.
- [0302] Bailey, M. J., Biely, P. and Poutanen, K. (1992) "Interlaboratory testing for assay of xylanase activity," *J. Biotechnol.*, 23:257-270.
- [0303] Bailey, M. J. and Linko, M. (1990) "Production of β -galactosidase by *Aspergillus oryzae* in submerged bioreactor cultivation," *J. Biotechnol.*, 16:57-66.
- [0304] Bailey M. J. and Poutanen K. (1989) "Production of xylanases by strains of *Aspergillus*," *Appl. Microbiol. Biotechnol.*, 30:5-10.
- [0305] Bailey M., Siika-aho M., Valkeajärvi A. and Penttilä M. (1993) "Hydrolytic properties of two cellulases of *Trichoderma reesei* expressed in yeast," *Biotechnol. Appl. Biochem.*, 17:65-76.
- [0306] Bendtsen J. D., Nielsen H., von Heijne G. and Brunak S. (2004) "Improved prediction of signal peptides: SignalP 3.0," *J. Mol. Biol.*, 340:783-795.
- [0307] Bernfeld, B. (1955) "Amylases, α and β ," In: *Methods in Enzymology*, vol. 1, Eds. Colowick, S. P. and Kaplan, N. O. Academic Press, New York, pp. 149-158.
- [0308] Biely P., Vrsanska M., Tenkanen M., Kluepfel D. (1997) "Endo-beta-1,4-xylanase families: differences in catalytic properties," *Journal of Biotechnology*, 57:151-166.
- [0309] Coen, D. M. (2001) "The polymerase chain reaction," In: *Current Protocols in Molecular Biology*, Ausubel, F. M., Brent, R., Kingston, R. E., More, D. D., Seidman, J. G., Smith, K. and Struhl, K. (eds.), John Wiley & Sons. Inc., Hoboken, USA.
- [0310] Gasteiger, E., Gattiker A., Hoogland C., Ivanyi I., Appel R. D. and Bairoch A. (2003) "ExpASY: the proteomics server for in-depth protein knowledge and analysis," *Nucleic Acids Res.*, 31:3784-3788.
- [0311] Gellissen, G. (ed.) (2005) "Production of recombinant proteins," *Novel Microbial and Eukaryotic Expression Systems*, Wiley-VCH Verlag GmbH & Co. Weinheim, Germany.
- [0312] Gill, S. C. and von Hippel, P. H. (1989) "Calculation of protein extinction coefficients from amino acid sequence data," *Anal. Biochem.*, 182:319-326.
- [0313] Gupta, R., E. Jung and S. Brunak. (2004) "Prediction of N-glycosylation sites in human proteins," in preparation: www.cbs.dtu.dk/services/NetNGIvc/.
- [0314] Haakana H., Miettinen-Oinonen A., Joutsjoki V., Mäntylä A., Suominen P. and Vehmaanperä J. (2004) "Cloning of cellulase genes from *Melanocarpus albomyces* and their efficient expression in *Trichoderma reesei*," *Enz. Microbiol. Technol.*, 34:159-167.
- [0315] Henrissat B. (1991) "A classification of glycosyl hydrolases based on amino acid sequence similarities," *Biochem. J.*, 280:309-316.
- [0316] Henrissat B. and Bairoch A. (1993) "New families in the classification of glycosyl hydrolases based on amino acid sequence similarities," *Biochem. J.*, 293:781-788.
- [0317] Henrissat B. and Bairoch A. (1996) "Updating the sequence-based classification of glycosyl hydrolases," *Biochem. J.*, 316:695-696.
- [0318] Henrissat B., Teeri T. T. and Warren R. A. J. (1998) "A scheme for designating enzymes that hydrolyse the polysaccharides in the cell wall of plants," *FEBS Letters*, 425:352-354.
- [0319] Hong J., H. Tamaki, K. Yamamoto, and Kumagai H. (2003a) "Cloning of a gene encoding a thermo-stable

- endo- β -1,4-glucanase from *Thermoascus aurantiacus* and its expression in yeast," *Biotech. Letters*, 25:657-661.
- [0320] Hong J., Tamaki H., Yamamoto K. and Kumagai H. (2003b) "Cloning of a gene encoding thermostable cellobiohydrolase from *Thermoascus aurantiacus* and its expression in yeast," *Appl. Microbiol. Biotechnol.*, 63:42-50.
- [0321] IUPAC (International Union of Pure and Applied Chemistry) (1987) "Measurement of cellulase activities," *Pure and Appl. Chem.*, 59:257-268.
- [0322] Joutsjoki, V. V., Torkkeli T. K. and Nevalainen K. M. H. (1993) "Transformation of *Trichoderma reesei* with the *Hormoconis resinae* glucoamylase P (gamP) gene: production of a heterologous glucoamylase by *Trichoderma reesei*," *Curr. Genet.*, 24:223-228.
- [0323] Karhunen T., Mäntylä A., Nevalainen K. M. H. and Suominen P. L. (1993) "High frequency one-step gene replacement in *Trichoderma reesei*. I. Endoglucanase I overproduction," *Mol. Gen. Genet.*, 241:515-522.
- [0324] Kurabi A., Berlin A., Gilkes N., Kilburn D., Markov A., Skomarovsky A., Gusakov A., Okunev O., Sinitsyn A., Gregg D. Xie D. and Saddler J. (2005) "Enzymatic hydrolysis of steam-exploded and ethanol organosolv-pre-treated Douglas-Fir by novel and commercial fungal cellulases," *Appl. Biochem and Biotechnol.* Vol. 121-124:219-229.
- [0325] Lever, M. (1972) "A new reaction for colorimetric determination of carbohydrates," *Anal. Biochem.*, 47:276-279.
- [0326] Lo Leggio, L., Kalogiannis S., Bhat M. K., and Pickersgill R. W. (1999) "High resolution structure and sequence of the *T. aurantiacus* xylanase I: implications for evolution of thermostability in family 10 xylanases and enzymes with (beta) alpha-barrel architecture," *Proteins* 36(3):295-306.
- [0327] Lowry, O., Rosenbrough, N., Farr, A. and Randall, R. (1951) "Protein measurement with the Folin phenol reagent," *J. Biol. Chem.* 193:265-275.
- [0328] Needleman S, and Wunsch C. (1970) "A general method applicable to the search for similarities in the amino acid sequence of two proteins," *Journal of Molecular Biology*, 48:443-453.
- [0329] Nielsen H., Engelbrecht J., Brunak S, and von Heijne G. (1997) "Identification of prokaryotic and eukaryotic signal peptides and prediction of their cleavage sites," *Protein Engineering*, 10:1-6.
- [0330] Paloheimo M., Mäntylä A., Kallio J., and Suominen P. (2003) "High-yield production of a bacterial xylanase in the filamentous fungus *Trichoderma reesei* requires a carrier polypeptide with an intact domain structure," *Appl. Env. Microbiol.*, 69:7073-7082.
- [0331] Parry N., Beever D., Owen E., Nerinckx W. Claeysens M, Van Beeumen J. and Bhat M. (2002) "Biochemical characterization and mode of action of a thermostable endoglucanase purified from *Thermoascus aurantiacus*," *Arch. of Biochem. and Biophys.*, 404:243-253.
- [0332] Penttilä M., Nevalainen H., Rättö M., Salminen E. and Knowles J. (1987) "A versatile transformation system for the cellulolytic filamentous fungus *Trichoderma reesei*," *Gene*, 61:155-164.
- [0333] Raeder U. and Broda P. (1985) "Rapid preparation of DNA from filamentous fungi," *Lett. Appl. Microbiol.*, 1:17-20.
- [0334] Rice P, Longden I and Bleasby A. (2000). "EMBOSS: The European Molecular Biology Open Software Suite," *Trends in Genetics*, 16:276-277.
- [0335] Sambrook J., Fritsch E. F. and Maniatis T. (1989) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, US.
- [0336] Sambrook J. and Russell D. W. (2001) *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, New York, US.
- [0337] Srisodsuk M, Reinikainen T, Penttilä M and Teeri T. (1993) "Role of the interdomain linker peptide of *Trichoderma reesei* cellobiohydrolase I in its interaction with crystalline cellulose," *J. Biol. Chem.*, Oct. 5, 268(28): 20756-20761.
- [0338] Sundberg, M., and Poutanen, K. (1991) "Purification and properties of two acetylxyloesterases of *Trichoderma reesei*," *Biotechnol. Appl. Biochem.*, 13:1-11.
- [0339] Suurnäkki, A., Tenkanen M., Siika-aho, M., Niku-Paavola, M.-L., Viikari, L. and Buchert, J. (2000) "*Trichoderma reesei* cellulases and their core domains in the hydrolysis and modification of chemical pulp," *Cellulose* 7:189-209.
- [0340] Tenkanen, M., Puls, J. and Poutanen, K (1992) Two major xylanases of *Trichoderma reesei*. *Enzyme Microbiol. Technol.* 14: 566-574.
- [0341] Tomme, P. McRae, S., Wood, T. and Claeysens, M. (1988) "Chromatographic separation of cellulolytic enzymes," *Methods in Enzymol.*, 160:187-192.
- [0342] Tuohy M., Walsh J., Murray P., Claeysens M., Cuffe M., Savage A. and Coughan M. (2002) "Kinetic parameters and mode of action of cellobiohydrolases produced by *Talaromyces emersonii*," *Biochem. Biophys. Acta*, 1596:366-380 (abstract).
- [0343] Van Petegem et al (2002) "Atomic resolution structure of major endoglucanase from *Thermoascus aurantiacus*," *Biochem. and Biophys. Res. Comm.*, 296:161-166.
- [0344] Van Tilbeurgh, H., Loontjens, F., de Bruyne, C. and Claeysens, M. (1988) "Fluorogenic and chromogenic glycosides as substrates and ligands of carbohydrases," *Methods Enzymol.*, 160:45-59.
- [0345] Wyman, C. E. (2001) "Twenty years of trials, tribulations, and research progress in bioethanol technology," *Applied Biochemistry and Biotechnology*, 91-93: 5-21.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 30

<210> SEQ ID NO 1

<211> LENGTH: 3192

<212> TYPE: DNA

<213> ORGANISM: *Thermoascus aurantiacus*

-continued

```

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1514)..(2122)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (2123)..(2187)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (2188)..(2949)

<400> SEQUENCE: 1

ctagaccttt atcctttcat ccgaccagac ttccccctttg accttggcgc cctggtgact    60
acctacctac ctaggtaggt aacgctcgtcg accctcttga atgatcctcg tcacactgca    120
aacatccgaa acatacggca aaagatgatt gggcatggat gcaggagaca tcgaatgagg    180
gcttagaagg aatgaaaac ctgggaccag gacgctaggt acgatgaaat ccgccaatgg    240
tgaaacttta agtcgtgctt acagcacagg ctctgtgaag attgcgctgt tcagacttaa    300
tcttctcatc acagtccaag tctttatgaa aaggaaaaga gagagaagag cgctatttcg    360
agctgtcggc ctcataggga gacagtcgag cataccagcg gtatcgacgt tagactcaac    420
caagaataat gacgagaata aacacagaag tcaaccttga actgtatatc agggttccag    480
cagcagatag ttacttgcat aaagacaact ccccgagggc tctctgcata caccaggatg    540
ttccggaatt attcactgct cgtttccgac gtggcgtcag tgatccgtct ccacagaacc    600
tctacctggg gaataacca ggggaggaat ctgcaagtaa gaacttaata ccaatccccg    660
gggctgccgg ggtgaatcaa atctccccgc ggaaattaa cccatacgat gtttttgcac    720
cacatgcatg cttggcacga tttctccgca agggagtcac agagaaagac atatttcgca    780
tactactgtg actctgcaga gttacatatc actcaggata cattgcagat cattgtccga    840
gcatcaaaca tggacctgca ggatcaacgg cccgacaaaa cacaagtggc taaagctggg    900
ggatgccccg acccgctcgc caatatcatt gatggatggt cccccacatt tttaaaacat    960
cgacggatcg gcccgcatac taatcctttt atcaacaaaa agttccactc gactagagaa   1020
aaaaaggcca aggccactaa ttgcagtcgg atactggtct tttcgccgtc caacaccttc   1080
atccatgatc cccttagcca ccaatgcccc acataataca tgttgacata ggtacgtagc   1140
tctgttatcc aatcgcattc gaacctcttt aacggacccc tcctacacac cttatcctaa   1200
cttcaggaga ctggtgcccc ttggggattg aggaggtccg ggttgcagga tgcggttctag   1260
gctaaattct cggccggtag ccatctcgaa tctctcgtga agccttcac tgaacggttg   1320
gcggcccgtc aagccgatga ccatgggttc ctgatagagc ttgtgcctga ccggccttgg   1380
cggcatagac gagctgaaca catcaggtat gaacagatca gatataaagt cggattgagt   1440
cctagtacga agcaatccgc caccacaaa tcaagcaacg agcgacagca ataacaatat   1500
caatcgaatc gca atg tat cag cgc gct ctt ctc ttc tct ttc ttc ctc   1549
Met Tyr Gln Arg Ala Leu Leu Phe Ser Phe Phe Leu
1           5           10

gcc gcc gcc cgc gcg cag cag gcc ggt acc gta acc gca gag aat cac   1597
Ala Ala Ala Arg Ala Gln Gln Ala Gly Thr Val Thr Ala Glu Asn His
15           20           25

cct tcc ctg acc tgg cag caa tgc tcc agc ggc ggt agt tgt acc acg   1645
Pro Ser Leu Thr Trp Gln Gln Cys Ser Ser Gly Gly Ser Cys Thr Thr
30           35           40

cag aat gga aaa gtc gtt atc gat gcg aac tgg cgt tgg gtc cat acc   1693
Gln Asn Gly Lys Val Val Ile Asp Ala Asn Trp Arg Trp Val His Thr

```


-continued

45	50	55	60	
acc tct gga tac acc	aac tgc tac acg	ggc aat acg tgg	gac acc agt	1741
Thr Ser Gly Tyr Thr	Asn Cys Tyr Thr	Gly Asn Thr Trp	Asp Thr Ser	
65	70	75		
atc tgt ccc gac gac	gtg acc tgc gct	cag aat tgt gcc	ttg gat gga	1789
Ile Cys Pro Asp Asp	Val Thr Cys Ala	Gln Asn Cys Ala	Leu Asp Gly	
80	85	90		
gcg gat tac agt ggc	acc tat ggt gtt	acg acc agt ggc	aac gcc ctg	1837
Ala Asp Tyr Ser Gly	Thr Tyr Gly Val	Thr Thr Ser Gly	Asn Ala Leu	
95	100	105		
aga ctg aac ttt gtc	acc caa agc tca	ggg aag aac att	ggc tcg cgc	1885
Arg Leu Asn Phe Val	Thr Gln Ser Ser	Gly Lys Asn Ile	Gly Ser Arg	
110	115	120		
ctg tac ctg ctg cag	gac gac acc act	tat cag atc ttc	aag ctg ctg	1933
Leu Tyr Leu Leu Gln	Asp Asp Thr Thr	Tyr Gln Ile Phe	Lys Leu Leu	
125	130	135	140	
ggc cag gag ttt acc	ttc gat gtc gac	gtc tcc aat ctc	cct tgc ggg	1981
Gly Gln Glu Phe Thr	Phe Asp Val Asp	Val Ser Asn Leu	Pro Cys Gly	
145	150	155		
ctg aac ggc gcc ctc	tac ttt gtg gcc	atg gac gcc gac	ggc gga ttg	2029
Leu Asn Gly Ala Leu	Tyr Phe Val Ala	Met Asp Ala Asp	Gly Gly Leu	
160	165	170		
tcc aaa tac cct ggc	aac aag gca ggc	gct aag tat ggc	act ggt tac	2077
Ser Lys Tyr Pro Gly	Asn Lys Ala Gly	Ala Lys Tyr Gly	Thr Gly Tyr	
175	180	185		
tgc gac tct cag tgc	cct cgg gat ctc	aag ttc atc aac	ggc cag	2122
Cys Asp Ser Gln Cys	Pro Arg Asp Leu	Lys Phe Ile Asn	Gly Gln	
190	195	200		
gtacgtcaga agtgataact	agccagcaga gcccatgaat	cattaactaa cgctgtcaaa		2182
tacag gcc aat gtt gaa	ggc tgg cag ccg tct	gcc aac gac cca	aat gcc	2232
Ala Asn Val Glu Gly	Trp Gln Pro Ser	Ala Asn Asp Pro	Asn Ala	
205	210	215		
ggc gtt ggt aac cac	ggt tcc tgc tgc	gct gag atg gat	gtc tgg gaa	2280
Gly Val Gly Asn His	Gly Ser Cys Cys	Ala Glu Met Asp	Val Trp Glu	
220	225	230		
gcc aac agc atc tct	act gcg gtg acg	cct cac cca tgc	gac acc ccc	2328
Ala Asn Ser Ile Ser	Thr Ala Val Thr	Pro His Pro Cys	Asp Thr Pro	
235	240	245	250	
ggc cag acc atg tgc	cag gga gac gac	tgt ggt gga acc	tac tcc tcc	2376
Gly Gln Thr Met Cys	Gln Gly Asp Asp	Cys Gly Gly Thr	Tyr Ser Ser	
255	260	265		
act cga tat gct ggt	acc tgc gac cct	gat ggc tgc gac	ttc aat cct	2424
Thr Arg Tyr Ala Gly	Thr Cys Asp Pro	Asp Gly Cys Asp	Phe Asn Pro	
270	275	280		
tac cgc cag ggc aac	cac tcg ttc tac	ggc ccc ggg cag	atc gtc gac	2472
Tyr Arg Gln Gly Asn	His Ser Phe Tyr	Gly Pro Gly Gln	Ile Val Asp	
285	290	295		
acc agc tcc aaa ttc	acc gtc gtc acc	cag ttc atc acc	gac gac ggg	2520
Thr Ser Ser Lys Phe	Thr Val Val Thr	Gln Phe Ile Thr	Asp Asp Gly	
300	305	310		
acc ccc tcc ggc acc	ctg acg gag atc	aaa cgc ttc tac	gtc cag aac	2568
Thr Pro Ser Gly Thr	Leu Thr Glu Ile	Lys Arg Phe Tyr	Val Gln Asn	
315	320	325	330	
ggc aag gta atc ccc	cag tcg gag tcg	acg atc agc ggc	gtc acc ggc	2616
Gly Lys Val Ile Pro	Gln Ser Glu Ser	Thr Ile Ser Gly	Val Thr Gly	
335	340	345		

-continued

```

aac tca atc acc acc gag tat tgc acg gcc cag aag gcc gcc ttc ggc 2664
Asn Ser Ile Thr Thr Glu Tyr Cys Thr Ala Gln Lys Ala Ala Phe Gly
350 355 360

gac aac acc ggc ttc ttc acg cac ggc ggg ctt cag aag atc agt cag 2712
Asp Asn Thr Gly Phe Phe Thr His Gly Gly Leu Gln Lys Ile Ser Gln
365 370 375

gct ctg gct cag ggc atg gtc ctc gtc atg agc ctg tgg gac gat cac 2760
Ala Leu Ala Gln Gly Met Val Leu Val Met Ser Leu Trp Asp Asp His
380 385 390

gcc gcc aac atg ctc tgg ctg gac agc acc tac ccg act gat gcg gac 2808
Ala Ala Asn Met Leu Trp Leu Asp Ser Thr Tyr Pro Thr Asp Ala Asp
395 400 405 410

ccg gac acc cct ggc gtc gcg cgc ggt acc tgc ccc acg acc tcc ggc 2856
Pro Asp Thr Pro Gly Val Ala Arg Gly Thr Cys Pro Thr Thr Ser Gly
415 420 425

gtc ccg gcc gac gtt gag tcg cag tac ccc aat tca tat gtt atc tac 2904
Val Pro Ala Asp Val Glu Ser Gln Tyr Pro Asn Ser Tyr Val Ile Tyr
430 435 440

tcc aac atc aag gtc gga ccc atc aac tcg acc ttc acc gcc aac 2949
Ser Asn Ile Lys Val Gly Pro Ile Asn Ser Thr Phe Thr Ala Asn
445 450 455

taagtaagta actggcactc taccaccgag agcttcgtga agatacaggg gtggttggga 3009

gattgtcgtg tacaggggac atgcatgct caaaaatcta catcagtttg ccaattgaac 3069

catgaaaaaa agggggagat caaagaagtc tgtcaaaaga ggggggctgt ggcagcttaa 3129

gccttggtgt agatcgagtc gacgcctat agtgagtcgt attagagctc gcggccgcca 3189

gct 3192

```

<210> SEQ ID NO 2

<211> LENGTH: 457

<212> TYPE: PRT

<213> ORGANISM: Thermoascus aurantiacus

<400> SEQUENCE: 2

```

Met Tyr Gln Arg Ala Leu Leu Phe Ser Phe Phe Leu Ala Ala Ala Arg
1 5 10 15

Ala Gln Gln Ala Gly Thr Val Thr Ala Glu Asn His Pro Ser Leu Thr
20 25 30

Trp Gln Gln Cys Ser Ser Gly Gly Ser Cys Thr Thr Gln Asn Gly Lys
35 40 45

Val Val Ile Asp Ala Asn Trp Arg Trp Val His Thr Thr Ser Gly Tyr
50 55 60

Thr Asn Cys Tyr Thr Gly Asn Thr Trp Asp Thr Ser Ile Cys Pro Asp
65 70 75 80

Asp Val Thr Cys Ala Gln Asn Cys Ala Leu Asp Gly Ala Asp Tyr Ser
85 90 95

Gly Thr Tyr Gly Val Thr Thr Ser Gly Asn Ala Leu Arg Leu Asn Phe
100 105 110

Val Thr Gln Ser Ser Gly Lys Asn Ile Gly Ser Arg Leu Tyr Leu Leu
115 120 125

Gln Asp Asp Thr Thr Tyr Gln Ile Phe Lys Leu Leu Gly Gln Glu Phe
130 135 140

Thr Phe Asp Val Asp Val Ser Asn Leu Pro Cys Gly Leu Asn Gly Ala
145 150 155 160

```


-continued

Leu Tyr Phe Val Ala Met Asp Ala Asp Gly Gly Leu Ser Lys Tyr Pro
 165 170 175
 Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp Ser Gln
 180 185 190
 Cys Pro Arg Asp Leu Lys Phe Ile Asn Gly Gln Ala Asn Val Glu Gly
 195 200 205
 Trp Gln Pro Ser Ala Asn Asp Pro Asn Ala Gly Val Gly Asn His Gly
 210 215 220
 Ser Cys Cys Ala Glu Met Asp Val Trp Glu Ala Asn Ser Ile Ser Thr
 225 230 235 240
 Ala Val Thr Pro His Pro Cys Asp Thr Pro Gly Gln Thr Met Cys Gln
 245 250 255
 Gly Asp Asp Cys Gly Gly Thr Tyr Ser Ser Thr Arg Tyr Ala Gly Thr
 260 265 270
 Cys Asp Pro Asp Gly Cys Asp Phe Asn Pro Tyr Arg Gln Gly Asn His
 275 280 285
 Ser Phe Tyr Gly Pro Gly Gln Ile Val Asp Thr Ser Ser Lys Phe Thr
 290 295 300
 Val Val Thr Gln Phe Ile Thr Asp Asp Gly Thr Pro Ser Gly Thr Leu
 305 310 315 320
 Thr Glu Ile Lys Arg Phe Tyr Val Gln Asn Gly Lys Val Ile Pro Gln
 325 330 335
 Ser Glu Ser Thr Ile Ser Gly Val Thr Gly Asn Ser Ile Thr Thr Glu
 340 345 350
 Tyr Cys Thr Ala Gln Lys Ala Ala Phe Gly Asp Asn Thr Gly Phe Phe
 355 360 365
 Thr His Gly Gly Leu Gln Lys Ile Ser Gln Ala Leu Ala Gln Gly Met
 370 375 380
 Val Leu Val Met Ser Leu Trp Asp Asp His Ala Ala Asn Met Leu Trp
 385 390 395 400
 Leu Asp Ser Thr Tyr Pro Thr Asp Ala Asp Pro Asp Thr Pro Gly Val
 405 410 415
 Ala Arg Gly Thr Cys Pro Thr Thr Ser Gly Val Pro Ala Asp Val Glu
 420 425 430
 Ser Gln Tyr Pro Asn Ser Tyr Val Ile Tyr Ser Asn Ile Lys Val Gly
 435 440 445
 Pro Ile Asn Ser Thr Phe Thr Ala Asn
 450 455

<210> SEQ ID NO 3
 <211> LENGTH: 3055
 <212> TYPE: DNA
 <213> ORGANISM: Acremonium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (972)..(1595)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1596)..(1729)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1730)..(2290)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (2291)..(2412)
 <220> FEATURE:
 <221> NAME/KEY: CDS

-continued

```

<222> LOCATION: (2413)..(2540)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (2541)..(2627)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (2628)..(2691)

<400> SEQUENCE: 3

gaattcggat cacaccgaga gcttcgcat ggccagctgt ctcagcttgt acccgtctac      60
caacgttccg catcttcggt accttgatag ctcgctgttg ctggactgct ttgtgagggg    120
actgtgccac gcctgggaga cgggtgccgt accatcggtt actgctcaga ctgagaaccg    180
tcgttgccga aacagccagg caggaagcct gtccaccttc atgtatcttc atatggaccc    240
cagcgcgccc ctctctttct cctcatttct tgcccaccac gatggacacc atgccaatct    300
atctcttgat cccttgactc ctcagccccc cagcagtcgg acaatgtaca gtgatgggca    360
tctctttctg tacatacgtc ccctctcgcg gtgtccacgc gcggccgggg atgcctggga    420
cggagtgccca cccgcaggga acgagacttg gctgatgggg tgcgggtgcat ggtggcacia    480
gagatccagg cccccgatc tegtctctgc acgtatcctt cccccgccgg cgatgcccaa    540
gtgggaagtc ttcggagcgg caccagggcc catcttgccg atgcccggca cggctctggc    600
ggttgccctc atctatcgtg gctgcacatc cggcgtgccc ccattgggaa agcaggcttt    660
gttcttcccg tctgtcgcac gtctcccacc taccctcctt cctcgcgaagg gcttaccctg    720
gcccctcaact gctgcttca ctcactgctg cttccccgca atgccccctc gccccccccc    780
ccccctctc ctttgagta cagatctaca taatatcgag acgccccca agctgtttct     840
ctggcacagc cctctcgcgc gtggtgcaag agcaagtcag agtatcaatt cccccatctc    900
tcactcagc ccttctgccc tgggtccacc gacattctgg gcccgtagcc aagaccgatc    960
cgctctcaac c atg cac aag cgg gcg gcc acc ctc tcc gcc ctc gtc gtc    1010
Met His Lys Arg Ala Ala Thr Leu Ser Ala Leu Val Val
1           5           10

gcc gcc gcc ggc ttc gcc cgc ggc cag ggc gtg ggc acg cag cag acg    1058
Ala Ala Ala Gly Phe Ala Arg Gly Gln Gly Val Gly Thr Gln Gln Thr
15          20          25

gag acg cac ccc aag ctc acc ttc cag aag tgc tcc gcc gcc ggc agc    1106
Glu Thr His Pro Lys Leu Thr Phe Gln Lys Cys Ser Ala Ala Gly Ser
30          35          40          45

tgc acg acc cag aac ggc gag gtg gtc atc gac gcc aac tgg cgc tgg    1154
Cys Thr Thr Gln Asn Gly Glu Val Val Ile Asp Ala Asn Trp Arg Trp
50          55          60

gtg cac gac aag aac ggc tac acc aac tgc tac acg ggc aac gag tgg    1202
Val His Asp Lys Asn Gly Tyr Thr Asn Cys Tyr Thr Gly Asn Glu Trp
65          70          75

aac acc acc atc tgc gcc gac gcc gcc tcg tgc gcc agc aac tgc gtc    1250
Asn Thr Thr Ile Cys Ala Asp Ala Ala Ser Cys Ala Ser Asn Cys Val
80          85          90

gtc gac ggc gcc gac tac cag ggc acc tac ggc gcc tcc acc tcc ggc    1298
Val Asp Gly Ala Asp Tyr Gln Gly Thr Tyr Gly Ala Ser Thr Ser Gly
95          100         105

aac gcc ctg acc ctc aag ttc gtc acc aag ggc agc tac gcc acc aac    1346
Asn Ala Leu Thr Leu Lys Phe Val Thr Lys Gly Ser Tyr Ala Thr Asn
110         115         120         125

atc ggc tcg cgc atg tac ctg atg gcc agc ccc acc aag tac gcc atg    1394
Ile Gly Ser Arg Met Tyr Leu Met Ala Ser Pro Thr Lys Tyr Ala Met

```


-continued

130	135	140	
ttc acc ctg ctg ggc cac gag ttc gcc ttc gac gtc gac ctg agc aag			1442
Phe Thr Leu Leu Gly His Glu Phe Ala Phe Asp Val Asp Leu Ser Lys			
145	150	155	
ctg ccc tgc ggc ctc aac ggc gcc gtc tac ttc gtc agc atg gac gag			1490
Leu Pro Cys Gly Leu Asn Gly Ala Val Tyr Phe Val Ser Met Asp Glu			
160	165	170	
gac ggc ggc acc agc aag tac ccc tcc aac aag gcc ggc gcc aag tac			1538
Asp Gly Gly Thr Ser Lys Tyr Pro Ser Asn Lys Ala Gly Ala Lys Tyr			
175	180	185	
ggc acg ggc tac tgc gac tcg cag tgt ccg cgc gac ctc aag ttt atc			1586
Gly Thr Gly Tyr Cys Asp Ser Gln Cys Pro Arg Asp Leu Lys Phe Ile			
190	195	200	205
gac ggc aag gtgagaacc gcactagcgt ccgccttcc gtgtccctcc			1635
Asp Gly Lys			
ttttgccttc ttcgaccgcc ctcttccttg cgggccaggg tcgctggggg gctgtcctcc			1695
tttctggtgg gcagcgtgc tgatcccgcg ccag gcc aac tcg gcc agc tgg cag			1750
Ala Asn Ser Ala Ser Trp Gln			
210	215		
ccc tcg tcc aac gac cag aac gcc ggc gtg ggc ggc atg ggc tcg tgc			1798
Pro Ser Ser Asn Asp Gln Asn Ala Gly Val Gly Gly Met Gly Ser Cys			
220	225	230	
tgc gcc gag atg gac atc tgg gag gcc aac tcc gtc tcc gcc gcc tac			1846
Cys Ala Glu Met Asp Ile Trp Glu Ala Asn Ser Val Ser Ala Ala Tyr			
235	240	245	
acg ccg cac ccg tgc cag aac tac cag cag cac agc tgc agc ggc gac			1894
Thr Pro His Pro Cys Gln Asn Tyr Gln Gln His Ser Cys Ser Gly Asp			
250	255	260	
gac tgc ggc ggc acc tac tcg gcc acc cgc ttc gcc ggc gac tgc gac			1942
Asp Cys Gly Gly Thr Tyr Ser Ala Thr Arg Phe Ala Gly Asp Cys Asp			
265	270	275	
ccg gac ggc tgc gac tgg aac gcc tac cgc atg ggc gtg cac gac ttc			1990
Pro Asp Gly Cys Asp Trp Asn Ala Tyr Arg Met Gly Val His Asp Phe			
280	285	290	295
tac ggc aac ggc aag acc gtc gac acc ggc aag aag ttc tcc atc gtc			2038
Tyr Gly Asn Gly Lys Thr Val Asp Thr Gly Lys Lys Phe Ser Ile Val			
300	305	310	
acc cag ttc aag ggc tcc ggc tcc acc ctg acc gag atc aag cag ttc			2086
Thr Gln Phe Lys Gly Ser Gly Ser Thr Leu Thr Glu Ile Lys Gln Phe			
315	320	325	
tac gtc cag gac ggc agg aag atc gag aac ccc aac gcc acc tgg ccc			2134
Tyr Val Gln Asp Gly Arg Lys Ile Glu Asn Pro Asn Ala Thr Trp Pro			
330	335	340	
ggc ctc gag ccc ttc aac tcc atc acc ccg gac ttc tgc aag gcc cag			2182
Gly Leu Glu Pro Phe Asn Ser Ile Thr Pro Asp Phe Cys Lys Ala Gln			
345	350	355	
aag cag gtc ttc ggc gac ccc gac cgc ttc aac gac atg ggc ggc ttc			2230
Lys Gln Val Phe Gly Asp Pro Asp Arg Phe Asn Asp Met Gly Gly Phe			
360	365	370	375
acc aac atg gcc aag gcc ctg gcc aac ccc atg gtc ctg gtg ctg tcg			2278
Thr Asn Met Ala Lys Ala Leu Ala Asn Pro Met Val Leu Val Leu Ser			
380	385	390	
ctg tgg gac gac gtgagccatt ttcgattct ctctgactc tctccgctg			2330
Leu Trp Asp Asp			
395			
ccatcaccac ctcttcacc accgccacga ggggtgtagct tgatctccgc tgactgacgt			2390

-continued

```

gtgcccacac ccccgtttct ag cac tac tcc aac atg ctg tgg ctc gac tct 2442
His Tyr Ser Asn Met Leu Trp Leu Asp Ser
400 405

acc tac ccg acc gac gcc gat ccc agc gcg ccc ggc aag gga cgt ggc 2490
Thr Tyr Pro Thr Asp Ala Asp Pro Ser Ala Pro Gly Lys Gly Arg Gly
410 415 420

acc tgc gac acc agc agc ggc gtg cca agc gac gtg gag tcg aag aat 2538
Thr Cys Asp Thr Ser Ser Gly Val Pro Ser Asp Val Glu Ser Lys Asn
425 430 435

gg gtgagtcgga tcttctgcat gcgcccgctt ttccgagcat tgcttggggt 2590
Gly
cctccctcag gctgacacac gcgcgccttc gatacag c gat gcg acc gtc atc 2643
Asp Ala Thr Val Ile
440

tac tcc aac atc aag ttt ggg ccg ctg gac tcc acc tac acg gct tcc 2691
Tyr Ser Asn Ile Lys Phe Gly Pro Leu Asp Ser Thr Tyr Thr Ala Ser
445 450 455

tgagcagccg ctttgggttc ggtggggccg aagcacaaca agtgtgtgcg tagctgagat 2751

gatggccgat ctctgtcctt tgtctcctag tgtctctctt atcgaacaac cccccgacct 2811

gcagcgtcgg cgggcatcgt atagtctggt gtaactgtat atagctctgt gcgtgtgaat 2871

cgaacgagca ccgacgaaat gtggtgtttc atgetatcgt acatgctctt gcgagatctg 2931

aagtcgtcaa ttagacattg ccaccatcca acttggcgac tgtccaccgg gtccatttgt 2991

atcactggct cttccgagac ccggtctctc tcacaccgta atcactgcaa gcagagttga 3051

attc 3055

```

```

<210> SEQ ID NO 4
<211> LENGTH: 459
<212> TYPE: PRT
<213> ORGANISM: Acremonium thermophilum

```

```

<400> SEQUENCE: 4

```

```

Met His Lys Arg Ala Ala Thr Leu Ser Ala Leu Val Val Ala Ala Ala
1 5 10 15

Gly Phe Ala Arg Gly Gln Gly Val Gly Thr Gln Gln Thr Glu Thr His
20 25 30

Pro Lys Leu Thr Phe Gln Lys Cys Ser Ala Ala Gly Ser Cys Thr Thr
35 40 45

Gln Asn Gly Glu Val Val Ile Asp Ala Asn Trp Arg Trp Val His Asp
50 55 60

Lys Asn Gly Tyr Thr Asn Cys Tyr Thr Gly Asn Glu Trp Asn Thr Thr
65 70 75 80

Ile Cys Ala Asp Ala Ala Ser Cys Ala Ser Asn Cys Val Val Asp Gly
85 90 95

Ala Asp Tyr Gln Gly Thr Tyr Gly Ala Ser Thr Ser Gly Asn Ala Leu
100 105 110

Thr Leu Lys Phe Val Thr Lys Gly Ser Tyr Ala Thr Asn Ile Gly Ser
115 120 125

Arg Met Tyr Leu Met Ala Ser Pro Thr Lys Tyr Ala Met Phe Thr Leu
130 135 140

Leu Gly His Glu Phe Ala Phe Asp Val Asp Leu Ser Lys Leu Pro Cys
145 150 155 160

Gly Leu Asn Gly Ala Val Tyr Phe Val Ser Met Asp Glu Asp Gly Gly

```


-continued

165		170		175
Thr Ser Lys Tyr Pro	Ser Asn Lys Ala Gly	Ala Lys Tyr Gly Thr Gly		
180	185	190		
Tyr Cys Asp Ser Gln	Cys Pro Arg Asp Leu	Lys Phe Ile Asp Gly Lys		
195	200	205		
Ala Asn Ser Ala Ser	Trp Gln Pro Ser Ser	Asn Asp Gln Asn Ala Gly		
210	215	220		
Val Gly Gly Met Gly	Ser Cys Cys Ala Glu	Met Asp Ile Trp Glu Ala		
225	230	235	240	
Asn Ser Val Ser Ala	Ala Tyr Thr Pro His	Pro Cys Gln Asn Tyr Gln		
245	250	255		
Gln His Ser Cys Ser	Gly Asp Asp Cys Gly	Gly Thr Tyr Ser Ala Thr		
260	265	270		
Arg Phe Ala Gly Asp	Cys Asp Pro Asp Gly	Cys Asp Trp Asn Ala Tyr		
275	280	285		
Arg Met Gly Val His	Asp Phe Tyr Gly Asn	Gly Lys Thr Val Asp Thr		
290	295	300		
Gly Lys Lys Phe Ser	Ile Val Thr Gln Phe	Lys Gly Ser Gly Ser Thr		
305	310	315	320	
Leu Thr Glu Ile Lys	Gln Phe Tyr Val Gln	Asp Gly Arg Lys Ile Glu		
325	330	335		
Asn Pro Asn Ala Thr	Trp Pro Gly Leu Glu	Pro Phe Asn Ser Ile Thr		
340	345	350		
Pro Asp Phe Cys Lys	Ala Gln Lys Gln Val	Phe Gly Asp Pro Asp Arg		
355	360	365		
Phe Asn Asp Met Gly	Gly Phe Thr Asn Met	Ala Lys Ala Leu Ala Asn		
370	375	380		
Pro Met Val Leu Val	Leu Ser Leu Trp Asp	Asp His Tyr Ser Asn Met		
385	390	395	400	
Leu Trp Leu Asp Ser	Thr Tyr Pro Thr Asp	Ala Asp Pro Ser Ala Pro		
405	410	415		
Gly Lys Gly Arg Gly	Thr Cys Asp Thr Ser	Ser Gly Val Pro Ser Asp		
420	425	430		
Val Glu Ser Lys Asn	Gly Asp Ala Thr Val	Ile Tyr Ser Asn Ile Lys		
435	440	445		
Phe Gly Pro Leu Asp	Ser Thr Tyr Thr Ala	Ser		
450	455			

<210> SEQ ID NO 5
 <211> LENGTH: 3401
 <212> TYPE: DNA
 <213> ORGANISM: Acremonium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (891)..(1299)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1300)..(1387)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1388)..(1442)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1443)..(1495)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1496)..(1643)

-continued

```

<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1644)..(1697)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1698)..(1928)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1929)..(2014)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (2015)..(2740)

<400> SEQUENCE: 5

ctcgagtttc cctggtcggc cactctctgc tcctctcgtc ctgcgccctt ggatgtgccg      60
tgtgtccagt cgtgtatctc ttgactgcac gacgtgttcc tcgcgactcg tctcgcgccg      120
gtggatgccc gtccactcat ttgtccgtct actgggtcag cctctcgtct cgaacgagct      180
tccacggccc actccccgga caacctcggc tctggatggc cctcctcccc ctccgtgtct      240
cccctcctgc ggggtccgtc gtgccctggc tgcattgctc acatcgcttg atcacgctgc      300
gagccaccgc agagccccc atccaaagcg accgtggcag cactacctct gtttctggga      360
tggggcccac gtcgatggc tggcatccct tgccaccctc ctccatcccc ctgacctcac      420
tcccaaccga taggagaagt ggatcatggc acgacccctg gcacgtcttg gactcgacga      480
gcttgatcgg gccggaagcc gtcaacgacg ggggagccgt gtcttgccac gcgtggccgt      540
ccttcgacag tggacagcga gaaaactggg ggggaagagg gctgctacag tcttgtcttg      600
cgaggcccga cgctcctagt ccgagaacca cctactgtgt tctcgcgaag acggggccag      660
cttagcggcc aaatttgccc cccgggcta gggctctagc atggggatga tgaactggtg      720
tcgacgatgt ctatataacg acggcgatct cctgtctctg agatcccatc ctttcatctc      780
caaccactt catcccttc tctctctctc ccctccctt ctctgacata ccgagtcctc      840
agaagcctcg tccgtcgtca cctattctca ctccccggg aactccggcc atg tat      896
Met Tyr
1

acc aag ttc gcc gcc ctc gcc gcc ctc gtg gcc acc gtc cgc ggc cag      944
Thr Lys Phe Ala Ala Leu Ala Ala Leu Val Ala Thr Val Arg Gly Gln
5 10 15

gcc gcc tgc tcg ctc acc gcc gag acc cac ccg tcg ctg cag tgg cag      992
Ala Ala Cys Ser Leu Thr Ala Glu Thr His Pro Ser Leu Gln Trp Gln
20 25 30

aag tgc acc gcg ccc gcc agc tgc acc acc gtc agc ggc cag gtc acc      1040
Lys Cys Thr Ala Pro Gly Ser Cys Thr Thr Val Ser Gly Gln Val Thr
35 40 45 50

atc gac gcc aac tgg cgc tgg ctg cac cag acc aac agc agc acc aac      1088
Ile Asp Ala Asn Trp Arg Trp Leu His Gln Thr Asn Ser Ser Thr Asn
55 60 65

tgc tac acc ggc aac gag tgg gac acc agc atc tgc agc tcc gac acc      1136
Cys Tyr Thr Gly Asn Glu Trp Asp Thr Ser Ile Cys Ser Ser Asp Thr
70 75 80

gac tgc gcc acc aag tgc tgc ctc gac gcc gcc gac tac acc ggc acc      1184
Asp Cys Ala Thr Lys Cys Cys Leu Asp Gly Ala Asp Tyr Thr Gly Thr
85 90 95

tac ggc gtc acc gcc agc gcc aac tcg ctc aac ctc aag ttc gtc acc      1232
Tyr Gly Val Thr Ala Ser Gly Asn Ser Leu Asn Leu Lys Phe Val Thr
100 105 110

cag ggg ccc tac tcc aag aac atc gcc tcg cgc atg tac ctc atg gag      1280

```


-continued

Gln Gly Pro Tyr Ser Lys Asn Ile Gly Ser Arg Met Tyr Leu Met Glu	
115	120 125 130
tcg gag tcc aag tac cag g gtagagcatat agatcacatc tttcgtcact	1329
Ser Glu Ser Lys Tyr Gln	
135	
tgcgtccggtt tcgcacggca agcgggtccag acgctaacgg gacggttctc ttctctag	1387
gc ttc act ctc ctc ggt cag gag ttt acc ttt gac gtg gac gtc tcc	1434
Gly Phe Thr Leu Leu Gly Gln Glu Phe Thr Phe Asp Val Asp Val Ser	
140	145 150
aac ctc gg gtaggtgatg acttctcccg catgagaaga gctctgctaa	1482
Asn Leu Gly	
155	
ccgtgttgtc cag c tgc ggt ctg aac gga gcg ctc tac ttc gtg tcc atg	1532
Cys Gly Leu Asn Gly Ala Leu Tyr Phe Val Ser Met	
160	165
gac ctc gac ggc ggc gtg tcc aag tac acc acc aac aag gcc ggc gcc	1580
Asp Leu Asp Gly Gly Val Ser Lys Tyr Thr Thr Asn Lys Ala Gly Ala	
170	175 180
aag tac ggc acc ggc tac tgc gac tcc cag tgc ccg cgg gat ctc aag	1628
Lys Tyr Gly Thr Gly Tyr Cys Asp Ser Gln Cys Pro Arg Asp Leu Lys	
185	190 195
ttc atc aac ggc cag gtgggtcagag agaccctctt cccctctcag tgaacgatgt	1683
Phe Ile Asn Gly Gln	
200	
ctgaccctct ctag gcc aac atc gac ggc tgg caa ccg tcg tcc aac gac	1733
Ala Asn Ile Asp Gly Trp Gln Pro Ser Ser Asn Asp	
205	210 215
gcc aac gcc ggc ctc ggg aac cac ggc agc tgc tgc tcc gag atg gac	1781
Ala Asn Ala Gly Leu Gly Asn His Gly Ser Cys Cys Ser Glu Met Asp	
220	225 230
atc tgg gag gcc aac aag gtc tcc gcc gcc tac acg ccg cac ccc tgc	1829
Ile Trp Glu Ala Asn Lys Val Ser Ala Ala Tyr Thr Pro His Pro Cys	
235	240 245
acc acc atc ggc cag acc atg tgc acc ggc gac gac tgc ggc ggc acc	1877
Thr Thr Ile Gly Gln Thr Met Cys Thr Gly Asp Asp Cys Gly Gly Thr	
250	255 260
tat tcg tcg gac cgc tat gcc ggc atc tgc gac ccc gac ggt tgc gat	1925
Tyr Ser Ser Asp Arg Tyr Ala Gly Ile Cys Asp Pro Asp Gly Cys Asp	
265	270 275 280
ttt gtaggttctt tctctcgccg ctccctgacg acctatatgt gtgaagggac	1978
Phe	
gcacagaaaa gacaaggtca aagctgacca gagcag aac tcg tac cgc atg ggc	2032
Asn Ser Tyr Arg Met Gly	
285	
gac acc agc ttc tac ggc ccc ggc aag acg gtc gac acc ggc tcc aag	2080
Asp Thr Ser Phe Tyr Gly Pro Gly Lys Thr Val Asp Thr Gly Ser Lys	
290	295 300
ttc acc gtc gtg acc cag ttc ctc acg ggc tcc gac ggc aac ctc agc	2128
Phe Thr Val Val Thr Gln Phe Leu Thr Gly Ser Asp Gly Asn Leu Ser	
305	310 315
gag atc aag cgc ttc tac gtg cag aac ggc aag gtc atc ccc aac tcc	2176
Glu Ile Lys Arg Phe Tyr Val Gln Asn Gly Lys Val Ile Pro Asn Ser	
320	325 330 335
gag tcc aag atc gcc ggc gtc tcc ggc aac tcc atc acc acc gac ttc	2224
Glu Ser Lys Ile Ala Gly Val Ser Gly Asn Ser Ile Thr Thr Asp Phe	
340	345 350

-continued

tgc acc gcc cag aag acc gcc ttc ggc gac acc aac gtc ttc gag gag Cys Thr Ala Gln Lys Thr Ala Phe Gly Asp Thr Asn Val Phe Glu Glu 355 360 365	2272
cgc ggc ggc ctc gcc cag atg ggc aag gcc ctg gcc gag ccc atg gtc Arg Gly Gly Leu Ala Gln Met Gly Lys Ala Leu Ala Glu Pro Met Val 370 375 380	2320
ctg gtc ctg tcc gtc tgg gac gac cac gcc gtc aac atg ctc tgg ctc Leu Val Leu Ser Val Trp Asp Asp His Ala Val Asn Met Leu Trp Leu 385 390 395	2368
gac tcc acc tac ccc acc gac agc acc aag ccc ggc gcc gcc cgc ggc Asp Ser Thr Tyr Pro Thr Asp Ser Thr Lys Pro Gly Ala Ala Arg Gly 400 405 410 415	2416
gac tgc ccc atc acc tcc ggc gtg ccc gcc gac gtc gag tcc cag gcg Asp Cys Pro Ile Thr Ser Gly Val Pro Ala Asp Val Glu Ser Gln Ala 420 425 430	2464
ccc aac tcc aac gtc atc tac tcc aac atc cgc ttc ggc ccc atc aac Pro Asn Ser Asn Val Ile Tyr Ser Asn Ile Arg Phe Gly Pro Ile Asn 435 440 445	2512
tcc acc tac acc ggc acc ccc agc ggc ggc aac ccc ccc ggc ggc ggc Ser Thr Tyr Thr Gly Thr Pro Ser Gly Gly Asn Pro Pro Gly Gly Gly 450 455 460	2560
acc acc acc acc acc acc acc acc acc tcc aag ccc tcc ggc ccc acc Thr Thr Thr Thr Thr Thr Thr Thr Thr Ser Lys Pro Ser Gly Pro Thr 465 470 475	2608
acc acc acc aac ccc tcg ggt ccg cag cag acg cac tgg ggt cag tgc Thr Thr Thr Asn Pro Ser Gly Pro Gln Gln Thr His Trp Gly Gln Cys 480 485 490 495	2656
ggc ggc cag gga tgg acc ggc ccc acg gtc tgc cag agc ccc tac acc Gly Gly Gln Gly Trp Thr Gly Pro Thr Val Cys Gln Ser Pro Tyr Thr 500 505 510	2704
tgc aag tac tcc aac gac tgg tac tcg cag tgc ctg taagccataa Cys Lys Tyr Ser Asn Asp Trp Tyr Ser Gln Cys Leu 515 520	2750
gccccctgta cgttcggaag acggtggcaa cagacaaacc cctccccga gcacaccccc	2810
cagggatcta agggggttgt ggtaagaca taagaatgcg ccgtggcttg gcctacgcca	2870
cggtcatgaa agtgcagtga aaatgggggc aagagtcgga aaaagtgagt ttgcttgcaa	2930
gggagagagg atgtcgagag gtgatgactt cgtttgtaca tagttggtc ttcgtgattg	2990
ggaacgggag gagggtcggg gggagccctc cagactcctt ggctctccg ctcgttccat	3050
ctttctcagt acatatacat ctgcattttc atccacgtct ctggcgtctc tggatgtgaa	3110
cgaatccgac aactggtggg ctgagatgaa tcgcaaggag agtatcttgc gaggatatca	3170
cagtcagaaa gtagcatttg agccactact aaaaggtaaa ccagtatgcg aagcttagca	3230
attatataca gcagctcaac ttcagaacga agtattgcat gtggcagaga atcttgggaa	3290
atgagccatg aagacctcgt cgagagagta cctctcaccg ccaaataacc agctagcggg	3350
ttgggagagg agcaatagga cgagcgcgat ggacagatat acgaactcga g	3401

<210> SEQ ID NO 6
 <211> LENGTH: 523
 <212> TYPE: PRT
 <213> ORGANISM: Acremonium thermophilum

<400> SEQUENCE: 6

Met Tyr Thr Lys Phe Ala Ala Leu Ala Ala Leu Val Ala Thr Val Arg

-continued

1	5	10	15
Gly 20	Gln Ala Ala Cys	Ser Leu Thr Ala Glu	Thr His Pro Ser Leu Gln 30
Trp 35	Gln Lys Cys Thr	Ala Pro Gly Ser Cys	Thr Thr Val Ser Gly Gln 45
Val 50	Thr Ile Asp Ala	Asn Trp Arg Trp Leu	His Gln Thr Asn Ser Ser 60
Thr 65	Asn Cys Tyr Thr	Gly Asn Glu Trp Asp	Thr Ser Ile Cys Ser Ser 80
Asp 85	Thr Asp Cys Ala	Thr Lys Cys Cys Leu	Asp Gly Ala Asp Tyr Thr 95
Gly 100	Thr Tyr Gly Val	Thr Ala Ser Gly Asn	Ser Leu Asn Leu Lys Phe 110
Val 115	Thr Gln Gly Pro	Tyr Ser Lys Asn Ile	Gly Ser Arg Met Tyr Leu 125
Met 130	Glu Ser Glu Ser	Lys Tyr Gln Gly Phe	Thr Leu Leu Gly Gln Glu 140
Phe 145	Thr Phe Asp Val	Asp Val Ser Asn Leu	Gly Cys Gly Leu Asn Gly 160
Ala 165	Leu Tyr Phe Val	Ser Met Asp Leu Asp	Gly Gly Val Ser Lys Tyr 175
Thr 180	Thr Asn Lys Ala	Gly Ala Lys Tyr Gly	Thr Gly Tyr Cys Asp Ser 190
Gln 195	Cys Pro Arg Asp	Leu Lys Phe Ile Asn	Gly Gln Ala Asn Ile Asp 205
Gly 210	Trp Gln Pro Ser	Ser Asn Asp Ala Asn	Ala Gly Leu Gly Asn His 220
Gly 225	Ser Cys Cys Ser	Glu Met Asp Ile Trp	Glu Ala Asn Lys Val Ser 240
Ala 245	Ala Tyr Thr Pro	His Pro Cys Thr Thr	Ile Gly Gln Thr Met Cys 255
Thr 260	Gly Asp Asp Cys	Gly Gly Thr Tyr Ser	Ser Asp Arg Tyr Ala Gly 270
Ile 275	Cys Asp Pro Asp	Gly Cys Asp Phe Asn	Ser Tyr Arg Met Gly Asp 285
Thr 290	Ser Phe Tyr Gly	Pro Gly Lys Thr Val	Asp Thr Gly Ser Lys Phe 300
Thr 305	Val Val Thr Gln	Phe Leu Thr Gly Ser	Asp Gly Asn Leu Ser Glu 320
Ile 325	Lys Arg Phe Tyr	Val Gln Asn Gly Lys	Val Ile Pro Asn Ser Glu 335
Ser 340	Lys Ile Ala Gly	Val Ser Gly Asn Ser	Ile Thr Thr Asp Phe Cys 350
Thr 355	Ala Gln Lys Thr	Ala Phe Gly Asp Thr	Asn Val Phe Glu Glu Arg 365
Gly 370	Gly Leu Ala Gln	Met Gly Lys Ala Leu	Ala Glu Pro Met Val Leu 380
Val 385	Leu Ser Val Trp	Asp Asp His Ala Val	Asn Met Leu Trp Leu Asp 400
Ser 405	Thr Tyr Pro Thr	Asp Ser Thr Lys Pro	Gly Ala Ala Arg Gly Asp 415

-continued

Cys Pro Ile Thr Ser Gly Val Pro Ala Asp Val Glu Ser Gln Ala Pro
 420 425 430

Asn Ser Asn Val Ile Tyr Ser Asn Ile Arg Phe Gly Pro Ile Asn Ser
 435 440 445

Thr Tyr Thr Gly Thr Pro Ser Gly Gly Asn Pro Pro Gly Gly Gly Thr
 450 455 460

Thr Thr Thr Thr Thr Thr Thr Thr Ser Lys Pro Ser Gly Pro Thr Thr
 465 470 475 480

Thr Thr Asn Pro Ser Gly Pro Gln Gln Thr His Trp Gly Gln Cys Gly
 485 490 495

Gly Gln Gly Trp Thr Gly Pro Thr Val Cys Gln Ser Pro Tyr Thr Cys
 500 505 510

Lys Tyr Ser Asn Asp Trp Tyr Ser Gln Cys Leu
 515 520

<210> SEQ ID NO 7
 <211> LENGTH: 3649
 <212> TYPE: DNA
 <213> ORGANISM: Chaetomium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1290)..(2879)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (2880)..(2943)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (2944)..(2949)

<400> SEQUENCE: 7

tctagagctg tgcacgcggc cgcgtaatac gactcactat agggcgaaga attcggatcg 60
 gactagagct cgtcacgggc tcgcgccgac gaggcgatga ggacgaaggc cgcacataat 120
 ccgtacttta cgctacatga cgactctcga aaattgtaa gggccggcat ttcggagcga 180
 gtgctgcgag ggcgcattcg cggcgctacct ggaattcctg gaatggtaag caatggccag 240
 caatgggcca ggtatggacc agcttgaatc ctggttgccg cgtcaccagg cccagcatgg 300
 tgcccagaat ggcccaccgt ggcccacgt cctaagaaac aagctgcgtc ccgcatcca 360
 aaaacgtcgt ctteggcgca cgctctcgt ggtcccccg gctggacacc ctggtggcc 420
 ctccaatgag cggcatttgc cctgtcgcg cgtgtcggca acctaatcg actccatctc 480
 tcggctccac gccgtccatc ctgtctcga cctcgtcacc tgtgctcccc ttgcccctcc 540
 ttgcccctcc ttgctccgc caccgctgc cacaatgtga ccctgctgcc cggagcgccc 600
 agcgcctatgc accgtttggg cttgtcgcgc gtgtcgcagc ctccatcgag cgattcgacc 660
 gtgtgctct ctccaccagc gttccccgcg ctctccatag tccatgctac ttgagccgt 720
 tgctcaciaa gctgccagcg gcatggctct gtcggtctcg cctctcctt tcccggaag 780
 cgctgccata caattctccg tctgccccag tccttgaggc gccgctattc ccaatcggcc 840
 atggcactgg ccagcccgat ccatgttcga tcgagcttcg acgggcccgtg agccgtctgc 900
 acggaggagc ttgcgagcct gcgaacctgg cggacctgga gaagcctggc ccatctccct 960
 ggatggagat actgggtgcg ctagcaccac ggctgcccac ggccaagctc cggcccaccc 1020
 ggaggcggga agagggttgc gttgctgtct tcggcggtcg tcagggcaa gggtaatcgt 1080
 caatgtggga aaagggctc atctccatga gattcatgac tcggacatcg tctatataag 1140

-continued

tcgagtcgcc	catcctccaa	cagccgattc	tgctcctcat	cccatcacca	ccctcgtcca	1200
caaccacgca	gttgtgtaca	tcaaaacaag	ttcgctcctt	ttacatcttc	accacaacaa	1260
cagcacatcc	tctcctttcg	gctttcaag	atg atg tat	aag aag ttc	gcc gct	1313
Met Met Tyr	Lys Lys Phe	Ala Ala				
1	5					
ctc gcc gcc	ctc gtg gct	ggc gcc tcc	gcc cag cag	gct tgc tcc	ctc	1361
Leu Ala Ala	Leu Val Ala	Gly Ala Ser	Ala Gln Gln	Ala Cys Ser	Leu	
10	15	20				
acc gct gag	aac cac cct	agc ctc acc	tgg aag cgc	tgc acc tct	ggc	1409
Thr Ala Glu	Asn His Pro	Ser Leu Thr	Trp Lys Arg	Cys Thr Ser	Gly	
25	30	35	40			
ggc agc tgc	tcg acc gtg	aac ggc gcc	gtc acc atc	gat gcc aac	tgg	1457
Gly Ser Cys	Ser Thr Val	Asn Gly Ala	Val Thr Ile	Asp Ala Asn	Trp	
45	50	55				
cgc tgg act	cac acc gtc	tcc ggc tcg	acc aac tgc	tac acc ggc	aac	1505
Arg Trp Thr	His Thr Val	Ser Gly Ser	Thr Asn Cys	Tyr Thr Gly	Asn	
60	65	70				
cag tgg gat	acc tcc ctc	tgc act gat	ggc aag agc	tgc gcc cag	acc	1553
Gln Trp Asp	Thr Ser Leu	Cys Thr Asp	Gly Lys Ser	Cys Ala Gln	Thr	
75	80	85				
tgc tgc gtc	gat ggc gct	gac tac tct	tcg acc tat	ggt atc acc	acc	1601
Cys Cys Val	Asp Gly Ala	Asp Tyr Ser	Ser Thr Tyr	Gly Ile Thr	Thr	
90	95	100				
agc ggt gac	tcc ctg aac	ctc aag ttc	gtc acc aag	cac cag tac	ggc	1649
Ser Gly Asp	Ser Leu Asn	Leu Lys Phe	Val Thr Lys	His Gln Tyr	Gly	
105	110	115	120			
acc aac gtc	ggc tcc cgt	gtc tat ctg	atg gag aac	gac acc aag	tac	1697
Thr Asn Val	Gly Ser Arg	Val Tyr Leu	Met Glu Asn	Asp Thr Lys	Tyr	
125	130	135				
cag atg ttc	gag ctc ctc	ggc aac gag	ttc acc ttc	gat gtc gat	gtc	1745
Gln Met Phe	Glu Leu Leu	Gly Asn Glu	Phe Thr Phe	Asp Val Asp	Val	
140	145	150				
tcc aac ctg	ggc tgc ggt	ctc aac ggc	gcc ctc tac	ttc gtt tcc	atg	1793
Ser Asn Leu	Gly Cys Gly	Leu Asn Gly	Ala Leu Tyr	Phe Val Ser	Met	
155	160	165				
gat gct gat	ggt ggc atg	agc aaa tac	tct ggc aac	aag gct ggc	gcc	1841
Asp Ala Asp	Gly Gly Met	Ser Lys Tyr	Ser Gly Asn	Lys Ala Gly	Ala	
170	175	180				
aag tac ggt	acc ggc tac	tgc gat gct	cag tgc ccg	cgc gac ctc	aag	1889
Lys Tyr Gly	Thr Gly Tyr	Cys Asp Ala	Gln Cys Pro	Arg Asp Leu	Lys	
185	190	195	200			
ttc atc aac	ggc gag gcc	aac aac atg	gct act gcc	ttc act cct	cac cct	1937
Phe Ile Asn	Gly Glu Ala	Asn Val Gly	Asn Trp Thr	Pro Ser Thr	Asn	
205	210	215				
gat gcc aac	gcc ggc ttc	ggc cgc tat	ggc agc tgc	tgc tct gag	atg	1985
Asp Ala Asn	Ala Gly Phe	Gly Arg Tyr	Gly Ser Cys	Cys Ser Glu	Met	
220	225	230				
gat gtc tgg	gag gcc aac	aac aac atg	gct act gcc	ttc act cct	cac cct	2033
Asp Val Trp	Glu Ala Asn	Asn Met Ala	Thr Ala Phe	Thr Pro His	Pro	
235	240	245				
tgc acc acc	ggt ggc cag	agc cgc tgc	gag gcc gac	acc tgc ggt	ggc	2081
Cys Thr Thr	Val Gly Gln	Ser Arg Cys	Glu Ala Asp	Thr Cys Gly	Gly	
250	255	260				
acc tac agc	tct gac cgc	tat gct ggt	ggt gtt tgc	gac cct gat	ggc tgc	2129
Thr Tyr Ser	Ser Asp Arg	Tyr Ala Gly	Val Cys Asp	Pro Asp Gly	Cys	
265	270	275	280			

-continued

gac ttc aac gcc tac cgc caa ggc gac aag acc ttc tac ggc aag ggc Asp Phe Asn Ala Tyr Arg Gln Gly Asp Lys Thr Phe Tyr Gly Lys Gly 285 290 295	2177
atg act gtc gac acc aac aag aag atg acc gtc gtc acc cag ttc cac Met Thr Val Asp Thr Asn Lys Lys Met Thr Val Val Thr Gln Phe His 300 305 310	2225
aag aac tcg gct ggc gtc ctc agc gag atc aag cgc ttc tac gtc cag Lys Asn Ser Ala Gly Val Leu Ser Glu Ile Lys Arg Phe Tyr Val Gln 315 320 325	2273
gac ggc aag atc att gcc aac gct gag tcc aag atc ccc ggc aac ccc Asp Gly Lys Ile Ile Ala Asn Ala Glu Ser Lys Ile Pro Gly Asn Pro 330 335 340	2321
gga aac tcc att acc cag gag tat tgc gat gcc cag aag gtc gcc ttc Gly Asn Ser Ile Thr Gln Glu Tyr Cys Asp Ala Gln Lys Val Ala Phe 345 350 355 360	2369
agt aac acc gat gac ttc aac cgc aag ggc ggt atg gct cag atg agc Ser Asn Thr Asp Asp Phe Asn Arg Lys Gly Gly Met Ala Gln Met Ser 365 370 375	2417
aag gcc ctc gca ggc ccc atg gtc ctg gtc atg tcc gtc tgg gat gac Lys Ala Leu Ala Gly Pro Met Val Leu Val Met Ser Val Trp Asp Asp 380 385 390	2465
cac tac gcc aac atg ctc tgg ctc gac tcg acc tac ccc atc gac cag His Tyr Ala Asn Met Leu Trp Leu Asp Ser Thr Tyr Pro Ile Asp Gln 395 400 405	2513
gcc ggc gcc ccc ggc gcc gag cgc ggt gct tgc ccg acc acc tcc ggt Ala Gly Ala Pro Gly Ala Glu Arg Gly Ala Cys Pro Thr Thr Ser Gly 410 415 420	2561
gtc cct gcc gag atc gag gcc cag gtc ccc aac agc aac gtc atc ttc Val Pro Ala Glu Ile Glu Ala Gln Val Pro Asn Ser Asn Val Ile Phe 425 430 435 440	2609
tcc aac atc cgt ttc ggc ccc atc ggc tcg acc gtc cct ggc ctt gac Ser Asn Ile Arg Phe Gly Pro Ile Gly Ser Thr Val Pro Gly Leu Asp 445 450 455	2657
ggc agc aac ccc ggc aac ccg acc acc acc gtc gtt cct ccc gct tct Gly Ser Asn Pro Gly Asn Pro Thr Thr Thr Val Val Pro Pro Ala Ser 460 465 470	2705
acc tcc acc tcc cgt ccg acc agc agc act agc tct ccc gtt tcg acc Thr Ser Thr Ser Arg Pro Thr Ser Ser Thr Ser Ser Pro Val Ser Thr 475 480 485	2753
ccg act ggc cag ccc ggc ggc tgc acc acc cag aag tgg ggc cag tgc Pro Thr Gly Gln Pro Gly Gly Cys Thr Thr Gln Lys Trp Gly Gln Cys 490 495 500	2801
ggc ggt atc ggc tac acc ggc tgc act aac tgc gtt gct ggc acc acc Gly Gly Ile Gly Tyr Thr Gly Cys Thr Asn Cys Val Ala Gly Thr Thr 505 510 515 520	2849
tgc act cag ctc aac ccc tgg tac agc cag gtatgtttct cttccccctt Cys Thr Gln Leu Asn Pro Trp Tyr Ser Gln 525 530	2899
ctagactcgc ttggatttga cagttgctaa catctgctca acag tgc ctg Cys Leu	2949
taaacaactc gcttcgtccg cacgacggag gagggccatg agaaagaatg ggcaacatag	3009
attctttgcg cggttgtgga ctacttgggt attttctgga tgtacatagt tttatcacgt	3069
catgaggctg tcatgtggg atgtgtatct ttttcgcttc ttcgtacata aatttacgca	3129
ttgagctttt cccccccaa aaacagttcc ctgatttget ggagtaactt gatggtaaag	3189

-continued

```

cttggtcata agctcttcaa tggaaaaaac gatacagtca tgccttgaca catcctccca 3249
aagtcttcgt ccatgacatc acggtcgatc cctaagcaca agttcaataa ccccatgtgg 3309
cgttgcccttg tcctgaaaca cagatgagat cttcagccca gccgcatcgg ccacttcctt 3369
gaactgagcc aacgagcgtt ccttcccgcc gattgagagc atcgcatagt ccttgaaggc 3429
tgcatagaga ggaatagggg gcttgtttcc ggtagtggg ctgccggaac tcggatctgt 3489
tggcgcaagg gggtcagggt tgatctgctc ggcgatgagg acgcgtccat cggggtttgt 3549
tagtgcacga gcgacattgc gcaggatggt gactgccaca gggtcggagt aatcgcgagg 3609
gatgtggcgg aggtagtaga ccagtgcacc tggaatcgat 3649

```

```

<210> SEQ ID NO 8
<211> LENGTH: 532
<212> TYPE: PRT
<213> ORGANISM: Chaetomium thermophilum

```

```

<400> SEQUENCE: 8

```

```

Met Met Tyr Lys Lys Phe Ala Ala Leu Ala Ala Leu Val Ala Gly Ala
1           5           10           15
Ser Ala Gln Gln Ala Cys Ser Leu Thr Ala Glu Asn His Pro Ser Leu
20           25           30
Thr Trp Lys Arg Cys Thr Ser Gly Gly Ser Cys Ser Thr Val Asn Gly
35           40           45
Ala Val Thr Ile Asp Ala Asn Trp Arg Trp Thr His Thr Val Ser Gly
50           55           60
Ser Thr Asn Cys Tyr Thr Gly Asn Gln Trp Asp Thr Ser Leu Cys Thr
65           70           75           80
Asp Gly Lys Ser Cys Ala Gln Thr Cys Cys Val Asp Gly Ala Asp Tyr
85           90           95
Ser Ser Thr Tyr Gly Ile Thr Thr Ser Gly Asp Ser Leu Asn Leu Lys
100          105          110
Phe Val Thr Lys His Gln Tyr Gly Thr Asn Val Gly Ser Arg Val Tyr
115          120          125
Leu Met Glu Asn Asp Thr Lys Tyr Gln Met Phe Glu Leu Leu Gly Asn
130          135          140
Glu Phe Thr Phe Asp Val Asp Val Ser Asn Leu Gly Cys Gly Leu Asn
145          150          155          160
Gly Ala Leu Tyr Phe Val Ser Met Asp Ala Asp Gly Gly Met Ser Lys
165          170          175
Tyr Ser Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp
180          185          190
Ala Gln Cys Pro Arg Asp Leu Lys Phe Ile Asn Gly Glu Ala Asn Val
195          200          205
Gly Asn Trp Thr Pro Ser Thr Asn Asp Ala Asn Ala Gly Phe Gly Arg
210          215          220
Tyr Gly Ser Cys Cys Ser Glu Met Asp Val Trp Glu Ala Asn Asn Met
225          230          235          240
Ala Thr Ala Phe Thr Pro His Pro Cys Thr Thr Val Gly Gln Ser Arg
245          250          255
Cys Glu Ala Asp Thr Cys Gly Gly Thr Tyr Ser Ser Asp Arg Tyr Ala
260          265          270
Gly Val Cys Asp Pro Asp Gly Cys Asp Phe Asn Ala Tyr Arg Gln Gly

```

-continued

275		280		285
Asp Lys Thr Phe Tyr	Gly Lys Gly Met Thr	Val Asp Thr Asn Lys Lys		
290	295	300		
Met Thr Val Val Thr	Gln Phe His Lys Asn Ser	Ala Gly Val Leu Ser		
305	310	315		320
Glu Ile Lys Arg Phe	Tyr Val Gln Asp Gly Lys	Ile Ile Ala Asn Ala		
325	330	335		
Glu Ser Lys Ile Pro	Gly Asn Pro Gly Asn Ser	Ile Thr Gln Glu Tyr		
340	345	350		
Cys Asp Ala Gln Lys	Val Ala Phe Ser Asn Thr	Asp Asp Phe Asn Arg		
355	360	365		
Lys Gly Gly Met Ala	Gln Met Ser Lys Ala	Leu Ala Gly Pro Met Val		
370	375	380		
Leu Val Met Ser Val	Trp Asp Asp His Tyr	Ala Asn Met Leu Trp Leu		
385	390	395		400
Asp Ser Thr Tyr Pro	Ile Asp Gln Ala Gly	Ala Pro Gly Ala Glu Arg		
405	410	415		
Gly Ala Cys Pro Thr	Thr Ser Gly Val Pro	Ala Glu Ile Glu Ala Gln		
420	425	430		
Val Pro Asn Ser Asn	Val Ile Phe Ser Asn	Ile Arg Phe Gly Pro Ile		
435	440	445		
Gly Ser Thr Val Pro	Gly Leu Asp Gly Ser	Asn Pro Gly Asn Pro Thr		
450	455	460		
Thr Thr Val Val Pro	Pro Ala Ser Thr Ser	Thr Ser Arg Pro Thr Ser		
465	470	475		480
Ser Thr Ser Ser Pro	Val Ser Thr Pro Thr	Gly Gln Pro Gly Gly Cys		
485	490	495		
Thr Thr Gln Lys Trp	Gly Gln Cys Gly Gly	Ile Gly Tyr Thr Gly Cys		
500	505	510		
Thr Asn Cys Val Ala	Gly Thr Thr Cys Thr	Gln Leu Asn Pro Trp Tyr		
515	520	525		
Ser Gln Cys Leu				
530				

<210> SEQ ID NO 9
<211> LENGTH: 1339
<212> TYPE: DNA
<213> ORGANISM: *Thermoascus aurantiacus*
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (17)..(122)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (123)..(177)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (178)..(236)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (237)..(296)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (297)..(449)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (450)..(508)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (509)..(573)

-continued

```

<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (574)..(647)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (648)..(745)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (746)..(806)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (807)..(1330)

<400> SEQUENCE: 9

ccgcgactg cgcac atg aag ctc ggc tct ctc gtg ctc gct ctc agc gca 52
Met Lys Leu Gly Ser Leu Val Leu Ala Leu Ser Ala
1 5 10

gct agg ctt aca ctg tcg gcc cct ctc gca gac agg aag cag gag acc 100
Ala Arg Leu Thr Leu Ser Ala Pro Leu Ala Asp Arg Lys Gln Glu Thr
15 20 25

aag cgt gcg aaa gta ttc caa t gttcgtaca tccacgtctg gcttgctggc 152
Lys Arg Ala Lys Val Phe Gln
30 35

ttactggcaa ctgacaatgg cgaag gg ttc ggt tca aac gag tcc ggt gct 203
Trp Phe Gly Ser Asn Glu Ser Gly Ala
40

gaa ttc gga agc cag aac ctt cca gga gtc gag gtcagcatgc ctgtactctc 256
Glu Phe Gly Ser Gln Asn Leu Pro Gly Val Glu
45 50 55

tgcattatat taatatctca agaggcttac tctttcgcag gga aag gat tat ata 311
Gly Lys Asp Tyr Ile
60

tgg cct gat ccc aac acc att gac aca ttg atc agc aag ggg atg aac 359
Trp Pro Asp Pro Asn Thr Ile Asp Thr Leu Ile Ser Lys Gly Met Asn
65 70 75

atc ttt cgt gtc ccc ttt atg atg gag aga ttg gtt ccc aac tca atg 407
Ile Phe Arg Val Pro Phe Met Met Glu Arg Leu Val Pro Asn Ser Met
80 85 90

acc ggc tct ccg gat ccg aac tac ctg gca gat ctc ata gcg 449
Thr Gly Ser Pro Asp Pro Asn Tyr Leu Ala Asp Leu Ile Ala
95 100 105

gtacatttca attccaccat gtttgagct gtcttcgctg tgctgacatt taatggtag 508

act gta aat gca atc acc cag aaa ggt gcc tac gcc gtc gtc gat cct 556
Thr Val Asn Ala Ile Thr Gln Lys Gly Ala Tyr Ala Val Val Asp Pro
110 115 120

cat aac tac ggc aga ta gtgaggtccc cggttctggt attgctgctg 603
His Asn Tyr Gly Arg Tyr
125

tatatctaag tagatatgtg tttctaacat ttccacgatt tcag c tac aat tct 657
Tyr Asn Ser
130

ata atc tcg agc cct tcc gat ttc cag acc ttc tgg aaa acg gtc gcc 705
Ile Ile Ser Ser Pro Ser Asp Phe Gln Thr Phe Trp Lys Thr Val Ala
135 140 145

tca cag ttt gct tcg aat cca ctg gtc atc ttc gac act a gtaagctgaa 755
Ser Gln Phe Ala Ser Asn Pro Leu Val Ile Phe Asp Thr
150 155 160

cacccgaaat taactgagtc tgagcatgtc tgacaagacg atccatgaaa g at aac 811
Asn Asn

```

-continued

```

gaa tac cac gat atg gac cag acc tta gtc ctc aat ctc aac cag gcc      859
Glu Tyr His Asp Met Asp Gln Thr Leu Val Leu Asn Leu Asn Gln Ala
165                      170                      175

gct atc gac ggc atc cgt tcc gcc gga gcc act tcc cag tac atc ttt      907
Ala Ile Asp Gly Ile Arg Ser Ala Gly Ala Thr Ser Gln Tyr Ile Phe
180                      185                      190

gtc gag ggc aat tcg tgg acc ggg gca tgg acc tgg acg aac gtg aac      955
Val Glu Gly Asn Ser Trp Thr Gly Ala Trp Thr Trp Thr Asn Val Asn
195                      200                      205                      210

gat aac atg aaa agc ctg acc gac cca tct gac aag atc ata tac gag      1003
Asp Asn Met Lys Ser Leu Thr Asp Pro Ser Asp Lys Ile Ile Tyr Glu
215                      220                      225

atg cac cag tac ctg gac tct gac gga tcc ggg aca tca gcg acc tgc      1051
Met His Gln Tyr Leu Asp Ser Asp Gly Ser Gly Thr Ser Ala Thr Cys
230                      235                      240

gta tct tcg acc atc ggt caa gag cga atc acc agc gca acg caa tgg      1099
Val Ser Ser Thr Ile Gly Gln Glu Arg Ile Thr Ser Ala Thr Gln Trp
245                      250                      255

ctc agg gcc aac ggg aag aag ggc atc atc ggc gag ttt gcg ggc gga      1147
Leu Arg Ala Asn Gly Lys Lys Gly Ile Ile Gly Glu Phe Ala Gly Gly
260                      265                      270

gcc aac gac gtc tgc gag acg gcc atc acg ggc atg ctg gac tac atg      1195
Ala Asn Asp Val Cys Glu Thr Ala Ile Thr Gly Met Leu Asp Tyr Met
275                      280                      285                      290

gcc cag aac acg gac gtc tgg act ggc gcc atc tgg tgg gcg gcc ggg      1243
Ala Gln Asn Thr Asp Val Trp Thr Gly Ala Ile Trp Trp Ala Ala Gly
295                      300                      305

ccg tgg tgg gga gac tac ata ttc tcc atg gag ccg gac aat ggc atc      1291
Pro Trp Trp Gly Asp Tyr Ile Phe Ser Met Glu Pro Asp Asn Gly Ile
310                      315                      320

gcg tat cag cag ata ctt cct att ttg act ccg tat ctt tgactgcag      1339
Ala Tyr Gln Gln Ile Leu Pro Ile Leu Thr Pro Tyr Leu
325                      330                      335

```

<210> SEQ ID NO 10

<211> LENGTH: 335

<212> TYPE: PRT

<213> ORGANISM: *Thermoascus aurantiacus*

<400> SEQUENCE: 10

```

Met Lys Leu Gly Ser Leu Val Leu Ala Leu Ser Ala Ala Arg Leu Thr
 1                      5                      10                      15

Leu Ser Ala Pro Leu Ala Asp Arg Lys Gln Glu Thr Lys Arg Ala Lys
20                      25                      30

Val Phe Gln Trp Phe Gly Ser Asn Glu Ser Gly Ala Glu Phe Gly Ser
35                      40                      45

Gln Asn Leu Pro Gly Val Glu Gly Lys Asp Tyr Ile Trp Pro Asp Pro
50                      55                      60

Asn Thr Ile Asp Thr Leu Ile Ser Lys Gly Met Asn Ile Phe Arg Val
65                      70                      75                      80

Pro Phe Met Met Glu Arg Leu Val Pro Asn Ser Met Thr Gly Ser Pro
85                      90                      95

Asp Pro Asn Tyr Leu Ala Asp Leu Ile Ala Thr Val Asn Ala Ile Thr
100                      105                      110

Gln Lys Gly Ala Tyr Ala Val Val Asp Pro His Asn Tyr Gly Arg Tyr
115                      120                      125

```


-continued

Tyr Asn Ser Ile Ile Ser Ser Pro Ser Asp Phe Gln Thr Phe Trp Lys
 130 135 140
 Thr Val Ala Ser Gln Phe Ala Ser Asn Pro Leu Val Ile Phe Asp Thr
 145 150 155 160
 Asn Asn Glu Tyr His Asp Met Asp Gln Thr Leu Val Leu Asn Leu Asn
 165 170 175
 Gln Ala Ala Ile Asp Gly Ile Arg Ser Ala Gly Ala Thr Ser Gln Tyr
 180 185 190
 Ile Phe Val Glu Gly Asn Ser Trp Thr Gly Ala Trp Thr Trp Thr Asn
 195 200 205
 Val Asn Asp Asn Met Lys Ser Leu Thr Asp Pro Ser Asp Lys Ile Ile
 210 215 220
 Tyr Glu Met His Gln Tyr Leu Asp Ser Asp Gly Ser Gly Thr Ser Ala
 225 230 235 240
 Thr Cys Val Ser Ser Thr Ile Gly Gln Glu Arg Ile Thr Ser Ala Thr
 245 250 255
 Gln Trp Leu Arg Ala Asn Gly Lys Lys Gly Ile Ile Gly Glu Phe Ala
 260 265 270
 Gly Gly Ala Asn Asp Val Cys Glu Thr Ala Ile Thr Gly Met Leu Asp
 275 280 285
 Tyr Met Ala Gln Asn Thr Asp Val Trp Thr Gly Ala Ile Trp Trp Ala
 290 295 300
 Ala Gly Pro Trp Trp Gly Asp Tyr Ile Phe Ser Met Glu Pro Asp Asn
 305 310 315 320
 Gly Ile Ala Tyr Gln Gln Ile Leu Pro Ile Leu Thr Pro Tyr Leu
 325 330 335

<210> SEQ ID NO 11
 <211> LENGTH: 2334
 <212> TYPE: DNA
 <213> ORGANISM: Acremonium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (13)..(13)
 <223> OTHER INFORMATION: N = unknown
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (715)..(797)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (798)..(856)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (857)..(1105)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1106)..(1228)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1229)..(1787)

<400> SEQUENCE: 11

tctgtctctt gtntcagaac agatctcctg gcggcctgct ttgccggtcc gaattgcat 60
 cgatgcaacg tcgattgcat acgagctaag cccgtctcgt gataaccgca aggggtcttc 120
 cgagtttctg tctgcgacc aggcattttc cgatttgtgt gcggggaccc aactgtcttc 180
 tggggagtac ctggtgacaa aagcacagat aaacagatgg atgacggtat tgctgtgata 240
 tcgccgtggc gctgaatcct ttctcttcgc taccaagata tttattcccc gttgtgaaat 300

-continued

cttctattca gcccatcca tccggcaaca cgcattctgct tttcgttccg gcattccgat	360
acctgggtcc tggagtgcct accgagcctc gcttctctggg atcgggcggt gcaccccgcc	420
aaaccctatg ccccaaacgg tacggacaag gatgccggac cccggttttg tccagaaagg	480
ttgcattcct acccacctcg ctggagccac aacatgcaga tcaccgcccg agggaggaca	540
tgtgtggtgc agggacgttg gcaactctgc tgtgtctgaa gtatatgagg ccgatggttc	600
tccttgcaaca aagcagagaa tggagttagcc agctcctcct caccagagtc gcctttgcag	660
cgtctcggca ttgcaggctc cccatcgtca gcatttcaact tctcagcaac gaac atg	717
Met	
1	
cgc tcc tca ccc ttt ctc cgc gca gct ctg gct gcc gct ctg cct ctg	765
Arg Ser Ser Pro Phe Leu Arg Ala Ala Leu Ala Ala Ala Leu Pro Leu	
5 10 15	
agc gcc cat gcc ctc gac gga aag tcg acg ag gtatgccaat cctcgtacct	817
Ser Ala His Ala Leu Asp Gly Lys Ser Thr Arg	
20 25	
ctgccctctg tagaaacaag tgaccgactg caaagacag a tac tgg gac tgc tgc	872
Tyr Trp Asp Cys Cys	
30	
aag ccg tcc tgc ggc tgg ccg gga aag gcc tcg gtg aac cag ccc gtc	920
Lys Pro Ser Cys Gly Trp Pro Gly Lys Ala Ser Val Asn Gln Pro Val	
35 40 45	
ttc tcg tgc tcg gcc gac tgg cag cgc atc agc gac ttc aac gcg aag	968
Phe Ser Cys Ser Ala Asp Trp Gln Arg Ile Ser Asp Phe Asn Ala Lys	
50 55 60 65	
tcg ggc tgc gac gga ggc tcc gcc tac tcg tgc gcc gac cag acg ccc	1016
Ser Gly Cys Asp Gly Gly Ser Ala Tyr Ser Cys Ala Asp Gln Thr Pro	
70 75 80	
tgg gcg gtc aac gac aac ttc tcg tac ggc ttc gca gcc acg gcc atc	1064
Trp Ala Val Asn Asp Asn Phe Ser Tyr Gly Phe Ala Ala Thr Ala Ile	
85 90 95	
gcc ggc ggc tcc gag tcc agc tgg tgc tgc gcc tgc tat gc	1105
Ala Gly Gly Ser Glu Ser Ser Trp Cys Cys Ala Cys Tyr Ala	
100 105 110	
gtgagttctc tgcaagccgc ttcccacccc cgctttctgt gcaggccgct tccccctac	1165
ccaccactt ccccccccc gcctctgtga tcgggcatcc gagctaagtt gcgtgtcgtc	1225
cag a ctc acc ttc aac tcg ggc ccc gtc gcg ggc aag acc atg gtg gtg	1274
Leu Thr Phe Asn Ser Gly Pro Val Ala Gly Lys Thr Met Val Val	
115 120 125	
cag tcg acc agc acc ggc ggc gac ctg ggc agc aac cag ttc gac ctc	1322
Gln Ser Thr Ser Thr Gly Gly Asp Leu Gly Ser Asn Gln Phe Asp Leu	
130 135 140	
gcc atc ccc ggc ggc ggc gtg ggc atc ttc aac ggc tgc gcc tcc cag	1370
Ala Ile Pro Gly Gly Gly Val Gly Ile Phe Asn Gly Cys Ala Ser Gln	
145 150 155	
ttc ggc ggc ctc ccc ggc gcc cag tac ggc ggc atc agc gac cgc agc	1418
Phe Gly Gly Leu Pro Gly Ala Gln Tyr Gly Gly Ile Ser Asp Arg Ser	
160 165 170	
cag tgc tcg tcc ttc ccc gcg ccg ctc cag ccg ggc tgc cag tgg cgc	1466
Gln Cys Ser Ser Phe Pro Ala Pro Leu Gln Pro Gly Cys Gln Trp Arg	
175 180 185 190	
ttc gac tgg ttc cag aac gcc gac aac ccc acc ttc acc ttc cag cgc	1514
Phe Asp Trp Phe Gln Asn Ala Asp Asn Pro Thr Phe Thr Phe Gln Arg	
195 200 205	

-continued

```

gtg cag tgc ccg tcc gag ctc acg tcc cgc acg ggc tgt aag cgc gac 1562
Val Gln Cys Pro Ser Glu Leu Thr Ser Arg Thr Gly Cys Lys Arg Asp
210                215                220

gac gac gcc agc tat ccc gtc ttc aac ccg cct agc ggt ggc tcc ccc 1610
Asp Asp Ala Ser Tyr Pro Val Phe Asn Pro Pro Ser Gly Gly Ser Pro
225                230                235

agc acc acc agc acc acc acc agc tcc ccg tcc ggt ccc acg ggc aac 1658
Ser Thr Thr Ser Thr Thr Thr Ser Ser Pro Ser Gly Pro Thr Gly Asn
240                245                250

cct cct gga ggc ggt ggc tgc act gcc cag aag tgg gcc cag tgc ggc 1706
Pro Pro Gly Gly Gly Gly Cys Thr Ala Gln Lys Trp Ala Gln Cys Gly
255                260                265                270

ggc act ggc ttc acg ggc tgc acc acc tgc gtc tgc ggc acc acc tgc 1754
Gly Thr Gly Phe Thr Gly Cys Thr Thr Cys Val Ser Gly Thr Thr Cys
275                280                285

cag gtg cag aac cag tgg tat tcc cag tgt ctg tgagcgggag gggtggtggg 1807
Gln Val Gln Asn Gln Trp Tyr Ser Gln Cys Leu
290                295

gtccgtttcc ctagggtga ggctgacgtg aactgggtcc tcttgccgc cccatcacgg 1867

gttcgtattc gcgcgcttag ggagaggagg atgcagtttg agggggccac attttgaggg 1927

ggacgcagtc tggggtcgaa gcttgctcgg tagggctgcc gtgacgtggt agagcagatg 1987

ggaccaagtg cggagctagg caggtgggtg gttgtggtgg tggcttacct tctgtaacgc 2047

aatggcatct catctcactc gcctgctccc tgattggtgg ctctgttcgg cctggcgctt 2107

tttgggaccg ctggctggaa tggattgctc cggaacgcca gggtgagctg ggctggcgcg 2167

agtagattgg ccgctccgag ctgcaacat aataaaattt tcggaccctg taagccgcac 2227

ccgaccaggt ctccattggc ggacatgcac gacgtccttc gcaggcacgg cctgcccgcc 2287

tctgatcacc cgcagtttcc gtaccgtcag accagataca agccccg 2334

```

```

<210> SEQ ID NO 12
<211> LENGTH: 297
<212> TYPE: PRT
<213> ORGANISM: Acremonium thermophilum
<221> NAME/KEY: misc_feature
<222> LOCATION: (13)..(13)

```

```

<400> SEQUENCE: 12

```

```

Met Arg Ser Ser Pro Phe Leu Arg Ala Ala Leu Ala Ala Ala Leu Pro
1          5          10          15

Leu Ser Ala His Ala Leu Asp Gly Lys Ser Thr Arg Tyr Trp Asp Cys
20                25                30

Cys Lys Pro Ser Cys Gly Trp Pro Gly Lys Ala Ser Val Asn Gln Pro
35                40                45

Val Phe Ser Cys Ser Ala Asp Trp Gln Arg Ile Ser Asp Phe Asn Ala
50                55                60

Lys Ser Gly Cys Asp Gly Gly Ser Ala Tyr Ser Cys Ala Asp Gln Thr
65                70                75                80

Pro Trp Ala Val Asn Asp Asn Phe Ser Tyr Gly Phe Ala Ala Thr Ala
85                90                95

Ile Ala Gly Gly Ser Glu Ser Ser Trp Cys Cys Ala Cys Tyr Ala Leu
100               105               110

Thr Phe Asn Ser Gly Pro Val Ala Gly Lys Thr Met Val Val Gln Ser
115               120               125

```

-continued

Thr	Ser	Thr	Gly	Gly	Asp	Leu	Gly	Ser	Asn	Gln	Phe	Asp	Leu	Ala	Ile
130					135					140					
Pro	Gly	Gly	Gly	Val	Gly	Ile	Phe	Asn	Gly	Cys	Ala	Ser	Gln	Phe	Gly
145					150					155					160
Gly	Leu	Pro	Gly	Ala	Gln	Tyr	Gly	Gly	Ile	Ser	Asp	Arg	Ser	Gln	Cys
165					170					175					
Ser	Ser	Phe	Pro	Ala	Pro	Leu	Gln	Pro	Gly	Cys	Gln	Trp	Arg	Phe	Asp
180					185					190					
Trp	Phe	Gln	Asn	Ala	Asp	Asn	Pro	Thr	Phe	Thr	Phe	Gln	Arg	Val	Gln
195					200					205					
Cys	Pro	Ser	Glu	Leu	Thr	Ser	Arg	Thr	Gly	Cys	Lys	Arg	Asp	Asp	Asp
210					215					220					
Ala	Ser	Tyr	Pro	Val	Phe	Asn	Pro	Pro	Ser	Gly	Gly	Ser	Pro	Ser	Thr
225					230					235					240
Thr	Ser	Thr	Thr	Thr	Ser	Ser	Pro	Ser	Gly	Pro	Thr	Gly	Asn	Pro	Pro
245					250					255					
Gly	Gly	Gly	Gly	Cys	Thr	Ala	Gln	Lys	Trp	Ala	Gln	Cys	Gly	Gly	Thr
260					265					270					
Gly	Phe	Thr	Gly	Cys	Thr	Thr	Cys	Val	Ser	Gly	Thr	Thr	Cys	Gln	Val
275					280					285					
Gln	Asn	Gln	Trp	Tyr	Ser	Gln	Cys	Leu							
290					295										

```

<210> SEQ ID NO 13
<211> LENGTH: 2033
<212> TYPE: DNA
<213> ORGANISM: Acremonium thermophilum
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (259)..(702)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (703)..(857)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (858)..(888)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (889)..(990)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (991)..(1268)

```

<400> SEQUENCE: 13

```

ctcgaggaga ggaaccgagt ttgaaagatg ctatatatcg atagactacc ggcgtcgct      60
cgccctgtcc gctctcttgc attccccctg ttgatgagac gagacaaaat tcttggttag    120
aaaagatccg tcgccgagat ttcaccagtg gtaagtcccg agaattggtc attcgacgtt    180
caatatgagt gtcaaagcta tgggtcctaa caaagaagga agcaagagct ttaaagagac    240
agaataacag cagcaaag atg cgt ctc cca cta ccg act ctg ctc gcc ctc      291
Met Arg Leu Pro Leu Pro Thr Leu Leu Ala Leu
1           5           10
ttg ccc tac tac ctc gaa gtg tcc gct cag ggg gca tcc gga acc ggc      339
Leu Pro Tyr Tyr Leu Glu Val Ser Ala Gln Gly Ala Ser Gly Thr Gly
15           20           25
acg aca aca cgt tac tgg gat tgc tgc aag ccg agc tgc gcg tgg cct      387
Thr Thr Thr Arg Tyr Trp Asp Cys Cys Lys Pro Ser Cys Ala Trp Pro
30           35           40

```


-continued

ctg aag ggc aat tcg ccc agc ccg gtg cag act tgc gac aag aat gac	435
Leu Lys Gly Asn Ser Pro Ser Pro Val Gln Thr Cys Asp Lys Asn Asp	
45 50 55	
agg ccg ctg aac gat ggg gga aac acc aag tcc ggc tgc gac aac ggt	483
Arg Pro Leu Asn Asp Gly Gly Asn Thr Lys Ser Gly Cys Asp Asn Gly	
60 65 70 75	
ggc ggg gcc ttc atg tgc tca tcc cag agt ccc tgg gcc gtc aat gag	531
Gly Gly Ala Phe Met Cys Ser Ser Gln Ser Pro Trp Ala Val Asn Glu	
80 85 90	
acc acc agc tac ggc tgg gca gcc gtt cgt atc gcc ggc agt acc gag	579
Thr Thr Ser Tyr Gly Trp Ala Ala Val Arg Ile Ala Gly Ser Thr Glu	
95 100 105	
tcg gcc tgg tgc tgt gcc tgc tac gag ctc acc ttc acc agt ggg ccc	627
Ser Ala Trp Cys Cys Ala Cys Tyr Glu Leu Thr Phe Thr Ser Gly Pro	
110 115 120	
gtc agt gga aag aag ctc ata gtc cag gcc acg aac act ggt gga gac	675
Val Ser Gly Lys Lys Leu Ile Val Gln Ala Thr Asn Thr Gly Gly Asp	
125 130 135	
ctt ggg agc aac cac ttt gac ctt gcg gtatgtgggg tttttcttc	722
Leu Gly Ser Asn His Phe Asp Leu Ala	
140 145	
ttcatcatcg ctctcaccat ggattcctcg gcgcaaggac caagattgag aagcgtcaat	782
gccgggttgg acacgggagc cgggatagga acacagaggc cgtttaagac cgtcagctga	842
cagcagagca attag att ccc gga ggt ggt gtt ggt cag tcc aat g	888
Ile Pro Gly Gly Gly Val Gly Gln Ser Asn	
150 155	
gtaggttctt tccctgaagt accggcaaca gcctgtgcgt tgctgtatac cccttttaat	948
catagcatct tctgtctgga tacaagccaa cccattttct ag ct tgc acg aac	1001
Ala Cys Thr Asn	
160	
cag tat ggt gcg ccc ccg aac ggc tgg gcc gac agg tat ggt ggc gtg	1049
Gln Tyr Gly Ala Pro Pro Asn Gly Trp Gly Asp Arg Tyr Gly Gly Val	
165 170 175	
cac tcg cgg agc gac tgc gac agc ttc ccc gcg gcg ctc aag gcc ggc	1097
His Ser Arg Ser Asp Cys Asp Ser Phe Pro Ala Ala Leu Lys Ala Gly	
180 185 190	
tgc tac tgg cga ttc gac tgg ttc cag gcc gcc gac aac ccg tcc gtg	1145
Cys Tyr Trp Arg Phe Asp Trp Phe Gln Gly Ala Asp Asn Pro Ser Val	
195 200 205 210	
agc ttc aaa cag gta gcc tgc ccg gca gcc atc aca gct aag agc ggc	1193
Ser Phe Lys Gln Val Ala Cys Pro Ala Ala Ile Thr Ala Lys Ser Gly	
215 220 225	
tgt act cgc cag aac gat gcc atc aac gag act ccg act ggg ccc agc	1241
Cys Thr Arg Gln Asn Asp Ala Ile Asn Glu Thr Pro Thr Gly Pro Ser	
230 235 240	
act gtg cct acc tac acc gcg tca gcc tgaaagtcgg ctggggcacc	1288
Thr Val Pro Thr Tyr Thr Ala Ser Gly	
245 250	
attgccaggg tgatggttgg gcatgtgta gtctcactca ccaggacat ttgtcgcgac	1348
ctgatcatag gcgccagggg agttgaaagg ggttgccgta cgagaagaca ttttgtcgcc	1408
gtcttactcc cagccacttc tgtacatatt caatgacatt acatagcccg caaatatggt	1468
catatatcgt ggccgccc aa accgccccgg tttgcttagg ctggagctga agtggctcgc	1528
cgatggctgt caaaggcagt cggaatattc ctcggtgctt cggcaacacg gtagctgctt	1588

-continued

```

gaaccgtacc cagcattaga acaccccccg ccgagggtt gctacgtcaa tggcggggtc 1648
tccaaccctt gcgcggcaca aaaccaacca cgcctcgtc ttttatgatg tcctcgctca 1708
aacgtcccgt gacgacactc cgctcatggt ctggctctct gatgtagaag gggtagggtca 1768
gccgatggtc gtcaccgtcg tcaatgcttc cctcaagctt cttgcggcct ttatcctcca 1828
actcttccca catgagaact ccactcttcc gccttttcc aaagccactg ccctccttgt 1888
caagggccaa aaaccaacgc cgctgatgaa tgcttccgat cgtgtttgac gcgcccgggg 1948
tatgcatttg gttcggcgca ctttttctgt cctccagctc ccttaactcc cgttccatct 2008
gagaggggtga ctcgtctact cgact 2033

```

```

<210> SEQ ID NO 14
<211> LENGTH: 251
<212> TYPE: PRT
<213> ORGANISM: Acremonium thermophilum

```

```

<400> SEQUENCE: 14

```

```

Met Arg Leu Pro Leu Pro Thr Leu Leu Ala Leu Leu Pro Tyr Tyr Leu
1           5           10           15
Glu Val Ser Ala Gln Gly Ala Ser Gly Thr Gly Thr Thr Thr Arg Tyr
20           25           30
Trp Asp Cys Cys Lys Pro Ser Cys Ala Trp Pro Leu Lys Gly Asn Ser
35           40           45
Pro Ser Pro Val Gln Thr Cys Asp Lys Asn Asp Arg Pro Leu Asn Asp
50           55           60
Gly Gly Asn Thr Lys Ser Gly Cys Asp Asn Gly Gly Gly Ala Phe Met
65           70           75           80
Cys Ser Ser Gln Ser Pro Trp Ala Val Asn Glu Thr Thr Ser Tyr Gly
85           90           95
Trp Ala Ala Val Arg Ile Ala Gly Ser Thr Glu Ser Ala Trp Cys Cys
100          105          110
Ala Cys Tyr Glu Leu Thr Phe Thr Ser Gly Pro Val Ser Gly Lys Lys
115          120          125
Leu Ile Val Gln Ala Thr Asn Thr Gly Gly Asp Leu Gly Ser Asn His
130          135          140
Phe Asp Leu Ala Ile Pro Gly Gly Gly Val Gly Gln Ser Asn Ala Cys
145          150          155          160
Thr Asn Gln Tyr Gly Ala Pro Pro Asn Gly Trp Gly Asp Arg Tyr Gly
165          170          175
Gly Val His Ser Arg Ser Asp Cys Asp Ser Phe Pro Ala Ala Leu Lys
180          185          190
Ala Gly Cys Tyr Trp Arg Phe Asp Trp Phe Gln Gly Ala Asp Asn Pro
195          200          205
Ser Val Ser Phe Lys Gln Val Ala Cys Pro Ala Ala Ile Thr Ala Lys
210          215          220
Ser Gly Cys Thr Arg Gln Asn Asp Ala Ile Asn Glu Thr Pro Thr Gly
225          230          235          240
Pro Ser Thr Val Pro Thr Tyr Thr Ala Ser Gly
245          250

```

```

<210> SEQ ID NO 15
<211> LENGTH: 2800
<212> TYPE: DNA

```


-continued

```

<213> ORGANISM: Chaetomium thermophilum
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (2786)..(2786)
<223> OTHER INFORMATION: N = unknown
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (768)..(2042)

<400> SEQUENCE: 15

ggatccaaga ccgatcccga ggattctcgg attatgtttg catctcacc tccgaaaccg      60
catgaaaaat tgaatgggc aactgtcgtc gtgtttaatg ctttgcacat catgggatca      120
tgttcaccgc ctctaatttc tcatactcca gatcctatct atcctccgca tctagccggc      180
ttcttgcttg tgatccaaag cctgatccc acgcggttc tagacgcttt agaaattaca      240
ccgaatctcc ccatgccctt cttgcaatat ctccccgacc aggaacttcg ggtgctcaac      300
atccgcgagc ttgacgacga cccttcttgg ccggcttggc atgacgactct gttcgggact      360
caatgcaact ctgggccctt caatgccgcg catgaccggt actgaggctt agccgccccca      420
atcgcttggc acggtacctt gcagacggaa tcccggggcc gttgtccgat ctgctttggt      480
tccggtagag aagcctcggg ggaagagaca cacggacaca acgattgcgg gccccaatgc      540
gctgctccta attgaggctc cgaggctcgtg tgccgtgtgg agaggccgcg actgggtctg      600
gggtgccggg gattgcccgg atgaagataa tctgggtgca accgtggata cataaaaggg      660
agtagttctc ccctctgtga aaccttcttc ccaggatctc tcctcgctc taagagtcca      720
aagtcattca agacatccta cagcgggggtc agtgagattc cataatc atg act cgc      776
Met Thr Arg
1

aag ttc gca ctc gtt ccc ctc ctt ctg ggt ctt gcc tcg gcc cag aaa      824
Lys Phe Ala Leu Val Pro Leu Leu Leu Gly Leu Ala Ser Ala Gln Lys
5 10 15

ccc ggc aac act cca gaa gtc cac ccc aag atc acc act tac cgc tgc      872
Pro Gly Asn Thr Pro Glu Val His Pro Lys Ile Thr Thr Tyr Arg Cys
20 25 30 35

agc cac cgc cag gga tgc cgc ccg gag acg aac tac atc gtc ctc gac      920
Ser His Arg Gln Gly Cys Arg Pro Glu Thr Asn Tyr Ile Val Leu Asp
40 45 50

tcc ctc acc cat ccc gtg cac cag ttg aac tcc aac gcg aac tgc ggc      968
Ser Leu Thr His Pro Val His Gln Leu Asn Ser Asn Ala Asn Cys Gly
55 60 65

gac tgg ggt aac ccg ccc ccg cgc agc gtc tgc cct gat gtc gag acc      1016
Asp Trp Gly Asn Pro Pro Pro Arg Ser Val Cys Pro Asp Val Glu Thr
70 75 80

tgc gcg cag aat tgc atc atg gag ggc atc caa gac tac tcc acc tac      1064
Cys Ala Gln Asn Cys Ile Met Glu Gly Ile Gln Asp Tyr Ser Thr Tyr
85 90 95

ggc gtg acc acc tct ggc tct tcc ctt cgc ctg aag cag atc cac cag      1112
Gly Val Thr Thr Ser Gly Ser Ser Leu Arg Leu Lys Gln Ile His Gln
100 105 110 115

ggc cgc gtc acc tct cct cgt gtc tac ctc ctc gac aag acg gag cag      1160
Gly Arg Val Thr Ser Pro Arg Val Tyr Leu Leu Asp Lys Thr Glu Gln
120 125 130

cag tat gag atg atg cgt ctc acc ggc ttc gag ttc act ttc gac gtc      1208
Gln Tyr Glu Met Met Arg Leu Thr Gly Phe Glu Phe Thr Phe Asp Val
135 140 145

gac acc acc aag ctc ccc tgc ggc atg aac gct gcg ctc tat ctc tcc      1256

```

-continued

Asp Thr Thr Lys Leu Pro Cys Gly Met Asn Ala Ala Leu Tyr Leu Ser	
150	155 160
gag atg gac gct acc ggc gct cgc tcc cgc ctc aac cct ggc ggt gcc	1304
Glu Met Asp Ala Thr Gly Ala Arg Ser Arg Leu Asn Pro Gly Gly Ala	
165	170 175
tac tac ggc acg ggt tac tgc gat gca cag tgc ttc gtc acc ccc ttc	1352
Tyr Tyr Gly Thr Gly Tyr Cys Asp Ala Gln Cys Phe Val Thr Pro Phe	
180	185 190 195
atc aat ggc atc ggc aac atc gag ggc aag ggc tgc tgc tgc aac gag	1400
Ile Asn Gly Ile Gly Asn Ile Glu Gly Lys Gly Ser Cys Cys Asn Glu	
200	205 210
atg gac att tgg gag gcc aac tgc cgt agt cag tcc att gct ccg cac	1448
Met Asp Ile Trp Glu Ala Asn Ser Arg Ser Gln Ser Ile Ala Pro His	
215	220 225
ccc tgc aac aag cag ggt ctg tac atg tgc tcc ggc cag gag tgc gag	1496
Pro Cys Asn Lys Gln Gly Leu Tyr Met Cys Ser Gly Gln Glu Cys Glu	
230	235 240
ttc gac ggc gtc tgc gac gag tgg gga tgc aca tgg aac ccg tac aag	1544
Phe Asp Gly Val Cys Asp Glu Trp Gly Cys Thr Trp Asn Pro Tyr Lys	
245	250 255
gtc aac gtt acc gac tac tat ggc cgc ggt ccg cag ttc aag gtc gac	1592
Val Asn Val Thr Asp Tyr Tyr Gly Arg Gly Pro Gln Phe Lys Val Asp	
260	265 270 275
acg acc cgt ccc ttc acc gtc atc aca cag ttt cca gcc gac cag aac	1640
Thr Thr Arg Pro Phe Thr Val Ile Thr Gln Phe Pro Ala Asp Gln Asn	
280	285 290
ggc aag ctg acg tgc atc cat cgc atg tat gtg caa gat ggc aag ttg	1688
Gly Lys Leu Thr Ser Ile His Arg Met Tyr Val Gln Asp Gly Lys Leu	
295	300 305
atc gag gcg cat acc gtc aac ctg ccg ggt tat cct caa gtg aac gcg	1736
Ile Glu Ala His Thr Val Asn Leu Pro Gly Tyr Pro Gln Val Asn Ala	
310	315 320
ctg aac gat gac ttc tgc cgt gcc acg gga gcc gcg acg aag tat ctt	1784
Leu Asn Asp Asp Phe Cys Arg Ala Thr Gly Ala Ala Thr Lys Tyr Leu	
325	330 335
gaa ctg ggt gcc act gcg ggt atg ggc gag gct ctg agg cgt ggt atg	1832
Glu Leu Gly Ala Thr Ala Gly Met Gly Glu Ala Leu Arg Arg Gly Met	
340	345 350 355
gtg ctg gct atg agc atc tgg tgg gat gag agc ggc ttc atg aac tgg	1880
Val Leu Ala Met Ser Ile Trp Trp Asp Glu Ser Gly Phe Met Asn Trp	
360	365 370
ctt gat agc ggc gag tct ggg ccg tgc aac ccg aac gag ggt aac cca	1928
Leu Asp Ser Gly Glu Ser Gly Pro Cys Asn Pro Asn Glu Gly Asn Pro	
375	380 385
cag aac att cgc cag att gag ccc gag ccg gag gtt acc tat agc aac	1976
Gln Asn Ile Arg Gln Ile Glu Pro Glu Pro Glu Val Thr Tyr Ser Asn	
390	395 400
ctg cgc tgg ggt gag att ggg tgc act tat aag cac aat ctg aag ggc	2024
Leu Arg Trp Gly Glu Ile Gly Ser Thr Tyr Lys His Asn Leu Lys Gly	
405	410 415
ggg tgg act ggc agg aac taagtgttg ggattagagc ctgtgattgg	2072
Gly Trp Thr Gly Arg Asn	
420	425
atacctgtgg gttaaaccggg gctcggtttg agagggttg tgaatttat ttctcgtaca	2132
tagttggcgt cttggcgaat atatgcccc aggactttga tccagtcttc gtccatttct	2192
ctgtgactta gttggtgcaa gtatcattgt tatgtcctgg gtgagacaaa gcaatctctt	2252

-continued

```

cagtggatcat gggtaaataa tctacaggct gtgaatggcg ttgcgtcagc ctcattaact 2312
taaacgattg gactcccctt ttctaataca tcgccgttgc cgtgtaactc tcttagatct 2372
cttgttgtat atggcttcaa ctogaagtga agaaaaatgg atacggcgac ctctttgtgc 2432
caattttctt gctgttcttc cggattgac cctcggcaag acaactatgg ccaatattct 2492
ggtatagtcg gcagttagtg ttgtgctgta caagtcgtgc gggagcaata ctcaacagcc 2552
gcccttaata tggttattta cgccacgacg cacttcatta cacggctttg gggggtatat 2612
attccgttca actctatccc tcattcgggtg tgattgaacg tctccaacag tgaaagtata 2672
agtctgacaa aaatgcccaa ccgccatgcc actgatgatc ctggtgagat gctcgtggtc 2732
tataacatcc tgtctaagtg ttacctcctt aatgtagcc ccagttctgc tctncttctc 2792
tcgacagc 2800

```

```

<210> SEQ ID NO 16
<211> LENGTH: 425
<212> TYPE: PRT
<213> ORGANISM: Chaetomium thermophilum
<221> NAME/KEY: misc_feature
<222> LOCATION: (2786)..(2786)

```

```

<400> SEQUENCE: 16

```

```

Met Thr Arg Lys Phe Ala Leu Val Pro Leu Leu Leu Gly Leu Ala Ser
1          5          10          15
Ala Gln Lys Pro Gly Asn Thr Pro Glu Val His Pro Lys Ile Thr Thr
20          25          30
Tyr Arg Cys Ser His Arg Gln Gly Cys Arg Pro Glu Thr Asn Tyr Ile
35          40          45
Val Leu Asp Ser Leu Thr His Pro Val His Gln Leu Asn Ser Asn Ala
50          55          60
Asn Cys Gly Asp Trp Gly Asn Pro Pro Pro Arg Ser Val Cys Pro Asp
65          70          75          80
Val Glu Thr Cys Ala Gln Asn Cys Ile Met Glu Gly Ile Gln Asp Tyr
85          90          95
Ser Thr Tyr Gly Val Thr Thr Ser Gly Ser Ser Leu Arg Leu Lys Gln
100         105         110
Ile His Gln Gly Arg Val Thr Ser Pro Arg Val Tyr Leu Leu Asp Lys
115         120         125
Thr Glu Gln Gln Tyr Glu Met Met Arg Leu Thr Gly Phe Glu Phe Thr
130         135         140
Phe Asp Val Asp Thr Thr Lys Leu Pro Cys Gly Met Asn Ala Ala Leu
145         150         155         160
Tyr Leu Ser Glu Met Asp Ala Thr Gly Ala Arg Ser Arg Leu Asn Pro
165         170         175
Gly Gly Ala Tyr Tyr Gly Thr Gly Tyr Cys Asp Ala Gln Cys Phe Val
180         185         190
Thr Pro Phe Ile Asn Gly Ile Gly Asn Ile Glu Gly Lys Gly Ser Cys
195         200         205
Cys Asn Glu Met Asp Ile Trp Glu Ala Asn Ser Arg Ser Gln Ser Ile
210         215         220
Ala Pro His Pro Cys Asn Lys Gln Gly Leu Tyr Met Cys Ser Gly Gln
225         230         235         240

```

-continued

Glu Cys Glu Phe Asp Gly Val Cys Asp Glu Trp Gly Cys Thr Trp Asn
 245 250 255
 Pro Tyr Lys Val Asn Val Thr Asp Tyr Tyr Gly Arg Gly Pro Gln Phe
 260 265 270
 Lys Val Asp Thr Thr Arg Pro Phe Thr Val Ile Thr Gln Phe Pro Ala
 275 280 285
 Asp Gln Asn Gly Lys Leu Thr Ser Ile His Arg Met Tyr Val Gln Asp
 290 295 300
 Gly Lys Leu Ile Glu Ala His Thr Val Asn Leu Pro Gly Tyr Pro Gln
 305 310 315 320
 Val Asn Ala Leu Asn Asp Asp Phe Cys Arg Ala Thr Gly Ala Ala Thr
 325 330 335
 Lys Tyr Leu Glu Leu Gly Ala Thr Ala Gly Met Gly Glu Ala Leu Arg
 340 345 350
 Arg Gly Met Val Leu Ala Met Ser Ile Trp Trp Asp Glu Ser Gly Phe
 355 360 365
 Met Asn Trp Leu Asp Ser Gly Glu Ser Gly Pro Cys Asn Pro Asn Glu
 370 375 380
 Gly Asn Pro Gln Asn Ile Arg Gln Ile Glu Pro Glu Pro Glu Val Thr
 385 390 395 400
 Tyr Ser Asn Leu Arg Trp Gly Glu Ile Gly Ser Thr Tyr Lys His Asn
 405 410 415
 Leu Lys Gly Gly Trp Thr Gly Arg Asn
 420 425

<210> SEQ ID NO 17
 <211> LENGTH: 1943
 <212> TYPE: DNA
 <213> ORGANISM: *Thermoascus aurantiacus*
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (13)..(256)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (257)..(329)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (330)..(370)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (371)..(444)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (445)..(493)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (494)..(561)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (562)..(683)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (684)..(786)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (787)..(932)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (933)..(1001)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1002)..(1090)
 <220> FEATURE:
 <221> NAME/KEY: Intron

-continued

```

<222> LOCATION: (1091)..(1155)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1156)..(1174)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1175)..(1267)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1268)..(1295)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1296)..(1361)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1362)..(1451)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1452)..(1551)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1552)..(1617)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1618)..(1829)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1830)..(1922)

<400> SEQUENCE: 17

ccgcggaag cc atg gtt cga cca acg atc cta ctt act tca ctc ctg cta 51
Met Val Arg Pro Thr Ile Leu Leu Thr Ser Leu Leu Leu
1 5 10

gct ccc ttc gca gct gcg agc cct atc ctc gag gaa cgc caa gct gca 99
Ala Pro Phe Ala Ala Ala Ser Pro Ile Leu Glu Glu Arg Gln Ala Ala
15 20 25

cag agt gtc gac caa ctg atc aag gct cgc ggc aag gtg tac ttt ggc 147
Gln Ser Val Asp Gln Leu Ile Lys Ala Arg Gly Lys Val Tyr Phe Gly
30 35 40 45

gtc gcc acg gac caa aac cgg ctg acg acc ggc aag aat gcg gct atc 195
Val Ala Thr Asp Gln Asn Arg Leu Thr Thr Gly Lys Asn Ala Ala Ile
50 55 60

atc cag gct gat ttc ggc cag gtc acg ccg gag aat agt atg aaa tgg 243
Ile Gln Ala Asp Phe Gly Gln Val Thr Pro Glu Asn Ser Met Lys Trp
65 70 75

gac gct act gaa c gtgcgtgaga aagataattt gatttttttc ttctatgacc 296
Asp Ala Thr Glu
80

gctcggaccg ttctgactag gtttataata tag ct tct caa gga aac ttc aac 349
Pro Ser Gln Gly Asn Phe Asn
85

ttt gcc ggt gct gat tac ctt gtacgtacat acgaccactt gacgtttctt 400
Phe Ala Gly Ala Asp Tyr Leu
90 95

gcacgcaact gcgattgagg agaagataact aatcttcttg aaag gtc aat tgg gcc 456
Val Asn Trp Ala

cag caa aat gga aag ctg atc cgt ggc cat act ctt g gttagtagaa 503
Gln Gln Asn Gly Lys Leu Ile Arg Gly His Thr Leu
100 105 110

cgccaacctg cttcctaac ttactgaaga aggaaaaccg aattgaccgt cccccaag 561

ta tgg cac tcg cag ctg ccc tcg tgg gtg agc tcc atc acc gac aag 608
Val Trp His Ser Gln Leu Pro Ser Trp Val Ser Ser Ile Thr Asp Lys
115 120 125

```

-continued

aat acg ctg acc aac gtg atg aaa aat cac atc acc acc ttg atg acc Asn Thr Leu Thr Asn Val Met Lys Asn His Ile Thr Thr Leu Met Thr 130 135 140	656
cgg tac aag ggc aag atc cgt gca tgg gtcagtcac ctaccctaag Arg Tyr Lys Gly Lys Ile Arg Ala Trp 145 150	703
ctgctgtttca atgaagagac aaataagaac acacgtatatt gcccgggcgt ttcagaatca	763
gaactgacag aatcactgaa tag gac gtg gtg aac gag gca ttc aac gag gat Asp Val Val Asn Glu Ala Phe Asn Glu Asp 155 160	816
ggc tcc ctc cgc cag act gtc ttc ctc aac gtc atc ggg gag gat tac Gly Ser Leu Arg Gln Thr Val Phe Leu Asn Val Ile Gly Glu Asp Tyr 165 170 175	864
atc ccg att gct ttc cag acc gcc cgc gcc gct gac ccg aat gcc aag Ile Pro Ile Ala Phe Gln Thr Ala Arg Ala Ala Asp Pro Asn Ala Lys 180 185 190	912
ctg tac atc aac gat tac aa gtaagattta aggctcagtg atattccatt Leu Tyr Ile Asn Asp Tyr Asn 195 200	962
tagtgtgaga agcattgctt atgagcatct gtattacag c ctc gac agt gcc tcg Leu Asp Ser Ala Ser 205	1017
tac ccc aag acg cag gcc att gtc aac cgc gtc aag caa tgg cgt gca Tyr Pro Lys Thr Gln Ala Ile Val Asn Arg Val Lys Gln Trp Arg Ala 210 215 220	1065
gct gga gtc ccg att gac ggc ata g gtatgtctct ctttctgttt Ala Gly Val Pro Ile Asp Gly Ile 225 230	1110
gtgatgtgac cgatttgaaa ccagtctaac gttagctggg tctag ga tcg caa acg Gly Ser Gln Thr	1166
cac ctc ag gtaaataatc gggaatgcct cggagaataa aagagaaaa His Leu Ser 235	1214
aaatgattgt cttatcagat cgtatcgact gactcatggc ttgtccaaaa tag c gct Ala	1271
ggt cag gga gcc ggt gtt cta caa taagtgcccc cctcccctat tttttactat Gly Gln Gly Ala Gly Val Leu Gln 240 245	1325
tattgagaga gcggaatagg ctgacaaccc caaacg gct ctt ccg ctc ctt gct Ala Leu Pro Leu Leu Ala 250	1379
agt gcc gga act ccc gag gtc gct atc acg gaa ctg gac gtg gct ggt Ser Ala Gly Thr Pro Glu Val Ala Ile Thr Glu Leu Asp Val Ala Gly 255 260 265	1427
gct agc ccg acg gat tac gtc aat gtatgtacct cgttgteect atcccccttg Ala Ser Pro Thr Asp Tyr Val Asn 270 275	1481
gatactttgt ataattatta tcttcccgga gctgttgat cagatctgac gatcatttct	1541
cgtttttttag gtc gtg aac gct tgc ctc aac gtg cag tcc tgc gtg ggc Val Val Asn Ala Cys Leu Asn Val Gln Ser Cys Val Gly 280 285	1590
atc acc gtc tgg ggc gtg gca gat ccg gtaagcgcgg ttcttccgta Ile Thr Val Trp Gly Val Ala Asp Pro 290 295	1637
ctccgtaccc aactagagtt cgggctgtca cgtcatgtct tagtctctt cagtcaggcc	1697

-continued

```

aaggccaaga cacaggacct gaaacgggca ggacgagct gctagcagcc caagaagcag 1757
ccacatgatg catgattatt attattatat ctccgagttc tgggctaacg attggtgata 1817
ataaataaat ag gac tca tgg cgt gct agc acg acg cct ctc ctc ttc gac 1868
Asp Ser Trp Arg Ala Ser Thr Thr Pro Leu Leu Phe Asp
300                305                310
ggc aac ttc aac ccg aag ccg gcg tac aac gcc att gtg cag gac ctg 1916
Gly Asn Phe Asn Pro Lys Pro Ala Tyr Asn Ala Ile Val Gln Asp Leu
315                320                325
cag cag tgagtataga ccggtggatc c 1943
Gln Gln

```

```

<210> SEQ ID NO 18
<211> LENGTH: 329
<212> TYPE: PRT
<213> ORGANISM: Thermoascus aurantiacus

```

```

<400> SEQUENCE: 18

```

```

Met Val Arg Pro Thr Ile Leu Leu Thr Ser Leu Leu Leu Ala Pro Phe
1                5                10                15
Ala Ala Ala Ser Pro Ile Leu Glu Glu Arg Gln Ala Ala Gln Ser Val
20                25                30
Asp Gln Leu Ile Lys Ala Arg Gly Lys Val Tyr Phe Gly Val Ala Thr
35                40                45
Asp Gln Asn Arg Leu Thr Thr Gly Lys Asn Ala Ala Ile Ile Gln Ala
50                55                60
Asp Phe Gly Gln Val Thr Pro Glu Asn Ser Met Lys Trp Asp Ala Thr
65                70                75                80
Glu Pro Ser Gln Gly Asn Phe Asn Phe Ala Gly Ala Asp Tyr Leu Val
85                90                95
Asn Trp Ala Gln Gln Asn Gly Lys Leu Ile Arg Gly His Thr Leu Val
100               105               110
Trp His Ser Gln Leu Pro Ser Trp Val Ser Ser Ile Thr Asp Lys Asn
115               120               125
Thr Leu Thr Asn Val Met Lys Asn His Ile Thr Thr Leu Met Thr Arg
130               135               140
Tyr Lys Gly Lys Ile Arg Ala Trp Asp Val Val Asn Glu Ala Phe Asn
145               150               155               160
Glu Asp Gly Ser Leu Arg Gln Thr Val Phe Leu Asn Val Ile Gly Glu
165               170               175
Asp Tyr Ile Pro Ile Ala Phe Gln Thr Ala Arg Ala Ala Asp Pro Asn
180               185               190
Ala Lys Leu Tyr Ile Asn Asp Tyr Asn Leu Asp Ser Ala Ser Tyr Pro
195               200               205
Lys Thr Gln Ala Ile Val Asn Arg Val Lys Gln Trp Arg Ala Ala Gly
210               215               220
Val Pro Ile Asp Gly Ile Gly Ser Gln Thr His Leu Ser Ala Gly Gln
225               230               235               240
Gly Ala Gly Val Leu Gln Ala Leu Pro Leu Leu Ala Ser Ala Gly Thr
245               250               255
Pro Glu Val Ala Ile Thr Glu Leu Asp Val Ala Gly Ala Ser Pro Thr
260               265               270
Asp Tyr Val Asn Val Val Asn Ala Cys Leu Asn Val Gln Ser Cys Val
275               280               285

```

-continued

Gly Ile Thr Val Trp Gly Val Ala Asp Pro Asp Ser Trp Arg Ala Ser
 290 295 300

Thr Thr Pro Leu Leu Phe Asp Gly Asn Phe Asn Pro Lys Pro Ala Tyr
 305 310 315 320

Asn Ala Ile Val Gln Asp Leu Gln Gln
 325

<210> SEQ ID NO 19
 <211> LENGTH: 2955
 <212> TYPE: DNA
 <213> ORGANISM: Acremonium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1335)..(1671)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1672)..(1806)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1807)..(2032)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (2033)..(2117)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (2118)..(2802)

<400> SEQUENCE: 19

tctagagctg tcgacgcggc cgcgtaatac gactcaactat agggcgaaga attcggatca 60
 cgtttgcttc agcaagtcgt tcgctacgac accacgtcca tgatggaggc cctgattcaa 120
 tcataccaag gacggggcat gatggctgat ggctggactc gaagtgagtg gcccgaggct 180
 gaatthtctt tcccgttctc tacagtcctt ccctcagcga cacatccgca gttttgacag 240
 cggaaatcgt caggatgctc cgccttctct cgcaacctga gtgccaggc gtctcggcca 300
 ccgtctctta tatatggccg ctgggtccgc ctttcgatcg gttttcgatt tggctctctc 360
 tagttccctc agctgacctg ggatctcgtc tgtggctccg aaacctcacc atcccagacg 420
 agcaagttct ccgcagttca cctcagctca tccggccctt ggtagcatcg cagcagcccc 480
 agacgaaggc accaagaag catactatat attaggctaa atcgagcccc acgtggaata 540
 tttgccatcg aggaggggtg gttgggcttc ttgtcctcgc aggtgctgcg cctgtaccta 600
 cctggtgctc cagctggtgc tcccgtggt gctgttccag tcgccgtctg gcccgaatgc 660
 tctgtatctc ggttcgtccc gcaactcctt cgccaagcgc taccaatgct ttgacgaacc 720
 cggtaaattt gcagtgacc tgcagctggg caaacccgca gtgggaacca cagacctggt 780
 tcgttcgaca cactccaatc gcaacccgc ccgcgcaaac cttgcaccac atgtcgcccc 840
 tttcccagtt gggctccctga agacacggag ccaacttccgt gatcgctcggc tcccgaagcc 900
 gacagtcgga cgctgcaata ggatgccagc acccgtggat ccaagggcca gtgaccccaa 960
 ctctttcgcg gtattctggc cctcccagaag gtatgccagg acttcctgt ctttgctacc 1020
 accagctctc ctccacggcg gaacggatac gccgtctcgc cggctcttgc tcgacaacat 1080
 gcgagggggc gcgaaggcta ggttgtgacg atgcgacggt gcgatgtcac catttggcag 1140
 tgatgttttc cgttgtcccc ttctccacc tgcccggtt cctcaaagac gcccgaacca 1200
 taaatacgat gcgacgcaa ccttcatgtg ttcgtggcat cttgcctgac cagtctcagc 1260
 aagaaacctg tggcggcgcg attgtcttga cttctgatt gaaaacggat ctgcgtctctc 1320

-continued

ctcgatagcc gacc atg cgc gcc aag caa ctc ctg gcg gcc ggc ctg ctg	1370
Met Arg Ala Lys Gln Leu Leu Ala Ala Gly Leu Leu	
1 5 10	
gcc ccc gcg tcc gtc tcg gcc cag ctc aac agc ctc gcc gtg gcg gct	1418
Ala Pro Ala Ser Val Ser Ala Gln Leu Asn Ser Leu Ala Val Ala Ala	
15 20 25	
ggc ctc aag tac ttc ggc acg gcc gtg cgg gag gcc aac gtc aac ggc	1466
Gly Leu Lys Tyr Phe Gly Thr Ala Val Arg Glu Ala Asn Val Asn Gly	
30 35 40	
gac gcc acc tac atg tcg tac gtc aac aac aag tcc gag ttc ggc cag	1514
Asp Ala Thr Tyr Met Ser Tyr Val Asn Asn Lys Ser Glu Phe Gly Gln	
45 50 55 60	
gtg acg ccc gag aac ggc cag aag tgg gat tcc acc gag ccc agc cag	1562
Val Thr Pro Glu Asn Gly Gln Lys Trp Asp Ser Thr Glu Pro Ser Gln	
65 70 75	
ggc cag ttc agc tac agc cag ggc gac atc gtc ccc ggc gtc gcg aag	1610
Gly Gln Phe Ser Tyr Ser Gln Gly Asp Ile Val Pro Gly Val Ala Lys	
80 85 90	
aag aac ggc cag gtg ctg cgc tgc cac acc ctg gtg tgg tac agc cag	1658
Lys Asn Gly Gln Val Leu Arg Cys His Thr Leu Val Trp Tyr Ser Gln	
95 100 105	
ctc ccc agc tgg g gtcagtgact ctctctttct ctctgtcttt ctctttgtct	1711
Leu Pro Ser Trp	
110	
ttctctcttt ctctctctct ctctctctct ctctctctct ctctctccca tccagcateg	1771
actgctgac ttgctgacca gaagctcgtg tgcag tg tca tcc gga agt tgg	1823
Val Ser Ser Gly Ser Trp	
115	
acc cgc gcg acg ctt cag tcc gtc atc gag acg cac atc tcg aac gtg	1871
Thr Arg Ala Thr Leu Gln Ser Val Ile Glu Thr His Ile Ser Asn Val	
120 125 130	
atg ggc cac tac aag ggc cag tgc tac gcc tgg gac gtg gtc aac gag	1919
Met Gly His Tyr Lys Gly Gln Cys Tyr Ala Trp Asp Val Val Asn Glu	
135 140 145 150	
gcc atc aac gac gac ggc acg tgg cgg acc agc gtc ttc tac aac acc	1967
Ala Ile Asn Asp Asp Gly Thr Trp Arg Thr Ser Val Phe Tyr Asn Thr	
155 160 165	
ttc aac acc gac tac ctg gcc att gcc ttc aac gcc gcg aag aag gcc	2015
Phe Asn Thr Asp Tyr Leu Ala Ile Ala Phe Asn Ala Ala Lys Lys Ala	
170 175 180	
gat gcg ggc gcg aag ct gtaggtgctg gcctttacgt tgccgcagcg	2062
Asp Ala Gly Ala Lys Leu	
185	
cacctccgcg acatgagccc cagagcgcgt ggctaatagt tcctcagca cgcag g	2118
tac tac aac gac tac aat ctc gag tac aac ggc gcc aag acc aac acg	2166
Tyr Tyr Asn Asp Tyr Asn Leu Glu Tyr Asn Gly Ala Lys Thr Asn Thr	
190 195 200	
gcc gtg cag ctg gtg cag atc gtg cag cag gcc ggc gcg ccc atc gac	2214
Ala Val Gln Leu Val Gln Ile Val Gln Gln Ala Gly Ala Pro Ile Asp	
205 210 215 220	
ggg gtg ggc ttc cag ggc cac ctg atc gtg ggg tca acg ccg tcg cgc	2262
Gly Val Gly Phe Gln Gly His Leu Ile Val Gly Ser Thr Pro Ser Arg	
225 230 235	
agc tcc ctg gcc acg gcg ctg aag cgc ttc acg gcg ctt ggc ctg gag	2310
Ser Ser Leu Ala Thr Ala Leu Lys Arg Phe Thr Ala Leu Gly Leu Glu	
240 245 250	

-continued

```

gtg gcg tac acg gag ctg gac atc cgg cac tcg agc ctg ccg ccg tcg      2358
Val Ala Tyr Thr Glu Leu Asp Ile Arg His Ser Ser Leu Pro Pro Ser
255                      260                      265

tcg gcg gcg ctg gcg acg cag ggc aac gac ttc gcc agc gtg gtg ggc      2406
Ser Ala Ala Leu Ala Thr Gln Gly Asn Asp Phe Ala Ser Val Val Gly
270                      275                      280

tcg tgc ctc gac gtg gcg ggc tgc gtg ggc atc acc atc tgg ggg ttc      2454
Ser Cys Leu Asp Val Ala Gly Cys Val Gly Ile Thr Ile Trp Gly Phe
285                      290                      295                      300

acg gac aag tac agc tgg gtg ccc gac acg ttc ccc gcc tcg gcc gcg      2502
Thr Asp Lys Tyr Ser Trp Val Pro Asp Thr Phe Pro Gly Ser Gly Ala
305                      310                      315

gcg ctg ctg tac gac gcg aac tac agc aag aag ccg gcg tgg acg tcg      2550
Ala Leu Leu Tyr Asp Ala Asn Tyr Ser Lys Lys Pro Ala Trp Thr Ser
320                      325                      330

gtc tcg tcg gtg ctg gcg gcc aag gcg acg aac ccg ccc gcc gcc ggg      2598
Val Ser Ser Val Leu Ala Ala Lys Ala Thr Asn Pro Pro Gly Gly Gly
335                      340                      345

aac cca ccc ccc gtc acc acc acg acc acg acc acc acc acg tcg aag      2646
Asn Pro Pro Pro Val Thr Thr Thr Thr Thr Thr Thr Thr Thr Ser Lys
350                      355                      360

ccg tcg cag ccc acc acc acg acc acg acc acc agc ccg cag ggt ccg      2694
Pro Ser Gln Pro Thr Thr Thr Thr Thr Thr Thr Ser Pro Gln Gly Pro
365                      370                      375                      380

cag cag acg cac tgg gcc cag tgc ggc ggg atc gcc tgg acg ggg ccg      2742
Gln Gln Thr His Trp Gly Gln Cys Gly Gly Ile Gly Trp Thr Gly Pro
385                      390                      395

cag tcg tgc cag agc ccg tgg acg tgc cag aag cag aac gac tgg tac      2790
Gln Ser Cys Gln Ser Pro Trp Thr Cys Gln Lys Gln Asn Asp Trp Tyr
400                      405                      410

tct cag tgc ctg tgaccaccac ggctgaccag ctgccattcc gaccacgggg      2842
Ser Gln Cys Leu
415

cccgactac aaaaagaggg gacgggtgtaa ataaagagcc gaacgggtct acgtacactg      2902

ttttgacctt ttctccgcag acgtatatta tcaattatag ttggatttct aga      2955

```

<210> SEQ ID NO 20

<211> LENGTH: 416

<212> TYPE: PRT

<213> ORGANISM: Acremonium thermophilum

<400> SEQUENCE: 20

```

Met Arg Ala Lys Gln Leu Leu Ala Ala Gly Leu Leu Ala Pro Ala Ser
1                      5                      10                      15

Val Ser Ala Gln Leu Asn Ser Leu Ala Val Ala Ala Gly Leu Lys Tyr
20                      25                      30

Phe Gly Thr Ala Val Arg Glu Ala Asn Val Asn Gly Asp Ala Thr Tyr
35                      40                      45

Met Ser Tyr Val Asn Asn Lys Ser Glu Phe Gly Gln Val Thr Pro Glu
50                      55                      60

Asn Gly Gln Lys Trp Asp Ser Thr Glu Pro Ser Gln Gly Gln Phe Ser
65                      70                      75                      80

Tyr Ser Gln Gly Asp Ile Val Pro Gly Val Ala Lys Lys Asn Gly Gln
85                      90                      95

Val Leu Arg Cys His Thr Leu Val Trp Tyr Ser Gln Leu Pro Ser Trp

```


-continued

100	105	110													
Val Ser Ser Gly Ser Trp Thr Arg Ala Thr Leu Gln Ser Val Ile Glu															
115	120	125													
Thr His Ile Ser Asn Val Met Gly His Tyr Lys Gly Gln Cys Tyr Ala															
130	135	140													
Trp Asp Val Val Asn Glu Ala Ile Asn Asp Asp Gly Thr Trp Arg Thr															
145	150	155	160												
Ser Val Phe Tyr Asn Thr Phe Asn Thr Asp Tyr Leu Ala Ile Ala Phe															
165	170	175													
Asn Ala Ala Lys Lys Ala Asp Ala Gly Ala Lys Leu Tyr Tyr Asn Asp															
180	185	190													
Tyr Asn Leu Glu Tyr Asn Gly Ala Lys Thr Asn Thr Ala Val Gln Leu															
195	200	205													
Val Gln Ile Val Gln Gln Ala Gly Ala Pro Ile Asp Gly Val Gly Phe															
210	215	220													
Gln Gly His Leu Ile Val Gly Ser Thr Pro Ser Arg Ser Ser Leu Ala															
225	230	235	240												
Thr Ala Leu Lys Arg Phe Thr Ala Leu Gly Leu Glu Val Ala Tyr Thr															
245	250	255													
Glu Leu Asp Ile Arg His Ser Ser Leu Pro Pro Ser Ser Ala Ala Leu															
260	265	270													
Ala Thr Gln Gly Asn Asp Phe Ala Ser Val Val Gly Ser Cys Leu Asp															
275	280	285													
Val Ala Gly Cys Val Gly Ile Thr Ile Trp Gly Phe Thr Asp Lys Tyr															
290	295	300													
Ser Trp Val Pro Asp Thr Phe Pro Gly Ser Gly Ala Ala Leu Leu Tyr															
305	310	315	320												
Asp Ala Asn Tyr Ser Lys Lys Pro Ala Trp Thr Ser Val Ser Ser Val															
325	330	335													
Leu Ala Ala Lys Ala Thr Asn Pro Pro Gly Gly Gly Asn Pro Pro Pro															
340	345	350													
Val Thr Thr Thr Thr Thr Thr Thr Thr Thr Ser Lys Pro Ser Gln Pro															
355	360	365													
Thr Thr Thr Thr Thr Thr Thr Ser Pro Gln Gly Pro Gln Gln Thr His															
370	375	380													
Trp Gly Gln Cys Gly Gly Ile Gly Trp Thr Gly Pro Gln Ser Cys Gln															
385	390	395	400												
Ser Pro Trp Thr Cys Gln Lys Gln Asn Asp Trp Tyr Ser Gln Cys Leu															
405	410	415													

<210> SEQ ID NO 21
 <211> LENGTH: 5092
 <212> TYPE: DNA
 <213> ORGANISM: *Thermoascus aurantiacus*
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (669)..(728)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (729)..(872)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (873)..(1015)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1016)..(1082)

-continued

```

<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1083)..(1127)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1128)..(1183)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1184)..(1236)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1237)..(1300)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1301)..(1717)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (1718)..(1776)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1777)..(2489)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (2490)..(2599)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (2600)..(3469)
<220> FEATURE:
<221> NAME/KEY: Intron
<222> LOCATION: (3470)..(3531)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (3532)..(3759)

<400> SEQUENCE: 21

ggatccgtcc gcgacacag gcagagagac ggcacgggga ctcgacctga tcctcccagg      60
gcggggtggt gtttgtggcg agggagcgat getgatgttc ttccagctcc gttgetacct      120
tcccacggcc atttagccgg cggacggcat gtaacatgtc aaacatgtgg gctcggcagt      180
ggggcggtga gacgcagcac ctgaccggc ggcgcggcgc ttgcagggtc cagggacagc      240
cggccgtggt cgtttgctgg gaaggcgaca cagacgactt ggcgcggccc gccggaaggc      300
gaggaatcat gagtgcgacg gagacatggc aagaccacgg ccttctctggc gaagaagaag      360
atgaataatc gcaggggagc tgtggcatgg accgcacggc cgccaggagc ctgccccgtg      420
aggtttctcg ggtgtttcca ctggttccat cgctgggggc gatccccgagc ccgtgtgccc      480
gtgtaactat tattgacgat caacatgcca tggccagcca gcttctataa taatcatata      540
taacaccccc cgttctcccg ctgccttgct ccgtggtctt cctggtcctg cttgaggttc      600
acgagtctcc ttgcatggtc aactcgctct ctgcttcctc cgctgcttga ctccgtacct      660
cagcaacc atg agg ctt ggg tgg ctg gag ctg gcc gtc gcg gcc gca      710
Met Arg Leu Gly Trp Leu Glu Leu Ala Val Ala Ala Ala Ala
1           5           10

acc gtc gcc agc gcc aag gtgcgtcaga cctcccccg gatcgacctt      758
Thr Val Ala Ser Ala Lys
15           20

taggtgcttc ttcagcaagt ggcgcccggc cgcgacatcc gccgcccgtg cctcaccga      818

cgcagcacc atagcagca ggagagaagg catctctgac gaaagctccc ccag gat      875
Asp

gac ttg gcc tac tcg ccg cct ttc tac ccg tcg cca tgg atg aac gga      923
Asp Leu Ala Tyr Ser Pro Pro Phe Tyr Pro Ser Pro Trp Met Asn Gly
25           30           35

aac gga gag tgg gcg gag gcc tac cgc agg gct gtc gac ttc gtc tcg      971

```


-continued

Asn Gly Glu Trp Ala Glu Ala Tyr Arg Arg Ala Val Asp Phe Val Ser	
40	45 50
cag ctg acc ctc gcg gag aag gtc aac ctg acg acc ggt gtc gg	1015
Gln Leu Thr Leu Ala Glu Lys Val Asn Leu Thr Thr Gly Val Gly	
55	60 65
gtgagtcacat tgacctctac cgagcccccg ttccatgtcc attgagcaat tggctgacgt	1075
cttgaag c tgg atg cag gag aaa tgt gtc ggt gaa acg ggc agc att ccg	1125
Trp Met Gln Glu Lys Cys Val Gly Glu Thr Gly Ser Ile Pro	
70	75 80
ag gtaggctcac ttcccaatgc cgctgcaaag gaggtgtcta aactggaata aatcag	1183
Arg	
a ctg ggg ttc cgt gga ctg tgc ctc caa gac tcg ccc ctt ggt gtc aga	1232
Leu Gly Phe Arg Gly Leu Cys Leu Gln Asp Ser Pro Leu Gly Val Arg	
85	90 95
ttt g gtaggtcttt caacagagaa caagggtcgt cgcgaggagag atgctgatcg	1286
Phe	
100	
atacctactt ttag ct gac tac gtt tct gcc ttc ccc gcc ggt gtc aat	1335
Ala Asp Tyr Val Ser Ala Phe Pro Ala Gly Val Asn	
105	110
gtc gct gca acg tgg gat aag aac ctc gcc tac ctt cgt ggg aag gcg	1383
Val Ala Ala Thr Trp Asp Lys Asn Leu Ala Tyr Leu Arg Gly Lys Ala	
115	120 125
atg ggt gag gaa cac cgt ggt aag ggc gtc gac gtc cag ctg gga cct	1431
Met Gly Glu Glu His Arg Gly Lys Gly Val Asp Val Gln Leu Gly Pro	
130	135 140
gtc gcc ggc cct ctt ggc aga cac ccc gac ggt ggc aga aac tgg gag	1479
Val Ala Gly Pro Leu Gly Arg His Pro Asp Gly Gly Arg Asn Trp Glu	
145	150 155 160
ggt ttc tct cct gac ccc gtc ctg acc ggt gtg ctt atg gcg gag acg	1527
Gly Phe Ser Pro Asp Pro Val Leu Thr Gly Val Leu Met Ala Glu Thr	
165	170 175
atc aag ggt atc cag gat gcc ggt gtg att gct tgc gcc aag cac ttc	1575
Ile Lys Gly Ile Gln Asp Ala Gly Val Ile Ala Cys Ala Lys His Phe	
180	185 190
att ggt aac gag atg gag cac ttc cgg caa gcc ggt gag gct gtt ggc	1623
Ile Gly Asn Glu Met Glu His Phe Arg Gln Ala Gly Glu Ala Val Gly	
195	200 205
tat ggt ttc gat att acc gag agt gtc agc tca aat atc gac gac aag	1671
Tyr Gly Phe Asp Ile Thr Glu Ser Val Ser Ser Asn Ile Asp Asp Lys	
210	215 220
acg ctt cac gag ctg tac ctt tgg ccc ttt gcg gat gct gtt cgc g	1717
Thr Leu His Glu Leu Tyr Leu Trp Pro Phe Ala Asp Ala Val Arg	
225	230 235
gtaagcagtc ccccccctcat aggtgattgt acatgtgtat ttctgactcg ctttcaaag	1776
ct ggc gtt ggt tcg ttc atg tgc tcc tac aac cag gtt aac aac agc	1823
Ala Gly Val Gly Ser Phe Met Cys Ser Tyr Asn Gln Val Asn Asn Ser	
240	245 250 255
tac agc tgc tcg aac agc tac ctc cta aac aag ttg ctc aaa tcg gag	1871
Tyr Ser Cys Ser Asn Ser Tyr Leu Leu Asn Lys Leu Leu Lys Ser Glu	
260	265 270
ctt gat ttt cag ggc ttc gtg atg agt gac tgg gga gcg cac cac agc	1919
Leu Asp Phe Gln Gly Phe Val Met Ser Asp Trp Gly Ala His His Ser	
275	280 285
ggc gtt gga gct gcc ctg gct ggc ctt gac atg tcg atg cca gga gac	1967
Gly Val Gly Ala Ala Leu Ala Gly Leu Asp Met Ser Met Pro Gly Asp	

-continued

290	295	300	
acc gcc ttt ggt acc ggc aaa tcc ttc tgg gga acc aac ctg acc atc			2015
Thr Ala Phe Gly Thr Gly Lys Ser Phe Trp Gly Thr Asn Leu Thr Ile			
305	310	315	
gcc gtt ctc aac ggt act gtt ccg gaa tgg cgt gtg gat gac atg gct			2063
Ala Val Leu Asn Gly Thr Val Pro Glu Trp Arg Val Asp Asp Met Ala			
320	325	330	335
gtt cgc atc atg gcg gcc ttt tac aag gtt ggt cgc gac cgt tac cag			2111
Val Arg Ile Met Ala Ala Phe Tyr Lys Val Gly Arg Asp Arg Tyr Gln			
340	345	350	
gtg ccg gtc aac ttc gac tcg tgg acg aag gat gaa tac ggt tac gag			2159
Val Pro Val Asn Phe Asp Ser Trp Thr Lys Asp Glu Tyr Gly Tyr Glu			
355	360	365	
cac gca ctg gtt ggc cag aac tat gtc aag gtc aat gac aag gtg gat			2207
His Ala Leu Val Gly Gln Asn Tyr Val Lys Val Asn Asp Lys Val Asp			
370	375	380	
gtt cgt gcc gac cat gcg gac atc atc cgt caa att ggg tct gct agt			2255
Val Arg Ala Asp His Ala Asp Ile Ile Arg Gln Ile Gly Ser Ala Ser			
385	390	395	
gtt gtc ctt ctt aag aac gat gga gga ctc cca ttg acc ggc tat gaa			2303
Val Val Leu Leu Lys Asn Asp Gly Gly Leu Pro Leu Thr Gly Tyr Glu			
400	405	410	415
aag ttc acc gga gtt ttt gga gag gat gcc gga tcg aac cgt tgg ggc			2351
Lys Phe Thr Gly Val Phe Gly Glu Asp Ala Gly Ser Asn Arg Trp Gly			
420	425	430	
gct gac ggc tgc tct gat cgt ggt tgc gac aac ggc acg ttg gca atg			2399
Ala Asp Gly Cys Ser Asp Arg Gly Cys Asp Asn Gly Thr Leu Ala Met			
435	440	445	
ggt tgg ggc agt ggc act gct gac ttc ccc tac ctt gtc act ccc gag			2447
Gly Trp Gly Ser Gly Thr Ala Asp Phe Pro Tyr Leu Val Thr Pro Glu			
450	455	460	
cag gca atc cag aat gaa atc ctt tcc aag ggg aag ggg tta			2489
Gln Ala Ile Gln Asn Glu Ile Leu Ser Lys Gly Lys Gly Leu			
465	470	475	
gtgagtgctg tcaccgacaa tgggtgccctt gaccagatgg aacaggttgc gtctcaggcc			2549
aggtattcct tcctccgtat cccatagcaat cgaatctcca ctgactttag gac agc			2605
Asp Ser			
gtt tct atc gtt ttc gtc aac gcc gac tct ggt gaa ggc tac atc aac			2653
Val Ser Ile Val Phe Val Asn Ala Asp Ser Gly Glu Gly Tyr Ile Asn			
480	485	490	495
gtt gat ggc aac gaa ggt gat cgg aag aac ctc acc ctc tgg aaa gga			2701
Val Asp Gly Asn Glu Gly Asp Arg Lys Asn Leu Thr Leu Trp Lys Gly			
500	505	510	
ggc gag gag gtg atc aag act gtt gca gcc aac tgc aac aac acc att			2749
Gly Glu Glu Val Ile Lys Thr Val Ala Ala Asn Cys Asn Asn Thr Ile			
515	520	525	
gtt gtg atg cac act gtg gga cct gtc ttg atc gat gag tgg tat gac			2797
Val Val Met His Thr Val Gly Pro Val Leu Ile Asp Glu Trp Tyr Asp			
530	535	540	
aac ccc aac gtc acc gcc atc gtc tgg gcc ggt ctt cca ggc cag gag			2845
Asn Pro Asn Val Thr Ala Ile Val Trp Ala Gly Leu Pro Gly Gln Glu			
545	550	555	
agc ggc aac agt ctc gtc gat gtg ctc tac ggc cgt gtc agc ccc gga			2893
Ser Gly Asn Ser Leu Val Asp Val Leu Tyr Gly Arg Val Ser Pro Gly			
560	565	570	575
gga aag acg ccg ttt acg tgg gga aag act cgc gag tcg tac ggc gct			2941

-continued

```

ggttcagaaa aagaaaatgt agaggtttca aacacgctag ttgaccctga taggaattga 4069
gcatgaatgc ctacacattc caagtcgtgt tagcgagtcg atagccgatg aacctattcc 4129
gtaggttgag gttcaccta caaataagcc aggatttaag taaatacctg ctcgtgaaat 4189
ctacaacgca tcagatcaga ggaaaattca aatggcagaa gtgcgagcac ctcggtgaga 4249
agagatcgag ctgtcgaagt cggctggaac acaggtaaag agaagtaata caattcattg 4309
atTTTTacat cgtttaacat gtagaaggta tctaaaatag taagtccaga tatgggcat 4369
ggagatcgcc tcggcgatct tcgggagat ctctgggagac gcacatgacc gcgcttaacc 4429
ctgtcggttg gacccgagtc cgaccgacgt catcagcgc ggcgaggta ggctgcgcgc 4489
aacgtcaatg ccagggggtg ctgggacagt tgcatatcaa tcgatcagtc aattaaagca 4549
tctgctttcc acgttctttt tttatcacct ttcacttccc ctgtcccact tgccttggga 4609
ttgttgagcc caaagaagaa ggagaagaaa atgggctcga cccccggaa cgggtggtcg 4669
acgagcacat catcagcagc gtcttattat caacattccc aaccaccggc cctcgttctc 4729
ctcgtctacc cgctcactct cctcctcggc tccctgtaca gagccatttc cccaccgcg 4789
cgggtgaggc acgatgctgc agaccctgct ctggccccga ccatagcgtc cgacatcaac 4849
ctgtcccagt catcccggta tcccattcc catagcaaca gcaacagccc ggtcaattac 4909
ttcggccgca aggacaacat ctttaacgtc tacttctgca agatcggctg gttctggacg 4969
accctcgctt tctcactgtt actcctcacc cagcctgctt acacaaacgc cggteccctg 5029
cgcgcccagc gcaccctcca agcctgtcc cgctacgcca tcgtcacctt actacctgga 5089
tcc 5092

```

<210> SEQ ID NO 22

<211> LENGTH: 843

<212> TYPE: PRT

<213> ORGANISM: *Thermoascus aurantiacus*

<400> SEQUENCE: 22

```

Met Arg Leu Gly Trp Leu Glu Leu Ala Val Ala Ala Ala Ala Thr Val
1           5           10           15
Ala Ser Ala Lys Asp Asp Leu Ala Tyr Ser Pro Pro Phe Tyr Pro Ser
20           25           30
Pro Trp Met Asn Gly Asn Gly Glu Trp Ala Glu Ala Tyr Arg Arg Ala
35           40           45
Val Asp Phe Val Ser Gln Leu Thr Leu Ala Glu Lys Val Asn Leu Thr
50           55           60
Thr Gly Val Gly Trp Met Gln Glu Lys Cys Val Gly Glu Thr Gly Ser
65           70           75           80
Ile Pro Arg Leu Gly Phe Arg Gly Leu Cys Leu Gln Asp Ser Pro Leu
85           90           95
Gly Val Arg Phe Ala Asp Tyr Val Ser Ala Phe Pro Ala Gly Val Asn
100          105          110
Val Ala Ala Thr Trp Asp Lys Asn Leu Ala Tyr Leu Arg Gly Lys Ala
115          120          125
Met Gly Glu Glu His Arg Gly Lys Gly Val Asp Val Gln Leu Gly Pro
130          135          140
Val Ala Gly Pro Leu Gly Arg His Pro Asp Gly Gly Arg Asn Trp Glu
145          150          155          160

```


-continued

Gly	Phe	Ser	Pro	Asp	Pro	Val	Leu	Thr	Gly	Val	Leu	Met	Ala	Glu	Thr	165	170	175	
Ile	Lys	Gly	Ile	Gln	Asp	Ala	Gly	Val	Ile	Ala	Cys	Ala	Lys	His	Phe	180	185	190	
Ile	Gly	Asn	Glu	Met	Glu	His	Phe	Arg	Gln	Ala	Gly	Glu	Ala	Val	Gly	195	200	205	
Tyr	Gly	Phe	Asp	Ile	Thr	Glu	Ser	Val	Ser	Ser	Asn	Ile	Asp	Asp	Lys	210	215	220	
Thr	Leu	His	Glu	Leu	Tyr	Leu	Trp	Pro	Phe	Ala	Asp	Ala	Val	Arg	Ala	225	230	235	240
Gly	Val	Gly	Ser	Phe	Met	Cys	Ser	Tyr	Asn	Gln	Val	Asn	Asn	Ser	Tyr	245	250	255	
Ser	Cys	Ser	Asn	Ser	Tyr	Leu	Leu	Asn	Lys	Leu	Leu	Lys	Ser	Glu	Leu	260	265	270	
Asp	Phe	Gln	Gly	Phe	Val	Met	Ser	Asp	Trp	Gly	Ala	His	His	Ser	Gly	275	280	285	
Val	Gly	Ala	Ala	Leu	Ala	Gly	Leu	Asp	Met	Ser	Met	Pro	Gly	Asp	Thr	290	295	300	
Ala	Phe	Gly	Thr	Gly	Lys	Ser	Phe	Trp	Gly	Thr	Asn	Leu	Thr	Ile	Ala	305	310	315	320
Val	Leu	Asn	Gly	Thr	Val	Pro	Glu	Trp	Arg	Val	Asp	Asp	Met	Ala	Val	325	330	335	
Arg	Ile	Met	Ala	Ala	Phe	Tyr	Lys	Val	Gly	Arg	Asp	Arg	Tyr	Gln	Val	340	345	350	
Pro	Val	Asn	Phe	Asp	Ser	Trp	Thr	Lys	Asp	Glu	Tyr	Gly	Tyr	Glu	His	355	360	365	
Ala	Leu	Val	Gly	Gln	Asn	Tyr	Val	Lys	Val	Asn	Asp	Lys	Val	Asp	Val	370	375	380	
Arg	Ala	Asp	His	Ala	Asp	Ile	Ile	Arg	Gln	Ile	Gly	Ser	Ala	Ser	Val	385	390	395	400
Val	Leu	Leu	Lys	Asn	Asp	Gly	Gly	Leu	Pro	Leu	Thr	Gly	Tyr	Glu	Lys	405	410	415	
Phe	Thr	Gly	Val	Phe	Gly	Glu	Asp	Ala	Gly	Ser	Asn	Arg	Trp	Gly	Ala	420	425	430	
Asp	Gly	Cys	Ser	Asp	Arg	Gly	Cys	Asp	Asn	Gly	Thr	Leu	Ala	Met	Gly	435	440	445	
Trp	Gly	Ser	Gly	Thr	Ala	Asp	Phe	Pro	Tyr	Leu	Val	Thr	Pro	Glu	Gln	450	455	460	
Ala	Ile	Gln	Asn	Glu	Ile	Leu	Ser	Lys	Gly	Lys	Gly	Leu	Asp	Ser	Val	465	470	475	480
Ser	Ile	Val	Phe	Val	Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile	Asn	Val	485	490	495	
Asp	Gly	Asn	Glu	Gly	Asp	Arg	Lys	Asn	Leu	Thr	Leu	Trp	Lys	Gly	Gly	500	505	510	
Glu	Glu	Val	Ile	Lys	Thr	Val	Ala	Ala	Asn	Cys	Asn	Asn	Thr	Ile	Val	515	520	525	
Val	Met	His	Thr	Val	Gly	Pro	Val	Leu	Ile	Asp	Glu	Trp	Tyr	Asp	Asn	530	535	540	
Pro	Asn	Val	Thr	Ala	Ile	Val	Trp	Ala	Gly	Leu	Pro	Gly	Gln	Glu	Ser	545	550	555	560

-continued

Gly Asn Ser Leu Val Asp Val Leu Tyr Gly Arg Val Ser Pro Gly Gly
 565 570 575
 Lys Thr Pro Phe Thr Trp Gly Lys Thr Arg Glu Ser Tyr Gly Ala Pro
 580 585 590
 Leu Leu Thr Lys Pro Asn Asn Gly Lys Gly Ala Pro Gln Asp Asp Phe
 595 600 605
 Thr Glu Gly Val Phe Ile Asp Tyr Arg Arg Phe Asp Lys Tyr Asn Glu
 610 615 620
 Thr Pro Ile Tyr Glu Phe Gly Phe Gly Leu Ser Tyr Thr Thr Phe Glu
 625 630 635 640
 Tyr Ser Asn Ile Tyr Val Gln Pro Leu Asn Ala Arg Pro Tyr Thr Pro
 645 650 655
 Ala Ser Gly Ser Thr Lys Ala Ala Pro Thr Phe Gly Asn Ile Ser Thr
 660 665 670
 Asp Tyr Ala Asp Tyr Leu Tyr Pro Glu Asp Ile His Lys Val Pro Leu
 675 680 685
 Tyr Ile Tyr Pro Trp Leu Asn Thr Thr Asp Pro Glu Glu Val Leu Arg
 690 695 700
 Arg Ser Arg Leu Thr Glu Met Lys Ala Glu Asp Tyr Ile Pro Ser Gly
 705 710 715 720
 Ala Thr Asp Gly Ser Pro Gln Pro Ile Leu Pro Ala Gly Gly Ala Pro
 725 730 735
 Gly Gly Asn Pro Gly Leu Tyr Asp Glu Met Tyr Arg Val Ser Ala Ile
 740 745 750
 Ile Thr Asn Thr Gly Asn Val Val Gly Asp Glu Val Pro Gln Leu Tyr
 755 760 765
 Val Ser Leu Gly Gly Pro Asp Asp Pro Lys Val Val Leu Arg Asn Phe
 770 775 780
 Asp Arg Ile Thr Leu His Pro Gly Gln Gln Thr Met Trp Thr Thr Thr
 785 790 795 800
 Leu Thr Arg Arg Asp Ile Ser Asn Trp Asp Pro Ala Ser Gln Asn Trp
 805 810 815
 Val Val Thr Lys Tyr Pro Lys Thr Val Tyr Ile Gly Ser Ser Ser Arg
 820 825 830
 Lys Leu His Leu Gln Ala Pro Leu Pro Pro Tyr
 835 840

<210> SEQ ID NO 23
 <211> LENGTH: 3510
 <212> TYPE: DNA
 <213> ORGANISM: Acremonium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (391)..(447)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (448)..(539)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (540)..(685)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (686)..(759)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (760)..(1148)
 <220> FEATURE:
 <221> NAME/KEY: Intron

-continued

```

<222> LOCATION: (1149)..(1217)
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (1218)..(3208)

<400> SEQUENCE: 23

gcaggtagct acgacattcg acggtccacg cccagtggcg tctgctcggc cgtctgggaa      60
ccatgcacgc ccgcctctta ggtcgagcga ggtataacat actatctgca cggctaccta      120
tatattacgt cgatgtcacc cgcaggatgc gagcaccatt acttctgtgc tcaccgccc      180
ttccgctccg catctcgtga acctaaaccc acgcgggcac actgcttctt gtgagagcct      240
ctaccogttc cacaagagcc atagctagag agagaagggc agccaaggga ccggtcaagc      300
ggcgctcttc atcgaccaa tctcgacaac ccggcagacg tcaccaccgg ctcccgcgcg      360

acgacgtcac acgggactga ctacgaagac atg agg cag gcc ctt gtt tcg ctg      414
Met Arg Gln Ala Leu Val Ser Leu
1          5

gcc ttg ctg gcc agc agc cct gtt tcg gcg gcg gtgaccgcca gggacgcca      467
Ala Leu Leu Ala Ser Ser Pro Val Ser Ala Ala
10         15

ggtatggtec caactgctct tcctccctgt ttctcctct accggtgctg acaacgacaa      527

tagctgcacc ag cga gaa ctc gcc act tcc gac cct ttc tat cct tcg cca      578
Arg Glu Leu Ala Thr Ser Asp Pro Phe Tyr Pro Ser Pro
20         25         30

tgg atg aac cct gaa gcc aat ggc tgg gag gac gcc tac gcc aag gcc      626
Trp Met Asn Pro Glu Ala Asn Gly Trp Glu Asp Ala Tyr Ala Lys Ala
35         40         45

aag gcg ttc gtt tcc cag ctg acg ctc ttg gaa aag gtc aac ctg acg      674
Lys Ala Phe Val Ser Gln Leu Thr Leu Leu Glu Lys Val Asn Leu Thr
50         55         60

act ggc atc gg  gtgagtcttg ttctctcctg tagaaccgcc taccagaaga      725
Thr Gly Ile Gly
65

cattcaggaa gtgctaata tgggcgggtg acag c tgg caa gga gga caa tgc      778
Trp Gln Gly Gly Gln Cys
70

gtg ggc aac gtc ggt tcc gtc ccg cgt ctc ggc ctt cgc agc ctg tgc      826
Val Gly Asn Val Gly Ser Val Pro Arg Leu Gly Leu Arg Ser Leu Cys
75         80         85         90

atg cag gac tcc ccc gtg ggt atc cgc ttt ggg gac tac gtc tcc gtc      874
Met Gln Asp Ser Pro Val Gly Ile Arg Phe Gly Asp Tyr Val Ser Val
95         100        105

ttc ccc tct ggt cag acc acg gct gcc acc ttc gac aag ggt ctg atg      922
Phe Pro Ser Gly Gln Thr Thr Ala Ala Thr Phe Asp Lys Gly Leu Met
110        115        120

aac cgt cgc ggc aat gcc atg ggc cag gag cac aaa gga aag ggt gtc      970
Asn Arg Arg Gly Asn Ala Met Gly Gln Glu His Lys Gly Lys Gly Val
125        130        135

aac gtc ctg ctc ggc ccg gtc gct ggc ccc att ggc cgt acg ccc gag      1018
Asn Val Leu Leu Gly Pro Val Ala Gly Pro Ile Gly Arg Thr Pro Glu
140        145        150

ggg gga cga aac tgg gag ggc ttc tcc ccc gac ccc gtc cta acg ggt      1066
Gly Gly Arg Asn Trp Glu Gly Phe Ser Pro Asp Pro Val Leu Thr Gly
155        160        165        170

att gcc ttg gcc gaa acg atc aag gga atc cag gat gct ggt gtc att      1114
Ile Ala Leu Ala Glu Thr Ile Lys Gly Ile Gln Asp Ala Gly Val Ile
175        180        185

```

-continued

gct tgc gcc aag cat ttc atc gcg aac gaa cag g	gtgctgatg	1158
Ala Cys Ala Lys His Phe Ile Ala Asn Glu Gln		
190	195	
gaacgcggga cgtgctctga tgcaaaccca cgagcactga	ccacgctttc ctogaacag	1217
aa cac ttc cgc cag tcc ggc gag gcc cag ggc	tac ggc ttt gac atc	1264
Glu His Phe Arg Gln Ser Gly Glu Ala Gln Gly	Tyr Gly Phe Asp Ile	
200	205 210	
tcc gag tcg ctg tcg tcc aac atc gac gac aag	acc atg cac gag ctg	1312
Ser Glu Ser Leu Ser Ser Asn Ile Asp Asp Lys	Thr Met His Glu Leu	
215	220 225	
tat ctg tgg ccc ttc gcc gac ggc gtg cgt gcc	ggc gtc ggc gcc atc	1360
Tyr Leu Trp Pro Phe Ala Asp Gly Val Arg Ala	Gly Val Gly Ala Ile	
230	235 240 245	
atg tgc tcg tac aac cag atc aac aac tcg tac	ggg tgc cag aac tcc	1408
Met Cys Ser Tyr Asn Gln Ile Asn Asn Ser Tyr	Gly Cys Gln Asn Ser	
250	255 260	
aag acc ctg aac aac ctg ctc aag aac gag ctc	ggc ttc cag ggc ttc	1456
Lys Thr Leu Asn Asn Leu Leu Lys Asn Glu Leu	Gly Phe Gln Gly Phe	
265	270 275	
gtc atg agc gac tgg cag gcc cag cac acc ggc	gcg gcc agc gcc gtc	1504
Val Met Ser Asp Trp Gln Ala Gln His Thr Gly	Ala Ala Ser Ala Val	
280	285 290	
gcc ggc ctg gac atg acc atg ccc ggc gac acc	agc ttc aac acc ggc	1552
Ala Gly Leu Asp Met Thr Met Pro Gly Asp Thr	Ser Phe Asn Thr Gly	
295	300 305	
ctc agc tac tgg ggc acg aac ctc acc ctc gcc	gtc ctg aac ggc acc	1600
Leu Ser Tyr Trp Gly Thr Asn Leu Thr Leu Ala	Val Leu Asn Gly Thr	
310	315 320 325	
gtc ccc gag tac cgc atc gac gac atg gtc atg	cgc atc atg gcc gcc	1648
Val Pro Glu Tyr Arg Ile Asp Asp Met Val Met	Arg Ile Met Ala Ala	
330	335 340	
ttc ttc aag acc ggc cag acc ctg gac ctg ccg	ccc atc aac ttc gac	1696
Phe Phe Lys Thr Gly Gln Thr Leu Asp Leu Pro	Pro Ile Asn Phe Asp	
345	350 355	
tcg tgg acc acc gac acc ttc ggc ccg ctc cac	ttc gcc gtc aac gag	1744
Ser Trp Thr Thr Asp Thr Phe Gly Pro Leu His	Phe Ala Val Asn Glu	
360	365 370	
gac cgc cag cag atc aac tgg cac gtc aac gtc	cag gac aac cat ggc	1792
Asp Arg Gln Gln Ile Asn Trp His Val Asn Val	Gln Asp Asn His Gly	
375	380 385	
agc ctc atc cgc gag atc gcg gcc aag gga acc	gtc ctg ctg aag aac	1840
Ser Leu Ile Arg Glu Ile Ala Ala Lys Gly Thr	Val Leu Leu Lys Asn	
390	395 400 405	
acc ggg tcc ctc ccg ctc aac aag ccc aag ttc	ctc gtc gtg gtc ggc	1888
Thr Gly Ser Leu Pro Leu Asn Lys Pro Lys Phe	Leu Val Val Val Gly	
410	415 420	
gac gac gcg ggc ccc aac ccg gcg gga ccc aac	gcc tgc ccc gac cgc	1936
Asp Asp Ala Gly Pro Asn Pro Ala Gly Pro Asn	Ala Cys Pro Asp Arg	
425	430 435	
gga tgc gac gtc ggc acc ctc ggc atg gcc tgg	ggc tcc ggc tcg gcc	1984
Gly Cys Asp Val Gly Thr Leu Gly Met Ala Trp	Gly Ser Gly Ser Ala	
440	445 450	
aac ttc ccc tac ctg atc acc ccg gac gcc gcg	ctg cag gcg cag gcg	2032
Asn Phe Pro Tyr Leu Ile Thr Pro Asp Ala Ala	Leu Gln Ala Gln Ala	
455	460 465	
atc aag gac ggc acc cgc tac gag agc gtg ctg	tcc aac tac cag ctc	2080

-continued

Ile	Lys	Asp	Gly	Thr	Arg	Tyr	Glu	Ser	Val	Leu	Ser	Asn	Tyr	Gln	Leu	
470					475					480					485	
gac	cag	acc	aag	gcg	ctg	gtc	acc	cag	gcc	aac	gcc	acg	gcc	atc	gtc	2128
Asp	Gln	Thr	Lys	Ala	Leu	Val	Thr	Gln	Ala	Asn	Ala	Thr	Ala	Ile	Val	
490					495					500						
ttc	gtc	aac	gcc	gac	tcg	ggc	gag	ggc	tac	atc	aac	gtc	gac	ggc	aac	2176
Phe	Val	Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile	Asn	Val	Asp	Gly	Asn	
505					510					515						
gag	ggc	gac	cgc	aag	aac	ctc	acg	ctc	tgg	cac	gac	ggc	gac	gcc	ctg	2224
Glu	Gly	Asp	Arg	Lys	Asn	Leu	Thr	Leu	Trp	His	Asp	Gly	Asp	Ala	Leu	
520					525					530						
atc	aag	agc	gtg	gcc	ggc	tgg	aac	ccg	aac	acc	atc	gtc	gtc	atc	cac	2272
Ile	Lys	Ser	Val	Ala	Gly	Trp	Asn	Pro	Asn	Thr	Ile	Val	Val	Ile	His	
535					540					545						
tcg	acc	ggc	ccc	gtc	ctc	gtg	acc	gac	tgg	tac	gac	cac	ccc	aac	atc	2320
Ser	Thr	Gly	Pro	Val	Leu	Val	Thr	Asp	Trp	Tyr	Asp	His	Pro	Asn	Ile	
550					555					560					565	
acc	gcc	atc	ctg	tgg	gcc	ggc	gtg	ccc	ggg	cag	gag	tcc	ggc	aac	gcc	2368
Thr	Ala	Ile	Leu	Trp	Ala	Gly	Val	Pro	Gly	Gln	Glu	Ser	Gly	Asn	Ala	
570					575					580						
atc	acc	gac	gtc	ctc	tac	gga	aaa	gtc	aac	ccg	tcg	ggc	cgc	agc	ccc	2416
Ile	Thr	Asp	Val	Leu	Tyr	Gly	Lys	Val	Asn	Pro	Ser	Gly	Arg	Ser	Pro	
585					590					595						
ttc	acc	tgg	ggt	ccg	acc	cgc	gag	agc	tac	ggc	acc	gac	gtc	ctc	tac	2464
Phe	Thr	Trp	Gly	Pro	Thr	Arg	Glu	Ser	Tyr	Gly	Thr	Asp	Val	Leu	Tyr	
600					605					610						
act	ccc	aac	aac	ggc	aag	ggc	gcg	ccg	cag	cag	gcc	ttc	tcc	gag	ggc	2512
Thr	Pro	Asn	Asn	Gly	Lys	Gly	Ala	Pro	Gln	Gln	Ala	Phe	Ser	Glu	Gly	
615					620					625						
gtc	ttc	atc	gac	tac	cgc	cac	ttc	gac	cgc	acc	aac	gcg	tcc	gtc	atc	2560
Val	Phe	Ile	Asp	Tyr	Arg	His	Phe	Asp	Arg	Thr	Asn	Ala	Ser	Val	Ile	
630					635					640					645	
tac	gag	ttc	ggc	cac	ggc	ctc	agc	tac	acg	acg	ttc	cag	tac	agc	aac	2608
Tyr	Glu	Phe	Gly	His	Gly	Leu	Ser	Tyr	Thr	Thr	Phe	Gln	Tyr	Ser	Asn	
650					655					660						
atc	cag	gtg	gtc	aag	tcc	aac	gcc	ggc	gcg	tac	aag	ccc	acg	acg	ggc	2656
Ile	Gln	Val	Val	Lys	Ser	Asn	Ala	Gly	Ala	Tyr	Lys	Pro	Thr	Thr	Gly	
665					670					675						
acg	acc	atc	ccc	gcg	ccc	acg	ttt	ggc	agc	ttc	tcc	aag	gac	ctc	aag	2704
Thr	Thr	Ile	Pro	Ala	Pro	Thr	Phe	Gly	Ser	Phe	Ser	Lys	Asp	Leu	Lys	
680					685					690						
gac	tac	ctc	ttc	ccg	tcg	gac	cag	ttc	cgc	tac	atc	acc	cag	tac	atc	2752
Asp	Tyr	Leu	Phe	Pro	Ser	Asp	Gln	Phe	Arg	Tyr	Ile	Thr	Gln	Tyr	Ile	
695					700					705						
tac	ccg	tac	ctc	aac	tcc	acc	gac	ccg	gcc	aag	gcg	tcg	ctc	gac	ccg	2800
Tyr	Pro	Tyr	Leu	Asn	Ser	Thr	Asp	Pro	Ala	Lys	Ala	Ser	Leu	Asp	Pro	
710					715					720					725	
cac	tac	ggc	aag	acg	gcg	gcc	gag	ttt	ctg	ccg	ccg	cac	gcg	ctg	gac	2848
His	Tyr	Gly	Lys	Thr	Ala	Ala	Glu	Phe	Leu	Pro	Pro	His	Ala	Leu	Asp	
730					735					740						
agc	aac	ccg	cag	ccg	ctg	ctg	cgg	tcg	tcg	ggc	aag	aac	gag	ccc	ggc	2896
Ser	Asn	Pro	Gln	Pro	Leu	Leu	Arg	Ser	Ser	Gly	Lys	Asn	Glu	Pro	Gly	
745					750					755						
ggc	aac	cgc	cag	ctg	tac	gac	atc	ctg	tac	acg	gtg	acg	gcg	gac	atc	2944
Gly	Asn	Arg	Gln	Leu	Tyr	Asp	Ile	Leu	Tyr	Thr	Val	Thr	Ala	Asp	Ile	
760					765					770						
acc	aac	acg	ggc	agc	atc	gtg	ggt	gcg	gag	gtg	ccg	cag	ctg	tac	gtg	2992

-continued

Thr Asn Thr Gly Ser Ile Val Gly Ala Glu Val Pro Gln Leu Tyr Val	
775	780 785
tcg ctg ggc ggg ccc gac gac ccc aaa gtg gtc ctg cgc ggg ttc gac	3040
Ser Leu Gly Gly Pro Asp Asp Pro Lys Val Val Leu Arg Gly Phe Asp	
790	795 800 805
cgc atc cgc atc gac ccg ggc aag acg gcg cag ttc cgc gtc acc ctg	3088
Arg Ile Arg Ile Asp Pro Gly Lys Thr Ala Gln Phe Arg Val Thr Leu	
810	815 820
acc cgc cgg gat ctc agc aac tgg gac ccg gcg atc cag gac tgg gtc	3136
Thr Arg Arg Asp Leu Ser Asn Trp Asp Pro Ala Ile Gln Asp Trp Val	
825	830 835
atc agc aag tac ccc aag aag gtg tac atc ggc cgg agc agc agg aag	3184
Ile Ser Lys Tyr Pro Lys Lys Val Tyr Ile Gly Arg Ser Ser Arg Lys	
840	845 850
ctg gaa ctc tcc gcc gac ctc gcg tgatccggcg acggccaagt acgtatgtgg	3238
Leu Glu Leu Ser Ala Asp Leu Ala	
855	860
actgccatcc gaacacctat actttttggc taggtagggg gagcagcaag gcctgagcat	3298
atactctctc cattgcacat ttctaatagta aatatatata tcattaattg ggagacccaa	3358
actcgaattt atgcatgcgt acaaagtgtg ttgaacaagt ttcggtccag cagatagtaa	3418
ccgtcttagt tcgtccatcc ctctctcgaa tgcgctgtat acacatgcgt atatagacgt	3478
tgtataggtg ccattgctag caatgcaagc tt	3510

<210> SEQ ID NO 24
 <211> LENGTH: 861
 <212> TYPE: PRT
 <213> ORGANISM: Acremonium thermophilum

<400> SEQUENCE: 24

Met Arg Gln Ala Leu Val Ser Leu Ala Leu Leu Ala Ser Ser Pro Val	
1	5 10 15
Ser Ala Ala Arg Glu Leu Ala Thr Ser Asp Pro Phe Tyr Pro Ser Pro	
20	25 30
Trp Met Asn Pro Glu Ala Asn Gly Trp Glu Asp Ala Tyr Ala Lys Ala	
35	40 45
Lys Ala Phe Val Ser Gln Leu Thr Leu Leu Glu Lys Val Asn Leu Thr	
50	55 60
Thr Gly Ile Gly Trp Gln Gly Gly Gln Cys Val Gly Asn Val Gly Ser	
65	70 75 80
Val Pro Arg Leu Gly Leu Arg Ser Leu Cys Met Gln Asp Ser Pro Val	
85	90 95
Gly Ile Arg Phe Gly Asp Tyr Val Ser Val Phe Pro Ser Gly Gln Thr	
100	105 110
Thr Ala Ala Thr Phe Asp Lys Gly Leu Met Asn Arg Arg Gly Asn Ala	
115	120 125
Met Gly Gln Glu His Lys Gly Lys Gly Val Asn Val Leu Leu Gly Pro	
130	135 140
Val Ala Gly Pro Ile Gly Arg Thr Pro Glu Gly Gly Arg Asn Trp Glu	
145	150 155 160
Gly Phe Ser Pro Asp Pro Val Leu Thr Gly Ile Ala Leu Ala Glu Thr	
165	170 175
Ile Lys Gly Ile Gln Asp Ala Gly Val Ile Ala Cys Ala Lys His Phe	
180	185 190

-continued

Ile	Ala	Asn	Glu	Gln	Glu	His	Phe	Arg	Gln	Ser	Gly	Glu	Ala	Gln	Gly	195	200	205	
Tyr	Gly	Phe	Asp	Ile	Ser	Glu	Ser	Leu	Ser	Ser	Asn	Ile	Asp	Asp	Lys	210	215	220	
Thr	Met	His	Glu	Leu	Tyr	Leu	Trp	Pro	Phe	Ala	Asp	Gly	Val	Arg	Ala	225	230	235	240
Gly	Val	Gly	Ala	Ile	Met	Cys	Ser	Tyr	Asn	Gln	Ile	Asn	Asn	Ser	Tyr	245	250	255	
Gly	Cys	Gln	Asn	Ser	Lys	Thr	Leu	Asn	Asn	Leu	Leu	Lys	Asn	Glu	Leu	260	265	270	
Gly	Phe	Gln	Gly	Phe	Val	Met	Ser	Asp	Trp	Gln	Ala	Gln	His	Thr	Gly	275	280	285	
Ala	Ala	Ser	Ala	Val	Ala	Gly	Leu	Asp	Met	Thr	Met	Pro	Gly	Asp	Thr	290	295	300	
Ser	Phe	Asn	Thr	Gly	Leu	Ser	Tyr	Trp	Gly	Thr	Asn	Leu	Thr	Leu	Ala	305	310	315	320
Val	Leu	Asn	Gly	Thr	Val	Pro	Glu	Tyr	Arg	Ile	Asp	Asp	Met	Val	Met	325	330	335	
Arg	Ile	Met	Ala	Ala	Phe	Phe	Lys	Thr	Gly	Gln	Thr	Leu	Asp	Leu	Pro	340	345	350	
Pro	Ile	Asn	Phe	Asp	Ser	Trp	Thr	Thr	Asp	Thr	Phe	Gly	Pro	Leu	His	355	360	365	
Phe	Ala	Val	Asn	Glu	Asp	Arg	Gln	Gln	Ile	Asn	Trp	His	Val	Asn	Val	370	375	380	
Gln	Asp	Asn	His	Gly	Ser	Leu	Ile	Arg	Glu	Ile	Ala	Ala	Lys	Gly	Thr	385	390	395	400
Val	Leu	Leu	Lys	Asn	Thr	Gly	Ser	Leu	Pro	Leu	Asn	Lys	Pro	Lys	Phe	405	410	415	
Leu	Val	Val	Val	Gly	Asp	Asp	Ala	Gly	Pro	Asn	Pro	Ala	Gly	Pro	Asn	420	425	430	
Ala	Cys	Pro	Asp	Arg	Gly	Cys	Asp	Val	Gly	Thr	Leu	Gly	Met	Ala	Trp	435	440	445	
Gly	Ser	Gly	Ser	Ala	Asn	Phe	Pro	Tyr	Leu	Ile	Thr	Pro	Asp	Ala	Ala	450	455	460	
Leu	Gln	Ala	Gln	Ala	Ile	Lys	Asp	Gly	Thr	Arg	Tyr	Glu	Ser	Val	Leu	465	470	475	480
Ser	Asn	Tyr	Gln	Leu	Asp	Gln	Thr	Lys	Ala	Leu	Val	Thr	Gln	Ala	Asn	485	490	495	
Ala	Thr	Ala	Ile	Val	Phe	Val	Asn	Ala	Asp	Ser	Gly	Glu	Gly	Tyr	Ile	500	505	510	
Asn	Val	Asp	Gly	Asn	Glu	Gly	Asp	Arg	Lys	Asn	Leu	Thr	Leu	Trp	His	515	520	525	
Asp	Gly	Asp	Ala	Leu	Ile	Lys	Ser	Val	Ala	Gly	Trp	Asn	Pro	Asn	Thr	530	535	540	
Ile	Val	Val	Ile	His	Ser	Thr	Gly	Pro	Val	Leu	Val	Thr	Asp	Trp	Tyr	545	550	555	560
Asp	His	Pro	Asn	Ile	Thr	Ala	Ile	Leu	Trp	Ala	Gly	Val	Pro	Gly	Gln	565	570	575	
Glu	Ser	Gly	Asn	Ala	Ile	Thr	Asp	Val	Leu	Tyr	Gly	Lys	Val	Asn	Pro	580	585	590	

-continued

Ser Gly Arg Ser Pro Phe Thr Trp Gly Pro Thr Arg Glu Ser Tyr Gly
 595 600 605
 Thr Asp Val Leu Tyr Thr Pro Asn Asn Gly Lys Gly Ala Pro Gln Gln
 610 615 620
 Ala Phe Ser Glu Gly Val Phe Ile Asp Tyr Arg His Phe Asp Arg Thr
 625 630 635 640
 Asn Ala Ser Val Ile Tyr Glu Phe Gly His Gly Leu Ser Tyr Thr Thr
 645 650 655
 Phe Gln Tyr Ser Asn Ile Gln Val Val Lys Ser Asn Ala Gly Ala Tyr
 660 665 670
 Lys Pro Thr Thr Gly Thr Thr Ile Pro Ala Pro Thr Phe Gly Ser Phe
 675 680 685
 Ser Lys Asp Leu Lys Asp Tyr Leu Phe Pro Ser Asp Gln Phe Arg Tyr
 690 695 700
 Ile Thr Gln Tyr Ile Tyr Pro Tyr Leu Asn Ser Thr Asp Pro Ala Lys
 705 710 715 720
 Ala Ser Leu Asp Pro His Tyr Gly Lys Thr Ala Ala Glu Phe Leu Pro
 725 730 735
 Pro His Ala Leu Asp Ser Asn Pro Gln Pro Leu Leu Arg Ser Ser Gly
 740 745 750
 Lys Asn Glu Pro Gly Gly Asn Arg Gln Leu Tyr Asp Ile Leu Tyr Thr
 755 760 765
 Val Thr Ala Asp Ile Thr Asn Thr Gly Ser Ile Val Gly Ala Glu Val
 770 775 780
 Pro Gln Leu Tyr Val Ser Leu Gly Gly Pro Asp Asp Pro Lys Val Val
 785 790 795 800
 Leu Arg Gly Phe Asp Arg Ile Arg Ile Asp Pro Gly Lys Thr Ala Gln
 805 810 815
 Phe Arg Val Thr Leu Thr Arg Arg Asp Leu Ser Asn Trp Asp Pro Ala
 820 825 830
 Ile Gln Asp Trp Val Ile Ser Lys Tyr Pro Lys Lys Val Tyr Ile Gly
 835 840 845
 Arg Ser Ser Arg Lys Leu Glu Leu Ser Ala Asp Leu Ala
 850 855 860

<210> SEQ ID NO 25
 <211> LENGTH: 3392
 <212> TYPE: DNA
 <213> ORGANISM: Chaetomium thermophilum
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (608)..(2405)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (2406)..(2457)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (2458)..(2861)

<400> SEQUENCE: 25

tgcgggggttg ctgcgactta attaataact ggcaaaacgg cccggagctc agctctgacc 60
 tccgccacat ccgctcggca ccatgccagc gcgttgcaac ggcatagaagc gctcagggttt 120
 ttcttccgcc tgctccccac tgccgatggc catctgcacc ccagctcgtc acatttatct 180
 cgcgcacagc gtcttccccac cagttgcctt gctcatgacg ctgttaaaga tggccctacc 240

-continued

tagccgctga	gtcccacaac	gccgagatgt	ctttggccct	ttacaaggca	cgccatggcc	300
gtccaaggtc	tgttcatgag	tgtgtttg	gggccgaagg	acacctcagt	ggccacgaaa	360
tgccgccgag	cgggccagca	catgtcgaga	gagacatgga	catttatccc	cgagatgctg	420
tattagggaa	cgggtccttt	tctcggagcc	gtgatccgag	agcgttcggg	agtcgttgag	480
taaaagatgt	cgagttgccc	ttatatatcg	cgggcctgta	gctatgtgcc	ctctattctc	540
acaggttcaa	tcatcagtcc	tcgccgtgag	acgtagcggc	ctgaactagc	gctcगतatc	600
ttccgtc	atg gct ctt	cat gcc ttc	ttg ttg ctg	gca tca gca	ttg ctg	649
Met Ala Leu	His Ala Phe	Leu Leu Leu	Ala Ala Ser	Ala Leu Leu		
1	5		10			
gcc cgg ggt	gcc ctg agc	caa cct gac	aac gtc cgt	cgc gct gct	ccg	697
Ala Arg Gly	Ala Leu Ser	Gln Pro Asp	Asn Val Arg	Arg Arg Ala	Ala Pro	
15	20		25		30	
acc ggg acg	gcc gcc tgg	gat gcc gcc	cac tcg cag	gct gcc gct	gcc	745
Thr Gly Thr	Ala Ala Trp	Asp Ala Ala	His Ser Gln	Ala Ala Ala	Ala Ala	
35	40		45			
gtg tcg aga	tta tca cag	caa gac aag	atc aac att	gtc acc ggc	gtt	793
Val Ser Arg	Leu Ser Gln	Gln Gln Asp	Lys Ile Asn	Ile Val Thr	Gly Val	
50	55		60			
ggc tgg ggt	aag ggt cct	tgc gtc ggc	aat acg aac	cct gtc tac	agc	841
Gly Trp Gly	Lys Gly Pro	Cys Val Gly	Asn Thr Asn	Pro Val Tyr	Ser	
65	70		75			
atc aac tac	cca cag ctc	tgc ctg cag	gat ggc cca	ctg ggt atc	cgc	889
Ile Asn Tyr	Pro Gln Leu	Cys Leu Gln	Asp Gly Pro	Leu Gly Ile	Arg	
80	85		90			
tcc gcc acc	agc gtc acg	gcc ttc acg	ccg ggc att	caa gcc gcg	tcg	937
Ser Ala Thr	Ser Val Thr	Ala Phe Thr	Pro Gly Ile	Gln Ala Ala	Ser	
95	100		105		110	
acc tgg gat	gtg gag ttg	atc cgg cag	cgt ggt gtc	tac cta gga	cag	985
Thr Trp Asp	Val Glu Leu	Ile Arg Gln	Arg Gly Val	Tyr Leu Gly	Gln	
115	120		125			
gag gcc cgg	gga act ggc	gtg cat gtc	ctg ctc ggc	ccc gtg gcc	ggt	1033
Glu Ala Arg	Gly Thr Gly	Val His Val	Leu Leu Gly	Pro Val Ala	Gly	
130	135		140			
gct ctt ggc	aag atc ccg	cac gga ggc	cgt aac tgg	gaa gcc ttc	ggc	1081
Ala Leu Gly	Lys Ile Pro	His Gly Gly	Arg Asn Trp	Glu Ala Phe	Gly	
145	150		155			
tcc gac ccc	tac ttg gcc	ggt atc gct	atg tcc gag	acc atc gag	ggc	1129
Ser Asp Pro	Tyr Leu Ala	Gly Ile Ala	Met Ser Glu	Thr Ile Glu	Gly	
160	165		170			
att cag tcg	gag ggt gtg	cag gct tgc	gcg aag cac	tac atc gcc	aat	1177
Ile Gln Ser	Glu Gly Val	Gln Ala Cys	Ala Lys His	Tyr Ile Ala	Asn	
175	180		185		190	
gag cag gaa	ctc aac cgc	gag aca atg	agc agc aac	gtc gac gac	cgc	1225
Glu Gln Glu	Leu Asn Arg	Glu Thr Met	Ser Ser Asn	Val Asp Asp	Arg	
195	200		205			
act atg cac	gag cta tac	ctc tgg ccg	ttc gcc gac	gcc gtg cat	tcc	1273
Thr Met His	Glu Leu Tyr	Leu Trp Pro	Phe Ala Asp	Ala Val His	Ser	
210	215		220			
aac gtg gcc	agc gtc atg	tgc agc tac	aac aag ctc	aac ggc acc	tgg	1321
Asn Val Ala	Ser Val Met	Cys Ser Tyr	Asn Lys Leu	Asn Gly Thr	Trp	
225	230		235			
ctc tgc gag	aac gat agg	gcc caa aac	cag ctg ctt	aag agg gag	ctc	1369
Leu Cys Glu	Asn Asp Arg	Ala Gln Asn	Gln Leu Leu	Lys Arg Glu	Leu	
240	245		250			

-continued

ggc ttc cgc ggc tac atc gtg agc gac tgg aac gcg cag cac acc acc	1417
Gly Phe Arg Gly Tyr Ile Val Ser Asp Trp Asn Ala Gln His Thr Thr	
255 260 265 270	
gtg ggc tcg gcc aac agt ggc atg gac atg acc atg cct ggc agc gac	1465
Val Gly Ser Ala Asn Ser Gly Met Asp Met Thr Met Pro Gly Ser Asp	
275 280 285	
ttc aac ggc tgg aac gtc ctc tgg ggt ccg cag ctc aac aac gcc gtc	1513
Phe Asn Gly Trp Asn Val Leu Trp Gly Pro Gln Leu Asn Asn Ala Val	
290 295 300	
aac agc ggc cag gtc tcg cag tcc cgc ctc aac gac atg gtc cag cgc	1561
Asn Ser Gly Gln Val Ser Gln Ser Arg Leu Asn Asp Met Val Gln Arg	
305 310 315	
att ctt gct gcg tgg tac ctc ctc ggc cag aac tcc gga tac ccg tcc	1609
Ile Leu Ala Ala Trp Tyr Leu Leu Gly Gln Asn Ser Gly Tyr Pro Ser	
320 325 330	
atc aac ctg cgt gcc aac gtc caa gcc aac cac aag gag aat gtg cgt	1657
Ile Asn Leu Arg Ala Asn Val Gln Ala Asn His Lys Glu Asn Val Arg	
335 340 345 350	
gcc gta gcc cgc gat ggc atc gtc ctc ctc aag aac gac gcc att ctg	1705
Ala Val Ala Arg Asp Gly Ile Val Leu Leu Lys Asn Asp Gly Ile Leu	
355 360 365	
cct ctt cag cgt ccc aat aag att gct ctt gtc ggc tcc gcc gca gtc	1753
Pro Leu Gln Arg Pro Asn Lys Ile Ala Leu Val Gly Ser Ala Ala Val	
370 375 380	
gtc aac ccc cgt ggt atg aac gcc tgc gtg gac cgt ggc tgc aac gag	1801
Val Asn Pro Arg Gly Met Asn Ala Cys Val Asp Arg Gly Cys Asn Glu	
385 390 395	
ggt gcc ctt ggc atg ggc tgg ggc tca ggc acg gtc gag tat ccc tac	1849
Gly Ala Leu Gly Met Gly Trp Gly Ser Gly Thr Val Glu Tyr Pro Tyr	
400 405 410	
ttt gtt gcg ccg tat gat gct ctg cgt gag cgg gca cag cgc gat ggc	1897
Phe Val Ala Pro Tyr Asp Ala Leu Arg Glu Arg Ala Gln Arg Asp Gly	
415 420 425 430	
acg cag atc agt ctg cat gca tcg gac aat aca aac ggg gtt aac aac	1945
Thr Gln Ile Ser Leu His Ala Ser Asp Asn Thr Asn Gly Val Asn Asn	
435 440 445	
gcc gtg cag ggc gct gac gcg gcg ttt gtg ttc atc act gct gac tcc	1993
Ala Val Gln Gly Ala Asp Ala Ala Phe Val Phe Ile Thr Ala Asp Ser	
450 455 460	
ggc gaa ggg tac att acc gtt gag ggc cat gct ggc gac cgg aat cat	2041
Gly Glu Gly Tyr Ile Thr Val Glu Gly His Ala Gly Asp Arg Asn His	
465 470 475	
ctg gat cct tgg cat aat ggt aac cag ctt gtg cag gct gtt gcg cag	2089
Leu Asp Pro Trp His Asn Gly Asn Gln Leu Val Gln Ala Val Ala Gln	
480 485 490	
gca aat aag aac gtc att gtg gtt gtg cac agc gtt ggg ccg gtt att	2137
Ala Asn Lys Asn Val Ile Val Val Val His Ser Val Gly Pro Val Ile	
495 500 505 510	
ctg gag acg atc ctc aat acg ccc ggt gtg agg gct gtt gtt tgg gct	2185
Leu Glu Thr Ile Leu Asn Thr Pro Gly Val Arg Ala Val Val Trp Ala	
515 520 525	
ggc ttg ccg agc cag gag agc ggt aac gcg ctg gtt gat gtg ctg tac	2233
Gly Leu Pro Ser Gln Glu Ser Gly Asn Ala Leu Val Asp Val Leu Tyr	
530 535 540	
ggc ctt gtt tcg ccg tcg ggc aag ctt gtc tac acc att gcg aag agc	2281
Gly Leu Val Ser Pro Ser Gly Lys Leu Val Tyr Thr Ile Ala Lys Ser	
545 550 555	

-continued

```

ccg agc gac tac ccg act agc att gtc cgt ggc gat gat aac ttc cgc 2329
Pro Ser Asp Tyr Pro Thr Ser Ile Val Arg Gly Asp Asp Asn Phe Arg
560 565 570

gag ggt ctg ttc atc gac tac agg cac ttc gat aac gcc cgg atc gag 2377
Glu Gly Leu Phe Ile Asp Tyr Arg His Phe Asp Asn Ala Arg Ile Glu
575 580 585 590

ccc cgt ttc gag ttt ggc ttc ggt ctc t gtaagtctct taccactccg 2425
Pro Arg Phe Glu Phe Gly Phe Gly Leu
595

ttttgtaaca acccgattct aacatccccc ag ca tac acc aac ttc agc tat 2477
Ser Tyr Thr Asn Phe Ser Tyr
600 605

tcc aac ctg ggc atc tcc tcg tcc gca acc gcc ggc cca gcc acg ggc 2525
Ser Asn Leu Gly Ile Ser Ser Ser Ala Thr Ala Gly Pro Ala Thr Gly
610 615 620

ccc acc gtc ccc ggc ggc ccg gcc gac ctc tgg aac tat gtc gcg acc 2573
Pro Thr Val Pro Gly Gly Pro Ala Asp Leu Trp Asn Tyr Val Ala Thr
625 630 635

gtc acg gcg acc gtt acc aac acc ggc ggc gtg gaa ggt gcc gag gtc 2621
Val Thr Ala Thr Val Thr Asn Thr Gly Gly Val Glu Gly Ala Glu Val
640 645 650

gct cag ctg tac atc tct ttg cca tct tcg gct cct gca tcg cca ccg 2669
Ala Gln Leu Tyr Ile Ser Leu Pro Ser Ser Ala Pro Ala Ser Pro Pro
655 660 665 670

aag cag ctt cgt ggc ttt gtc aag ctt aag ttg gcg cct ggt caa agc 2717
Lys Gln Leu Arg Gly Phe Val Lys Leu Lys Leu Ala Pro Gly Gln Ser
675 680 685

ggg acg gca acg ttt aga cta agg aag agg gat ttg gct tat tgg gat 2765
Gly Thr Ala Thr Phe Arg Leu Arg Lys Arg Asp Leu Ala Tyr Trp Asp
690 695 700

gtg ggg agg cag aat tgg gtt gtt cct tcg ggg agg ttt ggc gtg ctt 2813
Val Gly Arg Gln Asn Trp Val Val Pro Ser Gly Arg Phe Gly Val Leu
705 710 715

gtg ggg gct agt tcg agg gat att agg ttg cag ggg gag att gtt gtt 2861
Val Gly Ala Ser Ser Arg Asp Ile Arg Leu Gln Gly Glu Ile Val Val
720 725 730

taggggggta tggtcagcac ctagtggggg aattgatgtg taagttggag taggggtttt 2921

cgtgtacata cataccattt ggtcaatggt acgacattta gtttatgaag tttcctgggtg 2981

gctaccgctg atgagccctc gtatgatacc cacaatctat atgttttact cttctctttc 3041

cttttttctc ttccttttcc tttattactt cattccttgt gtactttctg tgaacctcca 3101

gtcgaccatc cgacceatt cgaaagtctt tcctgacctg gttcaggttg gcatattctc 3161

gaaaggatgt cgaccttct gaccctactg ggctaccggg aaagccctag gatggctgat 3221

ggacagatct ggtgatcaac tatgggaaca ctccggagat ggtgactaat atgcgatggt 3281

catttaaaga gcaccgctc cagcgatctc cccagttgct cctcaacgat tgacacggcc 3341

aatttatcca gattccggga ttctctgagt gagctgtccc tttttctag a 3392

```

<210> SEQ ID NO 26

<211> LENGTH: 734

<212> TYPE: PRT

<213> ORGANISM: Chaetomium thermophilum

<400> SEQUENCE: 26

```

Met Ala Leu His Ala Phe Leu Leu Leu Ala Ser Ala Leu Leu Ala Arg
1 5 10 15

```

-continued

Gly Ala Leu Ser Gln Pro Asp Asn Val Arg Arg Ala Ala Pro Thr Gly
 20 25 30
 Thr Ala Ala Trp Asp Ala Ala His Ser Gln Ala Ala Ala Ala Val Ser
 35 40 45
 Arg Leu Ser Gln Gln Asp Lys Ile Asn Ile Val Thr Gly Val Gly Trp
 50 55 60
 Gly Lys Gly Pro Cys Val Gly Asn Thr Asn Pro Val Tyr Ser Ile Asn
 65 70 75 80
 Tyr Pro Gln Leu Cys Leu Gln Asp Gly Pro Leu Gly Ile Arg Ser Ala
 85 90 95
 Thr Ser Val Thr Ala Phe Thr Pro Gly Ile Gln Ala Ala Ser Thr Trp
 100 105 110
 Asp Val Glu Leu Ile Arg Gln Arg Gly Val Tyr Leu Gly Gln Glu Ala
 115 120 125
 Arg Gly Thr Gly Val His Val Leu Leu Gly Pro Val Ala Gly Ala Leu
 130 135 140
 Gly Lys Ile Pro His Gly Gly Arg Asn Trp Glu Ala Phe Gly Ser Asp
 145 150 155 160
 Pro Tyr Leu Ala Gly Ile Ala Met Ser Glu Thr Ile Glu Gly Ile Gln
 165 170 175
 Ser Glu Gly Val Gln Ala Cys Ala Lys His Tyr Ile Ala Asn Glu Gln
 180 185 190
 Glu Leu Asn Arg Glu Thr Met Ser Ser Asn Val Asp Asp Arg Thr Met
 195 200 205
 His Glu Leu Tyr Leu Trp Pro Phe Ala Asp Ala Val His Ser Asn Val
 210 215 220
 Ala Ser Val Met Cys Ser Tyr Asn Lys Leu Asn Gly Thr Trp Leu Cys
 225 230 235 240
 Glu Asn Asp Arg Ala Gln Asn Gln Leu Leu Lys Arg Glu Leu Gly Phe
 245 250 255
 Arg Gly Tyr Ile Val Ser Asp Trp Asn Ala Gln His Thr Thr Val Gly
 260 265 270
 Ser Ala Asn Ser Gly Met Asp Met Thr Met Pro Gly Ser Asp Phe Asn
 275 280 285
 Gly Trp Asn Val Leu Trp Gly Pro Gln Leu Asn Asn Ala Val Asn Ser
 290 295 300
 Gly Gln Val Ser Gln Ser Arg Leu Asn Asp Met Val Gln Arg Ile Leu
 305 310 315 320
 Ala Ala Trp Tyr Leu Leu Gly Gln Asn Ser Gly Tyr Pro Ser Ile Asn
 325 330 335
 Leu Arg Ala Asn Val Gln Ala Asn His Lys Glu Asn Val Arg Ala Val
 340 345 350
 Ala Arg Asp Gly Ile Val Leu Leu Lys Asn Asp Gly Ile Leu Pro Leu
 355 360 365
 Gln Arg Pro Asn Lys Ile Ala Leu Val Gly Ser Ala Ala Val Val Asn
 370 375 380
 Pro Arg Gly Met Asn Ala Cys Val Asp Arg Gly Cys Asn Glu Gly Ala
 385 390 395 400
 Leu Gly Met Gly Trp Gly Ser Gly Thr Val Glu Tyr Pro Tyr Phe Val
 405 410 415

-continued

Ala Pro Tyr Asp Ala Leu Arg Glu Arg Ala Gln Arg Asp Gly Thr Gln
420 425 430

Ile Ser Leu His Ala Ser Asp Asn Thr Asn Gly Val Asn Asn Ala Val
435 440 445

Gln Gly Ala Asp Ala Ala Phe Val Phe Ile Thr Ala Asp Ser Gly Glu
450 455 460

Gly Tyr Ile Thr Val Glu Gly His Ala Gly Asp Arg Asn His Leu Asp
465 470 475 480

Pro Trp His Asn Gly Asn Gln Leu Val Gln Ala Val Ala Gln Ala Asn
485 490 495

Lys Asn Val Ile Val Val Val His Ser Val Gly Pro Val Ile Leu Glu
500 505 510

Thr Ile Leu Asn Thr Pro Gly Val Arg Ala Val Val Trp Ala Gly Leu
515 520 525

Pro Ser Gln Glu Ser Gly Asn Ala Leu Val Asp Val Leu Tyr Gly Leu
530 535 540

Val Ser Pro Ser Gly Lys Leu Val Tyr Thr Ile Ala Lys Ser Pro Ser
545 550 555 560

Asp Tyr Pro Thr Ser Ile Val Arg Gly Asp Asp Asn Phe Arg Glu Gly
565 570 575

Leu Phe Ile Asp Tyr Arg His Phe Asp Asn Ala Arg Ile Glu Pro Arg
580 585 590

Phe Glu Phe Gly Phe Gly Leu Ser Tyr Thr Asn Phe Ser Tyr Ser Asn
595 600 605

Leu Gly Ile Ser Ser Ser Ala Thr Ala Gly Pro Ala Thr Gly Pro Thr
610 615 620

Val Pro Gly Gly Pro Ala Asp Leu Trp Asn Tyr Val Ala Thr Val Thr
625 630 635 640

Ala Thr Val Thr Asn Thr Gly Gly Val Glu Gly Ala Glu Val Ala Gln
645 650 655

Leu Tyr Ile Ser Leu Pro Ser Ser Ala Pro Ala Ser Pro Pro Lys Gln
660 665 670

Leu Arg Gly Phe Val Lys Leu Lys Leu Ala Pro Gly Gln Ser Gly Thr
675 680 685

Ala Thr Phe Arg Leu Arg Lys Arg Asp Leu Ala Tyr Trp Asp Val Gly
690 695 700

Arg Gln Asn Trp Val Val Pro Ser Gly Arg Phe Gly Val Leu Val Gly
705 710 715 720

Ala Ser Ser Arg Asp Ile Arg Leu Gln Gly Glu Ile Val Val
725 730

<210> SEQ ID NO 27
 <211> LENGTH: 1631
 <212> TYPE: DNA
 <213> ORGANISM: Thermoascus aurantiacus
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (610)..(674)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (675)..(1628)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(609)
 <400> SEQUENCE: 27

-continued

atg tat cag cgc gct ctt ctc ttc tct ttc ttc ctc gcc gcc gcc cgc	48
Met Tyr Gln Arg Ala Leu Leu Phe Ser Phe Phe Leu Ala Ala Ala Arg	
1 5 10 15	
gcg cag cag gcc ggt acc gta acc gca gag aat cac cct tcc ctg acc	96
Ala Gln Gln Ala Gly Thr Val Thr Ala Glu Asn His Pro Ser Leu Thr	
20 25 30	
tgg cag caa tgc tcc agc ggc ggt agt tgt acc acg cag aat gga aaa	144
Trp Gln Gln Cys Ser Ser Gly Gly Ser Cys Thr Thr Gln Asn Gly Lys	
35 40 45	
gtc gtt atc gat gcg aac tgg cgt tgg gtc cat acc acc tct gga tac	192
Val Val Ile Asp Ala Asn Trp Arg Trp Val His Thr Thr Ser Gly Tyr	
50 55 60	
acc aac tgc tac acg ggc aat acg tgg gac acc agt atc tgt ccc gac	240
Thr Asn Cys Tyr Thr Gly Asn Thr Trp Asp Thr Ser Ile Cys Pro Asp	
65 70 75 80	
gac gtg acc tgc gct cag aat tgt gcc ttg gat gga gcg gat tac agt	288
Asp Val Thr Cys Ala Gln Asn Cys Ala Leu Asp Gly Ala Asp Tyr Ser	
85 90 95	
ggc acc tat ggt gtt acg acc agt ggc aac gcc ctg aga ctg aac ttt	336
Gly Thr Tyr Gly Val Thr Thr Ser Gly Asn Ala Leu Arg Leu Asn Phe	
100 105 110	
gtc acc caa agc tca ggg aag aac att ggc tcg cgc ctg tac ctg ctg	384
Val Thr Gln Ser Ser Gly Lys Asn Ile Gly Ser Arg Leu Tyr Leu Leu	
115 120 125	
cag gac gac acc act tat cag atc ttc aag ctg ctg ggt cag gag ttt	432
Gln Asp Asp Thr Thr Tyr Gln Ile Phe Lys Leu Leu Gly Gln Glu Phe	
130 135 140	
acc ttc gat gtc gac gtc tcc aat ctc cct tgc ggg ctg aac ggc gcc	480
Thr Phe Asp Val Asp Val Ser Asn Leu Pro Cys Gly Leu Asn Gly Ala	
145 150 155 160	
ctc tac ttt gtg gcc atg gac gcc gac ggc gga ttg tcc aaa tac cct	528
Leu Tyr Phe Val Ala Met Asp Ala Asp Gly Gly Leu Ser Lys Tyr Pro	
165 170 175	
ggc aac aag gca ggc gct aag tat ggc act ggt tac tgc gac tct cag	576
Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp Ser Gln	
180 185 190	
tgc cct cgg gat ctc aag ttc atc aac ggt cag gtacgtcaga agtgataact	629
Cys Pro Arg Asp Leu Lys Phe Ile Asn Gly Gln	
195 200	
agccagcaga gcccatgaat cattaactaa cgctgtcaaa tacag gcc aat gtt gaa	686
Ala Asn Val Glu	
205	
ggc tgg cag ccg tct gcc aac gac cca aat gcc ggc gtt ggt aac cac	734
Gly Trp Gln Pro Ser Ala Asn Asp Pro Asn Ala Gly Val Gly Asn His	
210 215 220	
ggt tcc tgc tgc gct gag atg gat gtc tgg gaa gcc aac agc atc tct	782
Gly Ser Cys Cys Ala Glu Met Asp Val Trp Glu Ala Asn Ser Ile Ser	
225 230 235	
act gcg gtg acg cct cac cca tgc gac acc ccc ggc cag acc atg tgc	830
Thr Ala Val Thr Pro His Pro Cys Asp Thr Pro Gly Gln Thr Met Cys	
240 245 250 255	
cag gga gac gac tgt ggt gga acc tac tcc tcc act cga tat gct ggt	878
Gln Gly Asp Asp Cys Gly Gly Thr Tyr Ser Ser Thr Arg Tyr Ala Gly	
260 265 270	
acc tgc gac cct gat ggc tgc gac ttc aat cct tac cgc cag ggc aac	926
Thr Cys Asp Pro Asp Gly Cys Asp Phe Asn Pro Tyr Arg Gln Gly Asn	
275 280 285	

-continued

```

cac tcg ttc tac ggc ccc ggg cag atc gtc gac acc agc tcc aaa ttc      974
His Ser Phe Tyr Gly Pro Gly Gln Ile Val Asp Thr Ser Ser Lys Phe
290                               295                               300

acc gtc gtc acc cag ttc atc acc gac gac ggg acc ccc tcc ggc acc      1022
Thr Val Val Thr Gln Phe Ile Thr Asp Asp Gly Thr Pro Ser Gly Thr
305                               310                               315

ctg acg gag atc aaa cgc ttc tac gtc cag aac ggc aag gta atc ccc      1070
Leu Thr Glu Ile Lys Arg Phe Tyr Val Gln Asn Gly Lys Val Ile Pro
320                               325                               330                               335

cag tcg gag tcg acg atc agc ggc gtc acc ggc aac tca atc acc acc      1118
Gln Ser Glu Ser Thr Ile Ser Gly Val Thr Gly Asn Ser Ile Thr Thr
340                               345                               350

gag tat tgc acg gcc cag aag gcc gcc ttc ggc gac aac acc ggc ttc      1166
Glu Tyr Cys Thr Ala Gln Lys Ala Ala Phe Gly Asp Asn Thr Gly Phe
355                               360                               365

ttc acg cac ggc ggg ctt cag aag atc agt cag gct ctg gct cag ggc      1214
Phe Thr His Gly Gly Leu Gln Lys Ile Ser Gln Ala Leu Ala Gln Gly
370                               375                               380

atg gtc ctc gtc atg agc ctg tgg gac gat cac gcc gcc aac atg ctc      1262
Met Val Leu Val Met Ser Leu Trp Asp Asp His Ala Ala Asn Met Leu
385                               390                               395

tgg ctg gac agc acc tac ccg act gat gcg gac ccg gac acc cct ggc      1310
Trp Leu Asp Ser Thr Tyr Pro Thr Asp Ala Asp Pro Asp Thr Pro Gly
400                               405                               410                               415

gtc gcg cgc ggt acc tgc ccc acg acc tcc ggc gtc ccg gcc gac gtt      1358
Val Ala Arg Gly Thr Cys Pro Thr Thr Ser Gly Val Pro Ala Asp Val
420                               425                               430

gag tcg cag tac ccc aat tca tat gtt atc tac tcc aac atc aag gtc      1406
Glu Ser Gln Tyr Pro Asn Ser Tyr Val Ile Tyr Ser Asn Ile Lys Val
435                               440                               445

gga ccc att ggc agc acc ggc aac cct agc ggc ggc aac cct ccc ggc      1454
Gly Pro Ile Gly Ser Thr Gly Asn Pro Ser Gly Gly Asn Pro Pro Gly
450                               455                               460

gga aac ccg cct ggc acc acc acc acc cgc cgc cca gcc act acc act      1502
Gly Asn Pro Pro Gly Thr Thr Thr Thr Arg Arg Pro Ala Thr Thr Thr
465                               470                               475

gga agc tct ccc gga cct acc cag tct cac tac ggc cag tgc ggc ggt      1550
Gly Ser Ser Pro Gly Pro Thr Gln Ser His Tyr Gly Gln Cys Gly Gly
480                               485                               490                               495

att ggc tac agc ggc ccc acg gtc tgc gcc agc ggc aca act tgc cag      1598
Ile Gly Tyr Ser Gly Pro Thr Val Cys Ala Ser Gly Thr Thr Cys Gln
500                               505                               510

gtc ctg aac cct tac tac tct cag tgc ctg taa                          1631
Val Leu Asn Pro Tyr Tyr Ser Gln Cys Leu
515                               520

```

<210> SEQ ID NO 28

<211> LENGTH: 521

<212> TYPE: PRT

<213> ORGANISM: Thermoascus aurantiacus

<400> SEQUENCE: 28

```

Met Tyr Gln Arg Ala Leu Leu Phe Ser Phe Phe Leu Ala Ala Ala Arg
1                               5                               10                               15

Ala Gln Gln Ala Gly Thr Val Thr Ala Glu Asn His Pro Ser Leu Thr
20                               25                               30

Trp Gln Gln Cys Ser Ser Gly Gly Ser Cys Thr Thr Gln Asn Gly Lys

```


-continued

Pro Ile Gly Ser Thr Gly Asn Pro Ser Gly Gly Asn Pro Pro Gly Gly
450 455 460

Asn Pro Pro Gly Thr Thr Thr Thr Arg Arg Pro Ala Thr Thr Thr Gly
465 470 475 480

Ser Ser Pro Gly Pro Thr Gln Ser His Tyr Gly Gln Cys Gly Gly Ile
485 490 495

Gly Tyr Ser Gly Pro Thr Val Cys Ala Ser Gly Thr Thr Cys Gln Val
500 505 510

Leu Asn Pro Tyr Tyr Ser Gln Cys Leu
515 520

<210> SEQ ID NO 29
 <211> LENGTH: 1734
 <212> TYPE: DNA
 <213> ORGANISM: Thermoascus aurantiacus
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (610)..(674)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1726)..(1731)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (675)..(1661)
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1)..(609)
 <220> FEATURE:
 <221> NAME/KEY: Intron
 <222> LOCATION: (1662)..(1725)

<400> SEQUENCE: 29

atg tat cag cgc gct ctt ctc ttc tct ttc ttc ctc gcc gcc gcc cgc	48
Met Tyr Gln Arg Ala Leu Leu Phe Ser Phe Phe Leu Ala Ala Ala Arg	
1 5 10 15	
gcg cag cag gcc ggt acc gta acc gca gag aat cac cct tcc ctg acc	96
Ala Gln Gln Ala Gly Thr Val Thr Ala Glu Asn His Pro Ser Leu Thr	
20 25 30	
tgg cag caa tgc tcc agc ggc ggt agt tgt acc acg cag aat gga aaa	144
Trp Gln Gln Cys Ser Ser Gly Gly Ser Cys Thr Thr Gln Asn Gly Lys	
35 40 45	
gtc gtt atc gat gcg aac tgg cgt tgg gtc cat acc acc tct gga tac	192
Val Val Ile Asp Ala Asn Trp Arg Trp Val His Thr Thr Ser Gly Tyr	
50 55 60	
acc aac tgc tac acg ggc aat acg tgg gac acc agt atc tgt ccc gac	240
Thr Asn Cys Tyr Thr Gly Asn Thr Trp Asp Thr Ser Ile Cys Pro Asp	
65 70 75 80	
gac gtg acc tgc gct cag aat tgt gcc ttg gat gga gcg gat tac agt	288
Asp Val Thr Cys Ala Gln Asn Cys Ala Leu Asp Gly Ala Asp Tyr Ser	
85 90 95	
ggc acc tat ggt gtt acg acc agt ggc aac gcc ctg aga ctg aac ttt	336
Gly Thr Tyr Gly Val Thr Thr Ser Gly Asn Ala Leu Arg Leu Asn Phe	
100 105 110	
gtc acc caa agc tca ggg aag aac att ggc tcg cgc ctg tac ctg ctg	384
Val Thr Gln Ser Ser Gly Lys Asn Ile Gly Ser Arg Leu Tyr Leu Leu	
115 120 125	
cag gac gac acc act tat cag atc ttc aag ctg ctg ggt cag gag ttt	432
Gln Asp Asp Thr Thr Tyr Gln Ile Phe Lys Leu Leu Gly Gln Glu Phe	
130 135 140	
acc ttc gat gtc gac gtc tcc aat ctc cct tgc ggg ctg aac ggc gcc	480

-continued

Thr	Phe	Asp	Val	Asp	Val	Ser	Asn	Leu	Pro	Cys	Gly	Leu	Asn	Gly	Ala		
145					150					155					160		
ctc	tac	ttt	gtg	gcc	atg	gac	gcc	gac	ggc	gga	ttg	tcc	aaa	tac	cct		528
Leu	Tyr	Phe	Val	Ala	Met	Asp	Ala	Asp	Gly	Gly	Leu	Ser	Lys	Tyr	Pro		
165					170					175							
ggc	aac	aag	gca	ggc	gct	aag	tat	ggc	act	ggg	tac	tgc	gac	tct	cag		576
Gly	Asn	Lys	Ala	Gly	Ala	Lys	Tyr	Gly	Thr	Gly	Tyr	Cys	Asp	Ser	Gln		
180					185					190							
tgc	cct	cgg	gat	ctc	aag	ttc	atc	aac	ggg	cag	gtacg	tcaga	agtgataact				629
Cys	Pro	Arg	Asp	Leu	Lys	Phe	Ile	Asn	Gly	Gln							
195					200												
agccagcaga	gccccatgaat	cattaactaa	cgctgtcaaaa	tacag	gcc	aat	ggt	gaa									686
Ala	Asn	Val	Glu														
205																	
ggc	tgg	cag	ccg	tct	gcc	aac	gac	cca	aat	gcc	ggc	ggt	ggt	aac	cac		734
Gly	Trp	Gln	Pro	Ser	Ala	Asn	Asp	Pro	Asn	Ala	Gly	Val	Gly	Asn	His		
210					215					220							
ggg	tcc	tgc	tgc	gct	gag	atg	gat	gtc	tgg	gaa	gcc	aac	agc	atc	tct		782
Gly	Ser	Cys	Cys	Ala	Glu	Met	Asp	Val	Trp	Glu	Ala	Asn	Ser	Ile	Ser		
225					230					235							
act	gcg	gtg	acg	cct	cac	cca	tgc	gac	acc	ccc	ggc	cag	acc	atg	tgc		830
Thr	Ala	Val	Thr	Pro	His	Pro	Cys	Asp	Thr	Pro	Gly	Gln	Thr	Met	Cys		
240					245					250					255		
cag	gga	gac	gac	tgt	ggg	gga	acc	tac	tcc	tcc	act	cga	tat	gct	ggg		878
Gln	Gly	Asp	Asp	Cys	Gly	Gly	Thr	Tyr	Ser	Ser	Thr	Arg	Tyr	Ala	Gly		
260					265					270							
acc	tgc	gac	cct	gat	ggc	tgc	gac	ttc	aat	cct	tac	cgc	cag	ggc	aac		926
Thr	Cys	Asp	Pro	Asp	Gly	Cys	Asp	Phe	Asn	Pro	Tyr	Arg	Gln	Gly	Asn		
275					280					285							
cac	tcg	ttc	tac	ggc	ccc	ggg	cag	atc	gtc	gac	acc	agc	tcc	aaa	ttc		974
His	Ser	Phe	Tyr	Gly	Pro	Gly	Gln	Ile	Val	Asp	Thr	Ser	Ser	Lys	Phe		
290					295					300							
acc	gtc	gtc	acc	cag	ttc	atc	acc	gac	gac	ggg	acc	ccc	tcc	ggc	acc		1022
Thr	Val	Val	Thr	Gln	Phe	Ile	Thr	Asp	Asp	Gly	Thr	Pro	Ser	Gly	Thr		
305					310					315							
ctg	acg	gag	atc	aaa	cgc	ttc	tac	gtc	cag	aac	ggc	aag	gta	atc	ccc		1070
Leu	Thr	Glu	Ile	Lys	Arg	Phe	Tyr	Val	Gln	Asn	Gly	Lys	Val	Ile	Pro		
320					325					330					335		
cag	tcg	gag	tcg	acg	atc	agc	ggc	gtc	acc	ggc	aac	tca	atc	acc	acc		1118
Gln	Ser	Glu	Ser	Thr	Ile	Ser	Gly	Val	Thr	Gly	Asn	Ser	Ile	Thr	Thr		
340					345					350							
gag	tat	tgc	acg	gcc	cag	aag	gcc	gcc	ttc	ggc	gac	aac	acc	ggc	ttc		1166
Glu	Tyr	Cys	Thr	Ala	Gln	Lys	Ala	Ala	Phe	Gly	Asp	Asn	Thr	Gly	Phe		
355					360					365							
ttc	acg	cac	ggc	ggg	ctt	cag	aag	atc	agt	cag	gct	ctg	gct	cag	ggc		1214
Phe	Thr	His	Gly	Gly	Leu	Gln	Lys	Ile	Ser	Gln	Ala	Leu	Ala	Gln	Gly		
370					375					380							
atg	gtc	ctc	gtc	atg	agc	ctg	tgg	gac	gat	cac	gcc	gcc	aac	atg	ctc		1262
Met	Val	Leu	Val	Met	Ser	Leu	Trp	Asp	Asp	His	Ala	Ala	Asn	Met	Leu		
385					390					395							
tgg	ctg	gac	agc	acc	tac	ccg	act	gat	gcg	gac	ccg	gac	acc	cct	ggc		1310
Trp	Leu	Asp	Ser	Thr	Tyr	Pro	Thr	Asp	Ala	Asp	Pro	Asp	Thr	Pro	Gly		
400					405					410					415		
gtc	gcg	cgc	ggg	acc	tgc	ccc	acg	acc	tcc	ggc	gtc	ccg	gcc	gac	ggt		1358
Val	Ala	Arg	Gly	Thr	Cys	Pro	Thr	Thr	Ser	Gly	Val	Pro	Ala	Asp	Val		
420					425					430							
gag	tcg	cag	tac	ccc	aat	tca	tat	ggt	atc	tac	tcc	aac	atc	aag	gtc		1406

-continued

Glu Ser Gln Tyr Pro Asn Ser Tyr Val Ile Tyr Ser Asn Ile Lys Val
435 440 445

gga ccc atc ggc tcg acc gtc cct ggc ctt gac ggc agc aac ccc ggc 1454
Gly Pro Ile Gly Ser Thr Val Pro Gly Leu Asp Gly Ser Asn Pro Gly
450 455 460

aac ccg acc acc acc gtc gtt cct ccc gct tct acc tcc acc tcc cgt 1502
Asn Pro Thr Thr Thr Val Val Pro Pro Ala Ser Thr Ser Thr Ser Arg
465 470 475

ccg acc agc agc act agc tct ccc gtt tcg acc ccg act ggc cag ccc 1550
Pro Thr Ser Ser Thr Ser Ser Pro Val Ser Thr Pro Thr Gly Gln Pro
480 485 490 495

ggc ggc tgc acc acc cag aag tgg ggc cag tgc ggc ggt atc ggc tac 1598
Gly Gly Cys Thr Thr Gln Lys Trp Gly Gln Cys Gly Gly Ile Gly Tyr
500 505 510

acc ggc tgc act aac tgc gtt gct ggc acc acc tgc act cag ctc aac 1646
Thr Gly Cys Thr Asn Cys Val Ala Gly Thr Thr Cys Thr Gln Leu Asn
515 520 525

ccc tgg tac agc cag gtatgtttct cttccccctt ctagactcgc ttggatttga 1701
Pro Trp Tyr Ser Gln
530

cagttgctaa catctgctca acag tgc ctg taa 1734
Cys Leu

<210> SEQ ID NO 30

<211> LENGTH: 534

<212> TYPE: PRT

<213> ORGANISM: *Thermoascus aurantiacus*

<400> SEQUENCE: 30

Met Tyr Gln Arg Ala Leu Leu Phe Ser Phe Phe Leu Ala Ala Ala Arg
1 5 10 15

Ala Gln Gln Ala Gly Thr Val Thr Ala Glu Asn His Pro Ser Leu Thr
20 25 30

Trp Gln Gln Cys Ser Ser Gly Gly Ser Cys Thr Thr Gln Asn Gly Lys
35 40 45

Val Val Ile Asp Ala Asn Trp Arg Trp Val His Thr Thr Ser Gly Tyr
50 55 60

Thr Asn Cys Tyr Thr Gly Asn Thr Trp Asp Thr Ser Ile Cys Pro Asp
65 70 75 80

Asp Val Thr Cys Ala Gln Asn Cys Ala Leu Asp Gly Ala Asp Tyr Ser
85 90 95

Gly Thr Tyr Gly Val Thr Thr Ser Gly Asn Ala Leu Arg Leu Asn Phe
100 105 110

Val Thr Gln Ser Ser Gly Lys Asn Ile Gly Ser Arg Leu Tyr Leu Leu
115 120 125

Gln Asp Asp Thr Thr Tyr Gln Ile Phe Lys Leu Leu Gly Gln Glu Phe
130 135 140

Thr Phe Asp Val Asp Val Ser Asn Leu Pro Cys Gly Leu Asn Gly Ala
145 150 155 160

Leu Tyr Phe Val Ala Met Asp Ala Asp Gly Gly Leu Ser Lys Tyr Pro
165 170 175

Gly Asn Lys Ala Gly Ala Lys Tyr Gly Thr Gly Tyr Cys Asp Ser Gln
180 185 190

Cys Pro Arg Asp Leu Lys Phe Ile Asn Gly Gln Ala Asn Val Glu Gly
195 200 205

-continued

Trp Gln Pro Ser Ala Asn Asp Pro Asn Ala Gly Val Gly Asn His Gly
 210 215 220
 Ser Cys Cys Ala Glu Met Asp Val Trp Glu Ala Asn Ser Ile Ser Thr
 225 230 235 240
 Ala Val Thr Pro His Pro Cys Asp Thr Pro Gly Gln Thr Met Cys Gln
 245 250 255
 Gly Asp Asp Cys Gly Gly Thr Tyr Ser Ser Thr Arg Tyr Ala Gly Thr
 260 265 270
 Cys Asp Pro Asp Gly Cys Asp Phe Asn Pro Tyr Arg Gln Gly Asn His
 275 280 285
 Ser Phe Tyr Gly Pro Gly Gln Ile Val Asp Thr Ser Ser Lys Phe Thr
 290 295 300
 Val Val Thr Gln Phe Ile Thr Asp Asp Gly Thr Pro Ser Gly Thr Leu
 305 310 315 320
 Thr Glu Ile Lys Arg Phe Tyr Val Gln Asn Gly Lys Val Ile Pro Gln
 325 330 335
 Ser Glu Ser Thr Ile Ser Gly Val Thr Gly Asn Ser Ile Thr Thr Glu
 340 345 350
 Tyr Cys Thr Ala Gln Lys Ala Ala Phe Gly Asp Asn Thr Gly Phe Phe
 355 360 365
 Thr His Gly Gly Leu Gln Lys Ile Ser Gln Ala Leu Ala Gln Gly Met
 370 375 380
 Val Leu Val Met Ser Leu Trp Asp Asp His Ala Ala Asn Met Leu Trp
 385 390 395 400
 Leu Asp Ser Thr Tyr Pro Thr Asp Ala Asp Pro Asp Thr Pro Gly Val
 405 410 415
 Ala Arg Gly Thr Cys Pro Thr Thr Ser Gly Val Pro Ala Asp Val Glu
 420 425 430
 Ser Gln Tyr Pro Asn Ser Tyr Val Ile Tyr Ser Asn Ile Lys Val Gly
 435 440 445
 Pro Ile Gly Ser Thr Val Pro Gly Leu Asp Gly Ser Asn Pro Gly Asn
 450 455 460
 Pro Thr Thr Thr Val Val Pro Pro Ala Ser Thr Ser Thr Ser Arg Pro
 465 470 475 480
 Thr Ser Ser Thr Ser Ser Pro Val Ser Thr Pro Thr Gly Gln Pro Gly
 485 490 495
 Gly Cys Thr Thr Gln Lys Trp Gly Gln Cys Gly Gly Ile Gly Tyr Thr
 500 505 510
 Gly Cys Thr Asn Cys Val Ala Gly Thr Thr Cys Thr Gln Leu Asn Pro
 515 520 525
 Trp Tyr Ser Gln Cys Leu
 530

1. A method for treating cellulosic material with cellobiohydrolase, endoglucanase and beta-glucosidase, whereby said cellobiohydrolase comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 2, 4, 6 or 8, or to an enzymatically active fragment thereof.

2-4. (canceled)

5. The method of claim 1, comprising using a cellobiohydrolase obtainable from *Thermoascus aurantiacus* or *Acremonium thermophilum* to which a cellulose binding domain,

derived from *Trichoderma reesei* or *Chaetomium thermophilum*, has been genetically attached, and the resulting fusion protein comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 28 or 30.

6. (canceled)

7. The method of claim 1, wherein the endoglucanase comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 10, 12, 14 or 16, or to an enzymatically active fragment thereof.

- 8-9.** (canceled)
- 10.** The method of claim **1**, wherein the beta-glucosidase comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 22, 24 or 26, or to an enzymatically active fragment thereof.
- 11.** (canceled)
- 12.** The method of claim **1**, wherein the cellulosic material is lignocellulosic material.
- 13.** The method of claim **1**, comprising treating lignocellulosic material with at least one further enzyme having an amino acid sequence having at least 80% identity to SEQ ID NOS: 18 or 20, or to an enzymatically active fragment thereof.
- 14-17.** (canceled)
- 18.** An enzyme preparation comprising cellobiohydrolase, endoglucanase and beta-glucosidase, wherein said cellobiohydrolase comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 2, 4, 6 or 8, or to an enzymatically active fragment thereof.
- 19-20.** (canceled)
- 21.** The enzyme preparation of claim **18**, wherein the enzymes are recombinant enzymes, comprising a cellobiohydrolase obtainable from *Thermoascus aurantiacus* to which a cellulose binding domain has been genetically attached.
- 22.** (canceled)
- 23.** The enzyme preparation of claim **21**, wherein the attached cellulose binding domain is derived from *Trichoderma reesei* or *Chaetomium termophilum*, and the resulting fusion protein comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 28 or 30.
- 24.** The enzyme preparation of claim **18**, wherein the endoglucanase comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 10, 12, 14 or 16, or to an enzymatically active fragment thereof.
- 25.** (canceled)
- 26.** The enzyme preparation of claim **18**, wherein the beta-glucosidase comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 22, 24 or 26, or to an enzymatically active fragment thereof.
- 27.** (canceled)
- 28.** The enzyme preparation of claim **18**, comprising at least one xylanase, which comprises an amino acid sequence having at least 80% identity to SEQ ID NOS: 18 or 20, or to an enzymatically active fragment thereof.
- 29.** (canceled)
- 30.** The enzyme preparation of claim **18**, wherein at least one of the enzymes is encoded by a gene similar to that included in a microorganism having accession number DSM 16723, DSM 16728, DSM 16729, DSM 16727, DSM 17326, DSM 17324, DSM 17323, DSM 17729, DSM 16723, DSM 16726, DSM 16725, DSM 17325 or DSM 17667.
- 31.** (canceled)
- 32.** Use of an enzyme preparation according to claim **18** for degrading cellulosic material.
- 33.** (canceled)
- 34.** Use of the method according to claim **1** in a process for preparing ethanol from cellulosic material.
- 35.** A polypeptide comprising a fragment having cellulolytic activity and being selected from the group consisting of:
- a polypeptide comprising an amino acid sequence having at least 66% identity to SEQ ID NO:4, 79% identity to SEQ ID NO:6, 78% identity to SEQ ID NO:12, 68% identity to SEQ ID NO: 14, 72% identity to SEQ ID NO: 16, 68% identity to SEQ ID NO:20, 74% identity to SEQ ID NOS:22 or 24, or 78% identity to SEQ ID NO:26;
 - a variant of a) comprising a fragment having cellulolytic activity; and
 - a fragment of a) or b) having cellulolytic activity.
- 36.** (canceled)
- 37.** An isolated polynucleotide selected from the group consisting of:
- a nucleotide sequence of SEQ ID NOS: 3, 5, 11, 13, 15, 19, 21, 23 or 25, or a sequence encoding a polypeptide of claim **35**;
 - a complementary strand of a)
 - a fragment of a) or b) comprising at least 20 nucleotides; and
 - a sequence that is degenerate as a result of the genetic code to any one of the sequences as defined in a), b) or c).
- 38-39.** (canceled)
- 40.** A vector, which comprises as a heterologous sequence a polynucleotide of claim **37**.
- 41.** The vector of claim **40**, which is capable of expressing a polypeptide comprising a fragment having cellulolytic activity and being selected from the group consisting of:
- a polypeptide comprising an amino acid sequence having at least 66% identity to SEQ ID NO:4, 79% identity to SEQ ID NO:6, 78% identity to SEQ ID NO:12, 68% identity to SEQ ID NO:14, 72% identity to SEQ ID NO:16, 68% identity to SEQ ID NO:20, 74% identity to SEQ ID NOS:22 or 24, or 78% identity to SEQ ID NO:26;
 - a variant of a) comprising a fragment having cellulolytic activity; and
 - a fragment of a) or b) having cellulolytic activity.
- 42.** A host cell comprising the vector of claim **40**.
- 43-44.** (canceled)
- 45.** An *Escherichia coli* strain having accession number DSM 16728, DSM 16729, DSM 17324, DSM 17323, DSM 17729, DSM 16726, DSM 16725, DSM 17325 or DSM 17667.
- 46.** An enzyme preparation comprising a polypeptide of claim **35**.
- 47-49.** (canceled)
- 50.** Use of a polypeptide according to claim **35** or an enzyme preparation comprising a polypeptide of claim **35** in an industry selected from the group consisting of fuel industry, textile industry, detergent industry, pulp and paper industry, food industry, feed industry and beverage industry.
- 51.** (canceled)
- 52.** The use according to claim **50**, wherein the enzyme preparation is spent culture medium.
- 53.** A method for preparing a polypeptide comprising a fragment having cellulolytic activity and being selected from the group consisting of:
- a polypeptide comprising an amino acid sequence having at least 66% identity to SEQ ID NO:4, 79% identity to SEQ ID NO:6, 78% identity to SEQ ID NO:12, 68% identity to SEQ ID NO:14, 72% identity to SEQ ID NO:16, 68% identity to SEQ ID NO:20, 74% identity to SEQ ID NOS:22 or 24, or 78% identity to SEQ ID NO:26;
 - a variant of a) comprising a fragment having cellulolytic activity; and
 - a fragment of a) or b) having cellulolytic activity, said method comprising transforming a host cell with a vector encoding said polypeptide, and culturing said

host cell under conditions enabling expression of said polypeptide, and optionally recovering and purifying the polypeptide produced.

54. A method of treating cellulosic material with a spent culture medium of at least one microorganism capable of producing a polypeptide comprising a fragment having cellulolytic activity and being selected from the group consisting of:

a) a polypeptide comprising an amino acid sequence having at least 66% identity to SEQ ID NO:4, 79% identity to SEQ ID NO:6, 78% identity to SEQ ID NO:12, 68%

identity to SEQ ID NO:14, 72% identity to SEQ ID NO:16, 68% identity to SEQ ID NO:20, 74% identity to SEQ ID NO:22 or 24, or 78% identity to SEQ ID NO:26;
b) a variant of a) comprising a fragment having cellulolytic activity; and
c) a fragment of a) or b) having cellulolytic activity,
said method comprising reacting the cellulosic material with the spent culture medium to obtain hydrolysed cellulosic material.

* * * * *