



US 20100086965A1

(19) **United States**

(12) **Patent Application Publication**
Van Maris et al.

(10) **Pub. No.: US 2010/0086965 A1**

(43) **Pub. Date: Apr. 8, 2010**

(54) **METABOLIC ENGINEERING OF ARABINOSE-FERMENTING YEAST CELLS**

(30) **Foreign Application Priority Data**

Oct. 2, 2006 (EP) 06121633.9

(76) Inventors: **Antonius Jeroen Adriaan Van Maris**, Delft (NL); **Jacobus Thomas Pronk**, Schipluiden (NL); **Hendrik Wouter Wisselink**, Culemborg (NL); **Johannes Pieter Van Dijken**, Leidschendam (NL); **Aaron Adriaan Winkler**, Den Haag (NL); **Johannes Hendrik De Winde**, Voorhout (NL)

Publication Classification

(51) **Int. Cl.**
C12P 35/00 (2006.01)
C12N 1/19 (2006.01)
C12N 15/74 (2006.01)
C12P 7/40 (2006.01)
C12P 7/54 (2006.01)
C12P 7/52 (2006.01)
C12P 7/48 (2006.01)
C12P 7/46 (2006.01)
C12P 7/06 (2006.01)
C12P 7/04 (2006.01)
C12P 7/16 (2006.01)
C12P 13/04 (2006.01)
C12P 5/02 (2006.01)
C12P 19/44 (2006.01)

Correspondence Address:
NIXON & VANDERHYE, PC
901 NORTH GLEBE ROAD, 11TH FLOOR
ARLINGTON, VA 22203 (US)

(52) **U.S. Cl.** **435/47**; 435/254.2; 435/254.21; 435/254.22; 435/254.23; 435/320.1; 435/136; 435/140; 435/141; 435/144; 435/145; 435/161; 435/157; 435/160; 435/106; 435/167; 435/74

(21) Appl. No.: **12/442,013**

(22) PCT Filed: **Oct. 1, 2007**

(86) PCT No.: **PCT/NL07/00246**

§ 371 (c)(1),
(2), (4) Date: **Jun. 12, 2009**

(57) **ABSTRACT**

The invention relates to an eukaryotic cell expressing nucleotide sequences encoding the ara A, ara B and ara D enzymes whereby the expression of these nucleotide sequences confers on the cell the ability to use L-arabinose and/or convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product such as ethanol. Optionally, the eukaryotic cell is also able to convert xylose into ethanol.

Related U.S. Application Data

(60) Provisional application No. 60/848,357, filed on Oct. 2, 2006.





Fig 1




Fig 2




Fig 3




Fig 4




Fig 5a




Fig 5b




Fig 5c




Fig 5d




Fig 6





Fig 7



METABOLIC ENGINEERING OF ARABINOSE-FERMENTING YEAST CELLS

FIELD OF THE INVENTION

[0001] The invention relates to an eukaryotic cell having the ability to use L-arabinose and/or to convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product and to a process for producing a fermentation product wherein this cell is used.

BACKGROUND OF THE INVENTION

[0002] Fuel ethanol is acknowledged as a valuable alternative to fossil fuels. Economically viable ethanol production from the hemicellulose fraction of plant biomass requires the simultaneous fermentative conversion of both pentoses and hexoses at comparable rates and with high yields. Yeasts, in particular *Saccharomyces* spp., are the most appropriate candidates for this process since they can grow and ferment fast on hexoses, both aerobically and anaerobically. Furthermore they are much more resistant to the toxic environment of lignocellulose hydrolysates than (genetically modified) bacteria.

[0003] EP 1 499 708 describes a process for making *S. cerevisiae* strains able to produce ethanol from L-arabinose. These strains were modified by introducing the araA (L-arabinose isomerase) gene from *Bacillus subtilis*, the araB (L-ribulokinase) and araD (L-ribulose-5-P4-epimerase) genes from *Escherichia coli*. Furthermore, these strains were either carrying additional mutations in their genome or over-expressing a TAL1 (transaldolase) gene. However, these strains have several drawbacks. They ferment arabinose in oxygen limited conditions. In addition, they have a low ethanol production rate of $0.05 \text{ g}\cdot\text{g}^{-1}\cdot\text{h}^{-1}$ (Becker and Boles, 2003). Furthermore, these strains are not able to use L-arabinose under anaerobic conditions. Finally, these *S. cerevisiae* strains have a wild type background, therefore they can not be used to co-ferment several C5 sugars.

[0004] WO 03/062430 and WO 06/009434 disclose yeast strains able to convert xylose into ethanol. These yeast strains are able to directly isomerise xylose into xylulose.

[0005] Still, there is a need for alternative strains for producing ethanol, which perform better and are more robust and resistant to relatively harsh production conditions.

DESCRIPTION OF THE FIGURES

[0006] FIG. 1. Plasmid maps of pRW231 and pRW243.

[0007] FIG. 2. Growth pattern of shake flask cultivations of strain RWB219 (○) and IMS0001 (●) in synthetic medium containing 0.5% galactose (A) and 0.1% galactose +2% L-arabinose (B). Cultures were grown for 72 hours in synthetic medium with galactose (A) and then transferred to synthetic medium with galactose and arabinose (B). Growth was determined by measuring the OD_{660} .

[0008] FIG. 3. Growth rate during serial transfers of *S. cerevisiae* IMS0001 in shake flask cultures containing synthetic medium with 2% (w/v) L-arabinose. Each datapoint represents the growth rate estimated from the OD_{660} measured during (exponential) growth. The closed and open circles represent duplicate serial transfer experiments.

[0009] FIG. 4. Growth rate during an anaerobic SBR fermentation of *S. cerevisiae* IMS0001 in synthetic medium

with 2% (w/v) L-arabinose. Each datapoint represents the growth rate estimated from the CO_2 profile (solid line) during exponential growth.

[0010] FIG. 5. Sugar consumption and product formation during anaerobic batch fermentations of strain IMS0002. The fermentations were performed in 1 synthetic medium supplemented with: 20 g l^{-1} arabinose (A); 20 g l^{-1} glucose and 20 g l^{-1} arabinose (B); 30 g l^{-1} glucose, 15 g l^{-1} xylose, and 15 g l^{-1} arabinose (C); Sugar consumption and product formation during anaerobic batch fermentations with a mixture of strains IMS0002 and RWB218. The fermentations were performed in 1 liter of synthetic medium supplemented with 30 g l^{-1} glucose, 15 g l^{-1} xylose, and 15 g l^{-1} arabinose (D). Symbols: glucose (●); xylose (○); arabinose (■); ethanol calculated from cumulative CO_2 production (□); ethanol measured by HPLC (▲); cumulative CO_2 production (Δ); xylitol (▼)

[0011] FIG. 6. Sugar consumption and product formation during an anaerobic batch fermentation of strain IMS0002 cells selected for anaerobic growth on xylose. The fermentation was performed in 1 liter of synthetic medium supplemented with 20 g l^{-1} xylose and 20 g l^{-1} arabinose. Symbols: xylose (○); arabinose (■); ethanol measured by HPLC (▲); cumulative CO_2 production (Δ); xylitol (▼).

[0012] FIG. 7. Sugar consumption and product formation during an anaerobic batch fermentation of strain IMS0003. The fermentation was performed in 1 liter of synthetic medium supplemented with: 30 g l^{-1} glucose, 15 g l^{-1} xylose, and 15 g l^{-1} arabinose. Symbols: glucose (●); xylose (○); arabinose (■); ethanol calculated from cumulative CO_2 production (□); ethanol measured by HPLC (▲); cumulative CO_2 production (Δ);

DESCRIPTION OF THE INVENTION

Eukaryotic Cell

[0013] In a first aspect, the invention relates to a eukaryotic cell capable of expressing the following nucleotide sequences, whereby the expression of these nucleotide sequences confers on the cell the ability to use L-arabinose and/or to convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product such as ethanol:

[0014] (a) a nucleotide sequence encoding an arabinose isomerase (araA), wherein said nucleotide sequence is selected from the group consisting of:

[0015] (i) nucleotide sequences encoding an araA, said araA comprising an amino acid sequence that has at least 55% sequence identity with the amino acid sequence of SEQ ID NO:1.

[0016] (ii) nucleotide sequences comprising a nucleotide sequence that has at least 60% sequence identity with the nucleotide sequence of SEQ ID NO:2.

[0017] (iii) nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);

[0018] (iv) nucleotide sequences the sequences of which differ from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code,

[0019] (b) a nucleotide sequence encoding a L-ribulokinase (araB), wherein said nucleotide sequence is selected from the group consisting of:

[0020] (i) nucleotide sequences encoding an araB, said araB comprising an amino acid sequence that has

- at least 20% sequence identity with the amino acid sequence of SEQ ID NO:3.
- [0021] (ii) nucleotide sequences comprising a nucleotide sequence that has at least 50% sequence identity with the nucleotide sequence of SEQ ID NO:4.
- [0022] (iii) nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);
- [0023] (iv) nucleotide sequences the sequences of which differ from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code,
- [0024] (c) a nucleotide sequence encoding an L-ribulose-5-P-4-epimerase (araD), wherein said nucleotide sequence is selected from the group consisting of:
- [0025] (i) nucleotide sequences encoding an araD, said araD comprising an amino acid sequence that has at least 60% sequence identity with the amino acid sequence of SEQ ID NO:5.
- [0026] (ii) nucleotide sequences comprising a nucleotide sequence that has at least 60% sequence identity with the nucleotide sequence of SEQ ID NO:6.
- [0027] (iii) nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);
- [0028] (iv) nucleotide sequences the sequences of which differ from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code.
- [0029] A preferred embodiment relates to an eukaryotic cell capable of expressing the following nucleotide sequences, whereby the expression of these nucleotide sequences confers on the cell the ability to use L-arabinose and/or to convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product such as ethanol:
- [0030] (a) a nucleotide sequence encoding an arabinose isomerase (araA), wherein said nucleotide sequence is selected from the group consisting of:
- [0031] (i) nucleotide sequences comprising a nucleotide sequence that has at least 60% sequence identity with the nucleotide sequence of SEQ ID NO:2,
- [0032] (ii) nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i);
- [0033] (iii) nucleotide sequences the sequences of which differ from the sequence of a nucleic acid molecule of (ii) due to the degeneracy of the genetic code,
- [0034] (b) a nucleotide sequence encoding a L-ribulokinase (araB), wherein said nucleotide sequence is selected from the group consisting of:
- [0035] (i) nucleotide sequences encoding an araB, said araB comprising an amino acid sequence that has at least 20% sequence identity with the amino acid sequence of SEQ ID NO:3.
- [0036] (ii) nucleotide sequences comprising a nucleotide sequence that has at least 50% sequence identity with the nucleotide sequence of SEQ ID NO:4.
- [0037] (iii) nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);
- [0038] (iv) nucleotide sequences the sequences of which differ from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code,

- [0039] (c) a nucleotide sequence encoding an L-ribulose-5-P-4-epimerase (araD), wherein said nucleotide sequence is selected from the group consisting of:
- [0040] (i) nucleotide sequences encoding an araD, said araD comprising an amino acid sequence that has at least 60% sequence identity with the amino acid sequence of SEQ ID NO:5.
- [0041] (ii) nucleotide sequences comprising a nucleotide sequence that has at least 60% sequence identity with the nucleotide sequence of SEQ ID NO:6.
- [0042] (iii) nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);
- [0043] (iv) nucleotide sequences the sequences of which differ from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code.

Sequence Identity and Similarity

[0044] Sequence identity is herein defined as a relationship between two or more amino acid (polypeptide or protein) sequences or two or more nucleic acid (polynucleotide) sequences, as determined by comparing the sequences. Usually, sequence identities or similarities are compared over the whole length of the sequences compared. In the art, "identity" also means the degree of sequence relatedness between amino acid or nucleic acid sequences, as the case may be, as determined by the match between strings of such sequences. "Similarity" between two amino acid sequences is determined by comparing the amino acid sequence and its conserved amino acid substitutes of one polypeptide to the sequence of a second polypeptide. "Identity" and "similarity" can be readily calculated by various methods, known to those skilled in the art.

[0045] Preferred methods to determine identity are designed to give the largest match between the sequences tested. Methods to determine identity and similarity are codified in publicly available computer programs. Preferred computer program methods to determine identity and similarity between two sequences include e.g. the BestFit, BLASTP, BLASTN, and FASTA (Altschul, S. F. et al., *J. Mol. Biol.* 215:403-410 (1990), publicly available from NCBI and other sources (BLAST Manual, Altschul, S., et al., NCBI NLM NIH Bethesda, Md. 20894). A most preferred algorithm used is EMBOSS (<http://www.ebi.ac.uk/emboss/align>). Preferred parameters for amino acid sequences comparison using EMBOSS are gap open 10.0, gap extend 0.5, Blosum 62 matrix. Preferred parameters for nucleic acid sequences comparison using EMBOSS are gap open 10.0, gap extend 0.5, DNA full matrix (DNA identity matrix).

[0046] Optionally, in determining the degree of amino acid similarity, the skilled person may also take into account so-called "conservative" amino acid substitutions, as will be clear to the skilled person. Conservative amino acid substitutions refer to the interchangeability of residues having similar side chains. For example, a group of amino acids having aliphatic side chains is glycine, alanine, valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl side chains is serine and threonine; a group of amino acids having amide-containing side chains is asparagine and glutamine; a group of amino acids having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group of amino acids having basic side chains is lysine, arginine, and histidine; and a group of amino acids having sulphur-containing side chains is cysteine and methionine. Preferred conser-

vative amino acids substitution groups are: valine-leucine-isoleucine, phenylalanine-tyrosine, lysine-arginine, alanine-valine, and asparagine-glutamine. Substitutional variants of the amino acid sequence disclosed herein are those in which at least one residue in the disclosed sequences has been removed and a different residue inserted in its place. Preferably, the amino acid change is conservative. Preferred conservative substitutions for each of the naturally occurring amino acids are as follows: Ala to ser; Arg to lys; Asn to gln or his; Asp to glu; Cys to ser or ala; Gln to asn; Glu to asp; Gly to pro; His to asn or gln; Ile to leu or val; Leu to ile or val; Lys to arg; gin or glu; Met to leu or ile; Phe to met, leu or tyr; Ser to thr; Thr to ser; Trp to tyr; Tyr to trp or phe; and, Val to ile or leu.

Hybridising Nucleic Acid Sequences

[0047] Nucleotide sequences encoding the enzymes expressed in the cell of the invention may also be defined by their capability to hybridise with the nucleotide sequences of SEQ ID NO.'s 2, 4, 6, 8, 16, 18, 20, 22, 24, 26, 28, 30 respectively, under moderate, or preferably under stringent hybridisation conditions. Stringent hybridisation conditions are herein defined as conditions that allow a nucleic acid sequence of at least about 25, preferably about 50 nucleotides, 75 or 100 and most preferably of about 200 or more nucleotides, to hybridise at a temperature of about 65° C. in a solution comprising about 1 M salt, preferably 6×SSC or any other solution having a comparable ionic strength, and washing at 65° C. in a solution comprising about 0.1 M salt, or less, preferably 0.2×SSC or any other solution having a comparable ionic strength. Preferably, the hybridisation is performed overnight, i.e. at least for 10 hours and preferably washing is performed for at least one hour with at least two changes of the washing solution. These conditions will usually allow the specific hybridisation of sequences having about 90% or more sequence identity.

[0048] Moderate conditions are herein defined as conditions that allow a nucleic acid sequences of at least 50 nucleotides, preferably of about 200 or more nucleotides, to hybridise at a temperature of about 45° C. in a solution comprising about 1 M salt, preferably 6×SSC or any other solution having a comparable ionic strength, and washing at room temperature in a solution comprising about 1 M salt, preferably 6×SSC or any other solution having a comparable ionic strength. Preferably, the hybridisation is performed overnight, i.e. at least for 10 hours, and preferably washing is performed for at least one hour with at least two changes of the washing solution. These conditions will usually allow the specific hybridisation of sequences having up to 50% sequence identity. The person skilled in the art will be able to modify these hybridisation conditions in order to specifically identify sequences varying in identity between 50% and 90%.

AraA

[0049] A preferred nucleotide sequence encoding a arabinose isomerase (araA) expressed in the cell of the invention is selected from the group consisting of:

[0050] (a) nucleotide sequences encoding an araA polypeptide said araA comprising an amino acid sequence that has at least 55, 60, 65, 70, 75, 80, 85, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 1;

[0051] (b) nucleotide sequences comprising a nucleotide sequence that has at least 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO. 2;

[0052] (c) nucleotide sequences the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);

[0053] (d) nucleotide sequences the sequence of which differ from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

The nucleotide sequence encoding an araA may encode either a prokaryotic or an eukaryotic araA, i.e. an araA with an amino acid sequence that is identical to that of an araA that naturally occurs in the prokaryotic or eukaryotic organism. The present inventors have found that the ability of a particular araA to confer to a eukaryotic host cell the ability to use arabinose and/or to convert arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product such as ethanol when co-expressed with araB and araD does not depend so much on whether the araA is of prokaryotic or eukaryotic origin. Rather this depends on the relatedness of the araA's amino acid sequence to that of the sequence SEQ ID NO. 1.

AraB

[0054] A preferred nucleotide sequence encoding a L-ribulokinase (AraB) expressed in the cell of the invention is selected from the group consisting of:

[0055] (a) nucleotide sequences encoding a polypeptide comprising an amino acid sequence that has at least 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 3;

[0056] (b) nucleotide sequences comprising a nucleotide sequence that has at least 50, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO.4;

[0057] (c) nucleotide sequences the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);

[0058] (d) nucleotide sequences the sequence of which differ from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

The nucleotide sequence encoding an araB may encode either a prokaryotic or an eukaryotic araB, i.e. an araB with an amino acid sequence that is identical to that of a araB that naturally occurs in the prokaryotic or eukaryotic organism. The present inventors have found that the ability of a particular araB to confer to a eukaryotic host cell the ability to use arabinose and/or to convert arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product when co-expressed with araA and araD does not depend so much on whether the araB is of prokaryotic or eukaryotic origin. Rather this depends on the relatedness of the araB's amino acid sequence to that of the sequence SEQ ID NO. 3.

AraD

[0059] A preferred nucleotide sequence encoding a L-ribulose-5-P-4-epimerase (araD) expressed in the cell of the invention is selected from the group consisting of:

[0060] (e) nucleotide sequences encoding a polypeptide comprising an amino acid sequence that has at least 60,

65, 70, 75, 80, 85, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 5;

[0061] (f) nucleotide sequences comprising a nucleotide sequence that has at least 60, 65, 70, 75, 80, 85, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO.6;

[0062] (g) nucleotide sequences the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);

[0063] (h) nucleotide sequences the sequence of which differs from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

The nucleotide sequence encoding an araD may encode either a prokaryotic or an eukaryotic araD, i.e. an araD with an amino acid sequence that is identical to that of a araD that naturally occurs in the prokaryotic or eukaryotic organism. The present inventors have found that the ability of a particular araD to confer to a eukaryotic host cell the ability to use arabinose and/or to convert arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product when co-expressed with araA and araB does not depend so much on whether the araD is of prokaryotic or eukaryotic origin. Rather this depends on the relatedness of the araD's amino acid sequence to that of the sequence SEQ ID NO. 5.

[0064] Surprisingly, the codon bias index indicated that expression of the *Lactobacillus plantarum* araA, araB and araD genes were more favorable for expression in yeast than the prokaryotic araA, araB and araD genes described in EP 1 499 708.

[0065] It is to be noted that *L. plantarum* is a Generally Regarded As Safe (GRAS) organism, which is recognized as safe by food registration authorities. Therefore, a preferred nucleotide sequence encodes an araA, araB or araD respectively having an amino acid sequence that is related to the sequences SEQ ID NO: 1, 3, or 5 respectively as defined above. A preferred nucleotide sequence encodes a fungal araA, araB or araD respectively (e.g. from a Basidiomycete), more preferably an araA, araB or araD respectively from an anaerobic fungus, e.g. an anaerobic fungus that belongs to the families Neocallimastix, Caecomyces, Pfomyces, Orpino-mycetes, or Ruminomyces. Alternatively, a preferred nucleotide sequence encodes a bacterial araA, araB or araD respectively, preferably from a Gram-positive bacterium, more preferably from the genus *Lactobacillus*, most preferably from *Lactobacillus plantarum* species. Preferably, one, two or three or the araA, araB and araD nucleotide sequences originate from a *Lactobacillus* genus, more preferably a *Lactobacillus plantarum* species. The bacterial araA expressed in the cell of the invention is not the *Bacillus subtilis* araA disclosed in EP 1 499 708 and given as SEQ ID NO:9. SEQ ID NO:10 represents the nucleotide acid sequence coding for SEQ ID NO:9. The bacterial araB and araD expressed in the cell of the invention are not the ones of *Escherichia coli* (*E. coli*) as disclosed in EP 1 499 708 and given as SEQ ID NO: 11 and SEQ ID NO:13. SEQ ID NO: 12 represents the nucleotide acid sequence coding for SEQ ID NO:11. SEQ ID NO:14 represents the nucleotide acid sequence coding for SEQ ID NO:13.

[0066] To increase the likelihood that the (bacterial) araA, araB and araD enzymes respectively are expressed in active form in a eukaryotic host cell of the invention such as yeast, the corresponding encoding nucleotide sequence may be adapted to optimise its codon usage to that of the chosen

eukaryotic host cell. The adaptiveness of a nucleotide sequence encoding the araA, araB, and araD enzymes (or other enzymes of the invention, see below) to the codon usage of the chosen host cell may be expressed as codon adaptation index (CAI). The codon adaptation index is herein defined as a measurement of the relative adaptiveness of the codon usage of a gene towards the codon usage of highly expressed genes. The relative adaptiveness (w) of each codon is the ratio of the usage of each codon, to that of the most abundant codon for the same amino acid. The CAI index is defined as the geometric mean of these relative adaptiveness values. Non-synonymous codons and termination codons (dependent on genetic code) are excluded. CAI values range from 0 to 1, with higher values indicating a higher proportion of the most abundant codons (see Sharp and Li, 1987, *Nucleic Acids Research* 15: 1281-1295; also see: Jansen et al., 2003, *Nucleic Acids Res.* 31(8):2242-51). An adapted nucleotide sequence preferably has a CAI of at least 0.2, 0.3, 0.4, 0.5, 0.6 or 0.7.

[0067] In a preferred embodiment, expression of the nucleotide sequences encoding an ara A, an ara B and an ara D as defined earlier herein confers to the cell the ability to use L-arabinose and/or to convert it into L-ribulose, and/or xylulose 5-phosphate. Without wishing to be bound by any theory, L-arabinose is expected to be first converted into L-ribulose, which is subsequently converted into xylulose 5-phosphate which is the main molecule entering the pentose phosphate pathway. In the context of the invention, "using L-arabinose" preferably means that the optical density measured at 660 nm (OD₆₆₀) of transformed cells cultured under aerobic or anaerobic conditions in the presence of at least 0.5% L-arabinose during at least 20 days is increased from approximately 0.5 till 1.0 or more. More preferably, the OD₆₆₀ is increased from 0.5 till 1.5 or more. More preferably, the cells are cultured in the presence of at least 1%, at least 1.5%, at least 2% L-arabinose. Most preferably, the cells are cultured in the presence of approximately 2% L-arabinose.

[0068] In the context of the invention, a cell is able "to convert L-arabinose into L-ribulose" when detectable amounts of L-ribulose are detected in cells cultured under aerobic or anaerobic conditions in the presence of L-arabinose (same preferred concentrations as in previous paragraph) during at least 20 days using a suitable assay. Preferably the assay is HPLC for L-ribulose.

[0069] In the context of the invention, a cell is able "to convert L-arabinose into xylulose 5-phosphate" when an increase of at least 2% of xylulose 5-phosphate is detected in cells cultured under aerobic or anaerobic conditions in the presence of L-arabinose (same preferred concentrations as in previous paragraph) during at least 20 days using a suitable assay. Preferably, an HPCL-based assay for xylulose 5-phosphate has been described in Zaldivar J., et al ((2002), *Appl. Microbiol. Biotechnol.*, 59:436-442). This assay is briefly described in the experimental part. More preferably, the increase is of at least 5%, 10%, 15%, 20%, 25% or more.

[0070] In another preferred embodiment, expression of the nucleotide sequences encoding an ara A, ara B and ara D as defined earlier herein confers to the cell the ability to convert L-arabinose into a desired fermentation product when cultured under aerobic or anaerobic conditions in the presence of L-arabinose (same preferred concentrations as in previous paragraph) during at least one month till one year. More preferably, a cell is able to convert L-arabinose into a desired fermentation product when detectable amounts of a desired

fermentation product are detected using a suitable assay and when the cells are cultured under the conditions given in previous sentence. Even more preferably, the assay is HPLC. Even more preferably, the fermentation product is ethanol.

[0071] A cell for transformation with the nucleotide sequences encoding the araA, araB, and araD enzymes respectively as described above, preferably is a host cell capable of active or passive xylose transport into and xylose isomerisation within the cell. The cell preferably is capable of active glycolysis. The cell may further contain an endogenous pentose phosphate pathway and may contain endogenous xylulose kinase activity so that xylulose isomerised from xylose may be metabolised to pyruvate. The cell further preferably contains enzymes for conversion of pyruvate to a desired fermentation product such as ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic or a cephalosporin. The cell may be made capable of producing butanol by introduction of one or more genes of the butanol pathway as disclosed in WO2007/041269.

[0072] A preferred cell is naturally capable of alcoholic fermentation, preferably, anaerobic alcoholic fermentation. The host cell further preferably has a high tolerance to ethanol, a high tolerance to low pH (i.e. capable of growth at a pH lower than 5, 4, 3, or 2.5) and towards organic acids like lactic acid, acetic acid or formic acid and sugar degradation products such as furfural and hydroxy-methylfurfural, and a high tolerance to elevated temperatures. Any of these characteristics or activities of the host cell may be naturally present in the host cell or may be introduced or modified through genetic selection or by genetic modification. A suitable host cell is a eukaryotic microorganism like e.g. a fungus, however, most suitable as host cell are yeasts or filamentous fungi.

[0073] Yeasts are herein defined as eukaryotic microorganisms and include all species of the subdivision Eumycotina (Alexopoulos, C. J., 1962, In: Introductory Mycology, John Wiley & Sons, Inc., New York) that predominantly grow in unicellular form. Yeasts may either grow by budding of a unicellular thallus or may grow by fission of the organism. Preferred yeasts as host cells belong to one of the genera *Saccharomyces*, *Kluyveromyces*, *Candida*, *Pichia*, *Schizosaccharomyces*, *Hansenula*, *Kloeckera*, *Schwanniomycetes*, or *Yarrowia*. Preferably the yeast is capable of anaerobic fermentation, more preferably anaerobic alcoholic fermentation.

[0074] Filamentous fungi are herein defined as eukaryotic microorganisms that include all filamentous forms of the subdivision Eumycotina. These fungi are characterized by a vegetative mycelium composed of chitin, cellulose, and other complex polysaccharides. The filamentous fungi of the present invention are morphologically, physiologically, and genetically distinct from yeasts. Vegetative growth by filamentous fungi is by hyphal elongation and carbon catabolism of most filamentous fungi is obligately aerobic. Preferred filamentous fungi as host cells belong to one of the genera *Aspergillus*, *Trichoderma*, *Humicola*, *Acremonium*, *Fusarium*, or *Penicillium*.

[0075] Over the years suggestions have been made for the introduction of various organisms for the production of bio-ethanol from crop sugars. In practice, however, all major bio-ethanol production processes have continued to use the yeasts of the genus *Saccharomyces* as ethanol producer. This is due to the many attractive features of *Saccharomyces* spe-

cies for industrial processes, i.e., a high acid-, ethanol- and osmo-tolerance, capability of anaerobic growth, and of course its high alcoholic fermentative capacity. Preferred yeast species as host cells include *S. cerevisiae*, *S. bulderi*, *S. barnetti*, *S. exiguus*, *S. uvarum*, *S. diastaticus*, *K. lactis*, *K. marxianus*, *K. fragilis*.

[0076] In a preferred embodiment, the host cell of the invention is a host cell that has been transformed with a nucleic acid construct comprising the nucleotide sequence encoding the araA, araB, and araD enzymes as defined above. In one more preferred embodiment, the host cell is co-transformed with three nucleic acid constructs, each nucleic acid construct comprising the nucleotide sequence encoding araA, araB or araD. The nucleic acid construct comprising the araA, araB, and/or araD coding sequence is capable of expression of the araA, araB, and/or araD enzymes in the host cell. To this end the nucleic acid construct may be constructed as described in e.g. WO 03/0624430. The host cell may comprise a single but preferably comprises multiple copies of each nucleic acid construct. The nucleic acid construct may be maintained episomally and thus comprise a sequence for autonomous replication, such as an ARS sequence. Suitable episomal nucleic acid constructs may e.g. be based on the yeast 2 μ or pKD1 (Fleer et al., 1991, Biotechnology 9:968-975) plasmids. Preferably, however, each nucleic acid construct is integrated in one or more copies into the genome of the host cell. Integration into the host cell's genome may occur at random by illegitimate recombination but preferably nucleic acid construct is integrated into the host cell's genome by homologous recombination as is well known in the art of fungal molecular genetics (see e.g. WO 90/14423, EP-A-0 481 008, EP-A-0 635 574 and U.S. Pat. No. 6,265,186). Accordingly, in a more preferred embodiment, the cell of the invention comprises a nucleic acid construct comprising the araA, araB, and/or araD coding sequence and is capable of expression of the araA, araB, and/or araD enzymes. In an even more preferred embodiment, the araA, araB, and/or araD coding sequences are each operably linked to a promoter that causes sufficient expression of the corresponding nucleotide sequences in a cell to confer to the cell the ability to use L-arabinose, and/or to convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate. Preferably the cell is a yeast cell. Accordingly, in a further aspect, the invention also encompasses a nucleic acid construct as earlier outlined herein. Preferably, a nucleic acid construct comprises a nucleic acid sequence encoding an araA, araB and/or araD. Nucleic acid sequences encoding an araA, araB, or araD have been all earlier defined herein. Even more preferably, the expression of the corresponding nucleotide sequences in a cell confer to the cell the ability to convert L-arabinose into a desired fermentation product as defined later herein. In an even more preferred embodiment, the fermentation product is ethanol. Even more preferably, the cell is a yeast cell.

[0077] As used herein, the term "operably linked" refers to a linkage of polynucleotide elements (or coding sequences or nucleic acid sequence) in a functional relationship. A nucleic acid sequence is "operably linked" when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the coding sequence. Operably linked means that the nucleic acid sequences being linked are typically contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame.

[0078] As used herein, the term “promoter” refers to a nucleic acid fragment that functions to control the transcription of one or more genes, located upstream with respect to the direction of transcription of the transcription initiation site of the gene, and is structurally identified by the presence of a binding site for DNA-dependent RNA polymerase, transcription initiation sites and any other DNA sequences, including, but not limited to transcription factor binding sites, repressor and activator protein binding sites, and any other sequences of nucleotides known to one of skill in the art to act directly or indirectly to regulate the amount of transcription from the promoter. A “constitutive” promoter is a promoter that is active under most environmental and developmental conditions. An “inducible” promoter is a promoter that is active under environmental or developmental regulation.

[0079] The promoter that could be used to achieve the expression of the nucleotide sequences coding for araA, araB and/or araD may be not native to the nucleotide sequence coding for the enzyme to be expressed, i.e. a promoter that is heterologous to the nucleotide sequence (coding sequence) to which it is operably linked. Although the promoter preferably is heterologous to the coding sequence to which it is operably linked, it is also preferred that the promoter is homologous, i.e. endogenous to the host cell. Preferably the heterologous promoter (to the nucleotide sequence) is capable of producing a higher steady state level of the transcript comprising the coding sequence (or is capable of producing more transcript molecules, i.e. mRNA molecules, per unit of time) than is the promoter that is native to the coding sequence, preferably under conditions where arabinose, or arabinose and glucose, or xylose and arabinose or xylose and arabinose and glucose are available as carbon sources, more preferably as major carbon sources (i.e. more than 50% of the available carbon source consists of arabinose, or arabinose and glucose, or xylose and arabinose or xylose and arabinose and glucose), most preferably as sole carbon sources. Suitable promoters in this context include both constitutive and inducible natural promoters as well as engineered promoters. A preferred promoter for use in the present invention will in addition be insensitive to catabolite (glucose) repression and/or will preferably not require arabinose and/or xylose for induction.

[0080] Promoters having these characteristics are widely available and known to the skilled person. Suitable examples of such promoters include e.g. promoters from glycolytic genes, such as the phosphofructokinase (PPK), triose phosphate isomerase (TPI), glyceraldehyde-3-phosphate dehydrogenase (GPD, TDH3 or GAPDH), pyruvate kinase (PYK), phosphoglycerate kinase (PGK) promoters from yeasts or filamentous fungi; more details about such promoters from yeast may be found in (WO 93/03159). Other useful promoters are ribosomal protein encoding gene promoters, the lactase gene promoter (LAC4), alcohol dehydrogenase promoters (ADH1, ADH4, and the like), the enolase promoter (ENO), the glucose-6-phosphate isomerase promoter (PGI1, Hauf et al, 2000) or the hexose(glucose) transporter promoter (HXT7) or the glyceraldehyde-3-phosphate dehydrogenase (TDH3). The sequence of the PGI1 promoter is given in SEQ ID NO:51. The sequence of the HXT7 promoter is given in SEQ ID NO:52. The sequence of the TDH3 promoter is given in SEQ ID NO:49. Other promoters, both constitutive and inducible, and enhancers or upstream activating sequences will be known to those of skill in the art. The promoters used in the host cells of the invention may be modified, if desired, to affect their control characteristics. A preferred cell of the

invention is a eukaryotic cell transformed with the araA, araB and araD genes of *L. plantarum*. More preferably, the eukaryotic cell is a yeast cell, even more preferably a *S. cerevisiae* strain transformed with the araA, araB and araD genes of *L. plantarum*. Most preferably, the cell is either CBS120327 or CBS120328 both deposited at the CBS Institute (The Netherlands) on Sep. 27, 2006.

[0081] The term “homologous” when used to indicate the relation between a given (recombinant) nucleic acid or polypeptide molecule and a given host organism or host cell, is understood to mean that in nature the nucleic acid or polypeptide molecule is produced by a host cell or organisms of the same species, preferably of the same variety or strain. If homologous to a host cell, a nucleic acid sequence encoding a polypeptide will typically be operably linked to another promoter sequence or, if applicable, another secretory signal sequence and/or terminator sequence than in its natural environment. When used to indicate the relatedness of two nucleic acid sequences the term “homologous” means that one single-stranded nucleic acid sequence may hybridize to a complementary single-stranded nucleic acid sequence. The degree of hybridization may depend on a number of factors including the amount of identity between the sequences and the hybridization conditions such as temperature and salt concentration as earlier presented. Preferably the region of identity is greater than about 5 bp, more preferably the region of identity is greater than 10 bp.

[0082] The term “heterologous” when used with respect to a nucleic acid (DNA or RNA) or protein refers to a nucleic acid or protein that does not occur naturally as part of the organism, cell, genome or DNA or RNA sequence in which it is present, or that is found in a cell or location or locations in the genome or DNA or RNA sequence that differ from that in which it is found in nature. Heterologous nucleic acids or proteins are not endogenous to the cell into which it is introduced, but has been obtained from another cell or synthetically or recombinantly produced. Generally, though not necessarily, such nucleic acids encode proteins that are not normally produced by the cell in which the DNA is transcribed or expressed. Similarly exogenous RNA encodes for proteins not normally expressed in the cell in which the exogenous RNA is present. Heterologous nucleic acids and proteins may also be referred to as foreign nucleic acids or proteins. Any nucleic acid or protein that one of skill in the art would recognize as heterologous or foreign to the cell in which it is expressed is herein encompassed by the term heterologous nucleic acid or protein. The term heterologous also applies to non-natural combinations of nucleic acid or amino acid sequences, i.e. combinations where at least two of the combined sequences are foreign with respect to each other.

Preferred Eukaryotic Cell Able to Use and/or Convert L-Arabinose and Xylose

[0083] In a more preferred embodiment, the cell of the invention that expresses araA, araB and araD is able to use L-arabinose and/or to convert it into L-ribulose, and/or xylulose 5-phosphate and/or a desired fermentation product as earlier defined herein and additionally exhibits the ability to use xylose and/or convert xylose into xylulose. The conversion of xylose into xylulose is preferably a one step isomerisation step (direct isomerisation of xylose into xylulose). This type of cell is therefore able to use both L-arabinose and xylose. “Using” xylose has preferably the same meaning as “using” L-arabinose as earlier defined herein.

[0084] Enzyme definitions are as used in WO 06/009434, for xylose isomerase (EC 5.3.1.5), xylulose kinase (EC 2.7.1.17), ribulose 5-phosphate epimerase (5.1.3.1), ribulose 5-phosphate isomerase (EC 5.3.1.6), transketolase (EC 2.2.1.1), transaldolase (EC 2.2.1.2), and aldose reductase" (EC 1.1.1.21).

[0085] In a preferred embodiment, the eukaryotic cell of the invention expressing araA, araB and araD as earlier defined herein has the ability of isomerising xylose to xylulose as e.g. described in WO 03/0624430 or in WO 06/009434. The ability of isomerising xylose to xylulose is conferred to the host cell by transformation of the host cell with a nucleic acid construct comprising a nucleotide sequence encoding a xylose isomerase. The transformed host cell's ability to isomerise xylose into xylulose is the direct isomerisation of xylose to xylulose. This is understood to mean that xylose isomerised into xylulose in a single reaction catalysed by a xylose isomerase, as opposed to the two step conversion of xylose into xylulose via a xylitol intermediate as catalysed by xylose reductase and xylitol dehydrogenase, respectively.

[0086] The nucleotide sequence encodes a xylose isomerase that is preferably expressed in active form in the transformed host cell of the invention. Thus, expression of the nucleotide sequence in the host cell produces a xylose isomerase with a specific activity of at least 10 U xylose isomerase activity per mg protein at 30° C., preferably at least 20, 25, 30, 50, 100, 200, 300 or 500 U per mg at 30° C. The specific activity of the xylose isomerase expressed in the transformed host cell is herein defined as the amount of xylose isomerase activity units per mg protein of cell free lysate of the host cell, e.g. a yeast cell free lysate. Determination of the xylose isomerase activity has already been described earlier herein.

[0087] Preferably, expression of the nucleotide sequence encoding the xylose isomerase in the host cell produces a xylose isomerase with a K_m for xylose that is less than 50, 40, 30 or 25 mM, more preferably, the K_m for xylose is about 20 mM or less.

[0088] A preferred nucleotide sequence encoding the xylose isomerase may be selected from the group consisting of:

[0089] (e) nucleotide sequences encoding a polypeptide comprising an amino acid sequence that has at least 60, 65, 70, 75, 80, 85, 90, 95, 97, 98, or 99% sequence identity with the amino acid sequence of SEQ ID NO. 7 or SEQ ID NO:15;

[0090] (f) nucleotide sequences comprising a nucleotide sequence that has at least 40, 50, 60, 70, 80, 90, 95, 97, 98, or 99% sequence identity with the nucleotide sequence of SEQ ID NO. 8 or SEQ ID NO:16;

[0091] (g) nucleotide sequences the complementary strand of which hybridises to a nucleic acid molecule sequence of (a) or (b);

[0092] (h) nucleotide sequences the sequence of which differs from the sequence of a nucleic acid molecule of (c) due to the degeneracy of the genetic code.

[0093] The nucleotide sequence encoding the xylose isomerase may encode either a prokaryotic or an eukaryotic xylose isomerase, i.e. a xylose isomerase with an amino acid sequence that is identical to that of a xylose isomerase that naturally occurs in the prokaryotic or eukaryotic organism. The present inventors have found that the ability of a particular xylose isomerase to confer to a eukaryotic host cell the ability to isomerise xylose into xylulose does not depend so

much on whether the isomerase is of prokaryotic or eukaryotic origin. Rather this depends on the relatedness of the isomerase's amino acid sequence to that of the *Piromyces* sequence (SEQ ID NO. 7). Surprisingly, the eukaryotic *Piromyces* isomerase is more related to prokaryotic isomerases than to other known eukaryotic isomerases. Therefore, a preferred nucleotide sequence encodes a xylose isomerase having an amino acid sequence that is related to the *Piromyces* sequence as defined above. A preferred nucleotide sequence encodes a fungal xylose isomerase (e.g. from a Basidiomycete), more preferably a xylose isomerase from an anaerobic fungus, e.g. a xylose isomerase from an anaerobic fungus that belongs to the families Neocallimastix, Caecomyces, *Piromyces*, Orpinomyces, or Ruminomyces. Alternatively, a preferred nucleotide sequence encodes a bacterial xylose isomerase, preferably a Gram-negative bacterium, more preferably an isomerase from the class *Bacteroides*, or from the genus *Bacteroides*, most preferably from *B. thetaio-taomicron* (SEQ ID NO. 15).

[0094] To increase the likelihood that the xylose isomerase is expressed in active form in a eukaryotic host cell such as yeast, the nucleotide sequence encoding the xylose isomerase may be adapted to optimise its codon usage to that of the eukaryotic host cell as earlier defined herein.

[0095] A host cell for transformation with the nucleotide sequence encoding the xylose isomerase as described above, preferably is a host capable of active or passive xylose transport into the cell. The host cell preferably contains active glycolysis. The host cell may further contain an endogenous pentose phosphate pathway and may contain endogenous xylulose kinase activity so that xylulose isomerised from xylose may be metabolised to pyruvate. The host further preferably contains enzymes for conversion of pyruvate to a desired fermentation product such as ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic or a cephalosporin. A preferred host cell is a host cell that is naturally capable of alcoholic fermentation, preferably, anaerobic alcoholic fermentation. The host cell further preferably has a high tolerance to ethanol, a high tolerance to low pH (i.e. capable of growth at a pH lower than 5, 4, 3, or 2.5) and towards organic acids like lactic acid, acetic acid or formic acid and sugar degradation products such as furfural and hydroxy-methylfurfural, and a high tolerance to elevated temperatures. Any of these characteristics or activities of the host cell may be naturally present in the host cell or may be introduced or modified by genetic modification. A suitable cell is a eukaryotic microorganism like e.g. a fungus, however, most suitable as host cell are yeasts or filamentous fungi. Preferred yeasts and filamentous fungi have already been defined herein.

[0096] As used herein the wording host cell has the same meaning as cell.

[0097] The cell of the invention is preferably transformed with a nucleic acid construct comprising the nucleotide sequence encoding the xylose isomerase. The nucleic acid construct that is preferably used is the same as the one used comprising the nucleotide sequence encoding araA, araB or araD.

[0098] In another preferred embodiment of the invention, the cell of the invention:

[0099] expressing *araA*, *araB* and *araD*, and exhibiting the ability to directly isomerise xylose into xylulose, as earlier defined herein

further comprises a genetic modification that increases the flux of the pentose phosphate pathway, as described in WO 06/009434. In particular, the genetic modification causes an increased flux of the non-oxidative part pentose phosphate pathway. A genetic modification that causes an increased flux of the non-oxidative part of the pentose phosphate pathway is herein understood to mean a modification that increases the flux by at least a factor 1.1, 1.2, 1.5, 2, 5, 10 or 20 as compared to the flux in a strain which is genetically identical except for the genetic modification causing the increased flux. The flux of the non-oxidative part of the pentose phosphate pathway may be measured by growing the modified host on xylose as sole carbon source, determining the specific xylose consumption rate and subtracting the specific xylitol production rate from the specific xylose consumption rate, if any xylitol is produced. However, the flux of the non-oxidative part of the pentose phosphate pathway is proportional with the growth rate on xylose as sole carbon source, preferably with the anaerobic growth rate on xylose as sole carbon source. There is a linear relation between the growth rate on xylose as sole carbon source (μ_{max}) and the flux of the non-oxidative part of the pentose phosphate pathway. The specific xylose consumption rate (Q_s) is equal to the growth rate (μ) divided by the yield of biomass on sugar (Y_{xs}) because the yield of biomass on sugar is constant (under a given set of Conditions: anaerobic, growth medium, pH, genetic background of the strain, etc.; i.e. $Q_s = \mu/Y_{xs}$). Therefore the increased flux of the non-oxidative part of the pentose phosphate pathway may be deduced from the increase in maximum growth rate under these conditions. In a preferred embodiment, the cell comprises a genetic modification that increases the flux of the pentose phosphate pathway and has a specific xylose consumption rate of at least 346 mg xylose/g biomass/h.

[0100] Genetic modifications that increase the flux of the pentose phosphate pathway may be introduced in the host cell in various ways. These including e.g. achieving higher steady state activity levels of xylulose kinase and/or one or more of the enzymes of the non-oxidative part pentose phosphate pathway and/or a reduced steady state level of unspecific aldose reductase activity. These changes in steady state activity levels may be effected by selection of mutants (spontaneous or induced by chemicals or radiation) and/or by recombinant DNA technology e.g. by overexpression or inactivation, respectively, of genes encoding the enzymes or factors regulating these genes.

[0101] In a more preferred host cell, the genetic modification comprises overexpression of at least one enzyme of the (non-oxidative part) pentose phosphate pathway. Preferably the enzyme is selected from the group consisting of the enzymes encoding for ribulose-5-phosphate isomerase, ribulose-5-phosphate epimerase, transketolase and transaldolase, as described in WO 06/009434.

[0102] Various combinations of enzymes of the (non-oxidative part) pentose phosphate pathway may be overexpressed. E.g. the enzymes that are overexpressed may be at least the enzymes ribulose-5-phosphate isomerase and ribulose-5-phosphate epimerase; or at least the enzymes ribulose-5-phosphate isomerase and transketolase; or at least the enzymes ribulose-5-phosphate isomerase and transaldolase;

or at least the enzymes ribulose-5-phosphate epimerase and transketolase; or at least the enzymes ribulose-5-phosphate epimerase and transaldolase; or at least the enzymes transketolase and transaldolase; or at least the enzymes ribulose-5-phosphate epimerase, transketolase and transaldolase; or at least the enzymes ribulose-5-phosphate isomerase, transketolase and transaldolase; or at least the enzymes ribulose-5-phosphate isomerase, ribulose-5-phosphate epimerase, and transaldolase; or at least the enzymes ribulose-5-phosphate isomerase, ribulose-5-phosphate epimerase, and transketolase. In one embodiment of the invention each of the enzymes ribulose-5-phosphate isomerase, ribulose-5-phosphate epimerase, transketolase and transaldolase are overexpressed in the host cell. More preferred is a host cell in which the genetic modification comprises at least overexpression of both the enzymes transketolase and transaldolase as such a host cell is already capable of anaerobic growth on xylose. In fact, under some conditions we have found that host cells overexpressing only the transketolase and the transaldolase already have the same anaerobic growth rate on xylose as do host cells that overexpress all four of the enzymes, i.e. the ribulose-5-phosphate isomerase, ribulose-5-phosphate epimerase, transketolase and transaldolase. Moreover, host cells overexpressing both of the enzymes ribulose-5-phosphate isomerase and ribulose-5-phosphate epimerase are preferred over host cells overexpressing only the isomerase or only the epimerase as overexpression of only one of these enzymes may produce metabolic imbalances.

[0103] There are various means available in the art for overexpression of enzymes in the cells of the invention. In particular, an enzyme may be overexpressed by increasing the copy number of the gene coding for the enzyme in the host cell, e.g. by integrating additional copies of the gene in the host cell's genome, by expressing the gene from an episomal multicopy expression vector or by introducing a episomal expression vector that comprises multiple copies of the gene.

[0104] Alternatively overexpression of enzymes in the host cells of the invention may be achieved by using a promoter that is not native to the sequence coding for the enzyme to be overexpressed, i.e. a promoter that is heterologous to the coding sequence to which it is operably linked. Suitable promoters to this end have already been defined herein.

[0105] The coding sequence used for overexpression of the enzymes preferably is homologous to the host cell of the invention. However, coding sequences that are heterologous to the host cell of the invention may likewise be applied, as mentioned in WO 06/009434.

[0106] A nucleotide sequence used for overexpression of ribulose-5-phosphate isomerase in the host cell of the invention is a nucleotide sequence encoding a polypeptide with ribulose-5-phosphate isomerase activity, whereby preferably the polypeptide has an amino acid sequence having at least 50, 60, 70, 80, 90 or 95% identity with SEQ ID NO. 17 or whereby the nucleotide sequence is capable of hybridising with the nucleotide sequence of SEQ ID NO. 18, under moderate conditions, preferably under stringent conditions.

[0107] A nucleotide sequence used for overexpression of ribulose-5-phosphate epimerase in the host cell of the invention is a nucleotide sequence encoding a polypeptide with ribulose-5-phosphate epimerase activity, whereby preferably the polypeptide has an amino acid sequence having at least 50, 60, 70, 80, 90 or 95% identity with SEQ ID NO. 19 or whereby the nucleotide sequence is capable of hybridising

with the nucleotide sequence of SEQ ID NO. 20, under moderate conditions, preferably under stringent conditions.

[0108] A nucleotide sequence used for overexpression of transketolase in the host cell of the invention is a nucleotide sequence encoding a polypeptide with transketolase activity, whereby preferably the polypeptide has an amino acid sequence having at least 50, 60, 70, 80, 90 or 95% identity with SEQ ID NO. 21 or whereby the nucleotide sequence is capable of hybridising with the nucleotide sequence of SEQ ID NO. 22, under moderate conditions, preferably under stringent conditions.

[0109] A nucleotide sequence used for overexpression of transaldolase in the host cell of the invention is a nucleotide sequence encoding a polypeptide with transaldolase activity, whereby preferably the polypeptide has an amino acid sequence having at least 50, 60, 70, 80, 90 or 95% identity with SEQ ID NO. 23 or whereby the nucleotide sequence is capable of hybridising with the nucleotide sequence of SEQ ID NO. 24, under moderate conditions, preferably under stringent conditions.

[0110] Overexpression of an enzyme, when referring to the production of the enzyme in a genetically modified host cell, means that the enzyme is produced at a higher level of specific enzymatic activity as compared to the unmodified host cell under identical conditions. Usually this means that the enzymatically active protein (or proteins in case of multi-subunit enzymes) is produced in greater amounts, or rather at a higher steady state level as compared to the unmodified host cell under identical conditions. Similarly this usually means that the mRNA coding for the enzymatically active protein is produced in greater amounts, or again rather at a higher steady state level as compared to the unmodified host cell under identical conditions. Overexpression of an enzyme is thus preferably determined by measuring the level of the enzyme's specific activity in the host cell using appropriate enzyme assays as described herein. Alternatively, overexpression of the enzyme may be determined indirectly by quantifying the specific steady state level of enzyme protein, e.g. using antibodies specific for the enzyme, or by quantifying the specific steady level of the mRNA coding for the enzyme. The latter may particularly be suitable for enzymes of the pentose phosphate pathway for which enzymatic assays are not easily feasible as substrates for the enzymes are not commercially available. Preferably in the host cells of the invention, an enzyme to be overexpressed is overexpressed by at least a factor 1.1, 1.2, 1.5, 2, 5, 10 or 20 as compared to a strain which is genetically identical except for the genetic modification causing the overexpression. It is to be understood that these levels of overexpression may apply to the steady state level of the enzyme's activity, the steady state level of the enzyme's protein as well as to the steady state level of the transcript coding for the enzyme.

[0111] In a further preferred embodiment, the host cell of the invention:

[0112] expressing araA, araB and araD, and exhibiting the ability to directly isomerise xylose into xylulose, and optionally

[0113] comprising a genetic modification that increases the flux of the pentose pathway as earlier defined herein further comprises a genetic modification that increases the specific xylulose kinase activity. Preferably the genetic modification causes overexpression of a xylulose kinase, e.g. by overexpression of a nucleotide sequence encoding a xylulose kinase. The gene encoding the xylulose kinase may be endog-

enous to the host cell or may be a xylulose kinase that is heterologous to the host cell. A nucleotide sequence used for overexpression of xylulose kinase in the host cell of the invention is a nucleotide sequence encoding a polypeptide with xylulose kinase activity, whereby preferably the polypeptide has an amino acid sequence having at least 50, 60, 70, 80, 90 or 95% identity with SEQ ID NO. 25 or whereby the nucleotide sequence is capable of hybridising with the nucleotide sequence of SEQ ID NO. 26, under moderate conditions, preferably under stringent conditions.

[0114] A particularly preferred xylulose kinase is a xylose kinase that is related to the xylulose kinase xylB from *Pirromyces* as mentioned in WO 03/0624430. A more preferred nucleotide sequence for use in overexpression of xylulose kinase in the host cell of the invention is a nucleotide sequence encoding a polypeptide with xylulose kinase activity, whereby preferably the polypeptide has an amino acid sequence having at least 45, 50, 55, 60, 65, 70, 80, 90 or 95% identity with SEQ ID NO. 27 or whereby the nucleotide sequence is capable of hybridising with the nucleotide sequence of SEQ ID NO. 28, under moderate conditions, preferably under stringent conditions.

[0115] In the host cells of the invention, genetic modification that increases the specific xylulose kinase activity may be combined with any of the modifications increasing the flux of the pentose phosphate pathway as described above, but this combination is not essential for the invention. Thus, a host cell of the invention comprising a genetic modification that increases the specific xylulose kinase activity in addition to the expression of the araA, araB and araD enzymes as defined herein is specifically included in the invention. The various means available in the art for achieving and analysing overexpression of a xylulose kinase in the host cells of the invention are the same as described above for enzymes of the pentose phosphate pathway. Preferably in the host cells of the invention, a xylulose kinase to be overexpressed is overexpressed by at least a factor 1.1, 1.2, 1.5, 2, 5, 10 or 20 as compared to a strain which is genetically identical except for the genetic modification causing the overexpression. It is to be understood that these levels of overexpression may apply to the steady state level of the enzyme's activity, the steady state level of the enzyme's protein as well as to the steady state level of the transcript coding for the enzyme.

[0116] In a further preferred embodiment, the host cell of the invention:

[0117] expressing araA, araB and araD, and exhibiting the ability to directly isomerise xylose into xylulose, and optionally

[0118] comprising a genetic modification that increases the flux of the pentose pathway and/or

[0119] further comprising a genetic modification that increases the specific xylulose kinase activity all as earlier defined herein

further comprises a genetic modification that reduces unspecific aldose reductase activity in the host cell. Preferably, unspecific aldose reductase activity is reduced in the host cell by one or more genetic modifications that reduce the expression of or inactivate a gene encoding an unspecific aldose reductase, as described in WO 06/009434. Preferably, the genetic modifications reduce or inactivate the expression of each endogenous copy of a gene encoding an unspecific aldose reductase in the host cell. Host cells may comprise multiple copies of genes encoding unspecific aldose reductases as a result of di-, poly- or aneu-ploidy, and/or the host

cell may contain several different (iso)enzymes with aldose reductase activity that differ in amino acid sequence and that are each encoded by a different gene. Also in such instances preferably the expression of each gene that encodes an unspecific aldose reductase is reduced or inactivated. Preferably, the gene is inactivated by deletion of at least part of the gene or by disruption of the gene, whereby in this context the term gene also includes any non-coding sequence up- or downstream of the coding sequence, the (partial) deletion or inactivation of which results in a reduction of expression of unspecific aldose reductase activity in the host cell. A nucleotide sequence encoding an aldose reductase whose activity is to be reduced in the host cell of the invention is a nucleotide sequence encoding a polypeptide with aldose reductase activity, whereby preferably the polypeptide has an amino acid sequence having at least 50, 60, 70, 80, 90 or 95% identity with SEQ ID NO. 29 or whereby the nucleotide sequence is capable of hybridising with the nucleotide sequence of SEQ ID NO. 30 under moderate conditions, preferably under stringent conditions.

[0120] In the host cells of the invention, the expression of the *araA*, *araB* and *araD* enzymes as defined herein is combined with genetic modification that reduces unspecific aldose reductase activity. The genetic modification leading to the reduction of unspecific aldose reductase activity may be combined with any of the modifications increasing the flux of the pentose phosphate pathway and/or with any of the modifications increasing the specific xylulose kinase activity in the host cells as described above, but these combinations are not essential for the invention. Thus, a host cell expressing *araA*, *araB*, and *araD*, comprising an additional genetic modification that reduces unspecific aldose reductase activity is specifically included in the invention.

[0121] In a preferred embodiment, the host cell is CBS120327 deposited at the CBS Institute (The Netherlands) on Sep. 27, 2006.

[0122] In a further preferred embodiment, the invention relates to modified host cells that are further adapted to L-arabinose (use L-arabinose and/or convert it into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product and optionally xylose utilisation by selection of mutants, either spontaneous or induced (e.g. by radiation or chemicals), for growth on L-arabinose and optionally xylose, preferably on L-arabinose and optionally xylose as sole carbon source, and more preferably under anaerobic conditions. Selection of mutants may be performed by serial passaging of cultures as e.g. described by Kuyper et al. (2004, FEMS Yeast Res. 4: 655-664) and/or by cultivation under selective pressure in a chemostat culture as is described in Example 4 of WO 06/009434. This selection process may be continued as long as necessary. This selection process is preferably carried out during one week till one year. However, the selection process may be carried out for a longer period of time if necessary. During the selection process, the cells are preferably cultured in the presence of approximately 20 g/l L-arabinose and/or approximately 20 g/l xylose. The cell obtained at the end of this selection process is expected to be improved as to its capacities of using L-arabinose and/or xylose, and/or converting L-arabinose into L-ribulose and/or xylulose 5-phosphate and/or a desired fermentation product such as ethanol. In this context "improved cell" may mean that the obtained cell is able to use L-arabinose and/or xylose in a more efficient way than the cell it derives from. For example, the obtained cell is expected to better grow: increase of the

specific growth rate of at least 2% than the cell it derives from under the same conditions. Preferably, the increase is of at least 4%, 6%, 8%, 10%, 15%, 20%, 25% or more. The specific growth rate may be calculated from OD_{660} as known to the skilled person. Therefore, by monitoring the OD_{660} , one can deduce the specific growth rate. In this context "improved cell" may also mean that the obtained cell converts L-arabinose into L-ribulose and/or xylulose 5-phosphate and/or a desired fermentation product such as ethanol in a more efficient way than the cell it derives from. For example, the obtained cell is expected to produce higher amounts of L-ribulose and/or xylulose 5-phosphate and/or a desired fermentation product such as ethanol: increase of at least one of these compounds of at least 2% than the cell it derives from under the same conditions. Preferably, the increase is of at least 4%, 6%, 8%, 10%, 15%, 20%, 25% or more. In this context "improved cell" may also mean that the obtained cell converts xylose into xylulose and/or a desired fermentation product such as ethanol in a more efficient way than the cell it derives from. For example, the obtained cell is expected to produce higher amounts of xylulose and/or a desired fermentation product such as ethanol: increase of at least one of these compounds of at least 2% than the cell it derives from under the same conditions. Preferably, the increase is of at least 4%, 6%, 8%, 10%, 15%, 20%, 25% or more.

[0123] In a preferred host cell of the invention at least one of the genetic modifications described above, including modifications obtained by selection of mutants, confer to the host cell the ability to grow on L-arabinose and optionally xylose as carbon source, preferably as sole carbon source, and preferably under anaerobic conditions. Preferably the modified host cell produce essentially no xylitol, e.g. the xylitol produced is below the detection limit or e.g. less than 5, 2, 1, 0.5, or 0.3% of the carbon consumed on a molar basis.

[0124] Preferably the modified host cell has the ability to grow on L-arabinose and optionally xylose as sole carbon source at a rate of at least 0.001, 0.005, 0.01, 0.03, 0.05, 0.1, 0.2, 0.25 or 0.3 h^{-1} under aerobic conditions, or, if applicable, at a rate of at least 0.001, 0.005, 0.01, 0.03, 0.05, 0.07, 0.08, 0.09, 0.1, 0.12, 0.15 or 0.2 h^{-1} under anaerobic conditions. Preferably the modified host cell has the ability to grow on a mixture of glucose and L-arabinose and optionally xylose (in a 1:1 weight ratio) as sole carbon source at a rate of at least 0.001, 0.005, 0.01, 0.03, 0.05, 0.1, 0.2, 0.25 or 0.3 h^{-1} under aerobic conditions, or, if applicable, at a rate of at least 0.001, 0.005, 0.01, 0.03, 0.05, 0.1, 0.12, 0.15, or 0.2 h^{-1} under anaerobic conditions.

[0125] Preferably, the modified host cell has a specific L-arabinose and optionally xylose consumption rate of at least 346, 350, 400, 500, 600, 650, 700, 750, 800, 900 or 1000 mg/g cells/h. Preferably, the modified host cell has a yield of fermentation product (such as ethanol) on L-arabinose and optionally xylose that is at least 20, 25, 30, 35, 40, 45, 50, 55, 60, 70, 80, 85, 90, 95 or 98% of the host cell's yield of fermentation product (such as ethanol) on glucose. More preferably, the modified host cell's yield of fermentation product (such as ethanol) on L-arabinose and optionally xylose is equal to the host cell's yield of fermentation product (such as ethanol) on glucose. Likewise, the modified host cell's biomass yield on L-arabinose and optionally xylose is preferably at least 55, 60, 70, 80, 85, 90, 95 or 98% of the host cell's biomass yield on glucose. More preferably, the modified host cell's biomass yield on L-arabinose and optionally xylose is equal to the host cell's biomass yield on glucose. It

is understood that in the comparison of yields on glucose and L-arabinose and optionally xylose both yields are compared under aerobic conditions or both under anaerobic conditions.

[0126] In a more preferred embodiment, the host cell is CBS120328 deposited at the CBS Institute (The Netherlands) on Sep. 27, 2006 or CBS121879 deposited at the CBS Institute (The Netherlands) on Sep. 20, 2007.

[0127] In a preferred embodiment, the cell expresses one or more enzymes that confer to the cell the ability to produce at least one fermentation product selected from the group consisting of ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic and a cephalosporin. In a more preferred embodiment, the host cell of the invention is a host cell for the production of ethanol. In another preferred embodiment, the invention relates to a transformed host cell for the production of fermentation products other than ethanol. Such non-ethanolic fermentation products include in principle any bulk or fine chemical that is producible by a eukaryotic microorganism such as a yeast or a filamentous fungus. Such fermentation products include e.g. lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic and a cephalosporin. A preferred host cell of the invention for production of non-ethanolic fermentation products is a host cell that contains a genetic modification that results in decreased alcohol dehydrogenase activity.

Method

[0128] In a further aspect, the invention relates to fermentation processes in which a host cell of the invention is used for the fermentation of a carbon source comprising a source of L-arabinose and optionally a source of xylose. Preferably, the source of L-arabinose and the source of xylose are L-arabinose and xylose. In addition, the carbon source in the fermentation medium may also comprise a source of glucose. The source of L-arabinose, xylose or glucose may be L-arabinose, xylose or glucose as such or may be any carbohydrate oligo- or polymer comprising L-arabinose, xylose or glucose units, such as e.g. lignocellulose, xylans, cellulose, starch, arabinan and the like. For release of xylose or glucose units from such carbohydrates, appropriate carbohydrases (such as xylanases, glucanases, amylases and the like) may be added to the fermentation medium or may be produced by the modified host cell. In the latter case the modified host cell may be genetically engineered to produce and excrete such carbohydrases. An additional advantage of using oligo- or polymeric sources of glucose is that it enables to maintain a low(er) concentration of free glucose during the fermentation, e.g. by using rate-limiting amounts of the carbohydrases. This, in turn, will prevent repression of systems required for metabolism and transport of non-glucose sugars such as xylose. In a preferred process the modified host cell ferments both the L-arabinose (optionally xylose) and glucose, preferably simultaneously in which case preferably a modified host cell is used which is insensitive to glucose repression to prevent diauxic growth. In addition to a source of L-arabinose, optionally xylose (and glucose) as carbon source, the fermentation medium will further comprise the appropriate ingredient required for growth of the modified host cell. Compositions of fermentation media for growth of microorganisms such as yeasts or filamentous fungi are well known in the art.

[0129] In a preferred process, there is provided a process for producing a fermentation product selected from the group consisting of ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic and a cephalosporin whereby the process comprises the steps of:

[0130] (a) fermenting a medium containing a source of L-arabinose and optionally xylose with a modified host cell as defined herein, whereby the host cell ferments L-arabinose and optionally xylose to the fermentation product, and optionally,

[0131] (b) recovering the fermentation product.

[0132] The fermentation process is a process for the production of a fermentation product such as e.g. ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic, such as Penicillin G or Penicillin V and fermentative derivatives thereof, and/or a cephalosporin. The fermentation process may be an aerobic or an anaerobic fermentation process. An anaerobic fermentation process is herein defined as a fermentation process run in the absence of oxygen or in which substantially no oxygen is consumed, preferably less than 5, 2.5 or 1 mmol/L/h, more preferably 0 mmol/L/h is consumed (i.e. oxygen consumption is not detectable), and wherein organic molecules serve as both electron donor and electron acceptors. In the absence of oxygen, NADH produced in glycolysis and biomass formation, cannot be oxidised by oxidative phosphorylation. To solve this problem many microorganisms use pyruvate or one of its derivatives as an electron and hydrogen acceptor thereby regenerating NAD⁺. Thus, in a preferred anaerobic fermentation process pyruvate is used as an electron (and hydrogen acceptor) and is reduced to fermentation products such as ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotics and a cephalosporin. In a preferred embodiment, the fermentation process is anaerobic. An anaerobic process is advantageous since it is cheaper than aerobic processes: less special equipment is needed. Furthermore, anaerobic processes are expected to give a higher product yield than aerobic processes. Under aerobic conditions, usually the biomass yield is higher than under anaerobic conditions. As a consequence, usually under aerobic conditions, the expected product yield is lower than under anaerobic conditions. According to the inventors, the process of the invention is the first anaerobic fermentation process with a medium comprising a source of L-arabinose that has been developed so far.

[0133] In another preferred embodiment, the fermentation process is under oxygen-limited conditions. More preferably, the fermentation process is aerobic and under oxygen-limited conditions. An oxygen-limited fermentation process is a process in which the oxygen consumption is limited by the oxygen transfer from the gas to the liquid. The degree of oxygen limitation is determined by the amount and composition of the ingoing gasflow as well as the actual mixing/mass transfer properties of the fermentation equipment used. Preferably, in a process under oxygen-limited conditions, the rate of oxygen consumption is at least 5.5, more preferably at least 6 and even more preferably at least 7 mmol/L/h.

[0134] The fermentation process is preferably run at a temperature that is optimal for the modified cell. Thus, for most

yeasts or fungal cells, the fermentation process is performed at a temperature which is less than 42° C., preferably less than 38° C. For yeast or filamentous fungal host cells, the fermentation process is preferably performed at a temperature which is lower than 35, 33, 30 or 28° C. and at a temperature which is higher than 20, 22, or 25° C.

[0135] A preferred process is a process for the production of ethanol, whereby the process comprises the steps of: (a) fermenting a medium containing a source of L-arabinose and optionally xylose with a modified host cell as defined herein, whereby the host cell ferments L-arabinose and optionally xylose to ethanol; and optionally, (b) recovery of the ethanol. The fermentation medium may also comprise a source of glucose that is also fermented to ethanol. In a preferred embodiment, the fermentation process for the production of ethanol is anaerobic. Anaerobic has already been defined earlier herein. In another preferred embodiment, the fermentation process for the production of ethanol is aerobic. In another preferred embodiment, the fermentation process for the production of ethanol is under oxygen-limited conditions, more preferably aerobic and under oxygen-limited conditions. Oxygen-limited conditions have already been defined earlier herein.

[0136] In the process, the volumetric ethanol productivity is preferably at least 0.5, 1.0, 1.5, 2.0, 2.5, 3.0, 5.0 or 10.0 g ethanol per litre per hour. The ethanol yield on L-arabinose and optionally xylose and/or glucose in the process preferably is at least 20, 25, 30, 35, 40, 45, 50, 60, 70, 80, 90, 95 or 98%. The ethanol yield is herein defined as a percentage of the theoretical maximum yield, which, for glucose and L-arabinose and optionally xylose is 0.51 g ethanol per g glucose or xylose. In another preferred embodiment, the invention relates to a process for producing a fermentation product selected from the group consisting of lactic acid, 3-hydroxypropionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propanediol, ethylene, glycerol, butanol, a β -lactam antibiotic and a cephalosporin. The process preferably comprises the steps of (a) fermenting a medium containing a source of L-arabinose and optionally xylose with a modified host cell as defined herein above, whereby the host cell ferments L-arabinose and optionally xylose to the fermentation product, and optionally, (b) recovery of the fermentation product. In a preferred process, the medium also contains a source of glucose.

[0137] In the fermentation process of the invention leading to the production of ethanol, several advantages can be cited by comparison to known ethanol fermentations processes:

[0138] anaerobic processes are possible.

[0139] oxygen limited conditions are also possible.

[0140] higher ethanol yields and ethanol production rates can be obtained.

[0141] the strain used may be able to use L-arabinose and optionally xylose.

[0142] Alternatively to the fermentation processes described above, another fermentation process is provided as a further aspect of the invention wherein, at least two distinct cells are used for the fermentation of a carbon source comprising at least two sources of carbon selected from the group consisting of but not limited thereto: a source of L-arabinose, a source of xylose and a source of glucose. In this fermentation process, "at least two distinct cells" means this process is preferably a co-fermentation process. In one preferred embodiment, two distinct cells are used: one being the one of the invention as earlier defined able to use L-arabinose, and/or

to convert it into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product such as ethanol and optionally being able to use xylose, the other one being for example a strain which is able to use xylose and/or convert it into a desired fermentation product such as ethanol as defined in WO 03/062430 and/or WO 06/009434. A cell which is able to use xylose is preferably a strain which exhibits the ability of directly isomerising xylose into xylulose (in one step) as earlier defined herein. These two distinct strains are preferably cultivated in the presence of a source of L-arabinose, a source of xylose and optionally a source of glucose. Three distinct cells or more may be co-cultivated and/or three or more sources of carbon may be used, provided at least one cell is able to use at least one source of carbon present and/or to convert it into a desired fermentation product such as ethanol. The expression "use at least one source of carbon" has the same meaning as the expression "use of L-arabinose". The expression "convert it (i.e. a source of carbon) into a desired fermentation product" has the same meaning as the expression "convert L-arabinose into a desired fermentation product".

[0143] In a preferred embodiment, the invention relates to a process for producing a fermentation product selected from the group consisting of ethanol, lactic acid, 3-hydroxypropionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, amino acids, 1,3-propanediol, ethylene, glycerol, butanol, β -lactam antibiotics and cephalosporins, whereby the process comprises the steps of:

[0144] (a) fermenting a medium containing at least a source of L-arabinose and a source of xylose with a cell of the invention as earlier defined herein and a cell able to use xylose and/or exhibiting the ability to directly isomerise xylose into xylulose, whereby each cell ferments L-arabinose and/or xylose to the fermentation product, and optionally,

[0145] (b) recovering the fermentation product.

All preferred embodiments of the fermentation processes as described above are also preferred embodiments of this further fermentation processes: identity of the fermentation product, identity of source of L-arabinose and source of xylose, conditions of fermentation (aerobical or anaerobical conditions, oxygen-limited conditions, temperature at which the process is being carried out, productivity of ethanol, yield of ethanol).

Genetic Modifications

[0146] For overexpression of enzymes in the host cells of the inventions as described above, as well as for additional genetic modification of host cells, preferably yeasts, host cells are transformed with the various nucleic acid constructs of the invention by methods well known in the art. Such methods are e.g. known from standard handbooks, such as Sambrook and Russel (2001) "Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, or F. Ausubel et al, eds., "Current protocols in molecular biology", Green Publishing and Wiley Interscience, New York (1987). Methods for transformation and genetic modification of fungal host cells are known from e.g. EP-A-0 635 574, WO 98/46772, WO 99/60102 and WO 00/37671.

[0147] Promoters for use in the nucleic acid constructs for overexpression of enzymes in the host cells of the invention have been described above. In the nucleic acid constructs for overexpression, the 3'-end of the nucleotide acid sequence

encoding the enzyme(s) preferably is operably linked to a transcription terminator sequence. Preferably the terminator sequence is operable in a host cell of choice, such as e.g. the yeast species of choice. In any case the choice of the terminator is not critical; it may e.g. be from any yeast gene, although terminators may sometimes work if from a non-yeast, eukaryotic, gene. The transcription termination sequence further preferably comprises a polyadenylation signal. Preferred terminator sequences are the alcohol dehydrogenase (ADH1) and the PGI1 terminators. More preferably, the ADH1 and the PGI1 terminators are both from *S. cerevisiae* (SEQ ID NO:50 and SEQ ID NO:53 respectively).

[0148] Optionally, a selectable marker may be present in the nucleic acid construct. As used herein, the term “marker” refers to a gene encoding a trait or a phenotype which permits the selection of, or the screening for, a host cell containing the marker. The marker gene may be an antibiotic resistance gene whereby the appropriate antibiotic can be used to select for transformed cells from among cells that are not transformed. Preferably however, non-antibiotic resistance markers are used, such as auxotrophic markers (URA3, TRP1, LEU). In a preferred embodiment the host cells transformed with the nucleic acid constructs are marker gene free. Methods for constructing recombinant marker gene free microbial host cells are disclosed in EP-A-0 635 574 and are based on the use of bidirectional markers. Alternatively, a screenable marker such as Green Fluorescent Protein, lacZ, luciferase, chloramphenicol acetyltransferase, beta-glucuronidase may be incorporated into the nucleic acid constructs of the invention allowing to screen for transformed cells.

[0149] Optional further elements that may be present in the nucleic acid constructs of the invention include, but are not limited to, one or more leader sequences, enhancers, integration factors, and/or reporter genes, intron sequences, centromeres, telomeres and/or matrix attachment (MAR) sequences. The nucleic acid constructs of the invention may further comprise a sequence for autonomous replication, such as an ARS sequence. Suitable episomal nucleic acid constructs may e.g. be based on the yeast 2 μ or pKD1 (Fleer et al., 1991, Biotechnology 9:968-975) plasmids. Alternatively the nucleic acid construct may comprise sequences for integration, preferably by homologous recombination. Such sequences may thus be sequences homologous to the target site for integration in the host cell's genome. The nucleic acid constructs of the invention can be provided in a manner known per se, which generally involves techniques such as restricting and linking nucleic acids/nucleic acid sequences, for which reference is made to the standard handbooks, such as Sambrook and Russel (2001) “Molecular Cloning: A Laboratory Manual (3rd edition), Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press.

[0150] Methods for inactivation and gene disruption in yeast or fungi are well known in the art (see e.g. Fincham, 1989, Microbiol Rev. 53(1):148-70 and EP-A-0 635 574).

[0151] In this document and in its claims, the verb “to comprise” and its conjugations is used in its non-limiting sense to mean that items following the word are included, but items not specifically mentioned are not excluded. In addition, reference to an element by the indefinite article “a” or “an” does not exclude the possibility that more than one of the element is present, unless the context clearly requires that there be one and only one of the elements. The indefinite article “a” or “an” thus usually means “at least one”.

[0152] The invention is further described by the following examples, which should not be construed as limiting the scope of the invention.

EXAMPLES

Plasmid and Strain Construction

Strains

[0153] The L-arabinose consuming *Sachharomyces cerevisiae* strain described in this work is based on strain RWB220, which is itself a derivative of RWB217. RWB217 is a CEN. PK strain in which four genes coding for the expression of enzymes in the pentose phosphate pathway have been overexpressed, TAL1, TKL1, RPE1, RKI1 (Kuyper et al., 2005a). In addition the gene coding for an aldose reductase (GRE3), has been deleted. Strain RWB217 also contains two plasmids, a single copy plasmid with a LEU2 marker for overexpression of the xylulokinase (XKS1) and an episomal, multicopy plasmid with URA3 as the marker for the expression of the xylose isomerase, XylA. RWB217 was subjected to a selection procedure for improved growth on xylose which is described in Kuyper et al. (2005b). The procedure resulted in two pure strains, RWB218 (Kuyper et al., 2005b) and RWB219. The difference between RWB218 and RWB219 is that after the selection procedure, RWB218 was obtained by plating and restreaking on mineral medium with glucose as the carbon source, while for RWB219 xylose was used.

[0154] Strain RWB219 was grown non-selectively on YP with glucose (YPD) as the carbon source in order to facilitate the loss of both plasmids. After plating on YPD single colonies were tested for plasmid loss by looking at uracil and leucine auxotrophy. A strain that had lost both plasmids was transformed with pSH47, containing the cre recombinase, in order to remove a KanMX cassette (Guldener et al., 1996), still present after integrating the RKI1 overexpression construct. Colonies with the plasmid were resuspended in Yeast Peptone medium (YP) (10 g/l yeast extract and 20 g/l peptone both from BD Difco Belgium) with 1% galactose and incubated for 1 hour at 30° C. About 200 cells were plated on YPD. The resulting colonies were checked for loss of the KanMX marker (G418 resistance) and pSH47 (URA3). A strain that had lost both the KanMX marker and the pSH47 plasmid was then named RWB220. To obtain the strain tested in this patent, RWB220 was transformed with pRW231 and pRW243 (table 2), resulting in strain IMS0001.

[0155] During construction strains were maintained on complex YP: 10 g l⁻¹ yeast extract (BD Difco), 20 g l⁻¹ peptone (BD Difco) or synthetic medium (MY) (Verduyn et al., 1992) supplemented with glucose (2%) as carbon source (YPD or MYD) and 1.5% agar in the case of plates. After transformation with plasmids strains were plated on MYD. Transformations of yeast were done according to Gietz and Woods (2002). Plasmids were amplified in *Escherichia coli* strain XL-1 blue (Stratagene, La Jolla, Calif., USA). Transformation was performed according to Inoue et al. (1990). *E. coli* was grown on LB (Luria-Bertani) plates or in liquid TB (Terrific Broth) medium for the isolation of plasmids (Sambrook et al, 1989).

Plasmids

[0156] In order to grow on L-arabinose, yeast needs to express three different genes, an L-arabinose isomerase (AraA), a L-ribulokinase (AraB), and a L-ribulose-5-P 4-epi-

merase (AraD) (Becker and Boles, 2003). In this work we have chosen to express AraA, AraB, and AraD from the lactic acid bacterium *Lactobacillus plantarum* in *S. cerevisiae*. Because the eventual aim is to consume L-arabinose in combination with other sugars, like D-xylose, the genes encoding the bacterial L-arabinose pathway were combined on the same plasmid with the genes coding for D-xylose consumption.

[0157] In order to get a high level of expression, the *L. plantarum* AraA and AraD genes were ligated into plasmid pAKX002, the 2 μ XylA bearing plasmid.

[0158] The AraA cassette was constructed by amplifying a truncated version of the TDH3 promoter with SpeI-5'Ptdh3 and 5'AraAPtdh3 (SEQ ID NO: 49), the AraA gene with Ptdh5'AraA and Tadh3'AraA and the ADH1 terminator (SEQ ID NO:50) with 3'AraATadh1 and 3'Tadh1-SpeI. The three fragments were extracted from gel and mixed in roughly equimolar amounts. On this mixture a PCR was performed using the SpeI-5'Ptdh3 and 3'Tadh1SpeI oligos. The resulting P_{TDH3}-AraA-T_{ADH1} cassette was gel purified, cut at the 5' and 3' SpeI sites and then ligated into pAKX002 cut with NheI, resulting in plasmid pRW230.

[0159] The AraD construct was made by first amplifying a truncated version of the HXT7 promoter (SEQ ID NO:52) with oligos SalI5'Phxt7 and 5'AraDPhxt, the AraD gene with Phxt5'AraD and Tpgi3'AraD and the GPII terminator (SEQ ID NO:53) region with the 3'AraDTpgi and 3'TpgiSalI oligos. The resulting fragments were extracted from gel and mixed in roughly equimolar amounts, after which a PCR was performed using the SalI5'Phxt7 and 3'Tpgi1SalI oligos. The resulting P_{HXT7}-AraD-T_{PGI1} cassette was gel purified, cut at the 5' and 3' SalI sites and then ligated into pRW230 cut with XhoI, resulting in plasmid pRW231 (FIG. 1).

[0160] Since too high an expression of the L-ribulokinase is detrimental to growth (Becker and Boles, 2003), the AraB gene was combined with the XKS1 gene, coding for xylulokinase, on an integration plasmid. For this, p415ADHXKS (Kuyper et al., 2005a) was first changed into pRW229, by cutting both p415ADHXKS and pRS305 with PvuI and ligating the ADHXKS-containing PvuI fragment from p415ADHXKS to the vector backbone from pRS305, resulting in pRW229.

[0161] A cassette, containing the *L. plantarum* AraB gene between the PGI1 promoter (SEQ ID NO:51) and ADH1 terminator (SEQ ID NO:50) was made by amplifying the PGI1 promoter with the SacI5'Ppgi1 and 5'AraBPpgi1 oligos, the AraB gene with the Ppgi5'AraB and Tadh3'AraB oligos and the ADH1 terminator with 3'AraBTadh1 and 3'Tadh1SacI oligos. The three fragments were extracted from gel and mixed in roughly equimolar amounts. On this mixture a PCR was performed using the SacI-5'Ppgi 1 and 3'Tadh1SacI oligos. The resulting P_{PGI1}-AraB-T_{ADH1} cassette was gel purified, cut at the 5' and 3' Sad sites and then ligated into pRW229 cut with SacI, resulting in plasmid pRW243 (FIG. 1).

[0162] Strain RWB220 was transformed with pRW231 and pRW243 (table 2), resulting in strain IMS0001.

[0163] Restriction endonucleases (New England Biolabs, Beverly, Mass., USA and Roche, Basel, Switzerland) and DNA ligase (Roche) were used according to the manufacturers' specifications. Plasmid isolation from *E. coli* was performed with the Qiaprep spin miniprep kit (Qiagen, Hilden, Germany). DNA fragments were separated on a 1% agarose (Sigma, St. Louis, Mo., USA) gel in 1 \times TBE (Sambrook et al, 1989). Isolation of fragments from gel was carried out with

the Qiaquick gel extraction kit (Quiagen). Amplification of the (elements of the) AraA, AraB and AraD cassettes was done with Vent_R DNA polymerase (New England Biolabs) according to the manufacturer's specification. The template was chromosomal DNA of *S. cerevisiae* CEN.PK113-7D for the promoters and terminators, or *Lactobacillus plantarum* DSM20205 for the Ara genes. The polymerase chain reaction (PCR) was performed in a Biometra TGradient Thermocycler (Biometra, Gottingen, Germany) with the following settings: 30 cycles of 1 min annealing at 55 $^{\circ}$ C., 60 $^{\circ}$ C. or 65 $^{\circ}$ C., 1 to 3 min extension at 75 $^{\circ}$ C., depending on expected fragment size, and 1 min denaturing at 94 $^{\circ}$ C.

Cultivation and Media

[0164] Shake-flask cultivations were performed at 30 $^{\circ}$ C. in a synthetic medium (Verduyn et al., 1992). The pH of the medium was adjusted to 6.0 with 2 M KOH prior to sterilisation. For solid synthetic medium, 1.5% of agar was added.

[0165] Pre-cultures were prepared by inoculating 100 ml medium containing the appropriate sugar in a 500-ml shake flask with a frozen stock culture. After incubation at 30 $^{\circ}$ C. in an orbital shaker (200 rpm), this culture was used to inoculate either shake-flask cultures or fermenter cultures. The synthetic medium for anaerobic cultivation was supplemented with 0.01 g l⁻¹ ergosterol and 0.42 g Tween 80 dissolved in ethanol (Andreasen and Stier, 1953; Andreasen and Stier, 1954). Anaerobic (sequencing) batch cultivation was carried out at 30 $^{\circ}$ C. in 2-1 laboratory fermenters (Applikon, Schiedam, The Netherlands) with a working volume of 1 l. The culture pH was maintained at pH 5.0 by automatic addition of 2 M KOH. Cultures were stirred at 800 rpm and sparged with 0.5 l min⁻¹ nitrogen gas (<10 ppm oxygen). To minimise diffusion of oxygen, fermenters were equipped with Norprene tubing (Cole Palmer Instrument company, Vernon Hills, USA). Dissolved oxygen was monitored with an oxygen electrode (Applisens, Schiedam, The Netherlands). Oxygen-limited conditions were achieved in the same experimental set-up by headspace aeration at approximately 0.05 l min⁻¹.

Determination of Dry Weight

[0166] Culture samples (10.0 ml) were filtered over pre-weighed nitrocellulose filters (pore size 0.45 μ m; Gelman laboratory, Ann Arbor, USA). After removal of medium, the filters were washed with demineralised water and dried in a microwave oven (Bosch, Stuttgart, Germany) for 20 min at 360 W and weighed. Duplicate determinations varied by less than 1%.

Gas Analysis

[0167] Exhaust gas was cooled in a condensor (2 $^{\circ}$ C.) and dried with a Permapure dryer type MD-110-48P-4 (Permapure, Toms River, USA). O₂ and CO₂ concentrations were determined with a NGA 2000 analyser (Rosemount Analytical, Orrville, USA). Exhaust gasflow rate and specific oxygen-consumption and carbondioxide production rates were determined as described previously (Van Urk et al., 1988; Weusthuis et al., 1994). In calculating these biomass-specific rates, volume changes caused by withdrawing culture samples were taken account for.

Metabolite Analysis

[0168] Glucose, xylose, arabinose, xylitol, organic acids, glycerol and ethanol were analysed by HPLC using a Waters

Alliance 2690 HPLC (Waters, Milford, USA) supplied with a BioRad HPX 87H column (BioRad, Hercules, USA), a Waters 2410 refractive-index detector and a Waters 2487 UV detector. The column was eluted at 60° C. with 0.5 g l⁻¹ sulphuric acid at a flow rate of 0.6 ml min⁻¹.

Assay for Xylulose 5-Phosphate (Zaldivar J., et al, Appl. Microbiol. Biotechnol., (2002), 59:436-442)

[0169] For the analysis of intracellular metabolites such as xylulose 5-phosphate, 5 ml broth was harvested in duplicate from the reactors, before glucose exhaustion (at 22 and 26 h of cultivation) and after glucose exhaustion (42, 79 and 131 h of cultivation). Procedures for metabolic arrest, solid-phase extraction of metabolites and analysis have been described in detail by Smits H. P. et al. (Anal. Biochem., 261:36-42, (1998)). However, the analysis by high-pressure ion exchange chromatography coupled to pulsed amperometric detection used to analyze cell extracts, was slightly modified. Solutions used were eluent A, 75 mM NaOH, and eluent B, 500 mM NaAc. To prevent contamination of carbonate in the eluent solutions, a 50% NaOH solution with low carbonate concentration (Baker Analysed, Deventer, The Netherlands) was used instead of NaOH pellets. The eluents were degassed with Helium (He) for 30 min and then kept under a He atmosphere. The gradient pump was programmed to generate the following gradients: 100% A and 0% B (0 min), a linear decrease of A to 70% and a linear increase of B to 30% (0-30 min), a linear decrease of A to 30% and a linear increase of B to 70% (30-70 min), a linear decrease of A to 0% and a linear increase of B to 100% (70-75 min), 0% A and 100% B (75-85 min), a linear increase of A to 100% and a linear decrease of B to 0% (85-95 min). The mobile phase was run at a flow rate of 1 ml/min. Other conditions were according to Smits et al. (1998).

Carbon Recovery

[0170] Carbon recoveries were calculated as carbon in products formed, divided by the total amount of sugar carbon consumed, and were based on a carbon content of biomass of 48%. To correct for ethanol evaporation during the fermentations, the amount of ethanol produced was assumed to be equal to the measured cumulative production of CO₂ minus the CO₂ production that occurred due to biomass synthesis (5.85 mmol CO₂ per gram biomass (Verduyn et al., 1990)) and the CO₂ associated with acetate formation.

Selection for Growth on L-Arabinose

[0171] Strain IMS0001 (CBS120327 deposited at the CBS on 27/09/06), containing the genes encoding the pathways for both xylose (XylA and XKS1) and arabinose (AraA, AraB, AraD) metabolization, was constructed according the procedure described above. Although capable of growing on xylose (data not shown), strain IMS0001 did not seem to be capable of growing on solid synthetic medium supplemented with 2% L-arabinose. Mutants of IMS0001 capable of utilizing L-arabinose as carbon source for growth were selected by serial transfer in shake flasks and by sequencing-batch cultivation in fermenters (SBR).

[0172] For the serial transfer experiments, a 500-ml shake flask containing 100 ml synthetic medium containing 0.5% galactose were inoculated with either strain IMS0001, or the reference strain RWB219. After 72 hours, at an optical density at 660 nm of 3.0, the cultures were used to inoculate a new shake flask containing 0.1% galactose and 2% arabinose.

Based on HPLC determination with D-ribulose as calibration standard, it was determined that already in the first cultivations of strain IMS0001, on medium containing a galactose/arabinose mixture, part of the arabinose was converted into ribulose and subsequently excreted to the supernatant. These HPLC analyses were performed using a Waters Alliance 2690 HPLC (Waters, Milford, USA) supplied with a BioRad HPX 87H column (BioRad, Hercules, USA), a Waters 2410 refractive-index detector and a Waters 2487 UV detector. The column was eluted at 60° C. with 0.5 g sulphuric acid at a flow rate of 0.6 ml min⁻¹. In contrast to the reference strain RWB219, the OD₆₆₀ of the culture of strain IMS0001 increased after depletion of the galactose. When after approximately 850 hours growth on arabinose by strain IMS0001 was observed (FIG. 2), this culture was transferred at an OD₆₆₀ of 1.7 to a shake flask containing 2% arabinose. Cultures were then sequentially transferred to fresh medium containing 2% arabinose at an OD₆₆₀ of 2-3. Utilization of arabinose was confirmed by occasionally measuring arabinose concentrations by HPLC (data not shown). The growth rate of these cultures increased from 0 to 0.15 h⁻¹ in approximately 3600 hours (FIG. 3).

[0173] A batch fermentation under oxygen limited conditions was started by inoculating 1 l of synthetic medium supplemented with 2% of arabinose with a 100 ml shake flask culture of arabinose-grown IMS0001 cells with a maximum growth rate on 2% of L-arabinose of approximately 0.12 h⁻¹. When growth on arabinose was observed, the culture was subjected to anaerobic conditions by sparging with nitrogen gas. The sequential cycles of anaerobic batch cultivation were started by either manual or automated replacement of 90% of the culture with synthetic medium with 20 g l⁻¹ arabinose. For each cycle during the SBR fermentation, the exponential growth rate was estimated from the CO₂ profile (FIG. 4). In 13 cycles, the exponential growth rate increased from 0.025 to 0.08 h⁻¹. After 20 cycles a sample was taken, and plated on solid synthetic medium supplemented with 2% of L-arabinose and incubated at 30° C. for several days. Separate colonies were re-streaked twice on solid synthetic medium with L-arabinose. Finally, a shake flask containing synthetic medium with 2% of L-L-arabinose was inoculated with a single colony, and incubated for 5 days at 30° C. This culture was designated as strain IMS0002 (CBS120328 deposited at the Centraal Bureau voor Schimmelculturen (CBS) on 27/09/06). Culture samples were taken, 30% of glycerol was added and samples were stored at -80° C.

Mixed Culture Fermentation

[0174] Biomass hydrolysates, a desired feedstock for industrial biotechnology, contain complex mixtures consisting of various sugars amongst which glucose, xylose and arabinose are commonly present in significant fractions. To accomplish ethanolic fermentation of not only glucose and arabinose, but also xylose, an anaerobic batch fermentation was performed with a mixed culture of the arabinose-fermenting strain IMS0002, and the xylose-fermenting strain RWB218. An anaerobic batch fermenter containing 800 ml of synthetic medium supplied with 30 g l⁻¹ D-glucose, 15 g l⁻¹ D-xylose, and 15 g l⁻¹ L-arabinose was inoculated with 100 ml of pre-culture of strain IMS0002. After 10 hours, a 100 ml inoculum of RWB218 was added. In contrast to the mixed sugar fermentation with only strain IMS0002, both xylose and arabinose were consumed after glucose depletion (FIG. 5D). The mixed culture completely consumed all sugars, and

within 80 hours 564.0 ± 6.3 mmol l⁻¹ ethanol (calculated from the CO₂ production) was produced with a high overall yield of 0.42 g g⁻¹ sugar. Xylitol was produced only in small amounts, to a concentration of 4.7 mmol l⁻¹.

Characterization of Strain IMS0002

[0175] Growth and product formation of strain IMS0002 was determined during anaerobic batch fermentations on synthetic medium with either L-arabinose as the sole carbon source, or a mixture of glucose, xylose and L-arabinose. The pre-cultures for these anaerobic batch fermentations were prepared in shake flasks containing 100 ml of synthetic medium with 2% L-arabinose, by inoculating with -80° C. frozen stocks of strain IMS0002, and incubating for 48 hours at 30° C.

[0176] FIG. 5A shows that strain IMS0002 is capable of fermenting 20 g l⁻¹ L-arabinose to ethanol during an anaerobic batch fermentation of approximately 70 hours. The specific growth rate under anaerobic conditions with L-arabinose as sole carbon source was 0.05 ± 0.001 h⁻¹. Taking into account the ethanol evaporation during the batch fermentation, the ethanol yield from 20 g l⁻¹ arabinose was 0.43 ± 0.003 g g⁻¹. Without evaporation correction the ethanol yield was 0.35 ± 0.01 g g⁻¹ of arabinose. No formation of arabinitol was observed during anaerobic growth on arabinose. In FIG. 5B, the ethanolic fermentation of a mixture of 20 g l⁻¹ glucose and 20 g l⁻¹ L-arabinose by strain IMS0002 is shown. L-arabinose consumption started after glucose depletion. Within 70 hours, both the glucose and L-arabinose were completely consumed. The ethanol yield from the total of sugars was 0.42 ± 0.003 g g⁻¹.

[0177] In FIG. 5C, the fermentation profile of a mixture of 30 g l⁻¹ glucose, 15 g l⁻¹ D-xylose, and 15 g l⁻¹ L-arabinose by strain IMS0002 is shown. Arabinose consumption started after glucose depletion. Within 80 hours, both the glucose and arabinose were completely consumed. Only 20 mM from 100 mM of xylose was consumed by strain IMS0002. In addition, the formation of 20 mM of xylitol was observed. Apparently, the xylose was converted into xylitol by strain IMS0002. Hence, the ethanol yield from the total of sugars was lower than for the above described fermentations: 0.38 ± 0.001 g g⁻¹. The ethanol yield from the total of glucose and arabinose was similar to the other fermentations: 0.43 ± 0.001 g g⁻¹.

[0178] Table 1 shows the arabinose consumption rates and the ethanol production rates observed for the anaerobic batch fermentation of strain IMS0002. Arabinose was consumed with a rate of 0.23-0.75 g h⁻¹ g⁻¹ biomass dry weight. The rate of ethanol produced from arabinose varied from 0.08-0.31 g h⁻¹ g⁻¹ biomass dry weight.

[0179] Initially, the constructed strain IMS0001 was able to ferment xylose (data not shown). In contrast to our expectations, the selected strain IMS0002 was not capable of fermenting xylose to ethanol (FIG. 5C). To regain the capability of fermenting xylose, a colony of strain IMS0002 was transferred to solid synthetic medium with 2% of D-xylose, and incubated in an anaerobic jar at 30° C. for 25 days. Subsequently, a colony was again transferred to solid synthetic medium with 2% of arabinose. After 4 days of incubation at 30° C., a colony was transferred to a shake flask containing synthetic medium with 2% arabinose. After incubation at 30° C. for 6 days, 30% of glycerol was added, samples were taken and stored at -80° C. A shake flask containing 100 ml of synthetic medium with 2% arabinose was inoculated with such a frozen stock, and was used as preculture for an anaero-

bic batch fermentation on synthetic medium with 20 g l⁻¹ xylose and 20 g l⁻¹ arabinose. In FIG. 6, the fermentation profile of this batch fermentation is shown. Xylose and arabinose were consumed simultaneously. The arabinose was completed within 70 hours, whereas the xylose was completely consumed in 120 hours. At least 250 mM of ethanol was produced from the total of sugars, not taking into account the evaporation of the ethanol. Assuming an end biomass dry weight of 3.2 g l⁻¹ (assuming a biomass yield of 0.08 g g⁻¹ sugar), the end ethanol concentration estimated from the cumulative CO₂ production (355 mmol l⁻¹) was approximately 330 mmol l⁻¹, corresponding to a ethanol yield of 0.41 g g⁻¹ pentose sugar. In addition to ethanol, glycerol, and organic acids, a small amount of xylitol was produced (approximately 5 mM).

Selection of Strain IMS0003

[0180] Initially, the constructed strain IMS0001 was able to ferment xylose (data not shown). In contrast to our expectations, the selected strain IMS0002 was not capable of fermenting xylose to ethanol (FIG. 5C). To regain the capability of fermenting xylose, a colony of strain IMS0002 was transferred to solid synthetic medium with 2% of D-xylose, and incubated in an anaerobic jar at 30° C. for 25 days. Subsequently, a colony was again transferred to solid synthetic medium with 2% of arabinose. After 4 days of incubation at 30° C., a colony was transferred to a shake flask containing synthetic medium with 2% arabinose. After incubation at 30° C. for 6 days, 30% of glycerol was added, samples were taken and stored at -80° C.

[0181] From this frozen stock, samples were spread on solid synthetic medium with 2% of L-arabinose and incubated at 30° C. for several days. Separate colonies were re-streaked twice on solid synthetic medium with L-arabinose. Finally, a shake flask containing synthetic medium with 2% of L-arabinose was inoculated with a single colony, and incubated for 4 days at 30° C. This culture was designated as strain IMS0003 (CBS 121879 deposited at the CBS on 20/09/07). Culture samples were taken, 30% of glycerol was added and samples were stored at -80° C.

Characterization of Strain IMS0003

[0182] Growth and product formation of strain IMS0003 was determined during an anaerobic batch fermentation on synthetic medium with a mixture of 30 g l⁻¹ glucose, 15 g l⁻¹ D-xylose and 15 g l⁻¹ L-arabinose. The pre-culture for this anaerobic batch fermentation was prepared in a shake flasks containing 100 ml of synthetic medium with 2% L-arabinose, by inoculating with a -80° C. frozen stock of strain IMS0003, and incubated for 48 hours at 30° C.

[0183] In FIG. 7, the fermentation profile of a mixture of 30 g l⁻¹ glucose, 15 g l⁻¹ D-xylose, and 15 g l⁻¹ L-arabinose by strain IMS0003 is shown. Arabinose consumption started after glucose depletion. Within 70 hours, the glucose, xylose and arabinose were completely consumed. Xylose and arabinose were consumed simultaneously. At least 406 mM of ethanol was produced from the total of sugars, not taking into account the evaporation of the ethanol. The final ethanol concentration calculated from the cumulative CO₂ production was 572 mmol l⁻¹, corresponding to an ethanol yield of 0.46 g g⁻¹ of total sugar. In contrast to the fermentation of a mixture of glucose, xylose and arabinose by strain IMS0002 (FIG. 5C) or a mixed culture of strains IMS0002 and RWB218 (FIG. 5D), strain IMS0003 did not produce detectable amounts of xylitol.

TABLE 1

<u>S. cerevisiae strains used.</u>		
Strain	Characteristics	Reference
RWB217	MATA ura3-52 leu2-112 loxP-P _{TPI} ::(-266, -1)TAL1 gre3::hphMX pUGP _{TPI} TKL1 pUGP _{TPI} -RPEI KanloxP-P _{TPI} ::(-?, -1)RKII {p415ADHXKS, PAKX002}	Kuyper et al. 2005a
RWB218	MATA ura3-52 leu2-112 loxP-P _{TPI} ::(-266, -1)TAL1 gre3::hphMX pUGP _{TPI} TKL1 pUGP _{TPI} -RPEI KanloxP-P _{TPI} ::(-?, -1)RKII {p415ADHXKS1, pAKX002}	Kuyper et al. 2005b
RWB219	MATA ura3-52 leu2-112 loxP-P _{TPI} ::(-266, -1)TAL1 gre3::hphMX pUGP _{TPI} TKL1 pUGP _{TPI} -RPEI KanloxP-P _{TPI} ::(-?, -1)RKII {p415ADHXKS1, pAKX002}	This work
RWB220	MATA ura3-52 leu2-112 loxP-P _{TPI} ::(-266, -1)TAL1 gre3::hphMX pUGP _{TPI} TKL1 pUGP _{TPI} -RPEI loxP-P _{TPI} ::(-?, -1)RKII	This work
IMS0001	MATA ura3-52 leu2-112 loxP-P _{TPI} ::(-266, -1)TAL1 gre3::hphMX pUGP _{TPI} TKL1 pUGP _{TPI} -RPEI loxP-P _{TPI} ::(-?, -1)RKII {pRW231, PRW243}	This work
IMS0002	MATA ura3-52 leu2-112 loxP-P _{TPI} ::(-266, -1)TAL1 gre3::hphMX pUGP _{TPI} TKL1 pUGP _{TPI} -RPEI loxP-P _{TPI} ::(-?, -1)RKII {pRW231, PRW243} selected for anaerobic growth on L-arabinose	This work

TABLE 2

<u>Plasmids used</u>		
plasmid	characteristics	Reference
pRS305	Integration, LEU2	Gietz and Sugino, 1988
pAKX002	2 μ , URA3, P _{TPI} Piromyces xyla	Kuyper et al. 2003
p415ADHXKS1	CEN, LEU2, P _{ADHI} S.cerXKS1	Kuyper et al., 2005a
pRW229	integration, LEU2, P _{ADHI} S.cerXKS1	This work
pRW230	pAKX002 with P _{IDH3} -AraA	This work
pRW231	pAKX002 with P _{IDH3} -AraA and P _{HXT1} -AraD	This work
pRW243	LEU2, integration, P _{ADHI} ScXKS1-T _{CYC} , P _{PGI} -L.plantarumAraB-T _{ADHI}	This work

TABLE 3

<u>oligos used in this work</u>	
Oligo	DNA sequence
<u>AraA expression cassette</u>	
SpeI5' Ptdh3	5' GACTAGTCGAGTTTATCATTATCAATACTGC3'
SEQ ID NO:	31
5' AraAPtdh	5' CTCATAATCAGGTACTGATAACATTTTGTGTTTATGTGTGTTTATTTC3'
SEQ ID NO:	32
Ptdh5' AraA	5' GAATAAACACACATAAACAAACAAAATGTTATCAGTACCTGATTATGAG3
SEQ ID NO:	33
Tadh3' AraA	5' AATCATAAATCATAAGAAATTCGCTTACTTTAAGAATGCCTTAGTCAT3'
SEQ ID NO:	34
3' AraATadh 1	5' ATGACTAAGGCATTCTTAAAGTAAGCGAATTTCTTATGATTTATGATT3'
SEQ ID NO:	35
3' Tadh 1 SpeI	5' CACTAGTCTCGAGTGTGGAAGAACGATTACAACAGG3'
SEQ ID NO:	36
<u>AraB expression cassette</u>	
SacI5' Ppgi1	5' CGAGCTCGTGGGTGATTGGATTATAGGAAG3'
SEQ ID NO:	37
5' AraBPpgi1	5' TTGGGCTGTTTCAACTAAATTCATTTTGGCTGGTATCTTGATTCTA3'
SEQ ID NO:	38
Ppgi5' AraB	5' TAGAATCAAGATACCAGCCTAAAAATGAATTTAGTTGAAACAGCCCAA3'
SEQ ID NO:	39

TABLE 3-continued

<u>oligos used in this work</u>	
Oligo	DNA sequence
Tadh3'AraB	5'AATCATAAATCATAAGAAATTCGCTCTAATATTTGATTGCTTGCCAG3'
SEQ ID NO: 40	
3'AraBTadh 1	5'CTGGCAAGCAATCAAATATTAGAGCGAATTTCTTATGATTTATGATT3'
SEQ ID NO: 41	
3'Tadh 1 SacI	5'TGAGCTCGTGTGGAAGAACGATTACAACAGG3'
SEQ ID NO: 42	
<u>AraD expression cassette</u>	
SalI5'Phxt7	5'ACGCGTCGACTCGTAGGAACAATTTTCGG3'
SEQ ID NO: 43	
5'AraDPhxt	5'CTTCTTGTTTTAATGCTTCTAGCATTTTTGGATTAAAATTAATAAACTT3'
SEQ ID NO: 44	
Phxt5'AraD	5'AAGTTTTTTTAATTTTAATCAAAAAATGCTAGAAGCATTAAACAAGAAG3'
SEQ ID NO: 45	
Tpgi3'AraD	5'GGTATATATTTAAGAGCGATTTGTTTACTTGCGAACTGCATGATCC3'
SEQ ID NO: 46	
3'AraDTpgi	5'GGATCATGCAGTTCGCAAGTAAACAAATCGCTCTTAAATATATACC3'
SEQ ID NO: 47	
3'TpgiSalI	5'CGCAGTCGACCTTTTAAACAGTTGATGAGAACC3'
SEQ ID NO: 48	

TABLE 4

Maximum observed specific glucose and arabinose consumption rates and ethanol production rates during anaerobic batch fermentations of <i>S. cerevisiae</i> IMS0002.				
C-source	q_{glu} g h ⁻¹ g ⁻¹ DW	q_{ara} g h ⁻¹ g ⁻¹ DW	$q_{eth, glu}$ g h ⁻¹ g ⁻¹ DW	$q_{eth, ara}$ g h ⁻¹ g ⁻¹ DW
20 g l ⁻¹ arabinose	—	0.75 ± 0.04	—	0.31 ± 0.02
20 g l ⁻¹ glucose	2.08 ± 0.09	0.41 ± 0.01	0.69 ± 0.00	0.19 ± 0.00
20 g l ⁻¹ arabinose				
30 g l ⁻¹ glucose	1.84 ± 0.04	0.23 ± 0.01	0.64 ± 0.03	0.08 ± 0.01
15 g l ⁻¹ xylose				
15 g l ⁻¹ arabinose				

q_{glu} : specific glucose consumption rate

q_{ara} : specific arabinose consumption rate

$q_{eth, glu}$: specific ethanol production rate during growth on glucose

$q_{eth, ara}$: specific ethanol production rate during growth on arabinose

REFERENCE LIST

- [0184] Andreasen AA, Stier T J (1954) Anaerobic nutrition of *Saccharomyces cerevisiae*. II. Unsaturated fatty acid requirement for growth in a defined medium. *J Cell Physiol* 43:271-281
- [0185] Andreasen AA, Stier T J (1953) Anaerobic nutrition of *Saccharomyces cerevisiae*. I. Ergosterol requirement for growth in a defined medium. *J Cell Physiol* 41:23-36
- [0186] Becker J, Boles E (2003) A modified *Saccharomyces cerevisiae* strain that consumes L-Arabinose and produces ethanol, *Appl Environ Microbiol* 69:4144-4150
- [0187] Gietz R. D., Sugino A. (1988). New yeast-*Escherichia coli* shuttle vectors constructed with in vitro mutagenized yeast genes lacking six-base pair restriction sites. *Gene* 74:527-534.
- [0188] Gietz, R. D., and R. A. Woods. 2002. Transformation of yeast by lithium acetate/single-stranded carrier DNA/polyethylene glycol method. *Methods Enzymol.* 350:87-96.
- [0189] Guldener U, Heck S, Fielder T, Beinhauer J, Hegemann J H. (1996) A new efficient gene disruption cassette for repeated use in budding yeast. *Nucleic Acids Res.* 1996 Jul. 1; 24(13):2519-24.
- [0190] Hauf J, Zimmermann F K, Muller S. Simultaneous genomic overexpression of seven glycolytic enzymes in the yeast *Saccharomyces cerevisiae*. *Enzyme Microb Technol.* 2000 Jun. 1; 26(9-10):688-698.
- [0191] Inoue H., H. Nojima and H. Okayama, High efficiency transformation of *Escherichia coli* with plasmids. *Gene* 96 (1990), pp. 23-28

- [0192] Kuyper M, Hartog M M P, Toirkens M J, Almering M J H, Winkler A A, Van Dijken J P, Pronk J T (2005a) Metabolic engineering of a xylose-isomerase-expressing *Saccharomyces cerevisiae* strain for rapid anaerobic xylose fermentation. *Ferns Yeast Research* 5:399-409
- [0193] Kuyper M, Toirkens M J, Diderich J A, Winkler A A, Van Dijken J P, Pronk J T (2005b) Evolutionary engineering of mixed-sugar utilization by a xylose-fermenting *Saccharomyces cerevisiae* strain. *Ferns Yeast Research* 5:925-934
- [0194] Sambrook, K., Fritsch, E. F. and Maniatis, I. (1989) *Molecular Cloning: A Laboratory Manual*, 2nd edn. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
- [0195] Van Urk H, Mak P R, Scheffers W A, Van Dijken J P (1988) Metabolic responses of *Saccharomyces cerevisiae* CBS 8066 and *Candida utilis* CBS 621 upon transition from glucose limitation to glucose excess. *Yeast* 4:283-291
- [0196] Verduyn C, Postma E, Scheffers W A, Van Dijken J P (1990) Physiology of *Saccharomyces cerevisiae* in anaerobic glucose-limited chemostat cultures. *J Gen Microbiol* 136:395-403
- [0197] Verduyn C, Postma E, Scheffers W A, Van Dijken J P (1992) Effect of benzoic acid on metabolic fluxes in yeasts: a continuous-culture study on the regulation of respiration and alcoholic fermentation. *Yeast* 8:501-517
- [0198] Weusthuis R A, Visser W, Pronk J T, Scheffers W A, Van Dijken J P (1994) Effects of oxygen limitation on sugar metabolism in yeasts—a continuous-culture study of the Kluyver effect. *Microbiology* 140:703-715

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 53

<210> SEQ ID NO 1

<211> LENGTH: 474

<212> TYPE: PRT

<213> ORGANISM: *Lactobacillus plantarum*

<400> SEQUENCE: 1

```

Met Leu Ser Val Pro Asp Tyr Glu Phe Trp Phe Val Thr Gly Ser Gln
1          5          10          15
His Leu Tyr Gly Glu Glu Gln Leu Lys Ser Val Ala Lys Asp Ala Gln
20          25          30
Asp Ile Ala Asp Lys Leu Asn Ala Ser Gly Lys Leu Pro Tyr Lys Val
35          40          45
Val Phe Lys Asp Val Met Thr Thr Ala Glu Ser Ile Thr Asn Phe Met
50          55          60
Lys Glu Val Asn Tyr Asn Asp Lys Val Ala Gly Val Ile Thr Trp Met
65          70          75          80
His Thr Phe Ser Pro Ala Lys Asn Trp Ile Arg Gly Thr Glu Leu Leu
85          90          95
Gln Lys Pro Leu Leu His Leu Ala Thr Gln Tyr Leu Asn Asn Ile Pro
100         105         110
Tyr Ala Asp Ile Asp Phe Asp Tyr Met Asn Leu Asn Gln Ser Ala His
115         120         125
Gly Asp Arg Glu Tyr Ala Tyr Ile Asn Ala Arg Leu Gln Lys His Asn
130         135         140
Lys Ile Val Tyr Gly Tyr Trp Gly Asp Glu Asp Val Gln Glu Gln Ile
145         150         155         160
Ala Arg Trp Glu Asp Val Ala Val Ala Tyr Asn Glu Ser Phe Lys Val
165         170         175
Lys Val Ala Arg Phe Gly Asp Thr Met Arg Asn Val Ala Val Thr Glu
180         185         190
Gly Asp Lys Val Glu Ala Gln Ile Lys Met Gly Trp Thr Val Asp Tyr
195         200         205
Tyr Gly Ile Gly Asp Leu Val Glu Glu Ile Asn Lys Val Ser Asp Ala
210         215         220
Asp Val Asp Lys Glu Tyr Ala Asp Leu Glu Ser Arg Tyr Glu Met Val
225         230         235         240

```

-continued

Gln Val Asp Asn Asp Ala Asp Thr Tyr Lys His Ser Val Arg Val Gln
245 250 255

Leu Ala Gln Tyr Leu Gly Ile Lys Arg Phe Leu Glu Arg Gly Gly Tyr
260 265 270

Thr Ala Phe Thr Thr Asn Phe Glu Asp Leu Trp Gly Met Glu Gln Leu
275 280 285

Pro Gly Leu Ala Ser Gln Leu Leu Ile Arg Asp Gly Tyr Gly Phe Gly
290 295 300

Ala Glu Gly Asp Trp Lys Thr Ala Ala Leu Gly Arg Val Met Lys Ile
305 310 315 320

Met Ser His Asn Lys Gln Thr Ala Phe Met Glu Asp Tyr Thr Leu Asp
325 330 335

Leu Arg His Gly His Glu Ala Ile Leu Gly Ser His Met Leu Glu Val
340 345 350

Asp Pro Ser Ile Ala Ser Asp Lys Pro Arg Val Glu Val His Pro Leu
355 360 365

Asp Ile Gly Gly Lys Asp Asp Pro Ala Arg Leu Val Phe Thr Gly Ser
370 375 380

Glu Gly Glu Ala Ile Asp Val Thr Val Ala Asp Phe Arg Asp Gly Phe
385 390 395 400

Lys Met Ile Ser Tyr Ala Val Asp Ala Asn Lys Pro Glu Ala Glu Thr
405 410 415

Pro Asn Leu Pro Val Ala Lys Gln Leu Trp Thr Pro Lys Met Gly Leu
420 425 430

Lys Lys Gly Ala Leu Glu Trp Met Gln Ala Gly Gly Gly His His Thr
435 440 445

Met Leu Ser Phe Ser Leu Thr Glu Glu Gln Met Glu Asp Tyr Ala Thr
450 455 460

Met Val Gly Met Thr Lys Ala Phe Leu Lys
465 470

<210> SEQ ID NO 2

<211> LENGTH: 1425

<212> TYPE: DNA

<213> ORGANISM: Lactobacillus plantarum

<400> SEQUENCE: 2

```

atgttatcag tacctgatta tgagttttgg tttgttaccg gttcacaaca cctttatggt    60
gaagaacaat tgaagtctgt tgctaaggat gcgcaagata ttgctgataa attgaatgca    120
agcggcaagt tacctataa agtagtcttt aaggatgta tgacgacggc tgaaagtatc    180
accaacttta tgaagaagt taattacaat gataaggtag ccggtggtat tacttggatg    240
cacacattct caccagctaa gaactggatt cgtggaactg aactgttaca aaaaccatta    300
ttacacttag caacgcaata tttgaataat attccatag cagacattga ctttgattac    360
atgaacctta accaaagtgc ccatggcgac cgcgagtatg cctacattaa cgcccggttg    420
cagaaacata ataagattgt ttacggctat tggggcgatg aagatgtgca agagcagatt    480
gcacgttggg aagacgtcgc cgtagcgtac aatgagagct ttaaagttaa gggtgctcgc    540
ttggcgaca caatgcgtaa tgtggccggt actgaaggtg acaaggttga agctcaaatt    600
aagatgggct ggacagttga ctattatggt atcggtgact tagttgaaga gatcaataag    660
gtttcggatg ctgatgttga taaggaatac gctgacttgg agtctcggta tgaaatggtc    720

```


-continued

```

caagttgata acgatgcgga cacgtataaa cattcagttc gggttcaatt ggcacaatat 780
ctgggtatta agcggttctt agaaagagggc ggttacacag cctttaccac gaactttgaa 840
gatctttggg ggatggagca attacctggg ctagcttcac aattattaat tcgtgatggg 900
tatggttttg gtgctgaagg tgactggaag acggctgctt taggacgggt tatgaagatt 960
atgtctcaca acaagcaaac cgcctttatg gaagactaca cgntagactt gcgtcatggt 1020
catgaagcga tcttaggttc acacatggtg gaagttgatc cgtctatcgc aagtgataaa 1080
ccacgggtcg aagttcatcc attggatatt gggggtaaag atgatcctgc tcgcctagta 1140
tttactgggt cagaagggtg agcaattgat gtcaccgttg ccgatttccg tgatgggttc 1200
aagatgatta gctacgcggt agatgcgaat aagccagaag ccgaaacacc taatttacca 1260
gttgctaagc aattatggac cccaaagatg ggcttgaaga aggggtgact agaatggatg 1320
caagctggtg gtggtcacca cacgatgctg tccttctcgt taactgaaga acaaatggaa 1380
gactatgcaa ccatggttgg catgactaag gcattcttaa agtaa 1425

```

```

<210> SEQ ID NO 3
<211> LENGTH: 533
<212> TYPE: PRT
<213> ORGANISM: Lactobacillus plantarum

```

```

<400> SEQUENCE: 3

```

```

Met Asn Leu Val Glu Thr Ala Gln Ala Ile Lys Thr Gly Lys Val Ser
1          5          10          15

Leu Gly Ile Glu Leu Gly Ser Thr Arg Ile Lys Ala Val Leu Ile Thr
20          25          30

Asp Asp Phe Asn Thr Ile Ala Ser Gly Ser Tyr Val Trp Glu Asn Gln
35          40          45

Phe Val Asp Gly Thr Trp Thr Tyr Ala Leu Glu Asp Val Trp Thr Gly
50          55          60

Ile Gln Gln Ser Tyr Thr Gln Leu Ala Ala Asp Val Arg Ser Lys Tyr
65          70          75          80

His Met Ser Leu Lys His Ile Asn Ala Ile Gly Ile Ser Ala Met Met
85          90          95

His Gly Tyr Leu Ala Phe Asp Gln Gln Ala Lys Leu Leu Val Pro Phe
100         105         110

Arg Thr Trp Arg Asn Asn Ile Thr Gly Gln Ala Ala Asp Glu Leu Thr
115         120         125

Glu Leu Phe Asp Phe Asn Ile Pro Gln Arg Trp Ser Ile Ala His Leu
130         135         140

Tyr Gln Ala Ile Leu Asn Asn Glu Ala His Val Lys Gln Val Asp Phe
145         150         155         160

Ile Thr Thr Leu Ala Gly Tyr Val Thr Trp Lys Leu Ser Gly Glu Lys
165         170         175

Val Leu Gly Ile Gly Asp Ala Ser Gly Val Phe Pro Ile Asp Glu Thr
180         185         190

Thr Asp Thr Tyr Asn Gln Thr Met Leu Thr Lys Phe Ser Gln Leu Asp
195         200         205

Lys Val Lys Pro Tyr Ser Trp Asp Ile Arg His Ile Leu Pro Arg Val
210         215         220

Leu Pro Ala Gly Ala Ile Ala Gly Lys Leu Thr Ala Ala Gly Ala Ser

```

-continued

225	230	235	240
Leu Leu Asp Gln Ser Gly Thr Leu Asp Ala Gly Ser Val Ile Ala Pro	245	250	255
Pro Glu Gly Asp Ala Gly Thr Gly Met Val Gly Thr Asn Ser Val Arg	260	265	270
Lys Arg Thr Gly Asn Ile Ser Val Gly Thr Ser Ala Phe Ser Met Asn	275	280	285
Val Leu Asp Lys Pro Leu Ser Lys Val Tyr Arg Asp Ile Asp Ile Val	290	295	300
Met Thr Pro Asp Gly Ser Pro Val Ala Met Val His Val Asn Asn Cys	305	310	315
Ser Ser Asp Ile Asn Ala Trp Ala Thr Ile Phe Arg Glu Phe Ala Ala	325	330	335
Arg Leu Gly Met Glu Leu Lys Pro Asp Arg Leu Tyr Glu Thr Leu Phe	340	345	350
Leu Glu Ser Thr Arg Ala Asp Ala Asp Ala Gly Gly Leu Ala Asn Tyr	355	360	365
Ser Tyr Gln Ser Gly Glu Asn Ile Thr Lys Ile Gln Ala Gly Arg Pro	370	375	380
Leu Phe Val Arg Thr Pro Asn Ser Lys Phe Ser Leu Pro Asn Phe Met	385	390	395
Leu Thr Gln Leu Tyr Ala Ala Phe Ala Pro Leu Gln Leu Gly Met Asp	405	410	415
Ile Leu Val Asn Glu Glu His Val Gln Thr Asp Val Met Ile Ala Gln	420	425	430
Gly Gly Leu Phe Arg Thr Pro Val Ile Gly Gln Gln Val Leu Ala Asn	435	440	445
Ala Leu Asn Ile Pro Ile Thr Val Met Ser Thr Ala Gly Glu Gly Gly	450	455	460
Pro Trp Gly Met Ala Val Leu Ala Asn Phe Ala Cys Arg Gln Thr Ala	465	470	475
Met Asn Leu Glu Asp Phe Leu Asp Gln Glu Val Phe Lys Glu Pro Glu	485	490	495
Ser Met Thr Leu Ser Pro Glu Pro Glu Arg Val Ala Gly Tyr Arg Glu	500	505	510
Phe Ile Gln Arg Tyr Gln Ala Gly Leu Pro Val Glu Ala Ala Ala Gly	515	520	525
Gln Ala Ile Lys Tyr	530		

<210> SEQ ID NO 4

<211> LENGTH: 1602

<212> TYPE: DNA

<213> ORGANISM: Lactobacillus plantarum

<400> SEQUENCE: 4

```

atgaatttag ttgaaacagc ccaagcgatt aaaactggca aagtttcttt aggaattgag      60
cttggtcaa ctcgaattaa agccgttttg atcacggagc attttaatac gattgcttcg      120
ggaagttacg tttgggaaa ccaatttggt gatggtactt ggacttacgc acttgaagat      180
gtctggaccg gaattcaaca aagttatacg caattagcag cagatgtccg cagtaaatat      240
cacatgagtt tgaagcatat caatgctatt ggcattagtg ccatgatgca cggataccta      300

```

-continued

```

gcatttgatc aacaagcga attattagtt ccgtttcgga cttggcgtaa taacattacg 360
gggcaagcag cagatgaatt gaccgaatta tttgatttca acattccaca acggtggagt 420
atcgcgcact tataccaggc aatcttaa atgaagcgc acgttaaaca ggtggacttc 480
ataacaacgc tggctggcta tgtaacctgg aaattgtcgg gtgagaaagt tctaggaatc 540
ggtgatgcgt ctggcgtttt cccaattgat gaaacgactg acacatacaa tcagacgatg 600
ttaaccaagt ttagccaact tgacaaagtt aaaccgtatt catgggatat cggcatatt 660
ttaccgctgg tttaccagc gggagccatt gctggaaagt taacggctgc cggggcgagc 720
ttacttgatc agagcggcac gctcgacgct ggcagtgtta ttgcaccgcc agaaggggat 780
gctggaacag gaatggtcgg tacgaacagc gtccgtaa gcacgggtaa catctcgggtg 840
ggaacctcag cattttcgat gaacgttcta gataaacat tgtctaaagt ctatcgcgat 900
attgatattg ttatgacgcc agatgggtca ccagttgcaa tggatgatgt taataattgt 960
tcatcagata ttaatgcgtg ggcaacgatt ttcgtgagt ttgcagccc gttgggaatg 1020
gaattgaaac cggatcgatt atatgaaacg ttattcttgg aatcaactcg cgctgatgcg 1080
gatgctggag ggttggctaa ttatagttat caatccggtg agaatattac taagattcaa 1140
gctggtcggc cgctatttgt acggacacca aacagtaaat ttagtttacc gaactttatg 1200
ttgaccaat tatatgcggc gttcgcaccc ctccaacttg gtatggatat tcttgtaac 1260
gaagaacatg ttcaaacgga cgttatgatt gcacagggtg gattgttccg aacgcggta 1320
attggccaac aagtattggc caacgcactg aacattccga ttactgtaat gagtactgct 1380
ggtgaaggcg gccatgggg gatggcagtg ttagccaact ttgcttgcg gcaaactgca 1440
atgaacctag aagatttctt agatcaagaa gtctttaag agccagaaag tatgacgctg 1500
agtccagaac cggaacgggt ggccggatat cgtgaattta ttcaacgta tcaagctggc 1560
ttaccagttg aagcagcggc tgggcaagca atcaaatatt ag 1602

```

<210> SEQ ID NO 5

<211> LENGTH: 242

<212> TYPE: PRT

<213> ORGANISM: Lactobacillus plantarum

<400> SEQUENCE: 5

```

Met Leu Glu Ala Leu Lys Gln Glu Val Tyr Glu Ala Asn Met Gln Leu
1           5           10           15

Pro Lys Leu Gly Leu Val Thr Phe Thr Trp Gly Asn Val Ser Gly Ile
          20           25           30

Asp Arg Glu Lys Gly Leu Phe Val Ile Lys Pro Ser Gly Val Asp Tyr
          35           40           45

Gly Glu Leu Lys Pro Ser Asp Leu Val Val Val Asn Leu Gln Gly Glu
          50           55           60

Val Val Glu Gly Lys Leu Asn Pro Ser Ser Asp Thr Pro Thr His Thr
65           70           75           80

Val Leu Tyr Asn Ala Phe Pro Asn Ile Gly Gly Ile Val His Thr His
          85           90           95

Ser Pro Trp Ala Val Ala Tyr Ala Ala Ala Gln Met Asp Val Pro Ala
          100          105          110

Met Asn Thr Thr His Ala Asp Thr Phe Tyr Gly Asp Val Pro Ala Ala
          115          120          125

```

-continued

Asp Ala Leu Thr Lys Glu Glu Ile Glu Ala Asp Tyr Glu Gly Asn Thr
 130 135 140
 Gly Lys Thr Ile Val Lys Thr Phe Gln Glu Arg Gly Leu Asp Tyr Glu
 145 150 155 160
 Ala Val Pro Ala Ser Leu Val Ser Gln His Gly Pro Phe Ala Trp Gly
 165 170 175
 Pro Thr Pro Ala Lys Ala Val Tyr Asn Ala Lys Val Leu Glu Val Val
 180 185 190
 Ala Glu Glu Asp Tyr His Thr Ala Gln Leu Thr Arg Ala Ser Ser Glu
 195 200 205
 Leu Pro Gln Tyr Leu Leu Asp Lys His Tyr Leu Arg Lys His Gly Ala
 210 215 220
 Ser Ala Tyr Tyr Gly Gln Asn Asn Ala His Ser Lys Asp His Ala Val
 225 230 235 240
 Arg Lys

<210> SEQ ID NO 6
 <211> LENGTH: 729
 <212> TYPE: DNA
 <213> ORGANISM: Lactobacillus plantarum

<400> SEQUENCE: 6

atgctagaag cattaanaaca agaagtttat gaggctaaca tgcagcttcc aaagctgggc 60
 ctggttactt ttacctgggg caatgtctcg ggcattgacc gggaaaaagg cctattcgtg 120
 atcaagccat ctggtgttga ttatgggtgaa ttaaaaccaa gcgatttagt cgttggttaac 180
 ttacagggtg aagtgggtga aggtaaacta aatccgtcta gtgatacgcc gactcatacg 240
 gtgttatata acgcttttcc taatattggc ggaattgtcc atactcattc gccatgggca 300
 gttgcctatg cagctgctca aatggatgtg ccagctatga acacgaccca tgctgatacg 360
 ttctatggtg acgtgccggc cgcggatgcg ctgactaagg aagaaattga agcagattat 420
 gaaggcaaca cgggtaaac cattgtgaag acgttccaag aacggggcct cgattatgaa 480
 gctgtaccag cctcattagt cagccagcac ggcccatttg cttggggacc aacgccagct 540
 aaagccgttt acaatgctaa agtggtggaa gtggttgccg aagaagatta tcatactgcg 600
 caattgaccg gtgcaagtag cgaattacca caatatttat tagataagca ttatttacgt 660
 aagcatggtg caagtgccta ttatgggtcaa aataatgcgc attctaagga tcatgcagtt 720
 cgcaagtaa 729

<210> SEQ ID NO 7
 <211> LENGTH: 437
 <212> TYPE: PRT
 <213> ORGANISM: Piromyces sp.

<400> SEQUENCE: 7

Met Ala Lys Glu Tyr Phe Pro Gln Ile Gln Lys Ile Lys Phe Glu Gly
 1 5 10 15
 Lys Asp Ser Lys Asn Pro Leu Ala Phe His Tyr Tyr Asp Ala Glu Lys
 20 25 30
 Glu Val Met Gly Lys Lys Met Lys Asp Trp Leu Arg Phe Ala Met Ala
 35 40 45
 Trp Trp His Thr Leu Cys Ala Glu Gly Ala Asp Gln Phe Gly Gly Gly

-continued

50					55					60					
Thr	Lys	Ser	Phe	Pro	Trp	Asn	Glu	Gly	Thr	Asp	Ala	Ile	Glu	Ile	Ala
65					70					75					80
Lys	Gln	Lys	Val	Asp	Ala	Gly	Phe	Glu	Ile	Met	Gln	Lys	Leu	Gly	Ile
				85					90					95	
Pro	Tyr	Tyr	Cys	Phe	His	Asp	Val	Asp	Leu	Val	Ser	Glu	Gly	Asn	Ser
			100					105					110		
Ile	Glu	Glu	Tyr	Glu	Ser	Asn	Leu	Lys	Ala	Val	Val	Ala	Tyr	Leu	Lys
			115				120					125			
Glu	Lys	Gln	Lys	Glu	Thr	Gly	Ile	Lys	Leu	Leu	Trp	Ser	Thr	Ala	Asn
						135					140				
Val	Phe	Gly	His	Lys	Arg	Tyr	Met	Asn	Gly	Ala	Ser	Thr	Asn	Pro	Asp
145						150					155				160
Phe	Asp	Val	Val	Ala	Arg	Ala	Ile	Val	Gln	Ile	Lys	Asn	Ala	Ile	Asp
				165					170					175	
Ala	Gly	Ile	Glu	Leu	Gly	Ala	Glu	Asn	Tyr	Val	Phe	Trp	Gly	Gly	Arg
			180					185					190		
Glu	Gly	Tyr	Met	Ser	Leu	Leu	Asn	Thr	Asp	Gln	Lys	Arg	Glu	Lys	Glu
			195				200					205			
His	Met	Ala	Thr	Met	Leu	Thr	Met	Ala	Arg	Asp	Tyr	Ala	Arg	Ser	Lys
	210					215					220				
Gly	Phe	Lys	Gly	Thr	Phe	Leu	Ile	Glu	Pro	Lys	Pro	Met	Glu	Pro	Thr
225						230					235				240
Lys	His	Gln	Tyr	Asp	Val	Asp	Thr	Glu	Thr	Ala	Ile	Gly	Phe	Leu	Lys
				245					250					255	
Ala	His	Asn	Leu	Asp	Lys	Asp	Phe	Lys	Val	Asn	Ile	Glu	Val	Asn	His
			260					265					270		
Ala	Thr	Leu	Ala	Gly	His	Thr	Phe	Glu	His	Glu	Leu	Ala	Cys	Ala	Val
			275				280					285			
Asp	Ala	Gly	Met	Leu	Gly	Ser	Ile	Asp	Ala	Asn	Arg	Gly	Asp	Tyr	Gln
	290					295					300				
Asn	Gly	Trp	Asp	Thr	Asp	Gln	Phe	Pro	Ile	Asp	Gln	Tyr	Glu	Leu	Val
305						310					315				320
Gln	Ala	Trp	Met	Glu	Ile	Ile	Arg	Gly	Gly	Gly	Phe	Val	Thr	Gly	Gly
				325					330					335	
Thr	Asn	Phe	Asp	Ala	Lys	Thr	Arg	Arg	Asn	Ser	Thr	Asp	Leu	Glu	Asp
			340					345					350		
Ile	Ile	Ile	Ala	His	Val	Ser	Gly	Met	Asp	Ala	Met	Ala	Arg	Ala	Leu
			355				360					365			
Glu	Asn	Ala	Ala	Lys	Leu	Leu	Gln	Glu	Ser	Pro	Tyr	Thr	Lys	Met	Lys
	370						375				380				
Lys	Glu	Arg	Tyr	Ala	Ser	Phe	Asp	Ser	Gly	Ile	Gly	Lys	Asp	Phe	Glu
385						390					395				400
Asp	Gly	Lys	Leu	Thr	Leu	Glu	Gln	Val	Tyr	Glu	Tyr	Gly	Lys	Lys	Asn
				405					410					415	
Gly	Glu	Pro	Lys	Gln	Thr	Ser	Gly	Lys	Gln	Glu	Leu	Tyr	Glu	Ala	Ile
			420					425						430	
Val	Ala	Met	Tyr	Gln											
			435												

-continued

<211> LENGTH: 1669

<212> TYPE: DNA

<213> ORGANISM: *Piromyces* sp.

<400> SEQUENCE: 8

```

gtaaatggct aaggaatatt tcccacaaat tcaaaagatt aagttcgaag gtaaggattc   60
taagaatcca ttagccttcc actactacga tgctgaaaag gaagtcattg gtaagaaaat   120
gaaggattgg ttacgtttcg ccatggcctg gtggcacact ctttgcgccg aagggtgctga   180
ccaattcggg ggaggtacaa agtctttccc atggaacgaa ggtactgatg ctattgaaat   240
tgccaagcaa aaggttgatg ctggtttcca aatcatgcaa aagcttggtg ttccatacta   300
ctgtttccac gatgttgatc ttgtttccga aggtaactct attgaagaat acgaatccaa   360
ccttaaggct gtcgttgctt acctcaagga aaagcaaaag gaaaccggtg ttaagcttct   420
ctggagtact gctaactgct tcggtcacaa gcgttacatg aacggtgctt ccaactaacc   480
agactttgat gttgtcgccc gtgctattgt tcaaattaag aacgccatag acgccggtat   540
tgaacttggg gctgaaaact acgtcttctg ggggtggtcg gaaggttaca tgagtctcct   600
taacactgac caaaagcgtg aaaaggaaca catggccact atgcttacca tggctcgtga   660
ctacgctcgt tccaagggat tcaagggtac tttcctcatt gaaccaaagc caatggaacc   720
aaccaagcac caatacgatg ttgacactga aaccgctatt ggtttcctta aggcccacaa   780
cttagacaag gacttcaagg tcaacattga agttaaccac gctactcttg ctggtcacac   840
tttcgaacac gaacttgctt gtgctgttga tgctgggatg ctcggttcca ttgatgctaa   900
ccgtggtgac taccaaaacg gttgggatac tgatcaattc ccaattgatc aatacgaact   960
cgtccaagct tggatggaaa tcatccgtgg tgggtggttc gttactggtg gtaccaactt  1020
cgatgccaaag actcgtcgtg actctactga cctcgaagac atcatcattg cccacgtttc  1080
tggatgggat gctatggctc gtgctcttga aaacgtgcc aagctcctcc aagaatctcc  1140
atacaccaag atgaagaagg aacgttacgc ttccttcgac agtggatttg gtaaggactt  1200
tgaagatggg aagctcacc cogaacaagt ttacgaatac ggtaagaaga acggtgaacc  1260
aaagcaaaact tctggtgagc aagaactcta cgaagctatt gttgccatgt accaataagt  1320
taatcgtagt taaattggta aaataattgt aaaatcaata aacttgctca tctccaatc  1380
aagtttaaaa gatcctatct ctgtactaat taaatatagt acaaaaaaaaa atgtataaac  1440
aaaaaaaaagt ctaaaagacg gaagaattta atttagggaa aaaataaaaa taataataaa  1500
caatagataa atcctttata ttaggaaaat gtcccattgt attattttca tttctactaa  1560
aaaagaaagt aaataaaaca caagaggaaa tttcccttt ttttttttt tgtaataaat  1620
tttatgcaaa tataaatata aataaaataa taaaaaaaaa aaaaaaaaaa  1669

```

<210> SEQ ID NO 9

<211> LENGTH: 496

<212> TYPE: PRT

<213> ORGANISM: *Bacillus subtilis*

<400> SEQUENCE: 9

```

Met Leu Gln Thr Lys Asp Tyr Glu Phe Trp Phe Val Thr Gly Ser Gln
1           5           10           15

His Leu Tyr Gly Glu Glu Thr Leu Glu Leu Val Asp Gln His Ala Lys
          20           25           30

```

-continued

Ser Ile Cys Glu Gly Leu Ser Gly Ile Ser Ser Arg Tyr Lys Ile Thr
 35 40 45
 His Lys Pro Val Val Thr Ser Pro Glu Thr Ile Arg Glu Leu Leu Arg
 50 55 60
 Glu Ala Glu Tyr Ser Glu Thr Cys Ala Gly Ile Ile Thr Trp Met His
 65 70 75 80
 Thr Phe Ser Pro Ala Lys Met Trp Ile Glu Gly Leu Ser Ser Tyr Gln
 85 90 95
 Lys Pro Leu Met His Leu His Thr Gln Tyr Asn Arg Asp Ile Pro Trp
 100 105 110
 Gly Thr Ile Asp Met Asp Phe Met Asn Ser Asn Gln Ser Ala His Gly
 115 120 125
 Asp Arg Glu Tyr Gly Tyr Ile Asn Ser Arg Met Gly Leu Ser Arg Lys
 130 135 140
 Val Ile Ala Gly Tyr Trp Asp Asp Glu Glu Val Lys Lys Glu Met Ser
 145 150 155 160
 Gln Trp Met Asp Thr Ala Ala Ala Leu Asn Glu Ser Arg His Ile Lys
 165 170 175
 Val Ala Arg Phe Gly Asp Asn Met Arg His Val Ala Val Thr Asp Gly
 180 185 190
 Asp Lys Val Gly Ala His Ile Gln Phe Gly Trp Gln Val Asp Gly Tyr
 195 200 205
 Gly Ile Gly Asp Leu Val Glu Val Met Asp Arg Ile Thr Asp Asp Glu
 210 215 220
 Val Asp Thr Leu Tyr Ala Glu Tyr Asp Arg Leu Tyr Val Ile Ser Glu
 225 230 235 240
 Glu Thr Lys Arg Asp Glu Ala Lys Val Ala Ser Ile Lys Glu Gln Ala
 245 250 255
 Lys Ile Glu Leu Gly Leu Thr Ala Phe Leu Glu Gln Gly Gly Tyr Thr
 260 265 270
 Ala Phe Thr Thr Ser Phe Glu Val Leu His Gly Met Lys Gln Leu Pro
 275 280 285
 Gly Leu Ala Val Gln Arg Leu Met Glu Lys Gly Tyr Gly Phe Ala Gly
 290 295 300
 Glu Gly Asp Trp Lys Thr Ala Ala Leu Val Arg Met Met Lys Ile Met
 305 310 315 320
 Ala Lys Gly Lys Arg Thr Ser Phe Met Glu Asp Tyr Thr Tyr His Phe
 325 330 335
 Glu Pro Gly Asn Glu Met Ile Leu Gly Ser His Met Leu Glu Val Cys
 340 345 350
 Pro Thr Val Ala Leu Asp Gln Pro Lys Ile Glu Val His Ser Leu Ser
 355 360 365
 Ile Gly Gly Lys Glu Asp Pro Ala Arg Leu Val Phe Asn Gly Ile Ser
 370 375 380
 Gly Ser Ala Ile Gln Ala Ser Ile Val Asp Ile Gly Gly Arg Phe Arg
 385 390 395 400
 Leu Val Leu Asn Glu Val Asn Gly Gln Glu Ile Glu Lys Asp Met Pro
 405 410 415
 Asn Leu Pro Val Ala Arg Val Leu Trp Lys Pro Glu Pro Ser Leu Lys
 420 425 430
 Thr Ala Ala Glu Ala Trp Ile Leu Ala Gly Gly Ala His His Thr Cys

-continued

435	440	445	
Leu Ser Tyr Glu Leu Thr Ala Glu Gln Met Leu Asp Trp Ala Glu Met			
450	455	460	
Ala Gly Ile Glu Ser Val Leu Ile Ser Arg Asp Thr Thr Ile His Lys			
465	470	475	480
Leu Lys His Glu Leu Lys Trp Asn Glu Ala Leu Tyr Arg Leu Gln Lys			
485	490	495	

<210> SEQ ID NO 10
 <211> LENGTH: 1511
 <212> TYPE: DNA
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 10

```

atgagaaagg ggcagtttac atgcttcaga caaaggatta tgaattctgg tttgtgacag    60
gaagccagca cctatacggg gaagagacgc tggaaactcgt agatcagcat gctaaaagca    120
tttgtgaggg gctcagcggg atttcttcca gatataaaat cactcataag cccgtcgtca    180
cttcaccgga aaccattaga gagctgtaa gagaagcgga gtacagtgag acatgtgctg    240
gcatcattac atggatgcac acattttccc ctgcaaaaat gtggatagaa ggcctttcct    300
cttatcaaaa accgcttatg catttgcata cccaatataa tcgcgatata ccgtagggta    360
cgattgacat ggattttatg aacagcaacc aatccgcgca tggcgatcga gactacgggt    420
acatcaactc gagaatgggg cttagccgaa aagtcaattgc cggctattgg gatgatgaag    480
aagtgaaaaa agaatgtcc cagtggatgg atacggcggc tgcattaaat gaaagcagac    540
atattaaggt tgccagattt ggagataaca tgcgtcatgt cgcggtaacg gacggagaca    600
aggtgggagc gcatattcaa tttggctggc aggttgacgg atatggcatc ggggatctcg    660
ttgaagtgat ggatcgcatt acggacgacg aggttgacac gctttatgcc gagtatgaca    720
gactatatgt gatcagtgag gaaacaaaac gtgacgaagc aaaggtagcg tccattaaag    780
aacaggcgaa aattgaactt ggattaaccg cttttcttga gcaaggcgga tacacagcgt    840
ttacgacatc gtttgaagtg ctgcacggaa tgaaacagct gccgggactt gccgttcagc    900
gcctgatgga gaaaggctat gggtttgccg gtgaaggaga ttggaagaca gcggcccttg    960
tacggatgat gaaaatcatg gctaaaggaa aaagaacttc cttcatggaa gattacacgt   1020
accattttga accgggaaat gaaatgattc tgggctctca catgcttgaa gtgtgtccga   1080
ctgtcgcttt ggatcagccg aaaatcgagg ttcattcgct ttcgattggc ggcaaagagg   1140
accctgcgcg tttggtattt aacggcatca gcggttctgc cattcaagct agcattgttg   1200
atattggcgg gcgtttccgc cttgtgctga atgaagtcaa cggccaggaa attgaaaaag   1260
acatgccgaa tttaccggtt gcccggtgtt tctggaagcc ggagccgtca ttgaaaacag   1320
cagcggaggc atggatttta gccggcggtg cacaccatac ctgcctgtct tatgaactga   1380
cagcggagca aatgcttgat tgggcggaaa tggcgggaat cgaaagtgtt ctcatttccc   1440
gtgatacgac aattcataaa ctgaaacacg agttaaatag gaacgagcgc ctttaccggc   1500
ttcaaaagta g                                     1511
    
```

<210> SEQ ID NO 11
 <211> LENGTH: 566
 <212> TYPE: PRT
 <213> ORGANISM: E. coli

-continued

<400> SEQUENCE: 11

Met Ala Ile Ala Ile Gly Leu Asp Phe Gly Ser Asp Ser Val Arg Ala
1 5 10 15
Leu Ala Val Asp Cys Ala Ser Gly Glu Glu Ile Ala Thr Ser Val Glu
20 25 30
Trp Tyr Pro Arg Trp Gln Lys Gly Gln Phe Cys Asp Ala Pro Asn Asn
35 40 45
Gln Phe Arg His His Pro Arg Asp Tyr Ile Glu Ser Met Glu Ala Ala
50 55 60
Leu Lys Thr Val Leu Ala Glu Leu Ser Val Glu Gln Arg Ala Ala Val
65 70 75 80
Val Gly Ile Gly Val Asp Ser Thr Gly Ser Thr Pro Ala Pro Ile Asp
85 90 95
Ala Asp Gly Asn Val Leu Ala Leu Arg Pro Glu Phe Ala Glu Asn Pro
100 105 110
Asn Ala Met Phe Val Leu Trp Lys Asp His Thr Ala Val Glu Arg Ser
115 120 125
Glu Glu Ile Thr Arg Leu Cys His Ala Pro Gly Asn Val Asp Tyr Ser
130 135 140
Arg Tyr Ile Gly Gly Ile Tyr Ser Ser Glu Trp Phe Trp Ala Lys Ile
145 150 155 160
Leu His Val Thr Arg Gln Asp Ser Ala Val Ala Gln Ser Ala Ala Ser
165 170 175
Trp Ile Glu Leu Cys Asp Trp Val Pro Ala Leu Leu Ser Gly Thr Thr
180 185 190
Arg Pro Gln Asp Ile Arg Arg Gly Arg Cys Ser Ala Gly His Lys Ser
195 200 205
Leu Trp His Glu Ser Trp Gly Gly Leu Pro Pro Ala Ser Phe Phe Asp
210 215 220
Glu Leu Asp Pro Ile Leu Asn Arg His Leu Pro Ser Pro Leu Phe Thr
225 230 235 240
Asp Thr Trp Thr Ala Asp Ile Pro Val Gly Thr Leu Cys Pro Glu Trp
245 250 255
Ala Gln Arg Leu Gly Leu Pro Glu Ser Val Val Ile Ser Gly Gly Ala
260 265 270
Phe Asp Cys His Met Gly Ala Val Gly Ala Gly Ala Gln Pro Asn Ala
275 280 285
Leu Val Lys Val Ile Gly Thr Ser Thr Cys Asp Ile Leu Ile Ala Asp
290 295 300
Lys Gln Ser Val Gly Glu Arg Ala Val Lys Gly Ile Cys Gly Gln Val
305 310 315 320
Asp Gly Ser Val Val Pro Gly Phe Ile Gly Leu Glu Ala Gly Gln Ser
325 330 335
Ala Phe Gly Asp Ile Tyr Ala Trp Phe Gly Arg Val Leu Ser Trp Pro
340 345 350
Leu Glu Gln Leu Ala Ala Gln His Pro Glu Leu Lys Ala Gln Ile Asn
355 360 365
Ala Ser Gln Lys Gln Leu Leu Pro Ala Leu Thr Glu Ala Trp Ala Lys
370 375 380
Asn Pro Ser Leu Asp His Leu Pro Val Val Leu Asp Trp Phe Asn Gly

-continued

385	390	395	400
Arg Arg Ser Pro	Asn Ala Asn Gln Arg	Leu Lys Gly Val Ile Thr Asp	
	405	410	415
Leu Asn Leu Ala Thr Asp Ala Pro Leu Leu Phe Gly Gly Leu Ile Ala		425	430
	420		
Ala Thr Ala Phe Gly Ala Arg Ala Ile Met Glu Cys Phe Thr Asp Gln		440	445
	435		
Gly Ile Ala Val Asn Asn Val Met Ala Leu Gly Gly Ile Ala Arg Lys		455	460
	450		
Asn Gln Val Ile Met Gln Ala Cys Cys Asp Val Leu Asn Arg Pro Leu		470	475
	465		480
Gln Ile Val Ala Ser Asp Gln Cys Cys Ala Leu Gly Ala Ala Ile Phe		490	495
	485		
Ala Ala Val Ala Ala Lys Val His Ala Asp Ile Pro Ser Ala Gln Gln		505	510
	500		
Lys Met Ala Ser Ala Val Glu Lys Thr Leu Gln Pro Arg Ser Glu Gln		520	525
	515		
Ala Gln Arg Phe Glu Gln Leu Tyr Arg Arg Tyr Gln Gln Trp Ala Met		535	540
	530		
Ser Ala Glu Gln His Tyr Leu Pro Thr Ser Ala Pro Ala Gln Ala Ala		555	560
	545		565
Gln Ala Val Ala Thr Leu			
	565		

<210> SEQ ID NO 12
 <211> LENGTH: 1453
 <212> TYPE: DNA
 <213> ORGANISM: E. coli

<400> SEQUENCE: 12

```

atggcgattg caattggcct cgattttggc agtgattctg tgcgagcttt ggcggtggac    60
tgcgccagcg gtgaagagat cgccaccagc gtagagtggg atccccgttg gcaaaaaggg    120
caattttgtg atgccccgaa taaccagttc cgatcatcgc cgcgtgacta cattgagtca    180
atggaagcgg cactgaaaac cgtgcttgca gagcttagcg tcgaacagcg cgcagctgtg    240
gtcgggattg gcggtgacag taccggctcg acgcccgcac cgattgatgc cgacggtaac    300
gtgctggcgc tgcgcccgga gtttgccgaa aaccggaacg cgatgttcgt attgtggaaa    360
gaccacactg cggttgaaag aagcgaagag attaccogtt tgtgccacgc gccgggcaat    420
gttgactact cccgctatat tggcggattt tattccagcg aatggttctg ggcaaaaatc    480
ctgcatgtga ctgccagga cagcgccgtg gcgcaatctg ccgcatcgtg gattgagctg    540
tgcgactggg tgccagctct gctttccggt accaccgcc cgcaggatat tcgtcgcgga    600
cgttgcagcg ccgggcataa atctctgtgg cacgaaagct ggggcggtt gccgccagcc    660
agtttctttg atgagctgga cccgatcctc aatcgccatt tgccttcccc gctgttccact    720
gacacctgga ctgccgatat tccgggtggc accttatgcc cggaatgggc gcagcgtctc    780
ggcctgcctg aaagcgtggg gatttccggc ggccgctttg actgccatat gggcgcagtt    840
ggcgcagcgc cacagcctaa cgcactggta aaagttatcg gtacttccac ctgcgacatt    900
ctgattgccg acaaacagag cgttggcgag cgggcagtta aaggtatttg cggtcaggtt    960
gatggcagcg tgggtgctgg atttatcggg ctggaagcag gccaatcggc gtttggtgat   1020
    
```

-continued

```

atctacgect ggttcggtcg cgtactcagc tggccgctgg aacagcttgc cgcccagcat 1080
ccggaactga aagcgcaaat caacgccagc cagaaacaac tgcttccggc gctgaccgaa 1140
gcatggggcca aaaatccgtc tctggatcac ctgccggtgg tgctcgactg gtttaacggt 1200
cgctcgctgc caaacgctaa ccaacgcctg aaaggggtga ttaccgatct taacctcgct 1260
accgacgctc cgctgctggt cggcgggttg attgctgcca ccgcctttgg cgcacgcgca 1320
atcatggagt gctttaccga tcaggggatc gccgtcaata acgtgatggc gctggggcggc 1380
atcgcgcgga aaaaccaagt cattatgcag gcctgctgcg acgtgctgaa tcgcccgtg 1440
caaattggtg cct 1453

```

```

<210> SEQ ID NO 13
<211> LENGTH: 231
<212> TYPE: PRT
<213> ORGANISM: E. coli

```

```

<400> SEQUENCE: 13

```

```

Met Leu Glu Asp Leu Lys Arg Gln Val Leu Glu Ala Asn Leu Ala Leu
1           5           10           15

Pro Lys His Asn Leu Val Thr Leu Thr Trp Gly Asn Val Ser Ala Val
          20           25           30

Asp Arg Glu Arg Gly Val Phe Val Ile Lys Pro Ser Gly Val Asp Tyr
          35           40           45

Ser Ile Met Thr Ala Asp Asp Met Val Val Val Ser Ile Glu Thr Gly
          50           55           60

Glu Val Val Glu Gly Ala Lys Lys Pro Ser Ser Asp Thr Pro Thr His
65           70           75           80

Arg Leu Leu Tyr Gln Ala Phe Pro Ser Ile Gly Gly Ile Val His Thr
          85           90           95

His Ser Arg His Ala Thr Ile Trp Ala Gln Ala Gly Gln Ser Ile Pro
          100          105          110

Ala Thr Gly Thr Thr His Ala Asp Tyr Phe Tyr Gly Thr Ile Pro Cys
          115          120          125

Thr Arg Lys Met Thr Asp Ala Glu Ile Asn Gly Glu Tyr Glu Trp Glu
          130          135          140

Thr Gly Asn Val Ile Val Glu Thr Phe Glu Lys Gln Gly Ile Asp Ala
145          150          155          160

Ala Gln Met Pro Gly Val Leu Val His Ser His Gly Pro Phe Ala Trp
          165          170          175

Gly Lys Asn Ala Glu Asp Ala Val His Asn Ala Ile Val Leu Glu Glu
          180          185          190

Val Ala Tyr Met Gly Ile Phe Cys Arg Gln Leu Ala Pro Gln Leu Pro
          195          200          205

Asp Met Gln Gln Thr Leu Leu Asn Lys His Tyr Leu Arg Lys His Gly
          210          215          220

Ala Lys Ala Tyr Tyr Gly Gln
225          230

```

```

<210> SEQ ID NO 14
<211> LENGTH: 696
<212> TYPE: DNA
<213> ORGANISM: E. coli

```

-continued

<400> SEQUENCE: 14

```

atgtagaag atctcaaacg ccaggtatta gaggccaacc tggcgctgcc aaaacataac    60
ctggtcacgc tcacatgggg caacgtcagc gccgttgatc gcgagcgcgg cgtctttgtg   120
atcaaaccctt cggcgctcga ttacagcatc atgaccgctg acgatatggt cgtgggttagc   180
atcgaaaccg gtgaagtggg tgaaggtgcg aaaaagccct cctccgatac gccaactcac   240
cgactgctct atcaggcatt cccgtccatt ggcggcattg tgcacacaca ctcgcgccac   300
gccactatct gggcgagcgg gggccagtcg attccagcaa ccggcaccac ccacgcccagc   360
tatttctacg gcaccattcc ctgcaccgca aaaatgaccg acgcagaaat caacgggtgaa   420
tatgagtggg aaaccggtaa cgtcatcgta gaaaccttcg aaaaacaggg tatcgatgca   480
gcgcaaatgc ccggcgctct ggtccattct cacggcccat ttgcatgggg caaaaatgcc   540
gaagatgagg tgcataacgc catcgtgctg gaagaggtcg cttatatggg gatattctgc   600
cgtcagttag cgccgagtt accggatagc cagcaaaccg tgctgaataa aactatctg    660
cgtaagcatg gcgcaaggc atattacggg cagtaa                                696

```

<210> SEQ ID NO 15

<211> LENGTH: 438

<212> TYPE: PRT

<213> ORGANISM: Bacteroides thetaiotaomicron

<400> SEQUENCE: 15

```

Met Ala Thr Lys Glu Phe Phe Pro Gly Ile Glu Lys Ile Lys Phe Glu
1           5           10          15
Gly Lys Asp Ser Lys Asn Pro Met Ala Phe Arg Tyr Tyr Asp Ala Glu
          20          25          30
Lys Val Ile Asn Gly Lys Lys Met Lys Asp Trp Leu Arg Phe Ala Met
          35          40          45
Ala Trp Trp His Thr Leu Cys Ala Glu Gly Gly Asp Gln Phe Gly Gly
          50          55          60
Gly Thr Lys Gln Phe Pro Trp Asn Gly Asn Ala Asp Ala Ile Gln Ala
65          70          75          80
Ala Lys Asp Lys Met Asp Ala Gly Phe Glu Phe Met Gln Lys Met Gly
          85          90          95
Ile Glu Tyr Tyr Cys Phe His Asp Val Asp Leu Val Ser Glu Gly Ala
          100         105         110
Ser Val Glu Glu Tyr Glu Ala Asn Leu Lys Glu Ile Val Ala Tyr Ala
          115         120         125
Lys Gln Lys Gln Ala Glu Thr Gly Ile Lys Leu Leu Trp Gly Thr Ala
          130         135         140
Asn Val Phe Gly His Ala Arg Tyr Met Asn Gly Ala Ala Thr Asn Pro
145         150         155         160
Asp Phe Asp Val Val Ala Arg Ala Ala Val Gln Ile Lys Asn Ala Ile
          165         170         175
Asp Ala Thr Ile Glu Leu Gly Gly Glu Asn Tyr Val Phe Trp Gly Gly
          180         185         190
Arg Glu Gly Tyr Met Ser Leu Leu Asn Thr Asp Gln Lys Arg Glu Lys
          195         200         205
Glu His Leu Ala Gln Met Leu Thr Ile Ala Arg Asp Tyr Ala Arg Ala
210         215         220

```

-continued

Arg Gly Phe Lys Gly Thr Phe Leu Ile Glu Pro Lys Pro Met Glu Pro
 225 230 235 240

Thr Lys His Gln Tyr Asp Val Asp Thr Glu Thr Val Ile Gly Phe Leu
 245 250 255

Lys Ala His Gly Leu Asp Lys Asp Phe Lys Val Asn Ile Glu Val Asn
 260 265 270

His Ala Thr Leu Ala Gly His Thr Phe Glu His Glu Leu Ala Val Ala
 275 280 285

Val Asp Asn Gly Met Leu Gly Ser Ile Asp Ala Asn Arg Gly Asp Tyr
 290 295 300

Gln Asn Gly Trp Asp Thr Asp Gln Phe Pro Ile Asp Asn Tyr Glu Leu
 305 310 315 320

Thr Gln Ala Met Met Gln Ile Ile Arg Asn Gly Gly Leu Gly Thr Gly
 325 330 335

Gly Thr Asn Phe Asp Ala Lys Thr Arg Arg Asn Ser Thr Asp Leu Glu
 340 345 350

Asp Ile Phe Ile Ala His Ile Ala Gly Met Asp Ala Met Ala Arg Ala
 355 360 365

Leu Glu Ser Ala Ala Ala Leu Leu Asp Glu Ser Pro Tyr Lys Lys Met
 370 375 380

Leu Ala Asp Arg Tyr Ala Ser Phe Asp Gly Gly Lys Gly Lys Glu Phe
 385 390 395 400

Glu Asp Gly Lys Leu Thr Leu Glu Asp Val Val Ala Tyr Ala Lys Thr
 405 410 415

Lys Gly Glu Pro Lys Gln Thr Ser Gly Lys Gln Glu Leu Tyr Glu Ala
 420 425 430

Ile Leu Asn Met Tyr Cys
 435

<210> SEQ ID NO 16
 <211> LENGTH: 1317
 <212> TYPE: DNA
 <213> ORGANISM: Bacteroides thetaiotaomicron

<400> SEQUENCE: 16

atggcaacaa aagaattttt tccgggaatt gaaaagatta aatttgaagg taaagatagt 60
 aagaacccga tggcattccg ttattacgat gcagagaagg tgattaatgg taaaaagatg 120
 aaggattggc tgagattcgc tatggcatgg tggcacacat tgtgcgctga aggtggtgat 180
 cagttcgggtg gcggaacaaa gcaattccca tggaatggta atgcagatgc tatacaggca 240
 gcaaagata agatggatgc aggatttgaa ttcatgcaga agatgggtat cgaatactat 300
 tgcttccatg acgtagactt ggtttcggaa ggtgccagtg tagaagaata cgaagctaac 360
 ctgaaagaaa tcgtagctta tgcaaacag aaacaggcag aaaccggtat caaactactg 420
 tggggactg ctaatgtatt cggtcacgcc cgctatatga acggtgcagc taccaatcct 480
 gacttcgatg tagtagctcg tgctgctggt cagatcaaaa atgcgattga tgcaacgatt 540
 gaacttggcg gagagaatta tgtgttttgg ggtggctgtg aaggctatat gtctcttctg 600
 aacacagatc agaaacgtga aaaagaacac cttgcacaga tgttgacgat tgctcgtgac 660
 tatgcccgtg cccgtggttt caaaggact ttctgatcg aaccgaaacc gatggaaccg 720
 actaaacatc aatatgacgt agatacggaa actgtaatcg gcttctgaa agctcatggt 780

-continued

```

ctggataagg atttcaaagt aatatcgag gtgaatcacg caactttggc aggtcacact 840
ttcgagcatg aattggctgt agctgtagac aatggtatgt tgggctcaat tgacgccaat 900
cgtggtgact atcagaatgg ctgggataca gaccaattcc cgatcgacaa ttatgaactg 960
actcaggcta tgatgcagat tatccgtaat ggtggtctcg gtaccggtgg tacgaacttt 1020
gatgctaaaa cccgtcgtaa ttctactgat ctggaagata tctttattgc tcacatcgca 1080
ggtatggacg ctatggcccg tgcactcgaa agtgcagcgg ctctgctcga cgaatctccc 1140
tataagaaga tgctggctga ccgttatgct tcatttgatg ggggcaaagg taaagaattt 1200
gaagacggca agctgactct ggaggatgtg gttgcttatg caaaaacaaa aggcgaaccg 1260
aacagacta gcggaagca agaactttat gaggcaattc tgaatatgta ttgctaa 1317

```

<210> SEQ ID NO 17

<211> LENGTH: 258

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 17

```

Met Ala Ala Gly Val Pro Lys Ile Asp Ala Leu Glu Ser Leu Gly Asn
1           5           10          15

Pro Leu Glu Asp Ala Lys Arg Ala Ala Tyr Arg Ala Val Asp Glu
          20          25          30

Asn Leu Lys Phe Asp Asp His Lys Ile Ile Gly Ile Gly Ser Gly Ser
          35          40          45

Thr Val Val Tyr Val Ala Glu Arg Ile Gly Gln Tyr Leu His Asp Pro
          50          55          60

Lys Phe Tyr Glu Val Ala Ser Lys Phe Ile Cys Ile Pro Thr Gly Phe
65          70          75          80

Gln Ser Arg Asn Leu Ile Leu Asp Asn Lys Leu Gln Leu Gly Ser Ile
          85          90          95

Glu Gln Tyr Pro Arg Ile Asp Ile Ala Phe Asp Gly Ala Asp Glu Val
          100         105         110

Asp Glu Asn Leu Gln Leu Ile Lys Gly Gly Gly Ala Cys Leu Phe Gln
          115         120         125

Glu Lys Leu Val Ser Thr Ser Ala Lys Thr Phe Ile Val Val Ala Asp
          130         135         140

Ser Arg Lys Lys Ser Pro Lys His Leu Gly Lys Asn Trp Arg Gln Gly
          145         150         155         160

Val Pro Ile Glu Ile Val Pro Ser Ser Tyr Val Arg Val Lys Asn Asp
          165         170         175

Leu Leu Glu Gln Leu His Ala Glu Lys Val Asp Ile Arg Gln Gly Gly
          180         185         190

Ser Ala Lys Ala Gly Pro Val Val Thr Asp Asn Asn Asn Phe Ile Ile
          195         200         205

Asp Ala Asp Phe Gly Glu Ile Ser Asp Pro Arg Lys Leu His Arg Glu
          210         215         220

Ile Lys Leu Leu Val Gly Val Val Glu Thr Gly Leu Phe Ile Asp Asn
          225         230         235         240

Ala Ser Lys Ala Tyr Phe Gly Asn Ser Asp Gly Ser Val Glu Val Thr
          245         250         255

Glu Lys

```

-continued

<210> SEQ ID NO 18
 <211> LENGTH: 2467
 <212> TYPE: DNA
 <213> ORGANISM: *Saccharomyces cerevisiae*
 <400> SEQUENCE: 18

```

ggatccaaga ccattattcc atcagaatgg aaaaaagttt aaaagatcac ggagatthttg    60
ttctttctgag cttctgctgt ccttgaaaac aaattattcc gctggccgcc ccaaacaaaa    120
acaacccccga ttaataaca ttgtcacagt attagaaatt ttctttttac aaattaccat    180
ttccagctta ctacttcta taatcctcaa tcttcagcaa gcgacgcagg gaatagccgc    240
tgaggatgcat aactgtcact tttcaattcg gccaatgcaa tctcaggcgg acgaataagg    300
gggccctctc gagaaaaaca aaaggaggat gagattagta ctttaatgtt gtgttcagta    360
attcagagac agacaagaga ggtttccaac acaatgtctt tagactcata ctatcttggg    420
tttgatcttt cgaccaaca actgaaatgt ctgcatta accaggacct aaaaattgtc    480
cattcagaaa cagtgaatt tgaaaaggat ctccgcatt atcacacaaa gaagggtgtc    540
tatatacacg gcgacactat cgaatgtccc gtagccatgt ggtagggggc tctagatctg    600
gttctctcga aatatcgca ggctaaattt ccattgaaca aagttatggc cgtctcaggg    660
tcttcccagc agcacgggtc tgtctactgg tcttccaag ccgaatctct gttagagcaa    720
ttgaataaga aaccggaaaa agatttattg cactacgtga gctctgtagc atttgcaagg    780
caaaccgccc ccaattggca agaccacagt actgcaaagc aatgtcaaga gtttgaagag    840
tgcataaggat ggccctgaaa aatggctcaa ttaacagggt ccagagccca ttttagattt    900
actggtcctc aaattctgaa aattgcacaa ttagaaccag aagcttacga aaaaacaaag    960
accatttctt tagtgtctaa ttttttgact tctatcttag tgggccatct tgttgaatta   1020
gaggaggcag atgcctgtgg tatgaacctt tatgatatac gtgaaagaaa attcatgtat   1080
gagctactac atctaattga tagttcttct aaggataaaa ctatcagaca aaaattaatg   1140
agagcaccca tgaaaaattt gatagcgggt accatctgta aatattttat tgagaagtac   1200
ggtttcaata caaactgcaa ggtctctccc atgactgggg ataatttagc cactatatgt   1260
tctttacccc tgcggaagaa tgacgttctc gtttccctag gaacaagtac tacagttctt   1320
ctggtcaccg ataagtatca cccctctccg aactatcatc ttttcattca tccaactctg   1380
ccaaaccatt atatgggtat gatttggtat tgtaatgggt ctttggcaag ggagaggata   1440
agagacgagt taaacaaaga acgggaaaat aattatgaga agactaacga ttggactctt   1500
tttaatcaag ctgtgctaga tgactcagaa agtagtgaaa atgaattagg tgtatatttt   1560
cctctggggg agatcgttcc tagcgtaaaa gccataaaca aaagggttat cttcaatcca   1620
aaaacgggta tgattgaaag agagggtggc aagttcaaag acaagaggca cgatgccaaa   1680
aatattgtag aatcacaggc ttaagttgc agggtaagaa tatctcccct gctttcggat   1740
tcaaacgcaa gctcacaaca gagactgaac gaagatacaa tcgtgaagtt tgattacgat   1800
gaatctccgc tgcgggacta cctaaataaa aggcagaaa ggactttttt tgtagggtgg   1860
gcttctaaaa acgatgctat tgtgaagaag tttgctcaag tcattggtgc taaaagggt   1920
aattttaggc tagaaacacc aaactcatgt gcccttggtg gttgttataa ggccatgtgg   1980
tcattggtat atgactctaa taaaattgca gttccttttg ataaatttct gaatgacaat   2040
  
```

-continued

```

tttccatggc atgtaatgga aagcatatcc gatgtggata atgaaaattg gatcgctata 2100
attccaagat tgtcccctta agcgaactgg aaaagactct catctaaaat atgtttgaat 2160
aatttatcat gccctgacaa gtacacacaa acacagacac ataatataca tacatatata 2220
tatatcaccg ttattatgcg tgcacatgac aatgcccttg tatgtttcgt atactgtagc 2280
aagtagtcat cattttgttc cccgttcgga aatgacaaa aagtaaaatc aataaatgaa 2340
gagtaaaaaa caatttatga aagggtgagc gaccagcaac gagagagaca aatcaaatta 2400
gcgctttcca gtgagaatat aagagagcat tgaaagagct aggttattgt taaatcatct 2460
cgagctc 2467

```

```

<210> SEQ ID NO 19
<211> LENGTH: 238
<212> TYPE: PRT
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 19

```

```

Met Val Lys Pro Ile Ile Ala Pro Ser Ile Leu Ala Ser Asp Phe Ala
1           5           10          15
Asn Leu Gly Cys Glu Cys His Lys Val Ile Asn Ala Gly Ala Asp Trp
20          25          30
Leu His Ile Asp Val Met Asp Gly His Phe Val Pro Asn Ile Thr Leu
35          40          45
Gly Gln Pro Ile Val Thr Ser Leu Arg Arg Ser Val Pro Arg Pro Gly
50          55          60
Asp Ala Ser Asn Thr Glu Lys Lys Pro Thr Ala Phe Phe Asp Cys His
65          70          75          80
Met Met Val Glu Asn Pro Glu Lys Trp Val Asp Asp Phe Ala Lys Cys
85          90          95
Gly Ala Asp Gln Phe Thr Phe His Tyr Glu Ala Thr Gln Asp Pro Leu
100         105         110
His Leu Val Lys Leu Ile Lys Ser Lys Gly Ile Lys Ala Ala Cys Ala
115        120        125
Ile Lys Pro Gly Thr Ser Val Asp Val Leu Phe Glu Leu Ala Pro His
130        135        140
Leu Asp Met Ala Leu Val Met Thr Val Glu Pro Gly Phe Gly Gly Gln
145        150        155        160
Lys Phe Met Glu Asp Met Met Pro Lys Val Glu Thr Leu Arg Ala Lys
165        170        175
Phe Pro His Leu Asn Ile Gln Val Asp Gly Gly Leu Gly Lys Glu Thr
180        185        190
Ile Pro Lys Ala Ala Lys Ala Gly Ala Asn Val Ile Val Ala Gly Thr
195        200        205
Ser Val Phe Thr Ala Ala Asp Pro His Asp Val Ile Ser Phe Met Lys
210        215        220
Glu Glu Val Ser Lys Glu Leu Arg Ser Arg Asp Leu Leu Asp
225        230        235

```

```

<210> SEQ ID NO 20
<211> LENGTH: 1328
<212> TYPE: DNA
<213> ORGANISM: Saccharomyces cerevisiae

```

```

<400> SEQUENCE: 20

```


-continued

```

gtaggcaact tacgtatctt gtatagtagg aatggctcgg tttatgtata ttaggagatc    60
aaaacgagaa aaaaatacca tatcgtatag tatagagagt ataaatataa gaaatgccgc    120
atatgtacaa ctaatctagc aaatctctag aacgcaattc cttagagact tcttctttca    180
tgaaggagat aacatcgtgc gggtcagctg cagtgaaaac actggtacca gcgacaataa    240
cgttggcacc ggctttggcg gctttcggga tggctcctt gcccaaacca ccatcgactt    300
ggatattcaa atgggggaaac ttggctctca aagtttccac ttttggcatc atgtcttcca    360
tgaatTTTTG gcctccaaac ccaggttcca cagtcataac aagagccata tccaaatgag    420
gagctagttc aaataaaacg tcaacagaag taccaggttt gatggcgcat gcagctttga    480
tgccttaga cttaatcaac ttaactaaat gcaaagggtc ttgtgtggcc tcgtagtgga    540
acgtaaattg gtcagcacca catttagcaa aatcgtcgac ccatttttca ggattttcaa    600
ccatcatgtg acaatcgaag aacgcagtgg gcttcttttc tgtgttgcta gcacgccag    660
ggcgtggcac agaacgacgt agggaggtaa caattggttg gccagagta atgtttggaa    720
caaaatggcc gtccatgaca tcgatatgta accaatctgc gccggcgttg atgaccttat    780
gacattcgca acccaagttg gcgaagtcag aagcaaggat actgggagct ataattggtt    840
tgaccatttt ttcttgtgtg tttacctcgc tcttgaatt agcaaattggc cttcttgcac    900
gaaattgtat cgagtttgct ttatttttct ttttacgggc ggattctttc tattctggct    960
ttcctataac agagatcatg aaagaagttc cagcttacgg atcaagaaag tacctataca   1020
tatacaaaaa tctgattact ttcccagctc gacttgata gctgttcttg ttttctcttg   1080
gcgacacatt ttttgttct gaagccacgt cctgctttat aagaggacat ttaaagttgc   1140
aggacttgaa tgcaattacc ggaagaagca accaaccggc atggttcagc atacaataca   1200
catttgatta gaaaagcaga gaataaatag acatgatacc tctcttttta tcctctgcag   1260
cgtattattg tttattccac gcaggcatcg gtcgttggct gttgttatgt ctcagataag   1320
cgcgtttg                                     1328

```

<210> SEQ ID NO 21

<211> LENGTH: 680

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 21

```

Met Thr Gln Phe Thr Asp Ile Asp Lys Leu Ala Val Ser Thr Ile Arg
1           5           10           15
Ile Leu Ala Val Asp Thr Val Ser Lys Ala Asn Ser Gly His Pro Gly
20           25           30
Ala Pro Leu Gly Met Ala Pro Ala Ala His Val Leu Trp Ser Gln Met
35           40           45
Arg Met Asn Pro Thr Asn Pro Asp Trp Ile Asn Arg Asp Arg Phe Val
50           55           60
Leu Ser Asn Gly His Ala Val Ala Leu Leu Tyr Ser Met Leu His Leu
65           70           75           80
Thr Gly Tyr Asp Leu Ser Ile Glu Asp Leu Lys Gln Phe Arg Gln Leu
85           90           95
Gly Ser Arg Thr Pro Gly His Pro Glu Phe Glu Leu Pro Gly Val Glu
100          105          110

```

-continued

Val	Thr	Thr	Gly	Pro	Leu	Gly	Gln	Gly	Ile	Ser	Asn	Ala	Val	Gly	Met
		115					120					125			
Ala	Met	Ala	Gln	Ala	Asn	Leu	Ala	Ala	Thr	Tyr	Asn	Lys	Pro	Gly	Phe
	130					135					140				
Thr	Leu	Ser	Asp	Asn	Tyr	Thr	Tyr	Val	Phe	Leu	Gly	Asp	Gly	Cys	Leu
145				150						155					160
Gln	Glu	Gly	Ile	Ser	Ser	Glu	Ala	Ser	Ser	Leu	Ala	Gly	His	Leu	Lys
				165					170					175	
Leu	Gly	Asn	Leu	Ile	Ala	Ile	Tyr	Asp	Asp	Asn	Lys	Ile	Thr	Ile	Asp
		180						185					190		
Gly	Ala	Thr	Ser	Ile	Ser	Phe	Asp	Glu	Asp	Val	Ala	Lys	Arg	Tyr	Glu
		195					200					205			
Ala	Tyr	Gly	Trp	Glu	Val	Leu	Tyr	Val	Glu	Asn	Gly	Asn	Glu	Asp	Leu
	210					215					220				
Ala	Gly	Ile	Ala	Lys	Ala	Ile	Ala	Gln	Ala	Lys	Leu	Ser	Lys	Asp	Lys
225				230						235					240
Pro	Thr	Leu	Ile	Lys	Met	Thr	Thr	Thr	Ile	Gly	Tyr	Gly	Ser	Leu	His
				245					250					255	
Ala	Gly	Ser	His	Ser	Val	His	Gly	Ala	Pro	Leu	Lys	Ala	Asp	Asp	Val
			260					265					270		
Lys	Gln	Leu	Lys	Ser	Lys	Phe	Gly	Phe	Asn	Pro	Asp	Lys	Ser	Phe	Val
		275					280					285			
Val	Pro	Gln	Glu	Val	Tyr	Asp	His	Tyr	Gln	Lys	Thr	Ile	Leu	Lys	Pro
	290					295					300				
Gly	Val	Glu	Ala	Asn	Asn	Lys	Trp	Asn	Lys	Leu	Phe	Ser	Glu	Tyr	Gln
305					310					315					320
Lys	Lys	Phe	Pro	Glu	Leu	Gly	Ala	Glu	Leu	Ala	Arg	Arg	Leu	Ser	Gly
				325					330					335	
Gln	Leu	Pro	Ala	Asn	Trp	Glu	Ser	Lys	Leu	Pro	Thr	Tyr	Thr	Ala	Lys
			340					345						350	
Asp	Ser	Ala	Val	Ala	Thr	Arg	Lys	Leu	Ser	Glu	Thr	Val	Leu	Glu	Asp
		355					360					365			
Val	Tyr	Asn	Gln	Leu	Pro	Glu	Leu	Ile	Gly	Gly	Ser	Ala	Asp	Leu	Thr
	370					375					380				
Pro	Ser	Asn	Leu	Thr	Arg	Trp	Lys	Glu	Ala	Leu	Asp	Phe	Gln	Pro	Pro
385					390					395					400
Ser	Ser	Gly	Ser	Gly	Asn	Tyr	Ser	Gly	Arg	Tyr	Ile	Arg	Tyr	Gly	Ile
				405					410					415	
Arg	Glu	His	Ala	Met	Gly	Ala	Ile	Met	Asn	Gly	Ile	Ser	Ala	Phe	Gly
			420					425					430		
Ala	Asn	Tyr	Lys	Pro	Tyr	Gly	Gly	Thr	Phe	Leu	Asn	Phe	Val	Ser	Tyr
		435					440					445			
Ala	Ala	Gly	Ala	Val	Arg	Leu	Ser	Ala	Leu	Ser	Gly	His	Pro	Val	Ile
	450					455					460				
Trp	Val	Ala	Thr	His	Asp	Ser	Ile	Gly	Val	Gly	Glu	Asp	Gly	Pro	Thr
465					470					475					480
His	Gln	Pro	Ile	Glu	Thr	Leu	Ala	His	Phe	Arg	Ser	Leu	Pro	Asn	Ile
				485					490					495	
Gln	Val	Trp	Arg	Pro	Ala	Asp	Gly	Asn	Glu	Val	Ser	Ala	Ala	Tyr	Lys
			500					505					510		
Asn	Ser	Leu	Glu	Ser	Lys	His	Thr	Pro	Ser	Ile	Ile	Ala	Leu	Ser	Arg

-continued

515			520			525									
Gln	Asn	Leu	Pro	Gln	Leu	Glu	Gly	Ser	Ser	Ile	Glu	Ser	Ala	Ser	Lys
530						535					540				
Gly	Gly	Tyr	Val	Leu	Gln	Asp	Val	Ala	Asn	Pro	Asp	Ile	Ile	Leu	Val
545					550					555					560
Ala	Thr	Gly	Ser	Glu	Val	Ser	Leu	Ser	Val	Glu	Ala	Ala	Lys	Thr	Leu
				565					570					575	
Ala	Ala	Lys	Asn	Ile	Lys	Ala	Arg	Val	Val	Ser	Leu	Pro	Asp	Phe	Phe
			580					585					590		
Thr	Phe	Asp	Lys	Gln	Pro	Leu	Glu	Tyr	Arg	Leu	Ser	Val	Leu	Pro	Asp
		595					600					605			
Asn	Val	Pro	Ile	Met	Ser	Val	Glu	Val	Leu	Ala	Thr	Thr	Cys	Trp	Gly
	610					615						620			
Lys	Tyr	Ala	His	Gln	Ser	Phe	Gly	Ile	Asp	Arg	Phe	Gly	Ala	Ser	Gly
625					630					635					640
Lys	Ala	Pro	Glu	Val	Phe	Lys	Phe	Phe	Gly	Phe	Thr	Pro	Glu	Gly	Val
				645					650					655	
Ala	Glu	Arg	Ala	Gln	Lys	Thr	Ile	Ala	Phe	Tyr	Lys	Gly	Asp	Lys	Leu
			660					665					670		
Ile	Ser	Pro	Leu	Lys	Lys	Ala	Phe								
	675					680									

<210> SEQ ID NO 22

<211> LENGTH: 2046

<212> TYPE: DNA

<213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 22

```

atggcacagt tctccgacat tgataaactt gcggtttcca ctttaagatt actttccggt      60
gaccaggtag aaagcgcaca atctggccac ccagggtcac cactaggatt ggcaccagtt      120
gcccatgtaa ttttcaagca actgcgctgt aaccctaaca atgaacattg gatcaataga      180
gacaggtttg ttctgtcgaa cggtcactca tgcgctcttc tgtactcaat gctccatcta      240
ttaggatacg attactctat cgaggacttg agacaattta gacaagtaaa ctcaaggaca      300
ccgggtcatc cagaattcca ctcagcggga gtggaaatca cttccggtcc gctaggccag      360
ggtatctcaa atgctgttgg tatggcaata gcgcaggcca actttgccgc cacttataac      420
gaggatggct ttcccatttc cgactcatat acgtttgcta ttgtagggga tggttgctta      480
caagagggtg tttcttcgga gacctcttcc ttagcgggac atctgcaatt gggtaacttg      540
attacgtttt atgacagtaa tagcatttcc attgacggta aaacctcgta ctcgttcgac      600
gaagatgttt tgaagcgata cgaggcatat ggttgggaag tcatggaagt cgataaagga      660
gacgacgata tggaatccat ttctagcgtt ttggaaaagg caaaactatc gaaggacaag      720
ccaaccataa tcaaggtaac tactacaatt ggatttgggt ccctacaaca gggactgct      780
ggtgttcatg ggtccgcttt gaaggcagat gatgttaaac agttgaagaa gaggtggggg      840
tttgacccaa ataatcatt ttagtagcct caagaggtgt acgattatta taagaagact      900
gttgtggaac ccggtcaaaa acttaatgag gaatgggata ggatgtttga agaatacaaa      960
accaaatttc ccgagaaggg taaagaattg caaagaagat tgaatggtga gttaccggaa     1020
ggttgggaaa agcatttacc gaagtttact ccggacgacg atgctctggc aacaagaaag     1080

```

-continued

```

acatcccagc aggtgctgac gaacatggtc caagttttgc ctgaattgat cggtggttct 1140
gccgatttga caccttcgaa tctgacaagg tgggaaggcg cggtagattt ccaacctccc 1200
attacccaac taggtaacta tgcaggaagg tacattagat acggtgtgag ggaacacgga 1260
atgggtgcca ttatgaacgg tatctctgcc tttggtgcaa actacaagcc ttacggtggt 1320
acctttttga acttcgtctc ttatgctgca ggagccgta ggtagccgc cttgtctggt 1380
aatccagtca tttgggttgc aacacatgac tctatcgggc ttggtgagga tggccaacg 1440
caccaacctt ttgaaactct ggctcacttg agggctattc caaacatgca tgtatggaga 1500
cctgctgatg gtaacgaaac ttctgctgcg tattattctg ctatcaaadc tggcgaaca 1560
ccatctggtt tggctttatc acgacagaat cttcctcaat tggagcattc ctcttttgaa 1620
aaagccttga aggggtggcta tgtgatccat gacgtggaga atcctgatat taccctggtg 1680
tcaacaggat cagaagtctc catttctata gatgcagcca aaaaattgta cgatactaaa 1740
aaaatcaaag caagagttgt ttccctgcca gacttttata cttttgacag gcaaagtgaa 1800
gaatacagat tctctgttct accagacggg gtccgatca tgccttttga agtattggct 1860
acttcaagct ggggtaagta tgctcatcaa tcgttcggac tgcacgaatt tggcgttca 1920
ggcaaggggc ctgaaattta caaattgttc gatttcacag cggacggtgt tgcgtcaagg 1980
gctgaaaaga caatcaatta ctacaaagga aagcagttgc tttctcctat gggaagagct 2040
ttctaa 2046

```

<210> SEQ ID NO 23

<211> LENGTH: 335

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 23

```

Met Ser Glu Pro Ala Gln Lys Lys Gln Lys Val Ala Asn Asn Ser Leu
1           5           10           15
Glu Gln Leu Lys Ala Ser Gly Thr Val Val Val Ala Asp Thr Gly Asp
20           25           30
Phe Gly Ser Ile Ala Lys Phe Gln Pro Gln Asp Ser Thr Thr Asn Pro
35           40           45
Ser Leu Ile Leu Ala Ala Ala Lys Gln Pro Thr Tyr Ala Lys Leu Ile
50           55           60
Asp Val Ala Val Glu Tyr Gly Lys Lys His Gly Lys Thr Thr Glu Glu
65           70           75           80
Gln Val Glu Asn Ala Val Asp Arg Leu Leu Val Glu Phe Gly Lys Glu
85           90           95
Ile Leu Lys Ile Val Pro Gly Arg Val Ser Thr Glu Val Asp Ala Arg
100          105          110
Leu Ser Phe Asp Thr Gln Ala Thr Ile Glu Lys Ala Arg His Ile Ile
115          120          125
Lys Leu Phe Glu Gln Glu Gly Val Ser Lys Glu Arg Val Leu Ile Lys
130          135          140
Ile Ala Ser Thr Trp Glu Gly Ile Gln Ala Ala Lys Glu Leu Glu Glu
145          150          155          160
Lys Asp Gly Ile His Cys Asn Leu Thr Leu Leu Phe Ser Phe Val Gln
165          170          175
Ala Val Ala Cys Ala Glu Ala Gln Val Thr Leu Ile Ser Pro Phe Val

```

-continued

180										185					190					
Gly	Arg	Ile	Leu	Asp	Trp	Tyr	Lys	Ser	Ser	Thr	Gly	Lys	Asp	Tyr	Lys					
		195					200					205								
Gly	Glu	Ala	Asp	Pro	Gly	Val	Ile	Ser	Val	Lys	Lys	Ile	Tyr	Asn	Tyr					
	210					215					220									
Tyr	Lys	Lys	Tyr	Gly	Tyr	Lys	Thr	Ile	Val	Met	Gly	Ala	Ser	Phe	Arg					
225					230					235					240					
Ser	Thr	Asp	Glu	Ile	Lys	Asn	Leu	Ala	Gly	Val	Asp	Tyr	Leu	Thr	Ile					
			245						250					255						
Ser	Pro	Ala	Leu	Leu	Asp	Lys	Leu	Met	Asn	Ser	Thr	Glu	Pro	Phe	Pro					
			260					265					270							
Arg	Val	Leu	Asp	Pro	Val	Ser	Ala	Lys	Lys	Glu	Ala	Gly	Asp	Lys	Ile					
	275						280					285								
Ser	Tyr	Ile	Ser	Asp	Glu	Ser	Lys	Phe	Arg	Phe	Asp	Leu	Asn	Glu	Asp					
	290					295					300									
Ala	Met	Ala	Thr	Glu	Lys	Leu	Ser	Glu	Gly	Ile	Arg	Lys	Phe	Ser	Ala					
305					310					315					320					
Asp	Ile	Val	Thr	Leu	Phe	Asp	Leu	Ile	Glu	Lys	Lys	Val	Thr	Ala						
				325					330					335						

<210> SEQ ID NO 24
 <211> LENGTH: 2046
 <212> TYPE: DNA
 <213> ORGANISM: Saccharomyces cerevisiae

<400> SEQUENCE: 24

```

atggcacagt tctccgacat tgataaactt gcggtttcca ctttaagatt actttccgtt    60
gaccaggtgg aaagcgaca atctggccac ccaggtgcac cactaggatt ggcaccagtt    120
gcccatgtaa ttttcaagca actgcgctgt aaccctaaca atgaacattg gatcaataga    180
gacaggtttg ttctgtcgaa cggtcactca tgcgctcttc tgtactcaat gctccatcta    240
ttaggatacg attactctat cgaggacttg agacaattta gacaagtaa ctcaaggaca    300
ccgggtcacc cagaattcca ctcagcggga gtggaaatca cttccggtec gctaggccag    360
ggtatctcaa atgctgttgg tatggcaata ggcagggcca actttgccgc cacttataac    420
gaggatggct ttcccatttc cgactcatat acgtttgcta ttgtagggga tggttgctta    480
caagaggggtg tttcttcgga gacctcttcc ttagcgggac atctgcaatt gggtaacttg    540
attacgtttt atgacagtaa tagcatttcc attgacggta aaacctcgta ctcgttcgac    600
gaagatgttt tgaagcgata cgaggcatat ggttgggaag tcatggaagt cgataaagga    660
gacgacgata tggaatccat ttctagcgtt ttggaaaagg caaaactatc gaaggacaag    720
ccaaccataa tcaaggtaac tactacaatt ggatttgggt ccctacaaca gggtagtctg    780
ggtgttcatg ggtccgcttt gaaggcagat gatgttaaag agttgaagaa gaggtggggg    840
tttgacccaa ataatcatt tgtagtacct caagaggtgt acgattatta taagaagact    900
gttgtggaac ccggtcaaaa acttaatgag gaatgggata ggatgtttga agaatacaaa    960
accaaatttc ccgagaaggg taaagaattg caaagaagat tgaatggtga gttaccggaa   1020
ggttgggaaa agcatttacc gaagtttact ccggacgacg atgctctggc aacaagaaag   1080
acatcccagc aggtgctgac gaacatggtc caagttttgc ctgaattgat cggtggttct   1140
gccgatttga caccttcgaa tctgacaagg tgggaaggcg cggtagattt ccaacctccc   1200
    
```

-continued

```

attacccaac taggtaacta tgcaggaagg tacattagat acggtgtgag ggaacacgga 1260
atgggtgcca ttatgaacgg tatctctgcc tttggtgcaa actacaagcc ttacggtggt 1320
acctttttga acttcgtctc ttatgctgca ggagccgta ggtagccgc cttgtctggt 1380
aatccagtca tttgggttgc aacacatgac tctatcgggc ttggtgagga tggccaacg 1440
caccaaccta ttgaaactct ggctcacttg agggctattc caaacatgca tgtatggaga 1500
cctgctgatg gtaacgaaac ttctgctgcg tattattctg ctatcaaac tggtcgaaca 1560
ccatctggtg tggctttatc acgacagaat cttcctcaat tggagcattc ctcttttgaa 1620
aaagccttga aggggtggta tgtgatccat gacgtggaga atcctgatat tatcctggtg 1680
tcaacaggat cagaagtctc catttctata gatgcagcca aaaaattgta cgatactaaa 1740
aaaatcaaag caagagttgt ttcctgcca gacttttata cttttgacag gcaaagtgaa 1800
gaatacagat tctctgttct accagacggg gttccgatca tgtcctttga agtattggct 1860
acttcaagct ggggtaagta tgctcatcaa tcgttcggac tcgacgaatt tggtcggtca 1920
ggcaaggggc ctgaaattta caaattgttc gatttcacag cggacgggtg tgcgtcaagg 1980
gctgaaaaga caatcaatta ctacaaagga aagcagttgc tttctcctat gggaagagct 2040
ttctaa 2046

```

<210> SEQ ID NO 25

<211> LENGTH: 600

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 25

```

Met Leu Cys Ser Val Ile Gln Arg Gln Thr Arg Glu Val Ser Asn Thr
1           5           10          15
Met Ser Leu Asp Ser Tyr Tyr Leu Gly Phe Asp Leu Ser Thr Gln Gln
20          25          30
Leu Lys Cys Leu Ala Ile Asn Gln Asp Leu Lys Ile Val His Ser Glu
35          40          45
Thr Val Glu Phe Glu Lys Asp Leu Pro His Tyr His Thr Lys Lys Gly
50          55          60
Val Tyr Ile His Gly Asp Thr Ile Glu Cys Pro Val Ala Met Trp Leu
65          70          75          80
Glu Ala Leu Asp Leu Val Leu Ser Lys Tyr Arg Glu Ala Lys Phe Pro
85          90          95
Leu Asn Lys Val Met Ala Val Ser Gly Ser Cys Gln Gln His Gly Ser
100         105         110
Val Tyr Trp Ser Ser Gln Ala Glu Ser Leu Leu Glu Gln Leu Asn Lys
115        120        125
Lys Pro Glu Lys Asp Leu Leu His Tyr Val Ser Ser Val Ala Phe Ala
130        135        140
Arg Gln Thr Ala Pro Asn Trp Gln Asp His Ser Thr Ala Lys Gln Cys
145        150        155        160
Gln Glu Phe Glu Glu Cys Ile Gly Gly Pro Glu Lys Met Ala Gln Leu
165        170        175
Thr Gly Ser Arg Ala His Phe Arg Phe Thr Gly Pro Gln Ile Leu Lys
180        185        190
Ile Ala Gln Leu Glu Pro Glu Ala Tyr Glu Lys Thr Lys Thr Ile Ser

```

-continued

195					200					205					
Leu	Val	Ser	Asn	Phe	Leu	Thr	Ser	Ile	Leu	Val	Gly	His	Leu	Val	Glu
210					215					220					
Leu	Glu	Glu	Ala	Asp	Ala	Cys	Gly	Met	Asn	Leu	Tyr	Asp	Ile	Arg	Glu
225					230					235					240
Arg	Lys	Phe	Ser	Asp	Glu	Leu	Leu	His	Leu	Ile	Asp	Ser	Ser	Ser	Lys
				245					250					255	
Asp	Lys	Thr	Ile	Arg	Gln	Lys	Leu	Met	Arg	Ala	Pro	Met	Lys	Asn	Leu
			260					265					270		
Ile	Ala	Gly	Thr	Ile	Cys	Lys	Tyr	Phe	Ile	Glu	Lys	Tyr	Gly	Phe	Asn
		275					280					285			
Thr	Asn	Cys	Lys	Val	Ser	Pro	Met	Thr	Gly	Asp	Asn	Leu	Ala	Thr	Ile
	290					295					300				
Cys	Ser	Leu	Pro	Leu	Arg	Lys	Asn	Asp	Val	Leu	Val	Ser	Leu	Gly	Thr
305					310					315					320
Ser	Thr	Thr	Val	Leu	Leu	Val	Thr	Asp	Lys	Tyr	His	Pro	Ser	Pro	Asn
				325					330					335	
Tyr	His	Leu	Phe	Ile	His	Pro	Thr	Leu	Pro	Asn	His	Tyr	Met	Gly	Met
			340					345					350		
Ile	Cys	Tyr	Cys	Asn	Gly	Ser	Leu	Ala	Arg	Glu	Arg	Ile	Arg	Asp	Glu
		355					360					365			
Leu	Asn	Lys	Glu	Arg	Glu	Asn	Asn	Tyr	Glu	Lys	Thr	Asn	Asp	Trp	Thr
	370					375					380				
Leu	Phe	Asn	Gln	Ala	Val	Leu	Asp	Asp	Ser	Glu	Ser	Ser	Glu	Asn	Glu
385					390					395					400
Leu	Gly	Val	Tyr	Phe	Pro	Leu	Gly	Glu	Ile	Val	Pro	Ser	Val	Lys	Ala
				405					410					415	
Ile	Asn	Lys	Arg	Val	Ile	Phe	Asn	Pro	Lys	Thr	Gly	Met	Ile	Glu	Arg
			420					425					430		
Glu	Val	Ala	Lys	Phe	Lys	Asp	Lys	Arg	His	Asp	Ala	Lys	Asn	Ile	Val
		435					440					445			
Glu	Ser	Gln	Ala	Leu	Ser	Cys	Arg	Val	Arg	Ile	Ser	Pro	Leu	Leu	Ser
	450					455					460				
Asp	Ser	Asn	Ala	Ser	Ser	Gln	Gln	Arg	Leu	Asn	Glu	Asp	Thr	Ile	Val
465					470					475					480
Lys	Phe	Asp	Tyr	Asp	Glu	Ser	Pro	Leu	Arg	Asp	Tyr	Leu	Asn	Lys	Arg
			485						490					495	
Pro	Glu	Arg	Thr	Phe	Phe	Val	Gly	Gly	Ala	Ser	Lys	Asn	Asp	Ala	Ile
			500					505					510		
Val	Lys	Lys	Phe	Ala	Gln	Val	Ile	Gly	Ala	Thr	Lys	Gly	Asn	Phe	Arg
		515					520					525			
Leu	Glu	Thr	Pro	Asn	Ser	Cys	Ala	Leu	Gly	Gly	Cys	Tyr	Lys	Ala	Met
	530					535					540				
Trp	Ser	Leu	Leu	Tyr	Asp	Ser	Asn	Lys	Ile	Ala	Val	Pro	Phe	Asp	Lys
545					550					555					560
Phe	Leu	Asn	Asp	Asn	Phe	Pro	Trp	His	Val	Met	Glu	Ser	Ile	Ser	Asp
				565					570					575	
Val	Asp	Asn	Glu	Asn	Trp	Asp	Arg	Tyr	Asn	Ser	Lys	Ile	Val	Pro	Leu
			580					585					590		
Ser	Glu	Leu	Glu	Lys	Thr	Leu	Ile								
	595					600									

-continued

<210> SEQ ID NO 26
<211> LENGTH: 2467
<212> TYPE: DNA
<213> ORGANISM: *Saccharomyces cerevisiae*
<400> SEQUENCE: 26

ggatccaaga ccattattcc atcagaatgg aaaaagttt aaaagatcac ggagattttg 60
ttcttctgag cttctgctgt ccttgaaaac aaattattcc gctggccgcc ccaaacaaaa 120
acaaccccgga ttaataaca ttgtcacagt attagaaatt ttctttttac aaattacat 180
ttccagctta ctacttcta taatcctcaa tcttcagcaa gcgacgcagg gaatagccgc 240
tgaggtgcat aactgtcact tttcaattcg gccaatgcaa tctcaggcgg acgaataagg 300
gggccctctc gagaaaaaca aaaggaggat gagattagta ctttaatgtt gtgttcagta 360
attcagagac agacaagaga ggtttccaac acaatgtctt tagactcata ctatcttggg 420
tttgatcttt cgaccaaca actgaaatgt ctcgccatta accaggacct aaaaattgtc 480
cattcagaaa cagtggaatt tgaaaaggat cttccgcatt atcacacaaa gaaggggtgc 540
tatatacacg gcgacactat cgaatgtccc gtagccatgt ggtaggggc tctagatctg 600
gttctctcga aatatcgga ggctaaatth ccattgaaca aagttatggc cgtctcaggg 660
tcttgccagc agcacgggtc tgtctactgg tcttccaag ccgaatctct gttagagcaa 720
ttgaataaga aaccggaaaa agatttattg cactacgtga gctctgtagc atttgcaagg 780
caaaccgccc ccaattggca agaccacagt actgcaagc aatgtcaaga gtttgaagag 840
tgcataagtg ggctgaaaa aatggctcaa ttaacagggt ccagagccca ttttagattt 900
actggtcctc aaattctgaa aattgcacaa ttagaaccag aagcttacga aaaaacaaag 960
accatttctt tagtgtctaa ttttttgact tctatcttag tgggcatct tgttgaatta 1020
gaggaggcag atgcctgtgg tatgaacctt tatgatatac gtgaaagaaa attcatgtat 1080
gagctactac atctaattga tagttcttct aaggataaaa ctatcagaca aaaattaatg 1140
agagcaccca tgaaaaatth gatagcgggt accatctgta aatattttat tgagaagtac 1200
ggtttcaata caaactgcaa ggtctctccc atgactgggg ataatttagc cactatatgt 1260
tctttacccc tgcggaagaa tgacgttctc gtttccctag gaacaagtac tacagttctt 1320
ctggtcaccg ataagtatca cccctctccg aactatcatc ttttcattca tccaactctg 1380
ccaaaccatt atatgggtat gatttgttat tgtaatgggt ctttggaag ggagaggata 1440
agagacgagt taaacaaaga acgggaaaat aattatgaga agactaacga ttggactctt 1500
tttaatcaag ctgtgctaga tgactcagaa agtagtgaaa atgaattagg tgtatatttt 1560
cctctggggg agatcgttcc tagcgtaaaa gccataaaca aaagggttat cttcaatcca 1620
aaaacgggta tgattgaaag agaggtggcc aagttcaaag acaagaggca cgatgccaaa 1680
aatattgtag aatcacaggc ttttaagttgc agggtaagaa tatctcccct gctttcggat 1740
tcaaacgcaa gctcacaaca gagactgaac gaagatacaa tcgtgaagtt tgattacgat 1800
gaatctccgc tgcgggacta cctaaataaa aggccagaaa ggactttttt tgtaggtggg 1860
gcttctaaaa acgatgctat tgtgaagaag tttgctcaag tcattggtgc taaaagggt 1920
aattttaggc tagaaacacc aaactcatgt gcccttggtg gttgttataa ggccatgtgg 1980
tcattgttat atgactctaa taaaattgca gttccttttg ataaatttct gaatgacaat 2040

-continued

```

tttccatggc atgtaatgga aagcatatcc gatgtggata atgaaaattg gatcgctata 2100
attccaagat tgtcccctta agcgaactgg aaaagactct catctaaaat atgtttgaat 2160
aatttatcat gccctgacaa gtacacacaa acacagacac ataatataca tacatatata 2220
tatatcaccg ttattatgcg tgcacatgac aatgcccttg tatgtttcgt atactgtagc 2280
aagtagtcat cattttgttc cccgttcgga aatgacaaa aagtaaaatc aataaatgaa 2340
gagtaaaaaa caatttatga aagggtgagc gaccagcaac gagagagaca aatcaaatta 2400
gcgctttcca gtgagaatat aagagagcat tgaagagct aggttattgt taaatcatct 2460
cgagctc 2467

```

<210> SEQ ID NO 27

<211> LENGTH: 494

<212> TYPE: PRT

<213> ORGANISM: Piromyces species

<400> SEQUENCE: 27

```

Met Lys Thr Val Ala Gly Ile Asp Leu Gly Thr Gln Ser Met Lys Val
1           5           10           15
Val Ile Tyr Asp Tyr Glu Lys Lys Glu Ile Ile Glu Ser Ala Ser Cys
                20           25           30
Pro Met Glu Leu Ile Ser Glu Ser Asp Gly Thr Arg Glu Gln Thr Thr
        35           40           45
Glu Trp Phe Asp Lys Gly Leu Glu Val Cys Phe Gly Lys Leu Ser Ala
        50           55           60
Asp Asn Lys Lys Thr Ile Glu Ala Ile Gly Ile Ser Gly Gln Leu His
65           70           75           80
Gly Phe Val Pro Leu Asp Ala Asn Gly Lys Ala Leu Tyr Asn Ile Lys
                85           90           95
Leu Trp Cys Asp Thr Ala Thr Val Glu Glu Cys Lys Ile Ile Thr Asp
        100          105          110
Ala Ala Gly Gly Asp Lys Ala Val Ile Asp Ala Leu Gly Asn Leu Met
        115          120          125
Leu Thr Gly Phe Thr Ala Pro Lys Ile Leu Trp Leu Lys Arg Asn Lys
        130          135          140
Pro Glu Ala Phe Ala Asn Leu Lys Tyr Ile Met Leu Pro His Asp Tyr
145          150          155          160
Leu Asn Trp Lys Leu Thr Gly Asp Tyr Val Met Glu Tyr Gly Asp Ala
        165          170          175
Ser Gly Thr Ala Leu Phe Asp Ser Lys Asn Arg Cys Trp Ser Lys Lys
        180          185          190
Ile Cys Asp Ile Ile Asp Pro Lys Leu Leu Asp Leu Leu Pro Lys Leu
        195          200          205
Ile Glu Pro Ser Ala Pro Ala Gly Lys Val Asn Asp Glu Ala Ala Lys
210          215          220
Ala Tyr Gly Ile Pro Ala Gly Ile Pro Val Ser Ala Gly Gly Gly Asp
225          230          235          240
Asn Met Met Gly Ala Val Gly Thr Gly Thr Val Ala Asp Gly Phe Leu
        245          250          255
Thr Met Ser Met Gly Thr Ser Gly Thr Leu Tyr Gly Tyr Ser Asp Lys
        260          265          270

```

-continued

Pro Ile Ser Asp Pro Ala Asn Gly Leu Ser Gly Phe Cys Ser Ser Thr
 275 280 285

Gly Gly Trp Leu Pro Leu Leu Cys Thr Met Asn Cys Thr Val Ala Thr
 290 295 300

Glu Phe Val Arg Asn Leu Phe Gln Met Asp Ile Lys Glu Leu Asn Val
 305 310 315 320

Glu Ala Ala Lys Ser Pro Cys Gly Ser Glu Gly Val Leu Val Ile Pro
 325 330 335

Phe Phe Asn Gly Glu Arg Thr Pro Asn Leu Pro Asn Gly Arg Ala Ser
 340 345 350

Ile Thr Gly Leu Thr Ser Ala Asn Thr Ser Arg Ala Asn Ile Ala Arg
 355 360 365

Ala Ser Phe Glu Ser Ala Val Phe Ala Met Arg Gly Gly Leu Asp Ala
 370 375 380

Phe Arg Lys Leu Gly Phe Gln Pro Lys Glu Ile Arg Leu Ile Gly Gly
 385 390 395 400

Gly Ser Lys Ser Asp Leu Trp Arg Gln Ile Ala Ala Asp Ile Met Asn
 405 410 415

Leu Pro Ile Arg Val Pro Leu Leu Glu Glu Ala Ala Ala Leu Gly Gly
 420 425 430

Ala Val Gln Ala Leu Trp Cys Leu Lys Asn Gln Ser Gly Lys Cys Asp
 435 440 445

Ile Val Glu Leu Cys Lys Glu His Ile Lys Ile Asp Glu Ser Lys Asn
 450 455 460

Ala Asn Pro Ile Ala Glu Asn Val Ala Val Tyr Asp Lys Ala Tyr Asp
 465 470 475 480

Glu Tyr Cys Lys Val Val Asn Thr Leu Ser Pro Leu Tyr Ala
 485 490

<210> SEQ ID NO 28
 <211> LENGTH: 2041
 <212> TYPE: DNA
 <213> ORGANISM: Piromyces sp.

<400> SEQUENCE: 28

attatataaa ataactttaa ataaaacaat ttttatttgt ttatttaatt attcaaaaaa 60
 aattaaagta aaagaaaaat aatacagtag aacaatagta ataatatcaa aatgaagact 120
 gttgctggta ttgatcttgg aactcaaagt atgaaagtcg ttatttacga ctatgaaaag 180
 aaagaaatta ttgaaagtgc tagctgtcca atggaattga tttccgaaag tgacgggtacc 240
 cgtgaacaaa ccactgaatg gtttgacaag ggtcttgaag tttgttttgg taagcttagt 300
 gctgataaca aaaagactat tgaagctatt ggtatttctg gtcaattaca cggttttggt 360
 cctcttgatg ctaacggtaa ggctttatac aacatcaaac tttggtgtga tactgctacc 420
 gttgaagaat gtaagattat cactgatgct gccggtggtyg acaaggctgt tattgatgcc 480
 cttggtaacc ttatgctcac cggtttcacc gctccaaaga tcctctggct caagcgcaac 540
 aagccagaag ctttcgctaa cttaaagtac attatgcttc cacacgatta cttaaactgg 600
 aagcttactg gtgattacgt tatggaatac ggtgatgcct ctggtaccgc tctcttcgat 660
 tctaagaacc gttgctggtc taagaagatt tgcgatatca ttgacccaaa acttttagat 720
 ttacttccaa agttaattga accaagcgct ccagctggta aggttaatga tgaagccgct 780

-continued

```

aaggcttacg gtattccagc cggattcca gttccgctg gtgggtgga taacatgatg 840
gggctggtg gtactggtag tggtagtgat ggttccctta ccatgtctat gggacttct 900
ggtagctttt acggttacag tgacaagcca attagtacc cagctaagtg ttaagtgg 960
ttctgttctt ctactggtag atggctcca ttactttgta ctatgaactg tactgttgcc 1020
actgaattcg ttcgtaacct cttccaaatg gatattaagg aacttaatgt tgaagctgcc 1080
aagtctccat gtggtagtga aggtggttta gttattccat tcttcaatgg tgaagaact 1140
ccaaacttac caaacggctg tgctagtatt actggtctta cttctgctaa caccagccgt 1200
gtaacattg ctcgtgctag tttcgaatcc gccgtttctg ctatgcctgg tggtttagat 1260
gctttccgta agttaggttt ccaaccaaag gaaattcgtc ttattggtgg tggttctaag 1320
tctgatctct ggagacaaat tgccgctgat atcatgaacc ttccaatcag agttccactt 1380
ttagaagaag ctgctgctct tgggtggtgct gttcaagctt tatggtgtct taagaaccaa 1440
tctggtaagt gtgatattgt tgaactttgc aaagaacaca ttaagattga tgaatctaag 1500
aatgctaacc caattgccga aaatggtgct gtttacgaca aggettacga tgaatactgc 1560
aaggttgtaa atactctttc tccattatat gcttaaattg ccaatgtaa aaaaaatata 1620
atgccatata attgccttgt caatacactg ttcatgttca tataatcata ggacattgaa 1680
tttacaaggt ttatacaatt aatatctatt atcatattat tatacagcat ttcattttct 1740
aagattagac gaaacaattc ttggttcctt gcaatataca aaatttacat gaatttttag 1800
aatagtctcg tatttatgcc caataatcag gaaaattacc taatgctgga ttcttgtaa 1860
taaaaacaaa ataaataaat taaataaaca aataaaaatt ataagtaaat ataaatata 1920
aagtaatata aaaaaaaagt aaataaataa ataaataaat aaaaattttt tgcaaatata 1980
taaataaata aataaaatat aaaaataatt tagcaataa attaaaaaaa aaaaaaaaaa 2040
a 2041

```

<210> SEQ ID NO 29

<211> LENGTH: 327

<212> TYPE: PRT

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 29

```

Met Ser Ser Leu Val Thr Leu Asn Asn Gly Leu Lys Met Pro Leu Val
1           5           10           15

Gly Leu Gly Cys Trp Lys Ile Asp Lys Lys Val Cys Ala Asn Gln Ile
20           25           30

Tyr Glu Ala Ile Lys Leu Gly Tyr Arg Leu Phe Asp Gly Ala Cys Asp
35           40           45

Tyr Gly Asn Glu Lys Glu Val Gly Glu Gly Ile Arg Lys Ala Ile Ser
50           55           60

Glu Gly Leu Val Ser Arg Lys Asp Ile Phe Val Val Ser Lys Leu Trp
65           70           75           80

Asn Asn Phe His His Pro Asp His Val Lys Leu Ala Leu Lys Lys Thr
85           90           95

Leu Ser Asp Met Gly Leu Asp Tyr Leu Asp Leu Tyr Tyr Ile His Phe
100          105          110

Pro Ile Ala Phe Lys Tyr Val Pro Phe Glu Glu Lys Tyr Pro Pro Gly
115          120          125

```

-continued

Phe Tyr Thr Gly Ala Asp Asp Glu Lys Lys Gly His Ile Thr Glu Ala
 130 135 140
 His Val Pro Ile Ile Asp Thr Tyr Arg Ala Leu Glu Glu Cys Val Asp
 145 150 155 160
 Glu Gly Leu Ile Lys Ser Ile Gly Val Ser Asn Phe Gln Gly Ser Leu
 165 170 175
 Ile Gln Asp Leu Leu Arg Gly Cys Arg Ile Lys Pro Val Ala Leu Gln
 180 185 190
 Ile Glu His His Pro Tyr Leu Thr Gln Glu His Leu Val Glu Phe Cys
 195 200 205
 Lys Leu His Asp Ile Gln Val Val Ala Tyr Ser Ser Phe Gly Pro Gln
 210 215 220
 Ser Phe Ile Glu Met Asp Leu Gln Leu Ala Lys Thr Thr Pro Thr Leu
 225 230 235 240
 Phe Glu Asn Asp Val Ile Lys Lys Val Ser Gln Asn His Pro Gly Ser
 245 250 255
 Thr Thr Ser Gln Val Leu Leu Arg Trp Ala Thr Gln Arg Gly Ile Ala
 260 265 270
 Val Ile Pro Lys Ser Ser Lys Lys Glu Arg Leu Leu Gly Asn Leu Glu
 275 280 285
 Ile Glu Lys Lys Phe Thr Leu Thr Glu Gln Glu Leu Lys Asp Ile Ser
 290 295 300
 Ala Leu Asn Ala Asn Ile Arg Phe Asn Asp Pro Trp Thr Trp Leu Asp
 305 310 315 320
 Gly Lys Phe Pro Thr Phe Ala
 325

<210> SEQ ID NO 30

<211> LENGTH: 984

<212> TYPE: DNA

<213> ORGANISM: *Saccharomyces cerevisiae*

<400> SEQUENCE: 30

```

atgtcttcac tggttactct taataacggt ctgaaaatgc ccctagtcgg cttaggggtgc      60
tggaaaattg acaaaaaagt ctgtgccaat caaatattatg aagctatcaa attaggctac      120
cgtttattcg atggtgcttg cgactacggc aacgaaaagg aagttggtga aggtatcagg      180
aaagccatct ccgaaggtct tgtttctaga aaggatatat ttgttgtttc aaagttatgg      240
aacaattttc accatcctga tcatgtaaaa ttagctttaa agaagacctt aagcgatatg      300
ggacttgatt atttagacct gtattatatt cacttcccaa tcgccttcaa atatggtcca      360
tttgaagaga aataccctcc aggattctat acgggcgcag atgacgagaa gaaaggctcac      420
atcaccgaag cacatgtacc aatcatagat acgtaccggg ctctggaaga atgtggtgat      480
gaaggcttga ttaagtctat tgggtgttcc aactttcagg gaagcttgat tcaagattta      540
ttacgtgggt gtagaatcaa gcccggtggc ttgcaaattg aacaccatcc ttatttgact      600
caagaacacc tagttgagtt ttgtaaatta cacgatatcc aagtagttgc ttactcctcc      660
ttcggtcctc aatcattcat tgagatggac ttacagttgg caaaaaccac gccaaactctg      720
ttcgagaatg atgtaatcaa gaaggtctca caaaaccatc caggcagtac cacttcccaa      780
gtattgctta gatgggcaac tcagagaggc attgccgtca ttccaaaatc ttccaagaag      840
gaaaggttac ttggcaacct agaaatcgaa aaaaagttca ctttaacgga gcaagaattg      900

```

-continued

aaggatattt ctgcactaaa tgccaacatc agatttaatg atccatggac ctggttggat 960

ggtaaattcc ccacttttgc ctga 984

<210> SEQ ID NO 31
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 31

gactagtcga gtttatcatt atcaatactg c 31

<210> SEQ ID NO 32
 <211> LENGTH: 49
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 32

ctcataatca ggtactgata acattttggt tgtttatgtg tgtttattc 49

<210> SEQ ID NO 33
 <211> LENGTH: 49
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 33

gaataaacac acataaaca acaaaatggt atcagtacct gattatgag 49

<210> SEQ ID NO 34
 <211> LENGTH: 48
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 34

aatcataaat cataagaaat tcgcttactt taagaatgcc ttagtcat 48

<210> SEQ ID NO 35
 <211> LENGTH: 48
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 35

atgactaagg cattcttaaa gtaagcgaat ttcttatgat ttatgatt 48

<210> SEQ ID NO 36
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: primer

<400> SEQUENCE: 36

cactagtctc gagtgtggaa gaacgattac aacagg 36

-continued

<210> SEQ ID NO 37
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 37

cgagctcgtg ggtgtattgg attataggaa g 31

<210> SEQ ID NO 38
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 38

ttgggctggtt tcaactaaat tcatttttag gctggatatct tgattcta 48

<210> SEQ ID NO 39
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 39

tagaatcaag ataccagcct aaaaatgaat ttagttgaaa cagcccaa 48

<210> SEQ ID NO 40
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 40

aatcataaat cataagaaat tcgctctaatt atttgattgc ttgccag 48

<210> SEQ ID NO 41
<211> LENGTH: 48
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 41

ctgggcaagc aatcaaatat tagagcgaat ttcttatgat ttatgatt 48

<210> SEQ ID NO 42
<211> LENGTH: 31
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 42

tgagctcgtg tggaagaacg attacaacag g 31

<210> SEQ ID NO 43
<211> LENGTH: 28

-continued

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 43

acgcgtcgac tcgtaggaac aatttcgg 28

<210> SEQ ID NO 44
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 44

cttcttggtt taatgcttct agcatttttt gattaaaatt aaaaaaactt 50

<210> SEQ ID NO 45
<211> LENGTH: 50
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 45

aagttttttt aattttaatc aaaaaatgct agaagcatta aaacaagaag 50

<210> SEQ ID NO 46
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 46

ggtatatatt taagagcgat ttgtttactt gcgaactgca tgatcc 46

<210> SEQ ID NO 47
<211> LENGTH: 46
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 47

ggatcatgca gttcgcaagt aaacaaatcg ctcttaaata tatacc 46

<210> SEQ ID NO 48
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: primer

<400> SEQUENCE: 48

cgcagtcgac cttttaaaca gttgatgaga acc 33

<210> SEQ ID NO 49
<211> LENGTH: 676
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: promoter

-continued

<400> SEQUENCE: 49

```

tcgagtttat cattatcaat actgccattt caaagaatac gtaaataatt aatagtagtg      60
atthtcttaa ctttatttag tcaaaaaatt agccttttaa ttctgctgta acccgtacat     120
gccccaaata gggggcgggt tacacagaat atataacatc gtaggtgtct ggggtaacag     180
tttattcctg gcatccacta aatataatgg agcccgtttt ttaagctggc atccagaaaa     240
aaaaagaatc ccagcaccaa aatattgttt tcttcaccaa ccatcagttc ataggtccat     300
tctcttagcg caactacaga gaacaggggc acaaacaggc aaaaaacggg cacaacctca     360
atggagtgat gcaacctgcc tggagtaa at gatgacacaa ggcaattgac ccacgcatgt     420
atctatctca ttttcttaca ctttctatta ctttctgctc tctctgattt ggaaaaagct     480
gaaaaaaaaa gttgaaacca gttccctgaa attattcccc tacttgacta ataagtatat     540
aaagacggta ggtattgatt gtaattctgt aaatctattt cttaaacttc ttaaattcta     600
cttttatagt tagtcttttt tttagtttta aaacaccaag aacttagttt cgaataaaca     660
cacataaaca aacaaa                                     676

```

<210> SEQ ID NO 50

<211> LENGTH: 326

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: terminator

<400> SEQUENCE: 50

```

gcaatttct tatgatttat gatttttatt attaaataag ttataaaaa aataagtgta      60
taciaatttt aaagtgactc ttaggtttta aaacgaaaat tcttattctt gagtaactct     120
ttcctgtagg tcaggttctt ttctcaggta tagcatgagg tcgctcttat tgaccacacc     180
tctaccggca tgccgagcaa atgctgcaa atcgctcccc atttcacca attgtagata     240
tgctaactcc agcaatgagt tgatgaatct cgggtgtgat tttatgtcct cagaggacaa     300
cacctgttgt aatcgttctt ccacac                                     326

```

<210> SEQ ID NO 51

<211> LENGTH: 374

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: promoter

<400> SEQUENCE: 51

```

gtgggtgat tggattatag gaagccacgc gctcaacctg gaattacagg aagctggtaa      60
ttttttgggt ttgcaatcat caccatctgc acgttggtat aatgtcccgt gtctatatat     120
atccattgac ggtattctat ttttttgcta ttgaaatgag cgttttttgt tactacaatt     180
ggttttacag acggaatttt cctatttctt ttcgtcccat ttttcctttt ctctattgtc     240
tcatatctta aaaaggtcct ttcttcataa tcaatgcttt cttttactta atattttact     300
tgcatcagtg gaattttaat acatattcct ctagtcttgc aaaatcgatt tagaatcaag     360
ataccagcct aaaa                                     374

```

<210> SEQ ID NO 52

<211> LENGTH: 390

-continued

```

<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: promoter

<400> SEQUENCE: 52
ctcgtaggaa caatttcggg cccctgctg ttctctgag gttcatcttt tacatttgct    60
tctgctggat aattttcaga ggcaacaagg aaaattaga tggcaaaaag tcgtctttca    120
aggaaaaatc cccaccatct ttcgagatcc cctgtaactt attggcaact gaaagaatga    180
aaaggaggaa aatacaaaat atactagaac tgaaaaaaaa aaagtataaa tagagacgat    240
atatgccaat acttcacaat gttcgaatct attcttcatt tgcagctatt gtaaaataat    300
aaaacatcaa gaacaaacaa gctcaacttg tcttttctaa gaacaaagaa taaacacaaa    360
aacaaaaagt tttttaatt ttaatcaaaa                                390

<210> SEQ ID NO 53
<211> LENGTH: 302
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: terminator

<400> SEQUENCE: 53
acaaatcgct cttaaatata tacctaaaga acattaaagc tatattataa gcaagatac    60
gtaaatTTTTG cttatattat tatacacata tcatatttct atatttttaa gatttggtta    120
tataatgtac gtaatgcaaa ggaaataaat tttatacatt attgaacagc gtccaagtaa    180
ctacattatg tgcactaata gtttagcgtc gtgaagactt tattgtgtcg cgaaaagtaa    240
aaattttaaa aattagagca cttgaactt gcgaaaagg ttctcatcaa ctgttttaaaa    300
gg                                                    302

```

1. A eukaryotic cell capable of expressing the following nucleotide sequences, wherein the expression of these nucleotide sequences confers on the cell the ability to use L-arabinose and/or to convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product:

- (a) a nucleotide sequence encoding an arabinose isomerase (araA), wherein said nucleotide sequence is selected from the group consisting of:
 - i. nucleotide sequences encoding an araA, said araA comprising an amino acid sequence that has at least 55% sequence identity with the amino acid sequence of SEQ ID NO:1,
 - ii. nucleotide sequences comprising a nucleotide sequence that has at least 60% sequence identity with the nucleotide sequence of SEQ ID NO:2,
 - iii. nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);
 - iv. nucleotide sequences the sequences of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code,
- (b) a nucleotide sequence encoding a L-ribulokinase (araB), wherein said nucleotide sequence is selected from the group consisting of:

- i. nucleotide sequences encoding an araB, said araB comprising an amino acid sequence that has at least 20% sequence identity with the amino acid sequence of SEQ ID NO:3,
- ii. nucleotide sequences comprising a nucleotide sequence that has at least 50% sequence identity with the nucleotide sequence of SEQ ID NO:4,
- iii. nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);
- iv. nucleotide sequences the sequences of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code,
- (c) a nucleotide sequence encoding an L-ribulose-5-P-4-epimerase (araD), wherein said nucleotide sequence is selected from the group consisting of:
 - i. nucleotide sequences encoding an araD, said araD comprising an amino acid sequence that has at least 60% sequence identity with the amino acid sequence of SEQ ID NO:5,
 - ii. nucleotide sequences comprising a nucleotide sequence that has at least 60% sequence identity with the nucleotide sequence of SEQ ID NO:6,
 - iii. nucleotide sequences the complementary strand of which hybridizes to a nucleic acid molecule of sequence of (i) or (ii);

- iv. nucleotide sequences the sequences of which differs from the sequence of a nucleic acid molecule of (iii) due to the degeneracy of the genetic code.
2. A cell according to claim 1, wherein one, two or three of the araA, araB and araD nucleotide sequences originate from a *Lactobacillus* genus, preferably a *Lactobacillus plantarum* species.
3. A cell according to claim 1, wherein the cell is a yeast cell, preferably belonging to one of the genera: *Saccharomyces*, *Kluyveromyces*, *Candida*, *Pichia*, *Schizosaccharomyces*, *Hansenula*, *Kloeckera*, *Schwanniomyces* or *Yarrowia*.
4. A cell according to claim 3, wherein the yeast cell belongs to one of the species: *S. cerevisiae*, *S. bulderi*, *S. barnetti*, *S. exiguus*, *S. uvarum*, *S. diastaticus*, *K. lactis*, *K. marxianus* or *K. fragilis*.
5. A cell according to claim 1, wherein the nucleotide sequences encoding the araA, araB and/or araD are operably linked to a promoter that causes sufficient expression of the corresponding nucleotide sequences in the cell to confer to the cell the ability to use L-arabinose and/or to convert L-arabinose into L-ribulose, and/or xylulose 5-phosphate and/or into a desired fermentation product.
6. A cell according to claim 1, wherein the cell exhibits the ability to directly isomerise xylose into xylulose.
7. A cell according to claim 6, wherein the cell comprises a genetic modification that increases the flux of the pentose phosphate pathway.
8. A cell according to claim 6, wherein the genetic modification comprises overexpression of at least one gene of the non-oxidative part of the pentose phosphate pathway.
9. A cell according to claim 8, wherein the gene is selected from the group consisting of the genes encoding ribulose-5-phosphate isomerase, ribulose-5-phosphate epimerase, transketolase and transaldolase.
10. A cell according to claim 8, wherein the genetic modification comprises overexpression of at least the genes coding for a transketolase and a transaldolase.
11. A cell according to claim 1, wherein the cell further comprises a genetic modification that increases the specific xylulose kinase activity.
12. A cell according to claim 11, wherein the genetic modification comprises overexpression of a gene encoding a xylulose kinase.
13. A cell according to claim 8, wherein the gene that is overexpressed is endogenous to the cell.
14. A cell according to claim 5, wherein the cell comprises a genetic modification that reduces unspecific aldose reductase activity in the cell.
15. A cell according to claim 14, wherein the genetic modification reduces the expression of, or inactivates a gene encoding an unspecific aldose reductase.
16. A cell according to claim 15, wherein the gene is inactivated by deletion of at least part of the gene or by disruption of the gene.
17. A cell according to claim 14, wherein the expression of each gene in the cell that encodes an unspecific aldose reductase is reduced or inactivated.
18. A cell according to claim 1, wherein the fermentation product is selected from the group consisting of ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic and a cephalosporin.
19. A nucleic acid construct comprising a nucleic acid sequence encoding an araA, a nucleic acid sequence encoding an araB and/or a nucleic acid sequence encoding an araD all as defined in claim 1.
20. A process for producing a fermentation product selected from the group consisting of ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam, antibiotic and a cephalosporin, whereby the process comprises:
- fermenting a medium containing a source of arabinose and optionally xylose with a modified cell as defined in claim 1, whereby the cell ferments arabinose and optionally xylose to the fermentation product; and optionally,
 - recovering the fermentation product.
21. A process for producing a fermentation product selected from the group consisting of ethanol, lactic acid, 3-hydroxy-propionic acid, acrylic acid, acetic acid, succinic acid, citric acid, malic acid, fumaric acid, an amino acid, 1,3-propane-diol, ethylene, glycerol, butanol, a β -lactam antibiotic and a cephalosporin, wherein the process comprises:
- fermenting a medium containing at least a source of L-arabinose and a source of xylose with a cell as defined in claim 1 and a cell able to use xylose and/or exhibiting the ability to directly isomerise xylose into xylulose, whereby each cell ferments L-arabinose and/or xylose to the fermentation product; and optionally,
 - recovering the fermentation product.
22. A process according to claim 20, wherein the medium also contains a source of glucose.
23. A process according to claim 20, wherein the fermentation product is ethanol.
24. A process according to claim 23, wherein the volumetric ethanol productivity is at least 0.5 g ethanol per litre per hour.
25. A process according to claim 23, wherein the ethanol yield is at least 30%.
26. A process according to claim 20, wherein the process is anaerobic.
27. A process according to claim 20, wherein the process is aerobic, preferably performed under oxygen limited conditions.

* * * * *