



US 20110014610A1

(19) **United States**

(12) **Patent Application Publication**
Akil et al.

(10) **Pub. No.: US 2011/0014610 A1**

(43) **Pub. Date: Jan. 20, 2011**

(54) **COMPOSITIONS AND METHODS FOR
DIAGNOSING AND TREATING
NEUROPSYCHIATRIC DISORDERS**

(75) Inventors: **Huda Akil**, Ann Arbor, MI (US);
Mary Atz, Tustin, CA (US);
William E. Bunney, JR., Laguna
Hills, CA (US); **William Byerley**,
San Francisco, CA (US); **Kathleen
Casey**, Santa Ana, CA (US);
Prabhakara V. Choudary, Davis,
CA (US); **Simon J. Evans**, Milan,
MI (US); **Edward G. Jones**,
Winters, CA (US); **Jun Li**, Palo
Alto, CA (US); **Juan F. Lopez**, Ann
Arbor, MI (US); **Richard M.
Myers**, Stanford, CA (US); **Brandi
Rollins**, Laguna Hills, CA (US);
Robert C. Thompson, Ann Arbor,
MI (US); **Hiroaki Tomita**, Irvine,
CA (US); **Marquis P. Vawter**,
Laguna Niguel, CA (US); **Stanley
J. Watson**, Ann Arbor, MI (US)

Correspondence Address:

**TOWNSEND AND TOWNSEND AND CREW,
LLP
TWO EMBARCADERO CENTER, EIGHTH
FLOOR
SAN FRANCISCO, CA 94111-3834 (US)**

(73) Assignee: **Board of Trustees of the Leland
Stanford Junior University**, Palo
Alto, CA (US)

(21) Appl. No.: **12/701,177**

(22) Filed: **Feb. 5, 2010**

Related U.S. Application Data

(63) Continuation of application No. 11/396,050, filed on
Mar. 31, 2006, now Pat. No. 7,687,235.

(60) Provisional application No. 60/667,299, filed on Mar.
31, 2005, provisional application No. 60/776,103,
filed on Feb. 22, 2006.

Publication Classification

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
G01N 33/53 (2006.01)

(52) **U.S. Cl.** **435/6; 435/7.1; 436/501**

(57) **ABSTRACT**

The present invention provides methods for diagnosing men-
tal disorders (e.g., psychotic disorders such as schizophrenia
and mood disorders such as major depression disorder and
bipolar disorder). The invention also provides methods of
identifying modulators of such mental disorders as well as
methods of using these modulators to treat patients suffering
from such mental disorders.

Annotation				Direction of Change in SZ relative to Controls					Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.444558	AA007604	KHDRBS3	8			up				KH domain containing, RNA binding, signal transduction associated 3
Hs.101139	AA017257		19	down						Transcribed locus
Hs.443049	AA251773		7		down					Transcribed locus
Hs.192788	AA314461	LOC389677	8						up	Similar to RIKEN cDNA 3000004N20
Hs.485557	AA418695	GSTA4	6					down		Glutathione S-transferase A4
Hs.535419	AA447681		18			up				Hypothetical LOC388459
Hs.36958	AA496799	BCAR3	1						down	Breast cancer anti-estrogen resistance 3
Hs.4204	AA700440		1					down		CDNA FLJ30779 fis, clone FEBRA2000815
Hs.241548	AA743526	RASA2	3		up					RAS p21 protein activator 2
Hs.524367	AA779991	CBARA1	10	down						Calcium binding atopy-related autoantigen 1
Hs.532987	AA827062	C18orf22	18	down						Chromosome 18 open reading frame 22
Hs.136905	AB002310	UREB1						down		Upstream regulatory element binding protein 1
Hs.114169	AB007876	LRRTM2	5				up			Leucine rich repeat transmembrane neuronal 2
Hs.435557	AB011103	KIF5C	2				up			Kinesin family member 5C
Hs.383564	AB011146	KIAA0574	15				up			KIAA0574 protein
Hs.146007	AB011154	KIAA0582	2			up				KIAA0582
Hs.118140	AB018259	DOCK4	7						down	Dedicator of cytokinesis 4
Hs.153610	AB018294	RIMS2	8				up			Regulating synaptic membrane exocytosis 2
Hs.178471	AB018341	ZNF432	19					down		Zinc finger protein 432
Hs.466261	AB021644	ZNF14	19		down					Zinc finger protein 14 (KOX 6)
Hs.303454	AB023136	SEC15L2	2	down						SEC15-like 2 (S. cerevisiae)
Hs.298658	AB024523	KLF3	4		up					Kruppel-like factor 3 (basic)
Hs.476399	AB029028	RAP140	3				down	down		Retinoblastoma-associated protein 140
Hs.132813	AB029033	IQSEC3	12			down				IQ motif and Sec7 domain 3
Hs.12264	AB032989	KIAA1163		down				down		Amphoterin-induced gene
Hs.50823	AB033060	PDCD6	5	down						Programmed cell death 6
Hs.472285	AB033098	KIAA1272	20					down		KIAA1272 protein
Hs.537101	AB033338					down				KRMP1 mRNA for mitotic kinesin-related protein, partial cds, alternative exon sequence.
Hs.140903	AB037744	MIB1	18			up				Mindbomb homolog 1 (Drosophila)
Hs.3346	AB037811	FLJ11280	1		up					Hypothetical protein FLJ11280
Hs.377588	AB040883	KIAA1450	4	up						KIAA1450 protein
Hs.287362	AB046767	TLE3	15						up	Transducin-like enhancer of split 3 (E(sp1) homolog,

FIG. 1 (Sheet 1 of 45)

UniGene ID	Annotation			Direction of Change in SZ relative to Controls							Name
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG		
Hs.13305	AB046788	ROBO2	3		up					Drosophila)	
Hs.474914	AB051446	RUTBC3	22	down						Roundabout, axon guidance receptor, homolog 2 (Drosophila)	
Hs.510745	AB097009	HNLF	7							RUN and TBC1 domain containing 3	
Hs.523789	AF001893	TncRNA	11						down	Putative NFkB activating protein HNLF	
Hs.352614	AF007155	LOC254531	15			down				Trophoblast-derived noncoding RNA	
Hs.147770	AF012023	ITGB1BP1	2	up						PLSC domain containing protein	
Hs.42400	AF022789	USP12	13	up						Integrin beta 1 binding protein 1	
Hs.465784	AF026030	TIMM44	19		up					Ubiquitin specific protease 12	
Hs.12451	AF035276	EML1	14	up						Translocase of inner mitochondrial membrane 44 homolog (yeast)	
Hs.12473	AF052109		1				up			Echinoderm microtubule associated protein like 1	
Hs.13438	AF052141		4				up			CDNA clone IMAGE:5260262, partial cds	
Hs.368046	AF054993	SNAP91	6				up			Clone 24626 mRNA sequence	
Hs.129997	AF070534		13				up			Synaptosomal-associated protein, 91kDa homolog (mouse)	
Hs.48372	AF086092		4					down		Clones 24632 and 24634 mRNA sequence	
Hs.513509	AF086220	FLJ32130	16	down						Full length insert cDNA clone YZ87G11	
Hs.131133	AF087875	PRKAG2	7		up					Hypothetical protein FLJ32130	
Hs.34560	AF116668	LMO2	11			down		down		Protein kinase, AMP-activated, gamma 2 non-catalytic subunit	
Hs.142245	AF126163	HHLA3	1	up						LIM domain only 2 (rhombotin-like 1)	
Hs.446240	AF144233	PRKCBP1	20	down						HERV-H LTR-associated 3	
Hs.47261	AF260704	SLCO1C1	12				down			Protein kinase C binding protein 1	
Hs.371788	AF318362	DKFZP547E1010	1	up					down	Solute carrier organic anion transporter family, member 1C1	
Hs.125747	AF381172		17		up					DKFZP547E1010 protein	
Hs.187946	AI240538		2							AA02 pseudogene mRNA, partial sequence, mRNA sequence	
Hs.23187	AI242476		7		down			down		Hypothetical gene supported by AK124342	
Hs.328801	AI308898		8			down				Transcribed locus	
Hs.542164	AI417157		19	up						Transcribed locus	
Hs.421200	AI480014		5	down		up				CDNA FLJ41369 fis, clone BRCAN2006117	
										Clone 24571 mRNA sequence	

FIG. 1 (Sheet 2 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.445066	AI499801	GRIN2B	12					down		Glutamate receptor, ionotropic, N-methyl D-aspartate 2B
Hs.47995	AI610676		10		down					Full-length cDNA clone CS0DC007YK10 of Neuroblastoma Cot 25-normalized of Homo sapiens (human)
Hs.503584	AI638679	PANX1	11			up				Pannexin 1
Hs.174746	AI670992		3					down		Transcribed locus
Hs.440729	AI671849		3			down				Transcribed locus
Hs.404218	AI674644		8		down					Transcribed locus
Hs.530218	AI679805		8						down	Transcribed locus
Hs.537738	AI692882		16		up					Full length insert cDNA clone ZD81C11
Hs.513356	AI694544		16	up	down					Transcribed locus, moderately similar to XP_498452.1 hypothetical gene supported by NM_173697 [Homo sapiens]
Hs.356084	AI703476	GPR27	3	up					up	G protein-coupled receptor 27
Hs.201106	AI758946		1		up					Transcribed locus
Hs.192788	AI768185	LOC389677	8						up	Similar to RIKEN cDNA 3000004N20
Hs.202054	AI796835					down				Transcribed locus
Hs.536522	AI822096	SMAD4							up	SMAD, mothers against DPP homolog 4 (Drosophila)
Hs.97104	AI823879		5		up					Transcribed locus
Hs.94949	AI934339	MCEE	2	up						Methylmalonyl CoA epimerase
Hs.142869	AI935586		4		up					Transcribed locus
Hs.459255	AI935701	NTRK3	15	up						Neurotrophic tyrosine kinase, receptor, type 3
Hs.114033	AJ420492	SSR1	6						down	Signal sequence receptor, alpha (translocon-associated protein alpha)
Hs.445098	AK000490	DEPDC1	1		up					DEP domain containing 1
Hs.517134	AK000586	C20orf43	20	up						Chromosome 20 open reading frame 43
Hs.288995	AK000820	ZNF587	19					down		Zinc finger protein 587
Hs.471221	AK000969	KLF7	2			up			up	Kruppel-like factor 7 (ubiquitous)
Hs.301943	AK001249	KIAA0467	1		down					KIAA0467 protein
Hs.144055	AK001563	SBNO1	12		up				up	Sno, strawberry notch homolog 1 (Drosophila)
Hs.27021	AK001697	RIOK2	5		down				down	RIO kinase 2 (yeast)
Hs.443260	AK001776	C20orf20	20		down		down		down	Chromosome 20 open reading frame 20
Hs.516311	AK001856		2					down	down	Homo sapiens, clone IMAGE:4826696, mRNA
Hs.92308	AK002085	LOC144438	12			up				Hypothetical protein LOC144438
Hs.477693	AK021677	NCK1	3	down						NCK adaptor protein 1

FIG. 1 (Sheet 3 of 45)

Annotation				Direction of Change in SZ relative to Controls					Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPRC	Nacc	PC	STG	Name
Hs.481819	AK021922	PDZK3	5						down	PDZ domain containing 3
Hs.476982	AK022155	CPOX	3		up					Coproporphyrinogen oxidase
Hs.492445	AK022204	EDD	8					down		E3 identified by differential display
Hs.31431	AK022233	FN3KRP	17;20;9	up						Fructosamine-3-kinase-related protein
Hs.285782	AK022630	LRRTM4	2			up	up			Leucine rich repeat transmembrane neuronal 4
Hs.132794	AK022741	PCYT1B				up				Phosphate cytidylyltransferase 1, choline, beta isoform
Hs.321653	AK022832	FLJ12770	1	up						Hypothetical protein FLJ12770
Hs.167165	AK023037	FLJ12975	12					down	down	Hypothetical protein FLJ12975
Hs.440833	AK023692	PKN2	1					down		Protein kinase N2
Hs.274422	AK024220	C20orf27	20	up						Chromosome 20 open reading frame 27
Hs.473374	AK025196	PTPRK	6			down				Protein tyrosine phosphatase, receptor type, K
Hs.431081	AK025233	USP53	4		down					Ubiquitin specific protease 53
Hs.499483	AK025626	LOC83693	16	up						Steroid dehydrogenase-like
Hs.465295	AK025773	LMAN1	18				up			Lectin, mannose-binding, 1
Hs.289092	AK026033	COTL1	16;11					up		Coactosin-like 1 (Dictyostelium)
Hs.537428	AK026035							down		CDNA FLJ34018 fis, clone FCBBF2002801
Hs.426324	AK026149	TUSC3	8		up					Tumor suppressor candidate 3
Hs.272759	AK026156	PITPNM2		up						Phosphatidylinositol transfer protein, membrane-associated 2
Hs.535771	AK026466	CYFIP2	5	down						Cytoplasmic FMR1 interacting protein 2
Hs.524341	AK026495	SLC2A13	12		up					Solute carrier family 2 (facilitated glucose transporter), member 13
Hs.79881	AK026659		3		down					CDNA: FLJ23006 fis, clone LNG00414
Hs.91521	AK026748	DKFZP761M1511	5	up					up	Hypothetical protein DKFZP761M1511
Hs.301296	AK026784		13		down					CDNA: FLJ23131 fis, clone LNG08502
Hs.415842	AK026870	RBM18	9				up			RNA binding motif protein 18
Hs.292575	AK026980	ZNF37B	10;7;1	down						Zinc finger protein 37b (KOX 21)
Hs.111286	AK027059	MRPS11	15		up					Mitochondrial ribosomal protein S11
Hs.130692	AK027273	MGC10946	12		down					Hypothetical protein MGC10946
Hs.452398	AK055302		2	down						MRNA; cDNA DKFZp564E143 (from clone DKFZp564E143)
Hs.532786	AK055378	LOC339287	17	up						Hypothetical protein LOC339287

FIG. 1 (Sheet 4 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.148105	AK055479	PRICKLE2	3	up						Prickle-like 2 (Drosophila)
Hs.536251	AK056079	JAM2	21					down		Junctional adhesion molecule 2
Hs.47918	AK056549	CNKSR2	X				up			Connector enhancer of kinase suppressor of Ras 2
Hs.193115	AK057990	KIAA1764	8	down						KIAA1764 protein
Hs.251399	AK074730	HRH3	20	up						Histamine receptor H3
Hs.254414	AK090803	SRrp35	6	up						Serine-arginine repressor protein (35 kDa)
Hs.242947	AK091081	DGKI	7			up				Diacylglycerol kinase, iota
Hs.443731	AK091775	USP8	15		up					Ubiquitin specific protease 8
Hs.494804	AK091948	LTB4DH	9			up				Leukotriene B4 12-hydroxydehydrogenase
Hs.179153	AK092145		9		up					CDNA FLJ34826 fis, clone NT2NE2008803
Hs.177275	AK092235	ANKRD6	6	up						Ankyrin repeat domain 6
Hs.433956	AK092711		2		up					Hypothetical LOC400944
Hs.144447	AK092984	WDR11	10					down		WD repeat domain 11
Hs.293077	AK093067	CHPT1	12			up				Choline phosphotransferase 1
Hs.523467	AK093229	NRIP3	11						up	Nuclear receptor interacting protein 3
Hs.465462	AK093588		18						up	CDNA FLJ36269 fis, clone THYMU2003012
Hs.12248	AK093871	CXXC4	4		up					CXXC finger 4
Hs.451846	AK093939	RGPR	1	down						Regucalcin gene promotor region related protein
Hs.331667	AK094282	MGC3200	1			down				Hypothetical protein LOC284615
Hs.22654	AK094487	SCN1A	2				up			Sodium channel, voltage-gated, type I, alpha
Hs.26479	AK094535		3			up			down	CDNA FLJ37216 fis, clone BRALZ2008696
Hs.438695	AK094876	FKBP11	12	down						FK506 binding protein 11, 19 kDa
Hs.380250	AK094968	IFI16	1						down	Interferon, gamma-inducible protein 16
Hs.466987	AK095884	PRKD2	19			down				Protein kinase D2
Hs.128686	AK097398	NUCB2	11			up				Nucleobindin 2
Hs.169378	AK098775	MPDZ	9				down	down		Multiple PDZ domain protein
Hs.494312	AK123824	NTRK2	9		up					Neurotrophic tyrosine kinase, receptor, type 2
Hs.525589	AK123878	MEG3	14					up		Maternally expressed 3
Hs.164649	AK124629	IXL	19					down		Intersex-like (Drosophila)
Hs.476782	AK126999	EIF4E3	3			up			up	Eukaryotic translation initiation factor 4E member 3
Hs.297044	AK127019	RUFY2	10			down				RUN and FYVE domain containing 2
Hs.256801	AK127476	ZNF493		up						Zinc finger protein 493
Hs.520804	AK128034		7						down	Similar to cell division cycle 10 homolog
Hs.435767	AK129838	PCYT1A	3		down					Phosphate cytidylyltransferase 1, choline, alpha isoform
Hs.126914	AK130263	KIAA1430	4	up						KIAA1430

FIG. 1 (Sheet 5 of 45)

Annotation			Direction of Change in SZ relative to Controls							
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.512181	AK130506	CXorf33		down						Chromosome X open reading frame 33
Hs.480825	AK130520	RNF150	4			up				Ring finger protein 150
Hs.184216	AL049980	DKFZP564C152	11						down	DKFZP564C152 protein
Hs.155090	AL117471	GNB5	15				up			Guanine nucleotide binding protein (G protein), beta 5
Hs.421907	AL122063	GLTSCR2				down				Glioma tumor suppressor candidate region gene 2
Hs.158688	AL133563	EIF5B	2	down						Eukaryotic translation initiation factor 5B
Hs.435700	AL137537	ATP8B2	1		up					ATPase, Class I, type 8B, member 2
Hs.478125	AL137692	INADL	1						up	InaD-like protein
Hs.26815	AL360202	THAP10	15			up				THAP domain containing 10
Hs.475103	AL389951	NUP50	22						up	Nucleoporin 50kDa
Hs.536940	AL512705	APEG1	2			down				Aortic preferentially expressed protein 1
Hs.292549	AL831922	DLG1	3	up						Discs, large homolog 1 (Drosophila)
Hs.268675	AL831995	MEF2A	15		up					MADS box transcription enhancer factor 2, polypeptide A (myocyte enhancer factor 2A)
Hs.518099	AL832398	MGC26717	3		down	up				Hypothetical protein MGC26717
Hs.30141	AL832603	FLJ20313	15		down					Hypothetical protein FLJ20313
Hs.175343	AL832699	PIK3C2A	11						down	Phosphoinositide-3-kinase, class 2, alpha polypeptide
Hs.490790	AL833137	THAP5	7					down		THAP domain containing 5
Hs.430300	AL833360		11		down					MRNA; cDNA DKFZp667E0114 (from clone DKFZp667E0114)
Hs.477921	AL833852	WWTR1	3		down					WW domain containing transcription regulator 1
Hs.282855	AL834302	FLJ12994	15		up					Hypothetical protein FLJ12994
Hs.32468	AV718518		5			up				Transcribed locus
Hs.163924	AW080999	NR3C2	4					down		Nuclear receptor subfamily 3, group C, member 2
Hs.418198	AW086065	PAPD4	5					down		PAP associated domain containing 4
Hs.446340	AW291402					up				Transcribed locus
Hs.293379	AW302422		19		up					Transcribed locus
Hs.429	AW438709	ATP5G3	2			down				ATP synthase, H+ transporting, mitochondrial FO complex, subunit c (subunit 9) isoform 3
Hs.408427	AW451792	COMM7	20	up						COMM domain containing 7
Hs.436556	AW571739		7	down						Transcribed locus
Hs.257786	AW964241		2					up		Transcribed locus
Hs.306423	BC000825	LOC339524	1						down	Hypothetical protein LOC339524
Hs.434953	BC001063	HMGB2	4			up				High-mobility group box 2

FIG. 1 (Sheet 6 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.535659	BC002458	MCM3AP	21		up					MCM3 minichromosome maintenance deficient 3 (S. cerevisiae) associated protein
Hs.483239	BC002515	ALDH7A1	5	up			down			Aldehyde dehydrogenase 7 family, member A1
Hs.7001	BC002867	C14orf9		up						Chromosome 14 open reading frame 9
Hs.224282	BC004887	LANCL2	7			up				LanC lantibiotic synthetase component C-like 2 (bacterial)
Hs.471582	BC004921	LOC93349	2			down				Hypothetical protein BC004921
Hs.516859	BC005095	PANK2	20		down					Pantothenate kinase 2 (Hallervorden-Spatz syndrome)
Hs.153546	BC005258	CDC23	5			up				CDC23 (cell division cycle 23, yeast, homolog)
Hs.18788	BC006283	DHRS10	19					down		Dehydrogenase/reductase (SDR family) member 10
Hs.443673	BC008143	KIAA1002	12;1			down				KIAA1002 protein
Hs.529630	BC008625		3						down	Homo sapiens, clone IMAGE:4183899, mRNA
Hs.534483	BC008630	MGC2941	17	up						Hypothetical protein MGC2941
Hs.516859	BC008667	PANK2	20		down					Pantothenate kinase 2 (Hallervorden-Spatz syndrome)
Hs.11923	BC009674	DJ167A19.1	1;2	up			up			Hypothetical protein DJ167A19.1
Hs.447579	BC010538	LOC339290	18					up		Hypothetical protein LOC339290
Hs.103555	BC011054	FLJ14775	17			down				Hypothetical protein FLJ14775
Hs.194408	BC014227	KIAA1244	6	up						KIAA1244
Hs.50823	BC014604	PDCD6	5	down						Programmed cell death 6
Hs.129837	BC015067	ZBTB8	1	up				down		Zinc finger and BTB domain containing 8
Hs.445113	BC015910	MARCH-II	19	up						Membrane-associated RING-CH protein II
Hs.487325	BC016285	PRKACB	1		up				down	Protein kinase, cAMP-dependent, catalytic, beta
Hs.536470	BC016735	LOC63929	22		up				down	Hypothetical protein LOC63929
Hs.279908	BC017788	TFB1M	6					down		Transcription factor B1, mitochondrial
Hs.517792	BC019303	C3orf10	3	up				up		Chromosome 3 open reading frame 10
Hs.499145	BC019602	YME1L1	10	down						YME1-like 1 (S. cerevisiae)
Hs.350268	BC020516	IRF2BP2				up				Interferon regulatory factor 2 binding protein 2
Hs.197922	BC020630	CaMKII α	1				up			Calcium/calmodulin-dependent protein kinase II
Hs.208961	BC021961	NSD1	5					down		Nuclear receptor binding SET domain protein 1
Hs.525588	BC023543	MEG3	14						up	Maternally expressed 3
Hs.436568	BC024272	CD74	5		down		down			CD74 antigen (invariant polypeptide of major histocompatibility complex, class II antigen-associated)
Hs.180933	BC029922	CXXC1	18					up		CXXC finger 1 (PHD domain)
Hs.535810	BC030122		5	up						CDNA clone IMAGE:4814828, partial cds
Hs.74655	BC033998	LOC124512	17	up						Hypothetical protein LOC124512
Hs.479099	BC035257	SORCS2	4	up						Sortilin-related VPS10 domain containing receptor 2

FIG. 1 (Sheet 7 of 45)

Annotation			Direction of Change in SZ relative to Controls									
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name		
Hs.459070	BC036099	ARNT2	15		down					Aryl-hydrocarbon receptor nuclear translocator 2		
Hs.302631	BC036622		7					down		CDNA clone IMAGE:5286843, partial cds		
Hs.98132	BC036875	LCN6	9	down						Lipocalin 10		
Hs.127951	BC037165	FLJ14503		down						Hypothetical protein FLJ14503		
Hs.90242	BC037800		1		up					Homo sapiens, clone IMAGE:4796172, mRNA		
Hs.212151	BC039075	CLSTN3	12			down				Calsynenin 3		
Hs.461074	BC040486	ZFP90	16			up				Zinc finger protein 90 homolog (mouse)		
Hs.516853	BC041916	C20orf194	20	up		up				Chromosome 20 open reading frame 194		
Hs.506309	BC041930	EEA1	12					down		Early endosome antigen 1, 162kD		
Hs.536567	BC042073								up	Data not found		
Hs.205865	BC042754	LOC143458							down	Hypothetical protein LOC143458		
Hs.434418	BC042833	MYT1L	2				up			Myelin transcription factor 1-like		
Hs.122110	BC043568	FLJ33718	4		up					Hypothetical protein FLJ33718		
Hs.413416	BC047331	JMJD1C	10					down		Jumonji domain containing 1C		
Hs.136888	BC047477	LOC338758	12		up					Hypothetical protein LOC338758		
Hs.443258	BC051799	SREBF2	22	up						Sterol regulatory element binding transcription factor 2		
Hs.169182	BC052964	KIF21B	1;19; 16						up	Kinesin family member 21B		
Hs.509314	BC059410	LOC285148	2			down		down		Hypothetical protein LOC285148		
Hs.180714	BC064523	15E1.2	12	down						Hypothetical protein 15E1.2		
Hs.519904	BC065748	RBM24	6					down		RNA binding motif protein 24		
Hs.459590	BC066358	CCL27	9			down				Chemokine (C-C motif) ligand 27		
Hs.433150	BC067884		1						up	CDNA clone IMAGE:4480427, partial cds		
Hs.506458	BC068451	EB-1	12		up					E2a-Pbx 1-associated protein		
Hs.86508	BC079833		5		down					CDNA clone IMAGE:4841343, partial cds		
Hs.12862	BC080181	RFNG	17	up						Radical fringe homolog (Drosophila)		
Hs.115284	BC080572	ZNF213	16						up	Zinc finger protein 213		
Hs.143587	BE467201							down		Transcribed locus		
Hs.102471	BE538923	PHACTR2	6	down						Phosphatase and actin regulator 2		
Hs.127486	BE551624		2					down		Transcribed locus		
Hs.21691	BE671266	GPR75	2	down						G protein-coupled receptor 75		
Hs.188594	BE889301		9		up					Transcribed locus		
Hs.118526	BF056604			down						Transcribed locus		
Hs.225598	BF059209		6						up	Transcribed locus		
Hs.144022	BF109731	FDFT1	8		up					Farnesyl-diphosphate farnesyltransferase 1		

FIG. 1 (Sheet 8 of 45)

Annotation			Direction of Change in SZ relative to Controls						Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.464848	BF378154	B4GALT6	18		up					UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 6
Hs.432792	BF439526	CBLL1	7			up				Cas-Br-M (murine) ecotropic retroviral transforming sequence-like 1
Hs.478465	BG707584	FLJ12748	3		up					Hypothetical protein FLJ12748
Hs.434375	BI820698	PTPRB	12					down	down	Protein tyrosine phosphatase, receptor type, B
Hs.414028	BM455428	C9orf116	9;3	up						Chromosome 9 open reading frame 116
Hs.469967	BM557121					up				Transcribed locus
Hs.445247	BM682460		5					down		Transcribed locus, weakly similar to NP_703324.1 Plasmodium falciparum 3D7 MAL1P3.06 gene
Hs.133469	BM719738	GOLGA1	9			down				Golgi autoantigen, golgin subfamily a, 1
Hs.479766	BP225938	TPARL	4						down	TPA regulated locus
Hs.470882	BQ002778	CALCRL	2		down					Calcitonin receptor-like
Hs.118769	BQ003501		2					down		Transcribed locus
Hs.229304	BQ007533								down	Transcribed locus
Hs.487648	BQ285965	SNX13	7					down		Sorting nexin 13
Hs.104980	BQ331336					up				Data not found
Hs.479808	BQ477415	IGFBP7	4				down	down		Insulin-like growth factor binding protein 7
Hs.179238	BU674160	LRRC6	8			up				Leucine rich repeat containing 6
Hs.31903	BU685761		3					down		Transcribed locus
Hs.28199	BX091447		7					down		Transcribed locus
Hs.445015	BX093081	GRIN2D							up	Glutamate receptor, ionotropic, N-methyl D-aspartate 2D
Hs.7413	BX097190		3		up					Transcribed locus
Hs.452398	BX099722		2	down				up		MRNA; cDNA DKFZp564E143 (from clone DKFZp564E143)
Hs.4817	BX537377	OPCML	11				up			Opioid binding protein/cell adhesion molecule-like
Hs.482363	BX537394	SLC30A5	5		down					Solute carrier family 30 (zinc transporter), member 5
Hs.343522	BX537745	ATP2B4	1;14		up					ATPase, Ca++ transporting, plasma membrane 4
Hs.479853	BX537946	EPHA5	4	up	up	up				EPH receptor A5
Hs.370510	BX538269	IGSF4	11		up					Immunoglobulin superfamily, member 4
Hs.192221	BX538289	ELL2		up						Elongation factor, RNA polymerase II, 2
Hs.494178	BX647070	RORB	9							RAR-related orphan receptor B
Hs.433381	BX647220	C6orf89	6			up				Chromosome 6 open reading frame 89
Hs.310545	BX647240	SYT1	12			up				Synaptotagmin I
Hs.436142	BX647689	PTPN13	4	up			up			Protein tyrosine phosphatase, non-receptor type 13 (APO-

FIG. 1 (Sheet 9 of 45)

UniGene ID	Annotation			Direction of Change in SZ relative to Controls						
	Acc	Symbol	Chr.	AnCg	CB	DLFFC	Nacc	PC	STG	Name
Hs.369978	BX647773	GTDC1	2					up		1/CD95 (Fas)-associated phosphatase)
Hs.146542	BX648027	NEGR1	1		up					Glycosyltransferase-like 1
Hs.213050	BX648050	ELAVL4	1		up					Neuronal growth regulator 1
Hs.175343	BX648778	PIK3C2A	11		down		down			ELAV (embryonic lethal, abnormal vision, Drosophila)-like 4 (Hu antigen D)
Hs.306423	CA440056	LOC339524	1						up	Phosphoinositide-3-kinase, class 2, alpha polypeptide
Hs.327736	CA442378	KIF5B	10			up				Hypothetical protein LOC339524
Hs.537332	CB996893	CNOT6L	4					down		Kinesin family member 5B
Hs.105636	CD639734		1		down					CCR4-NOT transcription complex, subunit 6-like
Hs.443031	CK300949	GLB1	3						up	Transcribed locus
Hs.437611	CK820590		X				up			Galactosidase, beta 1
Hs.29802	CN260580	SLIT2	4					down		Transcribed locus
Hs.444818	CR456854	CGGBP1	3;12	down						Slit homolog 2 (Drosophila)
Hs.500333	CR596764	C10orf58	10					up		CGG triplet repeat binding protein 1
Hs.301296	CR616826		13		down					Chromosome 10 open reading frame 58
Hs.120446	CR623819	ZCWCC1	22				down			CDNA: FLJ23131 fis, clone LNG08502
Hs.250072	CR627428	SLC4A7	3		up					Zinc finger, CW type with coiled-coil domain 1
Hs.477134	CR749341	DKFZP434 F2021	3		down					Solute carrier family 4, sodium bicarbonate cotransporter, member 7
Hs.118351	D13635	UBE3C	7				up			DKFZP434F2021 protein
Hs.35804	D25215	HERC3	4;8		up					Ubiquitin protein ligase E3C
Hs.423163	D87969	SLC35A1	6	up						Hect domain and RLD 3
Hs.536256	F01952	GNAZ	22			down		up		Solute carrier family 35 (CMP-sialic acid transporter), member A1
Hs.538896	F02333	ANKRD10	13					down		Guanine nucleotide binding protein (G protein), alpha z polypeptide
Hs.27996	F10010		4		down			down		Ankyrin repeat domain 10
Hs.492212	H95037	DECR1	8		up					Transcribed locus
Hs.131711	J03620	DLD	7		down					2,4-dienoyl CoA reductase 1, mitochondrial
Hs.449451	K02885			down						Dihydroipoamide dehydrogenase (E3 component of pyruvate dehydrogenase complex, 2-oxo-glutarate complex, branched chain keto acid dehydrogenase complex)
										Isolate 971.4 G01 T cell receptor beta (TCBRV)

FIG. 1 (Sheet 10 of 45)

Annotation			Direction of Change in SZ relative to Controls							
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.274873	L06845	CARS	11					up		Cysteinyl-tRNA synthetase
Hs.220629	L17000	CAMK4	5				up			Calcium/calmodulin-dependent protein kinase IV
Hs.196983	M61199	SSFA2	2					down		Sperm specific antigen 2
Hs.212838	NM_000014	A2M	12			down		down		Alpha-2-macroglobulin
Hs.232375	NM_000019	ACAT1	11	up						Acetyl-Coenzyme A acetyltransferase 1 (acetoacetyl Coenzyme A thiolase)
Hs.207776	NM_000027	AGA	4				down		down	Aspartylglucosaminidase
Hs.445358	NM_000042	APOH	17	down	up					Apolipoprotein H (beta-2-glycoprotein I)
Hs.160786	NM_000050	ASS	9	up						Argininosuccinate synthetase
Hs.169348	NM_000057	BLM	15	down						Bloom syndrome
Hs.476218	NM_000094	COL7A1	3	down						Collagen, type VII, alpha 1 (epidermolysis bullosa, dystrophic, dominant and recessive)
Hs.304682	NM_000099	CST3	20;11;22;12		down					Cystatin C (amyloid angiopathy and cerebral hemorrhage)
Hs.335513	NM_000129	F13A1	6		down					Coagulation factor XIII, A1 polypeptide
Hs.255230	NM_000181	GUSB	7						down	Glucuronidase, beta
Hs.303154	NM_000202	IDS	X;12			up				Iduronate 2-sulfatase (Hunter syndrome)
Hs.224012	NM_000214	JAG1	20		down	down				Jagged 1 (Alagille syndrome)
Hs.156519	NM_000251	MSH2	2		down					MutS homolog 2, colon cancer, nonpolyposis type 1 (E. coli)
Hs.21213	NM_000259	MYO5A	15		up		up			Myosin VA (heavy polypeptide 12, myosin)
Hs.181272	NM_000297	PKD2	4			down	down			Polycystic kidney disease 2 (autosomal dominant)
Hs.476595	NM_000333	ATXN7	3		down		down		down	Ataxin 7
Hs.271771	NM_000345	SNCA	4					up		Synuclein, alpha (non A4 component of amyloid precursor)
Hs.370771	NM_000389	CDKN1A	6;9				down			Cyclin-dependent kinase inhibitor 1A (p21, Cip1)
Hs.2785	NM_000422	KRT17	17	down						Keratin 17
Hs.250769	NM_000478	ALPL	1	down						Alkaline phosphatase, liver/bone/kidney
Hs.24422	NM_000538	RFXAP	13			up				Regulatory factor X-associated protein
Hs.529400	NM_000629	IFNAR1	21						up	Interferon (alpha, beta and omega) receptor 1
Hs.150749	NM_000633	BCL2	18						up	B-cell CLL/lymphoma 2
Hs.336046	NM_000640	IL13RA2	X				up			Interleukin 13 receptor, alpha 2
Hs.150749	NM_000657	BCL2	18						up	B-cell CLL/lymphoma 2
Hs.334707	NM_000666	ACY1	3;1	down					down	Aminoacylase 1

FIG. 1 (Sheet 11 of 45)

UniGene ID	Annotation		Direction of Change in SZ relative to Controls							Name
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	
Hs.197029	NM_000675	ADORA2A	22				up			Adenosine A2a receptor
Hs.388004	NM_000687	AHCY	20;16	up				up		S-adenosylhomocysteine hydrolase
Hs.34114	NM_000702	ATPIA2	1		down					ATPase, Na+/K+ transporting, alpha 2 (+) polypeptide
Hs.433307	NM_000709	BCKDHA	19;1		down					Branched chain keto acid dehydrogenase E1, alpha polypeptide (maple syrup urine disease)
Hs.282871	NM_000770	CYP2C8	10		up					Cytochrome P450, family 2, subfamily C, polypeptide 8
Hs.152096	NM_000775	CYP2J2	1					up		Cytochrome P450, family 2, subfamily J, polypeptide 2
Hs.202354	NM_000793	DIO2	14	down						Deiodinase, iodothyronine, type II
Hs.175934	NM_000806	GABRA1	5;17				up			Gamma-aminobutyric acid (GABA) A receptor, alpha 1
Hs.473648	NM_000819	GART	21				up			Phosphoribosylglycinamide formyltransferase, phosphoribosylglycinamide synthetase, phosphoribosylaminoimidazole synthetase
Hs.32945	NM_000838	GRM1	6		up					Glutamate receptor, metabotropic 1
Hs.268573	NM_000853	GSTT1	22			down				Glutathione S-transferase theta 1
Hs.376933	NM_000858	GUK1	1				up			Guanylate kinase 1
Hs.46732	NM_000898	MAOB	X	up						Monoamine oxidase B
Hs.2820	NM_000916	OXTR	3						up	Oxytocin receptor
Hs.354056	NM_000941	POR	7					up		P450 (cytochrome) oxidoreductase
Hs.446429	NM_000954	PTGDS	9;11;1; 7;6;14; 2;19;1 7;3;16; 22		down		down			Prostaglandin D2 synthase 21kDa (brain)
Hs.374588	NM_000985	RPL17	18;1; 15;3	up						Ribosomal protein L17
Hs.529631	NM_000996	RPL35A	3	up						Ribosomal protein L35a
Hs.134846	NM_001001 410	MGC24381	16	up						Hypothetical protein MGC24381
Hs.492031	NM_001001 482	FLJ11011	8						up	Hypothetical protein FLJ11011
Hs.188569	NM_001001 483	ZDHHC13	11					down		Zinc finger, DHHC domain containing 13
Hs.429	NM_001002 258	ATP5G3	2	up						ATP synthase, H+ transporting, mitochondrial F0 complex, subunit c (subunit 9) isoform 3
Hs.430439	NM_001002 757	HIRIP5	2				up			HIRA interacting protein 5

FIG. 1 (Sheet 12 of 45)

UniGene ID	Annotation			Direction of Change in SZ relative to Controls							Name
	Acc	Symbol	Chr.	AnCg	CB	DLFPC	Nacc	PC	STG		
Hs.92423	NM_001002838	PRKWINK3		down							Protein kinase, lysine deficient 3
Hs.522418	NM_001003722	GLE1L	9	up							GLE1 RNA export mediator-like (yeast)
Hs.444445	NM_001003802	SMARCD3	7				up				SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily d, member 3
Hs.528826	NM_001004300	LOC124411	16	down							Hypothetical protein LOC124411
Hs.462230	NM_001004313	LOC388335	17					down			Similar to RIKEN cDNA A730055C05 gene
Hs.405925	NM_001005290	DDA3	1	up							Differential display and activated by p53
Hs.446623	NM_001005335	HNRPL	19;15;2	down			down				Heterogeneous nuclear ribonucleoprotein L
Hs.438219	NM_001005408	KIAA1787						up			G protein pathway suppressor 2
Hs.334873	NM_001005502	CPM	12		up			up			Carboxypeptidase M
Hs.445841	NM_001036	RYR3	15		down						Ryanodine receptor 3
Hs.162585	NM_001046	SLC12A2	5	down	up				down		Solute carrier family 12 (sodium/potassium/chloride transporters), member 2
Hs.12409	NM_001048	SST	3						down		Somatostatin
Hs.159306	NM_001092	ABR	17				up				Active BCR-related gene
Hs.274361	NM_001095	ACCN2	12				down				Amiloride-sensitive cation channel 2, neuronal
Hs.474982	NM_001098	ACO2	22				up				Aconitase 2, mitochondrial
Hs.470316	NM_001105	ACVRI	2	down							Activin A receptor, type I
Hs.461253	NM_001128	APIG1	16	up							Adaptor-related protein complex 1, gamma 1 subunit
Hs.480653	NM_001154	ANXA5	4				down				Annexin A5
Hs.483239	NM_001182	ALDH7A1	5	up			down				Aldehyde dehydrogenase 7 family, member A1
Hs.25447	NM_001268	CHC1L	13				down				Chromosome condensation 1-like
Hs.162233	NM_001273	CHD4	12			up					Chromodomain helicase DNA binding protein 4
Hs.150793	NM_001275	CHGA	14				up				Chromogranin A (parathyroid secretory protein 1)
Hs.249129	NM_001279	CIDEA	18					up			Cell death-inducing DFFA-like effector a
Hs.23748	NM_001290	LDB2	4				up				LIM domain binding 2
Hs.13313	NM_001310	CREBL2	12		down						CAMP responsive element binding protein-like 2
Hs.166011	NM_001331	CTNND1	11;16			down		down			Catenin (cadherin-associated protein), delta 1

FIG. 1 (Sheet 13 of 45)

Annotation			Direction of Change in SZ relative to Controls							
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLFC	Nacc	PC	STG	Name
Hs.336916	NM_001350	DAXX	19							
Hs.159195	NM_001380	DOCK1	6		down			up		Death-associated protein 6
Hs.117060	NM_001393	ECM2	10				down			Dedicator of cytokinesis 1
Hs.196176	NM_001398	ECH1	9		down					Extracellular matrix protein 2, female organ and adipocyte specific
Hs.132483	NM_001410	EGFL4	19;X	up						Enoyl Coenzyme A hydratase 1, peroxisomal
Hs.299002	NM_001436	FBL	19	up						EGF-like-domain, multiple 4
Hs.357637	NM_001439	EXTL2	1			down	down			Fibrillarlin
Hs.272011	NM_001497	B4GALT1	9		up					Exostosins (multiple)-like 2
Hs.47338	NM_001549	IFIT3	10	down						UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 1
Hs.469386	NM_001566	INPP4A	2			up				Interferon-induced protein with tetratricopeptide repeats 3
Hs.289795	NM_001584	C11orf8	11		up			down		Inositol polyphosphate-4-phosphatase, type I, 107kDa
Hs.166160	NM_001607	ACAA1	3						down	Chromosome 11 open reading frame 8
Hs.368486	NM_001649	APXL	X	up	up					Acetyl-Coenzyme A acyltransferase 1 (peroxisomal 3-oxoacyl-Coenzyme A thiolase)
Hs.286221	NM_001658	ARF1	1;19		down					Apical protein-like (Xenopus laevis)
Hs.3109	NM_001666	ARHGAP4	X		up					ADP-ribosylation factor 1
Hs.413137	NM_001680	FXYD2	11					up		Rho GTPase activating protein 4
Hs.198365	NM_001724	BPGM	7			up				FXYD domain containing ion transport regulator 2
Hs.274873	NM_001751	CARS	11					up		2,3-bisphosphoglycerate mutase
Hs.445570	NM_001780	CD63	12;X				down			CysteinyI-tRNA synthetase
Hs.170129	NM_001821	CHML	1					down		CD63 antigen (melanoma 1 antigen)
Hs.129966	NM_001842	CNTFR	9				down			Choroideremia-like (Rab escort protein 2)
Hs.421621	NM_001864	COX7A1	19						down	Ciliary neurotrophic factor receptor
Hs.2242	NM_001891	CSN2	4	down						Cytochrome c oxidase subunit VIIa polypeptide 1 (muscle)
Hs.410037	NM_001901	CTGF	6;16		down					Casein beta
Hs.465413	NM_001914	CYB5	18			up				Connective tissue growth factor
Hs.113227	NM_001917	DAO	12		down					Cytochrome b-5
Hs.433839	NM_001958	EEF1A2	20;19;17;9				up			D-amino-acid oxidase
Hs.326035	NM_001964	EGR1	5	down						Eukaryotic translation elongation factor 1 alpha 2
Hs.306251	NM_001982	ERBB3	12				down			Early growth response 1
									down	V-erb-b2 erythroblastic leukemia viral oncogene homolog

FIG. 1 (Sheet 14 of 45)

UniGene ID	Annotation				Direction of Change in SZ relative to Controls						
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name	
Hs.370666	NM_002015	FOXO1A	13	down			down			3 (avian)	
Hs.103183	NM_002024	FMR1	X		down			down		Forkhead box O1A (rhabdomyosarcoma)	
Hs.62661	NM_002053	GBP1	1			down				Fragile X mental retardation 1	
Hs.430425	NM_002074	GNB1	1				up			Guanylate binding protein 1, interferon-inducible, 67kDa polypeptide 1	
Hs.309763	NM_002092	GRSF1	4;17	up	down					G-rich RNA sequence binding factor 1	
Hs.445733	NM_002093	GSK3B	3	down						Glycogen synthase kinase 3 beta	
Hs.181244	NM_002116	HLA-A	6;2;19		down		down			Major histocompatibility complex, class I, A	
Hs.32309	NM_002194	INPP1	2						down	Inositol polyphosphate-1-phosphatase	
Hs.374097	NM_002199	IRF2	4				down			Interferon regulatory factor 2	
Hs.25292	NM_002229	JUNB	19		down				down	Jun B proto-oncogene	
Hs.408960	NM_002241	KCNJ10	1						down	Potassium inwardly-rectifying channel, subfamily J, member 10	
Hs.41696	NM_002277	KRTHA1	17		down					Keratin, hair, acidic, 1	
Hs.159590	NM_002347	LY6H	8		up		up			Lymphocyte antigen 6 complex, locus H	
Hs.388613	NM_002499	NEO1	15					up		Neogenin homolog 1 (chicken)	
Hs.444934	NM_002525	NRD1	1	up						Nardilysin (N-arginine dibasic convertase)	
Hs.153952	NM_002526	NT5E	6;3						down	5'-nucleotidase, ecto (CD73)	
Hs.467701	NM_002539	ODC1	2	up						Ornithine decarboxylase 1	
Hs.435714	NM_002576	PAK1	11		up					P21/Cdc42/Rac1-activated kinase 1 (STE20 homolog, yeast)	
Hs.483564	NM_002622	PFDN1	5			down				Prefoldin 1	
Hs.464071	NM_002631	PGD	1	up						Phosphogluconate dehydrogenase	
Hs.468415	NM_002643	PIGF	2	up		up				Phosphatidylinositol glycan, class F	
Hs.444975	NM_002656	PLAGL1	6		up				up	Pleiomorphic adenoma gene-like 1	
Hs.2868	NM_002677	PMP2	8		down					Peripherial myelin protein 2	
Hs.321234	NM_002685	EXOSC10	1	up						Exosome component 10	
Hs.331420	NM_002703	PPAT	4		down				down	Phosphoribosyl pyrophosphate amidotransferase	
Hs.484371	NM_002752	MAPK9	5	down		down				Mitogen-activated protein kinase 9	
Hs.461777	NM_002768	PCOLN3	16	up	up			up		Procollagen (type III) N-endopeptidase	
Hs.368121	NM_002788	PSMA3	14		down					Proteasome (prosome, macropain) subunit, alpha type, 3	
Hs.446260	NM_002791	PSMA6	14	up						Proteasome (prosome, macropain) subunit, alpha type, 6	
Hs.422990	NM_002797	PSMB5	14;Y; X					up		Proteasome (prosome, macropain) subunit, beta type, 5	

FIG. 1 (Sheet 15 of 45)

Annotation			Direction of Change in SZ relative to Controls							
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.386866	NM_002841	PTPRG	3						down	Protein tyrosine phosphatase, receptor type, G
Hs.127657	NM_002852	PTX3	3		up					Pentaxin-related gene, rapidly induced by IL-1 beta
Hs.521640	NM_002874	RAD23B	9		up			down		RAD23 homolog B (S. cerevisiae)
Hs.148178	NM_002885	RAP1GAI	1		down					RAP1, GTPase activating protein 1
Hs.423935	NM_002904	RDBP	11;6	up	down					RD RNA binding protein
Hs.370620	NM_002908	REL	2		down					V-rel reticuloendotheliosis viral oncogene homolog (avian)
Hs.115474	NM_002915	RFC3	13		up					Replication factor C (activator 1) 3, 38kDa
Hs.23978	NM_002967	SAFB	19;Y		down					Scaffold attachment factor B
Hs.135787	NM_002968	SALL1	16				down			Sal-like 1 (Drosophila)
Hs.280202	NM_002972	SBF1	22	up						SET binding factor 1
Hs.465924	NM_003000	SDHB	1	up						Succinate dehydrogenase complex, subunit B, iron sulfur (Ip)
Hs.433795	NM_003029	SHC1	1;6	up						SHC (Src homology 2 domain containing) transforming protein 1
Hs.323878	NM_003038	SLC1A4	2				down			Solute carrier family 1 (glutamate/neutral amino acid transporter), member 4
Hs.443874	NM_003042	SLC6A1	3					down		Solute carrier family 6 (neurotransmitter transporter, GABA), member 1
Hs.334629	NM_003063	SLN	11		up					Sarcosin
Hs.360174	NM_003068	SNAI2	8		up					Snail homolog 2 (Drosophila)
Hs.1063	NM_003093	SNRPC	6	up						Small nuclear ribonucleoprotein polypeptide C
Hs.185597	NM_003119	SPG7	16		down					Spastic paraplegia 7, paraplegin (pure and complicated autosomal recessive)
Hs.443861	NM_003137	SRPK1	6			up				SFRS protein kinase 1
Hs.288229	NM_003165	STXBP1	9				up			Syntaxin binding protein 1
Hs.482390	NM_003243	TGFBR3	1	down						Transforming growth factor, beta receptor III (betaglycan, 300kDa)
Hs.104839	NM_003255	TIMP2	17	up						Tissue inhibitor of metalloproteinase 2
Hs.332173	NM_003260	TLE2	19						up	Transducin-like enhancer of split 2 (E(sp1) homolog, Drosophila)
Hs.432424	NM_003291	TPP2	13		down					Tripeptidyl peptidase II
Hs.12084	NM_003321	TUFM	16	up						Tu translation elongation factor, mitochondrial
Hs.439672	NM_003378	VGF	7			down			down	VGF nerve growth factor inducible
Hs.388927	NM_003403	YY1	14			up			up	YY1 transcription factor

FIG. 1 (Sheet 16 of 45)

UniGene ID	Annotation				Direction of Change in SZ relative to Controls						
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name	
Hs.399810	NM_003422	ZNF42	19		up					Zinc finger protein 42 (myeloid-specific retinoic acid-responsive)	
Hs.172979	NM_003451	ZNF177	19	up						Zinc finger protein 177	
Hs.144442	NM_003561	PLA2G10	16	up						Phospholipase A2, group X	
Hs.36958	NM_003567	BCAR3	1		down		down			Breast cancer anti-estrogen resistance 3	
Hs.371698	NM_003610	RAE1	20	up						RAE1 RNA export 1 homolog (S. pombe)	
Hs.250072	NM_003615	SLC4A7	3		up					Solute carrier family 4, sodium bicarbonate cotransporter, member 7	
Hs.104925	NM_003633	ENC1	5				up			Ectodermal-neural cortex (with BTB-like domain)	
Hs.104576	NM_003654	CHST1	11	up						Carbohydrate (keratan sulfate Gal-6) sulfotransferase 1	
Hs.400556	NM_003657	BCAS1	20		up					Breast carcinoma amplified sequence 1	
Hs.161181	NM_003675	PRPF18	10		down		down			PRP18 pre-mRNA processing factor 18 homolog (yeast)	
Hs.22393	NM_003677	DENR	12							Density-regulated protein	
Hs.213264	NM_003680	YARS	1				up			Tyrosyl-tRNA synthetase	
Hs.284491	NM_003681	PDXX	21				up			Pyridoxal (pyridoxine, vitamin B6) kinase	
Hs.470608	NM_003705	SLC25A12	2							Solute carrier family 25 (mitochondrial carrier, Aralar), member 12	
Hs.484222	NM_003729	RTCD1	1		down					RNA terminal phosphate cyclase domain 1	
Hs.4742	NM_003801	GPAAI	8;6	up						GPAAI anchor attachment protein 1 homolog (yeast)	
Hs.178748	NM_003813	ADAM21	14	down						A disintegrin and metalloproteinase domain 21	
Hs.169900	NM_003819	PABPC4	1;15;X		up					Poly(A) binding protein, cytoplasmic 4 (inducible form)	
Hs.445511	NM_003831	RIOK3	18		down					RIO kinase 3 (yeast)	
Hs.430551	NM_003870	IQGAP1	15				down			IQ motif containing GTPase activating protein 1	
Hs.7165	NM_003904	ZNF259			up					Zinc finger protein 259	
Hs.429180	NM_003908	EIF2S2	20;2	up	down				up	Eukaryotic translation initiation factor 2, subunit 2 beta, 38kDa	
Hs.121592	NM_003916	AP1S2	X	down						Adaptor-related protein complex 1, sigma 2 subunit	
Hs.371199	NM_003919	SGCE	7				down			Sarcoglycan, epsilon	
Hs.200770	NM_003930	SCAP2	7			up				Src family associated phosphoprotein 2	
Hs.158460	NM_003936	CDK5R2	2	up	down					Cyclin-dependent kinase 5, regulatory subunit 2 (p39)	
Hs.143728	NM_003941	WASL	7			up				Wiskott-Aldrich syndrome-like	
Hs.47357	NM_003956	CH25H	10	down						Cholesterol 25-hydroxylase	
Hs.463439	NM_003971	SPAG9	17		down					Sperm associated antigen 9	
Hs.315369	NM_004028	AQP4	18		down					Aquaporin 4	

FIG. 1 (Sheet 17 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.412117	NM_004033	ANXA6	5;3				up			Annexin A6
Hs.467898	NM_004036	ADCY3	2				up			Adenylate cyclase 3
Hs.433732	NM_004071	CLK1	2				down			CDC-like kinase 1
Hs.292549	NM_004087	DLG1	3	up	down		down			Discs, large homolog 1 (Drosophila)
Hs.202095	NM_004098	EMX2	10					down		Empty spiracles homolog 2 (Drosophila)
Hs.7557	NM_004117	FKBP5	6					up		FK506 binding protein 5
Hs.172791	NM_004182	UXT	X					up		Ubiquitously-expressed transcript
Hs.484703	NM_004233	CD83	6	down						CD83 antigen (activated B lymphocytes, immunoglobulin superfamily)
Hs.376206	NM_004235	KLF4	9						down	Kruppel-like factor 4 (gut)
Hs.465985	NM_004317	ASNA1	19				up			ArsA arsenite transporter, ATP-binding, homolog 1 (bacterial)
Hs.471401	NM_004328	BCS1L	2	up		up				BCS1-like (yeast)
Hs.131226	NM_004331	BNIP3L	8			up				BCL2/adenovirus E1B 19kDa interacting protein 3-like
Hs.13291	NM_004354	CCNG2	4			up				Cyclin G2
Hs.220529	NM_004363	CEACAM5	19;4		up			up		Carcinoembryonic antigen-related cell adhesion molecule 5
Hs.129452	NM_004392	DACHI	13		up					Dachshund homolog 1 (Drosophila)
Hs.408461	NM_004397	DDX6	11			down				DEAD (Asp-Glu-Ala-Asp) box polypeptide 6
Hs.171695	NM_004417	DUSP1	5;16;11		down				down	Dual specificity phosphatase 1
Hs.2128	NM_004419	DUSP5	10						down	Dual specificity phosphatase 5
Hs.371218	NM_004438	EPHA4	2				up			EPH receptor A4
Hs.213389	NM_004487	GOLGB1	3	up						Golgi autoantigen, golgin subfamily b, macrogolgin (with transmembrane signal), 1
Hs.248746	NM_004499	HNRPAB	5		down					Heterogeneous nuclear ribonucleoprotein A/B
Hs.472185	NM_004552	NDUFS5	1				up			NADH dehydrogenase (ubiquinone) Fe-S protein 5, 15kDa (NADH-coenzyme Q reductase)
Hs.408257	NM_004553	NDUFS6	5	up						NADH dehydrogenase (ubiquinone) Fe-S protein 6, 13kDa (NADH-coenzyme Q reductase)
Hs.119316	NM_004564	PET112L	4	up						PET112-like (yeast)
Hs.283454	NM_004569	PIGH	14			up		up		Phosphatidylinositol glycan, class H
Hs.296169	NM_004578	RAB4A	1	up		up				RAB4A, member RAS oncogene family
Hs.129783	NM_004588	SCN2B	11						up	Sodium channel, voltage-gated, type II, beta
Hs.433201	NM_004642	CDK2AP1	12;2		down					CDK2-associated protein 1

FIG. 1 (Sheet 18 of 45)

Annotation				Direction of Change in SZ relative to Controls					Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLFFC	Nacc	PC	STG	Name
Hs.106674	NM_004656	BAP1	3	up						BRCA1 associated protein-1 (ubiquitin carboxy-terminal hydrolase)
Hs.194695	NM_004675	ARHI	1						up	Ras homolog gene family, member 1
Hs.106876	NM_004691	ATP6V0D1	16				up			ATPase, H ⁺ transporting, lysosomal 38kDa, V0 subunit d isoform 1
Hs.471779	NM_004735	LRRFIP1	2		up					Leucine rich repeat (in FLII) interacting protein 1
Hs.2210	NM_004773	TRIP3	17				up			Thyroid hormone receptor interactor 3
Hs.464848	NM_004775	B4GALT6	18			up				UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 6
Hs.370487	NM_004776	B4GALT5	20				up			UDP-Gal:betaGlcNAc beta 1,4- galactosyltransferase, polypeptide 5
Hs.26703	NM_004779	CNOT8	5	up	down					CCR4-NOT transcription complex, subunit 8
Hs.29802	NM_004787	SLIT2	4				up			Slit homolog 2 (Drosophila)
Hs.483238	NM_004815	PARG1	1	down				down		PTPL1-associated RhoGAP 1
Hs.240395	NM_004823	KCNK6	19					up		Potassium channel, subfamily K, member 6
Hs.480218	NM_004827	ABCG2	4	down						ATP-binding cassette, sub-family G (WHITE), member 2
Hs.431109	NM_004853	STX8	17	up	down					Syntaxin 8
Hs.408515	NM_004883	NRG2	5	down				up		Neuregulin 2
Hs.282901	NM_004902	RNPC2	20;X	up						RNA-binding region (RNP1, RRM) containing 2
Hs.446091	NM_004906	WTAP	6					down		Wilms tumor 1 associated protein
Hs.164410	NM_004913	C16orf7	16;6	up						Chromosome 16 open reading frame 7
Hs.2171	NM_004962	GDF10	10		down		down			Growth differentiation factor 10
Hs.303870	NM_004976	KCNC1	11	down						Potassium voltage-gated channel, Shaw-related subfamily, member 1
Hs.32505	NM_004981	KCNJ4	22				down			Potassium inwardly-rectifying channel, subfamily J, member 4
Hs.151219	NM_004984	KIF5A	12				up			Kinesin family member 5A
Hs.149387	NM_004999	MYO6	6		down		down		down	Myosin VI
Hs.45002	NM_005052	RAC3	17						up	Ras-related C3 botulinum toxin substrate 3 (rho family, small GTP binding protein Rac3)
Hs.46440	NM_005075	SLCO1A2	12				down			Solute carrier organic anion transporter family, member 1A2
Hs.287362	NM_005078	TLE3	15						up	Transducin-like enhancer of split 3 (E(sp1) homolog, Drosophila)
Hs.183428	NM_005086	SSPN	12		down					Sarcospan (Kras oncogene-associated gene)

FIG. 1 (Sheet 19 of 45)

Annotation		Direction of Change in SZ relative to Controls									
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name	
Hs.356820	NM_005108	XYLB	3		up					Xylulokinase homolog (H. influenzae)	
Hs.37288	NM_005126	NR1D2	3			up				Nuclear receptor subfamily 1, group D, member 2	
Hs.1540	NM_005131	THOC1	18				down		down	THO complex 1	
Hs.464595	NM_005134	PPP4R1	18		down					Protein phosphatase 4, regulatory subunit 1	
Hs.410944	NM_005138	SCO2	22					up		SCO cytochrome oxidase deficient homolog 2 (yeast)	
Hs.502883	NM_005146	SART1	11	up						Squamous cell carcinoma antigen recognised by T cells	
Hs.369438	NM_005238	ETS1	11					down		V-ets erythroblastosis virus E26 oncogene homolog 1 (avian)	
Hs.481371	NM_005245	FAT	4					down		FAT tumor suppressor homolog 1 (Drosophila)	
Hs.25647	NM_005252	FOS	14;22;11		down				down	V-fos FBJ murine osteosarcoma viral oncogene homolog	
Hs.483305	NM_005340	HINT1	5	up				up		Histidine triad nucleotide binding protein 1	
Hs.2780	NM_005354	JUND	19;5					down		Jun D proto-oncogene	
Hs.380742	NM_005393	PLXNB3	X						down	Plexin B3	
Hs.303090	NM_005398	PPP1R3C	10				down			Protein phosphatase 1, regulatory (inhibitor) subunit 3C	
Hs.50732	NM_005399	PRKAB2	1					up	up	Protein kinase, AMP-activated, beta 2 non-catalytic subunit	
Hs.129727	NM_005431	XRCC2	7		down					X-ray repair complementing defective repair in Chinese hamster cells 2	
Hs.194148	NM_005433	YES1	18;2				down			V-yes-1 Yamaguchi sarcoma viral oncogene homolog 1	
Hs.368610	NM_005443	PAPSS1	4	up						3'-phosphoadenosine 5'-phosphosulfate synthase 1	
Hs.198612	NM_005458	GPR51	9				up			G protein-coupled receptor 51	
Hs.278500	NM_005471	GNPDA1	5;9			up				Glucosamine-6-phosphate deaminase 1	
Hs.429294	NM_005502	ABCA1	9				down			ATP-binding cassette, sub-family A (ABCI), member 1	
Hs.153299	NM_005510	DOM3Z	6					up		Dom-3 homolog Z (C. elegans)	
Hs.250666	NM_005524	HES1	3						down	Hairy and enhancer of split 1, (Drosophila)	
Hs.471508	NM_005544	IRS1	2				up			Insulin receptor substrate 1	
Hs.2795	NM_005566	LDHA	11;17;22;10;2				up			Lactate dehydrogenase A	
Hs.34560	NM_005574	LMO2	11			down		down		LIM domain only 2 (rhombotin-like 1)	
Hs.18069	NM_005606	LGMN	14			up				Legumain	
Hs.484324	NM_005649	ZNF354A	5				down			Zinc finger protein 354A	
Hs.183671	NM_005651	TDO2	4					up		Tryptophan 2,3-dioxygenase	
Hs.458917	NM_005697	SCAMP2	15						down	Secretory carrier membrane protein 2	

FIG. 1 (Sheet 20 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.21577	NM_005701	RNUT1	15					up		RNA, U transporter 1
Hs.118118	NM_005723	TM4SF9	4			up				Transmembrane 4 superfamily member 9
Hs.381072	NM_005729	PIPF	10	up	down					Peptidylprolyl isomerase F (cyclophilin F)
Hs.124553	NM_005741	ZNF263	16		down					Zinc finger protein 263
Hs.123464	NM_005767	P2RY5	13		down		down			Purinergic receptor P2Y, G-protein coupled, 5
Hs.424126	NM_005770	SERF2	15;17; 7;6	up						Small EDRK-rich factor 2
Hs.294603	NM_005776	CNIH1	14	up						Cornichon homolog (Drosophila)
Hs.188879	NM_005777	RBM6	3;19						down	RNA binding motif protein 6
Hs.436922	NM_005798	RFP2	13	down	down			down	down	Ret finger protein 2
Hs.440168	NM_005822	DSCR1L1	6				up			Down syndrome critical region gene 1-like 1
Hs.436944	NM_005841	SPRY1	4	down	down					Sprouty homolog 1, antagonist of FGF signaling (Drosophila)
Hs.18676	NM_005842	SPRY2	13;3	down						Sprouty homolog 2 (Drosophila)
Hs.432862	NM_005885	MARCH-VI	5			up				Membrane-associated RING-CH protein VI
Hs.186486	NM_005923	MAP3K5	6					down		Mitogen-activated protein kinase kinase kinase 5
Hs.433391	NM_005950	MT1G	16;12				down	down		Metallothionein 1G
Hs.435974	NM_005956	MTHFD1	14;2		down					Methylenetetrahydrofolate dehydrogenase (NADP+ dependent), methylenetetrahydrofolate cyclohydrolase, formyltetrahydrofolate synthetase
Hs.107474	NM_005966	NAB1	2;4;3			up				NGFI-A binding protein 1 (EGR1 binding protein 1)
Hs.2430	NM_005997	TCFL1	1					up		Transcription factor-like 1
Hs.170107	NM_006003	UQCRFS1	19					up		Ubiquinol-cytochrome c reductase, Rieske iron-sulfur polypeptide 1
Hs.406096	NM_006007	ZA20D2	9;19; 6;17	up						Zinc finger, A20 domain containing 2
Hs.436446	NM_006010	ARMET	3	up				up		Arginine-rich, mutated in early stage tumors
Hs.412842	NM_006023	C10orf7	10		down					Chromosome 10 open reading frame 7
Hs.203620	NM_006040	HS3ST4							up	Heparan sulfate (glucosamine) 3-O-sulfotransferase 4
Hs.115830	NM_006043	HS3ST2	16					up		Heparan sulfate (glucosamine) 3-O-sulfotransferase 2
Hs.503692	NM_006106	YAP1	11		down					Yes-associated protein 1, 65kDa
Hs.367854	NM_006109	SKB1	14	up						SKB1 homolog (S. pombe)
Hs.15250	NM_006117	PBC1	6		down					Peroxisomal D3,D2-enoyl-CoA isomerase
Hs.369068	NM_006141	DNCL12	16;2			up				Dynein, cytoplasmic, light intermediate polypeptide 2
Hs.1565	NM_006154	NEDD4	15	down	up					Neural precursor cell expressed, developmentally down-

FIG. 1 (Sheet 21 of 45)

UniGene ID	Annotation		Direction of Change in SZ relative to Controls							Name
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	
Hs.339831	NM_006211	PENK	8							regulated 4
Hs.38449	NM_006216	SERPINE2	2					up		Proenkephalin
Hs.334868	NM_006246	PPP2R5E	14					up		Serine (or cysteine) proteinase inhibitor, clade E (nexin, plasminogen activator inhibitor type 1), member 2
Hs.43322	NM_006251	PRKAA1	5		down				down	Protein phosphatase 2, regulatory subunit B (B56), epsilon isoform
Hs.375001	NM_006289	TLN1	9				down			Protein kinase, AMP-activated, alpha 1 catalytic subunit
Hs.25313	NM_006337	MCRS1	12			down				Talin 1
Hs.465784	NM_006351	TIMM44	19		up					Microspherule protein 1
Hs.132902	NM_006366	CAP2	6					up		Translocase of inner mitochondrial membrane 44 homolog (yeast)
Hs.376064	NM_006392	NOL5A	20;6		down					CAP, adenylate cyclase-associated protein, 2 (yeast)
Hs.11417	NM_006423	RABAC1	19			up				Nucleolar protein 5A (56kDa with KKE/D repeat)
Hs.435342	NM_006425	SLU7	5		up					Rab acceptor 1 (prenylated)
Hs.421509	NM_006430	CCT4	2	up					up	Step II splicing factor SLU7
Hs.433222	NM_006432	NPC2	14;11;3					down		Chaperonin containing TCP1, subunit 4 (delta)
Hs.109752	NM_006443	C6orf108	6;20	up						Niemann-Pick disease, type C2
Hs.14894	NM_006464	TGOLN2	2			up				Chromosome 6 open reading frame 108
Hs.439153	NM_006502	POLH	6		up					Trans-golgi network protein 2
Hs.530045	NM_006524	ZNF138	7	down						Polymerase (DNA directed), eta
Hs.444558	NM_006558	KHDRBS3	8					up		Zinc finger protein 138 (clone pHZ-32)
Hs.309288	NM_006561	CUGBP2	10					up		KH domain containing, RNA binding, signal transduction associated 3
Hs.368367	NM_006565	CTCF	16		down					CUG triplet repeat, RNA binding protein 2
Hs.159525	NM_006569	CGREF1	2					up		CCCCTC-binding factor (zinc finger protein)
Hs.45127	NM_006574	CSPG5	3					up	down	Cell growth regulator with EF hand domain 1
Hs.155090	NM_006578	GNB5	15					up		Chondroitin sulfate proteoglycan 5 (neuroglycan C)
Hs.30696	NM_006602	TCFL5	20					down		Guanine nucleotide binding protein (G protein), beta 5
Hs.412870	NM_006627	POP4	19;9					up		Transcription factor-like 5 (basic helix-loop-helix)
Hs.415846	NM_006657	FTCD	21	down						Processing of precursor 4, ribonuclease P/MRP subunit (S. cerevisiae)
Hs.227011	NM_006658	C7orf16	7	up					up	Formiminotransferase cyclodeaminase
Hs.199743	NM_006680	ME3	11					up		Chromosome 7 open reading frame 16
										Malic enzyme 3, NADP(+)-dependent, mitochondrial

FIG. 1 (Sheet 22 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.2207	NM_006685	PROL3	4	down						Proline rich 5 (salivary)
Hs.6396	NM_006694	JTB	1	up						Jumping translocation breakpoint
Hs.225949	NM_006707	BTNL3	5	down	down					Butyrophilin-like 3
Hs.146804	NM_006717	SPIN	9				up			Spindlin
Hs.505662	NM_006741	PPP1R1A	12			up				Protein phosphatase 1, regulatory (inhibitor) subunit 1A
Hs.301404	NM_006743	RBM3	X		down					RNA binding motif (RNPI, RRM) protein 3
Hs.109655	NM_006746	SCML1	X						down	Sex comb on midleg-like 1 (Drosophila)
Hs.192686	NM_006749	SLC20A2	8		down		down			Solute carrier family 20 (phosphate transporter), member 2
Hs.337295	NM_006819	STIP1	11						up	Stress-induced-phosphoprotein 1 (Hsp70/Hsp90-organizing protein)
Hs.260903	NM_006874	ELF2	4	up						E74-like factor 2 (ets domain transcription factor)
Hs.436405	NM_006899	IDH3B	20;6;1		down					Isocitrate dehydrogenase 3 (NAD+) beta
Hs.466848	NM_006905	PSG1	19		up					Pregnancy specific beta-1-glycoprotein 1
Hs.435274	NM_006922	SCN3A	2					down		Sodium channel, voltage-gated, type III, alpha
Hs.376984	NM_006941	SOX10	22				down			SRX (sex determining region Y)-box 10
Hs.237825	NM_006947	SRP72	4;16			up				Signal recognition particle 72kDa
Hs.534115	NM_006988	ADAMTS1	21		down				down	A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1
Hs.30246	NM_006996	SLC19A2	1	down						Solute carrier family 19 (thiamine transporter), member 2
Hs.29353	NM_007030	TPPP	5;20		down					Brain-specific protein p25 alpha
Hs.125750	NM_007035	KERA	12		up					Keratocan
Hs.43670	NM_007054	KIF3A	5				up			Kinesin family member 3A
Hs.436896	NM_007055	POLR3A	10			up				Polymerase (RNA) III (DNA directed) polypeptide A, 155kDa
Hs.474797	NM_007061	CDC42EP1	22			up				CDC42 effector protein (Rho GTPase binding) 1
Hs.160958	NM_007065	CDC37	19				up			CDC37 cell division cycle 37 homolog (S. cerevisiae)
Hs.142245	NM_007071	HHLA3	1	up						HERV-H LTR-associating 3
Hs.269512	NM_007085	FSTL1	3		down					Follistatin-like 1
Hs.106857	NM_007088	CALB2	16;19;15				up			Calbindin 2, 29kDa (calretinin)
Hs.484241	NM_007097	CLTB	5				up			Clathrin, light polypeptide (Lcb)
Hs.119014	NM_007147	ZNF175	19		down			down		Zinc finger protein 175
Hs.158174	NM_007149	ZNF184	6		down					Zinc finger protein 184 (Kruppel-like)
Hs.434283	NM_007167	ZNF258	1		up					Zinc finger protein 258

FIG. 1 (Sheet 23 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.205163	NM_007208	MRPL3	3		down					Mitochondrial ribosomal protein L3
Hs.388668	NM_007246	KLHL2	4				up			Kelch-like 2, Mayven (<i>Drosophila</i>)
Hs.377070	NM_007325	GRIA3	X				up			Glutamate receptor, ionotropic, AMPA 3
Hs.189826	NM_007345	ZNF236	18						up	Zinc finger protein 236
Hs.240770	NM_007362	NCBP2	3:1		down					Nuclear cap binding protein subunit 2, 20kDa
Hs.500104	NM_012095	AP3M1	10	up						Adaptor-related protein complex 3, mu 1 subunit
Hs.204041	NM_012111	AHSA1	14:9	up	up			up		AHA1, activator of heat shock 90kDa protein ATPase homolog 1 (yeast)
Hs.212395	NM_012127	CIZ1	9	up						CDKN1A interacting zinc finger protein 1
Hs.24178	NM_012155	EML2	19		up					Echinoderm microtubule associated protein like 2
Hs.438454	NM_012173	FBXO25	8			up		up		F-box protein 25
Hs.22867	NM_012199	EIF2C1	1	down						Eukaryotic translation initiation factor 2C, 1
Hs.179915	NM_012202	GNG3	11				up			Guanine nucleotide binding protein (G protein), gamma 3
Hs.155742	NM_012203	GRHRP	9:17		down					Glyoxylate reductase/hydroxypyruvate reductase
Hs.494804	NM_012212	LTB4DH	9			up				Leukotriene B4 12-hydroxydehydrogenase
Hs.481181	NM_012224	NEK1	4	down						NIMA (never in mitosis gene a)-related kinase 1
Hs.369779	NM_012238	SIRT1	10					down		Sirtuin (silent mating type information regulation 2 homolog) 1 (<i>S. cerevisiae</i>)
Hs.309583	NM_012254	SLC27A5	19	up	up					Solute carrier family 27 (fatty acid transporter), member 5
Hs.144287	NM_012259	HEY2	6	down						Hairy/enhancer-of-split related with YRPW motif 2
Hs.22920	NM_012261	C20orf103	20						down	Chromosome 20 open reading frame 103
Hs.21703	NM_012281	KCND2	7				up			Potassium voltage-gated channel, Shal-related subfamily, member 2
Hs.433057	NM_012304	FBXL7	5:6		up					F-box and leucine-rich repeat protein 7
Hs.397918	NM_012311	KIN	10						down	KIN, antigenic determinant of recA protein homolog (mouse)
Hs.370040	NM_012333	MYCBP	1	down	up					C-myc binding protein
Hs.470417	NM_012392	PEF	1	up						PEF protein with a long N-terminal hydrophobic domain (peflin)
Hs.430742	NM_012395	PFTK1	7		up					PFTAIRE protein kinase 1
Hs.167496	NM_012416	RANBP6	9				up			RAN binding protein 6
Hs.166313	NM_012419	RGS17	6			up				Regulator of G-protein signaling 17
Hs.471011	NM_012433	SF3B1	2			up				Splicing factor 3b, subunit 1, 155kDa
Hs.102735	NM_012446	SSBP2	5	down						Single-stranded DNA binding protein 2
Hs.235750	NM_012456	TIMM10	11:17		down					Translocase of inner mitochondrial membrane 10 homolog

FIG. 1 (Sheet 24 of 45)

UniGene ID		Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name		
Hs.59757	NM_012482	ZNF281	1							(yeast)		
Hs.115721	NM_013247	PRSS25	2	up				down		Zinc finger protein 281		
Hs.292156	NM_013253	DKK3	11		up					Protease, serine, 25		
Hs.476291	NM_013286	RBM15B	3	down						Dickkopf homolog 3 (Xenopus laevis)		
Hs.517436	NM_013313	YPBL1	22	up		up				RNA binding motif protein 15B		
Hs.433151	NM_013343	LOH3CR2A	3;1		up					Yippee-like 1 (Drosophila)		
Hs.40098	NM_013372	GREM1	15		up					Loss of heterozygosity, 3, chromosomal region 2, gene A		
Hs.279877	NM_013393	FTSJ2	7		down					Gremlin 1 homolog, cysteine knot superfamily (Xenopus laevis)		
Hs.7765	NM_013399	C16orf5	16		up					FtsJ homolog 2 (E. coli)		
Hs.416049	NM_013433	TNPO2	19			down				Chromosome 16 open reading frame 5		
Hs.408515	NM_013985	NRG2	5	down				up		Transporthin 2 (importin 3, karyopherin beta 2b)		
Hs.15400	NM_014015	DEXI	16				up			Neuregulin 2		
Hs.13370	NM_014044	UNC50	2		down					Dexamethasone-induced transcript		
Hs.436500	NM_014063	DBNL	7						up	Unc-50 homolog (C. elegans)		
Hs.368971	NM_014071	NCOA6	20	up						Drebrin-like		
Hs.333823	NM_014078	MRPL13	8;18		down	up	up			Nuclear receptor coactivator 6		
Hs.272215	NM_014079	KLF15	3		down					Mitochondrial ribosomal protein L13		
Hs.11614	NM_014157	HSPC065			down	down				Kruppel-like factor 15		
Hs.127496	NM_014170	HSPC135	3				up			HSPC065 protein		
Hs.18349	NM_014175	MRPL15	8;12		down					HSPC135 protein		
Hs.445890	NM_014184	HSPC163	1		down	up				Mitochondrial ribosomal protein L15		
Hs.233458	NM_014223	NFYC	1	up						HSPC163 protein		
Hs.467192	NM_014225	PPP2R1A	19				up			Nuclear transcription factor Y, gamma		
Hs.114062	NM_014241	PTPLA	10		up					Protein phosphatase 2 (formerly 2A), regulatory subunit A (PR 65), alpha isoform		
Hs.437277	NM_014275	MGAT4B	5;19;10	up						Protein tyrosine phosphatase-like (proline instead of catalytic arginine), member a		
Hs.301760	NM_014286	FREQ	9	down		down				Mannosyl (alpha-1,3-)-glycoprotein beta-1,4-N-acetylglucosaminyltransferase, isoenzyme B		
Hs.193842	NM_014290	TDRD7	9		down					Frequenin homolog (Drosophila)		
Hs.282998	NM_014309	RBM9	22				up			Tudor domain containing 7		
Hs.523054	NM_014313	SMP1	11	up						RNA binding motif protein 9		
Hs.330384	NM_014325	CORO1C	12	up						NPD014 protein		
										Coronin, actin binding protein, 1C		

FIG. 1 (Sheet 25 of 45)

Annotation				Direction of Change in SZ relative to Controls					Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPEC	Nacc	PC	STG	Name
Hs.370510	NM_014333	IGSF4	11	down	up					Immunoglobulin superfamily, member 4
Hs.189810	NM_014351	SULT4A1	22				up			Sulfotransferase family 4A, member 1
Hs.310431	NM_014386	PKD2L2	5	down						Polycystic kidney disease 2-like 2
Hs.316890	NM_014426	SNX5	20				down			Sorting nexin 5
Hs.443577	NM_014452	TNFRSF21	6					up		Tumor necrosis factor receptor superfamily, member 21
Hs.232543	NM_014456	PDCD4	10		down					Programmed cell death 4 (neoplastic transformation inhibitor)
Hs.111632	NM_014463	LSM3	3			up				LSM3 homolog, U6 small nuclear RNA associated (S. cerevisiae)
Hs.438994	NM_014480	ZNF544	19				up			Zinc finger protein 544
Hs.221436	NM_014483	RBMS3	3	down						RNA binding motif, single stranded interacting protein
Hs.143600	NM_014498	GOLPH4	3	down						Golgi phosphoprotein 4
Hs.252682	NM_014506	TOR1B	9	up						Torsin family 1, member B (torsin B)
Hs.13014	NM_014570	ARFGAP3	22				down			ADP-ribosylation factor GTPase activating protein 3
Hs.525339	NM_014584	ERO1L	14			up				ERO1-like (S. cerevisiae)
Hs.180933	NM_014593	CXXC1	18					up		CXXC finger 1 (PHD domain)
Hs.368808	NM_014600	EHD3	2;6				up			EH-domain containing 3
Hs.26704	NM_014608	CYFIP1	15;9			down				Cytoplasmic FMR1 interacting protein 1
Hs.413801	NM_014614	PSME4	2		up					Proteasome (prosome, macropain) activator subunit 4
Hs.330073	NM_014686	KIAA0355	19	up						KIAA0355
Hs.196054	NM_014707	HDAC9	7					down		Histone deacetylase 9
Hs.410092	NM_014741	KIAA0652	11				down			KIAA0652 gene product
Hs.484349	NM_014757	MAML1	5			down				Mastermind-like 1 (Drosophila)
Hs.44024	NM_014763	MRPL19	2			up				Mitochondrial ribosomal protein L19
Hs.370530	NM_014788	TRIM14	9	down						Tripartite motif-containing 14
Hs.533245	NM_014829	DDX46	5					down		DEAD (Asp-Glu-Ala-Asp) box polypeptide 46
Hs.3094	NM_014876	KIAA0063	22	up						KIAA0063 gene product
Hs.130014	NM_014880	DCL-1	2				down			Type I transmembrane C-type lectin receptor DCL-1
Hs.227602	NM_014892	RBM16	6		down					RNA binding motif protein 16
Hs.131683	NM_014912	CPEB3	10			up				Cytoplasmic polyadenylation element binding protein 3
Hs.22109	NM_014952	BAHD1	15	up						Bromo adjacent homology domain containing 1
Hs.124490	NM_014977	ACIN1	14					down		Apoptotic chromatin condensation inducer 1
Hs.270499	NM_015001	SHARP	1			up				SMART/HDAC1 associated repressor protein
Hs.159699	NM_015002	FBXO21	12			up				F-box protein 21
Hs.189409	NM_015033	FNBP1	9						down	Formin binding protein 1

FIG. 1 (Sheet 26 of 45)

UniGene ID	Annotation				Direction of Change in SZ relative to Controls						
	Acc	Symbol	Chr.	AnCg	CB	DLFPC	Nacc	PC	STG	Name	
Hs.293593	NM_015071	ARHGAP26	5		up					Rho GTPase activating protein 26	
Hs.440414	NM_015087	SPG20	13		down					Spastic paraplegia 20, spartin (Troyer syndrome)	
Hs.269775	NM_015093	MAP3K7IP2	6	up						Mitogen-activated protein kinase kinase 7 interacting protein 2	
Hs.331431	NM_015200	SCC-112	4	down						SCC-112 protein	
Hs.410497	NM_015379	BRI3		up						Brain protein I3	
Hs.105547	NM_015392	NPDC1	9	up						Neural proliferation, differentiation and control, 1	
Hs.25956	NM_015464	SOSTDC1	7						up	Sclerostin domain containing 1	
Hs.472630	NM_015474	SAMHD1	20			down				SAM domain and HD domain 1	
Hs.146100	NM_015553	PIP3-E	6				up			Phosphoinositide-binding protein PIP3-E	
Hs.127401	NM_015662	SLB				up				Selective LIM binding factor, rat homolog	
Hs.369144	NM_015665	AAAS	12							Achalasia, adrenocortical insufficiency, alacrimia (Allgrove, triple-A)	
Hs.391481	NM_015693	PDZK6	4	up						PDZ domain containing 6	
Hs.474914	NM_015705	RUTBC3	22	down						RUN and TBC1 domain containing 3	
Hs.512592	NM_015713	RRM2B	8		up					Ribonucleotide reductase M2 B (TP53 inducible)	
Hs.235368	NM_015719	COL5A3	19	down			down			Collagen, type V, alpha 3	
Hs.250693	NM_015852	ZNF117	7;19	down						Kruppel-related zinc finger protein	
Hs.128959	NM_015885	PCF11	11				down			Pre-mRNA cleavage complex II protein Pcf11	
Hs.348326	NM_015894	STMN3	20	down						Stathmin-like 3	
Hs.414952	NM_015910	LOC51057	2		up					Hypothetical protein LOC51057	
Hs.16606	NM_015960	CUTC	10		down					CutC copper transporter homolog (E.coli)	
Hs.435759	NM_015963	THAP4	2	up						THAP domain containing 4	
Hs.44298	NM_015969	MRPS17	7	up						Mitochondrial ribosomal protein S17	
Hs.370703	NM_015974	CRYL1	13				down			Crystallin, lambda 1	
Hs.370168	NM_015986	CRLF3	17							Cytokine receptor-like factor 3	
Hs.525752	NM_015995	KLF13	15	up						Kruppel-like factor 13	
Hs.271876	NM_016010	CGI-62	8	down						CGI-62 protein	
Hs.183646	NM_016011	CGI-63	1			down				Nuclear receptor binding factor 1	
Hs.106529	NM_016013	NDUFAF1	15	up						NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, assembly factor 1	
Hs.514216	NM_016016	CGI-69	17	up						CGI-69 protein	
Hs.145386	NM_016033	CGI-90	8	down			down			CGI-90 protein	
Hs.3945	NM_016045	C20orf45	20				down			Chromosome 20 open reading frame 45	
Hs.483296	NM_016048	CGI-111	5						up	CGI-111 protein	

FIG. 1 (Sheet 27 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.271614	NM_016049	C14orf122	14	up						Chromosome 14 open reading frame 122
Hs.157401	NM_016058	CGI-121	2		up		up			CGI-121 protein
Hs.436161	NM_016067	MRPS18C	4		up					Mitochondrial ribosomal protein S18C
Hs.435952	NM_016082	CDK5RAP1	20			down				CDK5 regulatory subunit associated protein 1
Hs.25829	NM_016084	RASD1	17					up		RAS, dexamethasone-induced 1
Hs.108969	NM_016145	PTD008	19	up						PTD008 protein
Hs.159581	NM_016155	MMP17	12	up						Matrix metalloproteinase 17 (membrane-inserted)
Hs.446179	NM_016200	LSM8	7		up					LSM8 homolog, U6 small nuclear RNA associated (S. cerevisiae)
Hs.131133	NM_016203	PRKAG2	7		up				up	Protein kinase, AMP-activated, gamma 2 non-catalytic subunit
Hs.135756	NM_016218	POLK	5			up				Polymerase (DNA directed) kappa
Hs.328865	NM_016221	DCTN4	5		down					Dynactin 4 (p62)
Hs.191213	NM_016224	SNX9	6			up				Sorting nexin 9
Hs.208759	NM_016231	NLK	17		up					Nemo like kinase
Hs.148685	NM_016235	GPRC5B	16				down			G protein-coupled receptor, family C, group 5, member B
Hs.334832	NM_016243	NQO3A2	1				down			NAD(P)H:quinone oxidoreductase type 3, polypeptide A2
Hs.18788	NM_016246	DHRS10	19				down	down		Dehydrogenase/reductase (SDR family) member 10
Hs.125132	NM_016269	LEF1	4		up	down				Lymphoid enhancer-binding factor 1
Hs.278627	NM_016297	PCYOX1	2			up				Prenylcysteine oxidase 1
Hs.470887	NM_016315	GULP1	2					down		GULP, engulfment adaptor PTB domain containing 1
Hs.158530	NM_016339	RAPGEFL1	17	up						Rap guanine nucleotide exchange factor (GEF)-like 1
Hs.483329	NM_016340	RAPGEF6	5		up					KIAA1961 protein
Hs.224137	NM_016390	C9orf114	9		down					Chromosome 9 open reading frame 114
Hs.497967	NM_016396	HSPC129	15			up				Hypothetical protein HSPC129
Hs.283322	NM_016401	HSPC138	11			up			up	Hypothetical protein HSPC138
Hs.436502	NM_016486	LOC51249	1	down						Hypothetical protein LOC51249
Hs.408233	NM_016492	RANGNRF	17				up			RAN guanine nucleotide release factor
Hs.524094	NM_016505	PS1D	1	up						Putative S1 RNA binding domain protein
Hs.29549	NM_016511	CLEC1	12;X	down						C-type lectin-like receptor-1
Hs.478393	NM_016559	PEX5L	3					up		Peroxisomal biogenesis factor 5-like
Hs.475387	NM_016565	E2IG2	11			down				E2IG2 protein
Hs.200063	NM_016596	HDAC7A	12						down	Histone deacetylase 7A
Hs.433439	NM_016622	MRPL35				up				Mitochondrial ribosomal protein L35
Hs.478064	NM_016625	MGC12197	3					up		BM-011 protein

FIG. 1 (Sheet 28 of 45)

Annotation			Direction of Change in SZ relative to Controls							
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLFPC	Nacc	PC	STG	Name
Hs.465144	NM_016626	RKHD2	18			up				Ring finger and KH domain containing 2
Hs.278635	NM_016648	HDCMA18P	4		down					HDCMA18P protein
Hs.369284	NM_016649	C20orf6	20		down					Chromosome 20 open reading frame 6
Hs.171342	NM_016652	CRNKL1	20			up		up		Ctn, crooked neck-like 1 (Drosophila)
Hs.127792	NM_016941	DLL3	19						up	Delta-like 3 (Drosophila)
Hs.40735	NM_017412	FZD3	8					down		Frizzled homolog 3 (Drosophila)
Hs.286233	NM_017425	SPAI7	11;19						up	Sperm autoantigenic protein 17
Hs.108112	NM_017443	POLE3	9		down					Polymerase (DNA directed), epsilon 3 (p17 subunit)
Hs.437084	NM_017544	NKRF	X				up			NF-kappa B repressing factor
Hs.458304	NM_017578	ROPNI	3	down						Ropporin, raphilin associated protein 1
Hs.165636	NM_017594	DIRAS2	9			up	up			DIRAS family, GTP-binding RAS-like 2
Hs.258798	NM_017615	C10orf86	10	up	down		down			Chromosome 10 open reading frame 86
Hs.29700	NM_017665	ZCCHC10	5	down						Zinc finger, CCHC domain containing 10
Hs.369932	NM_017714	C20orf13	20						up	Chromosome 20 open reading frame 13
Hs.483993	NM_017734	PALMD	1						down	Palmdelphin
Hs.440401	NM_017750	FLJ20296	2		down				down	All-trans-13,14-dihydroretinol saturose
Hs.147836	NM_017768	FLJ20331	1					down		Hypothetical protein FLJ20331
Hs.444269	NM_017776	FLJ20344	X		up	up				Hypothetical protein FLJ20344
Hs.150122	NM_017784	OSBPL10	3			up	up			Oxysterol binding protein-like 10
Hs.368710	NM_017785	FLJ20364	5		down					Hypothetical protein FLJ20364
Hs.408652	NM_017794	KIAA1797	9				up	up		KIAA1797
Hs.371210	NM_017847	C1orf27	1			up			down	Chromosome 1 open reading frame 27
Hs.134406	NM_017853	TXNL4B	16		up				up	Thioredoxin-like 4B
Hs.105606	NM_017854	FLJ20512	19	up			up	down		Hypothetical protein FLJ20512
Hs.377705	NM_017865	FLJ20531	1	up						Hypothetical protein FLJ20531
Hs.525238	NM_017924	C14orf119	14	up		up		up		Chromosome 14 open reading frame 119
Hs.249591	NM_017925	C9orf55	9	down						Chromosome 9 open reading frame 55
Hs.30783	NM_017967	FLJ20850	19	up						Hypothetical protein FLJ20850
Hs.297044	NM_017987	RUFY2	10			down				RUN and FYVE domain containing 2
Hs.516341	NM_017991	FLJ10081	2						down	Hypothetical protein FLJ10081
Hs.128258	NM_017993	FLJ10094	13	up		up	up			Hypothetical protein FLJ10094
Hs.532296	NM_017998	C9orf40	9					up		Chromosome 9 open reading frame 40
Hs.241523	NM_018008	ZNF312	3					up	up	Zinc finger protein 312
Hs.445244	NM_018013	FLJ10159	6				up			Hypothetical protein FLJ10159
Hs.353454	NM_018045	FLJ10276	1	up						Hypothetical protein FLJ10276

FIG. 1 (Sheet 29 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.213393	NM_018046	VG5Q	5			up				Angiogenic factor VG5Q
Hs.262823	NM_018060	FLJ10326	1		down					Mitochondrial isoleucine tRNA synthetase
Hs.356096	NM_018067	FLJ10350	1	up						Hypothetical protein FLJ10350
Hs.270107	NM_018115	SDAD1	4			up				SDA1 domain containing 1
Hs.447458	NM_018121	C10orf6	10	up						Hypothetical protein LOC143286
Hs.31082	NM_018126	TMEM33	4					up		Transmembrane protein 33
Hs.26156	NM_018216	PANK4	1;2	up						Pantothenate kinase 4
Hs.174021	NM_018225	SMU1	9					up		Smu-1 suppressor of mec-8 and unc-52 homolog (C. elegans)
Hs.260238	NM_018238	FLJ10842	7;12;2			down				Hypothetical protein FLJ10842
Hs.502773	NM_018269	MTCBP-1	2	up						Membrane-type 1 matrix metalloproteinase cytoplasmic tail binding protein-1
Hs.368960	NM_018297	NGLY1	3				up			N-glycanase 1
Hs.503022	NM_018312	C11orf23	11					down		Chromosome 11 open reading frame 23
Hs.58382	NM_018322	C6orf64	6		down					Chromosome 6 open reading frame 64
Hs.458312	NM_018328	MBD5	2	down						Methyl-CpG binding domain protein 5
Hs.37558	NM_018339	RFK	9				up			Riboflavin kinase
Hs.176227	NM_018342	FLJ11155	4				down			Hypothetical protein FLJ11155
Hs.416755	NM_018357	FLJ11196	15		down					Acheron
Hs.271643	NM_018368	C6orf209	6		down					Chromosome 6 open reading frame 209
Hs.211828	NM_018428	HCA66	17	up						Hepatocellular carcinoma-associated antigen 66
Hs.32148	NM_018445	SELS	15		down					Selenoprotein S
Hs.477325	NM_018456	EAF2	3		up					ELL associated factor 2
Hs.370102	NM_018464	C10orf70	10				up			Chromosome 10 open reading frame 70
Hs.479766	NM_018475	TPARL	4						down	TPA regulated locus
Hs.12126	NM_018487	HCA112	7;20;12	down			down			Hepatocellular carcinoma-associated antigen 112
Hs.207433	NM_018557	LRP1B	2		down					Low density lipoprotein-related protein 1B (deleted in tumors)
Hs.435991	NM_018569	C4orf16	4						up	Chromosome 4 open reading frame 16
Hs.519346	NM_018695	ERBB2IP	5		down					ErbB2 interacting protein
Hs.47572	NM_018696	ELAC1	18		up					ElaC homolog 1 (E. coli)
Hs.104980	NM_018706	DHTKD1	10			up				Dehydrogenase E1 and transketolase domain containing 1
Hs.184062	NM_018840	C20orf24	20;10	up						Chromosome 20 open reading frame 24
Hs.145256	NM_018930	PCDHB10	5					down		Protocadherin beta 10

FIG. 1 (Sheet 30 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.190518	NM_018944	C21orf45	21					up		Chromosome 21 open reading frame 45
Hs.484686	NM_018988	GFOD1	6	up					up	Glucose-fructose oxidoreductase domain containing 1
Hs.440534	NM_019022	FLJ20793	18		down					FLJ20793 protein
Hs.288224	NM_019051	MRPL50	9			up				Mitochondrial ribosomal protein L50
Hs.481836	NM_019061	PIP3AP	5	up						Phosphatidylinositol-3-phosphate associated protein
Hs.140950	NM_019065	EFCBP2	16			down				EF hand calcium binding protein 2
Hs.173524	NM_019081	LKAP	16						up	Limkain b1
Hs.466714	NM_019088	PD2	19		down	down				Hypothetical protein F23149 1
Hs.323396	NM_019557	LOC56181	1	up						Hypothetical protein RP1-317E23
Hs.178011	NM_019606	FLJ20257	7	up						Hypothetical protein FLJ20257
Hs.443529	NM_019607	FLJ11267	8						down	Hypothetical protein FLJ11267
Hs.100890	NM_019845	RPRM	2						up	Reprimo, TP53 dependant G2 arrest mediator candidate
Hs.413083	NM_019863	F8	X		up					Coagulation factor VIII, procoagulant component (hemophilia A)
Hs.133512	NM_020119	ZC3HAV1		up						Zinc finger CCH type, antiviral 1
Hs.193226	NM_020121	UGCGL2	13					down		UDP-glucose ceramide glucosyltransferase-like 2
Hs.109929	NM_020137	GRIPAP1				down				GRIP1 associated protein 1
Hs.262858	NM_020143	LOC56902	2						up	Putative 28 kDa protein
Hs.460242	NM_020145	SH3GLB2	9			down				SH3-domain GRB2-like endophilin B2
Hs.160565	NM_020154	C15orf24	15		down					Chromosome 15 open reading frame 24
Hs.250456	NM_020162	DHX33	17	up						DEAH (Asp-Glu-Ala-His) box polypeptide 33
Hs.47649	NM_020166	MCCC1	3		down					Methylcrotonoyl-Coenzyme A carboxylase 1 (alpha)
Hs.42785	NM_020186	ACN9	7						up	ACN9 homolog (S. cerevisiae)
Hs.202011	NM_020198	GK001	17						down	GK001 protein
Hs.6434	NM_020215	C14orf132	14	up						Chromosome 14 open reading frame 132
Hs.127432	NM_020234	MDS009	15		up					X 009 protein
Hs.22065	NM_020239	CDC42SE1	1	up		up				CDC42 small effector 1
Hs.118241	NM_020247	CABC1	1	up						Chaperone, ABC1 activity of bcl complex like (S. pombe)
Hs.300404	NM_020314	MGC16824	16		down					Esophageal cancer associated protein
Hs.194408	NM_020340	KIAA1244	6	up						KIAA1244
Hs.477869	NM_020353	PLSCR4	3					down		Phospholipid scramblase 4
Hs.322901	NM_020368	SAS10	4			up				Disrupter of silencing 10
Hs.390623	NM_020383	XPNPEP1	10				up			X-prolyl aminopeptidase (aminopeptidase P) 1, soluble
Hs.387755	NM_020408	C6orf149	6			down				Chromosome 6 open reading frame 149
Hs.483841	NM_020466	DJ122O8.2	6		up				down	Hypothetical protein dJ122O8.2

FIG. 1 (Sheet 31 of 45)

Annotation		Direction of Change in SZ relative to Controls									
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPPC	Nacc	PC	STG	Name	
Hs.481545	NM_020546	ADCY2	5			up		up	up	Adenylate cyclase 2 (brain)	
Hs.104613	NM_020640	RP42	3			up				RP42 homolog	
Hs.529551	NM_020654	SENP7	3					down		SUMO1/sentrin specific protease 7	
Hs.123450	NM_020655	JPH3	16		down					Kelch domain containing 4	
Hs.287374	NM_020657	ZNF304	19		down					Zinc finger protein 304	
Hs.283816	NM_020660	CX36	15			down				Connexin-36	
Hs.47166	NM_020685	C3orf14	3;11				up			Chromosome 3 open reading frame 14	
Hs.211252	NM_020689	SLC24A3	20				up			Solute carrier family 24 (sodium/potassium/calcium exchanger), member 3	
Hs.526401	NM_020695	TCEB3BP1	19					down		Transcription elongation factor B polypeptide 3 binding protein 1	
Hs.17255	NM_020706	SFRS15	21		up					Splicing factor, arginine/serine-rich 15	
Hs.434947	NM_020727	ZNF295	21					down		Zinc finger protein 295	
Hs.211751	NM_020836	KIAA1446	14			down				Brain-enriched guanylate kinase-associated protein	
Hs.515351	NM_020895	KIAA1533	19	up						KIAA1533	
Hs.443891	NM_020925	KIAA1573	1		down					KIAA1573 protein	
Hs.8453	NM_020932	MAGEE1				down				Melanoma antigen, family E, 1	
Hs.368525	NM_020992	PDLIM1	10		up					PDZ and LIM domain 1 (elfin)	
Hs.243994	NM_021008	DEAF1	11				up			Deformed epidermal autoregulatory factor 1 (Drosophila)	
Hs.198760	NM_021076	NEFH	22;20				up			Neurofilament, heavy polypeptide 200kDa	
Hs.192215	NM_021116	ADCY1	7				up		up	Adenylate cyclase 1 (brain)	
Hs.437403	NM_021129	PP	10			up				Pyrophosphatase (inorganic)	
Hs.147119	NM_021135	RPS6KA2	6	up						Ribosomal protein S6 kinase, 90kDa, polypeptide 2	
Hs.154029	NM_021170	HES4	1			down				Hairy and enhancer of split 4 (Drosophila)	
Hs.119889	NM_021183	RAP2C	X					down		RAP2C, member of RAS oncogene family	
Hs.445489	NM_021200	PLEKHB1	11;8;12;15;20				down			Pleckstrin homology domain containing, family B (evecitins) member 1	
Hs.435106	NM_021205	RHOU	1						down	Ras homolog gene family, member U	
Hs.494854	NM_021218	C9orf80	9			up				Chromosome 9 open reading frame 80	
Hs.234282	NM_021729	VPS11	11				down			Vacuolar protein sorting 11 (yeast)	
Hs.301062	NM_021808	GALNT9	12	up	up					UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetyl-galactosaminyltransferase 9 (GalNAc-T9)	
Hs.269764	NM_021813	BACH2	6		up					BTB and CNC homology 1, basic leucine zipper transcription factor 2	

FIG. 1 (Sheet 32 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.257341	NM_021818	SAV1	14				down			Salvador homolog 1 (Drosophila)
Hs.461954	NM_021947	SRR	17	down	down				down	Serine racemase
Hs.213050	NM_021952	ELAVL4	1		up					ELAV (embryonic lethal, abnormal vision, Drosophila)-like 4 (Hu antigen D)
Hs.329978	NM_022087	GALNT11	7				up			UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetyl-galactosaminyltransferase 11 (GalNAc-T11)
Hs.128866	NM_022089	HSA9947	1				up			Putative ATPase
Hs.247324	NM_022100	MRPS14	1			up				Mitochondrial ribosomal protein S14
Hs.198158	NM_022129	MAWBP	10	down						MAWD binding protein
Hs.488591	NM_022479	WBSCR17	7		up					Williams-Beuren syndrome chromosome region 17
Hs.480356	NM_022553	VPS52	6	down		up				Vacuolar protein sorting 52 (yeast)
Hs.510265	NM_022731	NUCKS	1	down						Nuclear ubiquitously casein kinase and cyclin-dependent kinase substrate
Hs.440880	NM_022753	FLJ12903	1		down					Hypothetical protein FLJ12903
Hs.369440	NM_022754	SFXN1	5				up		down	Sideroflexin 1
Hs.284630	NM_022771	TBC1D15	12				down			TBC1 domain family, member 15
Hs.27836	NM_022823	FNDC4	2	down		down				Fibronectin type III domain containing 4
Hs.370549	NM_022893	BCL11A	2					down		B-cell CLL/lymphoma 11A (zinc finger protein)
Hs.203559	NM_022915	MRPL44	2						up	Mitochondrial ribosomal protein L44
Hs.239154	NM_023039	ANKRA2	5;3				down			Ankyrin repeat, family A (RFXANK-like), 2
Hs.146679	NM_023071	SPATS2	12				up			Spermatogenesis associated, serine-rich 2
Hs.169615	NM_023080	FLJ20989	8	up						Hypothetical protein FLJ20989
Hs.225641	NM_023923	PHACTR4	1						down	Phosphatase and actin regulator 4
Hs.169054	NM_023928	AACS	12				up	up		Acetoacetyl-CoA synthetase
Hs.157160	NM_023936	MRPS34	16	up						Mitochondrial ribosomal protein S34
Hs.458367	NM_024026	MRP63	13;8	up		up			up	Mitochondrial ribosomal protein 63
Hs.15580	NM_024075	LENG5	19			up				Leukocyte receptor cluster (LRC) member 5
Hs.59804	NM_024077	SECISBP2	9;17			up				SECIS binding protein 2
Hs.412939	NM_024090	ELOVL6	4					up		ELOVL family member 6, elongation of long chain fatty acids (FEN1/Elo2, SUR4/Elo3-like, yeast)
Hs.441734	NM_024091	MGC5297	5	down						Hypothetical protein MGC5297
Hs.177688	NM_024303	ZSCAN5	19	down						Zinc finger and SCAN domain containing 5
Hs.28144	NM_024333	FSD1	19				up			Fibronectin type 3 and SPRY domain containing 1
Hs.499205	NM_024336	IRX3	16		down					Iroquois homeobox protein 3
Hs.118394	NM_024345	MGC10765	9		down					Hypothetical protein MGC10765

FIG. 1 (Sheet 33 of 45)

UniGene ID	Annotation		Direction of Change in SZ relative to Controls									
	Acc	Symbol	Chr.	AnCg	CB	DLPPC	Nacc	PC	STG	Name		
Hs.211914	NM_024407	NDUFS7	19	up						NADH dehydrogenase (ubiquinone) Fe-S protein 7, 20kDa (NADH-coenzyme Q reductase)		
Hs.485004	NM_024493	ZNF306	6	down			down			Zinc finger protein 306		
Hs.250693	NM_024498	ZNF117	7;19	down						Kruppel-related zinc finger protein		
Hs.460568	NM_024516	MGC4606	16		down					Hypothetical protein MGC4606		
Hs.211441	NM_024517	PHF2	9					down		PHD finger protein 2		
Hs.443789	NM_024581	C6orf60	6						down	Chromosome 6 open reading frame 60		
Hs.470679	NM_024583	SCRN3	2		down					Secernin 3		
Hs.39311	NM_024592	SRD5A2L	4			up				Steroid 5 alpha-reductase 2-like		
Hs.293563	NM_024595	FLJ12666	1		down					Hypothetical protein FLJ12666		
Hs.200943	NM_024611	NARG2	15	up						NMDA receptor regulated 2		
Hs.317340	NM_024683	FLJ22729	17	up						Hypothetical protein FLJ22729		
Hs.445826	NM_024686	FLJ23033	1	up						Hypothetical protein FLJ23033		
Hs.121915	NM_024742	ARMCS	16			down				Armadillo repeat containing 5		
Hs.468349	NM_024766	FLJ23451	2	down						Hypothetical protein FLJ23451		
Hs.443061	NM_024810	CXorf45	X					down		Chromosome X open reading frame 45		
Hs.533446	NM_024812	BAALC	8	up						Brain and acute leukemia, cytoplasmic		
Hs.187377	NM_024847	TMC7	16		up					Transmembrane channel-like 7		
Hs.193170	NM_024859	FLJ21687	X		up					PDZ domain containing, X chromosome		
Hs.478465	NM_024871	FLJ12748	3				down			Hypothetical protein FLJ12748		
Hs.456507	NM_024874	PKD1-like	1	up						Polycystic kidney disease 1-like		
Hs.130712	NM_024876	ADCK4	19		up					AarF domain containing kinase 4		
Hs.374147	NM_024943	FLJ23235	4			down				Hypothetical protein FLJ23235		
Hs.371096	NM_025021	MECT1	19	up						Mucoepidermoid carcinoma translocated 1		
Hs.260555	NM_025057	C14orf45	14		up					Chromosome 14 open reading frame 45		
Hs.24808	NM_025073	FLJ21168	1		up					Hypothetical protein FLJ21168		
Hs.302051	NM_025109	MYOHD1							down	Myosin head domain containing 1		
Hs.288945	NM_025147	FLJ13448	2			down				Hypothetical protein FLJ13448		
Hs.288981	NM_025152	C14orf127	14			up				Chromosome 14 open reading frame 127		
Hs.301526	NM_025188	TRIM45	1			up				Tripartite motif-containing 45		
Hs.443723	NM_025196	GRPEL1	4							GrpE-like 1, mitochondrial (E. coli)		
Hs.434075	NM_025205	MED28	4;1		down				up	Mediator of RNA polymerase II transcription, subunit 28 homolog (yeast)		
Hs.329327	NM_025235	TNKS2	10				down			Tankyrase, TRF1-interacting ankyrin-related ADP-ribose polymerase 2		

FIG. 1 (Sheet 34 of 45)

UniGene ID	Annotation				Direction of Change in SZ relative to Controls						
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name	
Hs.118354	NM_025263	PRR3	6;4	up						Proline rich 3	
Hs.189445	NM_030583	MA1N2	8					down		Matrilin 2	
Hs.127126	NM_030627	CPEB4	5					down		Cytoplasmic polyadenylation element binding protein 4	
Hs.11067	NM_030630	C17orf28	17	down						Chromosome 17 open reading frame 28	
Hs.209561	NM_030650	KIAA1715	2					down		KIAA1715	
Hs.449628	NM_030759	NRBF2	10;8; 1	up						Nuclear receptor binding factor 2	
Hs.177841	NM_030762	BHLHB3	12				down			Basic helix-loop-helix domain containing, class B, 3	
Hs.480519	NM_030821	PLA2G12A	4					up		Phospholipase A2, group XIIA	
Hs.538547	NM_030920	ANP32E	1		down					Acidic (leucine-rich) nuclear phosphoprotein 32 family, member E	
Hs.436996	NM_030948	PHACTR1	6	down						Phosphatase and actin regulator 1	
Hs.267120	NM_030963	RNF146	6		down					Ring finger protein 146	
Hs.300816	NM_030981	RAB1B	11;9; 2	up			up			RAB1B, member RAS oncogene family	
Hs.301048	NM_031216	SEH1L	18			up				SEH1-like (<i>S. cerevisiae</i>)	
Hs.247280	NM_031227	C20orf18	20;18		up					Chromosome 20 open reading frame 18	
Hs.109051	NM_031286	SH3BGRL3	1	up						SH3 domain binding glutamic acid-rich protein like 3	
Hs.110695	NM_031287	SF3B5	6;20; 12	up					up	Splicing factor 3b, subunit 5, 10kDa	
Hs.270437	NM_031361	COL4A3BP	5				up			Collagen, type IV, alpha 3 (Goodpasture antigen) binding protein	
Hs.378808	NM_032025	eIF2A	3	down	down	down		down		Eukaryotic translation initiation factor (eIF) 2A	
Hs.465642	NM_032108	SEMA6B	19	up						Sema domain, transmembrane domain (TM), and cytoplasmic domain, (semaphorin) 6B	
Hs.501793	NM_032127	DKFZP566M1046	11			down				Hypothetical protein DKFZp566M1046	
Hs.100914	NM_032142	Cep192	18					down		Centrosomal protein 192 kDa	
Hs.519326	NM_032151	DCO1M	5		up					Dimerization cofactor of hepatocyte nuclear factor 1 (HNF1) from muscle	
Hs.399984	NM_032168	FLJ12519						down	down	Hypothetical protein FLJ12519	
Hs.441378	NM_032169	FLJ12592	3	up						Putative acyl-CoA dehydrogenase	
Hs.339612	NM_032177	PHAX	5			up			up	RNA U, small nuclear RNA export adaptor (phosphorylation regulated)	
Hs.381214	NM_032261	C21orf56	21						up	Chromosome 21 open reading frame 56	

FIG. 1 (Sheet 35 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.19673	NM_032272	MAF1		up						Homolog of yeast MAF1
Hs.124015	NM_032304	HAGHL	16						down	Hydroxyacylglutathione hydrolase-like
Hs.9088	NM_032305	MGC3200	1						up	Hypothetical protein LOC284615
Hs.513315	NM_032349	SDOS	16	up						Hypothetical protein MGC11275
Hs.239500	NM_032366	MGC13114	16	up						Hypothetical protein MGC13114
Hs.277154	NM_032380	GFM2	5						up	G elongation factor, mitochondrial 2
Hs.438709	NM_032446	MEGF10	5		up				down	MEGF10 protein
Hs.437126	NM_032496	ARHGAP9	12						up	Rho GTPase activating protein 9
Hs.144527	NM_032507	PGBD1	6	down						PiggyBac transposable element derived 1
Hs.486010	NM_032511	C6orf168	6			down				Chromosome 6 open reading frame 168
Hs.293753	NM_032515	BOK	2		up					BCL2-related ovarian killer
Hs.163642	NM_032536	NTNG2	9		down					Netrin G2
Hs.501106	NM_032550	KIAA1914	10	down						KIAA1914
Hs.387255	NM_032569	N-PAC	16			up				Cytokine-like nuclear factor n-pac
Hs.145010	NM_032576	CYorf15B	Y		up					Chromosome Y open reading frame 15B
Hs.132868	NM_032582	USP32	17	up						Ubiquitin specific protease 32
Hs.277101	NM_032609	COX4I2	20	up						Cytochrome c oxidase subunit IV isoform 2 (lung)
Hs.154140	NM_032623	OSAP	4		up					Ovary-specific acidic protein
Hs.129634	NM_032630	CINP	14;12			up				Cyclin-dependent kinase 2-interacting protein
Hs.459379	NM_032687	CYHR1	8	up						Cysteine and histidine rich 1
Hs.436035	NM_032704	TUBA6	12	down						Tubulin alpha 6
Hs.476972	NM_032778	MINA	3		up			up		MYC induced nuclear antigen
Hs.401537	NM_032802	SPLL2A	15						down	Putative intramembrane cleaving protease
Hs.435948	NM_032810	ATAD1	10						down	ATPase family, AAA domain containing 1
Hs.190983	NM_032813	FLJ14624	13						down	Hypothetical protein FLJ14624
Hs.461113	NM_032830	CIRH1A	16		down					Cirrhosis, autosomal recessive 1A (cirhin)
Hs.520287	NM_032870	C6orf111	6	down						Chromosome 6 open reading frame 111
Hs.388645	NM_032901	MGC14288	12	up						Hypothetical protein MGC14288
Hs.406788	NM_032932	RAB11FIP4	4	up						RAB11 family interacting protein 4 (class II)
Hs.505676	NM_033082	CIP29	12	up						Cytokine induced protein 29 kDa
Hs.292986	NM_033115	MGC16169	4			up				Hypothetical protein MGC16169
Hs.224843	NM_033210	ZNF502	3							Zinc finger protein 502
Hs.370530	NM_033220	TRIM14	9	down						Tripartite motif-containing 14
Hs.297452	NM_033260	FOXQ1	6						down	Forkhead box Q1
Hs.27695	NM_033291	MID1	X		up					Midline 1 (Opitz/BBB syndrome)

FIG. 1 (Sheet 36 of 45)

Annotation				Direction of Change in SZ relative to Controls					Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLFC	Nacc	PC		STG
Hs.335033	NM_033427	CTTNBP2	7		up					Cortactin binding protein 2
Hs.348390	NM_033439	C9orf26	9		down			down		Chromosome 9 open reading frame 26 (NF-HEV)
Hs.348262	NM_033495	KLHL13			up					Kelch-like 13 (Drosophila)
Hs.347270	NM_033554	HLA-DPA1	6		down		down			Major histocompatibility complex, class II, DP alpha 1
Hs.200600	NM_052837	SCAMP3	1	up						Secretory carrier membrane protein 3
Hs.440092	NM_052839	PANX2				down				Pannexin 2
Hs.320823	NM_052865	C20orf72	20					down		Chromosome 20 open reading frame 72
Hs.12082	NM_053000	TIGA1		up						TIGA1
Hs.408427	NM_053041	COMMD7	20	up						COMM domain containing 7
Hs.231029	NM_053045	MGC14327	9		down					Hypothetical protein MGC14327
Hs.203717	NM_054034	FN1	2						down	Fibronectin 1
Hs.410830	NM_058190	C21orf70	21	up						Chromosome 21 open reading frame 70
Hs.287518	NM_080415	PNUTL2	17				down			Peanut-like 2 (Drosophila)
Hs.55940	NM_080430	SELM		up						Selenoprotein M
Hs.179080	NM_080552	SLC32A1	20					up		Solute carrier family 32 (GABA vesicular transporter), member 1
Hs.156506	NM_080656	MGC13017	5		up					Similar to RIKEN cDNA A430101B06 gene
Hs.264208	NM_080667	MGC15407	2		down					Similar to RIKEN cDNA 4931428D14 gene
Hs.269577	NM_080841	VPS16	20		down	up				Protein tyrosine phosphatase, receptor type, A
Hs.135805	NM_080875	LOC142678	1	up						Skeletrophin
Hs.464422	NM_130386	COLEC12	18		down					Collectin sub-family member 12
Hs.304578	NM_130442	ELMO1	7			down				Engulfment and cell motility 1 (ced-12 homolog, C. elegans)
Hs.196482	NM_130469	JDP2	14			down				Jun dimerization protein 2
Hs.16258	NM_130781	RAB24	5	up						RAB24, member RAS oncogene family
Hs.483259	NM_130809	LOC133619	5						down	Hypothetical protein MGC12103
Hs.103315	NM_133476	ZNF384	12	down						Zinc finger protein 384
Hs.156316	NM_133503	DCN	12		down					Decorin
Hs.156316	NM_133504	DCN	12		down					Decorin
Hs.411488	NM_138290	RPIB9	7		up					Rap2-binding protein 9
Hs.264345	NM_138330	TIZ	19					up		TRAF6-inhibitory zinc finger protein
Hs.29645	NM_138364	LOC90826	4			up				Hypothetical protein BC004337
Hs.370055	NM_138409	C6orf117	6					up		Chromosome 6 open reading frame 117
Hs.460487	NM_138414	LOC112869	16						up	Hypothetical protein BC011981
Hs.348411	NM_138467	LOC127253	1			up				Hypothetical protein BC009514

FIG. 1 (Sheet 37 of 45)

Annotation		Direction of Change in SZ relative to Controls									
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLFC	Nacc	PC	STG	Name	
Hs.444338	NM_138698	LOC91431	4			down				Prematurely terminated mRNA decay factor-like	
Hs.129159	NM_138701	C7orf1	7	up						Chromosome 7 open reading frame 11	
Hs.484371	NM_139068	MAPK9	5	down		down				Mitogen-activated protein kinase 9	
Hs.356523	NM_139124	MAPK8IP2	22; 20;12		down					Mitogen-activated protein kinase 8 interacting protein 2	
Hs.21187	NM_139169	TRUB1	10			up				TruB pseudouridine (psi) synthase homolog 1 (E. coli)	
Hs.33470	NM_139278	LGI3	8					up		Leucine-rich repeat LGI family, member 3	
Hs.65256	NM_139284	LGI4	19		down					Leucine-rich repeat LGI family, member 4	
Hs.173034	NM_139316	AMPH	7				up			Amphiphysin (Stiff-Man syndrome with breast cancer 128kDa autoantigen)	
Hs.193163	NM_139351	BIN1	2;11		up					Bridging integrator 1	
Hs.371240	NM_144497	AKAP12	6						up	A kinase (PRKA) anchor protein (gravin) 12	
Hs.469264	NM_144563	RPIA	2		down					Ribose 5-phosphate isomerase A (ribose 5-phosphate epimerase)	
Hs.7962	NM_144584	FLJ30525	1			up				Hypothetical protein FLJ30525	
Hs.432780	NM_144638	MGC29956	3	up						Hypothetical protein MGC29956	
Hs.55150	NM_144647	MGC26610	5	down						Hypothetical protein MGC26610	
Hs.12381	NM_144669	FLJ31978	12					up		Hypothetical protein FLJ31978	
Hs.283828	NM_144770	RBM11	21	down						RNA binding motif protein 11	
Hs.502223	NM_144981	FLJ25059	11	up	up					Hypothetical protein FLJ25059	
Hs.533086	NM_144996	ARL2L1	3	up						ADP-ribosylation factor-like 2-like 1	
Hs.413359	NM_145030	MGC22793	7					down	down	Hypothetical protein MGC22793	
Hs.293818	NM_145043	NEIL2	8						up	Nei like 2 (E. coli)	
Hs.202207	NM_145047	NOR1	1			up				Oxidored-nitro domain-containing protein	
Hs.515490	NM_145056	MGC15476	19		down					Thymus expressed gene 3-like	
Hs.294145	NM_145265	LOC133957				up				Similar to RIKEN cDNA 0610011N22	
Hs.418495	NM_145267	C6orf57	6	down						Chromosome 6 open reading frame 57	
Hs.522992	NM_145306	C10orf35	10		up					Chromosome 10 open reading frame 35	
Hs.21938	NM_148907	OSBPL9	1					down		Oxysterol binding protein-like 9	
Hs.15783	NM_152302	C20orf158	20		down					Chromosome 20 open reading frame 158	
Hs.434914	NM_152330	C14orf31	14					down		Chromosome 14 open reading frame 31	
Hs.374556	NM_152339	MGC26885	16			down				Hypothetical protein MGC26885	
Hs.135181	NM_152361	FLJ38944	19		up					Hypothetical protein FLJ38944	
Hs.462033	NM_152371	MGC26818	1			down			down	Hypothetical protein MGC26818	
Hs.523413	NM_152372	MYOM3	1			up				Myomesin family, member 3	

FIG. 1 (Sheet 38 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.434945	NM_152379	DKFZp547B1713	1						down	Hypothetical protein DKFZp547B1713
Hs.524828	NM_152437	DKFZp761B128	12	up						Hypothetical protein DKFZp761B128
Hs.520192	NM_152608	FLJ35382	1					down		Hypothetical protein FLJ35382
Hs.400698	NM_152618	FLJ35630	4	up						Hypothetical protein FLJ35630
Hs.507584	NM_152705	MGC9850	13						up	Polymerase (RNA) I polypeptide D, 16kDa
Hs.487564	NM_152745	NXPH1	7	down						Neurexophilin 1
Hs.534591	NM_152766	MGC40107	17	up				up		Hypothetical protein MGC40107
Hs.135997	NM_152773	LOC116211	3						up	Hypothetical protein BC013113
Hs.200100	NM_152793	Ells1	7	up		up				Hypothetical protein Ells1
Hs.12102	NM_152827	SNX3	6;7	up						Sorting nexin 3
Hs.27788	NM_153034	ZNF488	10	up						Zinc finger protein 488
Hs.390567	NM_153048	FYN	6;20	up						FYN oncogene related to SRC, FGR, YES
Hs.504943	NM_153207	AEBP2	12					down		AE binding protein 2
Hs.460217	NM_153208	MGC35048	16		down					Hypothetical protein MGC35048
Hs.180257	NM_153231	ZNF550	19		down					Zinc finger protein 550
Hs.436743	NM_153233	FLJ36445	19		up					Hypothetical protein FLJ36445
Hs.128188	NM_153234	C5orf11	5		down				down	Chromosome 5 open reading frame 11
Hs.511991	NM_153240	NPHP3	3							Nephronophthisis 3 (adolescent)
Hs.399779	NM_153266	MGC33486		up						Hypothetical protein MGC33486
Hs.171001	NM_153456	HS6ST3	13			down				Heparan sulfate 6-O-sulfotransferase 3
Hs.156723	NM_153498	CAMK1D	10	down						Calcium/calmodulin-dependent protein kinase ID
Hs.484195	NM_153607	LOC153222	5			up				Adult retina protein
Hs.40910	NM_153634	CPNE8	12			up				Copine VIII
Hs.435080	NM_153690	FAM43A	3		up					Family with sequence similarity 43, member A
Hs.110477	NM_153741	DPM3	1	up						Dolichyl-phosphate mannosyltransferase polypeptide 3
Hs.511774	NM_153750	C21orf81	18	down					down	Chromosome 21 open reading frame 81
Hs.463985	NM_170741	KCNJ16	17						down	Potassium inwardly-rectifying channel, subfamily J, member 16
Hs.297343	NM_172226	CAMKK2	12				up			Calcium/calmodulin-dependent protein kinase 2, beta
Hs.131686	NM_172386	ABCA9	17	down	down					ATP-binding cassette, sub-family A (ABC1), member 9
Hs.132439	NM_173054	RBLN	7	down		down				Reelin
Hs.165258	NM_173173	NR4A2	2	down						Nuclear receptor subfamily 4, group A, member 2

FIG. 1 (Sheet 39 of 45)

Annotation			Direction of Change in SZ relative to Controls									
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name		
Hs.31422	NM_173468	MOBKLLA	4	up						MOB1, Mps One Binder kinase activator-like 1A (yeast)		
Hs.513424	NM_173501	LOC146174	16	up						Hypothetical protein LOC146174		
Hs.420244	NM_173642	MGC47816	1				up			Hypothetical protein MGC47816		
Hs.323482	NM_173680	MGC33584	7						up	Hypothetical protein MGC33584		
Hs.418198	NM_173797	PAPD4	5					down		PAP associated domain containing 4		
Hs.356697	NM_173827	FLJ38991	4	up						Hypothetical protein FLJ38991		
Hs.121663	NM_173848	LOC138046	8				up			Hypothetical protein LOC138046		
Hs.436405	NM_174855	IDH3B	20; 6:1		down					Isocitrate dehydrogenase 3 (NAD+) beta		
Hs.4295	NM_174871	PSMD12	17			down				Proteasome (prosome, macropain) 26S subunit, non-ATPase, 12		
Hs.355606	NM_174909	MGC23909		up						Hypothetical protein MGC23909		
Hs.534579	NM_174923	MGC31967	9			down				Hypothetical protein MGC31967		
Hs.321709	NM_175567	P2RX4	12;9						up	Purinergic receptor P2X, ligand-gated ion channel, 4		
Hs.406062	NM_175614	NDUFA11	19	up		up				NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 11, 14.7kDa		
Hs.251673	NM_175849	DNMT3B	20		up					DNA (cytosine-5-)-methyltransferase 3 beta		
Hs.448218	NM_175885	MGC33846	11		down					Hypothetical protein MGC33846		
Hs.203	NM_176875	CCKBR	11			down				Cholecystokinin B receptor		
Hs.287636	NM_177963	SYT12	11						down	Synaptotagmin XII		
Hs.425383	NM_178124	CXorf40	X;12							Chromosome X open reading frame 40		
Hs.471917	NM_178579	PSMF1	20			up				Proteasome (prosome, macropain) inhibitor subunit 1 (PI31)		
Hs.63236	NM_181465	MRPL55	1	up						Mitochondrial ribosomal protein L55		
Hs.464697	NM_181483	C18orf1	18		up					Chromosome 18 open reading frame 1		
Hs.11900	NM_181485	ZGPAT	20		down					Zinc finger, CCH-type with G patch domain		
Hs.433422	NM_181716	PRR6		up		up			up	Proline rich 6		
Hs.126137	NM_181866	BACH	1				up			Brain acyl-CoA hydrolase		
Hs.479853	NM_182472	EPHA5	4		up	up				EPH receptor A5		
Hs.116567	NM_182491	LOC90637	7	up						Hypothetical protein LOC90637		
Hs.437336	NM_182523	MGC61571	3			up				Hypothetical protein MGC61571		
Hs.200668	NM_182661	CERK	22						up	Ceramide kinase		
Hs.529735	NM_182662	AADAT	4	down						Amino adipate aminotransferase		
Hs.497579	NM_182665	RASSF5	1		up					Ras association (RalGDS/AF-6) domain family 5		
Hs.207157	NM_182703	LOC348094	15	down						Hypothetical protein LOC348094		

FIG. 1 (Sheet 40 of 45)

Annotation				Direction of Change in SZ relative to Controls						
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name
Hs.310537	NM_182734	PLCB1	20				up			Phospholipase C, beta 1 (phosphoinositide-specific)
Hs.472101	NM_182797	PLCB4	20			up				Phospholipase C, beta 4
Hs.380021	NM_182931	MLL5				up				Myeloid/lymphoid or mixed-lineage leukemia 5 (trithorax homolog, Drosophila)
Hs.414099	NM_183010	TNRC5	6;20			down		up		Trinucleotide repeat containing 5
Hs.446240	NM_183048	PRKCBP1	20		up		up			Protein kinase C binding protein 1
Hs.109087	NM_183075	CYP2U1	4	down						Cytochrome P450, family 2, subfamily U, polypeptide 1
Hs.298651	NM_183236	RAB27A	15	down						RAB27A, member RAS oncogene family
Hs.203634	NM_183239	GSTO2	10			up				Glutathione S-transferase omega 2
Hs.317095	NM_183243	IMPDH1	7;10				up		up	IMP (inosine monophosphate) dehydrogenase 1
Hs.191540	NM_184042	COH1	8	down						Cohen syndrome 1
Hs.291079	NM_194279	HBLD1	14	up						HESB like domain containing 1
Hs.268668	NM_194285	FLJ39441	11		down					Hypothetical protein FLJ39441
Hs.465337	NM_194449	PLEKHE1	18						down	Pleckstrin homology domain containing, family E (with leucine rich repeats) member 1
Hs.274479	NM_197972	NME7	1	up	up	up		up		Non-metastatic cells 7, protein expressed in (nucleoside-diphosphate kinase)
Hs.149500	NM_198038	NUDT9	4	up						Nudix (nucleoside diphosphate linked moiety X)-type motif 9
Hs.76662	NM_198045	ZDHHC16	10	up		up				Zinc finger, DHHC domain containing 16
Hs.465838	NM_198058	ZNF266	19		down		down			Zinc finger protein 266
Hs.20521	NM_198318	HRMT1L2	19				up			HMT1 hnRNP methyltransferase-like 2 (S. cerevisiae)
Hs.494155	NM_198394	C9orf85	9	up						Chromosome 9 open reading frame 85
Hs.446414	NM_198793	CD47	3				up			CD47 antigen (Rh-related antigen, integrin-associated signal transducer)
Hs.71941	NM_198867	MGC15677						down		Hypothetical protein MGC15677
Hs.515032	NM_199054	MKMK2	19						down	MAP kinase interacting serine/threonine kinase 2
Hs.449009	NM_199121	WARP	1				down			Von Willebrand factor A domain-related protein
Hs.528335	NM_199138	FLJ25477	13						down	Hypothetical protein FLJ25477
Hs.274959	NM_199167	CLUL1	18		up					Clusterin-like 1 (retinal)
Hs.517155	NM_199170	TMEPAI	20	up						Transmembrane, prostate androgen induced RNA
Hs.13645	NM_199297	THY28		up						Thymocyte protein thy28
Hs.13645	NM_199298	THY28	11;1	up	down					Thymocyte protein thy28
Hs.113919	NM_199342	LOC374969	1	up						Hypothetical protein LOC374969
Hs.188746	NM_199355	ADAMTS18	16		up					A disintegrin-like and metalloprotease (reprolysin type)

FIG. 1 (Sheet 41 of 45)

UniGene ID	Annotation		Direction of Change in SZ relative to Controls							Name
	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	
Hs.459362	NM_199414	PRC1	15		down					with thrombospondin type 1 motif, 18
Hs.351851	NM_199461	NANOS1	10						up	Protein regulator of cytokinesis 1
Hs.18128	NM_199513	C20orf44	20				down			Nanos homolog 1 (Drosophila)
Hs.16803	NM_201412	LUC7L	16;8;5	up						Chromosome 20 open reading frame 44
Hs.220950	NM_201559	FOXO3A	6	up	up	up				LUC7-like (S. cerevisiae)
Hs.259750	NM_201998	SF1	11	up						Forkhead box O3A
Hs.475848	NM_202758	HSRTSBETA	18				down			Splicing factor 1
Hs.434406	NM_203282	ZNF539	19	down						RTS beta protein
Hs.64004	NM_203406	LOC153364	5					down		Zinc finger protein 539
Hs.417029	NM_203415	DERP6	17;11						up	Similar to metallo-beta-lactamase superfamily protein
Hs.473838	NM_203433	DSCR2	21		down					S-phase 2 protein
Hs.508266	NM_203495	COMMMD6	13	up		up				Down syndrome critical region gene 2
Hs.20013	NM_207170	P29	1		down					COMM domain containing 6
Hs.369763	NM_207311	LOC92558	12			down				GCIIP-interacting protein p29
Hs.306423	NM_207357	LOC339524	1						down	Hypothetical protein LOC92558
Hs.203717	NM_212474	FN1	2						down	Hypothetical protein LOC339524
Hs.153752	NM_212530	CDC25B	20	up						Fibronectin 1
Hs.76122	R05810		1						down	Cell division cycle 25B
Hs.283401	U22030	CYP2A7	19				down			Transcribed locus, moderately similar to NP_055301.1 neuronal thread protein AD7c-NTP [Homo sapiens]
Hs.292156	U32331	DKK3	11		up					Cytochrome P450, family 2, subfamily A, polypeptide 7
Hs.174312	U69193	TLR4	9		down					Dickkopf homolog 3 (Xenopus laevis)
Hs.209226	U85992	BMPER	7		up					Toll-like receptor 4
Hs.465529	XM_028067	MIDN	19		down	down				BMP-binding endothelial regulator precursor protein
Hs.534513	XM_028217		4		up				up	Midnolin
Hs.311363	XM_034594			down						Hypothetical LOC90024
Hs.18564	XM_035299	ZSWIM6	5						down	Data not found
Hs.292575	XM_035863			down						Zinc finger, SWIM domain containing 6
Hs.476164	XM_038288	ZCCHC11	1							Data not found
Hs.314436	XM_038999					up				Zinc finger, CCHC domain containing 11
Hs.471504	XM_041126	KIAA1486	2		up			down		Data not found
Hs.225974	XM_043493	SV2C	5	down						KIAA1486 protein
Hs.184736	XM_043653	BEXL1	X		down					Synaptic vesicle glycoprotein 2C
Hs.387336	XM_045423				up					Brain expressed X-linked-like 1

FIG. 1 (Sheet 42 of 45)

Annotation		Direction of Change in SZ relative to Controls									
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Name	
Hs.472285	XM_046600	KIAA1272	20					down		KIAA1272 protein	
Hs.298382	XM_051197	KIAA1005	16			up				KIAA1005 protein	
Hs.534526	XM_057296	LOC116064	3	down						Hypothetical protein LOC116064	
Hs.187636	XM_058513	LRRK2	12					down		Leucine-rich repeat kinase 2	
Hs.220594	XM_059492		3					down		Hypothetical LOC131076	
Hs.380923	XM_085929	MEIS3	19	up						Meis1, myeloid ecotropic viral integration site 1 homolog 3 (mouse)	
Hs.335413	XM_086879		22						up	Hypothetical LOC150371	
Hs.169863	XM_087089	WDR43	2				up			WD repeat domain 43	
Hs.180663	XM_166451					up				Data not found	
Hs.423725	XM_168302						down			Data not found	
Hs.196647	XM_171054	KIAA0527	3					down		KIAA0527 protein	
Hs.162902	XM_173173	AOF1	6			up				Amine oxidase (flavin containing) domain 1	
Hs.441783	XM_290629	C14orf78	14;12			down				Chromosome 14 open reading frame 78	
Hs.132497	XM_290941	PRNP	1	up						Prion protein interacting protein	
Hs.165762	XM_291142	FCHO2	5					down		FCH domain only 2	
Hs.29068	XM_291277	DKFZp761P0423	8					down		Hypothetical protein DKFZp761P0423	
Hs.124128	XM_291326	KIAA2022			up					KIAA2022 protein	
Hs.380081	XM_370575	FBXL15	10;12	up						F-box and leucine-rich repeat protein 15	
Hs.9587	XM_370878	KIAA2002	15		down					KIAA2002 protein	
Hs.410889	XM_371074	DKFZP564D166	17				up			Putative ankyrin-repeat containing protein	
Hs.445218	XM_371116	MYO5B	18					up		Myosin VB	
Hs.288164	XM_371614	FLJ10707	3	up						Hypothetical protein FLJ10707	
Hs.136235	XM_371760	LOC116068	5		down					Hypothetical protein LOC116068	
Hs.458358	XM_371844	TSPYL1	6					down		TSPY-like 1	
Hs.136398	XM_372124	ZCCHC6	9	up						Zinc finger, CCHC domain containing 6	
Hs.149940	XM_372128		9		up					Similar to Osteotesticular phosphatase; protein tyrosine phosphatase receptor type V; protein tyrosine phosphatase receptor type W; protein tyrosine phosphatase, receptor type, V	
Hs.532698	XM_373630	LOC145842	15		down					Hypothetical protein LOC145842	
Hs.97540	XM_374002		2		up					Hypothetical gene supported by BC032913; BC048425	
Hs.389638	XM_374317		7		down					Hypothetical gene supported by AL713796	

FIG. 1 (Sheet 43 of 45)

Annotation			Direction of Change in SZ relative to Controls							
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLFC	Nacc	PC	STG	Name
Hs.447579	XM_375527	LOC339290	18					up		Hypothetical protein LOC339290
Hs.408708	XM_375553			down				down		Data not found
Hs.35524	XM_375604					down			up	Data not found
Hs.172884	XM_375633	SLC8A2	19				up			Solute carrier family 8 (sodium-calcium exchanger), member 2
Hs.416553	XM_375714		1		up					Similar to RIKEN cDNA 1700025K23
Hs.474836	XM_376010	LOC387593	22			down				TPTE/TPIP pseudogene
Hs.502948	XM_376043			up						Similar to RIKEN cDNA 2310016E02
Hs.301283	XM_376193				up					Data not found
Hs.113912	XM_376350	RAPGEF2	4	down					down	Rap guanine nucleotide exchange factor (GEF) 2
Hs.148956	XM_376436	LOC134466	5			down				Hypothetical protein LOC134466
Hs.520638	XM_376567	KIAA1856	7					down		KIAA1856 protein
Hs.308710	XM_376680	KIAA1718	7	up						KIAA1718 protein
Hs.23133	XM_378178	MGC9913	19	up	up				up	Hypothetical protein MGC9913
Hs.503862	XM_378309		11						up	Hypothetical LOC399951
Hs.434271	XM_378706		17					up		Hypothetical LOC400621
Hs.477007	XM_379203	LOC348801	3		up					Hypothetical protein LOC348801
Hs.446474	XM_379250				up					Hypothetical gene supported by BC038466; BC062790
Hs.432656	XM_379438	LOC285740	6	up						Hypothetical protein LOC285740
Hs.489988	XM_379923							down		Data not found
Hs.494204	XM_495929	DKFZp434N2030	12			up				Hypothetical protein DKFZp434N2030
Hs.17250	XM_495935	MGC4767	12	up	up				up	Hypothetical protein MGC4767
Hs.435761	XM_496395	FLJ34433	1		down					Hypothetical protein FLJ34433
Hs.143840	XM_496525						down			Data not found
Hs.99488	XM_496588	LOC130355	2	up						Hypothetical protein LOC130355
Hs.428360	XM_496681	KIAA1982	4	up						KIAA1982 protein
Hs.148988	XM_496999	KIAA1688	8			down				KIAA1688 protein
Hs.112622	XM_497679		1			up				Similar to Laminin receptor 1
Hs.528187	XM_498917		2					down		Hypothetical gene supported by AK096649
Hs.329512	XM_498955		3	up	up					Hypothetical gene supported by BC034933; BC068085
Hs.171132	XM_499008		5			down				Hypothetical gene supported by AK124699
Hs.31917	XM_499048	C6orf176	6		down					Chromosome 6 open reading frame 176
Hs.308222	XM_499514	LOC401321	7	down						Hypothetical LOC401321
Hs.21925	Z39995		8	down		down				Transcribed locus

FIG. 1 (Sheet 44 of 45)

Annotation		Direction of Change in SZ relative to Controls							Name	
UniGene ID	Acc	Symbol	Chr.	AnCg	CB	DLPFC	Nacc	PC	STG	Transcribed locus
Hs.22697	Z40025		10	down						

FIG. 1 (Sheet 45 of 45)

GO: 0050874 organismal physiological process

LocusLink ID	UniGene ID	Symbol
10175	Hs.294603	CNIH
10560	Hs.30246	SLC19A2
11255	Hs.251399	HRH3
2009	Hs.12451	EML1
28639	Hs.449451	TRBC1
3437	Hs.47338	IFIT3
4690	Hs.477693	NCK1
4929	Hs.165258	NR4A2
501	Hs.483239	ALDH7A1
641	Hs.169348	BLM
9308	Hs.484703	CD83

GO: 0058550 eukaryotic translation initiation factor 2 complex

LocusLink ID	UniGene ID	Symbol
26523	Hs.22867	EIF2C1
83939	Hs.378808	eIF2A
8894	Hs.429180	EIF2S2
GO: 0005739	mitochondrion	
LocusLink ID	UniGene ID	Symbol

10105	Hs.381072	PPIF
10730	Hs.499145	YME1L1
126328	Hs.406062	NDUFA11
1891	Hs.196176	ECH1
26292	Hs.370040	MYCBP
27429	Hs.115721	PRSS25
285521	Hs.356697	FLJ38991
374291	Hs.211914	NDUFS7
38	Hs.232375	ACAT1
4129	Hs.46732	MAOB
4726	Hs.408257	NDUFS6
51103	Hs.106529	NDUFAF1
51373	Hs.44298	MRPS17
51629	Hs.514216	CGI-69
518	Hs.429	ATP5G3
5188	Hs.119316	PET112L
53343	Hs.149500	NUDT9
56997	Hs.118241	CABC1
617	Hs.471401	BCS1L
6390	Hs.465924	SDHB
65993	Hs.157160	MRPS34
7284	Hs.12084	TUFM
84134	Hs.321653	FLJ12770
84693	Hs.94949	MCEE
84701	Hs.277101	COX4I2

FIG. 2

GO: 0005622 intracellular

LocusLink ID	UniGene ID	Symbol	Description
50	Hs.474982	ACO2	Aconitase 2, mitochondrial
518	Hs.429	ATP5G3	ATP synthase, H ⁺ transporting, mitochondrial F0 complex, subunit c (subunit 9) isoform 3
9331	Hs.464848	B4GALT6	UDP-Gal:betaGlcNAc beta 1,4-galactosyltransferase, polypeptide 6
617	Hs.471401	BCS1L	BCS1-like (yeast)
665	Hs.131226	BNIP3L	BCL2/adenovirus E1B 19kDa interacting protein 3-like
57128	Hs.387755	C6orf149	Chromosome 6 open reading frame 149
8697	Hs.153546	CDC23	CDC23 (cell division cycle 23, yeast, homolog)
8941	Hs.158460	CDK5R2	Cyclin-dependent kinase 5, regulatory subunit 2 (p39)
1108	Hs.162233	CHD4	Chromodomain helicase DNA binding protein 4
51340	Hs.171342	CRNKL1	Crn, crooked neck-like 1 (Drosophila)
1500	Hs.166011	CTNND1	Catenin (cadherin-associated protein), delta 1
1528	Hs.465413	CYB5	Cytochrome b-5
1602	Hs.129452	DACH1	Dachshund homolog 1 (Drosophila)
1656	Hs.408461	DDX6	DEAD (Asp-Glu-Ala-Asp) box polypeptide 6
9162	Hs.242947	DGKI	Diacylglycerol kinase, iota
1783	Hs.369068	DNCL12	Dynein, cytoplasmic, light intermediate polypeptide 2
54550	Hs.140950	EFCBP2	EF hand calcium binding protein 2
83939	Hs.378808	eIF2A	Eukaryotic translation initiation factor (eIF) 2A
317649	Hs.476782	EIF4E3	Eukaryotic translation initiation factor 4E member 3
9844	Hs.304578	ELMO1	Engulfment and cell motility 1 (ced-12 homolog, C. elegans)
30001	Hs.525339	ERO1L	ERO1-like (S. cerevisiae)
2135	Hs.357637	EXTL2	Exostoses (multiple)-like 2
23014	Hs.159699	FBXO21	F-box protein 21
55634	Hs.444269	FLJ20344	Hypothetical protein FLJ20344
29997	Hs.421907	GLTSCR2	Glioma tumor suppressor candidate region gene 2
56850	Hs.109929	GRIPAP1	GRIP1 associated protein 1
119391	Hs.203634	GSTO2	Glutathione S-transferase omega 2
57801	Hs.154029	HES4	Hairy and enhancer of split 4 (Drosophila)
3148	Hs.434953	HMGB2	High-mobility group box 2
3423	Hs.303154	IDS	Iduronate 2-sulfatase (Hunter syndrome)
122953	Hs.196482	JDP2	Jun dimerization protein 2
10656	Hs.444558	KHDRBS3	KH domain containing, RNA binding, signal transduction associated 3
57677	Hs.35524	KIAA1559	Mouse zinc finger protein 14-like

FIG. 3 (Sheet 1 of 3)

GO: 0005622	intracellular		
LocusLink ID	UniGene ID	Symbol	Description
3799	Hs.327736	KIF5B	Kinesin family member 5B
8609	Hs.471221	KLF7	Kruppel-like factor 7 (ubiquitous)
55915	Hs.224282	LANCL2	LanC lantibiotic synthetase component C-like 2 (bacterial)
51176	Hs.125132	LEF1	Lymphoid enhancer-binding factor 1
5641	Hs.18069	LGMN	Legumain
3998	Hs.465295	LMAN1	Lectin, mannose-binding, 1
4005	Hs.34560	LMO2	LIM domain only 2 (rhombotin-like 1)
6218	Hs.294145	LOC133957	Similar to RIKEN cDNA 0610011N22
93349	Hs.471582	LOC93349	Hypothetical protein BC004921
27258	Hs.111632	LSM3	LSM3 homolog, U6 small nuclear RNA associated (S. cerevisiae)
4129	Hs.46732	MAOB	Monoamine oxidase B
51257	Hs.445113	MARCH-II	Membrane-associated RING-CH protein II
10445	Hs.25313	MCRS1	Microspherule protein 1
1072	Hs.25313	MCRS1	Microspherule protein 1
4205	Hs.268675	MEF2A	MADS box transcription enhancer factor 2, polypeptide A (myocyte enhancer factor 2A)
28998	Hs.333823	MRPL13	Mitochondrial ribosomal protein L13
9801	Hs.44024	MRPL19	Mitochondrial ribosomal protein L19
63931	Hs.247324	MRPS14	Mitochondrial ribosomal protein S14
4664	Hs.107474	NAB1	NGFI-A binding protein 1 (EGR1 binding protein 1)
126328	Hs.406062	NDUFA11	NADH dehydrogenase (ubiquinone) 1 alpha subcomplex, 11, 14.7kDa
57520	Hs.314436	NEDL2	NEDD4-related E3 ubiquitin ligase NEDL2
84656	Hs.387255	N-PAC	Cytokine-like nuclear factor n-pac
4925	Hs.128686	NUCB2	Nucleobindin 2
51449	Hs.278627	PCYOX1	Prenylcysteine oxidase 1
9468	Hs.132794	PCYT1B	Phosphate cytidyltransferase 1, choline, beta isoform
5281	Hs.468415	PIGF	Phosphatidylinositol glycan, class F
5311	Hs.181272	PKD2	Polycystic kidney disease 2 (autosomal dominant)
11128	Hs.436896	POLR3A	Polymerase (RNA) III (DNA directed) polypeptide A, 155kDa
5464	Hs.437403	Previously presented	Pyrophosphatase (inorganic)
25865	Hs.466987	PRKD2	Protein kinase D2
8559	Hs.161181	PRPF18	PRP18 pre-mRNA processing factor 18 homolog (yeast)

FIG. 3 (Sheet 2 of 3)

GO: 0005622	intracellular		
LocusLink ID	UniGene ID	Symbol	Description
201161	Hs.433422	PRR6	Proline rich 6
5718	Hs.4295	PSMD12	Proteasome (prosome, macropain) 26S subunit, non-ATPase, 12
9491	Hs.471917	PSMF1	Proteasome (prosome, macropain) inhibitor subunit 1 (PI31)
10567	Hs.11417	RABAC1	Rab acceptor 1 (prenylated)
5994	Hs.24422	RFXAP	Regulatory factor X-associated protein
51320	Hs.465144	RKHD2	Ring finger and KH domain containing 2
57484	Hs.480825	RNF150	Ring finger protein 150
6096	Hs.494178	RORB	RAR-related orphan receptor B
6165	Hs.529631	RPL35A	Ribosomal protein L35a
25939	Hs.472630	SAMHD1	SAM domain and HD domain 1
79048	Hs.59804	SECISBP2	SECIS binding protein 2
23013	Hs.270499	SHARP	SMART/HDAC1 associated repressor protein
6732	Hs.443861	SRPK1	SFRS protein kinase 1
10618	Hs.14894	TGOLN2	Trans-golgi network protein 2
30000	Hs.416049	TNPO2	Transportin 2 (importin 3, karyopherin beta 2b)
80263	Hs.301526	TRIM45	Tripartite motif-containing 45
55109	Hs.213393	VG5Q	Angiogenic factor VG5Q
51564	Hs.269577	VPS16	Protein tyrosine phosphatase, receptor type, A
64601	Hs.269577	VPS16	Protein tyrosine phosphatase, receptor type, A
8976	Hs.143728	WASL	Wiskott-Aldrich syndrome-like
29799	Hs.517436	YPEL1	Yippee-like 1 (Drosophila)
7528	Hs.388927	YY1	YY1 transcription factor
146198	Hs.461074	ZFP90	Zinc finger protein 90 homolog (mouse)

FIG. 3 (Sheet 3 of 3)

UniGene ID	Acc	Gene Symbol	Chromosomal Location	Lymphocyte Direction (Schizophrenia/Control)	Fold change	p-value	Fold Change Agree Brain/Lymphocyte	Fold Change Agree Brain/Lymphocyte	Brain Direction (Schizophrenia/Control)	Gene Name
Hs.232375	NM_000019	ACAT1	11q22.3-q23.1	UP	1.15	0.024	yes	yes	UP	Acetyl-Coenzyme A acetyltransferase 1 (acetoacetyl Coenzyme A thiolase)
Hs.130712	NM_024876	ADCK4	19q13.2	DOWN	-1.11	0.042	No	No	UP	AarF domain containing kinase 4
Hs.207776	NM_000027	AGA	4q32-q33	DOWN	-1.11	0.038	yes	yes	DOWN	Aspartylglucosaminidase
Hs.368486	NM_001649	APXL	Xp22.3	DOWN	-1.14	0.024	No	No	UP	Apical protein-like (Xenopus laevis)
Hs.126137	NM_181866	BACH	1p36.31-p36.11	DOWN	-1.11	0.049	No	No	UP	Brain acyl-CoA hydrolase
Hs.293753	NM_032515	BOK	2q37.3	DOWN	-1.12	0.020	No	No	UP	BCL2-related ovarian killer
Hs.288981	NM_025152	C14orf127	14q12	DOWN	-1.16	0.019	No	No	UP	Chromosome 14 open reading frame 127
Hs.443789	NM_024581	C6orf60	6q22.31	UP	1.46	0.018	No	No	DOWN	Chromosome 6 open reading frame 60
Hs.532296	NM_017998	C9orf40	9q21.31	UP	1.13	0.036	yes	yes	UP	Chromosome 9 open reading frame 40

FIG. 4 (Sheet 1 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/ Control)	Fold change	p-value	Fold Change Agree Brain/ Lympho- cyte	Fold Change Agree Brain/ Lymphocyte	Brain Direction (Schizo- phrenia/ Control)	Gene Name
Hs.297343	NM_172226	CAMKK2	12q24.2	DOWN	-1.15	0.030	No	No	UP	Calcium/calmodulin -dependent protein kinase kinase 2, beta
Hs.297343	NM_172226	CAMKK2	12q24.2	DOWN	-1.13	0.018	No	No	UP	Calcium/calmodulin -dependent protein kinase kinase 2, beta
Hs.474797	NM_007061	CDC42EP1	22q13.1	DOWN	-1.10	0.008	No	No	UP	CDC42 effector protein (Rho GTPase binding) 1
Hs.249129	NM_001279	CIDEA	18p11.21	DOWN	-1.16	0.016	No	No	UP	Cell death-inducing DFFA-like effector a
Hs.29549	NM_016511	CLEC1	12p13.31	DOWN	-1.14	0.025	yes	yes	DOWN	C-type lectin-like receptor-1
Hs.270437	NM_031361	COL4A3BP	5q13.3	UP	1.27	0.047	yes	yes	UP	Collagen, type IV, alpha 3 (Goodpasture antigen) binding protein
Hs.464422	NM_130386	COLEC12	18pter-p11.3	DOWN	-1.15	0.044	yes	yes	DOWN	Collectin sub-family member 12
Hs.330384	NM_014325	CORO1C	12q24.1	DOWN	-1.17	0.040	No	No	UP	Coronin, actin binding protein, 1C

FIG. 4 (Sheet 2 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/Control)	Fold change	p-value	Fold Change Agree Brain/Lymphocyte	Fold Change Agree Brain/Lymphocyte	Brain Direction (Schizophrenia/Control)	Gene Name
Hs.45127	NM_006574	CSPG5	3p21.3	UP	1.16	0.023	No	No	DOWN	Chondroitin sulfate proteoglycan 5 (neuroglycan C)
Hs.304682	NM_000099	CST3	20p11.21	UP	1.74	0.026	No	No	DOWN	Cystatin C (amyloid angiopathy and cerebral hemorrhage)
Hs.26704	NM_014608	CYFIP1	15q11	DOWN	-1.42	0.010	yes	yes	DOWN	Cytoplasmic FMR1 interacting protein 1
Hs.283401	U22030	CYP2A7	19q13.2	DOWN	-1.13	0.006	yes	yes	DOWN	Cytochrome P450, family 2, subfamily A, polypeptide 7
Hs.12451	AF035276	EML1	14q32	DOWN	-1.11	0.047	No	No	UP	Echinoderm microtubule associated protein like 1
Hs.104925	NM_003633	ENC1	5q12-q13.3	UP	1.45	0.046	yes	yes	UP	Ectodermal-neural cortex (with BTB-like domain)
Hs.104925	NM_003633	ENC1	5q12-q13.3	UP	1.42	0.033	yes	yes	UP	Ectodermal-neural cortex (with BTB-like domain)
Hs.306251	NM_001982	ERBB3	12q13	DOWN	-1.45	0.015	yes	yes	DOWN	V-erb-b2 erythroblastic leukemia viral oncogene homolog 3 (avian)

FIG. 4 (Sheet 3 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/ Control)	Fold change	p-value	Fold Change Agree Brain/ Lympho- cyte	Fold Change Agree Brain/ Lymphocyte	Brain Direction (Schizo- phrenia/ Control)	Gene Name
Hs.438695	AK094876	FKBP11	12q13.12	UP	1.34	0.041	No	No	DOWN	FK506 binding protein 11, 19 kDa
Hs.443529	NM_019607	FLJ11267	8q13.1	DOWN	-1.13	0.040	yes	yes	DOWN	Hypothetical protein FLJ11267
Hs.147836	NM_017768	FLJ20331	1p31.2	UP	1.38	0.034	No	No	DOWN	Hypothetical protein FLJ20331
Hs.413137	NM_001680	FXYD2	11q23	DOWN	-1.19	0.023	No	No	UP	FXYD domain containing ion transport regulator 2
Hs.473648	NM_000819	GART	21q22.1	UP	1.32	0.029	yes	yes	UP	Phosphoribosylglyci namide formyltransferase, phosphoribosylglyci namide synthetase, phosphoribosylamin oimidazole synthetase
Hs.522418	NM_0010037 22	GLEIL	9q34.13	UP	1.28	0.045	yes	yes	UP	GLE1 RNA export mediator-like (yeast)
Hs.522418	NM_0010037 22	GLEIL	9q34.13	DOWN	-1.12	0.015	No	No	UP	GLE1 RNA export mediator-like (yeast)
Hs.430425	NM_002074	GNBI	1p36.33	DOWN	-1.20	0.034	No	No	UP	Guanine nucleotide binding protein (G protein), beta polypeptide 1

FIG. 4 (Sheet 4 of 10)

UniGene ID	Acc	Gene Symbol	Chromosoma Location	Lymphocyte Direction (Schizophrenia/ Control)	Fold change	p-value	Fold Change Agree Brain/ Lympho- cyte	Fold Change Agree Brain/ Lymphocyte	Brain Direction (Schizo- phrenia/ Control)	Gene Name
Hs.155090	AL117471	GNB5	15q21.1	DOWN	-1.13	0.030	No	No	UP	Guanine nucleotide binding protein (G protein), beta 5
Hs.155090	NM_006578	GNB5	15q21.1	DOWN	-1.10	0.005	No	No	UP	Guanine nucleotide binding protein (G protein), beta 5
Hs.198612	NM_005458	GPR51	9q22.1-q22.3	UP	1.17	0.044	yes	yes	UP	G protein-coupled receptor 51
Hs.445066	A1499801	GRIN2B	12p12	DOWN	-1.16	0.034	yes	yes	DOWN	Glutamate receptor, ionotropic, N-methyl D-aspartate 2B
Hs.12126	NM_018487	HCA112	7q36.1	DOWN	-1.34	0.007	yes	yes	DOWN	Hepatocellular carcinoma-associated antigen 112
Hs.278635	NM_016648	HDCMA18P	4q26	DOWN	-1.40	0.046	yes	yes	DOWN	HDCMA18P protein
Hs.35804	D25215	HERC3	4q21	DOWN	-1.13	0.012	No	No	UP	Hect domain and RLD 3
Hs.380250	AK094968	IFI16	1q22	UP	1.19	0.038	No	No	DOWN	Interferon, gamma-inducible protein 16
Hs.430551	NM_003870	IQGAP1	15q26.1	UP	1.24	0.021	No	No	DOWN	IQ motif containing GTPase activating protein 1
Hs.6396	NM_006694	JTB	1q21	UP	1.53	0.048	yes	yes	UP	Jumping translocation breakpoint
Hs.21703	NM_012281	KCND2	7q31	UP	1.97	0.038	yes	yes	UP	Potassium voltage-gated channel, Shal-related subfamily, member 2

FIG. 4 (Sheet 5 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/Control)	Fold change	p-value	Fold Change Agree Brain/Lymphocyte	Fold Change Agree Brain/Lymphocyte	Brain Direction (Schizophrenia/Control)	Gene Name
Hs.408960	NM_002241	KCNJ10	1q22-q23	DOWN	-1.14	0.005	yes	yes	DOWN	Potassium inwardly-rectifying channel, subfamily J, member 10
Hs.32505	NM_004981	KCNJ4	22q13.1	DOWN	-1.10	0.028	yes	yes	DOWN	Potassium inwardly-rectifying channel, subfamily J, member 4
Hs.2785	NM_000422	KRT17	17q12-q21	DOWN	-1.15	0.035	yes	yes	DOWN	Keratin 17
Hs.23748	NM_001290	LDB2	4p16	DOWN	-1.20	0.011	No	No	UP	LIM domain binding 2
Hs.352614	AF007155	LOC254531	15q13.2	UP	1.95	0.036	No	No	DOWN	PLSC domain containing protein
Hs.47649	NM_020166	MCCC1	3q27	DOWN	-1.16	0.022	yes	yes	DOWN	Methylcrotonoyl-Coenzyme A carboxylase 1 (alpha)
Hs.535659	BC002458	MCM3AP	21q22.3	DOWN	-1.16	0.046	No	No	UP	MCM3 minichromosome maintenance deficient 3 (S. cerevisiae) associated protein
Hs.460217	NM_153208	MGC35048	16p13.11	DOWN	-1.18	0.013	yes	yes	DOWN	Hypothetical protein MGC35048

FIG. 4 (Sheet 6 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/ Control)	Fold change	p-value	Fold Change Agree Brain/ Lympho- cyte	Fold Change Agree Brain/ Lymphocyte	Brain Direction (Schizo- phrenia/ Control)	Gene Name
Hs.21213	NM_000259	MYO5A	15q21	DOWN	-1.14	0.027	No	No	UP	Myosin VA (heavy polypeptide 12, myosin)
Hs.472185	NM_004552	NDUFS5	1p34.2-p33	UP	1.21	0.035	yes	yes	UP	NADH dehydrogenase (ubiquinone) Fe-S protein 5, 15kDa (NADH-coenzyme Q reductase)
Hs.211914	NM_024407	NDUFS7	19p13.3	DOWN	-1.11	0.039	No	No	UP	NADH dehydrogenase (ubiquinone) Fe-S protein 7, 20kDa (NADH-coenzyme Q reductase)
Hs.459255	AI935701	NTRK3	15q25	DOWN	-1.14	0.040	No	No	UP	Neurotrophic tyrosine kinase, receptor, type 3
Hs.459255	AI935701	NTRK3	15q25	DOWN	-1.12	0.021	No	No	UP	Neurotrophic tyrosine kinase, receptor, type 3
Hs.435714	NM_002576	PAK1	11q13-q14	UP	1.23	0.030	yes	yes	UP	P21/Cdc42/Rac1-activated kinase 1 (STE20 homolog, yeast)
Hs.503584	AI638679	PANX1	11q21	UP	1.27	0.036	yes	yes	UP	Pannexin 1
Hs.278627	NM_016297	PCYOX1	2p13.3	UP	1.14	0.033	yes	yes	UP	Prenylcysteine oxidase 1

FIG. 4 (Sheet 7 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/Control)	Fold change	p-value	Fold Change Agree Brain/Lymphocyte	Fold Change Agree Brain/Lymphocyte	Brain Direction (Schizophrenia/Control)	Gene Name
Hs.481819	AK021922	PDZK3	5p13.3	UP	1.24	0.009	No	No	DOWN	PDZ domain containing 3
Hs.468415	NM_002643	PIGF	2p21-p16	DOWN	-1.12	0.020	No	No	UP	Phosphatidylinositol glycan, class F
Hs.466848	NM_006905	PSG1	19q13.2	DOWN	-1.11	0.036	No	No	UP	Pregnancy specific beta-1-glycoprotein 1
Hs.413801	NM_014614	PSME4	2p16.3	DOWN	-3.66	0.003	No	No	UP	Proteasome (prosome, macropain) activator subunit 4
Hs.446429	NM_000954	PTGDS	9q34.2-q34.3	DOWN	-2.93	0.016	yes	yes	DOWN	Prostaglandin D2 synthase 21kDa (brain)
Hs.114062	NM_014241	PTPLA	10p14-p13	DOWN	-1.12	0.034	No	No	UP	Protein tyrosine phosphatase-like (proline instead of catalytic arginine), member a
Hs.434375	BI820698	PTPRB	12q15-q21	UP	1.32	0.019	No	No	DOWN	Protein tyrosine phosphatase, receptor type, B
Hs.296169	NM_004578	RAB4A	1q42-q43	UP	1.13	0.031	yes	yes	UP	RAB4A, member RAS oncogene family
Hs.411488	NM_138290	RPIB9	7q21.13	DOWN	-1.16	0.001	No	No	UP	Rap2-binding protein 9

FIG. 4 (Sheet 8 of 10)

UniGene ID	Acc	Gene Symbol	Chromosomal Location	Lymphocyte Direction (Schizophrenia/Control)	Fold change	p-value	Fold Change Agree Brain/Lymphocyte	Fold Change Agree Brain/Lymphocyte	Brain Direction (Schizophrenia/Control)	Gene Name
Hs.374588	NM_000985	RPL17	18q21	DOWN	-1.17	0.012	No	No	UP	Ribosomal protein L17
Hs.374588	NM_000985	RPL17	18q21	DOWN	-1.13	0.026	No	No	UP	Ribosomal protein L17
Hs.465924	NM_003000	SDHB	1p36.1-p35	DOWN	-1.12	0.038	No	No	UP	Succinate dehydrogenase complex, subunit B, iron sulfur (Ip)
Hs.146804	NM_006717	SPIN	9q22.1-q22.3	DOWN	-1.32	0.039	No	No	UP	Spindlin
Hs.237825	NM_006947	SRP72	4q11	DOWN	-1.13	0.044	No	No	UP	Signal recognition particle 72kDa
Hs.102735	NM_012446	SSBP2	5q14.1	UP	1.22	0.050	No	No	DOWN	Single-stranded DNA binding protein 2
Hs.12409	NM_001048	SST	3q28	DOWN	-1.15	0.045	yes	yes	DOWN	Somatostatin
Hs.482390	NM_003243	TGFBR3	1p33-p32	UP	1.18	0.017	No	No	DOWN	Transforming growth factor, beta receptor III (betaglycan, 300kDa)
Hs.14894	NM_006464	TGOLN2	2p11.2	UP	1.15	0.016	yes	yes	UP	Trans-golgi network protein 2
Hs.465784	AF026030	TIMM44	19p13.3-p13.2	DOWN	-1.15	0.033	No	No	UP	Translocase of inner mitochondrial membrane 44 homolog (yeast)
Hs.465784	NM_006351	TIMM44	19p13.3-p13.2	DOWN	-1.15	0.033	No	No	UP	Translocase of inner mitochondrial membrane 44 homolog (yeast)

FIG. 4 (Sheet 9 of 10)

UniGene ID	Acc	Gene Symbol	Chromosome Location	Lymphocyte Direction (Schizophrenia/ Control)	Fold change	p-value	Fold Change Agree Brain/ Lympho- cyte	Fold Change Agree Brain/ Lymphocyte	Brain Direction (Schizo- phrenia/ Control)	Gene Name
Hs.287362	AB046767	TLE3	15q22	DOWN	-1.19	0.009	No	No	UP	Transducin-like enhancer of split 3 (E(sp1) homolog, Drosophila)
Hs.287362	NM_005078	TLE3	15q22	DOWN	-1.19	0.009	No	No	UP	Transducin-like enhancer of split 3 (E(sp1) homolog, Drosophila)
Hs.499145	BC019602	YME1L1	10p14	UP	1.13	0.043	No	No	DOWN	YME1-like 1 (S. cerevisiae)
Hs.172979	NM_003451	ZNF177	19p13.2	DOWN	-1.17	0.021	No	No	UP	Zinc finger protein 177

FIG. 4 (Sheet 10 of 10)

Accession Number	Gene Symbol	Microarray Fold Change	Q-PCR Fold Change	Microarray (p-value t-test)	Q-PCR (p-value t-test)	Cytogenetic Band	Gene Name
NM_002110	HCK	2.58	4.71	0.030	0.02	20q11-q12	hemopoietic cell kinase
NM_021005	NR2F2	1.74	4.15	0.047	0.02	15q26.2	nuclear receptor subfamily 2, group F, member 2
NM_017423	GALNT7	1.34	2.81	0.041	0.46	4q34.1	UDP-N-acetyl-alpha-D-galactosamine:polypeptide N-acetylgalactosaminyltransferase 7 (GalNAc-T7)
NM_139045	SMARCA2	1.75	2.27	0.028	0.03	9p22.3	SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 2
NM_018639	WSB2	1.49	2.18	0.003	0.02	12q24.23	WD repeat and SOCS box containing protein 2
NM_203504	G3BP2	1.28	2.04	0.045	0.07	4q21.21	Ras-GTPase activating protein SH3 domain-binding protein 2
NM_023927	NS3TP2	1.27	1.75	0.030	0.08	5q23.3	HCV NS3-transactivated protein 2
NM_003816	ADAM9	1.45	1.70	0.024	0.04	8p11.22	a disintegrin and metalloproteinase domain 9 (meltrin gamma)
BQ000126	TMOD3	1.21	1.39	0.031	0.12	15q21.1-q21.2	tropomodulin 3 (ubiquitous)
NM_003100	SNX2	1.43	0.95	0.042	0.45	5q23	sorting nexin 2
AL041379	HERC3	0.89	0.90	0.023	0.34	4q21	hect domain and RLD 3
AA487462	FLJ20637	0.81	0.68	0.014	0.08	4q22.1	hect domain and RLD 6
BM855607	IGJ	0.45	0.63	0.038	0.21	4q21	immunoglobulin J polypeptide, linker protein for immunoglobulin alpha and mu polypeptides
NM_005214	CTLA4	0.82	0.14	0.036	0.04	2q33	cytotoxic T-lymphocyte-associated protein 4
AV700777	ADH1B	0.89	0.10	0.023	0.02	4q21-q23	alcohol dehydrogenase IB (class I), beta polypeptide

FIG. 5

db SNP RS ID	NCBI (Mb)	Cis < 5	Regression (p-value)
rs1318822	170460379	8.6	0.0015
rs1403225	171867785	7.2	0.0158
rs1812424	173851484	5.2	0.0158
rs2119788	175143279	3.9	0.0020
rs723820	177930796	1.1	0.0020
rs723819	177930836	1.1	0.0020
rs1112286	178134007	0.9	0.0037
rs1375749	178157255	0.9	0.0037
rs1902018	179382986	0.3	0.0016
rs722387	179808887	0.8	0.0400
rs1112857	181655677	2.6	0.0400

FIG. 6

Genes dysregulated in MD, BP, and schizophrenia

1	UniGene ID	probe set	Acc	Name	Symbol	Direction of Change
2	Hs.282878	Hs.282878_at	NM_021633	Kelch-like 12 (Drosophila)	KLHL12	DLPFC-BPD&MDD&SCZ-U
3	Hs.435039	Hs.435039_at	NM_014914	Trinucleotide repeat containing 17	CENTG2	DLPFC-BPD&MDD&SCZ-U
4	Hs.530712	Hs.530712_at	NM_017917	Chromosome 14 open reading frame 10	C14orf10	DLPFC-BPD&MDD&SCZ-U
5	Hs.56294	Hs.56294_at	NM_004794	RAB33A, member RAS oncogene family	RAB33A	DLPFC-BPD&MDD&SCZ-U
6	Hs.21577	Hs.21577_at	NM_005701	RNA, U transporter 1	RNUT1	DLPFC-BPD&SCZ-U
7	Hs.274479	Hs.274479_at	NM_197972	Non-metastatic cells 7, protein expressed in (nucleoside-diphosphate kinase)	NME7	DLPFC-BPD&SCZ-U
8	Hs.368486	Hs.368486_at	NM_001649	Apical protein-like (Xenopus laevis)	APXL	DLPFC-BPD&SCZ-U
9	Hs.468415	Hs.468415_at	NM_002643	Phosphatidylinositol glycan, class F	PIGF	DLPFC-BPD&SCZ-U
10	Hs.471401	Hs.471401_at	NM_004328	BCS1-like (yeast)	BCS1L	DLPFC-BPD&SCZ-U
11	Hs.502145	Hs.502145_at	NM_006157	NEL-like 1 (chicken)	NELL1	DLPFC-BPD&SCZ-U
12	Hs.514036	Hs.514036_at	NM_006923	Stromal cell-derived factor 2	SDF2	DLPFC-BPD&SCZ-U
13	Hs.532853	Hs.532853+_at	NM_004146	NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7, 18kDa	NDUFB7	DLPFC-BPD&SCZ-U
14	Hs.111779	Hs.111779_at	NM_003118	Secreted protein, acidic, cysteine-rich (osteonectin)	SPARC	DLPFC-BPD&MDD&SCZ-D
15	Hs.171695	Hs.171695_at	NM_004417	Dual specificity phosphatase 1	DUSP1	DLPFC-BPD&MDD&SCZ-D

FIG. 7 (Sheet 1 of 2)

1 6	Hs.21283 8	Hs.212838- _at	NM_00001 4	Alpha-2- macroglobulin	A2M	DLPFC- BPD&MDD&SCZ -D
1 7	Hs.34560	Hs.34560- _at	NM_00557 4	LIM domain only 2 (rhombotin-like 1)	LMO2	DLPFC- BPD&MDD&SCZ -D
1 8	Hs.34727 0	Hs.347270- _at	NM_03355 4	Major histocompatibilit y complex, class II, DP alpha 1	HLA- DPA1	DLPFC- BPD&MDD&SCZ -D
1 9	Hs.43656 8	Hs.436568_a t	BC024272	CD74 antigen (invariant polypeptide of major histocompatibilit y complex, class II antigen- associated)	CD74	DLPFC- BPD&MDD&SCZ -D
2 0	Hs.49158 2	Hs.491582_a t	NM_00093 1	Plasminogen activator, tissue	PLAT	DLPFC- BPD&MDD&SCZ -D
2 1	Hs.53411 5	Hs.534115_a t	NM_00698 8	A disintegrin- like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1	ADAMTS 1	DLPFC- BPD&MDD&SCZ -D
2 2	Hs.17109	Hs.17109_at	NM_00486 7	Integral membrane protein 2A	ITM2A	DLPFC- BPD&SCZ-D
2 3	Hs.48513 0	Hs.485130_a t	K01615	Major histocompatibilit y complex, class II, DP beta 1	HLA- DPB1	DLPFC- BPD&SCZ-D
2 4	Hs.50487 7	Hs.504877_a t	X69549	Rho GDP dissociation inhibitor (GDI) beta	ARHGDI B	DLPFC- BPD&SCZ-D
2 5	Hs.52004 8	Hs.520048_a t	NM_01911 1	Major histocompatibilit y complex, class II, DR alpha	HLA-DRA	DLPFC- BPD&SCZ-D

FIG. 7 (Sheet 2 of 2)


Genes dysregulated in MD, BP, and schizophrenia

Symbol	Name	UniGene ID	AnCg	DLPFC	CB	nAcc
PTGDS	Prostaglandin	Hs.446429			CB-D	nAcc-D
PLAT	Plasminogen activator, tissue	Hs. 491582	AnCg-D	DLPFC-D		
ADAMTS1	Disintegrin-like and metalloprotease	Hs. 534115		DLPFC-D		nAcc-D

FIG. 8

Differentially Expressed Genes in Amygdala
 ~ Passed 4 Analysis Platforms ~
 O BPD O MDD O SZ

Amygdala
 (51 genes)



Dysregulated in Amygdala of BPD and SZ

Hs.460	Activating transcription factor 3	O	ATF3	BPD-D, SCZ-D
Hs.2128	Dual specificity phosphatase 5		DUSP5	BPD-D, SCZ-D
Hs.525704	V-jun sarcoma virus 17 oncogene homolog	O	JUN	BPD-D, SCZ-D
Hs.534052	Zinc finger protein 36, C3H type homolog		ZFP36	BPD-D, SCZ-D

FIG. 9

Differentially Expressed Genes in Amygdala
 ~ Passed 4 Analysis Platforms ~
 O BPD O MDD O SZ

Amygdala
(51 genes)

Dysregulated in Amygdala of BPD

UGCluster	Name	Symbol	Pass 4 Platforms
Hs.27283	GABA-A receptor, beta 1	GABRB1	BPD-I
Hs.24907	Coronin, actin binding protein, 2B	CORO2B	BPD-I




FIG. 10




FIG. 11

Differentially Expressed Genes in Amygdala
 ~ Passed 4 Analysis Platforms ~
 O BPD O MDD O SZ

Dysregulated in Amygdala of BPD, MDD and SCZ

UGCluster	Name	Symbol	Pass 4 Platforms
Hs.23748	LIM domain binding 2	LDB2O	BPD-I, MDD-I, SCZ-I
Hs.380089	EPH receptor B6	EPHB6	BPD-I, MDD-I, SCZ-I
Hs.376206	Kruppel-like factor 4 (gut)	KLF4O	BPD-D, MDD-D, SCZ-D

Amygdala
 (51 genes)




FIG. 12

**Genes Dysregulated in both SZ and BPD
DLPFC**

NAME	SYMBOL	NAME	SYMBOL
Kelch-like 12 (Drosophila)	KLHL12	LIM domain only 2 (rhombotin-like 1)	LMO2
Trinucleotide repeat containing 17	CENTG2	Major histocompatibility complex, class II, DP alpha 1	HLA-DPA1
Chromosome 14 open reading frame 10	C14orf10	CD74 antigen (invariant polypeptide of major histocompatibility complex, class II antigen-associated)	CD74
RAB33A, member RAS oncogene family	RAB33A	Plasminogen activator, tissue	PLAT
RNA, U transporter 1	RNUT1	A disintegrin-like and metalloprotease (reprolysin type) with thrombospondin type 1 motif, 1	ADAMTS1
Non-metastatic cells 7, protein expressed in (nucleoside-diphosphate kinase)	NME7	Integral membrane protein 2A	ITM2A
Apical protein-like (Xenopus laevis)	APXL	Major histocompatibility complex, class II, DP beta 1	HLA-DPB1
Phosphatidylinositol glycan, class F	PIGF	Rho GDP dissociation inhibitor (GDI) beta	ARHGDIB
BCS1-like (yeast)	BCS1L	Major histocompatibility complex, class II, DR alpha	HLA-DRA
NEL-like 1 (chicken)	NELL1	Major histocompatibility complex, class II, DR alpha	HLA-DRA
Stromal cell-derived factor 2	SDF2		
NADH dehydrogenase (ubiquinone) 1 beta subcomplex, 7, 18kDa	NDUFB7		
Secreted protein, acidic, cysteine-rich (osteonectin)	SPARC		
Dual specificity phosphatase 1	DUSP1		
Alpha-2-macroglobulin	A2M		

FIG. 13

LI_UP_DOWN_candidates_AND_BPD_26 JAN 06

???			
AMY			
NM_002781	19q13.2	PSG5	Pregnancy specific beta-1-glycoprotein 5
NM_003998	4q24		Nuclear factor of kappa light polypeptide gene enhance
NM_004898	4q12	CLOCK	Clock homolog (mouse)
NM_013263	16q12	BRD7	Bromodomain containing 7
AnCg			
NM_001048	3q28	SST	Somatostatin
NM_002093	3q13.3		Glycogen synthase kinase 3 beta
NM_003489	21q11.2	NRIP1	Nuclear receptor interacting protein 1
NM_003936	2q35	CDK5R2	Cyclin-dependent kinase 5, regulatory subunit 2 (p39)
NM_013444	Xp11.23-p11.1	UBQLN2	Ubiquilin 2
NM_017993	13q14.11	FLJ10094	Hypothetical protein FLJ10094
NM_020178	17q21	CA10	Carbonic anhydrase X
NM_021952	1p34	ELAVL4	ELAV (embryonic lethal, abnormal vision, Drosophila)-like 4 (Hu antigen D)

DLPFC

NM_025098	11q13.5	MOGAT2	Monoacylglycerol O-acyltransferase 2
-----------	---------	--------	--------------------------------------

HC

None

UP

AMY

NM_002065	1q31	GLUL	Glutamate-ammonia ligase (glutamine synthetase) (not in BPD)
NM_000014	12p13.3-p12.3	A2M	Alpha-2-macroglobulin
NM_001004	11p15.5-p15.4	RPLP2	Ribosomal protein, large, P2
NM_001283	7q22.1	AP1S1	Adaptor-related protein complex 1, sigma 1 subunit
NM_004355	5q32	CD74	CD74 antigen (invariant polypeptide of major histocompatibility complex, class II antigen-associated)
NM_004368	21q11.1	CNN2	Calponin 2
NM_005617	5q31-q33	RPS14	Ribosomal protein S14
NM_006119	10q24	???	Fibroblast growth factor 8 (androgen-induced)
NM_006870	20p12.1	DSTN	Destrin (actin depolymerizing factor)
NM_080391	1p35	PIP4A2	Protein tyrosine phosphatase type IVA, member 2
NM_153477	Xp11.23-p11.22	UXT	Ubiquitously-expressed transcript

AnCg

None

DLPFC2

NM_000014	12p13.3-p12.3	A2M	Alpha-2-macroglobulin
NM_001004	11p15.5-p15.4	RPLP2	Ribosomal protein, large, P2
NM_001013	19q13.4	RPS9	Ribosomal protein S9
NM_001283	7q22.1	AP1S1	Adaptor-related protein complex 1, sigma 1 subunit
NM_002178	12q13	IGFBP6	Insulin-like growth factor binding protein 6
NM_019597	Xq22	HNRPH2	Heterogeneous nuclear ribonucleoprotein H2 (H')
NM_018584	1p36.12	CaMKINalpha	Calcium/calmodulin-dependent protein kinase II inhibitor 1 (not in BPD)

HC

NM_053025	3q21	MYLK	Myosin, light polypeptide kinase
NM_006693	2q24	??	T-box, brain, 1 (not in BPD)

FIG. 14

Diagnosis	Total	PSPHL locus genotype			PSPHL mRNA expression	
		Del/Del	Del/Ins	Ins/Ins	No Expression	Expressed
BPD	19	18	1	0	18	1
MDD	22	8	13	1	8	14
SCZ	20	12	8	0	12	8
Control	64	43	18	3	43	21




FIG. 15

**COMPOSITIONS AND METHODS FOR
DIAGNOSING AND TREATING
NEUROPSYCHIATRIC DISORDERS**

**CROSS-REFERENCES TO RELATED
APPLICATIONS**

[0001] The present application is a continuation of U.S. application Ser. No. 11/396,050, which application claims priority to U.S. Ser. No. 60/667,299, filed Mar. 31, 2005, and U.S. Ser. No. 60/776,103, filed Feb. 22, 2006, which are herein incorporated by reference in their entirety.

**STATEMENT AS TO RIGHTS TO INVENTIONS
MADE UNDER FEDERALLY SPONSORED
RESEARCH AND DEVELOPMENT**

[0002] Not applicable.

BACKGROUND OF THE INVENTION

[0003] Psychotic disorders such as schizophrenia and mood disorders such as major depression and bipolar disorder are a major public health problem, affecting a significant portion of the adult population of the United States each year. While it has been hypothesized that mental disorders, including psychotic disorders such as schizophrenia as well as mood disorders such as major depression and bipolar disorder have genetic roots, little progress has been made in identifying gene sequences and gene products that play a role in causing these disorders, as is true for many diseases with a complex genetic origin (see, e.g., Burmeister, *Biol. Psychiatry* 45:522-532 (1999)). Relying on the discovery that certain genes expressed in particular brain pathways and regions are likely involved in the development of mental disorders, the present invention provides methods for diagnosis and treatment of mental disorders such as schizophrenia, as well as methods for identifying compounds effective in treating mental disorders.

BRIEF SUMMARY OF THE INVENTION

[0004] In order to further understand the neurobiology of psychotic disorders such as schizophrenia, the inventors of the present application have used DNA microarrays to study expression profiles of human post-mortem brains from patients diagnosed with schizophrenia. The work has focused on six brain regions that are pathways or circuits involved in schizophrenia: the anterior cingulate cortex (AnCg), dorso-lateral prefrontal cortex (DLPFC), cerebellar cortex (CB), superior temporal gyms (STG), parietal cortex (PC), and nucleus accumbens (nAcc).

[0005] The present invention demonstrates differential expression of genes in selected regions of brains of patients suffering from schizophrenia in comparison with normal control subjects. These genes include the transcripts listed in FIG. 1; the genes listed in FIG. 2 which are differentially expressed in the AnCg using Affymetrix chips and using brains with no agonal factors; the genes listed in FIG. 3 which are differentially expressed in the DLPFC using Affymetrix chips and using brains with no agonal factors; and the genes listed in FIG. 4 which are significantly dysregulated in both lymphoblastic and brain tissues.

[0006] In addition, the present invention identifies genes which are not differentially regulated in brain tissue but which are differentially regulated in lymphocytes of schizophrenic patients (FIG. 5). Also provided is a list (FIG. 6) of

single nucleotide polymorphic markers which are related to aspartylglucosaminuria (AGA), a gene which is dysregulated in both brain and lymphocytes of schizophrenic patients. FIGS. 7 and 8 show genes that are dysregulated in schizophrenia, major depression, and bipolar disorder.

[0007] The present invention also provides genes that are differentially expressed in the amygdala in patients diagnosed with major depression disorder, bipolar disorder, and/or schizophrenia (FIGS. 9-13).

[0008] The present invention also provides lithium responsive genes that are differentially expressed in the amygdala of lithium treated bipolar subjects and lithium treated non-human primates (FIG. 14).

[0009] The present invention also provides validation of a variant version of PSPHL with an insertion deletion mutation as a useful diagnostic tool to distinguish patients with bipolar disorder among patients presenting with depression, or for diagnosis of bipolar disorder.

[0010] Genes that are differentially expressed in neuropsychiatric disorders are useful in diagnosing psychotic and mood disorders, e.g., providing SNPs, biomarkers, diagnostic probe sets for PCR and chip assays, and antigens and antibodies for immunoassays such as ELISA and immunohistochemical assays. Differential expression by brain region similarly is a useful diagnostic and therapeutic tool, as psychotic and mood disorders primarily affect certain brain regions that are part of circuits or pathways involved in the disorder. Imaging brain endogenous gene expression with sequence-specific antisense radiopharmaceuticals and novel aptamer-based probes is a powerful diagnostic tool. Those probes can be detected using both fluorescent- and radiolabels that can be used in conjunction with single-photon emission computed tomography (SPECT) or positron emission tomography (PET) imaging modalities. The identification of genes, proteins, and biochemical assays involved in psychotic and mood disorders also provides the means for drug discovery for anti-psychotic therapeutics, such as small molecules, siRNA, and antibodies.

[0011] This invention thus provides methods for determining whether a subject has or is predisposed for a mental disorder. The invention also provides methods of providing a prognosis and for monitoring disease progression and treatment. Furthermore, the present invention provides nucleic acid and protein targets for assays for drugs for the treatment of mental disorders.

[0012] In one aspect, the methods comprise the steps of: (i) obtaining a biological sample from a subject; (ii) contacting the sample with a reagent that selectively associates with a polynucleotide or polypeptide encoded by a nucleic acid that hybridizes under stringent conditions to a nucleotide sequence listed in FIGS. 1-14; and (iii) detecting the level of reagent that selectively associates with the sample, thereby determining whether the subject has or is predisposed for a mental disorder.

[0013] In some embodiments, the reagent is an antibody. In some embodiments, the reagent is a nucleic acid. In some embodiments, the reagent associates with a polynucleotide. In some embodiments, the reagent associates with a polypeptide. In some embodiments, the polynucleotide comprises a nucleotide sequence listed in FIGS. 1-14. In some embodiments, the polypeptide comprises an amino acid sequence of a gene listed in FIGS. 1-14. In some embodiments, the level of reagent that associates with the sample is different (i.e., higher or lower) from a level associated with humans without

a mental disorder. In some embodiments, the biological sample is obtained from lymphocytes, amniotic fluid, spinal fluid, or saliva. In some embodiments, the mental disorder is a mood disorder. In some embodiments, the mental disorder is a psychotic disorder such as schizophrenia.

[0014] The invention also provides methods of identifying a compound for treatment of a mental disorder. In some embodiments, the methods comprises the steps of: (i) contacting the compound with a polypeptide, which is encoded by a polynucleotide that hybridizes under stringent conditions to a nucleic acid comprising a nucleotide sequence of FIGS. 1-14; and (ii) determining the functional effect of the compound upon the polypeptide, thereby identifying a compound for treatment of a mental disorder, e.g., schizophrenia.

[0015] In some embodiments, the contacting step is performed in vitro. In some embodiment, the polypeptide comprises an amino acid sequence of a gene listed in FIGS. 1-14. In some embodiments, the polypeptide is expressed in a cell or biological sample, and the cell or biological sample is contacted with the compound. In some embodiments, the methods further comprise administering the compound to an animal and determining the effect on the animal, e.g., an invertebrate, a vertebrate, or a mammal. In some embodiments, the determining step comprises testing the animal's mental function.

[0016] In some embodiments, the methods comprise the steps of (i) contacting the compound to a cell, the cell comprising a polynucleotide that hybridizes under stringent conditions to a nucleotide sequence of FIGS. 1-14; and (ii) selecting a compound that modulates expression of the polynucleotide, thereby identifying a compound for treatment of a mental disorder. In some embodiments, the polynucleotide comprises a nucleotide sequence listed in FIGS. 1-14. In some embodiment, the expression of the polynucleotide is enhanced. In some embodiments, the expression of the polynucleotide is decreased. In some embodiments, the methods further comprise administering the compound to an animal and determining the effect on the animal. In some embodiments, the determining step comprises testing the animal's mental function. In some embodiments, the mental disorder is a mood disorder or a psychotic disorder. In some embodiments, the psychotic disorder is schizophrenia. In some embodiments, the mood disorder is major depression disorder or bipolar disorder.

[0017] The invention also provides methods of treating a mental disorder in a subject. In some embodiments, the methods comprise the step of administering to the subject a therapeutically effective amount of a compound identified using the methods described above. In some embodiments, the mental disorder is a mood disorder or a psychotic disorder. In some embodiments, the compound is a small organic molecule, an antibody, an antisense molecule, an aptamer, an siRNA molecule, or a peptide.

[0018] The invention also provides methods of treating mental disorder in a subject, comprising the step of administering to the subject a therapeutically effective amount of a polypeptide, which is encoded by a polynucleotide that hybridizes under stringent conditions to a nucleic acid of FIGS. 1-14. In some embodiments, the polypeptide comprises an amino acid sequence encoded by a gene sequence listed in FIGS. 1-14. In some embodiments, the mental disorder is a mood disorder or a psychotic disorder.

[0019] The invention also provides methods of treating mental disorder in a subject, comprising the step of adminis-

tering to the subject a therapeutically effective amount of a polynucleotide, which hybridizes under stringent conditions to a nucleic acid of FIGS. 1-14. In some embodiments, the mental disorder is a mood disorder or a psychotic disorder. In some embodiments, the psychotic disorder is schizophrenia. In some embodiments, the mood disorder is a bipolar disorder or major depression.

BRIEF DESCRIPTION OF THE DRAWINGS

[0020] FIG. 1 lists genes that are differentially expressed in schizophrenic versus control patients in each of the 6 brain regions. For every gene (row) listed: the UniGene ID, GenBank Accession # ("Acc"), Gene Symbol, Chromosome # ("Chr"), and direction of change (up or down) in expression levels are listed in successive columns. The last column provides the name of the differentially expressed gene and related information, where available.

[0021] FIG. 2 shows gene ontology (GO) terms enriched in AnCg. Probe sets that showed GO term enrichment and met default FDR correction threshold criteria (<http://brainarray.mhri.med.umich.edu/Brainarray/>) are listed here. Under each enriched GO term, individual genes are listed in the rows. Information related to the LocusLink ID #, UniGene ID #, Gene Symbol and Gene Description is provided in successive columns.

[0022] FIG. 3 shows GO terms enriched in DLPFC. Probe sets that showed GO-term enrichment and met default FDR correction threshold criteria (<http://brainarray.mhri.med.umich.edu/Brainarray/>) are listed here. Under each enriched GO term, individual genes are listed in the rows. A LocusLink ID #, UniGene ID #, Gene Symbol and Gene Description are indicated in successive columns.

[0023] FIG. 4 lists 84 genes which are significantly dysregulated in both lymphoblastic and brain tissues.

[0024] FIG. 5 lists 16 genes which are significantly dysregulated in lymphoblasts only. These genes were significant by microarray and the amount and direction of change was confirmed using Q-PCR. Seven of these genes (bold in Accession Number column) exhibited statistically significant dysregulation when examined by Q-PCR and evaluated using the two-tailed t-test.

[0025] FIG. 6 lists 11 single nucleotide polymorphic markers correlated with aspartylglucosaminuria (AGA) gene expression which is dysregulated in both brain and lymphocytes of individuals with schizophrenia (see FIG. 4, infra). The regression p-values of genotype with lymphocyte gene expression are shown in the last column.

[0026] FIG. 7 lists genes involved in mood disorders and psychotic disorders.

[0027] FIG. 8 lists genes involved in mood disorders and psychotic disorders.

[0028] FIGS. 9-12 list genes differentially expressed in the amygdala that are involved in mood disorders and psychotic disorders;

[0029] FIG. 13 lists genes differentially expressed in the dorsolateral prefrontal cortex (DLPFC) that are involved in mood disorders and psychotic disorders.

[0030] FIG. 14 lists lithium responsive genes expressed in the amygdala.

[0031] FIG. 15 shows PSPHL locus genotype and PSPHL mRNA expression in BPD, MDD, SCZ, and control subjects.

DEFINITIONS

[0032] A “mental disorder” or “mental illness” or “mental disease” or “psychiatric or neuropsychiatric disease or illness or disorder” refers to mood disorders (e.g., major depression, mania, and bipolar disorders), psychotic disorders (e.g., schizophrenia, schizoaffective disorder, schizophreniform disorder, delusional disorder, brief psychotic disorder, and shared psychotic disorder), personality disorders, anxiety disorders (e.g., obsessive-compulsive disorder) as well as other mental disorders such as substance-related disorders, childhood disorders, dementia, autistic disorder, adjustment disorder, delirium, multi-infarct dementia, and Tourette’s disorder as described in Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, (DSM IV). Typically, such disorders have a complex genetic and/or a biochemical component.

[0033] “A psychotic disorder” refers to a condition that affects the mind, resulting in at least some loss of contact with reality. Symptoms of a psychotic disorder include, e.g., hallucinations, changed behavior that is not based on reality, delusions and the like. See, e.g., DSM IV. Schizophrenia, schizoaffective disorder, schizophreniform disorder, delusional disorder, brief psychotic disorder, substance-induced psychotic disorder, and shared psychotic disorder are examples of psychotic disorders.

[0034] “Schizophrenia” refers to a psychotic disorder involving a withdrawal from reality by an individual. Symptoms comprise for at least a part of a month two or more of the following symptoms: delusions (only one symptom is required if a delusion is bizarre, such as being abducted in a space ship from the sun); hallucinations (only one symptom is required if hallucinations are of at least two voices talking to one another or of a voice that keeps up a running commentary on the patient’s thoughts or actions); disorganized speech (e.g., frequent derailment or incoherence); grossly disorganized or catatonic behavior; or negative symptoms, i.e., affective flattening, alogia, or avolition. Schizophrenia encompasses disorders such as, e.g., schizoaffective disorders. Diagnosis of schizophrenia is described in, e.g., Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition (DSM IV). Types of schizophrenia include, e.g., paranoid, disorganized, catatonic, undifferentiated, and residual.

[0035] A “mood disorder” refers to disruption of feeling tone or emotional state experienced by an individual for an extensive period of time. Mood disorders include major depression disorder (i.e., unipolar disorder), mania, dysphoria, bipolar disorder, dysthymia, cyclothymia and many others. See, e.g., DSM IV.

[0036] “Major depression disorder,” “major depressive disorder,” or “unipolar disorder” refers to a mood disorder involving any of the following symptoms: persistent sad, anxious, or “empty” mood; feelings of hopelessness or pessimism; feelings of guilt, worthlessness, or helplessness; loss of interest or pleasure in hobbies and activities that were once enjoyed, including sex; decreased energy, fatigue, being “slowed down”; difficulty concentrating, remembering, or making decisions; insomnia, early-morning awakening, or oversleeping; appetite and/or weight loss or overeating and weight gain; thoughts of death or suicide or suicide attempts; restlessness or irritability; or persistent physical symptoms that do not respond to treatment, such as headaches, digestive

disorders, and chronic pain. Various subtypes of depression are described in, e.g., DSM IV.

[0037] “Bipolar disorder” is a mood disorder characterized by alternating periods of extreme moods. A person with bipolar disorder experiences cycling of moods that usually swing from being overly elated or irritable (mania) to sad and hopeless (depression) and then back again, with periods of normal mood in between. Diagnosis of bipolar disorder is described in, e.g., DSM IV. Bipolar disorders include bipolar disorder I (mania with or without major depression) and bipolar disorder II (hypomania with major depression), see, e.g., DSM IV.

[0038] Anxiety disorders, learning and memory disorders or cognitive disorders are described in DSM IV. Anxiety disorders display co-morbidity with depression, and learning and memory disorders display co-morbidity with schizophrenia.

[0039] An “agonist” refers to an agent that binds to a polypeptide or polynucleotide of the invention, stimulates, increases, activates, facilitates, enhances activation, sensitizes or up regulates the activity or expression of a polypeptide or polynucleotide of the invention.

[0040] An “antagonist” refers to an agent that inhibits expression of a polypeptide or polynucleotide of the invention or binds to, partially or totally blocks stimulation, decreases, prevents, delays activation, inactivates, desensitizes, or down regulates the activity of a polypeptide or polynucleotide of the invention.

[0041] “Inhibitors,” “activators,” and “modulators” of expression or of activity are used to refer to inhibitory, activating, or modulating molecules, respectively, identified using in vitro and in vivo assays for expression or activity, e.g., ligands, agonists, antagonists, and their homologs and mimetics. The term “modulator” includes inhibitors and activators. Inhibitors are agents that, e.g., inhibit expression of a polypeptide or polynucleotide of the invention or bind to, partially or totally block stimulation or enzymatic activity, decrease, prevent, delay activation, inactivate, desensitize, or down regulate the activity of a polypeptide or polynucleotide of the invention, e.g., antagonists. Activators are agents that, e.g., induce or activate the expression of a polypeptide or polynucleotide of the invention or bind to, stimulate, increase, open, activate, facilitate, enhance activation or enzymatic activity, sensitize or up regulate the activity of a polypeptide or polynucleotide of the invention, e.g., agonists. Modulators include naturally occurring and synthetic ligands, antagonists, agonists, small chemical molecules and the like. Assays to identify inhibitors and activators include, e.g., applying putative modulator compounds to cells, in the presence or absence of a polypeptide or polynucleotide of the invention and then determining the functional effects on a polypeptide or polynucleotide of the invention activity. Samples or assays comprising a polypeptide or polynucleotide of the invention that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of effect. Control samples (untreated with modulators) are assigned a relative activity value of 100%. Inhibition is achieved when the activity value of a polypeptide or polynucleotide of the invention relative to the control is about 80%, optionally 50% or 25-1%. Activation is achieved when the activity value of a polypeptide or polynucleotide of the invention relative to the control is 110%, optionally 150%, optionally 200-500%, or 1000-3000% higher.

[0042] The term “test compound” or “drug candidate” or “modulator” or grammatical equivalents as used herein describes any molecule, either naturally occurring or synthetic, e.g., protein, oligopeptide (e.g., from about 5 to about 25 amino acids in length, preferably from about 10 to 20 or 12 to 18 amino acids in length, preferably 12, 15, or 18 amino acids in length), small organic molecule, polysaccharide, lipid, fatty acid, polynucleotide, RNAi, oligonucleotide, etc. The test compound can be in the form of a library of test compounds, such as a combinatorial or randomized library that provides a sufficient range of diversity. Test compounds are optionally linked to a fusion partner, e.g., targeting compounds, rescue compounds, dimerization compounds, stabilizing compounds, addressable compounds, and other functional moieties. Conventionally, new chemical entities with useful properties are generated by identifying a test compound (called a “lead compound”) with some desirable property or activity, e.g., inhibiting activity, creating variants of the lead compound, and evaluating the property and activity of those variant compounds. Often, high throughput screening (HTS) methods are employed for such an analysis.

[0043] A “small organic molecule” refers to an organic molecule, either naturally occurring or synthetic, that has a molecular weight of more than about 50 Daltons and less than about 2500 Daltons, preferably less than about 2000 Daltons, preferably between about 100 to about 1000 Daltons, more preferably between about 200 to about 500 Daltons.

An “siRNA” or “RNAi” refers to a nucleic acid that forms a double stranded RNA, which double stranded RNA has the ability to reduce or inhibit expression of a gene or target gene when the siRNA expressed in the same cell as the gene or target gene. “siRNA” or “RNAi” thus refers to the double stranded RNA formed by the complementary strands. The complementary portions of the siRNA that hybridize to form the double stranded molecule typically have substantial or complete identity. In one embodiment, an siRNA refers to a nucleic acid that has substantial or complete identity to a target gene and forms a double stranded siRNA. Typically, the siRNA is at least about 15-50 nucleotides in length (e.g., each complementary sequence of the double stranded siRNA is 15-50 nucleotides in length, and the double stranded siRNA is about 15-50 base pairs in length, preferable about preferably about 20-30 base nucleotides, preferably about 20-25 or about 24-29 nucleotides in length, e.g., 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in length).

[0044] “Determining the functional effect” refers to assaying for a compound that increases or decreases a parameter that is indirectly or directly under the influence of a polynucleotide or polypeptide of the invention (such as a polynucleotide of FIGS. 1-6 or a polypeptide encoded by a gene of FIGS. 1-6), e.g., measuring physical and chemical or phenotypic effects. Such functional effects can be measured by any means known to those skilled in the art, e.g., changes in spectroscopic (e.g., fluorescence, absorbance, refractive index), hydrodynamic (e.g., shape), chromatographic, or solubility properties for the protein; measuring inducible markers or transcriptional activation of the protein; measuring binding activity or binding assays, e.g. binding to antibodies; measuring changes in ligand binding affinity; measurement of calcium influx; measurement of the accumulation of an enzymatic product of a polypeptide of the invention or depletion of an substrate; measurement of changes in protein levels of a polypeptide of the invention; measurement of RNA stability; G-protein binding; GPCR

phosphorylation or dephosphorylation; signal transduction, e.g., receptor-ligand interactions, second messenger concentrations (e.g., cAMP, IP3, or intracellular Ca^{2+}); identification of downstream or reporter gene expression (CAT, luciferase, β -gal, GFP and the like), e.g., via chemiluminescence, fluorescence, colorimetric reactions, antibody binding, inducible markers, and ligand binding assays.

[0045] Samples or assays comprising a nucleic acid or protein disclosed herein that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of inhibition. Control samples (untreated with inhibitors) are assigned a relative protein activity value of 100%. Inhibition is achieved when the activity value relative to the control is about 80%, preferably 50%, more preferably 25-0%. Activation is achieved when the activity value relative to the control (untreated with activators) is 110%, more preferably 150%, more preferably 200-500% (i.e., two to five fold higher relative to the control), more preferably 1000-3000% higher.

[0046] “Biological sample” includes sections of tissues such as biopsy and autopsy samples, and frozen sections taken for histologic purposes. Such samples include blood, serum, lymphocytes, spinal fluid, sputum, tissue, lysed cells, brain biopsy, cultured cells, e.g., primary cultures, explants, and transformed cells, stool, urine, etc. A biological sample is typically obtained from a eukaryotic organism, most preferably a mammal such as a primate, e.g., chimpanzee or human; cow; dog; cat; a rodent, e.g., guinea pig, rat, mouse; rabbit; or a bird; reptile; or fish. Biological samples can be used to examine nucleic acids and proteins.

[0047] “Antibody” refers to a polypeptide substantially encoded by an immunoglobulin gene or immunoglobulin genes, or fragments thereof which specifically bind and recognize an analyte (antigen). The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

[0048] An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one “light” (about 25 kD) and one “heavy” chain (about 50-70 kD). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (V_L) and variable heavy chain (V_H) refer to these light and heavy chains respectively.

[0049] Antibodies exist, e.g., as intact immunoglobulins or as a number of well-characterized fragments produced by digestion with various peptidases. Thus, for example, pepsin digests an antibody below the disulfide linkages in the hinge region to produce $F(ab)'_2$, a dimer of Fab which itself is a light chain joined to V_H-C_H1 by a disulfide bond. The $F(ab)'_2$ may be reduced under mild conditions to break the disulfide linkage in the hinge region, thereby converting the $F(ab)'_2$ dimer into an Fab' monomer. The Fab' monomer is essentially an Fab with part of the hinge region (see, Paul (Ed.) *Fundamental Immunology*, Third Edition, Raven Press, NY (1993)). While various antibody fragments are defined in terms of the digestion of an intact antibody, one of skill will appreciate that such fragments may be synthesized de novo either chemi-

cally or by utilizing recombinant DNA methodology. Thus, the term antibody, as used herein, also includes antibody fragments either produced by the modification of whole antibodies or those synthesized de novo using recombinant DNA methodologies (e.g., single chain Fv).

[0050] The terms “peptidomimetic” and “mimetic” refer to a synthetic chemical compound that has substantially the same structural and functional characteristics of the polynucleotides, polypeptides, antagonists or agonists of the invention. Peptide analogs are commonly used in the pharmaceutical industry as non-peptide drugs with properties analogous to those of the template peptide. These types of non-peptide compound are termed “peptide mimetics” or “peptidomimetics” (Fauchere, *Adv. Drug Res.* 15:29 (1986); Veber and Freidinger *TINS* p. 392 (1985); and Evans et al., *J. Med. Chem.* 30:1229 (1987), which are incorporated herein by reference). Peptide mimetics that are structurally similar to therapeutically useful peptides may be used to produce an equivalent or enhanced therapeutic or prophylactic effect. Generally, peptidomimetics are structurally similar to a paradigm polypeptide (i.e., a polypeptide that has a biological or pharmacological activity), such as a CCX CKR, but have one or more peptide linkages optionally replaced by a linkage selected from the group consisting of, e.g., $-\text{CH}_2\text{NH}-$, $-\text{CH}_2\text{S}-$, $-\text{CH}_2-\text{CH}_2-$, $-\text{CH}=\text{CH}-$ (cis and trans), $-\text{COCH}_2-$, $-\text{CH}(\text{OH})\text{CH}_2-$, and $-\text{CH}_2\text{SO}-$. The mimetic can be either entirely composed of synthetic, non-natural analogues of amino acids, or, is a chimeric molecule of partly natural peptide amino acids and partly non-natural analogs of amino acids. The mimetic can also incorporate any amount of natural amino acid conservative substitutions as long as such substitutions also do not substantially alter the mimetic’s structure and/or activity. For example, a mimetic composition is within the scope of the invention if it is capable of carrying out the binding or enzymatic activities of a polypeptide or polynucleotide of the invention or inhibiting or increasing the enzymatic activity or expression of a polypeptide or polynucleotide of the invention.

[0051] The term “gene” means the segment of DNA involved in producing a polypeptide chain; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons).

[0052] The term “isolated,” when applied to a nucleic acid or protein, denotes that the nucleic acid or protein is essentially free of other cellular components with which it is associated in the natural state. It is preferably in a homogeneous state although it can be in either a dry or aqueous solution. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein that is the predominant species present in a preparation is substantially purified. In particular, an isolated gene is separated from open reading frames that flank the gene and encode a protein other than the gene of interest. The term “purified” denotes that a nucleic acid or protein gives rise to essentially one band in an electrophoretic gel. Particularly, it means that the nucleic acid or protein is at least 85% pure, more preferably at least 95% pure, and most preferably at least 99% pure.

[0053] The term “nucleic acid” or “polynucleotide” refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form. Unless specifically limited, the term encompasses nucleic acids contain-

ing known analogues of natural nucleotides that have similar binding properties as the reference nucleic acid and are metabolized in a manner similar to naturally occurring nucleotides. Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions), alleles, orthologs, SNPs, haplotypes, and complementary sequences as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating sequences in which the third position of one or more selected (or all) codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., *Nucleic Acid Res.* 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.* 260:2605-2608 (1985); and Cassol et al. (1992); Rossolini et al., *Mol. Cell. Probes* 8:91-98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, and mRNA encoded by a gene.

[0054] The terms “polypeptide,” “peptide,” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymers. As used herein, the terms encompass amino acid chains of any length, including full-length proteins (i.e., antigens), wherein the amino acid residues are linked by covalent peptide bonds.

[0055] The term “amino acid” refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, i.e., an α carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. “Amino acid mimetics” refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid.

[0056] Amino acids may be referred to herein by either the commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

[0057] “Conservatively modified variants” applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, “conservatively modified variants” refers to those nucleic acids that encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are “silent variations,”

which are one species of conservatively modified variations. Every nucleic acid sequence herein that encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid that encodes a polypeptide is implicit in each described sequence.

[0058] As to amino acid sequences, one of skill will recognize that individual substitutions, deletions or additions to a nucleic acid, peptide, polypeptide, or protein sequence which alters, adds or deletes a single amino acid or a small percentage of amino acids in the encoded sequence is a “conservatively modified variant” where the alteration results in the substitution of an amino acid with a chemically similar amino acid. Conservative substitution tables providing functionally similar amino acids are well known in the art. Such conservatively modified variants are in addition to and do not exclude polymorphic variants, interspecies homologs, and alleles of the invention.

[0059] The following eight groups each contain amino acids that are conservative substitutions for one another:

1) Alanine (A), Glycine (G);

[0060] 2) Aspartic acid (D), Glutamic acid (E);

3) Asparagine (N), Glutamine (Q);

4) Arginine (R), Lysine (K);

5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V);

6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W);

7) Serine (S), Threonine (T); and

8) Cysteine (C), Methionine (M)

[0061] (see, e.g., Creighton, *Proteins* (1984)).

[0062] “Percentage of sequence identity” is determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the result by 100 to yield the percentage of sequence identity.

[0063] The terms “identical” or percent “identity,” in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., 60% identity, optionally 65%, 70%, 75%, 80%, 85%, 90%, or 95% identity over a specified region), when compared and aligned for maximum correspondence over a comparison window, or designated region as measured using one of the following sequence comparison algorithms or by manual alignment and visual inspection. Such sequences are then said to be “substantially identical.” This definition also refers to the complement of a test sequence. Optionally, the identity exists over a

region that is at least about 50 nucleotides in length, or more preferably over a region that is 100 to 500 or 1000 or more nucleotides in length.

[0064] For sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

[0065] A “comparison window”, as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith and Waterman (1970) *Adv. Appl. Math.* 2:482c, by the homology alignment algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443, by the search for similarity method of Pearson and Lipman (1988) *Proc. Nat’l. Acad. Sci. USA* 85:2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by manual alignment and visual inspection (see, e.g., Ausubel et al., *Current Protocols in Molecular Biology* (1995 supplement)).

[0066] An example of an algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al. (1977) *Nuc. Acids Res.* 25:3389-3402, and Altschul et al. (1990) *J. Mol. Biol.* 215:403-410, respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database sequence. T is referred to as the neighborhood word score threshold (Altschul et al., supra). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always >0) and N (penalty score for mismatching residues; always <0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucle-

otide sequences) uses as defaults a wordlength (W) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci. USA* 89:10915) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

[0067] The BLAST algorithm also performs a statistical analysis of the similarity between two sequences (see, e.g., Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-5787). One measure of similarity provided by the BLAST algorithm is the smallest sum probability (P(N)), which provides an indication of the probability by which a match between two nucleotide or amino acid sequences would occur by chance. For example, a nucleic acid is considered similar to a reference sequence if the smallest sum probability in a comparison of the test nucleic acid to the reference nucleic acid is less than about 0.2, more preferably less than about 0.01, and most preferably less than about 0.001.

[0068] An indication that two nucleic acid sequences or polypeptides are substantially identical is that the polypeptide encoded by the first nucleic acid is immunologically cross reactive with the antibodies raised against the polypeptide encoded by the second nucleic acid, as described below. Thus, a polypeptide is typically substantially identical to a second polypeptide, for example, where the two peptides differ only by conservative substitutions. Another indication that two nucleic acid sequences are substantially identical is that the two molecules or their complements hybridize to each other under stringent conditions, as described below. Yet another indication that two nucleic acid sequences are substantially identical is that the same primers can be used to amplify the sequence.

[0069] The phrase “selectively (or specifically) hybridizes to” refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

[0070] The phrase “stringent hybridization conditions” refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology—Hybridization with Nucleic Probes*, “Overview of principles of hybridization and the strategy of nucleic acid assays” (1993). Generally, stringent conditions are selected to be about 5-10° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m , 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent con-

ditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, optionally 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, 5×SSC, and 1% SDS, incubating at 42° C., or 5×SSC, 1% SDS, incubating at 65° C., with wash in 0.2×SSC, and 0.1% SDS at 65° C. Such washes can be performed for 5, 15, 30, 60, 120, or more minutes. Nucleic acids that hybridize to the genes listed in FIGS. 1-6 are encompassed by the invention.

[0071] Nucleic acids that do not hybridize to each other under stringent conditions are still substantially identical if the polypeptides that they encode are substantially identical. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Exemplary “moderately stringent hybridization conditions” include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 1×SSC at 45° C. Such washes can be performed for 5, 15, 30, 60, 120, or more minutes. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

[0072] For PCR, a temperature of about 36° C. is typical for low stringency amplification, although annealing temperatures may vary between about 32° C. and 48° C. depending on primer length. For high stringency PCR amplification, a temperature of about 62° C. is typical, although high stringency annealing temperatures can range from about 50° C. to about 65° C., depending on the primer length and specificity. Typical cycle conditions for both high and low stringency amplifications include a denaturation phase of 90° C.-95° C. for 30 sec.-2 min., an annealing phase lasting 30 sec.-2 min., and an extension phase of about 72° C. for 1-2 min. Protocols and guidelines for low and high stringency amplification reactions are provided, e.g., in Innis et al., *PCR Protocols, A Guide to Methods and Applications* (1990).

[0073] The phrase “a nucleic acid sequence encoding” refers to a nucleic acid that contains sequence information for a structural RNA such as rRNA, a tRNA, or the primary amino acid sequence of a specific protein or peptide, or a binding site for a trans-acting regulatory agent. This phrase specifically encompasses degenerate codons (i.e., different codons which encode a single amino acid) of the native sequence or sequences which may be introduced to conform with codon preference in a specific host cell.

[0074] The term “recombinant” when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (nonrecombinant) form of the cell or express native genes that are otherwise abnormally expressed, under-expressed or not expressed at all.

[0075] The term “heterologous” when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two

or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

[0076] An “expression vector” is a nucleic acid construct, generated recombinantly or synthetically, with a series of specified nucleic acid elements that permit transcription of a particular nucleic acid in a host cell. The expression vector can be part of a plasmid, virus, or nucleic acid fragment. Typically, the expression vector includes a nucleic acid to be transcribed operably linked to a promoter.

[0077] The phrase “specifically (or selectively) binds to an antibody” or “specifically (or selectively) immunoreactive with”, when referring to a protein or peptide, refers to a binding reaction which is determinative of the presence of the protein in the presence of a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein and do not bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, antibodies raised against a protein having an amino acid sequence encoded by any of the polynucleotides of the invention can be selected to obtain antibodies specifically immunoreactive with that protein and not with other proteins, except for polymorphic variants. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays, Western blots, or immunohistochemistry are routinely used to select monoclonal antibodies specifically immunoreactive with a protein. See, Harlow and Lane *Antibodies, A Laboratory Manual*, Cold Spring Harbor Publications, N.Y. (1988) for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity. Typically, a specific or selective reaction will be at least twice the background signal or noise and more typically more than 10 to 100 times background.

[0078] One who is “predisposed for a mental disorder” as used herein means a person who has an inclination or a higher likelihood of developing a mental disorder when compared to an average person in the general population.

DETAILED DESCRIPTION OF THE INVENTION

I. Introduction

[0079] To understand the complex genetic basis of mental disorders, the present invention provides studies that have been conducted to investigate the expression patterns of genes that are differentially expressed specifically in central nervous system of subjects with psychotic and mood disorders. The large spectrum of symptoms associated with mental disorders is a reflection of the complex genetic basis and complex gene expression patterns in patients with mental disorders. Different combinations of the genes disclosed herein can be responsible for one or more mental disorders. Furthermore, brain pathways or circuits as well as subcellular pathways are important for understanding the development and diagnosis of mental disorders. The selected brain regions described herein (anterior cingulate cortex (AnCg), dorsolateral prefrontal cortex (DLPFC), cerebellar cortex (CB), entorhinal cortex (ERC), superior temporal gyms (STG),

parietal cortex (PC), nucleus accumbens (nAcc), ventral thalamus (VThal), medial thalamus (MThal), amygdala (AMY) and/or the hippocampus (HC)) are implicated in the clinical symptoms of mental disorders such as psychotic and mood disorders. Brain imaging studies focusing on particular brain regions, cytoarchitectural changes in brain regions, expression of key neurotransmitters or related molecules in brain regions, and subcellular pathways in brain regions all contribute to the development of mental disorders, and thus are an important consideration in the diagnosis and therapeutic uses described herein.

[0080] The present invention demonstrates the altered expression (either higher or lower) of the genes of FIGS. 1-14 at the mRNA or protein level in various regions of the brain (e.g., FIGS. 1-4 and 6) or lymphocytes (e.g., FIGS. 4-6) of patients with mental disorders (e.g., schizophrenia, MDD and BPD) in comparison with normal individuals. This invention thus provides methods for diagnosis of mental disorders, e.g., schizophrenia, MDD and BPD, and the like, and other mental disorders by detecting the level of a transcript or translation product of the genes listed in FIGS. 1-14 as well as their corresponding biochemical pathways. The chromosomal location of such genes can be used to discover other genes in the region that are linked to development of a particular disorder. FIG. 6 of the invention also provides single nucleotide polymorphic markers which are related (in cis or trans) to regulatory sites associated with AGA.

[0081] The invention further provides methods of identifying a compound useful for the treatment of such disorders by selecting compounds that modulates the functional effect of the translation products or the expression of the transcripts described herein. The invention also provides for methods of treating patients with such mental disorders, e.g., by administering the compounds of the invention or by gene therapy. Therapeutic compounds include antibodies, peptides, antisense molecules, siRNA, and small organic molecules.

[0082] The genes and the polypeptides that they encode, which are associated with psychotic and mood disorders, are useful for facilitating the design and development of various molecular diagnostic tools such as GeneChips™ containing probe sets specific for all or selected mental disorders, including but not limited to psychotic and mood disorders, and as an ante- and/or post-natal diagnostic tool for screening newborns in concert with genetic counseling. Other diagnostic applications include evaluation of disease susceptibility, prognosis, and monitoring of disease or treatment process, as well as providing individualized medicine via predictive drug profiling systems, e.g., by correlating specific genomic motifs with the clinical response of a patient to individual drugs. In addition, the present invention is useful for multiplex SNP or haplotype profiling, including but not limited to the identification of pharmacogenetic targets at the gene, mRNA, protein, and pathway level (see, e.g, Basile V S, Masellis M, Potkin S G, Kennedy J L. Pharmacogenomics in schizophrenia: the quest for individualized therapy. *Hum Mol Genet.* 2002 Oct. 1; 11(20):2517-30). Profiling of splice variants is also useful for diagnostic and therapeutic applications. Diagnostic kits are contemplated by the present invention, and include arrays, nanoparticles, and magnetic beads. Marker combinations can provide useful diagnosis. Brain expression patterns, regions, pathways, and circuits can be used for in vivo imaging and diagnosis.

[0083] The genes and the polypeptides that they encode, described herein, as also useful as drug targets for the devel-

opment of therapeutic drugs for the treatment or prevention of mental disorders including, but not limited to, psychotic and mood disorders. Mental disorders have a high co-morbidity with other neurological disorders, such as Parkinson's disease or Alzheimer's. Therefore, the present invention can be used for diagnosis and treatment of patients with multiple disease states that include a mental disorder such as a psychotic disorder.

[0084] Antipsychotic medicines are in general equally effect for the treatment of schizophrenia, but act by different mechanisms. The similar effectiveness of the drugs for treatment of schizophrenia suggests that they act through a yet as unidentified common pathway. As demonstrated by the results shown herein, these drugs regulate a common gene, and/or a common group of genes as well as a unique set of genes.

[0085] The genes listed herein can be used to provide a differential diagnosis or prognosis of mood and psychotic disorders. In some cases, differentially expressed genes can be used to predict and treat particular symptoms or outcomes, such as suicide attempt. The therapeutic agents described herein can be used in combination with known therapeutics. Nucleic acid therapeutics can be delivered using adenoviral vectors, while peptides, nucleic acids, and other therapeutic molecules can be delivered using nanoparticles and translocation peptides. Orally available peptides can be made using D-amino acids or pegylation, and serum half life can be extended using albumin conjugation and the like.

II. General Recombinant Nucleic Acid Methods for Use with the Invention

[0086] In numerous embodiments of the present invention, polynucleotides of the invention will be isolated and cloned using recombinant methods. Such polynucleotides include, e.g., those listed in FIGS. 1-14, which can be used for, e.g., protein expression or during the generation of variants, derivatives, expression cassettes, to monitor gene expression, for the isolation or detection of sequences of the invention in different species, for diagnostic purposes in a patient, e.g., to detect mutations or to detect expression levels of nucleic acids or polypeptides of the invention. In some embodiments, the sequences of the invention are operably linked to a heterologous promoter. In one embodiment, the nucleic acids of the invention are from any mammal, including, in particular, e.g., a human, a mouse, a rat, a primate, etc.

A. General Recombinant Nucleic Acids Methods

[0087] This invention relies on routine techniques in the field of recombinant genetics. Basic texts disclosing the general methods of use in this invention include Sambrook et al., *Molecular Cloning, A Laboratory Manual* (3rd ed. 2001); Kriegler, *Gene Transfer and Expression: A Laboratory Manual* (1990); and *Current Protocols in Molecular Biology* (Ausubel et al., eds., 1994).

[0088] For nucleic acids, sizes are given in either kilobases (kb) or base pairs (bp). These are estimates derived from agarose or acrylamide gel electrophoresis, from sequenced nucleic acids, or from published DNA sequences. For proteins, sizes are given in kilodaltons (kDa) or amino acid residue numbers. Proteins sizes are estimated from gel electrophoresis, from sequenced proteins, from derived amino acid sequences, or from published protein sequences.

[0089] Oligonucleotides that are not commercially available can be chemically synthesized according to the solid phase phosphoramidite triester method first described by Beaucage & Caruthers, *Tetrahedron Letts.* 22:1859-1862 (1981), using an automated synthesizer, as described in Van Devanter et. al., *Nucleic Acids Res.* 12:6159-6168 (1984). Purification of oligonucleotides is by either native acrylamide gel electrophoresis or by anion-exchange HPLC as described in Pearson & Reanier, *J. Chrom.* 255:137-149 (1983).

[0090] The sequence of the cloned genes and synthetic oligonucleotides can be verified after cloning using, e.g., the chain termination method for sequencing double-stranded templates of Wallace et al., *Gene* 16:21-26 (1981).

B. Cloning Methods for the Isolation of Nucleotide Sequences Encoding Desired Proteins

[0091] In general, the nucleic acids encoding the subject proteins are cloned from DNA sequence libraries that are made to encode cDNA or genomic DNA. The particular sequences can be located by hybridizing with an oligonucleotide probe, the sequence of which can be derived from the sequences of the genes and/or SNPs listed in FIGS. 1-6, which provide a reference for PCR primers and defines suitable regions for isolating specific probes. Alternatively, where the sequence is cloned into an expression library, the expressed recombinant protein can be detected immunologically with antisera or purified antibodies made against a polypeptide comprising an amino acid sequence encoded by a gene listed in FIGS. 1-14.

[0092] Methods for making and screening genomic and cDNA libraries are well known to those of skill in the art (see, e.g., Gubler and Hoffman *Gene* 25:263-269 (1983); Benton and Davis *Science*, 196:180-182 (1977); and Sambrook, supra). Brain cells are an example of suitable cells to isolate RNA and cDNA sequences of the invention.

[0093] Briefly, to make the cDNA library, one should choose a source that is rich in mRNA. The mRNA can then be made into cDNA, ligated into a recombinant vector, and transfected into a recombinant host for propagation, screening and cloning. For a genomic library, the DNA is extracted from a suitable tissue and either mechanically sheared or enzymatically digested to yield fragments of preferably about 5-100 kb. The fragments are then separated by gradient centrifugation from undesired sizes and are constructed in bacteriophage lambda vectors. These vectors and phage are packaged in vitro, and the recombinant phages are analyzed by plaque hybridization. Colony hybridization is carried out as generally described in Grunstein et al., *Proc. Natl. Acad. Sci. USA.*, 72:3961-3965 (1975).

[0094] An alternative method combines the use of synthetic oligonucleotide primers with polymerase extension on an mRNA or DNA template. Suitable primers can be designed from specific sequences of the invention. This polymerase chain reaction (PCR) method amplifies the nucleic acids encoding the protein of interest directly from mRNA, cDNA, genomic libraries or cDNA libraries. Restriction endonuclease sites can be incorporated into the primers. Polymerase chain reaction or other in vitro amplification methods may also be useful, for example, to clone nucleic acids encoding specific proteins and express said proteins, to synthesize nucleic acids that will be used as probes for detecting the presence of mRNA encoding a polypeptide of the invention in physiological samples, for nucleic acid sequencing, or for other purposes (see, U.S. Pat. Nos. 4,683,195 and 4,683,202).

Genes amplified by a PCR reaction can be purified from agarose gels and cloned into an appropriate vector.

[0095] Appropriate primers and probes for identifying polynucleotides of the invention from mammalian tissues can be derived from the sequences provided herein. For a general overview of PCR, see, Innis et al. *PCR Protocols: A Guide to Methods and Applications*, Academic Press, San Diego (1990).

[0096] Synthetic oligonucleotides can be used to construct genes. This is done using a series of overlapping oligonucleotides, usually 40-120 bp in length, representing both the sense and anti-sense strands of the gene. These DNA fragments are then annealed, ligated and cloned.

[0097] A gene encoding a polypeptide of the invention can be cloned using intermediate vectors before transformation into mammalian cells for expression. These intermediate vectors are typically prokaryote vectors or shuttle vectors. The proteins can be expressed in either prokaryotes, using standard methods well known to those of skill in the art, or eukaryotes as described infra.

III. Purification of Proteins of the Invention

[0098] Either naturally occurring or recombinant polypeptides of the invention can be purified for use in functional assays. Naturally occurring polypeptides, e.g., polypeptides encoded by genes listed in FIGS. 1-14, can be purified, for example, from mouse or human tissue such as brain or any other source of an ortholog. Recombinant polypeptides can be purified from any suitable expression system.

[0099] The polypeptides of the invention may be purified to substantial purity by standard techniques, including selective precipitation with such substances as ammonium sulfate; column chromatography, immunopurification methods, and others (see, e.g., Scopes, *Protein Purification Principles and Practice* (1982); U.S. Pat. No. 4,673,641; Ausubel et al., supra; and Sambrook et al., supra).

[0100] A number of procedures can be employed when recombinant polypeptides are purified. For example, proteins having established molecular adhesion properties can be reversibly fused to polypeptides of the invention. With the appropriate ligand, the polypeptides can be selectively adsorbed to a purification column and then freed from the column in a relatively pure form. The fused protein is then removed by enzymatic activity. Finally the polypeptide can be purified using immunoaffinity columns.

A. Purification of Proteins from Recombinant Bacteria

[0101] When recombinant proteins are expressed by the transformed bacteria in large amounts, typically after promoter induction, although expression can be constitutive, the proteins may form insoluble aggregates. There are several protocols that are suitable for purification of protein inclusion bodies. For example, purification of aggregate proteins (hereinafter referred to as inclusion bodies) typically involves the extraction, separation and/or purification of inclusion bodies by disruption of bacterial cells typically, but not limited to, by incubation in a buffer of about 100-150 µg/ml lysozyme and 0.1% Nonidet P40, a non-ionic detergent. The cell suspension can be ground using a Polytron grinder (Brinkman Instruments, Westbury, N.Y.). Alternatively, the cells can be sonicated on ice. Alternate methods of lysing bacteria are described in Ausubel et al. and Sambrook et al., both supra, and will be apparent to those of skill in the art.

[0102] The cell suspension is generally centrifuged and the pellet containing the inclusion bodies resuspended in buffer

which does not dissolve but washes the inclusion bodies, e.g., 20 mM Tris-HCl (pH 7.2), 1 mM EDTA, 150 mM NaCl and 2% Triton-X 100, a non-ionic detergent. It may be necessary to repeat the wash step to remove as much cellular debris as possible. The remaining pellet of inclusion bodies may be resuspended in an appropriate buffer (e.g., 20 mM sodium phosphate, pH 6.8, 150 mM NaCl). Other appropriate buffers will be apparent to those of skill in the art.

[0103] Following the washing step, the inclusion bodies are solubilized by the addition of a solvent that is both a strong hydrogen acceptor and a strong hydrogen donor (or a combination of solvents each having one of these properties). The proteins that formed the inclusion bodies may then be renatured by dilution or dialysis with a compatible buffer. Suitable solvents include, but are not limited to, urea (from about 4 M to about 8 M), formamide (at least about 80%, volume/volume basis), and guanidine hydrochloride (from about 4 M to about 8 M). Some solvents that are capable of solubilizing aggregate-forming proteins, such as SDS (sodium dodecyl sulfate) and 70% formic acid, are inappropriate for use in this procedure due to the possibility of irreversible denaturation of the proteins, accompanied by a lack of immunogenicity and/or activity. Although guanidine hydrochloride and similar agents are denaturants, this denaturation is not irreversible and renaturation may occur upon removal (by dialysis, for example) or dilution of the denaturant, allowing re-formation of the immunologically and/or biologically active protein of interest. After solubilization, the protein can be separated from other bacterial proteins by standard separation techniques.

[0104] Alternatively, it is possible to purify proteins from bacteria periplasm. Where the protein is exported into the periplasm of the bacteria, the periplasmic fraction of the bacteria can be isolated by cold osmotic shock in addition to other methods known to those of skill in the art (see, Ausubel et al., supra). To isolate recombinant proteins from the periplasm, the bacterial cells are centrifuged to form a pellet. The pellet is resuspended in a buffer containing 20% sucrose. To lyse the cells, the bacteria are centrifuged and the pellet is resuspended in ice-cold 5 mM MgSO₄ and kept in an ice bath for approximately 10 minutes. The cell suspension is centrifuged and the supernatant decanted and saved. The recombinant proteins present in the supernatant can be separated from the host proteins by standard separation techniques well known to those of skill in the art.

B. Standard Protein Separation Techniques for Purifying Proteins

1. Solubility Fractionation

[0105] Often as an initial step, and if the protein mixture is complex, an initial salt fractionation can separate many of the unwanted host cell proteins (or proteins derived from the cell culture media) from the recombinant protein of interest. The preferred salt is ammonium sulfate. Ammonium sulfate precipitates proteins by effectively reducing the amount of water in the protein mixture. Proteins then precipitate on the basis of their solubility. The more hydrophobic a protein is, the more likely it is to precipitate at lower ammonium sulfate concentrations. A typical protocol is to add saturated ammonium sulfate to a protein solution so that the resultant ammonium sulfate concentration is between 20-30%. This will precipitate the most hydrophobic proteins. The precipitate is discarded (unless the protein of interest is hydrophobic) and

ammonium sulfate is added to the supernatant to a concentration known to precipitate the protein of interest. The precipitate is then solubilized in buffer and the excess salt removed if necessary, through either dialysis or diafiltration. Other methods that rely on solubility of proteins, such as cold ethanol precipitation, are well known to those of skill in the art and can be used to fractionate complex protein mixtures.

2. Size Differential Filtration

[0106] Based on a calculated molecular weight, a protein of greater and lesser size can be isolated using ultrafiltration through membranes of different pore sizes (for example, Amicon or Millipore membranes). As a first step, the protein mixture is ultrafiltered through a membrane with a pore size that has a lower molecular weight cut-off than the molecular weight of the protein of interest. The retentate of the ultrafiltration is then ultrafiltered against a membrane with a molecular cut off greater than the molecular weight of the protein of interest. The recombinant protein will pass through the membrane into the filtrate. The filtrate can then be chromatographed as described below.

3. Column Chromatography

[0107] The proteins of interest can also be separated from other proteins on the basis of their size, net surface charge, hydrophobicity and affinity for ligands. In addition, antibodies raised against proteins can be conjugated to column matrices and the proteins immunopurified. All of these methods are well known in the art.

[0108] It will be apparent to one of skill that chromatographic techniques can be performed at any scale and using equipment from many different manufacturers (e.g., Pharmacia Biotech).

IV. Detection of Gene Expression

[0109] Those of skill in the art will recognize that detection of expression of polynucleotides of the invention has many uses. For example, as discussed herein, detection of the level of polypeptides or polynucleotides of the invention in a patient is useful for diagnosing mood disorders or psychotic disorder or a predisposition for a mood disorder or psychotic disorder. Moreover, detection of gene expression is useful to identify modulators of expression of the polypeptides or polynucleotides of the invention.

[0110] A variety of methods of specific DNA and RNA measurement using nucleic acid hybridization techniques are known to those of skill in the art (see, Sambrook, supra). Some methods involve an electrophoretic separation (e.g., Southern blot for detecting DNA, and Northern blot for detecting RNA), but measurement of DNA and RNA can also be carried out in the absence of electrophoretic separation (e.g., by dot blot). Southern blot of genomic DNA (e.g., from a human) can be used for screening for restriction fragment length polymorphism (RFLP) to detect the presence of a genetic disorder affecting a polypeptide of the invention.

[0111] The selection of a nucleic acid hybridization format is not critical. A variety of nucleic acid hybridization formats are known to those skilled in the art. For example, common formats include sandwich assays and competition or displacement assays. Hybridization techniques are generally described in Hames and Higgins *Nucleic Acid Hybridization, A Practical Approach*, IRL Press (1985); Gall and Pardue,

Proc. Natl. Acad. Sci. U.S.A., 63:378-383 (1969); and John et al. *Nature*, 223:582-587 (1969).

[0112] Detection of a hybridization complex may require the binding of a signal-generating complex to a duplex of target and probe polynucleotides or nucleic acids. Typically, such binding occurs through ligand and anti-ligand interactions as between a ligand-conjugated probe and an anti-ligand conjugated with a signal. The binding of the signal generation complex is also readily amenable to accelerations by exposure to ultrasonic energy.

[0113] The label may also allow indirect detection of the hybridization complex. For example, where the label is a hapten or antigen, the sample can be detected by using antibodies. In these systems, a signal is generated by attaching fluorescent or enzyme molecules to the antibodies or in some cases, by attachment to a radioactive label (see, e.g., Tijssen, "Practice and Theory of Enzyme Immunoassays," *Laboratory Techniques in Biochemistry and Molecular Biology*, Burdon and van Knippenberg Eds., Elsevier (1985), pp. 9-20).

[0114] The probes are typically labeled either directly, as with isotopes, chromophores, lumiphores, chromogens, or indirectly, such as with biotin, to which a streptavidin complex may later bind. Thus, the detectable labels used in the assays of the present invention can be primary labels (where the label comprises an element that is detected directly or that produces a directly detectable element) or secondary labels (where the detected label binds to a primary label, e.g., as is common in immunological labeling). Typically, labeled signal nucleic acids are used to detect hybridization. Complementary nucleic acids or signal nucleic acids may be labeled by any one of several methods typically used to detect the presence of hybridized polynucleotides. The most common method of detection is the use of autoradiography with ^3H , ^{125}I , ^{35}S , ^{14}C , or ^{32}P -labeled probes or the like.

[0115] Other labels include, e.g., ligands that bind to labeled antibodies, fluorophores, chemiluminescent agents, enzymes, and antibodies which can serve as specific binding pair members for a labeled ligand. An introduction to labels, labeling procedures and detection of labels is found in Polak and Van Noorden *Introduction to Immunocytochemistry*, 2nd ed., Springer Verlag, NY (1997); and in Haugland *Handbook of Fluorescent Probes and Research Chemicals*, a combined handbook and catalogue Published by Molecular Probes, Inc. (1996).

[0116] In general, a detector which monitors a particular probe or probe combination is used to detect the detection reagent label. Typical detectors include spectrophotometers, phototubes and photodiodes, microscopes, scintillation counters, cameras, film and the like, as well as combinations thereof. Examples of suitable detectors are widely available from a variety of commercial sources known to persons of skill in the art. Commonly, an optical image of a substrate comprising bound labeling moieties is digitized for subsequent computer analysis.

[0117] Most typically, the amount of RNA is measured by quantifying the amount of label fixed to the solid support by binding of the detection reagent. Typically, the presence of a modulator during incubation will increase or decrease the amount of label fixed to the solid support relative to a control incubation which does not comprise the modulator, or as compared to a baseline established for a particular reaction type. Means of detecting and quantifying labels are well known to those of skill in the art.

[0118] In preferred embodiments, the target nucleic acid or the probe is immobilized on a solid support. Solid supports suitable for use in the assays of the invention are known to those of skill in the art. As used herein, a solid support is a matrix of material in a substantially fixed arrangement.

[0119] A variety of automated solid-phase assay techniques are also appropriate. For instance, very large scale immobilized polymer arrays (VLSIPS™), available from Affymetrix, Inc. (Santa Clara, Calif.) can be used to detect changes in expression levels of a plurality of genes involved in the same regulatory pathways simultaneously. See, Tijssen, supra., Fodor et al. (1991) *Science*, 251: 767-777; Sheldon et al. (1993) *Clinical Chemistry* 39(4): 718-719, and Kozal et al. (1996) *Nature Medicine* 2(7): 753-759.

[0120] Detection can be accomplished, for example, by using a labeled detection moiety that binds specifically to duplex nucleic acids (e.g., an antibody that is specific for RNA-DNA duplexes). One preferred example uses an antibody that recognizes DNA-RNA heteroduplexes in which the antibody is linked to an enzyme (typically by recombinant or covalent chemical bonding). The antibody is detected when the enzyme reacts with its substrate, producing a detectable product. Coutlee et al. (1989) *Analytical Biochemistry* 181: 153-162; Bogulayski (1986) et al. *J. Immunol. Methods* 89:123-130; Prooijen-Knegt (1982) *Exp. Cell Res.* 141:397-407; Rudkin (1976) *Nature* 265:472-473, Stollar (1970) *Proc. Nat'l Acad. Sci. USA* 65:993-1000; Ballard (1982) *Mol. Immunol.* 19:793-799; Pisetsky and Caster (1982) *Mol. Immunol.* 19:645-650; Viscidi et al. (1988) *J. Clin. Microbiol.* 41:199-209; and Kiney et al. (1989) *J. Clin. Microbiol.* 27:6-12 describe antibodies to RNA duplexes, including homo and heteroduplexes. Kits comprising antibodies specific for DNA:RNA hybrids are available, e.g., from Digene Diagnostics, Inc. (Beltsville, Md.).

[0121] In addition to available antibodies, one of skill in the art can easily make antibodies specific for nucleic acid duplexes using existing techniques, or modify those antibodies that are commercially or publicly available. In addition to the art referenced above, general methods for producing polyclonal and monoclonal antibodies are known to those of skill in the art (see, e.g., Paul (3rd ed.) *Fundamental Immunology* Raven Press, Ltd., NY (1993); Coligan *Current Protocols in Immunology* Wiley/Greene, N.Y. (1991); Harlow and Lane *Antibodies: A Laboratory Manual* Cold Spring Harbor Press, NY (1988); Stites et al. (eds.) *Basic and Clinical Immunology* (4th ed.) Lange Medical Publications, Los Altos, Calif., and references cited therein; Goding *Monoclonal Antibodies: Principles and Practice* (2d ed.) Academic Press, New York, N.Y., (1986); and Kohler and Milstein *Nature* 256: 495-497 (1975)). Other suitable techniques for antibody preparation include selection of libraries of recombinant antibodies in phage or similar vectors (see, Huse et al. *Science* 246:1275-1281 (1989); and Ward et al. *Nature* 341:544-546 (1989)). Specific monoclonal and polyclonal antibodies and antisera will usually bind with a K_D of at least about 0.1 μM , preferably at least about 0.01 μM or better, and most typically and preferably, 0.001 μM or better.

[0122] The nucleic acids used in this invention can be either positive or negative probes. Positive probes bind to their targets and the presence of duplex formation is evidence of the presence of the target. Negative probes fail to bind to the suspect target and the absence of duplex formation is evidence of the presence of the target. For example, the use of a wild type specific nucleic acid probe or PCR primers may

serve as a negative probe in an assay sample where only the nucleotide sequence of interest is present.

[0123] The sensitivity of the hybridization assays may be enhanced through use of a nucleic acid amplification system that multiplies the target nucleic acid being detected. Examples of such systems include the polymerase chain reaction (PCR) system, in particular RT-PCR or real time PCR, and the ligase chain reaction (LCR) system. Other methods recently described in the art are the nucleic acid sequence based amplification (NASBA, Cangene, Mississauga, Ontario) and Q Beta Replicase systems. These systems can be used to directly identify mutants where the PCR or LCR primers are designed to be extended or ligated only when a selected sequence is present. Alternatively, the selected sequences can be generally amplified using, for example, nonspecific PCR primers and the amplified target region later probed for a specific sequence indicative of a mutation.

[0124] An alternative means for determining the level of expression of the nucleic acids of the present invention is in situ hybridization. In situ hybridization assays are well known and are generally described in Angerer et al., *Methods Enzymol.* 152:649-660 (1987). In an in situ hybridization assay, cells or tissue, preferentially human cells or tissue from a selected brain region, are fixed to a solid support, typically a glass slide. If DNA is to be probed, the cells are denatured with heat or alkali. The cells are then contacted with a hybridization solution at a moderate temperature to permit annealing of specific probes that are labeled. The probes are preferably labeled with radioisotopes or fluorescent reporters.

V. Immunological Detection of the Polypeptides of the Invention

[0125] In addition to the detection of polynucleotide expression using nucleic acid hybridization technology, one can also use immunoassays to detect polypeptides of the invention. Immunoassays can be used to qualitatively or quantitatively analyze polypeptides. A general overview of the applicable technology can be found in Harlow & Lane, *Antibodies: A Laboratory Manual* (1988).

A. Antibodies to Target Polypeptides or Other Immunogens

[0126] Methods for producing therapeutic and diagnostic polyclonal and monoclonal antibodies that react specifically with a protein of interest or other immunogen are known to those of skill in the art (see, e.g., Coligan, supra; and Harlow and Lane, supra; Stites et al., supra and references cited therein; Goding, supra; and Kohler and Milstein *Nature*, 256: 495-497 (1975)). Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors (see, Huse et al., supra; and Ward et al., supra). For example, in order to produce antisera for use in an immunoassay, the protein of interest or an antigenic fragment thereof, is isolated as described herein. For example, a recombinant protein is produced in a transformed cell line. An inbred strain of mice or rabbits is immunized with the protein using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen.

[0127] Polyclonal sera are collected and titered against the immunogen in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid

support. Polyclonal antisera with a titer of 10^4 or greater are selected and tested for their cross-reactivity against unrelated proteins or even other homologous proteins from other organisms, using a competitive binding immunoassay. Specific monoclonal and polyclonal antibodies and antisera will usually bind with a K_D of at least about 0.1 mM, more usually at least about 1 μ M, preferably at least about 0.1 μ M or better, and most preferably, 0.01 μ M or better.

[0128] A number of proteins of the invention comprising immunogens may be used to produce antibodies specifically or selectively reactive with the proteins of interest. Recombinant protein is the preferred immunogen for the production of monoclonal or polyclonal antibodies. Naturally occurring proteins, such as one comprising an amino acid sequence encoded by a gene listed in FIGS. 1-14, may also be used either in pure or impure form. Synthetic peptides made using the protein sequences described herein may also be used as an immunogen for the production of antibodies to the protein. Recombinant protein can be expressed in eukaryotic or prokaryotic cells and purified as generally described supra. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated for subsequent use in immunoassays to measure the protein.

[0129] Methods of production of polyclonal antibodies are known to those of skill in the art. In brief, an immunogen, preferably a purified protein, is mixed with an adjuvant and animals are immunized. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to the polypeptide of interest. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired (see, Harlow and Lane, supra).

[0130] Monoclonal antibodies may be obtained using various techniques familiar to those of skill in the art. Typically, spleen cells from an animal immunized with a desired antigen are immortalized, commonly by fusion with a myeloma cell (see, Kohler and Milstein, *Eur. J. Immunol.* 6:511-519 (1976)). Alternative methods of immortalization include, e.g., transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse et al., supra.

[0131] Once target protein specific antibodies are available, the protein can be measured by a variety of immunoassay methods with qualitative and quantitative results available to the clinician. For a review of immunological and immunoassay procedures in general see, Stites, supra. Moreover, the immunoassays of the present invention can be performed in any of several configurations, which are reviewed extensively in Maggio *Enzyme Immunoassay*, CRC Press, Boca Raton, Fla. (1980); Tijssen, supra; and Harlow and Lane, supra.

[0132] Immunoassays to measure target proteins in a human sample may use a polyclonal antiserum that was raised to the protein (e.g., one has an amino acid sequence encoded

by a gene listed in FIG. 1-14) or a fragment thereof. This antiserum is selected to have low cross-reactivity against different proteins and any such cross-reactivity is removed by immunoabsorption prior to use in the immunoassay.

B. Immunological Binding Assays

[0133] In a preferred embodiment, a protein of interest is detected and/or quantified using any of a number of well-known immunological binding assays (see, e.g., U.S. Pat. Nos. 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, see also Asai *Methods in Cell Biology Volume 37: Antibodies in Cell Biology*, Academic Press, Inc. NY (1993); Stites, supra. Immunological binding assays (or immunoassays) typically utilize a "capture agent" to specifically bind to and often immobilize the analyte (in this case a polypeptide of the present invention or antigenic subsequences thereof). The capture agent is a moiety that specifically binds to the analyte. In a preferred embodiment, the capture agent is an antibody that specifically binds, for example, a polypeptide of the invention. The antibody may be produced by any of a number of means well known to those of skill in the art and as described above.

[0134] Immunoassays also often utilize a labeling agent to specifically bind to and label the binding complex formed by the capture agent and the analyte. The labeling agent may itself be one of the moieties comprising the antibody/analyte complex. Alternatively, the labeling agent may be a third moiety, such as another antibody, that specifically binds to the antibody/protein complex.

[0135] In a preferred embodiment, the labeling agent is a second antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second antibody can be modified with a detectable moiety, such as biotin, to which a third labeled molecule can specifically bind, such as enzyme-labeled streptavidin.

[0136] Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G, can also be used as the label agents. These proteins are normal constituents of the cell walls of streptococcal bacteria. They exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (see, generally, Kronval, et al. *J. Immunol.*, 111:1401-1406 (1973); and Akerstrom, et al. *J. Immunol.*, 135:2589-2542 (1985)).

[0137] Throughout the assays, incubation and/or washing steps may be required after each combination of reagents. Incubation steps can vary from about 5 seconds to several hours, preferably from about 5 minutes to about 24 hours. The incubation time will depend upon the assay format, analyte, volume of solution, concentrations, and the like. Usually, the assays will be carried out at ambient temperature, although they can be conducted over a range of temperatures, such as 10° C. to 40° C.

1. Non-Competitive Assay Formats

[0138] Immunoassays for detecting proteins of interest from tissue samples may be either competitive or noncompetitive. Noncompetitive immunoassays are assays in which the amount of captured analyte (in this case the protein) is directly measured. In one preferred "sandwich" assay, for example, the capture agent (e.g., antibodies specific for a

polypeptide encoded by a gene listed in FIG. 1-14) can be bound directly to a solid substrate where it is immobilized. These immobilized antibodies then capture the polypeptide present in the test sample. The polypeptide thus immobilized is then bound by a labeling agent, such as a second antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second can be modified with a detectable moiety, such as biotin, to which a third labeled molecule can specifically bind, such as enzyme-labeled streptavidin.

2. Competitive Assay Formats

[0139] In competitive assays, the amount of analyte (such as a polypeptide encoded by a gene listed in FIGS. 1-14) present in the sample is measured indirectly by measuring the amount of an added (exogenous) analyte displaced (or competed away) from a capture agent (e.g., an antibody specific for the analyte) by the analyte present in the sample. In one competitive assay, a known amount of, in this case, the protein of interest is added to the sample and the sample is then contacted with a capture agent, in this case an antibody that specifically binds to a polypeptide of the invention. The amount of immunogen bound to the antibody is inversely proportional to the concentration of immunogen present in the sample. In a particularly preferred embodiment, the antibody is immobilized on a solid substrate. For example, the amount of the polypeptide bound to the antibody may be determined either by measuring the amount of subject protein present in a protein/antibody complex or, alternatively, by measuring the amount of remaining uncomplexed protein. The amount of protein may be detected by providing a labeled protein molecule.

[0140] Immunoassays in the competitive binding format can be used for cross-reactivity determinations. For example, a protein of interest can be immobilized on a solid support. Proteins are added to the assay which compete with the binding of the antisera to the immobilized antigen. The ability of the above proteins to compete with the binding of the antisera to the immobilized protein is compared to that of the protein of interest. The percent cross-reactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% cross-reactivity with each of the proteins listed above are selected and pooled. The cross-reacting antibodies are optionally removed from the pooled antisera by immunosorption with the considered proteins, e.g., distantly related homologs.

[0141] The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein, thought to be perhaps a protein of the present invention, to the immunogen protein. In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required is less than 10 times the amount of the protein partially encoded by a sequence herein that is required, then the second protein is said to specifically bind to an antibody generated to an immunogen consisting of the target protein.

3. Other Assay Formats

[0142] In a particularly preferred embodiment, western blot (immunoblot) analysis is used to detect and quantify the

presence of a polypeptide of the invention in the sample. The technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support (such as, e.g., a nitrocellulose filter, a nylon filter, or a derivatized nylon filter) and incubating the sample with the antibodies that specifically bind the protein of interest. For example, the antibodies specifically bind to a polypeptide of interest on the solid support. These antibodies may be directly labeled or alternatively may be subsequently detected using labeled antibodies (e.g., labeled sheep anti-mouse antibodies) that specifically bind to the antibodies against the protein of interest.

[0143] Other assay formats include liposome immunoassays (LIA), which use liposomes designed to bind specific molecules (e.g., antibodies) and release encapsulated reagents or markers. The released chemicals are then detected according to standard techniques (see, Monroe et al. (1986) *Amer. Clin. Prod. Rev.* 5:34-41).

4. Labels

[0144] The particular label or detectable group used in the assay is not a critical aspect of the invention, as long as it does not significantly interfere with the specific binding of the antibody used in the assay. The detectable group can be any material having a detectable physical or chemical property. Such detectable labels have been well developed in the field of immunoassays and, in general, most labels useful in such methods can be applied to the present invention. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical or chemical means. Useful labels in the present invention include magnetic beads (e.g., Dynabeads™), fluorescent dyes (e.g., fluorescein isothiocyanate, Texas red, rhodamine, and the like), radiolabels (e.g., ³H, ¹²⁵I, ³⁵S, ¹⁴C, or ³²P), enzymes (e.g., horse radish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold or colored glass or plastic (e.g., polystyrene, polypropylene, latex, etc.) beads.

[0145] The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide variety of labels may be used, with the choice of label depending on the sensitivity required, the ease of conjugation with the compound, stability requirements, available instrumentation, and disposal provisions.

[0146] Non-radioactive labels are often attached by indirect means. The molecules can also be conjugated directly to signal generating compounds, e.g., by conjugation with an enzyme or fluorescent compound. A variety of enzymes and fluorescent compounds can be used with the methods of the present invention and are well-known to those of skill in the art (for a review of various labeling or signal producing systems which may be used, see, e.g., U.S. Pat. No. 4,391,904). Means of detecting labels are well known to those of skill in the art. Thus, for example, where the label is a radioactive label, means for detection include a scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence. The fluorescence may be detected visually, by means of photographic film, by the use of electronic detectors such as charge-coupled devices (CCDs) or photomultipliers and the like. Similarly, enzy-

matic labels may be detected by providing the appropriate substrates for the enzyme and detecting the resulting reaction product. Finally simple colorimetric labels may be detected directly by observing the color associated with the label. Thus, in various dipstick assays, conjugated gold often appears pink, while various conjugated beads appear the color of the bead.

[0147] Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies. In this case, antigen-coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need to be labeled and the presence of the target antibody is detected by simple visual inspection.

VI. Screening for Modulators of Polypeptides and Polynucleotides of the Invention

[0148] Modulators of polypeptides or polynucleotides of the invention, i.e. agonists or antagonists of their activity or modulators of polypeptide or polynucleotide expression, are useful for treating a number of human diseases, including mood disorders or psychotic disorders. Administration of agonists, antagonists or other agents that modulate expression of the polynucleotides or polypeptides of the invention can be used to treat patients with mood disorders or psychotic disorders.

A. Screening Methods

[0149] A number of different screening protocols can be utilized to identify agents that modulate the level of expression or activity of polypeptides and polynucleotides of the invention in cells, particularly mammalian cells, and especially human cells. In general terms, the screening methods involve screening a plurality of agents to identify an agent that modulates the polypeptide activity by binding to a polypeptide of the invention, modulating inhibitor binding to the polypeptide or activating expression of the polypeptide or polynucleotide, for example.

1. Binding Assays

[0150] Preliminary screens can be conducted by screening for agents capable of binding to a polypeptide of the invention, as at least some of the agents so identified are likely modulators of polypeptide activity. The binding assays usually involve contacting a polypeptide of the invention with one or more test agents and allowing sufficient time for the protein and test agents to form a binding complex. Any binding complexes formed can be detected using any of a number of established analytical techniques. Protein binding assays include, but are not limited to, methods that measure coprecipitation, co-migration on non-denaturing SDS-polyacrylamide gels, and co-migration on Western blots (see, e.g., Bennet and Yamamura, (1985) "Neurotransmitter, Hormone or Drug Receptor Binding Methods," in *Neurotransmitter Receptor Binding* (Yamamura, H. I., et al., eds.), pp. 61-89. The protein utilized in such assays can be naturally expressed, cloned or synthesized.

[0151] Binding assays are also useful, e.g., for identifying endogenous proteins that interact with a polypeptide of the

invention. For example, antibodies, receptors or other molecules that bind a polypeptide of the invention can be identified in binding assays.

2. Expression Assays

[0152] Certain screening methods involve screening for a compound that up or down-regulates the expression of a polypeptide or polynucleotide of the invention. Such methods generally involve conducting cell-based assays in which test compounds are contacted with one or more cells expressing a polypeptide or polynucleotide of the invention and then detecting an increase or decrease in expression (either transcript, translation product, or catalytic product). Some assays are performed with peripheral cells, or other cells, that express an endogenous polypeptide or polynucleotide of the invention.

[0153] Polypeptide or polynucleotide expression can be detected in a number of different ways. As described infra, the expression level of a polynucleotide of the invention in a cell can be determined by probing the mRNA expressed in a cell with a probe that specifically hybridizes with a transcript (or complementary nucleic acid derived therefrom) of a polynucleotide of the invention. Probing can be conducted by lysing the cells and conducting Northern blots or without lysing the cells using in situ-hybridization techniques. Alternatively, a polypeptide of the invention can be detected using immunological methods in which a cell lysate is probed with antibodies that specifically bind to a polypeptide of the invention.

[0154] Other cell-based assays are reporter assays conducted with cells that do not express a polypeptide or polynucleotide of the invention. Certain of these assays are conducted with a heterologous nucleic acid construct that includes a promoter of a polynucleotide of the invention that is operably linked to a reporter gene that encodes a detectable product. A number of different reporter genes can be utilized. Some reporters are inherently detectable. An example of such a reporter is green fluorescent protein that emits fluorescence that can be detected with a fluorescence detector. Other reporters generate a detectable product. Often such reporters are enzymes. Exemplary enzyme reporters include, but are not limited to, β -glucuronidase, chloramphenicol acetyl transferase (CAT); Alton and Vapnek (1979) *Nature* 282:864-869), luciferase, β -galactosidase, green fluorescent protein (GFP) and alkaline phosphatase (Toh, et al. (1980) *Eur. J. Biochem.* 182:231-238; and Hall et al. (1983) *J. Mol. Appl. Gen.* 2:101).

[0155] In these assays, cells harboring the reporter construct are contacted with a test compound. A test compound that either activates the promoter by binding to it or triggers a cascade that produces a molecule that activates the promoter causes expression of the detectable reporter. Certain other reporter assays are conducted with cells that harbor a heterologous construct that includes a transcriptional control element that activates expression of a polynucleotide of the invention and a reporter operably linked thereto. Here, too, an agent that binds to the transcriptional control element to activate expression of the reporter or that triggers the formation of an agent that binds to the transcriptional control element to activate reporter expression, can be identified by the generation of signal associated with reporter expression.

[0156] The level of expression or activity can be compared to a baseline value. As indicated above, the baseline value can be a value for a control sample or a statistical value that is

representative of expression levels for a control population (e.g., healthy individuals not having or at risk for mood disorders or psychotic disorders). Expression levels can also be determined for cells that do not express a polynucleotide of the invention as a negative control. Such cells generally are otherwise substantially genetically the same as the test cells.

[0157] A variety of different types of cells can be utilized in the reporter assays. Cells that express an endogenous polypeptide or polynucleotide of the invention include, e.g., brain cells, including cells from the cerebellum, anterior cingulate cortex, or dorsolateral prefrontal cortex. Cells that do not endogenously express polynucleotides of the invention can be prokaryotic, but are preferably eukaryotic. The eukaryotic cells can be any of the cells typically utilized in generating cells that harbor recombinant nucleic acid constructs. Exemplary eukaryotic cells include, but are not limited to, yeast, and various higher eukaryotic cells such as the COS, CHO and HeLa cell lines and stem cells, e.g., neural stem cells.

[0158] Various controls can be conducted to ensure that an observed activity is authentic including running parallel reactions with cells that lack the reporter construct or by not contacting a cell harboring the reporter construct with test compound. Compounds can also be further validated as described below.

3. Catalytic Activity

[0159] Catalytic activity of polypeptides of the invention can be determined by measuring the production of enzymatic products or by measuring the consumption of substrates. Activity refers to either the rate of catalysis or the ability to the polypeptide to bind (K_m) the substrate or release the catalytic product (K_d).

[0160] Analysis of the activity of polypeptides of the invention are performed according to general biochemical analyses. Such assays include cell-based assays as well as in vitro assays involving purified or partially purified polypeptides or crude cell lysates. The assays generally involve providing a known quantity of substrate and quantifying product as a function of time.

4. Validation

[0161] Agents that are initially identified by any of the foregoing screening methods can be further tested to validate the apparent activity. Preferably such studies are conducted with suitable animal models. The basic format of such methods involves administering a lead compound identified during an initial screen to an animal that serves as a model for humans and then determining if expression or activity of a polynucleotide or polypeptide of the invention is in fact upregulated. The animal models utilized in validation studies generally are mammals of any kind. Specific examples of suitable animals include, but are not limited to, primates, mice, and rats.

5. Animal Models

[0162] Animal models of mental disorders also find use in screening for modulators. In one embodiment, rat models of schizophrenia or other mental disorder, such as depression, are used for screening. In one embodiment, invertebrate models such as *Drosophila* models can be used, screening for modulators of *Drosophila* orthologs of the human genes disclosed herein. In another embodiment, transgenic animal

technology including gene knockout technology, for example as a result of homologous recombination with an appropriate gene targeting vector, or gene overexpression, will result in the absence, decreased or increased expression of a polynucleotide or polypeptide of the invention. The same technology can also be applied to make knockout cells. When desired, tissue-specific expression or knockout of a polynucleotide or polypeptide of the invention may be necessary. Transgenic animals generated by such methods find use as animal models of mental disorder and are useful in screening for modulators of mental disorder.

[0163] Knockout cells and transgenic mice can be made by insertion of a marker gene or other heterologous gene into an endogenous gene site in the mouse genome via homologous recombination. Such mice can also be made by substituting an endogenous polynucleotide of the invention with a mutated version of the polynucleotide, or by mutating an endogenous polynucleotide, e.g., by exposure to carcinogens.

[0164] For development of appropriate stem cells, a DNA construct is introduced into the nuclei of embryonic stem cells. Cells containing the newly engineered genetic lesion are injected into a host mouse embryo, which is re-implanted into a recipient female. Some of these embryos develop into chimeric mice that possess germ cells partially derived from the mutant cell line. Therefore, by breeding the chimeric mice it is possible to obtain a new line of mice containing the introduced genetic lesion (see, e.g., Capecchi et al., *Science* 244:1288 (1989)). Chimeric targeted mice can be derived according to Hogan et al., *Manipulating the Mouse Embryo: A Laboratory Manual*, Cold Spring Harbor Laboratory (1988) and *Teratocarcinomas and Embryonic Stem Cells: A Practical Approach*, Robertson, ed., IRL Press, Washington, D.C., (1987).

B. Modulators of Polypeptides or Polynucleotides of the Invention

[0165] The agents tested as modulators of the polypeptides or polynucleotides of the invention can be any small chemical compound, or a biological entity, such as a protein, sugar, nucleic acid or lipid. Alternatively, modulators can be genetically altered versions of a polypeptide or polynucleotide of the invention. Typically, test compounds will be small chemical molecules and peptides. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds that can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St. Louis, Mo.), Aldrich (St. Louis, Mo.), Sigma-Aldrich (St. Louis, Mo.), Fluka Chemika-Biochemica Analytika (Buchs, Switzerland) and the like. Modulators also include agents designed to reduce the level of mRNA of the invention (e.g. antisense molecules, ribozymes, DNazymes and the like) or the level of translation from an mRNA.

[0166] In one preferred embodiment, high throughput screening methods involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such "combinatorial chemical libraries" or "ligand libraries" are then screened in one or more assays, as

described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

[0167] A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

[0168] Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Pat. No. 5,010,175, Furka, *Int. J. Pept. Prot. Res.* 37:487-493 (1991) and Houghton et al., *Nature* 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (e.g., PCT Publication No. WO 91/19735), encoded peptides (e.g., PCT Publication WO 93/20242), random bio-oligomers (e.g., PCT Publication No. WO 92/00091), benzodiazepines (e.g., U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs et al., *Proc. Nat. Acad. Sci. USA* 90:6909-6913 (1993)), vinyllogous polypeptides (Hagihara et al., *J. Amer. Chem. Soc.* 114:6568 (1992)), nonpeptidic peptidomimetics with glucose scaffolding (Hirschmann et al., *J. Amer. Chem. Soc.* 114:9217-9218 (1992)), analogous organic syntheses of small compound libraries (Chen et al., *J. Amer. Chem. Soc.* 116:2661 (1994)), oligocarbamates (Cho et al., *Science* 261:1303 (1993)), and/or peptidyl phosphonates (Campbell et al., *J. Org. Chem.* 59:658 (1994)), nucleic acid libraries (see Ausubel, Berger and Sambrook, all supra), peptide nucleic acid libraries (see, e.g., U.S. Pat. No. 5,539,083), antibody libraries (see, e.g., Vaughn et al., *Nature Biotechnology*, 14(3):309-314 (1996) and PCT/US96/10287), carbohydrate libraries (see, e.g., Liang et al., *Science*, 274:1520-1522 (1996) and U.S. Pat. No. 5,593,853), small organic molecule libraries (see, e.g., benzodiazepines, Baum C&EN, January 18, page 33 (1993); isoprenoids, U.S. Pat. No. 5,569,588; thiazolidinones and metathiazanones, U.S. Pat. No. 5,549,974; pyrrolidines, U.S. Pat. Nos. 5,525,735 and 5,519,134; morpholino compounds, U.S. Pat. No. 5,506,337; benzodiazepines, U.S. Pat. No. 5,288,514, and the like).

[0169] Devices for the preparation of combinatorial libraries are commercially available (see, e.g., 357 MPS, 390 MPS, Advanced Chem Tech, Louisville Ky.; Symphony, Rainin, Woburn, Mass.; 433A Applied Biosystems, Foster City, Calif.; 9050 Plus, Millipore, Bedford, Mass.). In addition, numerous combinatorial libraries are themselves commercially available (see, e.g., ComGenex, Princeton, N.J.; Tripos, Inc., St. Louis, Mo.; 3D Pharmaceuticals, Exton, Pa.; Martek Biosciences, Columbia, Md., etc.).

C. Solid State and Soluble High Throughput Assays

[0170] In the high throughput assays of the invention, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each well of a microtiter plate can be used to run a separate assay against a

selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5-10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) modulators. If 1536 well plates are used, then a single plate can easily assay from about 100 to about 1500 different compounds. It is possible to assay several different plates per day; assay screens for up to about 6,000-20,000 different compounds are possible using the integrated systems of the invention. More recently, microfluidic approaches to reagent manipulation have been developed.

[0171] The molecule of interest can be bound to the solid state component, directly or indirectly, via covalent or non-covalent linkage, e.g., via a tag. The tag can be any of a variety of components. In general, a molecule that binds the tag (a tag binder) is fixed to a solid support, and the tagged molecule of interest is attached to the solid support by interaction of the tag and the tag binder.

[0172] A number of tags and tag binders can be used, based upon known molecular interactions well described in the literature. For example, where a tag has a natural binder, for example, biotin, protein A, or protein G, it can be used in conjunction with appropriate tag binders (avidin, streptavidin, neutravidin, the Fc region of an immunoglobulin, etc.). Antibodies to molecules with natural binders such as biotin are also widely available and appropriate tag binders (see, SIGMA Immunochemicals 1998 catalogue SIGMA, St. Louis Mo.).

[0173] Similarly, any haptenic or antigenic compound can be used in combination with an appropriate antibody to form a tag/tag binder pair. Thousands of specific antibodies are commercially available and many additional antibodies are described in the literature. For example, in one common configuration, the tag is a first antibody and the tag binder is a second antibody which recognizes the first antibody. In addition to antibody-antigen interactions, receptor-ligand interactions are also appropriate as tag and tag-binder pairs, such as agonists and antagonists of cell membrane receptors (e.g., cell receptor-ligand interactions such as transferrin, c-kit, viral receptor ligands, cytokine receptors, chemokine receptors, interleukin receptors, immunoglobulin receptors and antibodies, the cadherin family, the integrin family, the selectin family, and the like; see, e.g., Pigott & Power, *The Adhesion Molecule Facts Book I* (1993)). Similarly, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids, etc.), intracellular receptors (e.g., which mediate the effects of various small ligands, including steroids, thyroid hormone, retinoids and vitamin D; peptides), drugs, lectins, sugars, nucleic acids (both linear and cyclic polymer configurations), oligosaccharides, proteins, phospholipids and antibodies can all interact with various cell receptors.

[0174] Synthetic polymers, such as polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, and polyacetates can also form an appropriate tag or tag binder. Many other tag/tag binder pairs are also useful in assay systems described herein, as would be apparent to one of skill upon review of this disclosure.

[0175] Common linkers such as peptides, polyethers, and the like can also serve as tags, and include polypeptide sequences, such as poly-Gly sequences of between about 5 and 200 amino acids. Such flexible linkers are known to those of skill in the art. For example, poly(ethylene glycol) linkers are available from Shearwater Polymers, Inc., Huntsville,

Ala. These linkers optionally have amide linkages, sulfhydryl linkages, or hetero functional linkages.

[0176] Tag binders are fixed to solid substrates using any of a variety of methods currently available. Solid substrates are commonly derivatized or functionalized by exposing all or a portion of the substrate to a chemical reagent which fixes a chemical group to the surface which is reactive with a portion of the tag binder. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl groups. Aminoalkylsilanes and hydroxyalkylsilanes can be used to functionalize a variety of surfaces, such as glass surfaces. The construction of such solid phase biopolymer arrays is well described in the literature (see, e.g., Merrifield, *J. Am. Chem. Soc.* 85:2149-2154 (1963) (describing solid phase synthesis of, e.g., peptides); Geysen et al., *J. Immun. Meth.* 102:259-274 (1987) (describing synthesis of solid phase components on pins); Frank and Doring, *Tetrahedron* 44:60316040 (1988) (describing synthesis of various peptide sequences on cellulose disks); Fodor et al., *Science*, 251:767-777 (1991); Sheldon et al., *Clinical Chemistry* 39(4):718-719 (1993); and Kozal et al., *Nature Medicine* 2(7):753759 (1996) (all describing arrays of biopolymers fixed to solid substrates). Non-chemical approaches for fixing tag binders to substrates include other common methods, such as heat, cross-linking by UV radiation, and the like.

[0177] The invention provides in vitro assays for identifying, in a high throughput format, compounds that can modulate the expression or activity of the polynucleotides or polypeptides of the invention. In a preferred embodiment, the methods of the invention include such a control reaction. For each of the assay formats described, "no modulator" control reactions that do not include a modulator provide a background level of binding activity.

[0178] In some assays it will be desirable to have positive controls to ensure that the components of the assays are working properly. At least two types of positive controls are appropriate. First, a known activator of a polynucleotide or polypeptide of the invention can be incubated with one sample of the assay, and the resulting increase in signal resulting from an increased expression level or activity of polynucleotide or polypeptide determined according to the methods herein. Second, a known inhibitor of a polynucleotide or polypeptide of the invention can be added, and the resulting decrease in signal for the expression or activity can be similarly detected.

D. Computer-Based Assays

[0179] Yet another assay for compounds that modulate the activity of a polypeptide or polynucleotide of the invention involves computer assisted drug design, in which a computer system is used to generate a three-dimensional structure of the polypeptide or polynucleotide based on the structural information encoded by its amino acid or nucleotide sequence. The input sequence interacts directly and actively with a pre-established algorithm in a computer program to yield secondary, tertiary, and quaternary structural models of the molecule. Similar analyses can be performed on potential receptors or binding partners of the polypeptides or polynucleotides of the invention. The models of the protein or nucleotide structure are then examined to identify regions of the structure that have the ability to bind, e.g., a polypeptide or polynucleotide of the invention. These regions are then

used to identify polypeptides that bind to a polypeptide or polynucleotide of the invention.

[0180] The three-dimensional structural model of a protein is generated by entering protein amino acid sequences of at least 10 amino acid residues or corresponding nucleic acid sequences encoding a potential receptor into the computer system. The amino acid sequences encoded by the nucleic acid sequences provided herein represent the primary sequences or subsequences of the proteins, which encode the structural information of the proteins. At least 10 residues of an amino acid sequence (or a nucleotide sequence encoding 10 amino acids) are entered into the computer system from computer keyboards, computer readable substrates that include, but are not limited to, electronic storage media (e.g., magnetic diskettes, tapes, cartridges, and chips), optical media (e.g., CD ROM), information distributed by internet sites, and by RAM. The three-dimensional structural model of the protein is then generated by the interaction of the amino acid sequence and the computer system, using software known to those of skill in the art.

[0181] The amino acid sequence represents a primary structure that encodes the information necessary to form the secondary, tertiary, and quaternary structure of the protein of interest. The software looks at certain parameters encoded by the primary sequence to generate the structural model. These parameters are referred to as "energy terms," and primarily include electrostatic potentials, hydrophobic potentials, solvent accessible surfaces, and hydrogen bonding. Secondary energy terms include van der Waals potentials. Biological molecules form the structures that minimize the energy terms in a cumulative fashion. The computer program is therefore using these terms encoded by the primary structure or amino acid sequence to create the secondary structural model.

[0182] The tertiary structure of the protein encoded by the secondary structure is then formed on the basis of the energy terms of the secondary structure. The user at this point can enter additional variables such as whether the protein is membrane bound or soluble, its location in the body, and its cellular location, e.g., cytoplasmic, surface, or nuclear. These variables along with the energy terms of the secondary structure are used to form the model of the tertiary structure. In modeling the tertiary structure, the computer program matches hydrophobic faces of secondary structure with like, and hydrophilic faces of secondary structure with like.

[0183] Once the structure has been generated, potential ligand binding regions are identified by the computer system. Three-dimensional structures for potential ligands are generated by entering amino acid or nucleotide sequences or chemical formulas of compounds, as described above. The three-dimensional structure of the potential ligand is then compared to that of a polypeptide or polynucleotide of the invention to identify binding sites of the polypeptide or polynucleotide of the invention. Binding affinity between the protein and ligands is determined using energy terms to determine which ligands have an enhanced probability of binding to the protein.

[0184] Computer systems are also used to screen for mutations, polymorphic variants, alleles and interspecies homologs of genes encoding a polypeptide or polynucleotide of the invention. Such mutations can be associated with disease states or genetic traits and can be used for diagnosis. As described above, GeneChip™ and related technology can also be used to screen for mutations, polymorphic variants, alleles and interspecies homologs. Once the variants are iden-

tified, diagnostic assays can be used to identify patients having such mutated genes. Identification of the mutated a polypeptide or polynucleotide of the invention involves receiving input of a first amino acid sequence of a polypeptide of the invention (or of a first nucleic acid sequence encoding a polypeptide of the invention), e.g., any amino acid sequence having at least 60%, optionally at least 70% or 85%, identity with the amino acid sequence of interest, or conservatively modified versions thereof. The sequence is entered into the computer system as described above. The first nucleic acid or amino acid sequence is then compared to a second nucleic acid or amino acid sequence that has substantial identity to the first sequence. The second sequence is entered into the computer system in the manner described above. Once the first and second sequences are compared, nucleotide or amino acid differences between the sequences are identified. Such sequences can represent allelic differences in various polynucleotides, including SNPs and/or haplotypes, of the invention, and mutations associated with disease states and genetic traits.

VII. Compositions, Kits and Integrated Systems

[0185] The invention provides compositions, kits and integrated systems for practicing the assays described herein using polypeptides or polynucleotides of the invention, antibodies specific for polypeptides or polynucleotides of the invention, etc.

[0186] The invention provides assay compositions for use in solid phase assays; such compositions can include, for example, one or more polynucleotides or polypeptides of the invention immobilized on a solid support, and a labeling reagent. In each case, the assay compositions can also include additional reagents that are desirable for hybridization. Modulators of expression or activity of polynucleotides or polypeptides of the invention can also be included in the assay compositions.

[0187] The invention also provides kits for carrying out the therapeutic and diagnostic assays of the invention. The kits typically include a probe that comprises an antibody that specifically binds to polypeptides or polynucleotides of the invention, and a label for detecting the presence of the probe. The kits may include several polynucleotide sequences encoding polypeptides of the invention. Kits can include any of the compositions noted above, and optionally further include additional components such as instructions to practice a high-throughput method of assaying for an effect on expression of the genes encoding the polypeptides of the invention, or on activity of the polypeptides of the invention, one or more containers or compartments (e.g., to hold the probe, labels, or the like), a control modulator of the expression or activity of polypeptides of the invention, a robotic armature for mixing kit components or the like.

[0188] The invention also provides integrated systems for high-throughput screening of potential modulators for an effect on the expression or activity of the polypeptides of the invention. The systems typically include a robotic armature which transfers fluid from a source to a destination, a controller which controls the robotic armature, a label detector, a data storage unit which records label detection, and an assay component such as a microtiter dish comprising a well having a reaction mixture or a substrate comprising a fixed nucleic acid or immobilization moiety.

[0189] A number of robotic fluid transfer systems are available, or can easily be made from existing components. For

example, a Zymate XP (Zymark Corporation; Hopkinton, Mass.) automated robot using a Microlab 2200 (Hamilton; Reno, Nev.) pipetting station can be used to transfer parallel samples to 96 well microtiter plates to set up several parallel simultaneous STAT binding assays.

[0190] Optical images viewed (and, optionally, recorded) by a camera or other recording device (e.g., a photodiode and data storage device) are optionally further processed in any of the embodiments herein, e.g., by digitizing the image and storing and analyzing the image on a computer. A variety of commercially available peripheral equipment and software is available for digitizing, storing and analyzing a digitized video or digitized optical image, e.g., using PC, MACINTOSH®, or UNIX® based (e.g., SUN® work station) computers.

[0191] One conventional system carries light from the specimen field to a cooled charge-coupled device (CCD) camera, in common use in the art. A CCD camera includes an array of picture elements (pixels). The light from the specimen is imaged on the CCD. Particular pixels corresponding to regions of the specimen (e.g., individual hybridization sites on an array of biological polymers) are sampled to obtain light intensity readings for each position. Multiple pixels are processed in parallel to increase speed. The apparatus and methods of the invention are easily used for viewing any sample, e.g., by fluorescent or dark field microscopic techniques. Laser based systems can also be used.

VIII. Administration and Pharmaceutical Compositions

[0192] Modulators of the polynucleotides or polypeptides of the invention (e.g., antagonists or agonists) can be administered directly to a mammalian subject for modulation of activity of those molecules in vivo. Administration is by any of the routes normally used for introducing a modulator compound into ultimate contact with the tissue to be treated and is well known to those of skill in the art. Although more than one route can be used to administer a particular composition, a particular route can often provide a more immediate and more effective reaction than another route.

[0193] Diseases that can be treated include the following, which include the corresponding reference number from Morrison, *DSM-IV Made Easy*, 1995: Schizophrenia, Catatonic, Subchronic, (295.21); Schizophrenia, Catatonic, Chronic (295.22); Schizophrenia, Catatonic, Subchronic with Acute Exacerbation (295.23); Schizophrenia, Catatonic, Chronic with Acute Exacerbation (295.24); Schizophrenia, Catatonic, in Remission (295.55); Schizophrenia, Catatonic, Unspecified (295.20); Schizophrenia, Disorganized, Subchronic (295.11); Schizophrenia, Disorganized, Chronic (295.12); Schizophrenia, Disorganized, Subchronic with Acute Exacerbation (295.13); Schizophrenia, Disorganized, Chronic with Acute Exacerbation (295.14); Schizophrenia, Disorganized, in Remission (295.15); Schizophrenia, Disorganized, Unspecified (295.10); Schizophrenia, Paranoid, Subchronic (295.31); Schizophrenia, Paranoid, Chronic (295.32); Schizophrenia, Paranoid, Subchronic with Acute Exacerbation (295.33); Schizophrenia, Paranoid, Chronic with Acute Exacerbation (295.34); Schizophrenia, Paranoid, in Remission (295.35); Schizophrenia, Paranoid, Unspecified (295.30); Schizophrenia, Undifferentiated, Subchronic (295.91); Schizophrenia, Undifferentiated, Chronic (295.92); Schizophrenia, Undifferentiated, Subchronic with Acute Exacerbation (295.93); Schizophrenia, Undifferentiated,

Chronic with Acute Exacerbation (295.94); Schizophrenia, Undifferentiated, in Remission (295.95); Schizophrenia, Undifferentiated, Unspecified (295.90); Schizophrenia, Residual, Subchronic (295.61); Schizophrenia, Residual, Chronic (295.62); Schizophrenia, Residual, Subchronic with Acute Exacerbation (295.63); Schizophrenia, Residual, Chronic with Acute Exacerbation (295.94); Schizophrenia, Residual, in Remission (295.65); Schizophrenia, Residual, Unspecified (295.60); Delusional (Paranoid) Disorder (297.10); Brief Reactive Psychosis (298.80); Schizophreniform Disorder (295.40); Schizoaffective Disorder (295.70); Induced Psychotic Disorder (297.30); Psychotic Disorder NOS (Atypical Psychosis) (298.90); Personality Disorders, Paranoid (301.00); Personality Disorders, Schizoid (301.20); Personality Disorders, Schizotypal (301.22); Personality Disorders, Antisocial (301.70); Personality Disorders, Borderline (301.83) and bipolar disorders, maniac, hypomanic, dysthymic or cyclothymic disorders, substance-induced major depression, psychotic disorder, including schizophrenia (paranoid, catatonic, delusional) having schizoaffective disorder, and substance-induced psychotic disorder.

[0194] In some embodiments, modulators of polynucleotides or polypeptides of the invention can be combined with other drugs useful for treating mental disorders including psychotic disorders, e.g., schizophrenia; and mood disorders, e.g., bipolar disorders, or major depression. In some preferred embodiments, pharmaceutical compositions of the invention comprise a modulator of a polypeptide of polynucleotide of the invention combined with at least one of the compounds useful for treating schizophrenia, bipolar disorder, or major depression, e.g., such as those described in U.S. Pat. Nos. 6,297,262; 6,284,760; 6,284,771; 6,232,326; 6,187,752; 6,117,890; 6,239,162 or 6,166,008.

[0195] The pharmaceutical compositions of the invention may comprise a pharmaceutically acceptable carrier. Pharmaceutically acceptable carriers are determined in part by the particular composition being administered, as well as by the particular method used to administer the composition. Accordingly, there is a wide variety of suitable formulations of pharmaceutical compositions of the present invention (see, e.g., *Remington's Pharmaceutical Sciences*, 17th ed. 1985)).

[0196] The modulators (e.g., agonists or antagonists) of the expression or activity of the a polypeptide or polynucleotide of the invention, alone or in combination with other suitable components, can be made into aerosol formulations (i.e., they can be "nebulized") to be administered via inhalation or in compositions useful for injection. Aerosol formulations can be placed into pressurized acceptable propellants, such as dichlorodifluoromethane, propane, nitrogen, and the like.

[0197] Formulations suitable for administration include aqueous and non-aqueous solutions, isotonic sterile solutions, which can contain antioxidants, buffers, bacteriostats, and solutes that render the formulation isotonic, and aqueous and non-aqueous sterile suspensions that can include suspending agents, solubilizers, thickening agents, stabilizers, and preservatives. In the practice of this invention, compositions can be administered, for example, orally, nasally, topically, intravenously, intraperitoneally, or intrathecally. The formulations of compounds can be presented in unit-dose or multi-dose sealed containers, such as ampoules and vials. Solutions and suspensions can be prepared from sterile powders, granules, and tablets of the kind previously described. The modulators can also be administered as part of a prepared food or drug.

[0198] The dose administered to a patient, in the context of the present invention should be sufficient to effect a beneficial response in the subject over time. The optimal dose level for any patient will depend on a variety of factors including the efficacy of the specific modulator employed, the age, body weight, physical activity, and diet of the patient, on a possible combination with other drugs, and on the severity of the mental disorder. The size of the dose also will be determined by the existence, nature, and extent of any adverse side effects that accompany the administration of a particular compound or vector in a particular subject.

[0199] In determining the effective amount of the modulator to be administered a physician may evaluate circulating plasma levels of the modulator, modulator toxicity, and the production of anti-modulator antibodies. In general, the dose equivalent of a modulator is from about 1 ng/kg to 10 mg/kg for a typical subject.

[0200] For administration, modulators of the present invention can be administered at a rate determined by the LD-50 of the modulator, and the side effects of the modulator at various concentrations, as applied to the mass and overall health of the subject. Administration can be accomplished via single or divided doses.

IX. Gene Therapy Applications

[0201] A variety of human diseases can be treated by therapeutic approaches that involve stably introducing a gene into a human cell such that the gene is transcribed and the gene product is produced in the cell. Diseases amenable to treatment by this approach include inherited diseases, including those in which the defect is in a single or multiple genes. Gene therapy is also useful for treatment of acquired diseases and other conditions. For discussions on the application of gene therapy towards the treatment of genetic as well as acquired diseases, see, Miller, *Nature* 357:455-460 (1992); and Mulligan, *Science* 260:926-932 (1993).

[0202] In the context of the present invention, gene therapy can be used for treating a variety of disorders and/or diseases in which the polynucleotides and polypeptides of the invention has been implicated. For example, compounds, including polynucleotides, can be identified by the methods of the present invention as effective in treating a mental disorder. Introduction by gene therapy of these polynucleotides can then be used to treat, e.g., mental disorders including mood disorders or psychotic disorders (e.g., schizophrenia).

A. Vectors for Gene Delivery

[0203] For delivery to a cell or organism, the polynucleotides of the invention can be incorporated into a vector. Examples of vectors used for such purposes include expression plasmids capable of directing the expression of the nucleic acids in the target cell. In other instances, the vector is a viral vector system wherein the nucleic acids are incorporated into a viral genome that is capable of transfecting the target cell. In a preferred embodiment, the polynucleotides can be operably linked to expression and control sequences that can direct expression of the gene in the desired target host cells. Thus, one can achieve expression of the nucleic acid under appropriate conditions in the target cell.

B. Gene Delivery Systems

[0204] Viral vector systems useful in the expression of the nucleic acids include, for example, naturally occurring or

recombinant viral vector systems. Depending upon the particular application, suitable viral vectors include replication competent, replication deficient, and conditionally replicating viral vectors. For example, viral vectors can be derived from the genome of human or bovine adenoviruses, vaccinia virus, herpes virus, adeno-associated virus, minute virus of mice (MVM), HIV, sindbis virus, and retroviruses (including but not limited to Rous sarcoma virus), and MoMLV. Typically, the genes of interest are inserted into such vectors to allow packaging of the gene construct, typically with accompanying viral DNA, followed by infection of a sensitive host cell and expression of the gene of interest.

[0205] As used herein, “gene delivery system” refers to any means for the delivery of a nucleic acid of the invention to a target cell. In some embodiments of the invention, nucleic acids are conjugated to a cell receptor ligand for facilitated uptake (e.g., invagination of coated pits and internalization of the endosome) through an appropriate linking moiety, such as a DNA linking moiety (Wu et al., *J. Biol. Chem.* 263:14621-14624 (1988); WO 92/06180). For example, nucleic acids can be linked through a polylysine moiety to asialo-oromucoid, which is a ligand for the asialoglycoprotein receptor of hepatocytes.

[0206] Similarly, viral envelopes used for packaging gene constructs that include the nucleic acids of the invention can be modified by the addition of receptor ligands or antibodies specific for a receptor to permit receptor-mediated endocytosis into specific cells (see, e.g., WO 93/20221, WO 93/14188, and WO 94/06923). In some embodiments of the invention, the DNA constructs of the invention are linked to viral proteins, such as adenovirus particles, to facilitate endocytosis (Curiel et al., *Proc. Natl. Acad. Sci. U.S.A.* 88:8850-8854 (1991)). In other embodiments, molecular conjugates of the instant invention can include microtubule inhibitors (WO/9406922), synthetic peptides mimicking influenza virus hemagglutinin (Plank et al., *J. Biol. Chem.* 269:12918-12924 (1994)), and nuclear localization signals such as SV40 T antigen (WO93/19768).

[0207] Retroviral vectors are also useful for introducing the nucleic acids of the invention into target cells or organisms. Retroviral vectors are produced by genetically manipulating retroviruses. The viral genome of retroviruses is RNA. Upon infection, this genomic RNA is reverse transcribed into a DNA copy which is integrated into the chromosomal DNA of transduced cells with a high degree of stability and efficiency. The integrated DNA copy is referred to as a provirus and is inherited by daughter cells as is any other gene. The wild type retroviral genome and the proviral DNA have three genes: the gag, the pol and the env genes, which are flanked by two long terminal repeat (LTR) sequences. The gag gene encodes the internal structural (nucleocapsid) proteins; the pol gene encodes the RNA directed DNA polymerase (reverse transcriptase); and the env gene encodes viral envelope glycoproteins. The 5' and 3' LTRs serve to promote transcription and polyadenylation of virion RNAs. Adjacent to the 5' LTR are sequences necessary for reverse transcription of the genome (the tRNA primer binding site) and for efficient encapsulation of viral RNA into particles (the Psi site) (see, Mulligan, In: *Experimental Manipulation of Gene Expression*, Inouye (ed), 155-173 (1983); Mann et al., *Cell* 33:153-159 (1983); Cone and Mulligan, *Proceedings of the National Academy of Sciences, U.S.A.*, 81:6349-6353 (1984)).

[0208] The design of retroviral vectors is well known to those of ordinary skill in the art. In brief, if the sequences

necessary for encapsidation (or packaging of retroviral RNA into infectious virions) are missing from the viral genome, the result is a cis-acting defect which prevents encapsidation of genomic RNA. However, the resulting mutant is still capable of directing the synthesis of all virion proteins. Retroviral genomes from which these sequences have been deleted, as well as cell lines containing the mutant genome stably integrated into the chromosome are well known in the art and are used to construct retroviral vectors. Preparation of retroviral vectors and their uses are described in many publications including, e.g., European Patent Application EPA 0 178 220; U.S. Pat. No. 4,405,712, Gilboa *Biotechniques* 4:504-512 (1986); Mann et al., *Cell* 33:153-159 (1983); Cone and Mulligan *Proc. Natl. Acad. Sci. USA* 81:6349-6353 (1984); Eglitis et al. *Biotechniques* 6:608-614 (1988); Miller et al. *Biotechniques* 7:981-990 (1989); Miller (1992) supra; Mulligan (1993), supra; and WO 92/07943.

[0209] The retroviral vector particles are prepared by recombinantly inserting the desired nucleotide sequence into a retrovirus vector and packaging the vector with retroviral capsid proteins by use of a packaging cell line. The resultant retroviral vector particle is incapable of replication in the host cell but is capable of integrating into the host cell genome as a proviral sequence containing the desired nucleotide sequence. As a result, the patient is capable of producing, for example, a polypeptide or polynucleotide of the invention and thus restore the cells to a normal phenotype.

[0210] Packaging cell lines that are used to prepare the retroviral vector particles are typically recombinant mammalian tissue culture cell lines that produce the necessary viral structural proteins required for packaging, but which are incapable of producing infectious virions. The defective retroviral vectors that are used, on the other hand, lack these structural genes but encode the remaining proteins necessary for packaging. To prepare a packaging cell line, one can construct an infectious clone of a desired retrovirus in which the packaging site has been deleted. Cells comprising this construct will express all structural viral proteins, but the introduced DNA will be incapable of being packaged. Alternatively, packaging cell lines can be produced by transforming a cell line with one or more expression plasmids encoding the appropriate core and envelope proteins. In these cells, the gag, pol, and env genes can be derived from the same or different retroviruses.

[0211] A number of packaging cell lines suitable for the present invention are also available in the prior art. Examples of these cell lines include Crip, GPE86, PA317 and PG13 (see Miller et al., *J. Virol.* 65:2220-2224 (1991)). Examples of other packaging cell lines are described in Cone and Mulligan *Proceedings of the National Academy of Sciences, USA*, 81:6349-6353 (1984); Danos and Mulligan *Proceedings of the National Academy of Sciences, USA*, 85:6460-6464 (1988); Eglitis et al. (1988), supra; and Miller (1990), supra.

[0212] Packaging cell lines capable of producing retroviral vector particles with chimeric envelope proteins may be used. Alternatively, amphotropic or xenotropic envelope proteins, such as those produced by PA317 and GPX packaging cell lines may be used to package the retroviral vectors.

[0213] In some embodiments of the invention, an antisense polynucleotide is administered which hybridizes to a gene encoding a polypeptide of the invention. The antisense polypeptide can be provided as an antisense oligonucleotide (see, e.g., Murayama et al., *Antisense Nucleic Acid Drug Dev.* 7:109-114 (1997)). Genes encoding an antisense nucleic acid can also be provided; such genes can be introduced into cells

by methods known to those of skill in the art. For example, one can introduce an antisense nucleotide sequence in a viral vector, such as, for example, in hepatitis B virus (see, e.g., Ji et al., *J. Viral Hepat.* 4:167-173 (1997)), in adeno-associated virus (see, e.g., Xiao et al., *Brain Res.* 756:76-83 (1997)), or in other systems including, but not limited, to an HVJ (Sendai virus)-liposome gene delivery system (see, e.g., Kaneda et al., *Ann. NY Acad. Sci.* 811:299-308 (1997)), a "peptide vector" (see, e.g., Vidal et al., *CR Acad. Sci III* 32:279-287 (1997)), as a gene in an episomal or plasmid vector (see, e.g., Cooper et al., *Proc. Natl. Acad. Sci. U.S.A.* 94:6450-6455 (1997), Yew et al. *Hum Gene Ther.* 8:575-584 (1997)), as a gene in a peptide-DNA aggregate (see, e.g., Niidome et al., *J. Biol. Chem.* 272:15307-15312 (1997)), as "naked DNA" (see, e.g., U.S. Pat. Nos. 5,580,859 and 5,589,466), in lipidic vector systems (see, e.g., Lee et al., *Crit Rev Ther Drug Carrier Syst.* 14:173-206 (1997)), polymer coated liposomes (U.S. Pat. Nos. 5,213,804 and 5,013,556), cationic liposomes (Epand et al., U.S. Pat. Nos. 5,283,185; 5,578,475; 5,279,833; and 5,334,761), gas filled microspheres (U.S. Pat. No. 5,542,935), ligand-targeted encapsulated macromolecules (U.S. Pat. Nos. 5,108,921; 5,521,291; 5,554,386; and 5,166,320).

[0214] Upregulated transcripts listed in FIGS. 1-14 which are correlated with schizophrenia may be targeted with one or more short interfering RNA (siRNA) sequences that hybridize to specific sequences in the target, as described above. Targeting of certain brain transcripts with siRNA in vivo has been reported, for example, by Zhang et al., *J. Gene. Med.*, 12:1039-45 (2003), who utilized monoclonal antibodies against the transferrin receptor to facilitate passage of liposome-encapsulated siRNA molecules through the blood brain barrier. Targeted siRNAs represent useful therapeutic compounds for attenuating the over-expressed transcripts that are associated with disease states, e.g., schizophrenia.

[0215] In another embodiment, conditional expression systems, such as those typified by the tet-regulated systems and the RU-486 system, can be used (see, e.g., Gossen & Bujard, *PNAS* 89:5547 (1992); Oligino et al., *Gene Ther.* 5:491-496 (1998); Wang et al., *Gene Ther.* 4:432-441 (1997); Neering et al., *Blood* 88:1147-1155 (1996); and Rendahl et al., *Nat. Biotechnol.* 16:757-761 (1998)). These systems impart small molecule control on the expression of the target gene(s) of interest.

C. Pharmaceutical Formulations

[0216] When used for pharmaceutical purposes, the vectors used for gene therapy are formulated in a suitable buffer, which can be any pharmaceutically acceptable buffer, such as phosphate buffered saline or sodium phosphate/sodium sulfate, Tris buffer, glycine buffer, sterile water, and other buffers known to the ordinarily skilled artisan such as those described by Good et al. *Biochemistry* 5:467 (1966).

[0217] The compositions can additionally include a stabilizer, enhancer, or other pharmaceutically acceptable carriers or vehicles. A pharmaceutically acceptable carrier can contain a physiologically acceptable compound that acts, for example, to stabilize the nucleic acids of the invention and any associated vector. A physiologically acceptable compound can include, for example, carbohydrates, such as glucose, sucrose or dextrans; antioxidants, such as ascorbic acid or glutathione; chelating agents; low molecular weight proteins or other stabilizers or excipients. Other physiologically acceptable compounds include wetting agents, emulsifying agents, dispersing agents, or preservatives, which are particu-

larly useful for preventing the growth or action of microorganisms. Various preservatives are well known and include, for example, phenol and ascorbic acid. Examples of carriers, stabilizers, or adjuvants can be found in *Remington's Pharmaceutical Sciences*, Mack Publishing Company, Philadelphia, Pa., 17th ed. (1985).

D. Administration of Formulations

[0218] The formulations of the invention can be delivered to any tissue or organ using any delivery method known to the ordinarily skilled artisan. In some embodiments of the invention, the nucleic acids of the invention are formulated in mucosal, topical, and/or buccal formulations, particularly mucoadhesive gel and topical gel formulations. Exemplary permeation enhancing compositions, polymer matrices, and mucoadhesive gel preparations for transdermal delivery are disclosed in U.S. Pat. No. 5,346,701.

E. Methods of Treatment

[0219] The gene therapy formulations of the invention are typically administered to a cell. The cell can be provided as part of a tissue, such as an epithelial membrane, or as an isolated cell, such as in tissue culture. The cell can be provided in vivo, ex vivo, or in vitro.

[0220] The formulations can be introduced into the tissue of interest in vivo or ex vivo by a variety of methods. In some embodiments of the invention, the nucleic acids of the invention are introduced into cells by such methods as microinjection, calcium phosphate precipitation, liposome fusion, or biolistics. In further embodiments, the nucleic acids are taken up directly by the tissue of interest.

[0221] In some embodiments of the invention, the nucleic acids of the invention are administered ex vivo to cells or tissues explanted from a patient, then returned to the patient. Examples of ex vivo administration of therapeutic gene constructs include Nolte et al., *Proc Natl. Acad. Sci. USA* 93(6):2414-9 (1996); Koc et al., *Seminars in Oncology* 23 (1):46-65 (1996); Raper et al., *Annals of Surgery* 223(2):116-26 (1996); Dalesandro et al., *J. Thorac. Cardi. Surg.*, 11(2):416-22 (1996); and Makarov et al., *Proc. Natl. Acad. Sci. USA* 93(1):402-6 (1996).

X. Diagnosis of Mood and Psychotic Disorders

[0222] The present invention also provides methods of diagnosing mood disorders (such as major depression or bipolar disorder), psychotic disorders (such as schizophrenia). In one preferred embodiment, the disease state encompasses psychotic disorders. Diagnosis involves determining the level of a polypeptide or polynucleotide of the invention in a patient and then comparing the level to a baseline or range. Typically, the baseline value is representative of a polypeptide or polynucleotide of the invention in a healthy person not suffering from a mood disorder or psychotic disorder or under the effects of medication or other drugs. Variation of levels of a polypeptide or polynucleotide of the invention from the baseline range (either up or down) indicates that the patient has a mood disorder or psychotic disorder or at risk of developing at least some aspects of a mood disorder or psychotic disorder. In some embodiments, the level of a polypeptide or polynucleotide of the invention are measured by taking a blood, urine or tissue sample from a patient and measuring the amount of a polypeptide or polynucleotide of the invention in the sample using any number of detection methods, such as

those discussed herein, e.g., detection of expression levels or SNPs or haplotypes associated with these genes. The genes provided herein also can be used to develop probe sets for PCR and chip assays.

[0223] Single nucleotide polymorphism (SNP) analysis is also useful for detecting differences between alleles of the polynucleotides (e.g., genes) of the invention. SNPs linked to genes encoding polypeptides of the invention are useful, for instance, for diagnosis of diseases (e.g., mood disorders such as bipolar disease, major depression, and schizophrenia disorders) whose occurrence is linked to the gene sequences of the invention. For example, if an individual carries at least one SNP linked to a disease-associated allele of the gene sequences of the invention, the individual is likely predisposed for one or more of those diseases. If the individual is homozygous for a disease-linked SNP, the individual is particularly predisposed for occurrence of that disease. In some embodiments, the SNP associated with the gene sequences of the invention is located within 300,000; 200,000; 100,000; 75,000; 50,000; or 10,000 base pairs from the gene sequence.

[0224] Various real-time PCR methods can be used to detect SNPs, including, e.g., Taqman or molecular beacon-based assays (e.g., U.S. Pat. Nos. 5,210,015; 5,487,972; Tyagi et al., *Nature Biotechnology* 14:303 (1996); and PCT WO 95/13399) are useful to monitor for the presence or absence of a SNP. Additional SNP detection methods include, e.g., DNA sequencing, sequencing by hybridization, dot blotting, oligonucleotide array (DNA Chip) hybridization analysis, or are described in, e.g., U.S. Pat. No. 6,177,249; Landegren et al., *Genome Research*, 8:769-776 (1998); Botstein et al., *Am J Human Genetics* 32:314-331 (1980); Meyers et al., *Methods in Enzymology* 155:501-527 (1987); Keen et al., *Trends in Genetics* 7:5 (1991); Myers et al., *Science* 230:1242-1246 (1985); and Kwok et al., *Genomics* 23:138-144 (1994).

[0225] In some embodiments, the level of the enzymatic product of a polypeptide or polynucleotide of the invention is measured and compared to a baseline value of a healthy person or persons. Modulated levels of the product compared to the baseline indicates that the patient has a mood disorder or psychotic disorder or is at risk of developing at least some aspects of a mood disorder or psychotic disorder. Patient samples, for example, can be blood, PBS, lymphocytes, saliva, CSF, urine or tissue samples.

[0226] Immunoassays using antigens and antibodies for genes differentially expressed in psychotic disorders are also useful for immunoassays such as ELISA and immunohistochemical assays. The genes described herein are also useful for making differential diagnoses for psychiatric disorders.

[0227] In some embodiments, schizophrenia in a patient may be diagnosed or otherwise evaluated by visualizing expression in situ of one or more of the gene sequences in FIGS. 1-14. Those skilled in the art of visualizing the presence or expression of molecules including nucleic acids, polypeptides and other biochemicals in the brains of living patients will appreciate that the gene expression information described herein may be utilized in the context of a variety of visualization methods. Such methods include, but are not limited to, single-photon emission-computed tomography (SPECT) and positron-emitting tomography (PET) methods. See, e.g., Vassaux and Groot-wassink, "In Vivo Noninvasive Imaging for Gene Therapy," *J. Biomedicine and Biotechnology*, 2: 92-101 (2003); Turner, J., Smyth, P., Fallon, J. F., Kennedy, J. L., Potkin, S. G., *FIRST BIRN* (2006). Imaging and genetics in schizophrenia. *Neuroinformatics*, in press.

[0228] PET and SPECT imaging shows the chemical functioning of organs and tissues, while other imaging techniques—such as X-ray, CT and MRI—show structure. The use of PET and SPECT imaging is useful for qualifying and monitoring the development of brain diseases, including schizophrenia and related disorders. In some instances, the use of PET or SPECT imaging allows diseases to be detected years earlier than the onset of symptoms. The use of small molecules for labeling and visualizing the presence or expression of polypeptides and nucleotides has had success, for example, in visualizing proteins in the brains of Alzheimer's patients, as described by, e.g., Herholz K et al., *Mol Imaging Biol.*, 6(4):239-69 (2004); Nordberg A, *Lancet Neurol.*, 3(9): 519-27 (2004); *Neuropsychol Rev.*, Zakzanis K K et al., 13(1):1-18 (2003); Kung M P et al, *Brain Res.*, 1025(1-2):98-105 (2004); and Herholz K, *Ann Nucl Med.*, 17(2):79-89 (2003).

[0229] The dysregulated genes disclosed in FIGS. 1-14, or their encoded peptides (if any), or fragments thereof, can be used in the context of PET and SPECT imaging applications. After modification with appropriate tracer residues for PET or SPECT applications, molecules which interact or bind with the transcripts in FIGS. 1-14 or with any polypeptides encoded by those transcripts may be used to visualize the patterns of gene expression and facilitate diagnosis of schizophrenia as described herein. Similarly, if the encoded polypeptides encode enzymes, labeled molecules which interact with the products of catalysis by the enzyme may be used for the in vivo imaging and diagnostic application described herein.

[0230] Antisense technology is particularly suitable for detecting the transcripts identified in FIGS. 1-14 herein. For example, the use of antisense peptide nucleic acid (PNA) labeled with an appropriate radionuclide, such as ¹¹¹In, and conjugated to a brain drug-targeting system to enable transport across biologic membrane barriers, has been demonstrated to allow imaging of endogenous gene expression in brain cancer. See Suzuki et al., *Journal of Nuclear Medicine*, 10:1766-1775 (2004). Suzuki et al. utilize a delivery system comprising monoclonal antibodies that target transferring receptors at the blood-brain barrier and facilitate transport of the PNA across that barrier. Modified embodiments of this technique may be used to target upregulated genes associated with schizophrenia, such as the upregulated genes which appear in FIGS. 1-5, in methods of treating schizophrenic patients.

[0231] In other embodiments, the dysregulated genes listed in FIGS. 1-14 may be used in the context of prenatal and neonatal diagnostic methods. For example, fetal or neonatal lymphocytes can be isolated and the expression levels of appropriate transcripts (e.g., the transcripts in FIGS. 4-5) may be measured and correlated with the presence or increased likelihood of a mental disorder, e.g., schizophrenia. Similarly, the presence of one or more of the SNPs identified in FIG. 6 may be used to infer or corroborate dysregulated expression of ASG and the likelihood of schizophrenia in prenatal, neonatal, children and adult patients.

[0232] In other embodiments, the brain labeling and imaging techniques described herein or variants thereof may be used in conjunction with any of the dysregulated gene sequences in FIG. 1-4 or 6 in a forensic analysis, i.e., to determine whether deceased individual suffered from schizophrenia. Similarly, forensic examination of lymphocyte expression of any of the genes identified in FIGS. 4-5 may be

used alone or in conjunction with other methods to determine whether a deceased individual suffered from schizophrenia.

[0233] It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims.

EXAMPLES

Example 1

Identification of Genes Dysregulated in Schizophrenic Patients

[0234] Post mortem mental disorder brains (i.e., from schizophrenia patients) and control brains were used in this study. Each brain pair (case and control) was matched on the basis of gender, age, and postmortem interval. The patient's particular conditions in their terminal phase (agonal factors, e.g., seizure, coma, hypoxia, dehydration, and pyrexia) and the conditions of the brain tissue after death (postmortem factors, e.g., postmortem interval, and freezer interval) are two major influences on RNA preservation in postmortem brain tissue. Brain pH has been evaluated as an indicator for agonal status, and as an indicator of RNA preservation. Subjects with agonal factors and low pH samples, in which RNA quality was found to be compromised were eliminated from the study.

[0235] In this study, dysregulation of gene expression was studied in six brain regions: the anterior cingulate cortex (AnCg), dorsolateral prefrontal cortex (DLPFC), cerebellar cortex (CB), superior temporal gyms (STG), parietal cortex (PC), and nucleus accumbens (nAcc). Gene expression patterns in schizophrenic patients versus healthy controls were analyzed by using Affymetrix GeneChips {(HG_U133 set (A+B))} to interrogate the transcriptome from human post-mortem brains that met the strict quality control criteria referred to above, i.e., the agonal factor index of the donors was 0.0 (zero), and the pH of each individual brain was at least 6.4 (see Tomita et al., 2004; Li et al., 2004).

[0236] The probe set sequences used in data analysis are as defined in the CDF files created by the UMICH bioinformatics group (<http://brainarray.mhri.med.umich.edu/Brainarray/>) eliminating sequence redundancy that is inherent to UniGene definitions. The GeneChip hybridization data were processed, background-corrected, and normalized first with GCRMA using diagnosis (SZ, and control) and site {(University of California at Irvine (UCI), University of Michigan (UM) and University of California at Davis (UCD))} as factors in the analysis of sample replicates. The probe sets that showed (i) statistical significance of $p < 0.05$; (ii) an at least 1.2-fold change (FC) in expression in cases relative to controls in either direction; and (iii) Present (P) call of at least 10% in any given brain region out of the six analyzed (listed above) were selected as differentially expressed, i.e., dys-regulated genes.

[0237] A total of 1336 transcripts/genes were identified as differentially expressed in schizophrenic versus control brains in one or more of the six brain regions analyzed. Of those, 246 genes were upregulated in AnCg; 166 in CB; 156 in DLPFC; 124 in nAcc; 76 in PC; and 84 in STG. 138 genes were downregulated in AnCg; 187 in CB; 94 in DLPFC; 83 in nAcc; 125 in PC; and 105 in STG. This data is presented in FIG. 1.

[0238] Out of the six regions analyzed, only AnCg and DLPFC showed probe sets with Gene Ontology (GO) term enrichments and no significant enrichment of KEGG pathways. GO terms enriched in AnCg are listed in FIG. 2 shows that 3 GO terms were enriched in AnCg, specifically: GO:0050874, organismal physiological process (with 11 probe sets); GO:0058550, eukaryotic translation factor 2 complex (with 3 probe sets); and GO:0005739, mitochondrion (with 25 probe sets).

[0239] FIG. 3 shows that a single GO term was enriched in DLPFC, specifically: GO:0005622 intracellular (with 90 probe sets).

Example 2

Peripheral Biomarker Expression of Dysregulated Genes Found in Brain

[0240] For this study, a separate cohort of individuals with schizophrenia (n=5) were matched for gender and age to unaffected (n=5) members of a pedigree. Freshly isolated lymphocytes from each individual were transformed using the Epstein-Barr Virus and grown until confluent in RPMI-1640 media supplemented with 15% fetal bovine serum (heat-inactivated), 2 mM L-glutamine and 25 mg of gentamicin. RNA was extracted from $\sim 5 \times 10^7$ lymphoblastic cells using the standard TRIzol isolation protocol (Invitrogen, Carlsbad, Calif.). Affymetrix Human Genome U133A Arrays were used for gene expression according to the manufacturer's protocol. The gene expression traits were derived from the U133A chips and analyzed by robust multiarray condensation algorithm (RMA). Differential gene expression (gene expression trait for the purpose of this analysis) was defined as a gene that displayed a significant two-tailed t-test ($p < 0.05$) in schizophrenia versus unaffected family members. There were 1344 genes that passed the t-test for dysregulation in lymphocytes in schizophrenia compared to unaffecteds. This list was compared to genes in FIG. 1 (showing brain dysregulated genes in schizophrenia). The genes that were dysregulated in both brain and lymphocytes are shown in FIG. 4.

[0241] The list of 84 dysregulated genes in FIG. 4 may be grouped into those genes that agree in direction between brain and lymphoblasts, and those genes that disagree in direction between brain and lymphoblasts. Both dysregulated gene sets are biomarkers. The lymphoblasts do not have agonal factors, pH, or medication effects such as commonly seen in brain tissue. Thus, the gene transcripts in FIG. 4 may be used for monitoring lymphoblasts during treatment or for diagnostic purposes.

[0242] The subset of the 1344 genes identified by microarray analysis as significantly dysregulated in lymphoblasts only (i.e., not brains) is shown in FIG. 5. The microarray data was validated using Q-PCR to determine the fold change and direction of gene expression in the lymphocyte samples. Eight of these genes meet statistical significance in Q-PCR by t-test (two-tailed).

[0243] Aspartylglucosaminuria (AGA) gene expression is dysregulated in both the brain and lymphocytes of individuals with schizophrenia (FIG. 4). Eleven single nucleotide polymorphic markers were identified which correlate with AGA gene expression are shown in FIG. 6 also shows the regression p-values of genotype with lymphocyte gene expression. Of the 11 markers, 8 are associated with a cis-regulatory site (i.e., the Cis value is less than 5 Mb) and 3 are related to a trans-regulatory site (the Cis value is greater than 5 Mb). Detecting

these SNPs can facilitate the prediction of AGA gene expression in lymphoblasts. Similarly, detecting SNPs correlated with the expression of other dysregulated genes can facilitate the prediction of expression levels of those genes. The SNPs in FIG. 6 also represent targets for controlling expression of AGA, for diagnosing and treating schizophrenia, or for diagnosing and treating other disorders associated with altered AGA expression. For SNPs rs723820, rs723819, rs1112286, rs1375749, FIG. 6 shows that the minor alleles are associated with the decreased AGA expression in schizophrenia.

Example 3

Validation of PSPHL Insertion Deletion Mutation

[0244] The present invention extended the previous findings regarding the insertion-deletion polymorphism of phosphoserine phosphatase-like gene, and the association between deletion allele of PSPHL and susceptibility to bipolar disorder (BPD).

[0245] We previously determined 1) PSPHL gene consists of 4 exons. Exons 1, 2, 3 and 4 are 213 bp, 114 bp, 122 bp and 501 bp, in length, respectively, and span introns 1, 2 and 3 (3221 bp, 829 bp and 11939 bp, in length, respectively). 2) PSPHL and PSPH are highly homologous, which locate 200 kb apart from each other on chromosome 7p11.2 region. 3) PSPHL gene has two alternative transcripts, one of which utilizes the exons 1-4 (PSPHL-A), while another utilizes the exons 1, 2 and 4 (PSPHL-B). Predicted proteins of PSPHL-A and PSPHL-B share N-terminal 57 common amino acids, transcribed from exons 1 and 2. PSPH and the predicted PSPHL-A&B have 31 amino acids in common. 4) There locates an insertion/deletion polymorphism at the PSPHL locus. The deleted genomic region spans more than 30 kb, including the promoter region and the exons 1, 2 and 3 of PSPHL gene. 5) PSPHL shows a dichotomous (present or absent) pattern of expression among human population, which may due to the insertion/deletion polymorphism at the PSPHL locus. 6) Number of individuals who expresses PSPHL was significantly smaller in BPD patient group compared to control group. 7) Since PSPH is the rate limiting enzyme for serine synthesis, PSPHL may be involved in serine amino acid metabolic pathway, but might be involved in other pathways.

[0246] In the present invention, we further determined the followings:

[0247] 1) We verified involvement of the insertion-deletion polymorphism at the PSPHL locus on the expression pattern of the gene.

[0248] Among the 125 human postmortem brain tissues (19 BPD, 22 MDD, 20 SCZ patients and 64 controls) analyzed regarding the PSPHL genotype and PSPHL mRNA expression, 81 subjects (18 BPD, 8 MDD, 12 SCZ, and 43 Controls) showed homozygous pattern of the deletion allele (Del/Del) for the PSPHL locus, and all of the 81 Del/Del individuals lacked PSPHL mRNA expression. On the other hand, 40 subjects (1 BPD, 13 MDD, 8 SCZ, and 18 Controls) showed heterozygous pattern of the insertion and deletion alleles (Ins/Del) for the PSPHL locus, and 4 subjects (1 MDD and 3 Controls) showed homozygous pattern of the insertion allele (Ins/Ins). All of the 44 subjects (40 Ins/Del and 4 Ins/Ins), which have at least one insertion allele, showed PSPHL mRNA expression. This findings support that the presence/absence of PSPHL mRNA expression is due to the insertion/deletion polymorphism at the PSPHL locus. Our observations

on the 64 control subjects exactly matched Hardy Weinberg expectations. Allele frequencies for the insertion and deletion alleles for the PSPHL locus were estimated 0.18 and 0.82, respectively.

[0249] 2) We verified that number of individuals who expresses PSPHL was significantly smaller in BPD patient group compared to control group, and significantly larger in MDD patient group compared to the control group.

[0250] Based on hypergeometric distribution, the cumulative p value for observing 1 or less PSPHL non-expressed individuals in the 19 BPD patients is 0.0015. Also, the cumulative p value for observing 14 or more PSPHL-expressed individuals in the 22 MDD patients is 0.0002. There is no significant difference in the distribution between SCZ patients and controls. It is noteworthy that PSPHL non-expressed individuals are predominant in the BPD subjects, whereas PSPHL-expressed individuals are predominant in the MDD subjects. The probability of the observed difference in the distribution between BPD and MDD is 0.000098 based on the Fisher's exact test. These findings could be applicable for genetic testing to predict potential BPD patients among depressed patients who come to see physicians in their early stage of the chronic illnesses.

[0251] 3) We characterized that PSPHL-B mRNA expression level was also about 10 times higher than PSPHL-A in human postmortem brain tissue and cell lines derived from human brain.

[0252] PSPHL has at least two alternative transcripts; PSPHL-A (consists of the exons 1, 2, 3 and 4) and PSPHL-B (consists of the exons 1, 2 and 4). Based on quantitative RT-PCR evaluation with primer sets and TaqMan probes specific to PSPHL-A and PSPHL-B, both PSPHL-A and PSPHL-B were expressed in human postmortem brain cortices, including anterior cingulate and cerebellar cortices, from subjects which have at least one PSPHL insertion allele. PSPHL-B mRNA expression level was about 10 times higher than PSPHL-A in the brain tissues analyzed. Also, both PSPHL-A and PSPHL-B were expressed in cell lines derived from human brain, including human neuroblastoma cell lines, SK-N-SH, human glioma cell line, Hs 683, and human oligodendrocyte-derived cell line, OL, which have at least one PSPHL insertion allele. PSPHL-B mRNA expression level was also about 10 times higher than PSPHL-A in these cell lines. The glioblastoma cell line, U87-MG, which has homozygous PSPHL deletion allele, lack the expression of PSPHL-A and PSPHL-B.

[0253] 4) We verified promoter activity of 5' region of the PSPHL gene. The 5'-region of PSPHL (1015 bp fragment) was cloned into pGL-basic vector show sufficient promoter activity at least in the Hela cells and human oligodendrocyte cell line, OL. The vector contains the same region in opposite direction (negative control) did not show promoter activity.

[0254] The above examples are provided to illustrate the invention but not to limit its scope. Other variants of the invention will be readily apparent to one of ordinary skill in the art and are encompassed by the appended claims. All publications, databases, Genbank sequences, GO terms, patents, and patent applications cited herein are hereby incorporated by reference.

What is claimed is:

1. A method for determining whether a subject has or is predisposed for a mental disorder, the method comprising the steps of:

- (i) obtaining a biological sample from a subject;
 - (ii) contacting the sample with a reagent that selectively associates with a polynucleotide or polypeptide encoded by a nucleic acid that hybridizes under stringent conditions to a nucleotide sequence of Tables 1-14; and
 - (iii) detecting the level of reagent that selectively associates with the sample, thereby determining whether the subject has or is predisposed for a mental disorder.
- 2.** The method of claim **1**, wherein the reagent is an antibody.
- 3.** The method of claim **1**, wherein the reagent is a nucleic acid.
- 4.** The method of claim **1**, wherein the reagent associates with a polynucleotide.
- 5.** The method of claim **4**, wherein the reagent selectively associates with a polynucleotide comprising at least one of the SNPs listed in Table 6.
- 6.** The method of claim **1**, wherein the reagent associates with a polypeptide.
- 7.** The method of claim **1**, wherein the level of reagent that associates with the sample is different from a level associated with humans without a mental disorder.
- 8.** The method of claim **1**, wherein the biological sample is obtained from amniotic fluid.
- 9.** The method of claim **1**, wherein the mental disorder is a psychotic disorder or a mood disorder.
- 10.** The method of claim **1**, wherein the sample is derived from brain tissue or lymphocytes.
- 11.** The method of claim **10**, wherein the sample is derived from brain tissue.
- 12.** The method of claim **11**, wherein the sample is derived from a brain region selected from the group consisting of the anterior cingulate cortex (AnCg), dorsolateral prefrontal cortex (DLPFC), cerebellar cortex (CB), superior temporal gyrus (STG), parietal cortex (PC), nucleus accumbens (nAcc), and amygdala (amy).
- 13.** The method of claim **7**, wherein the level of reagent that associates with the sample is higher than a level associated with humans without a mental disorder.
- 14.** The method of claim **9**, wherein the psychotic disorder is schizophrenia.
- 15.** The method of claim **9**, wherein the mood disorder is major depression disorder or bipolar disorder.
- 16-46.** (canceled)

* * * * *