



US 20140135526A1

(19) **United States**

(12) **Patent Application Publication**
Lynch

(10) **Pub. No.: US 2014/0135526 A1**

(43) **Pub. Date: May 15, 2014**

(54) **COMPOSITIONS AND METHODS FOR
3-HYDROXYPROPIONATE
BIO-PRODUCTION FROM BIOMASS**

(60) Provisional application No. 60/992,290, filed on Dec. 4, 2007.

(71) Applicant: **OPX Biotechnologies, Inc.**, Boulder, CO (US)

Publication Classification

(72) Inventor: **Michael D. Lynch**, Boulder, CO (US)

(51) **Int. Cl.**
C12P 7/40 (2006.01)
C07C 57/04 (2006.01)

(73) Assignee: **OPX Biotechnologies, Inc.**, Boulder, CO (US)

(52) **U.S. Cl.**
CPC .. **C12P 7/40** (2013.01); **C07C 57/04** (2013.01)
USPC **562/598**; 435/136

(21) Appl. No.: **14/067,838**

(22) Filed: **Oct. 30, 2013**

(57) **ABSTRACT**

Related U.S. Application Data

(63) Continuation of application No. 13/284,337, filed on Oct. 28, 2011, now Pat. No. 8,652,816, which is a continuation of application No. 12/328,588, filed on Dec. 4, 2008, now Pat. No. 8,048,624.

Methods of obtaining mutant nucleic acid sequences that demonstrate elevated oxaloacetate a-decarboxylase activity are provided. Compositions, such as genetically modified microorganisms that comprise such mutant nucleic acid sequences, are described, as are methods to obtain the same.




FIG. 1




FIG 2




FIG. 3





FIG. 4A

FIG. 4B

NUMBER	GENE
1	aroA
2	aroB
3	aroC
4	aroD
5	aroE
6	aroF
7	aroK
8	aroL
9	aspC
10	entC
11	entD
12	entE
13	folA
14	folD
15	menD
16	menE
17	menF
18	pabA
19	pabB
20	pheA
21	purN
22	trpA
23	trpD
24	tyrA
25	tyrB
26	ubiA
27	ubiB
28	ubiC
29	ubiE
30	ubiG
31	ubiH
32	ubiX
33	ydiB

FIG. 4C-1






FIG. 4C-2





FIG 5A






FIG 6




FIG. 7A and 7B





FIG 8

FIG 9A






FIG 9B









	Aerobic Conditions	Anaerobic Conditions
E. coli NZN111	 <p>visible colonies</p>	 <p>no growth</p>
E. coli NZN111 + mmsA	 <p>visible colonies</p>	 <p>no growth</p>
E. coli NZN111 + mmsA + KGD	 <p>visible colonies</p>	 <p>no growth</p>
E. coli NZN111 + mmsA + mutant KGD pools	 <p>visible colonies</p>	 <p>positive clones grow</p>

FIG. 10A





	Minimal Media
E. coli AB354	 no growth
E. coli AB354 + gabT	 no growth
E. coli AB354 + gabT +kgd	 no growth
E. coli AB354 + gabT +mutant kgd pools	 positive clones grow

FIG. 10B




FIG. 11




FIG 12

**COMPOSITIONS AND METHODS FOR
3-HYDROXYPROPIONATE
BIO-PRODUCTION FROM BIOMASS**

RELATED APPLICATIONS

[0001] This application is a continuation application which claims priority under 35 USC 120 to U.S. patent application Ser. No. 13/284,337, filed Oct. 28, 2011, which is a continuation of Ser. No. 12/328,588, filed Dec. 4, 2008, and this application also claims priority under 35 USC 119 to U.S. Provisional Patent Application No. 60/992,290, filed Dec. 4, 2007. Both referenced patent applications are incorporated by reference in their respective entireties herewith.

REFERENCE TO A SEQUENCE LISTING

[0002] An electronically filed sequence listing is provided herewith which has been submitted in ASCII format and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Dec. 27, 2013, is named 34246-703.302_SL.txt and is 72,182 bytes in size.

FIELD OF THE INVENTION

[0003] The present invention relates to methods, systems and compositions, including genetically modified microorganisms, i.e., recombinant microorganisms, adapted to exhibit elevated oxaloacetate alpha-oxo decarboxylase activity (also referred to herein as oxaloacetate alpha-decarboxylase activity).

BACKGROUND OF THE INVENTION

[0004] 3-hydroxypropionate ("3-HP", CAS No. 503-66-2) has been identified as a highly attractive potential chemical feedstock for the production of many large market commodity chemicals that are currently derived from petroleum derivatives. For example, commodity products that can be readily produced using 3-HP include acrylic acid, 1,3-propanediol, methyl-acrylate, and acrylamide, as shown in FIG. 1. The sum value of these commodity chemicals is currently estimated to exceed several billions of dollars annually in the US. However, the current petrochemical manufacturing techniques for these commodities adverse impact the environment via the pollutants generated and the energy used in their production. Manufacture of these same commodities via the clean, cost-effective, production of 3-HP from biomass will simultaneously reduce toxic waste and substitute renewable feed stocks for non-renewable resources. In addition to the environmental benefits associated with bio-based production of 3-HP, if the production cost of the derived commodities is substantially reduced relative to petroleum-based production, this would make a biorefining industry not only environmentally beneficial but also a very attractive investment.

[0005] Previous attempts to produce 3-HP via biological pathways provide product titers which have been low and these processes have required the use of expensive, rich media. Both of these factors limit commercial feasibility and profitability. The use of rich media was necessary due to the toxicity of 3-HP when fermented with the more economical minimal media. For example, in wild type *E. coli*, metabolic activity is significantly inhibited at levels of 3-HP that are 5-10 times lower than the approximate 100 g/L titer needed for economic feasibility using the more economical minimal media. In fact, toxic effects have also been observed in rich media at product titers which are approximately two times

lower than desired titers for commercial feasibility (Refer to FIG. 2). Further, the fermentative pathways reported by other investigators have not addressed and resolved the toxicity mechanisms of 3-HP to the host organisms. Further to issues related to commodity chemical production, which largely relies on petroleum-based starting materials, there is an increasing need to reduce the domestic usage of petroleum and natural gas. The numerous motivating factors for this increasing need include, but are not limited to: pollutant reduction (such as greenhouse gases), environmental protection, and reducing the dependence on foreign oil. These issues not only impact fuel markets, but also the markets of numerous other products that are currently derived from oil. Biorefining promises the development of efficient biological processes allowing for the conversion of renewable sources of carbon and energy into large volume commodity chemicals.

[0006] A biosynthetic route to 3-HP as a platform chemical would be of benefit to the public, not only in terms of reduced dependence on petroleum, but also by a reduction in the amount of pollutants that are generated by current non-biosynthetic processes. Because 3-HP is not currently used as a building block for the aforementioned commodity chemicals, technical hurdles must be surmounted to achieve low cost biological routes to 3-HP. These hurdles include the development of a new organism that not only has a metabolic pathway enabling the production of 3-HP, but is also tolerant to the toxic effects of 3-HP thus enabling the sustained production of 3-HP at economically desired levels.

[0007] There are numerous motivating factors to reduce the domestic usage of petroleum. These factors include, but are not limited to: 1) the negative environmental impacts of petroleum refining such as production of greenhouse gases and the emission of a wide variety of pollutants; 2) the national security issues that are associated with the current dependence on foreign oil such as price instability and future availability; and 3) the long term economic concerns with the ever-increasing price of crude oil. These issues not only impact fuel markets, but also the multi-billion dollar commodity petro-chemical market

[0008] One potential method to alleviate these issues is the implementation of bioprocessing for the conversion of renewable feed stocks (e.g. agricultural wastes) to large volume commodity chemicals. It has been estimated that such bioprocesses already account for 5% of the 1.2 trillion dollar US chemical market. Furthermore, some experts are projecting that up to 50% of the total US chemical market will ultimately be generated through biological means.

[0009] While the attractiveness of such bioprocesses has been recognized for some time, recent advances in biological engineering, including several bio-refining success stories, have accelerated interest in the large scale production of chemicals through biological routes. However, many challenges still remain for the economical bio-production of commodity chemicals. These challenges include the need to convert biomass into usable feed stocks, the engineering of microbes to produce relevant chemicals at high titers and productivities, the improvement of the microbes' tolerance to the desired product, and the need to minimize the generation of byproducts that might affect downstream processes. Finally, the product must be economically competitive in the marketplace.

[0010] The contributions of bioprocessing are expected to grow in the future as existing biological methods become more efficient and as new bioprocesses are developed. A

recent analysis by the U.S. Department of Energy identified a list of the Top Value Added Chemicals from Biomass that are good candidates for biosynthetic production. Eight of the top value added chemicals were organic acids, including 3-hydroxypropionic acid (3-HP). As depicted in FIG. 1, 3-HP is considered to be a platform chemical, capable of yielding valuable derivative commodity chemicals including acrylic acid and acrylic acid polymers, acrylate esters, acrylate polymers (plastics), acrylamide, and 1,3-propanediol. Presently, these high value chemicals are produced from petroleum.

[0011] One method to efficiently generate 3-HP by a bioprocess approach would be the microbial biosynthesis of renewable biomass sugars to 3-HP. According to the DOE Report (Werpy, T.; Petersen, G. Volume 1: Results of Screening for Potential Candidates from Sugars and Synthetic Gas. Oak Ridge, Tenn., U.S. Department of Energy; 2004. Top Value Added Chemicals from Biomass), a number of factors will need to be addressed, including: identifying the appropriate biosynthetic pathway, improving the reactions to reduce other acid co-products, increasing microbial yields and productivities, reducing the unwanted salts, and scale-up and integration of the system. Additionally, as noted above, it is critical to engineer the microbial organism to be tolerant to the potential toxicity of the desired product at commercially significant concentrations.

[0012] The production of acrylic acid from 3-HP is of particular interest because of the high market value of acrylic acid and its numerous derivatives. In 2005, the estimated annual production capacity for acrylic acid was approximately 4.2 million metric tons, which places it among the top 25 organic chemical products. Also, this figure is increasing annually. The demand for acrylic acid may exceed \$2 billion by 2010. The primary application of acrylic acid is the synthesis of acrylic esters, such as methyl, butyl or ethyl acrylate. When polymerized, these acrylates are ingredients in numerous consumer products, such as paints, coatings, plastics, adhesives, dispersives and binders for paper, textiles and leather. Acrylates account for 55% of the world demand for acrylic acid products, with butyl acrylate and ethyl acrylate having the highest production volumes. The other key use of the acrylic acid is through polymerization to polyacrylic acid, which is used in hygiene products, detergents, and waste water treatment chemicals. Acrylic acid polymers can also be converted into super absorbent materials (which account for 32% of worldwide acrylic acid demand) or developed into replacement materials for phosphates in detergents. Both of these are fast growing applications for acrylic acid.

[0013] Today, acrylic acid is made in a two step catalytic oxidation of propylene (a petroleum product) to acrolein, and acrolein to acrylic acid, using a molybdenum/vanadium based catalyst, with optimized yields of approximately 90%. It should be noted that several commercial manufacturers of acrylic acid are exploring the use of propane instead of propylene. The use of propane is projected to be more environmentally friendly by reducing energy consumption during production. However, propane is petroleum based, and while its use is a step in the right direction from an energy consumption standpoint, it does not offer the benefits afforded by the bioprocessing route.

[0014] In addition to acrylic acid, acrylates, and acrylic acid polymers, another emerging high value derivative of 3-HP is 1,3-propanediol (1,3-PD). 1,3-PD has recently been used in carpet fiber production for carpets. Further applications of 1,3-PD are expected to include cosmetics, liquid

detergents, and anti-freeze. The market for 1,3-PD is expected to grow rapidly as it becomes more routinely used in commercial products.

[0015] Pursuing a cleaner, renewable carbon source route to commodity chemicals through 3-HP will require downstream optimization of the chemical reactions, depending on the desired end product. 3-HP production through bioprocesses directly, or through reaction routes to the high-value chemical derivatives of 3-HP will provide for large scale manufacture of acrylic acid, as well reduction of environmental pollution, the reduction in dependence on foreign oil, and the improvement in the domestic usage of clean methods of manufacturing. Furthermore, the products produced will be of the same quality but at a competitive cost and purity compared to the current petroleum based product.

[0016] Thus, notwithstanding various advances in the art, there remains a need for methods that identify and/or provide, and compositions directed to recombinant microorganisms that have improved 3-HP production capabilities, so that increased 3-HP titers are achieved.

BRIEF DESCRIPTION OF THE DRAWINGS

[0017] FIG. 1 depicts how biomass derived 3-HP can serve as a chemical feedstock to many major chemical commodities worth billions of dollars. (Adapted from Werpy et al. US Dept. Energy, 2004).

[0018] FIG. 2 depicts 3-hydroxypropionate toxicity in *E. coli* K12. The minimum concentration of 3-HP that is required to inhibit visible growth after 24 hrs in minimal media is shown for wild type *E. coli* K12 grown in both minimal media and rich media which contains more complex nutrients.

[0019] FIG. 3 is an overview of SCALES. (a) Genomic DNA is fragmented to several specific sizes and ligated into vectors creating libraries with defined insert sizes. (b) These libraries are individually transformed into the host cell line used for selections. (c) The pools of transformants are mixed and subjected to selection. Clones bearing inserts with increasing fitness in a given selection have a growth advantage. (d) Enriched plasmids are purified from the selected population, prepared for hybridization, and applied to a microarray. (e) The processed microarray signal is analyzed as a function of genomic sequence position. (f) A nonlinear multi-scale analysis decomposition gives signal not only as a function of position but as a function of scale or library size. (g) Data are visualized and analyzed as a function of genomic position and scale. (for the circular chromosome of *E. coli* shown, genomic position correlates to position around the circle and scale is represented by color. The height of the peak above the circle correlates to population frequency or fitness of a given scale at a given position.)

[0020] FIGS. 4A-4C-2 depict the SCALES data identifying the chorismate superpathway as a 3-HP target. (A) Fitness data for positions and scales conferring increased fitness of *E. coli* in the presence of 3-HP. Genomic position correlates to position around the circle and scale is represented by color (red=500-1000 bp, yellow=1000-2000 bp, green=4000-8000 bp, blue=8000-10000 bp). The height of the peak above the circle correlates to the fitness of a given scale at a given position. Peaks corresponding to genes involved in the chorismate superpathway are numbered. (B) List of genes in the chorismate superpathway identified in (A). (C) The fitness of each gene identified in (A) is color coded and identified in the chorismate superpathway.

[0021] FIG. 5A This figure depicts the natural metabolic pathways utilized by *E. coli* during bio-production which results in the natural products lactate, formate and acetate FIG. 5B. The proposed metabolic pathway to produce 3-HP as a bio-production product. Arrows represent enzymatic activities. The non natural enzymatic function to be evolved in this Phase I project is colored in red. Enzyme activities are as follows [i] glucokinase, [ii] phosphoglucose isomerase, [iii] 6-phosphofructose kinase, [iv] fructose bisphosphate aldolase, [v] triose-phosphate isomerase, [vi] glyceraldehyde 3-phosphate dehydrogenase, [vii] phosphoglycerate kinase, [viii] phosphoglycerate mutase, [ix] enolase, [x] pyruvate kinase, [xi] lactate dehydrogenase, [xii] pyruvate oxidase, [xiii] pyruvate-formate lyase, [xiv] phosphate acetyltransferase, [xv] acetate kinase, [xvi] phosphoenolpyruvate carboxykinase [xvii] the proposed oxaloacetate alpha-oxo decarboxylase, [xviii] 3-hydroxypropionate dehydrogenase and [xix] malonate semialdehyde dehydrogenase

[0022] FIG. 6 This figure depicts the chemical reaction performed by 2-oxo acid decarboxylases. R can be any group.

[0023] FIG. 7A This figure depicts the chemical reaction performed by alpha-ketoglutarate decarboxylase encoded by the *kgd* gene from *M. tuberculosis*. FIG. 7B depicts the proposed reaction performed by the newly evolved enzyme, oxaloacetate alpha-oxo-decarboxylase. The proposed enzyme will be encoded by the *oad-2* gene which will be evolved by mutation from the *kgd* gene.

[0024] FIG. 8 This figure depicts an overview of the methods to select a diverse library of 2-oxo acid decarboxylases for oxaloacetate alpha-oxo-decarboxylase activity. [i] A natural 2-oxo acid decarboxylase is mutated to create a variant library, [ii] this library is introduced into a microbial host that will not survive in a given environment without the presence of the product of the alpha-oxo-decarboxylase, malonate semialdehyde. [iii]. Positive mutants are identified by growth under selective conditions.

[0025] FIG. 9A depicts the proposed selection of the metabolism of *E. coli* strain NZN111 is shown in the left box. The *pflB* gene is disrupted blocking the formation of acetyl-coA in anaerobic conditions. The lack of acetyl-coA formation severely inhibits growth. The proposed additional enzymatic path to acetyl-coA is outlined in the right box. The characterized *mmsA* gene can supply acetyl-coA under anaerobic conditions if it is supplied with malonate semialdehyde by an oxaloacetate alpha-oxo decarboxylase. *Kgd* mutants with this activity will allow the strain to grow under anaerobic conditions.

[0026] FIG. 9B depicts the proposed selection of the relevant metabolism of *E. coli* strain AB354 is summarized in the left box. The *panD* gene is mutated blocking the synthesis of beta-alanine, an essential precursor for pantothenate (coA). The lack of pantothenate formation abolishes growth on minimal media. The proposed additional enzymatic path to beta-alanine is outlined in the right box. The characterized *R. norvegicus* beta-alanine aminotransferase gene (*gabT*) can supply beta-alanine if it is supplied with malonate semialdehyde as a substrate. An active oxaloacetate alpha-decarboxylase will supply this substrate and enable growth on minimal media. *Kgd* mutants with this activity will allow the strain to grow on minimal media.

[0027] FIG. 10A depicts the anticipated Selection Results of mutant colonies expressing the desired oxaloacetate alpha-oxo-decarboxylase will grow under anaerobic conditions when expressed in *E. coli* NZN111 expressing *mmsA*. No

growth will be observed under these conditions in the *E. coli* NZN111, *E. coli* NZN111+*mmsA* controls. Or in mutants not expressing the desired activity.

[0028] FIG. 10B depicts the anticipated Selection Results of mutant colonies expressing the desired oxaloacetate alpha-decarboxylase will grow on minimal media when expressed in *E. coli* AB354 expressing *gabT*. No growth will be observed under these conditions in the *E. coli* AB354, *E. coli* AB354+*gabT* controls, or in *kgd* mutants not expressing the desired activity.

[0029] FIG. 11 depicts the screening Protocol. Purified enzyme will be mixed in vitro with the appropriate substrate and reagents. A) The control reaction for the native alpha-ketoglutarate decarboxylase. B) Predicted results for the native alpha-ketoglutarate decarboxylase with oxaloacetate as a substrate. C) Predicted results for *kgd* mutants, both positive and negative, for oxaloacetate alpha-decarboxylase activity.

[0030] FIG. 12 Expression and Purification results of pKK223-Cterm-5xHis-*kgd* ('5xHis' disclosed as SEQ ID NO: 24). Lane 1=marker; lane 2=uninduced culture; lane 3=induced culture; lane 4=native lysate; lane 5=flowthrough; lane 6=first wash (wash 1); lane 7=last wash (wash 3); lane 8=first elution; lane 9=second elution, purified *kgd*; lane 10=pelleted cell debris. The arrow points to the band comprising purified alpha-ketoglutarate decarboxylase.

DETAILED DESCRIPTION OF EMBODIMENTS OF THE INVENTION

[0031] Generally the invention is directed to compositions and methods for production of target chemical compounds in an organism. Various aspects of the invention are directed to providing altered/modified proteins having different enzymatic activity/function as compared to the unaltered protein. Further aspects, of the invention are directed to recombinant organisms comprising altered/modified pathways which are enhanced for production of a target compound (e.g., 3-HP). In some embodiments, a recombinant organism of the invention is a microorganism or algae. In further embodiments, a recombinant organism is a bacterium (e.g., *E. coli*). In one aspect of the invention, an organism is modified to include one or more genes encoding a protein involved in biosynthesis to enhance production of a target chemical compound (e.g., 3-HP). In further embodiments, such one or more genes encode one or more proteins which enhance the capability of the organism to produce a target chemical compound in culture. In one embodiment, such a chemical compound is 3-HP. In yet a further embodiment, the organism comprises at least one recombinant gene resulting in pyruvate, oxaloacetate and acetyl-coA production without committed formate production.

[0032] In another embodiment, the recombinant organism comprises acetyl-coA that is produced via the intermediate malonate semialdehyde. In yet another embodiment, acetyl-coA is produced via the intermediate pyruvate through pyruvate synthase.

[0033] Another aspect of the invention is directed to a method for producing 3-HP comprising growing a recombinant organism of the invention, where the organism comprises an enzyme which converts oxaloacetate to malonate semialdehyde. In further embodiments, the recombinant organism is engineered to delete or substantially reduce activity of one or more genes, where the gene(s) include but are not limited to *pfkA*, *pfkB*, *ldhA*, *pta*, *poxB*, *pflB* or a combination

thereof. In yet a further embodiment, the recombinant organism is modified to enhance the activity (such as by increasing expression or improving the relevant functioning) of one or more enzymes including but not limited to *pck*, *mmsA*, *mmsB*, *oad-2*, homologs thereof, or any combination thereof.

[0034] In one embodiment, a method is provided for producing 3-HP comprising growing an organism under a condition which enhance said 3-HP production, wherein said condition is selected from acetyl-coA production via malonate semialdehyde, acetyl-coA production via pyruvate by pyruvate synthase, without committed production of formate, homologs thereof and any combination thereof.

[0035] In a further aspect of the invention a recombinant microorganism is provided capable of producing 3-HP at quantities greater than about 10, 15, 20, 30, 40, 50, 60, 65, 70, 75, 80, 85, 90, 95, 100, 105, 110, 115, 120, 125, 130, 135, 140, 145 or 150 g/L. In one embodiment, the recombinant organism is capable of producing 3-HP from about 30 to about 100 g/L of biomass/culture.

[0036] In a further aspect of the invention, a bio-production mixture is provided for producing 3HP, said mixture comprising a recombinant microorganism; one or more products selected from a group consisting tyrosine, phenylalanine, para-aminobenzoate, para-hydroxy-benzoate, 2,3,-dihydrobenzoate and shikimate.

[0037] In further embodiments, the mixture comprises a microorganism which is engineered to produce *pck*, *mmsA*, *mmsB*, *oad-2*, homologs thereof, or a combination thereof. In further embodiments, the microorganism does not produce enzymes selected from a group consisting of *pfkA*, *pfkB*, *ldhA*, *pta*, *poxB*, *pflB*, homologs thereof and a combination thereof. In various embodiments, the microorganism is *E. coli*.

[0038] In one aspect of the invention, an isolated polypeptide is provided possessing oxaloacetate alpha oxo-decarboxylase activity, converting oxaloacetate to malonate semialdehyde. Furthermore, a nucleic acid encoding the polypeptide is provided. In yet a further embodiments, a functional variant for the polypeptide or nucleic acid sequence is provided which is homologous to the reference polypeptide and/or nucleic acid and functions as an oxaloacetate alpha oxo-decarboxylase. In some embodiments such a functional variant has at least about 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96, 97, 98, or 99% identity with alpha-ketoglutarate decarboxylase.

[0039] Accordingly, in various aspects of the invention improved methods for biomass production of 3-HP at higher concentrations are disclosed. With this development, it is feasible to construct *E. coli* strains that are highly tolerant to 3-HP and that will maintain robust metabolic activity in the presence of higher concentrations of 3-HP. In various embodiments, metabolic pathways which support the bio-production of 3-HP are manipulated to increase 3-HP production. In some embodiments, such metabolic pathways do not rely upon or are not affected by metabolic processes that are themselves inhibited by 3-HP.

[0040] Utilizing processes for identification of a 3-HP insensitive bio-production pathway (infra, under "Metabolic Toxicity of 3-HP"), in various embodiments of the invention a bio-production pathway is characterized for the synthesis of 3-HP in *E. coli*. In a further embodiment, an altered 2-oxo acid decarboxylase is utilized in a bio-production pathway to produce 3-HP. In yet further embodiments, a bio-production pathway is utilized incorporating previously characterized

and sequenced enzymes that have been reported in the literature, as discussed below under "Previously Characterized Enzymes".

[0041] In various embodiments, a bio-production pathway (shown in FIG. 6) relies directly or indirectly on the metabolite oxaloacetate through the intermediate malonate semialdehyde. The desired enzymatic activity carries out the conversion of oxaloacetate to malonate semialdehyde. This can be accomplished via a decarboxylation reaction not previously reported by a particular enzyme. More specifically, the decarboxylation of 2-oxo acids, such as oxaloacetate, is accomplished by a well understood set of thiamine pyrophosphate dependant decarboxylases, including pyruvate decarboxylases and branched chain 2-oxo acid decarboxylases. A more recently characterized enzyme from *M. tuberculosis*, alpha ketoglutarate decarboxylase, coded by the *kgd* gene, possesses catalytic activity with a primary substrate very similar to oxaloacetate, decarboxylating the metabolite alpha-ketoglutarate to succinate semialdehyde. As described in greater detail below, an alpha-ketoglutarate decarboxylase from *M. tuberculosis* is modified into an oxaloacetate alpha-oxo-decarboxylase or a functional variant thereof. In various embodiments, any 2-oxo acid decarboxylase including but not limited to pyruvate decarboxylases from various sources or branched chain 2-oxo acid decarboxylases are modified into an oxaloacetate alpha-oxo decarboxylase or a functional variant thereof. In various embodiments, a "functional variant" is a protein encoded by a sequence having about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or 99 percent identity with the nucleic acid sequence encoding the modified/altered oxaloacetate alpha-oxo-decarboxylase. In further embodiments, sequence identity can be on the amino acid sequence level, where a functional variant has a sequence identity with the reference sequence of about 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, 96, 97, 98, or 99. For example, a functional variant can have sequence identity that is 90 percent, or 95 percent, but where the enzyme still functions as an oxaloacetate alpha-oxo-decarboxylase when expressed in an organism (e.g., microorganism, algae, plant), such as *E. coli*.

[0042] In other embodiments, a microorganism or algae is engineered to follow a preferred 3-HP bio-production pathway and also to enhance tolerance to 3-HP production at commercially viable levels. In some such embodiments, the microorganism is a bacterium, such as *E. coli*. Thus, as one example an *E. coli* strain is constructed and optimized for a desired pathway as discussed herein, wherein enhanced tolerance to 3-HP also is established so as to produce commercially viable titers of product. Accordingly, it is within the conception of the present invention that its teachings, methods and compositions may be combined with other teachings, methods and compositions more specifically directed to 3-HP tolerance improvement, including co-owned and/or licensed inventions.

Metabolic Toxicity of 3-HP

[0043] Severe growth inhibition has been observed for extracellular 3-HP levels as low as 10 g/L in minimal media (pH 7.0), which limits the economic feasibility of 3-HP production as a platform chemical. FIG. 2 demonstrates the toxic effects of 3-HP on *E. coli* when grown in minimal media. These toxic effects have been observed to be far greater when the strains are grown in minimal media as compared to growth in rich media (containing a mixture of all nutrients, amino acids and vitamins). However toxicity at levels below

required titers (100 g/L) are still observed in rich media. These data alone indicate that 3-HP may be exerting toxic effects by suppressing central metabolic pathways essential to amino acid metabolism.

Diagnosis of 3-HP Toxicity Mechanisms

[0044] To better understand the toxic effects of 3-HP on *E. coli*, a genome-wide technology is used (multi-Scale Analysis of Library Enrichments (SCALES)), such as disclosed in U.S. Patent Application Publication No. 20060084098, with related inventions described in U.S. Patent Application Publication Nos. 20080103060 and 200702185333 (the latter entitled “Enhanced Alcohol Tolerant Microorganism and Methods of Use Thereof,”) published Sep. 20, 2007), which are incorporated by reference herein in their entirety for their respective teachings of methods that provide important information which may be analyzed to make a discovery of previously unappreciated metabolic relationships. An overview of the SCALES approach as well as sample data are depicted in FIG. 3.

[0045] This genome-wide approach allows identification of numerous genetic changes that can reduce the toxic effects of 3-HP. The results of our studies (shown in FIG. 4) identified hundreds of genes and other genetic elements that when at increased copy confer varying levels of tolerance to the presence of 3-HP in *E. coli*. When applied alone, these genetic changes may allow for small increases in tolerance; but when applied together they allow for insight into the 3-HP toxicity mechanisms. By grouping genetic elements that confer tolerance by their metabolic roles key metabolic pathways that are inhibited by 3-HP were identified.

[0046] The data shown in FIG. 4 depict identification of the chorismate superpathway as a target of 3-HP toxicity. In some embodiments, toxicity is alleviated by several processes. For example, the addition of the downstream products of branches of the chorismate superpathway, tyrosine, phenylalanine, para-aminobenzoate (a tetrahydrofolate precursor), para-hydroxy-benzoate (a precursor of ubiquinone) and 2,3-dihydroxybenzoate (an enterobactin precursor) all alleviate toxicity to a degree.

A 3-HP bio-Production Pathway

[0047] The genetic modifications conferring a 3-HP tolerant phenotype can enhance a 3-HP bio-production process utilizing *E. coli*. In addition, the mechanisms identified indicate that several current pathways under consideration for the production of 3-HP may not be viable routes at high levels of production

[0048] In various embodiments, a bio-production pathway is utilized which uses one or more metabolic pathways not negatively affected by 3-HP. Therefore, in some embodiments one or more traditional fermentation pathways in *E. coli* as well as pathways involving amino acid intermediates that are currently being explored by others [9,10] are bypassed in order to enhance production. In certain embodiments, a pathway to produce 3-HP is that depicted in FIG. 5. Also, one or more gene deletions in *E. coli* are effectuated as well as the expression of several enzymatic functions new to *E. coli*. In some embodiments, the one or more gene deletions are selected genes including but not limited to gene(s) encoding pyruvate kinase (pfkA and pfkB), lactate dehydrogenase (ldhA), phosphate acetyltransferase (pta), pyruvate oxidase (poxB) and pyruvate-formate lyase (pflB) enzymes. In further embodiments, any of the one or more deletions in the preceding are combined with one or more enzyme modifications,

where the enzymes include but are not limited to phosphoenolpyruvate carboxykinase (pck), malonate semialdehyde dehydrogenase A (mmsA), malonate semialdehyde dehydrogenase B (mmsB) and oxaloacetate alpha-oxo-decarboxylase (oad-2) enzymes are expressed. It should be understood that the term “deletion” in this context does not necessarily require an entire gene deletion, but rather, a modification sufficient to knock out or effectively reduce function.

[0049] The enzymatic activity (oxaloacetate alpha-oxo-decarboxylase) utilized in the proposed pathway has not been reported in the known scientific literature. The enzyme oxaloacetate alpha-oxo-decarboxylase enhances 3-HP production.

[0050] In various embodiments, a pathway having features valuable for bio-production of organic acids in general and can be viewed as a metabolic starting point for numerous other products and in various different organisms (e.g., bacteria, yeast, algae). In various embodiments, such a pathway enhancer allows intracellular production of the key intermediate acetyl-coA without the committed production of the fermentative byproduct formate normally produced in microorganisms (e.g., *E. coli*) with acetyl-coA under fermentative conditions.

Previously Characterized Enzymes

[0051] In various embodiments, an engineered pathway of the invention comprises several genetic modifications to wild type microorganisms (e.g., *E. coli*), in addition to the expression of the oxaloacetate alpha-oxo decarboxylase. For example, one or more mutations in a microorganism (e.g., *E. coli*) can include but not limited to genes: pykA, pykF, ldhA, pflB, pta and poxB genes. Standard methodologies can be used to generate these gene deletions and such methods are routine in the art (See, for example, Sambrook and Russell, Molecular Cloning: A Laboratory Manual, Third Edition 2001 (volumes 1-3), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., hereinafter “Sambrook and Russell”).

[0052] In addition to these one or more genetic deletions, the following enzymatic activities can be expressed to enhance 3-HP production (e.g., in *E. coli*): phosphoenolpyruvate carboxykinase and malonate semialdehyde dehydrogenase. In a further embodiment, the mmsA gene is expressed (e.g., mmsA from *Rattus norvegicus* which has been shown to possess malonate semialdehyde dehydrogenase activity and converts malonate semialdehyde to acetyl-coA). In yet a further embodiment, the mmsB gene is expressed (e.g., mmsB gene from *Pseudomonas aeruginosa* which has been shown to have 3-hydroxypropionate dehydrogenase activity). In another further embodiment, a GDP dependant phosphoenolpyruvate carboxykinase is expressed (e.g., gene from *Alcaligenes eutrophus* which has been characterized with kinetics favoring the desired direction producing oxaloacetate). Any genes disclosed herein can be readily synthesized using standard methodologies.

2-Oxo Acid Decarboxylases.

[0053] Several 2-oxo decarboxylases (also referred to as 2-keto acid decarboxylases, alpha-oxo decarboxylases, or alpha-keto acid decarboxylases) with a broad substrate range have been previously characterized, including several pyruvate and branched chain 2-keto-acid decarboxylases. In various embodiments, enzymes from this class of decarboxylases are utilized. The reaction carried out by these enzymes is

depicted in FIG. 6. Of additional interest is that a convenient colorimetric method has been developed to assay this enzymatic activity by detection of the products of this enzyme class which are all aldehydes. In one embodiment, a previously characterized enzyme, alpha ketoglutarate decarboxylase, encoded by the *kgd* gene from *Mycobacterium tuberculosis* is used. The enzymatic reaction performed by this enzyme is depicted in FIG. 7A, which is very similar to the desired enzymatic activity, the decarboxylation of oxaloacetate to malonate semialdehyde depicted in FIG. 7B.

Altered Enzyme Activity

[0054] In one embodiment, clones comprising enhanced oxaloacetate alpha-oxo-decarboxylase activity are obtained by mutation of a gene encoding an enzyme having a similar catalytic activity, namely 2-oxo acid decarboxylases. For example, mutant libraries of a 2-oxo acid decarboxylase gene are constructed. Oxaloacetate alpha-oxo-decarboxylase activity is selected from a mutant library of a 2-oxo acid decarboxylase genes and in one embodiment from a mutant library of the *kgd* gene encoding an alpha-ketoglutarate decarboxylase. In further embodiments, mutant genes encoding enzymes that modulate or enhance the desired activity are identified.

Overview

[0055] To obtain the desired altered enzyme, a mutant library of a 2-oxo acid decarboxylase gene is constructed, which will be used for selections. In various embodiments, various 2-oxo acid decarboxylase genes are cloned into an appropriate expression system for *E. coli*. Several 2-keto acid decarboxylases with a broad substrate range have been previously characterized (Pohl, M., Sprenger, G. A., Muller, M., A new perspective on thiamine catalysis. *Current Opinion in Biotechnology*, 15(4), 335-342 (2004)). Of particular interest is an enzyme from *M. tuberculosis*, alpha-ketoglutarate decarboxylase, *kgd*, which has been purified and characterized (Tian, J Bryk, R. Itoh, M., Suematsu, M., and Carl Nathan, C. Variant tricarboxylic acid cycle in *Mycobacterium tuberculosis*: Identification of alpha-ketoglutarate decarboxylase. *PNAS*. Jul. 26, 2005 vol. 102(30): 10670-10677; Stephanopoulos, G., Challenges in engineering microbes for biofuels production. *Science*, 2007. 315(5813):801-804). Numerous 2-oxo acid decarboxylase genes are known in the art, including but limited to pyruvate decarboxylases from several sources, branched-chain 2-keto acid decarboxylases from various sources, benzylformate decarboxylases from various sources and phenylpyruvate decarboxylases from several sources (refer to www.metacyc.org for a more complete list). In one embodiment, the *kgd* gene, encoding and alpha-ketoglutarate decarboxylase from *M. tuberculosis* is cloned into an appropriate expression system for *E. coli*. Subsequently, this expression clone is mutated to create a library of mutant clones.

Cloning an 2-Oxo Acid Decarboxylase Gene

[0056] Cloning and expression of any 2 oxo-acid decarboxylase gene including but limited to the *kgd* gene is performed via gene synthesis supplied from a commercial supplier using standard or conventional techniques. Therefore, no culturing or manipulating of *M. tuberculosis* is required in the case of *kgd*. In addition, gene synthesis allows for codon optimization for a particular host. Once obtained using stan-

dard methodology, the gene is cloned into an expression system using standard techniques.

Construction of a 2-Oxo Acid Decarboxylase Gene Library

[0057] The plasmid containing the cloned 2-oxo acid decarboxylase gene, including but limited to the *kgd* gene is mutated by standard methods resulting in a large library of mutants. Generally, any of a number of well-known standard methods may be used (See, for example, chapters 1-19 of *Directed Evolution Library Creation Methods and Protocols*, F. H. Arnold & G. Georgiou, Eds., *Methods in Molecular Biology*, Vol. 231, Humana Press (2003)). The mutant sequences are introduced into a new host cell line, generating a final library for subsequent selection.

Selection of Altered Activity

[0058] A selection based approach such as described herein can result in the rapid identification of a t-oxo acid decarboxylase mutant with oxaloacetate alpha-oxo-decarboxylase activity. In one example, an available strain of *E. coli*, strain NZN111 is utilized as a host for the selection. This *E. coli* strain has deletions in both the *ldhA* and *pflB* genes resulting in severely limited growth (-10 hr doubling time) under anaerobic conditions (See right side of FIG. 5). This growth limitation is due in part to the inability to produce the necessary metabolite acetyl-coA under these conditions. (See FIG. 9A below.) A strain of *E. coli* NZN111 expressing *mmsA* (*E. coli* NZN111+*mmsA*) in addition to a mutant 2-oxo acid decarboxylase gene, including but limited to the *kgd* gene, having oxaloacetate alpha-oxo-decarboxylase activity is capable of producing the metabolite acetyl-coA from the metabolic intermediate malonate semialdehyde in media supplemented with tartrate (tartate can be used as a supplement and is readily converted to oxaloacetate in *E. coli*). This proposed strain has increased growth under anaerobic conditions when compared to both *E. coli* NZN111 and *E. coli* NZN111+*mmsA*, controls. For example, such a selection is depicted in FIG. 10A. In one embodiment, *E. coli* NZN111 is constructed to express an acetylating malonate semialdehyde dehydrogenase.

[0059] Similar to the 2-oxo acid decarboxylase gene, an acetylating malonate semialdehyde dehydrogenase gene, including but not limited to *mmsA*, from *Pseudomonas aeruginosa PAOJ*, is obtained via gene synthesis from the commercial provider. It is subsequently be cloned into an expression plasmid.

[0060] In another example, an available strain of *E. coli*, strain AB354 is utilized as a host for the selection. This *E. coli* strain has a mutation in the *panD* genes resulting in severely limited growth in minimal media conditions, without the supplementation of beta-alanine (See right side of FIG. 5). This growth limitation is due to the inability to produce beta-alanine under these conditions. (See FIG. 9B below.) A strain of *E. coli* AB354 expressing a beta alanine aminotransferase (*E. coli* AB354+beta alanine aminotransferase) in addition to a mutant 2-oxo acid decarboxylase gene, including but limited to the *kgd* gene, having oxaloacetate alpha-oxo-decarboxylase activity is capable of producing the metabolite beta-alanine from the metabolic intermediate malonate semialdehyde in minimal media. This proposed strain has a recovered ability to grow in minimal media with supplementation of beta-alanine. For example, such a selection is

depicted in FIG. 10B. In one embodiment, *E. coli* AB354 is constructed to express a beta-alanine pyruvate aminotransferase.

[0061] Similar to the 2-oxo acid decarboxylase gene, a beta-alanine pyruvate aminotransferase gene, including but not limited to PA0132 from *Pseudomonas aeruginosa* PAOJ, is obtained via gene synthesis from the commercial provider. It is subsequently be cloned into an expression plasmid.

Selection of Oxaloacetate Alpha-Oxo-Decarboxylase Activity

[0062] The mutant library of *kgd* genes is introduced into *E. coli* strain NZN111 expressing the *mmsA* gene. This population is grown under anaerobic conditions in media supplemented with oxaloacetate. Individual mutants expressing the desired oxaloacetate alpha-oxo-decarboxylase activity show increased growth rates compared to the control strains. These clones are isolated and the mutant protein they express subsequently screened for oxaloacetate alpha-oxo-decarboxylase activity as described above.

Colorimetric Confirmation of Decarboxylase Activity

[0063] A colorimetric approach is taken from current standard methodologies. This approach necessitates the expression and purification of the mutant enzymes and reaction with the purified enzyme, its cofactor (thiamin pyrophosphate) and the appropriate substrate. Protein expression and purification are performed with standard methodologies.

[0064] The above description of an approach using NZN111 is meant to be exemplary and not limiting. Its teachings may be applied to other microorganism systems to achieve the desired results. For example, and also not meant to be limiting, use of metabolic features of another *E. coli* strain, AB354, is explained in some of the examples below.

Examples Section

[0065] The following examples disclose specific methods for providing an *E. coli* cell with heterologous nucleic acid sequences that encode for enzymes or other polypeptides that confer increased tolerance to 3-HP. Where there is a method to achieve a certain result that is commonly practiced in two or more specific examples (or for other reasons), that method may be provided in a separate Common Methods section that follows the examples. Each such common method is incorporated by reference into the respective specific example that so refers to it. Also, where supplier information is not complete in a particular example, additional manufacturer information may be found in a separate Summary of Suppliers section that may also include product code, catalog number, or other information. This information is intended to be incorporated in respective specific examples that refer to such supplier and/or product. In the following examples, efforts have been made to ensure accuracy with respect to numbers used (e.g., amounts, temperatures, etc.), but some experimental error and deviation should be accounted for. Unless indicated otherwise, temperature is in degrees Celsius and pressure is at or near atmospheric pressure at approximately 5340 feet (1628 meters) above sea level. It is noted that work done at external analytical and synthetic facilities was not conducted at or near atmospheric pressure at approximately 5340 feet (1628 meters) above sea level. All reagents, unless otherwise indicated, were obtained commercially.

[0066] The meaning of abbreviations is as follows: “C” means Celsius or degrees Celsius, as is clear from its usage, “s” means second(s), “min” means minute(s), “h,” “hr,” or “hrs” means hour(s), “psi” means pounds per square inch, “nm” means nanometers, “d” means day(s), “μ.LL” or “uL” or “ul” means microliter(s), “mL” means milliliter(s), “L” means liter(s), “mm” means millimeter(s), “nm” means nanometers, “mM” means millimolar, “μ.LM” or “uM” means micromolar, “M” means molar, “mmol” means millimole(s), “fmol” or “uMol” means micromole(s), “g” means gram(s), “μ.Lg” or “ug” means microgram(s) and “ng” means nanogram(s), “PCR” means polymerase chain reaction, “OD” means optical density, “OD₆₀₀” means the optical density measured at a wavelength of 600 nm, “kDa” means kilodaltons, “g” means the gravitation constant, “bp” means base pair(s), “kbp” means kilobase pair(s), “% w/v” means weight/volume percent, % v/v” means volume/volume percent, “IPTG” means isopropyl-μ.L-D-thiogalactopyranoside, “RBS” means ribosome binding site, “rpm” means revolutions per minute, “HPLC” means high performance liquid chromatography, and “GC” means gas chromatography. Also, 10⁵ and the like are taken to mean 10⁵ and the like.

Example 1

Development of a Plasmid Comprising *Kgd*

[0067] The nucleic acid sequence for the alpha-ketoglutarate decarboxylase (*kgd*) from *M. tuberculosis* was codon optimized for *E. coli* according to a service from DNA 2.0 (Menlo Park, Calif. USA), a commercial DNA gene synthesis provider. The nucleic acid sequence was synthesized with an eight amino acid N-terminal tag to enable affinity based protein purification. This nucleic acid sequence incorporated an NcoI restriction site overlapping the gene start codon and was followed by a HindIII restriction site. In addition a Shine Delgarno sequence (i.e., a ribosomal binding site) was placed in front of the start codon preceded by an EcorI restriction site. This codon optimized *kgd* nucleic acid sequence construct (SEQ ID NO: 1), which is designed to encode for the native *kgd* protein (SEQ ID NO: 2) was synthesized by DNA 2.0 and then provided in a pJ206 vector backbone (SEQ ID NO: 3).

[0068] A circular plasmid based cloning vector termed pKK223-*kgd* for expression of the alpha-ketoglutarate decarboxylase in *E. coli* was constructed as follows. The *kgd* gene in the pJ206 vector was amplified via a polymerase chain reaction with the forward primer being TTTTTTGTATAC-CATGGATCGTAAATTTTCGTGATGATC (SEQ ID NO: 4) containing a NcoI site that incorporates the start methionine for the protein sequence, and the reverse primer being CCCG-GTGAGATCTAGATCCGAACGCTTTCGTC-CAAGATTTCTT (SEQ ID NO: 5) containing a XbaI site and a BglII site that replaces the stop codon of the *kgd* gene with an in-30 frame protein linker sequence SRS. Also, these primers effectively removed the eight amino acid N-terminal tag. The amplified *kgd* nucleic acid sequence was subjected to enzymatic restriction digestion with the enzymes NcoI and BglII obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer’s instructions. The digestion mixture was separated by agarose gel electrophoresis, and visualized under UV transillumination as described in Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the amplified *kgd* nucleic acid sequence was cut from the gel and the

DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions.

[0069] An *E. coli* cloning strain bearing pKK223-3 was grown by standard methodologies and plasmid DNA was prepared by a commercial miniprep column from Qiagen.

[0070] A new DNA vector was created by amplifying a pKK223-3 template by polymerase chain reaction with a forward primer being CGGATCTAGATCTCACCATCAC-CACCATTAGTCGACCTGCAGCCAAG (SEQ ID NO: 6) and a reverse primer being TGAGATCTAGATCCGTTAT-GTCCCATGGTTCTGTTTCCTGTGTG (SEQ ID NO: 7). The product was prepared by a commercial PCR-purification column from Qiagen. Both primers contain Xba1 restriction sites that allowed for the linear polymerase chain reaction product to be circularized after restriction digestion with Xba1 with enzymes obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions, and subsequent self-ligation. The new vector, named pKK223-ct-his (SEQ ID NO: 8), contained a multiple cloning region containing the a protein coding cassette under control of a IPTG-inducible promoter with an Nco1 site that incorporate the start methionine and with a Xba1 site and a Bg111 site that code for the in-frame protein sequence SRSHHHHH (SEQ ID NO: 9), a multi-histidine tag that allows for metal-affinity protein purification of the expressed protein.

[0071] To insert the gene of interest, *kgd*, this vector was prepared by restriction digestion with the enzymes Nco1 and Bg111 obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture was separated by agarose gel electrophoresis, and visualized under UV transillumination as described under Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the amplified *kgd* gene product was cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. Pieces of purified DNA corresponding to the amplified *kgd* gene product and the pKK223-cterm-5xhis vector backbone ('5xhis' disclosed as SEQ ID NO: 24) were ligated and the ligation product was transformed and electroporated according to manufacturer's instructions. The sequence of the resulting vector, termed pKK223-cterm-5xhis-*kgd* (SEQ ID NO: 10, and simply pKK223-*kgd* such as in the electronic sequence listing, '5xhis' disclosed as SEQ ID NO: 24), was confirmed by routine sequencing performed by the commercial service provided by Macrogen (USA). pKK223-cterm-5xhis-*kgd* ('5xhis' disclosed as SEQ ID NO: 24) confers resistance to beta-lactamase and contains the *kgd* gene of *M. tuberculosis* under control of a *ptac* promoter inducible in *E. coli* hosts by IPTG.

Example 2

Development of a Plasmid Comprising Mer (Partial Prophetic)

[0072] The nucleic acid sequence for the malonyl-coA reductase gene (*mer*) from *Chloroflexus auranticus* was codon optimized for *E. coli* according to a service from DNA 2.0 (Menlo Park, Calif. USA), a commercial DNA gene synthesis provider. Attached and extending beyond the ends of this codon optimized *mer* nucleic acid sequence (SEQ ID NO: 11) were an EcoRI restriction site before the start codon and a Hind111 restriction site. In addition a Shine Delgamo

sequence (i.e., a ribosomal binding site) was placed in front of the start codon preceded by an EcoRI restriction site. This gene construct was synthesized by DNA 2.0 and provided in a pJ206 vector backbone.

[0073] A circular plasmid based cloning vector termed pKK223-*mcr* for expression of the malonyl-CoA reductase in *E. coli* was constructed as follows. The *mer* gene in the pJ206 vector was amplified via a polymerase chain reaction with the forward primer being TCGTACCAACCATGGCCGG-TACGGGTCGTTTGGCTGGTAAAATTG (SEQ ID NO: 12) containing a Nco1 site that incorporates the start methionine for the protein sequence, and the reverse primer being CGGTGTGAGATCTAGATCCGACGG-TAATCGCACGACCGCGGT ID NO: 13) containing a Xba1 site and a Bg111 site that replaces the stop codon of the *mcr* gene with an in-frame protein linker sequence SRS. The amplified *mer* nucleic acid sequence was subjected to enzymatic restriction digestion with the enzymes Nco1 and Xba1 obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture was separated by agarose gel electrophoresis, and visualized under UV transillumination as described under Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the amplified *mer* nucleic acid sequence was cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions.

[0074] An *E. coli* cloning strain bearing pKK223-3 was grown by standard methodologies and plasmid DNA was prepared by a commercial miniprep column from Qiagen.

[0075] As described in Example 1 above, a new DNA vector was created by amplifying a pKK223-3 template by polymerase chain reaction with a forward primer being CGGATCTAGATCTCACCATCAC-CATTAGTCGACCTGCAGCCAAG (SEQ ID NO: 6) and a reverse primer being TGAGATCTAGATCCGTTATGTC-CCATGGTTCTGTTTCCTGTGTG (SEQ ID NO: 7). The product was prepared by a commercial PCR-purification column from Qiagen. Both primers contain Xba1 restriction sites that allowed for the linear polymerase chain reaction product to be circularized after restriction digestion with Xba1 and subsequent self-ligation with enzymes obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The vector, named pKK223-ct-his (SEQ ID NO: 8), contained a multiple cloning region containing the a protein coding cassette under control of a IPTG-inducible promoter with an Nco1 site that incorporates the start methionine and with a Xba1 site and a Bg111 site that codes for the in-frame protein sequence SRSHHHHH (SEQ ID NO: 9). The latter multi-histidine sequence allows for metal-affinity protein purification of the expressed protein.

[0076] To insert the gene of interest, *mer*, this vector was prepared by restriction digestion with the enzymes Nco1 and Xba1 obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture was separated by agarose gel electrophoresis, and visualized under UV transillumination as described under Subsection II of the Common Methods Section.

[0077] Pieces of purified DNA corresponding to the amplified codon optimized *mer* nucleic acid sequence and the pKK223-ct-his vector backbone were ligated and the ligation product was transformed and electroporated according to manufacturer's instructions. The sequence of the resulting

vector termed pKK223-mcr (SEQ ID NO: 14) is confirmed by routine sequencing performed by the commercial service provided by MacroGen(USA). pKK223-mcr confers resistance to beta-lactamase and contains the mer gene of *M. tuberculosis* under control of a ptac promoter inducible in *E. coli* hosts by IPTG.

Example 3

Development of a Plasmid Comprising a Beta Alanine-Pyruvate Aminotransferase Gene (Prophetic)

[0078] Introduction of a gene, such as the beta alanine pyruvate aminotransferase gene, into bacterial cells requires the addition of transcriptional (promoters) and translational (ribosome binding site) elements for controlled expression and production of proteins encoded by the gene. A nucleic acid sequence for a gene, whether obtained by gene synthesis or by amplification by polymerase chain reaction from genomic sources, can be ligated to nucleic acid sequences defining these transcriptional and translational elements. The present example discloses the addition of an *E. coli* minimal promoter and ribosome binding site properly oriented in the nucleic acid sequence before a gene of interest.

[0079] The beta alanine pyruvate aminotransferase gene from *Pseudomonas aeruginosa* PAO1 (locus_tag="PA0132") is amplified by polymerase chain reaction from a genomic DNA template with the forward primer being GGGTTTCCATGGACCAGCCGCTCAACGTGG (SEQ ID NO: 15) and the reverse primer being GGGTTTTTCAGGC-GATGCCGTTGAGCGCTTCGCC (SEQ ID NO: 16). The forward primer incorporates an NcoI restriction site at the start methionine codon of the gene and the reverse primer includes a stop codon for the gene. The amplified nucleic acid sequence is subjected to enzymatic restriction digestion with the restriction enzyme NcoI from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture is separated by agarose gel electrophoresis, and is visualized under UV transillumination as described under Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the restricted nucleic acid sequence is cut from the gel and the DNA is recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. An *E. coli* tpiA promoter and ribosome binding site is produced by polymerase chain reaction using a forward primer GGGAACGGCGGGGAAAAA-CAAACGTT (SEQ ID NO: 17) and a reverse primer GGTC-CATGGTAATTCTCCACGCTTATAAGC (SEQ ID NO: 18). Using genomic *E. Coli* K12 DNA as the template, a PCR reaction was conducted using these primers.

[0080] The forward primer is complimentary to the nucleic acid sequence upstream of the minimal tpiA promoter region (SEQ ID NO: 19), which is the minimal promoter sequence of the *E. coli* K12 tpi gene. The reverse primer is located just downstream of the minimal promoter region and includes an NcoI restriction site at the location of the start methionine and also includes a ribosome binding site. The PCR-amplified nucleic acid sequence is subjected to enzymatic restriction digestion with the restriction enzyme NcoI from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture is separated by agarose gel electrophoresis, and is visualized under UV transillumination as described in Subsection II of the Common

Methods Section. An agarose gel slice containing a DNA piece corresponding to the restricted nucleic acid sequence is cut from the gel and the DNA is recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The restricted, purified nucleic acid piece containing the transcriptional and translational elements is ligated to the recovered DNA containing the gene of interest. The ligation product is used as a template for a subsequent polymerase chain reaction using the forward primer GGGAACGGCGGGGAAAAA-CAAACGTT (SEQ ID NO: 20). Alternatively, any other forward primer may be use so long as it includes sufficient nucleic acid sequences upstream of the minimal tpiA promoter sequence (SEQ ID NO: 19). In the present specific example, the reverse primer is GGGTTTTTCAGGCATGCCGTTGAGCGCTTCGCC (SEQ ID NO: 21). The amplified nucleic acid product is separated by agarose gel electrophoresis, and is visualized under UV transillumination as described in Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the restricted nucleic acid sequence is cut from the gel and the DNA is recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions.

[0081] The resulting nucleic acid piece then is ligated into a suitable plasmid or other vector or transposon or other system, for example pSMART (Lucigen Corp, Middleton, Wis., USA), StrataClone (Stratagene, La Jolla, Calif., USA) or pCR2.1-TOPO TA (Invitrogen Corp, Carlsbad, Calif., USA) according to manufacturer's instructions. These methods also are described in the Subsection II of the Common Methods Section. Accordingly, the resulting nucleic acid piece can be restriction digested and purified and re-ligated into any other vector as is standard in the art. A similar method can be used to combine any gene with any transcriptional and translational elements with variation of restriction sites and primers.

[0082] The resulting nucleic acid is cloned using standard methodologies into the multiple cloning site of plasmid pBT-3, resulting in pBT-3-BAAT. This plasmid expresses the beta-alanine aminotransferase has a replicon compatible with pKK223 based vectors and confers chloramphenicol resistance.

Example 4

Development of a Plasmid Comprising an Acetylating Malonate Semialdehyde Dehydrogenase (Prophetic)

[0083] Introduction of a gene, such as an acetylating malonate semialdehyde dehydrogenase gene, into bacterial cells requires the addition of transcriptional (promoters) and translational (ribosome binding site) elements for controlled expression and production of proteins encoded by the gene. A nucleic acid sequence for a gene, whether obtained by gene synthesis or by amplification by polymerase chain reaction from genomic sources, can be ligated to nucleic acid sequences defining these transcriptional and translational elements. The present example discloses the addition of an *E. coli* minimal promoter and ribosome binding site properly oriented in the nucleic acid sequence before a gene of interest.

[0084] The acetylating malonate semialdehyde dehydrogenase gene, such as is readily available from several sources (e.g., <http://ca.expasy.org/cgi-bin/nicezyme.pl?1.2.1.18>) is amplified by polymerase chain reaction from a genomic DNA

template by standard PCR methodology. The forward primer incorporates an NcoI restriction site at the start methionine codon of the gene and the reverse primer includes a stop codon for the gene. The amplified nucleic acid sequence is subjected to enzymatic restriction digestion with the restriction enzyme NcoI from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture is separated by agarose gel electrophoresis, and is visualized under UV transillumination as described under Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the restricted nucleic acid sequence is cut from the gel and the DNA is recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions.

[0085] An *E. coli* *tpiA* promoter and ribosome binding site is produced by polymerase chain reaction using a forward primer GGGAACGGCGGGGAAAAACAAACGTT (SEQ ID NO: 17) and a reverse primer GGTCCATGGTAATTCTCACGCTTATAAGC (SEQ ID NO: 18). Using genomic *E. Coli* K12 DNA as the template, a PCR reaction was conducted using these primers. The forward primer is complementary to the nucleic acid sequence upstream of the minimal *tpiA* promoter region (SEQ ID NO: 19). The reverse primer is located just downstream of the minimal promoter region and includes an NcoI restriction site at the location of the start methionine and also includes a ribosome binding site. The PCR-amplified nucleic acid sequence is subjected to enzymatic restriction digestion with the restriction enzyme NcoI from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. The digestion mixture is separated by agarose gel electrophoresis, and is visualized under UV transillumination as described in Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the restricted nucleic acid sequence is cut from the gel and the DNA is recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions.

[0086] The restricted, purified nucleic acid piece containing the transcriptional and translational elements is ligated to the recovered DNA containing the gene of interest. The ligation product is used as a template for a subsequent polymerase chain reaction using the forward primer GGGAACGGCGGGGAAAAACAAACGTT (SEQ ID NO: 17). Alternatively, any other forward primer may be used so long as it includes sufficient nucleic acid sequences upstream of the minimal *tpiA* promoter sequence (SEQ ID NO: 19). In the present specific example, the reverse primer is GGGTTTTTCAGGCGATGCCGTTGAGCGCTTCGCC (SEQ ID NO: 21). The amplified nucleic acid product is separated by agarose gel electrophoresis, and is visualized under UV transillumination as described in Subsection II of the Common Methods Section. An agarose gel slice containing a DNA piece corresponding to the restricted nucleic acid sequence is cut from the gel and the DNA is recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions.

[0087] The resulting nucleic acid piece then is ligated into a suitable plasmid or other vector or transposon or other system, for example pSMART (Lucigen Corp, Middleton, Wis., USA), StrataClone (Stratagene, La Jolla, Calif., USA) or pCR2.1-TOPO TA (Invitrogen Corp, Carlsbad, Calif., USA) according to manufacturer's instructions. These methods also are described in the Subsection II of the Common

Methods Section. Accordingly, the resulting nucleic acid piece can be restriction digested and purified and re-ligated into any other vector as is standard in the art. A similar method can be used to combine any gene with any transcriptional and translational elements with variation of restriction sites and primers.

[0088] The resulting nucleic acid is cloned using standard methodologies into the multiple cloning site of plasmid pBT-3, resulting in pBT-3-mmsA. This plasmid expresses an acetylating malonate semialdehyde dehydrogenase has a replicon compatible with pKK223 based vectors and confers chloramphenicol resistance.

Example 5

Development of a Plasmid Comprising a Pyruvate Decarboxylase. Evolution of Pyruvate Decarboxylase Enzymes for the Enzymatic Conversion of Oxaloacetate to Malonate Semialdehyde (Prophetic)

[0089] Similarly to alpha-ketoglutarate dehydrogenase from *Mycobacterium tuberculosis*, the pyruvate decarboxylase from *Zymomonas mobilis* can be evolved to perform the conversion of oxaloacetate to malonate semialdehyde. The pyruvate decarboxylase enzyme is a thiamine diphosphate-dependent enzyme that decarboxylates 2-keto acids and has been shown to prefer short aliphatic substrates (Siegert P et al. (2005). Exchanging the substrate specificities of pyruvate decarboxylase from *Zymomonas mobilis* and benzoylformate decarboxylase from *Pseudomonas putida*. Protein Eng Des Sel 18, 345-357). Additionally, this enzyme does not require substrate activation by pyruvamide (Hoppner, T. C. & Doelle, H. W. (1983). Purification and kinetic characteristics of pyruvate decarboxylase and ethanol dehydrogenase from *Zymomonas mobilis* in relation to ethanol production. Eur J Appl Microbial Biotechnol 17, 152-157), and a structure of the protein characterized by x-ray crystallography shows the residues responsible for formation of the substrate and cofactor binding pockets (Dobritzsch D et al. (1998). High resolution crystal structure of pyruvate decarboxylase from *Zymomonas mobilis*. Implications for substrate activation in pyruvate decarboxylases. J Biol Chem 273, 20196-20204). Furthermore, alteration of the substrate specificity of this enzyme by specific amino acid changes have previously been reported (Siegert P et al. (2005). Exchanging the substrate specificities of pyruvate decarboxylase from *Zymomonas mobilis* and benzoylformate decarboxylase from *Pseudomonas putida*. Protein Eng Des Sel 18, 345-357). An example of a process for randomly mutating specific amino acid regions of this protein follows.

[0090] To evolve the binding pocket of the protein for performing the oxaloacetate to malonate semialdehyde conversion, specific regions of the nucleic acid sequence comprising regions of the protein's amino acid sequence will be mutated. Identification of specific amino acid regions within the protein that are involved in the binding pocket interactions is performed by examining the previously determined crystal structure and also by comparing the protein sequence of the *Zymomonas mobilis* pyruvate decarboxylase with pyruvate decarboxylase from other species showing strong sequence similarity. Using this information, the nucleotide sequence of the gene is examined in order to place restriction sites within the nucleotide sequence at the boundaries of the corresponding amino acid regions identified previously. From this nucle-

otide sequence, the *Zymomonas mobilis* pyruvate decarboxylase gene with these restriction sites is codon optimized for *E. coli* according to a service from DNA 2.0 (Menlo Park, Calif. USA), a commercial DNA gene synthesis provider (SEQ ID NO: 22). This gene construct is synthesized by DNA 2.0 and provided in a pJ206 vector backbone. Additionally, the protein sequence includes the addition of a hepta-histidine purification tag (SEQ ID NO: 25), which can be easily removed by restriction digestion of the plasmid with HindIII followed by self-ligation. The protein for which SEQ ID NO: 22 encodes is provided as SEQ ID NO: 23.

[0091] To specifically mutate amino acids in the pyruvate decarboxylase protein, the plasmid containing the codon-optimized sequence is cut at regions of interest via the incorporated restriction sites. Nucleotide sequences is synthesized or produced by polymerase chain reaction with oligonucleotides designed to incorporate specific or random changes at these regions of interest. These nucleotide sequences will incorporate restriction sites or overhanging ends complementary to the restriction sites used to cut the plasmid such that the new sequences are ligated into the plasmid to create the desired changes in the protein. These changes can be performed singly or multiply. If these changes are performed multiply, the resulting plasmids are transformed into a panD deleted *E. coli* strain and screened in a manner such as depicted in FIGS. 10A and 10B. Additionally, the protein produced by these changes may be assayed in a manner such as depicted in FIG. 11.

Example 6

Development of a Nucleic Acid Sequence Encoding a Protein Sequence Demonstrating Elevated Oxaloacetate Alpha-Decarboxylase Activity (Partial Prophetic)

[0092] Oxaloacetate alpha-decarboxylase activity is selected from a pool of alpha-ketoglutarate decarboxylase

and DNA sequencing (Macrogen, Korea). Expression and purification of his-tagged-kgd was performed as described in Subsection III of the Common Methods Section. SDS-PAGE results of expression and purification are shown in FIG. 12.

[0093] *E. coli* AB354 (Δ panD) was transformed with the vector controls, pKK223, pKK223-Cterm-5xhis ('5xHis' disclosed as SEQ ID NO: 24), as well as the test vectors pKK223-mcr and pKK223-Cterm-5xHis-kgd ('5xHis' disclosed as SEQ ID NO: 24), according to standard methods described below. Each of the strains were grown overnight in LB rich media supplemented with 200 mg/L ampicillin (according to standard protocols). Following overnight growth, cells twice were harvested by centrifugation and washed by resuspension in M9 minimal media (standard protocol), diluted 1:10,000 and plated on M9 minimal media plates with 0.05 g/L threonine, 0.1 g/L leucine, 0.067 g/L thiamine, with the additional appropriate supplements, where indicated at the following concentrations (10 g/L beta-alanine (Sigma Aldrich, St. Louis, Mo.), 1 mM Isopropyl β -D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 0.2 g/L putrescine (MPBiomedicals, Santa Ana, Calif.), 200 mg/L ampicillin (Research Products International Corp., Mt. Prospect, Ill.) After plating, agarose plates were incubated at 37 C overnight by standard methods. Table 1 depicts the results of these selection controls. A plus (+) indicates growth on a plate, minus (-) indicates no growth. These data confirm the absence of growth in the selection hosts. Putrescine is known to induce the expression of gamma-aminobutyrate transaminase in *E. coli*. This enzyme has been shown in some species including *Rattus norvegicus* to also have beta-alanine aminotransferase activity. The mer gene encoding the malonyl-coA reductase, has been shown to produce malonate semialdehyde. The lack of growth on the strain expressing malonyl-coA reductase in the presence of putrescine indicates the need for the co-expression of a beta-alanine aminotransferase in *E. coli* AB354 for the selection.

TABLE 1

Strain	Supplements						
	None	β -alanine	Amp	Amp + β -alanine	IPTG + Amp	AMP + Put	Amp + IPTG + Put
K12	+	+	-	-	-	-	-
AB354 (Δ panD)	-	+	-	-	-	-	-
AB354 (Δ panD) + pKK223	-	+	-	+	-	-	-
AB354 (Δ panD) + pKK223-mcr	-	+	-	+	-	-	-
AB354 (Δ panD) + pKK223-kgd	-	+	-	+	-	-	-

(kgd) mutants by selection in an *E. coli* AB354 host expressing a beta-alanine pyruvate aminotransferase. pKK223-cterm-5xhis-kgd ('5xhis' disclosed as SEQ ID NO: 24) encoding the kgd gene was constructed as described above. Confirmation of alpha-ketoglutarate decarboxylase protein expression and enzymatic activity with appropriate controls were as follows. *E. Coloni* 10GF' electrocompetent cells (Lucigen, Cat.#60061-1) were transformed with the pKK223-Cterm-5xHis-kgd ('5xHis' disclosed as SEQ ID NO: 24), plasmid containing sequence for 5xHIS-tagged kgd protein ('5xHIS' disclosed as SEQ ID NO: 24) behind a pTAC promoter. Transformants were confirmed using restriction digest

[0094] Mutant libraries of pKK223-cterm-5xhis-kgd ('5xhis' disclosed as SEQ ID NO: 24) were constructed as follows. Plasmid DNA of pKK223-cterm-5xhis-kgd ('5xhis' disclosed as SEQ ID NO: 24) was purified by standard methods and transformed in the mutator strain *E. coli* XL1-Red (Stratagene, La Jolla, Calif.) according to manufacturer's protocols. Cells were harvested according to manufacturer's protocols and mutated plasmid DNA purified by standard methods.

[0095] Mutant pKK223-cterm-5xhis-kgd ('5xhis' disclosed as SEQ ID NO: 24) DNA is used to transform an *E. coli* host, AB354+pBT-3-BAAT, described above. Greater than

10^5 transformants are collected from LB ampicillin (200 g/L), Chloramphenicol (40 g/L) agarose plates. Cells are washed in M9 minimal media, diluted 1:10,000 and plated on M9 minimal media plates with 0.05 g/L threonine, 0.1 g/L leucine,

[0096] 0.067 g/L thiamine, with 1 mM Isopropyl-D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 200 g/L ampicillin and 40 g/L chloramphenicol. Plates are incubated at 37 C for several days. Colonies that grow are individually collected as positives clones bearing oxaloacetate alpha-decarboxylase activity.

Example 7

Development of a Nucleic Acid Sequence Encoding a Protein Sequence Demonstrating Elevated Oxaloacetate Alpha-Decarboxylase Activity (Prophetic)

[0097] Oxaloacetate alpha-decarboxylase activity is selected from a pool of pyruvate decarboxylase (pdc) mutants by selection in an *E. coli* AB354 host expressing a beta-alanine pyruvate aminotransferase. pKK223-cterm-5xhis-pdc ('5xhis' disclosed as SEQ ID NO: 24) encoding the pdc gene is constructed as described above. Confirmation of pyruvate decarboxylase protein expression and enzymatic activity with appropriate controls are as follows. *E. Cloni* 10GF' electrocompetent cells (Lucigen, Cat.#60061-1) are transformed with the pKK223-Cterm-5xHis-pdc ('5xHis' disclosed as SEQ ID NO: 24), plasmid containing sequence for 5xHIS-tagged pdc protein ('5xHis' disclosed as SEQ ID NO: 24) behind a pTAC promoter. Transformants are confirmed using restriction digest and DNA sequencing (Macrogen, Korea). Expression and purification of his-tagged-pdc are performed as described in Subsection III of the Common Methods Section.

[0098] *E. coli* AB354 (panD) is transformed with the vector controls, pKK223, pKK223-Cterm-5xhis ('5xHis' disclosed as SEQ ID NO: 24), as well as the test vectors pKK223-mcr and pKK223-Cterm-5xHis-pdc ('5xHis' disclosed as SEQ ID NO: 24), according to standard methods described below. Each of the strains is grown overnight in LB rich media supplemented with 200 mg/L ampicillin (according to standard protocols). Following overnight growth, cells twice are harvested by centrifugation and washed by resuspension in M9 minimal media (standard protocol), diluted 1:10,000 and plated on M9 minimal media plates with 0.05 g/L threonine, 0.1 g/L leucine, 0.067 g/L thiamine, with the additional appropriate supplements, where indicated at the following concentrations (10 g/L beta-alanine (Sigma Aldrich, St. Louis, Mo.), 1 mM Isopropyl β -D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 0.2 g/L putrescine (MPBiomedicals, Santa Ana, Calif.), 200 mg/L ampicillin (Research Products International Corp., Mt. Prospect, Ill.) After plating, agarose plates were incubated at 37 C overnight by standard methods. Putrescine is known to induce the expression of gamma-aminobutyrate transaminase in *E. coli*. This enzyme has been shown in some species including *Rattus norvegicus* to also have beta-alanine aminotransferase activity. The mer gene encoding the malonyl-coA reductase, has been shown to produce malonate semialdehyde. The lack of growth on the strain expressing malonyl-coA reductase in the presence of putrescine indicates the need for the co-expression of a beta-alanine aminotransferase in *E. coli* AB354 for the selection.

[0099] Mutant libraries of pKK223-cterm-5xhis-pdc ('5xhis' disclosed as SEQ ID NO: 24) are constructed as follows. Plasmid DNA of pKK223-cterm-5xhis-pdc ('5xhis' disclosed as SEQ ID NO: 24) are purified by standard methods and transformed in the mutator strain *E. coli* XL1-Red (Stratagene, La Jolla, Calif.) according to manufacturer's protocols. Cells are harvested according to manufacturer's protocols and mutated plasmid DNA purified by standard methods.

[0100] Mutant pKK223-cterm-5xhis-pdc ('5xhis' disclosed as SEQ ID NO: 24) DNA is used to transform an *E. coli* host, AB354+pBT-3-BAAT, described above. Greater than 10^5 transformants are collected from LB ampicillin (200 g/L), Chloramphenicol (40 g/L) agarose plates. Cells are washed in M9 minimal media, diluted 1:10,000 and plated on M9 minimal media plates with 0.05 g/L threonine, 0.1 g/L leucine,

[0101] 0.067 g/L thiamine, with 1 mM Isopropyl β -D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 200 g/L ampicillin and 40 g/L chloramphenicol. Plates are incubated at 37 C for several days. Colonies that grow are individually collected as positives clones bearing oxaloacetate alpha-decarboxylase activity.

Example 8

Development of a Nucleic Acid Sequence Encoding a Protein Sequence Demonstrating Elevated Oxaloacetate Alpha-Decarboxylase Activity (Partial Prophetic)

[0102] Oxaloacetate alpha-decarboxylase activity is selected from a pool of alpha-ketoglutarate decarboxylase (kgd) mutants by selection in an *E. coli* NZN111 host expressing an acetylating malonate semialdehyde dehydrogenase. pKK223-cterm-5xhis-kgd ('5xhis' disclosed as SEQ ID NO: 24) encoding the kgd gene was constructed as described above. Confirmation of alpha-ketoglutarate decarboxylase protein expression and enzymatic activity with appropriate controls were as follows. *E. Cloni* 10GF' electrocompetent cells (Lucigen, Cat.#60061-1) were transformed with the pKK223-Cterm-5xHis-kgd ('5xHis' disclosed as SEQ ID NO: 24), plasmid containing sequence for 5xHIS-tagged kgd protein ('5xHIS' disclosed as SEQ ID NO: 24) behind a pTAC promoter. Transformants were confirmed using restriction digest and DNA sequencing (Macrogen, Korea). Expression and purification of his-tagged-kgd were performed as described in Subsection III of the Common Methods Section.

[0103] *E. coli* NZN111 is transformed with the vector controls, pKK223, pKK223-Cterm-5xhis ('5xHis' disclosed as SEQ ID NO: 24), as well as the test vectors pKK223-mcr and pKK223-Cterm-5xHis-kgd ('5xHis' disclosed as SEQ ID NO: 24), according to standard methods described below. Each of the strains is grown overnight in LB rich media supplemented with 200 mg/L ampicillin (according to standard protocols). Following overnight growth, cells twice are harvested by centrifugation and washed by resuspension in LB media (standard protocol), diluted 1:10,000 and plated on LB media plates with the additional appropriate supplements, where indicated at the following concentrations 1 mM Isopropyl β -D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 200 mg/L ampicillin (Research Products International Corp., Mt. Prospect, Ill.) After plating, agarose plates are incubated at 37 C overnight anaerobically in BD type A Bio-Bags according to manufacturer's instruc-

tions (BD Biosciences, Franklin Lakes, N.J., Catalog #261214). The *mer* gene encoding the malonyl-coA reductase, has been shown to produce malonate semialdehyde. The presence of growth of the strain expressing malonyl-coA reductase in the presence of the co expressed acetylating malonate semialdehyde dehydrogenase in *E. coli* NZN111 serves as a positive control for the selection.

[0104] Mutant libraries of pKK223-cterm-5×his-kgd ('5×his' disclosed as SEQ ID NO: 24) were constructed as follows. Plasmid DNA of pKK223-cterm-5×his-kgd ('5×his' disclosed as SEQ ID NO: 24) were purified by standard methods and transformed into the mutator strain *E. coli* XL1-Red (Stratagene, La Jolla, Calif.) according to manufacturer's protocols. Cells were harvested according to manufacturer's protocols and mutated plasmid DNA purified by standard methods.

[0105] Mutant pKK223-cterm-5×his-kgd ('5×his' disclosed as SEQ ID NO: 24) DNA is used to transform an *E. coli* host, NZN111+pBT-3-mmsA, described above. Greater than 10⁵ transformants are collected from LB ampicillin (200 g/L), Chloramphenicol (40 g/L) agarose plates. Cells are washed in LB media, diluted 1:10,000 and plated on LB media plates with 1 mM Isopropyl β-D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 200 g/L ampicillin and 40 g/L chloramphenicol. Plates are incubated at 37 C for several days anaerobically in BD type A Bio-Bags according to manufacturer's instructions (BD Biosciences, Franklin Lakes, N.J., Catalog #261214). Colonies that grow are individually collected as positives clones bearing oxaloacetate alpha-decarboxylase activity.

Example 9

Development of a Nucleic Acid Sequence Encoding a Protein Sequence Demonstrating Elevated Oxaloacetate Alpha-Decarboxylase Activity (Prophetic)

[0106] Oxaloacetate alpha-decarboxylase activity is selected from a pool of pyruvate decarboxylase (pdc) mutants by selection in an *E. coli* NZN111 host expressing an acetylating malonate semialdehyde dehydrogenase. pKK223-cterm-5×his-pdc ('5×his' disclosed as SEQ ID NO: 24) encoding the pdc gene is constructed as described above. Confirmation of pyruvate decarboxylase protein expression and enzymatic activity with appropriate controls are as follows. *E. Cloni* 10GF' electrocompetent cells (Lucigen, Cat. #60061-1) are transformed with the pKK223-Cterm-5×His-pdc ('5×His' disclosed as SEQ ID NO: 24), plasmid containing sequence for 5×HIS-tagged pdc protein ('5×His' disclosed as SEQ ID NO: 24) behind a pTAC promoter. Transformants are confirmed using restriction digest and DNA sequencing (Macrogen, Korea). Expression and purification of his-tagged-pdc are performed as described in Subsection III of the Common Methods Section. *E. coli* NZN111 and *E. coli* NZN111+pBT3-mmsA is transformed with the vector controls, pKK223, pKK223-Cterm-5×his ('5×His' disclosed as SEQ ID NO: 24), as well as the test vectors pKK223-mcr and pKK223-Cterm-5×His-pdc ('5×His' disclosed as SEQ ID NO: 24), according to standard methods described below. Each of the strains is grown overnight in LB rich media supplemented with 200 mg/L ampicillin (according to standard protocols). Following overnight growth, cells twice are harvested by centrifugation and washed by resuspension in LB media (standard protocol), diluted 1:10,000 and plated on

LB media plates with the additional appropriate supplements, where indicated at the following concentrations 1 mM Isopropyl β-D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 200 mg/L ampicillin (Research Products International Corp., Mt. Prospect, Ill.) After plating, agarose plates were incubated at 37 C overnight anaerobically in BD type A Bio-Bags according to manufacturer's instructions (BD Biosciences, Franklin Lakes, N.J., Catalog #261214). The *mcr* gene encoding the malonyl-coA reductase, has been shown to produce malonate semialdehyde. The presence of growth of the strain expressing malonyl-coA reductase in the presence of the co-expressed acetylating malonate semialdehyde in *E. coli* NZN111 serves as a positive control for the selection.

[0107] Mutant libraries of pKK223-cterm-5×his-pdc ('5×his' disclosed as SEQ ID NO: 24) are constructed as follows. Plasmid DNA of pKK223-cterm-5×his-pdc ('5×his' disclosed as SEQ ID NO: 24) are purified by standard methods and transformed into the mutator strain *E. coli* XL1-Red (Stratagene, La Jolla, Calif.) according to manufacturer's protocols. Cells are harvested according to manufacturer's protocols and mutated plasmid DNA purified by standard methods. Mutant pKK223-cterm-5×his-pdc ('5×his' disclosed as SEQ ID NO: 24) DNA is used to transform an *E. coli* host, NZN111+pBT-3-mmsA, described above. Greater than 10⁵ transformants are collected from LB ampicillin (200 g/L), chloramphenicol (40 g/L) agarose plates. Cells are washed in LB media, diluted 1:10,000 and plated on LB media plates with 1 mM Isopropyl β-D-1-thiogalactopyranoside (Thermo Fisher Scientific, Fairlawn, N.J.), 200 g/L ampicillin and 40 g/L chloramphenicol. Plates are incubated at 37 C for several days anaerobically in BD type A Bio-Bags according to manufacturer's instructions (BD Biosciences, Franklin Lakes, N.J., Catalog #261214). Colonies that grow are individually collected as positives clones bearing oxaloacetate alpha-decarboxylase activity.

Example 10

Confirmation of Oxaloacetate Alpha-Decarboxylase Activity (Partial Prophetic)

[0108] The colorimetric to confirm enzymatic decarboxylation of 2-oxo-acid substrates is adapted from current standard methodologies and is illustrated below in FIG. 11. This approach necessitates the expression and purification of the mutant enzymes and reaction with the purified enzyme, its cofactor (thiamin pyrophosphate) and the appropriate substrate. Protein expression and purification are performed with standard methodologies. This colorimetric screening method will be used both to conduct broad screening for positive oxaloacetate alpha-decarboxylase mutants, and also to conduct confirmatory testing of the positive clones identified in a selection method described above.

[0109] Transformants containing a gene cloned into the pKK223-Cterm-5×his expression vector ('5×his' disclosed as SEQ ID NO: 24) are grown overnight in LB+0.2% glucose+200 ug/mL Ampicillin, diluted 1:20 and grown (LB+0.2% glucose+200 ug/mL Ampicillin) to OD600 of 0.4. IPTG is added at 1 mM final concentration to induce protein expression. Cultures are then allowed to grow at 37 degrees C. for four hours. Cells were harvested by centrifugation at 4 degrees C. for 10 minutes at 4000 rpm. Pellets are resuspended and concentrated 50× (e.g. 500 mL culture resuspended in 10 mL buffer) in Qiagen Ni-NTA Lysis Buffer (50

mM Na₂HPO₄, 300 mM NaCl, 10 mM imidazole, pH 8.0)+1 mM PMSF. Lysozyme is added to a final concentration of 1 mg/mL; cells are incubated on ice for 30 minutes. Cells are lysed using a French Press (cell pressure=2000 psi) three times. Lysates are cleared by centrifugation at 4 degrees C. for 20 minutes, applied to Qiagen Ni-NTA columns, washed and eluted as specified by Qiagen (cat.#31314). Samples are analyzed by SDS-PAGE by routine protocols.

[0110] 100 uL reaction mixtures contain 50 mM Potassium phosphate (pH 7.0), 0.2 mM TPP, 1 mM MgCh, 10 mM of the appropriate substrate. 300 pg of purified enzyme is added to the reaction and incubated 16 hours at 37 degrees C. After 16 hours at 37 degrees C., 100 uL of Purpald colorimetric indicator (as per Sigma-Aldrich, cat.#162892) is added to each well in order to detect formation of corresponding aldehyde product. After addition of the Purpald, reactions are incubated at room temperature for 1 hour and read at a wavelength of 540 nm in a Thermomax Microplate Reader (Molecular Devices) using SOFTMax Pro Microplate Reader software, Ver. 4.0. Absorbances greater than control reactions without substrate are used to determine the presence of decarboxylation.

Common Methods Section

[0111] All methods in this Section are provided for incorporation into the above methods where so referenced therein and/or below.

[0112] Subsection I. Bacterial Growth Methods:

[0113] Bacterial growth culture methods, and associated materials and conditions, are disclosed for respective species that may be utilized as needed, as follows:

[0114] *Escherichia coli* K12 is a gift from the Gill lab (University of Colorado at Boulder) and is obtained as an actively growing culture. Serial dilutions of the actively growing *E. coli* K12 culture are made into Luria Broth (RPI Corp, Mt. Prospect, Ill., USA) and are allowed to grow for aerobically for 24 hours at 37° C. at 250 rpm until saturated.

[0115] *Pseudomonas aeruginosa* genomic DNA is a gift from the Gill lab (University of Colorado at Boulder).

[0116] Subsection II: Gel Preparation, DNA Separation, Extraction, Ligation, and Transformation Methods:

[0117] Molecular biology grade agarose (RPI Corp, Mt. Prospect, Ill., USA) is added to 1×TAE to make a 1% Agarose: TAE solution. To obtain 50×TAE add the following to 900 mL of distilled water: add the following to 900 ml distilled H₂O: 242 g Tris base (RPI Corp, Mt. Prospect, Ill., USA), 57.1 ml Glacial Acetic Acid (Sigma-Aldrich, St. Louis, Mo., USA) and 18.6 g EDTA (Fisher Scientific, Pittsburgh, Pa. USA) and adjust volume to 1 L with additional distilled water. To obtain 1×TAE, add 20 mL of 50×TAE to 980 mL of distilled water. The agarose-TAE solution is then heated until boiling occurred and the agarose is fully dissolved. The solution is allowed to cool to 50° C. before 10 mg/mL ethidium bromide (Acros Organics, Morris Plains, N.J., USA) is added at a concentration of 5 ul per 100 mL of 1% agarose solution. Once the ethidium bromide is added, the solution is briefly mixed and poured into a gel casting tray with the appropriate number of combs (Idea Scientific Co., Minneapolis, Minn., USA) per sample analysis. DNA samples are then mixed accordingly with 5×TAE loading buffer. 5×TAE loading buffer consists of 5×TAE(diluted from SOXTAE as described above), 20% glycerol (Acros Organics, Morris Plains, N.J., USA), 0.125% Bromophenol Blue (Alfa Aesar, Ward Hill, Mass., USA), and adjust volume to 50

mL with distilled water. Loaded gels are then run in gel rigs (Idea Scientific Co., Minneapolis, Minn., USA) filled with 1×TAE at a constant voltage of 125 volts for 25-30 minutes. At this point, the gels are removed from the gel boxes with voltage and visualized under a UV transilluminator (FOTO-DYNE Inc., Hartland, Wis., USA).

[0118] The DNA isolated through gel extraction is then extracted using the QIAquick Gel Extraction Kit following manufacturer's instructions (Qiagen (Valencia Calif. USA)). Similar methods are known to those skilled in the art.

[0119] The thus-extracted DNA then may be ligated into pSMART (Lucigen Corp, Middleton, Wis., USA), StrataClone (Stratagene, La Jolla, Calif., USA) or pCR2.1-TOPO TA (Invitrogen Corp, Carlsbad, Calif., USA) according to manufacturer's instructions. These methods are described in the next subsection of Common Methods.

Ligation Methods:

[0120] For Ligations into pSMART Vectors:

[0121] Gel extracted DNA is blunted using PCR Terminator (Lucigen Corp, Middleton, Wis., USA) according to manufacturer's instructions. Then 500 ng of DNA is added to 2.5 uL 4× CloneSmart vector premix, 1 ul CloneSmart DNA ligase (Lucigen Corp, Middleton, Wis., USA) and distilled water is added for a total volume of 10 ul. The reaction is then allowed to sit at room temperature for 30 minutes and then heat inactivated at 70° C. for 15 minutes and then placed on ice. *E. coli* 10G Chemically Competent cells (Lucigen Corp, Middleton, Wis., USA) are thawed for 20 minutes on ice. 40 ul of chemically competent cells are placed into a microcentrifuge tube and 1 ul of heat inactivated CloneSmart Ligation is added to the tube. The whole reaction is stirred briefly with a pipette tip. The ligation and cells are incubated on ice for 30 minutes and then the cells are heat shocked for 45 seconds at 42° C. and then put back onto ice for 2 minutes. 960 ul of room temperature Recovery media (Lucigen Corp, Middleton, Wis., USA) and places into microcentrifuge tubes. Shake tubes at 250 rpm for 1 hour at 37° C. Plate 100 ul of transformed cells on Luria Broth plates (RPI Corp, Mt. Prospect, Ill., USA) plus appropriate antibiotics depending on the pSMART vector used. Incubate plates overnight at 37° C.

For Ligations into StrataClone:

[0122] Gel extracted DNA is blunted using PCR Terminator (Lucigen Corp, Middleton, Wis., USA) according to manufacturer's instructions. Then 2 ul of DNA is added to 3 ul StrataClone Blunt Cloning buffer and 1 ul StrataClone Blunt vector mix amp/kan (Stratagene, La Jolla, Calif., USA) for a total of 6 ul. Mix the reaction by gently pipeting up at down and incubate the reaction at room temperature for 30 minutes then place onto ice. Thaw a tube of StrataClone chemically competent cells (Stratagene, La Jolla, Calif., USA) on ice for 20 minutes. Add 1 ul of the cloning reaction to the tube of chemically competent cells and gently mix with a pipette tip and incubate on ice for 20 minutes. Heat shock the transformation at 42° C. for 45 seconds then put on ice for 2 minutes. Add 250 ul pre-warmed Luria Broth (RPI Corp, Mt. Prospect, Ill., USA) and shake at 250 rpm for 37° C. for 2 hour. Plate 100 ul of the transformation mixture onto Luria Broth plates (RPI Corp, Mt. Prospect, Ill., USA) plus appropriate antibiotics. Incubate plates overnight at 37° C.

For Ligations into pCR2.1-TOPO TA:

[0123] Add 1 ul TOPO vector, 1 ul Salt Solution (Invitrogen Corp, Carlsbad, Calif., USA) and 3 ul gel extracted DNA into a microcentrifuge tube. Allow the tube to incubate at room

temperature for 30 minutes then place the reaction on ice. Thaw one tube of TOPIOF¹ chemically competent cells (Invitrogen Corp, Carlsbad, Calif., USA) per reaction. Add 1 ul of reaction mixture into the thawed TOPIOF¹ cells and mix gently by swirling the cells with a pipette tip and incubate on ice for 20 minutes. Heat shock the transformation at 42° C. for 45 seconds then put on ice for 2 minutes. Add 250 ul pre-warmed SOC media (Invitrogen Corp, Carlsbad, Calif., USA) and shake at 250 rpm for 37° C. for 1 hour. Plate 100 ul of the transformation mixture onto Luria Broth plates (RPI Corp, Mt. Prospect, Ill., USA) plus appropriate antibiotics. Incubate plates overnight at 37° C.

General Transformation and Related Culture Methodologies:

[0124] Chemically competent transformation protocols are carried out according to the manufactures instructions or according to the literature contained in Molecular Cloning (Sambrook and Russell). Generally, plasmid DNA or ligation products are chilled on ice for 5 to 30 min. in solution with chemically competent cells. Chemically competent cells are a widely used product in the field of biotechnology and are available from multiple vendors, such as those indicated above in this Subsection. Following the chilling period cells generally are heat-shocked for 30 seconds at 42° C. without shaking, re-chilled and combined with 250 microliters of rich media, such as S.O.C. Cells are then incubated at 37° C. while shaking at 250 rpm for 1 hour. Finally, the cells are screened for successful transformations by plating on media containing the appropriate antibiotics.

[0125] The choice of an *E. coli* host strain for plasmid transformation is determined by considering factors such as plasmid stability, plasmid compatibility, plasmid screening methods and protein expression. Strain backgrounds can be changed by simply purifying plasmid DNA as described above and transforming the plasmid into a desired or otherwise appropriate *E. coli* host strain such as determined by experimental necessities, such as any commonly used cloning strain (e.g., DH5a, Top IOP¹, *E. doni* 10G, etc.).

To Make 1 L M9 Minimal Media:

[0126] M9 minimal media was made by combining 5×M9 salts, 1M MgSO₄, 20% glucose, 1M CaCl₂ and sterile deionized water. The 5×M9 salts are made by dissolving the following salts in deionized water to a final volume of 1 L: 64 g Na₂HPO₄ · 7H₂O, 15 g KH₂PO₄, 2.5 g NaCl, 5.0 g NH₄Cl. The salt solution was divided into 200 mL aliquots and sterilized by autoclaving for 15 minutes at 15 psi on the liquid cycle. A 1M solution of MgSO₄ and 1M CaCl₂ were made separately, then sterilized by autoclaving. The glucose was filter sterilized by passing it through a 0.22 μm filter. All of the components are combined as follows to make 1 L of M9: 750 mL sterile water, 200 mL 5×M9 salts, 2 mL of 1M MgSO₄, 20 mL 20% glucose, 0.1 mL CaCl₂, Q.S. to a final volume of 1 L.

To Make EZ Rich Media:

[0127] All media components were obtained from TEKnova (Hollister Calif. USA) and combined in the following volumes. 100 mL 10×MOPS mixture, 10 mL 0.132M K₂HP0₄, 100 mL 10×ACGU, 200 mL 5× Supplement EZ, 10 mL 20% glucose, 580 mL sterile water.

Subsection III. Additional Methods Related to Enzyme Evaluation Expression and Purification of Proteins Expressed in pKK223-Cterm-5×his (‘5×his’ Disclosed as SEQ ID NO: 24) by Expression Plasmids

[0128] Transformants containing a gene cloned into the pKK223-Cterm-5×his (‘5×his’ disclosed as SEQ ID NO: 24) expression vector were grown overnight in LB+0.2% glucose+200 ug/mL Ampicillin, diluted 1:20 and grown (LB+0.2% glucose+200 ug/mL Ampicillin) to OD600 of 0.4. IPTG was added at 1 mM final concentration to induce protein expression. Cultures were then allowed to grow at 37 degrees C. for four hours. Cells were harvested by centrifugation at 4 degrees C. for 10 minutes at 4000 rpm. Pellets were resuspended and concentrated 50× (e.g. pellet from 500 mL culture resuspended in 10 mL buffer) in Qiagen Ni-NTA Lysis Buffer (50 mM Na₂HP0₄, 300 mM NaCl, 10 mM imidazole, pH 8.0)+1 mM PMSF. Lysozyme was added to a final concentration of 1 mg/mL; cells were incubated on ice for 30 minutes. Cells were lysed using a French Press (cell pressure=2000 psi) three times. Lysates were cleared by centrifugation at 4 degrees C. for 20 minutes, applied to Qiagen Ni-NTA columns, washed and eluted as specified by Qiagen (cat.#31314). Samples were analyzed by SDS-PAGE by routine protocols.

Decarboxylation Enzyme Reactions:

[0129] 100 uL reaction mixtures were added to microwells. Each 100 uL of reaction mixture contained 50 mM Potassium Phosphate (pH 7.0), 0.2 mM TPP, 1 mM MgCl₂, and 10 mM of the appropriate substrate. 300 pg of purified enzyme was added to a respective microwell and incubated 16 hours at 37 degrees C. After 16 hours at 37 degrees C., 100 uL of Purpald® colorimetric indicator (Sigma-Aldrich, cat.#162892), prepared per manufacturer’s instructions, was added to each microwell in order to detect formation of corresponding aldehyde product. After addition of the Purpald®, the microwells were incubated at room temperature for 1 hour and read at a wavelength of 540 nm in a Thermomax Microplate Reader (Molecular Devices) using SOFTMax Pro Microplate Reader software, Ver. 4.0.

Summary of Suppliers Section

[0130] The names and city addresses of major suppliers are provided in the methods above. In addition, as to Qiagen products, the DNeasy® Blood and Tissue Kit, Cat. No. 69506, is used in the methods for genomic DNA preparation; the QIAprep® Spin (“mini prep”), Cat. No. 27106, is used for plasmid DNA purification, and the QIAquick® Gel Extraction Kit, Cat. No. 28706, is used for gel extractions as described above.

Bio-Production Media

[0131] Bio-production media, which is used in the present invention with recombinant microorganisms having a biosynthetic pathway for 3-HP (and optionally products further downstream of 3-HP), must contain suitable carbon substrates. Suitable substrates may include, but are not limited to, monosaccharides such as glucose and fructose, oligosaccharides such as lactose or sucrose, polysaccharides such as starch or cellulose or mixtures thereof and unpurified mixtures from renewable feed stocks such as cheese whey permeate, comsteep liquor, sugar beet molasses, and barley malt. Additionally the carbon substrate may also be one-carbon

substrates such as carbon dioxide, or methanol for which metabolic conversion into key biochemical intermediates has been demonstrated. In addition to one and two carbon substrates methylotrophic organisms are also known to utilize a number of other carbon containing compounds such as methylamine, glucosamine and a variety of amino acids for metabolic activity. For example, methylotrophic yeast are known to utilize the carbon from methylamine to form trehalose or glycerol (Bellion et al., *Microb. Growth Cl Compd.*, [Int. Symp.], 7th (1993), 415-32. Editor(s): Murrell, J. Collin; Kelly, Don P. Publisher: Intercept, Andover, UK). Similarly, various species of *Candida* will metabolize alanine or oleic acid (Suiter et al., *Arch. Microbial.* 153:485-489 (1990)). Hence it is contemplated that the source of carbon utilized in the present invention may encompass a wide variety of carbon containing substrates and will only be limited by the choice of organism. Although it is contemplated that all of the above mentioned carbon substrates and mixtures thereof are suitable in the present invention as a carbon source, common carbon substrates used as carbon sources are glucose, fructose, and sucrose, as well as mixtures of any of these sugars. Sucrose may be obtained from feed stocks such as sugar cane, sugar beets, cassava, and sweet sorghum. Glucose and dextrose may be obtained through saccharification of starch based feed stocks including grains such as corn, wheat, rye, barley, and oats.

[0132] In addition, sugars may be obtained from cellulosic and lignocellulosic biomass through processes of pretreatment and saccharification, as described, for example, in US patent application US20070031918A1, which is herein incorporated by reference. Biomass refers to any cellulosic or lignocellulosic material and includes materials comprising cellulose, and optionally further comprising hemicellulose, lignin, starch, oligosaccharides and/or monosaccharides. Biomass may also comprise additional components, such as protein and/or lipid. Biomass may be derived from a single source, or biomass can comprise a mixture derived from more than one source; for example, biomass could comprise a mixture of corn cobs and corn stover, or a mixture of grass and leaves. Biomass includes, but is not limited to, bioenergy crops, agricultural residues, municipal solid waste, industrial solid waste, sludge from paper manufacture, yard waste, wood and forestry waste. Examples of biomass include, but are not limited to, corn grain, corn cobs, crop residues such as corn husks, corn stover, grasses, wheat, wheat straw, barley, barley straw, hay, rice straw, switchgrass, waste paper, sugar cane bagasse, sorghum, soy, components obtained from milling of grains, trees, branches, roots, leaves, wood chips, sawdust, shrubs and bushes, vegetables, fruits, flowers and animal manure. Any such biomass may be used in a bio-production method or system to provide a carbon source. In addition to an appropriate carbon source, such as selected from one of the above-disclosed types, bio-production media must contain suitable minerals, salts, cofactors, buffers and other components, known to those skilled in the art, suitable for the growth of the cultures and promotion of the enzymatic pathway necessary for 3-HP (and optionally products further downstream of 3-HP) production.

Culture Conditions

[0133] Typically cells are grown at a temperature in the range of about 25° C. to about 40° C. in an appropriate medium. Suitable growth media in the present invention are common commercially prepared media such as Luria Bertani

(LB) broth, M9 minimal media, Sabouraud Dextrose (SD) broth, Yeast medium (YM) broth or (Ymin) yeast synthetic minimal media. Other defined or synthetic growth media may also be used, and the appropriate medium for growth of the particular microorganism will be known by one skilled in the art of microbiology or bio-production science.

[0134] Suitable pH ranges for the bio-production are between pH 5.0 to pH 9.0, where pH 6.0 to pH 8.0 is a typical pH range for the initial condition.

[0135] Bio-productions may be performed under aerobic, microaerobic, or anaerobic conditions, with or without agitation.

[0136] The amount of 3-HP (and optionally products further downstream of 3-HP) produced in a bio-production media generally can be determined using a number of methods known in the art, for example, high performance liquid chromatography (HPLC) or gas chromatography (GC). Specific HPLC methods for the specific examples are provided herein.

Bio-Production Reactors and Systems:

[0137] Any of the recombinant microorganisms as described and/or referred to above may be introduced into an industrial bio-production system where the microorganisms convert a carbon source into 3-HP (and optionally products further downstream of 3-HP) in a commercially viable operation. The bio-production system includes the introduction of such a recombinant microorganism into a bioreactor vessel, with a carbon source substrate and bio-production media suitable for growing the recombinant microorganism, and maintaining the bio-production system within a suitable temperature range (and dissolved oxygen concentration range if the reaction is aerobic or microaerobic) for a suitable time to obtain a desired conversion of a portion of the substrate molecules to 3-HP (and optionally products further downstream of 3-HP). Industrial bio-production systems and their operation are well-known to those skilled in the arts of chemical engineering and bioprocess engineering. The following paragraphs provide an overview of the methods and aspects of industrial systems that may be used for the bio-production of 3-HP (and optionally products further downstream of 3-HP).

[0138] In various embodiments, any of a wide range of sugars, including, but not limited to sucrose, glucose, xylose, cellulose or hemicellulose, are provided to a microorganism, such as in an industrial system comprising a reactor vessel in which a defined media (such as a minimal salts media including but not limited to M9 minimal media, potassium sulfate minimal media, yeast synthetic minimal media and many others or variations of these), an inoculum of a microorganism providing one or more of the 3-HP (and optionally products further downstream of 3-HP) biosynthetic pathway alternatives, and the a carbon source may be combined. The carbon source enters the cell and is catabolized by well-known and common metabolic pathways to yield common metabolic intermediates, including phosphoenolpyruvate (PEP). (See *Molecular Biology of the Cell*, 3rd Ed., B. Alberts et al. Garland Publishing, New York, 1994, pp. 42-45, 66-74, incorporated by reference for the teachings of basic metabolic catabolic pathways for sugars; *Principles of Biochemistry*, 3rd Ed., D. L. Nelson & M. M. Cox, Worth Publishers, New York, 2000, pp 527-658, incorporated by reference for the teachings of major metabolic pathways; and *Biochemistry*, 4th Ed., L. Stryer, W. H. Freeman and Co., New York, 1995, pp. 463-650, also incorporated by reference for the teachings

of major metabolic pathways.). Further to types of industrial bio-production, various embodiments of the present invention may employ a batch type of industrial bioreactor. A classical batch bioreactor system is considered “closed” meaning that the composition of the medium is established at the beginning of a respective bio-production event and not subject to artificial alterations and additions during the time period ending substantially with the end of the bio-production event. Thus, at the beginning of the bio-production event the medium is inoculated with the desired organism or organisms, and bio-production is permitted to occur without adding anything to the system. Typically, however, a “batch” type of bio-production event is batch with respect to the addition of carbon source and attempts are often made at controlling factors such as pH and oxygen concentration. In batch systems the metabolite and biomass compositions of the system change constantly up to the time the bio-production event is stopped. Within batch cultures cells moderate through a static lag phase to a high growth log phase and finally to a stationary phase where growth rate is diminished or halted. If untreated, cells in the stationary phase will eventually die. Cells in log phase generally are responsible for the bulk of production of a desired end product or intermediate.

[0139] A variation on the standard batch system is the Fed-Batch system. Fed-Batch bio-production processes are also suitable in the present invention and comprise a typical batch system with the exception that the substrate is added in increments as the bio-production progresses. Fed-Batch systems are useful when catabolite repression is apt to inhibit the metabolism of the cells and where it is desirable to have limited amounts of substrate in the media. Measurement of the actual substrate concentration in Fed-Batch systems may be measured directly, such as by sample analysis at different times, or estimated on the basis of the changes of measurable factors such as pH, dissolved oxygen and the partial pressure of waste gases such as CO₂. Batch and Fed-Batch approaches are common and well known in the art and examples may be found in Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition (1989) Sinauer Associates, Inc., Sunderland, Mass., Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36:227, (1992), and *Biochemical Engineering Fundamentals*, 2nd Ed. J. E. Bailey and D. F. Ollis, McGraw Hill, New York, 1986, herein incorporated by reference for general instruction on bio-production, which as used herein may be aerobic, microaerobic, or anaerobic. Although the present invention may be performed in batch mode, as provided in Example 8, or in fed-batch mode, it is contemplated that the method would be adaptable to continuous bio-production methods. Continuous bio-production is considered an “open” system where a defined bio-production medium is added continuously to a bioreactor and an equal amount of conditioned media is removed simultaneously for processing. Continuous bio-production generally maintains the cultures within a controlled density range where cells are primarily in log phase growth. Two types of continuous bioreactor operation include: 1) Chemostat—where fresh media is fed to the vessel while simultaneously removing an equal rate of the vessel contents. The limitation of this approach is that cells are lost and high cell density generally is not achievable. In fact, typically one can obtain much higher cell density with a fed-batch process. 2) Perfusion culture, which is similar to the chemostat approach except that the stream that is removed from the vessel is subjected to a separation technique which recycles viable cells back to the vessel. This type of continu-

ous bioreactor operation has been shown to yield significantly higher cell densities than fed-batch and can be operated continuously. Continuous bio-production is particularly advantageous for industrial operations because it has less down time associated with draining, cleaning and preparing the equipment for the next bio-production event. Furthermore, it is typically more economical to continuously operate downstream unit operations, such as distillation, than to run them in batch mode.

[0140] Continuous bio-production allows for the modulation of one factor or any number of factors that affect cell growth or end product concentration. For example, one method will maintain a limiting nutrient such as the carbon source or nitrogen level at a fixed rate and allow all other parameters to moderate. In other systems a number of factors affecting growth can be altered continuously while the cell concentration, measured by media turbidity, is kept constant. Continuous systems strive to maintain steady state growth conditions and thus the cell loss due to the medium being drawn off must be balanced against the cell growth rate in the bio-production. Methods of modulating nutrients and growth factors for continuous bio-production processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology and a variety of methods are detailed by Brock, supra.

[0141] It is contemplated that embodiments of the present invention may be practiced using either batch, fed-batch or continuous processes and that any known mode of bio-production would be suitable. Additionally, it is contemplated that cells may be immobilized on an inert scaffold as whole cell catalysts and subjected to suitable bio-production conditions for 3-HP (and optionally products further downstream of 3-HP) production.

[0142] The following published resources are incorporated by reference herein for their respective teachings to indicate the level of skill in these relevant arts, and as needed to support a disclosure that teaches how to make and use methods of industrial bio-production of 3-HP (and optionally products further downstream of 3-HP) from sugar sources, and also industrial systems that may be used to achieve such conversion with any of the recombinant microorganisms of the present invention (*Biochemical Engineering Fundamentals*, 2nd Ed. J. E. Bailey and D. F. Ollis, McGraw Hill, New York, 1986, entire book for purposes indicated and Chapter 9, pages 533-657 in particular for biological reactor design; *Unit Operations of Chemical Engineering*, 5th Ed., W. L. McCabe et al., McGraw Hill, New York 1993, entire book for purposes indicated, and particularly for process and separation technologies analyses; *Equilibrium Staged Separations*, P. C. Wankat, Prentice Hall, Englewood Cliffs, N.J. USA, 1988, entire book for separation technologies teachings).

[0143] The scope of the present invention is not meant to be limited to the exact sequences provided herein. It is appreciated that a range of modifications to nucleic acid and to amino acid sequences may be made and still provide a desired functionality. The following discussion is provided to more clearly define ranges of variation that may be practiced and still remain within the scope of the present invention.

[0144] It is recognized in the art that some amino acid sequences of the present invention can be varied without significant effect of the structure or function of the proteins disclosed herein. Variants included can constitute deletions, insertions, inversions, repeats, and type substitutions so long as the indicated enzyme activity is not significantly affected.

Guidance concerning which amino acid changes are likely to be phenotypically silent can be found in Bowie, J. U., et Al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990).

[0145] In various embodiments polypeptides obtained by the expression of the polynucleotide molecules of the present invention may have at least approximately 80%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identity to one or more amino acid sequences encoded by the genes and/or nucleic acid sequences described herein for the 3-HP (and optionally products further downstream of 3-HP) biosynthesis pathways. A truncated respective polypeptide has at least about 90% of the full length of a polypeptide encoded by a nucleic acid sequence encoding the respective native enzyme, and more particularly at least 95% of the full length of a polypeptide encoded by a nucleic acid sequence encoding the respective native enzyme. By a polypeptide having an amino acid sequence at least, for example, 95% "identical" to a reference amino acid sequence of a polypeptide is intended that the amino acid sequence of the claimed polypeptide is identical to the reference sequence except that the claimed polypeptide sequence can include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the polypeptide. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence can be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence can be inserted into the reference sequence. These alterations of the reference sequence can occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence.

[0146] As a practical matter, whether any particular polypeptide is at least 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98% or 99% identical to any reference amino acid sequence of any polypeptide described herein (which may correspond with a particular nucleic acid sequence described herein), such particular polypeptide sequence can be determined conventionally using known computer programs such as the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, Wis. 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference sequence according to the present invention, the parameters are set, of course, such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed. For example, in a specific embodiment the identity between a reference sequence (query sequence, a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, may be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (*Comp. App. Biosci.* 6:237-245 (1990)). Preferred parameters used in a FASTDB amino acid alignment are: Matrix=PAM 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.

05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter. According to this embodiment, if the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction is made to the results to take into consideration the fact that the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. A determination of whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the above FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of this embodiment. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence. For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for.

[0147] The above descriptions and methods for sequence homology are intended to be exemplary and it is recognized that this concept is well-understood in the art. Further, it is appreciated that nucleic acid sequences may be varied and still provide a functional enzyme, and such variations are within the scope of the present invention. Nucleic acid sequences that encode polypeptides that provide the indicated functions for 3-HP (and optionally products further downstream of 3-HP) that increase tolerance or production are considered within the scope of the present invention. These may be further defined by the stringency of hybridization, described below, but this is not meant to be limiting when a function of an encoded polypeptide matches a specified 3-HP (and optionally products further downstream of 3-HP) tolerance-related or biosynthesis pathway enzyme activity.

[0148] Further to nucleic acid sequences, "hybridization" refers to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded

polynucleotide. The term “hybridization” may also refer to triple-stranded hybridization. The resulting (usually) double-stranded polynucleotide is a “hybrid” or “duplex.” “Hybridization conditions” will typically include salt concentrations of less than about 1M, more usually less than about 500 mM and less than about 200 mM. Hybridization temperatures can be as low as 5° C., but are typically greater than 22° C., more typically greater than about 30° C., and often are in excess of about 37° C. Hybridizations are usually performed under stringent conditions, i.e. conditions under which a probe will hybridize to its target subsequence. Stringent conditions are sequence-dependent and are different in different circumstances. Longer fragments may require higher hybridization temperatures for specific hybridization. As other factors may affect the stringency of hybridization, including base composition and length of the complementary strands, presence of organic solvents and extent of base mismatching, the combination of parameters is more important than the absolute measure of any one alone. Generally, stringent conditions are selected to be about 5° C. lower than the T_m for the specific sequence at a defined ionic strength and pH. Exemplary stringent conditions include salt concentration of at least 0.01 M to no more than 1M Na ion concentration (or other salts) at a pH 7.0 to 8.3 and a temperature of at least 25° C. For example, conditions of 5×SSPE (750 mM NaCl, 50 mM Na Phosphate, 5 mM EDTA, pH 7.4) and a temperature of 25-30° C. are suitable for allele-specific probe hybridizations. For stringent conditions, see for example, Sambrook and Russell and Anderson “Nucleic Acid Hybridization” 1st Ed., BIOS Scientific Publishers Limited (1999), which are hereby incorporated by reference for hybridization protocols. “Hybridizing specifically to” or “specifically hybridizing to” or like expressions refer to the binding, duplexing, or hybridizing of a molecule substantially to or only to a particular nucleotide sequence or sequences under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular DNA or RNA).

[0149] Having so described the present invention and provided examples, and further discussion, and in view of the above paragraphs, it is appreciated that various non-limiting aspects of the present invention may include:

[0150] A genetically modified (recombinant) microorganism comprising a nucleic acid sequence that encodes a polypeptide with at least 85% amino acid sequence identity to any of the enzymes of any of 3-HP tolerance-related or biosynthetic pathways, wherein the polypeptide has enzymatic activity effective to perform the enzymatic reaction of the respective 3-HP biosynthetic pathway enzyme, and the recombinant microorganism exhibits greater 3-H tolerance and/or 3-HP bio-production.

[0151] A genetically modified (recombinant) microorganism comprising a nucleic acid sequence that encodes a polypeptide with at least 90% amino acid sequence identity to any of the enzymes of any of 3-HP tolerance-related or biosynthetic pathways, wherein the polypeptide has enzymatic activity effective to perform the enzymatic reaction of the respective 3-HP tolerance-related or biosynthetic pathway enzyme, and the recombinant microorganism exhibits greater 3-HP tolerance and/or 3-HP bio-production.

[0152] A genetically modified (recombinant) microorganism comprising a nucleic acid sequence that encodes a polypeptide with at least 95% amino acid sequence identity to any of the enzymes of any of 3-HP tolerance-related or biosynthetic pathways, wherein the polypeptide has enzymatic

activity effective to perform the enzymatic reaction of the respective 3-HP tolerance-related or biosynthetic pathway enzyme, and the recombinant microorganism exhibits greater 3-HP tolerance and/or 3-HP bio-production.

[0153] The above paragraphs are meant to indicate modifications in the nucleic acid sequences may be made and a respective polypeptide encoded therefrom remains functional so as to perform an enzymatic catalysis along one of the 3-HP tolerance-related and/or biosynthetic pathways described above.

[0154] The term “heterologous DNA,” “heterologous nucleic acid sequence,” and the like as used herein refers to a nucleic acid sequence wherein at least one of the following is true: (a) the sequence of nucleic acids is foreign to (i.e., not naturally found in) a given host microorganism; (b) the sequence may be naturally found in a given host microorganism, but in an unnatural (e.g., greater than expected) amount; or (c) the sequence of nucleic acids comprises two or more subsequences that are not found in the same relationship to each other in nature. For example, regarding instance (c), a heterologous nucleic acid sequence that is recombinantly produced will have two or more sequences from unrelated genes arranged to make a new functional nucleic acid. Embodiments of the present invention may result from introduction of an expression vector into a host microorganism, wherein the expression vector contains a nucleic acid sequence coding for an enzyme that is, or is not, normally found in a host microorganism. With reference to the host microorganism’s genome, then, the nucleic acid sequence that codes for the enzyme is heterologous.

[0155] Also, and more generally, in accordance with examples and embodiments herein, there may be employed conventional molecular biology, cellular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. (See, e.g., Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Third Edition 2001 (volumes 1-3), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.; *Animal Cell Culture*, R. I. Freshney, ed., 1986). These published resources are incorporated by reference herein for their respective teachings of standard laboratory methods found therein. Further, all patents, patent applications, patent publications, and other publications referenced herein (collectively, “published resource(s)”) are hereby incorporated by reference in this application. Such incorporation, at a minimum, is for the specific teaching and/or other purpose that may be noted when citing the reference herein. If a specific teaching and/or other purpose is not so noted, then the published resource is specifically incorporated for the teaching(s) indicated by one or more of the title, abstract, and/or summary of the reference. If no such specifically identified teaching and/or other purpose may be so relevant, then the published resource is incorporated in order to more fully describe the state of the art to which the present invention pertains, and/or to provide such teachings as are generally known to those skilled in the art, as may be applicable. However, it is specifically stated that a citation of a published resource herein shall not be construed as an admission that such is prior art to the present invention.

[0156] Thus, based on the above disclosure, it is appreciated that within the scope of the present invention are methods for selection and identification of mutant polynucleotides comprising nucleic acid sequences that encode mutant polypeptides that demonstrate elevated activity of oxaloac-

etate alpha-oxo decarboxylase activity (also referred to herein as oxaloacetate alpha-decarboxylase activity). Also within the scope of the present invention may be compositions that comprise such identified mutant polynucleotides and polypeptides. In various embodiments, these methods are directed for the specific purpose of obtaining recombinant microorganisms that have capacity for increased bio-production of 3-HP. Although specific genes, enzymes, plasmids and other constructs are described in the above examples, these are not meant to limit the scope of the invention, particularly in view of the level of skill in the art.

[0157] Thus, while various embodiments of the present invention have been shown and described herein, it will be obvious that such embodiments are provided by way of example only. Numerous variations, changes and substitutions may be made without departing from the invention herein in its various embodiments. Specifically, and for what-

ever reason, for any grouping of compounds, nucleic acid sequences, polypeptides including specific proteins including functional enzymes, metabolic pathway enzymes or intermediates, elements, or other compositions, or concentrations stated or otherwise presented herein in a list, table, or other grouping (such as metabolic pathway enzymes shown in a figure), unless clearly stated otherwise, it is intended that each such grouping provides the basis for and serves to identify various subset embodiments, the subset embodiments in their broadest scope comprising every subset of such grouping by exclusion of one or more members (or subsets) of the respective stated grouping. Moreover, when any range is described herein, unless clearly stated otherwise, that range includes all values therein and all sub-ranges therein. Accordingly, it is intended that the invention be limited only by the spirit and scope of appended claims, and of later claims, and of either such claims as they may be amended during prosecution of this or a later application claiming priority hereto.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 25

<210> SEQ ID NO 1

<211> LENGTH: 3736

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Codon optimized kgd gene sequence with customized ends

<400> SEQUENCE: 1

```

ttgacggata tcaagcttct attaaccgaa cgcttcgtcc aagatttctt gctgctccac      60
ggcatgaacc ttcgaggaac cgctgctcgg cgagacatc gcgagacggg agatacgctt      120
gatgcccgcc aacttgctcg gcagcaactc cggcaactcc aaaccgaaac gcggccaggg      180
gccctgggtc gctgggttct cctggacca aaagaactct ttgacatttt cgtaacgatc      240
cagggtttca cgcagacgac gacgcggcaa cggcgccagt tgctccagac gcacaattgc      300
cagatcatta cggttgtctt tcgccttgcg tgccgccaat tcataataca acttaccgga      360
ggtcaacaga atacggctaa ctttattgcg gtcgccgata ccatacctct aggtcgggtc      420
ctccaggacg ctacggaatt taatctcggg aaaatcctta atctccgaaa ccgccgcctt      480
gtggcgcagc atggatttcg gggtaaacac gatcaacgga cgttggatgc cgtccaacgc      540
gtggcggcgc agcaagtga agtaattgga cggggtgctc ggcacgcga tcgtcatgct      600
accctcagcc cacagctgca aaaagcgtc aatgcgcgcc gacgtgtggt ccgggccctg      660
accctcgtgg ccgtgtggca gcaacagaac aacgttgga agctgacccc atttcgcttc      720
gcccgagctg atgaactcgt caatgatcga ctgggcacca ttaacaaagt cgccgaattg      780
cgctcccac agcacaaccg cgtctggatt gccaacggta taaccgtact caaaaccac      840
agccgcgtat tccgacaacg gogaatcata aaccaggaac ttaccaccgg tcgggctgcc      900
gtcgtgttta gtcgccagca gctgcagcgg ggtgaactcc tcgccggtgt gacggtcgat      960
cagcacagaa tgacgctggc taaaggtgcc acgacgagag tcttgaccgc tcagacggac     1020
cagcttgccc tcagcaacca ggctgcccag cgcagcaac tcaccaaacg ccagtcaat     1080
cttaccctca tacgccatct cagcagctt ttccagcacc ggctgaacgc gcgggtgtgc     1140
cgtgaaaccg ttaggcagcg ccaggaaggc atcaccgata cgtgccagca gggatttgtc     1200

```

-continued

aacagcggtc gccaggcccg ctgggatcat ttggtccgac tcgacgctct ccgacggttg 1260
gacgcggtgc ttctccagtt cgcgcacttc gttgaacaca cgctccagtt ggccctggta 1320
atcgcgcagc gcacccctccg cctctttcat gctgatgtcg ccacgaccga tcagagcctc 1380
ggtgtaggat ttacggggcg cacgcttggg gtccacgacg tcatagacat acggattggg 1440
catagatgga tcgtcacctt cattatgacc acgacgacga tagcacagca tgtcaataac 1500
aacgtccttt ttaaagcgtt ggcggaagtc aactgccaaa cgcgcaacc agacacacgc 1560
ttctggatcg tcgccgttca cgtggaagat tggcgcaccg atcatcttcg ccacatccgt 1620
gcagtactcg ctggaacgag agtattccgg agcgggtggg aagccgattt ggttggtcac 1680
aatgatgtga atcgtaccac ccacacgata gccaggcaga tttgccagat tcagcgtctc 1740
cgcaacaacg ccttgaccgg cgaaggccgc atcacctgac agcatcaacg gaacgacgga 1800
gaatgcacgt tggccgtcgg agtcaatgga accgtggtec agcaggtctt gcttcgcacg 1860
caccaaacct tccagcactg gatcgaccgc ttccaaatgg gacggatttg ccgtcagggg 1920
aacctgaata tcggttatcg caaacatttg cagatacaga cccgtcgcac ccaggtggta 1980
cttgacatcg ccggaaccgt gagcctggga cgggttcaga ttgccttcaa actccgtaaa 2040
gatttgcgaa tacggtttgc ccacgatgtt cgcaggaca ttcaagcgac cacggtgccg 2100
catgccaatg accacttcat ccaaaccatg ctccggacat tggtaaatcg ccgctccat 2160
cattggaata acagattccg caccctccag gctaaaacgc ttttgccca catacttggg 2220
ttgcaggaag gtttcaaacg cctccgctgc gttcagtttc gacaagatgt acttttggtg 2280
agcaacggtc ggtttgacgt gcttcgtctc gacacgctgc tccagccact ccttttggtc 2340
cgggtccaga atgtgcgct actcaacacc gatgtgacgg cagtacgctg cgcgcagcaa 2400
accagcacg tcacgcagct ttttgatttg agcaccgcg aaaccgtcaa ccttaaagac 2460
gcgggtccagg tcccacagag tcaggccatg cgtcaacacc tccaaatccg gatgccaacg 2520
aaagcgcgcc ttatccaagc gcaacgggtc ggtgtccgcc atcagatggc cgcgggtgcg 2580
ataggccgcg atcaggttca tcacacgtgc gttcttgcga acgatcgagt ccggattatc 2640
ggtgctccaa cgcactggca ggtacgggat gctcagctcg ccgaagacct catcccagaa 2700
gccatcagac aacagcagtt catgaatggg acgcaggaag tcaccgcttt ccgcaccttg 2760
aatgatcagg tggctgtagg tagaagtcag ggtaatcagt ttgccaatac ccagctcagc 2820
gatgcgttcc tcggacgctg cttggaactc cgcgggatat tccatcgcac cgacaccgat 2880
gatagcaact tgacctggca tcaggcgtgg cacagagtgc accgtgcaa tcgtgcccgg 2940
attcgtcagc gaaatcgtaa cgcagcga gtcctcggtg gtcagtttac catcacgagc 3000
acgacggacg atgtcctcgt acgaggtaac gaactgcgag aagcgcacg tctcgcaacg 3060
tttgataccg gccaccacca gagagcgctt gccatcttta ccttgacaggt caatagccag 3120
accagattg gtgtgcgag gagtaaccgc cgtcggctta ccgtccacct ccgtgtagtg 3180
gcgattcata ttcgggaact tcttaaccgc ctgaaccaga gcataacca gcaaatgggt 3240
aaagctgatt ttaccaccac gcgtgcgctt caactgatta ttgatcacga tacgattatc 3300
gatcaacaat ttcgctggca cagcacgcac cgaggtcgcc gtaggcactt ccagcgcgc 3360
gctcatgttc ttacgacag cagccgccc accacgcagg acagcaactt catcgcttc 3420
ggctggcgga ggcaccgagg tcttggcggc cagagccgcc acgacaccgt tgcccgtgc 3480

-continued

```

ggcgtgctcg gccggtttcg gggcgcttg cggagccgcc gctgccgcac gctcagcgac 3540
caaagggctg gtcacacgag tcggctcagc agccggttgg gaagtcggct ccgggctata 3600
gtccaccaga aactcatgcc agcttgggtc aacggaagac ggatcatcac gaaatttacg 3660
atacatacca acacgagacg ggtcctgagt caccatggat atatctcctt cttaaagaat 3720
tcgatatctc agcgac 3736

```

```

<210> SEQ ID NO 2
<211> LENGTH: 1224
<212> TYPE: PRT
<213> ORGANISM: Mycobacterium tuberculosis

```

```

<400> SEQUENCE: 2

```

```

Met Val Thr Gln Asp Pro Ser Arg Val Gly Met Tyr Arg Lys Phe Arg
1          5          10          15
Asp Asp Pro Ser Ser Val Asp Pro Ser Trp His Glu Phe Leu Val Asp
20          25          30
Tyr Ser Pro Glu Pro Thr Ser Gln Pro Ala Ala Glu Pro Thr Arg Val
35          40          45
Thr Ser Pro Leu Val Ala Glu Arg Ala Ala Ala Ala Ala Pro Gln Ala
50          55          60
Pro Pro Lys Pro Ala Asp Thr Ala Ala Ala Gly Asn Gly Val Val Ala
65          70          75          80
Ala Leu Ala Ala Lys Thr Ala Val Pro Pro Pro Ala Glu Gly Asp Glu
85          90          95
Val Ala Val Leu Arg Gly Ala Ala Ala Val Val Lys Asn Met Ser
100         105         110
Ala Ser Leu Glu Val Pro Thr Ala Thr Ser Val Arg Ala Val Pro Ala
115        120        125
Lys Leu Leu Ile Asp Asn Arg Ile Val Ile Asn Asn Gln Leu Lys Arg
130        135        140
Thr Arg Gly Gly Lys Ile Ser Phe Thr His Leu Leu Gly Tyr Ala Leu
145        150        155        160
Val Gln Ala Val Lys Lys Phe Pro Asn Met Asn Arg His Tyr Thr Glu
165        170        175
Val Asp Gly Lys Pro Thr Ala Val Thr Pro Ala His Thr Asn Leu Gly
180        185        190
Leu Ala Ile Asp Leu Gln Gly Lys Asp Gly Lys Arg Ser Leu Val Val
195        200        205
Ala Gly Ile Lys Arg Cys Glu Thr Met Arg Phe Ala Gln Phe Val Thr
210        215        220
Ala Tyr Glu Asp Ile Val Arg Arg Ala Arg Asp Gly Lys Leu Thr Thr
225        230        235        240
Glu Asp Phe Ala Gly Val Thr Ile Ser Leu Thr Asn Pro Gly Thr Ile
245        250        255
Gly Thr Val His Ser Val Pro Arg Leu Met Pro Gly Gln Gly Ala Ile
260        265        270
Ile Gly Val Gly Ala Met Glu Tyr Pro Ala Glu Phe Gln Gly Ala Ser
275        280        285
Glu Glu Arg Ile Ala Glu Leu Gly Ile Gly Lys Leu Ile Thr Leu Thr
290        295        300
Ser Thr Tyr Asp His Arg Ile Ile Gln Gly Ala Glu Ser Gly Asp Phe
305        310        315        320

```

-continued

Leu Arg Thr Ile His Glu Leu Leu Leu Ser Asp Gly Phe Trp Asp Glu
 325 330 335

Val Phe Arg Glu Leu Ser Ile Pro Tyr Leu Pro Val Arg Trp Ser Thr
 340 345 350

Asp Asn Pro Asp Ser Ile Val Asp Lys Asn Ala Arg Val Met Asn Leu
 355 360 365

Ile Ala Ala Tyr Arg Asn Arg Gly His Leu Met Ala Asp Thr Asp Pro
 370 375 380

Leu Arg Leu Asp Lys Ala Arg Phe Arg Ser His Pro Asp Leu Glu Val
 385 390 395 400

Leu Thr His Gly Leu Thr Leu Trp Asp Leu Asp Arg Val Phe Lys Val
 405 410 415

Asp Gly Phe Ala Gly Ala Gln Tyr Lys Lys Leu Arg Asp Val Leu Gly
 420 425 430

Leu Leu Arg Asp Ala Tyr Cys Arg His Ile Gly Val Glu Tyr Ala His
 435 440 445

Ile Leu Asp Pro Glu Gln Lys Glu Trp Leu Glu Gln Arg Val Glu Thr
 450 455 460

Lys His Val Lys Pro Thr Val Ala Gln Gln Lys Tyr Ile Leu Ser Lys
 465 470 475 480

Leu Asn Ala Ala Glu Ala Phe Glu Thr Phe Leu Gln Thr Lys Tyr Val
 485 490 495

Gly Gln Lys Arg Phe Ser Leu Glu Gly Ala Glu Ser Val Ile Pro Met
 500 505 510

Met Asp Ala Ala Ile Asp Gln Cys Ala Glu His Gly Leu Asp Glu Val
 515 520 525

Val Ile Gly Met Pro His Arg Gly Arg Leu Asn Val Leu Ala Asn Ile
 530 535 540

Val Gly Lys Pro Tyr Ser Gln Ile Phe Thr Glu Phe Glu Gly Asn Leu
 545 550 555 560

Asn Pro Ser Gln Ala His Gly Ser Gly Asp Val Lys Tyr His Leu Gly
 565 570 575

Ala Thr Gly Leu Tyr Leu Gln Met Phe Gly Asp Asn Asp Ile Gln Val
 580 585 590

Ser Leu Thr Ala Asn Pro Ser His Leu Glu Ala Val Asp Pro Val Leu
 595 600 605

Glu Gly Leu Val Arg Ala Lys Gln Asp Leu Leu Asp His Gly Ser Ile
 610 615 620

Asp Ser Asp Gly Gln Arg Ala Phe Ser Val Val Pro Leu Met Leu His
 625 630 635 640

Gly Asp Ala Ala Phe Ala Gly Gln Gly Val Val Ala Glu Thr Leu Asn
 645 650 655

Leu Ala Asn Leu Pro Gly Tyr Arg Val Gly Gly Thr Ile His Ile Ile
 660 665 670

Val Asn Asn Gln Ile Gly Phe Thr Thr Ala Pro Glu Tyr Ser Arg Ser
 675 680 685

Ser Glu Tyr Cys Thr Asp Val Ala Lys Met Ile Gly Ala Pro Ile Phe
 690 695 700

His Val Asn Gly Asp Asp Pro Glu Ala Cys Val Trp Val Ala Arg Leu
 705 710 715 720

Ala Val Asp Phe Arg Gln Arg Phe Lys Lys Asp Val Val Ile Asp Met

-continued

			725				730				735				
Leu	Cys	Tyr	Arg	Arg	Arg	Gly	His	Asn	Glu	Gly	Asp	Asp	Pro	Ser	Met
			740					745					750		
Thr	Asn	Pro	Tyr	Val	Tyr	Asp	Val	Val	Asp	Thr	Lys	Arg	Gly	Ala	Arg
		755					760					765			
Lys	Ser	Tyr	Thr	Glu	Ala	Leu	Ile	Gly	Arg	Gly	Asp	Ile	Ser	Met	Lys
	770					775					780				
Glu	Ala	Glu	Asp	Ala	Leu	Arg	Asp	Tyr	Gln	Gly	Gln	Leu	Glu	Arg	Val
785					790					795					800
Phe	Asn	Glu	Val	Arg	Glu	Leu	Glu	Lys	His	Gly	Val	Gln	Pro	Ser	Glu
				805					810						815
Ser	Val	Glu	Ser	Asp	Gln	Met	Ile	Pro	Ala	Gly	Leu	Ala	Thr	Ala	Val
			820					825					830		
Asp	Lys	Ser	Leu	Leu	Ala	Arg	Ile	Gly	Asp	Ala	Phe	Leu	Ala	Leu	Pro
		835					840					845			
Asn	Gly	Phe	Thr	Ala	His	Pro	Arg	Val	Gln	Pro	Val	Leu	Glu	Lys	Arg
850						855					860				
Arg	Glu	Met	Ala	Tyr	Glu	Gly	Lys	Ile	Asp	Trp	Ala	Phe	Gly	Glu	Leu
865					870					875					880
Leu	Ala	Leu	Gly	Ser	Leu	Val	Ala	Glu	Gly	Lys	Leu	Val	Arg	Leu	Ser
				885					890					895	
Gly	Gln	Asp	Ser	Arg	Arg	Gly	Thr	Phe	Ser	Gln	Arg	His	Ser	Val	Leu
			900					905					910		
Ile	Asp	Arg	His	Thr	Gly	Glu	Glu	Phe	Thr	Pro	Leu	Gln	Leu	Leu	Ala
		915					920						925		
Thr	Asn	Ser	Asp	Gly	Ser	Pro	Thr	Gly	Gly	Lys	Phe	Leu	Val	Tyr	Asp
	930					935					940				
Ser	Pro	Leu	Ser	Glu	Tyr	Ala	Ala	Val	Gly	Phe	Glu	Tyr	Gly	Tyr	Thr
945					950					955					960
Val	Gly	Asn	Pro	Asp	Ala	Val	Val	Leu	Trp	Glu	Ala	Gln	Phe	Gly	Asp
				965					970					975	
Phe	Val	Asn	Gly	Ala	Gln	Ser	Ile	Ile	Asp	Glu	Phe	Ile	Ser	Ser	Gly
		980						985					990		
Glu	Ala	Lys	Trp	Gly	Gln	Leu	Ser	Asn	Val	Val	Leu	Leu	Leu	Pro	His
		995					1000						1005		
Gly	His	Glu	Gly	Gln	Gly	Pro	Asp	His	Thr	Ser	Ala	Arg	Ile	Glu	
	1010					1015					1020				
Arg	Phe	Leu	Gln	Leu	Trp	Ala	Glu	Gly	Ser	Met	Thr	Ile	Ala	Met	
	1025					1030					1035				
Pro	Ser	Thr	Pro	Ser	Asn	Tyr	Phe	His	Leu	Leu	Arg	Arg	His	Ala	
	1040					1045					1050				
Leu	Asp	Gly	Ile	Gln	Arg	Pro	Leu	Ile	Val	Phe	Thr	Pro	Lys	Ser	
	1055					1060					1065				
Met	Leu	Arg	His	Lys	Ala	Ala	Val	Ser	Glu	Ile	Lys	Asp	Phe	Thr	
	1070					1075					1080				
Glu	Ile	Lys	Phe	Arg	Ser	Val	Leu	Glu	Glu	Pro	Thr	Tyr	Glu	Asp	
	1085					1090					1095				
Gly	Ile	Gly	Asp	Arg	Asn	Lys	Val	Ser	Arg	Ile	Leu	Leu	Thr	Ser	
	1100					1105					1110				
Gly	Lys	Leu	Tyr	Tyr	Glu	Leu	Ala	Ala	Arg	Lys	Ala	Lys	Asp	Asn	
	1115					1120					1125				

-continued

Arg Asn Asp Leu Ala Ile Val Arg Leu Glu Gln Leu Ala Pro Leu
 1130 1135 1140

Pro Arg Arg Arg Leu Arg Glu Thr Leu Asp Arg Tyr Glu Asn Val
 1145 1150 1155

Lys Glu Phe Phe Trp Val Gln Glu Glu Pro Ala Asn Gln Gly Ala
 1160 1165 1170

Trp Pro Arg Phe Gly Leu Glu Leu Pro Glu Leu Leu Pro Asp Lys
 1175 1180 1185

Leu Ala Gly Ile Lys Arg Ile Ser Arg Arg Ala Met Ser Ala Pro
 1190 1195 1200

Ser Ser Gly Ser Ser Lys Val His Ala Val Glu Gln Gln Glu Ile
 1205 1210 1215

Leu Asp Glu Ala Phe Gly
 1220

<210> SEQ ID NO 3

<211> LENGTH: 8621

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 p206 plasmid with codon optimized kgd gene nucleic
 acid sequence and softag

<400> SEQUENCE: 3

```

ggtagcggtg cttgggtcga tatcaaagtg catcacttct tcccgtatgc ccaactttgt      60
atagagagcc actgcgggat cgtcaccgta atctgcttgc acgtagatca cataagcacc      120
aagcgcggtg gcctcatgct tgaggagatt gatgagcgcg gtggcaatgc cctgcctccg      180
gtgctcgccg gagactgcca gatcatagat atagatctca ctacgcggct gctcaaactt      240
gggcagaacg taagccgcga gagcgccaac aaccgcttct tggtcgaagg cagcaagcgc      300
gatgaatgtc ttactacgga gcaagttccc gaggtaatcg gagtccggct gatggtggga      360
gtaggtggct acgtcaccga actcacgacc gaaaagatca agagcagccc gcatggattt      420
gacttggcca gggccgagcc tacatgtgcg aatgatgccc atacttgagc cacctaactt      480
tgttttaggg cgactgcctt gctgcgtaac atcgttctgt ctccataaca tcaaactcgc      540
accacggcgc taacgcgctt gctgcttggg tgcccggagg atagactgta caaaaaaaca      600
gtcataaaca gccatgaaaa ccgccactgc gccgttacca ccgctgcgct cggcgaaggt      660
tctggaccag ttgcgtgagc gcattttttt ttctcctcgc gcgcttacgc cccgcctcgc      720
cactcatcgc agtactgttg taattcatta agcattctgc cgacatggaa gccatcacag      780
acggcatgat gaacctgaat cggcagcggc atcagcacct tgtcgccttg cgtataatat      840
ttgccatag tgaaaacggg ggccaagaag ttgtccatat tggccacggt taaatcaaaa      900
ctggtgaaac tcaccaggg attggcgctg acgaaaaaca tattctcaat aaacccttta      960
gggaaatagg ccaggttttc accgtaacac gccacatctt gcgaatatat gtgtagaaac     1020
tgccggaaat cgtcgtggta ttcactccag agcgatgaaa acgtttcagt ttgctcatgg     1080
aaaacgggtg aacaagggtg aacactatcc catatcacca gctcacgctc tttcattgcc     1140
atacggaaact ccgatgagc atcatcagg cgggcaagaa tgtgaataaa ggccggataa     1200
aacttgctgt tatttttctt tacggtcttt aaaaaggccg taatatccag ctgaacggtc     1260
tggttatagg tacattgagc aactgactga aatgcctcaa aatggtcttt acgatgccat     1320

```

-continued

tgggatatat	caacggtggt	atatccagt	atTTTTTct	ccattTTTT	ttcctccttt	1380
agaaaaactc	atcgagcatc	aatgaaact	gcaatttatt	catatcagga	ttatcaatac	1440
catatTTTTg	aaaaagccgt	ttctgtaatg	aaggagaaaa	ctcaccgagg	cagttccata	1500
ggatggcaag	atcctggtat	cggctctgca	ttccgactcg	tccaacatca	atacaaccta	1560
ttaatttccc	ctcgtcaaaa	ataaggttat	caagtgagaa	atcaccatga	gtgacgactg	1620
aatccggtga	gaatggcaaa	agtttatgca	tttctttcca	gacttgttca	acaggccagc	1680
cattacgctc	gtcatcaaaa	tactcgcgat	caaccaaacc	gttattcatt	cgtgattgcg	1740
cctgagcgag	gcaaaatcgc	cgatcgctgt	taaaaggaca	attacaaaca	ggaatcgagt	1800
gcaaccggcg	caggaacact	gccagcgcat	caacaatatt	ttcacctgaa	tcaggatatt	1860
cttctaatac	ctggaacgct	gtttttccgg	ggatcgcagt	ggtgagtaac	catgcatcat	1920
caggagtacg	gataaaatgc	ttgatggctg	gaagtggcat	aaattccgtc	agccagttta	1980
gtctgaccat	ctcatctgta	acatcattgg	caacgctacc	tttgccatgt	ttcagaaaca	2040
actctggcgc	atcgggcttc	ccatacaagc	gatagattgt	cgcacctgat	tgccccacat	2100
tatcgcgagc	ccatttatac	ccatataaat	cagcatccat	gttgaattt	aatcgcggcc	2160
tcgacgtttc	ccgttgaata	tggtcattt	ttttttcctc	ctttaccaat	gcttaatcag	2220
tgaggcacct	atctcagcga	tctgtctatt	tcgttcatcc	atagttgcct	gactccccgt	2280
cgtgtagata	actacgatac	gggagggcct	accatctggc	cccagcgtg	cgatgatacc	2340
gcgagaacca	cgctcaccgg	ctccggattt	atcagcaata	aaccagccag	ccggaagggc	2400
cgagcgcaga	agtggctcctg	caactttatc	cgctccatc	cagtctatta	attggtgccg	2460
ggaagctaga	gtaagtagtt	cgccagttaa	tagtttgccg	aacgttggtg	ccatcgctac	2520
aggcatcgtg	gtgtcacgct	cgctgtttgg	tatggcttca	ttcagctccg	gttcccaacg	2580
atcaaggcga	gttacatgat	cccccatggt	gtgcaaaaa	gcggttagct	ccttcgggtcc	2640
tccgatcgtt	gtcagaagta	agttggccgc	agtgttatca	ctcatggta	tggcagcact	2700
gcataattct	cttactgtca	tgccatccgt	aagatgcttt	tctgtgactg	gtgagtactc	2760
aaccaagtca	ttctgagaat	agtgtatgcg	gcgaccgagt	tgctcttgcc	cggcgtcaat	2820
acgggataat	accgcgccac	atagcagaac	tttaaaagtg	ctcatcattg	gaaaacgttc	2880
ttcggggcga	aaactctcaa	ggatcttacc	gctgttgaga	tccagttcga	tgtaaccac	2940
tcgtgcaccc	aactgatctt	cagcatcttt	tactttcacc	agcgtttctg	ggtgagcaaa	3000
aacaggaagg	caaatgcccg	caaaaaagg	aataagggcg	acacggaaat	gttgaatact	3060
catattcttc	ctttttcaat	attattgaag	catttatcag	ggttattgtc	tcatgagcgg	3120
atacatattt	gaatgtattt	agaaaaataa	acaaataggg	gtcagtgtta	caaccaatta	3180
accaattctg	aacattatcg	cgagcccatt	tatacctgaa	tatggctcat	aacaccctt	3240
gtttgcctgg	cggcagtagc	gcggtggctc	cacctgaccc	catgccgaac	tcagaagtga	3300
aacgccgtag	cgccgatggt	agtgtgggga	ctccccatgc	gagagttagg	aactgccagg	3360
catcaataaa	aacgaaaggc	tcagtcgaaa	gactgggcct	ttcgcccggg	ctaattgagg	3420
ggtgtcgccc	ttttgacgga	tatcaagctt	ctattaaccg	aacgcttcgt	ccaagatttc	3480
ttgctgctcc	acggcatgaa	ccttcgagga	accgctgctc	ggcgcagaca	tcgctgcgacg	3540
ggagatacgc	ttgatgcccg	ccaacttgtc	cggcagcaac	tccggcaact	ccaaccgaa	3600
acgcggccag	gcccctggt	tcgctggttc	ctcctggacc	caaagaact	ctttgacatt	3660

-continued

ttcgtaacga tccagggttt cacgcagacg acgacgcggc aacggcgcca gttgctccag 3720
acgcacaatt gccagatcat tacggttgtc tttcgcttg cgtgcccga attcataata 3780
caacttaccg gaggtcaaca gaatacggct aactttattg cggtcgcca taccatcctc 3840
gtaggteggg tcctccagga cgctacggaa tttaatctcg gtaaaatcct taatctccga 3900
aacgcgcgcc ttgtggcgca gcatggattt cggggtaaac acgatcaacg gacgttggat 3960
gccgtccaac gcgtggcggc gcagcaagtg aaagtaattg gacgggggtg tcggcatcgc 4020
gatcgtcatg ctaccctcag cccacagctg caaaaagcgc tcaatgcgcg ccgacgtgtg 4080
gtccggggccc tgaccctcgt ggccgtgtgg cagcaacaga acaacgttgg acagctgacc 4140
ccatttcgct tcgcccagc tgatgaactc gtcaatgatc gactggggcac cattaacaaa 4200
gtcgccgaat tgcgcctccc acagcacaac cgcgtctgga ttgccaacgg tataaccgta 4260
ctcaaaaccc acagcccgct attccgacaa cggcgaatca taaaccagga acttaccacc 4320
ggtcgggctg ccgctcgtgt tagtcgccag cagctgcagc ggggtgaact cctcgccggt 4380
gtgacggctg atcagcacag aatgacgctg gctaaagggtg ccacgacgag agtcttgacc 4440
gctcagacgg accagcttgc cctcagcaac caggctgccc agcgcagca actcaccaaa 4500
cgcccagtea atcttaccct catacgccat ctacgacgc ttttcagca ccggtgaac 4560
gcgcgggtgt gccgtgaaac cgttaggcag cgcaggaag gcatcaccga tacgtgccag 4620
cagggatttg tcaacagcgg tcgccaggcc cgtgggatc atttggctcg actcgacgct 4680
ctccgacggt tggacgccgt gcttctccag ttcgcgcaact tcggtgaaca cacgctccag 4740
ttggccctgg taatcgcgca gcgcacctc cgcctcttcc atgctgatgt cgccacgacc 4800
gatcagagcc tcggtgtagg atttacgggc gccacgcttg gtgtccacga cgtcatagac 4860
atacggattg gtcatagatg gatcgtcacc ctattatga ccacgacgac gatagcacag 4920
catgtcaata acaacgtcct ttttaaagcg ttggcggagg tcaactgcca aacgcgcaac 4980
ccagacacac gcttctggat cgtcgccggt cacgtggaag attggcgcac cgatcatctt 5040
cgccacatcc gtgcagtact cgctcgaacg agagtattcc ggagcgggtg tgaagccgat 5100
ttggttgttc acaatgatgt gaatcgtacc acccacacga tagccaggca gatttgccag 5160
attcagcgtc tccgcaacaa cgcttgacc cgcgaaggcc gcatcaccgt gcagcatcaa 5220
cggaacgacg gagaatgcac gttggccgct ggagtcaatg gaaccgtggt ccagcaggtc 5280
ttgcttcgca cgcaccaaac cttccagcac tggatcgacc gcttccaaat gggacggatt 5340
tgccgtcagg gaaacctgaa tatcgttatc gccaaacatt tgcagataca gaccgctcgc 5400
accaggtgg tacttgacat cgccggaacc gtgagcctgg gacgggttca gattgccttc 5460
aaactccgta aagatttgcg aatacggttt gccacgatg ttcgccagga cattcaagcg 5520
accacggtgc ggcattgcaa tgaccacttc atccaaacca tgctcggcac attggtcaat 5580
cgcccgctcc atcattggaa taacagattc cgcaccctcc aggctaaaac gcttttggcc 5640
cacatacttg gtttgagga aggtttcaaa cgctccgct gcgctcagtt tcgacaagat 5700
gtacttttgt tgagcaacgg tcggtttgac gtgcttcgct tcgacacgct gctccagcca 5760
ctccttttgt tccgggtcca gaatgtgcgc gtactcaaca ccgatgtgac ggcagtacgc 5820
gtcgcgcagc aaaccagca cgtcacgcag ctttttgtat tgagcaccg cgaaaccgctc 5880
aaccttaaag acgcggtcca ggtcccacag agtcaggcca tgcgtcaaca cctccaaatc 5940

-continued

cggatgcgaa	cgaaagcgcg	ccttatccaa	gcgcaacggg	tcggtgtccg	ccatcagatg	6000
gccgcggttg	cgataggccg	cgatcagggt	catcacacgt	gcgttcttgt	caacgatcga	6060
gtccggatta	tcggtgctcc	aacgcactgg	caggtacggg	atgctcagct	cgcggaagac	6120
ctcatcccag	aagccatcag	acaacagcag	ttcatgaatg	gtacgcagga	agtcaccgct	6180
ttccgcacct	tgaatgatac	ggtggctgta	ggtagaagtc	agggtaatca	gtttgccaat	6240
accagctca	gcgatgcgtt	cctcggacgc	gccttggaac	tccgccggat	attccatcgc	6300
accgacaccg	atgatagcac	cttgacctgg	catcaggcgt	ggcacagagt	gcaccgtgcc	6360
aatcgtgccc	ggattcgtca	gcgaaatcgt	aacgccagcg	aagtcctcgg	tggtcagttt	6420
accatcacga	gcacgacgga	cgatgtcctc	gtacgcggta	acgaactgcg	cgaagcgcac	6480
cgctcgcgaa	cgtttgatac	cggccaccac	cagagagcgc	ttgccatctt	taccttgacg	6540
gtcaatagcc	agaccagat	tgggtgtgccc	aggagtaacc	gccgtcggct	taccgtccac	6600
ctccgtgtag	tggcgattca	tattcgggaa	cttcttaacc	gcctgaacca	gagcataacc	6660
cagcaaatgg	gtaaagctga	ttttaccacc	acgcgtgctg	ttcaactgat	tattgatcac	6720
gatacgatta	tcgatcaaca	atttcgctgg	cacagcacgc	accgaggtcg	ccgtaggcac	6780
ttccagcgac	gcgctcatgt	tcttcacgac	agcagccgcc	gcaccacgca	ggacagcaac	6840
ttcatcgect	tcggctggcg	gaggcaccgc	ggtcttggcg	gccagagccg	ccacgacacc	6900
gttgcccget	gcggccgtgt	cggccggttt	cggcggcget	tgcggagccg	ccgctgccgc	6960
acgctcagcg	accaaagggc	tggtcacacg	agtcggctca	gcagccggtt	gggaagtccg	7020
ctccgggcta	tagtccacca	gaaactcatg	ccagcttggg	tcaacggaag	acggatcatc	7080
acgaaattta	cgatacatac	caacacgaga	cgggtcctga	gtcaccatgg	atatatctcc	7140
ttcttaaaga	attcgatatc	tcagcgacaa	ggcgacaca	aaatttattc	taaatagcata	7200
ataaatactg	ataacatctt	atagtttgta	ttatatcttg	tattatcgtt	gacatgtata	7260
atcttgatat	caaaaactga	ttttcccttt	attatcttcg	agatttattt	tcttaattct	7320
ctttaacaaa	ctagaaatat	tgtatataca	aaaaatcata	aataatagat	gaatagttta	7380
attataggtg	ttcatcaatc	gaaaaagcaa	cgtatcttat	ttaaagtgcg	ttgctttttt	7440
ctcatttata	aggttaaata	attctcatat	atcaagcaaa	gtgacaggcg	cccttaaata	7500
ttctgacaaa	tgctctttcc	ctaaactccc	cccataaaaa	aaccgcgccg	agcggggttt	7560
tacgttattt	gcgattaac	gattactcgt	tatcagaacc	gccagggggg	cccagactta	7620
agactggccg	tcgttttaca	acacagaaaag	agttttaga	aacgcaaaaa	ggccatccgt	7680
caggggcctt	ctgcttagtt	tgatgcctgg	cagttcccta	ctctcgcctt	ccgcttcctc	7740
gctcactgac	tcgctgcgct	cggctcgttcg	gctgcggcga	gcggtatcag	ctcactcaaa	7800
ggcggtaata	cggttatcca	cagaatcagg	ggataacgca	ggaaagaaca	tgtgagcaaa	7860
aggccagcaa	aaggccagga	accgtaaaaa	ggccgcgttg	ctggcgtttt	tccataggct	7920
ccgccccctt	gacgagcatc	acaaaaatcg	acgctcaagt	cagaggtggc	gaaaccgcac	7980
aggactataa	agataccagg	cgtttccccc	tggaagctcc	ctcgtgcgct	ctcctgttcc	8040
gaccctgccg	cttaccggat	acctgtccgc	ctttctccct	tcgggaagcg	tggcgctttc	8100
tcatagctca	cgctgtaggt	atctcagttc	ggtgtaggtc	gttcgctcca	agctgggctg	8160
tgtgcacgaa	ccccccgttc	agcccagaccg	ctgcgcctta	tccggtaact	atcgtcttga	8220
gtccaaccgc	gtaagacacg	acttatcgcc	actggcagca	gccactggta	acaggattag	8280

-continued

```

cagagcagagg tatgtaggcg gtgctacaga gttcttgaag tgggtgggcta actacggcta 8340
cactagaaga acagtatttg gtatctgccc tctgctgaag ccagttacct tcggaaaaag 8400
agttggtagc tcttgatccg gcaaaaaaac caccgctggt agcgggtggt tttttgttg 8460
caagcagcag attacgcgca gaaaaaaagg atctcaagaa gatcctttga tcttttctac 8520
ggggtctgac gctcagtga acgacgcgcg cgtaactcac gttaaggat tttggtcatg 8580
agcttgcgcc gtccegtcaa gtcagcgtaa tgctctgctt a 8621

```

```

<210> SEQ ID NO 4
<211> LENGTH: 39
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Forward primer for PCR amplification of codon optimized kgd
nucleic acid sequence

```

```

<400> SEQUENCE: 4

```

```

tttttttgta taccatggat cgtaaatttc gtgatgatc 39

```

```

<210> SEQ ID NO 5
<211> LENGTH: 42
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Reverse primer for PCR amplification of codon optimized kgd
nucleic acid sequence

```

```

<400> SEQUENCE: 5

```

```

cccggtgaga tctagatccg aacgcttcgt ccaagatttc tt 42

```

```

<210> SEQ ID NO 6
<211> LENGTH: 47
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Forward primer for PCR amplification of pKK223-3 template

```

```

<400> SEQUENCE: 6

```

```

cggatctaga tctcaccatc accaccatta gtcgacctgc agccaag 47

```

```

<210> SEQ ID NO 7
<211> LENGTH: 44
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Reverse primer for PCR amplification of pKK223-3 template

```

```

<400> SEQUENCE: 7

```

```

tgagatctag atccggttatg tcccatgggt ctgtttcctg tgtg 44

```

```

<210> SEQ ID NO 8
<211> LENGTH: 4614
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
pKK223-ct-his vector sequence

```

```

<400> SEQUENCE: 8

```


-continued

ttgacaatta atcatcggct cgtataatgt gtggaattgt gagcggataa caatttcaca	60
caggaaacag aaccatggga cataacggat ctagatctca ccatcaccac cattagtcga	120
cctgcagcca agcttggctg ttttggcgga tgagagaaga ttttcagcct gatacagatt	180
aaatcagaac gcagaagcgg tctgataaaa cagaatattgc ctggcggcag tagcgcggtg	240
gtcccacctg accccatgcc gaactcagaa gtgaaacgcc gtagcgcga tggtagtgtg	300
gggtctcccc atgcgagagt agggaactgc caggcatcaa ataaaacgaa aggctcagtc	360
gaaagactgg gcctttcgtt ttatctgttg tttgtcggtg aacgctctcc tgagtaggac	420
aaatccgccg ggagcggatt tgaacgttgc gaagcaacgg cccggagggt ggcgggcagg	480
acgcccgcca taaactgcc ggcatacaat taagcagaag gccatcctga cggatggcct	540
ttttgcgttt ctacaaactc ttttgtttat ttttctaaat acattcaaat atgtatccgc	600
tcatgagaca ataaccctga taaatgcttc aataatattg aaaaaggaag agtatgagta	660
ttcaacattt ccgtgtcgc cttattccct tttttgcggc attttgcctt cctgtttttg	720
ctcaccaga aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtgg	780
gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac	840
gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccgtggtg	900
acgccgggca agagcaactc ggtcgcgca tacaacttcc tcagaatgac ttggttgagt	960
actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg	1020
ctgccataac catgagtgat aacactgagg ccaacttact tctgacaacg atcggaggac	1080
cgaaggagct aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt	1140
gggaaccgga gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgctgtagc	1200
aatggcaaca acggtgcgca aactattaac tggcgaacta cttactctag cttcccggca	1260
acaattaata gactggatgg agggcgataa agttgcagga ccacttctgc gctcggccct	1320
tccggctggc tggtttattg ctgataaatc tggagccggg gagcgtgggt ctgcgggtat	1380
cattgcagca ctggggccag atggtaagcc ctcccgtatc gtagttatct acacgacggg	1440
gagtcaggca actatggatg aacgaaatag acagatcgcg gagatagggt cctcactgat	1500
taagcattgg taactgtcag accaagttta ctcatatata ctttagattg atttaaaact	1560
tcatttttaa tttaaaagga tctaggtgaa gatccttttt gataatctca tgaccaaaat	1620
cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga tcaaaggatc	1680
ttcttgagat cctttttttc tgccgcgtaat ctgctgcttg caaacaaaaa aaccaccgct	1740
accagcggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga aggtaactgg	1800
cttcagcaga gcgcagatac caaatactgt ccttctagtg tagccgtagt taggccacca	1860
cttcaagaac tctgtagcac cgctacata cctcgctctg ctaatcctgt taccagtggc	1920
tgctgccagt ggcgataagt cgtgtcttac cgggttgac tcaagacgat agttaccgga	1980
taaggcgcag cggcgggct gaacgggggg ttcgtgcaca cagcccagct tggagcgaac	2040
gacctacacc gaactgagat acctacagcg tgagcattga gaaagcgcca cgcttcccga	2100
aggagaaaag gcggacaggt atccggtaag cggcagggtc ggaacaggag agcgcacgag	2160
ggagcttcca gggggaaacg cctggatctt ttatagtcct gtcgggtttc gccacctctg	2220
acttgagcgt cgatttttgt gatgctcgtc agggggggcg agcctatgga aaaacgccag	2280

-continued

caacgcggcc	tttttacggt	tcttggcctt	ttgctggcct	tttgctcaca	tgttctttcc	2340
tgcgttatcc	cctgattctg	tggataaccg	tattaccgcc	tttgagtgag	ctgataaccg	2400
tcgcccagc	cgaacgaccg	agcgcagcga	gtcagtgagc	gaggaagcgg	aagagcgcct	2460
gatgcggtat	tttctcctta	cgcactctgtg	cggatattca	caccgcatat	ggtgcactct	2520
cagtacaatc	tgctctgatg	cgcgatagtt	aagccagtat	acactccgct	atcgctacgt	2580
gactgggtca	tggctgcgcc	cgcacaccgc	ccaacaccgc	ctgacgcgcc	ctgacgggct	2640
tgtctgctcc	cggcatccgc	ttacagacaa	gctgtgaccg	tctccgggag	ctgcatgtgt	2700
cagaggtttt	caccgtcatc	accgaaacgc	gcgaggcagc	tgcggtaaag	ctcatcagcg	2760
tggtcgtgaa	gcgattcaca	gatgtctgcc	tgttcatccg	cgtccagctc	gttgagtttc	2820
tccagaagcg	ttaatgtctg	gcttctgata	aagcgggcca	tgtaagggc	ggttttttcc	2880
tgtttggtca	ctgatgcctc	cgtgtaaggg	ggatttctgt	tcatgggggt	aatgataaccg	2940
atgaaacgag	agaggatgct	cacgatacgg	gttactgatg	atgaacatgc	ccggttactg	3000
gaacgttggt	agggtaaaca	actggcggta	tggatgcggc	gggaccagag	aaaaatcact	3060
caggttcaat	gccagcgtt	cgttaataca	gatgtagggt	ttccacaggg	tagccagcag	3120
catcctgcga	tgcatatccg	gaacataatg	gtgcagggcg	ctgacttccg	cgtttccaga	3180
ctttacgaaa	cacggaaacc	gaagaccatt	catgttggtg	ctcaggtcgc	agacgttttg	3240
cagcagcagt	cgcttcacgt	tcgctcgcgt	atcggtgatt	cattctgcta	accagtaagg	3300
caaccccgcc	agcctagccg	ggtcctcaac	gacaggagca	cgatcatgcg	cacccgtggc	3360
caggacccaa	cgctgcccga	gatgcgccgc	gtgcggctgc	tggagatggc	ggacgcgatg	3420
gatatggtct	gccaaggggt	ggtttgcgca	ttcacagttc	tccgcaagaa	ttgattggct	3480
ccaattcttg	gagtggtgaa	tccgttagcg	aggtgccgcc	ggcttccatt	caggtcgagg	3540
tggcccggct	ccatgcaccg	cgacgcaacg	cggggaggca	gacaaggat	agggcggcgc	3600
ctacaatcca	tgccaaccgc	ttccatgtgc	tcgccgaggc	ggcataaatc	gccgtgacga	3660
tcagcggtec	agtgatcgaa	gttaggctgg	taagagccgc	gagcgatcct	tgaagctgtc	3720
cctgatggtc	gtcatctacc	tgcttgaca	gcatggcctg	caacgcgggc	atcccgatgc	3780
cgccggaagc	gagaagaatc	ataatgggga	aggccatcca	gcctcgcgtc	gcgaacgcca	3840
gcaagacgta	gcccagcgcg	tcggccgcca	tgccggcgat	aatggcctgc	ttctcgccga	3900
aacgtttggt	ggcgggacca	gtgacgaagg	cttgagcagc	ggcgtgcaag	attccgaata	3960
ccgcaagcga	caggccgatc	atcgtcgcgc	tccagcgaaa	gcggtcctcg	ccgaaaatga	4020
cccagagcgc	tgccggcacc	tgtcctacga	gttgcgatg	aaagaagaca	gtcataagtg	4080
cggcgacgat	agtcatgccc	cgcgcccacc	ggaaggagct	gactgggttg	aaggctctca	4140
agggcatcgg	tcgacgctct	cccttatgcg	actcctgcat	taggaagcag	cccagtagta	4200
ggttgaggcc	gttgagcacc	gccgcgcgca	ggaatgggtg	atgcaaggag	atggcgccca	4260
acagtcccc	ggccacgggg	cctgccacca	taccacgcc	gaaacaagcg	ctcatgagcc	4320
cgaagtggcg	agcccgatct	tcccacgcg	tgatgtcgcc	gatataggcg	ccagcaaccg	4380
cacctgtggc	gccgggtgatg	cgggccacga	tgctccggc	gtagaggatc	cgggcttatc	4440
gactgcacgg	tgaccaatg	cttctggcgt	caggcagcca	tcggaagctg	tggatggct	4500
gtgcaggtcg	taaatcactg	cataattcgt	gtcgtcaag	gcgcactccc	gttctggata	4560
atgtttttg	cgccgacatc	ataacggttc	tggcaaatat	tctgaaatga	gctg	4614

-continued

<210> SEQ ID NO 9
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Histidine tag peptide sequence

<400> SEQUENCE: 9

Ser Arg Ser His His His His His
1 5

<210> SEQ ID NO 10
<211> LENGTH: 8241
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
pKK223-kgd plasmid with codon optimized kgd gene nucleic acid
sequence

<400> SEQUENCE: 10

ttgacaatta atcatcggct cgtataatgt gtggaattgt gagcggataa caatttcaca 60
caggaaacag aaccatggat cgtaaatttc gtgatgatcc gtcttccggt gacccaagct 120
ggcatgagtt tctggtggac tatagcccgg agccgacttc ccaaccggct gctgagccga 180
ctcgtgtgac cagccctttg gtcgctgagc gtgcggcagc ggcggtccg caagcgccgc 240
cgaaaccggc cgacacggcc gcagcgggca acggtgtcgt ggcggtctcg gccgccaaga 300
ccgcggtgcc tccgccagcc gaaggcagatg aagttgctgt cctgcgtggt gcggcggtcg 360
ctgtcgtgaa gaacatgagc gcgtcgtgag aagtgcctac ggcgacctcg gtgcgtgctg 420
tgccagcgaa attggtgatc gataatcgta tcgtgatcaa taatcagttg aaacgcacgc 480
gtggtggtaa aatcagcttt acccatttgc tgggttatgc tctggttcag gcggttaaga 540
agttcccgaa tatgaatcgc cactacacgg aggtggacgg taagccgacg gcggttactc 600
ctgcgcacac caatctgggt ctggctattg acctgcaagg taaagatggc aagcgctctc 660
tggtggtggc cggatcaaaa cgttgcgaga cgatgcgctt cgcgcagttc gttaccgctg 720
acgaggacat cgtccgctgt gctcgtgatg gtaaaactgac caccgaggac ttcgctggcg 780
ttacgatttc gctgacgaat ccgggacagc ttggcacggt gcaactctgt ccacgcctga 840
tgccaggtca aggtgctatc atcgggtgctg gtgcgatgga atatccggcg gagttccaag 900
gcgcgtccga ggaacgcacg gctgagctgg gtattggcaa actgattacc ctgacttcta 960
cctacgacca ccgtatcatt caagggtcgg aaagcgggtga cttcctgcgt accattcatg 1020
aactgctggt gtctgatggc ttctgggatg aggtcttccg cgagctgagc atcccgtacc 1080
tgccagtgcg ttggagcacc gataatccgg actcgatcgt tgacaagaac gcacgtgtga 1140
tgaacctgat cgcggcctat cgcaaccgcg gccatctgat ggcgacacc gaccggttgc 1200
gcttgataaa ggccgccttt cgttcgcatc cggatttggg ggtggtgacg catggcctga 1260
ctctgtggga cctggaccgc gtctttaaag ttgacgggtt cgcggtgct caatacaaaa 1320
agctgcgtga cgtgctgggt ttgctgcgag acgcgtactg ccgtcacatc ggtggtgagt 1380
acgcgcacat tctggaccgg gaacaaaagg agtggtgga gcagcgtgtc gagacgaagc 1440
acgtcaaacc gaccgttgc caacaaaagt acatcttgtc gaaactgaac gcagcggagg 1500

-continued

cgtttgaaac	cttcctgcaa	accaagtatg	tgggcacaaa	gcgttttagc	ctggaggggtg	1560
cggaatctgt	tattccaatg	atggacgegg	cgattgacca	atgtgccgag	catggtttgg	1620
atgaagtggg	cattggcatg	cgcacccgtg	gtcgcttgaa	tgtcctggcg	aacatcgtgg	1680
gcaaaccgta	ttcgaaaatc	tttacggagt	ttgaaggcaa	tctgaaccgg	tcccaggctc	1740
acggttccgg	cgatgtcaag	taccacctgg	gtgcgacggg	tctgtatctg	caaatgtttg	1800
gcgataacga	tattcaggtt	tocctgacgg	caaatccgtc	ccatttgga	gcggtcgatc	1860
cagtgtgga	aggtttgggtg	cgtgcgaagc	aagacctgct	ggaccacggg	tccattgact	1920
ccgacggcca	acgtgcattc	tccgtcgttc	cgttgatgct	gcacgggat	gcggccttcg	1980
cggttcaagg	cgttgttgcg	gagacgctga	atctggcaaa	tctgcctggc	tatcgtgtgg	2040
gtggtacgat	tcacatcatt	gtgaacaacc	aaatcggctt	caccaccgct	ccggaatact	2100
ctcgttcgag	cgagtactgc	acggatgtgg	cgaagatgat	cggtgcgcca	atcttccacg	2160
tgaacggcga	cgatccagaa	gcgtgtgtct	gggttgcgcg	tttggcagtt	gacttccgcc	2220
aacgctttaa	aaaggacggt	gttattgaca	tgtgtgtgta	tcgtcgtcgt	ggtcataatg	2280
agggtgacga	tccatctatg	accaatccgt	atgtctatga	cgctcgtggc	accaagcgtg	2340
gcgcccgtaa	atcctacacc	gaggctctga	tcggtcgtgg	cgacatcagc	atgaaagagg	2400
cggaggatgc	gctgcgcat	taccagggcc	aactggagcg	tgtgttcaac	gaagtgcgcg	2460
aactggagaa	gcacggcgtc	caaccgtcgg	agagcgtcga	gtcggaccaa	atgatcccag	2520
cgggcctggc	gaccgctggt	gacaaatccc	tgtctggcag	tatcgggtgat	gccttccctgg	2580
cgctgcctaa	cggtttcacg	gcacacccgc	gcgttcagcc	gggtcgtgaa	aagcgtcgtg	2640
agatggcgta	tgagggttaag	attgactggg	cgtttgggtg	gttgctggcg	ctgggcagcc	2700
tggttgctga	gggcaagctg	gtccgtctga	gcggtcaaga	ctctcgtcgt	ggcaccttta	2760
gccagcgtca	ttctgtgctg	atcgaccgtc	acaccggcga	ggagttcacc	ccgctgcagc	2820
tgtggcgac	taacagcgac	ggcagcccga	ccggtggtaa	gttcctgggt	tatgattcgc	2880
cgttgtcgga	atacgcggct	gtgggttttg	agtacggtta	taccgttggc	aatccagacg	2940
cggttgtgct	gtgggaggcg	caattcggcg	actttgttaa	tgggtgccag	tcgatcattg	3000
acgagttcat	cagctcgggc	gaagcgaat	gggtcagct	gtccaacgtt	gttctgttgc	3060
tgccacacgg	ccacgagggt	cagggcccgg	accacacgtc	ggcgcgcatt	gagcgtttt	3120
tgacgctgtg	ggctgagggt	agcatgacga	tcgcgatgcc	gagcaccccg	tccaattact	3180
ttcaactgct	gcgcccacc	gcgttggacg	gcatccaacg	tccgttgatc	gtgtttacc	3240
cgaaatccat	gctgcgccac	aaggcggcgg	tttcggagat	taaggatttt	accgagatta	3300
aattccgtag	cgctcgggag	gaaccgacct	acgaggatgg	tatcggcgac	cgcaataaag	3360
ttagccgtat	tctgttgacc	tccgtaagt	tgtattatga	attggcggca	cgcaaggcga	3420
aagacaaccg	taatgatctg	gcaattgtgc	gtctggagca	actggcgccg	ttgccgcgtc	3480
gtcgtctgcg	tgaaccctg	gatcgttacg	aaaatgtcaa	agagttcttt	tgggtccagg	3540
aggaaccagc	gaaccagggc	gcctggccgc	gttccggttt	ggagttgccg	gagttgctgc	3600
cggacaagtt	ggcgggcatc	aagcgtatct	cccgtcgcgc	gatgtctgcg	ccgagcagcg	3660
gttcctcgaa	ggttcatgcc	gtggagcagc	aagaaatctt	ggacgaagcg	ttcggatcta	3720
gatctcacca	tcaccaccat	tagtcgacct	gcagccaagc	ttggctgttt	tggcggatga	3780
gagaagattt	tcagcctgat	acagatataa	tcagaacgca	gaagcgtct	gataaaacag	3840

-continued

aatttgectg gcggcagtag cgcgggtggc ccacctgacc ccatgccgaa ctcagaagtg 3900
aaacgccgta gcgccgatgg tagtgtgggg tctccccatg cgagagtagg gaactgccag 3960
gcatcaaata aaacgaaagg ctcagtcgaa agactgggccc tttcgtttta tctggtgttt 4020
gtcggtgaac gctctcctga gtaggacaaa tccgcgggga gcggtattga acgttgcgaa 4080
gcaacggccc ggagggtggc gggcaggacg cccgccataa actgccaggc atcaaattaa 4140
gcagaaggcc atcctgacgg atggcctttt tgcgtttcta caaactcttt tgtttatttt 4200
tctaaataca ttcaaatatg tatccgctca tgagacaata accctgataa atgcttcaat 4260
aatattgaaa aaggaagagt atgagtattc aacatttccg tgtcgccctt attccctttt 4320
ttgcggcatt ttgccttctt gtttttgctc acccagaaac gctggtgaaa gtaaaagatg 4380
ctgaagatca gttgggtgca cgagtggggtt acatcgaact ggatctcaac agcggtaaga 4440
tccttgagag ttttcgcccc gaagaacggt ttccaatgat gagcactttt aaagttctgc 4500
tatgtggcgc ggtattatcc cgtggtgacg ccgggcaaga gcaactcggc cgccgcatac 4560
actattctca gaatgacttg gttgagtact caccagtcac agaaaagcat cttacggatg 4620
gcatgacagt aagagaatta tgcagtgctg ccataacatc gatgataac actgcccgca 4680
acttacttct gacaacgatc ggaggaccga aggagctaac cgcttttttg cacaaactgg 4740
gggatcatgt aactcgcctt gatcgttggg aaccggagct gaatgaagcc ataccaaacg 4800
acgagcgtga caccacgatg ctgtagcaat ggcaacaacg ttgcgcaaac tattaactgg 4860
cgaactactt actctagctt cccggcaaca attaatagac tggatggagg cggataaagt 4920
tgcaggacca cttctgcgct cggcccttcc ggctggctgg tttattgctg ataaatctgg 4980
agccggtgag cgtgggtctc gcggtatcat tgcagcactg gggccagatg gtaagccctc 5040
ccgtatcgta gttatctaca cgacggggag tcaggcaact atggatgaac gaaatagaca 5100
gatcgcgtgag ataggtgcct cactgattaa gcattggtaa ctgtcagacc aagtttactc 5160
atatatactt tagattgatt taaaacttca tttttaattt aaaaggatct aggtgaagat 5220
cctttttgat aatctcatga ccaaaatccc ttaacgtgag ttttcgctcc actgagcgtc 5280
agaccccgta gaaaagatca aaggatcttc ttgagatcct ttttttctgc gcgtaatctg 5340
ctgcttgcaa acaaaaaaac caccgctacc agcgggtggtt tgtttgccgg atcaagagct 5400
accaactctt tttccgaagg taactggctt cagcagagcg cagataccaa atactgtcct 5460
tctagtgtag ccgtagttag gccaccactt caagaactct gtagcaccgc ctacatacct 5520
cgctctgcta atcctgttac cagtggctgc tgccagtggc gataagtcgt gtcttaccgg 5580
gttgactca agacgatagt taccggataa ggcgcagcgg tcgggctgaa cgggggggttc 5640
gtgcacacag cccagcttgg agcgaacgac ctacaccgaa ctgagatacc tacagcgtga 5700
gcatthagaa agcgcacgc tccccgaagg gagaaaggcg gacaggtatc cggtaagcgg 5760
cagggtcgga acaggagagc gcacgagggg gcttccaggg ggaaacgcct ggtatcttta 5820
tagtctgtc gggtttcgcc acctctgact tgagcgtcga tttttgtgat gctcgtcagg 5880
ggggcggagc ctatggaaaa acgccagcaa cgcggccttt ttacggttcc tggccttttg 5940
ctggcctttt gctcacatgt tctttcctgc gttatccctt gattctgtgg ataaccgtat 6000
taccgccttt gagtgagctg ataccgctcg ccgcagccga acgaccgagc gcagcagatc 6060
agtgagcgag gaagcggag agcgcctgat gcggtatttt ctccttacgc atctgtgagg 6120

-continued

tatttcacac	cgcatatggt	gcactctcag	tacaatctgc	tctgatgccg	catagttaag	6180
ccagtataca	ctecgctatc	gctacgtgac	tgggtcatgg	ctgcgccccg	acacccgcca	6240
acacccgctg	acgcgccttg	acgggcttgt	ctgctcccgg	catccgctta	cagacaagct	6300
gtgaccgtct	ccgggagctg	catgtgtcag	aggttttcac	cgatcatcacc	gaaacgcgcg	6360
aggcagctgc	ggtaaagctc	atcagcgtgg	tcgtgaagcg	attcacagat	gtctgcctgt	6420
tcatccgcgt	ccagctcgtt	gagtttctcc	agaagcgtaa	atgtctggct	tctgataaag	6480
cgggccatgt	taagggcggt	tttttctgt	ttggtcactg	atgcctccgt	gtaaggggga	6540
tttctgttca	tgggggtaat	gataccgatg	aaacgagaga	ggatgctcac	gatacggggt	6600
actgatgatg	aacatgcccc	gttactggaa	cgttgtgagg	gtaaacaact	ggcggtatgg	6660
atgcggcggg	accagagaaa	aatcactcag	ggtcaatgcc	agcgcttcgt	taatacagat	6720
gtaggtgttc	cacagggtag	ccagcagcat	cctgcgatgc	agatccggaa	cataatgggtg	6780
cagggcgctg	acttccgcgt	ttccagactt	tacgaaacac	ggaaaccgaa	gaccattcat	6840
gttgttgctc	aggtegcaga	cgttttgacg	cagcagtcgc	ttcacgttcg	ctcgcgtatc	6900
ggtgattcat	tctgctaacc	agtaaggcaa	ccccgccagc	ctagccgggt	cctcaacgac	6960
aggagcacga	tcatgcgcac	ccgtggccag	gacccaacgc	tgccccgagat	gcgcccgcgtg	7020
cggctgctgg	agatggcgga	cgcgatggat	atgttctgcc	aagggttggg	ttgcgcattc	7080
acagttctcc	gcaagaattg	attggctcca	attcttgag	tggatgaatc	gttagcgagg	7140
tgccgcggc	ttccattcag	gtcaggtg	ccccgctcca	tgcaccgca	cgcaacgcgg	7200
ggaggcagac	aaggtatagg	gcggcgccca	caatccatgc	caacccgttc	catgtgctcg	7260
ccgaggcggc	ataaatcgcc	gtgacgatca	gcggtccagt	gatcgaagtt	aggctggtaa	7320
gagccgcgag	cgatccttga	agctgtccct	gatggctg	atctacctgc	ctggacagca	7380
tggcctgcaa	cgcgggcac	ccgatgccgc	cggaagcgag	aagaatcata	atggggaagg	7440
ccatccagcc	tcgcgtcgcg	aacgccagca	agacgtagcc	cagcgcgtcg	gccgccatgc	7500
cggcgataat	ggcctgcttc	tcgccgaaac	gtttggtggc	gggaccagtg	acgaaggctt	7560
gagcgagggc	gtgcaagatt	ccgaataacc	caagcgacag	gccgatcatc	gtcgcgtccc	7620
agcgaaagcg	gtcctcgccg	aaaatgaccc	agagcgctgc	cggcacctgt	cctacgagtt	7680
gcatgataaa	gaagacagtc	ataagtgcgg	cgacgatagt	catgccccgc	gcccaccgga	7740
aggagctgac	tgggttgaag	gctctcaagg	gcatcggtcg	acgctctccc	ttatgcgact	7800
cctgcattag	gaagcagccc	agtagtaggt	tgaggccggt	gagcaccgcc	gccgcaagga	7860
atggtgcatg	caaggagatg	gcgcccaaca	gtccccggc	cacggggcct	gccaccatac	7920
ccacgcgaa	acaagcgtc	atgagccccg	agtggcgagc	ccgatcttcc	ccatcgggtga	7980
tgtcggcgat	ataggcgcca	gcaaccgcac	ctgtggcgcc	ggtgatgccg	gccacgatgc	8040
gtccggcgta	gaggatccgg	gcttatcgac	tgcacgggtg	accaatgctt	ctggcgtcag	8100
gcagccatcg	gaagctgtgg	tatggctgtg	caggtcgtaa	atcactgcat	aattcgtgtc	8160
gctcaaggcg	cactcccgtt	ctggataatg	ttttttgcgc	cgacatcata	acggttctgg	8220
caaatattct	gaaatgagct	g				8241

<210> SEQ ID NO 11

<211> LENGTH: 3716

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Codon-optimized mcr nucleic acid sequence

<400> SEQUENCE: 11

```

gatatcgaat tccgctagca ggagctaagg aagctaaaat gtccggtacg ggtcgtttgg      60
ctggtaaaat tgcattgatc accggtgggtg ctggtaacat tggttccgag ctgacccgcc      120
gttttctggc cgaggggtgcg acggttatta tcagcggccg taaccgtgcg aagctgaccg      180
cgctggccga gcgcatgcaa gccgaggccg gcgtgccggc caagcgcatt gatttggagg      240
tgatggatgg ttccgacct gtggctgtcc gtgccggtat cgaggcaatc gtcgctcgcc      300
acggtcagat tgacattctg gttaacaacg cgggctccgc cggtgcccaa cgtcgtttgg      360
cggaaattcc gctgacggag gcagaattgg gtccgggtgc ggaggagact ttgcacgctt      420
cgatcgcgaa tctgttgggc atgggttggc acctgatgcg tattgeggct ccgcacatgc      480
cagttggctc cgcagttatc aacgtttcga ctattttctc gcgcgcagag tactatggtc      540
gcatccgta  cgttaccccg aaggcagcgc tgaacgcttt gtcccagctg gctgcccgcg      600
agctgggcgc tcgtggcatc cgcgttaaca ctattttccc aggtcctatt gagtccgacc      660
gcatccgtac cgtgtttcaa cgtatggatc aactgaaggg tcgcccggag ggcgacaccg      720
cccatcactt tttgaacacc atgcgcctgt gccgcgcaa cgaccaaggc gctttggaac      780
gccgctttcc gtccgttggc gatggtgctg atgcggctgt gtttctggct tctgctgaga      840
gcgcggcact gtcgggtgag acgattgagg tcacccacgg tatggaactg ccggcgtgta      900
gcgaaacctc cttgttggcg cgtaccgatc tgcgtaccat cgacgcgagc ggtcgcacta      960
ccctgatttg cgctggcgat caaattgaag aagttagtgc cctgacgggc atgctgctga     1020
cgtgcggtag cgaagtgatt atcggcttcc gttctgcggc tgccctggcg caatttgagc     1080
aggcagtgaa tgaatctcgc cgtctggcag gtgcggatct caccccgccg atcgttttgc     1140
cgttggaacc acgtgaccgg gccaccattg atgcggtttt cgattggggc gcaggcgaga     1200
atacgggtgg catccatgcg gcggtcattc tgccggcaac ctcccacgaa ccggctccgt     1260
gcgtgattga agtcgatgac gaacgcgtcc tgaatttccg ggccgatgaa attaccggca     1320
ccatcgttat tgcgagccgt ttggcgcgct attggcaatc ccaacgcctg accccgggtg     1380
cccgtgcccg cggtcgcgct gttatctttc tgagcaacgg tgccgatcaa aatggtaatg     1440
tttacggtcg tattcaatct gcggcgatcg gtcaattgat tcgcgtttgg cgtcacgagg     1500
cggagttgga ctatcaacgt gcatccgccc caggcgatca cgttctgccg ccggtttggg     1560
cgaaccagat tgtccgtttc gctaaccgct ccctggaagg tctggagttc gcgtgcgctg     1620
ggaccgcaca gctgctgcac agccaacgtc atattaacga aattacgctg aacattccag     1680
ccaatattag cgcgaccacg ggcgcacgct ccgcccagct cggctggggc gagtccttga     1740
ttggtctgca cctgggcaag gtggctctga ttaccgggtg ttcggcgggc atcgggtggtc     1800
aaatcggctc tctgctggcc ttgtctggcg cgcgtgtgat gctggccgct cgcgatcgcc     1860
ataaattgga acagatgcaa gccatgattc aaagcgaatt ggccgagggt ggttataaccg     1920
atgtggagga ccgtgtgcac atcgcctccg gttcgcgatg gagcagcgag gcgcagctgg     1980
cagatctggg ggaacgtacg ctgtccgcat tcggtaccgt ggattatttg attaataacg     2040
ccggtattgc gggcgtggag gagatggtga tcgacatgcc ggtggaaggc tggcgtcaca     2100
ccctgtttgc caacctgatt tcgaattatt cgctgatgcg caagttggcg ccgctgatga     2160

```

-continued

```

agaagcaagg tagcggttac atcctgaacg tttcttcta ttttggcgtg gagaaggacg 2220
cggcgattcc ttatccgaac cggcgccgact acgcccgtctc caaggctggc caacgcgcga 2280
tggcggaagt gttcgctcgt ttctggggtc cagagattca gatcaatgct attgccccag 2340
gtccggttga aggegaccgc ctgctgggta ccggtgagcg tccgggctg tttgctcgtc 2400
gcgcccgtct gatcttgag aataaacgcc tgaacgaatt gcacgcggct ttgattgctg 2460
cggcccgcac cgatgagcgc tcgatgcacg agttggttga attggtgctg ccgaacgacg 2520
tggccgcggt ggagcagaac ccagcggccc ctaccgcgct gcgtgagctg gcacgcgct 2580
tccgtagcga aggtgatccg ggggcaagct cctcgtccgc cttgctgaat cgctccatcg 2640
ctgccaaagt gttggctcgc ttgcataacg gtggctatgt gctgccggcg gatatttttg 2700
caaatctgcc taatccgcgc gaccggttct ttaccggtgc gcaaattgac cgccaagctc 2760
gcaaggtgcg tgatggtatt atgggtatgc tgtatctgca gcgtatgcca accgagtttg 2820
acgtcgctat ggcaaccgtg tactatctgg ccgatcgtaa cgtgagcggc gaaactttcc 2880
atccgtctgg tggtttgcgc tacgagcgtc ccccgaccgg tggcgagctg ttcggcctgc 2940
catcgccgga acgtctggcg gagctgggtg gtagcacggt gtacctgatc ggtgaacacc 3000
tgaccgagca cctgaacctg ctggctcgtg cctatttggg gcgctacggt gcccgtaag 3060
tggatgatgat tgttgagacg gaaaccggtg cggaaaccat gcgtcgtctg ttgcatgatc 3120
acgtcgagggc aggtcgctg atgactattg tggcaggtga tcagattgag gcagcgattg 3180
accaagcgat cacgcgctat ggccgtccgg gtccggtggt gtgcactcca ttcgctccac 3240
tgccaaccgt tccgctggtc ggtcgtaaag actccgattg gagcacggtt ttgagcgagg 3300
cggaaattgc ggaactgtgt gagcatcagc tgaccacca tttccgtgtt gctcgtaaga 3360
tcgccttgtc ggatggcgcg tcgctggcgt tggttacccc ggaaacgact gcgactagca 3420
ccacggagca atttgcctg gcgaacttca tcaagaccac cctgcacgcg ttcaccgcca 3480
ccatcggtgt tgagtcggag cgcaccgcgc aacgtattct gattaaccag gttgatctga 3540
cgcgcgcgcg ccgtgcggaa gagccgcgtg acccgcacga gcgtcagcag gaattggaac 3600
gcttcattga agccgttctg ctggttaccg ctccgctgcc tcctgaggca gacacgcgct 3660
acgcaggccg tattcaccgc ggtcgtgcga ttaccgtcta atagaagctt gatata 3716

```

<210> SEQ ID NO 12

<211> LENGTH: 45

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic PCR primer

<400> SEQUENCE: 12

```

tcgtaccaac catggccggt acgggtcgtt tggctggtaa aattg 45

```

<210> SEQ ID NO 13

<211> LENGTH: 42

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic PCR primer

<400> SEQUENCE: 13

-continued

 cgggtgtgaga tctagatccg acggtaatcg cacgaccgcg gt 42

<210> SEQ ID NO 14

<211> LENGTH: 8262

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 pkk223 plasmid comprising codon-optimized mcr nucleic acid
 sequence

<400> SEQUENCE: 14

 ttgacaatta atcatcggct cgtataatgt gtggaattgt gagcggataa caatttcaca 60
 caggaaacag aaccatggcc ggtacggggtc gtttggtggtg taaaattgca ttgatcaccg 120
 gtggtgctgg taacattggt tccgagctga cccgccgttt tctggccgag ggtgcgacgg 180
 ttattatcag cggccgtaac cgtgcgaagc tgaccgcgct ggccgagcgc atgcaagccg 240
 aggccggcgt gccggccaag cgcattgatt tggaggtgat ggatggttcc gaccctgtgg 300
 ctgtccgtgc cggatcagag gcaatcgtcg ctgccacgg tcagattgac attctggtta 360
 acaacgcggg ctccgccggt gcccaacgtc gcttggcgga aattccgctg acggaggcag 420
 aattgggtcc ggggtgaggag gagactttgc acgcttcgat cgcgaaatctg ttgggcatgg 480
 gttggcacct gatgcgtatt ggggctccgc acatgccagt tggctccgca gttatcaacg 540
 tttcgactat tttctcgcgc gcagagtact atggtcgcac tccgtacgtt accccgaagg 600
 cagcgtgaa cgctttgtcc cagctggctg cccgcgagct gggcgctcgt ggcacccgcg 660
 ttaacactat tttcccaggt cctattgagt ccgaccgat ccgtaccgtg tttcaacgta 720
 tggatcaact gaagggtcgc ccggagggcg acaccgcca tcaacttttg aacaccatgc 780
 gcctgtgccg cgcaaacgac caaggcgctt tggaaacgac ctttccgtcc gttggcgatg 840
 ttgctgatgc ggctgtggtt ctggcttctg ctgagagcgc ggcactgtcg ggtgagacga 900
 ttgaggtcac ccacggtatg gaactgccgg cgtgtagcga aacctccttg ttggcgcgta 960
 ccgatctgag taccatcagc gcgagcggtc gactaccct gatttgcgct ggcgatcaaa 1020
 ttgaagaagt tatggccctg acgggcatgc tgcgtacgtg cggtagcga gtgattatcg 1080
 gcttccgttc tgcggctgcc ctggcgcaat ttgagcaggc agtgaatgaa tctcgcgctc 1140
 tggcagggtc ggatttcacc ccgccgatcg ctttgccggt ggaccacagt gaccgggcca 1200
 ccattgatgc ggttttcgat tggggcgag gcgagaatac ggggtggcatc catgcccggc 1260
 tcattctgcc ggcaacctcc cacgaaccgg ctccgtgctg gattgaagtc gatgacgaac 1320
 gcgtcctgaa tttcctggcc gatgaaatta ccggcaccat cgttattgag agccggttgg 1380
 cgcgctattg gcaatcccaa cgctgaccc cgggtgcccg tgcccgcggt ccgctgttta 1440
 tctttctgag caacggtgcc gatcaaatg gtaatgttta cggctcgtatt caatctgagg 1500
 cgatcggtea attgatcgc gtttgccgct acgagggcga gttggactat caacgtgcat 1560
 ccgccgagc cgatcacgtt ctgccgccgg tttggcgcaa ccagattgtc cgtttcgcta 1620
 accgctccct ggaaggctct gagttcgcgt gcgctggac cgcacagctg ctgcacagcc 1680
 aacgtcatat taacgaaatt acgctgaaca ttccagcaa tattagcgcg accacgggag 1740
 cacgttcgca cagcgtcggc tggggcgagt ccttgattgg tctgcacctg ggcaagggtg 1800
 ctctgattac cgggtggttcg gcgggcatcg gtggtcaaat cggctcgtctg ctggccttgt 1860
 ctggcgcgcg tgtgatgctg gccgctcgcg atcgccataa attggaacag atgcaagcca 1920

-continued

tgattcaaag cgaattggcg gaggttggtt ataccgatgt ggaggaccgt gtgcacatcg 1980
ctccgggttg cgatgtgagc agcggggcgc agctggcaga tctggtggaa cgtacgctgt 2040
ccgcattcgg taccgtggat tatttgatta ataacgccgg tattgcgggc gtggaggaga 2100
tggtgatcga catgccggtg gaaggctggc gtcacacct gtttgccaac ctgatttcga 2160
attattcgct gatgcgcaag ttggcgccgc tgatgaagaa gcaaggtagc ggttacatcc 2220
tgaacgtttc ttcctathtt ggcggtgaga aggacgcggc gattccttat ccgaaccgcg 2280
ccgactacgc cgtctccaag gctggccaac gcgcgatggc ggaagtgttc gctcgtttcc 2340
tgggtccaga gattcagatc aatgctattg cccaggtcc ggttgaaggc gaccgcctgc 2400
gtggtaccgg tgagcgtccg ggctgtttg ctcgctcgcg ccgtctgatc ttggagaata 2460
aacgcctgaa cgaattgcac gcggctttga ttgctcggc ccgcaccgat gagcgctcga 2520
tgcacgagtt ggttgaattg ttgctgccga acgacgtggc cgcgttgag cagaaccag 2580
cggccctac cgcgctcgt gagctggcac gcccttccg tagcgaaggt gatccggcgg 2640
caagctctc gtcgccttg ctgaatcgct ccacgctgc caagctgttg gctcgtttgc 2700
ataacggtgg ctatgtgctg ccggcggata tttttgaaa tctgcctaat ccgcccggacc 2760
cgttctttac ccgtgcgcaa attgaccgcg aagctcgcaa ggtgcgtgat ggtattatgg 2820
gtatgctgta tctgcagcgt atgccaaccg agtttgacgt cgctatggca accgtgtact 2880
atctggccga tctaacgtg agcggcgaaa ctttccatcc gtctggtggt ttgcgctacg 2940
agcgtacccc gaccggtggc gagctgttcg gcctgccatc gccggaacgt ctggcggagc 3000
tggttggtag cacggtgtac ctgatcggtg aacacctgac cgagcacctg aacctgctgg 3060
ctcgtgecta tttggagcgc tacggtgccc gtcaagtggg gatgattgtt gagacggaaa 3120
ccggtgcgga aacctgcgt cgtctgttgc atgatcacgt cgaggcaggt cgctgatga 3180
ctattgtggc aggtgatcag attgaggcag cgattgacca agcgatcacg cgctatggcc 3240
gtccgggtcc ggtggtgtgc actccattcc gtccactgcc aaccgttccg ctggtcggtc 3300
gtaaagactc cgattggagc accgttttga gcgaggcgga atttgcggaa ctgtgtgagc 3360
atcagctgac ccaccatttc cgtgttgctc gtaagatcgc cttgtcggat ggcgcgtcgc 3420
tggcgttggg taccocggaa acgactgcca ctagcaccac ggagcaattt gctctggcga 3480
acttcatcaa gaccaccctg cacgcgttca ccgcgaccat cgggtgtgag tcggagcgca 3540
ccgcgcaacg tattctgatt aaccaggttg atctgacgcg ccgcgccctg gcggaagagc 3600
cgcgtgacct gcacgagcgt cagcaggaat tggaacgctt cattgaagcc gttctgctgg 3660
ttaccgctcc gctgctcct gaggcagaca cgcgctacgc aggccgtatt caccgcggtc 3720
gtgcgattac cgtcggatct agatctcacc atcaccacca ttagtcgacc tgcagccaag 3780
cttggctggt ttggcggatg agagaagatt ttcagcctga tacagattaa atcagaacgc 3840
agaagcggtc tgataaaaca gaatttgctt ggccgagta gcgcgggtgg cccacctgac 3900
cccatgccga actcagaagt gaaacgccgt agcgcgatg gtagtgtggg gtctcccat 3960
gcgagagtag ggaactgcca ggcatcaaat aaaaagaaag gctcagtcga aagactgggc 4020
ctttcgtttt atctgttgtt tgtcggtgaa cgctctcctg agtaggaaa atccgcccgg 4080
agcggatttg aacgttgcca agcaacggcc cggaggggtg cgggcaggac gcccgccata 4140
aactgccagg catcaaatta agcagaaggc catcctgacg gatggccttt ttgcgtttct 4200

-continued

acaaactctt	ttgtttat	ttctaatac	attcaaatat	gtatccgctc	atgagacaat	4260
aaccctgata	aatgcttcaa	taatattgaa	aaaggaagag	tatgagtatt	caacatttcc	4320
gtgtcgccct	tattcccttt	tttgccgcat	tttgccctcc	tgtttttgct	caccagaaaa	4380
cgctggtgaa	agtaaaagat	gctgaagatc	agttgggtgc	acgagtgggt	tacatcgaac	4440
tggatctcaa	cagcggtaag	atccttgaga	gttttcgccc	cgaagaacgt	tttccaatga	4500
tgagcacttt	taaagttctg	ctatgtggcg	cggtattatc	ccgtgttgac	gccgggcaag	4560
agcaactcgg	tcgccgcata	cactattctc	agaatgactt	ggttgagtac	tcaccagtca	4620
cagaaaagca	tcttacggat	ggcatgacag	taagagaatt	atgcagtgct	gccataacca	4680
tgagtgataa	cactgcggcc	aacttacttc	tgacaacgat	cggaggaccg	aaggagctaa	4740
ccgctttttt	gcacaacatg	ggggatcatg	taactcgcct	tgatcgttgg	gaaccggagc	4800
tgaatgaagc	cataccaaac	gacgagcgtg	acaccacgat	gctgtagcaa	tggaacaac	4860
gttgcgcaaa	ctattaactg	gcgaactact	tactctagct	tcccggcaac	aattaataga	4920
ctggatggag	gcgataaaag	ttgcaggacc	acttctgcgc	tcggcccttc	cggtggctg	4980
gtttattgct	gataaatctg	gagccgggtga	gcgtgggtct	cgcggtatca	ttgcagcact	5040
ggggccagat	ggtaagccct	cccgtatcgt	agttatctac	acgacgggga	gtcaggcaac	5100
tatggatgaa	cgaaatagac	agatcgctga	gataggtgcc	tcactgatta	agcattggta	5160
actgtcagac	caagtttact	catatatact	ttagattgat	ttaaaacttc	atttttaatt	5220
taaaaggatc	taggtgaaga	tcctttttga	taatctcatg	acaaaaatcc	cttaacgtga	5280
gttttcgctc	cactgagcgt	cagaccccgt	agaaaagatc	aaaggatctt	cttgagatcc	5340
ttttttctg	cgcgtaatct	gctgcttgca	aacaaaaaaaa	ccaccgctac	cagcgggtgg	5400
ttgtttgccg	gatcaagagc	taccaactct	ttttccgaag	gtaactggct	tcagcagagc	5460
gcagatacca	aatactgtcc	ttctagtgtg	gccgtagtta	ggccaccact	tcaagaactc	5520
tgtagcaccg	cctacatacc	tcgctctgct	aatcctgtta	ccagtggctg	ctgccagtgg	5580
cgataagtcg	tgtcttaccg	ggttggactc	aagacgatag	ttaccggata	aggcgcagcg	5640
gtcgggctga	acggggggtt	cgtgcacaca	gccagcttg	gagcgaacga	cctacaccga	5700
actgagatac	ctacagcgtg	agcattgaga	aagcggccag	cttcccgaag	ggagaaaggc	5760
ggacaggat	ccggtaagcg	gcagggctcg	aacaggagag	cgcacgaggg	agcttccagg	5820
gggaaacgcc	tggtatcttt	atagtcctgt	cgggtttcgc	cacctctgac	ttgagcgtcg	5880
atttttgtga	tgctcgtcag	gggggcccag	cctatggaaa	aacgccagca	acgcccctt	5940
tttacggctc	ctggcctttt	gctggccttt	tgctcacatg	ttctttctg	cgttatcccc	6000
tgattctgtg	gataaccgta	ttaccgctt	tgagtgagct	gataccgctc	gccgcagccg	6060
aacgaccgag	cgcagcagct	cagtgagcga	ggaagcggaa	gagcgcctga	tgccgtatct	6120
tctccttacg	catctgtgct	gtatttcaca	ccgcatatgg	tgactctca	gtacaatctg	6180
ctctgatgcc	gcatagttaa	gccagtatac	actccgctat	cgctacgtga	ctgggtcatg	6240
gctgcgcccc	gacaccgctc	aacaccgct	gacgcgcctc	gacgggcttg	tctgctcccc	6300
gcatccgctt	acagacaagc	tgtgaccgtc	tccgggagct	gcatgtgtca	gaggttttca	6360
ccgtcatcac	cgaaacgcgc	gaggcagctg	cggtaaagct	catcagcgtg	gtcgtgaagc	6420
gattcacaga	tgtctgcctg	ttcatccgct	tccagctcgt	tgagtttctc	cagaagcgtt	6480
aatgtctggc	ttctgataaa	gcccccatg	ttaagggcgg	ttttttctg	tttggctcact	6540

-continued

```

gatgcctccg tgtaagggg atttctgttc atgggggtaa tgataccgat gaaacgagag 6600
aggatgctca cgatacgggt tactgatgat gaacatgccc ggttactgga acgttgtgag 6660
ggtaaacaac tggcgggatg gatgcggcgg gaccagagaa aaatcactca gggccaatgc 6720
cagcgcttcg ttaatacaga tgtaggtggt ccacagggta gccagcagca tctgcgatg 6780
cagatccgga acataatggt gcagggcgct gacttccgcy tttccagact ttacgaaaca 6840
cggaaaccga agaccattca tgttggtgct caggtcgcag acgttttca gcagcagtcg 6900
cttcacgttc gctcgcgat cggtgattca ttctgctaac cagtaaggca accccgccag 6960
cctagccggg tctcaacga caggagcacg atcatgcgca cccgtggcca ggaccaacg 7020
ctgcccgaga tgcgcccgt gcggctgctg gagatggcgg acgcgatgga tatgttctgc 7080
caagggttg tttgcgcatt cacagttctc cgcaagaatt gattggctcc aattcttga 7140
gtggtgaatc cgtagcgag gtgcccggc cttccattca ggtcgaggtg gcccggctcc 7200
atgcaccgcy acgcaacgcy gggaggcaga caaggtatag ggcggcgcct acaatccatg 7260
ccaaccggtt ccatgtgctc gccgaggcgg cataaatgcy cgtgacgatc agcggctccag 7320
tgatcgaagt taggctggtg agagccgcy gcgatccttg aagctgtccc tgatggctgt 7380
catctacctg cctggacagc atggcctgca acgcgggcat cccgatgccg ccggaagcga 7440
gaagaatcat aatggggaag gccatccagc ctgcgctgc gaacgccagc aagacgtagc 7500
ccagcgcgct gcccgccatg ccggcgataa tggcctgctt ctgcgcgaaa cgtttggtgg 7560
cgggaccagt gacgaaggct tgagcggagg cgtgcaagat tccgaatacc gcaagcgaca 7620
ggccgatcat cgtcgcgctc cagcgaagc ggtcctcgc gaaaatgacc cagagcgtg 7680
ccggcacctg tctacgagt tgcatgataa agaagacagt cataagtgcg gcgacgatag 7740
tcatgccccg cccccaccg aaggagctga ctgggtgaa ggctctcaag ggcatcggtc 7800
gacgctctcc cttatgcgac tctgcatta ggaagcagcc cagtagtagg ttgaggccgt 7860
tgagcaccgc cgcgcaagg aatggtgcat gcaaggagat ggcgccaac agtcccccg 7920
ccacggggcc tgccaccata cccacgccga aacaagcgt catgagcccg aagtggcgag 7980
cccgatcttc cccatcggtg atgtcggcga tataggcgc agcaaccgca cctgtggcgc 8040
cgtgatgcc gccacgatg cgtccggcgt agaggatccg ggcttatcga ctgcacggtg 8100
caccaatgct tctggcgtca ggcagccatc ggaagctgtg gtatggctgt gcaggtcgta 8160
aatcactgca taattcgtgt cgetcaaggc gactcccgt tctggataat gttttttgcy 8220
ccgacatcat aacggttctg gcaaatattc tgaatgagc tg 8262

```

```

<210> SEQ ID NO 15
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
PCR primer

```

```

<400> SEQUENCE: 15

```

```

gggtttccat ggaccagccg ctcaacgtgg 30

```

```

<210> SEQ ID NO 16
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

```

-continued

<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
PCR primer

<400> SEQUENCE: 16

gggttttcag gcgatgccgt tgagcgcttc gcc 33

<210> SEQ ID NO 17
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
PCR primer

<400> SEQUENCE: 17

gggaacggcg gggaaaaaca aacggt 26

<210> SEQ ID NO 18
<211> LENGTH: 30
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
PCR primer

<400> SEQUENCE: 18

gtccatggt aattctccac gcttataagc 30

<210> SEQ ID NO 19
<211> LENGTH: 85
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 19

ggtttgaata aatgacaaaa agcaaagcct ttgtgccgat gaatctctat actgtttcac 60
agacctgctg cctgcgggg cggcc 85

<210> SEQ ID NO 20
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
PCR primer

<400> SEQUENCE: 20

gggaacggcg gggaaaaaca aacggt 26

<210> SEQ ID NO 21
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
PCR primer

<400> SEQUENCE: 21

gggttttcag gcgatgccgt tgagcgcttc gcc 33

<210> SEQ ID NO 22
<211> LENGTH: 1754
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence

-continued

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
Codon-optimized pyruvate decarboxylase DNA sequence based on gene
of *Zymomonas mobilis*

<400> SEQUENCE: 22

```

catatgtcct aactgttgg tacttatctg gctgaacgtc tggttcaaat tggctgaag    60
catcactttg cggtagcggg cgattacaac ctggtgctgc tggataatct gctgctgaac    120
aaaaacatgg aacaggctca ctggtgtaac gaactgaact gcggtctctc tgctgaaggt    180
tatgcccggg ccaaaggcgc agctgcgggc gtagtgacct actccgttgg tgctctgtcc    240
gcatttgatg caattggtgg cgcctacgca gaaaacctgc cggtaattct gatctctggc    300
gctccgaaca acaacgatca cgcagctggt cacgtgctgc accatgctct gggcaaaact    360
gattatcatt accagctgga aatggcgaag aacatcactg ccgcagcaga agctatctat    420
actccagaag aagcggcggc aaaaatcgat catgtaatca aaacggccct gcgtgagaag    480
aaaccggtgt atctgaaaat tgcttgtaat atcgcgtcta tgccgtgtgc ggccccaggg    540
ccgcatctct ctctgtttaa cgatgaagct agcgatgaag cctctctgaa cgcagctgtg    600
gaagaaacct tgaaattcat tgcaaaccgt gacaaagttg cggactggtt tggctctaaa    660
ctgcgtgccg cgggtgcaga agaagcggcg gttaaattcg ctgacgccct ggggtggtgct    720
gtggccacca tggctgcggc taaatccttt ttcccgaag aaaatccgca ttacatcggg    780
acttctctgg gcgaggtttc ttaccaggt gtcgagaaaa ccatgaagga agctgacggc    840
gtgatcgccc tggccccggg tttcaatgac tactccacta ctggttgac cgacatcccg    900
gacccaaaga aactggttct ggcagagccg cgctccgttg ttgtaacgg tattcgcttt    960
ccgtccgtac acctgaagga ttatctgact cgtctggcgc agaaagtgag caagaaaacc   1020
ggcgtctctg atttctttaa atctctgaat gcgggtgagc tgaagaaagc cgcaccggcg   1080
gacccttctg ctccgctggt taacgccgaa attgcgcgcc aggtagaagc gctgctgact   1140
ccgaatacta ccgtaattgc ggagactggc gattcctggt tcaacgcaca acgtatgaag   1200
ctgcctaacg gcgctcgagt tgaatacгаа atgcagtggt gccacatcgg ctggtctggt   1260
cctgcagcct tcggctacgc cgtaggtgct ccggaacgtc gtaacatcct gatggtcggg   1320
gacggctctt tccaactgac cgcgcaggaa gtagcacaga tggttcgtct gaaactgccg   1380
gtaatcatct tcctgattaa caactacggc tataccattg aggtcatgat tcatgatggt   1440
ccgtataata acatcaaaaa ctgggattat gctggtctga tggaaagttt caacggcaac   1500
ggcggctacg attctggtgc tggtaaaggc ctgaaagcaa agacgggtgg cgagctcgca   1560
gaagcgatca aggttgctct ggctaacacc gatggtccga ctctgatcga atgttttatc   1620
ggtcgtgaag attgactga ggaactggtg aagtggggtg agcgtgtggc tgcccggaat   1680
tcccgtaaac cggtaaataa gcttctcggc catcaccatc accatcacta gaagcttctc   1740
tagagaacta tttc                                     1754

```

<210> SEQ ID NO 23

<211> LENGTH: 575

<212> TYPE: PRT

<213> ORGANISM: *Zymomonas mobilis*

<400> SEQUENCE: 23

Met Ser Tyr Thr Val Gly Thr Tyr Leu Ala Glu Arg Leu Val Gln Ile
1 5 10 15

-continued

Gly Leu Lys His His Phe Ala Val Ala Gly Asp Tyr Asn Leu Val Leu
 20 25 30
 Leu Asp Asn Leu Leu Leu Asn Lys Asn Met Glu Gln Val Tyr Cys Cys
 35 40 45
 Asn Glu Leu Asn Cys Gly Phe Ser Ala Glu Gly Tyr Ala Arg Ala Lys
 50 55 60
 Gly Ala Ala Ala Ala Val Val Thr Tyr Ser Val Gly Ala Leu Ser Ala
 65 70 75 80
 Phe Asp Ala Ile Gly Gly Ala Tyr Ala Glu Asn Leu Pro Val Ile Leu
 85 90 95
 Ile Ser Gly Ala Pro Asn Asn Asn Asp His Ala Ala Gly His Val Leu
 100 105 110
 His His Ala Leu Gly Lys Thr Asp Tyr His Tyr Gln Leu Glu Met Ala
 115 120 125
 Lys Asn Ile Thr Ala Ala Ala Glu Ala Ile Tyr Thr Pro Glu Glu Ala
 130 135 140
 Pro Ala Lys Ile Asp His Val Ile Lys Thr Ala Leu Arg Glu Lys Lys
 145 150 155 160
 Pro Val Tyr Leu Glu Ile Ala Cys Asn Ile Ala Ser Met Pro Cys Ala
 165 170 175
 Ala Pro Gly Pro Ala Ser Ala Leu Phe Asn Asp Glu Ala Ser Asp Glu
 180 185 190
 Ala Ser Leu Asn Ala Ala Val Glu Glu Thr Leu Lys Phe Ile Ala Asn
 195 200 205
 Arg Asp Lys Val Ala Val Leu Val Gly Ser Lys Leu Arg Ala Ala Gly
 210 215 220
 Ala Glu Glu Ala Ala Val Lys Phe Ala Asp Ala Leu Gly Gly Ala Val
 225 230 235 240
 Ala Thr Met Ala Ala Ala Lys Ser Phe Phe Pro Glu Glu Asn Pro His
 245 250 255
 Tyr Ile Gly Thr Ser Trp Gly Glu Val Ser Tyr Pro Gly Val Glu Lys
 260 265 270
 Thr Met Lys Glu Ala Asp Ala Val Ile Ala Leu Ala Pro Val Phe Asn
 275 280 285
 Asp Tyr Ser Thr Thr Gly Trp Thr Asp Ile Pro Asp Pro Lys Lys Leu
 290 295 300
 Val Leu Ala Glu Pro Arg Ser Val Val Val Asn Gly Ile Arg Phe Pro
 305 310 315 320
 Ser Val His Leu Lys Asp Tyr Leu Thr Arg Leu Ala Gln Lys Val Ser
 325 330 335
 Lys Lys Thr Gly Ala Leu Asp Phe Phe Lys Ser Leu Asn Ala Gly Glu
 340 345 350
 Leu Lys Lys Ala Ala Pro Ala Asp Pro Ser Ala Pro Leu Val Asn Ala
 355 360 365
 Glu Ile Ala Arg Gln Val Glu Ala Leu Leu Thr Pro Asn Thr Thr Val
 370 375 380
 Ile Ala Glu Thr Gly Asp Ser Trp Phe Asn Ala Gln Arg Met Lys Leu
 385 390 395 400
 Pro Asn Gly Ala Arg Val Glu Tyr Glu Met Gln Trp Gly His Ile Gly
 405 410 415
 Trp Ser Val Pro Ala Ala Phe Gly Tyr Ala Val Gly Ala Pro Glu Arg

-continued

	420	425	430		435	440	445								
Arg	Asn	Ile	Leu	Met	Val	Gly	Asp	Gly	Ser	Phe	Gln	Leu	Thr	Ala	Gln
Glu	Val	Ala	Gln	Met	Val	Arg	Leu	Lys	Leu	Pro	Val	Ile	Ile	Phe	Leu
Ile	Asn	Asn	Tyr	Gly	Tyr	Thr	Ile	Glu	Val	Met	Ile	His	Asp	Gly	Pro
Tyr	Asn	Asn	Ile	Lys	Asn	Trp	Asp	Tyr	Ala	Gly	Leu	Met	Glu	Val	Phe
Asn	Gly	Asn	Gly	Gly	Tyr	Asp	Ser	Gly	Ala	Gly	Lys	Gly	Leu	Lys	Ala
Lys	Thr	Gly	Gly	Glu	Leu	Ala	Glu	Ala	Ile	Lys	Val	Ala	Leu	Ala	Asn
Thr	Asp	Gly	Pro	Thr	Leu	Ile	Glu	Cys	Phe	Ile	Gly	Arg	Glu	Asp	Cys
Thr	Glu	Glu	Leu	Val	Lys	Trp	Gly	Lys	Arg	Val	Ala	Ala	Ala	Asn	Ser
Arg	Lys	Pro	Val	Asn	Lys	Leu	Leu	Gly	His	His	His	His	His	His	His

<210> SEQ ID NO 24
 <211> LENGTH: 5
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Histidine tag peptide sequence

<400> SEQUENCE: 24

His His His His His
 1 5

<210> SEQ ID NO 25
 <211> LENGTH: 7
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 Histidine tag peptide sequence

<400> SEQUENCE: 25

His His His His His His His
 1 5

What is claimed is:

1. A method for producing an acrylic acid-based consumer product, said method comprising

- i) combining a carbon source and a microorganism cell culture to produce 3-hydroxypropionic acid;
- ii) converting said 3-hydroxypropionic acid to acrylic acid; and
- iii) processing said acrylic acid into a consumer product.

2. The method of claim 1, wherein said cell culture comprises a genetically modified microorganism.

3. The method of claim 2, wherein said microorganism is modified for increased tolerance to 3-hydroxypropionic acid.

4. The method of claim 3, wherein said modification modulates one or more components of the chorismate superpathway.

5. The method of claim 2, wherein said microorganism is modified for increased production of 3-hydroxypropionic acid.

6. The method of claim 5, wherein said modification comprises reduction in activity of one or more enzymes selected from pyruvate kinase, phosphofructokinase, lactate dehydrogenase, phosphate acetyltransferase, pyruvate oxidase, pyruvate-formate lyase, and homologs thereof.

7. The method of claim 6, wherein said pyruvate kinase is selected from pykA and pykF, said phosphofructokinase is selected from pfkA and pfkB, said lactate dehydrogenase is selected from ldhA, said phosphate acetyltransferase is selected from pta, said pyruvate oxidase is selected from poxB, said pyruvate-formate lyase is selected from pflB, and homologs thereof.

8. The method of claim **5**, wherein said modification comprises increase in activity of one or more enzymes selected from phosphoenolpyruvate carboxykinase, malonyl-CoA reductase, 3-hydroxypropionate dehydrogenase, malonate semialdehyde dehydrogenase A, alpha-ketoglutarate decarboxylase, oxaloacetate alpha-oxo-decarboxylase, and homologs thereof.

9. The method of claim **8**, wherein said phosphoenolpyruvate carboxykinase is selected from pck, said malonyl-CoA reductase is selected from mer, said malonate semialdehyde dehydrogenase A is selected from mmsA, said 3-hydroxypropionic acid dehydrogenase is selected from mmsB, said alpha-ketoglutarate decarboxylase is selected from kgd, said oxaloacetate alpha-oxo-decarboxylase is selected from oad, and homologs thereof.

10. The method of claim **8**, wherein said modification comprises an increase in activity in a malonyl-CoA reductase enzyme.

11. The method of claim **8**, wherein said modification comprises an increase in activity in an oxaloacetate alpha-oxo-decarboxylase enzyme.

12. The method of claim **2**, wherein said microorganism is modified for increased tolerance to 3-hydroxypropionic acid, and wherein said microorganism is modified for increased production of 3-hydroxypropionic acid.

13. A method for producing an acrylic acid-based consumer product, said method comprising

- i) combining a carbon source and a genetically modified microorganism in cell culture to produce 3-hydroxypropionic acid;
 - ii) converting said 3-hydroxypropionic acid to acrylic acid; and
 - iii) processing said acrylic acid into a consumer product;
- wherein said microorganism is modified for increased production of 3-hydroxypropionic acid via an increase in activity in an oxaloacetate alpha decarboxylase enzyme or homolog thereof.

14. A method for producing acrylic acid, said method comprising

- i) combining a carbon source and a microorganism cell culture to produce 3-hydroxypropionic acid in a concentration of at least 10 g/L; and
- ii) converting said 3-hydroxypropionic acid to acrylic acid.

15. The method of claim **14**, wherein the combining comprises combining a carbon source and a microorganism cell culture comprising a microorganism genetically modified to increase activity of one or more enzymes selected from phosphoenolpyruvate carboxykinase, malonyl-CoA reductase, malonate semialdehyde dehydrogenase A, 3-HP dehydrogenase, alpha-ketoglutarate decarboxylase, oxaloacetate alpha-oxo-decarboxylase, and homologs thereof.

16. The method of claim **15**, wherein said phosphoenolpyruvate carboxykinase is selected from pck, said malonyl-CoA reductase is selected from mer, said malonate semialdehyde dehydrogenase A is selected from mmsA, said 3-hydroxypropionic acid dehydrogenase is selected from mmsB, said alpha-ketoglutarate decarboxylase is selected from kgd, said oxaloacetate alpha-oxo-decarboxylase is selected from oad, and homologs thereof.

17. The method of claim **14**, wherein the combining comprises combining a carbon source and a microorganism cell culture comprising a microorganism genetically modified to increase activity of malonyl-CoA reductase enzyme increasing production of 3-hydroxypropionic acid.

18. The method of claim **14**, wherein the combining comprises combining a carbon source and a microorganism cell culture comprising a microorganism genetically modified to increase activity of an oxaloacetate alpha-oxo-decarboxylase enzyme, increasing production of 3-hydroxypropionic acid.

19. Biologically-produced acrylic acid, wherein said acrylic acid is produced according to claim **14**.

20. A consumer product produced with acrylic acid according to claim **19**.

* * * * *