



US 20150072384A1

(19) **United States**

(12) **Patent Application Publication**  
**Lynch et al.**

(10) **Pub. No.: US 2015/0072384 A1**

(43) **Pub. Date: Mar. 12, 2015**

(54) **GENETICALLY MODIFIED ORGANISMS  
FOR INCREASED MICROBIAL  
PRODUCTION OF 3-HYDROXYPROPIONIC  
ACID INVOLVING AN OXALOACETATE  
ALPHA-DECARBOXYLASE**

**Publication Classification**

(71) Applicant: **OPX Biotechnologies, Inc.**, Boulder,  
CO (US)

(51) **Int. Cl.**  
*C12P 7/42* (2006.01)  
*C08F 20/06* (2006.01)  
*C07C 51/377* (2006.01)

(72) Inventors: **Michael D. Lynch**, Durham, NC (US);  
**Christopher P. Mercogliano**,  
Minneapolis, MN (US)

(52) **U.S. Cl.**  
CPC ..... *C12P 7/42* (2013.01); *C07C 51/377*  
(2013.01); *C08F 20/06* (2013.01)  
USPC ..... **435/135**

(21) Appl. No.: **14/341,223**

(57) **ABSTRACT**

(22) Filed: **Jul. 25, 2014**

**Related U.S. Application Data**

(63) Continuation of application No. 12/891,760, filed on  
Sep. 27, 2010, now Pat. No. 8,809,027.

(60) Provisional application No. 61/246,140, filed on Sep.  
27, 2009.

Microorganism compositions are described that comprise combinations of genetic modifications that include a genetic modification to increase oxaloacetate alpha-decarboxylase enzymatic activity. By such genetic modification a 3-hydroxypropionic acid ("3-HP") production pathway is provided or improved. In various embodiments, comprising other genetic modifications, including selected gene disruptions, 3-HP production is greater than in a control microorganism lacking such combinations of genetic modifications

Figure 1: CLUSTAL 2.0.12 multiple sequence alignment of Carbonic Anhydrase Polypeptides

		SEQ ID NO:
gil70728869 reflYYP_258618.1	MQNIIDGFLKFQREAFQPSSELFKQLASTQNPGLFVTCSDSRVVPPELLT 50	59
gil238754662 reflZP_04616015.1	MQDIIDGFLKFQREVFPQPSSELFKRLADTQHPGALFVTCSDSRVVPPELLT 50	59
gil83646817 reflYYP_435252.1	MKDIEGFLKFQREAFPERKELFKDLANQQPRTLFISSCSDSRLVPELVT 50	60
gil206562261 reflYYP_002233024.	MKDIEGFLKFQRDAYPARAALFRDLARSQNPRLFISCSDSLVPPELVT 50	61
gil15800068 reflNP_286080.1	MKEIIDGFLKFQREAFKREALFKQLATQQSPRTLFISSCSDSRLVPELVT 50	57
gil238790503 reflZP_04634271.1	MKEIIDGFLKFQRDAFPERAELFRSLATQQSPKTLFISSCSDSRMVPELVT 50	62
gil104782623 reflYYP_609121.1	MQDIIDGFLKFQRDAFPERVKLFDLQATQQSPRALFISSCSDSRLVPELVT 50	63
gil170722264 reflYYP_001749952.	MKAIIDGFLKFQKNAFPERVKLFDLQATQQAPKALFISSCSDSRLVPELVT 50	64
gil157369777 reflYYP_001477766.	MKEVIEGFLKFQREAFVERTALFQRLATQQSPRTLFISSCSDSRLVPELIT 50	65
gil188533851 reflYYP_001907648.	MQHIVEGFLNFQKDFPEQKELFRSLASSQNPKALFISSCSDSRLVPELVT 50	66
gil152985230 reflYYP_001348595.	MRDIIDGFLRFQRDAYPARSQLFKSLATRQAPKALFIACSDSRVVPPELLT 50	67
gil271966225 reflYYP_003340421.	MQDLEEGVARFQRDVPAKTELFRSLATAHQPATLFISSCDARVPELIT 50	68
	*:.*.***: : ** ** ;* :*:*:*:*:*:*:*	
gil70728869 reflYYP_258618.1	QREPGLDFVIRNAGNIVPSYGP-EPGGVSATVEYAVAVLGVSDIVICGHS 99	
gil238754662 reflZP_04616015.1	QREPGLDFVIRNAGNIVPSYGP-EPGGVSATVEYAVAVLGVSDIVICGHS 99	
gil83646817 reflYYP_435252.1	QREPGLDFVIRNAGNIVPPYGP-EPGGVSASVEYAVAAALRVADIVVCGHS 99	
gil206562261 reflYYP_002233024.	QREPGLDFVIRNAGNIVPSYGP-EPGGVSASVEYAVAAALRVADIVVCGHS 99	
gil15800068 reflNP_286080.1	QREPGLDFVIRNAGNIVPSYGP-EPGGVSASVEYAVAAALRVSDIVICGHS 99	
gil238790503 reflZP_04634271.1	QREPGLDFVIRNAGNIVPSYGP-EPGGISASVEYAVTALKVTDIVICGHS 99	
gil104782623 reflYYP_609121.1	QREPGLDFVIRNAGNIVPSYGP-EPGGVSASVEYAVAAALQVADIVICGHS 99	
gil170722264 reflYYP_001749952.	QREPGLDFVIRNAGNIVPSYGP-EPGGVSASVEYAVAGLNVADIVICGHS 99	
gil157369777 reflYYP_001477766.	QREPGLDFVIRNAGNIVPSFGP-EPGGVSASVEYAVSALGVSDIVICGHS 99	
gil188533851 reflYYP_001907648.	QQDPGQLFVIRNAGNIVPSFGP-EPGGVSATIEYAVVALGVSDIVICGHS 99	
gil152985230 reflYYP_001348595.	QREPGLDFVIRNAGNIVPGYGP-QPGGVSASVEYAVAVLGVADIVVCGHS 99	
gil271966225 reflYYP_003340421.	QSEPGELFVIRTAGNLVPA YAPGSADGVAAGIEYAVAVLGVSDIVVCGHS 100	
	*: **:*:*:*:*:*:* :*.. * :*: : **** *:* : * : * **	
gil70728869 reflYYP_258618.1	DCGAMTAISTCKCLDHLPAVANWLRHAESA KVINAAARQHASP AEHL DALV 149	
gil238754662 reflZP_04616015.1	NCGAMSAIAEQCLDHLPAVAAWLRHADSAKLVNAALPHASPKDRLNSLV 149	
gil83646817 reflYYP_435252.1	NCGAMTAVATCQCIDHMPAVAHWLR YADSAKVVNQARKHASERAKIEDMV 149	
gil206562261 reflYYP_002233024.	DCGAMTAIATCQCMDHMPAVGHWLR YAD SARVVNEARTHRSERERIDSMV 149	
gil15800068 reflNP_286080.1	NCGAMTAIASCQCMDHMPAVSHWLR YAD SARVVNEARPHSDLP SKAAAMV 149	
gil238790503 reflZP_04634271.1	DCGAMTAIAKCHCLDHMPAVKHWLQYADSAKVVNESREYKNIHDKTISMV 149	
gil104782623 reflYYP_609121.1	DCGAMTAIATCKCLDHMPAVAGWLR YAD SARVVNEARQHQS PHAKVEAMV 149	
gil170722264 reflYYP_001749952.	DCGAMTAIATCKCLDHMPAVAGWLR YADSAKVVNEARHHVDKPSKVASMV 149	
gil157369777 reflYYP_001477766.	DCGAMTAIATCQCLQHMPVANWLR YADSAKVVNQAYQHASENEKVSSMV 149	
gil188533851 reflYYP_001907648.	NCGAMKAIATCQCLAPMPAVEHWLR YADAAKAVVEKKNYDTEEDKVNAMV 149	
gil152985230 reflYYP_001348595.	DCGAMGAIASCA CLDHLPAVAGWLRHAEAA RAMNSAHEHSSDAARLDALV 149	
gil271966225 reflYYP_003340421.	GCGAMTAVADGLDPAALPAVAGWLRHADASRRARVTTTETGTG--EVAALV 148	
	**** *: :*:* **:: *::: . :*	
gil70728869 reflYYP_258618.1	RDNVIAQLANLKTHTPSVALALEQGRLNLHG WVVYDIESGAI VALDGNTQRF 199	
gil238754662 reflZP_04616015.1	RENVIAQLANIKTHTPSVALACAQGR LRLHG WVVYDIETGSIDVLDELTRTF 199	
gil83646817 reflYYP_435252.1	RENVIAQLANLQTHPSVRLALQEGRLTMHG WVFYDIESGGIDAYDGSRHAF 199	
gil206562261 reflYYP_002233024.	RENVVIAQLANLKTHTPAVRLALEEGRLALHG WVVYDIESGCIDAYDGATGRF 199	
gil15800068 reflNP_286080.1	RENVIAQLANLQTHPSVRLALEEGRIALHG WVVYDIESGSIAAFD GATRQF 199	
gil238790503 reflZP_04634271.1	HENVVIAQLANIQTHTPSVRLALEEGRLTIHG WVVYDIESGLISAFDRASRQF 199	
gil104782623 reflYYP_609121.1	RENVIAQLANIQTHPSVRLALEEGRVLHG WVIYDIESGRIDAFDGRGTGQF 199	
gil170722264 reflYYP_001749952.	RENVIAQLANIKTHTPSVALALEQGRLKLHG WVVYDIASGGIEALDGETRRF 199	
gil157369777 reflYYP_001477766.	QENVIAQLNNTHTPSVAVGLRNNALRLHG WVVYDIESGAIRALDKDSKKF 199	
gil188533851 reflYYP_001907648.	RHNVIAQLANLRTHTPSVARALEQGRLNLHG WVVYDIESGRIDALDGASRRF 199	
gil152985230 reflYYP_001348595.	RQNVLTQLANLATHPSVAHALAGKTVTLHG WVIYDIGTGTVAELD-ATGRP 197	
gil271966225 reflYYP_003340421.	..*:*:* * : **:* . :***:** :* : *	
gil70728869 reflYYP_258618.1	VSLAEYPHTCALASQASSAA- 219	
gil238754662 reflZP_04616015.1	SPLSAY---SVVSKPTE--- 213	
gil83646817 reflYYP_435252.1	VPLAEHPPEARAI PGKLSHAV- 219	
gil206562261 reflYYP_002233024.	VSLADHPGVRATPATLPVAA- 219	
gil15800068 reflNP_286080.1	VPLAANPRVCAIPLRQPTAA- 219	
gil238790503 reflZP_04634271.1	VSLAANPNVRAVPAHN---- 215	
gil104782623 reflYYP_609121.1	VSLADNPEVRAVSHSRHVA- 219	
gil170722264 reflYYP_001749952.	VSLAENPEVHVAVSQARHVA- 219	
gil157369777 reflYYP_001477766.	IPLATNPEVTATPAVSRF--- 217	
gil188533851 reflYYP_001907648.	VLLSDNPQVHFE----- 211	
gil152985230 reflYYP_001348595.	VSLAHPGVRAVGGEPGQAVA 220	
gil271966225 reflYYP_003340421.	SALAV----- 202	
	*:	

Figure 2




Figure 3A

Natural mixed acid fermentation in *E. coli*





Figure 3B

Proposed 3-HP fermentation



Gene Deletions:

- ldhA: pyruvate ~~→~~ lactate
- pflB: pyruvate ~~→~~ formate + acetyl coA
- poxB: pyruvate ~~→~~ acetate
- pta: acetyl-coA ~~→~~ acetate



FIG. 6A




FIG. 6B




Figure 7




Figure 8: Western blot of purified proteins probed with anti-6xHis antibody







Figure 9: GC-MS results at various pH with and without addition of Oad-2 candidate enzymes



**GENETICALLY MODIFIED ORGANISMS  
FOR INCREASED MICROBIAL  
PRODUCTION OF 3-HYDROXYPROPIONIC  
ACID INVOLVING AN OXALOACETATE  
ALPHA-DECARBOXYLASE**

CROSS-REFERENCE TO RELATED  
APPLICATIONS

[0001] This application claims priority from U.S. Provisional Application 61/246,140, filed Sep. 27, 2009. The entire contents of this application are hereby incorporated by reference in its entirety.

SEQUENCE LISTING

[0002] The instant application contains a Sequence Listing which has been submitted in ASCII format via EFS-Web and is hereby incorporated by reference in its entirety. Said ASCII copy, created on Feb. 10, 2011, is named OPXX2005.txt and is 234,626 bytes in size.

FIELD OF THE INVENTION

[0003] This invention relates to metabolically engineered microorganisms, such as bacterial strains, in which an identified functional variant of oxaloacetate alpha-decarboxylase is provided for production of a chemical product, 3-hydroxypropionic acid (3-HP) and products made from 3-HP. The metabolically engineered microorganisms may be adapted to exhibit increased tolerance to 3-HP. Production of products made from 3-HP so produced also is disclosed.

SEQUENCE LISTING

[0004] This provisional patent application provides a paper copy of sequence listings that are to be provided on compact disk in appropriate format in a later filing or submission.

BACKGROUND OF THE INVENTION

[0005] There are various approaches to increasing a genetically modified microorganism's productivity of 3-hydroxypropionic acid ("3-HP"). These approaches may be applied to a microorganism intended to be used in a production strain having the purpose of 3-HP production in an industrial microbial production system, whether or not 3-HP is the intended final product.

[0006] For example, without being limiting, various 3-HP production pathways are described in U.S. Pat. No. 6,852,517, WO2002/042418 (PCT/US01/43607), and U.S. Patent Publication No. US2008/0199926.

[0007] Nonetheless, notwithstanding these and other various microbial 3-HP production pathways, there remains a need in the art for alternative 3-HP production pathways.

SUMMARY OF THE INVENTION

[0008] According to one embodiment, the invention is directed to a method for producing an acrylic acid-based consumer product, said method comprising i) combining a carbon source and a microorganism cell culture to produce 3-hydroxypropionic acid, wherein a) wherein said microorganism is genetically modified for increased enzymatic activity in the organism by introduction of a heterologous nucleic acid sequence coding for a polypeptide having oxaloacetate alpha-decarboxylase enzymatic activity, the heterologous

nucleic acid sequence expressing a mutated oxaloacetate alpha-decarboxylase/oxaloacetate alpha-decarboxylase.

[0009] The carbon source according to the invention may be predominantly glucose, sucrose, fructose, dextrose, lactose, or a combination thereof. Alternatively, the carbon source is glycerol.

[0010] Included within the invention are embodiments where the cell culture comprises a genetically modified microorganism. The genetically modified microorganism may be modified for increased activity and specificity to convert oxaloacetate to malonate semialdehyde, increased tolerance to 3-hydroxypropionic acid, increased enzymatic activity in the organism's NADPH-dependent transhydrogenase pathway, increased intracellular bicarbonate levels, and combinations thereof.

[0011] In various embodiments, the genetically modified microorganism is modified for increased tolerance to 3-hydroxypropionic acid. The increase in tolerance to 3-hydroxypropionic acid may occur in one or more components of the 3-HP toleragenic complex (3HPTGC) complex.

[0012] The genetically modified bacteria may be further modified to decrease activity of, lactate dehydrogenase, phosphate acetyltransferase, pyruvate oxidase, or pyruvate-formate lyase, alcohol dehydrogenase, and combinations thereof.

[0013] The method according to the invention may further comprise separating and/or purifying 3-hydroxypropionic acid from said cell culture by extraction of 3-hydroxypropionic acid from said culture in the presence of a tertiary amine.

[0014] The method of the invention may include production of a consumer product, such as diapers, carpet, paint, adhesives, and acrylic glass. The invention includes biologically-produced 3-hydroxypropionic acid, where the 3-hydroxypropionic acid is produced according to the method of the invention. Such 3-hydroxypropionic acid may be essentially free of chemical catalyst, including a molybdenum and/or vanadium based catalyst. The 3-hydroxypropionic acid is produced according to the method of the invention may have a ratio of carbon-14 to carbon-12 of about  $1.0 \times 10^{-14}$  or greater. In various aspects, the 3-hydroxypropionic acid contains less than about 10% carbon derived from petroleum. In addition, 3-hydroxypropionic acid according to the invention may contain a residual amount of organic material related to its method of production. In various embodiments, the 3-hydroxypropionic acid contains a residual amount of organic material in an amount between 1 and 1,000 parts per million of the 3-hydroxypropionic acid.

[0015] Acrylic acid and a polymer produced from acrylic acid, where such are produced according to the method of the invention, are also included within the invention. Products, including commercial and consumer products, obtained from the polymers are also encompassed. For example, diapers, carpet, paint, adhesives, and acrylic glass are encompassed.

[0016] In addition, the invention encompasses a system for bioproduction of acrylic acid according to claim 40, said system comprising: a tank for saccharification of biomass; a line for passing the product of saccharification to a fermentation tank optionally via a pre-fermentation tank; a fermentation tank suitable for microorganism cell culture; a line for discharging contents from the fermentation tank to an extraction and/or separation vessel; an extraction and/or separation vessel suitable for removal of 3-hydroxypropionic acid from cell culture waste; a line for transferring 3-hydroxypropionic acid to a dehydration vessel; and a dehydration vessel suitable

for conversion of 3-hydroxypropionic acid to acrylic acid. In various embodiments, the system further comprises one or more pre-fermentation tanks, distillation columns, centrifuge vessels, back extraction columns, mixing vessels, or combinations thereof. In various embodiments, the system has a minimum production capacity of at least 1 ton acrylic acid per year.

**[0017]** In various embodiments, a further genetic modification has been made that increases NADH/NADPH transhydrogenase activity. For example, the transhydrogenase activity may be soluble, may be membrane bound, may have a further genetic modification that has been made that increases cyanase activity, may include a further genetic modification that increases carbonic anhydrase activity, and/or may include a further genetic modification that increases pyruvate dehydrogenase activity.

**[0018]** In various embodiments, the invention includes a culture system comprising a carbon source in an aqueous medium and a genetically modified microorganism according to any one of claims, wherein said genetically modified microorganism is present in an amount selected from greater than 0.05 gDCW/L, 0.1 gDCW/L, greater than 1 gDCW/L, greater than 5 gDCW/L, greater than 10 gDCW/L, greater than 15 gDCW/L or greater than 20 gDCW/L, such as when the volume of the aqueous medium is selected from greater than 5 mL, greater than 100 mL, greater than 0.5 L, greater than 1 L, greater than 2 L, greater than 10 L, greater than 250 L, greater than 1000 L, greater than 10,000 L, greater than 50,000 L, greater than 100,000 L or greater than 200,000 L, and such as when the volume of the aqueous medium is greater than 250 L and contained within a steel vessel.

**[0019]** Various, the carbon source for such culture systems is selected from dextrose, sucrose, a pentose, a polyol, a hexose, both a hexose and a pentose, and combinations thereof.

**[0020]** In various embodiments, the invention is an aqueous broth obtained from a culture system according to any one of claims, wherein said aqueous broth comprises i) a concentration of 3-hydroxypropionate selected from greater than 5 g/L, greater than 10 g/L, greater than 15 g/L, greater than 20 g/L, greater than 25 g/L, greater than 30 g/L, greater than 35 g/L, greater than 40 g/L, greater than 50 g/L, greater than 60 g/L, greater than 70 g/L, greater than 80 g/L, greater than 90 g/L, or greater than 100 g/L 3-hydroxypropionate; and ii) a concentration of 1,3-propanediol selected from less than 30 g/L; less than 20 g/L; less than 10 g/L; less than 5 g/L; less than 1 g/L; or less than 0.5 g/L. In some aspects, the aqueous broth comprises an amount of biomass selected from less than 20 gDCW/L biomass, less than 15 gDCW/L biomass, less than 10 gDCW/L biomass, less than 5 gDCW/L biomass or less than 1 gDCW/L biomass. Alternatively, the aqueous broth according to the invention is such that the 3-HP/succinate ratio (g3-HP/g succinate) is greater than 3, greater than 10 greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200. In various aspects, the 3-HP/fumarate ratio (g3-HP/g fumarate) is greater than 3, greater than 10 greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/glycerol ratio (g3-HP/g glycerol) is greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/acetate ratio (g3-HP/g acetate) is greater than 1.5, greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/alanine ratio (g3-HP/g alanine)

is greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/beta-alanine ratio (g3-HP/g beta-alanine) is greater than 1.5, greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/glutamate ratio (g3-HP/g glutamate) is greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/glutamine ratio (g3-HP/g glutamine) is greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/3-hydroxypropionaldehyde ratio (g3-HP/g 3-hydroxypropionaldehyde) is greater than 1.5, greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, or the 3-HP/1,3-propanediol ratio (g3-HP/g 1,3-propanediol) is greater than 1.5, greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200, and/or the 3-HP/lactate ratio (g3-HP/g lactate) is greater than 3, greater than 10, greater than 30, greater than 60, greater than 100, greater than 150 or greater than 200.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0021]** The novel features of the invention are set forth with particularity in the appended claims. A better understanding of the features and advantages of the present invention will be obtained by reference to the following detailed description that sets forth illustrative embodiments, in which the principles of the invention are utilized, and the accompanying drawings of which:

**[0022]** FIG. 1 depicts a CLUSTAL 2.0.12 multiple sequence alignment of Carbonic Anhydrase Polypeptides.

**[0023]** FIG. 2 depicts the reaction catalyzed by an oxaloacetate alpha-decarboxylase enzyme.

**[0024]** FIG. 3A depicts the natural mixed acid fermentation routes in *E. coli*

**[0025]** FIG. 3B depicts the proposed 3-HP fermentation pathway of the present invention

**[0026]** FIG. 4 depicts metabolic pathways of a microorganism related to aspects of the present invention, more particularly related to 3-HP production, with gene names of *E. coli* shown at certain enzymatic steps, the latter for example and not meant to be limiting.

**[0027]** FIG. 5 depicts the reaction catalyzed by the YdfG enzyme.

**[0028]** FIG. 6A depicts diaper manufacture.

**[0029]** FIG. 6B depicts diaper manufacture and diaper wear.

**[0030]** FIG. 7 provides a schematic of processing from biomass to polymerized acrylic acid.

**[0031]** FIG. 8 is a Western blot of purified Kgd, Oad, and Pdc proteins

**[0032]** FIG. 9 are the results of GC-MS analysis of 3-HP production by various Oad-2 candidate enzymes

**[0033]** Tables also are provided herein and are part of the specification.

#### DETAILED DESCRIPTION OF THE INVENTION

**[0034]** The present invention is related to various production methods and/or genetically modified microorganisms that have utility for fermentative production of 3-hydroxypropionic acid ("3-HP", CAS No. 503-66-2), which may be converted to a number of basic building blocks for polymers

used in a wide range of industrial and consumer products. The present invention also is directed to methods of making 3-HP that utilize populations of these microorganisms in vessels, and to systems for chemical production that employ these microorganisms and methods. As noted herein, various aspects of the present invention are directed to a microorganism cell comprises a metabolic pathway from oxaloacetate to malonate semialdehyde, and also in various embodiments the capability to convert malonate semialdehyde to 3-HP, whether native and/or provided by a heterologous nucleic acid sequence encoding a protein having such activity.

**[0035]** Functional variants of an oxaloacetate alpha-decarboxylase/oxaloacetate alpha-decarboxylase are provided herein, with a demonstration of improved enzymatic conversion using these variant forms. In particular, three identified functional variant forms of an oxaloacetate alpha-decarboxylase from *Leuconostoc mesenteroides* are shown to have increased enzymatic activity. These may be provided in microorganisms that may also be provided with other genetic modifications described herein, resulting in improved capacity to produce 3-HP, which thereafter is converted to other chemicals, including acrylic acid, which is utilized for a number of industrial and consumer products.

**[0036]** Further as to the microorganisms of the present invention, in various embodiments additional genetic modifications may be made, such as to 1) increase intracellular bicarbonate levels, such as by increasing carbonic anhydrase, 2) increase enzymatic activity of NADPH-dependent transhydrogenase.

**[0037]** Additionally, genetic modifications for increasing tolerance may be combined with the present invention. Moreover, genetic modifications to increase expression and/or enzymatic activity of carbonic anhydrase and/or cyanase may provide dual-functions to advantageously improve both 3-HP production and 3-HP tolerance.

**[0038]** Other additional genetic modifications are disclosed herein for various embodiments.

#### Definitions

**[0039]** As used in the specification and the claims, the singular forms “a,” “an,” and “the” include plural referents unless the context clearly dictates otherwise. Thus, for example, reference to an “expression vector” includes a single expression vector as well as a plurality of expression vectors, either the same (e.g., the same operon) or different; reference to “microorganism” includes a single microorganism as well as a plurality of microorganisms; and the like.

**[0040]** As used herein, dry cell weight (DCW) for *E. coli* strains is calculated as 0.33 times the measured OD<sub>600</sub> value, based on baseline DCW to OD<sub>600</sub> determinations.

**[0041]** As used herein, “reduced enzymatic activity,” “reducing enzymatic activity,” and the like is meant to indicate that a microorganism cell’s, or an isolated enzyme, exhibits a lower level of activity than that measured in a comparable cell of the same species or its native enzyme. That is, enzymatic conversion of the indicated substrate(s) to indicated product(s) under known standard conditions for that enzyme is at least 10, at least 20, at least 30, at least 40, at least 50, at least 60, at least 70, at least 80, or at least 90 percent less than the enzymatic activity for the same biochemical conversion by a native (non-modified) enzyme under a standard specified condition. This term also can include elimination of that enzymatic activity. A cell having reduced enzymatic activity of an enzyme can be identified using any method

known in the art. For example, enzyme activity assays can be used to identify cells having reduced enzyme activity. See, for example, *Enzyme Nomenclature*, Academic Press, Inc., New York 2007.

**[0042]** The term “heterologous DNA,” “heterologous nucleic acid sequence,” and the like as used herein refers to a nucleic acid sequence wherein at least one of the following is true: (a) the sequence of nucleic acids is foreign to (i.e., not naturally found in) a given host microorganism; (b) the sequence may be naturally found in a given host microorganism, but in an unnatural (e.g., greater than expected) amount; or (c) the sequence of nucleic acids comprises two or more subsequences that are not found in the same relationship to each other in nature. For example, regarding instance (c), a heterologous nucleic acid sequence that is recombinantly produced will have two or more sequences from unrelated genes arranged to make a new functional nucleic acid.

**[0043]** The term “heterologous” is intended to include the term “exogenous” as the latter term is generally used in the art. With reference to the host microorganism’s genome prior to the introduction of a heterologous nucleic acid sequence, the nucleic acid sequence that codes for the enzyme is heterologous (whether or not the heterologous nucleic acid sequence is introduced into that genome).

**[0044]** As used herein, the term “gene disruption,” or grammatical equivalents thereof (and including “to disrupt enzymatic function,” “disruption of enzymatic function,” and the like), is intended to mean a genetic modification to a microorganism that renders the encoded gene product as having a reduced polypeptide activity compared with polypeptide activity in or from a microorganism cell not so modified. The genetic modification can be, for example, deletion of the entire gene, deletion or other modification of a regulatory sequence required for transcription or translation, deletion of a portion of the gene which results in a truncated gene product (e.g., enzyme) or by any of various mutation strategies that reduces activity (including to no detectable activity level) the encoded gene product. A disruption may broadly include a deletion of all or part of the nucleic acid sequence encoding the enzyme, and also includes, but is not limited to other types of genetic modifications, e.g., introduction of stop codons, frame shift mutations, introduction or removal of portions of the gene, and introduction of a degradation signal, those genetic modifications affecting mRNA transcription levels and/or stability, and altering the promoter or repressor upstream of the gene encoding the enzyme.

**[0045]** In various contexts, a gene disruption is taken to mean any genetic modification to the DNA, mRNA encoded from the DNA, and the corresponding amino acid sequence that results in reduced polypeptide activity. Many different methods can be used to make a cell having reduced polypeptide activity. For example, a cell can be engineered to have a disrupted regulatory sequence or polypeptide-encoding sequence using common mutagenesis or knock-out technology. See, e.g., *Methods in Yeast Genetics* (1997 edition), Adams et al., Cold Spring Harbor Press (1998). One particularly useful method of gene disruption is complete gene deletion because it reduces or eliminates the occurrence of genetic reversions in the genetically modified microorganisms of the invention. Accordingly, a disruption of a gene whose product is an enzyme thereby disrupts enzymatic function. Alternatively, antisense technology can be used to reduce the activity of a particular polypeptide. For example, a cell can be engineered to contain a cDNA that encodes an antisense molecule

that prevents a polypeptide from being translated. Further, gene silencing can be used to reduce the activity of a particular polypeptide.

**[0046]** The term “antisense molecule” as used herein encompasses any nucleic acid molecule or nucleic acid analog (e.g., peptide nucleic acids) that contains a sequence that corresponds to the coding strand of an endogenous polypeptide. An antisense molecule also can have flanking sequences (e.g., regulatory sequences). Thus, antisense molecules can be ribozymes or antisense oligonucleotides.

**[0047]** As used herein, a ribozyme can have any general structure including, without limitation, hairpin, hammerhead, or axhead structures, provided the molecule cleaves RNA.

**[0048]** The term “reduction” or “to reduce” when used in such phrase and its grammatical equivalents are intended to encompass a complete elimination of such conversion(s).

**[0049]** Bio-production, as used herein, may be aerobic, microaerobic, or anaerobic. Also as used herein, the terms “production” and “bio-production” are used interchangeably when referring to microbial synthesis of 3-HP.

**[0050]** As used herein, the language “sufficiently homologous” refers to proteins or portions thereof that have amino acid sequences that include a minimum number of identical or equivalent amino acid residues when compared to an amino acid sequence of the amino acid sequences provided in this application (including the SEQ ID Nos./sequence listings) such that the protein or portion thereof is able to achieve the respective enzymatic reaction and/or other function. To determine whether a particular protein or portion thereof is sufficiently homologous may be determined by an assay of enzymatic activity, such as those commonly known in the art.

**[0051]** Descriptions and methods for sequence identity and homology are intended to be exemplary and it is recognized that these concepts are well-understood in the art. Further, it is appreciated that nucleic acid sequences may be varied and still encode an enzyme or other polypeptide exhibiting a desired functionality, and such variations are within the scope of the present invention.

**[0052]** Further to nucleic acid sequences, “hybridization” refers to the process in which two single-stranded polynucleotides bind non-covalently to form a stable double-stranded polynucleotide. The term “hybridization” may also refer to triple-stranded hybridization. The resulting (usually) double-stranded polynucleotide is a “hybrid” or “duplex.” “Hybridization conditions” will typically include salt concentrations of less than about 1M, more usually less than about 500 mM and less than about 200 mM. Hybridization temperatures can be as low as 5° C., but are typically greater than 22° C., more typically greater than about 30° C., and often are in excess of about 37° C. Hybridizations are usually performed under stringent conditions, i.e. conditions under which a probe will hybridize to its target subsequence. Stringent conditions are sequence-dependent and are different in different circumstances. Longer fragments may require higher hybridization temperatures for specific hybridization. As other factors may affect the stringency of hybridization, including base composition and length of the complementary strands, presence of organic solvents and extent of base mismatching, the combination of parameters is more important than the absolute measure of any one alone. Generally, stringent conditions are selected to be about 5° C. lower than the  $T_m$  for the specific sequence at a defined ionic strength and pH. Exemplary stringent conditions include salt concentration of at least 0.01 M to no more than 1 M Na ion concentration (or other salts) at a pH

7.0 to 8.3 and a temperature of at least 25° C. For example, conditions of 5×SSPE (750 mM NaCl, 50 mM NaPhosphate, 5 mM EDTA, pH 7.4) and a temperature of 25-30° C. are suitable for allele-specific probe hybridizations. For stringent conditions, see for example, Sambrook and Russell and Anderson “Nucleic Acid Hybridization” 1<sup>st</sup> Ed., BIOS Scientific Publishers Limited (1999), which are hereby incorporated by reference for hybridization protocols. “Hybridizing specifically to” or “specifically hybridizing to” or like expressions refer to the binding, duplexing, or hybridizing of a molecule substantially to or only to a particular nucleotide sequence or sequences under stringent conditions when that sequence is present in a complex mixture (e.g., total cellular) DNA or RNA.

**[0053]** The term “identified enzymatic functional variant” means a polypeptide that is determined to possess an enzymatic activity and specificity of an enzyme of interest but which has an amino acid sequence different from such enzyme of interest. A corresponding “variant nucleic acid sequence” may be constructed that is determined to encode such an identified enzymatic functional variant. For a particular purpose, such as increased tolerance to 3-HP via genetic modification to increase enzymatic conversion at one or more of the enzymatic conversion steps of the 3HPTGC in a microorganism, one or more genetic modifications may be made to provide one or more heterologous nucleic acid sequence(s) that encode one or more identified 3HPTGC enzymatic functional variant(s). That is, each such nucleic acid sequence encodes a polypeptide that is not exactly the known polypeptide of an enzyme of the 3HPTGC, but which nonetheless is shown to exhibit enzymatic activity of such enzyme. Such nucleic acid sequence, and the polypeptide it encodes, may not fall within a specified limit of homology or identity yet by its provision in a cell nonetheless provide for a desired enzymatic activity and specificity. The ability to obtain such variant nucleic acid sequences and identified enzymatic functional variants is supported by recent advances in the states of the art in bioinformatics and protein engineering and design, including advances in computational, predictive and high-throughput methodologies. Functional variants more generally include enzymatic functional variants, and the nucleic acids sequences that encode them, as well as variants of non-enzymatic polypeptides, wherein the variant exhibits the function of the original (target) sequence.

**[0054]** The use of the phrase “segment of interest” is meant to include both a gene and any other nucleic acid sequence segment of interest. One example of a method used to obtain a segment of interest is to acquire a culture of a microorganism, where that microorganism’s genome includes the gene or nucleic acid sequence segment of interest.

**[0055]** When the genetic modification of a gene product, i.e., an enzyme, is referred to herein, including the claims, it is understood that the genetic modification is of a nucleic acid sequence, such as or including the gene, that normally encodes the stated gene product, i.e., the enzyme.

**[0056]** In some embodiments a truncated respective polypeptide has at least about 90% of the full length of a polypeptide encoded by a nucleic acid sequence encoding the respective native enzyme, and more particularly at least 95% of the full length of a polypeptide encoded by a nucleic acid sequence encoding the respective native enzyme. By a polypeptide having an amino acid sequence at least, for example, 95% “identical” to a reference amino acid sequence of a polypeptide is intended that the amino acid sequence of

the claimed polypeptide is identical to the reference sequence except that the claimed polypeptide sequence can include up to five amino acid alterations per each 100 amino acids of the reference amino acid of the polypeptide. In other words, to obtain a polypeptide having an amino acid sequence at least 95% identical to a reference amino acid sequence, up to 5% of the amino acid residues in the reference sequence can be deleted or substituted with another amino acid, or a number of amino acids up to 5% of the total amino acid residues in the reference sequence can be inserted into the reference sequence. These alterations of the reference sequence can occur at the amino or carboxy terminal positions of the reference amino acid sequence or anywhere between those terminal positions, interspersed either individually among residues in the reference sequence or in one or more contiguous groups within the reference sequence. In other embodiments truncation may be more substantial, as described elsewhere herein.

**[0057]** Species and other phylogenetic identifications are according to the classification known to a person skilled in the art of microbiology.

**[0058]** Where methods and steps described herein indicate certain events occurring in certain order, those of ordinary skill in the art will recognize that the ordering of certain steps may be modified and that such modifications are in accordance with the variations of the invention. Additionally, certain steps may be performed concurrently in a parallel process when possible, as well as performed sequentially.

**[0059]** Prophetic examples provided herein are meant to be broadly exemplary and not limiting in any way. This applies to the examples regarding separation and purification of 3-HP, and conversions of 3-HP to downstream compounds, since there are numerous possible approaches to such steps and conversions, including those disclosed in references recited and incorporated herein.

**[0060]** The meaning of abbreviations is as follows: “C” means Celsius or degrees Celsius, as is clear from its usage, DCW means dry cell weight, “s” means second(s), “min” means minute(s), “h,” “hr,” or “hrs” means hour(s), “psi” means pounds per square inch, “nm” means nanometers, “d” means day(s), “μL” or “uL” or “ul” means microliter(s), “mL” means milliliter(s), “L” means liter(s), “mm” means millimeter(s), “nm” means nanometers, “mM” means millimolar, “μM” or “uM” means micromolar, “M” means molar, “mmol” means millimole(s), “μmol” or “uMol” means micromole(s), “g” means gram(s), “μg” or “ug” means microgram(s) and “ng” means nanogram(s), “PCR” means polymerase chain reaction, “OD” means optical density, “OD<sub>600</sub>” means the optical density measured at a photon wavelength of 600 nm, “kDa” means kilodaltons, “g” means the gravitation constant, “bp” means base pair(s), “kbp” means kilobase pair(s), “% w/v” means weight/volume percent, “% v/v” means volume/volume percent, “IPTG” means isopropyl-μ-D-thiogalactopyranoside, “RBS” means ribosome binding site, “rpm” means revolutions per minute, “HPLC” means high performance liquid chromatography, and “GC” means gas chromatography. As disclosed herein, “3-HP” means 3-hydroxypropionic acid and “3HPTGC” means the 3-HP toleragenic complex. Also, 10<sup>5</sup> and the like are taken to mean 10<sup>5</sup> and the like.

**[0061]** I. Carbon Sources

**[0062]** Bio-production media, which is used in the present invention with recombinant microorganisms having a biosynthetic pathway for 3-HP, must contain suitable carbon sources

or substrates for the intended metabolic pathways. Suitable substrates may include, but are not limited to, monosaccharides such as glucose and fructose, oligosaccharides such as lactose or sucrose, polysaccharides such as starch or cellulose or mixtures thereof and unpurified mixtures from renewable feedstocks such as cheese whey permeate, cornsteep liquor, sugar beet molasses, and barley malt. Additionally the carbon substrate may also be one-carbon substrates such as carbon dioxide, carbon monoxide, or methanol for which metabolic conversion into key biochemical intermediates has been demonstrated. In addition to one and two carbon substrates methylotrophic organisms are also known to utilize a number of other carbon containing compounds such as methylamine, glucosamine and a variety of amino acids for metabolic activity.

**[0063]** Although it is contemplated that all of the above mentioned carbon substrates and mixtures thereof are suitable in the present invention as a carbon source, common carbon substrates used as carbon sources are glucose, fructose, and sucrose, as well as mixtures of any of these sugars. Other suitable substrates include xylose, arabinose, other cellulose-based C-5 sugars, high-fructose corn syrup, and various other sugars and sugar mixtures as are available commercially. Sucrose may be obtained from feedstocks such as sugar cane, sugar beets, cassava, bananas or other fruit, and sweet *sorghum*. Glucose and dextrose may be obtained through saccharification of starch based feedstocks including grains such as corn, wheat, rye, barley, and oats. Also, in some embodiments all or a portion of the carbon source may be glycerol. Alternatively, glycerol may be excluded as an added carbon source.

**[0064]** In one embodiment, the carbon source is selected from glucose, fructose, sucrose, dextrose, lactose, glycerol, and mixtures thereof. Variously, the amount of these components in the carbon source may be greater than about 50%, greater than about 60%, greater than about 70%, greater than about 80%, greater than about 90%, or more, up to 100% or essentially 100% of the carbon source.

**[0065]** In addition, methylotrophic organisms are known to utilize a number of other carbon containing compounds such as methylamine, glucosamine and a variety of amino acids for metabolic activity. For example, methylotrophic yeast are known to utilize the carbon from methylamine to form trehalose or glycerol (Hellion et al., *Microb. Growth CI Compd. (Int. Symp.)*, 7th (1993), 415-32. Editor(s): Murrell, J. Collin; Kelly, Don P. Publisher: Intercept, Andover, UK). Similarly, various species of *Candida* will metabolize alanine or oleic acid (Sulter et al., *Arch. Microbiol.* 153:485-489 (1990)). Hence it is contemplated that the source of carbon utilized in embodiments of the present invention may encompass a wide variety of carbon-containing substrates.

**[0066]** In addition, fermentable sugars may be obtained from cellulosic and lignocellulosic biomass through processes of pretreatment and saccharification, as described, for example, in U.S. Patent Publication No. 2007/0031918A1, which is herein incorporated by reference. Biomass refers to any cellulosic or lignocellulosic material and includes materials comprising cellulose, and optionally further comprising hemicellulose, lignin, starch, oligosaccharides and/or monosaccharides. Biomass may also comprise additional components, such as protein and/or lipid. Biomass may be derived from a single source, or biomass can comprise a mixture derived from more than one source; for example, biomass could comprise a mixture of corn cobs and corn

stover, or a mixture of grass and leaves. Biomass includes, but is not limited to, bioenergy crops, agricultural residues, municipal solid waste, industrial solid waste, sludge from paper manufacture, yard waste, wood and forestry waste. Examples of biomass include, but are not limited to, corn grain, corn cobs, crop residues such as corn husks, corn stover, grasses, wheat, wheat straw, barley, barley straw, hay, rice straw, switchgrass, waste paper, sugar cane bagasse, *sorghum*, soy, components obtained from milling of grains, trees, branches, roots, leaves, wood chips, sawdust, shrubs and bushes, vegetables, fruits, flowers and animal manure. Any such biomass may be used in a bio-production method or system to provide a carbon source. Various approaches to breaking down cellulosic biomass to mixtures of more available and utilizable carbon molecules, including sugars, include: heating in the presence of concentrated or dilute acid (e.g., <1% sulfuric acid); treating with ammonia; treatment with ionic salts; enzymatic degradation; and combinations of these. These methods normally follow mechanical separation and milling, and are followed by appropriate separation processes.

[0067] In various embodiments, any of a wide range of sugars, including, but not limited to sucrose, glucose, xylose, cellulose or hemicellulose, are provided to a microorganism, such as in an industrial system comprising a reactor vessel in which a defined media (such as a minimal salts media including but not limited to M9 minimal media, potassium sulfate minimal media, yeast synthetic minimal media and many others or variations of these), an inoculum of a microorganism providing one or more of the 3-HP biosynthetic pathway alternatives, and the a carbon source may be combined. The carbon source enters the cell and is catabolized by well-known and common metabolic pathways to yield common metabolic intermediates, including phosphoenolpyruvate (PEP). (See Molecular Biology of the Cell, 3rd Ed., B. Alberts et al. Garland Publishing, New York, 1994, pp. 42-45, 66-74, incorporated by reference for the teachings of basic metabolic catabolic pathways for sugars; Principles of Biochemistry, 3rd Ed., D. L. Nelson & M. M. Cox, Worth Publishers, New York, 2000, pp 527-658, incorporated by reference for the teachings of major metabolic pathways; and Biochemistry, 4th Ed., L. Stryer, W. H. Freeman and Co., New York, 1995, pp. 463-650, also incorporated by reference for the teachings of major metabolic pathways.)

[0068] Bio-based carbon can be distinguished from petroleum-based carbon according to a variety of methods, including without limitation ASTM D6866, or various other techniques. For example, carbon-14 and carbon-12 ratios differ in bio-based carbon sources versus petroleum-based sources, where higher carbon-14 ratios are found in bio-based carbon sources. In various embodiments, the carbon source is not petroleum-based, or is not predominantly petroleum based. In various embodiments, the carbon source is greater than about 50% non-petroleum based, greater than about 60% non-petroleum based, greater than about 70% non-petroleum based, greater than about 80% non-petroleum based, greater than about 90% non-petroleum based, or more. In various embodiments, the carbon source has a carbon-14 to carbon-12 ratio of about  $1.0 \times 10^{-14}$  or greater.

[0069] Various components may be excluded from the carbon source. For example, in some embodiments, acrylic acid, 1,4-butanediol, and/or glycerol are excluded or essentially excluded from the carbon source. As such, the carbon source according to some embodiments of the invention may be less

than about 50% glycerol, less than about 40% glycerol, less than about 30% glycerol, less than about 20% glycerol, less than about 10% glycerol, less than about 5% glycerol, less than about 1% glycerol, or less. For example, the carbon source may be essentially glycerol-free. By essentially glycerol-free is meant that any glycerol that may be present in a residual amount does not contribute substantially to the production of the target chemical compound.

[0070] II. Microorganisms

[0071] Features as described and claimed herein may be provided in a microorganism selected from the listing herein, or another suitable microorganism, that also comprises one or more natural, introduced, or enhanced 3-HP bio-production pathways. Thus, in some embodiments the microorganism comprises an endogenous 3-HP production pathway (which may, in some such embodiments, be enhanced), whereas in other embodiments the microorganism does not comprise an endogenous 3-HP production pathway.

[0072] Varieties of these genetically modified microorganisms may comprise genetic modifications and/or other system alterations as may be described in other patent applications of one or more of the present inventor(s) and/or subject to assignment to the owner of the present patent application.

[0073] The examples describe specific modifications and evaluations to certain bacterial and yeast microorganisms. The scope of the invention is not meant to be limited to such species, but to be generally applicable to a wide range of suitable microorganisms. Generally, a microorganism used for the present invention may be selected from bacteria, cyanobacteria, filamentous fungi and yeasts.

[0074] For some embodiments, microbial hosts initially selected for 3-HP toleragenic bio-production should also utilize sugars including glucose at a high rate. Most microbes are capable of utilizing carbohydrates. However, certain environmental microbes cannot utilize carbohydrates to high efficiency, and therefore would not be suitable hosts for such embodiments that are intended for glucose or other carbohydrates as the principal added carbon source.

[0075] As the genomes of various species become known, the present invention easily may be applied to an ever-increasing range of suitable microorganisms. Further, given the relatively low cost of genetic sequencing, the genetic sequence of a species of interest may readily be determined to make application of aspects of the present invention more readily obtainable (based on the ease of application of genetic modifications to an organism having a known genomic sequence).

[0076] More particularly, based on the various criteria described herein, suitable microbial hosts for the bio-production of 3-HP that comprise tolerance aspects provided herein generally may include, but are not limited to, any gram negative organisms, more particularly a member of the family Enterobacteriaceae, such as *E. coli*, or *Oligotropha carboxidovorans*, or *Pseudomonas* sp.; any gram positive microorganism, for example *Bacillus subtilis*, *Lactobacillus* sp. or *Lactococcus* sp.; a yeast, for example *Saccharomyces cerevisiae*, *Pichia pastoris* or *Pichia stipitis*; and other groups or microbial species. More particularly, suitable microbial hosts for the bio-production of 3-HP generally include, but are not limited to, members of the genera *Clostridium*, *Zymomonas*, *Escherichia*, *Salmonella*, *Rhodococcus*, *Pseudomonas*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Alcaligenes*, *Klebsiella*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, *Brevibacterium*, *Pichia*, *Candida*, *Hansenula* and *Saccharomyces*. Hosts that may be particularly of interest include: *Oligotro-*

*pha carboxidovorans* (such as strain OM5), *Escherichia coli*, *Alcaligenes eutrophus* (*Cupriavidus necator*), *Bacillus licheniformis*, *Paenibacillus macerans*, *Rhodococcus erythropolis*, *Pseudomonas putida*, *Lactobacillus plantarum*, *Enterococcus faecium*, *Enterococcus gallinarium*, *Enterococcus faecalis*, *Bacillus subtilis* and *Saccharomyces cerevisiae*.

[0077] More particularly, suitable microbial hosts for the bio-production of 3-HP generally include, but are not limited to, members of the genera *Clostridium*, *Zymomonas*, *Escherichia*, *Salmonella*, *Rhodococcus*, *Pseudomonas*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Alcaligenes*, *Klebsiella*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, *Brevibacterium*, *Pichia*, *Candida*, *Hansenula* and *Saccharomyces*.

[0078] Hosts that may be particularly of interest include: *Oligotropha carboxidovorans* (such as strain OM5<sup>T</sup>), *Escherichia coli*, *Alcaligenes eutrophus* (*Cupriavidus necator*), *Bacillus licheniformis*, *Paenibacillus macerans*, *Rhodococcus erythropolis*, *Pseudomonas putida*, *Lactobacillus plantarum*, *Enterococcus faecium*, *Enterococcus gallinarium*, *Enterococcus faecalis*, *Bacillus subtilis* and *Saccharomyces cerevisiae*. Also, any of the known strains of these species may be utilized as a starting microorganism, as may any of the following species including respective strains thereof—*Cupriavidus basilensis*, *Cupriavidus campinensis*, *Cupriavidus gilardi*, *Cupriavidus laharsis*, *Cupriavidus metallidurans*, *Cupriavidus oxalaticus*, *Cupriavidus pauculus*, *Cupriavidus pinatubonensis*, *Cupriavidus respiraculi*, and *Cupriavidus taiwanensis*.

[0079] In some embodiments, the recombinant microorganism is a gram-negative bacterium. In some embodiments, the recombinant microorganism is selected from the genera *Zymomonas*, *Escherichia*, *Pseudomonas*, *Alcaligenes*, and *Klebsiella*. In some embodiments, the recombinant microorganism is selected from the species *Escherichia coli*, *Cupriavidus necator*, *Oligotropha carboxidovorans*, and *Pseudomonas putida*. In some embodiments, the recombinant microorganism is an *E. coli* strain.

[0080] In some embodiments, the recombinant microorganism is a gram-positive bacterium. In some embodiments, the recombinant microorganism is selected from the genera *Clostridium*, *Salmonella*, *Rhodococcus*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, and *Brevibacterium*. In some embodiments, the recombinant microorganism is selected from the species *Bacillus licheniformis*, *Paenibacillus macerans*, *Rhodococcus erythropolis*, *Lactobacillus plantarum*, *Enterococcus faecium*, *Enterococcus gallinarium*, *Enterococcus faecalis*, and *Bacillus subtilis*. In particular embodiments, the recombinant microorganism is a *B. subtilis* strain.

[0081] In some embodiments, the recombinant microorganism is a yeast. In some embodiments, the recombinant microorganism is selected from the genera *Pichia*, *Candida*, *Hansenula* and *Saccharomyces*. In particular embodiments, the recombinant microorganism is *Saccharomyces cerevisiae*.

[0082] III. Media and Culture Conditions

[0083] In addition to an appropriate carbon source, such as selected from one of the herein-disclosed types, bio-production media must contain suitable minerals, salts, cofactors, buffers and other components, known to those skilled in the art, suitable for the growth of the cultures and promotion of the enzymatic pathway necessary for 3-HP production, or other products made under the present invention.

[0084] Another aspect of the invention regards media and culture conditions that comprise genetically modified microorganisms of the invention and optionally supplements.

[0085] Typically cells are grown at a temperature in the range of about 25° C. to about 40° C. in an appropriate medium, as well as up to 70° C. for thermophilic microorganisms. Suitable growth media in the present invention are common commercially prepared media such as Luria Bertani (LB) broth, M9 minimal media, Sabouraud Dextrose (SD) broth, Yeast medium (YM) broth, (Ymin) yeast synthetic minimal media, and minimal media as described herein, such as M9 minimal media. Other defined or synthetic growth media may also be used, and the appropriate medium for growth of the particular microorganism will be known by one skilled in the art of microbiology or bio-production science. In various embodiments a minimal media may be developed and used that does not comprise, or that has a low level of addition of various components, for example less than 10, 5, 2 or 1 g/L of a complex nitrogen source including but not limited to yeast extract, peptone, tryptone, soy flour, corn steep liquor, or casein. These minimal medias may also have limited supplementation of vitamin mixtures including biotin, vitamin B12 and derivatives of vitamin B12, thiamin, pantothenate and other vitamins. Minimal medias may also have limited simple inorganic nutrient sources containing less than 28, 17, or 2.5 mM phosphate, less than 25 or 4 mM sulfate, and less than 130 or 50 mM total nitrogen.

[0086] Bio-production media, which is used in embodiments of the present invention with genetically modified microorganisms, must contain suitable carbon substrates for the intended metabolic pathways. As described hereinbefore, suitable carbon substrates include carbon monoxide, carbon dioxide, and various monomeric and oligomeric sugars.

[0087] Suitable pH ranges for the bio-production are between pH 3.0 to pH 10.0, where pH 6.0 to pH 8.0 is a typical pH range for the initial condition. However, the actual culture conditions for a particular embodiment are not meant to be limited by these pH ranges.

[0088] Bio-productions may be performed under aerobic, microaerobic, or anaerobic conditions, with or without agitation.

[0089] The amount of 3-HP or other product(s) produced in a bio-production media generally can be determined using a number of methods known in the art, for example, high performance liquid chromatography (HPLC), gas chromatography (GC), or GC/Mass Spectroscopy (MS). Specific HPLC methods for the specific examples are provided herein.

[0090] IV. Bio-Production Reactors and Systems

[0091] Fermentation systems utilizing methods and/or compositions according to the invention are also within the scope of the invention.

[0092] Any of the recombinant microorganisms as described and/or referred to herein may be introduced into an industrial bio-production system where the microorganisms convert a carbon source into 3-HP in a commercially viable operation. The bio-production system includes the introduction of such a recombinant microorganism into a bioreactor vessel, with a carbon source substrate and bio-production media suitable for growing the recombinant microorganism, and maintaining the bio-production system within a suitable temperature range (and dissolved oxygen concentration range if the reaction is aerobic or microaerobic) for a suitable time to obtain a desired conversion of a portion of the substrate molecules to 3-HP. Industrial bio-production systems



and their operation are well-known to those skilled in the arts of chemical engineering and bioprocess engineering.

**[0093]** Bio-productions may be performed under aerobic, microaerobic, or anaerobic conditions, with or without agitation. The operation of cultures and populations of microorganisms to achieve aerobic, microaerobic and anaerobic conditions are known in the art, and dissolved oxygen levels of a liquid culture comprising a nutrient media and such microorganism populations may be monitored to maintain or confirm a desired aerobic, microaerobic or anaerobic condition. When syngas is used as a feedstock, aerobic, microaerobic, or anaerobic conditions may be utilized. When sugars are used, anaerobic, aerobic or microaerobic conditions can be implemented in various embodiments.

**[0094]** Any of the recombinant microorganisms as described and/or referred to herein may be introduced into an industrial bio-production system where the microorganisms convert a carbon source into 3-HP, and optionally in various embodiments also to one or more downstream compounds of 3-HP in a commercially viable operation. The bio-production system includes the introduction of such a recombinant microorganism into a bioreactor vessel, with a carbon source substrate and bio-production media suitable for growing the recombinant microorganism, and maintaining the bio-production system within a suitable temperature range (and dissolved oxygen concentration range if the reaction is aerobic or microaerobic) for a suitable time to obtain a desired conversion of a portion of the substrate molecules to 3-HP.

**[0095]** In various embodiments, syngas components or sugars are provided to a microorganism, such as in an industrial system comprising a reactor vessel in which a defined media (such as a minimal salts media including but not limited to M9 minimal media, potassium sulfate minimal media, yeast synthetic minimal media and many others or variations of these), an inoculum of a microorganism providing an embodiment of the biosynthetic pathway(s) taught herein, and the carbon source may be combined. The carbon source enters the cell and is catabolized by well-known and common metabolic pathways to yield common metabolic intermediates, including phosphoenolpyruvate (PEP). (See *Molecular Biology of the Cell*, 3<sup>rd</sup> Ed., B. Alberts et al. Garland Publishing, New York, 1994, pp. 42-45, 66-74, incorporated by reference for the teachings of basic metabolic catabolic pathways for sugars; *Principles of Biochemistry*, 3<sup>rd</sup> Ed., D. L. Nelson & M. M. Cox, Worth Publishers, New York, 2000, pp. 527-658, incorporated by reference for the teachings of major metabolic pathways; and *Biochemistry*, 4<sup>th</sup> Ed., L. Stryer, W. H. Freeman and Co., New York, 1995, pp. 463-650, also incorporated by reference for the teachings of major metabolic pathways.)

**[0096]** Further to types of industrial bio-production, various embodiments of the present invention may employ a batch type of industrial bioreactor. A classical batch bioreactor system is considered “closed” meaning that the composition of the medium is established at the beginning of a respective bio-production event and not subject to artificial alterations and additions during the time period ending substantially with the end of the bio-production event. Thus, at the beginning of the bio-production event the medium is inoculated with the desired organism or organisms, and bio-production is permitted to occur without adding anything to the system. Typically, however, a “batch” type of bio-production event is batch with respect to the addition of carbon source and attempts are often made at controlling factors such

as pH and oxygen concentration. In batch systems the metabolite and biomass compositions of the system change constantly up to the time the bio-production event is stopped. Within batch cultures cells moderate through a static lag phase to a high growth log phase and finally to a stationary phase where growth rate is diminished or halted. If untreated, cells in the stationary phase will eventually die. Cells in log phase generally are responsible for the bulk of production of a desired end product or intermediate.

**[0097]** A variation on the standard batch system is the fed-batch system. Fed-batch bio-production processes are also suitable in the present invention and comprise a typical batch system with the exception that the nutrients, including the substrate, are added in increments as the bio-production progresses. Fed-Batch systems are useful when catabolite repression is apt to inhibit the metabolism of the cells and where it is desirable to have limited amounts of substrate in the media. Measurement of the actual nutrient concentration in Fed-Batch systems may be measured directly, such as by sample analysis at different times, or estimated on the basis of the changes of measurable factors such as pH, dissolved oxygen and the partial pressure of waste gases such as CO<sub>2</sub>. Batch and fed-batch approaches are common and well known in the art and examples may be found in Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition (1989) Sinauer Associates, Inc., Sunderland, Mass., Deshpande, Mukund V., *Appl. Biochem. Biotechnol.*, 36:227, (1992), and *Biochemical Engineering Fundamentals*, 2<sup>nd</sup> Ed. J. E. Bailey and D. F. Ollis, McGraw Hill, New York, 1986, herein incorporated by reference for general instruction on bio-production.

**[0098]** Although embodiments of the present invention may be performed in batch mode, or in fed-batch mode, it is contemplated that the invention would be adaptable to continuous bio-production methods. Continuous bio-production is considered an “open” system where a defined bio-production medium is added continuously to a bioreactor and an equal amount of conditioned media is removed simultaneously for processing. Continuous bio-production generally maintains the cultures within a controlled density range where cells are primarily in log phase growth. Two types of continuous bioreactor operation include a chemostat, wherein fresh media is fed to the vessel while simultaneously removing an equal rate of the vessel contents. The limitation of this approach is that cells are lost and high cell density generally is not achievable. In fact, typically one can obtain much higher cell density with a fed-batch process. Another continuous bioreactor utilizes perfusion culture, which is similar to the chemostat approach except that the stream that is removed from the vessel is subjected to a separation technique which recycles viable cells back to the vessel. This type of continuous bioreactor operation has been shown to yield significantly higher cell densities than fed-batch and can be operated continuously. Continuous bio-production is particularly advantageous for industrial operations because it has less down time associated with draining, cleaning and preparing the equipment for the next bio-production event. Furthermore, it is typically more economical to continuously operate downstream unit operations, such as distillation, than to run them in batch mode.

**[0099]** Continuous bio-production allows for the modulation of one factor or any number of factors that affect cell growth or end product concentration. For example, one method will maintain a limiting nutrient such as the carbon

source or nitrogen level at a fixed rate and allow all other parameters to moderate. In other systems a number of factors affecting growth can be altered continuously while the cell concentration, measured by media turbidity, is kept constant. Methods of modulating nutrients and growth factors for continuous bio-production processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology and a variety of methods are detailed by Brock, supra.

**[0100]** It is contemplated that embodiments of the present invention may be practiced using either batch, fed-batch or continuous processes and that any known mode of bio-production would be suitable. It is contemplated that cells may be immobilized on an inert scaffold as whole cell catalysts and subjected to suitable bio-production conditions for 3-HP production, or be cultured in liquid media in a vessel, such as a culture vessel. Thus, embodiments used in such processes, and in bio-production systems using these processes, include a population of genetically modified microorganisms of the present invention, a culture system comprising such population in a media comprising nutrients for the population, and methods of making 3-HP and thereafter, a downstream product of 3-HP.

**[0101]** Embodiments of the invention include methods of making 3-HP in a bio-production system, some of which methods may include obtaining 3-HP after such bio-production event. For example, a method of making 3-HP may comprise: providing to a culture vessel a media comprising suitable nutrients; providing to the culture vessel an inoculum of a genetically modified microorganism comprising genetic modifications described herein such that the microorganism produces 3-HP from syngas and/or a sugar molecule; and maintaining the culture vessel under suitable conditions for the genetically modified microorganism to produce 3-HP.

**[0102]** It is within the scope of the present invention to produce, and to utilize in bio-production methods and systems, including industrial bio-production systems for production of 3-HP, a recombinant microorganism genetically engineered to modify one or more aspects effective to increase tolerance to 3-HP (and, in some embodiments, also 3-HP bio-production) by at least 20 percent over control microorganism lacking the one or more modifications.

**[0103]** In various embodiments, the invention is directed to a system for bioproduction of acrylic acid as described herein, said system comprising: a tank for saccharification of biomass; a line for passing the product of saccharification to a fermentation tank optionally via a pre-fermentation tank; a fermentation tank suitable for microorganism cell culture; a line for discharging contents from the fermentation tank to an extraction and/or separation vessel; an extraction and/or separation vessel suitable for removal of 3-hydroxypropionic acid from cell culture waste; a line for transferring 3-hydroxypropionic acid to a dehydration vessel; and a dehydration vessel suitable for conversion of 3-hydroxypropionic acid to acrylic acid. In various embodiments, the system includes one or more pre-fermentation tanks, distillation columns, centrifuge vessels, back extraction columns, mixing vessels, or combinations thereof.

**[0104]** The following published resources are incorporated by reference herein for their respective teachings to indicate the level of skill in these relevant arts, and as needed to support a disclosure that teaches how to make and use methods of industrial bio-production of 3-HP, or other product(s) produced under the invention, from sugar sources, and also

industrial systems that may be used to achieve such conversion with any of the recombinant microorganisms of the present invention (Biochemical Engineering Fundamentals, 2<sup>nd</sup> Ed. J. E. Bailey and D. F. Ollis, McGraw Hill, New York, 1986, entire book for purposes indicated and Chapter 9, pages 533-657 in particular for biological reactor design; Unit Operations of Chemical Engineering, 5<sup>th</sup> Ed., W. L. McCabe et al., McGraw Hill, New York 1993, entire book for purposes indicated, and particularly for process and separation technologies analyses; Equilibrium Staged Separations, P. C. Wankat, Prentice Hall, Englewood Cliffs, N.J. USA, 1988, entire book for separation technologies teachings).

**[0105]** V. Genetic Modifications, Nucleotide Sequences, and Amino Acid Sequences

**[0106]** Embodiments of the present invention may result from introduction of an expression vector into a host microorganism, wherein the expression vector contains a nucleic acid sequence coding for an enzyme that is, or is not, normally found in a host microorganism.

**[0107]** The ability to genetically modify a host cell is essential for the production of any genetically modified (recombinant) microorganism. The mode of gene transfer technology may be by electroporation, conjugation, transduction, or natural transformation. A broad range of host conjugative plasmids and drug resistance markers are available. The cloning vectors are tailored to the host organisms based on the nature of antibiotic resistance markers that can function in that host. Also, as disclosed herein, a genetically modified (recombinant) microorganism may comprise modifications other than via plasmid introduction, including modifications to its genomic DNA.

**[0108]** It has long been recognized in the art that some amino acids in amino acid sequences can be varied without significant effect on the structure or function of proteins. Variants included can constitute deletions, insertions, inversions, repeats, and type substitutions so long as the indicated enzyme activity is not significantly adversely affected. Guidance concerning which amino acid changes are likely to be phenotypically silent can be found, inter alia, in Bowie, J. U., et al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990). This reference is incorporated by reference for such teachings, which are, however, also generally known to those skilled in the art.

**[0109]** In various embodiments polypeptides obtained by the expression of the polynucleotide molecules of the present invention may have at least approximately 50%, 60%, 70%, 80%, 90%, 95%, 96%, 97%, 98%, 99% or 100% identity to one or more amino acid sequences encoded by the genes and/or nucleic acid sequences described herein for the 3-HP tolerance-related and biosynthesis pathways.

**[0110]** As a practical matter, whether any particular polypeptide is at least 50%, 60%, 70%, 80%, 85%, 90%, 92%, 95%, 96%, 97%, 98% or 99% identical to any reference amino acid sequence of any polypeptide described herein (which may correspond with a particular nucleic acid sequence described herein), such particular polypeptide sequence can be determined conventionally using known computer programs such the Bestfit program (Wisconsin Sequence Analysis Package, Version 8 for Unix, Genetics Computer Group, University Research Park, 575 Science Drive, Madison, Wis. 53711). When using Bestfit or any other sequence alignment program to determine whether a particular sequence is, for instance, 95% identical to a reference

sequence according to the present invention, the parameters are set such that the percentage of identity is calculated over the full length of the reference amino acid sequence and that gaps in homology of up to 5% of the total number of amino acid residues in the reference sequence are allowed.

**[0111]** For example, in a specific embodiment the identity between a reference sequence (query sequence, i.e., a sequence of the present invention) and a subject sequence, also referred to as a global sequence alignment, may be determined using the FASTDB computer program based on the algorithm of Brutlag et al. (Comp. App. Biosci. 6:237-245 (1990)). Preferred parameters for a particular embodiment in which identity is narrowly construed, used in a FASTDB amino acid alignment, are: Scoring Scheme=PAM (Percent Accepted Mutations) 0, k-tuple=2, Mismatch Penalty=1, Joining Penalty=20, Randomization Group Length=0, Cutoff Score=1, Window Size=sequence length, Gap Penalty=5, Gap Size Penalty=0.05, Window Size=500 or the length of the subject amino acid sequence, whichever is shorter. According to this embodiment, if the subject sequence is shorter than the query sequence due to N- or C-terminal deletions, not because of internal deletions, a manual correction is made to the results to take into consideration the fact that the FASTDB program does not account for N- and C-terminal truncations of the subject sequence when calculating global percent identity. For subject sequences truncated at the N- and C-termini, relative to the query sequence, the percent identity is corrected by calculating the number of residues of the query sequence that are lateral to the N- and C-terminal of the subject sequence, which are not matched/aligned with a corresponding subject residue, as a percent of the total bases of the query sequence. A determination of whether a residue is matched/aligned is determined by results of the FASTDB sequence alignment. This percentage is then subtracted from the percent identity, calculated by the FASTDB program using the specified parameters, to arrive at a final percent identity score. This final percent identity score is what is used for the purposes of this embodiment. Only residues to the N- and C-termini of the subject sequence, which are not matched/aligned with the query sequence, are considered for the purposes of manually adjusting the percent identity score. That is, only query residue positions outside the farthest N- and C-terminal residues of the subject sequence are considered for this manual correction. For example, a 90 amino acid residue subject sequence is aligned with a 100 residue query sequence to determine percent identity. The deletion occurs at the N-terminus of the subject sequence and therefore, the FASTDB alignment does not show a matching/alignment of the first 10 residues at the N-terminus. The 10 unpaired residues represent 10% of the sequence (number of residues at the N- and C-termini not matched/total number of residues in the query sequence) so 10% is subtracted from the percent identity score calculated by the FASTDB program. If the remaining 90 residues were perfectly matched the final percent identity would be 90%. In another example, a 90 residue subject sequence is compared with a 100 residue query sequence. This time the deletions are internal deletions so there are no residues at the N- or C-termini of the subject sequence which are not matched/aligned with the query. In this case the percent identity calculated by FASTDB is not manually corrected. Once again, only residue positions outside the N- and C-terminal ends of the subject sequence, as displayed in the FASTDB alignment, which are not matched/aligned with the query sequence are manually corrected for.

**[0112]** More generally, nucleic acid constructs can be prepared comprising an isolated polynucleotide encoding a polypeptide having enzyme activity operably linked to one or more (several) control sequences that direct the expression of the coding sequence in a microorganism, such as *E. coli*, under conditions compatible with the control sequences. The isolated polynucleotide may be manipulated to provide for expression of the polypeptide. Manipulation of the polynucleotide's sequence prior to its insertion into a vector may be desirable or necessary depending on the expression vector. The techniques for modifying polynucleotide sequences utilizing recombinant DNA methods are well established in the art.

**[0113]** The control sequence may be an appropriate promoter sequence, a nucleotide sequence that is recognized by a host cell for expression of a polynucleotide encoding a polypeptide of the present invention. The promoter sequence contains transcriptional control sequences that mediate the expression of the polypeptide. The promoter may be any nucleotide sequence that shows transcriptional activity in the host cell of choice including mutant, truncated, and hybrid promoters, and may be obtained from genes encoding extracellular or intracellular polypeptides either homologous or heterologous to the host cell. Examples of suitable promoters for directing transcription of the nucleic acid constructs, especially in an *E. coli* host cell, are the lac promoter (Gronenborn, 1976, Mol. Gen. Genet. 148: 243-250), tac promoter (DeBoer et al., 1983, Proceedings of the National Academy of Sciences USA 80: 21-25), trc promoter (Brosius et al, 1985, J. Biol. Chem. 260: 3539-3541), T7 RNA polymerase promoter (Studier and Moffatt, 1986, J. Mol. Biol. 189: 113-130), phage promoter p<sub>L</sub> (Elvin et al., 1990, Gene 87: 123-126), tetA promoter (Skerra, 1994, Gene 151: 131-135), araBAD promoter (Guzman et al., 1995, J. Bacteriol. 177: 4121-4130), and rhaP<sub>BAD</sub> promoter (Haldimann et al., 1998, J. Bacteriol. 180: 1277-1286). Other promoters are described in "Useful proteins from recombinant bacteria" in Scientific American, 1980, 242: 74-94; and in Sambrook and Russell, "Molecular Cloning: A Laboratory Manual," Third Edition 2001 (volumes 1-3), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.

**[0114]** The control sequence may also be a suitable transcription terminator sequence, a sequence recognized by a host cell to terminate transcription. The terminator sequence is operably linked to the 3' terminus of the nucleotide sequence encoding the polypeptide. Any terminator that is functional in an *E. coli* cell may be used in the present invention. It may also be desirable to add regulatory sequences that allow regulation of the expression of the polypeptide relative to the growth of the host cell. Examples of regulatory systems are those that cause the expression of the gene to be turned on or off in response to a chemical or physical stimulus, including the presence of a regulatory compound. Regulatory systems in prokaryotic systems include the lac, tac, and trp operator systems.

**[0115]** For various embodiments of the invention the genetic manipulations may be described to include various genetic manipulations, including those directed to change regulation of, and therefore ultimate activity of, an enzyme or enzymatic activity of an enzyme identified in any of the respective pathways. Such genetic modifications may be directed to transcriptional, translational, and post-translational modifications that result in a change of enzyme activity and/or selectivity under selected and/or identified culture

conditions and/or to provision of additional nucleic acid sequences such as to increase copy number and/or mutants of an enzyme related to 3-HP production. Specific methodologies and approaches to achieve such genetic modification are well known to one skilled in the art, and include, but are not limited to: increasing expression of an endogenous genetic element; decreasing functionality of a repressor gene; introducing a heterologous genetic element; increasing copy number of a nucleic acid sequence encoding a polypeptide catalyzing an enzymatic conversion step to produce 3-HP; mutating a genetic element to provide a mutated protein to increase specific enzymatic activity; over-expressing; under-expressing; over-expressing a chaperone; knocking out a protease; altering or modifying feedback inhibition; providing an enzyme variant comprising one or more of an impaired binding site for a repressor and/or competitive inhibitor; knocking out a repressor gene; evolution, selection and/or other approaches to improve mRNA stability as well as use of plasmids having an effective copy number and promoters to achieve an effective level of improvement. Random mutagenesis may be practiced to provide genetic modifications that may fall into any of these or other stated approaches. The genetic modifications further broadly fall into additions (including insertions), deletions (such as by a mutation) and substitutions of one or more nucleic acids in a nucleic acid of interest. In various embodiments a genetic modification results in improved enzymatic specific activity and/or turnover number of an enzyme. Without being limited, changes may be measured by one or more of the following:  $K_M$ ;  $k_{cat}$ ; and  $K_{avidity}$ .

**[0116]** In various embodiments, to function more efficiently, a microorganism may comprise one or more gene deletions. For example, in *E. coli*, the genes encoding the lactate dehydrogenase (*ldhA*), phosphate acetyltransferase (*pta*), pyruvate oxidase (*poxB*), and pyruvate-formate lyase (*pflB*) may be disrupted, including deleted. Such gene disruptions, including deletions, are not meant to be limiting, and may be implemented in various combinations in various embodiments. Gene deletions may be accomplished by mutational gene deletion approaches, and/or starting with a mutant strain having reduced or no expression of one or more of these enzymes, and/or other methods known to those skilled in the art. Gene deletions may be effectuated by any of a number of known specific methodologies, including but not limited to the RED/ET methods using kits and other reagents sold by Gene Bridges (Gene Bridges GmbH, Dresden, Germany, <<[www.genebridges.com](http://www.genebridges.com)>>).

**[0117]** More particularly as to the latter method, use of Red/ET recombination, is known to those of ordinary skill in the art and described in U.S. Pat. Nos. 6,355,412 and 6,509,156, issued to Stewart et al. and incorporated by reference herein for its teachings of this method. Material and kits for such method are available from Gene Bridges (Gene Bridges GmbH, Dresden, Germany, <<[www.genebridges.com](http://www.genebridges.com)>>), and the method may proceed by following the manufacturer's instructions. The method involves replacement of the target gene by a selectable marker via homologous recombination performed by the recombinase from X-phage. The host organism expressing  $\lambda$ -red recombinase is transformed with a linear DNA product coding for a selectable marker flanked by the terminal regions (generally ~50 bp, and alternatively up to about ~300 bp) homologous with the target gene. The marker could then be removed by another recombination step per-

formed by a plasmid vector carrying the FLP-recombinase, or another recombinase, such as Cre.

**[0118]** Targeted deletion of parts of microbial chromosomal DNA or the addition of foreign genetic material to microbial chromosomes may be practiced to alter a host cell's metabolism so as to reduce or eliminate production of undesired metabolic products. This may be used in combination with other genetic modifications such as described herein in this general example. In this detailed description, reference has been made to multiple embodiments and to the accompanying drawings in which is shown by way of illustration specific exemplary embodiments in which the invention may be practiced. These embodiments are described in sufficient detail to enable those skilled in the art to practice the invention, and it is to be understood that modifications to the various disclosed embodiments may be made by a skilled artisan.

**[0119]** Further, for 3-HP production, such genetic modifications may be chosen and/or selected for to achieve a higher flux rate through certain enzymatic conversion steps within the respective 3-HP production pathway and so may affect general cellular metabolism in fundamental and/or major ways.

**[0120]** It will be appreciated that amino acid "homology" includes conservative substitutions, i.e. those that substitute a given amino acid in a polypeptide by another amino acid of similar characteristics. Typically seen as conservative substitutions are the following replacements: replacements of an aliphatic amino acid such as Ala, Val, Leu and Ile with another aliphatic amino acid; replacement of a Ser with a Thr or vice versa; replacement of an acidic residue such as Asp or Glu with another acidic residue; replacement of a residue bearing an amide group, such as Asn or Gln, with another residue bearing an amide group; exchange of a basic residue such as Lys or Arg with another basic residue; and replacement of an aromatic residue such as Phe or Tyr with another aromatic residue.

**[0121]** For all nucleic acid and amino acid sequences provided herein, it is appreciated that conservatively modified variants of these sequences are included, and are within the scope of the invention in its various embodiments. Functionally equivalent nucleic acid and amino acid sequences (functional variants), which may include conservatively modified variants as well as more extensively varied sequences, which are well within the skill of the person of ordinary skill in the art, and microorganisms comprising these, also are within the scope of various embodiments of the invention, as are methods and systems comprising such sequences and/or microorganisms. In various embodiments, nucleic acid sequences encoding sufficiently homologous proteins or portions thereof are within the scope of the invention. More generally, nucleic acid sequences that encode a particular amino acid sequence employed in the invention may vary due to the degeneracy of the genetic code, and nonetheless fall within the scope of the invention. The following table provides a summary of similarities among amino acids, upon which conservative and less conservative substitutions may be based, and also various codon redundancies that reflect this degeneracy.

**[0122]** Degeneracy of the Amino Acid Code

Amino Acid	Relationships	DNA codons
Alanine	N, Ali	GCT, GCC, GCA, GCG
Proline	N	CCT, CCC, CCA, CCG
Valine	N, Ali	GTT, GTC, GTA, GTG
Leucine	N, Ali	CTT, CTC, CTA, CTG, TTA, TTG
Isoleucine	N, Ali	ATT, ATC, ATA
Methionine	N	ATG
Phenylalanine	N, Aro	TTT, TTC
Tryptophan	N	TGG
Glycine	PU	GGT, GGC, GGA, GGG
Serine	PU	TCT, TCC, TCA, TCG, AGT, AGC
Threonine	PU	ACT, ACC, ACA, ACG
Asparagine	PU, Ami	AAT, AAC
Glutamine	PU, Ami	CAA, CAG
Cysteine	PU	TGT, TGC
Aspartic acid	NEG, A	GAT, GAC
Glutamic acid	NEG, A	GAA, GAG
Arginine	POS, B	CGT, CGC, CGA, CGG, AGA, AGG
Lysine	POS, B	AAA, AAG
Histidine	POS	CAT, CAC
Tyrosine	Aro	TAT, TAC
Stop Codons		TAA, TAG, TGA

Legend: side groups and other related properties: A = acidic; B = basic; Ali = aliphatic; Ami = amine; Aro = aromatic; N = nonpolar; PU = polar uncharged; NEG = negatively charged; POS = positively charged.

**[0123]** Also, variants and portions of particular nucleic acid sequences, and respective encoded amino acid sequences recited herein may exhibit a desired functionality, e.g., enzymatic activity at a selected level, when such nucleic acid sequence variant and/or portion contains a 15 nucleotide sequence identical to any 15 nucleotide sequence set forth in the nucleic acid sequences recited herein including, without limitation, the sequence starting at nucleotide number 1 and ending at nucleotide number 15, the sequence starting at nucleotide number 2 and ending at nucleotide number 16, the sequence starting at nucleotide number 3 and ending at nucleotide number 17, and so forth. It will be appreciated that the invention also provides isolated nucleic acid that contains a nucleotide sequence that is greater than 15 nucleotides (e.g., 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more nucleotides) in length and identical to any portion of the sequence set forth in nucleic acid sequences recited herein. For example, the invention provides isolated nucleic acid that contains a 25 nucleotide sequence identical to any 25 nucleotide sequence set forth in any one or more (including any grouping of) nucleic acid sequences recited herein including, without limitation, the sequence starting at nucleotide number 1 and ending at nucleotide number 25, the sequence starting at nucleotide number 2 and ending at nucleotide number 26, the sequence starting at nucleotide number 3 and ending at nucleotide number 27, and so forth. Additional examples include, without limitation, isolated nucleic acids that contain a nucleotide sequence that is 50 or more nucleotides (e.g., 100, 150, 200, 250, 300, or more nucleotides) in length and identical to any portion of any of the sequences disclosed herein. Such isolated nucleic acids can include, without limitation, those isolated nucleic acids containing a nucleic acid sequence represented in any one section of discussion and/or examples, such as regarding 3-HP production pathways, nucleic acid sequences encoding enzymes of the fatty acid synthase system, or 3-HP tolerance. For example, the invention provides an isolated nucleic acid containing a nucleic acid sequence listed herein that contains a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any

combination thereof (e.g., single deletion together with multiple insertions). Such isolated nucleic acid molecules can share at least 60, 65, 70, 75, 80, 85, 90, 95, 97, 98, or 99 percent sequence identity with a nucleic acid sequence listed herein (i.e., in the sequence listing).

**[0124]** Additional examples include, without limitation, isolated nucleic acids that contain a nucleic acid sequence that encodes an amino acid sequence that is 50 or more amino acid residues (e.g., 100, 150, 200, 250, 300, or more amino acid residues) in length and identical to any portion of an amino acid sequence listed or otherwise disclosed herein.

**[0125]** In addition, the invention provides isolated nucleic acid that contains a nucleic acid sequence that encodes an amino acid sequence having a variation of an amino acid sequence listed or otherwise disclosed herein. For example, the invention provides isolated nucleic acid containing a nucleic acid sequence encoding an amino acid sequence listed or otherwise disclosed herein that contains a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such isolated nucleic acid molecules can contain a nucleic acid sequence encoding an amino acid sequence that shares at least 60, 65, 70, 75, 80, 85, 90, 95, 97, 98, or 99 percent sequence identity with an amino acid sequence listed or otherwise disclosed herein.

**[0126]** Examples of properties that provide the bases for conservative and other amino acid substitutions are exemplified in Table 1. Accordingly, one skilled in the art may make numerous substitutions to obtain an amino acid sequence variant that exhibits a desired functionality. BLASTP, CLUSTALP, and other alignment and comparison tools may be used to assess highly conserved regions, to which fewer substitutions may be made (unless directed to alter activity to a selected level, which may require multiple substitutions). More substitutions may be made in regions recognized or believed to not be involved with an active site or other binding or structural motif. In accordance with Table 1, for example, substitutions may be made of one polar uncharged (PU) amino acid for a polar uncharged amino acid of a listed sequence, optionally considering size/molecular weight (i.e., substituting a serine for a threonine). Guidance concerning which amino acid changes are likely to be phenotypically silent can be found, inter alia, in Bowie, J. U., et Al., "Deciphering the Message in Protein Sequences: Tolerance to Amino Acid Substitutions," *Science* 247:1306-1310 (1990). This reference is incorporated by reference for such teachings, which are, however, also generally known to those skilled in the art. Recognized conservative amino acid substitutions comprise (substitutable amino acids following each colon of a set): ala:ser; arg:lys; asn:gln or his; asp:glu; cys:ser; gln:asn; glu:asp; gly:pro; his:asn or gln; ile:leu or val; leu:ile or val; lys:arg or gln or glu; met:leu or ile; phe:met or leu or tyr; ser:thr; thr:ser; trp:tyr; tyr:trp or phe; val:ile or leu.

**[0127]** It is noted that codon preferences and codon usage tables for a particular species can be used to engineer isolated nucleic acid molecules that take advantage of the codon usage preferences of that particular species. For example, the isolated nucleic acid provided herein can be designed to have codons that are preferentially used by a particular organism of interest. Numerous software and sequencing services are available for such codon-optimizing of sequences.

**[0128]** The invention provides polypeptides that contain the entire amino acid sequence of an amino acid sequence

listed or otherwise disclosed herein. In addition, the invention provides polypeptides that contain a portion of an amino acid sequence listed or otherwise disclosed herein. For example, the invention provides polypeptides that contain a 15 amino acid sequence identical to any 15 amino acid sequence of an amino acid sequence listed or otherwise disclosed herein including, without limitation, the sequence starting at amino acid residue number 1 and ending at amino acid residue number 15, the sequence starting at amino acid residue number 2 and ending at amino acid residue number 16, the sequence starting at amino acid residue number 3 and ending at amino acid residue number 17, and so forth. It will be appreciated that the invention also provides polypeptides that contain an amino acid sequence that is greater than 15 amino acid residues (e. g., 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30 or more amino acid residues) in length and identical to any portion of an amino acid sequence listed or otherwise disclosed herein. For example, the invention provides polypeptides that contain a 25 amino acid sequence identical to any 25 amino acid sequence of an amino acid sequence listed or otherwise disclosed herein including, without limitation, the sequence starting at amino acid residue number 1 and ending at amino acid residue number 25, the sequence starting at amino acid residue number 2 and ending at amino acid residue number 26, the sequence starting at amino acid residue number 3 and ending at amino acid residue number 27, and so forth. Additional examples include, without limitation, polypeptides that contain an amino acid sequence that is 50 or more amino acid residues (e.g., 100, 150, 200, 250, 300 or more amino acid residues) in length and identical to any portion of an amino acid sequence listed or otherwise disclosed herein. Further, it is appreciated that, per above, a 15 nucleotide sequence will provide a 5 amino acid sequence, so that the latter, and higher-length amino acid sequences, may be defined by the above-described nucleotide sequence lengths having identity with a sequence provided herein.

**[0129]** In addition, the invention provides polypeptides that an amino acid sequence having a variation of the amino acid sequence set forth in an amino acid sequence listed or otherwise disclosed herein. For example, the invention provides polypeptides containing an amino acid sequence listed or otherwise disclosed herein that contains a single insertion, a single deletion, a single substitution, multiple insertions, multiple deletions, multiple substitutions, or any combination thereof (e.g., single deletion together with multiple insertions). Such polypeptides can contain an amino acid sequence that shares at least 60, 65, 70, 75, 80, 85, 90, 95, 97, 98 or 99 percent sequence identity with an amino acid sequence listed or otherwise disclosed herein. A particular variant amino acid sequence may comprise any number of variations as well as any combination of types of variations.

**[0130]** The invention includes, in various embodiments, an amino acid sequence having a variation of any of the polynucleotide and polypeptide sequences disclosed herein. As one example, variations are exemplified for the carbonic anhydrase (*E. coli* cynT) amino acid sequence set forth in SEQ ID NO:57. FIG. 1 provides a CLUSTAL multiple sequence alignment of the *E. coli* carbonic anhydrase aligned with carbonic anhydrases of eleven other species that had relatively high homology, based on low E values, in a BLASTP comparison. SEQ ID NO:57 is the fifth sequence shown. Multiple conservative and less conservative substitutions are shown (i.e., by the “:” and “.” designations, respec-

tively), which can lead to additional modifications by one skilled in the art. Thus, examples of variations of the sequence set forth in SEQ ID NO:57 include, without limitation, any variation of the sequences as set forth in FIG. 1. Such variations are provided in FIG. 1 in that a comparison of the amino acid residue (or lack thereof) at a particular position of the sequence set forth in SEQ ID NO:57 with the amino acid residue (or lack thereof) at the same aligned position of any of the other eleven amino acid sequences of FIG. 1 provides a list of specific changes for the sequence set forth in SEQ ID NO:57. For example, the “E” glutamic acid at position 14 of SEQ ID NO:57 can be substituted with a “D” aspartic acid or “N” asparagine as indicated in FIG. 1. It will be appreciated that the sequence set forth in SEQ ID NO:57 can contain any number of variations as well as any combination of types of variations. It is noted that the amino acid sequences provided in FIG. 1 can be polypeptides having carbonic anhydrase activity.

**[0131]** As indicated herein, polypeptides having a variant amino acid sequence can retain enzymatic activity. Such polypeptides can be produced by manipulating the nucleotide sequence encoding a polypeptide using standard procedures such as site-directed mutagenesis or various PCR techniques. As noted herein, one type of modification includes the substitution of one or more amino acid residues for amino acid residues having a similar chemical and/or biochemical property. For example, a polypeptide can have an amino acid sequence set forth in an amino acid sequence listed or otherwise disclosed herein comprising one or more conservative substitutions.

**[0132]** More substantial changes can be obtained by selecting substitutions that are less conservative, and/or in areas of the sequence that may be more critical, for example selecting residues that differ more significantly in their effect on maintaining: (a) the structure of the polypeptide backbone in the area of the substitution, for example, as a sheet or helical conformation; (b) the charge or hydrophobicity of the polypeptide at the target site; or (c) the bulk of the side chain. The substitutions that in general are expected to produce the greatest changes in polypeptide function are those in which: (a) a hydrophilic residue, e.g., serine or threonine, is substituted for (or by) a hydrophobic residue, e.g., leucine, isoleucine, phenylalanine, valine or alanine; (b) a cysteine or proline is substituted for (or by) any other residue; (c) a residue having an electropositive side chain, e.g., lysine, arginine, or histidine, is substituted for (or by) an electronegative residue, e.g., glutamic acid or aspartic acid; or (d) a residue having a bulky side chain, e.g., phenylalanine, is substituted for (or by) one not having a side chain, e.g., glycine. The effects of these amino acid substitutions (or other deletions or additions) can be assessed for polypeptides having enzymatic activity by analyzing the ability of the polypeptide to catalyze the conversion of the same substrate as the related native polypeptide to the same product as the related native polypeptide. Accordingly, polypeptides having 5, 10, 20, 30, 40, 50 or less conservative substitutions are provided by the invention.

**[0133]** Polypeptides and nucleic acids encoding polypeptides can be produced by standard DNA mutagenesis techniques, for example, M13 primer mutagenesis. Details of these techniques are provided in Sambrook and Russell, 2001. Nucleic acid molecules can contain changes of a coding region to fit the codon usage bias of the particular organism into which the molecule is to be introduced.

**[0134]** Alternatively, the coding region can be altered by taking advantage of the degeneracy of the genetic code to alter the coding sequence in such a way that, while the nucleic acid sequence is substantially altered, it nevertheless encodes a polypeptide having an amino acid sequence identical or substantially similar to the native amino acid sequence. For example, alanine is encoded in the open reading frame by the nucleotide codon triplet GCT. Because of the degeneracy of the genetic code, three other nucleotide codon triplets—GCA, GCC, and GCG—also code for alanine. Thus, the nucleic acid sequence of the open reading frame can be changed at this position to any of these three codons without affecting the amino acid sequence of the encoded polypeptide or the characteristics of the polypeptide. Based upon the degeneracy of the genetic code, nucleic acid variants can be derived from a nucleic acid sequence disclosed herein using standard DNA mutagenesis techniques as described herein, or by synthesis of nucleic acid sequences. Thus, for various embodiments the invention encompasses nucleic acid molecules that encode the same polypeptide but vary in nucleic acid sequence by virtue of the degeneracy of the genetic code.

**[0135]** The invention also provides an isolated nucleic acid that is at least about 12 bases in length (e.g., at least about 13, 14, 15, 16, 17, 18, 19, 20, 25, 30, 40, 50, 60, 100, 250, 500, 750, 1000, 1500, 2000, 3000, 4000, or 5000 bases in length) and hybridizes, under hybridization conditions, to the sense or antisense strand of a nucleic acid having a sequence listed or otherwise disclosed herein. The hybridization conditions can be moderately or highly stringent hybridization conditions. Also, in some embodiments the microorganism comprises an endogenous 3-HP production pathway (which may, in some such embodiments, be enhanced), whereas in other embodiments the microorganism does not comprise a 3-HP production pathway, but is provided with one or more nucleic acid sequences encoding polypeptides having enzymatic activity or activities to complete a pathway, described herein, resulting in production of 3-HP. In some embodiments, the particular sequences disclosed herein, or conservatively modified variants thereof, are provided to a selected microorganism, such as selected from one or more of the species and groups of species or other taxonomic groups listed herein.

**[0136]** VI. Specific Mutations, Genetic Modifications, and Combinations Thereof

**[0137]** The present invention, in its various embodiments, is directed to making one, or a combination of, genetic modifications in a microorganism to increase production therein of the commercially important compound 3-hydroxypropionic acid (“3-HP,” CAS No. 503-66-2). More generally the present invention relates to methods, systems and compositions, including genetically modified microorganisms, e.g., recombinant microorganisms, comprising one or more genetic modifications directed to increased production of 3-HP based on introduction of a polynucleotide encoding identified functional variant polypeptides exhibiting increased oxaloacetate alpha-decarboxylase/oxaloacetate alpha-decarboxylase activity, and methods of making various products from 3-HP made by cultures of such microorganisms.

**[0138]** Choice of candidate oxaloacetate alpha-decarboxylase enzymes: The desired oxaloacetate alpha-decarboxylase activity catalyzes the conversion of oxaloacetate to malonate semialdehyde (see FIG. 2), and may be referred to more explicitly as an oxaloacetate alpha-decarboxylase. Candidate enzymes were identified and chosen according to their reported activities as decarboxylases of alpha-ketoacids and

by their reported substrate preference for shorter chain molecules. The genes for pyruvate decarboxylase from *Zymomonas mobilis* (pdc), 2-oxoglutarate decarboxylase (SEQ ID NO:035), (oad) from *Leuconostoc mesenteroides* (SEQ ID NO:036) and alpha-ketoglutarate decarboxylase (kgd) from *Mycobacterium tuberculosis* (SEQ ID NO:037) were chosen for further evaluation.

**[0139]** Through such evaluation, three functional variants were identified from the 2-oxoglutarate decarboxylase (oad) of *Leuconostoc mesenteroides* (SEQ ID NO:036) These, respectively, have the following mutations of that sequence: N45T, R249L, D302G, V418A and L476Q (SEQ ID NO:054), T479N (SEQ ID NO:055), and R394C, D434G and T511A (SEQ ID NO:056). These respectively showed a 1.6, 2.2 and 2.8 fold improvement in enzymatic conversion of oxaloacetate to malonate semialdehyde compared with a control native enzyme lacking mutations in an in vitro system (see Examples).

**[0140]** In various embodiments additional genetic modifications may be made to increase metabolic flux at one or more specified enzymatic conversion steps, and/or to reduce or eliminate an enzymatic conversion along certain metabolic pathways. In some embodiments, a genetic modification is provided to a selected microorganism cell to increase an oxaloacetate alpha-decarboxylase/oxaloacetate alpha-decarboxylase enzyme activity. Such genetic modification may supplement an existing enzymatic activity, or may provide such activity in a selected microorganism that previously lacked such enzymatic activity (whether inherently or due or other genetic modifications).

**[0141]** As disclosed in U.S. Provisional Patent Application No. 61/246,140, filed Sep. 27, 2009 and to which priority is claimed, SEQ ID NO:001 provides one non-limiting example of an amino acid sequence of an oxaloacetate alpha-decarboxylase/oxaloacetate alpha-decarboxylase enzyme, obtained from *Leuconostoc mesenteroides*. This sequence is observed to have lacked a small portion of the N-terminal, and SEQ ID NO:036 is provided herein as the entire sequence.

**[0142]** FIG. 2 graphically depicts a reaction catalyzed by an oxaloacetate alpha-decarboxylase. A polynucleotide encoding such polypeptide sequence, or variants, including conservative variants thereof, exhibiting such enzymatic activity (hereinafter referred to as “oad-2”), may be provided into a selected microorganism to increase 3-HP production. This is described in more detail below. In various embodiments, such provision, by genetic modification, is practiced in combination with other genetic modifications so as to further increase 3-HP production.

**[0143]** Such sequence may be encoded by a polynucleotide that may be provided to a selected microorganism by a genetic modification, so as to provide or increase such enzymatic activity in the selected microorganism. For example, in some embodiments such genetic modification may comprise providing a plasmid, or other vector (e.g., cosmids, bacteria artificial chromosome (“BAC”), viruses (e.g., bacteriophage, animal viruses, plant viruses), and artificial chromosomes (e.g., yeast artificial chromosomes (YAC)), that comprises a polynucleotide that encodes a sequence as described herein and further comprises appropriate promoter(s), binding site(s), and stop codon, such that the amino acid sequence is expressed to a desirable level in the selected microorganism.

**[0144]** FIG. 3A depicts the natural metabolic pathways utilized by *E. coli* during bio-production which results in the

natural products lactate, formate and acetate. By disruptions of appropriate genes, the production of these natural products are reduced or eliminated.

**[0145]** FIG. 3B depicts a proposed metabolic pathway to produce 3-HP as a bio-production product. Arrows represent enzymatic activities. Enzyme activities for FIGS. 3A and 3B are as follows: [i] glucokinase, [ii] phosphoglucose isomerase, [iii] 6-phosphofructose kinase, [iv] fructose bisphosphate aldolase, [v] triose-phosphate isomerase, [vi] glyceraldehyde 3-phosphate dehydrogenase, [vii] phosphoglycerate kinase, [viii] phosphoglycerate mutase, [ix] enolase, [x] pyruvate dehydrogenase, -, [xi] lactate dehydrogenase, [xii] pyruvate oxidase, [xiii] pyruvate-formate lyase, [xiv] phosphate acetyltransferase, [xv] acetate kinase, [xvi] phosphoenolpyruvate carboxykinase, [xvii] the proposed oxaloacetate alpha-decarboxylase (oad-2), [xviii] 3-hydroxypropionate dehydrogenase, [xix] pyruvate carboxylase.

**[0146]** FIG. 4 depicts a generalized embodiment comprising various metabolic modifications, some of which are optional for various embodiments of a genetically modified microorganism of the present invention. For various embodiments, an underlined gene name indicates that genetic modification is made to increase expression and/or enzymatic activity of the respective gene product (i.e., enzyme), a “Δ” before a gene name signifies that this gene is disrupted to decrease or eliminate expression and/or enzymatic activity of the respective gene product, and an asterisk indicates that this gene may be mutated to obtain a mutated form of the encoded gene product that exhibits increased enzymatic activity and/or specificity. Not all such genetic modifications need be made in a particular embodiment, and this full set of genetic modifications would be made for culture under anaerobic conditions.

**[0147]** As noted, the oxaloacetate alpha-decarboxylase is identified as “oad-2” or a grammatically equivalent identification. As shown in FIG. 2, oad-2 converts an oxaloacetate molecule to form malonate semialdehyde (“MSA”). The MSA is converted to 3-HP by a suitable enzyme (native, supplemented, or added), such as mmsB from *Pseudomonas aeruginosa* (SEQ ID NO:002) noting that the predominant reported reaction for this enzyme is conversion of 3-hydroxyisobutyrate to methylmalonate semialdehyde, and that in some embodiments mutant forms of mmsB may be produced, evaluated and identified that have greater, or more specific, activity for the conversion of MSA to 3-HP, such as by use of mutation and selection approaches described herein and/or known to those skilled in the art. Other enzymes for this latter conversion may be selected from enzymes capable of reductive conversions from MSA to 3-HP, such as a native or mutated form of ydfG or its functional equivalent in other species. This reaction is depicted in FIG.

**[0148]** The types of additional genetic modifications are generally divided into two groups—those that are made to increase enzymatic activity and/or specificity, generally so as to increase flux through a particular pathway and/or enzymatic conversion step, and those that are made to decrease or eliminate enzymatic activity at particular enzymatic conversion steps, thereby decreasing or eliminating conversion to particular metabolic intermediates or products. Various embodiments may include combinations of such genetic modifications from these two groups, in combination with the above-described genetic modification of a decarboxylase able to convert oxaloacetate to MSA.

**[0149]** Accordingly, in some embodiments, one or more of the genetic modifications in Table 1 may be provided to the selected microorganism that also is provided with the oxaloacetate alpha-decarboxylase enzymatic activity. Genetic modifications to the microorganism of these enzymes are made to increase enzymatic activity and/or specificity. One or more of these enzymatic functions may be native, and/or genetic modifications may be provided to supplement such native activity, or to provide such activity to a microorganism not previously demonstrating such activity. As described in a section below, there are various approaches to obtaining homologies that may be determined to exhibit a desired functional equivalence.

**[0150]** In some embodiments, one of the enzymatic activities of Table 1, the conversion of phosphoenolpyruvate (“PEP”) to OAA, is provided via providing a mutated enzyme exhibiting such activity to a great level than a non-mutated enzyme. For example, the Ppc enzyme of *E. coli*, phosphoenolpyruvate carboxylase, may be mutated, such as by constructing a mutant library of ppc by use of an error-inducing PCR site-directed mutagenesis method. For example, use of the XL1-Red mutator strain, which is deficient in several repair mechanisms necessary for accurate DNA replication and generates mutations in plasmids at a rate 5,000 times that of the wild-type mutation rate, may be employed using appropriate materials following a manufacturer’s instructions (see Stratagene QuikChange Mutagenesis Kit, Stratagene, La Jolla, Calif. USA). This technique or other techniques known to those skilled in the art, may be employed and then a population of such mutants, e.g., in a library, is evaluated, such as by a screening or selection method, to identify clones having a suitable or favorable mutation.

**[0151]** In other cases, such as for galP and pckA, genetic modifications can be made to increase overall expression of these protein functions in a microorganism cell. Various methods are known in the art for such types of genetic modifications, and are described in a section below.

**[0152]** In some embodiments a genetic modification provides increased enzymatic activity of pyruvate carboxylase (e.g., pyc from *Corynebacterium glutamicum* or *Rhizobium etli*).

**[0153]** Table 2 lists a number of protein functions, also providing enzyme classes and specific examples (with corresponding SEQ ID NOs.) in the group for which genetic modifications are made to reduce or eliminate the respective enzymatic activity. Once these protein functions are reduced or eliminated, more carbon and energy may flow to production of 3-HP (and also to biomass formation, to an extent, in some embodiments) rather than to the formation of metabolic products such as lactate, acetate, and formate.

**[0154]** Also, as gleaned from Tables 1 and 2, in various embodiments genetic modifications are made to increase enzymatic expression of galP and to decrease or eliminate enzymatic expression of all or part of the ptsHlcr operon, or their equivalents in a selected species.

**[0155]** Deletions of the polynucleotides encoding the polypeptides exhibiting enzymatic activities or other protein functions in Table 2 may be made in a selected *E. coli* strain using a RED/ET homologous recombination method, such as with kits supplied by Gene Bridges (Gene Bridges GmbH, Heidelberg, Germany, www.genebridges.com) according to manufacturer’s instructions. The successful deletion of these genes, as confirmed by standard methodologies, such as PCR, or DNA sequencing, results in a genetically modified micro-



organism having reduced or eliminated respective enzymatic activities or other protein functions. Methods for gene disruption in other species are known to those skilled in the art.

**[0156]** In view of Tables 1 and 2, it will be recognized that there are many possible combinations of increases in one or more protein functions, optionally with reductions in one or more protein functions. Protein functions can be independently varied, and any combination (i.e., a full factorial) of protein functions in Table 1 and Table 2 herein can be adjusted in various embodiments. In various embodiments of the present invention, these various combinations are provided in combination with one or more genetic modifications to provide, or to increase existing levels of, oxaloacetate alpha-decarboxylase enzymatic activity.

**[0157]** In various independent groupings of such embodiments, one or more protein functions selected from Table 1 may be added or increased, however excluding any substantial addition or change to any one of galP and its homologues, Ppc and its homologues, pckA and its homologues, or excluding two, three, four, or more, of such protein functions and their homologues from such smaller list or sub-list. In other independent groupings of embodiments, the various sub-lists developed from the list of Table 1 exclude one or more of the above-indicated protein functions but not their homologues.

**[0158]** In various independent groupings of such embodiments, one or more protein functions selected from Table 2 may be deleted or disrupted, however excluding any substantial reduction or change to any one of aceE and its homologues, aceF and its homologues, lpd and its homologues, ldhA and its homologues, pflB and its homologues, poxB and its homologues, Pta and its homologues, ptsH and its homologues, ptsI and its homologues, Crr and its homologues, pykA and its homologues, pykF and its homologues, or excluding two, three, four, five, six, seven, eight, nine, ten, eleven, twelve, or more, of such protein functions and their homologues from such smaller list or sub-list. In other independent groupings of embodiments, the various sub-lists developed from the list of Table 2 exclude one or more of the above-indicated protein functions but not their homologues.

**[0159]** In addition to the above-described genetic modifications, in various embodiments genetic modifications also are provided to increase the pool and availability of the cofactor NADPH, and/or, consequently, the NADPH/NADP<sup>+</sup> ratio. For example, in various embodiments for *E. coli*, this may be done by increasing activity, such as by genetic modification, of one or more of the following genes: *pgi* (in a mutated form), *pntAB*, overexpressed, *gapA:gapN* substitution/replacement, and disrupting or modifying a soluble transhydrogenase such as *sthA*, and/or genetic modifications of one or more of *zwf*, *gnd*, and *edd*.

**[0160]** Specific nucleic acid and amino acid sequences corresponding to the enzyme names and activities provided herein (e.g., for 3-HP production, and in Tables 1 and 2), including the claims, are readily found at widely used databases including [www.metacyc.org](http://www.metacyc.org), [www.brenda-enzymes.org](http://www.brenda-enzymes.org), and [www.ncbi.gov](http://www.ncbi.gov). Also, the particular enzymatic activities by enzymes, and other protein functions (and the nucleic acid sequences encoding them) that are disclosed herein are not meant to be limiting particularly in view of the various known approaches, standard in the art, to achieve desired metabolic conversions and to identify functionally analogous enzymes in different species. Different enzymes for different species may also be found on these web sites. The enzyme functions disclosed herein may be related to the stated E.C.

numbers provided herein, including Table 1 and 2, which are incorporated into this section. However, it is noted that some protein functions disclosed herein, including in the Tables 1 and 2, are not enzyme functions, and those functions are only stated in the "Protein Function" columns, and in various embodiments may be further limited by the function of the stated *E. coli* gene (which may be applied, in various embodiments, to functionally equivalent homologues in other species). Also, all information of Tables 1 and 2 are incorporated into the claims referring to these tables. Also, based on the teachings provided herein, it is appreciated that the members of a respective table are functionally related as to their overall metabolic effects on production of 3-HP through the 3-HP production pathway that includes oxaloacetate alpha-decarboxylase.

**[0161]** Thus, in various embodiments a genetically modified microorganism comprises a first set of genetic modification(s) to introduce and/or increase enzymatic activity of an oxaloacetate alpha-decarboxylase (such as SEQ ID NO:001 or SEQ ID NO:036), and optionally one or more enzymes capable of converting MSA to 3-HP (such as *mmsB* and a native or mutated *ydfG*). In some embodiments, a genetically modified microorganism demonstrates increased production of 3-HP compared to a control microorganism lacking such genetic modification(s). In some embodiments, a genetically modified microorganism comprises one or more genetic modifications to increase enzymatic activities or other protein functions identified in Table 1, and/or comprising one or more genetic modifications to reduce or eliminate enzymatic activities or other protein functions identified in Table 2, and optionally also comprising genetic modifications disclosed herein to increase microorganism NADPH pools and/or availability. Using approaches described herein, genes encoding enzymes having the enzyme activities described herein may be identified in other species, and evaluated as may be appropriate to the circumstance, toward obtaining a genetically modified microorganism of that species that comprises the genetic modifications taught herein to obtain a microorganism that exhibits higher production of 3-HP compared to a relevant control microorganism lacking the genetic modifications.

**[0162]** In some particular embodiments, a combination of genetic modifications are made to an *E. coli* strain, the combination comprising providing a polynucleotide encoding *oad-2* and the combination also comprising other genetic modifications shown in FIGS. 3B and 4, as well as described for Tables 1 and 2. In some of these embodiments, at least one of additions of *mmsB* or a native or mutated *ydfG* are provided to increase enzymatic conversion of MSA to 3-HP.

**[0163]** Other aspects of the scope of the invention are described in the following paragraphs.

**[0164]** In various embodiments, to function more efficiently, a microorganism may comprise one or more gene deletions. These are summarized in Table 2, described above. For example, in *E. coli*, the genes encoding lactate dehydrogenase (*ldhA*), phosphate acetyltransferase (*pta*), pyruvate oxidase (*poxB*) and pyruvate-formate lyase (*pflB*), may be disrupted, including deleted. Such gene disruptions, including deletions, are not meant to be limiting, and may be implemented in various combinations in various embodiments. Gene deletions may be accomplished by mutational gene deletion approaches, and/or starting with a mutant strain having reduced or no expression of one or more of these enzymes, and/or other methods known to those skilled in the art. Gene

deletions may be effectuated by any of a number of known specific methodologies, including but not limited to the RED/ET methods using kits and other reagents sold by Gene Bridges (Gene Bridges GmbH, Heidelberg, Germany, [www.genebridges.com](http://www.genebridges.com)). Further, for 3-HP production, such genetic modifications may be chosen and/or selected for to achieve a higher flux rate through certain basic pathways within the respective 3-HP production pathway and so may affect general cellular metabolism in fundamental and/or major ways.

**[0165]** In some embodiments, the genetically modified microorganism that so possesses oxaloacetate alpha-decarboxylase genetic modification(s) additionally comprises at least one genetic modification to increase, in the genetically modified microorganism, a protein function selected from the protein functions of Table 1 (Glucose transporter function (such as by galP), phosphoenolpyruvate carboxylase, and phosphoenolpyruvate carboxykinase). In certain embodiments, the genetically modified microorganism comprises at least one genetic modification to increase two or three protein functions selected from the protein functions of Table 1.

**[0166]** In some embodiments, such genetically modified microorganism additionally comprises at least one genetic modification to decrease protein functions selected from the protein functions of Table 2 (pyruvate dehydrogenase E1p, dihydrolipoamide acetyltransferase, pyruvate dehydrogenase E3, lactate dehydrogenase, pyruvate formate lyase, pyruvate oxidase, phosphate acetyltransferase, histidyl phosphorylatable protein (of PTS), phosphoryl transfer protein (of PTS), polypeptide chain (of PTS), pyruvate kinase I, and pyruvate kinase II).

**[0167]** In various embodiments, such genetically modified microorganism comprises at least one genetic modification to decrease enzymatic activity of two, three, four, five, six, seven, eight, nine, ten, eleven, or twelve protein functions selected from the protein functions of Table 2.

**[0168]** It will be appreciated that, in various embodiments, there can be many possible combinations of increases in one or more protein functions of Table 1, with reductions in one or more protein functions of Table 2 in the genetically modified microorganism comprising at least one genetic modification to provide or increase oxaloacetate alpha-decarboxylase protein function (i.e., enzymatic activity). Protein functions can be independently varied, and any combination (i.e., a full factorial) of genetic modifications of protein functions in Table 1 and Table 2 herein can be adjusted by the methods taught and provided into the genetically modified microorganism.

**[0169]** In some embodiments, at least one genetic modification to decrease enzymatic activity is a gene disruption. In some embodiments, at least one genetic modification to decrease enzymatic activity is a gene deletion.

**[0170]** In various embodiments, to obtain 3-hydroxypropionic acid (3-HP) as a desired product, the genetically modified microorganism comprises a protein function effective for converting malonate semialdehyde to 3-HP. The protein function effective for converting malonate semialdehyde to 3-HP can be native to the microorganism, but that is by no means necessary.

**[0171]** In some embodiments, the protein function effective for converting malonate semialdehyde to 3-HP is a native or mutated form of mmsB from *Pseudomonas aeruginosa*, or a functional equivalent thereof. Alternatively, or additionally,

this protein function can be a native or mutated form of ydfG, or a functional equivalent thereof.

**[0172]** Certain embodiments of the invention additionally comprise a genetic modification to increase the availability of the cofactor NADPH, which can increase the NADPH/NADP<sup>+</sup> ratio as may be desired. Non-limiting examples for such genetic modification are *pgi* (E.C. 5.3.1.9, in a mutated form), *pntAB* (E.C. 1.6.1.2), overexpressed, *gapA* (E.C. 1.2.1.12):*gapN* (E.C. 1.2.1.9, from *Streptococcus mutans*) substitution/replacement, and disrupting or modifying a soluble transhydrogenase such as *sthA* (E.C. 1.6.1.2), and/or genetic modifications of one or more of *zwf* (E.C. 1.1.1.49), *gnd* (E.C. 1.1.1.44), and *edd* (E.C. 4.2.1.12). Sequences of these genes are available at [www.metacyc.org](http://www.metacyc.org).

**[0173]** In some embodiments, the genetic modification increases microbial synthesis of 3-HP above a rate or titer of a control microorganism lacking said at least one genetic modification to produce 3-HP. In some embodiments, the genetic modification is effective to increase enzymatic conversions to 3-HP by at least about 5 percent, at least about 10 percent, at least about 20 percent, at least about 30 percent, or at least about 50 percent above the enzymatic conversion of a control microorganism lacking the genetic modification.

**[0174]** Variations of the present invention also provide a culture system comprising a population of a genetically modified microorganism as described herein, and a media comprising nutrients for the population.

**[0175]** Variations of this invention additionally include methods. In some variations, the invention provides a method of making a genetically modified microorganism comprising providing to a selected microorganism at least one genetic modification to introduce oxaloacetate alpha-decarboxylase enzymatic activity.

**[0176]** In some embodiments, the method additionally comprises introducing at least one genetic modification to increase enzymatic activity of a protein function selected from the protein functions of Table 1. In various embodiments, the method comprises introducing at least one genetic modification to increase two or three protein functions selected from the protein functions of Table 1 (Glucose transporter function (such as by galP), phosphoenolpyruvate carboxylase, and phosphoenolpyruvate carboxykinase).

**[0177]** In some embodiments, the method additionally comprises introducing at least one genetic modification to decrease a protein function selected from the protein functions of Table 2. In various embodiments, the method comprises introducing at least one genetic modification to decrease two, three, four, five, six, seven, eight, nine, ten, eleven, or twelve protein functions selected from the protein functions of Table 2 (pyruvate dehydrogenase E1p, dihydrolipoamide acetyltransferase, pyruvate dehydrogenase E3, lactate dehydrogenase, pyruvate formate lyase, pyruvate oxidase, phosphate acetyltransferase, histidyl phosphorylatable protein (of PTS), phosphoryl transfer protein (of PTS), polypeptide chain (of PTS), pyruvate kinase I, and pyruvate kinase II).

**[0178]** Again, there are many possible combinations of increases in one or more protein functions, optionally with reductions in one or more protein functions, that may be provided in combination with at least one genetic modification to provide or increase oxaloacetate alpha-decarboxylase enzymatic function, so as to increase 3-HP production in a genetically modified microorganism. Protein functions can be independently varied, and any combination (i.e., a full

factorial) of protein functions in Table 1 and Table 2 herein can be adjusted in various methods of the invention. Enzyme-activity and other protein function reductions can be accomplished by gene disruptions, such as gene deletions, or other modifications.

**[0179]** Additionally, genetic modifications and/or media supplements directed to improving tolerance to 3-HP may be provided, such as are taught in PCT Patent Publication No. US/2010/052748, published Jan. 28, 2010, and incorporated by reference for its teachings of tolerance-related genetic modifications and media supplements.

**[0180]** In some embodiments, the invention contemplates a culture system comprising: a) a population of a genetically modified microorganism as described herein; and b) a media comprising nutrients for the population. In some such embodiments the media additionally comprises at least 1 gram/liter of 3-HP.

**[0181]** As described in U.S. Provisional Patent Application No. 61/246,140, filed Sep. 27, 2009, incorporated by reference and to which priority is claimed, various combinations of genetic modifications may be implemented in various embodiments of the invention. These are described in the following paragraphs and also in Tables 1 and 2.

**[0182]** In some embodiments, the genetically modified microorganism that so possesses oxaloacetate alpha-decarboxylase genetic modification(s) additionally comprises at least one genetic modification to increase, in the genetically modified microorganism, a protein function selected from the protein functions of Table 1 (Glucose transporter function (such as by galP in *E. coli*), phosphoenolpyruvate carboxykinase (such as by pckA in *E. coli*), and phosphoenolpyruvate carboxylase (such as by ppc in *E. coli*). In certain embodiments, the genetically modified microorganism comprises at least one genetic modification to increase two, three, or four protein functions selected from the protein functions of Table 2.

**[0183]** In some embodiments, such genetically modified microorganism additionally comprises at least one genetic modification to decrease protein functions selected from the protein functions of Table 2, pyruvate dehydrogenase E1p, lipoate acetyltransferase/dihydrolipoamide acetyltransferase, pyruvate dehydrogenase E3 (lipoamide dehydrogenase), lactate dehydrogenase, pyruvate formate lyase, pyruvate oxidase, phosphate acetyltransferase, histidyl phosphorylatable protein (of PTS), phosphoryl transfer protein (of PTS), the polypeptide chain (of PTS), pyruvate kinase I, and pyruvate kinase II.

**[0184]** In various embodiments, such genetically modified microorganism comprises at least one genetic modification to decrease enzymatic activity of two, three, four, five, six, or seven protein functions selected from the protein functions of Table 2.

**[0185]** It will be appreciated that, in various embodiments, there can be many possible combinations of increases in one or more protein functions of Table 1, with reductions in one or more protein functions of Table 2 in the genetically modified microorganism comprising at least one genetic modification to provide or increase oxaloacetate alpha-decarboxylase protein function (i.e., enzymatic activity). Protein functions can be independently varied, and any combination (i.e., a full factorial) of genetic modifications of protein functions in Tables 1 and 2 herein can be adjusted by the methods taught and provided into said genetically modified microorganism.

**[0186]** In some embodiments, at least one genetic modification to decrease enzymatic activity is a gene disruption. In some embodiments, at least one genetic modification to decrease enzymatic activity is a gene deletion.

**[0187]** In various embodiments, to obtain 3-hydroxypropionic acid (3-HP) as a desired product, the genetically modified microorganism comprises a protein function effective for converting oxaloacetate to malonate semialdehyde, i.e., an oxaloacetate alpha-decarboxylase, and also a protein function effective for converting malonate semialdehyde to 3-HP. The latter can be native to the microorganism, but that is by no means necessary.

**[0188]** In some embodiments, the protein function effective for converting malonate semialdehyde to 3-HP is a native or mutated form of mmsB from *Pseudomonas aeruginosa*, or a functional equivalent thereof. Alternatively, or additionally, this protein function can be a native or mutated form of ydfG, or a functional equivalent thereof.

**[0189]** Certain embodiments of the invention additionally comprise a genetic modification to increase the availability of the cofactor NADPH, which can increase the NADPH/NADP+ ratio as may be desired. Non-limiting examples for such genetic modification are pgi (E.C. 5.3.1.9, in a mutated form), pntAB (E.C. 1.6.1.2), overexpressed, gapA (E.C. 1.2.1.12):gapN (E.C. 1.2.1.9, from *Streptococcus mutans*) substitution/replacement, and disrupting or modifying a soluble transhydrogenase such as sthA (E.C. 1.6.1.2), and/or genetic modifications of one or more of zwf (E.C. 1.1.1.49), gnd (E.C. 1.1.1.44), and edd (E.C. 4.2.1.12). Sequences of these genes are available at [www.metacyc.org](http://www.metacyc.org), and also are available at [www.ncbi.gov](http://www.ncbi.gov).

**[0190]** It is appreciated that any combination of genetic modifications described herein may be used in embodiments of the present invention. Also included within the scope of the invention are isolated nucleic acid and amino acid sequences encoding or comprising the mutations to the polypeptide sequences described herein.

**[0191]** Additional genetic modifications may be provided in a microorganism strain of the present invention. Many such modifications may be provided to impart a particular phenotype.

**[0192]** As one example, a deletion, of multifunctional 2-keto-3-deoxygluconate 6-phosphate aldolase and 2-keto-4-hydroxyglutarate aldolase and oxaloacetate alpha-decarboxylase (eda in *E. coli*), may be provided to various strains.

**[0193]** For example, the ability to utilize sucrose may be provided, and this would expand the range of feed stocks that can be utilized to produce 3-HP. Common laboratory and industrial strains of *E. coli*, such as the strains described herein, are not capable of utilizing sucrose as the sole carbon source. Since sucrose, and sucrose-containing feed stocks such as molasses, are abundant and often used as feed stocks for the production by microbial fermentation, adding appropriate genetic modifications to permit uptake and use of sucrose may be practiced in strains having other features as provided herein. Various sucrose uptake and metabolism systems are known in the art (for example, U.S. Pat. No. 6,960,455), incorporated by reference for such teachings. These and other approaches may be provided in strains of the present invention. The examples provide at least two approaches.

**[0194]** Also, genetic modifications may be provided to add functionality for breakdown of more complex carbon sources, such as cellulosic biomass or products thereof, for uptake, and/or for utilization of such carbon sources. For

example, numerous cellulases and cellulase-based cellulose degradation systems have been studied and characterized (see, for example, and incorporated by reference herein for such teachings, Beguin, P and Aubert, J-P (1994) FEMS Microbiol. Rev. 13: 25-58; Ohima, K. et al. (1997) Biotechnol. Genet. Eng. Rev. 14: 365414).

**[0195]** In addition to the above-described genetic modifications, in various embodiments genetic modifications also are provided to increase the pool and availability of the cofactor NADPH, and/or, consequently, the NADPH/NADP<sup>+</sup> ratio. For example, in various embodiments for *E. coli*, this may be done by increasing activity, such as by genetic modification, of one or more of the following genes—*pgi* (in a mutated form), *pntAB*, overexpressed, *gapA:gapN* substitution/replacement, and disrupting or modifying a soluble transhydrogenase such as *sthA*, and/or genetic modifications of one or more of *zwf*, *gnd*, and *edd*.

**[0196]** Any such genetic modifications may be provided to species not having such functionality, or having a less than desired level of such functionality.

**[0197]** More generally, and depending on the particular metabolic pathways of a microorganism selected for genetic modification, any subgroup of genetic modifications may be made to decrease cellular production of fermentation product (s) selected from the group consisting of acetate, acetoin, acetone, acrylic, malate, fatty acid ethyl esters, isoprenoids, glycerol, ethylene glycol, ethylene, propylene, butylene, isobutylene, ethyl acetate, vinyl acetate, other acetates, 1,4-butanediol, 2,3-butanediol, butanol, isobutanol, sec-butanol, butyrate, isobutyrate, 2-OH-isobutyrate, 3-OH-butyrate, ethanol, isopropanol, D-lactate, L-lactate, pyruvate, itaconate, levulinate, glucarate, glutarate, caprolactam, adipic acid, propanol, isopropanol, fusel alcohols, and 1,2-propanediol, 1,3-propanediol, formate, fumaric acid, propionic acid, succinic acid, valeric acid, and maleic acid. Gene deletions may be made as disclosed generally herein, and other approaches may also be used to achieve a desired decreased cellular production of selected fermentation products.

**[0198]** VII. Separation and Purification of the Chemical Product 3-HP

**[0199]** When 3-HP is the chemical product, the 3-HP may be separated and purified by the approaches described in the following paragraphs, taking into account that many methods of separation and purification are known in the art and the following disclosure is not meant to be limiting. Osmotic shock, sonication, homogenization, and/or a repeated freeze-thaw cycle followed by filtration and/or centrifugation, among other methods, such as pH adjustment and heat treatment, may be used to produce a cell-free extract from intact cells. Any one or more of these methods also may be employed to release 3-HP from cells as an extraction step.

**[0200]** Further as to general processing of a bio-production broth comprising 3-HP, various methods may be practiced to remove biomass and/or separate 3-HP from the culture broth and its components. Methods to separate and/or concentrate the 3-HP include centrifugation, filtration, extraction, chemical conversion such as esterification, distillation (which may result in chemical conversion, such as dehydration to acrylic acid, under some reactive-distillation conditions), crystallization, chromatography, and ion-exchange, in various forms. Additionally, cell rupture may be conducted as needed to release 3-HP from the cell mass, such as by sonication, homogenization, pH adjustment or heating. 3-HP may be further separated and/or purified by methods known in the art,

including any combination of one or more of centrifugation, liquid-liquid separations, including extractions such as solvent extraction, reactive extraction, two-phase aqueous extraction and two-phase solvent extraction, membrane separation technologies, distillation, evaporation, ion-exchange chromatography, adsorption chromatography, reverse phase chromatography and crystallization. Any of the above methods may be applied to a portion of a bio-production broth (i.e., a fermentation broth, whether made under aerobic, anaerobic, or microaerobic conditions), such as may be removed from a bio-production event gradually or periodically, or to the broth at termination of a bio-production event. Conversion of 3-HP to downstream products, such as described herein, may proceed after separation and purification, or, such as with distillation, thin-film evaporation, or wiped-film evaporation optionally also in part as a separation means.

**[0201]** For various of these approaches, one may apply a counter-current strategy, or a sequential or iterative strategy, such as multi-pass extractions. For example, a given aqueous solution comprising 3-HP may be repeatedly extracted with a non-polar phase comprising an amine to achieve multiple reactive extractions.

**[0202]** When a culture event (fermentation event) is at a point of completion, the spent broth may be transferred to a separate tank, or remain in the culture vessel, and in either case the temperature may be elevated to at least 60° C. for a minimum of one hour in order to kill the microorganisms. (Alternatively, other approaches to killing the microorganisms may be practiced.) By spent broth is meant the final liquid volume comprising the initial nutrient media, cells grown from the microorganism inoculum (and possibly including some original cells of the inoculum), 3-HP, and optionally liquid additions made after providing the initial nutrient media, such as periodic additions to provide additional carbon source, etc. It is noted that the spent broth may comprise organic acids other than 3-HP, such as for example acetic acid and/or lactic acid.

**[0203]** A centrifugation step may then be practiced to filter out the biomass solids (e.g., dead microorganism cells). This may be achieved in a continuous or batch centrifuge, and solids removal may be at least about 80%, 85%, 90%, or 95% in a single pass, or cumulatively after two or more serial centrifugations.

**[0204]** An optional step is to polish the centrifuged liquid through a filter, such as microfiltration or ultrafiltration, or may comprise a filter press or other filter device to which is added a filter aid such as diatomaceous earth. Alternative or supplemental approaches to this and the centrifugation may include removal of cells by a flocculent, where the cells floc and are allowed to settle, and the liquid is drawn off or otherwise removed. A flocculent may be added to a fermentation broth after which settling of material is allowed for a time, and then separations may be applied, including but not limited to centrifugation.

**[0205]** After such steps, a spent broth comprising 3-HP and substantially free of solids is obtained for further processing. By “substantially free of solids” is meant that greater than 98%, 99%, or 99.5% of the solids have been removed.

**[0206]** In various embodiments this spent broth comprises various ions of salts, such as Na, Cl, SO<sub>4</sub>, and PO<sub>4</sub>. In some embodiments these ions may be removed by passing this spent broth through ion exchange columns, or otherwise contacting the spent broth with appropriate ion exchange material. Here and elsewhere in this document, “contacting” is

taken to mean a contacting for the stated purpose by any way known to persons skilled in the art, such as, for example, in a column, under appropriate conditions that are well within the ability of persons of ordinary skill in the relevant art to determine. As but one example, these may comprise sequential contacting with anion and cation exchange materials (in any order), or with a mixed anion/cation material. This demineralization step should remove most such inorganic ions without removing the 3-HP. This may be achieved, for example, by lowering the pH sufficiently to protonate 3-HP and similar organic acids so that these acids are not bound to the anion exchange material, whereas anions, such as Cl and SO<sub>4</sub>, that remain charged at such pH are removed from the solution by binding to the resin. Likewise, positively charged ions are removed by contacting with cation exchange material. Such removal of ions may be assessed by a decrease in conductivity of the solution. Such ion exchange materials may be regenerated by methods known to those skilled in the art.

[0207] In some embodiments, the spent broth (such as but not necessarily after the previous demineralization step) is subjected to a pH elevation, after which it is passed through an ion exchange column, or otherwise contacted with an ion exchange resin, that comprises anionic groups, such as amines, to which organic acids, ionic at this pH, associate. Other organics that do not so associate with amines at this pH (which may be over 6.5, over 7.5, over 8.5, over 9.5, over 10.5, or higher pH) may be separated from the organic acids at this stage, such as by flushing with an elevated pH rinse. Thereafter elution with a lower pH and/or elevated salt content rinse may remove the organic acids. Eluting with a gradient of decreasing pH and/or increasing salt content rinses may allow more distinct separation of 3-HP from other organic acids, thereafter simplifying further processing.

[0208] This latter step of anion-exchange resin retention of organic acids may be practiced before or after the demineralization step. However, the following two approaches are alternatives to the anion-exchange resin step.

[0209] A first alternative approach comprises reactive extraction (a form of liquid-liquid extraction) as exemplified in this and the following paragraphs. The spent broth, which may be at a stage before or after the demineralization step above, is combined with a quantity of a tertiary amine such as Alamine336® (Cognis Corp., Cincinnati, Ohio USA) at low pH. Co-solvents for the Alamine336 or other tertiary amine may be added and include, but are not limited to benzene, carbon tetrachloride, chloroform, cyclohexane, disobutyl ketone, ethanol, #2 fuel oil, isopropanol, kerosene, n-butanol, isobutanol, octanol, and n-decanol that increase the partition coefficient when combined with the amine. After appropriate mixing a period of time for phase separation transpires, after which the non-polar phase, which comprises 3-HP associated with the Alamine336 or other tertiary amine, is separated from the aqueous phase.

[0210] When a co-solvent is used that has a lower boiling point than the 3-HP/tertiary amine, a distilling step may be used to remove the co-solvent, thereby leaving the 3-HP-tertiary amine complex in the non-polar phase.

[0211] Whether or not there is such a distillation step, a stripping or recovery step may be used to separate the 3-HP from the tertiary amine. An inorganic salt, such as ammonium sulfate, sodium chloride, or sodium carbonate, or a base such as sodium hydroxide or ammonium hydroxide, is added to the 3-HP/tertiary amine to reverse the amine protonation reaction, and a second phase is provided by addition of an aqueous

solution (which may be the vehicle for provision of the inorganic salt). After suitable mixing, two phases result and this allows for tertiary amine regeneration and re-use, and provides the 3-HP in an aqueous solution. Alternatively, hot water may also be used without a salt or base to recover the 3HP from the amine.

[0212] In the above approach the phase separation and extraction of 3-HP to the aqueous phase can serve to concentrate the 3-HP. It is noted that chromatographic separation of respective organic acids also can serve to concentrate such acids, such as 3-HP. In similar approaches other suitable, non-polar amines, which may include primary, secondary and quaternary amines, may be used instead of and/or in combination with a tertiary amine.

[0213] A second alternative approach is crystallization. For example, the spent broth (such as free of biomass solids) may be contacted with a strong base such as ammonium hydroxide, which results in formation of an ammonium salt of 3-HP. This may be concentrated, and then ammonium-3-HP crystals are formed and may be separated, such as by filtration, from the aqueous phase. Once collected, ammonium-3-HP crystals may be treated with an acid, such as sulfuric acid, so that ammonium sulfate is regenerated, so that 3-HP and ammonium sulfate result.

[0214] Also, various aqueous two-phase extraction methods may be utilized to separate and/or concentrate a desired chemical product from a fermentation broth or later-obtained solution. It is known that the addition of polymers, such as dextran and glycol polymers, such as polyethylene glycol (PEG) and polypropylene glycol (PPG) to an aqueous solution may result in formation of two aqueous phases. In such systems a desired chemical product may segregate to one phase while cells and other chemicals partition to the other phase, thus providing for a separation without use of organic solvents. This approach has been demonstrated for some chemical products, but challenges associated with chemical product recovery from a polymer solution and low selectivities are recognized (See "Extractive Recovery of Products from Fermentation Broths," Joong Kyun Kim et al., *Biotechnol. Bioprocess Eng.*, 1999(4)1-11, incorporated by reference for all of its teachings of extractive recovery methods).

[0215] Various substitutions and combinations of the above steps and processes may be made to obtain a relatively purified 3-HP solution. Also, methods of separation and purification disclosed in U.S. Pat. No. 6,534,679, issued Mar. 18, 2003, and incorporated by reference herein for such methods disclosures, may be considered based on a particular processing scheme. Also, in some culture events periodic removal of a portion of the liquid volume may be made, and processing of such portion(s) may be made to recover the 3-HP, including by any combination of the approaches disclosed above.

[0216] As noted, solvent extraction is another alternative. This may use any of a number of and/or combinations of solvents, including alcohols, esters, ketones, and various organic solvents. Without being limiting, after phase separation a distillation step or a secondary extraction may be employed to separate 3-HP from the organic phase.

[0217] The following published resources are incorporated by reference herein for their respective teachings to indicate the level of skill in these relevant arts, and as needed to support a disclosure that teaches how to make and use methods of industrial bio-production of 3-HP, and also industrial systems that may be used to achieve such conversion with any of the recombinant microorganisms of the present invention

(*Biochemical Engineering Fundamentals*, 2<sup>nd</sup> Ed. J. E. Bailey and D. F. Ollis, McGraw Hill, New York, 1986, entire book for purposes indicated and Chapter 9, pp. 533-657 in particular for biological reactor design; *Unit Operations of Chemical Engineering*, 5<sup>th</sup> Ed., W. L. McCabe et al., McGraw Hill, New York 1993, entire book for purposes indicated, and particularly for process and separation technologies analyses; *Equilibrium Staged Separations*, P. C. Wankat, Prentice Hall, Englewood Cliffs, N.J. USA, 1988, entire book for separation technologies teachings).

**[0218]** VII. Conversion of 3-HP to Acrylic Acid and Downstream Products

**[0219]** As discussed herein, various embodiments described herein are related to production of a particular chemical product, 3-hydroxypropionic acid (3-HP). This organic acid, 3-HP, may be converted to various other products having industrial uses, such as but not limited to acrylic acid, esters of acrylic acid, and other chemicals obtained from 3-HP, referred to as “downstream products.” Under some approaches the 3-HP may be converted to acrylic acid, acrylamide, and/or other downstream chemical products, in some instances the conversion being associated with the separation and/or purification steps. Many conversions to such downstream products are described herein. The methods of the invention include steps to produce downstream products of 3-HP.

**[0220]** As a C<sub>3</sub> building block, 3-HP offers much potential in a variety of chemical conversions to commercially important intermediates, industrial end products, and consumer products. For example, 3-HP may be converted to acrylic acid, acrylates (e.g., acrylic acid salts and esters), 1,3-propanediol, malonic acid, ethyl-3-hydroxypropionate, ethyl ethoxy propionate, propiolactone, acrylamide, or acrylonitrile.

**[0221]** For example, methyl acrylate may be made from 3-HP via dehydration and esterification, the latter to add a methyl group (such as using methanol); acrylamide may be made from 3-HP via dehydration and amidation reactions; acrylonitrile may be made via a dehydration reaction and forming a nitrile moiety; propiolactone may be made from 3-HP via a ring-forming internal esterification reaction (eliminating a water molecule); ethyl-3-HP may be made from 3-HP via esterification with ethanol; malonic acid may be made from 3-HP via an oxidation reaction; and 1,3-propanediol may be made from 3-HP via a reduction reaction. Also, acrylic acid, first converted from 3-HP by dehydration, may be esterified with appropriate compounds to form a number of commercially important acrylate-based esters, including but not limited to methyl acrylate, ethyl acrylate, methyl acrylate, 2-ethylhexyl acrylate, butyl acrylate, and lauryl acrylate. Alternatively, 3HP may be esterified to form an ester of 3HP and then dehydrated to form the acrylate ester.

**[0222]** Additionally, 3-HP may be oligomerized or polymerized to form poly(3-hydroxypropionate) homopolymers, or co-polymerized with one or more other monomers to form various co-polymers. Because 3-HP has only a single stereoisomer, polymerization of 3-HP is not complicated by the stereo-specificity of monomers during chain growth. This is in contrast to (S)-2-Hydroxypropanoic acid (also known as lactic acid), which has two (D, L) stereoisomers that must be considered during its polymerizations.

**[0223]** As will be further described, 3-HP can be converted into derivatives starting (i) substantially as the protonated form of 3-hydroxypropionic acid; (ii) substantially as the

deprotonated form, 3-hydroxypropionate; or (iii) as mixtures of the protonated and deprotonated forms. Generally, the fraction of 3-HP present as the acid versus the salt will depend on the pH, the presence of other ionic species in solution, temperature (which changes the equilibrium constant relating the acid and salt forms), and to some extent pressure. Many chemical conversions may be carried out from either of the 3-HP forms, and overall process economics will typically dictate the form of 3-HP for downstream conversion.

**[0224]** Also, as an example of a conversion during separation, 3-HP in an amine salt form, such as in the extraction step herein disclosed using Alamine 336 as the amine, may be converted to acrylic acid by contacting a solution comprising the 3-HP amine salt with a dehydration catalyst, such as aluminum oxide, at an elevated temperature, such as 170 to 180 C, or 180 to 190 C, or 190 to 200 C, and passing the collected vapor phase over a low temperature condenser. Operating conditions, including 3-HP concentration, organic amine, co-solvent (if any), temperature, flow rates, dehydration catalyst, and condenser temperature, are evaluated and improved for commercial purposes. Conversion of 3-HP to acrylic acid is expected to exceed at least 80 percent, or at least 90 percent, in a single conversion event. The amine may be re-used, optionally after clean-up. Other dehydration catalysts, as provided herein, may be evaluated. It is noted that U.S. Pat. No. 7,186,856 discloses data regarding this conversion approach, albeit as part of an extractive salt-splitting conversion that differs from the teachings herein. However, U.S. Pat. No. 7,186,856 is incorporated by reference for its methods, including extractive salt-splitting, the latter to further indicate the various ways 3-HP may be extracted from a microbial fermentation broth.

**[0225]** Further as to embodiments in which the chemical product being synthesized by the microorganism host cell is 3-HP, made as provided herein and optionally purified to a selected purity prior to conversion, the methods of the present invention can also be used to produce “downstream” compounds derived from 3-HP, such as polymerized-3-HP (poly-3-HP), acrylic acid, polyacrylic acid (polymerized acrylic acid, in various forms), methyl acrylate, acrylamide, acrylonitrile, propiolactone, ethyl 3-HP, malonic acid, and 1,3-propanediol. Numerous approaches may be employed for such downstream conversions, generally falling into enzymatic, catalytic (chemical conversion process using a catalyst), thermal, and combinations thereof (including some wherein a desired pressure is applied to accelerate a reaction).

**[0226]** As noted, an important industrial chemical product that may be produced from 3-HP is acrylic acid. Chemically, one of the carbon-carbon single bonds in 3-HP must undergo a dehydration reaction, converting to a carbon-carbon double bond and rejecting a water molecule. Dehydration of 3-HP in principle can be carried out in the liquid phase or in the gas phase. In some embodiments, the dehydration takes place in the presence of a suitable homogeneous or heterogeneous catalyst. Suitable dehydration catalysts are both acid and alkaline catalysts. Following dehydration, an acrylic acid-containing phase is obtained and can be purified where appropriate by further purification steps, such as by distillation methods, extraction methods, or crystallization methods, or combinations thereof.

**[0227]** Making acrylic acid from 3-HP via a dehydration reaction may be achieved by a number of commercial methodologies including via a distillation process, which may be part of the separation regime and which may include an acid

and/or a metal ion as catalyst. More broadly, incorporated herein for its teachings of conversion of 3-HP, and other  $\beta$ -hydroxy carbonyl compounds, to acrylic acid and other related downstream compounds, is U.S. Patent Publication No. 2007/0219390 A1, published Sep. 20, 2007, now abandoned. This publication lists numerous catalysts and provides examples of conversions, which are specifically incorporated herein. Also among the various specific methods to dehydrate 3-HP to produce acrylic acid is an older method, described in U.S. Pat. No. 2,469,701 (Redmon). This reference teaches a method for the preparation of acrylic acid by heating 3-HP to a temperature between 130 and 190° C., in the presence of a dehydration catalyst, such as sulfuric acid or phosphoric acid, under reduced pressure. U.S. Patent Publication No. 2005/0222458 A1 (Craciun et al.) also provides a process for the preparation of acrylic acid by heating 3-HP or its derivatives. Vapor-phase dehydration of 3-HP occurs in the presence of dehydration catalysts, such as packed beds of silica, alumina, or titania. These patent publications are incorporated by reference for their methods relating to converting 3-HP to acrylic acid.

**[0228]** The dehydration catalyst may comprise one or more metal oxides, such as  $\text{Al}_2\text{O}_3$ ,  $\text{SiO}_2$ , or  $\text{TiO}_2$ . In some embodiments, the dehydration catalyst is a high surface area  $\text{Al}_2\text{O}_3$  or a high surface area silica wherein the silica is substantially  $\text{SiO}_2$ . High surface area for the purposes of the invention means a surface area of at least about 50, 75, 100  $\text{m}^2/\text{g}$ , or more. In some embodiments, the dehydration catalyst may comprise an aluminosilicate, such as a zeolite.

**[0229]** For example, including as exemplified from such incorporated references, 3-HP may be dehydrated to acrylic acid via various specific methods, each often involving one or more dehydration catalysts. One catalyst of particular apparent value is titanium, such as in the form of titanium oxide,  $\text{TiO}_2$ . A titanium dioxide catalyst may be provided in a dehydration system that distills an aqueous solution comprising 3-HP, wherein the 3-HP dehydrates, such as upon volatilization, converting to acrylic acid, and the acrylic acid is collected by condensation from the vapor phase.

**[0230]** As but one specific method, an aqueous solution of 3-HP is passed through a reactor column packed with a titanium oxide catalyst maintained at a temperature between 170 and 190 C and at ambient atmospheric pressure. Vapors leaving the reactor column are passed over a low temperature condenser, where acrylic acid is collected. The low temperature condenser may be cooled to 30 C or less, 2 C or less, or at any suitable temperature for efficient condensation based on the flow rate and design of the system. Also, the reactor column temperatures may be lower, for instance when operating at a pressure lower than ambient atmospheric pressure. It is noted that Example 1 of U.S. Patent Publication No. 2007/0219390, published Sep. 20, 2007, now abandoned, provides specific parameters that employs the approach of this method. As noted, this publication is incorporated by reference for this teaching and also for its listing of catalysts that may be used in a 3-HP to acrylic acid dehydration reaction.

**[0231]** Further as to dehydration catalysts, the following table summarizes a number of catalysts (including chemical classes) that may be used in a dehydration reaction from 3-HP (or its esters) to acrylic acid (or acrylate esters). Such catalysts, some of which may be used in any of solid, liquid or gaseous forms, may be used individually or in any combination. This listing of catalysts is not intended to be limiting, and

many specific catalysts not listed may be used for specific dehydration reactions. Further without being limiting, catalyst selection may depend on the solution pH and/or the form of 3-HP in a particular conversion, so that an acidic catalyst may be used when 3-HP is in acidic form, and a basic catalyst may be used when the ammonium salt of 3-HP is being converted to acrylic acid. Also, some catalysts may be in the form of ion exchange resins.

TABLE 4

Dehydration Catalysts	
Catalyst by Chemical Class	Non-limiting Examples
Acids (including weak and strong)	$\text{H}_2\text{SO}_4$ , HCl, titanitic acids, metal oxide hydrates, metal sulfates ( $\text{MSO}_4$ , where M = Zn, Sn, Ca, Ba, Ni, Co, or other transition metals), metal oxide sulfates, metal phosphates (e.g., $\text{M}_3(\text{PO}_4)_2$ , where M = Ca, Ba), metal phosphates, metal oxide phosphates, carbon (e.g., transition metals on a carbon support), mineral acids, carboxylic acids, salts thereof, acidic resins, acidic zeolites, clays, $\text{SiO}_2/\text{H}_3\text{PO}_4$ , fluorinated $\text{Al}_2\text{O}_3$ , $\text{Nb}_2\text{O}_3/\text{PO}_5^{-3}$ , $\text{Nb}_2\text{O}_3/\text{SO}_4^{-2}$ , $\text{Nb}_2\text{O}_5\text{H}_2\text{O}$ , phosphotungstic acids, phosphomolybdic acids, silicomolybdic acids, silicotungstic acids, carbon dioxide
Bases (including weak and strong)	NaOH, ammonia, polyvinylpyridine, metal hydroxides, $\text{Zr}(\text{OH})_4$ , and substituted amines
Oxides (generally metal oxides)	$\text{TiO}_2$ , $\text{ZrO}_2$ , $\text{Al}_2\text{O}_3$ , $\text{SiO}_2$ , $\text{ZnO}_2$ , $\text{SnO}_2$ , $\text{WO}_3$ , $\text{MnO}_2$ , $\text{Fe}_2\text{O}_3$ , $\text{V}_2\text{O}_5$

**[0232]** As to another specific method using one of these catalysts, concentrated sulfuric acid and an aqueous solution comprising 3-HP are separately flowed into a reactor maintained at 150 to 165° C. at a reduced pressure of 100 mm Hg. Flowing from the reactor is a solution comprising acrylic acid. A specific embodiment of this method, disclosed in Example 1 of US2009/0076297, incorporated by reference herein, indicates a yield of acrylic acid exceeding 95 percent.

**[0233]** Based on the wide range of possible catalysts and knowledge in the art of dehydration reactions of this type, numerous other specific dehydration methods may be evaluated and implemented for commercial production.

**[0234]** The dehydration of 3-HP may also take place in the absence of a dehydration catalyst. For example, the reaction may be run in the vapor phase in the presence of a nominally inert packing such as glass, ceramic, a resin, porcelain, plastic, metallic or brick dust packing and still form acrylic acid in reasonable yields and purity. The catalyst particles can be sized and configured such that the chemistry is, in some embodiments, mass-transfer-limited or kinetically limited. The catalyst can take the form of powder, pellets, granules, beads, extrudates, and so on. When a catalyst support is optionally employed, the support may assume any physical form such as pellets, spheres, monolithic channels, etc. The supports may be co-precipitated with active metal species; or the support may be treated with the catalytic metal species and then used as is or formed into the aforementioned shapes; or the support may be formed into the aforementioned shapes and then treated with the catalytic species.

**[0235]** A reactor for dehydration of 3-HP may be engineered and operated in a wide variety of ways. The reactor operation can be continuous, semi-continuous, or batch. It is perceived that an operation that is substantially continuous and at steady state is advantageous from operations and economics perspectives. The flow pattern can be substantially

plug flow, substantially well-mixed, or a flow pattern between these extremes. A “reactor” can actually be a series or network of several reactors in various arrangements.

**[0236]** For example, without being limiting, acrylic acid may be made from 3-HP via a dehydration reaction, which may be achieved by a number of commercial methodologies including via a distillation process, which may be part of the separation regime and which may include an acid and/or a metal ion as catalyst. More broadly, incorporated herein for its teachings of conversion of 3-HP, and other 0-hydroxy carbonyl compounds, to acrylic acid and other related downstream compounds, is U.S. Patent Publication No. 2007/0219390 A1, published Sep. 20, 2007, now abandoned. This publication lists numerous catalysts and provides examples of conversions, which are specifically incorporated herein.

**[0237]** For example, including as exemplified from such incorporated references, 3-HP may be dehydrated to acrylic acid via various specific methods, each often involving one or more dehydration catalysts. One catalyst of particular apparent value is titanium, such as in the form of titanium oxide, TiO<sub>2</sub>. A titanium dioxide catalyst may be provided in a dehydration system that distills an aqueous solution comprising 3-HP, wherein the 3-HP dehydrates, such as upon volatilization, converting to acrylic acid, and the acrylic acid is collected by condensation from the vapor phase.

**[0238]** As but one specific method, an aqueous solution of 3-HP is passed through a reactor column packed with a titanium oxide catalyst maintained at a temperature between 170 and 190° C. and at ambient atmospheric pressure. Vapors leaving the reactor column are passed over a low temperature condenser, where acrylic acid is collected. The low temperature condenser may be cooled to 30° C. or less, 20° C. or less, 2° C. or less, or at any suitable temperature for efficient condensation based on the flow rate and design of the system. Also, the reactor column temperatures may be lower, for instance when operating at a pressure lower than ambient atmospheric pressure. It is noted that Example 1 of U.S. Patent Publication No. 2007/0219390, published Sep. 20, 2007, now abandoned, provides specific parameters that employs the approach of this method. As noted, this publication is incorporated by reference for this teaching and also for its listing of catalysts that may be used in a 3-HP to acrylic acid dehydration reaction.

**[0239]** Crystallization of the acrylic acid obtained by dehydration of 3-HP may be used as one of the final separation/purification steps. Various approaches to crystallization are known in the art, including crystallization of esters.

**[0240]** As noted above, in some embodiments, a salt of 3-HP is converted to acrylic acid or an ester or salt thereof. For example, U.S. Pat. No. 7,186,856 (Meng et al.) teaches a process for producing acrylic acid from the ammonium salt of 3-HP, which involves a first step of heating the ammonium salt of 3-HP in the presence of an organic amine or solvent that is immiscible with water, to form a two-phase solution and split the 3-HP salt into its respective ionic constituents under conditions which transfer 3-HP from the aqueous phase to the organic phase of the solution, leaving ammonia and ammonium cations in the aqueous phase. The organic phase is then back-extracted to separate the 3-HP, followed by a second step of heating the 3-HP-containing solution in the presence of a dehydration catalyst to produce acrylic acid. U.S. Pat. No. 7,186,856 is incorporated by reference for its methods for producing acrylic acid from salts of 3-HP. Various

alternatives to the particular approach disclosed in this patent may be developed for suitable extraction and conversion processes.

**[0241]** Methyl acrylate may be made from 3-HP via dehydration and esterification, the latter to add a methyl group (such as using methanol), acrylamide may be made from 3-HP via dehydration and amidation reactions, acrylonitrile may be made via a dehydration reaction and forming a nitrile moiety, propiolactone may be made from 3-HP via a ring-forming internal esterification reaction (eliminating a water molecule), ethyl-3-HP may be made from 3-HP via esterification with ethanol, malonic acid may be made from 3-HP via an oxidation reaction, and 1,3-propanediol may be made from 3-HP via a reduction reaction.

**[0242]** Malonic acid may be produced from oxidation of 3-HP as produced herein. U.S. Pat. No. 5,817,870 (Haas et al.) discloses catalytic oxidation of 3-HP by a precious metal selected from Ru, Rh, Pd, Os, Ir or Pt. These can be pure metal catalysts or supported catalysts. The catalytic oxidation can be carried out using a suspension catalyst in a suspension reactor or using a fixed-bed catalyst in a fixed-bed reactor. If the catalyst, preferably a supported catalyst, is disposed in a fixed-bed reactor, the latter can be operated in a trickle-bed procedure as well as also in a liquid-phase procedure. In the trickle-bed procedure the aqueous phase comprising the 3-HP starting material, as well as the oxidation products of the same and means for the adjustment of pH, and oxygen or an oxygen-containing gas can be conducted in parallel flow or counter-flow. In the liquid-phase procedure the liquid phase and the gas phase are conveniently conducted in parallel flow.

**[0243]** In order to achieve a sufficiently short reaction time, the conversion is carried out at a pH equal or greater than 6, preferably at least 7, and in particular between 7.5 and 9. According to a preferred embodiment, during the oxidation reaction the pH is kept constant, preferably at a pH in the range between 7.5 and 9, by adding a base, such as an alkaline or alkaline earth hydroxide solution. The oxidation is usefully carried out at a temperature of at least 10° C. and maximally 70° C. The flow of oxygen is not limited. In the suspension method it is important that the liquid and the gaseous phase are brought into contact by stirring vigorously. Malonic acid can be obtained in nearly quantitative yields. U.S. Pat. No. 5,817,870 is incorporated by reference herein for its methods to oxidize 3-HP to malonic acid.

**[0244]** 1,3-Propanediol may be produced from hydrogenation of 3-HP as produced herein. U.S. Patent Publication No. 2005/0283029 (Meng et al.) is incorporated by reference herein for its methods to hydrogenation of 3-HP, or esters of the acid or mixtures, in the presence of a specific catalyst, in a liquid phase, to prepare 1,3-propanediol. Possible catalysts include ruthenium metal, or compounds of ruthenium, supported or unsupported, alone or in combination with at least one or more additional metal(s) selected from molybdenum, tungsten, titanium, zirconium, niobium, vanadium or chromium. The ruthenium metal or compound thereof, and/or the additional metal(s), or compound thereof, may be utilized in supported or unsupported form. If utilized in supported form, the method of preparing the supported catalyst is not critical and can be any technique such as impregnation of the support or deposition on the support. Any suitable support may be utilized. Supports that may be used include, but are not limited to, alumina, titania, silica, zirconia, carbons, carbon blacks, graphites, silicates, zeolites, aluminosilicate zeolites, aluminosilicate clays, and the like.



[0245] The hydrogenation process may be carried out in liquid phase. The liquid phase includes water, organic solvents that are not hydrogenatable, such as any aliphatic or aromatic hydrocarbon, alcohols, ethers, toluene, decalin, dioxane, diglyme, n-heptane, hexane, xylene, benzene, tetrahydrofuran, cyclohexane, methylcyclohexane, and the like, and mixtures of water and organic solvent(s). The hydrogenation process may be carried out batch wise, semi-continuously, or continuously. The hydrogenation process may be carried out in any suitable apparatus. Exemplary of such apparatus are stirred tank reactors, trickle-bed reactors, high pressure hydrogenation reactors, and the like.

[0246] The hydrogenation process is generally carried out at a temperature ranging from about 20 to about 250° C., more particularly from about 100 to about 200° C. Further, the hydrogenation process is generally carried out in a pressure range of from about 20 psi to about 4000 psi. The hydrogen containing gas utilized in the hydrogenation process is, optionally, commercially pure hydrogen. The hydrogen containing gas is usable if nitrogen, gaseous hydrocarbons, or oxides of carbon, and similar materials, are present in the hydrogen containing gas. For example, hydrogen from synthesis gas (hydrogen and carbon monoxide) may be employed, such synthesis gas potentially further including carbon dioxide, water, and various impurities.

[0247] As is known in the art, it is also possible to convert 3-HP to 1,3-propanediol using biological methods. For example, 1,3-propanediol can be created from either 3-HP-CoA or 3-HP via the use of polypeptides having enzymatic activity. These polypeptides can be used either in vitro or in vivo. When converting 3-HP-CoA to 1,3-propanediol, polypeptides having oxidoreductase activity or reductase activity (e.g., enzymes from the 1.1.1.-class of enzymes) can be used. Alternatively, when creating 1,3-propanediol from 3-HP, a combination of a polypeptide having aldehyde dehydrogenase activity (e.g., an enzyme from the 1.1.1.34 class) and a polypeptide having alcohol dehydrogenase activity (e.g., an enzyme from the 1.1.1.32 class) can be used.

[0248] Another downstream production of 3-HP, acrylonitrile, may be converted from acrylic acid by various organic syntheses, including by not limited to the Sohio acrylonitrile process, a single-step method of production known in the chemical manufacturing industry

[0249] Also, addition reactions may yield acrylic acid or acrylate derivatives having alkyl or aryl groups at the carbonyl hydroxyl group. Such additions may be catalyzed chemically, such as by hydrogen, hydrogen halides, hydrogen cyanide, or Michael additions under alkaline conditions optionally in the presence of basic catalysts. Alcohols, phenols, hydrogen sulfide, and thiols are known to add under basic conditions. Aromatic amines or amides, and aromatic hydrocarbons, may be added under acidic conditions. These and other reactions are described in *Ulmann's Encyclopedia of Industrial Chemistry, Acrylic Acid and Derivatives*, WileyVCH Verlag GmbH, Weinham (2005), incorporated by reference for its teachings of conversion reactions for acrylic acid and its derivatives.

[0250] Acrylic acid obtained from 3-HP made by the present invention may be further converted to various chemicals, including polymers, which are also considered downstream products in some embodiments. Acrylic acid esters may be formed from acrylic acid (or directly from 3-HP) such as by condensation esterification reactions with an alcohol, releasing water. This chemistry described in *Monomeric*

*Acrylic Esters*, E. H. Riddle, Reinhold, N.Y. (1954), incorporated by reference for its esterification teachings. Among esters that are formed are methyl acrylate, ethyl acrylate, n-butyl acrylate, hydroxypropyl acrylate, hydroxyethyl acrylate, isobutyl acrylate, and 2-ethylhexyl acrylate, and these and/or other acrylic acid and/or other acrylate esters may be combined, including with other compounds, to form various known acrylic acid-based polymers. Although acrylamide is produced in chemical syntheses by hydration of acrylonitrile, herein a conversion may convert acrylic acid to acrylamide by amidation.

[0251] Acrylic acid obtained from 3-HP made by the present invention may be further converted to various chemicals, including polymers, which are also considered downstream products in some embodiments. Acrylic acid esters may be formed from acrylic acid (or directly from 3-HP) such as by condensation esterification reactions with an alcohol, releasing water. This chemistry is described in *Monomeric Acrylic Esters*, E. H. Riddle, Reinhold, N.Y. (1954), incorporated by reference for its esterification teachings. Among esters that are formed are methyl acrylate, ethyl acrylate, n-butyl acrylate, hydroxypropyl acrylate, hydroxyethyl acrylate, isobutyl acrylate, and 2-ethylhexyl acrylate, and these and/or other acrylic acid and/or other acrylate esters may be combined, including with other compounds, to form various known acrylic acid-based polymers. Although acrylamide is produced in chemical syntheses by hydration of acrylonitrile, herein a conversion may convert acrylic acid to acrylamide by amidation.

[0252] Direct esterification of acrylic acid can take place by esterification methods known to the person skilled in the art, by contacting the acrylic acid obtained from 3-HP dehydration with one or more alcohols, such as methanol, ethanol, 1-propanol, 2-propanol, n-butanol, tert-butanol or isobutanol, and heating to a temperature of at least 50, 75, 100, 125, or 150° C. The water formed during esterification may be removed from the reaction mixture, such as by azeotropic distillation through the addition of suitable separation aids, or by another means of separation. Conversions up to 95%, or more, may be realized, as is known in the art.

[0253] Several suitable esterification catalysts are commercially available, such as from Dow Chemical (Midland, Mich. US). For example, Amberlyst™ 131Wet Monodisperse gel catalyst confers enhanced hydraulic and reactivity properties and is suitable for fixed bed reactors. Amberlyst™ 39Wet is a macroreticular catalyst suitable particularly for stirred and slurry loop reactors. Amberlyst™ 46 is a macroporous catalyst producing less ether byproducts than conventional catalyst (as described in U.S. Pat. No. 5,426,199 to Rohm and Haas, which patent is incorporated by reference for its teachings of esterification catalyst compositions and selection considerations).

[0254] Acrylic acid, and any of its esters, may be further converted into various polymers. Polymerization may proceed by any of heat, light, other radiation of sufficient energy, and free radical generating compounds, such as azo compounds or peroxides, to produce a desired polymer of acrylic acid or acrylic acid esters. As one example, an aqueous acrylic acid solution's temperature raised to a temperature known to start polymerization (in part based on the initial acrylic acid concentration), and the reaction proceeds, the process frequently involving heat removal given the high exothermicity of the reaction. Many other methods of polymerization are known in the art. Some are described in *Ulmann's Encyclo-*

pedia of Industrial Chemistry, Polyacrylamides and Poly (Acrylic Acids), WileyVCH Verlag GmbH, Wienham (2005), incorporated by reference for its teachings of polymerization reactions.

**[0255]** For example, the free-radical polymerization of acrylic acid takes place by polymerization methods known to the skilled worker and can be carried out either in an emulsion or suspension in aqueous solution or another solvent. Initiators, such as but not limited to organic peroxides, often are added to aid in the polymerization. Among the classes of organic peroxides that may be used as initiators are diacyls, peroxydicarbonates, monoperoxydicarbonates, peroxyketals, peroxyesters, dialkyls, and hydroperoxides. Another class of initiators is azo initiators, which may be used for acrylate polymerization as well as co-polymerization with other monomers. U.S. Pat. Nos. 5,470,928; 5,510,307; 6,709,919; and 7,678,869 teach various approaches to polymerization using a number of initiators, including organic peroxides, azo compounds, and other chemical types, and are incorporated by reference for such teachings as applicable to the polymers described herein.

**[0256]** Accordingly, it is further possible for co-monomers, such as crosslinkers, to be present during the polymerization. The free-radical polymerization of the acrylic acid obtained from dehydration of 3-HP, as produced herein, in at least partly neutralized form and in the presence of crosslinkers is practiced in certain embodiments. This polymerization may result in hydrogels which can then be comminuted, ground and, where appropriate, surface-modified, by known techniques.

**[0257]** An important commercial use of polyacrylic acid is for superabsorbent polymers. This specification hereby incorporates by reference Modern Superabsorbent Polymer Technology, Buchholz and Graham (Editors), Wiley-VCH, 1997, in its entirety for its teachings regarding superabsorbent polymers components, manufacture, properties and uses. Superabsorbent polymers are primarily used as absorbents for water and aqueous solutions for diapers, adult incontinence products, feminine hygiene products, and similar consumer products. In such consumer products, superabsorbent materials can replace traditional absorbent materials such as cloth, cotton, paper wadding, and cellulose fiber. Superabsorbent polymers absorb, and retain under a slight mechanical pressure, up to 25 times or their weight in liquid. The swollen gel holds the liquid in a solid, rubbery state and prevents the liquid from leaking. Superabsorbent polymer particles can be surface-modified to produce a shell structure with the shell being more highly crosslinked. This technique improves the balance of absorption, absorption under load, and resistance to gel-blocking. It is recognized that superabsorbent polymers have uses in fields other than consumer products, including agriculture, horticulture, and medicine.

**[0258]** Superabsorbent polymers are prepared from acrylic acid (such as acrylic acid derived from 3-HP provided herein) and a crosslinker, by solution or suspension polymerization. Exemplary methods include U.S. Pat. Nos. 5,145,906; 5,350,799; 5,342,899; 4,857,610; 4,985,518; 4,708,997; 5,180,798; 4,666,983; 4,734,478; and 5,331,059, each incorporated by reference for their teachings relating to superabsorbent polymers.

**[0259]** Among consumer products, a diaper, a feminine hygiene product, and an adult incontinence product are made

with superabsorbent polymer that itself is made substantially from acrylic acid converted from 3-HP made in accordance with the present invention.

**[0260]** Diapers and other personal hygiene products may be produced that incorporate superabsorbent polymer made from acrylic acid made from 3-HP which is bio-produced by the teachings of the present application. The following provides general guidance for making a diaper that incorporates such superabsorbent polymer. The superabsorbent polymer first is prepared into an absorbent pad that may be vacuum formed, and in which other materials, such as a fibrous material (e.g., wood pulp) are added. The absorbent pad then is assembled with sheet(s) of fabric, generally a nonwoven fabric (e.g., made from one or more of nylon, polyester, polyethylene, and polypropylene plastics) to form diapers.

**[0261]** More particularly, in one non-limiting process, above a conveyor belt multiple pressurized nozzles spray superabsorbent polymer particles (such as about 400 micron size or larger), fibrous material, and/or a combination of these onto the conveyor belt at designated spaces/intervals. The conveyor belt is perforated and under vacuum from below, so that the sprayed on materials are pulled toward the belt surface to form a flat pad. In various embodiments, fibrous material is applied first on the belt, followed by a mixture of fibrous material and the superabsorbent polymer particles, followed by fibrous material, so that the superabsorbent polymer is concentrated in the middle of the pad. A leveling roller may be used toward the end of the belt path to yield pads of uniform thickness. Each pad thereafter may be further processed, such as to cut it to a proper shape for the diaper, or the pad may be in the form of a long roll sufficient for multiple diapers. Thereafter, the pad is sandwiched between a top sheet and a bottom sheet of fabric (one generally being liquid pervious, the other liquid impervious), such as on a conveyor belt, and these are attached together such as by gluing, heating or ultrasonic welding, and cut into diaper-sized units (if not previously so cut). Additional features may be provided, such as elastic components, strips of tape, etc., for fit and ease of wearing by a person. FIG. 6A depicts aspects of diaper manufacture and FIG. B depicts aspects of diaper manufacture and diaper wear. FIG. 7 provides a general, non-limiting depiction of the overall processes leading from biomass to a polymerized acrylic acid-based polymer such as may be used in diapers in accordance with embodiments of the present invention.

**[0262]** The ratio of the fibrous material to polymer particles is known to effect performance characteristics. In some embodiments, this ratio is between 75:25 and 90:10 (see U.S. Pat. No. 4,685,915, incorporated by reference for its teachings of diaper manufacture). Other disposable absorbent articles may be constructed in a similar fashion, such as for adult incontinence, feminine hygiene (sanitary napkins), tampons, etc. (see, for example, U.S. Pat. Nos. 5,009,653, 5,558,656, and 5,827,255 incorporated by reference for their teachings of sanitary napkin manufacture).

**[0263]** Low molecular-weight polyacrylic acid has uses for water treatment, flocculants, and thickeners for various applications including cosmetics and pharmaceutical preparations. For these applications, the polymer may be uncrosslinked or lightly crosslinked, depending on the specific application. The molecular weights are typically from about 200 to about 1,000,000 g/mol. Preparation of these low molecular-weight polyacrylic acid polymers is described in U.S. Pat. Nos. 3,904,685; 4,301,266; 2,798,053; and 5,093,

472, each of which is incorporated by reference for its teachings relating to methods to produce these polymers.

**[0264]** Acrylic acid may be co-polymerized with one or more other monomers selected from acrylamide, 2-acrylamido-2-methylpropanesulfonic acid, N,N-dimethylacrylamide, N-isopropylacrylamide, methacrylic acid, and methacrylamide, to name a few. The relative reactivities of the monomers affect the microstructure and thus the physical properties of the polymer. Co-monomers may be derived from 3-HP, or otherwise provided, to produce co-polymers. *Ulmann's Encyclopedia of Industrial Chemistry, Polyacrylamides and Poly(Acrylic Acids)*, WileyVCH Verlag GmbH, Weinham (2005), is incorporated by reference herein for its teachings of polymer and co-polymer processing.

**[0265]** Acrylic acid can in principle be copolymerized with almost any free-radically polymerizable monomers including styrene, butadiene, acrylonitrile, acrylic esters, maleic acid, maleic anhydride, vinyl chloride, acrylamide, itaconic acid, and so on. End-use applications typically dictate the co-polymer composition, which influences properties. Acrylic acid also may have a number of optional substitutions on it, and after such substitutions be used as a monomer for polymerization, or co-polymerization reactions. As a general rule, acrylic acid (or one of its co-polymerization monomers) may be substituted by any substituent that does not interfere with the polymerization process, such as alkyl, alkoxy, aryl, heteroaryl, benzyl, vinyl, allyl, hydroxy, epoxy, amide, ethers, esters, ketones, maleimides, succinimides, sulfoxides, glycidyl and silyl (see U.S. Pat. No. 7,678,869, incorporated by reference above, for further discussion). The following paragraphs provide a few non-limiting examples of copolymerization applications.

**[0266]** Paints that comprise polymers and copolymers of acrylic acid and its esters are in wide use as industrial and consumer products. Aspects of the technology for making such paints can be found in U.S. Pat. Nos. 3,687,885 and 3,891,591, incorporated by reference for its teachings of such paint manufacture. Generally, acrylic acid and its esters may form homopolymers or copolymers among themselves or with other monomers, such as amides, methacrylates, acrylonitrile, vinyl, styrene and butadiene. A desired mixture of homopolymers and/or copolymers, referred to in the paint industry as 'vehicle' (or 'binder') are added to an aqueous solution and agitated sufficiently to form an aqueous dispersion that includes sub-micrometer sized polymer particles. The paint cures by coalescence of these 'vehicle' particles as the water and any other solvent evaporate. Other additives to the aqueous dispersion may include pigment, filler (e.g., calcium carbonate, aluminum silicate), solvent (e.g., acetone, benzol, alcohols, etc., although these are not found in certain no VOC paints), thickener, and additional additives depending on the conditions, applications, intended surfaces, etc. In many paints, the weight percent of the vehicle portion may range from about nine to about 26 percent, but for other paints the weight percent may vary beyond this range.

**[0267]** Acrylic-based polymers are used for many coatings in addition to paints. For example, for paper coating latexes, acrylic acid is used from 0.1-5.0%, along with styrene and butadiene, to enhance binding to the paper and modify rheology, freeze-thaw stability and shear stability. In this context, U.S. Pat. Nos. 3,875,101 and 3,872,037 are incorporated by reference for their teachings regarding such latexes. Acrylate-based polymers also are used in many inks, particularly UV curable printing inks. For water treatment, acrylamide

and/or hydroxy ethyl acrylate are commonly co-polymerized with acrylic acid to produce low molecular-weight linear polymers. In this context, U.S. Pat. Nos. 4,431,547 and 4,029,577 are incorporated by reference for their teachings of such polymers. Co-polymers of acrylic acid with maleic acid or itaconic acid are also produced for water-treatment applications, as described in U.S. Pat. No. 5,135,677, incorporated by reference for that teaching. Sodium acrylate (the sodium salt of glacial acrylic acid) can be co-polymerized with acrylamide (which may be derived from acrylic acid via amidation chemistry) to make an anionic co-polymer that is used as a flocculant in water treatment.

**[0268]** For thickening agents, a variety of co-monomers can be used, such as described in U.S. Pat. Nos. 4,268,641 and 3,915,921, incorporated by reference for description of these co-monomers. U.S. Pat. No. 5,135,677 describes a number of co-monomers that can be used with acrylic acid to produce water-soluble polymers, and is incorporated by reference for such description.

**[0269]** Also as noted, some conversions to downstream products may be made enzymatically. For example, 3-HP may be converted to 3-HP-CoA, which then may be converted into polymerized 3-HP with an enzyme having polyhydroxy-acid synthase activity (EC 2.3.1.-). Also, 1,3-propanediol can be made using polypeptides having oxidoreductase activity or reductase activity (e.g., enzymes in the EC 1.1.1.-class of enzymes). Alternatively, when creating 1,3-propanediol from 3HP, a combination of (1) a polypeptide having aldehyde dehydrogenase activity (e.g., an enzyme from the 1.1.1.34 class) and (2) a polypeptide having alcohol dehydrogenase activity (e.g., an enzyme from the 1.1.1.32 class) can be used. Polypeptides having lipase activity may be used to form esters. Enzymatic reactions such as these may be conducted in vitro, such as using cell-free extracts, or in vivo.

**[0270]** Thus, various embodiments of the present invention, such as methods of making a chemical, include conversion steps to any such noted downstream products of microbially produced 3-HP, including but not limited to those chemicals described herein and in the incorporated references (the latter for jurisdictions allowing this). For example, one embodiment is making 3-HP molecules by the teachings herein and further converting the 3-HP molecules to polymerized-3-HP (poly-3-HP) or acrylic acid, and such as from acrylic acid then producing from the 3-HP molecules any one of polyacrylic acid (polymerized acrylic acid, in various forms), methyl acrylate, acrylamide, acrylonitrile, propiolactone, ethyl 3-HP, malonic acid, 1,3-propanediol, ethyl acrylate, n-butyl acrylate, hydroxypropyl acrylate, hydroxyethyl acrylate, isobutyl acrylate, 2-ethylhexyl acrylate, and acrylic acid or an acrylic acid ester to which an alkyl or aryl addition is made, and/or to which halogens, aromatic amines or amides, and aromatic hydrocarbons are added.

**[0271]** Also as noted, some conversions to downstream products may be made enzymatically. For example, 3-HP may be converted to 3-HP-CoA, which then may be converted into polymerized 3-HP with an enzyme having polyhydroxy-acid synthase activity (EC 2.3.1.-). Also, 1,3-propanediol can be made using polypeptides having oxidoreductase activity or reductase activity (e.g., enzymes in the EC 1.1.1.-class of enzymes). Alternatively, when creating 1,3-propanediol from 3HP, a combination of (1) a polypeptide having aldehyde dehydrogenase activity (e.g., an enzyme from the 1.1.1.34 class) and (2) a polypeptide having alcohol dehydrogenase activity (e.g., an enzyme from the 1.1.1.32 class) can be used.

Polypeptides having lipase activity may be used to form esters. Enzymatic reactions such as these may be conducted in vitro, such as using cell-free extracts, or in vivo.

**[0272]** Thus, various embodiments of the present invention, such as methods of making a chemical, include conversion steps to any such noted downstream products of microbially produced 3-HP, including but not limited to those chemicals described herein and in the incorporated references (the latter for jurisdictions allowing this). For example, one embodiment is making 3-HP molecules by the teachings herein and further converting the 3-HP molecules to polymerized-3-HP (poly-3-HP) or acrylic acid, and such as from acrylic acid then producing from the 3-HP molecules any one of polyacrylic acid (polymerized acrylic acid, in various forms), methyl acrylate, acrylamide, acrylonitrile, propiolactone, ethyl 3-HP, malonic acid, 1,3-propanediol, ethyl acrylate, n-butyl acrylate, hydroxypropyl acrylate, hydroxyethyl acrylate, isobutyl acrylate, 2-ethylhexyl acrylate, and acrylic acid or an acrylic acid ester to which an alkyl or aryl addition is made, and/or to which halogens, aromatic amines or amides, and aromatic hydrocarbons are added.

**[0273]** Reactions that form downstream compounds such as acrylates or acrylamides can be conducted in conjunction with use of suitable stabilizing agents or inhibiting agents reducing likelihood of polymer formation. See, for example, U.S. Patent Publication No. 2007/0219390 A1. Stabilizing agents and/or inhibiting agents include, but are not limited to, e.g., phenolic compounds (e.g., dimethoxyphenol (DMP) or alkylated phenolic compounds such as di-tert-butyl phenol), quinones (e.g., t-butyl hydroquinone or the monomethyl ether of hydroquinone (MEHQ)), and/or metallic copper or copper salts (e.g., copper sulfate, copper chloride, or copper acetate). Inhibitors and/or stabilizers can be used individually or in combinations as will be known by those of skill in the art. Also, in various embodiments, the one or more downstream compounds is/are recovered at a molar yield of up to about 100 percent, or a molar yield in the range from about 70 percent to about 90 percent, or a molar yield in the range from about 80 percent to about 100 percent, or a molar yield in the range from about 90 percent to about 100 percent. Such yields may be the result of single-pass (batch or continuous) or iterative separation and purification steps in a particular process.

**[0274]** Acrylic acid and other downstream products are useful as commodities in manufacturing, such as in the manufacture of consumer goods, including diapers, textiles, carpets, paint, adhesives, and acrylic glass.

**[0275]** In some embodiments, the invention contemplates a culture system comprising: a) a population of a genetically modified microorganism as described herein; and b) a media comprising nutrients for the population. In some such embodiments the media additionally comprises at least 1 gram/liter of 3-HP.

**[0276]** The teachings and results of the Examples that follow are hereby incorporated into this section for all applicable purposes.

**[0277]** Also, and more generally, in accordance with disclosures, discussions, examples and embodiments herein, there may be employed conventional molecular biology, cellular biology, microbiology, and recombinant DNA techniques within the skill of the art. Such techniques are explained fully in the literature. (See, e.g., Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Third Edition 2001 (volumes 1-3), Cold Spring Harbor Laboratory

Press, Cold Spring Harbor, N.Y.; *Animal Cell Culture*, R. I. Freshney, ed., 1986). These published resources are incorporated by reference herein for their respective teachings of standard laboratory methods found therein. Further, all patents, patent applications, patent publications, and other publications referenced herein (collectively, “published resource(s)”) are hereby incorporated by reference in this application. Such incorporation, at a minimum, is for the specific teaching and/or other purpose that may be noted when citing the reference herein. If a specific teaching and/or other purpose is not so noted, then the published resource is specifically incorporated for the teaching(s) indicated by one or more of the title, abstract, and/or summary of the reference. If no such specifically identified teaching and/or other purpose may be so relevant, then the published resource is incorporated in order to more fully describe the state of the art to which the present invention pertains, and/or to provide such teachings as are generally known to those skilled in the art, as may be applicable. However, it is specifically stated that a citation of a published resource herein shall not be construed as an admission that such is prior art to the present invention. Also, in the event that one or more of the incorporated published resources differs from or contradicts this application, including but not limited to defined terms, term usage, described techniques, or the like, this application controls.

**[0278]** While various embodiments of the present invention have been shown and described herein, it is emphasized that such embodiments are provided by way of example only. Numerous variations, changes and substitutions may be made without departing from the invention herein in its various embodiments. Specifically, and for whatever reason, for any grouping of compounds, nucleic acid sequences, polypeptides including specific proteins including functional enzymes, metabolic pathway enzymes or intermediates, elements, or other compositions, or concentrations stated or otherwise presented herein in a list, table, or other grouping (such as metabolic pathway enzymes shown in a figure), unless clearly stated otherwise, it is intended that each such grouping provides the basis for and serves to identify various subset embodiments, the subset embodiments in their broadest scope comprising every subset of such grouping by exclusion of one or more members (or subsets) of the respective stated grouping. Moreover, when any range is described herein, unless clearly stated otherwise, that range includes all values therein and all sub-ranges therein. Accordingly, it is intended that the invention be limited only by the spirit and scope of appended claims, and of later claims, and of either such claims as they may be amended during prosecution of this or a later application claiming priority hereto.

## EXAMPLES

### Example 1

Preparing a Genetically Modified *E. Coli* Host Cell Comprising an Oxaloacetate Alpha-Decarboxylase (oad-2) in Combination with Other Genetic Modifications to Increase 3-HP Production Relative to a Control *E. Coli* Cell (Prophetic)

**[0279]** Referring to FIG. 4 and Tables 1 and 2, genetic modifications are made to introduce a polypeptide that encodes SEQ ID NO:001 or SEQ ID NO:036, or its functional equivalents, and also to introduce a vector that encodes mmsB (SEQ ID NO:002). Vectors comprising galP and a native or

mutated Ppc also may be introduced by methods known to those skilled in the art (see, e.g., Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Third Edition 2001 (volumes 1-3), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., “Sambrook and Russell, 2001”), additionally recognizing that mutations may be made by a method using the XL1-Red mutator strain, using appropriate materials following a manufacturer’s instructions (Stratagene QuikChange Mutagenesis Kit, Stratagene, La Jolla, Calif. USA) and selected for or screened under standard protocols.

**[0280]** Also, genetic modifications are made to reduce or eliminate the enzymatic activities of the *E. coli* genes listed in Table 2. These genetic modifications are achieved by using the RED/ET homologous recombination method with kits supplied by Gene Bridges (Gene Bridges GmbH, Dresden, Germany, www.genebridges.com) according to manufacturer’s instructions.

**[0281]** The so-genetically modified microorganism is evaluated and found to exhibit higher productivity of 3-HP compared with a control *E. coli* lacking said genetic modifications. Productivity is measured by standard metrics, such as volumetric productivity (grams of 3-HP/hour) under similar culture conditions.

#### Example 2

##### General Example of Genetic Modification to a Host Cell (Prophetic and Non-Specific)

**[0282]** This example is meant to describe a non-limiting approach to genetic modification of a selected microorganism to introduce a nucleic acid sequence of interest. Alternatives and variations are provided within this general example. The methods of this Example are conducted to achieve a combination of desired genetic modifications in a selected microorganism species, such as a combination of genetic modifications selected from those shown in FIG. 4, and their equivalents in species other than *E. coli*.

**[0283]** A gene or other nucleic acid sequence segment of interest is identified in a particular species (such as *E. coli* as described above) and a nucleic acid sequence comprising that gene or segment is obtained. For clarity below the use of the term “segment of interest” below is meant to include both a gene and any other nucleic acid sequence segment of interest. One example of a method used to obtain a segment of interest is to acquire a culture of a microorganism, where that microorganism’s genome includes the gene or nucleic acid sequence segment of interest.

**[0284]** Based on the nucleic acid sequences at the ends of or adjacent the ends of the segment of interest, 5' and 3' nucleic acid primers are prepared. Each primer is designed to have a sufficient overlap section that hybridizes with such ends or adjacent regions. Such primers may include enzyme recognition sites for restriction digest or transposase insertion that could be used for subsequent vector incorporation or genomic insertion. These sites are typically designed to be outward of the hybridizing overlap sections. Numerous contract services are known that prepare primer sequences to order (e.g., Integrated DNA Technologies, Coralville, Iowa USA).

**[0285]** Once primers are designed and prepared, polymerase chain reaction (PCR) is conducted to specifically amplify the desired segment of interest. This method results in multiple copies of the region of interest separated from the microorganism’s genome. The microorganism’s DNA, the primers, and a thermophilic polymerase are combined in a

buffer solution with potassium and divalent cations (e.g., Mg or Mn) and with sufficient quantities of deoxynucleoside triphosphate molecules. This mixture is exposed to a standard regimen of temperature increases and decreases. However, temperatures, components, concentrations, and cycle times may vary according to the reaction according to length of the sequence to be copied, annealing temperature approximations and other factors known or readily learned through routine experimentation by one skilled in the art.

**[0286]** In an alternative embodiment the segment of interest may be synthesized, such as by a commercial vendor, and prepared via PCR per above, rather than obtaining from a microorganism or other natural source of DNA.

**[0287]** The nucleic acid sequences then are purified and separated, such as on an agarose gel via electrophoresis. Optionally, once the region is purified it can be validated by standard DNA sequencing methodology and may be introduced into a vector. Any of a number of vectors may be used, which generally comprise markers known to those skilled in the art, and standard methodologies are routinely employed for such introduction. Commonly used vector systems are pSMART (Lucigen, Middleton, Wis.), pET *E. coli* EXPRESSION SYSTEM (Stratagene, La Jolla, Calif.), pSC-B StrataClone Vector (Stratagene, La Jolla, Calif.), pRANGER-BTB vectors (Lucigen, Middleton, Wis.), and TOPO vector (Invitrogen Corp, Carlsbad, Calif., USA). Similarly, the vector then is introduced into any of a number of host cells. Commonly used host cells are *E. coli* 10G (Lucigen, Middleton, Wis.), *E. coli* 10GF' (Lucigen, Middleton, Wis.), StrataClone Competent cells (Stratagene, La Jolla, Calif.), *E. coli* BL21, *E. coli* BW25113, and *E. coli* K12 MG1655. Some of these vectors possess promoters, such as inducible promoters, adjacent the region into which the sequence of interest is inserted (such as into a multiple cloning site), while other vectors, such as pSMART vectors (Lucigen, Middleton, Wis.), are provided without promoters and with dephosphorylated blunt ends. The culturing of such plasmid-laden cells permits plasmid replication and thus replication of the segment of interest, which often corresponds to expression of the segment of interest.

**[0288]** Various vector systems comprise a selectable marker, such as an expressible gene encoding a protein needed for growth or survival under defined conditions. Common selectable markers contained on backbone vector sequences include genes that encode for one or more proteins required for antibiotic resistance as well as genes required to complement auxotrophic deficiencies or supply critical nutrients not present or available in a particular culture media. Vectors also comprise a replication system suitable for a host cell of interest.

**[0289]** The plasmids containing the segment of interest can then be isolated by routine methods and are available for introduction into other microorganism host cells of interest. Various methods of introduction are known in the art and can include vector introduction or genomic integration. In various alternative embodiments the DNA segment of interest may be separated from other plasmid DNA if the former will be introduced into a host cell of interest by means other than such plasmid.

**[0290]** While steps of the above general prophetic example involve use of plasmids, other vectors known in the art may be used instead. These include cosmids, viruses (e.g., bacteriophage, animal viruses, plant viruses), and artificial chro-

mosomes (e.g., yeast artificial chromosomes (YAC) and bacteria artificial chromosomes (BAC)).

**[0291]** Host cells into which the segment of interest is introduced may be evaluated for performance as to a particular enzymatic step, and/or tolerance or bio-production of a chemical compound of interest. Selections of better performing genetically modified host cells may be made, selecting for overall performance, tolerance, or production or accumulation of the chemical of interest.

**[0292]** It is noted that this procedure may incorporate a nucleic acid sequence for a single gene (or other nucleic acid sequence segment of interest), or multiple genes (under control of separate promoters or a single promoter), and the procedure may be repeated to create the desired heterologous nucleic acid sequences in expression vectors, which are then supplied to a selected microorganism so as to have, for example, a desired complement of enzymatic conversion step functionality for any of the herein-disclosed metabolic pathways. However, it is noted that although many approaches rely on expression via transcription of all or part of the sequence of interest, and then translation of the transcribed mRNA to yield a polypeptide such as an enzyme, certain sequences of interest may exert an effect by means other than such expression.

**[0293]** The specific laboratory methods used for the above approaches are well-known in the art and may be found in various references known to those skilled in the art, such as Sambrook and Russell, *Molecular Cloning: A Laboratory Manual*, Third Edition 2001 (volumes 1-3), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y. (hereinafter, Sambrook and Russell, 2001).

**[0294]** As an alternative to the above, other genetic modifications may also be practiced, such as a deletion of a nucleic acid sequence of the host cell's genome. One non-limiting method to achieve this is by use of Red/ET recombination, known to those of ordinary skill in the art and described in U.S. Pat. Nos. 6,355,412 and 6,509,156, issued to Stewart et al. and incorporated by reference herein for its teachings of this method. Material and kits for such method are available from Gene Bridges (Gene Bridges GmbH, Dresden, Germany, www.genebridges.com), and the method may proceed by following the manufacturer's instructions. Targeted deletion of genomic DNA may be practiced to alter a host cell's metabolism so as to reduce or eliminate production of undesired metabolic products, such as gene deletions for particular genes shown in FIG. 4. This may be used in combination with other genetic modifications such as described above in this general example.

**[0295]** The above-described approaches and compositions may be combined with other approaches and compositions that are directed to providing, or to improving, a desired property (e.g. productivity improvements in a microorganism). Various bio-production pathways, and methods of incorporating them into a microorganism, are known in the art and also are described in other patent applications having a common co-inventor with the present patent application. Any of such may be combined with any of the above-described approaches and combinations.

### Example 3

#### Mutational Development of Selected Polynucleotides (Prophetic)

**[0296]** A selected gene sequence is subjected to a mutation development protocol, starting by constructing a mutant

library of a native or previously evolved and/or codon-optimized polynucleotide by use of an error-inducing PCR site-directed mutagenesis method.

**[0297]** A polynucleotide exhibiting enzymatic activity of the selected gene (which may be any disclosed herein, e.g., a decarboxylase or mmsB) will be cloned into an appropriate expression system for *E. coli*. This sequence may be codon optimized. Cloning of a codon-optimized polynucleotide and its adequate expression of the will be accomplished via gene synthesis supplied from a commercial supplier using standard techniques. The gene will be synthesized with an eight amino acid C-terminal tag to enable affinity based protein purification. Once obtained using standard methodology, the gene will be cloned into an expression system using standard techniques.

**[0298]** The plasmid containing the above-described polynucleotide will be mutated by standard methods resulting in a large library of mutants ( $>10^6$ ). The mutant sequences will be excised from these plasmids and again cloned into an expression vector, generating a final library of greater than  $10^6$  clones for subsequent screening. These numbers ensure a greater than 99% probability that the library will contain a mutation in every amino acid encoded by sequence. It is acknowledged that each method of creating a mutational library has its own biases, including transformation into mutator strains of *E. coli*, error prone PCR, and in addition more site directed mutagenesis. In some embodiments, various methods may be considered and possibly several explored in parallel.

**[0299]** One such method is the use of the XL1-Red mutator strain, which is deficient in several repair mechanisms necessary for accurate DNA replication and generates mutations in plasmids at a rate 5,000 times that of the wild-type mutation rate, may be employed using appropriate materials following a manufacturer's instructions (See Stratagene QuikChange Mutagenesis Kit, Stratagene, La Jolla, Calif. USA). This technique or other techniques known to those skilled in the art, may be employed and then a population of such mutants, e.g., in a library, is evaluated, such as by a screening or selection method, to identify clones having a suitable or favorable mutation.

**[0300]** With the successful construction of a mutant library, it will be possible to screen this library for increased activity. The screening process will be designed to screen the entire library of greater than  $10^6$  mutants. This is done by screening methods suited to the particular enzymatic reaction.

### Example 4

#### Cloning of Oxaloacetate Alpha-Decarboxylase Candidate Enzymes

**[0301]** Candidate oxaloacetate alpha-decarboxylase genes were synthesized using codons optimized for expression in *E. coli* and sub-cloned into expression vectors to provide protein for oxaloacetate alpha-decarboxylase assays. For gene synthesis, the protein sequences for pyruvate decarboxylase (pdc) from *Zymomonas mobilis* (SEQ ID NO:035), 2-oxoglutarate decarboxylase (oad) from *Leuconostoc mesenteroides* (SEQ ID NO:036), and alpha-ketoglutarate decarboxylase (kgd) from *Mycobacterium tuberculosis* (SEQ ID NO:037) were used to create codon optimized genes for expression in *E. coli* by the service provided by DNA 2.0 (Menlo Park, Calif. USA). Additionally, the protein coding regions of the codon optimized PDC and OAD genes were augmented with

six histidines and five histidines, respectively. These additional amino acids provide a C-terminal histidine-tag for protein purification via immobilized metal-affinity chromatography. These plasmids were designated pJ201:pdC (SEQ ID NO:038), pJ251:oad (SEQ ID NO:039), and pJ206:kgd (SEQ ID NO:040), respectively.

**[0302]** The pdC and oad genes were individually sub-cloned into a pTrc-HisB expression vector obtained from Invitrogen (SEQ ID NO:041) (Carlsbad, Calif.). Sub-cloning for pdC gene was accomplished as follows: First, the expression vector was subjected to enzymatic restriction digestion with the restriction enzymes NcoI and NheI. Restriction enzymes were obtained from New England BioLabs (Ipswich, Mass. USA), and used according to manufacturer's instructions. The digestion mixtures were separated by agarose gel electrophoresis, and visualized under UV transillumination as described in the Common Methods Section. Agarose gel slices containing the DNA piece corresponding to the cut vector product were cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The gene inserts were amplified by polymerase chain reactions, cut with restriction enzymes, purified, and ligated to the cut vector. For the pdC gene containing insert, a polymerase chain reaction was performed with the forward primer being GGGATATCAT GAGCTATAACC GTTGG (SEQ ID NO:042), and the reverse primer being GAAATAGTTC TCTAGAGAAG CTTC (SEQ ID NO:043) and the pJ201:pdC plasmid was used as template. These primers provide a BspHI site at the start codon and an XbaI site after the stop codon. This PCR product was prepared for restriction digest using a PCR purification kit from Qiagen Corporation (Valencia, Calif., USA) using the manufacturer's instructions. The PCR product was prepared by restriction digest with BspHI and XbaI obtained from New England BioLabs (Ipswich, Mass. USA), and used according to manufacturer's instructions. After digestion, mixtures were separated by agarose gel electrophoresis, and visualized under UV transillumination as described in the Common Methods Section. An agarose gel slice containing the DNA piece corresponding to the amplified pdC gene fusion was cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The recovered product and the prepared vector backbone were ligated together with T4 DNA ligase obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. Ligation mixtures were transformed into chemically competent 100 cells (Lucigen, Middleton, Wis.) according to the manufacturer's instructions. Colonies possibly containing the new plasmid were cultured, and their DNA was isolated using a standard miniprep protocol and components from Qiagen (Valencia, Calif. USA) according to the manufacturer's instruction. Isolated plasmids were checked by restriction digests and confirmed by sequencing. The sequenced-verified isolated plasmids produced with this procedure were designated pTrc:pdC-his (SEQ ID NO:044).

**[0303]** Sub-cloning for oad gene was accomplished as follows: First, the expression vector was subjected to enzymatic restriction digestion with the restriction enzymes NcoI and NheI. Restriction enzymes were obtained from New England BioLabs (Ipswich, Mass. USA), and used according to manufacturer's instructions. The digestion mixtures were separated by agarose gel electrophoresis, and visualized under UV

transillumination as described in the Common Methods Section. Agarose gel slices containing the DNA piece corresponding to the cut vector product were cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The gene inserts were amplified by polymerase chain reactions, cut with restriction enzymes, purified, and ligated to the cut vector. For the oad gene containing insert, a polymerase chain reaction was performed with the forward primer being GGAGAATTACCATGGCGGATACCCTG (SEQ ID NO:045), and the reverse primer being GGGAACTAG ACTAATGATG ATGGTGG (SEQ ID NO:046), and the pJ251:oad plasmid was used as template. These primers provide an NcoI site and the start codon and an XbaI site after the stop codon. This PCR product was prepared for restriction digest purified using a PCR purification kit from Qiagen Corporation (Valencia, Calif., USA) using the manufacturer's instructions. The PCR product was prepared by restriction digest with NcoI and XbaI obtained from New England BioLabs (Ipswich, Mass. USA), and used according to manufacturer's instructions. After digestion, mixtures were separated by agarose gel electrophoresis, and visualized under UV transillumination as described in the Common Methods Section. An agarose gel slice containing the DNA piece corresponding to the amplified pdC gene fusion was cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The recovered product and the prepared vector backbone were ligated together with T4 DNA ligase obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. Ligations were transformed into chemically competent 100 cells from Lucigen according to the manufacturer's instructions. Colonies possibly containing the new plasmid were cultured, and their DNA was isolated using a standard miniprep protocol and components from Qiagen (Valencia, Calif. USA) according to the manufacturer's instruction. Isolated plasmids were checked by restriction digests and confirmed by sequencing. The sequenced-verified isolated plasmids produced with this procedure were designated pTrc:oad-his (SEQ ID NO:047).

**[0304]** The kgd gene was sub-cloned into a pKK223:c-term-His expression vector. Sub-cloning for kgd gene was accomplished as follows: First, the pKK223:c-term-His expression vector (SEQ ID NO:050) was subjected to enzymatic restriction digestion with the restriction enzymes NcoI and HindIII. Restriction enzymes were obtained from New England BioLabs (Ipswich, Mass. USA), and used according to manufacturer's instructions. The digestion mixtures were separated by agarose gel electrophoresis, and visualized under UV transillumination as described in the Common Methods Section. Agarose gel slices containing the DNA piece corresponding to the cut vector product were cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The gene inserts were amplified by polymerase chain reactions, cut with restriction enzymes, purified, and ligated to the cut vector. For the kgd gene containing insert, a polymerase chain reaction was performed with the forward primer being AGGTTCCCA TGGTGACTCA GGACCCG (SEQ ID NO:048), and the reverse primer being GTAAGCTTAG TGGTGATGGT GATGACCGAA CGCTTCGTCC (SEQ ID NO:049), and the pJ206:kgd plasmid was used as template. These primers provide an NcoI site and the start

codon, and penta-histidine coding sequence for affinity purification of protein followed by a stop codon as well as a HindIII site. This PCR product was prepared for restriction digest purified using a PCR purification kit from Qiagen Corporation (Valencia, Calif., USA) using the manufacturer's instructions. The PCR product was prepared by restriction digest with NcoI and HindIII obtained from New England BioLabs (Ipswich, Mass. USA), and used according to manufacturer's instructions. After digestion, mixtures were separated by agarose gel electrophoresis, and visualized under UV transillumination as described in the Common Methods Section. An agarose gel slice containing the DNA piece corresponding to the amplified *pdc* gene fusion was cut from the gel and the DNA recovered with a standard gel extraction protocol and components from Qiagen according to manufacturer's instructions. The recovered product and the prepared vector backbone were ligated together with T4 DNA ligase obtained from New England BioLabs (Ipswich, Mass. USA) according to manufacturer's instructions. Ligations were transformed into chemically competent 100 cells from Lucigen according to the manufacturer's instructions. Colonies possibly containing the new plasmid were cultured, and their DNA was isolated using a standard miniprep protocol and components from Qiagen (Valencia, Calif. USA) according to the manufacturer's instruction. Isolated plasmids were checked by restriction digests and confirmed by sequencing. The sequenced-verified isolated plasmids produced with this procedure were designated pKK223:kgd-his (SEQ ID NO:053).

#### Example 5

##### Preparation and Evaluation of Over-Expressed Oxaloacetate Alpha-Decarboxylase Candidate Enzymes and Selected Mutant Enzymes

**[0305]** In order to evaluate candidate enzymes for native oxaloacetate alpha-decarboxylase function, each of the protein was purified. Plasmids containing each of the genes were individually transformed into electro-competent BW25113 cells. For each enzyme that was tested, starter cultures from single colonies were grown overnight at 37° C. with 250 rpm agitation in 20 mL of LB media with 200 ug/mL ampicillin selection. The next morning, these cultures were used to inoculate a 1 L of TB media with 200 ug/mL ampicillin selection. Cultures were grown at 37° C. with 250 rpm agitation. Protein induction was initiated when the optical density of the culture measured at 600 nm reached between 0.5 and 0.7—(about 4 to 6 hr) with addition of IPTG to a final concentration of 1 mM. The cultures allowed to grow overnight at 30° C. The cells were harvested by centrifugation (6000×g for 15 minutes) and kept frozen at -70° C. until purification.

**[0306]** Purification was performed as follows: Cells were thawed on ice and then resuspended in 40 mL of Buffer A (25 mM Trizma base pH 8.0, 500 mM NaCl, and 1 mM Imidazole) supplemented with lysozyme (Sigma-Aldrich, St. Louis, Mo.), Dnase I (Sigma-Aldrich, St. Louis, Mo.), and Complete EDTA-free protease inhibitor cocktail (EMD, Gibbstown, N.J.). Resuspend cells were incubated on ice for at least 5 minutes or until the culture showed signs of lysis. To completely lyse the cells, the suspension was sonicated (85% power on a Branson probe tip sonicator) with five 10 second burst intervened by resting on ice for at least a minute. The

cell lysate was diluted to 50 mL with Buffer A and clarified by centrifugation at 10000×g for 15 minutes.

**[0307]** Proteins were purified from the lysate as follows: The clarified lysate was subjected to affinity purification with a nickel-charged fast flow column (Pharmacia). Clarified cell lysates was loaded on to the column at 1 mL per minute. The column was then washed with at least 50 column volumes of Buffer A until the flow through had an absorption at 260 nm equal to that of Buffer A. The protein was eluted with Elution Buffer (25 mM Trizma base pH 8.0, 100 mM NaCl, and 300 mM Imidazole) at 1 mL per minute and collected in 1.0 mL fractions. Fractions containing protein were identified using the BioRad Total Protein Assay method. Fractions showing protein were combined and concentrated by centrifugation using a Amicon ultra concentration unit (10 kDa cutoff) (Millipore, Milford Mass.). Concentrated proteins were further purified to remove imidazole using a pre-equilibrated (25 mM Trizma base pH 8.0, 100 mM NaCl) PD Miditrap G25 column (GE Lifesciences, Piscataway, N.J.) according the manufacturer's instructions. Proteins were concentrated a final time, glycerol was added to about 10% volume, protein aliquots were snap frozen on dry ice, and the proteins were stored at -70° C. until use. Final protein concentrations were measured using BioRad Total Protein Assay (BioRad, Hercules, Calif.)

**[0308]** The purification of each candidate protein was confirmed by SDS-PAGE and by SDS-PAGE followed by visualization via western blot analysis. Protein samples were prepared for SDS-PAGE by dilution in 100 uL of SDS sample buffer (Tris-Cl pH6.8, SDS, glycerol, β-mercaptoethanol, Bromophenol blue), boiled for 5 minutes and spun at 17,000G for 5 minutes. A portion of the sample was loaded on a 10% pre-cast SDS-PAGE gel (BioRad Ready Gel Tris-HCl Gel-161-1101). Electrophoresis was carried out using a BioRad Mini-Protean II system according to manufacturer's instructions. SDS gels were either stained with Coomassie Stain or transferred to nitrocellulose membrane using the same BioRad Mini-Protean II wet transfer system according to manufacturer's specifications for western blot visualization. Membranes were blocked for 1 hour at room temperature using PBST (NaCl, KCl, Na<sub>2</sub>HPO<sub>4</sub>, KH<sub>2</sub>PO<sub>4</sub>, Tween 20)+5% w/v nonfat dry milk Blots were then probed with a rabbit polyclonal anti-6× HIS-HRP antibody (AbCam Ab1187, 1:5000 dilution) in PBST+5% w/v nonfat dry milk for 1 hour at room temperature, washed 4 times in PBST for 5 minutes, and followed by developing with TMB substrate (Promega TMB Stabilized Substrate for HRP, cat #W4121). Protein expression was assessed by the presence or absence of bands at the expected molecular weight for each proteins of interest. The presence of each enzyme was confirmed in this manner (FIG. 8).

#### Example 6

##### Evaluation of Native Oxaloacetate Alpha-Decarboxylase Function Via a Spectrophotometric Assay to Determine the Specific Activities of the Native Candidate Enzymes

**[0309]** A coupled assay was developed in order to biochemically assess the oxaloacetate alpha-decarboxylase activity of the candidate enzymes. The coupled assay used exogenously-added purified *E. coli* YdfG protein, which converts malonate semialdehyde to 3-HP with the concomitant conversion of NADPH to NADP+ (FIG. 5). As the reduced form, NADPH, has a strong absorbance peak at 340 nm and



the oxidized form does not, it is possible to monitor the progress of these coupled reactions via a spectrophotometric assay. 3-HP is also detectable by standard analytical methods, for example HPLC and GS-MS.

**[0310]** Assays were performed as 200 microliter reactions in a 96-well plate format using a Molecular Dynamics SpectraMax 384 microplate reader with SoftmaxPro software (Molecular Dynamics, Sunnyvale, Calif.) to quantitate the rate of change in the 340 nm absorbance. All assays were conducted at 37° C., and the instrument was allowed to mix the plate for 1 second prior to each measurements. The progress of each reaction was monitored for 20 minutes during which measurements were made every 20 seconds. Reaction conditions consisted of 50 mM PIPES pH 6.8 (unless otherwise stated), 2 mM thymidine pyrophosphate, 10 mM dithiothreitol, 4 mM magnesium chloride, 1 mM NADPH (EMD Bioscience), 0.075 mg/mL purified YdfG (Molecular Throughput), and 5 mM oxaloacetate (unless otherwise stated). All chemicals were obtained from Sigma-Aldrich (St. Louis, Mo.) unless otherwise stated. Reactions were initiated with the addition of 10 microliters of concentrated purified protein to each reaction well. In addition, a negative control was performed with each experimental set to control for the rate of background NADPH oxidation. Once the reaction time course was read and the slopes of each well were calculated, the specific activities over the negative control of each candidate protein were calculated. All values reported are the average specific activities measured in triplicate. The following table shows the native specific activities of PDC and OAD determined by this method at varying oxaloacetate concentrations.

Specific Activities as Measured by the Spectrophotometric Assay at pH 6.8				
[OAA] (mM)	average PDC Specific activity (units/mg)	standard deviation	Average OAD Specific activity (units/mg)	standard deviation
10	0.0076	0.0024	0.0144	0.0012.
5	0.003	0.0001	0.006	0.0003
2.5 m	0.0025	0.0003	0.003	0.0007
1	0.0015	0.0002	0.0017	0.0009
0.5	0.002	0.0002	0	0.0004
0.1	0.0004	0.0003	0.0014	0.001

**[0311]** Activity of KGD was not measured by the spectrophotometric assay. These reactions were subjected to GC-MS analysis to confirm that 3-HP was produced, and that the candidate oxaloacetate alpha-decarboxylase enzymes were able to convert oxaloacetate to malonate semialdehyde. These results are shown in the following table. 3-HP was produced by PDC, OAD, and KGD in the presence of 5 mM oxaloacetate. These results show that these enzymes possess low levels of native oxaloacetate alpha-decarboxylase function.

Production of 3HP from spectrophotometric assay samples		
Enzyme	Beginning [OAA]	Final [3HP] (g/L)
Pdc (pH 6.0)	5 mM	0.041 +/- 0.003
Oad2 (pH 6.8)	5 mM	0.004 +/- 0.001
Kgd (pH 6.8)	5 mM	0.004 +/- 0.001

### Example 7

#### Confirmation of Oxaloacetate Alpha-Decarboxylase Enzyme Activity—Using GC-MS—

**[0312]** Confirmation of 3-HP production as well as assessment of the pH dependence of each candidate enzyme was performed as described above, except that coupled reactions with YdfG were evaluated for 3-HP production by GC-MS. Reaction conditions consisted of 50 mM PIPES pH 6.8, 2 mM thymidine pyrophosphate, 10 mM dithiothreitol, 4 mM magnesium chloride, 1 mM NADPH (EMD Bioscience), 0.075 mg/mL purified YdfG (Molecular Throughput), and 5 mM oxaloacetate (unless otherwise stated). All chemicals were obtained from Sigma-Aldrich unless otherwise stated. Reactions were performed at 0.6 mL volumes in triplicate and were initiated with the addition of 600 micrograms of purified protein. In addition, a negative control lacking protein was also performed with each set. Reactions were allowed to proceed at 37° C. for at least 12 hours.

**[0313]** The following method is used for GC-MS analysis of 3-HP. Soluble monomeric 3-HP is quantified using GC-MS after a single extraction with ethyl acetate from a reaction sample. Once the 3-HP has been extracted into the ethyl acetate, the active hydrogens on the 3-HP are replaced with trimethylsilyl groups using N,O-Bis-(Trimethylsilyl)trifluoroacetamide to make the compound volatile for GC analysis. A standard curve of known 3-HP concentrations is prepared at the beginning of the run and a known quantity of ketohexanoic acid (1 g/L) is added to both the standards and the samples to act as an internal standard for Quantitation, with tropic acid as an additional internal standard. The 3-HP content of individual samples is then assayed by examining the ratio of the ketohexanoic acid ion (m/z=247) to the 3-HP ion (219) and compared to the standard curve. 3-HP is quantified using a 3HP standard curve at the beginning of the run and the data are analyzed using HP Chemstation. The GC-MS system consists of a Hewlett Packard model 5890 GC and Hewlett Packard model 5972 MS. The column is Supelco SPB-1 (60 m×0.32 mm×0.25 um film thickness). The capillary coating is a non-polar methylsilicone. The carrier gas is helium at a flow rate of 1 mL/min. The 3-HP as derivatized is separated from other components in the ethyl acetate extract using either of two similar temperature regimes. In a first temperature gradient regime, the column temperature starts with 40° C. for 1 minute, then is raised at a rate of 10° C./minute to 235° C., and then is raised at a rate of 50° C./minute to 300° C. In a second temperature regime, which was demonstrated to process samples more quickly, the column temperature starts with 70° C. which is held for 1 min, followed by a ramp-up of 10° C./minute to 235° C. which is followed by a ramp-up of 50° C./minute to 300° C. FIG. 9 shows the detected amount of each candidate enzyme at pH 5.5, pH 6.0, and pH 6.8. PDC showed optimal activity at pH 5.5 while OAD and KGD showed optimal activity at pH 6.8.

### Example 8

#### Variants with Increased Oxaloacetate Alpha-Decarboxylase Activity

**[0314]** Three variants derived from the oad gene were identified that had an increase in oxaloacetate alpha-decarboxylase activity when evaluated in the spectrophotometric assay. Results are provided in the following table. The variants had

1.6 fold, 2.2 fold, and 2.8 fold increases in oxaloacetate alpha-decarboxylase specific activity, respectively. The variants carried the following changes relative to the parent oad gene: N45T, R249L, D302G, V418A, and L476Q; T479N; and: R394C, D434G, and T511A, respectively, where the first amino acid is the one found in the parent sequence at the specified site, and the second amino acid is the one found at that site in the variant (using single-letter codes for the amino acid). It is known in the art that various combinations of any and all of these mutations may be used to produce enzymes with increased oxaloacetate alpha-decarboxylase activity over the parental OAD enzyme. As is also known in the art, substitutions of other amino acids at these sites could also increase oxaloacetate alpha-decarboxylase specific activity relative to that of the parental enzyme.

Enzymes	Mutations	Average Specific Activity (units/mg)	Standard Deviation	Fold Increase
Native Oad	None	0.0078	0.0008	—
Mutant Oad 1	N45T, R249L, D302G, V418A, and L476Q	0.0121	0.0001	1.6
Mutant Oad2	T479N	0.0173	0.0009	2.2
Mutant Oad 3	R394C, D434G, and T511A	0.0222	0.0012	2.8

**[0315]** The following are non-limiting general prophetic examples directed to practicing the present invention in other microorganism species.

**[0316]** General Prophetic Example 9

Improvement of 3-HP Tolerance and/or  
Bio-Production in *Rhodococcus Erythropolis*

**[0317]** A series of *E. coli*-*Rhodococcus* shuttle vectors are available for expression in *R. erythropolis*, including, but not limited to, pRhBR17 and pDA71 (Kostichka et al., Appl. Microbiol. Biotechnol. 62:61-68(2003)). Additionally, a series of promoters are available for heterologous gene expression in *R. erythropolis* (see for example Nakashima et al., Appl. Environ. Microbiol. 70:5557-5568 (2004), and Tao et al., Appl. Microbiol. Biotechnol. 2005, DOI 10.1007/s00253-005-0064). Targeted gene disruption of chromosomal genes in *R. erythropolis* may be created using the method described by Tao et al., supra, and Brans et al. (Appl. Environ. Microbiol. 66: 2029-2036 (2000)). These published resources are incorporated by reference for their respective indicated teachings and compositions.

**[0318]** The nucleic acid sequences required for providing an increase in 3-HP tolerance, as described herein, optionally with nucleic acid sequences to provide and/or improve a 3-HP biosynthesis pathway, are cloned initially in pDA71 or pRhBR71 and transformed into *E. coli*. The vectors are then transformed into *R. erythropolis* by electroporation, as described by Kostichka et al., supra. The recombinants are grown in synthetic medium containing glucose and the tolerance to and/or bio-production of 3-HP are followed using methods known in the art or described herein.

General Prophetic Example 10

Improvement of 3-HP Tolerance and/or  
Bio-Production in *B. Licheniformis*

**[0319]** Most of the plasmids and shuttle vectors that replicate in *B. subtilis* are used to transform *B. licheniformis* by

either protoplast transformation or electroporation. The nucleic acid sequences required for improvement of 3-HP tolerance, and/or for 3-HP biosynthesis are isolated from various sources, codon optimized as appropriate, and cloned in plasmids pBE20 or pBE60 derivatives (Nagarajan et al., Gene 114:121-126 (1992)). Methods to transform *B. licheniformis* are known in the art (for example see Fleming et al. Appl. Environ. Microbiol., 61(11):3775-3780 (1995)). These published resources are incorporated by reference for their respective indicated teachings and compositions.

**[0320]** The plasmids constructed for expression in *B. subtilis* are transformed into *B. licheniformis* to produce a recombinant microorganism that then demonstrates improved 3-HP tolerance, and, optionally, 3-HP bio-production.

General Prophetic Example 11

Improvement of 3-HP Tolerance and/or  
Bio-Production in *Paenibacillus Macerans*

**[0321]** Plasmids are constructed as described herein for expression in *B. subtilis* and used to transform *Paenibacillus macerans* by protoplast transformation to produce a recombinant microorganism that demonstrates improved 3-HP tolerance, and, optionally, 3-HP bio-production.

General Prophetic Example 12

Expression of 3-HP Tolerance and/or Bio-Production  
in *Alcaligenes (Ralstonia) Eutrophus* (Currently  
Referred to as *Cupriavidus Necator*)

**[0322]** Methods for gene expression and creation of mutations in *Alcaligenes eutrophus* are known in the art (see for example Taghavi et al., Appl. Environ. Microbiol., 60(10): 3585-3591 (1994)). This published resource is incorporated by reference for its indicated teachings and compositions. Any of the nucleic acid sequences identified to improve 3-HP tolerance, and/or for 3-HP biosynthesis are isolated from various sources, codon optimized as appropriate, and cloned in any of the broad host range vectors described herein, and electroporated to generate recombinant microorganisms that demonstrate improved 3-HP tolerance, and, optionally, 3-HP bio-production. The poly(hydroxybutyrate) pathway in *Alcaligenes* has been described in detail, a variety of genetic techniques to modify the *Alcaligenes eutrophus* genome is known, and those tools can be applied for engineering a 3-HP toleragenic or, optionally, a 3-HP-gena-toleragenic recombinant microorganism.

General Prophetic Example 13

Improvement of 3-HP Tolerance and/or  
Bio-Production in *Pseudomonas Putida*

**[0323]** Methods for gene expression in *Pseudomonas putida* are known in the art (see for example Ben-Bassat et al., U.S. Pat. No. 6,586,229, which is incorporated herein by reference for these teachings). Any of the nucleic acid sequences identified to improve 3-HP tolerance, and/or for 3-HP biosynthesis are isolated from various sources, codon optimized as appropriate, and cloned in any of the broad host range vectors described herein, and electroporated to generate recombinant microorganisms that demonstrate improved 3-HP tolerance, and, optionally, 3-HP biosynthetic production. For example, these nucleic acid sequences are inserted

into pUCP18 and this ligated DNA are electroporated into electrocompetent *Pseudomonas putida* KT2440 cells to generate recombinant *P. putida* microorganisms that exhibit increased 3-HP tolerance and optionally also comprise 3-HP biosynthesis pathways comprised at least in part of introduced nucleic acid sequences.

#### General Prophetic Example 14

##### Improvement of 3-HP Tolerance and/or Bio-Production in *Lactobacillus Plantarum*

**[0324]** The *Lactobacillus* genus belongs to the Lactobacillales family and many plasmids and vectors used in the transformation of *Bacillus subtilis* and *Streptococcus* are used for *Lactobacillus*. Non-limiting examples of suitable vectors include pAM.beta.1 and derivatives thereof (Renault et al., Gene 183:175-182 (1996); and O'Sullivan et al., Gene 137:227-231 (1993)); pMBB1 and pHW800, a derivative of pMBB1 (Wyckoff et al. Appl. Environ. Microbiol. 62:1481-1486 (1996)); pMG1, a conjugative plasmid (Tanimoto et al., J. Bacteriol. 184:5800-5804 (2002)); pNZ9520 (Kleerebezem et al., Appl. Environ. Microbiol. 63:4581-4584 (1997)); pAM401 (Fujimoto et al., Appl. Environ. Microbiol. 67:1262-1267 (2001)); and pAT392 (Arthur et al., Antimicrob. Agents Chemother. 38:1899-1903 (1994)). Several plasmids from *Lactobacillus plantarum* have also been reported (e.g., van Kranenburg R, Golic N, Bongers R, Leer R J, de Vos W M, Siezen R J, Kleerebezem M. Appl. Environ. Microbiol. 2005 March; 71(3): 1223-1230).

#### General Prophetic Example 15

##### Improvement of 3-HP Tolerance and/or Bio-Production in *Enterococcus faecium*, *Enterococcus gallinarum*, and *Enterococcus faecalis*

**[0325]** The *Enterococcus* genus belongs to the Lactobacillales family and many plasmids and vectors used in the transformation of *Lactobacillus*, *Bacillus subtilis*, and *Streptococcus* are used for *Enterococcus*. Non-limiting examples of suitable vectors include pAM.beta.1 and derivatives thereof (Renault et al., Gene 183:175-182 (1996); and O'Sullivan et al., Gene 137:227-231 (1993)); pMBB1 and pHW800, a derivative of pMBB1 (Wyckoff et al. Appl. Environ. Microbiol. 62:1481-1486 (1996)); pMG1, a conjugative plasmid (Tanimoto et al., J. Bacteriol. 184:5800-5804 (2002)); pNZ9520 (Kleerebezem et al., Appl. Environ. Microbiol. 63:4581-4584 (1997)); pAM401 (Fujimoto et al., Appl. Environ. Microbiol. 67:1262-1267 (2001)); and pAT392 (Arthur et al., Antimicrob. Agents Chemother. 38:1899-1903 (1994)). Expression vectors for *E. faecalis* using the nisA gene from *Lactococcus* may also be used (Eichenbaum et al., Appl. Environ. Microbiol. 64:2763-2769 (1998)). Additionally, vectors for gene replacement in the *E. faecium* chromosome are used (Nallaapareddy et al., Appl. Environ. Microbiol. 72:334-345 (2006)).

**[0326]** For each of the General Prophetic Examples 65-71, the following 3-HP bio-production comparison may be incorporated thereto: Using analytical methods for 3-HP such as are described in Subsection III of Common Methods Section, 3-HP is obtained in a measurable quantity at the conclusion of a respective bio-production event conducted with the respective recombinant microorganism (see types of bio-production events, incorporated by reference into each respective General Prophetic Example). That measurable quantity is sub-

stantially greater than a quantity of 3-HP produced in a control bio-production event using a suitable respective control microorganism lacking the functional 3-HP pathway so provided in the respective General Prophetic Example. Tolerance improvements also may be assessed by any recognized comparative measurement technique, such as by using a MIC protocol provided in the Common Methods Section.

**[0327]** Common Methods Section

**[0328]** All methods in this Section are provided for incorporation into the Examples where so referenced.

**[0329]** Subsection I. Microorganism Species and Strains, Cultures, and Growth Media

**[0330]** Bacterial species, that may be utilized as needed, are as follows:

**[0331]** *Acinetobacter calcoaceticus* (DSMZ #1139) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in Brain Heart Infusion (BHI) Broth (RPI Corp, Mt. Prospect, Ill., USA). Serial dilutions of the resuspended *A. calcoaceticus* culture are made into BHI and are allowed to grow aerobically for 48 hours at 37° C. at 250 rpm until saturated.

**[0332]** *Bacillus subtilis* is a gift from the Gill lab (University of Colorado at Boulder) and is obtained as an actively growing culture. Serial dilutions of the actively growing *B. subtilis* culture are made into Luria Broth (RPI Corp, Mt. Prospect, Ill., USA) and are allowed to grow aerobically for 24 hours at 37° C. at 250 rpm until saturated.

**[0333]** *Chlorobium limicola* (DSMZ #245) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended using Pfennig's Medium I and II (#28 and 29) as described per DSMZ instructions. *C. limicola* is grown at 25° C. under constant vortexing.

**[0334]** *Citrobacter braakii* (DSMZ #30040) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in Brain Heart Infusion (BHI) Broth (RPI Corp, Mt. Prospect, Ill., USA). Serial dilutions of the resuspended *C. braakii* culture are made into BHI and are allowed to grow aerobically for 48 hours at 30° C. at 250 rpm until saturated.

**[0335]** *Clostridium acetobutylicum* (DSMZ #792) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in *Clostridium acetobutylicum* medium (#411) as described per DSMZ instructions. *C. acetobutylicum* is grown anaerobically at 37° C. at 250 rpm until saturated.

**[0336]** *Clostridium aminobutyricum* (DSMZ #2634) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in *Clostridium aminobutyricum* medium (#286) as described per DSMZ instructions. *C. aminobutyricum* is grown anaerobically at 37° C. at 250 rpm until saturated.

**[0337]** *Clostridium kluyveri* (DSMZ #555) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as an actively growing culture. Serial dilutions of *C. kluyveri* culture are made into *Clostridium kluyveri* medium (#286) as described per DSMZ instructions. *C. kluyveri* is grown anaerobically at 37° C. at 250 rpm until saturated.

[0338] *Cupriavidus metallidurans* (DMSZ #2839) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in Brain Heart Infusion (BHI) Broth (RPI Corp, Mt. Prospect, Ill., USA). Serial dilutions of the resuspended *C. metallidurans* culture are made into BHI and are allowed to grow for aerobically for 48 hours at 30° C. at 250 rpm until saturated.

[0339] *Cupriavidus necator* (DSMZ #428) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in Brain Heart Infusion (BHI) Broth (RPI Corp, Mt. Prospect, Ill., USA). Serial dilutions of the resuspended *C. necator* culture are made into BHI and are allowed to grow for aerobically for 48 hours at 30° C. at 250 rpm until saturated. As noted elsewhere, previous names for this species are *Alcaligenes eutrophus* and *Ralstonia eutrophus*.

[0340] *Desulfovibrio fructosovorans* (DSMZ #3604) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in *Desulfovibrio fructosovorans* medium (#63) as described per DSMZ instructions. *D. fructosovorans* is grown anaerobically at 37° C. at 250 rpm until saturated.

[0341] *Escherichia coli* Crooks (DSMZ #1576) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in Brain Heart Infusion (BHI) Broth (RPI Corp, Mt. Prospect, Ill., USA). Serial dilutions of the resuspended *E. coli* Crooks culture are made into BHI and are allowed to grow for aerobically for 48 hours at 37° C. at 250 rpm until saturated.

[0342] *Escherichia coli* K12 is a gift from the Gill lab (University of Colorado at Boulder) and is obtained as an actively growing culture. Serial dilutions of the actively growing *E. coli* K12 culture are made into Luria Broth (RPI Corp, Mt. Prospect, Ill., USA) and are allowed to grow for aerobically for 24 hours at 37° C. at 250 rpm until saturated.

[0343] *Halobacterium salinarum* (DSMZ #1576) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in *Halobacterium* medium (#97) as described per DSMZ instructions. *H. salinarum* is grown aerobically at 37° C. at 250 rpm until saturated.

[0344] *Lactobacillus delbrueckii* (#4335) is obtained from WYEAST USA (Odell, Oreg., USA) as an actively growing culture. Serial dilutions of the actively growing *L. delbrueckii* culture are made into Brain Heart Infusion (BHI) broth (RPI Corp, Mt. Prospect, Ill., USA) and are allowed to grow for aerobically for 24 hours at 30° C. at 250 rpm until saturated.

[0345] *Metallosphaera sedula* (DSMZ #5348) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as an actively growing culture. Serial dilutions of *M. sedula* culture are made into *Metallosphaera* medium (#485) as described per DSMZ instructions. *M. sedula* is grown aerobically at 65° C. at 250 rpm until saturated.

[0346] *Propionibacterium freudenreichii* subsp. *shermanii* (DSMZ #4902) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in PYG-medium (#104) as described per DSMZ

instructions. *P. freudenreichii* subsp. *shermanii* is grown anaerobically at 30° C. at 250 rpm until saturated.

[0347] *Pseudomonas putida* is a gift from the Gill lab (University of Colorado at Boulder) and is obtained as an actively growing culture. Serial dilutions of the actively growing *P. putida* culture are made into Luria Broth (RPI Corp, Mt. Prospect, Ill., USA) and are allowed to grow for aerobically for 24 hours at 37° C. at 250 rpm until saturated.

[0348] *Streptococcus mutans* (DSMZ #6178) is obtained from the German Collection of Microorganisms and Cell Cultures (Braunschweig, Germany) as a vacuum dried culture. Cultures are then resuspended in Luria Broth (RPI Corp, Mt. Prospect, Ill., USA). *S. mutans* is grown aerobically at 37° C. at 250 rpm until saturated.

[0349] Subsection II: Gel Preparation, DNA Separation, Extraction, Ligation, and Transformation Methods:

[0350] Molecular biology grade agarose (RPI Corp, Mt. Prospect, Ill., USA) is added to 1× TAE to make a 1% Agarose in TAE. To obtain 50× TAE add the following to 900 ml distilled H<sub>2</sub>O: 242 g Tris base (RPI Corp, Mt. Prospect, Ill., USA), 57.1 ml Glacial Acetic Acid (Sigma-Aldrich, St. Louis, Mo., USA), 18.6 g EDTA (Fisher Scientific, Pittsburgh, Pa. USA), and adjust volume to 1 L with additional distilled water. To obtain 1× TAE, add 20 mL of 50× TAE to 980 mL of distilled water. The agarose-TAE solution is then heated until boiling occurred and the agarose is fully dissolved. The solution is allowed to cool to 50° C. before 10 mg/mL ethidium bromide (Acros Organics, Morris Plains, N.J., USA) is added at a concentration of 5 ul per 100 mL of 1% agarose solution. Once the ethidium bromide is added, the solution is briefly mixed and poured into a gel casting tray with the appropriate number of combs (Idea Scientific Co., Minneapolis, Minn., USA) per sample analysis. DNA samples are then mixed accordingly with 5× TAE loading buffer. 5× TAE loading buffer consists of 5× TAE (diluted from 50× TAE as described herein), 20% glycerol (Acros Organics, Morris Plains, N.J., USA), 0.125% Bromophenol Blue (Alfa Aesar, Ward Hill, Mass., USA), and adjust volume to 50 mL with distilled water. Loaded gels are then run in gel rigs (Idea Scientific Co., Minneapolis, Minn., USA) filled with 1× TAE at a constant voltage of 125 volts for 25-30 minutes. At this point, the gels are removed from the gel boxes with voltage and visualized under a UV transilluminator (FO-TODYNE Inc., Hartland, Wis., USA).

[0351] The DNA isolated through gel extraction is then extracted using the QIAquick Gel Extraction Kit following manufacturer's instructions (Qiagen (Valencia, Calif. USA)). Similar methods are known to those skilled in the art.

[0352] The thus-extracted DNA then may be ligated into pSMART (Lucigen Corp, Middleton, Wis., USA), StrataClone (Stratagene, La Jolla, Calif., USA) or pCR2.1-TOPO TA (Invitrogen Corp, Carlsbad, Calif., USA) according to manufacturer's instructions. These methods are described in the next subsection of Common Methods.

[0353] Ligation Methods:

[0354] For ligations into pSMART Vectors:

[0355] Gel extracted DNA is blunted using PCRTerminator (Lucigen Corp, Middleton, Wis., USA) according to manufacturer's instructions. Then 500 ng of DNA is added to 2.5 uL 4× CloneSmart vector premix, 1 ul CloneSmart DNA ligase (Lucigen Corp, Middleton, Wis., USA) and distilled water is added for a total volume of 10 ul. The reaction is then allowed to sit at room temperature for 30 minutes and then heat inactivated at 70° C. for 15 minutes and then placed on

ice. *E. coli* 10G Chemically Competent cells (Lucigen Corp, Middleton, Wis., USA) are thawed for 20 minutes on ice. 40 ul of chemically competent cells are placed into a microcentrifuge tube and 1 ul of heat inactivated CloneSmart Ligation is added to the tube. The whole reaction is stirred briefly with a pipette tip. The ligation and cells are incubated on ice for 30 minutes and then the cells are heat shocked for 45 seconds at 42° C. and then put back onto ice for 2 minutes. 960 ul of room temperature Recovery media (Lucigen Corp, Middleton, Wis., USA) and places into microcentrifuge tubes. Shake tubes at 250 rpm for 1 hour at 37° C. Plate 100 ul of transformed cells on Luria Broth plates (RPI Corp, Mt. Prospect, Ill., USA) plus appropriate antibiotics depending on the pSMART vector used. Incubate plates overnight at 37° C.

**[0356]** General Transformation and Related Culture Methodologies:

**[0357]** Chemically competent transformation protocols are carried out according to the manufacturer's instructions or according to the literature contained in Molecular Cloning (Sambrook and Russell, 2001). Generally, plasmid DNA or ligation products are chilled on ice for 5 to 30 min. in solution with chemically competent cells. Chemically competent cells are a widely used product in the field of biotechnology and are available from multiple vendors, such as those indicated in this Subsection. Following the chilling period cells generally are heat-shocked for 30 seconds at 42° C. without shaking, re-chilled and combined with 250 microliters of rich media, such as S.O.C. Cells are then incubated at 37° C. while shaking at 250 rpm for 1 hour. Finally, the cells are screened for successful transformations by plating on media containing the appropriate antibiotics.

**[0358]** Alternatively, selected cells may be transformed by electroporation methods such as are known to those skilled in the art.

**[0359]** The choice of an *E. coli* host strain for plasmid transformation is determined by considering factors such as plasmid stability, plasmid compatibility, plasmid screening methods and protein expression. Strain backgrounds can be changed by simply purifying plasmid DNA as described herein and transforming the plasmid into a desired or otherwise appropriate *E. coli* host strain such as determined by experimental necessities, such as any commonly used cloning strain (e.g., DH5 $\alpha$ , Top10F', *E. coli* 10G, etc.).

**[0360]** Plasmid DNA was prepared using the commercial miniprep kit from Qiagen (Valencia, Calif. USA) according to manufacturer's instructions.

**[0361]** Subsection IIIa. 3-HP Preparation

**[0362]** A 3-HP stock solution was prepared as follows. A vial of  $\beta$ -propiolactone (Sigma-Aldrich, St. Louis, Mo., USA) was opened under a fume hood and the entire bottle contents was transferred to a new container sequentially using a 25-mL glass pipette. The vial was rinsed with 50 mL of HPLC grade water and this rinse was poured into the new container. Two additional rinses were performed and added to the new container. Additional HPLC grade water was added to the new container to reach a ratio of 50 mL water per 5 mL  $\beta$ -propiolactone. The new container was capped tightly and allowed to remain in the fume hood at room temperature for 72 hours. After 72 hours the contents were transferred to centrifuge tubes and centrifuged for 10 minutes at 4,000 rpm. Then the solution was filtered to remove particulates and, as needed, concentrated by use of a rotary evaporator at room

temperature. Assay for concentration was conducted, and dilution to make a standard concentration stock solution was made as needed.

**[0363]** Subsection IIIb. HPLC, GC/MS and Other Analytical Methods for 3-HP Detection (Analysis of Cultures for 3-HP Production)

**[0364]** For HPLC analysis of 3-HP, the Waters chromatography system (Milford, Mass.) consisted of the following: 600S Controller, 616 Pump, 717 Plus Autosampler, 486 Tunable UV Detector, and an in-line mobile phase Degas ser. In addition, an Eppendorf external column heater is used and the data are collected using an SRI (Torrance, Calif.) analog-to-digital converter linked to a standard desk top computer. Data are analyzed using the SRI Peak Simple software. A Coregel 64H ion exclusion column (Transgenomic, Inc., San Jose, Calif.) is employed. The column resin is a sulfonated polystyrene divinyl benzene with a particle size of 10  $\mu$ m and column dimensions are 300 $\times$ 7.8 mm. The mobile phase consisted of sulfuric acid (Fisher Scientific, Pittsburgh, Pa. USA) diluted with deionized (18 MS $\Omega$ m) water to a concentration of 0.02 N and vacuum filtered through a 0.2  $\mu$ m nylon filter. The flow rate of the mobile phase is 0.6 mL/min. The UV detector is operated at a wavelength of 210 nm and the column is heated to 60° C. The same equipment and method as described herein is used for 3-HP analyses for relevant prophetic examples.

**[0365]** The following method is used for GC-MS analysis of 3-HP. Soluble monomeric 3-HP is quantified using GC-MS after a single extraction of the fermentation media with ethyl acetate. Once the 3-HP has been extracted into the ethyl acetate, the active hydrogens on the 3-HP are replaced with trimethylsilyl groups using N,O-Bis-(Trimethylsilyl)trifluoroacetamide to make the compound volatile for GC analysis. A standard curve of known 3-HP concentrations is prepared at the beginning of the run and a known quantity of ketohexanoic acid (1 g/L) is added to both the standards and the samples to act as an internal standard for Quantitation, with tropic acid as an additional internal standard. The 3-HP content of individual samples is then assayed by examining the ratio of the ketohexanoic acid ion ( $m/z=247$ ) to the 3-HP ion (219) and compared to the standard curve. 3-HP is quantified using a 3HP standard curve at the beginning of the run and the data are analyzed using HP Chemstation. The GC-MS system consists of a Hewlett Packard model 5890 GC and Hewlett Packard model 5972 MS. The column is Supelco SPB-1 (60 m $\times$ 0.32 mm $\times$ 0.25  $\mu$ m film thickness). The capillary coating is a non-polar methylsilicone. The carrier gas is helium at a flow rate of 1 mL/min. The 3-HP as derivatized is separated from other components in the ethyl acetate extract using either of two similar temperature regimes. In a first temperature gradient regime, the column temperature starts with 40° C. for 1 minute, then is raised at a rate of 10° C./minute to 235° C., and then is raised at a rate of 50° C./minute to 300° C. In a second temperature regime, which was demonstrated to process samples more quickly, the column temperature starts with 70° C. which is held for 1 min, followed by a ramp-up of 10° C./minute to 235° C. which is followed by a ramp-up of 50° C./minute to 300° C.

**[0366]** The embodiments, variations, sequences, and figures described herein should provide an indication of the utility and versatility of the present invention. Other embodiments that do not provide all of the features and advantages set forth herein may also be utilized, without departing from the spirit and scope of the present invention. Such modifications and variations are considered to be within the scope of the invention.

TABLE 1

Protein Function	E.C. Classification	Gene Name in <i>E. coli</i>	SEQ ID NO. of <i>E. coli</i> gene	SEQ ID NO. of Expressed Enzyme
Glucose transport	N/A	galP	003	004
Phosphoenolpyruvate carboxykinase	4.1.1.49	pckA	007	008
Phosphoenolpyruvate carboxylase	4.1.1.49	ppc	009	010

TABLE 2

Protein Function	E.C. Classification	Gene Name in <i>E. coli</i>	SEQ ID NO. of <i>E. coli</i> gene	SEQ ID NO. of Expressed Enzyme
Pyruvate dehydrogenase E1p	1.2.4.1	aceE	011	012
lipoate acetyltransferase/dihydrolipoamide acetyltransferase	2.3.1.12	aceF	013	014

TABLE 2-continued

Protein Function	E.C. Classification	Gene Name in <i>E. coli</i>	SEQ ID NO. of <i>E. coli</i> gene	SEQ ID NO. of Expressed Enzyme
Pyruvate dehydrogenase E3 (lipoamide dehydrogenase)	1.8.1.4	l pd	015	016
Lactate dehydrogenase	1.1.1.28	ldhA	017	018
Pyruvate formate lyase (B "inactive")	2.3.1.—	pfIB	019	020
Pyruvate oxidase	1.2.2.2	poxB	021	022
Phosphate acetyltransferase	2.3.1.8	Pta	023	024
Heat stable, histidyl phosphorylatable protein (of PTS)	N/A	ptsH (HPr)	025	026
Phosphoryl transfer protein (of PTS)	N/A	ptsI	027	028
Polypeptide chain (of PTS)	N/A	Crr	029	030
Pyruvate kinase I	2.7.1.40	pykA	031	032
Pyruvate kinase II	2.7.1.40	pykF	033	034

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 68

<210> SEQ ID NO 1

<211> LENGTH: 452

<212> TYPE: PRT

<213> ORGANISM: *Bacillus cereus*

<400> SEQUENCE: 1

Met Met Met Met Lys Thr Lys Gln Thr Asp Glu Leu Leu Ala Lys Asp  
1 5 10 15

Glu Gln Tyr Val Trp His Gly Met Arg Pro Phe Ser Pro Asn Ser Thr  
20 25 30

Met Val Gly Ala Lys Ala Glu Gly Cys Trp Val Glu Asp Ile Gln Gly  
35 40 45

Lys Arg Tyr Leu Asp Gly Met Ser Gly Leu Trp Cys Val Asn Ser Gly  
50 55 60

Tyr Gly Arg Lys Glu Leu Ala Glu Ala Ala Tyr Lys Gln Leu Gln Thr  
65 70 75 80

Leu Ser Tyr Phe Pro Met Ser Gln Ser His Glu Pro Ala Ile Lys Leu  
85 90 95

Ala Glu Lys Leu Asn Glu Trp Leu Gly Gly Glu Tyr Val Ile Phe Phe  
100 105 110

Ser Asn Ser Gly Ser Glu Ala Asn Glu Thr Ala Phe Lys Ile Ala Arg  
115 120 125

Gln Tyr Tyr Ala Gln Lys Gly Glu Pro His Arg Tyr Lys Phe Met Ser  
130 135 140

Arg Tyr Arg Gly Tyr His Gly Asn Thr Met Ala Thr Met Ala Ala Thr  
145 150 155 160

Gly Gln Ala Gln Arg Arg Tyr Gln Tyr Glu Pro Phe Ala Ser Gly Phe  
165 170 175

-continued

---

Leu His Val Thr Pro Pro Asp Cys Tyr Arg Met Pro Glu Ile Glu Gly  
                   180                  185                  190

Gln His Ile Tyr Asp Val Glu Cys Val Lys Glu Val Asp Arg Val Met  
                   195                  200                  205

Thr Trp Glu Leu Ser Glu Thr Ile Ala Ala Phe Ile Met Glu Pro Ile  
                   210                  215                  220

Ile Thr Gly Gly Gly Ile Leu Met Pro Pro Gln Asp Tyr Met Lys Ala  
 225                  230                  235                  240

Val His Glu Met Cys Gln Lys His Gly Ala Leu Leu Ile Ser Asp Glu  
                   245                  250                  255

Val Ile Cys Gly Phe Gly Arg Thr Gly Lys Ala Phe Gly Phe Met Asn  
                   260                  265                  270

Tyr Asp Val Lys Pro Asp Ile Ile Thr Met Ala Lys Gly Ile Thr Ser  
                   275                  280                  285

Ala Tyr Leu Pro Leu Ser Ala Thr Ala Val Lys Lys Glu Ile Tyr Glu  
                   290                  295                  300

Ala Phe Lys Gly Lys Gly Glu Tyr Glu Phe Phe Arg His Ile Asn Thr  
 305                  310                  315                  320

Phe Gly Gly Asn Pro Ala Ala Cys Ala Leu Ala Leu Lys Asn Leu Glu  
                   325                  330                  335

Ile Met Glu Asn Glu Asn Leu Ile Glu Arg Ser Ala Gln Met Gly Ser  
                   340                  345                  350

Leu Leu Leu Glu Gln Leu Lys Asp Glu Ile Gly Glu His Pro Leu Val  
                   355                  360                  365

Gly Asn Ile Arg Gly Lys Gly Leu Leu Val Gly Ile Glu Leu Val Asn  
                   370                  375                  380

Asp Lys Glu Thr Lys Glu Pro Ile Asp Asn Asp Lys Ile Ala Ser Val  
 385                  390                  395                  400

Val Asn Ala Cys Lys Glu Lys Gly Leu Ile Ile Gly Arg Asn Gly Met  
                   405                  410                  415

Thr Thr Ala Gly Tyr Asn Asn Val Leu Thr Leu Ala Pro Pro Leu Val  
                   420                  425                  430

Ile Ser Ser Glu Glu Ile Ala Phe Val Val Gly Thr Leu Lys Thr Ala  
                   435                  440                  445

Met Glu Arg Ile  
                   450

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 1371

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

 <223> OTHER INFORMATION: Description of Artificial Sequence:  
 Codon-optimized (for E. Coli) synthetic polynucleotide

&lt;400&gt; SEQUENCE: 2

```

tgatgatgat gaaaaccaag cagacggatg agctgctggc gaaggacgaa cagtatgtgt    60
ggcatggcat ggcgccggtt agcccgaatt ccactatggt tggtgcaaaa gcagaaggtt    120
gctgggtcga ggacattcag ggcaagcgtt atctggatgg catgagcggc ctgtggtgcg    180
tgaattcggg ttatggccgt aaagagttgg cggaggcggc gtacaagcag ctgcaaacce    240
tgagctatct cccgatgtct cagagccacg aaccggcgat caaactggcg gagaaactga    300
atgaatggtt ggggtgtgaa tatgtgatct ttttcagcaa ttccggtagc gaggcaaacg    360

```

-continued

---

```

aacggcctt caagattgcg cgccaatact acgcgcaaaa aggcgaaccg catcgttaca 420
agttcatgtc tcgttaccgc ggctatcacg gcaataccat ggccaccatg gccgccaccg 480
gtcaagcgca acgccgttac caatacgagc cgtttgcttc tggttttctg catggttacgc 540
ctccggattg ttaccgcatg ccggaaatcg aaggccagca catctatgac gtogaatgcg 600
tgaaagaagt ggatcgtgtg atgacctggg aactgtccga aaccatcgca gccttcatca 660
tggagccgat taccacgggc ggtggtattc tgatgccacc gcaggactac atgaaggcag 720
ttcacgagat gtgccagaag cacggcgctc tgctgattag cgacgaagtc atctgcggtt 780
tcggccgtac gggtaaggcg tttggtttca tgaactacga tgtaaaccg gacatcatta 840
cgatggcgaa aggtattacg agcgcatatc tgctctgag cgcgactgcg gttaagaaag 900
agatctacga agccttcaag ggtaagggtg agtatgaatt ctttcgtcac atcaatacct 960
ttggtggtaa tccagccgcy tgtgcattgg cactgaaaaa cttggagatt atggagaatg 1020
aaaacctgat tgaacgcagc gcccaaatgg gtagcctgct gctggagcag ctgaaggatg 1080
agatcggcga gcaccgctg gttggtaaca tccgcggcaa gggcctgctg gtccgcatcg 1140
agctgggtcaa cgacaaagaa accaaagaac cgatcgataa tgacaagatt gctagcgtcg 1200
tgaatgcttg taaagagaag ggtctgatta tcggtcgtaa cggcatgacc accgcggggt 1260
acaacaacgt tctgacctg gcaccgccgc tggatgatcag ctccgaagag attgcgtttg 1320
tggttggtac gctgaaaacc gcaatggagc gtattcacca tcatcatcac t 1371

```

```

<210> SEQ ID NO 3
<211> LENGTH: 298
<212> TYPE: PRT
<213> ORGANISM: Pseudomonas aeruginosa

```

```

<400> SEQUENCE: 3

```

```

Met Thr Asp Ile Ala Phe Leu Gly Leu Gly Asn Met Gly Gly Pro Met
1           5           10           15
Ala Ala Asn Leu Leu Lys Ala Gly His Arg Val Asn Val Phe Asp Leu
20          25          30
Gln Pro Lys Ala Val Leu Gly Leu Val Glu Gln Gly Ala Gln Gly Ala
35          40          45
Asp Ser Ala Leu Gln Cys Cys Glu Gly Ala Glu Val Val Ile Ser Met
50          55          60
Leu Pro Ala Gly Gln His Val Glu Ser Leu Tyr Leu Gly Asp Asp Gly
65          70          75          80
Leu Leu Ala Arg Val Ala Gly Lys Pro Leu Leu Ile Asp Cys Ser Thr
85          90          95
Ile Ala Pro Glu Thr Ala Arg Lys Val Ala Glu Ala Ala Ala Ala Lys
100         105         110
Gly Leu Thr Leu Leu Asp Ala Pro Val Ser Gly Gly Val Gly Gly Ala
115        120        125
Arg Ala Gly Thr Leu Ser Phe Ile Val Gly Gly Pro Ala Glu Gly Phe
130        135        140
Ala Arg Ala Arg Pro Val Leu Glu Asn Met Gly Arg Asn Ile Phe His
145        150        155        160
Ala Gly Asp His Gly Ala Gly Gln Val Ala Lys Ile Cys Asn Asn Met
165        170        175

```



-continued

---

Leu Leu Gly Ile Leu Met Ala Gly Thr Ala Glu Ala Leu Ala Leu Gly  
                   180                                  185                                  190

Val Lys Asn Gly Leu Asp Pro Ala Val Leu Ser Glu Val Met Lys Gln  
                   195                                  200                                  205

Ser Ser Gly Gly Asn Trp Ala Leu Asn Leu Tyr Asn Pro Trp Pro Gly  
                   210                                  215                                  220

Val Met Pro Gln Ala Pro Ala Ser Asn Gly Tyr Ala Gly Gly Phe Gln  
                   225                                  230                                  235                                  240

Val Arg Leu Met Asn Lys Asp Leu Gly Leu Ala Leu Ala Asn Ala Gln  
                                   245                                  250                                  255

Ala Val Gln Ala Ser Thr Pro Leu Gly Ala Leu Ala Arg Asn Leu Phe  
                                   260                                  265                                  270

Ser Leu His Ala Gln Ala Asp Ala Glu His Glu Gly Leu Asp Phe Ser  
                   275                                  280                                  285

Ser Ile Gln Lys Leu Tyr Arg Gly Lys Asp  
                   290                                  295

<210> SEQ ID NO 4  
 <211> LENGTH: 895  
 <212> TYPE: DNA  
 <213> ORGANISM: Pseudomonas aeruginosa

<400> SEQUENCE: 4

```

atgaccgaca ttgcgtttct gggctctgggc aatatgggcg gtccgatggc cgccaacctg    60
ctgaaagccg gccaccgtgt gaatgtgttc gacctgcaac caaaagcggc cctgggcttg    120
gttgagcaag gcgctgaggg cgcagactct gctctgcaat gttgtgaggg tgcggaggtc    180
gtgatttcta tgctgccagc aggccagcat gtggaaagcc tgtacctggg cgatgatggg    240
ctgctggcac gcgtggcggg caagcctttg ctgattgact gtagcaccat cgcaccggaa    300
acggcgcgta aggtggcggg ggcagccgca gcaaagggcc tgacgctgct ggatgccccg    360
gtttcggggc gtgtcgggtg tgcccgtgca ggtacgctgt cgtttatcgt ggggtggctcg    420
gcggagggtt ttgcgcgtgc gcgtccggtt ctggagaata tgggtcgcaa cattttccac    480
gcgggtgatc acggcgtggt tcaggtggcg aaaatctgta acaacatgct gctgggtatc    540
ttgatggcgg gcaccgccga agccttggcg ctgggcgtca aaaacggtct ggaccgccga    600
gtgctgtccg aagtgatgaa acagagcagc ggtggtaact gggcgtgaa tctgtacaat    660
ccgtggccgg gtgtgatgcc gcaggcccca gcctctaata gctacgcagg cggcttccaa    720
gtgcgcctga tgaacaaaga cctgggcctg gcgtggcga atgcgcaagc ggtccaagcg    780
agcaccgccg tgggcgcact ggcccgtaac ctgtttagcc tgcacgctca agccgacgcc    840
gagcacgaag gtctggactt cagctctatt caaaaactgt atcgcggtaa ggatt          895

```

<210> SEQ ID NO 5  
 <211> LENGTH: 1392  
 <212> TYPE: DNA  
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 5

```

cctgacgcta aaaaacaggg gcgggtcaaac aaggcaatga cgtttttcgt ctgcttcctt    60
gccgctctgg cgggattact ctttggcctg gatatcggtg taattgctgg cgcactgccg    120
tttattgcag atgaattcca gattacttcg cacacgcaag aatgggtcgt aagctccatg    180

```

-continued

---

```

atgttcggtg cggcagtcgg tgcggtgggc agcggctggc tctcctttaa actcgggggc 240
aaaaagagcc tgatgatcgg cgcaattttg tttgttgccg gttcgcctgt ctctgcggct 300
gcgccaaacg ttgaagtact gattctttcc cgcgttctac tggggctggc ggtgggtgtg 360
gcctcttata ccgcaccgct gtacctctct gaaattgccc cggaaaaaat tcgtggcagt 420
atgatctcga tgtatcagtt gatgatcact atcgggatcc tcggtgctta tctttctgat 480
accgccttca gctacaccgg tgcattggcg tggatgctgg gtgtgattat catcccggca 540
athttgetgc tgattggtgt cttcttctcg ccagacagcc cacgttggtt tgcgcgcaaa 600
cgccgttttg ttgatgccga acgctgctg ctacgcctgc gtgacaccag cgcggaagcg 660
aaacgcgaac tggatgaaat ccgtgaaagt ttgcaggta aacagagtgg ctgggctgctg 720
tttaaagaga acagcaactt ccgcccgcgc gtgttccttg gcgtactgtt gcaggtaatg 780
cagcaattca ccgggatgaa cgtcatcatg tattacgcgc cgaaaatctt cgaactggcg 840
ggttatacca aactaccga gcaaatgtgg gggaccgtga ttgtcggcct gaccaacgta 900
cttgccacct ttatcgcaat cggccttggt gaccgctggg gacgtaaacc aacgctaacg 960
ctgggcttcc tggatgaggc tgctggcatg ggcgtactcg gtacaatgat gcatatcggg 1020
attcactctc cgtcggcgca gtatttcgcc atcgccatgc tgctgatgtt tattgtcggg 1080
tttgccatga gtgccggtcc gctgatttgg gtactgtgct ccgaaattca gccgctgaaa 1140
ggccgcgatt ttggcatcac ctgctccact gccaccaact ggattgcaa catgatcgtt 1200
ggcgcaacgt tcctgacct gctcaacacg ctgggtaacg ccaacacctt ctgggtgtat 1260
gcggtctga acgtactgtt tatcctgctg acattgtggc tggtagcgga aaccaaacac 1320
gtttcgtgg aacatattga acgtaatctg atgaaaggtc gtaaactgcg cgaaataggc 1380
gctcacgatt aa 1392

```

```

<210> SEQ ID NO 6
<211> LENGTH: 464
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli

```

```

<400> SEQUENCE: 6

```

```

Met Pro Asp Ala Lys Lys Gln Gly Arg Ser Asn Lys Ala Met Thr Phe
1           5           10           15
Phe Val Cys Phe Leu Ala Ala Leu Ala Gly Leu Leu Phe Gly Leu Asp
20           25           30
Ile Gly Val Ile Ala Gly Ala Leu Pro Phe Ile Ala Asp Glu Phe Gln
35           40           45
Ile Thr Ser His Thr Gln Glu Trp Val Val Ser Ser Met Met Phe Gly
50           55           60
Ala Ala Val Gly Ala Val Gly Ser Gly Trp Leu Ser Phe Lys Leu Gly
65           70           75           80
Arg Lys Lys Ser Leu Met Ile Gly Ala Ile Leu Phe Val Ala Gly Ser
85           90           95
Leu Phe Ser Ala Ala Ala Pro Asn Val Glu Val Leu Ile Leu Ser Arg
100          105          110
Val Leu Leu Gly Leu Ala Val Gly Val Ala Ser Tyr Thr Ala Pro Leu
115          120          125
Tyr Leu Ser Glu Ile Ala Pro Glu Lys Ile Arg Gly Ser Met Ile Ser
130          135          140

```

-continued

---

Met Tyr Gln Leu Met Ile Thr Ile Gly Ile Leu Gly Ala Tyr Leu Ser  
 145 150 155 160

Asp Thr Ala Phe Ser Tyr Thr Gly Ala Trp Arg Trp Met Leu Gly Val  
 165 170 175

Ile Ile Ile Pro Ala Ile Leu Leu Leu Ile Gly Val Phe Phe Leu Pro  
 180 185 190

Asp Ser Pro Arg Trp Phe Ala Ala Lys Arg Arg Phe Val Asp Ala Glu  
 195 200 205

Arg Val Leu Leu Arg Leu Arg Asp Thr Ser Ala Glu Ala Lys Arg Glu  
 210 215 220

Leu Asp Glu Ile Arg Glu Ser Leu Gln Val Lys Gln Ser Gly Trp Ala  
 225 230 235 240

Leu Phe Lys Glu Asn Ser Asn Phe Arg Arg Ala Val Phe Leu Gly Val  
 245 250 255

Leu Leu Gln Val Met Gln Gln Phe Thr Gly Met Asn Val Ile Met Tyr  
 260 265 270

Tyr Ala Pro Lys Ile Phe Glu Leu Ala Gly Tyr Thr Asn Thr Thr Glu  
 275 280 285

Gln Met Trp Gly Thr Val Ile Val Gly Leu Thr Asn Val Leu Ala Thr  
 290 295 300

Phe Ile Ala Ile Gly Leu Val Asp Arg Trp Gly Arg Lys Pro Thr Leu  
 305 310 315 320

Thr Leu Gly Phe Leu Val Met Ala Ala Gly Met Gly Val Leu Gly Thr  
 325 330 335

Met Met His Ile Gly Ile His Ser Pro Ser Ala Gln Tyr Phe Ala Ile  
 340 345 350

Ala Met Leu Leu Met Phe Ile Val Gly Phe Ala Met Ser Ala Gly Pro  
 355 360 365

Leu Ile Trp Val Leu Cys Ser Glu Ile Gln Pro Leu Lys Gly Arg Asp  
 370 375 380

Phe Gly Ile Thr Cys Ser Thr Ala Thr Asn Trp Ile Ala Asn Met Ile  
 385 390 395 400

Val Gly Ala Thr Phe Leu Thr Met Leu Asn Thr Leu Gly Asn Ala Asn  
 405 410 415

Thr Phe Trp Val Tyr Ala Ala Leu Asn Val Leu Phe Ile Leu Leu Thr  
 420 425 430

Leu Trp Leu Val Pro Glu Thr Lys His Val Ser Leu Glu His Ile Glu  
 435 440 445

Arg Asn Leu Met Lys Gly Arg Lys Leu Arg Glu Ile Gly Ala His Asp  
 450 455 460

<210> SEQ ID NO 7  
 <211> LENGTH: 1620  
 <212> TYPE: DNA  
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 7

cgcgtaaca atggtttgac cccgcaagaa ctcgaggctt atggtatcag tgacgtacat 60  
 gatatcgttt acaaccaag ctacgacctg ctgtatcagg aagagctcga tccgagcctg 120  
 acaggttatg agcgcggggt gttactaat ctgggtgccg ttgccgtcga taccgggatc 180  
 ttcaccggtc gttcaccaaa agataagtat atcgtccgtg acgataccac tcgcatact 240

-continued

---

```

ttctggtggg cagacaaagg caaaggtaag aacgacaaca aacctctctc tccggaacc 300
tggcagcatc tgaaggcct ggtgaccagg cagctttccg gcaaactctt gttcgttgtc 360
gacgctttct gtggtgcgaa cccggatact cgtctttccg tccgtttcat caccgaagtg 420
gcctggcagg cgcattttgt caaaaacatg tttattcgcc cgagcgatga agaactggca 480
ggtttcaaac cagactttat cgttatgaac ggcgcgaaagt gactaacc gcagtggaaa 540
gaacagggtc tcaactccga aaacttcgtg gcgtttaacc tgaccgagcg catgcagctg 600
attggcggca cctggtacgg cggcgaaatg aagaaaggga tgttctcgat gatgaactac 660
ctgctgccgc tgaaggatc cgcttctatg cactgctccg ccaacgcttg tgagaaaggc 720
gatggtgagg tgttcttcgg cctttccggc accggtaaaa ccacccttc caccgacccg 780
aaacgtgcc tgaattggcg tgacgaacac ggctgggacg atgacggcgt gtttaacttc 840
gaaggcggct gctacgcaa aactatcaag ctgtcgaaag aagcggaaacc tgaaatctac 900
aacgctatcc gtcgtgatgc gttgctggaa aacgtcaccg tgcgtgaaga tggcactatc 960
gactttgatg atggttcaaa aaccgagaac acccgcttt cttatccgat ctatcacatc 1020
gataacattg ttaagccggt ttccaaagcg ggccacgca ctaaggttat cttctgact 1080
gctgatgctt tcggcgtgtt gccgcgggt tctcgctga ctgccgatca aaccagtat 1140
cacttctct ctggttcac cgccaaactg gccgtactg agcgtggcat caccgaaccg 1200
acgccaacct tctccgcttg cttcggcgcg gcattcctgt cgctgcacc gactcagtac 1260
gcagaagtgc tggtgaaacg tatgcagcgc gcggcgcgc aggettatct ggtaaacact 1320
ggctggaacg gactggcaa acgtatctcg attaaagata cccgcgccat tatcgacgcc 1380
atcctcaacg gttcgtgga taatgcagaa accttcactc tgccgatgtt taacctggcg 1440
atcccaaccg aactgccggg cgtagacacg aagattctcg atccgcgtaa cacctacgct 1500
tctccggaac agtggcagga aaaagccgaa acctggcga aactgtttat cgacaacttc 1560
gataaataca ccgacacccc tgcgggtgcc gcgctggtag cggctggtcc gaaactgtaa 1620

```

&lt;210&gt; SEQ ID NO 8

&lt;211&gt; LENGTH: 540

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 8

```

Met Arg Val Asn Asn Gly Leu Thr Pro Gln Glu Leu Glu Ala Tyr Gly
1           5           10           15

Ile Ser Asp Val His Asp Ile Val Tyr Asn Pro Ser Tyr Asp Leu Leu
          20           25           30

Tyr Gln Glu Glu Leu Asp Pro Ser Leu Thr Gly Tyr Glu Arg Gly Val
          35           40           45

Leu Thr Asn Leu Gly Ala Val Ala Val Asp Thr Gly Ile Phe Thr Gly
          50           55           60

Arg Ser Pro Lys Asp Lys Tyr Ile Val Arg Asp Asp Thr Thr Arg Asp
65           70           75           80

Thr Phe Trp Trp Ala Asp Lys Gly Lys Gly Lys Asn Asp Asn Lys Pro
          85           90           95

Leu Ser Pro Glu Thr Trp Gln His Leu Lys Gly Leu Val Thr Arg Gln
          100          105          110

```



-continued

---

515	520	525	
Ala Gly Ala Ala Leu Val Ala Ala Gly Pro Lys Leu			
530	535	540	
<210> SEQ ID NO 9			
<211> LENGTH: 2647			
<212> TYPE: DNA			
<213> ORGANISM: Escherichia coli			
<400> SEQUENCE: 9			
aacgaacaat attccgcatt gcgtagtaat gtcagtatgc tccgcaaagt gctgggagaa			60
accatcaagg atgcgttggg agaacacatt cttgaacgcg tagaaactat ccgtaagttg			120
tcgaaatctt cacgcgctgg caatgatgct aaccgccagg agttgctcac caccttacia			180
aatttgtcga acgacgagct gctgcccgtt gcgctgctgt ttagtcagtt cctgaacctg			240
gccaacaccg ccgagcaata ccacagcatt tcgccgaaag gcgaagctgc cagcaaccgg			300
gaagtgatcg cccgcaccct gcgtaaaactg aaaaaccagc cggaaactgag cgaagacacc			360
atcaaaaaag cagtggaatc gctgtcgtcg gaactggtcc tcacggctca cccaaccgaa			420
attaccgctc gtacactgat ccacaaaatg gtggaagtga acgctctgtt aaaacagctc			480
gataacaaag atatcgctga ctacgaacac aaccagctga tgcgtcgcct gcgccagttg			540
atcgcccagt catggcatac cgatgaaatc cgtaagctgc gtccaagccc ggtagatgaa			600
gccaaatggg gctttgccgt agtggaaaac agcctgtggc aaggcgtacc aaattacctg			660
cgcgaactga acgaacaact ggaagagaac ctccgctaca aactgccctg cgaatttggt			720
ccggtccgtt ttacttcgtg gatggggcgc gaccgcgacg gcaaccggaa cgtcactgcc			780
gatatacccc gccacgtcct gctactcagc cgctggaaaag ccaccgattt gttcctgaaa			840
gatattcagg tgctggtttc tgaactgtcg atggttgaag cgaccctga actgctggcg			900
ctggttggcg aagaaggtgc cgcagaaccg tatcgctatc tgatgaaaaa cctgcttctt			960
cgctgatgg cgacacaggc atggctggaa gcgcgcctga aaggcgaaga actgccaaaa			1020
ccagaaggcc tgctgacaca aaacgaagaa ctgtgggaac cgctctacgc ttgctaccag			1080
tcacttcagg cgtgtggcat gggattatc gccaacggcg atctgctcga caccctgcgc			1140
cgctgaaat gtttcggcgt accgctggtc cgtattgata tccgtcagga gagcacgcgt			1200
cataccgaag cgctgggcga gctgaccgcg tacctcggtg tccggcacta cgaaagctgg			1260
tcagaggccg acaaacaggc gttcctgatc cggaactga actccaaacg tccgcttctg			1320
ccgcgcaact ggcaaccaag cgccgaaacg cgcgaagtgc tcgatacctg ccaggtgatt			1380
gccgaagcac cgcaaggctc cattgccgcc tacgtgatct cgatggcgaa aacgccgtcc			1440
gacgtactgg ctgtccacct gctgctgaaa gaagcgggta tccggtttgc gatgccggtt			1500
gctccgctgt ttgaaaccct cgatgatctg aacaacgcca acgatgtcat gaccagctg			1560
ctcaatattg actggtatcg tggcctgatt cagggcaaac agatggtgat gattggctat			1620
tccgactcag caaaagatgc gggagtgatg gcagcttccg gggcgcaata tcaggcacag			1680
gatgcattaa tcaaacctg cgaaaaagcg ggtattgagc tgacgttgtt ccacggctgc			1740
ggcggttcca ttggctcggg cggcgcacct gctcatgccc cgctgctgtc acaaccgcca			1800
ggaagcctga aaggcggcct gcgcgtaacc gaacagggcg agatgatccg ctttaaatat			1860
ggtctgccag aatcacctg cagcagcctg tcgctttata ccggggcgat tctggaagcc			1920

-continued

---

```

aacctgctgc caccgccgga gccgaaagag agctggcgtc gcattatgga tgaactgtca 1980
gtcatctcct gcgatgtcta ccgcggttac gtacgtgaaa acaaagattt tgtgccttac 2040
ttccgctccg ctacgccgga acaagaactg ggcaaactgc cgttgggttc acgtccggcg 2100
aaacgtogcc caaccggcgg cgtcgagtca ctacgcgcca ttccgtggat ctccgcctgg 2160
acgcaaaacc gtctgatget ccccgctcgg ctgggtgcag gtacggcgct gcaaaaagtg 2220
gtcgaagacg gcaaacagag cgagctggag gctatgtgcc gcgattggcc attcttctcg 2280
acgcgtctcg gcatgctgga gatggctctc gccaaagcag acctgtggct ggcggaatac 2340
tatgaccaac gcctggtaga caaagcactg tggccgtag gtaaagagtt acgcaacctg 2400
caagaagaag acatcaaagt ggtgctggcg attgccaacg attcccatct gatggccgat 2460
ctgccgtgga ttgcagagtc tattcagcta cggaatattt acaccgaccc gctgaacgta 2520
ttgcaggccg agttgctgca ccgctcccgc caggcagaaa aagaaggcca ggaaccggat 2580
cctcgcgtcg aacaagcgtt aatggctact attgccggga ttgcggcagg tatgcgtaat 2640
accggct 2647

```

&lt;210&gt; SEQ ID NO 10

&lt;211&gt; LENGTH: 883

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 10

```

Met Asn Glu Gln Tyr Ser Ala Leu Arg Ser Asn Val Ser Met Leu Gly
1           5           10           15

Lys Val Leu Gly Glu Thr Ile Lys Asp Ala Leu Gly Glu His Ile Leu
20           25           30

Glu Arg Val Glu Thr Ile Arg Lys Leu Ser Lys Ser Ser Arg Ala Gly
35           40           45

Asn Asp Ala Asn Arg Gln Glu Leu Leu Thr Thr Leu Gln Asn Leu Ser
50           55           60

Asn Asp Glu Leu Leu Pro Val Ala Arg Ala Phe Ser Gln Phe Leu Asn
65           70           75           80

Leu Ala Asn Thr Ala Glu Gln Tyr His Ser Ile Ser Pro Lys Gly Glu
85           90           95

Ala Ala Ser Asn Pro Glu Val Ile Ala Arg Thr Leu Arg Lys Leu Lys
100          105          110

Asn Gln Pro Glu Leu Ser Glu Asp Thr Ile Lys Lys Ala Val Glu Ser
115          120          125

Leu Ser Leu Glu Leu Val Leu Thr Ala His Pro Thr Glu Ile Thr Arg
130          135          140

Arg Thr Leu Ile His Lys Met Val Glu Val Asn Ala Cys Leu Lys Gln
145          150          155          160

Leu Asp Asn Lys Asp Ile Ala Asp Tyr Glu His Asn Gln Leu Met Arg
165          170          175

Arg Leu Arg Gln Leu Ile Ala Gln Ser Trp His Thr Asp Glu Ile Arg
180          185          190

Lys Leu Arg Pro Ser Pro Val Asp Glu Ala Lys Trp Gly Phe Ala Val
195          200          205

Val Glu Asn Ser Leu Trp Gln Gly Val Pro Asn Tyr Leu Arg Glu Leu
210          215          220

```

-continued

---

Asn Glu Gln Leu Glu Glu Asn Leu Gly Tyr Lys Leu Pro Val Glu Phe  
 225 230 235 240  
 Val Pro Val Arg Phe Thr Ser Trp Met Gly Gly Asp Arg Asp Gly Asn  
 245 250 255  
 Pro Asn Val Thr Ala Asp Ile Thr Arg His Val Leu Leu Leu Ser Arg  
 260 265 270  
 Trp Lys Ala Thr Asp Leu Phe Leu Lys Asp Ile Gln Val Leu Val Ser  
 275 280 285  
 Glu Leu Ser Met Val Glu Ala Thr Pro Glu Leu Leu Ala Leu Val Gly  
 290 295 300  
 Glu Glu Gly Ala Ala Glu Pro Tyr Arg Tyr Leu Met Lys Asn Leu Arg  
 305 310 315 320  
 Ser Arg Leu Met Ala Thr Gln Ala Trp Leu Glu Ala Arg Leu Lys Gly  
 325 330 335  
 Glu Glu Leu Pro Lys Pro Glu Gly Leu Leu Thr Gln Asn Glu Glu Leu  
 340 345 350  
 Trp Glu Pro Leu Tyr Ala Cys Tyr Gln Ser Leu Gln Ala Cys Gly Met  
 355 360 365  
 Gly Ile Ile Ala Asn Gly Asp Leu Leu Asp Thr Leu Arg Arg Val Lys  
 370 375 380  
 Cys Phe Gly Val Pro Leu Val Arg Ile Asp Ile Arg Gln Glu Ser Thr  
 385 390 395 400  
 Arg His Thr Glu Ala Leu Gly Glu Leu Thr Arg Tyr Leu Gly Ile Gly  
 405 410 415  
 Asp Tyr Glu Ser Trp Ser Glu Ala Asp Lys Gln Ala Phe Leu Ile Arg  
 420 425 430  
 Glu Leu Asn Ser Lys Arg Pro Leu Leu Pro Arg Asn Trp Gln Pro Ser  
 435 440 445  
 Ala Glu Thr Arg Glu Val Leu Asp Thr Cys Gln Val Ile Ala Glu Ala  
 450 455 460  
 Pro Gln Gly Ser Ile Ala Ala Tyr Val Ile Ser Met Ala Lys Thr Pro  
 465 470 475 480  
 Ser Asp Val Leu Ala Val His Leu Leu Leu Lys Glu Ala Gly Ile Gly  
 485 490 495  
 Phe Ala Met Pro Val Ala Pro Leu Phe Glu Thr Leu Asp Asp Leu Asn  
 500 505 510  
 Asn Ala Asn Asp Val Met Thr Gln Leu Leu Asn Ile Asp Trp Tyr Arg  
 515 520 525  
 Gly Leu Ile Gln Gly Lys Gln Met Val Met Ile Gly Tyr Ser Asp Ser  
 530 535 540  
 Ala Lys Asp Ala Gly Val Met Ala Ala Ser Trp Ala Gln Tyr Gln Ala  
 545 550 555 560  
 Gln Asp Ala Leu Ile Lys Thr Cys Glu Lys Ala Gly Ile Glu Leu Thr  
 565 570 575  
 Leu Phe His Gly Arg Gly Gly Ser Ile Gly Arg Gly Gly Ala Pro Ala  
 580 585 590  
 His Ala Ala Leu Leu Ser Gln Pro Pro Gly Ser Leu Lys Gly Gly Leu  
 595 600 605  
 Arg Val Thr Glu Gln Gly Glu Met Ile Arg Phe Lys Tyr Gly Leu Pro  
 610 615 620



-continued

---

Glu Ile Thr Val Ser Ser Leu Ser Leu Tyr Thr Gly Ala Ile Leu Glu  
 625 630 635 640  
 Ala Asn Leu Leu Pro Pro Pro Glu Pro Lys Glu Ser Trp Arg Arg Ile  
 645 650 655  
 Met Asp Glu Leu Ser Val Ile Ser Cys Asp Val Tyr Arg Gly Tyr Val  
 660 665 670  
 Arg Glu Asn Lys Asp Phe Val Pro Tyr Phe Arg Ser Ala Thr Pro Glu  
 675 680 685  
 Gln Glu Leu Gly Lys Leu Pro Leu Gly Ser Arg Pro Ala Lys Arg Arg  
 690 695 700  
 Pro Thr Gly Gly Val Glu Ser Leu Arg Ala Ile Pro Trp Ile Phe Ala  
 705 710 715 720  
 Trp Thr Gln Asn Arg Leu Met Leu Pro Ala Trp Leu Gly Ala Gly Thr  
 725 730 735  
 Ala Leu Gln Lys Val Val Glu Asp Gly Lys Gln Ser Glu Leu Glu Ala  
 740 745 750  
 Met Cys Arg Asp Trp Pro Phe Phe Ser Thr Arg Leu Gly Met Leu Glu  
 755 760 765  
 Met Val Phe Ala Lys Ala Asp Leu Trp Leu Ala Glu Tyr Tyr Asp Gln  
 770 775 780  
 Arg Leu Val Asp Lys Ala Leu Trp Pro Leu Gly Lys Glu Leu Arg Asn  
 785 790 795 800  
 Leu Gln Glu Glu Asp Ile Lys Val Val Leu Ala Ile Ala Asn Asp Ser  
 805 810 815  
 His Leu Met Ala Asp Leu Pro Trp Ile Ala Glu Ser Ile Gln Leu Arg  
 820 825 830  
 Asn Ile Tyr Thr Asp Pro Leu Asn Val Leu Gln Ala Glu Leu Leu His  
 835 840 845  
 Arg Ser Arg Gln Ala Glu Lys Glu Gly Gln Glu Pro Asp Pro Arg Val  
 850 855 860  
 Glu Gln Ala Leu Met Val Thr Ile Ala Gly Ile Ala Ala Gly Met Arg  
 865 870 875 880  
 Asn Thr Gly

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 2661

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 11

```

tcagaacggt tcccaaatga cgtggatccg atcgaaactc gcgactggct ccaggcgatc    60
gaatcgggtca tccgtgaaga aggtggtgag cgtgctcagt atctgatcga ccaactgctt    120
gctgaagccc gcaaaggcgg tgtaaacgta gccgcaggca caggtatcag caactacatc    180
aacaccatcc ccgttgaaga acaaccggag tatccgggta atctggaact ggaacgccgt    240
attcgttcag ctatccgctg gaacgccatc atgacgggtc tgcgtgcgtc gaaaaaagac    300
ctcgaactgg gcggccatat ggcgtccttc cagtcttccg caaccattta tgatgtgtgc    360
tttaaccact tcttccgtgc acgcaacgag caggatggcg gcgacctggt ttacttcag    420
ggccacatct cccggggcgt gtacgctcgt gctttcctgg aaggtegtet gactcaggag    480
cagctggata acttccgtca ggaagttcac ggcaatggcc tctcttccta tccgcaccgg    540

```

-continued

---

aaactgatgc	cggattctg	gcagttcccg	accgtatcta	tgggtctggg	tccgattggt	600
gctathtacc	aggctaaatt	cctgaaatat	ctggaacacc	gtggcctgaa	agatacctct	660
aaacaaaccg	tttacgcggt	cctcgggtgac	ggtgaaatgg	acgaaccgga	atccaaaggt	720
gcgatcacca	tcgctaccgg	tgaaaaactg	gataacctgg	tcttcggtat	caactgtaac	780
ctgcagcgtc	ttgacggccc	ggtcaccggg	aacggcaaga	tcatcaacga	actggaaggc	840
atcttcgaag	gtgctggctg	gaacgtgatc	aaagtgatgt	ggggtagccg	ttgggatgaa	900
ctgctgcgta	aggataccag	cggtaaactg	atccagctga	tgaacgaaac	cggtgacggc	960
gactaccaga	ccttcaaate	gaaagatggg	gcgtacgttc	gtgaacactt	cttcggtaaa	1020
tatcctgaaa	ccgcagcact	ggttgcagac	tggactgacg	agcagatctg	ggcactgaac	1080
cgtggtggtc	acgatccgaa	gaaaatctac	gctgcattca	agaaagcgca	ggaaacccaa	1140
ggcaaagcga	cagtaatcct	tgctcatacc	attaaagggt	acggcatggg	cgacgcggct	1200
gaaggtaaaa	acatcgcgca	ccaggttaag	aaaatgaaca	tggacgggtg	gcgtcatatc	1260
cgcgaccggt	tcaatgtgcc	ggtgtctgat	gcagatatcg	aaaaactgcc	gtacatcacc	1320
ttcccgaag	gttctgaaga	gcatacctat	ctgcacgctc	agcgtcagaa	actgcacggg	1380
tatctgccc	gccgtcagcc	gaacttcacc	gagaagcttg	agctgccgag	cctgcaagac	1440
ttcggcgcgc	tggtggaaga	gcagagcaaa	gagatctcta	ccactatcgc	tttcggtcgt	1500
gctctgaacg	tgatgctgaa	gaacaagtcg	atcaaagatc	gtctggtacc	gatcatcgcc	1560
gacgaagcgc	gtactttcgg	tatggaaggt	ctgttccgtc	agattggtat	ttacagcccg	1620
aacggtcagc	agtacacccc	gcaggaccgc	gagcagggtg	cttactataa	agaagacgag	1680
aaaggtcaga	ttctgcagga	agggatcaac	gagctgggcg	caggttgttc	ctggctggca	1740
gcggcgacct	cttacagcac	caacaatctg	ccgatgatcc	cgttctacat	ctattactcg	1800
atgttcggct	tccagcgtat	tggcgatctg	tgctgggccc	ctggcgacca	gcaagcgcgt	1860
ggcttctcga	tcggcggtac	ttccggtcgt	accaccctga	acggcgaagg	tctgcagcac	1920
gaagatggtc	acagccacat	tcagtcgctg	actatcccga	actgtatctc	ttacgacccc	1980
gcttacgctt	acgaagttgc	tgatcatcatg	catgacggtc	tggagcgtat	gtacggtgaa	2040
aaacaagaga	acgtttacta	ctacatcact	acgctgaacg	aaaactacca	catgccggca	2100
atgccggaag	gtgctgagga	aggatccggt	aaaggatatct	acaaactcga	aactattgaa	2160
ggtagcaaaag	gtaaagttca	gctgctcggc	tccggttcta	tcttcgctca	cgctccgtgaa	2220
gcagctgaga	tcctggcgaa	agattacggc	gtaggttctg	acgtttatag	cgtgacctcc	2280
ttcaccgagc	tggcgcgtga	tggtcaggat	tgtgaaacgct	ggaacatgct	gcacccgctg	2340
gaaactccgc	gcgttccgta	tatcgctcag	gtgatgaacg	acgctccggc	agtggcatct	2400
accgactata	tgaaactggt	cgctgagcag	gtccgtactt	acgtaccggc	tgacgactac	2460
cgcgtactgg	gtactgatgg	cttcggtcgt	tccgacagcc	gtgagaacct	gcgtcaccac	2520
ttcgaagttg	atgcttctta	tgctcgtggt	gcggcgtcgg	gcgaactggc	taaactgggc	2580
gaaatcgata	agaaagtggg	tgctgacgca	atcgccaaat	tcaacatcga	tgacagataaa	2640
gttaacccgc	gtctggcgta	a				2661

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 887

&lt;212&gt; TYPE: PRT

-continued

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 12

```

Met Ser Glu Arg Phe Pro Asn Asp Val Asp Pro Ile Glu Thr Arg Asp
1          5          10          15
Trp Leu Gln Ala Ile Glu Ser Val Ile Arg Glu Glu Gly Val Glu Arg
20          25          30
Ala Gln Tyr Leu Ile Asp Gln Leu Leu Ala Glu Ala Arg Lys Gly Gly
35          40          45
Val Asn Val Ala Ala Gly Thr Gly Ile Ser Asn Tyr Ile Asn Thr Ile
50          55          60
Pro Val Glu Glu Gln Pro Glu Tyr Pro Gly Asn Leu Glu Leu Glu Arg
65          70          75          80
Arg Ile Arg Ser Ala Ile Arg Trp Asn Ala Ile Met Thr Val Leu Arg
85          90          95
Ala Ser Lys Lys Asp Leu Glu Leu Gly Gly His Met Ala Ser Phe Gln
100         105         110
Ser Ser Ala Thr Ile Tyr Asp Val Cys Phe Asn His Phe Phe Arg Ala
115         120         125
Arg Asn Glu Gln Asp Gly Gly Asp Leu Val Tyr Phe Gln Gly His Ile
130         135         140
Ser Pro Gly Val Tyr Ala Arg Ala Phe Leu Glu Gly Arg Leu Thr Gln
145         150         155         160
Glu Gln Leu Asp Asn Phe Arg Gln Glu Val His Gly Asn Gly Leu Ser
165         170         175
Ser Tyr Pro His Pro Lys Leu Met Pro Glu Phe Trp Gln Phe Pro Thr
180         185         190
Val Ser Met Gly Leu Gly Pro Ile Gly Ala Ile Tyr Gln Ala Lys Phe
195         200         205
Leu Lys Tyr Leu Glu His Arg Gly Leu Lys Asp Thr Ser Lys Gln Thr
210         215         220
Val Tyr Ala Phe Leu Gly Asp Gly Glu Met Asp Glu Pro Glu Ser Lys
225         230         235         240
Gly Ala Ile Thr Ile Ala Thr Arg Glu Lys Leu Asp Asn Leu Val Phe
245         250         255
Val Ile Asn Cys Asn Leu Gln Arg Leu Asp Gly Pro Val Thr Gly Asn
260         265         270
Gly Lys Ile Ile Asn Glu Leu Glu Gly Ile Phe Glu Gly Ala Gly Trp
275         280         285
Asn Val Ile Lys Val Met Trp Gly Ser Arg Trp Asp Glu Leu Leu Arg
290         295         300
Lys Asp Thr Ser Gly Lys Leu Ile Gln Leu Met Asn Glu Thr Val Asp
305         310         315         320
Gly Asp Tyr Gln Thr Phe Lys Ser Lys Asp Gly Ala Tyr Val Arg Glu
325         330         335
His Phe Phe Gly Lys Tyr Pro Glu Thr Ala Ala Leu Val Ala Asp Trp
340         345         350
Thr Asp Glu Gln Ile Trp Ala Leu Asn Arg Gly Gly His Asp Pro Lys
355         360         365
Lys Ile Tyr Ala Ala Phe Lys Lys Ala Gln Glu Thr Lys Gly Lys Ala
370         375         380

```

-continued

---

Thr	Val	Ile	Leu	Ala	His	Thr	Ile	Lys	Gly	Tyr	Gly	Met	Gly	Asp	Ala
385					390					395					400
Ala	Glu	Gly	Lys	Asn	Ile	Ala	His	Gln	Val	Lys	Lys	Met	Asn	Met	Asp
				405					410					415	
Gly	Val	Arg	His	Ile	Arg	Asp	Arg	Phe	Asn	Val	Pro	Val	Ser	Asp	Ala
			420					425					430		
Asp	Ile	Glu	Lys	Leu	Pro	Tyr	Ile	Thr	Phe	Pro	Glu	Gly	Ser	Glu	Glu
		435					440					445			
His	Thr	Tyr	Leu	His	Ala	Gln	Arg	Gln	Lys	Leu	His	Gly	Tyr	Leu	Pro
	450					455					460				
Ser	Arg	Gln	Pro	Asn	Phe	Thr	Glu	Lys	Leu	Glu	Leu	Pro	Ser	Leu	Gln
465					470					475					480
Asp	Phe	Gly	Ala	Leu	Leu	Glu	Glu	Gln	Ser	Lys	Glu	Ile	Ser	Thr	Thr
				485					490					495	
Ile	Ala	Phe	Val	Arg	Ala	Leu	Asn	Val	Met	Leu	Lys	Asn	Lys	Ser	Ile
			500					505					510		
Lys	Asp	Arg	Leu	Val	Pro	Ile	Ile	Ala	Asp	Glu	Ala	Arg	Thr	Phe	Gly
		515					520						525		
Met	Glu	Gly	Leu	Phe	Arg	Gln	Ile	Gly	Ile	Tyr	Ser	Pro	Asn	Gly	Gln
	530					535					540				
Gln	Tyr	Thr	Pro	Gln	Asp	Arg	Glu	Gln	Val	Ala	Tyr	Tyr	Lys	Glu	Asp
545					550					555					560
Glu	Lys	Gly	Gln	Ile	Leu	Gln	Glu	Gly	Ile	Asn	Glu	Leu	Gly	Ala	Gly
				565					570					575	
Cys	Ser	Trp	Leu	Ala	Ala	Ala	Thr	Ser	Tyr	Ser	Thr	Asn	Asn	Leu	Pro
			580					585					590		
Met	Ile	Pro	Phe	Tyr	Ile	Tyr	Tyr	Ser	Met	Phe	Gly	Phe	Gln	Arg	Ile
		595					600					605			
Gly	Asp	Leu	Cys	Trp	Ala	Ala	Gly	Asp	Gln	Gln	Ala	Arg	Gly	Phe	Leu
	610					615					620				
Ile	Gly	Gly	Thr	Ser	Gly	Arg	Thr	Thr	Leu	Asn	Gly	Glu	Gly	Leu	Gln
625					630					635					640
His	Glu	Asp	Gly	His	Ser	His	Ile	Gln	Ser	Leu	Thr	Ile	Pro	Asn	Cys
				645					650					655	
Ile	Ser	Tyr	Asp	Pro	Ala	Tyr	Ala	Tyr	Glu	Val	Ala	Val	Ile	Met	His
			660					665					670		
Asp	Gly	Leu	Glu	Arg	Met	Tyr	Gly	Glu	Lys	Gln	Glu	Asn	Val	Tyr	Tyr
		675					680					685			
Tyr	Ile	Thr	Thr	Leu	Asn	Glu	Asn	Tyr	His	Met	Pro	Ala	Met	Pro	Glu
	690					695					700				
Gly	Ala	Glu	Glu	Gly	Ile	Arg	Lys	Gly	Ile	Tyr	Lys	Leu	Glu	Thr	Ile
705					710					715					720
Glu	Gly	Ser	Lys	Gly	Lys	Val	Gln	Leu	Leu	Gly	Ser	Gly	Ser	Ile	Leu
				725					730					735	
Arg	His	Val	Arg	Glu	Ala	Ala	Glu	Ile	Leu	Ala	Lys	Asp	Tyr	Gly	Val
			740					745					750		
Gly	Ser	Asp	Val	Tyr	Ser	Val	Thr	Ser	Phe	Thr	Glu	Leu	Ala	Arg	Asp
		755					760					765			
Gly	Gln	Asp	Cys	Glu	Arg	Trp	Asn	Met	Leu	His	Pro	Leu	Glu	Thr	Pro
	770					775					780				
Arg	Val	Pro	Tyr	Ile	Ala	Gln	Val	Met	Asn	Asp	Ala	Pro	Ala	Val	Ala

-continued

785	790	795	800
Ser Thr Asp Tyr Met Lys Leu Phe Ala Glu Gln Val Arg Thr Tyr Val	805	810	815
Pro Ala Asp Asp Tyr Arg Val Leu Gly Thr Asp Gly Phe Gly Arg Ser	820	825	830
Asp Ser Arg Glu Asn Leu Arg His His Phe Glu Val Asp Ala Ser Tyr	835	840	845
Val Val Val Ala Ala Leu Gly Glu Leu Ala Lys Arg Gly Glu Ile Asp	850	855	860
Lys Lys Val Val Ala Asp Ala Ile Ala Lys Phe Asn Ile Asp Ala Asp	865	870	875
Lys Val Asn Pro Arg Leu Ala	885		

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 1887

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 13

```

gctatcgaaa tcaaagtacc ggacatcggg gctgatgaag ttgaaatcac cgagatcctg    60
gtcaaagtgg gcgacaaagt tgaagccgaa cagtcgctga tcaccgtaga aggcgacaaa    120
gcctctatgg aagttccgtc tccgcaggcg ggtatcgтта aagagatcaa agtctctggt    180
ggcgataaaa cccagaccgg cgcactgatt atgattttcg attccgccga cggtgagca    240
gacgctgcac ctgctcaggc agaagagaag aaagaagcag ctccggcagc agcaccagcg    300
gctgcggcgg caaaagacgt taacgttccg gatatcggca gcgacgaagt tgaagtgacc    360
gaaatcctgg tgaaagttgg cgataaagtt gaagctgaac agtcgctgat caccgtagaa    420
ggcgacaagg cttctatgga agttccggct ccgттtгctg gcaccgtgaa agagatcaaa    480
gtgaacgtgg gtgacaaagt gtctaccggc tcgctgatta tggтcttcga agtcgcgggт    540
gaagcaggcg cggcagctcc ggccgctaaa caggaagcag ctccggcagc ggccccгca    600
ccagcggctg gcgtgaaaga agttaacgtt ccggatatcg gcggtgacga agттgaaгtг    660
actgaagtga tggтgaaagt gggcgacaaa gттccгctg aacagтcact gatcaccgта    720
gaaggcgaca aagctтctat ggaagттccg gcgccгттg caggcгtсгt gaaggaactg    780
aaagtcaacg ttggcgataa agtgaaaact ggctcгctga ttatgatctt cgaagттgaa    840
ggcgcagcgc ctgcgгcagc тcctгcгaaa caggaagcгg cagcгccгgc accгgгcagca    900
aaagctgaag ccccггcagc agcaccagct gcgaaagcгg aaggcaaatc tgaattгct    960
gaaaacgacg cttatгттca cгcгactccg ctgatccгcc gtctггcagc cгagттгггт    1020
gttaacctгt cгaaagtгaa gggcactггc cгtaaagгtc gtatcctгcг cгaagacгтт    1080
cagгcttacg tgaaagaagc tatcaaacгt gcagaagcag ctccггcagc gactггcггт    1140
ggtatccctg гcatгctгcc gtggccgaag gtгgactтca gcaagттггt tgaaatcгaa    1200
gaagtгgaaс tggгccгcat ccagaaaatc тctггтгcга acctгagccг таactгггта    1260
atgatccccгc atгттactca cттcгacaaa accgatatca ccгagттгга agcгттccгт    1320
aaacagcaga acgaaгaagc ggcgaaacгt aagctггatg tgaagatcac cccггттгtc    1380
ttcatcatga aagccгттгc тgcagctctt gagcagatgc ctгcгттcaа tagттcгctg    1440

```

-continued

---

```

tcggaagacg gtcagcgtct gaccctgaag aaatacatca acatcggtgt ggcgggtggat 1500
accccgaaacg gtctggttgt tccggtattc aaagacgtca acaagaaagg catcatcgag 1560
ctgtctcgcg agctgatgac tatttctaag aaagcgcgtg acggtaagct gactgcgggc 1620
gaaatgcagg gcggttgctt caccatctcc agcatcggcg gcctgggtac taccacttc 1680
gcgcccattg tgaacgcgcc ggaagtggct atcctcggcg tttccaagtc cgcgatggag 1740
ccggtgtgga atggtaaaga gttcgtgccg cgtctgatgc tgccgatttc tctctcttc 1800
gaccaccgcg tgatcgacgg tgetgatggt gcccgtttca ttaccatcat taacaacacg 1860
ctgtctgaca ttcgccgtct ggtgatg 1887

```

```

<210> SEQ ID NO 14
<211> LENGTH: 630
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli

```

```

<400> SEQUENCE: 14

```

```

Met Ala Ile Glu Ile Lys Val Pro Asp Ile Gly Ala Asp Glu Val Glu
1           5           10           15
Ile Thr Glu Ile Leu Val Lys Val Gly Asp Lys Val Glu Ala Glu Gln
          20           25           30
Ser Leu Ile Thr Val Glu Gly Asp Lys Ala Ser Met Glu Val Pro Ser
          35           40           45
Pro Gln Ala Gly Ile Val Lys Glu Ile Lys Val Ser Val Gly Asp Lys
          50           55           60
Thr Gln Thr Gly Ala Leu Ile Met Ile Phe Asp Ser Ala Asp Gly Ala
65           70           75           80
Ala Asp Ala Ala Pro Ala Gln Ala Glu Glu Lys Lys Glu Ala Ala Pro
          85           90           95
Ala Ala Ala Pro Ala Ala Ala Ala Ala Lys Asp Val Asn Val Pro Asp
          100          105          110
Ile Gly Ser Asp Glu Val Glu Val Thr Glu Ile Leu Val Lys Val Gly
          115          120          125
Asp Lys Val Glu Ala Glu Gln Ser Leu Ile Thr Val Glu Gly Asp Lys
          130          135          140
Ala Ser Met Glu Val Pro Ala Pro Phe Ala Gly Thr Val Lys Glu Ile
145          150          155          160
Lys Val Asn Val Gly Asp Lys Val Ser Thr Gly Ser Leu Ile Met Val
          165          170          175
Phe Glu Val Ala Gly Glu Ala Gly Ala Ala Ala Pro Ala Ala Lys Gln
          180          185          190
Glu Ala Ala Pro Ala Ala Ala Pro Ala Pro Ala Ala Gly Val Lys Glu
          195          200          205
Val Asn Val Pro Asp Ile Gly Gly Asp Glu Val Glu Val Thr Glu Val
          210          215          220
Met Val Lys Val Gly Asp Lys Val Ala Ala Glu Gln Ser Leu Ile Thr
225          230          235          240
Val Glu Gly Asp Lys Ala Ser Met Glu Val Pro Ala Pro Phe Ala Gly
          245          250          255
Val Val Lys Glu Leu Lys Val Asn Val Gly Asp Lys Val Lys Thr Gly
          260          265          270
Ser Leu Ile Met Ile Phe Glu Val Glu Gly Ala Ala Pro Ala Ala Ala

```

-continued

275					280					285					
Pro	Ala	Lys	Gln	Glu	Ala	Ala	Ala	Pro	Ala	Pro	Ala	Ala	Lys	Ala	Glu
290					295					300					
Ala	Pro	Ala	Ala	Ala	Pro	Ala	Ala	Lys	Ala	Glu	Gly	Lys	Ser	Glu	Phe
305					310					315					320
Ala	Glu	Asn	Asp	Ala	Tyr	Val	His	Ala	Thr	Pro	Leu	Ile	Arg	Arg	Leu
				325					330					335	
Ala	Arg	Glu	Phe	Gly	Val	Asn	Leu	Ala	Lys	Val	Lys	Gly	Thr	Gly	Arg
				340				345						350	
Lys	Gly	Arg	Ile	Leu	Arg	Glu	Asp	Val	Gln	Ala	Tyr	Val	Lys	Glu	Ala
				355				360						365	
Ile	Lys	Arg	Ala	Glu	Ala	Ala	Pro	Ala	Ala	Thr	Gly	Gly	Gly	Ile	Pro
				370				375						380	
Gly	Met	Leu	Pro	Trp	Pro	Lys	Val	Asp	Phe	Ser	Lys	Phe	Gly	Glu	Ile
385					390					395					400
Glu	Glu	Val	Glu	Leu	Gly	Arg	Ile	Gln	Lys	Ile	Ser	Gly	Ala	Asn	Leu
				405					410					415	
Ser	Arg	Asn	Trp	Val	Met	Ile	Pro	His	Val	Thr	His	Phe	Asp	Lys	Thr
				420					425					430	
Asp	Ile	Thr	Glu	Leu	Glu	Ala	Phe	Arg	Lys	Gln	Gln	Asn	Glu	Glu	Ala
				435					440					445	
Ala	Lys	Arg	Lys	Leu	Asp	Val	Lys	Ile	Thr	Pro	Val	Val	Phe	Ile	Met
				450					455					460	
Lys	Ala	Val	Ala	Ala	Ala	Leu	Glu	Gln	Met	Pro	Arg	Phe	Asn	Ser	Ser
465					470					475					480
Leu	Ser	Glu	Asp	Gly	Gln	Arg	Leu	Thr	Leu	Lys	Lys	Tyr	Ile	Asn	Ile
				485					490					495	
Gly	Val	Ala	Val	Asp	Thr	Pro	Asn	Gly	Leu	Val	Val	Pro	Val	Phe	Lys
				500					505					510	
Asp	Val	Asn	Lys	Lys	Gly	Ile	Ile	Glu	Leu	Ser	Arg	Glu	Leu	Met	Thr
				515					520					525	
Ile	Ser	Lys	Lys	Ala	Arg	Asp	Gly	Lys	Leu	Thr	Ala	Gly	Glu	Met	Gln
				530					535					540	
Gly	Gly	Cys	Phe	Thr	Ile	Ser	Ser	Ile	Gly	Gly	Leu	Gly	Thr	Thr	His
545					550					555					560
Phe	Ala	Pro	Ile	Val	Asn	Ala	Pro	Glu	Val	Ala	Ile	Leu	Gly	Val	Ser
				565					570					575	
Lys	Ser	Ala	Met	Glu	Pro	Val	Trp	Asn	Gly	Lys	Glu	Phe	Val	Pro	Arg
				580					585					590	
Leu	Met	Leu	Pro	Ile	Ser	Leu	Ser	Phe	Asp	His	Arg	Val	Ile	Asp	Gly
				595					600					605	
Ala	Asp	Gly	Ala	Arg	Phe	Ile	Thr	Ile	Ile	Asn	Asn	Thr	Leu	Ser	Asp
				610										615	
Ile	Arg	Arg	Leu	Val	Met										
625					630										

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 1422

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 15

-continued

---

```

agtactgaaa tcaaaactca ggtcgtggta cttggggcag gccccgcagg ttactccgct    60
gccttccggt ggcctgattt aggtctggaa accgtaatcg tagaacgta caacaccctt    120
ggcggtggtt gcctgaacgt cggctgtatc ctttctaaag cactgctgca cgtagcaaaa    180
gttatcgaag aagccaaagc gctggctgaa cacggtatcg tcttcggcga accgaaaacc    240
gatatcgaca agattcgtac ctggaaagag aaagtgatca atcagctgac cggtggtctg    300
gctggtatgg cgaaaggccg caaagtcaaa gtggtcaacg gtctgggtaa attcaccggg    360
gctaacaccc tggaagtga aggtgagaac ggcaaaaccg tgatcaactt cgacaacgcg    420
atcattgcag cgggttctcg cccgatccaa ctgccgttta ttccgcatga agatccgcgt    480
atctgggact cactgacgc gctggaactg aaagaagtac cagaacgcct gctggtaatg    540
ggtggcggtg tcatcgggtc ggaaatgggc accgtttacc acgcgctggg ttcacagatt    600
gacgtggttg aatgttcga ccaggttatc ccggcagctg acaaagacat cgttaaagtc    660
ttaccaagc gtatcagcaa gaaattcaac ctgatgctgg aaaccaagt taccgccgtt    720
gaagcgaaag aagacggcat ttatgtgacg atggaaggca aaaaagcacc cgctgaaccg    780
cagcgttacg acgccgtgct ggtagcgatt ggtcgtgtgc cgaacggtaa aaacctcgac    840
gcaggcaaag caggcgtgga agttgacgac cgtggtttca tccgcgttga caaacagctg    900
cgtaccaacg taccgcacat ctttgctatc ggcgatatcg tcggtcaacc gatgctggca    960
caciaaagtg ttcacgaagg tcacgttgcc gctgaagtta tcgccggtaa gaaacactac   1020
ttcgatccga aagttatccc gtccatcgcc tataccgaac cagaagttgc atgggtgggt   1080
ctgactgaga aagaagcgaa agagaaaggc atcagctatg aaaccgccac cttcccgtag   1140
gctgcttctg gtcgtgctat cgcttccgac tgcgcagacg gtatgaccaa gctgattttc   1200
gacaaagaat ctcaccgtgt gatcggtggt gcgattgtcg gtactaacgg cggcgagctg   1260
ctgggtgaaa tcggcctggc aatcgaaatg ggtgtgatg ctgaagacat cgcactgacc   1320
atccacgcgc acccgactct gcacgagtct gtggcctgg cggcagaagt gttcgaaggt   1380
agcattaccg acctgccgaa cccgaaagcg aagaagaagt aa                          1422

```

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 474

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 16

```

Met Ser Thr Glu Ile Lys Thr Gln Val Val Val Leu Gly Ala Gly Pro
1           5           10           15

Ala Gly Tyr Ser Ala Ala Phe Arg Cys Ala Asp Leu Gly Leu Glu Thr
20           25           30

Val Ile Val Glu Arg Tyr Asn Thr Leu Gly Gly Val Cys Leu Asn Val
35           40           45

Gly Cys Ile Pro Ser Lys Ala Leu Leu His Val Ala Lys Val Ile Glu
50           55           60

Glu Ala Lys Ala Leu Ala Glu His Gly Ile Val Phe Gly Glu Pro Lys
65           70           75           80

Thr Asp Ile Asp Lys Ile Arg Thr Trp Lys Glu Lys Val Ile Asn Gln
85           90           95

Leu Thr Gly Gly Leu Ala Gly Met Ala Lys Gly Arg Lys Val Lys Val
100          105          110

```



-continued

---

Val Asn Gly Leu Gly Lys Phe Thr Gly Ala Asn Thr Leu Glu Val Glu  
 115 120 125  
 Gly Glu Asn Gly Lys Thr Val Ile Asn Phe Asp Asn Ala Ile Ile Ala  
 130 135 140  
 Ala Gly Ser Arg Pro Ile Gln Leu Pro Phe Ile Pro His Glu Asp Pro  
 145 150 155 160  
 Arg Ile Trp Asp Ser Thr Asp Ala Leu Glu Leu Lys Glu Val Pro Glu  
 165 170 175  
 Arg Leu Leu Val Met Gly Gly Gly Ile Ile Gly Leu Glu Met Gly Thr  
 180 185 190  
 Val Tyr His Ala Leu Gly Ser Gln Ile Asp Val Val Glu Met Phe Asp  
 195 200 205  
 Gln Val Ile Pro Ala Ala Asp Lys Asp Ile Val Lys Val Phe Thr Lys  
 210 215 220  
 Arg Ile Ser Lys Lys Phe Asn Leu Met Leu Glu Thr Lys Val Thr Ala  
 225 230 235 240  
 Val Glu Ala Lys Glu Asp Gly Ile Tyr Val Thr Met Glu Gly Lys Lys  
 245 250 255  
 Ala Pro Ala Glu Pro Gln Arg Tyr Asp Ala Val Leu Val Ala Ile Gly  
 260 265 270  
 Arg Val Pro Asn Gly Lys Asn Leu Asp Ala Gly Lys Ala Gly Val Glu  
 275 280 285  
 Val Asp Asp Arg Gly Phe Ile Arg Val Asp Lys Gln Leu Arg Thr Asn  
 290 295 300  
 Val Pro His Ile Phe Ala Ile Gly Asp Ile Val Gly Gln Pro Met Leu  
 305 310 315 320  
 Ala His Lys Gly Val His Glu Gly His Val Ala Ala Glu Val Ile Ala  
 325 330 335  
 Gly Lys Lys His Tyr Phe Asp Pro Lys Val Ile Pro Ser Ile Ala Tyr  
 340 345 350  
 Thr Glu Pro Glu Val Ala Trp Val Gly Leu Thr Glu Lys Glu Ala Lys  
 355 360 365  
 Glu Lys Gly Ile Ser Tyr Glu Thr Ala Thr Phe Pro Trp Ala Ala Ser  
 370 375 380  
 Gly Arg Ala Ile Ala Ser Asp Cys Ala Asp Gly Met Thr Lys Leu Ile  
 385 390 395 400  
 Phe Asp Lys Glu Ser His Arg Val Ile Gly Gly Ala Ile Val Gly Thr  
 405 410 415  
 Asn Gly Gly Glu Leu Leu Gly Glu Ile Gly Leu Ala Ile Glu Met Gly  
 420 425 430  
 Cys Asp Ala Glu Asp Ile Ala Leu Thr Ile His Ala His Pro Thr Leu  
 435 440 445  
 His Glu Ser Val Gly Leu Ala Ala Glu Val Phe Glu Gly Ser Ile Thr  
 450 455 460  
 Asp Leu Pro Asn Pro Lys Ala Lys Lys Lys  
 465 470

&lt;210&gt; SEQ ID NO 17

&lt;211&gt; LENGTH: 987

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

-continued

&lt;400&gt; SEQUENCE: 17

```

aaactcgccg tttatagcac aaaacagtac gacaagaagt acctgcaaca ggtgaacgag    60
tcctttggct ttgagctgga attttttgac tttctgctga cggaaaaaac cgctaaaact    120
gccaatggct gcgaagcggc atgtattttc gtaaacgatg acggcagccg cccgggtgctg    180
gaagagctga aaaagcacgg cgtaaataat atcgccctgc gctgtgcccg tttcaataac    240
gtcgaccttg acgcgggcaaa agaactgggg ctgaaagtag tccgtgttcc agcctatgat    300
ccagaggccg ttgctgaaca cgccatcggc atgatgatga cgctgaaccg ccgtattcac    360
cgcgcgatc agcgtaccgg tgatgctaac ttctctctgg aaggctctgac cggctttact    420
atgtatggca aaacggcagg cgttatcggc accggtaaaa tcggtgtggc gatgctgcgc    480
attctgaaag gttttggtat gcgtctgctg gcgttcgatc cgtatccaag tgcagcggcg    540
ctggaactcg gtgtggagta tgctgatctg ccaaccctgt tctctgaatc agacgttatc    600
tctctgcaat gcccgtgac accggaaaaac tatcatctgt tgaacgaagc cgcttcgaa    660
cagatgaaaa atggcgtgat gatcgtcaat accagtcgcg gtgcattgat tgattctcag    720
gcagcaattg aagcgtgaa aatcagaaa attggttcgt tgggtatgga cgtgtatgag    780
aacgaacgag atctattctt tgaagataaa tccaacgacg tgatccagga tgacgtattc    840
cgtcgctgt ctgctgcca caacgtgctg ttaccggggc accaggcatt cctgacagca    900
gaagctctga ccagtatttc tcagactacg ctgcaaaact taagcaatct ggaaaaaggc    960
gaaacctgcc cgaacgaact ggtttaa                                     987

```

&lt;210&gt; SEQ ID NO 18

&lt;211&gt; LENGTH: 329

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 18

```

Met Lys Leu Ala Val Tyr Ser Thr Lys Gln Tyr Asp Lys Lys Tyr Leu
 1           5           10           15

Gln Gln Val Asn Glu Ser Phe Gly Phe Glu Leu Glu Phe Phe Asp Phe
 20           25           30

Leu Leu Thr Glu Lys Thr Ala Lys Thr Ala Asn Gly Cys Glu Ala Val
 35           40           45

Cys Ile Phe Val Asn Asp Asp Gly Ser Arg Pro Val Leu Glu Glu Leu
 50           55           60

Lys Lys His Gly Val Lys Tyr Ile Ala Leu Arg Cys Ala Gly Phe Asn
 65           70           75           80

Asn Val Asp Leu Asp Ala Ala Lys Glu Leu Gly Leu Lys Val Val Arg
 85           90           95

Val Pro Ala Tyr Asp Pro Glu Ala Val Ala Glu His Ala Ile Gly Met
 100          105          110

Met Met Thr Leu Asn Arg Arg Ile His Arg Ala Tyr Gln Arg Thr Arg
 115          120          125

Asp Ala Asn Phe Ser Leu Glu Gly Leu Thr Gly Phe Thr Met Tyr Gly
 130          135          140

Lys Thr Ala Gly Val Ile Gly Thr Gly Lys Ile Gly Val Ala Met Leu
 145          150          155          160

Arg Ile Leu Lys Gly Phe Gly Met Arg Leu Leu Ala Phe Asp Pro Tyr
 165          170          175

```

-continued

---

Pro Ser Ala Ala Ala Leu Glu Leu Gly Val Glu Tyr Val Asp Leu Pro  
180 185 190

Thr Leu Phe Ser Glu Ser Asp Val Ile Ser Leu His Cys Pro Leu Thr  
195 200 205

Pro Glu Asn Tyr His Leu Leu Asn Glu Ala Ala Phe Glu Gln Met Lys  
210 215 220

Asn Gly Val Met Ile Val Asn Thr Ser Arg Gly Ala Leu Ile Asp Ser  
225 230 235 240

Gln Ala Ala Ile Glu Ala Leu Lys Asn Gln Lys Ile Gly Ser Leu Gly  
245 250 255

Met Asp Val Tyr Glu Asn Glu Arg Asp Leu Phe Phe Glu Asp Lys Ser  
260 265 270

Asn Asp Val Ile Gln Asp Asp Val Phe Arg Arg Leu Ser Ala Cys His  
275 280 285

Asn Val Leu Phe Thr Gly His Gln Ala Phe Leu Thr Ala Glu Ala Leu  
290 295 300

Thr Ser Ile Ser Gln Thr Thr Leu Gln Asn Leu Ser Asn Leu Glu Lys  
305 310 315 320

Gly Glu Thr Cys Pro Asn Glu Leu Val  
325

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 2277

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 19

```

tccgagctta atgaaaagtt agccacagcc tgggaaggtt ttaccaaagg tgactggcag      60
aatgaagtaa acgtccgtga cttcattcag aaaaactaca ctccgtacga gggtgacgag      120
tccttctctgg ctggcgctac tgaagcgacc accaccctgt gggacaaaagt aatggaaggg      180
gttaaactgg aaaaccgcac tcacgcgcca gttgactttg acaccgctgt tgcttccacc      240
atcacctctc acgacgctgg ctacatcaac aagcagcttg agaaaatcgt tggctctgcag      300
actgaagctc cgctgaaaacg tgctcttata ccgttcggtg gtatcaaaat gatcgaaggt      360
tcttgcaaag cgtacaaccg cgaactggat ccgatgatca aaaaaatctt cactgaatac      420
cgtaaaaactc acaaccaggg cgtgttcgac gtttacactc cggacatcct gcggttgcgt      480
aaatctgggtg ttctgaccgg tctgccagat gcatatggcc gtggccgtat catcggtgac      540
taccgtcgcy ttgctctgta cggtatcgac tacctgatga aagacaaact ggcacagttc      600
acttctctgc aggctgatct ggaaaacggc gtaaacctgg aacagactat cgtctctgcgc      660
gaagaaatcg ctgaacagca ccgcgctctg ggtcagatga aagaaatggc tgcgaaatac      720
ggctacgaca tctctgggtcc ggctaccaac gctcaggaag ctatccagtg gacttacttc      780
ggctacctgg ctgctgttaa gtctcagaac ggtgctgcaa tgtccttcgg tcgtacctcc      840
accttctctgg atgtgtacat cgaacgtgac ctgaaagctg gcaagatcac cgaacaagaa      900
gcgcaggaaa tggttgacca cctggctcatg aaactgcgta tggttcgctt cctgcgtact      960
ccggaatacg atgaactgtt ctctggcgac ccgatctggg caaccgaatc tatcggtggg     1020
atgggcctcg acggtcgtac cctggttacc aaaaacagct tccgtttcct gaacaccctg     1080
tacaccatgg gtccgtctcc ggaaccgaac atgaccattc tgtggtctga aaaactgccg     1140

```

-continued

---

```

ctgaacttca agaaattcgc cgctaaagtg tccatcgaca cctcttctct gcagtatgag 1200
aacgatgacc tgatgcgtcc ggacttcaac aacgatgact acgctattgc ttgctgcgta 1260
agcccgatga tcgttggtaa acaaatgcag ttcttcggtg cgcgtgcaaa cctggcgaaa 1320
accatgctgt acgcaatcaa cggcggcggt gacgaaaaac tgaaaatgca ggttggtccg 1380
aagtctgaac cgatcaaagg cgatgtcctg aactatgatg aagtgatgga gcgcatggat 1440
cacttcatgg actggctggc taaacagtac atcactgcac tgaacatcat ccaactacatg 1500
cacgacaagt acagctacga agcctctctg atggcgctgc acgaccgtga cgttatccgc 1560
accatggcgt gtggtatcgc tggctctgctc gttgctgctg actccctgtc tgcaatcaaa 1620
tatgcgaaag ttaaaccgat tcgtgacgaa gacggtctgg ctatcgactt cgaaatcgaa 1680
ggcgaatacc cgcagtttgg taacaatgat ccgctgtag atgacctggc tgttgacctg 1740
gtagaacggt tcatgaagaa aattcagaaa ctgcacacct accgtgacgc tatcccgact 1800
cagtctgttc tgaccatcac ttctaacggt gtgtatggta agaaaacggg taacacccca 1860
gacggtcgtc gtgctggcgc gccgttcgga ccgggtgcta acccgatgca cggctcgtgac 1920
cagaaagggt cagtagctc tctgacttcc gttgctaaac tgccgtttgc ttacgctaaa 1980
gatggtatct cctacacctt ctctatcggt ccgaacgcac tgggtaaaga cgacgaagtt 2040
cgtaagacca acctggetgg tctgatggat ggttacttcc accacgaagc atccatcgaa 2100
ggtggtcagc acctgaacgt taacgtgatg aaccgtgaaa tgctgctcga cgcgatggaa 2160
aaccgggaaa aatatccgca gctgaccatc cgtgtatctg gctacgcagt acgtttcaac 2220
tcgctgacta aagaacagca gcaggacggtt attactcgta ccttcaactca atctatg 2277

```

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 760

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 20

```

Met Ser Glu Leu Asn Glu Lys Leu Ala Thr Ala Trp Glu Gly Phe Thr
1           5           10           15
Lys Gly Asp Trp Gln Asn Glu Val Asn Val Arg Asp Phe Ile Gln Lys
20          25          30
Asn Tyr Thr Pro Tyr Glu Gly Asp Glu Ser Phe Leu Ala Gly Ala Thr
35          40          45
Glu Ala Thr Thr Thr Leu Trp Asp Lys Val Met Glu Gly Val Lys Leu
50          55          60
Glu Asn Arg Thr His Ala Pro Val Asp Phe Asp Thr Ala Val Ala Ser
65          70          75          80
Thr Ile Thr Ser His Asp Ala Gly Tyr Ile Asn Lys Gln Leu Glu Lys
85          90          95
Ile Val Gly Leu Gln Thr Glu Ala Pro Leu Lys Arg Ala Leu Ile Pro
100         105         110
Phe Gly Gly Ile Lys Met Ile Glu Gly Ser Cys Lys Ala Tyr Asn Arg
115         120         125
Glu Leu Asp Pro Met Ile Lys Lys Ile Phe Thr Glu Tyr Arg Lys Thr
130         135         140
His Asn Gln Gly Val Phe Asp Val Tyr Thr Pro Asp Ile Leu Arg Cys
145         150         155         160

```

-continued

---

Arg Lys Ser Gly Val Leu Thr Gly Leu Pro Asp Ala Tyr Gly Arg Gly  
 165 170 175

Arg Ile Ile Gly Asp Tyr Arg Arg Val Ala Leu Tyr Gly Ile Asp Tyr  
 180 185 190

Leu Met Lys Asp Lys Leu Ala Gln Phe Thr Ser Leu Gln Ala Asp Leu  
 195 200 205

Glu Asn Gly Val Asn Leu Glu Gln Thr Ile Arg Leu Arg Glu Glu Ile  
 210 215 220

Ala Glu Gln His Arg Ala Leu Gly Gln Met Lys Glu Met Ala Ala Lys  
 225 230 235 240

Tyr Gly Tyr Asp Ile Ser Gly Pro Ala Thr Asn Ala Gln Glu Ala Ile  
 245 250 255

Gln Trp Thr Tyr Phe Gly Tyr Leu Ala Ala Val Lys Ser Gln Asn Gly  
 260 265 270

Ala Ala Met Ser Phe Gly Arg Thr Ser Thr Phe Leu Asp Val Tyr Ile  
 275 280 285

Glu Arg Asp Leu Lys Ala Gly Lys Ile Thr Glu Gln Glu Ala Gln Glu  
 290 295 300

Met Val Asp His Leu Val Met Lys Leu Arg Met Val Arg Phe Leu Arg  
 305 310 315 320

Thr Pro Glu Tyr Asp Glu Leu Phe Ser Gly Asp Pro Ile Trp Ala Thr  
 325 330 335

Glu Ser Ile Gly Gly Met Gly Leu Asp Gly Arg Thr Leu Val Thr Lys  
 340 345 350

Asn Ser Phe Arg Phe Leu Asn Thr Leu Tyr Thr Met Gly Pro Ser Pro  
 355 360 365

Glu Pro Asn Met Thr Ile Leu Trp Ser Glu Lys Leu Pro Leu Asn Phe  
 370 375 380

Lys Lys Phe Ala Ala Lys Val Ser Ile Asp Thr Ser Ser Leu Gln Tyr  
 385 390 395 400

Glu Asn Asp Asp Leu Met Arg Pro Asp Phe Asn Asn Asp Asp Tyr Ala  
 405 410 415

Ile Ala Cys Cys Val Ser Pro Met Ile Val Gly Lys Gln Met Gln Phe  
 420 425 430

Phe Gly Ala Arg Ala Asn Leu Ala Lys Thr Met Leu Tyr Ala Ile Asn  
 435 440 445

Gly Gly Val Asp Glu Lys Leu Lys Met Gln Val Gly Pro Lys Ser Glu  
 450 455 460

Pro Ile Lys Gly Asp Val Leu Asn Tyr Asp Glu Val Met Glu Arg Met  
 465 470 475 480

Asp His Phe Met Asp Trp Leu Ala Lys Gln Tyr Ile Thr Ala Leu Asn  
 485 490 495

Ile Ile His Tyr Met His Asp Lys Tyr Ser Tyr Glu Ala Ser Leu Met  
 500 505 510

Ala Leu His Asp Arg Asp Val Ile Arg Thr Met Ala Cys Gly Ile Ala  
 515 520 525

Gly Leu Ser Val Ala Ala Asp Ser Leu Ser Ala Ile Lys Tyr Ala Lys  
 530 535 540

Val Lys Pro Ile Arg Asp Glu Asp Gly Leu Ala Ile Asp Phe Glu Ile  
 545 550 555 560



-continued

---

```

agcaaggtgg atatggcact ggtcggcgat atcaagtcca ctctgcgtgc attgcttcca 960
ttggtggaag aaaaagccga tcgcaagttt ctggataaag cgctggaaga ttaccgagac 1020
gcccgcaaag ggctggacga ttttagctaaa ccgagcgaga aagccattca cccgcaatat 1080
ctggcgcagc aaattagtca ttttgccgcc gatgacgcta ttttcacctg tgacgttggt 1140
acgccaacgg tgtggggcgc acgttatcta aaaatgaacg gcaagegctc cctggttaggt 1200
tcgtttaacc acggttcgat ggctaacgcc atgccgcagg cgctgggtgc gcaggcgaca 1260
gagccagaac gtcaggtggt cgccatgtgc ggcgatggcg gttttagcat gttgatgggc 1320
gatttcctct cagtagtgca gatgaaactg ccagtgaaaa ttgtcgtctt taacaacagc 1380
gtgctgggct ttgtggcgat ggagatgaaa gctggtggct atttgactga cggcaccgaa 1440
ctacacgaca caaactttgc ccgcattgcc gaagcgtgcg gcattacggg tatccgtgta 1500
gaaaaagcgt ctgaagttga tgaagccctg caacgcgcct tctccatcga cggtcgggtg 1560
ttggtggatg tgggtggtgc caaagaagag ttagccattc caccgcagat caaactcgaa 1620
caggccaaag gtttcagcct gtatatgctg cgcgcaatca tcagcggacg cggtgatgaa 1680
gtgatcgaac tggcgaaaac aaactggcta aggtaa 1716

```

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 572

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 22

```

Met Lys Gln Thr Val Ala Ala Tyr Ile Ala Lys Thr Leu Glu Ser Ala
1           5           10           15

Gly Val Lys Arg Ile Trp Gly Val Thr Gly Asp Ser Leu Asn Gly Leu
20          25          30

Ser Asp Ser Leu Asn Arg Met Gly Thr Ile Glu Trp Met Ser Thr Arg
35          40          45

His Glu Glu Val Ala Ala Phe Ala Ala Gly Ala Glu Ala Gln Leu Ser
50          55          60

Gly Glu Leu Ala Val Cys Ala Gly Ser Cys Gly Pro Gly Asn Leu His
65          70          75          80

Leu Ile Asn Gly Leu Phe Asp Cys His Arg Asn His Val Pro Val Leu
85          90          95

Ala Ile Ala Ala His Ile Pro Ser Ser Glu Ile Gly Ser Gly Tyr Phe
100         105         110

Gln Glu Thr His Pro Gln Glu Leu Phe Arg Glu Cys Ser His Tyr Cys
115        120        125

Glu Leu Val Ser Ser Pro Glu Gln Ile Pro Gln Val Leu Ala Ile Ala
130        135        140

Met Arg Lys Ala Val Leu Asn Arg Gly Val Ser Val Val Val Leu Pro
145        150        155        160

Gly Asp Val Ala Leu Lys Pro Ala Pro Glu Gly Ala Thr Met His Trp
165        170        175

Tyr His Ala Pro Gln Pro Val Val Thr Pro Glu Glu Glu Glu Leu Arg
180        185        190

Lys Leu Ala Gln Leu Leu Arg Tyr Ser Ser Asn Ile Ala Leu Met Cys
195        200        205

```

-continued

---

Gly Ser Gly Cys Ala Gly Ala His Lys Glu Leu Val Glu Phe Ala Gly  
 210 215 220  
 Lys Ile Lys Ala Pro Ile Val His Ala Leu Arg Gly Lys Glu His Val  
 225 230 235 240  
 Glu Tyr Asp Asn Pro Tyr Asp Val Gly Met Thr Gly Leu Ile Gly Phe  
 245 250 255  
 Ser Ser Gly Phe His Thr Met Met Asn Ala Asp Thr Leu Val Leu Leu  
 260 265 270  
 Gly Thr Gln Phe Pro Tyr Arg Ala Phe Tyr Pro Thr Asp Ala Lys Ile  
 275 280 285  
 Ile Gln Ile Asp Ile Asn Pro Ala Ser Ile Gly Ala His Ser Lys Val  
 290 295 300  
 Asp Met Ala Leu Val Gly Asp Ile Lys Ser Thr Leu Arg Ala Leu Leu  
 305 310 315 320  
 Pro Leu Val Glu Glu Lys Ala Asp Arg Lys Phe Leu Asp Lys Ala Leu  
 325 330 335  
 Glu Asp Tyr Arg Asp Ala Arg Lys Gly Leu Asp Asp Leu Ala Lys Pro  
 340 345 350  
 Ser Glu Lys Ala Ile His Pro Gln Tyr Leu Ala Gln Gln Ile Ser His  
 355 360 365  
 Phe Ala Ala Asp Asp Ala Ile Phe Thr Cys Asp Val Gly Thr Pro Thr  
 370 375 380  
 Val Trp Ala Ala Arg Tyr Leu Lys Met Asn Gly Lys Arg Arg Leu Leu  
 385 390 395 400  
 Gly Ser Phe Asn His Gly Ser Met Ala Asn Ala Met Pro Gln Ala Leu  
 405 410 415  
 Gly Ala Gln Ala Thr Glu Pro Glu Arg Gln Val Val Ala Met Cys Gly  
 420 425 430  
 Asp Gly Gly Phe Ser Met Leu Met Gly Asp Phe Leu Ser Val Val Gln  
 435 440 445  
 Met Lys Leu Pro Val Lys Ile Val Val Phe Asn Asn Ser Val Leu Gly  
 450 455 460  
 Phe Val Ala Met Glu Met Lys Ala Gly Gly Tyr Leu Thr Asp Gly Thr  
 465 470 475 480  
 Glu Leu His Asp Thr Asn Phe Ala Arg Ile Ala Glu Ala Cys Gly Ile  
 485 490 495  
 Thr Gly Ile Arg Val Glu Lys Ala Ser Glu Val Asp Glu Ala Leu Gln  
 500 505 510  
 Arg Ala Phe Ser Ile Asp Gly Pro Val Leu Val Asp Val Val Val Ala  
 515 520 525  
 Lys Glu Glu Leu Ala Ile Pro Pro Gln Ile Lys Leu Glu Gln Ala Lys  
 530 535 540  
 Gly Phe Ser Leu Tyr Met Leu Arg Ala Ile Ile Ser Gly Arg Gly Asp  
 545 550 555 560  
 Glu Val Ile Glu Leu Ala Lys Thr Asn Trp Leu Arg  
 565 570

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 2142

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 23



-continued

---

tcccgta	ttatgctgat	ccctaccgga	accagcgtcg	gtctgaccag	cgtcagcctt	60
ggcgtgatcc	gtgcaatgga	acgcaaaggc	gttcgtctga	gcgttttcaa	acctatcgct	120
cagccgcgta	ccggtggcga	tgcgcccgat	cagactacga	ctatcgtgcg	tgccaactct	180
tccaccacga	cggccgctga	accgctgaaa	atgagctacg	ttgaaggtct	gctttccagc	240
aatcagaaaag	atgtgctgat	ggaagagatc	gtcgcaaact	accacgctaa	caccaaagac	300
gctgaagtcg	ttctggttga	aggctctggc	ccgacacgta	agcaccagtt	tgcccagtct	360
ctgaactacg	aaatcgctaa	aacgctgaat	gcgaaaatcg	tcttcgttat	gtctcagggc	420
actgacaccc	cggaacagct	gaaagagcgt	atcgaactga	cccgaacag	cttcggcggc	480
gccaaaaaca	ccaacatcac	cggcgttatc	gttaacaaac	tgaacgcacc	ggttgatgaa	540
cagggtcgta	ctcgcccgga	tctgtccgag	atcttcgacg	actcttccaa	agctaaagta	600
aacaatggtg	atccggcgaa	gctgcaagaa	tccagcccgc	tgccggttct	cggcgctgtg	660
ccgtggagct	ttgacctgat	cgcgactcgt	gcgatcgata	tggtctcgca	cctgaatgcg	720
accatcatca	acgaaggcga	catcaatact	cgcgcggtta	aatccgtcac	ttctctgcga	780
cgcagcattc	cgcacatgct	ggagcacttc	cgtgcgggtt	ctctgctggt	gacttccgca	840
gaccgtcctg	acgtgctggt	ggccgcttgc	ctggcagcca	tgaacggcgt	agaaatcggc	900
gccctgctgc	tgactggcgg	ttacgaaatg	gacgcgcgca	tttctaaact	gtgcgaacgt	960
gctttcgcta	ccggcctgcc	ggtatztatg	gtgaacacca	acacctggca	gacctctctg	1020
agcctgcaga	gcttcaacct	ggaagttccg	gttgacgac	acgaacgtat	cgagaaagtt	1080
caggaatacg	ttgctaacta	catcaacgct	gactggatcg	aatctctgac	tgccacttct	1140
gagcgcagcc	gtcgtctgtc	tccgcctgcg	ttccgttatc	agctgactga	acttgccgcg	1200
aaagcgggca	aacgtatcgt	actgccggaa	ggtgacgaac	cgcgtaccgt	taaagcagcc	1260
gctatctgtg	ctgaacgtgg	tatcgcaact	tgcgtactgc	tggttaatcc	ggcagagatc	1320
aaccgtggtg	cagcgtctca	gggtgtagaa	ctgggtgcag	ggattgaaat	cgttgatcca	1380
gaagtgggtc	gcgaaagcta	tgttggtcgt	ctggtcgaac	tgcgtaagaa	caaaggcatg	1440
accgaaaccg	ttgcccgcga	acagctggaa	gacaacgtgg	tgctcggtac	gctgatgctg	1500
gaacaggatg	aagttgatgg	tctggtttcc	ggtgctgttc	acactaccgc	aaacaccatc	1560
cgtccgcgcg	tgcagctgat	caaaaactgca	ccgggcagct	ccctggtatc	ttccgtgttc	1620
ttcatgctgc	tgccggaaca	ggtttacggt	tacggtgact	gtgcgatcaa	cccggatccg	1680
accgctgaac	agctggcaga	aatcgcgatt	cagtccgctg	attccgctgc	ggccttcggc	1740
atcgaaccgc	gcgttgctat	gctctcctac	tccaccggta	cttctggtgc	aggtagcgac	1800
gtagaaaaag	ttcgcaagc	aactcgtctg	gcgaggaaa	aacgtcctga	cctgatgatc	1860
gacggtccgc	tgcagtacga	cgtgcccgtg	atggctgacg	ttgcgaaatc	caaagcggcc	1920
aactctccgg	ttgcaggtcg	cgtaccgtg	ttcatcttcc	cggatctgaa	caccggtaac	1980
accacctaca	aagcgtgaca	gcgttctgcc	gacctgatct	ccatcgggcc	gatgctgcag	2040
ggtatgcgca	agccggttaa	cgacctgtcc	cgtggcgcac	tggttgacga	tatcgtctac	2100
accatcgcgc	tgactgcgat	tcagtctgca	cagcagcagt	aa		2142

&lt;210&gt; SEQ ID NO 24

&lt;211&gt; LENGTH: 714

-continued

---

```

<212> TYPE: PRT
<213> ORGANISM: Escherichia coli

<400> SEQUENCE: 24

Met Ser Arg Ile Ile Met Leu Ile Pro Thr Gly Thr Ser Val Gly Leu
1          5          10          15

Thr Ser Val Ser Leu Gly Val Ile Arg Ala Met Glu Arg Lys Gly Val
          20          25          30

Arg Leu Ser Val Phe Lys Pro Ile Ala Gln Pro Arg Thr Gly Gly Asp
          35          40          45

Ala Pro Asp Gln Thr Thr Thr Ile Val Arg Ala Asn Ser Ser Thr Thr
          50          55          60

Thr Ala Ala Glu Pro Leu Lys Met Ser Tyr Val Glu Gly Leu Leu Ser
65          70          75          80

Ser Asn Gln Lys Asp Val Leu Met Glu Glu Ile Val Ala Asn Tyr His
          85          90          95

Ala Asn Thr Lys Asp Ala Glu Val Val Leu Val Glu Gly Leu Val Pro
          100         105         110

Thr Arg Lys His Gln Phe Ala Gln Ser Leu Asn Tyr Glu Ile Ala Lys
          115         120         125

Thr Leu Asn Ala Glu Ile Val Phe Val Met Ser Gln Gly Thr Asp Thr
          130         135         140

Pro Glu Gln Leu Lys Glu Arg Ile Glu Leu Thr Arg Asn Ser Phe Gly
145         150         155         160

Gly Ala Lys Asn Thr Asn Ile Thr Gly Val Ile Val Asn Lys Leu Asn
          165         170         175

Ala Pro Val Asp Glu Gln Gly Arg Thr Arg Pro Asp Leu Ser Glu Ile
          180         185         190

Phe Asp Asp Ser Ser Lys Ala Lys Val Asn Asn Val Asp Pro Ala Lys
          195         200         205

Leu Gln Glu Ser Ser Pro Leu Pro Val Leu Gly Ala Val Pro Trp Ser
          210         215         220

Phe Asp Leu Ile Ala Thr Arg Ala Ile Asp Met Ala Arg His Leu Asn
225         230         235         240

Ala Thr Ile Ile Asn Glu Gly Asp Ile Asn Thr Arg Arg Val Lys Ser
          245         250         255

Val Thr Phe Cys Ala Arg Ser Ile Pro His Met Leu Glu His Phe Arg
          260         265         270

Ala Gly Ser Leu Leu Val Thr Ser Ala Asp Arg Pro Asp Val Leu Val
          275         280         285

Ala Ala Cys Leu Ala Ala Met Asn Gly Val Glu Ile Gly Ala Leu Leu
          290         295         300

Leu Thr Gly Gly Tyr Glu Met Asp Ala Arg Ile Ser Lys Leu Cys Glu
305         310         315         320

Arg Ala Phe Ala Thr Gly Leu Pro Val Phe Met Val Asn Thr Asn Thr
          325         330         335

Trp Gln Thr Ser Leu Ser Leu Gln Ser Phe Asn Leu Glu Val Pro Val
          340         345         350

Asp Asp His Glu Arg Ile Glu Lys Val Gln Glu Tyr Val Ala Asn Tyr
          355         360         365

Ile Asn Ala Asp Trp Ile Glu Ser Leu Thr Ala Thr Ser Glu Arg Ser
          370         375         380

```

-continued

---

Arg Arg Leu Ser Pro Pro Ala Phe Arg Tyr Gln Leu Thr Glu Leu Ala  
 385 390 395 400  
 Arg Lys Ala Gly Lys Arg Ile Val Leu Pro Glu Gly Asp Glu Pro Arg  
 405 410 415  
 Thr Val Lys Ala Ala Ala Ile Cys Ala Glu Arg Gly Ile Ala Thr Cys  
 420 425 430  
 Val Leu Leu Gly Asn Pro Ala Glu Ile Asn Arg Val Ala Ala Ser Gln  
 435 440 445  
 Gly Val Glu Leu Gly Ala Gly Ile Glu Ile Val Asp Pro Glu Val Val  
 450 455 460  
 Arg Glu Ser Tyr Val Gly Arg Leu Val Glu Leu Arg Lys Asn Lys Gly  
 465 470 475 480  
 Met Thr Glu Thr Val Ala Arg Glu Gln Leu Glu Asp Asn Val Val Leu  
 485 490 495  
 Gly Thr Leu Met Leu Glu Gln Asp Glu Val Asp Gly Leu Val Ser Gly  
 500 505 510  
 Ala Val His Thr Thr Ala Asn Thr Ile Arg Pro Pro Leu Gln Leu Ile  
 515 520 525  
 Lys Thr Ala Pro Gly Ser Ser Leu Val Ser Ser Val Phe Phe Met Leu  
 530 535 540  
 Leu Pro Glu Gln Val Tyr Val Tyr Gly Asp Cys Ala Ile Asn Pro Asp  
 545 550 555 560  
 Pro Thr Ala Glu Gln Leu Ala Glu Ile Ala Ile Gln Ser Ala Asp Ser  
 565 570 575  
 Ala Ala Ala Phe Gly Ile Glu Pro Arg Val Ala Met Leu Ser Tyr Ser  
 580 585 590  
 Thr Gly Thr Ser Gly Ala Gly Ser Asp Val Glu Lys Val Arg Glu Ala  
 595 600 605  
 Thr Arg Leu Ala Gln Glu Lys Arg Pro Asp Leu Met Ile Asp Gly Pro  
 610 615 620  
 Leu Gln Tyr Asp Ala Ala Val Met Ala Asp Val Ala Lys Ser Lys Ala  
 625 630 635 640  
 Pro Asn Ser Pro Val Ala Gly Arg Ala Thr Val Phe Ile Phe Pro Asp  
 645 650 655  
 Leu Asn Thr Gly Asn Thr Thr Tyr Lys Ala Val Gln Arg Ser Ala Asp  
 660 665 670  
 Leu Ile Ser Ile Gly Pro Met Leu Gln Gly Met Arg Lys Pro Val Asn  
 675 680 685  
 Asp Leu Ser Arg Gly Ala Leu Val Asp Asp Ile Val Tyr Thr Ile Ala  
 690 695 700  
 Leu Thr Ala Ile Gln Ser Ala Gln Gln Gln  
 705 710

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 255

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 25

ttccagcaag aagttaccat taccgctccg aacgggtctgc acaccegccg tgctgcccag 60

tttgtaaaag aagctaaggg cttcacttct gaaattactg tgacttccaa cggcaaaagc 120

-continued

---

```

gccagcgcga aaagcctggt taaactgcag actctgggccc tgactcaagg taccgttggtg 180
actatctccg cagaaggcga agacgagcag aaagcgggtg aacatctggt taaactgatg 240
gcggaactcg agtaa 255

```

```

<210> SEQ ID NO 26
<211> LENGTH: 85
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli

```

```

<400> SEQUENCE: 26

```

```

Met Phe Gln Gln Glu Val Thr Ile Thr Ala Pro Asn Gly Leu His Thr
1           5           10           15
Arg Pro Ala Ala Gln Phe Val Lys Glu Ala Lys Gly Phe Thr Ser Glu
          20           25           30
Ile Thr Val Thr Ser Asn Gly Lys Ser Ala Ser Ala Lys Ser Leu Phe
          35           40           45
Lys Leu Gln Thr Leu Gly Leu Thr Gln Gly Thr Val Val Thr Ile Ser
          50           55           60
Ala Glu Gly Glu Asp Glu Gln Lys Ala Val Glu His Leu Val Lys Leu
65           70           75           80
Met Ala Glu Leu Glu
          85

```

```

<210> SEQ ID NO 27
<211> LENGTH: 1725
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli

```

```

<400> SEQUENCE: 27

```

```

atttcaggca ttttagcatc cccgggtatc gctttcggta aagctctgct tctgaaagaa 60
gacgaaattg tcattgaccg gaaaaaatt tctgccgacc aggttgatca ggaagttgaa 120
cgttttctga gcggtcgtgc caaggcatca gccagctgg aaacgatcaa aacgaaagct 180
ggtgaaacgt tcggtgaaga aaaagaagcc atctttgaag ggcattat gctgctcgaa 240
gatgaggagc tggagcagga aatcatagcc ctgattaaag ataagcacat gacagctgac 300
gcagctgctc atgaagttat cgaaggtcag gcttctgccc tggaagagct ggatgatgaa 360
tacctgaaag aacgtgcggc tgacgtacgt gatatcggta agcgcctgct gcgcaacatc 420
ctgggctga agattatcga cctgagcgcc attcaggatg aagtcattct ggttgccgct 480
gacctgacgc cgtccgaaac cgcacagctg aacctgaaga aggtgctggg tttcatcacc 540
gacgcgggtg gccgtacttc ccacacctct atcatggcgc gttctctgga actacctgct 600
atcgtgggta ccggtagcgt cacctctcag gtgaaaaatg acgactatct gattctggat 660
gccgtaaata atcaggttta cgtcaatcca accaacgaag ttattgataa aatgcgcgct 720
gttcaggagc aagtggcttc tgaaaaagca gagcttgcta aactgaaaga tctgccagct 780
attacgctgg acggtcacca ggtagaagta tgcgctaaca ttggtacggt tcgtgacggt 840
gaaggtgcag agcgtaacgg cgctgaaggc gttggtctgt atcgtactga gttcctgttc 900
atggaccgcy acgactgcc cactgaagaa gaacagtttg ctgcttaca agcagtggtc 960
gaagcgtgtg gctcgcaagc ggttatcgtt cgtaccatgg acatcggcgg cgacaaagag 1020
ctgccataca tgaacttccc gaaagaagag aacccttcc tcggctggcg cgtatccgt 1080

```

-continued

---

```

atcgcgatgg atcgtagaga gatcctgcgc gatcagctcc gcgctatcct gcgtgcctcg 1140
gctttcggta aattgcgcat tatgttcccc atgatcatct ctggtgaaga agtgcgtgca 1200
ctgcgcaaag agatcgaaat ctacaaacag gaactgcgcg acgaaggtaa agcgtttgac 1260
gagtcaattg aaatcggcgt aatggtggaa acaccggctg ccgcaacaat tgcacgtcat 1320
ttagccaaag aagttgattt ctttagtatac ggcaccaatg atttaacgca gtacactctg 1380
gcagttgacc gtggtaatga tatgatttca cacctttacc agccaatgtc accgtccgtg 1440
ctgaacttga tcaagcaagt tattgatgct tctcatgctg aaggcaaatg gactggcatg 1500
tgtggtgagc ttgctggcga tgaacgtgct acacttctgt tgctggggat gggctctggac 1560
gaattctcta tgagcgccat ttctatcccg cgcattaaga agattatccg taacacgaac 1620
ttcgaagatg cgaaggtgtt agcagagcag gctcttgctc aaccgacaac ggacgagtta 1680
atgacgctgg ttaacaagtt cattgaagaa aaaacaatct gctaa 1725

```

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 575

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 28

```

Met Ile Ser Gly Ile Leu Ala Ser Pro Gly Ile Ala Phe Gly Lys Ala
1           5           10           15
Leu Leu Leu Lys Glu Asp Glu Ile Val Ile Asp Arg Lys Lys Ile Ser
20          25          30
Ala Asp Gln Val Asp Gln Glu Val Glu Arg Phe Leu Ser Gly Arg Ala
35          40          45
Lys Ala Ser Ala Gln Leu Glu Thr Ile Lys Thr Lys Ala Gly Glu Thr
50          55          60
Phe Gly Glu Glu Lys Glu Ala Ile Phe Glu Gly His Ile Met Leu Leu
65          70          75          80
Glu Asp Glu Glu Leu Glu Gln Glu Ile Ile Ala Leu Ile Lys Asp Lys
85          90          95
His Met Thr Ala Asp Ala Ala Ala His Glu Val Ile Glu Gly Gln Ala
100         105         110
Ser Ala Leu Glu Glu Leu Asp Asp Glu Tyr Leu Lys Glu Arg Ala Ala
115         120         125
Asp Val Arg Asp Ile Gly Lys Arg Leu Leu Arg Asn Ile Leu Gly Leu
130         135         140
Lys Ile Ile Asp Leu Ser Ala Ile Gln Asp Glu Val Ile Leu Val Ala
145         150         155         160
Ala Asp Leu Thr Pro Ser Glu Thr Ala Gln Leu Asn Leu Lys Lys Val
165         170         175
Leu Gly Phe Ile Thr Asp Ala Gly Gly Arg Thr Ser His Thr Ser Ile
180         185         190
Met Ala Arg Ser Leu Glu Leu Pro Ala Ile Val Gly Thr Gly Ser Val
195         200         205
Thr Ser Gln Val Lys Asn Asp Asp Tyr Leu Ile Leu Asp Ala Val Asn
210         215         220
Asn Gln Val Tyr Val Asn Pro Thr Asn Glu Val Ile Asp Lys Met Arg
225         230         235         240
Ala Val Gln Glu Gln Val Ala Ser Glu Lys Ala Glu Leu Ala Lys Leu

```

-continued

	245		250		255										
Lys	Asp	Leu	Pro	Ala	Ile	Thr	Leu	Asp	Gly	His	Gln	Val	Glu	Val	Cys
	260							265					270		
Ala	Asn	Ile	Gly	Thr	Val	Arg	Asp	Val	Glu	Gly	Ala	Glu	Arg	Asn	Gly
	275						280					285			
Ala	Glu	Gly	Val	Gly	Leu	Tyr	Arg	Thr	Glu	Phe	Leu	Phe	Met	Asp	Arg
	290					295					300				
Asp	Ala	Leu	Pro	Thr	Glu	Glu	Glu	Gln	Phe	Ala	Ala	Tyr	Lys	Ala	Val
305					310				315						320
Ala	Glu	Ala	Cys	Gly	Ser	Gln	Ala	Val	Ile	Val	Arg	Thr	Met	Asp	Ile
				325					330					335	
Gly	Gly	Asp	Lys	Glu	Leu	Pro	Tyr	Met	Asn	Phe	Pro	Lys	Glu	Glu	Asn
			340					345					350		
Pro	Phe	Leu	Gly	Trp	Arg	Ala	Ile	Arg	Ile	Ala	Met	Asp	Arg	Arg	Glu
	355						360					365			
Ile	Leu	Arg	Asp	Gln	Leu	Arg	Ala	Ile	Leu	Arg	Ala	Ser	Ala	Phe	Gly
	370					375					380				
Lys	Leu	Arg	Ile	Met	Phe	Pro	Met	Ile	Ile	Ser	Val	Glu	Glu	Val	Arg
385					390					395					400
Ala	Leu	Arg	Lys	Glu	Ile	Glu	Ile	Tyr	Lys	Gln	Glu	Leu	Arg	Asp	Glu
			405						410					415	
Gly	Lys	Ala	Phe	Asp	Glu	Ser	Ile	Glu	Ile	Gly	Val	Met	Val	Glu	Thr
			420					425					430		
Pro	Ala	Ala	Ala	Thr	Ile	Ala	Arg	His	Leu	Ala	Lys	Glu	Val	Asp	Phe
	435						440					445			
Phe	Ser	Ile	Gly	Thr	Asn	Asp	Leu	Thr	Gln	Tyr	Thr	Leu	Ala	Val	Asp
	450					455					460				
Arg	Gly	Asn	Asp	Met	Ile	Ser	His	Leu	Tyr	Gln	Pro	Met	Ser	Pro	Ser
465					470					475					480
Val	Leu	Asn	Leu	Ile	Lys	Gln	Val	Ile	Asp	Ala	Ser	His	Ala	Glu	Gly
			485						490					495	
Lys	Trp	Thr	Gly	Met	Cys	Gly	Glu	Leu	Ala	Gly	Asp	Glu	Arg	Ala	Thr
			500					505					510		
Leu	Leu	Leu	Leu	Gly	Met	Gly	Leu	Asp	Glu	Phe	Ser	Met	Ser	Ala	Ile
			515				520					525			
Ser	Ile	Pro	Arg	Ile	Lys	Lys	Ile	Ile	Arg	Asn	Thr	Asn	Phe	Glu	Asp
	530					535					540				
Ala	Lys	Val	Leu	Ala	Glu	Gln	Ala	Leu	Ala	Gln	Pro	Thr	Thr	Asp	Glu
545					550					555					560
Leu	Met	Thr	Leu	Val	Asn	Lys	Phe	Ile	Glu	Glu	Lys	Thr	Ile	Cys	
				565					570					575	

<210> SEQ ID NO 29  
 <211> LENGTH: 507  
 <212> TYPE: DNA  
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 29

ggtttggttcg ataaactgaa atctctgggt tccgacgaca agaaggatac cggaactatt 60  
 gagatcattg ctccgctctc tggcgagatc gtcaatatcg aagacgtgcc ggatgtcgtt 120  
 tttgcggaaa aaatcggttg tgatggtatt gctatcaaac caacgggtaa caaatggtc 180

-continued

---

```

gcgccagtag acggcaccat tggtaaaatc ttgaaacca accacgcatt ctctatcgaa 240
tctgatagcg gcgttgaact gttcgtccac ttcggtatcg acaccgttga actgaaaggc 300
gaaggcttca agcgtattgc tgaagaaggt cagcgcgtga aagttggcga tactgtcatt 360
gaatttgatc tgccgctgct ggaagagaaa gccaaagtcta ccctgactcc ggttgttatc 420
tccaacatgg acgaaatcaa agaactgatc aaactgtccg gtagcgtaac cgtgggtgaa 480
accccggtta tccgcatcaa gaagtaa 507

```

```

<210> SEQ ID NO 30
<211> LENGTH: 169
<212> TYPE: PRT
<213> ORGANISM: Escherichia coli
<400> SEQUENCE: 30

```

```

Met Gly Leu Phe Asp Lys Leu Lys Ser Leu Val Ser Asp Asp Lys Lys
1           5           10           15
Asp Thr Gly Thr Ile Glu Ile Ile Ala Pro Leu Ser Gly Glu Ile Val
20           25           30
Asn Ile Glu Asp Val Pro Asp Val Val Phe Ala Glu Lys Ile Val Gly
35           40           45
Asp Gly Ile Ala Ile Lys Pro Thr Gly Asn Lys Met Val Ala Pro Val
50           55           60
Asp Gly Thr Ile Gly Lys Ile Phe Glu Thr Asn His Ala Phe Ser Ile
65           70           75           80
Glu Ser Asp Ser Gly Val Glu Leu Phe Val His Phe Gly Ile Asp Thr
85           90           95
Val Glu Leu Lys Gly Glu Gly Phe Lys Arg Ile Ala Glu Glu Gly Gln
100          105          110
Arg Val Lys Val Gly Asp Thr Val Ile Glu Phe Asp Leu Pro Leu Leu
115          120          125
Glu Glu Lys Ala Lys Ser Thr Leu Thr Pro Val Val Ile Ser Asn Met
130          135          140
Asp Glu Ile Lys Glu Leu Ile Lys Leu Ser Gly Ser Val Thr Val Gly
145          150          155          160
Glu Thr Pro Val Ile Arg Ile Lys Lys
165

```

```

<210> SEQ ID NO 31
<211> LENGTH: 1443
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli
<400> SEQUENCE: 31

```

```

atgtccagaa ggcttcgacg aacaaaaatc gttaccacgt taggcccagc aacagatcgc 60
gataataatc ttgaaaaagt tatcgcgggc ggtgccaacg ttgtacgtat gaacttttct 120
cacggctcgc ctgaagatca caaatgcgc gcggataaag ttcgtgagat tgccgcaaaa 180
ctggggcgctc atgtggctat tctgggtgac ctccaggggc ccaaatccg tgtatccacc 240
tttaaagaag gcaaagtttt cctcaatatt ggggataaat tcctgctcga cgccaacctg 300
ggtaaaggty aaggcgacaa agaaaaagtc ggtatcgact acaaaggcct gcctgctgac 360
gtcgtgcctg gtgacatcct gctgctggac gatggctcgc tccagttaaa agtactggaa 420
gttcagggca tgaaagtgtt caccgaagtc accgtcggty gtcccctctc caacaataaa 480

```

-continued

---

```

ggtatcaaca aacttggcgg cggtttgctg gctgaagcgc tgaccgaaaa agacaaagca 540
gacattaaga ctgcggcggt gattggcgta gattacctgg ctgtctcctt cccacgctgt 600
ggcgaagatc tgaactatgc ccgtcgctcg gcacgcgatg caggatgtga tgcgaaaatt 660
gttgccaagg ttgaactgct ggaagccggt tgcagccagg atgcaatgga tgacatcatc 720
ctgcgctctg acgtggtaat ggttgcacgt ggcgacctcg gtgtggaaat tggcgacctg 780
gaactggctg gcattcagaa agcgttgatc cgtcgtgcgc gtcagctaaa ccgagcggta 840
atcacggcga cccagatgat ggagtcaatg attactaacc cgatgccgac gcgtgcagaa 900
gtcatggacg tagcaaacgc cgttctggat ggtactgacg ctgtgatgct gtctgcagaa 960
actgccgctg ggcagtatcc gtcagaaacc gttgcagcca tggcgcgcgt ttgcctgggt 1020
gcggaaaaaa tcccagcat caacgtttct aaacaccgtc tggacgttca gttcgacaat 1080
gtggaagaag ctattgcat gtcagcaatg tacgcagcta accacctgaa aggcgttacg 1140
gcgatcatca ccatgaccga atcgggtcgt accgcgctga tgacctcccg tatcagctct 1200
ggtctgcaa ttttcgcat gtcgcgcat gaacgtacgc tgaacctgac tgctctctat 1260
cgtggcgta cgccggtgca ctttgatagc gctaatagac gcgtagcagc tgccagcgaa 1320
gcggttaatc tgctgcgca taaaggttac ttgatgtctg gtgacctggt gattgtcacc 1380
cagggcgacg tgatgagtac cgtgggttct actaatacca cgcgtatattt aacggtagag 1440
taa 1443

```

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 480

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 32

```

Met Ser Arg Arg Leu Arg Arg Thr Lys Ile Val Thr Thr Leu Gly Pro
1          5          10          15
Ala Thr Asp Arg Asp Asn Asn Leu Glu Lys Val Ile Ala Ala Gly Ala
20          25          30
Asn Val Val Arg Met Asn Phe Ser His Gly Ser Pro Glu Asp His Lys
35          40          45
Met Arg Ala Asp Lys Val Arg Glu Ile Ala Ala Lys Leu Gly Arg His
50          55          60
Val Ala Ile Leu Gly Asp Leu Gln Gly Pro Lys Ile Arg Val Ser Thr
65          70          75          80
Phe Lys Glu Gly Lys Val Phe Leu Asn Ile Gly Asp Lys Phe Leu Leu
85          90          95
Asp Ala Asn Leu Gly Lys Gly Glu Gly Asp Lys Glu Lys Val Gly Ile
100         105         110
Asp Tyr Lys Gly Leu Pro Ala Asp Val Val Pro Gly Asp Ile Leu Leu
115        120        125
Leu Asp Asp Gly Arg Val Gln Leu Lys Val Leu Glu Val Gln Gly Met
130        135        140
Lys Val Phe Thr Glu Val Thr Val Gly Gly Pro Leu Ser Asn Asn Lys
145        150        155        160
Gly Ile Asn Lys Leu Gly Gly Gly Leu Ser Ala Glu Ala Leu Thr Glu
165        170        175

```



-continued

---

Lys Asp Lys Ala Asp Ile Lys Thr Ala Ala Leu Ile Gly Val Asp Tyr  
 180 185 190  
 Leu Ala Val Ser Phe Pro Arg Cys Gly Glu Asp Leu Asn Tyr Ala Arg  
 195 200 205  
 Arg Leu Ala Arg Asp Ala Gly Cys Asp Ala Lys Ile Val Ala Lys Val  
 210 215 220  
 Glu Arg Ala Glu Ala Val Cys Ser Gln Asp Ala Met Asp Asp Ile Ile  
 225 230 235 240  
 Leu Ala Ser Asp Val Val Met Val Ala Arg Gly Asp Leu Gly Val Glu  
 245 250 255  
 Ile Gly Asp Pro Glu Leu Val Gly Ile Gln Lys Ala Leu Ile Arg Arg  
 260 265 270  
 Ala Arg Gln Leu Asn Arg Ala Val Ile Thr Ala Thr Gln Met Met Glu  
 275 280 285  
 Ser Met Ile Thr Asn Pro Met Pro Thr Arg Ala Glu Val Met Asp Val  
 290 295 300  
 Ala Asn Ala Val Leu Asp Gly Thr Asp Ala Val Met Leu Ser Ala Glu  
 305 310 315 320  
 Thr Ala Ala Gly Gln Tyr Pro Ser Glu Thr Val Ala Ala Met Ala Arg  
 325 330 335  
 Val Cys Leu Gly Ala Glu Lys Ile Pro Ser Ile Asn Val Ser Lys His  
 340 345 350  
 Arg Leu Asp Val Gln Phe Asp Asn Val Glu Glu Ala Ile Ala Met Ser  
 355 360 365  
 Ala Met Tyr Ala Ala Asn His Leu Lys Gly Val Thr Ala Ile Ile Thr  
 370 375 380  
 Met Thr Glu Ser Gly Arg Thr Ala Leu Met Thr Ser Arg Ile Ser Ser  
 385 390 395 400  
 Gly Leu Pro Ile Phe Ala Met Ser Arg His Glu Arg Thr Leu Asn Leu  
 405 410 415  
 Thr Ala Leu Tyr Arg Gly Val Thr Pro Val His Phe Asp Ser Ala Asn  
 420 425 430  
 Asp Gly Val Ala Ala Ala Ser Glu Ala Val Asn Leu Leu Arg Asp Lys  
 435 440 445  
 Gly Tyr Leu Met Ser Gly Asp Leu Val Ile Val Thr Gln Gly Asp Val  
 450 455 460  
 Met Ser Thr Val Gly Ser Thr Asn Thr Thr Arg Ile Leu Thr Val Glu  
 465 470 475 480

<210> SEQ ID NO 33  
 <211> LENGTH: 1410  
 <212> TYPE: DNA  
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 33

```

aaaaagacca aaattgtttg caccatcgga ccgaaaaccg aatctgaaga gatgtagct    60
aaaatgctgg acgctggcat gaacgttatg cgtctgaact tctctcatgg tgactatgca    120
gaacacggtc agcgattca gaatctgctc aacgtgatga gcaaaactgg taaaaccgcc    180
gctatcctgc ttgataccaa aggtccggaa atccgcacca tgaaactgga aggcggtaac    240
gacgtttctc tgaaagctgg tcagacctt actttcacca ctgataaatc tgttatcggc    300
aacagcgaaa tggttgcggt aacgtatgaa ggtttcaacta ctgacctgct tgttggcaac    360
  
```

-continued

---

```

accgtactgg ttgacgatgg tctgatcggg atggaagtta cgcaccattga aggtaacaaa 420
gttatctgta aagtgctgaa caacgggtgac ctgggacgaaa acaaagggtgt gaacctgcct 480
ggcgtttcca ttgctctgcc agcactggct gaaaaagaca aacaggacct gatctttggt 540
tgcgaacaag gcgtagactt tgttgctgct tcctttattc gtaagcgctt tgacggtatc 600
gaaatccgtg agcacctgaa agcgcacggc ggcgaaaaca tccacatcat ctccaaaatc 660
gaaaaccagg aaggcctcaa caacttcgac gaaatcctcg aagcctctga cggcatcatg 720
gttgcgcggt ggcacctggg tgtagaaatc ccggtagaag aagttatctt cgcccagaag 780
atgatgatcg aaaaatgtat ccgtgcacgt aaagtcgta tcaactgcgac ccagatgctg 840
gattccatga tcaaaaacc acgcccact cgcgcagaag ccggtgacgt tgcaaacgcc 900
atcctcgacg gtactgacgc agtgatgctg tctggtgaat ccgcaaaagg taaatacccc 960
ctggaagcgg tttctatcat ggcgaccatc tgcgaacgta ccgaccgct gatgaacagc 1020
cgtctcgagt tcaacaatga caaccgtaaa ctgctgatta ccgaagcggg atgccgtggt 1080
gccgttgaaa ctgctgaaaa actggatgct ccgctgatcg tggttgctac tcagggcggt 1140
aaatctgctc ggcagctacg taaatacttc ccggatgcca ccatcctggc actgaccacc 1200
aacgaaaaaa cggctcatca gttggtactg agcaaaggcg ttgtgccgca gcttggtaaa 1260
gagatcactt ctactgatga tttctaccgt ctgggtaaag aactggctct gcagagcggt 1320
ctggcacaca aaggtgacgt tgtagttatg gtttctggtg cactggtacc gagcggcact 1380
actaacaccg catctgttca cgctctgtaa 1410

```

&lt;210&gt; SEQ ID NO 34

&lt;211&gt; LENGTH: 470

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 34

```

Met Lys Lys Thr Lys Ile Val Cys Thr Ile Gly Pro Lys Thr Glu Ser
1           5           10           15

Glu Glu Met Leu Ala Lys Met Leu Asp Ala Gly Met Asn Val Met Arg
          20           25           30

Leu Asn Phe Ser His Gly Asp Tyr Ala Glu His Gly Gln Arg Ile Gln
          35           40           45

Asn Leu Arg Asn Val Met Ser Lys Thr Gly Lys Thr Ala Ala Ile Leu
          50           55           60

Leu Asp Thr Lys Gly Pro Glu Ile Arg Thr Met Lys Leu Glu Gly Gly
65           70           75           80

Asn Asp Val Ser Leu Lys Ala Gly Gln Thr Phe Thr Phe Thr Thr Asp
          85           90           95

Lys Ser Val Ile Gly Asn Ser Glu Met Val Ala Val Thr Tyr Glu Gly
          100          105          110

Phe Thr Thr Asp Leu Ser Val Gly Asn Thr Val Leu Val Asp Asp Gly
          115          120          125

Leu Ile Gly Met Glu Val Thr Ala Ile Glu Gly Asn Lys Val Ile Cys
          130          135          140

Lys Val Leu Asn Asn Gly Asp Leu Gly Glu Asn Lys Gly Val Asn Leu
145          150          155          160

Pro Gly Val Ser Ile Ala Leu Pro Ala Leu Ala Glu Lys Asp Lys Gln

```

-continued

---

	165		170		175										
Asp	Leu	Ile	Phe	Gly	Cys	Glu	Gln	Gly	Val	Asp	Phe	Val	Ala	Ala	Ser
	180							185					190		
Phe	Ile	Arg	Lys	Arg	Ser	Asp	Val	Ile	Glu	Ile	Arg	Glu	His	Leu	Lys
	195						200					205			
Ala	His	Gly	Gly	Glu	Asn	Ile	His	Ile	Ile	Ser	Lys	Ile	Glu	Asn	Gln
	210					215					220				
Glu	Gly	Leu	Asn	Asn	Phe	Asp	Glu	Ile	Leu	Glu	Ala	Ser	Asp	Gly	Ile
225					230					235					240
Met	Val	Ala	Arg	Gly	Asp	Leu	Gly	Val	Glu	Ile	Pro	Val	Glu	Glu	Val
				245					250					255	
Ile	Phe	Ala	Gln	Lys	Met	Met	Ile	Glu	Lys	Cys	Ile	Arg	Ala	Arg	Lys
			260					265					270		
Val	Val	Ile	Thr	Ala	Thr	Gln	Met	Leu	Asp	Ser	Met	Ile	Lys	Asn	Pro
		275					280					285			
Arg	Pro	Thr	Arg	Ala	Glu	Ala	Gly	Asp	Val	Ala	Asn	Ala	Ile	Leu	Asp
	290					295					300				
Gly	Thr	Asp	Ala	Val	Met	Leu	Ser	Gly	Glu	Ser	Ala	Lys	Gly	Lys	Tyr
305					310					315					320
Pro	Leu	Glu	Ala	Val	Ser	Ile	Met	Ala	Thr	Ile	Cys	Glu	Arg	Thr	Asp
				325					330					335	
Arg	Val	Met	Asn	Ser	Arg	Leu	Glu	Phe	Asn	Asn	Asp	Asn	Arg	Lys	Leu
			340					345					350		
Arg	Ile	Thr	Glu	Ala	Val	Cys	Arg	Gly	Ala	Val	Glu	Thr	Ala	Glu	Lys
		355					360					365			
Leu	Asp	Ala	Pro	Leu	Ile	Val	Val	Ala	Thr	Gln	Gly	Gly	Lys	Ser	Ala
370						375					380				
Arg	Ala	Val	Arg	Lys	Tyr	Phe	Pro	Asp	Ala	Thr	Ile	Leu	Ala	Leu	Thr
385					390					395					400
Thr	Asn	Glu	Lys	Thr	Ala	His	Gln	Leu	Val	Leu	Ser	Lys	Gly	Val	Val
				405					410					415	
Pro	Gln	Leu	Val	Lys	Glu	Ile	Thr	Ser	Thr	Asp	Asp	Phe	Tyr	Arg	Leu
			420					425					430		
Gly	Lys	Glu	Leu	Ala	Leu	Gln	Ser	Gly	Leu	Ala	His	Lys	Gly	Asp	Val
		435					440					445			
Val	Val	Met	Val	Ser	Gly	Ala	Leu	Val	Pro	Ser	Gly	Thr	Thr	Asn	Thr
	450					455					460				
Ala	Ser	Val	His	Val	Leu										
465					470										

&lt;210&gt; SEQ ID NO 35

&lt;211&gt; LENGTH: 569

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Zymomonasmobilis

&lt;400&gt; SEQUENCE: 35

Met	Ser	Tyr	Thr	Val	Gly	Thr	Tyr	Leu	Ala	Glu	Arg	Leu	Val	Gln	Ile
1				5					10					15	
Gly	Leu	Lys	His	His	Phe	Ala	Val	Ala	Gly	Asp	Tyr	Asn	Leu	Val	Leu
			20					25					30		
Leu	Asp	Asn	Leu	Leu	Leu	Asn	Lys	Asn	Met	Glu	Gln	Val	Tyr	Cys	Cys
	35						40					45			

-continued

---

Asn	Glu	Leu	Asn	Cys	Gly	Phe	Ser	Ala	Glu	Gly	Tyr	Ala	Arg	Ala	Lys
50						55					60				
Gly	Ala	Ala	Ala	Ala	Val	Val	Thr	Tyr	Ser	Val	Gly	Ala	Leu	Ser	Ala
65					70					75					80
Phe	Asp	Ala	Ile	Gly	Gly	Ala	Tyr	Ala	Glu	Asn	Leu	Pro	Val	Ile	Leu
				85					90					95	
Ile	Ser	Gly	Ala	Pro	Asn	Asn	Asn	Asp	His	Ala	Ala	Gly	His	Val	Leu
			100					105					110		
His	His	Ala	Leu	Gly	Lys	Thr	Asp	Tyr	His	Tyr	Gln	Leu	Glu	Met	Ala
		115					120					125			
Lys	Asn	Ile	Thr	Ala	Ala	Ala	Glu	Ala	Ile	Tyr	Thr	Pro	Glu	Glu	Ala
	130					135					140				
Pro	Ala	Lys	Ile	Asp	His	Val	Ile	Lys	Thr	Ala	Leu	Arg	Glu	Lys	Lys
145					150					155					160
Pro	Val	Tyr	Leu	Glu	Ile	Ala	Cys	Asn	Ile	Ala	Ser	Met	Pro	Cys	Ala
				165					170					175	
Ala	Pro	Gly	Pro	Ala	Ser	Ala	Leu	Phe	Asn	Asp	Glu	Ala	Ser	Asp	Glu
			180					185						190	
Ala	Ser	Leu	Asn	Ala	Ala	Val	Glu	Glu	Thr	Leu	Lys	Phe	Ile	Ala	Asn
		195					200					205			
Arg	Asp	Lys	Val	Ala	Val	Leu	Val	Gly	Ser	Lys	Leu	Arg	Ala	Ala	Gly
	210					215					220				
Ala	Glu	Glu	Ala	Ala	Val	Lys	Phe	Ala	Asp	Ala	Leu	Gly	Gly	Ala	Val
225					230					235					240
Ala	Thr	Met	Ala	Ala	Ala	Lys	Ser	Phe	Phe	Pro	Glu	Glu	Asn	Pro	His
				245					250					255	
Tyr	Ile	Gly	Thr	Ser	Trp	Gly	Glu	Val	Ser	Tyr	Pro	Gly	Val	Glu	Lys
			260					265					270		
Thr	Met	Lys	Glu	Ala	Asp	Ala	Val	Ile	Ala	Leu	Ala	Pro	Val	Phe	Asn
		275					280						285		
Asp	Tyr	Ser	Thr	Thr	Gly	Trp	Thr	Asp	Ile	Pro	Asp	Pro	Lys	Lys	Leu
	290					295					300				
Val	Leu	Ala	Glu	Pro	Arg	Ser	Val	Val	Val	Asn	Gly	Ile	Arg	Phe	Pro
305					310					315					320
Ser	Val	His	Leu	Lys	Asp	Tyr	Leu	Thr	Arg	Leu	Ala	Gln	Lys	Val	Ser
				325					330					335	
Lys	Lys	Thr	Gly	Ala	Leu	Asp	Phe	Phe	Lys	Ser	Leu	Asn	Ala	Gly	Glu
			340					345					350		
Leu	Lys	Lys	Ala	Ala	Pro	Ala	Asp	Pro	Ser	Ala	Pro	Leu	Val	Asn	Ala
		355					360					365			
Glu	Ile	Ala	Arg	Gln	Val	Glu	Ala	Leu	Leu	Thr	Pro	Asn	Thr	Thr	Val
		370				375					380				
Ile	Ala	Glu	Thr	Gly	Asp	Ser	Trp	Phe	Asn	Ala	Gln	Arg	Met	Lys	Leu
385					390					395					400
Pro	Asn	Gly	Ala	Arg	Val	Glu	Tyr	Glu	Met	Gln	Trp	Gly	His	Ile	Gly
				405					410					415	
Trp	Ser	Val	Pro	Ala	Ala	Phe	Gly	Tyr	Ala	Val	Gly	Ala	Pro	Glu	Arg
			420					425					430		
Arg	Asn	Ile	Leu	Met	Val	Gly	Asp	Gly	Ser	Phe	Gln	Leu	Thr	Ala	Gln
		435					440					445			
Glu	Val	Ala	Gln	Met	Val	Arg	Leu	Lys	Leu	Pro	Val	Ile	Ile	Phe	Leu



-continued

---

Val Ile Ala Asp Val Leu Ser Gln Ser Arg His Thr Tyr Thr Ile His  
 245 250 255  
 Gly Ile Asp Val Leu Leu Gln Ala His Lys Ile Asn Ala Asp Leu Lys  
 260 265 270  
 Pro Asp Leu Val Val Arg Phe Gly Lys Thr Pro Val Ser Ala Arg Val  
 275 280 285  
 Leu Gln Trp Leu Lys Glu Glu Asn Ile Leu Thr Trp His Val Asp Glu  
 290 295 300  
 Asp Ala Gly Val Asp His Thr Arg His Ile Val Arg Ala Ile Lys Met  
 305 310 315 320  
 Ala Pro His Asp Phe Leu Glu Ser Met His Leu Thr Leu Ser Lys Asn  
 325 330 335  
 Gln Ile Asp Phe Asn Gln Lys Trp Leu Ser Leu Pro Lys Val Ile Lys  
 340 345 350  
 Thr Arg Asn Glu Met Asn Ile Ile Thr Ala Leu Asp Asp Ala Val Pro  
 355 360 365  
 Asp Asp Thr His Ile Phe Val Ala Asn Ser Met Pro Ile Arg Asp Met  
 370 375 380  
 Asp Asn Phe Phe Thr Gly Asn His Thr Gln Arg Ile Tyr Ala Asn Arg  
 385 390 395 400  
 Gly Ala Asn Gly Ile Asp Gly Val Ile Ser Ser Ala Leu Gly Met Ser  
 405 410 415  
 Ala Val Val Lys Gln Arg Ser Val Leu Leu Thr Gly Asp Leu Thr Leu  
 420 425 430  
 Phe His Asp Met Asn Gly Leu Met Met Ala Lys Asn Tyr Gln Leu Pro  
 435 440 445  
 Leu Asp Ile Ile Val Ile Asn Asn Asn Gly Gly Gly Ile Phe Ser Phe  
 450 455 460  
 Leu Pro Gln Ala Gly Ala Pro Lys Tyr Phe Glu Gln Leu Phe Gly Thr  
 465 470 475 480  
 Pro Leu Asn Ile Asp Ile Lys Lys Ile Ala Asp Leu Tyr Tyr Ile Asp  
 485 490 495  
 Tyr His Gln Leu Asn Val Pro Glu Ala Leu Ser Gln Ile Leu Gln Thr  
 500 505 510  
 Pro Ser Lys Thr Thr Arg Leu Ile Glu Tyr Lys Ser Asp His Gln Arg  
 515 520 525  
 Asn Arg Asp Asp His Arg Glu Val Leu Glu Met Leu Lys  
 530 535 540

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 1242

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Mycobacterium tuberculosis

&lt;400&gt; SEQUENCE: 37

Met Asn Leu Arg Gly Thr Ala Ala Arg Arg Arg His Arg Ala Thr Gly  
 1 5 10 15  
 Asp Thr Leu Asp Ala Arg Gln Leu Val Arg Gln Gln Leu Arg Gln Leu  
 20 25 30  
 Gln Thr Glu Thr Arg Pro Gly Ala Leu Val Arg Trp Phe Leu Leu Asp  
 35 40 45  
 Pro Lys Glu Leu Phe Asp Ile Phe Val Thr Ile Gln Gly Phe Thr Gln  
 50 55 60

-continued

---

Thr Thr Thr Arg Gln Arg Arg Gln Leu Leu Gln Thr His Asn Cys Gln  
 65 70 75 80  
 Ile Ile Thr Val Val Phe Arg Leu Ala Cys Arg Gln Phe Ile Ile Gln  
 85 90 95  
 Leu Thr Gly Gly Gln Gln Asn Thr Ala Asn Phe Ile Ala Val Ala Asp  
 100 105 110  
 Thr Ile Leu Val Gly Arg Phe Leu Gln Asp Ala Thr Glu Phe Asn Leu  
 115 120 125  
 Gly Lys Ile Leu Asn Leu Arg Asn Arg Arg Leu Val Ala Gln His Gly  
 130 135 140  
 Phe Arg Gly Lys His Asp Gln Arg Thr Leu Asp Ala Val Gln Arg Val  
 145 150 155 160  
 Ala Ala Gln Gln Val Lys Val Ile Gly Arg Gly Ala Arg His Arg Asp  
 165 170 175  
 Arg His Ala Thr Leu Ser Pro Gln Leu Gln Lys Ala Leu Asn Ala Arg  
 180 185 190  
 Arg Arg Val Val Arg Ala Leu Thr Leu Val Ala Val Trp Gln Gln Gln  
 195 200 205  
 Asn Asn Val Gly Gln Leu Thr Pro Phe Arg Phe Ala Arg Ala Asp Glu  
 210 215 220  
 Leu Val Asn Asp Arg Leu Gly Thr Ile Asn Lys Val Ala Glu Leu Arg  
 225 230 235 240  
 Leu Pro Gln His Asn Arg Val Trp Ile Ala Asn Gly Ile Thr Val Leu  
 245 250 255  
 Lys Thr His Ser Arg Val Phe Arg Gln Arg Arg Ile Ile Asn Gln Glu  
 260 265 270  
 Leu Thr Thr Gly Arg Ala Ala Val Ala Val Ser Arg Gln Gln Leu Gln  
 275 280 285  
 Arg Gly Glu Leu Leu Ala Gly Val Thr Val Asp Gln His Arg Met Thr  
 290 295 300  
 Leu Ala Lys Gly Ala Thr Thr Arg Val Leu Thr Ala Gln Thr Asp Gln  
 305 310 315 320  
 Leu Ala Leu Ser Asn Gln Ala Ala Gln Arg Gln Gln Leu Thr Lys Arg  
 325 330 335  
 Pro Val Asn Leu Thr Leu Ile Arg His Leu Thr Thr Leu Phe Gln His  
 340 345 350  
 Arg Leu Asn Ala Arg Val Cys Arg Glu Thr Val Arg Gln Arg Gln Glu  
 355 360 365  
 Gly Ile Thr Asp Thr Cys Gln Gln Gly Phe Val Asn Ser Gly Arg Gln  
 370 375 380  
 Ala Arg Trp Asp His Leu Val Arg Leu Asp Ala Leu Arg Arg Leu Asp  
 385 390 395 400  
 Ala Val Leu Leu Gln Phe Ala His Phe Val Glu His Thr Leu Gln Leu  
 405 410 415  
 Ala Leu Val Ile Ala Gln Arg Ile Leu Arg Leu Phe His Ala Asp Val  
 420 425 430  
 Ala Thr Thr Asp Gln Ser Leu Gly Val Gly Phe Thr Gly Ala Thr Leu  
 435 440 445  
 Gly Val His Asp Val Ile Asp Ile Arg Ile Gly His Arg Trp Ile Val  
 450 455 460

-continued

---

Thr	Leu	Ile	Met	Thr	Thr	Thr	Thr	Ile	Ala	Gln	His	Val	Asn	Asn	Asn			
465					470					475					480			
Val	Leu	Phe	Lys	Ala	Leu	Ala	Glu	Val	Asn	Cys	Gln	Thr	Arg	Asn	Pro			
				485					490					495				
Asp	Thr	Arg	Phe	Trp	Ile	Val	Ala	Val	His	Val	Glu	Asp	Trp	Arg	Thr			
			500					505					510					
Asp	His	Leu	Arg	His	Ile	Arg	Ala	Val	Leu	Ala	Arg	Thr	Arg	Val	Phe			
		515					520					525						
Arg	Ser	Gly	Gly	Glu	Ala	Asp	Leu	Val	Val	His	Asn	Asp	Val	Asn	Arg			
	530					535					540							
Thr	Thr	His	Thr	Ile	Ala	Arg	Gln	Ile	Cys	Gln	Ile	Gln	Arg	Leu	Arg			
545				550						555					560			
Asn	Asn	Ala	Leu	Thr	Arg	Glu	Gly	Arg	Ile	Thr	Val	Gln	His	Gln	Arg			
				565					570					575				
Asn	Asp	Gly	Glu	Cys	Thr	Leu	Ala	Val	Gly	Val	Asn	Gly	Thr	Val	Val			
		580						585				590						
Gln	Gln	Val	Leu	Leu	Arg	Thr	His	Gln	Thr	Phe	Gln	His	Trp	Ile	Asp			
		595					600					605						
Arg	Phe	Gln	Met	Gly	Arg	Ile	Cys	Arg	Gln	Gly	Asn	Leu	Asn	Ile	Val			
	610					615					620							
Ile	Ala	Lys	His	Leu	Gln	Ile	Gln	Thr	Arg	Arg	Thr	Gln	Val	Val	Leu			
625					630					635					640			
Asp	Ile	Ala	Gly	Thr	Val	Ser	Leu	Gly	Arg	Val	Gln	Ile	Ala	Phe	Lys			
			645					650						655				
Leu	Arg	Lys	Asp	Leu	Arg	Ile	Arg	Phe	Ala	His	Asp	Val	Arg	Gln	Asp			
			660					665					670					
Ile	Gln	Ala	Thr	Thr	Val	Arg	His	Ala	Asn	Asp	His	Phe	Ile	Gln	Thr			
		675					680					685						
Met	Leu	Gly	Thr	Leu	Val	Asn	Arg	Arg	Val	His	His	Trp	Asn	Asn	Arg			
	690					695					700							
Phe	Arg	Thr	Leu	Gln	Ala	Lys	Thr	Leu	Leu	Ala	His	Ile	Leu	Gly	Leu			
705					710					715					720			
Gln	Glu	Gly	Phe	Lys	Arg	Leu	Arg	Cys	Val	Gln	Phe	Arg	Gln	Asp	Val			
			725						730					735				
Leu	Leu	Leu	Ser	Asn	Gly	Arg	Phe	Asp	Val	Leu	Arg	Leu	Asp	Thr	Leu			
			740					745					750					
Leu	Gln	Pro	Leu	Leu	Leu	Phe	Arg	Val	Gln	Asn	Val	Arg	Val	Leu	Asn			
		755					760					765						
Thr	Asp	Val	Thr	Ala	Val	Arg	Val	Ala	Gln	Gln	Thr	Gln	His	Val	Thr			
	770					775						780						
Gln	Leu	Phe	Val	Leu	Ser	Thr	Arg	Glu	Thr	Val	Asn	Leu	Lys	Asp	Ala			
785					790					795					800			
Val	Gln	Val	Pro	Gln	Ser	Gln	Ala	Met	Arg	Gln	His	Leu	Gln	Ile	Arg			
				805					810					815				
Met	Arg	Thr	Lys	Ala	Arg	Leu	Ile	Gln	Ala	Gln	Arg	Val	Gly	Val	Arg			
			820					825					830					
His	Gln	Met	Ala	Ala	Val	Ala	Ile	Gly	Arg	Asp	Gln	Val	His	His	Thr			
		835					840					845						
Cys	Val	Leu	Val	Asn	Asp	Arg	Val	Arg	Ile	Ile	Gly	Ala	Pro	Thr	His			
	850					855					860							
Trp	Gln	Val	Arg	Asp	Ala	Gln	Leu	Ala	Glu	Asp	Leu	Ile	Pro	Glu	Ala			



-continued

865	870	875	880
Ile Arg Gln Gln Gln Phe Met Asn Gly Thr Gln Glu Val Thr Ala Phe	885	890	895
Arg Thr Leu Asn Asp Thr Val Val Val Gly Arg Ser Gln Gly Asn Gln	900	905	910
Phe Ala Asn Thr Gln Leu Ser Asp Ala Phe Leu Gly Arg Ala Leu Glu	915	920	925
Leu Arg Arg Ile Phe His Arg Thr Asp Thr Asp Asp Ser Thr Leu Thr	930	935	940
Trp His Gln Ala Trp His Arg Val His Arg Ala Asn Arg Ala Arg Ile	945	950	955
Arg Gln Arg Asn Arg Asn Ala Ser Glu Val Leu Gly Gly Gln Phe Thr	965	970	975
Ile Thr Ser Thr Thr Asp Asp Val Leu Val Arg Gly Asn Glu Leu Arg	980	985	990
Glu Ala His Arg Leu Ala Thr Phe Asp Thr Gly His His Gln Arg Ala	995	1000	1005
Leu Ala Ile Phe Thr Leu Gln Val Asn Ser Gln Thr Gln Ile Gly	1010	1015	1020
Val Arg Arg Ser Asn Arg Arg Arg Leu Thr Val His Leu Arg Val	1025	1030	1035
Val Ala Ile His Ile Arg Glu Leu Leu Asn Arg Leu Asn Gln Ser	1040	1045	1050
Ile Thr Gln Gln Met Gly Lys Ala Asp Phe Thr Thr Thr Arg Ala	1055	1060	1065
Phe Gln Leu Ile Ile Asp His Asp Thr Ile Ile Asp Gln Gln Phe	1070	1075	1080
Arg Trp His Ser Thr His Arg Gly Arg Arg Arg His Phe Gln Arg	1085	1090	1095
Arg Ala His Val Leu His Asp Ser Ser Arg Arg Thr Thr Gln Asp	1100	1105	1110
Ser Asn Phe Ile Ala Phe Gly Trp Arg Arg His Arg Gly Leu Gly	1115	1120	1125
Gly Gln Ser Arg His Asp Thr Val Ala Arg Cys Gly Arg Val Gly	1130	1135	1140
Arg Phe Arg Arg Arg Leu Arg Ser Arg Arg Cys Arg Thr Leu Ser	1145	1150	1155
Asp Gln Arg Ala Gly His Thr Ser Arg Leu Ser Ser Arg Leu Gly	1160	1165	1170
Ser Arg Leu Arg Ala Ile Val His Gln Lys Leu Met Pro Ala Trp	1175	1180	1185
Val Asn Gly Arg Arg Ile Ile Thr Lys Phe Thr Ile His Thr Asn	1190	1195	1200
Thr Arg Arg Val Leu Ser His His Gly Tyr Ile Ser Phe Leu Lys	1205	1210	1215
Asn Ser Ile Ser Gln Arg Gln Gly Arg His Lys Ile Tyr Ser Lys	1220	1225	1230
Cys Ile Ile Asn Thr Asp Asn Ile Leu	1235	1240	

-continued

---

<211> LENGTH: 4412  
<212> TYPE: DNA  
<213> ORGANISM: Artificial Sequence  
<220> FEATURE:  
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
          plasmid pJ201:pdc

<400> SEQUENCE: 38

tagaaaaact catcgagcat caaatgaaac tgcaatztat tcatatcagg attatcaata	60
ccatattttt gaaaaagccg tttctgtaat gaaggagaaa actcaccgag gcagttccat	120
aggatggcaa gatcctggta tcggctctgcg attccgactc gtccaacatc aatacaacct	180
attaatttcc cctcgtcaaa aataaggtta tcaagtgaga aatcaccatg agtgacgact	240
gaatccgggtg agaatggcaa aagtttatgc atttctttcc agacttgttc aacaggccag	300
ccattacgct cgatcatcaa atcaactcga tcaaccaaac cgttattcat tcgtgattgc	360
gcctgagcga ggcgaaatac gcgatcgcgt ttaaaaggac aattacaaac aggaatcgag	420
tgcaaccggc gcaggaacac tgccagcgc tcaacaatat tttcacctga atcaggatat	480
tcttctaata cctggaacgc tgtttttccg gggatcgcag tggtagtaaa ccatgcatca	540
tcaggagtac ggataaaatg cttgatggtc ggaagtggca taaattccgt cagccagttt	600
agtctgacca tctcatctgt aacatcattg gcaacgctac ctttgccatg tttcagaaac	660
aactctggcg catcgggctt cccatacaag cgatagattg tcgcacctga ttgcccgaca	720
ttatcgcgag cccatttata cccatataaa tcagcatcca tgttggaatt taatcgcggc	780
ctcgaagttt cccggtgaat atggctcata ttcttccttt ttcaatatta ttgaagcatt	840
tatcagggtt attgtctcat gagcggatac atatttgaat gtatttagaa aaataaacia	900
ataggggtca gtgttacaac caattaacca attctgaaca ttatcgcgag cccatttata	960
cctgaatatg gctcataaca ccccttgttt gcctggcggc agtagcgcgg tggcccacc	1020
tgaccccatg ccgaactcag aagtgaaacg ccgtagcggc gatggtagtg tggggactcc	1080
ccatgcgaga gtagggaact gccaggcatc aaataaaacg aaaggctcag tcgaaagact	1140
ggcctttcg cccgggctaa ttaggggggtg tcgcccctcg ctgaacatat gagctatacc	1200
gttggcacct atctggcaga acgctgggtg cagatcggcc tgaaacacca cttcgcgggtg	1260
gcgggtgact acaacctggt gctgctggac aacttgctgt tgaacaagaa tatggaacag	1320
gtgtactgct gtaatgagct gaattgtggc ttcagcgcgg agggttacgc ccgggcaaaa	1380
ggtgcagccg cagcggttgt gacctatagc gttggcgcgc tgagcgcatt tgatgcgatt	1440
ggtggtgctg acgctgagaa tttgcccgtc atcctgatca gcggcgcacc gaacaacaac	1500
gacctgctg cgggccacgt gctgcatcac gcgctgggta agaccgacta tcattatcag	1560
ttggagatgg cgaagaatat cacggccgct gcggaagcta tctatacgcc ggaagaggct	1620
cctgcaaaaa tcgatcacgt tatcaaaacc gactgcgtg agaagaagcc agtgtacctg	1680
gagattgctt gcaacattgc gtctatgccg tgcgcagcac cgggtcctgc gagegcgttg	1740
tttaacgatg aggccagcga cgaagcgagc ctgaacgcgg ctggtgaaga aacgctgaaa	1800
ttcattgcaa atcgtgataa ggttgctgct ctggttgcca gcaagctgcg tgccgcaggc	1860
gcggaagaag cagcggtgaa gttcgcagat gcctgggtg gcgcggtcgc aacctggct	1920
gccgcaaaa ctttctttcc ggaagagaat ccacactata tcggcactag ctgggggtgag	1980
gtgtcttatc cgggcgtcga gaaaacctg aaagaagcgg atgcggtgat cgccctggct	2040

-continued

---

ccggttttca	acgattacag	caccactggc	tggacggata	ttccggaccc	gaaaaagctg	2100
gtgctggcgg	agccgcgtag	cgtggtggtc	aatggtatcc	gttttccgtc	cgtccatctg	2160
aaggactacc	tgacccgcct	ggcgcagaag	gtagcaaga	aaacgggtgc	gctggatttc	2220
ttcaaatcgc	tgaatgcagg	tgaactgaaa	aaggcagccc	cagcagatcc	gtccgcaccg	2280
ctggtgaatg	cggagatcgc	gcgtaagtc	gaagctctgc	tgacgccgaa	cacgaccgtt	2340
atcgcggaga	cgggtgacag	ctggtttaac	gcgcaacgta	tgaagttgcc	gaatggtgct	2400
cgagttgagt	acgagatgca	atggggccac	attggtggt	ccgtgcctgc	agcctttggt	2460
tacgccgttg	gtgccccgga	gcgtcgcaac	attctgatgg	tgggtgacgg	tagctttcaa	2520
ctgaccgccc	aagaggtcgc	tcagatggtc	cgctgaaac	tgectgtcat	cattttcctg	2580
atcaataact	acggttacac	gatcgagggt	atgatccatg	atggcccgta	caacaacatt	2640
aagaattggg	actacgcggg	ttgatggag	gttttcaatg	gcaacggtgg	ttatgactcc	2700
ggtgcccgta	aaggtctgaa	agcaaagacg	ggtggcgagc	tcgcagaggc	gattaagggt	2760
gcgctggcga	ataccgacgg	cccgccttg	attgagtgtt	tcattggtcg	cgaggattgc	2820
accgaagaac	tgggtcaaatg	gggcaagcgc	gtcgcagccg	cgaattcccg	taaaccggtt	2880
aataagcttc	tcggccatca	ccatcaccat	cactagaagc	ttctctagag	aactatttcc	2940
gtcaatcgag	ttcgtacct	agggcgacac	aaaatttatt	ctaatgcat	aataaatact	3000
gataacatct	tatagtttgt	attatatttt	gtattatcgt	tgacatgtat	aattttgata	3060
tcaaaaactg	attttccctt	tattattttc	gagatttatt	ttcttaattc	tctttaacaa	3120
actagaaata	ttgtatatac	aaaaaatcat	aaataataga	tgaatagttt	aattataggt	3180
gttcatcaat	cgaaaaagca	acgtatctta	tttaaagtgc	gttgcttttt	tctcatttat	3240
aaggttaaat	aattctcata	tatcaagcaa	agtgcagggc	gcccttaaat	attctgacaa	3300
atgctctttc	cctaaactcc	cccataaaa	aaaccgcg	aagcgggttt	ttacgttatt	3360
tgccgattaa	cgattactcg	ttatcagaac	cgcccagggg	gcccagactt	aagactggcc	3420
gtcgttttac	aacacagaaa	gagttttag	aaacgcaaaa	aggccatccg	tcaggggcct	3480
tctgcttagt	ttgatgcctg	gcagttccct	actctgcct	tcgcttccct	cgctcactga	3540
ctcgtgcgc	tcggtcgttc	ggctgcccgc	agcggatca	gctcactcaa	aggcggtaat	3600
acggttatcc	acagaatcag	gggataacgc	aggaaagaac	atgtgagcaa	aaggccagca	3660
aaaggccagg	aaccgtaaaa	aggccgcggt	gctggcggtt	ttccataggc	tcgccccccc	3720
tgacgagcat	cacaaaaatc	gacgctcaag	tcagaggtgg	cgaaaccgca	caggactata	3780
aagataccag	gcgtttcccc	ctggaagctc	cctcgtgcgc	tctcctgttc	cgaccctgcc	3840
gcttacogga	tacctgtccg	cctttctccc	ttcgggaagc	gtggcgcttt	ctcatagctc	3900
acgctgtagg	tatctcagtt	cgggtgtagt	cgttcgcctc	aagctgggct	gtgtgcacga	3960
acccccggtt	cagcccagcc	gctgcgcctt	atccggtaac	tatcgtcttg	agtccaaccc	4020
ggtaagacac	gacttatcgc	cactggcagc	agccactggt	aacaggatta	gcagagcgag	4080
gtatgtaggc	ggtgctacag	agttcttgaa	gtggtgggct	aactacggct	acactagaag	4140
aacagtatth	ggtatctgcg	ctctgctgaa	gccagttacc	ttcggaaaaa	gagttggtag	4200
ctcttgatcc	ggcaaacaaa	ccaccgctgg	tagcgggtgg	ttttttgttt	gcaagcagca	4260
gattacgcgc	agaaaaaag	gatctcaaga	agatcctttg	atcttttcta	cggggtctga	4320

-continued

---

 cgctcagtgg aacgacgcgc gcgtaactca cgtaagga ttttggcat gagcttgcgc 4380

cgtcccgta agtcagcgta atgctctgct tt 4412

&lt;210&gt; SEQ ID NO 39

&lt;211&gt; LENGTH: 4418

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic plasmid pJ251:oad

&lt;400&gt; SEQUENCE: 39

gcttttagaa aaactcatcg agcatcaaat gaaactgcaa tttattcata tcaggattat 60

caataccata tttttgaaaa agccgtttct gtaatgaagg agaaaactca ccgaggcagt 120

tccataggat ggcaagatcc tggatcggt ctgcatcc gactcgtcca acatcaatac 180

aacctattaa tttccctcg tcaaaaataa gggtatcaag tgagaaatca ccatgagtga 240

cgactgaatc cggtgagaat ggcaaaagtt tatgcatttc tttccagact tgttcaacag 300

gccagccatt acgctcgta tcaaaatcac tcgcatcaac caaacctta ttcattcgtg 360

attgcgctg agcgaggcga aatagcgat cgctgttaa aggacaatta caaacaggaa 420

tcgagtgcaa ccggcgagg aacactgcca gcgcatcaac aatattttca cctgaatcag 480

gatattcttc taatacctgg aacgctgttt ttccgggat cgcagtggg agtaaccatg 540

catcatcagg agtacggata aaatgcttga tggtcggaag tggcataaat tccgtcagcc 600

agtttagtct gaccatctca tctgtaacat cattggcaac gctacctttg ccatgtttca 660

gaaacaactc tggcgcatcg ggcttcccat acaagcgata gattgtcgca cctgattgcc 720

cgacattatc gcgagcccat ttatacccat ataaatcagc atccatgttg gaatttaatc 780

gcggcctcga cgtttccgt tgaatatggc tcatattctt ctttttcaa tattattgaa 840

gcatttatca gggttattgt ctcatgagcg gatacatatt tgaatgtatt tagaaaaata 900

aacaatagg ggtcagtgtt acaaccaatt aaccaattct gaacattatc gcgagcccat 960

ttatacctga atatggctca taacaccct tgtttgcctg gcggcagtag cgcgggtggc 1020

ccacctgacc ccatgccgaa ctgagaagt aaacgcgta gcgccgatgg tagtgtgggg 1080

actcccatg cgagagtagg gaactgccag gcatcaaata aaacgaaagg ctgagtcgaa 1140

agactgggcc tttcgcccg gctaattagg ggggtgctgc cttattcgac tctataggg 1200

agttcctatt ctctagaaag tataggaact tctgaagggg ggtttcgccc ggggacagct 1260

gtcggtttga ataatgaca aaaagcaaag cttttgtgcc gatgaatctc tatactgttt 1320

cacagacctg ctgccctgc gggcgcccat cttcctttat tcgcttataa gcgtggagaa 1380

ttaaactgac ggatacctg accttcaaca ccaaacattt gttggaagcg ctgtttgaaa 1440

gcggtattcg tcaactcatt gtttctccgg gttctgtag caccctgatt gcactgctgc 1500

tggcagaata cgcagagcag aataacgaga tcaagctgtt cgttgacgtg gatgagcgca 1560

gcgcggttt cttcgactg ggcacgcga aaacgttgc ggagccggt gttttgctgg 1620

gtacctggg taccgccatc gctgaataca tgctgcggg ggcggaagcc tatgcccga 1680

atatccgct ggtcgtgttg agcaccgatc gtccacagga attgcagttc aacggcgac 1740

cgcaactat tccgcaaagc aacctgttcg gtcaactgac gaaacaggca gtctgattc 1800

-continued

---

gcctgcaaga	catgcacagc	gatgtgaccg	agtacatcga	ctttatcgtt	cagaaggctc	1860
ttaatctgtc	cattaccgcy	cctcgtggcc	cgattcagat	caacctgccg	ctgcygcaaac	1920
cgctgatgcc	ggctcctgat	cgccaagacg	aggtgcacgt	tcagcgtgtc	gtgtttgaca	1980
aacaaagcgt	gcaatatcgc	ttgccgccga	ttacggcgaa	acgcctgctg	atcctggcag	2040
gtccgaatgt	cctgaatagc	tacgatgata	gcctgaagaa	gtttgcgatc	aaaaacaatg	2100
ttccagttat	cgcggtatgt	ctgtcccaaa	gccgccacac	gtataccatt	cacggatttg	2160
acgttctgct	gcaagcccac	aagatcaacg	ccgacctgaa	accggatctg	gtggttcgtt	2220
ttggcaagac	gccggttagc	gcgcgtgtcc	tgcaatggct	gaaagaagag	aacattctga	2280
cctggcacgt	ggacgaggac	gcaggcgtgg	atcacaccgc	tcatatcgtc	cgtgccatca	2340
agatggcgcc	tcgatgtttt	ctggaatcca	tgcatctgac	gttgagcaaa	aaccagattg	2400
acttcaacca	gaagtggctg	tctctgccga	aagtgatcaa	aactcgtaac	gagatgaata	2460
tcataccgc	attggacgac	gcgggtcccg	atgacactca	catcttcgtg	gcgaacagca	2520
tgccgattcg	tgacatggac	aactttttca	cgggtaatca	caccagcgc	atttacgcta	2580
atcgtgggtc	aatgggtatt	gatggcgtga	ttagctccgc	cctgggcatg	agcgcggctc	2640
tcaagcaacg	ttcggctcctg	ctgacgggtg	acctgacgct	gtttcatgac	atgaacggtc	2700
tgatgatggc	gaagaattac	cagctgccgc	tgacatcat	tgtgattaac	aacaatggcg	2760
gtggcatttt	cagcttcttg	ccacaagccg	gtgctccgaa	gtactttgag	cagctgttcg	2820
gtaccccgct	gaacattgac	atcaaaaaga	ttgctgacct	gtattacatc	gattaccatc	2880
agttgaatgt	gccggaggcc	ctgagccaga	tcctgcagac	gccgagcaaa	accacccgcc	2940
tgatcgagta	taagtccgat	caccagcgtg	accgtgacga	tcatacgcga	gtgctggaga	3000
tgctgaaaca	ccaccatcat	cattagcacg	tgaaaatgaa	gggaagttcc	tatactttct	3060
agagaatagg	aacttctata	gggagtcgaa	taagggcgac	acaaaaggta	ttctaaatgc	3120
ataataaata	ctgataacat	cttatagttt	gtattatatt	ttgtattatc	gttgacatgt	3180
ataattttga	tatcaaaaac	tgattttccc	tttattattt	tcgagattta	ttttcttaat	3240
tctctttaac	aaactagaaa	tattgtatat	acaaaaaatc	ataaataata	gatgaatagt	3300
ttaattatag	gtgttcatca	atcgaaaaag	caacgtatct	tatttaaagt	gcgttgcttt	3360
tttctcattt	ataaggttaa	ataattctca	tatatcaagc	aaagtgacag	gcgcccttaa	3420
atattctgac	aatgctctt	tcctaaact	ccccccataa	aaaaaccgc	cgaagcgggt	3480
ttttacgtta	tttgcggatt	aacgattact	cgttatcaga	accgccagc	atgcctggca	3540
gttcctact	ctcgcctctg	cgctcggctc	ttcggctgcg	ggacctcagc	gctagcggag	3600
tgataactgg	cttactatgt	tggcactgat	gaggggtgca	gtgaagtgct	tcagtggca	3660
ggagaaaaaa	ggctgcaccg	gtgcgtcagc	agaatatgtg	atacaggata	tattccgctt	3720
cctcgtcac	tgactcgcta	cgctcggctc	ttcgaactgc	gcgagcggaa	atggcttacg	3780
aacggggcgg	agatttcctg	gaagatgcca	ggaagatact	taacaggaa	gtgagagggc	3840
cgcgcaaaag	ccgtttttcc	ataggctccg	ccccctgac	aagcatcacg	aatctgacg	3900
ctcaaatcag	tgggtggcgaa	acccgacagg	actataaaga	taccaggcgt	ttccccctgg	3960
cggtccctc	gtgcctctc	ctgttctctc	ctttcggttt	accgggtgca	ttccgctgtt	4020
atggcccgct	ttgtctcatt	ccacgcctga	cactcagttc	cgggtaggca	gttcgctcca	4080

-continued

---

```

agctggactg tatgcacgaa cccccgttc agtccgaccg ctgcgectta tccggtaact 4140
atcgtcttga gtccaaccg gaaagacatg caaaagcacc actggcagca gccactggta 4200
attgatttag aggagttagt cttgaagtca tgcgccggtt aaggctaaac tgaaaggaca 4260
agttttggtg actgcgctcc tccaagccag ttacctcggg tcaaagagtt ggtagctcag 4320
agaaccttcg aaaaaccgcc ctgcaaggcg gttttttcgt tttcagagca agagattacg 4380
cgcagaccaa aacgatctca agaagatcat cttattaa 4418

```

&lt;210&gt; SEQ ID NO 40

&lt;211&gt; LENGTH: 8621

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Synthetic plasmid pJ206:kgd

&lt;400&gt; SEQUENCE: 40

```

ggtggcggta cttgggtcga tatcaaagt catcacttct tcccgtatgc ccaactttgt 60
atagagagcc actgcgggat cgtcaccgta atctgcttgc acgtagatca cataagcacc 120
aagcgcggtg gcctcatgct tgaggagatt gatgagcggg gtggcaatgc cctgcctccg 180
gtgctcgcgg gagactgcca gatcatagat atagatctca ctacgcggct gctcaaactt 240
gggcagaacg taagccgcca gagcgccaac aaccgcttct tggtcgaagg cagcaagcgc 300
gatgaatgct ttactacgga gcaagttccc gaggtaatcg gagtccggct gatggtggga 360
gtaggtggct acgtcaccga actcaccgac gaaaagatca agagcagccc gcatggattt 420
gacttggtca gggccgagcc tacatgtgcg aatgatgccc atacttgagc cacctaactt 480
tgttttaggg cgactgccct gctgcgtaac atcgttgctg ctccataaca tcaaaccatcg 540
accacggcgg taacgcgctt gctgcttggg tgcccaggcc atagactgta caaaaaaaca 600
gtcataaaca gccatgaaaa ccgccactgc gccgttacca ccgctgcgct cgggtcaaggt 660
tctggaccag ttgcgtgagc gcattttttt ttctctctcg gcgtttacgc cccgccctgc 720
cactcatcgc agtactgttg taattcatta agcattctgc cgacatggaa gccatcacag 780
acggcatgat gaacctgaat cggcagcggc atcagcacct tgtcgccttg cgtataatat 840
ttgcccatag tgaaaacggg ggccaagaag ttgtccatat tggccacgct taaatcaaaa 900
ctgggtgaaac taccacggg attggcgctg acgaaaaaca tattctcaat aaacccttta 960
gggaaatagg ccaggttttc accgtaacac gccacatctt gcgaatatat gtgtagaaac 1020
tgccggaaat cgtcgtggta ttcactccag agcgatgaaa acgtttcagt ttgctcatgg 1080
aaaacgggtg aacaagggtg aacactatcc catatcacca gctcaccgct tttcattgcc 1140
atacggaaact ccggatgagc attcatcagg cgggcaagaa tgtgaataaa ggccggataa 1200
aacttggtgt tatttttctt tacgggtctt aaaaaggccg taatatccag ctgaacgggtc 1260
tggttatagg tacattgagc aactgactga aatgcctcaa aatgttcttt acgatgccat 1320
tgggatatat caacggtggt atatccagtg atttttttct ccattttttt ttctctcttt 1380
agaaaaactc atcgagcatc aaatgaaact gcaatttatt catatcagga ttatcaatac 1440
catatTTTTG aaaaagccgt ttctgtaatg aaggagaaaa ctaccgagg cagttccata 1500
ggatggcaag atcctggtat cggctctgca ttccgactcg tccaacatca atacaaccta 1560
ttaatttccc ctctcaaaa ataaggttat caagtgagaa atccatga gtgacgactg 1620

```

-continued

---

aatccggtga	gaatggcaaa	agtttatgca	tttctttcca	gacttggtca	acaggccagc	1680
cattacgctc	gtcatcaaaa	tactcgcac	caaccaaacc	gttattcatt	cgtgattgcg	1740
cctgagcgag	gcgaaatacg	cgatcgctgt	taaaaggaca	attacaaaca	ggaatcgagt	1800
gcaaccggcg	caggaacact	gccagcgcat	caacaatatt	ttcacctgaa	tcaggatatt	1860
cttctaatac	ctggaacgct	gtttttccgg	ggatcgcagt	ggtgagtaac	catgcatcat	1920
caggagtacg	gataaaatgc	ttgatggctg	gaagtggcat	aaattccgct	agccagttta	1980
gtctgaccat	ctcatctgta	acatcattgg	caacgctacc	tttgccatgt	ttcagaaaca	2040
actctggcgc	atcgggcttc	ccatacaagc	gatagattgt	cgcacctgat	tgcccacat	2100
tatcgcgagc	ccatttatac	ccatataaat	cagcatccat	gttggaaatt	aatcgcgccc	2160
tcgacgtttc	ccgttgaata	tggtcatttt	ttttttcttc	ctttaccaat	gcttaatcag	2220
tgaggcacct	atctcagcga	tctgtctatt	tcgttcatcc	atagttgcct	gactccccgt	2280
cgtgtagata	actacgatac	gggagggctt	accatctggc	cccagcctg	cgatgatacc	2340
gcgagaacca	cgctcaccgg	ctccggattt	atcagcaata	aaccagccag	ccggaagggc	2400
cgagcgcaga	agtggctctg	caactttatc	cgctccatc	cagtctatta	attggtgccg	2460
ggaagctaga	gtaagtagtt	cgccagttaa	tagtttgcgc	aacgttgttg	ccatcgctac	2520
aggcatcgtg	gtgtcacgct	cgctgtttgg	tatggcttca	ttcagctccg	gttcccaacg	2580
atcaaggcga	gttacctgat	ccccatggt	gtgcaaaaa	gcggttagct	ccttcggctc	2640
tccgatcgtt	gtcagaagta	agttggccgc	agtgttatca	ctcatggtta	tggcagcact	2700
gcataattct	cttactgtca	tgccatccgt	aagatgcttt	tctgtgactg	gtgagtactc	2760
aaccaagtca	ttctgagaat	agtgtatgcg	gcgaccgagt	tgctcttgcc	cggcgtcaat	2820
acgggataat	accgcgccac	atagcagaac	tttaaaagtg	ctcatcattg	gaaaacgttc	2880
ttcggggcga	aaactctcaa	ggatcttacc	gctgttgaga	tccagttcga	tgtaaccac	2940
tcgtgcaccc	aactgatctt	cagcatcttt	tactttcacc	agcgtttctg	ggtgagcaaa	3000
aacaggaagg	caaatgccc	caaaaaagg	aataagggcg	acacggaaat	gttgaatact	3060
catattcttc	ctttttcaat	attattgaag	catttatcag	ggttattgtc	tcatgagcgg	3120
atacatattt	gaatgtattt	agaaaaataa	acaaatagg	gtcagtggtta	caaccaatta	3180
accaattctg	aacattatcg	cgagcccatt	tatactgaa	tatggctcat	aacaccctt	3240
gtttgcctgg	cggcagtagc	gcggtggctc	cacctgacct	catgccgaac	tcagaagtga	3300
aacgcctag	cgccgatggt	agtgtgggga	ctccccatgc	gagagtaggg	aactgccagg	3360
catcaataaa	aacgaaaggc	tcagtcgaaa	gactgggctt	ttcgcccggg	ctaattgagg	3420
ggtgtcgccc	ttttgacgga	tatcaagctt	ctattaaccg	aacgcttctg	ccaagatttc	3480
ttgctgctcc	acggcatgaa	ccttcgagga	accgctgctc	ggcgcagaca	tcgcgcgacg	3540
ggagatacgc	ttgatgccc	ccaacttgtc	cggcagcaac	tccggcaact	ccaaccgaa	3600
acgcggccag	gcgccctggt	tcgctggttc	ctcctggacc	caaaagaact	ctttgacatt	3660
ttcgtaacga	tccagggttt	cacgcagacg	acgacgcggc	aacggcgcca	gttgctccag	3720
acgcacaatt	gccagatcat	tacggttgtc	tttcgccttg	cgtgcccga	attcataata	3780
caacttaccg	gaggtcaaca	gaatacggct	aactttattg	cggtcgccga	taccatcttc	3840
gtaggtcggg	tcctccagga	cgctacgga	tttaatctcg	gtaaatcct	taatctccga	3900

-continued

---

aaccgcccgc ttgtggcgca gcatggattt cggggtaaac acgatcaacg gacgttggat 3960  
gccgtccaac gcgtggcggc gcagcaagtg aaagtaattg gacgggggtgc tcggcatcgc 4020  
gatcgtcatg ctaccctcag cccacagctg caaaaagcgc tcaatgcgcg ccgacgtgtg 4080  
gtccgggccc tgaccctcgt ggccgtgtgg cagcaacaga acaacgttgg acagctgacc 4140  
ccatttcgct tcgcccgagc tgatgaactc gtcaatgatc gactggggcac cattaacaaa 4200  
gtcgccgaat tgcgcctccc acagcacaac cgcgtctgga ttgccaacgg tataaccgta 4260  
ctcaaaaccc acagccgcgt attccgacaa cggcgaatca taaaccagga acttaccacc 4320  
ggtcgggctg ccgtcgtctg tagtcgccag cagctgcagc ggggtgaact cctcgcgggt 4380  
gtgacggctg atcagcacag aatgacgctg gctaaagggt ccacgacgag agtcttgacc 4440  
gtcagacgg accagcttgc cctcagcaac caggctgccc agcgcagca actcaccaaa 4500  
cgcccagtca atcttaccct catacgccat ctcacgacgc tttccagca ccggctgaac 4560  
gcgcggtgt gccgtgaaac cgtaggcag gccaggaag gcatcaccga tacgtgccag 4620  
cagggatttg tcaacagcgg tcgccaggcc cgctgggatc atttgggtccg actcgcgct 4680  
ctccgacggg tggacgcccgt gcttctccag ttcgcgcact tcggtgaaca cacgctccag 4740  
ttggccctgg taatcgcgca gcgcacctc cgctcttcc atgctgatgt cgccacgacc 4800  
gatcagagcc tcgggtgtagg atttacgggc gccacgcttg gtgtccacga cgtcatagac 4860  
atacggattg gtcatagatg gatcgtcacc ctattatga ccacgacgac gatagcacag 4920  
catgtcaata acaacgtcct ttttaaagcg ttggcgggaag tcaactgcc aacgcgcaac 4980  
ccagacacac gcttctggat cgtcgcccgt cacgtggaag attggcgcac cgatcatctt 5040  
cgccacatcc gtgcagtact cgctcgaacg agagtattcc ggagcgggtg tgaagccgat 5100  
ttggttgctt acaatgatgt gaatcgtacc acccacacga tagccaggca gatttgccag 5160  
atcagcgtc tccgcaacaa cgcttgacc cgcaaggcc gcatcacctg gcagcatcaa 5220  
cggaacgacg gagaatgcac gttggccgtc ggagtcaatg gaaccgtggt ccagcaggtc 5280  
ttgcttogca cgcaccaaac cttccagcac tggatcgacc gcttccaaat gggacggatt 5340  
tgccgtcagg gaaacctgaa tatcgttatc gccaaacatt tgcagataca gaccgctcgc 5400  
accaggtgg tacttgacat cgccggaacc gtgagcctgg gacgggttca gattgccttc 5460  
aaactccgta aagatttgcg aatacggttt gccacgatg ttcgccagga cattcaagcg 5520  
accacggtgc ggcattgcaa tgaccacttc atccaaacca tgctcggcac attggtcaat 5580  
cgccgcgtcc atcattgga taacagattc cgcaccctcc aggtctaaac gcttttgccc 5640  
cacatacttg gtttgacgga aggtttcaaa cgctccgct gcggtcagtt tcgacaagat 5700  
gtacttttgt tgagcaacgg tcggtttgac gtgcttcgct tcgacacgct gctccagcca 5760  
ctccttttgt tccgggtcca gaatgtgcgc gtactcaaca ccgatgtgac ggcagtacgc 5820  
gtcgcgcagc aaaccagca cgtcacgcag ctttttgtat tgagcaccgg cgaaaccgct 5880  
aaccttaaag acgcggtcca ggtcccacag agtcaggcca tgcgtcaaca cctccaaatc 5940  
cggatgcgaa cgaaagcgcg ccttatccaa gcgcaacggg tcgggtgtccg ccatcagatg 6000  
gccgcggttg cgataggccg cgatcagggt catcacacgt gcgcttctgt caacgatcga 6060  
gtccggatta tcggtgctcc aacgcactgg caggtacggg atgctcagct cgcggaagac 6120  
ctcatcccag aagccatcag acaacagcag ttcatgaatg gtacgcagga agtcaccgct 6180



-continued

---

ttccgcacct	tgaatgatac	ggtggctgta	ggtagaagtc	agggtaatca	gtttgccaat	6240
accagctca	gcatgctgt	cctcggacgc	gccttgaac	tccgccgat	attccatcgc	6300
accgacaccg	atgatagcac	cttgacctgg	catcaggcgt	ggcacagagt	gcaccgtgcc	6360
aatcgtgccc	ggattcgtca	gcaaatcgt	aacgccagcg	aagtcctcgg	tggtcagttt	6420
accatcacga	gcacgacgga	cgatgtcctc	gtacgcggta	acgaactgcg	cgaagcgcac	6480
cgctctgcaa	cgtttgatac	cggccaccac	cagagagcgc	ttgccatctt	taccttgacg	6540
gtcaatagcc	agaccagat	tgggtgtgct	aggagtaacc	gccgtcggct	taccgtccac	6600
ctccgtgtag	tggcgattca	tattcgggaa	cttcttaacc	gcctgaacca	gagcataacc	6660
cagcaaatgg	gtaaagctga	ttttaccacc	acgcgtgcgt	ttcaactgat	tattgatcac	6720
gatacgatta	tcgatcaaca	atttcgctgg	cacagcacgc	accgaggtcg	ccgtaggcac	6780
ttccagcgac	gcgctcatgt	tcttcacgac	agcagccgcc	gcaccacgca	ggacagcaac	6840
ttcatcgctt	tcggctggcg	gaggcaccgc	ggtcttggcg	gccagagccg	ccacgacacc	6900
gttgcocgct	gcggccgtgt	cggccggttt	cggcggcgt	tgcggagccg	ccgctgccgc	6960
acgctcagcg	accaaagggc	tggtcacacg	agtcggctca	gcagccgggt	gggaagtcgg	7020
ctccgggcta	tagtccacca	gaaactcatg	ccagcttggg	tcaacggaag	acggatcatc	7080
acgaaattta	cgatacatac	caacacgaga	cgggtcctga	gtcaccatgg	atatatctcc	7140
ttcttaaaga	attcgatatc	tcagcgacaa	ggcgacaca	aaatttattc	taaatgcata	7200
ataaatactg	ataacatctt	atagtttgta	ttatattttg	tattatcgtt	gacatgtata	7260
atthtgatat	caaaaactga	ttttcccttt	attatthtcg	agatttattt	tcttaattct	7320
ctttaacaaa	ctagaaatat	tgtatataca	aaaaatcata	aataatagat	gaatagttta	7380
attataggtg	ttcatcaatc	gaaaaagcaa	cgtatcttat	ttaaagtgcg	ttgctttttt	7440
ctcatttata	aggttaaata	attctcatat	atcaagcaaa	gtgacaggcg	cccttaaata	7500
ttctgacaaa	tgctctttcc	ctaaactccc	cccataaaaa	aaccgcccca	agcggggttt	7560
tacgttattt	gcgattaac	gattactcgt	tatcagaacc	gccagggggg	cccagactta	7620
agactggccg	tcgthttaca	acacagaaaag	agthttgtaga	aacgcaaaaa	ggccatccgt	7680
caggggcctt	ctgcttagtt	tgatgcctgg	cagttcccta	ctctcgcctt	ccgcttccct	7740
gctcaactgac	tcgctgcgct	cggctcgttcg	gctgcggcga	gcggtatcag	ctcaactcaa	7800
ggcggtaata	cggttatcca	cagaatcagg	ggataacgca	ggaaagaaca	tgtgagcaaa	7860
aggccagcaa	aaggccagga	accgtaaaaa	ggccgcgttg	ctggcgthtt	tccataggtt	7920
ccgccccctt	gacgagcatc	acaaaaatcg	acgctcaagt	cagaggtggc	gaaaccggac	7980
aggactataa	agataccagg	cgtttccccc	tggaaagctcc	ctcgtgcgct	ctcctgttcc	8040
gacctgccc	cttaccggat	acctgtccgc	ctttctccct	tcgggaagcg	tggcgctttc	8100
tcatagetca	cgctgtaggt	atctcagttc	ggtgtaggtc	gttcgctcca	agctgggctg	8160
tgtgcacgaa	cccccgctc	agcccgaccg	ctgcgcctta	tccggtaact	atcgtcttga	8220
gtccaaccgg	gtaagacacg	acttatcgcc	actggcagca	gccactggta	acaggattag	8280
cagagcgagg	tatgtaggcg	gtgctacaga	gttcttgaag	tggtagggcta	actacggcta	8340
cactagaaga	acagtatttg	gtatctgcgc	tctgctgaag	ccagttacct	tcggaaaaag	8400
agttgtagc	tcttgatccg	gcaaacaaac	caccgctggt	agcggtggtt	ttttgtttg	8460

-continued

---

caagcagcag attacgcgca gaaaaaaagg atctcaagaa gatcctttga tcttttctac 8520  
 ggggtctgac gctcagtga acgacgcgcg cgtaactcac gttaaggat tttggtcag 8580  
 agcttgccgc gtccegtcaa gtcagcgtaa tgctctgctt a 8621

<210> SEQ ID NO 41  
 <211> LENGTH: 4401  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic  
 plasmid pTrc-HisB

<400> SEQUENCE: 41

gtttgacagc ttatcatcga ctgcacgggtg caccaatgct tctggcgtca ggcagccatc 60  
 ggaagctgtg gtatggctgt gcaggctgta aatcactgca taattcgtgt cgctcaaggc 120  
 gcaactccgt tctggataat gttttttgcg ccgacatcat aacggttctg gcaaatattc 180  
 tgaaatgagc tgttgacaat taatcatccg gctcgtataa tgtgtggaat tgtgagcgga 240  
 taacaatttc acacaggaaa cagcgccgct gagaaaaagc gaagcggcac tgctctttaa 300  
 caatttatca gacaatctgt gtgggcactc gaccggaatt atcgattaac tttattatta 360  
 aaaattaaag aggtatata taatgtatcg attaaataag gaggaataaa ccatgggggg 420  
 ttctcatcat catcatcatc atggatggc tagcatgact ggtggacagc aaatgggtcg 480  
 ggatctgtac gacgatgacg ataaggatcc gagctcgaga tctgcagctg gtaccatag 540  
 ggaattcgaa gcttggctgt tttggcggat gagagaagat tttcagcctg atacagatta 600  
 aatcagaacg cagaagcggc ctgataaaaac agaatttgcc tggcggcagc agcgcgggtg 660  
 tcccacctga ccccatgccg aactcagaag tgaaacgccg tagcgcgat ggtagtgtgg 720  
 ggtctcccca tgcgagagta gggaaactgcc aggcataaaa taaaacgaaa ggctcagtcg 780  
 aaagactggg cctttgctt tatctgttgt ttgtcgggta acgctctcct gagtaggaca 840  
 aatccgccgg gagcggattt gaacgcttgc aagcaacggc ccggagggtg gcgggcagga 900  
 cgcccgccat aaactgccag gcatcaaatt aagcagaagg ccatcctgac ggatggcctt 960  
 tttgcgtttc taaaaactct ttttgtttat ttttctaaat acattcaaat atgtatccgc 1020  
 tcatgagaca ataaccctga taaatgcttc aataatattg aaaaaggaag agtatgagta 1080  
 ttcaacattt ccgtgtcgcc cttattccct tttttgcggc attttgcctt cctgtttttg 1140  
 ctcaccaga aacgctggtg aaagtaaaag atgctgaaga tcagttgggt gcacgagtg 1200  
 gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac 1260  
 gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccggttg 1320  
 acgcccggca agagcaactc ggtcgcccga tacactatc tcagaatgac ttggttgagt 1380  
 actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg 1440  
 ctgccataac catgagtgat aacctgccc ccaacttact tctgacaacg atcggaggac 1500  
 cgaaggagct aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt 1560  
 gggaaaccgga gctgaatgaa gccatacaca acgacgagcg tgacaccag atgcctgtag 1620  
 caatggcaac aacggtgccc aaactattaa ctggcgaact acttacteta gcttcccggc 1680  
 aacaattaat agactggatg gaggcggata aagttgcagg accacttctg cgctcggccc 1740

-continued

---

ttccggctgg	ctggtttatt	gctgataaat	ctggagccgg	tgagcgtggg	tctcgcggta	1800
tcattgcagc	actggggcca	gatggtaagc	cctcccgtat	cgtagttatc	tacacgacgg	1860
ggagtccaggc	aactatggat	gaacgaaata	gacagatcgc	tgagataggt	gcctcactga	1920
ttaagcattg	gtaactgtca	gaccaagttt	actcatatat	acttttagatt	gattttaaac	1980
ttcattttta	attttaaagg	atctaggtga	agatcctttt	tgataatctc	atgacaaaa	2040
tcccttaacg	tgagttttcg	ttccactgag	cgtcagaccc	cgtagaaaag	atcaaaggat	2100
cttcttgaga	tcctttttt	ctgcgcgtaa	tctgctgctt	gcaaacaaaa	aaaccaccgc	2160
taccagcggg	ggtttgttg	ccggatcaag	agctaccaac	tctttttccg	aaggtaactg	2220
gcttcagcag	agcgcagata	caaataactg	tccttctagt	gtagccgtag	ttaggccacc	2280
acttcaagaa	ctctgtagca	ccgcctacat	acctcgctct	gctaactctg	ttaccagtgg	2340
ctgctgccag	tggcgataag	tcgtgtctta	ccgggttggg	ctcaagacga	tagttaccgg	2400
ataaggcgca	gcggtcgggc	tgaacggggg	gttcgtgcac	acagcccagc	ttggagcgaa	2460
cgacctacac	cgaactgaga	tacctacagc	gtgagctatg	agaaagcgc	acgcttcccg	2520
aagggagaaa	ggcggacagg	tatccggtaa	gcggcagggt	cggaacagga	gagcgcacga	2580
gggagcttcc	agggggaac	gcctgggtatc	tttatagtcc	tgctcgggtt	cgccacctct	2640
gacttgagcg	tcgattttg	tgatgctcgt	cagggggcg	gagcctatgg	aaaaacgcca	2700
gcaacgcggc	ctttttacgg	ttcctggcct	tttgctggcc	ttttgctcac	atgttctttc	2760
ctgcgttata	ccctgattct	gtggataacc	gtattaccgc	ctttgagtga	gctgataccg	2820
ctcgcgcag	ccgaacgacc	gagcgcagcg	agtcagttag	cgaggaagcg	gaagagcgcc	2880
tgatgcggta	ttttctcctt	acgcatctgt	gcggtatttc	acaccgata	tggtgcactc	2940
tcagtacaat	ctgctctgat	gccgcatagt	taagccagta	tacactccgc	tatcgctacg	3000
tgactgggtc	atggctgccc	cccgcacccc	gccaaccccc	gctgacgcgc	cctgacgggc	3060
ttgtctgctc	ccggcatccg	cttacagaca	agctgtgacc	gtctccggga	gctgcatgtg	3120
tcagaggttt	tcaccgtcat	caccgaaacg	cgcgaggcag	cagatcaatt	cgcgcgcgaa	3180
ggcgaagcgg	catgcattta	cgttgacacc	atcgaatggc	gcaaacctt	tcgcggtatg	3240
gcatgatagc	gcccggaaga	gagtcaattc	agggtggtga	atgtgaaacc	agtaacgtta	3300
tacgatgtcg	cagagtatgc	cgggtgtctct	tatcagaccg	tttcccgcgt	ggtgaaccag	3360
gccagcccag	tttctgcgaa	aacgcgggaa	aaagtgggag	cgccgatggc	ggagctgaat	3420
tacattocca	accgcgtggc	acaacaactg	gcgggcaaac	agtcgctgct	gattggcggt	3480
gccacctcca	gtctggccct	gcacgcgcgc	tcgcaaattg	tcgcgccgat	taaatctcgc	3540
gccgatcaac	tgggtgccag	cgtgggtggg	tcgatggtag	aacgaagcgg	cgctgaagcc	3600
tgtaaagcgg	cgggtgcacaa	tcttctcgcg	caacgcgtca	gtgggctgat	cattaactat	3660
ccgctggatg	accaggatgc	cattgctgtg	gaagctgctt	gcactaatgt	tccggcgttt	3720
tttcttgatg	tctctgacca	gacacccatc	aacagtatta	tttctocca	tgaagacggt	3780
acgcgactgg	gcgtggagca	tctggctgca	ttgggtcacc	agcaaatcgc	gctgttagcg	3840
ggcccattaa	gttctgtctc	ggcgcgtctg	cgtctggctg	gctggcataa	atatctcact	3900
cgcaatcaaa	ttcagccgat	agcggaaacg	gaaggcgact	ggagtgccat	gtccggtttt	3960
caacaaacca	tgcaaatgct	gaatgagggc	atcgttccca	ctgcgatgct	ggttgccaac	4020

-continued

---

```

gatcagatgg cgctgggccc aatgcgcgcc attaccgagt cggggtgcg cggtgggtgcg 4080
gatatctcgg tagtgggata cgacgatacc gaagacagct catgttatat cccgccgtca 4140
accaccatca aacaggattt tcgcctgctg gggcaaacca gcgtggaccg cttgctgcaa 4200
ctctctcagg gccaggcggg gaagggcaat cagctgttgc ccgtctcact ggtgaaaaga 4260
aaaaccaccc tggcgcccaa tacgcaaacc gcctctcccc gcgcgttggc cgattcatta 4320
atgcagctgg cacgacaggt ttcccgactg gaaagcgggc agtgagcgca acgcaattaa 4380
tgtgagttag cgcaattga g 4401

```

```

<210> SEQ ID NO 42
<211> LENGTH: 25
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 42

```

```

gggatatcat gagctatacc gttgg 25

```

```

<210> SEQ ID NO 43
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 43

```

```

gaaatagttc tctagagaag ctcc 24

```

```

<210> SEQ ID NO 44
<211> LENGTH: 6123
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        plasmid pTrc:pdc-his

```

```

<400> SEQUENCE: 44

```

```

gtttgacagc ttatcatcga ctgcacggtg caccaatgct tctggcgtca ggcagccatc 60
ggaagctgtg gtatggctgt gcaggctgta aatcactgca taattegtgt cgctcaaggc 120
gcactcccgt tctggataat gttttttgcg ccgacatcat aacggttctg gcaaatattc 180
tgaaatgagc tgttgacaat taatcatccg gctcgtataa tgtgtggaat tgtgagcgga 240
taacaatttc acacaggaaa cagcgcgct gagaaaaagc gaagcggcac tgctctttaa 300
caatttatca gacaatctgt gtgggcactc gaccggaatt atcgattaac tttattatta 360
aaaattaaag aggtatataat taatgtatcg attaaataag gaggaataaa ccatgagcta 420
taccgttggc acctatctgg cagaacgcct ggtgcagatc ggcctgaaac accacttcgc 480
ggtggcgggt gactacaacc tgggtgctgct ggacaacttg ctggtgaaca agaatatgga 540
acaggtgtac tgctgtaatg agctgaattg tggcttcagc gcggagggtt acgcccgggc 600
caaaggtgca gccgcagcgg ttgtgacctg tagcgttggc gcgctgagcg catttgatgc 660
gattggtggt gcgtacgctg agaatttgcc ggtcatcctg atcagcggcg caccgaacaa 720
caacgaccat gctgcgggcc acgtgctgca tcacgcgctg ggtaagaccg actatcatta 780

```

-continued

---

tcagttggag	atggcgaaga	atatcacggc	cgctgcggaa	gctatctata	cgccggaaga	840
ggctcctgca	aaaatcgatc	acgttatcaa	aaccgcactg	cgtgagaaga	agccagtgta	900
cctggagatt	gcctgcaaca	ttgcttctat	gccgtgcgca	gcaccgggtc	ctgcgagcgc	960
gttgtttaac	gatgaggcca	gcgacgaagc	gagcctgaac	gcggctgttg	aagaaacgct	1020
gaaattcatt	gcaaatcgtg	ataaggttgc	tgtcctgggt	ggcagcaagc	tgcgtgccgc	1080
aggcgcggaa	gaagcagcgg	tgaagttcgc	agatgccctg	ggtggcgcgg	tcgcaacat	1140
ggctgccgca	aaatctttct	ttccggaaga	gaatccacac	tatatcggca	ctagctgggg	1200
tgaggtgtct	tatccgggcg	tcgagaaaac	catgaaagaa	gcggatgcgg	tgatcgcct	1260
ggctccgggt	ttcaacgatt	acagcaccac	tggtcggacg	gatattccgg	acccgaaaa	1320
gctggtgctg	gcggagccgc	gtagcgtggg	ggtcaatggg	atccgttttc	cgcccgcca	1380
tctgaaggac	tacctgacc	gcctggcgca	gaaggttagc	aagaaaacgg	gtgcgctgga	1440
tttcttcaaa	tcgctgaatg	caggtgaact	gaaaaaggca	gccccagcag	atccgtccgc	1500
accgctgggtg	aatgcggaga	tcgcgctca	agtcgaagct	ctgctgacgc	cgaacacgac	1560
cgttatcgcg	gagacgggtg	acagctgggt	taacgcgcaa	cgtatgaagt	tgccgaatgg	1620
tgctcgagtt	gagtacgaga	tgcaatgggg	ccacattggc	tggtccgtgc	ctgcagcctt	1680
tggttacgcc	gttggtgcc	cggagcgtcg	caacattctg	atggtgggtg	acggtagctt	1740
tcaactgacc	gccaagagg	tcgctcagat	ggtccgcctg	aaactgcctg	tcatcatttt	1800
cctgatcaat	aactacggtt	acacgatcga	ggttatgatc	catgatggcc	cgtacaacaa	1860
cattaagaat	tgggactacg	cgggtttgat	ggaggttttc	aatggcaacg	gtggttatga	1920
ctccggtgcg	ggtaaaggtc	tgaaagcaaa	gacgggtggc	gagctcgag	aggcgattaa	1980
ggttgcgctg	gcgaataccg	acggccccgac	cttgattgag	tgtttcattg	gtcgcgagga	2040
ttgcaccgaa	gaactggtca	aatggggcaa	gcgcgtcgca	gccgcgaatt	cccgtaaacc	2100
ggttaataag	cttctcggcc	atcaccatca	ccatcactag	aagcttctct	aggtatggct	2160
agcatgactg	gtggacagca	aatgggtcgg	gatctgtacg	acgatgacga	taaggatcga	2220
tggggatccg	agctcgagat	ctgcagctgg	taccatatgg	gaattcgaag	cttggctggt	2280
ttggcggatg	agagaagatt	ttcagcctga	tacagattaa	atcagaacgc	agaagcggtc	2340
tgataaaaca	gaatttgctt	ggcggcagta	gcgcggtggg	cccacctgac	cccattgccga	2400
actcagaagt	gaaacgccgt	agcgcctgatg	gtagtgtggg	gtctccccat	gcgagagtag	2460
ggaactgcc	ggcatcaaat	aaaacgaaag	gctcagtcga	aagactgggc	ctttcgtttt	2520
atctgttggt	tgtcggtgaa	cgctctcctg	agtaggacaa	atccgccggg	agcggatttg	2580
aacggtgcga	agcaacggcc	cggaggggtg	cgggcaggac	gcccgcata	aactgccagg	2640
catcaaat	agcagaaggc	catcctgacg	gatggccttt	ttgcgtttct	acaaactctt	2700
tttgtttatt	tttctaaata	cattcaata	tgtatccgct	catgagacaa	taaccctgat	2760
aaatgcttca	ataatattga	aaaaggaaga	gtatgagtat	tcaacatttc	cgtgtcgcct	2820
ttattccctt	tttgccgca	ttttgccttc	ctgtttttgc	tcaccagaa	acgctgggtga	2880
aagtaaaaga	tgctgaagat	cagttgggtg	cacgagtggt	ttacatcgaa	ctggatctca	2940
acagcggtaa	gatccttgag	agttttcgcc	ccgaagaacg	ttttccaatg	atgagcactt	3000
ttaaagttct	gctatgtggc	gcggtattat	cccgtgttga	cgccgggcaa	gagcaactcg	3060

-continued

---

gtcgcgcat	acactattct	cagaatgact	tggttgagta	ctcaccagtc	acagaaaagc	3120
atcttacgga	tggcatgaca	gtaagagaat	tatgcagtgc	tgccataacc	atgagtgata	3180
acactgcggc	caacttactt	ctgacaacga	tcgaggagacc	gaaggagcta	accgcttttt	3240
tgacacaacat	gggggatcat	gtaactcgcc	ttgatcgttg	ggaaccggag	ctgaatgaag	3300
ccataccaaa	cgacgagcgt	gacaccacga	tgctgtagc	aatggcaaca	acgttgcgca	3360
aactattaac	tggcgaacta	cttactctag	cttcccggca	acaattaata	gactggatgg	3420
aggcggataa	agttgcagga	ccacttctgc	gctcggccct	tccggctggc	tggtttattg	3480
ctgataaatc	tggagccggt	gagcgtgggt	ctcgcggtat	cattgcagca	ctggggccag	3540
atggtaagcc	ctcccgtatc	gtagttatct	acacgacggg	gagtcaggca	actatggatg	3600
aacgaaatag	acagatcgct	gagatagggt	cctcactgat	taagcattgg	taactgtcag	3660
accaagttta	ctcatatata	ctttagattg	atttaaaact	tcatttttaa	tttaaaagga	3720
tctaggtgaa	gatccttttt	gataatctca	tgacccaaaat	cccttaacgt	gagttttcgt	3780
tccactgagc	gtcagacccc	gtagaaaaga	tcaaaggatc	ttcttgagat	cctttttttc	3840
tgcgcgtaat	ctgctgcttg	caaacaaaaa	aaccaccgct	accagcgggtg	gtttgtttgc	3900
cggatcaaga	gctaccaact	ctttttccga	aggtaactgg	cttcagcaga	gcgcagatac	3960
caaatactgt	ccttctagtg	tagccgtagt	taggccacca	cttcaagaac	tctgtagcac	4020
cgctacata	cctcgtcttg	ctaatacctgt	taccagtggc	tgctgccagt	ggcgataagt	4080
cgtgtcttac	cgggttgac	tcaagacgat	agttaccgga	taaggcgcag	cggtcgggct	4140
gaacggggggg	ttcgtgcaca	cagcccagct	tggagcgaac	gacctacacc	gaactgagat	4200
acctacagcg	tgagctatga	gaaagcgcca	cgcttcccga	aggagaaaag	gcggacaggt	4260
atccggtaag	cggcagggtc	ggaacaggag	agcgcacgag	ggagcttcca	gggggaaacg	4320
cctggtatct	ttatagtcct	gtcgggtttc	gccacctctg	acttgagcgt	cgatttttgt	4380
gatgctcgtc	agggggggcg	agcctatgga	aaaacgccag	caacgcggcc	tttttacggt	4440
tcttgccctt	ttgctggcct	tttgetcaca	tgtctttcc	tgcgttatcc	cctgattctg	4500
tggataaccg	tattaccgcc	tttgagtgag	ctgataccgc	tcgccgcagc	cgaacgaccg	4560
agcgcagcga	gtcagtgagc	gaggaagcgg	aagagcgccct	gatgcggtat	tttctcctta	4620
cgcatctgtg	cggatattca	caccgcatac	ggtgcactct	cagtacaatc	tgctctgatg	4680
ccgcatagtt	aagccagtat	acactccgct	atcgctacgt	gactgggtca	tggtgcgcc	4740
ccgacacccg	ccaacacccg	ctgacgcgcc	ctgacgggct	tgtctgctcc	cggcatccgc	4800
ttacagacaa	gctgtgaccg	tctccgggag	ctgcatgtgt	cagaggtttt	caccgtcatc	4860
accgaaaacgc	gcgaggcagc	agatcaattc	gcgcgcgaag	gcgaagcggc	atgcatttac	4920
ggtgacacca	tcgaatggcg	caaaaccttt	cgcggtatgg	catgatagcg	cccggaagag	4980
agtcaattca	gggtggtgaa	tgtgaaacca	gtaacgttat	acgatgtcgc	agagtatgcc	5040
ggtgtctctt	atcagaccgt	ttcccgcgtg	gtgaaccagg	ccagccacgt	ttctgcgaaa	5100
acgcgggaaa	aagtggaagc	ggcgatggcg	gagctgaatt	acattcccaa	ccgcgtggca	5160
caacaactgg	cgggcaaacg	gtcgttgctg	attggcggtg	ccacctccag	tctggccctg	5220
cacgcgcgct	cgcaaatgtg	cgcggcgatt	aaatctcgcg	ccgatcaact	gggtgccagc	5280
gtggtggtgt	cgatggtaga	acgaagcggc	gtcgaagcct	gtaaagcggc	ggtgcacaat	5340

-continued

---

```

cttctcgcgc aacgcgtcag tgggctgac attaactatc cgctggatga ccaggatgcc 5400
attgctgtgg aagctgctg cactaatggt cggcggttat ttcttgatgt ctctgaccag 5460
acacccatca acagtattat tttctcccat gaagacggta cgcgactggg cgtggagcat 5520
ctggtcgcat tgggtcacca gcaaactcgc ctgttagcgg gccattaag ttctgtctcg 5580
gcgctgtctg gtctggtggt ctggcataaa tatctcactc gcaatcaaat tcagccgata 5640
gcggaacggg aaggcgactg gagtgccatg tccggttttc aacaaacct gcaaatgctg 5700
aatgagggca tcgttccac tgcgatgctg gttgccaacg atcagatggc gctgggagca 5760
atgcgcgcca ttaccgagtc cgggctgcgc gttggtgcgg atatctcggg agtgggatac 5820
gacgataccg aagacagctc atggtatata ccgcccga ccaccatcaa acaggatatt 5880
cgctgtctgg ggcaaaccag cgtggaccgc ttgctgcaac tctctcaggg ccaggcgggtg 5940
aagggcaatc agctgttgcc cgtctcactg gtgaaaagaa aaaccacct ggcgccaat 6000
acgcaaaccg cctctccccg cgcgttgccc gattcattaa tgcagctggc acgacagggt 6060
tcccgactgg aaagcgggca gtgagcgcaa cgcaattaat gtgagttagc gcgaattgat 6120
ctg 6123

```

```

<210> SEQ ID NO 45
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 45

```

```

ggagaattac catggcggat accctg 26

```

```

<210> SEQ ID NO 46
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 46

```

```

gggaatctag actaatgatg atggtgg 27

```

```

<210> SEQ ID NO 47
<211> LENGTH: 5787
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        plasmid pTrc:oat-his

```

```

<400> SEQUENCE: 47

```

```

gtttgacagc ttatcatcga ctgcacgggt caccaatgct tctggcgtca ggcagccatc 60
ggaagctgtg gtatggctgt gcaggtcgta aatcactgca taattcgtgt cgctcaaggc 120
gcactcccgt tctggataat gttttttgcg ccgacatcat aacggttctg gcaaatatc 180
tgaaatgagc tgttgacaat taatcatccg gctcgtataa tgtgtggaat tgtgagcgga 240
taacaatttc acacagaaa cagaccatgg cggataccct gaccttcaac accaaacatt 300

```

-continued

---

tgttgaagc	gctgtttgaa	agcgggtatc	gtcaattcat	tgttttctccg	ggttctcgta	360
gcaccccgat	tgactgctg	ctggcagaat	acgcagagca	gaataacgag	atcaagctgt	420
tcgttgacgt	ggatgagcgc	agcgcgggtt	tcttcgcaact	gggcatcgcg	aaaacggtgc	480
tggagccggt	tgttttgctg	ggtacctcgg	gtaccgccat	cgctgaatac	atgctcgcg	540
tggcgggaagc	ctatgcggcg	aatatcccgc	tggtcgtggt	gagcaccgat	cgccacag	600
aattgcagtt	caacggcgca	ccgcaaacta	ttccgcaaag	caacctgttc	ggccaactga	660
cgaaacaggc	agtcctgatt	cgctgcaag	acatgcacag	cgatgtgacc	gagtacatcg	720
actttatcgt	tcagaaggtc	gttaatctgt	ccattaccgc	gcctcgtggc	ccgattcaga	780
tcaacctgcc	gctgcgcaa	ccgctgatgc	cggtcctgga	tcgtcaagac	gaggtgcacg	840
ttcagcgtgt	cgtgtttgac	aaacaaagcg	tgcaatatcg	cttgccgccc	attacggcga	900
aacgcctgct	gatcctggca	ggtccgaatg	tctgaatag	ctacgatgat	agcctgaaga	960
agtttgcat	caaaaacaat	gttccagtta	tcgctgatgt	tctgtccca	agccgccaca	1020
cgtataccat	tcacgtatt	gacgttctgc	tgcaagccca	caagatcaac	gccgacctga	1080
aaccggatct	ggtggttcgt	tttggaaga	cgccggttag	cgccgctgtc	ctgcaatggc	1140
tgaagaaga	gaacattctg	acctggcacg	tggacgagga	cgccagcgtg	gatcacaccc	1200
gtcatatcgt	ccgtgccatc	aagatggcgc	ctcatgattt	tctggaatcc	atgcatctga	1260
cgttgagcaa	aaaccagatt	gacttcaacc	agaagtggct	gtctctgccg	aaagtgatca	1320
aaactcgtaa	cgagatgaat	atcatcaccg	cattggacga	cgccgtccc	gatgacactc	1380
acatcttcgt	ggcgaacagc	atgccgattc	gtgacatgga	caactttttc	acgggtaatc	1440
accccagcg	catttacgct	aatcgtggtg	caaatggat	tgatggcgtg	attagctccg	1500
ccctgggcat	gagcgcggtc	gtcaagcaac	gttcggtcct	gctgacgggt	gacctgacgc	1560
tgtttcatga	catgaacggt	ctgatgatgg	cgaagaatta	ccagctgccg	ctggacatca	1620
ttgtgattaa	caacaatggc	ggtggcattt	tcagcttctt	gccacaagcc	ggtgctccga	1680
agtactttga	gcagctgttc	ggtaccccgc	tgaacattga	catcaaaaag	attgctgacc	1740
tgtattacat	cgattaccat	cagttgaatg	tgccggaggc	cctgagccag	atcctgcaga	1800
cgccgagcaa	aaccaccgc	ctgatcgagt	ataagtccga	tcaccagcgt	aaccgtgacg	1860
atcatcgcga	agtgctggag	atgctgaaac	accaccatca	tcattagtct	agagtcgacc	1920
tgcaggcatg	caagcttggc	tgttttggcg	gatgagagaa	gattttcagc	ctgatacaga	1980
ttaaatcaga	acgcagaagc	ggtctgataa	aacagaattt	gcctggcggc	agtagcgcgg	2040
tggccccacc	tgaccccatg	ccgaactcag	aagtgaaacg	ccgtagcggc	gatggttagtg	2100
tggggtctcc	ccatgcgaga	gtagggaaact	gccaggcatc	aaataaaacg	aaaggctcag	2160
tcgaaagact	gggcctttcg	ttttatctgt	tgtttgctgg	tgaacgctct	cctgagtagg	2220
acaaatccgc	cgggagcggg	tttgaacggt	gcgaagcaac	ggcccggagg	gtggcgggca	2280
ggacgcccgc	cataaactgc	caggcatcaa	attaagcaga	aggccatcct	gacggatggc	2340
ctttttgcgt	ttctacaaac	tctttttggt	tatttttcta	aatacattca	aatatgtatc	2400
cgctcatgag	acaataacc	tgataaatgc	ttcaataata	ttgaaaagg	aagagtatga	2460
gtattcaaca	tttcctgttc	gcccttattc	ccttttttgc	ggcattttgc	cttctgttt	2520
ttgctcacc	agaaacgctg	gtgaaagtaa	aagatgctga	agatcagttg	ggtgcacgag	2580



-continued

---

tgggttacat	cgaactggat	ctcaacagcg	gtaagatcct	tgagagtttt	cgccccgaag	2640
aacgttttcc	aatgatgagc	acttttaaag	ttctgctatg	tggcgcggta	ttatcccgtg	2700
ttgacgccgg	gcaagagcaa	ctcggtcgcc	gcatacacta	ttctcagaat	gacttggttg	2760
agtactcacc	agtcacagaa	aagcatctta	cggatggcat	gacagtaaga	gaattatgca	2820
gtgctgccat	aacctagagt	gataaacctg	cggccaactt	acttctgaca	acgatcggag	2880
gaccgaagga	gctaaccgct	tttttgcaca	acatggggga	tcatgtaact	cgcttgatc	2940
gttgggaacc	ggagctgaat	gaagccatac	caaacgacga	gcgtgacacc	acgatgccta	3000
cagcaatggc	aacaacgttg	cgcaaactat	taactggcga	actacttact	ctagcttccc	3060
ggcaacaatt	aatagactgg	atggaggcgg	ataaagttgc	aggaccactt	ctgcgctcgg	3120
cccttcgggc	tggttggttt	attgctgata	aatctggagc	cggtgagcgt	gggtctcgcg	3180
gtatcattgc	agcactgggg	ccagatggta	agccctcccg	tatcgtagtt	atctacacga	3240
cggggagtca	ggcaactatg	gatgaacgaa	atagacagat	cgctgagata	ggtgcctcac	3300
tgattaagca	ttggtaactg	tcagaccaag	tttactcata	tatactttag	attgatttaa	3360
aacttcattt	ttaatttaa	aggatctagg	tgaagatcct	ttttgataat	ctcatgacca	3420
aaatccctta	acgtgagttt	tcgttccact	gagcgtcaga	ccccgtagaa	aagatcaaag	3480
gatcttcttg	agatcctttt	tttctgcgcg	taatctgctg	cttgcaaaca	aaaaaacacc	3540
cgctaccagc	ggtggtttgt	ttgccggatc	aagagctacc	aactcttttt	ccgaaggtaa	3600
ctggcttcag	cagagcgcag	ataccaaata	ctgtccttct	agtgtagccg	tagttaggcc	3660
accacttcaa	gaactctgta	gcaccgccta	catacctcgc	tctgctaata	ctgttaccag	3720
tggtctgctg	cagtggcgat	aagtcgtgtc	ttaccggggt	ggactcaaga	cgatagttac	3780
cggataaggc	gcagcggtcg	ggctgaacgg	ggggttcgtg	cacacagccc	agcttgagac	3840
gaacgacct	caccgaactg	agatacctac	agcgtgagct	atgagaaagc	gccacgcttc	3900
ccgaagggag	aaaggcggac	aggatccggg	taagcggcag	ggtcgggaaca	ggagagcgca	3960
cgagggagct	tccaggggga	aacgcctggg	atctttatag	tccgtcggg	tttcgccacc	4020
tctgacttga	gcgctgattt	ttgtgatgct	cgtcaggggg	gcgagccta	tggaaaaacg	4080
ccagcaacgc	ggccttttta	cggttcctgg	ccttttctg	gccttttctg	cacatgttct	4140
ttcctgcggt	atcccctgat	tctgtggata	accgtattac	cgcttttgag	tgagctgata	4200
ccgctcgccg	cagccgaacg	accgagcgca	gcgagtcagt	gagcgaggaa	gcggaagagc	4260
gcctgatgcg	gtattttctc	cttacgcata	tgtcgggtat	ttcacaccgc	atatggtgca	4320
ctctcagtac	aatctgctct	gatgccgcat	agttaagcca	gtatacactc	cgctatcgct	4380
acgtgactgg	gtcatggctg	cgccccgaca	cccgccaaaca	cccgctgacg	cgccctgacg	4440
ggcttgctctg	ctcccggcat	ccgcttacag	acaagctgtg	accgtctccg	ggagctgcat	4500
gtgtcagagg	ttttcaccgt	catcaccgaa	acgcgcgagg	cagcagatca	attcgcgcgc	4560
gaaggcgaag	cggcatgcat	ttacgttgac	accatcgaat	ggtgcaaac	ctttcgcggt	4620
atggcatgat	agcgcgccga	agagagtcaa	ttcaggggtg	tgaatgtgaa	accagtaacg	4680
ttatacgatg	tcgcagagta	tgccgggtgc	tcttatcaga	ccgtttcccg	cgtgggtgaa	4740
caggccagcc	acgtttctgc	gaaaacgcgg	gaaaaagtgg	aagcggcgat	ggcggagctg	4800
aattacattc	ccaaccgctg	ggcacaacaa	ctggcgggca	aacagtcggt	gctgattggc	4860

-continued

---

```

gttgccacct ccagtctggc cctgcacgcg ccgtcgcaaa ttgtcgcggc gattaaatct 4920
cgcgccgatc aactgggtgc cagcgtggtg gtgtcgatgg tagaacgaag cggcgtcgaa 4980
gcctgtaaag cggcgggtgca caatcttctc gcgcaacgcg tcagtgggct gatcattaac 5040
tatccgctgg atgaccagga tgccattgct gtggaagctg cctgcactaa tgttccggcg 5100
ttattttctg atgtctctga ccagacaccc atcaacagta ttattttctc ccatgaagac 5160
ggtacgcgac tgggcgtgga gcactctggc gcattgggtc accagcaaat cgcgctgtta 5220
gcgggccccat taagttctgt ctcggcgctg ctgctctctg ctggctggca taaatatctc 5280
actcgcaatc aaattcagcc gatagcggaa cgggaaggcg actggagtgc catgtccggt 5340
tttcaacaaa ccatgcaaat gctgaatgag ggcacgttc cactgcat gctggttgcc 5400
aacgatcaga tggcgtggg cgcaatgccc gccattaccg agtccgggct gcgcttggt 5460
gcggatatct cggtagtggg atacgacgat accgaagaca gctcatgtta tatcccgccg 5520
tcaaccacca tcaaacagga ttttcgctg ctggggcaaa ccagcgtgga ccgcttgctg 5580
caactctctc agggccaggc ggtgaagggc aatcagctgt tgcccgtctc actggtgaaa 5640
agaaaaacca ccctggcgcc caatacgcaa accgcctctc cccgcgctt ggccgattca 5700
ttaatgcagc tggcacgaca ggtttcccga ctggaaagcg ggcagtgagc gcaacgcaat 5760
taatgtgagt tagcgcgaat tgatctg 5787

```

```

<210> SEQ ID NO 48
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 48

```

```

aggttcccat ggtgactcag gacccg 26

```

```

<210> SEQ ID NO 49
<211> LENGTH: 40
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        primer

```

```

<400> SEQUENCE: 49

```

```

gtaagcttag tggatgatgt gatgaccgaa cgcttcgtcc 40

```

```

<210> SEQ ID NO 50
<211> LENGTH: 4614
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
        plasmid pKK223:cterm-His

```

```

<400> SEQUENCE: 50

```

```

ttgacaatta atcatcggct cgtataatgt gtggaattgt gagcggataa caatttcaca 60
caggaaacag aacctggga cataacggat ctagatctca ccatcaccac cattagtcga 120
cctgcagcca agcttggtg ttttggcgga tgagagaaga ttttcagcct gatacagatt 180
aaatcagaac gcagaagcgg tctgataaaa cagaatttgc ctggcggcag tagcgcggtg 240

```

-continued

---

gtcccacctg accccatgcc gaactcagaa gtgaaacgcc gtagcgccga tggtagtggtg 300  
gggtctcccc atgcgagagt agggaaactgc caggcatcaa ataaaacgaa aggctcagtc 360  
gaaagactgg gcctttcgtt ttatctgttg tttgtcggtg aacgctctcc tgagtaggac 420  
aaatccgccg ggagcggatt tgaacgttgc gaagcaacgg cccggagggt ggcgggcagg 480  
acgcccgccca taaactgccca ggcatacaat taagcagaag gccatcctga cggatggcct 540  
ttttgcgttt ctacaaactc ttttgtttat ttttctaaat acattcaaat atgtatccgc 600  
tcatgagaca ataacctga taaatgcttc aataatattg aaaaaggaag agtatgagta 660  
ttcaacattt ccgtgtcgc cttattccct tttttgcggc attttgccct cctgtttttg 720  
ctcaccacaga aacgctgggtg aaagtaaaag atgtgaaga tcagttgggt gcacgagtg 780  
gttacatcga actggatctc aacagcggta agatccttga gagttttcgc cccgaagaac 840  
gttttccaat gatgagcact tttaaagttc tgctatgtgg cgcggtatta tcccgtgttg 900  
acgcccggca agagcaactc ggtcgcggca tacactattc tcagaatgac ttggttgagt 960  
actcaccagt cacagaaaag catcttacgg atggcatgac agtaagagaa ttatgcagtg 1020  
ctgccataac catgagtgat aacactgagg ccaacttact tctgacaacg atcggaggac 1080  
cgaaggagct aaccgctttt ttgcacaaca tgggggatca tgtaactcgc cttgatcgtt 1140  
gggaaccgga gctgaatgaa gccataccaa acgacgagcg tgacaccacg atgctgtagc 1200  
aatggcaaca acgttgcgca aactattaac tggcgaacta cttactctag cttcccggca 1260  
acaattaata gactggatgg aggcggataa agttgcagga ccacttctgc gctcggccct 1320  
tccggctggc tggtttattg ctgataaatc tggagccggg gagcgtgggt ctccgggtat 1380  
cattgcagca ctggggccag atggtaagcc ctcccgtatc gtagttatct acacgacggg 1440  
gagtcaggca actatggatg aacgaaatag acagatcgtc gagatagggt cctcactgat 1500  
taagcattgg taactgtcag accaagtta ctcatatata ctttagattg atttaaaact 1560  
tcatttttaa tttaaaagga tctaggtgaa gatccttttt gataatctca tgacaaaaat 1620  
cccttaacgt gagttttcgt tccactgagc gtcagacccc gtagaaaaga tcaaaggatc 1680  
ttcttgagat cctttttttc tgccgcgtaat ctgctgcttg caaacaaaaa aaccaccgct 1740  
accagcgggtg gtttgtttgc cggatcaaga gctaccaact ctttttccga aggtaactgg 1800  
cttcagcaga ggcagatac caaatactgt ccttctagtg tagccgtagt taggccacca 1860  
cttcaagaac tctgtagcac cgcctacata cctcgtctcg ctaatcctgt taccagtggc 1920  
tgctgccagt ggcgataagt cgtgtcttac cgggttgac tcaagacgat agttaccgga 1980  
taaggcgcag cggtcgggct gaacggggggg ttcgtgcaca cagcccagct tggagcgaac 2040  
gacctacacc gaactgagat acctacagcg tgagcattga gaaagcgcca cgcttcccga 2100  
aggagaaaag gcgacaggt atccggtaag cggcagggtc ggaacaggag agcgcacgag 2160  
ggagcttcca gggggaaacg cctggtatct ttatagtcct gtcgggtttc gccacctctg 2220  
acttgagcgt cgatttttgt gatgctcgtc agggggcgg agcctatgga aaaacgccag 2280  
caacgcggcc tttttacggt toctggcctt ttgctggcct tttgctcaca tgttctttcc 2340  
tgcgttatcc cctgattctg tggataaccg tattaccgcc tttgagtgag ctgataccgc 2400  
tcgccgcagc cgaacgaccg agcgcagcga gtcagtgagc gaggaagcgg aagagcgcct 2460  
gatgcggtat tttctcctta cgcactctgt cggtatttca caccgcatat ggtgcactct 2520

-continued

---

```

cagtacaatc tgctctgatg ccgcatagtt aagccagtat aactccgct atcgctacgt 2580
gactgggtca tggctgcgcc ccgacacccg ccaacacccg ctgacgcgcc ctgacgggct 2640
tgtctgctcc cggcatccgc ttacagacaa gctgtgaccg tctccgggag ctgcatgtgt 2700
cagaggtttt caccgtcatc accgaaacgc gcgaggcagc tgcggtaaag ctcacagcg 2760
tggtcgtgaa gcgattcaca gatgtctgcc tgttcatccg cgtccagctc gttgagtttc 2820
tccagaagcg ttaatgtctg gcttctgata aagcgggcca tgtaagggc ggttttttcc 2880
tgtttggtca ctgatgcctc cgtgtaaggg ggatttctgt tcatgggggt aatgataccg 2940
atgaaacgag agaggatgct cacgatacgg gttactgatg atgaacatgc ccggttactg 3000
gaacgttggt agggtaaaca actggcggta tggatgcggc gggaccagag aaaaatcact 3060
caggttcaat gccagcgctt cgtaataca gatgtaggtg ttccacaggg tagccagcag 3120
catcctgcga tgcagatccg gaacataatg gtgcagggcg ctgacttccg cgtttccaga 3180
ctttacgaaa cacggaaacc gaagaccatt catgttggtg ctcaggctgc agacgttttg 3240
cagcagcagt cgcttcacgt tcgctcgcgt atcggtgatt cattctgcta accagtaagg 3300
caaccccgcc agcctagccg ggtcctcaac gacaggagca cgatcatgcg caccctggc 3360
caggacccaa cgctgcccga gatgcgccgc gtgcggctgc tggagatggc ggacgcgatg 3420
gatatggtct gccaaagggt ggtttgccga ttcacagttc tccgcaagaa ttgattggct 3480
ccaattcttg gagtggtgaa tccgtagcg aggtgccgcc ggcttccatt caggtcgagg 3540
tgccccggt ccatgcaccg cgacgcaacg cggggaggca gacaaggat agggcggcgc 3600
ctacaatcca tgccaacccg ttccatgtgc tcgccaggc ggcataaatc gccgtgacga 3660
tcagcgggcc agtgatcgaa gttaggctgg taagagccgc gagcgatcct tgaagctgtc 3720
cctgatggtc gtcactacc tgccctggaca gcatggcctg caacgcgggc atcccgatgc 3780
cgccggaagc gagaagaatc ataatgggga aggccatcca gcctcgcgtc gcgaacgcca 3840
gcaagacgta gccagcgcg tcggccgcca tgccgggat aatggcctgc ttctcgccga 3900
aacgtttggt ggcgggacca gtgacgaagg cttgagcgag ggcgtgcaag attccgaata 3960
ccgcaagcga caggccgatc atcgtcgcgc tccagcgaaa gcggtcctcg ccgaaaatga 4020
cccagagcgc tgccggcacc tgcctacga gttgcatgat aaagaagaca gtcataagtg 4080
cggcgacgat agtcatgccc cgcgcccacc ggaaggagct gactgggttg aaggtctca 4140
agggcatcgg tcgacgctct cccttatgcg actcctgcat taggaagcag cccagtagta 4200
ggttgaggcc gttgagcacc gccgcgcaa ggaatggtgc atgcaaggag atggcgccca 4260
acagtcccc gccacgggg cctgccacca taccacgcc gaaacaagcg ctcacgagcc 4320
cgaagtggcg agccgatct tcccacggtg tgatgtcggc gatataggcg ccagcaaccg 4380
cacctgtggc gccggtgatg ccggccacga tgcgtccggc gtagaggatc cgggcttacc 4440
gactgcacgg tgcaccaatg cttctggcgt caggcagcca tcggaagctg tggatggct 4500
gtgcaggtcg taaatcactg cataattcgt gtcgctcaag gcgcactccc gttctggata 4560
atgttttttg cgccgacatc ataacggttc tggcaaatat tctgaaatga gctg 4614

```

&lt;210&gt; SEQ ID NO 51

&lt;211&gt; LENGTH: 26

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 51

aggttcccat ggtgactcag gacccg 26

<210> SEQ ID NO 52  
 <211> LENGTH: 40  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic primer

<400> SEQUENCE: 52

gtaagcttag tggatgatgt gatgaccgaa cgcttcgtcc 40

<210> SEQ ID NO 53  
 <211> LENGTH: 8241  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic plasmid pKK223:kgd-his  
 <220> FEATURE:  
 <221> NAME/KEY: modified\_base  
 <222> LOCATION: (1425)..(1425)  
 <223> OTHER INFORMATION: a, c, t, g, unknown or other

<400> SEQUENCE: 53

atggatcgta aatttcgtga tgatccgtct tccgttgacc caagctggca tgagtttctg 60  
 gtggactata gcccggagcc gacttcccaa ccgctgctg agccgactcg tgtgaccagc 120  
 cctttggtcg ctgagcgtgc ggcagcggcg gctccgcaag cgccgccgaa accggccgac 180  
 acggccgcag cgggcaacgg tgcctggcg gctctggccg ccaagaccgc ggtgcctccg 240  
 ccagccgaag gcgatgaagt tgctgtcctg cgtggtgctg cggtgctgt cgtgaagaac 300  
 atgagcgcgt cgctggaagt gcctacggcg acctcggctg gtgctgtgcc agcgaattg 360  
 ttgatcgata atcgatcgt gatcaataat cagttgaaac gcacgcgtgg tggtaaaatc 420  
 agctttaccc atttgcctgg ttatgctctg gttcaggcgg ttaagaagt cccgaatatg 480  
 aatcgccact acacggaggt ggacggtaag ccgacggcgg ttactcctgc gcacaccaat 540  
 ctgggtctgg ctattgacct gcaaggtaaa gatggcaagc gctctctggt ggtggccggt 600  
 atcaaacggt gcgagacgat gcgcttcgcg cagttcgtaa ccgctacga ggacatcgtc 660  
 cgtcgtgctc gtgatggtaa actgaccacc gaggacttgc ctggcggttac gatttcgctg 720  
 acgaatccgg gcacgattgg cacgggtgcac tctgtgccac gcctgatgcc aggtcaaggt 780  
 gctatcatcg gtgctggtgc gatggaatat ccggcggagt tccaaggcgc gtccgaggaa 840  
 cgcacgctg agctgggtat tggcaaactg attacctga cttctaccta cgaccaccgt 900  
 atcattcaag gtgcggaag cggtgacttc ctgctacca ttcataact gctgtgtct 960  
 gatggcttct gggatgaggt cttccgcgag ctgacatcc cgtacctgcc agtgcgttgg 1020  
 agcaccgata atccgactc gatcgttgac aagaacgcac gtgtgatgaa cctgatcgcg 1080  
 gcctatcgca accgcggcca tctgatggcg gacaccgacc cgttgcgctt ggataaggcg 1140  
 cgctttcgtt cgcacccgga tttggaggtg ttgacgcatg gcctgactct gtgggacctg 1200

-continued

---

gaccgcgtct	ttaaggttga	cggtttcgcg	ggtgctcaat	acaaaaagct	gcgtgacgtg	1260
ctgggtttgc	tgcgcgacgc	gtactgccgt	cacatcggtg	ttgagtacgc	gcacattctg	1320
gacccggaac	aaaaggagtg	gctggagcag	cgtgtcgaga	cgaagcacgt	caaaccgacc	1380
gttgctcaac	aaaagtacat	cttgctgaaa	ctgaacgcag	cggangcgtt	tgaaaccttc	1440
ctgcaaacca	agtatgtggg	ccaaaagcgt	tttagcctgg	agggtgcgga	atctgttatt	1500
ccaatgatgg	acgcggcgat	tgaccaatgt	gccgagcatg	gtttggatga	agtggtcatt	1560
ggcatgccgc	accgtggtcg	cttgaatgtc	ctggcgaaca	tcgtgggcaa	accgtattcg	1620
caaatcttta	cggagtttga	aggcaatctg	aaccctgcc	aggctcacgg	ttccggcgat	1680
gtcaagtacc	acctgggtgc	gacgggtctg	tatctgcaaa	tgtttggcga	taacgatatt	1740
caggtttccc	tgacggcaaa	tccgtcccat	ttggaagcgg	tcgatccagt	gctggaaggt	1800
ttggtgcgtg	cgaagcaaga	cctgctggac	cacggttcca	ttgactccga	cggccaacgt	1860
gcattctccg	tcgttccgtt	gatgctgcac	ggtgatgcgg	ccttcgctgg	tcaaggcgtt	1920
gttgcggaga	cgctgaatct	ggcaaatctg	cctggctatc	gtgtgggtgg	tacgattcac	1980
atcattgtga	acaaccaa	cggcttcacc	accgctccgg	aatactctcg	ttcgagcgag	2040
tactgcacgg	atgtggcgaa	gatgatcgg	gcgccaatct	tccacgtgaa	cggcgacgat	2100
ccagaagcgt	gtgtctgggt	tgcgcgtttg	gcagttgact	tccgccaacg	ctttaaaaag	2160
gacgttgta	ttgacatgct	gtgctatcgt	cgtcgtggtc	ataatgagg	tgacgatcca	2220
tctatgacca	atccgtatgt	ctatgacgtc	gtggacacca	agcgtggcgc	ccgtaaatcc	2280
tacaccgagg	ctctgatcgg	tcgtggcgac	atcagcatga	aagaggcgga	ggatgcgctg	2340
cgcgattacc	agggccaa	ggagcgtgtg	ttcaacgaag	tgcgcgaact	ggagaagcac	2400
ggcgtccaac	cgteggagag	cgtcgagtcg	gaccaa	tcccagcggg	cctggcgacc	2460
gctgttgaca	aatccctgct	ggcacgtatc	ggtgatgcct	tcctggcgct	gcctaacggt	2520
ttcacggcac	accgcgcgt	tcagccggtg	ctggaaaagc	gtcgtgagat	ggcgtatgag	2580
ggtaagattg	actgggcgtt	tggtgagttg	ctggcgtcgg	gcagcctggt	tgctgagggc	2640
aagctggtec	gtctgagcgg	tcaagactct	cgtcgtggca	cctttagcca	gcgtcattct	2700
gtgctgatcg	accgtcacac	cggcgaggag	ttcacccgc	tgcagctgct	ggcgactaac	2760
agcgacggca	gcccgaccg	tggttaagttc	ctggtttatg	attcgccgtt	gtcggaatac	2820
gcggctgtgg	gttttgagta	cggttatacc	gttggaatc	cagacgcggt	tgtgctgtgg	2880
gaggcgcaat	tcggcgactt	tgtaaatggt	gccagtcga	tcattgacga	gttcatcagc	2940
tcgggccaag	cgaaatgggg	tcagctgtcc	aacgttggtc	tgttgctgcc	acacggccac	3000
gagggtcagg	gcccggacca	cacgtcggcg	cgcattgagc	gctttttgca	gctgtgggct	3060
gagggtagca	tgacgatcgc	gatgccgagc	acccgtcca	attactttca	cttgctgcgc	3120
cgccacgcgt	tggacggcat	ccaacgtccg	ttgatcgtgt	ttaccccgaa	atccatgctg	3180
cgccacaagg	cggcggtttc	ggagattaag	gattttaccg	agattaaatt	ccgtagcgtc	3240
ctggaggaac	cgacctacga	ggatggatc	ggcgaccgca	ataaagttag	ccgtattctg	3300
ttgacctccg	gtaagttgta	ttatgaattg	gcggcacgca	aggcgaaaga	caaccgtaat	3360
gatctggcaa	ttgtgcgtct	ggagcaactg	gcgccgttgc	cgcgtcgtcg	tctgcgtgaa	3420
accctggatc	gttacgaaaa	tgtcaaagag	ttcttttggg	tccaggagga	accagcgaac	3480

-continued

---

cagggcgect	ggcccgcttt	cggtttggag	ttgccggagt	tgctgcegga	caagttggcg	3540
ggcatcaagc	gtatctcccg	tcgcgcatg	tctgcccga	gcagcggttc	ctcgaaggtt	3600
catgccgtgg	agcagcaaga	aatcttggac	gaagcgttcg	gatctagatc	tcaccatcac	3660
caccattagt	cgacctgcag	ccaagcttgg	ctgttttggc	ggatgagaga	agattttcag	3720
cctgatacag	attaatcag	aacgcagaag	cggtctgata	aaacagaatt	tgctggcgg	3780
cagtagcgcg	gtgggtcccac	ctgacccccat	gccgaactca	gaagtgaaac	gccgtagcgc	3840
cgatggtagt	gtgggtctc	cccatgagag	agtagggaac	tgccaggcat	caaataaaac	3900
gaaaggctca	gtcgaaagac	tgggcctttc	gttttatctg	ttgtttgtcg	gtgaacgctc	3960
tctgagtag	gacaaatccg	ccgggagcgg	atttgaacgt	tgcaagcaa	cggcccggag	4020
ggtggcgggc	aggacgcccg	ccataaactg	ccaggcatca	aattaagcag	aaggccatcc	4080
tgacggatgg	cctttttgcg	tttctacaaa	ctcttttgtt	tatttttcta	aatacattca	4140
aatatgtatc	cgctcatgag	acaataacc	tgataaatgc	ttcaataata	ttgaaaaagg	4200
aagagtatga	gtattcaaca	tttccgtgtc	gccttattc	ccttttttgc	ggcattttgc	4260
cttctgttt	ttgctcacc	agaaacgctg	gtgaaagtaa	aagatgctga	agatcagttg	4320
ggtgcacgag	tgggttacat	cgaactggat	ctcaacagcg	gtaagatcct	tgagagtttt	4380
cgccccgaag	aacgttttcc	aatgatgagc	acttttaaag	ttctgctatg	tggcgcggta	4440
ttatcccgtg	ttgacgccgg	gcaagagcaa	ctcggtcgcc	gcatacacta	ttctcagaat	4500
gacttggttg	agtactcacc	agtcacagaa	aagcatctta	cggatggcat	gacagtaaga	4560
gaattatgca	gtgctgccat	aaccatgagt	gataaacctg	cggccaactt	acttctgaca	4620
acgatcggag	gaccgaagga	gctaaccgct	tttttgcaca	acatggggga	tcatgtaact	4680
cgccttgatc	gttgggaacc	ggagctgaat	gaagccatac	caaacgacga	gcgtgacacc	4740
acgatgctgt	agcaatggca	acaacgcttc	gcaaacatt	aactggcgaa	ctacttactc	4800
tagcttcccg	gcaacaatta	atagactgga	tggaggcggg	taaagttgca	ggaccacttc	4860
tgcgctcggc	ccttccggct	ggctggttta	ttgctgataa	atctggagcc	ggtgagcgtg	4920
ggtctcggcg	tatcattgca	gcaactgggc	cagatggtaa	gcctcccgt	atcgtagtta	4980
tctacacgac	ggggagtcag	gcaactatgg	atgaacgaaa	tagacagatc	gctgagatag	5040
gtgcctcact	gattaagcat	tggtaactgt	cagaccaagt	ttactcatat	atactttaga	5100
ttgatttaaa	acttcatttt	taatttaaaa	ggatctaggt	gaagatcctt	tttgataatc	5160
tcatgaccaa	aatcccttaa	cgtgagtttt	cgttccactg	agcgtcagac	cccgtagaaa	5220
agatcaaagg	atcttcttga	gatccttttt	ttctgcgctg	aatctgctgc	ttgcaaacaa	5280
aaaaaccacc	gctaccagcg	gtggtttgtt	tgccggatca	agagctacca	actctttttc	5340
cgaaggtaac	tggcttcagc	agagcgcaga	taccaaatac	tgctcttcta	gtgtagccgt	5400
agttaggcca	ccacttcaag	aactctgtag	caccgcctac	atacctcgct	ctgctaatec	5460
tgttaccagt	ggctgctgcc	agtggcgata	agtcgtgtct	taccgggttg	gactcaagac	5520
gatagttacc	ggataaggcg	cagcggtcgg	gctgaacggg	gggttcgtgc	acacagccca	5580
gcttggagcg	aacgacctac	accgaactga	gatacctaca	gcgtgagcat	tgagaaagcg	5640
ccacgcttcc	cgaagggaga	aaggcggaca	ggatccgggt	aagcggcagg	gtcggaacag	5700
gagagcgcac	gagggagctt	ccagggggaa	acgcctggta	tctttatagt	cctgtcgggt	5760

-continued

---

ttcgccacct	ctgacttgag	cgctgatttt	tgtgatgctc	gtcagggggg	cggagcctat	5820
ggaaaaacgc	cagcaacgcg	gcctttttac	ggttcctggc	cttttgctgg	ccttttgctc	5880
acatgttctt	tectgcgtta	tcccctgatt	ctgtggataa	ccgtattacc	gcctttgagt	5940
gagctgatac	cgctcgccgc	agccgaacga	ccgagcgcag	cgagtcagtg	agcgaggaag	6000
cggaagagcg	cctgatgcgg	tattttctcc	ttacgcatct	gtgcggtatt	tcacaccgca	6060
tatggtgcac	tctcagtaca	atctgctctg	atgccgcata	gttaagccag	tatacactcc	6120
gctatcgcta	cgtgactggg	tcatggctgc	gccccgacac	ccgccaacac	ccgctgacgc	6180
gccctgacgg	gcttgtctgc	tcccggcctc	cgcttacaga	caagctgtga	ccgtctccgg	6240
gagctgcatg	tgtcagaggt	tttcaccgtc	atcaccgaaa	cgcgcgaggc	agctgcggta	6300
aagctcatca	gcgtggctgt	gaagcgattc	acagatgtct	gcctgttcat	ccgctccag	6360
ctcgttgagt	ttctccagaa	gcgttaatgt	ctggcttctg	ataaagcggg	ccatgttaag	6420
ggcggttttt	tectgtttgg	tcaactgatgc	ctccgtgtaa	gggggatttc	tgttcatggg	6480
ggtaatgata	ccgatgaaac	gagagaggat	gctcacgata	cgggttactg	atgatgaaca	6540
tgcccggtta	ctggaacggt	gtgagggtaa	acaactggcg	gtatggatgc	ggcgggacca	6600
gagaaaaatc	actcagggtc	aatgccagcg	cttcgttaat	acagatgtag	gtgttccaca	6660
gggtagccag	cagcatcctg	cgatgcagat	ccggaacata	atgggtgcagg	gcgctgactt	6720
ccgctgttcc	agactttacg	aaacacggaa	accgaagacc	attcatgttg	ttgctcaggt	6780
cgcagacggt	ttgcagcagc	agtcgcttca	cgctcgctcg	cgatcggtg	attcattctg	6840
ctaaccagta	aggcaacccc	gccagcctag	ccgggtcctc	aacgacagga	gcacgatcat	6900
gcgcacccgt	ggccaggacc	caacgctgcc	cgagatgcgc	cgctgcggc	tgctggagat	6960
ggcggacgcg	atggatatgt	tctgccaaag	gttggtttgc	gcattcacag	ttctccgcaa	7020
gaattgattg	gctccaattc	ttggagtggg	gaatccgtta	gcgaggtgcc	gccggcttcc	7080
attcaggtcg	aggtggcccg	gctccatgca	ccgcgacgca	acgcggggag	gcagacaagg	7140
tatagggcgg	cgctacaat	ccatgccaac	ccgttccatg	tgctcgccga	ggcggcataa	7200
atcgccgtga	cgatcagcgg	tccagtgatc	gaagttaggc	tggttaagagc	cgcgagcgat	7260
ccttgaagct	gtccctgatg	gtcgtcatct	acctgcctgg	acagcatggc	ctgcaacgcg	7320
ggcatcccga	tgccgccgga	agcgagaaga	atcataatgg	ggaaggccat	ccagcctcgc	7380
gtcgcgaacg	ccagcaagac	gtagcccagc	gcgtcgcccg	ccatgccggc	gataatggcc	7440
tgcttctcgc	cgaaacgttt	ggtaggggga	ccagtgcgca	aggcttgagc	gagggcgtgc	7500
aagattccga	ataccgcaag	cgacaggccg	atcatcgctg	cgctccagcg	aaagcgggtcc	7560
tcgccgaaaa	tgaccagag	cgctgccggc	acctgtccta	cgagttgcat	gataaagaag	7620
acagtcataa	gtgcggcgac	gatagtcatg	ccccgcgccc	accggaagga	gctgactggg	7680
ttgaaggctc	tcaagggcat	cggtcgacgc	tctcccttat	gcgactcctg	cattaggaag	7740
cagcccagta	gtaggttgag	gccgttgagc	accgcccggc	caaggaatgg	tgcatgcaag	7800
gagatggcgc	ccaacagtcc	cccggccacg	gggctgcca	ccataccac	gccgaaacaa	7860
gcgctcatga	gcccgaagtg	gcgagcccga	tcttccccat	cggtgatgct	ggcgatatag	7920
gcgccagcaa	ccgcacctgt	ggcgccgggtg	atgccggcca	cgatgcgtcc	ggcgtagagg	7980
atccgggctt	atcgactgca	cggtgcacca	atgcttctgg	cgtcaggcag	ccatcggaag	8040



-continued

---

```

ctgtggtatg gctgtgcagg tcgtaaataca ctgcataatt cgtgtegetc aaggcgact 8100
cccgttctgg ataatgtttt ttgcgccgac atcataacgg ttctggcaaa tattctgaaa 8160
tgagctggtg acaattaatc atcggtcgtg ataatgtgtg gaattgtgag cggataacaa 8220
tttcacacag gaaacagaac c 8241

```

```

<210> SEQ ID NO 54
<211> LENGTH: 541
<212> TYPE: PRT
<213> ORGANISM: Leuconostoc mesenteroides

```

```

<400> SEQUENCE: 54

```

```

Met Thr Asp Thr Leu Thr Phe Asn Thr Lys His Leu Leu Glu Ala Leu
1          5          10          15
Phe Glu Ser Gly Ile Arg His Phe Ile Val Ser Pro Gly Ser Arg Ser
20          25          30
Thr Pro Ile Ala Leu Leu Leu Ala Glu Tyr Ala Glu Gln Thr Asn Glu
35          40          45
Ile Lys Leu Phe Val Asp Val Asp Glu Arg Ser Ala Gly Phe Phe Ala
50          55          60
Leu Gly Ile Ala Lys Thr Leu Leu Glu Pro Val Val Leu Leu Gly Thr
65          70          75          80
Ser Gly Thr Ala Ile Ala Glu Tyr Met Pro Ala Val Ala Glu Ala Tyr
85          90          95
Ala Ala Asn Ile Pro Leu Val Val Leu Ser Thr Asp Arg Pro Gln Glu
100         105         110
Leu Gln Phe Asn Gly Ala Pro Gln Thr Ile Pro Gln Ser Asn Leu Phe
115        120        125
Gly Gln Leu Thr Lys Gln Ala Val Leu Ile Arg Leu Gln Asp Met His
130        135        140
Ser Asp Val Thr Glu Tyr Ile Asp Phe Ile Val Gln Lys Val Val Asn
145        150        155        160
Leu Ser Ile Thr Ala Pro Arg Gly Pro Ile Gln Ile Asn Leu Pro Leu
165        170        175
Arg Lys Pro Leu Met Pro Val Leu Asp Arg Gln Asp Glu Val His Val
180        185        190
Gln Arg Val Val Phe Asp Lys Gln Ser Val Gln Tyr Arg Leu Pro Pro
195        200        205
Ile Thr Ala Lys Arg Leu Leu Ile Leu Ala Gly Pro Asn Val Leu Asn
210        215        220
Ser Tyr Asp Asp Ser Leu Lys Lys Phe Ala Ile Lys Asn Asn Val Pro
225        230        235        240
Val Ile Ala Asp Val Leu Ser Gln Ser Leu His Thr Tyr Thr Ile His
245        250        255
Gly Ile Asp Val Leu Leu Gln Ala His Lys Ile Asn Ala Asp Leu Lys
260        265        270
Pro Asp Leu Val Val Arg Phe Gly Lys Thr Pro Val Ser Ala Arg Val
275        280        285
Leu Gln Trp Leu Lys Glu Glu Asn Ile Leu Thr Trp His Val Gly Glu
290        295        300
Asp Ala Gly Val Asp His Thr Arg His Ile Val Arg Ala Ile Lys Met
305        310        315        320

```

-continued

---

Ala Pro His Asp Phe Leu Glu Ser Met His Leu Thr Leu Ser Lys Asn  
 325 330 335

Gln Ile Asp Phe Asn Gln Lys Trp Leu Ser Leu Pro Lys Val Ile Lys  
 340 345 350

Thr Arg Asn Glu Met Asn Ile Ile Thr Ala Leu Asp Asp Ala Val Pro  
 355 360 365

Asp Asp Thr His Ile Phe Val Ala Asn Ser Met Pro Ile Arg Asp Met  
 370 375 380

Asp Asn Phe Phe Thr Gly Asn His Thr Gln Arg Ile Tyr Ala Asn Arg  
 385 390 395 400

Gly Ala Asn Gly Ile Asp Gly Val Ile Ser Ser Ala Leu Gly Met Ser  
 405 410 415

Ala Val Ala Lys Gln Arg Ser Val Leu Leu Thr Gly Asp Leu Thr Leu  
 420 425 430

Phe His Asp Met Asn Gly Leu Met Met Ala Lys Asn Tyr Gln Leu Pro  
 435 440 445

Leu Asp Ile Ile Val Ile Asn Asn Asn Gly Gly Gly Ile Phe Ser Phe  
 450 455 460

Leu Pro Gln Ala Gly Ala Pro Lys Tyr Phe Glu Gln Gln Phe Gly Thr  
 465 470 475 480

Pro Leu Asn Ile Asp Ile Lys Lys Ile Ala Asp Leu Tyr Tyr Ile Asp  
 485 490 495

Tyr His Gln Leu Asn Val Pro Glu Ala Leu Ser Gln Ile Leu Gln Thr  
 500 505 510

Pro Ser Lys Thr Thr Arg Leu Ile Glu Tyr Lys Ser Asp His Gln Arg  
 515 520 525

Asn Arg Asp Asp His Arg Glu Val Leu Glu Met Leu Lys  
 530 535 540

<210> SEQ ID NO 55  
 <211> LENGTH: 541  
 <212> TYPE: PRT  
 <213> ORGANISM: Leuconostoc mesenteroides

<400> SEQUENCE: 55

Met Thr Asp Thr Leu Thr Phe Asn Thr Lys His Leu Leu Glu Ala Leu  
 1 5 10 15

Phe Glu Ser Gly Ile Arg His Phe Ile Val Ser Pro Gly Ser Arg Ser  
 20 25 30

Thr Pro Ile Ala Leu Leu Leu Ala Glu Tyr Ala Glu Gln Asn Asn Glu  
 35 40 45

Ile Lys Leu Phe Val Asp Val Asp Glu Arg Ser Ala Gly Phe Phe Ala  
 50 55 60

Leu Gly Ile Ala Lys Thr Leu Leu Glu Pro Val Val Leu Leu Gly Thr  
 65 70 75 80

Ser Gly Thr Ala Ile Ala Glu Tyr Met Pro Ala Val Ala Glu Ala Tyr  
 85 90 95

Ala Ala Asn Ile Pro Leu Val Val Leu Ser Thr Asp Arg Pro Gln Glu  
 100 105 110

Leu Gln Phe Asn Gly Ala Pro Gln Thr Ile Pro Gln Ser Asn Leu Phe  
 115 120 125

Gly Gln Leu Thr Lys Gln Ala Val Leu Ile Arg Leu Gln Asp Met His  
 130 135 140

-continued

---

Ser	Asp	Val	Thr	Glu	Tyr	Ile	Asp	Phe	Ile	Val	Gln	Lys	Val	Val	Asn	145	150	155	160
Leu	Ser	Ile	Thr	Ala	Pro	Arg	Gly	Pro	Ile	Gln	Ile	Asn	Leu	Pro	Leu	165	170	175	
Arg	Lys	Pro	Leu	Met	Pro	Val	Leu	Asp	Arg	Gln	Asp	Glu	Val	His	Val	180	185	190	
Gln	Arg	Val	Val	Phe	Asp	Lys	Gln	Ser	Val	Gln	Tyr	Arg	Leu	Pro	Pro	195	200	205	
Ile	Thr	Ala	Lys	Arg	Leu	Leu	Ile	Leu	Ala	Gly	Pro	Asn	Val	Leu	Asn	210	215	220	
Ser	Tyr	Asp	Asp	Ser	Leu	Lys	Lys	Phe	Ala	Ile	Lys	Asn	Asn	Val	Pro	225	230	235	240
Val	Ile	Ala	Asp	Val	Leu	Ser	Gln	Ser	Arg	His	Thr	Tyr	Thr	Ile	His	245	250	255	
Gly	Ile	Asp	Val	Leu	Leu	Gln	Ala	His	Lys	Ile	Asn	Ala	Asp	Leu	Lys	260	265	270	
Pro	Asp	Leu	Val	Val	Arg	Phe	Gly	Lys	Thr	Pro	Val	Ser	Ala	Arg	Val	275	280	285	
Leu	Gln	Trp	Leu	Lys	Glu	Glu	Asn	Ile	Leu	Thr	Trp	His	Val	Asp	Glu	290	295	300	
Asp	Ala	Gly	Val	Asp	His	Thr	Arg	His	Ile	Val	Arg	Ala	Ile	Lys	Met	305	310	315	320
Ala	Pro	His	Asp	Phe	Leu	Glu	Ser	Met	His	Leu	Thr	Leu	Ser	Lys	Asn	325	330	335	
Gln	Ile	Asp	Phe	Asn	Gln	Lys	Trp	Leu	Ser	Leu	Pro	Lys	Val	Ile	Lys	340	345	350	
Thr	Arg	Asn	Glu	Met	Asn	Ile	Ile	Thr	Ala	Leu	Asp	Asp	Ala	Val	Pro	355	360	365	
Asp	Asp	Thr	His	Ile	Phe	Val	Ala	Asn	Ser	Met	Pro	Ile	Arg	Asp	Met	370	375	380	
Asp	Asn	Phe	Phe	Thr	Gly	Asn	His	Thr	Gln	Arg	Ile	Tyr	Ala	Asn	Arg	385	390	395	400
Gly	Ala	Asn	Gly	Ile	Asp	Gly	Val	Ile	Ser	Ser	Ala	Leu	Gly	Met	Ser	405	410	415	
Ala	Val	Val	Lys	Gln	Arg	Ser	Val	Leu	Leu	Thr	Gly	Asp	Leu	Thr	Leu	420	425	430	
Phe	His	Asp	Met	Asn	Gly	Leu	Met	Met	Ala	Lys	Asn	Tyr	Gln	Leu	Pro	435	440	445	
Leu	Asp	Ile	Ile	Val	Ile	Asn	Asn	Asn	Gly	Gly	Gly	Ile	Phe	Ser	Phe	450	455	460	
Leu	Pro	Gln	Ala	Gly	Ala	Pro	Lys	Tyr	Phe	Glu	Gln	Leu	Phe	Gly	Asn	465	470	475	480
Pro	Leu	Asn	Ile	Asp	Ile	Lys	Lys	Ile	Ala	Asp	Leu	Tyr	Tyr	Ile	Asp	485	490	495	
Tyr	His	Gln	Leu	Asn	Val	Pro	Glu	Ala	Leu	Ser	Gln	Ile	Leu	Gln	Thr	500	505	510	
Pro	Ser	Lys	Thr	Thr	Arg	Leu	Ile	Glu	Tyr	Lys	Ser	Asp	His	Gln	Arg	515	520	525	
Asn	Arg	Asp	Asp	His	Arg	Glu	Val	Leu	Glu	Met	Leu	Lys	530	535	540				

-continued

---

```

<210> SEQ ID NO 56
<211> LENGTH: 541
<212> TYPE: PRT
<213> ORGANISM: Leuconostoc mesenteroides

<400> SEQUENCE: 56

Met Thr Asp Thr Leu Thr Phe Asn Thr Lys His Leu Leu Glu Ala Leu
1          5          10          15
Phe Glu Ser Gly Ile Arg His Phe Ile Val Ser Pro Gly Ser Arg Ser
20          25          30
Thr Pro Ile Ala Leu Leu Leu Ala Glu Tyr Ala Glu Gln Asn Asn Glu
35          40          45
Ile Lys Leu Phe Val Asp Val Asp Glu Arg Ser Ala Gly Phe Phe Ala
50          55          60
Leu Gly Ile Ala Lys Thr Leu Leu Glu Pro Val Val Leu Leu Gly Thr
65          70          75          80
Ser Gly Thr Ala Ile Ala Glu Tyr Met Pro Ala Val Ala Glu Ala Tyr
85          90          95
Ala Ala Asn Ile Pro Leu Val Val Leu Ser Thr Asp Arg Pro Gln Glu
100         105         110
Leu Gln Phe Asn Gly Ala Pro Gln Thr Ile Pro Gln Ser Asn Leu Phe
115        120        125
Gly Gln Leu Thr Lys Gln Ala Val Leu Ile Arg Leu Gln Asp Met His
130        135        140
Ser Asp Val Thr Glu Tyr Ile Asp Phe Ile Val Gln Lys Val Val Asn
145        150        155        160
Leu Ser Ile Thr Ala Pro Arg Gly Pro Ile Gln Ile Asn Leu Pro Leu
165        170        175
Arg Lys Pro Leu Met Pro Val Leu Asp Arg Gln Asp Glu Val His Val
180        185        190
Gln Arg Val Val Phe Asp Lys Gln Ser Val Gln Tyr Arg Leu Pro Pro
195        200        205
Ile Thr Ala Lys Arg Leu Leu Ile Leu Ala Gly Pro Asn Val Leu Asn
210        215        220
Ser Tyr Asp Asp Ser Leu Lys Lys Phe Ala Ile Lys Asn Asn Val Pro
225        230        235        240
Val Ile Ala Asp Val Leu Ser Gln Ser Arg His Thr Tyr Thr Ile His
245        250        255
Gly Ile Asp Val Leu Leu Gln Ala His Lys Ile Asn Ala Asp Leu Lys
260        265        270
Pro Asp Leu Val Val Arg Phe Gly Lys Thr Pro Val Ser Ala Arg Val
275        280        285
Leu Gln Trp Leu Lys Glu Glu Asn Ile Leu Thr Trp His Val Asp Glu
290        295        300
Asp Ala Gly Val Asp His Thr Arg His Ile Val Arg Ala Ile Lys Met
305        310        315        320
Ala Pro His Asp Phe Leu Glu Ser Met His Leu Thr Leu Ser Lys Asn
325        330        335
Gln Ile Asp Phe Asn Gln Lys Trp Leu Ser Leu Pro Lys Val Ile Lys
340        345        350
Thr Arg Asn Glu Met Asn Ile Ile Thr Ala Leu Asp Asp Ala Val Pro
355        360        365

```

-continued

---

Asp Asp Thr His Ile Phe Val Ala Asn Ser Met Pro Ile Arg Asp Met  
 370 375 380

Asp Asn Phe Phe Thr Gly Asn His Thr Gln Cys Ile Tyr Ala Asn Arg  
 385 390 395 400

Gly Ala Asn Gly Ile Asp Gly Val Ile Ser Ser Ala Leu Gly Met Ser  
 405 410 415

Ala Val Val Lys Gln Arg Ser Val Leu Leu Thr Gly Asp Leu Thr Leu  
 420 425 430

Phe His Gly Met Asn Gly Leu Met Met Ala Lys Asn Tyr Gln Leu Pro  
 435 440 445

Leu Asp Ile Ile Val Ile Asn Asn Asn Gly Gly Gly Ile Phe Ser Phe  
 450 455 460

Leu Pro Gln Ala Gly Ala Pro Lys Tyr Phe Glu Gln Leu Phe Gly Thr  
 465 470 475 480

Pro Leu Asn Ile Asp Ile Lys Lys Ile Ala Asp Leu Tyr Tyr Ile Asp  
 485 490 495

Tyr His Gln Leu Asn Val Pro Glu Ala Leu Ser Gln Ile Leu Gln Ala  
 500 505 510

Pro Ser Lys Thr Thr Arg Leu Ile Glu Tyr Lys Ser Asp His Gln Arg  
 515 520 525

Asn Arg Asp Asp His Arg Glu Val Leu Glu Met Leu Lys  
 530 535 540

<210> SEQ ID NO 57  
 <211> LENGTH: 219  
 <212> TYPE: PRT  
 <213> ORGANISM: Escherichia coli

<400> SEQUENCE: 57

Met Lys Glu Ile Ile Asp Gly Phe Leu Lys Phe Gln Arg Glu Ala Phe  
 1 5 10 15

Pro Lys Arg Glu Ala Leu Phe Lys Gln Leu Ala Thr Gln Gln Ser Pro  
 20 25 30

Arg Thr Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu  
 35 40 45

Val Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly  
 50 55 60

Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Ser  
 65 70 75 80

Val Glu Tyr Ala Val Ala Ala Leu Arg Val Ser Asp Ile Val Ile Cys  
 85 90 95

Gly His Ser Asn Cys Gly Ala Met Thr Ala Ile Ala Ser Cys Gln Cys  
 100 105 110

Met Asp His Met Pro Ala Val Ser His Trp Leu Arg Tyr Ala Asp Ser  
 115 120 125

Ala Arg Val Val Asn Glu Ala Arg Pro His Ser Asp Leu Pro Ser Lys  
 130 135 140

Ala Ala Ala Met Val Arg Glu Asn Val Ile Ala Gln Leu Ala Asn Leu  
 145 150 155 160

Gln Thr His Pro Ser Val Arg Leu Ala Leu Glu Glu Gly Arg Ile Ala  
 165 170 175

Leu His Gly Trp Val Tyr Asp Ile Glu Ser Gly Ser Ile Ala Ala Phe

-continued

---

180	185	190
Asp Gly Ala Thr Arg Gln Phe Val Pro Leu Ala Ala Asn Pro Arg Val		
195	200	205
Cys Ala Ile Pro Leu Arg Gln Pro Thr Ala Ala		
210	215	

<210> SEQ ID NO 58  
 <211> LENGTH: 219  
 <212> TYPE: PRT  
 <213> ORGANISM: *Psuedomonas fluorescens*  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Carbonic anhydrase P. fluorescens Pf-5

<400> SEQUENCE: 58

Met Gln Asn Ile Ile Asp Gly Phe Leu Lys Phe Gln Arg Glu Ala Phe		
1	5	10 15
Pro Gln Arg Ser Glu Leu Phe Lys Gln Leu Ala Ser Thr Gln Asn Pro		
20	25	30
Gly Thr Leu Phe Val Thr Cys Ser Asp Ser Arg Val Val Pro Glu Leu		
35	40	45
Leu Thr Gln Gln Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly		
50	55	60
Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Thr		
65	70	75 80
Val Glu Tyr Ala Val Ala Val Leu Gly Val Ser Asp Ile Val Ile Cys		
85	90	95
Gly His Ser Asp Cys Gly Ala Met Thr Ala Ile Ser Thr Cys Lys Cys		
100	105	110
Leu Asp His Leu Pro Ala Val Ala Asn Trp Leu Arg His Ala Glu Ser		
115	120	125
Ala Lys Val Ile Asn Ala Ala Arg Gln His Ala Ser Pro Ala Glu His		
130	135	140
Leu Asp Ala Leu Val Arg Asp Asn Val Ile Ala Gln Leu Ala Asn Leu		
145	150	155 160
Lys Thr His Pro Ser Val Ala Leu Ala Leu Glu Gln Gly Arg Leu Asn		
165	170	175
Leu His Gly Trp Val Tyr Asp Ile Glu Ser Gly Ala Ile Val Ala Leu		
180	185	190
Asp Gly Asn Thr Gln Arg Phe Val Ser Leu Ala Glu Tyr Pro His Thr		
195	200	205
Cys Ala Leu Ala Ser Gln Ala Ser Ser Ala Ala		
210	215	

<210> SEQ ID NO 59  
 <211> LENGTH: 213  
 <212> TYPE: PRT  
 <213> ORGANISM: *Yersinia ruckeri*  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Carbonic anhydrase Y. ruckeri ATCC 29473

<400> SEQUENCE: 59

Met Gln Asp Ile Ile Asp Gly Phe Leu Lys Phe Gln Arg Glu Val Phe		
1	5	10 15
Pro Gln Arg Ser Glu Leu Phe Lys Arg Leu Ala Asp Thr Gln His Pro		
20	25	30

-continued

---

Gly Ala Leu Phe Val Thr Cys Ser Asp Ser Arg Val Val Pro Glu Leu  
35 40 45

Leu Thr Gln Arg Glu Pro Gly Glu Leu Phe Val Ile Arg Asn Ala Gly  
50 55 60

Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Thr  
65 70 75 80

Val Glu Tyr Ala Val Ala Val Leu Gly Val Thr Asp Val Val Ile Cys  
85 90 95

Gly His Ser Asn Cys Gly Ala Met Ser Ala Ile Ala Glu Cys Gln Cys  
100 105 110

Leu Asp His Leu Pro Ala Val Ala Ala Trp Leu Arg His Ala Asp Ser  
115 120 125

Ala Lys Leu Val Asn Ala Ala Leu Pro His Ala Ser Pro Lys Asp Arg  
130 135 140

Leu Asn Ser Leu Val Arg Glu Asn Val Ile Ala Gln Leu Ala Asn Ile  
145 150 155 160

Lys Thr His Pro Ser Val Ala Leu Ala Cys Ala Gln Gly Arg Leu Arg  
165 170 175

Leu His Gly Trp Val Tyr Asp Ile Glu Thr Gly Ser Ile Asp Val Leu  
180 185 190

Asp Glu Leu Thr Arg Thr Phe Ser Pro Leu Ser Ala Tyr Ser Val Val  
195 200 205

Ser Lys Pro Thr Glu  
210

&lt;210&gt; SEQ ID NO 60

&lt;211&gt; LENGTH: 219

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Hahella chejuensis*

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Carbonic anhydrase *Hahella chejuensis* KCTC 2396

&lt;400&gt; SEQUENCE: 60

Met Lys Asp Ile Ile Glu Gly Phe Leu Lys Phe Gln Arg Glu Ala Phe  
1 5 10 15

Pro Glu Arg Lys Glu Leu Phe Lys Asp Leu Ala Asn Gln Gln Gln Pro  
20 25 30

Arg Thr Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu  
35 40 45

Val Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly  
50 55 60

Asn Ile Val Pro Pro Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Ser  
65 70 75 80

Val Glu Tyr Ala Val Ala Ala Leu Arg Val Ala Asp Ile Val Val Cys  
85 90 95

Gly His Ser Asn Cys Gly Ala Met Thr Ala Val Ala Thr Cys Gln Cys  
100 105 110

Ile Asp His Met Pro Ala Val Ala His Trp Leu Arg Tyr Ala Asp Ser  
115 120 125

Ala Lys Val Val Asn Gln Ala Arg Lys His Ala Ser Glu Arg Ala Lys  
130 135 140

Ile Glu Asp Met Val Arg Glu Asn Val Ile Ala Gln Leu Ala Asn Leu  
145 150 155 160

-continued

---

Gln Thr His Pro Ser Val Arg Leu Ala Leu Gln Glu Gly Arg Leu Thr  
                           165                          170                          175

Met His Gly Trp Phe Tyr Asp Ile Glu Ser Gly Gly Ile Asp Ala Tyr  
                           180                          185                          190

Asp Gly Ser Arg His Ala Phe Val Pro Leu Ala Glu His Pro Glu Ala  
                           195                          200                          205

Arg Ala Ile Pro Gly Lys Leu Ser His Ala Val  
                           210                          215

<210> SEQ ID NO 61  
 <211> LENGTH: 219  
 <212> TYPE: PRT  
 <213> ORGANISM: Burkholderia cenocepacia  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Carbonic anhydrase Burkholderia cenocepacia  
 J23152

<400> SEQUENCE: 61

Met Lys Asp Ile Ile Glu Gly Phe Leu Lys Phe Gln Arg Asp Ala Tyr  
 1                          5                          10                          15

Pro Ala Arg Ala Ala Leu Phe Arg Asp Leu Ala Arg Ser Gln Asn Pro  
                           20                          25                          30

Arg Ala Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu  
                           35                          40                          45

Val Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly  
                           50                          55                          60

Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Ser  
 65                          70                          75                          80

Val Glu Tyr Ala Val Ala Ala Leu Arg Val Thr Asp Val Val Ile Cys  
                           85                          90                          95

Gly His Ser Asp Cys Gly Ala Met Thr Ala Ile Ala Thr Cys Gln Cys  
                           100                          105                          110

Met Asp His Met Pro Ala Val Gly His Trp Leu Arg Tyr Ala Asp Ser  
                           115                          120                          125

Ala Arg Val Val Asn Glu Ala Arg Thr His Arg Ser Glu Arg Glu Arg  
                           130                          135                          140

Ile Asp Ser Met Val Arg Glu Asn Val Val Ala Gln Leu Ala Asn Leu  
 145                          150                          155                          160

Lys Thr His Pro Ala Val Arg Leu Ala Leu Glu Glu Gly Arg Leu Ala  
                           165                          170                          175

Leu His Gly Trp Val Tyr Asp Ile Glu Ser Gly Cys Ile Asp Ala Tyr  
                           180                          185                          190

Asp Gly Ala Thr Gly Arg Phe Val Ser Leu Ala Asp His Pro Gly Val  
                           195                          200                          205

Arg Ala Thr Pro Ala Thr Leu Pro Val Ala Ala  
                           210                          215

<210> SEQ ID NO 62  
 <211> LENGTH: 215  
 <212> TYPE: PRT  
 <213> ORGANISM: Yersinia frederiksenii  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Carbonic anhydrase Yersinia frederiksenii ATCC  
 33461

<400> SEQUENCE: 62



-continued

---

```

Met Lys Glu Ile Ile Asp Gly Phe Leu Lys Phe Gln Arg Asp Ala Phe
1          5          10          15

Pro Glu Arg Ala Glu Leu Phe Arg Ser Leu Ala Thr Gln Gln Ser Pro
          20          25          30

Lys Thr Leu Phe Ile Ser Cys Ser Asp Ser Arg Met Val Pro Glu Leu
          35          40          45

Val Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly
          50          55          60

Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Ile Ser Ala Ser
65          70          75          80

Val Glu Tyr Ala Val Thr Ala Leu Lys Val Thr Asp Ile Val Ile Cys
          85          90          95

Gly His Ser Asp Cys Gly Ala Met Thr Ala Ile Ala Lys Cys His Cys
          100          105          110

Leu Asp His Met Pro Ala Val Lys His Trp Leu Gln Tyr Ala Asp Ser
          115          120          125

Ala Lys Val Val Asn Glu Ser Arg Glu Tyr Lys Asn Ile His Asp Lys
          130          135          140

Thr Ile Ser Met Val His Glu Asn Val Val Ala Gln Leu Ala Asn Ile
145          150          155          160

Gln Thr His Pro Ser Val Arg Leu Ala Leu Glu Glu Gly Arg Leu Thr
          165          170          175

Ile His Gly Trp Val Tyr Asp Ile Glu Ser Gly Leu Ile Ser Ala Phe
          180          185          190

Asp Arg Ala Ser Arg Gln Phe Val Ser Leu Ala Ala Asn Pro Asn Val
          195          200          205

Arg Ala Val Pro Ala His Asn
          210          215

```

&lt;210&gt; SEQ ID NO 63

&lt;211&gt; LENGTH: 219

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Pseudomonas entomophila

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Carbonic anhydrase Pseudomonas entomophila L48

&lt;400&gt; SEQUENCE: 63

```

Met Gln Asp Ile Ile Asp Gly Phe Leu Lys Phe Gln Arg Asp Ala Phe
1          5          10          15

Pro Glu Arg Val Lys Leu Phe Lys Asp Leu Ala Thr Gln Gln Ser Pro
          20          25          30

Arg Ala Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu
          35          40          45

Val Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly
          50          55          60

Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Ser
65          70          75          80

Val Glu Tyr Ala Val Ala Ala Leu Gln Val Ala Asp Ile Val Ile Cys
          85          90          95

Gly His Ser Asp Cys Gly Ala Met Thr Ala Ile Ala Thr Cys Lys Cys
          100          105          110

Leu Asp His Met Pro Ala Val Ala Gly Trp Leu Arg Tyr Ala Asp Ser
          115          120          125

```

-continued

---

Ala Arg Val Val Asn Glu Ala Arg Gln His Gln Ser Pro His Ala Lys  
 130 135 140

Val Glu Ala Met Val Arg Glu Asn Val Ile Ala Gln Leu Ala Asn Ile  
 145 150 155 160

Gln Thr His Pro Ser Val Arg Leu Ala Leu Glu Glu Gly Arg Val Ala  
 165 170 175

Leu His Gly Trp Ile Tyr Asp Ile Glu Ser Gly Arg Ile Asp Ala Phe  
 180 185 190

Asp Gly Arg Thr Gly Gln Phe Val Ser Leu Ala Asp Asn Pro Glu Val  
 195 200 205

Arg Ala Val Ser His Ala Ser Arg His Val Ala  
 210 215

<210> SEQ ID NO 64  
 <211> LENGTH: 219  
 <212> TYPE: PRT  
 <213> ORGANISM: Pseudomonas putida  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Carbonate dehydratase Pseudomonas putida W619  
 <400> SEQUENCE: 64

Met Lys Ala Ile Ile Asp Gly Phe Leu Lys Phe Gln Lys Asn Ala Phe  
 1 5 10 15

Pro Glu Arg Val Lys Leu Phe Lys Asp Leu Ala Asn Gln Gln Ala Pro  
 20 25 30

Lys Ala Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu  
 35 40 45

Val Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly  
 50 55 60

Asn Ile Val Pro Ser Tyr Gly Pro Glu Pro Gly Gly Val Ser Ala Ser  
 65 70 75 80

Val Glu Tyr Ala Val Ala Gly Leu Asn Val Ala Asp Ile Val Ile Cys  
 85 90 95

Gly His Ser Asp Cys Gly Ala Met Thr Ala Ile Ala Thr Cys Lys Cys  
 100 105 110

Leu Asp His Met Pro Ala Val Ala Gly Trp Leu Arg Tyr Ala Asp Ser  
 115 120 125

Ala Lys Val Val Asn Glu Ala Arg His His Val Asp Lys Pro Ser Lys  
 130 135 140

Val Ala Ser Met Val Arg Glu Asn Val Ile Ala Gln Leu Ala Asn Ile  
 145 150 155 160

Gln Thr His Pro Ser Val Arg Leu Ala Leu Glu Glu Gly Arg Val Thr  
 165 170 175

Leu His Gly Trp Ile Tyr Asp Ile Glu Thr Gly Gly Ile Asp Ala Phe  
 180 185 190

Asp Gly Ser Thr Gly Thr Phe Val Ser Leu Ala Glu Asn Pro Glu Val  
 195 200 205

His Ala Val Ser Gln Gln Ala Arg His Val Ala  
 210 215

<210> SEQ ID NO 65  
 <211> LENGTH: 217  
 <212> TYPE: PRT  
 <213> ORGANISM: Serratia proteamaculans  
 <220> FEATURE:

-continued

---

 <223> OTHER INFORMATION: Carbonate dehydratase *Serratia proteamaculans*  
568

&lt;400&gt; SEQUENCE: 65

Met Lys Glu Val Ile Glu Gly Phe Leu Lys Phe Gln Arg Glu Ala Phe  
 1 5 10 15  
 Val Glu Arg Thr Ala Leu Phe Gln Arg Leu Ala Thr Gln Gln Ser Pro  
 20 25 30  
 Arg Thr Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu  
 35 40 45  
 Ile Thr Gln Arg Glu Pro Gly Asp Leu Phe Val Ile Arg Asn Ala Gly  
 50 55 60  
 Asn Ile Val Pro Ser Phe Gly Pro Glu Pro Gly Gly Val Ser Ala Ser  
 65 70 75 80  
 Val Glu Tyr Ala Val Ser Ala Leu Gly Val Glu Asp Ile Val Ile Cys  
 85 90 95  
 Gly His Ser Asp Cys Gly Ala Met Thr Ala Ile Ala Thr Cys Gln Cys  
 100 105 110  
 Leu Gln His Met Pro Thr Val Ala Asn Trp Leu Arg Tyr Ala Asp Ser  
 115 120 125  
 Ala Lys Val Val Asn Gln Ala Tyr Gln His Ala Ser Glu Asn Glu Lys  
 130 135 140  
 Val Ser Ser Met Val Arg Glu Asn Val Ile Ala Gln Leu Asn Asn Ile  
 145 150 155 160  
 Lys Thr His Pro Ser Val Ala Leu Ala Leu Glu Gln Gly Arg Leu Lys  
 165 170 175  
 Leu His Gly Trp Val Tyr Asp Ile Ala Ser Gly Gly Ile Glu Ala Leu  
 180 185 190  
 Asp Gly Glu Thr Arg Arg Phe Ile Pro Leu Ala Thr Asn Pro Glu Val  
 195 200 205  
 Thr Ala Thr Pro Ala Val Ser Arg Phe  
 210 215

&lt;210&gt; SEQ ID NO 66

&lt;211&gt; LENGTH: 211

&lt;212&gt; TYPE: PRT

<213> ORGANISM: *Erwinia tasmaniensis*

&lt;220&gt; FEATURE:

 <223> OTHER INFORMATION: Carbonate dehydratase *Erwinia tasmaniensis*  
Et1/99

&lt;400&gt; SEQUENCE: 66

Met Gln His Ile Val Glu Gly Phe Leu Asn Phe Gln Lys Asp Ile Phe  
 1 5 10 15  
 Pro Glu Gln Lys Glu Leu Phe Arg Ser Leu Ala Ser Ser Gln Asn Pro  
 20 25 30  
 Lys Ala Leu Phe Ile Ser Cys Ser Asp Ser Arg Leu Val Pro Glu Leu  
 35 40 45  
 Val Thr Gln Gln Asp Pro Gly Gln Leu Phe Val Ile Arg Asn Ala Gly  
 50 55 60  
 Asn Ile Val Pro Ser Phe Gly Pro Glu Pro Gly Gly Val Ser Ala Thr  
 65 70 75 80  
 Ile Glu Tyr Ala Val Val Ala Leu Gly Val Ser Asp Ile Val Ile Cys  
 85 90 95

-continued

---

Gly His Ser Asn Cys Gly Ala Met Lys Ala Ile Ala Thr Cys Gln Cys  
 100 105 110

Leu Ala Pro Met Pro Ala Val Glu His Trp Leu Arg Tyr Ala Asp Ala  
 115 120 125

Ala Lys Ala Val Val Glu Lys Lys Asn Tyr Asp Thr Glu Glu Asp Lys  
 130 135 140

Val Asn Ala Met Val Gln Glu Asn Val Ile Ala Gln Leu Asn Asn Ile  
 145 150 155 160

Lys Thr His Pro Ser Val Ala Val Gly Leu Arg Asn Asn Ala Leu Arg  
 165 170 175

Leu His Gly Trp Val Tyr Asp Ile Glu Ser Gly Ala Ile Arg Ala Leu  
 180 185 190

Asp Lys Asp Ser Lys Lys Phe Val Leu Leu Ser Asp Asn Pro Gln Val  
 195 200 205

His Phe Glu  
 210

&lt;210&gt; SEQ ID NO 67

&lt;211&gt; LENGTH: 220

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Pseudomonas aeruginosa

&lt;220&gt; FEATURE:

<223> OTHER INFORMATION: Carbonate dehydratase Pseudomonas aeruginosa  
 PA7

&lt;400&gt; SEQUENCE: 67

Met Arg Asp Ile Ile Asp Gly Phe Leu Arg Phe Gln Arg Asp Ala Tyr  
 1 5 10 15

Pro Ala Arg Ser Gln Leu Phe Lys Ser Leu Ala Thr Arg Gln Ala Pro  
 20 25 30

Lys Ala Leu Phe Ile Ala Cys Ser Asp Ser Arg Val Val Pro Glu Leu  
 35 40 45

Leu Thr Gln Arg Glu Pro Gly Glu Leu Phe Val Ile Arg Asn Ala Gly  
 50 55 60

Asn Ile Val Pro Gly Tyr Gly Pro Gln Pro Gly Gly Val Ser Ala Ser  
 65 70 75 80

Val Glu Tyr Ala Val Ala Val Leu Gly Val Ala Asp Ile Val Val Cys  
 85 90 95

Gly His Ser Asp Cys Gly Ala Met Gly Ala Ile Ala Ser Cys Ala Cys  
 100 105 110

Leu Asp His Leu Pro Ala Val Ala Gly Trp Leu Arg His Ala Glu Ala  
 115 120 125

Ala Arg Ala Met Asn Ser Ala His Glu His Ser Ser Asp Ala Ala Arg  
 130 135 140

Leu Asp Ala Leu Val Arg His Asn Val Ile Ala Gln Leu Ala Asn Leu  
 145 150 155 160

Arg Thr His Pro Ser Val Ala Arg Ala Leu Glu Gln Gly Arg Leu Asn  
 165 170 175

Leu His Gly Trp Val Tyr Asp Ile Glu Ser Gly Arg Ile Asp Ala Leu  
 180 185 190

Asp Gly Ala Ser Arg Arg Phe Val Ser Leu Ala Glu His Pro Gly Val  
 195 200 205

Arg Ala Val Gly Gly Glu Pro Gly Gln Ala Val Ala  
 210 215 220

-continued

---

<210> SEQ ID NO 68  
 <211> LENGTH: 202  
 <212> TYPE: PRT  
 <213> ORGANISM: Streptosporangium roseum  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Carbonate dehydratase Streptosporangium roseum  
 DSM 43021

<400> SEQUENCE: 68

Met Gln Asp Leu Glu Glu Gly Val Ala Arg Phe Gln Arg Asp Val Phe  
 1 5 10 15  
 Pro Ala Lys Thr Glu Leu Phe Thr Arg Leu Ala Thr Ala His Gln Pro  
 20 25 30  
 Ala Thr Leu Phe Ile Ser Cys Ser Asp Ala Arg Val Val Pro Glu Leu  
 35 40 45  
 Ile Thr Gln Ser Glu Pro Gly Glu Leu Phe Val Ile Arg Thr Ala Gly  
 50 55 60  
 Asn Leu Val Pro Ala Tyr Ala Pro Gly Ser Ala Asp Gly Val Ala Ala  
 65 70 75 80  
 Gly Ile Glu Tyr Ala Val Ala Val Leu Gly Val Ser Asp Ile Val Val  
 85 90 95  
 Cys Gly His Ser Gly Cys Gly Ala Met Thr Ala Val Ala Asp Gly Leu  
 100 105 110  
 Asp Pro Ala Ala Leu Pro Ala Val Ala Gly Trp Leu Arg His Ala Asp  
 115 120 125  
 Ala Ser Arg Ala Arg Val Thr Thr Thr Glu Thr Gly Thr Gly Glu Val  
 130 135 140  
 Ala Ala Leu Val Arg Gln Asn Val Leu Thr Gln Leu Ala Asn Leu Ala  
 145 150 155 160  
 Thr His Pro Ser Val Ala His Ala Leu Ala Gly Lys Thr Val Thr Leu  
 165 170 175  
 His Gly Trp Ile Tyr Asp Ile Gly Thr Gly Thr Val Ala Glu Leu Asp  
 180 185 190  
 Ala Thr Gly Arg Pro Ser Ala Leu Ala Val  
 195 200

---

What is claimed is:

**1-43.** (canceled)

**44.** A method for producing an acrylic acid-based consumer product comprising

- i) combining a carbon source and a microorganism to produce 3-hydroxypropionic acid, wherein said microorganism comprises
  - a) a heterologous nucleic acid sequence encoding an enzyme selected from the group consisting of: a native or mutated form of a mmsB protein, and a native or mutated form of a ydfG protein;
  - b) a genetic modification to decrease or eliminate an enzymatic activity selected from the group consisting of: lactate dehydrogenase, phosphate acetyltransferase, and pyruvate-formate lyase;
  - c) a heterologous nucleic acid sequence encoding an acetyl-CoA carboxylase; and
  - d) a heterologous nucleic acid sequence encoding a cyanase,

ii) converting said 3-hydroxypropionic acid to acrylic acid; and

iii) processing said acrylic acid into a consumer product.

**45.** The method of claim **44**, wherein said carbon source has a ratio of carbon-14 to carbon-12 of about  $1.0 \times 10^{-14}$  or greater.

**46.** The method of claim **44**, wherein following (i) said method further comprises separating or purifying 3-hydroxypropionic acid from said cell culture by extraction of 3-hydroxypropionic acid from said culture in the presence of a tertiary amine.

**47.** The method of claim **44**, wherein said 3-hydroxypropionic acid is produced at a higher level than in a non-genetically modified microorganism.

**48.** The method of claim **44**, wherein said consumer product is selected from the group consisting of diapers, carpet, paint, adhesives, and acrylic glass.

**49.** The method of claim **48**, wherein said consumer product is diapers.

**50.** The method of claim **44**, wherein said carbon source is predominantly glucose, sucrose, fructose, dextrose, lactose, or a combination thereof.

**51.** The method of claim **44**, wherein said carbon source is less than 50% glycerol.

**52.** The method of claim **44**, wherein said lactate dehydrogenase is encoded by a nucleic acid sequence having at least 90% sequence identity to SEQ ID NO:19 or has an amino acid sequence having at least 90% sequence identity to SEQ ID NO: 20.

**53.** The method of claim **44** wherein said pyruvate-formate lyase is encoded by a nucleic acid sequence having at least 90% sequence identity to SEQ ID NOs: 21 or has an amino acid sequence having at least 90% sequence identity to SEQ ID NO: 22.

**54.** The method of claim **44**, wherein said genetically modified microorganism further comprises a heterologous nucleic acid sequence encoding an acetyl-CoA carboxylase.

**55.** A method for producing an acrylic acid-based consumer product comprising

- i) combining a carbon source and a microorganism to produce 3-hydroxypropionic acid, wherein said microorganism comprises
  - a) a heterologous nucleic acid sequence encoding a 3-hydroxypropionate dehydrogenase;
  - b) a heterologous nucleic acid sequence encoding a cyanase or a carbonic anhydrase; and
  - c) a heterologous nucleic acid sequence encoding an NADPH-dependent transhydrogenase or an acetyl-CoA carboxylase;
- ii) converting said 3-hydroxypropionic acid to acrylic acid; and
- iii) processing said acrylic acid into a consumer product.

**56.** The method of claim **55**, wherein said genetically modified microorganism further comprises at least one heterologous nucleic acid sequence encoding an oxaloacetate alpha-decarboxylase.

**57.** The method of claim **56**, wherein said oxaloacetate alpha-decarboxylase is encoded by a sequence selected from the group consisting of: SEQ ID NOs: 54, 55, and 56.

**58.** The method of claim **56**, wherein said genetically modified microorganism further comprises at least one heterologous nucleic acid sequence encoding a phosphoenolpyruvate carboxykinase or a phosphoenol pyruvate carboxylase.

**59.** The method of claims **56**, wherein said genetically modified microorganism further comprises at least one genetic modification to reduce enzymatic activity of a protein selected from the group consisting of: lactate dehydrogenase, pyruvate formate lyase, phosphate acetyltransferase, heat stable, histidyl phosphorylatable protein, phosphoryl transfer protein, polypeptide chain, pyruvate kinase I, and pyruvate kinase II.

**60.** The method of claim **55**, wherein said genetically modified microorganism further comprises a modification of a gene to increase the NADPH/NADP<sup>+</sup> ratio, wherein the modification is selected from the group consisting of: increasing activity of *pgi*, increasing activity of *pntAB*, *gapA:gapN* substitution or replacement, and disrupting *sthA*.

**61.** The method of claim **55**, wherein the 3-hydroxypropionate dehydrogenase is a native or mutated form of a *mmsB* protein or a native or mutated form of a *ydfG* protein.

**62.** The method of claim **55**, wherein the said genetically modified microorganism further comprises a heterologous nucleic acid sequence encoding an NADPH-dependent transhydrogenase.

**63.** The method of claim **55**, wherein said genetically modified microorganism further comprises a heterologous nucleic acid sequence encoding a cyanase or a carbonic anhydrase that increases intracellular bicarbonate levels.

**64.** The method of claim **55**, wherein said genetically modified microorganism further comprises a heterologous nucleic acid sequence encoding an acetyl-CoA carboxylase.

**65.** The method of claim **55**, wherein said genetically modified microorganism is selected from the group consisting of: *Zymomonas*, *Escherichia*, *Pseudomonas*, *Alcaligenes*, *Salmonella*, *Shigella*, *Burkholderia*, *Oligotropha*, and *Klebsiella*.

**66.** The method of claim **55**, wherein said genetically modified microorganism is selected from the group consisting of: *Escherichia coli*, *Cupriavidus necator*, *Oligotropha carboxidovorans*, and *Pseudomonas putida*.

**67.** The method of claim **55**, wherein said genetically modified microorganism is an *Escherichia coli*.

**68.** The method of claim **55**, wherein said genetically modified microorganism is a gram-positive bacterium.

**69.** The method of claim **55**, wherein said genetically modified microorganism is selected from the group consisting of: *Clostridium*, *Rhodococcus*, *Bacillus*, *Lactobacillus*, *Enterococcus*, *Paenibacillus*, *Arthrobacter*, *Corynebacterium*, and *Brevibacterium*.

**70.** The method of claim **55**, wherein said genetically modified microorganism is selected from the group consisting of: *Bacillus licheniformis*, *Paenibacillus macerans*, *Rhodococcus erythropolis*, *Lactobacillus plantarum*, *Enterococcus faecium*, *Enterococcus gallinarum*, *Enterococcus faecalis*, and *Bacillus subtilis*.

**71.** The method of claim **55**, wherein said genetically modified microorganism is a fungus.

**72.** The method of claim **55**, wherein said genetically modified microorganism is a yeast.

**73.** The method of claim **55**, wherein said yeast is selected from the group consisting of: *Pichia*, *Candida*, *Hansenula*, and *Saccharomyces*.

**74.** A method for producing an acrylic acid-based consumer product comprising

- i) combining a carbon source and a microorganism to produce 3-hydroxypropionic acid, wherein said microorganism comprises
  - a) a heterologous nucleic acid sequence encoding a 3-hydroxypropionate dehydrogenase; and
  - b) a heterologous nucleic acid sequence encoding a cyanase or a carbonic anhydrase,
- ii) converting said 3-hydroxypropionic acid to acrylic acid; and
- iii) processing said acrylic acid into a consumer product.

**75.** The method of claim **72**, wherein said genetically modified microorganism further comprises a heterologous nucleic acid sequence encoding an acetyl-CoA carboxylase.