

US 20150335029A1

(19) **United States**

(12) **Patent Application Publication**
MITTER et al.

(10) **Pub. No.: US 2015/0335029 A1**

(43) **Pub. Date: Nov. 26, 2015**

(54) **METHOD FOR PRODUCING PLANT SEED
CONTAINING ENDOPHYTIC
MICRO-ORGANISMS**

(30) **Foreign Application Priority Data**

Jun. 22, 2012 (EP) 12173124.4

Publication Classification

(71) Applicant: **AIT AUSTRIAN INSTITUTE OF
TECHNOLOGY GMBH**, Vienna (AT)

(51) **Int. Cl.**
A01N 63/00 (2006.01)

(72) Inventors: **Birgit MITTER**, Giesshübl (AT);
Angela SESSITSCH, Vienna (AT);
Muhammad NAVEED, Tulln (AT)

(52) **U.S. Cl.**
CPC **A01N 63/00** (2013.01)

(57) **ABSTRACT**

(21) Appl. No.: **14/410,537**

The invention discloses a method for producing plant seed containing endophytic micro-organisms characterised in by the following steps: —contacting a flowering plant with a preparation of endophytic microorganisms, whereby the endophytic microorganisms enter the plant via the flowers and are conveyed to seed produced by the plant; and—obtaining the plant seed containing endophytic microorganisms from the plant.

(22) PCT Filed: **Jun. 21, 2013**

(86) PCT No.: **PCT/EP2013/062976**

§ 371 (c)(1),

(2) Date: **Dec. 22, 2014**

Fig. 1




Fig. 2a





Fig. 2b




Fig. 2c




Fig. 3



Fig. 4a




Fig. 4b



Fig. 4c




Fig. 5a



Fig. 5b



Fig. 5c




Fig. 6




Fig. 7

METHOD FOR PRODUCING PLANT SEED CONTAINING ENDOPHYTIC MICRO-ORGANISMS

[0001] The present invention relates to the production of plant seeds comprising endophytes.

[0002] In spite of limited arable land coupled with rising demand of a steadily increasing human population which could hit 9 billion by 2050, food supply is a global challenge making production of economically attractive and high quality food, free from unacceptable levels of chemicals, a dire need. The use of microorganisms with the aim of improving plant growth and health is an important practice and necessary for agriculture.

[0003] During the past couple of decades, plant growth promoting rhizobacteria (PGPR) have received worldwide importance and acceptance in agricultural practice. These microorganisms are the potential tools for sustainable agriculture because they not only ensure the availability of essential nutrients to plants but also enhance the nutrient use efficiency.

[0004] Endophytic bacteria may in future be even more important than rhizosphere bacteria in promoting plant growth because they escape competition with rhizosphere microorganisms and establish a more intimate contact with plant tissues. In addition, the inherent nature of certain endophytes to potentially colonize plants in a systematic manner provides a novel approach as a delivery system to plants for various beneficial traits.

[0005] Bacterial mechanisms of plant growth promotion include biological nitrogen fixation (BNF), synthesis of phytohormones and vitamins, environmental stress relief, synergism with other bacteria-plant interactions, inhibition of plant ethylene synthesis, as well as increasing availability of nutrients like phosphorus, iron and other micro-elements, and growth enhancement by volatile compounds.

[0006] Numerous application strategies have been used for PGPR/endophytic bacteria at the experimental levels, ranging from seed treatment and soil application to stem injection and foliar spray. Seed treatment (soaking and embedding in carrier material) with bacterial inocula prior to sowing is the traditional, most commonly used and easiest means of inoculation. Peat (carrier based mixing) inoculants have been the standard for the inoculation industry; however, several other commercial preparations have been marketed. Crop Genetics International Ltd. developed a seed inoculation technique by applying a pressure differential to infuse the bacterial suspension into imbibed seeds and re-drying the seeds (U.S. Pat. No. 5,415,672 A).

[0007] To get benefits from bacterial inocula, it is crucial to apply (technique and timing) bacterial inocula in a viable way. In addition, it is equally important that the microorganisms remain viable during several months of seed storage and are easily activated and colonize the plant environment. However, by using conventional methods (carrier based, liquid broth and soil application; see also: U.S. Pat. No. 7,084,331 B2, U.S. Pat. No. 7,906,313 B2, U.S. Pat. No. 7,037,879 B2), the viability of bacteria is subjected to the hazards of drying, fertilizer contact, seed coat toxicity, incompatible pesticidal and mineral additives. Besides this, several soil and environmental stresses affect the survival/colonization efficiency of the inoculant strains. Bacterial population density, the host plant species, agronomic practices and climatic conditions are among the important factors for the success of biological plant fertilization. Examples for use of endophytes as plant

growth enhancers, bio-pesticides, pathogen treatment or pest tolerance agents are disclosed e.g. in WO 00/29607 A1, WO 2011/117351 A1, WO 2010/115156 A2, WO 2007/107000 A1, WO 2007/021200 A1, US 2012/144533 A1, U.S. Pat. No. 4,940,834 A, CA 2562175 A1 and WO 2011/082455 A1.

[0008] With current inoculation methods, however, colonisation of the plants with the desired endophytic microorganisms is difficult and often not reproducible, which makes it difficult to apply this technology on an industrial scale. For example, microorganisms used in seed coating often do not survive well or are unable to colonize the plant (because the microorganisms on the outside cannot enter the seed or plant). If the plant is mechanically (or otherwise) wounded to provide an entry, this puts the health of the seed, seedlings or plant at risk, because harmful microorganisms could enter the plant as well in an unprotected manner. Moreover, even if the microorganisms can colonise a given plant, there can be a natural loss of viability and the efficiency of colonization can be low. More complex inoculation techniques (e.g. by applying vacuum or pressure infiltration, inoculation by injection, etc.) are also causing risk for the plant and are—most importantly—not transferable to a large scale or industrial applicability and are therefore not effective.

[0009] It is an object of the present invention to provide an improved method for producing seeds containing endophytic microorganisms. The method should provide seeds with a reproducible and defined composition of endophytic microorganisms enabling the growth of plants with the desired properties due to the presence of such endophytic microorganisms. It is another object to provide methods for introducing endophytic microorganisms into plant seeds for microorganisms that are not or are not necessarily occurring in seeds.

[0010] Therefore, the invention provides a method for producing plant seed containing endophytic microorganisms which is characterised by the following steps:

[0011] contacting a flowering plant with a preparation of endophytic microorganisms, whereby the endophytic microorganisms enter the plant via the flowers and are conveyed to seed produced by the plant; and

[0012] obtaining the plant seed containing endophytic microorganisms from the plant.

[0013] The term “endophyte” means—in its broadest meaning—the location of an organism, with “endo” means “inside” and “phyte” means “plants”. Therefore, endophyte—in its broadest meaning—refers to organisms that live within plants. Fungi and bacteria are the most common organisms associated with the term endophyte.

[0014] An important feature of endophytic microorganisms is that they occupy internal tissues of plants without causing substantive damage to their hosts. In many cases endophytes are responsible for conferring one or more advantages to the plant. For the present invention, an “endophytic microorganism” is defined in this usual way: as a microorganism that lives within a plant and is responsible for plant beneficial effects, for example tolerance to drought, metals, disease (e.g. increasing resistance to pathogens and parasites), and herbivory, and/or growth promotion and nutrient acquisition, production of phytohormones, antibiotics (protection against microorganisms being harmful for seeds and plants) or siderophores, pesticides; promoting biological nitrogen fixation, etc. (as some (of many) examples: chilling tolerance (*Burkholderia*), salinity stress (*Achrobacter*, *Azospirillum*), tolerance to drought (*Burkholderia*, *Pantoea*), metals, disease (*Bacillus*, *Pseudomonas*, *Xanthomonas*), growth promotion

(*Azospirillum*, *Azotobacter*, *Bacillus*, *Burkholderia*, *Enterobacter*, *Klebsiella*, *Pantoea* and *Pseudomonas*) and nutrient acquisition (*Pseudomonas*, *Bacillus*, *Rhizobium*, *Micrococcus*, *Flavobacterium*, *Burkholderia*, *Achromobacter*, *Erwinia*, and *Agrobacterium*) (U.S. Pat. No. 7,906,313 B2)).

[0015] Endophytic organisms associated with plants are varied and complex. Endophytic microbes occupy a relatively privileged niche within a plant and frequently contribute to plant health or growth. Co-evolution may exist between endophytes and their host e.g. in resisting to environmental stresses. Endophytes have been targeted as valuable sources of new bioactive compounds. Endophytes inhabit plant tissues, particularly the so-called intercellular space, space between cells. Endophytic microorganisms have been found in virtually every plant studied, where they colonize the internal tissues of their host plant and can form a range of different relationships including symbiotic, mutualistic, commensalistic and trophobiotic. Most endophytes appear to originate from the rhizosphere or phyllosphere; however, some may be transmitted through the seed. Endophytic microorganisms can promote plant growth and yield and can act as biocontrol agents. Endophytes can also be beneficial to their host by producing a range of natural products that are not only beneficial to the plant but could also be harnessed for potential use in medicine, agriculture or industry. In addition, it has been shown that they have the potential to remove soil contaminants by enhancing phytoremediation and may play a role in soil fertility through phosphate solubilisation and nitrogen fixation. There is increasing interest in developing the potential biotechnological applications of endophytes for improving phytoremediation and the sustainable production of non-food crops for biomass and biofuel production.

[0016] The method according to the present invention uses a completely new and diligent way for introducing endophytes into plants which turned out in the course of the present invention to be very effective, yet without harming the plants or seeds during or after inoculation. Applying the microorganisms to the flowering plants uses a natural entry into the plant which introduces the endophyte in an efficient manner into the next generation plant seeds. Within the course of the present invention it also turned out that when the microorganisms are applied to the plant at the time of flowering (e.g. by simple spraying), the microorganisms get entry when grain formation starts and establish populations inside the seed. The method of the present invention can facilitate the crop productivity by enhancing germination, seedling vigour and biomass in comparison with non-treated control. Moreover, the introduction of the beneficial microorganisms inside seed instead of external application by e.g. seed coating makes the inocula less susceptible to environmental perturbation and better compatible to chemical seed coatings (pesticides and herbicides). Using bacterial colonized seeds, the plant growth and biomass are statistically similar as the conventional inoculation method e.g. exogenous seed soaking and soil inoculation (that are more laborious and less practicable in certain circumstances).

[0017] Accordingly, the present invention provides a new concept of applying endophyte strains for improved plant growth and vitality—the integration of the bacteria or fungus strain inside the plant seed. The microorganisms are e.g. sprayed on the parent flowering plants, enter the plants and colonize the emerging seeds. The microorganisms may also be applied by specific instruments to the flower, e.g. by a spatula, a syringe or an inoculating loop. Another preferred

embodiment for administering the endophytes to the flower of a plant is performed by employing pollen-feeding insects, preferably humble-bees, that carry the endophytic microorganisms. Such insects (besides humble-bees also honey-bees, butterflies, some wasp and fly species or other “pollinators” may be used) can even be provided from commercial sources and contacted with the endophytes before they are released to contact the flowering plants. The microorganisms are preferably provided at a body part of these insects that has the highest probability to contact the flower of the plant (e.g. the legs or the ventral part of the body).

[0018] By planting the internally colonized seeds the endophytes get activated and proliferate and colonize the offspring generation plants. Internally colonized seeds may result (depending on the nature of the endophyte) in improved biomass production and plant vitality in the subsequent plant generation. The positive effects are at least comparable (if not improved) to that observed after external application of endophytes, but makes the inocula less susceptible to environmental perturbation and better compatible to chemical seed coatings (pesticides and herbicides). With the present invention it is also possible to introduce endophytes into seeds that are not or are not necessarily present in seeds. Virtually any kind of endophytes can be introduced into seeds by the method according to the present, provided that these endophytes have at least a basic power to establish themselves in the seeds.

[0019] None of the prior art methods, especially not the methods disclosed in WO 00/29607 A1, WO 2011/117351 A1, WO 2010/115156 A2, WO 2007/107000 A1, WO 2007/021200 A1, US 2012/144533 A1, U.S. Pat. No. 4,940,834 A, CA 2562175 A1 and WO 2011/082455 A1, aim at providing methods for providing seeds comprising selected endophytes. The main goal of these methods according to the prior art is always the provision of the endophytes to the very plant treated and not—as in the present invention—to supply a mother plant with the endophytes of interest and to obtain endophyte containing seeds from this mother plant for rising daughter plants already containing the endophytes and, optionally, passing the endophytes further to their own daughter generation. Accordingly, the technology provided with the present invention can provide seeds with completely novel characteristics, e.g. having a unique set-up of endophytes (for example by having one single endophyte species being predominantly present in the seeds (e.g. representing more than 50%, or more than 70% or even more than 80% of the total of endophytes in the seed)).

[0020] Suitable plants include both monocots and dicots (including eudicots) that can be colonized by the endophytic microorganisms according to the present invention. Of course, the plant has to be a flowering plant (angiosperm) in order to transfer the microorganisms to the plant in the course of the flowering phase. The resulting seeds contain the inoculated endophytes in an efficient concentration. Plants grown from such seeds contain the endophytes and the beneficial properties of the endophyte can develop in the seeds or plants. Accordingly, the plants arising from such seeds—wherein the endophyte can express its beneficial function to the plant—may be at any stage of growth, including seeds, seedlings, or full plants. The present invention is therefore not simply about spraying the microorganisms to a given plant (or seed) in order to provide the beneficial endophytic effect to this plant, but it provides a method which safeguards presence of endophytes in the seeds generated from this plant and therefore for the next generations of the plant. This essentially

differs from all inoculation strategies applied before (seed impregnation, spraying the microorganisms to the seeds, germs or the whole plants), because the present method deals with the production of seeds which contain a reproducible endophyte set-up.

[0021] In a preferred embodiment, the target plant is a plant of the family Graminae (grasses). The grass plants into which these endophytes are introduced may be any of the useful grasses belonging to the genera *Agropyron*, *Agrostis*, *Andropogon*, *Anthoxanthum*, *Arrhenatherum*, *Avena*, *Brachypodium*, *Bromus*, *Chloris*, *Cynodon*, *Dactylis*, *Elymus*, *Eragrostis*, *Festuca*, *Glyceria*, *Hierochloe*, *Hordeum*, *Lolium*, *Oryza*, *Panicum*, *Paspalum*, *Phalaris*, *Phleum*, *Poa*, *Setaria*, *Sorghum*, *Triticum*, *Zea* and *Zoysia*.

[0022] In a preferred embodiment, the target plant is selected from the wheats, including, *Triticum monococcum*, *Triticum turgidum*, *Triticum timopheevi* (Timopheev's Wheat) and *Triticum aestivum* (Bread Wheat).

[0023] In another preferred embodiment, the target plant is a corn of the genus *Zea*. *Zea* is a genus of the family Gramineae (Poaceae), commonly known as the grass family. The genus consists of some four species: *Zea mays*, cultivated corn and teosinte; *Zea diploperennis* Iltis et al., diploperennial teosinte; *Zea luxurians* (Durieu et Asch.) Bird; and *Zea perennis* (Hitchc.) Reeves et Mangelsd., perennial teosinte.

[0024] Other useful grasses which may be used on an industrial basis are rye grasses and bluegrasses. Bluegrasses known in the art include Kentucky bluegrass, Canada bluegrass, rough meadow grass, bulbous meadow grass, alpine meadow grass, wavy meadow grass, wood meadow grass, Balforth meadow grass, swamp meadow grass, broad leaf meadow grass, narrow leaf meadow grass, smooth meadow grass, spreading meadow grass and flattened meadow grass.

[0025] In a preferred embodiment, the plants for which seeds are produced by the method according to the present invention are dicots, including eudicots such as tomato, watermelon, squash, cucumber, strawberry, pepper, soybean, peanut, Brassicaceae, especially rape, sunflower, sugar beet, cotton, alfalfa and arabidopsis.

[0026] Accordingly, the plant is preferably selected from the group of Graminae (grasses), preferably grasses of the genera *Agropyron*, *Agrostis*, *Andropogon*, *Anthoxanthum*, *Arrhenatherum*, *Avena*, *Brachypodium*, *Bromus*, *Chloris*, *Cynodon*, *Dactylis*, *Elymus*, *Eragrostis*, *Festuca*, *Glyceria*, *Hierochloe*, *Hordeum*, *Lolium*, *Oryza*, *Panicum*, *Paspalum*, *Phalaris*, *Phleum*, *Poa*, *Setaria*, *Sorghum*, *Triticum*, *Zea*, especially *Zea mays*, cultivated corn and teosinte, *Zea diploperennis* Iltis et al., diploperennial teosinte, *Zea luxurians* (Durieu et Asch.) Bird; and *Zea perennis* (Hitchc.) Reeves et Mangelsd., perennial teosinte. and *Zoysia*; wheats, preferably *Triticum monococcum*, *Triticum turgidum*, *Triticum timopheevi* (Timopheev's Wheat) and *Triticum aestivum* (Bread Wheat); preferably rye grasses and bluegrasses, especially Kentucky bluegrass, Canada bluegrass, rough meadow grass, bulbous meadow grass, alpine meadow grass, wavy meadow grass, wood meadow grass, Balforth meadow grass, swamp meadow grass, broad leaf meadow grass, narrow leaf meadow grass, smooth meadow grass, spreading meadow grass and flattened meadow grass; dicots, preferably eudicots, especially tomato, watermelon, squash, cucumber, strawberry, pepper, soybean, peanut, Brassicaceae, especially rape, sunflower, sugar beet, cotton, alfalfa and arabidopsis.

[0027] The method according to the present invention is specifically suitable for providing seeds of transgenic plants.

By the present invention, transgenic plants are obtainable that—besides their advantageous properties provided by the transgene—also contain “tailored” endophyte properties that can selectively be construed and provided by the present invention.

[0028] According to a preferred embodiment of the present method the endophytic microorganism is an endophytic bacterium, preferably selected from *Burkholderia*, *Rhizobium*, *Bradyrhizobium*, *Mesorhizobium*, and *Sinorhizobium*, *Herbaspirillum*, *Azospirillum*, *Acetobacter*, *Arthrobacter*, *Bacillus*, *Paenibacillus*, *Streptomyces*, *Enterobacter*, and *Pseudomonas*, *Pantoea* and *Enterobacter*, especially *Burkholderia phytofirmans*.

[0029] According to another preferred embodiment, the endophytic microorganism is an endophytic fungus, preferably selected from *Curvularia*, *Mycorrhiza*, *Pisifimospora*, *Trichoderma*, and *Colletotrichum*.

[0030] In a preferred embodiment according to the present invention, contacting the flower of a plant with a preparation of endophytic microorganisms is performed via spraying the microorganisms at the time of flowering. Spraying is specifically useful in an industrial production method. Other methods include the inoculation by brushing, by an inoculating loop, by applying droplets, etc.; however, spraying can be easily automated, e.g. in glasshouse cultures.

[0031] Inoculation is done by applying the culture of the endophyte to the flowering plant. It is recommendable to safeguard conditions which are favourable to the microorganisms used. The microorganisms are usually applied in suspension at a suitable concentration. Accordingly, it is preferred to contact the flower of a plant with a preparation of endophytic microorganisms by applying the microorganisms in a suspension of 10^6 to 10^{10} cfu/mL, preferably of 10^7 to 10^9 cfu/mL, especially of 10^8 to 10^9 cfu/mL.

[0032] The seeds obtained by the present method can be treated like normal seeds. The beneficial properties (the endophytes) remain safely packed inside the seed preventing the exposure of hazards from outside (which usually causes damage to cultures which are applied when the seeds are only coated). Accordingly, the seeds may be stored for considerable time without significant loss of their endophytic activity. Preferably, the plant seed obtained by the present method containing endophytic microorganisms from the plant is stored for at least 1 month, preferably for at least 3 months, especially for at least 6 months.

[0033] Also much longer storage times are, of course, possible for the seeds produced according to the present invention. It is therefore also preferred that the plant seed obtained by the present method containing endophytic microorganisms from the plant is stored for at least 12 months, preferably for at least 2 years, especially for at least 3 years.

[0034] The method according to the present invention is suitable for providing virtually any endophyte-containing seed, because the transfer of the microorganisms from the flower to the seed is a way with low hazard exposure (to plant and endophyte). It is specifically suitable for producing seeds with an endophyte which is in principle known to naturally proliferate in plants, especially in the given plant, i.e. a “naturally obtainable endophyte”. These endophytes are derivable from natural sources from the same plant type or from other plant types. According to a preferred embodiment, the endophytic microorganism is therefore a naturally obtainable endophyte.

[0035] It is also possible to use the present method for providing seeds with artificially created or optimised microorganisms, e.g. recombinantly engineered bacteria or fungi; or strains which have been optimised by various culture techniques and/or selection rounds. Another preferred embodiment of the present invention is therefore to use a recombinantly produced bacterium as the endophytic microorganism.

[0036] As already mentioned, the seeds obtained by the present method can be further processed in usual ways. For example, it can be treated with various substances which further promote the plants to be produced from the seeds, e.g. by impregnating the seeds with growth promoting agents or other chemicals beneficial for plant health, such as herbicides, pesticides, antibiotics, etc. It is, of course, also possible to provide a coating with further (or the same) endophytic microorganisms as the microorganism according to the present invention. According to a preferred embodiment of the present invention, the obtained plant seed containing endophytic microorganisms is therefore subjected to a seed impregnation step.

[0037] This invention also relates to the seeds obtainable by the method according to the present invention which show—compared to seeds according to the prior art—a unique endophyte set-up.

[0038] According to a preferred embodiment, the present invention provides seeds which can be grown to plants that are improved (compared to the wild type plants) with respect to stress tolerance. “Stress” in this context may be an environmental stress, including, high temperature, drought, metals and metal ions, which cause a variety of plant problems and/or death, and abnormal pH (including both acidic and/or alkaline). With the seeds produced by the present invention plants can be obtained that have reproducibly improved stress resistance, e.g. at least about a 5, 10, 20, 25 and 50% change in thermotolerance, at least about a 5, 10, 20, 25 and 50% change in drought tolerance, at least about a 5, 10, 20, 25 and 50% change in metal tolerance, or at least about a 5, 10, 20, 25 and 50% change in pH tolerance (each as measured according to U.S. Pat. No. 7,906,313 B2, and compared to controls without the method according to the present invention applied).

[0039] According to a preferred embodiment, the seeds according to the present invention can be grown to plants with increased growth. Growth enhancement is generally measured as a comparison of plants cultured from seeds made according to the present invention with control plants lacking this endophyte composition. Differences in plant size, including leaf, root and stems are generally measured by weight, with increased growth being measured as at least about an at least 2% difference, preferably an at least 3% difference (which can already be regarded as a very significant gain in yield. Even more preferred, in some instances, a 5-10% difference between control plants and the plants grown from the seeds according to the present invention may be obtained, with at least about a 25% difference being specifically preferred.

[0040] The method according to the present invention enables the creation of completely new seeds/endophyte combinations. One of the most significant properties of preferred seeds obtainable by the present invention is the possibility to provide predominant endophyte populations in the seeds. Normally, seeds containing endophytes contain a diverse population of many different endophytic microorganisms with usually more than 10 or even more than 20 different

identifiable culturable strains (or even more than 30)(but none of these strains being predominant), the method according to the present invention enables the production of seeds with a predominant species of endophytic microorganism. Accordingly, preferred seed preparations which are provided by the present invention contain seeds having an endophytic microorganism population wherein more than 30%, preferably more than 40%, especially more than 50%, of the endophytic microorganisms represent the inoculant strain. This means that most (more than 50%, preferably more than 60%, especially more than 70%) of the seeds in the preparation contain more than 30%, preferably more than 40%, especially more than 50%, endophytic microorganisms comprising the inoculant strain.

[0041] It is even possible to provide a seed preparation containing seeds, wherein more than 60%, preferably more than 70%, more preferred more than 80%, especially more than 90%, endophytic microorganisms of a single species (the endophytic microorganism of the inoculant strain). This enables the production of seeds containing e.g. more than 60%, preferably more than 70%, especially more than 80%, of the applied endophytic strain (e.g. within a single field).

[0042] A specific embodiment of the present invention is therefore a seed preparation obtainable by a method according to the present method.

[0043] According to a preferred embodiment, the present invention provides a seed preparation containing seeds having more than 30%, preferably more than 40%, especially more than 50%, of the endophytic microorganisms are *Burkholderia phytofirmans*, especially *Burkholderia phytofirmans* PsJN (DSM17436); *Pantoea* sp. FD17 or *Paenibacillus* sp. S10., *Actinobacter* sp. S9, *Bradyrhizobium* sp. NC92 and *Bradyrhizobium japonicum* TAL379.

[0044] The present invention also provides seeds obtainable by the method according to the present invention with unique characteristics, e.g. with a predominant endophyte species as disclosed above. A preferred embodiment of the present invention is therefore drawn to seeds, especially maize seeds, obtainable by a method according to the present invention, wherein the endophytic microorganisms are preferably present in a population density of 10^2 to 10^5 cfu/g fresh weight.

[0045] According to a preferred embodiment, the present invention provides maize seed obtainable by a method according to the present invention, preferably wherein the endophytic microorganisms are *Burkholderia phytofirmans*, especially in a population density of 10^2 to 10^5 cfu/g fresh weight of seed. It is known that in maize, usually the viable population densities are much lower (for sweet corn, it was reported that such concentrations are below 10^1 cfu/g fresh weight (Kaga et al. Microbes Environ 24 (2009), 154-162)); in contrast thereto, the seeds according to this preferred embodiment contain at least 10^2 , preferably at least 10^3 , especially at least 10^4 , cfu/g fresh weight of one species, especially of *Burkholderia phytofirmans* (strain PsJN). Accordingly, the endophyte concentration of such seeds contains a predominant strain, which is not the case in natural plants or plants having been inoculated with prior art inoculation methods.

[0046] The seeds according to the present invention provide a marketable seed product containing a predetermined weight or volume of seeds with a uniform endophyte composition. For example, a marketable seed product containing at least 100 g seeds, preferably at least 1 kg seeds, more pre-

ferred at least 5 kg seeds, especially at least 10 kg seeds, can be provided by the method according to the present invention that contains—as a whole product—more than 30%, preferably more than 40%, especially more than 50%, of a single species of an endophytic microorganism, i.e. the inoculant strain. According to a preferred embodiment, the present invention provides a marketable seed product containing at least 100 g seeds, preferably at least 1 kg seeds, more preferred at least 5 kg seeds, especially at least 10 kg seeds, wherein—as a whole product—more than 50%, preferably more than 60%, especially more than 70% of a single species of an endophytic microorganism, i.e. the inoculant strain, are contained. According to an even more preferred embodiment, the present invention provides a marketable seed product containing at least 100 g seeds, preferably at least 1 kg seeds, more preferred at least 5 kg seeds, especially at least 10 kg seeds, wherein—as a whole product—more than 75%, more preferably more than 80%, especially more than 90%, endophytic microorganism of a single species (the endophytic microorganism of the inoculant strain) are contained.

[0047] Such uniformity in endophytic composition is unique and is extremely advantageous for high-tech and/or industrial agriculture. It allows significant standardisation with respect to qualitative endophyte load of seed products. The term “marketable seed product” means any commercially usable product containing plant seeds in a suitable package (e.g. a box, a bag, an envelope or any other container used for storing, shipping or offering plant seeds for sale). Suitable volumes or weights are those that are currently used for plant seeds (i.e. the at least 100 g, at least 1, 5 or 10 kg; but also 25 or more, 40 or more, 50 kg or more, even 100 kg or more, 500 kg or more, 1 t or more, etc.). Suitable containers or packages are those traditionally used in plant seed commercialisation; however, also other containers with more sophisticated storage capabilities (e.g. with microbiologically tight wrappings or with gas- or water-proof containments) can be used. The amount of endophytes (qualitatively and quantitatively) contained in the seeds or in the marketable seed product as a whole can be determined by standard techniques in microbiology readily available to any person skilled in the art of plant endophyte analysis.

[0048] The invention is further described by way of the following examples and the drawing figures, yet without being restricted thereto.

[0049] FIG. 1 shows cob sheath, grain and cob interior colonization of *Burkholderia phytofirmans* strain PsJN in maize cvs Peso and Morignon (x-axis shows CFU/g dry weight);

[0050] FIG. 2 shows light microscopy images of a mature seed colonized by *Burkholderia phytofirmans* strain PsJN::gusA; the blue colour is due to gusA-marked bacterial cells; strain PsJN is present inside the embryo (a,b) and in radicals (c); PsJN starts moving from embryo to germinated parts (c);

[0051] FIG. 3 shows *Burkholderia phytofirmans* strain PsJN recovery from the grain interior at different time intervals after harvesting (DAH; Days after harvesting);

[0052] FIG. 4 shows the effect of *Burkholderia phytofirmans* strain PsJN colonized/non-colonized seeds on germination and seedling growth of maize (a,b,c); and

[0053] FIG. 5 shows the effect of *Burkholderia phytofirmans* strain PsJN colonized/non-colonized seeds on shoot growth of maize (a,b,c; 30, 45, 60 days after sowing).

[0054] FIG. 6 shows representative results of GUS-staining in pepper treated with PsJN::gusA110 15 days p.i. GUS activity was found in all fruit parts including seeds.

[0055] FIG. 7 shows FISH analyses of pepper seeds colonized by *B. phytofirmans* PsJN using a general probe targeting eubacteria and a 23S rDNA probe specific for *B. phytofirmans*. Bacteria other than *B. phytofirmans* (eubmix-FITC) are indicated with a small arrow and *B. phytofirmans* PsJN is indicated with a longer arrow.

EXAMPLES

Example 1

Introducing *Burkholderia phytofirmans* Strain PsJN into Maize Seeds

[0056] The concept of internal seed colonization with plant growth promoting microorganisms according to the present invention was tested with the endophytic bacterium *Burkholderia phytofirmans* strain PsJN and two plant varieties of maize. Strain PsJN was applied by spraying female flowers with a suspension of 10^8 - 10^9 cfu mL⁻¹. At maturity, PsJN cells were detected within maize seeds at viable population densities that ranged from 10^2 - 10^5 CFU g⁻¹ fresh weight. Strain PsJN was not recovered from plants of the seed inoculation trial. After 12 months of storage 10^2 viable cells per g seeds were still recovered. Experiments were performed to determine the effects of internally colonized maize seeds on offspring plant biomass and vigor as compared to non-treated controls and external application of the same bacterial strain.

Experimental Description

[0057] The present invention provides seeds having beneficial microorganisms, especially bacteria, inside, enabling improved plant biomass equally over control as employing the same microorganisms (in the present case: bacteria) exogenously to seeds. A variant of the bacterium *Burkholderia phytofirmans* strain PsJN chromosomally tagged with the beta-glucuronidase gene (gusA, reporter gene for detection and monitoring of the strain by color formation) was used as a test strain in to maize cultivars (Peso and Morignon). For this, series of experiments were performed and the experimental setup was divided into two categories (1st and 2nd year experiments).

[0058] A) Evaluation of strain PsJN colonization potential in different tissues of maize plants (particularly grains).

[0059] B) Follow-up evaluation of strain PsJN colonized seed and strain PsJN inoculation (exogenously) to improve plant productivity over control.

Growth of PsJN Strain as Bacterial Inoculum

[0060] The bacterial strain was grown by loop-inoculating one single colony in LB broth amended with spectinomycin ($100 \mu\text{g mL}^{-1}$) in 100 mL flasks. The bacterial culture was incubated at $28 \pm 2^\circ \text{C}$. for 2 days at 180 rpm in a shaking incubator. The bacterial inoculum was applied in two different ways i.e. seed soaking and spraying inoculum at flowering stage. Maize seeds were surface sterilized by dipping for 5 and 3 min in 70% ethanol and NaOCl following 3 washings with sterilized water. There were three treatments, 1) seed inoculation 2) specific spraying of flowers and 3) seed inoculation combined with flower inoculation. Plants grown from

seeds treated with sterile culture broth only served as control. For inoculation, seeds of two maize cultivars were dipped for 3-4 hours in bacterial inoculum (10^8 - 10^9 cfu mL⁻¹). Likewise, bacterial inoculum was specifically sprayed to the female flower when the crop reached flowering stage. Seeds were sown in plastic trays filled with soil and 12 days old seedlings were transferred into 50 kg soil container (2 plants in each container) under wirehouse conditions.

Endophytic Colonization by PsJN Strain (Particularly Grain Colonization)

[0061] The rhizosphere and endophytic colonization of root, stem and leaves by the gusA-labeled variant of *B. phytofirmans* strains PsJN was determined by plate counting using LB plates amended with 5-Bromo-4-chloro-3-indolyl β -D-glucuronide (X-glcA, 50 μ g mL⁻¹), IPTG (50 μ g mL⁻¹) and the antibiotic spectinomycin (100 μ g mL⁻¹). Root, stem and leaf samples were washed, surface sterilized (as described above) and used for PsJN recovery (colonization). For this, samples were crushed in 0.9% saline buffer, subjected to oscillation in a pulsifier for 30 sec and dilution series were spread on agar plates. Beta-glucuronidase positive cells appear blue on media containing X-glcA. The blue colonies were counted after 3 days of incubation at 30° C. and the original cell number per g plant tissue was calculated. Similarly, PsJN colonization was also observed from different cob parts i.e. sheath, grains and cob interior. The identity of the blue colonies was further confirmed by RFLP analysis of the 16S23S rRNA intergenic spacer region.

Follow-Up Experiments were Performed in the 2nd Year to Evaluate the

[0062] 1. Viability, activation and colonization ability of strain PsJN colonizing maize seeds.

[0063] 2. Effect of strain PsJN colonized seed on germination and seedling vigor compared to untreated control (plastic tray assay).

[0064] 3. Effect of strain PsJN colonized seed on plant biomass compared to untreated control (pot trials).

[0065] Prior to the plant experiments, PsJN colonized seeds of both cultivars were tested to see whether PsJN cells are present and still alive inside. For this purpose, 20 seeds were imbibed in saline buffer for 2-3 days and subsequently crushed in 0.9% saline buffer, shaken for 45 second with a pulsifier and spread in dilutions on LB plates amended with X-glcA, IPTG and spectinomycin.

[0066] Bacterial inoculum was prepared as described above and three experiments were performed with four treatments i.e. control, seed inoculation with strain PsJN (exogenously), PsJN colonized seeds (produced in 1st year by spraying), PsJN colonized seed+inoculation.

[0067] For testing the germination performance, seeds (45) were surface sterilized and inoculated as described earlier, and were sown in plastic tray (diameter 30 cm) with three replicates. Data regarding time to start germination, mean germination time, time to 50% and final germination, germination index and energy, coefficient of uniform germination, and skewness were recorded of PsJN colonized over control.

[0068] Two pot experiments were performed to evaluate the performance of PsJN colonized seeds concerning plant biomass production as compared to control. Surface sterilized seeds were directly sown in pots with soil (first pot trial) or alternatively sown in plastic trays, and after 10 days seedlings were transferred to 5 kg pots (2nd pot trial). All plants were harvested after 60 days and data of plant height, number of

leaves per plant and root-shoot biomass were recorded. The data were subjected to analyses of variance using SPSS software package version 19 (SPSS Inc, Chicago, Ill.).

Results

Experiment A (1st Year): Seed Colonization by Strain PsJN

[0069] The ability of strain PsJN to colonize maize cobs (cob sheath, cob interior and grains) was analyzed in plants treated by specific flower inoculation (by spraying) only or by seed inoculation (FIG. 1). Only inoculation of flowers resulted in internal colonization of seeds. Internal seed colonization by strain PsJN was observed in both cultivars and both flower inoculation treatments. PsJN cells were detected in maize seeds at viable population densities that ranged from 10^2 - 10^5 CFU g⁻¹ fresh weight.

Experiment B1 (2nd Year): Viability, Activation and Colonization Ability of Strain PsJN Colonizing Maize Seeds.

[0070] PsJN colonized seeds, recovered from the first year experiment were tested to see whether PsJN cells survive inside of dormant seed and have the ability to colonize the plants emerging from the seeds what is very important as seeds may be stored for several months till planting. 10^2 viable cells were detected in two months old dormant seeds (FIG. 1). Imbibing in saline buffer for 2-3 days activated the 6 months old seeds and together with the seeds beginning to germinate PsJN started to proliferate resulting in a recovery of 10^4 viable cells (FIG. 4). Sprouts the emerged of 420 day old seeds were colonized by 10^5 PsJN cells and the bacteria was found all over inside the sprouts (FIGS. 1 and 2).

Experiment B2 (2nd Year): Effect of PsJN Colonized Seeds on Germination and Seedling Vigor as Compared to Untreated Control

[0071] The data summarized in table 1 and 4 revealed that PsJN colonized seeds showed significant improved germination ability. PsJN colonized seeds of both cultivars started to germinate 36-48 hours early than the control. PsJN colonized seed showed almost 100% final germination rate and required less mean germination time as compared to the control seeds. Consequently, the colonized seeds have better germination index as compared to control.

[0072] Moreover PsJN colonized seeds of both cultivars showed significantly higher maize seedling biomass as compared to untreated control seeds (Tables 2 and 5; FIGS. 3 and 4) but nonsignificantly higher seedling biomass as compared to seeds exogenously inoculated with PsJN.

Experiment B3 (2nd Year): Effect of PsJN Colonized Seed on Plant Biomass Compared to Untreated Control (Pot Trials)

[0073] The data of the pot trials (Table 3 and 6) revealed that PsJN colonized maize seeds had a positive effect on plant biomass production comparable to seeds externally coated with PsJN cells with cv Morignon being more responsive than cv Peso in both treatments (Tables 3 and 6; FIG. 5). The PsJN colonized seeds showed 38% increase in plant biomass production and a significant increase in root biomass as compared to the control. Moreover, the number of leaves per plant was higher in plants of PsJN colonized seed as compared to the control.

Conclusions

- [0074] *Burkholderia phytofirmans* PsJN can be introduced into maize seeds by spraying cells onto flowers.
- [0075] Seed inoculation only does not allow colonization of maize seeds of the next generation.
- [0076] Strain PsJN can survive inside maize seeds for at least 12 months
- [0077] Seed-colonizing bacterial cells are rapidly activated, proliferate and colonize emerging sprouts
- [0078] Seed-colonizing PsJN shows substantial plant growth promotion
- [0079] The present example therefore shows that the method according to the present invention enables an effective and reliable way to generate seeds with endophytes in a controlled and reproducible manner.

Example 2

Introducing *B. Phytofirmans* PsJN and *Enterobacter* sp. FD17 into Wheat and Barley Seeds

Experimental Description

[0080] Seeds of wheat (*Triticum* spp. cvs Collada and Monsun) and barley (*Hordeum vulgare* L. cvs Victoriana and Totum) were surface sterilized by dipping for 5 and 3 min in 70% ethanol and NaOCl following 3 washings with sterilized water. Seeds were sown in plastic trays and 12 days old seedlings were transferred into 20 kg soil containers and grown under green-house conditions. The soil has been collected from an agricultural field in Tulln, Lower Austria, and sieved to remove plant material. Bacterial strains (gusA-labelled variants of *B. phytofirmans* PsJN and *Enterobacter* sp. FD17) were grown by loop inoculation in LB broth amended with spectinomycin ($100 \mu\text{g mL}^{-1}$) in 100 mL Erlenmeyer flask. Bacterial cultures were incubated at $28 \pm 2^\circ \text{C}$. for 2 days at 180 rpm in a shaking incubator. Bacterial inoculum was applied by spraying exclusively flowers. Control plants were treated with sterilized broth.

Endophytic Colonization of Wheat and Barley Seeds

[0081] Plants were harvested at ripening stage and seeds were collected. Seed colonization by the inoculant strains was determined by GUS-staining. Therefore, seeds were cut in two pieces and incubated in GUS-staining solution (1 mM EDTA, 5 mM potassium ferricyanide, 5 mM potassium ferrocyanide, 100 mM sodium phosphate, pH 7.0, 1% Triton-X-100, 0.1 mg/mL X-Gluc predissolved in $5 \mu\text{L}/\text{mg}$ N,N-dimethylformamide, 0.1% IPTG) directly after harvesting at 37°C . for 20 hours. Afterwards, samples were rinsed with % ethanol. The ethanol was then discarded and samples were fixed in paraformaldehyde solution (4% paraformaldehyde dissolved in PBS at 60°C . with constant stirring until clarifying of the solution) overnight at 4°C . Finally, the fixed samples were rinsed 3 times in PBS and stored in the last rinse at 4°C . until further processing. In parallel, seeds were manually crushed under sterile conditions and used for bacterial community DNA isolation employing standard procedures. The presence of the inoculant strains was confirmed by sequence analysis of the 16S-23S rRNA intergenic spacer region (IGS) of single clones and subsequent comparison with those from the inoculants strains.

Results

Experiment A (1st Year):

[0082] Both seeds of wheat and barley were found to be internally colonized by the inoculants strains. Sequence analysis of the IGS region confirmed the presence of *Enterobacter* sp. FD17 and *B. phytofirmans* PsJN.

Conclusions

[0083] *Burkholderia phytofirmans* PsJN and *Enterobacter* sp. FD17 can be introduced into barley and wheat seeds by spraying cells onto flowers.

Example 3

Introducing *B. Phytofirmans* PsJN into Tomato and Pepper Seeds

Experimental Description

[0084] The colonization behavior of *Burkholderia phytofirmans* PsJN during transmission from flowers to seeds was studied with tomato (*Solanum lycopersicum* cv. Micro Tom and Matina) and pepper (*Capsicum annuum* cv. Feher). Presence of PsJN was investigated at 3 different time points. Detection of bacteria in the seed interior of harvested samples was conducted by GUS-staining and microscopy on the one hand and strain-specific quantitative PCR on the other hand. For detection by visual observation of staining and microscopy, the gusA-labelled variant of the strain PsJN, *Burkholderia phytofirmans* PsJN::gusA110, was used in parallel to the wild-strain that was detected via qPCR.

[0085] The ability of PsJN to survive in the seed and proliferate with the emerging seedling was studied in a subsequent germination experiment. Hereby, the harvested seeds from the previously treated plants were sown and nursed for a certain period. Afterwards the seedlings were examined regarding their presence of PsJN by GUS-staining and quantitative PCR of PsJN-specific genes.

[0086] The bacterial strains were grown by loop-inoculating one single colony in LB broth containing 0.1% of the antibiotic spectinomycin in case of *B. phytofirmans* PsJN::gusA110 and without antibiotics in case of the wild-type strain and incubated at 28°C . on a shaker (160 rpm) overnight. The overnight culture was transferred to 500 mL Erlenmeyer flasks containing 250 mL liquid LB medium. They were incubated on a shaker (120 rpm) at 28°C . for 2 days to allow for growth of bacteria. Subsequently, aliquots of 40 mL of the incubated medium containing the bacterial culture were filled in 50 mL plastic tubes and centrifuged at 4500 rpm and 4°C . for 10 minutes (Megafuge 40R, Heraeus, Hanau, Germany). Afterwards, the supernatant was discarded and the bacterial pellet re-suspended by vortexing in 20 mL PBS (0.2 g/L KCl, 1.44 g/L Na_2HPO_4 and 0.24 g/L KH_2PO_4 , in dH_2O , pH 7.4, autoclaved). The control suspension was treated accordingly. The aliquots of each bacterial suspension were then pooled in 500 mL Schott bottles. The concentration of the suspensions was measured by help of spectrophotometry (NanoDrop 1000 3.7.1., Wilmington, Del., USA) and adjusted to 10^8 CFU/mL .

[0087] Specific inoculation of tomato and pepper flowers was conducted when the plants reached growth stage 61-63 on the BBCH scale (for tomato: first inflorescence: first

flower open—third inflorescence: first flower open; for pepper: first flower open—third flower open) (FELLER et al., 1995b).

[0088] The bacterial inoculants and buffer only for the control were filled in a 50 mL glass pump spray bottle previously sterilized with 70% ethanol. The plants to be inoculated were spatially separated from the others to avoid contamination by drift. One single flower or 2 to 3 immediately adjacent flowers were sprayed with 675 µL of the inoculum. A filter paper was used to shield the surrounding plant parts such as leaves and stem from drift and take up surplus inoculum to avoid dripping on the soil. The treated inflorescences/flowers were marked with a twist tie to allow for later identification.

[0089] Six replicates of the inoculated plants were analyzed at 3 different developmental stages. Pepper samples were taken 3 days and 15 days after spraying as well as at full ripeness. The plant material (buds, flowers, fertilized flowers, developing fruits, immature fruits, ripe fruits and seeds) was cut with a sterile scalpel and subsequently incubated in GUS-staining solution (1 mM EDTA, 5 mM potassium ferricyanide, 5 mM potassium ferrocyanide, 100 mM sodium phosphate, pH 7.0, 1% Triton-X-100, 0.1 mg/mL X-Gluc predissolved in 5 µL/mg N,N-dimethylformamide, 0.1% IPTG) directly after harvesting at 37° C. for 20 hours. Afterwards, destaining was done by rinsing the samples with 70% ethanol. The ethanol was then discarded and the samples fixed in paraformaldehyde solution (4% paraformaldehyde dissolved in PBS at 60° C. with constant stirring until clarifying

of the solution) overnight at 4° C. Finally, the fixed samples were rinsed 3 times in PBS and stored in the last rinse at 4° C. until further processing.

[0090] Material of plants inoculated with PsJN wild-type and control samples were immediately after harvest frozen in liquid nitrogen and transferred for storage at –80° C. Afterwards, DNA was isolated using standard procedures and used as described above for Example 2.

Results

Experiment A (1st Year):

[0091] Upon flower spraying *B. phytofirmans* PsJN colonized seeds and pericarp of fruits of tomato and pepper (FIG. 6). The colonization process was monitored by GUS-staining and microscopy. The cell number of strain *B. phytofirmans* PsJN during transmission from flowers into seeds was tested by TaqMan-quantitative PCR using primers and probe targeting a gene encoding glutamine synthetase. The amount of *B. phytofirmans* PsJN cells applied on one flower was roughly 10⁸ and the cell number calculated per mg plant material during the process of colonization dropped from about 3000 cells per flowers to a few dozen cells in seeds. Results were confirmed by fluorescence in situ hybridization (FIG. 7).

Conclusions

[0092] *Burkholderia phytofirmans* PsJN can be introduced into tomato and pepper by spraying cells onto flowers.

TABLE 1

Comparative performance of PsJN colonized seed and PsJN inoculation (exogenously) on germination of maize cv Peso (Data are average of three replicate)								
Treatment	Time to Start Germination	Time to 50% Germination (T50)	Mean emergence Time (MET)	Final Germination % (FGP)	Germination Energy (GE)	Coefficient of uniform emergence (CUE)	Germination index (GI)	Skewness
Control [‡]	4a [†]	5.20b	6.74a	83.33bc	72.92ab	0.80NS	6.45bc	0.77bc
PsJN Inoculation [‡]	3.33ab	4.80c	6.55a	100a	85.42a	0.67	8.82a	0.73c
Control [§]	4a	5.60a	6.83a	77.08c	64.58b	0.85	5.45c	0.82a
PsJN Inoculation [§]	3.33ab	5.30ab	6.73a	89.58b	68.75ab	0.74	6.85b	0.78ab
PsJN colonized seed [‡]	2.33bc	4.33d	5.49b	100a	69ab	0.77	8.75a	0.79ab

[†]Values sharing similar letter(s) do not differ significantly at P < 0.05, according to Duncan's Multiple Range Test.
[‡]Seeds prepared by spraying PsJN inoculum (10⁸-10⁹ cfu mL⁻¹)
[‡]Parent seed used for first year experiment
[§]Offspring seed produced from first year experiment

TABLE 2

Comparative difference of PsJN inoculated and PsJN colonized seed on biomass of maize cv Peso in plastic tray experiment (data are average of three replicate).										
Treatment	Fresh Plant biomass (g)				Dry Plant biomass (g)				Plant height (cm)	No. of leaves per plant
	Stem	Leaves	Root	Total biomass	Stem	Leaves	Root	Total biomass		
Control	79.37 c [†]	95.70 b	37.20 b	212.27 c	3.63 c	9.65 b	1.39 b	14.67 c	93.37 b	6.58 c
PsJN Inoculation	93.77 b	111.03 a	38.4 ab	244.43 b	4.22 b	10.65 ab	1.73 a	16.90 b	95.87 a	7.04 b

TABLE 2-continued

Comparative difference of PsJN inoculated and PsJN colonized seed on biomass of maize cv Peso in plastic tray experiment (data are average of three replicate).										
Treatment	Fresh Plant biomass (g)				Dry Plant biomass (g)				Plant	No. of
	Stem	Leaves	Root	Total biomass	Stem	Leaves	Root	Total biomass	height (cm)	leaves per plant
PsJN colonized seed‡	99.70 b	113.33 a	39.63 a	251.43 ab	4.39 b	11.17 a	1.79 a	17.35 b	97.33 a	7.20 b

†Values sharing similar letter(s) do not differ significantly at P < 0.05, according to Duncan’s Multiple Range Test.

‡Seeds prepared by spraying PsJN inoculum (10⁸-10⁹ cfu mL⁻¹)

TABLE 3

Comparative performance of PsJN colonized seed and PsJN inoculation (exogenously) on plant biomass of maize cv Peso under potted conditions (data are average of three replicate).						
Treatment	Pot trial I (Direct sowing)				Pot trial II (Nursery sowing)	
	Plant	No. of	Shoot biomass	Root bio-mass	Shoot bio-mass	Root bio-mass
	height (cm)	leaves per plant				
Control	96.42 c†	6.98 c	5.32 c	0.82 c	1.29 c	0.28 c
PsJN Inoculation	108.01 ab	9.04 ab	8.80 ab	1.42 a	2.37 b	0.423 ab
PsJN colonized seed‡	104.62 b	8.42 b	7.17 b	1.12 b	2.16 b	0.358 b

†Values sharing similar letter(s) do not differ significantly at P < 0.05, according to Duncan’s Multiple Range Test.

‡Seeds prepared by spraying PsJN inoculum (10⁸-10⁹ cfu mL⁻¹)

TABLE 4

Comparative performance of PsJN colonized seed and PsJN inoculation (exogenously) on germination of maize cv Morignon (data are average of three replicate).								
Treatment	Time to Start Germination	Time to 50% Germination (T50)	Mean emergence Time (MET)	Final Germination % (FGP)	Germination Energy (GE)	Coefficient of uniform emergence (CUE)	Germination index (GI)	Skewness
Control‡	4.33a†	4.98a	6.72a	85.42bc	79.17ab	0.81NS	6.66b	0.74NS
PsJN Inoculation‡	3.67a-c	4.96a	6.65a	95.83ab	89.58a	0.78	8.25a	0.75
Control§	4ab	5.02a	6.65a	79.17c	75b	0.74	6.65b	0.76
PsJN Inoculation§	3.33bc	5.07a	6.59a	91.67ab	75b	0.65	7.88ab	0.77
PsJN colonized seed‡	3c	4.10b	5.69b	100a	83.33ab	0.69	9.06a	0.72

†Values sharing similar letter(s) do not differ significantly at P < 0.05, according to Duncan’s Multiple Range Test.

‡Seeds prepared by spraying PsJN inoculum (10⁸-10⁹ cfu mL⁻¹)

§Parent seed used for first year experiment

§Offspring seed produced from first year experiment

TABLE 5

Comparative performance of PsJN colonized seed and PsJN inoculation (exogenously) on seedling biomass of maize cv Morignon in plastic tray experiment (data are average of three replicate).										
Treatment	Fresh Plant biomass (g)				Dry Plant biomass (g)				Plant	No. of leaves
	Stem	Leaves	Root	Total biomass	Stem	Leaves	Root	Total biomass	height (cm)	per plant
Control	81.07 c†	97.70 b	38.43 b	215.93 c	3.83 c	9.67 c	1.76 b	15.26 c	94.76NS	6.53 c
PsJN Inoculation	92.67 b	104.80 a	42.40 a	239.23 b	4.64 b	10.57 b	2.34 a	17.67 b	95.00	6.87 b
PsJN colonized seed‡	92.90 b	105.07 a	41.93 a	240.13 b	4.66 b	11.25 ab	2.35 a	18.24 ab	95.02	6.84 b

†Values sharing similar letter(s) do not differ significantly at $P < 0.05$, according to Duncan's Multiple Range Test.

‡Seeds prepared by spraying PsJN inoculum (10^8 - 10^9 cfu mL⁻¹)

TABLE 6

Comparative performance of PsJN colonized seed vs PsJN inoculation (exogenously) on plant biomass of maize cv Morignon under potted conditions (data are average of three replicate).						
Treatment	Pot trial I (Direct sowing)				Pot trial II (Nursery sowing)	
	Plant	No. of	Shoot	Root bio-	Shoot bio-	Root bio-
	height (cm)	leaves per plant	biomass	mass	mass	mass
Control	101.42 c†	7.98 c	6.36 c	1.12 c	3.29 c	0.41 c
PsJN Inoculation	110.67 b	9.47 b	8.17 b	1.42 b	4.37 b	0.623 ab
PsJN colonized seed‡	113.01 ab	9.83 b	8.80 b	1.56 ab	4.26 b	0.558 b

†Values sharing similar letter(s) do not differ significantly at $P < 0.05$, according to Duncan's Multiple Range Test.

‡Seeds prepared by spraying PsJN inoculum (10^8 - 10^9 cfu mL⁻¹)

1-27. (canceled)

28. A plant seed population produced by the process of i) providing an agriculturally-acceptable preparation comprising a population of isolated bacterial endophytes; ii) specifically contacting at least one flower of a flowering plant with the preparation under conditions such that the bacterial endophytes are localized to a plurality of seeds derived from the flowering plant, wherein the plurality of seeds individually contain at least about 100 bacterial endophytes per seed; and iii) collecting the plurality of derived seeds, thereby producing a plant seed population.

29. The seed population of claim 28, wherein the isolated bacterial endophytes are heterologous to the flowering plant.

30. The seed population of claim 28, wherein the isolated bacterial endophytes are localized to the seed interior.

31. The seed population of claim 28, wherein the derived seeds have one or more improved traits as compared to an untreated control seed, wherein the one or more improved traits are individually selected from the group consisting of increased plant growth, increased yield, increased biomass, improved nutrient acquisition, increased resistance to environmental stresses, increased tolerance to drought, increased tolerance to heavy metals, increased tolerance to disease, and increased tolerance to herbivory, increased production of phytohormones, increased production of antibiotics, increased production of siderophores, increased production of pesticides, and increased promotion of nitrogen fixation.

32. The population of claim 28, wherein the step of contacting the at least one flower is performed in a non-sterile environment.

33. The seed population of claim 28, wherein the seeds are collected in a manner suitable for storage for at least about 1 month.

34. A field of plants at least 1 acre in size derived from the seed population of claim 28.

35. A method for producing a synthetic combination comprising a plant seed and endophytic microorganisms, comprising contacting at least one flower of a flowering plant with a preparation comprising a purified population of microorganisms, the population comprising an endophytic microorganism, wherein the endophytic microorganisms are present in the interior of the plant seed.

36. The method of claim 35, further comprising obtaining the seed produced by the flowering plant.

37. The method of claim 35, wherein contacting the flower of a plant with a preparation of endophytic microorganisms is performed by specifically spraying the population at the time of flowering.

38. The method of claim 36, wherein the seed has an improved trait as compared to an untreated control seed, wherein the improved trait is selected from the group consisting of increased germination rate, increased germination index, increased germination energy, and more uniform germination as compared to an untreated control seed.

39. The method of claim **36**, further comprising storing the seed produced by the plant for at least 1 month.

40. The method of claim **39**, wherein the seed is stored for at least 6 months.

41. The method of claim **36**, further comprising growing the plant seed to a progeny plant, wherein the progeny plant has an improved trait as compared to an untreated control plant.

42. The method of claim **41**, wherein the improved trait is selected from the group consisting of increased plant growth, increased yield, increased biomass, improved nutrient acquisition, increased resistance to environmental stresses, increased tolerance to drought, increased tolerance to heavy metals, increased tolerance to disease, and increased tolerance to herbivory, increased production of phytohormones, increased production of antibiotics, increased production of siderophores, increased production of pesticides, and increased promotion of nitrogen fixation as compared to an untreated control plant.

43. The method of claim **35**, wherein the at least one flower of the flowering plant is contacted with a preparation of microorganisms in a suspension of at least 10^6 cfu/mL.

44. A method of improving a trait in a seed or plant produced by the seed comprising:

contacting at least one flower of a flowering plant with a preparation comprising a purified population of microorganisms, the population comprising an endophytic microorganism, and

obtaining the seed produced by the plant.

45. The method of claim **44**, wherein the improved trait is selected from the group consisting of increased germination rate, increased germination index, increased germination energy, more uniform germination increased plant growth, increased yield, increased biomass, improved nutrient acquisition, increased resistance to environmental stresses, increased tolerance to drought, increased tolerance to heavy metals, increased tolerance to disease, and increased tolerance to herbivory, increased production of phytohormones, increased production of antibiotics, increased production of siderophores, increased production of pesticides, and increased promotion of nitrogen fixation as compared to untreated control seed or plant.

46. The method of claim **44**, wherein the endophytic microorganisms are present in the interior of the plant seed.

47. The synthetic combination produced by the method of claim **35**.

* * * * *