



US 20160160245A1

(19) **United States**

(12) **Patent Application Publication**
Yocum et al.

(10) **Pub. No.: US 2016/0160245 A1**

(43) **Pub. Date: Jun. 9, 2016**

(54) **METHOD OF PRODUCING SUCCINIC ACID AND OTHER CHEMICALS USING FACILITATED DIFFUSION FOR SUGAR IMPORT**

(71) Applicant: **MYRIANT CORPORATION**, Woburn, MA (US)

(72) Inventors: **R. Rogers Yocum**, Lexington, MA (US); **Andrew Christopher Collard**, New Haven, CT (US); **Theron Hermann**, Arlington, MA (US); **Xiaohui Yu**, Woburn, MA (US); **Wei Gong**, Woburn, MA (US)

(73) Assignee: **MYRIANT CORPORATION**, Woburn, MA (US)

(21) Appl. No.: **14/906,501**

(22) PCT Filed: **Jul. 22, 2014**

(86) PCT No.: **PCT/US2014/047696**

§ 371 (c)(1),

(2) Date: **Jan. 20, 2016**

Related U.S. Application Data

(60) Provisional application No. 61/857,300, filed on Jul. 23, 2013.

Publication Classification

(51) **Int. Cl.**

C12P 7/46 (2006.01)

C07K 14/195 (2006.01)

(52) **U.S. Cl.**

CPC **C12P 7/46** (2013.01); **C07K 14/195** (2013.01)

(57) **ABSTRACT**

This invention relates to the production of succinic acid and other chemicals derived from phosphoenolpyruvate (PEP) by fermentation with a microorganism in which the fermentation medium contains one or more sugars, and in which one or more of the sugars is imported into the cell by facilitated diffusion. As a specific example, succinic acid is produced from a glucose-containing renewable feedstock through fermentation using a biocatalyst. Examples of such a biocatalyst include microorganisms that have been enhanced in their ability to utilize glucose as a carbon and energy source. The biocatalysts of the present invention are derived from the genetic manipulation of parental strains that were originally constructed with the goal to produce one or more chemicals (for example succinic acid and/or a salt of succinic acid) at a commercial scale using feedstocks that include, for example, glucose, fructose, or sucrose. The genetic manipulations of the present invention involve the introduction of exogenous genes involved in the transport and metabolism of glucose or fructose into the parental strains. The genes involved in the transport and metabolism of glucose or fructose can also be introduced into a microorganism prior to developing the organism to produce a particular chemical. The genes involved in the transport and metabolism of sucrose can also be used to augment or improve the efficiency of sugar transport and metabolism by strains already known to have some ability for glucose utilization in biological fermentations.

Figure 1




Figure 2




Figure 3





Figure 4



**METHOD OF PRODUCING SUCCINIC ACID
AND OTHER CHEMICALS USING
FACILITATED DIFFUSION FOR SUGAR
IMPORT**

CROSS-REFERENCE TO RELATED
APPLICATION

[0001] This application claims the priority of the U.S. Provisional Application Ser. No. 61/857,300, filed on Jul. 23, 2013.

FIELD OF THE INVENTION

[0002] The present invention is in the field of producing specialty and commodity organic chemicals using biocatalysts (bacteria and other microorganisms) that can be modified to increase their efficiency in using sugar-containing feedstocks. More specifically, the present invention is related to the genetic modifications of genes that encode functions involving transport and metabolism of sugars for the biological production of succinic acid and other chemicals.

BACKGROUND OF THE INVENTION

[0003] A large number of organic chemicals are currently derived from petrochemical feedstocks. There is a growing interest in producing many of these petrochemical-derived organic compounds through biological fermentation processes using renewable feedstocks. The list of organic compounds that can be derived from renewable feedstocks includes α,ω -diacids (succinic, fumaric, malic, glucaric, malonic, and maleic), 2,5-furan dicarboxylic acid, propionic acid, 3-hydroxy propionic acid, aspartic acid, glucaric acid, glutamic acid, itaconic acid, levulinic acid, 3-hydroxybutyrolactone, glycerol, and butanediols such as 1,4 butanediol (US Patent Application 20090047719), 1,3-butanediol (US Patent Application 20090253192), and 2,3-butanediol. Many other types of organic compounds, including, but not limited to, amino acids, vitamins, alcohols (such as ethanol, n-propanol, isopropanol, n-butanol, isobutanol, and higher alcohols), fatty acids, esters of fatty acids, hydrocarbons, isoprenoids, turpenes, carotenoids, amines, can also be produced using renewable feedstocks. Any such compound shall be referred to herein as a “desired compound”. Although fermentation processes for many of these desired compounds have been developed, in order to compete with petrochemical processes, there is a constant need to improve the overall economics of fermentation, for example to improve product titer (final concentration in grams per liter of product) and product yield (grams of product per gram of carbon source such as glucose), and to reduce the titer of unwanted byproducts, such as acetate.

[0004] Many bacteria, including *Escherichia coli*, use a system for actively transporting glucose and other sugars into the cell called a phosphotransferase system (PTS). This system uses PEP (phosphoenol pyruvate) as the source of energy and phosphate for simultaneously transporting and phosphorylating the sugar. PTS systems usually require four or more proteins that together function to import and phosphorylate the incoming sugar. Some of these proteins are common to all of the sugars that a given organism imports by a PTS, while other protein components of the PTS are specific for one or more particular sugars.

[0005] For example, in *E. coli*, the proteins that are common to all PTS pathways are PtsH and PtsI, encoded by the

genes ptsH and ptsI, respectively. In addition to these two “common” PTS proteins, one or more additional sugar-specific PTS proteins are required to import and phosphorylate particular sugars. For example, import of glucose by the PTS requires two additional proteins named Crr and PtsG. Crr is a cytoplasmic protein with a single domain called A, and PtsG is a membrane protein with two domains named B and C. The phosphate group from PEP is relayed from protein to protein and is finally transferred to glucose as it is imported, at the 6 position to give glucose-6-phosphate inside the cell. The order of the relay starting with PEP is PtsI, PtsH, Crr, and finally PtsG. Historically, these proteins have also been called by other names, such as EI, HPr, EIIA^{Glc}, and EIIBC, respectively. As another example from *E. coli*, fructose is imported by a similar relay using PtsI, PtsH, FruA, and FruB, the last two of which are also known as EII^{Fru} and EII^{Fru}, respectively. For some sugars, for example mannitol, the sugar-specific protein domains corresponding to A, B, and C as mentioned above for glucose are fused into one membrane bound polypeptide, while for other sugars, for example mannose, the A and B domains are fused into one cytoplasmic polypeptide, while the membrane bound component is comprised of two subunits called C and D.

[0006] In all cases, the system relies on the “common subunits” (PtsI and PtsH in *E. coli*), and PEP is the source of energy and phosphate. As a result, every molecule of sugar imported by a PTS system results in the utilization of one molecule of PEP and the production of one molecule of phosphorylated sugar and one molecule of pyruvate. However, PEP is also an obligate intermediate in several biochemical pathways, such as 1) formation of pyruvate and ATP by pyruvate kinase, 2) the anapleurotic pathways catalyzed by PEP carboxykinase and PEP carboxylase, which both feed carbon into the TCA (tricarboxylic acid) cycle, and 3) the entry into the common aromatic amino acid and aromatic vitamin biosynthetic pathway catalyzed by one or more isozymes of 3-deoxy-D-arabinoheptulosonate-7-phosphate synthase (DAHP synthase). Thus there is an inevitable competition for PEP between the PTS system for sugar import and the other pathways just mentioned.

[0007] Since many bacteria, including both Gram positives and Gram negatives, use a PTS system, it is obviously a system that has prevailed under many circumstances throughout evolution. However, under anaerobic conditions, production of ATP from sugars such as glucose is much less efficient than under aerobic conditions, and the so-called “substrate level” phosphorylation, for example, by pyruvate kinase, becomes a larger portion of the ATP production budget than under aerobic conditions where oxidative phosphorylation provides the majority of the ATP budget. As such, it is noteworthy that some organisms, such as *Saccharomyces cerevisiae* and *Zymomonas mobilis*, both of which are well adapted to anaerobic growth on glucose and other sugars, do not have a PTS system, but instead use a facilitated diffusion protein (also called a permease) to import glucose and other sugars. Furthermore, when organisms that natively use a PTS are genetically engineered to overproduce particular compounds by fermentation, the pathways in many cases use PEP as an intermediate, so that the PTS competes with the desired biosynthetic pathway for PEP. Alleviation from this competition by reducing the activity of the PTS is known to increase flux to a desired biosynthetic pathway.

[0008] For example, PEP is an intermediate in the reductive branch of the tricarboxylic acid (TCA) cycle that leads to

succinate. During the metabolic evolution of KJ122, an *E. coli* succinate producer, a frameshift mutation arose in the *ptsI* gene, which resulted in an increase in succinate production from glucose. Reinstalling a wild type *ptsI* gene caused a drastic reduction in succinate production, proving that the *ptsI* mutation contributed strongly to the strain improvement.

[0009] For another example, aromatic amino acids are built from PEP and erythrose-4-phosphate. Deletion of three *pts* genes (Δ *ptsHI*, *crr*) in an *E. coli* strain was shown to increase flux to the aromatic amino acid biosynthetic pathway when cells are grown on glucose as the carbon source.

[0010] In both of the above examples, import of glucose is presumably still accomplished at some level by the so-called galactose permease (GalP, encoded by the *galP* gene). In the first example, a mutation that reduced the activity of a repressor (GalS) of the *galP* gene was found to result from metabolic evolution (WO2011/123154). In the second example, one or more mutations occurred after deletion of *pts* genes that resulted in an increase in growth rate. The resulting strain depended on GalP for significant growth on glucose, and one or more mutations in the strain could have been related to an increase in expression of *galP* (U.S. Pat. No. 6,962,794). However, the strains from this second example produced only low titers of aromatic amino acids after engineering the “Pts-/*Glu+*” strains for aromatic amino acid production. Phenylalanine, tyrosine, and tryptophan were produced at 1.7, 0.8, and 2.2 g/l respectively. Since these titers are nowhere near high enough to support an economically attractive commercial process, it is not clear that the invention disclosed in U.S. Pat. No. 6,962,794 is useful for commercial production. As such, there is still a need to improve fermentation parameters for economically viable commercial production of chemicals by fermentation.

[0011] Although the use of GalP for glucose import conserves PEP, it is a proton symporter, so it consumes about $\frac{1}{3}$ of an ATP for each glucose molecule transported. Some microorganisms, for example the bacterium *Zymomonas mobilis* and the yeast *Saccharomyces cerevisiae* use facilitated diffusion for importing glucose. *Z. mobilis* has one facilitator protein that functions to import both glucose and fructose. *S. cerevisiae* has at least 14 different hexose importers, many of which import glucose and at least some of which import fructose as well. This mode for glucose import requires no ATP expenditure until the sugar is inside the cytoplasm, after which an ATP is consumed to form glucose-6-phosphate to allow the sugar to enter glycolysis. Most importantly, unlike for the PTS system, no PEP is consumed. As such, facilitated diffusion clearly works well for some organisms, and costs the cell less in terms of PEP and ATP than either a PTS system or a proton symporter such as GalP. Ingram et al. (U.S. Pat. No. 5,602,030) demonstrated that the facilitated diffusion protein (Glf, encoded by the *glf* gene) from *Zymomonas mobilis*, together with a glucokinase (Glc, encoded by the *glk* gene), also from *Zymomonas mobilis*, expressed from those genes on a multicopy plasmid, could functionally replace the PTS to support growth in a minimal glucose medium of an *E. coli* strain, where the parent had no native glucose facilitated diffusion capability, and other glucose import systems had been disabled by mutation. The recombinant *E. coli* *ptsG*⁻, *ptsM*⁻, *glk*⁻ strain ZSC113 containing the two *Z. mobilis* genes *glf* and *glk* on a plasmid could grow aerobically on minimal glucose medium.

[0012] These disclosures proved that the *Z. mobilis* proteins could function in *E. coli* enough to support growth aerobi-

cally with a specific growth rate of 0.53 hr⁻¹. However, wild type *E. coli* using the native PTS for glucose import has an aerobic specific growth rate of 1.0 to 1.2 hr⁻¹, so the strains engineered in U.S. Pat. No. 5,602,030 to use *glf* appear to be severely limited by glucose uptake. Moreover, the disclosures did not show that the facilitated diffusion system could support anaerobic growth. A number of important chemicals produced by fermentation require robust anaerobic growth to support an economically attractive commercial production system (WO2012/018699). The examples in U.S. Pat. No. 5,602,030 and Snoep et al (1994) showed that modest growth could be obtained by expressing *glf* and *glk* from a multicopy plasmid, but it was not demonstrated that growth could be supported by integrated copies of the *glf* and *glk* genes, yet it is often desirable for commercial scale production to use strains that do not contain a plasmid. Finally, U.S. Pat. No. 5,602,030 did not demonstrate that a *glf*-based system could support high titer production of a commodity chemical such as ethanol or succinate in *E. coli* or any other organism that does not natively use facilitated diffusion. As such, it was not clear from the disclosure of U.S. Pat. No. 5,602,030 alone that a *glf* could replace the PTS and result in an economically attractive fermentation processes for producing a desired chemical in a host strain that does not have a native facilitated diffusion system.

[0013] Tang et al (2013) went a couple steps further to show that anaerobic production of succinate could be achieved by expression of *Z. mobilis* *glf* in combination with a glucokinase in an *E. coli* strain background that was Δ *ptsI*, Δ *aldhA*, Δ *pflB*, *pck*^{*}. However, the best succinate production in this system was modest, only 220 mM (26 g/l) in 96 hours. Despite having optimized by combinatorial modulation the expression of *glf* and *glk*, this titer and productivity is nowhere near that of previously published strains that produced 83 g/l succinate without the use of *glf*. Thus, despite the more advanced work of Tang et al., it had still not been demonstrated that the use of facilitated diffusion for glucose import was useful for actually improving fermentation production parameters at levels that would be necessary for economically attractive commercial production, which would be at the benchmark of at least 83 g/l (WO2012/018699). To further complicate the potential replacement of a PTS by *glf*, in *E. coli*, and presumably in other bacteria, the components of the PTS have many diverse regulatory functions that affect many different metabolic pathways, so it is impossible to predict what the effects will be of a deletion in any one or more of the PTS genes on the overall physiology and fermentative properties of any resulting modified strain. Native *Z. mobilis* strains, which naturally use facilitated diffusion for glucose uptake, are capable of producing up to about 60 g/l ethanol and a similar quantity of carbon dioxide from glucose. An engineered strain of *Z. mobilis* is reported to produce 64 g/l succinate from glucose (EP20070715351). However, this fermentation required 10 of yeast extract in the fermentation medium, which is undesirable for commercial production of succinic acid, both because of its expense and the increased cost required for downstream purification of the succinate from the yeast extract components. Furthermore, *Z. mobilis* is often not a convenient or optimal host organism for use in fermentative processes.

[0014] Thus, to summarize the prior art, it had been shown that *E. coli* can be engineered to use facilitated diffusion of glucose to support aerobic growth to a modest rate, and to support a modest level of succinate production anaerobically,

but there has been no disclosure of any bacterial strain or process that has been engineered to confer the non-native use of facilitated diffusion for glucose import and that is improved over strains using native glucose import systems such as PTS and/or GalP for production of a chemical by fermentation. Furthermore, there has been no disclosure of any bacterial strain or process that uses facilitated diffusion for glucose import and that is capable of producing succinate or any chemical other than ethanol and carbon dioxide at a titer, yield, and rate that is high enough in a medium that would be commercially attractive, such as a minimal glucose medium. As such, there is still a need for improved strains that can produce succinate and chemicals in a process that is economically attractive when all factors including productivity, cost of the medium, and downstream purification are taken into account.

SUMMARY OF THE INVENTION

[0015] This present invention provides biocatalysts (for example genetically engineered microorganisms) and methods for using facilitated diffusion of glucose for improving the fermentative production of commercially important products, for example, but not limited to, specialty and commodity chemicals. Specifically, the present invention is useful in the fermentative production of organic acids, amino acids, and other biochemicals that have PEP as a biochemical intermediate in their biosynthetic pathway, using sugar-containing renewable feedstocks. As a specific example, the present invention is useful in the fermentative production of succinic acid from a glucose, fructose, or sucrose-containing renewable feedstock using biocatalysts that have been constructed to use facilitated diffusion of a sugar. The principles of the present invention can be applied to many other desired chemical compounds that can be produced by fermentation, particularly chemical intermediates of the TCA cycle or derivatives thereof, such as fumaric acid, malic acid, glutamate, derivatives of glutamate, aspartate, derivatives of aspartate, aromatic amino acids (phenylalanine, tyrosine, tryptophan), and compounds derived from intermediates in the central aromatic pathway, such as vitamins and cis, cis-muconic acid.

[0016] According to the present invention, genes coding for the proteins involved in facilitated diffusion of sugars such as glucose can be introduced into a wide variety of biocatalysts either to confer a new ability to the biocatalyst to import a sugar as a source of carbon and energy from the fermentation medium by facilitated diffusion, or to augment or improve an already existing capacity of the biocatalysts for sugar transport and metabolism. Strains engineered to have the added ability to import sugars by facilitated diffusion can have improved fermentation parameters when compared to parameters of the parent strain, such as increased titer (g/l of desired chemical product), increased yield (grams of product produced per gram of sugar consumed), increased specific productivity (g/l-hr of product formation), and/or decreased titer of unwanted byproducts such as acetate, pyruvate and/or amino acids. These improved parameters can result from conservation of energy (for example use of less ATP for formation of proton gradients to drive proton symporters such as GalP), conservation of PEP for pathways that use PEP as an intermediate, such as the succinate pathway(s), and decreasing of overflow metabolism into acetate production pathways or other unwanted pathways.

[0017] This approach is particularly advantageous for production of chemicals that are derived at least in part from or

through PEP, such as succinate, malate, fumarate, lactate, ethanol, butanols, propane diols, 3-hydroxypropionic acid, acrylic acid, propionic acid, lactic acid, amino acids such as glutamate, aspartate, methionine, lysine, threonine, and isoleucine, compounds derived from the central aromatic pathway such as phenylalanine, tyrosine, tryptophan, aromatic vitamins, aromatic vitamin-like compounds, and any other compound that is derived from PEP as a biosynthetic intermediate.

[0018] In one embodiment, the present invention provides biocatalysts that do not natively have the ability to import a sugar by facilitated diffusion with an added heterologous gene (or genes) that confers a new ability to import a sugar by facilitated diffusion. In another embodiment, the present invention provides novel biocatalysts that produce a higher titer of a desired fermentation product than the parent biocatalyst. In another embodiment, the present invention provides novel biocatalysts that produce a higher yield of a desired fermentation product than the parent biocatalyst. In another embodiment, the present invention provides novel biocatalysts that produce a higher specific productivity for a desired fermentation product than the parent biocatalyst. In another embodiment, the present invention provides a novel biocatalyst that produces a lower titer of an undesired byproduct than the parent biocatalyst.

[0019] The gene or genes that code for the protein or proteins that function in the facilitated diffusion of a sugar can be derived from any organism that has the native ability to carry out facilitated diffusion of a sugar, the only requirement being that the protein or proteins are able to function in the new host. The gene encoding a sugar kinase, for example a glucokinase, that is required to phosphorylate the sugar after it enters the cytoplasm can be derived from the same donor from which came the gene(s) for facilitated diffusion, or a native sugar kinase gene from the recipient host can be used, or a combination of both sugar kinases can be used.

[0020] In another embodiment, the present invention provides for methods for producing a desired fermentation product comprising cultivating a genetically engineered microorganism that used facilitated diffusion to import a sugar.

[0021] In another embodiment, the present invention provides for methods for improving fermentation performance parameters (titer, yield, specific productivity, minimizing byproduct formation) of strains engineered to use facilitated diffusion.

[0022] In another embodiment, the present invention provides for methods for achieving an improved balance of facilitated diffusion and sugar kinase activity leading to improved growth and fermentation parameters in genetically engineered microorganism that used facilitated diffusion to import a sugar.

[0023] According to the present invention, one approach is to genetically transfer a facilitated diffusion system for importing a sugar from a second donor organism that naturally contains the relevant genes (for example glf or glk or a combination thereof) into a first recipient organism that does not naturally contain said relevant genes, so as to confer on said first recipient organism a new ability to import said sugar by facilitated diffusion. In a preferred embodiment, the first recipient has been previously engineered or constructed to be devoid of, or substantially reduced in, its ability to import said sugar by any native system or systems present in a parent or

ancestor of said first recipient strain. In such an embodiment, the resulting strain is in effect forced to use facilitated diffusion for growth on said sugar.

[0024] In a preferred embodiment, the first recipient strain is an *E. coli* strain, and the second donor strain is *Zymomonas mobilis* CP4. In a more preferred embodiment, said first strain is WG53, which in turn is derived from KJ122 by deletion of ptsH, ptsI, and galP. The exact nature of the deletions of ptsH, ptsI, and galP can vary widely, the only important criterion being that the activities of the PtsH, PtsI, and GalP proteins are eliminated or substantially reduced.

[0025] The first recipient organism of the invention can vary widely, the only criterion being that it does not natively contain a protein that functions in facilitated diffusion for a sugar such as glucose. In addition to *E. coli*, examples of first recipient organisms include, but are not limited to: *Gluconobacter oxydans*, *Gluconobacter asaii*, *Achromobacter delmarvae*, *Achromobacter viscosus*, *Achromobacter lacticum*, *Agrobacterium tumefaciens*, *Agrobacterium radiobacter*, *Alcaligenes faecalis*, *Arthrobacter citreus*, *Arthrobacter tumescens*, *Arthrobacter paraffineus*, *Arthrobacter hydrocarboglutamicus*, *Arthrobacter oxydans*, *Aureobacterium saperae*, *Azotobacter indicus*, *Brevibacterium ammoniagenes*, *divaricatum*, *Brevibacterium lactofermentum*, *Brevibacterium flavum*, *Brevibacterium globosum*, *Brevibacterium fuscum*, *Brevibacterium ketoglutamicum*, *Brevibacterium helcolum*, *Brevibacterium pusillum*, *Brevibacterium testaceum*, *Brevibacterium roseum*, *Brevibacterium immariophilum*, *Brevibacterium linens*, *Brevibacterium protopharmiae*, *Corynebacterium acetophilum*, *Corynebacterium glutamicum*, *Corynebacterium callunae*, *Corynebacterium acetoacidophilum*, *Corynebacterium acetoglutamicum*, *Enterobacter aerogenes*, *Erwinia amylovora*, *Erwinia carotovora*, *Erwinia herbicola*, *Erwinia chrysanthemi*, *Flavobacterium peregrinum*, *Flavobacterium fucatum*, *Flavobacterium aurantium*, *Flavobacterium rhenanum*, *Flavobacterium sewanense*, *Flavobacterium breve*, *Flavobacterium meningosepticum*, *Micrococcus* sp. CCM825, *Morganella morganii*, *Nocardia opaca*, *Nocardia rugosa*, *Planococcus eucinatus*, *Proteus rettgeri*, *Propionibacterium shermanii*, *Pseudomonas synxantha*, *Pseudomonas azotoformans*, *Pseudomonas fluorescens*, *Pseudomonas ovalis*, *Pseudomonas stutzeri*, *Pseudomonas acidovolans*, *Pseudomonas mucidolens*, *Pseudomonas testosteroni*, *Pseudomonas aeruginosa*, *Rhodococcus erythropolis*, *Rhodococcus rhodochrous*, *Rhodococcus* sp. ATCC 15592, *Rhodococcus* sp. ATCC 19070, *Sporosarcina ureae*, *Staphylococcus aureus*, *Vibrio metschnikovii*, *Vibrio tyrogenes*, *Actinomyces madurae*, *Actinomyces violaceochromogenes*, *Kitasatosporia parulosa*, *Streptomyces coelicolor*, *Streptomyces flavelus*, *Streptomyces griseolus*, *Streptomyces lividans*, *Streptomyces olivaceus*, *Streptomyces tanashiensis*, *Streptomyces virginiae*, *Streptomyces antibioticus*, *Streptomyces cacaoi*, *Streptomyces lavendulae*, *Streptomyces viridochromogenes*, *Aeromonas salmonicida*, *Bacillus pumilus*, *Bacillus circulans*, *Bacillus thiaminolyticus*, *Escherichia freundii*, *Microbacterium ammoniophilum*, *Serratia marcescens*, *Salmonella typhimurium*, *Salmonella schottmulleri*, *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus amyloliquefaciens*, *Klebsiella oxytoca*, *Klebsiella pneumoniae*, *Acinetobacter baylyi*, *Corynebacterium glutamicum*, *Brevibacterium flavum*, *Mannheimia succiniproducens* and *Anaerobiospirillum succiniproducens*, and *Xanthomonas citri*.

[0026] Examples of second donor organisms are any strain or species that has a native facilitated diffusion system for a sugar, for example *Zymomonas mobilis* strains (in addition to strain CP4), *Homo sapiens*, *Azospirillum amazonense*, *Flavobacteriaceae* bacterium S85, *Saccharomyces cerevisiae* or other yeast genera.

[0027] In another embodiment, a first parent strain is first constructed to use facilitated diffusion for importing a sugar, and then the resulting strain is further engineered to overproduce a chemical of commercial interest such as succinic acid.

[0028] Novel aspects of this invention are that the glf gene from a non-pathogenic, robust sugar utilizer has been stably integrated into the chromosome of a bacterium, such that the newly constructed bacterium can produce a commercially viable product with an economically viable process. The titer, yield and/or specific productivity of product from glucose or another sugar is greater than those parameters of the parent organism. The glf gene is integrated at a site in the chromosome that does not interfere with any relevant aspect of growth or product production. The acetate titer is less than that of the parent strain at about 45 to 48 hours in a representative fermentation, allowing a 2 day fermentation cycle time, unlike a prior art example. Strains in the prior art that used facilitated diffusion for sugar import did not produce sufficient titers of the desired product to be economically attractive. Another novel aspect of this invention is that by using facilitated diffusion for sugar import, it was unexpectedly found that the production of the unwanted byproduct acetate or acetic acid was significantly reduced. The prior art strain KJ122 produces about 5 to 7 g/l acetate in a typical fed glucose fermentation (WO2012/018699), while new strains of the invention produce only about 4.2 g/l or less.

[0029] Additional advantages of this invention will become readily apparent from the ensuing description.

BRIEF DESCRIPTION OF THE DRAWINGS

[0030] FIG. 1 Structure of plasmid pAC19, a source of an expression cassette for *Z. mobilis* glf and glk.

[0031] FIG. 2 Structure of plasmid pAC21, a source of a selectable and counter-selectable cassette containing cat (chloramphenicol resistance) and sacB (levan sucrase) genes.

[0032] FIG. 3 Structure of plasmid pSS2, a source of an expression cassette for *Z. mobilis* glf without glk.

[0033] FIG. 4 Structure of plasmid pMH68, a source of an expression cassette for integration of a second copy of the *E. coli* crr gene at the pflD locus.

[0034] Table 1. Production of succinate by AC15 in 7 liter fermentors

[0035] Table 2. Production of succinate by red mutants of AC15 in 500 ml microaerobic fermentors.

[0036] Table 3. Production of succinate by two isolates of SS8 in 500 ml microaerobic fermentors.

[0037] Table 4. Production of succinate by YSS41 in 20 liter microaerobic fermentors.

[0038] Table 5. Production of succinate by MH141 in 500 ml microaerobic fermentors.

[0039] Table 6. Succinate production by *E. coli* strains KJ122 and YSS41 in 20 liter fermentors under optimized aeration conditions for both strains.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Definitions

[0040] When the phrase “for example” or “such as” is used, the subsequently mentioned items are meant to be illustrative examples for the idea or concept being disclosed. The subsequently mentioned items are not meant to be limited to the examples given, since any other specific item or example that would fall under the generalization of the idea or concept is meant to be included. For any given compound, it might be more appropriate to produce a salt of said compound, so for example, succinic acid might be produced at pH near 7 as a salt of sodium, potassium, calcium, magnesium, ammonium, etc., while lysine might be produced as a salt of chloride, sulfate, bicarbonate, etc. As such, any time a compound is named herein, any salt of said compound is meant to be included, and any time a salt is named, the free acid or free base is also meant to be included. Thus, for example, “succinate” is meant to include “succinic acid” and vice versa, and “acetate” is meant to include “acetic acid” and vice versa.

[0041] “Facilitated diffusion” means the action of a system, typically comprising an integral membrane protein situated in a biological membrane (for example the inner membrane of a Gram negative bacterium or the single membrane of a Gram positive bacterium), or a complex of more than one protein molecules situated in a biological membrane, that functions to specifically allow one or more chemicals called the “substrate” (for example glucose and/or fructose), but not chemicals in general (for example water and cytoplasmic metabolites other than the specific substrate), to cross through the membrane without any energy (such as that provided by hydrolysis of ATP or PEP) or gradient of a different chemical (for example a proton gradient) provided directly to the system by the cell. If there is a concentration gradient, for example if the concentration of a substrate is higher outside the cell than inside the cell, there will be a net flux of that substrate into the cell at a rate that is faster than would occur if the facilitated diffusion system were absent. The protein(s) that function for facilitated diffusion typically have a binding affinity that is specific for one or more substrates and allows the system to assist passing the substrate across the membrane at relatively low concentrations of several millimolar or less. Some types of facilitated diffusion can function by creating a pore or channel through the membrane that discriminates in favor of a substrate, and in other types the protein(s) can bind the substrate on one side of the membrane and then rotate through the membrane to release the substrate on the opposite side of the membrane. A facilitated diffusion protein (sometimes called simply a facilitator) is a protein component of such a system. Thus, the thermodynamic driving force for facilitated diffusion is a gradient of substrate concentration, in which the substrate (for example a sugar) flows from a higher concentration outside of a cell to a lower concentration inside the cell. We shall use the genetic symbols Glf and glf to respectively mean a facilitated diffusion protein and a gene encoding such a protein that has specificity for glucose. We usually consider Glf to be a comprised of a single polypeptide chain, but a Glf could be a complex comprised of more than one polypeptide chain. Although the specific examples of Glf written herein are bacterial in origin, our definition is meant to include facilitated diffusion system derived from any organism. For example, it is well known that the yeast *Saccharomyces cerevisiae* and other yeasts have one or more facilitated

diffusion proteins for importing hexoses (for example glucose and fructose) named HXT1, HXT2, HTX3, HTX4, HTX5, HTX6, HTX7, etc.), and human erythrocytes use facilitated diffusion to import and export glucose via a protein named GLUT1. The mechanism of action of Glf's can vary widely, including pore-facilitated transport and carrier-facilitated transport. Although the specific examples given in this specification disclose a Glf that has good specificity for glucose, it is known in the art that a Glf protein can be active on more than one sugar, for example Glf from *Zymomonas mobilis* and *Saccharomyces cerevisiae* can be active on fructose as well as glucose.

[0042] Proton symport is defined as a system for importing a substrate across a biological membrane that uses a proton gradient as a driving force. A higher concentration of protons outside of the cell has a thermodynamic tendency to diffuse back into the cell. This thermodynamic pressure is used to carry in a substrate such as a sugar. A proton symporter is a protein or complex of proteins that functions to provide proton symport. An example of a proton symporter is the GalP protein of *E. coli*, which is well known to function in the import of galactose, glucose, and other sugars.

[0043] A glucokinase and a fructokinase are enzymes that catalyze phosphorylation of glucose, fructose, or other sugar, usually at the 6th carbon position, but alternatively possibly at the 1st carbon or another position. We shall use the genetic symbols Glk and glk to respectively mean a glucokinase and a gene that encodes a glucokinase. Frk and frk mean a fructokinase and a gene that encodes a fructokinase, respectively.

[0044] A crr gene is a gene that encodes an EIIA^{glc} component of a PTS, such as the crr gene of an *E. coli* strain or of a *Bacillus subtilis* strain or a homolog of such a crr gene.

[0045] A PTS (phosphotransferase system) is a group of proteins that act together to pump a sugar into a cytoplasm and simultaneously phosphorylate the sugar, using PEP as the source of phosphate and energy. Examples of genes encoding PTS proteins from *E. coli* include ptsH, ptsI, crr, ptsG, fruA, fruB, manX, manY, and manZ. The corresponding proteins are named PtsH, PtsI, Crr, PtsG, FruA, FruB, ManX, ManY, and ManZ. However, there are many more examples from *E. coli* and other prokaryotes, and these proteins can have alternate names, for example Crr is sometimes named EIIA^{glc}. Some of the PTS proteins are more specific to one or more particular sugars than to other sugars, while some PTS proteins, for example PtsH and PtsI from *E. coli*, are used in common for many different sugars.

[0046] In this specification, the term “microaerobic” means that the feed rate of air is less than 0.1 volume of air per volume of liquid culture per minute. In 7 and 20 liter fermentor examples disclosed herein, this is accomplished with a sparger and flow meter, or by allowing the tank to breathe through a sterile membrane attached to the top of the tank without any forced air flow. In 500 ml fermentor tank examples disclosed herein, no air is deliberately introduced, but a small amount of air is introduced from leakage, feeding of the base solution, and taking of samples.

[0047] A “minimal medium” is a microbial growth medium comprised of water, a pure carbon source (such as a substantially pure sugar or mixture of substantially pure sugars), mineral salts (for example potassium, sodium, magnesium, calcium, bicarbonate plus carbonate, phosphate, sulfate and chloride), a pure nitrogen source such as ammonium or urea, trace metals (iron, copper, zinc, manganese, cobalt, molybdenum, and optionally borate), optionally glycine betaine

(also known as simply betaine), and optionally an antifoam agent. Minimal media do not contain any complex (also known as “rich”) nutrient source such as yeast extract, corn steep liquor, soy hydrolysate, broth, casein hydrolysate, grain, legume, or any other “undefined” mixture of nutrients that typically would be derived from an agricultural source without any physical or chemical purification or separation steps. Reasonably pure sugars derived from sugar cane, corn starch, sorghum starch, tapioca starch, or any other reasonably pure starch source is considered to be acceptable for a minimal medium. A minimal medium can contain one or a few pure chemicals to satisfy a particular growth requirement (auxotrophy or bradytrophly) or to enhance a biochemical pathway. For example, some strains require a vitamin such as biotin, which can be added at small concentrations without a significant negative impact on a process. As another example, addition of a vitamin such as thiamine, while not absolutely required for growth, can nonetheless enhance growth or a biochemical pathway. Minimal media are preferable for fermentative production of many chemicals due to the relatively low cost of the components, and due producing cleaner fermentations broths that allows for more favorable economics for downstream purification of the desired chemical. Ethanol production is an exception, since downstream purification can be accomplished with distillation, an economically attractive method for purification of the desired product even from complex media.

[0048] An aromatic biochemical means any one or more of the following: phenylalanine, tyrosine, or tryptophan, or any derivative thereof (such as L-dihydroxyphenylalanine, melatonin, indole, indole acetic acid, indigo, serotonin, cinnamic acid, hydroxy styrene), a vitamin or vitamin-like compound containing an aromatic moiety (such as p-hydroxybenzoic acid, 2,3-dihydroxybenzoic acid, p-amino benzoic acid, folate, tocopherol, pyrroloquinoline quinone).

[0049] A homolog of a first gene or protein is defined as a second gene or protein in which the second protein or the protein inferred to be translated from the second gene has the same or a similar biochemical function as the first protein or protein inferred to be translated from the first gene, and in which an alignment of the first and second proteins or first and second inferred translated proteins results in a 25% or greater identity or similarity for a region of at least 50 amino acids in length, when using the default parameters of a publically available computer alignment program such as BLAST.

[0050] A mutation is any change in a DNA sequence relative to the DNA sequence of the related wild type or native gene. A mutation can be a single or multiple base change that introduces a premature stop codon or an amino acid that is different from the wild type amino acid at that position. A mutation can be an insertion or deletion of one or more bases that creates a frame shift that results in a protein that is significantly different from the wild type protein. A mutation can be a deletion that removes much, most, or all of a coding region (also known as an open reading frame or orf). One type of mutation removes one or more entire orfs plus additional non-coding DNA either upstream or downstream from the coding region, or both. A mutation can result from insertion of a relatively large DNA sequence (more than about 100 bases), for example an insertion element (for example IS186 or IS4) or a transposon (for example Tn10). When the intent is to remove a function, a preferable mutation is a deletion of all or most of an orf, however, smaller mutations such as single base changes or insertions can often accomplish removal of a

function for all practical purposes. Mutations can be spontaneous, induced by mutagenesis, or constructed by genetic engineering. Some mutations, when desired to accomplish a strain improvement, are mutations that decrease or eliminate a biological function, such as particular elements of a PTS. However, some mutations, when desired to accomplish a strain improvement, are mutations that increase a biological function, for example a “promoter up mutation” can increase the expression of a desired gene, such as a *glf* gene.

[0051] “Exogenous” means a gene or protein derived from a second genus that has been installed in a first genus, where said second genus is a different genus from said first genus.

[0052] A gene is defined as a region of a chromosome that encodes a protein or enzyme, and is meant to include both the open reading frame that corresponds to the protein or enzyme and any DNA sequences surrounding the open reading frame that contribute to controlling the level or rate of production of the protein or enzyme, such as promoters, ribosome binding sites, operators, regulatory protein binding sites, DNA corresponding to 5' untranslated mRNA leader sequences, terminators, and antiterminator sites. When two or more open reading frames that correspond to protein coding DNA sequences are under the control of a single promoter and a single terminator, the whole region encompassing the promoter, open reading frames corresponding to protein coding DNA sequences and the terminator is referred as an operon. For example, when the exogenous genes *glf* and *glk* are under the control of a single promoter and a single terminator, it is referred as *glf-glk* operon.

[0053] The present invention provides biocatalysts for succinic acid production in high titer, yield and productivity using a minimal medium with a sugar as a carbon source. The term “yield” as defined in this invention refers to the number of grams of product (such as succinic acid) produced per gram of sugar (such as glucose or sucrose) consumed. The term “productivity” as defined in this present invention refers to the number of grams of product (such as succinic acid) produced per liter of culture per hour. The term “titer” is defined as the concentration of product (such as succinic acid) in the fermentation broth in grams per liter. The desirable yield for succinic acid is in the range of 0.8-1.2 grams of succinic acid produced per gram of sugar consumed. The desirable productivity for succinic acid in this present invention is in the range of 1 gram or more of succinic acid produced per liter per hour. The desirable titer of succinic acid is greater than 26 g/l, or more preferably greater than 64 g/l, and most preferably greater than 83 g/l in a fermentation time of 48 hours or less.

[0054] The bacterial growth rate is measured in terms of the rate of increase in the optical density at 550 or 600 nanometers of a liquid culture resulting from the bacterial multiplication. The bacterial growth rate is also expressed in terms of time required for doubling of bacterial cells. In the bacterial cells suitable for the present invention, the bacterial cells are expected to have a doubling time of between 20 minutes and 3 hours.

[0055] According to the present invention, the biocatalyst for succinic acid production can be developed in two different ways. Under the first approach, a wild type bacterial species is genetically manipulated and, optionally, evolved, to grow efficiently using facilitated diffusion for import of glucose or other sugar. Once such a strain is constructed, subsequent genetic manipulations are carried out in the metabolic pathways to obtain a bacterial strain that produces succinic acid or

another desired chemical with high titer, yield and productivity, for example, by following methods known in the art.

[0056] The patent applications published under Patent Cooperation Treaty with the publication No. WO 2010/115067 and United States Patent Application Publication No. US 20100184171 provide the details about the genetic engineering techniques useful in generating a strain of *E. coli* with improved succinic acid production capacity. These two patent applications are incorporated herein by reference.

[0057] Under the second approach, a bacterial strain already developed to have a commercially attractive yield and productivity for a chemical such as succinic acid as described in the patent application publications US 20100184171 and WO 2010/115067 is used as a parental strain. Further genetic manipulations, and optionally, evolution, are then carried out with this strain to obtain a bacterial strain that has the ability to use facilitated diffusion to import glucose or another sugar to produce succinic acid at a commercially attractive titer, yield, and productivity.

[0058] As a specific example, this present invention discloses biocatalysts and methods that have improved ability over that of the prior art to produce succinic acid at high enough titer, yield and productivity while gaining the new ability to import a sugar by facilitated diffusion. For example, the KJ122 strain of *E. coli* described by Jantama et al. can be selected as the starting strain for the present invention. The KJ122 strain of *E. coli* is reported to have the ability to produce succinic acid in a minimal medium at high titer and productivity.

[0059] The KJ122 strain of *E. coli* was derived from the *E. coli* C strain through gene deletions and metabolic evolution as described in US Patent Application Publication No. 20100184171 and in the International Patent Application Publication No. WO 2010/115067. These two patent application publication documents providing details about the genetic changes that led to the development of the KJ122 strain of *E. coli* are incorporated herein by reference. KJ122 does not have any substantial ability to import glucose as a source of carbon by facilitated diffusion in the production of succinic acid. The absence of this function in KJ122 is attributable to the lack of a gene that encodes a Glf protein. The inventors have discovered genetic approaches that enable KJ122 to more efficiently use glucose as a source of carbohydrate while retaining or improving its original ability to produce succinic acid at high titer, yield, and productivity in a minimal medium.

[0060] The term “carbohydrate” as used in this invention includes mono-saccharides such as glucose, fructose, xylose, and arabinose, disaccharides such as sucrose, melibiose, maltose and lactose, trisaccharides such as raffinose and maltotriose, and higher oligosaccharides, and hydrolysates derived from the enzymatic or chemical digestion of polysaccharides such as starch, cellulose, and biomass. Simple carbohydrates, those with from one to three saccharide units, are referred to herein as “sugars”, for example glucose, fructose, sucrose, maltose, etc.

[0061] The terms “PTS⁺ organism” or “PTS⁺ bacterium” refers to a bacterium which has the capacity for a carbohydrate transport based on a PTS. The term “non-PTS organism,” or “non-PTS bacterium” or “PTS⁻” bacterium refers to bacterial cells that are mutated in one or more genes that encode a PTS function, such that the activity of the PTS is decreased relative to that of the wild type PTS.

[0062] In one aspect, the present invention discloses the addition of genes to an organism in order to install or increase the activity of one or more proteins and/or enzymes involved in the import and conversion of a sugar into metabolic intermediates such as glucose 6-phosphate, glucose 1-phosphate, fructose 6-phosphate, or fructose 1-phosphate that can be further metabolized by the cell. The genes that encode relevant proteins or enzymes are chosen from a group consisting of a glf gene, an HXT gene, a glk gene, and a frk gene.

[0063] In another embodiment, the present invention provides a process for producing succinic acid or another chemical using facilitated diffusion to import a sugar such as glucose as a renewable feedstock. In one aspect, the present invention provides a process for producing succinic acid from a sugar-containing medium that makes use of a biocatalyst that has a decreased activity in at least one protein of the organism’s native PTS system relative to that of the ancestral or parental strain. In another aspect, the present invention provides a process for producing succinic acid or other chemical in a sugar-containing medium that makes use of a biocatalyst that has a decreased activity in at least one protein of the organism’s native sugar import system relative to that of the ancestral or parental strain involving use of a protein symport system, such as GalP.

[0064] The present invention provides ways to manipulate a PTS and in turn the bacterial carbohydrate uptake system. Since EI and HPr proteins function as “general” or “common” components of the PTS system, inactivation of either the ptsI gene coding for EI protein or the ptsH gene coding for HPr protein would lead to the complete inactivation of a PTS. There will be substantially less carbohydrate transport through the PTS system in bacterial cells where the activity of ptsH or ptsI or both has been decreased or eliminated. When the PTS is partially or completely inactivated, the bacterial cell has to depend on one or more other alternative permease systems for carbohydrate transport.

[0065] When there is active glucose transport through PTS, the EIIA^{Glc} remains unphosphorylated as there is a carbohydrate substrate for accepting its phosphate group. However, when there is no glucose in the medium, the phosphorylated form of EIIA^{Glc} cannot transfer its phosphate group to glucose and therefore it remains in its phosphorylated state. The unphosphorylated EIIA^{Glc} mediates the phenomenon generally known as carbon catabolite repression (CCR). Under CCR, when glucose is present in the growth medium, the transport and/or utilization of other carbohydrates in the medium is prevented until the glucose in the medium is decreased to a low concentration. The carbon catabolite repression results from the inhibitory effect of unphosphorylated EIIA^{Glc} on permease systems or other systems of carbon source utilization. A number of permeases involved in the carbohydrate transport are known to be inhibited by unphosphorylated EIIA^{Glc}, for example, LacY or lactose permease. In addition, the unphosphorylated EIIA^{Glc} is known to have a negative effect on the transcription of number of genes involved in carbohydrate transport and metabolism through its influence on the adenylate cyclase system.

[0066] Strain KJ122, good succinate producer, contains a frameshift mutation in the ptsI gene, and this mutation is important for good succinate production. Thus it was surprising in the context of the current invention that further improvements in succinate production could be made by deleting ptsHI and galP, and then installing a facilitated diffusion system.

[0067] In another embodiment, the present invention provides a non-naturally occurring duplication of the *crr* gene that encodes the EIIA^{Glc} protein. The inventors discovered that strains containing a *ptsHI* deletion, a *galP* mutation, and an installation of a functional *glf* gene, have an unexpected tendency to acquire a mutation in the *crr* gene which causes a decrease or elimination in function of the EIIA^{Glc} protein, which in turn causes an unexpected undesirable decrease in succinate production parameters. Duplication of the *crr* gene by integrating a second copy of *crr* at a locus separate from the native *crr* locus solves this problem by greatly reducing the frequency of mutants that become phenotypically *crr* negative.

[0068] The present invention will be explained in detail below. An example bacterium belonging to the genus *Escherichia* of the present invention is a strain which is constructed from a parental strain that is not initially capable of using facilitated diffusion for sugar import, but which after genetic engineering as disclosed herein harbors a *glf* gene, and optionally an exogenous *glk* gene, and has the ability to use facilitated diffusion for import of glucose and fructose.

[0069] The exogenous genes introduced into the cell can be maintained within the cell on a self-replicating plasmid. A plasmid can be maintained through antibiotic selection or complementation of a chromosomal mutation. However, when the exogenous genes are maintained within the biocatalyst on a self-replicating plasmid within the cell, it is necessary to assure there is no unnecessary waste of energy and materials leading to the inhibition of growth, and a decrease in the yield or productivity of the organic material being manufactured using the biocatalyst. Preferentially, the exogenous genes are integrated into the host chromosome so that there is no need to add any antibiotics to maintain the plasmids within the cell, and little or no metabolic burden is placed on the cell for plasmid maintenance. There are many possible locations within the cell for the integration of the exogenous genes. The preferential locations for integrating the exogenous genes within the *E. coli* chromosomal DNA include regions that do not encode an essential function for growth and product formation under commercial fermentation conditions.

[0070] When the exogenous genes are obtained as an operon, it is preferable to remove any possible negative regulatory genes or proteins from the operon. It is ideal to have only the genes and proteins that function positively in facilitated diffusion and metabolism. Thus, expression of a facilitated diffusion gene is preferably not inhibited by a repressor or by carbon catabolite repression.

[0071] The following examples are provided as a way of illustrating the present invention and not as a limitation.

[0072] Any bacterium that does not natively use a facilitated diffusion system for sugar import can be improved according to the present invention.

[0073] A bacterium of the present invention may be obtained by introduction of one or more genes that enables utilization of facilitated diffusion into a succinic acid producing strain such as KJ122 or other strain previously engineered to produce a desired chemical. Alternatively a bacterium of the present invention may be obtained by conferring an ability to produce succinic acid or other desired chemical to a bacterium in which utilization of facilitated diffusion has already been enabled by genetic engineering, and optionally by evolution. This latter alternative can be accomplished, for example, by following all the steps used for constructing

KJ122 but starting with strain ATCC 9637 or a K-12 type *E. coli* strain, or any other safe *E. coli* strain, instead of starting with strain ATCC 8739.

Example 1

Construction of AC15, a Derivative of KJ122 that Contains the *Glf* and *Glk* Genes from Gene Cluster from *Zymomonas mobilis* CP4

[0074] All manipulations of DNA and plasmids, polymerase chain reaction (PCR), transformation, and chromosomal integration were accomplished by standard methods that are well known in the art. It is well known that DNA sequences can be cloned and joined together to form new combinations that cannot be easily found in nature. In addition to the more traditional methods involving restriction enzymes and DNA ligase, newer methods using recombinering in yeast, the so-called "Gibson Method" of in vitro splicing of DNA, or any other appropriate method can be used to construct such novel DNA sequences. The DNA fragments needed can be obtained from libraries of clones or by PCR from appropriate template DNA. It is also understood that many desired DNA sequences can be designed and synthesized from chemical precursors. Such a service is supplied by a number of commercial companies, for example DNA 2.0 and GeneArt (Invitrogen).

[0075] Plasmid pAC19 was constructed to contain an artificial operon containing the *glf* and *glk* genes from *Z. mobilis*, driven by the P₂₆ promoter from the *Bacillus subtilis* phage SP01. This operon was embedded between an upstream sequence homologous to the *E. coli* C *tdcC* gene and a downstream sequence homologous to the *E. coli* C *tdcE* gene, to foster integration into the *tdcCDE* locus of strains to be engineered. The cassette described above is carried on a low copy plasmid vector derived from pCL1921, which contains the pSC101 origin of replication and a spectinomycin resistance gene. The components for the cassette were obtained by PCR using appropriate synthetic DNA primers obtained from commercial suppliers such as Sigma and Integrated DNA Technologies (IDT). The source for the *Zymomonas* genes was pLO11740, which originally contained a *zwf* and *edd* gene in addition to the desired *glf* and *glk* genes. The *glf*, *zwf*, *edd*, *glk* cluster was transferred to pCL1921, and then the unnecessary *zwf* and *edd* genes were deleted by inside out PCR. The upstream and downstream *tdc* sequences were obtained by PCR from KJ122 chromosomal DNA as template. The P₂₆ promoter was obtained from bacteriophage SP01. The sequence of pAC19 is given as SEQ ID #1.

[0076] All constructions were done while growing strains on LB medium (10 grams Bacto-tryptone, 5 grams Bacto-yeast extract, and 5 grams sodium chloride) supplemented as appropriate with antibiotic or sucrose. To construct strain AC15, the cassette containing the artificial operon of pAC19 was integrated into the chromosome of strain WG53, using a two step gene replacement method previously described. The *cat*, *sacB* cassette for the first step was contained on plasmid pAC21, SEQ ID #2. pAC21 is similar to pAC19, except that the artificial operon is replaced with a *cat*, *sacB* cassette that contains a chloramphenicol resistance gene and a counterselectable *sacB* gene encoding levan sucrose. The transforming DNA was obtained by PCR from pAC21 for the first step and by PCR from pAC19 for the second step.

[0077] Strain WG53 was obtained by deleting the *ptsH*, *ptsI*, and *galP* genes from succinate producing strain KJ122,

using a two step gene replacement method similar to that described in the above paragraph. The DNA sequence spanning the ptsHI deletion is given as SEQ ID #3. Note that this deletion leaves the crr gene intact, as well as native promoters that naturally exist upstream from the ptsH gene. The DNA sequence spanning the galP deletion is given as SEQ ID #4.

[0078] While intermediate strain WG53 grew extremely poorly on minimal glucose medium, strains KJ122 and AC15 grew well on minimal glucose medium, demonstrating that 1) the ptsHI and galP genes had been successfully deleted in WG53, and 2) the glf, glk cassette was functional in AC15 allowing glucose to be imported.

Example 2

Strain AC15 Produces Succinate as Well as Parent KJ122

[0079] Strains KJ122 and AC15 were grown under microaerobic condition in 7 liter fermentors (New Brunswick Scientific) at 39° C. using a minimal medium with glucose fed batch system. The starting volume of 3 liters contained potassium phosphate monobasic (18 mM), magnesium sulfate (2 mM), betaine (1.33 mM), trace elements, Antifoam 204 (8 ppm) and 25 g/l glucose. The pH was adjusted initially to pH 7.0 and thereafter was maintained at pH 6.5 as acid was produced by addition of the ammonium hydroxide/ammonium bicarbonate solution described below. The 150 ml inocula were grown aerobically and contained a minimal medium similar to the above described medium, except that glucose was at 20 g/l and calcium chloride was added to a final concentration of 0.1 mM. Agitation was set at 750 RPM (revolutions per minute). When glucose decreased to 5 g/l, a 650 g/l glucose feed was started and maintained at a rate aimed to keep the glucose concentration at about 5 g/l or less. The stock solution used for neutralization contained both ammonium hydroxide and ammonium bicarbonate (7 N NH₄OH and 3M NH₄HCO₃). AC15 was aerated at 35 ml/min, while KJ122 was not given air other than what was present in the head space, which was equilibrated with the atmosphere through a breathable sterile membrane filter. These were conditions that had been shown to work well for each strain. Sugars, succinate, and byproducts from 48 hour samples were assayed by HPLC. The results of averaged duplicates are shown in Table 1. AC15 produced about the same titer as parent KJ122, but the acetate byproduct was significantly lower, and the yield on glucose was higher for AC15.

Example 3

Spontaneous “Red Mutants” Derived from AC15

[0080] KJ122 is able to ferment lactose, as evidenced by formation of red colonies on MacConkey lactose plates (Beckton-Dickinson, Franklin Lakes, N.J.). However, AC15 does not ferment lactose, as evidenced by producing “white” (beige) colonies on MacConkey lactose plates. This white colony phenotype of AC15 results from binding and inhibition of lactose permease (LacY) by unphosphorylated EIIA^{Glc} protein. This white colony phenotype is present in all strains deleted for ptsHI, since the enzymes required to phosphorylate EIIA^{Glc} are absent, and as a result, all EIIA^{Glc} present in the cells remains unphosphorylated. Thus, ironically, *E. coli* ptsHI mutants are phenotypically Lac⁻, even though lactose is not imported by the PTS system in *E. coli*.

[0081] The inventors noticed by chance that when MacConkey lactose plates were streaked with AC15 and allowed to incubate overnight at 37° C., and then for an extra day at room temperature (about 22° C.), a large number of red colonies emerged from the lawn of white colonies that had grown over the denser part of the streak. Upon restreaking of several of the red colonies, it was observed that two classes of red colonies had evolved. We shall call the first class “solid red”, since the individual colonies were uniformly red across the entire colony. A second class shall be called “fried egg red”, since the individual colonies were red in the center, but the outer portion of the colonies were white or beige. We shall call the strains giving rise to all types of red colonies on MacConkey lactose collectively “red mutants”.

[0082] A white colony of AC15, and four red mutants, named AC15-R1, -R2, -R3, and -R4 (two of which are solid red and two of which are fried egg red), were tested for succinate production in 500 ml microaerobic fermentors (Fleakers, Corning Glass, Corning, N.Y.) using a medium and method similar to those described above for the 7 liter fermentors, with the differences being that the starting volume of the minimal medium was 200 ml, the glucose was all batched in the starting medium at 100 g/l, no glucose was fed, agitation was with a magnetic stirring bar at 350 RPM, and no air was deliberately introduced or removed. The results are shown in Table 2. The two fried egg mutants performed similarly to parent AC15, while the two solid red mutants performed significantly worse than parent AC15.

[0083] Genome sequencing of the parent AC15 and the four red mutants, using the Illumina HiSeq2000 system, revealed that both solid red mutants had acquired one mutation each, and both of these mutations were in the crr gene, which encodes EIIA^{Glc}. Both were judged to be null mutations. Both fried egg red mutants had acquired one mutation each, and both of these mutations were in the lactose operon. Both of these were judged to be mutations that would lead to a higher level of expression of the lactose operon (one was a mutation in the lacO operator, and the other was a frameshift in lacI, the gene that encodes the Lac repressor. All four mutations made sense in that they could explain the observed phenotype of increased ability to ferment lactose. The crr null mutations relieved the inhibition of the LacY permease, as would be expected, while the lactose operon mutations would be expected to overproduce LacY, allowing at least some escape from the inhibition. However, the crr null mutations clearly had an additional pleiotropic effect, causing a decrease in the cells’ ability to produce succinate under our fermentation conditions. This was an unexpected effect that was not predicted.

Example 4

[0084] The *Zymomonas mobilis* glk gene is not essential for functioning of the glf gene in *E. coli* (084) Plasmid pSS2 was constructed using methods similar to those described above for pAC19. The only differences between pSS2 and pAC19 is that the, *Z. mobilis* glk gene was deleted from the artificial operon. In other aspects, such as vector backbone, the promoter driving expression of glf, embedding the artificial operon in the tdc flanking sequences, and orientation of the various components, pSS2 is similar to pAC19. The DNA sequence of pSS2 is given as SEQ ID #5.

[0085] The artificial operon from pSS2 was integrated at the tdc locus of KJ122 as described above for the operon from pAC19, using the two step gene replacement method. Two

isolates, which are presumably identical to each other were named SS8-9 and SS8-11. These two new strains were compared to AC15 in 500 ml microaerobic fermentors as described above in Example 3. The results, which are averages of duplicate fermentors assayed at 48 hours, are shown in Table 3. SS8-9 and SS8-11 both gave growth and succinate titers similar to that of AC15, while the acetate production of both SS8 isolates was somewhat lower than that of AC15. Thus, the *Z. mobilis* glk gene is unnecessary for functioning of the glf gene in this context, and the *Z. mobilis* glk gene might even be slightly harmful to the fermentation parameters. Presumably, the SS8 isolates are using the endogenous *E. coli* glk gene to phosphorylate glucose.

Example 5

Metabolic Evolution of Strain AC15

[0086] As noted above in Example 3, strain AC15 preferred to receive a higher level of aeration than parent KJ122 in 7 liter fermentors. In order to obtain a derivative of AC15 that could thrive on less air, AC15 was subjected to metabolic evolution in 500 ml fermentors with a starting volume of 200 ml and no deliberate supply of aeration. The conditions were microaerobic, since no measures were taken to remove oxygen. A small amount of air was assumed to leak into the fermentation vessels during the course of the evolution. The conditions for growth were as described in Example 3. After 48 hours of growth, the culture was diluted 1:100 into a fresh fermentor containing 200 ml of fresh medium, and this step was then repeated 40 more times. Each one of these inoculations to fresh medium shall be called a “transfer”. Thus, the strain was subjected to a total of 41 transfers to fresh medium. Each transfer corresponds to about 7 generations of cell division. A sample of the liquid culture from the last transfer was plated on a MacConkey lactose agar petri plate, and a single white colony was chosen and named YSS41.

[0087] By varying the rate of aeration in 7 liter fermentors, it was determined that YSS41 performed well for succinate production with 5 ml/min of air, which was substantially less than the 35 ml/min required for optimal performance of the parent AC15. With 5 ml/min air flow, YSS41 produced 94 g/l succinate and 1.3 g/l acetate, for a succinate yield of 0.95 g/g glucose in 48 hours in a 7 liter fermentor.

[0088] YSS41 was compared to KJ122 for succinate production in a 20 liter fermentor. The fermentation protocol was similar to that described above for 7 liter fermentors, except that the starting volume was 9 liters, and the aeration rate was 25 ml/min for both strains, conditions that had been determined to be productive for both strains. The results for 48 hour samples are shown in Table 4. The succinate titer for YSS41 was 100 g/l (significantly higher than for KJ122), the acetate as 2.2 g/l (significantly lower than for KJ122), and the succinate yield was 0.95 g/g glucose (a little lower than for KJ122). Thus, the evolved strain YSS41 was able to perform well in a 20 liter fermentor with an aeration requirement that was no higher than for the ancestor strain KJ122.

Example 6

Stabilizing YSS41 Against Mutations in the Crr Gene

[0089] When streaked on MacConkey lactose plates, YSS41 still gave rise to red mutants, both of the solid red type and of the fried egg red type. The crr gene was sequenced for

one isolate of each type. Strain MYR222, a fried egg type had a wild type crr gene sequence. MYR223, a solid red type, had an insertion element inserted in the crr open reading frame. The DNA sequence of the insertion element matched that of IS186. Thus, the pattern established for AC15 red mutants appeared to apply also to YSS41 red mutants. In 500 ml microaerobic fermentors, grown as in Example 4, MYR222 performed similarly to YSS41, while MYR223 performed more poorly (see Table 5). Thus the potential loss of performance due to accumulation of solid red mutants in a population remained a possibility with strain YSS41.

[0090] In order to solve this potential loss, a second copy of the crr gene was integrated into a site distant from the native crr locus. The crr gene, together with its flanking promoters and terminator were amplified by PCR using YSS41 chromosomal DNA as a template, and primers BY249 (SEQ ID #6) and BY250 (SEQ ID #7). The resulting blunt fragment was then ligated into a low copy plasmid derived from pCL1921 that contained a clone of a portion of the pflDC region from *E. coli* C at the unique BstZ171 restriction site in the pflD open reading frame. The pflDC genes are homologous to the pflBA genes that encode pyruvate-formate lyase and the pyruvate-formate lyase activating enzyme. The pflDC genes are not essential for *E. coli*, and deletion of either pflD or pflC has no significant effect on growth, so it was reasoned that insertion of a cassette at that locus would not have any negative consequence for growth or succinate production. The resulting low copy plasmid, pMH68, contains the crr gene from YSS41 embedded in flanking sequences from pflDC, in a low copy plasmid. The DNA sequence of pMH68 is given as SEQ ID #8.

[0091] The integration cassette from pMH68 was amplified by PCR using primers BY124 (SEQ ID #9) and BY125 (SEQ ID #10), which were the same primers used to clone the pflDC genes to begin with. The integration cassette was then integrated into the chromosome of YSS41, using the two step gene replacement method. The resulting strain was named MH141, which is now a merodiploid for crr, meaning that it contains two copies of a wild type crr gene in two distant locations on the chromosome, one at its native locus, and the second inserted in the pflD open reading frame.

[0092] As expected, strain MH141 produced white colonies on MacConkey lactose plates. If a heavy streak is made, and the plates are allowed to incubate overnight at 37° C., and then for an extra day at room temperature, red colonies emerged from the lawn of white colonies that had grown over the denser part of the streak. However, the number of red mutants arising from MH141 was significantly lower than for a similar streak of YSS41 made on the same plate. 23 red mutants were picked from YSS41 and 12 red mutants were picked from MH141, and all were restreaked on MacConkey lactose plates. When scored for the type of red mutant, 12 of the 23 YSS41 red mutants were of the solid red type, while the other 11 of the 23 were of the fried egg type. In contrast, all 12 of the MH141 red mutants were of the fried egg type. Thus, by duplicating the crr gene in the chromosome, the rate of formation of the solid red mutants has been decreased by at least a factor of ten. One fried egg red mutant isolated from MH141 was named MH141-R1 and tested in 500 ml microaerobic fermentors as described above (see Table 5). Both MH141 and MH141-R1 performed similarly to parent YSS41 with respect to growth, succinate titer, and acetate titer. Thus, a more stable strain, MH141, has been constructed that uses facilitated diffusion for glucose import, and which produces a

higher titer of succinate and a lower titer of the byproduct acetate when compared to the ancestor strain KJ122, which uses a the GalP system for glucose import.

Example 7

YSS41 Acquired Mutations in the Glf, Glk Cassette During Metabolic Evolution

[0093] The DNA sequences of the glf, glk expression cassettes in AC15 and YSS41 were determined. The regions were amplified by PCR and the resulting fragments were sequenced over the glf and glk genes and more than 200 base pairs upstream and downstream, by the dideoxy chain termination method. The sequenced region corresponds to bases 4976 to 7920 of pAC19, given in SEQ ID #1. Two mutations were found that were acquired during the evolution of YSS41. The first mutation was a G to A change at base number 7742 of SEQ ID #1. This base is in the 5' untranslated region of the glf, glk mRNA transcript, just upstream from the glf open reading frame, and results in a C to U change at base -22 relative to the ATG start codon, or +15 relative to the start of transcription, in the glf mRNA (messenger RNA). This mutation is expected to increase or decrease the rate of translation of the glf open reading frame. The second mutation was a G to A change at base number 6173 of SEQ ID #1. This base is in the 5' untranslated region just upstream from the glk open reading frame, and results in a C to U change at base -15 relative to the ATG start codon in the glk mRNA. This mutation is expected to increase or decrease the rate of translation of the glk open reading frame. Thus, the evolution of YSS41 resulted in a more optimal balance of expression between the glf and glk open reading frames, to result in a strain that outgrew and outperformed the parent strain AC15.

[0094] Other mutations that alter the rate of transcription or expression of the glf and glk genes, or that alter the concentration, specific activity, or stability of the glf and glk proteins, can similarly achieve a more optimal balance between the two encoded proteins will also benefit growth and production of a desired chemical. These other alternative mutations can be obtained by the using the method described above for YSS41. This method can also be applied to strains engineered to produce products other than succinate, where the ability to use facilitated diffusion or sugar import has been engineered into the strain.

Example 8

[0095] Fermentation of KJ122 and YSS41 after optimization of air flow rate for YSS41 (095) The optimum air flow rate for parent strain KJ122 had been determined to be 25 ml/minute in a 20 liter fermentor. At the air flow rate of 25 ml/min, YSS41 strain showed better succinate titer and yield when compared to that of KJ122. Further improvement in succinate yield and titer with YSS41 strain was obtained by increasing the air flow rate to 50 ml/min. Thus the optimal air flow rate for YSS41 strain with reference to succinate yield and titer seems to be different from that of KJ122. Table 6 provides fermentation results in the 20 liter fermentor under the optimized air flow conditions for each strain. YSS41 outperformed parent KJ122 in titer, yield, and acetate byproduct formation. The initial volume of the fermentation was 9500 ml. After feeding glucose and neutralizing with base the final volume was 12500 ml.

REFERENCES

- [0096] All references are listed for the convenience of the reader. Each reference is incorporated by reference in its entirety.
- [0097] U.S. Pat. No. 5,602,030
- [0098] U.S. Pat. No. 6,962,794
- [0099] U.S. Pat. No. 7,220,561
- [0100] U.S. Pat. No. 8,389,214
- [0101] U.S. Pat. No. 8,476,041
- [0102] US Patent Application Publication No. 20050079617
- [0103] US Patent Application Publication No. 20090047719
- [0104] US Patent Application Publication No. 20090253192
- [0105] US Patent Application Publication No. 20100184171
- [0106] European Patent Document No. EP20070715351
- [0107] European Patent Document No. EP0785275B1
- [0108] International Patent Application Publication No. WO 2010/115067
- [0109] International Patent Application Publication No. WO2011/123154
- [0110] International Patent Application Publication No. WO2012/018699
- [0111] Altschul, S. F., Gish, W., Miller, W., Myers, E. W., and Lipman, D. J. (1990) Basic local alignment search tool, *J Mol Biol* 215, 403-410.
- [0112] Altschul, S. F., Madden, T. L., Schaffer, A. A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. J. (1997) Gapped BLAST and PSI-BLAST: a new generation of protein database search programs, *Nucleic Acids Res* 25, 3389-3402.
- [0113] Altschul, S. F., and Koonin, E. V. (1998) Iterated profile searches with PSI-BLAST—a tool for discovery in protein databases, *Trends Biochem Sci* 23, 444-447.
- [0114] Altschul, S. F., Wootton, J. C., Gertz, E. M., Agarwala, R., Morgulis, A., Schaffer, A. A., and Yu, Y. K. (2005) Protein database searches using compositionally adjusted substitution matrices, *FEBS J* 272, 5101-5109.
- [0115] Baba, T., Ara, T., Hasegawa, M., Takai, Y., Okumura, Y., Baba, M., Datsenko, K. A., Tomita, M., Wanner, B. L., and Mori, H. (2006) Construction of *Escherichia coli* K-12 in-frame, single-gene knockout mutants: the Keio collection, *Mol Syst Biol* 2, 2006 0008
- [0116] Barrell, W. O., Yi, K. C., and Conway, T. (1990) Sequence and genetic organization of a *Zymomonas mobilis* gene cluster that encodes several enzymes of glucose metabolism, *J Bacteriol* 172, 7227-7240.
- [0117] Gosset, G. (2005) Improvement of *Escherichia coli* production strains by modification of the phosphoenolpyruvate:sugar phosphotransferase system, *Microb Cell Fact* 4, 14.
- [0118] Hogema, B. M., Arents, J. C., Bader, R., and Postma, P. W. (1999) Autoregulation of lactose uptake through the LacY permease by enzyme IIAGlc of the PTS in *Escherichia coli* K-12, *Mol Microbiol* 31, 1825-1833.
- [0119] Jantama, K., Haupt, M. J., Svoronos, S. A., Zhang, X., Moore, J. C., Shanmugam, K. T., and Ingram, L. O. (2008) Combining metabolic engineering and metabolic evolution to develop nonrecombinant strains of *Escherichia coli* C that produce succinate and malate, *Biotechnol Bioeng* 99, 1140-1153

- [0120] Jantama, K., Zhang, X., Moore, J. C., Shanmugam, K. T., Svoronos, S. A., and Ingram, L. O. (2008) Eliminating side products and increasing succinate yields in engineered strains of *Escherichia coli* C, *Biotechnol Bioeng* 101, 881-893.
- [0121] Lee, G., Talkington, C., and Pero, J. (1980) Nucleotide sequence of a promoter recognized by *Bacillus subtilis* RNA polymerase, *Mol Gen Genet* 180, 57-65.
- [0122] Lee, G., Hannett, N. M., Korman, A., and Pero, J. (1980) Transcription of cloned DNA from *Bacillus subtilis* phage SP01. Requirement for hydroxymethyluracil-containing DNA by phage-modified RNA polymerase, *J Mol Biol* 139, 407-422.
- [0123] Lerner, C. G., and Inouye, M. (1990) Low copy number plasmids for regulated low-level expression of cloned genes in *Escherichia coli* with blue/white insert screening capability, *Nucleic Acids Res* 18, 4631.
- [0124] Merryman, C., and Gibson, D. G. (2012) Methods and applications for assembling large DNA constructs, *Metab Eng* 14, 196-204.
- [0125] Moran, C. P., Jr., Lang, N., LeGrice, S. F., Lee, G., Stephens, M., Sonenshein, A. L., Pero, J., and Losick, R. (1982) Nucleotide sequences that signal the initiation of transcription and translation in *Bacillus subtilis*, *Mol Gen Genet* 186, 339-346.
- [0126] Mueckler, M., Caruso, C., Baldwin, S. A., Panico, M., Blench, I., Morris, H. R., Allard, W. J., Lienhard, G. E., and Lodish, H. F. (1985) Sequence and structure of a human glucose transporter, *Science* 229, 941-945.
- [0127] Neidhardt, F. C., and Curtiss, R. (1996) *Escherichia coli and Salmonella: cellular and molecular biology*, 2nd ed., ASM Press, Washington, D.C.
- [0128] Niu, W., Draths, K. M., and Frost, J. W. (2002) Benzene-free synthesis of adipic acid, *Biotechnol Prog* 18, 201-211.
- [0129] Pao, S. S., Paulsen, I. T., and Saier, M. H., Jr. (1998) Major facilitator superfamily, *Microbiol Mol Biol Rev* 62, 1-34.
- [0130] Parker, C., Peekhaus, N., Zhang, X., and Conway, T. (1997) Kinetics of Sugar Transport and Phosphorylation Influence Glucose and Fructose Cometabolism by *Zymomonas mobilis*, *Appl Environ Microbiol* 63, 3519-3525.
- [0131] Rogers, P. L., Jeon, Y. J., Lee, K. J., and Lawford, H. G. (2007) *Zymomonas mobilis* for fuel ethanol and higher value products, *Adv Biochem Eng Biotechnol* 108, 263-288.
- [0132] Sambrook, J. (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y.
- [0133] Shanks, R. M., Kadouri, D. E., MacEachran, D. P., and O'Toole, G. A. (2009) New yeast recombineering tools for bacteria, *Plasmid* 62, 88-97.
- [0134] Snoep, J. L., Arfman, N., Yomano, L. P., Fliege, R. K., Conway, T., and Ingram, L. O. (1994) Reconstruction of glucose uptake and phosphorylation in a glucose-negative mutant of *Escherichia coli* by using *Zymomonas mobilis* genes encoding the glucose facilitator protein and glucokinase, *J Bacteriol* 176, 2133-2135.
- [0135] Tang, J., Zhu, X., Lu, J., Liu, P., Xu, H., Tan, Z., and Zhang, X. (2013) Recruiting alternative glucose utilization pathways for improving succinate production, *Appl Microbiol Biotechnol* 97, 2513-2520.

- [0136] Weisser, P., Kramer, R., Sahn, H., and Sprenger, G. A. (1995) Functional expression of the glucose transporter of *Zymomonas mobilis* leads to restoration of glucose and fructose uptake in *Escherichia coli* mutants and provides evidence for its facilitator action, *J Bacteriol* 177, 3351-3354.

- [0137] Zhang, X., Jantama, K., Shanmugam, K. T., and Ingram, L. O. (2009) Reengineering *Escherichia coli* for Succinate Production in Mineral Salts Medium, *Appl Environ Microbiol* 75, 7807-7813.

- [0138] Zhang, X., Jantama, K., Moore, J. C., Jarboe, L. R., Shanmugam, K. T., and Ingram, L. O. (2009) Metabolic evolution of energy-conserving pathways for succinate production in *Escherichia coli*, *Proc Natl Acad Sci USA* 106, 20180-20185.

TABLE 1

Production of Succinate by AC15 in 7 Liter Fermentors					
Strain	Relevant genotype	Aeration ml/min	Succinate g/l	Acetate g/l	Yield g/g glucose
KJ122	parent, ptsI*, galP+	0	87	5.2	0.83
AC15	KJ122, ΔptsHI, ΔgalP, P ₂₆ -glf, glk	35	87	2.7	0.88

TABLE 2

Production of succinate by AC15 red mutants in 500 ml microaerobic fermentors					
Strain	Colony phenotype on MacConkey lactose	Succinate g/l	Acetate g/l	OD ₆₀₀	Mutation found
AC15	white	74	2.4	7.5	none
AC15-R1	solid red	51	9.0	6.5	crr Lys16 frameshift
AC15-R3	solid red	65	6.6	7.0	crr Met1Ile
AC15-R2	fried egg red	73	2.4	8.0	lacO G11A
AC15-R4	fried egg red	73	3.0	7.5	lacI Asp300 frameshift

TABLE 3

Succinate production by SS8 isolates in 500 ml microaerobic fermentors				
Strain	Relevant genotype	Succinate g/l	Acetate g/l	OD ₆₀₀
AC15	KJ122, ΔptsHI, ΔgalP, P ₂₆ -glf, glk	64	6.0	7.5
SS8-9	KJ122, ΔptsHI, ΔgalP, P ₂₆ -glf	64	4.2	8.5
SS8-11	KJ122, ΔptsHI, ΔgalP, P ₂₆ -glf	64	3.6	8.2

TABLE 4

Succinic acid production by YSS-41, in a 20 liter fermentor					
Strain	Relevant genotype	Air flow rate ml/min	Succinate g/l	Acetate g/l	Yield on glucose g/g
KJ122	ptsI*	25	87	6.8	1.00
KJ122	ptsI*	25	85	6.8	0.98
YSS41	KJ122□□ΔptsHI, ΔgalP, P ₂₆ -glf, glk, evolved	25	100.	2.2	0.95

TABLE 5

Succinate production in 500 ml microaerobic fermentors by MH141, a merodiploid for crr+.					
Strain	Relevant genotype	Colony phenotype on MacConkey lactose	Succinate g/l	Acetate g/l	OD ₆₀₀
YSS41	AC15, evolved, crr ⁺	white	67	3.9	10.0

TABLE 5-continued

Succinate production in 500 ml microaerobic fermentors by MH141, a merodiploid for crr+.					
Strain	Relevant genotype	Colony phenotype on MacConkey lactose	Succinate g/l	Acetate g/l	OD ₆₀₀
MH141	YSS41, ΔpflD::crr ⁺	white	68	3.2	8.5

TABLE 6

Succinate production by <i>E. coli</i> strains KJ122 and YSS41 in 20 liter fermentors under optimized aeration conditions for both strains					
Strain	Air flow rate (ml/min)	Succinate titer (g/l)	Succinate yield on glucose (g/g)	Acetate titer (g/l)	Cell mass as OD ₆₀₀
KJ122	25	81	0.86	3.9	12
YSS41	50	96	0.98	2.5	13
YSS41	25	93	0.98	2.5	13

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 12

<210> SEQ ID NO 1

<211> LENGTH: 9430

<212> TYPE: DNA

<213> ORGANISM: Escherichia coli

<220> FEATURE:

<221> NAME/KEY: pAC19

<222> LOCATION: (1)..(9430)

<223> OTHER INFORMATION: Plasmid used to install a glf-glk cassette at the tdc locus of bacterial strain KJ122

<400> SEQUENCE: 1

```

gttgacagta agacgggtaa gcctggtgat gataccgctg ccttactggg tgcattagcc      60
agtctgaatg acctgtcagc ggataatccg aagtggtcag actggaaaat cagagggcag      120
gaactgctga acagcaaaaa gtcagatagc accacatagc agaccgcgca taaaacgccc      180
tgagaagccc gtgacgggct tttcttgat tatgggtagt ttccctgcat gaatccataa      240
aaggcgctg tagtgccatt taccgccatt cactgccaga gccgtgagcg cagcgaactg      300
aatgtcacga aaaagacagc gactcaggtg cctgatggtc ggagacaaaa ggaatattca      360
gcgatttgcc cgagcttgcg aggggtgctac ttaagccttt agggttttaa ggtctgtttt      420
gtagaggagc aaacagcgtt tgcgacatcc ttttgtaata ctgcggaact gactaaagta      480
gtgagttata cacagggctg ggatctattc tttttatctt tttttattct ttctttattc      540
tataaattat aaccacttga atataaaca aaaaaacaca caaaggctca gcggaattta      600
cagagggctc agcagaattt acaagttttc cagcaaaggc ctagcagaat ttacagatac      660
ccacaactca aaggaaaagg actagtaatt atcattgact agcccatctc aattggtata      720
gtgattaata tcacctagac caattgagat gtatgtctga attagttgtt ttcaaagcaa      780
atgaactagc gattagtgcg tatgacttaa cggagcatga aaccaagcta attttatgct      840
gtgtggcact actcaacccc acgattgaaa accctacaag gaaagaacgg acggtatcgt      900

```


-continued

tcacttataa	ccaatacget	cagatgatga	acatcagtag	ggaaaatgct	tatgggtgat	960
tagctaaagc	aaccagagag	ctgatgacga	gaactgtgga	aatcaggaat	cctttgggta	1020
aaggctttga	gattttccag	tggacaaact	atgccaagtt	ctcaagcgaa	aaattagaat	1080
tagtttttag	tgaagagata	ttgccttate	ttttccagtt	aaaaaaattc	ataaaatata	1140
atctggaaca	tgtaagtct	tttgaaaaca	aatactctat	gaggatttat	gagtgggtat	1200
taaaagaact	aacacaaaag	aaaactcaca	aggcaaatat	agagattagc	cttgatgaat	1260
ttaagtcat	gttaatgctt	gaaaataact	accatgagtt	taaaaggctt	aaccaatggg	1320
ttttgaaacc	aataagtaaa	gatttaaaca	cttacagcaa	tatgaaattg	gtggttgata	1380
agcgaggccg	cccgactgat	acgttgattt	tccaagttga	actagataga	caaattggatc	1440
tcgtaaccga	acttgagaac	aaccagataa	aatgaatgg	tgacaaaata	ccaacaacca	1500
ttacatcaga	ttcctaccta	cgtaacggac	taagaaaaac	actacacgat	gctttaactg	1560
caaaaattca	gctcaccagt	tttgaggcaa	aatttttgag	tgacatgcaa	agtaagcatg	1620
atctcaatgg	ttcgttctca	tggctcacgc	aaaaacaacg	aaccacacta	gagaacatac	1680
tggctaaata	cggaaggatc	tgaggttctt	atggctcttg	tatctatcag	tgaagcatca	1740
agactaacia	acaaaagtag	aacaactggt	caccgttaca	tatcaaaggg	aaaactgtcc	1800
atatgcacag	atgaaaacgg	tgtaaaaaag	atagatacat	cagagctttt	acgagttttt	1860
ggtgcattta	aagctgttca	ccatgaacag	atcgacaatg	taacagatga	acagcatgta	1920
acacctaata	gaacaggtga	aaccagtaaa	acaagcaac	tagaacatga	aattgaacac	1980
ctgagacaac	ttgttacagc	tcaacagtca	cacatagaca	gcctgaaaca	ggcgatgctg	2040
cttatcgaat	caaagctgcc	gacaacacgg	gagccagtga	cgctcccgt	ggggaaaaaa	2100
tcatggcaat	tctggaagaa	atagcgcttt	cagccggcaa	acctgaagcc	ggatctgcca	2160
ttctgataac	aaactagcaa	caccagaaca	gccgtttgc	gggcagcaaa	accggttatg	2220
cttgtaaacc	gttttgtaa	aaaattttta	aaataaaaaa	ggggacctct	agggtcccca	2280
attaattagt	aatataatct	attaaggctc	attcaaaagg	tcatccaccg	gatcaattcc	2340
cctgctcgcg	caggctgggt	gccaagctct	cgggtaacat	caaggcccga	tccttgagac	2400
ccttgccctc	ccgcacgatg	atcgtgccgt	gatcgaaatc	cagatccttg	accgcagtt	2460
gcaaaccctc	actgatccgc	atgccgcttc	catacagaag	ctgggcgaac	aaacgatgct	2520
cgcttccag	aaaaccgagg	atgcgaacca	cttcatccgg	ggtcagcacc	accggcaagc	2580
gccgcgacgg	ccgaggtctt	cogatctcct	gaagccaggg	cagatccgtg	cacagcacct	2640
tgccgtagaa	gaacagcaag	gccgccaatg	cctgacgatg	cgtggagacc	gaaaccttgc	2700
gctcgttcgc	cagccaggac	agaaatgcct	cgacttcgct	gctgccaag	gttgccgggt	2760
gacgcacacc	gtggaacgg	atgaaggcac	gaaccagtg	gacataagcc	tggtcggttc	2820
gtaagctgta	atgcaagtag	cgtatgcgct	cacgcaactg	gtccagaacc	ttgaccgaac	2880
gcagcgggtg	taacggcgca	gtggcggttt	tcatggcttg	ttatgactgt	ttttttgggg	2940
tacagtctat	gcctcgggca	tccaagcagc	aagcgcgtta	cgccgtgggt	cgatgtttga	3000
tgttatggag	cagcaacgat	gttacgcagc	agggcagtcg	ccctaaaaca	aagttaaaca	3060
tcatgagggg	agcgggtgatc	gccgaagtat	cgactcaact	atcagaggta	gttggcgctca	3120
tcgagcgcca	tctcgaaccg	acgttgctgg	ccgtacattt	gtacggctcc	gcagtggatg	3180

-continued

gcggcctgaa gccacacagt gatattgatt tgctggttac ggtgaccgta aggcttgatg	3240
aaacaacgcg gcgagctttg atcaacgacc ttttgaaaac ttcggcttcc cctggagaga	3300
gcgagattct ccgcgctgta gaagtcacca ttgttggtgca cgacgacatc attccgtggc	3360
gttatccagc taagcgcgaa ctgcaatttg gagaatggca gcgcaatgac attcctgcag	3420
gtatcttcga gccagccacg atcgacattg atctggctat cttgctgaca aaagcaagag	3480
aacatagcgt tgccttggtg ggtccagcgg cggaggaact ctttgatccg gttcctgaac	3540
aggatctatt tgaggcgcta aatgaaacct taacgctatg gaactcgccg cccgactggg	3600
ctggcgatga gcgaaatgta gtgcttacgt tgtcccgcac ttggtacagc gcagtaaccg	3660
gcaaaatcgc gccgaaggat gtcgctgccc actgggcaat ggagcgcctg ccggcccagt	3720
atcagcccgt catacttgaa gctagacagg cttatcttgg acaagaagaa gatcgcttgg	3780
cctcgcgcgc agatcagttg gaagaatttg tccactacgt gaaaggcgag atcaccaagg	3840
tagtcggcaa ataattgcta acaattcgtt caagccgacg ccgcttcgcg gcgcggtta	3900
actcaagcgt tagatgcact aagcacataa ttgctcacag ccaaactatc aggtcaagtc	3960
tgcttttatt atttttaagc gtgcataata agccctacac aaattgggag atatatcatg	4020
aaaggctggc ttttcttgt tatcgcaata gttggcgaag taatcgcaac atccgcatta	4080
aaatctagcg agggctttac taagctgatc cggtgatga ccttttgaat gacctttaat	4140
agattatatt actaattaat tggggaccct agaggtcccc ttttttattt taaaaatttt	4200
ttcacaaaac ggtttacaag catacgttgg ccgattcatt aatgcagctg gcacgacagg	4260
tttcccgact ggaaagcggg cagtgagcgc aacgcaatta atgtgagtta gctcactcat	4320
taggcacccc aggctttaca ctttatgctt ccggctcgta tgttggtgag aattgtgagc	4380
ggataacaat ttcacacagg aaacagctat gaccatgatt acgccaagct tgcattgctg	4440
caggctgact ctagaggatc cccccgcgcg ccgacagagt aataggtttt acttaatagc	4500
tcttcctgtc ccttcaggc agtgatccgc attccgttct catggcgagg caacatttcg	4560
ggatggaaga taatgttctt tgctacagga aatcaacaa tatgcccacc agatgccact	4620
ggcagccgcc cgtgcgcggt tactaactct ataatgcag ggatctcatc aatgacaaca	4680
tcttcgggac tgtttcctgc cagtcccatg atgatggcga catccgtggc atggcctttg	4740
cccgtcagtg acaacgacct gtacagatcg accacaatat ggctcgtcgc ggttaataag	4800
ccgctacttt ccagccgatc aataaaaactt tttccggcat tcattggccc caccggtatgc	4860
gaactggagg gaccaatccc aattttgaaa atatcgaatg cactaatcat atccacaccc	4920
tcggattgcc gttcagtgaa gtggagcggg acgaccttac gaccgtcccg ctcacgaggc	4980
tttacgcact acgtactgcg atggcttcaa tttccagcgg gagggcggat ccaactaatac	5040
aaaatatatc aaaagttaat aataatatta ttcttactta agactttttt gtcttcattt	5100
tttagtaaaa aatataaaaa aggccacctc ccgattttat cggaaggcag cctcttaaat	5160
tcagttcata atattaataa atattattca acttcagaat atttggtggc ataggcagct	5220
gccgcacca acagtccagg ctgcccgataa gtaatcaact taaccggaat cttggacatg	5280
acgcgttcaa agcgtccttt tgaaacaaag cgctgacgga aaccagattc tggcaaatgg	5340
gaagcgatac gaagaccgac accaccgcca ataacaacac tggttcgacc ctgtgccaaa	5400
gcaagatcac cagcgatagc gccaaaggctc aagcagaagc gatccaaagc ggcttcagca	5460

-continued

aggttgctt	taccttccaa	agccatctgc	cataatntaa	tatcatccag	caagctgaac	5520
ggaacgcctt	caatggcagc	cagtgtctcg	tagatattac	caagaccg	gccagaaata	5580
atgcgttcga	tagaaacgcg	gcggaacgt	tcacgtaaac	gtgccagaat	tttgtcttca	5640
agtctgtcaa	gcgagcaaaa	gtcgatatga	ccgccttcag	tttcgatgac	gaaataacgg	5700
ccttcagtc	gcaacagatg	ggcaacaccc	aagcccgctt	ccggaccaag	aatagtgata	5760
acaccatcgc	taggaagcgc	ttcatcagga	ccacaaatat	gatccagata	agaagaatcc	5820
atatgcgcaa	ccgcgtggc	aaccgcgcg	aagtcattga	tcagaacatg	cgtatcgatg	5880
tccagctttt	cattcagagt	agctggctt	aatacccaag	ggttattggt	aagttttaa	5940
acttcacat	gaaccgggc	agccatgca	atagctgcg	cacgtggcag	aggacgacc	6000
agtttttcac	cgaaacgtt	ccaagctaac	tgcaagctag	catgttctgc	cgttttaaa	6060
gtgtttctt	ctccaagaga	aagaaccga	ccattgctta	cttccgcaat	agagaaacgc	6120
gcatgcgttc	caccgatgtc	aatcgcaaca	atttcataa	taattcctt	ctgaaatcag	6180
aaggctacc	aacaggtaaa	ataagtccg	ccgctttata	ccatcgttgt	aaacaaaaag	6240
tataattggt	taagacttat	ctaaaaaga	caaaaggatt	cagccaaagc	aagtttaact	6300
acttctggga	gcccacatc	tcctcgattt	catccaggt	ccgaccttg	gtttccggca	6360
cgaagcgagc	aacaatcaag	ccacctaaga	tacttaatgc	tgcgaaaacg	agataggaga	6420
aaccgtggtt	gaaagtctga	ttcaatgctg	gagaaccatc	ggcaacctta	aacaggaagt	6480
taaccaagat	attagctaac	cattgtccg	taacagcgat	aggcatagct	gcgcccttga	6540
tggaactcgg	gaacatttct	gacagaacaa	cccagcagac	agggcccat	gacataccaa	6600
agactgcaat	ataagaagc	acagaagcca	aaggcaaac	accaccgact	ttgaaccaga	6660
aacagcagcc	taaacagcc	atcattgcag	ccataccgag	agcacccaa	ataagcagag	6720
gtttacggcc	gaagcgtca	acaacacggg	aagcaatcat	ggtgaagatg	aagttcacia	6780
caccgataga	gatggtctgc	aataatgccg	tatcagctcc	aaaacctaa	ttctggaaca	6840
tctgcggtgc	ataatacagc	acggcgtaa	taccgactaa	ctgctggaag	gcagcaacgg	6900
atacaccggc	aaaaacaacg	gtgataccaa	aagcaaacaa	acctgcgctg	cttttgtcca	6960
tggctttatc	aaagccagct	ttaatctttt	gaatcgctag	attaggatcg	gcttgcggtt	7020
ccagacgagc	aaggattttg	ctagcctcgg	aatgacgtcc	cttcatcacc	aaccaatgcg	7080
gcgtatccgg	tgcggttaac	agcagcaata	agaaggcaat	accgatcagg	ccttctgaag	7140
ccggagacca	gcaccaacca	ctggcattaa	ccaatcgat	agaaccgaaa	tgagccagta	7200
accaggtaaa	gatataaccg	gttaaagcac	ccgtcacaat	ggccatctgc	tgaccagaaa	7260
ccatctgacc	acgtttgtct	ggcggagcaa	tttcagcaat	atagggtggg	gtcaagggtg	7320
aaacgacacc	gatacctaaa	ccggcaagaa	accggaaaaa	gcaaaaaatt	tgtaaagccg	7380
aaccaccggt	tccaataat	ttttcggtta	acgcagcacc	aaaaccggcg	gcgacgaaac	7440
aaatggaact	catcaacaat	ccgccgcgac	gaccgaagcg	aataccaatc	cagccagaca	7500
gaaagaacc	ggtaacacia	ccgaccaaaa	cagcaacaac	gaccatcca	gaaaggaag	7560
ccgcagccgt	agcagacag	tgacgagggg	caataaaatg	gatatcaacc	ggtgtaccga	7620
ttgcagcgat	aaccgctgaa	tcgtaaccga	aaagcaagcc	gcctatagca	gcgattaggg	7680
ctagtcgctg	gactagacc	tgactacttt	cagaactcat	ggcgattcct	ctcctctag	7740

-continued

```

agcgtcctgc tgttgtaag attattatac cacaccttgt agataaagtc aacaactttt 7800
tgcaaaattt ttcaggaatt ttagcagagg ttgttctgga tgtagaacaa aacatctttc 7860
cgctcttggt ctgtaggat atctttcttg gaagctaggt aggcctcgag ttatggcagt 7920
tggttaaaag gaaacaaaaa gaccgttttc acacaaaacg gtctttttcg atttcttttt 7980
acagtcacag ccacttttgc accaattaag gccacgctgt catttaaact ccgtttttcc 8040
agttcaaagt caattgcctt caatgcacct tcgtagctgt ggtgagccag cggtgctggc 8100
tctcccccat ttacggataa gaatgcattt tccgagtaa taccgctggc aatacctgac 8160
attaatactt cacagtcgct ggcatcgagt acggaact taatcgaaga cgaaccacag 8220
ttaataacca aaacaaccgg aaattcattc atctcttttc tcatcctgag ttacggatta 8280
aaacagtttg tatacgatgt tcaggatggt cagcagacca atcacggtaa caaacacggt 8340
atccagacga ccacggtatt tcgccagaga cggcgttta cggatggcat acatcggcaa 8400
caggcacagc agggatgcca taatcgggtc gccatggct tcaatcaggt cgaggatgtt 8460
cgggttggtg taggcaacaa cccaggtgga gccatgatg aagatcatgc tgagagtatt 8520
cagtttacc agcgacctt tggttttgtc acctttataa ccgaacttca gaatcagacc 8580
attcaagcct tccagcgtcc ccagatagtg accgaagaaa gatttgaaga tagccacgag 8640
tgcgatgatg gaagccgat attccagtgt aatcgcgaac gttgttttgg taccggcat 8700
ggacgcaaag tggtagcca gataagaaag cactggaata ttctgcgctt tggcttccgc 8760
catggtggcc ggagacagag taaacaggca gctaaaggca aagaacatca ccaactgcaac 8820
catcagcatg ctggcacgag aatgatttg ggaacattta cgttcggtga agtcgagacc 8880
gaagtctttc tcatactctt cacgtttaga aaccacgaag gaagagacga ttggcgagaa 8940
gttaaaggag aaaacctga tggaaatccc cagccagaca gtgatcagga taccgtcatg 9000
accggttaac gacagcgaac cgaggtcaac ctggtcgata actgcagagt tccagtaagg 9060
gatcagcgac aaagaaatca gcaccaggct ggcgataaac ggccatacca ggtagctcag 9120
ggtaccgagc tcgaattcac tggccgtcgt tttacaacgt cgtgactggg aaaaccctgg 9180
cgttacccaa cttaatcggc ttgcagcaca tcccccttcc gccagctggc gtaatagcga 9240
agaggcccgc accgatcgcc cttcccaca gttgcgcagc ctgaatggcg aatggcgctt 9300
gatgcggtat tttctcctta cgcactctgt cggtatttca caccgcatat ggtgcactct 9360
cagtacaatc tgctctgatg ccgcatagtt aagccagccc cgacacccgc caacacccgc 9420
tgacgaattc 9430

```

```

<210> SEQ ID NO 2
<211> LENGTH: 9462
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: pAC21
<222> LOCATION: (1)..(9462)
<223> OTHER INFORMATION: Pplasmid used to install a slectable and
counterselectable cat-sacB cassette at the tdc locus of strain
KJ122
<220> FEATURE:
<221> NAME/KEY: pAC21
<222> LOCATION: (1)..(9462)
<223> OTHER INFORMATION: Plasmid used to install a slectable and
counterselectable cat-sacB cassette at the tdc locus of strain
KJ122
<220> FEATURE:

```


-continued

```

<221> NAME/KEY: misc_feature
<222> LOCATION: (5924)..(5924)
<223> OTHER INFORMATION: n i s a , c , g , o r t

<400> SEQUENCE: 2
gttgacagta agacgggtaa gcctgttgat gataccgctg ccttactggg tgcattagcc 60
agtctgaatg acctgtcacg ggataatccg aagtggtcag actggaaaat cagagggcag 120
gaactgctga acagcaaaaa gtcagatagc accacatagc agaccgcgca taaaacgccc 180
tgagaagccc gtgacgggct tttcttgat tatgggtagt ttccttgcat gaatccataa 240
aaggcgctg tagtgccatt taccgccatt cactgccaga gccgtgagcg cagcgaactg 300
aatgtcacga aaaagacagc gactcaggtg cctgatggtc ggagacaaaa ggaatattca 360
gcgatttgcc cgagcttgcg agggtgctac ttaagccttt agggttttaa ggtctgtttt 420
gtagaggagc aaacagcgtt tgcgacatcc ttttgaata ctgcggaact gactaaagta 480
gtgagttata cacagggctg ggatctattc tttttatctt tttttattct ttctttattc 540
tataaattat aaccacttga atataaaca aaaaaacaca caaaggctca gcggaattta 600
cagagggctc agcagaattt acaagttttc cagcaaaggt ctagcagaat ttacagatac 660
ccacaactca aaggaaaagg actagtaatt atcattgact agcccatctc aattggtata 720
gtgattaata tcacctagac caattgagat gtatgtctga attagttgtt ttcaaagcaa 780
atgaactagc gattagtgcg tatgacttaa cggagcatga aaccaagcta attttatgct 840
gtgtggcact actcaacccc acgattgaaa acctacaag gaaagaacgg acggtatcgt 900
tcacttataa ccaatacgtc cagatgatga acatcagtag ggaaaatgct tatggtgtat 960
tagctaaagc aaccagagag ctgatgacga gaactgtgga aatcaggaat cctttgggta 1020
aaggctttga gattttccag tggacaaact atgccaaagt ctcaagcga aaattagaat 1080
tagtttttag tgaagagata ttgccttatc tttccagtt aaaaaattc ataaaatata 1140
atctggaaca tgtaagtct tttgaaaaca aatactctat gaggatttat gagtggttat 1200
taaaagaact aacacaaaag aaaactcaca aggcaatat agagattagc cttgatgaat 1260
ttaagtcat gttaatgctt gaaaataact accatgagtt taaaaggctt aaccaatggg 1320
ttttgaaacc aataagtaaa gatttaaca cttacagcaa tatgaaattg gtggttgata 1380
agcgaggccg cccgactgat acgttgattt tccaagttga actagataga caaatggatc 1440
tcgtaaccga acttgagaac aaccagataa aatgaatgg tgacaaaata ccaacaacca 1500
ttacatcaga ttctaccta cgtaacggac taagaaaaac actacacgat gctttaactg 1560
caaaaattca gctcaccagt tttgaggcaa aatttttgag tgacatgcaa agtaagcatg 1620
atctcaatgg ttcgttctca tggctcacgc aaaaacaacg aaccacacta gagaacatac 1680
tggttaaata cggaaggatc tgaggttctt atggctcttg tatctatcag tgaagcatca 1740
agactaaca acaaaagtag aacaactggt caccgttaca tatcaaaggg aaaactgtcc 1800
atatgcacag atgaaaacgg tgtaaaaaag atagatacat cagagctttt acgagttttt 1860
ggtgcattta aagctgttca ccatgaacag atcgacaatg taacagatga acagcatgta 1920
acacctaata gaacaggtga aaccagtaaa acaaagcaac tagaacatga aattgaacac 1980
ctgagacaac ttgttacagc tcaacagtca cacatagaca gcctgaaaca ggcgatgctg 2040
cttatcgaat caaagctgcc gacaacacgg gagccagtga cgctcccgt ggggaaaaaa 2100

```


-continued

tcatggcaat	tctggaagaa	atagcgcttt	cagccggcaa	acctgaagcc	ggatctgcca	2160
ttctgataac	aaactagcaa	caccagaaca	gcccgtttgc	gggcagcaaa	acccgttatg	2220
cttgtaaacc	gttttgtgaa	aaaatthtta	aaataaaaaa	ggggacctct	agggtcccca	2280
attaattagt	aatataatct	attaaaggtc	attcaaaagg	tcatccaccg	gatcaattcc	2340
cctgctcgcg	caggctgggt	gccaagctct	cgggtaacat	caaggcccga	tccttgagac	2400
ccttgccctc	ccgcacgatg	atcgtgccgt	gatcgaaatc	cagatccttg	accgcagtt	2460
gcaaacctc	actgatccgc	atgccggttc	catacagaag	ctgggcgaac	aaacgatgct	2520
cgcttccag	aaaaccgagg	atgcgaacca	cttcatccgg	ggtcagcacc	accggcaagc	2580
gccgcgacgg	ccgaggtctt	ccgatctcct	gaagccaggg	cagatccgtg	cacagcacct	2640
tgccgtagaa	gaacagcaag	gccgccaatg	cctgacgatg	cgtggagacc	gaaaccttgc	2700
gctcgttcgc	cagccaggac	agaaatgcct	cgacttcgct	gctgcccaag	gttgccgggt	2760
gacgcacacc	gtggaaacgg	atgaaggcac	gaaccagtg	gacataagcc	tgttcggttc	2820
gtaagctgta	atgcaagtag	cgtatgcgct	cacgcaactg	gtccagaacc	ttgaccgaac	2880
gcagcggtag	taacggcgca	gtggcggttt	tcatggcttg	ttatgactgt	ttttttgggg	2940
tacagtctat	gcctcgggca	tccaagcagc	aagcgcgcta	cgccgtgggt	cgatgtttga	3000
tgttatggag	cagcaacgat	gttacgcagc	agggcagtcg	ccctaaaaca	aagttaaaca	3060
tcatgagggg	agcggtagtc	gccgaagtat	cgactcaact	atcagaggta	gttggcgtca	3120
tcgagcgcca	tctcgaaccg	acgttgctgg	ccgtacattt	gtacggctcc	gcagtggatg	3180
gcggcctgaa	gccacacagt	gatattgatt	tgctggttac	ggtgaccgta	aggcttgatg	3240
aaacaacgcg	gcgagctttg	atcaacgacc	ttttgaaaac	ttcggcttcc	cctggagaga	3300
gcgagattct	ccgcgctgta	gaagtcacca	ttgttgtagc	cgacgacatc	attccgtggc	3360
gttatccagc	taagcgcgaa	ctgcaatttg	gagaatggca	gcgcaatgac	attcttgacg	3420
gtatcttcga	gccagccacg	atcgacattg	atctggctat	cttgctgaca	aaagcaagag	3480
aacatagcgt	tgccttggtg	ggtccagcgg	cggaggaact	ctttgatccg	gttcctgaac	3540
aggatctatt	tgaggcgcta	aatgaaacct	taacgctatg	gaactcgccg	cccgactggg	3600
ctggcgatga	gcgaaatgta	gtgcttacgt	tgtcccgcac	ttggtacagc	gcagtaaccg	3660
gcaaaaatcgc	gccgaaggat	gtcgctgccc	actgggcaat	ggagcgccctg	ccggcccagt	3720
atcagcccgt	catacttgaa	gctagacagg	cttatcttgg	acaagaagaa	gatcgcttgg	3780
cctcgcgcgc	agatcagttg	gaagaatttg	tccactacgt	gaaaggcgag	atcaccaagg	3840
tagtcggcaa	ataatgtcta	acaattcggt	caagccgacg	ccgcttcgcg	gcgcggttta	3900
actcaagcgt	tagatgcact	aagcacataa	ttgctcacag	ccaaactatc	aggtaagtc	3960
tgtttttatt	atthtttaagc	gtgcataata	agccctacac	aaattgggag	atatatcatg	4020
aaaggctggc	tttttcttgt	tatcgcaata	gttggcgaag	taatcgcaac	atccgcatta	4080
aaatctagcg	aggcctttac	taagctgatc	cggtaggatg	ccttttgaat	gacctttaat	4140
agattatatt	actaattaat	tggggaccct	agaggcccc	ttttttattt	taaaaatttt	4200
ttcacaaaac	ggtttacaag	catacgttgg	ccgattcatt	aatgcagctg	gcacgacagg	4260
tttcccgact	ggaaagcggg	cagtgagcgc	aacgcaatta	atgtgagtta	gctcactcat	4320
taggcacccc	aggctttaca	ctttatgctt	ccggctcgta	tgttgtgtgg	aattgtgagc	4380

-continued

ggataacaat	ttcacacagg	aaacagctat	gaccatgatt	acgccaaagct	tgcattgctg	4440
caggtcgact	ctagaggatc	ccccccgccc	ccgacagagt	aataggtttt	acttaatagc	4500
tcttcctgtc	ccttccaggc	agtgatccgc	attccgttct	catggcgagg	caacatttcg	4560
ggatggaaga	taatgttctt	tgctacagga	aatcaacaa	tatgcgacc	agatgccact	4620
ggcagccgcc	cgctgcgct	tactaactct	ataaatgcag	ggatctcatc	aatgacaaca	4680
tctcgccgac	tgtttctg	cagtcccatg	atgatggcga	catccgtggc	atggcctttg	4740
cccgtcagtg	acaacgacc	gtacagatcg	accacaatat	ggctcgtcgc	ggtaataag	4800
ccgctacttt	ccagccgatc	aataaaactt	tttccggcat	tcattggccc	cacggtatgc	4860
gaactggagg	gaccaatccc	aattttgaaa	atctcgaatg	cactaatcat	gtgacggaag	4920
atcacttcgc	agaataaata	aatcctgggtg	tccctgttga	taccgggaag	ccctgggcca	4980
acttttggcg	aaaatgagac	gttgatccgc	acgtaagagg	ttccaacttt	caccataatg	5040
aaataagatc	actaccgggc	gtattttttg	agttatcgag	attttcagga	gctaaggaag	5100
ctaaaatgga	gaaaaaatc	actggatata	ccaccgttga	tatatcccaa	tggcatcgta	5160
aagaacattt	tgaggcattt	cagtcagttg	ctcaatgtac	ctataaccag	accgttcagc	5220
tgatattac	ggccttttta	aagaccgtaa	agaaaaataa	gcacaagttt	tatccggcct	5280
ttattcacat	tcttgcccgc	ctgatgaatg	ctcatccgga	attccgtatg	gcaatgaaag	5340
acggtgagct	ggatgatgg	gatagtgttc	acccttggtt	caccgttttc	catgagcaaa	5400
ctgaaacggt	ttcatcgctc	tggagtgaat	accacgacga	tttccggcag	tttctacaca	5460
tatattcgca	agatgtggcg	tgttacgggtg	aaaaactggc	ctatttcctt	aaaggggtta	5520
ttgagaatat	gttttctg	tcagccaatc	cctgggtgag	tttcaccagt	tttgatttaa	5580
acgtggccaa	tatggacaac	ttcttccccc	ccgttttcac	catgggcaaa	tattatacgc	5640
aaggcgacaa	ggtgctgatg	ccgctggcga	ttcaggttca	tcattgccgt	tgtgatggct	5700
tccatgctgg	cagaatgctt	aatgaattac	aacagtactg	cgatgagtgg	cagggcgggg	5760
cgtaattttt	ttaaggcagt	tattgggtgcc	cttaaacgcc	tggtgctacg	cctgaataag	5820
tgataataag	cggatgaatg	gcagaaatc	gaaagcaaat	tcgaccgggt	cgctcggttc	5880
ggcaggggtc	gttaaatagc	cgcttatgct	tattgctggt	ttantcggt	cccggggatc	5940
gcgcccgccg	accggatccc	atcacatata	cctgccgttc	actattattt	agtgaaatga	6000
gatattatga	tattttctga	attgtgatta	aaaaggcaac	tttatgcca	tgcaacagaa	6060
actataaaaa	atacagagaa	tgaaaagaaa	cagatagatt	tttagttct	ttaggcccg	6120
agtctgcaaa	tccttttatg	atcttctatc	aaacaaaaga	ggaaaataga	ccagttgcaa	6180
tccaaacgag	agtctaatag	aatgaggtcg	aaaagtaaat	cgcgcggtt	tgttactgat	6240
aaagcaggca	agacctaaaa	tgtgtaaagg	gcaaagtgt	tactttggcg	tcaccctta	6300
catattttag	gtctttttt	attgtgcgta	actaacttgc	catcttcaaa	caggagggt	6360
ggaagaagca	gaccgctaac	acagtacata	aaaaaggaga	catgaacgat	gaacatcaaa	6420
aagtttgcaa	aacaagcaac	agtattaacc	tttactaccg	cactgctggc	aggaggcgca	6480
actcaagcgt	ttgcgaaaga	aacgaaccaa	aagccatata	aggaaacata	cggcatttcc	6540
catattacac	gccatgatat	gctgcaaatc	cctgaacagc	aaaaaatga	aaaatatcaa	6600
gttctgaat	tcgattcgtc	cacaattaa	aatatctctt	ctgcaaaagg	cctggacgtt	6660

-continued

tgggacagct	ggccattaca	aaacgctgac	ggcactgtcg	caaactatca	cggtaccac	6720
atcgtctttg	cattagccgg	agatcctaaa	aatgcggatg	acacatcgat	ttacatgttc	6780
tatcaaaaag	tggcgaaac	ttctattgac	agctggaaaa	acgctggccg	cgctctttaa	6840
gacagcgaca	aattcgatgc	aatgattct	atcctaaaag	accaaacaca	agaatggcca	6900
ggttcagcca	catttacatc	tgacggaaaa	atccgtttat	tctacactga	tttctccggt	6960
aaacattacg	gcaaacaac	actgacaact	gcacaagtta	acgtatcagc	atcagacagc	7020
tctttgaaca	tcaacggtgt	agaggattat	aatcaatct	ttgacggtga	cggaaaaacg	7080
tatcaaaatg	tacagcagtt	catcgatgaa	ggcaactaca	gctcaggcga	caaccatacg	7140
ctgagagatc	ctcactacgt	agaagataaa	ggccacaaat	acttagtatt	tgaagcaaac	7200
actggaactg	aagatggcta	ccaaggcgaa	gaatctttat	ttaacaaagc	atactatggc	7260
aaaagcacat	cattcttccg	tcaagaaagt	caaaaacttc	tgcaaagcga	taaaaaacgc	7320
acggctgagt	tagcaaacgg	cgctctcggg	atgattgagc	taaacgatga	ttacacactg	7380
aaaaaagtga	tgaaccgct	gattgcatct	aacacagtaa	cagatgaaat	tgaacgcgcg	7440
aacgtcttta	aatgaacgg	caaatggtac	ctgttcactg	actcccgcgg	atcaaaaatg	7500
acgattgacg	gcattacgtc	taacgatatt	tacatgcttg	gttatgtttc	taattcttta	7560
actggcccat	acaagccgct	gaacaaaact	ggccttgtgt	taaaaatgga	tcttgatcct	7620
aacgatgtaa	cctttactta	ctcacacttc	gctgtacctc	aagcgaaagg	aaacaatgtc	7680
gtgattacaa	gctatatgac	aaacagagga	ttctacgcag	acaacaatc	aacgtttgcg	7740
ccgagcttcc	tgctgaacat	caaaggcaag	aaaacatctg	ttgtcaaaga	cagcatcctt	7800
gaacaaggac	aattaacagt	taacaataaa	aaacgcaaaa	gaaaatgcca	ataticctatt	7860
ggcattttct	tttatttctt	ccatttaaat	ggatgcatgc	gctagcggag	tgtatactgg	7920
cttactatgt	tggcactgat	gaggggtgca	gtgaagtgct	tcagcctcgt	gagcgggacg	7980
gtcgtaaggt	cgttccgctc	cacttcactg	aacggcaatc	cgaggggtgtg	gatccaatta	8040
aggccacgct	gtcattttaa	ttccgttttt	ccagttcaaa	tgcaattgcc	ttcaatgcac	8100
cttcgtagct	gtggtgagcc	agcgggtgctg	gctctcccc	atttacggat	aagaatgcat	8160
tttccgagtt	aataccgctg	gcaatacctg	acattaatac	ttcacagtcg	ctggcatcga	8220
gtacggaaaa	cttaatcgaa	gacgaaccac	agttaataac	caaaacaacc	ggaaattcat	8280
tcatctcttt	tctcatcctg	agttacggat	taaaacagtt	tgtatacgat	gttcaggatg	8340
gtcagcagac	caatcacggt	aacaaacacg	ttatccagac	gaccacggta	tttcgccaga	8400
gacggcgctt	tacggatggc	atacatcggc	aacaggcaca	gcagggatgc	gataatcggg	8460
gcgcccattg	cttcaatcag	gtcgaggatg	ttcgggttgg	cgtaggcaac	aaccaggtg	8520
gagcccatga	tgaagatcat	gctgagagta	ttcagtttac	ccagcgcac	tttggttttg	8580
tcacctttat	aaccgaactt	cagaatcaga	ccattcaagc	cttcagcgt	ccccagatag	8640
tgaccgaaga	aagatttgaa	gatagccacg	agtgcgatga	tggaagccgc	atattccagt	8700
gtaatcgca	acgttgtttt	ggtaccggtc	atggacgcaa	agtggttagc	cagataagaa	8760
agcactggaa	tattctgcgc	tttggttcc	gccatgttgg	ccggagacag	agtaaacagg	8820
cagctaaagg	caaagaacat	caccactgca	accatcagca	tgctggcacg	agaaatgatt	8880
tgggaacatt	tacgttcggt	gaagtcgca	ccgaagtctt	tctcactc	ttcacgttta	8940

-continued

```

gaaaccacga aggaagagac gattggcgag aagttaaagg agaaaacat gatggaaatc 9000
cccagccaga cagtgatcag gataccgtca tgaccgggta acgacagcga accgaggtca 9060
acctggtcga taactgcaga gttccagtaa gggatcagcg acaaagaaat cagcaccagg 9120
ctggcgataa acggccatac caggtagctc agggtagcga gctcgaattc actggccgctc 9180
gttttacaac gtcgtgactg ggaaaaccct ggcgttacc aacttaatcg ccttgagca 9240
catccccctt tcgccagctg gcgtaatagc gaagaggccc gcaccgatcg cccttcccaa 9300
cagttgcgca gcctgaatgg cgaatggcgc ctgatgcggg attttctcct tacgcatctg 9360
tgcggtatctt cacaccgat atggtgcact ctcagtacaa tctgctctga tgccgatag 9420
ttaagccagc cccgacacc gccaacacc gctgacgaat tc 9462

```

```

<210> SEQ ID NO 3
<211> LENGTH: 1091
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli
<220> FEATURE:
<221> NAME/KEY: ptsHI
<222> LOCATION: (1)..(1091)
<223> OTHER INFORMATION: Sequence surrounding the deletion of ptsHI. The
fusion junction is in between the nucleotide positions 425 and
426.

```

```

<400> SEQUENCE: 3
gcaacagtaa tgccagcttg ttaaaaatgc gtaaaaaagc accttttttag gtgctttttt 60
gtggcctgct tcaaactttc gcccctcctg gcattgattc agcctgtcgg aactgggtatt 120
taaccagact aattatcttg atgcgcgaaa ttaatcgta caggaaaagc caaagctgaa 180
tcgattttat gatttggtc aattcttctt ttagcggcat aatgtttaat gacgtacgaa 240
acgtcagcgg tcaacaccgg ccagcaatgg actgtattgc gctcttcgtg cgtcgcgtct 300
gttaaaaact ggcgctaaca atacaggcta aagtgaacc gccaggctag actttagttc 360
cacaacacta aacctataag ttggggaaat acaatgttcc agcaagaagt taccattacc 420
gctccacaat ctgctaacc acgagatgcg gcccaattta ctgcttagga gaagatcatg 480
ggtttgctcg ataaactgaa atctctgggt tccgacgaca agaaggatac cggaactatt 540
gagatcattg ctccgctctc tggcgagatc gtcaatatcg aagacgtgcc ggatgtcgtt 600
tttgccgaaa aaatcgttgg tgatgggtatt gctatcaaac caacgggtaa caaatggtc 660
gcgccagtag acggcaccat tggtaaaatc tttgaaacca accacgcatt ctctatcgaa 720
tctgatagcg gcgttgaact gttcgtccac ttcggtatcg acaccgttga actgaaaggc 780
gaaggcttca agcgtattgc tgaagaaggc cagcgcgtga aagttggcga tactgtcatt 840
gaatttgatc tgccgctgct ggaagagaaa gccaaagtcta ccctgactcc ggttggtatc 900
tccaacatgg acgaaatcaa agaactgatc aaactgtccg gtagecgtaac cgtgggtgaa 960
accccggtta tccgcatcaa gaagtaattc tgccgcagtg aaaaatggcg cccatcggcg 1020
ccattttttt atgcttccgc cagcggcggc aaaatcaatt catcgtctct atgctgctgg 1080
gtgtagcgca t 1091

```

```

<210> SEQ ID NO 4
<211> LENGTH: 66
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli
<220> FEATURE:

```

-continued

<221> NAME/KEY: galP
 <222> LOCATION: (1)..(66)
 <223> OTHER INFORMATION: Sequence surrounding the deletion of galP gene.
 The fusion junction is in between the nucleotide positions 48 and
 49.

<400> SEQUENCE: 4

atgcctgacg ctaaaaaaca ggggcggtca aacaaggcaa tgacgtttat aggcgctcac 60
 gattaa 66

<210> SEQ ID NO 5
 <211> LENGTH: 8446
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: pSS2
 <222> LOCATION: (1)..(8446)
 <223> OTHER INFORMATION: Plasmid used to install a glf cassette at the
 tdc locus of bacterial strain KJ122.

<400> SEQUENCE: 5

gttgacagta agacgggtaa gcctgttgat gataccgctg cttactggg tgcattagcc 60
 agtctgaatg acctgtcacg ggataatccg aagtggtcag actggaaaat cagagggcag 120
 gaactgctga acagcaaaaa gtcagatagc accacatagc agaccgcca taaaacgccc 180
 tgagaagccc gtgacgggct tttcttgat tatgggtagt ttccttgcg gaatccataa 240
 aaggcgctg tagtgccatt taccgccatt cactgccaga gccgtgagcg cagcgaactg 300
 aatgtcacga aaaagacagc gactcaggtg cctgatggtc ggagacaaaa ggaatattca 360
 gcgatttgcc cgagcttgcg aggggtgctac ttaagccttt agggttttaa ggtctgtttt 420
 gtagaggagc aaacagcgtt tgcgacatcc ttttgaata ctgcggaact gactaaagta 480
 gtgagttata cacagggctg ggatctattc tttttatctt tttttattct ttctttattc 540
 tataaattat aaccacttga atataaaca aaaaaacaca caaaggctca gcggaattta 600
 cagaggtct agcagaattt acaagttttc cagcaaaggc ctagcagaat ttacagatac 660
 ccacaactca aaggaaaagg actagtaatt atcattgact agcccatctc aattggtata 720
 gtgattaata tcacctagac caattgagat gtatgtctga attagttgtt ttcaaagcaa 780
 atgaactagc gattagtcgc tatgacttaa cggagcatga aaccaagcta attttatgct 840
 gtgtggcact actcaacccc acgattgaaa acctacaag gaaagaacgg acggtatcgt 900
 tcacttataa ccaatacgtc cagatgatga acatcagtag ggaaaatgct tatgggtgat 960
 tagctaaagc aaccagagag ctgatgacga gaactgtgga aatcaggaat cctttggtta 1020
 aaggctttga gattttccag tggacaaaact atgccaagtt ctcaagcga aaattagaat 1080
 tagtttttag tgaagagata ttgccttacc tttccagtt aaaaaattc ataaaatata 1140
 atctggaaca tgttaagtct tttgaaaaca aatactctat gaggatttat gagggttat 1200
 taaaagaact aacacaaaag aaaactcaca agcacaatat agagattagc cttgatgaat 1260
 ttaagtcat gtaaatgctt gaaaataact accatgagtt taaaaggctt aaccaatggg 1320
 ttttgaaacc aataagtaaa gatttaaca cttacagcaa tatgaaattg gtggttgata 1380
 agcgaggccg cccgactgat acggtgattt tccaagttga actagataga caaatggatc 1440
 tcgtaaccga acttgagaac aaccagataa aatgaatgg tgacaaaata ccaacaacca 1500
 ttacatcaga ttctaccta cgtaacggac taagaaaac actacacgat gctttaactg 1560

-continued

caaaaattca gctcaccagt tttgaggcaa aatTTTTgag tgacatgcaa agtaagcatg	1620
atctcaatgg ttcgTtctca tggctcacgc aaaaacaacg aaccacacta gagaacatac	1680
tggctaaata cggaaggatc tgaggTtctt atggctcttg tatctatcag tgaagcatca	1740
agactaacia acaaaagtag aacaactgTt caccgTtaga tatcaaaggg aaaactgtcc	1800
atatgcacag atgaaaacgg tgtaaaaaag atagatacat cagagctTtt acgagTtttt	1860
ggtgcattta aagctgttca ccatgaacag atcgacaatg taacagatga acagcatgta	1920
acacctaata gaacaggtga aaccagtaaa acaaagcaac tagaacatga aattgaacac	1980
ctgagacaac ttgttacagc tcaacagtca cacatagaca gcctgaaaca ggcgatgctg	2040
cttatogaat caaagctgcc gacaacacgg gagccagtga cgctcccgT ggggaaaaaa	2100
tcatggcaat tctggaagaa atagcgctTt cagccggcaa acctgaagcc ggatctgcga	2160
ttctgataac aaactagcaa caccagaaca gcccgTttgc gggcagcaaa acccgTtatg	2220
cttgtaaacc gTtttTtgaa aaaatTttta aaataaaaaa ggggacctct agggTcccca	2280
attaattagt aatataatct attaaaggTc attcaaaagg tcatccaccg gatcaattcc	2340
cctgctcgcg caggctgggt gccaaagctct cgggtaacat caaggcccgga tccTtgagc	2400
cctgcccctc ccgcacgatg atcgTgccgT gatcgaaatc cagatcctTg acccgcagTt	2460
gcaaacctc actgatccgc atgcccgTtc catacagaag ctgggcgaac aaacgatgct	2520
cgcttccag aaaaccgagg atgcgaacca ctTcatccgg ggtcagcacc accggcaagc	2580
gccgcgacgg ccgaggtctt ccgatctcct gaagccaggg cagatccgTg cacagcacct	2640
tgccgtagaa gaacagcaag gccgccaatg cctgacgatg cgtggagacc gaaacctTgc	2700
gctcgTtcgc cagccaggac agaaatgcct cgacttcgct gctgcccag gTtgccgggt	2760
gacgcacacc gtggaaacgg atgaaggcac gaaccagTg gacataagcc TgttcggTtc	2820
gtaagctgta atgcaagtag cgtatgcgct cacgcaactg gtccagaacc ttgaccgaac	2880
gcagcggtgg taacggcgca gtggcggtTt tcatggctTg ttatgactgt tTttTgggg	2940
tacagtctat gcctcgggca tccaagcagc aagcgcgTta cgccgTgggt cgatgTttga	3000
Tgttatggag cagcaacgat gttacgcagc agggcagTcg ccctaaaaca aagTtaaaca	3060
tcatgagggga agcggTgatc gccgaagtat cgactcaact atcagaggta gTtggcgTca	3120
tcgagcgcca tctegaaccg acgTtTgctgg ccgtacattt gtacggctcc gcagTggatg	3180
gcggcctgaa gccacacagt gatattgatt TgctggTtac ggtgaccgta aggctTgatg	3240
aaacaacgcg gcgagctTtg atcaacgacc tTttgaaac ttcggctTtc cctggagaga	3300
gcgagattct ccgcgctgta gaagTcacca ttgtTgtgca cgacgacatc attccgTggc	3360
gttatccagc taagcgcgaa ctgcaattTg gagaatggca gcgcaatgac attctTgcag	3420
gtatcttoga gccagccagc atcgacattg atctggctat ctTgctgaca aaagcaagag	3480
aacatagcgt TgcctTggta ggtccagcgg cggaggaact ctTtgatccg gTtccTgaac	3540
aggatctatt tgaggcgcta aatgaaacct taacgctatg gaactcgccg cccgactggg	3600
ctggcgatga gcgaaatgta gtgctTacgt Tgtcccgat ttggtacagc gcagTaacgg	3660
gcaaaatcgc gccgaaggat gTcgctgccg actgggcaat ggagcgctg ccggcccagT	3720
atcagcccgt catactTgaa gctagacagg ctTatctTgg acaagaagaa gatcgctTgg	3780
cctcgcgcgc agatcagTtg gaagaattTg tccactacgt gaaaggcgag atcaccaagg	3840

-continued

tagtcggcaa	ataatgtcta	acaattcggt	caagccgacg	ccgcttcgcg	gcgaggctta	3900
actcaagcgt	tagatgcact	aagcacataa	ttgctcacag	ccaaactatc	aggtaagtc	3960
tgcttttatt	atthtttaagc	gtgcataata	agccctacac	aaattgggag	atatatcatg	4020
aaaggctggc	tttttcttgt	tatcgcaata	gttggcgaag	taatcgcaac	atccgcatta	4080
aaatctagcg	agggttttac	taagctgatc	cggtggatga	ccttttgaat	gacctttaat	4140
agattatatt	actaattaat	tggggaccct	agaggtcccc	ttttttatth	taaaaattht	4200
ttcacaaaa	ggtttacaag	catacgttgg	ccgattcatt	aatgcagctg	gcacgacagg	4260
tttcccgact	ggaaagcggg	cagtgagcgc	aacgcaatta	atgtgagtta	gctcactcat	4320
taggcacccc	aggctttaca	ctttatgctt	ccggctcgta	tgthgtgtgg	aattgtgagc	4380
ggataacaat	ttcacacagg	aaacagctat	gaccatgatt	acgccaagct	tgcatgcctg	4440
caggctgact	ctagaggatc	ccccccgccc	ccgacagagt	aataggthtt	acttaatagc	4500
tcttcctgtc	ccttccaggc	agtgatccgc	attccgttct	catggcgagg	caacatthtcg	4560
ggatggaaga	taatgttctt	tgctacagga	aatcaacaa	tatgcgacc	agatgccact	4620
ggcagccgcc	cgctgcgcgt	tactaactct	ataaatgcag	ggatctcatc	aatgacaaca	4680
tcttcgggac	tgthtctgtc	cagtcccatt	atgatggcga	catccgtggc	atggcctthg	4740
cccgtcagtg	acaacgacc	gtacagatcg	accacaatat	ggctcgtcgc	ggthtaataag	4800
ccgctactth	ccagccgatc	aataaaaact	tttccggcat	tcattggccc	cacggtatgc	4860
gaactggagg	gaccaatccc	aatthtgaaa	atategaatg	cactaatcat	atccacaccc	4920
tcggattgcc	gthcagtgaa	gtggagcggg	acgaccttac	gaccgtccc	ctcacgaggc	4980
thtacgcact	acgtactgcg	atggctthca	thtccagcgg	gagggcggat	ccactaatac	5040
aaaatatatc	aaaagthaat	aataatatta	thcttactta	agactththt	gtthtcattt	5100
thtagtaaaa	aatataaaaa	aggccacctc	ccgattthtat	cggaaggcag	cctcttaaat	5160
tcagthcata	atathtaaaa	atathattca	acttcagatt	atthgttggc	ataggcagcg	5220
ctccgcccgc	thtataccat	cgthgtaaac	aaaaagtata	atthgttaag	acttatctaa	5280
aaaagacaaa	aggattcagc	caaagcaagt	thtaactact	ctgggagcgc	cacatctcct	5340
cgattthcatc	caggctccga	cctthggtht	ccggcacgaa	gcgagcaaca	atcaagccac	5400
ctaagatact	taatgtctgc	aaaacgagat	aggagaaaacc	gtggttgaaa	gtctgattca	5460
atgctggaga	accatcggca	accttaacaa	ggaagthaac	caagatatta	gctaaccatt	5520
gtccggtaac	agcgataggc	atagctgcgc	ccttgatgga	actcgggaac	atthctgaca	5580
gaacaaccca	gcagacaggg	ccccatgaca	taccaaagac	tgcaatataa	agaagcacag	5640
aagccaaaagg	caaaacacca	ccgactthtga	accagaaaaca	gcagcctaaa	acagccatca	5700
thgcagccat	accgagagca	ccccaaataa	gcagaggtht	acggccgaag	cggtcaacaa	5760
cacgggaagc	aatcatggtg	aagatgaagt	tcacaacacc	gatagagatg	gtctgcaata	5820
atgccgtatc	agctccaaaa	cctaaattct	ggaacatctg	cggtgcataa	tacagcacgg	5880
cgthaatacc	gactaactgc	tggaaggcag	caacggatac	accggcaaaa	acaacggthg	5940
tacaaaaagc	aaacaaacct	gcgctgctth	tgthccatggc	thtatcaaac	ccagctthta	6000
tctthtgaat	cgthcagatta	ggatcggctt	gcggtthccag	acgagcaagg	atthtgctag	6060
cctcggaatg	acgthccctc	atcaccaacc	aatgcggcgt	atccggtgcg	gthaacagca	6120

-continued

gcaataagaa	ggcaataaccg	atcaggcctt	ctgaagccgg	agaccagcac	caaccactgg	6180
cattaacca	atcgatagaa	ccgaaatgag	ccagtaacca	ggtaaagata	taaccgggta	6240
aagcaccctg	cacaatggcc	atctgctgac	cagaaacat	ctgaccacgt	ttgtctggcg	6300
gagcaatttc	agcaatatag	gttgggggtca	aggttgaaac	gacaccgata	cctaaaccgg	6360
caagaaaccg	gaaaaagcaa	aaaatttgta	aagccgaacc	accggttcca	aataatTTTT	6420
cggttaacgc	agcaccaaaa	ccggcggcga	cgaaacaaat	ggaactcatc	aacaatccgc	6480
cgcgacgacc	gaagcgaata	ccaatccagc	cagacagcaa	agaaccggta	acacaaccga	6540
ccaaaacagc	aacaacgacc	atcccagaaa	gggaagccgc	agccgtagca	gacagggtgac	6600
gaggggcaat	aaaatggata	tcaaccgggtg	taccgattgc	agcgataacc	gctgaatcgt	6660
aaccgaaaag	caagccgcct	atagcagcga	ttagggctag	tcgcgtgact	agaccctgac	6720
tactttcaga	actcatggcg	attcctctcc	ctctagagcg	tctgctgtt	gttaagatta	6780
ttataccaca	ccttgtagat	aaagtcaaca	actttttgca	aaatTTTTca	ggaatTTtag	6840
cagaggttgt	tctggatgta	gaacaaaaca	tctttccgct	cttgtgctgt	taggatatct	6900
ttcttggaag	ctaggtagc	ctcgagttat	ggcagttggt	taaaaggaaa	caaaaagacc	6960
gttttcacac	aaaacggtct	ttttcgattt	ctttttacag	tcacagccac	ttttgcacca	7020
attaaggcca	cgctgtcatt	taaactccgt	ttttccagtt	caaagtcaat	tgcttcaat	7080
gcaccttcgt	agctgtggtg	agccagcggg	gctggctctc	ccccatttac	ggataagaat	7140
gcattttccg	agttaatacc	gtcggcaata	cctgacatta	atacttcaca	gtcgctggca	7200
tcgagtacgg	aaaacttaat	cgaagacgaa	ccacagttaa	taacccaaac	aaccggaaat	7260
tcatatcat	cttttctcat	cctgagttac	ggattaaaac	agtttgata	cgatgttcag	7320
gatggtcagc	agaccaatca	cggtaacaaa	cacgttatcc	agacgaccac	ggtatTTcgc	7380
cagagacggc	gctttacgga	tggcatacat	cggcaacagg	cacagcaggg	atgcgataat	7440
cggtgcgccc	atggcttcaa	tcaggctgag	gatgttcggg	ttggcgtagg	caacaacca	7500
ggtggagccc	atgatgaaga	tcatgctgag	agtattcagt	ttaccagcg	acactttggt	7560
tttgtcacct	ttataaccga	acttcagaat	cagaccatc	aagccttcca	gcgtccccag	7620
atagtaccg	aagaaagatt	tgaagatagc	cacgagtgcg	atgatggaag	ccgcatattc	7680
cagtgtaatc	gcgaacgttg	ttttggtagc	ggatcatggc	gcaaagtggg	tagccagata	7740
agaaagcact	ggaatattct	gcgctttggc	ttccgccatg	ttggccggag	acagagtaaa	7800
caggcagcta	aaggcaaaga	acatcaccac	tgcaaccatc	agcatgctgg	cacgagaaat	7860
gatttgggaa	catttacgtt	cggtgaagtc	gcgaccgaag	tctttctcat	actcttcacg	7920
tttagaaacc	acgaaggaag	agacgattgg	cgagaagtta	aaggagaaaa	ccatgatgga	7980
aatccccagc	cagacagtga	tcaggatacc	gtcatgaccg	gttaacgaca	gcgaaccgag	8040
gtcaacctgg	tcgataactg	cagagttcca	gtaagggatc	agcgacaaag	aaatcagcac	8100
caggctggcg	ataaacggcc	ataccaggta	gctcagggta	ccgagctcga	attcactggc	8160
cgctgTTTTa	caacgtcgtg	actgggaaaa	cctggcggtt	acccaactta	atcgcttgc	8220
agcacatccc	cctttcgcca	gctggcgtaa	tagcgaagag	gcccgcaccg	atcgcccttc	8280
ccaacagttg	cgcagcctga	atggcgaatg	gcgcctgatg	cggtatTTtc	tccttacgca	8340
tctgtgcggg	atttcacacc	gcatatggtg	cactctcagt	acaatctgct	ctgatgccgc	8400

-continued

 atagttaagc cagccccgac acccgccaac acccgctgac gaattc 8446

<210> SEQ ID NO 6
 <211> LENGTH: 31
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR Primer BY249 for amplifying and cloning the
 crr gene of YSS41, together with BY249.

<400> SEQUENCE: 6

gggcaacagt aatgccagct tgtaaaaaat g 31

<210> SEQ ID NO 7
 <211> LENGTH: 25
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR Primer BY250 for amplifying and cloning the
 crr gene of YSS41 together with BY249.

<400> SEQUENCE: 7

gggcgctaca cccagcagca tgaga 25

<210> SEQ ID NO 8
 <211> LENGTH: 8937
 <212> TYPE: DNA
 <213> ORGANISM: Escherichia coli
 <220> FEATURE:
 <221> NAME/KEY: pMH68
 <222> LOCATION: (1)..(8937)
 <223> OTHER INFORMATION: Plasmid used to install a second copy of teh
 crr gene from YSS41 at the pflD locus.

<400> SEQUENCE: 8

 gttgacagta agacgggtaa gcctgttgat gataccgctg ccttactggg tgcattagcc 60
 agtctgaatg acctgtcacg ggataatccg aagtggtcag actggaaaat cagagggcag 120
 gaactgctga acagcaaaaa gtcagatagc accacatagc agaccgcca taaaacgccc 180
 tgagaagccc gtgacgggct tttcttgat tatgggtagt ttccttgcat gaatccataa 240
 aaggcgectg tagtgccatt taccgccatt cactgccaga gccgtgagcg cagcgaactg 300
 aatgtcacga aaaagacagc gactcaggtg cctgatggtc ggagacaaaa ggaatattca 360
 gcgatttgcc cgagcttgcg aggggtgctac ttaagccttt agggttttaa ggtctgtttt 420
 gtagaggagc aaacagcgtt tgcgacatcc ttttgaata ctgcggaact gactaaagta 480
 gtgagttata cacagggctg ggatctattc tttttatctt tttttattct ttctttattc 540
 tataaattat aaccacttga atataaaca aaaaaacaca caaaggtcta gcggaattta 600
 cagagggctc agcagaattt acaagttttc cagcaaaggc ctagcagaat ttacagatac 660
 ccacaactca aaggaaaagg actagtaatt atcattgact agcccatctc aattggtata 720
 gtgattaata tcacctagc caattgagat gtatgtctga attagttgtt ttcaaagcaa 780
 atgaactagc gattagtcgc tatgacttaa cggagcatga aaccaagcta attttatgct 840
 gtgtggcact actcaacccc acgattgaaa accctacaag gaaagaacgg acggtatcgt 900
 tcacttataa ccaatacgtc cagatgatga acatcagtag ggaaaatgct tatgggtgat 960
 tagctaaagc aaccagagag ctgatgacga gaactgtgga aatcaggaat cctttgggta 1020

-continued

aaggctttga gattttccag tggacaaact atgccaagtt ctcaagcgaa aaattagaat	1080
tagtttttag tgaagagata ttgccttata ttttccagtt aaaaaaattc ataaaatata	1140
atctggaaca tgtaagtct tttgaaaaca aatactctat gaggatttat gagtggttat	1200
taaaagaact aacacaaaag aaaactcaca aggcaaatat agagattagc cttgatgaat	1260
ttaagtccat gttaatgctt gaaaataact accatgagtt taaaaggctt aaccaatggg	1320
ttttgaaacc aataagtaa gatttaaca cttacagcaa tatgaaattg gtggttgata	1380
agcgaggccg cccgactgat acggtgattt tccaagttga actagataga caaatggatc	1440
tcgtaaccga acttgagaac aaccagataa aatgaatgg tgacaaaata ccaacaacca	1500
ttacatcaga ttctacctt cgtaacggac taagaaaaac actacacgat gctttaactg	1560
caaaaattca gctcaccagt tttgaggcaa aatttttgag tgacatgcaa agtaagcatg	1620
atctcaatgg ttcgttctca tggctcacgc aaaaacaacg aaccacacta gagaacatac	1680
tggctaaata cggaaggatc tgaggttctt atggctcttg tatctatcag tgaagcatca	1740
agactaacia acaaaagtag aacaactggt caccgtaga tatcaaaggg aaaactgtcc	1800
atatgcacag atgaaaacgg tgtaaaaaag atagatacat cagagctttt acgagttttt	1860
ggtgcattta aagctgttca ccatgaacag atcgacaatg taacagatga acagcatgta	1920
acacctaata gaacaggtga aaccagtaaa acaaagcaac tagaacatga aattgaacac	1980
ctgagacaac ttgttacagc tcaacagtca cacatagaca gcctgaaaca ggcgatgctg	2040
cttatcgaat caaagctgcc gacaacacgg gagccagtga cgcctccctg ggggaaaaaa	2100
tcattggcaat tctggaagaa atagcgcttt cagccggcaa acctgaagcc ggatctgcca	2160
ttctgataac aaactagcaa caccagaaca gcccgtttgc gggcagcaaa acccgttatg	2220
cttgtaaacc gttttgtgaa aaaattttta aaataaaaaa ggggacctct agggccccca	2280
attaattagt aatataatct attaaaggtc attcaaaagg tcatccaccg gatcaattcc	2340
cctgctcgcg caggctgggt gccaaagctct cgggtaacat caaggcccga tccctggagc	2400
ccttgccctc ccgcacgat atcgtgccgt gatcgaaatc cagatccttg acccgcatgt	2460
gaaaaccctc actgatccgc atgccgctt cacaagaag ctgggcgaac aaacgatgct	2520
cgcttccag aaaaccgagg atgcgaacca cttcatccgg ggtcagcacc accggcaagc	2580
gccgcgacgg ccgaggtctt ccgatctcct gaagccaggg cagatccgtg cacagcacct	2640
tgccgtagaa gaacagcaag gccgccaatg cctgacgatg cgtggagacc gaaaccttgc	2700
gctcgttcgc cagccaggac agaaatgcct cgacttcgct gctgcccag gttgccgggt	2760
gacgcacacc gtggaaacgg atgaaggcac gaaccagtg gacataagcc tgttcggttc	2820
gtaagctgta atgcaagtag cgtatgcgct cacgcaactg gtccagaacc ttgaccgaac	2880
gcagcgggtg taacggcgca gtggcggttt tcatggcttg ttatgactgt ttttttgggg	2940
tacagtctat gcctcgggca tccaagcagc aagcgcgcta cgccgtgggt cgatgtttga	3000
tgttatggag cagcaacgat gttacgcagc agggcagtcg ccctaaaaca aagttaaaca	3060
tcatgagggg agcgggtgat gccgaagtat cgactcaact atcagaggtg gttggcgtca	3120
tcgagcgcca tctcgaaccg acggtgctgg ccgtacattt gtacggctcc gcagtggatg	3180
gcggcctgaa gccacacagt gatattgatt tgctggttac ggtgaccgta aggcttgatg	3240
aaacaacgcg gcgagctttg atcaacgacc ttttgaaac ttcggcttcc cctggagaga	3300

-continued

gcgagattct	ccgcgctgta	gaagtcacca	ttgttggtgca	cgacgacatc	attccgtggc	3360
gttatccagc	taagcgcgaa	ctgcaatttg	gagaatggca	gcgcaatgac	attcctgcag	3420
gtatcttcga	gccagccacg	atcgacattg	atctggctat	cttgctgaca	aaagcaagag	3480
aacatagcgt	tgccttggtg	ggtccagcgg	cggaggaact	ctttgatccg	gttcctgaac	3540
aggatctatt	tgaggcgcta	aatgaaacct	taacgctatg	gaactcgccg	cccgactggg	3600
ctggcgatga	gcgaaatgta	gtgcttacgt	tgtcccgcac	ttggtacagc	gcagtaaccg	3660
gcaaaatcgc	gccgaaggat	gtcgctgccg	actgggcaat	ggagcgcctg	ccggcccagt	3720
atcagcccgt	catacttgaa	gctagacagg	cttatcttgg	acaagaagaa	gatcgcttgg	3780
cctcgcgcgc	agatcagttg	gaagaatttg	tccactacgt	gaaaggcgag	atcaccaagg	3840
tagtcggcaa	ataatgtcta	acaattcggt	caagccgacg	ccgcttcgcg	gcgcggtta	3900
actcaagcgt	tagatgcact	aagcacataa	ttgctcacag	ccaaactatc	aggtcaagtc	3960
tgettttatt	atthttaagc	gtgcataata	agccctacac	aaattgggag	atatatcatg	4020
aaaggctggc	tttttcttgt	tatcgcaata	gttggcgaag	taatcgcaac	atccgcatta	4080
aatctagcgc	agggctttac	taagctgatc	cggtggtatg	ccttttgaat	gacctttaat	4140
agattatatt	actaattaat	tggggaccct	agaggtcccc	ttttttatth	taaaaattht	4200
ttcacaaaac	ggtttacaag	catacgttgg	ccgattcatt	aatgcagctg	gcacgacagg	4260
tttcccgcgc	ggaaagcggg	cagtgagcgc	aacgcaatta	atgtgagtta	gctcactcat	4320
taggcacccc	aggctttaca	ctttatgctt	ccggctcgta	tgttggtggtg	aattgtgagt	4380
gccaatacc	gtgaaatctc	gctggagcgg	gcgctgcttt	ataccgccag	ccatcggcaa	4440
accgaaggcg	aaccggtgat	attgcgccgg	gcgaaagcaa	cagcgtatat	ccttgaacat	4500
gttgaaatth	cgattcgtga	tgaagaactg	attgccggta	accgcaccgt	aaaaccgcgc	4560
gccgggatta	tgtcgcggga	aatggaccct	tactggctgc	tgaaagagct	ggatcaattc	4620
ccgacgcgtc	cgcaggaccg	ctttgctatc	agcgaagaag	ataaacgtat	ctaccgcgaa	4680
gagttgttcc	cgtactggga	aaaacgttcg	atgaaagatt	tcatcaacgg	gcagatgaca	4740
gatgaagtaa	aagccgcgac	cagcacgcag	atthtcagca	tcaaccagac	agataaaggc	4800
caggggcaca	ttattattga	ttaccacgc	ctgctgaatc	acgggctggg	ggagctggta	4860
gcacagatgc	agcaacattg	tcagcaacag	ccggagaatc	actthtatca	ggcagcgcgtg	4920
ttactgctgg	aagcctcgca	gaaacacatt	ttgcgttacg	ccgaactggc	ggaaacgatg	4980
gcgcaaaact	gcacagatgc	ccagcgtcgc	gaagagctgc	tgactattgc	ggagatctcc	5040
cgccataacg	cgcaacataa	gccgcagacg	ttctggcagg	cgtgccagtt	attctgggtac	5100
atgaacatca	ttctgcaata	cgaatccaac	gccagttcgc	tatcgttggg	gcgcttcgac	5160
cagtatatgt	tgcggttcta	tcagacatca	ttaaccagg	gcgaagatgc	ggcgttcctg	5220
aaagaactgc	tcgaatctth	atgggtgaaa	tgcaacgaca	tcgtgctggt	gcgctccacc	5280
agcagcgcgc	gthatttcgc	cggthtcccg	accggctata	ccgcactgct	cggcggtta	5340
accgagaacg	gacgtagcgc	ggtgaacgtg	ctthcgttcc	thtgccttga	cgctatcaa	5400
agcgtgcaat	taccgcaacc	gaacctcggc	gtgcgcaacta	acgccttgat	cgacacgcgc	5460
thcctgatga	aaaccgcccga	aaccattcgc	ctcggcaccg	gtattccgca	aatctthaac	5520
gatgaagtgg	tggtgccagc	gthcctcaac	cgtggcgtth	cgctggaaga	tgcgcgcgac	5580

-continued

tattccgtag	tgggctgtgt	ggaattatct	attcccggca	gaacctacgg	cttgcgatgac	5640
atcgcgatgt	ttaacctgct	gaaagtgatg	gaaatctgcc	tgcatgaaaa	tgaaggcaat	5700
gccgcgctga	cttatgaagg	tttactggaa	cagatccgtg	ccaagatcag	ccactacatc	5760
accctgatgg	ttgaaggcag	taatatttgc	gatatcggcc	atcgcgactg	ggcacctgta	5820
ccgctgctct	cgtcttttat	cagcgattgt	ctggaaaaag	gccgcgatat	taccgatggc	5880
ggcgcgctgt	ataacttctc	cggcgtacag	gggatcggtg	tcgccaacct	gagcgattct	5940
ctccatgcgt	tgaagggat	ggtttttgat	caacagcgtt	taagttttga	cgaattgctg	6000
tcggtattaa	aagccaactt	tgcaacgcca	gaaggcgaaa	aagtccgcgc	tcgcttaatt	6060
aaccgctttg	agaaatacgg	taacgatatc	gacgaggtgg	ataacattag	cgccgaactg	6120
ttgcgccact	actgcaaaga	agtggaaaaa	taccagaacc	cgcgcgccgg	ctacttcacg	6180
ccgggatcgt	agggcttcaa	actttcgccc	ctcctggcat	tgattcagcc	tgctcggaact	6240
ggtatttaac	cagactaatt	atthttgatgc	gcgaaattaa	tcgttacagg	aaaagccaaa	6300
gctgaatcga	ttttatgatt	tggttcaatt	cttcctttag	cggcataatg	tttaatgacg	6360
tacgaaacgt	cagcggtcaa	cacccgccag	caatggactg	tattgcgctc	ttcgtgcgctc	6420
gcgtctgtta	aaaactggcg	ctaacaatac	aggctaaagt	cgaaccgcca	ggctagactt	6480
tagttccaca	acactaaacc	tataagttgg	ggaaatacaa	tgttccagca	agaagttacc	6540
attaccgctc	cgacaatctg	ctaataccacg	agatgcggcc	caatttactg	cttaggagaa	6600
gatcatgggt	ttgttcgata	aactgaaatc	tctggtttcc	gacgacaaga	aggataccgg	6660
aactattgag	atcattgctc	cgctctctgg	cgagatcgtc	aatatcgaag	acgtgccgga	6720
tgctgthttt	gcggaaaaaa	tcgthttgta	tggtattgct	atcaaaccaa	cgggtaaaaa	6780
aatggtcgcg	ccagtagacg	gcaccattgg	taaaatcttt	gaaaccaacc	acgcattctc	6840
tatcgaatct	gatagcggcg	ttgaactggt	cgtccacttc	ggatcgcaca	ccgthgaact	6900
gaaaggcgaa	ggcttcaagc	gtattgctga	agaaggtcag	cgctgaaag	ttggcgatac	6960
tgctattgaa	ttgatctgc	cgctgctgga	agagaaagcc	aagtctaccc	tgactccggt	7020
tgthtatctc	aacatggacg	aatcaaaaga	actgatcaaa	ctgtccggtg	gcgthaacgt	7080
gggtgaaacc	ccgthtatcc	gcatcaagaa	gtaattcttg	ccgcagtgaa	aatggcgcc	7140
catcgcgccc	atthttttat	gcttccgcca	gcggcgccaa	aatcaattca	tcgctctcat	7200
gctgctgggt	gtagcgcctt	accgthttctg	ctcacgttcc	gthgggatcg	gtgthttggcg	7260
cgacgccaga	cggctgthtt	gccggagaac	agctggcaga	cggcggtctg	tcacctatgc	7320
tggttcagga	cgcacaaggg	ccaacggcgg	tactgaagtc	agtcagtaag	ctcgataaca	7380
cactgctgtc	taacggtaca	ttgctgaacg	tgaattcac	tccggcgacc	ctggaagggtg	7440
aagcgggatt	acgcaaactg	gccgacttct	tacggcggtt	taccagctt	aagthacaac	7500
atattcagtt	taacgtggtg	aacgcccaga	cgttgccgga	agcgcacacg	cgcccacaag	7560
attatgccgg	gctggtggtg	cgcgthtccg	gatacagcgc	cttctthtgc	gaactgtcga	7620
aggagatcca	ggatgacatc	atccgccgga	cagcgcacca	gctgthaacgt	tgtggaaacg	7680
cgccgcaatg	atgtggcgcg	cathttcaac	atcagcgtt	atcactgaa	tgacgggtgag	7740
ggcattcgta	cggthgtctt	thttaaaggc	tgtccgcac	thtgcccgtg	gtgtgctaat	7800
ccggagtcga	tctccggcaa	aatccagacg	gtacgcagag	aggcgaaatg	tctgcactgt	7860

-continued

```

gcgaaatggt tgcgtgatgc ggatgaatgc cctccgggg cgtttgaacg gattggtcgc 7920
gatatcagcc ttgacgctct ggaacgggaa gtgatgaaag atgacatddd ttttcgcacg 7980
tccggcggcg gcgtcacgct ttctggcggc gaagtgttaa tgcaggcggg gtttgctacc 8040
cgtdttttac agcgactgca gctgtgggggt gtctcatgtg ccattgaaac tgccggagac 8100
gcgccagcca gcaagctggt accgctggcg aaattgtgcg atgaagtgtt gttcgattta 8160
aaaatcatgg acgcgactca ggcgcgggat gtggtgaaga tgaacctgcc acgctgctg 8220
gagaatctgc gtttgctggt gagtgagggc gtcaactgta tcccgcgtdt accgctgatc 8280
cctggtttca cgctcagccg ggagaatatg cagcagggcg tggatgtgct gatcccgtg 8340
aatatcaggg agatccatct gttaccgttt catcagtacg gcgaaccgaa ataccgctg 8400
ctggggaaaa catggtcgat gaaagaggtg cctgcgcgct cgtcagccga tgtggcaacg 8460
atgcgcgaaa tggcagaacg ggccggattd caggttaccg tgggaggtta aaatggcata 8520
cctggtggca gtaaccgctt gcgtcagtgg cgtggcgcct acttatatgg cggcggaacg 8580
gctggaaaag ttgtgctgt tagagaagtg gggagtcagc attgaaactc agggcgctcg 8640
aattcactgg cgtcgttdt acaacgtcgt gactgggaaa acctggcgt tacccaactt 8700
aatcgcttg cagcacatcc ccttttcgcc agctggcgta atagcgaaga ggcccgcacc 8760
gatcgccctt cccaacagtt gcgcagcctg aatggcgaat ggccgctgat gcggtatddd 8820
ctccttacgc atctgtgctg tatttcacac cgcatatggt gcaactctag tacaatctgc 8880
tctgatgccg catagttaag ccagccccga caccgcgcaa caccgctga cgaatc 8937

```

```

<210> SEQ ID NO 9
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer BY124 for amplifying and cloning teh
pflDC geen of Escherichia coli C, or cassettes embedded in pflD,
together with BY125.

```

```

<400> SEQUENCE: 9

```

```

tgccaatacc cgtgaaatct cgc 23

```

```

<210> SEQ ID NO 10
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR Primer BY125 for amplifying and cloning the
pflDC genes of E. coli C, or cassettes embedded in pflD, together
with BY124.

```

```

<400> SEQUENCE: 10

```

```

cgccctgagt ttcaatgctg act 23

```

```

<210> SEQ ID NO 11
<211> LENGTH: 26
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: PCR primer AC74 for amplifying and clonig the
tdcC-tdcG region of E. coli C and derivatives or cassettes
embedded in teh tdcC-tdcG region,together with AC75.

```

```

<400> SEQUENCE: 11

```


-continued

 tgagctacct ggtatggccg tttatc

26

<210> SEQ ID NO 12
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: PCR primer Ac75 for amplifying and cloning the tdcC-tdcG region of E. coli C and derivatives, or cassettes embedded in the tdcC-tdcG region, together with AC74.

<400> SEQUENCE: 12

cccgccgccg acagagtaat ag

22

What is claimed is:

1. A bacterium producing more than 30 grams per liter of a desired chemical, wherein one of the biosynthetic intermediates for said desired chemical is phosphoenolpyruvate, and said bacterium contains at least one exogenous gene that encodes a protein that functions in the facilitated diffusion of a sugar.

2. The bacterium of claim 1, wherein the bacterium is selected from a group consisting of *Gluconobacter oxydans*, *Gluconobacter asaii*, *Achromobacter delmarvae*, *Achromobacter viscosus*, *Achromobacter lacticum*, *Agrobacterium tumefaciens*, *Agrobacterium radiobacter*, *Alcaligenes faecalis*, *Arthrobacter citreus*, *Arthrobacter tumescens*, *Arthrobacter paraffineus*, *Arthrobacter hydrocarboglutamicus*, *Arthrobacter oxydans*, *Aureobacterium saperdae*, *Azotobacter indicus*, *Brevibacterium ammoniagenes*, *divaricatum*, *Brevibacterium lactofermentum*, *Brevibacterium flavum*, *Brevibacterium globosum*, *Brevibacterium fuscum*, *Brevibacterium ketoglutamicum*, *Brevibacterium helcolum*, *Brevibacterium pusillum*, *Brevibacterium testaceum*, *Brevibacterium roseum*, *Brevibacterium immariophilum*, *Brevibacterium linens*, *Brevibacterium protopharmiae*, *Corynebacterium acetophilum*, *Corynebacterium glutamicum*, *Corynebacterium callunae*, *Corynebacterium acetoacidophilum*, *Corynebacterium acetoglutamicum*, *Enterobacter aerogenes*, *Erwinia amylovora*, *Erwinia carotovora*, *Erwinia herbicola*, *Erwinia chrysanthemi*, *Flavobacterium peregrinum*, *Flavobacterium fucatum*, *Flavobacterium aurantium*, *Flavobacterium rhenanum*, *Flavobacterium sewanense*, *Flavobacterium breve*, *Flavobacterium meningosepticum*, *Micrococcus* sp. CCM825, *Morganella morganii*, *Nocardia opaca*, *Nocardia rugosa*, *Planococcus eucinatus*, *Proteus rettgeri*, *Propionibacterium shermanii*, *Pseudomonas synxantha*, *Pseudomonas azotoformans*, *Pseudomonas fluorescens*, *Pseudomonas ovalis*, *Pseudomonas stutzeri*, *Pseudomonas acidovolans*, *Pseudomonas mucidolens*, *Pseudomonas testosteroni*, *Pseudomonas aeruginosa*, *Rhodococcus erythropolis*, *Rhodococcus rhodochrous*, *Rhodococcus* sp. ATCC 15592, *Rhodococcus* sp. ATCC 19070, *Sporosarcina ureae*, *Staphylococcus aureus*, *Vibrio metschnikovii*, *Vibrio tyrogenes*, *Actinomyces madurae*, *Actinomyces violaceochromogenes*, *Kitasatosporia parulosa*, *Streptomyces coelicolor*, *Streptomyces flavelus*, *Streptomyces griseolus*, *Streptomyces lividans*, *Streptomyces olivaceus*, *Streptomyces tanashiensis*, *Streptomyces virginiae*, *Streptomyces antibioticus*, *Streptomyces cacaoi*, *Streptomyces lavendulae*, *Streptomyces viridochromogenes*, *Aeromonas salmonicida*, *Bacillus pumilus*,

Bacillus circulans, *Bacillus thiaminolyticus*, *Escherichia freundii*, *Microbacterium ammoniophilum*, *Serratia marcescens*, *Salmonella typhimurium*, *Salmonella schottmulleri*, *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus amyloliquefaciens* and *Xanthomonas citri*.

3. The bacterium of claim 1, wherein the bacterium is selected from a group consisting of *Escherichia coli*, *Corynebacterium glutamicum*, *Brevibacterium flavum*, *Mannheimia succiniproducens* and *Anaerobiospirillum succiniproducens*.

4. The bacterium according to claim 1, wherein said desired chemical is selected from a group consisting of succinic acid, fumaric acid, glucaric acid, malonic acid, maleic acid, 2,5-furan dicarboxylic acid, propionic acid, 3-hydroxypropionic acid, aspartic acid, glucaric acid, glutamic acid, itaconic acid, levulinic acid, 3-hydroxybutyrolactone, and butanediols such as 1,4 butanediol, 1,3-butanediol and 2,3-butanediol.

5. The bacterium according to claim 1, wherein said desired chemical is succinate.

6. The bacterium according to claim 1, wherein said desired chemical is cis, cis-muconic acid.

7. The bacterium according to claim 1, wherein said desired chemical is an aromatic biochemical.

8. A bacterium according to claim 1, wherein the bacterium is a PTS⁻ bacterial strain.

9. A bacterium according to claim 1, wherein the bacterium has a reduced level of phosphotransferase activity compared to a related wild type strain.

10. A bacterium of claim 1 further comprising a mutation or deletion in one or more genes that encode proteins that functions in a phosphotransferase system for sugar import, said mutated or deleted gene being other than a crr gene.

11. A bacterium of claim 10 further comprising a deletion in a gene that encodes a sugar importer that functions using proton symport.

12. A bacterium of claim 10 wherein said mutated or deleted gene is a ptsH gene or a homolog thereof.

13. A bacterium of claim 10 wherein said mutated or deleted gene ptsI gene or a homolog thereof.

14. A bacterium of claim 10 wherein said mutated or deleted genes are selected from a group consisting of ptsH, ptsI, homolog of ptsH and homolog of ptsI.

15. A bacterium according to claim 1, wherein the bacterium is a galP⁻ bacterial strain.

16. The bacterium of claim 1, wherein said exogenous gene is contained on a replicating plasmid.

17. The bacterium of claim 1, wherein said exogenous gene is integrated into the host chromosome.

18. The bacterium of claim 1 wherein said exogenous gene is a glf gene.

19. The bacterium of claim 1 wherein said exogenous genes are a glf and a glk genes.

20. The bacterium of claim 1 in which said exogenous genes are a glf gene and a frk gene.

21. The bacterium of claim 1 in which said exogenous gene is derived from a yeast.

22. The bacterium of claim 1 in which said exogenous gene or genes are derived from a strain of *Zymomonas mobilis*.

23. A bacterium of claim 1, wherein said bacterium is grown in minimal medium.

24. A bacterium of claim 1 wherein said bacterium produces more than 64 grams per liter of a desired chemical.

25. A bacterium of claim 1 wherein said bacterium produces more than 83 grams per liter of a desired chemical.

26. A bacterium containing two or more copies of a functional crr gene or a functional homolog of a crr gene.

27. A method for producing a desired chemical comprising the steps of:

growing a bacterium of claim 1 in a minimal fermentation medium; and

optionally purifying said chemical from the fermentation medium.

28. A method for producing succinic acid comprising the steps of:

growing a bacterium comprising at least one exogenous gene that encodes a protein that function in the facilitated diffusion of sugar and producing at least 60 grams of succinate per liter and less than 4.2 grams of acetate per liter in a minimal fermentation medium; and

optionally purifying succinic acid from the fermentation medium.

29. A method for improving the titer and yield of a desired chemical produced by a bacterial strain that has been engineered to use facilitated diffusion for import of a sugar, comprising the steps of:

subjecting a parent strain to serial transfers into fresh liquid medium;

plating the resulting culture for single colonies on a petri plate; and

choosing a single colony.

30. A bacterial strain engineered to use facilitated diffusion for import of a sugar comprising a glf-glk operon, wherein said bacterial strain has been evolved for improved titer and yield for succinate production, and further comprising one or more mutations in the glf-glk operon.

31. A bacterial strain of claim 30 in which said one or more mutations alter bases in the DNA sequence that corresponds to the 5' untranslated leader region of an mRNA that encodes either of said glf or glk gene.

32. A bacterial strain engineered to use facilitated diffusion for import of a sugar comprising a glf gene, wherein said bacterial strain has been evolved for improved titer and yield for succinate production, and said glf gene comprises one or more mutations.

33. A bacterial strain engineered to use facilitated diffusion for import of a sugar comprising a glk gene, wherein said bacterial strain has been evolved for improved titer and yield for succinate production, and said glk gene comprises one or more mutations.

* * * * *