

(19) **United States**(12) **Plant Patent Application Publication**  
**Jacobs**(10) **Pub. No.: US 2016/0192565 P1**  
(43) **Pub. Date: Jun. 30, 2016**(54) **CHRYSANTHEMUM PLANT NAMED**  
**'ZANMUCUP'****Publication Classification**(71) Applicant: **Chrysanthemum Breeders Association**  
**Research B.V.**, Valkenburg ZH (NL)(51) **Int. Cl.**  
**A01H 5/00** (2006.01)(72) Inventor: **Henricus Cornelius Maria Jacobs**,  
Valkenburg (NL)(52) **U.S. Cl.**  
USPC ..... **PLT/295**(21) Appl. No.: **14/544,362**(57) **ABSTRACT**(22) Filed: **Dec. 30, 2014**

A *chrysanthemum* plant named 'Zanmucup' characterized by its medium sized blooms with intense yellow ray florets and prolific branching; natural season flowering date September 22; blooming for a period of 4 weeks.

## BOTANICAL DESIGNATION

[0001] *Chrysanthemum* × *morifolium* Ramat.

[0009] FIG. 3 shows the various stages of foliage of the new cultivar.

## CULTIVAR DENOMINATION

[0002] 'Zanmucup'

## DESCRIPTION OF THE INVENTION

## RELATED CULTIVARS

[0003] This new plant cultivar 'Zanmucup' is of the botanical classification *Chrysanthemum* × *morifolium* Ramat.

[0010] The observations and measurements were gathered from plants grown out door in Rijsenhout, The Netherlands, under natural day length and temperature and planted in week 25 in 2014, receiving drip irrigation. The natural blooming date of this crop was September 22. The average height of the plants was 30 cm. No growth retardants were used. No tests were done on disease or insect resistance or susceptibility. No tests were done on cold or drought tolerance. This new variety produces medium sized blooms with intense yellow ray florets blooming for a period of 4 weeks.

## BACKGROUND OF THE INVENTION

[0004] The present invention relates to a new and distinct cultivar of *chrysanthemum* plant, botanically known as *Chrysanthemum* × *morifolium* Ramat., commercially known as a garden mum, and hereinafter referred to by the cultivar denomination 'Zanmucup'. 'Zanmucup' is a product of a breeding and selection program for outdoor pot mums (garden mums) which had the objective of creating new cultivars with a double type inflorescence, a natural season flowering date around September 22, blooming for a period of 5 weeks. 'Zanmucup' is a seedling resulting from a cross of the female parent 21570 with the male parent 24605. Plants of the new cultivar 'Zanmucup' differ from plants of the female parent in the natural season flowering date; plants of the seedling flower one week earlier than those of the female parent. Plants of the seedling differ from plants of the male parent in inflorescence type; inflorescences of the seedling are double, while those of the male parent are semi-double.

[0011] From the cultivars known to inventor the most similar existing cultivar in comparison to 'Zanmucup' is 'Zanmustone' (U.S. Plant Pat. No. 22,619). When 'Zanmustone' and 'Zanmucup' are being compared the following difference is noticed: The differences of 'Zanmustone' and 'Zanmucup' are (1) Color ray florets. And (2) Leaf color. (1) Ray-florets of 'Zanmucup' are more deep yellow than those of 'Zanmustone'. (2) Leaves of 'Zanmustone' are more greyish green than those of 'Zanmucup'.

[0005] The new and distinct cultivar was discovered and selected as a flowering plant by Henricus Cornelius Maria Jacobs on a cultivated field in Rijsenhout, The Netherlands in 2010. The first act of asexual production of 'Zanmucup' was accomplished when vegetative cuttings from the initial selection in 2010 were propagated further in a controlled environment in Rijsenhout, The Netherlands. The new cultivar has been found to retain its distinctive characteristics through successive propagations.

[0012] The following is a description of the plant and characteristics that distinguish 'Zanmucup' as a new and distinct variety.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0006] The present invention of a new and distinct variety of *chrysanthemum* (of about 16 weeks) is shown in the accompanying drawings, the color being as nearly true as possible with color photographs of this type.

[0007] FIG. 1 shows a plant of the cultivar in full bloom.

[0008] FIG. 2 shows the various stages of bloom of the new cultivar.

[0013] The color designations are taken from the plant itself. Accordingly, any discrepancies between the color designations and the colors depicted in the photographs are due to photographic tolerances. The color chart used in this description is: The Royal Horticultural Society Colour Chart, edition 2001.

TABLE 1

Botanical Description of <i>Chrysanthemum</i> × <i>morifolium</i> Ramat. 'Zanmucup'	
Bud	
Size:	Small; cross-section 4 mm, height 3 mm
Shape:	Round
Texture:	Pubescent
Outside color:	Greyed-green 191A
Phyllaries	
Number:	24, arranged in 3 rows
Shape:	Elliptic
Apex:	Acute
Base:	Truncate

TABLE 1-continued

Botanical Description of <i>Chrysanthemum × morifolium</i> Ramat. 'Zanmucup'	
Margin:	Entire
Color:	Upper side: Greyed-green 189A Lower side: Green 136A
Length and width:	4 mm; 3 mm
Texture:	Pubescent
Inflorescences	
Type:	Double
Height:	2 cm
Diameter:	4 cm
Peduncle length:	4-6.5 cm
Peduncle color:	Green 136C
Peduncle diameter:	2 mm
Peduncle texture:	Pubescent
Number per branch:	About 5-6 .inflorescences
Duration of flowering:	5 weeks
Seeds:	Produced in small quantities, ovate, Greyed-brown 199A, length 1.5 mm, diameter 0.5 mm
Fragrance:	Faint chrysanthemum odor
Color inflorescences	
Center of inflorescence:	Immature stage: Yellow-orange 17A.; Mature stage: Yellow-orange 17A Yellow 12a
Color of upper side of ray florets:	
Color of lower side of ray florets:	Yellow 8A
Tonality from distance:	A garden mum with yellow blooms and a dark yellow center
Color of ray florets after aging of the plant: Ray florets	Yellow-orange 15A
Texture:	Upper and lower side smooth
Number:	160
Shape:	elliptic
Apex:	dentate
Base:	acute
Cross-section:	flat
Longitudinal axis:	straight
Length of corolla tube:	3-4. mm
Ray floret margin:	Entire
Ray floret length:	1-2 cm
Ray floret width:	2-5 mm
Ratio length/width:	medium
Disc florets	absent
Receptacle	
Color:	Yellow-green 145D
Shape:	Domed raised
Height:	3 mm
Diameter:	3 mm
Reproductive Organs	
Androecium:	absent
Gynoecium:	Present on ray florets
Style color:	Yellow-green 154C
Style length:	4 mm
Stigma color:	Yellow 7A

TABLE 1-continued

Botanical Description of <i>Chrysanthemum × morifolium</i> Ramat. 'Zanmucup'	
Stigma width:	1 mm
Ovary:	Enclosed in calyx
Plant	
Form:	Grown as a spray type pot mum, outdoor raised and mounded
Growth habit:	Spherical shape
Growth rate:	Medium
Height:	30 cm
Width:	40-45 cm
Stem color:	Greyed-brown 199A
Stem strength:	strong
Stem brittleness:	not brittle
Stem anthocyanin coloration:	absent
Internode length:	1 cm
Length of lateral branch:	From top to bottom 20-22 cm
Lateral branch color:	Green 137D
Lateral branch, attachment:	Weak
Lateral branch diameter:	2 mm
Branching (average number of lateral branches):	Prolific with 8 breaks after pinching
Natural season blooming date:	September 22 (week 39)
Foliage	
Leaf color:	Upper side: Green 136B Lower side: Green 138B
Color midvein:	Upper side: Yellow-green 147D Lower side: Yellow-green 148D
Size:	Small; length 3-5.5 cm, width 1.5-2.6 cm
Quantity (number per lateral branches):	20
Shape:	Elliptic
Texture upper side:	Sparsely pubescent
Texture under side:	Pubescent
Venation arrangement:	Pinnate
Shape of margin:	Serrated
Shape of base of sinus between lateral lobes:	Rounded
Margin of sinus between lateral lobes:	Diverging
Shape of base:	Truncate to asymmetric
Apex:	Pointed
Petiole length:	0.5-1.2 cm
Petiole diameter:	2 mm
Petiole color:	Yellow-green 147D

TABLE 2

	Differences with the comparison variety, when grown under identical conditions	
	'Zanmucup'	'Zanmustone'
Ray floret upperside color	Yellow 12A	Yellow 8A
Leaf upperside color	Green 136B	Green 139A-B

## I claim:

1. A new and distinct *chrysanthemum* plant named 'Zanmucup' as described and illustrated.

\* \* \* \* \*






FIG. 1



**FIG. 2**



**FIG. 3**