



US 20230105479A1

(19) **United States**

(12) **Patent Application Publication**

Lee et al.

(10) **Pub. No.: US 2023/0105479 A1**

(43) **Pub. Date: Apr. 6, 2023**

(54) **OPTICALLY CONTROLLABLE FGFR STIMULATION USING WIRELESS CONTROLLED CELLULAR LIGHTING SYSTEM**

(71) Applicant: **The Johns Hopkins University**,
Baltimore, MD (US)

(72) Inventors: **Gabsang Lee**, Ellicott City, MD (US);
Hyesoo Kim, Baltimore, MD (US);
InYoung Choi, Baltimore, MD (US);
HoTae Lim, Baltimore, MD (US); **Alex V. Huynh**, Baltimore, MD (US)

(73) Assignee: **The Johns Hopkins University**,
Baltimore, MD (US)

(21) Appl. No.: **17/957,998**

(22) Filed: **Sep. 30, 2022**

Related U.S. Application Data

(63) Continuation of application No. 17/152,220, filed on Jan. 19, 2021.

(60) Provisional application No. 62/962,246, filed on Jan. 17, 2020.

Publication Classification


(51) **Int. Cl.**
C07K 14/405 (2006.01)
C07K 14/71 (2006.01)
C12N 5/074 (2006.01)
C12N 13/00 (2006.01)
C12N 5/0735 (2006.01)
C12M 1/00 (2006.01)



(52) **U.S. Cl.**
CPC *C07K 14/405* (2013.01); *C07K 14/71* (2013.01); *C12N 5/0696* (2013.01); *C12N 13/00* (2013.01); *C12N 5/0606* (2013.01); *C12M 31/10* (2013.01); *C07K 2319/02* (2013.01)



(57) **ABSTRACT**




The present invention relates to the field of stem cells. More specifically, the present invention provides compositions and methods for using optogenetics to sustain the pluripotency of stem cells. In one embodiment, a vector comprises a nucleotide sequencing encoding a fusion protein comprising the intracellular domain of fibroblast growth factor 1 receptor (FGFR1) and a photoactivatable domain.

Specification includes a Sequence Listing.












FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E





FIG. 2F






FIG. 4A




FIG. 4B




FIG. 4C




FIG. 4D





FIG. 4G




FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D

IP Address: 10.16.109.103

No.	LED On	Time	Unit	Notes
1	1	1	MIN	
2	NONE	119	MIN	

Cycles: 10

Single Well Mode

Start	Pause	Add	Edit	Clear	Delete
-------	-------	-----	------	-------	--------

6-Well Plate
ID: 6W-001

0%

0%

0%

0%

0%

0%

Plate 1
12-Well

Plate 2
6-Well

+ Add Plate

Blue Screen Board v1.0

FIG. 5E




FIG. 6A




FIG. 6B




FIG. 6C



FIG. 6D




FIG. 6E




FIG. 6F




FIG. 7A




FIG. 7B




FIG. 8A




FIG. 8B




FIG. 8C




FIG. 9A




FIG. 9B




FIG. 9C




FIG. 9D




FIG. 9E




FIG. 9F




FIG. 10A




FIG. 10B




FIG. 11A




FIG. 11B




FIG. 12A




FIG. 12B




FIG. 12C

gRNA	No. of transfected cells	No. of colonies after selection
#1	8 million	6
#91	4 million	7

FIG. 13A




FIG. 13B




FIG. 13C



FIG. 13D




FIG. 13E



FIG. 13F




FIG. 14C

FIG. 14B

FIG. 14A





FIG. 14E

FIG. 14D






FIG. 15B




FIG. 15C




FIG. 16




FIG. 17




FIG. 18

OPTICALLY CONTROLLABLE FGFR STIMULATION USING WIRELESS CONTROLLED CELLULAR LIGHTING SYSTEM

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application claims the benefit of U.S. Provisional Application No. 62/962,246, filed Jan. 17, 2020, which is incorporated herein by reference in its entirety.

GOVERNMENT SUPPORT CLAUSE

[0002] This invention was made with government support under grant nos. NS093213 and AR070751 awarded by the National Institutes of Health, and grant no. 1547515 awarded by the National Science Foundation. The government has certain rights in the invention.

FIELD OF THE INVENTION

[0003] The present invention relates to the field of stem cells. More specifically, the present invention provides compositions and methods for using optogenetics to sustain the pluripotency of stem cells.

INCORPORATION-BY-REFERENCE OF MATERIAL SUBMITTED ELECTRONICALLY

[0004] This application contains a sequence listing. It has been submitted electronically via EFS-Web as an ASCII text file entitled "P15986-02_ST25.txt." The sequence listing is 26,756 bytes in size, and was created on Jan. 19, 2021. It is hereby incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

[0005] Identification of channel rhodopsins as light-gated ion channels in algal system [1], followed by various sensory photoreceptors, has opened a new chapter of optogenetics [2] in biological processes. Optogenetics utilized key properties of sensory photoreceptors by sophisticated modulation of molecular targets with high spatial and temporal control, which certainly overcomes many drawbacks of conventional approaches. Current application of optogenetics is mostly limited to neuroscience field relying on ion channels and light-driven ion pumps [3], but there should be plenty of other applications in many biology and medicine. **[0006]** Previous pioneering studies have discovered several photoactivatable proteins, such as LOV (light-oxygen-voltage-sensing) domain [4,5], phytochrome B (PhyB) [6,7], and cryptochrome 2 (Cry2) [8]. Such photoactivatable proteins are key players for the optogenetic control of intracellular signal transduction because, by absorbing energy from the photons in excitation light, they can undergo conformational changes, rearrange inter- or intra-protein contacts, and modulate inter- or intra-protein interactions [9]. Such changes in photoactivatable proteins can mimic the dimerization of a signaling pathway, simply by light illumination at a certain wavelength. For example, a photosensitive protein Cry2 homo-oligomerizes, and LOV domain homo-dimerizes, both within seconds when illuminated with stimulatory light (~470 nm) [9,10]. Therefore, photoactivatable proteins are enticing light-sensing modules for use in optogenetic studies of intracellular signaling in mammalian systems [11] because they have rapid responsiveness with

high temporal (sub-second time resolution) and spatial (sub-cellular spatial resolution) precision, and they do not need exogenous cofactors or signaling peptides.

[0007] Fibroblast growth factor 2 (FGF2) is a key component that promotes self-renewal and inhibits spontaneous differentiation of mammalian pluripotent stem cells (PSCs), including human PSC (hPSC) and porcine induced PSC (piPSC) [12-19]. The human FGF family consists of 22 members (FGF1 to 23, except FGF15), which are structurally related signaling molecules and subdivided into 7 subfamilies [20]. The mammalian FGF receptor (FGFR) family, however, has only 4 highly conserved receptor tyrosine kinases (FGFR1 to 4) [7]. Among them, FGFR1 predominantly interacts with FGF1 and FGF2 [21-23]. Like other receptor tyrosine kinases, FGFR1 dimerizes and trans-autophosphorylates a large number of tyrosine residues upon binding with FGF2, consequently activating downstream signaling pathways [24].

[0008] PSCs can be proliferated indefinitely and have the potential to differentiate into many different cell types with promising applications. The establishment of a high-quality, high-precision culture system is necessary for PSC applications; the existing culture protocols are completely dependent on recombinant proteins that are critical culture medium components for the maintenance of PSCs [12-16, 19]. However, high cost, thermal instability [25], and random distribution of the recombinant proteins impede large-scale manufacturing. Here, we have developed a novel culture system for PSCs using optical induction of the FGF signaling pathway. Our system maintained pluripotency of PSCs similar to the conventional culture system. Furthermore, the optically maintained PSCs displayed the ability to differentiate into three germ layers, demonstrating that the optical culture system can sustain the pluripotency of PSCs without exogenous FGF2 protein supplementation.

SUMMARY OF THE INVENTION

[0009] The present invention provides one or more fusion protein embodiments having a cytoplasmic region of the FGF receptor-1 and a light-oxygen-voltage domain, which creates a tunable, blue light-dependent activation of FGF signaling in human and porcine pluripotent stem cells (PSCs). The present invention also provides a novel PSC culture system which can include a wired or wirelessly controllable optical activation of the fibroblast growth factor (FGF) signaling pathway in the cells in culture without the need for daily supplementation of recombinant FGF2 protein, a key molecule for maintaining pluripotency of PSCs. The present invention provides a highly controllable optical stimulation of the FGF signaling pathway sufficient for long-term maintenance of PSCs, without the loss of differentiation potential into three germ layers. These embodiments allow for culture systems that are a cost-effective platform for human clinical trials and animal cellular agriculture

[0010] Therefore, in accordance with an embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain.

[0011] In accordance with another embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the

fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain comprising the nucleotide sequence of SEQ ID NO:6.

[0012] In accordance with an embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain, a promoter sequence and a sequence encoding a myristolation signal peptide (Myr).

[0013] In accordance with an embodiment, the present invention provides an expression vector comprising synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain.

[0014] In accordance with an embodiment, the present invention provides an expression vector comprising synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain.

[0015] In accordance with another embodiment, the present invention provides a an expression vector comprising synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain, a promoter sequence and a sequence encoding a myristolation signal peptide (Myr).

[0016] In accordance with another embodiment, the present invention provides a an expression vector comprising synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain, a promoter sequence and a sequence encoding a myristolation signal peptide (Myr) comprising the nucleotide sequence of SEQ ID NO:1.

[0017] In accordance with an embodiment, the present invention provides expression vectors comprising the synthetic polynucleotides described herein.

[0018] In accordance with an embodiment, the present invention provides a transformant transformed by the vectors described herein, including, for example, mammalian cells, and specifically, pluripotent stem cells.

[0019] In accordance with an embodiment, the present invention provides a fibroblast growth factor free pluripotent stem cell culture system comprising one or more transformed cells as described herein and an illumination source.

[0020] In accordance with an embodiment, the present invention provides a method for maintaining a pluripotent stem cell line in culture comprising: a) a plurality of pluripotent stem cells in a fibroblast growth factor free culture medium, wherein said are transformed with the expression vectors described herein; b) said transformed cells are maintained in the cell culture system as described herein; and c) periodically illuminating the cells of b) at a wavelength of about 470 nm for a sufficient time and intensity such that the cells maintain their pluripotent capability.

[0021] Accordingly, in one aspect, the present invention provides a vector. In one embodiment, a vector comprises a nucleotide sequencing encoding a fusion protein comprising

the intracellular domain of fibroblast growth factor 1 receptor (FGFR1) and a photoactivatable domain. In another embodiment, the fusion protein further comprises a signal peptide. In a specific embodiment, the signal peptide comprises a myristolation signal peptide (Myr). In a more specific embodiment, the Myr comprises the amino acid sequence of SEQ ID NO:4.

[0022] In another specific embodiment, the intracellular domain of FGFR1 comprises the amino acid sequence of SEQ ID NO:6. In particular embodiments, the photoactivatable domain comprises a light-oxygen-voltage sensing (LOV) domain. In a specific embodiment, the LOV domain comprises the amino acid sequence of SEQ ID NO:8. In yet another embodiment, the vector comprises the nucleotide sequence of SEQ ID NO:1. In certain embodiments, the present invention provides a pluripotent stem cell (PSC) comprising a vector described herein.

[0023] In another aspect, the present invention provides a modified pluripotent stem cell. In one embodiment, the present invention provides a PSC whose genome comprises a nucleotide sequence encoding a fusion protein comprising the intracellular domain of FGFR1 and a photoactivatable domain. In another embodiment, the present invention provides a PSC that expresses a fusion protein comprising the intracellular domain of fibroblast growth factor 1 receptor (FGFR1) and a photoactivatable domain.

[0024] In another embodiment, the fusion protein further comprises a signal peptide. In a specific embodiment, the signal peptide comprises a myristolation signal peptide (Myr). In a more specific embodiment, the Myr comprises the amino acid sequence of SEQ ID NO:4. In another specific embodiment, the intracellular domain of FGFR1 comprises the amino acid sequence of SEQ ID NO:6. In particular embodiments, the photoactivatable domain comprises a light-oxygen-voltage sensing (LOV) domain. In a specific embodiment, the LOV domain comprises the amino acid sequence of SEQ ID NO:8.

[0025] In a further aspect, the present invention provides a fibroblast growth factor 2 (FGF2) free PSC culture system. In one embodiment, the culture system comprises a PSC described herein and a cell culture illumination source capable of illuminating the PSC with a selected wavelength and intensity of light. In specific embodiments, the cell culture illumination source comprises a cell culture substrate, an upper and lower enclosure, one or more illumination sources, one or more circuit boards, and a microcontroller.

[0026] In particular embodiments, the illumination source can vary the wavelength of the illumination. In a specific embodiment, the illumination source emits light at a wavelength of about 470 nm. In other embodiments, the illumination source comprises one or more light emitting diodes (LEDs). In a specific embodiment, the illumination source continuously illuminates the cells for a period of time between about 1 minute and about 120 minutes. In another specific embodiment, the illumination source illuminates the cells in a time interval of between about 30 minutes to about 4 hours. In certain embodiments, the illumination source illuminates the cells at an intensity of between about 0.1 $\mu\text{W}/\text{mm}^2$ to about 25 $\mu\text{W}/\text{mm}^2$. In particular embodiments, the microcontroller is controlled through a computer server via either a wireless or wired computer client.

[0027] In a further aspect, the present invention provides methods for maintaining the pluripotency of PSCs without

exogenous FGF2 supplementation. In a specific embodiment, the method comprises the step of illuminating a PSC described herein with a light having a wavelength of about 470 nm.

BRIEF DESCRIPTION OF THE FIGURES

[0028] FIG. 1A-1H. Establishment of a novel and efficient FGF2-free hPSC culture system. FIG. 1A, 1B: A schematic illustration of the FGF signaling pathway by (FIG. 1A) FGF ligand-elicited activation or by (FIG. 1B) an opto-FGFR activation via blue light illumination. FIG. 1C: A schematic illustration of the AAVS1 locus targeting using homologous recombination enhanced by the CRISPR/Cas9 system. P1-P4 indicate the PCR primer locations. Myr, myristoylation signal peptide; cytoFGFR1, the cytoplasmic region of FGFR1. FIG. 1D: A schematic illustration of Opto-FGFR PSC maintenance (+F, 10 ng/mL FGF2 protein treatment, daily; +L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h). FIG. 1E: Representative colony morphologies of Opto-FGFR hESCs maintained in either +L condition for 54 weeks (left) or dark for 2 weeks (right). FIG. 1F, 1G: qRT-PCR results of Opto-FGFR hPSCs cultured for 3 weeks (FIG. 1F, hESCs) or 12 weeks (FIG. 1G, iPSCs) in either +F or +L condition (n=4; ns, not significant; unpaired t-test). All error bars represent mean $\pm$ s.e.m. FIG. 1H: Representative images of Opto-FGFR hiPSCs maintained in +L condition for 12 weeks showed undifferentiated colony morphology (left) and expression of pluripotency markers TRA-1-81 (middle) and NANOG (right). Scale bars, 100 μm .

[0029] FIG. 2A-2F. The maintenance of pluripotency of Opto-FGFR hESCs cultured with blue light illumination. FIG. 2A: A schematic illustration of Opto-FGFR hESCs maintenance (+F, 10 ng/mL FGF2 protein treatment supplied daily; +L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 1-10 min. every 2 h; Diff. Ctrl, hESC-derived myotubes). FIG. 2B, 2C: Heat maps showing expression levels, (FIG. 2B) in log 2(FPKM) or (FIG. 2C) in FPKM, of specific genes in Opto-FGFR hESCs cultured in either +F, +L, or Diff. Ctrl condition for 3 weeks. FPKM, fragments per kilobase of transcript per million mapped reads. FIG. 2D: A weighted Venn diagram between +F and +L groups, showing overlap of significantly differentially expressed genes compared to the Diff. Ctrl group. FIG. 2E, 2F: Gene ontology (GO) analyses of (FIG. 2E) significantly upregulated genes and (FIG. 2F) down-regulated genes in the +L group compared to the Diff. Ctrl group.

[0030] FIG. 3A-3G. In vitro and in vivo differentiation abilities of Opto-FGFR hPSCs maintained with blue light illumination into three germ layers. FIG. 3A: A schematic illustration of the differentiation of optically maintained Opto-FGFR hPSCs (1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h) into three germ layers. FIG. 3B-3G: Germ layer differentiation of Opto-FGFR (FIG. 3B-3E, hESCs; fig, hiPSCs) in which cells were cultured with blue light illumination. FIG. 3B, 3F: qRT-PCR results of the Opto-FGFR hPSCs, which were cultured with blue light illumination for 6 weeks and then differentiated into three germ layers (n=4). All error bars represent mean $\pm$ s.e.m. FIG. 3C: Representative images of the Opto-FGFR hESCs, which were optically maintained for 10 weeks and then differentiated into three germ layers and immunostained with the indicated antibodies. FIG. 3D: Representative images of the Opto-FGFR hESCs, which were cultured with blue light illumination for 13 weeks, differentiated into dopaminergic neurons (left)

and skeletal muscles (right), and immunostained with the indicated antibodies. FIG. 3E, 3G: Representative images of Opto-FGFR hPSC-derived tissues of three germ layers in in vivo teratoma assay. Scale bars, 100 μm .

[0031] FIG. 4A-4G. Establishment of a novel and efficient FGF2-free piPSC culture system that maintains pluripotency. FIG. 4A: A schematic illustration of the pRosa26 locus targeting using homologous recombination enhanced by the CRISPR/Cas9 system. cytoFGFR1, the cytoplasmic region of the FGFR1. FIG. 4B: A schematic illustration of Opto-FGFR piPSCs maintenance (+F, 10 ng/mL FGF2 protein treatment, daily; +L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h). FIG. 4C: Representative colony morphologies of Opto-FGFR piPSCs maintained in either +L condition (left) or dark (right) for 3 weeks.

[0032] FIG. 4D: A representative image of the Opto-FGFR piPSCs immunostained with OCT4 antibody after 3 weeks maintenance in +L condition. FIG. 4E: qRT-PCR results of Opto-FGFR piPSCs cultured in either +F or +L condition for 1 week (n=4; **, p value <0.01; *, p value <0.05; n.s., not significant; unpaired t-test). FIG. 4F: A schematic illustration of the differentiation of optically maintained Opto-FGFR piPSCs in +L condition into three germ layers. FIG. 4G: qRT-PCR results of the Opto-FGFR piPSCs, which were maintained in +L condition for 3 weeks and then differentiated into three germ layers (n=6). All error bars represent mean $\pm$ s.e.m. Scale bars, 100 μm .

[0033] FIG. 5A-5F. Simple and cost-efficient LED illumination system for cell culture. FIG. 5A: A blown-out diagram of the LED illuminating device, consisting of the base casing, the microcontroller board (Arduino Uno), the LED circuit boards, and the top casing, with a standard 6-well tissue culture plate. FIG. 5B: Wire diagram between Arduino Uno and LED circuit boards. FIG. 5C: A fully-assembled LED illuminating device for 6-well plates. FIG. 5D: Schematic overview of the wireless optogenetic system. FIG. 5E: The server computer's user interface (UI) where users can control the phases of illumination and LED intensity and view current status of LED illumination. FIG. 5F: The mobile client's UI where users can wirelessly control phases of illumination and LED intensity and view current status of LED illumination.

[0034] FIG. 6A-6F. The generation of opto-FGFR knock-in hPSCs. FIG. 6A: Validation of opto-FGFR gene cassette knock-in in hPSCs. Genomic DNA from the parental hESCs, Opto-FGFR hESCs, and Opto-FGFR hiPSCs was amplified using the indicated primer pairs (Endogenous, P1 and P2; Left Arm, P1 and P3; Right Arm, P4 and P2; primer pair locations are shown in FIG. 1c). FIG. 6B, 6C: Representative colony morphologies and NANOG expression of opto-FGFR knock-in (FIG. 6B) hESCs and (FIG. 6C) hiPSCs. FIG. 6D: PSC culture plates for FGF ligand treatment (left) and blue light illumination (right). FIGS. 6E, 6F: The phosphorylation kinetics of (FIG. 6E) ERK1/2 and (FIG. 6F) MEK in the Opto-FGFR hESCs treated with FGF2 protein (red circles, n=4) or blue light (blue squares, n=4). All error bars represent mean $\pm$ s.e.m. Scale bars, 100 μm .

[0035] FIG. 7A-7B. Phosphorylation of ERK1/2 in Opto-FGFR hESCs by blue light illumination. FIG. 7A, 7B: The effect of duration (a, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 0-10 min) and intensity (b, 0-1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min) of illumination on ERK1/2 phosphorylation (n=4; **, p value <0.01; *, p value <0.05; n.s., not significant; unpaired t-test). All error bars represent mean $\pm$ s.e.m.

[0036] FIG. 8A-8C. Profiles of protein phosphorylation upon activation of the FGF signaling pathway. FIG. 8A: The heatmap shows comparable levels of phosphorylation in multiple protein substrates in the FGF signaling pathway activated by FGF2 protein treatment (FGF, 10 ng/mL for 30 min; n=4) and blue light illumination (Opto, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 30 min; n=4). FIG. 8B: The relative value of phosphorylation in the blue light illumination and FGF2 protein treated groups. FIG. 8C: The ratio of phosphorylated versus non-phosphorylated proteins in the blue light illumination and FGF2 protein treated groups. All error bars represent mean $\pm$ s.e.m.

[0037] FIG. 9A-9F. Optogenetic activation of FGF signaling is sufficient to maintain pluripotency in hESCs. FIG. 9A: A schematic illustration of Opto-FGFR hESCs maintenance (+F, 10 ng/mL FGF2 protein treatment supplied daily; +L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h; Untreated, maintained without FGF2 protein or blue light). FIG. 9B: Quantification of FACS analyses of SSEA4⁺ cells from Opto-FGFR hESCs maintained in +F, +L, and Untreated conditions for 1 week (passage 1) or 3 weeks (passage 3) (n=3; ***, p value <0.001; ns, not significant; unpaired t-test). The colored lines indicate each mean value. FIG. 9C: Representative FACS plots of OCT4::EGFP⁺ cells from Opto-FGFR hESCs cultured in either +F or +L condition for 3 weeks. FIG. 9D, 9E: Representative images of the Opto-FGFR hESCs immunostained with the indicated antibodies after (FIG. 9D) 6 weeks or (FIG. 9E) 54 weeks maintenance in +L condition. FIG. 9F: A karyotype of Opto-FGFR hESCs cultured with blue light for 4 weeks showed no aneuploidy and no structural rearrangements. Scale bars, 100 μm .

[0038] FIG. 10A-10B. The freezing and thawing of Opto-FGFR hESCs cultured with blue light illumination. FIG. 10A: A schematic illustration of the freezing and thawing procedure of optically maintained hESCs (1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h). Opto-FGFR hESCs cultured with blue light illumination for 2 passages (2 weeks) were frozen into stock vials (stored in LN2 tank for 2 weeks) then thawed and maintained for 3 weeks with blue light illumination. FIG. 10B: A representative morphology of the resulting Opto-FGFR hESC colony (left) and the expression of NANOG (right). Scale bars, 100 μm .

[0039] FIG. 11A-11B. Gene expression profiles of pluripotency markers in Opto-FGFR hESCs at a single-cell level. FIG. 11A: A schematic illustration of Opto-FGFR hESC maintenance (+F, 10 ng/mL FGF2 protein treatment supplied daily; +L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h) and their single-cell qRT-PCR. OCT4::EGFP⁺ single-cells were captured by FACS sorting into a 96-well plate for qRT-PCR. FIG. 11B: qRT-PCR results of Opto-FGFR hESCs cultured in either +F or +L condition for 45 weeks (n=78 for +F, n=88 for +L; n.s., not significant; unpaired t-test). For box and whisker plots, the black line across the box represents the median, the box represents interquartile range and the whiskers represent the minimum and maximum.

[0040] FIG. 12A-12C. Scatter and volcano plots of global transcriptome analyses (RNA-seq). FIG. 12A: A schematic illustration of Opto-FGFR hESCs maintenance (+F, 10 ng/mL FGF2 protein treatment supplied daily; +L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 1 min. every 2 h). FIG. 12B: Scatter plot of global transcriptome changes between +F and +L conditions. The loge expression levels of genes from RNA-seq data are plotted. FPKM, fragments per kilobase of transcript per million mapped reads. FIG. 12C: Volcano plot

of global transcriptome changes between +F and +L conditions. The loge (fold change) expression levels (+F vs +L) and $-\log_{10}(\text{p value})$ of genes from RNA-seq data are plotted.

[0041] FIG. 13A-13F. The generation of opto-FGFR knock-in piPSCs. FIG. 13A: The number of transfected cells and colonies formed after puromycin selection in each trial.

[0042] FIG. 13B: Representative colony morphology of opto-FGFR knock-in piPSCs. FIG. 13C: Phosphorylation of pErk1/2 in Opto-FGFR piPSCs in response to blue light illumination (1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min). A.U., arbitrary unit. FIG. 13D: A schematic illustration of Opto-FGFR piPSCs maintenance (+L, 1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h). FIG. 13E: A representative image of alkaline phosphatase staining of undifferentiated piPSCs maintained with blue light illumination for 1 week. FIG. 13F: A karyotype of Opto-FGFR piPSC cultured with blue light for 3 weeks showed no aneuploidy and no structural rearrangements. Scale bars, 100 μm .

[0043] FIG. 14A-14E. Optogenetic activation of FGF signaling is sufficient to maintain pluripotency of hESCs even in a feeder-free culture system. FIG. 14A-14C: Representative images of optically maintained Opto-FGFR hESCs (1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h) without feeder cells for 3 weeks showed (FIG. 14A) undifferentiated colony morphology and expression of pluripotency markers (FIG. 14B) OCT4 and (FIG. 14C) NANOG. FIG. 14D, 14E: Representative images of optically maintained Opto-FGFR hESCs (1 $\mu\text{W}/\text{mm}^2$ at 470 nm for 5 min. every 2 h) without feeder cells for 8 weeks showed (FIG. 14D) undifferentiated colony morphology and (FIG. 14E) no aneuploidy and no structural rearrangements in their karyotype. Scale bars, 100 μm .

[0044] FIG. 15A-15C. Cry2 PHR-based opto-FGFR system. FIG. 15A: Phosphorylation of ERK1/2 in response to blue light illumination (3.4-34 $\mu\text{W}/\text{mm}^2$ at 470 nm for 1 min. every 10 min. for 3 cycles) in human embryonic kidney cells (293T) transiently transfected with Cry2 PHR-based opto-FGFR. PD173074 (0.2 μM) was used as an FGF signaling inhibitor. FIG. 15B: A representative image of hESCs transiently transfected with Cry2 PHR-based opto-FGFR. FIG. 15C: Phosphorylation of ERK1/2 in response to blue light illumination (34 $\mu\text{W}/\text{mm}^2$ at 470 nm for 1 min. every 10 min. for 3 cycles) in hESCs transiently transfected with Cry2 PHR-based opto-FGFR. A.U., arbitrary unit. Scale bars, 100 μm .

[0045] FIG. 16 illustrates a schematic diagram of a general embodiment of the illumination system 100 of the present invention. The illumination system 100 is adapted to illuminate cells, in particular, transformed PSCs with the vectors of the present invention, to stimulate the FGF pathway and maintain the cells as pluripotent. The illumination system 100 comprises, in one embodiment, LED illumination sources 101 which are embedded in the upper enclosure 104 which is shaped to accommodate a cell culture substrate 106 in which the transformed PSCs are maintained. In an embodiment, the LEDs are individually connected to a printed circuit board 102 which is connected to a microcontroller 103 placed in the lower enclosure 105 and can be fastened together with fasteners.

[0046] FIG. 17 illustrates a schematic diagram of an embodiment of the circuit board 102 and microcontroller 103 controlling the illumination sources 101 of the present illumination system.

[0047] FIG. 18 illustrates a schematic diagram of an embodiment of the fibroblast growth factor free pluripotent stem cell culture system 200. The stem cell culture system 200 comprises at least one illumination device 100 which is connected to a computer server 201 which controls the illumination parameters of the illumination sources 101 of the device 100 via wireless or wired connection to either a mobile client computer 202 or client computer 203 with additional control software.

DETAILED DESCRIPTION OF THE INVENTION

[0048] It is understood that the present invention is not limited to the particular methods and components, etc., described herein, as these may vary. It is also to be understood that the terminology used herein is used for the purpose of describing particular embodiments only, and is not intended to limit the scope of the present invention. It must be noted that as used herein and in the appended claims, the singular forms “a,” “an,” and “the” include the plural reference unless the context clearly dictates otherwise. Thus, for example, a reference to a “protein” is a reference to one or more proteins, and includes equivalents thereof known to those skilled in the art and so forth.

[0049] Unless defined otherwise, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Specific methods, devices, and materials are described, although any methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention.

[0050] All publications cited herein are hereby incorporated by reference including all journal articles, books, manuals, published patent applications, and issued patents. In addition, the meaning of certain terms and phrases employed in the specification, examples, and appended claims are provided. The definitions are not meant to be limiting in nature and serve to provide a clearer understanding of certain aspects of the present invention.

[0051] The present inventors have now developed a novel culture system for PSCs using optical induction of the FGF signaling pathway. The inventive fusion protein and illumination system maintained pluripotency of PSCs similar to the conventional culture system. Furthermore, the optically maintained PSCs displayed the ability to differentiate into three germ layers, demonstrating that the optical culture system can sustain the pluripotency of PSCs without exogenous FGF2 protein supplementation.

[0052] Previous reports have demonstrated the optical activation of receptor tyrosine kinases in immortalized cell lines and developing embryos as an initial step in evolving such optical technologies. Although a study using the optical control of ERK phosphorylation reported a perturbation of patterning and morphogenesis in *Drosophila* embryos, other studies did not report any biologically relevant phenotypes; therefore, translating these approaches for use with PSCs has remained challenging.

[0053] The present inventors have now extended the approach to human and porcine PSCs and provide clear evidence that the inventive opto-FGFR PSCs are sufficiently maintained by blue light illumination and without the need for FGF2 protein supplementation. Since there is no need to supplement FGF2 protein, the media does not need to be changed on a daily basis in the optical PSC culture system.

In feeder-free culture systems, very high concentrations of FGF2 (up to 100 ng/mL) are required for promoting self-renewal and inhibiting spontaneous differentiation of hPSCs.

[0054] Even in a feeder-free culture system, the inventors succeeded in maintaining pluripotency of hESCs using the inventive illuminating culture system (FIG. 14). Moreover, the optically maintained PSCs appeared to retain the normal differentiation potential of traditional PSCs.

[0055] There are various kinds of photoactivatable proteins available for optogenetic approaches. Using the photolyase homology region of the Cry2 protein (Cry2PHR39) and LOV domain. In the present invention, the photosensory LOV domain served as an initiator in the light-induced intermolecular signal transduction cascade of the FGF signaling pathway (termed as opto-FGFR signaling) for maintaining the pluripotency of mammalian PSCs, without FGF2 protein supplementation. Although the inventors showed that they could optically activate FGF signaling in human embryonic kidney cells (HEK293T) and hESCs transiently transfected with the Cry2 PHR-based opto-FGFR (FIG. 15), the LOV domain was selected as a photodimerizable domain because of two reasons: the LOV domain has a shorter length (LOV, 432 bp or 144 aa26 vs Cry2 PHR, 1,506 bp or 502 aa35), which is more favorable for knock-in, and LOV domain has a shorter dissociation time after light withdrawal (LOV $t_{112}=2.5$ min vs Cry2 PHR $t_{112}=5-6$ min), which is more advantageous for accurate adjustment of the opto-FGFR signaling in PSC, than others. It is important to select an appropriate photosensory domain as the light-sensing actuator module for a given optogenetic modulation system in PSCs.

[0056] While it is a widely accepted idea that FGF2 protein directly stimulates PSCs for their self-renewal, there are other possible mechanisms of action of FGF2, such as direct and indirect interaction and association with the IGF pathway. However, until now, there has been no experimental tool to assess this possibility. The present invention herein provides the first direct evidence that the molecular mechanism underlying the daily supplementation with recombinant FGF2 protein to maintain the pluripotency of PSC is solely through the activation of the FGF signaling pathway.

[0057] The prior art stem cell culture systems are heavily dependent on random distribution of expensive and thermo-unstable recombinant proteins in a dish, which always can jeopardize future mass production of human stem cells for a population-wide cell therapy. The present inventive opto-FGFR PSCs enable optical activation without any exogenous FGF2 recombinant proteins and offer spatiotemporally precise optical control of stem cell behaviors. Beyond implementing a new technology in the stem cell field, the present experimental data will help lead to the development of a potential therapeutic cure against many human diseases by modulating multiple signaling pathways other than FGF2 and maximizing the potential of human PSCs.

[0058] The prior art animal agriculture and husbandry methods cause significant environmental issues such as greenhouse gas emissions, fresh water consumption, and arable land usage. As an alternative, cellular agriculture or lab-grown meats have been introduced and social consent is growing due to ethical and environmentally friendly concepts. However, the production costs and use of animal-derived thermo-unstable recombinant proteins during the

cellular agriculture process are barriers to entering the market for mass consumption. Based on the inventors' calculation, recombinant FGF2 protein for 1 billion of PSC culture costs \$14,000 weekly. In addition, due to thermal and chemo-physical instabilities, FGF2 is recommended to be freshly added into the media and it requires additional labor and miscellaneous fees. In conclusion, the present inventive opto-FGFR system in livestock PSCs can be a realistic solution to decrease these production costs.

[0059] Therefore, in accordance with an embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain.

[0060] In an embodiment the LOV domain has the following nucleotide sequence (SEQ ID NO:7):

```
CCTGACTACAGTCTCGTGAAGGCTCTGCAAATGGCACAACAGAATTTGT
CATTACAGACGCCTCCCTCCCAGACAACCTATCGTCTACGCCAGTAGAG
GGTTTCTGACACTGACAGGCTATTCTCTCGACCAGATCCTGGGCAGGAAC
TGCAGGTTTCTGCAAGGGCCAGAAACAGACCCAAGAGCTGTGGATAAGAT
CAGGAATGCCATCACCAAGGCGTTGATACAGTGTCTGTCTGCTGAATT
ATAGACAGGATGGCACAACCTTCTGGAATCTCTTCTCGTGGCTGGACTC
AGAGATTCTAAGGGCAATATTGTCAACTACGTCGGAGTGCAGTCAAAGGT
GAGCGAAGATTATGCCAAGCTGCTGGTCAACGAGCAGAACATTGAGTACA
AAGGTGTGCGCACCAGTAACATGCTGCGCAGAAAG.
```

[0061] In accordance with another embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain comprising the nucleotide sequence of SEQ ID NO:6.

[0062] The term “chimeric” or “fusion” polypeptide or protein refers to a composition comprising at least one polypeptide or peptide sequence or domain that is chemically bound in a linear fashion with a second polypeptide or peptide domain. One embodiment of this invention is an isolated or recombinant nucleic acid molecule encoding a fusion protein comprising at least two domains, wherein the first domain comprises a polypeptide encoding FGFR1 protein, and the second domain comprising a polypeptide encoding a LOV protein.

[0063] In some embodiments, the fusion protein is designed for human PSCs.

[0064] In some alternative embodiments, the fusion protein is designed for porcine induced PSCs.

[0065] In accordance with an embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain, a promoter sequence and a sequence encoding a myristolation signal peptide (Myr).

[0066] In accordance with an embodiment, the present invention provides an expression vector comprising synthetic polynucleotide comprising a sequence encoding the

intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable proteins, including cryptochrome 2 (CRY2), light, oxygen, and voltage (LOV) domains, phytochrome B (PhyB) and UV-resistance locus 8 (UVR8) and its derivatives.

[0067] In some embodiments, the expression vector can be a plasmid.

[0068] In accordance with an embodiment, the present invention provides an expression vector comprising synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain comprising the nucleotide sequence of SEQ ID NO:6.

[0069] In accordance with another embodiment, the present invention provides a synthetic polynucleotide comprising a sequence encoding the intracellular domain of the fibroblast growth factor 1 receptor (FGFR1) fused to a sequence encoding a photoactivatable light-oxygen-voltage sensing (LOV) domain, a promoter sequence and a sequence encoding a myristolation signal peptide (Myr). In an embodiment, the AAV-Myr-opto-FGFR-LOV comprising the nucleotide sequence of SEQ ID NO:1:

```
tgcctttctctgaccagcattctctccctgggectgtgcegetttctgtc
tgcagcttgtggcctgggtcacctctacggctggcccagatccttccctg
ccgectccttcagggttcgctcttccctccactccctcttcccttgcctc
tgctgtgttgctgcccaggatgctcttccggagcacttccctctcgge
gctgcaccacgtgatgtcctctgagcggatcctccccgtgtctgggtcct
ctccgggcatctctcctccctcacccaaccccatgccgtcttcaactcgt
gggttcccttttccctctccttctggtggcctgtgccatctctcgcttctt
aggatggccttctccgacggatgtctcccttgcgteccgectccctctct
tgtaggcctgcatcatcaccgtttttctggacaaccccaaagtacccgt
ctccctggcttttagccactctccatcctcttgccttcttgcctggaca
ccccgttctcctgtggattcgggtcacctctcactcctttcatttgggca
gctccctaccccccttacctctctagtctgtgctagctcttccagcccc
ctgtcatggcatcttccaggggtccgagagctcagctagtcttcttctc
caaccggggccctatgtccacttcaggacagcatgtttgctgcctccag
ggatcctgtgtccccgagctgggaccaccttatattcccagggccggtta
atgtggctctggttctggttacttttatctgtccctccacccacagtg
gggcaagcttctgacctcttcttcttccacagggcctcgagagatct
ggcagcggagAGGGGCAGAGGAAGTCTTCTAACATGCGGTGACGTGGAGGA
GAATCCCGGCCCTAGGCTCGAGATGACCGAGTACAAGCCCACGGTGCGCC
TCGCCACCCGCGACGACGTCCCAGGGCCGTACGCACCTCGCCGCCGCG
TTCGCCGACTACCCCGCCACGCGCCACCGTCGATCCGGACCGCCACAT
CGAGCGGGTACCCGAGCTGCAAGAACTCTTCTCACGCGCGTCGGGCTCG
ACATCGGCAAGGTGTGGGTGCGGACGACGGCGCGCGGTGGCGGTCTGG
ACCACGCCGGAGAGCGTCAAGCGGGGGCGGTGTTGCCGAGATCGGCCC
```

- continued

GCGCATGGCCGAGTTGAGCGGTTCCCGGCTGGCCGCGCAGCAACAGATGG
AAGGCCTCCTGGCGCCGCACCCGGCCCAAGGAGCCCGCGTGGTTCTTGGCC
ACCGTCGGCGTCTCGCCCGACCACCAGGGCAAGGGTCTGGGCAGCGCCGT
CGTGCTCCCCGAGTGGAGGCGGCCGAGCGCGCCGGGTGCCCCGCTTCC
TGGAGACCTCCGCGCCCCGCAACCTCCCCTTCTACGAGCGGCTCGGCTTC
ACCGTCACCGCCGACGTGAGGTGCCCCAAGGACCGCGCACCTGGTGCAT
GACCCGCAAGCCCGGTGCCtgatctagagggccccgtttaaacccgctgat
cagcctcgactgtgccttctagttgccagccatctgttgtttgccctcc
cccggtgccttccttgaccctggaaggtgccactcccactgtcctttccta
ataaaatgaggaaattgcatcgcatgtgtctgagttaggtgtcattctattc
tgggggggtgggggtggggcaggacagcaagggggaggattgggaagacaat
agcaggcatgctggggatgcggtgggctctatgggtctcgacattgatta
ttgactagttattaatagtaataatcaattacggggtcattagttcatagecc
atatatggagttccgcgttacataaacttacggtaaatggccccgcctggct
gaccgccccaacgacccccgcccattgacgtcaataatgacgtatgttccc
atagtaacgccaatagggactttccattgacgtcaatgggtggagtat
acggtaaaactgcccacttggcagtacatcaagtgtatcatatgccaaagta
cgccccctattgacgtcaatgacggtaaatggccccgcctggcattatgcc
cagtacatgaccttatgggactttcctacttggcagtacatctacgtatt
agtcatcgctattaccatgggtgatgcgggttttggcagtacatcaatgggc
gtggatagcggtttgactcacggggatttccaagtctccaccccatgac
gtcaatgggagtttgttttggcaccaaaatcaacgggactttccaaaatg
tcgtaacaactccgccccattgacgcaaatgggcggtaggcggtgacggt
gggagggtctatataagcagagctctctggctaactagagaacccactgct
tactggcttatcgaaattaatacgactcactatagggagacacaagctgg
ctagcggtttaaacggggccctctagactcgagcggccgcggccaccATGGG
GAGTAGCAAGAGCAAGCCTAAGGACCCCAGCCAGCGCTCGACATGAAGA
GCGGCACCAAGAAGAGCGACTTCCATAGCCAGATGGCTGTGCACAAGCTG
GCCAAGAGCATCCCTCTGCGCAGACAGGTAACAGTGTGAGCTGACTCCAG
TGCATCCATGAACTCTGGGGTTCTCCTGGTTCGGCCCTCACGGCTCTCCT
CCAGCGGGACCCCCATGCTGGCTGGAGTCTCCGAATATGAGCTCCCTGAG
GATCCCCGCTGGGAGCTGCCACGAGACAGACTGGTCTTAGGCCAAACCACT
TGGCGAGGGCTGCTTCGGGCAGGTGGTGTGGCTGAGGCCATCGGGCTGG
ATAAGGACAAACCAACCGTGTGACCAAAGTGGCCGTGAAGATGTTGAAG
TCCGACGCAACGGAGAAGGACCTGTGCGATCTGATCTCGGAGATGGAGAT
GATGAAAATGATTGGGAAGCACAGAATATCATCAACCTTCTGGGAGCGT
GCACACAGGATGGTCCTCTTTATGTATTGTGGAGTACGCCTCCAAGGC
AATCTCCGGGAGTATCTACAGGCCCGGAGGCCCTCTGGGCTGGAGTACTG
CTATAACCCAGCCACAACCCCGAGGAACAGCTGTCTTCAAAGATCTGG

- continued

TATCCTGTGCCTATCAGGTGGCTCGGGGCATGGAGTATCTTGCCTCTAAG
AAGTGTATACACCGAGACCTGGCTGCTAGGAACGTCCTGGTGACCGAGGA
TAACGTAATGAAGATCGCAGACTTTGGCTTAGCTCGAGACATTATCATA
TCGACTACTACAAGAAAACCACCAACGGCCGGCTGCCTGTGAAGTGGATG
GCCCCTGAGGCGTTGTTTGACCGGATCTACACACACCAGAGCGATGTGTG
GTCTTTTGGAGTGCTCTTGTGGGAGATCTTCACTCTGGGTGGCTCCCCAT
ACCCCGGTGTGCCTGTGGAGGAACTTTCAAGCTGCTGAAGGAGGGTCAT
CGAATGGACAAGCCCAGTAACGTACCAATGAGCTGTACATGATGATGCG
GGACTGCTGGCATGCAGTGCCCTCTCAGAGACCTACGTTCAAGCAGTTGG
TGGAAGACCTGGACCGCATTGTGGCCTTGACCTCCAACCAGGAGTATCTG
GACCTGTCCATACCGCTGGACCAGTACTCACCCAGCTTTCCCGACACACG
GAGCTCCACCTGCTCCTCAGGGGAGGACTCTGTCTTCTCTCATGAGCCGT
ATACCTGGGAGCCCTGTCTGCCTCGACACCCCCACCCAGCTTGCCAACAGT
GGACTCAAACGGCGCGTTCGAGACCGGTCCTGACTACAGTCTCGTGAAGGC
TCTGCAAATGGCACAACAGAATTTTGTCTATTACAGACGCCTCCCTCCCAG
ACAACCCTATCGTCTACGCCAGTAGAGGGTTTCTGACACTGACAGGCTAT
TCTCTCGACCAGATCCTGGGCAGGAAGTGCAGGTTTCTGCAAGGGCCAGA
AACAGACCCCAAGAGCTGTGGATAAGATCAGGAATGCCATCACCAAAGGCG
TTGATACCAGTGTCTGTCTGCTGAATTATAGACAGGATGGCACAACTTTC
TGGAATCTCTTCTTCGTGGCTGGACTCAGAGATTCTAAGGGCAATATTGT
CAACTACGTCGGAGTGCAGTCAAAGGTGAGCGAAGATTATGCCAAGCTGC
TGGTCAACGAGCAGAACATTGAGTACAAAGGTGTGCGCACCAGTAACATG
CTGCGCAGAAAGCCCCGTGGATCCGGAGTCGACTATCCGTACGACGTACC
AGACTACGCACTCGACtaagaattccaccacactggactagtggatccga
gctcggtagccaagcttaagactagggacaggattggtgacagaaaagccc
catccttaggcctcctccttcctagtcctcctgatattgggtctaaccccc
acctcctgttaggcagattccttatctggtgacacacccccatttcctgg
agccatctctctccttgccagaaaccttaaggtttgcttacgatggagcc
agagaggatcctgggagggagagcttggcaggggggtgggagggaaagggg
ggatgcgtgacctgcccggttctcagtggccaccctgcgctacctctcc
cagaacctgagctgctctgacgcggctgtctgggtgcgtttcactgatcct
gggtgctgcagcttccttacacttcccaagaggagaagcagtttgga
caaatcagaataagttggctcctgagttctaactttggctcttcaccttt
ctagtccccaatttatattgttcctccgtgcgtcagttttacctgtgaga
taaggccagtagccagccccgccttgccagggctgtggtgaggagggggg
tgtccgtgtggaaaactccctttgtgagaatggtgcgtcctaggtgttca
ccaggctcgtggccgcctctactccctttctctttctccatccttctttcc
ttaaagagtccccagtgctatctgggacatattcctccgcccagagcagg
gtcccgttccttaaggccctgctctgggcttctgggtttgagtccttgg

-continued

caagcccaggagagggcgctcaggtctccctgtcccccttccctcggtccacc
atctcatgccccctgggtctcctgcccccttccctacaggggttccctgggtc
tgctct.

[0070] In some embodiments, the plasmid comprises a AAV-CAGGS-eGFP vector where the eGFP region was replaced by a region of the opto-FGFR1 domain from mFGFR1-LOV vector to create a AAV-opto-FGFR.

[0071] It will be understood by those of ordinary skill in the art that in one or more embodiments, one can knock-in the AAV-Myr-opto-FGFR-LOV plasmid into the AAV1 locus (safe harbor area with constitutively expressed gene). It will also be understood that in other embodiments, the plasmid can be incorporated into the genetically active loci of stem cells, including human stem cells.

[0072] As used herein, the term FGFRF1 means the protein encoded by the FGFR1 gene, and which is a member of the fibroblast growth factor receptor (FGFR) family. The amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein would consist of an extracellular region, composed of three immunoglobulin-like domains, a single hydrophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular portion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. A marked difference between this gene product and the other family members is its lack of a cytoplasmic tyrosine kinase domain. The result is a transmembrane receptor that could interact with other family members and potentially inhibit signaling. Multiple alternatively spliced transcript variants encoding the same isoform have been found for this gene

[0073] N-myristoylation is the attachment of a 14-carbon fatty acid, myristate, onto the N-terminal glycine residue of target proteins, catalyzed by N-myristoyltransferase (NMT), a ubiquitous and essential enzyme in eukaryotes. Many of the target proteins of NMT are crucial components of signalling pathways, and myristoylation typically promotes membrane binding that is essential for proper protein localization or biological function.

[0074] The term “variants” as used herein, means that the wild type amino acid sequences comprising the polypeptides of the compositions, may include substituted amino acids.

[0075] The term, “amino acid” includes the residues of the natural α -amino acids (e.g., Ala, Arg, Asn, Asp, Cys, Glu, Gln, Gly, His, Lys, Ile, Leu, Met, Phe, Pro, Ser, Thr, Trp, Tyr, and Val) in D or L form, as well as β -amino acids, synthetic and unnatural amino acids. Many types of amino acid residues are useful in the polypeptides and the invention is not limited to natural, genetically-encoded amino acids. Examples of amino acids that can be utilized in the peptides described herein can be found, for example, in Fasman, 1989, CRC Practical Handbook of Biochemistry and Molecular Biology, CRC Press, Inc., and the reference cited therein. Another source of a wide array of amino acid residues is provided by the website of RSP Amino Acids LLC.

[0076] The term “peptide” as used herein, includes a sequence of from four to sixteen amino acid residues in

which the α -carboxyl group of one amino acid is joined by an amide bond to the main chain (α - or β -)amino group of the adjacent amino acid. The peptides provided herein for use in the described and claimed methods and compositions can be cyclic.

[0077] In reference to the fusion polypeptide composition of the present invention, the functional portion can comprise, for instance, about 90%, 95%, or more, of the FGFR1 and/or LOV polypeptide.

[0078] The functional portion of the fusion polypeptide composition of the present invention can comprise additional amino acids at the amino or carboxy terminus of the portion, or at both termini, which additional amino acids are not found in the amino acid sequence of either of the wild type FGFR1 and/or LOV polypeptides. Desirably, the additional amino acids do not interfere with the biological function of the functional portion.

[0079] Included in the scope of the invention are functional variants of the inventive polypeptides, and proteins described herein. The term “functional variant” as used herein refers to either the FGFR1 and/or LOV polypeptide, or fusion protein having substantial or significant sequence identity or similarity to the FGFR1 and/or LOV polypeptide, or fusion protein, which functional variant retains the biological activity of the FGFR1 and/or LOV polypeptide, or fusion protein of which it is a variant. In reference to the original FGFR1 and/or LOV polypeptide, or protein, the functional variant can, for instance, be at least about 30%, 50%, 75%, 80%, 90%, 98% or more identical in amino acid sequence to the FGFR1 and/or LOV polypeptide, or protein.

[0080] The functional variant can, for example, comprise the amino acid sequence of the FGFR1 and/or LOV polypeptide fusion protein with at least one or more conservative amino acid substitutions. Conservative amino acid substitutions are known in the art, and include amino acid substitutions in which one amino acid having certain physical and/or chemical properties is exchanged for another amino acid that has the same chemical or physical properties. For instance, the conservative amino acid substitution can be an acidic amino acid substituted for another acidic amino acid (e.g., Asp or Glu), an amino acid with a nonpolar side chain substituted for another amino acid with a nonpolar side chain (e.g., Ala, Gly, Val, Ile, Leu, Met, Phe, Pro, Trp, Val, etc.), a basic amino acid substituted for another basic amino acid (Lys, Arg, etc.), an amino acid with a polar side chain substituted for another amino acid with a polar side chain (Asn, Cys, Gln, Ser, Thr, Tyr, etc.), etc.

[0081] Functional variants can also include extensions of the FGFR1 and/or LOV polypeptide fusion protein. For example, a functional variant of the FGFR1 and/or LOV polypeptide fusion protein can include 1, 2, 3, 4 and 5 additional amino acids from either the N-terminal or C-terminal end of the FGFR1 and/or LOV polypeptide fusion protein.

[0082] Alternatively or additionally, the functional variants can comprise the amino acid sequence of the FGFR1 and/or LOV polypeptide fusion protein with at least one non-conservative amino acid substitution. In this case, it is preferable for the non-conservative amino acid substitution to not interfere with or inhibit the biological activity of the functional variant. Preferably, the non-conservative amino acid substitution enhances the biological activity of the functional variant, such that the biological activity of the

functional variant is increased as compared to the FGFR1 and/or LOV polypeptide fusion protein.

[0083] The FGFR1 and/or LOV polypeptide fusion protein can consist essentially of the specified amino acid sequence or sequences described herein, such that other components of the functional variant, e.g., other amino acids, do not materially change the biological activity of the functional variant.

[0084] It will be understood by those of ordinary skill in the art that the orientation of the two proteins in the fusion protein construct can be reversed, i.e., the N-terminal protein can comprise the LOV protein and the C-terminal protein can comprise the FGFR1 protein.

[0085] Analogs and mimetics include molecules which include molecules which contain non-naturally occurring amino acids or which do not contain amino acids but nevertheless behave functionally the same as the peptide. Natural product screening is one useful strategy for identifying analogs and mimetics.

[0086] Examples of incorporating non-natural amino acids and derivatives during peptide synthesis include, but are not limited to, use of norleucine, 4-amino butyric acid, 4-amino-3-hydroxy-5-phenylpentanoic acid, 6-aminohexanoic acid, t-butylglycine, norvaline, phenylglycine, omithine, sarcosine, 4-amino-3-hydroxy-6-methylheptanoic acid, 2-thienyl alanine and/or D-isomers of amino acids. A partial list of known non-natural amino acid contemplated herein is shown in Table 1.

TABLE 1

Non-natural Amino Acids			
Non-conventional amino acid	Code	Non-conventional amino acid	Code
α-aminobutyric acid	Abu	L-N-methylalanine	Nmala
α-amino-α-methylbutyrate	Mgab	L-N-methylarginine	Nmarg
aminocyclopropane-carboxylate	Cpro	L-N-methylasparagine	Nmasn
aminoisobutyric acid	Aib	L-N-methylaspartic acid	Nmasp
aminonorbomyl-carboxylate	Norb	L-N-methyleysteine	Nmcys
cyclohexylalanine		L-N-methylglutamine	Nmgln
cyclopentylalanine	Cpen	L-N-methylglutamic acid	Nmglu
D-alanine	Dal	Chexa L-N-methylhistidine	Nmhis
D-arginine	Darg	L-N-methylisoleucine	Nmile
D-aspartic acid	Dasp	L-N-methylleucine	Nmleu
D-cysteine	Deys	L-N-methyllysine	Nmlys
D-glutamine	Dgln	L-N-methylmethionine	Nmmt
D-glutamic acid	Dglu	L-N-methylnorleucine	Nmnle
D-histidine	Dhis	L-N-methylnorvaline	Nmnva
D-isoleucine	Dile	L-N-methylornithine	Nmorn
D-leucine	Dleu	L-N-methylphenylalanine	Nmphe
D-lysine	Dlys	L-N-methylproline	Nmpro
D-methionine	Dmet	L-N-methylserine	Nmser
D-ornithine	Dorn	L-N-methylthreonine	Nmthr
D-phenylalanine	Dphe	L-N-methyltryptophan	Nmtrp
D-proline	Dpro	L-N-methyltyrosine	Nmtyr
D-serine	Dser	L-N-methylvaline	Nruval
D-threonine	Dthr	L-N-methylethylglycine	Nmetg
D-tryptophan	Dtrp	L-N-methyl-t-butylglycine	Nmtbug
D-tyrosine	Dtyr	L-norleucine	Nle
D-valine	Dval	L-norvaline	Nva
D-α-methylalanine	Dmala	α-methyl-aminoisobutyrate	Maib
D-α-methylarginine	Dmarg	α-methyl-γ-aminobutyrate	Mgab
D-α-methylasparagine	Dmasn	α-methylcyclohexylalanine	Mchexa
D-α-methylaspartate	Dmasp	α-methylcyclopentylalanine	Mcpen
D-α-methylcysteine	Dmcys	α-methyl-α-naphthylalanine	Manap
D-α-methylglutamine	Dmgln	α-methylpenicillamine	Mpen
D-α-methylhistidine	Dmhis	N-(4-aminobutyl)glycine	Nglu
D-α-methylisoleucine	Dmile	N-(2-aminoethyl)glycine	Naeg
D-α-methylleucine	Dmleu	N-(3-aminopropyl)glycine	Norn
D-α-methyllysine	Dmlys	N-amino-α-methylbutyrate	Nmaabu
D-α-methylmethionine	Dmmt	α-naphthylalanine	Anap
D-α-methylornithine	Dmorn	N-benzylglycine	Nphe
D-α-methylphenylalanine	Dmphe	N-(2-carbamylethyl)glycine	Ngln
D-α-methylproline	Dmpro	N-(carbamylmethyl)glycine	Nasn
D-α-methylserine	Dmser	N-(2-carboxyethyl)glycine	Nglu
D-α-methylthreonine	Dmthr	N-(carboxymethyl)glycine	Nasp
D-α-methyltryptophan	Dmtrp	N-cyclobutylglycine	Ncbut
D-α-methyltyrosine	Dmty	N-cycloheptylglycine	Nchep
D-α-methylvaline	Dmval	N-cyclohexylglycine	Nchex
D-N-methylalanine	Dnmala	N-cyclodecylglycine	Ncdec
D-N-methylarginine	Dnmarg	N-cyclododecylglycine	Ncdod
D-N-methylasparagine	Dnmasn	N-cyclooctylglycine	Ncoct
D-N-methylaspartate	Dnmasp	N-cyclopropylglycine	Nepro
D-N-methylcysteine	Druncys	N-cycloundecylglycine	Ncund
D-N-methylglutamine	Dnmgln	N-(2,2-diphenylethyl)glycine	Nbhm
D-N-methylglutamate	Dnmglu	N-(3,3-diphenylpropyl)glycine	Nbhe
		N-(3-guanidinopropyl)glycine	Narg
		N-(1-hydroxyethyl)glycine	Nthr

TABLE 1-continued

Non-natural Amino Acids			
Non-conventional amino acid	Code	Non-conventional amino acid	Code
D-N-methylhistidine	Dnmhis	N-(hydroxyethyl)glycine	Nser
D-N-methylisoleucine	Dnmile	N-(imidazolylethyl)glycine	Nhis
D-N-methylleucine	Dnmleu	N-(3-indolylethyl)glycine	Nhtrp
D-N-methyllysine	Dnmlys	N-methyl-γ-aminobutyrate	Nmgabu
N-methylcyclohexylalanine	Nmchexa	D-N-methylmethionine	Dnmmet
D-N-methylornithine	Dnmorn	N-methylcyclopentylalanine	Nmcpen
N-methylglycine	Nala	D-N-methylphenylalanine	Dumphe
N-methylaminoisobutyrate	Nmaib	D-N-methylproline	Dnmpro
N-(1-methylpropyl)glycine	Nile	D-N-methylserine	Dnmser
N-(2-methylpropyl)glycine	Nieu	D-N-methylthreonine	Dnmthr
D-N-methyltryptophan	Dnmtrp	N-(1-methylethyl)glycine	Nval
D-N-methyltyrosine	Druntyr	N-methyl-α-naphthylalanine	Nmanap
D-N-methylvaline	Dnmval	N-methylpenicillamine	Nmpen
γ-aminobutyric acid	Gabu	N-(p-hydroxyphenyl)glycine	Nhtyr
L-t-butylglycine	Tbug	N-(thiomethyl)glycine	Neys
L-ethylglycine	Etg	penicillamine	Pen
L-homophenylalanine	Hphe	L-α-methylalanine	Mala
L-α-methylarginine	Marg	L-α-methylasparagine	Masn
L-α-methylaspartate	Masp	L-α-methyl-t-butylglycine	Mtbug
L-α-methyleysteine	Meys	L-methylethylglycine	Metg
L-α-methylglutamine	Mgln	L-α-methylglutamate	Mglu
L-α-methylhistidine	Mhis	L-α-methylhomophenylalanine	Mhphe
L-α-methylisoleucine	Mile	N-(2-methylthioethyl)glycine	Nmet
L-α-methylleucine	Mien	L-α-methyllysine	Miys
L-α-methylmethionine	Mmet	L-α-methylnorleucine	Mule
L-α-methylnorvaline	Mnva	L-α-methylornithine	Morn
L-α-methylphenylalanine	Mphe	L-α-methylproline	Mpro
L-α-methylserine	Mser	L-α-methylthreonine	Mthr
L-α-methyltryptophan	Mtrp	L-α-methyltyrosine	Mtyr
L-α-methylvaline	Mval	L-N-methylhomophenylalanine	Nruhpe
N-(N-(2,2-diphenylethyl) carbamylmethyl)glycine	Nnbhm	N-(N-(3,3-diphenylpropyl) carbamylmethyl)glycine	Nnbhe
1-carboxy-1-(2,2-diphenyl-ethylamino)cyclopropane	Nmbc		

[0087] Analogs of the subject peptides contemplated herein include modifications to side chains, incorporation of non-natural amino acids and/or their derivatives during peptide synthesis and the use of crosslinkers and other methods which impose conformational constraints on the peptide molecule or their analogs.

[0088] Examples of side chain modifications contemplated by the present invention include modifications of amino groups such as by reductive alkylation by reaction with an aldehyde followed by reduction with NaBH₄; amidination with methylacetimidate; acylation with acetic anhydride; carbamoylation of amino groups with cyanate; trinitrobenzylation of amino groups with 2, 4, 6-trinitrobenzene sulphonic acid (TNBS); acylation of amino groups with succinic anhydride and tetrahydrophthalic anhydride; and pyridoxylation of lysine with pyridoxal-5-phosphate followed by reduction with NaBH₄.

[0089] The guanidine group of arginine residues may be modified by the formation of heterocyclic condensation products with reagents such as 2,3-butanedione, phenylglyoxal and glyoxal.

[0090] The carboxyl group may be modified by carbodiimide activation via O-acylisourea formation followed by subsequent derivitization, for example, to a corresponding amide.

[0091] Sulphydryl groups may be modified by methods such as carboxymethylation with iodoacetic acid or iodoacetamide; performic acid oxidation to cysteic acid; formation of a mixed disulphides with other thiol compounds; reaction with maleimide, maleic anhydride or other substituted male-

imide; formation of mercurial derivatives using 4-chloromercuribenzoate, 4-chloromercuriphenylsulphonic acid, phenylmercury chloride, 2-chloromercuri-4-nitrophenol and other mercurials; carbamoylation with cyanate at alkaline pH.

[0092] Tryptophan residues may be modified by, for example, oxidation with N-bromosuccinimide or alkylation of the indole ring with 2-hydroxy-5-nitrobenzyl bromide or sulphenyl halides. Tyrosine residues on the other hand, may be altered by nitration with tetranitromethane to form a 3-nitrotyrosine derivative.

[0093] Modification of the imidazole ring of a histidine residue may be accomplished by alkylation with iodoacetic acid derivatives or N-carbethoxylation with diethylpyrocarbonate.

[0094] Crosslinkers can be used, for example, to stabilize 3D conformations, using homo-bifunctional crosslinkers such as the bifunctional imido esters having (CH₂)_n spacer groups with n=1 to n=6, glutaraldehyde, N-hydroxysuccinimide esters and hetero-bifunctional reagents which usually contain an amino-reactive moiety such as N-hydroxysuccinimide and another group specific-reactive moiety such as maleimido or dithio moiety (SH) or carbodiimide (COOH). In addition, peptides can be conformationally constrained by, for example, incorporation of C_α and N_α-methylamino acids, introduction of double bonds between C_α and C_β atoms of amino acids and the formation of cyclic peptides or analogues by introducing covalent bonds such as forming an amide bond between the N and C termini, between two side chains or between a side chain and the N or C terminus.

[0095] In some embodiments, the FGFR1 protein in the fusion protein is mammalian. In certain embodiments, the FGFR1 protein can be murine, porcine, ovine, bovine, human, or combinations thereof.

[0096] In accordance with an embodiment, the present invention provides a composition comprising a polypeptide encoding human FGFR1 protein, or a functional portion or fragment, or variant thereof, linked to a polypeptide encoding LOV protein, or a functional portion or fragment, or variant thereof.

[0097] In accordance with an embodiment, the present invention provides expression vectors comprising the synthetic polynucleotides described herein. In one embodiment, the present invention provides an expression cassette specific for use with the CRISPR/Cas9 system. In an embodiment, a gRNA sequence (GTCCCCTCCACCCACAGTG) (SEQ ID NO:9 is used to target AAVS1 locus.

[0098] A “CRISPR,” “CRISPR system system,” or “CRISPR nuclease system” and their grammatical equivalents can include a non-coding RNA molecule (e.g., guide RNA) that binds to DNA and Cas proteins (e.g., Cas9) with nuclease functionality (e.g., two nuclease domains). See, e.g., Sander, J. D., et al., “CRISPR-Cas systems for editing, regulating and targeting genomes,” *Nature Biotechnology*, 32:347-355 (2014); see also e.g., Hsu, P. D., et al., “Development and applications of CRISPR-Cas9 for genome engineering,” *Cell* 157(6):1262-1278 (2014).

[0099] The term “gene editing” and its grammatical equivalents as used herein can refer to genetic engineering in which one or more nucleotides are inserted, replaced, or removed from a genome. Gene editing can be performed using a nuclease (e.g., a natural-existing nuclease or an artificially engineered nuclease).

[0100] The term “safe harbor” and “immune safe harbor”, and their grammatical equivalents as used herein can refer to a location within a genome that can be used for integrating exogenous nucleic acids wherein the integration does not cause any significant effect on the growth of the host cell by the addition of the nucleic acid alone. Non-limiting examples of safe harbors can include HPRT, AAVS SITE (e.g., AAVS1, AAVS2, etc.), CCR5, or Rosa26.

[0101] Site specific gene editing can be achieved using non-viral gene editing such as CRISPR, TALEN (see U.S. patent application Ser. No. 14/193,037), transposon-based, ZEN, meganuclease, or Mega-TAL, or Transposon-based system. For example, PiggyBac (see Moriarty, B. S., et al., “Modular assembly of transposon integratable multigene vectors using RecWay assembly,” *Nucleic Acids Research* (8):e92 (2013) or sleeping beauty (see Aronovich, E. L, et al., “The Sleeping Beauty transposon system: a non-viral vector for gene therapy,” *Hum. Mol. Genet.*, 20(R1): R14-R20. (2011) transposon systems can be used.

[0102] Site specific gene editing can also be achieved without homologous recombination. An exogenous polynucleic acid can be introduced into a cell genome without the use of homologous recombination. In some cases, a transgene can be flanked by engineered sites that are complementary to a targeted double strand break region in a genome. A transgene can be excised from a polynucleic acid so it can be inserted at a double strand break region without homologous recombination.

CRISPR System

[0103] Methods described herein can take advantage of a CRISPR system. There are at least five types of CRISPR systems which all incorporate RNAs and Cas proteins. Types I, III, and IV assemble a multi-Cas protein complex that is capable of cleaving nucleic acids that are complementary to the crRNA. Types I and III both require pre-crRNA processing prior to assembling the processed crRNA into the multi-Cas protein complex. Types II and V CRISPR systems comprise a single Cas protein complexed with at least one guiding RNA.

[0104] The general mechanism and recent advances of CRISPR system is discussed in Cong, L. et al., “Multiplex genome engineering using CRISPR systems,” *Science*, 339 (6121): 819-823 (2013); Fu, et al., “High-frequency off-target mutagenesis induced by CRISPR-Cas nucleases in human cells,” *Nature Biotechnology*, 31, 822-826 (2013); Chu, V T et al. “Increasing the efficiency of homology-directed repair for CRISPR-Cas9-induced precise gene editing in mammalian cells,” *Nature Biotechnology* 33, 543-548 (2015); Shmakov, S. et al., “Discovery and functional characterization of diverse Class 2 CRISPR-Cas systems,” *Molecular Cell*, 60, 1-13 (2015); Makarova, K S et al., “An updated evolutionary classification of CRISPR-Cas systems,” *Nature Reviews Microbiology*, 13, 1-15 (2015). Site-specific cleavage of a target DNA occurs at locations determined by both 1) base-pairing complementarity between the guide RNA and the target DNA (also called a protospacer) and 2) a short motif in the target DNA referred to as the protospacer adjacent motif (PAM). For example, an engineered cell can be generated using a CRISPR system, e.g., a type II CRISPR system. A Cas enzyme used in the methods disclosed herein can be Cas9, which catalyzes DNA cleavage. Enzymatic action by Cas9 derived from *Streptococcus pyogenes* or any closely related Cas9 can generate double stranded breaks at target site sequences which hybridize to 20 nucleotides of a guide sequence and that have a protospacer-adjacent motif (PAM) following the 20 nucleotides of the target sequence.

[0105] Cas Protein

[0106] A vector can be operably linked to an enzyme-coding sequence encoding a CRISPR enzyme, such as a Cas protein (CRISPR-associated protein). Non-limiting examples of Cas proteins can include Cas 1, Cas 1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9 (also known as Csn 1 or Csx12), Cas 10, Csy1, Csy2, Csy3, Cse 1, Cse2, Csc 1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb 1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx1S, Csf1, Csf2, CsO, Csf4, Cpf1, c2c 1, c2c3, Cas9HiFi, homologues thereof, or modified versions thereof. An unmodified CRISPR enzyme can have DNA cleavage activity, such as Cas9. A CRISPR enzyme can direct cleavage of one or both strands at a target sequence, such as within a target sequence and/or within a complement of a target sequence. For example, a CRISPR enzyme can direct cleavage of one or both strands within or within about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 50, 100, 200, 500, or more base pairs from the first or last nucleotide of a target sequence. A vector that encodes a CRISPR enzyme that is mutated with respect to a corresponding wild-type enzyme such that the mutated CRISPR enzyme lacks the ability to cleave one or both strands of a target polynucleotide containing a target

sequence can be used. A Cas protein can be a high fidelity cas protein such as Cas9HiFi.

[0107] A vector that encodes a CRISPR enzyme comprising one or more nuclear localization sequences (NLSs), such as more than or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, NLSs can be used. For example, a CRISPR enzyme can comprise more than or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, NLSs at or near the ammo-terminus, more than or more than about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, NLSs at or near the carboxyl-terminus, or any combination of these (e.g., one or more NLS at the ammo-terminus and one or more NLS at the carboxyl terminus). When more than one NLS is present, each can be selected independently of others, such that a single NLS can be present in more than one copy and/or in combination with one or more other NLSs present in one or more copies.

[0108] Cas9 can refer to a polypeptide with at least or at least about 50%, 60%, 70%, 80%, 90%, 100% sequence identity and/or sequence similarity to a wild type exemplary Cas9 polypeptide (e.g., Cas9 from *S. pyogenes*). Cas9 can refer to a polypeptide with at most or at most about 50%, 60%, 70%, 80%, 90%, 100% sequence identity and/or sequence similarity to a wild type exemplary Cas9 polypeptide (e.g., from *S. pyogenes*). Cas9 can refer to the wild type or a modified form of the Cas9 protein that can comprise an amino acid change such as a deletion, insertion, substitution, variant, mutation, fusion, chimera, or any combination thereof.

[0109] A polynucleotide encoding an endonuclease (e.g., a Cas protein such as Cas9) can be codon optimized for expression in particular cells, such as eukaryotic cells. This type of optimization can entail the mutation of foreign-derived (e.g., recombinant) DNA to mimic the codon preferences of the intended host organism or cell while encoding the same protein.

[0110] CRISPR enzymes used in the methods can comprise NLSs. The NLS can be located anywhere within the polypeptide chain, e.g., near the N- or C-terminus. For example, the NLS can be within or within about 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50 amino acids along a polypeptide chain from the N- or C-terminus. Sometimes the NLS can be within or within about 50 amino acids or more, e.g., 100, 200, 300, 400, 500, 600, 700, 800, 900, or 1000 amino acids from the N- or C-terminus.

[0111] An endonuclease can comprise an amino acid sequence having at least or at least about 50%, 60%, 70%, 75%, 80%, 85%, 90%, 95%, 99%, or 100%, amino acid sequence identity to the nuclease domain of a wild type exemplary site-directed polypeptide (e.g., Cas9 from *S. pyogenes*).

[0112] While *S. pyogenes* Cas9 (SpCas9) is commonly used as a CRISPR endonuclease for genome engineering, it may not be the best endonuclease for every target excision site. For example, the PAM sequence for SpCas9 (5' NGG 3') is abundant throughout the human genome, but a NGG sequence may not be positioned correctly to target a desired gene for modification. In some cases, a different endonuclease may be used to target certain genomic targets. In some cases, synthetic SpCas9-derived variants with non-NGG PAM sequences may be used. Additionally, other Cas9 orthologues from various species have been identified and these "non-SpCas9s" bind a variety of PAM sequences that could also be useful for the present invention. For example, the relatively large size of SpCas9 (approximately 4 kb

coding sequence) means that plasmids carrying the SpCas9 cDNA may not be efficiently expressed in a cell. Conversely, the coding sequence for *Staphylococcus aureus* Cas9 (Sa-Cas9) is approximately 1 kilobase shorter than SpCas9, possibly allowing it to be efficiently expressed in a cell. Similar to SpCas9, the SaCas9 endonuclease is capable of modifying target genes in mammalian cells in vitro and in mice in vivo.

[0113] Alternatives to *S. pyogenes* Cas9 may include RNA-guided endonucleases from the Cpf1 family that display cleavage activity in mammalian cells. Unlike Cas9 nucleases, the result of Cpf1-mediated DNA cleavage is a double-strand break with a short 3' overhang. Cpf1's staggered cleavage pattern may open up the possibility of directional gene transfer, analogous to traditional restriction enzyme cloning, which may increase the efficiency of gene editing. Like the Cas9 variants and orthologues described above, Cpf1 may also expand the number of sites that can be targeted by CRISPR to AT-rich regions or AT-rich genomes that lack the NGG PAM sites favored by SpCas9.

[0114] Any functional concentration of Cas protein can be introduced to a cell. For example, 15 micrograms of Cas mRNA can be introduced to a cell. In other cases, a Cas mRNA can be introduced from 0.5 micrograms to 100 micrograms. A Cas mRNA can be introduced from 0.5, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 micrograms.

[0115] Guide RNA

[0116] As used herein, the term "guide RNA (gRNA)", and its grammatical equivalents can refer to an RNA which can be specific for a target DNA and can form a complex with a Cas protein. A guide RNA can comprise a guide sequence, or spacer sequence, that specifies a target site and guides an RNA/Cas complex to a specified target DNA for cleavage. For example, FIG. 15, demonstrates that guide RNA can target a CRISPR complex to three genes and perform a targeted double strand break. Site-specific cleavage of a target DNA occurs at locations determined by both 1) base-pairing complementarity between a guide RNA and a target DNA (also called a protospacer) and 2) a short motif in a target DNA referred to as a protospacer adjacent motif (PAM).

[0117] A method disclosed herein also can comprise introducing into a cell or embryo at least one guide RNA or nucleic acid, e.g., DNA encoding at least one guide RNA. A guide RNA can interact with a RNA-guided endonuclease to direct the endonuclease to a specific target site, at which site the 5' end of the guide RNA base pairs with a specific protospacer sequence in a chromosomal sequence.

[0118] A guide RNA can comprise two RNAs, e.g., CRISPR RNA (crRNA) and transactivating crRNA (tracrRNA). A guide RNA can sometimes comprise a single-guide RNA (sgRNA) formed by fusion of a portion (e.g., a functional portion) of crRNA and tracrRNA. A guide RNA can also be a dual RNA comprising a crRNA and a tracrRNA. A guide RNA can comprise a crRNA and lack a tracrRNA. Furthermore, a crRNA can hybridize with a target DNA or protospacer sequence.

[0119] As discussed above, a guide RNA can be an expression product. For example, a DNA that encodes a guide RNA can be a vector comprising a sequence coding for the guide RNA. A guide RNA can be transferred into a cell or organism by transfecting the cell or organism with an isolated guide RNA or plasmid DNA comprising a sequence

coding for the guide RNA and a promoter. A guide RNA can also be transferred into a cell or organism in other way, such as using virus-mediated gene delivery.

[0120] A guide RNA can be isolated. For example, a guide RNA can be transfected in the form of an isolated RNA into a cell or organism. A guide RNA can be prepared by in vitro transcription using any in vitro transcription system. A guide RNA can be transferred to a cell in the form of isolated RNA rather than in the form of plasmid comprising encoding sequence for a guide RNA.

[0121] A guide RNA can comprise a DNA-targeting segment and a protein binding segment. A DNA-targeting segment (or DNA-targeting sequence, or spacer sequence) comprises a nucleotide sequence that can be complementary to a specific sequence within a target DNA (e.g., a proto-spacer). A protein-binding segment (or protein-binding sequence) can interact with a site-directed modifying polypeptide, e.g., an RNA-guided endonuclease such as a Cas protein. By “segment” it is meant a segment/section/region of a molecule, e.g., a contiguous stretch of nucleotides in an RNA. A segment can also mean a region/section of a complex such that a segment may comprise regions of more than one molecule. For example, in some cases a protein-binding segment of a DNA-targeting RNA is one RNA molecule and the protein-binding segment therefore comprises a region of that RNA molecule. In other cases, the protein-binding segment of a DNA-targeting RNA comprises two separate molecules that are hybridized along a region of complementarity.

[0122] A guide RNA can comprise two separate RNA molecules or a single RNA molecule. An exemplary single molecule guide RNA comprises both a DNA-targeting segment and a protein-binding segment.

[0123] An exemplary two-molecule DNA-targeting RNA can comprise a crRNA-like (“CRISPR RNA” or “targeter-RNA” or “crRNA” or “crRNA repeat”) molecule and a corresponding tracrRNA-like (“trans-acting CRISPR RNA” or “activator-RNA” or “tracrRNA”) molecule. A first RNA molecule can be a crRNA-like molecule (targeter-RNA), that can comprise a DNA-targeting segment (e.g., spacer) and a stretch of nucleotides that can form one half of a double-stranded RNA (dsRNA) duplex comprising the protein-binding segment of a guide RNA. A second RNA molecule can be a corresponding tracrRNA-like molecule (activator-RNA) that can comprise a stretch of nucleotides that can form the other half of a dsRNA duplex of a protein-binding segment of a guide RNA. In other words, a stretch of nucleotides of a crRNA-like molecule can be complementary to and can hybridize with a stretch of nucleotides of a tracrRNA-like molecule to form a dsRNA duplex of a protein-binding domain of a guide RNA. As such, each crRNA-like molecule can be said to have a corresponding tracrRNA-like molecule. A crRNA-like molecule additionally can provide a single stranded DNA-targeting segment, or spacer sequence. Thus, a crRNA-like and a tracrRNA-like molecule (as a corresponding pair) can hybridize to form a guide RNA. A subject two-molecule guide RNA can comprise any corresponding crRNA and tracrRNA pair.

[0124] A DNA-targeting segment or spacer sequence of a guide RNA can be complementary to sequence at a target site in a chromosomal sequence, e.g., protospacer sequence) such that the DNA-targeting segment of the guide RNA can base pair with the target site or protospacer. In some cases,

a DNA-targeting segment of a guide RNA can comprise from or from about 10 nucleotides to from or from about 25 nucleotides or more. For example, a region of base pairing between a first region of a guide RNA and a target site in a chromosomal sequence can be or can be about 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 22, 23, 24, 25, or more than 25 nucleotides in length. Sometimes, a first region of a guide RNA can be or can be about 19, 20, or 21 nucleotides in length.

[0125] A guide RNA can target a nucleic acid sequence of or of about 20 nucleotides. A target nucleic acid can be less than or less than about 20 nucleotides. A target nucleic acid can be at least or at least about 5, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30 or more nucleotides. A target nucleic acid can be at most or at most about 5, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30 or more nucleotides. A target nucleic acid sequence can be or can be about 20 bases immediately 5' of the first nucleotide of the PAM. A guide RNA can target the nucleic acid sequence.

[0126] A guide nucleic acid, for example, a guide RNA, can refer to a nucleic acid that can hybridize to another nucleic acid, for example, the target nucleic acid or proto-spacer in a genome of a cell. A guide nucleic acid can be RNA. A guide nucleic acid can be DNA. The guide nucleic acid can be programmed or designed to bind to a sequence of nucleic acid site-specifically. A guide nucleic acid can comprise a polynucleotide chain and can be called a single guide nucleic acid. A guide nucleic acid can comprise two polynucleotide chains and can be called a double guide nucleic acid.

[0127] A guide nucleic acid can comprise one or more modifications to provide a nucleic acid with a new or enhanced feature. A guide nucleic acid can comprise a nucleic acid affinity tag. A guide nucleic acid can comprise synthetic nucleotide, synthetic nucleotide analog, nucleotide derivatives, and/or modified nucleotides.

[0128] A guide nucleic acid can comprise a nucleotide sequence (e.g., a spacer), for example, at or near the 5' end or 3' end, that can hybridize to a sequence in a target nucleic acid (e.g., a protospacer). A spacer of a guide nucleic acid can interact with a target nucleic acid in a sequence-specific manner via hybridization (i.e., base pairing). A spacer sequence can hybridize to a target nucleic acid that is located 5' or 3' of a protospacer adjacent motif (PAM). The length of a spacer sequence can be at least or at least about 5, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30 or more nucleotides. The length of a spacer sequence can be at most or at most about 5, 10, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 30 or more nucleotides.

[0129] A guide RNA can also comprises a dsRNA duplex region that forms a secondary structure. For example, a secondary structure formed by a guide RNA can comprise a stem (or hairpin) and a loop. A length of a loop and a stem can vary. For example, a loop can range from about 3 to about 10 nucleotides in length, and a stem can range from about 6 to about 20 base pairs in length. A stem can comprise one or more bulges of 1 to about 10 nucleotides. The overall length of a second region can range from about 16 to about 60 nucleotides in length. For example, a loop can be or can be about 4 nucleotides in length and a stem can be or can be about 12 base pairs. A dsRNA duplex region can comprise a protein-binding segment that can form a complex with an RNA-binding protein, such as a RNA-guided endonuclease, e.g., Cas protein.

[0130] A guide RNA can also comprise a tail region at the 5' or 3' end that can be essentially single-stranded. For example, a tail region is sometimes not complementary to any chromosomal sequence in a cell of interest and is sometimes not complementary to the rest of a guide RNA. Further, the length of a tail region can vary. A tail region can be more than or more than about 4 nucleotides in length. For example, the length of a tail region can range from or from about 5 to from or from about 60 nucleotides in length.

[0131] A guide RNA can be introduced into a cell or embryo as an RNA molecule. For example, a RNA molecule can be transcribed in vitro and/or can be chemically synthesized. A guide RNA can then be introduced into a cell or embryo as an RNA molecule. A guide RNA can also be introduced into a cell or embryo in the form of a non-RNA nucleic acid molecule, e.g., DNA molecule. For example, a DNA encoding a guide RNA can be operably linked to promoter control sequence for expression of the guide RNA in a cell or embryo of interest. A RNA coding sequence can be operably linked to a promoter sequence that is recognized by RNA polymerase III (Pol III).

[0132] A DNA molecule encoding a guide RNA can also be linear. A DNA molecule encoding a guide RNA can also be circular.

[0133] A DNA sequence encoding a guide RNA can also be part of a vector. Some examples of vectors can include plasmid vectors, phagemids, cosmids, artificial/mini-chromosomes, transposons, and viral vectors. For example, a DNA encoding a RNA-guided endonuclease is present in a plasmid vector. Other non-limiting examples of suitable plasmid vectors include pUC, pBR322, pET, pBluescript, and variants thereof. Further, a vector can comprise additional expression control sequences (e.g., enhancer sequences, Kozak sequences, polyadenylation sequences, transcriptional termination sequences, etc.), selectable marker sequences (e.g., antibiotic resistance genes), origins of replication, and the like.

[0134] When both a RNA-guided endonuclease and a guide RNA are introduced into a cell as DNA molecules, each can be part of a separate molecule (e.g., one vector containing fusion protein coding sequence and a second vector containing guide RNA coding sequence) or both can be part of a same molecule (e.g., one vector containing coding (and regulatory) sequence for both a fusion protein and a guide RNA).

[0135] A Cas protein, such as a Cas9 protein or any derivative thereof, can be pre-complexed with a guide RNA to form a ribonucleoprotein (RNP) complex. The RNP complex can be introduced into primary immune cells. Introduction of the RNP complex can be timed. The cell can be synchronized with other cells at G1, S, and/or M phases of the cell cycle. The RNP complex can be delivered at a cell phase such that HDR is enhanced. The RNP complex can facilitate homology directed repair.

[0136] A guide RNA can also be modified. The modifications can comprise chemical alterations, synthetic modifications, nucleotide additions, and/or nucleotide subtractions. The modifications can also enhance CRISPR genome engineering. A modification can alter chirality of a gRNA. In some cases, chirality may be uniform or stereopure after a modification. A guide RNA can be synthesized. The synthesized guide RNA can enhance CRISPR genome engineering. A guide RNA can also be truncated. Truncation can be used to reduce undesired off-target mutagenesis. The truncation

can comprise any number of nucleotide deletions. For example, the truncation can comprise 1, 2, 3, 4, 5, 10, 15, 20, 25, 30, 40, 50 or more nucleotides. A guide RNA can comprise a region of target complementarity of any length. For example, a region of target complementarity can be less than 20 nucleotides in length. A region of target complementarity can be more than 20 nucleotides in length.

[0137] In some cases, a dual nickase approach may be used to introduce a double stranded break. Cas proteins can be mutated at known amino acids within either nuclease domains, thereby deleting activity of one nuclease domain and generating a nickase Cas protein capable of generating a single strand break. A nickase along with two distinct guide RNAs targeting opposite strands may be utilized to generate a DSB within a target site (often referred to as a “double nick” or “dual nickase” CRISPR system). This approach may dramatically increase target specificity, since it is unlikely that two off-target nicks will be generated within close enough proximity to cause a DSB.

[0138] In some cases, a GUIDE-Seq analysis can be performed to determine the specificity of engineered guide RNAs. The general mechanism and protocol of GUIDE-Seq profiling of off-target cleavage by CRISPR system nucleases is discussed in Tsai, S. et al., “GUIDE-Seq enables genome-wide profiling of off-target cleavage by CRISPR system nucleases,” *Nature*, 33: 187-197 (2015).

[0139] A gRNA can be introduced at any functional concentration. For example, a gRNA can be introduced to a cell at 10 micrograms. In other cases, a gRNA can be introduced from 0.5 micrograms to 100 micrograms. A gRNA can be introduced from 0.5, 5, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65, 70, 75, 80, 85, 90, 95, or 100 micrograms.

[0140] In some cases, a method can comprise an endonuclease selected from the group consisting of Cas 1, Cas1B, Cas2, Cas3, Cas4, Cas5, Cas6, Cas7, Cas8, Cas9, Cas 10, Csy 1, Csy2, Csy3, Cse 1, Cse2, Csc1, Csc2, Csa5, Csn2, Csm2, Csm3, Csm4, Csm5, Csm6, Cmr1, Cmr3, Cmr4, Cmr5, Cmr6, Csb 1, Csb2, Csb3, Csx17, Csx14, Csx10, Csx16, CsaX, Csx3, Csx1, Csx1S, Csf1, Csf2, CsO, Csf4, Cpf1, c2c1, c2c3, Cas9HiFi, homologues thereof or modified versions thereof. A Cas protein can be Cas9. In some cases, a method can further comprise at least one guide RNA (gRNA). A gRNA can comprise at least one modification. An exogenous TCR can bind a cancer neo-antigen.

[0141] Disclosed herein is a method of making an engineered cell comprising: introducing at least one polynucleic acid encoding at least one exogenous LOV receptor sequence; introducing at least one guide RNA (gRNA) comprising at least one modification; and introducing at least one endonuclease; wherein the gRNA comprises at least one sequence complementary to at least one endogenous genome. In some cases, a modification is on a 5' end, a 3' end, from a 5' end to a 3' end, a single base modification, a 2'-ribose modification, or any combination thereof. A modification can be selected from a group consisting of base substitutions, insertions, deletions, chemical modifications, physical modifications, stabilization, purification, and any combination thereof.

[0142] In accordance with an embodiment, the present invention provides a transformant transformed by the vectors described herein, including, for example, mammalian cells, and specifically, pluripotent stem cells.

[0143] In some embodiments, the transformant is a stem cell. In other embodiments, the transformant is a human derived stem cell.

[0144] In further embodiments, the transformant is a stem cell from a mammal, such as a pig, cow, sheep, goat, chicken or other livestock animal.

[0145] The invention further provides a host cell comprising any of the recombinant expression vectors described herein. As used herein, the term “host cell” refers to any type of cell that can contain the inventive recombinant expression vector. The host cell can be a eukaryotic cell, e.g., plant, animal, fungi, or algae. The host cell can be a cultured cell or a primary cell, i.e., isolated directly from an organism, e.g., a human or pig. The host cell can be an adherent cell or a suspended cell, i.e., a cell that grows in suspension. Suitable host cells are known in the art and include, for instance, stem cells. In some preferred embodiments, the host cells are pluripotent stem cells of human or porcine derivation.

[0146] Also provided by the invention is a population of cells comprising at least one host cell described herein. The population of cells can be a substantially homogeneous population, in which the population comprises mainly of host cells (e.g., consisting essentially of) comprising the recombinant expression vector. The population also can be a clonal population of cells, in which all cells of the population are clones of a single host cell comprising a recombinant expression vector, such that all cells of the population comprise the recombinant expression vector. In one embodiment of the invention, the population of cells is a clonal population comprising host cells comprising a recombinant expression vector as described herein.

[0147] The host referred to in the inventive methods can be any host. Preferably, the host is a mammal. As used herein, the term “mammal” refers to any mammal, including, but not limited to, mammals of the order Rodentia, such as mice and hamsters, and mammals of the order Lagomorpha, such as rabbits. It is preferred that the mammals are from the order Carnivora, including Felines (cats) and Canines (dogs). It is more preferred that the mammals are from the order Artiodactyla, including Bovine (cows) and Swine (pigs) or of the order Perissodactyla, including Equine (horses). It is most preferred that the mammals are of the order Primates, Ceboidea, or Simiiformes (monkeys) or of the order Anthropoidea (humans and apes). An especially preferred mammal is the pig.

[0148] Stem cells for use in the methods disclosed herein are not limited in any way and may be obtained from any source. Preferably the stem cells are pluripotent stem cells that are obtained from a mammalian source.

[0149] The term “nucleic acid” as used herein, includes “polynucleotide,” “oligonucleotide,” and “nucleic acid molecule,” and generally means an isolated or purified polymer of DNA or RNA, which can be single-stranded or double-stranded, synthesized or obtained (e.g., isolated and/or purified) from natural sources, which can contain natural, non-natural or altered nucleotides, and which can contain a natural, non-natural or altered internucleotide linkage, such as a phosphoramidate linkage or a phosphorothioate linkage, instead of the phosphodiester found between the nucleotides of an unmodified oligonucleotide. In some embodiments, the nucleic acid does not comprise any insertions, deletions, inversions, and/or substitutions. However, it may be suitable in some instances, as discussed herein, for

the nucleic acid to comprise one or more insertions, deletions, inversions, and/or substitutions.

[0150] Preferably, the nucleic acids of the invention are recombinant. As used herein, the term “recombinant” refers to (i) molecules that are constructed outside living cells by joining natural or synthetic nucleic acid segments to nucleic acid molecules that can replicate in a living cell, or (ii) molecules that result from the replication of those described in (i) above. For purposes herein, the replication can be in vitro replication or in vivo replication.

[0151] The nucleic acids can be constructed based on chemical synthesis and/or enzymatic ligation reactions using procedures known in the art. See, for example, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 3rd ed., Cold Spring Harbor Press, Cold Spring Harbor, N.Y. 2001; and Ausubel et al., *Current Protocols in Molecular Biology*, Greene Publishing Associates and John Wiley & Sons, NY, 1994. For example, a nucleic acid can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed upon hybridization (e.g., phosphorothioate derivatives and acridine substituted nucleotides). Examples of modified nucleotides that can be used to generate the nucleic acids include, but are not limited to, 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N⁶-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N⁶-substituted adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N⁶-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, 3-(3-amino-3-N-2-carboxypropyl) uracil, and 2,6-diaminopurine. Alternatively, one or more of the nucleic acids of the invention can be purchased from companies, such as Macromolecular Resources (Fort Collins, Colo.) and Synthesgen (Houston, Tex.).

[0152] The invention also provides an isolated or purified nucleic acid comprising a nucleotide sequence which is complementary to the nucleotide sequence of any of the nucleic acids described herein or a nucleotide sequence which hybridizes under stringent conditions to the nucleotide sequence of any of the nucleic acids described herein.

[0153] The term “isolated” as used herein means having been removed from its natural environment. The term “purified” as used herein means having been increased in purity, wherein “purity” is a relative term, and not to be necessarily construed as absolute purity. For example, the purity can be at least about 50%, can be greater than 60%, 70% or 80%, or can be 100%.

[0154] The term “primer” refers to an oligonucleotide, whether natural or synthetic, capable of acting as a point of initiation of DNA synthesis under conditions in which synthesis of a primer extension product complementary to a nucleic acid strand is induced, i.e., in the presence of four different nucleoside triphosphates and an agent for polym-

erization (i.e., DNA polymerase or reverse transcriptase) in an appropriate buffer and at a suitable temperature. A primer is preferably a single-stranded oligodeoxyribonucleotide. The appropriate length of a primer depends on the intended use of the primer but typically ranges from about 10 to about 30 nucleotides. Short primer molecules generally require cooler temperatures to form sufficiently stable hybrid complexes with the template. A primer need not reflect the exact sequence of the template but must be sufficiently complementary to specifically hybridize with a template. When primer pairs are referred to herein, the pair is meant to include one forward primer which is capable of hybridizing to the sense strand of a double-stranded target nucleic acid (the “sense primer”) and one reverse primer which is capable of hybridizing to the antisense strand of a double-stranded target nucleic acid (the “antisense primer”).

[0155] “Probe” refers to an oligonucleotide which binds through complementary base pairing to a sub-sequence of a target nucleic acid. A primer may be a probe. It will be understood by one of skill in the art that probes will typically substantially bind target sequences lacking complete complementarity with the probe sequence depending upon the stringency of the hybridization conditions. The probes are typically directly labeled (e.g., with isotopes or fluorescent moieties) or indirectly labeled such as with biotin to which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the target, by Southern blot for example.

[0156] The recombinant expression vectors of the invention can be prepared using standard recombinant DNA techniques described in, for example, Sambrook et al., supra, and Ausubel et al., supra. Constructs of expression vectors, which are circular or linear, can be prepared to contain a replication system functional in a prokaryotic or eukaryotic host cell. Replication systems can be derived, e.g., from ColEI, 2 μ plasmid, λ , SV40, bovine papilloma virus, and the like.

[0157] In some exemplary embodiments, the CRISPR system has been used in this system to permanently incorporate the opto-FGFR system into the human stem cells. We specifically target the AAVS1 locus (a safe-harbor region with constitutive expression).

[0158] Desirably, the recombinant expression vector comprises regulatory sequences, such as transcription and translation initiation and termination codons, which are specific to the type of host (e.g., bacterium, fungus, plant, or animal) into which the vector is to be introduced, as appropriate and taking into consideration whether the vector is DNA or RNA based.

[0159] The recombinant expression vector can include one or more marker genes, which allow for selection of transformed or transfected hosts. Marker genes include biocide resistance, e.g., resistance to antibiotics, heavy metals, etc., complementation in an auxotrophic host to provide prototrophy, and the like. Suitable marker genes for the inventive expression vectors include, for instance, neomycin/G418 resistance genes, hygromycin resistance genes, histidinol resistance genes, tetracycline resistance genes, puromycin resistance genes and ampicillin resistance genes.

[0160] The selection of promoters, e.g., strong, weak, inducible, tissue-specific and developmental-specific, is within the ordinary skill of the artisan. Similarly, the combining of a nucleotide sequence with a promoter is also

within the skill of the artisan. The promoter can be a non-viral promoter or a viral promoter, e.g., a cytomegalovirus (CMV) promoter, the tet-on promoter, or the ubiquitin C promoter, for example.

[0161] Fibroblast growth factor (FGF) free pluripotent stem cell culture systems.

[0162] Currently, there are a few prior art culture optogenetics systems on the market, but these systems are all very expensive. High cost commercial products prevent large volume optogenetics studies and thus, limit the rate of research in this field. There is also a lack of instrumentation that allows for precise control of illumination and remote monitoring of multiple culture plates.

[0163] In accordance with some embodiments, the fibroblast growth factor free pluripotent stem cell culture system of the present invention is a hardware, software, and mobile application system capable of precise control and real-time monitoring of cell culture illumination that can serve as an inexpensive alternative to high cost commercial products. The inventive illumination system is designed to be easy-to-assemble, easy-to-use, and low-cost, utilizing 3D printing of thermoplastic materials, programmable open-source electronics, and other readily-available electronic parts. Additionally, the inventive illumination system was designed to be adjustable and be customized to the individual researcher. The 3D models and source code are publicly available to be edited for any specific needs. Additionally, the LED illuminating device of the inventive illumination system can be customized with any color LED to be used with any photosynthetic protein of interest. The inventive illumination system provides the opportunity to apply a flexible optical culture system to cell cultures in any laboratory.

[0164] As such, in accordance with an embodiment, the present invention provides a fibroblast growth factor free pluripotent stem cell culture system comprising one or more transformed cells as described herein and an illumination source.

[0165] In accordance with an embodiment, the present invention provides a cell culture illumination system comprising a cell culture substrate, an upper and lower enclosure, one or more illumination sources, one or more circuit boards, a power source, and a microcontroller.

[0166] Referring now to FIG. 16, in general, the cell culture illumination system 100 is designed in such a way that it is easily implemented, for example for use with standard size cell culture plates. The illumination system comprises one or more illumination sources 101 of any type which are capable of irradiating light of a tunable frequency into each well, and to the cells contained within each well of a cell culture plate 106. In some embodiments the illumination source is one or more tunable LEDs. The illumination sources are connected electronically to at least one circuit board 102 and are arranged inside an upper enclosure 104 which is designed to accept the cell culture plate. The circuit board 102 is connected electronically to a microcontroller 103 which controls the light frequency and intensity. The microcontroller 103 and upper enclosure 104 are designed to fit into a lower enclosure 105 and can be held together with fasteners such as screws. The microcontroller 103 is coupled to a power source 107 (see FIG. 17) such as a USB port from a computer. Other power sources could also be used, including for example, wired to a common power outlet or that can be re-chargeable such as a battery.

[0167] As shown in FIGS. 16-18, the microcontroller 103 is connected either by wire or wirelessly to a main computer server 200. The main computer server 200 can be programmed with information related to the illumination frequency, intensity and duration.

[0168] It should be noted that the microcontroller 103 and the main computer server 200 can all include a computing device such as a microprocessor, hard drive, solid state drive, smartphone or any other suitable computing device known to or conceivable by one of skill in the art. As shown in FIG. 18, the main computer server 200 can be controlled by a computing client device 203, or a smartphone or mobile computing device 202 which is programmed with a non-transitory computer readable medium that is programmed with steps to execute the different illumination frequencies, intensities, and durations of irradiation.

[0169] Any such computer application will be fixed on a non-transitory computer readable medium. It should be noted that the computer application is programmed onto a non-transitory computer readable medium that can be read and executed by any of the computing devices mentioned in this application. The non-transitory computer readable medium can take any suitable form known to one of skill in the art. The non-transitory computer readable medium is understood to be any article of manufacture readable by a computer or smartphone. Such non-transitory computer readable media includes, but is not limited to, magnetic media, such as floppy disk, flexible disk, hard, disk, reel-to-reel tape, cartridge tape, cassette tapes or cards, optical media such as CD-ROM, DVD, blu-ray, writable compact discs, magneto-optical media in disc, tape, or card form, and paper media such as punch cards or paper tape. Alternately, the program for executing the method and algorithms of the present invention can reside on a remote server or other networked device. Any databases associated with the present invention can be housed on a central computing device, server(s), in cloud storage, or any other suitable means known to or conceivable by one of skill in the art. All of the information associated with the application is transmitted either wired or wirelessly over a network, via the internet, cellular telephone network, or any other suitable data transmission means known to or conceivable by one of skill in the art.

[0170] In accordance with an embodiment, the present invention provides a method for maintaining a pluripotent stem cell line in culture comprising: a) a plurality of pluripotent stem cells in a fibroblast growth factor free culture medium, wherein said are transformed with the expression vectors described herein; b) said transformed cells are maintained in the cell culture system as described herein; and c) periodically irradiating the cells of b) at a wavelength of about 470 nm for a sufficient time and intensity such that the cells maintain their pluripotent capability.

[0171] In some exemplary embodiments the intensity at the maximal output in each well of a 6-well plate was between 1 and 23 $\mu\text{W}/\text{mm}^2$, with the ability to vary the intensity from 2, 3, 4, 5, 10, 15, 20 or greater $\mu\text{W}/\text{mm}^2$. The LEDs can be illuminated for different lengths of time (for example, but without limitation, 0 to 10 min at 1 $\mu\text{W}/\text{mm}^2$) and intensity (for example, but without limitation, 0 to 1 $\mu\text{W}/\text{mm}^2$ for 5 min). For the long term maintenance of Opto-FGFR hPSCs, the blue light illumination can be

repeated 1 min, 5 min, or 10 min or longer in every 1-10 hours, preferably about 2 hours at 1 $\mu\text{W}/\text{mm}^2$ during cell culture.

[0172] The present inventive approaches defined herein for the optical maintenance of human and porcine PSCs provides new insights for an alternative synthetic approach for the next generation in stem cell research. Furthermore, considering the number of signaling pathways that are crucial for controlling cell specification processes, the opto-FGFR PSCs of the present invention are a proof-of-concept for the optical control over a plethora of mammalian stem cell fates.

[0173] The following examples have been included to provide guidance to one of ordinary skill in the art for practicing representative embodiments of the presently disclosed subject matter. In light of the present disclosure and the general level of skill in the art, those of skill can appreciate that the following examples are intended to be exemplary only and that numerous changes, modifications, and alterations can be employed without departing from the scope of the presently disclosed subject matter. The synthetic descriptions and specific examples that follow are only intended for the purposes of illustration, and are not to be construed as limiting in any manner to make compounds of the disclosure by other methods.

[0174] Without further elaboration, it is believed that one skilled in the art, using the preceding description, can utilize the present invention to the fullest extent. The following examples are illustrative only, and not limiting of the remainder of the disclosure in any way whatsoever.

EXAMPLES

[0175] The following examples are put forth so as to provide those of ordinary skill in the art with a complete disclosure and description of how the compounds, compositions, articles, devices, and/or methods described and claimed herein are made and evaluated, and are intended to be purely illustrative and are not intended to limit the scope of what the inventors regard as their invention. Efforts have been made to ensure accuracy with respect to numbers (e.g., amounts, temperature, etc.) but some errors and deviations should be accounted for herein. Unless indicated otherwise, parts are parts by weight, temperature is in degrees Celsius or is at ambient temperature, and pressure is at or near atmospheric. There are numerous variations and combinations of reaction conditions, e.g., component concentrations, desired solvents, solvent mixtures, temperatures, pressures and other reaction ranges and conditions that can be used to optimize the product purity and yield obtained from the described process. Only reasonable and routine experimentation will be required to optimize such process conditions.

Materials and Methods

[0176] Pluripotent cell line culture. The human embryonic stem cells (H9, WiCell) and human induced pluripotent stem cells (induced from GM00024C, fibroblasts from Coriell) were cultured using standard protocols [26]. The hESCs and hiPSCs were cultured on mouse embryonic fibroblasts (MEFs, Applied StemCell) and pre-plated at 12,000 to 15,000 cells/cm². The medium contained DMEM/F12, 20% knockout serum replacement, 1 mM L-glutamine, 100 μM MEM non-essential amino acids, and 0.1 mM β -mercaptoethanol (Life Technologies). Ten ng/mL of FGF2 (R&D

Systems) was added after sterile filtration, and the cells were fed daily and passaged weekly using either 6 U/mL Dispase (Life Technologies) or mechanical means. For feeder-free culture conditions, hPSCs were cultured onto recombinant human Laminin 521 (Life Technologies)-coated plates with StemFit® Basic02 medium (Ajinomoto). The porcine induced pluripotent stem cells (induced from P350-05; PCASMC, porcine coronary artery smooth muscle cells from Sigma-Aldrich) were cultured with mitotically inactivated MEFs in the PSC medium, and the cells were fed daily and passaged weekly using either 6 U/mL Dispase or mechanical means. For routine cell culture, all experiments including passaging were performed in a dark culture room with safelight that does not affect LOV photodimerization.

[0177] Plasmid construct. To establish the Opto-FGFR hESC and hiPSC lines, the EGFP region of the AAV-CAGGS-EGFP vector (Addgene plasmid #22212) was replaced by a region of the opto-FGFR1 domain. The PHR domain from the cytoplasmic FGFR1-Cry2 PHR-mCitrine vector and the LOV domain from the mFGFR1-LOV vector (Addgene plasmid #58745) were used for the construction of AAV-opto-FGFR. AAV-opto-FGFRs were then used to establish the opto-FGFR targeted to the AAVS1 region. Two guide RNAs (gRNAs) sequences for opto-FGFR knock-in at the pRosa26 region were chosen to target the first intron of pRosa26 and subcloned into gRNA Cloning Vector (Addgene plasmid #41824) as described previously [27]. The sequences for pRosa26 gRNA constructs were as follows: pRosa26-gRNA #1, 5'-CCTGGCTTAAACCTGATTCTT-3' (SEQ ID NO:10); pRosa26-gRNA #91, 50-GTGAGAGT-TATCTGACCGTA-3' (SEQ ID NO:11). All insert sequences were verified by Sanger DNA sequencing (JHU Synthesis & Sequencing Facility).

TABLE 2		
Sequences of Primer Used in This Study		
Genes	Forward	Reverse
DPPA3	5'-TTAATCCAACCTA CATCCCAGGG-3' (SEQ ID NO: 12)	5'-AGGGGAAACAGAT TCGCTACTA-3' (SEQ ID NO: 13)
ESG1	5'-ATATCCC GCCGTG GGTGAAAGTTC-3' (SEQ ID NO: 14)	5'-ACTCAGCCATGGA CTGGAGCATCC-3' (SEQ ID NO: 15)
FOXA2	5'-CGACTGGAGCAGC TACTATGC-3' (SEQ ID NO: 16)	5'-TACGTGTTTCATGC CGTTCAT-3' (SEQ ID NO: 17)
GAPDH	5'-CGAGATCCCTCCA AAATCAA-3' (SEQ ID NO: 18)	5'-TTCTAGACGGCAG GTCAGGT-3' (SEQ ID NO: 19)
KLF4	5'-TCCCATCTTTCTC CACGTTT-3' (SEQ ID NO: 20)	5'-GGATCGGATAGGT GAAGCTG-3' (SEQ ID NO: 21)
MSGN1	5'-AGCGAAGGCTGCA GTGTC-3' (SEQ ID NO: 22)	5'-TGGCCTCTCTGGC TGTAGAC-3' (SEQ ID NO: 23)
NANOG	5'-CATGAGTGTGGAT CCAGCTTG-3' (SEQ ID NO: 24)	5'-CCTGAATAAGCAG ATCCATGG-3' (SEQ ID NO: 25)
OCT4	5'-AGTGAGAGGCAAC CTGGAGA-3' (SEQ ID NO: 26)	5'-ACACTCGGACCAC ATCCTTC-3' (SEQ ID NO: 27)

TABLE 2-continued		
Sequences of Primer Used in This Study		
Genes	Forward	Reverse
REX1	5'-CAGATCCTAAACA GCTCGCAGAAT-3' (SEQ ID NO: 28)	5'-GCGTACGCAAATT AAAGTCCAGA-3' (SEQ ID NO: 29)
PAX6	5'-CTTTGCTTGGGAA ATCCGAG-3' (SEQ ID NO: 30)	5'-AGCCAGGTTGCGA AGAACTC-3' (SEQ ID NO: 31)
SOX1	5'-CCTCGGATCTCTG GTCAAGT-3' (SEQ ID NO: 32)	5'-GCAGGTACATGCT GATCATCTC-3' (SEQ ID NO: 33)
SOX2	5'-GGGAAATGGGAGG GGTGCAAAGAGG-3' (SEQ ID NO: 34)	5'-TTGCGTGAGTGTG GATGGGATTGGTG-3' (SEQ ID NO: 35)
SOX17	5'-CGCACGGAATTTG AACAGTA-3' (SEQ ID NO: 36)	5'-GGATCAGGGACCT GTCACAC-3' (SEQ ID NO: 37)
TBX6	5'-AAGTACCAACCCC GCATACA-3' (SEQ ID NO: 38)	5'-TAGGCTGTCACGG AGATGAA-3' (SEQ ID NO: 39)
pGapdh	5'-CTCAACGGGAAGC TCACTG-3' (SEQ ID NO: 40)	5'-CCCTGTTGCTGTA GCCAAAT-3' (SEQ ID NO: 41)
pGata4	5'-GCTTCGCGGGCTC CTACT-3' (SEQ ID NO: 42)	5'-CCGGTTGATGCCA TTCAT-3' (SEQ ID NO: 43)
pLin28a	5'-TGCCGGCATCTGT AAATGGT-3' (SEQ ID NO: 44)	5'-GCAGTTTGCATTC CTTGGCA-3' (SEQ ID NO: 45)
pMsgn1	5'-CGCTGGAGTCCTA TTCTTCG-3' (SEQ ID NO: 46)	5'-GTCTGTGAGTTCC CCGATGT-3' (SEQ ID NO: 47)
pNanog	5'-TTGCCCCGAAGCA TCCATT-3' (SEQ ID NO: 48)	5'-CCAGCTCTGATTA CCCCACA-3' (SEQ ID NO: 49)
pPax6	5'-AGTTCTTCGCAAC CTGGCTA-3' (SEQ ID NO: 50)	5'-CATTTGGCCCTTC GATTAGA-3' (SEQ ID NO: 51)
pOct4	5'-TGAGGCTTTGCAG CTCAGTT-3' (SEQ ID NO: 52)	5'-ACTGCTTGATCGT TTGCCCT-3' (SEQ ID NO: 53)
pSox1	5'-CACCCGGATTACA AGTACCG-3' (SEQ ID NO: 54)	5'-GAGTTGGAGATGG GGCTGTA-3' (SEQ ID NO: 55)
pSox2	5'-TAAGTACACACTG CCCGGAG-3' (SEQ ID NO: 56)	5'-CATGGAACCGAG CGTCATGC-3' (SEQ ID NO: 57)
pSox17	5'-TCGGGGACATGAA GATGAAG-3' (SEQ ID NO: 58)	5'-GCGGCCGGTACTT GTAGTT-3' (SEQ ID NO: 59)
pTbx6	5'-AGTATCAGCCCCG CATACAC-3' (SEQ ID NO: 60)	5'-CTGCTCGGGATCT GACTCTC-3' (SEQ ID NO: 61)

[0178] Gene targeting. The hESC, hiPSC, and piPSC lines were cultured in Rho kinase (ROCK) inhibitor (Calbiochem) for 24 h before nucleofection [28]. The cells were harvested

using Accutase (Innovative Cell Technologies), and 2×10^6 cells were resuspended in the appropriate reagent (Lonza) with 1 μ g Cas9 plasmid (Addgene plasmid #41815), 1 μ g guide RNA (Addgene plasmid #41817, for opto-FGFR knock-in at AAVS1 locus; pRosa26-gRNA #1 or #91, for opto-FGFR knock-in at pRosa26 locus), and 1 μ g donor plasmid. Nucleofection was performed according to the manufacturer's instructions (Lonza). The cells were subsequently plated on puro-resistant MEFs (Applied StemCell), and puromycin was used for colony selection. The puromycin-resistant colonies were manually picked and expanded.

[0179] Blue light illumination. A customized blue light illumination plate (TouchBright W-Series) was designed and manufactured by Live Cell Instrument (Seoul, Korea). This plate contained 42 individual LEDs (70 mW per LED) per well on a 6-well plate. The light intensity was controlled by software (Live Cell Instrument), and the actual light intensity at 470 nm to the cell plate was measured by LaserCheck (Coherent). The intensity at the maximal output in each well of a 6-well plate was 23 μ W/mm². The LEDs were illuminated for different lengths of time (0-10 min. at 1 μ W/mm²) and intensity (0-1 μ W/mm² for 5 min). For the long-term maintenance of Opto-FGFR hPSCs with blue light during the passaging, the blue light illumination was repeated at 1, 5, or 10 min. every 2 h at 1 μ W/mm² during cell culture. The DIY LED illuminating device contained 1 individual LED (60 mW per LED) per well on a 6- or 12-well plate. The maximal output in each well of a 6-well plate was 7 μ W/mm².

[0180] Western blot analysis. The phosphorylation status of MEK and ERK1/2 were determined by Western blot analysis. The cells were starved from FGF2 protein for 24 h, and then exposed to either FGF2 protein or blue light illumination. One μ g aliquots of whole cell lysates were loaded onto 4-20% gradient SDS-PAGE gels and then transferred to nitrocellulose membranes (Bio-Rad) using the Trans-Blot® Turbo™ Transfer System (Bio-Rad). The detection of phosphorylated ERK1/2 proteins (42 and 44 kDa) was performed using a phospho-ERK1/2 antibody (Cell Signaling), total ERK1/2 expression level was determined using a total phospho-ERK1/2 antibody (Cell Signaling), detection of pMEK protein (45 kDa) was performed using a pMEK1/2 antibody (Cell Signaling), and total MEK expression level was determined using a total MEK1/2 antibody (Cell Signaling). The expression levels are quantified by normalizing to GAPDH expression (Cell Signaling). Following the primary antibody incubation, the membranes were incubated with the appropriate secondary antibodies and subsequently visualized with an ECL kit (Pierce).

[0181] Transcriptome analysis. Briefly, total RNA was extracted from cell pellets using TRIzol Reagent (Life Technologies), and 1 μ g of RNA was reverse transcribed using the High Capacity cDNA Reverse Transcription Kit (Applied Biosystems). The qRT-PCR mixtures were prepared with SYBR Green PCR Master Mix (Kapa Biosystem), and the reactions were performed using the Mastercycler ep Realplex2 (Eppendorf). The transcription levels were assessed by normalizing to GAPDH expression. The RNA-seq libraries were constructed using TruSeq Stranded Total RNA RiboZero Gold sample Prep Kit (Illumina) following the manufacturer's protocol. After the sequencing run, the Illumina Real Time Analysis (RTA) module was used to perform image analysis and base calling, and the

BCL Converter Software (bcl2fastq v2.17.1.14) was used to generate FASTQ files that contained the sequence reads. The pair-end reads of cDNA sequences were aligned back to the human genome (UCSC hg19 from Illumina iGenome) by the spliced read mapper TopHat (v2.0.9). The sequencing depth is over 90 million (45 million paired-ends) mappable sequencing reads. The alignment statistics and Q/C was achieved by SAMtools (v0.1.18) and RSeQC (v2.3.5) to calculate quality control metrics on the resulting aligned reads; this provides information on the mappability, uniformity of gene body coverage, insert length distributions, and junction annotation. The heatmap from the RNA-seq data was generated followed the manufacturer's instructions (Partek® Genomics Suite®, version 6.6). To create the heatmap, the undifferentiated-experimental and differentiated-control groups were compared to generate the 8132 differentially expressed gene set, which was corrected to a fold change of greater or less than two and a false discovery rate (FDR) with p value < 0.05 . Next, a clustering analysis of this heatmap was performed using the hierarchical clustering method with standardized expression normalization (a default process in the Partek® Genomics Suite®, version 6.6). The single-cell qRT-PCR was performed as previously described [29]. Briefly, OCT4::EGFP+single-cells were isolated by fluorescence-activated cell sorting (FACS) into a 96-well plate containing 10 μ g of lysis buffer. These plates were then sealed for single-cell reverse transcription and cDNA amplification using a universal sequence from the adaptor. The final amplified cDNA was used for qRT-PCR.

[0182] FACS analysis. The cells were dissociated with Accutase for flow cytometry and analyzed using BD FACSCalibur (Becton Dickinson), and the data was visualized using FlowJo software (Tree Star Inc.).

[0183] In vitro differentiation into three germ layers. Opto-FGFR hPSCs cultured with blue light for more than 6 weeks were differentiated into three germ layer lineages following the manufacturer's protocol (STEMdiff™ Trilineage Differentiation Kit, Catalog #05230), and the immunofluorescent staining and RT-PCR for the representative markers of each lineage were performed at the recommended time point.

[0184] Directed differentiation of hPSCs. Undifferentiated hPSCs were dissociated into single cells using Accutase to differentiate into desired cell types. For directed differentiation of hPSC-derived myotubes and hPSC-derived dopaminergic neurons, we followed previously described protocols [26,30].

[0185] Teratoma formation assay. The teratomas were generated by the intramuscular injection of Opto-FGFR hESCs and Opto-FGFR hiPSCs cultured with blue light for more than 8 weeks into NOD SCID mice. In brief, 5×10^6 cells were harvested and injected into the hindlimb of 6-week-old male NOD SCID mice. Approximately 7 weeks after injection, the tumors were dissected and fixed in 10% formalin (Sigma-Aldrich); the paraffin sections were prepared and stained with hematoxylin/eosin. The differentiated tissues that represent the three embryonic germ layers were identified.

[0186] LED illuminating device and software. The LED illuminating device consists of 6 or 12 LED diodes (EDGELEC), a printed circuit board (Elegoo), a microcontroller (Elegoo), and a 3D printed plastic casing. The LEDs were soldered onto the circuit board, which was wired to the microcontroller. The plastic casing was designed using com-

puter-aided design software (AutoCAD) and printed with a 3D plastic printer (Monoprice 3D Select) using polylactic acid plastic (HATCHBOX). The casing was secured together using metal screws (Snug Fastener). The components for the LED illuminating device should be accurately printed and assembled to fix the distance between the cells and the light, among the devices. The software to control the LED illuminating device and the mobile application were developed in Processing 3.5.3.

[0187] Statistical analysis. All statistical analyses were performed using Prism 6 (GraphPad). The values were the results of at least three independent experiments, with multiple replicates of each, and reported as the mean $\pm$ SEM. Differences between two samples were analyzed for significance using the unpaired two-tailed Student's t-test in Prism (GraphPad).

Results

Generation of a Novel FGF2-Free Human PSC Culture System

[0188] Generation of a light-inducible FGF signaling system. To create a light-inducible FGF signaling system for maintaining the pluripotency of PSCs, the intracellular domain of FGFR1 was fused with a photoactivatable LOV domain (discovered in *Vaucheria fridida*) and a myristoylation signal peptide (Myr) [31-35] (FIGS. 1A and B). This opto-FGFR cassette was inserted into the AAVS1 safe-harbor locus in OCT4::EGFP human embryonic stem cells (hESCs) [36] and human iPSCs (hiPSCs) using the CRISPR/Cas9 system [27,37] (FIG. 1C and FIG. 6A; Opto-FGFR hESC/hiPSC lines). The isolated Opto-FGFR hESC/hiPSC clones exhibited a typical colony morphology and expressed the pluripotency marker NANOG (FIGS. 6B and 6C). To test the effects of blue light illumination on the established Opto-FGFR hPSCs (FIG. 6D), we compared the phosphorylation levels of ERK1/2 and MEK after treatment with FGF2 protein or blue light (470 nm) illumination because phosphorylation of MEK and ERK1/2 are markers for activation of the FGF signaling pathway in hESCs [12,15]. After 24 h of FGF2 protein starvation, the cells were harvested at 15 min. intervals up to 120 min. following 10 ng/mL of FGF2 supplementation. For the blue light illumination, cells were collected at the same time points following 1 μ W/mm² of blue light illumination for 5 min. Western blot analyses demonstrated activated FGF signaling in both conditions (FIGS. 6E and 6F).

[0189] Optimization of the optical induction of FGF signaling. Next, we optimized the light illumination conditions and found that 5 min. of illumination was sufficient to induce FGF signaling, and the blue light exposure resulted in faster activation of FGF signaling than FGF2 supplementation (FIGS. 6E and 6F). The effects of the different exposure times (0-10 min. at 1 μ W/mm²) and intensities of blue light (0-1 μ W/mm² for 5 min) on ERK1/2 phosphorylation were measured, and the results showed the saturation and tunability of blue light in opto-FGFR (FIG. 7). In addition, we found that the blue light illumination-based activation of FGF signaling in Opto-FGFR hESCs showed similar levels of protein phosphorylation in multiple downstream targets as those treated with recombinant FGF2 protein (FIG. 8). These data suggest that the optical induction of FGF sig-

naling via the LOV module has effects comparable to the activation via FGF2 protein supplementation in hPSC culture.

[0190] Long-term validation of the light-inducible FGF signaling system. The FGF2 protein is a critical medium component for maintaining the undifferentiated state in hPSCs; however, in the daily feeding condition, its concentration decreased by approximately 40% in 4 h and approximately 10% in 24 h after the initial feeding [38] (FIG. 1D, +F, positive control; FGF2 protein treatment). To repeatedly stimulate the FGF signaling at the appropriate level, we maintained Opto-FGFR hPSCs with a pulsed (not continued) blue light (FIG. 1D, +L, blue light illumination; 1 μ W/mm², illumination of blue light for 1 min. or 5 min. in every 2 h). While the Opto-FGFR hESCs cultured in the dark were fully differentiated after 2 weeks, the cells cultured with pulsed blue light maintained hESC-colony morphologies for more than 1 year without FGF2 supplementation (FIG. 1D). The optically maintained hESCs from multiple passages showed a normal karyotype and comparable expression levels of pluripotency markers, including OCT4 and NANOG mRNAs and OCT4, NANOG, and SSEA4 proteins, to that of the cells maintained with FGF2 supplementation (FIGS. 1F and 1G and FIG. 9). Notably, the Opto-FGFR hiPSCs optically maintained for 12 weeks still expressed multiple pluripotency markers, OCT4 and NANOG mRNAs, and TRA-1-81 and NANOG proteins (FIGS. 1G and 1H). Moreover, the Opto-FGFR hESCs maintained with blue light illumination were successfully recovered after a freeze/thaw cycle (FIG. 10). To compare the transcriptional distribution of key pluripotency markers in hPSCs maintained with either pulsed blue light or FGF2 protein in more detail, we further profiled the gene expression of pluripotency markers at a single-cell level. The cells maintained with either condition for over 45 weeks were purified to OCT4::EGFP+single-cells by cell sorting (FIG. 11A). The expression levels of key pluripotency marker genes, such as OCT4, NANOG, SOX2, REX1, ESG1, and KLF4, in the optically maintained single-PSCs were similar to the levels in the single-PSCs maintained with FGF2 supplementation (FIG. 11B). These data demonstrate that our optical induction system for FGF signaling is a suitable long-term culture system for maintaining the pluripotency of hPSCs.

[0191] Conservation of pluripotency in optically maintained hPSC. To investigate the similarities at the molecular level between the light illumination and FGF2 treatment conditions that could conserve the pluripotency of hPSCs, we employed global transcriptome analyses (RNA-seq) (FIG. 12). The hierarchical clustering analysis demonstrated that the optically maintained hESCs (1, 5, or 10 min. of blue light illumination every 2 h for 3 weeks) had comparable transcriptional characteristics to that of the cells maintained with FGF2 supplementation, but not to that of the differentiated cells (FIGS. 2A and 2B). Moreover, the expression levels of pluripotency marker genes in the optically maintained hPSCs were similar to the levels in the hPSCs maintained with FGF2 supplementation (FIG. 2C). The majority of significantly differentially expressed genes either in the optically maintained hPSCs or in the hPSCs maintained with FGF2 protein overlapped with those of the differentiated cells (FIG. 2D). In a gene ontology (GO) analysis, the number of stem cell maintenance-related GO terms was significantly over-represented in the upregulated

genes of the optically maintained hESCs over the differentiated cells (FIG. 2E), which emphasizes that illumination is sufficient to maintain stem cell pluripotency. In contrast, various cell differentiation-related GO terms were significantly enriched in the downregulated genes of the optically maintained hESCs over the differentiated cells (FIG. 2F), which confirms that the light illumination condition can maintain the transcriptional profiles of human pluripotency similar to that of the FGF2 treatment group and also be sufficient to prevent any spontaneous differentiation of PSCs.

[0192] Preserved pluripotency in optically maintained hPSC. To find the direct differentiation potentials of hPSCs optically maintained, we differentiated these cells into three germ layers (FIG. 3A). Quantitative real-time PCR (qRT-PCR) results showed that the expression levels of PAX6 and SOX/ectoderm), TBX6 and MSGN1 (mesoderm), and FOXA2 and SOX/7 (endoderm) were significantly increased in the differentiated Opto-FGFR hPSCs compared to the undifferentiated cells (FIG. 3B, 3F). The differentiation abilities were also confirmed by immunostaining with NES-TIN (ectoderm), TBX6 (mesoderm), and FOXA2 (endoderm) antibodies (FIG. 3C). We further verified the directed differentiation abilities of optically maintained hPSCs into TH+/TJU1+dopaminergic neuron (ectoderm) or MF20+ skeletal muscle (mesoderm) (FIG. 3D). Additionally, the in vivo differentiation capacity of Opto-FGFR hPSCs was confirmed by a teratoma formation assay. The three germ layers were observed in teratoma tissue via hematoxylin/eosin staining (FIG. 3E, 3G). These results showed that long-term culture conditions along with the optical induction of the FGF signaling pathway maintained the pluripotency of PSCs without losing the cell specification capability toward the three germ layers.

[0193] Generation of a novel FGF2-free porcine PSC culture system. Porcine PSCs also show an FGF2-dependency when maintaining their pluripotencies [17,18]. To expand our opto-FGFR system to piPSCs, the opto-FGFR cassette was inserted into the porcine Rosa26 (pRosa26) safe-harbor locus in piPSCs using the CRISPR/Cas9 system (FIG. 4A). For the opto-FGFR knock-in at pRosa26 locus, two gRNAs were designed and applied to target the first intron of pRosa26 [39], and the opto-FGFR knock-in efficiencies were comparable with each other (FIG. 13A). The selected Opto-FGFR piPSC line showed a typical colony morphology and enabled the optical activation of the FGF signaling pathway (FIGS. 13B and 13C). While the Opto-FGFR piPSCs cultured with pulsed blue light maintained the piPSC-colony morphologies for more than 20 weeks, the cells cultured in the dark were fully differentiated after 3 weeks (FIGS. 4B and 4C). The optically maintained alkaline phosphatase-positive piPSCs exhibited a normal karyotype and comparable expression levels of pluripotency markers, such as pOct4, pNanog, pSox2, pLin28a mRNAs and pOct4 protein, to that of the cells maintained with FGF2 supplementation (FIGS. 4D and 4E and FIG. 13D-13F). Furthermore, the optically maintained piPSCs possessed the differentiation potential into the three germ layers (FIGS. 4F and 4G), revealing that our optical induction system for FGF signaling is a suitable culture system for maintaining the pluripotency of piPSC. Overall, these results support that we successfully developed an efficient and economical FGF2-free PSC culture system.

[0194] Simple and cost-efficient LED illumination system. To lower the entry barriers and increase control of the optical culture system, we next developed a do it yourself (DIY) light illumination system. We engineered a multi-well LED illuminating device to optically stimulate standard 6-well or 12-well tissue culture plates in a conventional CO2 incubator (FIG. 5A). The LED illuminating device consists of 6 or 12 LEDs of any color, a circuit board, an open-source microcontroller, and an enclosing plastic casing. The LEDs were soldered onto the circuit board and are controlled by the microcontroller (FIGS. 5A and 5B). The casing was designed using computer-aided design software and printed using a 3D plastic printer (FIG. 5A, 5C). Through our developed software and mobile application, the LED illuminating devices are remotely controlled by any computer or smartphone via a wireless connection (FIG. 5D). Users can easily control the pulse time length, frequency, and intensity of light illumination, and check current status of the LED illumination system (FIGS. 5E and 5F), which is seamlessly adopted in our routine stem cell culture. This DIY system provides the opportunity to apply optical culture systems to cell cultures in any laboratory.

DISCUSSION

[0195] We describe a novel and efficient culture system for PSCs via the activation of the FGF pathway to maintain PSCs without the need for supplementation of FGF2 recombinant protein. Previous reports have demonstrated the optical activation of receptor tyrosine kinases in immortalized cell lines [31,40,41] and developing embryos [42,43] as an initial step in evolving such optical technologies. Although a study using the optical control of ERK phosphorylation reported a perturbation of patterning and morphogenesis in *Drosophila* embryos [42], other studies did not report any biologically relevant phenotypes; therefore, translating these approaches for use with PSCs has remained challenging. Here, we extend the approach to human and porcine PSCs and provide clear evidence that our Opto-FGFR PSCs are sufficiently maintained by blue light illumination and without the need for FGF2 protein supplementation. Although overall patterns of protein phosphorylation in the blue light illuminated group were similar to FGF2 protein treated group, MSK1/2, eNOS, and HSP27, which are known to regulate cell proliferation [44-46], were slightly more phosphorylated in the blue light exposed group than FGF2 treated group (Supplementary FIG. 3). Since there is no need to supplement FGF2 protein, the media does not need to be frequently changed in the optical PSC culture system. In feeder-free culture systems, very high concentrations of FGF2 (up to 100 ng/mL) are required for promoting self-renewal and inhibiting spontaneous differentiation of hPSCs [15]. Even in a feeder-free culture system, we succeeded in maintaining pluripotency of hESCs using our optical culture system (FIG. 14). Moreover, the optically maintained PSCs appeared to retain the normal differentiation potential of traditional PSCs.

[0196] There are various kinds of photoactivatable proteins available for optogenetic approaches [9,10]. Using the photolyase homology region of the Cry2 protein (Cry2 PHR [47]) and LOV domain [35], two pioneer researches have established the optical activation systems of the FGF signaling pathway [31,40]. In the present study, the photosensory LOV domain served as an initiator in the light-induced intermolecular signal transduction cascade of the FGF sig-

naling pathway (termed as opto-FGFR signaling) for maintaining the pluripotency of mammalian PSCs, without FGF2 protein supplementation. Although we could optically activate FGF signaling in human embryonic kidney cells (HEK293T) and hESCs transiently transfected with the Cry2 PHR-based opto-FGFR (FIG. 15), we selected the LOV domain as a photodimerizable domain because of two reasons: the LOV domain has a shorter length (LOV, 432 bp or 144 aa [31] vs Cry2 PHR, 1506 bp or 502 aa [40]), which is more favorable for knock-in, and LOV domain has a shorter dissociation time after light withdrawal (LOV $t_{1/2}$ $\frac{1}{4}$ 2.5 min. [31] vs Cry2 PHR $t_{1/2}$ $\frac{1}{4}$ 5-6 min. [9,10]), which is more advantageous for accurate adjustment of the opto-FGFR signaling in PSC, than others. It is important to select an appropriate photosensory domain as the light-sensing actuator module for a given optogenetic modulation system in PSCs. In future work, it might also be important to build a systematic approach to select an appropriate photosensory domain for a given signaling pathway in a given cell type.

[0197] The current stem cell culture system is heavily dependent on random distribution of expensive and thermo-unstable recombinant proteins in a dish, which could jeopardize future mass production of human stem cells for a population-wide cell therapy. Our Opto-FGFR PSCs enable optical activation without any exogenous FGF2 recombinant proteins and offer spatiotemporally precise optical control of stem cell behaviors. Beyond implementing a new technology in the stem cell field, our experimental data could help lead to the development of a potential therapeutic cure against many human diseases by modulating multiple signaling pathways other than FGF2 and maximizing the potential of human PSCs. While our optogenetic culture system to stimulate the FGF signaling pathway is suitable for maintaining the pluripotency of PSCs, further studies to activate the LIF and/or Activin signaling via optogenetic stimulation for generating PSCs in a naïve state of pluripotency [48] will be of great interest. Furthermore, it will be interesting to see whether other signaling molecules to maintain the pluripotency of PSCs, such as insulin and/or TGF β 1 (or Nodal) [49], can be replaceable by the photo-activatable system. Regarding hiPSC culture for potential therapeutic application, developing an optogenetic system to activate insulin and/or TGF β 1 (or Nodal) signaling would be one way to reduce its production costs. In addition, because the FGF signaling pathway is a key factor for several differentiation protocols of PSCs, including ectodermal and neuronal differentiation protocols, our optogenetic system could be applied to the differentiation researches.

[0198] As many researchers adopted optogenetic techniques particularly for neuroscience field, there is a growing concern of a possible phototoxicity. For example, other group recently reported a phototoxicity caused by blue light illumination in *in vitro* culture of neural cells [50]. However, the light intensity (1 mW/mm²) in their illumination condition is 1000-fold higher than that of our culture condition (1 μ W/mm²), and the frequency (1 Hz) in their condition is 7200-fold higher than that of our condition (0.14 mHz); which are why we did not detect any noticeable toxicity during over 1-year culture. It will be important to know what the susceptible ranges of the light illumination conditions in different cell types in future is. While it still needs to study more to find an ideal illumination conditions, '5 min. blue light illumination in every 2 h' was sufficient to maintain the pluripotency of PSCs; further studies to exquisitely optimize

the illuminating conditions, such as pulse width, frequency, and intensity of illumination, will be of great interest.

[0199] The current animal agriculture and husbandry methods cause significant environmental issues such as greenhouse gas emissions, fresh water consumption, and arable land usage. As an alternative, cellular agriculture [51] or lab-grown meats [52] have been introduced and social consent is growing due to ethical and environmentally friendly concepts. However, the production costs and use of animal-derived thermo-unstable recombinant proteins during the cellular agriculture process are barriers to entering the market for mass consumption. Based on our calculation, recombinant FGF2 protein for 1 billion of PSC culture costs \$14,000 weekly. In addition, due to thermal and chemical instabilities, FGF2 is recommended to be freshly added into the media and it requires additional labor fees. In conclusion, our opto-FGFR system in livestock PSCs can be a realistic solution to decrease these production costs.

[0200] Although researchers may consider that it is too early to know whether genetic engineering techniques will be safe in the clinic, the results of clinical trials of CRISPR/Cas9-mediated genome editing have been promising up to now [53-55]. We carefully speculate that the CRISPR-based genome editing of PSCs would be clinically possible with additional development and troubleshooting. Based on our karyotypic analysis results, our CRISPR-based genome editing of PSCs did not cause detectable karyotypic abnormalities (FIG. 11F FIG. 13F, and FIG. 14E); however, we will study much details of any possible genetic aberrations in future. If we employ newly developed genetic engineering techniques, such as prime editors [56], high-fidelity CRISPR/Cas9 nucleases [57], and CRISPR/Cas9 nickase [58], to increase accuracy and to reduce off-target effects, the safety issues can be mitigated. While it is hard to precisely estimate the cost to produce genetically engineered cell lines through the CRISPR system, there would not be an additional expense after upfront cost after production.

[0201] Currently, there are a few culture optogenetics systems on the market, but these systems are all very expensive. High cost commercial products prevent large volume optogenetics studies and thus, limit the rate of research in this field. There is also a lack of instrumentation that allows for precise control of illumination and remote monitoring of multiple culture plates. Our DIY illumination system is a hardware, software, and mobile application system capable of precise control and real-time monitoring of cell culture illumination that can serve as an inexpensive alternative to high cost commercial products. This illumination system was designed to be easy-to-assemble, easy-to-use, and low-cost, utilizing 3D printing of thermoplastic materials, programmable open-source electronics, and other readily-available electronic parts. Additionally, this system was designed to be adjustable and be customized to the individual researcher. The 3D models and source code are publicly available to be edited for any specific needs. Our cost saving PSC culture system with illumination system could help understanding disease mechanism. Furthermore, we did not find any notable safety issue, even in the feeder-free condition during the culture period with blue light illumination. Additionally, the LED illuminating device can be customized with any color LED to be used with any photosynthetic protein of interest. This DIY system provides the opportunity to apply a flexible optical culture system to cell cultures in any laboratory.

CONCLUSION

[0202] Our approach for the optical maintenance of human and porcine PSCs provides new insights for an alternative synthetic approach for the next generation in stem cell research. Furthermore, considering the number of signaling pathways that are crucial for controlling cell specification processes, our Opto-FGFR PSCs are a proof-of-concept for the optical control over a plethora of mammalian stem cell fates.

REFERENCES

- [0203] 1. G. Nagel, D. Ollig, M. Fuhrmann, S. Kateriya, A. M. Musti, E. Bamberg, P. Hegemann, Channelrhodopsin-1: a light-gated proton channel in green algae, *Science* 296 (5577) (2002) 2395-2398.
- [0204] 2. K. Deisseroth, G. Feng, A. K. Majewska, G. Miesenböck, A. Ting, M. J. Schnitzer, Next-generation optical technologies for illuminating genetically targeted brain circuits, *J. Neurosci.* 26 (41) (2006) 10380-10386.
- [0205] 3. L. Fenno, O. Yizhar, K. Deisseroth, The development and application of optogenetics, *Annu. Rev. Neurosci.* 34 (2011) 389-412.
- [0206] 4. M. Yazawa, A. M. Sadaghiani, B. Hsueh, R. E. Dolmetsch, Induction of protein-protein interactions in live cells using light, *Nat. Biotechnol.* 27 (10) (2009) 941-945.
- [0207] 5. Y. I. Wu, D. Frey, O. I. Lungu, A. Jaehrig, I. Schlichting, B. Kuhlman, K. M. Hahn, A genetically encoded photoactivatable Rac controls the motility of living cells, *Nature* 461 (7260) (2009) 104-108.
- [0208] 6. A. Levskaya, O. D. Weiner, W. A. Lim, C. A. Voigt, Spatiotemporal control of cell signalling using a light-switchable protein interaction, *Nature* 461 (7266) (2009) 997-1001.
- [0209] 7. S. Shimizu-Sato, E. Huq, J. M. Tepperman, P. H. Quail, A light-switchable gene promoter system, *Nat. Biotechnol.* 20 (10) (2002) 1041-1044.
- [0210] 8. M. J. Kennedy, R. M. Hughes, L. A. Peteya, J. W. Schwartz, M. D. Ehlers, C. L. Tucker, Rapid blue-light-mediated induction of protein interactions in living cells, *Nat. Methods* 7 (12) (2010) 973-975.
- [0211] 9. K. Zhang, B. Cui, Optogenetic control of intracellular signaling pathways, *Trends Biotechnol.* 33 (2) (2015) 92-100.
- [0212] 10. D. Tischer, O. D. Weiner, Illuminating cell signalling with optogenetic tools, *Nat. Rev. Mol. Cell Biol.* 15 (8) (2014) 551-558.
- [0213] 11. L. J. Bugaj, A. T. Choksi, C. K. Mesuda, R. S. Kane, D. V. Schaffer, Optogenetic protein clustering and signaling activation in mammalian cells, *Nat. Methods* 10 (3) (2013) 249-252.
- [0214] 12. M. K. Furue, J. Na, J. P. Jackson, T. Okamoto, M. Jones, D. Baker, R. Hata, H. D. Moore, J. D. Sato, P. W. Andrews, Heparin promotes the growth of human embryonic stem cells in a defined serum-free medium, *Proc. Natl. Acad. Sci. U.S.A.* 105 (36) (2008) 13409-13414.
- [0215] 13. P. J. Tesar, J. G. Chenoweth, F. A. Brook, T. J. Davies, E. P. Evans, D. L. Mack, R. L. Gardner, R. D. McKay, New cell lines from mouse epiblast share defining features with human embryonic stem cells, *Nature* 448 (7150) (2007) 196-199.
- [0216] 14. K. Takahashi, K. Tanabe, M. Ohnuki, M. Narita, T. Ichisaka, K. Tomoda, S. Yamanaka, Induction of pluripotent stem cells from adult human fibroblasts by defined factors, *Cell* 131 (5) (2007) 861-872.
- [0217] 15. M. E. Levenstein, T. E. Ludwig, R. H. Xu, R. A. Llanas, K. VanDenHeuvel-Kramer, D. Manning, J. A. Thomson, Basic fibroblast growth factor support of human embryonic stem cell self-renewal, *Stem Cell.* 24 (3) (2006) 568-574.
- [0218] 16. J. A. Thomson, J. Itskovitz-Eldor, S. S. Shapiro, M. A. Waknitz, J. J. Swiergiel, V. S. Marshall, J. M. Jones, Embryonic stem cell lines derived from human blastocysts, *Science* 282 (5391) (1998) 1145-1147.
- [0219] 17. F. D. West, S. L. Terlouw, D. J. Kwon, J. L. Mumaw, S. K. Dhara, K. Hasneen, J. R. Dobrinsky, S. L. Stice, Porcine induced pluripotent stem cells produce chimeric offspring, *Stem Cells Dev.* 19 (8) (2010) 1211-1220.
- [0220] 18. T. Ezashi, B. P. Telugu, A. P. Alexenko, S. Sachdev, S. Sinha, R. M. Roberts, Derivation of induced pluripotent stem cells from pig somatic cells, *Proc. Natl. Acad. Sci. U.S.A.* 106 (27) (2009) 10993-10998.
- [0221] 19. M. Amit, M. K. Carpenter, M. S. Inokuma, C. P. Chiu, C. P. Harris, M. A. Waknitz, J. Itskovitz-Eldor, J. A. Thomson, Clonally derived human embryonic stem cell lines maintain pluripotency and proliferative potential for prolonged periods of culture, *Dev. Biol.* 227 (2) (2000) 271-278.
- [0222] 20. N. Itoh, D. M. Ornitz, Evolution of the Fgf and Fgfr gene families, *Trends Genet.* 20 (11) (2004) 563-569.
- [0223] 21. H. D. Beer, L. Vindevoghel, M. J. Gait, J. M. Revest, D. R. Duan, I. Mason, C. Dickson, S. Werner, Fibroblast growth factor (FGF) receptor 1-IIIb is a naturally occurring functional receptor for FGFs that is preferentially expressed in the skin and the brain, *J. Biol. Chem.* 275 (21) (2000) 16091-16097.
- [0224] 22. I. Urakawa, Y. Yamazaki, T. Shimada, K. Iijima, H. Hasegawa, K. Okawa, T. Fujita, S. Fukumoto, T. Yamashita, Klotho converts canonical FGF receptor into a specific receptor for FGF23, *Nature* 444 (7120) (2006) 770-774.
- [0225] 23. D. M. Ornitz, A. Yayon, J. G. Flanagan, C. M. Svahn, E. Levi, P. Leder, Heparin is required for cell-free binding of basic fibroblast growth factor to a soluble receptor and for mitogenesis in whole cells, *Mol. Cell Biol.* 12 (1) (1992) 240-247.
- [0226] 24. M. Mohammadi, I. Dikic, A. Sorokin, W. H. Burgess, M. Jaye, J. Schlessinger, Identification of six novel autophosphorylation sites on fibroblast growth factor receptor 1 and elucidation of their importance in receptor activation and signal transduction, *Mol. Cell Biol.* 16 (3) (1996) 977-989.
- [0227] 25. G. Chen, D. R. Gulbranson, P. Yu, Z. Hou, J. A. Thomson, Thermal stability of fibroblast growth factor protein is a determinant factor in regulating self-renewal, differentiation, and reprogramming in human pluripotent stem cells, *Stem Cell.* 30 (4) (2012)
- [0228] 26. I. Y. Choi, H. Lim, K. Estrellas, J. Mula, T. V. Cohen, Y. Zhang, C. J. Donnelly, J. P. Richard, Y. J. Kim, H. Kim, Y. Kazuki, M. Oshimura, H. L. Li, A. Hotta, J. Rothstein, N. Maragakis, K. R. Wagner, G. Lee, Concordant but varied phenotypes among duchenne muscular

- dystrophy patient-specific myoblasts derived using a human iPSC-based model, *Cell Rep.* 15 (10) (2016) 2301-2312.
- [0229] 27. P. Mali, L. Yang, K. M. Esvelt, J. Aach, M. Guell, J. E. DiCarlo, J. E. Norville, G. M. Church, RNA-guided human genome engineering via Cas9, *Science* 339 (6121) (2013) 823-826.
- [0230] 28. I. Y. Choi, H. Lim, H. J. Cho, Y. Oh, B. K. Chou, H. Bai, L. Cheng, Y. J. Kim, S. Hyun, H. Kim, J. H. Shin, G. Lee, Transcriptional landscape of myogenesis from human pluripotent stem cells reveals a key role of TWIST1 in maintenance of skeletal muscle progenitors, *Elife* 9 (2020).
- [0231] 29. H. Lim, I. Y. Choi, G. Lee, Profiling individual human embryonic stem cells by quantitative RT-PCR, *J. Vis. Exp.* 87 (2014) e51408.
- [0232] 30. S. Kriks, J. W. Shim, J. Piao, Y. M. Ganat, D. R. Wakeman, Z. Xie, L. Carrillo-Reid, G. Auyeung, C. Antonacci, A. Buch, L. Yang, M. F. Beal, D. J. Surmeier, J. H. Kordower, V. Tabar, L. Studer, Dopamine neurons derived from human ES cells efficiently engraft in animal models of Parkinson's disease, *Nature* 480 (7378) (2011) 547-551.
- [0233] 31. M. Grusch, K. Schelch, R. Riedler, E. Reichhart, C. Differ, W. Berger, A. Ingles-Prieto, H. Janovjak, Spatio-temporally precise activation of engineered receptor tyrosine kinases by light, *EMBO J.* 33 (15) (2014) 1713-1726.
- [0234] 32. K. Huang, T. Merkle, C. F. Beck, Isolation and characterization of a *Chlamydomonas* gene that encodes a putative blue-light photoreceptor of the phototropin family, *Physiol. Plant* 115 (4) (2002) 613-622.
- [0235] 33. T. Kinoshita, M. Doi, N. Suetsugu, T. Kagawa, M. Wada, K. Shimazaki, Phot1 and phot2 mediate blue light regulation of stomatal opening, *Nature* 414 (6864) (2001) 656-660.
- [0236] 34. F. Takahashi, D. Yamagata, M. Ishikawa, Y. Fukamatsu, Y. Ogura, M. Kasahara, T. Kiyosue, M. Kikuyama, M. Wada, H. Kataoka, Aureochrome, a photoreceptor required for photomorphogenesis in stramenopiles, *Proc. Natl. Acad. Sci. U.S.A.* 104 (49) (2007) 19625-19630.
- [0237] 35. T. Toyooka, O. Hisatomi, F. Takahashi, H. Kataoka, M. Terazima, Photoreactions of aureochrome-1, *Biophys. J.* 100 (11) (2011) 2801-2809.
- [0238] 36. Y. Oh, G. S. Cho, Z. Li, I. Hong, R. Zhu, M. J. Kim, Y. J. Kim, E. Tampakakis, L. Tung, R. Haganir, X. Dong, C. Kwon, G. Lee, Functional coupling with cardiac muscle promotes maturation of hpSC-derived sympathetic neurons, *Cell Stem Cell* 19 (1) (2016) 95-106.
- [0239] 37. L. Cong, F. A. Ran, D. Cox, S. Lin, R. Barretto, N. Habib, P. D. Hsu, X. Wu, W. Jiang, L. A. Marraffini, F. Zhang, Multiplex genome engineering using CRISPR/Cas systems, *Science* 339 (6121) (2013) 819-823.
- [0240] 38. S. Lotz, S. Goderie, N. Tokas, S. E. Hirsch, F. Ahmad, B. Corneo, S. Le, A. Banerjee, R. S. Kane, J. H. Stern, S. Temple, C. A. Fasano, Sustained levels of FGF2 maintain undifferentiated stem cell cultures with biweekly feeding, *PLoS One* 8 (2) (2013) e56289.
- [0241] 39. Z. Xie, D. Pang, K. Wang, M. Li, N. Guo, H. Yuan, J. Li, X. Zou, H. Jiao, H. Ouyang, Z. Li, X. Tang, Optimization of a CRISPR/Cas9-mediated knock-in strategy at the porcine Rosa26 locus in porcine foetal fibroblasts, *Sci. Rep.* 7 (1) (2017) 3036.
- [0242] 40. N. Kim, J. M. Kim, M. Lee, C. Y. Kim, K. Y. Chang, W. D. Heo, Spatiotemporal control of fibroblast growth factor receptor signals by blue light, *Chem Biol.* 21 (7) (2014) 903-912.
- [0243] 41. Y. Li, M. Lee, N. Kim, G. Wu, D. Deng, J. M. Kim, X. Liu, W. D. Heo, Z. Zi, Spatiotemporal control of TGF-beta signaling with light, *ACS Synth. Biol.* 7 (2) (2018) 443-451.
- [0244] 42. H. E. Johnson, Y. Goyal, N. L. Pannucci, T. Schupbach, S. Y. Shvartsman, J. E. Toettcher, The Spatiotemporal Limits of Developmental Erk Signaling, *Dev. Cell* 40 (2) (2017) 185-192.
- [0245] 43. K. Sako, S. J. Pradhan, V. Barone, A. Ingles-Prieto, P. Muller, V. Ruprecht, D. Capek, S. Galande, H. Janovjak, C. P. Heisenberg, Optogenetic control of nodal signaling reveals a temporal pattern of nodal signaling regulating cell fate specification during gastrulation, *Cell Rep.* 16 (3) (2016) 866-877.
- [0246] 44. S. Zhang, Y. Hu, Y. Huang, H. Xu, G. Wu, H. Dai, Heat shock protein 27 promotes cell proliferation through activator protein-1 in lung cancer, *Oncol. Lett.* 9 (6) (2015) 2572-2576.
- [0247] 45. D. Reyes, C. Ballare, G. Castellano, D. Soronellas, J. R. Bago, J. Blanco, M. Beato, Activation of mitogen- and stress-activated kinase 1 is required for proliferation of breast cancer cells in response to estrogens or progestins, *Oncogene* 33 (12) (2014) 1555-1565.
- [0248] 46. C. Gentile, R. C. Muise-Helmericks, C. J. Drake, VEGF-mediated phosphorylation of eNOS regulates angioblast and embryonic endothelial cell proliferation, *Dev. Biol.* 373 (1) (2013) 163-175.
- [0249] 47. H. Liu, X. Yu, K. Li, J. Klejnot, H. Yang, D. Lisiero, C. Lin, Photoexcited CRY2 interacts with CIB1 to regulate transcription and floral initiation in *Arabidopsis*, *Science* 322 (5907) (2008) 1535-1539.
- [0250] 48. X. Gao, M. Nowak-Imialek, X. Chen, D. Chen, D. Herrmann, D. Ruan, A. C. H. Chen, M. A. Eckersley-Maslin, S. Ahmad, Y. L. Lee, T. Kobayashi, D. Ryan, J. Zhong, J. Zhu, J. Wu, G. Lan, S. Petkov, J. Yang, L. Antunes, L. S. Campos, B. Fu, S. Wang, Y. Yong, X. Wang, S. G. Xue, L. Ge, Z. Liu, Y. Huang, T. Nie, P. Li, D. Wu, D. Pei, Y. Zhang, L. Lu, F. Yang, S. J. Kimber, W. Reik, X. Zou, Z. Shang, L. Lai, A. Surani, P. P. L. Tam, A. Ahmed, W. S. B. Yeung, S. A. Teichmann, H. Niemann, P. Liu, Establishment of porcine and human expanded potential stem cells, *Nat. Cell Biol.* 21 (6) (2019) 687-699.
- [0251] 49. G. Chen, D. R. Gulbranson, Z. Hou, J. M. Bolin, V. Ruotti, M. D. Probasco, K. Smuga-Otto, S. E. Howden, N. R. Diol, N. E. Propson, R. Wagner, G. O. Lee, J. Antosiewicz-Bourget, J. M. Teng, J. A. Thomson, Chemically defined conditions for human iPSC derivation and culture, *Nat. Methods* 8 (5) (2011) 424-429.
- [0252] 50. J. H. Stockley, K. Evans, M. Matthey, K. Volbracht, S. Agathou, J. Mukanowa, J. Burrone, R. T. Karadottir, Surpassing light-induced cell damage in vitro with novel cell culture media, *Sci. Rep.* 7 (1) (2017) 849.
- [0253] 51. S. L. Liu, K. Gasteratos, Assessing cell-based animal proteins, *Science* 363 (6429) (2019) 826.
- [0254] 52. E. Dolgin, Sizzling interest in lab-grown meat belies lack of basic research, *Nature* 566 (7743) (2019) 161-162.
- [0255] 53. H. Ledford, Quest to use CRISPR against disease gains ground, *Nature* 577 (7789) (2020) 156.

[0256] 54. E. A. Stadtmauer, J. A. Fraietta, M. M. Davis, A. D. Cohen, K. L. Weber, E. Lancaster, P. A. Mangan, I. Kulikovskaya, M. Gupta, F. Chen, L. Tian, V. E. Gonzalez, J. Xu, I. Y. Jung, J. J. Melenhorst, G. Plesa, J. Shea, T. Matlawski, A. Cervini, A. L. Gaymon, S. Desjardins, A. Lamontagne, J. Salas-Mckee, A. Fesnak, D. L. Siegel, B. L. Levine, J. K. Jadowsky, R. M. Young, A. Chew, W. T. Hwang, E. O. Hexner, B. M. Carreno, C. L. Nobles, F. D. Bushman, K. R. Parker, Y. Qi, A. T. Satpathy, H. Y. Chang, Y. Zhao, S. F. Lacey, C. H. June, CRISPR-engineered T cells in patients with refractory cancer, *Science* 367 (6481)

[0257] 55. L. Xu, J. Wang, Y. Liu, L. Xie, B. Su, D. Mou, L. Wang, T. Liu, X. Wang, B. Zhang, L. Zhao, L. Hu, H. Ning, Y. Zhang, K. Deng, L. Liu, X. Lu, T. Zhang, J. Xu, C. Li, H. Wu, H. Deng, H. Chen, CRISPR-edited stem cells in a patient with hiv and acute lymphocytic leukemia, *N. Engl. J. Med.* 381 (13) (2019) 1240-1247.

[0258] 56. A. V. Anzalone, P. B. Randolph, J. R. Davis, A. A. Sousa, L. W. Koblan, J. M. Levy, P. J. Chen, C. Wilson, G. A. Newby, A. Raguram, D. R. Liu, Search-and-replace genome editing without double-strand breaks or donor DNA, *Nature* 576 (7785) (2019) 149-157.

[0259] 57. B. P. Kleinstiver, V. Pattanayak, M. S. Prew, S. Q. Tsai, N. T. Nguyen, Z. Zheng, J. K. Joung, High-fidelity CRISPR-Cas9 nucleases with no detectable genome-wide off-target effects, *Nature* 529 (7587) (2016) 490-495.

[0260] 58. F. A. Ran, P. D. Hsu, C. Y. Lin, J. S. Gootenberg, S. Konermann, A. E. Trevino, D. A. Scott, A. Inoue, S. Matoba, Y. Zhang, F. Zhang, Double nicking by RNA-guided CRISPR Cas9 for enhanced genome editing specificity, *Cell* 154 (6) (2013) 1380-1389.

SEQUENCE LISTING

Sequence total quantity: 61
SEQ ID NO: 1 moltype = DNA length = 5206
FEATURE Location/Qualifiers
misc_feature 1..5206
 note = AAV-Myr-opto-FGFR-LOV
misc_feature 1790..2373
 note = promoter
misc_feature 2496..2537
 note = Myr signal peptide
misc_feature 2544..3815
 note = FGFR1
misc_feature 3828..4262
 note = LOV
source 1..5206
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 1

tgctttctct	gaccagcatt	ctctcccctg	ggcctgtgcc	gctttctgtc	tgcagcttgt	60
ggcctggggt	acctctacgg	ctggcccaga	tccttcccctg	ccgcctcctt	caggttccgt	120
cttcctccac	tccctcttcc	ccttgctctc	tgctgtgttg	ctgcccagg	atgctctttc	180
cggagcactt	ccttctcggc	gctgcaccac	gtgatgtcct	ctgagcggat	cctccccgtg	240
tctgggtcct	ctccgggcat	ctctcctccc	tcacccaacc	ccatgccgtc	ttcactcgct	300
gggttccctt	ttccttctcc	ttctggggcc	tggtccatct	ctcgtttctt	aggatggcct	360
tctccgacgg	atgtctccct	tgcgtcccgc	ctccccttct	tgtaggcctg	catcatcacc	420
gtttttctgg	acaaccccaa	agtaccccg	ctccctggct	ttagccacct	ctccatcctc	480
ttgctttctt	tgcctggaca	ccccgttctc	ctgtggattc	gggtcacctc	tcactccttt	540
catttgggca	gctcccctac	cccccttacc	tctctagtct	gtgctagctc	ttccagcccc	600
ctgtcatggc	atcttccagg	ggtcgagag	ctcagctagt	cttcttccct	caacccgggc	660
ccctatgtcc	acttcaggac	agcatgtttg	ctgcctccag	ggatcctgtg	tccccgagct	720
gggaccacct	tatatccca	gggcgggtta	atgtggctct	ggttctgggt	acttttatct	780
gtcccctcca	ccccacagt	gggcaagctt	ctgacctctt	ctcttccctc	cacagggcct	840
cgagagatct	ggcagcggag	agggcagagg	aagtcttcta	acatgcggtg	acgtggagga	900
gaatcccggc	cctaggctcg	agatgaccga	gtacaagccc	acggtgcgcc	tcgccaccgc	960
cgacgacgtc	cccaggggcc	tacgcaccct	cgccgcgcgc	ttcgccgact	accccgccac	1020
gcgccacacc	gtcgatccgg	accgccacat	cgagcgggtc	accgagctgc	aagaactctt	1080
cctcacgcgc	gtcgggctcg	acatcgga	ggtgtgggtc	gcggacgacg	gcgccgcggt	1140
ggcgggtctg	accacgcccg	agagcgtcga	agcggggggc	gtgttcgccc	agatcgggcc	1200
gcgcatggcc	gagttgagcg	gttcccggct	ggccgcgcag	caacagatgg	aaggcctcct	1260
ggcgccgcac	cggcccgaag	agcccgcgtg	gttcttgccc	accgtcggcg	tctcgcccga	1320
ccaccagggc	aagggtctgg	gcagcgccgt	cgtgctcccc	ggagtggagg	cggccgagcg	1380
cgcgggggtg	ccgccttccc	tggagacctc	cgcgccccgc	aacctccctt	tctacgagcg	1440
gctcggtctc	accgtcaccg	ccgacgtcga	ggtgcccga	ggaccgcgca	cctgggtgcat	1500
gaccgcgaag	cccgggtgcct	gatctagagg	gcccgttta	accgctgat	cagcctcgac	1560
tgtgccttct	agttgccagc	catctgttgt	ttgcccctcc	cccgtgcctt	ccttgaccct	1620
ggaaggtgcc	actcccactg	tcctttccta	ataaaatgag	gaaattgcat	cgcattgtct	1680
gagtaggtgt	cattctattc	tgggggggtg	ggtggggcag	gacagcaagg	gggaggattg	1740
ggaagacaat	agcaggcatg	ctgggggatg	ggtgggctct	atgggtctcg	acattgatta	1800
ttgactagtt	attaatagta	atcaattacg	gggtcattag	ttcatagccc	atatatggag	1860
ttccgcgtta	cataacttac	ggtaaattgg	ccgcctggct	gaccgcccga	cgacccccgc	1920
ccattgacgt	caataatgac	gtatgttccc	atagtaacgc	caatagggac	tttccattga	1980
cgtcaatggg	tggagtattt	acggtaaa	gcccacttgg	cagtacatca	agtgatatcat	2040
atgccaaagta	cgccccctat	tgacgtcaat	gacggtaaat	ggcccgccctg	gcattatgcc	2100
cagtacatga	ccttatggga	ctttcctact	tggcagtaca	tctacgtatt	agtcactcgct	2160

-continued

attaccatgg	tgatgcggtt	ttggcagtac	atcaatgggc	gtggatagcg	gtttgactca	2220
cggggatttc	caagtctcca	ccccattgac	gtcaatggga	gtttgttttg	gcacccaaat	2280
caacgggact	ttccaaaatg	tcgtaacaac	tccgccccat	tgacgcaa	gggcggtagg	2340
cgtgtacggg	gggaggtcta	tataagcaga	gctctctggc	taactagaga	accactgct	2400
tactggctta	tcgaaattaa	tacgactcac	tatagggaga	cacaagctgg	ctagcgttta	2460
aacgggccc	ctagactcga	gcgccgcgg	ccaccatggg	gagtagcaag	agcaagccta	2520
aggaccccag	ccagcgctc	gacatgaaga	gcggcaccaa	gaagagcgac	ttccatagcc	2580
agatggctgt	gcacaagctg	gccaaagagc	tccctctgcg	cagacaggtg	acagtgtcag	2640
ctgactccag	tgcattccatg	aactctgggg	ttctctgggt	tcggccctca	cggtctcct	2700
ccagcgggac	ccccatgctg	gctggagctc	ccgaatatga	gctccctgag	gatccccgct	2760
gggagctgcc	acgagacaga	ctggctcttag	gcaaaccact	tggcgagggc	tgttcggggc	2820
aggtgggtgt	ggctgaggcc	atcgggctgg	ataaggacaa	acccaaccgt	gtgaccaaag	2880
tggccgtgaa	gatgttgaa	tcgacgcaa	cggagaagga	cctgtcggat	ctgatctcgg	2940
agatggagat	gatgaaaatg	attgggaagc	acaagaatat	catcaacctt	ctgggagcgt	3000
gcacacagga	tggtcctctt	tatgtcattg	tggagtacgc	ctccaaaggc	aatctccggg	3060
agtatctaca	ggcccggagg	cctcctgggg	tggagtactg	ctataacccc	agccacaacc	3120
ccgaggaaca	gctgtcttcc	aaagatctgg	tatcctgtgc	ctatcaggtg	gctcggggca	3180
tggagtatct	tgctcttaag	aagtgtatac	accgagacct	ggctgctagg	aacgtcctgg	3240
tgaccgagga	taacgtaatg	aagatcgtag	actttggctt	agctcgagac	attcatcata	3300
tcgactacta	caagaaaacc	accaacggcc	ggctgcctgt	gaagtggatg	gcccctgagg	3360
cgttggttga	ccgatctac	acacaccaga	gcgatgtgtg	gtcttttggg	gtgctcttgt	3420
gggagatctt	cactctgggt	ggctcccat	acccgggtgt	gcctgtggag	gaacttttca	3480
agctgctgaa	ggagggtcat	cgaatggaca	agcccagtaa	ctgtaccaat	gagctgtaca	3540
tgatgatgcg	ggactgctgg	catgcagtgc	cctctcagag	acctacgttc	aagcagttgg	3600
tggaagacct	ggacgcatt	gtggccttga	cctccaacca	ggagtatctg	gacctgtcca	3660
taccgctgga	ccagtactca	cccagcttcc	ccgacacacg	gagctccacc	tgctcctcag	3720
gggaggactc	tgtcttctct	catgagcctg	tacctgagga	gcctgtctg	cctcgacacc	3780
ccaccagct	tgccaacagt	ggactcaaac	ggcgctcga	gaccggtcct	gactacagtc	3840
tcgtgaaggc	tctgcaaatg	gcacaacaga	attttgtcat	tacagacgcc	tccctcccag	3900
acaaccctat	cgtctacgcc	agtagagggt	ttctgacact	gacaggctat	tctctcgacc	3960
agatcctggg	caggaactgc	aggtttctgc	aagggccaga	aacagaccca	agagctgtgg	4020
ataagatcag	gaatgccatc	accaaaggcg	ttgataccag	tgtctgtctg	ctgaattata	4080
gacaggatgg	cacaaccttc	tggaaatctc	tcttcgtggc	tggactcaga	gattctaagg	4140
gcaatattgt	caactacgtc	ggagtgcagt	caaagggtgag	cgaagattat	gccaagctgc	4200
tggccaacga	gcagaacatt	gagtacaaag	gtgtgcgcac	cagtaacatg	ctgcgcagaa	4260
agcccggtgg	atccggagtc	gactatccgt	acgacgtacc	agactacgca	ctcgactaag	4320
aattccacca	cactggacta	gtggatccga	gctcggtagc	aagcttaaga	ctagggacag	4380
gattgggtgac	agaaaagccc	catccttagg	cctcctcctt	cctagtctcc	tgatattggg	4440
tctaaccctc	acctcctgtt	aggcagattc	cttatctggg	gacacacccc	catttcctgg	4500
agccatctct	ctccttgcca	gaacctctaa	ggtttgctta	cgatggagcc	agagaggatc	4560
ctgggagggg	gagcttgcca	gggggtggga	gggaaggggg	ggatgcgtga	cctgcccggg	4620
tctcagtggc	cacctgcgc	tacctctctc	cagaacctga	gctgctctga	cgcggctgtc	4680
tggcgctgtt	cactgatcct	ggtgctgcag	cttccttaca	cttcccaaga	ggagaagcag	4740
tttggaaaaa	caaaatcaga	ataagttggg	cctgagttct	aactttgggt	cttcaccttt	4800
ctagtcccca	atttatattg	ttcctccgtg	cgtcagtttt	acctgtgaga	taaggccagt	4860
agccagcccc	gtcctggcag	ggctgtgggtg	aggagggggg	tgtccgtgtg	gaaaactccc	4920
tttgtgagaa	tgggtgcgtc	taggtgttca	ccaggctcgtg	gccgcctcta	ctccctttct	4980
ctttctccat	ccttctttcc	ttaaagagtc	ccaagtgcga	tctgggacat	attcctccgc	5040
ccagagcagg	gtcccgtctc	cctaaggccc	tgctctgggc	ttctgggttt	gagtccttgg	5100
caagcccagg	agaggcgctc	aggtctccct	gtcccccttc	ctcgtccacc	atctcatgcc	5160
cctggctctc	ctgccccttc	cctacagggg	ttcctggctc	tgctct		5206

SEQ ID NO: 2
 FEATURE
 misc_feature
 source

moltype = DNA length = 584
 Location/Qualifiers
 1..584
 note = Promoter of AAV-Myr-opto-FGFR-LOV
 1..584
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 2

gacattgatt	attgactagt	tattaatagt	aatcaattac	gggggtcatta	gttcatagcc	60
catatatgga	gttccgcgtt	acataactta	cggtaaatgg	cccgcctggc	tgaccgcccc	120
acgacccccg	cccatagacg	tcaataatga	cgtatgttcc	catagtaacg	ccaataggga	180
ctttccattg	acgtcaatgg	gtggagtatt	tacggtaaac	tgcccacttg	gcagtacatc	240
aagtgtatca	tatgccaa	acgcccccta	ttgacgtcaa	tgacggtaaa	tgccccgcct	300
ggcattatgc	ccagtacatg	accttatggg	actttcctac	ttggcagtag	atctacgtat	360
tagtcacgc	tattaccatg	gtgatgcggg	tttggcagta	catcaatggg	cgtggatagc	420
ggtttgactc	acggggattt	ccaagtctcc	accccattga	cgtcaatggg	agtttgtttt	480
ggcaccaaaa	tcaacgggac	tttccaaaat	gtcgtaacaa	ctccgcccc	ttgacgcaaa	540
tgggcggtag	gcgtgtacgg	tgggaggtct	atataagcag	agct		584

SEQ ID NO: 3
 FEATURE
 misc_feature
 source

moltype = DNA length = 42
 Location/Qualifiers
 1..42
 note = Nucleotide sequence of signal peptide
 1..42

-continued

	mol_type = other DNA organism = synthetic construct	
SEQUENCE: 3		
atgggggagta gcaagagcaa gcctaaggac cccagccagc gc		42
SEQ ID NO: 4	moltype = AA length = 14	
FEATURE	Location/Qualifiers	
REGION	1..14	
	note = Amino acid sequence of signal petpide	
source	1..14	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 4		
MGSSKSKPKD PSQR		14
SEQ ID NO: 5	moltype = DNA length = 1272	
FEATURE	Location/Qualifiers	
misc_feature	1..1272	
	note = Nucleotide sequence encoding FGFR1	
source	1..1272	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 5		
atgaagagcg gcaccaagaa gagcgacttc catagccaga tggctgtgca caagctggcc	60	
aagagcatcc ctctgcgag acaggtaca gtgtcagctg actccagtgc atccatgaac	120	
tctgggggttc tcctgggtcg gccctcacgg ctctcctcca gcgggacccc catgctggct	180	
ggagtctccg aatatgagct ccctgaggat ccccgctggg agctgccacg agacagactg	240	
gtcttaggca aaccacttgg cgagggctgc ttcgggcagg tgggtgttggc tgaggccatc	300	
gggctggata aggacaaacc caaccgtgtg accaaagtgg ccgtgaagat gttgaagtcc	360	
gacgcaacgg agaaggacct gtccgatctg atctcggaga tggagatgat gaaaatgatt	420	
gggaagcaca agaatatcat caaccttctg ggagcgtgca cacaggatgg tcctctttat	480	
gtcattgtgg agtacgcctc caaaggcaat ctccgggagt atctacaggc ccggaggcct	540	
cctgggctgg agtactgcta taaccccagc cacaaccccg aggaacagct gtcttccaaa	600	
gatctggtat cctgtgccta tcaggtggct cggggcatgg agtatcttgc ctctaagaag	660	
tgtatacacc gagacctggc tgctaggaac gtctgtgtga ccgaggataa cgtaatgaag	720	
atcgagactc ttggcttagc tcgagacatt catcatatcg actactacaa gaaaaccacc	780	
aacggccggc tgctgtgaa gtggatggcc cctgaggcgt tgtttgaccg gatctacaca	840	
caccagagcg atgtgtgttc ttttggagtg ctcttgtggg agatcttcac tctgggtggc	900	
tccccatacc ccggtgtgcc tgtggaggaa cttttcaagc tgctgaagga gggtcatcga	960	
atggacaagc ccagtaactg taccaatgag ctgtacatga tgatgcggga ctgctggcat	1020	
gcagtgcctc ctcagagacc tacgttcaag cagttgggtg aagacctgga ccgcattgtg	1080	
gccttgacct ccaaccagga gtatctggac ctgtccatac cgctggacca gtactcacc	1140	
agctttcccg acacacggag ctccacctgc tcctcagggg aggactctgt cttctctcat	1200	
gagccggtac ctgaggagcc ctgtctgcct cgacacccca cccagcttgc caacagtgga	1260	
ctcaaacggc gc	1272	
SEQ ID NO: 6	moltype = AA length = 424	
FEATURE	Location/Qualifiers	
REGION	1..424	
	note = Amino acid sequence of FGFR1	
source	1..424	
	mol_type = protein	
	organism = synthetic construct	
SEQUENCE: 6		
MKSGTKKSDF HSQMAVHKLA KSIPLRRQVT VSADSSASMN SGVLLVRPSR LSSSGTPMLA	60	
GVSEYELPED PRWELPRDL VLGKPLGEGC FGQVLAELAI GLDKDKPNRV TKVAVKMLKS	120	
DATEKDLSDL ISEMEMMKMI GKHKNIINLL GACTQDGPLY VIVEYASKGN LREYLQARRP	180	
PGLEYCYNPS HNPPEQLSSK DLVSCAYQVA RGMEYLASKK CIHRDLAARN VLVTEDNVMK	240	
IADFLGARDI HHIDYYKKT NGRLPVKWMA PEALFDRIYT HQSDVWSFGV LLWEIFTLGG	300	
SPYPGVPVEE LFKLLKEGHR MDKPSNCTNE LYMMMRDCWH AVPSQRPTFK QLVEDLDRIV	360	
ALTSNQEYLD LSIPLDQYSP SFPDTRSSTC SSGEDSVFSH EPLPEEPCLP RHPTQLANS	420	
LKRR	424	
SEQ ID NO: 7	moltype = DNA length = 435	
FEATURE	Location/Qualifiers	
misc_feature	1..435	
	note = Nucleotide sequence encoding LOV domain	
source	1..435	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 7		
cctgactaca gtctcgtgaa ggctctgcaa atggcacaa	agaattttgt cattacagac	60
gcctccctcc cagacaaccc tatcgtctac gccagtagag	ggtttctgac actgacaggc	120
tattctctcg accagatcct gggcaggaa	tgcaagggcc agaaacagac	180
ccaagagctg tggataagat caggaatgcc atcaccaaag	gcgttgatac cagtgtctgt	240
ctgctgaatt atagacagga tggcacaacc ttctggaatc	tcttcttcgt ggctggactc	300

-continued

```

agagattcta agggcaatat tgtcaactac gtcggagtgc agtcaaaggt gagegaagat 360
tatgccaaagc tgctggtcaa cgagcagaac attgagtaca aaggtgtgcg caccagtaac 420
atgctgcgca gaaag 435

SEQ ID NO: 8          moltype = AA  length = 145
FEATURE              Location/Qualifiers
REGION              1..145
                    note = Amino acid sequence of LOV domain
source              1..145
                    mol_type = protein
                    organism = synthetic construct

SEQUENCE: 8
PDYSLVKALQ MAQQNFVITD ASLPDNPIVY ASRGFLTLTG YSLDQILGRN CRFLQGPETD 60
PRAVDKIRNA ITKGVDTSVC LLNRYQDGTG FWNLFVAVGL RDSKGNIVNY VGVQSKVSED 120
YAKLLVNEQN IEYKGVRTSN MLRRK 145

SEQ ID NO: 9          moltype = DNA  length = 20
FEATURE              Location/Qualifiers
misc_feature        1..20
                    note = guide RNA
source              1..20
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 9
gtcccctcca cccacagtg 20

SEQ ID NO: 10         moltype = DNA  length = 20
FEATURE              Location/Qualifiers
misc_feature        1..20
                    note = pRosa26-gRNA#1
source              1..20
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 10
cctggcttaa cctgattctt 20

SEQ ID NO: 11         moltype = DNA  length = 20
FEATURE              Location/Qualifiers
misc_feature        1..20
                    note = pRosa26-gRNA#91
source              1..20
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 11
gtgagagtta tctgaccgta 20

SEQ ID NO: 12         moltype = DNA  length = 23
FEATURE              Location/Qualifiers
misc_feature        1..23
                    note = DPPA3 forward primer
source              1..23
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 12
ttaatccaac ctacatccca ggg 23

SEQ ID NO: 13         moltype = DNA  length = 22
FEATURE              Location/Qualifiers
misc_feature        1..22
                    note = DPPA3 reverse primer
source              1..22
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 13
aggggaaaca gattcgctac ta 22

SEQ ID NO: 14         moltype = DNA  length = 24
FEATURE              Location/Qualifiers
misc_feature        1..24
                    note = ESG1 forward primer
source              1..24
                    mol_type = other DNA
                    organism = synthetic construct

SEQUENCE: 14
atatcccgcc gtgggtgaaa gttc 24

```

-continued

SEQ ID NO: 15	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
misc_feature	1..24	
	note = ESG1 reverse primer	
source	1..24	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 15		
actcagccat ggactggagc atcc		24
SEQ ID NO: 16	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = FOXA2 forward primer	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 16		
cgactggagc agctactatg c		21
SEQ ID NO: 17	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = FOXA2 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 17		
tacgtgttca tgccgttcac		20
SEQ ID NO: 18	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = GAPDH forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 18		
cgagatccct ccaaaatcaa		20
SEQ ID NO: 19	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = GAPDH reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 19		
ttctagacgg caggtcaggt		20
SEQ ID NO: 20	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = KLF4 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 20		
tcccatcttt ctccacgttc		20
SEQ ID NO: 21	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = KLF4 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 21		
ggatcggata ggtgaagctg		20
SEQ ID NO: 22	moltype = DNA length = 18	
FEATURE	Location/Qualifiers	
misc_feature	1..18	
	note = MSGN1 forward primer	
source	1..18	
	mol_type = other DNA	

-continued

SEQUENCE: 22	organism = synthetic construct	
agcgaaggct gcagtgtc		18
SEQ ID NO: 23	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = MSGN1 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 23		
tggcctctct ggctgtagac		20
SEQ ID NO: 24	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = NANOG forward primer	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 24		
catgagtgtg gatccagctt g		21
SEQ ID NO: 25	moltype = DNA length = 21	
FEATURE	Location/Qualifiers	
misc_feature	1..21	
	note = NANOG reverse primer	
source	1..21	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 25		
cctgaataag cagatccatg g		21
SEQ ID NO: 26	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = OCT4 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 26		
agtgagaggc aacctggaga		20
SEQ ID NO: 27	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = OCT4 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 27		
acactcggac cacatccttc		20
SEQ ID NO: 28	moltype = DNA length = 24	
FEATURE	Location/Qualifiers	
misc_feature	1..24	
	note = REX1 forward primer	
source	1..24	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 28		
cagatcctaa acagctcgca gaat		24
SEQ ID NO: 29	moltype = DNA length = 23	
FEATURE	Location/Qualifiers	
misc_feature	1..23	
	note = REX1 reverse primer	
source	1..23	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 29		
gcgtacgcaa attaaagtcc aga		23
SEQ ID NO: 30	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	

-continued

misc_feature	1..20	
	note = PAX6 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 30		
ctttgcttgg gaaatccgag		20
SEQ ID NO: 31	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = PAX6 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 31		
agccagggttg cgaagaactc		20
SEQ ID NO: 32	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = SOX1 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 32		
cctcggatct ctggtcaagt		20
SEQ ID NO: 33	moltype = DNA length = 22	
FEATURE	Location/Qualifiers	
misc_feature	1..22	
	note = SOX1 reverse primer	
source	1..22	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 33		
gcaggtacat gctgatcatc tc		22
SEQ ID NO: 34	moltype = DNA length = 26	
FEATURE	Location/Qualifiers	
misc_feature	1..26	
	note = SOX2 forward primer	
source	1..26	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 34		
gggaaatggg aggggtgcaa aagagg		26
SEQ ID NO: 35	moltype = DNA length = 26	
FEATURE	Location/Qualifiers	
misc_feature	1..26	
	note = SOX2 reverse primer	
source	1..26	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 35		
ttgcgtgagt gtggatggga ttggtg		26
SEQ ID NO: 36	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = SOX17 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 36		
cgcacggaat ttgaacagta		20
SEQ ID NO: 37	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = SOX17 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 37		

-continued

ggatcagggg cctgtcacac	20
SEQ ID NO: 38	moltype = DNA length = 20
FEATURE	Location/Qualifiers
misc_feature	1..20
	note = TBX6 forward primer
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 38	
aagtaccaac cccgcataca	20
SEQ ID NO: 39	moltype = DNA length = 20
FEATURE	Location/Qualifiers
misc_feature	1..20
	note = TBX6 reverse primer
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 39	
taggctgtca cggagatgaa	20
SEQ ID NO: 40	moltype = DNA length = 19
FEATURE	Location/Qualifiers
misc_feature	1..19
	note = pGapdh forward primer
source	1..19
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 40	
ctcaacgggga agctcactg	19
SEQ ID NO: 41	moltype = DNA length = 20
FEATURE	Location/Qualifiers
misc_feature	1..20
	note = pGapdh reverse primer
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 41	
ccctgttgct gtagccaaat	20
SEQ ID NO: 42	moltype = DNA length = 18
FEATURE	Location/Qualifiers
misc_feature	1..18
	note = pGata4 forward primer
source	1..18
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 42	
gcttcgcggg ctctact	18
SEQ ID NO: 43	moltype = DNA length = 18
FEATURE	Location/Qualifiers
misc_feature	1..18
	note = pGata4 reverse primer
source	1..18
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 43	
ccggttgatg ccattcat	18
SEQ ID NO: 44	moltype = DNA length = 20
FEATURE	Location/Qualifiers
misc_feature	1..20
	note = pLin28a forward primer
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 44	
tgccggcatc tgtaaattggt	20
SEQ ID NO: 45	moltype = DNA length = 20
FEATURE	Location/Qualifiers
misc_feature	1..20
	note = pLin28a reverse primer

-continued

source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 45		
gcagtttgca ttccttgga		20
SEQ ID NO: 46	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20 note = pMsgn1 forward primer	
source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 46		
cgctggagtc ctattcttcg		20
SEQ ID NO: 47	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20 note = pMsgn1 reverse primer	
source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 47		
gtctgtgagt tccccgatgt		20
SEQ ID NO: 48	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
misc_feature	1..19 note = pNanog forward primer	
source	1..19 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 48		
ttgccccgaa gcatccatt		19
SEQ ID NO: 49	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20 note = pNanog reverse primer	
source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 49		
ccagctctga ttacccaca		20
SEQ ID NO: 50	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20 note = pPax6 forward primer	
source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 50		
agttcttcgc aacctggcta		20
SEQ ID NO: 51	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20 note = pPax6 reverse primer	
source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 51		
catttgcccc ttcgattaga		20
SEQ ID NO: 52	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20 note = pOct4 forward primer	
source	1..20 mol_type = other DNA organism = synthetic construct	
SEQUENCE: 52		
tgaggctttg cagctcagtt		20

-continued

SEQ ID NO: 53	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pOct4 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 53		
actgcttgat cgtttgcct		20
SEQ ID NO: 54	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pSox1 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 54		
cacccggatt acaagtaccg		20
SEQ ID NO: 55	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pSox1 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 55		
gagttggaga tggggctgta		20
SEQ ID NO: 56	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pSox2 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 56		
taagtacaca ctgcccggag		20
SEQ ID NO: 57	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pSox2 reverse primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 57		
catggaaccg agcgcatgc		20
SEQ ID NO: 58	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pSox17 forward primer	
source	1..20	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 58		
tcggggacat gaagatgaag		20
SEQ ID NO: 59	moltype = DNA length = 19	
FEATURE	Location/Qualifiers	
misc_feature	1..19	
	note = pSox17 reverse primer	
source	1..19	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 59		
gcggccggta cttgtagtt		19
SEQ ID NO: 60	moltype = DNA length = 20	
FEATURE	Location/Qualifiers	
misc_feature	1..20	
	note = pTbx6 forward primer	
source	1..20	
	mol_type = other DNA	

-continued

organism = synthetic construct	
SEQUENCE: 60	
agtatcagcc ccgcatacac	20
SEQ ID NO: 61	moltype = DNA length = 20
FEATURE	Location/Qualifiers
misc_feature	1..20
	note = pTbx6 reverse primer
source	1..20
	mol_type = other DNA
	organism = synthetic construct
SEQUENCE: 61	
ctgctcgga tctgactctc	20

1-33. (canceled)

34. A method for maintaining cells, the method comprising:

- (a) providing or obtaining, in a culture medium, cells genetically engineered to be maintained by exposure to light; and
- (b) illuminating the cells with light at a sufficient light intensity and time such that the cells are maintained, thereby maintaining the cells,

wherein the method does not require a step of supplementing the culture medium with at least one growth factor required for maintaining the cells.

35. The method of claim 34, wherein the illuminating of (b) replaces the step of supplementing the culture medium with the at least one growth factor required for maintaining the cells.

36. The method of claim 34, wherein the cells provided or obtained are capable of self-renewal.

37. The method of claim 36, wherein, after maintaining the cells, the cells are capable of self-renewal.

38. The method of claim 34, wherein the cells provided or obtained have the potential to differentiate or to further differentiate into different cell types.

39. The method of claim 38, wherein, after maintaining the cells, the cells have the potential to differentiate or to further differentiate into different cell types.

40. The method of claim 34, wherein the cells provided or obtained are stem cells.

41. The method of claim 40, wherein the stem cells are pluripotent stem cells.

42. The method of claim 40, wherein the stem cells are embryonic stem cells.

43. The method of claim 40, wherein the stem cells are induced pluripotent stem cells.

44. The method of claim 40, wherein the maintaining comprises maintaining pluripotency of the stem cells.

45. The method of claim 34, wherein the cells provided or obtained are genetically engineered to express a fusion protein comprising a growth factor receptor protein or a portion thereof fused to a photoactivatable domain.

46. The method of claim 45, wherein the photoactivatable domain is selected from the group consisting of: cryptochrome 2 (Cry2), light, oxygen, and voltage (LOV) domain, phytochrome B (PhyB), and UV-resistance locus 8 (UVR8).

47. The method of claim 45, wherein the growth factor receptor protein or portion thereof is a fibroblast growth factor receptor (FGFR), a transforming growth factor receptor (TGFR), a portion thereof, or a domain thereof.

48. The method of claim 47, wherein the growth factor receptor protein or portion thereof is FGFR or an intracellular domain of FGFR.

49. The method of claim 48, wherein the method results in phosphorylation of ERK1/2.

50. The method of claim 48, wherein the method does not require a step of supplementing the culture medium with exogenous fibroblast growth factor (FGF).

51. The method of claim 34, wherein the cells are maintained for at least 1 week.

52. The method of claim 34, wherein the cells provided or obtained are mammalian cells.

53. The method of claim 34, wherein the cells provided or obtained are from a livestock animal.

54. The method of claim 53, wherein the livestock animal is selected from the group consisting of: a pig, a cow, a sheep, and a goat.

55. The method of claim 34, wherein the cells provided or obtained are from a chicken.

56. The method of claim 34, wherein the cells provided or obtained are from a human.

57. The method of claim 34, wherein the light is produced by an illumination source.

58. The method of claim 57, wherein the illumination source comprises a light-emitting diode (LED).

59. The method of claim 34, wherein the intensity is from about 0.1 $\mu\text{W}/\text{mm}^2$ to about 25 $\mu\text{W}/\text{mm}^2$.

60. The method of claim 57, wherein the illumination source: (i) illuminates the cells for a period of time from about 1 minute to about 120 minutes; (ii) illuminates the cells in a time interval of from about 30 minutes to about 4 hours; or (iii) both (i) and (ii).

61. The method of claim 57, wherein the illumination source continuously illuminates the cells.

62. The method of claim 57, wherein the illumination source illuminates the cells with a pulsed light pattern.

63. The method of claim 57, wherein the illumination source spatiotemporally controls illumination of the cells.

* * * * *