



US 20230130218A1

(19) **United States**

(12) **Patent Application Publication**

WISNIEWSKI et al.

(10) **Pub. No.: US 2023/0130218 A1**

(43) **Pub. Date: Apr. 27, 2023**

(54) **IMMUNOTHERAPEUTIC MODULATION OF AMYLOIDOGENIC DISEASE USING NON-FIBRILLOGENIC, NON-AMYLOIDOGENIC POLYMERIZED PROTEINS AND PEPTIDES**

(71) Applicant: **NEW YORK UNIVERSITY**, New York, NY (US)

(72) Inventors: **Thomas M. WISNIEWSKI**, Staten Island, NY (US); **Fernando GONI**, New York, NY (US)

(21) Appl. No.: **17/743,854**

(22) Filed: **May 13, 2022**

Related U.S. Application Data

- (63) Continuation of application No. 15/933,146, filed on Mar. 22, 2018, now Pat. No. 11,332,506, which is a continuation of application No. 13/550,316, filed on Jul. 16, 2012, now Pat. No. 9,926,353.
- (60) Provisional application No. 61/509,320, filed on Jul. 19, 2011, provisional application No. 61/509,442, filed on Jul. 19, 2011.

Publication Classification


- (51) **Int. Cl.**
C07K 14/47 (2006.01)
G01N 33/68 (2006.01)
A61K 9/00 (2006.01)
A61K 39/00 (2006.01)
C07K 16/18 (2006.01)
- (52) **U.S. Cl.**
CPC *C07K 14/4711* (2013.01); *G01N 33/6896* (2013.01); *A61K 9/0056* (2013.01); *A61K 39/0007* (2013.01); *C07K 16/18* (2013.01); *G01N 2333/4709* (2013.01); *G01N 2800/28* (2013.01); *G01N 2800/2821* (2013.01); *A61K 2039/64* (2013.01); *C07K 2317/20* (2013.01); *A61K 2039/523* (2013.01)

(57) **ABSTRACT**

The present invention is directed to polymerized products and compositions useful for the treatment and prevention of amyloid disease in a subject. The invention further relates to isolated antibodies that recognize a common conformational epitope of amyloidogenic proteins or peptides that are useful for the diagnosis, treatment, and prevention of amyloid disease.


Specification includes a Sequence Listing.

Locomotor Testing TgSwDI



Figures 1A–1D


Locomotor Testing Tg 3x



Figures 2A–2D

Tg3x Vaccination

A.



B.






Figure 3A–3B



Figures 3C



Figures 4A–4B






Figure 4C


C.
3x Tg Mice Vaccinated with combined pABri and Aβ1-30KK




D.
3x Tg Mice Vehicle Vaccinated



Figures 5C–5D




Figures 6A–6C




Figures 7A–7B

A.



B.



Figures 8A–8B




Figure 9




Figure 10




Figure 11




Figure 12

**IMMUNOTHERAPEUTIC MODULATION OF
AMYLOIDOGENIC DISEASE USING
NON-FIBRILLOGENIC,
NON-AMYLOIDOGENIC POLYMERIZED
PROTEINS AND PEPTIDES**

[0001] This application is a continuation of U.S. patent application Ser. No. 13/550,316, filed Jul. 16, 2012, which claims the benefit of U.S. Provisional Patent Application Ser. Nos. 61/509,320 and 61/509,442, both filed Jul. 19, 2011, which are hereby incorporated by reference in their entirety.

[0002] This invention was made with government support under grant numbers AG20245 and NS073501 awarded by the National Institutes of Health. The government has certain rights in this invention.

FIELD OF THE INVENTION

[0003] The present invention relates to pharmaceutical agents and antibodies suitable for the diagnosis, prevention, and treatment of amyloid disease.

BACKGROUND OF THE INVENTION

[0004] Amyloidosis broadly encompasses a variety of diseases that are characterized by the extracellular or intracellular deposition of amyloid proteins in tissues and/or organs. Amyloids are insoluble fibrous protein/peptide aggregates and their deposition may occur in localized sites or systemically. The fibrillar composition of these deposits is an identifying characteristic for the various forms of amyloid disease. In some cases the amyloid protein/peptide accumulates intracellularly, resulting in cell dysfunction and ultimately cell death. Examples of intracellular amyloid proteins include α -synuclein, forming Lewy bodies in Parkinson's disease, and huntingtin, forming neuronal inclusions in Huntington disease. The pathogenesis of Alzheimer's disease (AD), the most common of the amyloid related neurodegenerative disorders, is linked to the cleavage of the amyloid precursor protein generating the amyloid- β (A β) peptide which undergoes a shape change into a pathological conformer having a high β -sheet content. Intracerebral and cerebrovascular deposits composed primarily of fibrils of the pathological A β peptide are characteristic of both familial and sporadic forms of AD. In addition to A β , abnormally folded and phosphorylated tau protein forms toxic oligomeric structures and neurofibrillary tangles in AD. Similar to AD, prion-associated diseases, such as Creutzfeld-Jacob disease, have also been characterized as amyloid diseases. The pathogenesis of prion disease is linked to a change of the cellular prion protein (PrP^C) into the disease associated PrP^{Sc} (Sc for scrapie). Currently, there is no effective therapy for any of these disorders.

[0005] An active area of translational research and current clinical trials for amyloid disease has focused on immunotherapy, using both passive and active immunization against amyloid proteins, particularly A β in AD (Wisniewski et al., "Amyloid- β Immunization for Alzheimer's Disease," *Lancet Neurol* 7:805-811 (2008)). Although immunotherapy holds great promise as a means of reducing amyloid deposition, it, unfortunately, has been accompanied by major obstacles. Specific problems associated with immunotherapy that were identified in a clinical trial for AD include the potential of toxicity from encephalitis (related to excessive cell mediated immunity), the immunological targeting

of both the normal and abnormal A β peptide, the failure to address tau related pathology, and the apparent poor efficacy. Moreover, although autopsy data from this early immunotherapy vaccine trial suggested that many patients had a significant reduction in amyloid burden, these patients exhibited only minor cognitive benefits (Wisniewski et al., "Amyloid- β Immunization for Alzheimer's Disease," *Lancet Neurol* 7:805-811 (2008) and Holmes et al., "Long Term Effects of A β 42 Immunization in Alzheimer's Disease: Immune Response, Plaque Removal and Clinical Function," *Lancet* 372:216-223 (2008)). Therefore, an immunotherapeutic approach that can effectively reduce amyloid burden and overcome the aforementioned problems is warranted.

[0006] The present invention is directed to overcoming these and other deficiencies in the art.

SUMMARY OF THE INVENTION

[0007] A first aspect of the present invention relates to a non-amyloidogenic, non-fibrillogenic polymerized product comprising two or more protein and/or peptide units, where each unit is independently selected from the group consisting of an amyloid-beta (A β) peptide, an α -synuclein protein or peptide, a tau protein or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a prion protein (PrP) protein or peptide, and any combination thereof.

[0008] A second aspect of the present invention is directed to an isolated antibody or binding portion thereof having antigenic specificity for an epitope a non-amyloidogenic, non-fibrillogenic polymerized product. The polymerized product comprises two or more protein and/or peptide units, each unit independently selected from the group consisting of an amyloid-beta (A β) peptide, an α -synuclein protein or peptide, a tau protein or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a prion protein (PrP) protein or peptide, and any combination thereof.

[0009] The development of an effective immunotherapeutic approach for the prevention and treatment of amyloid related diseases has been hindered by potential cell-mediated toxicity, non-specific immunological targeting of both normal and amyloidogenic proteins, and overall poor efficacy. The immunotherapeutic approach of the present invention employs non-amyloidogenic, non-fibrillogenic polymerized protein and/or peptides and co-polymerized proteins and/or peptides to generate a specific immunological response to conformational epitopes that are shared by various amyloidogenic proteins, thereby overcoming many of the above noted obstacles.

BRIEF DESCRIPTION OF THE DRAWINGS

[0010] FIGS. 1A-1D show the results of locomotor activity testing in SwDI transgenic (Tg) control mice and Tg mice vaccinated with synthetic polymerized A β peptide (A β 1-30₁₈K₁₉K), polymerized ABri peptide (ABri-Glut), or polymerized A β 1-42 peptide. No significant differences between the groups were noted in distance traveled (FIG. 1A) or mean velocity (V_{mean}) (FIG. 1C). A β 1-42 vaccinated Tg mice exhibited slower maximum velocity (V_{max}) (FIG. 1B) ($p=0.0234$ post-hoc only; A β 1-42 versus control $p<0.05$) and longer resting time (FIG. 1D) ($p=0.0311$ post-hoc only; A β 1-42 versus control $p<0.05$) compared to controls. No significant differences between the A β 1-30₁₈K₁₉K

and ABri treated Tg animals and controls in these parameters were observed (FIGS. 1B and 1D)

[0011] FIGS. 2A-2D show the results of locomotor activity testing in Tg3x transgenic control mice and Tg mice vaccinated with synthetic polymerized A β peptide (A β 1-30K₁₈K₁₉), polymerized ABri peptide (ABri-Glut), or the combination of A β 1-30K₁₈K₁₉ and ABri polymerized peptides. No significant differences between the groups were noted in distance traveled (FIG. 2A), maximum velocity (Vmax) (FIG. 2B), mean velocity (Vmean) (FIG. 2C), or in resting time (FIG. 2D).

[0012] FIGS. 3A-3C illustrate the effects of polymerized peptide vaccination on behavioral improvement in 3xTg and SwDI transgenic animal models of amyloidogenic disease. FIG. 3A is a graph depicting the results of the radial arm maze. The number of errors (y-axis) versus the day of testing (x-axis) are plotted for 3xTg mice vaccinated with polymerized ABri, transgenic control (Tg control), and wildtype. Vaccinated transgenic animals commit significantly fewer errors than non-treated transgenic and perform similarly to the wildtype animals. $p=0.0044$ one-way ANOVA; $p<0.01$ wildtype and Tg pABri treated vs. Tg control; $p=>0.05$ wildtype vs. Tg pABri treated. FIG. 3B is a similar graph showing the radial arm maze results for 3xTg animals vaccinated with polymerized ABri, polymerized A β 1-30K₁₈K₁₉, and the combination (polymerized ABri and A β 1-30K₁₈K₁₉). All treatment groups committed significantly fewer errors than the transgenic control animals. $p=0.004$ by one-way ANOVA; $p<0.01$ all Tg Treated groups and wild-type versus Tg Control; no significant difference between wild-type and Tg treated groups. FIG. 3C is another graph depicting the results of the radial arm maze cognitive testing. In this graph the number of errors (y-axis) versus the day of testing (x-axis) are plotted for SwDI transgenic mice vaccinated with polymerized ABri, polymerized A β 1-30K₁₈K₁₉, polymerized A β 1-42, transgenic control mice, and wildtype mice (WT). All treatment groups committed significantly fewer errors than the transgenic control animals ($p<0.01$ all Tg Treated groups and wild-type versus Tg Control).

[0013] FIGS. 4A-4C are bar graphs showing IgM and IgG antibody levels raised in TgSwDI mice vaccinated with polymerized ABri peptide (pABri) (FIG. 4A), polymerized A β 1-30K₁₈K₁₉ (FIG. 4B), and polymerized A β 1-42 peptide (FIG. 4C). Antibody titers against A β 1-40, A β 1-42, pABri, and a mutant A β 1-30KK were measured in vaccinated mice prior to the first inoculation (T0), after the 6th inoculation (T1) and at the time of sacrifice (TF). (** $p<0.0001$, ** $p<0.01$, * $p>0.05$ versus T0).

[0014] FIGS. 5A-5D are bar graphs showing IgM and IgG antibody levels raised in 3xTg mice vaccinated with polymerized ABri peptide (FIG. 5A), polymerized A β 1-30K₁₈K₁₉ (FIG. 5B), the combination of polymerized peptides (FIG. 5C), and vehicle (FIG. 5D). Antibody titers against A β 1-40, A β 1-42, pABri, and a mutant A β 1-30KK were measured in vaccinated mice prior to the first inoculation (T0), after the 6th inoculation (T1), and at the time of sacrifice (TF). (** $p<0.0001$, ** $p<0.01$, * $p>0.05$ versus T0).

[0015] FIGS. 6A-6C depict the amyloid and paired helical fiber (PHF1) burden in Tg3x transgenic control and vaccinated animals. FIG. 6A is a bar graph showing the reduction of hippocampal amyloid burden in Tg3x animals administered polymerized ABri (Tg-pABri), polymerized A β 1-30K₁₈K₁₉ (Tg-AB1-30KK), or the combination of polym-

erized peptides (Tg-combined) compared to transgenic control (Tg-control) animals ($p=0.001$ by one way ANOVA post-hoc testing; $p<0.01$ all treatment groups vs. control). FIGS. 6B and 6C show a reduction in PHF1 in the hippocampus and cortex, respectively, of Tg3x animals administered polymerized ABri or the combination of polymerized ABri and A β 1-30K₁₈K₁₉ ($p<0.0001$ one way ANOVA post hoc; $p<0.001$ Tg-ABri and Tg-combined vs. Tg-control and Tg-A β 1-30K₁₈K₁₉ for both hippocampal and cortical analyses). No significant difference was found between Tg-control and Tg-A β 1-30K₁₈K₁₉ treated animals.

[0016] FIGS. 7A-7B depict the amyloid burden in SwDI transgenic control and vaccinated animals. FIG. 7A is a bar graph showing the reduction in hippocampal amyloid burden in polymerized ABri vaccinated (Tg pABri), polymerized A β 1-30K₁₈K₁₉ vaccinated (Tg-A β 1-30KK), and polymerized A β 1-42 vaccinated (Tg-A β 42) animals compared to transgenic control (Tg Control). FIG. 7B is an electron micrograph (EM) image of negatively stained polymerized A β 1-42 peptide, which is predominately in the form of spherical particles of ~200 nm.

[0017] FIGS. 8A-8B are EM images of aged A β 1-42 peptides. In FIG. 8A A β 1-42 peptide was aged for 3 months after controlled polymerized with glutaraldehyde. There is no evidence of fibril formation, and the sample was typical of oligomerized peptides/proteins. FIG. 8B is an A β 1-42 peptide from the same synthesis batch as the peptide in FIG. 8A. This peptide was dissolved in saline and aged for 3 months before the EM. In contrast to the polymerized A β 1-42 peptide, the unpolymerized A β 1-42 sample was completely fibrillized.

[0018] FIG. 9 shows specific anti-PrP IgA antibodies in feces of control and deer vaccinated with prion protein (PrP) as measured by ELISA. White tail deer were orally inoculated with an attenuated salmonella carrying deer-PrP or an empty vector (control) four times over a period of a few months following the protocol for mucosal vaccination described in Goni et al., "High Titers of Mucosal and Systemic Anti-PrP Antibodies Abrogate Oral Prion Infection in Mucosal-Vaccinated Mice," *Neurosci.* 153:679-686 (2008), which is hereby incorporated by reference in its entirety. Animals were subsequently orally boosted with a mixture of polymerized PrP and polymerized recombinant PrP peptides as described in the Examples herein. Antibody titers were measure before the first inoculation (T0), after the fourth inoculation (T5), and ten days after the boost (T6). The boost with PrP and polymerized PrP fragments was repeated two months after the first boost, and titers were again measured 10 days after the second boost (T7). The control group did not develop any noticeable titer whereas the vaccinated group showed some increase in mucosa titer after the immune response was established with the salmonella oral delivery. The titers were very low (T5) but were greatly enhanced after the animals were boosted with the polymerized PrP and PrP fragments (T6 and T7) showing the importance of these antigenic preparation to have a sustainable immune response.

[0019] FIG. 10 shows serum anti-PrP IgM antibodies of control deer and deer vaccinated with PrP as described in FIG. 9. Again the control group did not show any noticeable increase in antibody titer; whereas the vaccinated group showed some concomitant IgM titer in serum at the same time of the mucosal response (T5). Anti-PrP antibody titers in the vaccinated animals greatly increased after the two

boosts with the polymerized PrP and PrP fragments showing that this type of boost could invoke a therapeutic serum response. Vaccinated vs. control at T0 not significant, * $p < 0.05$, ** $p = 0.0009$, *** $p < 0.0001$ (two-tailed test).

[0020] FIG. 11 shows immunoblots of salmonella lysate (lane 1), sheep PrP (lane 2), polymerized sheep PrP (lane 3), deer PrP (lane 4), and polymerized deer PrP (lane 5) that were developed using purified antibodies from vaccinated animal 781 at T7 and control animal 786 at T7. Both animals mounted a good immune response to salmonella (lane 1); however, only the vaccinated animal had antibodies against different deer PrP molecules (lane 4) and the oligomers present in the polymerized deer PrP (lane 5). Vaccinated vs. control at T0 not significant, * $p < 0.017$, ** $p = 0.0022$, *** $p < 0.01$ (two-tailed test).

[0021] FIG. 12 is a Kaplan and Meier survival curve showing the protective effect of PrP vaccination in White Tail Deer at 18 months after challenge. Three out of six control animals became sick with the prionoses Chronic Wasting Disease and were properly euthanized. None of the vaccinated animals had signs of the disease at 18-months. Protection for the progression of the disease is evident and due to the immune response elicited by the inoculations involving the polymerized PrP and PrP fragments.

DETAILED DESCRIPTION OF THE INVENTION

[0022] A first aspect of the present invention relates to a non-amyloidogenic, non-fibrillogenic polymerized product comprising two or more protein and/or peptide units, where each unit is independently selected from the group consisting of an amyloid-beta ($A\beta$) peptide, an α -synuclein protein or peptide, a tau peptide or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a PrP protein or peptide, and any combination thereof.

[0023] In accordance with this aspect of the invention suitable $A\beta$ peptides of the polymerized product include, without limitation, $A\beta$ 1-42 and peptides derived therefrom. The amino acid sequence of $A\beta$ 1-42 is provided as SEQ ID NO:1 below.

(SEQ ID NO: 1)
daefrhds gy evhhqklvff aedvgsnkga iiglmvggvv ia

[0024] Accordingly, in one embodiment of the invention, the $A\beta$ peptide of the polymerized product comprises amino acid residues 1-42 of SEQ ID NO:1. Alternatively suitable $A\beta$ peptides of the polymerized product include, without

[0025] In another embodiment of the present invention, the $A\beta$ peptide of the polymerized product contains two amino acid substitutions at amino acid residues 18 and 19 of SEQ ID NO:1. In one embodiment the amino acid substitutions comprise a valine to lysine substitution at position 18 and a phenylalanine to lysine substitution at position 19 of SEQ ID NO:1. Peptides having these two amino acid substitutions are referred to as $A\beta$ K₁₈K₁₉ peptides and are derived from the amino acid sequence of SEQ ID NO:2 below.

(SEQ ID NO: 2)
daefrhds gy evhhqklk~~kf~~ aedvgsnkga iiglmvggvv ia

[0026] Accordingly, suitable $A\beta$ peptides comprising the double lysine substitution include, without limitation, peptides comprising amino acid residues 1-30 of SEQ ID NO:2 ($A\beta$ 1-30K₁₈K₁₉), amino acid residues 1-40 of SEQ ID NO:2 ($A\beta$ 1-40K₁₈K₁₉), amino acid residues 1-42 of SEQ ID NO:2 ($A\beta$ 1-42K₁₈K₁₉), amino acid residues 1-20 of SEQ ID NO:2 ($A\beta$ 1-20K₁₈K₁₉), amino acid residues 10-30 of SEQ ID NO:2 ($A\beta$ 10-30K₁₈K₁₉), amino acid residues 10-40 of SEQ ID NO:2 ($A\beta$ 10-40K₁₈K₁₉), amino acid residues 10-42 of SEQ ID NO:2 ($A\beta$ 10-42K₁₈K₁₉), amino acid residues 20-40 of SEQ ID NO:2 ($A\beta$ 20-40K₁₈K₁₉), and amino acid residues 20-42 of SEQ ID NO:2 ($A\beta$ 20-42K₁₈K₁₉).

[0027] The polymerized product of the present invention may comprise a homopolymer of an $A\beta$ peptide or a copolymer of two or more different $A\beta$ peptides. Alternatively, the polymerized product of the present invention may comprise a copolymer of an $A\beta$ protein or peptide copolymerized with any one or more of an α -synuclein protein or peptide, a tau protein or peptide, a TDP-43 protein or peptide, an amylin protein or peptide, and/or a PrP protein or peptide. In one embodiment of the present invention, the polymerized product comprises a copolymer of one or more $A\beta$ peptides copolymerized with one or more α -synuclein proteins or peptides. In another embodiment of the present invention, the polymerized product comprises a copolymer of one or more $A\beta$ peptides copolymerized with one or more tau proteins or peptides. In another embodiment of the present invention, the polymerized product comprises a copolymer of one or more $A\beta$ peptides copolymerized with one or more TDP-43 proteins or peptides

[0028] The polymerized product of the present invention may also comprise an α -synuclein protein or peptide thereof. The α -synuclein protein comprises the amino acid sequence of SEQ ID NO:3.

(SEQ ID NO: 3)
1 mdvfmkglsk akegvvaaae ktkqgvaaee gsktkegvlyv gsktkegvvh gvatvaektk
61 eqvtnvvggav vtgvtavaqk tvegagsiaa atgfvkdkql gkneegapqe giledmpvdp
121 dneayempse egyqdyepa

limitation, peptides comprising amino acid residues 1-16 of SEQ ID NO:1, amino acid residues 1-20 of SEQ ID NO:1, amino acid residues 1-30 of SEQ ID NO:1, amino acid residues 1-40 of SEQ ID NO:1, amino acid residues 1-42 of SEQ ID NO:1, amino acid residues 10-30 of SEQ ID NO:1, amino acid residues 20-40 of SEQ ID NO:1, and amino acid residues 20-42 of SEQ ID NO:1.

[0029] Accordingly, in one embodiment of the invention, the α -synuclein protein of the polymerized product comprises amino acid residues 1-140 of SEQ ID NO:3. Peptides of α -synuclein derived from the amino acid sequence of SEQ ID NO:3 are also suitable for use in the polymerized product of the present invention. In one embodiment, the peptide comprises at least five contiguous amino acids of

SEQ ID NO:3. In another embodiment, the peptide comprises at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60, at least 65, at least 70, at least 75, at least 80, at least 90 or at least 100 contiguous amino acids of SEQ ID NO:3.

[0030] In one embodiment of the invention, the α -synuclein peptide is a C-terminal α -synuclein peptide as disclosed in WO 2004/041067 to Schenk et al., which is hereby incorporated by reference in its entirety. Suitable C-terminal α -synuclein peptides for use in the polymerized product of the present invention include, without limitation, peptides comprising amino acid residues 70-140 of SEQ ID NO:3, amino acid residues 100-140 of SEQ ID NO:3, amino acid residues 120-140 of SEQ ID NO:3, amino acid residues 130-140 of SEQ ID NO:3, or amino acid residues 135-140 of SEQ ID NO:3.

[0031] The polymerized product of the present invention may comprise a homopolymer of an α -synuclein protein or peptide or a copolymer of two or more different α -synuclein

proteins or peptides. Alternatively, the polymerized product of the present invention may comprise a copolymer of an α -synuclein protein or peptide co-polymerized with any one or more of an A β peptide, a tau protein or peptide, a TDP-43 protein or peptide, an amylin protein or peptide, and/or a PrP protein or peptide.

[0032] The pharmaceutical composition of the present invention may also comprise a tau protein or peptide thereof. Suitable tau proteins of the present invention include any of the eight isoforms of the human tau protein that are known in the art (see NCBI Accession Nos. NP_058519.3, NP_005901, NP_058518.1, NP_058525.1, NP_001116539.1, NP_001116538.2, NP_001190180.1, and NP_001190181.1, which are hereby incorporated by reference in their entirety). The tau isoforms vary in the number of N-terminal inserts resulting from the splicing of exons two and three, and microtubule-binding domains resulting from the splicing of exon ten. The amino acid sequence corresponding to human tau protein isoform 2 (441a.a), containing two N-terminal inserts and four microtubule binding (2N4R) domains is provided as SEQ ID NO:4 below.

(SEQ ID NO: 4)

Met	Ala	Glu	Pro	Arg	Gln	Glu	Phe	Glu	Val	Met	Glu	Asp	His	Ala	Gly	1	5	10	15
Thr	Tyr	Gly	Leu	Gly	Asp	Arg	Lys	Asp	Gln	Gly	Gly	Tyr	Thr	Met	His	20	25	30	
Gln	Asp	Gln	Glu	Gly	Asp	Thr	Asp	Ala	Gly	Leu	Lys	Glu	Ser	Pro	Leu	35	40	45	
Gln	Thr	Pro	Thr	Glu	Asp	Gly	Ser	Glu	Glu	Pro	Gly	Ser	Glu	Thr	Ser	50	55	60	
Asp	Ala	Lys	Ser	Thr	Pro	Thr	Ala	Glu	Asp	Val	Thr	Ala	Pro	Leu	Val	65	70	75	80
Asp	Glu	Gly	Ala	Pro	Gly	Lys	Gln	Ala	Ala	Ala	Gln	Pro	His	Thr	Glu	85	90	95	
Ile	Pro	Glu	Gly	Thr	Thr	Ala	Glu	Glu	Ala	Gly	Ile	Gly	Asp	Thr	Pro	100	105	110	
Ser	Leu	Glu	Asp	Glu	Ala	Ala	Gly	His	Val	Thr	Gln	Ala	Arg	Met	Val	115	120	125	
Ser	Lys	Ser	Lys	Asp	Gly	Thr	Gly	Ser	Asp	Asp	Lys	Lys	Ala	Lys	Gly	130	135	140	
Ala	Asp	Gly	Lys	Thr	Lys	Ile	Ala	Thr	Pro	Arg	Gly	Ala	Ala	Pro	Pro	145	150	155	160
Gly	Gln	Lys	Gly	Gln	Ala	Asn	Ala	Thr	Arg	Ile	Pro	Ala	Lys	Thr	Pro	165	170	175	
Pro	Ala	Pro	Lys	Thr	Pro	Pro	Ser	Ser	Gly	Glu	Pro	Pro	Lys	Ser	Gly	180	185	190	
Asp	Arg	Ser	Gly	Tyr	Ser	Ser	Pro	Gly	Ser	Pro	Gly	Thr	Pro	Gly	Ser	195	200	205	
Arg	Ser	Arg	Thr	Pro	Ser	Leu	Pro	Thr	Pro	Pro	Thr	Arg	Glu	Pro	Lys	210	215	220	
Lys	Val	Ala	Val	Val	Arg	Thr	Pro	Pro	Lys	Ser	Pro	Ser	Ser	Ala	Lys	225	230	235	240
Ser	Arg	Leu	Gln	Thr	Ala	Pro	Val	Pro	Met	Pro	Asp	Leu	Lys	Asn	Val	245	250	255	
Lys	Ser	Lys	Ile	Gly	Ser	Thr	Glu	Asn	Leu	Lys	His	Gln	Pro	Gly	Gly	260	265	270	

-continued

Gly Lys Val Gln Ile Ile Asn Lys Lys Leu Asp Leu Ser Asn Val Gln
275 280 285

Ser Lys Cys Gly Ser Lys Asp Asn Ile Lys His Val Pro Gly Gly Gly
290 295 300

Ser Val Gln Ile Val Tyr Lys Pro Val Asp Leu Ser Lys Val Thr Ser
305 310 315 320

Lys Cys Gly Ser Leu Gly Asn Ile His His Lys Pro Gly Gly Gly Gln
325 330 335

Val Glu Val Lys Ser Glu Lys Leu Asp Phe Lys Asp Arg Val Gln Ser
340 345 350

Lys Ile Gly Ser Leu Asp Asn Ile Thr His Val Pro Gly Gly Gly Asn
355 360 365

Lys Lys Ile Glu Thr His Lys Leu Thr Phe Arg Glu Asn Ala Lys Ala
370 375 380

Lys Thr Asp His Gly Ala Glu Ile Val Tyr Lys Ser Pro Val Val Ser
385 390 395 400

Gly Asp Thr Ser Pro Arg His Leu Ser Asn Val Ser Ser Thr Gly Ser
405 410 415

Ile Asp Met Val Asp Ser Pro Gln Leu Ala Thr Leu Ala Asp Glu Val
420 425 430

Ser Ala Ser Leu Ala Lys Gln Gly Leu

[0033] Accordingly, in one embodiment, a tau protein of the polymerized product of the present invention comprises amino acid residues 1-441 of SEQ ID NO:4. Alternatively, the composition of the invention comprises amino acid residues 1-758 of tau isoform 1 (NP_058519.3), amino acid residues 1-383 of tau isoform 3 (NP_058518.1), amino acid residues 1-352 of tau isoform 4 (NP_058525.1), amino acid residues 1-412 of tau isoform 5 (NP_001116539.1), amino acid residue 1-776 of tau isoform 6 (NP_001116538.2), amino acid residues 1-381 of tau isoform 7 (NP_001190180.1), and amino acid residues 1-410 of tau isoform 8 (NP_001190181.1).

[0034] The pharmaceutical composition of the present invention may also comprise tau peptides derived from the amino acid sequence of SEQ ID NO:4 or the amino acid sequence of any other tau isoform. In one embodiment, the tau peptide comprises at least five contiguous amino acids of a tau protein, e.g., the tau protein of SEQ ID NO:4. In another embodiment, the tau peptide comprises at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60, at least 65, at least 70, at least 75, at least 80, at least 90, or at least 100 contiguous amino acids of a tau protein, e.g., the tau protein of SEQ ID NO:4. In yet another embodiment, the peptide comprises at least 200, at least 300, at least 400, or at least 500 contiguous amino acids, of a tau protein.

[0035] A number of tau peptides that are suitable for polymerization and incorporation into the polymerized product of the present invention are disclosed in U.S. Pat.

No. 8,012,936 to Sigurdsson et al., which is hereby incorporated by reference in its entirety. Suitable peptides include, without limitation, amino acid residues 1-30 of SEQ ID NO:4, amino acid residues 30-60 of SEQ ID NO:4, amino acid residues 60-90 of SEQ ID NO:4, amino acid residues 90-120 of SEQ ID NO:4, amino acid residues 120-150 of SEQ ID NO:4, amino acid residues 150-180 of SEQ ID NO:4, amino acid residues 210-240 of SEQ ID NO:4, amino acid residues 270-300 of SEQ ID NO:4, amino acid residues 300-330 of SEQ ID NO:4, amino acid residues 330-360 of SEQ ID NO:4, amino acid residues 360-390 of SEQ ID NO:4, amino acid residues 390-420 of SEQ ID NO:4, amino acid residues 411-440 of SEQ ID NO:4, amino acid residues 133-162 of SEQ ID NO:4, amino acid residues 379-409 of SEQ ID NO:4, amino acid residues 192-221 of SEQ ID NO:4, amino acid residues 221-250 of SEQ ID NO:4, or amino acid residues 184-213 of SEQ ID NO:4.

[0036] The polymerized product of the present invention may comprise a homopolymer of tau protein or peptide or a copolymer of two or more different tau proteins or peptides. Alternatively, the polymerized product of the present invention may comprise a copolymer of a tau protein or peptide co-polymerized with any one or more of an A β peptide, an α -synuclein protein or peptide, a TDP-43 protein or peptide, an amylin protein or peptide, and/or a PrP protein or peptide.

[0037] The polymerized product of the present invention may also comprise a TAR DNA-binding protein 43 (TDP 43) or peptide thereof. The human TDP 43 protein comprises the amino acid sequence of SEQ ID NO:5.

(SEQ ID NO: 5)

1 mseyirvted endepieips eddgtvllst vtaqfpgacg lryrnpvsqc mrgvrlvegi

61 lhapdagwgn lvyvvnypkd nkrkmdetda ssavkvkrav qktsdlivlg lpwktteqdl

-continued
121 keyfstfgev lmvqvkkdlk tghskgfgfv rfteyetqvk vmsqqrhmidg rwccklplns
181 kqsqdeplrs rkfvvgrcte dmtedelref fsqygdvmdv fipkpfrafa fvtfaddqia
241 qslcgedlii kgisvhisna epkhnsnrql ersgrfggnp ggfgnqggfg nsrgggaglg
301 nnqgsnmggg mnfgafsinp ammaaqaal qsswgmmgm1 asqqnqsgps gnnqnqgnmq
361 repnqafgsg nnsysgsnsg aaigwgsasn agsgsgfngg fgssmdskss gwgm

[0038] Accordingly, in one embodiment of the invention, the TDP-43 protein of the polymerized product comprises amino acid residues 1-414 of SEQ ID NO:5. Peptides of TDP-43 derived from the amino acid sequence of SEQ ID NO:5 are also suitable for use in the pharmaceutical composition of the present invention. In one embodiment, the TDP-43 peptide comprises at least five contiguous amino acids of SEQ ID NO:5. In another embodiment, the TDP-43 peptide comprises at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60, at least 65, at least 70, at least 75, at least 80, at least 90, at least 100, at least 150, at least 200, at least 250, or at least 300 contiguous amino acids of SEQ ID NO:5.
[0039] The polymerized product of the present invention may comprise a homopolymer of a TDP-43 protein or

25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60 contiguous amino acids of SEQ ID NO:6.
[0042] The polymerized product of the present invention may comprise a homopolymer of an amylin protein or peptide or a copolymer of two or more different amylin proteins or peptides. Alternatively, the polymerized product of the present invention may comprise a copolymer of an amylin protein or peptide co-polymerized with any one or more of an Aβ peptide, a tau protein or peptide, an α-synuclein protein or peptide, a TDP-43 protein or peptide, and/or a PrP protein or peptide.
[0043] The polymerized product of the present invention may also comprise a prion protein or peptide thereof. The human prion protein comprises the amino acid sequence of SEQ ID NO:7.

(SEQ ID NO: 7)
1 manlgcwmlv lfvatwsdlg lckkrpkpgg wntggsrypg qgsppgnryp pqggggwgqpp
61 hggggwgqphg ggwgqphggg wgqphgggwg qgggthsqwn kpskpktnmk hmagaaaaga
121 vvgglggyvl gsamsrpiih fgsdyedryy renmhrypny vyyrpmdeys nqnnfvhdcv
181 nitikqhtvt tttkgenfte tdkmmervv eqmcitqyer esqayykrqs smvlfssppv
241 illisflifl ivg

peptide or a copolymer of two or more different TDP-43 proteins or peptides. Alternatively, the polymerized product of the present invention may comprise a copolymer of a TDP-43 protein or peptide co-polymerized with any one or more of an Aβ peptide, a tau protein or peptide, an α-synuclein protein or peptide, an amylin protein or peptide, and/or a PrP protein or peptide.
[0040] The polymerized product of the present invention can also comprise an amylin protein or peptide thereof. The human amylin protein comprises the amino acid sequence of SEQ ID NO:6.

[0044] Accordingly, in one embodiment of the invention, the prion protein of the polymerized product comprises amino acid residues 1-253 of SEQ ID NO:7 or analogs thereof. Peptides of prion protein derived from the amino acid sequence of SEQ ID NO:7 are also suitable for use in the polymerized product of the present invention. In one embodiment, the peptide comprises at least five contiguous amino acids of SEQ ID NO:7. In another embodiment, the peptide comprises at least 10, at least 15, at least 20, at least 25, at least 30, at least 35, at least 40, at least 45, at least 50, at least 55, at least 60, at least 65, at least 70, at least 75, at

(SEQ ID NO: 6)
1 hqvekrkent atcatqrlan flvhssnnfg ailsstnvgs ntygkrnave vlkreplnyl
61 pl

[0041] Accordingly, in one embodiment of the invention, the amylin protein of the polymerized product comprises amino acid residues 1-62 of SEQ ID NO:6. Peptides of amylin derived from the amino acid sequence of SEQ ID NO:6 are also suitable for use in the pharmaceutical composition of the present invention. In one embodiment, the amylin peptide comprises at least five contiguous amino acids of SEQ ID NO:6. In another embodiment, the amylin peptide comprises at least 10, at least 15, at least 20, at least

least 80, at least 90 or at least 100, at least 125, and least 150, at least 200, or at least 225 contiguous amino acids of SEQ ID NO:7.
[0045] The polymerized product of the present invention may comprise a homopolymer of an PrP protein or peptide or a copolymer of two or more different PrP proteins or peptides. Alternatively, the polymerized product of the present invention may comprise a copolymer of an PrP protein or peptide co-polymerized with any one or more of an Aβ peptide, a tau protein or peptide, an α-synuclein protein or

peptide, a TDP-43 protein or peptide, and/or an amylin protein or peptide as described herein.

[0046] The polymerized product of the present invention may further comprise one or more polymerized ABri peptides (CSRTVKKNIEEN; SEQ ID NO: 8), ADan peptides (CFNLFLNSQEKHY; SEQ ID NO:9), or ABri/ADan fusion peptides (CSRTVKKNIEENGSGSGCFNLFLNSQEKHY; SEQ ID NO:10) as disclosed in U.S. Patent Application Publication No. 20100284909 to Wisniewski et al., which is hereby incorporated by reference in its entirety.

[0047] The proteins and peptides of the polymerized product of the present invention may comprise naturally occurring peptides or analog peptides. Analog peptides typically differ from naturally occurring peptides at one, two, or a few positions, often by virtue of conservative substitutions. Analogs typically exhibit at least 80 or 90% sequence identity with natural peptides. Some analogs include unnatural amino acids or modifications of N or C terminal amino acids at one, two or a few positions. Examples of unnatural amino acids are D-amino acids, alpha amino acids, alpha-disubstituted amino acids, N-alkyl amino acids, lactic acid, 4-hydroxyproline, gamma-carboxyglutamate, epsilon-N,N, N-trimethyllysine, epsilon-N-acetyllysine, O-phosphoserine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine, omega-N-methylarginine, beta-alanine, ornithine, norleucine, norvaline, hydroxyproline, thyroxine, gamma-amino butyric acid, homoserine, citrulline, and isoaspartic acid. Analog peptides can be screened for prophylactic or therapeutic efficacy using animal models as described in the Examples herein.

[0048] In another embodiment of the present invention, the polymerized product comprises a polymer of a fusion peptide. Suitable fusion peptides contain any first Aβ, α-synuclein, tau, TDP-43, amylin, or prion protein peptide of the present invention fused to any second Aβ, α-synuclein, tau, TDP-43, amylin, or prion protein peptide of the present invention. The fusion peptide preferably contains a short linker sequence coupling the first peptide to a second peptide. Preferred linker sequences include glycine-rich (e.g. G₃₋₅) or serine-rich (e.g. GSG, GSGS (SEQ ID NO: 13), GSGSG (SEQ ID NO: 14), GS_NG) linker sequences.

[0049] In accordance with this aspect of the present invention the polymerized product may contain a polymer or copolymer of any one or more of the Aβ, α-synuclein, tau, TDP-43, amylin, or prion protein proteins or peptides described herein further linked in-frame to an adjuvant polypeptide. The adjuvant polypeptide can be any adjuvant polypeptide known in the art, including, but not limited to, cholera toxin B, flagellin, human papillomavirus L1 or L2 protein, herpes simplex glycoprotein D (gD), complement C4 binding protein, TL4 ligand, and IL-1β. The protein or peptides comprising the polymers of the polymerized product of the present invention may be linked directly to the adjuvant polypeptide or coupled to the adjuvant by way of a short linker sequence. Suitable linker sequences include glycine or serine-rich linkers described supra or other flexible immunoglobulin linkers as disclosed in U.S. Pat. No. 5,516,637 to Huang et al, which is hereby incorporated by reference in its entirety.

[0050] In another embodiment, the polymerized product of the present invention may contain a polymer or copolymer of any one of the Aβ, α-synuclein, tau, TDP-43, amylin, or prion protein proteins or peptides described supra conjugated to an immunogenic carrier molecule. The immuno-

genic carrier molecule can be covalently or non-covalently bonded to the proteins or peptides as described herein. Suitable immunogenic carrier molecules include, but are not limited to, serum albumins, chicken egg ovalbumin, keyhole limpet hemocyanin, tetanus toxoid, thyroglobulin, pneumococcal capsular polysaccharides, CRM 197, immunoglobulin molecules, alum, and meningococcal outer membrane proteins. Other suitable immunogenic carrier molecules include T-cell epitopes, such as tetanus toxoid (e.g., the P2 and P30 epitopes), Hepatitis B surface antigen, pertussis, toxoid, diphtheria toxoid, measles virus F protein, *Chlamydia trachomatis* major outer membrane protein, *Plasmodium falciparum* circumsporozoite T, *P. falciparum* CS antigen, *Schistosoma mansoni* triose phosphate isomerase, *Escherichia coli* TraT, and Influenza virus hemagglutinin (HA). Other suitable immunogenic carrier molecules include promiscuous T helper cell epitopes which are derived from hepatitis B virus, *Bordetella pertussis*, *Clostridium tetani*, *Pertusaria trachythallina*, *E. coli*, *Chlamydia trachomatis*, Diphtheria, *P. falciparum*, and *Schistosoma mansoni* (see U.S. Pat. No. 6,906,169 to Wang; U.S. Patent Application Publication No. 20030068325 to Wang, and WO 2002/096350 to Wang, which are hereby incorporated by reference in their entirety). Yet other suitable carriers include T-helper cell epitopes derived from tetanus toxin, cholera toxin B, pertussis toxin, diphtheria toxin, measles virus F protein, hepatitis B virus surface antigen, *C. trachomatis* major outer membrane protein, *P. falciparum* circumsporozoite, *S. mansoni* triose phosphate isomerase, or *E. coli* TraT (see WO01/42306 to Chain, which is hereby incorporated by reference in its entirety).

[0051] The peptides and proteins of the present invention can be linked to immunogenic carrier molecules by chemical crosslinking prior to polymerization. Techniques for linking a peptide immunogen to an immunogenic carrier molecule include the formation of disulfide linkages using N-succinimidyl-3-(2-pyridyl-thio) propionate (SPDP) and succinimidyl 4-(N-maleimidomethyl)cyclohexane-1-carboxylate (SMCC) (if the peptide lacks a sulfhydryl group, this can be provided by addition of a cysteine residue). These reagents create a disulfide linkage between themselves and peptide cysteine residues on one protein, and an amide linkage through the epsilon-amino on a lysine, or other free amino group in other amino acids. A variety of such disulfide/amide-forming agents are described by Jansen et al., "Immunotoxins: Hybrid Molecules Combining High Specificity and Potent Cytotoxicity," *Immun Rev* 62:185-216 (1982), which is hereby incorporated by reference in its entirety. Other bifunctional coupling agents form a thioether rather than a disulfide linkage. Many of these thio-ether-forming agents are commercially available and include reactive esters of 6-maleimidocaproic acid, 2-bromoacetic acid, 2-iodoacetic acid, and 4-(N-maleimido-methyl)cyclohexane-1-carboxylic acid. The carboxyl groups can be activated by combining them with succinimide or 1-hydroxyl-2-nitro-4-sulfonic acid, sodium salt.

[0052] The one or more proteins or peptides comprising the polymers or copolymers of the polymerized product of the present invention can be synthesized by solid phase or solution phase peptide synthesis, recombinant expression, or can be obtained from natural sources. Automatic peptide synthesizers are commercially available from numerous suppliers, such as Applied Biosystems, Foster City, Calif. Standard techniques of chemical peptide synthesis are well

known in the art (see e.g., SYNTHETIC PEPTIDES: A USERS GUIDE 93-210 (Gregory A. Grant ed., 1992), which is hereby incorporated by reference in its entirety). Protein or peptide production via recombinant expression can be carried out using bacteria, such as *E. coli*, yeast, insect or mammalian cells and expression systems. Procedures for recombinant protein/peptide expression are well known in the art and are described by Sambrook et al, Molecular Cloning: A Laboratory Manual (C.S.H.P. Press, NY 2d ed., 1989).

[0053] Recombinantly expressed peptides can be purified using any one of several methods readily known in the art, including ion exchange chromatography, hydrophobic interaction chromatography, affinity chromatography, gel filtration, and reverse phase chromatography. The peptide is preferably produced in purified form (preferably at least about 80% or 85% pure, more preferably at least about 90% or 95% pure) by conventional techniques. Depending on whether the recombinant host cell is made to secrete the peptide into growth medium (see U.S. Pat. No. 6,596,509 to Bauer et al., which is hereby incorporated by reference in its entirety), the peptide can be isolated and purified by centrifugation (to separate cellular components from supernatant containing the secreted peptide) followed by sequential ammonium sulfate precipitation of the supernatant. The fraction containing the peptide is subjected to gel filtration in an appropriately sized dextran or polyacrylamide column to separate the peptides from other proteins. If necessary, the peptide fraction may be further purified by HPLC.

[0054] Polymerization of the proteins or peptides alone or conjugated to an adjuvant polypeptide or immunogenic carrier molecule can be achieved using standard techniques known in the art. As used herein, polymerization refers to process of reacting two or more peptide and/or protein units together under suitable conditions to form three-dimensional networks or polymer chains. As described herein, the proteins and peptides can be polymerized by a reaction with a cross linking reagent. Suitable cross-linking reagents include, but are not limited to glutaraldehyde and 1-Ethyl-3-[3-dimethylaminopropyl]carbodiimide hydrochloride (EDC) (see Goni et al., "Immunomodulation Targeting Abnormal Protein Conformation Reduced Pathology in a Mouse Model of Alzheimer's Disease," *PLoS One* 5(10): e13391 (2010), which is hereby incorporated by reference in its entirety). Alternatively, the proteins and peptides can be polymerized by cysteine oxidation induced disulfide cross linking.

[0055] Another aspect of the present invention is directed to a pharmaceutical composition comprising the polymerized product of the invention and a pharmaceutical carrier.

[0056] In accordance with this aspect of the present invention, the pharmaceutical composition may contain a single homopolymer of an A β protein or peptide, α -synuclein protein or peptide, tau protein or peptide, TDP-43 protein or peptide, amylin protein or peptide, or prion protein or peptide. Alternatively, the pharmaceutical composition may contain a mixture of one or more proteins or peptides, i.e. heteropolymers of the one or more aforementioned proteins and/or peptides.

[0057] The pharmaceutical composition of the present invention can further contain, in addition to peptide polymers, other pharmaceutically acceptable components (see REMINGTON'S PHARMACEUTICAL SCIENCE (19th ed., 1995), which is hereby incorporated by reference in its entirety). The incorporation of such pharmaceutically acceptable compo-

nents depends on the intended mode of administration and therapeutic application of the pharmaceutical composition. Typically, however, the pharmaceutical composition will include a pharmaceutically-acceptable, non-toxic carrier or diluent, which are defined as vehicles commonly used to formulate pharmaceutical compositions for animal or human administration. The diluent is selected so as not to affect the biological activity of the composition. Exemplary carriers or diluents include distilled water, physiological phosphate-buffered saline, Ringer's solutions, dextrose solution, and Hank's solution.

[0058] Pharmaceutical compositions can also include large, slowly metabolized macromolecules such as proteins, polysaccharides such as chitosan, polylactic acids, polyglycolic acids and copolymers (such as latex functionalized sepharose, agarose, cellulose), polymeric amino acids, amino acid copolymers, and lipid aggregates (such as oil droplets or liposomes).

[0059] The pharmaceutical composition of the present invention can further contain an adjuvant. One class of preferred adjuvants is aluminum salts, such as aluminum hydroxide, aluminum phosphate, or aluminum sulfate. Such adjuvants can be used with or without other specific immunostimulating agents such as MPL or 3-DMP, QS-21, flagellin, attenuated *Salmonella* (e.g., *Salmonella typhimurium*), polymeric or monomeric amino acids such as polyglutamic acid or polylysine, or pluronic polyols. Oil-in-water emulsion formulations are also suitable adjuvants that can be used with or without other specific immunostimulating agents such as muramyl peptides (e.g., N-acetylmuramyl-L-threonyl-D-isoglutamine (thr-MDP), N-acetyl-normuramyl-L-alanyl-D-isoglutamine (nor-MDP), N-acetylmuramyl-L-alanyl-D-isoglutaminyl-L-alanine-2-(1'-2'dipalmitoyl-sn-glycero hydroxyphosphoryloxy)-ethylamine (MTP-PE), N-acetylglucosaminyl-N-acetylmuramyl-L-Al-D-isoglu-L-Ala-dipalmitoxy propylamide (DTP-DPP) Theramide™, or other bacterial cell wall components). A suitable oil-in-water emulsion is MF59 (containing 5% Squalene, 0.5% Tween 80, and 0.5% Span 85 (optionally containing various amounts of MTP-PE) formulated into submicron particles using a microfluidizer such as Model 110Y microfluidizer (Microfluidics, Newton Mass.) as described in WO90/14837 to Van Nest et al., which is hereby incorporated by reference in its entirety. Other suitable oil-in-water emulsions include SAF (containing 10% Squalene, 0.4% Tween 80, 5% pluronic-blocked polymer L121, and thr-MDP, either microfluidized into a submicron emulsion or vortexed to generate a larger particle size emulsion) and Ribi™ adjuvant system (RAS; containing 2% squalene, 0.2% Tween 80, and one or more bacterial cell wall components). Another class of preferred adjuvants is saponin adjuvants, such as Stimulon™ (QS-21) or particles generated therefrom such as ISCOMs (immunostimulating complexes) and ISCOMATRIX. Other suitable adjuvants include incomplete or complete Freund's Adjuvant (IFA), cytokines, such as interleukins (IL-1, IL-2, and IL-12), macrophage colony stimulating factor (M-CSF), lysolecithin, tumor necrosis factor (TNF), and liposome polycation DNA particles. Such adjuvants are generally available from commercial sources.

[0060] In another embodiment of the present invention, the pharmaceutical composition further includes a delivery vehicle. Suitable delivery vehicles include, but are not

limited to biodegradable microspheres, microparticles, nanoparticles, liposomes, collagen minipellets, and cochleates.

[0061] In one embodiment of this aspect of the invention, the pharmaceutical agent includes a mucosal delivery system. A preferred mucosal delivery system consists of attenuated *Salmonella* (e.g., *Salmonella typhimurium*) with a non-toxic fragment C of tetanus toxin (TetC) or glutathione S-transferase (GST). Methods of mucosal vaccination via oral administration of *S. typhimurium* are described in Goni et al., “Mucosal Vaccination Delays or Prevents Prion Infection via an Oral Route,” *Neuroscience* 133:413-21 (2005), and Goni et al., “High Titers of Mucosal and Systemic Anti-PrP Antibodies Abrogate Oral Prion Infection in Mucosal-Vaccinated Mice,” *Neuroscience* 153:679-686 (2008), which are hereby incorporated by reference in their entirety.

[0062] Another aspect of the present invention relates to a method of inducing an immune response against an amyloidogenic form of a protein or peptide in a subject. This method involves administering to the subject a polymerized product of the present invention under conditions effective to induce an immune response against the amyloidogenic form of the protein or peptide in the subject. In a preferred embodiment of this aspect of the present invention, a subject that would benefit from an immune response against an amyloidogenic form of a protein or peptide is selected prior to administering the polymerized product.

[0063] As used herein, an “amyloid protein”, “amyloidogenic protein”, and “amyloidogenic form of a protein” are used interchangeably to encompass any insoluble fibrous protein/peptide aggregate that is deposited intra- or extracellularly within the body. Amyloidogenic protein/peptide deposition may be organ-specific (e.g., central nervous system, pancreas, etc.) or systemic. In accordance with this aspect of the invention, amyloidogenic proteins/peptides subject to deposition include beta protein precursor, prion and prion proteins, α -synuclein, tau, ABri precursor protein, ADan precursor protein, amylin, apolipoprotein AI, apolipoprotein AII, lysozyme, cystatin C, gelsolin, protein, atrial natriuretic factor, calcitonin, keratopithelin, lactoferrin, immunoglobulin light chains, transthyretin, A amyloidosis, β 2-microglobulin, immunoglobulin heavy chains, fibrinogen alpha chains, prolactin, keratin, and medin. Amyloid deposition may occur as its own entity or as a result of another illness (e.g., multiple myeloma, chronic infection, or chronic inflammatory disease).

[0064] In accordance with this aspect of the present invention, an immune response is the development of a beneficial humoral (antibody mediated) and/or a cellular (mediated by antigen-specific T cells or their secretion products) response directed against the polymerized, non-fibrillogenic proteins or peptides of the polymerized product and cross-reactive with any amyloidogenic protein. Such a response is an active response induced by administration of the immunogenic polymerized protein and/or peptides and represents a therapeutic means for clearing or removing amyloid protein deposits from the body of the subject.

[0065] The presence of a humoral immunological response can be determined and monitored by testing a biological sample (e.g., blood, plasma, serum, urine, saliva feces, CSF or lymph fluid) from the subject for the presence of antibodies directed to the immunogenic component of the administered polymerized product. Methods for detecting antibodies in a biological sample are well known in the art,

e.g., ELISA, Dot blots, SDS-PAGE gels or ELISPOT. The presence of a cell-mediated immunological response can be determined by proliferation assays ($CD4^+$ T cells) or CTL (cytotoxic T lymphocyte) assays which are readily known in the art.

[0066] The present invention is further directed to a method of preventing and/or treating a condition mediated by an amyloidogenic protein or peptide in a subject. This method involves administering to the subject, a polymerized product of the present invention containing one or more polymers or co-polymers comprising an amyloid-beta ($A\beta$) peptide, an α -synuclein protein or peptide, a tau protein or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a prion protein (PrP) or peptide or any combination thereof as described supra. The polymerized product is administered under conditions effective to treat the condition mediated by the amyloidogenic protein or peptide in the subject. In a preferred embodiment of this aspect of the present invention, a subject at risk of having or having a condition mediated by an amyloidogenic protein or peptide is selected prior to administering the polymerized product of the present invention.

[0067] Conditions or diseases associated with, or resulting from, the deposition of amyloidogenic proteins or peptides include, but are not limited to, Alzheimer’s disease, diffuse Lewy body disease, Down’s syndrome, fronto-temporal dementia, Parkinson’s disease, hereditary cerebral hemorrhage with amyloidosis, kuru, Creutzfeldt-Jakob disease, Gerstmann-Straussler-Scheinker disease, fatal familial insomnia, British familial dementia, Danish familial dementia, familial corneal amyloidosis, Familial corneal dystrophies, medullary thyroid carcinoma, insulinoma, type 2 diabetes, isolated atrial amyloidosis, pituitary amyloidosis, aortic amyloidosis, plasma cell disorders, familial amyloidosis, senile cardiac amyloidosis, inflammation-associated amyloidosis, familial Mediterranean fever, dialysis-associated amyloidosis, systemic amyloidosis, and familial systemic amyloidosis. In accordance with this aspect of the present invention, administration of the polymerized product is effective to stimulate an immune response in the subject that is effective at reducing and/or clearing the amyloidogenic protein that is causing or exacerbating the aforementioned disease conditions.

[0068] A second aspect of the present invention is directed to an isolated antibody or binding portion thereof having antigenic specificity for an epitope a non-amyloidogenic, non-fibrillogenic polymerized product of the present invention. As described infra, the polymerized product of the present invention comprises two or more protein or peptide units, each unit independently selected from the group consisting of an amyloid-beta ($A\beta$) peptide, an α -synuclein protein or peptide, a tau protein or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a prion protein (PrP) protein or peptide, and any combination thereof.

[0069] As used herein, “epitope” refers to an antigenic determinant of the one or more polymerized proteins or peptides of the present invention that is recognized by the isolated antibody. The epitope recognized by the antibody of the present invention may be a linear epitope, i.e. the primary structure of the amino acid sequence of the target proteins or peptides. Preferably, the linear epitope recognized by the isolated antibody of the present invention does

not have amino acid sequence homology to a non-amyloid protein. Alternatively, the epitope recognized by the isolated antibody of the present invention is a non-linear or conformational epitope, i.e. the tertiary or quaternary structure of a polymerized protein or peptide. In one embodiment of the present invention, the non-linear or conformational epitope recognized by the isolated antibody of the present invention is a conformational epitope that is common to or shared with one or more, or all, amyloidogenic proteins. Accordingly, the isolated antibody of the present invention has antigenic specificity for a shared conformational epitope common to all amyloidogenic proteins known in the art.

[0070] An isolated antibody of the present invention encompasses any immunoglobulin molecule that specifically binds to an epitope of a polymerized product of the present invention. Preferably, the antibody of the present invention binds specifically to an epitope that is shared by a polymerized product of the present invention and one or more amyloidogenic proteins. As used herein, the term “antibody” is meant to include intact immunoglobulins derived from natural sources or from recombinant sources, as well as immunoreactive portions (i.e., antigen binding portions) of intact immunoglobulins. The antibodies of the present invention may exist in a variety of forms including, for example, polyclonal antibodies, monoclonal antibodies, intracellular antibodies (“intrabodies”), antibody fragments (e.g. Fv, Fab and F(ab)2), as well as single chain antibodies (scFv), chimeric antibodies, and humanized antibodies (Ed Harlow and David Lane, *USING ANTIBODIES: A LABORATORY MANUAL* (Cold Spring Harbor Laboratory Press, 1999); Houston et al., “Protein Engineering of Antibody Binding Sites: Recovery of Specific Activity in an Anti-Digoxin Single-Chain Fv Analogue Produced in *Escherichia coli*,” *Proc Natl Acad Sci USA* 85:5879-5883 (1988); Bird et al., “Single-Chain Antigen-Binding Proteins,” *Science* 242:423-426 (1988)).

[0071] Antibodies of the present invention may also be synthetic antibodies. A synthetic antibody is an antibody which is generated using recombinant DNA technology, such as, for example, an antibody expressed by a bacteriophage. Alternatively, the synthetic antibody is generated by the synthesis of a DNA molecule encoding and expressing the antibody of the invention or the synthesis of an amino acid specifying the antibody, where the DNA or amino acid sequence has been obtained using synthetic DNA or amino acid sequence technology which is available and well known in the art.

[0072] Methods for monoclonal antibody production may be carried out using the techniques described herein or other well-known in the art (*MONOCLONAL ANTIBODIES—PRODUCTION, ENGINEERING AND CLINICAL APPLICATIONS* (Mary A. Ritter and Heather M. Ladyman eds., 1995), which is hereby incorporated by reference in its entirety). Generally, the process involves obtaining immune cells (lymphocytes) from the spleen of a mammal which has been previously immunized with the antigen of interest (i.e., a polymerized protein or peptide product of the present invention) either in vivo or in vitro. Exemplary polymerized products comprising one or more A β peptides, α -synuclein proteins or peptides, tau proteins or peptides, TDP-43 proteins or peptides, amylin proteins or peptides, and a prion proteins or peptides are described supra.

[0073] The antibody-secreting lymphocytes are fused with myeloma cells or transformed cells, which are capable of

replicating indefinitely in cell culture, thereby producing an immortal, immunoglobulin-secreting cell line. Fusion with mammalian myeloma cells or other fusion partners capable of replicating indefinitely in cell culture is achieved by standard and well-known techniques, for example, by using polyethylene glycol (PEG) or other fusing agents (Milstein and Kohler, “Derivation of Specific Antibody-Producing Tissue Culture and Tumor Lines by Cell Fusion,” *Eur J Immunol* 6:511 (1976), which is hereby incorporated by reference in its entirety). The immortal cell line, which may be murine, but may also be derived from cells of other mammalian species, is selected to be deficient in enzymes necessary for the utilization of certain nutrients, to be capable of rapid growth, and have good fusion capability. The resulting fused cells, or hybridomas, are cultured, and the resulting colonies screened for the production of the desired monoclonal antibodies. Colonies producing such antibodies are cloned, and grown either in vivo or in vitro to produce large quantities of antibody.

[0074] Alternatively monoclonal antibodies can be made using recombinant DNA methods as described in U.S. Pat. No. 4,816,567 to Cabilly et al, which is hereby incorporated by reference in its entirety. The polynucleotides encoding a monoclonal antibody are isolated from mature B-cells or hybridoma cells, for example, by RT-PCR using oligonucleotide primers that specifically amplify the genes encoding the heavy and light chains of the antibody. The isolated polynucleotides encoding the heavy and light chains are then cloned into suitable expression vectors, which when transfected into host cells such as *E. coli* cells, simian COS cells, Chinese hamster ovary (CHO) cells, or myeloma cells that do not otherwise produce immunoglobulin protein, and monoclonal antibodies are generated by the host cells. Also, recombinant monoclonal antibodies or fragments thereof of the desired species can be isolated from phage display libraries (McCafferty et al., “Phage Antibodies: Filamentous Phage Displaying Antibody Variable Domains,” *Nature* 348: 552-554 (1990); Clackson et al., “Making Antibody Fragments using Phage Display Libraries,” *Nature* 352:624-628 (1991); and Marks et al., “By-Passing Immunization. Human Antibodies from V-Gene Libraries Displayed on Phage,” *J. Mol. Biol.* 222:581-597 (1991), which are hereby incorporated by reference in their entirety).

[0075] The polynucleotide(s) encoding a monoclonal antibody can further be modified using recombinant DNA technology to generate alternative antibodies. For example, the constant domains of the light and heavy chains of a mouse monoclonal antibody can be substituted for those regions of a human antibody to generate a chimeric antibody. Alternatively, the constant domains of the light and heavy chains of a mouse monoclonal antibody can be substituted for a non-immunoglobulin polypeptide to generate a fusion antibody. In other embodiments, the constant regions are truncated or removed to generate the desired antibody fragment of a monoclonal antibody. Furthermore, site-directed or high-density mutagenesis of the variable region can be used to optimize specificity and affinity of a monoclonal antibody.

[0076] The monoclonal antibody of the present invention can be a humanized antibody. Humanized antibodies are antibodies that contain minimal sequences from non-human (e.g., murine) antibodies within the variable regions. Such antibodies are used therapeutically to reduce antigenicity and human anti-mouse antibody responses when adminis-

tered to a human subject. In practice, humanized antibodies are typically human antibodies with minimal to no non-human sequences.

[0077] An antibody can be humanized by substituting the complementarity determining region (CDR) of a human antibody with that of a non-human antibody (e.g., mouse, rat, rabbit, hamster, etc.) having the desired specificity, affinity, and capability (Jones et al., "Replacing the Complementarity-Determining Regions in a Human Antibody With Those From a Mouse," *Nature* 321:522-525 (1986); Riechmann et al., "Reshaping Human Antibodies for Therapy," *Nature* 332:323-327 (1988); Verhoeven et al., "Reshaping Human Antibodies: Grafting an Antilysozyme Activity," *Science* 239:1534-1536 (1988), which are hereby incorporated by reference in their entirety). The humanized antibody can be further modified by the substitution of additional residues either in the Fv framework region and/or within the replaced non-human residues to refine and optimize antibody specificity, affinity, and/or capability.

[0078] The monoclonal A β of the present invention can also be a human monoclonal A β . A human antibody is an antibody produced by a human or an antibody having an amino acid sequence corresponding to an antibody produced by a human. Human antibodies can be produced using various techniques known in the art. Immortalized human B lymphocytes immunized in vitro or isolated from an immunized individual that produce an antibody directed against a target antigen can be generated (See e.g., Reisfeld et al., *MONOCLONAL ANTIBODIES AND CANCER THERAPY* 77 (Alan R. Liss ed., 1985) and U.S. Pat. No. 5,750,373 to Garrard, which are hereby incorporated by reference in their entirety). Also, the human antibody can be selected from a phage library, where that phage library expresses human antibodies (Vaughan et al., "Human Antibodies with Sub-Nanomolar Affinities Isolated from a Large Non-immunized Phage Display Library," *Nature Biotechnology*, 14:309-314 (1996); Sheets et al., "Efficient Construction of a Large Nonimmune Phage Antibody Library: The Production of High-Affinity Human Single-Chain Antibodies to Protein Antigens," *Proc. Natl. Acad. Sci. U.S.A.* 95:6157-6162 (1998); Hoogenboom et al., "By-passing Immunisation. Human Antibodies From Synthetic Repertoires of Germline VH Gene Segments Rearranged In Vitro," *J Mol Biol* 227:381-8 (1992); Marks et al., "By-passing Immunization. Human Antibodies from V-gene Libraries Displayed on Phage," *J Mol Biol* 222:581-97 (1991), which are hereby incorporated by reference in their entirety). Human antibodies can also be made in transgenic mice containing human immunoglobulin loci that are capable upon immunization of producing the full repertoire of human antibodies in the absence of endogenous immunoglobulin production. This approach is described in U.S. Pat. No. 5,545,807 to Surani et al.; U.S. Pat. No. 5,545,806 to Lonberg et al.; U.S. Pat. No. 5,569,825 to Lonberg et al.; U.S. Pat. No. 5,625,126 to Lonberg et al.; U.S. Pat. No. 5,633,425 to Lonberg et al.; and U.S. Pat. No. 5,661,016 to Lonberg et al., which are hereby incorporated by reference in their entirety.

[0079] Procedures for raising polyclonal antibodies are also well known in the art. Typically, such antibodies are raised by administering the polymerized product of the present invention subcutaneously to rabbits (e.g., New Zealand white rabbits), goats, sheep, swine, or donkeys which have been bled to obtain pre-immune serum. The polymerized product can be injected in combination with an adju-

vant. The rabbits are bled approximately every two weeks after the first injection and periodically boosted with the same antigen three times every six weeks. Polyclonal antibodies are recovered from the serum by affinity chromatography using the corresponding polymerized product to capture the antibody. This and other procedures for raising polyclonal antibodies are disclosed in Ed Harlow and David Lane, *USING ANTIBODIES: A LABORATORY MANUAL* (Cold Spring Harbor Laboratory Press, 1988), which is hereby incorporated by reference in its entirety.

[0080] As noted above, in addition to whole antibodies, the present invention encompasses binding portions of such antibodies. Such binding portions include the monovalent Fab fragments, Fv fragments (e.g., single-chain antibody, scFv), and single variable V_H and V_L domains, and the bivalent F(ab')₂ fragments, Bis-scFv, diabodies, triabodies, minibodies, etc. These antibody fragments can be made by conventional procedures, such as proteolytic fragmentation procedures, as described in James Goding, *MONOCLONAL ANTIBODIES: PRINCIPLES AND PRACTICE* 98-118 (Academic Press, 1983) and Ed Harlow and David Lane, *ANTIBODIES: A LABORATORY MANUAL* (Cold Spring Harbor Laboratory, 1988), which are hereby incorporated by reference in their entirety, or other methods known in the art.

[0081] Also suitable for use in the present invention are antibody fragments engineered to bind to intracellular proteins, i.e. intrabodies. Although amyloid protein deposits are generally extracellular, intracellular accumulation of certain amyloid proteins (e.g., A β 1-42) have been observed (D'Andrea et al., "Targeting Amyloid Beta: Targeting Intracellular A β 42 for Alzheimer's Disease Drug Discover," *Drug Development Research* 56:194-200 (2002); Knobloch et al., "Intracellular Abeta and Cognitive Deficits Precede beta-Amyloid Deposition in arcAbeta Mice," *Neurobiol Aging* 28(9):1297-306 (2007), which are hereby incorporated by reference in their entirety). Accordingly, an intrabody can be used to bind selectively to an epitope of an amyloid protein within a cell. In a preferred embodiment, the intrabody recognizes an epitope of the A β 1-42 oligomer accumulating within the perikaryon of affected neurons (e.g., pyramidal neurons) in AD.

[0082] Intrabodies are generally obtained by selecting a single variable domain from variable regions of an antibody having two variable domains (i.e., a heterodimer of a heavy chain variable domain and a light chain variable domain). Single chain Fv fragments, Fab fragments, ScFv-Ck fusion proteins, single chain diabodies, V_H-C_H1 fragments, and even whole IgG molecules are suitable formats for intrabody development (Kontermann R. E., "Intrabodies as Therapeutic Agents," *Methods* 34:163-70 (2004), which is here by incorporated by reference in its entirety).

[0083] Intrabodies having antigen specificity for a conformational epitope of an amyloidogenic protein can be obtained from phage display, yeast surface display, or ribosome surface display. Methods for producing libraries of intrabodies and isolating intrabodies of interest are further described in U.S. Published Patent Application No. 20030104402 to Zauderer and U.S. Published Patent Application No. 20050276800 to Rabbitts, which are hereby incorporated by reference in their entirety. Methods for improving the stability and affinity binding characteristics of intrabodies are described in WO2008070363 to Zhenping; Contreras-Martinez et al., "Intracellular Ribosome Display via SecM Translation Arrest as a Selection for Antibodies

with Enhanced Cytosolic Stability,” *J Mol Biol* 372(2):513-24 (2007), which are hereby incorporated by reference in their entirety.

[0084] It may further be desirable, especially in the case of antibody fragments, to modify the antibody in order to increase its serum half-life. This can be achieved, for example, by incorporation of a salvage receptor binding epitope into the antibody fragment by mutation of the appropriate region in the antibody fragment or by incorporating the epitope into a peptide tag that is then fused to the antibody fragment at either end or in the middle (e.g., by DNA or peptide synthesis).

[0085] Antibody mimics are also suitable for use in accordance with the present invention. A number of antibody mimics are known in the art including, without limitation, those known as monobodies, which are derived from the tenth human fibronectin type III domain (¹⁰Fn3) (Koide et al., “The Fibronectin Type III Domain as a Scaffold for Novel Binding Proteins,” *J Mol Biol* 284:1141-1151 (1998); Koide et al., “Probing Protein Conformational Changes in Living Cells by Using Designer Binding Proteins: Application to the Estrogen Receptor,” *Proc Natl Acad Sci USA* 99:1253-1258 (2002), each of which is hereby incorporated by reference in its entirety); and those known as affibodies, which are derived from the stable alpha-helical bacterial receptor domain Z of staphylococcal protein A (Nord et al., “Binding Proteins Selected from Combinatorial Libraries of an alpha-helical Bacterial Receptor Domain,” *Nature Biotechnol* 15(8):772-777 (1997), which is hereby incorporated by reference in its entirety).

[0086] The present invention is further directed to a pharmaceutical composition containing the isolated antibody of the present invention as described supra. In a preferred embodiment, the isolated antibody recognizes and binds to a shared conformational epitope common to one or more amyloid proteins. The pharmaceutical composition of the present invention may contain an antibody mixture where all antibodies recognize the same conformational epitope. Alternatively, the pharmaceutical composition may contain an antibody mixture where one or more antibodies recognize one or more different conformational epitopes of amyloid proteins. The pharmaceutical composition of the present invention further contains a pharmaceutically acceptable carrier or other pharmaceutically acceptable components as described supra.

[0087] Another aspect of the present invention relates to a method of treating a condition mediated by an amyloidogenic protein in a subject. This method involves administering to the subject an antibody of the present invention, where the antibody has antigen specificity for an epitope of a non-amyloidogenic, non-fibrillogenic polymerized product, wherein the polymerized product comprises two or more protein and/or peptide units, each unit independently selected from the group consisting of an amyloid-beta (A β) peptide, an α -synuclein protein or peptide, a tau protein or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a prion protein (PrP) protein or peptide, and any combination thereof. Preferably the antibody has antigen specificity for a shared conformational epitope that is common to one or more amyloidogenic proteins. The antibody or a pharmaceutical composition containing the antibody is administered in an amount effective to treat the condition involving the amyloidogenic protein in the subject. In accordance with this

aspect of the invention, the antibody or pharmaceutical composition containing the antibody is administered in an amount effective to generate passive immunity in the subject against one or more amyloidogenic proteins, thereby facilitating the clearance of amyloid deposits from the subject.

[0088] Conditions mediated by an amyloidogenic protein that are amenable to treatment in accordance with this aspect of the present invention are described supra.

[0089] In a preferred embodiment of this aspect of the present invention, a subject having a condition or at risk of developing a condition mediated by an amyloidogenic protein is selected prior to administration of the antibody of the present invention. Subjects amenable to treatment in accordance with the methods of the present invention include individuals at risk of developing an amyloid related disease but not showing symptoms, as well as subjects showing symptoms at the time of therapeutic intervention (i.e. antibody administration). Diseases subject to treatment include any disease associated with or caused by an amyloidogenic protein as described supra. The pharmaceutical compositions of the present invention contain polymerized products that are not endogenous to the body, or antibodies specific for only pathological protein conformations. Therefore, the risk of inducing an auto-immune response is avoided and prophylactic treatment using these pharmaceutical compositions of the present invention is particularly suitable.

[0090] In the case of Alzheimer’s disease, for example, virtually anyone is at risk of suffering from Alzheimer’s disease if he or she lives long enough. Therefore, the compositions of the present invention can be administered prophylactically to the general population without the need for any assessment of the risk of the subject patient. However, the present methods and compositions are especially suitable for prophylactic treatment of individuals who have a known genetic risk of Alzheimer’s disease or other condition related to an amyloidogenic protein. Genetic markers associated with a risk of Alzheimer’s disease include mutations in the APP gene, particularly mutations at position 717 and positions 670 and 671 referred to as the Hardy and Swedish mutations respectively. Other markers of risk are mutations in the presenilin genes, PS1 and PS2, and mutations in ApoE4, family history of AD, hypercholesterolemia, or atherosclerosis.

[0091] In asymptomatic patients, treatment can begin at any age (e.g., 10, 20, 30). Usually, however, it is not necessary to begin treatment until a patient reaches 40, 50, 60 or 70 years of age. Treatment typically entails multiple dosages over a period of time. Treatment can be monitored by assaying antibody, or activated T-cell or B-cell responses to the therapeutic agent (e.g., polymerized product) over time. If the response falls, a booster dosage is indicated. In the case of potential Down’s syndrome patients, treatment can begin antenatally by administering the therapeutic agent to the mother or shortly after birth.

[0092] In prophylactic applications, the pharmaceutical compositions of the present invention are administered to a patient susceptible to, or otherwise at risk of, a particular disease in an amount sufficient to eliminate or reduce the risk or delay the onset of the disease. In therapeutic applications, pharmaceutical compositions are administered to a patient suspected of, or already suffering from an amyloidogenic disease in an amount sufficient to cure, or at least partially arrest, the symptoms of the disease and its complications. An amount adequate to accomplish this is defined as a thera-

apeutically- or pharmaceutically-effective dose. In both prophylactic and therapeutic regimes, agents are usually administered in several dosages until a sufficient immune response has been achieved. Typically, the immune response is monitored and repeated dosages are given if the immune response starts to fade.

[0093] Effective doses of the compositions of the present invention, for the treatment of the above described conditions vary depending upon many different factors, including means of administration, target site, physiological state of the patient, whether the patient is human or an animal, other medications administered, and whether treatment is prophylactic or therapeutic. Usually, the subject is a human, but in some diseases, such as prion protein related diseases, the subject can be a nonhuman mammal, such as a bovine. Other non-human mammals amenable to treatment in accordance with the methods of the present invention include primates, dogs, cats, rodents (e.g., mouse, rat, guinea pig), horses, deer, cervids, cattle and cows, sheep, and pigs. Treatment dosages need to be titrated to optimize safety and efficacy, and could involve oral treatment.

[0094] When treatment of a subject involves the administration of a polymerized product of the present invention containing one or more polymerized or copolymerized A β , α -synuclein, tau, TDP-43, amylin, and/or PrP protein or peptide immunogens, the appropriate dosage will depend on whether adjuvant is co-administered, with higher dosages being required in the absence of adjuvant. The amount of an immunogen for administration sometimes varies from 1 μ g-500 μ g per patient and more usually from 5-500 μ g per injection for human administration. Occasionally, a higher dose of 1-2 mg per injection is used. Typically about 10, 20, 50 or 100 μ g is used for each human injection. The timing of injections can vary significantly from once a day, to once a week, to once a month, to once a year, to once a decade. Generally an effective dosage can be monitored by obtaining a fluid sample from the patient, generally a blood serum sample, and determining the titer of antibody developed against the immunogen, using methods well known in the art and readily adaptable to the specific antigen to be measured. Ideally, a sample is taken prior to initial dosing and subsequent samples are taken and titered after each immunization. Generally, a dose or dosing schedule which provides a detectable titer at least four times greater than control or "background" levels at a serum dilution of 1:100 is desirable, where background is defined relative to a control serum or relative to a plate background in ELISA assays.

[0095] On any given day that a dosage of immunogen is given, the dosage is greater than 1 μ g/patient and usually greater than 10 μ g/patient if adjuvant is also administered, and greater than 10 μ g/patient and usually greater than 100 μ g/patient in the absence of adjuvant. A typical regimen consists of an immunization followed by booster injections at 6 weekly intervals. Another regimen consists of an immunization followed by booster injections 1, 2 and 12 months later. Another regimen entails an injection every two months for life. Alternatively, booster injections can be on an irregular basis as indicated by monitoring of immune response.

[0096] For passive immunization with a composition comprising an antibody of the present invention, the dosage ranges from about 0.0001 to 100 mg/kg, and more usually 0.01 to 5 mg/kg of the host body weight. An exemplary treatment regime entails administration once per every two

weeks or once a month or once every 3 to 6 months. In some methods, two or more monoclonal antibodies with different binding specificities are administered simultaneously, in which case the dosage of each antibody administered falls within the ranges indicated. Antibody is usually administered on multiple occasions. Intervals between single dosages can be weekly, monthly or yearly. Intervals can also be irregular as indicated by measuring blood levels of antibody to polymerized proteins or peptide in the patient. Alternatively, antibody can be administered as a sustained release formulation, in which case less frequent administration is required. Dosage and frequency vary depending on the half-life of the antibody in the patient. In general, human antibodies show the longest half life, followed by humanized antibodies, chimeric antibodies, and nonhuman antibodies. The dosage and frequency of administration can vary depending on whether the treatment is prophylactic or therapeutic. In prophylactic applications, a relatively low dosage is administered at relatively infrequent intervals over a long period of time. Some patients continue to receive treatment for the rest of their lives. In therapeutic applications, a relatively high dosage at relatively short intervals is sometimes required until progression of the disease is reduced or terminated, and preferably until the patient shows partial or complete amelioration of symptoms of disease. Thereafter, the patient can be administered a prophylactic regime.

[0097] Compositions of the present invention, i.e., polymerized products and antibodies, can be administered by parenteral, topical, intravenous, oral, subcutaneous, intraperitoneal, intranasal or intramuscular means for prophylactic and/or therapeutic treatment. The most typical route of administration for compositions formulated to induce an immune response is subcutaneous although others can be equally effective. The next most common is intramuscular injection. This type of injection is most typically performed in the arm or leg muscles. Intravenous injections as well as intraperitoneal injections, intra-arterial, intracranial, or intradermal injections are also effective in generating an immune response. In some methods, agents such as antibodies are injected directly into a particular tissue where deposits have accumulated, for example intracranial injection. Intramuscular injection or intravenous infusion are preferred for administration of antibody. In some methods, particular therapeutic antibodies are injected directly into the cranium. In some methods, antibodies are administered as a sustained release composition or device, such as a MedipadTM device.

[0098] The pharmaceutical agents of the present invention may be formulated for parenteral administration. Solutions or suspensions of the agent can be prepared in water suitably mixed with a surfactant such as hydroxypropylcellulose. Dispersions can also be prepared in glycerol, liquid polyethylene glycols, and mixtures thereof in oils. Illustrative oils are those of petroleum, animal, vegetable, or synthetic origin, for example, peanut oil, soybean oil, or mineral oil. In general, water, saline, aqueous dextrose and related sugar solution, and glycols, such as propylene glycol or polyethylene glycol, are preferred liquid carriers, particularly for injectable solutions. Under ordinary conditions of storage and use, these preparations contain a preservative to prevent the growth of microorganisms.

[0099] Pharmaceutical formulations suitable for injectable use include sterile aqueous solutions or dispersions and

sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersions. In all cases, the form must be sterile and must be fluid to the extent that easy syringability exists. It must be stable under the conditions of manufacture and storage and must be preserved against the contaminating action of microorganisms, such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (e.g., glycerol, propylene glycol, and liquid polyethylene glycol), suitable mixtures thereof, and vegetable oils.

[0100] When it is desirable to deliver the pharmaceutical agents of the present invention systemically, they may be formulated for parenteral administration by injection, e.g., by bolus injection or continuous infusion. Formulations for injection may be presented in unit dosage form, e.g., in ampoules or in multi-dose containers, with an added preservative. The compositions may take such forms as suspensions, solutions or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents.

[0101] Intraperitoneal or intrathecal administration of the agents of the present invention can also be achieved using infusion pump devices such as those described by Medtronic, Northridge, Calif. Such devices allow continuous infusion of desired compounds avoiding multiple injections and multiple manipulations.

[0102] In addition to the formulations described previously, the compositions of the present invention may also be formulated as a depot preparation. Such long acting formulations may be formulated with suitable polymeric or hydrophobic materials (for example as an emulsion in an acceptable oil) or ion exchange resins, or as sparingly soluble derivatives, for example, as a sparingly soluble salt.

[0103] Another aspect of the present invention relates to a method of diagnosing an amyloid disease in a subject. This method involves detecting, in the subject, the presence of an amyloidogenic proteins or peptides using a diagnostic reagent, where the diagnostic reagent is an antibody, or active binding fragment thereof, of the present invention. As described supra, the antibody has antigenic specificity for a conformational epitope of an amyloidogenic form of a protein. Preferably the conformational epitope recognized by the antibody is shared with one or more other amyloidogenic protein or peptides. The diagnosis of the amyloid disease is based on the detection of an amyloidogenic protein or peptide in the subject.

[0104] Detecting the presence of amyloidogenic deposits in a subject using the diagnostic reagent can be achieved by obtaining a biological sample from the subject (e.g., blood, urine, cerebral spinal fluid), contacting the biological sample with the diagnostic antibody reagent, and detecting binding of the diagnostic antibody reagent to an amyloidogenic protein in the sample from the subject. Assays for carrying out the detection of an amyloid protein in a biological sample using a diagnostic antibody are well known in the art and include, without limitation, ELISA, immunohistochemistry, western blot.

[0105] Alternatively, detecting the presence of amyloid deposits in a subject using diagnostic antibody reagent of the present invention can be achieved using in vivo imaging techniques. In vivo imaging involves administering to the subject the diagnostic antibody having antigenic specificity for a conformational epitope of a polymerized product containing two or more protein and/or peptide units where

each unit is independently selected from the group consisting of A β , α -synuclein, tau, TDP-43, amylin, and/or PrP proteins or peptides, and detecting the binding of the diagnostic agent to the amyloidogenic protein in vivo. As described supra, preferred antibodies bind to a conformational epitope of an amyloidogenic form of a protein or peptide without binding to the non-amyloidogenic proteins and without binding to the non-pathological forms of the amyloidogenic proteins.

[0106] Diagnostic antibodies or similar reagents can be administered by intravenous injection into the body of the patient, or directly into the brain by intracranial injection or by drilling a hole through the skull. The dosage of antibody should be within the same ranges as for treatment methods. Typically, the antibody is labeled, although in some methods, the primary antibody with affinity for the conformational epitope of an amyloid protein is unlabelled and a secondary labeling agent is used to bind to the primary antibody. The choice of label depends on the means of detection. For example, a fluorescent label is suitable for optical detection. Use of paramagnetic labels is suitable for tomographic detection without surgical intervention. Radioactive labels can also be detected using PET or SPECT.

[0107] Diagnosis is performed by comparing the number, size, and/or intensity of labeled amyloid protein deposits in a sample from the subject or in the subject, to corresponding baseline values. An appropriate baseline value can be the average level of amyloid protein deposition in a population of undiseased individuals. Alternatively, an appropriate baseline value may be the level of amyloid protein deposition in the same subject determined at an earlier time.

[0108] The diagnostic methods described above can also be used to monitor a subject's response to therapy. In this embodiment, detection of amyloid deposits in the subject is determined prior to the commencement of treatment. The level of amyloid deposition in the subject at this timepoint is used as a baseline value. At various times during the course of treatment the detection of amyloid deposits can be repeated, and the measured values thereafter compared with the baseline values. A decrease in values relative to baseline signals a positive response to treatment.

[0109] The present invention is further directed to a kit for performing the above described diagnostic and monitoring methods. Typically, such kits contain a diagnostic antibody reagent, preferably the antibody of the present invention that has antigenic specificity for a polymerized A β , α -synuclein, tau, TDP-43, amylin, and/or PrP protein or peptide product. The kit can also include a detectable label. The diagnostic antibody itself may contain the detectable label (e.g., fluorescent molecule, biotin, etc.) which is directly detectable or detectable via a secondary reaction (e.g., reaction with streptavidin). Alternatively, a second reagent containing the detectable label may be utilized, where the second reagent has binding specificity for the primary antibody. In a diagnostic kit suitable for measuring amyloid in a biological sample, the antibodies of the kit may be supplied prebound to a solid phase, such as to the wells of a microtiter dish.

[0110] Diagnostic kits of the present invention also include kits that are useful for detecting antibody production in a subject following administration of a polymerized protein or peptide of the present invention. Typically, such kits include a reagent that contains the antigenic epitope of the antibodies generated by the subject in a polymerized product as described supra. The kit also includes a detectable

label. In a preferred embodiment, the label is typically in the form of labeled anti-idiotypic antibodies. The antigenic epitope reagents of the kit can be supplied prebound to a solid phase, such as to the wells of a microtiter dish.

EXAMPLES

[0111] The following examples are provided to illustrate embodiments of the present invention but they are by no means intended to limit its scope.

Materials and Methods for Examples 1-4

[0112] Peptide Synthesis: The 42 amino acid A β peptide (SEQ ID NO:1), the ABri peptide (Cys-Ser-Arg-Thr-Val-Lys-Lys-Asn-Ile-Ile-Glu-Glu-Asn) (SEQ ID NO:8), and A β 1-30K₁₈K₁₉ (daefrhdsy evhhqklkkf aedvgsnkg) (SEQ ID NO:12) were synthesized on an ABI 430A peptide synthesizer (AME Bioscience, Chicago, Ill.) at the Keck peptide synthesis facility at Yale University, CT, using a Vydac C18 preparative column, 2.5×30 cm (Vydac Separations, Hesperia, Calif.). Standard protocols for tBOC (tert-butyloxycarbonyl) chemistry were used. The peptides were subsequently cleaved from the resins using hydrofluoric acid and purified by high-pressure liquid chromatography (HPLC) on a Vydac C18 preparative column using linear gradients from 0-70% of acetonitrile in 0.1% trifluoroacetic acid. Mass spectroscopy of the lyophilized end-product was used to verify the expected molecular weight.

[0113] Peptide Polymerization and Assessment of Conformation. In order to make the peptides immunogenic and to potentially ensure a conformation specific immune response, the peptides were first subjected to controlled polymerization using the following protocol. Peptides were dissolved at 3 mg/ml, in 100 mM borate buffer saline (BBS), pH 7.4. Fresh 1% glutaraldehyde in BBS was prepared and added to the peptide to a final 5 mM glutaraldehyde concentration and incubated in an Eppendorf block at 800 rpm at 56° C. for 16 hrs. The solution was then quenched with 0.5 M glycine to make the solution 100 mM in glycine. After five minutes the solution was diluted 1:3 with BBS, dialyzed against 2 mM BBS overnight at 4° C., aliquoted, and lyophilized.

[0114] For electron microscopic studies of the polymerized A β peptide, the original and polymerized peptides were incubated at 1 mg/ml in phosphate buffered saline, pH 7.4. The sample (3 μ l) was put onto a carbon coated 400 mesh Cu/Rh grid (Ted Pella Inc., Redding, Calif.) and stained with 1% uranyl acetate in distilled water (Polysciences, Inc., Warrington, Pa.). Stained grids were examined under a Philips CM-12 electron microscope (FEI; Eindhoven, The Netherlands) and photographed with a (1 k×1 k) digital camera (Gatan, Inc., Pleasanton, Calif.).

[0115] Immunization of Mice. Animal studies were approved by the NYU School of Medicine Institutional Animal Care and Use Committee and were consistent with the recommendations of the American Veterinary Association. Two transgenic (Tg) mouse models were used in these experiments. The first model, the 3×Tg model, develops both plaque and tangle pathology (Oddo et al., "Triple-Transgenic Model of Alzheimer's Disease with Plaques and Tangles: Intracellular Abeta and Synaptic Dysfunction," *Neuron* 39:409-21 (2003), which is hereby incorporated by reference in its entirety). The second model, the TgSwDI model, develops extensive congophilic angiopathy (CAA) (Davis et al., "Early-Onset and Robust Cerebral Microvas-

cular Accumulation of Amyloid Beta-Protein in Transgenic Mice Expressing Low Levels of a Vasculotropic Dutch/Iowa Mutant Form of Amyloid Beta-Protein Precursor," *J. Biol. Chem.* 279:20296-306 (2004), which is hereby incorporated by reference in its entirety). Starting at the age of 3 months mice were immunized 4 times biweekly, subcutaneously with 50 μ g/animal of polymerized peptide in sterile saline: Alum 9:1, and thereafter 4 times bimonthly with 25 μ g/animal until the 12 month. At the age of 15-16 months mice were subject to locomotor and cognitive behavioral testing (radial arm maze), followed by histological and biochemical analysis. Animals were bled from the caudal vein prior to inoculation (T0), after the 6th inoculation (T6) and at the time of sacrifice (TF). The blood was collected in heparinized tubes and plasma separated and stored at -80° C.

[0116] Locomotor and Cognitive Behavioral Testing. Locomotor Activity: A Hamilton-Kinder Smart-frame Photobeam System was used to make a computerized recording of animal activity over a designated period of time. Exploratory locomotor activity is recorded in a circular open field activity chamber measuring (70×70 cm). A video camera mounted above the chamber automatically recorded horizontal movements in the open field in each dimension (i.e., x, y, and two z planes). Total distance was measured in centimeters (cm) traveled and is defined as sequential movement interruptions of the animal measured relative to the background. The duration of the behavior was timed for 15 min. Results were reported based on distance traveled (cm), mean resting time, and maximum velocity of the animal.

[0117] Radial Arm Maze: Prior to testing, the mice were adapted to the room with lights on for 15 min. Spatial learning was evaluated using an eight-arm radial maze with a water well at the end of each arm. Clear Plexiglas guillotine doors, operated by a remote pulley system, controlled access to the arms from a central area from which the animals entered and exited the apparatus. After 3-4 days of adaptation, water-restricted mice (2 hours daily access to water) were given one training session per day for ten consecutive days. For each session, all arms were baited with saccharine flavored water, and animals were permitted to enter all arms until the eight rewards had been consumed. The number of errors (entries to previously visited arms) and time to complete each session were recorded.

[0118] Antibody Levels. Antibody levels were determined in duplicate on 1:100 dilutions of plasma using ELISA as described previously (Goni et al., "Mucosal Vaccination Delays or Prevents Prion Infection Via an Oral Route," *Neurosci.* 133:413-421 (2005); Asuni et al., "A β Derivative Vaccination in Alum Adjuvant Prevents Amyloid Deposition and Does Not Cause Brain Microhemorrhages in Alzheimer's Model Mice," *Eur. J. Neurosci.* 24:2530-2542 (2006), which are hereby incorporated by reference in their entirety), in which 5 μ g/plate A β 1-40, A β 1-42, A β 1-30KK, or pABri was coated onto Immulon 2HB 96 well microtiter wells (Thermo, Waltham, Mass.). The bound antibodies were detected by a horseradish peroxidase labeled goat anti-mouse IgG (Amersham Biosciences, Piscataway, N.J.) or a peroxidase conjugated goat anti-mouse IgM (Sigma; A8786). Tetramethyl benzidine (TMB; Pierce, Rockford, Ill.) was the color developing substrate and the readings were done at 450 nm.

[0119] Histology. Mice were anesthetized with sodium pentobarbital (150 mg/kg, i.p.), perfused transaortically with phosphate buffer, and the brains processed as described

previously (Asuni et al., “A β Derivative Vaccination in Alum Adjuvant Prevents Amyloid Deposition and Does Not Cause Brain Microhemorrhages in Alzheimer’s Model Mice,” *Eur. J. Neurosci.* 24:2530-2542 (2006); Sigurdsson et al., “An Attenuated Immune Response is Sufficient to Enhance Cognition in an Alzheimer’s Disease Mouse Model Immunized With Amyloid- β Derivatives,” *J. Neurosci.* 24:6277-6282 (2004), which are hereby incorporated by reference in their entirety). Serial coronal sections (40 μ m) were stained with a mixture of 4G8/6E10, monoclonal antibodies that recognizes A β and stains both pre-amyloid and A β plaques (Sadowski et al., “Blocking the Apolipoprotein/Amyloid β Interaction Reduces the Parenchymal and Vascular Amyloid- β Deposition and Prevents Memory Deficit in AD Transgenic Mice,” *Proc. Natl. Acad. Sci. (USA)* 103:18787-18792 (2006); Scholtzova et al., “Induction of Toll-Like Receptor 9 Signaling as a Method for Ameliorating Alzheimer’s Disease Related Pathology,” *J. Neurosci.* 29:1846-1854 (2009), which are hereby incorporated by reference in their entirety). Immunostaining was performed as described previously (Sadowski et al., “Blocking the Apolipoprotein/Amyloid β Interaction Reduces the Parenchymal and Vascular Amyloid- β Deposition and Prevents Memory Deficit in AD Transgenic Mice,” *Proc. Natl. Acad. Sci. (USA)* 103:18787-18792 (2006); Scholtzova et al., “Induction of Toll-Like Receptor 9 Signaling as a Method for Ameliorating Alzheimer’s Disease Related Pathology,” *J. Neurosci.* 29:1846-1854 (2009), which are hereby incorporated by reference in their entirety). All procedures were performed by an individual blinded to the experimental conditions of the study. The A β burden is defined as the percentage of area in the measurement field occupied by reaction product.

Example 1—Locomotor and Cognitive Testing

[0120] In order to verify that cognitive testing was not confounded by differences in sensorimotor abilities in the polymerized peptide vaccinated versus control Tg3x and SwDI transgenic mice, locomotor testing was conducted first. There was no significant differences between SwDI Tg control mice and SwDI Tg mice vaccinated with polymerized A β 1-30_{K₁₈K₁₉}, polymerized ABri peptide (ABri-Glut), or polymerized A β 1-42 peptide in distance traveled (FIG. 1A) or mean velocity (Vmean) (FIG. 1C). A β 1-42 vaccinated Tg mice exhibited slower maximum velocity (Vmax) (FIG. 1B) ($p=0.0234$ post-hoc only; A β 1-42 versus control $p<0.05$) and longer resting time (FIG. 1D) ($p=0.0311$ post-hoc only; A β 1-42 versus control $p<0.05$) compared to controls. No significant differences between the A β 1-30_{K₁₈K₁₉} and ABri treated Tg animals versus controls in these parameters were observed (FIGS. 1B and 1D). There was no significant difference between Tg3x control mice and Tg3x mice vaccinated with a synthetic polymerized A β 1-30_{K₁₈K₁₉}, polymerized ABri peptide (ABri-Glut), or the combination of A β 1-30_{K₁₈K₁₉} and ABri polymerized peptides in distance traveled (FIG. 2A), maximum velocity (Vmax) (FIG. 2B), mean velocity (Vmean) (FIG. 2C), or in resting time (FIG. 2D).

[0121] Radial arm maze cognitive testing showed there were significant differences between the untreated control Tg3x mice (Tg control) versus ABri treated Tg mice and wild-type controls (FIG. 3A). There was no difference between the wild-type controls and the ABri vaccinated Tg3x mice. There was also a significant difference between

the untreated control Tg3x mice (Tg control) versus animals vaccinated with the polymerized A β 1-30_{K₁₈K₁₉} peptide and animals vaccinated with the combination of polymerized ABri and A β 1-30_{K₁₈K₁₉} peptides (FIG. 3B).

[0122] Radial arm maze cognitive testing in the SwDI Tg mice showed there were significant differences between the untreated control SwDI Tg mice (Tg Control) versus the polymerized A β 1-42, A β 1-30_{K₁₈K₁₉}, and ABri treated Tg mice and wild-type controls (FIG. 3C). There was no difference between the wild-type controls and the vaccinated SwDI Tg mice.

Example 2—Antibody Titers

[0123] In TgSwDI mice vaccinated with polymerized ABri, significant IgG and IgM titers were noted against A β 1-40, A β 1-42, and polymerized ABri at T1 and Tf (FIG. 4A). In TgSwDI mice vaccinated with polymerized A β 1-30_{K₁₈K₁₉}, significant IgG and IgM titers were noted against A β 1-40, A β 1-42, and A β 1-30_{K₁₈K₁₉} at T1 (FIG. 4B). Significant IgM titers against A β 1-40, A β 1-42, and A β 1-30_{K₁₈K₁₉}, and significant IgG titers against A β 1-42 were observed at Tf (FIG. 4B). In TgSwDI mice vaccinated with polymerized A β 1-42, significant IgG and IgM titers were also noted against A β 1-40, A β 1-42, and A β 1-30_{K₁₈K₁₉} at T1, and to a lesser extent at Tf (see FIG. 4C).

[0124] In Tg3x mice vaccinated with polymerized ABri, significant IgG and IgM titers were noted against A β 1-42 and polymerized ABri at T1 and Tf (FIG. 5A). Significant IgM titers against A β 1-40 were observed at T1, and significant IgG titers against A β 1-40 were observed at Tf in these animals. In the polymerized A β 1-30_{K₁₈K₁₉} vaccinated Tg3x mice, significant IgG and IgM titers were noted against A β 1-42 and A β 1-30_{K₁₈K₁₉} at T1 and Tf, and significant IgM and IgG antibodies against A β 1-40 were observed at Tf (FIG. 5B). In Tg3x animals vaccinated with the combination of polymerized A β 1-30_{K₁₈K₁₉} and ABri, significant IgG and IgM titers were also noted against A β 1-42, A β 1-30_{K₁₈K₁₉}, and ABri at T1 and Tf (see FIG. 5C). FIG. 5D shows that IgG and IgM antibody titer against A β 1-40 and A β 1-42 does not differ significantly in vehicle treated transgenic 3xTg animals.

Example 3—Amyloid Burden

[0125] There was a significant reduction in hippocampal amyloid burden in Tg3x animals administered polymerized ABri (Tg-pABri), polymerized A β 1-30_{K₁₈K₁₉} (Tg-AB1-30KK), or the combination of polymerized peptides (Tg-combined) compared to transgenic control (Tg-control) animals as shown in FIG. 6A. There was also a significant reduction in PHF1 in the hippocampus (FIG. 6B) and cortex (FIG. 6C), respectively, of Tg3x animals administered polymerized ABri or the combination of polymerized ABri and A β 1-30_{K₁₈K₁₉}. No significant difference in PHF1 burden in the hippocampus or cortex was found between Tg-control and Tg-A β 1-30_{K₁₈K₁₉} treated animals.

[0126] There was a significant reduction in the hippocampal amyloid burden in SwDI Tg animals vaccinated with polymerized ABri, A β 1-30_{K₁₈K₁₉}, or A β 1-42 compared to control Tg mice (see FIG. 7A).

Example 4—A β 1-42 Peptide Polymerization and Assessment of Conformation

[0127] To examine the stability of polymerized peptide conformation over time, aged A β 1-42 was analyzed by

electron microscopy. As shown in the EM photomicrograph of FIG. 7B the polymerized A β 1-42 peptide is predominately in the form of spherical particles of ~200 nm at 3 weeks post polymerization. Similarly, the EM photomicrograph of FIG. 8A shows that A β 1-42 peptide aged for 3

boosted with a mixture of recombinant polymerized deer PrP protein and polymerized PrP fragments corresponding to amino acid residues 121-166; 13-155; 155-200 and 169-230 of the mouse prion protein of SEQ ID NO: 11 below.

```

1   manlgywlla lfvtmwtdvg lckkrpkpgg wntggs rypg qgspggnryp pqggtwgqph
61  gggwgqphgg swgqphggsw gqphggwgq gggthnqwnk pskpktnlkh vagaaaagav
121 vgglggymlg samsrpmihf gndwedryyr enmyrypnqv yyrpvdqysn qnnfvhdcvn
181 itikqhtvtt ttkgenftet dvkmmervve qmcvtqyqke sqayydgrrs sstvlfsspp
241 villisflif livg

```

months after controlled polymerization with glutaraldehyde maintains the spherical particle form that is typical of oligomerized peptides/proteins without any evidence of fibril formation. In contrast, a non-polymerized A β 1-42 peptide from the same synthesis batch as the peptide in FIG. 8A shows complete fibrillization at 3 months.

Discussion of Examples 1-4

[0128] In summary, controlled polymerized forms of A β 1-42 and A β 1-30K₁₈K₁₉ have been obtained. As demonstrated by EM analysis the A β 1-42 polymerized form does not produce fibrils when aged. The polymerized peptides are stable immunogens that produce an immune response that more specifically targets the oligomeric pathogenic form of A β without targeting the normal, physiological conformers of A β . Vaccination of TgSwDI and Tg3x animals with polymerized ABri, A β 1-42 or A β 1-30K₁₈K₁₉ elicits a good antibody response and produces cognitive benefits. Administration of polymerized A β 1-42 reduces amyloid beta pathology, reduces A β oligomer levels, and diminishes congophilic angiopathy, which is extensive in the SwDI animal model. As demonstrated herein, polymeric synthetic peptides have been designed that mimic the secondary structure found in amyloid and oligomeric forms of A β and pathological forms of tau. The antibody response elicited can target both amyloid and tau pathology resulting in a cognitive benefit, pathology burden reduction, and lack of apparent autoimmune toxicity. Collectively, this data demonstrates these polymeric synthetic peptides are excellent candidates for comprehensive immunomodulation in AD.

Materials and Methods for Example 5

[0129] Five White Tail deer were inoculated (vaccinated group) orally with an attenuated salmonella carrying deer PrP to stimulate the mucosal immune system. At the same time six White Tail deer were inoculated with the same strain of attenuated salmonella but without any insert or foreign protein to be delivered (Control group). The inoculations were repeated 4 times over a period of a few months following the protocol described by Goni et al., “High Titers of Mucosal and Systemic Anti-PrP Antibodies Abrogate Oral Prion Infection in Mucosal-Vaccinated Mice,” *Neurosci.* 153:679-686 (2008), which is hereby incorporated by reference in its entirety.

[0130] Before the first inoculation (T0) and after the fourth inoculation (T5) samples of serum, saliva, and feces were taken and the specific antibodies against PrP were measured by ELISA. Subsequently vaccinated animals were orally

[0131] The PrP peptides and recombinant protein were subject to controlled polymerization as described herein and Goni et al, “Immunomodulation Targeting Abnormal Protein Conformation Reduces Pathology in a Mouse Model of Alzheimer’s Disease,” *PLoS One* 5(10):e13391 (2010), which is hereby incorporated by reference in its entirety. The polymerized PrP fragments and PrP protein were dissolved in sterile saline and delivered with 9:1 sodium hydroxide (Alum).

[0132] Ten days after delivery of the first booster, serum, saliva, and feces samples were taken (T6). Two months later the same boost with polymerized PrP and polymerized PrP fragments was repeated. Ten days after the second booster, serum, saliva, and feces were collected (T7).

[0133] The animals were subsequently boosted again and then orally challenged by delivering Chronic Wasting Disease (CWD) brain homogenate in bait. Samples were taken at different times and survival was assessed to the time the animals showed clear signs of prion disease.

Example 5—Administration of Polymerized PrP Protein/Peptides Protects Against Chronic Wasting Disease in Deer

[0134] FIGS. 9 and 10 show the anti-PrP IgA antibody titers in feces and anti-PrP IgM antibody titers in plasma, respectively, in both control and vaccinated deer over the course of the treatment regimen described above. The control group did not develop any noticeable IgA titer whereas the vaccinated group showed some increase in mucosa titer after the immune response was established with the salmonella oral delivery. The IgA titers were initially very low (T5) but were greatly enhanced after the animals were boosted with the polymerized PrP and PrP fragments (T6 and T7) showing the importance of these antigenic preparation on generating a sustainable immune response (FIG. 9). With regard to IgM antibody titers, the control group did not show any noticeable increase in antibody titer; whereas the vaccinated group showed some concomitant IgM titer in serum at the same time of the mucosal response (T5). Anti-PrP antibody titers in the vaccinated animals greatly increased after the two boosts with the polymerized PrP and PrP fragments showing that this type of boost could invoke a therapeutic serum response.

[0135] An analysis of the antibody specificity in the vaccinated deer found that the antibodies produced are specific to prion protein. FIG. 10 contains immunoblot data using purified antibodies from T7 in vaccinated animal 781 and control animal 786. Lane 1 of the blot contains salmonella lysate, to which antibodies in both animals were highly

reactive to. Lanes 4 and 5 contain deer PrP and polymerized deer PrP. The bands appearing in both lanes of the blot incubated with antibodies from the vaccinated animal but not in the lanes of the blot incubated with antibodies from the control animal indicate the generation and presence of deer PrP antibodies in the vaccinated, but not control animals. Lanes 2 and 3 contain sheep PrP and polymerized sheep PrP. The absence of bands in these lanes indicates the specificity of the antibody response observed in the vaccinated animals, i.e., the generated antibodies are specific for deer PrP.

[0136] PrP vaccination was clearly protective as indicated by the Kaplan Meier survival curve of FIG. 12. Both

vaccinated and control animals were challenged with exposure to CWD. Three out of the six control animals became with sick prionoses CWD and had to be properly euthanized. In contrast, none of the vaccinated animals had any signs of disease over the course of 18 months after challenge.

[0137] Although preferred embodiments have been depicted and described in detail herein, it will be apparent to those skilled in the relevant art that various modifications, additions, substitutions, and the like can be made without departing from the spirit of the invention and these are therefore considered to be within the scope of the invention as defined in the claims which follow.

SEQUENCE LISTING	
<160> NUMBER OF SEQ ID NOS: 14	
<210> SEQ ID NO 1	
<211> LENGTH: 42	
<212> TYPE: PRT	
<213> ORGANISM: Artificial	
<220> FEATURE:	
<223> OTHER INFORMATION: A-beta peptide	
<400> SEQUENCE: 1	
Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys	
1 5 10 15	
Leu Val Phe Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile	
20 25 30	
Gly Leu Met Val Gly Gly Val Val Ile Ala	
35 40	
<210> SEQ ID NO 2	
<211> LENGTH: 42	
<212> TYPE: PRT	
<213> ORGANISM: Artificial	
<220> FEATURE:	
<223> OTHER INFORMATION: A-beta peptide	
<400> SEQUENCE: 2	
Asp Ala Glu Phe Arg His Asp Ser Gly Tyr Glu Val His His Gln Lys	
1 5 10 15	
Leu Lys Lys Phe Ala Glu Asp Val Gly Ser Asn Lys Gly Ala Ile Ile	
20 25 30	
Gly Leu Met Val Gly Gly Val Val Ile Ala	
35 40	
<210> SEQ ID NO 3	
<211> LENGTH: 140	
<212> TYPE: PRT	
<213> ORGANISM: Homo sapiens	
<400> SEQUENCE: 3	
Met Asp Val Phe Met Lys Gly Leu Ser Lys Ala Lys Glu Gly Val Val	
1 5 10 15	
Ala Ala Ala Glu Lys Thr Lys Gln Gly Val Ala Glu Ala Ala Gly Lys	
20 25 30	
Thr Lys Glu Gly Val Leu Tyr Val Gly Ser Lys Thr Lys Glu Gly Val	
35 40 45	
Val His Gly Val Ala Thr Val Ala Glu Lys Thr Lys Glu Gln Val Thr	
50 55 60	

-continued

Asn	Val	Gly	Gly	Ala	Val	Val	Thr	Gly	Val	Thr	Ala	Val	Ala	Gln	Lys
65					70					75					80
Thr	Val	Glu	Gly	Ala	Gly	Ser	Ile	Ala	Ala	Ala	Thr	Gly	Phe	Val	Lys
				85					90					95	
Lys	Asp	Gln	Leu	Gly	Lys	Asn	Glu	Glu	Gly	Ala	Pro	Gln	Glu	Gly	Ile
			100					105					110		
Leu	Glu	Asp	Met	Pro	Val	Asp	Pro	Asp	Asn	Glu	Ala	Tyr	Glu	Met	Pro
		115					120					125			
Ser	Glu	Glu	Gly	Tyr	Gln	Asp	Tyr	Glu	Pro	Glu	Ala				
	130					135					140				
<210> SEQ ID NO 4															
<211> LENGTH: 441															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 4															
Met	Ala	Glu	Pro	Arg	Gln	Glu	Phe	Glu	Val	Met	Glu	Asp	His	Ala	Gly
1				5					10					15	
Thr	Tyr	Gly	Leu	Gly	Asp	Arg	Lys	Asp	Gln	Gly	Gly	Tyr	Thr	Met	His
			20					25					30		
Gln	Asp	Gln	Glu	Gly	Asp	Thr	Asp	Ala	Gly	Leu	Lys	Glu	Ser	Pro	Leu
		35					40					45			
Gln	Thr	Pro	Thr	Glu	Asp	Gly	Ser	Glu	Glu	Pro	Gly	Ser	Glu	Thr	Ser
	50					55					60				
Asp	Ala	Lys	Ser	Thr	Pro	Thr	Ala	Glu	Asp	Val	Thr	Ala	Pro	Leu	Val
65					70					75					80
Asp	Glu	Gly	Ala	Pro	Gly	Lys	Gln	Ala	Ala	Ala	Gln	Pro	His	Thr	Glu
			85						90					95	
Ile	Pro	Glu	Gly	Thr	Thr	Ala	Glu	Glu	Ala	Gly	Ile	Gly	Asp	Thr	Pro
			100					105					110		
Ser	Leu	Glu	Asp	Glu	Ala	Ala	Gly	His	Val	Thr	Gln	Ala	Arg	Met	Val
		115					120					125			
Ser	Lys	Ser	Lys	Asp	Gly	Thr	Gly	Ser	Asp	Asp	Lys	Lys	Ala	Lys	Gly
	130					135					140				
Ala	Asp	Gly	Lys	Thr	Lys	Ile	Ala	Thr	Pro	Arg	Gly	Ala	Ala	Pro	Pro
145					150					155					160
Gly	Gln	Lys	Gly	Gln	Ala	Asn	Ala	Thr	Arg	Ile	Pro	Ala	Lys	Thr	Pro
				165					170					175	
Pro	Ala	Pro	Lys	Thr	Pro	Pro	Ser	Ser	Gly	Glu	Pro	Pro	Lys	Ser	Gly
			180					185					190		
Asp	Arg	Ser	Gly	Tyr	Ser	Ser	Pro	Gly	Ser	Pro	Gly	Thr	Pro	Gly	Ser
		195					200					205			
Arg	Ser	Arg	Thr	Pro	Ser	Leu	Pro	Thr	Pro	Pro	Thr	Arg	Glu	Pro	Lys
	210					215					220				
Lys	Val	Ala	Val	Val	Arg	Thr	Pro	Pro	Lys	Ser	Pro	Ser	Ser	Ala	Lys
225					230					235					240
Ser	Arg	Leu	Gln	Thr	Ala	Pro	Val	Pro	Met	Pro	Asp	Leu	Lys	Asn	Val
			245						250					255	
Lys	Ser	Lys	Ile	Gly	Ser	Thr	Glu	Asn	Leu	Lys	His	Gln	Pro	Gly	Gly
			260					265					270		
Gly	Lys	Val	Gln	Ile	Ile	Asn	Lys	Lys	Leu	Asp	Leu	Ser	Asn	Val	Gln

-continued

275					280					285						
Ser	Lys	Cys	Gly	Ser	Lys	Asp	Asn	Ile	Lys	His	Val	Pro	Gly	Gly	Gly	
290					295					300						
Ser	Val	Gln	Ile	Val	Tyr	Lys	Pro	Val	Asp	Leu	Ser	Lys	Val	Thr	Ser	
305					310					315					320	
Lys	Cys	Gly	Ser	Leu	Gly	Asn	Ile	His	His	Lys	Pro	Gly	Gly	Gly	Gln	
					325					330					335	
Val	Glu	Val	Lys	Ser	Glu	Lys	Leu	Asp	Phe	Lys	Asp	Arg	Val	Gln	Ser	
					340					345					350	
Lys	Ile	Gly	Ser	Leu	Asp	Asn	Ile	Thr	His	Val	Pro	Gly	Gly	Gly	Asn	
					355					360					365	
Lys	Lys	Ile	Glu	Thr	His	Lys	Leu	Thr	Phe	Arg	Glu	Asn	Ala	Lys	Ala	
370					375					380						
Lys	Thr	Asp	His	Gly	Ala	Glu	Ile	Val	Tyr	Lys	Ser	Pro	Val	Val	Ser	
385					390					395					400	
Gly	Asp	Thr	Ser	Pro	Arg	His	Leu	Ser	Asn	Val	Ser	Ser	Thr	Gly	Ser	
					405					410					415	
Ile	Asp	Met	Val	Asp	Ser	Pro	Gln	Leu	Ala	Thr	Leu	Ala	Asp	Glu	Val	
					420					425					430	
Ser	Ala	Ser	Leu	Ala	Lys	Gln	Gly	Leu								
435					440											
<210> SEQ ID NO 5																
<211> LENGTH: 414																
<212> TYPE: PRT																
<213> ORGANISM: Homo sapiens																
<400> SEQUENCE: 5																
Met	Ser	Glu	Tyr	Ile	Arg	Val	Thr	Glu	Asp	Glu	Asn	Asp	Glu	Pro	Ile	
1	5					10					15					
Glu	Ile	Pro	Ser	Glu	Asp	Asp	Gly	Thr	Val	Leu	Leu	Ser	Thr	Val	Thr	
20					25					30						
Ala	Gln	Phe	Pro	Gly	Ala	Cys	Gly	Leu	Arg	Tyr	Arg	Asn	Pro	Val	Ser	
35					40					45						
Gln	Cys	Met	Arg	Gly	Val	Arg	Leu	Val	Glu	Gly	Ile	Leu	His	Ala	Pro	
50					55					60						
Asp	Ala	Gly	Trp	Gly	Asn	Leu	Val	Tyr	Val	Val	Asn	Tyr	Pro	Lys	Asp	
65					70					75					80	
Asn	Lys	Arg	Lys	Met	Asp	Glu	Thr	Asp	Ala	Ser	Ser	Ala	Val	Lys	Val	
					85					90					95	
Lys	Arg	Ala	Val	Gln	Lys	Thr	Ser	Asp	Leu	Ile	Val	Leu	Gly	Leu	Pro	
100					105					110						
Trp	Lys	Thr	Thr	Glu	Gln	Asp	Leu	Lys	Glu	Tyr	Phe	Ser	Thr	Phe	Gly	
115					120					125						
Glu	Val	Leu	Met	Val	Gln	Val	Lys	Lys	Asp	Leu	Lys	Thr	Gly	His	Ser	
130					135					140						
Lys	Gly	Phe	Gly	Phe	Val	Arg	Phe	Thr	Glu	Tyr	Glu	Thr	Gln	Val	Lys	
145					150					155					160	
Val	Met	Ser	Gln	Arg	His	Met	Ile	Asp	Gly	Arg	Trp	Cys	Asp	Cys	Lys	
					165					170					175	
Leu	Pro	Asn	Ser	Lys	Gln	Ser	Gln	Asp	Glu	Pro	Leu	Arg	Ser	Arg	Lys	
180					185					190						

-continued

Val	Phe	Val	Gly	Arg	Cys	Thr	Glu	Asp	Met	Thr	Glu	Asp	Glu	Leu	Arg
		195					200					205			
Glu	Phe	Phe	Ser	Gln	Tyr	Gly	Asp	Val	Met	Asp	Val	Phe	Ile	Pro	Lys
		210				215					220				
Pro	Phe	Arg	Ala	Phe	Ala	Phe	Val	Thr	Phe	Ala	Asp	Asp	Gln	Ile	Ala
225					230					235					240
Gln	Ser	Leu	Cys	Gly	Glu	Asp	Leu	Ile	Ile	Lys	Gly	Ile	Ser	Val	His
				245					250					255	
Ile	Ser	Asn	Ala	Glu	Pro	Lys	His	Asn	Ser	Asn	Arg	Gln	Leu	Glu	Arg
			260					265					270		
Ser	Gly	Arg	Phe	Gly	Gly	Asn	Pro	Gly	Gly	Phe	Gly	Asn	Gln	Gly	Gly
		275					280					285			
Phe	Gly	Asn	Ser	Arg	Gly	Gly	Gly	Ala	Gly	Leu	Gly	Asn	Asn	Gln	Gly
	290					295					300				
Ser	Asn	Met	Gly	Gly	Gly	Met	Asn	Phe	Gly	Ala	Phe	Ser	Ile	Asn	Pro
305					310					315					320
Ala	Met	Met	Ala	Ala	Ala	Gln	Ala	Ala	Leu	Gln	Ser	Ser	Trp	Gly	Met
				325					330					335	
Met	Gly	Met	Leu	Ala	Ser	Gln	Gln	Asn	Gln	Ser	Gly	Pro	Ser	Gly	Asn
			340					345					350		
Asn	Gln	Asn	Gln	Gly	Asn	Met	Gln	Arg	Glu	Pro	Asn	Gln	Ala	Phe	Gly
		355					360					365			
Ser	Gly	Asn	Asn	Ser	Tyr	Ser	Gly	Ser	Asn	Ser	Gly	Ala	Ala	Ile	Gly
	370					375					380				
Trp	Gly	Ser	Ala	Ser	Asn	Ala	Gly	Ser	Gly	Ser	Gly	Phe	Asn	Gly	Gly
385					390				395						400
Phe	Gly	Ser	Ser	Met	Asp	Ser	Lys	Ser	Ser	Gly	Trp	Gly	Met		
				405					410						
<210> SEQ ID NO 6															
<211> LENGTH: 62															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 6															
His	Gln	Val	Glu	Lys	Arg	Lys	Cys	Asn	Thr	Ala	Thr	Cys	Ala	Thr	Gln
1				5					10					15	
Arg	Leu	Ala	Asn	Phe	Leu	Val	His	Ser	Ser	Asn	Asn	Phe	Gly	Ala	Ile
			20					25					30		
Leu	Ser	Ser	Thr	Asn	Val	Gly	Ser	Asn	Thr	Tyr	Gly	Lys	Arg	Asn	Ala
		35					40					45			
Val	Glu	Val	Leu	Lys	Arg	Glu	Pro	Leu	Asn	Tyr	Leu	Pro	Leu		
	50					55					60				
<210> SEQ ID NO 7															
<211> LENGTH: 253															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 7															
Met	Ala	Asn	Leu	Gly	Cys	Trp	Met	Leu	Val	Leu	Phe	Val	Ala	Thr	Trp
1			5						10					15	
Ser	Asp	Leu	Gly	Leu	Cys	Lys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	Trp	Asn
		20						25					30		

-continued

Thr	Gly	Gly	Ser	Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	Gly	Asn	Arg
		35					40					45			
Tyr	Pro	Pro	Gln	Gly	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly
	50					55				60					
Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly
65					70					75					80
Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	Gly	Thr	His
				85					90					95	
Ser	Gln	Trp	Asn	Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	Met	Lys	His	Met
			100					105					110		
Ala	Gly	Ala	Ala	Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	Leu	Gly	Gly	Tyr
		115					120					125			
Val	Leu	Gly	Ser	Ala	Met	Ser	Arg	Pro	Ile	Ile	His	Phe	Gly	Ser	Asp
	130					135					140				
Tyr	Glu	Asp	Arg	Tyr	Tyr	Arg	Glu	Asn	Met	His	Arg	Tyr	Pro	Asn	Gln
145					150					155					160
Val	Tyr	Tyr	Arg	Pro	Met	Asp	Glu	Tyr	Ser	Asn	Gln	Asn	Asn	Phe	Val
				165					170					175	
His	Asp	Cys	Val	Asn	Ile	Thr	Ile	Lys	Gln	His	Thr	Val	Thr	Thr	Thr
			180					185					190		
Thr	Lys	Gly	Glu	Asn	Phe	Thr	Glu	Thr	Asp	Val	Lys	Met	Met	Glu	Arg
		195					200					205			
Val	Val	Glu	Gln	Met	Cys	Ile	Thr	Gln	Tyr	Glu	Arg	Glu	Ser	Gln	Ala
	210					215					220				
Tyr	Tyr	Lys	Arg	Gly	Ser	Ser	Met	Val	Leu	Phe	Ser	Ser	Pro	Pro	Val
225					230					235					240
Ile	Leu	Leu	Ile	Ser	Phe	Leu	Ile	Phe	Leu	Ile	Val	Gly			
				245					250						

<210> SEQ ID NO 8
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: ABri peptide

<400> SEQUENCE: 8

Cys	Ser	Arg	Thr	Val	Lys	Lys	Asn	Ile	Ile	Glu	Glu	Asn
1				5					10			

<210> SEQ ID NO 9
<211> LENGTH: 13
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: ADan peptide

<400> SEQUENCE: 9

Cys	Phe	Asn	Leu	Phe	Leu	Asn	Ser	Gln	Glu	Lys	His	Tyr
1				5					10			

<210> SEQ ID NO 10
<211> LENGTH: 31
<212> TYPE: PRT
<213> ORGANISM: Artificial
<220> FEATURE:
<223> OTHER INFORMATION: ABri/ADan fusion peptide

-continued

<400> SEQUENCE: 10															
Cys	Ser	Arg	Thr	Val	Lys	Lys	Asn	Ile	Ile	Glu	Glu	Asn	Gly	Ser	Gly
1				5					10					15	
Ser	Gly	Cys	Phe	Asn	Leu	Phe	Leu	Asn	Ser	Gln	Glu	Lys	His	Tyr	
			20					25					30		
<210> SEQ ID NO 11															
<211> LENGTH: 254															
<212> TYPE: PRT															
<213> ORGANISM: Mouse															
<400> SEQUENCE: 11															
Met	Ala	Asn	Leu	Gly	Tyr	Trp	Leu	Leu	Ala	Leu	Phe	Val	Thr	Met	Trp
1				5					10					15	
Thr	Asp	Val	Gly	Leu	Cys	Lys	Lys	Arg	Pro	Lys	Pro	Gly	Gly	Trp	Asn
			20					25					30		
Thr	Gly	Gly	Ser	Arg	Tyr	Pro	Gly	Gln	Gly	Ser	Pro	Gly	Gly	Asn	Arg
		35					40					45			
Tyr	Pro	Pro	Gln	Gly	Gly	Thr	Trp	Gly	Gln	Pro	His	Gly	Gly	Gly	Trp
	50					55					60				
Gly	Gln	Pro	His	Gly	Gly	Ser	Trp	Gly	Gln	Pro	His	Gly	Gly	Ser	Trp
65				70					75						80
Gly	Gln	Pro	His	Gly	Gly	Gly	Trp	Gly	Gln	Gly	Gly	Gly	Thr	His	Asn
			85					90						95	
Gln	Trp	Asn	Lys	Pro	Ser	Lys	Pro	Lys	Thr	Asn	Leu	Lys	His	Val	Ala
			100					105						110	
Gly	Ala	Ala	Ala	Ala	Gly	Ala	Val	Val	Gly	Gly	Leu	Gly	Gly	Tyr	Met
			115				120					125			
Leu	Gly	Ser	Ala	Met	Ser	Arg	Pro	Met	Ile	His	Phe	Gly	Asn	Asp	Trp
	130					135					140				
Glu	Asp	Arg	Tyr	Tyr	Arg	Glu	Asn	Met	Tyr	Arg	Tyr	Pro	Asn	Gln	Val
145					150					155					160
Tyr	Tyr	Arg	Pro	Val	Asp	Gln	Tyr	Ser	Asn	Gln	Asn	Asn	Phe	Val	His
				165					170					175	
Asp	Cys	Val	Asn	Ile	Thr	Ile	Lys	Gln	His	Thr	Val	Thr	Thr	Thr	Thr
			180					185					190		
Lys	Gly	Glu	Asn	Phe	Thr	Glu	Thr	Asp	Val	Lys	Met	Met	Glu	Arg	Val
		195					200					205			
Val	Glu	Gln	Met	Cys	Val	Thr	Gln	Tyr	Gln	Lys	Glu	Ser	Gln	Ala	Tyr
	210					215					220				
Tyr	Asp	Gly	Arg	Arg	Ser	Ser	Ser	Thr	Val	Leu	Phe	Ser	Ser	Pro	Pro
225					230					235					240
Val	Ile	Leu	Leu	Ile	Ser	Phe	Leu	Ile	Phe	Leu	Ile	Val	Gly		
			245					250							
<210> SEQ ID NO 12															
<211> LENGTH: 30															
<212> TYPE: PRT															
<213> ORGANISM: Artificial															
<220> FEATURE:															
<223> OTHER INFORMATION: A-beta1-30k18k19															
<400> SEQUENCE: 12															
Asp	Ala	Glu	Phe	Arg	His	Asp	Ser	Gly	Tyr	Glu	Val	His	His	Gln	Lys
1				5					10					15	

-continued

Leu	Lys	Lys	Phe	Ala	Glu	Asp	Val	Gly	Ser	Asn	Lys	Gly	Ala
			20					25				30	
<210> SEQ ID NO 13													
<211> LENGTH: 4													
<212> TYPE: PRT													
<213> ORGANISM: Artificial													
<220> FEATURE:													
<223> OTHER INFORMATION: Glycine-rich linker													
<400> SEQUENCE: 13													
Gly Ser Gly Ser													
1													
<210> SEQ ID NO 14													
<211> LENGTH: 5													
<212> TYPE: PRT													
<213> ORGANISM: Artificial													
<220> FEATURE:													
<223> OTHER INFORMATION: Glycine-rich linker													
<400> SEQUENCE: 14													
Gly Ser Gly Ser Gly													
1 5													

1. A non-amyloidogenic, non-fibrillogenic polymerized product comprising two or more protein and/or peptide units, wherein each unit is independently selected from the group consisting of an amyloid-beta (A β) peptide, an α -synuclein protein or peptide, a tau protein or peptide, a TAR DNA-binding protein 43 (TDP-43) protein or peptide, an amylin protein or peptide, a prion protein (PrP) protein or peptide, and any combination thereof.
2. The polymerized product of claim 1, wherein the A β peptide comprises an amino acid sequence selected from the group consisting of amino acid residues 1-16 of SEQ ID NO:1, amino acid residues 1-20 of SEQ ID NO: 1, amino acid residues 1-30 of SEQ ID NO:1, amino acid residues 1-40 of SEQ ID NO:1, amino acid residues 1-42 of SEQ ID NO:1, amino acid residues 10-30 of SEQ ID NO:1, amino acid residues 20-40 of SEQ ID NO:1, and amino acid residues 20-42 of SEQ ID NO:1.
3. The polymerized product of claim 2, wherein the A β peptide comprises amino acid residues 1-42 of SEQ ID NO:1.
4. The polymerized product of claim 1, wherein the A β peptide comprises an amino acid sequence selected from the group consisting of amino acid residues 1-30 of SEQ ID NO:2 (A β 1-30K₁₈K₁₉), amino acid residues 1-40 of SEQ ID NO:2 (A β 1-40K₁₈K₁₉), amino acid residues 1-42 of SEQ ID NO:2 (A β 1-42K₁₈K₁₉), amino acid residues 1-20 of SEQ ID NO:2 (A β 1-20K₁₈K₁₉), amino acid residues 10-30 of SEQ ID NO:2 (A β 10-30K₁₈K₁₉), amino acid residues 10-40 of SEQ ID NO:2 (A β 10-40K₁₈K₁₉), amino acid residues 10-42 of SEQ ID NO:2 (A β 10-42K₁₈K₁₉), amino acid residues 20-40 of SEQ ID NO:2 (A β 20-40K₁₈K₁₉), and amino acid residues 20-42 of SEQ ID NO:2 (A β 20-42K₁₈K₁₉).
5. The polymerized product of claim 4, wherein the A β peptide comprises amino acid residues 1-30 of SEQ ID NO:2 (A β 1-30K₁₈K₁₉).
6. The polymerized product of claim 1, wherein the α -synuclein protein or peptide comprises an amino acid

- sequence of SEQ ID NO:3 or a peptide derived from the amino acid sequence of SEQ ID NO:3.
7. The polymerized product of claim 1, wherein the tau protein or peptide comprises an amino acid sequence of SEQ ID NO: 4 or a peptide derived from the amino acid sequence of SEQ ID NO:4.
8. The polymerized product of claim 1, wherein the TDP-43 protein or peptide comprises an amino acid sequence of SEQ ID NO:5 or a peptide derived from the amino acid sequence of SEQ ID NO:5.
9. polymerized product of claim 1, wherein the amylin protein or peptide comprises an amino acid sequence of SEQ ID NO:6 or a peptide derived from the amino acid sequence of SEQ ID NO:6.
10. The polymerized product of claim 1, wherein the PrP protein or peptide comprising an amino acid sequence of SEQ ID NO:7 or a peptide derived from the amino acid sequence of SEQ ID NO:7.
11. The polymerized product of claim 1, wherein the polymerized product is a co-polymer comprising one or more amyloid-beta (A β) peptides copolymerized with one or more α -synuclein proteins or peptides.
12. The polymerized product of claim 1, wherein the polymerized product is a co-polymer comprising one or more amyloid-beta (A β) peptides co-polymerized with one or more tau peptides.
13. The polymerized product of claim 1, wherein the polymerized product is a co-polymer comprising one or more TDP-43 peptides co-polymerized with one or more amyloid-beta (A β) peptides and/or one or more tau peptides.
14. The polymerized product of claim 1, wherein an adjuvant polypeptide is linked in frame to one or more of the proteins or peptides of the polymerized product.
15. The polymerized product of claim 14, wherein the adjuvant polypeptide is selected from the group consisting of cholera toxin B, flagellin, human papillomavirus L1 or L2

protein, herpes simplex glycoprotein D (gD), complement C4 binding protein, TL4 ligand, and IL-1 β .

16. (canceled)

17. The polymerized product of claim **1**, wherein an immunogenic carrier molecule is conjugated to the one or more proteins or peptides of the polymerized product.

18. The polymerized product of claim **17**, wherein the immunogenic carrier molecule is covalently or non-covalently bonded to the one or more proteins or peptides of the polymerized product.

19. The polymerized product of claim **17**, wherein the immunogenic carrier molecule is selected from the group consisting of bovine serum albumin, chicken egg ovalbumin, keyhole limpet hemocyanin, tetanus toxoid, diphtheria toxoid, thyroglobulin, a pneumococcal capsular polysaccharide, CRM 197, and a meningococcal outer membrane protein.

20. A pharmaceutical composition comprising:
the polymerized product of claim **1** and
a pharmaceutically acceptable carrier.

21. The pharmaceutical composition of claim **20** further comprising:
an adjuvant.

* * * * *