

(19) United States

(12) Patent Application Publication

Samulski

(10) Pub. No.: US 2023/0250452 A1

(43) Pub. Date: Aug. 10, 2023

(54) METHODS AND COMPOSITIONS FOR DUAL GLYCAN BINDING AAV2.5 VECTOR

(71) Applicant: The University of North Carolina at Chapel Hill, Chapel Hill, NC (US)

(72) Inventor: Richard Jude Samulski, Hillsborough, NC (US)

(21) Appl. No.: 18/186,746

(22) Filed: Mar. 20, 2023

A61P 31/20 (2006.01)

A61P 1/16 (2006.01)

C07K 14/005 (2006.01)

C12N 7/00 (2006.01)

(52) U.S. Cl.

CPC C12N 15/86 (2013.01); A61P 31/18 (2018.01); A61P 31/20 (2018.01); A61P 1/16 (2018.01); C07K 14/005 (2013.01); C12N 7/00 (2013.01); C12N 2750/14142 (2013.01); C12N 2750/14143 (2013.01); C12N 2750/14122 (2013.01); C12N 2750/14171 (2013.01)

Related U.S. Application Data

(60) Division of application No. 17/581,639, filed on Jan. 21, 2022, now Pat. No. 11,639,509, which is a continuation of application No. PCT/US2021/056582, filed on Oct. 26, 2021.

(60) Provisional application No. 63/106,733, filed on Oct. 28, 2020.

Publication Classification

(51) Int. Cl.

C12N 15/86 (2006.01)

A61P 31/18 (2006.01)

(57) ABSTRACT

Disclosed herein are methods and compositions comprising an adeno-associated virus 2.5 (AAV2.5) capsid protein, comprising one or more amino acids substitutions, (e.g., which does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5) wherein the substitutions introduce a new glycan binding site into the AAV capsid protein.

Specification includes a Sequence Listing.

AAV9

#1

#2

2.8x10⁸ p/s

6.6x10⁷ p/s

AAV2.5-G9

#3

#4

7.9x10⁸ p/s

8.6x10⁹ p/s

AAV2.5-G9

#5

#6

#7

#8

#9

3.4x10⁸ p/s

5.3x10⁸ p/s

5.3x10⁹ p/s

3.1x10⁹ p/s

1.4x10⁷ p/s

FIG. 1





FIG. 2



METHODS AND COMPOSITIONS FOR DUAL GLYCAN BINDING AAV2.5 VECTOR

STATEMENT OF PRIORITY

[0001] This patent application is a divisional of U.S. patent application Ser. No. 17/581,639, filed on Jan. 21, 2022, which is a continuation under 35 U.S.C. § 111(a) of PCT Application No. PCT/US2021/056582, filed on Oct. 26, 2021, which claims the benefit, under 35 U.S.C. § 119(e), of U.S. Provisional Application No. 63/106,733, filed on Oct. 28, 2020, the entire contents of each of which are incorporated by reference herein.

STATEMENT OF GOVERNMENT SUPPORT

[0002] This invention was made with government support under Grant Numbers HL085794 and OD011107 awarded by the National Institutes of Health. The government has certain rights in the invention.

STATEMENT REGARDING ELECTRONIC FILING OF A SEQUENCE LISTING

[0003] A Sequence Listing in XML format, entitled 5470-891CTDV_ST26.xml, 4,941 bytes in size, generated on Mar. 20, 2023, and filed herewith, is hereby incorporated by reference in its entirety for its disclosures.

FIELD OF THE INVENTION

[0004] The present invention relates to modified capsid proteins from adeno-associated virus (AAV), virus capsids and virus vectors comprising the same, as well as methods of their use.

BACKGROUND OF THE INVENTION

[0005] Inborn errors of metabolism are responsible for a group of diverse congenital diseases where a single gene produces a dysfunctional enzyme necessary for the development and maintenance of neurological function as well as neuronal survival. Unfortunately, many of these are quickly progressing neurodegenerative disorders that result in premature death in the first or second decade of life. Loss-of-function mutations in these enzymatic genes typically follow autosomal recessive or X-linked modes of inheritance, which makes them attractive for simple gene replacement strategies. A major challenge to developing gene therapies for this group of diseases is that successful therapeutic intervention must occur very early in development because damage is potentially irreversible. Over the last two decades studies have consistently demonstrated that the greatest opportunity for success exists when intervening prior to initiation of the neurodegenerative process. For example, mucopolysaccharidosis (MPS) and leukodystrophies are rare pediatric neurodegenerative disorders that result from impaired metabolism of carbohydrate molecules or fatty acids, but are treatable by cellular therapies and enzyme replacement strategies. Several studies have demonstrated that intervention is unable to slow or reverse disease progression after behavioral or physical symptoms manifest. However, children who were asymptomatic at the time of treatment benefited most from the intervention and had the greatest odds of survival.

[0006] Clinically, onset of MPS and leukodystrophies is typically observed in infancy and early childhood with

progressive central and peripheral nervous system involvement. Primary symptoms include the loss of fine and gross motor movements, sensory impairment, distal muscle weakness, and tendon contractures. Affected patients frequently become wheelchair-dependent, hearing and vision impaired, and these diseases are often fatal by 2-8 years of life. Genes known to be involved in leukodystrophies contain mutations in arylsulfatase A (ASA), responsible for metachromatic leukodystrophy; galactosylceramidase (GALC), responsible for Krabbe disease; aspartoacylase (ASPA), responsible for Canavan disease; and a peroxisomal ATP-binding cassette (ABCD1), responsible for X-linked adrenoleukodystrophy. Adeno-associated viral (AAV) vectors provide an attractive option for efficient, targeted gene therapy because they are nonpathogenic with a strong safety profile in humans. Naturally occurring AAV serotypes have shown tropism for multiple tissues, and thus there is a need in the field for the development of methods to target AAV vectors to specifically desired target tissues with minimal off-target expression.

[0007] Virus-glycan interactions are critical determinants of host cell invasion. Cell surface carbohydrates such as sialic acids, gangliosides or heparan sulfate are exploited by a vast number of viruses such as influenza, herpesvirus, SV40, polyomavirus, papillomavirus and other pathogens. In most cases, a single class of glycans primarily serves as the cell surface attachment factor for viruses, leading to sequential or parallel engagement of other receptors/co-receptors for cell entry. Adeno-associated viruses (AAV) are helper-dependent parvoviruses that exploit heparan sulfate (HS), galactose (Gal) or sialic acids (Sia) as primary receptors for cell surface binding. For instance, AAV serotypes 2 and 3b utilize HS; AAV1, 4 and 5 bind Sia with different linkage specificities; while AAV9 exploits Gal for host cell attachment. Different AAV strains also require subsequent interaction with co-receptors such as integrin $\alpha V\beta 5$ or $\alpha 5\beta 1$, fibroblast growth factor receptor (FGFR), platelet-derived growth factor receptor (PDGFR), epidermal growth factor receptor (EGFR), hepatocyte growth factor receptor (HGFR), or the laminin receptor for cellular uptake.

[0008] A notable exception to the monogamous relationship between a specific AAV strain and a single class of carbohydrates is AAV serotype 6, which recognizes both Sia and HS. However, only Sia has been shown essential for viral transduction. The Sia binding footprints for AAV1, AAV4, AAV5 and AAV6 remain to be determined. More recently, key amino acid residues involved in Gal recognition by AAV9 capsids were identified by using a combination of molecular docking and site-directed mutagenesis. What is needed are virus vectors that have multiple glycan binding capability to exploit alternative pathways for cell entry and transduction.

[0009] The present invention overcomes previous shortcomings in the art by providing modified capsid proteins with multiple glycan binding sites, AAV vectors comprising these capsid proteins and methods for their use as therapeutic vectors in disorders such as neurodegenerative leukodystrophies and MPS.

SUMMARY OF THE INVENTION

[0010] Aspects of the invention relate to an adeno-associated virus (AAV) capsid protein that comprises an AAV2.5 capsid protein comprising one or more amino acid substitutions that introduce a new glycan binding site (e.g., which

does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5). In one embodiment, the one or more amino acid substitutions do not include A267S (sometimes referred to herein as AAV2.5G9 A267).

[0011] In embodiments of the capsid proteins, capsids, viral vectors and methods described herein, the one or more amino acid substitutions comprise SQAGASDIRDQSR464-476SX₁AGX₂SX₃X₄X₅X₆QX₇R, wherein X₁₋₇ can be any amino acid, and EYSW500-503EX₈X₉W, wherein X₈₋₉ can be any amino acid.

[0012] In embodiments of the capsid proteins, capsids, viral vectors and methods described herein, X₁ is V or a conservative substitution thereof; X₂ is P or a conservative substitution thereof; X₃ is N or a conservative substitution thereof; X₄ is M or a conservative substitution thereof; X₅ is A or a conservative substitution thereof; X₆ is V or a conservative substitution thereof; X₇ is G or a conservative substitution thereof; X₈ is F or a conservative substitution thereof; and/or X₉ is A or a conservative substitution thereof.

[0013] In embodiments of the capsid proteins, capsids, viral vectors and methods described herein, X₁ is V, X₂ is P, X₃ is N, X₄ is M, X₅ is A, X₆ is V, X₇ is G, X₈ is F, and X₉ is A, wherein the new glycan binding site is a galactose binding site.

[0014] In embodiments of the capsid proteins, capsids, viral vectors and methods described herein, the amino acid sequence of the AAV2.5 capsid protein is SEQ ID NO:1 or a functional derivative thereof (e.g., which does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5).

[0015] In embodiments of the capsid proteins, capsids, viral vectors and methods described herein, the amino acid sequence of the capsid protein is SEQ ID NO:2 or a functional derivative thereof (e.g., which does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5).

[0016] Aspects of the invention relate to a viral capsid comprising the AAV capsid protein described above.

[0017] Aspects of the invention relate to a virus vector comprising the viral capsid described above and a nucleic acid comprising at least one terminal repeat sequence, wherein the nucleic acid is encapsidated by the viral capsid.

[0018] In some embodiments, the AAV2.5G9 described herein evades immune response elicited by pre-existing antibody e.g., antibodies recognizing AAVrh10 or antibodies recognizing other AAV serotypes except AAV2.5G9.

[0019] Aspects of the invention relate to a composition comprising the AAV capsid protein described above, the viral capsid described above, and/or the virus vector described above, in a pharmaceutically acceptable carrier.

[0020] Aspects of the invention relate to a method of introducing a nucleic acid into a cell, comprising contacting the cell with the virus vector described above (e.g., AAV2.5G9 comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof). In some embodiments of the methods described herein, the cell is in neural tissue. In some embodiments of the methods described herein, the cell is a neuron or a glial cell. In some embodiments of the methods described herein, the glial cell is an astrocyte. In some embodiments of the methods described herein, the

virus vector has enhanced transduction of neural tissue as compared to an AAV1, AAV2, AAV9, or AAV2.5 virus vector. In some embodiments of the methods described herein the cell is in a subject. In some embodiments of the methods described herein the subject is a human subject. In some embodiments of the methods described herein the subject is a child. In some embodiments of the methods described herein the child is an infant. In some embodiments of the methods described herein the subject is in utero. In some embodiments of the methods described herein the subject has a reduced immunologic profile when contacted with the virus vector as compared to when contacted with an AAV1, AAV2, AAV9, or AAV2.5 virus vector.

[0021] In some embodiments, the AAV2.5G9 is used for repeat administration of a therapeutic in the method of introducing a nucleic acid into a cell in a subject (e.g., human, child, infant, in utero). In some embodiments, the AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) is used for one administration in the method wherein, AAVrh10 is used for another administration. In some embodiments, the AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) and other AAV vectors used for repeat administration in the method are used at a same viral titer. In some embodiments, AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) and other AAV vectors used for repeat administration in the method are used at different viral titers.

[0022] Aspects of the invention relate to a method of treating a disease or disorder in a subject in need thereof, comprising introducing a therapeutic nucleic acid into a cell of the subject by administering to the subject the virus vector described herein and/or the composition described herein, under conditions whereby the therapeutic nucleic acid is expressed in the cell of the subject. In some embodiments of the methods described herein the subject is a human. In some embodiments of the methods described herein the subject is in utero. In some embodiments of the methods described herein the subject has or is at risk for a CNS disease or disorder. In some embodiments of the methods described herein the subject has or is at risk for a congenital neurodegenerative disorder. In some embodiments of the methods described herein the subject has or is at risk for adult-onset autosomal dominant leukodystrophy (ADLD), Aicardi-Goutieres syndrome, Alexander disease, CADA-SIL, Canavan disease, CARASIL, cerebrotendinous xanthomatosis childhood ataxia and cerebral hypomyelination (CACH)/vanishing white matter disease (VWMD), Fabry disease, fucosidosis, GM1 gangliosidosis, Krabbe disease, L-2-hydroxyglutaric aciduria megalencephalic leukoencephalopathy with subcortical cysts, metachromatic leukodystrophy, multiple sulfatase deficiency, Pelizaeus-Merzbacher disease, Pol III-Related Leukodystrophies, Refsum disease, salla disease (free sialic acid storage disease), Sjogren-Larsson syndrome, X-linked adrenoleukodystrophy, Zellweger syndrome spectrum disorders, Mucopolysaccharidosis Type I, Mucopolysaccharidosis Type II, Mucopolysaccharidosis Type III, Mucopolysaccharidosis Type IV, Mucopolysaccharidosis Type V, Mucopolysaccharidosis Type VI, Mucopolysaccharidosis Type VII, Mucopolysaccharidosis Type IX and any combination thereof. In some embodiments of the methods described herein the subject has or is at risk of having pain associated with a disease or disorder. In some embodiments of the

methods described herein the virus vector or composition is delivered via an enteral, parenteral, intrathecal, intracisternal, intracerebral, intraventricular, intranasal, intra-aural, intra-ocular, peri-ocular, intrarectal, intramuscular, intraperitoneal, intravenous, oral, sublingual, subcutaneous and/or transdermal route. In some embodiments of the methods described herein the virus vector or composition is delivered intracranially and/or intraspinally.

[0023] In some embodiments, the AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) is used in the method of treating a disease or disorder for repeat administration/dosing of the therapeutic (e.g., to a human, child, infant, in utero). In some embodiments, the AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) is used for one administration wherein, AAVrh10 is used for another administration. In some embodiments, the AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) and other AAV vectors used for repeat administration in the method are used at a same viral titer. In some embodiments, AAV2.5G9 (e.g., comprising the capsid protein of SEQ ID NO:2 or a functional derivative thereof) and other AAV vectors used for repeat administration in the method are used at different viral titers.

[0024] To the extent that any disclosure in PCT/US2020/029493 filed Apr. 23, 2020 falls within the invention as defined in any one or more of the claims of this application, or within any invention to be defined in amended claims that may in the future be filed in this application or in any patent derived therefrom, and to the extent that the laws of any relevant country or countries to which that or those claims apply provide that the disclosure of PCT/US2020/029493, filed Apr. 23, 2020, is part of the state of the art against that or those claims in or for that or those countries, we hereby reserve the right to disclaim the said disclosure from the claims of the present application or any patent derived therefrom to the extent necessary to prevent invalidation of the present application or any patent derived therefrom.

[0025] These and other aspects of the invention are addressed in more detail in the description of the invention set forth below.

Definitions

[0026] The singular forms “a,” “an” and “the” are intended to include the plural forms as well, unless the context clearly indicates otherwise.

[0027] Furthermore, the term “about,” as used herein when referring to a measurable value such as an amount of the length of a polynucleotide or polypeptide sequence, dose, time, temperature, and the like, is meant to encompass variations of $\pm 20\%$, $\pm 10\%$, $\pm 5\%$, $\pm 1\%$, $\pm 0.5\%$, or even $\pm 0.1\%$ of the specified amount.

[0028] Also as used herein, “and/or” refers to and encompasses any and all possible combinations of one or more of the associated listed items, as well as the lack of combinations when interpreted in the alternative (“or”).

[0029] Unless the context indicates otherwise, it is specifically intended that the various features of the invention described herein can be used in any combination.

[0030] Moreover, the present invention also contemplates that in some embodiments of the invention, any feature or combination of features set forth herein can be excluded or omitted.

[0031] To illustrate further, if, for example, the specification indicates that a particular amino acid can be selected from A, G, I, L and/or V, this language also indicates that the amino acid can be selected from any subset of these amino acid(s) for example A, G, I or L; A, G, I or V; A or G; only L; etc., as if each such subcombination is expressly set forth herein. Moreover, such language also indicates that one or more of the specified amino acids can be disclaimed. For example, in particular embodiments the amino acid is not A, G or I; is not A; is not G or V; etc., as if each such possible disclaimer is expressly set forth herein.

[0032] The term “tropism” as used herein refers to preferential entry of the virus or viral vector into certain cell or tissue types or preferential interaction with the cell surface that facilitates entry into certain cell or tissue types, optionally and preferably followed by expression (e.g., transcription and, optionally, translation) of sequences carried by the viral genome in the cell.

[0033] The term “target cell” is used to refer to a cell that is infected by the viral vector described herein. In some embodiments, the “target cell” may refer to a cell type that is infected by the virus/viral vector and in which the regulatory elements on the heterologous nucleic acid within promote expression.

[0034] The term “conservative substitution” or “conservative substitution mutation” as used herein refers to a mutation where an amino acid is substituted for another amino acid that has similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure, chemical properties, and/or hydropathic nature of the polypeptide to be substantially unchanged. The following groups of amino acids have been historically substituted for one another as conservative changes: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, try, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. Other commonly accepted conservative substitutions are listed below:

Residue	Conservative Substitutions	Residue	Conservative Substitutions
Ala	Ser	Leu	Ile; Val
Arg	Lys	Lys	Arg; Gln
Asn	Gln; His	Met	Leu; Ile
Asp	Glu	Phe	Met; Leu; Tyr
Gln	Asn	Ser	Thr; Gly
Cys	Ser	Thr	Ser; Val
Glu	Asp	Trp	Tyr
Gly	Pro	Tyr	Trp; Phe
His	Asn; Gln	Val	Ile; Leu
Ile	Leu, Val		

[0035] As used herein, the terms “reduce,” “reduces,” “reduction” and similar terms mean a decrease of at least about 5%, 10%, 15%, 20%, 25%, 35%, 50%, 75%, 80%, 85%, 90%, 95%, 97%, 98%, 99%, 100% or more.

[0036] As used herein, the terms “enhance,” “enhances,” “enhancement” and similar terms indicate an increase of at least about 10%, 20%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400%, 500% or more.

[0037] As used herein, the term “polypeptide” encompasses both peptides and proteins, unless indicated otherwise.

[0038] A “polynucleotide,” “nucleic acid,” or “nucleic acid molecule” as used herein is a sequence of nucleotide

bases, and may be RNA, DNA or DNA-RNA hybrid sequences (including both naturally occurring and non-naturally occurring nucleotide), but in representative embodiments are either single or double stranded DNA sequences.

[0039] As used herein, an “isolated” polynucleotide (e.g., an “isolated DNA” or an “isolated RNA”) means a polynucleotide at least partially separated from at least some of the other components of the naturally occurring organism or virus, for example, the cell or viral structural components or other polypeptides or nucleic acids commonly found associated with the polynucleotide. In representative embodiments an “isolated” nucleotide is enriched by at least about 10-fold, 100-fold, 1000-fold, 10,000-fold or more as compared with the starting material.

[0040] Likewise, an “isolated” polypeptide means a polypeptide that is at least partially separated from at least some of the other components of the naturally occurring organism or virus, for example, the cell or viral structural components or other polypeptides or nucleic acids commonly found associated with the polypeptide. In representative embodiments an “isolated” polypeptide is enriched by at least about 10-fold, 100-fold, 1000-fold, 10,000-fold or more as compared with the starting material.

[0041] As used herein, by “isolate” or “purify” (or grammatical equivalents) a virus vector, it is meant that the virus vector is at least partially separated from at least some of the other components in the starting material. In representative embodiments an “isolated” or “purified” virus vector is enriched by at least about 10-fold, 100-fold, 1000-fold, 10,000-fold or more as compared with the starting material.

[0042] A “therapeutic molecule” (e.g., a nucleic acid or polypeptide) is a molecule that can alleviate, reduce, prevent, delay and/or stabilize symptoms that result from an absence or defect in a protein in a cell or subject and/or is a molecule that otherwise confers a benefit to a subject. Such therapeutic molecules may be encoded by a heterologous nucleic acid present in the viral vector described herein, and under the regulatory sequences that promote expression of the nucleic acid.

[0043] By the terms “treat,” “treating” or “treatment of” (and grammatical variations thereof) it is meant that the severity of the subject’s condition is reduced, at least partially improved or stabilized and/or that some alleviation, mitigation, decrease or stabilization in at least one clinical symptom is achieved and/or there is a delay in the progression of the disease or disorder.

[0044] The terms “prevent,” “preventing” and “prevention” (and grammatical variations thereof) refer to prevention and/or delay of the onset of a disease, disorder and/or a clinical symptom(s) in a subject and/or a reduction in the severity of the onset of the disease, disorder and/or clinical symptom(s) relative to what would occur in the absence of the methods of the invention. The prevention can be complete, e.g., the total absence of the disease, disorder and/or clinical symptom(s). The prevention can also be partial, such that the occurrence of the disease, disorder and/or clinical symptom(s) in the subject and/or the severity of onset is less than what would occur in the absence of the present invention.

[0045] A “treatment effective,” “therapeutic,” or “effective” amount as used herein is an amount that is sufficient to provide some improvement or benefit to the subject. Alternatively stated, a “treatment effective,” “therapeutic,” or

“effective” amount is an amount that will provide some alleviation, mitigation, decrease or stabilization in at least one clinical symptom in the subject. Those skilled in the art will appreciate that the therapeutic effects need not be complete or curative, as long as some benefit is provided to the subject.

[0046] A “prevention effective” amount as used herein is an amount that is sufficient to prevent and/or delay the onset of a disease, disorder and/or clinical symptoms in a subject and/or to reduce and/or delay the severity of the onset of a disease, disorder and/or clinical symptoms in a subject relative to what would occur in the absence of the methods of the invention. Those skilled in the art will appreciate that the level of prevention need not be complete, as long as some benefit is provided to the subject.

[0047] The terms “heterologous nucleotide sequence,” “heterologous nucleic acid,” or “heterologous nucleic acid molecule” are used interchangeably herein and refer to a sequence that is not naturally occurring in the virus. Generally, the heterologous nucleic acid comprises an open reading frame that encodes a polypeptide or nontranslated RNA of interest (e.g., for delivery to a cell or subject) such as a therapeutic or diagnostic molecule.

[0048] As used herein, the terms “virus vector,” “vector” or “gene delivery vector” refer to a virus (e.g., AAV) particle that functions as a nucleic acid delivery vehicle, and which comprises the vector genome (e.g., viral DNA [vDNA]) packaged within a virion. Alternatively, in some contexts, the term “vector” may be used to refer to the vector genome/vDNA alone.

[0049] As used herein when referring to viruses, the terms “vector,” “particle,” and “virion” may be used interchangeably.

[0050] The virus vectors of the invention can further be “targeted” virus vectors (e.g., having a directed tropism) and/or a “hybrid” parvovirus (i.e., in which the viral TRs and viral capsid are from different parvoviruses), e.g., as described in international patent publication WO 00/28004, the disclosure of which is incorporated herein by reference in its entirety.

[0051] The virus vectors of the invention can further be duplexed parvovirus particles as described in international patent publication WO 01/92551 (the disclosure of which is incorporated herein by reference in its entirety). Thus, in some embodiments, double stranded (duplex) genomes can be packaged into the virus capsids of the invention.

BRIEF DESCRIPTION OF THE DRAWINGS

[0052] FIG. 1 shows effects of AAV administration on weight. Fetal brain and body weights of AAV-treated (combined AAV9, AAV2G9, and AAV2.5G9; N=9) and control fetuses (N=36) were compared. No significant differences were observed between the groups ($p < 0.05$). Data are shown as mean \pm standard error of the mean. Significance was determined by Student’s t-test analysis at $p \leq 0.05$.

[0053] FIG. 2 shows detection of AAV-mediated firefly luciferase transduction and expression by bioluminescence. Individual sections of the right and left hemispheres (frontal, parietal, temporal, occipital lobes) from fetuses administered AAV9, AAV2G9, and AAV2.5G9 were imaged for bioluminescence. Each image corresponds to an animal number as noted in Table 3. Total bioluminescence is noted below each image in photons/second (p/s). Data are shown

as mean \pm standard error of the mean. Significance was determined by Student's t-test analysis at $p \leq 0.05$.

DETAILED DESCRIPTION OF THE INVENTION

[0054] The present invention will now be described with reference to the accompanying drawings, in which representative embodiments of the invention are shown. This invention may, however, be embodied in different forms and should not be construed as limited to the embodiments set forth herein. Rather, these embodiments are provided so that this disclosure will be thorough and complete, and will fully convey the scope of the invention to those skilled in the art.

[0055] Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. The terminology used in the description of the invention herein is for the purpose of describing particular embodiments only and is not intended to be limiting of the invention. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference herein in their entirety.

[0056] To the extent that any disclosure in PCT/US2020/029493 filed Apr. 23, 2020 falls within the invention as defined in any one or more of the claims of this application, or within any invention to be defined in amended claims that may in the future be filed in this application or in any patent derived therefrom, and to the extent that the laws of any relevant country or countries to which that or those claims apply provide that the disclosure of PCT/US2020/029493 is part of the state of the art against that or those claims in or for that or those countries, we hereby reserve the right to disclaim the said disclosure from the claims of the present application or any patent derived therefrom to the extent necessary to prevent invalidation of the present application or any patent derived therefrom.

[0057] Aspects of the invention relate to the discovery of a "pocket" on the AAV capsid protein that defines a glycan recognition footprint and the grafting of such a recognition footprint onto a heterologous capsid protein to thereby produce a novel AAV capsid protein. Specific amino acids that define this pocket have been identified and are described herein, for example for the galactose binding site of AAV9. Thus the present invention is directed to molecular grafting of a new glycan recognition footprint (e.g., that of a donor AAV strain capsid) onto a capsid protein to thereby modify the capsid protein. Such grafting is guided by structural modeling studies and achieved by site-directed mutagenesis. Recombinant vectors (having capsids derived from such grafting) carrying transgenes (e.g., reporter cassettes) display rapid onset and enhanced transgene expression in cell culture and animal models. Viral vectors generated from this strategy can address challenges such as dose-dependent immunotoxicity observed in human gene therapy clinical trials.

[0058] In some embodiments, the substitutions introduce a glycan binding site from a capsid protein of a first AAV serotype ("donor") into the capsid protein of a second AAV serotype ("template") that is different from said first AAV serotype. Thus, in one aspect, the present invention relates to an adeno-associated virus (AAV) capsid protein, which comprises an AAV2.5 capsid comprising one or more amino acids substitutions, wherein the substitutions introduce a glycan binding site into the AAV capsid protein, to thereby

produce a "modified capsid protein," or a "modified AAV2.5 capsid protein." In some embodiments, the glycan binding site can be a hexose binding site, wherein the hexose is a galactose (Gal), a mannose (Man), a glucose (Glu) or a fucose (fuc). In some embodiments, the glycan binding site can be a sialic acid (Sia) binding site, wherein the Sia residue is N-acetylneuraminic acid (Neu5Ac) or N-Glycolylneuraminic acid (Neu5Gc). In some embodiments, the glycan binding site can be a disaccharide binding site, wherein the disaccharide is a sialic acid linked to galactose in the form Sia(α 2,3)Gal or Sia(α 2,6)Gal.

[0059] In some embodiments, the glycan binding site is a galactose binding site. In some embodiments, the AAV9 galactose binding site (donor) is grafted into an AAV2.5 capsid protein template, resulting in the introduction of a new glycan binding site in the engrafted (modified) AAV2.5 capsid protein template. The new glycan binding site is generated by the introduction of one or more amino acid substitutions into the AAV2.5 capsid template.

[0060] Immunogenicity of a given AAV virus can be altered from even a slight change in amino acid sequence (e.g., of the capsid protein). In this way, the neutralizing antibody response of a subject exposed to a given AAV vector therapeutic can also differ dramatically to that of the same AAV vector therapeutic with altered amino acid sequence. Generally, mutants that have few amino acid substitutions are preferred for immunological reasons, as few mutations will lead to fewer reactive antibodies. In some instances, the ability to generate capsid mutants from different amino acid substitutions and/or insertions that exhibit the same phenotype will also allow the physician to draw on a plurality of such rationally designed viral vectors. This would be advantageous in reducing the impact of pre-existing neutralizing antibodies on the administered therapeutic, and could permit repeated therapeutic administration using the different vectors.

[0061] In some embodiments, AAV2.5G9 evades immune response elicited by pre-existing antibodies e.g., antibodies recognizing AAVrh10 or antibodies recognizing other AAV serotypes except AAV2.5G9. In some embodiments, the pre-existing antibodies are neutralizing antibodies. In some embodiments, AAV2.5G9 is used for repeat administration of a therapeutic. In some embodiments, AAV2.5G9 is used for one administration wherein, AAVrh10 is used for the other administration. In some embodiments, AAV2.5G9 and other AAV vectors used for repeat administration are used at a same viral titer. In some embodiments, AAV2.5G9 and other AAV vectors used for repeat administration are used at different viral titers.

[0062] In some embodiments, the amino acid substitutions comprise:

a) SQAGASDIRDQSR464-476SX₁AGX₂SX₃X₄X₅X₆QX₇R, wherein X₁₋₇ can be any amino acid; and b) EYSW500-503EX₈X₉W, wherein X₈₋₉ can be any amino acid. In some embodiments, the resulting modified AAV2.5 capsid protein does not contain a substitution at position 267. In some embodiments, the resulting modified AAV2.5 capsid protein does not contain a substitution of serine for alanine at position 267.

[0063] In some embodiments, the amino acid substitutions are in amino acids 464-476, and/or amino acids 500-503 in AAV2.5 (SEQ ID NO:1; VP1 numbering).

[0064] In some embodiments, X₁ is V or a conservative substitution thereof, X₂ is P or a conservative substitution

thereof, X_3 is N or a conservative substitution thereof, X_4 is M or a conservative substitution thereof, X_5 is A or a conservative substitution thereof, X_6 is V or a conservative substitution thereof, X_7 is G or a conservative substitution thereof, X_8 is F or a conservative substitution thereof, and/or X_9 is A or a conservative substitution thereof.

[0065] In some embodiments, X_1 is V, X_2 is P, X_3 is N, X_4 is M, X_5 is A, X_6 is V, X_7 is G, X_8 is F, and X_9 is A, to thereby result in a new glycan binding site that is a galactose binding site.

[0066] The AAV2.5 capsid template may have the amino acid sequence of SEQ ID NO:1, or a functional derivative thereof. A functional derivative of an amino acid sequence may have an amino acid substitution, insertion or deletion, which substantially preserves one or more properties or functions of the original sequence. In some embodiments, the functional derivative does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5. In some embodiments, the functional derivative does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

[0067] Functional derivatives have amino acid substitutions, insertions and/or deletions that do not substantially affect protein function such as the derivatives will retain one or more activities (properties or functions) when compared to that of the original protein (e.g., SEQ ID NO:1). Such derivatives will retain at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95%, 96%, 97%, 98% 99% or be indistinguishable (not significantly different) with respect to one or more activities of the original protein. Such activities include, without limitation, one or more cell type and/or tissue tropism.

[0068] In some embodiments, the functional derivative results from one or more conservative amino acid substitutions of SEQ ID NO:1. Examples of conservative amino acid substitutions are provided herein. In some embodiments, the functional derivative does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5. In some embodiments, the functional derivative does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

[0069] In some embodiments, the AAV capsid protein template/backbone is from AAV2.5 (SEQ ID NO:1; VP1

numbering), and the amino acid substitutions are a Q465V substitution, an A468P substitution, a D470N substitution, an I471M substitution, an R472A substitution, a D473V substitution, an S475G substitution, a Y501F substitution, and/or an S502A substitution, in any combination. Thus, in some embodiments, the present invention provides an AAV2.5 capsid protein, comprising an AAV capsid protein backbone from AAV2.5 (SEQ ID NO:1; VP1 numbering) comprising a Q465V substitution, an A468P substitution, a D470N substitution, an I471M substitution, an R472A substitution, a D473V substitution, an S475G substitution, a Y501F substitution, and an S502A substitution, wherein the substitutions introduce a glycan binding site into the AAV2.5 capsid protein. In some embodiments, the AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

[0070] The AAV2.5 capsid protein that serves as the template originated from specific mutations to the AAV2 capsid sequences as described in U.S. Pat. No. 9,012,224, the contents of which are incorporated herein by reference. This was generated by changing 5 amino acids in AAV2 to resemble AAV1 at those specific locations (Q263A; 265insT; N705A; V708A; T716N). The resulting amino acid modifications are shown in the below sequences (SEQ ID NO:1) as capital letters. The properties conferred to a viral particle from the resulting AAV2.5 capsid protein are well characterized (U.S. Pat. No. 9,012,224). Without limitation, the properties the AAV2.5 capsid confers to a viral particle include, enhanced skeletal muscle tropism, reduced liver-hepatocyte tropism as compared to AAV2, neural tropism, and glial tropism (e.g., astrocytes), as well as the ability to escape neutralization from existing AAV2 and AAV1 neutralizing antibodies. In some embodiments of the invention described herein, the amino acid sequence of the AAV2.5 capsid protein is that shown in SEQ ID NO:1, which utilizes VP1 numbering. In some embodiments, the AAV2.5 capsid protein is a functional derivative of the capsid protein having the amino acid sequence of SEQ ID NO:1. In some embodiments, the functional derivative does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

SEQ ID NO: 1. AAV2.5 capsid protein

```

1  maadgylpdw ledtlsegir qwwklkpgpp ppkpaerhkd dsrglvpky kylgpfngld
61  kgepvneada aalehdkayd rqlsdgdnpy lkynhadaef qerlkedtsf ggnlgravfq
121 akkrvleplg lveepvktap gkkrvvehsp vepdsssgtg kagqqparkr lnfgqtgdad
181 svdpqpplgq ppaapsglgt ntmatgsgap madnnegadg vgnssgnwhc dstwmgdrvi
241 ttstrtwalp tynnhlykqi ssAsTgasnd nhyfgystpw gyfdfnrfhc hfsprdwqrl
301 innnwgfprk rlnfklnfiq vkevtqndgt ttiannltst vqvftdseyq lpyvlgsahq
361 gclppfpadv fmvpygylyt lnngsqavgr ssfycleyfp sqmlrtggnf tfsytfedvp
421 fhssyahsqd ldrlmnlid qylyylsrtn tpsgtttqsr lqfsqagasd irdqsrnwlp
481 gpcyrqqrvs ktsadnnnse yswtgatkyh lngrdslvnp gpamashkdd eekffpqsgv
541 lifgkqgsek tnvdiekvmi tdeeeirttn pvategygs vstnlqrgrnq aatadvntqg

```


-continued

601 vlpqmvwqdr dvylqgpiwa kiphtdgfhf pspmggfgl khpppqilik ntpvpanpst

661 tfsaakfasf itqystgqvs veiewelqke nskrwnpeiq ytsnyAksAn vdftrvdNngv

721 YSeprpigtr yltrnl (AAV2.5)

[0071] In some embodiments, the AAV2.5 capsid has an amino acid sequence that is at least about 70%, 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% similar or identical to that shown in SEQ ID NO:1. In some embodiments, the AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

[0072] For example, in particular embodiments, an “AAV2.5” capsid protein encompasses the amino acid sequence shown in SEQ ID NO:1, as well as sequences that are at least about 75%, 80%, 85%, 90%, 95%, 97%, 98% or 99% similar or identical to the amino acid sequence of SEQ ID NO:1, wherein the AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

[0073] Methods of determining sequence similarity or identity between two or more amino acid sequences are known in the art. Sequence similarity or identity may be determined using standard techniques known in the art, including, but not limited to, the local sequence identity algorithm of Smith & Waterman. *Adv. Appl. Math.* 2:482 (1981), by the sequence identity alignment algorithm of Needleman & Wunsch. *J. Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman. *Proc. Natl. Acad. Sci. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Drive, Madison, Wis.), the Best Fit sequence program described by Devereux et al. *Nucl. Acid Res.* 12:387-395 (1984), or by inspection.

[0074] Another suitable algorithm is the BLAST algorithm, described in Altschul et al. *J. Mol. Biol.* 215:403-410 (1990) and Karlin et al. *Proc. Natl. Acad. Sci. USA* 90:5873-5787 (1993). A particularly useful BLAST program is the WU-BLAST-2 program which was obtained from Altschul et al. *Methods in Enzymology* 266:460-480 (1996); blast.wustl.edu/blast/README.html. WU-BLAST-2 uses several search parameters, which are optionally set to the default values. The parameters are dynamic values and are established by the program itself depending upon the composition of the particular sequence and composition of the particular database against which the sequence of interest is being searched; however, the values may be adjusted to increase sensitivity.

[0075] Further, an additional useful algorithm is gapped BLAST as reported by Altschul et al. *Nucleic Acids Res.* 25:3389-3402 (1997).

[0076] In some embodiments, a modified AAV capsid protein of the present invention has the amino acid sequence shown in SEQ ID NO:2, or is a functional derivative thereof, (e.g., wherein the AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5).

[0077] In some embodiments, the modified AAV capsid protein comprises, consists essentially of, or consists of the nucleotide sequence of SEQ ID NO:2 or a nucleotide sequence at least about 70% identical thereto, e.g., at least about 70, 75, 80, 85, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% identical thereto (e.g., wherein the AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5).

SEQ ID NO: 2. AAV2.5G9 A267 capsid protein

1 maadgyldpw ledtlsegir qwwklkpgpp ppkpaerhkd dsrglvlpqy kylgpfngld

61 kgepvneada aalehdkayd rqlsdgdnpy lkynhadaef qerlkedtsf ggnlgravfq

121 akkrvleplg lveepvktap gkkpvehsp vepdsssgtg kagqqparkr lnfgqtdgad

181 svdpdpplgq ppaapslgt ntmatgsgap madnnegadg vgnssgnwhc dstwmgdrvi

241 ttstrtwalp tynnhlykqi ssAsTgasnd nhyfgystpw gyfdfnrfhc hfsprdwqrl

301 innnwgfprk rlnfklfnig vkevtqndgt ttiannltst vqvftdseyq lpyvlgsahq

361 gclppfpadv fmvpygygt lnngsqavgr ssfycleyfp sqmlrtgnnf tfsytfedvp

421 fhssyahsqsl drlmnlid qlyylsrtn tpsgtttqsr lqfSVAGPSN MAVQGRnwl

481 gpcyrqgrvs ktsadnnnsE FAWtgatkyh lngrdslvnp gpamashkdd eekffpqsgv

541 lifgkqgsek tnvdiekvmi tdeeeirttn pvategygsv stnlqrgrnq aatadvntqg

-continued

601 vlpgmvwqdr dvylqgpiwa kiphtdghfh psplmggfgl khpppqilik ntpvpanpst

661 tfsaakfasf itqystgqvs veiewelqke nskrwnpeiq ytsnyAksAn vdftvdNngv

72lyseprpigtr yltrnl (AAV2.5G9 A267)

[0078] The examples provided herein indicate possible amino acid substitutions in an AAV2.5 template for introduction of a galactose binding site from an AAV9 donor. These examples, which are not intended to be limiting, demonstrate the principle that a glycan binding site from a donor AAV serotype can be introduced into a capsid protein template of a different AAV serotype by modifying residues that define the “pocket” described herein.

[0079] Such modified or chimeric capsid proteins comprising a new glycan binding site can be assembled into capsids that make up virus particles that can be used as virus vectors that have the beneficial phenotype of increased cell surface binding and more rapid and enhanced transgene expression in vivo.

[0080] Table 2 lists non-limiting exemplary serotypes of AAV and accession numbers of the genome and capsid sequences that may be encompassed by the invention. The AAV serotype of the donor and the template is not limited to human AAV, but may include non-human AAV, for example, Avian or Bovine AAV, as well as non-human primate AAV, examples of which are shown in Table 1. As used herein, the term “adeno-associated virus” (AAV), includes but is not limited to, AAV type 1, AAV type 2, AAV type 2.5, AAV type 3 (including types 3A and 3B), AAV type 4, AAV type 5, AAV type 6, AAV type 7, AAV type 8, AAV type 9, AAV type 10, AAV type 11, avian AAV, bovine AAV, canine AAV, equine AAV, ovine AAV, Clade F AAV and any other AAV now known or later discovered. See, e.g., BERNARD N. FIELDS et al. VIROLOGY, Volume 2, Chapter 69 (4th ed., Lippincott-Raven Publishers). A number of relatively new AAV serotypes and clades have been identified (see Table 1).

[0081] The genomic sequences of various serotypes of AAV, as well as the sequences of the native terminal repeats (TRs), Rep proteins, and capsid subunits are known in the art. Exemplary but non-limiting examples of such sequences may be found in the literature or in public databases such as GenBank® Database. See, e.g., GenBank® Database Accession Numbers NC_002077.1, NC_001401.2, NC_001729.1, NC_001863.1, NC_001829.1, NC_006152.1, NC_001862.1, AF513851.1, AF513852.1, the disclosures of which are incorporated by reference herein for teaching parvovirus and AAV nucleic acid and amino acid sequences. See also, e.g., Srivastava et al. *J. Virology* 45:555 (1983); Chiorini et al. *J. Virology* 71:6823 (1998); Chiorini et al. *J. Virology* 73:1309 (1999); Bantel-Schaal et al. *J. Virology* 73:939 (1999); Xiao et al. *J. Virology* 73:3994 (1999); Muramatsu et al. *Virology* 221:208 (1996); Shade et al. *J. Virol.* 58:921 (1986); Gao et al. (2002) *Proc. Nat. Acad. Sci. USA* 99:11854 (2002); International Patent Publications WO 00/28061, WO 99/6160 and WO 98/11244; and U.S. Pat. No. 6,156,303; the disclosures of which are incorporated by reference herein for teaching parvovirus and AAV nucleic acid and amino acid sequences.

[0082] The capsid structures of autonomous parvoviruses and AAV are described in more detail in BERNARD N. FIELDS et al. VIROLOGY, Volume 2, Chapters 69 & 70

(4th ed., Lippincott-Raven Publishers). See also, description of the crystal structure of AAV2 (Xie et al. *Proc. Nat. Acad. Sci.* 99:10405-10 (2002)), AAV4 (Padron et al. *J. Virol.* 79: 5047-58 (2005)), AAV5 (Walters et al. *J. Virol.* 78: 3361-71 (2004)) and CPV (Xie et al. *J. Mol. Biol.* 6:497-520 (1996) and Tsao et al. *Science* 251: 1456-64 (1991)).

[0083] Other aspects of the invention relate to a capsid comprising the modified AAV capsid protein of the invention, e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5.

[0084] Other aspects of the invention relate to a virus vector or particle comprising (a) the modified AAV capsid of the invention (e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5); and (b) a nucleic acid comprising at least one terminal repeat sequence, wherein the nucleic acid is encapsidated by the capsid.

[0085] Another aspect of the invention relates to a composition comprising the modified AAV2.5 capsid protein, and/or the AAV capsid and/or virus vector comprising the modified AAV2.5 capsid protein, in a pharmaceutically acceptable carrier (e.g. wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5).

[0086] The present invention additionally provides a method of introducing a nucleic acid into a cell, comprising contacting the cell with a virus vector comprising the modified AAV2.5 capsid protein (e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5). The cell can be in a subject and in some embodiments, the subject can be a human subject. In some embodiments, the subject may be in utero. In some embodiments, the cell may be a neural cell (e.g., a neuronal cell or a glial cell, e.g., a cell of neuronal tissue). In some embodiments, the resultant virus vectors have enhanced transduction (e.g., enhanced levels of nucleic acid expression in) a cell (e.g., a neural cell, e.g., a neuronal cell) when contacted to the cell as compared to transduction levels of virus vectors of the donor and template serotypes when contacted to the cell. For example, if the AAV capsid protein donor is AAV serotype 9 and the AAV capsid protein template is AAV serotype 2.5 (AAV2.5), the resulting virus vector would be compared to AAV1, AAV2, AAV9, or AAV2.5.

Methods of Producing Virus Vectors

[0087] The invention also encompasses virus vectors comprising the modified capsid proteins and capsids of the

invention. In particular embodiments, the virus vector is a parvovirus vector (e.g., comprising a parvovirus capsid and/or vector genome), for example, an AAV vector (e.g., comprising a AAV capsid and/or vector genome). In representative embodiments, the virus vector comprises a modified AAV capsid comprising a modified capsid protein subunit of the invention and a vector genome.

[0088] For example, in representative embodiments, the virus vector comprises: (a) a modified virus capsid (e.g., a modified AAV capsid) comprising a modified capsid protein of the invention; and (b) a nucleic acid comprising a terminal repeat sequence (e.g., an AAV TR), wherein the nucleic acid comprising the terminal repeat sequence is encapsidated by the modified virus capsid. The nucleic acid can optionally comprise two terminal repeats (e.g., two AAV TRs).

[0089] In representative embodiments, the virus vector is a recombinant virus vector wherein the genome of the virus comprises a heterologous nucleic acid of interest. The heterologous nucleic acid may encode a polypeptide or functional RNA of interest. Recombinant virus vectors are described in more detail below. The heterologous nucleic acid can be operably linked to appropriate control sequences to promote expression in the target cell. For example, the heterologous nucleic acid can be operably associated with expression control elements, such as transcription/translation control signals, origins of replication, polyadenylation signals, internal ribosome entry sites (IRES), promoters, and/or enhancers, and the like.

[0090] In particular embodiments, the virus vectors of the invention have altered (e.g., reduced) transduction of liver as compared with the level of transduction by a virus vector without the modified capsid protein (e.g., as compared to a virus vector with AAV2.5 capsid protein). In particular embodiments, the virus vector has systemic transduction toward muscle, e.g., the vector transduces multiple skeletal muscle groups throughout the body and optionally transduces cardiac muscle and/or diaphragm muscle. In some embodiments, the virus vectors of the invention have enhanced transduction (e.g., enhanced levels of nucleic acid expression in) neural (e.g., neuronal, glial such as astrocyte or oligodendrocyte) tissue, as compared to an appropriate control (e.g., as compared to other tissues, and/or as compared to transduction levels with other virus vectors, for example, a vector without the modified capsid protein, e.g., AAV1, AAV2, AAV9, AAV2.5, or any AAV serotype as listed in Table 1). An appropriate control may be an otherwise identical viral vector that has not received the grafted glycan binding site.

[0091] In some embodiments, the AAV2.5G9 exhibits substantially increased (e.g., over 10x, 20x, 30x, 40x, 50x, 60x, 70x, 80x, 90x, or over 100x) transduction in a given cell type as compared to AAVrh10. For example, the AAV2.5G9 may exhibit from over 10x to over 100x transduction in Human skin fibroblast GM16095 cells as compared to AAVrh10 as measured by Relative Luciferase unit RLU Luc. The AAV2.5G9 may exhibit from over 10x to over 100x transduction in neuronal cells or glial cells (such as astrocyte or oligodendrocyte) as compared to AAVrh10 as measured by Relative Luciferase unit RLU Luc.

[0092] It will be understood by those skilled in the art that the modified capsid proteins, virus capsids and virus vectors of the invention exclude those capsid proteins, capsids and virus vectors that have the indicated amino acids at the

specified positions in their native state (i.e., are not “modified” with the substitutions described herein that introduce a glycan binding site).

[0093] The present invention further provides methods of producing the inventive virus vectors. In one representative embodiment, the present invention provides a method of producing a virus vector, the method comprising providing to a cell: (a) a nucleic acid template comprising at least one TR sequence (e.g., AAV TR sequence), and (b) AAV sequences sufficient for replication of the nucleic acid template and encapsidation into AAV capsids (e.g., AAV rep sequences and AAV cap sequences encoding the AAV capsids of the invention). Optionally, the nucleic acid template further comprises at least one heterologous nucleic acid sequence. In particular embodiments, the nucleic acid template comprises two AAV ITR sequences, which are located 5' and 3' to the heterologous nucleic acid sequence (if present), although they need not be directly contiguous thereto.

[0094] The nucleic acid template and AAV rep and cap sequences are provided under conditions such that virus vector comprising the nucleic acid template packaged within the AAV capsid is produced in the cell. The method can further comprise the step of collecting the virus vector from the cell. The virus vector can be collected from the medium and/or by lysing the cells.

[0095] The cell can be a cell that is permissive for AAV viral replication. Any suitable cell known in the art may be employed. In particular embodiments, the cell is a mammalian cell. As another option, the cell can be a trans-complementing packaging cell line that provides functions deleted from a replication-defective helper virus, e.g., 293 cells or other Ela trans-complementing cells.

[0096] The AAV replication and capsid sequences may be provided by any method known in the art. Current protocols typically express the AAV replcap genes on a single plasmid. The AAV replication and packaging sequences need not be provided together, although it may be convenient to do so. The AAV rep and/or cap sequences may be provided by any viral or non-viral vector. For example, the rep/cap sequences may be provided by a hybrid adenovirus or herpesvirus vector (e.g., inserted into the Ela or E3 regions of a deleted adenovirus vector). EBV vectors may also be employed to express the AAV cap and rep genes. One advantage of this method is that EBV vectors are episomal, yet will maintain a high copy number throughout successive cell divisions (i.e., are stably integrated into the cell as extra-chromosomal elements, designated as an “EBV based nuclear episome.”

[0097] As a further alternative, the rep/cap sequences may be stably incorporated into a cell.

[0098] Typically the AAV rep/cap sequences will not be flanked by the TRs, to prevent rescue and/or packaging of these sequences.

[0099] The nucleic acid template can be provided to the cell using any method known in the art. For example, the template can be supplied by a non-viral (e.g., plasmid) or viral vector. In particular embodiments, the nucleic acid template is supplied by a herpesvirus or adenovirus vector (e.g., inserted into the Ela or E3 regions of a deleted adenovirus). As another example, a baculovirus vector carrying a reporter gene flanked by the AAV TRs can be used. EBV vectors may also be employed to deliver the template, as described above with respect to the replcap genes.

[0100] In another representative embodiment, the nucleic acid template is provided by a replicating rAAV virus. In still other embodiments, an AAV provirus comprising the nucleic acid template is stably integrated into the chromosome of the cell.

[0101] To enhance virus titers, helper virus functions (e.g., adenovirus or herpesvirus) that promote a productive AAV infection can be provided to the cell. Helper virus sequences necessary for AAV replication are known in the art. Typically, these sequences will be provided by a helper adenovirus or herpesvirus vector. Alternatively, the adenovirus or herpesvirus sequences can be provided by another non-viral or viral vector, e.g., as a non-infectious adenovirus miniplasmid that carries all of the helper genes that promote efficient AAV production.

[0102] Further, the helper virus functions may be provided by a packaging cell with the helper sequences embedded in the chromosome or maintained as a stable extrachromosomal element. Generally, the helper virus sequences cannot be packaged into AAV virions, e.g., are not flanked by TRs.

[0103] Those skilled in the art will appreciate that it may be advantageous to provide the AAV replication and capsid sequences and the helper virus sequences (e.g., adenovirus sequences) on a single helper construct. This helper construct may be a non-viral or viral construct. As one nonlimiting illustration, the helper construct can be a hybrid adenovirus or hybrid herpesvirus comprising the AAV replcap genes.

[0104] In one particular embodiment, the AAV rep/cap sequences and the adenovirus helper sequences are supplied by a single adenovirus helper vector. This vector can further comprise the nucleic acid template. The AAV replcap sequences and/or the rAAV template can be inserted into a deleted region (e.g., the Ela or E3 regions) of the adenovirus.

[0105] In a further embodiment, the AAV rep/cap sequences and the adenovirus helper sequences are supplied by a single adenovirus helper vector. According to this embodiment, the rAAV template can be provided as a plasmid template.

[0106] In another illustrative embodiment, the AAV rep/cap sequences and adenovirus helper sequences are provided by a single adenovirus helper vector, and the rAAV template is integrated into the cell as a provirus. Alternatively, the rAAV template is provided by an EBV vector that is maintained within the cell as an extrachromosomal element (e.g., as an EBV based nuclear episome).

[0107] In a further exemplary embodiment, the AAV rep/cap sequences and adenovirus helper sequences are provided by a single adenovirus helper. The rAAV template can be provided as a separate replicating viral vector. For example, the rAAV template can be provided by a rAAV particle or a second recombinant adenovirus particle.

[0108] According to the foregoing methods, the hybrid adenovirus vector typically comprises the adenovirus 5' and 3' cis sequences sufficient for adenovirus replication and packaging (i.e., the adenovirus terminal repeats and PAC sequence). The AAV replcap sequences and, if present, the rAAV template are embedded in the adenovirus backbone and are flanked by the 5' and 3' cis sequences, so that these sequences may be packaged into adenovirus capsids. As described above, the adenovirus helper sequences and the AAV rep/cap sequences are generally not flanked by TRs so that these sequences are not packaged into the AAV virions.

[0109] Zhang et al. (*Gene Ther.* 18:704-12 (2001)) describes a chimeric helper comprising both adenovirus and the AAV rep and cap genes.

[0110] Herpesvirus may also be used as a helper virus in AAV packaging methods. Hybrid herpesviruses encoding the AAV Rep protein(s) may advantageously facilitate scalable AAV vector production schemes. A hybrid herpes simplex virus type I (HSV-1) vector expressing the AAV2 rep and cap genes has been described, e.g., PCT Publication No. WO 00/17377, incorporated by reference herein.

[0111] As a further alternative, the virus vectors of the invention can be produced in insect cells using baculovirus vectors to deliver the rep/cap genes and rAAV template.

[0112] AAV vector stocks free of contaminating helper virus may be obtained by any method known in the art. For example, AAV and helper virus may be readily differentiated based on size. AAV may also be separated away from helper virus based on affinity for a heparin substrate. Deleted replication-defective helper viruses can be used so that any contaminating helper virus is not replication competent. As a further alternative, an adenovirus helper lacking late gene expression may be employed, as only adenovirus early gene expression is required to mediate packaging of AAV virus. Adenovirus mutants defective for late gene expression are known in the art (e.g., ts100K and ts149 adenovirus mutants).

Recombinant Virus Vectors

[0113] The virus vectors of the present invention are useful for the delivery of nucleic acids to cells in vitro, ex vivo, and in vivo. In particular, the virus vectors (comprising modified capsid protein AAV2.5, e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5) can be advantageously employed to deliver or transfer nucleic acids to animal cells, including e.g., mammalian cells.

[0114] Any heterologous nucleic acid sequence(s) of interest may be delivered in the virus vectors of the present invention. Nucleic acids of interest include nucleic acids encoding polypeptides, including therapeutic (e.g., for medical or veterinary uses) and/or immunogenic (e.g., for vaccines) polypeptides.

[0115] Therapeutic polypeptides include, but are not limited to, cystic fibrosis transmembrane regulator protein (CFTR), dystrophin (including mini- and micro-dystrophins, see, e.g., Vincent et al. *Nature Genetics* 5:130 (1993); U.S. Patent Publication No. 2003017131; PCT Publication No. WO/2008/088895, Wang et al. *Proc. Natl. Acad. Sci. USA* 97:13714-13719 (2000); and Gregorevic et al. *Mol. Ther.* 16:657-64 (2008)), myostatin propeptide, follistatin, activin type II soluble receptor, IGF-1, anti-inflammatory polypeptides such as the I kappa B dominant mutant, sarcospan, utrophin (Tinsley et al. *Nature* 384:349 (1996)), mini-utrophin, clotting factors (e.g., Factor VIII, Factor IX, Factor X, etc.), erythropoietin, angiostatin, endostatin, catalase, tyrosine hydroxylase, superoxide dismutase, leptin, the LDL receptor, lipoprotein lipase, ornithine transcarbamylase, β -globin, α -globin, spectrin, α_1 -antitrypsin, adenosine deaminase, hypoxanthine guanine phosphoribosyl transferase, β -glucocerebrosidase, sphingomyelinase, lysosomal hexosaminidase A, branched-chain keto acid dehydrogenase, RP65 protein, cytokines (e.g., α -interferon, β -inter-

feron, interferon- γ , interleukin-2, interleukin-4, granulocyte-macrophage colony stimulating factor, lymphotoxin, and the like), peptide growth factors, neurotrophic factors and hormones (e.g., somatotropin, insulin, insulin-like growth factors 1 and 2, platelet derived growth factor, epidermal growth factor, fibroblast growth factor, nerve growth factor, neurotrophic factor-3 and -4, brain-derived neurotrophic factor, bone morphogenic proteins [including RANKL and VEGF], glial derived growth factor, transforming growth factor- α and - β , and the like), lysosomal acid α -glucosidase, α -galactosidase A, receptors (e.g., the tumor necrosis growth factor α soluble receptor), S100A1, parvalbumin, adenylyl cyclase type 6, a molecule that modulates calcium handling (e.g., SERCA_{2A}, Inhibitor 1 of PP1 and fragments thereof [e.g., PCT Publication Nos. WO 2006/029319 and WO 2007/100465]), a molecule that effects G-protein coupled receptor kinase type 2 knockdown such as a truncated constitutively active bARKct, anti-inflammatory factors such as IRAP, anti-myostatin proteins, aspartoacylase, monoclonal antibodies (including single chain monoclonal antibodies; an exemplary Mab being the Herceptin® Mab), neuropeptides and fragments thereof (e.g., galanin, Neuropeptide Y (see U.S. Pat. No. 7,071,172), angiogenesis inhibitors such as Vasohibins and other VEGF inhibitors (e.g., Vasohibin 2 [see PCT Publication WO JP2006/073052])). Other illustrative heterologous nucleic acid sequences encode suicide gene products (e.g., thymidine kinase, cytosine deaminase, diphtheria toxin, and tumor necrosis factor), proteins conferring resistance to a drug used in cancer therapy, tumor suppressor gene products (e.g., p53, Rb, Wt-1), TRAIL, FAS-ligand, and any other polypeptide that has a therapeutic effect in a subject in need thereof. AAV vectors can also be used to deliver monoclonal antibodies and antibody fragments, for example, an antibody or antibody fragment directed against myostatin (see, e.g., Fang et al. *Nature Biotechnology* 23:584-590 (2005)).

[0116] Heterologous nucleic acid sequences encoding polypeptides include those encoding reporter polypeptides (e.g., an enzyme). Reporter polypeptides are known in the art and include, but are not limited to, green fluorescent protein (GFP), β -galactosidase, alkaline phosphatase, luciferase, and chloramphenicol acetyltransferase gene.

[0117] Optionally, the heterologous nucleic acid encodes a secreted polypeptide (e.g., a polypeptide that is a secreted polypeptide in its native state or that has been engineered to be secreted, for example, by operable association with a secretory signal sequence as is known in the art).

[0118] Alternatively, in particular embodiments of this invention, the heterologous nucleic acid may encode an antisense nucleic acid, a ribozyme (e.g., as described in U.S. Pat. No. 5,877,022), RNAs that effect spliceosome-mediated trans-splicing (see, Puttaraju et al. *Nature Biotech.* 17:246 (1999); U.S. Pat. Nos. 6,013,487; 6,083,702), interfering RNAs (RNAi) including siRNA, shRNA or miRNA that mediate gene silencing (see, Sharp et al. *Science* 287:2431 (2000)), and other non-translated RNAs, such as “guide” RNAs (Gorman et al. *Proc. Nat. Acad. Sci. USA* 95:4929 (1998); U.S. Pat. No. 5,869,248 to Yuan et al.), and the like. Exemplary untranslated RNAs include RNAi against a multiple drug resistance (MDR) gene product (e.g., to treat and/or prevent tumors and/or for administration to the heart to prevent damage by chemotherapy), RNAi against myostatin (e.g., for Duchenne muscular dystrophy), RNAi against VEGF (e.g., to treat and/or prevent tumors), RNAi

against phospholamban (e.g., to treat cardiovascular disease, see e.g., Andino et al. *J. Gene Med.* 10:132-142 (2008) and Li et al. *Acta Pharmacol Sin.* 26:51-55 (2005)); phospholamban inhibitory or dominant-negative molecules such as phospholamban S16E (e.g., to treat cardiovascular disease, see e.g., Hoshijima et al. *Nat. Med.* 8:864-871 (2002)), RNAi to adenosine kinase (e.g., for epilepsy), and RNAi directed against pathogenic organisms and viruses (e.g., hepatitis B and/or C virus, human immunodeficiency virus, CMV, herpes simplex virus, human papilloma virus, etc.).

[0119] Further, a nucleic acid sequence that directs alternative splicing can be delivered. To illustrate, an antisense sequence (or other inhibitory sequence) complementary to the 5' and/or 3' splice site of dystrophin exon 51 can be delivered in conjunction with a U1 or U7 small nuclear (sn) RNA promoter to induce skipping of this exon. For example, a DNA sequence comprising a U1 or U7 snRNA promoter located 5' to the antisense/inhibitory sequence(s) can be packaged and delivered in a modified capsid of the invention.

[0120] The virus vector may also comprise a heterologous nucleic acid that shares homology with and recombines with a locus on a host chromosome. This approach can be utilized, for example, to correct a genetic defect in the host cell.

[0121] The present invention also provides virus vectors that express an immunogenic polypeptide, e.g., for vaccination. The nucleic acid may encode any immunogen of interest known in the art including, but not limited to, immunogens from human immunodeficiency virus (HIV), simian immunodeficiency virus (SIV), influenza virus, HIV or SIV gag proteins, tumor antigens, cancer antigens, bacterial antigens, viral antigens, and the like.

[0122] The use of parvoviruses as vaccine vectors is known in the art (see, e.g., Miyamura et al. *Proc. Nat. Acad. Sci USA* 91:8507 (1994); U.S. Pat. No. 5,916,563 to Young et al., U.S. Pat. No. 5,905,040 to Mazzara et al., U.S. Pat. Nos. 5,882,652, 5,863,541 to Samulski et al.). The antigen may be presented in the parvovirus capsid. Alternatively, the antigen may be expressed from a heterologous nucleic acid introduced into a recombinant vector genome. Any immunogen of interest as described herein and/or as is known in the art can be provided by the virus vector of the present invention.

[0123] An immunogenic polypeptide can be any polypeptide suitable for eliciting an immune response and/or protecting the subject against an infection and/or disease, including, but not limited to, microbial, bacterial, protozoal, parasitic, fungal and/or viral infections and diseases. For example, the immunogenic polypeptide can be an orthomyxovirus immunogen (e.g., an influenza virus immunogen, such as the influenza virus hemagglutinin (HA) surface protein or the influenza virus nucleoprotein, or an equine influenza virus immunogen) or a lentivirus immunogen (e.g., an equine infectious anemia virus immunogen, a Simian Immunodeficiency Virus (SIV) immunogen, or a Human Immunodeficiency Virus (HIV) immunogen, such as the HIV or SIV envelope GP160 protein, the HIV or SIV matrix/capsid proteins, and the HIV or SIV gag, pol and env gene products). The immunogenic polypeptide can also be an arenavirus immunogen (e.g., Lassa fever virus immunogen, such as the Lassa fever virus nucleocapsid protein and/or the Lassa fever envelope glycoprotein), a poxvirus immunogen (e.g., a vaccinia virus immunogen, such as the

vaccinia L1 or L8 gene product), a flavivirus immunogen (e.g., a yellow fever virus immunogen or a Japanese encephalitis virus immunogen), a filovirus immunogen (e.g., an Ebola virus immunogen, or a Marburg virus immunogen, such as NP and GP gene products), a bunyavirus immunogen (e.g., RVFV, CCHF, and/or SFS virus immunogens), or a coronavirus immunogen (e.g., an infectious human coronavirus immunogen, such as the human coronavirus envelope glycoprotein, or a porcine transmissible gastroenteritis virus immunogen, or an avian infectious bronchitis virus immunogen). The immunogenic polypeptide can further be a polio immunogen, a herpesvirus immunogen (e.g., CMV, EBV, HSV immunogens) a mumps virus immunogen, a measles virus immunogen, a rubella virus immunogen, a diphtheria toxin or other diphtheria immunogen, a pertussis antigen, a hepatitis (e.g., hepatitis A, hepatitis B, hepatitis C, etc.) immunogen, and/or any other vaccine immunogen now known in the art or later identified as an immunogen.

[0124] Alternatively, the immunogenic polypeptide can be any tumor or cancer cell antigen. Optionally, the tumor or cancer antigen is expressed on the surface of the cancer cell. Exemplary cancer and tumor cell antigens are described in S.A. Rosenberg (Immunity 10:281 (1991)). Other illustrative cancer and tumor antigens include, but are not limited to: BRCA1 gene product, BRCA2 gene product, gp100, tyrosinase, GAGE-1/2, BAGE, RAGE, LAGE, NY-ESO-1, CDK-4, β -catenin, MUM-1, Caspase-8, KIAA0205, HPVE, SART-1, PRAME, p15, melanoma tumor antigens (Kawakami et al. *Proc. Natl. Acad. Sci. USA* 91:3515 (1994); Kawakami et al. *J. Exp. Med.*, 180:347 (1994); Kawakami et al. *Cancer Res.* 54:3124 (1994)), MART-1, gp100 MAGE-1, MAGE-2, MAGE-3, CEA, TRP-1, TRP-2, P-15, tyrosinase (Brichard et al. *J. Exp. Med.* 178:489 (1993)); HER-2/neu gene product (U.S. Pat. No. 4,968,603), CA125, LK26, FB5 (endosialin), TAG 72, AFP, CA19-9, NSE, DU-PAN-2, CA50, SPan-1, CA72-4, HCG, STN (sialyl Tn antigen), c-erbB-2 proteins, PSA, L-CanAg, estrogen receptor, milk fat globulin, p53 tumor suppressor protein (Levine. *Ann. Rev. Biochem.* 62:623 (1993)); mucin antigens (PCT Publication No. WO 90/05142); telomerases; nuclear matrix proteins; prostatic acid phosphatase; papilloma virus antigens; and/or antigens now known or later discovered to be associated with the following cancers: melanoma, adenocarcinoma, thymoma, lymphoma (e.g., non-Hodgkin's lymphoma, Hodgkin's lymphoma), sarcoma, lung cancer, liver cancer, colon cancer, leukemia, uterine cancer, breast cancer, prostate cancer, ovarian cancer, cervical cancer, bladder cancer, kidney cancer, pancreatic cancer, brain cancer and any other cancer or malignant condition now known or later identified (see, e.g., Rosenberg. *Ann. Rev. Med.* 47:481-91 (1996)).

[0125] As a further alternative, the heterologous nucleic acid can encode any polypeptide that is desirably produced in a cell in vitro, ex vivo, or in vivo. For example, the virus vectors may be introduced into cultured cells and the expressed nucleic acid product isolated therefrom.

[0126] It will be understood by those skilled in the art that the heterologous nucleic acid(s) of interest can be operably associated with appropriate control sequences. For example, the heterologous nucleic acid can be operably associated with expression control elements, such as transcription/translation control signals, origins of replication, polyadenylation signals, internal ribosome entry sites (IRES), promoters, and/or enhancers, and the like.

[0127] Further, regulated expression of the heterologous nucleic acid(s) of interest can be achieved at the post-transcriptional level, e.g., by regulating selective splicing of different introns by the presence or absence of an oligonucleotide, small molecule and/or other compound that selectively blocks splicing activity at specific sites (e.g., as described in PCT Publication No. WO 2006/119137).

[0128] Those skilled in the art will appreciate that a variety of promoter/enhancer elements can be used depending on the level and tissue-specific expression desired. The promoter/enhancer can be constitutive or inducible, depending on the pattern of expression desired. The promoter/enhancer can be native or foreign and can be a natural or a synthetic sequence. By foreign, it is intended that the transcriptional initiation region is not found in the wild-type host into which the transcriptional initiation region is introduced.

[0129] In particular embodiments, the promoter/enhancer elements can be native to the target cell or subject to be treated. In representative embodiments, the promoters/enhancer element can be native to the heterologous nucleic acid sequence. The promoter/enhancer element is generally chosen so that it functions in the target cell(s) of interest. Further, in particular embodiments the promoter/enhancer element is a mammalian promoter/enhancer element. The promoter/enhancer element may be constitutive or inducible.

[0130] Inducible expression control elements are typically advantageous in those applications in which it is desirable to provide regulation over expression of the heterologous nucleic acid sequence(s). Inducible promoters/enhancer elements for gene delivery can be tissue-specific or preferred promoter/enhancer elements, and include muscle specific or preferred (including cardiac, skeletal and/or smooth muscle specific or preferred), neural tissue specific or preferred (including brain-specific or preferred), eye specific or preferred (including retina-specific and cornea-specific), liver specific or preferred, bone marrow specific or preferred, pancreatic specific or preferred, spleen specific or preferred, and/or lung specific or preferred promoter/enhancer elements. Other inducible promoter/enhancer elements include hormone-inducible and metal-inducible elements. Exemplary inducible promoters/enhancer elements include, but are not limited to, a Tet on/off element, a RU486-inducible promoter, an ecdysone-inducible promoter, a rapamycin-inducible promoter, and a metallothionein promoter.

[0131] In embodiments wherein the heterologous nucleic acid sequence(s) is transcribed and then translated in the target cells, specific initiation signals are generally included for efficient translation of inserted protein coding sequences. These exogenous translational control sequences, which may include the ATG initiation codon and adjacent sequences, can be of a variety of origins, both natural and synthetic.

[0132] The virus vectors according to the present invention provide a means for delivering heterologous nucleic acids into a broad range of cells, including dividing and non-dividing cells. The virus vectors can be employed, for example, to deliver a nucleic acid of interest to a cell in vitro, e.g., to produce a polypeptide in vitro or for ex vivo gene therapy. The virus vectors are additionally useful in a method of delivering a nucleic acid to a subject in need thereof, e.g., to express an immunogenic and/or therapeutic polypeptide and/or a functional RNA. In this manner, the

polypeptide and/or functional RNA can be produced in vivo in the subject. The subject can be in need of the polypeptide because the subject has a deficiency of the polypeptide. Further, the method can be practiced because the production of the polypeptide and/or functional RNA in the subject may impart some beneficial effect.

[0133] The virus vectors can also be used to produce a polypeptide of interest and/or functional RNA in cultured cells or in a subject (e.g., using the subject as a bioreactor to produce the polypeptide and/or to observe the effects of the functional RNA on the subject, for example, in connection with screening methods).

[0134] In general, the virus vectors of the present invention can be employed to deliver a heterologous nucleic acid encoding a polypeptide and/or functional RNA (e.g., a therapeutic polypeptide, e.g., a therapeutic nucleic acid) to treat and/or prevent any disease state or disorder for which it is beneficial to deliver a therapeutic polypeptide and/or functional RNA, e.g., to a subject in need thereof, e.g., wherein the subject has or is at risk for a disease state or disorder. In some embodiments, the disease state is a CNS disease or disorder. In some embodiments, the subject has or is at risk of having pain associated with a disease or disorder. In some embodiments, the subject is a human. In some embodiments, the subject is in utero.

[0135] Illustrative disease states include, but are not limited to: cystic fibrosis (cystic fibrosis transmembrane regulator protein) and other diseases of the lung, hemophilia A (Factor VIII), hemophilia B (Factor IX), thalassemia (β -globin), anemia (erythropoietin) and other blood disorders, Alzheimer's disease (GDF; neprilysin), multiple sclerosis (β -interferon), Parkinson's disease (glial-cell line derived neurotrophic factor [GDNF]), Huntington's disease (RNAi to remove repeats), amyotrophic lateral sclerosis, epilepsy (galanin, neurotrophic factors), and other neurological disorders, cancer (endostatin, angiostatin, TRAIL, FAS-ligand, cytokines including interferons; RNAi including RNAi against VEGF or the multiple drug resistance gene product, mir-26a [e.g., for hepatocellular carcinoma]), diabetes mellitus (insulin), muscular dystrophies including Duchenne (dystrophin, mini-dystrophin, insulin-like growth factor I, a sarcoglycan [e.g., α , β , γ], RNAi against myostatin, myostatin propeptide, follistatin, activin type II soluble receptor, anti-inflammatory polypeptides such as the Ikappa B dominant mutant, sarcospan, utrophin, mini-utrophin, antisense or RNAi against splice junctions in the dystrophin gene to induce exon skipping [see e.g., PCT Publication No. WO/2003/095647], antisense against U7 snRNAs to induce exon skipping [see e.g., PCT Publication No. WO/2006/021724], and antibodies or antibody fragments against myostatin or myostatin propeptide) and Becker, Gaucher disease (glucocerebrosidase), Hurler's disease (α -L-iduronidase), adenosine deaminase deficiency (adenosine deaminase), glycogen storage diseases (e.g., Fabry disease [α -galactosidase] and Pompe disease [lysosomal acid α -glucosidase]) and other metabolic disorders, congenital emphysema (α 1-antitrypsin), Lesch-Nyhan Syndrome (hypoxanthine guanine phosphoribosyl transferase), Niemann-Pick disease (sphingomyelinase), Tay Sachs disease (lysosomal hexosaminidase A), Maple Syrup Urine Disease (branched-chain keto acid dehydrogenase), retinal degenerative diseases (and other diseases of the eye and retina; e.g., PDGF for macular degeneration and/or vasohibin or other inhibitors of VEGF or other angiogenesis

inhibitors to treat/prevent retinal disorders, e.g., in Type I diabetes), diseases of solid organs such as brain (including Parkinson's Disease [GDNF], astrocytomas [endostatin, angiostatin and/or RNAi against VEGF], glioblastomas [endostatin, angiostatin and/or RNAi against VEGF]), liver, kidney, heart including congestive heart failure or peripheral artery disease (PAD) (e.g., by delivering protein phosphatase inhibitor I (I-1) and fragments thereof (e.g., I1C), serca2a, zinc finger proteins that regulate the phospholamban gene, Barka, β 2-adrenergic receptor, β 2-adrenergic receptor kinase (BARK), phosphoinositide-3 kinase (PI3 kinase), S100A1, parvalbumin, adenylyl cyclase type 6, a molecule that effects G-protein coupled receptor kinase type 2 knock-down such as a truncated constitutively active bARKct; calsarcin, RNAi against phospholamban; phospholamban inhibitory or dominant-negative molecules such as phospholamban S16E, etc.), arthritis (insulin-like growth factors), joint disorders (insulin-like growth factor 1 and/or 2), intimal hyperplasia (e.g., by delivering enos, inos), improve survival of heart transplants (superoxide dismutase), AIDS (soluble CD4), muscle wasting (insulin-like growth factor I), kidney deficiency (erythropoietin), anemia (erythropoietin), arthritis (anti-inflammatory factors such as IRAP and TNF α soluble receptor), hepatitis (α -interferon), LDL receptor deficiency (LDL receptor), hyperammonemia (ornithine transcarbamylase), Krabbe's disease (galactocerebrosidase), Batten's disease, spinal cerebral ataxias including SCA1, SCA2 and SCA3, phenylketonuria (phenylalanine hydroxylase), autoimmune diseases, congenital neurodegenerative disorders (e.g., monogenic neurodegenerative disorders) such as mucopolysaccharidosis (including, but not limited to, Mucopolysaccharidosis Type I (also known as Hurler syndrome, Hurler-Scheie Syndrome, or Scheie syndrome, IDUA, alpha-L-iduronidase), Mucopolysaccharidosis Type II (also known as Hunter syndrome, IDS, I2L enzyme), Mucopolysaccharidosis Type III (also known as Sanfilippo syndrome, GNS [N-acetylglucosamine-6-sulfatase], HGSNAT [heparan-alpha-glucosaminide N-acetyltransferase], NAGLU [alpha-N-acetylglucosaminidase], and/or SGSH [sulfamidase]), Mucopolysaccharidosis Type IV (also known as Morquio syndrome, GALNS [galatosamine (N-acetyl)-6-sulfatase] and/or GLB1 [beta-galactosidase]), Mucopolysaccharidosis Type V (also known as Scheie syndrome, now a subgroup of type I, also IDUA, alpha-L-iduronidase), Mucopolysaccharidosis Type VI (also known as Maroteaux-Lamy syndrome, ARSB, arylsulfatase B), Mucopolysaccharidosis Type VII (also known as Sly syndrome, GUSB, beta-glucuronidase), Mucopolysaccharidosis Type IX (also known as Natowicz syndrome, HYAL1, hyaluronidase) and/or leukodystrophy (including, but not limited to, adult-onset autosomal dominant leukodystrophy (ADLD; LMNB1, lamin B1), Aicardi-Goutieres syndrome (TREX1, RNASEHSB, RNASEH2C, and/or RNASEH2A), Alexander disease (FRAP, glial fibrillary acidic protein), CADASIL (Notch3), Canavan disease (ASPA, aspartoacylase), CARASIL (HTRA1, serine protease HTRA1), cerebrotendinous xanthomatosis ("CTX," CYP27A1, sterol 27-hydroxylase) childhood ataxia and cerebral hypomyelination (CACH)/vanishing white matter disease (VWMD) (eIF2B, eukaryotic initiation factor 2B), Fabry disease (GLA, alpha-galactosidase A), fucosidosis (FUCA1, alpha-L-fucosidase), GM1 gangliosidosis (GLB1, beta-galactosidase), L-2-hydroxyglutaric aciduria (L2HDGH, L-2-hydroxyglutarate dehydrogenase), Krabbe disease (GALC,

galactocerebrosidase), megalencephalic leukoencephalopathy with subcortical cysts (“MLC,” MLC1 and/or HEP-ACAM), metachromatic leukodystrophy (ASA, arylsulfatase A), multiple sulfatase deficiency (“MSD,” SUMF1, sulfatase modifying factor 1 affecting all sulfatase enzymes), Pelizaeus-Merzbacher disease (also known as “X-linked spastic paraplegia,” PLP1 [X-linked proteolipid protein 1] and/or GJA12 [gap junction protein 12]), Pol III-Related Leukodystrophies (POLR3A and/or POLR3B), Refsum disease (PHYH, [phytanoyl-CoA hydroxylase] and/or Pex7 [PHYH importer into peroxisomes]), salla disease (also known as “free sialic acid storage disease,” SLC17A5, sialic acid transporter), Sjogren-Larsen syndrome (ALDH3A2, aldehyde dehydrogenase), X-linked adrenoleukodystrophy (“ALD,” ABCD1, peroxisomal ATPase Binding Cassette protein), Zellweger syndrome spectrum disorders (also known as peroxisomal biogenesis disorders, PEX1, PEX2, PEX3, PEX4, PEX5, PEX10, PEX11B, PEX12, PEX13, PEX14, PEX16, PEX19, PEX26), and the like. The invention can further be used following organ transplantation to increase the success of the transplant and/or to reduce the negative side effects of organ transplantation or adjunct therapies (e.g., by administering immunosuppressant agents or inhibitory nucleic acids to block cytokine production). As another example, bone morphogenic proteins (including BNP 2, 7, etc., RANKL and/or VEGF) can be administered with a bone allograft, for example, following a break or surgical removal in a cancer patient.

[0136] Thus, in some embodiments, the present invention provides a method of treating a disease in a subject in need thereof, comprising introducing a therapeutic nucleic acid into a cell of the subject by administering to the subject the virus vector and/or composition of the present invention, under conditions whereby the therapeutic nucleic acid is expressed in the subject.

[0137] The invention can also be used to produce induced pluripotent stem cells (iPS). For example, a virus vector of the invention can be used to deliver stem cell associated nucleic acid(s) into a non-pluripotent cell, such as adult fibroblasts, skin cells, liver cells, renal cells, adipose cells, cardiac cells, neural cells, epithelial cells, endothelial cells, and the like. Nucleic acids encoding factors associated with stem cells are known in the art. Nonlimiting examples of such factors associated with stem cells and pluripotency include Oct-3/4, the SOX family (e.g., SOX1, SOX2, SOX3 and/or SOX15), the Klf family (e.g., Klf1, Klf2, Klf4 and/or Klf5), the Myc family (e.g., C-myc, L-myc and/or N-myc), NANOG and/or LIN28.

[0138] The invention can also be practiced to treat and/or prevent a metabolic disorder such as diabetes (e.g., insulin), hemophilia (e.g., Factor IX or Factor VIII), a lysosomal storage disorder such as a mucopolysaccharidosis disorder (e.g., Sly syndrome [β -glucuronidase], Hurler Syndrome [α -L-iduronidase], Scheie Syndrome [α -L-iduronidase], Hurler-Scheie Syndrome [α -L-iduronidase], Hunter’s Syndrome [iduronate sulfatase], Sanfilippo Syndrome A [heparan sulfamidase], B [N-acetylglucosaminidase], C [acetyl-CoA: α -glucosaminide acetyltransferase], D [N-acetylglucosamine 6-sulfatase], Morquio Syndrome A [galactose-6-sulfate sulfatase], B [β -galactosidase], Maroteaux-Lamy Syndrome [N-acetylgalactosamine-4-sulfatase], etc.), Fabry disease (α -galactosidase), Gaucher’s disease (glucocerebrosidase), or a glycogen storage disorder (e.g., Pompe disease; lysosomal acid α -glucosidase).

[0139] Gene transfer has substantial potential use for understanding and providing therapy for disease states. There are a number of inherited diseases in which defective genes are known and have been cloned. In general, the above disease states fall into two classes: deficiency states, usually of enzymes, which are generally inherited in a recessive manner, and unbalanced states, which may involve regulatory or structural proteins, and which are typically inherited in a dominant manner. For deficiency state diseases, gene transfer can be used to bring a normal gene into affected tissues for replacement therapy, as well as to create animal models for the disease using antisense mutations. For unbalanced disease states, gene transfer can be used to create a disease state in a model system, which can then be used in efforts to counteract the disease state. Thus, virus vectors according to the present invention permit the treatment and/or prevention of genetic diseases.

[0140] The virus vectors comprising a modified AAV2.5 capsid protein (e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5) according to the present invention may also be employed to provide a functional RNA to a cell in vitro or in vivo. Expression of the functional RNA in the cell, for example, can diminish expression of a particular target protein by the cell. Accordingly, functional RNA can be administered to decrease expression of a particular protein in a subject in need thereof. Functional RNA can also be administered to cells in vitro to regulate gene expression and/or cell physiology, e.g., to optimize cell or tissue culture systems or in screening methods.

[0141] In addition, virus vectors comprising a modified AAV2.5 capsid protein (e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5) according to the instant invention find use in diagnostic and screening methods, whereby a nucleic acid of interest is transiently or stably expressed in a cell culture system, or alternatively, a transgenic animal model.

[0142] The virus vectors of the present invention can also be used for various non-therapeutic purposes, including but not limited to use in protocols to assess gene targeting, clearance, transcription, translation, etc., as would be apparent to one skilled in the art. The virus vectors can also be used for the purpose of evaluating safety (spread, toxicity, immunogenicity, etc.). Such data, for example, are considered by the United States Food and Drug Administration as part of the regulatory approval process prior to evaluation of clinical efficacy.

[0143] As a further aspect, the virus vectors of the present invention may be used to produce an immune response in a subject. According to this embodiment, a virus vector comprising a heterologous nucleic acid sequence encoding an immunogenic polypeptide can be administered to a subject, and an active immune response is mounted by the subject against the immunogenic polypeptide. Immunogenic polypeptides are as described hereinabove. In some embodiments, a protective immune response is elicited.

[0144] Alternatively, the virus vector may be administered to a cell ex vivo and the altered cell is administered to the subject. The virus vector comprising the heterologous

nucleic acid is introduced into the cell, and the cell is administered to the subject, where the heterologous nucleic acid encoding the immunogen can be expressed and induce an immune response in the subject against the immunogen. In particular embodiments, the cell is an antigen-presenting cell (e.g., a dendritic cell).

[0145] An “active immune response” or “active immunity” is characterized by “participation of host tissues and cells after an encounter with the immunogen. It involves differentiation and proliferation of immunocompetent cells in lymphoreticular tissues, which lead to synthesis of antibody or the development of cell-mediated reactivity, or both.” Herbert B. Herscovitz. *Immunophysiology: Cell Function and Cellular Interactions in Antibody Formation*, in IMMUNOLOGY: BASIC PROCESSES 117 (Joseph A. Bellanti ed., 1985). Alternatively stated, an active immune response is mounted by the host after exposure to an immunogen by infection or by vaccination. Active immunity can be contrasted with passive immunity, which is acquired through the “transfer of preformed substances (antibody, transfer factor, thymic graft, interleukin-2) from an actively immunized host to a non-immune host.” Id.

[0146] A “protective” immune response or “protective” immunity as used herein indicates that the immune response confers some benefit to the subject in that it prevents or reduces the incidence of disease. Alternatively, a protective immune response or protective immunity may be useful in the treatment and/or prevention of disease, in particular cancer or tumors (e.g., by preventing cancer or tumor formation, by causing regression of a cancer or tumor and/or by preventing metastasis and/or by preventing growth of metastatic nodules). The protective effects may be complete or partial, as long as the benefits of the treatment outweigh any disadvantages thereof.

[0147] In particular embodiments, the virus vector or cell comprising the heterologous nucleic acid can be administered in an immunogenically effective amount, as described herein.

[0148] The virus vectors of the present invention can also be administered for cancer immunotherapy by administration of a virus vector expressing one or more cancer cell antigens (or an immunologically similar molecule) or any other immunogen that produces an immune response against a cancer cell. To illustrate, an immune response can be produced against a cancer cell antigen in a subject by administering a virus vector comprising a heterologous nucleic acid encoding the cancer cell antigen, for example to treat a patient with cancer and/or to prevent cancer from developing in the subject. The virus vector may be administered to a subject in vivo or by using ex vivo methods, as described herein. Alternatively, the cancer antigen can be expressed as part of the virus capsid or be otherwise associated with the virus capsid (e.g., as described above).

[0149] As another alternative, any other therapeutic nucleic acid (e.g., RNAi) or polypeptide (e.g., cytokine) known in the art can be administered to treat and/or prevent cancer.

[0150] As used herein, the term “cancer” encompasses tumor-forming cancers. Likewise, the term “cancerous tissue” encompasses tumors. A “cancer cell antigen” encompasses tumor antigens.

[0151] The term “cancer” has its understood meaning in the art, for example, an uncontrolled growth of tissue that has the potential to spread to distant sites of the body (i.e.,

metastasize). Exemplary cancers include, but are not limited to melanoma, adenocarcinoma, thymoma, lymphoma (e.g., non-Hodgkin’s lymphoma, Hodgkin’s lymphoma), sarcoma, lung cancer, liver cancer, colon cancer, leukemia, uterine cancer, breast cancer, prostate cancer, ovarian cancer, cervical cancer, bladder cancer, kidney cancer, pancreatic cancer, brain cancer and any other cancer or malignant condition now known or later identified. In representative embodiments, the invention provides a method of treating and/or preventing tumor-forming cancers.

[0152] The term “tumor” is also understood in the art, for example, as an abnormal mass of undifferentiated cells within a multicellular organism. Tumors can be malignant or benign. In representative embodiments, the methods disclosed herein are used to prevent and treat malignant tumors.

[0153] By the terms “treating cancer,” “treatment of cancer” and equivalent terms it is intended that the severity of the cancer is reduced or at least partially eliminated and/or the progression of the disease is slowed and/or controlled and/or the disease is stabilized. In particular embodiments, these terms indicate that metastasis of the cancer is prevented or reduced or at least partially eliminated and/or that growth of metastatic nodules is prevented or reduced or at least partially eliminated.

[0154] By the terms “prevention of cancer” or “preventing cancer” and equivalent terms it is intended that the methods at least partially eliminate or reduce and/or delay the incidence and/or severity of the onset of cancer. Alternatively stated, the onset of cancer in the subject may be reduced in likelihood or probability and/or delayed.

[0155] In particular embodiments, cells may be removed from a subject with cancer and contacted with a virus vector expressing a cancer cell antigen according to the instant invention. The modified cell is then administered to the subject, whereby an immune response against the cancer cell antigen is elicited. This method can be advantageously employed with immunocompromised subjects that cannot mount a sufficient immune response in vivo (i.e., cannot produce enhancing antibodies in sufficient quantities).

[0156] It is known in the art that immune responses may be enhanced by immunomodulatory cytokines (e.g., α -interferon, β -interferon, γ -interferon, ω -interferon, τ -interferon, interleukin-1 α , interleukin-1 β , interleukin-2, interleukin-3, interleukin-4, interleukin 5, interleukin-6, interleukin-7, interleukin-8, interleukin-9, interleukin-10, interleukin-11, interleukin 12, interleukin-13, interleukin-14, interleukin-18, B cell Growth factor, CD40 Ligand, tumor necrosis factor- α , tumor necrosis factor- β , monocyte chemoattractant protein-1, granulocyte-macrophage colony stimulating factor, and lymphotoxin). Accordingly, immunomodulatory cytokines (preferably, CTL inductive cytokines) may be administered to a subject in conjunction with the virus vector.

[0157] Cytokines may be administered by any method known in the art. Exogenous cytokines may be administered to the subject, or alternatively, a nucleic acid encoding a cytokine may be delivered to the subject using a suitable vector, and the cytokine produced in vivo.

Subjects, Pharmaceutical Formulations, and Modes of Administration

[0158] Virus vectors and capsids comprising a modified AAV2.5 capsid protein (comprising a modified AAV2.5 capsid protein (e.g., wherein the modified AAV2.5 capsid

protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5)) according to the present invention find use in both veterinary and medical applications. Suitable subjects include both avians and mammals. The term “avian” as used herein includes, but is not limited to, chickens, ducks, geese, quail, turkeys, pheasant, parrots, parakeets, and the like. The term “mammal” as used herein includes, but is not limited to, humans, non-human primates, bovines, ovines, caprines, equines, felines, canines, lagomorphs, etc. A subject can be a fully developed subject (e.g., an adult) or a subject undergoing the developmental process (e.g., a child, infant or fetus). Human subjects include in utero (e.g., embryos, fetuses), neonates, infants, juveniles, adults and geriatric subjects.

[0159] In representative embodiments, the subject is “in need of” the methods of the invention and thus in some embodiments can be a “subject in need thereof.”

[0160] In particular embodiments, the present invention provides a pharmaceutical composition comprising a virus vector and/or capsid comprising a modified AAV2.5 capsid protein (comprising a modified AAV2.5 capsid protein (e.g., wherein the modified AAV2.5 capsid protein does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, and/or does not contain a serine substitution at the position corresponding to amino acid 267 of AAV2.5)) of the invention in a pharmaceutically acceptable carrier and, optionally, other medicinal agents, pharmaceutical agents, stabilizing agents, buffers, carriers, adjuvants, diluents, etc. For injection, the carrier will typically be a liquid. For other methods of administration, the carrier may be either solid or liquid. For inhalation administration, the carrier will be respirable, and optionally can be in solid or liquid particulate form.

[0161] By “pharmaceutically acceptable” it is meant a material that is not toxic or otherwise undesirable, i.e., the material may be administered to a subject without causing any undesirable biological effects.

[0162] One aspect of the present invention is a method of transferring a nucleic acid to a cell in vitro. The virus vector may be introduced into the cells at the appropriate multiplicity of infection according to standard transduction methods suitable for the particular target cells. Titers of virus vector to administer can vary, depending upon the target cell type and number, and the particular virus vector, and can be determined by those of skill in the art without undue experimentation. In representative embodiments, at least about 10^3 infectious units, optionally at least about 10^5 infectious units are introduced to the cell.

[0163] The cell(s) into which the virus vector is introduced can be of any type, including but not limited to neural cells (including cells of the peripheral and central nervous systems, in particular, brain cells such as neurons and glial cells such as astrocytes and oligodendrocytes), lung cells, cells of the eye (including retinal cells, retinal pigment epithelium, and corneal cells), epithelial cells (e.g., gut and respiratory epithelial cells), muscle cells (e.g., skeletal muscle cells, cardiac muscle cells, smooth muscle cells and/or diaphragm muscle cells), dendritic cells, pancreatic cells (including islet cells), hepatic cells, myocardial cells, bone cells (e.g., bone marrow stem cells), hematopoietic stem cells, spleen cells, keratinocytes, fibroblasts, endothelial cells, prostate cells, germ cells, and the like. In representative embodiments, the

cell can be any progenitor cell. As a further embodiment, the cell can be a stem cell (e.g., neural stem cell, liver stem cell). As still a further embodiment, the cell can be a cancer or tumor cell. Moreover, the cell can be from any species of origin, as indicated above.

[0164] The virus vector can be introduced into cells in vitro for the purpose of administering the modified cell to a subject. In particular embodiments, the cells have been removed from a subject, the virus vector is introduced therein, and the cells are then administered back into the subject. Methods of removing cells from subject for manipulation ex vivo, followed by introduction back into the subject are known in the art (see, e.g., U.S. Pat. No. 5,399,346). Alternatively, the recombinant virus vector can be introduced into cells from a donor subject, into cultured cells, or into cells from any other suitable source, and the cells are administered to a subject in need thereof (i.e., a “recipient” subject).

[0165] Suitable cells for ex vivo nucleic acid delivery are as described above. Dosages of the cells to administer to a subject will vary upon the age, condition and species of the subject, the type of cell, the nucleic acid being expressed by the cell, the mode of administration, and the like. Typically, at least about 10^2 to about 10^8 cells or at least about 10^3 to about 10^6 cells will be administered per dose in a pharmaceutically acceptable carrier. In particular embodiments, the cells transduced with the virus vector are administered to the subject in a treatment effective or prevention effective amount in combination with a pharmaceutical carrier.

[0166] In some embodiments, the virus vector is introduced into a cell and the cell can be administered to a subject to elicit an immunogenic response against the delivered polypeptide (e.g., expressed as a transgene or in the capsid). Typically, a quantity of cells expressing an immunogenically effective amount of the polypeptide in combination with a pharmaceutically acceptable carrier is administered. An “immunogenically effective amount” is an amount of the expressed polypeptide that is sufficient to evoke an active immune response against the polypeptide in the subject to which the pharmaceutical formulation is administered. In particular embodiments, the dosage is sufficient to produce a protective immune response (as defined above). The degree of protection conferred need not be complete or permanent, as long as the benefits of administering the immunogenic polypeptide outweigh any disadvantages thereof.

[0167] In some embodiments, the subject may have a reduced immunologic profile (e.g., immunologic response, e.g., antigenic cross-reactivity) when contacted with a virus vector of the present invention as compared to a control, e.g., when contacted with another AAV virus vector (e.g., AAV1, AAV2, AAV9, AAV2.5, or any AAV serotype listed in Table 1).

[0168] A further aspect of the invention is a method of administering the virus vector and/or virus capsid to a subject. Administration of the virus vectors and/or capsids according to the present invention to a human subject or an animal in need thereof can be by any means known in the art. Optionally, the virus vector and/or capsid can be delivered in a treatment effective or prevention effective dose in a pharmaceutically acceptable carrier.

[0169] The virus vectors and/or capsids of the invention can further be administered to elicit an immunogenic response (e.g., as a vaccine). Typically, immunogenic com-

positions of the present invention comprise an immunogenically effective amount of virus vector and/or capsid in combination with a pharmaceutically acceptable carrier. Optionally, the dosage is sufficient to produce a protective immune response (as defined above).

[0170] Dosages of the virus vector and/or capsid to be administered to a subject depend upon the mode of administration, the disease or condition to be treated and/or prevented, the individual subject's condition, the particular virus vector or capsid, the nucleic acid to be delivered, and the like, and can be determined in a routine manner. Exemplary doses for achieving therapeutic effects are titers of at least about 10^5 , 10^6 , 10^7 , 10^8 , 10^9 , 10^{10} , 10^{11} , 10^{12} , 10^3 , 10^{14} , 10^{15} transducing units, optionally about 10^8 - 10^{13} transducing units.

[0171] In particular embodiments, more than one administration (e.g., two, three, four or more administrations) may be employed to achieve the desired level of gene expression over a period of various intervals, e.g., daily, weekly, monthly, yearly, etc.

[0172] "Repeated administration", "repeat dose" or "repeat dosing" or the like means at least one additional dose or dosing that is administered to a subject subsequent to an earlier dose or dosing of the same or similar material. For example, a repeated dose of a viral vector and/or therapeutic is at least one additional dose of the viral vector and/or therapeutic after a prior dose of the same material. While the material may be the same, the amount of the material in the repeated dose may be different from the earlier dose. For example, in an embodiment of any one of the methods or compositions provided herein, the amount of the viral vector and/or therapeutic in the repeated dose may be less than the amount of the viral vector and/or therapeutic of the earlier dose. Alternatively, in an embodiment of any one of the methods or compositions provided herein, the repeated dose may be in an amount that is at least equal to the amount of the viral vector and/or therapeutic in the earlier dose. A repeat dose may be administered weeks, months or years after the prior dose. In some embodiments of any one of the methods provided herein, the repeat dose or dosing is administered at least 1 week after the dose or dosing that occurred just prior to the repeat dose or dosing. In some embodiments of any one of the methods provided herein, the repeat dose or dosing is administered at least 1 month after the dose or dosing that occurred just prior to the repeat dose or dosing. Repeat dosing is considered to be efficacious if it results in a beneficial effect for the subject. Preferably, efficacious repeat dosing results in a beneficial effect in conjunction with reduced immune response, such as to the viral vector and/or to the encoded transgene.

[0173] Exemplary modes of administration include oral, rectal, transmucosal, intranasal, inhalation (e.g., via an aerosol), buccal (e.g., sublingual), vaginal, intrathecal, intraocular, transdermal, in utero (or in ovo), parenteral (e.g., intravenous, subcutaneous, intradermal, intramuscular [including administration to skeletal, diaphragm and/or cardiac muscle], intradermal, intrapleural, intracerebral, and intraarticular), topical (e.g., to both skin and mucosal surfaces, including airway surfaces, and transdermal administration), intralymphatic, and the like, as well as direct tissue or organ injection (e.g., to liver, skeletal muscle, cardiac muscle, diaphragm muscle or brain). Administration can also be to a tumor (e.g., in or near a tumor or a lymph node). The most suitable route in any given case will depend on the nature

and severity of the condition being treated and/or prevented and on the nature of the particular vector that is being used.

[0174] Administration to skeletal muscle according to the present invention includes but is not limited to administration to skeletal muscle in the limbs (e.g., upper arm, lower arm, upper leg, and/or lower leg), back, neck, head (e.g., tongue), thorax, abdomen, pelvis/perineum, and/or digits. Suitable skeletal muscles include but are not limited to abductor digiti minimi (in the hand), abductor digiti minimi (in the foot), abductor hallucis, abductor ossis metatarsi quinti, abductor pollicis *brevis*, abductor pollicis longus, adductor *brevis*, adductor hallucis, adductor longus, adductor magnus, adductor pollicis, anconeus, anterior scalene, articularis genus, biceps brachii, biceps femoris, *brachialis*, brachioradialis, buccinator, coracobrachialis, corrugator supercilii, deltoid, depressor anguli oris, depressor labii inferioris, digastric, dorsal interossei (in the hand), dorsal interossei (in the foot), extensor carpi radialis *brevis*, extensor carpi radialis longus, extensor carpi ulnaris, extensor digiti minimi, extensor digitorum, extensor digitorum *brevis*, extensor digitorum longus, extensor hallucis *brevis*, extensor hallucis longus, extensor indicis, extensor pollicis *brevis*, extensor pollicis longus, flexor carpi radialis, flexor carpi ulnaris, flexor digiti minimi *brevis* (in the hand), flexor digiti minimi *brevis* (in the foot), flexor digitorum *brevis*, flexor digitorum longus, flexor digitorum *profundus*, flexor digitorum superficialis, flexor hallucis *brevis*, flexor hallucis longus, flexor pollicis *brevis*, flexor pollicis longus, *frontalis*, gastrocnemius, geniohyoid, gluteus maximus, gluteus medius, gluteus minimus, gracilis, iliocostalis cervicis, iliocostalis lumborum, iliocostalis thoracis, iliocostalis, inferior gemellus, inferior oblique, inferior rectus, infrapinnatus, interspinalis, intertransversi, lateral pterygoid, lateral rectus, latissimus dorsi, levator anguli oris, levator labii superioris, levator labii superioris alaeque nasi, levator palpebrae superioris, levator scapulae, long rotators, longissimus capitis, longissimus cervicis, longissimus thoracis, longus capitis, longus colli, lumbricals (in the hand), lumbricals (in the foot), masseter, medial pterygoid, medial rectus, middle scalene, multifidus, mylohyoid, obliquus capitis inferior, obliquus capitis superior, obturator externus, obturator internus, occipitalis, omohyoid, opponens digiti minimi, opponens pollicis, orbicularis oculi, orbicularis oris, palmar interossei, palmaris *brevis*, palmaris longus, pectineus, pectoralis major, pectoralis minor, peroneus *brevis*, peroneus longus, peroneus tertius, *piriformis*, plantar interossei, plantaris, platysma, popliteus, posterior scalene, pronator quadratus, pronator *teres*, psoas major, quadratus femoris, quadratus plantae, rectus capitis anterior, rectus capitis lateralis, rectus capitis posterior major, rectus capitis posterior minor, rectus femoris, rhomboid major, rhomboid minor, risorius, sartorius, scalenus minimus, semimembranosus, semispinalis capitis, semispinalis cervicis, semispinalis thoracis, semitendinosus, serratus anterior, short rotators, soleus, spinalis capitis, spinalis cervicis, spinalis thoracis, splenius capitis, splenius cervicis, sternocleidomastoid, sternohyoid, sternothyroid, stylohyoid, subclavius, subscapularis, superior gemellus, superior oblique, superior rectus, supinator, supraspinatus, temporalis, tensor fascia lata, *Teres major*, *Teres minor*, thoracis, thyrohyoid, tibialis anterior, tibialis posterior, trapezius, *Triceps brachii*, *Vastus intermedius*, *Vastus lateralis*, *Vastus medialis*, *Zygomaticus major*, and *Zygomaticus minor*, and any other suitable skeletal muscle as known in the art.

[0175] The virus vector and/or capsid can be delivered to skeletal muscle by intravenous administration, intra-arterial administration, intraperitoneal administration, limb perfusion, (optionally, isolated limb perfusion of a leg and/or arm; see e.g., Arruda et al. (2005) *Blood* 105:3458-3464), and/or direct intramuscular injection. In particular embodiments, the virus vector and/or capsid is administered to a limb (arm and/or leg) of a subject (e.g., a subject with muscular dystrophy such as DMD) by limb perfusion, optionally isolated limb perfusion (e.g., by intravenous or intra-articular administration). In embodiments of the invention, the virus vectors and/or capsids of the invention can advantageously be administered without employing “hydrodynamic” techniques. Tissue delivery (e.g., to muscle) of vectors is often enhanced by hydrodynamic techniques (e.g., intravenous/intravenous administration in a large volume), which increase pressure in the vasculature and facilitate the ability of the vector to cross the endothelial cell barrier. In particular embodiments, the viral vectors and/or capsids of the invention can be administered in the absence of hydrodynamic techniques such as high volume infusions and/or elevated intravascular pressure (e.g., greater than normal systolic pressure, for example, less than or equal to a 5%, 10%, 15%, 20%, 25% increase in intravascular pressure over normal systolic pressure). Such methods may reduce or avoid the side effects associated with hydrodynamic techniques such as edema, nerve damage and/or compartment syndrome.

[0176] Administration to cardiac muscle includes administration to the left atrium, right atrium, left ventricle, right ventricle and/or septum. The virus vector and/or capsid can be delivered to cardiac muscle by intravenous administration, intra-arterial administration such as intra-aortic administration, direct cardiac injection (e.g., into left atrium, right atrium, left ventricle, right ventricle), and/or coronary artery perfusion.

[0177] Administration to diaphragm muscle can be by any suitable method including intravenous administration, intra-arterial administration, and/or intra-peritoneal administration.

[0178] Delivery to a target tissue can also be achieved by delivering a depot comprising the virus vector and/or capsid. In representative embodiments, a depot comprising the virus vector and/or capsid is implanted into skeletal, cardiac and/or diaphragm muscle tissue or the tissue can be contacted with a film or other matrix comprising the virus vector and/or capsid. Such implantable matrices or substrates are described, e.g., in U.S. Pat. No. 7,201,898.

[0179] In particular embodiments, a virus vector and/or virus capsid according to the present invention is administered to skeletal muscle, diaphragm muscle and/or cardiac muscle (e.g., to treat and/or prevent muscular dystrophy, heart disease [for example, PAD or congestive heart failure]).

[0180] In representative embodiments, the invention is used to treat and/or prevent disorders of skeletal, cardiac and/or diaphragm muscle.

[0181] In a representative embodiment, the invention provides a method of treating and/or preventing muscular dystrophy in a subject in need thereof, the method comprising: administering a treatment or prevention effective amount of a virus vector of the invention to a mammalian subject, wherein the virus vector comprises a heterologous nucleic acid encoding dystrophin, a mini-dystrophin, a

micro-dystrophin, myostatin propeptide, follistatin, activin type II soluble receptor, IGF-1, anti-inflammatory polypeptides such as the Ikappa B dominant mutant, sarcospan, utrophin, a micro-dystrophin, laminin- α 2, α -sarcoglycan, β -sarcoglycan, γ -sarcoglycan, δ -sarcoglycan, IGF-1, an antibody or antibody fragment against myostatin or myostatin propeptide, and/or RNAi against myostatin. In particular embodiments, the virus vector can be administered to skeletal, diaphragm and/or cardiac muscle as described elsewhere herein.

[0182] Alternatively, the invention can be practiced to deliver a nucleic acid to skeletal, cardiac or diaphragm muscle, which is used as a platform for production of a polypeptide (e.g., an enzyme) or functional RNA (e.g., RNAi, microRNA, antisense RNA) that normally circulates in the blood or for systemic delivery to other tissues to treat and/or prevent a disorder (e.g., a metabolic disorder, such as diabetes [e.g., insulin], hemophilia [e.g., Factor IX or Factor VIII], a mucopolysaccharide disorder [e.g., Sly syndrome, Hurler Syndrome, Scheie Syndrome, Hurler-Scheie Syndrome, Hunter’s Syndrome, Sanfilippo Syndrome A, B, C, D, Morquio Syndrome, Maroteaux-Lamy Syndrome, etc.] or a lysosomal storage disorder such as Gaucher’s disease [glucocerebrosidase] or Fabry disease [α -galactosidase A] or a glycogen storage disorder such as Pompe disease [lysosomal acid α glucosidase]). Other suitable proteins for treating and/or preventing metabolic disorders are described herein. The use of muscle as a platform to express a nucleic acid of interest is described in U.S. Patent Publication No. 20020192189.

[0183] Thus, as one aspect, the invention further encompasses a method of treating and/or preventing a metabolic disorder in a subject in need thereof, the method comprising: administering a treatment or prevention effective amount of a virus vector of the invention to skeletal muscle of a subject, wherein the virus vector comprises a heterologous nucleic acid encoding a polypeptide, wherein the metabolic disorder is a result of a deficiency and/or defect in the polypeptide. Illustrative metabolic disorders and heterologous nucleic acids encoding polypeptides are described herein. Optionally, the polypeptide is secreted (e.g., a polypeptide that is a secreted polypeptide in its native state or that has been engineered to be secreted, for example, by operable association with a secretory signal sequence as is known in the art). Without being limited by any particular theory of the invention, according to this embodiment, administration to the skeletal muscle can result in secretion of the polypeptide into the systemic circulation and delivery to target tissue(s). Methods of delivering virus vectors to skeletal muscle are described in more detail herein.

[0184] The invention can also be practiced to produce antisense RNA, RNAi or other functional RNA (e.g., a ribozyme) for systemic delivery.

[0185] The invention also provides a method of treating and/or preventing congenital heart failure or PAD in a subject in need thereof, the method comprising administering a treatment or prevention effective amount of a virus vector of the invention to a mammalian subject, wherein the virus vector comprises a heterologous nucleic acid encoding, for example, a sarcoplasmic endoreticulum Ca^{2+} -ATPase (SERCA2a), an angiogenic factor, phosphatase inhibitor I (I-1) and fragments thereof (e.g., I1C), RNAi against phospholamban; a phospholamban inhibitory or dominant-negative molecule such as phospholamban S 16E, a zinc

finger protein that regulates the phospholamban gene, β 2-adrenergic receptor, β 2-adrenergic receptor kinase (BARK), PI3 kinase, calsarcin, a β -adrenergic receptor kinase inhibitor (β ARKct), inhibitor 1 of protein phosphatase 1 and fragments thereof (e.g., I1C), S100A1, parvalbumin, adenylyl cyclase type 6, a molecule that effects G-protein coupled receptor kinase type 2 knockdown such as a truncated constitutively active β ARKct, Pim-1, PGC-1 α , SOD-1, SOD-2, EC-SOD, kallikrein, HIF, thymosin- β 4, mir-1, mir-133, mir-206, mir-208 and/or mir-26a.

[0186] In some embodiments, the invention further encompasses a method of treating and/or preventing a congenital neurodegenerative disorder (e.g., monogenic neurodegenerative disorder) in a subject in need thereof, the method comprising: administering a treatment or prevention effective amount of a virus vector of the invention to neural tissue (e.g., neuronal cells) of a subject, wherein the virus vector comprises a heterologous nucleic acid encoding a polypeptide, wherein the congenital neurodegenerative disorder is a result of a deficiency and/or defect in the polypeptide. Illustrative congenital neurodegenerative disorders and heterologous nucleic acids encoding polypeptides are described herein. Optionally, the polypeptide is secreted (e.g., a polypeptide that is a secreted polypeptide in its native state or that has been engineered to be secreted, for example, by operable association with a secretory signal sequence as is known in the art). In some embodiments, the subject is a human. In some embodiments, the subject is in utero. In some embodiments, the subject has or is at risk for a congenital (e.g., monogenic) neurodegenerative disorder. In some embodiments, the subject has or is at risk for mucopolysaccharidosis or leukodystrophy.

[0187] Injectables can be prepared in conventional forms, either as liquid solutions or suspensions, solid forms suitable for solution or suspension in liquid prior to injection, or as emulsions. Alternatively, one may administer the virus vector and/or virus capsids of the invention in a local rather than systemic manner, for example, in a depot or sustained-release formulation. Further, the virus vector and/or virus capsid can be delivered adhered to a surgically implantable matrix (e.g., as described in U.S. Patent Publication No. 20040013645).

[0188] The virus vectors and/or virus capsids disclosed herein can be administered to the lungs of a subject by any suitable means, optionally by administering an aerosol suspension of respirable particles comprised of the virus vectors and/or virus capsids, which the subject inhales. The respirable particles can be liquid or solid. Aerosols of liquid particles comprising the virus vectors and/or virus capsids may be produced by any suitable means, such as with a pressure-driven aerosol nebulizer or an ultrasonic nebulizer, as is known to those of skill in the art. See e.g., U.S. Pat. No. 4,501,729. Aerosols of solid particles comprising the virus vectors and/or capsids may likewise be produced with any solid particulate medicament aerosol generator, by techniques known in the pharmaceutical art.

[0189] The virus vectors and virus capsids can be administered to tissues of the central nervous system (CNS) (e.g., brain, eye) and may advantageously result in broader distribution of the virus vector or capsid than would be observed in the absence of the present invention.

[0190] In particular embodiments, the delivery vectors of the invention may be administered to treat diseases of the CNS, including genetic disorders, neurodegenerative disorders,

psychiatric disorders and tumors. Illustrative diseases of the CNS include, but are not limited to Alzheimer's disease, Parkinson's disease, Huntington's disease, Canavan disease, Leigh's disease, Refsum disease, Tourette syndrome, primary lateral sclerosis, amyotrophic lateral sclerosis, progressive muscular atrophy, Pick's disease, muscular dystrophy, multiple sclerosis, myasthenia gravis, Binswanger's disease, trauma due to spinal cord or head injury, Tay Sachs disease, Lesch-Nyan disease, epilepsy, cerebral infarcts, psychiatric disorders including mood disorders (e.g., depression, bipolar affective disorder, persistent affective disorder, secondary mood disorder), schizophrenia, drug dependency (e.g., alcoholism and other substance dependencies), neuroses (e.g., anxiety, obsessional disorder, somatoform disorder, dissociative disorder, grief, post-partum depression), psychosis (e.g., hallucinations and delusions), dementia, paranoia, attention deficit disorder, psychosexual disorders, sleeping disorders, pain disorders, eating or weight disorders (e.g., obesity, cachexia, anorexia nervosa, and bulimia) cancers and tumors (e.g., pituitary tumors) of the CNS, and congenital neurodegenerative disorders such as mucopolysaccharidosis (including, but not limited to, Mucopolysaccharidosis Type I (also known as Hurler syndrome, Hurler-Scheie Syndrome, or Scheie syndrome, IDUA, alpha-L-iduronidase), Mucopolysaccharidosis Type II (also known as Hunter syndrome, IDS, I2L enzyme), Mucopolysaccharidosis Type III (also known as Sanfilippo syndrome, GNS [N-acetylglucosamine-6-sulfatase], HGSNAT [heparan-alpha-glucosaminide N-acetyltransferase], NAGLU [alpha-N-acetylglucosaminidase], and/or SGSH [sulfamidase]), Mucopolysaccharidosis Type IV (also known as Morquio syndrome, GALNS [galactosamine (N-acetyl)-6-sulfatase] and/or GLB1 [beta-galactosidase]), Mucopolysaccharidosis Type V (also known as Scheie syndrome, now a subgroup of type I, also IDUA, alpha-L-iduronidase), Mucopolysaccharidosis Type VI (also known as Maroteaux-Lamy syndrome, ARSB, arylsulfatase B), Mucopolysaccharidosis Type VII (also known as Sly syndrome, GUSB, beta-glucuronidase), Mucopolysaccharidosis Type IX (also known as Natowicz syndrome, HYAL1, hyaluronidase) and/or leukodystrophy (including, but not limited to, adult-onset autosomal dominant leukodystrophy (ADLD; LMNB1, lamin B1), Aicardi-Goutieres syndrome (TREX1, RNASEHSB, RNASEH2C, and/or RNASEH2A), Alexander disease (FRAP, glial fibrillary acidic protein), CADASIL (Notch3), Canavan disease (ASPA, aspartoacylase), CARASIL (HTRA1, serine protease HTRA1), cerebrotendinous xanthomatosis ("CTX," CYP27A1, sterol 27-hydroxylase) childhood ataxia and cerebral hypomyelination (CACH)/vanishing white matter disease (VWMD) (eIF2B, eukaryotic initiation factor 2B), Fabry disease (GM, alpha-galactosidase A), fucosidosis (FUCA1, alpha-L-fucosidase), GM1 gangliosidosis (GLB1, beta-galactosidase), L-2-hydroxyglutaric aciduria (L2HDGH, L-2-hydroxyglutarate dehydrogenase), Krabbe disease (GALC, galactocerebrosidase), megalencephalic leukoencephalopathy with subcortical cysts ("MLC," MLC1 and/or HEPACAM), metachromatic leukodystrophy (ASA, arylsulphatase A), multiple sulfatase deficiency ("MSD," SUMF1, sulfatase modifying factor 1 affecting all sulfatase enzymes), Pelizaeus-Merzbacher disease (also known as "X-linked spastic paraplegia," PLP1 [X-linked proteolipid protein 1] and/or GJA12 [gap junction protein 12]), Pol III-Related Leukodystrophies (POLR3A and/or POLR3B), Refsum dis-

ease (PHYH, [phytanoyl-CoA hydroxylase] and/or Pex7 [PHYH importer into peroxisomes]), salla disease (also known as “free sialic acid storage disease,” SLC17A5, sialic acid transporter), Sjogren-Larsen syndrome (ALDH3A2, aldehyde dehydrogenase), X-linked adrenoleukodystrophy (“ALD,” ABCD1, peroxisomal ATPase Binding Cassette protein), Zellweger syndrome spectrum disorders (also known as peroxisomal biogenesis disorders, PEX1, PEX2, PEX3, PEX4, PEX5, PEX10, PEX11B, PEX12, PEX13, PEX14, PEX16, PEX19, PEX26), and the like.

[0191] Disorders of the CNS include ophthalmic disorders involving the retina, posterior tract, and optic nerve (e.g., retinitis pigmentosa, diabetic retinopathy and other retinal degenerative diseases, uveitis, age-related macular degeneration, glaucoma).

[0192] Most, if not all, ophthalmic diseases and disorders are associated with one or more of three types of indications: (1) angiogenesis, (2) inflammation, and (3) degeneration. The delivery vectors of the present invention can be employed to deliver anti-angiogenic factors; anti-inflammatory factors; factors that retard cell degeneration, promote cell sparing, or promote cell growth and combinations of the foregoing.

[0193] Diabetic retinopathy, for example, is characterized by angiogenesis. Diabetic retinopathy can be treated by delivering one or more anti-angiogenic factors either intraocularly (e.g., in the vitreous) or periorcularly (e.g., in the sub-Tenon’s region). One or more neurotrophic factors may also be co-delivered, either intraocularly (e.g., intravitreally) or periorcularly.

[0194] Uveitis involves inflammation. One or more anti-inflammatory factors can be administered by intraocular (e.g., vitreous or anterior chamber) administration of a delivery vector of the invention.

[0195] Retinitis pigmentosa, by comparison, is characterized by retinal degeneration. In representative embodiments, retinitis pigmentosa can be treated by intraocular (e.g., vitreal administration) of a delivery vector encoding one or more neurotrophic factors.

[0196] Age-related macular degeneration involves both angiogenesis and retinal degeneration. This disorder can be treated by administering the inventive delivery vectors encoding one or more neurotrophic factors intraocularly (e.g., vitreous) and/or one or more anti-angiogenic factors intraocularly or periorcularly (e.g., in the sub-Tenon’s region).

[0197] Glaucoma is characterized by increased ocular pressure and loss of retinal ganglion cells. Treatments for glaucoma include administration of one or more neuroprotective agents that protect cells from excitotoxic damage using the inventive delivery vectors. Such agents include N-methyl-D-aspartate (NMDA) antagonists, cytokines, and neurotrophic factors, delivered intraocularly, optionally intravitreally.

[0198] In other embodiments, the present invention may be used to treat seizures, e.g., to reduce the onset, incidence and/or severity of seizures. The efficacy of a therapeutic treatment for seizures can be assessed by behavioral (e.g., shaking, ticks of the eye or mouth) and/or electrographic means (most seizures have signature electrographic abnormalities). Thus, the invention can also be used to treat epilepsy, which is marked by multiple seizures over time.

[0199] In one representative embodiment, somatostatin (or an active fragment thereof) is administered to the brain using a delivery vector of the invention to treat a pituitary tumor.

[0200] According to this embodiment, the delivery vector encoding somatostatin (or an active fragment thereof) is administered by microinfusion into the pituitary. Likewise, such treatment can be used to treat acromegaly (abnormal growth hormone secretion from the pituitary). The nucleic acid sequences (e.g., GenBank Accession No. J00306) and amino acid sequences (e.g., GenBank Accession No. P01166; contains processed active peptides somatostatin-28 and somatostatin-14) of somatostatins are known in the art.

[0201] In particular embodiments, the vector can comprise a secretory signal as described, e.g., in U.S. Pat. No. 7,071,172.

[0202] In representative embodiments of the invention, the virus vector and/or virus capsid is administered to the CNS (e.g., to the brain or to the eye). The virus vector and/or capsid may be introduced into the spinal cord, brainstem (medulla oblongata, pons), midbrain (hypothalamus, thalamus, epithalamus, pituitary gland, substantia nigra, pineal gland), cerebellum, telencephalon (corpus striatum, cerebrum including the occipital, temporal, parietal and frontal lobes, cortex, basal ganglia, hippocampus and portamygdala), limbic system, neocortex, corpus striatum, cerebrum, and/or inferior colliculus. The virus vector and/or capsid may also be administered to different regions of the eye such as the retina, cornea and/or optic nerve.

[0203] The virus vector and/or capsid may be delivered into the cerebrospinal fluid (e.g., by lumbar puncture) for more disperse administration of the delivery vector. The virus vector and/or capsid may further be administered intravascularly to the CNS in situations in which the blood-brain barrier has been perturbed (e.g., brain tumor or cerebral infarct).

[0204] The virus vector and/or capsid can be administered to the desired region(s) of the CNS by any route known in the art, including but not limited to, intrathecal, intracerebral, intraventricular, intravenous (e.g., in the presence of a sugar such as mannitol), intranasal, intra-aural, intra-ocular (e.g., intra-vitreous, sub-retinal, anterior chamber) and periorcular (e.g., sub-Tenon’s region) delivery as well as intramuscular delivery with retrograde delivery to motor neurons.

[0205] In some embodiments, the virus vector or composition of the present invention may be delivered via an enteral, parenteral, intrathecal, intracisternal, intracerebral, intraventricular, intranasal, intra-aural, intra-ocular, periorcular, intrarectal, intramuscular, intraperitoneal, intravenous, oral, sublingual, subcutaneous and/or transdermal route. In some embodiments, the virus vector or composition of the present invention may be delivered intracranially and/or intraspinally.

[0206] In particular embodiments, the virus vector and/or capsid is administered in a liquid formulation by direct injection (e.g., stereotactic injection) to the desired region or compartment in the CNS. In other embodiments, the virus vector and/or capsid may be provided by topical application to the desired region or by intra-nasal administration of an aerosol formulation. Administration to the eye may be by topical application of liquid droplets. As a further alterna-

tive, the virus vector and/or capsid may be administered as a solid, slow-release formulation (see, e.g., U.S. Pat. No. 7,201,898).

[0207] In yet additional embodiments, the virus vector can be used for retrograde transport to treat and/or prevent diseases and disorders involving motor neurons (e.g., amyotrophic lateral sclerosis (ALS); spinal muscular atrophy (SMA), etc.). For example, the virus vector can be delivered to muscle tissue from which it can migrate into neurons.

[0208] The present invention may be as defined in any one of the following numbered paragraphs.

1. An adeno-associated virus (AAV) capsid protein which comprises an AAV2.5 capsid protein comprising one or more amino acid substitutions that introduce a new glycan binding site (e.g., which does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5).

2. The AAV capsid protein of paragraph 1, wherein the one or more amino acid substitutions comprise:

[0209] a) SQAGASDIRDQSR464-476SX₁AGX₂SX₃X₄X₅X₆QX₇R, wherein X₁₋₇ can be any amino acid; and

[0210] b) EYSW500-503EX₈X₉W, wherein X₈₋₉ can be any amino acid.

3. The AAV capsid protein of paragraph 2, wherein:

[0211] X₁ is V or a conservative substitution thereof;

[0212] X₂ is P or a conservative substitution thereof;

[0213] X₃ is N or a conservative substitution thereof;

[0214] X₄ is M or a conservative substitution thereof;

[0215] X₅ is A or a conservative substitution thereof;

[0216] X₆ is V or a conservative substitution thereof;

[0217] X₇ is G or a conservative substitution thereof;

[0218] X₈ is F or a conservative substitution thereof; and/or

[0219] X₉ is A or a conservative substitution thereof.

4. The AAV capsid protein of paragraph 3, wherein X₁ is V, X₂ is P, X₃ is N, X₄ is M, X₅ is A, X₆ is V, X₇ is G, X₈ is F, and X₉ is A, wherein the new glycan binding site is a galactose binding site.

5. The AAV capsid protein of any one of paragraphs 1-4, wherein the amino acid sequence of the AAV2.5 capsid protein is SEQ ID NO:1 or a functional derivative thereof (e.g., which does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5).

6. The AAV capsid protein of any one of paragraphs 1-5, wherein the amino acid sequence is SEQ ID NO:2 or a functional derivative thereof (e.g., which does not contain a substitution at the position corresponding to amino acid 267 of AAV2.5, or does not contain a serine at the position corresponding to amino acid 267 of AAV2.5).

7. A viral capsid comprising the AAV capsid protein of any one of paragraphs 1-6.

8. A virus vector comprising:

[0220] (a) the viral capsid of paragraph 7; and

[0221] (b) a nucleic acid comprising at least one terminal repeat sequence,

[0222] wherein the nucleic acid is encapsidated by the viral capsid.

8a. The virus vector of paragraph 8, that exhibits substantially increased transduction in one or more of human fibroblasts, neuronal and glial cells as compared to AAV rh10, as measured by Relative Luciferase unit.

8b. The virus vector of any one of paragraphs 8 or 8a that evades pre-existing neutralizing antibodies in a subject.

9. A composition comprising the AAV capsid protein of any one of paragraphs 1-6, the viral capsid of paragraph 7 and/or the virus vector of any one of paragraphs 8-8b, in a pharmaceutically acceptable carrier.

10. A method of introducing a nucleic acid into a cell, comprising contacting the cell with the virus vector of any one of paragraphs 8-8b.

11. The method of paragraph 10, wherein the cell is in neural tissue.

12. The method of paragraph 11, wherein the cell is a neuron or a glial cell.

13. The method of paragraph 12, wherein the glial cell is an astrocyte.

14. The method of paragraph 11, wherein the virus vector has enhanced transduction of neural tissue as compared to an AAV1, AAV2, AAV9, or AAV2.5 virus vector.

15. The method of any one of paragraphs 10-14, wherein the cell is in a subject.

16. The method of paragraph 15, wherein the subject is a human subject.

17. The method of paragraph 16, wherein the subject is a child.

18. The method of paragraph 17, wherein the child is an infant.

19. The method of paragraph 15 or 16, wherein the subject is in utero.

20. The method of any one of paragraphs 15-19, wherein the subject has a reduced immunologic profile when contacted with the virus vector of paragraph 8 as compared to when contacted with an AAV1, AAV2, AAV9, or AAV2.5 virus vector.

21. A method of treating a disease or disorder in a subject in need thereof, comprising introducing a therapeutic nucleic acid into a cell of the subject by administering to the subject the virus vector of any one of paragraphs 8-8b and/or the composition of paragraph 9, under conditions whereby the therapeutic nucleic acid is expressed in the cell of the subject.

22. The method of paragraph 21, wherein the subject is a human.

23. The method of paragraph 21 or 22, wherein the subject is in utero.

24. The method of any one of paragraphs 21-23, wherein the subject has or is at risk for a CNS disease or disorder.

25. The method of any one of paragraphs 21-23, wherein the subject has or is at risk for a congenital neurodegenerative disorder.

26. The method of any one of paragraphs 21-23, wherein the subject has or is at risk for adult-onset autosomal dominant leukodystrophy (ADLD), Aicardi-Goutieres syndrome, Alexander disease, CADASIL, Canavan disease, CARASIL, cerebrotendinous xanthomatosis childhood ataxia and cerebral hypomyelination (CACH)/vanishing white matter disease (VWMD), Fabry disease, fucosidosis, GM1 gangliosidosis, Krabbe disease, L-2-hydroxyglutaric aciduria megalencephalic leukoencephalopathy with subcortical cysts, metachromatic leukodystrophy, multiple sulfatase deficiency, Pelizaeus-Merzbacher disease, Pol III-Related

Leukodystrophies, Refsum disease, salla disease (free sialic acid storage disease), Sjogren-Larsson syndrome, X-linked adrenoleukodystrophy, Zellweger syndrome spectrum disorders, Mucopolysaccharidosis Type I, Mucopolysaccharidosis Type II, Mucopolysaccharidosis Type III, Mucopolysaccharidosis Type IV, Mucopolysaccharidosis Type V, Mucopolysaccharidosis Type VI, Mucopolysaccharidosis Type VII, Mucopolysaccharidosis Type IX and any combination thereof.

27. The method of paragraph 21 or 22, wherein the subject has or is at risk of having pain associated with a disease or disorder.

28. The method of any one of paragraphs 21-27, wherein the virus vector or composition is delivered via an enteral, parenteral, intrathecal, intracisternal, intracerebral, intraventricular, intranasal, intra-aural, intra-ocular, peri-ocular, intrarectal, intramuscular, intraperitoneal, intravenous, oral, sublingual, subcutaneous and/or transdermal route.

29. The method of any one of paragraphs 21-27, wherein the virus vector or composition is delivered intracranially and/or intraspinally.

29a. The method of any one of paragraphs 15-29, wherein the virus vector is used for repeat dosing in the subject.

30. The AAV capsid protein, viral capsid, virus vector, composition or method of any one of paragraphs 1-29a, wherein Applicants disclaimer as follows: To the extent that any disclosure in PCT/US2020/029493 filed Apr. 23, 2020 falls within the invention as defined in any one or more of the claims of this application, or within any invention to be defined in amended claims that may in the future be filed in this application or in any patent derived therefrom, and to the extent that the laws of any relevant country or countries to which that or those claims apply provide that the disclosure of PCT/US2020/029493 is part of the state of the art against that or those claims in or for that or those countries, we hereby reserve the right to disclaim the said disclosure from the claims of the present application or any patent derived therefrom to the extent necessary to prevent invalidation of the present application or any patent derived therefrom. Without limitation, we state that the above reservation of a right of disclaimer applies at least to claims 1-29a as listed in this application and paragraphs 1-29a as set forth above.

[0223] Having described the present invention, the same will be explained in greater detail in the following examples, which are included herein for illustration purposes only and are not intended to be limiting to the invention.

EXAMPLES

Example 1: Rhesus Monkey in Utero Treatment of AAV2G9 and AAV2.5G9

[0224] Congenital monogenic neurodegenerative disorders such as mucopolysaccharidosis and leukodystrophy are prime candidates for targeted gene therapy, but successful interventions must occur prior to physical and behavioral manifestations of disease. In some cases, disease initiation occurs early during development and therefore treatment must be considered prior to birth. This study compares the safety, efficiency, and cell tropism of a naturally occurring AAV serotype (AAV9) with two novel chimeric AAV vectors (AAV2G9 and AAV2.5G9 (amino acid sequence of the capsid is show in SEQ ID NO:2)) following in utero intracranial administration into primates in the early second

trimester under ultrasound guidance. Tissues were harvested near term and transgene expression assessed by ex vivo bioluminescence imaging (BLI) and qPCR. BLI indicated high levels of firefly luciferase expression in the cerebral hemispheres and spinal cord with all vectors tested. qPCR was highly correlated with BLI findings. No adverse effects on fetal growth or development were observed. Tissues were within normal limits with expected populations of neurons, astrocytes, and oligodendrocytes confirmed by immunohistochemistry. These studies demonstrate the safety, efficacy, and tropism of chimeric AAV vectors for targeted gene therapy for congenital neurodegenerative disorders amenable to gene replacement strategies.

[0225] No adverse effects were detected sonographically or at fetal tissue harvest. Fetal body and organ weights at tissue harvest were within normal limits when compared to historical controls (N=36) (mean 465.1 ± 16.8 historical control mean 484.1 ± 14.2 g; FIG. 1). Brain weights were also comparable (mean 53.8 ± 1.7 g) to concurrent (mean 55.5 ± 1.9 g) and historical controls (mean 56.1 ± 0.6 g).

[0226] Bioluminescence Imaging (BLI) Results.

[0227] All fetuses administered AAV vectors were observed with high levels of firefly luciferase expression within the cerebral hemispheres and primarily correlated with the side of vector administration (FIG. 2). Fetuses administered AAV9 showed total bioluminescence of 2.8×10^8 p/s and 6.6×10^8 p/s in all brain lobes, and 7.9×10^8 p/s and 8.6×10^9 p/s was observed in fetuses administered AAV2G9. Fetuses administered AAV2.5G9 showed 1.4×10^7 to 5.3×10^9 p/s. Bioluminescence in individual brain lobes was in general greater with the chimeric vectors (AAV2G9, AAV2.5G9) when compared to AAV9 (Table 3). Firefly luciferase expression was also noted in the spinal cord. Very low or no bioluminescence was detected in tissues outside the central nervous system.

[0228] Vector Biodistribution Results.

[0229] Vector biodistribution was assessed by qPCR of firefly luciferase copies/50,000 cells in the fetal brain (all lobes of the cerebral hemispheres, cerebellum), spinal cord, and peripheral tissues at tissue harvest (Table 4). Presence of the vector was detected in all AAV-treated brains (9/9) and in all regions of the spinal cord. Presence of the vector was greater for AAV2.5G9 in all brain and spinal cord regions compared to AAV2G9 and AAV9. Overall, luciferase copy numbers varied between individual animals (Table 4). Compared to bioluminescence, high vector genome copies were observed in the spinal cord. Very low or no vector genome was amplified by qPCR in tissues outside the central nervous system.

[0230] This study addressed the transduction efficiency and biodistribution of AAV9 and two novel chimeric AAV vectors, AAV2G9 and AAV2.5G9. Fetal brain development was shown to follow normal developmental patterns after intracranial administration demonstrating the safety of this approach and the AAV vectors studied. The chimeric vectors tested were shown to have robust transduction efficiency in all brain lobes, cerebellum, and spinal cord with single site administration.

[0231] AAV administration in utero has been demonstrated to be safe in the developing eye and ear of the mouse, which contain sensory photoreceptors and sensory hair cells that are post-mitotic and do not regenerate. Post-natal assessments of visual and auditory function revealed that in utero injection of AAV vectors encoding a GFP reporter gene

had little to no effect on sensory thresholds. This is of significance because the death or dysfunction of even a small population of these sensory cells is readily detectable by functional analysis, therefore demonstrating a favorable safety profile. Further histological analysis revealed normal sensory cell density and morphology.

[0232] Rhesus Monkeys.

[0233] All animal procedures conformed to the requirements of the Animal Welfare Act and protocols were approved prior to implementation by the Institutional Animal Care and Use Committee at the University of California, Davis. Activities related to animal care (diet, housing) were performed per California National Primate Research Center standard operating procedures. Normally cycling, adult female rhesus monkeys (*Macaca mulatta*) (N=9 gene transfer; 3 controls) with a history of prior pregnancy were bred and identified as pregnant according to established methods. Pregnancy in the rhesus monkey is divided into trimesters by 55-day increments, with the first trimester (0-55 days gestation), second trimester (56-110 days gestation), and third trimester (111-165 days gestation). Parturition typically occurs at 165±10 days gestational age.

[0234] Vector Administration and Fetal Monitoring.

[0235] Dams were screened for AAV antibodies to select seronegative females for study assignment. AAV vectors were administered under ultrasound guidance in the early second trimester (65±5 days) using an intraventricular approach. Vector supernatant (1×10^{12} genome copies in 0.1 ml volume) was injected via intracranial administration into the right or left lateral ventricle (N=9). All pregnancies were sonographically monitored every 10-14 days during gestation according to established procedures.

[0236] Tissue Harvests.

[0237] Hysterotomies were performed near term according to established protocols. All tissues were removed and imaged for firefly luciferase expression. The fetal brain was weighed then the right and left hemispheres (frontal, parietal, temporal, occipital lobes), cerebellum, and midbrain sectioned. Regions of the spinal cord (cervical, thoracic, lumbar) were also assessed post-BLI for molecular and histological analysis. Samples of all tissues were fixed in formalin for histological analysis or snap-frozen in liquid nitrogen for molecular analysis. Frozen samples were stored at $\leq -80^\circ \text{C}$. until analysis.

[0238] BLI.

[0239] BLI for luciferase expression was performed immediately following an intravenous injection of D-luciferin (100 mg/kg) (IVIS 200® imaging system with Living Image software, Xenogen, Alameda, Calif.). Bioluminescence was assessed using semi-quantitative methods (photons/cm²; P/S) by placing regions of interest around sections with positive luminescence.

[0240] Vector Biodistribution.

[0241] To quantify vector biodistribution, genomic DNA was isolated from snap-frozen tissues using the Gentra Puregene Tissue kit (Qiagen, Valencia, Calif.). qPCR was conducted with primers for firefly luciferase to quantify vector presence and with the housekeeping gene epsilon-globin (Life Technologies) as an internal control for DNA isolation and PCR reactions. Real-Time qPCR analysis was run in 96 well optical plates using the 7900 ABI Sequence Detection System and TaqMan Universal PCR Master Mix

(Applied Biosystems). Genomic DNA expression was quantified relative to the housekeeping gene to normalize the amount of sample DNA.

[0242] Immunohistochemistry.

[0243] Formalin-fixed paraffin sections of the cerebellum and right and left frontal, parietal, temporal, and occipital lobes were assessed by hematoxylin and eosin (H&E) staining to evaluate tissue morphology. IHC was performed with markers of neurons (Neuro-Chrom™ pan-neuronal marker, EMD Millipore), astrocytes (Glial fibrillary acidic protein, Abcam), or oligodendrocytes (Cyclic-nucleotide phosphodiesterase, Abcam) according to established protocols. Briefly, sections were deparaffinized with xylene, then rehydrated in graded ethanol. Heat-mediated antigen retrieval was performed in citrate buffer prior to incubation with primary antibodies overnight at 4° C. Secondary antibodies were applied for 1 h at room temperature (AlexaFluor-488, Life Technologies) for visualization. ProLong Gold antifade reagent with 4',6-diamidino-2-phenylindole, dihydrochloride (DAPI; Life Technologies) was used for mounting and to identify nuclei (Molecular Probes).

[0244] RNA In Situ Hybridization.

[0245] Luciferase reporter RNA was visualized with the RNAscope® 2.5 Assay system (Advanced Cell Diagnostics, Hayward, Calif.) following the protocol described by the manufacturer. Target probes included luciferase (luc2) to visualize vector presence and FOX3 to visualize neurons. Negative control probes were targeted against the bacterial gene dapB while positive control probes targeted housekeeping genes Polr2a and PPIB. Sections were mounted with VectaMount (Vector Laboratories Inc, Burlingame, Calif.) and visualized with an Olympus BX61 microscope.

Example 2: AAV2.5G9 Chimera Retains Activities of AAV2.5

[0246] Preliminary analysis shown in Example 1 has indicated that the AAV2.5G9 chimera retains the various activities of the AAV2.5 chimera, such as the transduction of skeletal muscle, heparin binding, acquisition of a distinct immunological profile, reduced liver tropism, and neurologic cell tropism (neurons and glial cells such as astrocytes). This preliminary analysis will be confirmed by future experiments, examples of which are provided below.

Transduce Skeletal Muscle

[0247] The AAV2.5G9 A267 variant (e.g., SEQ ID NO:2) will be evaluated for the ability to transduce skeletal muscle by the following experiments. Following injection of 1×10^{10} genome containing viral particles into the gastrocnemius muscle of BALB/c mice and compared to control AAV serotypes AAV2.5 and AAV2. Each mouse is imaged at 7, 14, 28, 42, and 95 days post injection. The virus used in this experiment is purified using heparin HPLC or cesium chloride gradients. The amount of light emitted from each animal is calculated using CMIR_image software. The regions of interest (ROI) from each leg are defined and used to calculate total photons emitted. Data are to be represented as an average of all 6 limbs. The AAV2.5G9 A267 variant exhibits comparable to greater muscle transduction than AAV2.5 and/or AAV2.5G9 A267S, and both AAV2.5G9 A267 and AAV2.5 exhibit much higher muscle transduction than AAV2.

Heparin Binding

[0248] The ability of the different variants to be purified by heparin will also be examined by the following experiments. AAV2.5G9 A267 is compared to the AAV2.5, AAV2.5G9 A267S and AAV2 serotypes. Equivalent particles of each AAV variant are applied to heparin agarose type 1 and allowed to bind. The columns are washed with PBS, followed by elution in sodium chloride. The number of particles present in the flow thru, washes and elutions are then determined via dot blot hybridization. Data are to be depicted as percentage of unbound particles (wash and flow thru) and bound (elution). As in prior experiments (U.S. Pat. No. 9,012,224) the AAV2.5 variant exhibits heparin binding profiles similar to AAV2. AAV2.5G9 A267 also exhibits a similar heparin binding profile to that of AAV2 and AAV2.5, indicating that the grafting of the Gal binding pocket onto the AAV2.5 serotype preserves this binding activity and also preserves receptor binding involved in cell tropism.

Immunological Profile

[0249] Similar to other non-enveloped viruses, high doses of AAV generate neutralizing antibody that prevents repeated dosing. With the advent of new serotypes, repeat administration is possible. To explore the ability to avoid a pre-existing immune response to AAV1, AAV2 and/or AAV9, the chimeric AAV2.5G9 A267 vector will be tested for transgene expression in vitro after exposure to serum from animals pre-exposed to different AAV serotypes (1, 2, and 2.5, 9 respectively) by the following experiments.

[0250] To generate animals with a robust immune response to AAV virion shell, 4×10^{10} particles of AAV serotype 1, 2, 2.5, 2.5G9, and 9 vector are independently injected intramuscularly in C57bl/6 mice. Four weeks post-injection, blood is isolated and serum collected. Serum from these animals is then used in a neutralizing antibody assay using 293 cells and AAV specific serotype vectors expressing GFP as a reporter gene. In this assay, serum is sequentially diluted and then mixed with a known amount of serotype specific vector (1×10^8 particles) at 4°C . for 2 hr. This mixture of serum and vector is then added to 293 cells in 24-well plates in the presence of adenovirus helper virus at a multiplicity of infection of 5. Under these conditions, green fluorescent protein (GFP) expression is a measure of serotype-specific vector ability to infect cells in the presence of neutralizing antibodies. The neutralizing antibody titer is then calculated as the highest dilution where GFP expression is 50% or less than control vector (without pre-mixture with serotype specific serum).

[0251] Results will indicate that animals pre-exposed to AAV1 can neutralize AAV1 GFP transduction (e.g., with dilutions as high as 1:1000). However this serotype 1 specific neutralizing antibody requires more mouse serum to neutralize AAV chimeric 2.5 (e.g., 1:100 dilution), and AAV chimeric 2.5G9 A267. More importantly, this observation is true for mouse sera obtained from animals pre-exposed to AAV serotype 2 virion shells. In this assay, only after sera are diluted 1:10,000 is 50% GFP transduction observed when compared to AAV2 control. However, for chimeric 2.5 and 2.5G9, 50% GFP transduction is observed with far less dilution (e.g., only 1:100 dilution) of this mouse serum. Since only 0.6% of the amino acid changes differ from AAV2 in this chimeric vector, these alterations have profound effects on the ability of pre-existing AAV2 neutraliz-

ing antibody to recognize the AAV2.5 and AAV2.5G9 A267 capsid shell. Animals pre-exposed to 2.5 and AAV2.5G9 A267 and then assayed for neutralizing activity against AAV 1, 2, 2.5, 2.5G9 A267 and 9 yields expected results, with highest dilution required for the 2.5 and 2.5G9 A267 vector (e.g., 1:8000) followed by lower dilution (e.g., 1:1000) for AAV2 and even lower dilution (e.g., 1:100) for AAV serotype 1, respectively.

[0252] Similarly, animals pre-exposed to AAV9 can neutralize AAV9 GFP transduction with dilutions as high as 1:1000. However this serotype 9 specific neutralizing antibody will require more mouse serum to neutralize AAV chimeric 2.5G9 A267 (e.g., 1:100 dilution). More importantly, this observation is true for mouse sera obtained from animals pre-exposed to AAV serotype 9 virion shells. In this assay, only after sera are diluted considerably (e.g., 1:10,000) is 50% GFP transduction observed when compared to AAV9 control. However, for chimeric 2.5 and 2.5G9, 50% GFP transduction is observed with far less dilution (e.g., only 1:100 dilution) of this mouse serum. Animals pre-exposed to 2.5 and AAV2.5G9 A267 and then assayed for neutralizing activity against AAV1, 2, 2.5, 2.5G9 A267 and 9 yields expected results, with highest dilution required for the 2.5 and 2.5G9 vector (e.g., 1:8000) followed by lower dilution (e.g., 1:1000) for AAV2 and even lower dilution (e.g., 1:100) for AAV serotype 1 and AAV9 respectively.

[0253] The expected conclusions from these studies is that the amino acid alterations made in AAV2.5 to produce chimeric AAV2.5G9 A267, although small in number, are sufficient to significantly affect the immune profile for this virion when challenged with neutralizing antibodies specific for AAV2, AAV2.5, and AAV9.

[0254] These studies will indicate the AAV2.5G9 A267 vectors are suitable for transducing individuals pre-exposed to AAV1, AAV2, AAV2.5, AAV9, or combinations thereof, thereby providing greater versatility in available vectors. For example, this chimeric vector would allow for re-administration in animals and patients pre-exposed to AAV1, AAV2, AAV9 or AAV2.5. In addition, this demonstrates that selected amino acids can be changed in the AAV2.5 capsid amino acid sequence that significantly alter immune response.

Transduction of Brain and Liver is Also Preserved in the AAV2.5G9 Variant

[0255] Cell type and tissue tropism will also be confirmed by the following experiments. Six- to eight-week-old male C57bl/6 mice are utilized to determine efficiency of AAV2 and the 2.5 vector transduction in liver. The mice are anesthetized using 300 μL 2.5% Avertin, and 1×10^{11} particles of AAV2, AAV2.5, AAV2.5G9 A267 and AAV9 vector carrying the human Factor IX (hFIX) transgene virus are dissolved in 250 μL PBS and injected slowly through the portal vein. The vectors are duplexed virus particles as described in international patent publication WO 01/92551. After 1 and 6 weeks, 100 μL of blood from each mouse is collected from the tail vein using heparin-coated capillary glass tubes. Serum is collected by centrifuging the blood sample at 4°C , 8000 rpm for 20 min. Sera are stored at -80°C . until tested. Expression of hFIX in the serum is tested by standard ELISA methods. Serial dilutions of normal human serum with hFIX levels of 5 $\mu\text{g/mL}$ are used as a standard. Using this assay, it will be found that the 2.5 and the AAV2.5G9 vectors each have a reduced ability to transduce

liver as compared with the AAV2 virus. This experiment will demonstrate that the AAV2.5G9 A267 variant exhibits the muscle tropism of the 2.5 vector, and also preserves the loss of the liver specific tropism the 2.5 vector in turn lost when compared to the liver specific tropism characteristic of AAV2.

[0256] In another experiment, the duplexed AAV2.5 vector, duplexed AAV2.5G9 A267 vector, duplexed AAV9 vector, and duplexed AAV2 vector, each containing a green fluorescent protein (GFP) reporter transgene cassette, are respectively injected into the cortex region of the mouse brain under conditions previously established for AAV2. The vectors are then assayed for neuron specific transduction. It is well-established that AAV1 and AAV2 are specific for neuronal transduction and that AAV2.5 vector transduces neurons as well as non-neuronal cells (glial cells such as astrocytes).

[0257] The sum of these experiments when testing the AAV2.5G9 A267 vector for tissue-specific transduction in vivo will likewise demonstrate that in addition to preserving the gained tissue-specific tropism (e.g., muscle, skeletal or cardiac) of AAV2.5 (previously reported as derived from the AAV serotype 1 parent), and preserving the lost cell type specific transduction (e.g., liver-hepatocyte-specific transduction) of the AAV2.5, the AAV2.5G9 A267 vector also preserves the new tropism (non-neuronal/astrocytes) of the AAV2.5 that is not present in either the donor parent (AAV1) or recipient parent capsid (AAV2) and is totally unique to the chimeric 2.5 vector.

[0258] Heparin Binding Experiments.

[0259] Batch binding of rAAV to heparin agarose is performed as described previously (Rabinowitz (2004) *J. Virology* 78:4421-4432). Briefly, equivalent particles of rAAV virions are applied to heparin agarose type 1 (H-6508, Sigma, St. Louis, Mo.) in 1×PBS, allowed to bind for one hour at room temperature, centrifuged at low speed for 2 minutes, and supernatant (flow through) is then removed. Six washes of five bed-volumes of PBS 1 mM MgCl₂ are performed, followed by a three-step elution of five bed-volumes of PBS 1 mM MgCl containing 0.5 M NaCl (step 1), 1.0 M NaCl (step 2), or 1.5 M NaCl (step 3). The number of rAAV particles present in the washes and the 3-step elution is determined by dot blot hybridization.

[0260] Animal Imaging.

[0261] 1×10^{10} viral genome containing particles (vg) are injected into the gastrocnemius of 6-week-old male BALB/c mice. A total of 6 limbs are injected for each virus type using 25 ul of virus. Animals are imaged at different days post injection using the Roper Scientific Imaging (Princeton Instruments). Briefly, animals are anesthetized and injected IP with luciferin substrate. Ten minutes post-injection the animals are placed in the chamber and light emission is then determined. The average number of total pixels per region of interest is determined using the CMIR_Image software (Center for Molecular Imaging Research, Mass. General) and plotted over time.

Example 3: AAV2.5G9 Exhibits Dual Glycan Binding

[0262] AAV2.5G9 Exploits HS and Gal Receptors Interchangeably in Vitro Similar to AAV2G9

[0263] Competitive inhibition assays will provide evidence of the usage of dual glycan receptors by AAV2.5G9 variant by the following experiments. These assays utilize

virus binding on the cell surface involving soluble heparin and ECL, which selectively binds terminally galactosylated glycans. A mutant CHO cell line, CHO-Lec2, is deficient in transporting CMP-sialic acids from Golgi compartments to the cell surface (Deutscher et al. *J. Biol. Chem.* 261:96-100 (1986)). Therefore, the majority of terminal glycan moieties on the CHO-Lec2 surface are galactose. This unique galactosylation pattern on the surface of CHO-Lec2 and sialylation of wild-type CHO-Pro5 cells can be useful in studying AAV-galactose/AAV-sialic acid interactions (Shen et al. *J. Virol.* 86:10408-10417 (2012); Shen et al. *J. Biol. Chem.* 286:13532-13540 (2011)). HS, but not ECL, significantly inhibits AAV2 transduction in CHO-Lec2 cells, whereas ECL selectively blocks AAV9 transduction by nearly two log units. These results are consistent with the expected transduction profiles for AAV2 and AAV9 (Shen et al. *J. Biol. Chem.* 286:13532-13540 (2011); Summerford et al. *J. Virol.* 72:1438-1445 (1998); Bell et al. *J. Clin. Invest.* 121:2427-2435 (2011)). In contrast, AAV2G9 and AAV2.5G9 will only be effectively neutralized by pretreatment with a combination of both ECL and HS. A small but significant inhibitory effect may be observed for ECL.

[0264] The transduction profiles for AAV2, AAV2.5G9 A267, AAV9, and AAV2G9 are further corroborated by inhibition of cell surface binding of each strain using ECL or HS. The unique cell surface attachment of the chimeric AAV strain is further supported by competitive inhibition of cell surface attachment of AAV2.5G9 exclusively by a combination of ECL and HS but neither reagent alone. This is similar to AAV2G9 and will indicate the ability of AAV2.5G9 A267 to bind two different glycans interchangeably similar to AAV2G9.

[0265] In Vitro Characterization of the Dual Glycan-Binding AAV2.5G9 Chimera.

[0266] Assays are performed for inhibition of AAV2, AAV2.5G9 A267, AAV2G9, and AAV9 transduction on CHO Lec2 cells with FITC-ECL and soluble heparin. CHO Lec2 cells are prechilled at 4° C. and incubated with FITC-ECL, soluble heparin, or both prior to infection with AAV2, AAV2.5G9 A267, AAV2G9, or AAV9 packaging a CBA-luciferase reporter transgene cassette. Transduction efficiency is measured 24 h post-infection as luciferase activity in relative light units. The percentage of transgene expression is calculated by normalizing transduction efficiency to relative light units from controls. Assays for inhibition of cell surface binding are performed with AAV2, AAV2.5G9 A267, AAV2G9, and AAV9 on CHO Lec2 cells with FITC-ECL and soluble heparin. Different AAV particles are bound to cells prechilled at 4° C., and unbound virions are removed by washing with cold PBS. Bound virions are quantified using qPCR after viral genome extraction. The percentage of bound virions is determined by normalizing number of bound virions to that of corresponding controls.

[0267] In Vitro Binding and Transduction Assays.

[0268] CHO-Pro5 and CHO-Lec2 cells are cultured in a minimum Eagle's medium (Thermo Scientific) supplemented with 10% FBS, 100 units/ml of penicillin (Cellgro), 100 µg/ml of streptomycin (Cellgro), and 2.5 µg/ml of amphotericin B (Sigma). Cells are seeded at a density of 1×10^5 cells/well in 24-well plates.

[0269] Competitive Inhibition Assays.

[0270] CHO-Lec2 cells are prechilled at 4° C. for 30 min and incubated with 100 µg/ml of FITC-labeled *Erythrina crista-galli* lectin (FITC-ECL, Vector Laboratories) in α minimum Eagle’s medium at 4° C. for 1 h. Alternatively, different viral capsids are incubated with 100 µg/ml of soluble heparin (Sigma) or 1×PBS (control) at room temperature for 1 h. Mock-treated or FITC-ECL-treated cells are then infected with HS-bound or mock-treated AAV2, AAV2.5G9 A267, AAV2G9, or AAV9 capsids packaging a CBA-Luc transgene cassette at an MOI of 1000 vg copies/cell. Following incubation in the cold room for 1 hr, unbound virions are removed by three washes with ice-cold 1×PBS. For cell surface binding assays, the number of bound virions is measured by quantifying vector genome copy numbers/cell in each well using quantitative PCR. For transduction assays, infected Lec2 cells are moved to 37° C. and incubated for 24 h prior to quantitation of luciferase transgene expression from cell lysates.

Example 4: AAV2.5G9 Exhibits Unexpectedly Higher Transduction than AAVrh10

[0271] Human skin fibroblast GM16095 cells were transduced with either AAVrh10-luciferase reporter and AAV2.5G9-luciferase reporter. The multiplicity of infection (MOI) was 10,000. Transduction of GM16095 cells with AAV2.5G9 was substantially higher (118 fold) compared with AAVrh10 as measured by Relative Luciferase unit RLU Luc (data not shown). Similar experiments will be performed with various types of neuronal and glial cells to compare the transduction of the AAVrh10 with the AAV2.5G9. It is expected that the AAV2.5G9 will demonstrate significantly higher transduction in the neuronal and/or glial cells as compared to the AAVrh10.

[0272] The foregoing is illustrative of the present invention, and is not to be construed as limiting thereof.

TABLE 1					
AAV Genomes					
AAV Sero-types/ Isolates	GenBank Accession Number	AAV Sero- types/ Isolates	GenBank Accession Number	AAV Sero- types/ Isolates	GenBank Accession Number
Clonal Isolates		Hu S17	AY695376	Cy3	AY243019
		Hu T88	AY695375	Cy5	AY243017
Avian AAV	AY186198, AY629583,	Hu T71 Hu T70	AY695374 AY695373	Rh13 Clade E	AY243013
ATCC VR-865	NC_004828	Hu T40 Hu T32	AY695372 AY695371	Rh38 Hu66	AY530558 AY530626
Avian AAV strain	NC_006263, AY629583	Hu T17 Hu LG15 Clade C	AY695370 AY695377	Hu42 Hu67 Hu40	AY530605 AY530627 AY530603
DA-1 Bovine AAV	NC_005889, AY388617	AAV3 AAV3B Hu9	NC_001729 NC_001863 AY530629	Hu41 Hu37 Rh40	AY530604 AY530600 AY530559
AAV4	NC_001829	Hu10	AY530576	Rh2	AY243007
AAV5	AY18065, AF085716	Hu11 Hu53	AY530577 AY530615	Bb1 Bb2	AY243023 AY243022
Rh34	AY243001	Hu55	AY530617	Rh10	AY243015
Rh33	AY243002	Hu54	AY530616	Hu17	AY530582
Rh32	AY243003	Hu7	AY530628	Hu6	AY530621
AAV10	AY631965	Hu18	AY530583	Rh25	AY530557
AAV11	AY631966	Hu15	AY530580	Pi2	AY530554

TABLE 1-continued					
AAV Genomes					
AAV Sero- types/ Isolates	GenBank Accession Number	AAV Sero- types/ Isolates	GenBank Accession Number	AAV Sero- types/ Isolates	GenBank Accession Number
AAV12	DQ813647	Hu16	AY530581	Pi1	AY530553
AAV13	EU285562	Hu25	AY530591	Pi3	AY530555
Clade A		Hu60	AY530622	Rh57	AY530569
		Ch5	AY243021	Rh50	AY530563
AAV1	NC_002077, AF063497	Hu3	AY530595	Rh49	AY530562
AAV6	NC_001862	Hu1	AY530575	Hu39	AY530601
Hu.48	AY530611	Hu4	AY530602	Rh58	AY530570
Hu 43	AY530606	Hu2	AY530585	Rh61	AY530572
Hu 44	AY530607	Hu61	AY530623	Rh52	AY530565
Hu 46	AY530609	Clade D		Rh53	AY530566
		Rh62	AY530573	Rh51	AY530564
Clade B		Rh48	AY530561	Rh64	AY530574
Hu19	AY530584	Rh54	AY530567	Rh43	AY530560
Hu20	AY530586	Rh55	AY530568	AAV8	AF513852
Hu23	AY530589	Cy2	AY243020	Rh8	AY242997
Hu22	AY530588	AAV7	AF513851	Rh1	AY530556
Hu24	AY530590	Rh35	AY243000	Clade F	
Hu21	AY530587	Rh37	AY242998	AAV9	AY530579
Hu27	AY530592	Rh36	AY242999	(Hu14)	
Hu28	AY530593	Cy6	AY243016	Hu31	AY530596
Hu29	AY530594	Cy4	AY243018	Hu32	AY530597
Hu63	AY530624				
Hu64	AY530625				
Hu13	AY530578				
Hu56	AY530618				
Hu57	AY530619				
Hu49	AY530612				
Hu58	AY530620				
Hu34	AY530598				
Hu35	AY530599				
AAV2	NC_001401				
Hu45	AY530608				
Hu47	AY530610				
Hu51	AY530613				
Hu52	AY530614				
Hu T41	AY695378				

Virus and Serotype	Genome Accession No.	Capsid/VP1 Accession No.
AAV1	NC_002077.1	NP_049542.1
AAV2	NC_001401.2	YP_680426.1
AAV3A	NC_001729.1	NP_043941.1
AAV3B	NC_001863.1	NP_045760.1
AAV4	NC_001829.1	NP_044927.1
AAV5	NC_006152.1	YP_068409.1
AAV6	NC_001862.1	NP_045758.1
AAV7	AF513851.1	AAN03855.1
AAV8	AF513852.1	AAN03857.1
AAV9	AY530579.1	AAS99264.1
AAV10	AY631965.1*	AAT46337.1
AAV11	AY631966.1*	AAT46339.1
AAV13	EU285562.1	ABZ10812.1

*Incomplete sequence

TABLE 3							
Quantitative Assessment of Bioluminescence in the Central Nervous System.							
Group	Animal #	R. Frontal	L. Frontal	R. Parietal	L. Parietal	R. Temporal	L. Temporal
AAV9	#1	26.8	16.3	92.2	35.0	86.5	81.4
	#2	4.0	5.6	7.5	6.5	24.2	283.7
AAV2G9	#3	65.4	45.2	223.7	226.9	250.0	489.8
	#4	0.0	0.0	498.6	576.7	2,240.0	4,760.0
AAV2.5G9	#5	20.2	240.0	44.5	418.8	180.3	404.0
	#6	0.0	0.0	31.9	16.8	131.6	9.4
	#7	158.0	63.6	413.0	188.5	1,934.0	205.3
	#8	101.2	52.7	349.3	305.8	1,852.0	259.9
	#9	180.3	836.9	850.1	936.3	1,129.0	646.0

Group	Animal #	R. Occipital	L. Occipital	R. Cerebellum	L. Cerebellum	Spinal Cord	
AAV9	#1	63.1	18.0	86.9	14.9	4.2	
	#2	22.8	174.8	2.0	12.3	10.4	
AAV2G9	#3	232.5	555.0	0.0	0.0	20.3	
	#4	5,262.0	3,240.0	153.6	187.8	2.6	
AAV2.5G9	#5	82.7	1,985.0	1,067.0	1,098.0	5.1	
	#6	238.4	0.0	33.4	45.3	1.2	
	#7	808.4	203.7	98.5	81.7	46.6	
	#8	2,675.0	257.8	185.7	7.6	5.2	
	#9	793.1	181.9	2,978.0	268.9	50.3	

TABLE 4								
Evaluation of Vector Biodistribution in the Central Nervous System by Real-Time qPCR.								
Group	Animal #	R. Frontal	L. Frontal	R. Parietal	L. Parietal	R. Temporal	L. Temporal	R. Occipital
AAV9	#1	81.5	61.4	211.5	1,647.3	283.1	2,437.0	118.8
	#2	0.0	307.7	132.5	195.6	5,310.8	2,713.5	0.0
AAV2G9	#3	109.0	223.2	225.7	834.7	2,198.7	1,174.1	207.3
	#4	0.0	202.2	10,444.0	571.5	2,222.3	2,933.2	30,256.2
AAV2.5G9	#5	464.0	1,378.2	331.5	2,695.3	106.4	28,274.3	589.6
	#6	152.6	0.0	83.5	338.7	1,840.8	22.5	775.2
	#7	130.4	377.3	135.2	514.3	322.7	916.6	160.0
	#8	97.6	260.6	1,323.7	375.6	3,784.1	1,294.6	3,445.5
	#9	653.8	1,179.3	1,433.8	1,556.3	1,249.3	679.6	2,015.1

Group	Animal #	L. Occipital	R. Cerebellum	L. Cerebellum	Cervical SC	Thoracic SC	Lumbar SC	
AAV9	#1	271.4	150.7	487.7	1,562.4	651.3	809.4	
	#2	1,024.4	705.7	6,147.8	440.7	898.5	1,292.0	
AAV2G9	#3	826.8	0.0	312.6	549.4	317.7	527.7	
	#4	169.4	132.3	460.8	2,006.8	1,402.9	609.6	
AAV2.5G9	#5	3,490.1	79.5	36,558.6	779.7	9,167.3	9,272.1	
	#6	0.0	236.0	3,527.6	669.3	578.8	252.4	
	#7	757.2	309.2	178.5	126,384.3	7,228.8	6,107.2	
	#8	1,414.1	5,518.0	24.5	5,422.7	2,692.5	1,548.3	
	#9	772.2	647.3	59.6	11,304.7	0.0	89.5	

SEQUENCE LISTING

Sequence total quantity: 2

SEQ ID NO: 1moltype = AAlength = 736

FEATURELocation/Qualifiers

REGION1..736

note = AAV2.5 capsid protein

source1..736

mol_type = protein

organism = synthetic construct

SEQUENCE: 1

-continued

MAADGYLPDW	LEDTLSEGIR	QWWKLKPGPP	PPKPAERHKD	DSRGLVLPGY	KYLGPFNGLD	60
KGEPVNEADA	AALEHDKAYD	RQLDSGDNPY	LKYNHADAEF	QERLKEDTSF	GGNLGRAVFQ	120
AKKRVLEPLG	LVEEPVKTAP	GKKRPVEHSP	VEPDSSSGTG	KAGQQPARKR	LNFGQTGDAD	180
SVDPDQPLGQ	PPAAPSGLGT	NTMATGSGAP	MADNNEGADG	VGNSSGNWHC	DSTWMGDRVI	240
TTSTRTWALP	TYNNHLYKQI	SSASTGASND	NHYFGYSTPW	GYFDFNRFHC	HFSPRDWQRL	300
INNNWGFRPK	RLNFKLFNIQ	VKEVTQNDGT	TTIANNLTST	VQVFTDSEYQ	LPYVLGSAHQ	360
GCLPPFPADV	FMVPQYGYLT	LNNGSQAVGR	SSFYCLEYFP	SQMLRTGNNF	TFSYTFEDVP	420
FHSSYAHSQS	LDRLMNPLID	QYLYYLSRTN	TPSGTTTQSR	LQFSQAGASD	IRDQSRNWLP	480
GPCYRQQRVS	KTSADNNNSE	YSWTGATKYH	LNGRDSLVP	GPAMASHKDD	EEKFFPQSGV	540
LIFGKQGSEK	TNVDIEKVM	TDEEEIRTTN	PVATEQYGSV	STNLQRGNRQ	AATADVNTQG	600
VLPGMVWQDR	DVYLQGPWA	KIPHTDGHFH	PSPLMGGFGL	KHPPPQILIK	NTPVPANPST	660
TFSAAKFASF	ITQYSTGQVS	VEIEWELQKE	NSKRWNPEIQ	YTSNYAKSAN	VDFTVDNNGV	720
YSEPRPIGTR	YLTRNL					736
SEQ ID NO: 2		moltype = AA length = 736				
FEATURE		Location/Qualifiers				
REGION		1..736				
source		note = AAV2.5G9 A267 capsid protein				
		1..736				
		mol_type = protein				
		organism = synthetic construct				
SEQUENCE: 2						
MAADGYLPDW	LEDTLSEGIR	QWWKLKPGPP	PPKPAERHKD	DSRGLVLPGY	KYLGPFNGLD	60
KGEPVNEADA	AALEHDKAYD	RQLDSGDNPY	LKYNHADAEF	QERLKEDTSF	GGNLGRAVFQ	120
AKKRVLEPLG	LVEEPVKTAP	GKKRPVEHSP	VEPDSSSGTG	KAGQQPARKR	LNFGQTGDAD	180
SVDPDQPLGQ	PPAAPSGLGT	NTMATGSGAP	MADNNEGADG	VGNSSGNWHC	DSTWMGDRVI	240
TTSTRTWALP	TYNNHLYKQI	SSASTGASND	NHYFGYSTPW	GYFDFNRFHC	HFSPRDWQRL	300
INNNWGFRPK	RLNFKLFNIQ	VKEVTQNDGT	TTIANNLTST	VQVFTDSEYQ	LPYVLGSAHQ	360
GCLPPFPADV	FMVPQYGYLT	LNNGSQAVGR	SSFYCLEYFP	SQMLRTGNNF	TFSYTFEDVP	420
FHSSYAHSQS	LDRLMNPLID	QYLYYLSRTN	TPSGTTTQSR	LQFSVAGPSN	MAVQGRNWLP	480
GPCYRQQRVS	KTSADNNNSE	FAWTGATKYH	LNGRDSLVP	GPAMASHKDD	EEKFFPQSGV	540
LIFGKQGSEK	TNVDIEKVM	TDEEEIRTTN	PVATEQYGSV	STNLQRGNRQ	AATADVNTQG	600
VLPGMVWQDR	DVYLQGPWA	KIPHTDGHFH	PSPLMGGFGL	KHPPPQILIK	NTPVPANPST	660
TFSAAKFASF	ITQYSTGQVS	VEIEWELQKE	NSKRWNPEIQ	YTSNYAKSAN	VDFTVDNNGV	720
YSEPRPIGTR	YLTRNL					736

That which is claimed is:

1. A method of treating a disease or disorder in a subject in need thereof, comprising introducing a therapeutic nucleic acid into a cell of the subject by administering to the subject an AAV vector comprising:

- a) an AAV capsid protein that comprises an AAV2.5 VP1 capsid protein, further comprising amino acid substitutions that introduce a galactose-binding site, wherein the amino acid substitutions comprise Q465V, A468P, D470N, I471M, R472A, D473V, S475G, Y501F, and S502A, wherein the amino acid sequence is SEQ ID NO:2; and
- b) a therapeutic nucleic acid comprising at least one terminal repeat sequence,

wherein the nucleic acid is encapsidated by the AAV capsid, under conditions whereby the therapeutic nucleic acid is expressed in the cell of the subject.

2. The method of claim 1, wherein the AAV vector is formulated in a pharmaceutically acceptable carrier.

3. The method of claim 1, wherein the AAV capsid protein further comprises an amino acid substitution at the position corresponding to amino acid 267 of AAV2.5 which is not a serine.

4. The method of claim 1, wherein the capsid protein confers dual glycan binding ability to an AAV vector.

5. The method of claim 1, wherein the subject is a human.

6. The method of claim 5, wherein the subject is a child.

7. The method of claim 6, wherein the child is an infant.

8. The method of claim 1, wherein the subject is in utero.

9. The method of claim 1, wherein the subject has or is at risk for a central nervous system (CNS) disease or disorder.

10. The method of claim 1, wherein the subject has or is at risk for a congenital neurodegenerative disorder.

11. The method of claim 1, wherein the subject has or is at risk for adult-onset autosomal dominant leukodystrophy (ADLD), Aicardi-Goutieres syndrome, Alexander disease, CADASIL, Canavan disease, CARASIL, cerebrotendinous xanthomatosis childhood ataxia and cerebral hypomyelination (CACH)/vanishing white matter disease (VWMD), Fabry disease, fucosidosis, GM1 gangliosidosis, Krabbe disease, L-2-hydroxyglutaric aciduria megalencephalic leukoencephalopathy with subcortical cysts, metachromatic leukodystrophy, multiple sulfatase deficiency, Pelizaeus-Merzbacher disease, Pol III-Related Leukodystrophies, Refsum disease, salla disease (free sialic acid storage disease), Sjogren-Larsson syndrome, X-linked adrenoleukodystrophy, Zellweger syndrome spectrum disorders, Mucopolysaccharidosis Type I, Mucopolysaccharidosis Type II, Mucopolysaccharidosis Type III, Mucopolysaccharidosis Type IV, Mucopolysaccharidosis Type V, Mucopolysaccharidosis Type VI, Mucopolysaccharidosis Type VII, Mucopolysaccharidosis Type IX and any combination thereof.

12. The method of claim 1, wherein the subject has or is at risk of having pain associated with a disease or disorder.

13. The method of claim 1, wherein the virus vector or composition is administered to the subject via an enteral, parenteral, intrathecal, intracisternal, intracerebral, intraventricular, intranasal, intra-aural, intra-ocular, peri-ocular, intrarectal, intramuscular, intraperitoneal, intravenous, oral, sublingual, subcutaneous and/or transdermal route, to thereby contact the cell.

14. The method of claim **1**, wherein the virus vector or composition is administered to the subject intracranially and/or intraspinally, to thereby contact the cell.

15. The method of claim **1**, wherein the cell is in neural tissue.

16. The method of claim **15**, wherein the cell is a neuron or a glial cell.

17. The method of claim **16**, wherein the glial cell is an astrocyte.

18. The method of claim **15**, wherein the AAV vector has enhanced transduction of neural tissue as compared to an AAV1, AAV2, AAV9, or AAV2.5 virus vector.

19. The method of claim **15**, wherein the subject has a reduced immunologic profile when contacted with the AAV vector as compared to when contacted with an AAV1, AAV2, AAV9, or AAV2.5 virus vector.

* * * * *