



US 20240017097A1

(19) **United States**

(12) **Patent Application Publication**
CHALASANI et al.

(10) **Pub. No.: US 2024/0017097 A1**

(43) **Pub. Date: Jan. 18, 2024**

(54) **SONOGENIC STIMULATION OF CELLS**

Publication Classification

(71) Applicant: **Salk Institute for Biological Studies**,
La Jolla, CA (US)

(51) **Int. Cl.**
A61K 41/00 (2006.01)
C12N 15/85 (2006.01)
A61K 38/17 (2006.01)
A61N 7/00 (2006.01)

(72) Inventors: **Sreekanth H. CHALASANI**, La Jolla,
CA (US); **Stuart IBSEN**, La Jolla, CA
(US)

(73) Assignee: **Salk Institute for Biological Studies**,
La Jolla, CA (US)

(52) **U.S. Cl.**
CPC *A61K 41/0028* (2013.01); *C12N 15/85*
(2013.01); *A61K 38/1709* (2013.01); *A61N*
7/00 (2013.01); *C12N 2800/106* (2013.01);
A61N 2007/0026 (2013.01); *A61N 2007/0039*
(2013.01)

(21) Appl. No.: **18/459,143**

(22) Filed: **Aug. 31, 2023**

Related U.S. Application Data

(63) Continuation of application No. 17/029,143, filed on
Sep. 23, 2020, which is a continuation of application
No. 14/843,108, filed on Sep. 2, 2015, now Pat. No.
10,806,951.

(60) Provisional application No. 62/054,600, filed on Sep.
24, 2014.

(57) **ABSTRACT**

The invention provides compositions featuring TRP-4 poly-
peptides and polynucleotides, methods for expressing such
polypeptides and polynucleotides in a cell type of interest,
and methods for inducing the activation of the TRP-4
polypeptide in neurons and other cell types using ultrasound.

Specification includes a Sequence Listing.








FIG. 1A








FIG. 1D







FIG. 2A




FIG. 2B




FIG. 2C




FIG. 2D




FIG. 2E




FIG. 2F




FIG. 3A




FIG. 3B




FIG. 3C




FIG. 3D




FIG. 3E






FIG. 3F






FIG. 4A




FIG. 4B




FIG. 4C

FIG. 5A




FIG. 5B




FIG. 5C




FIG. 5D




FIG. 5E




FIG. 5F




FIG. 6A




FIG. 6B




FIG. 6C




FIG. 6D




FIG. 7A




FIG. 7B





FIG. 7C






FIG. 8B




FIG. 8A

FIG. 8C




FIG. 9




FIG. 10A




FIG. 10B




FIG. 10C




FIG. 10D




FIG. 11A




FIG. 11B




FIG. 11C




FIG. 11D




FIG. 11E




FIG. 11F




FIG. 11G




FIG. 11H





FIG. 12



Before Ultrasound



After Ultrasound




FIG. 13B




FIG. 13A

FIG. 14A




FIG. 14B




FIG. 14C




FIG. 14D




FIG. 14E




FIG. 14F






FIG. 15A




AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20) and *trp-4* specifically in AWC neurons (n=23)

FIG. 15B




AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20). Of these 23 animals tested, 3 reached the baseline

FIG. 15C



AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20). AWC calcium responses without ultrasound stimulus in wild-type (n=10)


FIG. 15D



AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20) and *AWC::trp-4* transgenics (n=6)

FIG. 15E

AWC::GCaMP2.2b; AWC::trp4
without Ultrasound (n = 6)




AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20). Each color represents the response of an individual neuron to ultrasound stimulus presented at t=5 seconds.

FIG. 15F


AWC Response without Ultrasound

— *AWC::GCaMP2.2b* (n=10)
— *AWC::GCaMP2.2b; AWC::trp4* (n=6)



AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20). Average AWC calcium responses without ultrasound stimulus in wild-type and *AWC::trp-4* transgenics.

FIG. 15G



AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20). Average AWC calcium data binned similarly to the data shown in Fig. 5F. Microbubbles are included in all AWC calcium recordings.

FIG. 16A




FIG. 16B




FIG. 17A




FIG. 17B




FIG. 17C




FIG. 17D




FIG. 18A




FIG. 18B





FIG. 18C



SONOGENIC STIMULATION OF CELLS

CROSS-REFERENCE TO RELATED APPLICATIONS

[0001] This application is a continuation of U.S. patent application Ser. No. 17/029,143, filed on Sep. 23, 2020, which is a continuation of U.S. patent application Ser. No. 14/843,108, filed on Sep. 2, 2015, now U.S. Pat. No. 10,806,951, which claims priority to and the benefit of U.S. Provisional Application No. 62/054,600, filed on Sep. 24, 2014, the entire contents of all of which are incorporated by reference herein.

STATEMENT OF RIGHTS TO INVENTIONS MADE UNDER FEDERALLY SPONSORED RESEARCH

[0002] This work was supported by Grant No: NIH R01NM096881-03 from the National Institutes of Health. The government has certain rights in the invention.

SEQUENCE LISTING

[0003] The present application contains a Sequence Listing which has been submitted electronically in XML format.

[0004] The content of the electronic XML Sequence Listing, (Date of creation: Aug. 30, 2023; Size: 17,389 bytes; Name: 167776-010203USCON2-Sequence_Listing.xml) is herein incorporated by reference in its entirety.

BACKGROUND OF THE INVENTION

[0005] Understanding how neural circuits generate specific behaviors requires an ability to identify the participating neurons, record and perturb their activity patterns. The best-understood motor circuit, the crab stomatogastric ganglion (STG) has benefited from electrophysiological access to well-defined cell types as well as an ability to manipulate them. A number of approaches have been developed for manipulating neuronal activity using light (optogenetics) or small molecules. While these methods have revealed insights into circuit computations in a number of model systems including mice, they suffer from one drawback: difficulty in delivering stimulus to the target neurons that are present in deeper brain regions. Methods for the non-invasive stimulation of target neurons and other cell types are required.

BRIEF SUMMARY OF THE INVENTION

[0006] The invention provides compositions featuring TRP-4 polypeptides and polynucleotides, methods for expressing such polypeptides and polynucleotides in a cell type of interest, and methods for inducing the activation of the TRP-4 polypeptide in neurons and other cell types using ultrasound.

[0007] In one aspect, the invention provides a method of stimulating a cell, the method involving contacting a TRP-4 polypeptide expressing cell with ultrasound, thereby stimulating the cell. In one embodiment, the TRP-4 polypeptide comprises the amino acid sequence of SEQ ID NO: 1.

[0008] In another aspect, the invention provides a method of inducing cation influx in a cell, the method involving expressing a heterologous TRP-4 polypeptide in a cell, and applying ultrasound to the cell, thereby inducing cation influx in the cell. In one embodiment, the cell is a mamma-

lian cell or bacterial cell (e.g, bacterial biofilm). In another embodiment, the TRP-4 polypeptide is encoded by a polynucleotide codon-optimized for expression in the cell. In another embodiment, the cell is muscle cell, cardiac muscle cell, neuron, motor neuron, sensory neuron, interneuron, or insulin secreting cell. In another embodiment, the ultrasound has a frequency of about 0.8 MHz to about 4 MHz. In another embodiment, the ultrasound has a focal zone of about 1 cubic millimeter to about 1 cubic centimeter. In another embodiment, the method further involves contacting the cell with a microbubble prior to applying ultrasound. In another embodiment, the cell is in vitro or in vivo.

[0009] In yet another aspect, the invention provides a method of treating a disease or disorder in a subject in need thereof, the method involving expressing in a cell (e.g., muscle cell, cardiac muscle cell, neuron, motor neuron, sensory neuron, interneuron, or insulin secreting cell) of a subject a heterologous nucleic acid molecule encoding a TRP-4 polypeptide; and applying ultrasound to the cell, thereby treating a disease or disorder in the subject. In one embodiment, the disease is a neurological disease selected from Parkinson Disease, depression, obsessive-compulsive disorder, chronic pain, epilepsy and cervical spinal cord injury. In another embodiment, the disorder is muscle weakness.

Definitions

[0010] Unless defined otherwise, technical and scientific terms used herein have the same meaning as commonly understood by a person of ordinary skill in the art. See, e.g., Singleton et al., *DICTIONARY OF MICROBIOLOGY AND MOLECULAR BIOLOGY* 2nd ed., J. Wiley & Sons (New York, NY 1994); Sambrook et al., *MOLECULAR CLONING, A LABORATORY MANUAL*, Cold Springs Harbor Press (Cold Springs Harbor, N Y 1989). Any methods, devices and materials similar or equivalent to those described herein can be used in the practice of this invention. The following definitions are provided to facilitate understanding of certain terms used frequently herein and are not meant to limit the scope of the present disclosure.

[0011] By TRP-4 polypeptide is meant a transient receptor potential cation channel capable of conferring ultrasound sensitivity on a neuron and having at least about 85% amino acid identity to SEQ ID NO: 1 or a human ortholog thereof.

SEQ ID NO: 1:
MDSPRGGILGRALREASQSTRQENDVMDQVPVRQMNRDYGGSRRTQMN
PHTSQPGPSHVSI VNVPERGGPTSSTSTHETEHTAHRTESGRFIRRRR
QSREVT TTTTRPYDPAPPTQTRTSSGSTVNGWGENRPKSADEEIKRRRR
SGGGILSRGLREMNKMVEELEQASEEPSTRKGILGTALKDMEGTTYQKI
YRKREETPKRSRSFDDQEMSNRVGMI EHLRLDKDPLELQQLGLTDLT
DTIPTDRPPLRRSSTHLQIGKNSRIIFVFKQPSRDSVTPDRLLGKPLF
RESLTSHASSHEEMSSEDLAMADPQTKILYFAKRDEWANVESEIETIKR
SDFSMADNHGFTAFLAVKAGKDQIVDKMIRK GARVDYSTKGRNATHI
AAMYSGVETLELILKRYSELLRKGAGPKKQLAIHVACERKSKKAFPIVK
RILEDTDQRM AEDGDGSLPIHLAFKFGNVNIVELL LSGPSDEQTRKADG

-continued

NGDTLLHLAARSGNIEAVRTAIAAGCDNANVQNRVGRTPLEHEVAEVDQ
 NMLKIMFKLRADANIHDKEDKTPVHVAERGDTSMVESLIDKFGGSIRA
 RTRDGSLLHIAACSGHTSTALAFKRGVPLEMPNKKGALGLHSAAAAG
 ENDVVKMLIARGTNVDVTRDNYTALHVAVQSGKASVVETLLGSGADIH
 VKGGELGQTALHIAASLNGAESRDCAMMLLKSGGQPDVAQMDGETCLHI
 AARSGNKDIMRLLLDENADSKISSKIGETPLQVAAKSCNFEEAASMILKH
 LSEVLTQEQLKEHVNHRNDGFTALHYAAEIEQRQLHFPGEDAKLVNLL
 IDYGGMVEMPSLNANETAMHMAARSGNQAVLLAMVNKI GAGAVQIVQNK
 QSKNGWSPLLEACARGHSGVANI LLKHHARIDVEDEMGR TALHLAAFNG
 HLSLVHLLLQHKAFVNSKSKTGEAPLHLAAQGHVQVNVLVQDHGAAL
 EAITLDNQ TALHFAAKFGQLAVSQ TLLALGANPNARDDKGQ TPLHLAAE
 NDFPDVVKFLKMRNNRSVLT AIDHNGFTCAHIAAMKGS LAVVRELM
 IDKPMVIQAKTKTLEAT TLHMAAAGGHANIVKILLENGANAEDENSHGM
 TALHLGAKNGFISILEAFDKI LWKRCSRKTGLNALHIAAFYGN SDFVNE
 MLKHVQATVRSEPP IYNHHVNKEFSTEYGF TPLHLAAQSGHDSLVRMLL
 NQGVQVDATSTTMNVIPLHLAAQQGHIAVVGMLLSRSTQ QHAKDWRGR
 TPLHLAAQNGHYEMVSLLI AQGSNINVMQNGWTGLHFATRAGHLSVVK
 LFIDSSADPLAETKEGKVP LCFAAAHNIECLRFL LKQKHDTHQLMEDR
 KFIFDLMVC GKINDNEPLQEFILQSPAPIETAVKLSALYRDMSEKEKER
 AKDLLNVAVESENMAVELLGI TATEYNAALLL KAKDNRGRPLLDVLIEN
 EQKEVVS YASVQRYL TEVWTARVDWSFGKFVAFSLFVLI CPPAWFYFSL
 PLDSRIGRAPIIKFVCHIVSHVYFTILLTIVVLNITHKMYEVT SVVNP
 VEWLLLLWLSGNLVSELSTVGGGSGLGIVKVLILVLSAMAIAVHVLAFL
 LPAVELTHLDNDEKLHFARTMLYLKNQLFAFALLFAFVEYLDDELTVHHL
 FGPWAI IIRDLMYDLARFLVILMLFVAGFTLHVTSIFQPAYQPVEDDSA

-continued

ELMRLASPSQTLEMLFFSLFGLVEPDSMPPLHLVPDFAKIILKLLFGIY
 MMVTLIVLINLLIAMMSDYQRIQAQSDKEWKFGRAILIRQMNKKSATP
 SPINMLTKLIVL RVAWRNRLRCMTRKAQDDLRFEEENIDAF SMGGGQGG
 RQSP TNEGREGQQELGNSADWNIETVIDWRKIVSMYYQANGKLT DGR TK
 EDVDLAMAVPTSF

For specific proteins described herein (e.g., TRP-4), the named protein includes any of the protein's naturally occurring forms, or variants or homologs that maintain the protein transcription factor activity (e.g., within at least 50%, 80%, 90%, 95%, 96%, 97%, 98%, 99% or 100% activity compared to the native protein). In some embodiments, variants or homologs have at least 90%, 95%, 96%, 97%, 98%, 99% or 100% amino acid sequence identity across the whole sequence or a portion of the sequence (e.g. a 50, 100, 150 or 200 continuous amino acid portion) compared to a naturally occurring form. In other embodiments, the protein is the protein as identified by its NCBI sequence reference. In other embodiments, the protein is the protein as identified by its NCBI sequence reference or functional fragment or homolog thereof. In embodiments, the TRP-4 polypeptide is substantially identical to the protein identified by the NCBI reference number Gene ID: GI: 193247904 or a variant or homolog having substantial identity thereto. In embodiments, the TRP-4 polypeptide is the protein as identified by the NCBI sequence reference GI: 193247904. In embodiments, the TRP-4 polypeptide is the protein as identified by the NCBI sequence reference GI:193247904, homolog or functional fragment thereof. In embodiments, the TRP-4 polypeptide includes the amino acid sequence of SEQ ID NO: 1. In embodiments, the TRP-4 polypeptide is the amino acid sequence of SEQ ID NO: 1.

[0012] By "TRP-4 polynucleotide" is meant a nucleic acid molecule encoding a TRP-4 polypeptide. In particular embodiments, the codons of the TRP-4 polynucleotide are optimized for expression in an organism of interest (e.g., optimized for human expression, bacterial expression, murine expression). The sequence of an exemplary TRP-4 polynucleotide is provided herein below

SEQ ID NO: 2:

ATGGATTCCGCCAGTGGCGGAATCCTGGGAAGAGCTTTACGAGAAGCATCACAATCGACTAGGC
 AAGAAAATGATGTTGATATGGATCAGGTACCCGTACGGCAGATGAACAGGGATTACGGTGGATC
 CAGGAGGACTCAGATGAATCCCCACACCTCCCAACCTGGTCCATCTCATGTATCAATTGTAAT
 GTCCCAGAACGCGGAGGACCCACATCTTCCACATCAACCACACATGAGACAGAGCACACGGCAC
 ATAGGACAGAGTCCGGGAGGTTTATCAGACGCCGTCGCCAATCTCGAGAGGTTACCACCACAAC
 CACAAGACCCATGACCCCGCTCCTCCAACCCAGACCCGAACAAGCTCCGGCTCAACAGTAAAT
 GGATGGGGGAGAAATCGACCGAAATCTGCTGATGAGGAGATCAAACGGCGGCGAAGAAGTGGCG
 GGGGAATCCTGTCTCGCGGGCTTCGAGAAATGAACAAAATGGTGAAGAGTTGGAGCAAGCAAG
 TGAAGAGCCAAGTACCAGGAAGGAATTCTGGGTACTGCGTTAAAGGATATGGAAGGGACCACT
 TATCAAAAGATTTACAGGAAAAGGGAGGAAACTCCAAGCGCTCCCGTTCATTTGACGATCAGG
 AGATGTGCAATCGAGTAGGAATGATCGAGCACTTGCTCCGAGACAAGGATCCTTTGGAGCTTCA
 GCAGTTGGGATTAACCGACCTCCTCACCACCGACACCATCCCAACTGACCGACCACCCCTCCGC

- continued

CGATCCTCGACCCATCTCCAAATCGGAAAGAACTCACGGATCATCTTCGTTCCGAAACAACCAT
CCCGTGATTCAGTCACCCCGCCGGATCGTCTTCTCGGGAAACCTCTGTTTCGAGAGAGTCTCAC
CTCCACGCATCGTCTCATGAGGAAATGTGAGTGAGGACTTGGCAATGGCCGGATCCTCAGACG
AAGATTTTGTATTTTCGCGAAGAGAGATGAGTGGGCGAATGTGGAGTCTGAGATAGAGACTATCA
AGCGGAGTGATTTTAGTATGGCTGATAATCACGGCTTACC GCCTTCCCTCCTAGCCGTCAAAGC
TGGCAAGGATCAAATCGTAGACAAGATGATCCGAAAAGGTGCTCGAGTGGACTATAGCACTAAA
GATGGCCGTAACGCGACTCATATTGCCGCCATGTACTCCGGAGTTGAAACTCTTGAGCTTATCC
TCAAGCGATACTCTGAGCTGCTCCGAAAAGGTGCGGGGCC TAAAAAGCAGCTGGCAATCCATGT
GGCTTGCAGAGAAAATCCAAGAAAGCATTTC CAATTGTGAAGCGGATTTTGGAAAGATACTGAT
CAAAGAATGGCAGAGGATGGGGATGGATCCTTGCCGATACACTTGGCATTCAAGTTTGGGAATG
TTAATATTGTGGAGCTTCTGCTAAGTGGGCCTTCGGATGAACAAACCAGGAAAGCTGATGGAAA
CGGGGATACCTTGCTTCATTTGGCCGCTCGGAGTGGGAATATCGAAGCGGTTCCGGACAGCGATT
GCGGCTGGATGTGATAATGCGAATGTGCAGAATAGGGTGGGAAGGACGCCGCTACATGAGGTAG
CCGAAAGTCGGAGATCAAAATATGCTAAAAATCATGTTCAAACCTCCGCGCCGACGCCAACATCCA
TGATAAGGAGGACAAGACTCCGGTACACGTTGCAGCGGAGCGAGGTGACACTTCGATGGTCGAG
TCACTAATTGACAAGTTTGGTGGCTCAATTCGCGCTAGGACCCGTGATGGGTGACGCTTCTGC
ATATTGCCGCATGTTTCAGGACATACTAGCACCGCATTGCCGTTTTTGAAGAGAGGAGTCCCCCT
CTTCATGCCCAACAAAAAGGAGCCCTGGGTCTTCACTCCGCAGCAGCTGCTGGCTTCAACGAC
GTCGTCAAAATGCTCATTGCTCGGGTACTAATGTAGATGTCCGTACACGAGACAACTACACCG
CTCTCCACGTAGCGGTTCAATCTGGCAAGGCTTCGGTTGTAGAGACCCTGCTGGGAAGTGGTGC
AGACATTCATGTGAAGGGCGGGAACTAGGACAACTGCACTGCACATTGCGGCAAGCTTGAAT
GGAGCCGAGAGTCCGGATTGTGCGATGATGTTGCTGAAAAGTGGAGGGCAGCCGGATGTTGCAC
AAATGGATGGGGAGACTTGTCTGCATATTGCTGCCAGGAGTGGGAATAAGGATATCATGAGGCT
CCTGCTTGACGAGAACGCCGACTCGAAAATAAGCTCAAAGATCGGAGAGACACCCCTCCAGGTG
GCCGCCAAGTCATGCAATTTTGAAGCAGCATCAATGATTTTGAAGCACCTTTCGGAAGTTCTGA
CCCAAGAACAGCTTAAGGAACATGTCAATCATAGAACCAATGACGGCTTACAGCTCTTCACTA
CGCCGCTGAAATCGAGCAGCGCCAGTTACACTTTCAGGAGAAGATGCCAAGCTAGTAAATCTT
CTGATCGACTACGGTGGAAATGGTAGAAATGCCATCACTCAATGCAAATGAGACGGCGATGCATA
TGGCGGCAAGATCCGGAAATCAAGCTGTACTCCTGGCGATGGTCAATAAGATCGGAGCCGGTGC
GGTGCAAATCGTGCAGAACAAGCAGAGCAAGAACGGATGGTCACCGCTGTTGGAAGCATGTGCC
AGAGGGCATTCTGGAGTGGCGAATATTTTGTGTAAGCACCACGCCGATTGATGTATTTCGATG
AAATGGGCCGTA CTGCTCTGCACCTGGCAGCTTTC AATGGGCATCTCTCCCTGGTTACCTTCT
TCTGCAGCACAAAGCATTCTGTGAACAGTAAATCGAAAACCGGAGAGGCACCGCTCCACTTAGCA
GCTCAGCATGGTCATGTGAAGGTGGTGAATGTCCGGTGCAGGATCATGGTGCAGCGCTGGAGG
CAATTACGCTGGATAATCAGACAGCCCTCCACTTTGCCGCAAATTCGGTCAGCTAGCTGTGAG
TCAAACCTTCTGGCTCTCGGAGCAAACCCCAATGCACGTGACGACAAGGGTCAAACCCCTCTC
CATCTGGCAGCTGAGAATGACTTCCCCGACGTTGTGAAGCTCTTCTGAAAATGAGAAATAACA
ACCGGAGTGTGTTGACCGCAATTGATCATAATGGATTACCTGCGCACATATTGCTGCGATGAA
GGGTTCCCTAGCCGTGGTCCGTGAGCTTATGATGATCGACAAGCCTATGGTAATCCAGGCAAAG

- continued

ACCAAAACACTGGAAGCCACTACACTTCATATGGCAGCTGCGGGAGGTCACGCGAACATTGTGA
AGATTCTGCTGGAGAATGGAGCAAACGCGGAAGATGAGAATTCGCACGGAATGACTGCTCTCCA
CCTTGGCGCCAAAACGGATTTCATATCGATTTTGGAGGCATTCGATAAGATCCTATGGAAACGG
TGTTTCGAGAAAGACCGGTCTCAACGCTCTCCACATCGCTGCGTTTACGGAAATTCGGATTTG
TCAATGAAATGCTCAAGCACGTACAAGCAACAGTCCGTTCCGAGCCGCCATCTACAATCACCA
TGTCAATAAGGAATTCTCAACTGAATACGGCTTCACACCTCTCCATTTAGCCGCTCAAAGTGGA
CACGACAGTCTTGTGCGGATGCTTCTGAATCAGGGAGTGCAAGTTGACGCGACCAGTACTACAA
TGAACGTGATCCCCCTCCATCTGGCTGCCAGCAAGGCCACATCGCAGTGGTAGGAATGCTCCT
GTCCAGATCTACTCAGCAGCAGCACGCCAAGGATTGGAGAGGCAGGACCCCGCTCCACCTAGCC
GCTCAGAATGGCCACTACGAGATGGTCTCACTTCTCATGCTCAGGGATCTAACATCAATGTCA
TGGATCAGAATGGCTGGACTGGTCTTCACTTTGCCACTCGTGCCGGGCACCTGAGTGTGTCGTC
GCTGTTTCATCGATAGTTTCAGCGGATCCATTGGCGGAGACCAAGGAGGGCAAAGTTCCATTGTG
TTTGCTGCAGCTCATAATCATATAGAATGTCTTCGATTCCTCCTGAAACAGAAGCATGACACAC
ATCAATTGATGGAAGATCGGAAGTTTCATATTCGACTTGATGGTTTGTGGTAAAACCAATGACAA
TGAGCCTCTACAAGAGTTTATCTTCAATCACCTGCTCCAATTGAGACGGCAGTCAAGTTGTCC
GCGTTGTACAGAGATATGTCGAGAAGGAGAAGGAGAGGGCGAAGGATCTGTTGAATGTGGCAG
TGTTTCAGTGAGAATATGGCTGTGGAGTTGTTAGGGATCACCGCCACCGAATACAATGCCGCTCT
TCTCCTGAAGGCTAAGGACAATCGAGGCCGGCCCCCTACTAGATGTTCTCATTGAAAATGAGCAG
AAAGAAGTAGTCTCCTACGCTCTGTCCAACGCTACCTGACAGAAGTATGGACTGCCCGTGTGCG
ACTGGTCATTTCGAAAGTTTGTGCGATTCTCCCTCTCGTGCTAATATGCCCCCGGCATGGTT
CTACTTCTCACTTCCACTGGATAGTCGGATCGGAAGAGCTCCGATTATTAATTTGTGTGCCAT
ATCGTGTCTCATGTCTATTTACGATACTGCTGACAATGTGGTGTGTAATATTACACATAAGA
TGACGAAGTAACTTCGGTGGTTCCAAACCTGTGGAATGGCTCCTGTTGCTCTGGCTCTCTGG
AAATCTGGTCTCCGAACCTCCTCACTGTGGTGGAGGATCTGGCCTAGGAATCGTAAAGGTCTTA
ATCCTAGTCCTTTCCCGATGGCGATAGCCGTCCATGTCTTAGCCTTCTGCTCCCGGCAGTAT
TCCTAACCCACCTGGATAACGATGAAAAGCTACATTTCCCGGACAATGCTTTATTTGAAAAA
TCAACTTTTCGCCTTTGCCCTGCTATTTGCTTTTGTAGAGTACCTGGATTTCTGACAGTGCAT
CATTTGTTCCGGTCCCTGGGCGATCATTATTCGAGATCTAATGTATGATTTGGCCGTTTCTTG
TGATCCTGATGTTGTTGCTGGCGGGCTTCACACTCCACGTGACGAGTATCTCCAGCCTGCCTA
CCAGCCTGTCGACGAGGACAGCGCCGAGCTGATGCGTCTGGCCTCCCGTCTCAAACCTCGAA
ATGCTCTTCTTCTCGCTCTTCGGACTCGTCGAGCCGATTCAATGCCCCGCTCCATCTAGTTC
CAGATTTTGCAAAAATCATCTTAAACTTCTATTTCGGAATCTACATGATGGTCCCTTGATTGT
GCTGATCAACTTGCTGATTGCTATGATGTCTGACACCTACCAACGAATTCAGGCACAGTCGGAT
AAGGAATGGAAGTTTCGGAAGAGCTATTCTGATCAGACAGATGAATAAGAAAAGCGCCACGCCGT
CGCCGATAAATATGTTAACAAAGTTGATAATTGTGCTGAGGGTAGCCTGGCGGAATCGGTTGAG
ATGCATGACCCGAAAAGCCCAAGACGATCTTCGTTTCGAGGAGAACATCGACGCTTCTCCATG

- continued

GGTGGCGCCAGCAGGGAAGGCAAAGTCCGACCAATGAAGGAAGAGAAGGCCAGCAAGAGCTTG
 GTAACCTCGGCTGACTGGAACATCGAGACAGTCATCGACTGGAGGAAGATTGTTTCAATGTACTA
 TCAGGCGAATGGGAAGCTTACAGACGGGCGAACCAAAGAGGATGTGGATTGGCAATGGCAGTA
 CCTACTAGTTTTTAG

[0013] For expression in a human cell, the following
 codon optimized sequence is used:

(SEQ ID NO: 3)

GTTTAAACGGCGCGCCGGTACCATGAATCCTCACACTTCTCAGCCAGGGCCAAGCCATGTCTCC
 ATTGTCAACGTGCCAGAGCGGGGGGACCAACCTCCTCAACCTCCACCACACACGAGACCGAAC
 ACACAGCCCATCGCACAGAGAGCGGCCGATTTCATCCGGAGAAGGCCAGTCCAGAGAAGTGAC
 TACCACAACCTACCAGGCCCTACGATCCTGCACCACCTACCAGACAAGAAGTAGCTCCGGCTCC
 ACCGTGAATGGGTGGGGCGAGAACAGGCCCAAGTCTGCCGACGAGGAGATCAAGCGACGGAGAA
 GGAGTGGCGGGGAATCCTGTCAAGAGGGCTGAGGGAGATGAACAAGATGGTGGAGGAACTGGA
 ACAGGCCTCTGAGGAACCCAGTACACGCAAGGGCATTCTGGGGACTGCTCTGAAAGACATGGAG
 GGCACAACCTTACCAGAAGATCTATCGGAAAAGAGAGGAAACCCCTAAAAGGTCTCGCAGTTTCG
 ACGATCAGGAGATGAGCAACAGAGTGGGGATGATCGAACATCTGCTGAGGGACAAGGACCCCT
 GGAGCTCCAGCAGCTGGGACTGACAGACCTGCTGACCACAGATACTATTCCAACCGACCGACCA
 CCACTGCGCCGATCTAGTACTCACCTCCAGATCGGCAAGAACAGCCGGATCATTTTCGTGCCAA
 AACAGCCCAGCCGCGATTCCGTCACCTCCTCCAGACCGACTGCTGGGCAAGCCTCTGTTTCGGGA
 GTCTCTGACCAGTCACGCCTCAAGCCATGAGGAAATGTCTCTGAAGATCTGGCTATGGCCGAC
 CCCAGACCAAGATCCTGTACTTCGCCAAACGCGACGAGTGGGCTAATGTGGAGTCCGAAATTG
 AGACAATCAAGCGGTCAGACTTCAGCATGGCCGACAACACGGATTCACTGCTTTTCTGCTGGC
 AGTGAAGGCCGGCAAAGACCAGATTGTGATAAGATGATCCGAAAAGGAGCACGGGTGGATTAT
 TCTACCAAGGACGGCAGAAACGCCACACATATTGCCGCTATGTACAGTGGCGTGGAGACACTGG
 AACTGATCCTGAAGAGGTATTTCAGAGCTGCTGCGCAAAGGCCGCGGGCTAAGAAACAGCTGGC
 AATCCACGTGGCCTGCGAAAGGAAGTCCAAGAAAGCCTTCCAATTGTGAAAAGAACTCTGGAG
 GACACCGATCAGAGGATGGCTGAAGACGGAGATGGCTCTCTGCCATTACCTGGCATTCAAAT
 TTGGGAATGTGAACATCGTTCGAGCTGCTGCTGTCGGACCTTCTGATGAACAGACTAGAAAGGC
 CGACGGGAATGGAGATACCTGCTGCATCTGGCAGCACGCTCCGGAACATTGAGGCTGTGCGA
 ACCGCAATCGCCGCGGATGCGACAATGCCAACGTGCAGAACCGCGTCCGGGCGAACCACTGC
 ACGAGGTGGCTGAAGTCGGAGATCAGAATATGCTGAAGATTATGTTCAAACGCGCGCAGACGC
 CAACATCCATGACAAGGAGGATAAAAACACAGTGCACGTCGCCGCTGAGCGAGGCGATACTTCA
 ATGGTGGAAAGCCTGATTGACAAGTTTGGCGGGTCCATCCGAGCCGGACAAGAGATGGCTCTA
 CTCTGCTGCATATCGCAGCTGTTCCGGGCACACCTCTACAGCTCTGGCATTCTGAAGAGAGG
 CGTGCTCTGTTTATGCCAAATAAGAAAGGAGCCCTGGGACTGCATAGCGCCGCGCCGCGGC
 TTCAACGACGTGGTCAAGATGCTGATCGCCAGGGGAACAAATGTGGATGTCAGGACCCGCGACA
 ACTACACAGCCCTGCACGTGGCTGTCCAGAGTGGCAAGGCCAGCGTGGTTCGAGACTCTGCTGGG
 CAGCGGAGCAGATATTCATGTGAAGGGAGGAGAAGTGGGACAGACCGCCCTGCACATCGCAGCC
 AGCCTGAACGGGGCAGAGTCCAGGGACTGCGCCATGATGCTGCTGAAAAGCGGGGGACAGCCTG

- continued

ATGTGGCCCAGATGGACGGAGAAACCTGTCTGCACATTGCTGCACGGTCTGGCAATAAGGATAT
CATGAGACTGCTGCTGGACGAGAACGCCGATAGTAAGATTAGTTCAAAAATCGGCGAAACTCCA
CTCCAGGTGGCCGCTAAGTCTTGCAACTTCGAGGCAGCCAGTATGATCCTGAAACACCTGTCAG
AAGTGCTGACCCAGGAGCAGCTGAAGGAACACGTCAATCATAGAACTAACGACGGCTTCACCCG
CCTGCATTACGCCGCCGAGATTGAGCAGAGGCAGCTGCACCTTCCAGGGGAGGATGCCAAGCTG
GTGAATCTGCTGATCGACTATGGCGGGATGGTTCGAGATGCCCTCACTGAATGCAAACGAAACCG
CCATGCACATGGCCGCTAGAAGCGGAAATCAGGCCTGTGCTGCTGGCAATGGTCAACAAGATTGG
AGCCGGCGCTGTGCAGATCGTCCAGAATAAGCAGTCAAAAAACGGCTGGAGCCCACTGCTGGAG
GCATGTGCCAGGGGGCATAGCGGAGTGGCTAACATTCTGCTGAAGCACCATGCACGCATCGACG
TGTTTCGATGAAATGGGGCGAACAGCCCTGCACCTGGCAGCCTTTAATGGACACCTGAGCCTGGT
GCATCTGCTGCTCCAGCACAAAAGCCTTCGTCAACTCAAAGAGCAAACCGGAGAGGCTCCACTG
CACCTGGCTGCACAGCACGGGCATGTGAAGGTGGTCAATGTGCTGGTCCAGGATCATGGGGCCG
CTCTGGAGGCCATCACACTGGACAACCAGACTGCTCTGCACCTTCGAGCCAAATTTGGACAGCT
GGCCGTGAGCCAGACACTGCTGGCTCTGGGGGCAAATCCTAACGCTAGAGACGATAAGGGACAG
ACTCCACTGCACCTGGCCGCCGAGAACGACTTCCCCGATGTGGTCAAGCTGTTTCTGAAAATGA
GAAACAATAACAGGAGCGTGTGACAGCAATTGATCATAATGGCTTCACCTGCGCCACATCGC
CGCTATGAAAGGCAGCCTGGCCGTGGTTCAGGGAGCTGATGATGATTGACAAGCCTATGGTCATC
CAGGCAAAGACTAAAACCTGGAAGCCACTACCTGCACATGGCAGCCGCTGGAGGACACGCCA
ACATTGTGAAGATCCTGCTGGAGAATGGCGCTAACGCAGAAGATGAGAACAGCCACGGCATGAC
CGCACTGCACCTGGGAGCCAAAAACGGATTCAATTCATCCTGGAGGCCCTTGACAAGATTCTG
TGGAAGCGGTGCAGCCGGAAGACAGGGCTGAATGCTCTGCATATCGCAGCCTTCTACGGAAATA
GCGACTTTGTGAACGAGATGCTGAAACACGTGCAGGCCACTGTCCGAGTGAACCCCTATCTA
CAATCACCATGTGAACAAGGAGTTCTCAACCGAATATGGCTTTACACCTCTGCATCTGGCTGCA
CAGAGCGGGCACGATTCCCTGGTGCAGGATGCTGCTGAATCAGGGCGTGCAGGTGCAGCCACCA
GCACAACTATGAACGTGATTCCACTGCATCTGGCAGCTCAGCAGGGACACATCGCAGTGGTCGG
AATGCTGCTGTCCCGCTCTACCCAGCAGCAGCACGCTAAGGATTGGCGAGGACGGACACCCCTG
CATCTGGCAGCCCAGAACGGCCACTATGAGATGGTGAGCCTGCTGATTGCCAGGGCTCCAATA
TCAACGTGATGGACCAGAAATGGCTGGACTGGACTGCATTTTCGCAACCCGGGCTGGACACCTGAG
CGTGGTCAAGCTGTTTATCGACAGCTCCGCCGATCCTCTGGCTGAGACCAAGGAAGGCAAAGTG
CCACTGTGCTTCGCTGCCGCCACAACCATATTGAGTGTCTGAGATTTCTGCTGAAGCAGAAAC
ACGATACACATCAGCTGATGGAAGATAGGAAGTTCACTTTGACCTGATGGTGTGCGGCAAAC
TAATGACAACGAGCCTCTCCAGGAGTTCATCTCCAGTCCCCGCTCCTATCGAGACCGCAGTG
AAACTGTCTGCCCTGTACAGAGATATGAGTGAAAAGGAGAAAGAAAGGGCTAAGGACCTGCTGA
ATGTGGCAGTCTTTTCTGAGAACATGGCCGTGGAACCTGCTGGGAATTACAGCAACTGAGTATAA
TGCTGCACTGCTGCTGAAGGCCAAAGATAACAGAGGCAGGCCACTGCTGGACGTGCTGATCGAG
AACGAACAGAAAGAGGTGGTTCAGTTACGCTCAGTGCAGAGATACCTGACAGAAGTGTGGACTG
CTCGGGTCGATTGGTCAATCGGGAAGTTTGTGGCATTCAGCCTGTTTGTCTGATTTGCCACC
CGCCTGGTTCTACTTTTCCCTGCCACTGGACTCTAGGATTGGACGCGCCCCCATCATCAAGTTC
GTGTGCCACATCGTGTCCCATGTCTACTTTACCATTCTGCTGACAATCGTGGTCTGAATATCA

- continued

CTCACAAGATGTATGAGGTGACCAGCGTGGTCCCAAATCCCGTCGAATGGCTGCTGCTGCTGTG
GCTGTCCGGCAACCTGGTGAGCGAGCTGTCCACCGTCGGAGGAGGCAGCGGACTGGGAATTGTG
AAGGTCCTGATCCTGGTGTGAGCGCAATGGCCATCGCAGTGCACGTCTGGCTTTCTGCTGC
CCGCAGTGTCTGACTCATCTGGACAATGATGAGAAGCTGCACTTCGCCCGCACCATGCTGTA
CCTGAAAAACCAGCTGTTTCGCTTTGCTCTGCTGTTTCGCTTTTGTGGAATATCTGGACTTCCTG
ACAGTCCACCATCTGTTTGGCCCTTGGGCTATCATTATTAGGGACCTGATGTACGATCTGGCAC
GGTTCCTGGTCATCTGATGCTGTTTCGTCGCCGGCTTCACCCTGCATGTGACCTCTATCTTTCA
GCCCGCTATCAGCCTGTGACGAGGATAGTGCTGAACTGATGCGGCTGGCAAGTCCCTCACAG
ACCCTGGAGATGCTGTTCTTTAGTCTGTTTCGGCTGGTGAACCCGATTCAATGCCTCCACTGC
ACCTGGTGCCTGACTTCGCCAAGATTATCCTGAACTGCTGTTTGGGATCTACATGATGGTGAC
CCTGATTGCTCTGATCAACCTGCTGATTGCTATGATGTCTGATACATATCAGCGCATCCAGGCA
CAGAGTGACAAGGAGTGGAAATTTGGCCGGCCATTCTGATCAGACAGATGAATAAGAAATCTG
CTACCCCTAGTCCAATTAACATGCTGACAAAACCTGATTATCGTGCTGCGGGTTCGCTTGGCGCAA
TCGACTGCGGTGTATGACCCGAAAGGCCAGGACGATCTGCGGTTTCGAGGAAAACATCGACGCT
TTTTCAATGGGGGAGGACAGCAGGGACGACAGAGCCCTACCAATGAGGGACGAGAAGGACAGC
AGGAGCTGGGCAATTCGCCGATTGGAACATTGAAACAGTATCGACTGGAGAAAGATCGTCTC
TATGTACTATCAGGCCAATGGCAAACCTGACTGACGGGCGAACCAAGGAGGATGTGATCTGGCT
ATGGCTGTCCCTACTTCTTTCTGAATTCCGATAACTTGTATTATGCAGCTTATAATGGTTACAA
ATAAAGCAATAGCATCACAATTTTCAAAATAAAGCATTTTTTTTCACTGCATTCTAGTTGTGGT
TTGTCCAAACTCATCAATGTATCTTATCATGTCTGGCGGCCGC

[0014] The term “amino acid” refers to naturally occurring and synthetic amino acids, as well as amino acid analogs and amino acid mimetics that function in a manner similar to the naturally occurring amino acids. Naturally occurring amino acids are those encoded by the genetic code, as well as those amino acids that are later modified, e.g., hydroxyproline, γ -carboxyglutamate, and O-phosphoserine. Amino acid analogs refers to compounds that have the same basic chemical structure as a naturally occurring amino acid, i.e., an a carbon that is bound to a hydrogen, a carboxyl group, an amino group, and an R group, e.g., homoserine, norleucine, methionine sulfoxide, methionine methyl sulfonium. Such analogs have modified R groups (e.g., norleucine) or modified peptide backbones, but retain the same basic chemical structure as a naturally occurring amino acid. Amino acid mimetics refers to chemical compounds that have a structure that is different from the general chemical structure of an amino acid, but that functions in a manner similar to a naturally occurring amino acid. The terms “non-naturally occurring amino acid” and “unnatural amino acid” refer to amino acid analogs, synthetic amino acids, and amino acid mimetics, which are not found in nature.

[0015] Amino acids may be referred to herein by either their commonly known three letter symbols or by the one-letter symbols recommended by the IUPAC-IUB Biochemical Nomenclature Commission. Nucleotides, likewise, may be referred to by their commonly accepted single-letter codes.

[0016] “Conservatively modified variants” applies to both amino acid and nucleic acid sequences. With respect to

particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein. For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide. Such nucleic acid variations are “silent variations,” which are one species of conservatively modified variations. Every nucleic acid sequence herein, which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid, which encodes a polypeptide is implicit in each described sequence with respect to the expression product, but not with respect to actual probe sequences.

[0017] By “altered” is meant an increase or decrease. An increase is any positive change, e.g., by at least about 5%, 10%, or 20%; preferably by about 25%, 50%, 75%, or even by 100%, 200%, 300% or more. A decrease is a negative change, e.g., a decrease by about 5%, 10%, or 20%; preferably by about 25%, 50%, 75%; or even an increase by 100%, 200%, 300% or more.

[0018] The terms “comprises”, “comprising”, and are intended to have the broad meaning ascribed to them in U.S. Patent Law and can mean “includes”, “including” and the like.

[0019] “Contacting” is used in accordance with its plain ordinary meaning and refers to the process of allowing at least two distinct species (e.g. chemical compounds including biomolecules, or cells) to become sufficiently proximal to react, interact, affect or physically touch. It should be appreciated, however, that the resulting reaction product can be produced directly from a reaction between the added reagents or from an intermediate from one or more of the added reagents, which can be produced in the reaction mixture. Contacting may include allowing two species to react, interact, or physically touch, wherein the two species may be a recombinant viral particle as described herein and a cell. In embodiments, the two species are an ultrasound contrast agent that is exposed to ultrasound and a cell.

[0020] The word “expression” or “expressed” as used herein in reference to a gene means the transcriptional and/or translational product of that gene. The level of expression of a DNA molecule in a cell may be determined on the basis of either the amount of corresponding mRNA that is present within the cell or the amount of protein encoded by that DNA produced by the cell. The level of expression of non-coding nucleic acid molecules (e.g., siRNA) may be detected by standard PCR or Northern blot methods well known in the art. See, Sambrook et al., 1989 *Molecular Cloning: A Laboratory Manual*, 18.1-18.88.

[0021] Expression of a transfected gene can occur transiently or stably in a cell. During “transient expression” the transfected gene is not transferred to the daughter cell during cell division. Since its expression is restricted to the transfected cell, expression of the gene is lost over time. In contrast, stable expression of a transfected gene can occur when the gene is co-transfected with another gene that confers a selection advantage to the transfected cell. Such a selection advantage may be a resistance towards a certain toxin that is presented to the cell. Expression of a transfected gene can further be accomplished by transposon-mediated insertion into the host genome. During transposon-mediated insertion, the gene is positioned in a predictable manner between two transposon linker sequences that allow insertion into the host genome as well as subsequent excision. Stable expression of a transfected gene can further be accomplished by infecting a cell with a lentiviral vector, which after infection forms part of (integrates into) the cellular genome thereby resulting in stable expression of the gene.

[0022] The term “exogenous” refers to a molecule or substance (e.g., a compound, nucleic acid or protein) that originates from outside a given cell or organism. For example, an “exogenous promoter” as referred to herein is a promoter that does not originate from the plant it is expressed by. Conversely, the term “endogenous” or “endogenous promoter” refers to a molecule or substance that is native to, or originates within, a given cell or organism.

[0023] The term “gene” means the segment of DNA involved in producing a protein; it includes regions preceding and following the coding region (leader and trailer) as well as intervening sequences (introns) between individual coding segments (exons). The leader, the trailer as well as the introns include regulatory elements that are necessary

during the transcription and the translation of a gene. Further, a “protein gene product” is a protein expressed from a particular gene.

[0024] The terms “identical” or percent “identity,” in the context of two or more nucleic acids or polypeptide sequences, refer to two or more sequences or subsequences that are the same or have a specified percentage of amino acid residues or nucleotides that are the same (i.e., about 60% identity, preferably 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99%, or higher identity over a specified region, when compared and aligned for maximum correspondence over a comparison window or designated region) as measured using a BLAST or BLAST 2.0 sequence comparison algorithms with default parameters described below, or by manual alignment and visual inspection (see, e.g., NCBI web site <http://www.ncbi.nlm.nih.gov/BLAST/> or the like). Such sequences are then said to be “substantially identical.” This definition also refers to, or may be applied to, the complement of a test sequence. The definition also includes sequences that have deletions and/or additions, as well as those that have substitutions. As described below, the preferred algorithms can account for gaps and the like. Preferably, identity exists over a region that is at least about 25 amino acids or nucleotides in length, or more preferably over a region that is 50-100 amino acids or nucleotides in length.

[0025] The term “isolated”, when applied to a nucleic acid or protein, denotes that the nucleic acid or protein is essentially free of other cellular components with which it is associated in the natural state. It can be, for example, in a homogeneous state and may be in either a dry or aqueous solution. Purity and homogeneity are typically determined using analytical chemistry techniques such as polyacrylamide gel electrophoresis or high performance liquid chromatography. A protein that is the predominant species present in a preparation is substantially purified.

[0026] By “mammal” is meant any warm-blooded animal including but not limited to a human, cow, horse, pig, sheep, goat, bird, mouse, rat, dog, cat, monkey, baboon, or the like. Preferably, the mammal is a human.

[0027] “Nucleic acid” refers to deoxyribonucleotides or ribonucleotides and polymers thereof in either single- or double-stranded form, or complements thereof. The term “polynucleotide” refers to a linear sequence of nucleotides. The term “nucleotide” typically refers to a single unit of a polynucleotide, i.e., a monomer. Nucleotides can be ribonucleotides, deoxyribonucleotides, or modified versions thereof. Examples of polynucleotides contemplated herein include single and double stranded DNA, single and double stranded RNA (including siRNA), and hybrid molecules having mixtures of single and double stranded DNA and RNA. The terms also encompass nucleic acids containing known nucleotide analogs or modified backbone residues or linkages, which are synthetic, naturally occurring, and non-naturally occurring, which have similar binding properties as the reference nucleic acid, and which are metabolized in a manner similar to the reference nucleotides. Examples of such analogs include, without limitation, phosphorothioates, phosphoramidates, methyl phosphonates, chiral-methyl phosphonates, and 2-O-methyl ribonucleotides.

[0028] Nucleic acid is “operably linked” when it is placed into a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a polypeptide if it is

expressed as a preprotein that participates in the secretion of the polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, “operably linked” means that the DNA sequences being linked are near each other, and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accordance with conventional practice.

[0029] By “positioned for expression” is meant that a polynucleotide (e.g., a DNA molecule) is positioned adjacent to a DNA sequence, which directs transcription, and, for proteins, translation of the sequence (i.e., facilitates the production of, for example, a recombinant polypeptide of the invention, or an RNA molecule).

[0030] The term “plasmid” or “vector” refers to a nucleic acid molecule that encodes for genes and/or regulatory elements necessary for the expression of genes. Expression of a gene from a plasmid or vector can occur in cis or in trans. If a gene is expressed in cis, the gene and the regulatory elements are encoded by the same plasmid and vector. Expression in trans refers to the instance where the gene and the regulatory elements are encoded by separate plasmids or vectors.

[0031] As used herein, the terms “prevent,” “preventing,” “prevention,” “prophylactic treatment” and the like refer to reducing the probability of developing a disorder or condition in a subject, who does not have, but is at risk of or susceptible to developing a disorder or condition.

[0032] By “reference” or “control” is meant a standard condition. For example, an untreated cell, tissue, or organ that is used as a reference.

[0033] The terms “protein”, “peptide”, and “polypeptide” are used interchangeably to denote an amino acid polymer or a set of two or more interacting or bound amino acid polymers. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer.

[0034] The term “recombinant” when used with reference, e.g., to a cell, or nucleic acid, protein, or vector, indicates that the cell, nucleic acid, protein or vector, has been modified by the introduction of a heterologous nucleic acid or protein or the alteration of a native nucleic acid or protein, or that the cell is derived from a cell so modified. Thus, for example, recombinant cells express genes that are not found within the native (non-recombinant) form of the cell or express native genes that are otherwise abnormally expressed, under expressed or not expressed at all. Transgenic cells and plants are those that express a heterologous gene or coding sequence, typically as a result of recombinant methods.

[0035] The term “subject” as used herein refers to a vertebrate, preferably a mammal (e.g., dog, cat, rodent, horse, bovine, rabbit, goat, or human).

[0036] By “transformed cell” is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a polynucleotide molecule encoding (as used herein) a polypeptide of the invention.

[0037] As used herein, the terms “treat,” “treating,” “treatment,” and the like refer to reducing or ameliorating a disorder and/or symptoms associated therewith. It will be appreciated that, although not precluded, treating a disorder or condition does not require that the disorder, condition or symptoms associated therewith be completely eliminated.

[0038] By “transformed cell” is meant a cell into which (or into an ancestor of which) has been introduced, by means of recombinant DNA techniques, a polynucleotide molecule encoding (as used herein) a polypeptide of the invention.

[0039] The terms “transfection”, “transduction”, “transfecting” or “transducing” can be used interchangeably and are defined as a process of introducing a nucleic acid molecule or a protein to a cell. Nucleic acids are introduced to a cell using non-viral or viral-based methods. The nucleic acid molecules may be gene sequences encoding complete proteins or functional portions thereof. Non-viral methods of transfection include any appropriate transfection method that does not use viral DNA or viral particles as a delivery system to introduce the nucleic acid molecule into the cell. Exemplary non-viral transfection methods include calcium phosphate transfection, liposomal transfection, nucleofection, sonoporation, transfection through heat shock, magnetofection and electroporation. In some embodiments, the nucleic acid molecules are introduced into a cell using electroporation following standard procedures well known in the art. For viral-based methods of transfection any useful viral vector may be used in the methods described herein. Examples for viral vectors include, but are not limited to retroviral, adenoviral, lentiviral and adeno-associated viral vectors. In some embodiments, the nucleic acid molecules are introduced into a cell using a retroviral vector following standard procedures well known in the art. The terms “transfection” or “transduction” also refer to introducing proteins into a cell from the external environment. Typically, transduction or transfection of a protein relies on attachment of a peptide or protein capable of crossing the cell membrane to the protein of interest. See, e.g., Ford et al. (2001) *Gene Therapy* 8:1-4 and Prochiantz (2007) *Nat. Methods* 4:119-20.

[0040] As used herein, the terms “treat,” “treating,” “treatment,” and the like refer to reducing or ameliorating a disorder and/or symptoms associated therewith. It will be appreciated that, although not precluded, treating a disorder or condition does not require that the disorder, condition or symptoms associated therewith be completely eliminated.

[0041] An “effective amount” is an amount sufficient to accomplish a stated purpose (e.g. achieve the effect for which it is administered, treat a disease, reduce enzyme activity, reduce one or more symptoms of a disease or condition, reduce viral replication in a cell). An example of an “effective amount” is an amount sufficient to contribute to the treatment, prevention, or reduction of a symptom or symptoms of a disease, which could also be referred to as a “therapeutically effective amount.” A “reduction” of a symptom or symptoms (and grammatical equivalents of this phrase) means decreasing of the severity or frequency of the symptom(s), or elimination of the symptom(s). A “prophylactically effective amount” of a drug is an amount of a drug that, when administered to a subject, will have the intended prophylactic effect, e.g., preventing or delaying the onset (or reoccurrence) of an injury, disease, pathology or condition, or reducing the likelihood of the onset (or reoccurrence) of an injury, disease, pathology, or condition, or their symp-

toms. The full prophylactic effect does not necessarily occur by administration of one dose, and may occur only after administration of a series of doses. Thus, a prophylactically effective amount may be administered in one or more administrations. An “activity decreasing amount,” as used herein, refers to an amount of antagonist required to decrease the activity of an enzyme or protein (e.g. Tat, Rev) relative to the absence of the antagonist. A “function disrupting amount,” as used herein, refers to the amount of antagonist required to disrupt the function of an enzyme or protein relative to the absence of the antagonist. The exact amounts will depend on the purpose of the treatment, and will be ascertainable by one skilled in the art using known techniques (see, e.g., Lieberman, *Pharmaceutical Dosage Forms* (vols. 1-3, 1992); Lloyd, *The Art, Science and Technology of Pharmaceutical Compounding* (1999); Pickar, *Dosage Calculations* (1999); and *Remington: The Science and Practice of Pharmacy*, 20th Edition, 2003, Gennaro, Ed., Lippincott, Williams & Wilkins).

[0042] “Patient” or “subject in need thereof” refers to a living organism suffering from or prone to a disease or condition that can be treated by using the methods provided herein. The term does not necessarily indicate that the subject has been diagnosed with a particular disease, but typically refers to an individual under medical supervision. Non-limiting examples include humans, other mammals, bovines, rats, mice, dogs, monkeys, goat, sheep, cows, deer, and other non-mammalian animals. In embodiments, a patient is human.

BRIEF DESCRIPTION OF THE DRAWINGS

[0043] FIGS. 1A-1F: Amplifying ultrasound signals using microbubbles modifies animal behavior. FIG. 1A is a schematic of the computer-controlled imaging and ultrasound exposure system, FIG. 1B shows the agar plate with animals and FIG. 1C shows a stabilized microbubble. FIG. 1D provides images showing that animals exhibit reversal and omega bends upon ultrasound stimulus (10 ms, 2.25 MHz with peak negative pressure of 0.9 MPa) only in the presence of microbubbles. Quantification of animal responses to ultrasound stimuli (10 ms, 2.25 MHz) with varying peak negative pressures without as shown in FIG. 1E and with microbubbles as shown in FIG. 1F. n=30 for each of the conditions. Averages and s.e.m. are shown. ** indicates p<0.01 and * indicates p<0.05 by Fisher’s exact t-test.

[0044] FIGS. 2A-2F: TRP-4 expression activates ASH and AWC neurons. FIG. 2A shows behavioral responses to low intensity ultrasound require the pore-forming TRP-4 channel. n=30 for each condition. Transgenic animals expressing TRP-4 in ASH neurons as shown in FIG. 2B and in AWC neurons as shown in FIG. 2C execute more reversals upon low intensity ultrasound stimulation (2.25 MHz, 10 ms). n>30 for each genotype and condition. Averages and s.e.m. are shown in all data panels. ** indicates p<0.05, while * indicates p<0.01 by Fisher’s exact t-test. FIG. 2D provides a schematic identifying chemosensory neurons ASH and AWC in *C. elegans*. False-colored images showing changes in GCaMP fluorescence in AWC neurons upon ultrasound stimulation. Warmer colors indicate increased calcium and neural activity. FIG. 2E provides average AWC calcium responses upon ultrasound stimulation (WT n=20, AWC::TRP-4 n=23). FIG. 2F provides average responses binned by distinct times for Control and AWC::TRP-4 animals. Averages and s.e.m. are shown. * indicates p<0.05 by t-test.

[0045] FIGS. 3A-3F: PVD neurons inhibit reversals. FIG. 3A provides a schematic showing PVD and FLP neurons in *C. elegans*. FIG. 3B shows two independent transgenics expressing TRP-4 in PVD neurons show reduced reversals when stimulated with a 2.25 MHz 0.47 MPa peak negative pressure ultrasound wave for 10 ms. n>46 for each genotype. Averages and s.e.m. are shown. ** indicates p<0.05, while * indicates p<0.01 by Fisher’s exact t-test. FIG. 3C provides false-colored images showing changes in GCaMP fluorescence in PVD neurons upon ultrasound stimulation. Warmer colors indicate increased calcium and neural activity. FIG. 3D shows average PVD calcium responses (n=16) along with distance moved by the animal shown as a function of time. Peak PVD response occurs when the animal has stopped moving. FIG. 3E provides a schematic showing the neural circuit that responds to ultrasound stimuli and microbubbles. ASH and AWC neurons promote reversals, while PVD neurons inhibit them. FIG. 3F shows AIY calcium responses to ultrasound stimuli. A representative trace showing the ratio of change in fluorescence to the baseline is shown. Ultrasound stimulus was given at t=5 s and neurons that responded within a 5.5 second window after the stimulus were counted as responders. Bar graphs show % responders with and without ultrasound stimuli for AIY::GCaMP and AIY::GCaMP;AIY::trp-4. Numbers on the bars indicate the number of animals analyzed in each condition. * indicates p<0.05 by fisher exact t-test.

[0046] FIGS. 4A-4C: Ultrasound stimulus modifies microbubble distribution around *C. elegans*. FIG. 4A shows microbubbles that were labeled such that DiI was stably incorporated into the lipid monolayer. FIG. 4B provides a whole animal view showing brightfield, fluorescence before and after ultrasound stimulus and finally, the difference in red. FIG. 4C shows a magnified view of the animal’s head showing the same frames as above. The white arrow points to a large microbubble that is destroyed upon ultrasound stimulation. The red images highlight those microbubbles that have been activated and destroyed by ultrasound stimulus.

[0047] FIGS. 5A-5F: Small reversal and omega bend responses to ultrasound stimuli. trp-4 mutants have altered number of small reversals as shown in FIG. 5A, but not omega bends as shown in FIG. 5B when compared to wild-type animals. Transgenics expressing trp-4 in ASH neurons also exhibit fewer numbers of small reversals as shown in FIG. 5C, but not omega bends as shown in FIG. 5D. AWC::trp-4 animals do not have any significant differences in their small reversal as shown in FIG. 5E or omega bend responses as shown in FIG. 5F upon ultrasound stimulation. Averages and 95% confidence intervals are shown. ** p<0.01 by Fisher’s exact t-test.

[0048] FIGS. 6A-6D: AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b as shown in FIG. 6A and without microbubbles as shown in FIG. 6B. AWC calcium responses in transgenic animals expressing trp-4 specifically in AWC neurons with as shown in FIG. 6C and without microbubbles as shown in FIG. 6D. Each color represents the response of an individual neuron to ultrasound stimulus presented at t=5 seconds.

[0049] FIGS. 7A-7C: PVD activity depends on worm movement. PVD activity is strongly correlated with movement (n=89) as shown in FIG. 7A, in the backward direction (Backward n=25, forward n=16) as shown in FIG. 7B and

when animals stop or slow down (stop or slow down n=22, not stopping or slowing down n=19) as shown in FIG. 7C. Averages and s.e.m are shown with ** indicating $p < 0.001$ by Fisher's exact t-test.

[0050] FIGS. 8A-8C: Ultrasound stimulus modifies microbubble distribution. FIG. 8A shows microbubbles that are uniformly distributed on an agar surface and appear white. FIG. 8B shows that ultrasound stimulus (10 pulses of 10 ms, 2.25 MHz with peak negative pressure of 0.9 MPa) activates the microbubbles specifically in an area of 1 mm diameter (white arrow). Microbubbles outside this focal zone appear undisturbed. FIG. 8C shows that the microbubble (i) expands and (ii) contracts in size with the rarefaction (low-pressure) and compression (high-pressure) portions of the ultrasound pressure wave. This oscillation behavior occurs at the frequency of the driving ultrasound resulting in a variety of behaviors including (iii) microbubble collapse, (iv) fluid microstreaming and (v) merging of microbubbles. These microbubble behaviors create mechanical distortions that can propagate through the agar and the body of the animal.

[0051] FIG. 9: Behavioral responses to ultrasound. FIG. 9 panels show an animal reversing and generating a high-angled omega bend upon ultrasound stimulus. Reversals with greater than two head bends were scored as large, while those with fewer than two head bends were counted as small. Reversals and omega bends are shown with a red line overlaid on the animal tracks on the agar surface.

[0052] FIGS. 10A-10D: Microbubbles transduce ultrasound stimuli. FIG. 10A provides a bar and whisker plot showing the distribution of microbubbles fractionated based on their size. A one-way ANOVA test shows significance in the distribution (** indicates $p < 0.001$). Mixed size population was used for all experiments shown to maintain consistency. FIG. 10B provides images showing animals incubated with small (top) and large (bottom) population of microbubbles. FIG. 10C shows behavioral responses of wildtype animals incubated with small and large microbubbles upon ultrasound stimulation of 10 ms pulse with 2.25 MHz with peak negative pressure of 0.9 MPa. Averages and 95% confidence limits are shown. ** indicates $p < 0.001$ fisher's exact t-test. FIG. 10D provides a graph showing the effect of external humidity levels on animal reversal behavior. It was observed that at different times of the year the animals had different reversal behavior in response to the same 0.47 MPa ultrasound exposure. Under low humidity levels the animals would undergo more large reversals than under high humidity conditions. Applicants accounted for this variable behavior by running a wild-type control for each of the genetically modified strains that was tested. These controls were run on the same day and under the same conditions as the tested strain. Error bars show standard error of the proportion.

[0053] FIGS. 11A-11H show AIY responses to ultrasound in FIGS. 11A-11D. Ratio of change in fluorescence in the AIY neurite without as shown in FIGS. 11A and 11B and with ultrasound as shown in FIGS. 11C and 11D. Neurons that responded in a 5.5 second window around $t = 5$ seconds are shown in FIGS. 11A and 11C and those that did not are shown in FIGS. 11B and 11D. Ultrasound stimulus was presented at $t = 5$ seconds in FIGS. 11C and 11D. FIGS. 11E-11H show the ratio of change in fluorescence to baseline fluorescence in the AIY neurite expressing *trp-4* in AIY interneurons specifically without as shown in FIGS. 11E and

11F and with ultrasound as shown in FIGS. 11G and 11H. Neurons that responded in the same 5.5 second window around $t = 5$ seconds are shown in FIGS. 11E and 11G and those that did not are shown in FIGS. 11F and 11H. Each trace represents data from a single neuron recorded once.

[0054] FIG. 12 provides two panels showing damage to worms through multiple exposures to high peak negative pressure ultrasound in the presence of microbubbles. The worm displays a normal curved sinusoidal body position before exposure to the ultrasound (left). After exposure to 10 pulses of 0.9 MPa peak negative pressure ultrasound with a 1 Hz repetition rate the worm displays abnormalities in maintaining a normal body position and locomotion behavior is inhibited indicating damage has occurred (right).

[0055] FIGS. 13A and 13B provide two panels showing a thermocouple used for measuring temperature increases on agar surfaces. Images showing the probe touching the agar surface without as shown in FIG. 13A and with microbubbles as shown in FIG. 13B.

[0056] FIGS. 14A-14F provide six graphs showing small reversal and omega bend responses to ultrasound stimuli in the presence of microbubbles. *trp-4* mutants have altered number of small reversals as shown in FIG. 14A, but not omega bends as shown in FIG. 14B when compared to wild-type animals. Transgenics expressing *trp-4* in ASH neurons also exhibit fewer numbers of small reversals as shown in FIG. 14C, but not omega bends as shown in FIG. 14D. AWC::*trp-4* animals do not have any significant differences in their small reversal as shown in FIG. 14E or omega bend responses as shown in FIG. 14F upon ultrasound stimulation. Proportions and standard error of the proportion are shown. ** $p < 0.01$ by Fisher's exact test.

[0057] FIGS. 15A-15G: AWC calcium responses to ultrasound stimulus. Ratio of change in fluorescence to baseline fluorescence in AWC neurons expressing the calcium sensor, GCaMP2.2b (n=20) as shown in FIG. 15A and *trp-4* specifically in AWC neurons (n=23) as shown in FIG. 15B. Of these 23 animals tested, 3 reached the baseline as shown in FIG. 15C. AWC calcium responses without ultrasound stimulus in wild-type (n=10) are shown in FIG. 15D and AWC::*trp-4* transgenics (n=6) are shown in FIG. 15E. Each color represents the response of an individual neuron to ultrasound stimulus presented at $t = 5$ seconds. FIG. 15F provides average AWC calcium responses without ultrasound stimulus in wild-type and AWC::*trp-4* transgenics. FIG. 15G provides average AWC calcium data binned similarly to the data shown in FIG. 5F. Microbubbles are included in all AWC calcium recordings.

[0058] FIGS. 16A and 16B provide two graphs showing that FLP neurons do not respond to ultrasound. FIG. 16A provides an average of 6 different FLP GCaMP responses to the ultrasound stimulus presented at $t = 5$ s. No response was observed. FIG. 16B provides the 6 individual FLP GCaMP traces shown in FIG. 16A. Microbubbles are present in all FLP recordings.

[0059] FIGS. 17A-17D provide four panels showing that PVD responses depend on worm movement. FIG. 17A provides an expanded view of average PVD trace in response to ultrasound and microbubbles shown in FIG. 6D. The animal was stimulated with a single ultrasound pulse at $t = 5$ s. There was an immediate decrease in fluorescence, which was then followed by a rapid increase. PVD activity is strongly correlated with movement (n=89) as shown in FIG. 17B, in the backward direction (Backward n=25,

forward n=16) as shown in FIG. 17C and in animals that stopped (stop or slow down n=22, not stopping or slowing down n=19) as shown in FIG. 17D. Proportions and standard error of the proportion are shown with *** indicating $p < 0.001$ by Fisher's exact t-test.

[0060] FIGS. 18A-18C show that AIY transgenic worms have normal local search. Animals were moved from food to a food-free plate and their reversals and omega bends were quantified. The two AIY::trp-4 transgenics executed normal number of small reversals as shown in FIG. 18A, large reversals as shown in FIG. 18B and omega bends as shown in FIG. 18C.

DETAILED DESCRIPTION OF THE INVENTION

[0061] The invention provides compositions featuring TRP-4 polypeptides and polynucleotides, methods for expressing such polypeptides and polynucleotides in a cell type of interest, and methods for inducing the activation of the TRP-4 polypeptide in neurons and other cell types using ultrasound.

[0062] The invention is based, at least in part, on the discovery that misexpression of TRP-4, a pore-forming subunit of a mechanotransduction channel, sensitizes cells to an ultrasound stimulus resulting in calcium influx and motor outputs. Accordingly, this approach can be used to alter cellular functions in vivo.

[0063] Accordingly, the invention provides polynucleotides encoding a TRP4 polypeptide, expression vectors comprising such polynucleotides, cells expressing a recombinant TRP4 polypeptide, and methods for stimulating such cells with ultrasound.

Ultrasound

[0064] Ultrasound is well suited for stimulating neuron populations as it focuses easily through intact thin bone and deep tissue (K. Hynynen and F. A. Jolesz, *Ultrasound Med Biol* 24 (2), 275 (1998)) to volumes of just a few cubic millimeters (G. T. Clement and K. Hynynen, *Phys Med Biol* 47 (8), 1219 (2002)). The non-invasive nature of ultrasound stimulation is particularly significant for manipulating vertebrate neurons including those in humans, as it eliminates the need for surgery to insert light fibers (required for some current optogenetic methods). Also, the small focal volume of the ultrasound wave compares well with light that is scattered by multiple layers of brain tissue (S. I. Al-Juboori, A. Dondzillo, E. A. Stubblefield et al., *PLoS ONE* 8 (7), e67626 (2013)). Moreover, ultrasound has been previously used to manipulate deep nerve structures in human hands and reduce chronic pain (W. D. O'Brien, Jr., *Prog Biophys Mol Biol* 93 (1-3), 212 (2007); L. R. Gavrilov, G. V. Gersuni, O. B. Ilyinsky et al., *Prog Brain Res* 43, 279 (1976)). The invention provides for novel non-invasive compositions for the expression of TRP4 in cells, and methods to stimulate cells expressing TRP4 using low-intensity ultrasound stimulation.

Cellular Compositions Comprising Recombinant TRP-4

[0065] The invention provides cells comprising a recombinant nucleic acid molecule encoding a TRP-4 polypeptide. In one embodiment, the invention provides a cardiac muscle cell comprising a TRP-4 polynucleotide under the control of a promoter suitable for expression in a cardiac cell (e.g.,

NCX1 promoter). In another embodiment, the invention provides a muscle cell comprising a TRP-4 polynucleotide under the control of a promoter suitable for expression in a muscle cell (e.g., myoD promoter). In another embodiment, the invention provides an insulin secreting cell (e.g., beta islet cell) comprising a TRP-4 polynucleotide under the control of a promoter suitable for expression in an insulin-secreting cell (e.g., Pdx1 promoter). In another embodiment, the invention provides an adipocyte comprising a TRP-4 polynucleotide under the control of a promoter suitable for expression in an adipocyte (e.g., iaP2). In another embodiment, the invention provides a neuron comprising a TRP-4 polynucleotide under the control of a promoter suitable for expression in a neuron (e.g., nestin, Tuj1 promoter), in a motor neuron (e.g., H2b promoter), in an interneuron (e.g., Islet 1 promoter), in a sensory neuron (e.g., OMP promoter, T1R, T2R promoter, rhodopsin promoter, Trp channel promoter). Such cells may be cells in vitro or in vivo. In particular embodiments, the cells express a mechanotransduction polypeptide that is a transient receptor potential channel-N (TRPN) polypeptide that is sensitive to ultrasound. In particular embodiments, the mechanotransduction polypeptide is TRP-4 or a functional portion or homolog thereof. In embodiments, the mechanotransduction polypeptide comprises or consists of the amino acid sequence of SEQ ID NO: 1.

Expression of Recombinant TRP-4

[0066] In one approach, a cell of interest (e.g., neuron, such as a motor neuron, sensory neuron, neuron of the central nervous system, or neuronal cell lines) is engineered to express a TRP-4 polynucleotide whose expression renders the cell responsive to ultrasound stimulation. Ultrasound stimulation of such cells induces cation influx.

[0067] TRP-4 may be constitutively expressed or its expression may be regulated by an inducible promoter or other control mechanism where conditions necessitate highly controlled regulation or timing of the expression of a TRP-4 protein. For example, heterologous DNA encoding a TRP4 gene to be expressed is inserted in one or more pre-selected DNA sequences. This can be accomplished by homologous recombination or by viral integration into the host cell genome. The desired gene sequence can also be incorporated into a cell, particularly into its nucleus, using a plasmid expression vector and a nuclear localization sequence. Methods for directing polynucleotides to the nucleus have been described in the art. The genetic material can be introduced using promoters that will allow for the gene of interest to be positively or negatively induced using certain chemicals/drugs, to be eliminated following administration of a given drug/chemical, or can be tagged to allow induction by chemicals, or expression in specific cell compartments.

[0068] Calcium phosphate transfection can be used to introduce plasmid DNA containing a target gene or polynucleotide into cells and is a standard method of DNA transfer to those of skill in the art. DEAE-dextran transfection, which is also known to those of skill in the art, may be preferred over calcium phosphate transfection where transient transfection is desired, as it is often more efficient. Since the cells of the present invention are isolated cells, microinjection can be particularly effective for transferring genetic material into the cells. This method is advantageous because it provides delivery of the desired genetic material

directly to the nucleus, avoiding both cytoplasmic and lysosomal degradation of the injected polynucleotide. Cells can also be genetically modified using electroporation.

[0069] Liposomal delivery of DNA or RNA to genetically modify the cells can be performed using cationic liposomes, which form a stable complex with the polynucleotide. For stabilization of the liposome complex, dioleoyl phosphatidylethanolamine (DOPE) or dioleoyl phosphatidylcholine (DOPC) can be added. Commercially available reagents for liposomal transfer include Lipofectin (Life Technologies). Lipofectin, for example, is a mixture of the cationic lipid N-[1-(2, 3-dioleoyloxy)propyl]-N—N—N-trimethyl ammonia chloride and DOPE. Liposomes can carry larger pieces of DNA, can generally protect the polynucleotide from degradation, and can be targeted to specific cells or tissues. Cationic lipid-mediated gene transfer efficiency can be enhanced by incorporating purified viral or cellular envelope components, such as the purified G glycoprotein of the vesicular stomatitis virus envelope (VSV-G). Gene transfer techniques which have been shown effective for delivery of DNA into primary and established mammalian cell lines using lipopolyamine-coated DNA can be used to introduce target DNA into the de-differentiated cells or reprogrammed cells described herein.

[0070] Naked plasmid DNA can be injected directly into a tissue comprising cells of interest. Microprojectile gene transfer can also be used to transfer genes into cells either in vitro or in vivo. The basic procedure for microprojectile gene transfer was described by J. Wolff in *Gene Therapeutics* (1994), page 195. Similarly, microparticle injection techniques have been described previously, and methods are known to those of skill in the art. Signal peptides can be also attached to plasmid DNA to direct the DNA to the nucleus for more efficient expression.

[0071] Viral vectors are used to genetically alter cells of the present invention and their progeny. Viral vectors are used, as are the physical methods previously described, to deliver one or more polynucleotide sequences encoding TRP4, for example, into the cells. Viral vectors and methods for using them to deliver DNA to cells are well known to those of skill in the art. Examples of viral vectors that can be used to genetically alter the cells of the present invention include, but are not limited to, adenoviral vectors, adeno-associated viral vectors, retroviral vectors (including lentiviral vectors), alphaviral vectors (e. g., Sindbis vectors), and herpes virus vectors.

Targeted Cell Types

[0072] TRP-4 can be expressed in virtually any eukaryotic or prokaryotic cell of interest. In one embodiment, the cell is a bacterial cell or other pathogenic cell type. In another embodiment, the cell is a mammalian cell, such as an adipocyte, muscle cell, cardiac muscle cell, insulin secreting cell (e.g., beta islet cell), and neuron (e.g., motor neuron, sensory neuron, neuron of the central nervous system, and neuronal cell line).

Methods of Stimulating a Neural Cell

[0073] The methods provided herein are, inter alia, useful for the stimulation (activation) of cells. In particular, ultrasound stimulation induces cation influx, thereby altering cell activity. Expression of TRP-4 in a pathogen cell (bacteria) and subsequent ultrasound stimulation induces cation influx

and bacterial cell killing. Ultrasound stimulation of a muscle cell expressing TRP-4 results in muscle contraction. This can be used to enhance muscle contraction or functionality in subjects in need thereof, including subjects suffering from muscle weakness, paralysis, or muscle wasting. Altering the intensity of the ultrasound modulates the extent of muscle activity.

[0074] The term “neural cell” as provided herein refers to a cell of the brain or nervous system. Non-limiting examples of neural cells include neurons, glia cells, astrocytes, oligodendrocytes and microglia cells. Where a neural cell is stimulated, a function or activity (e.g., excitability) of the neural cell is modulated by modulating, for example, the expression or activity of a given gene or protein (e.g., TRP-4) within said neural cell. The change in expression or activity may be 10%, 20%, 30%, 40%, 50%, 60%, 70%, 80%, 90% or more in comparison to a control (e.g., unstimulated cell). In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or higher than the expression or activity in the absence of stimulation. In certain instances, expression or activity is 1.5-fold, 2-fold, 3-fold, 4-fold, 5-fold, 10-fold or lower than the expression or activity in the absence of stimulation. The neural cell may be stimulated by applying an ultrasonic wave to the neural cell.

[0075] The term “applying” as provided herein is used in accordance with its plain ordinary meaning and includes the meaning of the terms contacting, introducing and exposing. An “ultrasonic wave” as provided herein is an oscillating sound pressure wave having a frequency greater than the upper limit of the human hearing range. Ultrasound (ultrasonic wave) is thus not separated from ‘normal’ (audible) sound by differences in physical properties, only by the fact that humans cannot hear it. Although this limit varies from person to person, it is approximately 20 kilohertz (20,000 hertz) in healthy, young adults. Ultrasound (ultrasonic wave) devices operate with frequencies from 20 kHz up to several gigahertz. The methods provided herein use the energy of an ultrasonic wave to stimulate a neural cell expressing an exogenous mechanotransduction protein. A mechanotransduction protein as provided herein refers to a cellular protein capable of converting a mechanical stimulus (e.g., sound, pressure, movement) into chemical activity. Cellular responses to mechanotransduction are variable and give rise to a variety of changes and sensations. In embodiments, the mechanotransduction protein is a mechanically gated ion channel, which makes it possible for sound, pressure, or movement to cause a change in the excitability of a cell (e.g., a sensory neuron). The stimulation of a mechanotransduction protein may cause mechanically sensitive ion channels to open and produce a transduction current that changes the membrane potential of a cell.

[0076] In one aspect, a method of stimulating a cell is provided. The method includes (i) transfecting a cell with a recombinant vector including a nucleic acid sequence encoding an exogenous mechanotransduction polypeptide, thereby forming a transfected cell. (ii) To the transfected cell an ultrasonic wave is applied, thereby stimulating a cell. In embodiments, the mechanotransduction polypeptide is a transient receptor potential channel-N (TRPN) polypeptide or homolog thereof. In embodiments, the mechanotransduction polypeptide is TRP-4 or a functional portion or homolog thereof. In embodiments, the mechanotransduction polypeptide includes the amino acid sequence of TRP4 SEQ ID NO:

1. In embodiments, the mechanotransduction polypeptide is the sequence of SEQ ID NO: 1. In embodiments, the ultrasonic wave has a frequency of about 0.8 MHz to about 4 MHz. In embodiments, the ultrasonic wave has a frequency of about 1 MHz to about 3 MHz. In embodiments, the ultrasonic wave has a focal zone of about 1 cubic millimeter to about 1 cubic centimeter.

[0077] In embodiments, the method further includes before the applying of step (ii) contacting the transfected neural cell with an ultrasound contrast agent. In embodiments, the ultrasound contrast agent is a microbubble. In embodiments, the microbubble has a diameter of about 1 μm to about 6 μm . In embodiments, the neural cell forms part of an organism. In embodiments, the organism is a bacterial cell or mammalian cell (e.g., human, murine, bovine, feline, canine).

Methods of Treatment

[0078] In another aspect, a method of treating a neurological disease in a subject in need thereof is provided. The method includes (i) administering to a subject a therapeutically effective amount of a recombinant nucleic acid encoding an exogenous mechanotransduction polypeptide (e.g., TRP-4). In step (ii) an ultrasonic wave is applied to the subject, resulting in a change in TRP-4 conductance, i.e., cation influx. In one embodiment, the methods treat a cardiac disease by enhancing cardiac muscle activity or neurological disease by altering neural activity in the subject. In embodiments, the neurological disease is Parkinson Disease, depression, obsessive-compulsive disorder, chronic pain, epilepsy or cervical spinal cord injury. In embodiments, the neurological disease is retinal degeneration or atrial fibrillation. In embodiments, the mechanotransduction polypeptide is a transient receptor potential channel-N (TRPN) polypeptide or homolog thereof. In embodiments, the mechanotransduction polypeptide is TRP-4 or a functional portion or homolog thereof. In embodiments, the method further includes before the applying of step (ii) administering to the subject an ultrasound contrast agent. In embodiments, the ultrasound contrast agent is a microbubble. In embodiments, the microbubble has a diameter of about 1 μm to about 6 μm , and is injected into the body (e.g., the brain) where it enhances ultrasound stimulation.

EXAMPLES

[0079] Reliable activation of identified neurons, particularly those in deeper brain regions remains a major challenge in neuroscience. Here, Applicants demonstrate low intensity ultrasound as a non-invasive trigger to activate neurons in the nematode, *Caenorhabditis elegans*. Applicants show that neuron-specific misexpression of TRP-4, the pore-forming subunit of a mechanotransduction channel, activates those cells in response to ultrasound stimuli and initiates behavior. Applicants suggest that this method can be broadly used to manipulate cellular functions in vivo.

[0080] To probe the effects of ultrasound on neuronal function, Applicants chose the nematode *C. elegans*, with its small nervous system consisting of just 302 neurons (J. G. White, E. Southgate, J. N. Thomson et al., *Phil. Transact. R. Soc. Lond. B* 314, 1 (1986)), and strong correlations between individual neurons and robust behaviors (M. de Bono and A. V. Maricq, *Annu Rev Neurosci* 28, 451 (2005); C. I. Barg-

mann, *WormBook*, 1 (2006); R. O'Hagan and M. Chalfie, *Int Rev Neurobiol* 69, 169 (2006)).

Example 1: Imaging Setup Delivers Ultrasound Waves to Animals

[0081] To investigate the role of ultrasound on *C. elegans* behavior, Applicants developed a novel imaging setup (FIG. 1A). Low intensity ultrasound was generated from a transducer and focused onto an agar plate where animals were corralled into a small area using a copper solution (FIG. 1i). Applicants' setup allowed for the ultrasound wave to be focused to a 1 mm diameter circular area at the agar surface (FIGS. 8A-8C). The whole setup was placed in a large tank filled with water to facilitate uniform transduction of the ultrasound wave. Previous studies have shown that at high ultrasound intensities (>2.5 MPa) water vapor bubbles would form spontaneously and collapse rapidly, initiating shockwaves that would compromise the integrity of cell membranes (termed "cavitation") (C. K. Holland and R. E. Apfel, *J Acoust Soc Am* 88 (5), 2059 (1990); S. Bao, B. D. Thrall, and D. L. Miller, *Ultrasound Med Biol* 23 (6), 953 (1997)). Applicants confirmed these results in Applicants' assay setup and also observed damage to animals at these high ultrasound intensities (data not shown). Applicants chose to focus on low intensity ultrasound to eliminate these damaging effects and found that at these intensities ultrasound had no effect on animal behavior (FIGS. 1D and 1E). The entire setup was placed in a large tank filled with water to facilitate uniform transduction of the ultrasound wave. Depending on solution or tissue gas concentrations, high ultrasound peak negative pressures (>2.5 MPa) can create inertial cavitation with the resulting shockwaves compromising the integrity of cell membranes. Consistently, Applicants observed that animals exposed to multiple pulses of high ultrasound pressures were unable to maintain their normal body posture (FIG. 12). Therefore, Applicants chose to use low-pressure ultrasound, which does not cause these damaging effects, to stimulate animal behavior.

[0082] Applicants used data from a previous study to estimate the mechanical deformation of the low intensity ultrasound wave (A. P. Brysev, A. F. Bunkin, R. V. Klopotov et al., *Opt. Spectrosc.* 93 (2), 282 (2002)). Applicants estimate that at this intensity, the ultrasound wave is likely to pass through *C. elegans* causing a mechanical deformation of 0.005 nm, and hypothesized that this small change is unlikely to influence cellular functions in vivo. This hypothesis is consistent with previous studies, which have shown that mechanical changes of this magnitude do not modify either neurons or non-neurons (S. Ito, H. Kume, K. Naruse et al., *Am J Respir Cell Mol Biol* 38 (4), 407 (2008); K. Shibasaki, N. Murayama, K. Ono et al., *J Neurosci* 30 (13), 4601 (2010)).

[0083] Moreover, Applicants found that a single 10 ms duration ultrasound pulse of 2.25 MHz and peak negative pressures below 0.9 MPa had no effect on animal behavior. The mechanical disturbances of the fluid and tissue in the ultrasound focal zone take the form of compression and expansion deformations as well as bulk tissue distortions caused by acoustic radiation forces, but at low-pressures they were not large enough to influence *C. elegans* locomotion. Previous studies have shown that ultrasound waves can cause temperature changes in the focal zone. Applicants first estimated the temperature increase as a result of ultrasound exposure. In a previous study, a continuous 1.1 MHz ultra-

sound pulse with a peak negative pressure of 2.6 MPa increased the temperature of the surrounding media at the rate of 35° C./sec. Using these data, Applicants estimated that the temperature increase around the worms on the agar surface to be 0.04° C. for single ultrasound pulse at 0.9 MPa. Moreover, Applicants directly measured the magnitude of temperature change on the agar surface using a miniature thermocouple and found that an ultrasound peak negative pressure of 0.7 MPa caused a temperature increase of less than 0.1° C. This is a temperature stimulus that animals including *C. elegans* are unlikely to detect. Together, these results show that *C. elegans* is unlikely to respond to the temperature and mechanical changes induced by the low-pressure ultrasound wave.

Example 2: Microbubbles Amplify the Mechanical Deformation of the Ultrasound Wave

[0084] To amplify the ultrasound wave, Applicants included gas-filled microbubbles in Applicants' assay (FIG. 1C). Previous studies have shown that the majority of the ultrasound energy propagates through water and soft tissue as a longitudinal wave with alternating compression and rarefaction phases. These two phases create pressures that are alternately higher and lower than the ambient pressure level respectively. Applicants designed the microbubbles to respond to the mechanical deformations induced by an ultrasound pulse. Applicants filled the microbubbles with a stabilizing mixture of perfluorohexane and air that allows the compression and rarefaction phases of the ultrasound wave to shrink and expand the microbubbles from one half to four times their original diameters in a process known as stable cavitation. This occurs at the driving frequency of the underlying ultrasound pulse. Applicants found that animals showed a dramatic response to ultrasound when surrounded by microbubbles (FIGS. 1D and 1F). When the ultrasound wave was focused on the head of a worm, the animal immediately initiated a backward movement (termed "reversal") followed by a high-angled turn (labeled "omega bend") (FIGS. 1D and 13). These behaviors were scored as previously described (FIG. 13) (J. M. Gray, J. J. Hill, and C. I. Bargmann, *Proc Natl Acad Sci USA* 102 (9), 3184 (2005)) and quantified as shown (FIGS. 1E and 1F). The animal's behavioral responses were correlated with the intensity of the ultrasound wave (FIG. 1F) and the size of the microbubbles (FIG. 14). Applicants suggest that microbubbles (1-3 μm in diameter, mixed size) are likely to resonate with the 2.25 MHz ultrasound pulse causing large mechanical fluctuations around the animal and in turn, reversal behavior. To probe how microbubbles transduce the ultrasound wave and modify animal behavior Applicants analyzed microbubbles labeled with fluorescent DiO (FIG. 1C). Applicants found that microbubbles are evenly distributed around the animal and upon ultrasound stimulation some are destroyed, while others fuse and yet others move (FIGS. 4A-4C). These results suggest that these fluctuations in microbubbles around the animal are sufficient to initiate reversal behavior. Ultrasound waves have been previously shown to cause an increase in temperature in the focal zone (C. H. Farny, R. G. Holt, and R. A. Roy, *Biomedical Engineering, IEEE Transactions on* 57 (1), 175 (2010)). Using this dataset, Applicants estimate that the low-intensity ultrasound pulse (2.25 MHz) might cause a temperature increase of 0.04° C. on the agar surface, a stimulus that animals including *C. elegans* are unlikely to detect (I. Mori,

H. Sasakura, and A. Kuhara, *Curr Opin Neurobiol* 17 (6), 712 (2007); D. A. Clark, C. V. Gabel, H. Gabel et al., *J Neurosci* 27 (23), 6083 (2007)). Taken together, these results suggest that mechanical distortions around the worm transduce the ultrasound stimulus and initiate behavioral changes.

Example 3: TRP-4 Stretch Sensitive Ion Channels Sensitize Neurons to Ultrasound

[0085] Applicants hypothesized that ultrasound is a mechanical stimulus that require specific mechanotransduction channels to transduce the signals in individual neurons. Applicants tested the ability of TRP-4, a pore forming cation-selective mechanotransduction channel (L. Kang, J. Gao, W. R. Schafer et al., *Neuron* 67 (3), 381 (2010); W. Li, Z. Feng, P. W. Sternberg et al., *Nature* 440 (7084), 684 (2006)), to transduce this ultrasound induced mechanical stimulus. This channel is specifically expressed in a few *C. elegans* neurons, the four CEPs (CEPDL, CEPDR, CEPVL and CEPVR) and the two ADE (ADEL and ADER) dopaminergic neurons and the DVA and DVC interneurons (L. Kang, J. Gao, W. R. Schafer et al., *Neuron* 67 (3), 381 (2010); W. Li, Z. Feng, P. W. Sternberg et al., *Nature* 440 (7084), 684 (2006)). TRP-4 is both necessary and sufficient to generate mechanoreceptor currents in CEP neurons. Applicants found that animals missing TRP-4 have reduced responses to specific intensities (0.41 and 0.47 MPa peak negative pressure) of ultrasound stimulation, which suggests that this channel is required to generate reversals (FIG. 2A). In contrast, *trp-4* mutants do not show any significant change in their omega bend behaviors upon ultrasound stimulation (FIGS. 5A-5F). At higher intensities, *trp-4* mutants have similar responses compared to wildtype, which suggests that there is an alternate pathway that detects ultrasound at these intensities. Collectively, these results suggest that TRP-4 might be activated in response to ultrasound with peak negative pressure levels less than 0.5 MPa and modifies neurons involved in generating small and large reversals.

[0086] To test whether ultrasound sensitivity could be conferred to additional neurons, Applicants analyzed the behavior of transgenic animals misexpressing TRP-4 in specific chemosensory neurons. Applicants initially misexpressed this channel in ASH, a well-studied polymodal nociceptive neuron (M. A. Hilliard, C. Bergamasco, S. Arbucci et al., *Embo J* 23 (5), 1101 (2004)), whose activation leads to reversals and omega bends (Z. V. Guo, A. C. Hart, and S. Ramanathan, *Nat Methods* 6 (12), 891 (2009)). Consistently, Applicants found that ASH expression of TRP-4 generated a significant increase in reversals at ultrasound intensity with a peak negative pressure of 0.47 MPa (FIG. 2B). Moreover, Applicants found that these ASH::*trp-4* transgenics do not show any change in their omega bend responses (FIG. 5), confirming that this channel specifically modifies the reversal neural circuit. Next, Applicants tested the effects of TRP-4 misexpression on function and behavior of the AWC sensory neuron. Previous results have implied that AWC activation is correlated with an increase in the animal's ability to generate reversals (S. H. Chalasani, N. Chronis, M. Tsunozaki et al., *Nature* 450 (7166), 63 (2007)). Applicants found that animals misexpressing TRP-4 in AWC neurons also initiated significantly more large reversals at the same ultrasound intensity of 0.47 MPa peak negative pressures, but not omega bends (FIGS. 2C and 5). To test whether ultrasound could directly stimu-

late AWC neurons, Applicants recorded the activity of these neurons in animals expressing the calcium indicator, GCaMP3 (L. Tian, S. A. Hires, T. Mao et al., *Nat Methods* 6 (12), 875 (2009)). Consistent with Applicants' behavioral data, Applicants found that ultrasound stimulation activated AWC neurons (FIGS. 2D-2F). Also, Applicants find that AWC responses are significantly reduced in the absence of microbubbles, which suggests that the ultrasound signals need to be amplified before they can modify neuronal functions (FIGS. 6A-6D). Consistent with the behavior data, Applicants observe that AWC neurons expressing TRP-4 show a significant increase in their activity a few seconds (t=12 to t=17 seconds) after the ultrasound stimulus (FIG. 2F). Both wild-type AWC neurons and those misexpressing TRP-4 showed a response lasting about 2-3 seconds immediately upon exposure to a single ultrasound pulse in the presence of microbubbles. However, Applicants also observed that AWC neurons misexpressing TRP-4 show a significant increase in their activity starting at 7 seconds after ultrasound exposure (t=12 seconds in FIG. 5F) and lasting for at least 5 seconds, which is not observed in wild-type neurons. This sustained increase in AWC calcium levels likely represents the activity of TRP-4, which could potentiate calcium entry into the neuron via other calcium channels. Interestingly, large reversals take approximately 10-20 seconds to complete, a time window where Applicants also observe sustained AWC calcium activity in the AWC::trp-4 transgenics. The sustained AWC calcium activity observed in these AWC::trp-4 transgenics is likely correlated with the increased frequency of large reversals generated by these animals after ultrasound stimulation. Taken together, these results show that TRP-4 channels are sensitive to low-pressure ultrasound, and ectopic expression of these channels in sensory neurons causes correlated changes in neuronal activity and behavior.

[0087] Interestingly, FLP neurons do not respond to ultrasound (FIG. 16). Microbubbles are present in all FLP recordings.

Example 4: Newly Identified Roles for PVD Sensory and AIY Interneurons in Generating Behavior in the Presence of Microbubbles

[0088] To test Applicants' approach of analyzing neuronal function by misexpressing TRP-4, Applicants probed the functions of poorly understood PVD neurons (FIG. 3A). PVD neurons have extensive dendritic processes that are regularly spaced and non-overlapping and cover most of the animal, excluding the head and the neck (A. Albeg, C. J. Smith, M. Chatzigeorgiou et al., *Mol Cell Neurosci* 46 (1), 308 (2011)). Applicants find that expressing TRP-4 in PVD neurons leads to a significant decrease in their reversal responses upon ultrasound stimulation (FIG. 3B). Applicants hypothesize that PVD neurons suppress reversals and misexpressing TRP-4 channels activates these neurons upon ultrasound stimulation, which in turn suppresses reversals. To test Applicants' hypothesis Applicants monitored PVD neuron activity in response to ultrasound stimulation. Applicants find that PVD neurons are more likely to be activated when the animal is moving backward than when moving forward (FIGS. 7A-7C). Also, Applicants find a strong correlation between PVD activity and animal movement. In particular, Applicants find that PVD neurons reach their maximum response when the animal has stopped reversing (FIGS. 3C and 3D). These results suggest that expressing

TRP-4 in PVD neurons activates them upon ultrasound stimulation and causes premature suppression of backward movement leading to fewer reversals. See also FIG. 17.

[0089] Applicants' studies show that *C. elegans* neural circuits can be probed by combining ultrasound stimulation with microbubbles that amplify the mechanical deformations. Specifically, Applicants find that upon activation ASH and AWC sensory neurons increase in reversals, while activating PVD neurons suppresses reversals (FIG. 3E). Interestingly, Applicants identify that persistent AWC neural activity might be required to drive reversal behavior providing a correlation between a distinct AWC neuronal activity pattern and whole animal behavior. Also, Applicants define a novel role for PVD neurons in suppressing reversal behavior. Taken together, these results and other studies (D. Tobin, D. Madsen, A. Kahn-Kirby et al., *Neuron* 35 (2), 307 (2002)) show that TRP channels can be used to manipulate neuronal functions and thus provide insight into how neural circuits transform environmental changes into behavior.

[0090] Applicants then tested whether this approach can manipulate the function of an interneuron, whose processes do not contact the external cuticle of the animal. Applicants misexpressed TRP-4 in AIY interneurons, which are at least 25 μ m from the cuticle, and analyzed the behavior of these animals upon ultrasound stimulation. Optogenetic studies have previously shown that activating AIY interneurons reduces turns. In contrast, Applicants find that AIY::trp-4 transgenics are significantly more likely to initiate high-angled omega bends upon ultrasound stimulation (two independent transgenics). It is possible that expressing TRP-4 in AIY neurons has altered that neuron's function, leading to increased turns. However, animals with genetically altered AIY function have been shown to have increased turns in a local search assay. Applicants found that these AIY::trp-4 transgenics did not show any defects in local search (FIG. 18) confirming that the AIY neurons were not altered in these animals. These data suggest that AIY can initiate different behaviors based on type of stimulation, ultrasound or light.

[0091] To confirm whether ultrasound stimulus is activating AIY interneurons, we used calcium imaging. AIY neural activity is typically measured from a bulb in the AIY neurite. Consistent with previous observation, Applicants found that AIY is a noisy neuron with a number of transients during recordings (FIG. 11). Applicants collected from a number of AIY recordings from wild-type animals and defined the relevant transient. Applicants counted all neurons that responded within a 5.5 second after the ultrasound pulse as responders. Using this criteria, AIY neurons in wild-type animals did not show a significant response to ultrasound stimulus (4/29) (FIGS. 11A and 11B). In contrast, Applicants observed a significant number of AIY neurons in AIY::trp-4 transgenics (11/28 animals) had a positive response (FIGS. 11E and 11F). In contrast, Applicants suggest that increased proportion of AIY responders in the AIY::trp-4 transgenics suggests that ultrasound stimulus activates AIY interneurons. These results show that mechanical deformations from the ultrasound-microbubble interaction can penetrate at least 25 μ m into the worm and influence the function of AIY interneurons. Moreover, Applicants find that misexpressing TRP-4 can influence both reversal and omega bend neural circuitry, suggesting that the sonogenetic approach is broadly applicable for manipulating circuit activity. Further, these results show that AIY interneurons likely have at least

three activity states with one suppressing turns, one promoting forward turns (as revealed by optogenetic stimulation) and one increasing omega turns (as revealed by ultrasound stimulation). These studies validate the approach of using sonogenetics to reveal novel roles for both PVD and AIY neurons in modifying turn behavior.

[0092] These studies show that *C. elegans* neural circuits can be probed by combining low-pressure ultrasound stimulation with microbubbles that amplify the mechanical deformations. Specifically, Applicants found that *C. elegans* are insensitive to low-pressure ultrasound but respond when surrounded by microbubbles. Applicants found that animals missing the TRP-4 mechanosensitive ion channel have significantly reduced sensitivity to the ultrasound-microbubble stimulation, indicating that mechanosensitive ion channels play an important role in the mechanism of ultrasound stimulation. Applicants also found that misexpressing the TRP-4 mechanosensitive ion channel in specific neurons modifies their neural activity upon ultrasound stimulation, resulting in altered animal behaviors. Specifically, misexpressing TRP-4 in ASH and AWC sensory neurons results in an increase in large reversals, while activating PVD neurons suppresses this behavior. Applicants also defined novel roles for PVD neurons in suppressing reversal behavior and AIY neurons in stimulating omega bend behavior.

[0093] These novel methods provide new insights into the neural activity patterns that drive whole-animal behavior. Persistent AWC neural activity might drive reversal behavior, providing a correlation between a distinct AWC neuronal activity pattern and whole-animal behavior. Ultrasound stimulation may activate neurons with different kinetics than what has been seen using optogenetics. For example, activating AIY interneurons using light leads to an increase in forward turns, while using low-pressure ultrasound increases omega bend frequency. These studies indicate an alternative role for AIY in promoting omega bends. The stimulation of AIY interneurons demonstrates that this ultrasound technique can also be applied to deep internal neurons that do not contact the skin of the worm. Taken together, these results and other studies show that TRP channels can be used to manipulate neuronal functions and thus provide insight into how neural circuits transform environmental changes into precise behaviors.

[0094] In order to target smaller groups of neurons, the resolution of the ultrasound focal zone can be made smaller than the 1 mm diameter. Frequencies above 2.25 MHz can produce sub-millimeter focal zone spot sizes. Higher frequency ultrasound waves with their smaller focal zones are better suited to targets that are closer to the body surface as these waves do not penetrate tissues as well. One of the advantages of ultrasound is that small focal zones can be maintained noninvasively even in deep brain tissue. Outside the focal zone the peak negative pressures are significantly lower and are unlikely to result in neuron activation. This was seen on the agar plates where only worms that were in the focal zone responded to the ultrasound and nearby worms that were outside the focal zone did not. Another advantage of ultrasound is that this focal zone can be moved arbitrarily within the tissue to simulate different regions without any invasive procedures. With an electronically steerable ultrasound beam, multiple different targets can be noninvasively manipulated either simultaneously or in rapid succession. Moreover, the genetic targeting of the stretch

sensitive ion channels to individual neurons allows for targeting well below the resolution of the ultrasound focal zone.

[0095] The use of ultrasound as a non-invasive neuronal activator can be broadly applied to decode neural circuits in larger vertebrate brains with opaque skin and intact skulls. Ultrasound waves with peak negative pressures of <1 MPa have been shown to penetrate through skull and brain tissue with very little impedance or tissue damage. These results show that low-pressure ultrasound (with peak negative pressures 0.4-0.6 MPa) specifically activates neurons expressing the TRP-4 channel. Moreover, TRP-4 channels do not have mammalian homologs, therefore, it is unlikely that expressing these channels in the mammalian brain would produce deleterious effects. This suggests that neurons in diverse model organisms misexpressing this channel can be activated by ultrasound stimulation, allowing scientists to probe their functions in influencing animal behavior. Additionally, other mechanosensitive channels can be explored that may be more sensitive to mechanic deformations than TRP-4. Of particular interest are the bacterial MscL and MscS channels that have different sensitivities to membrane stretch and are selective for different ions. Moreover, TRP-4 and other channels may be mutated in and around the pore region in order to change their ion selectivity as well as their sensitivity to mechanical stretch to broaden the utility of this method.

[0096] Furthermore, if low-pressure ultrasound stimulation by itself does not activate TRP-4 expressing neurons, the mechanical signals can be amplified by gas-filled microbubbles. Perfluorohexane microbubbles are well-established for use as ultrasound contrast agents in vivo and can be administered intravenously to circulate throughout the vertebrate body including the brain. They can remain active for up to 60 minutes providing a time window where they could be used safely to amplify the ultrasound stimulus and manipulate neural activity. Microbubbles have been shown to undergo inertial cavitation when exposed to ultrasound with peak negative pressure of 0.58 MPa and higher. Using ultrasound pressure levels lower than this will prevent damage to the brain from the microbubble-ultrasound interaction. Moreover, Applicants used a third of the number of microbubbles that has been previously used to successfully image the mouse brain showing that the required microbubble dose would not be prohibitive for in vivo administration. These experiments show that in the presence of microbubbles the low pressure ultrasound stimulated the deep AIY interneurons expressing TRP-4. This result enables Applicants to estimate the distances at which the mechanical deformations from the ultrasound-microbubble interaction can effectively penetrate into brain tissue from the vasculature. The *C. elegans* cuticle is 0.5 μm thick and the AIY interneurons are 25 μm from the cuticle, indicating that the mechanical deformations traveled at least 25.5 μm into the worm. In contrast, the mammalian blood-brain barrier is 0.2 μm thick and the average distance of a neuron from a capillary is less than 20 μm . These distances are well within the range of the sonogenetic approach. With the data presented in this paper, the invention provides a novel, non-invasive approach to activate genetically targeted neurons using low-pressure ultrasound stimulation

[0097] The results described herein above were carried out using the following materials and methods.

[0098] Behavioral Assay

[0099] Well-fed young adults were placed on an empty NGM agar plate and corralled into a small area using a filter paper soaked in copper solution (200 mM). A solution (15 μ l) of microbubbles at a density of 3.8×10^7 /ml was added to the plate with worms. The worms were allowed to crawl around for 10 minutes before being stimulated by ultrasound. An animal was moved into the fixed ultrasound focal zone and stimulated with one pulse and the resulting reversal and omega bend response is recorded. Reversals with fewer than two head bends were identified as small, while those with more than two were counted as large. High-angled turns that lead to a significant change in the direction of an animal's movement were identified as omega bends (FIG. 9) (J. M. Gray, J. J. Hill, and C. I. Bargmann, *Proc Natl Acad Sci USA* 102 (9), 3184 (2005)). Data was analyzed using SPSS software v22 (IBM, NY).

[0100] Imaging

[0101] Transgenic animals expressing GCaMP in specific neurons were corralled into a small area by filter paper soaked in copper solution (as described above). The acetylcholine agonist and paralytic, tetramisole (J. A. Lewis, C. H. Wu, J. H. Levine et al., *Neuroscience* 5 (6), 967 (1980)), was used at 1.3 mM to paralyze the animals to facilitate recording neural activity. These animals were surrounded by a solution of microbubbles and stimulated using ultrasound intensities as described. Fluorescence was recorded at 10 frames/second using an EMCCD camera (Photometrics, Quant-EM) and resulting movies were analyzed using Metamorph software (Molecular Devices) as described (S. H. Chalasani, N. Chronis, M. Tsunozaki et al., *Nature* 450 (7166), 63 (2007)). Briefly, a fluorescence baseline was calculated using a 3-second window from $t=1$ to $t=4$ seconds. The ratio of change in fluorescence to baseline fluorescence was plotted in all graphs using custom MATLAB scripts (S. H. Chalasani, N. Chronis, M. Tsunozaki et al., *Nature* 450 (7166), 63 (2007)). For imaging PVD neurons, the concentration of the paralytic was reduced to 1 mM, which allowed these animals greater movement. Their motion along with the corresponding fluorescent intensity changes was captured and analyzed using Metamorph software.

[0102] Microbubble Synthesis

[0103] Microbubbles were made using a probe sonication technique as described (C. E. Schutt, S. D. Ibsen, M. J. Benchimol et al., *Small* (2014)). The stabilizing lipid monolayer consisted of distearoyl phosphatidylcholine (DSPC, Avanti Polar Lipids Inc., Alabaster, AL), distearoyl phosphatidylethanolamine-methyl polyethylene glycol (mPEG-DSPC 5 k, Layson Bio Inc., Arab, AL) and DiO (Biotium Inc., CA) in 85:13:2 molar ratio. The gas core of the microbubble consisted of perfluorohexane (Sigma-Aldrich, St. Louis, MO) and air mixture designed to attain stability under atmospheric pressure. Microbubbles were fractionated based on size by their settling time (FIGS. 10A-10C). Applicants chose a mixed size of microbubbles to maintain uniformity across all the experiments. The microbubbles were shown to be stable on agar plates sealed with parafilm for up to 24 hours.

[0104] Molecular Biology and Transgenic Animals

[0105] All *C. elegans* strains were grown under standard conditions as described (S. Brenner, *Genetics* 77 (1), 71 (1974)). Cell-selective expression of TRP-4 was achieved by driving the full-length cDNA under *odr-3* (AWC), *sra-6*

(ASH) and *des-2* (PVD and FLP) promoters. Germline transformations were obtained using the methods previously described (C. C. Mello, J. M. Kramer, D. Stinchcomb et al., *Embo J* 10 (12), 3959 (1991)). Complete information for all strains is listed in Table 1.

[0106] Temperature Estimation

[0107] Ultrasound stimulation in combination with microbubbles has been previously shown to cause temperature changes in the surrounding media (D. Razansky, P. D. Einziger, and D. R. Adam, *IEEE Trans Ultrason Ferroelectr Freq Control* 53 (1), 137 (2006)). The authors experimentally found a temperature increase of $14.11^\circ \text{C}/\text{sec}$ using a continuous 1.1 MHz ultrasound pulse with a peak negative pressure of 2.8 MPa (D. Razansky, P. D. Einziger, and D. R. Adam, *IEEE Trans Ultrason Ferroelectr Freq Control* 53 (1), 137 (2006)). In Applicants' assays, Applicants used pulses of 10 ms and a maximum peak ultrasound pressure at 0.8 MPa.

[0108] Applicants assumed a linear relationship between energy deposition and peak ultrasound pressure and calculated the temperature increase around the worms on the agar surface to be 0.04°C .

[0109] Ultrasound and Microscopy Setup

[0110] A schematic of the ultrasound and microscopy setup is shown in FIG. 1A and previously described (S. Ibsen, M. Benchimol, and S. Esener, *Ultrasonics* 53 (1), 178 (2013)). 10 ms, 2.25 MHz sine wave ultrasound pulse was generated with a submersible 2.25 MHz transducer (V305-Su, Panametrics, Waltham, MA) using a waterproof connector cable (BCU-58-6W, Panametrics, Waltham, MA). The resulting sound field was quantified using a needle hydrophone (HNP-0400, Onda Corporation, Sunnyvale, CA). An arbitrary waveform generator (PCI5412, National Instruments, Austin, TX) controlled by a custom designed program (LabVIEW 8.2, National Instruments, Austin, TX) was used to create the desired ultrasound pulse. The peak negative pressure of the ultrasound pulse was adjusted from 0 to 0.9 MPa using a 300 W amplifier (VTC2057574, Vox Technologies, Richardson, TX). Ultrasound attenuation through the plastic and agar was found to be minimal.

[0111] White light illumination was achieved by reflecting light from an external light source up at the petri dish using a mirror mounted at 45° . Behavior was captured using a high-speed camera (FASTCAM, Photron, San Diego, CA). Fluorescent images were collected using a Nikon 1-FL EPI-fluorescence attachment on the same setup as described. GCaMP imaging was performed using a $40\times$ objective and the images were captured using a Quanti-EM 512C camera (Photometrics, Tucson, AZ).

[0112] The petri dish was held at the air-water interface with a three-prong clamp mounted to an XYZ micromanipulator stage allowing the dish to be scanned in the XY plane, while maintaining a constant Z distance between the objective and ultrasound transducer. This alignment positioned the agar surface in the focal zone of the ultrasound wave.

TABLE 1

Table showing list of all strains and their genotypes		
Strain	Genotype	Description
N2	wild-type	WT
VC1141	<i>trp-4(ok1605)</i>	<i>trp-4</i> mutant

TABLE 1-continued

Table showing list of all strains and their genotypes		
Strain	Genotype	Description
IV133	ueEx71 [sra-6::trp-4, elt-2::gfp]	ASH expression of trp-4 in wildtype background
IV157	ueEx85 [odr-3::trp-4, elt-2::gfp]	AWC expression of trp-4 in wildtype background
CX10536	kyEx2595 [str-2::GCaMP2.2b, unc-122::gfp]	AWC imaging line in wildtype background
IV344	ueEx219 [odr-3::trp-4, unc-122::rfp], kyEx2595 [str-2::GCaMP2.2b, unc-122::gfp]	AWC imaging line with trp-4 expressed in AWC
IV242	ueEx150 [des-2::trp-4; elt-2::gfp #3]	PVD expression of trp-4 in wildtype background
IV243	ueEx151 [des-2::trp-4; elt-2::gfp #4]	PVD expression of trp-4 in wildtype background

TABLE 1-continued

Table showing list of all strains and their genotypes		
Strain	Genotype	Description
IV219	ueEx134 [des-2::GCaMP3, unc-122::rfp]	PVD and FLP imaging line in wildtype background
IV494	ueEx307 [ttx-3::trp-4; elt-2::gfp #3]	AIY expression of trp-4 in wildtype background
IV495	ueEx308 [ttx-3::trp-4; elt-2::gfp #4]	AIY expression of trp-4 in wildtype background
CX8554	kyEx1489 [ttx-3::GCaMP1.0, unc-122::gfp]	AIY imaging line in wildtype background
IV646	kyEx1489[ttx-3::GCaMP1.0, unc-122::gfp]; ueEx440[ttx-3::trp-4, unc-122::rfp]	AIY imaging line with trp-4 expressed in AIY

SEQUENCE LISTING

Sequence total quantity: 3

SEQ ID NO: 1 moltype = AA length = 1924
 FEATURE Location/Qualifiers
 source 1..1924
 mol_type = protein
 organism = Caenorhabditis elegans

SEQUENCE: 1

```

MDSPRGGILG RALREASQST RQENDVDMQDQ VVVRQMNRDY GGSRRTOQNP HTSQPGPSHV 60
SIVNVPERGG PTSSTSTTHE TEHTAHRTEG GRFIRRRRQS REVTTTTTRP YDPAPPTQTR 120
TSSGSTVNGW GENRPKSADE EIKRRRRSGG GILSRGLREM NKMVEELEQA SEEPSTRKGI 180
LGTALKDMEG TTYQKIYRKR EETPKRSRSF DDQEMSNRVG MIEHLLRDKD PLELQQLGLT 240
DLLTTDTIPT DRPPLRRSST HLQIGKNSRI IFVFKQPSRD SVTPPDRLLG KPLFRESLTS 300
HASSHEEMSS EDLAMADPQT KILYFAKRDE WANVESEIET IKRSDFSMAD NHGFTAFLLA 360
VKAGKDQIVD KMIRK GARVD YSTKDGRNAT HIAAMYSGVE TLELILKRYE ELLRKGAGPK 420
KQLAIHVACE RSKKAFPIV KRILEDTDQR MAEDGDGSLP IHLAFKFGNV NIVELLLSGP 480
SDEQTRKADG NGDTLLHLAA RSGNIEAVRT AIAAGCDNAN VQNRVGRTPLE HEVAEVDQDN 540
MLKIMFKLRA DANIHKEDK TPVHVAERG DTSMVESLID KFGGSIRART RDGSTLLHIA 600
ACSGHTSTAL AFLKRGVPLF MPNKKGALGL HSAAAAGFND VVKMLIARGT NVDVTRTDNY 660
TALHVAVQSG KASVVTLLG SGADIVHVGK ELGQTLALHIA ASLNGAESRD CAMMLLKSGG 720
QPDVAQMDGE TCLHIAARSG NKDIMRLLLD ENADSKISSK IGETPLQVAA KSCNFEAASM 780
ILKHLSEVLT QEQLKEHVNH RTNDGFTALH YAAEIEQRQL HFPGEDAKLV NLLIDYGGMV 840
EMPSLNANET AMHMAARSGN QAVLLAMVNK IGAGAVQIVQ NKQSKNGWSP LLEACARGHS 900
GVANILLKHH ARIDVFDEMG RTALHLAFAFN GHLSLVHLLL QHKAFVNSKS KTGEAPLHLA 960
AQHGHVKVVN VLVQDHGAAL EAITLDNQTA LHFPAKFGQL AVSQTLLALG ANPNARDDKG 1020
QTPLHLAAEN DFPDVKLFL KMRNNNRSVL TAIDHNGFTC AHIAAMKGS L AVVRELM MID 1080
KPMVIQAKTK TLEATTLHMA AAGGHANIVK ILLENGANAE DENSHGMTAL HLGAKNGFIS 1140
ILEAFDKILW KRCSRKTGLN ALHIAAFYGN SDFVNEMLKH VQATVRSEPP IYNHHVNKEF 1200
STEYGFPLH LAAQSGHDSL VRMLLNQGVQ VDATSTTMNV IPLHLAAQQG HIAVVGMLLS 1260
RSTQQQAKD WRGRTPHLA AQNGHYEMVS LLIAQGSNIN VMDQNGWTGL HFATRAGHLS 1320
VVKLFIDSSA DPLAETKEGK VPLCFAAAHN HIECLRFLK QKHDTHQLME DRKFIFDLMV 1380
CGKTNDNEPL QEFILQSPAP IETAVKLSAL YRDMSEKEKE RAKDLLNAV FSENMAVELL 1440
GITATEYNAA LLLKAKDNRG RPLLDVLIEN EQKEVVSYS VQRYLTVWV ARVDWSFGKF 1500
VAFSLFVLIC PPAWFYFSLP LDSRIGRAPI IKFVCHIVSH VYFTILLTIV VLNI THKMYE 1560
VTSVVPNPVE WLLLLWLSGN LVSELSTVGG GSGLGIVKVL ILVLSAMAIA VHVLAFLPA 1620
VFLTHLDNDE KLHFARTMLY LKNQLFAFAL LFAFVEYLDL LTVHHLFGPW AIIIRDLMYD 1680
LARFLVILML FVAGFTLHVT SIFQPAYQPV DEDSAELMRL ASPSQTLEML FFSLFGLVEP 1740
DSMPPLHLVP DFAKIILKLL FGIYMMVTLI VLINLLIAMM SDTYQRIQAQ SDKEWKFGRA 1800
ILIRQMKNKS ATPSPINMLT KLIIIVLRVAV RNRLRCMTRK AQDDLRFEEEN IDAFSMGGGQ 1860
QGRQSPTNEG REGQOELGNS ADWNIETVID WRKIVSMYYQ ANGKLTGDRT KEDVDLAMA V 1920
PTSF 1924

```

SEQ ID NO: 2 moltype = DNA length = 5775
 FEATURE Location/Qualifiers
 source 1..5775
 mol_type = other DNA
 organism = Caenorhabditis elegans

SEQUENCE: 2

```

atggattcgc cacgtggcgg aatcctggga agagctttac gagaagcatc acaatcgact 60
aggcaagaaa atgatgttga tatggatcag gtacccttac ggcagatgaa cagggtattac 120
ggtggatcca ggaggactca gatgaatccc cacacctccc aacctggtcc atctcatgta 180
tcaattgtaa atgtcccaga acgcgaggga cccacatctt ccacatcaac cacacatgag 240
acagagcaca cggcacatag gacagagtcc gggaggttta tcagacgccg tcgccaatct 300

```

-continued

cgagaggta	ccaccacaac	cacaagacc	tatgacccg	ctcctccaac	ccagaccga	360
acaagctccg	gctcaacagt	aatggatgg	ggggagaatc	gaccgaaatc	tgctgatgag	420
gagatcaaac	ggcggcgaag	aagtggcggg	ggaatcctgt	ctcgcgggct	tcgagaaatg	480
aacaaaatgg	tggaagagt	ggagcaagca	agtgaagagc	caagtaccag	gaaggggaatt	540
ctgggtactg	cgtaaaggga	tatggaaggg	accacttatc	aaaagattta	caggaaaagg	600
gaggaaactc	ccaagcgtc	ccgttcattt	gacgatcagg	agatgtcgaa	tcgagtagga	660
atgatcgagc	acttgctccg	agacaaggat	cctttggagc	ttcagcagtt	gggattaacc	720
gacctcctca	ccaccgacac	catcccaact	gaccgaccac	ccctccgccg	atcctcgacc	780
catctccaaa	tcggaagaa	ctcacggatc	atcttcgctc	cgaaacaacc	atcccgtgat	840
tcagtcaccc	cgccggatcg	tcttctcggg	aaacctctgt	ttcgagagag	tctcacctcc	900
cacgcacgt	ctcatgagga	aatgtcgagt	gaggacttgg	caatggcggg	tctcagacg	960
aagattttgt	atctcgcgaa	gagagatgag	tgggcgaatg	tggagtctga	gatagagact	1020
atcaagcggg	gtgattttag	tatggctgat	aatcacggct	tcaccgcctt	cctcctagcc	1080
gtcaaagctg	gcaaggatca	aatcgtagac	aagatgatcc	gaaaaggtgc	tcgagtgagc	1140
tatagcacta	aagatggccg	taacgcgact	catattgccc	ccatgtactc	cggagttgaa	1200
actcttgagc	ttatcctcaa	gcgatactct	gagctgctcc	gaaaaggtgc	ggggcctaaa	1260
aagcagctgg	caatccatgt	ggcttgcgag	agaaaatcca	agaaagcatt	tccaattgtg	1320
aagcggattt	tggaagatac	tgatcaaaga	atggcagagg	atggggatgg	atccttgccg	1380
atacacttgg	cattcaagtt	tgggaatggt	aatattgtgg	agcttctgct	aagtgggctt	1440
tcggatgaac	aaaccaggaa	agctgatgga	aacggggata	ccttgcttca	tttggccgct	1500
cggagtggga	atatcgaagc	ggttcggaca	gcgattgccc	ctggatgtga	taatgcgaat	1560
gtgcagaata	gggtgggaag	gacgcgcgta	catgaggtag	ccgaagtcgg	agatcaaaa	1620
atgctaaaaa	tcatgttcaa	actccgcgcc	gacgccaaca	tccatgataa	ggaggacaag	1680
actccggtac	acgttgccagc	ggagcggagt	gacacttcca	tggtcgagtc	actaattgac	1740
aagtttgggt	gctcaattcg	cgctaggacc	cgtgatgggt	cgacgcttct	gcattattgct	1800
gcatgttcag	caccgactag	caccgactag	cgcttttga	agagaggagt	ccccctctt	1860
atgcccaca	aaaaaggagc	cctgggtctt	actccgcgag	cagctgctgg	cttcaacgac	1920
gtcgtcaaaa	tgctcattgc	tcggggact	aatgtagatg	tcctgacacg	agacaactac	1980
accgctctcc	acgtagcggg	tcaatctggc	aaggcttcgg	ttgtagagac	cctgctggga	2040
agtgggtcag	acattcatgt	gaagggcggg	gaactaggac	aaactgcact	gcacattgct	2100
gcaagcttga	atggagccga	gagtcgggat	tgtgcgatga	tgttgctgaa	aagtggaggg	2160
cagccggatg	ttgcacaaat	ggatggggag	acttgtctgc	atattgctgc	caggagtggg	2220
aataaggata	tcatgaggct	cctgcttgac	gagaacgccc	actcgaat	aagctcaaag	2280
atcggagaga	accctcca	ggtggccgcc	aagtcagca	atcttgaagc	agcatcaatg	2340
atcttgaagc	acccttcgga	agttctgacc	caagaacagc	ttaaggaaca	tgtcaatcat	2400
agaaccaatg	acggcttcc	agctcttcc	tacgcccgtg	aaatcgagca	gcgcccagtt	2460
cactttccag	gagaagatgc	caagctagta	aatctctgta	tcgactacgg	tggaaatggt	2520
gaaatgccat	cactcaatgc	aaatgagacg	gcgatgcata	tggcggcaag	atccggaaat	2580
caagctgtac	tcctggcgat	ggtcaataag	atcggagccc	gtgcggtgca	aatcgtgcag	2640
aacaagcaga	gcaagaacgg	atggtcaccg	ctgttggaa	catgtgccag	agggcattct	2700
ggagtggcga	atattttggt	gaagcaccac	gcccgtattg	atgtattcga	tgaaatgggc	2760
cgactgctc	tgacactggc	agctttcaat	ggcatctct	ccctgggtca	ccttcttctg	2820
cagcacaag	cattcgtgaa	cagtaaatcg	aaaaccggag	aggcaccgct	ccacttagca	2880
gctcagcatg	gtcatgtgaa	ggtgggtgaa	gtcctgggtg	aggatcatgg	tgacagcctg	2940
gaggcaatta	cgctggataa	tcagacagcc	ctccactttg	ccgcaaaat	cggtcagcta	3000
gctgtgagtc	aaacccttct	ggctctcggg	gcaaacccca	atgcacgtga	cgacaagggg	3060
caaaccctc	tccatctggc	agctgagaat	gacttccccg	acgttggtgaa	gctcttctctg	3120
aaaatgagaa	ataacaaccg	gagtggtgtg	accgcaattg	atcataatgg	atcacctgct	3180
gacatattg	ctgcatgaa	gggttcccta	gcccgtggct	gtgagcttat	gatgatcgac	3240
aagcctatgg	taatccaggc	aaagaccaa	acactggaag	ccactacact	tcatatggca	3300
gctgcccggg	gtcacgcgaa	cattgtgaa	attctgctgg	agaatggagc	aaacgcggaa	3360
gatgagaatt	cgacgcgaa	gactgctctc	caacttggcg	ccaaaaacgg	atcattatcg	3420
atcttggagg	cattcgataa	gatcctatgg	aaacgggtgt	cgagaaagac	cggctcctaac	3480
gctctccaca	tcgctgcggt	ctacggaaat	tcggatttctg	tcaatgaaat	gctcaagcac	3540
gtacaagcaa	cagtcgcttc	cgagccgccc	atctacaatc	accatgtcaa	taaggaattc	3600
tcaactgaat	acggcttcc	acctctccat	ttagccgctc	aaagtggaca	cgacagctct	3660
gtcgggatgc	ttctgaatca	gggagtgcaa	gtgacgcgca	ccagactac	aatgaacgtg	3720
atccccctc	atctggctgc	ccagcaaggc	cacatcgag	tggtaggaat	gctcctgtcc	3780
agatctactc	agcagcagca	cgccaaggat	tggagaggca	ggaccccgct	ccacctagcc	3840
gctcagaatg	gccactacga	gatggtctca	cttctcattg	ctcaggatc	taacatcaat	3900
gtcatggatc	agaatggctg	gactggctct	cacttggcca	ctcgtgcccg	gcacctgagt	3960
gtcgtcaagc	tgctcatcga	tgctcagcgc	gatccattgg	cggagaccga	ggagggcaaa	4020
gttccattgt	gctttgctgc	agctcataat	catatagaat	gtcttcgatt	cctcctgaaa	4080
cagaagcatg	acacacatca	attgatggaa	gatcggaaat	tcatattcga	cttgatgggt	4140
tgtggtaaaa	ccaatgacaa	tgagcctcta	caagagttaa	ttcttcaatc	acctgctcca	4200
attgagacgg	cagtcaagtt	gtccgcgctg	tacagagata	tgtcggagaa	ggagaaggag	4260
agggcgaagg	atctgttgaa	tgtggcagtg	ttcagtgaga	atatggctgt	ggagtgtta	4320
gggatcaccg	ccaccgaata	caatgccgct	cttctcctga	aggctaagga	caatcgaggc	4380
cggcccctac	tagatgttct	cattgaaaat	gagcagaaag	aagtagtctc	ctacgcgctc	4440
gtccaacgct	acctgacaga	agtatggact	gcccgtgtcg	actggctatt	cggaaagtct	4500
gtcgcattct	ccctctctgt	gctaatatgc	ccccggcat	ggttctactt	ctcacttcca	4560
ctggatagtc	ggatcggaa	agctccgatt	attaaaattg	tgtgccatat	cgtgtctcat	4620
gtctatttta	cgactactgct	gacaattgtg	gtgttgaata	ttacacataa	gatgtacgaa	4680
gtaacttcgg	tggttccaaa	ccctgtggaa	tggctcctgt	tgctctggct	ctctggaaat	4740
ctggctctccg	aactctccac	tgtcgggtgga	ggatctggcc	taggaatcgt	aaaggtccta	4800
atcctagtcc	tttccgcat	ggcagatagcc	gtccatgtcc	tagccttct	gctcccggca	4860

-continued

gtattcctaa	cccacctgga	taacgatgaa	aagctacatt	tgccecgac	aatgctttat	4920
ttgaaaaatc	aacttttcgc	ctttgccctg	ctatttgctt	ttgtagagta	cctggatttc	4980
ctgacagtgc	atcatttggt	cggtcctctg	gcgatcatta	ttcgagatct	aatgtatgat	5040
ttggcccgtt	tccttgatg	cctgatggtg	ttcgtggcgg	gcttcacact	ccacgtgacg	5100
agtatcttcc	agcctgccta	ccagcctgtc	gacgaggaca	gcgcecgact	gatgcgctctg	5160
gcctccccgt	ctcaaaccct	cgaaatgctc	ttcttctcgc	tcttcggact	cgctcgagccc	5220
gattcaatgc	ccccgctcca	tctagttcca	gattttgcaa	aaatcatctt	aaaacttcta	5280
ttcggaatct	acatgatggt	caccttgatt	gtgctgatca	acttgctgat	tgctatgatg	5340
tctgacacct	accaacgaat	tcaggcacag	tcggataagg	aatggaagt	cggaagagct	5400
attctgatca	gacagatgaa	taagaaaagc	gccacgcctg	cgccgataaa	tatgttaaca	5460
aagttgataa	ttgtgctgag	ggtagcctgg	cggaatcggt	tgagatgcat	gacccgaaaa	5520
gcccaagacg	atcttcgctt	cgaggagaac	atcgacgcgt	tctccatggg	tgccggccag	5580
caggggaaggc	aaagtccgac	caatgaagga	agagaaggcc	agcaagagct	tggttaactcg	5640
gctgactgga	acatcgagac	agtcacgcac	tggaagga	ttgtttcaat	gtactatcag	5700
gcgaatggga	agcttacaga	cgggcgaacc	aaagaggatg	tggtttggc	aatggcagta	5760
cctactagtt	tttag					5775

SEQ ID NO: 3
 FEATURE
 misc_feature
 source

moltype = DNA length = 5803
 Location/Qualifiers
 1..5803
 note = Description of Artificial Sequence: Synthetic
 polynucleotide
 1..5803
 mol_type = other DNA
 organism = synthetic construct

SEQUENCE: 3

gtttaaacgg	cgcgcccgtg	ccatgaatcc	tcacacttct	cagccagggc	caagccatgt	60
ctccattgtc	aacgtgccag	agcggggggg	accaacctcc	tcaacctcca	ccacacacga	120
gaccgaacac	acagcccatc	gcacagagag	cggccgattc	atccggagaa	ggcggcagtc	180
cagagaagtg	actaccacaa	ctaccaggcc	ctacgatcct	gcaccaccta	cccagacaag	240
aactagctcc	ggctccaccg	tgaatgggtg	ggcgagaaac	aggcccaagt	ctgccgacga	300
ggagatcaag	cgacggagaa	ggagtggcgg	gggaatcctg	tcaagagggc	tgagggagat	360
gaacaagatg	gtggaggaa	tggaaacaggc	ctctgaggaa	cccagtagac	gcaagggcat	420
tctggggact	gctctgaaag	acatggaggg	cacaacttac	cagaagatct	atcggaaaag	480
agaggaaacc	cctaaaaggt	ctcgcagttt	cgacgatcag	gagatgagca	acagagtggg	540
gatgatcgaa	catctgctga	gggacaagga	cccctggag	ctccagcagc	tgggactgac	600
agacctgctg	accacagata	ctattccaac	cgaccgacca	ccactgcgcc	gatctagtac	660
tcacctccag	atcggcaaga	acagccggat	catttctgtg	ccaaaacagc	ccagccgcca	720
ttccgtcact	cctccagacc	gactgctggg	caagcctctg	tttcgggagt	ctctgaccag	780
tcacgcctca	agccatgagg	aaatgtcctc	tgaagatctg	gctatggcgg	acccccagac	840
caagatcctg	tacttcgcca	aacgcgacga	gtgggcta	gtggagtccg	aaattgagac	900
aatcaagcgg	tcagacttca	gcatggccga	taaccacgga	ttcactgctt	ttctgctggc	960
agtgaagccc	ggcaaagacc	agattgtcga	taagatgatc	cgaaaaggag	cacgggtgga	1020
ttattctacc	aaggacgca	gaaacgccac	acatattgcc	gctatgtaca	gtggcgtgga	1080
gacactggaa	ctgatcctga	agaggtattc	agagctgtg	cgcaaaggcg	ccgggcttaa	1140
gaaacagctg	gcaatccacg	tggcctgcga	aaggaagtcc	aagaaagcct	tcccaattgt	1200
gaaaagaatc	ctggaggaca	ccgatcagag	gatggctgaa	gacggagatg	gctctctgcc	1260
cattcacctg	gcattcaa	ttgggaatgt	gaacatcgtc	gagctgctgc	tgtccggacc	1320
ttctgatgaa	cagactagaa	aggccgacgg	gaatggagat	accctgctgc	atctggcagc	1380
acgctccgga	aacattgagg	ctgtgcgaac	cgcaatcgcc	gccggatgcg	acaatgcca	1440
cgctgcagaac	cgcgtcgggc	gaacaccact	gcacgagggtg	gctgaagtgc	gagatcagaa	1500
tatgctgaag	attatgttca	aactgcgcgc	aacctgccaac	atccatgaca	aggaggataa	1560
aacaccagtg	cacgtcggcg	ctgagcaggg	cgatacttca	atgggtgaaa	gcctgattga	1620
caagtttggc	gggtccatcc	gagcccggac	aagagatggc	tctactctgc	tgcatatcgc	1680
agcctgttcc	gggcacacct	ctacagctct	ggcatctctg	aagagaggcg	tgctctgtt	1740
tatgccaat	aagaaaggag	ccctgggact	gcatagcgcc	gccgcccggc	gcttcaacga	1800
ctgggtcaag	atgctgatcg	ccaggggaac	aaatgtggat	gtcaggacc	gcgacaacta	1860
cacagccctg	cacgtggctg	tccagagtgg	caaggccagc	gtggctgaga	ctctgctggg	1920
cagcggagca	gatattcatg	tgaagggagg	agaactggga	cagaccgccc	tgacatcgc	1980
agccagcctg	aacggggcag	agtccaggga	ctgcgccatg	atgctgctga	aaagcggggg	2040
acagcctgat	gtggcccaga	tggacggaga	aacctgtctg	cacattgctg	cacggctctgg	2100
caataaggat	atcatgagac	tgctgctgga	cgagaacgcc	gatagtaaga	ttagttcaaa	2160
aatcggcgaa	actccactcc	aggtggccgc	taagtcttgc	aacttcgagg	cagccagtat	2220
gatcctgaaa	cacctgtcag	aagtgtgac	ccaggagcag	ctgaaggaac	acgtcaatca	2280
tagaactaac	gacggcttca	ccgccctgca	ttacgccgcc	gagattgagc	agaggcagct	2340
gcactttcca	ggggaggatg	ccaagctggg	gaatctgctg	atcgactatg	gcgggatggg	2400
cgagatgccc	tactgaaatg	caaacgaaac	cgccatgcac	atggccgcta	gaagcggaaa	2460
tcaggctgtg	ctgctggcaa	tggtaacaaa	gatggagacc	ggcgctgtgc	agatcgtcca	2520
gaataagcag	tcaaaaaacg	gctggagccc	actgctggag	gcatgtgcca	gggggcatag	2580
cggagtggct	aacattctgc	tgaagcacca	tgcacgcac	gacgtgttgc	atgaaatggg	2640
gcgaacagcc	ctgcacctgg	cagcctttaa	tggacacctg	agcctggtgc	atctgctgct	2700
ccagcacaaa	gccttcgtca	actcaaagag	caaaaccgga	gaggctccac	tgacacctggc	2760
tgacacagac	gggcatgtga	aggtgggtcaa	tgtgctggtc	caggatcatg	gggcccgtct	2820
ggaggccatc	acactggaca	accagactgc	tctgcacttc	gcagccaaat	ttggacagct	2880
ggccgtgagc	cagacactgc	tggctctggg	ggcaaatcct	aacgctagag	acgataaggg	2940
acagactcca	ctgcacctgg	ccgccgagaa	cgacttcccc	gatgtggtca	agctgtttct	3000

-continued

gaaaatgaga	aacaataaca	ggagcgtgct	gacagcaatt	gatcataatg	gcttcacctg	3060
cgcccacatc	gccgctatga	aaggcagcct	ggccgtggtc	agggagctga	tgatgattga	3120
caagcctatg	gtcatccagg	caaagactaa	aaccctggaa	gccactaccc	tgacacatggc	3180
agccgctgga	ggacacgcca	acattgtgaa	gatcctgctg	gagaatggcg	ctaacgcaga	3240
agatgagaac	agccacggca	tgaccgcact	gcacctggga	gccccaaaacg	gattcatttc	3300
catcctggag	gcctttgaca	agattctgtg	gaagcgggtg	agccggaaga	cagggctgaa	3360
tgctctgcat	atcgagcct	tctacggaaa	tagcgacttt	gtgaacgaga	tgctgaaaca	3420
cgtgcaggcc	actgtccgca	gtgaaccccc	tatctacaat	caccatgtga	acaaggagtt	3480
ctcaaccgaa	tatggcttta	cacctctgca	tctggctgca	cagagcgggc	acgattccct	3540
ggtgcggatg	ctgctgaatc	agggcgtgca	ggtcgagccc	accagcacia	ctatgaacgt	3600
gattccactg	catctggcag	ctcagcaggg	acacatcgca	gtggctggaa	tgctgctgtc	3660
ccgctctacc	cagcagcagc	acgctaagga	ttggcagagga	cggacacccc	tgcatctggc	3720
agcccagaac	ggccactatg	agatggtgag	cctgctgatt	gcccagggct	ccaatatcaa	3780
cgtgatggac	cagaatggct	ggactggact	gcatttcgca	acccgggctg	gacacctgag	3840
cgtggtcaag	ctgtttatcg	acagctccgc	cgatcctctg	gctgagacca	aggaaggcaa	3900
agtgccactg	tgcttcgctg	ccgcccacaa	ccatattgag	tgtctgagat	ttctgctgaa	3960
gcagaaacac	gatacacatc	agctgatgga	agataggaag	ttcatctttg	acctgatggt	4020
gtgcgcaaaa	actaatgaca	acgagcctct	ccaggagttc	atcctccagt	cccccgctcc	4080
tatcgagacc	gcagtgaaac	tgtctgccct	gtacagagat	atgagtgaaa	aggagaaaga	4140
aagggctaag	gacctgctga	atgtggcagt	ctttctgag	aacatggccg	tggaaactgct	4200
gggaattaca	gcaactgagt	ataatgctgc	actgctgctg	aaggccaaag	ataacagagg	4260
caggccactg	ctggacgtgc	tgatcgagaa	cgaacagaaa	gaggtggtca	gttacgcctc	4320
agtgcagaga	tacctgacag	aagtgtggac	tgctcgggtc	gatttggtcat	tccgggaagt	4380
tgtggcattc	agcctgtttg	tccctgattt	cccacccgcc	tggttctact	tttccctgcc	4440
actggactct	aggattggac	gcgcccccat	catcaagtcc	gtgtgccaca	tctgttccca	4500
tgtctacttt	accattctgc	tgacaatcgt	ggtcctgaat	atcactcaca	agatgtatga	4560
ggtgaccagc	gtggtcccaa	atcccgtcga	atggctgctg	ctgctgtggc	tgtccggcaa	4620
cctggtgagc	gagctgtcca	ccgtcggagg	aggcagcggg	ctgggaattg	tgaaggctct	4680
gatcctgggtg	ctgagcgcga	tggccatcgc	agtgcacgtc	ctggctttcc	tgctgcccgc	4740
agtgtttctg	actcatctgg	acaatgatga	gaagctgcac	ttcgcccgca	ccatgctgta	4800
cctgaaaaac	cagctgttcg	cctttgctct	gctgttcgct	tttgtggaat	atctggactt	4860
cctgacagtc	caccatctgt	ttgggccttg	ggctatcatt	attagggacc	tgatgtacga	4920
tctggcacgg	ttcctgggtca	tccctgatgt	gttcgtcgcc	ggcttcaccc	tgcatgtgac	4980
ctctatcttt	cagcccgctt	atcagcctgt	cgacgaggat	agtgtgtaac	tgatgctggc	5040
ggcaagtccc	tcacagacc	tggagatgct	gtcttttagt	ctggtcggcc	tggtggaacc	5100
cgattcaatg	cctccactgc	acctggtgcc	tgacttcgcc	aagattatcc	tgaactgct	5160
gtttgggatc	tacatgatgg	tgaccctgat	tgtcctgatc	aacctgctga	ttgctatgat	5220
gtctgataca	tatcagcgca	tccaggcaca	gagtgacaag	gagtggaaat	ttggccgggc	5280
cattctgatc	agacagatga	ataagaaatc	tgctacccct	agtccaatta	acatgctgac	5340
aaaactgatt	atcgtgctgc	gggtcgcttg	gcgcaatcga	ctgcggtgta	tgaccgaaa	5400
ggcccaggac	gatctgcggt	tccagggaaa	catcgacgct	ttttcaatgg	ggggaggaca	5460
gcagggacga	cagagcccta	ccaatgaggg	acgagaagga	cagcaggagc	tgggcaattc	5520
cgccgattgg	aacattgaaa	cagtgatcga	ctggagaaag	atcgtctcta	tgtactatca	5580
ggccaatggc	aaactgactg	acgggcgaac	caaggaggat	gtcgatctgg	ctatggctgt	5640
cctactttct	ttctgaattc	cgataacttg	tttattgcag	cttataatgg	ttacaaataa	5700
agcaatagca	tcacaaattt	cacaaataaa	gcattttttt	cactgcattc	tagttgtggt	5760
ttgtccaaac	tcacaaatgt	atcttatcat	gtctggcggc	cgc		5803

1. An imaging and ultrasound exposure system for altering the function of nerve cells that innervate a targeted tissue portion of a mammalian subject, the system comprising an imaging source and an ultrasound source comprising a ultrasound beam focusing transducer; wherein said ultrasound source of the system (i) delivers ultrasound to an exposed surface of the mammalian subject; (ii) provides ultrasound energy to the innervated targeted tissue portion, wherein the nerve cells that innervate the targeted tissue portion are configured to express an ultrasound-sensitive, exogenous TRP-N or TRP-4 transmembrane protein; and (iii) modulates the membrane potential of the nerve cells comprising the targeted tissue structure when the targeted tissue portion of the animal is exposed to ultrasound energy from the ultrasound source.

2. The system of claim 1, wherein the nerve cells are genetically modified to express the exogenous TRP-N transmembrane protein.

3. The system of claim 1, wherein the nerve cells are genetically modified to express the exogenous TRP-4 transmembrane protein.

4. The system of claim 2, wherein the exogenous TRP-N transmembrane protein is a non-mammalian TRP-N transmembrane protein.

5. The system of claim 3, wherein the exogenous TRP-4 transmembrane protein is a non-mammalian TRP-4 transmembrane protein.

6. The system of claim 1, wherein said TRP-4 polypeptide is encoded by a polynucleotide codon-optimized for expression in the cell.

7. The system of claim 1, wherein the exogenous TRP-4 polypeptide comprises an amino acid sequence having at least 85% identity to the amino acid sequence of SEQ ID NO: 1.

8. The system of claim 1, wherein the exogenous TRP-4 polypeptide comprises the amino acid sequence of SEQ ID NO: 1.

9. The system of claim 1, wherein the ultrasound source comprises an ultrasound transducer.

10. The system of claim 1, wherein the nerve cells are motor neurons, sensory neurons, or interneurons.

11. The system of claim 1, wherein the mammalian subject is a human subject.

12. The system of claim 1, wherein the nerve cells are transduced using a vector comprising a polynucleotide sequence encoding the exogenous TRP-N polypeptide or using a recombinant nucleic acid molecule encoding the exogenous TRP-N polypeptide.

13. The system of claim 1, wherein the nerve cells are transduced using a vector comprising a polynucleotide sequence encoding the exogenous TRP-4 polypeptide or using a recombinant nucleic acid molecule encoding the exogenous TRP-4 polypeptide.

14. The system of claim 1, wherein the ultrasound has a frequency of about 0.8 MHz to about 4 MHz.

15. The system of claim 1, wherein said ultrasound has a focal zone of about 1 cubic millimeter to about 1 cubic centimeter.

16. The system of claim 1, wherein the system operates with an ultrasound frequency of at least 20 kHz.

* * * * *