



US 20240076727A1

(19) **United States**

(12) **Patent Application Publication**  
**IAFRATE et al.**

(10) **Pub. No.: US 2024/0076727 A1**

(43) **Pub. Date: Mar. 7, 2024**

(54) **METHODS FOR DETERMINING A NUCLEOTIDE SEQUENCE CONTIGUOUS TO A KNOWN TARGET NUCLEOTIDE SEQUENCE**

Sep. 19, 2016, now Pat. No. 10,017,810, which is a continuation of application No. 13/793,564, filed on Mar. 11, 2013, now Pat. No. 9,487,828.

(71) Applicant: **THE GENERAL HOSPITAL CORPORATION**, Boston, MA (US)

(60) Provisional application No. 61/679,302, filed on Aug. 3, 2012, provisional application No. 61/645,364, filed on May 10, 2012.

(72) Inventors: **Anthony John IAFRATE**, Newton, MA (US); **Long Phi LE**, Boston, MA (US); **Zongli ZHENG**, Boston, MA (US)

**Publication Classification**

(51) **Int. Cl.**  
*C12Q 1/6855* (2006.01)  
*C12Q 1/6806* (2006.01)  
*C12Q 1/6874* (2006.01)

(73) Assignee: **THE GENERAL HOSPITAL CORPORATION**, Boston, MA (US)

(52) **U.S. Cl.**  
CPC ..... *C12Q 1/6855* (2013.01); *C12Q 1/6806* (2013.01); *C12Q 1/6874* (2013.01); *C12Q 2525/155* (2013.01); *C12Q 2525/161* (2013.01); *C12Q 2525/191* (2013.01); *C12Q 2531/113* (2013.01); *C12Q 2549/119* (2013.01)

(21) Appl. No.: **18/459,672**

(22) Filed: **Sep. 1, 2023**

**Related U.S. Application Data**


(63) Continuation of application No. 16/897,588, filed on Jun. 10, 2020, now Pat. No. 11,781,179, which is a continuation of application No. 15/984,612, filed on May 21, 2018, now Pat. No. 10,718,009, which is a continuation of application No. 15/269,448, filed on

(57) **ABSTRACT**


The technology described herein is directed to methods of determining oligonucleotide sequences, e.g. by enriching target sequences prior to sequencing the sequences.

**Specification includes a Sequence Listing.**


**TARGETED RNA & DNA SEQUENCING**




TARGETED RNA & DNA SEQUENCING




**FIG. 1**




**FIG. 2**



**FIG. 3A**



**FIG. 3B**



**FIG. 4A**

SLC34A2 Exon 13                      ROS1                      ALTERNATIVE SPLICING

```

AGGATGTCCTGTCAGGCTCCCTGAGACCTTTGATAACATAAACCATTAGCAGAGGCTCAGGCTGGAGTCCCAAATAAACCAGGCATTCCTCCAAATTACTAGA
TCAAGCCTCTGAGACCTTTGATAACATAAACCATTAGCAGAGGCTCAGGCTGGAGTCCCAAATAAACCAGGCATTCCTCCAAATTACTAGA
GAGACCTTTGATAACATAAACCATTAGCAGAGGCTCAGGCTGGAGTCCCAAATAAACCAGGCATTCCTCCAAATTACTAGA
GTCCGTCAAGGCTCTGAGACCTTTGATAACATAAACCATTAGCAGAGGCTCAGGCTGGAGTCCCAAATAAACCAGGCATTCCTCCAAATTACTAGA

GCAGGATGTCCTGTCAGGCTCCCTGAGACCTTTGATAACATAAACCATTAGCAGAGGCTCAGGATGATTTTGGATACCAGAAACAAGTTTCATCTACTACTA
GGCCGAGGAGGGCAGGATGTCCTGTCAGGCTCCCTGAGACCTTTGATAACATAAACCATTAGCAGAGGCTCAGGATGATTTTGGATACCAGAAACAAGTTTCATCTACTACTA

```

Exon 32                      Exon 34

**FIG. 4B**

```




> SAMPLE_03
FREQ          FUSION
164  SLC34A2+Ex13.f2 -- ROS1 - Ex32.f1
103  SLC34A2+Ex13.f2 -- ROS1 - Ex34.f1

> SAMPLE_15
FREQ          FUSION
41   EZR+Ex10.f2 --ROS1-Ex34.f1
8    EZR+Ex10.f2 --ROS1- Ex35.f1

```

**FIG. 4C**

TOTAL NUMBER OF BASES [Mbp]	49.19
• NUMBER OF Q17 BASES [Mbp]	43.86
• NUMBER OF Q20 BASES [Mbp]	41.58
TOTAL NUMBER OF READS	563,255
MEAN LENGTH [bp]	87
LONGEST READ [bp]	203



SEQUENCING RESULTS

FIG. 5

CCDC6 -- RET




FIG. 6

EM14 -- ALK




FIG. 7

TARGETED TNA SEQ




FIG. 8




FIG. 9

**METHODS FOR DETERMINING A  
NUCLEOTIDE SEQUENCE CONTIGUOUS  
TO A KNOWN TARGET NUCLEOTIDE  
SEQUENCE**

**CROSS-REFERENCE TO RELATED  
APPLICATIONS**

**[0001]** This application is a continuation under 35 U.S.C. § 120 of U.S. application Ser. No. 16/897,588, filed Jun. 10, 2020, which is a continuation under 35 U.S.C. § 120 of U.S. application Ser. No. 15/984,612 filed May 21, 2018, now U.S. Pat. No. 10,718,009, which is a continuation under 35 U.S.C. § 120 of U.S. application Ser. No. 15/269,448 filed Sep. 19, 2016, now U.S. Pat. No. 10,017,810, which is a continuation under 35 U.S.C. § 120 of U.S. application Ser. No. 13/793,564 filed Mar. 11, 2013 now U.S. Pat. No. 9,487,828B2, and which claims benefit under 35 U.S.C. § 119(e) of U.S. Provisional Application Nos. 61/645,364 filed May 10, 2012 and 61/679,302 filed Aug. 3, 2012, the contents of which are incorporated herein by reference in their entirety.

**GOVERNMENT SUPPORT**

**[0002]** This invention was made with federal funding under Grant No. 5R21CA161590 awarded by the National Institutes of Health. The U.S. government has certain rights in the invention.

**SEQUENCE LISTING**

**[0003]** The instant application contains a Sequence Listing which has been submitted in XML format via EFS-Web and is hereby incorporated by reference in its entirety. Said XML copy, created on Aug. 28, 2023, is named 030258-074968USC5\_SL.xml and is 102,006 bytes in size.

**TECHNICAL FIELD**

**[0004]** The technology described herein relates to methods of determining oligonucleotide sequences.

**BACKGROUND**

**[0005]** Target enrichment prior to next-generation sequencing is more cost-effective than whole genome, whole exome, and whole transcriptome sequencing and therefore more practical for broad implementation; both for research discovery and clinical applications. For example, high coverage depth afforded by target enrichment approaches enables a wider dynamic range for allele counting (in gene expression and copy number assessment) and detection of low frequency mutations, a critical feature for evaluating somatic mutations in cancer. Examples of current enrichment protocols for next generation sequencing include hybridization-based capture assays (TruSeq Capture™ Illumina; SureSelect Hybrid Capture™, Agilent) and polymerase chain reaction (PCR)-based assays (HaloPlex™, Agilent; AmpliSeq™, Ion Torrent™; TruSeq Amplicon™, Illumina; emulsion/digital PCR, Raindance). Hybridization-based approaches capture not only the targeted sequences covered by the capture probes but also near off-target bases that consume sequencing capacity. In addition, these methods are relatively time-consuming, labor-intensive, and suffer from a relatively low level of specificity. A PCR amplification based approach is simpler and faster but by

conventional design requires the use of both forward and reverse primers flanking the target loci. In particular, for detection of genomic rearrangements with unknown fusion partners, PCR is not applicable.

**SUMMARY**

**[0006]** The technology described herein is directed to methods of determining oligonucleotide sequences. In some embodiments, the methods described herein relate to enriching target sequences prior to sequencing the sequences.

**[0007]** In one aspect, the technology described herein relates to a method of determining the nucleotide sequence contiguous to a known target nucleotide sequence, the method comprising; (a) ligating a target nucleic acid comprising the known target nucleotide sequence with a universal oligonucleotide tail-adaptor; (b) amplifying a portion of the target nucleic acid and the amplification strand of the universal oligonucleotide tail-adaptor with a first adaptor primer and a first target-specific primer; (c) amplifying a portion of the amplicon resulting from step (b) with a second adaptor primer and a second target-specific primer; (d) sequencing the amplified portion from step (c) using a first and second sequencing primer; wherein the universal oligonucleotide tail-adaptor comprises a first ligatable duplex end and a second unpaired end; wherein the universal oligonucleotide tail-adaptor comprises a blocking strand and an amplification strand; wherein the blocking strand comprises a 5' duplex portion; wherein the amplification strand comprises an unpaired 5' portion, a 3' duplex portion, and a 3' T overhang; wherein the amplification strand comprises nucleic acid sequences identical to a first and second sequencing primers; wherein the duplex portions of the blocking strand and the amplification strand are substantially complementary and form the first ligatable duplex end comprising a 3' T overhang; wherein the duplex portion is of sufficient length to remain in duplex form at the ligation temperature; wherein the first target-specific primer comprises a nucleic acid sequence that can specifically anneal to the known target nucleotide sequence of the target nucleic acid at the annealing temperature; wherein the second target-specific primer comprises a 3' portion comprising a nucleic acid sequence that can specifically anneal to a portion of the known target nucleotide sequence comprised by the amplicon resulting from step (b), and a 5' portion comprising a nucleic acid sequence that is identical to a second sequencing primer and the second target-specific primer is nested with respect to the first target-specific primer; wherein the first adaptor primer comprises a nucleic acid sequence identical to a 5' portion of the first sequencing primer; and wherein the second adaptor primer comprises a nucleic acid sequence identical to a portion of the first sequencing primer and is nested with respect to the first adaptor primer.

**[0008]** In some embodiments, the blocking strand of the universal oligonucleotide tail-adaptor can further comprise a 3' unpaired portion which is not substantially complementary to the 5' unpaired portion of the amplification strand; and wherein the 3' unpaired portion of the blocking strand is not substantially complementary to or substantially identical to any of the primers. In some embodiments, the second adaptor primer can be nested with respect to the first adaptor primer by at least 3 nucleotides. In some embodiments, the portion of the amplification strand that comprises a nucleic acid sequence identical to a first and second sequencing

primers can be comprised, at least in part, by the 5' unpaired portion of the amplification strand.

**[0009]** In some embodiments, the first target-specific primer can further comprise a 5' tag sequence portion comprising a nucleic acid sequence of high GC content which is not substantially complementary to or substantially identical to any other portion of any of the primers. In some embodiments, the first target-specific primer can further comprise a 5' tag sequence portion comprising a nucleic acid sequence of high GC content which will not specifically anneal to any other portion of any of the primers or their complements at the annealing temperature. In some embodiments, the second adaptor primer can be identical to the full-length first sequencing primer. In some embodiments, the portions of the target-specific primers that specifically anneal to the known target can anneal specifically at a temperature of about 65° C. in a PCR buffer.

**[0010]** In some embodiments, the method can further; prior to step (a), the steps of: mechanically shearing the nucleic acid; subjecting the nucleic acid to end-repair; subjecting the nucleic acid to phosphorylation; and subjecting the nucleic acid to adenylation. In some embodiments, the sample can comprise genomic DNA. In some embodiments, the sample can comprise RNA and the method can further comprise a first step of subjecting the sample to a reverse transcriptase regimen. In some embodiments, the reverse transcriptase regimen can comprise the use of random hexamers.

**[0011]** In some embodiments, the known target sequence can be comprised by a gene rearrangement. In some embodiments, the gene rearrangement can be present in a nucleic acid selected from the group consisting of: genomic DNA; RNA; and cDNA. In some embodiments, the gene rearrangement can comprise an oncogene. In some embodiments, the gene rearrangement can comprise a fusion oncogene.

**[0012]** In some embodiments, the nucleic acid product can be sequenced by a next-generation sequencing method. In some embodiments, the next-generation sequencing method can comprise a method selected from the group consisting of: Ion Torrent™, Illumina, SOLiD™, 454™; Massively Parallel Signature Sequencing solid-phase, reversible dye-terminator sequencing; and DNA nanoball sequencing. In some embodiments, the first and second sequencing primers are compatible with the selected next-generation sequencing method.

**[0013]** In some embodiments, the method can comprise contacting the sample, or separate portions of the sample, with a plurality of sets of first and second target-specific primers. In some embodiments, the method can comprise contacting a single reaction mixture comprising the sample with a plurality of sets of first and second target-specific primers. In some embodiments, the plurality of sets of first and second target-specific primers can specifically anneal to known target nucleotide sequences comprised by separate genes. In some embodiments, at least two sets of first and second target-specific primers can specifically anneal to different portions of a known target nucleotide sequence. In some embodiments, at least two sets of first and second target-specific primers can specifically anneal to different portions of a single gene comprising a known target nucleotide sequence. In some embodiments, at least two sets of first and second target-specific primers can specifically anneal to different exons of a gene comprising a known nucleotide

target sequence. In some embodiments, the plurality of first target-specific primers can comprise identical 5' tag sequence portions. In some embodiments, the universal oligonucleotide tail-adaptor can further comprise a barcode portion. In some embodiments, multiple samples can each be contacted with a universal oligonucleotide tail-adaptor with a unique barcode portion and wherein the samples are pooled after step (a).

**[0014]** In some embodiments, each amplification step can comprise a set of cycles of a PCR amplification regimen from 5 cycles to 20 cycles in length. In some embodiments, the target-specific primers and the adaptor primers can be designed such that they will specifically anneal to their complementary sequences at an annealing temperature of from about 61 to 72° C. In some embodiments, the target-specific primers and the adaptor primers can be designed such that they will specifically anneal to their complementary sequences at an annealing temperature of about 65° C.

**[0015]** In some embodiments, the sample can comprise a biological sample obtained from a subject. In some embodiments, the sample can be obtained from a subject in need of treatment for a disease associated with a genetic alteration. In some embodiments, the disease can be cancer. In some embodiments, the sample can comprise a population of tumor cells. In some embodiments, the sample can comprise a tumor biopsy. In some embodiments, the cancer can be lung cancer.

**[0016]** In some embodiments, the known target sequence can be comprised by a disease-associated gene. In some embodiments, the known target sequence can be comprised by a gene rearrangement product in the sample. In some embodiments, the gene rearrangement product can be an oncogene.

**[0017]** In some embodiments, the known target sequence can comprise a sequence from a gene selected from the group of: ALK; ROS1; and RET. In some embodiments, at least one set of a first target-specific primer and a second target-specific primer can be selected from the group consisting of; SEQ ID NOs: 5 and 6; SEQ ID NOs: 7 and 8; SEQ ID NOs: 9 and 10; SEQ ID NOs: 11 and 12; SEQ ID NOs: 13 and 14; SEQ ID NOs: 15 and 16; SEQ ID NOs: 17 and 18; SEQ ID NOs: 19 and 20; SEQ ID NOs: 21 and 22; SEQ ID NOs: 23 and 24; SEQ ID NOs: 25 and 26; SEQ ID NOs: 27 and 28; SEQ ID NOs: 29 and 30; SEQ ID NOs: 31 and 32; SEQ ID NOs: 33 and 34; SEQ ID NOs: 35 and 36; and SEQ ID NOs: 37 and 38.

**[0018]** In some embodiments, the presence of a gene rearrangement of ALK in a sample obtained from a tumor in a subject can indicate that the tumor is susceptible to treatment with a treatment selected from the group consisting of: an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.

**[0019]** In some embodiments, the presence of a gene rearrangement of ROS1 in a sample obtained from a tumor in a subject can indicate that the tumor is susceptible to treatment with a treatment selected from the group consisting of: a ROS inhibitor; an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.

**[0020]** In some embodiments, the presence of a gene rearrangement of RET in a sample obtained from a tumor in

a subject can indicate that the tumor is susceptible to treatment with a treatment selected from the group consisting of: a RET inhibitor; DP-2490; DP-3636; SU5416; BAY 43-9006; BAY 73-4506 (regorafenib); ZD6474; NVP-AST487; sorafenib; RPI-1; XL184; vandetanib; sunitinib; imatinib; pazopanib; axitinib; motesanib; gefitinib; and withaferin A.

**[0021]** In one aspect, the technology described herein relates to a method of treating cancer, the method comprising; detecting, in a tumor sample obtained from a subject in need of treatment for cancer, the presence of one or more oncogene rearrangements according to the method described herein; administering a cancer treatment which is effective against tumors having any of the detected oncogene rearrangements. In some embodiments, a treatment selected from the group consisting of: an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684; can be effective against tumors having an ALK oncogene rearrangement. In some embodiments, a treatment selected from the group consisting of: a ROS1 inhibitor; an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684; can be effective against tumors having an ROS1 oncogene rearrangement. In some embodiments, a treatment selected from the group consisting of: a RET inhibitor; DP-2490; DP-3636; SU5416; BAY 43-9006; BAY 73-4506 (regorafenib); ZD6474; NVP-AST487; sorafenib; RPI-1; XL184; vandetanib; sunitinib; imatinib; pazopanib; axitinib; motesanib; gefitinib; and withaferin A; can be effective against tumors having an RET oncogene rearrangement.

**[0022]** In one aspect, the technology described herein relates to a method of determining if a subject in need of treatment for cancer will be responsive to a given treatment, the method comprising; detecting, in a tumor sample obtained from the subject, the presence of an oncogene rearrangement according to the method as described herein; wherein the subject is determined to be responsive to a treatment targeting an oncogene rearrangement product if the presence of the oncogene rearrangement is detected.

**[0023]** In some embodiments, when the presence of an ALK oncogene rearrangement is detected, the subject can be responsive to a treatment selected from the group consisting of: an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684. In some embodiments, when the presence of an ROS1 oncogene rearrangement is detected, the subject can be responsive to a treatment selected from the group consisting of: an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.

**[0024]** In some embodiments, when the presence of an RET oncogene rearrangement is detected, the subject will be responsive to a treatment selected from the group consisting of: a RET inhibitor; DP-2490; DP-3636; SU5416; BAY 43-9006; BAY 73-4506 (regorafenib); ZD6474; NVP-AST487; sorafenib; RPI-1; XL184; vandetanib; sunitinib; imatinib; pazopanib; axitinib; motesanib; gefitinib; and withaferin A.

**[0025]** In some embodiments, the cancer can be lung cancer.

#### BRIEF DESCRIPTION OF THE DRAWINGS

**[0026]** FIG. 1 depicts a schematic illustration of an example of library construction for targeted RNA and DNA sequencing. 1) A standard procedure of double-stranded cDNA synthesis is applied using total nucleic acid from FFPE specimen as starting material. Alternatively, the starting material may also be sheared gDNA. 2) After cleanup, the double stranded cDNA or gDNA is subjected to end-repair and dA tailing, directly followed by ligation of a half-truncated Y adapter without cleanup in between. 3) After SPRI cleanup, the ligated sample is subjected to 14 cycles of PCR amplification using multiplex gene specific primers (GSP1s) and Ion Torrent™ short length Forward primer A 5' 20-mer (A20), with annealing temperature at 65° C. 4) After a second SPRI cleanup, the sample is subjected to an additional 14 cycles of PCR amplification using multiplex nested gene specific primers (3' downstream of GSP1 and in the same direction) tagged with Ion Torrent™ Reverse primer (P1\_GSP2s) and Forward primer A (full length 30-mer), with annealing temperature at 65° C. 5) After a final third SPRI cleanup, the product is Ion Torrent™ library ready for downstream emulsion PCR and sequencing. Sample pooling may be carried out after Step 2.

**[0027]** FIG. 2 depicts the mapping results, demonstrating different primer (dots) extension products in one sequencing run of ROS1 sequences, including reads that corresponding to genomic DNA (spanning intron-exon boundaries), cDNA (spanning exons) and fusion cDNA (on exon 34, mapping on the fusion partner SLC34A2 not shown here). 91.8% specificity (127,446/138,787) was achieved.

**[0028]** FIG. 3A depicts a schematic presentation of nested primer targeting strategy using ROS1 as an example. The assay target panel includes a total of 17 pairs of GSP1s and GSP2s for ROS1, ALK and RET. FIG. 3B depicts a representation of the possible types of extension products using gDNA and cDNA templates.

**[0029]** FIG. 4A depicts the visualization of reads mapping using the Integrative Genomics Viewer. FIG. 4B depicts the sequence of two alternative spliced fusion sequences (SEQ ID NOs: 39-44, respectively, in order of appearance). FIG. 4C depicts a summary table reporting fusion transcripts in involved genes annotated with exon, frame, and fusion read coverage details.

**[0030]** FIG. 5 depicts the results of an example sequencing run. Sequences shown are ctaaggtaac (SEQ ID NO: 49), taaggagaac (SEQ ID NO: 50), aagaggattc (SEQ ID NO: 51), taccaagatc (SEQ ID NO: 52), cagaagggaac (SEQ ID NO: 53), ctgcaagttc (SEQ ID NO: 54) ttcgtgattc (SEQ ID NO: 55), ttccgataac (SEQ ID NO: 56), ctgaccgaac (SEQ ID NO: 57), tctcgaatc (SEQ ID NO: 58), taggtggttc (SEQ ID NO: 59) and tctaaccgac (SEQ ID NO: 60).

**[0031]** FIGS. 6 and 7 depict examples of the results of sequencing runs for ALK and RET sequences, respectively.

**[0032]** FIG. 8 depicts a schematic presentation of the targeted sequencing approach described herein.

**[0033]** FIG. 9 depicts schematics of illustrative universal oligonucleotide tail-adaptors (SEQ ID NOs: 45-48, respectively, in order of appearance).

#### DETAILED DESCRIPTION

**[0034]** Embodiments of the technology described herein relate to methods of determining (i.e sequencing) oligonucleotide sequences. In some embodiments, the methods

described herein relate to methods of enriching target sequences prior to a sequencing step. In some embodiments, the sequence of one end of the target sequence to be enriched is not known prior to the sequencing step.

**[0035]** For convenience, the meaning of some terms and phrases used in the specification, examples, and appended claims, are provided below. Unless stated otherwise, or implicit from context, the following terms and phrases include the meanings provided below. The definitions are provided to aid in describing particular embodiments, and are not intended to limit the claimed invention, because the scope of the invention is limited only by the claims. Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. If there is an apparent discrepancy between the usage of a term in the art and its definition provided herein, the definition provided within the specification shall prevail.

**[0036]** For convenience, certain terms employed herein, in the specification, examples and appended claims are collected here.

**[0037]** The terms “decrease”, “reduced”, “reduction”, or “inhibit” are all used herein generally to mean a decrease by a statistically significant amount. However, for avoidance of doubt, “reduced”, “reduction”, “decrease”, or “inhibit” means a decrease by at least 10% as compared to a reference level, for example a decrease by at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% decrease (e.g. absent level or non-detectable level as compared to a reference level), or any decrease between 10-100% as compared to a reference level. In the context of a marker or symptom is meant a statistically significant decrease in such level. The decrease can be, for example, at least 10%, at least 20%, at least 30%, at least 40% or more, and is preferably down to a level accepted as within the range of normal for an individual without such disorder.

**[0038]** The terms “increased”, “increase”, “enhance”, or “activate” are all used herein to generally mean an increase by a statistically significant amount; for the avoidance of doubt, the terms “increased”, “increase”, “enhance”, or “activate” mean an increase of at least 10% as compared to a reference level, for example an increase of at least about 20%, or at least about 30%, or at least about 40%, or at least about 50%, or at least about 60%, or at least about 70%, or at least about 80%, or at least about 90% or up to and including a 100% increase or any increase between 10-100% as compared to a reference level, or at least about a 2-fold, or at least about a 3-fold, or at least about a 4-fold, or at least about a 5-fold or at least about a 10-fold increase, or any increase between 2-fold and 10-fold or greater as compared to a reference level.

**[0039]** As used herein, a “subject” means a human or animal. Usually the animal is a vertebrate such as a primate, rodent, domestic animal or game animal. Primates include chimpanzees, cynomolgous monkeys, spider monkeys, and macaques, e.g., Rhesus. Rodents include mice, rats, woodchucks, ferrets, rabbits and hamsters. Domestic and game animals include cows, horses, pigs, deer, bison, buffalo, feline species, e.g., domestic cat, canine species, e.g., dog, fox, wolf, avian species, e.g., chicken, emu, ostrich, and fish, e.g., trout, catfish and salmon. In some embodiments, the

subject is a mammal, e.g., a primate, e.g., a human. The terms, “individual,” “patient” and “subject” are used interchangeably herein.

**[0040]** Preferably, the subject is a mammal. The mammal can be a human, non-human primate, mouse, rat, dog, cat, horse, or cow, but is not limited to these examples. Mammals other than humans can be advantageously used as subjects that represent animal models of, e.g. lung cancer. A subject can be male or female.

**[0041]** A subject can be one who has been previously diagnosed with or identified as suffering from or having a condition in need of treatment (e.g. cancer) or one or more complications related to such a condition, and optionally, have already undergone treatment for the condition or the one or more complications related to the condition. Alternatively, a subject can also be one who has not been previously diagnosed as having the condition (e.g. cancer) or one or more complications related to the condition. For example, a subject can be one who exhibits one or more risk factors for the condition or one or more complications related to the condition or a subject who does not exhibit risk factors.

**[0042]** A “subject in need” of treatment for a particular condition can be a subject having that condition, diagnosed as having that condition, or at risk of developing that condition.

**[0043]** As used herein, a “disease associated with a genetic alteration” refers to any disease which is caused by, at least in part, by an alteration in the genetic material of the subject as compared to a healthy wildtype subject, e.g. a deletion, an insertion, a SNP, a gene rearrangement. A disease can be caused by, at least in part, an alteration in the genetic material of the subject if the alteration increases the risk of the subject developing the disease, increases the subject’s susceptibility to a disease (including infectious diseases, or diseases with an infectious component), causes the production of a disease-associated molecule, or causes cells to become diseased or abnormal (e.g. loss of cell cycle regulation in cancer cells). Diseases can be associated with multiple genetic alterations, e.g. cancers.

**[0044]** As used herein, the term “nucleic acid” or “nucleic acid sequence” refers to any molecule, preferably a polymeric molecule, incorporating units of ribonucleic acid, deoxyribonucleic acid or an analog thereof. The nucleic acid can be either single-stranded or double-stranded. A single-stranded nucleic acid can be one strand nucleic acid of a denatured double-stranded DNA. Alternatively, it can be a single-stranded nucleic acid not derived from any double-stranded DNA. In one aspect, the template nucleic acid is DNA. In another aspect, the template is RNA. Suitable nucleic acid molecules are DNA, including genomic DNA or cDNA. Other suitable nucleic acid molecules are RNA, including mRNA.

**[0045]** The term “isolated” or “partially purified” as used herein refers, in the case of a nucleic acid, to a nucleic acid separated from at least one other component (e.g., nucleic acid or polypeptide) that is present with the nucleic acid as found in its natural source and/or that would be present with the nucleic acid when expressed by a cell. A chemically synthesized nucleic acid or one synthesized using in vitro transcription/translation is considered “isolated.”

**[0046]** The term “gene” means a nucleic acid sequence which is transcribed (DNA) to RNA in vitro or in vivo when operably linked to appropriate regulatory sequences. The

gene can include regulatory regions preceding and following the coding region, e.g. 5' untranslated (5'UTR) or "leader" sequences and 3' UTR or "trailer" sequences, as well as intervening sequences (introns) between individual coding segments (exons).

**[0047]** As used herein, the term "complementary" refers to the hierarchy of hydrogen-bonded base pair formation preferences between the nucleotide bases G, A, T, C and U, such that when two given polynucleotides or polynucleotide sequences anneal to each other, A pairs with T and G pairs with C in DNA, and G pairs with C and A pairs with U in RNA. As used herein, "substantially complementary" refers to a nucleic acid molecule or portion thereof (e.g. a primer) having at least 90% complementarity over the entire length of the molecule or portion thereof with a second nucleotide sequence, e.g. 90% complementary, 95% complementary, 98% complementary, 99% complementary, or 100% complementary. As used herein, "substantially identical" refers to a nucleic acid molecule or portion thereof having at least 90% identity over the entire length of a the molecule or portion thereof with a second nucleotide sequence, e.g. 90% identity, 95% identity, 98% identity, 99% identity, or 100% identity.

**[0048]** As used herein, "specific" when used in the context of a primer specific for a target nucleic acid refers to a level of complementarity between the primer and the target such that there exists an annealing temperature at which the primer will anneal to and mediate amplification of the target nucleic acid and will not anneal to or mediate amplification of non-target sequences present in a sample.

**[0049]** As used herein, "amplified product", "amplification product", or "amplicon" refers to oligonucleotides resulting from a PCR reaction that are copies of a portion of a particular target nucleic acid template strand and/or its complementary sequence, which correspond in nucleotide sequence to the template oligonucleotide sequence and/or its complementary sequence. An amplification product can further comprise sequence specific to the primers and which flanks sequence which is a portion of the target nucleic acid and/or its complement. An amplified product, as described herein will generally be double-stranded DNA, although reference can be made to individual strands thereof.

**[0050]** As used herein, a "portion" of a nucleic acid molecule refers to contiguous set of nucleotides comprised by that molecule. A portion can comprise all or only a subset of the nucleotides comprised by the molecule. A portion can be double-stranded or single-stranded.

**[0051]** As used herein, the terms "treat," "treatment," "treating," or "amelioration" refer to therapeutic treatments, wherein the object is to reverse, alleviate, ameliorate, inhibit, slow down or stop the progression or severity of a condition associated with a disease or disorder, e.g. lung cancer. The term "treating" includes reducing or alleviating at least one adverse effect or symptom of a condition, disease or disorder associated with a condition. Treatment is generally "effective" if one or more symptoms or clinical markers are reduced. Alternatively, treatment is "effective" if the progression of a disease is reduced or halted. That is, "treatment" includes not just the improvement of symptoms or markers, but also a cessation of, or at least slowing of, progress or worsening of symptoms compared to what would be expected in the absence of treatment. Beneficial or desired clinical results include, but are not limited to, alleviation of one or more symptom(s), diminishment of

extent of disease, stabilized (i.e., not worsening) state of disease, delay or slowing of disease progression, amelioration or palliation of the disease state, remission (whether partial or total), and/or decreased mortality, whether detectable or undetectable. The term "treatment" of a disease also includes providing relief from the symptoms or side-effects of the disease (including palliative treatment).

**[0052]** The term "statistically significant" or "significantly" refers to statistical significance and generally means a two standard deviation (2SD) below normal, or lower, concentration of the marker.

**[0053]** Other than in the operating examples, or where otherwise indicated, all numbers expressing quantities of ingredients or reaction conditions used herein should be understood as modified in all instances by the term "about." The term "about" when used in connection with percentages can mean  $\pm 1\%$ .

**[0054]** As used herein the term "comprising" or "comprises" is used in reference to compositions, methods, and respective component(s) thereof, that are essential to the method or composition, yet open to the inclusion of unspecified elements, whether essential or not.

**[0055]** The term "consisting of" refers to compositions, methods, and respective components thereof as described herein, which are exclusive of any element not recited in that description of the embodiment.

**[0056]** As used herein the term "consisting essentially of" refers to those elements required for a given embodiment. The term permits the presence of elements that do not materially affect the basic and novel or functional characteristic(s) of that embodiment.

**[0057]** The singular terms "a," "an," and "the" include plural referents unless context clearly indicates otherwise. Similarly, the word "or" is intended to include "and" unless the context clearly indicates otherwise. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of this disclosure, suitable methods and materials are described below. The abbreviation, "e.g." is derived from the Latin *exempli gratia*, and is used herein to indicate a non-limiting example. Thus, the abbreviation "e.g." is synonymous with the term "for example."

**[0058]** Definitions of common terms in cell biology and molecular biology can be found in "The Merck Manual of Diagnosis and Therapy", 19th Edition, published by Merck Research Laboratories, 2006 (ISBN 0-911910-19-0); Robert S. Porter et al. (eds.), The Encyclopedia of Molecular Biology, published by Blackwell Science Ltd., 1994 (ISBN 0-632-02182-9). Definitions of common terms in molecular biology can also be found in Benjamin Lewin, Genes X, published by Jones & Bartlett Publishing, 2009 (ISBN-10: 0763766321); Kendrew et al. (eds.), Molecular Biology and Biotechnology: a Comprehensive Desk Reference, published by VCH Publishers, Inc., 1995 (ISBN 1-56081-569-8) and Current Protocols in Protein Sciences 2009, Wiley Intersciences, Coligan et al., eds.

**[0059]** Unless otherwise stated, the present invention was performed using standard procedures, as described, for example in Sambrook et al., Molecular Cloning: A Laboratory Manual (3 ed.), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA (2001); and Davis et al., Basic Methods in Molecular Biology, Elsevier Science Publishing, Inc., New York, USA (1995) which are all incorporated by reference herein in their entireties.

**[0060]** Other terms are defined herein within the description of the various aspects of the invention.

**[0061]** Described herein are methods of determining the nucleotide sequence contiguous to a known target nucleotide sequence. Traditional sequencing methods generate sequence information randomly (e.g. “shotgun” sequencing) or between two known sequences which are used to design primers. In contrast, the methods described herein, in some embodiments, allow for determining the nucleotide sequence (e.g. sequencing) upstream or downstream of a single region of known sequence with a high level of specificity and sensitivity.

**[0062]** In some embodiments, the methods described herein relate to a method of enriching specific nucleotide sequences prior to determining the nucleotide sequence using a next-generation sequencing technology. In some embodiments, the methods of enriching specific nucleotide sequences do not comprise hybridization enrichment.

**[0063]** In some embodiments, the technology described herein can relate to a method of determining the nucleotide sequence contiguous to a known target nucleotide sequence, the method comprising; (a) ligating a target nucleic acid comprising the known target nucleotide sequence with a universal oligonucleotide tail-adaptor; (b) amplifying a portion of the target nucleic acid and the amplification strand of the universal oligonucleotide tail-adaptor with a first adaptor primer and a first target-specific primer; (c) amplifying a portion of the amplicon resulting from step (b) with a second adaptor primer and a second target-specific primer; and (d) sequencing the amplified portion from step (c) using a first and second sequencing primer. As used herein, the term “target nucleic acid” refers to a nucleic acid molecule comprising both the nucleic acid sequence which is to be determined and the known target nucleotide sequence. The target nucleic acid can be of any length and can be double-stranded or single-stranded. As used herein, the term “known target nucleotide sequence” refers to a portion of a target nucleic acid for which the sequence (e.g. the identity and order of the nucleotide bases comprises the nucleic acid) is known. The known target nucleotide sequence can be of any length of 10 or more nucleotides, preferably 30 or more nucleotides (e.g. 30 nucleotides, 40 nucleotides, 50 nucleotides or more). As used herein, the term “nucleotide sequence contiguous to” refers to a nucleotide sequence which is located on the same nucleic acid molecule (i.e. the target nucleic acid) as the known target nucleotide sequence and either upstream or downstream of the known target nucleotide sequence. The nucleotide sequence contiguous to can comprise any length of nucleotide sequence. In some embodiments, the nucleotide sequence contiguous to the known target nucleotide sequence comprises 1 kb or less of nucleotide sequence, e.g. 1 kb or less of nucleotide sequence, 750 bp or less of nucleotide sequence, 500 bp or less of nucleotide sequence, 400 bp or less of nucleotide sequence, 300 bp or less of nucleotide sequence, 200 bp or less of nucleotide sequence, 100 bp or less of nucleotide sequence. Where a sample comprises different target nucleic acids comprising the known target nucleotide sequence (e.g. a cell where the known target nucleotide sequence occurs multiple times in the genome, or on separate, non-identical chromosomes), there can be multiple sequences which comprise “nucleotide sequence contiguous to” the known target nucleotide sequence. As used herein, the term “determining

the nucleotide sequence”, refers to determining the identity and relative positions of the nucleotide bases comprising a nucleic acid.

**[0064]** In step (a) of the method described herein, the universal oligonucleotide tail-adaptor can be ligated to the target nucleic acid. In some embodiments, the target nucleic acid can be comprised by a sample comprising a plurality of nucleic acids, some of which do not comprise the target nucleic acid. In some embodiments, the universal oligonucleotide tail-adaptor can be ligated to substantially all of the nucleic acids in a sample. In some embodiments, the universal oligonucleotide tail-adaptor can be ligated to both nucleic acids which comprise the target nucleic acid sequence and to nucleic acids which do not comprise the target nucleic acid sequence.

**[0065]** As used herein, the term “universal oligonucleotide tail-adaptor” refers to a nucleic acid molecule comprised of two strands (a blocking strand and an amplification strand) and comprising a first ligatable duplex end and a second unpaired end. The blocking strand of the universal oligonucleotide tail-adaptor comprises a 5' duplex portion. The amplification strand comprises an unpaired 5' portion, a 3' duplex portion, and a 3' T overhang and nucleic acid sequences identical to a first and second sequencing primers. The duplex portions of the blocking strand and the amplification strand are substantially complementary and form the first ligatable duplex end comprising a 3' T overhang and the duplex portion is of sufficient length to remain in duplex form at the ligation temperature.

**[0066]** In some embodiments, the portion of the amplification strand that comprises a nucleic acid sequence identical to a first and second sequencing primers can be comprised, at least in part, by the 5' unpaired portion of the amplification strand.

**[0067]** In some embodiments, the universal oligonucleotide tail-adaptor can comprise a duplex portion and an unpaired portion, wherein the unpaired portion comprises only the 5' portion of the amplification strand, i.e. the entirety of the blocking strand is a duplex portion.

**[0068]** In some embodiments, the universal oligonucleotide tail-adaptor can have a “Y” shape, i.e. the unpaired portion can comprise portions of both the blocking strand and the amplification strand which are unpaired. The unpaired portion of the blocking strand can be shorter than, longer than, or equal in length to the unpaired portion of the amplification strand. In some embodiments, the unpaired portion of the blocking strand can be shorter than the unpaired portion of the amplification strand. Y shaped universal oligonucleotide tail-adaptors have the advantage that the unpaired portion of the blocking strand will not be subject to 3' extension during a PCR regimen.

**[0069]** In some embodiments, the blocking strand of the universal oligonucleotide tail-adaptor can further comprise a 3' unpaired portion which is not substantially complementary to the 5' unpaired portion of the amplification strand; and wherein the 3' unpaired portion of the blocking strand is not substantially complementary to or substantially identical to any of the primers. In some embodiments, the blocking strand of the universal oligonucleotide tail-adaptor can further comprise a 3' unpaired portion which will not specifically anneal to the 5' unpaired portion of the amplification strand at the annealing temperature; and wherein the 3'

unpaired portion of the blocking strand will not specifically anneal to any of the primers or the complements thereof at the annealing temperature.

**[0070]** In some embodiments, the duplex portion of the universal oligonucleotide tail-adaptor (e.g. the duplex portions of either or both of the strands) is at least 7 base pairs in length, e.g. 7 bp or more, 8 bp or more, 9 bp or more, 10 bp or more, 11 bp or more, 12 bp or more, 13 bp or more, or 14 bp or more in length. In some embodiments, the duplex portion of a universal oligonucleotide tail-adaptor can be at least 30 bp or longer, e.g. 30 bp or more, 31 bp or more, 32 bp or more, 33 bp or more, 34 bp or more, 35 bp or more, 40 bp or more, or 50 bp or more in length. The duplex portion of a universal oligonucleotide tail-adaptor should not be so long as to suppress PCR amplification of the desired amplicons in the PCR amplification regimen being used. Some next-generation sequencing methods use Y-shaped adaptor molecules. These Y-shaped adaptor molecules require duplex portions that are of limited length, (e.g. 17 bp or less) to avoid the formation of intramolecular hairpins during several PCR steps (e.g. library enrichment PCR, bridge PCR, or emulsion PCR). The Y-shaped universal oligonucleotide tail-adaptors of the methods described herein are not subject to this limitation of the duplex end as the two This PCR suppression effect resulting from the duplex end is not applicable in this invention since the two target primers can access the target fragment internally. In some embodiments, the duplex portion of a universal oligonucleotide tail-adaptor can be at least 18 bp or longer, e.g. 18 bp or more, 19 bp or more, 20 bp or more, 21 bp or more, 22 bp or more, 23 bp or more, 24 bp or more, or 25 bp or more in length.

**[0071]** Illustrative examples of universal oligonucleotide tail-adaptors are shown in FIG. 9.

**[0072]** Ligation of the universal oligonucleotide tail-adaptor can be accomplished by any method known in the art, e.g. blunt-end ligation or TA ligation. In some embodiments, prior to ligation of the Universal oligonucleotide tail-adaptor, the nucleic acids in a sample can be subjected to nucleic acid end-repair to blunt the ends of the nucleic acid. End-repair is well known in the art and relevant kits and/or enzymes are available commercially, (e.g. the NEBNEXT™ End Repair Module (Cat No. E6050L; New England Biolabs; Ipswich, MA).

**[0073]** In some embodiments, prior to ligation of the universal oligonucleotide tail-adaptor, the nucleic acids in a sample can be phosphorylated and/or adenylated. Adenylation can provide an adenosine overhang on the 3' end of a nucleic acid. A second nucleic acid with a thionine 3' overhang can then be ligated to the the first nucleic acid by TA ligation. Methods of ligation are well known in the art and relevant kits and/or enzymes are available commercially, e.g. the NEBNEXT™ da-Tailing module (Cat No. E6053L; New England Biolabs; Ipswich, MA) can be used to adenylate a blunt end of a nucleic acid. In some embodiments, Universal oligonucleotide tail-adaptors can be provided with a a thionine 3' overhang.

**[0074]** Steps (b) and (c) of the methods described herein can each comprise a PCR amplification regimen, i.e. a set of polymerase chain reaction (PCR) amplification cycles. As used herein, the term “amplification regimen” refers to a process of specifically amplifying, i.e., increasing the abundance of, a nucleic acid sequence of interest, and more particularly, the exponential amplification occurring when

the products of a previous polymerase extension serve as templates for the successive rounds of extension. A PCR amplification regimen according to the invention comprises at least one, and preferably at least 5 or more iterative cycles, where each cycle comprises the steps of: 1) strand separation (e.g., thermal denaturation); 2) oligonucleotide primer annealing to template molecules; and 3) nucleic acid polymerase extension of the annealed primers. Conditions and times necessary for each of these steps can be devised by one of ordinary skill in the art. An amplification regimen according to the methods described herein is preferably performed in a thermal cycler, many of which are commercially available.

**[0075]** PCR requires the use of a nucleic acid polymerase. As used herein, the phrase “nucleic acid polymerase” refers an enzyme that catalyzes the template-dependent polymerization of nucleoside triphosphates to form primer extension products that are complementary to the template nucleic acid sequence. A nucleic acid polymerase enzyme initiates synthesis at the 3' end of an annealed primer and proceeds in the direction toward the 5' end of the template. Numerous nucleic acid polymerases are known in the art and commercially available. One group of preferred nucleic acid polymerases are thermostable, i.e., they retain function after being subjected to temperatures sufficient to denature annealed strands of complementary nucleic acids, e.g. 94° C., or sometimes higher.

**[0076]** As understood in the art, PCR requires cycles including a strand separation step generally involving heating of the reaction mixture. As used herein, the term “strand separation” or “separating the strands” means treatment of a nucleic acid sample such that complementary double-stranded molecules are separated into two single strands available for annealing to an oligonucleotide primer. More specifically, strand separation according to the methods described herein is achieved by heating the nucleic acid sample above its  $T_m$ . Generally, for a sample containing nucleic acid molecules in buffer suitable for a nucleic acid polymerase, heating to 94° C. is sufficient to achieve strand separation. An exemplary buffer contains 50 mM KCl, 10 mM Tris-HCl (pH 8.8@25° C.), 0.5 to 3 mM MgCl<sub>2</sub>, and 0.1% BSA.

**[0077]** As also understood in the art, PCR requires annealing primers to template nucleic acids. Any strand of a target nucleic acid can be a template nucleic acid, as the template nucleic acid is defined as a single-strand nucleic acid to which a given primer will specifically anneal. As used herein, “anneal” refers to permitting two complementary or substantially complementary nucleic acids strands to hybridize, and more particularly, when used in the context of PCR, to hybridize such that a primer extension substrate for a template-dependent polymerase enzyme is formed. Conditions for primer-target nucleic acid annealing vary with the length and sequence of the primer and are based upon the calculated  $T_m$  for the primer. Generally, an annealing step in an amplification regimen involves reducing the temperature following the strand separation step to a temperature based on the calculated  $T_m$  for the primer sequence, for a time sufficient to permit such annealing.  $T_m$  can be readily predicted by one of skill in the art using any of a number of widely available algorithms (e.g., OLIGO™ (Molecular Biology Insights Inc. Colorado) primer design software and VENTRO NTI™ (Invitrogen, Inc. California) primer design software and programs available on the internet, including

Primer3, Oligo Calculator, and NetPrimer™ (Premier Bio-soft; Palo Alto, CA; and freely available on the world wide web). For example, the  $T_m$  of a primer can be calculated using following formula, which is used by NetPrimer™ software and is described in more detail in Frieir et al. PNAS 1986 83:9373-9377 which is incorporated by reference herein in its entirety.

$$T_m = \frac{\Delta H}{\Delta S + R \ln(C/4)} + 16.6 \log([K^+]/(1+0.7[K^+])) - 273.15$$

wherein,  $\Delta H$  is enthalpy for helix formation;  $\Delta S$  is entropy for helix formation;  $R$  is molar gas constant (1.987 cal/° C.\*mol);  $C$  is the nucleic acid concentration; and  $[K^+]$  is salt concentration. For most amplification regimens, the annealing temperature is selected to be about 5° C. below the predicted  $T_m$ , although temperatures closer to and above the  $T_m$  (e.g., between 1° C. and 5° C. below the predicted  $T_m$  or between 1° C. and 5° C. above the predicted  $T_m$ ) can be used, as can, for example, temperatures more than 5° C. below the predicted  $T_m$  (e.g., 6° C. below, 8° C. below, 10° C. below or lower). Generally, the closer the annealing temperature is to the  $T_m$ , the more specific is the annealing. The time allowed for primer annealing during a PCR amplification regimen depends largely upon the volume of the reaction, with larger volumes requiring longer times, but also depends upon primer and template concentrations, with higher relative concentrations of primer to template requiring less time than lower relative concentrations. Depending upon volume and relative primer/template concentration, primer annealing steps in an amplification regimen can be on the order of 1 second to 5 minutes, but will generally be between 10 seconds and 2 minutes, preferably on the order of 30 seconds to 2 minutes. As used herein, “substantially anneal” refers to a degree of annealing during a PCR amplification regimen which is sufficient to produce a detectable level of a specifically amplified product.

**[0078]** PCR also relies upon polymerase extension of annealed primers at each cycle. As used herein, the term “polymerase extension” means the template-dependent incorporation of at least one complementary nucleotide, by a nucleic acid polymerase, onto the 3' end of an annealed primer. Polymerase extension preferably adds more than one nucleotide, preferably up to and including nucleotides corresponding to the full length of the template. Conditions for polymerase extension vary with the identity of the polymerase. The temperature used for polymerase extension is generally based upon the known activity properties of the enzyme. Although, where annealing temperatures are required to be, for example, below the optimal temperatures for the enzyme, it will often be acceptable to use a lower extension temperature. In general, although the enzymes retain at least partial activity below their optimal extension temperatures, polymerase extension by the most commonly used thermostable polymerases (e.g., Taq polymerase and variants thereof) is performed at 65° C. to 75° C., preferably about 68-72° C.

**[0079]** Primer extension is performed under conditions that permit the extension of annealed oligonucleotide primers. As used herein, the term “conditions that permit the extension of an annealed oligonucleotide such that extension products are generated” refers to the set of conditions including, for example temperature, salt and co-factor concentrations, pH, and enzyme concentration under which a nucleic acid polymerase catalyzes primer extension. Such conditions will vary with the identity of the nucleic acid

polymerase being used, but the conditions for a large number of useful polymerase enzymes are well known to those skilled in the art. One exemplary set of conditions is 50 mM KCl, 10 mM Tris-HCl (pH 8.8@25° C.), 0.5 to 3 mM MgCl<sub>2</sub>, 200 uM each dNTP, and 0.1% BSA at 72° C., under which Taq polymerase catalyzes primer extension. The conditions for initiation and extension usually include the presence of at least one, but more preferably all four different deoxyribonucleoside triphosphates and a polymerization-inducing agent such as DNA polymerase or reverse transcriptase, in a suitable buffer (in this context “buffer” includes solvents (generally aqueous) plus necessary cofactors and reagents which affect pH, ionic strength, etc.) and at a suitable temperature.

**[0080]** In some embodiments, each amplification step can comprise a set of cycles of a PCR amplification regimen from 5 cycles to 20 cycles in length. In some embodiments, each amplification step can comprise a set of cycles of a PCR amplification regimen from 10 cycles to 20 cycles in length. In some embodiments, each amplification step can comprise a set of cycles of a PCR amplification regimen from 12 cycles to 16 cycles in length. In some embodiments, the annealing temperature can be less than 70° C. In some embodiments, the annealing temperature can be less than 72° C.

**[0081]** In various embodiments, the methods and compositions described herein relate to performing a PCR amplification regimen with one or more of the types of primers described herein. As used herein, “primer” refers to a DNA or RNA polynucleotide molecule or an analog thereof capable of specifically annealing to a polynucleotide template and providing a 3' end that serves as a substrate for a template-dependent polymerase to produce an extension product which is complementary to the polynucleotide template. A primer useful in the methods described herein is generally single-stranded, and a primer and its complement can anneal to form a double-stranded polynucleotide. Primers according to the methods and compositions described herein can be less than or equal to 300 nucleotides in length, e.g., less than or equal to 300, or 250, or 200, or 150, or 100, or 90, or 80, or 70, or 60, or 50, or 40, or 30 or fewer, or 20 or fewer, or 15 or fewer, but at least 10 nucleotides in length. Methods of making primers are well known in the art, and numerous commercial sources offer oligonucleotide synthesis services suitable for providing primers according to the methods and compositions described herein, e.g. INVITRO-GEN™ Custom DNA Oligos; Life Technologies; Grand Island, NY or custom DNA Oligos from IDT; Coralville, IA).

**[0082]** In some embodiments, after the Universal oligonucleotide tail-adaptor is ligated to the nucleic acids in a sample (e.g. the target nucleic acids), the target nucleic acid can be amplified in a first amplification step (i.e. step (b)). The first amplification step can be a set of PCR amplification cycles using a first target-specific primer and a first tail-adaptor primer.

**[0083]** As used herein, the term “first target-specific primer” refers to a single-stranded oligonucleotide comprising a nucleic acid sequence that can specifically anneal to the target nucleic acid at the annealing temperature.

**[0084]** In some embodiments, the first-target specific primer can comprise a 5' tag sequence portion. In some embodiments, all first-target specific primers present in a reaction can comprise identical 5' tag sequence portions. In

a multiplex PCR reaction, different primer species can interact with each other in an undesired off-target manner, leading to primer extension and subsequently amplification by DNA polymerase. These primer dimers tend to be short, and their efficient amplification can overtake the reaction and dominate resulting in poor amplification of desired target sequence. The inclusion of a 5' tag sequence on the first-target specific primer(s) causes any potential primer dimers that can result to contain the same complementary tails on both ends. In subsequent PCR cycles, the primer dimers would denature into single-stranded DNA primer dimers, each comprising complementary sequences on their two ends which are introduced by the 5' tag. Instead of primer annealing to these single stranded DNA primer dimers, an intra-molecular hairpin (a panhandle like structure) formation would preferentially occur due to the proximate accessibility of the complementary tags on the same primer dimer molecule instead of an inter-molecular interaction with new primers on separate molecules. As a result, these primer dimers are very inefficiently amplified, such that primers are not exponentially consumed by the undesired dimers for amplification. Instead the tagged primers can remain in high and sufficient concentration for desired specific amplification of target sequences. Accumulation of primer dimers can also be a detriment to multiplex PCR because they compete for and consume other reagents in the reaction. In some embodiments, the 5' tag sequence can be a GC-rich sequence, i.e. the tag sequence can comprise at least 50% GC content, at least 55% GC content, at least 60% GC content, at least 65% GC content, at least 70% GC content, at least 75% GC content, at least 80% GC content, or higher GC content. In some embodiments, the tag sequence can comprise at least 60% GC content. In some embodiments, the tag sequence can comprise at least 65% GC content.

**[0085]** As used herein, the term “second target-specific primer” refers to a single-stranded oligonucleotide comprising a 3' portion comprising a nucleic acid sequence that can specifically anneal to a portion of the known target nucleotide sequence comprised by the amplicon resulting from step (b), and a 5' portion comprising a nucleic acid sequence that is identical to a second sequencing primer. The second target-specific primer is nested with respect to the first target-specific primer. In some embodiments, the second target-specific primer is nested with respect to the first target-specific primer by at least 3 nucleotides, e.g. by 3 or more, 4 or more, 5 or more, 6 or more, 7 or more, 8 or more, 9 or more, 10 or more, or 15 or more nucleotides.

**[0086]** In some embodiments, all of the second target-specific primers present in a reaction comprise the same 5' portion. In some embodiments, the 5' portion of the can serve to suppress primer dimers as described for the 5' tag of the first target-specific primer described above herein.

**[0087]** In some embodiments, the first and second target-specific primers are substantially complementary to the same strand of the target nucleic acid. In some embodiments, the the portions of the first and second target-specific primers that specifically anneal to the known target sequence can comprise a total of at least 20 unique bases of the known target nucleotide sequence, e.g. 20 or more unique bases, 25 or more unique bases, 30 or more unique bases, 35 or more unique bases, 40 or more unique bases, or 50 or more unique bases. In some embodiments, the the portions of the first and

second target-specific primers that specifically anneal to the known target sequence can comprise a total of at least 30 unique bases of the known target nucleotide sequence,

**[0088]** As used herein, the term “first adaptor primer” refers to a nucleic acid molecule comprising a nucleic acid sequence identical to a 5' portion of the first sequencing primer. As the first tail-adaptor primer is therefor identical to at least a portion of the sequence of the amplification strand (as opposed to complementary), it will not be able to specifically anneal to any portion of the universal oligonucleotide tail-adaptor itself.

**[0089]** As used herein, the term “second adaptor primer” refers to a nucleic acid molecule comprising a nucleic acid sequence identical to a portion of the first sequencing primer and is nested with respect to the first adaptor primer. As the second tail-adaptor primer is therefor identical to at least a portion of the sequence of the amplification strand (as opposed to complementary), it will not be able to specifically anneal to any portion of the universal oligonucleotide tail-adaptor itself. In some embodiments, the second adaptor primer is identical to the first sequencing primer.

**[0090]** The second adaptor primer should be nested with respect to the first adaptor primer, that is, the first adaptor primer comprises a nucleic acid sequence identical to the amplification strand which is not comprised by the second adaptor primer and which is located closer to the 5' end of the amplification primer than any of the sequence identical to the amplification strand which is comprised by the second adaptor primer. In some embodiments, the second adaptor primer is nested by at least 3 nucleotides, e.g. by 3 nucleotides, by 4 nucleotides, by 5 nucleotides, by 6 nucleotides, by 7 nucleotides, by 8 nucleotides, by 9 nucleotides, by 10 nucleotides or more.

**[0091]** In some embodiments, the first adaptor primer can comprise a nucleic acid sequence identical to about the 20 5'-most bases of the amplification strand of the universal oligonucleotide tail-adaptor and the second adaptor primer can comprise a nucleic acid sequence identical to about 30 bases of the amplification strand of the universal oligonucleotide tail-adaptor, with a 5' base which is at least 3 nucleotides 3' of the 5' terminus of the amplification strand.

**[0092]** The use of nested adaptor primers eliminates the possibility of producing final amplicons that are amplifiable (e.g. during bridge PCR or emulsion PCR) but cannot be sequenced, a situation that can arise during hemi-nested methods. In other situations, hemi-nested approaches using a primer identical to a sequencing primer can result in the carry-over of undesired amplification products from the first PCR step to the second PCR step and would ultimately yield artificial sequencing reads. The use of two adaptor primers, as described herein can reduce, and in some embodiments eliminate, these problems.

**[0093]** In the first PCR amplification cycle of the first amplification step, the first target-specific primer can specifically anneal to a template strand of any nucleic acid comprising the known target nucleotide sequence. Depending upon the orientation with which the first target-specific primer was designed, sequence upstream or downstream of the known target nucleotide sequence, and complementary to the template strand will be synthesized. If, during the extension phase of PCR, the 5' end of the template strand terminates in a ligated Universal oligonucleotide tail-adaptor, the 3' end of the newly synthesized product strand will comprise sequence complementary to the first tail-adaptor

primer. In subsequent PCR amplification cycles, both the first target-specific primer and the first tail-adaptor primer will be able to specifically anneal to the appropriate strands of the target nucleic acid sequence and the sequence between the known nucleotide target sequence and the Universal oligonucleotide tail-adaptor can be amplified, (i.e. copied).

**[0094]** In the next step (i.e. step (c)) of the method described herein, a portion of the amplified portion resulting from step (b) is amplified in a second amplification step. The second amplification step can be a set of PCR amplification cycles using a second target-specific primer and a first sequencing primer. The second set of PCR amplification cycles can have PCR parameters identical to, or which differ from, those of the first set of PCR amplification cycles. E.g. the PCR amplification regimens of steps (b) and (c) can have the same or different annealing temperatures or the same or different extension step time lengths.

**[0095]** The methods described herein allow for determining the nucleotide sequence contiguous to a known target nucleotide sequence on either or both flanks of the known target nucleotide sequence. Regardless of whether the target nucleic acid normally exists as a single-stranded or double-stranded nucleic acid, sequence information is typically represented in a single-stranded format (Strand A), from 5' to 3'. If the sequence 5' of the known target nucleotide sequence of Strand A is to be determined, the gene-specific primers can be complementary (i.e. anneal to) Strand A. If the sequence 3' of the known target nucleotide sequence of Strand A is to be determined, the gene-specific primers can be identical to Strand A, such that they will anneal to the complementary strand of a double-stranded target nucleic acid. Such considerations of primer design are well known to those of ordinary skill in the art.

**[0096]** In some embodiments, the methods described herein, relating to the use of a first and second gene-specific primer can result in assays with a superior on-target rate, e.g. 70-90%. In some embodiments, the assays and methods described herein can have a target specificity rate of at least 85%.

**[0097]** In some embodiments, the four types of primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of from about 61 to 72° C., e.g. from about 61 to 69° C., from about 63 to 69° C., from about 63 to 67° C., from about 64 to 66° C. In some embodiments, the four types of primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of less than 72° C. In some embodiments, the four types of primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of less than 70° C. In some embodiments, the four types of primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of less than 68° C. In some embodiments, the four types of primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of about 65° C.

**[0098]** In some embodiments, the portions of the target-specific primers that specifically anneal to the known target nucleotide sequence will anneal specifically at a temperature of about 61 to 72° C., e.g. from about 61 to 69° C., from about 63 to 69° C., from about 63 to 67° C., from about 64 to 66° C. In some embodiments, the portions of the target-specific primers that specifically anneal to the known target

nucleotide sequence will anneal specifically at a temperature of about 65° C. in a PCR buffer.

**[0099]** In some embodiments, the primers and/or adaptors described herein can not comprise modified bases (e.g. the primers and/or adaptors can not comprise a blocking 3' amine).

**[0100]** In the next step (i.e. step (d)) of the methods described herein, the amplified portion resulting from step (c) can be sequenced. In some embodiments, the sequencing can be performed by a next-generation sequencing method. As used herein “next-generation sequencing” refers to oligonucleotide sequencing technologies that have the capacity to sequence oligonucleotides at speeds above those possible with conventional sequencing methods (e.g. Sanger sequencing), due to performing and reading out thousands to millions of sequencing reactions in parallel. Non-limiting examples of next-generation sequencing methods/platforms include Massively Parallel Signature Sequencing™ (Lynx Therapeutics); 454 Pyro-Sequencing™ (454 Life Sciences/Roche Diagnostics); solid-phase, reversible dye-terminator sequencing (Solexa/Illumina); SOLiD™ technology (Applied Biosystems); Ion semiconductor sequencing (ION Torrent™); DNA nanoball sequencing (Complete Genomics); and technologies available from Pacific Biosciences, Intelligen Bio-systems, Oxford Nanopore Technologies, and Helicos Biosciences. In some embodiments, the sequencing primers can comprise portions compatible with the selected next-generation sequencing method. Next-generation sequencing technologies and the constraints and design parameters of associated sequencing primers are well known in the art (see, e.g. Shendure, et al., “Next-generation DNA sequencing,” *Nature*, 2008, vol. 26, No. 10, 1135-1145; Mardis, “The impact of next-generation sequencing technology on genetics,” *Trends in Genetics*, 2007, vol. 24, No. 3, pp. 133-141; Su, et al., “Next-generation sequencing and its applications in molecular diagnostics” *Expert Rev Mol Diagn*, 2011, 11(3):333-43; Zhang et al., “The impact of next-generation sequencing on genomics”, *J Genet Genomics*, 2011, 38(3):95-109; (Nyren, P. et al. *Anal Biochem* 208: 17175 (1993); Bentley, D. R. *Curr Opin Genet Dev* 16:545-52 (2006); Strausberg, R. L., et al. *Drug Disc Today* 13:569-77 (2008); U.S. Pat. Nos. 7,282,337; 7,279,563; 7,226,720; 7,220,549; 7,169,560; 6,818,395; 6,911,345; US Pub. Nos. 2006/0252077; 2007/0070349; and 20070070349; which are incorporated by reference herein in their entireties).

**[0101]** In some embodiments, the sequencing step relies upon the use of a first and second sequencing primers. In some embodiments, the first and second sequencing primers are selected to be compatible with a next-generation sequencing method as described herein.

**[0102]** Methods of aligning sequencing reads to known sequence databases of genomic and/or cDNA sequences are well known in the art and software is commercially available for this process. In some embodiments, reads (less the sequencing primer and/or adaptor nucleotide sequence) which do not map, in their entirety, to wild-type sequence databases can be genomic rearrangements or large indel mutations. In some embodiments, reads (less the sequencing primer and/or adaptor nucleotide sequence) comprising sequences which map to multiple locations in the genome can be genomic rearrangements.

**[0103]** In some embodiments, target nucleic acids and/or amplification products thereof can be isolated from enzymes, primers, or buffer components before and/or after

any of steps a-d. Methods for isolating nucleic acids are well known in the art. In some embodiments, the isolation can comprise Solid Phase Reversible Immobilization (SPRI) cleanup. Methods for SPRI cleanup are well known in the art and kits are commercially available, e.g. Agencourt AMPure XP™-PCR Purification (Cat No. A63880, Beckman Coulter; Brea, CA). In some embodiments, enzymes can be inactivated by heat treatment.

**[0104]** In some embodiments, the target nucleic acid can be comprised by genomic DNA. In some embodiments, the target nucleic acid can be comprised by ribonucleic acid (RNA), e.g. mRNA. In some embodiments, the target nucleic acid can be comprised by cDNA. Many of the sequencing methods suitable for use in the methods described herein provide sequencing runs with optimal read lengths of tens to hundreds of nucleotide bases (e.g. Ion Torrent™ technology can produce read lengths of 200-400 bp). Target nucleic acids comprised, for example, by genomic DNA or mRNA, can be comprised by nucleic acid molecules which are substantially longer than this optimal read length. In order for the amplified nucleic acid portion resulting from step (c) to be of a suitable length for use in a particular sequencing technology, the average distance between the known target nucleotide sequence and an end of the target nucleic acid to which the Universal oligonucleotide tail-adaptor can be ligated should be as close to the optimal read length of the selected technology as possible. For example, if the optimal read-length of a given sequencing technology is 200 bp, then the nucleic acid molecules amplified in accordance with the methods described herein should have an average length of about 400 bp or less. Target nucleic acids comprised by, e.g., genomic DNA or mRNA, can be sheared, e.g. mechanically or enzymatically sheared, to generate fragments of any desired size prior to step (a). Non-limiting examples of mechanical shearing processes include sonication, nebulization, and AFA™ shearing technology available from Covaris (Woburn, MA). In some embodiments, a target nucleic acid comprised by genomic DNA can be mechanically sheared by sonication. In some embodiments, when the target nucleic acid is comprised by RNA, the sample can be subjected to a reverse transcriptase regimen to generate DNA template and the DNA template can then be sheared. In some embodiments, target RNA can be sheared before performing the reverse transcriptase regimen. In some embodiments, a sample comprising target RNA can be used in the methods described herein using total nucleic acids extracted from either fresh or degraded specimens; without the need of genomic DNA removal for cDNA sequencing; without the need of ribosomal RNA depletion for cDNA sequencing; without the need of mechanical or enzymatic shearing in any of the steps; by subjecting the RNA for double-stranded cDNA synthesis using random hexamers; and by subjecting the nucleic acid to end-repair, phosphorylation, and adenylation in a single tube.

**[0105]** In some embodiments, the known target nucleotide sequence can be comprised by a gene rearrangement. The methods described herein are suited for determining the presence and/or identity of a gene rearrangement as the identity of only one half of the gene rearrangement must be previously known (i.e. the half of the gene rearrangement which is to be targeted by the gene-specific primers). In some embodiments, the gene rearrangement can comprise an oncogene. In some embodiments, the gene rearrangement can comprise a fusion oncogene.

**[0106]** In some embodiments, the target nucleic acid can be comprised by a sample. In some embodiments, the target nucleic acid can be comprised by a sample obtained from a subject. In some embodiments a sample can be a diagnostic sample obtained from a subject. In some embodiments, a sample can further comprise proteins, cells, fluids, biological fluids, preservatives, and/or other substances. By way of non-limiting example, a sample can be a cheek swab, blood, serum, plasma, sputum, cerebrospinal fluid, urine, tears, alveolar isolates, pleural fluid, pericardial fluid, cyst fluid, tumor tissue, tissue, a biopsy, saliva, an aspirate, or combinations thereof. In some embodiments, a sample can be obtained by resection or biopsy.

**[0107]** In some embodiments, the sample can be obtained from a subject in need of treatment for cancer. In some embodiments, the sample can comprise a population of tumor cells, e.g. at least one tumor cell. In some embodiments, the sample can comprise a tumor biopsy, including but not limited to, untreated biopsy tissue or treated biopsy tissue (e.g. formalin-fixed and/or paraffin-embedded biopsy tissue).

**[0108]** In some embodiments, the sample can be freshly collected. In some embodiments, the sample can be stored prior to being used in the methods and compositions described herein. In some embodiments, the sample is an untreated sample. As used herein, “untreated sample” refers to a biological sample that has not had any prior sample pre-treatment except for dilution and/or suspension in a solution. In some embodiments, a sample can be obtained from a subject and preserved or processed prior to being utilized in the methods and compositions described herein. By way of non-limiting example, a sample can be embedded in paraffin wax, refrigerated, or frozen. A frozen sample can be thawed before determining the presence of a nucleic acid according to the methods and compositions described herein. In some embodiments, the sample can be a processed or treated sample. Exemplary methods for treating or processing a sample include, but are not limited to, centrifugation, filtration, sonication, homogenization, heating, freezing and thawing, contacting with a preservative (e.g. anti-coagulant or nuclease inhibitor) and any combination thereof. In some embodiments, the sample can be treated with a chemical and/or biological reagent. Chemical and/or biological reagents can be employed to protect and/or maintain the stability of the sample or nucleic acid comprised by the sample during processing and/or storage. In addition, or alternatively, chemical and/or biological reagents can be employed to release nucleic acids from other components of the sample. By way of non-limiting example, a blood sample can be treated with an anti-coagulant prior to being utilized in the methods and compositions described herein. The skilled artisan is well aware of methods and processes for processing, preservation, or treatment of samples for nucleic acid analysis. In some embodiments, the sample can be a clarified fluid sample, for example, by centrifugation. In some embodiments, the sample can be clarified by low-speed centrifugation (e.g. 3,000×g or less) and collection of the supernatant comprising the clarified fluid sample.

**[0109]** In some embodiments, the nucleic acid present in a sample can be isolated, enriched, or purified prior to being utilized in the methods and compositions described herein. Methods of isolating, enriching, or purifying nucleic acids from a sample are well known to one of ordinary skill in the art. By way of non-limiting example, kits for isolation of

genomic DNA from various sample types are commercially available (e.g. Catalog Nos. 51104, 51304, 56504, and 56404; Qiagen; Germantown, MD).

**[0110]** The methods described herein can be used in multiplex techniques. In embodiments of the methods described herein, multiplex applications can include determining the nucleotide sequence contiguous to one or more known target nucleotide sequences. As used herein, “multiplex PCR” refers to a variant of PCR where simultaneous amplification of more than one target nucleic acid in one reaction vessel and subsequent determination of the sequence of the amplification products by using more than one set of first and second gene-specific primers. Multiplex can refer to the detection of between about 2-1,000 different target sequences in a single reaction. As used herein, multiplex refers to the detection of any range between 2-1,000, e.g., between 5-500, 25-1000, or 10-100 different target sequences in a single reaction, etc. The term “multiplex” as applied to PCR implies that there are primers specific for at least two different target sequences in the same PCR reaction.

**[0111]** In some embodiments, the target nucleic acids in a sample, or separate portions of a sample, can be amplified with a plurality of first and second target-specific primers. In some embodiments, the plurality of first and second target-specific primers can be present in a single reaction mixture, e.g. multiple amplification products can be produced in the same reaction mixture. In some embodiments, the plurality of sets of first and second target-specific primers can specifically anneal to known target sequences comprised by separate genes. In some embodiments, at least two sets of first and second target-specific primers can specifically anneal to different portions of a known target sequence. In some embodiments, at least two sets of first and second target-specific primers can specifically anneal to different portions of a known target sequence comprised by a single gene. In some embodiments, at least two sets of first and second target-specific primers can specifically anneal to different exons of a gene comprising a known target sequence. In some embodiments, the plurality of first target-specific primers can comprise identical 5' tag sequence portions.

**[0112]** In embodiments of the methods described herein, multiplex applications can include determining the nucleotide sequence contiguous to one or more known target nucleotide sequences in multiple samples in one sequencing reaction or sequencing run. The multiple samples can be of different origins, e.g. from different tissues and/or different subjects. In such embodiments, the universal oligonucleotide tail-adaptor can further comprise a barcode portion. In some embodiments, a universal oligonucleotide tail-adaptor with a unique barcode portion can be added to each sample and ligated to the nucleic acids therein; the samples can then be pooled after step (a). Each resulting sequencing read of an amplification product will therefor comprise a barcode identifying which sample comprised the original template nucleic acid from which the amplification product is derived. The use of barcode portions in next-generation sequencing applications is well known in the art and described, for example, in Margulies, M. et al. “Genome Sequencing in Microfabricated High-Density Picolitre Reactors”, *Nature*, 437, 376-80 (2005); Mikkelsen, T. et al. “Genome-Wide Maps of Chromatin State in Pluripotent and Lineage-Committed Cells”, *Nature*, 448, 553-60 (2007);

McLaughlin, S. et al. “Whole-Genome Resequencing With Short Reads: Accurate Mutation Discovery With Mate Pairs and Quality Values”, ASHG Annual Meeting (2007); Shendure I. et al. “Accurate Multiplex Polony Sequencing of an Evolved Bacterial Genome”, *Science*, 309, 1728-32 (2005); Harris, T. et al. “Single-Molecule DNA Sequencing of a Viral Genome”, *Science*, 320, 106-9 (2008); Simen, B. et al. “Prevalence of LoW Abundance Drug Resistant Variants by Ultra Deep Sequencing in Chronically HIV-infected Antiretroviral (ARV) Naive Patients and the Impact on Virologic Outcomes”, 16th International HIV Drug Resistance Workshop, Barbados (2007); Thomas, R. et al. “Sensitive Mutation Detection in Heterogeneous Cancer Specimens by Massively Parallel Picoliter Reactor Sequencing”, *Nature Med.*, 12, 852-855 (2006); Mitsuya, Y et al. “Minority Human Immunodeficiency Virus Type 1 Variants in Antiretroviral-Naive Persons With Reverse Transcriptase Codon 215 Revertant Mutations”, *I. Vir.*, 82, 10747-10755 (2008); Binladen, J. et al. “The Use of Coded PCR Primers Enables High-Throughput Sequencing of Multiple Homolog Amplification Products by 454 Parallel Sequencing”, *PLoS ONE*, 2, e197 (2007); and Hoffmann, C. et al. “DNA Bar Coding and PyROSequencing to Identify Rare HIV Drug Resistance Mutations”, *Nuc. Acids Res.*, 35, e91 (2007), all of which are herein incorporated by reference.

**[0113]** In some embodiments of the technology described herein, determining the sequence contiguous to a known oligonucleotide target sequence can provide information relevant to treatment of disease, and/or can be comprised by a method of treating disease. In some embodiments, the sample can be from a subject in need of treatment for a disease associated with a genetic alteration. In some embodiments, the known oligonucleotide target sequence can be comprised by a disease-associated gene, e.g. an oncogene. In some embodiments, the sequence contiguous to a known oligonucleotide target sequence and/or the known oligonucleotide target sequence can comprise a mutation or genetic abnormality which is disease-associated, e.g. a SNP, an insertion, a deletion, and/or a gene rearrangement. In some embodiments, the sequence contiguous to a known oligonucleotide target sequence and/or the known oligonucleotide target sequence present in a sample can be comprised by a gene rearrangement product. In some embodiments, the gene rearrangement can be an oncogene, e.g. a fusion oncogene.

**[0114]** Certain treatments for cancer are particularly effective against tumors comprising certain oncogenes, e.g. a treatment agent which targets the action or expression of a given fusion oncogene can be effective against tumors comprising that fusion oncogene but not against tumors lacking the fusion oncogene. The methods described herein can allow the determination of specific sequences which reveal oncogene status (e.g. mutations, SNPs, and/or rearrangements). As described herein, the methods described herein can further allow the determination of specific sequences when the sequence of only one flank is known, e.g. the methods described herein can determine the presence and identity of gene rearrangements involving known oncogenes where the precise location and/or rearrangement partner are not known before the methods described herein are performed.

**[0115]** In some embodiments, the technology described herein relates to a method of treating cancer, the method comprising; detecting, in a tumor sample obtained from a

subject in need of treatment for cancer, the presence of one or more oncogene rearrangements according to the method described herein; administering a cancer treatment which is effective against tumors having any of the detected oncogene rearrangements. In some embodiments, the technology described herein relates to a method of determining if a subject in need of treatment for cancer will be responsive to a given treatment, the method comprising; detecting, in a tumor sample obtained from the subject, the presence of an oncogene rearrangement according to the method as described herein; wherein the subject is determined to be responsive to a treatment targeting an oncogene rearrangement product if the presence of the oncogene rearrangement is detected.

**[0116]** In some embodiments, e.g. when the sample is obtained from a subject in need of treatment for lung cancer, the known oligonucleotide target sequence can comprise sequence from a gene selected from the group of ALK; ROS1; and RET. Gene rearrangements involving the ALK, ROS1, and RET genes and which result in fusion oncogenes are well known in the art (see, e.g. Soda et al. *Nature* 2007 448:561-6; Rikova et al. *Cell* 2007 131:1190-1203; Kohno et al. *Nature Medicine* 2012 18:375-7; Takouchi et al. *Nature Medicine* 2012 18:378-81; which are incorporated by reference herein in their entireties. However, the precise location of the gene rearrangement (e.g. where in the ALK, ROS1, and/or RET gene the rearrangement has occurred), and the identity of the second gene involved in the rearrangement can vary. In the methods described herein, the presence and identity of such a rearrangement can be detected without having to know the location of the rearrangement or the identity of the second gene involved in the gene rearrangement.

**[0117]** In some embodiments, the known target sequence can comprise sequence from a gene selected from the group of: ALK; ROS1; and RET. In some embodiments, at least one set of a first target-specific primer and a second target-specific primer can be selected from the group consisting of; SEQ ID NOs: 5 and 6; SEQ ID NOs: 7 and 8; SEQ ID NOs: 9 and 10; SEQ ID NOs: 11 and 12; SEQ ID NOs: 13 and 14; SEQ ID NOs: 15 and 16; SEQ ID NOs: 17 and 18; SEQ ID NOs: 19 and 20; SEQ ID NOs: 21 and 22; SEQ ID NOs: 23 and 24; SEQ ID NOs: 25 and 26; SEQ ID NOs: 27 and 28; SEQ ID NOs: 29 and 30; SEQ ID NOs: 31 and 32; SEQ ID NOs: 33 and 34; SEQ ID NOs: 35 and 36; and SEQ ID NOs: 37 and 38.

**[0118]** In some embodiments, the presence of a gene rearrangement of ALK in a sample obtained from a tumor in a subject can indicate that the tumor is susceptible to treatment with a treatment selected from the group consisting of: an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; diamino and aminopyrimidine inhibitors of ALK kinase activity such as NVP-TAE684 and PF-02341066 (see, e.g. Galkin et al. *Proc Natl Acad Sci USA*, 2007, 104:270-275; Zou et al. *Cancer Res*, 2007, 67:4408-4417; Hallberg and Palmer *F1000 Med Reports* 2011 3:21; and Sakamoto et al. *Cancer Cell* 2011 19:679-690) and molecules disclosed in WO 04/079326. All of the foregoing references are incorporated by reference herein in their entireties. An ALK inhibitor can include any agent that reduces the expression and/or kinase activity of ALK or a portion thereof, including, e.g. oligonucleotides, small molecules, and/or peptides that reduce the

expression and/or activity of ALK or a portion thereof. As used herein “anaplastic lymphoma kinase” or “ALK” refers to a transmembrane tyrosine kinase typically involved in neuronal regulation in the wildtype form. The nucleotide sequence of the ALK gene and mRNA are known for a number of species, including human (e.g. SEQ ID NO: 2 (mRNA), NCBI Gene ID: 238).

**[0119]** In some embodiments, the presence of a gene rearrangement of ROS1 in a sample obtained from a tumor in a subject can indicate that the tumor is susceptible to treatment with a treatment selected from the group consisting of: a ROS1 inhibitor and an ALK inhibitor as described herein above (e.g. crizotinib). A ROS1 inhibitor can include any agent that reduces the expression and/or kinase activity of ROS1 or a portion thereof, including, e.g. oligonucleotides, small molecules, and/or peptides that reduce the expression and/or activity of ROS1 or a portion thereof. As used herein “c-ROS1 oncogene 1” or “ROS1” (also referred to in the art as ROS1-1) refers to a transmembrane tyrosine kinase of the sevenless subfamily and which interacts with PTPN6. The nucleotide sequence of the ROS1 gene and mRNA are known for a number of species, including human (e.g. SEQ ID NO: 1 (mRNA), NCBI Gene ID: 238).

**[0120]** In some embodiments, the presence of a gene rearrangement of RET in a sample obtained from a tumor in a subject can indicate that the tumor is susceptible to treatment with a treatment selected from the group consisting of: a RET inhibitor; DP-2490, DP-3636, SU5416; BAY 43-9006, BAY 73-4506 (regorafenib), ZD6474, NVP-AST487, sorafenib, RPM, XL184, vandetanib, sunitinib, imatinib, pazopanib, axitinib, motesanib, gefitinib, and withaferin A (see, e.g. Samadi et al. *Surgery* 2010 148:1228-36; Cuccuru et al. *JNCI* 2004 13:1006-1014; Akeno-Stuart et al. *Cancer Research* 2007 67:6956; Grazma et al. *J Clin Oncol* 2010 28:15s 5559; Mologni et al. *J Mol Endocrinol* 2006 37:199-212; Calmomagno et al. *Journal NCI* 2006 98:326-334; Mologni. *Curr Med Chem* 2011 18:162-175 and the compounds disclosed in WO 06/034833; US Patent Publication 2011/0201598 and U.S. Pat. No. 8,067,434). All of the foregoing references are incorporated by reference herein in their entireties. A RET inhibitor can include any agent that reduces the expression and/or kinase activity of RET or a portion thereof, including, e.g. oligonucleotides, small molecules, and/or peptides that reduce the expression and/or activity of RET or a portion thereof. As used herein “rearranged during transfection” or “RET” refers to a receptor tyrosine kinase of the cadherin superfamily which is involved in neural crest development and recognizes glial cell line-derived neurotrophic factor family signaling molecules. The nucleotide sequence of the ROS1 gene and mRNA are known for a number of species, including human (e.g. SEQ ID NOs: 3-4 (mRNA), NCBI Gene ID: 5979).

**[0121]** Further non-limiting examples of applications of the methods described herein include detection of hematological malignancy markers and panels thereof (e.g. including those to detect genomic rearrangements in lymphomas and leukemias), detection of sarcoma-related genomic rearrangements and panels thereof; and detection of IGH/TCR gene rearrangements and panels thereof for lymphoma testing.

**[0122]** In some embodiments, the methods described herein relate to treating a subject having or diagnosed as having, e.g. cancer with a treatment for cancer. Subjects

having cancer can be identified by a physician using current methods of diagnosing cancer. For example, symptoms and/or complications of lung cancer which characterize these conditions and aid in diagnosis are well known in the art and include but are not limited to, weak breathing, swollen lymph nodes above the collarbone, abnormal sounds in the lungs, dullness when the chest is tapped, and chest pain. Tests that may aid in a diagnosis of, e.g. lung cancer include, but are not limited to, x-rays, blood tests for high levels of certain substances (e.g. calcium), CT scans, and tumor biopsy. A family history of lung cancer, or exposure to risk factors for lung cancer (e.g. smoking or exposure to smoke and/or air pollution) can also aid in determining if a subject is likely to have lung cancer or in making a diagnosis of lung cancer.

**[0123]** Cancer can include, but is not limited to, carcinoma, including adenocarcinoma, lymphoma, blastoma, melanoma, sarcoma, leukemia, squamous cell cancer, small-cell lung cancer, non-small cell lung cancer, gastrointestinal cancer, Hodgkin's and non Hodgkin's lymphoma, pancreatic cancer, glioblastoma, basal cell carcinoma, biliary tract cancer, bladder cancer, brain cancer including glioblastomas and medulloblastomas; breast cancer, cervical cancer, choriocarcinoma; colon cancer, colorectal cancer, endometrial carcinoma, endometrial cancer; esophageal cancer, gastric cancer; various types of head and neck cancers, intraepithelial neoplasms including Bowen's disease and Paget's disease; hematological neoplasms including acute lymphocytic and myelogenous leukemia; Kaposi's sarcoma, hairy cell leukemia; chronic myelogenous leukemia, AIDS-associated leukemias and adult T-cell leukemia lymphoma; kidney cancer such as renal cell carcinoma, T-cell acute lymphoblastic leukemia/lymphoma, lymphomas including Hodgkin's disease and lymphocytic lymphomas; liver cancer such as hepatic carcinoma and hepatoma, Merkel cell carcinoma, melanoma, multiple myeloma; neuroblastomas; oral cancer including squamous cell carcinoma; ovarian cancer including those arising from epithelial cells, sarcomas including leiomyosarcoma, rhabdomyosarcoma, liposarcoma, fibrosarcoma, and osteosarcoma; pancreatic cancer; skin cancer including melanoma, stromal cells, germ cells and mesenchymal cells; prostate cancer, rectal cancer; vulval cancer, renal cancer including adenocarcinoma; testicular cancer including germinal tumors such as seminoma, non-seminoma (teratomas, choriocarcinomas), stromal tumors, and germ cell tumors; thyroid cancer including thyroid adenocarcinoma and medullar carcinoma; esophageal cancer, salivary gland carcinoma, and Wilms' tumors. In some embodiments, the cancer can be lung cancer.

**[0124]** In some embodiments, the methods described herein comprise administering an effective amount of compositions described herein, e.g. a treatment for cancer to a subject in order to alleviate a symptom of a cancer. As used herein, "alleviating a symptom of a cancer" is ameliorating any condition or symptom associated with the cancer. As compared with an equivalent untreated control, such reduction is by at least 5%, 10%, 20%, 40%, 50%, 60%, 80%, 90%, 95%, 99% or more as measured by any standard technique. A variety of means for administering the compositions described herein to subjects are known to those of skill in the art. Such methods can include, but are not limited to oral, parenteral, intravenous, intramuscular, subcutaneous, transdermal, airway (aerosol), pulmonary, cutaneous, topical, injection, or intratumoral administration.

Administration can be local or systemic. The term "effective amount" as used herein refers to the amount of a treatment needed to alleviate at least one or more symptom of the disease or disorder, and relates to a sufficient amount of pharmacological composition to provide the desired effect. The term "therapeutically effective amount" therefore refers to an amount that is sufficient to effect a particular anti-cancer effect when administered to a typical subject. An effective amount as used herein, in various contexts, would also include an amount sufficient to delay the development of a symptom of the disease, alter the course of a symptom of the disease (for example but not limited to, slowing the progression of a symptom of the disease), or reverse a symptom of the disease. Thus, it is not generally practicable to specify an exact "effective amount". However, for any given case, an appropriate "effective amount" can be determined by one of ordinary skill in the art using only routine experimentation. The effects of any particular dosage can be monitored by a suitable bioassay. The dosage can be determined by a physician and adjusted, as necessary, to suit observed effects of the treatment.

**[0125]** Non-limiting examples of a treatment for cancer can include radiation therapy, surgery, gemcitabine, cisplatin, paclitaxel, carboplatin, bortezomib, AMG479, vorinostat, rituximab, temozolomide, rapamycin, ABT-737, PI-103; alkylating agents such as thiotepa and CYTOXAN® cyclophosphamide; alkyl sulfonates such as busulfan, improsulfan and piposulfan; aziridines such as benzdopa, carboquone, meturedopa, and uredopa; ethylenimines and methylamelamines including altretamine, triethylenemelamine, triethylenephosphoramide, triethylenethiophosphoramide and trimethylolomelamine; acetogenins (especially bullatacin and bullatacinone); a camptothecin (including the synthetic analogue topotecan); bryostatin; calystatin; CC-1065 (including its adozelesin, carzelesin and bizelesin synthetic analogues); cryptophycins (particularly cryptophycin 1 and cryptophycin 8); dolastatin; duocarmycin (including the synthetic analogues, KW-2189 and CB1-TM1); eleutherobin; pancratistatin; a sarcodictyin; spongistatin; nitrogen mustards such as chlorambucil, chlornaphazine, cholophosphamide, estramustine, ifosfamide, mechlorethamine, mechlorethamine oxide hydrochloride, melphalan, novembichin, phenesterine, prednimustine, trofosfamide, uracil mustard; nitrosoureas such as carmustine, chlorozotocin, fotemustine, lomustine, nimustine, and ranimustine; antibiotics such as the enediyne antibiotics (e.g., calicheamicin, especially calicheamicin gammaII and calicheamicin omegaII (see, e.g., Agnew, Chem. Intl. Ed. Engl., 33: 183-186 (1994))); dynemicin, including dynemicin A; bisphosphonates, such as clodronate; an esperamicin; as well as neocarzinostatin chromophore and related chromoprotein enediyne antibiotic chromophores), aclacinomysins, actinomycin, authramycin, azaserine, bleomycins, cactinomycin, carabycin, caminomycin, carzinophilin, chromomycin, dactinomycin, daunorubicin, detorubicin, 6-di-azo-5-oxo-L-norleucine, ADRIAMYCIN® doxorubicin (including morpholino-doxorubicin, cyanomorpholino-doxorubicin, 2-pyrrolino-doxorubicin and deoxydoxorubicin), epirubicin, esorubicin, idarubicin, marcellomycin, mitomycins such as mitomycin C, mycophenolic acid, nogalamycin, olivomycins, peplomycin, potfiromycin, puromycin, quelamycin, rodorubicin, streptonigrin, streptozocin, tubercidin, ubenimex, zinostatin, zorubicin; anti-metabolites such as methotrexate and 5-fluorouracil

(5-FU); folic acid analogues such as denopterin, methotrexate, pteropterin, trimetrexate; purine analogs such as fludarabine, 6-mercaptopurine, thiamiprine, thioguanine; pyrimidine analogs such as ancitabine, azacitidine, 6-azauridine, carmofur, cytarabine, dideoxyuridine, doxifluridine, enocitabine, floxuridine; androgens such as calusterone, dromostanolone propionate, epitio stanol, mepitio stanol, testosterone; anti-adrenals such as aminoglutethimide, mitotane, trilostane; folic acid replenisher such as frolic acid; aceglatone; aldophosphamide glycoside; aminolevulinic acid; eniluracil; amsacrine; bestrabucil; bisantrene; edatraxate; defofamine; demecolcine; diaziquone; elformithine; elliptinium acetate; an epothilone; etoglucid; gallium nitrate; hydroxyurea; lentinan; lonidainine; maytansinoids such as maytansine and ansamitocins; mitoguazone; mitoxantrone; mopidanmol; nitraerine; pentostatin; phenamet; pirarubicin; losoxantrone; podophyllinic acid; 2-ethylhydrazide; procarbazine; PSK® polysaccharide complex (JHS Natural Products, Eugene, Oreg.); razoxane; rhizoxin; sizofuran; spirogermanium; tenuazonic acid; triaziquone; 2,2',2"-trichlorotriethylamine; trichothecenes (especially T-2 toxin, verracurin A, roridin A and anguidine); urethan; vindesine; dacarbazine; mannomustine; mitobronitol; mitolactol; pipobroman; gacytosine; arabinoside ("Ara-C"); cyclophosphamide; thiotepa; taxoids, e.g., TAXOL® paclitaxel (Bristol-Myers Squibb Oncology, Princeton, N.J.), ABRAXANE® Cremophor-free, albumin-engineered nanoparticle formulation of paclitaxel (American Pharmaceutical Partners, Schaumburg, Ill.), and TAXOTERE® doxetaxel (Rhone-Poulenc Rorer, Antony, France); chloranbucil; GEMZAR® gemcitabine; 6-thioguanine; mercaptopurine; methotrexate; platinum analogs such as cisplatin, oxaliplatin and carboplatin; vinblastine; platinum; etoposide (VP-16); ifosfamide; mitoxantrone; vincristine; NAVELBINE® vinorelbine; novantrone; teniposide; edatrexate; daunomycin; aminopterin; xeloda; ibandronate; irinotecan (Camptosar, CPT-11) (including the treatment regimen of irinotecan with 5-FU and leucovorin); topoisomerase inhibitor RFS 2000; difluoromethylornithine (DMFO); retinoids such as retinoic acid; capecitabine; combretastatin; leucovorin (LV); oxaliplatin, including the oxaliplatin treatment regimen (FOLFOX®); lapatinib (Tykerb®); inhibitors of PKC-alpha, Raf, H-Ras, EGFR (e.g., erlotinib (Tarceva®)) and VEGF-A that reduce cell proliferation and pharmaceutically acceptable salts, acids or derivatives of any of the above. In addition, the methods of treatment can further include the use of radiation or radiation therapy. Further, the methods of treatment can further include the use of surgical treatments.

**[0126]** In some embodiments, the methods described herein can be applicable for resequencing, e.g. for confirming particularly relevant, low-quality, and/or complex sequences obtained by non-directed sequencing of a large amount of nucleic acids. By way of non-limiting examples, the methods described herein can allow the directed and/or targeted resequencing of targeted disease gene panels (e.g. 10-100 genes), resequencing to confirm variants obtained in large scale sequencing projects, whole exome resequencing, and/or targeted resequencing for detection of single nucleotide variants, multiple nucleotide variants, insertions, deletions, copy number changes, and methylation status.

**[0127]** In some embodiments, the methods described herein can allow microbiota sequencing, ancient sample sequencing, and/or new variant virus genotyping.

**[0128]** All patents and other publications; including literature references, issued patents, published patent applications, and co-pending patent applications; cited throughout this application are expressly incorporated herein by reference for the purpose of describing and disclosing, for example, the methodologies described in such publications that might be used in connection with the technology described herein. These publications are provided solely for their disclosure prior to the filing date of the present application. Nothing in this regard should be construed as an admission that the inventors are not entitled to antedate such disclosure by virtue of prior invention or for any other reason. All statements as to the date or representation as to the contents of these documents is based on the information available to the applicants and does not constitute any admission as to the correctness of the dates or contents of these documents.

**[0129]** The description of embodiments of the disclosure is not intended to be exhaustive or to limit the disclosure to the precise form disclosed. While specific embodiments of, and examples for, the disclosure are described herein for illustrative purposes, various equivalent modifications are possible within the scope of the disclosure, as those skilled in the relevant art will recognize. For example, while method steps or functions are presented in a given order, alternative embodiments may perform functions in a different order, or functions may be performed substantially concurrently. The teachings of the disclosure provided herein can be applied to other procedures or methods as appropriate. The various embodiments described herein can be combined to provide further embodiments. Aspects of the disclosure can be modified, if necessary, to employ the compositions, functions and concepts of the above references and application to provide yet further embodiments of the disclosure. These and other changes can be made to the disclosure in light of the detailed description. All such modifications are intended to be included within the scope of the appended claims.

**[0130]** Specific elements of any of the foregoing embodiments can be combined or substituted for elements in other embodiments. Furthermore, while advantages associated with certain embodiments of the disclosure have been described in the context of these embodiments, other embodiments may also exhibit such advantages, and not all embodiments need necessarily exhibit such advantages to fall within the scope of the disclosure.

**[0131]** The technology described herein is further illustrated by the following examples which in no way should be construed as being further limiting.

**[0132]** Some embodiments of the technology described herein can be defined according to any of the following numbered paragraphs:

**[0133]** 1. A method of determining the nucleotide sequence contiguous to a known target nucleotide sequence, the method comprising;

**[0134]** (a) ligating a target nucleic acid comprising the known target nucleotide sequence with a universal oligonucleotide tail-adaptor;

**[0135]** (b) amplifying a portion of the target nucleic acid and the amplification strand of the universal oligonucleotide tail-adaptor with a first adaptor primer and a first target-specific primer;

- [0136] (c) amplifying a portion of the amplicon resulting from step (b) with a second adaptor primer and a second target-specific primer;
- [0137] (d) sequencing the amplified portion from step (c) using a first and second sequencing primer;
- [0138] wherein the universal oligonucleotide tail-adaptor comprises a first ligatable duplex end and a second unpaired end;
- [0139] wherein the universal oligonucleotide tail-adaptor comprises a blocking strand and an amplification strand;
- [0140] wherein the blocking strand comprises a 5' duplex portion;
- [0141] wherein the amplification strand comprises an unpaired 5' portion, a 3' duplex portion, and a 3' T overhang;
- [0142] wherein the amplification strand comprises nucleic acid sequences identical to a first and second sequencing primers;
- [0143] wherein the duplex portions of the blocking strand and the amplification strand are substantially complementary and form the first ligatable duplex end comprising a 3' T overhang;
- [0144] wherein the duplex portion is of sufficient length to remain in duplex form at the ligation temperature;
- [0145] wherein the first target-specific primer comprises a nucleic acid sequence that can specifically anneal to the known target nucleotide sequence of the target nucleic acid at the annealing temperature;
- [0146] wherein the second target-specific primer comprises a 3' portion comprising a nucleic acid sequence that can specifically anneal to a portion of the known target nucleotide sequence comprised by the amplicon resulting from step (b), and a 5' portion comprising a nucleic acid sequence that is identical to a second sequencing primer and the second target-specific primer is nested with respect to the first target-specific primer;
- [0147] wherein the first adaptor primer comprises a nucleic acid sequence identical to a 5' portion of the first sequencing primer; and
- [0148] wherein the second adaptor primer comprises a nucleic acid sequence identical to a portion of the first sequencing primer and is nested with respect to the first adaptor primer.
- [0149] 2. The method of paragraph 1, wherein the blocking strand of the universal oligonucleotide tail-adaptor further comprises a 3' unpaired portion which is not substantially complementary to the 5' unpaired portion of the amplification strand; and
- [0150] wherein the 3' unpaired portion of the blocking strand is not substantially complementary to or substantially identical to any of the primers.
- [0151] 3. The method of any of paragraphs 1-2, wherein the second adaptor primer is nested with respect to the first adaptor primer by at least 3 nucleotides.
- [0152] 4. The method of any of paragraphs 1-3, wherein the portion of the amplification strand that comprises a nucleic acid sequence identical to a first and second sequencing primers is comprised, at least in part, by the 5' unpaired portion of the amplification strand.
- [0153] 5. The method of any of paragraphs 1-4, wherein the first target-specific primer further comprises a 5' tag sequence portion comprising a nucleic acid sequence of high GC content which is not substantially complementary to or substantially identical to any other portion of any of the primers.
- [0154] 6. The method of any of paragraphs 1-5, wherein the second adaptor primer is identical to the full-length first sequencing primer.
- [0155] 7. The method of any of paragraphs 1-6, wherein the portions of the target-specific primers that specifically anneal to the known target will anneal specifically at a temperature of about 65° C. in a PCR buffer.
- [0156] 8. The method of any of paragraphs 1-7, wherein the method further comprises; prior to step (a), the steps of:
- [0157] mechanically shearing the nucleic acid;
- [0158] subjecting the nucleic acid to end-repair;
- [0159] subjecting the nucleic acid to phosphorylation;
- [0160] and subjecting the nucleic acid to adenylation.
- [0161] 9. The method of any of paragraphs 1-8, wherein the sample comprises genomic DNA.
- [0162] 10. The method of any of paragraphs 1-9, wherein the sample comprises RNA and the method further comprises a first step of subjecting the sample to a reverse transcriptase regimen.
- [0163] 11. The method of any of paragraphs 1-10, wherein the reverse transcriptase regimen comprises the use of random hexamers.
- [0164] 12. The method of any of paragraphs 1-11, wherein the known target sequence is comprised by a gene rearrangement.
- [0165] 13. The method of paragraph 12 wherein the gene rearrangement is present in a nucleic acid selected from the group consisting of:
- [0166] genomic DNA; RNA; and cDNA.
- [0167] 14. The method of any of paragraphs 12-13, wherein the gene rearrangement comprises an oncogene.
- [0168] 15. The method of paragraph 14, wherein the gene rearrangement comprises a fusion oncogene.
- [0169] 16. The method of any of paragraphs 1-15, wherein the nucleic acid product is sequenced by a next-generation sequencing method.
- [0170] 17. The method of paragraph 16, wherein the next-generation sequencing method comprises a method selected from the group consisting of:
- [0171] Ion Torrent™, Illumina, SOLiD™, 454™; Massively Parallel Signature Sequencing™ solid-phase, reversible dye-terminator sequencing; and DNA nanoball sequencing.
- [0172] 18. The method of any of paragraphs 1-17, wherein the first and second sequencing primers are compatible with the selected next-generation sequencing method.
- [0173] 19. The method of any of paragraphs 1-18, wherein the method comprises contacting the sample, or separate portions of the sample, with a plurality of sets of first and second target-specific primers.
- [0174] 20. The method of any of paragraphs 1-19, wherein the method comprises contacting a single reaction mixture comprising the sample with a plurality of sets of first and second target-specific primers.
- [0175] 21. The method of any of paragraphs 1-20, wherein the plurality of sets of first and second target-

- specific primers specifically anneal to known target nucleotide sequences comprised by separate genes.
- [0176] 22. The method of any of paragraphs 1-21, wherein at least two sets of first and second target-specific primers specifically anneal to different portions of a known target nucleotide sequence.
- [0177] 23. The method of any of paragraphs 1-22, wherein at least two sets of first and second target-specific primers specifically anneal to different portions of a single gene comprising a known target nucleotide sequence.
- [0178] 24. The method of any of paragraphs 1-23, wherein at least two sets of first and second target-specific primers specifically anneal to different exons of a gene comprising a known nucleotide target sequence.
- [0179] 25. The method of any of paragraphs 19-24, wherein the plurality of first target-specific primers comprise identical 5' tag sequence portions.
- [0180] 26. The method of any of paragraphs 1-25, wherein the universal oligonucleotide tail-adaptor further comprises a barcode portion.
- [0181] 27. The method of paragraph 26, wherein multiple samples are each contacted with a universal oligonucleotide tail-adaptor with a unique barcode portion and wherein the samples are pooled after step (a).
- [0182] 28. The method of any of paragraphs 1-27, wherein each amplification step comprises a set of cycles of a PCR amplification regimen from 5 cycles to 20 cycles in length.
- [0183] 29. The method of any of paragraphs 1-28, wherein the target-specific primers and the adaptor primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of from about 61 to 72° C.
- [0184] 30. The method of any of paragraphs 1-29, wherein the target-specific primers and the adaptor primers are designed such that they will specifically anneal to their complementary sequences at an annealing temperature of about 65° C.
- [0185] 31. The method of any of paragraphs 1-30, wherein the sample comprises a biological sample obtained from a subject.
- [0186] 32. The method of any of paragraphs 1-31, wherein the sample is obtained from a subject in need of treatment for a disease associated with a genetic alteration.
- [0187] 33. The method of paragraph 32, wherein the disease is cancer.
- [0188] 34. The method of any of paragraphs 1-33, wherein the sample comprises a population of tumor cells.
- [0189] 35. The method of any of paragraphs 1-34, wherein the sample comprises a tumor biopsy.
- [0190] 36. The method of any of paragraphs 1-35, wherein the cancer is lung cancer.
- [0191] 37. The method of any of paragraphs 1-36, wherein the known target sequence is comprised by a disease-associated gene.
- [0192] 38. The method of any of paragraphs 1-37, wherein the known target sequence is comprised by a gene rearrangement product in the sample.
- [0193] 39. The method of any of paragraphs 1-38, wherein gene rearrangement product is an oncogene.
- [0194] 40. The method of any of paragraphs 1-39, wherein the known target sequence comprises sequence from a gene selected from the group of:
- [0195] ALK; ROS1; and RET.
- [0196] 41. The method of paragraph 40, wherein at least one set of a first target-specific primer and a second target-specific primer are selected from the group consisting of;
- [0197] SEQ ID NOs: 5 and 6; SEQ ID NOs: 7 and 8; SEQ ID NOs: 9 and 10; SEQ ID NOs: 11 and 12; SEQ ID NOs: 13 and 14; SEQ ID NOs: 15 and 16; SEQ ID NOs: 17 and 18; SEQ ID NOs: 19 and 20; SEQ ID NOs: 21 and 22; SEQ ID NOs: 23 and 24; SEQ ID NOs: 25 and 26; SEQ ID NOs: 27 and 28; SEQ ID NOs: 29 and 30; SEQ ID NOs: 31 and 32; SEQ ID NOs: 33 and 34; SEQ ID NOs: 35 and 36; and SEQ ID NOs: 37 and 38.
- [0198] 42. The method of any of paragraphs 40-41, wherein the presence of a gene rearrangement of ALK in a sample obtained from a tumor in a subject indicates that the tumor is susceptible to treatment with a treatment selected from the group consisting of:
- [0199] an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.
- [0200] 43. The method of any of paragraphs 40-41, wherein the presence of a gene rearrangement of ROS1 in a sample obtained from a tumor in a subject indicates that the tumor is susceptible to treatment with a treatment selected from the group consisting of:
- [0201] a ROS inhibitor; an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.
- [0202] 44. The method of any of paragraphs 40-41, wherein the presence of a gene rearrangement of RET in a sample obtained from a tumor in a subject indicates that the tumor is susceptible to treatment with a treatment selected from the group consisting of:
- [0203] a RET inhibitor; DP-2490; DP-3636; SU5416; BAY 43-9006; BAY 73-4506 (regorafenib); ZD6474; NVP-AST487; sorafenib; RPI-1; XL184; vandetanib; sunitinib; imatinib; pazopanib; axitinib; motesanib; gefitinib; and withaferin A.
- [0204] 45. A method of treating cancer, the method comprising;
- [0205] detecting, in a tumor sample obtained from a subject in need of treatment for cancer, the presence of one or more oncogene rearrangements according to the method of any of paragraphs 1-44;
- [0206] administering a cancer treatment which is effective against tumors having any of the detected oncogene rearrangements.
- [0207] 46. The method of paragraph 45, wherein a treatment selected from the group consisting of:
- [0208] an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684;
- [0209] is effective against tumors having an ALK oncogene rearrangement.

- [0210]** 47. The method of paragraph 45, wherein a treatment selected from the group consisting of:
- [0211]** a ROS1 inhibitor; an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684;
- [0212]** is effective against tumors having an ROS1 oncogene rearrangement.
- [0213]** 48. The method of paragraph 45, wherein a treatment selected from the group consisting of:
- [0214]** a RET inhibitor; DP-2490; DP-3636; SU5416; BAY 43-9006; BAY 73-4506 (regorafenib); ZD6474; NVP-AST487; sorafenib; RPI-1; XL184; vandetanib; sunitinib; imatinib; pazopanib; axitinib; motesanib; gefitinib; and withaferin A;
- [0215]** is effective against tumors having an RET oncogene rearrangement.
- [0216]** 49. A method of determining if a subject in need of treatment for cancer will be responsive to a given treatment, the method comprising:
- [0217]** detecting, in a tumor sample obtained from the subject, the presence of an oncogene rearrangement according to the method of any of paragraphs 1-44;
- [0218]** wherein the subject is determined to be responsive to a treatment targeting an oncogene rearrangement product if the presence of the oncogene rearrangement is detected.
- [0219]** 50. The method of paragraph 49, wherein if the presence of an ALK oncogene rearrangement is detected, the subject will be responsive to a treatment selected from the group consisting of:
- [0220]** an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.
- [0221]** 51. The method of paragraph 49, wherein if the presence of an ROS1 oncogene rearrangement is detected, the subject will be responsive to a treatment selected from the group consisting of:
- [0222]** an ALK inhibitor; crizotinib (PF-02341066); AP26113; LDK378; 3-39; AF802; IPI-504; ASP3026; AP-26113; X-396; GSK-1838705A; CH5424802; and NVP-TAE684.
- [0223]** 52. The method of paragraph 49, wherein if the presence of an RET oncogene rearrangement is detected, the subject will be responsive to a treatment selected from the group consisting of:
- [0224]** a RET inhibitor; DP-2490; DP-3636; SU5416; BAY 43-9006; BAY 73-4506 (regorafenib); ZD6474; NVP-AST487; sorafenib; RPI-1; XL184; vandetanib; sunitinib; imatinib; pazopanib; axitinib; motesanib; gefitinib; and withaferin A.
- [0225]** 53. The method of any of paragraphs 44-52, wherein the cancer is lung cancer.

## EXAMPLES

### Example 1

**[0226]** In one embodiment, described herein is an assay using a novel half-truncated “Y” shape adapter for next generation sequencing library construction followed by two single-end nested polymerase chain reactions enables rapid and efficient target enrichment using RNA or genomic DNA

as template from fresh or formalin-fixed and paraffin-embedded (FFPE) specimens. This enrichment method enables targeted resequencing of cDNA or gDNA for potential detection of genetic alterations (single nucleotide variants, insertions/deletions, and copy number), epigenetic alterations (methylation), gene expression, and genomic rearrangements.

**[0227]** Target enrichment prior to next-generation sequencing is more cost-effective than whole genome, whole exome, and whole transcriptome sequencing and therefore more practical for broad implementation both for research discovery and clinical applications. High coverage depth afforded by target enrichment approaches enables wider dynamic range for allele counting (in gene expression and copy number assessment) and detection of low frequency mutations, a critical feature for evaluating somatic mutations in cancer. Before broad implementation of whole genome or whole exome sequencing is possible, the mainstay of clinical next generation sequencing will involve select disease panels with a discrete number of gene targets. Likewise, research studies for analysis of large sample sizes based on defined gene targets will also require an economical means for genotyping. The assays described herein would be applicable in both of these settings.

**[0228]** For the detection of inter and intra-chromosomal rearrangements, whole genome or transcriptome sequencing is useful for novel discovery of genomic rearrangements and does not require prior knowledge of the involved gene/chromosomal partners. However, these whole genome and transcriptome approaches are not practical currently in the clinical setting due to high cost, low sequencing depth resulting in poor sensitivity, and highly demanding bioinformatics analysis. Fluorescence in situ hybridization (FISH) has been the gold standard assay for the detection of genomic rearrangements in clinics; however, the assay is low-throughput and its implementation requires special equipment and expert knowledge/experience. RT-PCR is also effective at detecting such rearrangements but requires knowledge of both the 5' and 3' partners and is not scalable for a large number of targets/samples.

**[0229]** Examples of commonly used preparatory enrichment assays for next generation sequencing include hybridization-based capture assays (TruSeq Capture™, Illumina; SureSelect Hybrid Capture™, Agilent) and polymerase chain reaction (PCR)-based assays (HaloPlex™ Agilent; AmpliSeq™, Ion Torrent™; TruSeq Amplicon™, Illumina; emulsion/digital PCR, Raindance). Hybridization-based approaches capture not only the targeted sequences covered by the capture probes but also near off-target bases that consume additional sequencing capacity. In addition, these methods are relatively time-consuming, labor-intensive, and less specific. A PCR amplification based approach is simpler and faster but by conventional design requires both forward and reverse primers flanking target loci. In particular for detection of genomic rearrangements with unknown fusion partners, PCR is not applicable.

**[0230]** Described herein is a target enrichment assay using a novel half-truncated “Y” shape adapter for next generation sequencing library construction, enabling rapid sequencing of cDNA or gDNA from fresh or FFPE specimens. FIG. 1 outlines the library construction for target enrichment using a half-truncated Y adapter for Ion Torrent™ sequencing as an example. Importantly, this method is adaptable for any other next generation sequencing platform, including but not

limited to Illumina, SOLiD™, and 454™. In short a randomly sheared, double-stranded cDNA or gDNA template can be end-repaired, adenylated, and then ligated on one or both ends with the universal Y adapter to create a universal sequence for initiating PCR and subsequent sequencing. An initial round of PCR using a target specific primer tagged with a stuffer tail (the stuffer tail aids in multiplexing a high number of targets) and a primer same as the 20 bases at 5' Y adapter overhang. A second round of PCR is carried out with a primer same as the 30 bases at the 5' Y adapter overhang and a second tandem nested target specific primer that anneals 3' downstream of the first target specific primer. The second tandem nested primer is 5' tagged with the second primer sequence required for downstream emulsion PCR or clustering depending on the sequencing technology. High specificity in this system is achieved with the unidirectional tandem, nested primers which will effectively cover a target sequence of 36-40 bases or more (depending on the spacing between the primers). Of note, the number of PCR cycles may be optimized depending on how much starting nucleic acid material is used, the number of pooled samples, and the number of targets.

**[0231]** In summary, the method utilizes a half-truncated Y adapter to universally tag all dsDNA fragments with a common 3' end which is utilized for two rounds of PCR using two unidirectional, nested target specific primers for specificity. The application of tagged nested primers also avoids the effects of primer homodimerization and heterodimerization when targeting many different sites in the genome or transcriptome. In addition to multiplexing targets, the methods described herein also allow for multiplex sample pooling after the Y adapter ligation (Step 2) during which individual samples are ligated with unique barcoded adapters. Once barcoded via Y adapter ligation, multiple samples may be pooled into one reaction tube for downstream target enrichment.

**[0232]** In contrast to detection of genomic rearrangements which is more tolerant of insertion and deletion sequencing errors such as those found on Ion Torrent™ and 454™ sequencing as a result of homopolymers, detection of single base and multiple base mutations (including insertions and deletions) requires higher accuracy sequencing achievable with the Illumina sequencing platform. For this purpose, the half-truncated Y adapted library can be converted to an Illumina library, e.g. by amplifying the overhanging, truncated 3' arm of the Y adapter using a primer tagged with the Illumina forward adapter sequence (barcoded or non-barcoded), and tagged gene specific nested primers (GSP2s tagged with Illumina reversed primer). Similarly, Bi-directional sequencing is also achievable by introducing a sequencing primer via the overhanging, truncated 3' Y adapter arm using a tagged primer during the first PCR, Step 3.

**[0233]** The assays and methods described herein have been applied on cDNA derived from FFPE specimens for the detection of ROS1 gene fusions using seven gene specific primers targeting seven exons comprising the ROS1 kinase domain. ROS1 genomic rearrangements were detected with known and previously unknown fusion partners such as “SLC34A2 Exon 12-ROS1 Exon 34” and “EZR Exon 9-ROS1 Exon 34” fusions. The assay achieved high on-target specificity (~85-95% when mapping using human genome hg19 reference), which enables the sequencing of multiple samples even with the smallest scale sequencing

platforms (and thus least expensive) such as the Ion Torrent PGM 314™ chip with high coverage results (7 targeted loci, 5 samples, >1000× coverage for each target per sample). FIG. 2 shows mapping of sequencing reads to the target loci in gene ROS1 kinase domain.

**[0234]** Advantages of the methods described herein include:

**[0235]** 1. The methods described herein can be utilized for target enrichment of double stranded cDNA or gDNA from fresh or FFPE samples, allowing mapping of both 5' and 3' ends of target cDNA or gDNA fragments, using next-generation sequencing. Current hybridization-based approaches require days of preparation and hybridization, and show lower specificity with capture of near off-target sequences. Current amplification based approaches require known forward and reverse primers by design and are not amenable for high scale multiplexing of targets.

**[0236]** 2. Simple Bioinformatics analysis. Depending on the number of selected targets, analysis of data generated from this assay is simple and fast. In addition, it may be set up to be compatible with current existing bioinformatics tools. Thus, small-size clinical labs may be able to perform data analysis without the need for significant investment in bioinformatics, which will be a limitation for broad clinical implementation of next generation sequencing.

**[0237]** 3. High specificity (~85-95%). A conventional Y adapter construct having both components of forward and reverse primers (5' and 3' overhangs) for routine library construction will introduce a high level of background off-target sequencing due to carry over of “sequenceable” starting genomic or transcriptome library material. For instance, if the target is of size of 100 bp and present in two copies in one genome, the ratio of target-to-genome is about  $1:3 \times 10^7$ . Hybridization-based approaches normally employ biotinylated oligonucleotide baits to pull out hybridized target fragments by streptavidin coated magnetic beads. Just one non-specific binding event out of the possible  $3 \times 10^7$  possibilities could dramatically result in a 50% off-target rate. The use of a half-truncated Y adapter effectively avoids this background carry over issue, since the starting library material cannot be amplified in subsequent preparatory steps for sequencing. Additional specificity is achieved by using two unidirectional primers, with GSP2 3' downstream of GSP1. Effectively, the use of the two tandem primers yields a target priming site of 40 (assume two 20 base pair length primers) and higher specificity than with just one primer. The use of 5' 20-mer primer in the first PCR and the full 30-mer primer (serves as a nested primer for the universal priming site) in the second PCR further increases specificity. Finally, additional specificity is achieved with the use of tagged primers for both PCR steps. Tagged primers, through intramolecular hairpin formation, prevent the propagation of primer homodimers and heterodimers which could overtake the PCR reactions and lead to nonspecific and undesired artifactual products; therefore, tagged primers allow the ability to multiplex many targets while avoiding primer dimers.

**[0238]** 4. Economical cost. The key components in the methods described herein are conventional, unmodified tagged primers, standard PCR reagents, and routine thermocycling. Unlike hybridization based capture methods or microfluidic digital PCR setups, the described target enrichment protocol avoids the use of relatively expensive biotinylated oligonucleotides, streptavidin coated magnetic

beads, and special equipment. Once manufactured, pooled GSP1 and GSP2 primer mixes may be used for thousands of reactions.

**[0239]** 5. Automation. Because the method described herein relies on standard PCR techniques and SPRI cleanup, there is potential for facile automation in high throughput applications. Volumes may be adjusted for 384 well plates for ultra-high throughput implementation.

**[0240]** Applications of the methods and assays described herein include, but are not limited to: 1. A lung cancer translocation panel consisting of known therapeutic targets including genes ALK, ROS1 and RET; 2. hematological malignancy panels including those to detect genomic rearrangements in lymphomas and leukemias; 3. Sarcoma genomic rearrangement panel; 4. IGH/TCR gene rearrangement panel for lymphoma testing; 5. Targeted disease gene panel for resequencing (10-100 genes); 6. Targeted resequencing for confirming variants from large scale sequencing projects; 7. Potential for whole exome targeted resequencing; 8. Targeted resequencing for detection of single nucleotide variants, multiple nucleotide variants, insertions, deletions, copy number changes, and methylation status; 9. Microbiota sequencing; 10. Ancient sample sequencing; 11. New variant virus genotyping.

#### Example 2: Targeted Next-Generation Sequencing Assay for Simultaneous Detection of ALK, ROS1 and RET Gene Rearrangements

**[0241]** Knowledge of chromosomal rearrangement status in cancer is important for individualized targeted therapy. Recently, three major receptor tyrosine kinases involved in rearrangements in lung cancer have been described. Gene rearrangements involving the ALK gene has been established as a therapeutic target. Early clinical trial data also suggest that a ROS1 inhibitor is effective in treating patients testing positive for the ROS1 rearrangement. In vitro evidence has shown that tumor cells harbouring the RET rearrangement are responsive to a RET inhibitor. Thus, ALK, ROS1, and RET currently represent three important therapeutic targets in lung cancer.

**[0242]** Current clinical assays for the detection of gene rearrangements include fluorescence in situ hybridization (FISH), immunohistochemistry (IHC), and reverse transcription polymerase chain reaction (RT-PCR). FISH and IHC may not be able to accommodate the increasing demand for high volume testing/screening due to their low throughput, high cost, and complex interpretation. RT-PCR assays require full knowledge of both fusion partners which are sometimes unavailable and effectively impact clinical sensitivity. Multiplex RT-PCR has been used for detection of variant fusions involving different exons between two known fusion partners. In general, these assays are limited to a one gene target at a time approach requiring multiple reaction setups and analysis.

**[0243]** The latest next-generation sequencing-based assays for transcriptome or target capture sequencing have been applied largely for the purpose of research and discovery in large sequencing facilities. These assays in the research setting generally achieve low sequencing depth due

to the vast number of targets, thus yielding low analytical sensitivity and subsequent poor clinical sensitivity. Though affordable for large sequencing facilities, these assays are not yet within reach for many clinical laboratories.

**[0244]** Described above are methods and assays related to a targeted sequencing method which applies a novel half-truncated “Y” shape adapter for next generation sequencing library construction followed by two single-end nested polymerase chain reactions enabling rapid and efficient target enrichment using RNA or genomic DNA as template from fresh or formalin-fixed and paraffin-embedded (FFPE) specimens.

**[0245]** Based on this method, described in this Example is a specific assay for the detection of ALK, ROS1, and RET gene rearrangements (FIGS. 3A-3B). Gene specific primers (GSP1) are designed to prime the exons near or on the kinase domain. Nested primers (GSP2) are designed to prime downstream of GSP1 but within the same exons and proximity to their paired GSP1s. The panel currently includes seven pairs of primers for targeting ROS1 exons 31 to 37; four pairs of primers for targeting ALK exons 19 to 22; and six pairs of primers for targeting RET exons 8 to 13.

**[0246]** This assay can be adapted to different NGS sequencing platforms by using platform specific adaptor sequence following the half-functional “Y” adaptor configuration and the GSP2 primers. After sequencing, the reads are mapped to the human genome reference allowing the identification of fusion partner genes, alternative splicing, as well as frame status of the fusions using a bioinformatics algorithm developed by the inventors (e.g. FIGS. 4A and 4B). The resulting output is a simple annotated table for quick reporting (FIG. 4C). Using this three gene target panel assay with multi-sample barcoding, ROS1, ALK and RET rearrangement positive samples have been detected using one Ion Torrent™ sequencing run.

**[0247]** This assay is applicable for degraded RNA, such as RNA extracted from formalin-fixed and paraffin-embedded (FFPE) specimen, which is the most widely available clinical material for molecular diagnostic testing. This assay takes advantage of bench-top NGS platforms, a simple informatics analysis pipeline, and therefore relatively easy implementation for clinical laboratories. Of note, this assay does not require prior knowledge of the fusion partners and would yield high clinical sensitivity in the detection of various gene partners (and corresponding multiple exons) associated with the principle target gene. In addition, the assay will be functional for both unknown 5' upstream or 3' downstream fusion partners as long as one of the partners is known. Multiplexing and deep sequencing allows testing of samples with low tumor cellularity, rare fusions, and rare alternative splicing events. The detailed, patient specific rearrangement information afforded by this assay would be useful for evaluating genotype specific therapeutic response and perhaps a patient specific tumor marker for minimal residual disease monitoring. Based on conventional oligo synthesis reagents, off the shelf enzymes, and the ability to multiplex many samples in one run, this assay will be a cost-effective clinical assay for detecting gene rearrangements.

TABLE 1

Primers adapted for IonTorrent™ platform, v1. Primer names indicate target gene and target gene exon comprising the known target nucleotide sequence. R1 designates a first target-specific primer and R2 designates a second target-specific primer. Detailed in this table is one set of a first target specific primer and one second target-specific for each listed exon of each gene.

Primer Name	Sequence	SEQ ID NO
ALKex19_P1R2	CCTCTCTATGGGCAGTCGGTGATGCGAGAGTGGCAGGTG TGG	5
ALKex19_tag.R1	GGATCTCGACGCTCTCCCTCAGAGGTCACCACAGAGAGG ATCAG	6
ALKex20_P1R2	CCTCTCTATGGGCAGTCGGTGATCATGGCTTGCAGCTCC TGGT	7
ALKex20_tag.R1	GGATCTCGACGCTCTCCCTGCAGCTCCATCTGCATGGCTT G	8
ALKex21_P1R2	CCTCTCTATGGGCAGTCGGTGATGGCCTTCATACACCTC CCCAA	9
ALKex21_tag.R1	GGATCTCGACGCTCTCCCTTTGGGCATTCCGGACACC	10
ALKex22_P1R2	CCTCTCTATGGGCAGTCGGTGATAGGAAATCCAGTTCGT CCTGTT CAGA	11
ALKex22_tag.R1	GGATCTCGACGCTCTCCCTGATCAGGGCTTCATGAGGA AATC	12
RETex10_P1R2	CCTCTCTATGGGCAGTCGGTGATGGCTCCCCAGGCTCGT GT	13
RETex10_tag.R1	GGATCTCGACGCTCTCCCTAGGTGCCATAGCCAGCTTTA ATCC	14
RETex11_P1R2	CCTCTCTATGGGCAGTCGGTGATATCACCGTGCGGCACA GCTC	15
RETex11_tag.R1	GGATCTCGACGCTCTCCCTGAGGACAGCGCTGCGATCA	16
RETex12_P1R2	CCTCTCTATGGGCAGTCGGTGATAGAACCAAGTTCTTCC GAGGGAAT	17
RETex12_tag.R1	GGATCTCGACGCTCTCCCTTCCAAATTCGCCTTCTCCTAG AGTT	18
RETex13_P1R2	CCTCTCTATGGGCAGTCGGTGATACAGCAGGTCTCGCAG CTCAC	19
RETex13_tag.R1	GGATCTCGACGCTCTCCCTTGACCTGCTTCAGGACGTTG AA	20
RETex8_P1R2	CCTCTCTATGGGCAGTCGGTGATCTTGCTGACTGCACAG GACAGG	21
RETex8_tag.R1	GGATCTCGACGCTCTCCCTTCCCTCACACTCCAGCCGTCTC	22
RETex9_P1R2	CCTCTCTATGGGCAGTCGGTGATTGGTGTGGGAGAGCA GGT	23
RETex9_tag.R1	GGATCTCGACGCTCTCCCTCCGTGGGGCAGGTCTTG	24
ROS1Ex31_P1R2	CCTCTCTATGGGCAGTCGGTGATGGCTGCATGAAGTTTT AACATGG	25
ROS1Ex31_tag.R1	GGATCTCGACGCTCTCCCTTGATATTACAGACATAAGCA GGACCTTGG	26
ROS1Ex32_P1R2	CCTCTCTATGGGCAGTCGGTGATCTAGTAATTTGGGAAT GCCTGGTTT	27
ROS1Ex32_tag.R1	GGATCTCGACGCTCTCCCTTTCAGCTTTCTCCCACTGTAT TGAA	28

TABLE 1-continued

Primers adapted for IonTorrent™ platform, v1. Primer names indicate target gene and target gene exon comprising the known target nucleotide sequence. R1 designates a first target-specific primer and R2 designates a second target-specific primer. Detailed in this table is one set of a first target specific primer and one second target-specific for each listed exon of each gene.

Primer Name	Sequence	SEQ ID NO
ROS1Ex33_P1R2	CCTCTCTATGGGCAGTCGGTGATCATCTTCCACCTTAAAT TCTGGTCTGTA	29
ROS1Ex33_tag.R1	GGATCTCGACGCTCTCCCTCAGGATCCATTAAATGTCAT CTTCC	30
ROS1Ex34_P1R2	CCTCTCTATGGGCAGTCGGTGATAGTAAGTATGAAACTT GTTTCTGGTATCC	31
ROS1Ex34_tag.R1	GGATCTCGACGCTCTCCCTGGTCAGTGGGATTGTAACAA CCAG	32
ROS1Ex35_P1R2	CCTCTCTATGGGCAGTCGGTGATCACCCCTTCTTGGCAC TTT	33
ROS1Ex35_tag.R1	GGATCTCGACGCTCTCCCTTCTTTGTCTTCGTTTATAAGC ACTGTC	34
ROS1Ex36_P1R2	CCTCTCTATGGGCAGTCGGTGATTTCAATCTCTCTTGGG TTGGA	35
ROS1Ex36_tag.R1	GGATCTCGACGCTCTCCCTCCGAGGGAAGGCAGGAAGA TT	36
ROS1Ex37_P1R2	CCTCTCTATGGGCAGTCGGTGATCAGGAATTCAATCTTC TCCTGGTC	37
ROS1Ex37_tag.R1	GGATCTCGACGCTCTCCCTCTCATCAGATGTGCCTCCTTC AG	38

SEQUENCE LISTING

Sequence total quantity: 60  
 SEQ ID NO: 1 moltype = DNA length = 6267  
 FEATURE Location/Qualifiers  
 source 1..6267  
 mol\_type = genomic DNA  
 organism = Homo sapiens

SEQUENCE: 1

```

agctgcaagt ggcggggcgc caggcagatg cgatccagcg gctctggggg cggcagcggg 60
ggtagcagct ggtacctccc gccgcctctg ttcggagggt cgcggggcac cgaggtgctt 120
tccggccgcc ctctggtcgg ccacccaaag ccgcgggcgc tgatgatggg tgaggagggg 180
gcggaagat ttcgggcgcc cctgccctga acgcccctag ctgctgccgc cggggccgct 240
ccagtgcctg cgaactctga ggagccgagg cgcgggtgag agcaaggacg ctgcaaaact 300
gcgcagcgcg ggggctggga ttcacgcccc gaagtccagc aggcagacag tccgaagcct 360
tcccgcagcg gagagatagc ttgaggggtg gcaagacggc agcctccgcc ctgggttccc 420
gcccagaccg ggcagaagag cttggaggag ccaaaaggaa cgcaaaaggc ggccaggaca 480
gcgtgcagca gctgggagcc gccgttctca gccttaaaag ttgcagagat tggaggctgc 540
cccagagagg gacagacccc agctccgact gcggggggca ggagaggacg gtaccaact 600
gccacctccc ttcaaccata gtagttcttc tgtaccgagc gcagcagct acagacgggg 660
gcgcggcact cggcgcggag agcgggaggc tcaaggtccc agccagtgag cccagtgtgc 720
ttgagtgtct ctggactcgc ccctgagctt ccaggctctg ttcatttaga ctccctgctc 780
cctccgtgca gttgggggaa agcaagagac ttgcgcgcac gcacagtcct ctggagatca 840
ggtggaagga gccgctgggt accaaggact gttcagagcc tcttccatc tcggggagag 900
gaaaggggta ggctggggcc ggagagcagt gtaaacggcc tcctccggcg ggatgggagc 960
catcgggctc ctgtggctcc tgccgctgct gctttccacg gcagctgtgg gctccgggat 1020
ggggaccggc cagegcgcgg gctccccagc tgcggggcgc ccgctgcagc cccgggagcc 1080
actcagctac tcgcgcctgc agaggaagag tctggcagtt gacttcgtgg tgcctcgtc 1140
cttccgtgtc tacgcccggg acctactgct gccaccatcc tcctcggagc tgaaggctgg 1200
caggcccagag gcccgcggtc cgetagctct ggactgcgcc ccgctgctca ggttgctggg 1260
gccggcgccg ggggtctcct ggaccgcccg ttcaccagcc ccggcagagg cccggagcgt 1320
gtccagggtg ctgaagggcg gctccgtgcg caagctccgg cgtgccaaag agttgggtgt 1380
ggagctgggc gaggaggcga tcttggaggg ttgctgcggg cccccgggg aggcggctgt 1440
  
```

-continued

ggggtgctc	cagttcaatc	tcagcgagct	gttcagttgg	tggattegcc	aaggcgaagg	1500
gcgactgagg	atccgcctga	tgccccgagaa	gaaggcgtcg	gaagtgggca	gagaggggaag	1560
gctgtccgcg	gcaattcgcg	cctcccagcc	ccgccttctc	ttccagatct	tcgggactgg	1620
tcatagctcc	ttggaatcac	caacaaacat	gccttctcct	tctcctgat	atthttacatg	1680
gaatctcacc	tggataatga	aagactcctt	ccctttcctg	tctcatcgca	gccgatatgg	1740
tctggagtgc	agctttgact	tccccgtgta	gctggagtat	tccccctccac	tgcacgacct	1800
caggaaccag	agctggctct	ggcgccgcct	cccctccgag	gaggcctccc	agatggactt	1860
gctggatggg	cctggggcag	agcgttctaa	ggagatgccc	agaggctcct	ttctccttct	1920
caacacctca	gctgactcca	agcacaccat	actgtacccc	tggatgagga	gcagcagtga	1980
gcaactgcaca	ctggccgtct	cggtgcacag	gcacctgcag	ccctctggaa	ggtacattgc	2040
ccagctgctg	ccccacaacg	aggctgcaag	agagatcctc	ctgatgccc	ctccagggaa	2100
gcatggttgg	acagtgcctc	agggaagaat	cgggcgtcca	gacaacccat	ttcgagtggc	2160
cctggaatac	atctccagtg	gaaaccgcag	cttgtctgca	gtggacttct	ttgccctgaa	2220
gaactgcagt	gaaggaacat	ccccaggctc	caagatggcc	ctgcagagct	ccttcacttg	2280
ttggaatggg	acagtctctc	agcttgggca	ggcctgtgac	ttccaccagg	actgtgccc	2340
gggagaagat	gagagccaga	tgtgcccggaa	actgctgtg	ggttttact	gcaacttga	2400
agatggcttc	tgtggctgga	ccccaggcac	actgtcacc	cacactcctc	aatggcaggt	2460
caggacccta	aaggatgccc	ggttccagga	ccaccaagac	catgctctat	tgctcagtac	2520
caactgatgc	cccgtctctg	aaagtgtctc	agtaccagct	gctacgtttc	ctgcaccgat	2580
caagagctct	ccatgtgagc	tccgaatgtc	ctggctcatt	cgtggagtct	tgaggggaaa	2640
cgtgtccttg	gtgctagtgg	agaacaaaac	cgggaaggag	caaggcagga	tggctctggca	2700
tgtccgccc	tatgaaggct	tgagcctgtg	gcagtggatg	gtgttgctc	tcctcgatgt	2760
gtctgacagg	ttctggctgc	agatggctgc	atgggtggga	caaggatcca	gagccatcgt	2820
ggcttttgac	aatatctcca	tcagcctgga	ctgtacctc	accattagcg	gagaggacaa	2880
gatcctgcag	aatacagcac	ccaaatcaag	aaacctgttt	gagagaaacc	caaacaagga	2940
gctgaaaccc	ggggaataat	caccaagaca	gacccccatc	tttgacctca	cagttcattg	3000
gctgttcacc	acatgtgggg	ccagcgggccc	ccatggcccc	accaggcac	agtgcacaaa	3060
cgctaccag	aactccaacc	tgagcgtgga	gggtggggagc	gagggcccc	tgaaggcat	3120
ccagatctgg	aagggtgccag	ccaccgacac	ctacagcatc	tcgggctacg	gagctgctgg	3180
cgggaaaggc	gggaagaaca	ccatgatgcg	gtcccacggc	gtgtctgtgc	tgggcatctt	3240
caacctggag	aaggatgaca	tgtgtacat	cctggttggg	cagcagggag	aggaccctg	3300
ccccagtaca	aaccagttaa	tccagaaagt	ctgcattgga	gagaacaatg	tgatagaaga	3360
agaaatccgt	gtgaacagaa	gcgtgcatga	gtgggcagga	ggcggaggag	gagggggtgg	3420
agccacctac	gtatttaaga	tgaaggatgg	agtcccggtg	cccctgatca	ttgcagccgg	3480
aggtgggtgg	agggcctacg	gggccaagac	agacacgttc	caccagaga	gactggagaa	3540
tactcctcg	gttctagggc	taaacggcaa	ttccggagcc	gcaggtggtg	gaggtggctg	3600
gaatgataac	acttctctgc	tctgggcccg	aaaatctttg	caggaggggtg	ccaccggagg	3660
acattcctgc	ccccaggcca	tgaagaagtg	ggggtgggag	acaagagggg	gtttcggagg	3720
gggtggaggg	gggtgctcct	caggtggagg	aggcggagga	tatataggcg	gcaatgcagc	3780
ctcaaaacat	gaccccga	tggatgggga	agatgggggt	tccttcatca	gtccactggg	3840
catcctgtac	accccagctt	taaaagtgat	ggaaggccac	ggggaagtga	atattaagca	3900
ttatctaaac	tgcagtcact	gtgaggtaga	cgaatgtcac	atggacctg	aaagccacaa	3960
ggtcatctgc	ttctgtgacc	acgggacggg	gctggctgag	gatggcgtct	cctgcattgt	4020
gtcaccacc	ccggagccac	acctgccact	ctcctgac	ctctctgtg	tgacctctgc	4080
cctcgtggcc	gcccgtgtcc	tggtttctc	cgcatcctg	attgtgtacc	gccggaagca	4140
ccaggagctg	caagccatgc	agatggagct	gcagagccct	gagtacaagc	tgagcaagct	4200
ccgcacctcg	accatcatga	ccgactacaa	ccccactac	tgctttgctg	gcaagacctc	4260
ctccatcagt	gacctgaagg	aggtgcccgc	gaaaaacatc	accctcattc	ggggtctggg	4320
ccatggcgcc	tttggggagg	tgtatgaagg	ccaggtgtcc	ggaatgccc	acgacccaag	4380
ccccctgcaa	gtggctgtga	agacgtgcc	tgaagtgtgc	tctgaacagg	acgaactgga	4440
tttctcatg	gaagccctga	tcatcagcaa	attcaaccac	cagaacattg	ttcgtctgat	4500
tggggtgagc	ctgcaatccc	tgccccgggt	catcctgctg	gagctcatgg	cgggggggaga	4560
cctcaagtcc	ttcctccgag	agacccgccc	tcgcccagac	cagccctcct	ccctggccat	4620
gctggacctt	ctgcacgtgg	ctcgggacat	tgctgtggc	tgtcagtatt	tggaggaaaa	4680
ccacttcac	caccgagaca	ttgtgtccag	aaactgcctc	ttgacctgtc	caggccctgg	4740
aagagtggcc	aagattggag	acttcgggat	ggcccagac	atctacaggg	cgagctacta	4800
tagaaagggg	ggctgtgcca	tgtgtccagt	taagtggatg	ccccagagg	ccttcatgga	4860
aggaatattc	acttctaaaa	cagacacatg	gtcctttgga	gtgctgctat	gggaaatctt	4920
ttctcttgg	tatatgccat	accccagcaa	aagcaaccag	gaagttctgg	agtttctcac	4980
cagtggaggc	cggatggacc	cacccaagaa	ctgccctggg	cctgtatacc	ggataatgac	5040
tcagtgtctg	caacatcagc	ctgaagacag	gcccactttt	gccatcattt	tggagaggat	5100
tgaatactgc	acccaggacc	cggatgtaat	caacaccgct	ttgccgatag	aatatggctc	5160
acttgtggaa	gaggaagaga	aagtgcctgt	gaggcccaag	gacctgagg	gggttctctc	5220
tctcctggtc	tctcaacagg	caaaacggga	ggaggagcgc	agcccagctg	ccccaccacc	5280
tctgcctacc	acctcctctg	gcaaggctgc	aaagaaacc	acagctgcag	agatctctgt	5340
tcgagtcct	agagggcccg	ccgtggaagg	gggacacgtg	aatatggcat	tctctcagtc	5400
caacctcct	tcggagtgtc	acaaggtcca	cggatccaga	aacaagccc	ccagcttgtg	5460
gaacccaacg	tacggctcct	ggtttacaga	gaaacccacc	aaaaagaata	atcctatagc	5520
aaagaaggag	ccacacgaca	ggggtaacct	gggctggag	ggaagctgta	ctgtcccacc	5580
taacgttgca	actgggagac	ttccgggggc	ctcactgctc	ctagagccct	cttcgctgac	5640
tgccaatatg	aaggaggtac	ctctgttcag	gctacgtcac	ttcccttgtg	ggaatgtcaa	5700
ttacggctac	cagcaacagg	gcttgcctt	agaagccgct	actgcccctg	gagctggtca	5760
ttacgaggat	accattctga	aaagcaagaa	tagcatgaac	cagcctgggc	cctgagctcg	5820
gtcgcacact	cacttctctt	ccttgggatc	cctaagaccg	tggaggagag	agaggcaatg	5880
gctccttcac	aaaccagaga	ccaaatgtca	cgttttgttt	tgtgccaacc	tattttgaag	5940
taccacaaaa	aaagctgtat	tttgaataat	ctttagaaag	gttttgagca	tgggttcatc	6000

-continued

ctattctttc	gaaagaagaa	aatatcataa	aatgagtgga	taaatacaag	gccagatgt	6060
ggttgcataa	ggtttttatg	catgtttggt	gtatacttcc	ttatgcttct	ttcaaattgt	6120
gtgtgctctg	cttcaatgta	gtcagaatta	gctgcttcta	tgtttcatag	ttgggggtcat	6180
agatggttcc	ttgccttggt	gatgtggaca	tgagccattt	gaggggagag	ggaacggaaa	6240
taaaggaggt	atgtgtaatg	actaaaa				6267

SEQ ID NO: 2                   moltype = DNA   length = 7368  
 FEATURE                    Location/Qualifiers  
 source                     1..7368  
                           mol\_type = genomic DNA  
                           organism = Homo sapiens

SEQUENCE: 2

caagctttca	agcattcaaa	ggtctaaatg	aaaaaggcta	agtattattt	caaaaggcaa	60
gtatatccta	atatagcaaa	acaaacaaaag	caaaatccat	cagctactcc	tccaattgaa	120
gtgatgaagc	ccaaataatt	catatagcaa	aatggagaaa	attagaccgg	ccatctaaaa	180
atctgccatt	ggtgaagtga	tgaagaacat	ttactgtctt	attccgaagc	ttgtcaattt	240
tgcaactctt	ggctgcctat	ggatttctgt	ggtgcagtgt	acagttttaa	atagctgcct	300
aaagtcgtgt	gtaactaatc	tgggccagca	gcttgacctt	ggcacaccac	ataatctgag	360
tgaaccgtgt	atccaaggat	gtcacttttg	gaactctgta	gatcagaaaa	actgtgcttt	420
aaagtgtcgg	gagtcgtgtg	aggttggctg	tagcagcggc	gaaggtgcat	atgaagagga	480
agtactggaa	aatgcagacc	taccaactgc	tccctttgct	tcttccattg	gaagccacaa	540
tatgacatta	cgatggaaat	ctgcaaactt	ctctggagta	aaatacatca	ttcagtggaa	600
atatgcacaa	cttctgggaa	gctggactta	tactaagact	gtgtccagac	cgctcctatgt	660
ggtcaagccc	ctgcaccctt	tactgagta	cattttccga	gtggtttggg	tcttcacagc	720
gcagctgcag	ctctactccc	ctccaagtcc	cagttacagg	actcatcctc	atggagttcc	780
tgaactgca	cctttgatta	ggaatattga	gagctcaagt	cccgaactg	tgggaagtcag	840
ctgggatcca	cctcaattcc	caggtggacc	tattttgggt	tataacttaa	ggctgatcag	900
caaaaatcaa	aaattagatg	cagggacaca	gagaaccagt	ttccagtttt	actccacttt	960
accaaatact	atctacaggt	tttctattgc	agcagtaaat	gaagtgggtg	agggtccaga	1020
agcagaatct	agtattacca	cttcatcttc	agcagttcaa	caagaggaac	agtggctctt	1080
ttatccaga	aaaacttctc	taagaaagag	atctttaaaa	catttagtag	atgaagcaca	1140
ttgcttcgg	ttggatgcta	tataccataa	tattacagga	atatctgttg	atgtccacca	1200
gcaaatggtt	tatttctctg	aaggaactct	catatgggcg	agaaggctg	ccaacatgct	1260
tgatgtatct	gactgagaa	ttttttacag	aggttcagga	ttaatttctt	ctatctccat	1320
agattggctt	tatcaaagaa	tgtatttcat	catggatgaa	ctggatgtg	tctgtgattt	1380
agagaactgc	tcaaacatcg	aggaaattac	tccaccctct	attagtgcac	ctcaaaaaat	1440
tgtggctgat	tcatacaatg	ggtatgtctt	ttacctctg	agagatggca	tttatagagc	1500
agaccttctt	gtaccatctg	gccgggtgtg	agaagctgtg	cgtattgtgg	agagttgcac	1560
gttaaaggac	tttgcaatca	agccacaagc	caagcgaatc	atttacttca	atgacactgc	1620
ccaagtcttc	atgtcaacat	ttctggatgg	ctctgcttcc	catctcatcc	tacctcgcat	1680
cccctttgct	gatgtgaaaa	gttttgcttg	tgaaaacaat	gactttcttg	tcacagatgg	1740
caaggtcatt	ttccaacagg	atgctttgtc	ttttaatgaa	ttcatcgtgg	gatgtgacct	1800
gagtcacata	gaagaatttg	ggtttggtaa	cttggctatc	tttggctcat	cctcccagct	1860
gcacctctg	ccaggccgct	cgcaggagct	ttcggctctg	tttggctctc	accaggctct	1920
tgttcaatgg	aagcctcctg	cccttgccat	aggagccaat	gtcatcctga	tcagtgatat	1980
tattgaactc	tttgaattag	gcccttctgc	ctggcagaac	tggacctatg	agggtgaaagt	2040
atccaccocaa	gacctcctg	aagtcactca	tattttcttg	aacataagtg	gaacctatgct	2100
gaatgtacct	gagctgcaga	gtgctatgaa	atacaagggt	tctgtgagag	caagttctcc	2160
aaagaggcca	ggccctggtt	cagagccctc	agtgggtact	accctggtgc	cagctagtga	2220
accaccattt	atcatggctg	tgaagaaga	tgggctttgg	agtaaaccat	taaatagctt	2280
tggccagga	gagttcttat	cctctgatat	aggaaatgtg	tcagacatgg	attggtataa	2340
caacagcctc	tactacagtg	acacgaaagg	cgagtttttt	gtgtggctgc	tgaatgggac	2400
ggatatctca	gagaattatc	acctaccag	catgacagga	gcaggggctt	tagcttttga	2460
gtggctgggt	cactttctct	actgggctgg	aaagacatat	gtgatacaaa	ggcagtcctg	2520
ggtgacggga	cacacagaca	ttgttaccca	cgtgaagcta	ttggtgaatg	acatgggtgg	2580
ggattcagtt	ggtggatata	tctactggac	cacactctat	tcagtggaaa	gcaccagact	2640
aaatggggaa	agttcccttg	tactacagac	acagccttgg	ttttctggga	aaaaggtaat	2700
tgtctaaact	ttagacctca	gtgatgggct	cctgtattgg	ttggttcaag	acagtcaatg	2760
tattcacctg	tacacagctg	ttcttcgggg	acagagcact	ggggatacca	ccatcacaga	2820
atgtgcagcc	tggagtactt	ctgaaatttc	ccagaatgca	ctgatgtact	atagtggctg	2880
gtgttctgg	atcaattgct	ttaggattat	cacaactcaa	gaaataggtc	agaaaaccag	2940
tgtctctggt	ttggaaccag	ccagatttaa	tcagttcaca	attattcaga	catcccttaa	3000
gccctgcca	gggaactttt	cctttacccc	taaggttatt	ccagattctg	ttcaagagtc	3060
ttcatttagg	attgaaggaa	atgcttcaag	ttttcaaatc	ctgtggaatg	gtccccctgc	3120
ggtagactgg	ggtgtagttt	tctacagtgt	agaatttagt	gctcattcta	agttcttggc	3180
tagtgaacaa	cactctttac	ctgtattttac	tgtggaagga	ctggaacctt	atgctttatt	3240
taatctttct	gtcactcctt	atacctactg	gggaaagggc	cccaaaacat	ctctgtcact	3300
tcgagcacct	gaaacagttc	catcagcacc	agagaacccc	agaatattta	tattaccaag	3360
tggaaaatgc	tgcaacaaga	atgaagttgt	ggtggaattt	aggtggaaca	aacctaaagca	3420
tgaaaatggg	gtgttaacaa	aatttgaaat	ttctacaat	atatccaatc	aaagtattac	3480
aaacaaaaca	tgtgaagact	ggattgctgt	caatgtcact	ccctcagtga	tgtcttttca	3540
acttgaagge	atgagtccca	gatgctttat	tgcttccag	gttagggcct	ttacatctaa	3600
ggggccagga	ccatagctg	acgttgtaaa	gtctacaaca	tcagaaatca	accatttcc	3660
tcacctcata	actcttcttg	gtaacaagat	agttttttta	gatatggatc	aaaatcaagt	3720
tgtgtggacg	ttttcagcag	aaagagttat	cagtgcctgt	tgctacacag	ctgataatga	3780
gatgggatat	tatgctgaag	gggactcact	ctttctctg	cacttgcaca	atcgtcttag	3840

-continued

```

ctctgagctt ttccaagatt cactgggttt tgatatcaca gttattaca ttgactggat 3900
ttcaaggcac ctctactttg cactgaaaga atcacaaaat ggaatgcaag tatttgatgt 3960
tgatcttgaa cacaaggtga aatatcccag agaggtgaag attcacaata ggaattcaac 4020
aataatctct ttttctgtat atcctctttt aagtcgcttg tattggacag aagtttccaa 4080
ttttggctac cagatgttct actacagtat tatcagtcac accttgcacc gaattctgca 4140
accacagctt acaaaccaac aaaacaaaag gaatcaatgt tcttgtaatg tgactgaatt 4200
tgagttaagt ggagcaatgg ctattgatac ctctaaccta gagaaacct tgatatactt 4260
tgccaaagca caagagatct gggcaatgga tctggaaggc tgtcagtggt ggagagttat 4320
cacagtacct gctatgctcg caggaaaaac ccttgtagc ttaactgtgg atggagatct 4380
tatatactgg atcatcacag caaaggacag cacacagatt tatcaggcaa agaaaggaaa 4440
tggggccatc gtttcccagg tgaaggccct aaggagtagg catatcttgg cttacagttc 4500
agtatgcag ccttttccag ataaagcgtt tctgtctcta gcttcagaca ctgtggaacc 4560
aactatactt aatgccacta acactagcct cacaatcaga ttacctctgg ccaagacaaa 4620
cctcacatgg tatggcatca ccagccctac tccaacatac ctggtttat atgcagaagt 4680
taatgacagg aaaaacagct ctgacttgaa atatagaatt ctggaatttc aggacagtat 4740
agctcttatt gaagatttac aaccattttc aacatacatg atacagatag ctgtaaaaaa 4800
ttattattca gatccttgg aacatttacc accaggaaaa gagatttggg gaaaaactaa 4860
aaatggagta ccagaggcag tgcagctcat taatacaact gtgcggtcag acaccagcct 4920
cattatatct tggagagaaat ctcacaaagg aaatggacct aaagaatcag tccgttatca 4980
gttggaatc tcacacctgg ccctaattcc tgaacctcct ctaagacaaa gtgaatttcc 5040
aaatggaagg ctcaactctc ttgttactag actgtctggt ggaaatattt atgtgttaaa 5100
ggttcttgcc tgccactctg aggaaatgtg gtgtacagag agtcatcctg tcaactgtgga 5160
aatgtttaac acaccagaga aaccttattc cttggttcca gagaacacta gtttgcaatt 5220
taattggaag gctccattga atgttaacct catcagattt tgggttgagc tacagaagtg 5280
gaaatacaat gagttttacc atgttaaaac ttatagcagc caaggtcctg cttatgtctg 5340
taatacaca aatctacaac cttatacttc atataatgtc agagtgtggt tggtttataa 5400
gacgggagaa aatagcacct cacttccaga aagctttaag acaaaagctg gagtcccaa 5460
taaaccaggc attcccaaat tactagaagg gagtaaaaat tcaatacagt gggagaaagc 5520
tgaagataat ggatgtagaa ttacatacta tatccttgag ataagaaaga gcacttcaaa 5580
taatttacag aaccagaatt taaggtggaa gatgacattt aatggatcct gcagtgtgt 5640
ttgcacatgg aagtccaaaa acctgaaagg aatatttcag ttcagagtag tagctgcaaa 5700
taatctaggg tttggtgaat atagtggaaat cagtgagaat attatattag ttggagatga 5760
tttttgata ccagaaacaa gtttcatact tactattata gttggaatat ttctggttgt 5820
tacaatccca ctgacctttg tctggcatag aagataaag aatcaaaaaa gtgccaagga 5880
aggggtgaca gtgcttataa acgaagacaa agagtggct gagctgcgag gtctggcagc 5940
cggagtaggc ctggctaatt cctgctatgc aatcatact cttccaacc aagaggagat 6000
tgaaaatctt cctgccttcc ctcggaaaa actgactctg cgtctcttgc tgggaagtgg 6060
agcctttgga gaagtgtatg aaggaacagc agtggacatc ttaggagttg gaagtggaga 6120
aatcaaagta gcagtgaaga ctttgaagaa gggttccaca gaccaggaga agattgaatt 6180
cctgaaggag gcacatctga tgagcaaat taatcatccc aacattctga agcagcttgg 6240
agtttgtctg ctgaatgaac cccaatacat tatcctggaa ctgatggagg gaggagacct 6300
tcttacttat ttgcgtaaaag cccggatggc aacgttttat ggtcctttac tcaccttgg 6360
tgaccttcta gacctgtgtg tagatatttc aaaagctgt gtctacttgg aacggatgca 6420
ttcattcac agggatctgg cagctagaaa ttgccttgg tccgtgaaag actataccag 6480
tccacggata gtgaagattg gagactttgg actcgccaga gacatctata aaaatgatta 6540
ctatagaaag agaggggaag gcctgctccc agttcgggtg atggctccag aaagtttgat 6600
ggatggaatc ttcactactc aatctgatgt atggtctttt ggaattctga tttgggagat 6660
ttaactctt ggtcatcagc cttatccagc tcattccaac cttgatgtgt taaactatgt 6720
gcaaacagga gggagactgg agccaccaag aaattgtcct gatgatctgt ggaatttaat 6780
gaccagtgct tgggctcaag aacccgacca aagacctact tttcatagaa ttcaggacca 6840
acttcagtta ttcagaaatt ttttcttaaa tagcatttat aagtcagag atgaagcaaa 6900
caacagtgga gtcataaatg aaagctttga aggtgaagat ggcgatgtga tttgtttgaa 6960
ttcagatgac attatgccag ttgctttaat ggaaacgaaag aaccgagaag ggttaacta 7020
tatggtactt gctacagaat gtggccaagg tgaagaaaag tctgagggtc ctctaggctc 7080
ccaggaatct gaatcttgtg gtctgaggaa agaagagaag gaaccacatg cagacaaaga 7140
tttctgcaa gaaaaacaag tggcttactg ccctctggc aagcctgaag gcctgaacta 7200
tgctgtctc actcacagtg gatatggaga tgggtctgat taatagcgtt gtttgggaaa 7260
tagagagttg agataaacac tctcattcag tagttactga aagaaaactc tgctagaatg 7320
ataaatgtca tgggtgtcta taactccaaa taacaatgc aacgttcc 7368

```

```

SEQ ID NO: 3          moltype = DNA length = 4174
FEATURE              Location/Qualifiers
source                1..4174
                     mol_type = genomic DNA
                     organism = Homo sapiens

```

```

SEQUENCE: 3
agtcccgcga ccgaagcagg gcgcgcagca gcgctgagtg ccccggaacg tgcgtcgcgc 60
cccagtgctc cgtcgcgtcc gccgcgcccc gggcggggat gggcgggcca gactgagcgc 120
cgcaccgcgc atccagaccg gccggcccta gccgcagtc ctccagccgt ggcccagcgc 180
cgcacggggc atggcgaaag cgacgtccgg tgcgcgggg ctgcgtctgc tgttgctgct 240
gctgctgccg ctgctaggca aagtggcatt gggcctctac ttctcgagg atgcttactg 300
ggagaagctg tatgtggacc aggcggcccg cacgccctt ctgtaectcc atgccctgcg 360
ggacgcccc ctgagaggtgc ccagcttccg cctgggccc catctctacg gcacgtaccg 420
cacacggctg catgagaaca actggatctg catccaggag gacaccggcc tcctctacct 480
taaccggagc ctggaccata gctcctggga gaagctcagt gtccgcaacc gcggctttcc 540
cctgctcacc gtctacctca aggtcttctc gtcaccaca tcccttctg agggcgagtg 600

```

-continued

```

ccagtggcca ggctgtgcc gcgataactt ctcttttctt aacacctcct ttccagcctg 660
cagctccctc aagccccggg agctctgctt cccagagaca aggcctcctc tccgcattcg 720
ggagaaccga cccccaggca ccttccacca gttccgcctg ctgcctgtgc agttcttgtg 780
ccccaacatc agcgtggcct acaggctcct ggagggtgag ggtctgccct tccgctgcgc 840
cccgacagc ctggaggtga gcacgcctg ggccctggac cgcgagcagc gggagaagta 900
cgagctggtg gccgtgtgca ccgtgcacgc cggcgcgcgc gaggaggtgg tgatggtgcc 960
cttcccggtg accgtgtacg acgaggacga ctccggcgc accctccccg cgggcgtcga 1020
caccgccagc gccgtggtgg agtccaagcg gaaggaggac accgtggtgg ccacgctgcg 1080
tgtcttcgat gcagacgtgg tacctgcctc aggggagctg gtgaggcgtt acacaagcac 1140
gctgctcccc ggggacacct gggcccagca gaccttccgg gtggaacct ggccccacga 1200
gacctcggtc caggccaacg gcagcttctg gcggcgacc gtacatgact ataggctggt 1260
tctcaaccgg aacctctcca tctcggagaa ccgaccatg cagctggcgg tgctggtcaa 1320
tgactcagac ttccagggcc caggagcggg cgtcctcttg ctccacttca acgtgtcgg 1380
gctgccggtc agcctgcacc tgcccagtac ctactccctc tccgtgagca ggagggtcgc 1440
ccgatttgcc cagatcggga aagtctgtgt ggaaaactgc caggcattca gtggcatcaa 1500
cgtccagtac aagctgcatt cctctggtgc caactgcagc acgctagggg tggtcacctc 1560
agccgaggac acctcgggga tccgtttgtg gaatgacacc aaggccctgc ggcggcccaa 1620
gtgtgccgaa cttcactaca tgggtggtgg caccgaccag cagacctta ggcaggccca 1680
ggcccagctg cttgtaacag tggaggggtc atatgtggcc gaggaggcgg gctgccccct 1740
gtcctgtgca gtcagcaaga gacggctgga gtgtgaggag tgtggcggcc tgggctcccc 1800
aacaggcagg tgtgagtgga ggcaaggaga tggcaaagg atcaccagga acttctccac 1860
ctgctctccc agcaccaaga cctgccccga cggccactgc gatgttgtgg agaccaaga 1920
catcaacatt tgccctcagg actgcctccg gggcagcatt gttgggggac acgagcctgg 1980
ggagccccgg gggattaaag ctggctatgg cacctgcaac tgcttccctg aggaggagaa 2040
gtgcttctgc gagcccgaag acatccagga tccactgtgc gacgagctgt gccgcacggt 2100
gatcgcagcc cctgtcctct tctccttcat cgtctcggtg ctgctgtctg cctctctgcat 2160
ccactgctac cacaagtttg cccacaagcc acctatctcc tcagctgaga tgaccttccg 2220
gaggcccgcc caggccttcc cggtcagcta ctctcttcc ggtgcccgcc ggccctcgtc 2280
ggactccatg gagaaccagg tctccgtgga tgcttcaag atcctggagg atccaaagtg 2340
ggaattccct cggagaact tggttcttgg aaaaactcta ggagaaggcg aatttggaaa 2400
agtggtaaac gcaacggcct tccatctgaa aggcagagca gggtaacca cgggtggcct 2460
gaagatgctg aaagagaacg cctccccgag tgagcttcca gacctgctgt cagagttcaa 2520
cgtcctgaag caggtcaacc acccacatgt catcaaattg tatggggcct gcagccagga 2580
tggcccgtc ctccctcctc tggagtacgc caaatcggc tccctgcggg gcttctcctc 2640
cgagagccgc aaagtgggyc ctggctacct ggcagctgga ggcagccgca actccagctc 2700
cctggaccac cggatgagc gggccctcac catgggagc ctcatctcat ttgctggca 2760
gatctcacag gggatgcagt atctggccga gatgaagctc gttcatcggg acttggcagc 2820
cagaaacatc ctggtagctg aggggcccga gatgaagatt tcggatttcg gcttgtcccg 2880
agatgtttat gaagaggatt cctacgtgaa gaggagccag ggtcggattc cagttaaatg 2940
gatggcaatt gaatccctt ttgatcatat ctacaccacg caaagtgatg tatggtcttt 3000
tggtgtcctg ctgtgggaga tctgaccctc agggggaaac ccctatcctg ggattcctcc 3060
tgagcggctc ttcaacctc tgaagaccgg ccaccggatg gagaggccag acaactgcag 3120
cgaggagatg taccgctga tctgcaatg ctggaagcag gagccggaca aaaggccggt 3180
gtttgcccag atcagcaaag acctggagaa gatgatggtt aagaggagag actacttgg 3240
ccttgcggcg tccactccat ctgactccct gatattatgac gacggcctct cagaggagga 3300
gacaccgctg gtggactgta ataatgcccc cctccctcga gccctccctt ccacatggat 3360
tgaaaacaaa ctctatggtg gaatttccca tgcatctact agattctagc accgctgtcc 3420
cctctgcact atccttctc tctgtgatgc ttttataaaa tgtttctggt ctgaacaaaa 3480
ccaaagtctg ctctgaacct ttttatttgt aaatgtctga ctttgcctcc agtttacatt 3540
taggcattat tgcaactatg tttttctaaa aggaagtga aataagtgtg attaccacat 3600
tgcccagcaa cttaggatgg tagaggaaaa aacagatcag ggcggaactc tcaggggaga 3660
ccaagaacag gttgataag gcgcttctgg ggtgggaatc aagtcatagt acttctactt 3720
taactaagtg gataaataa caaatctggg gaggtattca gttgagaaag gagccaccag 3780
caccactcag cctgcaactg gagcacagcc aggttcccc agaccctcc tgggcaggca 3840
ggtgcctctc agaggccacc cggcactggc gagcagccac tggccaagcc tcagccccag 3900
tcccagccac atgtcctcca tcaggggtag cgaggttgc ggagctggct ggccctggga 3960
ggacgcaccc cactgctgt tttcacatcc tttcccttac ccacctcag gacggttgc 4020
acttatgaag tcagtgtctaa agctggagca gttgcttttt gaaagaacat ggtctgtggt 4080
gctgtggtct tacaatggac agtaaatatg gttcttgcca aaactccttc ttttgtcttt 4140
gattaatac tagaaattta aaaaaaaaaa aaaa 4174

```

```

SEQ ID NO: 4          moltype = DNA length = 5629
FEATURE              Location/Qualifiers
source                1..5629
                    mol_type = genomic DNA
                    organism = Homo sapiens

```

```

SEQUENCE: 4
agtcccgcga ccgaagcagg ggcgcgagca gcgctgagtg ccccggaacg tgcgtcgcgc 60
ccccagtgtc cgtcgcgtcc gccgcgcccc gggcggggat ggggcggcca gactgagcgc 120
cgcacccgcc atccagaccg gccggcccta gccgcagctc ctccagccgt ggccccagcg 180
cgcacggggc atggcgaagg cgacgtccgg tgccgcgggg ctgcgtctgc tgttgcgtgc 240
gctgctgccg ctgctaggca aagtggcatt gggcctctac ttctcagggg atgcttactg 300
ggagaagctg tatgtggacc aggcggcccg cacgccttg ctgtacgtcc atgccctgcg 360
ggacgcccc caggaggtgc ccagcttccg cctgggccag catctctacg gcacgtaccg 420
cacacggctg catgagaaca actggatctg catccaggag gacaccggcc tctctacct 480
taaccggagc ctggaccata gctcctggga gaagctcagt gtccgcaacc gcggctttcc 540

```

-continued

cctgctcacc	gtctacctca	aggttcttct	gtcaccacaca	tcccttctgt	agggcgagt	600
ccagtggcca	ggctgtgccc	gcgtatactt	ctccttcttc	aacacctcct	ttccagcctg	660
cagctccctc	aagccccggg	agctctgctt	cccagagaca	aggccctcct	tccgcattcg	720
ggagaaccga	cccccaggca	ccttccacca	gttccgcctg	ctgcctgtgc	agttcttgtg	780
ccccaacatc	agcgtggcct	acaggctcct	ggaggggtgag	ggctctgccct	tccgctgcgc	840
cccggacagc	ctggaggtga	gcacgcgctg	ggcctggac	cgcgagcagc	gggagaagta	900
cgagctgggtg	gccgtgtgca	ccgtgcacgc	cggcgcgcg	gaggaggtgg	tgatgggtgcc	960
cttcccgtgc	accgtgtacg	acgaggacga	ctcactgcgcc	accttccccg	cgggcgtcga	1020
caccgccagc	gccgtgggtg	agttcaagcg	gaaggaggac	accgtgggtg	ccacgctgcg	1080
tgtcttcgat	gcagacgtgg	tacctgcctc	aggggagctg	gtgaggcggg	acacaagcac	1140
gctgctcccc	ggggacacct	gggcccagca	gaccttccgg	gtggaacct	ggccccacga	1200
gacctcgggtc	caggccaacg	gcagcttctg	gcggcgacc	gtacatgact	ataggctggg	1260
tctcaaccgg	aacctctcca	tctcggagaa	ccgaccatg	cagctggcgg	tgctgggtcaa	1320
tgactcagac	ttccagggcc	caggagcggg	cgtcctcttg	ctccacttca	acgtgtcggg	1380
gctgccggtc	agcctgcacc	tgcccagtac	ctactccctc	tccgtgagca	ggagggctcg	1440
ccgatttgcc	cagatcggga	aagtctgtgt	ggaaaactgc	caggcattca	gtggcatcaa	1500
cgctccagtac	aagctgcatt	cctctgggtg	caactgcagc	acgctagggg	tggtcacctc	1560
agccgaggac	acctcgggga	tcctgtttgt	gaatgacacc	aaggccctgc	ggcggcccaa	1620
gtgtgccgaa	cttactaca	tggtgggtgg	caccgaccag	cagacctcta	ggcaggccca	1680
ggcccagctg	cttgtaacag	tggaggggtc	atatgtggcc	gaggaggcgg	gctgccccct	1740
gtcctgtgca	gtcagcaaga	gacggctgga	gtgtgaggag	tgtggcggcc	tggtctcccc	1800
aacaggcagg	tgtgagtggg	ggcaaggaga	tggcaaaggg	atcaccagga	acttctccac	1860
ctgctctccc	agcaccaaga	cctgccccga	cggccactgc	gatgttgtgg	agacccaaga	1920
catcaacatt	tgccctcagg	actgcctccg	ggcagcatt	gttgggggac	acgagcctgg	1980
ggagccccgg	gggattaaag	ctggctatgg	cactgcacac	tgcttccctg	aggaggagaa	2040
gtgcttctgc	gagcccgaag	acatccagga	tccactgtgc	gacgagctgt	gccgcacggt	2100
gatcgcagcc	gctgtcctct	tctccttcat	cgtctcgggtg	ctgctgtctg	ccttctgcat	2160
ccactgctac	cacaagtgtg	cccacaagcc	acccatctcc	tcaactgaga	tgaccttccg	2220
gagcccgcgc	caggccttcc	cggctcagct	ctcctcttcc	gggtcccggc	ggcctctgct	2280
ggactccatg	gagaaccagg	tctccgtgga	tgcttcaag	atcctggagg	atccaaagtg	2340
ggaattccct	cggagaact	tggttcttgg	aaaaactcta	ggagaaggcg	aatttggaaa	2400
agtgtgcaag	gcaacggcct	tccatctgaa	aggcagagca	gggtacacca	cggtggcctg	2460
gaagatgctg	aaagagaacg	cctccccgag	tgagcttcca	gacctgctgt	cagagttcaa	2520
gctcctgaag	caggtcaacc	acccacatgt	catcaaattg	tatggggcct	gcagccagga	2580
tgcccccgtc	ctcctcatcg	tggagtacgc	caaatacggc	tccctgcggg	gcttccctccg	2640
cgagagccgc	aaagtggggc	ctggctacct	gggcagtgga	ggcagccgca	actccagctc	2700
cctggaccac	ccggatgagc	gggcccctc	catggggcag	ctcatctcat	ttgctggca	2760
gatctcacag	gggatgcagt	atctggccga	gatgaagctc	gttcatcggg	acttggcagc	2820
cagaaacatc	ctggtagctg	aggggcccga	gatgaagatt	tccgatttccg	gcttgtccccg	2880
agatgtttat	gaagaggatt	cctacgtgaa	gaggagccag	ggctcggattc	cagttaaatg	2940
gatggcaatt	gaatcccttt	ttgatcatat	ctacaccacg	caaagtgatg	tatgggtcttt	3000
tggtgtcctg	ctgtgggaga	tgtgacctc	agggggaaac	ccctatcctg	ggattcctcc	3060
tgagcggctc	ttcaaccttc	tgaagaccgg	ccaccggatg	gagaggccag	acaactgcag	3120
cgaggagatg	taccgcctga	tgctgcaatg	ctggaagcag	gagccggaca	aaaggccggg	3180
gtttgcggac	atcagcaaa	acctggagaa	gatgatgggt	aagaggagag	actacttggg	3240
ccttgcggcg	tccactccat	ctgactccct	gatttatgac	gacggcctct	cagaggagga	3300
gacaccgctg	gtggactgta	ataatgcccc	cctcctcoga	gccctccctt	ccacatggat	3360
tgaaaaacaaa	ctctatggca	tgtcagaccc	gaaactggcct	ggagagagtc	ctgtaccact	3420
cacgagagct	gatggcacta	acactgggtt	tccaagatat	ccaaatgata	gtgtatatgc	3480
taactggatg	ctttcaccct	cagcggcaca	attaatggac	acgtttgata	gttaacattt	3540
ctttgtgaaa	ggtaatggac	tcacaagggg	aagaaacatg	ctgagaatgg	aaagtctacc	3600
ggccctttct	ttgtgaacgt	cacattggcc	gagccgtgtt	cagttcccag	gtggcagact	3660
ggtttttggg	agtttgggtt	aaactccaag	gtgtttttac	ttctgatagc	cgggtgatttt	3720
ccctcctagc	agacatgcca	caccgggtaa	gagctctgag	tcttagtggg	taagcattcc	3780
tttctcttca	gtgcccagca	gcacccagtg	ttggctctgtg	tccatcagtg	accaccaaca	3840
ttctgtgttc	acatgtgtgg	gtccaacact	tactacctgg	tgtatgaaat	tggaacctgaa	3900
ctgttggtat	tttctagtgt	ccgccaacaa	aggcaaaaaa	atttaaacat	gaagcacaca	3960
cacaaaaaag	gcagtaggaa	aaatgctggc	cctgatgacc	tgctccttatt	cagaatgaga	4020
gactgcgggg	ggggcctggg	ggtagtgtca	atgcccctcc	agggctggag	gggaagaggg	4080
gccccgagga	tgggcctggg	ctcagcattc	gagatcttga	gaatgatgtt	tttttaactca	4140
tgaaccttt	ccttaggaag	acatttgggt	ttcatcatga	ttaagatgat	tccatagattt	4200
agcacaatgg	agagattcca	tgccatcttt	actatgtgga	tggtggatc	aggggaagagg	4260
gctcacaaga	cacatttgtc	ccccgggccc	accacatcat	cctcacgtgt	tccggtactga	4320
gcagccacta	cccctgatga	gaacagtatg	aagaaagggg	gctgttggag	tcccagaatt	4380
gctgacagca	gaggctttgc	tgtgtgtaat	cccactgcc	accagcctgc	agcacacccc	4440
acagccaagt	agaggcgaaa	gcagtggctc	atcctacctg	ttaggagcag	gtagggcttg	4500
tactcacttt	aatttgaatc	ttatcaactt	actcataaag	ggacaggcta	gctagctgtg	4560
ttagaagttag	caatgacaat	gaccaaggac	tgctacacct	ctgattacaa	ttctgatgtg	4620
aaaaagatgg	tggttggctc	ttatagagcc	tgtgtgaaag	gcccattggat	cagctcttcc	4680
tgtgtttgta	atttaagtct	gctacaagat	gtttctgttt	cttagattct	gaccatgact	4740
cataagcttc	ttgtcattct	tcattgcttg	tttgtggtea	cagatgcaca	acactcctcc	4800
agtcttgtgg	gggcagcttt	tgggaagtct	cagcagctct	tctggctgtg	ttgtcagcac	4860
tgtaacttcg	cagaaaagag	tccgattacc	aaaacactgc	ctgctcttca	gacttaaaagc	4920
actgatagga	cttaaaatag	tctcattcaa	atactgtatt	ttatataggc	atttcacaaa	4980
aacagcaaaa	ttgtggcatt	ttgtgaggcc	aaggcttggg	tgctgtgta	atagagcctt	5040
tggtgtgtg	cgcacacacc	cagagggaga	gtttgaaaaa	tgcttattgg	acacgtaacc	5100

-continued

---

```

tggtctaat ttgggtgtt tttcagatac actgtgataa gttcttttac aaatatctat 5160
agacatggta aacttttggg tttcagatat gcttaatgat agtcttacta aatgcagaaa 5220
taagaataaa ctttctcaaa ttattaaaaa tgctacaca gtaagtgtga attgctgcaa 5280
caggtttggt ctcaggaggg taagaactcc aggtctaac agctgaccca gtgatgggga 5340
atztatcctt gaccaattta tccttgacca ataaccta atgtctattcc tgagttataa 5400
aagtcccat ccttattagc tctactggaa ttttcataca cgtaaagca gaagttacta 5460
agtattaagt attactgagt attaagtagt aatctgtcag ttattaaaat ttgtaaaatc 5520
tatttatgaa aggtcattaa accagatcat gttccttttt ttgtaataca ggtgactaag 5580
aaaatcagtt gtgtaataaa aatcatgtat cataaaaaaa aaaaaaaaaa 5629

SEQ ID NO: 5          moltype = DNA length = 42
FEATURE              Location/Qualifiers
misc_feature         1..42
                     note = Description of Artificial Sequence: Synthetic primer
source              1..42
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 5
cctctctatg ggcagtcggt gatgagagag tggcaggtgt gg 42

SEQ ID NO: 6          moltype = DNA length = 44
FEATURE              Location/Qualifiers
misc_feature         1..44
                     note = Description of Artificial Sequence: Synthetic primer
source              1..44
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 6
ggatctcgac gctctccctc agaggtcacc acagagagga tcag 44

SEQ ID NO: 7          moltype = DNA length = 43
FEATURE              Location/Qualifiers
misc_feature         1..43
                     note = Description of Artificial Sequence: Synthetic primer
source              1..43
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 7
cctctctatg ggcagtcggt gatcatggct tgcagctcct ggt 43

SEQ ID NO: 8          moltype = DNA length = 41
FEATURE              Location/Qualifiers
misc_feature         1..41
                     note = Description of Artificial Sequence: Synthetic primer
source              1..41
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 8
ggatctcgac gctctccctg cagctccatc tgcattgctt g 41

SEQ ID NO: 9          moltype = DNA length = 45
FEATURE              Location/Qualifiers
misc_feature         1..45
                     note = Description of Artificial Sequence: Synthetic primer
source              1..45
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 9
cctctctatg ggcagtcggt gatggccttc atacacctcc ccaa 45

SEQ ID NO: 10         moltype = DNA length = 37
FEATURE              Location/Qualifiers
misc_feature         1..37
                     note = Description of Artificial Sequence: Synthetic primer
source              1..37
                     mol_type = other DNA
                     organism = synthetic construct

SEQUENCE: 10
ggatctcgac gctctccctt tgggcattcc ggacacc 37

SEQ ID NO: 11         moltype = DNA length = 49
FEATURE              Location/Qualifiers
misc_feature         1..49
                     note = Description of Artificial Sequence: Synthetic primer
source              1..49
                     mol_type = other DNA

```

-continued

---

```

                organism = synthetic construct
SEQUENCE: 11
cctctctatg ggcagtcggt gataggaaat ccagttcgtc ctgttcaga          49

SEQ ID NO: 12      moltype = DNA length = 43
FEATURE           Location/Qualifiers
misc_feature      1..43
                  note = Description of Artificial Sequence: Synthetic primer
source           1..43
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 12
ggatctcgac gctctccctg atcagggctt ccatgaggaa atc          43

SEQ ID NO: 13      moltype = DNA length = 41
FEATURE           Location/Qualifiers
misc_feature      1..41
                  note = Description of Artificial Sequence: Synthetic primer
source           1..41
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 13
cctctctatg ggcagtcggt gatggctccc caggctcggt t          41

SEQ ID NO: 14      moltype = DNA length = 43
FEATURE           Location/Qualifiers
misc_feature      1..43
                  note = Description of Artificial Sequence: Synthetic primer
source           1..43
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 14
ggatctcgac gctctcccta ggtgccatag ccagctttaa tcc          43

SEQ ID NO: 15      moltype = DNA length = 43
FEATURE           Location/Qualifiers
misc_feature      1..43
                  note = Description of Artificial Sequence: Synthetic primer
source           1..43
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 15
cctctctatg ggcagtcggt gatatcaccg tgcggcacag ctc          43

SEQ ID NO: 16      moltype = DNA length = 39
FEATURE           Location/Qualifiers
misc_feature      1..39
                  note = Description of Artificial Sequence: Synthetic primer
source           1..39
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 16
ggatctcgac gctctccctg aggacagcgg ctgcatca          39

SEQ ID NO: 17      moltype = DNA length = 47
FEATURE           Location/Qualifiers
misc_feature      1..47
                  note = Description of Artificial Sequence: Synthetic primer
source           1..47
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 17
cctctctatg ggcagtcggt gatagaacca agttcttccg agggaat          47

SEQ ID NO: 18      moltype = DNA length = 44
FEATURE           Location/Qualifiers
misc_feature      1..44
                  note = Description of Artificial Sequence: Synthetic primer
source           1..44
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 18
ggatctcgac gctctccctt ccaaattcgc cttctcctag agtt          44

SEQ ID NO: 19      moltype = DNA length = 44
FEATURE           Location/Qualifiers

```

-continued

---

```

misc_feature      1..44
                  note = Description of Artificial Sequence: Synthetic primer
source            1..44
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 19
cctctctatg ggcagtcggt gatacagcag gtctcgcagc tcac                44

SEQ ID NO: 20      moltype = DNA length = 41
FEATURE           Location/Qualifiers
misc_feature      1..41
                  note = Description of Artificial Sequence: Synthetic primer
source            1..41
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 20
ggatctcgac gctctccctt gacctgcttc aggacgttga a                    41

SEQ ID NO: 21      moltype = DNA length = 45
FEATURE           Location/Qualifiers
misc_feature      1..45
                  note = Description of Artificial Sequence: Synthetic primer
source            1..45
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 21
cctctctatg ggcagtcggt gatcttgctg actgcacagg acagg                45

SEQ ID NO: 22      moltype = DNA length = 40
FEATURE           Location/Qualifiers
misc_feature      1..40
                  note = Description of Artificial Sequence: Synthetic primer
source            1..40
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 22
ggatctcgac gctctccctt cctcacactc cagccgtctc                    40

SEQ ID NO: 23      moltype = DNA length = 42
FEATURE           Location/Qualifiers
misc_feature      1..42
                  note = Description of Artificial Sequence: Synthetic primer
source            1..42
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 23
cctctctatg ggcagtcggt gattggtgct gggagagcag gt                    42

SEQ ID NO: 24      moltype = DNA length = 37
FEATURE           Location/Qualifiers
misc_feature      1..37
                  note = Description of Artificial Sequence: Synthetic primer
source            1..37
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 24
ggatctcgac gctctccctc cgtcggggca ggtcttg                        37

SEQ ID NO: 25      moltype = DNA length = 46
FEATURE           Location/Qualifiers
misc_feature      1..46
                  note = Description of Artificial Sequence: Synthetic primer
source            1..46
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 25
cctctctatg ggcagtcggt gatggctgca tgaagtttta acatgg                46

SEQ ID NO: 26      moltype = DNA length = 48
FEATURE           Location/Qualifiers
misc_feature      1..48
                  note = Description of Artificial Sequence: Synthetic primer
source            1..48
                  mol_type = other DNA
                  organism = synthetic construct

SEQUENCE: 26

```

-continued

---

```

ggatctcgac gctctcctt gatattacag acataagcag gaccttgg          48

SEQ ID NO: 27          moltype = DNA length = 48
FEATURE              Location/Qualifiers
misc_feature          1..48
                      note = Description of Artificial Sequence: Synthetic primer
source               1..48
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 27
cctctctatg ggcagtcggt gatctagtaa tttgggaatg cctggttt          48

SEQ ID NO: 28          moltype = DNA length = 44
FEATURE              Location/Qualifiers
misc_feature          1..44
                      note = Description of Artificial Sequence: Synthetic primer
source               1..44
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 28
ggatctcgac gctctcctt tcagctttct cccactgtat tgaatg          44

SEQ ID NO: 29          moltype = DNA length = 52
FEATURE              Location/Qualifiers
misc_feature          1..52
                      note = Description of Artificial Sequence: Synthetic primer
source               1..52
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 29
cctctctatg ggcagtcggt gatcatcttc caccttaaat tctggttctg ta          52

SEQ ID NO: 30          moltype = DNA length = 44
FEATURE              Location/Qualifiers
misc_feature          1..44
                      note = Description of Artificial Sequence: Synthetic primer
source               1..44
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 30
ggatctcgac gctctcctc aggatccatt aaatgtcatc ttcc          44

SEQ ID NO: 31          moltype = DNA length = 52
FEATURE              Location/Qualifiers
misc_feature          1..52
                      note = Description of Artificial Sequence: Synthetic primer
source               1..52
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 31
cctctctatg ggcagtcggt gatagtaagt atgaaacttg tttctggtat cc          52

SEQ ID NO: 32          moltype = DNA length = 43
FEATURE              Location/Qualifiers
misc_feature          1..43
                      note = Description of Artificial Sequence: Synthetic primer
source               1..43
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 32
ggatctcgac gctctcctg gtcagtgga ttgtaacaac cag          43

SEQ ID NO: 33          moltype = DNA length = 43
FEATURE              Location/Qualifiers
misc_feature          1..43
                      note = Description of Artificial Sequence: Synthetic primer
source               1..43
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 33
cctctctatg ggcagtcggt gatcaccctt tccttggcac ttt          43

SEQ ID NO: 34          moltype = DNA length = 46
FEATURE              Location/Qualifiers
misc_feature          1..46
                      note = Description of Artificial Sequence: Synthetic primer

```

-continued

---

```

source                1..46
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 34
ggatctcgac gctctcctt ctttgtcttc gttataaagc actgtc          46

SEQ ID NO: 35         moltype = DNA length = 45
FEATURE              Location/Qualifiers
misc_feature         1..45
                      note = Description of Artificial Sequence: Synthetic primer
source              1..45
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 35
cctctctatg ggcagtcggt gatttcaatc tcctcttggg ttgga          45

SEQ ID NO: 36         moltype = DNA length = 40
FEATURE              Location/Qualifiers
misc_feature         1..40
                      note = Description of Artificial Sequence: Synthetic primer
source              1..40
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 36
ggatctcgac gctctccctc cgaggggaagg caggaagatt          40

SEQ ID NO: 37         moltype = DNA length = 47
FEATURE              Location/Qualifiers
misc_feature         1..47
                      note = Description of Artificial Sequence: Synthetic primer
source              1..47
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 37
cctctctatg ggcagtcggt gatcaggaat tcaatcttct cctggtc          47

SEQ ID NO: 38         moltype = DNA length = 42
FEATURE              Location/Qualifiers
misc_feature         1..42
                      note = Description of Artificial Sequence: Synthetic primer
source              1..42
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 38
ggatctcgac gctctccctc tcatcagatg tgctccttc ag          42

SEQ ID NO: 39         moltype = DNA length = 103
FEATURE              Location/Qualifiers
misc_feature         1..103
                      note = Description of Artificial Sequence: Synthetic
                      polynucleotide
source              1..103
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 39
aggatgtccc tgtcaaggct cctgagacct ttgataacat aaccattagc agagaggctc  60
aggctggagt cccaataaa ccaggcattc ccaattact aga          103

SEQ ID NO: 40         moltype = DNA length = 88
FEATURE              Location/Qualifiers
misc_feature         1..88
                      note = Description of Artificial Sequence: Synthetic
                      oligonucleotide
source              1..88
                      mol_type = other DNA
                      organism = synthetic construct

SEQUENCE: 40
tcaaggctct gagaccttg ataacataac cattagcaga gaggctcagg ctggagtcca  60
aatcaaacag gcattcccaa atactaga          88

SEQ ID NO: 41         moltype = DNA length = 81
FEATURE              Location/Qualifiers
misc_feature         1..81
                      note = Description of Artificial Sequence: Synthetic
                      oligonucleotide
source              1..81

```

-continued

---

```

mol_type = other DNA
organism = synthetic construct
SEQUENCE: 41
gagacctttg ataacataac cattagcaga gaggctcagg ctggagtccc aaataaacca 60
ggcattccca aaattactag a 81

SEQ ID NO: 42      moltype = DNA length = 96
FEATURE          Location/Qualifiers
misc_feature     1..96
                 note = Description of Artificial Sequence: Synthetic
                 oligonucleotide
source          1..96
                 mol_type = other DNA
                 organism = synthetic construct
SEQUENCE: 42
gtcctgtcaa ggctctgaga cctttgataa cataaccatt agcagagagg ctcaggctgg 60
agtcccaaat aaaccaggca ttcccaaatt actaga 96

SEQ ID NO: 43      moltype = DNA length = 103
FEATURE          Location/Qualifiers
misc_feature     1..103
                 note = Description of Artificial Sequence: Synthetic
                 polynucleotide
source          1..103
                 mol_type = other DNA
                 organism = synthetic construct
SEQUENCE: 43
gcaggatgtc cctgtcaagg ctctgagacc ttgataaca taaccattag cagagaggct 60
caggatgatt tttgatacc agaaacaagt ttcatactta cta 103

SEQ ID NO: 44      moltype = DNA length = 115
FEATURE          Location/Qualifiers
misc_feature     1..115
                 note = Description of Artificial Sequence: Synthetic
                 polynucleotide
source          1..115
                 mol_type = other DNA
                 organism = synthetic construct
SEQUENCE: 44
ggcgcaggag gggcaggatg tccctgtcaa ggctctgaga cctttgataa cataaccatt 60
agcagagagg ctcaggatga tttttggata ccagaaacaa gtttcatact tacta 115

SEQ ID NO: 45      moltype = DNA length = 46
FEATURE          Location/Qualifiers
misc_feature     1..46
                 note = Description of Artificial Sequence: Synthetic
                 oligonucleotide
source          1..46
                 mol_type = other DNA
                 organism = synthetic construct
SEQUENCE: 45
ccatctcatc cctgcgtgtc tccgactcag ctaaggtaac tggcgt 46

SEQ ID NO: 46      moltype = DNA length = 20
FEATURE          Location/Qualifiers
misc_feature     1..20
                 note = Description of Artificial Sequence: Synthetic
                 oligonucleotide
source          1..20
                 mol_type = other DNA
                 organism = synthetic construct
SEQUENCE: 46
cgccaggttc cagtcacgac 20

SEQ ID NO: 47      moltype = DNA length = 70
FEATURE          Location/Qualifiers
misc_feature     1..70
                 note = Description of Artificial Sequence: Synthetic
                 oligonucleotide
source          1..70
                 mol_type = other DNA
                 organism = synthetic construct
SEQUENCE: 47
aatgataccg cgaccaccga gatctacact agatcgcaca ctctttccct acacgacgct 60
cttccgatct 70

```

-continued

---

SEQ ID NO: 48	moltype = DNA length = 54	
FEATURE	Location/Qualifiers	
misc_feature	1..54	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..54	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 48		
gatcggaaga gccagttacc ttagctgagt cggagacacg cagggatgag atgg		54
SEQ ID NO: 49	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 49		
ctaaggtaac		10
SEQ ID NO: 50	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 50		
taaggagaac		10
SEQ ID NO: 51	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 51		
aagaggattc		10
SEQ ID NO: 52	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 52		
taccaagatc		10
SEQ ID NO: 53	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 53		
cagaaggaac		10
SEQ ID NO: 54	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 54		
ctgcaagttc		10

-continued

---

SEQ ID NO: 55	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 55		
ttcgtgattc		10
SEQ ID NO: 56	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 56		
ttccgataac		10
SEQ ID NO: 57	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 57		
ctgaccgaac		10
SEQ ID NO: 58	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 58		
tcctcgaatc		10
SEQ ID NO: 59	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 59		
tagtggttc		10
SEQ ID NO: 60	moltype = DNA length = 10	
FEATURE	Location/Qualifiers	
misc_feature	1..10	
	note = Description of Artificial Sequence: Synthetic oligonucleotide	
source	1..10	
	mol_type = other DNA	
	organism = synthetic construct	
SEQUENCE: 60		
tctaaccggac		10

---

What is claimed is:

**1.** A method for preparing a nucleic acid for sequencing, the method comprising:

(a) amplifying a single strand of a nucleic acid template comprising a known target nucleotide sequence, a first nucleotide sequence contiguous to the known target nucleotide sequence, and a 5' portion comprising an amplification strand of a ligated universal oligonucleotide tail-adaptor, using (i) a first primer that anneals to a 3' portion of the nucleic acid template comprising the known target nucleotide sequence and (ii) a second primer that comprises a nucleotide sequence identical to a portion of the amplification strand of the ligated universal oligonucleotide tail-adaptor, thereby generating a first amplification product comprising the first nucleotide sequence and the known target nucleotide sequence;

(b) amplifying the first amplification product of (a) using (i) a third primer that comprises a nucleotide sequence identical to a portion of the first primer and a nucleotide sequence that is identical to a portion of a first sequencing primer, and (ii) a fourth primer that comprises a 3' nucleotide sequence identical to a portion of the second primer and a 5' portion comprising a nucleic acid sequence that is identical to a portion of a second sequencing primer,

thereby generating a second amplification product.

**2.** The method of claim **1**, further comprising

(c) determining the sequence of the first nucleotide sequence in the second amplification product.

**3.** The method of claim **2**, wherein step (c) comprises sequencing the second amplification product using at least the first sequencing primer.

**4.** The method of claim **1**, wherein the second primer comprises a nucleotide sequence identical to a portion of the second sequencing primer.

**5.** The method of claim **1**, wherein the amplification strand comprises a barcode portion and the second amplification product comprises the barcode portion.

**6.** The method of claim **1**, wherein the first nucleotide sequence comprises a genetic alteration.

**7.** The method of claim **1**, wherein the nucleic acid template comprises DNA.

**8.** The method of claim **1**, wherein the first nucleotide sequence comprises an insertion or deletion.

**9.** The method of claim **1**, wherein the nucleic acid template comprises cDNA.

**10.** The method of claim **1**, wherein the known target nucleotide sequence and/or the first nucleotide sequence comprises a sequence comprising a gene rearrangement.

**12.** The method of claim **10**, wherein the gene rearrangement comprises a fusion oncogene.

**13.** The method of claim **1**, wherein the nucleic acid template is prepared from a sample obtained from a subject having a condition associated with a genetic alteration.

**14.** The method of claim **1**, wherein the method further comprises, prior to (a), ligating the universal oligonucleotide tail adaptor to the nucleic acid template comprising the known target nucleotide sequence to produce a ligation product.

**15.** The method of claim **14**, wherein the universal oligonucleotide tail adaptor comprises a first ligatable duplex end and a second unpaired end.

**16.** The method of claim **14**, wherein the universal oligonucleotide tail adaptor comprises a blocking strand.

**17.** A method for preparing a nucleic acid for sequencing, the method comprising:

(a) amplifying a single strand of a nucleic acid template comprising a known target nucleotide sequence, a first nucleotide sequence contiguous to the known target nucleotide sequence, and a 5' portion comprising an amplification strand of a ligated universal oligonucleotide tail-adaptor, using (i) a target-specific primer that anneals to the 3' portion of the nucleic acid template comprising the known target nucleotide sequence and (ii) a first adaptor primer that comprises a nucleotide sequence identical to a portion of the amplification strand of the ligated universal oligonucleotide tail-adaptor, thereby generating a first amplification product comprising the first nucleotide sequence and the known target nucleotide sequence;

(b) amplifying the first amplification product of (a) using (i) a second target-specific primer that comprises a nucleotide sequence identical to a portion of the first primer and anneals to the first amplification product of (a), and

(ii) a second adaptor primer that comprises a 3' nucleotide sequence identical to a portion of the first adaptor primer,

thereby generating a second amplification product.

**18.** The method of claim **17**, further comprising:

(d) determining the sequence of the first nucleotide sequence in the second amplification product by a process comprising next-generation sequencing.

\* \* \* \* \*