



US005422104A

United States Patent [19]

[11] Patent Number: **5,422,104**

Fiers et al.

[45] Date of Patent: **Jun. 6, 1995**

[54] **TNF-MUTEINS**

[75] Inventors: **Walter Fiers, Destelbergen; Jan Tavernier, Balegem; Xaveer Van Ostade, Antwerp, all of Belgium**

[73] Assignee: **Hoffmann-La Roche Inc., Nutley, N.J.**

[21] Appl. No.: **794,400**

[22] Filed: **Nov. 20, 1991**

[30] **Foreign Application Priority Data**

Nov. 21, 1990 [EP] European Pat. Off. 90810901

[51] Int. Cl.⁶ **A61K 45/05; C07K 13/00; C12P 21/06**

[52] U.S. Cl. **424/85.1; 530/351; 435/69.5**

[58] Field of Search **530/351, 395, 402; 430/144; 435/69.5; 424/85.1; 574/2.8**

[56] **References Cited**

U.S. PATENT DOCUMENTS

4,650,674 3/1987 Aggarwal et al. .
4,948,875 8/1990 Tanaka et al. 530/350
4,990,455 2/1991 Yamagishi et al. 435/69.5

FOREIGN PATENT DOCUMENTS

40162 12/1989 Australia .
0168214 1/1986 European Pat. Off. .
3843534 7/1990 Germany .
88/06625 9/1988 WIPO .

OTHER PUBLICATIONS

Yamagishi et al, *Protein Eng.* 3(4), 1990 p. 372 (abst only).
Wells et al, *Science* 243, 1989, pp. 1330-1336.
Goh et al, *Protein Eng.* 4(7) 1991, pp. 785-791.
Carlino et al, *JBC* 262, 1987, pp. 958-961.
Tavernier et al, *J. Mol. Biol.* 211(2) 1990, pp. 493-502.
Eck et al, *JBC* 264, 1989, pp. 17595-17606.

Yamagishi et al *Protein Engineering* 3(8) 1990, pp. 713-719.

Yamagishi et al *Protein Engineering* 3(4) 1990 (previously cited).

Tsujimoto et al, *J. Biochem* 101, 1987, pp. 919-925.

Tartaglia, et al, *Immunology Today*, vol. 13, No. 5, (1992).

Barrett, et al, *Eur. Journal Immunology* vol. 21, 1649-1656 (1991).

Lewis, et al, *Proc. Natl. Acad. Science* vol. 88, 2830-2834 (1991).

Masegi, et al. *Protein Engineering Engineering* 375-376 (1889).

Tsukio, *Patent Abstract of Japan*, vol. 13, No. 119 (1989).

Ostade, et al. *The Embo Journal*, vol. 10, No. 4 827-836 (1991).

Primary Examiner—Garnette D. Draper

Assistant Examiner—K. Cochrane Carlson

Attorney, Agent, or Firm—George M. Gould; William H. Epstein; Catherine A. Picut

[57] **ABSTRACT**

It is an object of this invention to provide a human Tumor Necrosis Factor mutein or a pharmaceutically acceptable salt thereof characterized in that the TNF sequence is changed by a deletion, insertion, substitution or combinations thereof, of one or more amino acids so that the mutein shows a significant difference between its binding affinity to the human p75-Tumor-Necrosis-Factor-Receptor and to the human p55-Tumor-Necrosis-Factor-Receptor. The invention also includes DNA sequences coding for such muteins, vectors comprising such DNA sequences, host cells transformed with such vectors and a process for the production of such muteins employing such transformed host cells and pharmaceutical compositions containing such muteins and their use for the treatment of illnesses, for example cancer.

4 Claims, 16 Drawing Sheets




FIG. 1a




FIG. 1b




FIG. 2a

Hind III
1 AAGCTTCACG CTGCCGCAAG CACTCAGGGC GCAAGGGCTG CTAAAGGAAG
51 CGGAACACGT AGAAAGCCAG TCCGCAGAAA CGGTGCTGAC CCCGGATGAA
101 TGTCAGCTAC TGGGCTATCT GGACAAGGGA AAACGCAAGC GCAAAGAGAA
151 AGCAGGTAGC TTGCAGTGGG CTTACATGGC GATAGCTAGA CTGGGCGGTT
201 TTATGGACAG CAAGCGAACC GGAATTGCCA GCTGGGGCGC CCTCTGGTAA
251 GGTTGGGAAG CCCTGCAAAG TAAACTGGAT GGCTTTCTTG CCGCCAAGGA
301 TCTGATGGCG CAGGGGATCA AGATCTGATC AAGAGACAGG ATGACGGTCG
351 TTTCGCATGC TTGAACAAGA TGGATTGCAC GCAGGTTCTC CGGCCGCTTG
401 GGTGGAGAGG CTATTCGGCT ATGACTGGGC ACAACAGACA ATCGGCTGCT
451 CTGATGCCGC CGTGTTCCGG CTGTCAGCGC AGGGGCGCCC GGTTCTTTTT
501 GTCAAGACCG ACCTGTCCGG TGCCCTGAAT GAACTGCAGG ACGAGGCAGC
551 GCGGCTATCG TGGCTGGCCA CGACGGGCGT TCCTTGCGCA GCTGTGCTCG
601 ACGTTGTCAC TGAAGCGGGA AGGGACTGGC TGCTATTGGG CGAAGTGCCG
651 GGGCAGGATC TCCTGTCATC TCACCTTGCT CCTGCCGAGA AAGTATCCAT
701 CATGGCTGAT GCAATGCGGC GGCTGCATAC GCTTGATCCG GCTACCTGCC
751 CATTGACCA CCAAGCGAAA CATCGCATCG AGCGAGCACG TACTCGGATG
801 GAAGCCGGTC TTGTCGATCA GGATGATCTG GACGAAGAGC ATCAGGGGCT
851 CGCGCCAGCC GAACTGTTCG CCAGGCTCAA GCGCGCATG CCCGACGGCG
901 AGGATCTCGT CGTGACCCAT GGCGATGCCT GCTTGCCGAA TATCATGGTG
951 GAAAATGGCC GCTTTTCTGG ATTCATCGAC TGTGGCCGGC TGGGTGTGGC
1001 GGACCGCTAT CAGGACATAG CGTTGGCTAC CCGTGATATT GCTGAAGAGC
1051 TTGGCGGCGA ATGGGCTGAC CGCTTCCTCG TGCTTTACGG TATCGCCGCT
1101 CCCGATTCGC AGCGCATCGC CTTCTATCGC CTTCTTGACG AGTTCTTCTG
1151 AGCGGGACTC TGGGGTTCGA AATGACCGAC CAAGCGACGC CCAACCTGCC
1201 ATCACGAGAT TTCGATTCCA CCGCCGCCTT CTATGAAAGG TTGGGCTTCG
1251 GAATCGTTTT CCGGGACGCC GGCTGGATGA TCCTCCAGCG CGGGGATCTC
1301 ATGCTGGAGT TCTTCGCCCA CCCCAGGGCTC GATCCCCTCG CGAGTTGGTT

FIG. 2b

1351 CAGCTGCTGC CTGAGGCTGG ACGACCTCGC GGAGTTCTAC CGGCAGTGCA
1401 AATCCGTCGG CATCCAGGAA ACCAGCAGCG GCTATCCGCG CATCCATGCC
1451 CCCGAACTGC AGGAGTGGGG AGGCACGATG GCCGCTTTGG SalI TCGACAATTC
1501 GCGCTAACTT ACATTAATTG CGTTGCGCTC ACTGCCCGCT TTCCAGTCGG
1551 GAAACCTGTC GTGCCAGCTG CATTAAATGAA TCGGCCAACG CGCGGGGAGA
1601 GGCGGTTTGC GTATTGGGCG CCAGGGTGGT TTTTCTTTTC ACCAGTGAGA
1651 CGGGCAACAG CTGATTGCCC TTCACCGCCT GGCCCTGAGA GAGTTGCAGC
1701 AAGCGGTCCA CGCTGGTTTG CCCAGCAGG CGAAAATCCT GTTTGATGGT
1751 GGTTAACGGC GGGATATAAC ATGAGCTGTC TTCGGTATCG TCGTATCCCA
1801 CTACCGAGAT ATCCGCACCA ACGCGCAGCC CGGACTCGGT AATGGCGCGC
1851 ATTGCGCCCA GCGCCATCTG ATCGTTGGCA ACCAGCATCG CAGTGGGAAC
1901 GATGCCCTCA TTCAGCATT TGCATGGTTG TTGAAAACCG GACATGGCAC
1951 TCCAGTCGCC TTCCCGTTCC GCTATCGGCT GAATTTGATT GCGAGTGAGA
2001 TATTTATGCC AGCCAGCCAG ACGCAGACGC GCCGAGACAG AACTTAATGG
2051 GCCCGCTAAC AGCGCGATTT GCTGGTGACC CAATGCGACC AGATGCTCCA
2101 CGCCCAGTCG CGTACCGTCT TCATGGGAGA AAATAATACT GGTGATGGGT
2151 GTCTGGTCAG AGACATCAAG AAATAACGCC GGAACATTAG TGCAGGCAGC
2201 TTCCACAGCA ATGGCATCCT GGTCATCCAG CGGATAGTTA ATGATCAGCC
2251 CACTGACGCG TTGCGCGAGA AGATTGTGCA CCGCCGCTTT ACAGGCTTCG
2301 ACGCCGCTTC GTTCTACCAT CGACACCACC ACGCTGGCAC CCAGTTGATC
2351 GGCGCGAGAT TTAATCGCCG CGACAATTTG CGACGGCGCG TGCAGGGCCA
2401 GACTGGAGGT GGCAACGCCA ATCAGCAACG ACTGTTTGCC CGCCAGTTGT
2451 TGTGCCACGC GGTTGGGAAT GTAATTCAGC TCCGCCATCG CCGCTTCCAC
2501 TTTTCCCGC GTTTTCGCAG AACGTGGCT GGCCTGGTTC ACCACGCGGG
2551 AACGGTCTG ATAAGAGACA CCGGCATACT CTGCGACATC GTATAACGTT
2601 ACTGGTTTCA CATTACCCAC CCTGAATTGA CTCTCTTCCG GGCGCTATCA
2651 TGCCATACCG CGAAAGGTTT TGCGCCATTC SalI GATGGTGTCA ACGTAAATGC
2701 ATGCCGCTTC GCCTTCGCGC GCGAATTGTC GACCCTGTCC CTCCTGTTCA

FIG. 2c

2751 GCTACTGACG GGGTGGTGCG TAACGGCAAAG AGCACCGCCG GACATCAGCG
2801 CTAGCGGAGT GTATACTGGC TTACTATGTT GGCACCTGATG AGGGTGTCAG
2851 TGAAGTGCTT CATGTGGCAG GAGAAAAAAG GCTGCACCGG TGCATCAGCA
2901 GAATATGTGA TACAGGATAT ATTCCGCTTC CTCGCTCACT GACTCGCTAC
2951 GCTCGGTCGT TCGACTGCGG CGAGCGGAAA TGGCTTACGA ACGGGGCGGA
3001 GATTCCTGG AAGATGCCAG GAAGATACTT AACAGGGAAG TGAGAGGGCC
3051 GCGGCAAAGC CGTTTTTCCA TAGGCTCCGC CCCCTGACA AGCATCACGA
3101 AATCTGACGC TCAAATCAGT GGTGGCGAAA CCCGACAGGA CTATAAAGAT
3151 ACCAGGCGTT TCCCCTGGCG GCTCCCTCGT GCGCTCTCCT GTTCCTGCCT
3201 TTCGGTTTAC CGGTGTCATT CCGCTGTTAT GGCCGCGTTT GTCTCATTCC
3251 ACGCCTGACA CTCAGTTCCG GGTAGGCAGT TCGCTCCAAG CTGGACTGTA
3301 TGCACGAACC CCCCGTTCAG TCCGACCGCT GCGCCTTATC CGGTAAGTAT
3351 CGTCTTGAGT CCAACCCGGA AAGACATGCA AAAGCACCCAC TGGCAGCAGC
3401 CACTGGTAAT TGATTTAGAG GAGTTAGTCT TGAAGTCATG CGCCGGTTAA
3451 GGCTAAACTG AAAGGACAAG TTTTGGTGAC TGCGCTCCTC CAAGCCAGTT
3501 ACCTCGGTTC AAAGAGTTGG TAGCTCAGAG AACCTTCGAA AAACCGCCCT
3551 GCAAGGCGGT TTTTTCGTTT TCAGAGCAAG AGATTACGCG CAGACCAAAA
3601 CGATCTCAAG AAGATCATCT TATTAATCAG ATAAAATATT TCTAGATTTC
3651 AGTGCAATTT ATCTCTTCAA ATGTAGCACC TGAAGTCAGC CCCATACGAT
3701 ATAAGTTGTT AATTCTCATG TTTGACAGCT TATCATCGAT

FIG. 2d




FIG. 3a

^{XhoI}
1 CTCGAGAAAT CATAAAAAT TTATTTGCTT TGTGAGCGGA TAACAATTAT
51 AATAGATTCA ATTGTGAGCG GATAACAATT TCACACAGAA ^{EcoRI}TTCATTAAAG
101 AGGAGAAATT AAGCATGGTC AGATCATCTT CTCGAACCCC GAGTGACAAG
151 CCTGTAGCCC ATGTTGTCGC GAACCCTCAA GCTGAGGGGC AGCTCCAGTG
201 GCTGAACCGC CGGGCCAATG CCCTCCTGGC CAATGGCGTG GAGCTGAGAG
251 ATAACCAGCT GGTGGTGCCA TCAGAGGGCC TGTACCTCAT CTACTCCCAG
301 GTCCTCTTCA AGGGCCAAGG CTGCCCCTCC ACCCATGTGC TCCTCACCCA
351 CACCATCAGC CGCATCGCCG TCTCCTACCA GACCAAGGTC AACCTCCTCT
401 CTGCCATCAA GAGCCCCTGC CAGAGGGAGA CCCAGAGGG GGCTGAGGCC
451 AAGCCCTGGT ATGAGCCCAT CTATCTGGGA GGGGTCTTCC AGCTGGAGAA
501 GGGTGACCGA CTCAGCGCTG AGATCAATCG GCCCGACTAT CTCGACTTTG
551 CCGAGTCTGG GCAGGTCTAC TTTGGGATCA TTGCCCTGTG AGGAGGACGA
601 ACATCCAACC TTCCCAAACG CCTCCCCTGC CCCAATCCCT TTATTACCCC
651 CTCCTTCAGA CACCCTCAAC CTCTTCTGGC TCAAAAAGAG AATTGGGGGC
701 TTAGGGTCGG ^{HindIII}AACCCAAGCT TGGACTCCTG TTGATAGATC CAGTAATGAC
751 CTCAGAACTC CATCTGGATT TG TTCAGAAC GCTCGGTTGC CGCCGGGCGT
801 TTTTTATTGG TGAGAATCCA AGCTAGCTTG GCGAGATTTT CAGGAGCTAA
851 GGAAGCTAAA ATGGAGAAAA AAATCACTGG ATATACCACC GTTGATATAT
901 CCCAATGGCA TCGTAAAGAA CATTTTGAGG CATTTCAGTC AGTTGCTCAA
951 TGTACCTATA ACCAGACCGT TCAGCTGGAT ATTACGGCCT TTTTAAAGAC
1001 CGTAAAGAAA AATAAGCACA AGTTTTATCC GGCCTTTATT CACATTCTTG
1051 CCCGCCTGAT GAATGCTCAT CCGGAATTC GTATGGCAAT GAAAGACGGT
1101 GAGCTGGTGA TATGGGATAG TG TTCACCCT TGTTACACCG TTTTCCATGA
1151 GCAAACCTGAA ACGTTTTTCAT CGCTCTGGAG TGAATACCAC GACGATTTCC
1201 GGCAGTTTCT ACACATATAT TCGCAAGATG TGGCGTGTTA CGGTGAAAAC
1251 CTGGCCTATT TCCCTAAAGG GTTTATTGAG AATATGTTTT TCGTCTCAGC

FIG. 3b

1301 CAATCCCTGG GTGAGTTTCA CCAGTTTTGA TTAAACGTG GCCAATATGG
1351 ACAACTTCTT CGCCCCGTT TTCACCATGG GCAAATATTA TACGCAAGGC
1401 GACAAGGTGC TGATGCCGCT GGCGATTCAG GTTCATCATG CCGTCTGTGA
1451 TGGCTTCCAT GTCGGCAGAA TGCTTAATGA ATTACAACAG TACTGCGATG
1501 AGTGGCAGGG CGGGGCGTAA TTTTTTTAAG GCAGTTATTG GTGCCCTTAA
1551 ACGCCTGGGG TAATGACTCT CTAGCTTGAG GCATCAAATA AAACGAAAGG
1601 CTCAGTCGAA AGACTGGGCC TTTCGTTTTA TCTGTTGTTT GTCGGTGAAC
1651 GCTCTCCTGA GTAGGACAAA TCCGCCGCTC TAGAGCTGCC TCGCGCGTTT
1701 CGGTGATGAC GGTGAAAACC TCTGACACAT GCAGCTCCCG GAGACGGTCA
1751 CAGCTTGTCT GTAAGCGGAT GCCGGGAGCA GACAAGCCCG TCAGGGCGCG
1801 TCAGCGGGTG TTGGCGGGTG TCGGGGCGCA GCCATGACCC AGTCACGTAG
1851 CGATAGCGGA GTGTATACTG GCTTAACTAT GCGGCATCAG AGCAGATTGT
1901 ACTGAGAGTG CACCATATGC GGTGTGAAAT ACCGCACAGA TCGTAAGGA
1951 GAAAATACCG CATCAGGCGC TCTTCCGCTT CCTCGCTCAC TGA CTGCTG
2001 CGCTCGGTCT GTCGGCTGCG GCGAGCGGTA TCAGCTCACT CAAAGGCGGT
2051 AATACGGTTA TCCACAGAAT CAGGGGATAA CGCAGGAAAG AACATGTGAG
2101 CAAAAGGCCA GCAAAGGCC AGGAACCGTA AAAAGGCCGC GTTGCTGGCG
2151 TTTTTCCATA GGCTCCGCCC CCCTGACGAG CATCACAAA ATCGACGCTC
2201 AAGTCAGAGG TGGCGAAACC CGACAGGACT ATAAAGATAC CAGGCGTTTC
2251 CCCCTGGAAG CTCCTCGTG CGCTCTCCTG TTCCGACCCT GCCGCTTACC
2301 GGATACCTGT CCGCCTTTCT CCCTTCGGGA AGCGTGGCGC TTTCTCAATG
2351 CTCACGCTGT AGGTATCTCA GTTCGGTGTA GGTCGTTTCGC TCCAAGCTGG
2401 GCTGTGTGCA CGAACCCCC GTTCAGCCG ACCGCTGCGC CTTATCCGGT
2451 AACTATCGTC TTGAGTCCAA CCCGGTAAGA CACGACTTAT CGCCACTGGC
2501 AGCAGCCACT GGTAACAGGA TTAGCAGAGC GAGGTATGTA GCGGGTGCTA
2551 CAGAGTTCTT GAAGTGGTGG CCTAACTACG GCTACACTAG AAGGACAGTA
2601 TTTGGTATCT GCGCTCTGCT GAAGCCAGTT ACCTTCGGAA AAAGAGTTGG
2651 TAGCTCTTGA TCCGGCAAAC AAACCACCGC TGGTAGCGGT GGTTTTTTTG

FIG. 3c

2701 TTTGCAAGCA GCAGATTACG CGCAGAAAAA AAGGATCTCA AGAAGATCCT
2751 TTGATCTTTT CTACGGGGTC TGACGCTCAG TGGAACGAAA ACTCACGTTA
2801 AGGGATTTTG GTCATGAGAT TATCAAAAAG GATCTTCACC TAGATCCTTT
2851 TAAATTA AAA ATGAAGTTTT AAATCAATCT AAAGTATATA TGAGTAAACT
2901 TGGTCTGACA GTTACCAATG CTTAATCAGT GAGGCACCTA TCTCAGCGAT
2951 CTGTCTATTT CGTTCATCCA TAGCTGCCTG ACTCCCCGTC GTGTAGATAA
3001 CTACGATACG GGAGGGGCTTA CCATCTGGCC CCAGTGCTGC AATGATACCG
3051 CGAGACCCAC GCTCACCGGC TCCAGATTTA TCAGCAATAA ACCAGCCAGC
3101 CGGAAGGGCC GAGCGCAGAA GTGGTCCTGC AACTTTATCC GCCTCCATCC
3151 AGTCTATTAA TTGTTGCCGG GAAGCTAGAG TAAGTAGTTC GCCAGTTAAT
3201 AGTTTGCGCA ACGTTGTTGC CATTGCTACA GGCATCGTGG TGTCACGCTC
3251 GTCGTTTGGT ATGGCTTCAT TCAGCTCCGG TTCCAACGA TCAAGGCGAG
3301 TTACATGATC CCCCATGTTG TGCAAAAAAG CGGTTAGCTC CTTGGTCTCCT
3351 CCGATCGTTG TCAGAAGTAA GTTGGCCGCA GTGTTATCAC TCATGGTTAT
3401 GGCAGCACTG CATAATTCTC TTA CTGTCAT GCCATCCGTA AGATGCTTTT
3451 CTGTGACTGG TGAGTACTCA ACCAAGTCAT TCTGAGAATA GTGTATGCGG
3501 CGACCGAGTT GCTCTTGCCC GGCGTCAATA CGGGATAATA CCGCGCCACA
3551 TAGCAGA AACT TAAAAGTGC TCATCATTGG AAAACGTTCT TCGGGGCGAA
3601 AACTCTCAAG GATCTTACCG CTGTTGAGAT CCAGTTCGAT GTAACCCACT
3651 CGTGCACCCA ACTGATCTTC AGCATCTTTT ACTTTCACCA GCGTTTCTGG
3701 GTGAGCAAAA ACAGGAAGGC AAAATGCCGC AAAAAAGGGA ATAAGGGCGA
3751 CACGGAAATG TTGAATACTC ATACTCTTCC TTTTTC AATA TTATTGAAGC
3801 ATTTATCAGG GTTATTGTCT CATGAGCGGA TACATATTTG AATGTATTTA
3851 GAAAAATAAA CAAATAGGGG TTCCGCGCAC ATTTCCCGA AAAGTGCCAC
3901 CTGACGTCTA AGAAACCATT ATTATCATGA CATTAACCTA TAAAAATAGG
3951 CGTATCACGA GGCCCTTTCG TCTTCAC

FIG. 3d




FIG. 4




FIG. 5




FIG. 6(a)




FIG. 6(b)




FIG. 6(c)




FIG. 6(d)




FIG. 6(e)




FIG. 6(f)




FIG. 7(a)




FIG. 7(b)




FIG. 7(c)




FIG. 7(d)




FIG. 7(e)




FIG. 8

TNF-MUTEINS

BACKGROUND OF INVENTION

Tumor Necrosis Factor, or more specifically Tumor Necrosis Factor-alpha, is a cytokine, primarily produced by stimulated macrophages, that exhibits not only a striking cytotoxicity against various tumour cells [Carswell et al., *Proc. Nat. Acad. Sci., U.S.A.* 72, 3666-3670, (1975)] but also plays a multiple role as a mediator of inflammation and the immune response [See. Beutler and Cerami, *Ann. Rev. Immunol.* 7, 625-655 (1989); Bonavista and Granger (eds.) "Tumor Necrosis Factor: Structure, Mechanism of Action, Role in Disease and Therapy, Karger, Basel (1990)]. The primary structure of human Tumor Necrosis Factor-alpha (hTNF- α) has been deduced from the nucleotide sequence of a cDNA which has been cloned and expressed in *E. coli* [Pennica et al., *Nature* 312, 724-729 (1984); Marmenout et al., *Europ. J. Biochem.* 152, 515-522 (1985); Wang et al., *Science* 226, 149-154 (1985); Shirai et al., *Nature* 313, 803-806 (1985)]. A striking homology in amino acid sequence (30%) was found between hTNF- α and human Lymphotoxin, often referred to as human Tumor Necrosis Factor-beta (hTNF- β), a cytokine produced by a subset of lympho-

polyacrylamide gel electrophoresis (SDS-PAGE) (p55-TNF-R) and a receptor with an apparent molecular weight of 75 kD on SDS-PAGE (p75-TNF-R). Both forms of TNF-receptors have been cloned previously. The cloning of p55-TNF-R was done by Loetscher et al. [*Cell* 61, 351-359, (1990)] and the cloning of p75-TNF-R was done by Dembic et al. [*Cytokine* 2, 53-58, (1990)] See also European Patent Application No. 90116707.2 (both receptors). It was found more recently that both receptors bind not only TNF- α , but also TNF- β with high affinity [Schönfeld et al., *J. Biol. Chem.* 266, 3863-3869 (1991)].

SUMMARY OF THE INVENTION

An object of the present invention is a mutein or a pharmaceutically acceptable salt thereof of human Tumor Necrosis Factor having an amino acid sequence which is changed by deletion, insertion and/or substitution of one or more amino acids such that the mutein shows a significant difference between its binding affinity to the human p75-Tumor-Necrosis-Factor-Receptor and the human p55-Tumor-Necrosis-Factor-Receptor.

A preferred embodiment of the present invention is a mutein as defined above on the basis of the amino acid sequence of TNF- α as disclosed by Pennica et al. supra, namely [SEQ ID No: 1]

1	VAL	ARG	SER	SER	SER	ARG	THR	PRO	SER	10	ASP	LYS	PRO	VAL	ALA	HIS
					20											30
	VAL	VAL	ALA	ASN	PRO	GLN	ALA	GLU	GLY	GLN	LEU	GLN	TRP	LEU	ASN	
	ARG	ARG	ALA	ASN	ALA	LEU	LEU	ALA	ASN	40	GLY	VAL	GLU	LEU	ARG	ASP
	ASN	GLN	LEU	VAL	50	VAL	PRO	SER	GLU	GLY	LEU	TYR	LEU	ILE	TYR	SER
	GLN	VAL	LEU	PHE	LYS	GLY	GLN	GLY	CYS	70	PRO	SER	THR	HIS	VAL	LEU
	LEU	THR	HIS	THR	80	ILE	SER	ARG	ILE	ALA	VAL	SER	TYR	GLN	THR	LYS
	VAL	ASN	LEU	LEU	SER	ALA	ILE	LYS	SER	100	PRO	CYS	GLN	ARG	GLU	THR
	PRO	GLU	GLY	ALA	110	GLU	ALA	LYS	PRO	TRP	TYR	GLU	PRO	ILE	TYR	LEU
	GLY	GLY	VAL	PHE	GLN	LEU	GLU	LYS	GLY	130	ASP	ARG	LEU	SER	ALA	GLU
	ILE	ASN	ARG	PRO	140	ASP	TYR	LEU	ASP	PHE	ALA	GLU	SER	GLY	GLN	VAL
	TYR	PHE	GLY	ILE	157	ILE	ALA	LEU								

cytes [Gray et al., *Nature* 312, 721-724 (1984); Fiers et al., *Cold Spring Harbour Symp.* 51, 587-595 (1986)].

hTNF- α with modified amino acid sequences, so called TNF- α -muteins, have also been described in the art [See, e.g., Yamagishi et al., *Protein Engineering* 3, 713-719, (1990) or by Fiers in "Tumor Necrosis Factors: Structure, Function and Mechanism of Action", Aggarwal and Vilcek (eds.), Marcel Dekker, Inc., New York, (in press), or by Fiers et al. in Bonavista and Granger, pp. 77-81 supra. In addition TNF- α -muteins have also been the object of several patent applications, for example, International Patent Applications Publ. Nos. WO 86/02381, WO 86/04606, WO 88/06625 and European Patent Applications Publ. Nos. 155,549; 158,286; 168,214; 251,037 and 340,333, and Deutsche Offenlegungsschrift Nr. 3843534.

Muteins of Lymphotoxin have also been disclosed in the art, for example in European Patent Applications Publ. Nos. 250,000; 314,094 and 336,383.

The biological effects of TNF are mediated via specific receptors, namely a receptor with an apparent molecular weight of 55 kD on sodium dodecylsulfate

or as disclosed by Marmenout et al. supra or Wang et al. supra or Shirai et al. supra. More specifically muteins of deduced amino acid sequence as are coded for by the nucleotide sequence of the insert of the plasmid pDS56/RBSII,Sph1-TNF α [SEQ ID No: 2] (See also FIG. 3a and 3B) coding for mature TNF- α .

Another preferred embodiment of the present invention is a mutein as defined above wherein the TNF- α amino acid sequence is changed by substitution of one or more amino acids, preferably one or two by other amino acids, and preferably by naturally occurring amino acids.

Another preferred embodiment is a human Tumor Necrosis Factor mutein wherein SEQ ID NO: 1 is changed by deletion, insertion, substitution or combinations thereof, of between one and 10 amino acids.

A more preferred embodiment of the present invention are muteins as defined above wherein the TNF- α amino acid sequence is substituted at position 29 and/or 32 or position 31 and 32 or position 31 or position 29 and 31 whereby substitutions at position 29 and/or 32 or position 31 and 32 or position 31 are preferred (referring

to [SEQ ID No:1]) by other amino acids, preferably naturally occurring amino acids. Any amino acid, preferably any naturally occurring one, can be used at one or more of these positions which leads to a TNF-mutein showing a significant difference between its binding affinity to the human p75-TNF-R and the human p55o TNF-R. For substitutions at position 29 serine [SEQ ID No:4], glycine [SEQ ID No:5] or tyrosine [SEQ ID No:6] are preferred, serine is especially preferred, for example in case of a single position mutein at position 29 (Ser²⁹-TNF α) [SEQ ID No:4]. For substitutions at position 31 glutamic acid, for example Glu³¹-TNF α [SEQ ID No:7], or asparagine [SEQ ID No: 8] are preferred. For substitutions at position 32 tyrosine, for example Tyr³²-TNF α [SEQ ID No:10] or tryptophan, for example Trp³²-TNF α [SEQ ID No:9] are preferred, Trp³² is specifically preferred. Especially preferred substitutions in case of a double position mutein at positions 29 and 32 are Ser²⁹-Trp³²-TNF α [SEQ ID No: 12] and at position 31 and 32 are Asn³¹-Thr³²-TNF α . [SEQ ID No: 11]. It is understood that the muteins of the present invention can also be prepared by methods known in the art of chemical peptide and protein synthesis, for example by partial or total liquid or solid phase synthesis as described by Gross and Meyenhofer in "The Peptides" Vols. 1-9, Academic Press, Inc., Harcourt Brace Jovanovich, Publs., San Diego (1979-1987) or by Fields and Nobel, Int. J. Pept. Prot. Res. 35, 161-214 (1990).

Another preferred embodiment of the present invention is a mutein of TNF- α comprising the amino acid sequence set forth in SEQ ID No: 1 wherein at least one of the positions 29, 31 or 32 is substituted with any naturally occurring amino acid different from the corresponding amino acid in SEQ ID No: 1.

Analogs obtained by deletion, substitution or addition or combinations thereof of one or several amino acids from or to the muteins as defined in the previous paragraph, whereby position 29 and/or 32 or position 31 or position 31 and 32 in the mutein are not changed and which analogs still show a significant difference between its binding affinity to the human p75-TNF-R and the human p55-TNF-R are also an object of the present invention. With respect to such substitution analogs, amino acid substitutions in proteins which do not generally alter the activity are known in the state of the art and are described, for example, by H. Neurath and R. L. Hill in "The Proteins" (Academic Press, New York, 1979, see especially FIG. 6, page 14). The most commonly occurring exchanges are: Ala/Ser, Val/Ile, Asp/Glu, Thr/Ser, Ala/Gly, Ala/Thr, Ser/Asn, Ala/Val, Ser/Gly, Tyr/Phe, Ala/Pro, Lys/Arg, Asp/Asn, Leu/Ile, Leu/Val, Ala/Glu, Asp/Gly as well as these in reverse (the three letter abbreviations are used for amino acids and are standard and known in the art).

Analogs made by substitution, addition, deletion or combinations thereof can be produced by methods known in the art and described for example in Sambrook et al. [Molecular Cloning, A Laboratory Manual, 2nd ed., Cold Spring Harbour Laboratory, Cold Spring Harbour Laboratory Press, USA (1989)] or as described herein. Whether such an analog still shows the significant difference between its binding affinity to the p75-TNF-R and the p55-TNF-R can be determined as described below and more specifically in Examples II) and 2) or Example VIII. Furthermore, salts of such muteins and analogs are also an object of the present invention. Such salts can be produced by methods known in the art.

It is furthermore an object of the present invention to provide a mutein as described above for the treatment of illnesses, for example cancer.

It is well known in the art that on the basis of its biological activities TNF- α can be a valuable compound for the treatment of various disorders. For example TNF- α , alone or in combination with interferon, can be an effective antitumor agent [Brouckaert et al., Int. J. Cancer 38, 763-769 (1986)]. However, its systemic toxicity is a major limitation to its wider therapeutic use [Taguchi T. and Sohmura Y., Biotherapy 3, 177-186 (1991)].

The discovery of two TNF-receptors with (putatively) distinct functional roles should allow one to separate in a given disease state the beneficial and unwanted biological responses to TNF. There is circumstantial evidence supporting the feasibility of this approach. It has been shown for example [Brouckaert et al., Agents and Actions 26, 196-197 (1989); Everaerd, B. et al., Biochem. Biophys. Res. Comm. 163, 378-385 (1989)] that in mice, murine TNF- α (mTNF- α) is up to 50-fold more toxic than human TNF- α (hTNF- α), although when tested in cell culture (murine and human), both are equally active on sensitive cell lines.

It is believed that the strategy of separating beneficial and unwanted TNF α activities by using compounds specifically binding to one or the other TNF-receptor, such as the TNF-muteins of the present invention, can be used in general in other disease states where TNF plays a role.

DNA-sequences comprising a DNA-sequence coding for TNF-muteins as hereinbefore described are also an object of the present invention. Such DNA-sequences can be constructed starting from genomic-or cDNA-sequences coding for hTNF as disclosed in the art using known methods of in vitro mutagenesis [see e.g. Sambrook et al., 1989]. Such mutagenesis can be carried out at random in order to obtain a large number of mutants which can then be tested for their desired properties in appropriate assay systems or, in order to mutate defined positions in a given DNA-sequence, by so called site directed mutagenesis [see, e.g., Sambrook et al., 1989, 15.51-15.113] or by mutagenesis using the polymerase chain reaction [see, e.g., White et al., Trends in Genetics 5, 185-189 (1989)].

A preferred embodiment of the invention is a purified and isolated DNA sequence comprising positions 115 to 591 of SEQ ID NO:2 wherein the DNA sequence is changed by deletion, insertion, substitution or combinations thereof, such that the DNA sequence codes for a human Tumor Necrosis Factor mutein containing at least one amino acid different from SEQ ID No: 1 and the mutein shows a significant difference between its binding affinity to the human p75-(Tumor Necrosis Factor)-Receptor and to human p55-(Tumor Necrosis Factor)-Receptor.

Another preferred embodiment is a purified and isolated DNA sequence comprising positions 115 to 591 of SEQ ID No: 13 wherein at least one of the codons at positions 202 to 204, 208 to 210, or 211 to 213 codes for an amino acid different from the amino acid coded for by the corresponding condon in SEQ ID No: 2.

One chemical mutagen which is often used for random mutagenesis is sodium bisulfite which converts cytosin residues into uracil residue and hence leads to a transition of "C" to "T" (standard abbreviations for nucleotides) [for the method see e.g. Shortle and Nathans, Proc. Nat. Acad. Sci. U.S.A. 75, 2170-2174

(1978) or Pine and Huang, *Meth. Enzym.* 154, 415-430 (1987)]. This mutagen acts solely on single stranded DNA whereas the expression of the mutated target DNA sequence is achieved with a double stranded plasmid vector. One possibility to avoid the necessity of recloning in mutagenesis and expression vectors is the use of so called "phasmids". These are vectors which, in addition to a plasmid origin of replication, carry also an origin of replication derived from a filamentous phage. Examples of such phasmids are the pMa- and pMcphasmids as described by Stanssen et al. [*Nucleic Acids Res.* 17, 4441-4454, (1989)]. Using this expression system one can construct so called "gap-duplex"-structures [see also Kramer et al., *Nucl. Acids. Res.* 12, 9441-9456 (1984)] where only the TNF-coding sequence is in a single stranded configuration and therefore accessible for the specific chemical mutagen. "Gap-duplexes" to be used in at random mutagenesis can be constructed as described for site-specific mutagenesis by Stanssen et al. supra with the exception that the (-)strand contains the same active antibiotic resistance gene as the (+)strand. By making use of different restriction sites in the DNA-sequence encoding hTNF α [SEQ ID No:2], variation of the width of the gap is possible. Examples of such restriction sites are the C1a1-Sal1 sites (470 nucleotides), BstX1-BstX1 sites (237 nucleotides) or Styl-Styl sites (68 nucleotides). Such gap-duplex-constructs can then be treated with increasing concentrations (up to 4M) of bisulfite, followed by several dialysis steps, as described by Shortle and Nathans supra. A suitable procaryotic host cell can then be transformed by such phasmid constructs according to methods known in the art and described for example by Sambrook et al. supra. A suitable procaryotic host cell means in this context a host cell deficient in a specific repair function so that an uracil residue is maintained in the DNA during replication and which host cell is capable of expressing the corresponding mutated TNF. Such specific host strains are known in the art, for example for *E. coli* strains, e.g. *E. coli* BW 313 [Kunkel, T.A., *Proc. Natl. Acad. Sci. USA* 82, 488-492 (1985)]. The resulting clones can then be screened for those expressing a desired TNF-mutagen by appropriate assay systems. For example each colony can be inoculated in a microtiterplate in a suitable medium containing the relevant antibiotic. The cells may be lysed by addition of lysozyme, followed by sequential freeze-thaw cycles. After precipitation of nucleic acids and centrifugation, the supernatant of each colony can directly be used in appropriate assays as described, for example, in Example IIa and IIb or Example VIII measuring binding to the p75-TNF-R and the p55-TNF-R on the surface of living cells or in purified form.

If desired, the specific sites of mutation can be determined, for example by restriction fragment analysis [see, e.g., Sambrook et al. Supra]. By determination of the DNA-sequence of such fragments the exact position of the mutation can be determined and if such mutation leads to an amino acid replacement the new amino acid can be derived from the determined DNA-sequence. DNA-sequencing can be performed according to methods known in the art, for example by using T7 polymerase on supercoiled DNA with a commercially available sequencing kit (Pharmacia, Uppsala, Sweden).

As already mentioned above, another possibility of mutating a given DNA-sequence is by "site directed mutagenesis". A widely used strategy for such kind of mutagenesis as originally outlined by Hutchinson and Edgell [*J. Virol.* 8, 181 (1971)] involves the annealing of

a synthetic oligonucleotide carrying the desired nucleotide substitution to a target region of a single stranded DNA-sequence wherein the mutation should be introduced [for review see Smith, *Annual. Rev. Genet.* 19,423 (1985) and for improved methods see references 2-6 in Stanssen et al. supra.

One such preferred method is the one of Stanssen et al. supra (1989) using "gapped duplex DNA" as originally described by Kramer et al. supra (1984) [see also Kramer and Fritz, *Methods in Enzymology*, (1987), Academic Press, Inc., USA], but using antibiotic resistance genes instead of M13 functional genes for selection of the mutation containing strand as well as the phasmid-technology described by Stanssen et al. supra (1989). An advantage of this method lies also in the capability of performing successive cycles of mutagenesis without the need to transfer the gene to a new mutagenesis vector. The second round mutagenesis differs only in the selection using another antibiotic marker (Stanssen et al., supra). As a control, site-specific back mutagenesis of the mutant to the wild-type TNF can be used. In addition, the use of an oligonucleotide, creating or destroying a restriction site in the TNF gene, allows one to control the mutant not only by hybridization to the oligonucleotide used for site directed mutagenesis but also by the presence or absence of the restriction site. In order to create a set of TNF-mutagens wherein at a defined position of their amino acid sequence the wild-type amino acid, is replaced by any naturally occurring amino acid a set of oligonucleotides is used with all possible codons at the defined position.

As already mentioned above, another possibility of mutating a given DNA-sequence is the mutagenesis by using the polymerase chain reaction (PCR). The principle of this method is outlined by White et al. supra (1989), whereas improved methods are described in Innis et al. [*PCR Protocols: A Guide to Methods and Applications*, Academic Press, Inc. (1990)].

PCR is an in vitro method for producing large amounts of a specific DNA fragment of defined length and sequence from small amounts of a template DNA. PCR is based on the enzymatic amplification of the DNA fragment which is flanked by two oligonucleotide primers that hybridize to opposite strands of the target sequence. The primers are oriented with their 3' ends pointing towards each other. Repeated cycles of heat denaturation of the template, annealing of the primers to their complementary sequences and extension of the annealed primers with a DNA polymerase result in the amplification of the segment defined by the 5' ends of the PCR primers. Since the extension product of each primer can serve as a template for the other, each cycle essentially doubles the amount of the DNA fragment produced in the previous cycle. Since the primers are physically incorporated into the amplified product and mismatches between the 5' end of the primer and the template do not significantly affect the efficiency of the amplification, it is possible to alter the amplified sequence thereby introducing the desired mutation into the amplified DNA. By utilizing the thermostable Taq DNA polymerase isolated from the thermophilic bacteria *Thermus aquaticus*, it has been possible to avoid denaturation of the polymerase which necessitated the addition of enzyme after each heat denaturation step. This development has led to the automation of PCR by a variety of simple temperature-cycling devices. In addition, the specificity of the amplification reaction is increased by allowing the use of higher tem-

peratures for primer annealing and extension. The increased specificity improves the overall yield of amplified products by minimizing the competition from non-target fragments for enzyme and primers.

Design and synthesis of oligonucleotides can be effected as known in the art and described, for example, in Sambrook et al. supra (1989) or in one of the references cited above with respect to site-directed mutagenesis.

As soon as a DNA-sequence coding for a TNF-mutein of the present invention has been created, expression can be effected by the phasmid technology as described above or by use of any suitable pro- or eukaryotic expression system well known in the art [see, e.g., Sambrook et al., supra,].

Expression is effected preferably in prokaryotic cells, for example, in *E. coli*, *Bacillus subtilis* and so on, whereby *E. coli*, specifically *E. coli* K12 strains for example M15 [described as DZ 291 by Villarejo et al. in J. Bacteriol. 120 J, 466-474 (1974)], HB 101 [ATCC No. 33694], WK6 (Stanssens et al. supra) or *E. coli* SG13009 [Gottesman et al., J. Bacteriol. 148, 265-273 (1981)] are preferred. Expression of the muteins of the present invention can also be effected in lower or higher eukaryotic cells, like for example yeast cells (like *Saccharomyces*, *Pichia* etc.), filamentous fungi (like *Aspergillus* etc.) or cell lines (like chinese hamster ovary cell lines etc.), whereby expression in yeast cells is preferred [see Sreekrishna et al., Biochem. 28, 4117-4125, (1989); Hitzeman et al., Nature 293, 717-722 (1981); European Patent Application Publication No. 263 311]. Expression of the TNF-muteins of the present invention may occur in such systems either intracellularly, or, after suitable adaption of the gene, extracellularly (see Lee-mans et al., Gene 85, 99-108, 1989).

Suitable vectors used for expression in *E. coli* are mentioned e.g. by Sambrook et al. [supra] or by Fiers et al. in "Procd. 8th Int. Biotechnology Symposium" [Sot. Franc. de Microbiol., Paris, (Durand et al., eds.), pp. 680-697 (1988)] or and more specifically vectors of the pDS family [Bujard et al., Methods in Enzymology, eds. Wu and Grossmann, Academic Press, Inc. Vol. 155, 416-433 (1987); StOber et al., Immunological Methods, eds. Lefkovits and Pernis, Academic Press, Inc., Vol. IV, 121-152 (1990)] like, for example, pDS56/RBSII,Sph1-TNF α Ser29 or pDS56/RBSII,Sph1-TNF α Trp32 (see Example I) or pDS56/RBSII,Sph1-TNF α Glu31 or pDS56/RBSII,Sph1-TNF α Asn31Thr32 (see Example VII). The transformed *E. coli* strains M15 (pREP4;pDS56/RBSII,Sph1-TNF α Glu31) and M15 (PREP4;pDS56/RBSII,Sph1-TNF α Asn31Thr32) have been deposited under the Budapest Treaty for patent purposes at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH (DSMZ) in Braunschweig, BRD at September 8th, 1991 under accession numbers DSM 6714 and DSM 6715 respectively. These specific pDS56/RBSII-plasmids with their specific regulatable promoter/operator elements and ribosomal binding sites can achieve a high level of expression. Therefore, the plasmids can be maintained in *E. coli* cells only when the activity of the promoter/operator element is repressed by the binding of a lac repressor to the operator. The activity of the promoter can be restored when the culture has reached the desired cell density by addition of isopropyl- β -D-thio-galacto-pyranoside (IPTG), which inactivates the repressor and clears the promoter. Since most of the *E. coli* strains do not provide enough repressor molecules to completely repress the function

of the promoter sequences present in these high copy number plasmids, such *E. coli* strains, *E. coli* M15 or SG13009, have to be first transformed with a plasmid, such as pREP 4, which codes for the lac repressor, before being transformed with the specific pDS56/RBSII-plasmids of the invention which thereafter can be stably maintained in the *E. coli* cells. In addition to coding for the lac repressor, pREP4 also contains a region of the plasmid pACYC184 [Chang and Cohen, J. Bacteriol. 134, 1141-1156 (1978)], which contains all information required for replication and stable transmission to daughter cells. The DNA sequence of pREP4 is set out in FIG. 2b and SEQ ID No: 14 [see also "System for high level production in *E. coli* and rapid purification of recombinant proteins: application to epitope mapping, preparation of antibodies and structure function analysis" by Stüber et al. in Immunological Methods, Vol. IV, pp 121-152, Lefkovits and Pernis (eds.), Academic Press, New York (1990)].

A preferred embodiment of the present invention is an expression vector suitable for producing a human Tumor Necrosis Factor mutein comprising the amino acid sequence set forth in SEQ ID No: 1 wherein SEQ ID No: 1 is changed by deletion, insertion, substitution or combinations thereof, of at least one amino acid so that the mutein shows a significant difference between its binding affinity to the human p75-(Tumor Necrosis Factor)-Receptor and to human p55-Tumor Necrosis Factor)Receptor when the vector is stably transformed or transfected in a prokaryotic or lower eukaryotic host cell.

Another preferred embodiment of the present invention is a vector comprising SEQ ID No: 2 wherein the DNA sequence comprising positions 115 to 591 is changed by deletion, insertion, substitution or combinations thereof.

Transformation of the host cells by vectors as described above may be carried out by any conventional procedure [see, e.g., Sambrook et al. supra]. Where the host cell is a prokaryote, such as *E. coli* for example, competent cells which are capable of DNA uptake are prepared from cells harvested after exponential growth phase and subsequently treated according to the known CaCl₂-method. Transformation can also be performed after forming a protoplast of the host cell or by other methods known in the art and described, for example in Sambrook et al. Therefore a vector, especially for expression in a prokaryotic or lower eukaryotic host cell, comprising a DNA-sequence coding for a TNF-mutein as described above, and a host cell, especially a prokaryotic host cell, for example, *E. coli*, or a lower eukaryotic host cell, transformed by such a vector are also an object of the present invention.

Usually, the host organisms which contain a desired expression vector are grown under conditions which are optimal for their growth. In case of a prokaryotic host at the end of the exponential growth, when the increase in cell number per unit time decreases, the expression of the desired TNF-mutein is induced, that is the DNA coding for the desired TNF-mutein is transcribed and the transcribed mRNA is translated. The induction can be carried out by adding an inducer or a derepressor to the growth medium or by altering a physical parameter, for example a change in temperature. In the expression vectors used in the preferred embodiments of the present invention, the expression is controlled by the lac repressor. By adding IPTG, the

expression control sequence is derepressed and the synthesis of the desired TNF-mutein is thereby induced.

A preferred embodiment of the present invention is a prokaryotic or lower eukaryotic host cell stably transformed or transfected with a vector suitable for producing a human Tumor Necrosis Factor mutein comprising the amino acid sequence set forth in SEQ ID No: 1 wherein SEQ ID No: 1 is changed by deletion, insertion, substitution or combinations thereof, of at least one amino acid so that the mutein shows a significant difference between its binding affinity to the human p75-(Tumor Necrosis Factor)-Receptor and to human p55-(Tumor Necrosis Factor)-Receptor.

Another preferred embodiment of the present invention is a host cell which is stably transformed or transfected with an expression vector comprising positions 115 to 591 of SEQ ID No:2 and in which the DNA sequence is changed by deletion, insertion, substitution or combinations thereof, such that the DNA sequence codes for a human Tumor Necrosis Factor mutein containing at least one amino acid different from SEQ ID No:1.

TNF-muteins of the present invention produced by transformed host cells as stated above can be recovered from the culture medium or after opening the cells with or without extraction by any appropriate method known in protein and peptide chemistry such as, for example, precipitation with ammonium sulfate, dialysis, ultrafiltration, gel filtration or ion-exchange chromatography, gel electrophoresis, isoelectric focusing, affinity chromatography, like immunoaffinity chromatography, HPLC or the like. Specifically preferred methods are precipitation with ammonium sulfate and/or polyethyl-amine, dialysis, affinity chromatography, for example on phenyl-agarose, specifically phenyl-sepharose, or ion-exchange chromatography, specifically on a MONO-Q-and/or MONO-S-matrix (Pharmacia, Uppsala, Sweden) or more specifically preferred are those as described by Tavernier et al. [J. Mol. Biol. 211,493-501 (1990)] and those disclosed in Example I or Example III.

It is therefore also an object of the present invention to provide a process for the preparation of a compound as specified above which process comprises cultivating a transformed host cell as described above in a suitable medium and isolating a mutein from the culture supernatant or the host cell itself, and if desired converting said mutein into a 2.5 pharmaceutically acceptable salt. The compounds whenever prepared according to such a process are also an object of the present invention.

The muteins of the present invention are characterized by showing a significant difference between its binding affinity to the human p75-TNF-R and the human p55-TNF-R. Such property can be determined by any assay known in the art measuring binding affinities. For example, the binding of TNF itself and of the muteins of the present invention can be measured using cells in cell culture which express the two types of TNF-receptors to a different degree, for example Hep-2 cells which exclusively express the human p55-TNF-R and U937 or HL60 cells which express both the human p55-TNF-R and the human p75-TNF-R [see Brockhaus et al., Proc. Nat. Acad. Sci. U.S.A. 87, 3127-3131, (1990); Hohmann et al., J. Biol. Chem. 264, 14927-14934, (1989); Loetscher et al. (1990); Dembic et al. (1990)]. Of course binding affinities can also be determined directly by using purified native or recombinant p55-TNF-R and p75-TNF-R as specifically described in

Example 112, or by using the corresponding soluble analogs of such receptors.

The term "significant difference between its binding affinity to the human p75-Tumor-Necrosis-Factor-Receptor (p75-TNF-R) and to the human p55-Tumor-Necrosis-Factor-Receptor" (p55-TNF-R) refers, the context of the present invention, to a difference in binding affinities to the two types of TNF-receptors which is with respect to the assay system used, significant enough to say that a mutein of the present invention binds preferentially to one of the two TNF-receptors as compared to wild type TNF. The binding affinity for the p55-TNF-R expressed as a K_D -value is measured using Hep-2 cells which only carry that receptor. The binding affinity for the p75-TNF-R is measured using the U937 cells which predominantly, but not exclusively carry the p75 receptor. In terms of the assay system described in Example II (b)(iii)(Table E), the muteins of the present invention differ in their binding affinities to p55-TNF-R and p75-TNF-R by a factor in the range from about 10 to more than 200. A preferential upper limit of this range is 1000 and a most preferential upper limit of this range is 10000. More specifically this term means in the context of the assay-system of Example II (b)(iii) that a K_D -value of a specific TNF-mutein of the present invention is at least a factor of 10 or more, especially preferred at least a factor of 10^2 larger than for TNF- α itself determined by using U937 cells whereby its K_D -value determined by using Hep-2 cells for the same TNF-mutein is not larger than a factor of 2 as for TNF- α itself [for specific data see Table E]. It is however understood that these specific K_D -values are given for illustration and should not be considered as limiting in any manner. Since the purified receptors bind TNF α in the filter binding assays of the present invention with high affinity (see Schönfeld et al., J. Biol. Chem. 266, 3863-3869), namely for the p75-TNF-R with a K_D of 1.0×10^{-10} M and for the p55-TNF-R with a K_D of 16×10^{-11} M the preferential binding of the muteins of the present invention to one of the two TNF-receptors can be also illustrated by a so called selectivity factor "S" which is defined in the following manner:

$$S = \frac{IC_{50} \text{ p75-TNF-R}}{IC_{50} \text{ p55-TNF-R}}$$

"IC₅₀ p75-TNF-R" or "IC₅₀ p55-TNF-R" stands for the concentration of a mutein of the present invention which concentration leads to a 50% inhibition of the binding of TNF α to the p75-TNF-R or p55-TNF-R in a competition assay (such values can be calculated from the data shown in FIG. 1 and FIG. 7; see Table F). Accordingly the muteins of the present invention can show an S-value in the range of 10 to at least 500, preferentially 1000 (see Table G). In addition based on the IC₅₀-values the value of decrease of the affinity of the mutein for both receptors can be calculated (see Table F).

The muteins of the present invention can be characterized by their anti-tumour activity by methods known in the art and described for example in Example IV.

The muteins of the present invention may show considerably reduced cytotoxic activity in standard TNF-assays which are based on murine cell lines, such as L929 (see Table E) or L-M cell lines.

TNF-muteins of the present invention can be used for the treatment of illnesses, for example cancer.

A further object of the present invention is a pharmaceutical composition and a process for its preparation which composition contains one or more compounds of the invention, if desired in combination with additional pharmaceutically active substances with or without nontoxic, inert, therapeutically compatible carrier materials. For this purpose, one or more compounds of the invention, where desired or required in combination with other pharmaceutically active substances, can be processed in a known manner with the usually used solid or liquid carrier materials. The dosage of such preparations can be effected having regard to the usual criteria in analogy to already used preparations of similar activity and structure.

A preferred embodiment of the present invention is a pharmaceutical composition comprising an effective amount of a human Tumor Necrosis Factor mutein comprising SEQ ID No: 1 in which SEQ ID No: 1 is changed by deletion, insertion, substitution or combinations thereof, of at least one amino acid so that the mutein shows a significant difference between its binding affinity to the human p75-(Tumor Necrosis Factor)-Receptor and to human p55-(Tumor Necrosis Factor)-Receptor or a pharmaceutically acceptable salt thereof and an inert carrier.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1a is a graph showing the results of a competitive binding assay between ^{125}I -TNF and Trp 32 -TNF, Ser 29 -TNF and wild type-TNF for the p75 receptor.

FIG. 1b is a graph showing the results of a competitive binding assay between ^{125}I -TNF and Trp 32 -TNF, Ser 29 -TNF and wild type-TNF for the p55 receptor.

FIG. 2a is a schematic depiction of plasmid pREP4.

FIGS. 2b, 2c, and 2d are the nucleotide sequence of plasmid pREP4.

FIG. 3a is a schematic depiction of plasmid pDS56/RBSII,SphI-TNF α .

FIGS. 3b, 3c, and 3d are the nucleotide sequence of plasmid pDS56/RBSII,SphI-TNF α .

FIG. 4 is a graph showing the results of an assay measuring the antitumor effect of interferon-gamma and TNF, alone or in combination.

FIG. 5 is a graph showing the results of an assay measuring the antitumor effect of interferon-gamma and TNF, alone or in combination and Trp 32 -TNF alone or in combination with interferon-gamma.

FIG. 6 is a series of graphs showing the results of a competitive binding assay between ^{125}I -TNF and various muteins for the p75 receptor and the p55 receptor.

FIG. 7 is a series of graphs showing the results of a competitive binding assay between ^{125}I -TNF and various muteins for the p75 receptor and the p55 receptor.

FIG. 8 is a schematic representation of mutagenesis of the TNF β gene using PCR with primers containing the altered codons.

DETAILED DESCRIPTION OF THE INVENTION

After the invention has been described in general hereinbefore, the following Examples are intended to illustrate details of the invention, without thereby limiting it in any manner.

Example I

A. Preparation of Ser 29 -TNF α and Trp 32 -TNF α

(1) Construction of a mutagenesis vector

From the human TNF expression plasmid pDS56/RBSII,SphI-TNF α (see FIG. 3a: The expression plasmid contain the regulatable promoter/operator element N25OPSN25OP29 (), the synthetic ribosomal binding site RBSII (), genes () ribosomal binding S for β -actinase (bla), chloramphenicol acetyltransferase (cat), and transcriptional terminators () to of phage lambda (to) and T1 of rrnB operon of *E. coli* (T1), and the replication region of plasmid pBR322 (repl.). The coding region under control of N25OPSN25OP29 and RBSII is indicated by an arrow; for complete nucleotide sequence of the plasmid see [SEQ ID No: 2] FIG. 3b/1-3b/3 given by the one letter standard abbreviations for nucleotides), an EcoRI-HindIII fragment was isolated, containing the ribosome binding site RBSII, the mature TNF α coding sequence and a 130 bp 3' non-translated sequence. This fragment was cloned into the EcoRIHindIII opened pMac phasmids (Stanssens et al., supra), resulting in the constructions pMa/RBSII,SphI-TNF α and pMc/RBSII,SphI-TNF α .

(2) Isolation of single-stranded (ss)DNA

The pMa/RBSII,SphI-TNF α phasmid was transformed to *E. coli* WK6 (Stanssens et al., supra). One colony was picked up and cultured in 5 ml LB medium (Sambrook et al., supra 1989) with carbenicillin (50 $\mu\text{g}/\text{ml}$) at 37° C., overnight. 1 ml of this confluent culture was used to inoculate 200 ml LB containing carbenicillin. When the absorbance (650 nm) reached a value of 0.1, the culture was infected with M13K07 helper phage (Stanssens et al., (1989) at a m.o.i. of about 20 and further incubated overnight at 37° C. Then, the cells were spun down (10 min, 10,000 rpm) and the supernatant was transferred into another tube. 50 ml PEG-solution (20% polyethylene glycol 6000; 2.5 M NaCl) was added and the mixture was kept on ice for one hour to precipitate the phages. After centrifugation (10 min; 8000 rpm), the supernatant was removed and the tube was dried on paper towels for 10 min. The phage pellet was resuspended in 6 ml TE buffer (10 mM Tris-HCl, 0.1 mM EDTA, pH8). A first extraction was performed with 6 ml TE-saturated phenol, followed by vortexing for 3 min. After centrifugation (3 min) in an Eppendorf centrifuge, the aqueous phase was transferred to a fresh tube and a second extraction was carried out with chloroform:isoamylalcohol (24:1) in the same way as described. The single stranded DNA could be precipitated by adding 1/10 volume of 5M NaClO $_4$ and 1 volume of isopropanol (-20° C., 2 hours). This ssDNA was pelleted by centrifugation for 20 min in an Eppendorf centrifuge. The pellet was dried and dissolved in 500 μl TE buffer as a control, 5 μl of this mixture was run on an agarose gel, containing 1 $\mu\text{g}/\text{ml}$ ethidium bromide. Usually, the ratio of pMa/RBSII,SphI-TNF α ssDNA (= (+)strand) over helper phage ssDNA was between 2:1 and 20:1. The amount of total ssDNA was estimated to be at least 200 ng/ μl .

(3) Construction of a gap-duplex

From the phasmid pMc, the EcoRI-HindIII large fragment was isolated and used for hybridization to the pMa/RBSII, SphI-TNF α (+)strand. In a typical experiment, 15 μl ssDNA ($\pm 3 \mu\text{g}$), 15 μl of the double stranded, linear fragment ($\pm 1.5 \mu\text{g}$), 10 ml hybridization buffer (1.5 M KCl; 100 mM Tris-HCl, pH 7.5) and 40 μl H $_2$ O were mixed and incubated at 100° C. for 4 min, 65° C. for 8 min and room temperature for 15 min. An aliquot (10 ml) was electrophoresed on an agarose gel containing ethidium bromide, to check the forma-

tion of gap duplex DNA and, if so, to estimate its quantity (this usually amounted to 50 ng/10 ml hybridization mixture).

(4) Annealing of the mutant oligonucleotide and fill-in of the gap duplex

Oligonucleotides were synthesized containing the mutated codon and destroying or creating a restriction site in the TNF gene. The oligonucleotides 5'CCGGCGGTTGGACCACTGGAGC3'[SEQ ID No:15] and 5'CATTGGCCCAGCGGTTTCAG3'[SEQ ID No: 16] (mutated bases underlined) were used to create the Ser²⁹ and Trp³² mutations, respectively. After enzymatic phosphorylation, about 8 pmol was added to 40 ng of gapduplex. H₂O was added to a final volume of 10 ml. This mixture was heated to 65° C. for 5 min and allowed to cool to room temperature. Subsequently, 18 ml H₂O, 4 μl fill-in buffer 10 (625 mM KCl, 275 mM TrisHCl, 150 mM MgCl₂, 20 mM DTT pH 7.5), 2 μl ATP 1 mM, 4 μl of the four dNTP's 1 mM, 1 μl ligase and 1 ml Klenow polymerase were added and the mixture was incubated at room temperature for 45 min.

(5) Transformation to *E. coli* WK6 mutS and *E. coli* WK6

We used 10 IL1 of the filled-in gap duplex DNA to transform (Sambrook et al., 1989) *E. coli* WK6 mutS (Stanssens et al., supra). From this mixture (1.2 ml), 100 ml was plated out on agar plates containing 25 μg/ml chloramphenicol to check transformation efficiency. The remainder was used to inoculate 20 ml LB+chloramphenicol and further grown overnight at 25° C. . A small-scale plasmid DNA preparation [Birnboim, H. C. and Doly, J., Nucleic Acids Res., 7, 1513, (1979)] of this culture (not yet grown to confluency) resulted in a mixed phasmid population that could be transformed to *E. coli* WK6. Again, 100 μl transformation mixture was plated out on agar plates containing chloramphenicol.

(6) Screening by colony hybridization

About 100 colonies, resulting from the transformation to *E. coli* WK6, were streaked on a nylon filter (PALL, Glen Cove, New York) and incubated overnight at 37° C. . The filter was transferred (face up) to Whatmann 3MM papers which were soaked in 0.5 M NaOH (3 min). Neutralization was done by transfer to Whatmann 3MM sheets soaked in 1M Tris-HCl pH 7.4 (twice for 1 min) and 2XSSC (20xSSC=3M NaCl; 0.3M Na citrate, pH7) (5 min). After drying, the filter was baked at 80° C. between sheets of 3MM paper. Subsequently, the filter was prewetted in 6xSSC (5 min) and prehybridized at 67° C. for 5 min in 10x Denhardt solution (2% (w/v) Fico11 (400,000 MV), 2% (w/v) Polyvinylpyrrolidone (44,000 MW), 2% (w/v) Bovine Serum Albumin), 6xSSC buffer and 0.2% SDS. After rinsing in 6xSSC buffer, the filter was placed in a Petri dish containing 4 ml 6xSSC and 60 pmol of the ³²P-labeled mutant oligonucleotide for 1 hour at room temperature, and rinsed in 100 ml 6xSSC. The filter was covered with Saran® wrap or suitable plastic film and autoradiographed on preflashed films (Fuji) at -70° C. for 1 hour. Subsequently, the filter was again washed in 6xSSC buffer at increasing temperatures (varying between 51° C. and 75° C. , according to the length of the probe and its amount of G and C residues), followed each time by an autoradiography, as described above. For instance, a wash at 64° C. could clearly distinguish the Ser²⁹ mutants from the wild-type colonies, while the

Trp³² mutants were detected after two subsequent washes at 62° C. and 63° C. , respectively.

(7) Restriction fragment analysis

Because the Ser²⁹ mutation created an Ava2 restriction site and Arg32 destroyed the NciI restriction site, both corresponding endonucleases could be used for restriction fragment analysis to check once again the presence of the mutation. The colonies were picked up and grown to confluency in 5 ml LB medium containing chloramphenicol. From these cultures, plasmid DNA was prepared, digested with the appropriate restriction endonucleases and electrophoresed on agarose gels, according to classical procedures (Sambrook et al., 1989).

(8) Subcloning to a bacterial expression vector

Transfer of the mutated TNF gene to an expression vector was carried out exactly the opposite way as the construction of the mutagenesis vector. The phasmid pMc/RBSII,Sph1-TNFα Ser29 or pMc/RBSII,Sph1-TNFα Trp³² was digested with EcoR1-HindIII and the small fragment was inserted into the EcoR1-HindIII opened pDS56/RBSII,Sph1-TNFα vector generating plasmids pDS56/RBSII,Sph1-TNFα Ser29 and pDS56/RBSII,Sph1-TNFα Trp32 and transformed into *E. coli* M15 cells already containing plasmid pREP4 [SEQ ID No: 14] (encoding the lac repressor; see FIGS. 2a and 2b/1-2b/3 for a complete nucleotide sequence of the plasmid given by the one letter standard abbreviations for nucleotides) by standard methods. Such cultures of transformed *E. coli* M15 were grown at 37° C. in LB medium (10 g bacto tryptone, 5 g yeast extract, 5 g NaCl per liter) containing 100 mg/l ampicillin and 25 mg/l kanamycin. At an optical density at 600 nm of about 0.7 to 1.0 units, IPTG s5 was added to a final concentration of 2mM. After an additional 2.5 to 5 h at 37° C. the cells were harvested by centrifugation and the TNF muteins were purified according to Tavernier et al. [J. Mol. Biol. 211, 493-501, (1990)]. The transformed *E. coli* strains M15 (pREP4;pDS56/RBSII,Sph1-TNFα Ser29) and M15(pREP4;pDS56/RBSII,Sph1-TNFα Trp32) have been deposited under the Budapest Treaty for patent purposes at the Deutsche Sammlung von Mikroorganismen und Zellkulturen GmbH(DSM) in Braunschweig, BRD at November 19th, 1990 under accession numbers DSM 6240 and DSM 6241 respectively.

Example II

A. Characterization of Ser²⁹-TNFα and Trp³²-TNFα

1) Differential binding and biological activity on Hep2 and U937 cells

(a) Cell culture

Hep-2 [ATCC No. CCL 23], U937 [ATCC No. CRL 1593] and RAJI [ATCC No. CCL 86] cells were grown in RPMI 1640 medium, supplemented with 10% (v/v) inactivated fetal calf serum, L-glutamine (2 mM), sodium pyruvate (1mM), 2-mercaptoethanol (5×10⁻⁵M), 1% of a 100x mixture of non-essential amino acids [Gibco Laboratories, Paisley, GB] and gentamycine (25 mg/ml). The non-adherent cells (U937 and RAJI) were harvested after reaching a density of 1×10⁶ cells/ml. For binding experiments, the adherent Hep-2 cells were grown to confluency, trypsinized, collected and seeded in large Petri dishes (150 cm²) at a density of 2.5×10⁶ cells/ml. Subsequently, the dishes were placed in a CO₂-incubator overnight. Because Hep-2 cells are not strongly adherent, the cells could be harvested the same way as the non-adherent cells. Dulbecco's medium,

supplemented with 10% inactivated newborn calf serum was used for L929 cell growth.

(b). Determination of the specific activities on L929, Hep-2 and U937 cells.

The amount of protein was determined by the Biorad (Richmond, Calif., USA) protein dye reagent according to the instructions of the manufacturer. The purity of the TNF muteins was determined by SDS-PAGE.

The cytotoxic activity on mouse L929 cells was determined using the standard L929 assay (Ruff and Gifford in "Lymphokines", ed. by E. Pick, Vol. 2, 235-275, Academic Press, 1981, Orlando, USA). The cytotoxicity assay on Hep-2 cells was performed the same way as the L929 assay with the only exception that cycloheximide (50 ILg/ml) was added instead of actinomycin D.

(c.) Receptor binding assay.

(i.) -Iodination of TNF- α and Trp³²-TNF

5 μ g Iodogen (Pierce, USA) was dissolved in 10 μ l chloroform and dried under a nitrogen stream in a small glass tube. To this, 10 μ l Na¹²⁵I (Amersham, 100 mCi/ml in 0.1 M borate buffer, pH 8) was added and kept for 15 min. on ice. This solution was quickly pipetted to an Eppendorf tube, containing 5 μ g TNF- α [Penica et al., s.a.] or 3.2 μ g Of Trp³²-TNF in 10 μ l phosphate buffer pH 7. Again the reaction was kept for 15 min on ice. To separate the iodinated TNF- α from the Na¹²⁵I, a PD-10 gelfiltration column (Pharmacia) was first equilibrated with 0.1 M phosphate buffer + 0.25% gelatin and prerun with 1 μ g TNF- α or Trp³²-TNF, depending on the iodinated TNF species. Subsequently, the reaction mixture was loaded onto the column, and fractions of about 400 μ l were collected from which 2 μ l aliquots were counted in a γ -counter (LKB 1275 Minigamma, Pharmacia LKB, Uppsala, Sweden). A specific radioactivity of 10-75 and 80 μ Ci/mg was obtained for TNF- α and Trp³²-TNF, respectively.

(ii.)-Determination of the K_D-value of labeled TNF- α and Trp³²-TNF by Scatchard analysis

A dilution series in multiples of 2 in the range of 12.8nM to 0.006nM of the labeled TNF- α or Trp³²-TNF was made up in a microtiterplate. Each dilution was made in triplicate. Non-specific binding was measured by the same setup, wherein each point contained a 100 fold excess of unlabeled TNF (1.28 μ M to 0.6nM). To each well, approximately 2 \times 10⁶ cells (U937, Hep-2 or RAJI) were added. The reaction was performed in 0.2 ml tissue culture medium, containing 0.1% NaN₃ for 2-3 hours at 4° C. After this, samples were transferred from the microtiterplates to small plastic tubes (Mircronic systems), already containing 300 μ l phthalate oil (dinonylphthalate 33%, dibutylphthalate 66% (v/v)). The tubes were centrifuged in a microfuge (Eppendorf) for 10 min. to spin down the cells, thereby separating them from the supernatant, using the phthalate oil as a separation medium. After inversion of the tubes, the cell pellet (now on top) could easily be isolated by melting off the top of the tubes with a hot scalpel. The amount of radioactivity, bound on the cells, was measured by counting in a γ -counter. From these data, a Scatchard plot and, subsequently, the dissociation constant K_D was determined using the equilibrium binding type "HOT" in the EBDNLIGAND programm [Mc. Pher-son et al., J. Pharmacol. Methods 14, 213-228, (1985)].

(iii)-Determination of the K_D of mutant TNF [Ser²⁹-TNF- α and Trp³²-(iii.)

TNF- α] by competition analysis

The Scatchard data showed that a concentration of 0.4 nM radiolabeled TNF- α was high enough to show a clearly detectable signal and fell within the linear part of the saturation curves. This concentration, however, was also low enough to allow addition up to a 5000 fold excess of cold mutant TNF (2 μ M), necessary to perform a competition experiment in which ¹²⁵I-wild type TNF is the primary ligand and cold mutant the competitor.

A ten well dilution series of unlabeled mutant TNF (2 mM to 0.004 μ M) in concentration steps in multiples of 2 was set up in a microtiterplate. The two remaining wells contained no unlabeled TNF (total binding) and a 5000 fold excess of the wild-type, unlabeled TNF (background), respectively. To all wells, 0.4 nM of radiolabeled TNF- α (10-75 μ Ci/ μ g) was added. After addition of 2 \times 10⁶ cells, the total volume was 0.2 ml/well. The medium of incubation, reaction conditions and isolation of the cells were exactly the same as described above for the Scatchard analysis experiments. Each point was measured in triplicate and the dissociation experiments were done twice, the average of the two K_D's being indicated in Table E. Using the "DRUG" method of the EBDA/LIGAND program, competition curves were plotted and the K_D of the muteins was calculated. The following experimental data were used for such calculations:

1. Labeling of hTNF

first labeling (=batch 1):	1.2 \times 10 ⁸ dpm/5 μ g
	= 3.7 \times 10 ⁵ dpm/pmol
	= \pm 10 μ Ci/ μ g
second labeling (=batch 2):	5.3 \times 10 ⁸ dpm/3.2 μ g
	= 1.9 \times 10 ⁶ dpm/pmol
	= \pm 75 Ci/ μ g

2. Determination of the K_D of wild-type TNF

We measured the K_D of ¹²⁵I-TNF (batch 1) on Hep-2 and U937 cells by Scatchard analysis.

Hep-2: K_D=9.17 \times 10⁻¹⁰

U937: K_D=2.5 \times 10⁻¹⁰

3. Competition experiments

All displacement experiments were carried out, using ¹²⁵I-TNF (batch 1) as the primary ligand, except experiment B.3 (table B, 3.), where ¹²⁵I-TNF (batch 2) was used.

In each experiment, the binding at each concentration was measured in triplicate and only the averages are shown in the following tables (A-D).

From each experiment shown in these tables, the K_D value was calculated using the programm of Mc. Pher-son et al. (1985). The average of the K_D determinations (2 experiments for Ser²⁹-TNF α on Hep-2 cells and on U937 cells, two experiments for Trp³²-TNF α on Hep-2 cells and three on U937 cells) are shown in table E.

TABLE A

Competition with Ser ²⁹ -TNF α on U937 cells.	
Mean dpm	concentration of mutant [mol]
1. 2120	0
1869	1 \times 10 ⁻⁹
1779	2 \times 10 ⁻⁹
1719	4 \times 10 ⁻⁹
1708	8 \times 10 ⁻⁹
1575	1.6 \times 10 ⁻⁸

TABLE A-continued

Competition with Ser ²⁹ -TNF α on U937 cells.		
Mean dpm	concentration of mutant [mol]	
		5
1415	3.2×10^{-8}	
1320	6.4×10^{-8}	
1200	1.25×10^{-7}	
983	2.5×10^{-7}	
949	5×10^{-7}	
632	1×10^{-6}	
533	2×10^{-6}	10
Background: 299		
2. 1014	0	
635	4×10^{-9}	
603	8×10^{-9}	
641	1.5×10^{-8}	
572	3×10^{-8}	15
489	6×10^{-8}	
413	1.2×10^{-7}	
380	2.5×10^{-7}	
319	5×10^{-7}	
263	1×10^{-6}	
238	2×10^{-6}	20
Background: 205		

TABLE B

Competition with Trp ³² -TNF α on U937 cells		
		25
1. 2120	0	
1917	1×10^{-9}	
1698	2×10^{-9}	
1655	4×10^{-9}	
1585	8×10^{-9}	
1488	1.5×10^{-8}	30
1377	3×10^{-8}	
1333	6×10^{-8}	
1166	1.25×10^{-7}	
1026	2.5×10^{-7}	
953	5×10^{-7}	
777	1×10^{-6}	35
628	2×10^{-6}	
Background: 299		
2. 1047	0	
653	4×10^9	
629	8×10^{-9}	
636	1.5×10^{-8}	
585	3×10^{-8}	40
546	6×10^{-8}	
508	1.2×10^{-7}	
479	2.5×10^{-7}	
422	5×10^{-7}	
357	1.10^{-6}	
294	2×10^{-6}	45
Background: 214		
3. 8340	0	
(carried out with ¹²⁵ I-TNF, batch 2)	4×10^{-9}	
4759	8×10^{-9}	
4041	1.5×10^{-8}	
3620	3×10^{-8}	
3275	6×10^{-8}	
3034	1.25×10^{-7}	
2387	2.5×10^{-7}	
1981	5×10^{-7}	
1472	1×10^{-6}	
1192	2×10^{-6}	55
814		
Background: 307		

TABLE C

Competition with Ser ²⁹ -TNF α on Hep-2 cells		
		60
1. 938	0	
799	1×10^{-9}	
677	2×10^{-9}	
564	4×10^{-9}	
510	8×10^{-9}	
451	1.6×10^{-8}	
442	3.2×10^{-8}	
446	6.4×10^{-8}	
379	1.25×10^{-7}	65

TABLE C-continued

Competition with Ser ²⁹ -TNF α on Hep-2 cells		
		10
374	2.5×10^{-7}	
437	5×10^{-7}	
359	1×10^{-6}	
383	2×10^{-6}	
Background: 353		
2. 457	0	
273	4×10^{-9}	
240	8×10^{-9}	
253	1.5×10^{-8}	
235	3×10^{-8}	
207	6×10^{-8}	
239	1.2×10^{-7}	
215	2.5×10^{-7}	
211	5×10^{-7}	
193	1×10^{-6}	
238	2×10^{-6}	
Background: 215		

TABLE D

Competition with Trp ³² -TNF α on Hep-2 cells		
		25
1. 938	0	
742	1×10^{-9}	
608	2×10^{-9}	
537	4×10^{-9}	
547	8×10^{-9}	
397	1.6×10^{-8}	
394	3.2×10^{-8}	
405	6.4×10^{-8}	
395	1.25×10^{-7}	
388	2.5×10^{-7}	
379	5×10^{-7}	
353	1×10^{-6}	
386	2×10^{-6}	
Background: 353		
2. 445	0	
298	4×10^{-9}	
222	8×10^{-9}	
256	1.5×10^{-8}	
202	3×10^{-8}	
227	6×10^{-8}	
210	1.2×10^{-7}	
221	2.5×10^{-7}	
197	5×10^{-7}	
231	1×10^{-6}	
202	2×10^{-6}	
Background: 203		

TABLE E

	Hep-2		L929	
	affinity (K _D)	specific activity (U/mg)	U937 affinity (K _D)	specific activity (U/mg)
50 TNF- α	9.17×10^{-10} (*)	2.9×10^7 (100%)	2.5×10^{-10} (*)	2×10^7 (100%)
Ser ²⁹ -TNF- α	1.06×10^{-9} (86.5%)	9.3×10^6 (32%)	5.07×10^{-8} (0.49%)	10^5 (0.5%)
Trp ³² -TNF- α	1.06×10^{-9} (86.5%)	4.5×10^7 (155%)	3.53×10^{-8} (0.71%)	6.4×10^4 (0.32%)

K_D values indicated by an asterisk (*) were obtained by Scatchard analysis. All other K_D values were determined by competition analysis. Relative values (in percentage to TNF- α) are indicated between brackets.

60 It can be seen that the binding constant (K_D) of Ser²⁹-TNF-oc and Trp³²-TNF- α determined with Hep-2 cells (which only carry the p55-TNF-R) are almost the same as TNF- α . Also the biological activity (specific activity) on these cells is largely retained (note that the accuracy of this assay is only a factor of 3). Strikingly, the binding affinity (measured in the competition assay) of Ser²⁹-TNF- α and Trp³²-TNF- α to the U937 cells, which predominantly but not exclusively, carry the

high affinity receptor p75-TNF-R, has been largely lost (increase in K_D -value by a factor of more than 100). Thus, the binding affinity of the Ser²⁹-TNF- α for p75-TNF-R has been reduced approximately 50 fold to about 2% of its binding affinity to p55-TNF-R. The binding affinity of Trp³²-TNF- α for p75-TNF-R has been reduced approximately 33 fold to about 3% of its binding affinity to p55-TNF-R. It may also be noted that the biological activity of Ser²⁹-TNF- α and Trp³²-TNF- α , determined in the standard assay based on L929-cells, has been largely lost (decrease by a factor more than 100).

Differential binding to the human p75-TNF-R and the human p55-TNF-R

Competition of human ¹²⁵I-TNF- α binding by Trp³²- and Ser²⁹-TNF- α and human TNF- α to TNF-receptors purified from HL60 cells was determined as follows. 2 β l aliquots of the native p55-TNF-R and the p75-TNF-R purified as described in European Patent Application No. 90116707.2 dissolved at a concentration of about 0.3 μ g/ml in 20 mM Hepes, 50 mM Tris, 50 mM NaCl, 1 mM EDTA, 0.1% octylglucoside, 0.1 mg/ml BSA, pH 8.0, were spotted onto prewetted nitrocellulose filters in triplicate. The filters were blocked with blocking buffer (50 mM Tris, 140 mM NaCl, 5 mM EDTA, 0.02% NaN₃, 1% defatted milk powder) for 1.5 hours at room temperature. After washing with PBS the filters were incubated with 10 ng/ml ¹²⁵I-TNF α and varying concentrations of Trp³²-or Ser²⁹-TNF α , or TNF α overnight at 4° C. . The filters were washed with blocking buffer (2x for 5 min.) and with H₂O (1x for 5 min.), air dried, and counted in a γ -counter. Results are given in FIGS. 1a and b, whereby FIG. 1 shows binding of TNF α (open rectangle), Ser²⁹-TNF α (filled circles) and Trp³²-TNF α (filled rectangle) to human p75TNF-R in case of FIG. 1a to human p75-TNF-R and in case of FIG. 1b to human p55-TNF-R. Based on the data shown in FIG. 1 and in addition those of FIG. 7 the IC50-values were calculated and are listed for Ser²⁹- and Trp³²-TNF α in Table F. Values for the decrease in affinity for these muteins on both receptors with respect to TNF α are also given in Table F. Values for "S", the selectivity factor, based on IC50 values given in Table F and calculated from FIG. 7 are shown in Table G.

TABLE F

Receptor	Competitor	IC50 μ g/ml	Decrease in Affinity
p75-TNF-R	TNF α	0.010	—
	Ser ²⁹ -TNF α	2.5	250
	Trp ³² -TNF α	5	500
p55-TNF-R	TNF α	0.011	—
	Ser ²⁹ -TNF α	0.09	8.2
	Trp ³² -TNF α	0.017	1.5

TABLE G

Mutein	S = $\frac{\text{IC50 p75-TNF-R}}{\text{IC50 p55-TNF-R}}$
TNF α	1
Ser ²⁹ -TNF α	28
Trp ³² -TNF α	294
Gly ²⁹ -TNF α	80
Tyr ²⁹ -TNF α	110
Tyr ³² -TNF α	90
Ser ²⁹ -Trp ³² -TNF α	450

Example III

Purification of Trp³²-TNF α

Transformed cells obtained according to Example I were processed in the following manner:

- Opening by French press, addition of polyethyleneimine until a final concentration of 0.4%, pH 7.6; and removal of precipitate.
- Ammonium sulphate precipitation at pH 7.2; fraction 30-70%
- Dialysis against 25% ammonium sulphate in 10 mM Tris, pH 6.8
- Phenyl-Sepharose column CL-4B (35 \times 250 mm) Load in 25% ammonium sulphate - 10 mM Tris, pH 6.8

Elution: gradient 25% ammonium sulphate-Tris buffer to 20 mM ethanolamine, pH 9 (2 times 150 ml).

- Column Mono Q (HR 16/10). Load: in 20 mM ethanolamine, pH 9. Elution: gradient (2 times 300 ml) in the same buffer, from 0 to 1 M sodium chloride (Pharmacia, FPLC). Active fractions dialysed versus 0.01 M phosphate buffer pH 7
 - Column of Heparin Sepharose CL-6B (30 \times 80 mm) Load in 0.01 M phosphate buffer pH 7. Elute with a gradient in the same buffer from 0 to 1 M sodium chloride
 - Active fractions were concentrated on Amicon (micro-ultrafiltration system 8 MC; membrane $\text{\textcircled{C}}$ 25 mm; diaflo 10 YM10 - 25 mm) and separately loaded on a gelfiltration column (Ultrapac TSK G-2000 SWG; 21.5 \times 600 mm), equilibrated in 0.01 M phosphate pH 7 and 0.9% sodium chloride
- LPS (determined by test kit of Kabivitrum):
Most active fraction contained 5 mg/ml Trp³²-TNF α ; endotoxin content: 26 E.U./mg
The last active fraction contained 1.8 mg/ml TNF and 47 E.U./mg protein.

Example IV.

Anti-tumour effect of hTNF α and hIFN γ on subcutaneous HT-29 tumours in nude mice.

- 5 \times 10⁶ HT-29 human colon adenocarcinoma cells [ATCC HTB38] were subcutaneously injected in nude mice. Groups consisted of 5 mice. The treatment comprises daily perilesional injections during 6 days per week, followed by 1 day without treatment. Results are given in FIG. 4 whereby "PBS" refers to phosphate buffered saline as known in the art. The single arrow indicates the start of the treatment with 5 μ g hTNF α or 5000 IU human Interferon γ (hIFN γ) or both. The double arrow indicates the time that these doses were doubled and the crossed arrow indicates the end of the treatment.

- Comparison of the anti-tumour potential of hTNF α and Trp³²-TNF α

- 5 \times 10⁶ HT-29 human colon adenocarcinoma cells were subcutaneously injected in nude mice. Groups consisted of 5 mice. The treatment started on day 6 following inoculation and comprises daily perilesional injections during 6 days per week. Tumour volume was estimated every 3 or 4 days by measuring the larger (a) and the smaller (b) diameter and calculating the $a \times b^2 \times 0.4$ according to Attia and Weiss as known in the art. Results are given in FIG. 5 whereby the arrow indicates the start of the treatment and open triangles with tip down refers to 10⁴ IU of hIFN γ and 10 μ g hTNF α , filled triangles with tip down refer to 10⁴ IU of hIFN γ and 10 μ g Trp³²-TNF, filled rectangles refer to 10 μ g Trp³²-TNF α , open rectangles refer to 10 μ g hTNF α , open triangles refer to phosphate buffered

saline and filled circles refer to 10^4 IU of hIFN γ . In vitro, there is no difference in cytotoxicity for Hep or HT-29 cells between hTNF α and Trp³²-TNF α .

EXAMPLE V

Preparation of Ser²⁹-Trp³²-TNF α

Ser²⁹-Trp³²-TNF α was prepared as described in Example I with the following exceptions:

1. The oligonucleotide used, contains the following sequence [SEQ ID No: 17] (mutated bases underlined):

5'GGGCATTGGCCCAGCGGTTGGACCACT-GGAGC3'

2. An Nci I site was destroyed while an Ava II site was created, allowing for check of the presence of the mutation by restriction fragment analysis. No hybridization analysis was performed. 6 clones resulting from the WK6 transformation were grown up and DNA was prepared and analysed as described in Example I, 3 of the 6 clones contained the mutation.

This DNA sequence was subcloned into the pDS56 expression vector, generating the plasmid pDS56/RBSII,SphI-TNF α Ser²⁹Trp³², and transformed to the *E. coli* M15 strain. Expression and purification was performed as described in Example 1.

EXAMPLE VI

Preparation of Gly²⁹-TNF α , Tyr²⁹-TNF α and Tyr³²-TNF α

Gly²⁹-TNF α , Tyr²⁹-TNF α and Tyr³²-TNF α were prepared as described in Example I with the following exception. Oligonucleotides were used, containing a fully degenerated codon at position 29 or 32, resulting in a random insertion of all twenty amino acids at one of the two positions. The sequence of these oligonucleotides are as follows:

Position 29 [SEQ ID No: 18]

5'CCACGCCATTTCGCGAGGAGG-GCATIGGCCCGGCGGTNNNCCACT-GGAGC3'

Position 32 [SEQ ID No:19]:

5'CCACGCCATTTCGCGAGGAGG-GCATTGGCNNNGCGGTTTCAGCC3'

where N=A, C, G or T and mutated bases are underlined.

Together with the mutation, also a unique Nru-I site is introduced. Thus, instead of directly transforming the phasmid-pool, isolated from one of the WK6 muts strain, this DNA was first digested with Nru-I, the linear band eluted from the agarose gel, ligated and transformed to the SURE strain (Stratagene, La Jolla, Calif., USA). In this way, one can select only for phasmids, containing the mutations. 168 colonies obtained were inoculated in microtiterplates, grown to confluency and their lysates tested for biological activity towards Hep-2 cells in a manner as described in Example IIa and for differential binding as described in Example IIb or Example VIII. On the basis of the biological activity on the one side and differential binding as determined according to Example IIb or Example VIII colonies were selected and further characterized by DNA sequence analysis of corresponding inserts as known in the art. DNase-sequences coding for Gly²⁹-TNF α , Tyr²⁹-TNF α and Tyr³²-TNF α were isolated from corresponding colonies and cloned in bacterial expression vectors as described in Example I. Mutins expressed were purified

to more than 95% homogeneity by means of a MONO-Q ion exchange chromatography step.

EXAMPLE VII

5 Preparation of Glu³¹-TNF α and Asn³¹-Thr³²-TNF α

Mutagenesis of the TNF α gene using PCR

Three PCR reactions were performed with plasmid pDS56/RBSII,SphI-TNF α [SEQ ID No:2][FIG. 3] as the template DNA using a Perkin-Elmer Cetus GeneAmp™ DNA Amplification Reagent Kit with AmpliTaq™ Recombinant Taq DNA Polymerase (Perkin Elmer Cetus, Vaterstetten, BRD) [see FIG. 8]. In reaction I primers 17/F [SEQ ID No: 20](5'-GGCGTAT-CACGAGGCCCTTTCG3'; primer 17/F comprises nucleotides 3949-3970 of plasmid pDS56/RBSII,SphI-TNF α) and 21/M5 [SEQ ID No: 22] (5-ATTGGCCCGCTCGTTCAGCCACT-GGAGCTGCCCTC-3'; primer 21/M5 comprises nucleotides which are complementary to nucleotides 219-184 of plasmid pDS56/RBSII,SphI-TNF α , mutated bases are underlined) were used, reaction II contained primers 17/F and 21/M6 [SEQ ID No:23] (5'-ATTGGCAGTGTTCAGCCACTGGAG-CTGCCCTC-3'; primer 21/M6 comprises nucleotides which are complementary to nucleotides 219-184 of plasmid pDS56/RBSII,SphI-TNF α , mutated bases are underlined), and reaction III contained primers 21/MR [SEQ ID No: 24] (5'-GCCCTCCTGGCCAATGGCGTGG-3'; primer 21/MR comprises nucleotides 220-241 of plasmid pDS56/RBSII,SphI-TNF α) and 17/O [SEQ ID No: 21] (5'-CATTACTGGATCTATCAACAGG-3'; primer 17/O comprises nucleotides which are complementary to nucleotides 748-727 of plasmid pDS56/RBSII,SphI-TNF α). Therefore 10 μ l template DNA (10 g), 5 μ l each of the two primers (100 pmole each), 16 μ l dNTP's mix (1.25 mM of dATP, dGTP, dCTP, and dTTP), 10 μ l 10x reaction buffer (100 mM Tris-HCl pH8.3, 500 mM KCL, 15 mM MgCl₂ and 0.1% gelatin), 1 μ l (5 units) AmpliTaq™ DNA polymerase and 53 μ l H₂O were mixed in an Eppendorf tube and overlaid with 80 μ l mineral oil (Perkin-Elmer Cetus). The tubes were transferred to a DNA thermal cycler (TRIO-Thermoblock, Biometra) and kept for 1 min at 94° C., before 35 cycles of melting the DNA (1 min at 94° C.), annealing the primers (1 min at 50° C.), and extending the primers (3 min at 72° C.) were performed. After additional 2 min at 72° C., the reactions were cooled to room temperature and extracted with chloroform. The DNA present in the aqueous phase was precipitated with ethanol and subjected to electrophoresis in a 6 polyacrylamide gel [Sambrook et al., 1989]. After staining of the DNA with ethidium bromide, fragments I, II and III [see FIG. 8; these fragments originate from reactions I, II and III, respectively] were isolated from the gel and purified [Sambrook et al., 1989].

60 Preparation of DNA fragments encoding Glu³¹-TNF α and Asn³¹-Thr³²-TNF α

Fragments I, II and III were enzymatically phosphorylated, before in two parallel reactions fragments I and III and fragments II and III were ligated with each other [Sambrook et al., 1989]. After heat-inactivation of the ligase and digestion with restriction enzymes EcoRI and HindIII, the DNA was subjected to electrophoresis in a 6% polyacrylamide gel. After staining of the DNA

with ethidium bromide, the EcoRI-HindIII fragments A and B [see FIG. 4] were isolated from the gel and purified as previously described.

Preparation of plasmids encoding Glu³¹-TNF α and Asn³¹-Thr³²-TNF α

In separate experiments, the EcoRI-HindIII fragments A and B were inserted according to standard methods [Sambrook et al., 1989] into the EcoRI-HindIII opened plasmid pDS56/RBSII,Sph1-TNF α Ser29 generating plasmids pDS56/RBSII,Sph1-TNF α Glu31 and pDS56/RBSII,Sph1TNF α Asn31Thr32, respectively. Plasmid DNA was prepared [Birnboim et al., 1979] and the identity of the coding region for the TNF α muteins was confirmed by sequencing the double-stranded DNA [Sambrook et al., 1989].

Production of Glu³¹-TNF α and Asn³¹-Thr³²-TNF α

Plasmids pDS56/RBSII,Sph1-TNF α Glu31 and pDS56/RBSII,Sph1TNF α Asn31Thru32 were transformed into *E. coli* M15 cells containing already plasmid pREP4 by standard methods. Transformed cells were grown at 37° C. in LB medium containing 100 mg/l ampicillin and 25 mg/l kanamycin. At an optical density at 600 nm of about 0.7 to 1.0, IPTG was added to a final concentration of 2 mM. After additional 2.5 to 5 h at 37° C. the cells were harvested by centrifugation.

EXAMPLE VIII

Differential binding recombinant human p75-TNF-R and recombinant human p55-TNF-R

1. 10 ml suspensions of transformed and induced *E. coli* cells expressing recombinant human TNF α , Ser²⁹-TNF α , Trp³²-TNF α , Glu³¹-TNF α , and Asn³¹-Thr³²-TNF α [*E. coli* cells expressing recombinant dihydrofolate reductase (DHFR) were included as a control] were centrifuged at 4,000 rpm for 10 min and resuspended in 0.9 ml of lysis buffer (10 mM Tris-HCl pH 8.0, 5 mM EDTA, 2 mM PMSF, 10 mM benzamidine, 200 units/ml aprotinine and 0.1 mg/ml lysozyme). After

20 min incubation at room temperature 50 μ l of 1M MgCl₂, 20 μ l of 5 mg/ml DNaseI, 50 μ l of 5M NaCl and 50 μ l of 10% NP-40 were added and the mixture was further incubated at room temperature for 15 min. 0.5 ml of the lysate clarified by centrifugation at 13,000 rpm for 5 min was subjected to ammonium sulfate precipitation (30%–70% cut). The 70% ammonium sulfate pellet was dissolved in 0.2 ml PBS and analyzed by SDS-PAGE to confirm the presence of the recombinant proteins.

For the differential binding assay microtiter plates were coated with recombinant human p75-TNF-R-human IgG γ 3 and p55-TNF-R-human IgG γ 3 fusion proteins (European Patent Applications Publ. Nos. 417 563, 422 339) dissolved in PBS at 0.3 μ g/ml and 0.1 μ g/ml, respectively, (100 μ l/well, overnight at 4° C.). After blocking with blocking buffer (50 mM Tris pH 7.4, 140 mM NaCl, 5 mM EDTA, 0.02% NaN₃, 1% defatted milk powder) the microtiter plate was washed with PBS and incubated with 5 ng/ml human ¹²⁵I-TNF α (labelled by the Iodogen method to a specific activity of about 30 μ Ci/ μ g as described above) in the presence of different dilutions of the *E. coli* lysate partially purified by ammonium sulfate precipitation. The volume was 100 μ l/well and each dilution was assayed in duplicate. After three hours at room temperature the wells were thoroughly washed with PBS and counted in a γ -counter. Results are shown in FIG. 6 whereby closed circles refer to binding to p55-TNF-R-human IgG γ 3- and open circles refer to binding to p75-TNF-R-human IgG γ 3.

2. Determination of binding of Ser²⁹-Trp³²-TNF α , Gly²⁹-TNF α , Tyr²⁹-TNF α and Tyr³²-TNF α was performed as described under 1. with the only exception that MONO-Q ion exchange chromatography purified muteins were used. Results are shown in FIG. 7 whereby open and closed circles have the same meaning as in FIG. 6 and μ g/ml gives the amount of purified mutein/ml.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i i i) NUMBER OF SEQUENCES: 24

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 157 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(v i) ORIGINAL SOURCE:

- (A) ORGANISM: Homo sapiens
- (F) TISSUE TYPE: Blood
- (G) CELL TYPE: Macrophage

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His	Val
1				5					10					15	
Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Leu	Asn	Arg	Arg
			20					25					30		
Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln	Leu
		35					40					45			

-continued

Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu	Phe
	50					55					60				
Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr	Ile
65					70					75					80
Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser	Ala
				85					90					95	
Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala	Lys
			100					105					110		
Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu	Lys
		115					120					125			
Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp	Phe
	130					135					140				
Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu			
145					150					155					

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 3977 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (recombinant plasmid)

(v i i) IMMEDIATE SOURCE:

(B) CLONE: pDS56/RBSII,Sph1-TNF- alpha

(i x) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: 115..591

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:2:

CTCGAGAAAT	CATAAAAAAT	TTATTTGCTT	TGTGAGCGGA	TAACAATTAT	AATAGATTCA	60
ATTGTGAGCG	GATAACAATT	TCACACAGGA	TTCATTAAAG	AGGAGAAATT	AAGC ATG	117
					Met	
					1	
GTC AGA TCA TCT TCT CGA ACC CCG AGT GAC AAG CCT GTA GCC CAT GTT	165					
Val Arg Ser Ser Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His Val						
	5 10 15					
GTC GCG AAC CCT CAA GCT GAG GGG CAG CTC CAG TGG CTG AAC CGC CGG	213					
Val Ala Asn Pro Gln Ala Glu Gly Gln Leu Gln Trp Leu Asn Arg Arg						
	20 25 30					
GCC AAT GCC CTC CTG GCC AAT GGC GTG GAG CTG AGA GAT AAC CAG CTG	261					
Ala Asn Ala Leu Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln Leu						
	35 40 45					
GTG GTG CCA TCA GAG GGC CTG TAC CTC ATC TAC TCC CAG GTC CTC TTC	309					
Val Val Pro Ser Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu Phe						
	50 55 60 65					
AAG GGC CAA GGC TGC CCC TCC ACC CAT GTG CTC CTC ACC CAC ACC ATC	357					
Lys Gly Gln Gly Cys Pro Ser Thr His Val Leu Leu Thr His Thr Ile						
	70 75 80					
AGC CGC ATC GCC GTC TCC TAC CAG ACC AAG GTC AAC CTC CTC TCT GCC	405					
Ser Arg Ile Ala Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser Ala						
	85 90 95					
ATC AAG AGC CCC TGC CAG AGG GAG ACC CCA GAG GGG GCT GAG GCC AAG	453					
Ile Lys Ser Pro Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala Lys						
	100 105 110					
CCC TGG TAT GAG CCC ATC TAT CTG GGA GGG GTC TTC CAG CTG GAG AAG	501					
Pro Trp Tyr Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu Lys						
	115 120 125					
GGT GAC CGA CTC AGC GCT GAG ATC AAT CGG CCC GAC TAT CTC GAC TTT	549					
Gly Asp Arg Leu Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp Phe						
	130 135 140 145					

-continued

GCC GAG TCT GGG CAG GTC TAC TTT GGG ATC ATT GCC CTG TGAGGAGGAC	598
Ala Glu Ser Gly Gln Val Tyr Phe Gly Ile Ile Ala Leu	
150 155	
GAACATCCAA CCTTCCCAA CGCCTCCCCT GCCCCAATCC CTTTATTACC CCCTCCTTCA	658
GACACCCTCA ACCTCTTCTG GCTCAAAAAG AGAATTGGGG GCTTAGGGTC GGAACCCAAG	718
CTTGGACTCC TGTTGATAGA TCCAGTAATG ACCTCAGAAC TCCATCTGGA TTTGTTTCTG	778
ACGCTCGGTT GCCGCCGGGG GTTTTTTATT GGTGAGAATC CAAGCTAGCT TGGCGAGATT	838
TTCAGGAGCT AAGGAAGCTA AAATGGAGAA AAAAATCACT GGATATACCA CCGTTGATAT	898
ATCCCAATGG CATCGTAAAG AACATTTTGA GGCATTTTCTG TCAGTTGCTC AATGTACCTA	958
TAACCAGACC GTTACGCTGG ATATTACGGC CTTTTTAAAG ACCGTAAAGA AAAATAAGCA	1018
CAAGTTTTAT CCGGCCTTTA TTCACATTCT TGCCCGCCTG ATGAATGCTC ATCCGGAATT	1078
TCGTATGGCA ATGAAAGACG GTGAGCTGGT GATATGGGAT AGTGTTTACC CTTGTTACAC	1138
CGTTTTCCAT GAGCAAACCTG AAACGTTTTT C ATCGCTCTGG AGTGAATACC ACGACGATTT	1198
CCGGCAGTTT CTACACATAT ATTCGCAAGA TGTGGCGTGT TACGGTGAAA ACCTGGCCTA	1258
TTTCCCTAAA GGGTTTATTG AGAATATGTT TTTTCTCTCA GCCAATCCCT GGGTGAGTTT	1318
CACCAGTTTT GATTTAAACG TGGCCAATAT GGACAACCTT TTCGCCCCCG TTTTACCAT	1378
GGGCAAATAT TATACGCAAG GCGACAAGGT GCTGATGCCG CTGGCGATTC AGGTTTATCA	1438
TGCCGTCTGT GATGGCTTCC ATGTCGGCAG AATGCTTAAT GAATTACAAC AGTACTGCGA	1498
TGAGTGGCAG GGCGGGGCGT AATTTTTTTA AGGCAGTTAT TGGTGCCCTT AAACGCCTGG	1558
GGTAATGACT CTCTAGCTTG AGGCATCAAA TAAAACGAAA GGCTCAGTCG AAAGACTGGG	1618
CCTTTCGTTT TATCTGTTGT TTGTCCGGTGA ACGCTCTCCT GAGTAGGACA AATCCGCCGC	1678
TCTAGAGCTG CCTCGCGCGT TTCGGTGATG ACGGTGAAAA CCTCTGACAC ATGCAGCTCC	1738
CGGAGACGGT CACAGCTTGT CTGTAAGCGG ATGCCGGGAG CAGACAAGCC CGTCAGGGCG	1798
CGTCAGCGGG TGTTGGCGGG TGTCGGGGCG CAGCCATGAC CCAGTCACGT AGCGATAGCG	1858
GAGTGTATAC TGGCTTAACT ATGCCGCATC AGAGCAGATT GACTGAGAG TGCACCATAT	1918
GCGGTGTGAA ATACCGCACA GATGCGTAAG GAGAAAATAC CGCATCAGGC GCTCTTCCGC	1978
TTCTCGCTC ACTGACTCGC TGCGCTCGGT CTGTCCGGCTG CGGCGAGCGG TATCAGCTCA	2038
CTCAAAGGCG GTAATACGGT TATCCACAGA ATCAGGGGAT AACGCAGGAA AGAACATGTG	2098
AGCAAAGGC CAGCAAAGG CCAGGAACCG TAAAAGGCC GCGTTGCTGG CGTTTTTCCA	2158
TAGGCTCCGC CCCCTGACG AGCATCACAA AAATCGACGC TCAAGTCAGA GGTGGCGAAA	2218
CCCGACAGGA CTATAAAGAT ACCAGGCGTT TCCCCCTGGA AGCTCCCTCG TGCCTCTCC	2278
TGTTCCGACC CTGCCGCTTA CCGGATACCT GTCCGCCTTT CTCCCTTCGG GAAGCGTGGC	2338
GCTTTCTCAA TGCTCACGCT GTAGGTATCT CAGTTGCCTG TAGGTCGTTT GCTCCAAGCT	2398
GGGCTGTGTG CACGAACCC CCGTTCAGCC CGACCGCTGC GCCTTATCCG GTAACATCG	2458
TCTTGAGTCC AACCCGGTAA GACACGACTT ATCGCCACTG GCAGCAGCCA CTGGTAACAG	2518
GATTAGCAGA GCGAGGTATG TAGGGGGTGC TACAGAGTTC TTGAAGTGGT GGCCTAACTA	2578
CGGCTACACT AGAAGGACAG TATTTGGTAT CTGCGCTCTG CTGAAGCCAG TTACCTTCGG	2638
AAAAAGAGTT GGTAGCTCTT GATCCGGCAA ACAAACCACC GCTGGTAGCG GTGGTTTTTT	2698
TGTTTGCAAG CAGCAGATTA CGCGCAGAAA AAAAGGATCT CAAGAAGATC CTTTGATCTT	2758
TTCTACGGGG TCTGACGCTC AGTGGAACGA AAATCACGT TAAGGGATTT TGGTCATGAG	2818
ATTATCAAAA AGGATCTTCA CCTAGATCCT TTAAATTA AAATGAAGTT TAAATCAAT	2878
CTAAAGTATA TATGAGTAAA CTTGGTCTGA CAGTTACCAA TGCTTAATCA GTGAGGCACC	2938

-continued

```

TATCTCAGCG ATCTGTCTAT TTCGTTTCATC CATAGCTGCC TGACTIONCCCG TCGTG TAGAT 2998
AACTACGATA CGGGAGGGCT TACCATCTGG CCCAGTGCT GCAATGATAC CGCGAGACCC 3058
ACGCTCACCG GCTCCAGATT TATCAGCAAT AAACCAGCCA GCCGGAAGGG CCGAGCGCAG 3118
AAGTGGTCCT GCAACTTTAT CCGCCTCCAT CCAGTCTATT AATTGTTGCC GGAAGCTAG 3178
AGTAAGTAGT CCGCCAGTTA ATAGTTTGCG CAACGTTGTT GCCATTGCTA CAGGCATCGT 3238
GGTCTCACGC TCGTCGTTTG GTATGGCTTC ATTCAGCTCC GGTTCCCAAC GATCAAGGCG 3298
AGTTACATGA TCCCCCATGT TGTGCAAAAA AGCGGTTAGC TCCTTCGGTC CTCCGATCGT 3358
TGTCAGAAGT AAGTTGGCCG CAGTGTTATC ACTCATGGTT ATGGCAGCAC TGCATAATTC 3418
TCTTACTGTC ATGCCATCCG TAAGATGCTT TTCTGTGACT GGTGAGTACT CAACCAAGTC 3478
ATTCTGAGAA TAGTGTATGC GGCGACCGAG TTGCTCTTGC CCGGCGTCAA TACGGGATAA 3538
TACCGCGCCA CATAGCAGAA CTTTAAAAGT GCTCATCATT GGAAAACGTT CTTCGGGGCG 3598
AAAACCTCTCA AGGATCTTAC CGCTGTTGAG ATCCAGTTCG ATGTAACCCA CTCGTGCACC 3658
CAACTGATCT TCAGCATCTT TTTACTTTCAC CAGCGTTTCT GGCTGAGCAA AAACAGGAAG 3718
GCAAAAATGCC GCAAAAAAAGG GAATAAGGGC GACACGGAAA TGTTGAATAC TCATACTCTT 3778
CCTTTTTCAA TATTATTGAA GCATTTATCA GGGTTATTGT CTCATGAGCG GATACATATT 3838
TGAATGTATT TAGAAAAATA AACAAATAGG GGTTCCGCGC ACATTTCCCC GAAAAGTCCC 3898
ACCTGACGTC TAAGAAACCA TTATTATCAT GACATTAACC TATAAAAATA GGCATATCAC 3958
GAGGCCCTTT CGTCTTCAC 3977

```

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 158 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(i x) FEATURE:

- (A) NAME/KEY: Modified site
- (B) LOCATION: 29, 31 and 32
- (D) OTHER INFORMATION:/note="Xaa =any naturally occurring amino acid"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:3:

```

Met Val Arg Ser Ser Ser Arg Thr Pro Ser Asp Lys Pro Val Ala His
-1 +1 5 10 15
Val Val Ala Asn Pro Gln Ala Glu Gly Gln Leu Gln Trp Xaa Asn Xaa
20 25 30
Xaa Ala Asn Ala Leu Leu Ala Asn Gly Val Glu Leu Arg Asp Asn Gln
35 40 45
Leu Val Val Pro Ser Glu Gly Leu Tyr Leu Ile Tyr Ser Gln Val Leu
50 55 60
Phe Lys Gly Gln Gly Cys Pro Ser Thr His Val Leu Leu Thr His Thr
65 70 75
Ile Ser Arg Ile Ala Val Ser Tyr Gln Thr Lys Val Asn Leu Leu Ser
80 85 90 95
Ala Ile Lys Ser Pro Cys Gln Arg Glu Thr Pro Glu Gly Ala Glu Ala
100 105 110
Lys Pro Trp Tyr Glu Pro Ile Tyr Leu Gly Gly Val Phe Gln Leu Glu
115 120 125
Lys Gly Asp Arg Leu Ser Ala Glu Ile Asn Arg Pro Asp Tyr Leu Asp
130 135 140

```


-continued

Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu
	145					150					155		

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1				5					10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Ser	Asn	Arg
			20						25					30	
Arg	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
		35						40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1				5					10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Gly	Asn	Arg
			20						25					30	
Arg	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
		35						40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp

-continued

	130						135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1				5					10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Tyr	Asn	Arg
			20						25					30	
Arg	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1				5					10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Leu	Asn	Glu
			20						25					30	
Arg	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		

-continued

Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1				5					10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Leu	Asn	Asn
				20					25					30	
Arg	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1				5					10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Leu	Asn	Arg
				20					25					30	
Trp	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	

-continued

Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1					5				10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Leu	Asn	Arg
				20					25					30	
Tyr	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 158 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1					5				10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Leu	Asn	Asn
				20					25					30	
Thr	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala

-continued

100					105					110					
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 158 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met	Val	Arg	Ser	Ser	Ser	Arg	Thr	Pro	Ser	Asp	Lys	Pro	Val	Ala	His
-1	+1					5				10					15
Val	Val	Ala	Asn	Pro	Gln	Ala	Glu	Gly	Gln	Leu	Gln	Trp	Ser	Asn	Arg
				20					25					30	
Trp	Ala	Asn	Ala	Leu	Leu	Ala	Asn	Gly	Val	Glu	Leu	Arg	Asp	Asn	Gln
			35					40					45		
Leu	Val	Val	Pro	Ser	Glu	Gly	Leu	Tyr	Leu	Ile	Tyr	Ser	Gln	Val	Leu
		50					55					60			
Phe	Lys	Gly	Gln	Gly	Cys	Pro	Ser	Thr	His	Val	Leu	Leu	Thr	His	Thr
	65					70					75				
Ile	Ser	Arg	Ile	Ala	Val	Ser	Tyr	Gln	Thr	Lys	Val	Asn	Leu	Leu	Ser
80					85					90					95
Ala	Ile	Lys	Ser	Pro	Cys	Gln	Arg	Glu	Thr	Pro	Glu	Gly	Ala	Glu	Ala
				100					105					110	
Lys	Pro	Trp	Tyr	Glu	Pro	Ile	Tyr	Leu	Gly	Gly	Val	Phe	Gln	Leu	Glu
			115					120					125		
Lys	Gly	Asp	Arg	Leu	Ser	Ala	Glu	Ile	Asn	Arg	Pro	Asp	Tyr	Leu	Asp
		130					135					140			
Phe	Ala	Glu	Ser	Gly	Gln	Val	Tyr	Phe	Gly	Ile	Ile	Ala	Leu		
	145					150					155				

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 3977 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (recombinant plasmid)

(v i i) IMMEDIATE SOURCE:
 (B) CLONE: pDS56/RBSII,SphI-TNF- alpha

(i x) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 115..591

(i x) FEATURE:
 (A) NAME/KEY: Modified site
 (B) LOCATION: 202-204, 208-210 and 211-213
 (D) OTHER INFORMATION:/note="N =A, G, C or T"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CTCGAGAAAT	CATAAAAAAT	TTATTTGCTT	TGTGAGCGGA	TAACAATTAT	AATAGATTCA	60
ATTGTGAGCG	GATAACAATT	TCACACAGGA	TTCATTAAAG	AGGAGAAATT	AAGC ATG	117
					Met	

-continued

CGTCAGCGGG	TGTTGGCGGG	TGTCGGGGCG	CAGCCATGAC	CCAGTCACGT	AGCGATAGCG	1858
GAGTGTATAC	TGGCTTAACT	ATGCCGCATC	AGAGCAGATT	GTAAGTACAG	TGCACCATAT	1918
GCGGTGTGAA	ATACCGCACA	GATGCGTAAG	GAGAAAATAC	CGCATCAGGC	GCTCTTCCGC	1978
TTCTCGCTC	ACTGACTCGC	TGCGCTCGGT	CTGTCGGCTG	CGGCGAGCGG	TATCAGCTCA	2038
CTCAAAGGCG	GTAATACGGT	TATCCACAGA	ATCAGGGGAT	AACGCAGGAA	AGAACATGTG	2098
AGCAAAGGC	CAGCAAAGG	CCAGGAACCG	TAAAAAGGCC	GCGTTGCTGG	CGTTTTTCCA	2158
TAGGCTCCGC	CCCCCTGACG	AGCATCACAA	AAATCGACGC	TCAAGTCAGA	GGTGGCGAAA	2218
CCCGACAGGA	CTATAAAGAT	ACCAGGCGTT	TCCCCCTGGA	AGCTCCCTCG	TGCGCTCTCC	2278
TGTTCCGACC	CTGCCGCTTA	CCGGATACCT	GTCGGCCTTT	CTCCCTTCGG	GAAGCGTGGC	2338
GCTTTCTCAA	TGCTCACGCT	GTAGGTATCT	CAGTTGCCTG	TAGGTCGTTC	GCTCCAAGCT	2398
GGGCTGTGTG	CACGAACCCC	CCGTTCAGCC	CGACCGCTGC	GCCTTATCCG	GTAACATATCG	2458
TCTTGAGTCC	AACCCGGTAA	GACACGACTT	ATCGCCACTG	GCAGCAGCCA	CTGGTAACAG	2518
GATTAGCAGA	GCGAGGTATG	TAGGGGGTGC	TACAGAGTTC	TTGAAGTGGT	GGCCTAACTA	2578
CGGCTACACT	AGAAGGACAG	TATTTGGTAT	CTGCGCTCTG	CTGAAGCCAG	TTACCTTCGG	2638
AAAAAGAGTT	GGTAGCTCTT	GATCCGGCAA	ACAAACCACC	GCTGGTAGCG	GTGGTTTTTT	2698
TGTTTGCAAG	CAGCAGATTA	CGCGCAGAAA	AAAAGGATCT	CAAGAAGATC	CTTTGATCTT	2758
TTCTACGGGG	TCTGACGCTC	AGTGGAACGA	AAACTCACGT	TAAGGGATTT	TGGTCATGAG	2818
ATTATCAAAA	AGGATCTTCA	CCTAGATCCT	TTTAAATTAA	AAATGAAGTT	TTAAATCAAT	2878
CTAAAGTATA	TATGAGTAAA	CTTGGTCTGA	CAGTTACCAA	TGCTTAATCA	GTGAGGCACC	2938
TATCTCAGCG	ATCTGTCTAT	TTCGTTTATC	CATAGCTGCC	TGACTCCCCG	TCGTGTAGAT	2998
AACTACGATA	CGGGAGGGCT	TACCATCTGG	CCCAGTGCT	GCAATGATAC	CGCGAGACCC	3058
ACGCTCACCG	GCTCCAGATT	TATCAGCAAT	AAACCAGCCA	GCCGGAAGGG	CCGAGCGCAG	3118
AAGTGGTCCT	GCAACTTTAT	CCGCCTCCAT	CCAGTCTATT	AATTGTTGCC	GGGAAGCTAG	3178
AGTAAGTAGT	CCGCCAGTTA	ATAGTTTGCG	CAACGTTGTT	GCCATTGCTA	CAGGCATCGT	3238
GGTCTCACGC	TCGTCGTTTG	GTATGGCTTC	ATTCAGCTCC	GGTTCCCAAC	GATCAAGGCG	3298
AGTTACATGA	TCCCCCATGT	TGTGCAAAAA	AGCGGTTAGC	TCCTTCGGTC	CTCCGATCGT	3358
TGTCAGAAGT	AAGTTGGCCG	CAGTGTTATC	ACTCATGGTT	ATGGCAGCAC	TGCATAATTC	3418
TCTTACTGTC	ATGCCATCCG	TAAGATGCTT	TTCTGTGACT	GGTGAGTACT	CAACCAAGTC	3478
ATTCTGAGAA	TAGTGTATGC	GGCGACCGAG	TTGCTCTTGC	CCGGCGTCAA	TACGGGATAA	3538
TACCGCGCCA	CATAGCAGAA	CTTTAAAAGT	GCTCATCATT	GGAAAACGTT	CTTCGGGGCG	3598
AAAACCTCTCA	AGGATCTTAC	CGCTGTTGAG	ATCCAGTTCG	ATGTAACCCA	CTCGTGCACC	3658
CAACTGATCT	TCAGCATCTT	TTACTTTCAC	CAGCGTTTCT	GGCTGAGCAA	AAACAGGAAG	3718
GCAAAATGCC	GCAAAAAAGG	GAATAAGGGC	GACACGGAAA	TGTTGAATAC	TCATACTCTT	3778
CCTTTTTCAA	TATTATTGAA	GCATTTATCA	GGGTTATTGT	CTCATGAGCG	GATACATATT	3838
TGAATGTATT	TAGAAAAATA	AACAAATAGG	GGTTCCGCGC	ACATTTCCCC	GAAAAGTCCC	3898
ACCTGACGTC	TAAGAAACCA	TTATTATCAT	GACATTAACC	TATAAAAATA	GGCGTATCAC	3958
GAGGCCCTTT	CGTCTTCAC					3977

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 3740 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

-continued

(i i) MOLECULE TYPE: DNA (recombinant plasmid)

(i x) FEATURE:

(A) NAME/KEY: CDS

(B) LOCATION: complement (2613..one- of(1532))

(D) OTHER INFORMATION: /note="Contains coding region for the lacI gene beginning at residue 2613 to 1532"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:14:

AAGCTTCACG	CTGCCGCAAG	CACTCAGGGC	GCAAGGGCTG	CTAAAGGAAG	CGGAACACGT	60
AGAAAGCCAG	TCCGCAGAAA	CGGTGCTGAC	CCCGGATGAA	TGTCAGCTAC	TGGGCTATCT	120
GGACAAGGGA	AAACGCAAGC	GCAAAGAGAA	AGCAGGTAGC	TTGCAGTGGG	CTTACATGGC	180
GATAGCTAGA	CTGGGCGGTT	TTATGGACAG	CAAGCGAACC	GGAATTGCCA	GCTGGGGCGC	240
CCTCTGGTAA	CGTTGGGAAG	CCCTGCAAAG	TAAACTGGAT	GGCTTTCTTG	CCGCCAAGGA	300
TCTGATGGCG	CAGGGGATCA	AGATCTGATC	AAGAGACAGG	ATGACGGTCG	TTTCGCATGC	360
TTGAACAAGA	TGGATTGCAC	GCAGGTTCTC	CGGCCGCTTG	GGTCGAGAGG	CTATTCGGCT	420
ATGACTGGGC	ACAACAGACA	ATCCGCTGCT	CTGATGCCGC	CGTGTTCCGG	CTGTCAGCCC	480
AGGGGCGCCC	GGTTCTTTTT	GTCAAGACCG	ACCTGTCCGG	TGCCCTGAAT	GAAGTGCAGG	540
ACGAGGCAGC	GCGGCTATCG	TGGCTGGCCA	CGACGGGCGT	TCCTTGCGCA	GCTGTGCTCG	600
ACGTTGTAC	TGAAGCGGGA	AGGGACTGGC	TGCTATTGGG	CGAAGTGCCG	GGGCAGGATC	660
TCCTGTCATC	TCACCTTGCT	CCTGCCGAGA	AAGTATCCAT	CATGGCTGAT	GCAATGCGGC	720
GGCTGCATAC	GCTTGATCCG	GCTACCTGCC	CATTCGACCA	CCAAGCGAAA	CATCGCATCG	780
AGCGAGCACG	TACTCGGATG	GAAGCCGGTC	TTGTCGATCA	GGATCATCTG	GACGAAGAGC	840
ATCAGGGGCT	CGCGCCAGCC	GAAGTGTTCG	CCAGGCTCAA	GGCGCGCATG	CCCGACGGCG	900
AGGATCTCGT	CGTGACCCAT	GGCGATGCCT	GCTTGCCGAA	TATCATGGTG	GAAAATGGCC	960
GCTTTTCTGG	ATTCATCGAC	TGTGGCCGGC	TGGGTGTGGC	GGACCGCTAT	CAGGACATAG	1020
CGTTGGCTAC	CCGTGATATT	GCTGAAGAGC	TTGGCGGCGA	ATGGGCTGAC	CGCTTCCTCG	1080
TGCTTTACGG	TATCGCCGCT	CCCGATTTCG	AGCGCATCGC	CTTCTATCGC	CTTCTTGACG	1140
AGTTCTTCTG	AGCGGGACTC	TGGGGTTCGA	AATGACCGAC	CAAGCGACGC	CCAACCTGCC	1200
ATCACGAGAT	TTCGATTCCA	CCGCCGCCTT	CTATGAAAGG	TTGGGCTTCG	GAATCGTTTT	1260
CCGGGACGCC	GGCTGGATGA	TCCTCCAGCG	CGGGGATCTC	ATGCTGGAGT	TCTTCGCCCA	1320
CCCCGGGCTC	GATCCCCTCG	CGAGTTGGTT	CAGCTGCTGC	CTGAGGCTGG	ACGACCTCGC	1380
GGAGTTCTAC	CGGCAGTGCA	AATCCGTCGG	CATCCAGGAA	ACCAGCAGCG	GCTATCCGCG	1440
CATCCATGCC	CCCGAACTGC	AGGAGTGGGG	AGGCACGATG	GCCGCTTTGG	TCGACAATTC	1500
GCGCTAACTT	ACATTAATTG	CGTTGCGCTC	ACTGCCCGCT	TTCCAGTCGG	GAAACCTGTC	1560
GTGCCAGCTG	CATTAATGAA	TCGGCCAACG	CGCGGGGAGA	GGCGGTTTGC	GTATTGGGCG	1620
CCAGGGTGGT	TTTTCTTTTC	ACCAGTGAGA	CGGGCAACAG	CTGATTGCC	TTCACCGCCT	1680
GGCCCTGAGA	GAGTTGCAGC	AAGCGGTCCA	CGCTGGTTTG	CCCAGCAGG	CGAAAATCCT	1740
GTTTGATGGT	GGTTAACGGC	GGGATATAAC	ATGAGCTGTC	TTCGGTATCG	TCGTATCCCA	1800
CTACCGAGAT	ATCCGCACCA	ACGCGCAGCC	CGGACTCGGT	AATGGCGCGC	ATTGCGCCCA	1860
GCGCCATCTG	ATCGTTGGCA	ACCAGCATCG	CAGTGGGAAC	GATGCCCTCA	TTCAGCATTT	1920
GCATGGTTTG	TTGAAAACCG	GACATGGCAC	TCCAGTCGCC	TTCCCGTTCC	GCTATCGGCT	1980
GAATTTGATT	GCGAGTGAGA	TATTTATGCC	AGCCAGCCAG	ACGCAGACGC	GCCGAGACAG	2040
AACTTAATGG	GCCCGCTAAC	AGCGCGATTT	GCTGGTGACC	CAATGCGACC	AGATGCTCCA	2100
CGCCCAGTCG	CGTACCGTCT	TCATGGGAGA	AAATAATACT	GTTGATGGGT	GTCTGGTCAG	2160

-continued

AGACATCAAG	AAATAACGCC	GGAACATTAG	TGCAGGCAGC	TTCCACAGCA	ATGGCATCCT	2 2 2 0
GGTCATCCAG	CGGATAGTTA	ATGATCAGCC	CACTGACGCG	TTGCGCGAGA	AGATTGTGCA	2 2 8 0
CCGCCGCTTT	ACAGGCTTCG	ACGCCGCTTC	GTTCTACCAT	CGACACCACC	ACGCTGGCAC	2 3 4 0
CCAGTTGATC	GGCGCGAGAT	TTAATCGCCG	CGACAATTTG	CGACGGCGCG	TGCAGGGCCA	2 4 0 0
GACTGGAGGT	GGCAACGCCA	ATCAGCAACG	ACTGTTTGCC	CGCCAGTTGT	TGTGCCACGC	2 4 6 0
GGTTGGGAAT	GTAATTCAGC	TCCCCATCG	CCGCTTCCAC	TTTTTCCCGC	GTTTTCGCAG	2 5 2 0
AAACGTGGCT	GGCCTGGTTC	ACCACGCGGG	AAACGGTCTG	ATAAGAGACA	CCGGCATACT	2 5 8 0
CTGCGACATC	GTATAACGTT	ACTGGTTTCA	CATTCACCAC	CCTGAATTGA	CTCTCTTCCG	2 6 4 0
GGCGCTATCA	TGCCATAACG	CGAAAGGTTT	TGCGCCATTC	GATGGTGTCA	ACGTAAATGC	2 7 0 0
ATGCCGCTTC	GCCTTCGCCC	GCGAATTGTC	GACCCTGTCC	CTCCTGTTCA	GCTACTGACG	2 7 6 0
GGGTGGTGCG	TAACGGCAAA	AGCACCGCCG	GACATCAGCG	CTAGCGGAGT	GTATACTGGC	2 8 2 0
TTACTATGTT	GGCACTGATG	AGGGTGTGAG	TGAAGTGCTT	CATGTGGCAG	GAGAAAAAAG	2 8 8 0
GCTGCACCGG	TGCGTCAGCA	GAATATGTGA	TACAGGATAT	ATTCCGCTTC	CTCGCTCACT	2 9 4 0
GACTCGCTAC	GCTCGGTCGT	TCGACTGCGG	CGAGCGGAAA	TGGCTTACGA	ACGGGGCGGA	3 0 0 0
GATTTCTCTGG	AAGATGCCAG	GAAGATACTT	AACAGGGAAG	TGAGAGGGCC	GCGGCAAAGC	3 0 6 0
CGTTTTTCCA	TAGGCTCCGC	CCCCCTGACA	AGCATCACGA	AATCTGACGC	TCAAATCAGT	3 1 2 0
GGTGGCGAAA	CCCCACAGGA	CTATAAAGAT	ACCAGGCGTT	TCCCCTGGCG	GCTCCCTCGT	3 1 8 0
GCGCTCTCCT	GTTCTGCTT	TTCGTTTAC	CGGTGTCATT	CCGCTGTTAT	GGCCGCGTTT	3 2 4 0
GTCTCATTCC	ACGCCTGACA	CTCAGTTCCG	GGTAGGCAGT	TCGCTCCAAG	CTGGACTGTA	3 3 0 0
TGCACGAACC	CCCCGTTTCC	TCCGACCGCT	GCGCCTTATC	CGGTAACTAT	CGTCTTGAGT	3 3 6 0
CCAACCCGGA	AAGACATGCA	AAAGCACACC	TGGCAGCAGC	CACTGGTAAT	TGATTTAGAG	3 4 2 0
GAGTTAGTCT	TGAAGTCATG	CGCCGGTTAA	GGCTAAACTG	AAAGGACAAG	TTTTCGTCAC	3 4 8 0
TGCGCTCCTC	CAAGCCAGTT	ACCTCGGTTT	AAAGAGTTGG	TAGCTCAGAG	AACCTTCGAA	3 5 4 0
AAACCGCCCT	GCAAGGCGGT	TTTTTCGTTT	TCAGAGCAAG	AGATTACGCG	CAGACCAAAA	3 6 0 0
CGATCTCAAG	AAGATCATCT	TATTAATCAG	ATAAAATATT	TCTAGATTTC	AGTGCAATTT	3 6 6 0
ATCTCTTCAA	ATGTAGCACC	TGAAGTCAGC	CCCATACGAT	ATAAGTTGTT	AATTCTCATG	3 7 2 0
TTTGACAGCT	TATCATCGAT					3 7 4 0

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..22
- (D) OTHER INFORMATION: /function=
" Oligonucleotide used for gap
duplex mutagenesis"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCGGCGGTTG GACCACTGGA GC

2 2

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs

-continued

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..19
- (D) OTHER INFORMATION: /function=
" Oligonucleotide used for gap
duplex mutagenesis"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CATTGGCCCA GCGGTT CAG

19

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 32 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..32
- (D) OTHER INFORMATION: /function=
" Oligonucleotide used for gap
duplex mutagenesis"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GGGCATTGGC CCAGCGGTTG GACCACTGGA GC

32

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 50 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..50
- (D) OTHER INFORMATION: /function=
" Oligonucleotide used for gap
duplex mutagenesis"
/ note="Used to create any position 29-mutein of
TNF-alpha"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CCACGCCATT CGCGAGGAGG GCATTGGCCC GGCGGTTNNN CCACTGGAGC

50

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..42
- (D) OTHER INFORMATION: /function=
" Oligonucleotide used for gap
duplex mutagenesis"
/ note="Used to create any position 32-mutein of
TNF-alpha"

-continued

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CCACGCCATT CGCGAGGAGG GCATTGGCNN NGCGGTTTCAG CC

4 2

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: complement (1..22)
- (D) OTHER INFORMATION: /function="PCR primer"
/ note="Complementary to positions 3949 to 3970 of
Sequence ID No. 2"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGCGTATCAC GAGGCCCTTT CG

2 2

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..22
- (D) OTHER INFORMATION: /function="PCR primer"
/ note="Complementary to positions
748 - 727 of Seq. ID No. 2"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CATTACTGGA TCTATCAACA GG

2 2

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

- (A) NAME/KEY: misc_feature
- (B) LOCATION: 1..36
- (D) OTHER INFORMATION: /function="PCR primer"
/ product="primer 21/M5"
/ note="PCR primer which is complementary to
positions 219-184 of Seq. ID No. 2 with mismatched
residues at positions 10-12."

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:22:

ATTGGCCCGC TCGTTCAGCC ACTGGAGCTG CCCCTC

3 6

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 36 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

-continued

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..36

(D) OTHER INFORMATION: /function="PCR primer for mutagenesis"

/ note="PCR primer for mutagenesis which is complementary to positions 219-184 of Seq. ID No. 2 with mismatched bases at positions 7-9 and 11-12"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:23:

ATTGGCAGTG TTGTTTCAGCC ACTGGAGCTG CCCCTC

3 6

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(i i) MOLECULE TYPE: DNA (oligonucleotide)

(i x) FEATURE:

(A) NAME/KEY: misc_feature

(B) LOCATION: 1..22

(D) OTHER INFORMATION: /function="PCR primer"

/ product="primer 21/MR"

/ note="PCR primer used in conjunction with Seq.

ID Nos. 22 & 23 to create muteins of TNF-alpha"

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:24:

GCCCTCCTGG CCAATGGCGT GG

2 2

We claim:

1. A human Tumor Necrosis Factor mutein or a pharmaceutically acceptable salt thereof consisting of SEQ ID NO:4.

2. A human Tumor Necrosis Factor mutein or a pharmaceutically acceptable salt thereof consisting of SEQ ID NO: 5.

3. A human Tumor Necrosis Factor mutein or a pharmaceutically acceptable salt thereof consisting of SEQ ID NO: 6.

4. A human Tumor Necrosis Factor mutein or a pharmaceutically acceptable salt thereof consisting of SEQ ID NO: 12.

5. A human Tumor Necrosis Factor mutein or a pharmaceutically acceptable salt thereof consisting of SEQ ID NO: 12.

* * * * *

45

50

55

60

65