



US005683676A

United States Patent [19][11] **Patent Number:** **5,683,676**

Akehurst et al.

[45] **Date of Patent:** ***Nov. 4, 1997**[54] **CANISTER CONTAINING AEROSOL FORMULATIONS CONTAINING P134A AND PARTICULATE MEDICAMENTS**5,118,494 6/1992 Schultz et al. 424/45
5,126,123 6/1992 Johnson 424/45**FOREIGN PATENT DOCUMENTS**[75] **Inventors: Rachel Ann Akehurst; Anthony James Taylor; David Andrew Wyatt, all of Ware, Great Britain**0372777 6/1990 European Pat. Off. 424/45
0 504 112 9/1992 European Pat. Off. .
27 03 119 10/1990 Germany .
90/07333 7/1990 WIPO .[73] **Assignee: Glaxo Group Limited, London, England****OTHER PUBLICATIONS**[*] **Notice:** The term of this patent shall not extend beyond the expiration date of Pat. No. 5,676,929.

Genarro, A. R. (1985. Remington's Pharmaceutical Sciences, 17th edition, Mack Pub. Co., pp. 1670-1677.

Oberholz, *Frankfurter Allgemeine Zeitung*, Oct. 1989, vol. 25, No. 207, p. 7.Dalby et al., *Pharmaceutical Technology*, Mar. 1990, vol. 14, No. 3, pp. 26-33.[21] **Appl. No.: 444,926**

Amzacort™ carton, William H. Rorer, Inc., Fort Washington, Pennsylvania, USA 19034, 1986.

[22] **Filed: May 19, 1995***Pharmaceutical Journal*, Sep. 29, 1990, vol. 245, pp. 428-429.**Related U.S. Application Data***The Theory and Practice of Industrial Pharmacy*, 2nd Ed., 1976 (Philadelphia, PA: Lea and Febiger), pp. 270 and 276-278.[62] **Division of Ser. No. 328,959, Oct. 24, 1994, abandoned, which is a continuation of Ser. No. 102,241, Aug. 5, 1993, abandoned.***Handbook of Aerosol Technology*, 2nd Edition, 1979 (New York, New York: Van Nostrand Reinhold Company), pp. 30, 32, 33, 166, 167, 232, 233.**[30] Foreign Application Priority Data**Dec. 12, 1991 [GB] United Kingdom 9126378
Dec. 12, 1991 [GB] United Kingdom 9126405
Feb. 6, 1992 [GB] United Kingdom 9202522

U.S. Senate Hearings, 12-14 May 1987, 343-347, 437 (U.S. Government Printing Office, Washington, D.C., 1987), CIS:1987-S321-26.

[51] **Int. Cl.⁶ A61K 9/12***Hagers Handbook of Pharmaceutical Practice*, 1971, pp. 342-354 (Berlin: Springer-Verlag).[52] **U.S. Cl. 424/45; 424/46; 222/635**[58] **Field of Search 222/402.1, 635; 424/45, 46***Primary Examiner*—Raj Bawa*Attorney, Agent, or Firm*—Bacon & Thomas**[56] References Cited****[57] ABSTRACT****U.S. PATENT DOCUMENTS**

This invention relates to aerosol formulations of use for the administration of medicaments by inhalation, in particular a pharmaceutical aerosol formulation which comprises particulate salmeterol and physiologically acceptable salts and solvates thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant. A method of treating respiratory disorders which comprises administration by inhalation of an effective amount of a pharmaceutical aerosol formulation as defined is also described.

2,868,691 1/1959 Porush et al. 167/54
2,885,427 5/1959 Rob et al. 260/653.7
3,014,844 12/1961 Thiel et al. 167/82
3,219,533 11/1965 Mullins 167/82
3,320,125 5/1967 Grim 167/54
3,809,294 5/1974 Torgeson 222/182
3,897,779 8/1975 Hansen 128/266
4,174,295 11/1979 Bergigia et al. 252/305
4,405,598 9/1983 Brown 424/45
4,940,171 7/1990 Gilroy 222/402.18**32 Claims, No Drawings**

CANISTER CONTAINING AEROSOL FORMULATIONS CONTAINING P134A AND PARTICULATE MEDICAMENTS

The present application is a divisional of application Ser. No. 08/328,959, filed Oct. 24, 1994 (now abandoned) which is a continuation of application Ser. No. 08/102,241, filed Aug. 5, 1993 (now abandoned), which is a divisional of application Ser. No. PCT/EP92/02808, filed Dec. 4, 1992 (now abandoned).

BACKGROUND OF THE INVENTION

1. Field of the Invention

This invention relates to aerosol formulations of use for the administration of medicaments by inhalation.

The use of aerosols to administer medicaments has been known for several decades. Such aerosols generally comprise the medicament, one or more chlorofluorocarbon propellants and either a surfactant or a solvent, such as ethanol. The most commonly used aerosol propellants for medicaments have been propellant 11 (CCl₃F) and/or propellant 114 (CF₂ClCF₂Cl) with propellant 12 (CCl₂F₂). However these propellants are now believed to provoke the degradation of stratospheric ozone and there is thus a need to provide aerosol formulations for medicaments which employ so called "ozone-friendly" propellants.

2. Description of the Prior Art

A class of propellants which are believed to have minimal ozone-depleting effects in comparison to conventional chlorofluorocarbons comprise fluorocarbons and hydrogen-containing chlorofluorocarbons, and a number of medicinal aerosol formulations using such propellant systems are disclosed in, for example, EP 0372777, WO91/04011, WO91/11173, WO91/11495 and WO91/14422. These applications are all concerned with the preparation of pressurised aerosols for the administration of medicaments and seek to overcome the problems associated with the use of the new class of propellants, in particular the problems of stability associated with the pharmaceutical formulations prepared. The applications all propose the addition of one or more of adjuvants such as alcohols, alkanes, dimethyl ether, surfactants (including fluorinated and non-fluorinated surfactants, carboxylic acids, polyethoxylates etc) and even conventional chlorofluorocarbon propellants in small amounts intended to minimise potential ozone damage.

Thus, for example EP 0372777 requires the use of 1,1,1,2-tetrafluoroethane in combination with both a cosolvent having greater polarity than 1,1,1,2-tetrafluoroethane (e.g. an alcohol or a lower alkane) and a surfactant in order to achieve a stable formulation of a medicament powder. In particular it is noted in the specification at page 3, line 7 that "it has been found that the use of propellant 134a (1,1,1,2-tetrafluoroethane) and drug as a binary mixture or in combination with a conventional surfactant such as sorbitan trioleate does not provide formulations having suitable properties for use with pressurised inhalers". Surfactants are generally recognised by those skilled in the art to be essential components of aerosol formulations, required not only to reduce aggregation of the medicament but also to lubricate the valve employed, thereby ensuring consistent reproducibility of valve actuation and accuracy of dose dispensed. Whilst WO91/11173, WO91/11495 and WO91/14422 are concerned with formulations comprising an admixture of drug and surfactant, WO91/04011 discloses medicinal aerosol formulations in which the particulate medicaments are pre-coated with surfactant prior to dispersal in 1,1,1,2-tetrafluoroethane.

SUMMARY OF THE INVENTION

We have now surprisingly found that, in contradistinction to these teachings, it is in fact possible to obtain satisfactory dispersions of certain medicaments in fluorocarbon or hydrogen-containing chlorofluorocarbon propellants such as 1,1,1,2-tetrafluoroethane without recourse to the use of any surfactant or cosolvent in the composition, or the necessity to pre-treat the medicament prior to dispersal in the propellant. More particularly, satisfactory dispersions may be formed where the medicament is selected from salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof.

There is thus provided in one aspect of the invention a pharmaceutical aerosol formulation which comprises particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates (for example hydrates) thereof and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant. By "substantially free of surfactant" is meant formulations which contain no significant amounts of surfactant, for example less than 0.0001% by weight of the medicament.

In an alternative embodiment the present invention provides a pharmaceutical aerosol formulation as hereinbefore defined with the proviso that when said formulation consists essentially of salbutamol and 1,1,1,2-tetrafluoroethane in a weight ratio of 0.05:18, said salbutamol is present in the form of a physiologically acceptable salt.

The particle size of the particulate (e.g. micronised) medicament should be such as to permit inhalation of substantially all of the medicament into the lungs upon administration of the aerosol formulation and will thus be less than 100 microns, desirably less than 20 microns, and preferably in the range 1-10 microns, e.g. 1-5 microns.

Suitable pharmaceutically acceptable salts of the medicaments of use in the formulations of the present invention include acid addition salts such as for example sulphates, hydrochlorides and xinafoates (1-hydroxy-2-naphthoate), amine salts or alkali metal salts (e.g. sodium). Salmeterol will preferably be in the form of its xinafoate salt and salbutamol will preferably be in the form of its sulphate salt.

The final aerosol formulation desirably contains 0.005-10% w/w, preferably 0.005-5% w/w, especially 0.01-1.0% w/w, of medicament relative to the total weight of the formulation.

The propellants for use in the invention may be any fluorocarbon or hydrogen-containing chlorofluorocarbon or mixtures thereof having a sufficient vapour pressure to render them effective as propellants. Preferably the propellant will be a non-solvent for the medicament. Suitable propellants include, for example, C₁₋₄hydrogen-containing chlorofluorocarbons such as CH₂ClF, CCIF₂CHClF, CF₃CHClF, CHF₂CCIF₂, CHClFCHF₂, CF₃CH₂Cl and CCIF₂CH₃; C₁₋₄hydrogen-containing fluorocarbons such as CHF₂CHF₂, CF₃CH₂F, CHF₂CH₃ and CF₃CHF₂CF₃; and perfluorocarbons such as CF₃CF₃ and CF₃CF₂CF₃.

Where mixtures of the fluorocarbons or hydrogen-containing chlorofluorocarbons are employed they may be mixtures of the above identified compounds or mixtures, preferably binary mixtures, with other fluorocarbons or hydrogen-containing chloro-fluorocarbons for example CHClF₂, CH₂F₂ and CF₃CH₃. Preferably a single fluorocarbon or hydrogen-containing chlorofluorocarbon is

employed as the propellant. Particularly preferred as propellants are C₁₋₄ hydrogen-containing fluorocarbons such as 1,1,1,2-tetrafluoroethane (CF₃CH₂F) and 1,1,1,2,3,3,3-heptafluoro-n-propane (CF₃CHF₂CF₃).

It is desirable that the formulations of the invention contain no components which may provoke the degradation of stratospheric ozone. In particular it is desirable that the formulations are substantially free of chlorofluorocarbons such as CCl₃F, CCl₂F₂ and CF₃CCl₃.

The propellant may additionally contain a volatile adjuvant such as a saturated hydrocarbon for example propane, n-butane, isobutane, pentane and isopentane or a dialkyl ether for example dimethyl ether. In general, up to 50% w/w of the propellant may comprise a volatile hydrocarbon, for example 1 to 30% w/w. However, formulations which are substantially free of volatile adjuvants are preferred.

It is further desirable that the formulations of the invention are substantially free of liquid components of higher polarity than the propellant employed. Polarity may be determined for example, by the method described in European Patent Application Publication No. 0327777. In particular formulations which are substantially free of alcohols such as ethanol are preferable. As used herein "substantially free" means less than 1% w/w based upon the fluorocarbon or hydrogen-containing chlorofluorocarbon, in particular less than 0.5% for example 0.1% or less.

A particularly preferred embodiment of the invention provides a pharmaceutical aerosol formulation consisting essentially of one or more particulate medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof, and one or more fluorocarbon or hydrogen-containing chlorofluorocarbon propellant.

It will be appreciated by those skilled in the art that the aerosol formulations according to the invention may, if desired, contain a combination of two or more active ingredients. Aerosol compositions containing two active ingredients (in a conventional propellant system) are known, for example, for the treatment of respiratory disorders such as asthma. Accordingly the present invention further provides aerosol formulations in accordance with the invention which contain two or more particulate medicaments. Medicaments may be selected from suitable combinations of the medicaments mentioned hereinbefore or may be selected from any other suitable drug useful in inhalation therapy and which may be presented in a form which is substantially completely insoluble in the selected propellant. Appropriate medicaments may thus be selected from, for example, analgesics, e.g. codeine, dihydromorphine, ergotamine, fentanyl or morphine; anginal preparations, e.g. diltiazem; antiallergics, e.g. cromoglycate, ketotifen or nedocromil; anti-infectives e.g. cephalosporins, penicillins, streptomycin, sulphonamides, tetracyclines and pentamidine; antihistamines, e.g. methapyrilene; anti-inflammatories, e.g. flunisolide, budesonide, tipredane or triamcinolone acetone; antitussives, e.g. noscapine; bronchodilators, e.g. ephedrine, adrenaline, fenoterol, formoterol, isoprenaline, metaproterenol, phenylephrine, phenylpropanolamine, pirbuterol, reproterol, rimiterol, terbutaline, isoetharine, tulobuterol, orciprenaline, or (-)-4-amino-3,5-dichloro- α -[[6-[2-(2-pyridinyl)ethoxy]hexyl]amino]methyl]benzane-methanol; diuretics, e.g. amiloride; anticholinergics e.g. ipratropium, atropine or oxitropium; hormones, e.g. cortisone, hydrocortisone or prednisolone; xanthines e.g. aminophylline, choline theophyllinate, lysine theophyllinate

or theophylline; and therapeutic proteins and peptides, e.g. insulin or glucagon. It will be clear to a person skilled in the art that, where appropriate, the medicaments may be used in the form of salts (e.g. as alkali metal or amine salts or as acid addition salts) or as esters (e.g. lower alkyl esters) or as solvates (e.g. hydrates) to optimise the activity and/or stability of the medicament and/or to minimise the solubility of the medicament in the propellant.

Particularly preferred aerosol formulations contain salbutamol (e.g. as the free base or the sulphate salt) or salmeterol (e.g. as the xinafoate salt) in combination with an anti-inflammatory steroid such as a beclomethasone ester (e.g. the dipropionate) or a fluticasone ester (e.g. the propionate) or an antiallergic such as cromoglycate (e.g. the sodium salt). Combinations of salmeterol and fluticasone propionate or beclomethasone dipropionate, or salbutamol and fluticasone propionate or beclomethasone dipropionate are preferred, especially salmeterol xinafoate and fluticasone propionate or salbutamol and beclomethasone dipropionate.

The formulations of the invention may be prepared by dispersal of the medicament in the selected propellant in an appropriate container, e.g. with the aid of sonication. The process is desirably carried out under anhydrous conditions to obviate any adverse effects of moisture on suspension stability.

The formulations according to the invention form weakly flocculated suspensions on standing but, surprisingly, these suspensions have been found to be easily redispersed by mild agitation to provide suspensions with excellent delivery characteristics suitable for use in pressurised inhalers, even after prolonged storage. Minimising and preferably avoiding the use of formulation excipients e.g. surfactants, cosolvents etc in the aerosol formulations according to the invention is also advantageous since the formulations may be substantially taste and odour free, less irritant and less toxic than conventional formulations.

The chemical and physical stability and the pharmaceutical acceptability of the aerosol formulations according to the invention may be determined by techniques well known to those skilled in the art. Thus, for example, the chemical stability of the components may be determined by HPLC assay, for example, after prolonged storage of the product. Physical stability data may be gained from other conventional analytical techniques such as, for example, by leak testing, by valve delivery assay (average shot weights per actuation), by dose reproducibility assay (active ingredient per actuation) and spray distribution analysis.

The particle size distribution of the aerosol formulations according to the invention is particularly impressive and may be measured by conventional techniques; for example by cascade impaction or by the "Twin Impinger" analytical process. As used herein reference to the "Twin Impinger" assay means "Determination of the deposition of the emitted dose in pressurised inhalations using apparatus A" as defined in British Pharmacopoeia 1988, pages A204-207, Appendix XVII C. Such techniques enable the "respirable fraction" of the aerosol formulations to be calculated. As used herein reference to "respirable fraction" means the amount of active ingredient collected in the lower impingement chamber per actuation expressed as a percentage of the total amount of active ingredient delivered per actuation using the twin impinger method described above. The formulations according to the invention have been found to have a respirable fraction of 20% or more by weight of the medicament, preferably 25 to 70%, for example 30 to 60%.

Optionally, the medicament may be surface-modified prior to its dispersion in the propellant by treatment with a

substantially non-polar liquid medium which is a non-solvent for the medicament. There is thus provided in a further aspect of the invention an aerosol formulation comprising particulate, surface-modified medicament, as defined herein, and a fluorocarbon or hydrogen-containing chlorofluorocarbon propellant, which formulation is substantially free of surfactant. By "surface-modified medicament" is meant particles of medicament selected from the group consisting of salmeterol, salbutamol, fluticasone propionate, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof which have been surface-modified by admixture with a substantially non-polar non-solvent liquid, followed by removal of the liquid. The substantially non-polar non-solvent liquid medium is conveniently an aliphatic hydrocarbon, e.g. a lower alkane, which is sufficiently volatile to permit its ready evaporation, e.g. at ambient temperature and pressure, after slurring with the medicament. The use of isopentane as liquid medium is particularly advantageous in this respect.

The medicament is desirably slurried with the liquid medium under anhydrous conditions to obviate any adverse effects of moisture on suspension stability. The slurry may advantageously be sonicated to maximise the surface-modifying effect of the treatment. The liquid may be removed by any convenient means for example by evaporation or by filtration followed by evaporation, provided that following treatment the medicament is substantially free of the liquid. The formulations of the invention will be substantially free of the non-solvent non-polar liquid. Surface-modified medicament prepared by the above-described process comprises a further aspect of the present invention.

The formulations according to the invention may be filled into canisters suitable for delivering pharmaceutical aerosol formulations. Canisters generally comprise a container capable of withstanding the vapour pressure of the propellant used such as a plastic or plastic-coated glass bottle or preferably a metal can, for example an aluminium can which may optionally be anodised, lacquer-coated and/or plastic-coated, which container is closed with a metering valve. The metering valves are designed to deliver a metered amount of the formulation per actuation and incorporate a gasket to prevent leakage of propellant through the valve. The gasket may comprise any suitable elastomeric material such as for example low density polyethylene, chlorobutyl, black and white butadiene-acrylonitrile rubbers, butyl rubber and neoprene. Suitable valves are commercially available from manufacturers well known in the aerosol industry, for example, from Valois, France (e.g. DF10, DF30, DF60), Bepak plc, UK (e.g. BK300, BK356) and 3M-Neotechnic Ltd, UK (e.g. Spraymiser.®).

Conventional bulk manufacturing methods and machinery well known to those skilled in the art of pharmaceutical aerosol manufacture may be employed for the preparation of large scale batches for the commercial production of filled canisters. Thus, for example, in one bulk manufacturing method a metering valve is crimped onto an aluminium can to form an empty canister. The particulate medicament is added to a charge vessel and liquified propellant is pressure filled through the charge vessel into a manufacturing vessel. The drug suspension is mixed before recirculation to a filling machine and an aliquot of the drug suspension is then filled through the metering valve into the canister. Typically, in batches prepared for pharmaceutical use, each filled canister is check-weighed, coded with a batch number and packed into a tray for storage before release testing.

Each filled canister is conveniently fitted into a suitable channelling device prior to use to form a metered dose

inhaler for administration of the medicament into the lungs or nasal cavity of a patient. Suitable channelling devices comprise for example a valve actuator and a cylindrical or cone-like passage through which medicament may be delivered from the filled canister via the metering valve to the nose or mouth of a patient e.g. a mouthpiece actuator. Metered dose inhalers are designed to deliver a fixed unit dosage of medicament per actuation or "puff", for example in the range of 10 to 5000 microgram medicament per puff.

Administration of medicament may be indicated for the treatment of mild, moderate or severe acute or chronic symptoms or for prophylactic treatment. It will be appreciated that the precise dose administered will depend on the age and condition of the patient, the particular particulate medicament used and the frequency of administration and will ultimately be at the discretion of the attendant physician. When combinations of medicaments are employed the dose of each component of the combination will in general be that employed for each component when used alone. Typically, administration may be one or more times, for example from 1 to 8 times per day, giving for example 1,2,3 or 4 puffs each time.

Suitable daily doses, may be, for example in the range 50 to 200 microgram of salmeterol, 100 to 1000 microgram of salbutamol, 50 to 2000 microgram of fluticasone propionate or 100 to 2000 microgram of beclomethasone dipropionate, depending on the severity of the disease.

Thus, for example, each valve actuation may deliver 25 microgram salmeterol, 100 microgram salbutamol, 25, 50, 125 or 250 microgram fluticasone propionate or 50, 100, 200 or 250 microgram beclomethasone dipropionate. Typically each filled canister for use in a metered dose inhaler contains 100, 160 or 240 metered doses or puffs of medicament.

The filled canisters and metered dose inhalers described herein comprise further aspects of the present invention.

A still further aspect of the present invention comprises a method of treating respiratory disorders such as, for example, asthma, which comprises administration by inhalation of an effective amount of a formulation as herein described.

The following non-limitative Examples serve to illustrate the invention.

EXAMPLE 1

Micronised salmeterol xinafoate (24 mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2 g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w salmeterol xinafoate.

EXAMPLE 2

Micronised salmeterol xinafoate (38.28 g) and 1,1,1,2-tetrafluoroethane (36.36 kg) were added to a pressure vessel and mixed with a high shear mixer for 20 minutes. Aliquots (18.2 g) of the suspension were filled into aluminium cans closed with a metering valve, filling under pressure through the valve using conventional filling equipment. The resulting inhalers contained 9.57 mg salmeterol xinafoate and delivered 25 microgram salmeterol (39.9 microgram salt) per actuation.

EXAMPLE 3

Micronised fluticasone propionate (24 mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-

7

tetrafluoroethane (18.2 g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w fluticasone propionate.

EXAMPLES 4 AND 5

Micronised fluticasone propionate (66 mg or 6.6 mg) was weighed directly into each of 100 open aluminium cans and a metering valve was then crimped into place on each can. 1,1,1,2-Tetrafluoroethane (18.2 g) was then added to each canister under pressure, through the valve, and each filled canister shaken to disperse the drug. The resulting inhalers contained 66 or 6.6 mg fluticasone propionate and delivered 250 or 25 microgram fluticasone propionate per actuation (Examples 4 and 5 respectively).

EXAMPLE 6

Micronised salbutamol (24 mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2 g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.132% w/w salbutamol.

EXAMPLES 7 AND 8

Micronised salbutamol (24 mg or 48 mg) was weighed directly into each of 3 open aluminium cans. 1,1,1,2-Tetrafluoroethane (18.2 g) was added to each can from a vacuum flask and a metering valve was then crimped into place. Each filled canister was then shaken in an ultrasonic bath for 8 minutes. The resulting inhalers contained 24 mg or 48 mg salbutamol and delivered 100 or 200 microgram salbutamol per actuation (Examples 7 and 8 respectively).

EXAMPLE 9

Micronised salbutamol sulphate (31.7 mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (18.2 g) was added from a vacuum flask. The bottle was quickly sealed with a blank aluminium ferrule. The resulting aerosol contained 0.174% w/w salbutamol sulphate.

EXAMPLE 10

Micronised salbutamol sulphate (31.7 mg) was weighed directly into each of 4 open aluminium cans. 1,1,1,2-Tetrafluoroethane (18.2 g) was added to each can from a vacuum flask and a metering valve was then crimped into place. Each filled canister was then shaken in an ultrasonic bath for 5 minutes. The resulting inhalers contained 31.7 mg salbutamol sulphate and delivered 100 microgram salbutamol per actuation.

EXAMPLE 11

Isopentane (25 ml) was added to micronised salmeterol xinafoate (0.5 g) to form a slurry, which was sonicated for 3 minutes. The resulting suspension was dried by evaporating the isopentane at ambient temperature to yield surface-modified salmeterol xinafoate. Samples of this product (11.6 mg) were weighed into aluminium aerosol cans and 1,1,1,2-tetrafluoroethane (18.2 g—99.95% w/w of total fill weight) was added to each can, whereafter suitable metering valves were crimped onto the cans, which were then each sonicated for 5 minutes. The resulting aerosols contained salmeterol in an amount equivalent to 240 actuations at 25 microgram per actuation.

8

EXAMPLE 12

Micronised beclomethasone dipropionate monohydrate (68 mg) was weighed into a clean, dry, plastic-coated glass bottle and 1,1,1,2-tetrafluoroethane (to 18.2 g) was added from a vacuum flask. The bottle was quickly sealed with a metering valve. The resulting aerosol dispensed 250 microgram beclomethasone dipropionate (as the monohydrate) per 75.8 mg actuation.

EXAMPLE 13

Micronised salmeterol xinafoate (9.57 mg) is weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4 g) added from a vacuum flask. A metering valve is crimped into place and the filled canister sonicated for five minutes. The aerosol delivers 25 microgram salmeterol per actuation.

EXAMPLE 14

Micronised fluticasone propionate (13.3 mg) is weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4 g) added from a vacuum flask. A metering valve is crimped into place and the filled canister sonicated for five minutes. The aerosol delivers 50 microgram fluticasone propionate per actuation.

EXAMPLE 15

Micronised salbutamol sulphate (29 mg) was weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4 g) added from a vacuum flask. A metering valve was crimped into place and the filled canister sonicated for five minutes. The aerosol delivered 100 microgram salbutamol per actuation.

EXAMPLE 16

Micronised beclomethasone dipropionate monohydrate (62 mg) was weighed directly into an aluminium can and 1,1,1,2,3,3,3-heptafluoro-n-propane (to 21.4 g) added from a vacuum flask. A metering valve was crimped into place and the filled canister sonicated for five minutes. The aerosol delivered 250 microgram beclomethasone dipropionate per actuation.

EXAMPLE 17

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.066	50 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

Micronised medicaments were weighed into an aluminium can, 1,1,1,2-tetrafluoroethane (18.2 g) was added from a vacuum flask and a metering valve was crimped into place.

EXAMPLE 18

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.165	125 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

Micronised medicaments were weighed into an aluminium can, 1,1,1,2-tetrafluoroethane (18.2 g) was added from a vacuum flask and a metering valve was crimped into place.

9
EXAMPLE 19

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.132	100 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

EXAMPLE 20

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.330	250 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

EXAMPLE 21

	Per Inhaler % w/w	Per Actuation
Salbutamol*	0.132	100 microgram
Fluticasone propionate	0.132	100 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

*as free base or an equivalent weight of salt e.g. sulphate

EXAMPLE 22

	Per Inhaler % w/w	Per Actuation
Salbutamol*	0.264	200 microgram
Fluticasone propionate	0.330	250 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

*as free base or an equivalent weight of salt e.g. sulphate

EXAMPLE 23

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Beclomethasone dipropionate	0.066	50 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

EXAMPLE 24

	Per Inhaler % w/w	Per Actuation
Salmeterol xinafoate	0.048	36.25 microgram
Fluticasone propionate	0.264	200 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

EXAMPLE 25

	Per Inhaler % w/w	Per Actuation
Salbutamol*	0.132	100 microgram
Beclomethasone dipropionate	0.066	50 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

*as free base or an equivalent weight of salt e.g. sulphate

10
EXAMPLE 26

	Per Inhaler % w/w	Per Actuation
5 Salbutamol*	0.264	200 microgram
Beclomethasone dipropionate	0.264	200 microgram
1,1,1,2-Tetrafluoroethane	to 100	to 75.8 mg

*as free base or an equivalent weight of salt e.g. sulphate

10 In Examples 19 to 26 micronised medicaments are weighed into aluminium cans, 1,1,1,2-tetrafluoroethane (18.2 g) is added from a vacuum flask, and metering valves are crimped into place.

We claim:

15 1. A canister suitable for delivering a pharmaceutical aerosol formulation for inhalation therapy which comprises a container capable of withstanding the vapor pressure of the propellant used, which container is closed with a metering valve and contains a pharmaceutical aerosol formulation
20 consisting essentially of a particulate medicament which is salmeterol or a physiologically acceptable salt or solvate thereof and 1,1,1,2-tetrafluoroethane as propellant, which formulation contains less than 0.0001% w/w surfactant based upon the weight of medicament, the particulate medi-
25 cament being present in an amount of 0.005 to 5% w/w relative to the total weight of the formulation and having a particle size of less than 100 microns.

2. A canister as claimed in claim 1 wherein the particulate medicament is present in an amount of 0.01 to 1% w/w
30 relative to the total weight of the formulation.

3. A canister suitable for delivering a pharmaceutical aerosol formulation for inhalation therapy which comprises a container capable of withstanding the vapor pressure of the propellant used, which container is closed with a metering
35 valve and contains a pharmaceutical aerosol formulation consisting of a particulate medicament which is salmeterol or a physiologically acceptable salt or solvate thereof and 1,1,1,2-tetrafluoroethane as propellant, the particulate medi-
cament being present in an amount of 0.005 to 5% w/w
40 relative to the total weight of the formulation and having a particle size of less than 100 microns.

4. A canister suitable for delivering a pharmaceutical aerosol formulation for inhalation therapy which comprises a container capable of withstanding the vapor pressure of the propellant used, which container is a plastics-coated alumi-
45 num can and is closed with a metering valve and contains a pharmaceutical aerosol suspension formulation consisting essentially of a particulate medicament which is salmeterol xinafoate and 1,1,1,2-tetrafluoroethane as propellant, which
50 formulation contains less than 0.0001% w/w surfactant based upon the weight of medicament, the particulate medi-
cament being present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation and having a particle size of less than 100 microns.

55 5. A canister suitable for delivering a pharmaceutical aerosol formulation for inhalation therapy which comprises a container capable of withstanding the vapor pressure of the propellant used, which container is closed with a metering valve and contains a pharmaceutical aerosol formulation
60 consisting essentially of a particulate medicament which is salmeterol or a physiologically acceptable salt or solvate thereof and a particulate medicament selected from the group consisting of salbutamol, beclomethasone dipropi-
onate and physiologically acceptable salts and solvates
65 thereof, and 1,1,1,2-tetrafluoroethane as propellant, which formulation contains less than 0.0001% w/w surfactant based upon the weight of medicament, the particulate medi-

caments being present in an amount of 0.005 to 5% w/w relative to the total weight of the formulation and having a particle size of less than 100 microns.

6. A canister suitable for delivering a pharmaceutical aerosol formulation for inhalation therapy which comprises a container capable of withstanding the vapor pressure of the propellant used, which container is closed with a metering valve and contains a pharmaceutical aerosol formulation consisting of a particulate medicament which is salmeterol or a physiologically acceptable salt or solvate thereof and a particulate medicament selected from the group consisting of salbutamol, beclomethasone dipropionate and physiologically acceptable salts and solvates thereof, and 1,1,1,2-tetrafluoroethane as propellant, the particulate medicaments being present in an amount of 0.005 to 5% w/w relative to the total weight of the formulation and having a particle size of less than 100 microns.

7. A canister as claimed in claim 5 wherein the formulation is free of surfactant.

8. A canister as claimed in claim 5 wherein the particulate medicaments are present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.

9. A canister as claimed in claim 6 wherein the particulate medicaments are present in an amount of 0.01 to 1% w/w relative to the total weight of the formulation.

10. A canister as claimed in claim 5 wherein the salmeterol is in the form of its xinafoate salt.

11. A canister as claimed in claim 6 wherein the salmeterol is in the form of its xinafoate salt.

12. A canister as claimed in claim 5 wherein the formulation has a respirable fraction of 20% or more by weight of the particulate medicaments.

13. A canister as claimed in claim 6 wherein the formulation has a respirable fraction of 20% or more by weight of the particulate medicaments.

14. A canister as claimed in claim 1 wherein the formulation is free of surfactant.

15. A canister as claimed in claim 1 wherein the container is plastics-coated, lacquer-coated or anodized.

16. A canister as claimed in claim 1 wherein the container is a metal can.

17. A canister as claimed in claim 1 wherein the container is a metal can which is plastics-coated.

18. A canister as claimed in claim 1 wherein the container is an aluminum can.

19. A canister as claimed in claim 1 wherein the container is an aluminum can which is plastics-coated.

20. A canister as claimed in claim 1 wherein the salmeterol is in the form of its xinafoate salt.

21. A metered dose inhaler which comprises a canister as claimed in claim 1 fitted into a channelling device for nasal or oral inhalation of the pharmaceutical aerosol formulation.

22. A canister as claimed in claim 3 wherein the container is plastics-coated, lacquer-coated or anodized.

23. A canister as claimed in claim 3 wherein the container is a metal can.

24. A canister as claimed in claim 3 wherein the container is a metal can which is plastics-coated.

25. A canister as claimed in claim 3 wherein the container is an aluminum can.

26. A canister as claimed in claim 3 wherein the container is an aluminum can which is plastics-coated.

27. A canister as claimed in claim 3 wherein the salmeterol or physiologically acceptable salt or solvate thereof is present in an amount of 0.01 to 1% w/w based on total weight of the formulation.

28. A canister as claimed in claim 3 wherein the salmeterol is in the form of its xinafoate salt.

29. A canister as claimed in claim 3 wherein the formulation has a respirable fraction of 20% or more by weight of salmeterol or physiologically acceptable salt or solvate thereof.

30. A metered dose inhaler which comprises a canister as claimed in claim 3 fitted into a channelling device for nasal or oral inhalation of the pharmaceutical aerosol formulation.

31. A canister as claimed in claim 1 wherein the pharmaceutical aerosol formulation is in a form for inhalation into lungs of a patient.

32. A canister as claimed in claim 3 wherein the pharmaceutical aerosol formulation is in a form for inhalation into lungs of a patient.

* * * * *