



US005733343A

United States Patent [19]
Möckli

[11] **Patent Number:** **5,733,343**
[45] **Date of Patent:** **Mar. 31, 1998**

[54] **PROCESS FOR DYEING KERATIN-CONTAINING FIBRES WITH CATIONIC DYES**

[75] **Inventor:** Peter Möckli, Sandgrubenstrasse, Switzerland

[73] **Assignee:** Ciba Specialty Chemicals Corporation, Tarrytown, N.Y.

[21] **Appl. No.:** 756,448

[22] **Filed:** Nov. 26, 1996

Related U.S. Application Data

[63] Continuation of Ser. No. 392,783, filed as PCT/EP94/02077, Jun. 27, 1994, abandoned.

[30] **Foreign Application Priority Data**

Jul. 5, 1993 [CH] Switzerland 2020/93

[51] **Int. Cl.⁶** **A61K 7/13**

[52] **U.S. Cl.** **8/426; 8/568; 8/573; 8/639; 8/655; 8/657; 8/659; 8/692**

[58] **Field of Search** **8/404, 405, 428, 8/565, 568, 570, 571, 572, 573, 670, 690, 691, 692, 916, 917, 639, 655, 657, 659**

[56] **References Cited**

U.S. PATENT DOCUMENTS

2,708,671 5/1955 Forone et al. 8/691
3,221,005 11/1965 Moore et al. 8/691
3,336,286 8/1967 Sartori 8/691

3,660,008 5/1972 Kissa 8/589
3,765,835 10/1973 Clarke et al. 8/589
3,787,178 1/1974 Renfrew 8/691
3,824,074 7/1974 Bugaut et al. 8/10
3,856,788 12/1974 Corbett et al. 260/244 R
3,869,454 3/1975 Lang et al. 8/405
3,985,499 10/1976 Lang et al. 8/10.1
4,181,499 1/1980 Koller et al. 8/568
4,349,348 9/1982 Tappe et al. 8/691
4,705,525 11/1987 Abel et al. 8/555
4,705,526 11/1987 Abel et al. 8/555

FOREIGN PATENT DOCUMENTS

2856225 7/1979 Germany .
3287520 12/1991 Japan .
1373081 11/1974 United Kingdom .

OTHER PUBLICATIONS

Chem. Abstracts, 112 : 145333b, Abstract of DE 3.829.870 Apr. 13, 1989.

Chem. Abstracts, 93 : 53774g, Abstract of JP 80/22.638, Feb. 18, 1980.

Primary Examiner—Paul Lieberman
Assistant Examiner—Caroline L. Dusheck
Attorney, Agent, or Firm—Kevin T. Mansfield

[57] **ABSTRACT**

Keratin-containing fibres, in particular human hair, are dyed using dyes of formulae (1) to (6) indicated in claim 1. These dyes make it possible to dye by the trichromatic principle even in dark shades.

22 Claims, No Drawings

PROCESS FOR DYEING KERATIN-CONTAINING FIBRES WITH CATIONIC DYES

This is a continuation of application Ser. No. 08/392,783, granted 35 USC 371 filing date of Feb. 28, 1995, now abandoned, originally International application PCT/EP 94/02077 filed Jun. 27, 1994.


The present invention relates to a process for dyeing keratin-containing fibres, in particular human hair, with cationic dyes.

By far the largest proportion of all hair dyeings are carried out, even today, using so-called "oxidation colours", which involves applying small, colourless precursor molecules to the hair and reacting them by an oxidation process to form larger, coloured molecules. Although this produces the most durable ("permanent") colourings, increasing reservations are being voiced about possible toxicological risks posed not only by the substances used as starting materials but also by the oxidation intermediate and end products, whose precise composition is virtually uncontrollable. Further disadvantages are the relatively complicated use and in particular also the hair damage due to the aggressive chemicals used.

The other, so-called "semipermanent" and "temporary" colourings involve the use of ready-prepared dyes, primarily uncharged disperse dyes and relatively sparingly water-soluble acid dyes. Cationic dyes, by contrast, play only a very minor part. As the terms "semipermanent" and "temporary" indicate, these colourings only have a medium to poor fastness level. Especially the cationic dyes have a reputation for poor hydrolysis and light resistance and for uneven colouring of the hair, for example between root and tip (see: John F. Corbett: The Chemistry of Hair-care Products, JSDC August 1976, p. 290). In addition, the known cationic dyes have an insufficient build-up; i.e., even if increased amounts are used, it is impossible to exceed a certain, relatively low, colour strength. For instance, it is not possible to achieve a deep black coloration with the most important cationic hair dyes Basic Yellow 57, Basic Red 76, Basic Blue 99, Basic Brown 16 and Basic Brown 17 which are used in practice. For the same reason it is difficult to tint relatively dark natural hair with these dyes.

It has now been found that surprisingly cationic dyes of the below-indicated formulae have none of these disadvantages. They can be used to achieve in a very simple way and under gentle conditions very deep dyeings having excellent light, shampooing and crock fastness properties. Owing to their extremely clean shades, they also extend the range of possible mixed shades considerably, especially in the direction of the increasingly important brilliant fashion colours.


The present invention accordingly provides a process for dyeing keratin-containing fibres, which comprises treating the fibres with a dye of the formula



(1)


-continued

(2)




5

(3)




10

(4)



15

(5)




20

25

or


(6)




30

where

35 X is —O—, —S— or



40 Y is —CH=,




45

—N=,

R is hydrogen, C₁–C₄alkyl, Cl or nitro,R' is hydrogen, C₁–C₄alkyl, Cl, nitro, amino, C₁–C₄monoalkylamino or di-C₁–C₄alkylamino,50 R₁ and R₂ are each independently of the other unsubstituted or OH—, C₁–C₄alkoxy-, halogen-, CN—, amino-, C₁–C₄monoalkylamino- or di-C₁–C₄alkylamino-substituted C₁–C₄alkyl,R₃ is hydrogen, C₁–C₄alkyl or CN,R₄ is unsubstituted or OH— or CN-substituted C₁–C₄alkyl,55 R₅ is hydrogen or C₁–C₄alkyl,R₆ and R₇ are each independently of the other hydrogen, C₁–C₄alkyl or C₁–C₄alkoxy, orR₅ and R₆ are together with the nitrogen and carbon atoms joining them together a 5- or 6-membered ring,60 R₈, R₉, R₁₀ and R₁₁ are each independently of the others hydrogen or C₁–C₄alkyl, with the proviso that at least one of these 4 substituents is C₁–C₄alkyl and that not all four substituents are ethyl,R₁₂ and R₁₃ are each independently of the other hydrogen, C₁–C₄alkyl or C₁–C₄alkoxy,

65 K is the radical of a coupling component of the aniline or phenol series or the radical of a heterocyclic coupling component,

K_1 is the radical of an aromatic or heterocyclic amine, and An^\ominus is a colourless anion, with the proviso that, in the dyes of the formula (1), K is not a radical of *N,N*-dimethylaniline when X is



Y is $-N=$ and R and R_1 are each methyl.

For the purposes of the present invention, alkyl radicals are generally straight-chain or branched C_1 - C_4 alkyl groups. Suitable are for example methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, *sec*-butyl or *tert*-butyl.

Suitable alkoxy radicals are those having 1 to 4 carbon atoms, e.g. methoxy, ethoxy, propoxy, isopropoxy, *n*-butoxy, isobutoxy or *tert*-butoxy.

Halogen is to be understood as meaning fluorine, bromine, iodine or in particular chlorine.

If R_5 and R_6 are combined with the nitrogen atom and two carbon atoms joining them together into a 5- or 6-membered ring, this ring may contain a further heteroatom, for example oxygen or sulfur. Moreover, the ring may be substituted, for example by hydroxyl, alkoxy, alkyl, halogen, CN or phenyl, or carry a further fused-on benzene ring. Preferred rings formed by R_5 , R_6 , the linked carbon atoms and the nitrogen atom are pyrroline, dihydrooxazine and di- or tetrahydropyridine rings carrying 0 to 4 methyl groups.


Suitable anions An^\ominus include organic as well as inorganic anions, for example chloride, bromide, sulfate, hydrogensulfate, methosulfate, phosphate, borotetrafluoride, carbonate, bicarbonate, oxalate, formate, acetate, propionate, lactate or complex anions, such as the anion of zinc chloride double salts.

The anion is generally given by the method of preparation. Preferred anions are chloride, sulfate, hydrogensulfate, methosulfate, phosphate, formate, acetate or lactate.


To dye by the process of the invention it is preferable to use a dye of the formula (1) where R' is hydrogen, C_1 - C_2 alkyl, amino, C_1 - C_2 monoalkylamino or di- C_1 - C_2 alkylamino or a dye of the formula (1) where R_1 is unsubstituted C_1 - C_4 alkyl.

It is likewise preferable to use dyes of the formula (2) where R is hydrogen or C_1 - C_4 alkyl or a dye of the formula (2) where R_1 is unsubstituted C_1 - C_4 alkyl.

Of the dyes of the formula (1), preference is given to those where X is








and especially those where X is



and Y is $-CH=$.

In the dyes of the formula (1), K is in particular the radical of a coupling component of the formula



where

35 R_{14} is hydrogen or unsubstituted or OH-, C_1 - C_4 alkoxy-, halogen-, CN-, amino-, C_1 - C_4 monoalkylamino- or di- C_1 - C_4 alkylamino-substituted C_1 - C_4 alkyl,

R_{15} and R_{16} are each independently of the other hydrogen, C_1 - C_4 alkyl, C_1 - C_4 alkoxy or halogen,

40 R_{17} and R_{18} are each independently of the other hydrogen, unsubstituted or OH-, C_1 - C_4 alkoxy-, halogen-, CN-, amino-, C_1 - C_4 monoalkylamino- or di- C_1 - C_4 alkylamino-substituted C_1 - C_4 alkyl, or

45 R_{17} and R_{18} are together with the nitrogen atom joining them together a 5- or 6-membered ring, or

R_{15} and R_{17} are together with the nitrogen and carbon atoms joining them together a 5- or 6-membered ring, or

R_{16} and R_{18} are together with the nitrogen and carbon atoms joining them together a 5- or 6-membered ring, and

50 R_{19} is hydrogen or unsubstituted or OH-, C_1 - C_4 alkoxy-, halogen-, CN-, amino-, C_1 - C_4 monoalkylamino- or di- C_1 - C_4 alkylamino-substituted C_1 - C_4 alkyl.


If R_{17} and R_{18} are to combine with the nitrogen atom joining them together into a 5- or 6-membered ring, this ring is in particular a pyrrolidine, piperidine, morpholine or piperazine ring. These rings can be further substituted, for example by C_1 - C_4 alkyl or C_1 - C_4 alkoxy. Preference, however, is given to the unsubstituted rings.

If R_{15} and R_{17} or R_{16} and R_{18} are combined with the nitrogen atom and the two carbon atoms joining them together into a 5- or 6-membered ring, this ring may contain a further heteroatom, for example oxygen or sulfur. Moreover, the ring may be substituted, for example by hydroxyl, alkoxy, alkyl, halogen or CN, or carry a further fused-on benzene ring. Preferred rings formed by R_{15} and R_{17} or R_{16} and R_{18} and the carbon atoms joining them together and the nitrogen atom are pyrroline, dihydroox-

5

azine and di- or tetrahydropyridine rings carrying 0 to 4 methyl groups.

In particular K is the radical of a coupling component of the formula



where

R_{14} is hydrogen or unsubstituted C_1-C_4 alkyl,

R_{15} and R_{16} are each independently of the other hydrogen, C_1-C_4 alkyl, C_1-C_4 alkoxy or halogen,

R_{17} and R_{18} are each independently of the other hydrogen or unsubstituted C_1-C_4 alkyl, or

R_{17} and R_{18} are together with the nitrogen atom joining them together a pyrrolidine, piperidine, morpholine or piperazine ring, or

R_{15} and R_{17} are together with the nitrogen and carbon atom joining them together a pyrrolidine, piperidine, morpholine or piperazine ring, or

R_{16} and R_{18} are together with the nitrogen and carbon atom joining them together a pyrrolidine, piperidine, morpholine or piperazine ring, and

R_{19} is hydrogen or unsubstituted C_1-C_4 alkyl.

Of very particular interest for the process of the invention are dyes of the formula (1) or (2) where K is the radical of a coupling component of the formula (7) or (8) where

R_{14} is methyl or ethyl,


R_{15} and R_{16} are each independently of the other hydrogen, methyl, ethyl, methoxy, ethoxy or chlorine,

R_{17} and R_{18} are each independently of the other hydrogen, methyl or ethyl, and

R_{19} is hydrogen, methyl or ethyl.

Preference is also given to using a dye of the formula (3), (4) or (5) where R_3 is hydrogen or methyl or a dye of the formula (3), (4) or (5) where R_4 is unsubstituted or hydroxyl-substituted C_1-C_4 alkyl, in particular methyl.


In the dyes of the formula (3) and (4), K_1 is in particular the radical of an amine of the formula



6

-continued

(14)



where

R_{15} and R_{16} are each independently of the other hydrogen, C_1-C_4 alkyl, C_1-C_4 alkoxy or halogen,

R_{17} and R_{18} are each independently of the other hydrogen, unsubstituted or OH—, C_1-C_4 alkoxy-, halogen-, CN—, amino-, C_1-C_4 monoalkylamino- or di- C_1-C_4 alkylamino-substituted C_1-C_4 alkyl, or

R_{17} and R_{18} are together with the nitrogen atom joining them together a 5- or 6-membered ring, or

R_{15} and R_{17} are together with the nitrogen and carbon atoms joining them together a 5- or 6-membered ring, or

R_{16} and R_{18} are together with the nitrogen and carbon atoms joining them together a 5- or 6-membered ring, and

R_{19} is hydrogen or unsubstituted or OH—, C_1-C_4 alkoxy-, halogen-, CN—, amino-, C_1-C_4 monoalkylamino- or di- C_1-C_4 alkylamino-substituted C_1-C_4 alkyl, and in particular the radical of an amine of the formula (12), (13) or (14), where

R_{15} and R_{16} are each independently of the other hydrogen, methyl, ethyl, methoxy, ethoxy or chlorine, or

R_{15} and R_{17} are together with the nitrogen and carbon atoms joining them together a pyrrolidine, piperidine, morpholine or piperazine ring,

R_{17} and R_{18} are each independently of the other hydrogen, methyl or ethyl, and

R_{19} is hydrogen, methyl or ethyl.

If the process of the invention is carried out using a dye of the formula (5), it is in particular a dye of the formula (5) where

R_5 is hydrogen or methyl and R_6 and R_7 are each independently of the other hydrogen, C_1-C_2 alkyl or C_1-C_2 alkoxy,

or R_5 and R_6 are together with the nitrogen and carbon atoms joining them together a pyrrolidine, piperidine, morpholine or piperazine ring.

Of the dyes of the formula (6), preference is given to using those where

R_8 , R_9 , R_{10} and R_{11} are each independently of the others hydrogen or C_1-C_2 alkyl, with the proviso that at least one of these 4 substituents is C_1-C_2 alkyl and that not all four substituents are ethyl, and






R_{12} and R_{13} are each independently of the other hydrogen, C_1-C_2 alkyl or C_1-C_2 alkoxy.

The dyes used according to the invention are known or can be prepared in a manner known per se.


The present invention furthermore provides a process for dyeing keratin-containing fibres, which comprises treating the fibres with a mixture of at least two cationic dyes having a delocalized positive charge and a cation weight below 300, preferably below 280.

Preference is given to using a mixture of at least three cationic dyes with a delocalized positive charge and a cation weight below 280 and in particular a mixture of a yellow, a red and a blue cationic dye with delocalized positive charge and a cation weight below 280.


A very particularly preferred embodiment of the novel process for dyeing keratin-containing fibres comprises treating the fibres with a mixture of at least two cationic dyes of the formula




or



where
X is —O—, —S—, or



Y is —CH=,



or —N=,

R is hydrogen, C₁-C₄alkyl, Cl or nitro,
R' is hydrogen, C₁-C₄alkyl, Cl, nitro, amino,
C₁-C₄monoalkylamino or di-C₁-C₄alkylamino,
R₁ and R₂ are each independently of the other unsubstituted
or OH—, C₁-C₄alkoxy-, halogen-, CN—, amino-,
C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-
substituted C₁-C₄alkyl,
R₃ is hydrogen, C₁-C₄alkyl or CN,
R₄ is unsubstituted or OH— or CN-substituted C₁-C₄alkyl,
R₅ is hydrogen or C₁-C₄alkyl,
R₆ and R₇ are each independently of the other hydrogen,
C₁-C₄alkyl or C₁-C₄alkoxy, or
R₅ and R₆ are together with the nitrogen and carbon atoms
joining them together a 5- or 6-membered ring.

- (1) R₈, R₉, R₁₀ and R₁₁ are each independently of the others
hydrogen or C₁-C₄alkyl,
R₁₂ and R₁₃ are each independently of the other hydrogen,
C₁-C₄alkyl or C₁-C₄alkoxy,
5 K is the radical of a coupling component of the aniline series
or the radical of a heterocyclic coupling component,
K₁ is the radical of an aromatic or heterocyclic amine, and
An[⊖] is a colourless anion.

- (2) The process of the invention is suitable for dyeing furs
10 and also animal and human hair, especially live human hair
and domestic animals' hair. As a consequence of the high
affinity and the good water solubility of the dyes used, it is
possible to do the dyeing at room temperature from aqueous
solutions without any assistants whatsoever.

- 15 However, it is also possible to use any assistants custom-
ary for cationic dyes used in the dyeing of hair, for example
wetting agents, swelling agents, penetration aids or scents.
In addition, the dyes can be incorporated into shampoos,
creams, gels or pastes. Such cosmetic formulations for
20 dyeing hair comprising at least one dye of the above-
indicated formulae (1) to (6) and also assistants form a
further part of the subject-matter of the present invention.

- (4) It has been found that the dyeing effect of the dyes used
depends relatively little on the formulation of the dyes.

- 25 A particular advantage of the dyes used according to the
invention for dyeing hair is that, owing to the good build-up
of the dyes, the colourings can be prepared by the trichro-
matic principle; that is, it is possible by using a yellow, a red
and a blue dye in suitable mixtures of these dyes to achieve
30 virtually all shades. In addition, exact prediction of the
shades obtained is possible, which is not the case with the
so-called "oxidation dyes" owing to the varying composition
of the end products.


- Using colorimetric methods of measurement it is also
35 possible to obtain on natural, unbleached hair predicted
shades having regard to the hair's natural colour by deter-
mining its yellow, red and blue content and deducting it from
the recipe of the desired shade. This is not feasible with the
hair dyes previously used.

- 40 The colourings obtained are crock-, water-, wash- and
light-fast and stable to permanent-deformation agents, for
example thioglycolic acid.

- The Examples which follow illustrate the invention. Parts
and percentages are by weight. The temperatures are given
45 in degrees Celsius.

EXAMPLE 1

A braid-sewn strand of blond, natural, untreated human
hair is dyed at 25° C. for 5 minutes in a conventional manner
50 with a dye emulsion containing 0.1% of the dye of the
formula



- 55 3.5% of Cetearyl Alcohol
1.0% of Cetareth 80
0.5% of glyceryl mono-di-stearate
60 3.0% of stearamide DEA
1.0% of stearamphopropylsulfonate
0.5% of polyquaternium-6 and
65 water to 100%.


Then the hair is thoroughly rinsed with water and air-
dried. The result is an intensive brilliant yellow colouring

9

which is many times stronger than a colouring prepared with Basic Yellow 57 in the same way. The light, shampooing and friction fastness properties of the colouring according to the invention are excellent.

EXAMPLE 2


Example 1 is repeated with the dye of the formula



affording an intensively golden yellow colouring with likewise excellent fastness properties.

EXAMPLE 3


A 1% solution of the dye of the formula



in a surfactant base containing 10% of cocoamphoglycinate and 90% of water is applied to Chinese, bleached yak hair at 25° C. for 5 minutes, and then the hair is thoroughly rinsed and air-dried. The intensively scarlet red colouring obtained is many times stronger than a comparative dyeing with Basic Red 76 and also of distinctly better light fastness.

EXAMPLE 4

A strand of medium brown, untreated human hair is dyed for 5 minutes at room temperature with a dye emulsion containing 0.1% of the dye of the formula



and otherwise having the same composition as the dye emulsion of Example 1. Then the strand of hair is thoroughly


10

rinsed with water and air-dried. The result is a very attractive chestnut-brown shade of the kind which is frequently desired. This shade is impossible to achieve with Basic Red 76 on account of the insufficient build-up of this dye.

5

EXAMPLE 5

A strand of bleached yak hair is dyed for 5 minutes at 25° C. with a dye emulsion which contains 0.1% of the dye of the formula




and otherwise has the same composition as the dye emulsion of Example 3. Then the strand of hair is thoroughly rinsed with water and air-dried. The blue colouring obtained is very significantly stronger and more brilliant than a dyeing with Basic Blue 99 prepared in the same way.

25

EXAMPLE 6

Example 4 is repeated with the red dye replaced by the blue dye of the formula



This shifts the original brown of the hair to a mattish brown hue which hides very well undesirable rust-red shades as frequently obtained following oxidation dyeings and lightenings. The scope for these tinting uses is much less with Basic Blue 99.

45

EXAMPLES 7-70










The method of Examples 1-3 is applied with the dyes listed below in the table, affording colourings on the hair in the specified hues.

Example	Dye	Hue
7		yellow
8		yellow

-continued

Example	Dye	Hue
9		yellow
10		yellow
11		yellow
12		yellow
13		yellow
14		yellow
15		yellow
16		orange
17		greenish yellow
18		greenish yellow
19		orange

-continued

Example	Dye	Hue
20		yellowish orange Cl [⊖]
21		yellow CH ₃ SO ₄ [⊖]
22		greenish yellow Cl [⊖]
23		reddish orange Cl [⊖]
24		red Cl [⊖]
25		scarlet Cl [⊖]
26		golden yellow Cl [⊖]
27		red Cl [⊖]
28		red Cl [⊖]

-continued

Example	Dye	Hue
29		red
30		reddish orange
31		red
32		red
33		red
34		blue
35		blue
36		blue










-continued

Example	Dye	Hue
37		blue Cl [⊖]
38		blue Cl [⊖]
39		blue Cl [⊖]
40		blue Cl [⊖]
41		blue Cl [⊖]
42		blue Cl [⊖]
43		blue Cl [⊖]
44		blue CH ₃ SO ₄ [⊖]
45		blue Cl [⊖]

-continued

Example	Dye	Hue
46		orange
47		orange
48		orange
49		reddish orange
50		orange
51		ruby
52		scarlet
53		scarlet





-continued

Example	Dye	Hue
54		scarlet Cl [⊖]
55		scarlet Cl [⊖]
56		scarlet Cl [⊖]
57		scarlet Cl [⊖]
58		scarlet Cl [⊖]
59		violet Cl [⊖]
60		violet Cl [⊖]
61		violet Cl [⊖]
62		blue CH ₃ SO ₄ [⊖]

-continued

Example	Dye	Hue
63		blue
64		blue
65		bluish violet
66		bluish violet
67		blue
68		violet
69		violet
70		violet
71		blue

-continued

Example	Dye	Hue
72		blue
73		blue
74		blue
75		red

EXAMPLE 76

A braided strand of blond, natural, untreated human hair is treated at 25° C. for 5 minutes with a dye emulsion which has the same composition as the emulsion in Example 1 but contains as dyes 0.11% of the dye of Example 4 and 0.10% of the dye of Example 5. After the strand of hair has been thoroughly rinsed with water and dried, it has a deep violet colour with very good fastness properties.

EXAMPLE 77

Example 76 is repeated with the dyes replaced by 0.08% of the dye of Example 1 and 0.06% of the dye of Example 5, affording a very brilliant green colouring on the hair.

EXAMPLE 78

0.02% of the dye of Example 1 and 0.08% of the dye of Example 5 are dissolved in a surfactant base comprising a 10% aqueous solution of cocoamphoglycinate and this solution is used to dye a strand of bleached yak hair at room temperature for 5 minutes. A bright, brilliant turquoise shade is obtained on the hair.

EXAMPLE 79

Blond, untreated human hair is treated for 20 minutes at room temperature with a dye emulsion which has the same composition as the emulsion in Example 1 but contains as dyes 0.2% of the dye of Example 1, 0.1% of the dye of Example 4 and 0.17% of the dye of Example 6. Thorough rinsing and drying of the hair leaves a deep black colouring having good fastness properties.

35

EXAMPLE 80

Example 79 is repeated with the dyes replaced by a dye mixture containing

- 0.138% of the dye of Example 2,
 - 0.082% of the dye of Example 4 and
 - 0.026% of the dye of Example 6,
- affording a chestnut brown colouring.

EXAMPLE 81

Olive-coloured hair is obtained on repeating Example 79 with the following dye mixture:

- 0.13% of the dye of Example 2,
- 0.006% of the dye of Example 4 and
- 0.032% of the dye of Example 6.

EXAMPLE 82

Example 81 is repeated with a dye mixture containing

- 0.01% of the dye of Example 2,
 - 0.11% of the dye of Example 4 and
 - 0.21% of the dye of Example 6,
- affording a dark navy colouring on the hair.

EXAMPLE 83

A surfactant base comprising a 10% aqueous solution of cocoamphoglycinate is used to dissolve

- 0.036% of the dye of Example 1,
- 0.034% of the dye of Example 2 and
- 0.06% of the dye of Example 3






60

65


and this solution is used to treat a strand of bleached yak hair for 10 minutes at 25° C. Rinsing and drying leaves a luminously orange dyeing having excellent light, shampooing and friction fastness properties.

What is claimed is:

1. A process for dyeing fibres of human hair, which comprises treating the fibres with a tinctorially effective amount of a dye of the formula




or




where

X is



Y is $-\text{CH}=\text{}$ or



R is hydrogen, C₁-C₄alkyl, Cl or nitro,

R' is hydrogen, C₁-C₄alkyl, Cl, nitro, amino, C₁-C₄monoalkylamino or di-C₁-C₄alkylamino,

R₁ and R₂ are each independently of the other unsubstituted C₁-C₄alkyl or OH-, C₁-C₄alkoxy-, halogen-, CN-, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl,

R₃ is hydrogen, C₁-C₄alkyl or CN,

R₄ is unsubstituted C₁-C₄alkyl or OH-, or CN-substituted C₁-C₄alkyl,

R₅ is hydrogen or C₁-C₄alkyl,

R₆ and R₇ are each independently of the other hydrogen, C₁-C₄alkyl or C₁-C₄alkoxy, or



R₅ and R₆ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring,

R₈, R₉, R₁₀ and R₁₁ are each independently of the others hydrogen or C₁-C₄alkyl, with the proviso that at least one of these 4 substituents is C₁-C₄alkyl and that not all four substituents are ethyl,

R₁₂ and R₁₃ are each independently of the other hydrogen, C₁-C₄alkyl or C₁-C₄alkoxy,

K is the radical of a coupling component of the aniline or phenol series or the radical of a heterocyclic coupling component,

K₁ is the radical of an amine of the formula



where

R₁₅ and R₁₆ are each independently of the other hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy or halogen,

R₁₇ and R₁₈ are each independently of the other hydrogen, unsubstituted C₁-C₄alkyl or OH-, C₁-C₄alkoxy-, halogen-, CN-, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl, or

R₁₇ and R₁₈ together with the nitrogen atom joining them together form a 5- or 6-membered ring, or

R₁₅ and R₁₇ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring, or

R₁₆ and R₁₈ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring, and

R₁₉ is hydrogen or unsubstituted C₁-C₄alkyl or OH-, C₁-C₄alkoxy-, halogen-, CN-, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl, and

An[⊖] is a colourless anion.





2. A process according to claim 1, wherein the dye is of the formula (2) where R is hydrogen or C₁-C₄alkyl.

3. A process according to claim 1, wherein the dye is of the formula (1) or (2) where R₁ is unsubstituted C₁-C₄alkyl.


4. A process according to claim 1, wherein the dye is of the formula (1) where R₁ is unsubstituted C₁-C₄alkyl.

5. A process according to claim 1, wherein the dye is of the formula (1) where Y is $-\text{CH}=\text{}$.

6. A process according to claim 1, wherein the dye is of the formula (1) or (2) where K is the radical of a coupling component of the formula



or



where

R₁₄ is hydrogen or unsubstituted C₁-C₄alkyl or OH—, C₁-C₄alkoxy-, halogen-, CN—, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl,

R₁₅ and R₁₆ are each independently of the other hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy or halogen,

R₁₇ and R₁₈ are each independently of the other hydrogen, unsubstituted C₁-C₄alkyl or OH—, C₁-C₄alkoxy-, halogen-, CN—, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl, or


R₁₇ and R₁₈ together with the nitrogen atom joining them together form a 5- or 6-membered ring, or

R₁₅ and R₁₇ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring, or


R₁₆ and R₁₈ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring, and

R₁₉ is hydrogen or unsubstituted C₁-C₄alkyl or OH—, C₁-C₄alkoxy-, halogen-, CN—, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl.

7. A process according to claim 6, wherein the dye is of the formula (1) where K is the radical of a coupling component of the formula



or



where

R₁₄ is hydrogen or unsubstituted C₁-C₄alkyl, R₁₅ and R₁₆ are each independently of the other hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy or halogen,

R₁₇ and R₁₈ are each independently of the other hydrogen or unsubstituted C₁-C₄alkyl, or

R₁₇ and R₁₈ together with the nitrogen atom joining them together form a pyrrolidine, piperidine, morpholine or piperazine ring, or

R₁₅ and R₁₇ together with the nitrogen and carbon atom joining them together form a pyrrolidine, piperidine, morpholine or piperazine ring, or

R₁₆ and R₁₈ together with the nitrogen and carbon atom joining them together form a pyrrolidine, piperidine, morpholine or piperazine ring.

8. A process according to claim 7, wherein the dye is of the formula (1) or (2) where K is the radical of a coupling component of the formula (7) or (8) where

R₁₄ is methyl or ethyl,

R₁₅ and R₁₆ are each independently of the other hydrogen, methyl, ethyl, methoxy, ethoxy or chlorine, and

R₁₇ and R₁₈ are each independently of the other hydrogen, methyl or ethyl.

9. A process according to claim 1, wherein the dye is of the formula (3), (4) or (5) where R₃ is hydrogen or methyl.

10. A process according to claim 1, wherein the dye is of the formula (3), (4) or (5) where R₄ is unsubstituted C₁-C₄alkyl or hydroxyl-substituted C₁-C₄alkyl.

11. A process according to claim 1, wherein the dye is of the formula (3) or (4) where K₁ is the radical of an amine of the formula (12), (13) or (14) where

R₁₅ and R₁₆ are each independently of the other hydrogen, methyl, ethyl, methoxy, ethoxy or chlorine, or

R₁₅ and R₁₇ together with the nitrogen and carbon atoms joining them together form a pyrrolidine, piperidine, morpholine or piperazine ring,

R₁₇ and R₁₈ are each independently of the other hydrogen, methyl or ethyl, and

R₁₉ is hydrogen, methyl or ethyl.

12. A process according to claim 1, wherein the dye is of the formula (5) where

R₅ is hydrogen or methyl and R₆ and R₇ are each independently of the other hydrogen, C₁-C₂alkyl or C₁-C₂alkoxy, or

R₅ and R₆ together with the nitrogen and carbon atoms joining them together form a pyrrolidine, piperidine, morpholine or piperazine ring.

13. A process according to claim 1, wherein the dye is of the formula (6) where







R₈, R₉, R₁₀ and R₁₁ are each independently of the others hydrogen or C₁-C₂alkyl, with the proviso that at least one of these 4 substituents is C₁-C₂alkyl and that not all four substituents are ethyl, and

R₁₂ and R₁₃ are each independently of the other hydrogen, C₁-C₂alkyl or C₁-C₂alkoxy.

14. A process according to claim 1, wherein the dye is of the formula (1) where R' is hydrogen, C₁-C₂alkyl, amino, C₁-C₂monoalkylamino or di-C₁-C₂alkylamino.


15. A process for dyeing hairs of humans according to claim 1, wherein predetermined shades are determined with colorimetric methods of measurement.

16. A process for dyeing fibres of human hair, wherein the fibres are treated with a tinctorially effective amount of a mixture of cationic dyes of at least two of the formulae




where

X is



Y is —CH= or



R is hydrogen, C₁-C₄alkyl, Cl or nitro,

R' is hydrogen, C₁-C₄alkyl, Cl, nitro, amino, C₁-C₄monoalkylamino or di-C₁-C₄alkylamino,

R₁ and R₂ are each independently of the other unsubstituted C₁-C₄alkyl or OH—, C₁-C₄alkoxy-, halogen-, CN—, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl,

R₃ is hydrogen, C₁-C₄alkyl or CN,

R₄ is unsubstituted C₁-C₄alkyl or OH— or CN-substituted C₁-C₄alkyl,

R₅ is hydrogen or C₁-C₄alkyl,

R₆ and R₇ are each independently of the other hydrogen, C₁-C₄alkyl or C₁-C₄alkoxy, or




R₅ and R₆ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring,

R₈, R₉, R₁₀ and R₁₁ are each independently of the others hydrogen or C₁-C₄alkyl,

R₁₂ and R₁₃ are each independently of the other hydrogen, C₁-C₄alkyl or C₁-C₄alkoxy,

K is the radical of a coupling component of the aniline series or the radical of a heterocyclic coupling component,

K₁ the radical of an amine of the formula



where

R₁₅ and R₁₆ are each independently of the other hydrogen, C₁-C₄alkyl, C₁-C₄alkoxy or halogen,

R₁₇ and R₁₈ are each independently of the other hydrogen, unsubstituted C₁-C₄alkyl or OH—, C₁-C₄alkoxy-, halogen-, CN—, amino-, C₁-C₄monoalkylamino- or di-C₁-C₄alkylamino-substituted C₁-C₄alkyl, or

R₁₇ and R₁₈ together with the nitrogen atom joining them together form a 5- or 6-membered ring, or

R₁₅ and R₁₇ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring, or

R₁₆ and R₁₈ together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring, and

33

R_{19} is hydrogen or unsubstituted C_1-C_4 alkyl or $OH-$, C_1-C_4 alkoxy-, halogen-, $CN-$, amino-, C_1-C_4 monoalkylamino- or di- C_1-C_4 alkylamino-substituted C_1-C_4 alkyl, and

An^\ominus is a colourless anion.

17. A process according to claim 16, wherein the fibres are treated with a mixture of at least two cationic dyes having a delocalized positive charge and a cation molecular weight below 280.




18. A process according to claim 17, wherein the fibres are treated with a mixture of at least three cationic dyes having a delocalized positive charge and a cation molecular weight below 280.

19. A process according to claim 18, wherein the fibres are treated with a mixture of a yellow, a red and a blue cationic dye having a delocalized positive charge and a cation molecular weight below 280.

20. A process for dyeing hairs of humans according to claim 16, which comprises applying to the hairs a mixture of ready-prepared dyes of at least two of the formulae (1) to (6) and wherein predeterminable shades are determined with colorimetric methods of measurement.



21. A process for dyeing hairs of humans according to claim 20, which comprises applying to the hairs a mixture of a yellow, a red and a blue dye, and wherein predeterminable shades are determined with colorimetric methods of measurement.

22. A process for dyeing fibres of human hair, wherein the fibres are treated with a tinctorially effective amount of a mixture of cationic dyes of at least two of the formulae



34

-continued



where

R is hydrogen, C_1-C_4 alkyl, Cl or nitro,

R_1 and R_2 are each independently of the other unsubstituted C_1-C_4 alkyl or $OH-$, C_1-C_4 alkoxy-, halogen-, $CN-$, amino-, C_1-C_4 monoalkylamino- or di- C_1-C_4 alkylamino-substituted C_1-C_4 alkyl,

R_3 is hydrogen, C_1-C_4 alkyl or CN ,

R_4 is unsubstituted C_1-C_4 alkyl or $OH-$ or CN -substituted C_1-C_4 alkyl,

R_5 is hydrogen or C_1-C_4 alkyl,

R_6 and R_7 are each independently of the other hydrogen, C_1-C_4 alkyl or C_1-C_4 alkoxy, or

R_5 and R_6 together with the nitrogen and carbon atoms joining them together form a 5- or 6-membered ring,

R_8 , R_9 , R_{10} and R_{11} are each independently of the others hydrogen or C_1-C_4 alkyl,

R_{12} and R_{13} are each independently of the other hydrogen, C_1-C_4 alkyl or C_1-C_4 alkoxy,

K is the radical of a coupling component of the aniline series or the radical of a heterocyclic coupling component,

K_1 is the radical of an aromatic or heterocyclic amine, and An^\ominus is a colourless anion.

* * * * *