



US005744343A

United States Patent [19][11] **Patent Number:** **5,744,343****Draetta et al.**[45] **Date of Patent:** **Apr. 28, 1998**[54] **UBIQUITIN CONJUGATING ENZYMES**[75] Inventors: **Giulio Draetta**, Winchester; **Mark Rolfe**, Newton Upper Falls; **Jens W. Eckstein**, Cambridge; **Guillaume Cottarel**, Chestnut Hill, all of Mass.[73] Assignee: **Mitotix, Inc.**, Cambridge, Mass.[21] Appl. No.: **305,520**[22] Filed: **Sep. 13, 1994****Related U.S. Application Data**

[63] Continuation-in-part of Ser. No. 247,904, May 23, 1994, which is a continuation-in-part of Ser. No. 176,937, Jan. 4, 1994, abandoned.

[51] **Int. Cl.**⁶ **C12N 9/10**; C12N 1/20;
C12N 15/00; C07H 21/04[52] **U.S. Cl.** **435/193**; 435/252.3; 435/254.11;
435/320.1; 435/325; 536/23.2; 536/23.4[58] **Field of Search** 435/69.1, 193,
435/252.3, 320.1, 254.11, 325; 536/23.2,
24.3[56] **References Cited****FOREIGN PATENT DOCUMENTS**WO 89/09829 10/1989 WIPO .
WO 89/12678 12/1989 WIPO .
WO 91/17245 11/1991 WIPO .
WO 92/20804 11/1992 WIPO .
WO 93/09235 5/1993 WIPO .**OTHER PUBLICATIONS**Adams et al. (1993) "Rapid cDNA sequencing (expressed sequence tags) from a directionally cloned human infant brain cDNA library" *Nature Genetics* 4:373-380.Band et al. (1991) "Loss of p53 Protein in Human Papillomavirus Type 16 E6-Immortalized Human Mammary Epithelial Cells" *Journal of Virology*, 65(12):6671-6676.Berleth et al. (1992) "Inhibition of Ubiquitin-Protein Ligase (E3) by Mono- and Bifunctional Phenylarsenoxides" *The Journal of Biological Chemistry* 267(23):16403-16411.Bissonnette et al. (1992) "Apoptotic cell death induced by c-myc is inhibited by bcl-2" *Nature* 359:552-556.Chen et al. (1993) "Multiple Ubiquitin-Conjugating Enzymes Participate in the In Vivo Degradation of the Yeast MAT α 2 Repressor" *Cell* 74:357-369.Cook et al. (1992) "Structure of a Diubiquitin Conjugate and a Model for Interaction with Ubiquitin Conjugating Enzyme(E2)" *The Journal of Biological Chemistry* 267(23):16467-16471.Cook et al. (1993) "Tertiary Structures of Class I Ubiquitin-Conjugating Enzymes Are Highly Conserved: Crystal Structure of Yeast Ubc4" *Biochemistry* 32(50):13809-13817.Crook et al. (1991) "Degradation of p53 Can Be Targeted by HPV E6 Sequences Distinct from Those Required for p53 Binding and Trans-Activation" *Cell* 67:547-556.Dohmen et al. (1991) "The N-end rule is mediated by the UBC2(RAD6) ubiquitin-conjugating enzyme" *Proceedings of the National Academy of Sciences* 88:7351-7355.Ellison et al. (1991) "Epitope-tagged Ubiquitin" *J Biol Chem* 266(31):21150-21157.Evan et al. (1992) "Induction of Apoptosis in Fibroblasts by c-myc Protein" *Cell* 69:119-128.Girod et al. (1993) "Homologs of the essential ubiquitin-conjugating enzymes UBC1, 4, and 5 in yeast are encoded by a multigene family in *Arabidopsis thaliana*" *The Plant Journal* 3(4):545-552.Glotzer et al. (1991) "Cyclin is degraded by the ubiquitin pathway" *Nature* 349:132-138.Hartwell et al. (1992) "Defects in a Cell Cycle Checkpoint May Be Responsible for the Genomic Instability of Cancer Cells" *Cell* 71:543-546.Hershko et al. (1993) "Components of Ubiquitin-Protein Ligase System" *J Biol Chem* 258(13):8206-8214.Hochstrasser et al. (1990) "In Vivo Degradation of a Transcriptional Regulator: The Yeast α 2 Repressor" *Cell* 61:697-708.Hofer et al. (1991) "Purification and partial characterization of ubiquitin-activating enzyme from *Saccharomyces cerevisiae*" *FEBS Letters* 289(1):54-58.Holloway et al. (1993) "Anaphase Is Initiated by Proteolysis Rather Than by the Inactivation of Maturation-Promoting Factor" *Cell* 73:1393-1402.Huibregtse et al. (1993) "Localization of the E6-AP Regions That Direct Human Papillomavirus E6 Binding, Association with p53, and Ubiquitination of Associated Proteins" *Mol Cell Biol* 13(8):pp. 4918-4927.Huibregtse et al. (1993) "Cloning and Expression of the cDNA for E6-AP, a Protein That Mediates the Interaction of the Human Papillomavirus E6 Oncoprotein with p53" *Mol Cell Biol* 13(2):775-784.Huibregtse et al. (1991) "A cellular protein mediates association of p53 with the E6 oncoprotein of human papillomavirus types 16 or 18" *EMBO J* 10(3):4129-4135.Hupp et al. (1992) "Regulation of the Specific DNA Binding Function of p53" *Cell* 71:875-886.Jacobs et al. (1993) "Rapid Assessment of Drug Susceptibilities of *Mycobacterium tuberculosis* by Means of Luciferase Reporter Phages" *Science* 260:819-822.Jentsch et al. (1992) "The Ubiquitin-conjugation System" *Ann Rev Genet* 26:179-207.Klemperer et al. (1989) "A Novel, Arsenite-Sensitive E2 of the Ubiquitin Pathway: Purification and Properties" *Biochemistry* 28:6035-6041.Koken et al. (1991) "Structural and Functional conservation of two human homologs of the yeast DNA repair gene RAD6" *PNAS* 88:8865-8869.Milner et al. (1990) "p53 Is Associated with p34^{cdc2} Transformed Cells" *EMBO J* 9(9):2885-2889.

(List continued on next page.)

Primary Examiner—Robert A. Wax**Assistant Examiner**—Lisa J. Hobbs**Attorney, Agent, or Firm**—Matthew P. Vincent, Esq.; Beth E. Arnold, Esq.; Foley, Hoag & Eliot LLP[57] **ABSTRACT**

The present invention concerns three ubiquitin-conjugating enzymes.

27 Claims, 41 Drawing Sheets

OTHER PUBLICATIONS

- Münger et al. (1989) "The E6 and E7 Genes of the Human Papillomavirus Type 16 Together Are Necessary and Sufficient for Transformation of Primary Human Keratinocytes" *J Virol* 63(10):4417-4421.
- Nepveu et al. (1987) "Alternative Modes of c-myc regulation in growth factor-stimulated and differentiating cells" *Oncogene* 1:243-250.
- Rechsteiner et al. (1991) "Natural Substrates of the Ubiquitin Proteolytic Pathway" *Cell* 66:615-618.
- Schärer et al. (1992) "Mammalian p53 can function as a transcription factor in yeast" *Nucleic Acids Research* 20(7):1539-1545.
- Scheffner et al. (1992) "Interaction of the Human Papillomavirus Type 16 E6 Oncoprotein with Wild-Type and Mutant Human p53 Proteins" *J Virol* 66(8):5100-5105.
- Scheffner et al. (1990) "The E6 Oncoprotein Encoded by Human Papillomavirus Types 16 and 18 Promotes the Degradation of p53" *Cell* 63:1129-1136.
- Scheffner et al. (1992) "Targeted degradation of the retinoblastoma protein by human papillomavirus E7-E6 fusion proteins" *EMBO J* 11(7):2425-2431.
- Scheffner et al. (1993) "The HPV-16 E6 and E6-AP Complex Functions as a Ubiquitin-Protein Ligase in the Ubiquitination of p53" *Cell* 75:495-505.
- Seufert and Jentsch (1990) "Ubiquitin-conjugating enzymes UBC4 and UBC5 mediate selective degradation of short-lived and abnormal proteins" *EMBO J* 9(2):543-550.
- Treier et al. (1992) "*Drosophila UbcD1* encodes a highly conserved ubiquitin-conjugating enzyme involved in selective protein degradation" *EMBO J* 11(1):367-372.
- Tyers et al. (1992) "The Cin3-Cdc28 kinase complex of *S. cerevisiae* is regulated by proteolysis and phosphorylation" *EMBO J* 11(5):1773-1784.
- Watanabe et al. (1989) "Human Papillomavirus Type 16 Transformation of Primary Human embryonic Fibroblasts Requires Expression of Open Reading Frames E6 and E7" *J Virol* 63(2):965-969.
- Zhen et al. (1993) "The *ubc-2* Gene of *Caenorhabditis elegans* Encodes a Ubiquitin-Conjugating Enzyme Involved in Selective Protein Degradation" *Mol Cell Biol* 13(3):1371-1377.

ATOM	1	N	ARG	1	5	10.652	30.749	37.986	1.00	0.00
ATOM	2	CA	ARG	1	5	10.654	30.876	39.370	1.00	0.00
ATOM	3	C	ARG	1	5	9.386	31.560	39.753	1.00	0.00
ATOM	4	O	ARG	1	5	9.153	32.661	39.298	1.00	0.00
ATOM	5	CB	ARG	1	5	11.872	31.683	39.871	1.00	0.00
ATOM	6	CG	ARG	1	5	12.638	31.033	41.051	1.00	0.00
ATOM	7	CD	ARG	1	5	11.956	31.005	42.439	1.00	0.00
ATOM	8	NE	ARG	1	5	11.007	29.919	42.560	1.00	0.00
ATOM	9	CZ	ARG	1	5	10.281	29.677	43.686	1.00	0.00
ATOM	10	NH1	ARG	1	5	10.417	30.413	44.773	1.00	0.00
ATOM	11	NH2	ARG	1	5	9.411	28.694	43.623	1.00	0.00
ATOM	12	1H	ARG	1	5	10.497	29.737	37.803	1.00	0.00
ATOM	13	2H	ARG	1	5	9.927	31.348	37.541	1.00	0.00
ATOM	14	3H	ARG	1	T	11.573	31.167	37.744	1.00	0.00
ATOM	15	HE	ARG	1	5	10.888	29.459	41.680	1.00	0.00
ATOM	16	1HH1	ARG	1	5	9.832	30.497	45.580	1.00	0.00
ATOM	17	2HH1	ARG	1	5	11.297	30.876	44.878	1.00	0.00
ATOM	18	1HH2	ARG	1	5	8.878	28.410	44.420	1.00	0.00
ATOM	19	2HH2	ARG	1	5	9.283	28.223	42.749	1.00	0.00
ATOM	20	N	ILE	1	6	8.510	30.841	40.468	1.00	0.00
ATOM	21	CA	ILE	1	6	7.082	31.170	40.678	1.00	0.00
ATOM	22	C	ILE	1	6	6.442	31.065	39.284	1.00	0.00
ATOM	23	O	ILE	1	6	5.956	29.988	38.915	1.00	0.00
ATOM	24	CB	ILE	1	6	6.853	32.470	41.413	1.00	0.00
ATOM	25	CG1	ILE	1	6	7.651	32.561	42.727	1.00	0.00
ATOM	26	CG2	ILE	1	6	5.366	32.803	41.630	1.00	0.00
ATOM	27	CD1	ILE	1	6	6.834	32.009	43.915	1.00	0.00
ATOM	28	H	ILE	1	6	8.808	29.963	40.841	1.00	0.00
ATOM	29	N	HIS	1	7	6.540	32.257	38.639	1.00	0.00
ATOM	30	CA	HIS	1	7	6.197	32.320	37.173	1.00	0.00
ATOM	31	C	HIS	1	7	7.002	31.274	36.346	1.00	0.00
ATOM	32	O	HIS	1	7	8.174	30.919	36.611	1.00	0.00
ATOM	33	CB	HIS	1	7	6.233	33.740	36.613	1.00	0.00
ATOM	34	CG	HIS	1	7	7.507	34.441	37.147	1.00	0.00
ATOM	35	ND1	HIS	1	7	8.661	34.555	36.453	1.00	0.00
ATOM	36	CD2	HIS	1	7	7.660	35.160	38.373	1.00	0.00
ATOM	37	CE1	HIS	1	7	9.552	35.219	37.259	1.00	0.00
ATOM	38	NE2	HIS	1	7	8.924	35.642	38.428	1.00	0.00
ATOM	39	H	HIS	1	7	7.119	32.928	39.101	1.00	0.00
ATOM	40	1HD	HIS	1	7	8.806	34.258	35.530	1.00	0.00
ATOM	41	N	LYS	1	8	6.366	30.860	35.208	1.00	0.00
ATOM	42	CA	LYS	1	8	6.776	29.679	34.371	1.00	0.00
ATOM	43	C	LYS	1	8	6.529	28.399	35.097	1.00	0.00
ATOM	44	O	LYS	1	8	5.719	27.551	34.809	1.00	0.00
ATOM	45	CB	LYS	1	8	8.237	29.782	33.828	1.00	0.00

FIG. 1A

ATOM	46	CG	LYS	1	8	8.500	29.320	32.392	1.00	0.00
ATOM	47	CD	LYS	1	8	9.916	29.656	31.893	1.00	0.00
ATOM	48	CE	LYS	1	8	9.996	29.567	30.420	1.00	0.00
ATOM	49	NZ	LYS	1	8	11.060	30.278	29.662	1.00	0.00
ATOM	50	H	LYS	1	8	5.445	31.225	35.069	1.00	0.00
ATOM	51	1HZ	LYS	1	8	10.982	31.309	29.543	1.00	0.00
ATOM	52	1HZ	LYS	1	8	12.021	29.906	29.803	1.00	0.00
ATOM	53	3HZ	LYS	1	8	11.084	30.047	28.648	1.00	0.00
ATOM	54	N	GLU	1	9	7.274	28.215	36.215	1.00	0.00
ATOM	55	CA	GLU	1	9	6.995	27.070	37.159	1.00	0.00
ATOM	56	C	GLU	1	9	5.445	26.859	37.570	1.00	0.00
ATOM	57	O	GLU	1	9	4.977	25.769	37.789	1.00	0.00
ATOM	58	CB	GLU	1	9	7.951	27.203	38.425	1.00	0.00
ATOM	59	CG	GLU	1	9	9.433	26.780	38.356	1.00	0.00
ATOM	60	CD	GLU	1	9	10.169	27.525	39.767	1.00	0.00
ATOM	61	OE1	GLU	1	9	9.570	27.669	40.887	1.00	0.00
ATOM	62	OE2	GLU	1	9	11.354	27.904	39.613	1.00	0.00
ATOM	63	H	GLU	1	9	7.918	28.973	36.320	1.00	0.00
ATOM	64	N	LEU	1	10	4.816	27.970	37.728	1.00	0.00
ATOM	65	CA	LEU	1	10	3.434	28.097	38.068	1.00	0.00
ATOM	66	C	LEU	1	10	3.047	29.230	37.145	1.00	0.00
ATOM	67	O	LEU	1	10	3.769	30.204	37.045	1.00	0.00
ATOM	68	CB	LEU	1	10	3.176	28.409	39.531	1.00	0.00
ATOM	69	CG	LEU	1	10	2.041	27.485	40.074	1.00	0.00
ATOM	70	CD1	LEU	1	10	0.728	27.903	39.503	1.00	0.00
ATOM	71	CD2	LEU	1	10	2.366	26.000	39.832	1.00	0.00
ATOM	72	H	LEU	1	10	5.291	28.782	37.389	1.00	0.00
ATOM	73	N	LEU	1	11	1.850	29.001	36.566	1.00	0.00
ATOM	74	CA	ASN	1	11	1.355	29.849	35.463	1.00	0.00
ATOM	75	C	ASN	1	11	-0.107	30.195	35.375	1.00	0.00
ATOM	76	O	ASN	1	11	-0.745	29.756	34.411	1.00	0.00
ATOM	77	CB	ASN	1	11	2.047	29.486	34.124	1.00	0.00
ATOM	78	CG	ASN	1	11	1.754	30.546	33.096	1.00	0.00
ATOM	79	OD1	ASN	1	11	1.217	31.610	33.377	1.00	0.00
ATOM	80	ND2	ASN	1	11	1.971	30.162	31.811	1.00	0.00
ATOM	81	H	ASN	1	11	1.420	28.167	36.911	1.00	0.00
ATOM	82	1HD2	ASN	1	11	2.285	29.261	31.512	1.00	0.00
ATOM	83	2HD2	ASN	1	11	1.643	30.782	31.098	1.00	0.00
ATOM	84	N	ASP	1	12	-0.619	30.939	36.323	1.00	0.00
ATOM	85	CA	ASP	1	12	-2.064	31.262	36.250	1.00	0.00
ATOM	86	C	ASP	1	12	-2.668	31.635	34.910	1.00	0.00
ATOM	87	O	ASP	1	12	-3.709	31.008	34.539	1.00	0.00
ATOM	88	CB	ASP	1	12	-2.517	32.237	37.409	1.00	0.00
ATOM	89	CG	ASP	1	12	-1.293	32.675	38.157	1.00	0.00
ATOM	90	OD1	ASP	1	12	-0.497	33.446	37.645	1.00	0.00

FIG. 1B

ATOM	91	OD2	ASP	1	12	-1.057	32.123	39.266	1.00	0.00
ATOM	92	H	ASP	1	12	-0.119	31.179	37.155	1.00	0.00
ATOM	93	N	LEU	1	13	-2.099	32.601	34.132	1.00	0.00
ATOM	94	CA	LEU	1	13	-2.827	33.180	32.995	1.00	0.00
ATOM	95	C	LEU	1	13	-2.951	32.329	31.780	1.00	0.00
ATOM	96	O	LEU	1	13	-3.808	32.567	30.921	1.00	0.00
ATOM	97	CB	LEU	1	13	-2.337	34.600	32.584	1.00	0.00
ATOM	98	CG	LEU	1	13	-3.491	35.515	32.232	1.00	0.00
ATOM	99	CD1	LEU	1	13	-4.133	36.006	33.554	1.00	0.00
ATOM	100	CD2	LEU	1	13	-2.935	36.524	31.217	1.00	0.00
ATOM	101	H	LEU	1	13	-1.276	33.096	34.308	1.00	0.00
ATOM	102	N	ALA	1	14	-2.089	31.252	31.763	1.00	0.00
ATOM	103	CA	ALA	1	14	-2.227	30.131	30.792	1.00	0.00
ATOM	104	C	ALA	1	14	-3.449	29.293	31.048	1.00	0.00
ATOM	105	O	ALA	1	14	-3.724	28.399	30.301	1.00	0.00
ATOM	106	CB	ALA	1	14	-1.062	29.249	30.600	1.00	0.00
ATOM	107	H	ALA	1	14	-1.479	31.103	32.541	1.00	0.00
ATOM	108	N	ARG	1	15	-4.037	29.445	32.285	1.00	0.00
ATOM	109	CA	ARG	1	15	-5.241	28,633	32.573	1.00	0.00
ATOM	110	C	ARG	1	15	-6.478	29.572	32.619	1.00	0.00
ATOM	111	O	ARG	1	15	-7.563	29.184	32.147	1.00	0.00
ATOM	112	CB	ARG	1	15	-5.074	27.850	33.877	1.00	0.00
ATOM	113	CG	ARG	1	15	-5.256	26.348	33.808	1.00	0.00
ATOM	114	CD	ARG	1	15	-6.599	25.816	34.374	1.00	0.00
ATOM	115	NE	ARG	1	15	-7.738	26.515	33.767	1.00	0.00
ATOM	116	CZ	ARG	1	15	-8.322	27.475	34.535	1.00	0.00
ATOM	117	NH1	ARG	1	15	-9.426	28.064	34.120	1.00	0.00
ATOM	118	NH2	ARG	1	15	-7.720	27.973	35.652	1.00	0.00
ATOM	119	H	ARG	1	15	-3.720	30.097	32.975	1.00	0.00
ATOM	120	HE	ARG	1	15	-8.062	26.201	32.875	1.00	0.00
ATOM	121	1HH1	ARG	1	15	-9.878	28.795	34.631	1.00	0.00
ATOM	122	2HH1	ARG	1	15	-9.779	27,853	33.209	1.00	0.00
ATOM	123	1HH2	ARG	1	15	-8.137	28.803	36.022	1.00	0.00
ATOM	124	2HH2	ARG	1	15	-6.866	27.700	36.093	1.00	0.00
ATOM	125	N	ASP	1	16	-6.237	30.813	33.067	1.00	0.00
ATOM	126	CA	ASP	1	16	-7.250	31.856	32.803	1.00	0.00
ATOM	127	C	ASP	1	16	-6.824	33.136	32.067	1.00	0.00
ATOM	128	O	ASP	1	16	-6.532	34.221	32.610	1.00	0.00
ATOM	129	CB	ASP	1	16	-8.089	32.302	34.005	1.00	0.00
ATOM	130	CG	ASP	1	16	-8.462	31.097	34.974	1.00	0.00
ATOM	131	OD1	ASP	1	16	-9.602	30.692	34.912	1.00	0.00
ATOM	132	OD2	ASP	1	16	-7.672	30.572	35.769	1.00	0.00
ATOM	133	H	ASP	1	16	-5.451	31.212	33.539	1.00	0.00
ATOM	134	N	PRO	1	17	-6.956	33.072	30.746	1.00	0.00
ATOM	135	CA	PRO	1	17	-6.993	34.342	30.013	1.00	0.00

FIG. 1C

ATOM	136	C	PRO	1	17	-8.205	35.156	30.521	1.00	0.00
ATOM	137	O	PRO	1	17	-9.305	34.673	30.362	1.00	0.00
ATOM	138	CB	PRO	1	17	-7.265	33.841	28.530	1.00	0.00
ATOM	139	CG	PRO	1	17	-6.733	32.384	28.576	1.00	0.00
ATOM	140	CD	PRO	1	17	-7.148	31.899	29.962	1.00	0.00
ATOM	141	N	PRO	1	18	-7.965	36.355	31.086	1.00	0.00
ATOM	142	CA	PRO	1	18	-9.083	37.138	31.627	1.00	0.00
ATOM	143	C	PRO	1	18	-9.936	37.823	30.552	1.00	0.00
ATOM	144	O	PRO	1	18	-10.510	37.184	29.678	1.00	0.00
ATOM	145	CB	PRO	1	18	-8.415	37.993	32.696	1.00	0.00
ATOM	146	CG	PRO	1	18	-6.972	38.217	32.269	1.00	0.00
ATOM	147	CD	PRO	1	18	-6.703	36.974	31.360	1.00	0.00
ATOM	148	N	ALA	1	19	-10.151	39.105	30.790	1.00	0.00
ATOM	149	CA	ALA	1	19	-11.197	39.768	29.931	1.00	0.00
ATOM	150	C	ALA	1	19	-10.607	40.086	28.534	1.00	0.00
ATOM	151	O	ALA	1	19	-10.127	41.187	28.324	1.00	0.00
ATOM	152	CB	ALA	1	19	-11.617	41.179	30.598	1.00	0.00
ATOM	153	H	ALA	1	19	-9.561	39.653	31.383	1.00	0.00
ATOM	154	N	GLN	1	20	-10.496	39.036	27.758	1.00	0.00
ATOM	155	CA	GLN	1	20	-9.643	38.863	26.552	1.00	0.00
ATOM	156	C	GLN	1	20	-8.342	39.556	26.683	1.00	0.00
ATOM	157	O	GLN	1	20	-8.219	40.714	26.316	1.00	0.00
ATOM	158	CB	GLN	1	20	-10.452	39.059	25.250	1.00	0.00
ATOM	159	CG	GLN	1	20	-11.446	40.245	25.067	1.00	0.00
ATOM	160	CD	GLN	1	20	-12.575	40.215	26.077	1.00	0.00
ATOM	161	OE1	GLN	1	20	-12.883	41.218	26.660	1.00	0.00
ATOM	162	NE2	GLN	1	20	-13.154	39.042	26.289	1.00	0.00
ATOM	163	H	GLN	1	20	-11.149	38.324	28.017	1.00	0.00
ATOM	164	1HE2	GLN	1	20	-12.901	38.240	25.748	1.00	0.00
ATOM	165	2HE2	GLN	1	20	-13.792	38.993	27.058	1.00	0.00
ATOM	166	N	CYS	1	21	-7.381	38.866	27.312	1.00	0.00
ATOM	167	CA	CYS	1	21	-5.983	39.312	27.548	1.00	0.00
ATOM	168	C	CYS	1	21	-5.049	38.159	27.644	1.00	0.00
ATOM	169	O	CYS	1	21	-5.488	37.103	28.037	1.00	0.00
ATOM	170	CB	CYS	1	21	-5.939	40.115	28.837	1.00	0.00
ATOM	171	SG	CYS	1	21	-6.547	41.738	28.478	1.00	0.00
ATOM	172	H	CYS	1	21	-7.581	37.909	27.521	1.00	0.00
ATOM	173	N	SER	1	22	-3.768	38,479	27.242	1.00	0.00
ATOM	174	CA	SER	1	22	-2.909	37.373	26.846	1.00	0.00
ATOM	175	C	SER	1	22	-1.484	37.385	27.299	1.00	0.00
ATOM	176	O	SER	1	22	-0.890	38.502	27.246	1.00	0.00
ATOM	177	CB	SER	1	22	-3.010	37.109	25.307	1.00	0.00
ATOM	178	OG	SER	1	22	-4.179	37.764	24.726	1.00	0.00
ATOM	179	H	SER	1	22	-3.415	39.413	27.183	1.00	0.00
ATOM	180	HG	SER	1	22	-3.987	37.872	23.804	1.00	0.00

FIG. 1D

ATOM	181	N	ALA	1	23	-0.882	36.264	27.816	1.00	0.00
ATOM	182	CA	ALA	1	23	0.449	36.497	28.382	1.00	0.00
ATOM	183	C	ALA	1	23	1.533	35.875	27.516	1.00	0.00
ATOM	184	O	ALA	1	23	1.418	34.935	26.775	1.00	0.00
ATOM	185	CB	ALA	1	23	0.545	35.912	29.743	1.00	0.00
ATOM	186	H	ALA	1	23	-1.458	35.449	27.752	1.00	0.00
ATOM	187	N	GLY	1	24	2.746	36.445	27.787	1.00	0.00
ATOM	188	CA	GLY	1	24	3.850	35.741	27.233	1.00	0.00
ATOM	189	C	GLY	1	24	5.127	36.253	27.699	1.00	0.00
ATOM	190	O	GLY	1	24	5.075	37.310	28.265	1.00	0.00
ATOM	191	H	GLY	1	24	3.000	37.253	28.319	1.00	0.00
ATOM	192	N	PRO	1	25	6.264	35.552	27.463	1.00	0.00
ATOM	193	CA	PRO	1	25	7.528	36.069	27.834	1.00	0.00
ATOM	194	C	PRO	1	25	7.954	37.302	27.064	1.00	0.00
ATOM	195	O	PRO	1	25	7.102	37.967	26.519	1.00	0.00
ATOM	196	CB	PRO	1	25	8.536	34.960	27.498	1.00	0.00
ATOM	197	CG	PRO	1	25	7.766	33.902	26.624	1.00	0.00
ATOM	198	CD	PRO	1	25	6.327	34.297	26.774	1.00	0.00
ATOM	199	N	VAL	1	26	9.251	37.652	27.070	1.00	0.00
ATOM	200	CA	VAL	1	26	9.831	38.466	25.975	1.00	0.00
ATOM	201	C	VAL	1	26	10.054	37.488	24.900	1.00	0.00
ATOM	202	O	VAL	1	26	9.071	36.906	24.424	1.00	0.00
ATOM	203	CB	VAL	1	26	11.032	39.240	26.509	1.00	0.00
ATOM	204	CG1	VAL	1	26	10.726	40.772	26.799	1.00	0.00
ATOM	205	CG2	VAL	1	26	11.723	38.692	27.823	1.00	0.00
ATOM	206	H	VAL	1	26	9.681	37.030	27.724	1.00	0.00
ATOM	207	N	GLY	1	27	11.292	37.168	24.590	1.00	0.00
ATOM	208	CA	GLY	1	27	11.482	35.738	24.271	1.00	0.00
ATOM	209	C	GLY	1	27	11.647	34.818	25.477	1.00	0.00
ATOM	210	O	GLY	1	27	11.513	35.432	26.503	1.00	0.00
ATOM	211	H	GLY	1	27	12.099	37.604	24.988	1.00	0.00
ATOM	212	N	ASP	1	28	12.041	33.489	25.391	1.00	0.00
ATOM	213	CA	ASP	1	28	12.052	32.693	26.625	1.00	0.00
ATOM	214	C	ASP	1	28	12.264	33.254	28.042	1.00	0.00
ATOM	215	O	ASP	1	28	11.812	32.613	29.013	1.00	0.00
ATOM	216	CB	ASP	1	28	12.848	31.459	26.313	1.00	0.00
ATOM	217	CG	ASP	1	28	12.220	30.275	26.971	1.00	0.00
ATOM	218	OD1	ASP	1	28	11.048	29.929	26.718	1.00	0.00
ATOM	219	OD2	ASP	1	28	12.910	29.622	27.780	1.00	0.00
ATOM	220	H	ASP	1	28	12.089	33.023	24.507	1.00	0.00
ATOM	221	N	ASP	1	29	13.052	34.361	28.176	1.00	0.00
ATOM	222	CA	ASP	1	29	13.109	34.931	29.560	1.00	0.00
ATOM	223	C	ASP	1	29	11.716	35.121	30.289	1.00	0.00
ATOM	224	O	ASP	1	29	10.664	35.488	29.732	1.00	0.00
ATOM	225	CB	ASP	1	29	14.006	36.222	29.650	1.00	0.00

FIG. 1E

ATOM	226	CG	ASP	1	29	13.724	37.078	30.916	1.00	0.00
ATOM	227	OD1	ASP	1	29	13.034	38.080	30.842	1.00	0.00
ATOM	228	OD2	ASP	1	29	14.208	36.708	31.973	1.00	0.00
ATOM	229	H	ASP	1	29	13.367	34.766	27.317	1.00	0.00
ATOM	230	N	MET	1	30	11.742	34.874	31.593	1.00	0.00
ATOM	231	CA	MET	1	30	10.513	34.871	32.347	1.00	0.00
ATOM	232	C	MET	1	30	10.424	35.967	33.348	1.00	0.00
ATOM	233	O	MET	1	30	9.395	36.073	34.009	1.00	0.00
ATOM	234	CB	MET	1	30	10.024	33.562	32.904	1.00	0.00
ATOM	235	CG	MET	1	30	8.608	33.344	32.660	1.00	0.00
ATOM	236	SD	MET	1	30	8.121	33.393	30.892	1.00	0.00
ATOM	237	CE	MET	1	30	6.473	32.740	31.116	1.00	0.00
ATOM	238	H	MET	1	30	12.645	34.898	32.021	1.00	0.00
ATOM	239	N	PHE	1	31	11.476	36.831	33.387	1.00	0.00
ATOM	240	CA	PHE	1	31	11.426	37.966	34.248	1.00	0.00
ATOM	241	C	PHE	1	31	10.748	39.124	33.639	1.00	0.00
ATOM	242	O	PHE	1	31	10.171	39.983	34.265	1.00	0.00
ATOM	243	CB	PHE	1	31	12.847	38.280	34.753	1.00	0.00
ATOM	244	CG	PHE	1	31	13.502	36.936	35.115	1.00	0.00
ATOM	245	CD1	PHE	1	31	14.835	36.780	34.734	1.00	0.00
ATOM	246	CD2	PHE	1	31	12.856	35.900	35.819	1.00	0.00
ATOM	247	CE1	PHE	1	31	15.477	35.585	34.972	1.00	0.00
ATOM	248	CE2	PHE	1	31	13.540	34.632	36.102	1.00	0.00
ATOM	249	CZ	PHE	1	31	14.843	34.454	35.587	1.00	0.00
ATOM	250	H	PHE	1	31	12.277	36.803	32.788	1.00	0.00
ATOM	251	N	HIS	1	32	10.865	39.200	32.331	1.00	0.00
ATOM	252	CA	HIS	1	32	9.972	40.079	31.653	1.00	0.00
ATOM	253	C	HIS	1	32	8.924	39.415	30.769	1.00	0.00
ATOM	254	O	HIS	1	32	9.202	38.493	30.018	1.00	0.00
ATOM	255	CB	HIS	1	32	10.760	41.083	30.810	1.00	0.00
ATOM	256	CG	HIS	1	32	10.498	42.477	31.288	1.00	0.00
ATOM	257	ND1	HIS	1	32	11.346	43.479	31.266	1.00	0.00
ATOM	258	CD2	HIS	1	32	9.246	43.049	31.602	1.00	0.00
ATOM	259	CE1	HIS	1	32	10.678	44.659	31.574	1.00	0.00
ATOM	260	NE1	HIS	1	32	9.377	44.389	31.784	1.00	0.00
ATOM	261	H	HIS	1	32	11.489	38.555	31.889	1.00	0.00
ATOM	262	1HD	HIS	1	32	12.295	43.246	31.197	1.00	0.00
ATOM	263	N	TRP	1	33	7.790	39.969	30.994	1.00	0.00
ATOM	264	CA	TRP	1	33	6.604	39.493	30.279	1.00	0.00
ATOM	265	C	TRP	1	33	6.205	40.551	29.240	1.00	0.00
ATOM	266	O	TRP	1	33	6.438	41.689	29.500	1.00	0.00
ATOM	267	CB	TRP	1	33	5.415	39.260	31.209	1.00	0.00
ATOM	268	CG	TRP	1	33	5.641	38.164	32.176	1.00	0.00
ATOM	269	CD1	TRP	1	33	6.812	37.983	32.936	1.00	0.00
ATOM	270	CD2	TRP	1	33	4.801	37.018	32.425	1.00	0.00

FIG. 1F

ATOM	271	NE1	TRP	1	33	6.752	36.776	33.576	1.00	0.00
ATOM	272	CE2	TRP	1	33	5.536	36.139	33.304	1.00	0.00
ATOM	273	CE3	TRP	1	33	3.526	36.671	31.930	1.00	0.00
ATOM	274	CZ2	TRP	1	33	5.043	34.854	33.676	1.00	0.00
ATOM	275	CZ3	TRP	1	33	3.036	35.429	32.346	1.00	0.00
ATOM	276	CH2	TRP	1	33	3.758	34.524	33.185	1.00	0.00
ATOM	277	H	TRP	1	33	7.826	40.663	31.713	1.00	0.00
ATOM	278	1HE	TRP	1	33	7.482	36.492	34.165	1.00	0.00
ATOM	279	N	GLN	1	34	5.696	40.048	28.134	1.00	0.00
ATOM	280	CA	GLN	1	34	4.869	40.836	27.220	1.00	0.00
ATOM	281	C	GLN	1	34	3.424	40.408	27.177	1.00	0.00
ATOM	282	O	GLN	1	34	3.094	39.240	27.133	1.00	0.00
ATOM	283	CB	GLN	1	34	5.470	40.924	25.800	1.00	0.00
ATOM	284	CG	GLN	1	34	6.917	41.448	25.783	1.00	0.00
ATOM	285	CD	GLN	1	34	7.429	41.452	24.291	1.00	0.00
ATOM	286	OE1	GLN	1	34	7.612	42.380	23.559	1.00	0.00
ATOM	287	NE1	GLN	1	34	7.489	40.260	23.927	1.00	0.00
ATOM	288	H	GLN	1	34	5.517	39.064	28.146	1.00	0.00
ATOM	289	1HE2	GLN	1	34	7.323	39.442	24.478	1.00	0.00
ATOM	290	2HE2	GLN	1	34	7.792	40.263	22.974	1.00	0.00
ATOM	291	N	ALA	1	35	2.538	41.366	27.220	1.00	0.00
ATOM	292	CA	ALA	1	35	1.184	40.851	27.117	1.00	0.00
ATOM	293	C	ALA	1	35	0.289	41.720	26.273	1.00	0.00
ATOM	294	O	ALA	1	35	0.110	42.908	26.564	1.00	0.00
ATOM	295	CB	ALA	1	35	0.621	40.755	28.550	1.00	0.00
ATOM	296	H	ALA	1	35	2.893	42.286	27.389	1.00	0.00
ATOM	297	N	THR	1	36	-0.357	40.970	25.395	1.00	0.00
ATOM	298	CA	THR	1	36	-1.550	41.487	24.753	1.00	0.00
ATOM	299	C	THR	1	36	-2.696	41.829	25.712	1.00	0.00
ATOM	300	O	THR	1	36	-3.280	41.022	26.401	1.00	0.00
ATOM	301	CB	THR	1	36	-2.200	40.523	23.714	1.00	0.00
ATOM	302	OG1	THR	1	36	-1.176	39.765	22.985	1.00	0.00
ATOM	303	CG2	THR	1	36	-3.302	41.301	22.943	1.00	0.00
ATOM	304	H	THR	1	36	0.086	40.113	25.129	1.00	0.00
ATOM	305	1HG	THR	1	36	-1.335	40.089	22.109	1.00	0.00
ATOM	306	N	ILE	1	37	-3.026	43.130	25.648	1.00	0.00
ATOM	307	CA	ILE	1	37	-4.191	43.548	26.429	1.00	0.00
ATOM	308	C	ILE	1	37	-5.298	44.216	25.630	1.00	0.00
ATOM	309	O	ILE	1	37	-5.101	45.204	24.964	1.00	0.00
ATOM	310	CB	ILE	1	37	-3.797	44.351	27.724	1.00	0.00
ATOM	311	CG1	ILE	1	37	-4.897	45.181	28.449	1.00	0.00
ATOM	312	CG2	ILE	1	37	-2.542	45.130	27.448	1.00	0.00
ATOM	313	CD1	ILE	1	37	-4.556	46.013	29.659	1.00	0.00
ATOM	314	H	ILE	1	37	-2.472	43.740	25.081	1.00	0.00
ATOM	315	N	ILE	1	37	-6.458	43.574	25.878	1.00	0.00

FIG. 1G

ATOM	316	CA	MET	1	38	-7.586	44.220	25.347	1.00	0.00
ATOM	317	C	MET	1	38	8.605	44.872	26.322	1.00	0.00
ATOM	318	O	MET	1	38	-8.349	45.935	26.893	1.00	0.00
ATOM	319	CB	MET	1	38	-8.343	43.241	24.447	1.00	0.00
ATOM	320	CG	MET	1	38	-7.470	42.311	23.603	1.00	0.00
ATOM	321	SD	MET	1	38	-8.288	40.905	22.853	1.00	0.00
ATOM	322	CE	MET	1	38	-9.622	41.838	22.065	1.00	0.00
ATOM	323	H	MET	1	38	6.505	42.757	26.452	1.00	0.00
ATOM	324	N	GLY	1	39	-9.770	44.272	26.496	1.00	0.00
ATOM	325	CA	GLY	1	39	-11.056	44.966	26.850	1.00	0.00
ATOM	326	C	GLY	1	39	-12.298	44.411	26.038	1.00	0.00
ATOM	327	O	GLY	1	39	-12.214	43.940	24.916	1.00	0.00
ATOM	328	H	GLY	1	39	-9.966	43.294	26.425	1.00	0.00
ATOM	329	N	PRO	1	40	-13.477	44.500	26.766	1.00	0.00
ATOM	330	CA	PRO	1	40	-14.704	43.852	26.332	1.00	0.00
ATOM	331	C	PRO	1	40	-15.431	44.406	25.096	1.00	0.00
ATOM	332	O	PRO	1	40	-15.378	45.605	24.835	1.00	0.00
ATOM	333	CG	PRO	1	40	-15.528	43.836	27.644	1.00	0.00
ATOM	334	CG	PRO	1	40	-15.102	45.100	28.414	1.00	0.00
ATOM	335	CD	PRO	1	40	-13.575	45.060	28.122	1.00	0.00
ATOM	336	N	ASN	1	41	-16.104	43.478	24.432	1.00	0.00
ATOM	337	CA	ASN	1	41	-16.866	43.806	23.227	1.00	0.00
ATOM	338	C	ASN	1	41	-17.788	45.030	23.243	1.00	0.00
ATOM	339	O	ASN	1	41	-18.334	45.404	24.266	1.00	0.00
ATOM	340	CB	ASN	1	41	-17.505	42.566	22.629	1.00	0.00
ATOM	341	CG	ASN	1	41	-17.457	42.783	21.100	1.00	0.00
ATOM	342	OD1	ASN	1	41	-16.424	42.515	20.518	1.00	0.00
ATOM	343	ND2	ASN	1	41	-18.580	43.205	20.495	1.00	0.00
ATOM	344	H	ASN	1	41	-16.031	42.509	24.671	1.00	0.00
ATOM	345	1HD2	ASN	1	41	-19.340	43.519	21.065	1.00	0.00
ATOM	346	2HD2	ASN	1	41	-18.839	43.307	19.534	1.00	0.00
ATOM	347	N	ASP	1	42	-17.706	45.666	22.075	1.00	0.00
ATOM	348	CA	ASP	1	42	-18.549	46.827	21.646	1.00	0.00
ATOM	349	C	ASP	1	42	-17.990	48.094	22.183	1.00	0.00
ATOM	350	O	ASP	1	42	-17.904	49.099	21.503	1.00	0.00
ATOM	351	CB	ASP	1	42	-20.049	46.643	22.001	1.00	0.00
ATOM	352	CG	ASP	1	42	-20.617	45.386	21.248	1.00	0.00
ATOM	353	OD1	ASP	1	42	-21.189	44.546	22.019	1.00	0.00
ATOM	354	OD2	ASP	1	42	-20.505	45.329	19.972	1.00	0.00
ATOM	355	H	ASP	1	42	-17.172	45.239	21.345	1.00	0.00
ATOM	356	N	SER	1	43	-17.518	47.956	23.425	1.00	0.00
ATOM	357	CA	SER	1	43	-16.981	49.076	24.283	1.00	0.00
ATOM	358	C	SER	1	43	-15.645	49.654	23.703	1.00	0.00
ATOM	359	O	SER	1	43	-14.885	49.017	22.898	1.00	0.00
ATOM	360	CB	SER	1	43	-16.864	48.550	25.721	1.00	0.00

FIG. 1H

ATOM	361	OG	SER	1	43	-17.907	47.578	25.969	1.00	0.00
ATOM	362	H	SER	1	43	-17.691	47.082	23.879	1.00	0.00
ATOM	363	HG	SER	1	43	-17.668	46.770	25.535	1.00	0.00
ATOM	364	N	PRO	1	43	-15.449	50.978	24.037	1.00	0.00
ATOM	365	CA	PRO	1	44	-14.337	51.686	23.444	1.00	0.00
ATOM	366	C	PRO	1	44	-12.965	51.042	23.266	1.00	0.00
ATOM	367	O	PRO	1	44	-12.323	50.986	22.187	1.00	0.00
ATOM	368	CB	PRO	1	44	-14.294	52.931	24.332	1.00	0.00
ATOM	369	CG	PRO	1	44	-15.749	53.253	24.642	1.00	0.00
ATOM	370	CD	PRO	1	44	-16.388	51.873	24.723	1.00	0.00
ATOM	371	N	TYR	1	45	-12.499	50.612	24.469	1.00	0.00
ATOM	372	CA	TYR	1	45	-11.173	49.936	24.432	1.00	0.00
ATOM	373	C	TYR	1	45	-10.989	48.618	23.768	1.00	0.00
ATOM	374	O	TYR	1	45	-9.874	48.082	23.625	1.00	0.00
ATOM	375	CB	TYR	1	45	-10.642	49.821	25.898	1.00	0.00
ATOM	376	CG	TYR	1	45	-10.263	51.093	26.551	1.00	0.00
ATOM	377	CD1	TYR	1	45	-11.055	51.437	27.637	1.00	0.00
ATOM	378	CD2	TYR	1	45	-9.186	51.866	26.062	1.00	0.00
ATOM	379	CE1	TYR	1	45	-10.678	52.632	28.338	1.00	0.00
ATOM	380	CE2	TYR	1	45	-8.954	53.120	26.707	1.00	0.00
ATOM	381	CZ	TYR	1	45	-9.696	53.503	27.827	1.00	0.00
ATOM	382	OH	TYR	1	45	-9.427	54.731	28.452	1.00	0.00
ATOM	383	H	TYR	1	45	-12.786	50.950	25.365	1.00	0.00
ATOM	384	HH	TYR	1	45	-9.015	55.269	27.788	1.00	0.00
ATOM	385	N	GLN	1	46	-12.094	48.046	23.382	1.00	0.00
ATOM	386	CA	GLN	1	46	-12.061	46.725	22.755	1.00	0.00
ATOM	387	C	GLN	1	46	-11.454	46.729	21.459	1.00	0.00
ATOM	388	O	GLN	1	46	-11.537	47.737	20.760	1.00	0.00
ATOM	389	CB	GLN	1	46	-13.469	46.126	22.604	1.00	0.00
ATOM	390	CG	GLN	1	46	-13.789	45.049	21.539	1.00	0.00
ATOM	391	CD	GLN	1	46	-14.461	45.627	20.354	1.00	0.00
ATOM	392	OE1	GLN	1	46	-14.289	46.795	20.051	1.00	0.00
ATOM	393	NE2	GLN	1	46	-15.311	44.856	19.671	1.00	0.00
ATOM	394	H	GLN	1	46	-12.898	48.611	23.201	1.00	0.00
ATOM	395	1HE2	GLN	1	46	-15.436	43.884	19.867	1.00	0.00
ATOM	396	2HE2	GLN	1	46	-15.796	45.251	18.890	1.00	0.00
ATOM	397	N	GLY	1	47	-10.734	45.644	21.155	1.00	0.00
ATOM	398	CA	GLY	1	47	-10.114	45.605	19.848	1.00	0.00
ATOM	399	C	GLY	1	47	-8.629	45.843	19.905	1.00	0.00
ATOM	400	O	GLY	1	47	-7.769	45.119	19.319	1.00	0.00
ATOM	401	H	GLY	1	47	-10.775	44.822	21.723	1.00	0.00
ATOM	402	N	GLY	1	48	-8.271	46.947	20.695	1.00	0.00
ATOM	403	CA	GLY	1	48	-6.913	47.393	20.950	1.00	0.00
ATOM	404	C	GLY	1	48	-6.009	46.228	21.154	1.00	0.00
ATOM	405	O	GLY	1	48	-6.280	45.269	21.911	1.00	0.00

FIG. 11

ATOM	406	H	GLY	1	48	-9.072	47.442	21.032	1.00	0.00
ATOM	407	N	VAL	1	49	-4.877	46.300	20.420	1.00	0.00
ATOM	408	CA	VAL	1	49	-3.846	45.346	20.774	1.00	0.00
ATOM	409	C	VAL	1	49	-2.700	46.129	21.429	1.00	0.00
ATOM	410	O	VAL	1	49	-1.856	46.847	20.841	1.00	0.00
ATOM	411	CB	VAL	1	49	-3.358	44.738	19.453	1.00	0.00
ATOM	412	CG1	VAL	1	49	-2.394	43.577	19.728	1.00	0.00
ATOM	413	CG2	VAL	1	49	-4.498	44.517	18.390	1.00	0.00
ATOM	414	H	VAL	1	49	-4.707	47.023	19.750	1.00	0.00
ATOM	415	N	PHE	1	50	-2.811	46.022	22.734	1.00	0.00
ATOM	416	CA	PHE	1	50	-1.793	46.690	23.383	1.00	0.00
ATOM	417	C	PHE	1	50	-0.835	45.766	23.937	1.00	0.00
ATOM	418	O	PHE	1	50	-1.082	44.564	24.001	1.00	0.00
ATOM	419	CB	PHE	1	50	-2.343	47.588	24.462	1.00	0.00
ATOM	420	CG	PHE	1	50	-3.767	48.162	24.266	1.00	0.00
ATOM	421	CD1	PHE	1	50	-4.158	48.819	23.058	1.00	0.00
ATOM	422	CD2	PHE	1	50	-4.648	48.064	25.365	1.00	0.00
ATOM	423	CE1	PHE	1	50	-5.442	49.326	22.959	1.00	0.00
ATOM	424	CE2	PHE	1	50	-5.989	48.477	25.179	1.00	0.00
ATOM	425	CZ	PHE	1	50	-6.378	49.138	23.978	1.00	0.00
ATOM	426	H	PHE	1	50	-3.535	45.534	23.221	1.00	0.00
ATOM	427	N	PHE	1	51	0.269	46.369	24.384	1.00	0.00
ATOM	428	CA	PHE	1	51	1.219	45.510	25.011	1.00	0.00
ATOM	429	C	PHE	1	51	1.628	46.101	26.372	1.00	0.00
ATOM	430	O	PHE	1	51	1.290	47.244	26.681	1.00	0.00
ATOM	431	CB	PHE	1	51	2.558	45.235	24.183	1.00	0.00
ATOM	432	CG	PHE	1	51	2.071	44.870	22.735	1.00	0.00
ATOM	433	CD1	PHE	1	51	1.474	43.648	22.482	1.00	0.00
ATOM	434	CD2	PHE	1	51	2.253	45.869	21.762	1.00	0.00
ATOM	435	CE1	PHE	1	51	1.003	43.443	21.104	1.00	0.00
ATOM	436	CE2	PHE	1	51	1.858	45.668	20.423	1.00	0.00
ATOM	437	CZ	PHE	1	51	1.236	44.497	20.130	1.00	0.00
ATOM	438	H	PHE	1	51	0.598	47.248	24.040	1.00	0.00
ATOM	439	N	LEU	1	52	2.130	45.091	27.132	1.00	0.00
ATOM	440	CA	LEU	1	52	2.325	45.227	28.576	1.00	0.00
ATOM	441	C	LEU	1	52	3.524	44.445	29.054	1.00	0.00
ATOM	442	O	LEU	1	52	3.530	43.203	29.019	1.00	0.00
ATOM	443	CB	LEU	1	52	1.065	44.789	29.287	1.00	0.00
ATOM	444	CG	LEU	1	52	0.938	45.233	30.789	1.00	0.00
ATOM	445	CD1	LEU	1	52	-0.220	46.185	30.933	1.00	0.00
ATOM	446	CD2	LEU	1	52	0.653	44.204	31.814	1.00	0.00
ATOM	447	H	LEU	1	52	2.068	44.177	26.732	1.00	0.00
ATOM	448	N	THR	1	53	4.567	45.174	29.416	1.00	0.00
ATOM	449	CA	THR	1	53	5.541	44.339	30.004	1.00	0.00
ATOM	450	C	THR	1	53	5.394	44.249	31.552	1.00	0.00

FIG. 1J

ATOM	451	O	THR	1	53	5.698	45.208	32.281	1.00	0.00
ATOM	452	CB	THR	1	53	6.993	44.839	29.721	1.00	0.00
ATOM	453	OG1	THR	1	53	7.141	46.178	30.117	1.00	0.00
ATOM	454	OG2	THR	1	53	7.449	44.539	28.336	1.00	0.00
ATOM	455	H	THR	1	53	4.610	46.165	29.290	1.00	0.00
ATOM	456	1HG	THR	1	53	6.693	46.593	29.392	1.00	0.00
ATOM	457	N	ILE	1	54	5.055	43.019	32.006	1.00	0.00
ATOM	458	CA	ILE	1	54	5.201	42.975	33.500	1.00	0.00
ATOM	459	C	ILE	1	54	6.483	42.187	33.824	1.00	0.00
ATOM	460	O	ILE	1	54	6.624	41.049	33.449	1.00	0.00
ATOM	461	CB	ILE	1	54	3.895	42.546	34.056	1.00	0.00
ATOM	462	CG1	ILE	1	54	4.116	42.239	35.538	1.00	0.00
ATOM	463	CG2	ILE	1	54	3.293	41.330	33.449	1.00	0.00
ATOM	464	CD1	ILE	1	54	2.835	42.116	36.297	1.00	0.00
ATOM	465	H	ILE	1	54	4.883	42.200	31.459	1.00	0.00
ATOM	466	N	HIS	1	55	7.356	42.905	34.533	1.00	0.00
ATOM	467	CA	HIS	1	55	8.680	42.432	35.034	1.00	0.00
ATOM	468	C	HIS	1	55	8.488	41.851	36.430	1.00	0.00
ATOM	469	O	HIS	1	55	8.711	42.401	37.499	1.00	0.00
ATOM	470	CB	HIS	1	55	9.677	43.626	35.221	1.00	0.00
ATOM	471	CG	HIS	1	55	11.084	43.145	35.610	1.00	0.00
ATOM	472	ND1	HIS	1	55	11.463	42.849	36.889	1.00	0.00
ATOM	473	CD2	HIS	1	55	12.160	43.008	34.734	1.00	0.00
ATOM	474	CE1	HIS	1	55	12.798	42.553	36.874	1.00	0.00
ATOM	475	NE2	HIS	1	55	13.166	42.652	35.589	1.00	0.00
ATOM	476	H	HIS	1	55	6.985	43.825	34.656	1.00	0.00
ATOM	477	1HD	HIS	1	55	10.779	42.792	37.588	1.00	0.00
ATOM	478	N	PHE	1	56	8.004	40.579	36.294	1.00	0.00
ATOM	479	CA	PHE	1	56	8.124	39.740	37.478	1.00	0.00
ATOM	480	C	PHE	1	56	9.564	39.404	37.759	1.00	0.00
ATOM	481	O	PHE	1	56	10.111	38.735	36.928	1.00	0.00
ATOM	482	CB	PHE	1	56	7.431	38.414	37.221	1.00	0.00
ATOM	483	CG	PHE	1	56	5.957	38.636	37.218	1.00	0.00
ATOM	484	CD1	PHE	1	56	5.243	38.642	38.488	1.00	0.00
ATOM	485	CD2	PHE	1	56	5.258	38.651	36.021	1.00	0.00
ATOM	486	CE1	PHE	1	56	3.843	38.681	38.472	1.00	0.00
ATOM	487	CE2	PHE	1	56	3.883	38.677	36.013	1.00	0.00
ATOM	488	CZ	PHE	1	56	3.180	38.671	37.227	1.00	0.00
ATOM	489	H	PHE	1	56	8.146	40.122	35.416	1.00	0.00
ATOM	490	N	PRO	1	57	10.161	40.000	38.828	1.00	0.00
ATOM	491	CA	PRO	1	57	11.669	39.843	38.911	1.00	0.00
ATOM	492	C	PRO	1	57	12.092	38.428	39.104	1.00	0.00
ATOM	493	O	PRO	1	57	11.362	37.470	39.343	1.00	0.00
ATOM	494	CB	PRO	1	57	12.003	40.789	40.050	1.00	0.00
ATOM	495	CG	PRO	1	57	10.739	40.883	40.945	1.00	0.00

FIG. 1K

ATOM	496	CD	PRO	1	57	9.576	40.703	39.952	1.00	0.00
ATOM	497	N	THR	1	58	13.424	38.247	38.912	1.00	0.00
ATOM	498	CA	THR	1	58	14.071	37.056	39.470	1.00	0.00
ATOM	499	C	THR	1	58	13.788	36.737	40.841	1.00	0.00
ATOM	500	O	THR	1	58	13.926	35.629	41.294	1.00	0.00
ATOM	501	CB	THR	1	58	15.541	37.145	39.032	1.00	0.00
ATOM	502	OG1	THR	1	58	15.680	37.609	37.672	1.00	0.00
ATOM	503	CG2	THR	1	58	16.131	35.762	39.182	1.00	0.00
ATOM	504	H	THR	1	58	14.036	38.966	38.583	1.00	0.00
ATOM	505	1HG	THR	1	58	16.524	37.297	37.373	1.00	0.00
ATOM	506	N	ASP	1	59	13.457	37.777	41.583	1.00	0.00
ATOM	507	CA	ASP	1	59	13.181	37.664	43.011	1.00	0.00
ATOM	508	C	ASP	1	59	11.769	37.504	43.490	1.00	0.00
ATOM	509	O	ASP	1	59	11.159	38.230	44.245	1.00	0.00
ATOM	510	CB	ASP	1	59	13.889	38.750	43.734	1.00	0.00
ATOM	511	CG	ASP	1	59	13.491	40.164	43.290	1.00	0.00
ATOM	512	OD1	ASP	1	59	14.211	40.777	42.414	1.00	0.00
ATOM	513	OD2	ASP	1	59	12.597	40.779	43.786	1.00	0.00
ATOM	514	H	ASP	1	59	13.329	38.682	41.178	1.00	0.00
ATOM	515	N	TYR	1	60	11.264	36.412	43.011	1.00	0.00
ATOM	516	CA	TYR	1	60	9.821	36.218	43.226	1.00	0.00
ATOM	517	C	TYR	1	60	9.406	35.407	44.477	1.00	0.00
ATOM	518	O	TYR	1	60	10.033	34.441	44.781	1.00	0.00
ATOM	519	CB	TYR	1	60	9.157	35.843	41.829	1.00	0.00
ATOM	520	CG	TYR	1	60	7.838	36.612	41.789	1.00	0.00
ATOM	521	CD1	TYR	1	60	8.034	37.990	41.682	1.00	0.00
ATOM	522	CD2	TYR	1	60	6.546	36.030	41.874	1.00	0.00
ATOM	523	CE1	TYR	1	60	6.972	38.870	41.721	1.00	0.00
ATOM	524	CE2	TYR	1	60	5.438	36.950	41.740	1.00	0.00
ATOM	525	CZ	TYR	1	60	5.729	38.357	41.751	1.00	0.00
ATOM	526	OH	TYR	1	60	4.710	39.254	41.968	1.00	0.00
ATOM	527	H	TYR	1	60	11.803	35.930	42.321	1.00	0.00
ATOM	528	HH	TYR	1	60	4.929	39.846	42.676	1.00	0.00
ATOM	529	N	PRO	1	61	8.341	35.745	45.277	1.00	0.00
ATOM	530	CA	PRO	1	61	7.611	36.986	45.163	1.00	0.00
ATOM	531	C	PRO	1	61	8.031	38.127	46.103	1.00	0.00
ATOM	532	O	PRO	1	61	7.544	38.281	47.197	1.00	0.00
ATOM	533	CB	PRO	1	61	6.241	36.440	45.549	1.00	0.00
ATOM	534	CG	PRO	1	61	6.451	35.361	46.551	1.00	0.00
ATOM	535	CD	PRO	1	61	7.874	34.846	46.340	1.00	0.00
ATOM	536	N	PHE	1	62	8.921	39.015	45.665	1.00	0.00
ATOM	537	CA	PHE	1	62	9.316	39.953	46.675	1.00	0.00
ATOM	538	C	PHE	1	62	9.091	41.436	46.494	1.00	0.00
ATOM	539	O	PHE	1	62	8.277	42.242	47.033	1.00	0.00
ATOM	540	CB	PHE	1	62	10.835	39.726	47.093	1.00	0.00

FIG. 1L

ATOM	541	CG	PHE	1	62	11.170	38.343	47.627	1.00	0.00
ATOM	542	CD1	PHE	1	62	10.256	37.516	48.364	1.00	0.00
ATOM	543	CD2	PHE	1	62	12.483	37.929	47.366	1.00	0.00
ATOM	544	CD1	PHE	1	62	10.701	36.264	48.835	1.00	0.00
ATOM	545	CE2	PHE	1	62	12.959	36.670	47.863	1.00	0.00
ATOM	546	CZ	PHE	1	62	12.047	35.848	48.625	1.00	0.00
ATOM	547	H	PHE	1	62	9.595	38.814	44.954	1.00	0.00
ATOM	548	N	LYS	1	63	9.863	41.843	45.495	1.00	0.00
ATOM	549	CA	LYS	1	63	9.548	43.072	44.849	1.00	0.00
ATOM	550	C	LYS	1	63	8.397	43.143	43.796	1.00	0.00
ATOM	551	O	LYS	1	63	8.195	42.199	42.989	1.00	0.00
ATOM	552	CB	LYS	1	63	10.812	43.763	44.268	1.00	0.00
ATOM	553	CG	LYS	1	63	11.675	44.228	45.486	1.00	0.00
ATOM	554	CD	LYS	1	63	13.008	44.744	45.085	1.00	0.00
ATOM	555	CE	LYS	1	63	14.141	43.689	44.873	1.00	0.00
ATOM	556	NZ	LYS	1	63	13.885	43.084	43.628	1.00	0.00
ATOM	557	H	LYS	1	63	10.564	41.264	45.080	1.00	0.00
ATOM	558	1HZ	LYS	1	63	13.607	43.721	42.854	1.00	0.00
ATOM	559	2HZ	LYS	1	63	14.604	42.424	43.268	1.00	0.00
ATOM	560	3HZ	LYS	1	63	13.094	42.434	43.815	1.00	0.00
ATOM	561	N	PRO	1	64	7.612	44.296	44.014	1.00	0.00
ATOM	562	CA	PRO	1	64	6.552	44.555	43.047	1.00	0.00
ATOM	563	C	PRO	1	64	7.019	44.767	41.595	1.00	0.00
ATOM	564	O	PRO	1	64	7.827	45.556	41.196	1.00	0.00
ATOM	565	CB	PRO	1	64	5.886	45.784	43.752	1.00	0.00
ATOM	566	CG	PRO	1	64	6.176	45.623	45.213	1.00	0.00
ATOM	567	CD	PRO	1	64	7.675	45.291	45.131	1.00	0.00
ATOM	568	N	PRO	1	65	6.468	43.847	40.815	1.00	0.00
ATOM	569	CA	PRO	1	65	6.866	43.890	39.421	1.00	0.00
ATOM	570	C	PRO	1	65	6.657	45.256	38.732	1.00	0.00
ATOM	571	O	PRO	1	65	5.727	46.018	38.928	1.00	0.00
ATOM	572	CB	PRO	1	65	6.066	42.674	38.836	1.00	0.00
ATOM	573	CG	PRO	1	65	5.450	41.854	39.988	1.00	0.00
ATOM	574	CD	PRO	1	65	5.640	42.744	41.175	1.00	0.00
ATOM	575	N	LYS	1	66	7.616	45.558	37.920	1.00	0.00
ATOM	576	CA	LYS	1	66	7.582	46.638	36.943	1.00	0.00
ATOM	577	C	LYS	1	66	6.677	46.420	35.758	1.00	0.00
ATOM	578	O	LYS	1	66	6.975	45.573	34.895	1.00	0.00
ATOM	579	CB	LYS	1	66	9.042	47.057	36.632	1.00	0.00
ATOM	580	CG	LYS	1	66	9.026	47.916	35.359	1.00	0.00
ATOM	581	CD	LYS	1	66	9.426	47.170	34.096	1.00	0.00
ATOM	582	CE	LYS	1	66	8.561	47.715	32.960	1.00	0.00
ATOM	583	NZ	LYS	1	66	7.593	46.650	32.604	1.00	0.00
ATOM	584	H	LYS	1	66	8.315	44.850	37.823	1.00	0.00
ATOM	585	1HZ	LYS	1	66	7.202	46.096	33.393	1.00	0.00

FIG. 1M

ATOM	586	2HZ	LYS	1	66	6.840	46.784	31.900	1.00	0.00
ATOM	587	3HZ	LYS	1	66	8.051	45.940	31.997	1.00	0.00
ATOM	588	N	VAL	1	67	5.641	47.237	35.914	1.00	0.00
ATOM	589	CA	VAL	1	67	4.618	47.092	34.890	1.00	0.00
ATOM	590	C	VAL	1	67	4.669	48.202	33.838	1.00	0.00
ATOM	591	O	VAL	1	67	5.038	49.290	34.212	1.00	0.00
ATOM	592	CB	VAL	1	67	3.210	47.149	35.616	1.00	0.00
ATOM	593	CG1	VAL	1	67	2.008	47.020	34.713	1.00	0.00
ATOM	594	CG2	VAL	1	67	3.138	46.027	36.596	1.00	0.00
ATOM	595	H	VAL	1	67	5.652	47.816	36.730	1.00	0.00
ATOM	596	N	ALA	1	68	4.284	47.890	32.576	1.00	0.00
ATOM	597	CA	ALA	1	68	4.063	49.030	31.594	1.00	0.00
ATOM	598	C	ALA	1	68	3.210	48.782	30.295	1.00	0.00
ATOM	599	O	ALA	1	68	3.050	47.648	29.844	1.00	0.00
ATOM	600	CB	ALA	1	68	5.339	49.726	31.204	1.00	0.00
ATOM	601	H	ALA	1	68	3.936	46.964	32.435	1.00	0.00
ATOM	602	N	PHE	1	69	2.744	49.948	29.729	1.00	0.00
ATOM	603	CA	PHE	1	69	2.170	49.817	28.428	1.00	0.00
ATOM	604	C	PHE	1	69	3.003	49.993	27.243	1.00	0.00
ATOM	605	O	PHE	1	69	4.174	50.197	27.488	1.00	0.00
ATOM	606	CB	PHE	1	69	1.005	50.861	28.439	1.00	0.00
ATOM	607	CG	PHE	1	69	-0.236	50.170	29.024	1.00	0.00
ATOM	608	CD1	PHE	1	69	-0.571	50.303	30.370	1.00	0.00
ATOM	609	CD2	PHE	1	69	-1.184	49.627	28.119	1.00	0.00
ATOM	610	CE1	PHE	1	69	-1.950	49.879	30.749	1.00	0.00
ATOM	611	CE2	PHE	1	69	-2.510	49.317	28.496	1.00	0.00
ATOM	612	CZ	PHE	1	69	-2.919	49.553	29.836	1.00	0.00
ATOM	613	H	PHE	1	69	3.122	50.795	30.100	1.00	0.00
ATOM	614	N	THR	1	70	2.418	49.796	26.051	1.00	0.00
ATOM	615	CA	THR	1	70	3.174	50.059	24.865	1.00	0.00
ATOM	616	C	THR	1	70	2.540	50.869	23.691	1.00	0.00
ATOM	617	O	THR	1	70	3.101	51.454	22.774	1.00	0.00
ATOM	618	CB	THR	1	70	3.823	48.693	24.425	1.00	0.00
ATOM	619	OG1	THR	1	70	4.392	47.934	25.445	1.00	0.00
ATOM	620	OG2	THR	1	70	5.058	48.811	23.431	1.00	0.00
ATOM	621	H	THR	1	70	1.520	49.357	26.083	1.00	0.00
ATOM	622	1HG	THR	1	70	4.658	48.583	26.082	1.00	0.00
ATOM	623	N	THR	1	71	1.197	50.942	23.754	1.00	0.00
ATOM	624	CA	THR	1	71	0.474	51.874	22.977	1.00	0.00
ATOM	625	C	THR	1	71	0.307	53.101	23.855	1.00	0.00
ATOM	626	O	THR	1	71	0.482	52.993	25.082	1.00	0.00
ATOM	627	CB	THR	1	71	-0.890	51.230	22.586	1.00	0.00
ATOM	628	OG1	THR	1	71	-1.547	51.870	21.495	1.00	0.00
ATOM	629	OG2	THR	1	71	-1.785	51.219	23.771	1.00	0.00
ATOM	630	H	THR	1	71	0.626	50.488	24.439	1.00	0.00

FIG. 1N

ATOM	631	1HG	THR	1	71	-2.411	51.480	21.469	1.00	0.00
ATOM	632	N	ARG	1	72	-0.235	54.173	23.226	1.00	0.00
ATOM	633	CA	ARG	1	72	-0.698	55.208	24.180	1.00	0.00
ATOM	634	C	ARG	1	72	-2.115	55.840	23.974	1.00	0.00
ATOM	635	O	ARG	1	72	-2.217	57.065	23.776	1.00	0.00
ATOM	636	CB	ARG	1	72	0.288	56.308	24.307	1.00	0.00
ATOM	637	CG	ARG	1	72	0.826	56.516	25.720	1.00	0.00
ATOM	638	CD	ARG	1	72	-0.223	57.282	26.573	1.00	0.00
ATOM	639	NE	ARG	1	72	-0.810	58.398	25.780	1.00	0.00
ATOM	640	CZ	ARG	1	72	-1.820	59.012	26.451	1.00	0.00
ATOM	641	NH1	ARG	1	72	-2.145	58.515	27.586	1.00	0.00
ATOM	642	NH2	ARG	1	72	-2.355	60.127	25.927	1.00	0.00
ATOM	643	H	ARG	1	72	-0.234	54.315	22.236	1.00	0.00
ATOM	644	HE	ARG	1	72	-0.329	58.774	24.988	1.00	0.00
ATOM	645	1HH1	ARG	1	72	-2.785	59.051	28.137	1.00	0.00
ATOM	646	2HH1	ARG	1	72	-1.825	57.675	28.024	1.00	0.00
ATOM	647	1HH2	ARG	1	72	-3.097	60.500	26.484	1.00	0.00
ATOM	648	2HH2	ARG	1	72	-2.152	60.596	25.068	1.00	0.00
ATOM	649	N	ILE	1	73	-3.075	54.932	24.098	1.00	0.00
ATOM	650	CA	ILE	1	73	-4.455	55.301	23.856	1.00	0.00
ATOM	651	C	ILE	1	73	-5.133	55.989	25.011	1.00	0.00
ATOM	652	O	ILE	1	73	-5.141	55.445	26.100	1.00	0.00
ATOM	653	CB	ILE	1	73	-5.320	54.019	23.424	1.00	0.00
ATOM	654	CG1	ILE	1	73	-6.850	54.247	23.209	1.00	0.00
ATOM	655	CG2	ILE	1	73	-5.101	52.911	24.470	1.00	0.00
ATOM	656	CD1	ILE	1	73	-7.640	52.943	22.791	1.00	0.00
ATOM	657	H	ILE	1	73	-2.810	53.985	24.276	1.00	0.00
ATOM	658	N	TYR	1	74	-5.586	57.222	24.931	1.00	0.00
ATOM	659	CA	TYR	1	74	-6.264	58.008	25.978	1.00	0.00
ATOM	660	C	TYR	1	74	-6.963	57.448	27.192	1.00	0.00
ATOM	661	O	TYR	1	74	-8.075	56.932	27.127	1.00	0.00
ATOM	662	CB	TYR	1	74	-7.256	58.973	25.383	1.00	0.00
ATOM	663	CG	TYR	1	74	-6.545	60.286	24.901	1.00	0.00
ATOM	664	CD1	TYR	1	74	-6.956	61.516	25.449	1.00	0.00
ATOM	665	CD2	TYR	1	74	-5.569	60.181	23.917	1.00	0.00
ATOM	666	CE1	TYR	1	74	-6.330	62.685	25.001	1.00	0.00
ATOM	667	CE2	TYR	1	74	-4.903	61.360	23.491	1.00	0.00
ATOM	668	CZ	TYR	1	74	-5.267	62.542	24.077	1.00	0.00
ATOM	669	OH	TYR	1	74	-4.459	63.643	23.767	1.00	0.00
ATOM	670	H	TYR	1	74	-5.424	57.697	24.066	1.00	0.00
ATOM	671	HH	TYR	1	74	-3.605	63.327	23.505	1.00	0.00
ATOM	672	N	HIS	1	75	-6.244	57.542	28.315	1.00	0.00
ATOM	673	CA	HIS	1	75	-6.712	57.415	29.691	1.00	0.00
ATOM	674	C	HIS	1	75	-5.809	58.147	30.656	1.00	0.00
ATOM	675	O	HIS	1	75	-4.618	58.037	30.567	1.00	0.00

FIG. 10

ATOM	676	CB	HIS	1	75	-6.903	55.932	30.164	1.00	0.00
ATOM	677	CG	HIS	1	75	-7.735	55.806	31.446	1.00	0.00
ATOM	678	ND1	HIS	1	75	-7.320	55.110	32.508	1.00	0.00
ATOM	679	CD2	HIS	1	75	-9.064	56.208	31.661	1.00	0.00
ATOM	680	CE1	HIS	1	75	-8.337	55.061	33.430	1.00	0.00
ATOM	681	NE2	HIS	1	75	-9.429	55.714	32.910	1.00	0.00
ATOM	682	H	HIS	1	75	-5.350	57.933	28.095	1.00	0.00
ATOM	683	1HD	HIS	1	75	-6.505	54.575	32.609	1.00	0.00
ATOM	684	N	PRO	1	76	-6.406	58.853	31.622	1.00	0.00
ATOM	685	CA	PRO	1	76	-5.631	59.218	32.828	1.00	0.00
ATOM	686	C	PRO	1	76	-4.689	58.242	33.509	1.00	0.00
ATOM	687	O	PRO	1	76	-3.545	58.615	33.836	1.00	0.00
ATOM	688	CB	PRO	1	76	-6.733	59.744	33.703	1.00	0.00
ATOM	689	CG	PRO	1	76	-8.079	59.421	33.110	1.00	0.00
ATOM	690	CD	PRO	1	76	-7.793	59.273	31.629	1.00	0.00
ATOM	691	N	ASN	1	77	-5.120	56.969	33.592	1.00	0.00
ATOM	692	CA	ASN	1	77	-4.149	55.995	34.023	1.00	0.00
ATOM	693	C	ASN	1	77	-3.232	55.350	32.975	1.00	0.00
ATOM	694	O	ASN	1	77	-2.235	54.686	33.293	1.00	0.00
ATOM	695	CB	ASN	1	77	-4.544	55.069	35.221	1.00	0.00
ATOM	696	CG	ASN	1	77	-5.398	53.859	34.799	1.00	0.00
ATOM	697	OD1	ASN	1	77	-5.269	53.390	33.660	1.00	0.00
ATOM	698	ND2	ASN	1	77	-6.264	53.533	35.724	1.00	0.00
ATOM	699	H	ASN	1	77	-6.008	56.604	33.313	1.00	0.00
ATOM	700	1HD2	ASN	1	77	-6.190	53.971	36.620	1.00	0.00
ATOM	701	2HD2	ASN	1	77	-6.951	52.815	35.616	1.00	0.00
ATOM	702	N	ILE	1	78	-3.577	55.500	31.719	1.00	0.00
ATOM	703	CA	ILE	1	78	-2.428	55.111	30.819	1.00	0.00
ATOM	704	C	ILE	1	78	-1.366	56.277	30.720	1.00	0.00
ATOM	705	O	ILE	1	78	-1.137	56.911	29.708	1.00	0.00
ATOM	706	CB	ILE	1	78	-2.914	54.640	29.485	1.00	0.00
ATOM	707	CG1	ILE	1	78	-4.001	53.586	29.802	1.00	0.00
ATOM	708	CG2	ILE	1	78	-1.718	53.970	28.701	1.00	0.00
ATOM	709	CD1	ILE	1	78	-4.663	53.021	28.578	1.00	0.00
ATOM	710	H	ILE	1	78	-4.240	56.191	31.434	1.00	0.00
ATOM	711	N	ASN	1	79	-0.702	56.476	31.895	1.00	0.00
ATOM	712	CA	ASN	1	79	0.362	57.504	31.918	1.00	0.00
ATOM	713	C	ASN	1	79	1.462	57.177	30.980	1.00	0.00
ATOM	714	O	ASN	1	79	1.783	56.010	30.852	1.00	0.00
ATOM	715	CB	ASN	1	79	0.977	57.697	33.357	1.00	0.00
ATOM	716	CG	ASN	1	79	1.977	58.941	33.383	1.00	0.00
ATOM	717	OD1	ASN	1	79	3.102	58.857	32.913	1.00	0.00
ATOM	718	ND2	ASN	1	79	1.421	59.936	34.047	1.00	0.00
ATOM	719	H	ASN	1	79	-0.879	55.970	32.740	1.00	0.00
ATOM	720	1HD2	ASN	1	79	0.480	59.871	34.380	1.00	0.00

FIG. 1P

ATOM	721	2HD2	ASN	1	79	1.919	60.793	34.181	1.00	0.00
ATOM	722	N	SER	1	80	2.012	58.274	30.296	1.00	0.00
ATOM	723	CA	SER	1	80	3.020	58.037	29.307	1.00	0.00
ATOM	724	C	SER	1	80	4.213	57.172	29.722	1.00	0.00
ATOM	725	O	SER	1	80	4.736	56.463	28.906	1.00	0.00
ATOM	726	CB	SER	1	80	3.540	59.374	28.740	1.00	0.00
ATOM	727	OG	SER	1	80	2.464	60.229	28.410	1.00	0.00
ATOM	728	H	SER	1	80	1.733	59.222	30.444	1.00	0.00
ATOM	729	HG	SER	1	80	2.142	59.977	27.555	1.00	0.00
ATOM	730	N	ASN	1	81	4.499	57.059	31.048	1.00	0.00
ATOM	731	CA	ASN	1	81	5.499	56.121	31.571	1.00	0.00
ATOM	732	C	ASN	1	81	5.049	54.722	31.893	1.00	0.00
ATOM	733	O	ASN	1	81	5.800	53.806	32.205	1.00	0.00
ATOM	734	CB	ASN	1	81	6.239	56.638	32.776	1.00	0.00
ATOM	735	CG	ASN	1	81	5.190	57.043	33.835	1.00	0.00
ATOM	736	OD1	ASN	1	81	4.206	56.324	33.995	1.00	0.00
ATOM	737	ND2	ASN	1	81	5.364	58.167	34.505	1.00	0.00
ATOM	738	H	ASN	1	81	4.021	57.645	31.702	1.00	0.00
ATOM	739	1HD2	ASN	1	81	6.256	58.616	34.550	1.00	0.00
ATOM	740	2HD2	ASN	1	81	4.596	58.637	34.941	1.00	0.00
ATOM	741	N	GLY	1	82	3.733	54.599	31.889	1.00	0.00
ATOM	742	CA	GLY	1	82	3.169	53.260	32.012	1.00	0.00
ATOM	743	C	GLY	1	82	2.883	52.770	33.467	1.00	0.00
ATOM	744	O	GLY	1	82	2.757	51.537	33.677	1.00	0.00
ATOM	745	H	GLY	1	82	3.081	55.355	31.943	1.00	0.00
ATOM	746	N	SER	1	83	2.699	53.686	34.408	1.00	0.00
ATOM	747	CA	SER	1	83	2.182	53.325	35.707	1.00	0.00
ATOM	748	C	SER	1	83	0.653	53.291	35.565	1.00	0.00
ATOM	749	O	SER	1	83	0.005	53.943	34.723	1.00	0.00
ATOM	750	CB	SER	1	83	2.747	54.306	36.709	1.00	0.00
ATOM	751	OG	SER	1	83	4.143	54.436	36.565	1.00	0.00
ATOM	752	H	SER	1	83	2.728	54.661	34.188	1.00	0.00
ATOM	753	HG	SER	1	83	4.292	55.334	36.830	1.00	0.00
ATOM	754	N	ILE	1	84	0.097	52.428	36.385	1.00	0.00
ATOM	755	CA	ILE	1	84	-1.370	52.476	36.461	1.00	0.00
ATOM	756	C	ILE	1	84	-1.850	52.509	37.900	1.00	0.00
ATOM	757	O	ILE	1	84	-1.165	51.998	38.770	1.00	0.00
ATOM	758	CB	ILE	1	84	-1.937	51.310	35.649	1.00	0.00
ATOM	759	CG1	ILE	1	84	-0.929	50.116	35.676	1.00	0.00
ATOM	760	CG2	ILE	1	84	-2.392	51.652	34.199	1.00	0.00
ATOM	761	CD1	ILE	1	84	-0.862	49.238	36.911	1.00	0.00
ATOM	762	H	ILE	1	84	0.585	51.890	37.073	1.00	0.00
ATOM	763	N	CYS	1	85	-3.050	53.104	38.044	1.00	0.00
ATOM	764	CA	CYS	1	85	-3.456	53.377	39.430	1.00	0.00
ATOM	765	C	CYS	1	85	-3.946	52.219	40.300	1.00	0.00

FIG. 1Q

ATOM	766	O	CYS	1	85	-3.154	51.631	40.977	1.00	0.00
ATOM	767	CB	CYS	1	85	-4.494	54.530	39.493	1.00	0.00
ATOM	768	SG	CYS	1	85	-3.719	56.051	38.899	1.00	0.00
ATOM	769	H	CYS	1	85	-3.555	53.567	37.316	1.00	0.00
ATOM	770	N	LEU	1	86	-5.235	51.872	40.178	1.00	0.00
ATOM	771	CA	LEU	1	86	-5.836	50.570	40.669	1.00	0.00
ATOM	772	C	LEU	1	86	-5.198	49.716	41.738	1.00	0.00
ATOM	773	O	LEU	1	86	-4.347	48.854	41.497	1.00	0.00
ATOM	774	CB	LEU	1	86	-6.178	49.687	39.435	1.00	0.00
ATOM	775	CG	LEU	1	86	-5.062	49.502	38.390	1.00	0.00
ATOM	776	CD1	LEU	1	86	-4.529	48.072	38.326	1.00	0.00
ATOM	777	CD2	LEU	1	86	-5.429	49.889	36.984	1.00	0.00
ATOM	778	H	LEU	1	86	-5.624	52.275	39.349	1.00	0.00
ATOM	779	N	ASP	1	87	-5.646	49.975	42.971	1.00	0.00
ATOM	780	CA	ASP	1	87	-5.525	49.028	44.088	1.00	0.00
ATOM	781	C	ASP	1	87	-4.109	48.504	44.263	1.00	0.00
ATOM	782	O	ASP	1	87	-3.363	49.082	45.066	1.00	0.00
ATOM	783	CB	ASP	1	87	-6.527	47.893	43.964	1.00	0.00
ATOM	784	CG	ASP	1	87	-6.431	46.788	45.001	1.00	0.00
ATOM	785	OD1	ASP	1	87	-6.114	47.067	46.154	1.00	0.00
ATOM	786	OD2	ASP	1	87	-6.744	45.628	44.680	1.00	0.00
ATOM	787	H	ASP	1	87	-6.293	50.722	43.123	1.00	0.00
ATOM	788	N	ILE	1	88	-3.811	47.376	43.540	1.00	0.00
ATOM	789	CA	ILE	1	88	-2.601	46.573	43.737	1.00	0.00
ATOM	790	C	ILE	1	88	-1.296	47.330	43.558	1.00	0.00
ATOM	791	O	ILE	1	88	-0.504	47.300	44.477	1.00	0.00
ATOM	792	CB	ILE	1	88	-2.715	45.322	42.836	1.00	0.00
ATOM	793	CG1	ILE	1	88	-2.555	45.432	41.275	1.00	0.00
ATOM	794	CG2	ILE	1	88	-4.113	44.647	42.978	1.00	0.00
ATOM	795	CD1	ILE	1	88	-2.522	44.051	40.547	1.00	0.00
ATOM	796	H	ILE	1	88	-4.311	47.348	42.674	1.00	0.00
ATOM	797	N	LEU	1	89	-1.124	48.012	42.405	1.00	0.00
ATOM	798	CA	LEU	1	89	-0.037	48.937	42.290	1.00	0.00
ATOM	799	C	LEU	1	89	-0.284	50.287	42.862	1.00	0.00
ATOM	800	O	LEU	1	89	-0.472	51.269	42.113	1.00	0.00
ATOM	801	CB	LEU	1	89	-0.443	49.094	40.838	1.00	0.00
ATOM	802	CG	LEU	1	89	1.420	48.043	40.288	1.00	0.00
ATOM	803	CD1	LEU	1	89	2.636	47.915	41.237	1.00	0.00
ATOM	804	CD2	LEU	1	89	0.675	46.722	39.962	1.00	0.00
ATOM	805	H	LEU	1	89	-1.933	48.260	41.873	1.00	0.00
ATOM	806	N	ARG	1	90	-0.418	50.340	44.189	1.00	0.00
ATOM	807	CA	ARG	1	90	-0.026	51.384	45.135	1.00	0.00
ATOM	808	C	ARG	1	90	-0.473	51.119	46.614	1.00	0.00
ATOM	809	O	ARG	1	90	0.338	50.878	47.480	1.00	0.00
ATOM	810	CB	ARG	1	90	-0.202	52.892	44.694	1.00	0.00

FIG. 1R

ATOM	811	CG	ARG	1	90	-1.614	53.327	44.386	1.00	0.00
ATOM	812	CD	ARG	1	90	-1.771	54.741	43.914	1.00	0.00
ATOM	813	NE	ARG	1	90	-3.140	55.176	43.940	1.00	0.00
ATOM	814	CZ	ARG	1	90	-3.469	56.121	43.129	1.00	0.00
ATOM	815	NH1	ARG	1	90	-4.669	56.743	43.266	1.00	0.00
ATOM	816	NH2	ARG	1	90	-2.772	56.487	42.010	1.00	0.00
ATOM	817	H	ARG	1	90	-0.600	49.477	44.659	1.00	0.00
ATOM	818	HE	ARG	1	90	-3.649	54.776	44.702	1.00	0.00
ATOM	819	1HH1	ARG	1	90	-4.832	57.565	42.721	1.00	0.00
ATOM	820	2HH1	ARG	1	90	-5.240	56.526	44.058	1.00	0.00
ATOM	821	1HH2	ARG	1	90	-3.123	57.247	41.464	1.00	0.00
ATOM	822	2HH2	ARG	1	90	-1.861	56.172	41.743	1.00	0.00
ATOM	823	N	SER	1	91	-1.779	51.132	46.880	1.00	0.00
ATOM	824	CA	SER	1	91	-2.246	50.840	48.231	1.00	0.00
ATOM	825	C	SER	1	91	-2.039	49.496	48.865	1.00	0.00
ATOM	826	O	SER	1	91	-2.155	49.290	50.084	1.00	0.00
ATOM	827	CB	SER	1	91	-3.706	51.221	48.212	1.00	0.00
ATOM	828	OG	SER	1	91	-3.945	52.494	47.464	1.00	0.00
ATOM	829	H	SER	1	91	-2.454	51.157	46.142	1.00	0.00
ATOM	830	HG	SER	1	91	-4.217	53.128	48.114	1.00	0.00
ATOM	831	N	GLN	1	92	-1.695	48.524	47.995	1.00	0.00
ATOM	832	CA	GLN	1	92	-1.634	47.187	48.509	1.00	0.00
ATOM	833	C	GLN	1	92	-0.236	46.696	48.873	1.00	0.00
ATOM	834	O	GLN	1	92	0.410	46.150	47.994	1.00	0.00
ATOM	835	CB	GLN	1	92	-2.345	46.228	47.634	1.00	0.00
ATOM	836	CG	GLN	1	92	-2.649	44.788	48.100	1.00	0.00
ATOM	837	CD	GLN	1	92	-3.525	44.132	47.065	1.00	0.00
ATOM	838	OE1	GLN	1	92	-3.152	43.262	46.300	1.00	0.00
ATOM	839	NE2	GLN	1	92	-4.730	44.710	46.949	1.00	0.00
ATOM	840	H	GLN	1	92	-1.391	48.636	47.050	1.00	0.00
ATOM	841	1HE2	GLN	1	92	-5.066	45.404	47.585	1.00	0.00
ATOM	842	2HE2	GLN	1	92	-5.388	44.626	46.200	1.00	0.00
ATOM	843	N	TRP	1	93	0.004	46.795	50.177	1.00	0.00
ATOM	844	CA	TRP	1	93	1.331	46.337	50.664	1.00	0.00
ATOM	845	C	TRP	1	93	1.379	44.847	50.786	1.00	0.00
ATOM	846	O	TRP	1	93	1.413	44.260	51.903	1.00	0.00
ATOM	847	CB	TRP	1	93	1.747	47.092	51.956	1.00	0.00
ATOM	848	CG	TRP	1	93	3.202	46.849	52.298	1.00	0.00
ATOM	849	CD1	TRP	1	93	3.639	46.472	53.600	1.00	0.00
ATOM	850	CD2	TRP	1	93	4.403	46.830	51.462	1.00	0.00
ATOM	851	NE1	TRP	1	93	4.988	46.241	53.629	1.00	0.00
ATOM	852	CE2	TRP	1	93	5.507	46.452	52349	1.00	0.00
ATOM	853	CE3	TRP	1	93	4.729	47.037	50.081	1.00	0.00
ATOM	854	CZ2	TRP	1	93	6.827	46.319	51.969	1.00	0.00
ATOM	855	CZ3	TRP	1	93	6.089	46.803	49.652	1.00	0.00

FIG. 1S

ATOM	856	CH2	GLN	1	93	7.087	46.544	50.614	1.00	0.00
ATOM	857	H	TRP	1	93	-0.649	47.178	50.831	1.00	0.00
ATOM	858	1HE	TRP	1	93	5.547	46.033	54.407	1.00	0.00
ATOM	859	N	SER	1	94	1.490	44.234	49.587	1.00	0.00
ATOM	860	CA	SER	1	94	1.669	42.774	49.495	1.00	0.00
ATOM	861	C	SER	1	94	2.312	42.502	48.181	1.00	0.00
ATOM	862	O	SER	1	94	1.754	42.899	47.152	1.00	0.00
ATOM	863	CB	SER	1	94	0.262	42.108	49.593	1.00	0.00
ATOM	864	OG	SER	1	94	-0.412	42.387	48.358	1.00	0.00
ATOM	865	H	SER	1	94	1.514	44.787	48.753	1.00	0.00
ATOM	866	HG	SER	1	94	-0.092	43.219	48.036	1.00	0.00
ATOM	867	N	PRO	1	95	3.435	41.722	48.161	1.00	0.00
ATOM	868	CA	PRO	1	95	3.724	41.167	46.814	1.00	0.00
ATOM	869	C	PRO	1	95	2.942	39.914	46.539	1.00	0.00
ATOM	870	O	PRO	1	95	3.473	38.913	46.075	1.00	0.00
ATOM	871	CB	PRO	1	95	5.243	40.974	46.986	1.00	0.00
ATOM	872	CG	PRO	1	95	5.541	40.527	48.437	1.00	0.00
ATOM	873	CD	PRO	1	95	4.449	41.477	49.162	1.00	0.00
ATOM	874	N	ALA	1	96	1.631	39.963	46.751	1.00	0.00
ATOM	875	CA	ALA	1	96	0.742	38.929	46.296	1.00	0.00
ATOM	876	C	ALA	1	96	0.008	39.214	45.017	1.00	0.00
ATOM	877	O	ALA	1	96	-1.215	39.302	44.930	1.00	0.00
ATOM	878	CB	ALA	1	96	-0.170	38.368	47.410	1.00	0.00
ATOM	879	H	ALA	1	96	1.330	40.893	46.964	1.00	0.00
ATOM	880	N	LEU	1	97	0.914	39.429	44.050	1.00	0.00
ATOM	881	CA	LEU	1	97	-.494	39.917	42.699	1.00	0.00
ATOM	882	C	LEU	1	97	0.773	38.935	41.618	1.00	0.00
ATOM	883	O	LEU	1	97	1.827	38.934	40.926	1.00	0.00
ATOM	884	CB	LEU	1	97	1.151	41.281	42.350	1.00	0.00
ATOM	885	CG	LEU	1	97	1.297	42.490	43.278	1.00	0.00
ATOM	886	CD1	LEU	1	97	1.815	43.697	42.585	1.00	0.00
ATOM	887	CD2	LEU	1	97	-0.071	42.794	44.045	1.00	0.00
ATOM	888	H	LEU	1	97	1.874	39.401	44.330	1.00	0.00
ATOM	889	N	THR	1	98	-0.195	38.052	41.512	1.00	0.00
ATOM	890	CA	THR	1	98	-0.030	37.084	40.473	1.00	0.00
ATOM	891	C	THR	1	98	-0.990	37.340	39.335	1.00	0.00
ATOM	892	O	THR	1	98	-2.041	37.953	39.480	1.00	0.00
ATOM	893	CB	THR	1	98	-0.163	35.572	41.034	1.00	0.00
ATOM	894	OG1	THR	1	98	0.762	34.750	40.274	1.00	0.00
ATOM	895	CG2	THR	1	98	-1.504	35.029	41.173	1.00	0.00
ATOM	896	H	THR	1	98	-1.027	38.138	42.060	1.00	0.00
ATOM	897	1HG	THR	1	98	0.266	34.090	39.808	1.00	0.00
ATOM	898	N	ILE	1	99	-0.363	37.052	38.234	1.00	0.00
ATOM	899	CA	ILE	1	99	-0.894	37.527	37.020	1.00	0.00
ATOM	900	C	ILE	1	99	-2.380	37.698	36.697	1.00	0.00

FIG. 1T

ATOM	901	O	ILE	1	99	-2.724	38.764	36.246	1.00	0.00
ATOM	902	CB	ILE	1	99	-0.082	37.024	35.825	1.00	0.00
ATOM	903	CG1	ILE	1	99	-0.308	37.613	34.450	1.00	0.00
ATOM	904	CG2	ILE	1	99	-0.305	35.483	35.600	1.00	0.00
ATOM	905	CD1	ILE	1	99	0.494	38.884	34.205	1.00	0.00
ATOM	906	H	ILE	1	99	0.345	36.354	38.343	1.00	0.00
ATOM	907	N	SER	1	100	-3.270	36.706	36.849	1.00	0.00
ATOM	908	CA	SER	1	100	-4.724	36.906	36.744	1.00	0.00
ATOM	909	C	SER	1	100	-5.204	38.211	37.355	1.00	0.00
ATOM	910	O	SER	1	100	-5.742	39.096	36.685	1.00	0.00
ATOM	911	CB	SER	1	100	-5.415	35.624	37.222	1.00	0.00
ATOM	912	OG	SER	1	100	-4.736	35.233	38.459	1.00	0.00
ATOM	913	H	SER	1	100	-2.868	35.938	37.348	1.00	0.00
ATOM	914	HG	SER	1	100	-5.204	34.561	38.936	1.00	0.00
ATOM	915	N	LYS	1	101	-4.768	38.359	38.639	1.00	0.00
ATOM	916	CA	LYS	1	101	-5.113	39.564	39.413	1.00	0.00
ATOM	917	C	LYS	1	101	-4.402	40.879	39.024	1.00	0.00
ATOM	918	O	LYS	1	101	-4.754	41.953	39.522	1.00	0.00
ATOM	919	CB	LYS	1	101	-4.912	39.238	40.866	1.00	0.00
ATOM	920	CG	LYS	1	101	-5.274	40.259	42.002	1.00	0.00
ATOM	921	CD	LYS	1	101	-3.984	40.688	42.680	1.00	0.00
ATOM	922	CE	LYS	1	101	-4.365	40.819	44.136	1.00	0.00
ATOM	923	NZ	LYS	1	101	-3.203	41.272	44.871	1.00	0.00
ATOM	924	H	LYS	1	101	-4.189	37.616	38.973	1.00	0.00
ATOM	925	1HZ	LYS	1	101	-2.880	42.183	44.487	1.00	0.00
ATOM	926	2HZ	LYS	1	101	-3.296	41.474	45.887	1.00	0.00
ATOM	927	3HZ	LYS	1	101	-2.424	40.594	44.747	1.00	0.00
ATOM	928	N	VAL	1	102	-3.475	40.750	38.077	1.00	0.00
ATOM	929	CA	VAL	1	102	-2.860	41.961	37.557	1.00	0.00
ATOM	930	C	VAL	1	102	-3.510	42.388	36.239	1.00	0.00
ATOM	931	O	VAL	1	102	-3.849	43.567	35.957	1.00	0.00
ATOM	932	CB	VAL	1	102	-1.348	41.798	37.381	1.00	0.00
ATOM	933	CG1	VAL	1	102	-0.665	43.196	37.067	1.00	0.00
ATOM	934	CG2	VAL	1	102	-0.609	41.199	38.633	1.00	0.00
ATOM	935	H	VAL	1	102	-3.259	39.834	37.736	1.00	0.00
ATOM	936	N	LEU	1	103	-3.744	41.406	35.354	1.00	0.00
ATOM	937	CA	LEU	1	103	-4.192	41.772	34.076	1.00	0.00
ATOM	938	C	LEU	1	103	-5.690	42.032	34.111	1.00	0.00
ATOM	939	O	LEU	1	103	-6.211	42.990	33.448	1.00	0.00
ATOM	940	CB	LEU	1	103	-3.726	40.740	32.995	1.00	0.00
ATOM	941	CG	LEU	1	103	-2.507	41.346	32.267	1.00	0.00
ATOM	942	CD1	LEU	1	103	-1.795	40.254	31.459	1.00	0.00
ATOM	943	CD2	LEU	1	103	-2.903	42.587	31.541	1.00	0.00
ATOM	944	H	LEU	1	103	-3.911	40.514	35.773	1.00	0.00
ATOM	945	N	LEU	1	104	-6.424	41.145	34.856	1.00	0.00

FIG. 1U

ATOM	946	CA	LEU	1	104	-7.810	41.575	35.127	1.00	0.00
ATOM	947	C	LEU	1	104	-7.855	42.961	35.724	1.00	0.00
ATOM	948	O	LEU	1	104	-8.500	43.819	35.130	1.00	0.00
ATOM	949	CB	LEU	1	104	-8.506	40.520	35.983	1.00	0.00
ATOM	950	CG	LEU	1	104	-9.953	40.895	36.094	1.00	0.00
ATOM	951	CD1	LEU	1	104	-10.722	40.812	34.846	1.00	0.00
ATOM	952	CD2	LEU	1	104	-10.702	40.028	37.125	1.00	0.00
ATOM	953	H	LEU	1	104	-6.131	40.287	35.278	1.00	0.00
ATOM	954	N	SER	1	105	-7.118	43.132	36.833	1.00	0.00
ATOM	955	CA	SER	1	105	-7.386	44.418	37.550	1.00	0.00
ATOM	956	C	SER	1	105	-6.973	45.720	36.778	1.00	0.00
ATOM	957	O	SER	1	105	-7.602	46.7373	36.986	1.00	0.00
ATOM	958	CB	SER	1	105	-6.761	44.419	38.909	1.00	0.00
ATOM	959	OG	SER	1	105	-7.437	43.369	39.676	1.00	0.00
ATOM	960	H	SER	1	105	-6.570	42.378	37.194	1.00	0.00
ATOM	961	HG	SER	1	105	-7.053	42.532	39.449	1.00	0.00
ATOM	962	N	ILE	1	106	-5.953	45.587	35.942	1.00	0.00
ATOM	963	CA	ILE	1	106	-5.565	46.649	34.950	1.00	0.00
ATOM	964	C	ILE	1	106	-6.694	46.937	33.906	1.00	0.00
ATOM	965	O	ILE	1	106	-7.039	48.044	33.459	1.00	0.00
ATOM	966	CB	ILE	1	106	-4.240	46.330	34.320	1.00	0.00
ATOM	967	CG1	ILE	1	106	-3.013	46.371	35.246	1.00	0.00
ATOM	968	CG2	ILE	1	106	-3.926	47.235	33.094	1.00	0.00
ATOM	969	CD1	ILE	1	106	-1.919	45.558	34.629	1.00	0.00
ATOM	970	H	ILE	1	106	-5.752	44.610	35.861	1.00	0.00
ATOM	971	N	CYS	1	107	-7.274	45.781	33.409	1.00	0.00
ATOM	972	CA	CYS	1	107	-8.520	45.970	32.586	1.00	0.00
ATOM	973	C	CYS	1	107	-9.757	46.433	33.338	1.00	0.00
ATOM	974	O	CYS	1	107	-10.600	47.145	32.731	1.00	0.00
ATOM	975	CB	CYS	1	107	-8.856	44.670	31.741	1.00	0.00
ATOM	976	SG	CYS	1	107	-7.429	44.342	30.720	1.00	0.00
ATOM	977	H	CYS	1	107	-7.117	44.941	33.928	1.00	0.00
ATOM	978	N	SER	1	108	-9.830	46.108	34.695	1.00	0.00
ATOM	979	CA	SER	1	108	-11.060	46.479	35.328	1.00	0.00
ATOM	980	C	SER	1	108	-11.219	47.952	35.369	1.00	0.00
ATOM	981	O	SER	1	108	-12.337	48.405	35.150	1.00	0.00
ATOM	982	CB	SER	1	108	-11.152	45.799	36.770	1.00	0.00
ATOM	983	OG	SER	1	108	-11.173	44.322	36.587	1.00	0.00
ATOM	984	H	SER	1	108	-9.241	45.357	34.991	1.00	0.00
ATOM	985	HG	SER	1	108	-11.732	44.168	35.837	1.00	0.00
ATOM	986	N	LEU	1	109	-10.155	48.631	35.780	1.00	0.00
ATOM	987	CA	LEU	1	109	-10.220	50.104	35.653	1.00	0.00
ATOM	988	C	LEU	1	109	-9.817	50.687	34.332	1.00	0.00
ATOM	989	O	LEU	1	109	-9.444	51.821	34.154	1.00	0.00
ATOM	990	CB	LEU	1	109	-9.558	50.849	36.830	1.00	0.00

FIG. 1V

ATOM	991	CG	LEU	1	109	-10.330	50.611	38.192	1.00	0.00
ATOM	992	CD1	LEU	1	109	-9.530	51.170	39.297	1.00	0.00
ATOM	993	CD2	LEU	1	109	-11.727	51.119	38.162	1.00	0.00
ATOM	994	H	LEU	1	109	-9.232	48.266	35.900	1.00	0.00
ATOM	995	N	LEU	1	110	-9.934	49.766	33.374	1.00	0.00
ATOM	996	CA	LEU	1	110	-10.049	50.216	32.023	1.00	0.00
ATOM	997	C	LEU	1	110	-11.371	49.745	31.298	1.00	0.00
ATOM	998	O	LEU	1	110	-11.455	49.615	30.102	1.00	0.00
ATOM	999	CB	LEU	1	110	-8.751	49.816	31.312	1.00	0.00
ATOM	1000	CG	LEU	1	110	-7.942	50.875	30.483	1.00	0.00
ATOM	1001	CD1	LEU	1	110	-7.368	51.879	31.470	1.00	0.00
ATOM	1002	CD2	LEU	1	110	-6.829	50.314	29.656	1.00	0.00
ATOM	1003	H	LEU	1	110	-10.041	48.784	33.533	1.00	0.00
ATOM	1004	N	CYS	1	111	-12.308	49.338	32.152	1.00	0.00
ATOM	1005	CA	CYS	1	111	-13.548	48.999	31.551	1.00	0.00
ATOM	1006	C	CYS	1	111	-14.398	50.255	31.331	1.00	0.00
ATOM	1007	O	CYS	1	111	-14.130	50.937	30.370	1.00	0.00
ATOM	1008	CB	CYS	1	111	-14.416	48.026	32.492	1.00	0.00
ATOM	1009	SG	CYS	1	111	-14.116	46.278	32.379	1.00	0.00
ATOM	1010	H	CYS	1	111	-12.026	49.328	33.112	1.00	0.00
ATOM	1011	N	ASP	1	112	-15.364	50.587	32.178	1.00	0.00
ATOM	1012	CA	ASP	1	112	-15.593	51.987	32.403	1.00	0.00
ATOM	1013	C	ASP	1	112	-14.518	52.650	33.331	1.00	0.00
ATOM	1014	O	ASP	1	112	-14.507	52.386	34.545	1.00	0.00
ATOM	1015	CB	ASP	1	112	-17.121	52.182	32.807	1.00	0.00
ATOM	1016	CG	ASP	1	112	-17.888	52.937	31.766	1.00	0.00
ATOM	1017	OD1	ASP	1	112	-19.113	52.989	31.785	1.00	0.00
ATOM	1018	OD2	ASP	1	112	-17.199	53.537	30.912	1.00	0.00
ATOM	1019	H	ASP	1	112	-15.765	49.992	32.874	1.00	0.00
ATOM	1020	N	PRO	1	113	-13.531	53.288	32.615	1.00	0.00
ATOM	1021	CA	PRO	1	113	-12.199	53.281	33.187	1.00	0.00
ATOM	1022	C	PRO	1	113	-12.012	54.384	34.243	1.00	0.00
ATOM	1023	O	PRO	1	113	-12.637	55.445	34.025	1.00	0.00
ATOM	1024	CB	PRO	1	113	-11.398	53.419	31.897	1.00	0.00
ATOM	1025	CG	PRO	1	113	-12.171	54.275	30.881	1.00	0.00
ATOM	1026	CD	PRO	1	113	-13.610	53.875	31.275	1.00	0.00
ATOM	1027	N	ASN	1	114	-11.261	54.180	35.336	1.00	0.00
ATOM	1028	CA	ASN	1	114	-11.182	55.245	36.355	1.00	0.00
ATOM	1029	C	ASN1	1	114	-10.597	56.637	35.865	1.00	0.00
ATOM	1030	O	ASN	1	114	-9.378	56.803	35.714	1.00	0.00
ATOM	1031	CB	ASN	1	114	-10.330	54.764	37.541	1.00	0.00
ATOM	1032	CG	ASN	1	114	-10.137	55.772	38.647	1.00	0.00
ATOM	1033	OD1	ASN	1	114	-9.345	56.731	38.656	1.00	0.00
ATOM	1034	ND2	ASN	1	114	-10.980	55.677	39.678	1.00	0.00
ATOM	1035	H	ASN	1	114	-10.615	53.417	35.332	1.00	0.00

FIG. 1W

ATOM	1036	1HD2	ASN	1	114	-11.523	54.841	39.754	1.00	0.00
ATOM	1037	2HD2	ASN	1	114	-11.318	56.332	40.354	1.00	0.00
ATOM	1038	N	PRO	1	115	-11.568	57.547	35.930	1.00	0.00
ATOM	1039	CA	PRO	1	115	-11.202	58.938	35.634	1.00	0.00
ATOM	1040	C	PRO	1	115	-10.529	59.625	36.725	1.00	0.00
ATOM	1041	O	PRO	1	115	-9.921	60.690	36.497	1.00	0.00
ATOM	1042	CB	PRO	1	115	-12.673	59.470	35.330	1.00	0.00
ATOM	1043	CG	PRO	1	115	-13.720	58.740	58.740	1.00	0.00
ATOM	1044	CD	PRO	1	115	-12.975	57.417	36.159	1.00	0.00
ATOM	1045	N	ASP	1	116	-10.763	59.056	37.948	1.00	0.00
ATOM	1046	CA	ASP	1	116	-10.371	59.852	39.097	1.00	0.00
ATOM	1047	C	ASP	1	116	-8.965	60.130	39.381	1.00	0.00
ATOM	1048	O	ASP	1	116	-8.729	61.091	40.080	1.00	0.00
ATOM	1049	CB	ASP	1	116	-11.112	59.169	40.210	1.00	0.00
ATOM	1050	CG	ASP	1	116	-12.562	59.525	40.004	1.00	0.00
ATOM	1051	OD1	ASP	1	116	-12.897	60.693	40.229	1.00	0.00
ATOM	1052	OD2	ASP	1	116	-13.360	58.645	39.645	1.00	0.00
ATOM	1053	H	ASP	1	116	-11.087	58.122	38.101	1.00	0.00
ATOM	1054	N	ASP	1	117	-8.046	59.379	38.889	1.00	0.00
ATOM	1055	CA	ASP	1	117	-6.667	59.696	39.115	1.00	0.00
ATOM	1056	C	ASP	1	117	-6.000	60.362	37.956	1.00	0.00
ATOM	1057	O	ASP	1	117	-5.912	59.906	36.817	1.00	0.00
ATOM	1058	CB	ASP	1	117	-5.812	58.462	39.672	1.00	0.00
ATOM	1059	CG	ASP	1	117	-5.071	58.862	40.969	1.00	0.00
ATOM	1060	OD1	ASP	1	117	-5.706	59.224	41.919	1.00	0.00
ATOM	1061	OD2	ASP	1	117	-3.839	58.861	40.982	1.00	0.00
ATOM	1062	H	ASP	1	117	-8.297	58.676	38.225	1.00	0.00
ATOM	1063	N	PRO	1	118	-5.510	61.616	38.265	1.00	0.00
ATOM	1064	CA	PRO	1	118	-4.974	62.450	37.154	1.00	0.00
ATOM	1065	C	PRO	1	118	-3.551	62.218	36.754	1.00	0.00
ATOM	1066	O	PRO	1	118	-2.677	63.093	36.734	1.00	0.00
ATOM	1067	CB	PRO	1	118	-5.149	63.800	37.805	1.00	0.00
ATOM	1068	CG	PRO	1	118	-4.936	63.597	39.311	1.00	0.00
ATOM	1069	CD	PRO	1	118	-5.578	62.267	39.561	1.00	0.00
ATOM	1070	N	LEU	1	119	-3.257	60.906	36.463	1.00	0.00
ATOM	1071	CA	LEU	1	119	-1.905	60.688	35.982	1.00	0.00
ATOM	1072	C	LEU	1	119	-1.307	61.479	34.745	1.00	0.00
ATOM	1073	O	LEU	1	119	-0.441	62.318	34.908	1.00	0.00
ATOM	1074	CB	LEU	1	119	-1.456	59.254	35.930	1.00	0.00
ATOM	1075	CG	LEU	1	119	-1.532	58.349	37.167	1.00	0.00
ATOM	1076	CD1	LEU	1	119	-0.912	56.912	36.972	1.00	0.00
ATOM	1077	CD2	LEU	1	119	-0.790	58.990	38.341	1.00	0.00
ATOM	1078	H	LEU	1	119	-3.921	60.164	36.368	1.00	0.00
ATOM	1079	N	VAL	1	120	-1.971	61.216	33.617	1.00	0.00
ATOM	1080	CA	VAL	1	120	-1.959	62.209	32.538	1.00	0.00

FIG. 1X

ATOM	1081	C	VAL	1	120	3.329	62.894	32.518	1.00	0.00
ATOM	1082	O	VAL	1	120	-4.360	62.293	32.613	1.00	0.00
ATOM	1083	CB	VAL	1	120	-1.499	61.604	31.253	1.00	0.00
ATOM	1084	CG1	VAL	1	120	0.035	61.659	31.162	1.00	0.00
ATOM	1085	CG2	VAL	1	120	-2.097	60.128	30.928	1.00	0.00
ATOM	1086	H	VAL	1	120	-2.662	60.500	33.718	1.00	0.00
ATOM	1087	N	PRO	1	121	-3.277	64.280	32.514	1.00	0.00
ATOM	1088	CA	PRO	1	121	-4.501	65.112	32.553	1.00	0.00
ATOM	1089	C	PRO	1	121	-5.183	65.158	31.223	1.00	0.00
ATOM	1090	O	PRO	1	121	-5.005	66.142	30.408	1.00	0.00
ATOM	1091	CB	PRO	1	121	-4.078	66.545	32.890	1.00	0.00
ATOM	1092	CG	PRO	1	121	-2.593	66.574	32.951	1.00	0.00
ATOM	1093	CD	PRO	1	121	-2.099	65.167	32.581	1.00	0.00
ATOM	1094	N	GLU	1	122	-5.946	64.072	30.888	1.00	0.00
ATOM	1095	CA	GLU	1	122	-6.357	64.028	29.506	1.00	0.00
ATOM	1096	C	GLU	1	122	-7.790	64.010	29.329	1.00	0.00
ATOM	1097	O	GLU	1	122	-8.390	64.999	28.991	1.00	0.00
ATOM	1098	CB	GLU	1	122	-5.562	62.967	28.680	1.00	0.00
ATOM	1099	CG	GLU	1	122	-5.508	61.536	29.110	1.00	0.00
ATOM	1100	CD	GLU	1	122	-4.428	60.841	28.287	1.00	0.00
ATOM	1101	OE1	GLU	1	122	-4.533	59.662	28.084	1.00	0.00
ATOM	1102	OE2	GLU	1	122	-3.419	61.458	27.964	1.00	0.00
ATOM	1103	H	GLU	1	122	-5.977	63.325	31.552	1.00	0.00
ATOM	1104	N	ILE	1	123	-8.356	62.858	29.591	1.00	0.00
ATOM	1105	CA	ILE	1	123	-9.817	62.884	29.471	1.00	0.00
ATOM	1106	C	ILE	1	123	-10.637	62.807	30.714	1.00	0.00
ATOM	1107	O	ILE	1	123	-11.593	63.548	30.791	1.00	0.00
ATOM	1108	CB	ILE	1	123	-10.141	61.868	28.403	1.00	0.00
ATOM	1109	CG1	ILE	1	123	-11.547	62.072	27.779	1.00	0.00
ATOM	1110	CG2	ILE	1	123	-10.056	60.403	28.786	1.00	0.00
ATOM	1111	CD1	ILE	1	123	-11.761	63.221	26.711	1.00	0.00
ATOM	1112	H	ILE	1	123	-7.841	62.124	30.034	1.00	0.00
ATOM	1113	N	ALA	1	124	-10.185	61.872	31.630	1.00	0.00
ATOM	1114	CA	ALA	1	124	-10.610	61.878	33.046	1.00	0.00
ATOM	1115	C	ALA	1	124	-12.162	62.048	33.276	1.00	0.00
ATOM	1116	O	ALA	1	124	-13.048	61.516	32.542	1.00	0.00
ATOM	1117	CB	ALA	1	124	-9.795	63.009	33.690	1.00	0.00
ATOM	1118	H	ALA	1	124	-9.351	61.353	31.444	1.00	0.00
ATOM	1119	N	ARG	1	125	-12.434	62.735	34.377	1.00	0.00
ATOM	1120	CA	ARG	1	125	-13.795	63.212	34.682	1.00	0.00
ATOM	1121	C	ARG	1	125	-14.747	63.817	33.616	1.00	0.00
ATOM	1122	O	ARG	1	125	-15.960	63.759	33.744	1.00	0.00
ATOM	1123	CB	ARG	1	125	-13.769	64.204	35.773	1.00	0.00
ATOM	1124	CG	ARG	1	125	-13.675	63.377	37.072	1.00	0.00
ATOM	1125	CD	ARG	1	125	-14.972	62.537	37.191	1.00	0.00

FIG. 1Y

ATOM	1126	NE	ARG	1	125	-14.854	61.500	38.232	1.00	0.00
ATOM	1127	CZ	ARG	1	125	-15.861	60.967	38.847	1.00	0.00
ATOM	1128	NH1	ARG	1	125	-15.655	59.990	39.749	1.00	0.00
ATOM	1129	NH2	ARG	1	125	-17.122	61.349	38.534	1.00	0.00
ATOM	1130	H	ARG	1	125	-11.855	63.259	35.001	1.00	0.00
ATOM	1131	HE	ARG	1	125	-13.919	61.221	38.454	1.00	0.00
ATOM	1132	1HH1	ARG	1	125	-16.416	59.493	40.167	1.00	0.00
ATOM	1133	2HH1	ARG	1	125	-14.748	59.735	40.086	1.00	0.00
ATOM	1134	1HH2	ARG	1	125	-17.856	61.089	39.162	1.00	0.00
ATOM	1135	2HH2	ARG	1	125	-17.314	61.932	37.745	1.00	0.00
ATOM	1136	N	ILE	1	126	-14.127	64.432	32.608	1.00	0.00
ATOM	1137	CA	ILE	1	126	-14.788	64.980	31.466	1.00	0.00
ATOM	1138	C	ILE	1	126	-15.410	63.856	30.692	1.00	0.00
ATOM	1139	O	ILE	1	126	-16.523	63.970	30.167	1.00	0.00
ATOM	1140	CB	ILE	1	126	-13.766	65.724	30.522	1.00	0.00
ATOM	1141	CG1	ILE	1	126	-13.082	66.752	31.461	1.00	0.00
ATOM	1142	CG2	ILE	1	126	-14.467	66.385	29.299	1.00	0.00
ATOM	1143	CD1	ILE	1	126	-11.753	67.296	30.873	1.00	0.00
ATOM	1144	H	ILE	1	126	-13.135	64.303	32.601	1.00	0.00
ATOM	1145	N	TYR	1	127	-14.624	62.715	30.636	1.00	0.00
ATOM	1146	CA	TYR	1	127	-15.167	61.452	30.126	1.00	0.00
ATOM	1147	C	TYR	1	127	-16.529	61.083	30.734	1.00	0.00
ATOM	1148	O	TYR	1	127	-17.440	60.554	30.100	1.00	0.00
ATOM	1149	CB	TYR	1	127	-14.051	60.272	30.134	1.00	0.00
ATOM	1150	CG	TYR	1	127	-14.489	59.022	30.878	1.00	0.00
ATOM	1151	CD1	TYR	1	127	-15.254	58.078	30.129	1.00	0.00
ATOM	1152	CD2	TYR	1	127	-14.209	58.796	32.260	1.00	0.00
ATOM	1153	CE1	TYR	1	127	-15.911	56.995	30.838	1.00	0.00
ATOM	1154	CE2	TYR	1	127	-14.809	57.674	32.912	1.00	0.00
ATOM	1155	CZ	TYR	1	127	-15.750	56.817	32.219	1.00	0.00
ATOM	1156	OH	TYR	1	127	-16.381	55.818	32.877	1.00	0.00
ATOM	1157	H	TYR	1	127	-13.751	62.624	31.115	1.00	0.00
ATOM	1158	HH	TYR	1	127	-16.924	55.379	32.236	1.00	0.00
ATOM	1159	N	GLN	1	128	-16.689	61.447	32.006	1.00	0.00
ATOM	1160	CA	GLN	1	128	-17.972	61.076	32.613	1.00	0.00
ATOM	1161	C	GLN	1	128	-18.820	62.307	32.767	1.00	0.00
ATOM	1162	O	GLN	1	128	-19.700	62.437	33.627	1.00	0.00
ATOM	1163	CB	GLN	1	128	-17.573	60.544	33.940	1.00	0.00
ATOM	1164	CG	GLN	1	128	-18.401	59.339	34.361	1.00	0.00
ATOM	1165	CD	GLN	1	128	-17.669	58.576	35.474	1.00	0.00
ATOM	1166	OE1	GLN	1	128	-17.437	59.077	36.587	1.00	0.00
ATOM	1067	NE2	GLN	1	128	-17.060	57.418	35.047	1.00	0.00
ATOM	1068	H	GLN	1	128	-15.933	61.976	32.392	1.00	0.00
ATOM	1069	1HE2	GLN	1	128	-17.400	56.974	34.218	1.00	0.00
ATOM	1070	2HE2	GLN	1	128	-16.287	57.080	35.584	1.00	0.00

FIG. 1Z

ATOM	1171	N	THR	1	129	-18.610	63.261	31.790	1.00	0.00
ATOM	1172	CA	THR	1	129	-19.554	64.402	31.531	1.00	0.00
ATOM	1173	C	THR	1	129	-20.034	64.338	30.090	1.00	0.00
ATOM	1174	O	THR	1	129	-21.188	64.645	29.868	1.00	0.00
ATOM	1175	CB	THR	1	129	-18.694	65.689	31.759	1.00	0.00
ATOM	1176	OG1	THR	1	129	-17.843	65.533	32.883	1.00	0.00
ATOM	1177	CG2	THR	1	129	-19.575	66.877	31.994	1.00	0.00
ATOM	1178	H	THR	1	129	-17.822	63.069	31.205	1.00	0.00
ATOM	1179	1HG	THR	1	129	-17.276	64.776	32.808	1.00	0.00
ATOM	1180	N	ASP	1	130	-19.100	63.839	29.242	1.00	0.00
ATOM	1181	CA	ASP	1	130	-19.272	63.530	27.815	1.00	0.00
ATOM	1182	C	ASP	1	130	-18.547	62.268	27.468	1.00	0.00
ATOM	1183	O	ASP	1	130	-17.360	62.026	27.412	1.00	0.00
ATOM	1184	CB	ASP	1	130	-18.822	64.757	26.978	1.00	0.00
ATOM	1185	CG	ASP	1	130	-18.897	64.631	25.454	1.00	0.00
ATOM	1186	OD1	ASP	1	130	-19.920	64.947	24.888	1.00	0.00
ATOM	1187	OD2	ASP	1	130	017.905	64.242	24.882	1.00	0.00
ATOM	1188	H	ASP	1	130	-18.255	63.427	29.582	1.00	0.00
ATOM	1189	N	ARG	1	131	-19.410	61.274	27.309	1.00	0.00
ATOM	1190	CA	ARG	1	131	-18.935	59.959	26.768	1.00	0.00
ATOM	1191	C	ARG	1	131	-18.469	60.154	25.366	1.00	0.00
ATOM	1192	O	ARG	1	131	-17.594	59.446	24.815	1.00	0.00
ATOM	1193	CB	ARG	1	131	-20.032	58.855	26.756	1.00	0.00
ATOM	1194	CG	ARG	1	131	-20.416	58.478	28.193	1.00	0.00
ATOM	1195	CD	ARG	1	131	-19.194	57.782	28.955	1.00	0.00
ATOM	1196	NE	ARG	1	131	-18.986	56.441	28.423	1.00	0.00
ATOM	1197	CZ	ARG	1	131	-19.071	55.351	29.184	1.00	0.00
ATOM	1198	NH1	ARG	1	131	-18.900	54.185	28.671	1.00	0.00
ATOM	1199	NH2	ARG	1	131	-19.376	55.517	30.503	1.00	0.00
ATOM	1200	H	ARG	1	131	-20.369	61.548	27.370	1.00	0.00
ATOM	1201	HE	ARG	1	131	-18.632	56.469	27.488	1.00	0.00
ATOM	1202	1HH1	ARG	1	131	-18.956	53.332	29.191	1.00	0.00
ATOM	1203	2HH1	ARG	1	131	-18.792	54.024	27.690	1.00	0.00
ATOM	1204	1HH2	ARG	1	131	-19.481	54.718	31.095	1.00	0.00
ATOM	1205	2HH2	ARG	1	131	-19.542	56.412	30.918	1.00	0.00
ATOM	1206	N	GLU	1	132	-19.073	61.146	24.679	1.00	0.00
ATOM	1207	CA	GLU	1	132	-19.097	61.123	23.223	1.00	0.00
ATOM	1208	C	GLU	1	132	-17.704	61.256	22.550	1.00	0.00
ATOM	1209	O	GLU	1	132	-17.303	60.476	21.676	1.00	0.00
ATOM	1210	CB	GLU	1	132	-20.158	62.165	22.829	1.00	0.00
ATOM	1211	CG	GLU	1	132	-20.663	62.490	21.437	1.00	0.00
ATOM	1212	CD	GLU	1	132	-19.538	62.468	20.507	1.00	0.00
ATOM	1213	OE1	GLU	1	132	-18.751	63.417	20.464	1.00	0.00
ATOM	1214	OE2	GLU	1	132	-19.328	61.449	19.842	1.00	0.00
ATOM	1215	H	GLU	1	132	-19.447	61.944	25.151	1.00	0.00

FIG. 1AA

ATOM	1216	N	LYS	1	133	-16.973	62.276	23.084	1.00	0.00
ATOM	1217	CA	LYS	1	133	-15.616	62.452	22.586	1.00	0.00
ATOM	1218	C	LYS	1	133	-14.629	61.391	22.856	1.00	0.00
ATOM	1219	O	LYS	1	133	-13.837	60.980	22.044	1.00	0.00
ATOM	1220	CB	LYS	1	133	-15.115	63.846	22.950	1.00	0.00
ATOM	1221	CG	LYS	1	133	-15.678	64.917	22.035	1.00	0.00
ATOM	1222	CD	LYS	1	133	-16.243	66.106	22.825	1.00	0.00
ATOM	1223	CE	LYS	1	133	-17.623	66.499	22.255	1.00	0.00
ATOM	1224	NZ	LYS	1	133	-18.680	65.449	22.4551	1.00	0.00
ATOM	1225	H	LYS	1	133	-17.198	62.759	23.930	1.00	0.00
ATOM	1226	1HZ	LYS	1	133	-18.725	64.799	21.645	1.00	0.00
ATOM	1227	2HZ	LYS	1	133	-18.439	64.912	23.313	1.00	0.00
ATOM	1228	3HZ	LYS	1	133	-19.642	65.786	22.663	1.00	0.00
ATOM	1229	N	TYR	1	134	-14.790	60.794	24.068	1.00	0.00
ATOM	1230	CA	TYR	1	134	-14.077	59.548	24.365	1.00	0.00
ATOM	1231	C	TYR	1	134	-14.477	58.281	23.495	1.00	0.00
ATOM	1232	O	TYR	1	134	-13.564	57.463	23.191	1.00	0.00
ATOM	1233	CB	TYR	1	134	-14.163	59.285	25.901	1.00	0.00
ATOM	1234	CG	TYR	1	134	-13.902	57.807	26.279	1.00	0.00
ATOM	1235	CD1	TYR	1	134	-12.616	57.238	26.234	1.00	0.00
ATOM	1236	CD2	TYR	1	143	-14.952	56.954	26.703	1.00	0.00
ATOM	1237	CE1	TYR	1	134	-12.308	55.950	26.691	1.00	0.00
ATOM	1238	CE2	TYR	1	134	-14.739	55.641	27.144	1.00	0.00
ATOM	1239	CZ	TYR	1	134	-13.388	55.147	27.137	1.00	0.00
ATOM	1240	OH	TYR	1	134	-13.089	53.833	27.509	1.00	0.00
ATOM	1241	H	TYR	1	134	-15.605	61.016	24.604	1.00	0.00
ATOM	1242	HH	TYR	1	134	-12.629	53.955	28.329	1.00	0.00
ATOM	1243	N	ASN	1	135	-15.732	58.290	23.026	1.00	0.00
ATOM	1244	CA	ASN	1	135	-16.023	57.578	21.793	1.00	0.00
ATOM	1245	C	ASN	1	135	-15.345	58.003	20.446	1.00	0.00
ATOM	1246	O	ASN	1	135	-16.007	57.968	19.385	1.00	0.00
ATOM	1247	CB	ASN	1	135	-17.458	57.139	21.710	1.00	0.00
ATOM	1248	CG	ASN	1	135	-17.585	55.762	22.262	1.00	0.00
ATOM	1249	OD1	ASN	1	135	-17.810	55.462	23.457	1.00	0.00
ATOM	1250	ND2	ASN	1	135	-17.359	54.819	21.344	1.00	0.00
ATOM	1251	H	ASN	1	135	-16.422	58.886	23.436	1.00	0.00
ATOM	1252	1HD2	ASN	1	135	-17.077	54.977	20.398	1.00	0.00
ATOM	1253	2HD2	ASN	1	135	-17.430	53.873	21.660	1.00	0.00
ATOM	1254	N	ARG	1	136	-14.053	58.370	20.478	1.00	0.00
ATOM	1255	CA	ARG	1	136	-13.385	58.938	19.318	1.00	0.00
ATOM	1256	C	ARG	1	136	-11.907	59.136	19.706	1.00	0.00
ATOM	1257	O	ARG	1	136	-11.120	58.192	19.624	1.00	0.00
ATOM	1258	CB	ARG	1	136	-13.963	60.323	18.860	1.00	0.00
ATOM	1259	CG	ARG	1	136	-13.759	60.585	17.391	1.00	0.00
ATOM	1260	CD	ARG	1	136	-15.027	60.276	16.539	1.00	0.00

FIG. 1BB

ATOM	1261	NE	ARG	1	136	-16.001	61.343	16.710	1.00	0.00
ATOM	1262	CZ	ARG	1	136	-16.953	61.385	17.722	1.00	0.00
ATOM	1263	NH1	ARG	1	136	-17.634	62.501	17.758	1.00	0.00
ATOM	1264	NH2	ARG	1	136	-17.160	60.427	18.572	1.00	0.00
ATOM	1265	H	ARG	1	136	-13.577	58.270	21.352	1.00	0.00
ATOM	1266	HE	ARG	1	136	-15.978	62.141	16.108	1.00	0.00
ATOM	1267	1HH1	ARG	1	136	-18.194	62.691	18.565	1.00	0.00
ATOM	1268	2HH1	ARG	1	136	-17.585	63.094	16.955	1.00	0.00
ATOM	1269	1HH2	ARG	1	136	-17.955	60.445	19.178	1.00	0.00
ATOM	1270	2HH2	ARG	1	136	-16.527	59.655	18.631	1.00	0.00
ATOM	1271	N	ILE	1	137	-11.553	60.299	20.250	1.00	0.00
ATOM	1272	CA	ILE	1	137	-10.216	60.181	20.894	1.00	0.00
ATOM	1273	C	ILE	1	137	-10.249	59.245	22.086	1.00	0.00
ATOM	1274	O	ILE	1	137	-10.876	59.502	23.085	1.00	0.00
ATOM	1275	CB	ILE	1	137	-9.737	61.669	21.177	1.00	0.00
ATOM	1276	CG1	ILE	1	137	-10.160	62.400	22.437	1.00	0.00
ATOM	1277	CG2	ILE	1	137	-10.058	62.553	19.908	1.00	0.00
ATOM	1278	CD1	ILE	1	137	-9.379	63.684	22.863	1.00	0.00
ATOM	1279	H	ILE	1	137	-12.189	61.029	20.500	1.00	0.00
ATOM	1280	N	ALA	1	138	-9.665	58.091	21.783	1.00	0.00
ATOM	1281	CA	ALA	1	138	-9.599	56.705	22.426	1.00	0.00
ATOM	1282	C	ALA	1	138	-10.195	55.649	21.536	1.00	0.00
ATOM	1283	O	ALA	1	138	-9.601	54.689	21.054	1.00	0.00
ATOM	1284	CB	ALA	1	138	-10.297	56.767	23.804	1.00	0.00
ATOM	1285	H	ALA	1	138	-9.100	58.150	20.960	1.00	0.00
ATOM	1286	N	ARG	1	139	-11.487	55.821	21.280	1.00	0.00
ATOM	1287	CA	ARG	1	139	-11.974	55.035	20.159	1.00	0.00
ATOM	1288	C	ARG	1	139	-11.531	55.522	18.757	1.00	0.00
ATOM	1289	O	ARG	1	139	-12.365	55.995	17.960	1.00	0.00
ATOM	1290	CB	ARG	1	139	-13.446	54.877	20.345	1.00	0.00
ATOM	1291	CG	ARG	1	139	-14.037	53.544	20.350	1.00	0.00
ATOM	1292	CD	ARG	1	139	-13.777	52.739	19.100	1.00	0.00
ATOM	1293	NE	ARG	1	139	-13.294	51.422	19.446	1.00	0.00
ATOM	1294	CZ	ARG	1	139	-14.070	50.451	19.933	1.00	0.00
ATOM	1295	NH1	ARG	1	139	-15.376	50.557	20.176	1.00	0.00
ATOM	1296	NH2	ARG	1	139	-13.546	49.286	20.238	1.00	0.00
ATOM	1297	H	ARG	1	139	-11.953	56.599	21.701	1.00	0.00
ATOM	1298	HE	ARG	1	139	-12.325	51.217	19.213	1.00	0.00
ATOM	1299	1HH1	ARG	1	139	-15.852	49.744	20.511	1.00	0.00
ATOM	1300	2HH1	ARG	1	139	-15.778	51.473	20.172	1.00	0.00
ATOM	1301	1HH2	ARG	1	139	-14.068	48.524	20.623	1.00	0.00
ATOM	1302	2HH2	ARG	1	139	-12.550	49.203	20.209	1.00	0.00
ATOM	1303	N	GLU	1	140	-10.248	55.319	18.431	1.00	0.00
ATOM	1304	CA	GLU	1	140	-9.561	55.842	17.254	1.00	0.00
ATOM	1305	C	GLU	1	140	-8.281	54.950	17.238	1.00	0.00

FIG. 1CC


ATOM	1306	O	GLU	1	140	-8.151	54.134	16.315	1.00	0.00
ATOM	1307	CB	GLU	1	140	-9.381	57.331	17.443	1.00	0.00
ATOM	1308	CG	GLU	1	140	-9.055	58.269	16.321	1.00	0.00
ATOM	1309	CD	GLU	1	140	-7.581	57.782	15.971	1.00	0.00
ATOM	1310	OE1	GLU	1	140	-6.649	57.991	16.665	1.00	0.00
ATOM	1311	OE2	GLU	1	140	-7.440	57.213	14.940	1.00	0.00
ATOM	1312	H	GLU	1	140	-9.824	54.778	19.157	1.00	0.00
ATOM	1313	N	TRP	1	141	-7.528	55.060	18.298	1.00	0.00
ATOM	1314	CA	TRP	1	141	-6.370	54.173	18.571	1.00	0.00
ATOM	1315	C	TRP	1	141	-6.737	52.782	18.944	1.00	0.00
ATOM	1316	O	TRP	1	141	-5.982	51.968	19.452	1.00	0.00
ATOM	1317	CB	TRP	1	141	-5.483	54.663	19.722	1.00	0.00
ATOM	1318	CG	TRP	1	141	-4.774	55.962	19.326	1.00	0.00
ATOM	1319	CD1	TRP	1	141	-3.685	56.131	18.409	1.00	0.00
ATOM	1320	CD2	TRP	1	141	-5.095	57.289	19.700	1.00	0.00
ATOM	1321	NE1	TRP	1	141	-3.327	57.468	18.247	1.00	0.00
ATOM	1322	CE2	TRP	1	141	-4.213	58.240	19.050	1.00	0.00
ATOM	1323	CE3	TRP	1	141	-6.038	57.792	20.591	1.00	0.00
ATOM	1324	CZ2	TRP	1	141	-4.413	59.642	19.300	1.00	0.00
ATOM	1325	CZ3	TRP	1	141	-6.247	59.169	20.882	1.00	0.00
ATOM	1326	CH2	TRP	1	141	-5.415	60.068	20.154	1.00	0.00
ATOM	1327	H	TRP	1	141	-7.623	55.892	18.845	1.00	0.00
ATOM	1328	1HE	TRP	1	141	-2.570	57.837	17.746	1.00	0.00
ATOM	1329	N	THR	1	142	-7.945	52.388	18.669	1.00	0.00
ATOM	1330	CA	THR	1	142	-8.470	51.003	18.796	1.00	0.00
ATOM	1331	C	THR	1	142	-8.785	50.384	17.443	1.00	0.00
ATOM	1332	O	THR	1	142	-8.951	49.195	17.303	1.00	0.00
ATOM	1333	CB	THR	1	142	-9.836	51.112	19.635	1.00	0.00
ATOM	1334	OG1	THR	1	142	-10.085	52.512	20.129	1.00	0.00
ATOM	1335	CG2	THR	1	142	-9.646	50.189	20.902	1.00	0.00
ATOM	1336	H	THR	1	142	-8.557	53.070	18.268	1.00	0.00
ATOM	1337	1HG	THR	1	142	-9.297	52.761	20.593	1.00	0.00
ATOM	1338	N	GLN	1	143	-8.798	51.237	16.420	1.00	0.00
ATOM	1339	CA	GLN	1	143	-8.800	50.759	15.056	1.00	0.00
ATOM	1340	C	GLN	1	143	-7.468	50.990	14.393	1.00	0.00
ATOM	1341	O	GLN	1	143	-6.836	49.990	14.069	1.00	0.00
ATOM	1342	CB	GLN	1	143	-9.874	51.493	14.234	1.00	0.00
ATOM	1343	CG	GLN	1	143	-10.095	50.582	13.074	1.00	0.00
ATOM	1344	CD	GLN	1	143	-10.628	51.426	11.972	1.00	0.00
ATOM	1345	OE1	GLN	1	143	-11.716	51.960	12.091	1.00	0.00
ATOM	1346	NE2	GLN	1	143	-9.832	51.593	10.953	1.00	0.00
ATOM	1347	H	GLN	1	143	-8.441	52.158	16.577	1.00	0.00
ATOM	1348	1HE2	GLN	1	143	-8.858	51.392	10.856	1.00	0.00
ATOM	1349	2HE2	GLN	1	143	-10.262	52.137	10.232	1.00	0.00
ATOM	1350	N	LYS	1	144	-7.185	52.353	14.272	1.00	0.00

FIG. 1DD

ATOM	1351	CA	LYS	1	144	-5.872	52.774	13.923	1.00	0.00
ATOM	1352	C	LYS	1	144	-4.944	52.505	15.064	1.00	0.00
ATOM	1353	O	LYS	1	144	-4.948	53.182	16.090	1.00	0.00
ATOM	1354	CB	LYS	1	144	-6.052	54.219	13.415	1.00	0.00
ATOM	1355	CG	LYS	1	144	-4.756	54.831	12.839	1.00	0.00
ATOM	1356	CD	LYS	1	144	-4.529	56.258	13.351	1.00	0.00
ATOM	1357	CE	LYS	1	144	-4.289	56.135	14.821	1.00	0.00
ATOM	1358	NZ	LYS	1	144	-4.298	57.465	15.486	1.00	0.00
ATOM	1359	H	LYS	1	144	-7.735	53.032	14.760	1.00	0.00
ATOM	1360	1HZ	LYS	1	144	-5.272	57.787	15.657	1.00	0.00
ATOM	1361	2HZ	LYS	1	144	-3.709	57.497	16.342	1.00	0.00
ATOM	1362	3HZ	LYS	1	144	-3.674	58.052	14.895	1.00	0.00
ATOM	1363	N	TYR	1	145	-4.035	51.523	14.725	1.00	0.00
ATOM	1364	CA	TYR	1	145	-2.783	51.563	15.405	1.00	0.00
ATOM	1365	C	TYR	1	145	-1.705	51.911	14.373	1.00	0.00
ATOM	1366	O	TYR	1	145	-1.262	51.077	13.622	1.00	0.00
ATOM	1367	CB	TYR	1	145	-2.651	50.166	16.073	1.00	0.00
ATOM	1368	CG	TYR	1	145	-1.338	50.008	16.760	1.00	0.00
ATOM	1369	CD1	TYR	1	145	-1.119	50.509	18.043	1.00	0.00
ATOM	1370	CD2	TYR	1	145	-0.302	49.394	16.035	1.00	0.00
ATOM	1371	CE1	TYR	1	145	0.150	50.465	18.579	1.00	0.00
ATOM	1372	CE2	TYR	1	145	0.972	49.409	16.589	1.00	0.00
ATOM	1373	CZ	TYR	1	145	1.210	49.922	17.847	1.00	0.00
ATOM	1374	OH	TYR	1	145	2.486	49.934	18.410	1.00	0.00
ATOM	1375	H	TYR	1	145	-4.122	50.802	14.038	1.00	0.00
ATOM	1376	HH	TYR	1	145	2.907	50.776	18.294	1.00	0.00
ATOM	1377	N	ALA	1	146	-1.219	53.200	14.453	1.00	0.00
ATOM	1378	CA	ALA	1	146	-0.018	53.540	13.627	1.00	0.00
ATOM	1379	C	ALA	1	146	0.585	54.832	14.174	1.00	0.00
ATOM	1380	O	ALA	1	146	1.508	54.792	14.958	1.00	0.00
ATOM	1381	CB	ALA	1	146	-0.347	53.708	12.163	1.00	0.00
ATOM	1382	H	ALA	1	146	-1.694	53.960	14.896	1.00	0.00
ATOM	1383	N	MET	1	147	0.146	55.965	13.712	1.00	0.00
ATOM	1384	CA	MET	1	147	0.072	57.199	14.504	1.00	0.00
ATOM	1385	C	MET	1	147	-1.165	57.157	15.439	1.00	0.00
ATOM	1386	O	MET	1	147	-1.746	58.134	15.774	1.00	0.00
ATOM	1387	CB	MET	1	147	0.219	58.444	13.697	1.00	0.00
ATOM	1388	CG	MET	1	147	-0.769	58.439	12.499	1.00	0.00
ATOM	1389	SD	MET	1	147	-0.431	59.820	11.346	1.00	0.00
ATOM	1390	CE	MET	1	147	-2.099	59.928	10.616	1.00	0.00
ATOM	1391	OXT	MET	1	147	-1.521	56.064	15.926	1.00	0.00
ATOM	1392	H	MET	1	147	-0.461	55.994	12.919	1.00	0.00

END

FIG. 1EE



ATOM	1	N	CYS *1	85	12.813	-15.194	42.441	1.00	0.00
ATOM	2	H	CYS *1	85	13.027	-15.873	41.738	1.00	0.00
ATOM	3	CA	CYS *1	85	13.822	-14.593	43.296	1.00	0.00
ATOM	4	CB	CYS *1	85	14.815	-15.616	43.734	1.00	0.00
ATOM	5	SG	CYS *1	85	15.117	-15.447	45.437	1.00	0.00
ATOM	6	C	CYS *1	85	14.478	-13.341	42.778	1.00	0.00
ATOM	7	O	CYS *1	85	13.994	-12.195	43.023	1.00	0.00
ATOM	8	N	LEU 1	86	15.596	-13.553	42.021	1.00	0.00
ATOM	9	CA	LEU 1	86	16.482	-12.490	41.469	1.00	0.00
ATOM	10	C	LEU 1	86	17.155	-11.557	42.453	1.00	0.00
ATOM	11	O	LEU 1	86	16.664	-11.390	43.514	1.00	0.00
ATOM	12	CB	LEU 1	86	15.850	-11.805	40.281	1.00	0.00
ATOM	13	CG	LEU 1	86	14.624	-12.437	39.580	1.00	0.00
ATOM	14	CD1	LEU 1	86	14.040	-11.515	38.482	1.00	0.00
ATOM	15	CD2	LEU 1	86	14.916	-13.795	38.925	1.00	0.00
ATOM	16	H	LEU 1	86	15.932	-14.493	41.961	1.00	0.00
ATOM	17	N	ASP 1	87	18.301	-10.942	42.020	1.00	0.00
ATOM	18	CA	ASP 1	87	18.887	-9.697	42.577	1.00	0.00
ATOM	19	C	ASP 1	87	17.933	-8.538	42.215	1.00	0.00
ATOM	20	O	ASP 1	87	17.603	-7.688	42.977	1.00	0.00
ATOM	21	CB	ASP 1	87	20.280	-9.473	41.980	1.00	0.00
ATOM	22	CG	ASP 1	87	20.280	-9.437	40.434	1.00	0.00
ATOM	23	OD1	ASP 1	87	21.191	-8.936	39.856	1.00	0.00
ATOM	24	OD2	ASP 1	87	19.409	-9.977	39.792	1.00	0.00
ATOM	25	H	ASP 1	87	18.468	-11.284	41.095	1.00	0.00
ATOM	26	N	ILE 1	88	17.532	-8.458	40.922	1.00	0.00
ATOM	27	CA	ILE 1	88	16.874	-7.254	40.484	1.00	0.00
ATOM	28	C	ILE 1	88	15.486	-7.061	40.987	1.00	0.00
ATOM	29	O	ILE 1	88	14.834	-5.995	40.819	1.00	0.00
ATOM	30	CB	ILE 1	88	16.911	-7.193	38.904	1.00	0.00
ATOM	31	CG1	ILE 1	88	16.078	-8.320	38.291	1.00	0.00
ATOM	32	CG2	ILE 1	88	18.322	-7.055	38.472	1.00	0.00
ATOM	33	CD	ILE 1	88	14.676	-8.014	37.765	1.00	0.00
ATOM	34	H	ILE 1	88	17.739	-9.251	40.349	1.00	0.00
ATOM	35	N	ARG 1	90	14.426	-7.619	44.251	1.00	0.00
ATOM	36	CA	ARG 1	90	14.634	-7.627	45.688	1.00	0.00
ATOM	37	C	ARG 1	90	15.897	-6.845	46.216	1.00	0.00
ATOM	38	O	ARG 1	90	15.818	-5.995	47.079	1.00	0.00
ATOM	39	CB	ARG 1	90	14.624	-9.055	46.305	1.00	0.00
ATOM	40	CG	ARG 1	90	15.765	-10.003	46.039	1.00	0.00
ATOM	41	CD	ARG 1	90	15.221	-11.397	46.218	1.00	0.00
ATOM	42	NE	ARG 1	90	14.864	-11.584	47.622	1.00	0.00
ATOM	43	CZ	ARG 1	90	13.976	-12.617	47.838	1.00	0.00
ATOM	44	NH1	ARG 1	90	13.824	-13.004	49.063	1.00	0.00
ATOM	45	NH2	ARG 1	90	13.426	-13.227	46.779	1.00	0.00

FIG. 3A

ATOM	46	H	ARG	1	90	14.907	-6.934	43.703	1.00	0.00
ATOM	47	HE	ARG	1	90	15.187	-11.015	48.378	1.00	0.00
ATOM	48	1HH1	ARG	1	90	13.264	-13.807	49.262	1.00	0.00
ATOM	49	2HH1	ARG	1	90	14.153	-12.435	49.817	1.00	0.00
ATOM	50	2HH1	ARG	1	90	12.658	-13.824	47.013	1.00	0.00
ATOM	51	2HH2	ARG	1	90	13.676	-13.086	45.821	1.00	0.00
ATOM	52	N	SER	1	91	17.097	-7.261	45.695	1.00	0.00
ATOM	53	CA	SER	1	91	18.257	-6.643	46.299	1.00	0.00
ATOM	54	C	SER	1	91	19.180	-5.852	45.357	1.00	0.00
ATOM	55	O	SER	1	91	20.364	-5.736	45.486	1.00	0.00
ATOM	56	CB	SER	1	91	19.094	-7.651	47.083	1.00	0.00
ATOM	57	OG	SER	1	91	18.260	-8.522	47.886	1.00	0.00
ATOM	58	H	SER	1	91	17.222	-7.822	44.876	1.00	0.00
ATOM	59	HG	SER	1	91	17.859	-7.970	48.544	1.00	0.00
ATOM	60	N	LEU	1	109	18.852	-18.359	38.255	1.00	0.00
ATOM	61	CA	LEU	1	109	18.365	-18.942	39.568	1.00	0.00
ATOM	62	C	LEU	1	109	17.179	-19.841	39.495	1.00	0.00
ATOM	63	O	LEU	1	109	16.540	-20.333	40.413	1.00	0.00
ATOM	64	CB	LEU	1	109	18.387	-17.901	40.725	1.00	0.00
ATOM	65	CG	LEU	1	109	19.746	-17.216	40.897	1.00	0.00
ATOM	66	CD1	LEU	1	109	19.834	-15.819	40.391	1.00	0.00
ATOM	67	CD2	LEU	1	109	20.454	-17.598	42.195	1.00	0.00
ATOM	68	H	LEU	1	109	18.274	-17.735	37.728	1.00	0.00
ATOM	69	N	ASN	1	114	17.767	-21.723	43.157	1.00	0.00
ATOM	70	CA	ASN	1	114	17.585	-21.498	44.611	1.00	0.00
ATOM	71	C	ASN	1	114	16.288	-22.097	45.204	1.00	0.00
ATOM	72	O	ASN	1	114	15.110	-21.738	44.968	1.00	0.00
ATOM	73	CB	ASN	1	114	17.738	-20.004	44.811	1.00	0.00
ATOM	74	CG	ASN	1	114	17.761	-19.604	46.263	1.00	0.00
ATOM	75	OD1	ASN	1	114	17.013	-18.825	46.820	1.00	0.00
ATOM	76	ND2	ASN	1	114	18.754	-20.204	46.933	1.00	0.00
ATOM	77	H	ASN	1	114	17.244	-21.387	42.373	1.00	0.00
ATOM	78	1HD2	ASN	1	114	19.381	-20.849	46.496	1.00	0.00
ATOM	79	2HD2	ASN	1	114	19.079	-20.336	47.869	1.00	0.00
ATOM	80	N	ASP	1	116	16.009	-21.817	48.288	1.00	0.00
ATOM	81	CA	ASP	1	116	16.037	-21.119	49.574	1.00	0.00
ATOM	82	C	ASP	1	116	15.396	-19.793	49.630	1.00	0.00
ATOM	83	O	ASP	1	116	15.774	-18.837	50.337	1.00	0.00
ATOM	84	CB	ASP	1	116	17.464	-20.968	50.072	1.00	0.00
ATOM	85	CG	ASP	1	116	18.319	-22.224	49.976	1.00	0.00
ATOM	86	OD1	ASP	1	116	18.006	-23.130	50.730	1.00	0.00
ATOM	87	OD2	ASP	1	116	19.236	-22.263	49.180	1.00	0.00
ATOM	88	H	ASP	1	116	16.323	-21.489	47.396	1.00	0.00
ATOM	89	N	ASP	1	117	14.372	-19.664	48.769	1.00	0.00
ATOM	90	CA	ASP	1	117	13.409	-18.653	49.009	1.00	0.00

FIG. 3B

ATOM	91	C	ASP	1	117	12.024	-19.105	48.641	1.00	0.00
ATOM	92	O	ASP	1	117	11.645	-19.083	47.471	1.00	0.00
ATOM	93	CB	ASP	1	117	13.801	-17.459	48.181	1.00	0.00
ATOM	94	CG	ASP	1	117	13.342	-16.211	48.889	1.00	0.00
ATOM	95	OD1	ASP	1	117	13.889	-15.978	49.966	1.00	0.00
ATOM	96	OD2	ASP	1	117	12.577	-15.397	48.399	1.00	0.00
ATOM	97	H	ASP	1	117	14.092	-20.522	48.339	1.00	0.00
END										

FIG. 3C




FIG. 4

Human Met Ala Leu Lys Arg Ile His Lys Glu Leu Asn Asp Leu Ala Arg Asp Pro Pro Ala Gln Cys Ser Ala Gly Pro Val Gly
S Pombe Met Ala Leu Lys Arg Ile Asn Arg Glu Leu Ala Asp Leu Gly Lys Asp Pro Pro Ser Ser Cys Ser Ala Gly Pro Val Gly
C Albicans Met Ser Leu Lys Arg Ile Asn Lys Glu Leu Ser Asp Leu Gly Arg Asp Pro Pro Ser Ser Cys Ser Ala Gly Pro Val Gly

Asp Asp Met Phe His Trp Gln Ala Thr Ile Met Gly Pro Asn Asp Ser Pro Tyr Gln Gly Val Phe Phe Leu Thr Ile
Asp Asp Leu Phe His Trp Gln Ala Thr Ile Met Gly Pro Ala Asp Ser Pro Tyr Ala Gly Val Phe Leu Leu Ser Ile
Asp Asp Leu Tyr His Trp Gln Ala Ser Ile Met Gly Pro Pro Asp Ser Pro Tyr Ala Gly Val Phe Phe Leu Ser Ile

His Phe Pro Thr Asp Tyr Pro Phe Lys Pro Pro Lys Val Ala Phe Thr Thr Arg Ile Tyr His Pro Asn Ile Asn Ser Asn
His Phe Pro Thr Asp Tyr Pro Phe Lys Pro Pro Lys Val Asn Phe Thr Thr Arg Ile Tyr His Pro Asn Ile Asn Ser Asn
His Phe Pro Thr Asp Tyr Pro Leu Lys Pro Pro Lys Ile Ala Leu Thr Thr Lys Ile Tyr His Pro Asn Ile Asn Ser Asn

Gly Ser Ile Cys Leu Asp Ile Leu Arg Ser Gln Trp Ser Pro Ala Leu Thr Thr Ile Ser Lys Val Leu Ser Ile Cys Ser
Gly Ser Ile Cys Leu Asp Ile Leu Arg Asp Gln Trp Ser Pro Ala Leu Thr Thr Ile Ser Lys Val Leu Ser Ile Cys Ser
Gly Asn Ile Cys Leu Asp Ile Leu Lys Asp Gln Trp Ser Pro Ala Leu Thr Thr Ile Ser Lys Val Leu Ser Ile Cys Ser

Leu Leu Cys Asp Pro Asn Pro Asp Asp Pro Leu Val Pro Glu Ile Ala Arg Ile Tyr Gln Thr Thr Asp Arg Glu Lys Tyr Asn
Leu Leu Thr Asp Pro Asn Pro Asp Asp Pro Leu Val Pro Glu Ile Ala His Val Tyr Lys Thr Asp Arg Ser Arg Tyr Glu
Leu Leu Thr Asp Ala Asn Pro Asp Asp Pro Leu Val Pro Glu Ile Ala His Ile Tyr Lys Gln Asp Arg Lys Lys Tyr Glu

Arg Ile Ala ArgGlu Trp Thr Gln Lys Tyr Ala Met
Leu Ser Ala ArgGlu Trp Thr Arg Lys Tyr Ala Ile
Ala Thr Ala LysGlu Trp Thr Lys Lys Tyr Ala Val

FIG. 5




FIG. 6




FIG. 7A




FIG. 7B




FIG. 8

UBIQUITIN CONJUGATING ENZYMES

RELATED APPLICATIONS

This application is a continuation-in-part of U.S. Ser. No. 08/247,904, entitled "Human Ubiquitin Conjugating Enzyme", which is a continuation-in-part of U.S. Ser. No. 08/176,937 filed Jan. 4, 1994, now abandoned, entitled "Assay and Reagents for Detecting Inhibitors of Ubiquitin-dependent Degradation of Cell Cycle Regulatory Proteins", the specification of which are incorporated by reference herein.

BACKGROUND OF THE INVENTION

The ubiquitin-mediated proteolysis system is the major pathway for the selective, controlled degradation of intracellular proteins in eukaryotic cells. Ubiquitin modification of a variety of protein targets within the cell appears to be important in a number of basic cellular functions such as regulation of gene expression, regulation of the cell-cycle, modification of cell surface receptors, biogenesis of ribosomes, and DNA repair. One major function of the ubiquitin-mediated system is to control the half-lives of cellular proteins. The half-life of different proteins can range from a few minutes to several days, and can vary considerably depending on the cell-type, nutritional and environmental conditions, as well as the stage of the cell-cycle.

Targeted proteins undergoing selective degradation, presumably through the actions of a ubiquitin-dependent proteasome, are covalently tagged with ubiquitin through the formation of an isopeptide bond between the C-terminal glycyl residue of ubiquitin and a specific lysyl residue in the substrate protein. This process is catalyzed by a ubiquitin-activating enzyme (E1) and a ubiquitin-conjugating enzyme (E2), and in some instances may also require auxiliary substrate recognition proteins (E3s). Following the linkage of the first ubiquitin chain, additional molecules of ubiquitin may be attached to lysine side chains of the previously conjugated moiety to form branched multi-ubiquitin chains.

The conjugation of ubiquitin to protein substrates is a multi-step process. In an initial ATP requiring step, a thioester is formed between the C-terminus of ubiquitin and an internal cysteine residue of an E1 enzyme. Activated ubiquitin is then transferred to a specific cysteine on one of several E2 enzymes. Finally, these E2 enzymes donate ubiquitin to protein substrates. Substrates are recognized either directly by ubiquitin-conjugated enzymes or by associated substrate recognition proteins, the E3 proteins, also known as ubiquitin ligases.

Ubiquitin is itself a substrate for ubiquitination. Depending on the ubiquitin-conjugating enzyme and the nature of the substrate, specific lysine residues of ubiquitin are used as acceptor sites for further ubiquitinations. This can lead to either a linear multi-ubiquitin chain (when a single lysine residue of ubiquitin is used) or multi-ubiquitin "trees" (when more than one lysine residue of ubiquitin is used). Although the attachment of a single ubiquitin moiety to a substrate can be sufficient for degradation, multi-ubiquitination appears to be required in most cases.

Many proteins that control cell-cycle progression are short-lived. For example, regulation of oncoproteins and anti-oncoproteins clearly plays an important role in determining steady-state levels of protein expression, and alterations in protein degradation are as likely as changes in transcription and/or translation to cause either the proliferative arrest of cells, or alternatively, the transformation of cells.

For instance, the p53 protein is a key regulator of mammalian cell growth and its gene is frequently mutated in a wide range of human tumors (Hollstein et al. (1991) *Science* 253:49-53). Furthermore, many DNA tumor viruses encode viral antigens that inactivate p53 (e.g., see Vogelstein et al. (1992) *Cell* 70:523-526). The high risk human papillomaviruses, such as HPV-16 and -18, are strongly implicated in the pathogenesis of cervical carcinoma (zur Hansen et al. (1991) *Science* 254:1167-1173). These viruses encode two transforming proteins, E6 and E7, that target the cellular growth regulators p53 and pRb respectively. The mode of inactivation of p53 by E6 is apparently mediated by a ubiquitin-dependent pathway. Viral E6 and a cellular E6-associated protein (E6AP) combine to stimulate the ubiquitination of p53, thus targeting p53 for degradation (Scheffner et al. (1990) *Cell* 63:1129-1136. In this reaction, E6 and E6AP are thought to be providing a ubiquitin ligase, or E3-like activity (Scheffner et al. (1993) *Cell* 75:495-505). However, the ubiquitin-conjugating enzyme (E2) involved in p53 ubiquitination has not previously been characterized.

SUMMARY OF THE INVENTION

The present invention relates to the discovery in eukaryotic cells, particularly human cells and certain yeast cells, of a novel ubiquitin conjugating enzyme (hereinafter "UbCE"). In human cells, the enzyme can function to mediate ubiquitination of cell check regulatory proteins, e.g. p53, and is therefore involved in regulating cell cycle progression, e.g. cell growth.

One aspect of the invention features a substantially pure preparation of an a human UbCE polypeptide ("hUbCE"), or a fragment thereof, which can function as a ubiquitin conjugating enzyme. In a preferred embodiment: the polypeptide has an amino acid sequence at least 90% homologous to the amino acid sequence of SEQ ID No. 2; the polypeptide has an amino acid sequence at least 95% homologous to the amino acid sequence of SEQ ID No. 2; the polypeptide has an amino acid sequence at least 97% homologous to the amino acid sequence of SEQ ID No. 2; the polypeptide has an amino acid sequence identical to the amino acid sequence of SEQ ID No. 2. In a preferred embodiment: the fragment comprises at least 5 contiguous amino acid residues of SEQ ID No. 2; the fragment comprises at least 20 contiguous amino acid residues of SEQ ID No. 2; the fragment comprises at least 50 contiguous amino acid residues of SEQ ID No. 2. In a preferred embodiment, the fragment comprises at least a portion of amino acid residues Cys-107 through Met-147, e.g. 5 amino acid residues, e.g. 15 amino acid residues, e.g. 25 amino acid residues.

Another aspect of the invention features a substantially pure preparation of a *Candida* UbCE polypeptide ("caUbCE"), or a fragment thereof, which can function as a ubiquitin conjugating enzyme. In a preferred embodiment: the polypeptide has an amino acid sequence at least 90% homologous to the amino acid sequence of SEQ ID No. 4; the polypeptide has an amino acid sequence at least 95% homologous to the amino acid sequence of SEQ ID No. 4; the polypeptide has an amino acid sequence at least 97% homologous to the amino acid sequence of SEQ ID No. 4; the polypeptide has an amino acid sequence identical to the amino acid sequence of SEQ ID No. 4. In a preferred embodiment: the fragment comprises at least 5 contiguous amino acid residues of SEQ ID No. 4; the fragment comprises at least 20 contiguous amino acid residues of SEQ ID No. 4; the fragment comprises at least 50 contiguous amino acid residues of SEQ ID No. 4. In a preferred embodiment, the fragment comprises at least a portion of amino acid

residues Cys-107 through Val-147, e.g. 5 amino acid residues, e.g. 15 amino acid residues, e.g. 25 amino acid residues.

Another aspect of the invention features a substantially pure preparation of a *Schizosaccharomyces* UbCE polypeptide ("spUbCE"), or a fragment thereof, which can function as a ubiquitin conjugating enzyme. In a preferred embodiment: the polypeptide has an amino acid sequence at least 90% homologous to the amino acid sequence of SEQ ID No. 6; the polypeptide has an amino acid sequence at least 95% homologous to the amino acid sequence of SEQ ID No. 6; the polypeptide has an amino acid sequence at least 97% homologous to the amino acid sequence of SEQ ID No. 6; the polypeptide has an amino acid sequence identical to the amino acid sequence of SEQ ID No. 6. In a preferred embodiment: the fragment comprises at least 5 contiguous amino acid residues of SEQ ID No. 6; the fragment comprises at least 20 contiguous amino acid residues of SEQ ID No. 6; the fragment comprises at least 50 contiguous amino acid residues of SEQ ID No. 6. In a preferred embodiment, the fragment comprises at least a portion of amino acid residues Cys-107 through Ile-147, e.g. 5 amino acid residues, e.g. 15 amino acid residues, e.g. 25 amino acid residues.

Another aspect of the present invention features an hUbCE polypeptide which functions in one of either role of an agonist of cell-cycle regulation or an antagonist of cell-cycle regulation. In a preferred embodiment the hUbCE polypeptide has: an ability to mediate ubiquitination of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; an ability to mediate ubiquitin-dependent degradation of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; an ability to affect the cellular half-life of a cell-cycle regulatory protein, e.g. a cell-cycle checkpoint protein, e.g. p53, e.g. in normal cells, e.g. in normal proliferating cells, e.g. in virally-infected cells, e.g. in papillomavirus infected cells, e.g. in HPV-infected cells, e.g. in HPV-16, HPV-18, HPV-31, or HPV-33 infected cells, e.g. in cells expressing a papillomavirus E6 protein, e.g. in transformed cells, e.g. in cancerous cells. The biological activity can further include the ability to bind and conjugate ubiquitin, as well as bind and transfer ubiquitin to E6AP.

Yet another aspect of the present invention concerns an immunogen comprising an UbCE polypeptide, or a fragment thereof in an immunogenic preparation, the immunogen being capable of eliciting an immune response specific for the UbCE polypeptide; e.g. a humoral response, eg. an antibody response., e.g. a cellular response.

A still further aspect of the present invention features an antibody preparation specifically reactive with an epitope of the UbCE immunogen, e.g. reactive with hUbCE, e.g. reactive with caUbC, e.g. reactive with spUbCE.

Another aspect of the present invention features recombinant hUbCE polypeptide, or a fragment thereof, having an amino acid sequence preferably: at least 90% homologous to SEQ ID No. 2; at least 95% homologous to SEQ ID No. 2; at least 97% homologous to SEQ ID No. 2. In a preferred embodiment, the recombinant hUbCE protein functions in one of either role of an agonist of cell cycle regulation or an antagonist of cell cycle regulation. In a more preferred embodiment: the hUbCE polypeptide mediates ubiquitination of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; the hUbCE polypeptide mediates ubiquitin-dependent degradation of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; the hUbCE polypeptide affects the cellular half-life of a cell-cycle regulatory protein, e.g. a

cell-cycle checkpoint protein, e.g. p53, e.g. in normal cells, e.g. in normal proliferating cells, e.g. in virally-infected cells, e.g. in papillomavirus infected cells, e.g. in HPV-infected cells, e.g. in HPV-16, HPV-18, HPV-31 or HPV-33 infected cells, e.g. in cells expressing a papillomavirus E6 protein, e.g. in transformed cells, e.g. in cancerous cells.

Another aspect of the present invention features recombinant caUbCE polypeptide, or a fragment thereof, having an amino acid sequence preferably: at least 90% homologous to SEQ ID No. 4; at least 95% homologous to SEQ ID No. 4; at least 97% homologous to SEQ ID No. 4. In a preferred embodiment, the recombinant caUbCE protein functions in one of either role of an agonist of cell cycle regulation or an antagonist of cell cycle regulation. In a more preferred embodiment the caUbCE polypeptide mediates ubiquitination of cellular proteins of candida cells.

Another aspect of the present invention features recombinant spUbCE polypeptide, or a fragment thereof, having an amino acid sequence preferably: at least 90% homologous to SEQ ID No. 6; at least 95% homologous to SEQ ID No. 6; at least 97% homologous to SEQ ID No. 6. In a preferred embodiment, the recombinant spUbCE protein functions in one of either role of an agonist of cell cycle regulation or an antagonist of cell cycle regulation. In a more preferred embodiment the spUbCE polypeptide mediates ubiquitination of cellular proteins of *Schizosaccharomyces* cells.

In yet other preferred embodiments, the recombinant UbCE protein is a fusion protein further comprising a second polypeptide portion having an amino acid sequence from a protein unrelated the protein of SEQ ID No. 2, 4 or 6. Such fusion proteins can be functional in a two-hybrid assay.

Another aspect of the present invention provides a substantially pure nucleic acid having a nucleotide sequence which encodes an hUbCE polypeptide, or a fragment thereof, having an amino acid sequence at least 90% homologous to SEQ ID NO. 2. In a more preferred embodiment, the nucleic acid encodes a protein having an amino acid sequence at least 95% homologous to SEQ ID No. 2; and more preferably at least 97% homologous to SEQ ID No. 2. The nucleic acid preferably encodes: a hUbCE polypeptide which mediates ubiquitination of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; a hUbCE polypeptide which mediates ubiquitin-dependent degradation of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; a hUbCE polypeptide which affects the cellular half-life of a cell-cycle regulatory protein, e.g. a cell-cycle checkpoint protein, e.g. p53, e.g. in normal cells, e.g. in normal proliferating cells, e.g. in virally-infected cells, e.g. in papillomavirus infected cells, e.g. in HPV-infected cells, e.g. in HPV-16, HPV-18, HPV-31, or HPV-33 infected cells, e.g. in cells expressing a papillomavirus E6 protein, e.g. in transformed cells, e.g. in cancerous cells.

Another aspect of the present invention provides a substantially pure nucleic acid having a nucleotide sequence which encodes a caUbCE polypeptide, or a fragment thereof, having an amino acid sequence at least 90% homologous to SEQ ID NO. 4. In a more preferred embodiment, the nucleic acid encodes a protein having an amino acid sequence at least 95% homologous to SEQ ID No. 4; and more preferably at least 97% homologous to SEQ ID No. 4.

Another aspect of the present invention provides a substantially pure nucleic acid having a nucleotide sequence which encodes an spUbCE polypeptide, or a fragment

5

thereof, having an amino acid sequence at least 90% homologous to SEQ ID NO. 4. In a more preferred embodiment, the nucleic acid encodes a protein having an amino acid sequence at least 95% homologous to SEQ ID No. 4; and more preferably at least 97% homologous to SEQ ID No. 4.

In yet a further preferred embodiment, the nucleic acid which encodes a UbCE polypeptide of the present invention, or a fragment thereof, hybridizes under stringent conditions to a nucleic acid probe corresponding to at least 12 consecutive nucleotides of one of SEQ ID Nos. 1, 3 or 5; more preferably to at least 20 consecutive nucleotides of said sequences; more preferably to at least 40 consecutive nucleotides. In yet a further preferred embodiment, the UbCE encoding nucleic acid hybridizes to a nucleic acid probe corresponding to a subsequence encoding at least 4 consecutive amino acids between residues 107 and 147 of SEQ ID No. 2, 4 or 6, more preferably at least 10 consecutive amino acid residues, and even more preferably at least 20 amino acid residues. In yet a preferred embodiment the nucleic acid encodes an hUbCE polypeptide which includes Cys-107 through Cys-111.

Furthermore, in certain preferred embodiments, UbCE encoding nucleic acid will comprise a transcriptional regulatory sequence, e.g. at least one of a transcriptional promoter or transcriptional enhancer sequence, operably linked to the UbCE gene sequence so as to render the UbCE gene sequence suitable for use as an expression vector. In one embodiment, the UbCE gene is provided as a sense construct. In another embodiment, the UbCE gene is provided as an anti-sense construct.

The present invention also features transgenic non-human animals, e.g. mice, which either express a heterologous hUbCE gene, e.g. derived from humans, or which mis-express their own homolog of the subject human gene, e.g. expression of the mouse hUbCE homolog is disrupted. Such a transgenic animal can serve as an animal model for studying cellular disorders comprising mutated or mis-expressed hUbCE alleles.

The present invention also provides a probe/primer comprising a substantially purified oligonucleotide, wherein the oligonucleotide comprises a region of nucleotide sequence which hybridizes under stringent conditions to at least 10 consecutive nucleotides of sense or antisense sequence of SEQ ID No. 1 or naturally occurring mutants thereof. In preferred embodiments, the probe/primer further comprises a label group attached thereto and able to be detected, e.g. the label group is selected from a group consisting of radioisotopes, fluorescent compounds, enzymes, and enzyme co-factors. Such probes can be used as a part of a diagnostic test kit for identifying transformed cells, such as for measuring a level of a hUbCE nucleic acid in a sample of cells isolated from a patient; e.g. measuring the hUbCE mRNA level in a cell; e.g. determining whether the genomic hUbCE gene has been mutated or deleted.

The present invention also provides a method for treating an animal having unwanted cell growth characterized by a loss of wild-type p53 function, comprising administering a therapeutically effective amount of an agent able to inhibit a ubiquitin conjugating activity of the subject hUbCE protein.

The present invention also provides a method for treating an animal having an unwanted mycotic infection, comprising administering a therapeutically effective amount of an agent able to inhibit a ubiquitin conjugating activity of a fungal ubiquitin-conjugating enzyme, such as the subject

6

caUbCE protein or spUBC protein, without substantially inhibiting the hUbCE protein.

Another aspect of the present invention provides a method of determining if a subject, e.g. a human patient, is at risk for a disorder characterized by unwanted cell proliferation, comprising detecting, in a tissue of the subject, the presence or absence of a genetic lesion characterized by at least one of (i) a mutation of a gene encoding a protein represented by SEQ ID No. 2, or a homolog thereof; or (ii) the mis-expression of the hUbCE gene. In preferred embodiments: detecting the genetic lesion comprises ascertaining the existence of at least one of a deletion of one or more nucleotides from the gene, an addition of one or more nucleotides to the gene, an substitution of one or more nucleotides of the gene, a gross chromosomal rearrangement of the gene, a gross alteration in the level of a messenger RNA transcript of the gene, the presence of a non-wild type splicing pattern of a messenger RNA transcript of the gene, or a non-wild type level of the protein. For example, detecting the genetic lesion can comprise (i) providing a probe/primer comprising an oligonucleotide containing a region of nucleotide sequence which hybridizes to a sense or antisense sequence of SEQ ID No. 1 or naturally occurring mutants thereof or 5' or 3' flanking sequences naturally associated with the gene; (ii) exposing the probe/primer to nucleic acid of the tissue; and (iii) detecting, by hybridization of the probe/primer to the nucleic acid, the presence or absence of the genetic lesion; e.g. wherein detecting the lesion comprises utilizing the probe/primer to determine the nucleotide sequence of the hUbCE gene and, optionally, of the flanking nucleic acid sequences; e.g. wherein detecting the lesion comprises utilizing the probe/primer in a polymerase chain reaction (PCR); e.g. wherein detecting the lesion comprises utilizing the probe/primer in a ligation chain reaction (LCR). In alternate embodiments, the level of the protein is detected in an immunoassay.

Yet a further aspect of the present invention concerns three-dimensional molecular models of the subject UbCE proteins, and their use as templates for the design of agents able to inhibit at least one biological activity of the ubiquitin conjugating enzyme. In preferred embodiments, the molecular models can be used to design pharmacophores by rational drug design; e.g. agents which can inhibit binding of the subject hUbCE protein with any one of ubiquitin, an E1 enzyme, an E3 protein(s) such as E6 or E6AP, or the downstream target of the enzyme, such as p53.

For instance, one aspect of the present invention concerns a method for identifying inhibitors of the subject ubiquitin-conjugating enzyme by molecular modeling. In general, the method comprise providing a molecular model of the enzyme, such as the active site, as well as a molecular model of a candidate drug. The drug model is docked with the UbCE model and binding criteria, e.g. electrostatic interactions, hydrogen bonding, hydrophobic interactions, desolvation effects, cooperative motions of ligand and enzyme, of the docked models is determined. Based on the binding criteria of a particular candidate drug, the likelihood of the candidate drug being an inhibitor of said UbCE can be determined. Thus, the subject method can be used to design candidate agents, which when obtained, e.g. by chemical synthesis or from commercial sources, can be provided in an assay with the human ubiquitin-conjugating enzyme of the present invention in order to determine the actual inhibitory activity of the candidate drug. In preferred embodiments, the hUbCE model includes the amino acid residues Cys-85, Leu-86, Asp-87, Ile-88, Arg-90, Ser-91, Leu-109, Asn-114, Asp-116, and Asp-117, the atomic coordinates of these

residues, at 300° K, having an overall RMS within 2 Å of the atomic coordinates shown in FIG. 2, more preferably an overall RMS within 1 Å, and most preferably an overall RMS within 0.5 Å. Moreover, the hUbCE model can include amino acid residues Arg-5 through Met-147 of SEQ ID No. 2. In preferred embodiments, the atomic coordinates for the C-α carbon for each of these residues, at 300° K, have an overall RMS within 2 Å of the C-α atomic coordinates shown in FIG. 1, more preferably an overall RMS within 1 Å, and most preferably an overall RMS within 0.5 Å. Moreover, the hUbCE model can include the atomic coordinates for each atom of the amino acid residues Arg-5 through Met-147 of SEQ ID No. 2. In preferred embodiments, the atomic coordinates for each of these residues, at 300° K, have an overall RMS within 2 Å of the C-α atomic coordinates shown in FIG. 1, more preferably an overall RMS within 1 Å, and most preferably an overall RMS within 0.5 Å.

Yet a further aspect of this invention concerns addressable electronic memory means, e.g. RAM or ROM memory, magnetic disk devices, optical storage devices, having stored therein an addressable electronic representation of atomic coordinates of a molecular model of a human ubiquitin-conjugating enzyme. In preferred embodiments, the hUbCE model comprises the amino acid residues Cys-85, Leu-86, Asp-87, Ile-88, Arg-90, Ser-91, Leu-109, Asn-114, Asp-116, and Asp-117, the atomic coordinates of these residues, at 300° K, having an overall RMS within 2 Å of the atomic coordinates shown in FIG. 2. In another embodiment, the human ubiquitin-conjugating enzyme model comprises amino acid residues Arg-5 through Met-147 of SEQ ID No. 2, the atomic coordinates of the residues, at 300° K, having an overall RMS within 2 Å of the atomic coordinates shown in FIG. 1.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

The practice of the present invention will employ, unless otherwise indicated, conventional techniques of cell biology, cell culture, molecular biology, transgenic biology, microbiology, recombinant DNA, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature. See, for example, *Molecular Cloning A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press:1989); *DNA Cloning*, Volumes I and II (D. N. Glover ed., 1985); *Oligonucleotide Synthesis* (M. J. Gait ed., 1984); Mullis et al. U.S. Pat. No. 4,683,195; *Nucleic Acid Hybridization* (B. D. Hames & S. J. Higgins eds. 1984); *Transcription And Translation* (B. D. Hames & S. J. Higgins eds. 1984); *Culture Of Animal Cells* (R. I. Freshney, Alan R. Liss, Inc., 1987); *Immobilized Cells And Enzymes* (IRL Press, 1986); B. Perbal, *A Practical Guide To Molecular Cloning* (1984); the treatise, *Methods In Enzymology* (Academic Press, Inc., N.Y.); *Gene Transfer Vectors For Mammalian Cells* (J. H. Miller and M. P. Calos eds., 1987., Cold Spring Harbor Laboratory); *Methods In Enzymology*, Vols. 154 and 155 (Wu et al. eds.), *Immunochemical Methods In Cell And Molecular Biology* (Mayer and Walker, eds., Academic Press, London, 1987); *Handbook Of Experimental Immunology*, Volumes I-IV (D. M. Weir and C. C. Blackwell, eds., 1986); *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986).

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 is the atomic coordinates for Arg-5 through Met-147 of SEQ ID No. 2 in standard Brookhaven protein databank (pdb) format.

FIG. 2 is a stick figure illustrating the residues of the active site of hUbCE.

FIG. 3 is the atomic coordinates for Cys-85, Leu-86, Asp-87, Ile-88, Arg-90, Ser-91, Leu-109, Asn-114, Asp-116, and Asp-117 of SEQ ID No. 2 in standard Brookhaven protein databank (pdb) format.

FIG. 4 is the schematic overview of the operation of the GROW method of drug design. The site and seed coordinate file and command file are provided to the GROW procedure by the user. Growth can be visualized as a tree process in which each library template is attached to the seed (A) and then evaluated by the scoring function (e.g. binding criteria). Of the resulting constructs, a given number of best constructs (e.g. 10) are kept for the next level (B). To each retained monopeptide/seed construct are attached all library templates, which are again scored (C). After pruning(D), the process is repeated (E) until the specified peptide length is reached (F). In this tree diagram, circles represent those nodes selected (based on best binding criteria evaluation) for further growth. Uncircled nodes are pruned. Horizontal dots denote continuation across all template additions (e.g. other members of a series), and vertical dots represent the iterative process of tree growth.

FIG. 5 is a sequence alignment of hUbCE ("human"), spUbCE ("S pombe") and caUbCE ("C albicans").

FIG. 6 illustrate the hUbCE dependent ubiquitination of p53 in an in vitro ubiquitination reaction. A complete ubiquitination reaction shown in lane 6 contained E1, hUbCE, E6, E6AP, p53 and ubiquitin. The following changes were made in lanes 1-5: lane 1 no E6, lane 2 no E6AP, lane 3 UBC2 replaces hUbCE, lane 4 no E1, lane 5 no ubiquitin. In lane 7 mutant hUbCE (Cys85→Ser) replaces wild-type hUbCE.

FIG. 7A shows the ubiquitination of E6AP. Purified proteins were used in ubiquitination reactions containing biotinylated ubiquitin. Lane 1 ubiquitin, lane 2 E1, ubiquitin and hUbCE, lane 3 E1, ubiquitin, hUbCE and E6AP, lane 4 E1, ubiquitin, hUbCE, E6AP and E6, lane 5 E1, hUbCE, E6AP and E6, lane 6 ubiquitin, hUbCE and E6AP, lane 7 E1, ubiquitin and E6AP, lane 8 ubiquitin and E6AP.

FIG. 7B demonstrates the hUbCE-specific ubiquitination of E6AP. All lanes contained E1 and ubiquitin with the following additions: lane 1 nothing, lane 2 hUbCE, lane 3 hUbCE and E6AP, lane 4 GST.UBC8, lane 5 GST.UBC8 and E6AP, lane 6 GST.UBC2, lane 7 GST.UBC2 and E6AP, lane 8 GST.epiUBC, lane 9 GST.epiUBC and E6AP.

FIG. 8 shows the degree of inhibition of E6 stimulated p53 degradation in co-injection experiments. The indicated DNAs were co-injected with pX.E6. The levels of inhibition of the E6 stimulated p53 degradation are derived from an analysis of approx. 150 injected cells per experimental point in at least two independent experiments.

DETAILED DESCRIPTION OF THE INVENTION

The ubiquitin system is essential for a wide spectrum of cellular phenomena, and is a component of many biological regulatory mechanisms, including aspects of growth control, metabolic regulation, embryonic development, and cell-cycle progression.

The present invention relates to the discovery of a family of related ubiquitin-conjugating enzymes ("UbCE"). In particular, members of this family have been cloned from various eukaryotic sources, and include, for example, a human ubiquitin-conjugating enzyme ("hUbCE"), *C. albi-*

cans ubiquitin-conjugating enzyme ("caUbCE"), and an *S. pombe* ubiquitin-conjugating enzyme ("spUbCE"). The nucleotide sequences for the human UbCE, the *C. albicans* UbCE, and the *S. pombe* UbCE coding sequences are provided in SEQ ID Nos. 1, 3 and 5, respectively. The corresponding amino acid sequences are represented in SEQ ID Nos. 2, 4 and 6.

The biological activity of the UbCE proteins of the present invention is likely to be important in a number of basic cellular functions, such as regulation of gene expression, regulation of the cell-cycle, modification of cell surface receptors, biogenesis of ribosomes, and DNA repair. An apparent function of members of this family of enzymes in ubiquitin-mediated systems is to control the cellular half-lives of various proteins. For instance, as demonstrated in the Examples, hUbCE is implicated in the ubiquitin-mediated inactivation of cell-cycle regulatory proteins, particularly p53. As is generally known, p53 is a checkpoint protein that plays an important role in sensing DNA damage or regulating cellular response to stress. Moreover, lesions in the p53 gene have been shown to be associated with a wide variety of proliferative diseases. Consequently, the present invention identifies a potential molecular target, e.g., hUbCE, for regulating the cellular half-life of p53 and thereby modulating, for instance, cell proliferation, apoptosis and cellular sensitivity to chemotherapeutics and DNA damaging agents.

Accordingly, the present invention makes available diagnostic and therapeutic assays, reagents and kits for detecting and treating proliferative disorders arising from, for example, tumorigenic transformation of cells, or other hyperplastic or neoplastic transformation processes. For example, the present invention makes available reagents, such as antibodies and nucleic acid probes, for detecting altered complex formation, and/or altered levels of hUbCE expression, and/or hUbCE-gene deletion or mutation, in order to identify transformed cells. Moreover, the present invention provides a method of treating a wide variety of pathological cell proliferative conditions, such as by gene therapy utilizing recombinant gene constructs encoding the subject hUbCE proteins, by providing peptidomimetics which either inhibit or potentiate the interaction between hUbCE and E6AP or an E6AP-containing complexes, or by providing inhibitors of the catalytic activity of hUbCE. Such methods can also be used in tissue culture, such as to regulate the transformation of cells in vitro.

In similar fashion, the present invention also makes available diagnostic and therapeutic assays for detecting and treating yeast/fungal infections, where such infections occur in an animal, e.g. humans, or on a non-living object, such as food or medical instruments. For example, given the apparent role of the subject UbCEs, namely caUbCE and spUbCE, in regulation of proteins involved in growth, mating and proliferation of yeast, inhibitors of the subject ubiquitin conjugating enzyme can be used to treat mycotic infections, as disinfectants, or as food preservatives.

For convenience, certain terms employed in the specification, examples, and appended claims are collected here.

As used herein, the term "nucleic acid" refers to polynucleotides such as deoxyribonucleic acid (DNA), and, where appropriate, ribonucleic acid (RNA). The term should also be understood to include, as equivalents, analogs of either RNA or DNA made from nucleotide analogs, and, as applicable to the embodiment being described, single (sense or antisense) and double-stranded polynucleotides.

As used herein, the terms "gene", "recombinant gene" and "gene construct" refer to a nucleic acid comprising an open reading frame encoding a UbCE polypeptide of the present invention, including both exon and (optionally) intron sequences. In preferred embodiments, the nucleic acid is DNA or RNA. Exemplary recombinant genes include nucleic acids which encode all or a catalytically active portion of the hUbCE protein represented in SEQ ID No. 2, the caUbCE protein represented in SEQ ID No. 4, or the spUbCE protein represented in SEQ ID No. 6. The term "intron" refers to a DNA sequence present in a given UbCE-gene which is not translated into protein and is generally found between exons.

The term "transfection" refers to the introduction of a nucleic acid, e.g., an expression vector, into a recipient cell by nucleic acid-mediated gene transfer. "Transformation", as used herein, refers to a process in which a cell's genotype is changed as a result of the cellular uptake of exogenous nucleic acid, and, for example, the transformed cell expresses a recombinant form of one of the subject UbCE proteins.

"Cells" or "cell cultures" or "recombinant host cells" or "host cells" are often used interchangeably as will be clear from the context. These terms include the immediate subject cell which expresses a ubiquitin-conjugating enzyme of the present invention, and, of course, the progeny thereof. It is understood that not all progeny are exactly identical to the parental cell, due to chance mutations or difference in environment. However, such altered progeny are included in these terms, so long as the progeny retain the characteristics relevant to those conferred on the originally transformed cell. In the present case, such a characteristic might be the ability to produce a recombinant UbCE-protein.

As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting another nucleic acid to which it has been linked. The term "expression vector" includes plasmids, cosmids or phages capable of synthesizing the subject proteins encoded by their respective recombinant genes carried by the vector. Preferred vectors are those capable of autonomous replication and/or expression of nucleic acids to which they are linked. In the present specification, "plasmid" and "vector" are used interchangeably as the plasmid is the most commonly used form of vector. Moreover, the invention is intended to include such other forms of expression vectors which serve equivalent functions and which become known in the art subsequently hereto.

"Transcriptional regulatory sequence" is a generic term used throughout the specification to refer to DNA sequences, such as initiation signals, enhancers, and promoters, as well as polyadenylation sites, which induce or control transcription of protein coding sequences with which they are operably linked. In preferred embodiments, transcription of a recombinant UbCE-gene is under the control of a promoter sequence (or other transcriptional regulatory sequence) which controls the expression of the recombinant gene in a cell-type in which expression is intended. It will also be understood that the recombinant gene can be under the control of transcriptional regulatory sequences which are the same or which are different from those sequences which control transcription of the naturally-occurring form of the regulatory protein.

The term "tissue-specific promoters" means a DNA sequence that serves as a promoter, i.e., regulates expression of a selected DNA sequence operably linked to the promoter, and which effects expression of the selected DNA sequence

in specific cells of a tissue, such as cells of an epithelial lineage, e.g. cervical squamous cells. In an illustrative embodiment of epithelial-specific promoters, gene constructs can be used as a part of gene therapy to deliver, for example, genes encoding a dominant negative hUbCE mutant, in order to inhibit degradation of p53 required for the pathogenesis of certain papillomavirus-mediated disorders, e.g. papillomas, or to direct expression of an antisense construct of the subject ubiquitin-conjugating enzyme in only epithelial tissue. The term also covers so-called "leaky" promoters, which regulate expression of a selected DNA primarily in one tissue, but cause expression in other tissues as well.

As used herein, a "transgenic animal" is any animal, preferably a non-human mammal in which one or more of the cells of the animal contain heterologous nucleic acid introduced by way of human intervention, such as by transgenic techniques well known in the art. The nucleic acid is introduced into the cell, directly or indirectly by introduction into a precursor of the cell, by way of deliberate genetic manipulation, such as by microinjection or by infection with a recombinant virus. The term genetic manipulation does not include classical cross-breeding, or in vitro fertilization, but rather is directed to the introduction of a recombinant DNA molecule. This molecule may be integrated within a chromosome, or it may be extrachromosomally replicating DNA. In the typical transgenic animals described herein, the transgene causes cells to express a recombinant form of the subject UbCE protein, e.g. either agonistic or antagonistic forms, or in which the endogenous UbCE gene has been disrupted. However, transgenic animals in which the recombinant UbCE gene is silent are also contemplated, as for example, the FLP or CRE recombinase dependent constructs described below. The "non-human animals" of the invention include vertebrates such as rodents, non-human primates, sheep, dog, cow, amphibians, reptiles, etc. Preferred non-human animals are selected from the rodent family including rat and mouse, most preferably mouse. The term "chimeric animal" is used herein to refer to animals in which the recombinant gene is found, or in which the recombinant is expressed in some but not all cells of the animal. The term "tissue-specific chimeric animal" indicates that the recombinant UbCE gene is present and/or expressed in some tissues but not others.

As used herein, the term "transgene" means a nucleic acid sequence (encoding, e.g., a UbCE polypeptide), which is partly or entirely heterologous, i.e., foreign, to the transgenic animal or cell into which it is introduced, or, is homologous to an endogenous gene of the transgenic animal or cell into which it is introduced, but which is designed to be inserted, or is inserted, into the animal's genome in such a way as to alter the genome of the cell into which it is inserted (e.g., it is inserted at a location which differs from that of the natural gene or its insertion results in a knockout). A transgene can include one or more transcriptional regulatory sequences and any other nucleic acid, such as introns, that may be necessary for optimal expression of a selected nucleic acid.

"Homology" refers to sequence similarity between two peptides or between two nucleic acid molecules. Homology can be determined by comparing a position in each sequence which may be aligned for purposes of comparison. When a position in the compared sequence is occupied by the same base or amino acid, then the molecules are homologous at that position. A degree of homology between sequences is a function of the number of matching or homologous positions shared by the sequences.

The term "evolutionarily related to", with respect to nucleic acid sequences encoding the subject ubiquitin-

conjugating enzymes, refers to nucleic acid sequences which have arisen naturally in an organism, including naturally occurring mutants. The term also refers to nucleic acid sequences which, while derived from a naturally occurring enzymes, have been altered by mutagenesis, as for example, combinatorial mutagenesis described below, yet still encode polypeptides which have at least one activity of a UbCE protein.

As described below, one aspect of this invention pertains to an isolated nucleic acid comprising a nucleotide sequence encoding one of the subject UbCE proteins, fragments thereof encoding polypeptides having at least one biological activity of the UbCE protein, and/or equivalents of such nucleic acids. The term "nucleic acid" as used herein is intended to include such fragments and equivalents. The term "equivalent" is understood to include nucleotide sequences encoding functionally equivalent UbCE proteins or functionally equivalent peptides having an activity of a ubiquitin-conjugating enzyme such as described herein. Equivalent nucleotide sequences will include sequences that differ by one or more nucleotide substitutions, additions or deletions, such as allelic variants; and will also include sequences that differ from the nucleotide sequence encoding the hUbCE protein shown in SEQ ID No: 1, the caUbCE protein shown in SEQ ID No: 3, or the spUbCE protein shown in SEQ ID No: 5, due to the degeneracy of the genetic code. Equivalents will also include nucleotide sequences which hybridize under stringent conditions (i.e., equivalent to about 20°-27° C. below the melting temperature (T_m) of the DNA duplex formed in about 1M salt) to the nucleotide sequence represented in at least one of SEQ ID Nos: 1, 3 or 5. In one embodiment, equivalents will further include nucleic acid sequences derived from and evolutionarily related to the nucleotide sequences shown in any of SEQ ID Nos: 1, 3 or 5.

The term "isolated" as also used herein with respect to nucleic acids, such as DNA or RNA, refers to molecules separated from other DNAs, or RNAs, respectively, that are present in the natural source of the macromolecule. For example, an isolated nucleic acid encoding one of the subject UbCE-proteins preferably includes no more than 10 kilobases (kb) of nucleic acid sequence which naturally immediately flanks the UbCE gene in genomic DNA, more preferably no more than 5 kb of such naturally occurring flanking sequences, and most preferably less than 1.5 kb of such naturally occurring flanking sequence. The term isolated as used herein also refers to a nucleic acid or peptide that is substantially free of cellular material or culture medium when produced by recombinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized. Moreover, an "isolated nucleic acid" is meant to include nucleic acid fragments which are not naturally occurring as fragments and would not be found in the natural state.

Polypeptides referred to herein as possessing the activity of a ubiquitin-conjugating enzyme (UbCE), e.g. are UbCE agonists are understood to have an amino acid sequence identical to or homologous with the amino acid sequences shown in any one of SEQ ID Nos: 2, 4 or 6, and which are capable of forming a thiol ester adduct with the C-terminal carboxyl group of ubiquitin and transferring the ubiquitin to an ϵ -amino group in an acceptor protein by formation of an isopeptide bond. The biological activity of the subject UbCE proteins can include participation in degradative pathways for selective proteolysis of constitutively or conditionally short-lived proteins as well as abnormal proteins. Antagonistic forms of the subject UbCE proteins are defined as

proteins that are homologous, but not identical, to the UbCE proteins represented in SEQ ID Nos: 2, 4 or 6, or that are fragments of the wild-type proteins, which inhibit the transfer of ubiquitin by the naturally occurring form of the ubiquitin-conjugating enzyme. For instance, as described below, mutations in the active site of the enzyme, e.g. Cys-85, can produce dominant negative forms of the subject UbCEs which antagonize the action of the wild-type form of the protein.

Polypeptides referred to in particular as having an activity of an hUbCE protein are defined as peptides that have an amino acid sequence corresponding to all or a portion of the amino acid sequence of the human ubiquitin conjugating enzyme shown in SEQ ID No:2 and which have at least one biological activity of an hUbCE protein- such as an ability to mediate ubiquitination of cellular proteins, e.g. cell-cycle regulatory proteins, e.g., p53; an ability to mediate ubiquitin-dependent degradation of cellular proteins, e.g. cell-cycle regulatory proteins, e.g. p53; an ability to affect the cellular half-life of a cell-cycle regulatory protein, e.g. a cell-cycle checkpoint protein, e.g. p53, e.g. in normal cells, e.g. in normal proliferating cells, e.g. in virally-infected cells, e.g. in papillomavirus infected cells, e.g. in HPV-infected cells, e.g. in HPV-16, HPV-18, HPV-31, or HPV-33 infected cells, e.g. in cells expressing a papillomavirus E6 protein, e.g. in transformed cells, e.g. in cancerous cells. Other biological activities of the subject hUbCE proteins are described herein or will be reasonably apparent to those skilled in the art.

Moreover, it will be generally appreciated that, under certain circumstances, it will be advantageous to provide homologs of naturally-occurring forms of the subject UbCE proteins which are either agonists or antagonists of only a subset of that protein's biological activities. Thus, specific biological effects can be elicited by treatment with a homolog of limited function, and with fewer side effects relative to treatment with agonists or antagonists which are directed to all of the biological activities of that protein. For example, hUbCE homologs can be generated which bind to and inhibit activation of other proteins in the ubiquitin pathway of p53 without substantially interfering with the ubiquitination of other cellular proteins.

In one embodiment, the nucleic acid of the invention encodes a polypeptide which is either an agonist or antagonist the human UbCE protein and comprises an amino acid sequence represented by SEQ ID No: 2. Preferred nucleic acids encode a peptide having an hUbCE protein activity, or which is an antagonist thereof, and being at least 90% homologous, more preferably 95% homologous and most preferably 97% homologous with an amino acid sequence shown in SEQ ID No: 2. Nucleic acids which encode agonist or antagonist forms of an hUbCE protein and having at least about 98-99% homology with a sequence shown in SEQ ID No: 2 are also within the scope of the invention. Preferably, the nucleic acid is a cDNA molecule comprising at least a portion of the nucleotide sequence encoding an hUbCE protein shown in SEQ ID No. 1. A preferred portion of the cDNA molecule shown in SEQ ID No. 1 includes the coding region of the molecule.

In another embodiment, the nucleic acid of the invention encodes a polypeptide which is either an agonist or antagonist a *Candida* UbCE protein, e.g. a *C. albicans* UbCE, and comprises an amino acid sequence represented by SEQ ID No: 4. Preferred nucleic acids encode a peptide having an caUbCE protein activity, or which is an antagonist thereof, and being at least 90% homologous, more preferably 95% homologous and most preferably 97% homologous with an

amino acid sequence shown in SEQ ID No: 4. Nucleic acids which encode agonist or antagonist forms of an caUbCE protein and having at least about 98-99% homology with a sequence shown in SEQ ID No: 4 are also within the scope of the invention. Preferably, the nucleic acid is a cDNA molecule comprising at least a portion of the nucleotide sequence encoding an caUbCE protein shown in SEQ ID No. 3. A preferred portion of the cDNA molecule shown in SEQ ID No. 3 includes the coding region of the molecule. The present invention contemplates closely related homologs (orthologs) from other species of *Candida*, e.g. *Candida stellatoidea*, *Candida tropicalis*, *Candida parapsilosis*, *Candida krusei*, *Candida pseudotropicalis*, *Candida quillerrnondii*, or *Candida rugosa*.

In yet another embodiment, the nucleic acid of the invention encodes a polypeptide which is either an agonist or antagonist a *Schizosaccharomyces* UbCE protein, e.g. an *S. pombe* UbCE, and comprises an amino acid sequence represented by SEQ ID No: 6. Preferred nucleic acids encode a peptide having an spUbCE protein activity, or which is an antagonist thereof, and being at least 90% homologous, more preferably 95% homologous and most preferably 97% homologous with an amino acid sequence shown in SEQ ID No: 6. Nucleic acids which encode agonist or antagonist forms of an spUbCE protein and having at least about 98-99% homology with a sequence shown in SEQ ID No: 6 are also within the scope of the invention. Preferably, the nucleic acid its a cDNA molecule comprising at least a portion of the nucleotide sequence encoding an spUbCE protein shown in SEQ ID No. 5. A preferred portion of the cDNA molecule shown in SEQ ID No. 5 includes the coding region of the molecule.

Another aspect of the invention provides a nucleic acid which hybridizes under high or low stringency conditions to a nucleic acid which encodes a peptide having all or a portion of an amino acid sequence shown in one of SEQ ID Nos: 2, 4 or 6. Appropriate stringency conditions which promote DNA hybridization, for example, 6.0x sodium chloride/sodium citrate (SSC) at about 45° C., followed by a wash of 2.0xSSC at 50° C., are known to those skilled in the art or can be found in *Current Protocols in Molecular Biology*, John Wiley & Sons, N.Y. (1989), 6.3.1-6.3.6. For example, the salt concentration in the wash step can be selected from a low stringency of about 2.0xSSC at 50° C. to a high stringency of about 0.2xSSC at 50° C. In addition, the temperature in the wash step can be increased from low stringency conditions at room temperature, about 22° C., to high stringency conditions at about 65° C.

Isolated nucleic acids which differ in sequence from the nucleotide sequences represented in SEQ ID Nos: 1, 3 or 5 due to degeneracy in the genetic code are also within the scope of the invention. Such nucleic acids can encode functionally equivalent peptides (i.e., a peptide having a biological activity of a UbCE protein) but differ in sequence from the sequence shown in SEQ ID No: 1, 3 or 5 due to degeneracy in the genetic code. For example, a number of amino acids are designated by more than one triplet. Codons that specify the same amino acid, or synonyms (for example, CAU and CAC are synonyms for histidine) may result in "silent" mutations which do not affect the amino acid sequence of the subject UbCE protein. However, it is expected that DNA sequence polymorphisms that do lead to changes in the amino acid sequences of the present hUbCE protein will exist from one human subject to the next. One skilled in the art will appreciate that these variations in one or more nucleotides (up to about 3-4% of the nucleotides) of the nucleic acids encoding peptides having an activity of,

for example, an hUbCE protein may exist among individuals due to natural allelic variation. Any and all such nucleotide variations and resulting amino acid polymorphisms are within the scope of this invention.

Fragments of the nucleic acid encoding an active portion of one of the subject ubiquitin-conjugating enzymes are also within the scope of the invention. As used herein, a fragment of the nucleic acid encoding an active portion of a UbCE protein refers to a nucleotide sequence having fewer nucleotides than the nucleotide sequence encoding the entire amino acid sequence of the protein but which encodes a peptide which possess agonistic or antagonistic activity relative to a naturally occurring form of the enzyme.

Nucleic acid fragments within the scope of the invention also include those capable of hybridizing under high or low stringency conditions with nucleic acids from other species for use in screening protocols to detect UbCE homologs.

Nucleic acids within the scope of the invention may also contain linker sequences, modified restriction endonuclease sites and other sequences useful for molecular cloning, expression or purification of recombinant peptides having at least one biological activity of the subject UbCE ubiquitin-conjugating enzymes. In a preferred embodiment, the nucleic acid fragment comprises at least a portion of the nucleic acid sequence represented by nucleotide residues 319 through 441 of SEQ ID No. 1, corresponding to amino acid residues Cys-107 through Met-147. In preferred embodiments, the nucleic acid encodes an hUbCE polypeptide which includes Cys-107 through Cys-111, and more preferably includes Cys-107 through Asp-117. As illustrated by FIG. 2, certain of the residues from Cys-107 to Asp-111 are important members of the ubiquitin-binding site of hUbCE. Correspondingly, nucleic acid encoding caUbCE or spUbCE preferably include Cys-107 through Val-147 and Cys-107 through Ile-107, respectively.

As indicated by the examples set out below, a nucleic acid encoding a peptide having an activity of the subject ubiquitin-conjugating enzymes may be obtained from mRNA or genomic DNA present in any of a number of eukaryotic cells in accordance with protocols described herein, as well as those generally known to those skilled in the art. A cDNA encoding a homolog of the human UbCE protein, for example, can be obtained by isolating total mRNA from a cell, e.g. a mammalian cell. Double stranded cDNAs can then be prepared from the total mRNA, and subsequently inserted into a suitable plasmid or bacteriophage vector using any one of a number of known techniques. A gene encoding a UbCE protein can also be cloned using established polymerase chain reaction techniques in accordance with the nucleotide sequence information provided by the invention.

Another aspect of the invention relates to the use of the isolated nucleic acid in "antisense" therapy. As used herein, "antisense" therapy refers to administration or in situ generation of oligonucleotide probes or their derivatives which specifically hybridizes (e.g. binds) under cellular conditions, with the cellular mRNA and/or genomic DNA encoding a UbCE protein so as to inhibit expression of that protein, e.g. by inhibiting transcription and/or translation. The binding may be by conventional base pair complementarity, or, for example, in the case of binding to DNA duplexes, through specific interactions in the major groove of the double helix. In general, "antisense" therapy refers to the range of techniques generally employed in the art, and includes any therapy which relies on specific binding to oligonucleotide sequences.

An antisense construct of the present invention can be delivered, for example, as an expression plasmid which, when transcribed in the cell, produces RNA which is complementary to at least a unique portion of the cellular mRNA which encodes a UbCE-protein, e.g. the human UbCE gene represented in SEQ ID No. 1. Alternatively, the antisense construct can be an oligonucleotide probe which is generated ex vivo and which, when introduced into the cell causes inhibition of expression by hybridizing with the mRNA and/or genomic sequences encoding one of the subject UbCE proteins. Such oligonucleotide probes are preferably modified oligonucleotide which are resistant to endogenous nucleases, e.g. exonucleases and/or endonucleases, and are therefore stable in vivo. Exemplary nucleic acid molecules for use as antisense oligonucleotides are phosphoramidate, phosphothioate and methylphosphonate analogs of DNA (see also U.S. Pat. Nos. 5,176,996; 5,264,564; and 5,256,775). Additionally, general approaches to constructing oligomers useful in antisense therapy have been reviewed, for example, by van der Krol et al. (1988) *Biotechniques* 6:958-976; and Stein et al. (1988) *Cancer Res* 48:2659-2668.

Accordingly, the modified oligomers of the invention are useful in therapeutic, diagnostic, and research contexts. In therapeutic applications, the oligomers are utilized in a manner appropriate for antisense therapy in general. For such therapy, the oligomers of the invention can be formulated for a variety of loads of administration, including systemic and topical or localized administration. Techniques and formulations generally may be found in *Remington's Pharmaceutical Sciences*, Meade Publishing Co., Easton, Pa. For systemic administration, injection is preferred, including intramuscular, intravenous, intraperitoneal, and subcutaneous for injection, the oligomers of the invention can be formulated in liquid solutions, preferably in physiologically compatible buffers such as Hank's solution or Ringer's solution. In addition, the oligomers may be formulated in solid form and redissolved or suspended immediately prior to use. Lyophilized forms are also included.

Systemic administration can also be by transmucosal or transdermal means, or the compounds can be administered orally. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration bile salts and fusidic acid derivatives. In addition, detergents may be used to facilitate permeation. Transmucosal administration may be through nasal sprays or using suppositories. For oral administration, the oligomers are formulated into conventional oral administration forms such as capsules, tablets, and tonics. For topical administration, the oligomers of the invention are formulated into ointments, salves, gels, or creams as generally known in the art.

In addition to use in therapy, the oligomers of the invention may be used as diagnostic reagents to detect the presence or absence of the target DNA or RNA sequences to which they specifically bind. Such diagnostic tests are described in further detail below.

This invention also provides expression vectors containing a nucleic acid encoding the subject UbCE proteins, operably linked to at least one transcriptional regulatory sequence. Operably linked is intended to mean that the nucleic acid is linked to a transcriptional regulatory sequence in a manner which allows expression of the enzyme encoded by the nucleic acid, and that expression is, for example, either constitutively or inducibly controlled by

the transcriptional regulatory sequence. Regulatory sequences are art-recognized. Accordingly, the term regulatory sequence includes promoters, enhancers and other expression control elements. Such regulatory sequences are described in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, Calif. (1990). For instance, any of a wide variety of expression control sequences—sequences that control the expression of a DNA sequence when operatively linked to it may be used in these vectors to express DNA sequences encoding the UbCE proteins of this invention. Such useful expression control sequences, include, for example, the early and late promoters of SV40, adenovirus or cytomegalovirus immediate early promoter, the lac system, the trp system, the TAC or TRC system, T7 promoter whose expression is directed by T7 RNA polymerase, the major operator and promoter regions of phage lambda, the control regions for fd coat protein, the promoter for 3-phosphoglycerate kinase or other glycolytic enzymes, the promoters of acid phosphatase, e.g., Pho5, the promoters of the yeast alpha -mating factors, the polyhedron promoter of the baculovirus system and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells or their viruses, and various combinations thereof. It should be understood that the design of the expression vector may depend on such factors as the choice of the host cell to be transformed and/or the type of protein desired to be expressed. Moreover, the vector's copy number, the ability to control that copy number and the expression of any other proteins encoded by the vector, such as antibiotic markers, should also be considered. In one embodiment, the expression vector includes a DNA encoding the subject hUbCE protein, e.g. a recombinant hUbCE protein, e.g. a recombinant protein having an agonistic activity relative to a naturally-occurring form of hUbCE, e.g. a recombinant protein having an antagonistic activity relative to a naturally-occurring form of hUbCE. Similar expression vectors for producing recombinant forms of the caUbCE and spUbCE proteins are also contemplated. Such expression vectors can be used to transfect cells to thereby produce proteins or peptides, including fusion proteins or peptides encoded by nucleic acids as described herein.

Moreover, hUbCE-expression vectors can be used as a part of a gene therapy protocol to reconstitute hUbCE function in a cell in which hUbCE is misexpressed, or alternatively, to provide an antagonist of the naturally-occurring hUbCE or an antisense construct—such as to inhibit the hUbCE-mediated degradation of p53. For instance, expression constructs of the subject hUbCE-proteins may be administered in any biologically effective carrier, e.g. any formulation or composition capable of effectively transfecting cells in vivo with a recombinant hUbCE-gene. Approaches include insertion of the subject gene in viral vectors including recombinant retroviruses, adenovirus, adeno-associated virus, and herpes simplex virus-1, or recombinant bacterial or eukaryotic plasmids. Viral vectors can be used to transfect cells directly; plasmid DNA can be delivered with the help of, for example, cationic liposomes (lipofectin) or derivatized (e.g. antibody conjugated), polylysine conjugates, gramicidin S, artificial viral envelopes or other such intracellular carriers, as well as direct injection of the gene construct or CaPO₄ precipitation carried out in vivo. It will be appreciated that because transduction of appropriate target cells represents the critical first step in gene therapy, choice of the particular gene delivery system will depend on such factors as the phenotype of the intended target and the route of administration, e.g. locally or systemically.

A preferred approach for in vivo introduction of nucleic acid encoding one of the subject proteins into a cell is by use of a viral vector containing nucleic acid, e.g. a cDNA, encoding the gene product. Infection of cells with a viral vector has the advantage that a large proportion of the targeted cells can receive the nucleic acid. Additionally, molecules encoded within the viral vector, e.g., by a cDNA contained in the viral vector, are expressed efficiently in cells which have taken up viral vector nucleic acid.

Retrovirus vectors and adeno-associated virus vectors are generally understood to be the recombinant gene delivery system of choice for the transfer of exogenous genes in vivo, particularly into humans. These vectors provide efficient delivery of genes into cells, and the transferred nucleic acids are stably integrated into the chromosomal DNA of the host. A major prerequisite for the use of retroviruses is to ensure the safety of their use, particularly with regard to the possibility of the spread of wild-type virus in the cell population. The development of specialized cell lines (termed "packaging cells") which produce only replication-defective retroviruses has increased the utility of retroviruses for gene therapy, and defective retroviruses are well characterized for use in gene transfer for gene therapy purposes (for a review see Miller, A. D. (1990) *Blood* 76:271). Thus, recombinant retrovirus can be constructed in which part of the retroviral coding sequence (gag, pol, env) has been replaced by nucleic acid encoding one of the subject hUbCE-proteins rendering the retrovirus replication defective. The replication defective retrovirus is then packaged into virions which can be used to infect a target cell through the use of a helper virus by standard techniques. Protocols for producing recombinant retroviruses and for infecting cells in vitro or in vivo with such viruses can be found in *Current Protocols in Molecular Biology*, Ausubel, F. M. et al. (eds.) Greene Publishing Associates, (1989), Sections 9.10–9.14 and other standard laboratory manuals. Examples of suitable retroviruses include pLJ, pZIP, pWE and pEM which are well known to those skilled in the art. Examples of suitable packaging virus lines for preparing both ecotropic and amphotropic retroviral systems include ψ Crip, ψ Cre, ψ 2 and ψ Am. Retroviruses have been used to introduce a variety of genes into many different cell types, including neural cells, epithelial cells, endothelial cells, lymphocytes, myoblasts, hepatocytes, bone marrow cells, in vitro and/or in vivo (see for example Eglitis, et al. (1985) *Science* 230:1395–1398; Danos and Mulligan (1988) *Proc. Natl. Acad. Sci. USA* 85:6460–6464; Wilson et al. (1988) *Proc. Natl. Acad. Sci. USA* 85:3014–3018; Armentano et al. (1990) *Proc. Natl. Acad. Sci. USA* 87:6141–6145; Huber et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8039–8043; Ferry et al. (1991) *Proc. Natl. Acad. Sci. USA* 88:8377–8381; Chowdhury et al. (1991) *Science* 254:1802–1805; van Beusechem et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:7640–7644; Kay et al. (1992) *Human Gene Therapy* 3:641–647; Dai et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:10892–10895; Hwu et al. (1993) *J. Immunol.* 150:4104–4115; U.S. Pat. No. 4,868,116; U.S. Pat. No. 4,980,286; PCT Application WO 89/07136; PCT Application WO 89/02468; PCT Application WO 89/05345; and PCT Application WO 92/07573).

Furthermore, it has also been shown that it is possible to limit the infection spectrum of retroviruses and consequently of retroviral-based vectors, by modifying the viral packaging proteins on the surface of the viral particle (see, for example PCT publications WO93/25234, WO94/06920, and WO94/11524). For instance, strategies for the modification of the infection spectrum of retroviral vectors include:

coupling antibodies specific for cell surface antigens to the viral env protein (Roux et al. (1989) *PNAS* 86:9079-9083; Julan et al (1992) *J. Gen Virol* 73:3251-3255; and Goud et al. (1983) *Virology* 163:251-254); or coupling cell surface ligands to the viral env proteins (Neda et al. (1991) *J Biol Chem* 266:14143-14146). Coupling can be in the form of the chemical cross-linking with a protein or other variety (e.g. lactose to convert the env protein to an asialoglycoprotein), as well as by generating fusion proteins (e.g. single-chain antibody/env fusion proteins). This technique, while useful to limit or otherwise direct the infection to certain tissue types, and can also be used to convert an ecotropic vector in to an amphotropic vector.

Moreover, use of retroviral gene delivery can be further enhanced by the use of tissue- or cell-specific transcriptional regulatory sequences which control expression of the hUbCE-gene of the retroviral vector.

Another viral gene delivery system useful in the present invention utilizes adenovirus-derived vectors. The genome of an adenovirus can be manipulated such that it encodes a gene product of interest, but is inactivate in terms of its ability to replicate in a normal lytic viral life cycle (see, for example, Berkner et al. (1988) *BioTechniques* 6:616; Rosenfeld et al. (1991) *Science* 252:431-434; and Rosenfeld et al. (1992) *Cell* 68:143-155). Suitable adenoviral vectors derived from the adenovirus strain Ad type 5 dl324 or other strains of adenovirus (e.g. Ad2, Ad3, Ad7 etc.) are well known to those skilled in the art. Recombinant adenoviruses can be advantageous in certain circumstances in that they are not capable of infecting nondividing cells and can be used to infect a wide variety of cell types, including airway epithelium (Rosenfeld et al. (1992) cited supra), endothelial cells (Lemarchand et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:6482-6486), hepatocytes (Herz and Gerard (1993) *Proc. Natl. Acad. Sci. USA* 90:2812-2816) and muscle cells (Quantin et al. (1992) *Proc. Natl. Acad. Sci. USA* 89:2581-2584). Furthermore, the virus particle is relatively stable and amenable to purification and concentration, and as above, can be modified so as to affect the spectrum of infectivity. Additionally, introduced adenoviral DNA (and foreign DNA contained therein) is not integrated into the genome of a host cell but remains episomal, thereby avoiding potential problems that can occur as a result of insertional mutagenesis in situations where introduced DNA becomes integrated into the host genome (e.g., retroviral DNA). Moreover, the carrying capacity of the adenoviral genome for foreign DNA is large (up to 8 kilobases) relative to other gene delivery vectors (Berkner et al., supra; Haj-Ahmand and Graham (1986) *J. Virol.* 57:267). Most replication-defective adenoviral vectors currently in use and therefore flavored by the present invention are deleted for all or parts of the viral E1 and E3 genes but retain as much as 80% of the adenoviral genetic material (see, e.g., Jones et al. (1979) *Cell* 16:683; Berkner et al., supra; and Graham et al. in *Methods in Molecular Biology*, E. J. Murray, Ed. (*Humana, Clifton, N.J.*, 1991) vol.7. pp. 109-127). Expression of the inserted hUbCE-gene can be under control of, for example, the E1A promoter, the major late promoter (MLP) and associated leader sequences, the E3 promoter, or exogenously added promoter sequences.

Yet another viral vector system useful for delivery of the subject hUbCE-gene is the adeno-associated virus (AAV). Adeno-associated virus is a naturally occurring defective virus that requires another virus, such as an adenovirus or a herpes virus, as a helper virus for efficient replication and a productive life cycle. (For a review see Muzyczka et al. *Curr. Topics in Micro. and Immunol.* (1992) 158:97-129). It

is also one of the few viruses that may integrate its DNA into non-dividing cells, and exhibits a high frequency of stable integration (see for example Flotte et al. (1992) *Am. J. Respir. Cell. Mol. Biol.* 7:349-356; Samulski et al. (1989) *J. Virol.* 63:3822-3828; and McLaughlin et al. (1989) *J. Virol.* 62:1963-1973). Vectors containing as little as 300 base pairs of AAV can be packaged and can integrate. Space for exogenous DNA is limited to about 4.5 kb. An AAV vector such as that described in Tratschin et al. (1985) *Mol. Cell. Biol.* 5:3251-3260 can be used to introduce DNA into cells. A variety of nucleic acids have been introduced into different cell types using AAV vectors (see for example Hermonat et al. (1984) *Proc. Natl. Acad. Sci. USA* 81:6466-6470; Tratschin et al. (1985) *Mol. Cell. Biol.* 4:2072-2081; Wondisford et al. (1988) *Mol. Endocrinol.* 2:32-39; Tratschin et al. (1984) *J. Virol.* 51:611-619; and Flotte et al. (1993) *J. Biol. Chem.* 268:3781-3790).

Other viral vector systems that may have application in gene therapy have been derived from herpes virus, vaccinia virus, and several RNA viruses. In particular, herpes virus vectors may provide a unique strategy for persistence of the recombinant hUbCE-gene in cells of the central nervous system and ocular tissue (Pepose et al. (1994) *Invest Ophthalmol Vis Sci* 35:2662-2666).

In addition to viral transfer methods, such as those illustrated above, non-viral methods can also be employed to cause expression of an hUbCE-protein, or an hUbCE anti-sense molecule, in the tissue of an animal. Most nonviral methods of gene transfer rely on normal mechanisms used by mammalian cells for the uptake and intracellular transport of macromolecules. In preferred embodiments, non-viral gene delivery systems of the present invention rely on endocytic pathways for the uptake of the subject hUbCE-gene by the targeted cell. Exemplary gene delivery systems of this type include liposomal derived systems, poly-lysine conjugates, and artificial viral envelopes.

In a representative embodiment, a gene encoding one of the subject hUbCE-proteins can be entrapped in liposomes bearing positive charges on their surface (e.g., lipofectins) and (optionally) which are tagged with antibodies against cell surface antigens of the target tissue (Mizuno et al. (1992) *No Shinkei Geka* 20:547-551; PCT publication WO91/06309; Japanese patent application 1047381; and European patent publication EP-A-43075). For example, lipofection of papilloma-virus infected epithelial cells can be carried out using liposomes tagged with monoclonal antibodies against, for example, squamous cells.

In similar fashion, the gene delivery system comprises an antibody or cell surface ligand which is cross-linked with a gene binding agent such as poly-lysine (see, for example, PCT publications WO93/04701, WO92/22635, WO92/20316, WO92/19749, and WO92/06180). For example, the subject hUbCE-gene construct can be used to transfect HPV-infected squamous cells in vivo using a soluble poly-nucleotide carrier comprising an HPV viral coat protein conjugated to a polycation, e.g. poly-lysine (see U.S. Pat. No. 5,166,320). It will also be appreciated that effective delivery of the subject nucleic acid constructs via-mediated endocytosis can be improved using agents which enhance escape of the gene from the endosomal structures. For instance, whole adenovirus or fusogenic peptides of the influenza HA gene product can be used as part of the delivery system to induce efficient disruption of DNA-containing endosomes (Mulligan et al. (1993) *Science* 260-926; Wagner et al (1992) *PNAS* 89:7934; and Christiano et al. (1993) *PNAS* 90:2122).

In clinical settings, the gene delivery systems can be introduced into a patient by any of a number of methods,

each of which is familiar in the art. For instance, a pharmaceutical preparation of the gene delivery system can be introduced systemically, e.g. by intravenous injection, and specific transduction of the in the target cells occurs predominantly from specificity of transfection provided by the gene delivery vehicle, cell-type or tissue-type expression due to the transcriptional regulatory sequences controlling expression of the gene, or a combination thereof. In other embodiments, initial delivery of the recombinant gene is more limited with introduction into the animal being quite localized. For example, the gene delivery vehicle can be introduced by catheter (see U.S. Pat. No. 5,328,470) or by stereotactic injection (e.g. Chen et al. (1994) *PNAS* 91:3054-3057).

Moreover, the pharmaceutical preparation can consist essentially of the gene delivery system in an acceptable diluent, or can comprise a slow release matrix in which the gene delivery vehicle is imbedded. Alternatively, where the complete gene delivery system can be produced intact from recombinant cells, e.g. retroviral packages, the pharmaceutical preparation can comprise one or more cells which produce the gene delivery system. In the case of the latter, methods of introducing the viral packaging cells may be provided by, for example, rechargeable or biodegradable devices. The generation of such implants is generally known in the art. See, for example, *Concise Encyclopedia of Medical & Dental Materials*, ed. by David Williams (MIT Press: Cambridge, Mass., 1990); Sabel et al. U.S. Pat. No. 4,883,666; Aebischer et al. U.S. Pat. No. 4,892,538; Aebischer et al. U.S. Pat. No. 5,106,627; Lim U.S. Pat. No. 4,391,909; Sefton U.S. Pat. No. 4,353,888; and Aebischer et al. (1991) *Biomaterials* 12:50-55).

This invention also pertains to a host cell transfected or transformed to express a recombinant forms of the subject UbCE proteins. The host cell may be any prokaryotic or eukaryotic cell. For example, an hUbCE protein of the present invention may be expressed in bacterial cells such as *E. Coli*, insect cells (baculovirus), yeast, or mammalian cells. Other suitable host cells are known to those skilled in the art.

The term "recombinant protein" refers to a protein of the present invention which is produced by recombinant DNA techniques, wherein generally DNA encoding the UbCE protein is inserted into a suitable expression vector which is in turn used to transform a host cell to produce the heterologous protein. Moreover, the phrase "derived from", with respect to a recombinant gene encoding the recombinant UbCE, is meant to include within the meaning of "recombinant protein" those proteins having an amino acid sequence of a native UbCE, e.g. hUbCE, caUbCE or spUbCE, or an amino acid sequence similar thereto which is generated by mutations including substitutions and deletions of a naturally occurring form of the protein. Recombinant proteins preferred by the present invention, in addition to native proteins, are at least 90% homologous, more preferably 95% homologous and most preferably 97% homologous with an amino acid sequence shown in one of SEQ ID Nos: 2, 4 or 6. Polypeptides having an activity of an hUbCE protein, or which are antagonistic thereto, and which are at least about 90%, more preferably at least about 95%, and most preferably at least about 98-99% homologous with a sequence shown in SEQ ID No: 2, 4 or 6 are also within the scope of the invention.

The present invention further pertains to recombinant UbCE homologs which are encoded by genes derived from other non-human mammals, e.g. mouse, rat, rabbit, or pig, and which have amino acid sequences evolutionarily related

to an hUbCE protein. As described above, such recombinant hUbCE proteins preferably are capable of functioning in one of either role of an agonist or antagonist of at least one biological activity of an hUbCE. The term "evolutionarily related to", as set out above, refers to ubiquitin-conjugating enzymes having amino acid sequences which have arisen naturally, or which are mutationally derived, for example, by combinatorial mutagenesis or scanning mutagenesis, but which proteins are homologous to the human UbCE protein represented in SEQ ID No: 2.

The present invention further pertains to methods of producing the subject proteins. For example, a host cell transfected with an expression vector encoding one of the subject UbCE proteins can be cultured under appropriate conditions to allow expression of the peptide to occur. The peptide may be secreted (e.g. through use of recombinantly added signal sequence) and isolated from a mixture of cells and medium containing the secreted protein. Alternatively, the peptide may be retained cytoplasmically, as it presumably is its naturally occurring form, and the cells harvested, lysed and the protein isolated. A cell culture includes host cells, media and other byproducts. Suitable media for cell culture are well known in the art. The subject UbCE polypeptides can be isolated from cell culture medium, host cells, or both using techniques known in the art for purifying proteins including ion-exchange chromatography, gel filtration chromatography, ultrafiltration, electrophoresis, and immunoaffinity purification with antibodies raised against the protein. In a preferred embodiment, the UbCE protein is a fusion protein containing a domain which facilitates its purification, such as the hUbCE-GST fusion protein described below.

Thus, a nucleotide sequence derived from the cloning of a UbCE protein of the present invention, encoding all or a selected portion of the protein, can be used to produce a recombinant form of the enzyme via microbial or eukaryotic cellular processes. Ligating the polynucleotide sequence into a gene construct, such as an expression vector, and transforming or transfecting into hosts, either eukaryotic (yeast, avian, insect or mammalian) or prokaryotic (bacterial cells), are standard procedures used in producing other well-known proteins, e.g. insulin, interferons, human growth hormone, IL-1, IL-2, and the like. Similar procedures, or modifications thereof, can be employed to prepare recombinant proteins, or portions thereof, by microbial means or tissue-culture technology in accord with the subject invention.

The recombinant protein can be produced by ligating the cloned gene, or a portion thereof, into a vector suitable for expression in either prokaryotic cells, eukaryotic cells, or both. Expression vehicles for production of recombinant UbCEs include plasmids and other vectors. For instance, suitable vectors for the expression of the subject proteins include plasmids of the types: pBR322-derived plasmids, pEMBL-derived plasmids, pEX-derived plasmids, pBTac-derived plasmids and pUC-derived plasmids for expression in prokaryotic cells, such as *E. coli*.

A number of vectors exist for the expression of recombinant proteins in yeast. For instance, YEP24, YIP5, YEP51, YEP52, pYES2, and YRP17 are cloning and expression vehicles useful in the introduction of genetic constructs into *S. cerevisiae* (see, for example, Broach et al. (1983) in *Experimental Manipulation of Gene Expression*, ed. M. Inouye Academic Press, p. 83, incorporated by reference herein). These vectors can replicate in *E. coli* due the presence of the pBR322 ori, and in *S. cerevisiae* due to the replication determinant of the yeast 2 micron plasmid. In addition, drug resistance markers such as ampicillin can be used.

The preferred mammalian expression vectors contain both prokaryotic sequences to facilitate the propagation of the vector in bacteria, and one or more eukaryotic transcription units that are expressed in eukaryotic cells. The pcDNA1/amp, pcDNA1/neo, pRc/CMV, pSV2gpt, pSV2neo, pSV2-dhfr, pTk2, pRSVneo, pMSG, pSVT7, pko-neo and pHyg derived vectors are examples of mammalian expression vectors suitable for transfection of eukaryotic cells. Some of these vectors are modified with sequences from bacterial plasmids, such as pBR322, to facilitate replication and drug resistance selection in both prokaryotic and eukaryotic cells. Alternatively, derivatives of viruses such as the bovine papilloma virus (BPV-1), or Epstein-Barr virus (pHEBo, pREP-derived and p205) can be used for transient expression of proteins in eukaryotic cells. The various methods employed in the preparation of the plasmids and transformation of host organisms are well known in the art. For other suitable expression systems for both prokaryotic and eukaryotic cells, as well as general recombinant procedures, see *Molecular Cloning: A Laboratory Manual*, 2nd Ed., ed. by Sambrook, Fritsch and Maniatis (Cold Spring Harbor Laboratory Press:1989) Chapters 16 and 17.

In some instances, it may be desirable to express the recombinant UbCE by the use of a baculovirus expression system. Examples of such baculovirus expression systems include pVL-derived vectors (such as pVL1392, pVL1393 and pVL941), pAcUW-derived vectors (such as pAcUW1), and pBlueBac-derived vectors (such as the β -gal containing pBlueBac III).

When expression of a portion of the ubiquitin-conjugating enzyme is desired, i.e. a truncation mutant, it may be necessary to add, a start codon (ATG) to the oligonucleotide fragment containing the desired sequence to be expressed. It is well known in the art that a methionine at the N-terminal position can be enzymatically cleaved by the use of the enzyme methionine aminopeptidase (MAP). MAP has been cloned from *E. coli* (Ben-Bassat et al. (1987) *J. Bacteriol.* 169:751-757) and *Salmonella typhimurium* and its in vitro activity has been demonstrated on recombinant proteins (Miller et al. (1987) *PNAS* 84:2718-1722). Therefore, removal of an N-terminal methionine, if desired, can be achieved either in vivo by expressing UbCE-derived polypeptides in a host which produces MAP (e.g., *E. coli* or CM89 or *S. cerevisiae*), or in vitro by use of purified MAP (e.g., procedure of Miller et al.).

Alternatively, the coding sequences for the polypeptide can be incorporated as a part of a fusion gene including a nucleotide sequence encoding a different polypeptide. This type of expression system can be useful under conditions where it is desirable to produce an immunogenic fragment of a UbCE protein. In an exemplary embodiment, the VP6 capsid protein of rotavirus can be used as an immunologic carrier protein for portions of the hUbCE polypeptide, either in the monomeric form or in the form of a viral particle. The nucleic acid sequences corresponding to the portion of the hUbCE protein to which antibodies are to be raised can be incorporated into a fusion gene construct which includes coding sequences for a late vaccinia virus structural protein to produce a set of recombinant viruses expressing fusion proteins comprising a portion of the protein hUbCE as part of the virion. It has been demonstrated with the use of immunogenic fusion proteins utilizing the Hepatitis B surface antigen fusion proteins that recombinant Hepatitis B virions can be utilized in this role as well. Similarly, chimeric constructs coding for fusion proteins containing a portion of an UbCE protein and the poliovirus capsid protein can be created to enhance immunogenicity of the set of

polypeptide antigens (see, for example, EP Publication No. 0259149; and Evans et al. (1989) *Nature* 339:385; Huang et al. (1988) *J. Virol.* 62:3855; and Schlienger et al. (1992) *J. Virol.* 66:2).

5 The Multiple Antigen Peptide system for peptide-based immunization can also be utilized, wherein a desired portion of a UbCE protein is obtained directly from organo-chemical synthesis of the peptide onto an oligomeric branching lysine core (see, for example, Posnett et al. (1988) *J Biol Chem* 263:1719 and Nardelli et al. (1992) *J. Immunol.* 148:914). Antigenic determinants of the UbCE proteins can also be expressed and presented by bacterial cells.

10 In addition to utilizing fusion proteins to enhance immunogenicity, it is widely appreciated that fusion proteins can also facilitate the expression of proteins, such as the UbCE proteins of the present invention. For example, as described below, the hUbCE protein can be generated as a glutathione-S-transferase (GST) fusion protein. Such GST fusion proteins can enable purification of the hUbCE protein, such as by the use of glutathione-derivatized matrices (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. (NY: John Wiley & Sons, 1991); Smith et al. (1988) *Gene* 67:31; and Kaelin et al. (1992) *Cell* 70:351). In another embodiment, a fusion gene coding for a purification leader sequence, such as a poly-(His)/enterokinase cleavage site sequence at the N-terminus of the desired portion of the hUbCE protein, can allow purification of the expressed hUbCE-fusion protein by affinity chromatography using a Ni²⁺ metal resin. The purification leader sequence can then be subsequently removed by treatment with enterokinase (e.g., see Hochuli et al. (1987) *J. Chromatography* 411:177; and Janknecht et al. *PNAS* 88:8972). Similar constructs can be generated for expression of caUbCE and spUbCE.

15 Techniques for making fusion genes are well known. Essentially, the joining of various DNA fragments coding for different polypeptide sequences is performed in accordance with conventional techniques, employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. In another embodiment, the fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons 1992).

20 Another aspect of the invention pertains to isolated peptides having an activity of the naturally occurring form(s) of the subject UbCE proteins or which are antagonists of at least one biological activity of the naturally occurring form of the subject UbCE proteins. A peptide having an activity of an hUbCE protein has at least one biological activity of an hUbCE protein. As set out above, in preferred embodiments, a biological activity of an hUbCE protein includes an ability to mediate ubiquitination of cellular proteins, such as cell-cycle regulatory proteins, erg. p53. In particular, the hUbCE of the present invention is able to mediate ubiquitin-dependent degradation of p53. Other biological activities of the subject hUbCE protein are described herein or will be reasonably apparent to those skilled in the art. A peptide having at least one biological activity of the subject hUbCE protein may differ in amino acid sequence

from the sequence shown in SEQ ID No: 2 but preferably, such differences result in a modified protein which functions in the same or similar manner (e.g. agonist) as a native hUbCE protein or which has the same or similar characteristics of a native hUbCE protein. Furthermore, as described herein, peptides having amino acid sequences homologous to SEQ ID No. 2 yet which function as antagonists of a naturally occurring hUbCE protein are also contemplated by the present invention. Various modifications of the hUbCE protein to produce these and other functionally equivalent peptides are described in detail herein. In similar fashion, homologs of the subject caUBC and spUBC polypeptides are contemplated, including both agonistic and antagonistic forms. The term peptide, as used herein, refers to peptides, proteins, and polypeptides.

The present invention also makes available isolated UbCE proteins, which proteins are isolated from or otherwise substantially free of other extracellular proteins, especially other proteins of the ubiquitin conjugating system (i.e. other E1 or E2 enzymes, as well as E3 proteins or ubiquitin) normally associated with the ubiquitin-conjugating enzyme in the cellular milieu. The term "substantially free of other extracellular proteins" (also referred to herein as "contaminating proteins") is defined as encompassing preparations of the subject UbCE protein comprising less than 20% (by dry weight) contaminating protein, and preferably comprising less than 5% contaminating protein. Functional forms of the subject UbCE proteins can be prepared, for the first time, as purified preparations by using a cloned gene as described herein. By "purified", it is meant, when referring to a peptide or DNA or RNA sequence, that the indicated molecule is present in the substantial absence of other biological macromolecules, such as other proteins (particularly other enzymes of the ubiquitin system such as other E1 or E2 proteins, as well as other contaminating proteins). The term "purified" as used herein preferably means at least 80% by dry weight, more preferably in the range of 95-99% by weight, and most preferably at least 99.8% by weight, of biological macromolecules of the same type present (but water, buffers, and other small molecules, especially molecules having a molecular weight of less than 5000, can be present). The term "pure" as used herein preferably has the same numerical limits as "purified" immediately above. "Isolated" and "purified" do not encompass either natural materials in their native state or natural materials that have been separated into components (e.g., in an acrylamide gel) but not obtained either as pure (e.g. lacking contaminating proteins or chromatography reagents such as denaturing agents and polymers, e.g. acrylamide or agarose) substances or solutions.

Isolated peptide, having an activity of an UbCE protein, or which can function as antagonists of a naturally occurring form of the UbCE protein described herein can also be obtained by screening peptides recombinantly produced from the corresponding fragment of the nucleic acids encoding such peptides. In addition, fragments can be chemically synthesized using techniques known in the art such as conventional Merrifield solid phase f-Moc or t-Boc chemistry. For example, the hUbCE protein may be arbitrarily divided into fragments of desired length with no overlap of the fragments, or preferably divided into overlapping fragments of a desired length. The fragments can be produced (recombinantly or by chemical synthesis) and tested to identify those peptides having an hUbCE protein activity or alternatively to identify antagonists. Similar manipulation of the caUbCE and soUbCE proteins can be carried out.

Furthermore, it is also possible to modify the structure of a UbCE polypeptide for such purposes as enhancing thera-

peutic or prophylactic efficacy, or stability (e.g., shelf life *ex vivo* and resistance to proteolytic degradation *in vivo*). Such modified peptides are considered functional equivalents of peptides having an activity of, or which antagonize, a UbCE protein as defined herein. A modified polypeptide can be produced in which the amino acid sequence has been altered, such as by amino acid substitution, deletion, or addition.

For example, it is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid (i.e. conservative mutations) will not have a major effect on the biological activity of the resulting molecule. Conservative replacements are those that take place within a family of amino acids that are related in their side chains. Genetically encoded amino acids can be divided into four families: (1) acidic=aspartate, glutamate; (2) basic=lysine, arginine, histidine; (3) nonpolar=alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan; and (4) uncharged polar=glycine, asparagine, glutamine, cysteine, serine, threonine, tyrosine. Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids. In similar fashion, the amino acid repertoire can be grouped as (1) acidic=aspartate, glutamate; (2) basic=lysine, arginine, histidine, (3) aliphatic=glycine, alanine, valine, leucine, isoleucine, serine, threonine, with serine and threonine optionally be grouped separately as aliphatic-hydroxyl; (4) aromatic=phenylalanine, tyrosine, tryptophan; (5) amide=asparagine, glutamine; and (6) sulfur-containing=cysteine and methionine. (see, for example, *Biochemistry*, 2nd ed, Ed. by L. Stryer, WH Freeman and Co.: 1981). Whether a change in the amino acid sequence of a peptide results in a functional UbCE homolog can be readily determined by assessing the ability of the variant peptide to, for instance, mediate ubiquitination in a fashion similar to the wild-type UbCE. Peptides in which more than one replacement has taken place can readily be tested in the same manner.

The invention also includes a method of generating sets of combinatorial mutants of the subject UbCE proteins, as well as truncation and fragmentation mutants, and is especially useful for identifying potential variant sequences which are functional in ubiquitinating cellular proteins. One purpose for screening such combinatorial libraries is, for example, to isolate novel UbCE homologs which act as antagonist of the wild-type ("authentic") UbCE activity, e.g. an hUbCE homolog which inhibits p53 ubiquitination, or alternatively, possess novel activities all together. Such proteins, when expressed from recombinant DNA constructs, can be used in gene therapy protocols.

Likewise, mutagenesis can give rise to UbCE homologs which have intracellular half-lives dramatically different than the corresponding wild-type protein. For example, the altered protein can be rendered either more stable or less stable to proteolytic degradation or other cellular process which result in destruction of, or otherwise inactivation of, a naturally occurring form of the subject hUbCE protein. Such hUbCE homologs (either agonist or antagonist homologs), and the genes which encode them, can be utilized to alter the envelope of recombinant hUbCE expression by modulating The half-life of the protein. For instance, a short half-life for the recombinant hUbCE can give rise to more transient biological effects associated with that homolog and, when part of an inducible expression system, can allow tighter control of recombinant hUbCE levels within the cell. As above, such proteins, and particularly their recombinant nucleic acid constructs, can be used in gene therapy protocols.

In one aspect of this method, the amino acid sequences for a population of UbCE homologs or other related proteins are aligned, preferably to promote the highest homology possible. Such a population of variants can include, for example, hUbCE homologs from one or more species, or UbCE homologs from the same species but which differ due to mutation. Amino acids which appear at each position of the aligned sequences are selected to create a degenerate set of combinatorial sequences. For instance, alignment of the hUbCE, caUbCE and spUbCE sequences provided in the appended sequence listing (see also FIG. 5) can be used to generate a degenerate library of UbCE proteins represented by the general formula:

Met Xaa(1) Leu Lys Arg Ile Xaa(2) Xaa(3) Glu Leu
Xaa(4) Asp Leu Xaa(5) Xaa(6) Asp Pro Pro Xaa(7)
Xaa(8) Cys Ser Ala Gly Pro Val Gly Asp Asp Xaa(9)
Xaa(10) His Trp Gln Ala Xaa(11) Ile Met Gly Pro Asn
Asp Ser Pro Tyr Xaa(12) Gly Gly Val Phe Phe Leu
Xaa(13) Ile His Phe Pro Thr Asp Tyr Pro Xaa(14) Lys
Pro Pro Lys Xaa(15) Xaa(16) Xaa(17) Thr Thr Xaa(18)
Ile Tyr His Pro Asn Ile Asn Ser Asn Gly Xaa(19) Ile
Cys Leu Asp Ile Leu Xaa(20) Xaa(21) Gln Trp Ser Pro
Ala Leu Thr Ile Ser Lys Val Leu Leu Ser Ile Cys Ser
Leu Leu Xaa(22) Asp Xaa(23) Asn Pro Asp Asp Pro
Leu Val Pro Glu Ile Ala Xaa(24) Xaa(25) Tyr Xaa(26)
Xaa(27) Asp Arg Xaa(28) Xaa(29) Tyr Xaa(30) Xaa
(31) Xaa(32) Ala Xaa(33) Glu Trp Thr Xaa(34) Lys Tyr
Ala Xaa(35) (SEQ ID No. 7)

wherein Xaa(1) represents Ala or Ser; Xaa(2) represents His or Asn; Xaa(3) represents Lys or Arg; Xaa(4) represents Ala, Ser or Asn; Xaa(5) represents Gly or Ala; Xaa(6) represents Arg or Lys; Xaa(7) represents Ala or Ser; Xaa(8) represents Gln or Ser; Xaa(9) represents Leu or Met; Xaa(10) represents Phe or Tyr; Xaa(11) represents Ser or Thr; Xaa(12) represents Gln or Ala; Xaa(13) represents Ser or Thr; Xaa(14) represents Leu or Phe; Xaa(15) represents Val or Ile; Xaa(16) represents Ala or Asn; Xaa(17) represents Leu or Phe; Xaa(18) represents Arg or Lys; Xaa(19) represents Ser or Asn; Xaa(20) represents Arg or Lys; Xaa(21) represents Ser or Asp; Xaa(22) represents Thr or Cys; Xaa(23) represents Ala or Pro; Xaa(24) represents Arg or His; Xaa(25) represents Val or Ile; Xaa(26) represents Lys or Gln; Xaa(27) represents Thr or Gln; Xaa(28) represents Ser, Lys or Glu; Xaa(29) represents Arg or Lys; Xaa(30) represents Asn or Gln; Xaa(31) represents Ala, Leu or Arg; Xaa(32) represents Ile, Ser or Thr; Xaa(33) represents Arg or Lys; Xaa(34) represents Arg, Lys or Gln; Xaa(35) represents Val, Ile or Met.

To further expand the library, each of the degenerate positions (Xaa) can be rendered even more degenerate by including other amino acid residues which are of the same "family" as the residues which appear in each of the UbCEs, e.g. Xaa(1) can be Gly, Ala, Val, Leu, Ile, Ser or Thr (e.g. aliphatic), Xaa(22) can be Ser, Thr, Cys or Met (aliphatic-hydroxyl and sulfur-containing), etc. Alternatively, isosteric substitutions can be made without regard to, for example, charge or polarity of the amino acid sidechain. For instance, Xaa(17) can be Leu, Ile, Asn, Met, Phe or Tyr, as the sidechains of Ile, Asn and Met each occupy approximately the same steric space as Leu, and Tyr is isosteric for Phe. Likewise, where the degeneracy is conserved from the human and yeast homologs, the degenerate library can, at that position, only include, for example, the amino acid residue which occurs in the human UbCE. To illustrate, Xaa(3) is a Lysine in hUbCE and caUbCE, and Arginine in spUbCE. In a library which rejects conservative mutations of the human UbCE as equivalent, Xaa(3) would be Lys.

In a preferred embodiment, the combinatorial UbCE library is produced by way of a degenerate library of genes encoding a library of polypeptides which each include at least a portion of potential UbCE sequences. A mixture of synthetic oligonucleotides can be enzymatically ligated into gene sequences such that the degenerate set of potential UbCE sequences are expressible as individual polypeptides, or alternatively, as a set of larger fusion proteins (e.g. for phage display) containing the set of UbCE sequences therein.

There are many ways by which the library of potential UbCE homologs can be generated from a degenerate oligonucleotide sequence. Chemical synthesis of a degenerate gene sequence can be carried out in an automatic DNA synthesizer, and the synthetic genes then be ligated into an appropriate gene for expression. The purpose of a degenerate set of genes is to provide, in one mixture, all of the sequences encoding the desired set of potential UbCE sequences. The synthesis of degenerate oligonucleotides is well known in the art (see, for example, Narang, S. A. (1983) *Tetrahedron* 39:3; Itakura et al. (1981) *Recombinant DNA, Proc 3rd Cleveland Sympos. Macromolecules*, ed. A. G. Walton, Amsterdam: Elsevier pp273-289; Itakura et al. (1984) *Annu. Rev. Biochem.* 53:323; Itakura et al. (1984) *Science* 198:1056; Ike et al. (1983) *Nucleic Acid Res.* 11:477. Such techniques have been employed in the directed evolution of other proteins (see, for example, Scott et al. (1990) *Science* 249:386-390; Roberts et al. (1992) *PNAS* 89:2429-2433; Devlin et al. (1990) *Science* 249:404-406; Cwirla et al. (1990) *PNAS* 87: 6378-6382; as well as U.S. Pat. Nos. 5,223,409, 5,198,346, and 5,096,815).

A wide range of techniques are known in the art for screening gene products of combinatorial libraries made by point mutations, and for screening cDNA libraries for gene products having a certain property. Such techniques will be generally adaptable for rapid screening of the gene libraries generated by the combinatorial mutagenesis of UbCE homologs. The most widely used techniques for screening large gene libraries typically comprises cloning the gene library into replicable expression vectors, transforming appropriate cells with the resulting library of vectors, and expressing the combinatorial genes under conditions in which detection of a desired activity facilitates relatively easy isolation of the vector encoding the gene whose product was detected. Each of the illustrative assays described below are amenable to high through-put analysis as necessary to screen large numbers of, for example, degenerate hUbCE sequences created by combinatorial mutagenesis techniques.

In one illustrative screening assay, the candidate hUbCE gene products are displayed on the surface of a cell or viral particle, and the ability of particular cells or viral particles to bind other components of the ubiquitin pathway, e.g. E1 or E3 proteins (e.g. E6AP or E6AP complexes), ubiquitin, or p53, via this gene product is detected in a "palming assay". For instance, the gene library can be cloned into the gene for a surface membrane protein of a bacterial cell, and the resulting fusion protein detected by panning (Ladner et al., WO 88/06630; Fuchs et al. (1991) *Bio/Technology* 9:1370-1371; and Goward et al. (1992) *TIBS* 18:136-140). In a similar fashion, fluorescently labeled molecules which bind hUbCE can be used to score for potentially functional hUbCE homologs. Cells can be visually inspected and separated under a fluorescence microscope, or, where the morphology of the cell permits, separated by a fluorescence-activated cell sorter.

In an alternate embodiment, the gene library is expressed as a fusion protein on the surface of a viral particle. For

instance, in the filamentous phage system, foreign peptide sequences can be expressed on the surface of infectious phage, thereby conferring two significant benefits. First, since these phage can be applied to affinity matrices at very high concentrations, a large number of phage can be screened at one time. Second, since each infectious phage displays the combinatorial gene product on its surface, if a particular phage is recovered from an affinity matrix in low yield, the phage can be amplified by another round of infection. The group of almost identical *E. coli* filamentous phages M13, fd, and f1 are most often used in phage display libraries, as either of the phage gIII or gVIII coat proteins can be used to generate fusion proteins without disrupting the ultimate packaging of the viral particle (Ladner et al. PCT publication WO 90/02909; Garrard et al., PCT publication WO 92/09690; Marks et al. (1992) *J. Biol. Chem.* 267:16007-16010; Griffiths et al. (1993) *EMBO J* 12:725-734; Clackson et al. (1991) *Nature* 352:624-628; and Barbas et al. (1992) *PNAS* 89:4457-4461).

In an illustrative embodiment, the recombinant phage antibody system (RPAS, Pharmacia Catalog number 27-9400-01) can be easily modified for use in expressing and screening hUbCE combinatorial libraries. For instance, the pCANTAB 5 phagemid of the RPAS kit contains the gene which encodes the phage gIII coat protein. The hUbCE combinatorial gene library can be cloned into the phagemid adjacent to the gIII signal sequence such that it will be expressed as a gIII fusion protein. After ligation, the phagemid is used to transform competent *E. coli* TGI cells. Transformed cells are subsequently infected with M13KO7 helper phage to rescue the phagemid and its candidate hUbCE gene insert. The resulting recombinant phage contain phagemid DNA encoding a specific candidate hUbCE, and display one or more copies of the corresponding fusion coat protein. The phage-displayed candidate hUbCE which are capable of binding a particular target protein, such as an E1 enzyme, an E3 protein (i.e. E6 or E6-AP), or p53, are selected or enriched by panning. For instance, the phage library can be panned on glutathione immobilized p53-GST fusion proteins or E6-GST or E6-AP-GST fusion proteins (described, for example, in U.S. patent application Ser. No. 08/176,937), and unbound phage washed away from the cells. The bound phage is then isolated, and if the recombinant phage express at least one copy of the wild type gIII coat protein, they will retain their ability to infect *E. coli*. Thus, successive rounds of reinfection of *E. coli*, and panning can be employed to greatly enrich for hUbCE homologs that retain some ability to interact with normal targets of the wild-type hUbCE, and which can then be screened for further biological activities in order to differentiate agonists and antagonists. In an exemplary embodiment, by use of two or more target proteins in sequential panning steps, the phage display library can be used to isolate hUbCE homologs which are candidate antagonists of the normal cellular function of the naturally occurring hUbCE. For instance, isolating from the library those variants which retain the ability to bind, for example, either the papillomavirus E6 protein or the cellular E6-AP protein, but which are unable to bind p53, provides a set of hUbCE homologs some of which may be capable of antagonizing the ability of the wild-type hUbCE to mediate ubiquitination of p53.

In yet another illustrative embodiment, the p53-dependent reporter construct described in the 08/176,937 application can be used to identify antagonists through their ability to enhance expression of the reporter gene by inhibiting the degradation of p53 wild-type hUbCE. Thus, a combinatorial

library can be screened by a detecting expression of the reporter gene, and appropriate clones isolated for further manipulation.

Other forms of mutagenesis can also be utilized to generate a combinatorial library from the subject UbCE proteins. For example, hUbCE homologs (both agonist and antagonist forms) can be generated and isolated from a library by screening using, for example, alanine scanning mutagenesis and the like (Ruf et al. (1994) *Biochemistry* 33:1565-1572; Wang et al. (1994) *J. Biol. Chem.* 269:3095-3099; Balint et al. (1993) *Gene* 137:109-118; Grodberg et al. (1993) *Eur. J. Biochem.* 218:597-601; Nagashima et al. (1993) *J. Biol. Chem.* 268:2888-2892; Lowman et al. (1991) *Biochemistry* 30:10832-10838; and Cunningham et al. (1989) *Science* 244:1081-1085), by linker scanning mutagenesis (Gustin et al. (1993) *Virology* 193:653-660; Brown et al. (1992) *Mol. Cell Biol.* 12:2644-2652; McKnight et al. (1982) *Science* 232:316); by saturation mutagenesis (Meyers et al. (1986) *Science* 232:613); by PCR mutagenesis (Leung et al. (1989) *Method Cell Mol Biol* 1:11-19); or by random mutagenesis (Miller et al. (1992) *A Short Course in Bacterial Genetics*, CSHL Press, Cold Spring Harbor, N.Y.; and Greener et al. (1994) *Strategies in Mol Biol* 7:32-34).

An important goal of the present invention is to provide reduction of the UbCE proteins to small functional units that can be ultimately used to generate UbCE mimetics, e.g. peptide or non-peptide agents, which are able to disrupt binding of UbCE with other cellular and/or viral proteins. Thus, such mutagenic techniques as described herein are particularly useful to map the determinants of the hUbCE protein which participate in protein-protein interactions involved in, for example, binding of the subject hUbCE to other proteins of the ubiquitin-conjugating system (both cellular and viral), as well as the target protein itself (e.g. p53). To illustrate, the critical residues of hUbCE involved in molecular recognition of E6 and/or E6-AP can be determined and used to generate hUbCE-derived peptidomimetics which competitively inhibit hUbCE binding. By employing, for example, scanning mutagenesis to map the amino acid residues of hUbCE involved in binding E6AP, peptidomimetic compounds can be generated which mimic those residues in binding to E6AP, and which therefore can inhibit binding of the hUbCE to E6AP and interfere with the function of E6AP in regulating the cellular half-life of p53. For instance, non-hydrolyzable peptide analogs of such residues can be generated using benzodiazepine (e.g., see Freidinger et al. in *Peptides: Chemistry and Biology*, G. R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), azepine (e.g., see Huffman et al. in *Peptides: Chemistry and Biology*, G. R. Marshall ed., ESCOM Publisher Leiden, Netherlands, 1988), substituted gamma lactam rings (Garvey et al. in *Peptides. Chemistry and Biology*, G. R. Marshall ed., ESCOM Publisher: Leiden, Netherlands, 1988), keto-methylene pseudopeptides (Ewenson et al. (1986) *J Med Chem* 29:295; and Ewenson et al. in *Peptides: Structure and Function* (Proceedings of the 9th American Peptide Symposium) Pierce Chemical Co. Rockland, Ill., 1985), β -turn dipeptide cores (Nagai et al. (1985) *Tetrahedron Lett* 26:647; and Sato et al. (1986) *J Chem Soc Perkin Trans* 1:1231), and β -aminoalcohols (Gordon et al. (1985) *Biochem Biophys Res Commun* 126:419; and Dann et al. (1986) *Biochem Biophys Res Commun* 134:71). Such peptidomimetics can serve as drugs which prevent the action of hUbCE in the destruction of, for example, p53. Furthermore, such data concerning protein-protein interactions can be used in conjunction with the molecular model of hUbCE

described below for rational design of mimetics of this interaction. In like manner, peptidomimetics of caUbCE and spUbCE can be derived which may be useful in, for example, the generation of anti-mycotic agents.

Another aspect of the invention pertains to an antibody specifically reactive with the subject UbCE proteins. For example, by using immunogens derived from the hUbCE protein of the present invention, anti-protein/anti-peptide antisera or monoclonal antibodies can be made by standard protocols (See, for example, *Antibodies: A Laboratory Manual* ed. by Harlow and Lane (Cold Spring Harbor Press: 1988)). A mammal such as a mouse, a hamster or rabbit can be immunized with an immunogenic form of the peptide (e.g.) the whole hUbCE protein or an antigenic fragment which is capable of eliciting an antibody response). Techniques for conferring immunogenicity on a protein or peptide include conjugation to carriers or other techniques well known in the art. An immunogenic portion of the subject UbCE protein can be administered in the presence of adjuvant. The progress of immunization can be monitored by detection of antibody titers in plasma or serum. Standard ELISA or other immunoassays can be used with the immunogen as an antigen to assess the levels of antibodies. In a preferred embodiment, the subject antibodies are immunospecific for hUbCE antigenic determinants, e.g. antigenic determinants of a protein represented by SEQ ID No. 2 or a closely related human or non-human mammalian homolog (e.g. 90 percent homologous to SEQ ID No. 2, preferably at least 95 percent homologous and more preferably at least 97 percent homologous to SEQ ID No. 2). In yet a further preferred embodiment of the present invention, the anti-hUbCE antibodies does not substantially cross react with a protein which is: e.g. less than 90 percent homologous with SEQ ID No. 2; e.g. less than 95 percent homologous with SEQ ID No. 2; e.g. less than 98-99 percent homologous with SEQ ID No. 2. By "does not substantially cross-react", it is meant that: the antibody has a binding affinity for a non-homologous E2 enzyme which is less than 10 percent, more preferably less than 5 percent, and most preferably less than about 1-2 percent of the binding affinity of that antibody for the protein of SEQ ID No. 2; the antibody does not specifically bind a protein which is non-homologous to SEQ ID No. 2. Preferred antibodies against the subject caUbCE and spUbCE proteins have similar criteria, e.g., antibodies specific for caUbCE or spUbCE do not specifically bind proteins which do not share high sequence homology with SEQ ID No. 4 or 6, respectively.

Following immunization, anti-UbCE antisera can be obtained and, if desired, polyclonal anti-UbCE antibodies isolated from the serum. To produce monoclonal antibodies, antibody producing cells (lymphocytes) can be harvested from an immunized animal and fused by standard somatic cell fusion procedures with immortalizing cells such as myeloma cells to yield hybridoma cells. Such techniques are well known in the art, and include, for example, the hybridoma technique (originally developed by Kohler and Milstein, (1975) *Nature*, 256:495-497), the human B cell hybridoma technique (Kozbar et al., (1983) *Immunology Today*: 4:72), and the EBV-hybridoma technique to produce human monoclonal antibodies (Cole et al., (1985) *Monoclonal Antibodies and Cancer Therapy*, Alan R. Liss, Inc. pp. 77-96). Hybridoma cells can be screened immunochemically for production of antibodies specifically reactive with the subject UbCE protein and monoclonal antibodies isolated from a culture comprising such hybridoma cells.

The term antibody as used herein is intended to include fragments thereof which are also specifically reactive with

the UbCE proteins of the present invention. Antibodies can be fragmented using conventional techniques and the fragments screened for utility in the same manner as described above for whole antibodies. For example, F(ab')₂ fragments can be generated by treating antibody with pepsin. The resulting F(ab')₂ fragment can be treated to reduce disulfide bridges to produce Fab' fragments. The antibody of the present invention is further intended to include bispecific and chimeric molecules having an anti-UbCE portion.

Both monoclonal and polyclonal antibodies (Ab) directed against the subject ubiquitin conjugating enzymes, and antibody fragments such as Fab' and F(ab')₂, can be used as specialty chemicals to block the action of the enzyme and allow the study of, for example, the cell cycle or cell proliferation when the subject UbCE is inhibited, e.g. by microinjection of anti-UbCE antibodies.

Antibodies which specifically bind hUbCE epitopes can also be used in immunohistochemical staining of tissue samples in order to evaluate the abundance and pattern of expression of hUbCE. Anti-hUbCE antibodies can be used diagnostically in immuno-precipitation and immunoblotting to detect and evaluate hUbCE levels in tissue or bodily fluid as part of a clinical testing procedure. For instance, such measurements can be useful in predictive valuations of the onset or progression of tumors. Likewise, the ability to monitor hUbCE levels in an individual can allow determination of the efficacy of a given treatment regimen for an individual afflicted with such a disorder. The level of hUbCE can be measured in cells isolated from bodily fluid, such as in samples of cerebral spinal fluid or blood, or can be measured in tissue, such as produced by biopsy. Diagnostic assays using anti-hUbCE antibodies can include, for example, immunoassays designed to aid in early diagnosis of a neoplastic or hyperplastic disorder, e.g. the presence of cancerous cells in the sample, e.g. to detect cells in which a lesion of the hUbCE gene has occurred.

Another application of anti-UbCE antibodies is in the immunological screening of cDNA libraries constructed in expression vectors, such as λgt11, λt18-23, λZAP, and λORF8. Messenger libraries of this type, having coding sequences inserted in the correct reading frame and orientation, can produce fusion proteins. For instance, λgt11 will produce fusion proteins whose amino termini consist of β-galactosidase amino acid sequences and whose carboxy termini consist of a foreign polypeptide. Antigenic epitopes of UbCE can then be detected with antibodies, as, for example, reacting nitrocellulose filters lifted from infected plates with anti-UbCE antibodies. Phage, scored by this assay, can then be isolated from the infected plate. Thus, the presence of hUbCE homologs can be detected and cloned from other human sources, i.e. to identified other closely homologous human isoforms, as well as to identify hUbCE homologs in other mammals.

Moreover, the nucleotide sequence determined from the cloning of subject hUbCE from a human cell line will further allow for the generation of probes designed for use in identifying hUbCE homologs in other human cell-types, particularly cancer or other transformed or immortalized cells, as well as hUbCE homologs from other non-human mammals. Probes based on the yeast UbCE sequences, caUbCE and spUbCE can be generated and used to identify and phenotype mycotic infections.

In addition, nucleotide probes can be generated from the cloned sequence of the hUbCE protein, which allow for histological screening of intact tissue and tissue samples for the presence of hUbCE mRNA. Similar to the diagnostic uses of anti-hUbCE antibodies, the use of probes directed to

hUbCE mRNA, or to genomic hUbCE sequences, can be used for both predictive and therapeutic evaluation of allelic mutations which might be manifest in, for example, neoplastic or hyperplastic disorders (e.g. unwanted cell growth). Used in conjunction with anti-hUbCE antibody immunoassays, the nucleotide probes can help facilitate the determination of the molecular basis for a developmental disorder which may involve some abnormality associated with expression (or lack thereof) of an hUbCE protein. For instance, variation in hUbCE synthesis can be differentiated from a mutation in the hUbCE coding sequence.

For example, the present method provides a method for determining if a subject is at risk for a disorder characterized by unwanted cell proliferation. In preferred embodiments, the subject method can be generally characterized as comprising detecting, in a tissue of a subject (e.g. a human patient), the presence or absence of a genetic lesion characterized by at least one of (i) a mutation of a gene encoding hUbCE or (ii) the mis-expression of the hUbCE gene. To illustrate, such genetic lesions can be detected by ascertaining the existence of at least one of (i) a deletion of one or more nucleotides from the hUbCE gene, (ii) an addition of one or more nucleotides to the hUbCE gene, (iii) a substitution of one or more nucleotides of the hUbCE gene, (iv) a gross chromosomal rearrangement of the hUbCE gene, (v) a gross alteration in the level of a messenger RNA transcript of the hUbCE gene, (vi) the presence of a non-wild type splicing pattern of a messenger RNA transcript of the hUbCE gene, and (vii) a non-wild type level of the hUbCE protein. In one aspect of the invention there is provided a probe/primer comprising an oligonucleotide containing a region of nucleotide sequence which is capable of hybridizing to a sense or antisense sequence of SEQ ID No: 1 or naturally occurring mutants thereof; or 5 or 3 flanking sequences or intronic sequences naturally associated with the hUbCE gene. The probe is exposed to nucleic acid of a tissue sample; and the hybridization of the probe to the sample nucleic acid is detected. In certain embodiments, detection of the lesion comprises utilizing the probe/primer in, for example, a polymerase chain reaction (PCR) (see, e.g. U.S. Pat. Nos. 4,683,195 and 4,683,202), or, alternatively, in a ligation chain reaction (LCR) (see, e.g., Landegran et al. (1988) *Science* 241:1077-1080; and Nakazawa et al. (1994) *PNAS* 91:360-364), the later of which can be particularly useful for detecting even point mutations in the hUbCE gene. Alternatively, or additionally, the level of hUbCE protein can be detected in an immunoassay.

Also, the use of anti-sense techniques (e.g. microinjection of antisense molecules, or transfection with plasmids whose transcripts are anti-sense with regard to, e.g. UbCE mRNA) can be used to investigate the role of UbCE in the cell cycle and cell proliferation, by inhibiting endogenous UbCE production. Such techniques can be utilized in cell culture, but can also be used in the creation of transgenic animals

Another aspect of the present invention concerns transgenic animals, e.g. as animal models for developmental and proliferative diseases, which are comprised of cells (of that animal) which contain a transgene of the present invention and which preferably (though optionally) express the subject UbCE in one or more cells in the animal. In preferred embodiments, the expression of the transgene is restricted to specific subsets of cells, tissues or developmental stages utilizing, for example, cis-acting sequences that control expression in the desired pattern. In the present invention, such mosaic expression of the subject UbCE proteins can be essential for many forms of lineage analysis and can additionally provide a means to assess the effects of UbCE

mutations or overexpression that might grossly alter development in small patches of tissue within an otherwise normal embryo. Toward this end, tissue-specific regulatory sequences and conditional regulatory sequences can be used to control expression of the transgene in certain spatial patterns. Moreover, temporal patterns of expression can be provided by, for example, conditional recombination systems or prokaryotic transcriptional regulatory sequences.

Genetic techniques which allow for the expression of transgenes can be regulated via site-specific genetic manipulation in vivo are known to those skilled in the art. For instance, Genetic systems are available which allow for the regulated expression of a recombinase that catalyzes the genetic recombination a target sequence. As used herein, the phrase "target sequence" refers to a nucleotide sequence that is genetically recombined by a recombinase. The target sequence is flanked by recombinase recognition sequences and is generally either excised or inverted in cells expressing recombinase activity. Recombinase catalyzed recombination events can be designed such that recombination of the target sequence results in either the activation or repression of expression of the subject receptor. For example, excision of a target sequence which interferes with the expression of the receptor can be designed to activate expression of that protein. This interference with expression of the subject protein can result from a variety of mechanisms, such as spatial separation of the UbCE gene from the promoter element or an internal stop codon. Moreover, the transgene can be made wherein the coding sequence of the UbCE gene is flanked by recombinase recognition sequences and is initially transfected into cells in a 3' to 5' orientation with respect to the promoter element. In such an instance, inversion of the target sequence will reorient the subject UbCE gene by placing the 5' end of the coding sequence in an orientation with respect to the promoter element which allow for promoter driven transcriptional activation.

In an illustrative embodiment, either the creloxP recombinase system of bacteriophage P1 (Lakso et al. (1992) *PNAS* 89:6232-6236; Orban et al. (1992) *PNAS* 89:6861-6865) or the FLP recombinase system of *Saccharomyces cerevisiae* (O'Gorman et al. (1991) *Science* 251:1351-1355; PCT publication WO 92/15694) can be used to generate in vivo site-specific genetic recombination systems. Cre recombinase catalyzes the site-specific recombination of an intervening target sequence located between loxP sequences. loxP sequences are 34 base pair nucleotide repeat sequences to which the Cre recombinase binds and are required for Cre recombinase mediated genetic recombination. The orientation of loxP sequences determines whether the intervening target sequence is excised or inverted when Cre recombinase is present (Abremski et al (1984) *J. Biol. Chem.* 259:1509-1514); catalyzing the excision of the target sequence when the loxP sequences are oriented as direct repeats and catalyzes inversion of the target sequence when loxP sequences are oriented as inverted repeats.

Accordingly, genetic recombination of the target sequence is dependent on expression of the Cre recombinase. Expression of the recombinase can be regulated by promoter elements which are subject to regulatory control, e.g., tissue-specific, developmental stage-specific, inducible or repressible by externally added agents. This regulated control will result in genetic recombination of the target sequence only in cells where recombinase expression is mediated by the promoter element. Thus, the activation of expression of the recombinant UbCE gene can be regulated via regulation of recombinase expression.

Use of the these recombinase system to regulate expression of, for example, a dominant negative UbCE gene, such as the Cys85Ser mutant or an antisense gene, requires the construction of a transgenic animal containing transgenes encoding both the Cre recombinase and the subject gene. Animals containing both the Cre recombinase and the UbCE genes can be provided through the construction of "double" transgenic animals. A convenient method for providing such animals is to mate two transgenic animals each containing a transgene, e.g., the UbCE gene and recombinase gene.

One advantage derived from initially constructing transgenic animals containing a UbCE transgene in a recombinase-mediated expressible format derives from the likelihood that the subject UbCE protein, whether antagonistic or agonistic, will be deleterious upon expression in the transgenic animal. In such an instance, a founder population, in which the subject transgene is silent in all tissues can be propagated and maintained. Individuals of this founder population can be crossed with animals expressing the recombinase in, for example, one or more tissues. Thus, the creation of a founder population in which the UbCE transgene is silent will allow the study of, for example, the role of the p53 checkpoint in tissue or at developmental stages which can confer, for example, a lethal phenotype.

Similar conditional transgenes can be provided using prokaryotic promoter sequences which require prokaryotic proteins to be simultaneously expressed in order to facilitate expression of the transgene. Exemplary promoters and the corresponding trans-activating prokaryotic proteins are given in U.S. Pat. No. 4,833,080. Moreover, expression of the conditional transgenes can be induced by gene therapy-like methods wherein a gene encoding the trans-activating protein, e.g. a recombinase or a prokaryotic protein, is delivered to the tissue and caused to be expressed, such as in a cell-type specific manner. By this method, the transgene could remain silent into adulthood until "turned on" by the introduction of the trans-activator.

Methods of making knock-out or disruption transgenic animals are also generally known. See, for example, *Manipulating the Mouse Embryo*, (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1986).

Furthermore, the present invention, by making available purified and recombinant forms of the subject UbCE proteins, will allow the development of assays which can be used to screen for drugs which are either agonists or antagonists. For instance, in addition to agents which disrupt binding of the hUbCE protein to other cellular (or viral) proteins, inhibitors of the enzymatic activity of the subject hUbCE can be used to prevent transfer of ubiquitin to hUbCE and/or inhibit any downstream transfer of ubiquitin from hUbCE (e.g. to p53 or an intermediary E3 complex, e.g. E6/E6-AP). In a preferred embodiment, the hUbCE inhibitor is a mechanism based inhibitor which chemically alters the enzyme, e.g. covalently binds Cys-85, and which is a specific inhibitor of hUbCE, e.g. has an inhibition constant 10-fold, 100-fold, or more preferably, 1000-fold different for human E2 enzymes other than the subject hUbCE protein. Inhibitor specificity can be improved, for example, by utilizing specificity subsites of the hUbCE enzyme involved in interactions between hUbCE and p53 or hUbCE and E1, which are unique to one of those complexes relative to other human E2 enzymes.

Assays for the measurement of ubiquitination are disclosed in U.S. patent application Ser. No. 08/176,937, filed on Jan. 4, 1994, and herein incorporated by reference. Such assays can be used in conjunction with the subject hUbCE protein to generate a ubiquitin-conjugating system to detect

agents able to inhibit hUbCE-mediated ubiquitination of a cellular or target protein. Such agents can be used to, for example, treat papillomavirus infected cells. Similar assay systems can be constructed for the fungal UbCEs in order to detect inhibitors which may serve as anti-fungal agents. In preferred embodiments, the assay system employed for identifying anti-fungal agents are run side-by-side with the analogous assay system derived with hUbCE. Differential screening assays can be used to exploit any difference in mechanism or specificity between mammalian UbCEs and yeast UbCEs (including other yeast E2 enzymes) in order to identify agents which display a statistically significant increase in specificity for inhibiting the yeast enzymes relative to the mammalian enzymes. Thus, lead compounds which act specifically on pathogens, such as fungus involved in mycotic infections, can be developed.

Additionally, the subject ubiquitin conjugating enzyme can be used to generate an interaction trap assay for subsequently detecting inhibitors of hUbCE biological activity (see, for example, U.S. Pat. No: 5,283,317; PCT publication WO94/10300; Zervos et al. (1993) *Cell* 72:223-232; Madura et al. (1993) *J Biol Chem* 268:12046-12054; Bartel et al. (1993) *Biotechniques* 14:920-924; and Iwabuchi et al. (1993) *Oncogene* 8:1693-1696) In an illustrative embodiment, *Saccharomyces cerevisiae* YPB2 cells are transformed simultaneously with a plasmid encoding a GAL4db-hUbCE fusion and with a plasmid encoding the GAL4ad domain fused to p53t or E6AP. Moreover, the strain is transformed such that the GAL4-responsive promoter drives expression of a phenotypic marker. For example, the ability to grow in the absence of histidine can depend on the expression of the HIS3 gene if it is under control of a GAL4-responsive promoter and, therefore, indicates that a functional GAL4 activator has been reconstituted through the interaction of hUbCE and p53 or E6AP. Thus, agent able to inhibit hUbCE interaction with one of these proteins will result in yeast cells unable to grow in the absence of histidine. Alternatively, the phenotypic marker can be one which provides a negative selection when expressed such that agents which disrupt the hUbCE interactions confer positive growth selection to the cells.

Another aspect of the present invention concerns three-dimensional molecular models of the subject UbCE proteins, and their use as templates for the design of agents able to inhibit at least one biological activity of the ubiquitin conjugating enzyme. An integral step to our approach to designing inhibitors of the subject ubiquitin-conjugating enzyme involves construction of computer graphics models of the ubiquitin conjugating enzyme which can be used to design pharmacophores by rational drug design. For instance, for an inhibitor to interact optimally with the subject enzyme, it will generally be desirable that it have a shape which is at least partly complimentary to that of a particular binding site of the enzyme, as for example those portions of the human ubiquitin conjugating enzyme which are involved in recognition of ubiquitin, an E1 enzyme, an E3 protein(s) such as E6 or E6AP, or a downstream target of the pathway, such as p53. Additionally, other factors, including electrostatic interactions, hydrogen bonding, hydrophobic interactions, desolvation effects, and cooperative motions of ligand and enzyme, all influence the binding effect and should be taken into account in attempts to design bioactive inhibitors.

As described in Example 4, a computer-generated molecular model of the subject enzymes can be created. In preferred embodiments, at least the C α -carbon positions of the UbCE sequence of interest are mapped to a particular

coordinate pattern, such as the coordinates for hUbCE shown in FIG. 1, by homology modeling, and the structure of the protein and velocities of each atom are calculated at a simulation temperature (T_o) at which the docking simulation is to be determined. Typically, such a protocol involves primarily the prediction of side-chain conformations in the modeled protein, while assuming a main-chain trace taken from a tertiary structure such as provided in FIG. 1. Computer programs for performing energy minimization routines are commonly used to generate molecular models. For example, both the CHARMM (Brooks et al. (1983) *J Comput Chem* 4:187-217) and AMBER (Weiner et al (1981) *J. Comput. Chem.* 106:765) algorithms handle all of the molecular system setup, force field calculation, and analysis (see also, Eisenfield et al. (1991) *Am J Physiol* 261:C376-386; Lybrand (1991) *J Pharm Belg* 46:49-54; Froimowitz (1990) *Biotechniques* 8:640-644; Burbam et al. (1990) *Proteins* 7:99-111; Pedersen (1985) *Environ Health Perspect* 61:185-190; and Kini et al. (1991) *J Biomol Struct Dyn* 9:475-488). At the heart of these programs is a set of subroutines that, given the position of every atom in the model, calculate the total potential energy of the system and the force on each atom. These programs may utilize a starting set of atomic coordinates, such as the model coordinates provided in FIG. 1, the parameters for the various terms of the potential energy function, and a description of the molecular topology (the covalent structure). Common features of such molecular modeling methods include: provisions for handling hydrogen bonds and other constraint forces; the use of periodic boundary conditions; and provisions for occasionally adjusting positions, velocities, or other parameters in order to maintain or change temperature, pressure, volume, forces of constraint, or other externally controlled conditions.

Most conventional energy minimization methods use the input data described above and the fact that the potential energy function is an explicit, differentiable function of Cartesian coordinates, to calculate the potential energy and its gradient (which gives the force on each atom) for any set of atomic positions. This information can be used to generate a new set of coordinates in an effort to reduce the total potential energy and, by repeating this process over and over, to optimize the molecular structure under a given set of external conditions. These energy minimization methods are routinely applied to molecules similar to the subject UbCE proteins as well as nucleic acids, polymers and zeolites.

In general, energy minimization methods can be carried out for a given temperature, T_i , which may be different than the docking simulation temperature, T_o . Upon energy minimization of the molecule at T_i , coordinates and velocities of all the atoms in the system are computed. Additionally, the normal modes of the system are calculated. It will be appreciated by those skilled in the art that each normal mode is a collective, periodic motion, with all parts of the system moving in phase with each other, and that the motion of the molecule is the superposition of all normal model. For a given temperature, the mean square amplitude of motion in a particular mode is inversely proportional to the effective force constant for that mode, so that the motion of the molecule will often be dominated by the low frequency vibrations.

After the molecular model has been energy minimized at T_i , the system is "heated" or "cooled" to the simulation temperature, T_o , by carrying out an equilibration run where the velocities of the atoms are scaled in a step-wise manner until the desired temperature, T_o , is reached. The system is

further equilibrated for a specified period of time until certain properties of the system, such as average kinetic energy, remain constant. The coordinates and velocities of each atom are then obtained from the equilibrated system.

Further energy minimization routines can also be carried out. For example, a second class of methods involves calculating approximate solutions to the constrained EOM for the protein. These methods use an iterative approach to solve for the Lagrange multipliers and, typically, only need a few iterations if the corrections required are small. The most popular method of this type, SHAKE (Ryckaert et al. (1977) *J Comput Phys* 23:327; and Van Gunsteren et al. (1977) *Mol Phys* 34:1311) is easy to implement and scales as $O(N)$ as the number of constraints increases. Therefore, the method is applicable to macromolecules such as the UbCE proteins of the present invention. An alternative method, RATTLE (Anderson (1983) *J Comput Phys* 52:24) is based on the velocity version of the Verlet algorithm. Like SHAKE, RATTLE is an iterative algorithm and can be used to energy minimize the model of the subject protein.

The increasing availability of biomacromolecule structures of potential pharmacophoric molecules that have been solved crystallographically has prompted the development of a variety of direct computational methods for molecular design, in which the steric and electronic properties of substrate binding sites are used to guide the design of potential inhibitors (Cohen et al. (1990) *J. Med. Cam.* 33:883-894; Kuntz et al. (1982) *J. Mol. Biol* 161:269-288; DesJarlais (1988) *J. Med. Cam.* 31:722-729; Bartlett et al. (1989) (*Spec. Publ., Roy. Soc. Chem.*) 78:182-196; Goodford et al. (1985) *J. Med. Cam.* 28:849-857; DesJarlais et al. *J Med. Cam.* 29:2149-2153). Directed methods generally fall into two categories: (1) design by analogy in which 3-D structures of known molecules (such as from a crystallographic database) are docked to the enzyme structure and scored for goodness-of-fit; and (2) de novo design, in which the ligand model is constructed piece-wise in the enzyme. The latter approach, in particular, can facilitate the development of novel molecules, uniquely designed to bind to the subject human ubiquitin-conjugating enzyme.

In an illustrative embodiment, the design of potential hUbCE inhibitors begins from the general perspective of shape complimentary for the active site and substrate specificity subsites of the enzyme, and a search algorithm is employed which is capable of scanning a database of small molecules of known three-dimensional structure for candidates which fit geometrically into the target protein site. It is not expected that the molecules found in the shape search will necessarily be leads themselves, since no evaluation of chemical interaction necessarily be made during the initial search. Rather, it is anticipated that such candidates might act as the framework for further design, providing molecular skeletons to which appropriate atomic replacements can be made. Of course, the chemical complimentary of these molecules can be evaluated, but it is expected that atom types will be changed to maximize the electrostatic, hydrogen bonding, and hydrophobic interactions with the enzyme. Most algorithms of this type provide a method for finding a wide assortment of chemical structures that are complementary to the shape of a binding site of the subject enzyme. Each of a set of small molecules from a particular data-base, such as the Cambridge Crystallographic Data Bank (CCDB) (Allen et al. (1973) *J. Chem. Doc.* 13:119), is individually docked to the binding site of the hUbCE enzyme in a number of geometrically permissible orientations with use of a docking algorithm. In a preferred embodiment, a set of computer algorithms called DOCK, can be used to charac-

terize the shape of invaginations and grooves that form the active sites and recognition surfaces of the subject enzyme (Kuntz et al. (1982) *J. Mol. Biol* 161:269–288). The program can also search a database of small molecules for templates whose shapes are complementary to particular binding sites of the enzyme (DesJarlais et al. (1988) *J Med Chem* 31:722–729). These templates normally require modification to achieve good chemical and electrostatic interactions (DesJarlais et al. (1989) *ACS Symp Ser* 413:60–69). However, the program has been shown to position accurately known cofactors for inhibitors based on shape constraints alone.

The orientations are evaluated for goodness-of-fit and the best are kept for further examination using molecular mechanics programs, such as AMBER or CHARMM. Such algorithms have previously proven successful in finding a variety of molecules that are complementary in shape to a given binding site of a receptor-enzyme, and have been shown to have several attractive features. First, such algorithms can retrieve a remarkable diversity of molecular architectures. Second, the best structures have, in previous applications to other proteins, demonstrated impressive shape complementarity over an extended surface area. Third, the overall approach appears to be quite robust with respect to small uncertainties in positioning of the candidate atoms.

Goodford (1985, *J Med Chem* 28:849–857) and Boobbyer et al. (1989, *J Med Chem* 32:1083–1094) have produced a computer program (GRID) which seeks to determine regions of high affinity for different chemical groups (termed probes) on the molecular surface of the binding site. GRID hence provides a tool for suggesting modifications to known ligands that might enhance binding. It may be anticipated that some of the sites discerned by GRID as regions of high affinity correspond to “pharmacophoric patterns” determined inferentially from a series of known ligands. As used herein, a pharmacophoric pattern is a geometric arrangement of features of the anticipated ligand that is believed to be important for binding. Attempts have been made to use pharmacophoric patterns as a search screen for novel ligands (Jakes et al. (1987) *J Mol Graph* 5:41–48; Brint et al. (1987) *J Mol Graph* 5:49–56; Jakes et al. (1986) *J Mol Graph* 4:12–20); however, the constraint of steric and “chemical” fit in the putative (and possibly unknown) receptor binding site is ignored. Goodsell and Olson (1990, *Proteins: Struct Funct Genet* 8:195–202) have used the Metropolis (simulated annealing) algorithm to dock a single known ligand into a target protein. They allow torsional flexibility in the ligand and use GRID interaction energy maps as rapid lookup tables for computing approximate interaction energies. Given the large number of degrees of freedom available to the ligand, the Metropolis algorithm is time-consuming and is unsuited to searching a candidate database of a few thousand small molecules.

Yet a further embodiment of the present invention utilizes a computer algorithm such as CLIX which searches such databases as CCDB for small molecules which can be oriented in the receptor binding site in a way that is both sterically acceptable and has a high likelihood of achieving favorable chemical interactions between the candidate molecule and the surrounding amino acid residues. The method is based on characterizing the receptor site in terms of an ensemble of favorable binding positions for different chemical groups and then searching for orientations of the candidate molecules that cause maximum spatial coincidence of individual candidate chemical groups with members of the ensemble. The current availability of computer power dic-

tates that a computer-based search for novel ligands follows a breadth-first strategy. A breadth-first strategy aims to reduce progressively the size of the potential candidate search space by the application of increasingly stringent criteria, as opposed to a depth-first strategy wherein a maximally detailed analysis of one candidate is performed before proceeding to the next. CLIX conforms to this strategy in that its analysis of binding is rudimentary—it seeks to satisfy the necessary conditions of steric fit and of having individual groups in “correct” places for bonding, without imposing the sufficient condition that favorable bonding interactions actually occur. A ranked “shortlist” of molecules, in their favored orientations, is produced which can then be examined on a molecule-by-molecule basis, using computer graphics and more sophisticated molecular modeling techniques. CLIX is also capable of suggesting changes to the substituent chemical groups of the candidate molecules that might enhance binding.

The algorithmic details of CLIX is described in Lawrence et al. (1992) *Proteins* 12:31–41, and the CLIX algorithm can be summarized as follows. The GRID program is used to determine discrete favorable interaction positions (termed target sites) in the binding site of the protein for a wide variety of representative chemical groups. For each candidate ligand in the CCDB an exhaustive attempt is made to make coincident, in a spatial sense in the binding site of the protein, a pair of the candidate’s substituent chemical groups with a pair of corresponding favorable interaction sites proposed by GRID. All possible combinations of pairs of ligand groups with pairs of GRID sites are considered during this procedure. Upon locating such coincidence, the program rotates the candidate ligand about the two pairs of groups and checks for steric hindrance and coincidence of other candidate atomic groups with appropriate target sites. Particular candidate/orientation combinations that are good geometric fits in the binding site and show sufficient coincidence of atomic groups with GRID sites are retained.

Consistent with the breadth-first strategy, this approach involves simplifying assumptions. Rigid protein and small molecule geometry is maintained throughout. As a first approximation rigid geometry is acceptable as the energy minimized coordinates of the hUbCE deduced structure, as described in Example 4, describe an energy minimum for the molecule, albeit a local one. If the surface residues of the site of interest are not involved in crystal contacts then the crystal configuration of those residues. We believe that the deduced structure described in Example 4 should reasonably mimic the mean solution configuration. Moreover, the equivalent models of caUbCE and spUbCE can be derived by the same method.

A further assumption implicit in CLIX is that the potential ligand, when introduced into the binding site of ubiquitin-conjugating enzyme, does not induce change in the protein’s stereochemistry or partial charge distribution and so alter the basis on which the GRID interaction energy maps were computed. It must also be stressed that the interaction sites predicted by GRID are used in a positional and type sense only, i.e., when a candidate atomic group is placed at a site predicted as favorable by GRID, no check is made to ensure that the bond geometry, the state of protonation, or the partial charge distribution favors a strong interaction between the protein and that group. Such detailed analysis should form part of more advanced modeling of candidates identified in the CLIX shortlist.

Yet another embodiment of a computer-assisted molecular design method for identifying inhibitors of the subject ubiquitin-conjugating enzyme comprises the de novo syn-

thesis of potential inhibitors by algorithmic connection of small molecular fragments that will exhibit the desired structural and electrostatic complementarity with the active site of the enzyme. The methodology employs a large template set of small molecules which are iteratively pieced together in a model of the UbCE active site. Each stage of ligand growth is evaluated according to a molecular mechanics-based energy function, which considers van der Waals and coulombic interactions, internal strain energy of the lengthening ligand, and desolvation of both ligand and enzyme. The search space can be managed by use of a data tree which is kept under control by pruning according to the binding criteria.

In an illustrative embodiment the search space is limited to consider only amino acids and amino acid analogs as the molecular building blocks. Such a methodology generally employs a large template set of amino acid conformations, though need not be restricted to just the 20 natural amino acids, as it can easily be extended to include other related fragments of interest to the medicinal chemist, e.g. amino acid analogs. The putative ligands that result from this construction method are peptides and peptide-like compounds rather than the small organic molecules that are typically the goal of drug design research. The appeal of the peptide building approach is not that peptides are preferable to organics as potential pharmaceutical agents, but rather that: (1) they can be generated relatively rapidly de novo; (2) their energetics can be studied by well-parameterized force field methods; (3) they are much easier to synthesize than are most organics; and (4) they can be used in a variety of ways, for peptidomimetic inhibitor design, protein-protein binding studies, and even as shape templates in the more commonly used 3D organic database search approach described above.

Such a de novo peptide design method has been incorporated in a software package called GROW (Moon et al. (1991) *Proteins* 11:314-328). In a typical design session, standard interactive graphical modeling methods are employed to define the structural environment in which GROW is to operate. For instance, environment could be the active site cleft of hUbCE, or it could be a set of features on the protein's surface to which the user wishes to bind a peptide-like molecule, e.g. a ubiquitin, p53, E6 or E6AP mimetic. The GROW program then operates to generate a set of potential ligand molecules. Interactive modeling methods then come into play again, for examination of the resulting molecules, and for selection of one or more of them for further refinement.

To illustrate, GROW operates on an atomic coordinate file generated by the user in the interactive modeling session, such as the coordinates provided in FIG. 1, or the coordinates of the active site provided in FIG. 3, plus a small fragment (e.g., an acetyl group) positioned in the active site to provide a starting point for peptide growth. These are referred to as "site" atoms and "seed" atoms, respectively. A second file provided by the user contains a number of control parameters to guide the peptide growth (Moon et al. (1991) *Proteins* 11:314-328).

The operation of the GROW algorithm is conceptually fairly simple, and is summarized in FIG. 4. GROW proceeds in an iterative fashion, to systematically attach to the seed fragment each amino acid template in a large preconstructed library of amino acid conformations. When a template has been attached, it is scored for goodness-of-fit to the receptor site, and then the next template in the library is attached to the seed. After all the templates have been tested, only the highest scoring ones are retained for the next level of growth. This procedure is repeated for the second growth

level; each library template is attached in turn to each of the bonded seed/amino acid molecules that were retained from the first step, and is then scored. Again, only the best of the bonded seed/dipeptide molecules that result are retained for the third level of growth. The growth of peptides can proceed in the N-to-C direction only, the reverse direction only or in alternating directions, depending on the initial control specifications supplied by the user. Successive growth levels therefore generate peptides that are lengthened by one residue. The procedure terminates when the user-defined peptide length has been reached, at which point the user can select from the constructed peptides those to be studied further. The resulting data provided by the GROW procedure include not only residue sequences and scores, but also atomic coordinates of the peptides, related directly to the coordinate system of the receptor site atoms.

In yet another embodiment, potential, pharmacophoric compounds can be determined using a method based on an energy minimization-quenched molecular dynamics algorithm for determining energetically favorable positions of functional groups in the binding sites of the subject ubiquitin-conjugating enzyme. The method can aid in the design of molecules that incorporate such functional groups by modification of known ligands or de novo construction.

For example, the multiple copy simultaneous search method (MCSS) described by Miranker et al. (1991) *Proteins* 11:29-34. To determine and characterize a local minima of a functional group in the forcefield of the protein, multiple copies of selected functional groups are first distributed in a binding site of interest on the UbCE protein. Energy minimization of these copies by molecular mechanics or quenched dynamics yields the distinct local minima. The neighborhood of these minima can then be explored by a grid search or by constrained minimization. In one embodiment, the MCSS method uses the classical time dependent Hartree (TDH) approximation to simultaneously minimize or quench many identical groups in the forcefield of the protein.

Implementation of the MCSS algorithm requires a choice of functional groups and a molecular mechanics model for each of them. Groups must be simple enough to be easily characterized and manipulated (3-6 atoms, few or no dihedral degrees of freedom), yet complex enough to approximate the steric and electrostatic interactions that the functional group would have in binding to the site of interest in the UbCE protein. A preferred set is, for example, one in which most organic molecules can be described as a collection of such groups (*Patai's Guide to the Chemistry of Functional Groups*, ed. S. Patai (New York: John Wiley, and Sons, (1989)). This includes fragments such as acetonitrile, methanol, acetate, methyl ammonium, dimethyl ether, methane, and acetaldehyde.

Determination of the local energy minima in the binding site requires that many starting positions be sampled. This can be achieved by distributing, for example, 1,000-5,000 groups at random inside a sphere centered on the binding site; only the space not occupied by the protein needs to be considered. If the interaction energy of a particular group at a certain location with the protein is more positive than a given cut-off (e.g. 5.0 kcal/mole) the group is discarded from that site. Given the set of starting positions all the fragments are minimized simultaneously by use of the TDH approximation (Elber et al. (1990) *J Am Chem Soc* 112:9161-9175). In this method, the forces on each fragment consist of its internal forces and those due to the protein. The essential element of this method is that the interactions between the fragments are omitted and the forces on the protein are

normalized to those due to a single fragment. In this way simultaneous minimization or dynamics of any number of functional groups in the field of a single protein can be performed.

Minimization is performed successively on subsets of, eg. 100, of the randomly placed groups. After a certain number of step intervals, such as 1,000 intervals, the results can be examined to eliminate groups converging to the same minimum. This process is repeated until minimization is complete (e.g. RMS gradient of 0.01 kcal/mole/Å). Thus the resulting energy minimized set of molecules comprises what amounts to a set of disconnected fragments in three dimensions representing potential pharmacophores.

The next step then is to connect the pharmacophoric pieces with spacers assembled from small chemical entities (atoms, chains, or ring moieties). In a preferred embodiment, each of the disconnected can be linked in space to generate a single molecule using such computer programs as, for example, NEWLEAD (Tschinke et al. (1993) *J Med Chem* 36:3863,3870). The procedure adopted by NEWLEAD executes the following sequence of commands (1) connect two isolated moieties, (2) retain the intermediate solutions for further processing, (3) repeat the above steps for each of the intermediate solutions until no disconnected units are found, and (4) output the final solutions, each of which is single molecule. Such a program can use for example, three types of spacers: library spacers, single-atom spacers, and fuse-ring spacers. The library spacers are optimized structures of small molecules such as ethylene, benzene and methylamide. The output produced by programs such as NEWLEAD consist of a set of molecules containing the original fragments now connected by spacers. The atoms belonging to the input fragments maintain their original orientations in space. The molecules are chemically plausible because of the simple makeup of the spacers and functional groups, and energetically acceptable because of the rejection of solutions with van-der Waals radii violations.

In one embodiment of the invention, the target regulatory protein is the tumor suppressor p53, and any one of the above assays or molecular modeling protocols is used to identify inhibitors of ubiquitin-mediated destruction of p53, such as by disrupting interaction of hUbCE with p53, or interactions between hUbCE and other proteins of the ubiquitin system such as E6 or E6AP, or alternatively, by mechanistically inhibiting the enzymatic activity of the enzyme. Many lines of evidence point to the importance of p53 in human carcinogenesis. For instance, mutations within the p53 gene are the most frequent genetic aberration thus far associated with human cancer. Although p53 can block the progression of the cell cycle when artificially expressed at high levels, it appears to be dispensable for normal development. Thus, for mice containing homozygous deletions and humans harboring germline mutations of p53, development is normal and p53 protein is expressed at very low levels in most cell types. Emerging evidence, however, suggests that p53 is a checkpoint protein that plays an important role in sensing DNA damage or regulating cellular response to stress. Under normal conditions, p53 is an unstable protein and is present at very low levels in the cell, and the level of p53 in a cell appears to be controlled at least in part by degradation involving the ubiquitin system and, based on data presented herein, is likely to be mediated by the subject hUbCE. Treating cells with UV light or X rays dramatically reduces the rate of p53 degradation, leading to a rapid increase in its concentration in the cell and presumably inducing the transcription of genes that block passage

through the restriction point. However, while normal cell lines irradiated in G₁ fail to enter S phase, many tumor lines do not. In fact, there is a perfect correlation between cell lines that lack this feedback control and cells that have mutations in the p53 gene. These mutations are of two sorts: recessive mutations that inactivate the gene, and dominant mutations that produce abnormal proteins. An inhibitor developed using the subject hUbCE in a ubiquitin-conjugating assay or by rational drug design could subsequently be used therapeutically to enhance the function of the p53 checkpoint by increasing the steady state concentration of p53 in the treated cell. Given that elevated levels of wild-type p53 protein can lead to apoptosis in a variety of transformed cell types (Yonish-Rouach et al. (1991) *Nature* 352:345-347; Shaw et al. *PNAS* 89:4495-4499; and Caelles et al. (1994) *Nature* 370:220-223), inhibitors of hUbCE-mediated degradation of p53 may be attractive therapeutic agents not only in cervical cancer, but also other cancer types, by increasing the fortitude of the checkpoint in transformed cells which contain wild-type p53, or by offsetting a diminishment in p53 activity by increasing the level of (mutant) p53. Moreover, such agents can also be used prophylactically in normal cells to increase p53 levels and thereby enhance the protection against DNA damaging agents when it is known that exposure to damaging agents, such as radiation, is imminent.

Moreover, the oncogenic activity of certain viruses, such as the simian virus 40 (SV40), the adenovirus type 5 (Ad5), and the high human papilloma virus types 16 and 18 (HPV16 and HPV18), has been correlated with the virus' ability to interact with and inactivate the cellular p53 protein. In the instance of the high-risk papilloma viruses, the association of the viral oncoprotein E6 with p53 leads to the specific ubiquitination and degradation of p53. This has suggested a model in which E6 immortalizes cells by deregulating cell growth control through the elimination of the p53 tumor suppressor protein. This model accounts for the observations that p53 levels are very low in HPV-immortalized cells and that the half-life of p53 in HPV16-immortalized keratinocytes is shorter than in primary keratinocytes. Thus, the present invention can be employed in the identification of an agent that can block the ubiquitin dependent degradation of p53 as mediated by E6, and thereby block proliferation of HPV-transformed cells.

The subject human ubiquitin conjugating enzyme is likely to be involved in altering the activity of other cellular proteins, particularly proteins which seem to have short half-lives, and the present invention contemplates the use of hUbCE inhibitors, including antagonistic forms of the hUbCE protein, to inhibit the ubiquitination of other cellular proteins by hUbCE. For example, in another embodiment, the regulatory protein ubiquitinated by hUbCE is the myc oncoprotein. The myc regulatory protein is activated by translocation or mutation in many B-cell lymphomas or by amplification in tumor types, such as small cell lung cancer and breast cancer. The c-myc gene is the cellular homolog of the viral oncogene v-myc, which is found in a number of avian and feline retroviruses which induce leukemia and carcinomas. Myc has been implicated in the control of normal cell proliferation by many studies. In particular, it is one of the immediate early growth response genes that are rapidly induced in quiescent cells upon mitogenic induction, suggesting that it plays some role in mediating the transition from quiescence to proliferation. However, increased levels of myc itself is not sufficient to cause proliferation. In fact, in normal cells the opposite happens and the cell undergoes apoptosis. Therefore, inhibitors identified in the present

assay can be used to effectively induce apoptosis in cells which do not normally overexpress myc. For example, specific delivery of these agents to lymphocytes can be used to inhibit proliferation of B- and/or T-cells in order to induce clonal deletion and generate tolerance to particular antigens.

In tumor cells, on the other hand, elevated or deregulated expression of c-myc is so widespread as to suggest a critical role for myc gene activation in multi-stage carcinomas (Field et al. (1990) *Anticancer Res* 10:1-22; and Spencer et al. (1991) *Adv Cancer Res* 56:1-48). However, such over-expression of myc in these cells is typically believed to be accompanied by expression of other cellular proteins, such as bcl-2. Interestingly, however, almost all tumor cells tested that overexpress myc readily undergo apoptosis in the presence of cytotoxic and growth-inhibitory drugs (Cotter et al. (1990) *Anticancer Res* 10:1153-1159; and Lennon et al. (1990) *Biochem Soc Trans* 18:343-345). Therefore, inhibitors of the ubiquitin-mediated degradation of myc can be used to further deregulate the expression of myc in order to render the cells even more sensitive to a chemotherapeutic treatment, or to possibly upset the careful balance of the transformed cell and cause apoptosis to occur even in the absence of a second cytotoxic drug.

The regulation of cyclin by ubiquitination is yet another therapeutic target which may implicate hUbCE inhibitors. Cyclin degradation is a key step governing exit from mitosis and progression into the next cell-cycle. For example, the transition from metaphase to anaphase which marks the end of mitosis is induced by the degradation of cyclin by a ubiquitin-mediated pathway, which in turn leads to the inactivation of cyclin-dependent kinases (cdk) operational at that cycle-cycle stage. As cells enter interphase, cyclin degradation ceases, cyclin accumulates and, as a result of a complex series of post-translational modifications, cyclin/cdk complexes are activated as kinases which drive the cell through mitosis. Cyclin degradation is thus one of the crucial events in exiting mitosis. Indeed, cyclin mutants that retain the ability to activate the cdk complexes, but which cannot be degraded, arrest the cell-cycle in mitosis. Similar cyclin-dependence exists at other points of the cell-cycle as well. Thus, inhibitors of ubiquitin-mediated degradation of a cyclin (such as where the cyclin is chosen from cyclin A, B, C, D1, D2, D3, E or F) can be used as antiproliferative agents.

Yet a further possible substrate of the subject hUbCE is the fos oncogene product, which can undergo ubiquitin-mediated degradation in a cell and has been implicated in neoplastic transformation as well as in mediating the action of a variety of extracellular stimuli. The control of gene expression by c-fos is believed to play a critical role in cellular proliferation and developmental responses, and alterations in the normal pattern of c-fos can lead to oncogenesis. Given the prominence of c-fos as an early response gene, apparent over-expression and prolonged lifetime of c-fos, as may be caused by an inhibitor of the ubiquitin-mediated degradation of c-fos, might sufficiently unbalance the cell-cycle and cause cell death. Alternatively, such inhibitors can be used to mimic the effects of an external stimulus on the cell, such as treatment with a cytokine.

Exemplification

The invention now being generally described, it will be more readily understood by reference to the following examples which are included merely for purposes of illustration of certain aspects and embodiments of the present invention, and are not intended to limit the invention.

We have defined the biochemical roles of hUbCE and E6AP in the E6 stimulated ubiquitination of p53 in vitro and

have shown that inhibition of these enzymes in vivo can lead to an inhibition of E6-stimulated p53 degradation. As described in the examples below, inhibition of hUbCE and E6AP enzyme function in vivo causes an inhibition of E6-stimulated p53 degradation. The level of inhibition achieved in the micro-injection experiments in Example 3 was 25-30%. This may be a consequence of not every injected cell achieving high level expression of the injected construct, a phenomenon we have noted before in many different systems. It may also suggest that there is some redundancy in the cellular ubiquitin conjugation machinery, or that the intracellular concentrations of E1, hUbCE and E6AP are not rate-limiting for p53 degradation in the cell line used. All of our data suggest that E6 is absolutely required for ubiquitination of p53 in our in vitro and in vivo assay systems. We are currently investigating the possibility that hUbCE and E6AP are involved in the normal turnover of p53, with the possible involvement of an, as yet, unidentified cellular E6 homolog.

EXAMPLE 1

Cloning and Expression of a Novel Human Ubiquitin-conjugating Enzyme

The cDNA encoding the human ubiquitin-conjugating enzyme of the present invention was cloned from HeLa cells (ATCC CCL2). Briefly, polyadenylated RNA was isolated from cultured HeLa cells and first strand cDNA was prepared following standard protocols (c.f., Chomczynski U.S. Pat. No. 4,843,155; and Sambrook et al. *Molecular Cloning: A Laboratory Manual*, CSHL Press, Cold Spring Harbor, N.Y. (1989)). Using the nested PCR primer sets 5'-(GC)₃AAGCTTTAYGARGCGGWWGGWGTYYTTYTT-3' (SEQ ID No. 8), 5'-(GC)₃GAATTCACNGCRTAYTTYTTNGTCCCAYTC-3' (SEQ ID No. 9) and 5'-(GC)₃AAGCTTCCNGTNGGNGAYTTRTTYCAYTGGCA-3' (SEQ ID No. 10), 5-(GC)₃GAATTCATNGTNARNGCNGGCGACCA-3' (SEQ ID No. 11), which also provided convenient restriction sites in the PCR products, the coding sequences for the hUbCE gene was amplified from the HeLa cDNA library, and a HindIII-EcoRI fragment therefrom was subsequently ligated into a pBluescript II KS+ phagemid (pKS+ Stratagene catalog no. 212207) for further manipulation. The resulting pKS-hUbCE construct was amplified in XL1-Blue Cells (Stratagene Catalog no. 260268), and double stranded construct purified. The nucleic acid sequence determined for the hUbCE clone is represented in SEQ ID NO. 1, and the corresponding deduced amino acid sequence is provided in SEQ ID No. 2.

The hUbCE gene was subsequently sub-cloned from pKS+ into other expression vectors to generate gene constructs for producing the recombinant hUbCE protein in either bacterial or insect cells. In some instances, the recombinant hUbCE was provided with exogenous sequences to produce fusion proteins where the additional sequences of the fusion protein facilitate its purification. For example, after further amplification, the pKS-E2 construct was cut with XhoI and EcoRI, and the fragment containing the hUbCE coding sequence sub-cloned into a pGEX vector (Pharmacia catalog no. PGEX-4T) previously digested with SallI and EcoRI. The resulting pGEX-hUbCE construct encoded a glutathione-S-transferase (GST)/hUbCE fusion (Smith et al. (1988) *Gene* 67:31-40). The pGEX construct was introduced into *E. coli* by transformation, and the transformants grown in liquid media (LB) in the presence of IPTG. Purification of GST/hUbCE fusion protein was by

standard protocols (*Current Protocols in Molecular Biology*, eds. Ausubel et al. (NY:John Wiley & Sons, 1991); Pharmacia instruction booklet (for catalog no. 27-4570)) using a glutathione-sepharose column (Pharmacia catalog no. 27-4570). Treatment with thrombin removed the GST domain from the fusion protein.

Alternatively, the hUbCE coding sequence was excised from the pKS-hUbCE construct as a HindIII-EcoRI fragment and ligated into pVL1393 cut with Sma I and Eco I. Briefly, the hUbCE gene fragment was purified by agarose gel separation, and ligated into the baculovirus vector pVL1393 (Invitrogen catalog no. V1392-20) previously cut with Sma I and Bgl II. The pVL1393-hUbCE construct was then used to transfect *Spodoptera frugiperda* (Sf9 cells, ATCC CRL 1711), and the cells maintained in insect cell culture media (Grace's *Antheraea* medium) supplemented with 10% FBS, lactal bumin hydrolysate, TC yeastolate and glutamate (Invitrogen catalog no. B823) following standard protocols (Invitrogen product guide; Summers and Smith (1987); *Texas Agricultural Experiment Station Bulletin No. 1555*, College Station, Tex.; Luckow et al. (1988) *Bio/technology* 6:47-55; and Miller et al., in *Genetic Engineering*, Vol. 8 ed. Setlow and Hollaender (Plenum Press: New York) pages 277-298). Transfected cells are grown until cells begin to lose their adherence to the culture plate surface, at which time the cells are harvested, collected by centrifugation, and lysed. The lysate is clarified by centrifugation to remove the cell wall debris, and the hUbCE can be purified from the lysate.

For instance, the hUbCE protein was isolated on an E1:ubiquitin charged column. Isolation of enzymes of the ubiquitin-conjugating system has been greatly assisted by "covalent" ubiquitin-affinity chromatography (Crehanover et al. (1982) *J. Biol. Chem.* 257:2537-2542; and Pickart et al. (1985) *J. Biol. Chem.* 260:1573-1581). This method takes advantage of the fact that the E1 enzyme is capable of forming a thiol ester with immobilized ubiquitin (e.g. ubiquitin-Sepharose) in the presence of ATP. Moreover, E1 enzymes bound to the immobilized ubiquitin can be exchanged with the subject hUbCE protein. Thus, both E1 and the subject hUbCE protein can be specifically purified on such columns, and can be recovered after elution with, for example, dithiothreitol. Moreover, with minor changes, this protocol can be used to isolate hUbCE:Ub conjugates (e.g. activated ubiquitin conjugates) for use in therapeutic target assays.

As described in U.S. patent application Ser. No. 08/176, 9375 the an E1-containing lysate was applied to a sepharose-ubiquitin column (Hershko et al. (1983) *J Biol. Chem.* 257:2537-2542) in the presence of ATP (e.g. 5 mM ATP, 10 mM MgCl₂, and 0.2 mM dithiothreitol, 50 mM Tris-HCl (pH 7.2)). The column was washed several times with this buffer. A clarified lysate of the hUbCE -producing insect cells, adjusted to 50 mM Tris-HCl (pH 7.2), 5 mM ATP, 10 mM MgCl₂, and 0.2 mM dithiothreitol, was then applied to the Ub:E1 column, washed, then eluted to remove any remaining ub:E1 (e.g. hUbCE will be exchanged for E1 on the column). The subject hUbCE protein was then eluted from the column by washing with 50 mM Tris-HCl (pH 9.0) containing 2 mM dithiothreitol.

In another exemplary embodiment, the recombinant hUbCE protein is generated as a poly(His) fusion protein for purification on a Ni²⁺ metal column. An XhoI to EcoRI fragment of the pKS construct is cloned into the pBlueBac A baculovirus (Invitrogen catalog no. V360-20) previously digested with XhoI and EcoRI. Following the manufacturer's protocols, the His₆-hUbCE fusion protein is then

expressed in Sf9 insect cells, and purified on a Ni²⁺ charged sepharose resin (Invitrogen catalog no. R801; see also Hochuli et al. (1987) *J. Chromatography* 411:177-184; and Janknecht et al. (1991) *PNAS* 88:8972-8976). Following purification of the fusion protein, the His₆ tag can be removed by treatment with entokinase.

EXAMPLE 2

In vitro ubiquitination of p53

We describe the cloning of a new human ubiquitin-conjugation enzyme hUbCE in Example 1. In Examples 4 and 5, we show that hUbCE specifically ubiquitinylates E6AP and is involved in the turnover of p53 in vivo. We have defined several discrete biochemical steps in the activation and transfer of ubiquitin onto p53. These biochemical reactions provide two levels of specificity in the ubiquitination of p53; the hUbCE dependent ubiquitination of E6AP, and the E6-dependent transfer of ubiquitin from ubiquitinylated E6AP to p53.

Proteins

To perform an in vitro ubiquitination reaction, native hUbCE and UBC2, the human homolog of the *S. cerevisiae* DNA repair gene, Rad6 (Koken et al. (1991) *PNAS* 88:8865-8869) were expressed and purified from *E. coli* BL21(DE3). Both proteins are readily soluble and easily purified using standard procedures. The cloning and purification of each of the proteins hUbCE, UBC2, p53, human E1, E6, and E6AP have been described in U.S. Ser. No. 08/247,904. Briefly, native p53 was expressed from the baculoviral vector pVL1392 in Sf9 insect cells according to the manufacturer's instructions (Pharmingen) and purified on a p53 affinity column. HPV18 E6 was expressed *E. coli* BL21 as a GST fusion protein and purified on GSH-sepharose. Human E1 was cloned by PCR from the published cDNA sequence (Handley et al. (1991) *PNAS* 88:258-262), and native protein was expressed and purified from baculoviral infected cells. E6AP was expressed in *E. coli* JM109 as a GST fusion protein and purified on GSH-sepharose.

Ubiquitination reactions

Ubiquitination reactions contained 50-200 ng of the indicated proteins in 50 mM Tris pH 7.5, 5 mM MgCl₂, 2mM ATP-γ-S, 0.1 mM DTT and 5 μM ubiquitin. Total reactions (30 μl) were incubated at 25° C. for 3 hrs and then loaded on an 8% SDS gel for analysis of p53 ubiquitination or a 4-20% gradient gel for analysis of ubiquitination of the ubiquitin-conjugating enzymes and E6AP. The gels were run and proteins were electrophoretically transferred to nitrocellulose. p53 proteins were revealed with the monoclonal antibody DO-1 (Oncogene Science) and the ECL system from NEN. Ubiquitinylated proteins were visualized using Extravidin-HRP from Sigma and the ECL system from NEN.

As demonstrated in FIG. 6, the appearance of specific p53-ubiquitin conjugates requires hUbCE, HPV18-E6, E6AP, ubiquitin and E1, the ubiquitin activating enzyme. In contrast, UBC2 was active in a minimal conjugation reaction containing E1, ATP and ubiquitin, in that E1 could activate ubiquitin and transfer it onto UBC2. However, UBC2 could not substitute for hUbCE in the p53 conjugation reaction (FIG. 2, lane 3). In addition, we made an active site cysteine-to-serine mutation in hUbCE. Such active site E2 mutants should accept activated ubiquitin from E1 but should not ubiquitinylate their downstream substrates owing to the high stability of the ester linkage formed between the active site serine and the carboxy-terminus of ubiquitin. This

mutant was inactive in the p53 conjugation reaction (FIG. 6, lane 7). These results demonstrate that a catalytically active hUbCE is absolutely required for generation of ubiquitinated p53 in this in vitro system.

In FIG. 7A we show that ubiquitinated E1 could transfer ubiquitin efficiently to hUbCE but not directly to E6AP and that ubiquitinated hUbCE transferred ubiquitin to E6AP in a reaction that was not further stimulated by E6. All of these ubiquitination reactions required the presence of the ubiquitin-activating enzyme, E1, and ubiquitin.

To address the issue of the specificity of hUbCE-mediated ubiquitination of E6AP we performed ubiquitination reactions with purified recombinant hUbCE, GST-UBC2, GST-UBC8 (Kaiser et al. (1994) *J Biol. Chem.* 269:8797-8802) and a GST-fusion of the so-called epidermal ubiquitin conjugating enzyme (Liu et al. (1992) *J Biol Chem* 267:15829-15835). Each of these recombinant proteins could accept activated ubiquitin from E1, but only hUbCE could donate ubiquitin to E6AP (FIG. 7B). We also confirmed that native UBC2 could accept ubiquitin from E1 but could not donate ubiquitin to E6AP (data not shown).

We then purified the ubiquitinated E6AP by affinity chromatography on glutathione-Sepharose and demonstrated that it did not contain appreciable amounts of ubiquitinated E1, ubiquitinated hUbCE or free ubiquitin. We found that this purified, ubiquitinated E6AP could donate ubiquitin to p53 in an E6-dependent reaction.

EXAMPLE 3

Microinjection of Sense and Anti-sense constructs of the hUbCE Gene

To investigate the consequences of interfering with hUbCE and E6AP function in p53 degradation, we performed microinjection experiments using sense and anti-sense constructs of the hUbCE gene. To facilitate the detection of p53 by indirect immunofluorescence, the experiments were performed in the human tumor cell line MDA-MB-468 which contain high levels of mutant p53 (Arg273His). In this line, the degradation of p53 could be stimulated by microinjection of an HPV-18 E6 expression plasmid.

In order to determine whether hUbCE and E6AP mediate the E6-dependent ubiquitination and degradation of p53 in vivo co-injection experiments were performed. To briefly describe the experiments, the CMV expression vectors were obtained by inserting the entire open-reading frame of one of HPV-18 E6, human E1, human E6-AP, hUbCE, or a Cys-85 mutant of hUbCE, in either a sense or anti-sense orientation (as indicated in FIG. 8) in the pX-plasmid (Baldin et al. (1993) *Genes & Devel.*, 7:812-821). Plasmids were purified with a Promega Wizard Maxi-prep kit and injected at a concentration of 50 to 100 µg/µl in the presence of normal affinity-purified rabbit or mouse antibody (5 mg/ml in PBS) used as microinjection marker.

Cell monolayers of asynchronous MDA-MB-468 cells were injected with the indicated DNAs (FIG. 8) along with rabbit IgG to allow identification of injected cells with an automated microinjection system (AIS, Zeiss; Ansoorge et al. (1988) *J. Biochem. Biophys. Meth.*, 16:283-292). All microinjection experiments were carried out in 3.5 cm Petri dishes containing 3 ml of DMEM medium carbonate free, in order to avoid the decrease in pH of the medium during the injection. Each cell was injected at a pressure between 50 and 150 hPa. After 24 hrs the cells were fixed and stained with a p53 specific monoclonal antibody (DO-1; Oncogene Sciences) followed by a biotinylated horse anti-rabbit antibody and Texas red conjugated streptavidin. Injected cells

were identified by staining with an FITC conjugated goat anti-rabbit antibody (Baldin et al. (p993) *Genes and Dev* 7:812-821).

When either an anti-sense or mutant hUbCE expression plasmid or an expression plasmid encoding anti-sense E6AP was co-injected with the E6 expression plasmid, the E6 stimulated degradation of p53 was inhibited (FIG. 8). Similar results were obtained when polyclonal antibodies generated against human hUbCE or an expression plasmid encoding a mutant form of E6AP (Peter Howley, pers.comm.) were microinjected (not shown).

Co-injection of an E6 expression plasmid with an expression plasmid encoding anti-sense E1 also inhibited the E6 stimulated degradation of p53. Co-injection of anti-sense or mutant UBC2 expression plasmids had a negligible effect on the E6 stimulated degradation of p53 (FIG. 8).

Moreover, the data show that an hUbCE mutant, Cys-85→Ser, which produces an inactive form of the enzyme, is possibly a dominant negative mutant able to at least partially rescue p53.

EXAMPLE 4

Generating a Molecular Model of the hUbCE protein

The three dimensional coordinates of the protein backbone from the structure of UBC1 from *A. thaliana* (Brookhaven databank file 1AAK.pdb) were used for homology modeling of hUbCE. Modeling was performed with the Protein Workbench software package of QUANTA, version 4 (MSI, Burlington Mass.).

Briefly, the amino acid sequence of hUbCE (SEQ ID No. 2) and UBC1 were aligned using the alignment program in QUANTA. This alignment shows a 44% match of similar residues. The coordinates of the backbone non-hydrogen atoms were then copied onto the hUbCE sequence, sidechain coordinates for the hUbCE model were ignored at this point. The resulting hUbCE structure was then energy minimized using 200 steps of the steepest descent algorithm followed by 5000 steps of the adopted-base Newton Raphson algorithm. All atoms, including polar hydrogens and all side chains were allowed to move. The resulting CHARMM energy of the system was -7084.2 kcal.

In the next step, the structure was heated up to 500° K using 2000 steps or a total time of 2 psec. After heating, the system was then allowed to equilibrate for 9 psec (9000 steps). The final CHARMM energy after 10 psec was around -5750 kcal. Finally, the system was cooled down to 300° K in steps of 50° K (1 psec cooling, 4 psec equilibration) and finally equilibrated at 300° K for 6 psec. The final total CHARMM energy was around -6650 kcal. The final structure showed no serious conformational strains or improper angles. The atomic coordinates for the full length model are shown in FIG. 1.

In the next step we modeled the 4-meric peptide Ala-Ile-Arg-Gly into the active site. This peptide was derived from the c-terminal sequence of ubiquitin (RIRG). A thioester bond was manually constructed in both cases between the C-terminal Gly and the active site cysteines. The system was energy-minimized and subsequently subjected to molecular dynamics simulations. In both cases the Ile residue of the peptide settles into the hydrophobic pocket. There are two backbone-backbone hydrogen bonds between the loops and the peptide. The Arg of the peptide forms hydrogen bonds with a conserved Asp residue (between the conserved Val and Ile residues) in both cases.

-continued

145

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 522 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: both
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: cDNA

- (i x) FEATURE:
 (A) NAME/KEY: CDS
 (B) LOCATION: 22..462

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:5:

```

CGCAAAAGCA AACCCAGTAAC G ATG GCT TTG AAA AGA ATT AAC CGT GAA TTA      51
                    Met Ala Leu Lys Arg Ile Asn Arg Glu Leu
                    1                               5                               10

GCT GAT CTT GGA AAA GAC CCA CCG TCT TCT TGT TCC GCC GGC CCT GTT      99
Ala Asp Leu Gly Lys Asp Pro Pro Ser Ser Cys Ser Ala Gly Pro Val
                    15                               20                               25

GGC GAT GAT TTA TTC CAT TGG CAA GCT ACA ATC ATG GGT CCT GCT GAC     147
Gly Asp Asp Leu Phe His Trp Gln Ala Thr Ile Met Gly Pro Ala Asp
                    30                               35                               40

AGC CCT TAT GCG GGT GGT GTC TTC TTC TTG TCC ATT CAT TTC CCT ACG     195
Ser Pro Tyr Ala Gly Gly Val Phe Phe Leu Ser Ile His Phe Pro Thr
                    45                               50                               55

GAC TAC CCA TTC AAG CCA CCA AAG GTA AAC TTT ACA ACC AGA ATC TAT     243
Asp Tyr Pro Phe Lys Pro Pro Lys Val Asn Phe Thr Thr Arg Ile Tyr
                    60                               65                               70

CAT CCC AAC ATC AAT TCA AAC GGT AGC ATT TGT TTG GAT ATC CTT CGT     291
His Pro Asn Ile Asn Ser Asn Gly Ser Ile Cys Leu Asp Ile Leu Arg
                    75                               80                               85

GAC CAA TGG TCT CCA GCG TTG ACT ATA TCA AAG GTA TTA CTG TCT ATC     339
Asp Gln Trp Ser Pro Ala Leu Thr Ile Ser Lys Val Leu Leu Ser Ile
                    95                               100                               105

TGC TCA TTG TTG ACA GAT CCT AAT CCT GAT GAT CCG CTT GTG CCT GAA     387
Cys Ser Leu Leu Thr Asp Pro Asn Pro Asp Asp Pro Leu Val Pro Glu
                    110                               115                               120

ATT GCG CAC GTC TAC AAA ACT GAC AGA TCC CGT TAT GAA TTA AGT GCT     435
Ile Ala His Val Tyr Lys Thr Asp Arg Ser Arg Tyr Glu Leu Ser Ala
                    125                               130                               135

CGT GAA TGG ACT AGA AAA TAC GCA ATC TAGAGTTTGT TTCTGTGTTG         482
Arg Glu Trp Thr Arg Lys Tyr Ala Ile
                    140                               145

ATATTAAATA TTCATCTCTT AAAAAAAAAA AAAAAACTCG                       522

```

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 147 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:6:

```

Met Ala Leu Lys Arg Ile Asn Arg Glu Leu Ala Asp Leu Gly Lys Asp
 1                               5                               10                               15

Pro Pro Ser Ser Cys Ser Ala Gly Pro Val Gly Asp Asp Leu Phe His
                20                               25                               30

Trp Gln Ala Thr Ile Met Gly Pro Ala Asp Ser Pro Tyr Ala Gly Gly

```


-continued

	35		40		45														
Val	Phe	Phe	Leu	Ser	Ile	His	Phe	Pro	Thr	Asp	Tyr	Pro	Phe	Lys	Pro				
	50					55					60								
Pro	Lys	Val	Asn	Phe	Thr	Thr	Arg	Ile	Tyr	His	Pro	Asn	Ile	Asn	Ser				
65					70					75					80				
Asn	Gly	Ser	Ile	Cys	Leu	Asp	Ile	Leu	Arg	Asp	Gln	Trp	Ser	Pro	Ala				
				85					90					95					
Leu	Thr	Ile	Ser	Lys	Val	Leu	Leu	Ser	Ile	Cys	Ser	Leu	Leu	Thr	Asp				
			100					105						110					
Pro	Asn	Pro	Asp	Asp	Pro	Leu	Val	Pro	Glu	Ile	Ala	His	Val	Tyr	Lys				
		115					120					125							
Thr	Asp	Arg	Ser	Arg	Tyr	Glu	Leu	Ser	Ala	Arg	Glu	Trp	Thr	Arg	Lys				
	130					135					140								
Tyr	Ala	Ile																	
145																			

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 147 amino acids
 (B) TYPE: amino acid
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: protein

(v) FRAGMENT TYPE: internal

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Met	Xaa	Leu	Lys	Arg	Ile	Xaa	Xaa	Glu	Leu	Xaa	Asp	Leu	Xaa	Xaa	Asp				
1				5				10						15					
Pro	Pro	Xaa	Xaa	Cys	Ser	Ala	Gly	Pro	Val	Gly	Asp	Asp	Xaa	Xaa	His				
		20					25						30						
Trp	Gln	Ala	Xaa	Ile	Met	Gly	Pro	Asn	Asp	Ser	Pro	Tyr	Xaa	Gly	Gly				
		35				40						45							
Val	Phe	Phe	Leu	Xaa	Ile	His	Phe	Pro	Thr	Asp	Tyr	Pro	Xaa	Lys	Pro				
	50					55					60								
Pro	Lys	Xaa	Xaa	Xaa	Thr	Thr	Xaa	Ile	Tyr	His	Pro	Asn	Ile	Asn	Ser				
65					70				75					80					
Asn	Gly	Xaa	Ile	Cys	Leu	Asp	Ile	Leu	Xaa	Xaa	Gln	Trp	Ser	Pro	Ala				
				85					90					95					
Leu	Thr	Ile	Ser	Lys	Val	Leu	Leu	Ser	Ile	Cys	Ser	Leu	Leu	Xaa	Asp				
			100					105						110					
Xaa	Asn	Pro	Asp	Asp	Pro	Leu	Val	Pro	Glu	Ile	Ala	Xaa	Xaa	Tyr	Xaa				
		115					120					125							
Xaa	Asp	Arg	Xaa	Xaa	Tyr	Xaa	Xaa	Xaa	Ala	Xaa	Glu	Trp	Thr	Xaa	Lys				
	130					135					140								
Tyr	Ala	Xaa																	
145																			

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: other nucleic acid

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:8:

-continued

GCGCGCAAGC TTTAYGARGG WGGWGTYYTY TT

3 2

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 36 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: other nucleic acid

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:9:

GCGCGCGAAT TCACNGCRTA YTTYTTNGTC CCAYTC

3 6

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 38 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: other nucleic acid

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:10:

GCGCGCAAGC TTCNGTNGG NGAYTTRTTY CAYTGGCA

3 8

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(i i) MOLECULE TYPE: other nucleic acid

(x i) SEQUENCE DESCRIPTION: SEQ ID NO:11:

GCGCGCGAAT TCAATNGTNAR NGCNGGCGAC CA

3 2

We claim:

1. An isolated nucleic acid comprising a nucleotide sequence which encodes a ubiquitin conjugating enzyme (UbCE) polypeptide having an amino acid sequence at least 95% identical to at least one of SEQ ID Nos. 2, 4 or 6, which UbCE polypeptide catalyzes ubiquitin conjugation.

2. An isolated nucleic acid comprising a nucleotide sequence encoding an UbCE polypeptide comprising an amino acid sequence at least 95% homologous to SEQ ID NO. 2, which UbCE polypeptide specifically binds at least one of an E6 AP protein, a papillomavirus E6 protein, or p53.

3. The nucleic acid of claim 1 or 2, further comprising a transcriptional regulatory sequence operably linked to said nucleotide sequence so as to render said nucleotide sequence along with said transcriptional regulatory sequence suitable for use as an expression vector.

4. An expression vector, capable of replicating in at least one of prokaryotic cell or eukaryotic cell, comprising the nucleic acid of claim 1 or 2.

5. A host cell transfected with the expression vector of claim 4 and expressing said UbCE polypeptide.

6. A method of producing a recombinant ubiquitin conjugating enzyme comprising culturing the cell of claim 5 in a cell culture medium to express said UbCE polypeptide and isolating said UbCE polypeptide from said cell culture.

7. The nucleic acid of claim 1, wherein said UbCE polypeptide has an amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID No. 2.

8. The nucleic acid of claim 1 or 2, wherein said UbCE polypeptide has an amino acid sequence at least 97% identical to the amino acid sequence of SEQ ID No. 2.

9. The nucleic acid of claim 8, wherein said UbCE polypeptide has an amino acid sequence represented by SEQ ID No. 2.

10. The nucleic acid of claim 1, wherein the UbCE polypeptide has an amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID No. 4.

11. The nucleic acid of claim 1, wherein said UbCE polypeptide has an amino acid sequence at least 97% identical to the amino acid sequence of SEQ ID No. 4.

12. The nucleic acid of claim 1, wherein said UbCE polypeptide has an amino acid sequence represented by SEQ ID No. 4.

13. The nucleic acid of claim 1, wherein said UbCE polypeptide has an amino acid sequence at least 95% identical to the amino acid sequence of SEQ ID No. 6.

14. The nucleic acid of claim 1, wherein said UbCE polypeptide has an amino acid sequence at least 97% identical to the amino acid sequence of SEQ ID No. 6.

15. The nucleic acid of claim 1, wherein said UbCE polypeptide has an amino acid sequence represented by SEQ ID No. 6.

16. The nucleic acid of claim 2, wherein said UbCE polypeptide contains at least one amino acid residue different from SEQ ID No. 2 and which inhibits ubiquitination of a p53 or E6AP protein by an enzyme having a sequence represented by SEQ ID No. 2.

17. The nucleic acid of claim 2, wherein said UbCE polypeptide comprises a dominant negative mutation at the active site cysteine.

18. The nucleic acid of claim 16, wherein said UbCE polypeptide has an ubiquitin conjugating activity which is less than 5 percent of a ubiquitin conjugating enzyme identical to SEQ ID No. 2.

19. The nucleic acid of claim 1, wherein said UbCE polypeptide specifically binds at least one of an E6-AP protein, a papillomavirus E6 protein, or p53.

20. The nucleic acid of claim 1 or 2, wherein said UbCE polypeptide mediates ubiquitination of a p53 protein.

21. The nucleic acid of claim 1 or 2, wherein said UbCE polypeptide mediates ubiquitination of an E6-AP polypeptide.

22. The nucleic acid of claim 1, wherein said nucleotide sequence which encodes said UbCE polypeptide is the coding sequence of SEQ ID No. 1.

23. The nucleic acid of claim 1, wherein said nucleotide sequence which encodes said UbCE polypeptide is the coding sequence of SEQ ID No. 3.

24. The nucleic acid of claim 1, wherein said nucleotide sequence which encodes said UbCE polypeptide is the coding sequence of SEQ ID No. 5.

25. The nucleic acid of claim 1 or 2, wherein said UbCE polypeptide is comprises within a fusion protein.

26. The nucleic acid of claim 25, wherein said fusion protein is glutathione-S transferase (GST) fusion protein.

27. The nucleic acid of claim 25, wherein said fusion protein includes a purification leader sequence.

* * * * *