



US005942223A

# United States Patent [19]

Bazer et al.

[11] Patent Number: 5,942,223

[45] Date of Patent: Aug. 24, 1999

## [54] ANTIVIRAL THERAPY USING OVINE OR BOVINE INTERFERON-TAU

[75] Inventors: **Fuller Warren Bazer**, College Station, Tex.; **Howard Marcellus Johnson**, Gainesville, Fla.; **Carol Hanlon Pontzer**, Silver Spring, Md.; **Troy Lee Ott**, Bryan, Tex.; **Gino Van Heeke**, Witterswil, Switzerland

[73] Assignee: **University of Florida**, Gainesville, Fla.

[21] Appl. No.: **08/455,524**

[22] Filed: **May 31, 1995**

### Related U.S. Application Data

[63] Continuation of application No. 08/438,753, May 10, 1995, Pat. No. 5,705,363, which is a continuation-in-part of application No. 08/139,891, Oct. 19, 1993, abandoned, which is a continuation-in-part of application No. 07/847,741, Mar. 9, 1992, abandoned, which is a continuation-in-part of application No. 07/318,050, Mar. 2, 1989, abandoned, said application No. 08/139,891, is a continuation-in-part of application No. 07/969,890, Oct. 30, 1992, abandoned.

[51] Int. Cl.<sup>6</sup> ..... **A61K 38/21**; C12N 15/20; C07K 14/555

[52] U.S. Cl. .... **424/85.4**; 435/69.51

[58] Field of Search ..... 424/85.4, 85.7; 435/69.51

### [56] References Cited

#### U.S. PATENT DOCUMENTS

4,997,646 3/1991 Hansen et al. .  
5,019,382 5/1991 Cummins ..... 424/85.4

#### FOREIGN PATENT DOCUMENTS

0 367 063 5/1990 European Pat. Off. .  
WO 90/09806 9/1990 WIPO .  
WO 93/12146 12/1992 WIPO .  
WO 94/10113 5/1994 WIPO .

#### OTHER PUBLICATIONS

Imakawa, K., et al. (1987) *J. Cell Biol.* 105(4, part 2) : 11A, abst. No. 49.

Charpigny, G., et al. (1988) *FEBS Lett.* 228: 12-16.

Bazer, F.W., and Johnson, H.M., "Type I Conceptus Interferons: Maternal Recognition of Pregnancy Signal and Potential Therapeutic Agents", *American J. of Reproductive Immun.* 26:19-22 (1991).

Bazer, F.W., et al., "Roles of Ovine Trophoblast Protein-1 and Oestradiol/ Prolactin in the Establishment of Pregnancy in Sheep and Pigs", *Reprod. Fertil. Dev.* 4:335-340 (1992).

Farin, C.E., et al., "Expression of Trophoblastic Interferon Genes in Sheep and Cattle", *Biol. of Reprod.* 43:210-218 (1990).

Hansen, T.R., et al., "Complex Binding of the Embryonic Interferon, Ovine Trophoblast Protein-1, to Endometrial Receptors", *J. Interferon Res.* 9:215-225 (1989).

Hansen, T.R., et al., "The Genes for the Trophoblast Interferons and the Related Interferon- $\alpha_H$  Posses Distinct 5'-Promoter and 3'-Flanking Sequence", *J. Biol. Chem.* 266(5):3060-3067 (1991).

Imakawa, K., et al., "Interferon-like sequence of ovine trophoblast protein secreted by embryonic trophectoderm", *Nature* 330:377-379 (1987).

Imakawa, K., et al., "Molecular Cloning and Characterization of Complementary Deoxyribonucleic Acids Corresponding to Bovine Trophoblast Protein 1: A Comparison with Ovine Trophoblast Protein-1 and Bovine Interferon- $\alpha_H$ ", *Mol. Endocrin.* 3(1):127-139 (1989).

Leaman, D.W., et al., "Genes for the Trophoblast Interferons and their Distribution among Mammals", *Reprod. Fertil. Dev.* 4:349-353 (1992).

Leaman, D.W., and Roberts, R.M., "Genes for the Trophoblast Interferons in Sheep, Goat, and Musk Ox and Distribution of Related Genes Among Mammals", *J. Interferon Res.* 12:1-11 (1992).

Newton, G.R., et al., "Inhibition of Lymphocyte Proliferation by Ovine Trophoblast Protein-1 and a High Molecular Weight Glycoprotein Produced by the Peri-Implantation Sheep Conceptus", *Am. J. Reprod. Immunol.* 19:99-107 (1989).

Ott, T.L., et al., "Cloning and Expression in *Saccharomyces cerevisiae* of a Synthetic Gene for the Type-I Trophoblast Interferon Ovine Trophoblast Protein-1: Purification and Antiviral Activity", *J. Interferon Res.* 11:357-364 (1991).

Pontzer, C.H., et al., "Antiviral Activity of the Pregnancy Recognition Hormone Ovine Trophoblast Protein-1", *Biochem. and Biophys. Res. Commun.* 152(2):801-807 (1988).

Pontzer, C.H., et al., "Localization of an antiviral site on the pregnancy recognition hormone, ovine trophoblast protein-1", *Proc. Natl. Acad. Sci. USA* 87:5945-5949 (1990).

Pontzer, C.H., et al., "Antiproliferative Activity of a Pregnancy Recognition Hormone, Ovine Trophoblast Protein-1", *Cancer Res.* 51:5304-5307 (1991).

Roberts, R.M., et al., "Interferons at the placenta interface", *J. Reprod. Fert., Suppl.* 41:63-74 (1990).

Roberts, R.M., et al., "Unique Features of the Trophoblast Interferons", *Pharmac. Ther.* 51:329-345 (1991a).

Roberts, R.M., et al., "The polypeptides and genes for ovine and bovine trophoblast protein-1", *J. Reprod. Fert., Suppl.* 43:3-12 (1991b).

Roberts, R.M., et al., "Interferons as Hormones of Pregnancy", *Endocrine Rev.* 13(3):432-452 (1992).

Tuo, W., et al., "Natural Killer Cell Activity of Lymphocytes Exposed to Ovine, Type I, Trophoblast Interferon", *AJRI* 29:26-34 (1993).

(List continued on next page.)

Primary Examiner—David L. Fitzgerald

Attorney, Agent, or Firm—Charles K. Scholtz; Joanne R. Petithory; Dehlinger & Associates

### [57] ABSTRACT

The invention provides antiviral therapeutic methods employing bovine or ovine interferon-tau (IFN- $\tau$ ) proteins and polypeptides. The IFN- $\tau$  proteins exhibit the antiviral and antiproliferative properties characteristic of type I interferons. An advantage of the invention is that IFN- $\tau$  has essentially no cytotoxic effects on treated cells as does, for example, IFN- $\alpha$ .

8 Claims, 21 Drawing Sheets

## OTHER PUBLICATIONS

Vallet, J.L., et al., "The Effect of Ovine Trophoblast Protein-One on Endometrial Protein Secretion and Cyclic Nucleotides", *Biol. Reprod.* 37:1307-1316 (1987).

Vallet, J.L., et al., "Effect of ovine conceptus secretory proteins and purified ovine trophoblast protein-1 on inter-  
oestrous interval and plasma concentrations of prostaglan-  
dins F-2 $\alpha$  and E and 13,14-dihydro-15-keto-prostaglandin  
F-2 $\alpha$  in cyclic ewes", *J. Reprod. Fert.* 84:493-504 (1988).

Whaley, A.E., et al., "Molecular Cloning of Unique Inter-  
ferons from Human Placenta," in *Society of Reproduction*,  
*24th Annual Meeting*, p. 186 (1991).

Yong, K., et al., "Insulin-Like Growth Factors in Sheep  
Uterine Fluids: Concentrations and Relationship to Ovine  
Trophoblast Protein-1 Production during Early Pregnancy",  
*Biol. Reprod.* 45:135-142 (1991).





Fig. 1B

Fnu4H I		Mae III	
Bbv I		Hph I	
Alu I		BstE II	
BsmA I	Pyu II	Fnu4H I	
NsoB II	NsoB II		
TCGATTGCTCCGCACAGCTGCCTGCAAGACCGGAAAGACTTCGGTCTGCCCGCAGGAAATGGTTGAAGGTGACCAACTGC		160	
AGCTAACAGAGGCGTGTCGACGGACGTTCTGTGCCTTTCTGAAGCCAGACGGCGTCCTTTACCAACTTCCACTGGTTGACG		.	
81	96	148	
87		148	
97		149	
98			
98			
Msp I		Fnu4H I	
ScrF I		Bbv I	
Nci I		Pst I	
Bcn I		Fnu4H I	
Alu I	Hpa II	Bbv I	
HinD III	Rsa I		
AAAAAGACCAAGCTTTC		240	
TTTCTGTTTCGAAAGGCCCATGACATACTTTACGACGTCGTCAGAAAGTTGGACAAGATGTGACTTGTAAGAACCGG		.	
170	181	232	
171		236	
177		236	
177		236	
177		237	
178		238	

Fig. 1C







Fig. 1E






Fig. 2



-23 -9  
met ala phe val leu ser leu leu met ala leu val leu val ser  
oIFNt cccc ATG GCC TTC GTG CTC TCT CTA CTG ATG GCC CTG GTG CTG GTC AGC  
huIFNt1 cccc ATG GCC TTC GTG CTC TCT CTA CTC ATG GCC CTG GTG CTG GTC AGC

-8 -1 +1 11  
tyr gly pro gly gly ser leu gly cys tyr leu ser arg lys leu met leu asp ala  
TAT GGC CCA GGA GGA TCT CTG GGT TGT TAC CTA TCT CGG AAA CTC ATG CTG GAT GCC  
TAC GGC CCA GGA GGA TCC CTG GGT TGT GAC CTG TCT CAG AAC CAC GTG CTG GTT GGC  
asp gln asn his val val gly

12 30  
arg glu asn leu lys leu leu asp arg met asn arg leu ser pro his ser cys leu  
AGG GAG AAC CTC AAG CTC CTG GAC CGA ATG AAC AGA CTC TCC CCT CAT TCC TGT CTG  
AGG AAG AAC CTC AGG CTC CTG GAC GAA ATG AGG AGA CTC TCC CCT CGC TTT TGT CTG  
lys arg glu arg arg phe

31 49  
gln asp arg lys asp phe gly leu pro gln glu met val glu gly arg gln leu gln  
CAG GAC AGA AAA GAC TTT GGT CTT CCC CAG GAG ATG GTG GAG GGC GAC CAG CTC CAG  
CAG GAC AGA AAA GAC TTC GCT TTA CCC CAG GAA ATG GTG GAG GGC GGC CAG CTC CAG  
ala gly

50 68  
lys asp gln ala phe pro val leu tyr glu met leu gln gln ser phe asn leu phe  
AAG GAC CAG GCC TTC CCT GTG CTC TAC GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC  
GAG GCC CAG GCC ATC TCT GTG CTC CAT GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC  
glu ala ile ser his

69 87  
tyr thr glu his ser ser ala ala trp asp thr thr leu leu glu gln leu cys thr  
TAC ACA GAG CAC TCC TCT GCT GCC TGG GAC ACC ACC CTC CTG GAG CAG CTC TGC ACT  
CAC ACA GAG CAC TCC TCT GCT GCC TGG GAC ACC ACC CTC CTG GAG CAG CTC CGC ACT  
his arg

88 106  
gly leu gln gln gln leu asp his leu asp thr cys arg gly gln val met gly glu  
GGA CTC CAA CAG CAG CTG GAC CAC CTG GAC ACC TGC AGG GGT CAA GTG ATG GGA GAG  
GGA CTC CAT CAG CAG CTG GAC AAC CTG GAT GCC TGC CTG GGG CAG GTG ATG GGA GAG  
his asn ala leu

107 125  
glu asp ser glu leu gly asn met asp pro ile val thr val lys lys tyr phe gln  
GAA GAC TCT GAA CTG GGT AAC ATG GAC CCC ATT GTG ACC GTG AAG AAG TAC TTC CAG  
GAA GAC TCT GCC CTG GGA AGG ACG GGC CCC ACC CTG GCT CTG AAG AGG TAC TTC CAG  
ala arg thr gly thr leu ala leu arg

126 144  
gly ile tyr asp tyr leu gln glu lys gly tyr ser asp cys ala trp glu ile val  
GGC ATC TAT GAC TAC CTG CAA GAG AAG GGA TAC AGC GAC TGC GCC TGG GAA ATC GTC  
GGC ATC CAT GTC TAC CTG AAA GAG AAG GGA TAC AGC GAC TGC GCC TGG GAA ACC GTC  
his val lys thr

145 163  
arg val glu met met arg ala leu thr val ser thr thr leu gln lys arg leu thr  
AGA GTC GAG ATG ATG AGA GCC CTC ACT GTA TCA ACC ACC TTG CAA AAA AGG TTA ACA  
AGA CTG GAA ATC ATG AGA TCC TTC TCT TCA TTA ATC AGC TTG CAA GAA AGG TTA AGA  
leu ile ser phe ser ser leu ile ser glu arg

164 172  
lys met gly gly asp leu asn ser pro end  
AAG ATG GGT GGA GAT CTG AAC TCA CCT TGA  
ATG ATG GAT GGA GAC CTG AGC TCA CCT TGA  
met asp ser

Fig. 3




Fig. 4





Fig. 5

Peptides	MW	HI*	Sequence
IFNt(1-37) (SEQ ID NO:5)	4465	-0.78	CYSLRKLMLDARENKLLDRMNRLSPHSCLODRKDFG
IFNt(34-64) (SEQ ID NO:6)	3610	-0.72	KDFGLPQEMVEGDQLQKDQAFPVLYEMLQQS
IFNt(62-92) (SEQ ID NO:7)	3586	-0.53	QQSFNLFYTEHSSAAWDTTLLLEQLCTGLQQQ
IFNt(90-122) (SEQ ID NO:8)	3712	-0.86	QQQLDHLDTCRGQVMGEEDSELGNMDPIVTVKK
IFNt(119-150) (SEQ ID NO:9)	3948	-0.56	TVKKYFQGIYDYLQEKGYSDCAWEIVRVEMMR
IFNt(139-172) (SEQ ID NO:10)	3818	-0.11	CAWEIVRVEMMRALTVSTTLQKRLTKMGGDLNSP

\*Hydropathic Index

Fig. 6





1 80  
CTGAGATGGGATCAGAGAACCTACCTGAAGGTTCCCCCTGACCCCATCTCAGCCAGCCCAGCAGCAGCCGCATCTTCCCC

81 140  
ATG GCC TTC GTG CTC TCT CTA CTG ATG GCC CTG GTG CTG GTC AGC TAT GGC CCA GGA GGA  
S1 S20  
Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr Gly Pro Gly Gly  
141 200  
TCT CTG GGT TGT TAC CTA TCT CGG AAA CTC ATG CTG GAT GCC AGG GAG AAC CTC AAG CTC  
S21 17  
Ser Leu Gly Cys Tyr Leu Ser Arg Lys Leu Met Leu Asp Ala Arg Glu Asn Leu Lys Leu  
201 260  
CTG GAC CGA ATG AAC AGA CTC TCC CCT CAT TCC TGT CTG CAG GAC AGA AAA GAC TTT GGT  
18 37  
Leu Asp Arg Met Asn Arg Leu Ser Pro His Ser Cys Leu Gln Asp Arg Lys Asp Phe Gly  
261 320  
CTT CCC CAG GAG ATG GTG GAG GGC GAC CAG CTC CAG AAG GAC CAG GCC TTC CCT GTG CTC  
38 57  
Leu Pro Gln Glu Met Val Glu Gly Asp Gln Leu Gln Lys Asp Gln Ala Phe Pro Val Leu  
321 380  
TAC GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC TAC ACA GAG CAC TCC TCT GCT GCC TGG  
58 77  
Tyr Glu Met Leu Gln Gln Ser Phe Asn Leu Phe Tyr Thr Glu His Ser Ser Ala Ala Trp  
381 440  
GAC ACC ACC CTC CTG GAG CAG CTC TGC ACT GGA CTC CAA CAG CAG CTG GAC CAC CTG GAC  
78 97  
Asp Thr Thr Leu Leu Glu Gln Leu Cys Thr Gly Leu Gln Gln Gln Leu Asp His Leu Asp  
441 500  
ACC TGC AGG GGT CAA GTG ATG GGA GAG GAA GAC TCT GAA CTG GGT AAC ATG GAC CCC ATT  
98 117  
Thr Cys Arg Gly Gln Val Met Gly Glu Glu Asp Ser Glu Leu Gly Asn Met Asp Pro Ile  
501 560  
GTG ACC GTG AAG AAG TAC TTC CAG GGC ATC TAT GAC TAC CTG CAA GAG AAG GGA TAC AGC  
118 137  
Val Thr Val Lys Lys Tyr Phe Gln Gly Ile Tyr Asp Tyr Leu Gln Glu Lys Gly Tyr Ser  
561 620  
GAC TGC GCC TGG GAA ATC GTC AGA GTC GAG ATG ATG AGA GCC CTC ACT GTA TCA ACC ACC  
138 157  
Asp Cys Ala Trp Glu Ile Val Arg Val Glu Met Met Arg Ala Leu Thr Val Ser Thr Thr  
621 666  
TTG CAA AAA AGG TTA ACA AAG ATG GGT GGA GAT CTG AAC TCA CCT TGATGACTCTTGCCGACTA  
158 172  
Leu Gln Lys Arg Leu Thr Lys Met Gly Gly Asp Leu Asn Ser Pro



764  
AGATGCCACATCAGCCTCCTACACCCGCCTGTGTTTCATTTTCTGCTCCAGCCACCAAATTCATTG

844  
AATTACTTTAGCTGATACTTTGTCTAGTAGTAAAAAGCAAGTAGATATAAAAGTATTCAGCTGTAGGGGCATGAGTCCTGA

924  
AATGATGCCTTCCCTGATGTTATCTGTTGCTGATTTATTTATACCTTCTAGCATTTAACATACTTAAATATTAGGAAAT

972  
TTGTTAAGTTACATTTACATCTGTACATCATATTAATAATTTCTAAACAAAAA

Fig. 7









Fig. 12




Fig. 13




Fig. 14





Fig. 15




Fig. 16





Fig. 17



**Fig. 18**





88		90										100						106
gly	leu	gln	gln	gln	leu	asp	his	leu	asp	thr	cys	arg	gly	gln	val	met	gly	glu
GGA	CTC	CAA	CAG	CAG	CTG	GAC	CAC	CTG	GAC	ACC	TGC	AGG	GGT	CAA	GTG	ATG	GGA	GAG
		T					A		t	G		CT	g	g				
		T					A		t	G		CT	g	g				
		T				t	G		t	G		CT	g	g		C		
		T				t	G		t	G		CT	g	g		C		
		T				t	G		t	G		CT	g	g		C		
		T				t	G			G		CT	g	g		C		
		T				t	G		t	G	t	CT	g	g	T	CT		
107			110										120					125
glu	asp	ser	glu	leu	gly	asn	met	asp	pro	ile	val	thr	val	lys	lys	tyr	phe	gln
GAA	GAC	TCT	GAA	CTG	GGT	AAC	ATG	GAC	CCC	ATT	GTG	ACC	GTG	AAG	AAG	TAC	TTC	CAG
			CC		a	GG	C	G		CC	C	G	T	C		G		
			CC		a	GG	C	G		CC	C	G	T	C		G		
			CC		a	GA	C	G		CC	C	G		A		G	t	
			CC		a	GG	C	G		CC	C	G		A		C	t	
			CC		a	GG	C	G		CC	C	G		A		C	t	
			CC		a	GG	C	G		CC	C	G				GC		
			CC		a	GG	C	G		CC	C	G				GC		
126				130									140					144
gly	ile	tyr	asp	tyr	leu	gln	glu	lys	gly	tyr	ser	asp	cys	ala	trp	glu	ile	val
GGC	ATC	TAT	GAC	TAC	CTG	CAA	GAG	AAG	GGA	TAC	AGC	GAC	TGC	GCC	TGG	GAA	ATC	GTC
		C	T			A											C	
		C	T			A											C	
		C	T			A				t	t						t	
		C	T			A				t	t					!		
		C	T			A				t	!							
		C	AT											!				
		C	T						!									
145					150									160				163
arg	val	glu	met	met	arg	ala	leu	thr	val	ser	thr	thr	leu	gln	lys	arg	leu	thr
AGA	GTC	GAG	ATG	ATG	AGA	GCC	CTC	ACT	GTA	TCA	ACC	ACC	TTG	CAA	AAA	AGG	TTA	ACA
	C	G	a	C		T	T	T	TC	T	T	G			G			G
		g	a	C		T	T	T	TC	T	T	G			G			G
	C	G	a	C		T	t	g	T			G		C				G
164									172									
lys	met	gly	gly	asp	leu	asn	ser	pro										
AAG	ATG	GGT	GGA	GAT	CTG	AAC	TCA	CCT	TGA									
T		A		c		G												
T		A		c		G												
T		A		c		G												

Fig. 19B

-23

-9

oTP-1Met ala phe val leu ser leu leu met ala leu val leu val ser

LEXY.5

TOSHI.9

TOSHI.10

-8

-1 +1

11

tyr gly pro gly gly ser leu gly cys tyr leu ser arg lys leu met leu asp ala

asp

gln asn his val val gly

(---)asp

gln asn his val val gly

arg asp

gln asn his val val gly

12

20

30

arg glu asn leu lys leu leu asp arg met asn arg leu ser pro his ser cys leu

lys arg

glu arg

arg phe

lys arg

gln arg

arg phe

ser gln arg

gly gln arg

leu arg phe

31

40

49

gln asp arg lys asp phe gly leu pro gln glu met val glu gly asp gln leu gln

ala

gly

ala

(Stop)

gly

ala phe

gly

Clone 21!

gly

Clone 35!

gly

Clone 15!

gly

Clone 18!

val ser phe

50

60

68

lys asp gln ala phe pro val leu tyr glu met leu gln gln ser phe asn leu phe

glu ala

ile ser

his

glu ala

ile ser

his

glu ala

ile ser

his

glu ala

ile ser

his

glu ala

ile ser

his lys

glu ala

ile ser

his

69

80

87

tyr thr glu his ser ser ala ala try asp thr thr leu leu glu gln leu cys thr

his

arg

his

arg

his

arg

his

arg

his

arg

his arg

arg

his lys arg

leu

Fig. 20A

88	90											100						106
gly	leu	gln	gln	gln	leu	asp	his	leu	asp	thr	cys	arg	gly	gln	val	met	gly	glu
		his					asn			ala		leu						
		his					asn			ala		leu						
		his					asp			ala		leu			thr			
		his					asp			ala		leu			thr			
		his					asp			ala		leu			thr			
		his					asp			ala		leu			thr			
		his					asp			ala		leu			thr			
		his					asp			ala		leu			leu	thr		
107		110										120						125
glu	asp	ser	glu	leu	gly	asn	met	asp	pro	ile	val	thr	val	lys	lys	tyr	phe	gln
			ala			arg	thr	gly		thr	leu	ala	leu		arg			
			ala			arg	thr	gly		thr	leu	ala	leu		arg			
			ala			arg	thr	gly		thr	leu	ala	met		arg			
			ala			arg	thr	gly		thr	leu	ala	met		thr			
			ala			arg	thr	gly		thr	leu	ala	met		thr			
			ala			arg	thr	gly		thr	leu	ala			ser			
			ala			arg	thr	gly		thr	leu	ala			ser			
126			130									140						144
gly	ile	tyr	asp	tyr	leu	gln	glu	lys	gly	tyr	ser	asp	cys	ala	trp	glu	ile	val
		his	val			lys											thr	
		his	val			lys											thr	
		his	val			lys												
		his	val			lys												
		his	val			lys					!							
		his	ile									!						
		his	val						!				!					
145					150									160				163
arg	val	glu	met	met	arg	ala	leu	thr	val	ser	thr	thr	leu	gln	lys	arg	leu	thr
	leu		ile			ser	phe	ser	ser	leu	ile	ser			glu			arg
			ile			ser	phe	ser	ser	leu	ile	ser			glu			arg
	leu		ile			ser		ser	ser			ser		his				arg
164								172										
lys	met	gly	gly	asp	leu	asn	ser	pro	<u>stop</u>									
met		asp				ser			<u>stop</u>									
met		asp				ser			<u>stop</u>									
met		asp				ser			<u>stop</u>									

Fig. 20B




Fig. 21



# ANTIVIRAL THERAPY USING OVINE OR BOVINE INTERFERON-TAU

This application is a continuation of patent application Ser. No. 08/438,753, filed May 10, 1995, now U.S. Pat. No. 5,705,363, herein incorporated by reference, which is a continuation-in-part of patent application Ser. No. 08/139,891, filed Oct. 19, 1993, now abandoned, incorporated herein by reference, which is a continuation-in-part of patent application Ser. No. 07/847,741, filed Mar. 9, 1992, now abandoned, which is a continuation-in-part of application Ser. No. 07/318,050, filed Mar. 2, 1989, now abandoned. Application Ser. No. 08/139,891 is also a continuation-in-part of patent application Ser. No. 07/969,890, filed Oct. 30, 1992 and now abandoned.

This invention was made with government support under National Institutes of Health grants HD 10436, HD 26006, CA 38587, and CA 57084. Accordingly, the United States government has certain rights in this invention.

## FIELD OF THE INVENTION

The present invention relates to interferon- $\tau$  compositions and methods of use.

## REFERENCES

- Abraham, G., et al., *Clin. Immunol. Immunopathol.* 56:1-8 (1990).
- Ausubel, F. M., et al., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John Wiley & Sons, Inc., Media, PA.
- Bayne, M. L., et al., *Gene* 66:235 (1988).
- Bazer, F. W., et al., *J. Animal Sci.* 57(Supp. 2):425 (1983).
- Bazer, F. W., et al., *J. Reproduction and Fertility* 76:841 (1986).
- Bazer, F. W., et al., *Biology of Reproduction*, vol.40; (Supplement 1):63, (Abstract) (1989).
- Beames, et al., *Biotechniques* 11:378 (1991).
- Benoit, P., et al., *J. Immunol.* 150(3):707 (1993).
- Bonnem, E. M., et al., *J. Bio. Response Modifiers* 3:580 (1984).
- Boyer, S. J., et al., *J. Biol. Regul. Homeost. Agents.* 6(3):99-102 (1992).
- Chan, et al., *DNA* 4:165 (1985).
- Crea, R., U.S. Pat. No. 4,888,286, issued Dec. 19, 1989.
- Crowe, S. M., et al., *AIDS Res. Hum. Retroviruses* 3(2):135 (1987).
- Cumber, J. A., et al., *Methods in Enzymology*, 112:207 (1985).
- Dalgleish, et al., *Nature* 312:763 (1984).
- Davis, G. L., et al., *N. England J. Med.* 321:1501 (1989).
- Davis, G. L., et al., *Theriogenology* 38:867 (1992).
- Degre, M., *Int. J. Cancer* 14:699-703 (1974).
- DeMaeyer, E., et al., *Interferons and Other Regulatory Cytokines*, John Wiley and Sons, New York (1988).
- Dianzani, F., *J. Interferon Res., Special Issue*, 5/92:109 (1992).
- Duncan, R. J. S., et al., *Anal Biochem*, 182:68 (1983).
- Dusheiko, G. M., et al., *J. Hematology* 3(Supl. 2):S199 (1986).
- Eaton, M. A. W., et al., U.S. Pat. No. 4,719,180, issued Jan. 12, 1988.
- Ecker, D. J., et al., *J. Biol. Chem.* 264:7715-7719 (1989).
- Elliot, S., et al., *J. Biol. Chem.* 261:2936 (1986).
- Ernst, J. F., *DNA* 5:483 (1986).
- Familetti, P. C., et al., *Meth. Enzymol.* 78:387 (1981).
- Feher, Z., et al., *Curr. Genet.* 16:461 (1989).

- Fent, K., and Zbinden, G., *Trends Pharm. Sci.* 8:100-105 (1987).
- Figuro, F., et al., *Immunogenetics* 15:431 (1982).
- Finter, N. B., et al., *Drugs* 42(5):749 (1991).
- Foa, P., et al., *Cell Tissue Kinet.* 15(4):399-404 (1982).
- Francis, M. L., et al., *AIDS Res. and Human Retroviruses* 8(2):199 (1992).
- Frangioni, J. V., et al., *Anal. Biochem.* 210(1):179-187 (1993).
- Fritz, R. B., et al., *J. Immunol.* 134:2328 (1985).
- Godkin, J. D., et al., *J. Reprod. Fert.* 65:141 (1982).
- Griggs, N. D., et al., *J. Immunol.* 149:517 (1992).
- Guan, K. L., et al., *Anal. Biochem.* 192(2):262-267 (1991).
- Haahr, S., et al., *The Lancet* 337:863-864 (1991).
- Hakes, D. J., et al., *Anal. Biochem.* 202(2):293-298 (1992).
- Hansen, P. J., et al., U.S. Pat. No. 4,997,646, issued Mar. 5, 1991.
- Harlow, E., et al., *Antibodies: A Laboratory Manual*, Cold Spring Harbor Press, Cold Spring Harbor, N.Y. (1988).
- Helmer, S. D., et al., *J. Reprod. Fert.* 79:83-91 (1987).
- Hitzeman, R. A., et al., U.S. Pat. No. 4,775,622, issued Oct. 4, 1988.
- Hoffman, A. E., et al., *Virology* 147:326 (1985).
- Howatson, et al., *J. Endocrinol.* 119:531 (1988).
- Imakawa, K., et al., *Nature* 330:377 (1987).
- Imakawa, K., et al., *Mol. Endocrinol.* 3:127-139 (1989).
- Innis, M., U.S. Pat. No. 4,975,276, issued Dec. 4, 1990.
- Johnson, W. C., Jr., *Methods in Enzymology*, Vol. 210, pp. 426-447, (1992).
- Kashima, H., et al., *Laryngoscope* 98:334 (1988).
- Klein, J., et al., *Immunogenetics* 17:553 (1983).
- Kotzin, B. L., et al., *J. Exp. Med.* 165:1237 (1987).
- Krown, S. E., in "Mechanisms of Interferon Actions", (Pfeffer, L. M., ed.), *CRC Press Inc.*, Boca Raton, Vol. II, pp. 143-178, (1987).
- Kyte, J., et al., *J. Mol. Biol.* 157:105 (1982).
- Langford, M. P., et al., *Meth. Enzymol.* 78:339 (1981).
- Lawrence, et al., *Nucl. Acids. Res.* 13:1777 (1985).
- Lowry, O. H., et al., *J. Biol. Chem.* 193:265-275 (1951).
- Ludwig, D. L., et al., *Gene* 132:33 (1993).
- Maniatis, T., et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory (1982).
- Martin, E. W., "In: Dispensing of medication: a practical manual on the formulation and dispensing of pharmaceutical products", (Hoover, J. E. ed.), 8th edition, Mack Publishing Co., Easton, Pa., (1976).
- Meyer, F., et al., U.S. Pat. No. 4,885,166, issued Dec. 5, 1989.
- McInnes, B., et al., *J. Interferon Res.* 9(3), pp. 305-314, (1989).
- Mullis, K. B., U.S. Pat. No. 4,683,202, issued Jul. 28, 1987.
- Mullis, K. B., et al., U.S. Pat. No. 4,683,195, issued Jul. 28, 1987.
- Miyoshi, E., et al., *Int. J. Cancer* 52(1):137-140 (1992).
- Nagy, et al., *Int. J. Canc.* 32:321 (1983).
- Oeda, K., et al., U.S. Pat. No. 4,766,068, issued Aug. 23, 1988.
- Oldham, R. K., *Hospital Practice* 20:71 (1985).
- Paulesu, et al., *J. Biol. Regul. Homeost. Agents* 5:81 (1991).
- Pearson, W. R. and Lipman, D. J., *PNAS* 85:2444-2448 (1988).
- Pearson, W. R., *Methods in Enzymology* 183:63-98 (1990).
- Pederson, et al., *Science* 235:790 (1987).
- Perczel, A., et al., *Protein Engineering*, Vol. 4, (Supp. 6), pp. 669-679, (1991).
- Perron, H., et al., *The Lancet* 337:862-863 (1991).
- Pontzer, et al., *Biochem. Biophys. Res. Comm.* 152:801 (1988).



- Pontzer, C. H., et al., *Cancer Res.* 51:5304 (1991).
- Poste, G., et al., *Proc. Nat'l Acad. Sci., USA*, Vol. 78:6226 (1981).
- Quesada, J. R., et al., *N. England J. Med.* 310:15 (1984).
- Rasmussen, H. B., et al., *Acta. Neurol. Scand.* 88:190-198 (1993).
- Reilly, P. R., et al., *Baculovirus Expression Vectors: A Laboratory Manual* (1992).
- Roberts, R. M., et al., *Endocrine Reviews* 13:432 (1992).
- Rothstein, R., *DNA Cloning: A Practical Approach*, vol. II (Glover, D. M., Ed.) Oxford: IRL Press, pp. 46-66 (1986).
- Ruegg, C. L., et al., *J. Interferon Res.*, 10:621-626, (1990).
- Rutter, W. J., et al., U.S. Pat. No. 4,769,238, issued Sept. 6, 1988.
- Sabin, E., et al., *Bio/Technology* 7:705-709 (1989).
- Sambrook, J., et al., *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory, Cold Spring Harbor, N.Y.) (1989).
- Sanger, et al., *Proc. Natl. Acad. Sci. USA* 74:5463 (1977).
- Senda, T., et al., *EMBO J.*, Vol. 11, Supp. 9, pp. 3193-3201, (1992).
- Shaw, K. J., et al., *DNA* 7:117 (1988).
- Shen, L. P., et al., *Sci. Sin.* 29:856 (1986).
- Shoulders, C. C., et al., *Nucleic Acids Res.* 10(16):4873-4882 (1982).
- Smith, D. B., et al., *Gene* 67:31 (1988).
- Smith, P. K., et al., *Anal. Biochem.* 150:76 (1985).
- Stewart, H. J., et al., *J. Endocrinol.* 115:R13 (1987).
- Talal, N., et al., *Annals of Allergy* 69:221-224 (1992).
- Todd, R. F., et al., *Leuk. Res.* 5(6):491-495 (1981).
- Vallet, J. L., et al., *Biol. Reprod.* 37:1307 (1987).
- Vallet, J. L., et al., *J. Endocrinology* 117:R5-R8 (1988).
- Venables, P., et al., *Brit. J. Rheumatology* 31:841-846 (1992).
- von Eichborn, J. F., et al., U.S. Pat. No. 5,145,677, issued Sep. 8, 1992.
- Wallis, S. C., et al., *EMBO J.* 2(12):2369-2373 (1983).
- Whaley, A. E., et al., *J. Biol. Chem.* 269:10846-10868 (1994).
- Wang, C. Y., et al., U.S. Pat. No. 4,879,212, issued Nov. 7, 1989.
- Wang, C. Y., et al., U.S. Pat. No. 4,735,896, issue Apr. 5, 1988.
- Wilson, et al., *Biology of Reproduction* 20(Supp. 1):101A, Abstract (1979).
- Wraith, D. C., et al., *Cell* 59:247 (1989).
- Wu, D. A., et al., *DNA* 10:201 (1991).
- Yoshio, T., et al., U.S. Pat. No. 4,849,350, issued Jul. 18, 1989.
- Young, et al., *Proc. Natl. Acad. Sci. USA* 80:1194 (1983).
- Zamvil, S. S., et al., *Nature* 317:355 (1985).

### BACKGROUND OF THE INVENTION

Conceptus membranes, or trophoctoderm, of various mammals produce biochemical signals that allow for the establishment and maintenance of pregnancy (Bazer, et al., 1983). One such protein, ovine trophoblast protein-one (oTP-1), was identified as a low molecular weight protein secreted by sheep conceptuses between days 10 and 21 of pregnancy (Wilson, et al., 1979; Bazer, et al., 1986). The protein oTP-1 was shown to inhibit uterine secretion of prostaglandin F<sub>2</sub>-alpha, which causes the corpus luteum on the ovary to undergo physiological and endocrinological demise in nonpregnant sheep (Bazer, et al., 1986). Accordingly, oTP-1 has antiluteolytic biological activity. The primary role of oTP-1 was assumed to be associated with the establishment of pregnancy.

oTP-1 was subsequently found to (i) exhibit limited homology (50-70%) with interferon alphas (IFN $\alpha$ ) of various species (Imakawa, et al., 1987), and (ii) bind to a Type I interferon receptor (Stewart, et al., 1987). Despite some similarities with IFN $\alpha$ , oTP-1 has several features that distinguish it from IFN $\alpha$  including the following: oTP-1's role in reproductive biochemistry (other interferons are not known to have any role in the biochemical regulation of reproductive cycles), oTP-1's cellular source—trophoblast cells (IFN $\alpha$  is derived from lymphocyte cells), oTP-1's size—172 amino acids (IFN $\alpha$  is typically about 166 amino acids), and oTP-1 is weakly inducible by viruses (IFN $\alpha$  is highly inducible by viruses). The International Interferon Society recognizes oTP-1 as belonging to an entirely new class of interferons which have been named interferon-tau (IFN $\tau$ ). The Greek letter  $\tau$  stands for trophoblast.

The interferons have been classified into two distinct groups: type I interferons, including IFN $\alpha$ , IFN $\beta$ , and IFN $\omega$  (also known as IFN $\alpha$ II); and type II interferons, represented by IFN $\gamma$  (reviewed by DeMaeyer, et al.). In humans, it is estimated that there are at least 17 IFN $\alpha$  non-allelic genes, at least about 2 or 3 IFN $\beta$  non-allelic genes, and a single IFN $\gamma$  gene.

IFN $\alpha$ 's have been shown to inhibit various types of cellular proliferation. IFN $\alpha$ 's are especially useful against hematologic malignancies such as hairy-cell leukemia (Quesada, et al., 1984). Further, these proteins have also shown activity against multiple myeloma, chronic lymphocytic leukemia, low-grade lymphoma, Kaposi's sarcoma, chronic myelogenous leukemia, renal-cell carcinoma, urinary bladder tumors and ovarian cancers (Bonnem, et al., 1984; Oldham, 1985). The role of interferons and interferon receptors in the pathogenesis of certain autoimmune and inflammatory diseases has also been investigated (Benoit, et al., 1993).

IFN $\alpha$ 's are also useful against various types of viral infections (Finter, et al., 1991). Alpha interferons have shown activity against human papillomavirus infection, Hepatitis B, and Hepatitis C infections (Finter, et al., 1991; Kashima, et al., 1988; Dusheiko, et al., 1986; Davis, et al., 1989).

Significantly, however, the usefulness of IFN $\alpha$ 's has been limited by their toxicity: use of interferons in the treatment of cancer and viral disease has resulted in serious side effects, such as fever, chills, anorexia, weight loss, and fatigue (Pontzer, et al., 1991; Oldham, 1985). These side effects often require (i) the interferon dosage to be reduced to levels that limit the effectiveness of treatment, or (ii) the removal of the patient from treatment. Such toxicity has reduced the usefulness of these potent antiviral and antiproliferative proteins in the treatment of debilitating human and animal diseases.

### SUMMARY OF THE INVENTION

In a first aspect, the present invention relates to compositions of and methods employing ovine interferon- $\tau$ . The invention includes an isolated nucleic acid molecule that encodes an ovine interferon- $\tau$ . One embodiment of this nucleic acid molecule is a nucleic acid molecule having the sequence presented as SEQ ID NO:1. In another embodiment, the nucleic acid molecule encodes an ovine interferon- $\tau$  polypeptide having a sequence presented as SEQ ID NO:2. The ovine interferon- $\tau$  polypeptide may include an amino-terminal extension, such as, a leader sequence.

In another embodiment, the present invention includes an expression vector having a nucleic acid containing an open



reading frame (ORF) that encodes an ovine interferon- $\tau$ , including the nucleic acid and polypeptide sequences described above. The vector further includes regulatory sequences effective to express the open reading frame in a host cell. Further, the invention includes a method of recombinantly producing ovine interferon- $\tau$  using the expression vectors of the present invention. The expression vectors are introduced into suitable host cells. The host cells are then cultured under conditions that result in the expression of the ORF sequence.

In one embodiment, the present invention includes a recombinantly produced ovine interferon- $\tau$  protein.

Further, the invention includes a method of inhibiting tumor cell growth. In the method, the tumor cells are contacted with ovine interferon- $\tau$  at a concentration effective to inhibit growth of the tumor cells. Target tumor cells include, but are not limited to carcinoma cells, hematopoietic cancer cells, leukemia cells, lymphoma cells and melanoma cells.

The invention also includes a method of inhibiting viral replication. In this method, cells infected with a virus are contacted with ovine interferon- $\tau$  at a concentration effective to inhibit viral replication within said cells. Ovine interferon- $\tau$  may be used to inhibit the replication of both RNA and DNA viruses. Exemplary RNA viruses include feline leukemia virus, ovine progressive pneumonia virus, ovine lentivirus, equine infectious anemia virus, bovine immunodeficiency virus, visna-maedi virus, and caprine arthritis encephalitis virus.

In a second aspect, the present invention relates to compositions of and methods employing human interferon- $\tau$ 's. In one embodiment, the invention includes an isolated nucleic acid molecule that encodes a human interferon- $\tau$ . Several variants of human interferon- $\tau$  (HuIFN $\tau$ ) are disclosed herein, including HuIFN $\tau$ 1, HuIFN $\tau$ 2, HuIFN $\tau$ 3, HuIFN $\tau$ 4, HuIFN $\tau$ 5, HuIFN $\tau$ 6 and HuIFN $\tau$ 7. The nucleic acid molecules of the present invention include nucleic acid molecules having the following sequences: SEQ ID NO:43, SEQ ID NO:29, SEQ ID NO:25, SEQ ID NO:33, SEQ ID NO:27, SEQ ID NO:21 and SEQ ID NO:23.

The nucleic acids of the present invention also include nucleic acid molecules encoding the following polypeptide sequences: SEQ ID NO:44, SEQ ID NO:30, SEQ ID NO:34, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:22, and SEQ ID NO:24. The nucleic acids may further include sequences encoding leader sequences for the human interferon- $\tau$  which they encode, for example, SEQ ID NO:41 or SEQ ID NO:42.

The second aspect of the invention further includes an expression vector having a nucleic acid sequence containing an open reading frame that encodes a human interferon- $\tau$ , including the nucleic acid and polypeptide sequences described above. The vector further includes regulatory sequences effective to express said open reading frame in a host cell. The regulatory sequence may include sequences useful for targeting or secretion of the human IFN $\tau$  polypeptide: such sequences may be endogenous (such as the normally occurring human IFN $\tau$  leader sequences, present, for example, in SEQ ID NO:41) or heterologous (such as a secretory signal recognized in yeast, mammalian cell, insect cell, tissue culture or bacterial expression systems). In the expression vector, regulatory sequences may also include, 5' to said nucleic acid sequence, a promoter region and an ATG start codon in-frame with the human interferon- $\tau$  coding sequence, and 3' to said coding sequence, a translation termination signal followed by a transcription termination signal.

In a further embodiment, the invention includes a method of recombinantly producing human interferon- $\tau$ . In the method, the expression vector, containing sequences encoding a human interferon- $\tau$  open reading frame (ORF), is introduced into suitable host cells, where the vector is designed to express the ORF in the host cells. The transformed host cells are then cultured under conditions that result in the expression of the ORF sequence. Numerous vectors and their corresponding hosts are useful in the practice of this method of the invention, including, lambda gt11 phage vector and *E. coli* cells. Other host cells include, yeast, mammalian cell, insect cell, tissue culture, plant cell culture, transgenic plants or bacterial expression systems.

In another embodiment, the invention includes an isolated human interferon- $\tau$  protein or polypeptide. The protein may be recombinantly produced. Further, the protein or polypeptide may include any of the following human interferon- $\tau$  sequences: SEQ ID NO:44, SEQ ID NO:30, SEQ ID NO:34, SEQ ID NO:26, SEQ ID NO:28, SEQ ID NO:22, and SEQ ID NO:24.

The invention further includes a method of inhibiting tumor cell growth. In the method, the tumor cells are contacted with a human interferon- $\tau$  polypeptide at a concentration effective to inhibit growth of the tumor cells. The human interferon- $\tau$  may be a part of any acceptable pharmacological formulation. Tumor cells whose growth may be inhibited by human interferon- $\tau$  include, but are not limited to, human carcinoma cells, hematopoietic cancer cells, human leukemia cells, human lymphoma cells, and human melanoma cells. In one embodiment, the tumor cells are steroid-sensitive tumor cells, for example, mammary tumor cells.

In yet another embodiment of the present invention, human interferon- $\tau$  polypeptides are used in a method of inhibiting viral replication. In this method, cells infected with a virus are contacted with human interferon- $\tau$  at a concentration effective to inhibit viral replication within said cells. The human interferon- $\tau$  may be a part of any acceptable pharmacological formulation. The replication of both RNA and DNA viruses may be inhibited by human interferon- $\tau$  polypeptides. Exemplary RNA viruses include human immunodeficiency virus (HIV) or hepatitis c virus (HCV). An exemplary DNA virus is hepatitis B virus (HBV).

In yet another aspect, the present invention includes a method of enhancing fertility in a female mammal. In this method, an effective mammalian fertility enhancing amount of human interferon- $\tau$  is administered to the female mammal in a pharmaceutically acceptable carrier.

The invention also includes isolated human interferon- $\tau$  polypeptides. These polypeptides are derived from the interferon- $\tau$  amino acid sequence and are typically between about 15 and 172 amino acids in length.

Also included in the invention is a fusion polypeptide that contains a human interferon- $\tau$  polypeptide that is between 15 and 172 amino acids long and derived from a human interferon- $\tau$  amino acid coding sequence, and a second soluble polypeptide. In one embodiment, human interferon- $\tau$  sequences are used in fusion constructs with other types of interferons to reduce the toxicity of the other types of interferons, for example, interferon- $\alpha$  and interferon- $\beta$ .

The invention also includes a polypeptide composition having (a) a human interferon- $\tau$  polypeptide, where said polypeptide is (i) derived from an interferon- $\tau$  amino acid coding sequence, and (ii) between 15 and 172 amino acids



long, and (b) a second soluble polypeptide. Interferon- $\alpha$  and interferon- $\beta$  are examples of such second soluble polypeptides. This composition may be used to reduce the toxicity of the other types of interferons when the interferons are used in pharmaceutical formulations or in therapeutic applications.

The invention also includes purified antibodies that are immunoreactive with human interferon- $\tau$ . The antibodies may be polyclonal or monoclonal.

These and other objects and features of the invention will be more fully appreciated when the following detailed description of the invention is read in conjunction with the accompanying drawings.

#### BRIEF DESCRIPTION OF THE FIGURES

FIGS. 1A and 1B present the nucleic acid coding sequence of a synthetic gene of OvIFN $\tau$  designed to include 19 unique restriction enzyme sites spaced evenly throughout the coding sequence.

FIG. 2 shows the cloning strategy used for making a synthetic gene encoding OvIFN $\tau$ .

FIG. 3 shows a comparison of the predicted protein sequences of a human interferon- $\tau$  gene and an ovine interferon- $\tau$  gene. Divergent amino acids are indicated by presentation of the alternative amino acid on the line below the nucleic acid sequences.

FIG. 4 presents data demonstrating that both OvIFN $\tau$  and IFN $\alpha$  were able to drastically reduce growth of HL-60 cells.

FIG. 5 presents data demonstrating that rHuIFN $\alpha$  is cytotoxic and OvIFN $\tau$  is not. In the figure, results of one of three replicate experiments are presented as mean % viability  $\pm$  SD.

FIG. 6 presents the sequences of polypeptides derived from the IFN $\tau$  sequence.

FIG. 7 presents the complete nucleic acid and amino acid sequence of an OvIFN $\tau$  sequence.

FIG. 8 presents data supporting the lack of cytotoxicity, relative to IFN $\alpha$ , when IFN $\tau$  is used to treat peripheral blood mononuclear cells.

FIG. 9 shows the results of treatment of a human cutaneous T cell lymphoma line, HUT 78, with IFN $\tau$ .

FIG. 10 shows the results of treatment of a human T cell lymphoma line, H9, with IFN $\tau$ .

FIG. 11A presents data for the peptide inhibition, relative to FIV (feline immunodeficiency virus) replication, of polypeptides derived from OvIFN $\tau$  with whole OvIFN $\tau$ .

FIG. 11B presents data for the peptide inhibition, relative to HIV (human immunodeficiency virus) replication, of polypeptides derived from OvIFN $\tau$  with whole OvIFN $\tau$ .

FIG. 12 presents data demonstrating the inhibition of the antiviral activity of IFN $\tau$  by IFN $\tau$ -derived peptides.

FIG. 13 presents data demonstrating the inhibition by IFN $\tau$ -derived peptides of OvIFN $\tau$  antiviral activity.

FIG. 14 presents data demonstrating the inhibition by IFN $\tau$ -derived peptides of bovine IFN $\alpha$  antiviral activity.

FIG. 15 presents data demonstrating the inhibition by IFN $\tau$ -derived peptides of human IFN $\alpha$  antiviral activity.

FIG. 16 presents data evaluating the lack of inhibition by IFN $\tau$ -derived peptides of bovine IFN $\gamma$  antiviral activity.

FIG. 17 presents data demonstrating the anti-IFN $\tau$ -derived peptide antisera inhibition of the antiviral activity of IFN $\tau$ .

FIG. 18 presents data demonstrating the anti-IFN $\tau$ -derived peptide antisera inhibition of the binding of radio-labeled IFN $\tau$  to cells.

FIGS. 19A and 19B present an alignment of nucleic acid sequences encoding IFN $\tau$  polypeptides.

FIGS. 20A and 20B present an alignment of amino acid sequences of IFN $\tau$  polypeptides.

FIG. 21 presents data comparing the cytotoxicity of IFN $\tau$  with IFN $\beta$ .

#### BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is the nucleotide sequence of a synthetic gene encoding ovine interferon- $\tau$  (OvIFN $\tau$ ). Also shown is the encoded amino acid sequence.

SEQ ID NO:2 is an amino acid sequence of a mature OvIFN $\tau$  protein.

SEQ ID NO:3 is a synthetic nucleotide sequence encoding a mature human interferon- $\tau$  (HuIFN $\tau$ ) protein.

SEQ ID NO:4 is an amino acid sequence for a mature HuIFN $\tau$ 1 protein.

SEQ ID NO:5 is the amino acid sequence of fragment 1–37 of SEQ ID NO:2.

SEQ ID NO:6 is the amino acid sequence of fragment 34–64 of SEQ ID NO:2.

SEQ ID NO:7 is the amino acid sequence of fragment 62–92 of SEQ ID NO:2.

SEQ ID NO:8 is the amino acid sequence of fragment 90–122 of SEQ ID NO:2.

SEQ ID NO:9 is the amino acid sequence of fragment 119–150 of SEQ ID NO:2.

SEQ ID NO:10 is the amino acid sequence of fragment 139–172 of SEQ ID NO:2.

SEQ ID NO:11 is the nucleotide sequence of a natural HuIFN $\tau$ 1 gene with a leader sequence.

SEQ ID NO:12 is the predicted amino acid coding sequence of the SEQ ID NO:11.

SEQ ID NO:13 is a 25-mer synthetic oligonucleotide according to the subject invention.

SEQ ID NO:14 is a 25-mer synthetic oligonucleotide according to the subject invention.

SEQ ID NO:15 is the amino acid sequence of fragment 1–37 of SEQ ID NO:4.

SEQ ID NO:16 is the amino acid sequence of fragment 34–64 of SEQ ID NO:4.

SEQ ID NO:17 is the amino acid sequence of fragment 62–92 of SEQ ID NO:4.

SEQ ID NO:18 is the amino acid sequence of fragment 90–122 of SEQ ID NO:4.

SEQ ID NO:19 is the amino acid sequence of fragment 119–150 of SEQ ID NO:4.

SEQ ID NO:20 is the amino acid sequence of fragment 139–172 of SEQ ID NO:4.

SEQ ID NO:21 is the nucleotide sequence of cDNA HuIFN $\tau$ 6.

SEQ ID NO:22 is the predicted amino acid sequence encoded by the sequence represented as SEQ ID NO:21.

SEQ ID NO:23 is the nucleotide sequence of cDNA HuIFN $\tau$ 7.

SEQ ID NO:24 is the predicted amino acid sequence encoded by the sequence represented as SEQ ID NO:23.

SEQ ID NO:25 is the nucleotide sequence of cDNA HuIFN $\tau$ 4.

SEQ ID NO:26 is the predicted amino acid sequence encoded by the sequence represented as SEQ ID NO:25.

SEQ ID NO:27 is the nucleotide sequence of cDNA HuIFN $\tau$ 5.



SEQ ID NO:28 is the predicted amino acid sequence encoded by the sequence represented as SEQ ID NO:27.

SEQ ID NO:29 is the nucleotide sequence of genomic DNA clone HuIFN $\tau$ 2.

SEQ ID NO:30 is the predicted amino acid sequence encoded by the sequence represented as SEQ ID NO:29.

SEQ ID NO:31 is the nucleotide sequence, including leader sequence, of genomic DNA clone HuIFN $\tau$ 3, a natural HuIFN $\tau$  gene.

SEQ ID NO:32 is the predicted amino acid sequence (including leader sequence) encoded by the sequence represented as SEQ ID NO:31.

SEQ ID NO:33 is the nucleotide sequence, excluding leader sequence, of genomic DNA clone HuIFN $\tau$ 3, a natural HuIFN $\tau$  gene.

SEQ ID NO:34 is the predicted amino acid sequence of a mature human IFN $\tau$  protein encoded by HuIFN $\tau$ 3, encoded by the sequence represented as SEQ ID NO:33.

SEQ ID NO:35 is the amino acid sequence of fragment 1–37 of SEQ ID NO:33.

SEQ ID NO:36 is the amino acid sequence of fragment 34–64 of SEQ ID NO:33.

SEQ ID NO:37 is the amino acid sequence of fragment 62–92 of SEQ ID NO:33.

SEQ ID NO:38 is the amino acid sequence of fragment 90–122 of SEQ ID NO:33.

SEQ ID NO:39 is the amino acid sequence of fragment 119–150 of SEQ ID NO:33.

SEQ ID NO:40 is the amino acid sequence of fragment 139–172 of SEQ ID NO:33.

SEQ ID NO:41 is the amino acid sequence of fragment 1–23 of SEQ ID NO:32.

SEQ ID NO:42 is the amino acid sequence of fragment 1–23 of SEQ ID NO:11.

SEQ ID NO:43 is the nucleotide sequence, excluding leader sequence, of DNA clone HuIFN $\tau$ 1.

SEQ ID NO:44 is the predicted amino acid sequence of a mature human IFN $\tau$  protein encoded by HuIFN $\tau$ 1, encoded by the sequence represented as SEQ ID NO:43.

## DETAILED DESCRIPTION OF THE INVENTION

### I. Definitions.

Interferon- $\tau$  (IFN $\tau$ ) refers to any one of a family of interferon proteins having greater than 70%, or preferably greater than about 80%, or more preferably greater than about 90% amino acid homology to either the sequence presented as (a) SEQ ID NO:2 or (b) SEQ ID NO:34. Amino acid homology can be determined using, for example, the LALIGN program with default parameters. This program is found in the FASTA version 1.7 suite of sequence comparison programs (Pearson, et al., 1988; Pearson, 1990; program available from William R. Pearson, Department of Biological Chemistry, Box 440, Jordan Hall, Charlottesville, Va.). Typically, IFN $\tau$  has at least one characteristic from the following group of characteristics: (a) expressed during embryonic/fetal stages by trophectoderm/placenta, (b) anti-luteolytic properties, (c) anti-viral properties, and (d) anti-cellular proliferation properties. IFN $\tau$  can be obtained from a number of sources including cows, sheep, ox, and humans.

An interferon- $\tau$  polypeptide is a polypeptide having between about 15 and 172 amino acids derived from an interferon- $\tau$  amino acid coding sequence, where said 15 to

172 amino acids are contiguous in native interferon- $\tau$ . Such 15–172 amino acid regions can also be assembled into polypeptides where two or more such interferon- $\tau$  regions are joined that are normally discontinuous in the native protein.

### II. Isolation & Characterization of Interferon- $\tau$ .

#### A. Ovine and Bovine Interferon- $\tau$ .

##### 1. Interferon- $\tau$ Coding Sequences.

Ovine interferon- $\tau$  (OvIFN $\tau$ ) is a major conceptus secretory protein produced by the embryonic trophectoderm during the critical period of maternal recognition in sheep. One isolate of mature OvIFN $\tau$  is 172 amino acids in length (SEQ ID NO:2). The cDNA coding sequence contains an additional 23 amino acids at the amino-terminal end of the mature protein (Imakawa, et al., 1987). The coding sequence of this OvIFN $\tau$  isolate is presented as FIG. 7.

Relative to other interferons, oIFN $\tau$  shares about 45 to 68% amino acid homology with Interferon- $\alpha$  and the greatest sequence similarity with the interferon- $\omega$ s (IFN $\omega$ s) of about 68%.

For the isolation of OvIFN $\tau$  protein, conceptuses were collected from pregnant sheep and cultured in vitro in a modified Minimum Essential Medium as described previously (Godkin, et al., 1982). Conceptuses were collected on various days of pregnancy with the first day of mating being described as Day 0. OvIFN $\tau$  was purified from conceptus culture medium essentially as described by Vallet, et al., (1987) and Godkin, et al. (1982).

The homogeneity of OvIFN $\tau$  was assessed by sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE; Maniatis, et al.; Ausubel, et al.). Determination of protein concentration in purified OvIFN $\tau$  samples was performed using the bicinchoninic (BCA) assay (Pierce Chemical Co., Rockford, Ill.; Smith, et al., 1985).

A homologous protein to OvIFN $\tau$  was isolated from cows (BoIFN $\tau$ ; Helmer, et al., 1987; Imakawa, et al., 1989). OvIFN $\tau$  and BoIFN $\tau$  (i) have similar functions in maternal recognition of pregnancy, and (ii) share a high degree of amino acid and nucleotide sequence homology between mature proteins. The nucleic acid sequence homology between OvIFN $\tau$  and BoIFN $\tau$  is 76.3% for the 5' non-coding region, 89.7% for the coding region, and 91.9% for the 3' non-coding region. The amino acid sequence homology is 80.4%.

Example 1 describes the reproductive functions of OvIFN $\tau$ . OvIFN $\tau$  and recombinant human Interferon- $\alpha$ 2 (rHuIFN $\alpha$ ) were infused into uterine lumen of ewes at a variety of concentrations. The life span of the corpus luteum was assessed by examination of interestrous intervals, maintenance of progesterone secretion, and inhibition of prostaglandin secretion (Davis, et al., 1992). Comparison of the data resulting from these examinations demonstrated a considerable lengthening of the interestrous interval when OvIFN $\tau$  is administered at 100  $\mu$ g/day and no meaningful effect when rHuIFN $\alpha$  is administered. These data support the conclusion that OvIFN $\tau$  significantly influences the biochemical events of the estrous cycle.

The antiviral properties of interferon- $\tau$  at various stages of the reproductive cycle were also examined (Example 2). Conceptus cultures were established using conceptus obtained from sheep at days 12 through 16 of the estrus cycle. Antiviral activity of supernatant from each conceptus culture was assessed. Culture supernatants had increasing antiviral activity associated with advancing development of the conceptus up to Day 16 post estrus.



## 2. Recombinant Production of IFN $\tau$

Recombinant OvIFN $\tau$  was produced using bacterial and yeast cells. The amino acid coding sequence for OvIFN $\tau$  was used to generate a corresponding DNA coding sequence with codon usage optimized for expression in *E. coli* (Example 3). The DNA coding sequence was synthetically constructed by sequential addition of oligonucleotides. Cloned oligonucleotides were fused into a single polynucleotide using the restriction digestions and ligations outlined in FIG. 2. The polynucleotide coding sequence had the sequence presented as SEQ ID NO:1.

For expression of recombinant OvIFN $\tau$ , this synthetic coding sequence can be placed in a number of bacterial expression vectors: for example, lambda gt11 (Promega, Madison Wis.); pGEX (Smith, et al.); pGEMEX (Promega); and pBS (Stratagene, La Jolla Calif.) vectors. Other bacterial expression vectors containing suitable promoters, such as the T7 RNA polymerase promoter or the tac promoter, may also be used. Cloning of the OvIFN $\tau$  synthetic polynucleotide into a modified pIN III omp-A expression vector is described in Example 3. Production of the OvIFN $\tau$  protein was induced by the addition of IPTG. Soluble recombinant IFN $\tau$  was liberated from the cells by sonication or osmotic fractionation.

The protein can be further purified by standard methods, including size fractionation (column chromatography or preoperative gel electrophoresis) or affinity chromatography (using, for example, anti-OvIFN $\tau$  antibodies (solid support available from Pharmacia, Piscataway N.J.). Protein preparations can also be concentrated by, for example, filtration (Amicon, Danvers, Mass.).

The synthetic OvIFN $\tau$  gene was also cloned into the yeast cloning vector pBS24Ub (Example 4; Sabin, et al.; Ecker, et al.). Synthetic linkers were constructed to permit in-frame fusion of the OvIFN $\tau$  coding sequences with the ubiquitin coding sequences in the vector. The resulting junction allowed in vivo cleavage of the ubiquitin sequences from the OvIFN $\tau$  sequences.

The recombinant plasmid pBS24Ub-IFN $\tau$  was transformed into the yeast *S. cerevisiae*. Transformed yeast cells were cultured, lysed and the recombinant IFN $\tau$  (r-IFN $\tau$ ) protein isolated from the cell lysates.

The amount of r-IFN $\tau$  was quantified by radioimmunoassay. Microsequencing of the purified r-IFN $\tau$  was carried out. The results demonstrated identity with native OvIFN $\tau$  through the first 15 amino acids. The results also confirmed that the ubiquitin/IFN $\tau$  fusion protein was correctly processed in vivo.

Recombinant IFN $\tau$  obtained by this method exhibited antiviral activity similar to the antiviral activity of IFN $\tau$  purified from conceptus-conditioned culture medium.

Other yeast vectors can be used in the practice of the present invention. They include 2 micron plasmid vectors (Ludwig, et al.), yeast integrating plasmids (YIps; e.g., Shaw, et al.), YEP vectors (Shen, et al.), yeast centromere plasmids (YCps; e.g., Ernst), and the like. Preferably, the vectors include an expression cassette containing an effective yeast promoter, such as the MF $\alpha$ 1 promoter (Ernst, Bayne, et al.), GADPH promoter (glyceraldehyde-3-phosphate-dehydrogenase; Wu, et al.), the galactose-inducible GAL10 promoter (Ludwig, et al., Feher, et al., Shen, et al.), or the methanol-regulated alcohol oxidase (AOX) promoter (Tschopp, et al.). The AOX promoter is particularly useful in *Pichia pastoris* host cells (for example, the AOX promoter is used in pHIL and pPIC vectors included in the Pichia expression kit, available from Invitrogen, San Diego, Calif.).

The expression cassette may include additional elements to facilitate expression and purification of the recombinant protein, and/or to facilitate the insertion of the cassette into a vector or a yeast chromosome. For example, the cassette may include a signal sequence to direct secretion of the protein. An exemplary signal sequence suitable for use in a variety of yeast expression vectors is the MF $\alpha$ 1 pre-pro signal sequence (Bayne, et al., Ludwig, et al., Shaw, et al.). Other signal sequences may also be used. For example, the Pho1 signal sequence (Elliot, et al.) is particularly effective in *Pichia Pastoris* and *Schizosaccharomyces pombe* host cells.

Exemplary expression cassettes include (i) a cassette containing (5' to 3') the AOX promoter, the Pho1 signal sequence, and a DNA sequence encoding OvIFN $\tau$ , for expression in *P. pastoris* host cells, and (ii) a cassette containing (5' to 3') the MF $\alpha$ 1 promoter, the MF $\alpha$ 1 pre-pro signal sequence, and a DNA sequence encoding IFN $\tau$ , for expression in *S. cerevisiae* host cells.

Additional yeast vectors suitable for use with the present invention include, but are not limited to, other vectors with regulatable expression (Hitzeman, et al.; Rutter, et al.; Oeda, et al.). The yeast transformation host is typically *Saccharomyces cerevisiae*, however, as illustrated above, other yeast suitable for transformation can be used as well (e.g., *Schizosaccharomyces pombe*, *Pichia pastoris* and the like).

The DNA encoding the IFN $\tau$  polypeptide can be cloned into any number of commercially available vectors to generate expression of the polypeptide in the appropriate host system. These systems include the above described bacterial and yeast expression systems as well as the following: baculovirus expression (Reilly, et al.; Beames, et al.; Clontech, Palo Alto Calif.); plant cell expression, transgenic plant expression (e.g., S. B. Gelvin and R. A. Schilperoot, *Plant Molecular Biology*, 1988), and expression in mammalian cells (Clontech, Palo Alto Calif.; Gibco-BRL, Gaithersburg Md.). These recombinant polypeptides can be expressed as fusion proteins or as native proteins. A number of features can be engineered into the expression vectors, such as leader sequences which promote the secretion of the expressed sequences into culture medium. The recombinantly produced polypeptides are typically isolated from lysed cells or culture media. Purification can be carried out by methods known in the art including salt fractionation, ion exchange chromatography, and affinity chromatography. Immunoaffinity chromatography can be employed, as described above, using antibodies generated based on the IFN $\tau$  polypeptides.

## B. Human Interferon- $\tau$

### 1. Identification and Cloning of Human Genomic Sequences Encoding an Interferon- $\tau$ Protein.

Human genomic DNA was screened for sequences homologous to interferon- $\tau$  (Example 5). Several sequences that hybridized with the OvIFN $\tau$  cDNA probe were identified. Several clones containing partial sequences of human interferon- $\tau$  were then isolated (Example 6). Two synthetic 25-mer oligonucleotides, corresponding to sequences from the OvIFN $\tau$  cDNA (Imakawa, et al., 1987; Whaley, et al., 1994) were synthesized. These primers were employed in amplification reactions using DNA derived from the following two cDNA libraries: human placenta and human cytotrophoblast cells isolated from term placenta. The resulting amplified DNA fragments were electrophoretically separated and a band containing human IFN $\tau$  amplification products was isolated. The amplification products were subcloned and the inserted amplification products sequenced using the dideoxy termination method.



Comparison of sequences from five of these clones revealed a high degree of sequence homology between the isolates, but the sequences were not identical. This result suggests the existence of multiple variants of human interferon- $\tau$  genes. Analysis of the nucleotide and protein sequences suggests that human interferon- $\tau$  genes may be classified on the basis of sequence homology into at least three groups. The groups are presented below.

Example 7 describes the isolation of several full-length human IFN $\tau$  genes. High molecular weight DNA was isolated from human peripheral blood mononuclear cells (PBMCs) and size-fractionated. Fractions were tested for the presence of IFN $\tau$  sequences using polymerase chain reaction: DNA molecules from fractions that tested amplification positive were used to generate a subgenomic library in  $\lambda$ gt11.

This subgenomic library was plated and hybridized with an OvIFN $\tau$  cDNA probe (Example 7A). Approximately 20 clones were identified that hybridized to the probe. Plaques corresponding to the positive clones were passaged, DNA isolated and analyzed by amplification reactions using OvIFN $\tau$  primers. Of these twenty plaques, six plaques generated positive PCR signals. The phage from these six clones were purified and the inserts sequenced. One of the inserts from one of these six clones was used as a hybridization probe in the following screening.

Recombinant phage from the  $\lambda$ gt11 subgenomic library were screened using the hybridization probe just described (Example 7B). Five clones giving positive hybridization signals were isolated and the inserts sequenced. The sequences from three of the clones overlapped, and the resulting consensus nucleic acid sequence (HuIFN $\tau$ 1) is presented as SEQ ID NO:11 with the predicted protein coding sequence presented as SEQ ID NO:12. The predicted mature protein coding sequence is presented as SEQ ID NO:4. The sequences from the other two clones are presented as SEQ ID NO:29 (HuIFN $\tau$ 2) and SEQ ID NO:31 (HuIFN $\tau$ 3). The predicted mature amino acid sequence from HuIFN $\tau$ 2 is presented as SEQ ID NO:30. The predicted amino acid sequence from HuIFN $\tau$ 3 is presented as SEQ ID NO:32, and the mature amino acid sequence as SEQ ID NO:34.

Comparison of the predicted protein sequences (FIG. 3) of one of the human interferon- $\tau$  genes (SEQ ID NO:4) and the ovine interferon- $\tau$  gene demonstrates the levels of sequence homology and divergence at the amino acid level.

An alignment of the nucleic acid sequences of the seven human interferon- $\tau$  nucleic acid sequences, described herein (Examples 6 and 7), with ovine interferon- $\tau$  is shown in FIGS. 19A and 19B. Sequences of OvIFN $\tau$  (oIFN $\tau$ ), HuIFN $\tau$ 1, HuIFN $\tau$ 2, and HuIFN $\tau$ 3 start at the upper left corner of FIG. 19A with the initiation ATG codon and continue through the second page of the figure. Sequences of HuIFN $\tau$ 4, HuIFN $\tau$ 5, HuIFN $\tau$ 6 and HuIFN $\tau$ 7 start approximately half-way down FIG. 19A with the CAG codon at amino acid position 40 (to the right of the exclamation marks) and continue through the second page of the figure. The 5' and 3' ends of each of the clones for HuIFN $\tau$ 4, HuIFN $\tau$ 5, HuIFN $\tau$ 6 and HuIFN $\tau$ 7 are represented by exclamation marks.

The complete coding sequence of OvIFN $\tau$  is presented in the top row of each aligned set. Nucleotides in the other sequences are indicated only at positions where they differ from those of OvIFN $\tau$ . Lower case letters represent nucleotide changes that do not result in amino acid changes, while upper case letters represent those changes that result in an amino acid substitution.

An alignment of the seven corresponding amino acid sequences, constructed in essentially the same manner as described above, is presented in FIGS. 20A and 20B. As above, the complete amino acid sequence of OvIFN $\tau$  is presented in the top row, and amino acids of other sequences are indicated only at positions where they differ from the ovine sequence.

An examination of the alignments reveals that the seven sequences may be grouped into at least three groups. Group I contains HuIFN $\tau$ 1 and HuIFN $\tau$ 2, group II contains HuIFN $\tau$ 3, HuIFN $\tau$ 4 and HuIFN $\tau$ 5, and group III contains HuIFN $\tau$ 6 and HuIFN $\tau$ 7. These groups may represent families of interferon- $\tau$  genes having distinct cellular functions.

These groupings were established based on the following criteria. In mature proteins, Group I HuIFN $\tau$ s have an asparagine (ASN) at amino acid position number 95 (numbers in reference to FIGS. 20A to 20B), a methionine (MET) at amino acid position number 104, and a leucine (LEU) at amino acid position number 120; Group II HuIFN $\tau$ s have an aspartic acid (ASP) at amino acid position number 95, a threonine (THR) at amino acid position number 104, and a methionine (MET) at amino acid position number 120; and Group III HuIFN $\tau$ s have an arginine (ARG) at amino acid position number 72, a valine (VAL) at amino acid position number 120, and a serine (SER) at amino acid position number 122.

The nucleic acid and polypeptide human IFN $\tau$  sequences presented as SEQ ID NO:3, SEQ ID NO:4, SEQ ID NO:11, SEQ ID NO:12, SEQ ID NO:21, SEQ ID NO:22, SEQ ID NO:23, SEQ ID NO:24, SEQ ID NO:25, SEQ ID NO:26, SEQ ID NO:27, SEQ ID NO:28, SEQ ID NO:29, SEQ ID NO:30, SEQ ID NO:31, SEQ ID NO:32, SEQ ID NO:33, and SEQ ID NO:34 can be used as the source for specific primers and probes to detect isolates of further human IFN $\tau$  coding sequences and/or pseudogenes. Further, as described above, there may be more than one isoform of the IFN $\tau$  protein and more than one coding sequence per species. The specific nucleic acid probes used in the practice of the present invention and antibodies reactive with the IFN $\tau$  polypeptides of the present invention may be useful to isolate unidentified variants of interferon- $\tau$  in mammals, according to the methods of the invention disclosed herein.

## 2. Characterization of the Expression of Interferon- $\tau$ in Human Tissues.

Human placental cDNA libraries and an ovine cDNA library were analyzed by hybridization to the OvIFN $\tau$  cDNA probe (Example 8). This DNA hybridization analysis suggested that the IFN $\tau$ -signals from human cDNA libraries were approximately  $\frac{1}{100}$  of the signal obtained using the ovine cDNA library. OvIFN $\tau$  cDNAs constitute around 0.4% of the ovine cDNA library. Accordingly, the abundance of human cDNAs responding to the OvIFN $\tau$  probe appears to be low, at least in the term placenta from which the cDNA libraries were generated.

The presence of HuIFN $\tau$  mRNA in human term placenta and amniocytes were also analyzed. The results suggest the presence of HuIFN $\tau$  mRNA in the feto-placental annex. The amniocytes also expressed the messages corresponding to OvIFN $\tau$  primers and human probe, suggesting that the expression of IFN $\tau$  mRNA is not limited to the term placenta.

In addition, a RT-PCR analysis for the presence of HuIFN $\tau$  was applied to the total cellular RNA isolated from human adult lymphocytes: the results suggest that IFN $\tau$  mRNA exists in lymphocytes.

The expression of interferon- $\tau$  in human tissue was also examined using in situ hybridization (Example 9). Sections



from four healthy, different term and first trimester human placentas were examined. This analysis employed a cDNA probe derived from the OvIFN $\tau$  cDNA sequences (Example 9B). In situ hybridization was performed using an anti-sense RNA probe. In three separate experiments, specific hybridization was observed in all term and first trimester placental tissues.

First trimester placental villi (composed of an outer layer of syncytiotrophoblast, an underlying layer of cytotrophoblast, and a central stromal region with various types of mesenchymal cells) displayed the highest transcript level of IFN $\tau$  in the cytotrophoblast cells. Less intense but detectable levels were present in both the syncytiotrophoblast and stromal cells. A similar pattern of transcript expression was demonstrated in the placental villi of term tissue but the level of signal detection was low. First trimester extravillous trophoblasts displayed the highest amount of message and stained positive when present in the maternal blood spaces within the decidua.

Howatson, et al., (1988) noted IFN $\alpha$  production in the syncytiotrophoblast of chorionic villi in both first trimester and term tissues. Also, Paulesu, et al. (1991) observed IFN $\alpha$  in extravillous trophoblast as well as syncytiotrophoblasts, noting more intense and abundant reactivity in first trimester placental tissue when compared to those taken at term. These investigators employed antibodies raised against human IFN $\alpha$  subtypes, and none observed any IFN $\alpha$  in the villous cytotrophoblasts.

The present results demonstrate that the human IFN $\tau$  gene is highly expressed in early placental tissues by migrating extravillous trophoblasts, but is also expressed in villous syncytiotrophoblasts, cytotrophoblasts, and various stromal cells. These results demonstrate the detection of IFN $\tau$  transcripts in human pregnancy tissues, and IFN $\tau$  expression in the villous cytotrophoblasts as well as the extravillous trophoblast of first trimester placenta.

#### C. Antiviral Properties of Interferon- $\tau$ .

The antiviral activity of OvIFN $\tau$  has been evaluated against a number of viruses, including both RNA and DNA viruses. The relative specific activity of OvIFN $\tau$ , purified to homogeneity, was evaluated in antiviral assays (Example 10). OvIFN $\tau$  had a higher specific antiviral activity than either rBoIFN $\alpha$  or rBoIFN $\gamma$  (Example 10, Table 3).

One advantage of the present invention is that OvIFN $\tau$  has potent antiviral activity with limited cytotoxic effects. Highly purified OvIFN $\tau$  was tested for anti-retroviral and cytotoxic effects on peripheral blood lymphocytes exposed to feline AIDS and human AIDS retroviruses (Bazer, F. W., et al., 1989). The feline AIDS lentivirus produces a chronic AIDS-like syndrome in cats and is a model for human AIDS (Pederson, et al., 1987). Replication of either virus in peripheral blood lymphocytes (PBL) was monitored by reverse transcriptase (RT) activity in culture supernatants over time.

To determine IFN $\tau$  antiviral activity against FIV and HIV, RNA-dependent DNA polymerase RT activity was assayed in FIV- and HIV-infected feline and human PBL cultures treated with OvIFN $\tau$  (Example 11). Replication of FIV was reduced to about one-third of control values when cells were cultured in the presence of OvIFN $\tau$ . Addition of OvIFN $\tau$  produced a rapid, dose-dependent decrease in reverse transcriptase (RT) activity (Example 11, Table 4). While concentrations as low as 0.62 ng/ml of IFN $\tau$  inhibited viral replication, much higher concentrations (40 ng/ml) having greater effects on RT-activity were without toxic effects on the cells. The results suggest that replication of the feline

immunodeficiency virus was reduced significantly compared to control values when cells were cultured in the presence of OvIFN $\tau$ .

IFN $\tau$  appeared to exert no cytotoxic effect on the cells hosting the retrovirus. This was true even when IFN $\tau$  was present at 40 ng per ml of culture medium. This concentration of IFN $\tau$  is equivalent to about 8,000 antiviral units of alpha interferon—when OvIFN $\tau$  and HuIFN $\alpha$  are each assayed for their ability to protect Madin-Darby bovine kidney cells from lysis by vesicular stomatitis virus (lysis assay as described by Pontzer, et al., 1988).

IFN $\tau$  was also tested for activity against HIV replication in human cells. Human peripheral lymphocytes, which had been infected with HIV were treated with varying concentrations of IFN $\tau$  (Example 12). Replication of HIV in peripheral blood lymphocytes was monitored by reverse transcriptase activity in culture supernatants over time. Over a range of concentrations of IFN $\tau$  produced significant anti-HIV effects (Example 12, Table 5). A concentration of only 10 ng/ml resulted in over a 50% reduction in RT activity after only six days. A concentration of 500 ng/ml resulted in a 90% reduction in RT activity within 10 days. Further, there was no evidence of any cytotoxic effects attributable to the administration of IFN $\tau$  (Example 12, Table 5).

Further, the antiviral effects of IFN $\tau$  against HIV were evaluated by treating human PBMC cells with various amounts of either recombinant IFN $\tau$  or recombinant human IFN $\alpha$  at the time of infection with HIV (Example 18). The data from these experiments (Example 18, Table 11) support the conclusion that, at similar concentrations, IFN $\alpha$  and IFN $\tau$  are effective in reducing the replication of HIV in human lymphocytes. However, treatment of cells with IFN $\alpha$  resulted in cytotoxicity, whereas no such cytotoxicity was observed with treatment using IFN $\tau$ , even when IFN $\tau$  was used at much higher concentrations. No cytotoxicity was observed using IFN $\tau$ , even when IFN $\tau$  was used at 200 times the dosage of interferon-alpha II.

Both FIV and HIV reverse transcriptase themselves were unaffected by IFN $\tau$  in the absence of PBL. Therefore, the antiviral activity is not due to a direct effect on the viral RT.

Interferon- $\tau$  has also been shown to inhibit Hepatitis B Virus DNA replication in hepatocytes (Example 18). A human cell line derived from liver cells transfected with Hepatitis B Virus (HBV) was used to test the antiviral effects of IFN $\tau$ . The cells were treated with both the IFN $\alpha$  and IFN $\tau$  over a range of concentrations. Both IFN $\alpha$  and IFN $\tau$  reduced DNA production by approximately two-fold compared to the no interferon control.

To demonstrate that the effect of the interferons was specific to the infecting virus and not the result of an effect on general cell metabolism, the hepatocytes were examined for the effects of IFN $\alpha$  and IFN $\tau$  on hepatospecific mRNA production (Example 18). Two hepatocyte specific proteins, Apo E and Apo A1, were detected by hybridization analysis. There was no apparent reduction of mRNA production for either hepatospecific mRNA at concentrations up to 40,000 units/ml of either IFN $\alpha$  or IFN $\tau$ . Further, no evidence for hepatotoxicity with IFN $\tau$  was seen in this assay.

The effects of recombinant ovine interferon tau (roIFN $\tau$ ) on ovine lentivirus replication (OvLV) were also evaluated. In vitro effects were assayed by infecting goat synovial membrane cells with serial dilutions of OvLV. The infected cells were treated daily with roIFN $\tau$  (0–2,500 antiviral units/ml [AVU/ml]) for 6 to 12 days, and virus replication and cytopathic effect (CPE; e.g., as in Example 2) were evaluated.



Evaluation methods included viral growth curves, cell proliferation assay (e.g., as in Examples 13, 14 or 15), syncytia formation assay (e.g., as in Nagy, et al., Dalglish, et al.), and quantitation of proviral DNA by PCR and reverse transcriptase assay (Mullis, Mullis, et al.). A reduction (p<0.001) of OvLV titer and CPE (80–99%) was observed in the roIFN $\tau$ -treated cultures.

In vivo effects of roIFN $\tau$  were assayed by inoculating twenty-four newborn lambs with  $5 \times 10^6$  TCID<sub>50</sub> of OvLV strain 85/34. Eleven of these lambs were treated with  $10^5$ – $10^6$  AVU/Kg of roIFN $\tau$  once a day for 30 days post-inoculation (PI) and twice a week thereafter. Thirteen lambs were used as controls. Virus titers in blood, as determined by an end point dilution method, peaked at 4–6 weeks PI in both groups. OvLV titers in roIFN $\tau$ -treated lambs were reduced relative to control animals. The largest reduction, a 90% decrease in OvLV titer in treated animals relative to control animals (p<0.01), was obtained 4 weeks PI.

The OvLV studies described above indicate that recombinant ovIFN $\tau$  can significantly reduce OvLV replication, and suggest that IFN $\tau$  may be used to control clinical diseases caused by lentivirus infections. Taken together with the other antiviral data, these results suggest that IFN $\tau$  is an effective antiviral agent against a wide variety of viruses, including both RNA and DNA viruses.

Ovine interferon- $\tau$  may be useful in veterinary applications including, but not limited to, the treatment of the following viral diseases: feline leukemia virus, ovine progressive pneumonia virus, ovine lentivirus, equine infectious anemia virus, bovine immunodeficiency virus, visna-maedi virus, and caprine arthritis encephalitis.

Human interferon- $\tau$  may be used for the treatment of, for example, the following viral diseases: human immunodeficiency virus (HIV), hepatitis c virus (HCV) and hepatitis B virus (HBV).

#### D. Antiproliferative Properties of IFN $\tau$ .

The effects of IFN $\tau$  on cellular growth have also been examined. In one analysis, anti-cellular growth activity was examined using a colony inhibition assay (Example 13). Human amnion (WISH) or MDBK cells were plated at low cell densities to form colonies originating from single cells. Dilutions of interferons were added to triplicate wells and the plates were incubated to allow colony formation. IFN $\tau$  inhibited both colony size and number in these assays. IFN $\tau$  was more effective at inhibiting cell proliferation of the human cell line (WISH) than human IFN $\alpha$ . The antiproliferative activity of IFN $\tau$  was dose-dependent. High concentrations of IFN $\tau$  stopped proliferation, while cell viability was not impaired.

Based on cell cycle analysis, using flow cytometry, IFN $\tau$  appears to inhibit progress of cells through S phase. These results demonstrate the antiproliferative effect of IFN $\tau$ , and underscore its low cytotoxicity.

The antiproliferative effects of IFN $\tau$  were also studied for rat and bovine cell lines (Example 14). The rate of <sup>3</sup>H-thymidine incorporation was used to assess the rate of cellular proliferation. The data obtained demonstrate that IFN $\tau$  drastically reduced the rate of cellular proliferation (Example 14, Table 7) for each tested cell line.

The antiproliferative activity and lack of toxicity of IFN $\tau$  was further examined using a series of human tumor cell lines (Example 15). A variety of human tumor cell lines were selected from the standard lines used in NIH screening procedure for antineoplastic agents (Pontzer, C. H., et al., (1991)). At least one cell line from each major neoplastic category was examined.

The following cell lines were obtained from American Type Culture Collection (12301 Parklawn Dr., Rockville Md. 20852):

NCI-H460 human lung large cell carcinoma;  
DLD-1 human colon adenocarcinoma;  
SK-MEL-28 human malignant melanoma;  
ACHN human renal adenocarcinoma;  
HL-60 human promyelocytic leukemia;  
H9 human T cell lymphoma;  
HUT 78 human cutaneous T cell lymphoma; and  
MCF7 human breast adenocarcinoma.

As above, the antiproliferative activity was evaluated by measuring the rate of <sup>3</sup>H-thymidine incorporation into cells which have been treated with IFN $\tau$ . Significant differences between treatments were assessed by an analysis of variance followed by Scheffe's F-test. Cell cycle analysis was performed by flow cytometry.

Examination of IFN $\tau$  inhibition of MCF7 (breast adenocarcinoma) proliferation demonstrated that IFN $\tau$  reduced MCF7 proliferation in a dose-dependent manner. A 50% reduction in <sup>3</sup>H-thymidine was observed with 10,000 units/ml of IFN $\tau$  (Example 15, Table 8). This cell line had previously been found to be unresponsive to antiestrogen treatment.

A comparison of the antiproliferative effects of IFN $\tau$  and IFN $\alpha$  was conducted using HL-60 (human promyelocytic leukemia) cells. Results with the promyelocytic leukemia HL-60 are typical of those obtained comparing IFN $\tau$  with human IFN $\alpha$  (Example 15). Concentrations as low as 100 units/ml of both IFNs produced significant (>60%) growth reduction. Increasing amounts of IFNs further decreased tumor cell proliferation (FIG. 4). High doses of HuIFN $\alpha$ , but not OvIFN $\tau$ , were cytotoxic (FIG. 5). Cell viability was reduced by approximately 80% by IFN $\alpha$ . By contrast, nearly 100% of the IFN $\tau$ -treated cells remained viable when IFN $\tau$  was applied at 10,000 units/ml. Thus, while both interferons inhibit proliferation, only IFN $\tau$  is without cytotoxicity. This lack of toxicity provides an advantage of IFN $\tau$  for use in vivo therapies.

The human cutaneous T cell lymphoma, HUT 78, responded similarly to HL-60 when treated with IFN $\tau$  (Example 15, FIG. 9). Both OvIFN $\tau$  and rHuIFN $\alpha$  reduce HUT 78 cell growth, but IFN $\alpha$  demonstrated adverse effects on cell viability.

The T cell lymphoma H9 was less sensitive to the antiproliferative effects of IFN $\alpha$  than the tumor cell lines described above. While IFN $\alpha$  was not toxic to the H9 cells, it failed to inhibit cell division significantly at any of the concentrations examined (Example 15, FIG. 10). In contrast, IFN $\tau$  was observed to reduce H9 growth by approximately 60%. Thus, only OvIFN $\tau$  is an effective growth inhibitor of this T cell lymphoma.

In three additional tumor cell lines (NCI-H460, DLD-1 and SK-MEL-28) IFN $\tau$  and IFN $\alpha$  were equally efficacious antitumor agents. In the melanoma, SK-MEL-28, inhibition of proliferation by IFN $\alpha$  was accomplished by a 13% drop in viability, while IFN $\tau$  was not cytotoxic. In the majority of tumors examined, IFN $\tau$  is equal or preferable to IFN $\alpha$  as an antineoplastic agent against human tumors.

IFN $\tau$  exhibits antiproliferative activity against human tumor cells without toxicity and is as potent or more potent than human IFN $\alpha$ . Clinical trials of the IFN $\alpha$ 2s have shown them to be effective antitumor agents (Dianzani, F., 1992; Krown, 1987). One advantage of IFN $\tau$  as a therapeutic is the elimination of toxic effects seen with high doses IFN $\alpha$ s.

An additional application of the IFN $\tau$  is against tumors like Kaposi's sarcoma (associated with HIV infection)



where the antineoplastic effects of IFN $\tau$  are coupled with IFN $\tau$  ability to inhibit retroviral growth.

The in vivo efficacy of interferon- $\tau$  treatment was examined in a mouse system (Example 16). B16-F10 is a syngeneic mouse transplantable tumor selected because of its high incidence of pulmonary metastases (Poste, et al., 1981). Interferon treatment was initiated 3 days after the introduction of the tumor cells. The in vivo administration of IFN $\tau$  dramatically reduced B16-F10 pulmonary tumors. Thus, IFN $\tau$  appears to be an efficacious antineoplastic agent in vivo as well as in vitro.

These results support the usefulness of human IFN $\tau$  for use in methods to inhibit or reduced tumor cell growth, including, but are not limited to, the following types of tumor cells: human carcinoma cells, hematopoietic cancer cells, human leukemia cells, human lymphoma cells, human melanoma cells and steroid-sensitive tumor cells (for example, mammary tumor cells).

#### E. Cytotoxicity of Interferons.

One advantage of IFN $\tau$  over other interferons, such as IFN $\alpha$ , is that treatment of a subject with therapeutic doses of IFN $\tau$  does not appear to be associated with cytotoxicity. In particular, IFN- $\tau$  appears to be non-toxic at concentrations at which IFN- $\beta$  induces toxicity. This is demonstrated by experiments in which L929 cells were treated with various concentrations of either oIFN $\tau$  or MuIFN- $\beta$  (Lee Biomolecular, San Diego, Calif.), ranging from 6000 U/ml to 200,000 U/ml (Example 18E).

IFN $\tau$ , MuIFN- $\beta$  or medium (control) were added at time zero and the cells were incubated for 72 hours. The results of the experiments are presented in FIG. 21. The percent of live cells (relative to control) is indicated along the y-axis ( $\pm$ standard error). One hundred percent is equal to the viability of L929 cells treated with medium alone. The results indicate that oIFN $\tau$  is essentially non-toxic at concentrations up to 100,000 U/ml, and is significantly less toxic than MuIFN- $\beta$  over the entire therapeutic range of the compounds.

It has been previously demonstrated that in vivo treatment with both of the type I IFNs, IFN $\beta$  and IFN $\alpha$  in humans and animals causes toxicity manifested as a number of side effects including fever, lethargy, tachycardia, weight loss, and leukopenia (Degre, 1974; Fent and Zbinden, 1987). The effect of in vivo treatment with IFN $\tau$ , IFN $\beta$  and IFN $\alpha$  ( $10^5$  U/injection) on total white blood cell (WBC), total lymphocyte counts and weight measurements in NZW mice (Table 13) was examined as described in Example 18F. No significant difference between IFN $\tau$  treated and untreated mice was observed for WBC, lymphocyte counts or weight change.

In comparison, IFN $\beta$  treated mice exhibited a 31.7% depression in lymphocyte counts 12 hours after injection. Further, depression of lymphocyte counts continued 24 hours after IFN $\beta$  injection. IFN $\alpha$  treated mice exhibited a 55.8% lymphocyte depression and significant weight loss 12 hours after injection. Thus, IFN $\tau$  appears to lack toxicity in vivo unlike IFN $\beta$  and IFN $\alpha$  as evidenced by studies of peripheral blood and weight measurements.

oIFN $\tau$  is 172 amino acids long compared to 165 or 166 amino acids for IFN $\alpha$ . The carboxyl-terminal portion of oIFN $\tau$  is hydrophilic and thought to be surface accessible. To explore whether this carboxyl "tail" interacts with the binding epitope of oIFN $\tau$  to mediate the relative lack of cytotoxicity, a series of deletion mutants were generated.

The carboxyl terminal 2, 6 and 10 amino acids of oIFN $\tau$  were removed by cassette mutagenesis of a synthetic oIFN $\tau$  gene. The mutant (variant) synthetic genes were cloned into

the pHIL-S1 Pichia expression plasmid (Invitrogen, San Diego, Calif.), the plasmid was cut with BglII, and the linearized plasmid was used to transform *Pichia pastoris* (strain GS115; Invitrogen) spheroplasts according to the manufacturer's instructions.

Recombinant variant proteins expressed by transformed His<sup>+</sup> Mut<sup>-</sup> yeast cells were purified and used to determine in vitro antiviral activity and relative cytotoxicity of the variants compared to intact oIFN $\tau$  and IFN- $\alpha$ . The cytotoxicity of the variants was distributed between the that of oIFN $\tau$  and IFN- $\alpha$ . Variants with shorter deletions were more similar in their cytotoxic properties to oIFN $\tau$ , while those with longer deletions were more similar to IFN- $\alpha$ .

While not wishing to be bound by any specific molecular mechanisms underlying the properties of IFN $\tau$ , the results of the experiments suggest that the C-terminus 10 amino acids of IFN $\tau$  may play a role in the decreased cytotoxicity of IFN $\tau$  relative to other interferons.

#### III. Interferon- $\tau$ Polypeptide Fragments, Protein Modeling and Protein Modifications.

##### A. IFN $\tau$ Polypeptide Fragments.

The variety of IFN $\tau$  activities, its potency and lack of cytotoxicity, as taught by the present specification, suggest the importance of structure/function analysis for this novel interferon. The structural basis for OvIFN $\tau$  function has been examined using six overlapping synthetic peptides corresponding to the entire OvIFN $\tau$  sequence (FIG. 6). The corresponding polypeptides derived from the ovine IFN $\tau$  sequence are presented as SEQ ID NO:5 to SEQ ID NO:10. Three peptides representing amino acids 1-37, 62-92 and 139-172 have been shown to inhibit IFN $\tau$  antiviral activity (Example 17). The peptides were effective competitors at concentrations of 300  $\mu$ M and above.

The synthetic polypeptide representing the C-terminal region of ovIFN $\tau$ , OvIFN $\tau$  (139-172), and the internal peptide OvIFN $\tau$  (62-92), inhibited IFN $\tau$  and rBoIFN $\alpha$  antiviral activity to the same extent, while the N-terminal peptide OvIFN $\tau$  (1-37) was more effective in inhibiting OvIFN $\tau$  antiviral activity. Dose-response data indicated that IFN $\tau$  (62-92) and IFN $\tau$  (139-172) inhibited IFN $\tau$  antiviral activity to similar extents. The same peptides that blocked IFN $\tau$  antiviral activity also blocked the antiviral activity of recombinant bovine IFN $\alpha$  (rBoIFN $\alpha$ ); recombinant bovine IFN $\gamma$  was unaffected by the peptides. These two IFN $\tau$  peptides may represent common receptor binding regions for IFN $\tau$  and various IFN $\alpha$ s.

The two synthetic peptides OvIFN $\tau$  (1-37) and OvIFN $\tau$  (139-172) also blocked OvIFN $\tau$  anti-FIV and anti-HIV activity (Example 17; FIGS. 11A and 11B). While both peptides blocked FIV RT activity, only the C-terminal peptide, OvIFN $\tau$  (139-172), appeared to be an efficient inhibitor of vesicular stomatitis virus activity on the feline cell line, Fc9.

The above data taken together suggest that the C-terminal regions of type I interferons may bind to common site on the type I interferon receptor, while the N-terminal region may be involved in the elicitation of unique functions. These results suggest that portions of the IFN $\tau$  molecule may be used to substitute regions of interferon alpha molecules. For example, the region of an interferon alpha molecule that is responsible for increased cytotoxicity, relative to IFN $\tau$  treatment, can be identified by substituting polypeptide regions derived from IFN $\tau$  for regions of an interferon alpha molecule. Such substitutions can be carried out by manipulation of synthetic genes (see below) encoding the selected IFN $\tau$  and interferon alpha molecules, coupled to the functional assays described herein (such as, antiviral, antiproliferative and cytotoxicity assays).



Polyclonal anti-peptide antisera against the IFN $\tau$  peptides yielded similar results as the polypeptide inhibition studies, described above. Antibodies directed against the same three regions (OvIFN $\tau$  (1–37), IFN $\tau$  (62–92) and IFN $\tau$  (139–172)) blocked OvIFN $\tau$  function, confirming the importance of these three domains in antiviral activity (Example 17). These peptides, although apparently binding to the interferon receptor, did not in and of themselves elicit interferon-like effects in the cells.

The antiproliferative activity of IFN $\tau$  (Example 17, Table 11) involved a further region of the molecule, since IFN $\tau$  (119–150) was the most effective inhibitor of OvIFN $\tau$ -induced reduction of cell proliferation. This results suggests that the region of the molecule primarily responsible for inhibition of cell growth is the IFN $\tau$  (119–150) region. This region of the IFN $\tau$  molecule may be useful alone or fused to other proteins (such as serum albumin, an antibody or an interferon alpha polypeptide) as an antineoplastic agent. A conjugated protein between an N-terminal peptide derived from human interferon- $\alpha$  and serum albumin was shown to have anticellular proliferation activity (Ruegg, et al., 1990).

Finally, binding of  $^{125}\text{I}$ -OvIFN $\tau$  to its receptor on MDBK cells could be blocked by antisera to 4 of the 6 peptides; the 4 polypeptides representing amino acids 1–37, 62–92, 119–150 and 139–172 of OvIFN $\tau$ . This reflects the multiple binding domains as well as the functional significance of these regions. Since different regions of IFN $\tau$  are involved in elicitation of different functions, modification of selected amino acids could potentially result in IFN $\tau$ -like interferons with selective biological activity.

Polypeptide fragments of human IFN $\tau$  proteins, having similar properties to the OvIFN $\tau$  polypeptides just described, are proposed based on the data presented above for OvIFN $\tau$  polypeptide fragments combined with the HuIFN $\tau$  sequence information disclosed herein. Such human-sequence derived polypeptides include, but are not limited to, the following: SEQ ID NO:15 to SEQ ID NO:20, and SEQ ID NO:35 to SEQ ID NO:40.

The above data demonstrate the identification of synthetic peptides having four discontinuous sites on the OvIFN $\tau$  protein that are involved in receptor interaction and biological activity. In order to elucidate the structural relationship of these regions, modeling of the three dimensional structure of IFN $\tau$  was undertaken. A three dimensional model would be useful in interpretation of existing data and the design of future structure/function studies.

#### B. Molecular Modeling

Combining circular dichroism (CD) data of both the full length recombinant OvIFN $\tau$  and IFN $\beta$  (a protein of known three dimensional structure (Senda, et al., 1992)), a model of OvIFN $\tau$  was constructed. The most striking feature of this model is that IFN $\tau$  falls into a class of proteins with a four-helix bundle motif. The CD spectra of IFN $\tau$  was taken on an AVIV 60 S spectropolarimeter. Two different methods were employed for secondary structure estimations, the algorithm of Perczel, et al., (1991) and variable selection by W. C. Johnson, Jr. (1992).

Secondary structure estimations of the spectra indicate 70–75% alpha helix (characterized by minima at 222 and 208 nm and maximum at 190 nm). The variable selection algorithm estimates the remainder of the molecule to be 20% beta sheet and 10% turn. The Chang method estimates the remainder to be 30% random coil. Alignment of IFN $\tau$  and IFN $\beta$  sequences revealed homology between the two molecules, specifically in the regions of known helical structure in IFN $\beta$ . Sequence analysis of IFN $\tau$  also showed that proposed helical regions possess an apolar periodicity indicative of a four-helix bundle motif.

The final modeling step was to apply the IFN $\beta$  x-ray crystallographic coordinates of the IFN $\beta$  carbon backbone to the IFN $\tau$  sequence. The functionally active domains of IFN $\tau$ , identified above, were localized to one side of the molecule and found to be in close spatial proximity. This is consistent with multiple binding sites on IFN $\tau$  interacting simultaneously with the type I IFN receptor.

The three dimensional modeling data coupled with the function data described above, provides the information necessary to introduce sequence variations into specific regions of IFN $\tau$  to enhance selected functions (e.g., antiviral or anticellular proliferation) or to substitute a region(s) of selected function into other interferon molecules (e.g., antiviral, antineoplastic, or reduced cytotoxicity).

#### C. Recombinant and Synthetic Manipulations

The construction of a synthetic gene for OvIFN $\tau$  is described in Example 3. Briefly, an amino acid sequence of ovIFN $\tau$  was back-translated from an ovIFN $\tau$  cDNA (Imakawa, et al., 1987) using optimal codon usage for *E. coli*. The sequence was edited to include 20, unique, restriction sites spaced throughout the length of the construct. This 540 base pair synthetic gene sequence was divided into 11 oligonucleotide fragments. Individual fragments were synthesized and cloned, either single or double stranded, into either pTZ 19R, pTZ 18R or pBluescript, amplified and fused. The synthetic OvIFN $\tau$  construct was then cloned into a modified pIN-III-ompA expression vector for expression in bacteria and also cloned into a yeast expression plasmid. A similarly constructed human IFN $\tau$  synthetic gene (SEQ ID NO:3) has been designed, constructed and expressed in yeast cells.

Expression of the OvIFN $\tau$  synthetic gene in yeast (Example 4) allowed over production of recombinant IFN $\tau$  in *S. cerevisiae*: large quantities (5–20 mg/l) of recombinant IFN $\tau$  can be purified from soluble yeast extract using sequential ion exchange and molecular sieve chromatography. Recombinant IFN $\tau$  purified in this fashion exhibited potent antiviral activity (2 to  $3 \times 10^8$  units/mg) similar to native OvIFN $\tau$ .

The synthetic gene construct facilitates introduction of mutations for possible enhancement of antitumor (anticellular proliferative) and antiviral activities. Further, the disparate regions of the molecule responsible for different functions can be modified independently to generate a molecule with a desired function. For example, two deletion mutants, OvIFN $\tau$ (1–162) and OvIFN $\tau$  (1–166), have been constructed to examine the role of carboxy terminal sequences in IFN $\tau$  molecules.

Additional mutant IFN $\tau$  molecules have been constructed to identify residues critical for antiproliferative activity. For example, one particular residue, TYR 123 has been implicated in the anticellular proliferative activity of IFN $\alpha$  (McInnes, et al., 1989). The equivalent of TYR 123 in IFN $\tau$  is contained within peptide OvIFN $\tau$  (119–150): this polypeptide inhibits OvIFN $\tau$  and human IFN $\alpha$  antiproliferative activity. Mutations converting TYR 123 to conservative (TRP) and nonconservative (ASP) substitutions have been generated, as well as mutant sequences having deletion of this residue. The codon for TYR 123 is located within an *sspI* site; elimination of this site has been used for screening. The antiproliferative activity of these mutant IFN $\tau$  is evaluated as described herein.

Synthetic peptides can be generated which correspond to the IFN $\tau$  polypeptides of the present invention. Synthetic peptides can be commercially synthesized or prepared using standard methods and apparatus in the art (Applied Biosystems, Foster City Calif.).



Alternatively, oligonucleotide sequences encoding peptides can be either synthesized directly by standard methods of oligonucleotide synthesis, or, in the case of large coding sequences, synthesized by a series of cloning steps involving a tandem array of multiple oligonucleotide fragments corresponding to the coding sequence (Crea; Yoshio et al.; Eaton et al.). Oligonucleotide coding sequences can be expressed by standard recombinant procedures (Maniatis et al.; Ausubel et al.).

The biological activities of the interferon- $\tau$  polypeptides described above can be exploited using either the interferon- $\tau$  polypeptides alone or conjugated with other proteins (see below).

#### IV. Production of Fusion Proteins.

In another aspect, the present invention includes interferon- $\tau$  or interferon- $\tau$ -derived polypeptides covalently attached to a second polypeptide to form a fused, or hybrid, protein. The interferon- $\tau$  sequences making up such fused proteins can be recombinantly produced interferon- $\tau$  or a bioactive portion thereof, as described above.

For example, where interferon- $\tau$  is used to inhibit viral expression, polypeptides derived from IFN $\tau$  demonstrating antiviral activity may be advantageously fused with a soluble peptide, such as, serum albumin, an antibody (e.g., specific against a virus-specific cell surface antigen), or an interferon alpha polypeptide. In one embodiment, the IFN $\tau$  polypeptides provide a method of reducing the toxicity of other interferon molecules (e.g., IFN $\beta$  or IFN $\alpha$ ) by replacing toxicity-associated regions of such interferons with, for example, corresponding interferon- $\tau$  regions having lower toxicity. In another embodiment, fusion proteins are generated containing interferon- $\tau$  regions that have anticellular proliferation properties. Such regions may be obtained from, for example, the human interferon- $\tau$  sequences disclosed herein.

The fused proteins of the present invention may be formed by chemical conjugation or by recombinant techniques. In the former method, the interferon- $\tau$  and second selected polypeptide are modified by conventional coupling agents for covalent attachment. In one exemplary method for coupling soluble serum albumin to an interferon- $\tau$  polypeptide, serum albumin is derivatized with N-succinimidyl-S-acetyl thioacetate (Duncan), yielding thiolated serum albumin. The activated serum albumin polypeptide is then reacted with interferon- $\tau$  derivatized with N-succinimidyl 3-(2-pyridyldithio) propionate (Cumber), to produce the fused protein joined through a disulfide linkage.

As an alternative method, recombinant interferon- $\tau$  may be prepared with a cysteine residue to allow disulfide coupling of the interferon- $\tau$  to an activated ligand, thus simplifying the coupling reaction. An interferon- $\tau$  expression vector, used for production of recombinant interferon- $\tau$ , can be modified for insertion of an internal or a terminal cysteine codon according to standard methods of site-directed mutagenesis (Ausubel, et al.).

In one method, a fused protein is prepared recombinantly using an expression vector in which the coding sequence of a second selected polypeptide is joined to the interferon- $\tau$  coding sequence. For example, human serum albumin coding sequences can be fused in-frame to the coding sequence of an interferon- $\tau$  polypeptide, such as, SEQ ID NO:9, SEQ ID NO:19 or SEQ ID NO:39. The fused protein is then expressed using a suitable host cell. The fusion protein may be purified by molecular-sieve and ion-exchange chromatography methods, with additional purification by polyacrylamide gel electrophoretic separation and/or HPLC chromatography, if necessary.

It will be appreciated from the above how interferon- $\tau$ -containing fusion proteins may be prepared. One variation on the above fusion is to exchange positions of the interferon- $\tau$  and selected second protein molecules in the fusion protein (e.g., carboxy terminal versus amino terminal fusions). Further, internal portions of a native interferon- $\tau$  polypeptide (for example, amino acid regions of between 15 and 172 amino acids) can be assembled into polypeptides where two or more such interferon- $\tau$  portions are contiguous that are normally discontinuous in the native protein.

In addition to the above-described fusion proteins, the present invention also contemplates polypeptide compositions having (a) a human interferon- $\tau$  polypeptide, where said polypeptide is (i) derived from an interferon- $\tau$  amino acid coding sequence, and (ii) between 15 and 172 amino acids long, and (b) a second soluble polypeptide. Interferon- $\alpha$  and interferon- $\beta$  are examples of such second soluble polypeptides. IFN $\tau$  polypeptides associated with reduced toxicity may be co-administered with more toxic interferons to reduce the toxicity of the more toxic interferons when used in pharmaceutical formulations or in therapeutic applications. Such IFN $\tau$  polypeptides would, for example, reduce the toxicity of IFN $\alpha$  but not interfere with IFN $\alpha$  antiviral properties.

#### V. Antibodies Reactive with Interferon- $\tau$ .

Fusion proteins containing the polypeptide antigens of the present invention fused with the glutathione-S-transferase (Sj26) protein can be expressed using the pGEX-GLI vector system in *E. coli* JM101 cells. The fused Sj26 protein can be isolated readily by glutathione substrate affinity chromatography (Smith). Expression and partial purification of IFN $\tau$  proteins is described in (Example 20), and is applicable to any of the other soluble, induced polypeptides coded by sequences described by the present invention.

Insoluble GST (sj26) fusion proteins can be purified by preparative gel electrophoresis.

Alternatively, IFN $\tau$ - $\beta$ -galactosidase fusion proteins can be isolated as described in Example 19.

Also included in the invention is an expression vector, such as the lambda gt11 or pGEX vectors described above, containing IFN $\tau$  coding sequences and expression control elements which allow expression of the coding regions in a suitable host. The control elements generally include a promoter, translation initiation codon, and translation and transcription termination sequences, and an insertion site for introducing the insert into the vector.

The DNA encoding the desired polypeptide can be cloned into any number of vectors (discussed above) to generate expression of the polypeptide in the appropriate host system. These recombinant polypeptides can be expressed as fusion proteins or as native proteins. A number of features can be engineered into the expression vectors, such as leader sequences which promote the secretion of the expressed sequences into culture medium. Recombinantly produced IFN $\tau$ , and polypeptides derived therefrom, are typically isolated from lysed cells or culture media. Purification can be carried out by methods known in the art including salt fractionation, ion exchange chromatography, and affinity chromatography. Immunoaffinity chromatography can be employed using antibodies generated against selected IFN $\tau$  antigens.

In another aspect, the invention includes specific antibodies directed against the polypeptides of the present invention. Typically, to prepare antibodies, a host animal, such as a rabbit, is immunized with the purified antigen or fused protein antigen. Hybrid, or fused, proteins may be generated using a variety of coding sequences derived from other



proteins, such as  $\beta$ -galactosidase or glutathione-S-transferase. The host serum or plasma is collected following an appropriate time interval, and this serum is tested for antibodies specific against the antigen. Example 20 describes the production of rabbit serum antibodies which are specific against the IFN $\tau$  antigens in a Sj26/IFN $\tau$  hybrid protein. These techniques can be applied to the all of the IFN $\tau$  molecules and polypeptides derived therefrom.

The gamma globulin fraction or the IgG antibodies of immunized animals can be obtained, for example, by use of saturated ammonium sulfate or DEAE Sephadex, or other techniques known to those skilled in the art for producing polyclonal antibodies.

Alternatively, purified protein or fused protein may be used for producing monoclonal antibodies. Here the spleen or lymphocytes from a animal immunized with the selected polypeptide antigen are removed and immortalized or used to prepare hybridomas by methods known to those skilled in the art (Harlow, et al.). Lymphocytes can be isolated from a peripheral blood sample. Epstein-Barr virus (EBV) can be used to immortalize human lymphocytes or a fusion partner can be used to produce hybridomas.

Antibodies secreted by the immortalized cells are screened to determine the clones that secrete antibodies of the desired specificity, for example, by using the ELISA or Western blot method (Ausubel et al.). Experiments performed in support of the present invention have yielded four hybridomas producing monoclonal antibodies specific for ovine IFN $\tau$  have been isolated.

Antigenic regions of polypeptides are generally relatively small, typically 7 to 10 amino acids in length. Smaller fragments have been identified as antigenic regions. Interferon- $\tau$  polypeptide antigens are identified as described above. The resulting DNA coding regions can be expressed recombinantly either as fusion proteins or isolated polypeptides.

In addition, some amino acid sequences can be conveniently chemically synthesized (Applied Biosystems, Foster City Calif.). Antigens obtained by any of these methods may be directly used for the generation of antibodies or they may be coupled to appropriate carrier molecules. Many such carriers are known in the art and are commercially available (e.g., Pierce, Rockford Ill.).

Antibodies reactive with IFN $\tau$  are useful, for example, in the analysis of structure/function relationships.

## VI. Utility

### A. Reproductive.

Although IFN $\tau$  bears some similarity to the IFN $\alpha$  family based on structure and its potent antiviral properties, the IFN $\alpha$ s do not possess the reproductive properties associated with IFN $\tau$ . For example, recombinant human IFN $\alpha$  had no effect on interestrus interval compared to IFN $\tau$ , even when administered at twice the dose (Davis, et al., 1992).

Therefore, although IFN $\tau$  has some structural similarities to other interferons, it has very distinctive properties of its own: for example, the capability of significantly influencing the biochemical events of the estrous cycle.

The human IFN $\tau$  of the present invention can be used in methods of enhancing fertility and prolonging the life span of the corpus luteum in female mammals as generally described in Hansen, et al., herein incorporated by reference. Further, the human interferon- $\tau$  of the present invention could be used to regulate growth and development of uterine and/or fetal-placental tissues. The human IFN $\tau$  is particularly useful for treatment of humans, since potential antigenic responses are less likely using such a same-species protein.

### B. Antiviral Properties.

The antiviral activity of IFN $\tau$  has broad therapeutic applications without the toxic effects that are usually associated with IFN $\alpha$ s. Although the presence of IFN $\tau$  in culture medium inhibited reverse transcriptase activity of the feline immunodeficiency virus (Example 11), this is not due to a direct effect of IFN $\tau$  on the reverse transcriptase. Rather, IFN $\tau$  appears to induce the host cell to produce a factor(s) which is inhibitory to the reverse transcriptase of the virus.

IFN $\tau$  was found to exert its antiviral activity without adverse effects on the cells: no evidence of cytotoxic effects attributable to the administration of IFN $\tau$  was observed. It is the lack of cytotoxicity of IFN $\tau$  which makes it extremely valuable as an in vivo therapeutic agent. This lack of cytotoxicity sets IFN $\tau$  apart from most other known antiviral agents and all other known interferons.

Formulations comprising the IFN $\tau$  compounds of the present invention can be used to inhibit viral replication.

The human IFN $\tau$  of the present invention can be employed in methods for affecting the immune relationship between fetus and mother, for example, in preventing transmission of maternal viruses (e.g., HIV) to the developing fetus. The human interferon- $\tau$  is particularly useful for treatment of humans, since potential antigenic responses are less likely using a homologous protein.

### C. Anticellular Proliferation Properties.

IFN $\tau$  exhibits potent anticellular proliferation activity. IFN $\tau$  can also be used to inhibit cellular growth without the negative side effects associated with other interferons which are currently known. Formulations comprising the IFN $\tau$  compounds of the subject invention can be used to inhibit, prevent, or slow tumor growth.

The development of certain tumors is mediated by estrogen. Experiments performed in support of the present invention indicate that IFN $\tau$  can suppress estrogen receptor numbers. Therefore, IFN $\tau$  can be used in the treatment or prevention of estrogen-dependent tumors.

### D. Interfering with the Binding of Interferons to Receptors.

IFN $\tau$  appears to interact with the Type I IFN receptor via several epitopes on the molecule, and these regions either separately or in combination may affect distinct functions of IFN $\tau$  differently.

The polypeptides of the present invention are useful for the selective inhibition of binding of interferons to the interferon receptor. Specifically, as described herein, certain of the disclosed peptides selectively inhibit the antiviral activity of IFN $\tau$  while others inhibit the antiproliferative activity. Combinations of these peptides could be used to inhibit both activities. Advantageously, despite binding to the interferon receptor and blocking IFN $\tau$  activity, these peptides do not, themselves, elicit the antiviral or antiproliferative activity.

Therefore, such polypeptides can be used as immunoregulatory molecules when it is desired to prevent immune responses triggered by interferon molecules. These peptides could be used as immunosuppressants to prevent, for example, interferon-mediated immune responses to tissue transplants. Other types of interferon mediated responses may also be blocked, such as the cytotoxic effects of alpha interferon.

### E. Pharmaceutical Compositions.

IFN $\tau$  proteins can be formulated according to known methods for preparing pharmaceutically useful compositions. Formulations comprising interferons or interferon-like compounds have been previously described (for example, Martin, 1976). In general, the compositions of the subject



invention will be formulated such that an effective amount of the IFN $\tau$  is combined with a suitable carrier in order to facilitate effective administration of the composition.

The compositions used in these therapies may also be in a variety of forms. These include, for example, solid, semi-solid, and liquid dosage forms, such as tablets, pills, powders, liquid solutions or suspensions, liposomes, suppositories, injectable, and infusible solutions. The preferred form depends on the intended mode of administration and therapeutic application. The compositions also preferably include conventional pharmaceutically acceptable carriers and adjuvants which are known to those of skill in the art. Preferably, the compositions of the invention are in the form of a unit dose and will usually be administered to the patient one or more times a day.

IFN $\tau$ , or related polypeptides, may be administered to a patient in any pharmaceutically acceptable dosage form, including oral intake, inhalation, intranasal spray, intraperitoneal, intravenous, intramuscular, intralesional, or subcutaneous injection. Specifically, compositions and methods used for other interferon compounds can be used for the delivery of these compounds.

One primary advantage of the compounds of the subject invention, however, is the extremely low cytotoxicity of the IFN $\tau$  proteins. Because of this low cytotoxicity, it is possible to administer the IFN $\tau$  in concentrations which are greater than those which can generally be utilized for other interferon (e.g., IFN $\alpha$ ) compounds. Thus, IFN $\tau$  can be administered at rates from about  $5 \times 10^4$  to  $20 \times 10^6$  units/day to about  $500 \times 10^6$  units/day or more. In a preferred embodiment, the dosage is about  $20 \times 10^6$  units/day. High doses are preferred for systemic administration. It should, of course, be understood that the compositions and methods of this invention may be used in combination with other therapies.

Once improvement of a patient's condition has occurred, a maintenance dose is administered if necessary. Subsequently, the dosage or the frequency of administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained. When the symptoms have been alleviated to the desired level, treatment should cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

The compositions of the subject invention can be administered through standard procedures to treat a variety of cancers and viral diseases including those for which other interferons have previously shown activity. See, for example, Finter, et al. (1991); Dianzani, et al. (1992); Francis, et al. (1992) and U.S. Pat. Nos. 4,885,166 and 4,975,276. However, as discussed above, the compositions of the subject invention have unique features and advantages, including their ability to treat these conditions without toxicity.

#### F. Treatment of Skin Disorders.

Disorders of the skin can be treated intralesionally using IFN $\tau$ , wherein formulation and dose will depend on the method of administration and on the size and severity of the lesion to be treated. Preferred methods include intradermal and subcutaneous injection. Multiple injections into large lesions may be possible, and several lesions on the skin of a single patient may be treated at one time. The schedule for administration can be determined by a person skilled in the art. Formulations designed for sustained release can reduce the frequency of administration.

#### G. Systemic Treatment.

Systemic treatment is essentially equivalent for all applications. Multiple intravenous, subcutaneous and/or intramuscular doses are possible, and in the case of implantable methods for treatment, formulations designed for sustained release are particularly useful. Patients may also be treated using implantable subcutaneous portals, reservoirs, or pumps.

#### H. Regional Treatment.

Regional treatment with the IFN $\tau$  polypeptides of the present invention is useful for treatment of cancers in specific organs. Treatment can be accomplished by intraarterial infusion. A catheter can be surgically or angiographically implanted to direct treatment to the affected organ. A subcutaneous portal, connected to the catheter, can be used for chronic treatment, or an implantable, refillable pump may also be employed.

The following examples illustrate, but in no way are intended to limit the present invention.

#### Materials and Methods

Restriction endonucleases, T4 DNA ligase, T4 polynucleotide kinase, Taq DNA polymerase, and calf intestinal phosphatase were purchased from New England Biolabs (Beverly, Mass.) or Promega Biotech (Madison, Wis.); these reagents were used according to the manufacturer's instruction. For sequencing reactions, a "SEQUENASE DNA II" sequencing kit was used (United States Biochemical Corporation, Cleveland Ohio). Immunoblotting and other reagents were from Sigma Chemical Co. (St. Louis, Mo.) or Fisher Scientific (Needham, Mass.). Nitrocellulose filters are obtained from Schleicher and Schuell (Keene, N.H.).

Synthetic oligonucleotide linkers and primers are prepared using commercially available automated oligonucleotide synthesizers (e.g., an ABI model 380B-02 DNA synthesizer (Applied Biosystems, Foster City, Calif.)). Alternatively, custom designed synthetic oligonucleotides may be purchased, for example, from Synthetic Genetics (San Diego, Calif.). cDNA synthesis kit and random priming labeling kits are obtained from Boehringer-Mannheim Biochemical (BMB, Indianapolis, Ind.).

Oligonucleotide sequences encoding polypeptides can be either synthesized directly by standard methods of oligonucleotide synthesis, or, in the case of large coding sequences, synthesized by a series of cloning steps involving a tandem array of multiple oligonucleotide fragments corresponding to the coding sequence (Crea; Yoshio et al.; Eaton et al.). Oligonucleotide coding sequences can be expressed by standard recombinant procedures (Maniatis et al.; Ausubel et al.).

Alternatively, peptides can be synthesized directly by standard in vitro techniques (Applied Biosystems, Foster City Calif.).

Common manipulations involved in polyclonal and monoclonal antibody work, including antibody purification from sera, are performed by standard procedures (Harlow et al.). Pierce (Rockford, Ill.) is a source of many antibody reagents.

Recombinant human IFN $\alpha$  (rHuIFN $\alpha$ ) and rBoIFN $\gamma$  was obtained from Genentech Inc. (South San Francisco, Calif.). The reference preparation of recombinant human IFN $\alpha$  (rHuIFN $\alpha$ ) was obtained from the National Institutes of Health: rHuIFN $\alpha$  is commercially available from Lee Biomolecular (San Diego, Calif.).

All tissue culture media, sera and IFNs used in this study were negative for endotoxin, as determined by assay with Limulus amoebocyte lysate (Associates of Cape Cod, Woods Hole, Mass.) at a sensitivity level of 0.07 ng/ml.

#### General ELISA Protocol for Detection of Antibodies.

Polystyrene 96 well plates Immulon II (PGC) were coated with 5  $\mu$ g/mL (100  $\mu$ L per well) antigen in 0.1M carb/bicarbonate buffer, pH 9.5. Plates were sealed with parafilm and stored at 4° C. overnight.

Plates were aspirated and blocked with 300  $\mu$ L 10% NGS and incubated at 37° C. for 1 hr.

Plates were washed 5 times with PBS 0.5% "TWEEN-20".



Antisera were diluted in 0.1M PBS, pH 7.2. The desired dilution(s) of antisera (0.1 mL) were added to each well and the plate incubated 1 hours at 37° C. The plates was then washed 5 times with PBS 0.5% “TWEEN-20”.

Horseradish peroxidase (HRP) conjugated goat antihuman antiserum (Cappel) was diluted 1/5,000 in PBS. 0.1 mL of this solution was added to each well. The plate was incubated 30 min at 37° C., then washed 5 times with PBS.

Sigma ABTS (substrate) was prepared just prior to addition to the plate.

The reagent consists of 50 mL 0.05M citric acid, pH 4.2, 0.078 mL 30% hydrogen peroxide solution and 15 mg ABTS. 0.1 mL of the substrate was added to each well, then incubated for 30 min at room temperature. The reaction was stopped with the addition of 0.050 mL 5% SDS (w/v). The relative absorbance is determined at 410 nm.

EXAMPLE 1

Reproductive Functions of IFNτ

The effect of interferon-τ on the lifespan of the corpus luteum was examined.

IFNτ was infused into uterine lumen of ewes at the concentrations given in Table 1. Recombinant human IFNα (rHuIFNα) was infused at similar concentrations. In addition, control animals, which received control proteins, were also used. The life span of the corpus luteum was assessed by examination of interestrous intervals, maintenance of progesterone secretion, and inhibition of prostaglandin secretion (Davis, et al., 1992).

TABLE 1

Effect of Interferons on Reproductive Physiology		
Interferon	Treatment	Interestrous Interval (days) (Means)
Control	—	17.3
rHuIFNα	100 μg/day	16.0
	200 μg/day	16.0
	2000 μg/day	19.0
OvIFNτ	100 μg/day	27.2

Comparison of the interestrous intervals for the control animals and for animals receiving OvIFNτ demonstrate a considerable lengthening of the interval, when IFNτ is administered at 100 μg/day. On the other hand, comparison of the interestrous interval for the control animal and for animals receiving recombinant human IFNα, demonstrated that rHuIFNα had no meaningful effect.

These results demonstrate that interferon-τ has the capability of significantly influencing the biochemical events of the reproductive cycle.

EXAMPLE 2

Antiviral Properties of Interferon-τ at Various Stages of the Reproductive Cycle

Conceptus cultures were established using conceptus obtained from sheep at days 12 through 16 of the estrous cycle. Antiviral activity of supernatant from each conceptus culture was assessed using a cytopathic effect assay (Familetti, et al., 1981). Briefly, dilutions of IFNτ or other IFNs were incubated with Madin-Darby bovine kidney (MDBK) cells for 16–18 hours at 37° C. Following incubation, inhibition of viral replication was determined in

a cytopathic effect assay using vesicular stomatitis virus (VSV) as the challenge virus.

One antiviral unit caused a 50% reduction in destruction of the monolayer, relative to untreated MDBK cells infected with VSV (control plates). Specific activities were further evaluated using normal ovine fibroblasts (Shnf) in a plaque inhibition assay (Langford, et al., 1981). A minimum of three samples were examined at each time point, and each sample was assayed in triplicate. The results presented in Table 2 are expressed as mean units/ml.

TABLE 2

IFNτ Antiviral Activity of Conceptus Cultures and Allantoic and Amniotic Fluids			
	Day	Samples	Unit/ml
Conceptus Cultures	10	9	<3
	12	5	34
	13	6	4.5 × 10 <sup>3</sup>
	14	3	7.7 × 10 <sup>3</sup>
	16	12	2.0 × 10 <sup>6</sup>
Allantoic Fluid	60	3	1.4 × 10 <sup>3</sup>
	100	4	11
	140	3	<3
Amniotic Fluid	60	3	22
	100	4	<3

Culture supernatants had increasing antiviral activity associated with advancing development of the conceptus (Table 2).

EXAMPLE 3

Expression of IFNτ in Bacteria

The amino acid coding sequence for OvIFNτ (Imakawa, et al., 1987) was used to generate a corresponding DNA coding sequence with codon usage optimized for expression in *E. coli*. Linker sequences were added to the 5' and 3' ends to facilitate cloning in bacterial expression vectors. The nucleotide sequence was designed to include 19 unique restriction enzyme sites spaced evenly throughout the coding sequence (FIGS. 1A and 1B).

The nucleotide sequence was divided into eleven oligonucleotide fragments ranging in sizes of 33 to 75 bases. Each of the eleven oligonucleotides were synthesized on a 380-B 2-column DNA synthesizer (Applied Biosystems) and cloned single- or double-stranded into one of the following vectors: “pBLUESCRIPT+ (KS)” (Stratagene, LaJolla, Calif.), pTZ18R (Pharmacia, Piscataway, N.J.), or pTZ19R (Pharmacia, Piscataway, N.J.) cloning vectors.

The vectors were transformed into *E. coli* K. strain “XL1-BLUE” (recA1 endA1 gyrA96 thi hsdR17 (r<sub>k</sub><sup>-</sup>, m<sub>k</sub><sup>+</sup>) supE44 relA1 λ-(lac), {F<sup>'</sup>, proAB, lacZ<sup>'</sup>ZAM15, Tn10(tet<sup>R</sup>)}) which is commercially available from Stratagene (La Jolla, Calif.). Transformed cells were grown in L broth supplemented with ampicillin (50 μg/ml). Oligonucleotide cloning and fusion was performed using standard recombinant DNA techniques.

Cloning vectors were cut with the appropriate restriction enzymes to insert the synthetic oligonucleotides. The vectors were treated with calf intestine alkaline phosphatase (CIP) to remove terminal phosphate groups. Oligonucleotides were phosphorylated and cloned, as either single- or double-stranded molecules, into the appropriate vector using T4 DNA ligase. When single-strands were introduced into cloning vectors, the second strand was completed by the bacterial host following transfection.



For double-stranded cloning, oligonucleotides were first annealed with their synthetic complementary strand then ligated into the cloning vector. *E. coli* K12 strains SB221 or NM522 were then transformed with the ligation. *E. coli* strain GM119 was used for cloning when the methylation-sensitive *Stu*I and *Cla*I restriction sites were involved. Restriction analyses were performed on isolated DNA at each stage of the cloning procedure.

Cloned oligonucleotides were fused into a single polynucleotide using the restriction digestions and ligations outlined in FIG. 2. oligonucleotide-containing-DNA fragments were typically isolated after electrophoretic size fractionation on low-melting point agarose gels (Maniatis, et al.; Sambrook, et al.; Ausubel, et al.). The resulting IFN $\tau$  polynucleotide coding sequence spans position 16 through 531: a coding sequence of 172 amino acids.

The nucleotide sequence of the final polynucleotide was confirmed by DNA sequencing using the dideoxy chain termination method.

The full length *Stu*I/*Sst*I fragment (540 bp; FIG. 2) was cloned into a modified pIN III omp-A expression vector and transformed into a competent SB221 strain of *E. coli*. For expression of the IFN $\tau$  protein, cells carrying the expression vector were grown in L-broth containing ampicillin to an OD (550 nm) of 0.1–1, induced with IPTG for 3 hours and harvested by centrifugation. Soluble recombinant IFN $\tau$  was liberated from the cells by sonication or osmotic fractionation.

#### EXAMPLE 4

##### Expression of IFN $\tau$ in Yeast

The synthetic IFN $\tau$  gene, synthesized in Example 3, was flanked at the 5' end by an *Stu*I restriction site and at the 3' end by a *Sac*I restriction site.

##### A. Isolation of the Synthetic IFN $\tau$ Gene.

Two oligonucleotide primers (SEQ ID NO:13 and SEQ ID NO:14) were used to attach linkers to the synthetic IFN $\tau$  gene using polymerase chain reaction. The linker at the 5' end allowed the placement of the synthetic IFN $\tau$  gene in correct reading with the ubiquitin coding sequence present in the yeast cloning vector pBS24Ub (Chiron Corp., Emeryville, Calif.). The linker also constructed a ubiquitin-IFN $\tau$  junction region that allowed in vivo cleavage of the ubiquitin sequences from the IFN $\tau$  sequences. The 5' oligonucleotide also encoded a *Sac*II restriction endonuclease cleavage site. The 3' oligonucleotide contained a *Stu*I cleavage site.

The vector carrying the synthetic IFN $\tau$  gene (Example 3) was isolated from *E. coli* strain "XLI-BLUE" by the alkaline lysis method. Isolated vector was diluted 500-fold in 10 mM Tris, pH 8.0/1 mM EDTA/10 mM NaCl. The PCR reaction was performed in a 100  $\mu$ l volume using Taq DNA polymerase and primers SEQ ID NO:13/SEQ ID NO:14. The amplified fragments were digested with *Stu*I and *Sac*II. These digested fragments were ligated into the *Sac*II and *Sma*I sites of "pBLUESCRIPT+(KS)."

The resulting plasmid was named pBSY-IFN $\tau$ . The DNA sequence was verified using double stranded DNA as the template.

##### B. Construction of the Expression Plasmid.

Plasmid pBSY-IFN $\tau$  was digested with *Sac*II and *Eco*RV and the fragment containing the synthetic IFN $\tau$  gene was isolated. The yeast expression vector pBS24Ub (Sabin, et al.; Ecker, et al.) was digested with *Sac*II. Blunt ends were

generated using T4 DNA polymerase. The vector DNA was extracted with phenol and ethanol precipitated (Sambrook, et al., 1989). The recovered linearized plasmid was digested with *Sac*II, purified by agarose gel electrophoresis, and ligated to the *Sac*II-*Eco*RV fragment isolated from pBSY-IFN $\tau$ . The resulting recombinant plasmid was designated pBS24Ub-IFN $\tau$ .

The recombinant plasmid pBS24Ub-IFN $\tau$  was transformed into *E. coli*. Recombinant clones containing the IFN $\tau$  insert were isolated and identified by restriction enzyme analysis. Plasmid DNA from clones containing IFN $\tau$  coding sequences was used for transformation of *S. cerevisiae* (Rothstein, 1986). Transformation mixtures were plated on uracil omission medium and incubated for 3–5 days at 30° C. Colonies were then streaked and maintained on uracil and leucine omission medium (Rothstein, 1986).

##### C. Expression Experiments.

For small-scale expression, a single colony of *S. cerevisiae* AB116 containing pBS24Ub-IFN $\tau$  was picked from a leucine and uracil omission plate and grown at 30° C. in YEP medium (1% yeast extract, 2% peptone) containing 1% glucose for inducing conditions or 8% glucose for noninducing conditions. Cell lysates were recovered and subjected to SDS-PAGE in 15% acrylamide, 0.4% bisacrylamide (Sambrook, et al., 1989). The fractionated proteins were visualized by Coomassie blue staining.

Recombinant IFN $\tau$  was visualized specifically by immunoblotting with monoclonal antibody or polyclonal antiserum against ovine IFN $\tau$  upon electrotransfer of the fractionated cell extract to "NYTRAN" paper (Rothstein, 1986).

For large-scale expression, pBS24-IFN $\tau$  was grown for 24 hours at 30° C. in 5 $\times$ uracil and leucine omission medium containing 8% glucose. This culture was then diluted 20-fold in YEP medium containing 1% glucose and further incubated for another 24–36 hours.

Cells were harvested by centrifugation, washed in 50 mM Tris, pH 7.6, 1 mM EDTA and resuspended in wash buffer containing 1 mM PMSF. The cells were lysed using a Bead-beater apparatus (Biospec Products, Bartlesville, Okla.). The lysate was spun at 43,000 $\times$ g for 20 minutes. The supernatant fraction was recovered and subjected to the purification protocol described below.

##### D. Purification of roIFN $\tau$ from Yeast Cell Lysate.

The supernatant was loaded on a 1 $\times$ 10 cm DEAE column and washed with 10 mM Tris, pH 8.0. Retained proteins were eluted with a 300 ml, 0 to 0.5M NaCl gradient in 10 mM Tris, pH 8.0. Three-milliliter fractions were collected. Ten-microliter samples of fractions 17–26 containing the recombinant (roIFN $\tau$ ) were electrophoretically separated on 15% SDS-polyacrylamide gels. The gels were stained with Coomassie blue.

Fractions 18, 19, and 20 contained largest amount of roIFN $\tau$ . These fractions were loaded individually on a 1.5 $\times$ 90 cm Sephadex S-200 column and proteins were resolved in two peaks. Aliquots of each protein peak (25  $\mu$ l) were electrophoretically separated on 15% SDS-polyacrylamide gels and the proteins visualized with Coomassie staining.

Purified roIFN $\tau$ -containing fractions were combined and the amount of roIFN $\tau$  quantified by radioimmunoassay (Vallet, et al., 1988). Total protein concentration was determined by using the Lowry protein assay (Lowry, et al., 1951).

Microsequencing of purified roIFN $\tau$  demonstrated identity with native IFN $\tau$  through the first 15 amino acids,



confirming that the ubiquitin/roIFN $\tau$  fusion protein was correctly processed in vivo.

Purified roIFN $\tau$  exhibited 2 to  $3 \times 10^8$  units of antiviral activity per milligram of protein ( $n=3$  replicate plates) which is similar to the antiviral activity of IFN $\tau$  purified from conceptus-conditioned culture medium ( $2 \times 10^8$  U/mg).

#### EXAMPLE 5

##### Southern Blot Analysis of Human High Molecular Weight DNA

Human venous blood samples from healthy donors were collected in heparinized tubes and peripheral blood lymphocytes were isolated by density-gradient centrifugation using a Ficoll-Isopaque gradient (1.077 g/ml) (Sigma Chemical Co.). High molecular weight (HMW) DNA was isolated from these cells (Sambrook, et al., 1989).

Two 10  $\mu$ g samples of HMW DNA were digested with the restriction endonucleases HindIII or PstI (Promega) for 2 hours at 37° C., and the DNA fragments electrophoretically separated in a 0.8% agarose gel (Bio-Rad, Richmond, Calif.) at 75 volts for 8 hours. The DNA fragments were transferred onto a nylon membrane (IBI-International Biotechnologies, Inc., New Haven, Conn.). The membrane was baked at 80° C. for 2 hours and incubated at 42° C. for 4 hours in the following prehybridization solution: 5 $\times$ SSC (1 $\times$ SSC is 0.15M NaCl and 0.15M sodium citrate), 50% vol/vol formamide, 0.6% (wt/vol) SDS, 0.5% (wt/vol) nonfat dry milk, 20 mM Tris-HCl (pH 7.5), 4 mM EDTA, and 0.5 mg/ml single stranded herring sperm DNA (Promega).

The filter was then incubated in a hybridization solution (5 $\times$ SSC, 20% vol/vol formamide, 0.6% (wt/vol) SDS, 0.5% (wt/vol) nonfat dry milk, 20 mM Tris-HCl (pH 7.5), 4 mM EDTA, and  $2 \times 10^8$  cpm/ml  $^{32}$ P-labelled OvIFN $\tau$  cDNA (Imakawa, et al., 1987)) for 18 hours at 42° C. The filter was washed at 42° C. for 15 minutes with 2 $\times$ SSC and 0.1% (wt/vol) SDS and exposed to X-ray film (XAR, Eastman Kodak, Rochester, N.Y.) at -80° C. for 48 hours in the presence of an intensifying screen.

Autoradiography detected a hybridization signal at approximately 3.4 kb in DNA digested with PstI and a slightly smaller ( $\approx 3.0$  kb) fragment in the HindIII digested DNA. These results indicate the presence of human DNA sequences complementary to the OvIFN $\tau$  cDNA probe.

#### EXAMPLE 6

##### Isolation of Partial Sequence of Human IFN $\tau$ cDNA by PCR

Two synthetic oligonucleotides (each 25-mer), corresponding to the nucleotides in the DNA sequence from 231 to 255 (contained in SEQ ID NO:13) and 566 to 590 (contained in SEQ ID NO:14) of OvIFN $\tau$  cDNA (numbering relative to the cap site, Imakawa, et al., 1987) were synthesized. These primers contained, respectively, cleavage sites for the restriction endonucleases PstI and EcoRI. SEQ ID NO:13 was modified to contain the EcoRI site, which begins at position 569.

DNA was isolated from approximately  $1 \times 10^5$  plaque forming units (pfu) of the following two cDNA libraries: human term placenta (Clontech, Inc., Palo Alto, Calif.) and human term cytotrophoblast (Dr. J. F. Strauss, University of Pennsylvania, Philadelphia Pa.). The DNA was employed in polymerase chain reaction (PCR) amplifications (Mullis; Mullis, et al.; Perkin Elmer Cetus Corp. Norwalk Conn.).

Amplification reactions were carried out for 30 cycles (45° C., 1 m; 72° C., 2 m; 94° C., 1 m) (thermal cycler and reagents, Perkin Elmer Cetus) using primers SEQ ID NO:13/SEQ ID NO:14.

Amplification products were electrophoretically separated (100 volts in a 1.5% agarose gel (Bio-Rad)) and transferred onto a nylon membrane (IBI). The membrane was baked at 80° C. for 2 hours and prehybridized and hybridized with  $^{32}$ P-labelled OvIFN $\tau$  cDNA as described above. The membrane was washed in 5 $\times$ SSC/0.1% (wt/vol) SDS for 5 minutes at 42° C. and in 2 $\times$ SSC/0.1% (wt/vol) SDS for 2 minutes at 42° C. It was then exposed at -80° C. to "XAR" (Eastman Kodak) X-ray film for 24 hours in the presence of an intensifying screen. An amplification product that hybridized with the labelled probe DNA was detected.

PCR was performed again as directed above. Amplified products were digested with the restriction endonucleases EcoRI and PstI (Promega) for 90 minutes at 37° C. The resulting DNA fragments were electrophoretically separated as described above and the band containing the IFN $\tau$  amplification product was excised from the gel. DNA fragments were recovered by electroelution, subcloned into EcoRI/PstI digested-dephosphorylated plasmid pUC19 and transformed into *E. coli* strain JM101 (Promega) by calcium chloride method (Sambrook, et al., 1989). The plasmids were isolated and the inserted amplification product sequenced using the dideoxy termination method (Sanger, et al., 1977; "SEQUENCE" reactions, United States Biochemical, Cleveland, Ohio). Nucleotide sequences were determined, and comparison of these as well as the deduced amino acid sequences to other IFN sequences were performed using "DNA STAR SOFTWARE" (Madison, Wis.).

Comparison of the sequences of these clones revealed the following four different clones: from the human placental library, HuIFN $\tau$ 6 (299 bp), HuIFN $\tau$ 7 (288 bp) and HuIFN $\tau$ 4 (307 bp), which exhibit 95% identity in their nucleotide sequences; from the cytotrophoblast library clone CTB 35 (HuIFN $\tau$ 5; 294 basepairs), which shares 95% and 98% identity with HuIFN $\tau$ 6 and HuIFN $\tau$ 4, respectively.

#### EXAMPLE 7

##### Isolation of Full-Length Human IFN $\tau$ Genes

Ten micrograms PBMC HMW DNA was digested with restriction endonuclease EcoRI and subjected to electrophoretic analysis in a 0.8% agarose gel. A series of samples containing ranges of DNA fragments sized 1.5 to 10 kb (e.g., 1.5 to 2.5 kb, 2.5 kb to 3 kb) were excised from the gel. The DNAs were electroeluted and purified. Each DNA sample was amplified as described above using the OvIFN $\tau$  primers. The DNA molecules of any sample that yielded a positive PCR signal were cloned into  $\lambda$ gt11 (the subgenomic  $\lambda$ gt11 library).

A. PCR Identification of Clones Containing Sequences Complementary to OvIFN $\tau$ .

The  $\lambda$ gt11 phage were then plated for plaques and plaque-lift hybridization performed using the  $^{32}$ P-labelled OvIFN $\tau$  cDNA probe. Approximately 20 clones were identified that hybridized to the probe.

Plaques that hybridized to the probe were further analyzed by PCR using the OvIFN $\tau$  primers described above. Six plaques which generated positive PCR signals were purified. The phage DNA from these clones was isolated and digested with EcoRI restriction endonuclease. The DNA inserts were subcloned into pUC19 vectors and their nucleotide sequences determined by dideoxy nucleotide sequencings.



### B. Hybridization Identification of Clones Containing Sequences Complementary to PCR-Positive Phage.

Recombinant phage from the  $\lambda$ gt11 subgenomic library were propagated in *E. coli* Y1080 and plated with *E. coli* Y1090 at a density of about 20,000 plaques/150 mm plate. The plates were overlaid with duplicate nitrocellulose filters, which were hybridized with a  $^{32}$ P-labelled probe from one of the six human IFN $\tau$  cDNA clones isolated above.

Clones giving positive hybridization signals were further screened and purified. The phage DNAs from hybridization-positive clones were isolated, digested with EcoRI, subcloned into pUC19 vector and sequenced. The sequence information was then analyzed.

#### 1. HuIFN $\tau$ 1

Three clones yielded over-lapping sequence information for over 800 bases relative to the mRNA cap site (clones were sequenced in both orientations). The combined nucleic acid sequence information is presented as SEQ ID NO:11 and the predicted protein coding sequence is presented as SEQ ID NO:12. Comparison of the predicted mature protein sequence (SEQ ID NO:12) of this gene to the predicted protein sequence of OvIFN $\tau$  is shown in FIG. 3.

#### 2. HuIFN $\tau$ 2, HuIFN $\tau$ 3

Two additional clones giving positive hybridization signals (HuIFN $\tau$ 2 and HuIFN $\tau$ 3) were also screened, purified, and phage DNAs subcloned and sequenced as above. The sequences of these two clones are presented in FIGS. 19A and 19B. As can be appreciated in FIGS. 19A and 19B, the nucleotide sequence of both clones (HuIFN $\tau$ 2 and HuIFN $\tau$ 3) is homologous to that of HuIFN $\tau$ 1 and OvIFN $\tau$ .

HuIFN $\tau$ 2 (SEQ ID NO:29), may be a pseudo-gene, as it appears to contain a stop codon at position 115–117. The sequence, SEQ ID NO:29, is presented without the leader sequence. The leader sequence is shown in FIG. 20A. As can be seen from the HuIFN $\tau$ 2 sequence presented in FIG. 20A, the first amino acid present in mature HuIFN $\tau$ 1 (a CYS residue) is not present in the HuIFN $\tau$ 2 sequence. Accordingly, the predicted amino acid sequence presented as SEQ ID NO:29 corresponds to a mature IFN $\tau$  protein with the exceptions of the first CYS residue and the internal stop codon.

The internal stop codon in the nucleic acid coding sequence can be modified by standard methods to replace the stop codon with an amino acid codon, for example, encoding GLN. The amino acid GLN is present at this position in the other isolates of human IFN $\tau$  (HuIFN $\tau$ ). Standard recombinant manipulations also allow introduction of the initial CYS residue if so desired.

HuIFN $\tau$ 3 (SEQ ID NO:31), appears to encode a human IFN $\tau$  protein. The translated amino acid sequence of the entire protein, including the leader sequence, is presented as SEQ ID NO:32. The translated amino acid sequence of the mature protein is presented as SEQ ID NO:34.

### EXAMPLE 8

#### Analysis of the Presence of HuIFN $\tau$ mRNA by RT-PCR

Human placental cDNA libraries and an ovine cDNA library, constructed from day 15–16 conceptuses, were analyzed by hybridization to the OvIFN $\tau$  cDNA probe, described above. cDNAs were size-fractionated on agarose gels and transferred to filters (Maniatis, et al.; Sambrook, et al.). Southern blot analysis with OvIFN $\tau$  probe showed that the autoradiographic signals from human cDNA libraries

were approximately  $1/100$  of the signal obtained using the OvIFN $\tau$  cDNA library.

The presence of HuIFN $\tau$  mRNA in human term placenta and amniocytes (26 weeks, 2 million cells) was analyzed by using reverse transcriptase-PCR (RT-PCR) method (Clontech Laboratories, Palo Alto Calif.).

Total cellular RNA (tcRNA) isolated from human placenta, amniocytes and ovine conceptuses were reverse transcribed using the primer SEQ ID NO:14. The primer SEQ ID NO:13 was then added to the reaction and polymerase chain reaction carried out for 40 cycles. The PCR products were size fractionated on agarose gels and transferred to filters. The DNA on the filters was hybridized with  $^{32}$ P-labelled OvIFN $\tau$  and HuIFN $\tau$  cDNAs. The results of these analyses demonstrate the presence of human IFN $\tau$  mRNA in the feto-placental annex. The amniocytes also expressed the messages corresponding to OvIFN $\tau$  primers and human probe.

In addition, a RT-PCR analysis for the presence of HuIFN $\tau$  was applied to the tcRNA isolated from human adult lymphocytes. A densitometric analysis revealed that IFN $\tau$  mRNA exists in lymphocytes.

### EXAMPLE 9

#### In Situ Hybridization

##### A. Tissue

Slides of semiserial 5- $\mu$  paraffin embedded sections from four healthy, different term and first trimester human placentas were examined.

##### B. cRNA Probe Preparation

From the cDNA clone isolated from OvIFN $\tau$  amplified library a fragment corresponding to the OvIFN $\tau$  cDNA bases #77–736 (base #1 is cap site; open reading frame of OvIFN $\tau$  cDNA is base #81–665; FIG. 7) was subcloned into the transcription vector, pBS (New England Biolabs). Several pBS clones were isolated, subcloned, and their nucleotides sequenced. From this clone a 3' fragment (bases #425–736) was excised using the restriction endonucleases NlaIV and EcoRI and subcloned into the transcription vector pBS. This vector was designated pBS/OvIFN $\tau$ .

After linearization of the pBS/OvIFN $\tau$  plasmid, an anti-sense cRNA probe was synthesized by in vitro transcription (Sambrook, et al., 1989) using T<sub>7</sub> RNA polymerase (Stratagene). A trace amount of  $^3$ H-CTP (NEN-DuPont, Cambridge, Mass.) was used in the transcription reaction. dUTP labeled with digoxigenin (Boehringer-Mannheim, Indianapolis, Ind.) was incorporated into the cRNA and yield was estimated through TCA precipitation and scintillation counting.

##### C. Hybridization

In situ hybridization was performed using the anti-sense RNA probe, as described by Lawrence, et al. (1985) with the following modifications. Deparaffinized and hydrated sections were prehybridized for 10 minutes at room temperature in phosphate buffered saline (PBS) containing 5 mM MgCl<sub>2</sub>. Nucleic acids in the sections were denatured for 10 minutes at 65° C. in 50% formamide/2 $\times$ SSC. Sections were incubated overnight at 37° C. with a hybridization cocktail (30  $\mu$ l/slide) containing 0.3  $\mu$ g/ml digoxigenin-labelled cRNA probe and then washed for 30 minutes each at 37° C. in 50 formamide/1 $\times$ SSC. Final washes were performed for 30 minutes each at room temperature in 1 $\times$ SSC and 0.1 $\times$ SSC. The sections were blocked for 30 minutes with 0.5% Triton X-100 (Sigma) and 0.5% non-fat dry milk.



Hybridization signal was detected using purified sheep antidioxigenin Fab fragments conjugated to alkaline phosphatase (Boehringer-Mannheim). After unbound antibody was removed, nitroblue tetrazolium/5-bromo-4-chloro-3-indolyl-phosphate substrate (Promega) and levamisole (Bector Laboratories, Burlingame, Calif.) Were added for signal detection via calorimetric substrate generation. The tissues were counterstained in methyl green (Sigma), dehydrated, and mounted.

As a control, some tissue sections were pretreated with 100  $\mu$ g/ml of pancreatic RNaseA (Sigma) for 30 minutes at 37° C. The RNase was inactivated on the slide with 400 units of RNase inhibitor (Promega). The slides were then washed twice in 250 ml of PBS/5 mM MgCl<sub>2</sub>. In other control experiments, tRNA (Sigma) was substituted for the digoxigenin probes.

Specific hybridization was observed in all term and first trimester placental tissues in three separate experiments with various OvIFN $\tau$  cRNA probe concentrations and blocking reagents.

First trimester placental villi composed of an outer layer of syncytiotrophoblast, an underlying layer of cytotrophoblast, and a central stromal region with various types of mesenchymal cells, displayed the highest transcript level of IFN $\tau$  in the cytotrophoblast cells. Less intense but detectable levels were present in both the syncytiotrophoblast and stromal cells. A similar pattern of transcript expression was demonstrated in the placental villi of term tissue but the level of signal detection was low. First trimester extravillous trophoblast displayed the highest amount of message and stained positive when present in the maternal blood spaces.

EXAMPLE 10

Antiviral Activity of IFN $\tau$

The relative specific activity of OvIFN $\tau$ , purified to homogeneity, was evaluated in antiviral assays. The antiviral assays were performed essentially as described above in Example 2. Specific activities are expressed in antiviral units/mg protein obtained from antiviral assays using either Madin-Darby bovine kidney (MDBK) cells or sheep normal fibroblasts (Shnf). All samples were assayed simultaneously to eliminate interassay variability. The results, presented in Table 3, are the means of four determinations where the standard deviation was less than 10% of the mean.

TABLE 3

	Antiviral Activity of IFN $\tau$ and Known IFNS	
	Specific Activities	
	MDBK	Shnf
OvIFN $\tau$	2 $\times$ 10 <sup>8</sup>	3 $\times$ 10 <sup>8</sup>
rBoIFN $\alpha$	6 $\times$ 10 <sup>7</sup>	1 $\times$ 10 <sup>7</sup>
rBoIFN $\gamma$	4.5 $\times$ 10 <sup>6</sup>	3 $\times$ 10 <sup>6</sup>
NIH rHuIFN $\alpha$	2.2 $\times$ 10 <sup>8</sup>	2.2 $\times$ 10 <sup>8</sup>
rHuIFN $\alpha$	2.9 $\times$ 10 <sup>5</sup>	4.3 $\times$ 10 <sup>5</sup>

IFN $\tau$  had a higher specific activity than either rBoIFN $\alpha$  or rBoIFN $\tau$  (Table 3). The NIH standard preparation of rHuIFN $\alpha$  had a similar specific activity, while a commercial preparation of rHuIFN $\alpha$  exhibited low specific antiviral

activity. Comparable relative antiviral activity was demonstrated using either bovine or ovine cells.

EXAMPLE 11

Anti-Retroviral Activity and Cytotoxic Effects of IFN $\tau$

Highly purified OvIFN $\tau$  was tested for anti-retroviral and cytotoxic effects on feline peripheral blood lymphocytes exposed to the feline immunodeficiency retrovirus. This lentivirus produces a chronic AIDS-like syndrome in cats and is a model for human AIDS (Pederson, et al., 1987). Replication of the virus in peripheral blood lymphocytes is monitored by reverse transcriptase activity in culture supernatants over time. The data from these assays are presented in Table 4.

TABLE 4

Effect of OvIFN $\tau$ on FIV Replication						
IFN $\tau$ Concentration (ng/ml)	RT Activity (cpm/ml)					
	Harvest Days					
	Experiment 1	Day 2	Day 5	Day 8	Day 12	Day 15
0.00		93,908	363,042	289,874	171,185	125,400
0.62		77,243	179,842	172,100	218,281	73,039
1.25		94,587	101,873	122,216	71,916	50,038
2.50		63,676	72,320	140,783	75,001	36,105
5.00		69,348	82,928	90,737	49,546	36,299
	Harvest Days					
	Experiment 2	Day 2	Day 5	Day 8	Day 13	Day 17
0.0		210,569	305,048	279,556	500,634	611,542
2.5		121,082	106,815	108,882	201,676	195,356
5.0		223,975	185,579	108,114	175,196	173,881
10.0		167,425	113,631	125,131	131,649	129,364
20.0		204,879	80,399	59,458	78,277	72,179
40.0		133,768	54,905	31,606	72,580	53,493

Addition of OvIFN $\tau$  produced a rapid, dose-dependent decrease in reverse transcriptase (RT) activity (Table 4). While concentrations as low as 0.62 ng/ml of IFN $\tau$  inhibited viral replication, much higher concentrations (40 ng/ml) having greater effects on RT-activity were without toxic effects on the cells. The results suggest that replication of the feline immunodeficiency virus was reduced significantly compared to control values when cells were cultured in the presence of OvIFN $\tau$ .

IFN $\tau$  appeared to exert no cytotoxic effect on the cells hosting the retrovirus. This was true even when IFN $\tau$  was present at 40 ng per ml of culture medium.

EXAMPLE 12

Effects of IFN $\tau$  on HIV Infected Human Peripheral Lymphocytes

IFN $\tau$  was also tested for activity against HIV infection in human cells. Human peripheral blood lymphocytes, which had been infected with HIV (Crowe, et al.), were treated with varying concentrations of OvIFN $\tau$ . Replication of HIV in peripheral blood lymphocytes was monitored by reverse transcriptase activity in culture supernatants over time.



Reverse transcriptase activity was measured essentially by the method of Hoffman, et al. The data from these assays are presented in Table 5.

TABLE 5

Effect of OvIFN $\tau$ on HIV Replication in Human Peripheral Lymphocytes				
IFN $\tau$ Concentration (ng/ml)	RT Activity			
	Day 6		Day 10	
	cpm/ml	% Reduction	cpm/ml	% Reduction
0	4,214	—	25,994	—
10	2,046	51	9,883	62
50	1,794	57	4,962	81
100	1,770	58	3,012	88
500	1,686	60	2,670	90
1000	1,499	64	2,971	89

As shown in Table 5, concentrations of OvIFN $\tau$  produced significant antiviral effects. A concentration of only 10 ng/ml resulted in over a 50% reduction in RT activity after only six days. A concentration of 500 ng/ml resulted in a 90% reduction in RT activity within 10 days.

The viability of human peripheral blood lymphocytes after treatment with IFN $\tau$ , over a range of concentrations for 3–13 days, was evaluated by trypan blue exclusion. The results of this viability analysis are presented in Table 6.

TABLE 6

Effect of OvIFN $\tau$ on Viability of HIV Infected Human Peripheral Lymphocytes			
IFN $\tau$ Concentration (ng/ml)	Viable Cells/ml $\times 10^5$		
	Day 3	Day 6	Day 13
0	16.0	7.5	5.3
10	13.0	7.5	6.0
50	13.0	11.5	9.0
100	15.0	8.5	9.5
500	16.5	12.0	11.0
1000	21.9	9.5	8.5

The data presented in Table 6 show no evidence of cytotoxic effects attributable to the administration of IFN $\tau$ .

EXAMPLE 13

Inhibition of Cellular Growth

The effects of IFN $\tau$  on cellular growth were also examined. Anti-cellular growth activity was examined using a colony inhibition assay. Human amnion (WISH) or MDBK cells were plated at low cell densities to form colonies originating from single cells. Cells were cultured at 200 or 400 cells/well in 24 well plates in HMEM supplemented with 2% fetal bovine serum (FBS) and essential and non-essential amino acids. Various dilutions of interferons were added to triplicate wells, and the plates were incubated for 8 days to allow colony formation. Colonies were visualized after staining with crystal violet, and counted. Cell cycle analysis was performed with HMEM containing 0.5% “spent” media for an additional 7 days. WISH cells were used without being synchronized.

For examination of IFN $\tau$  activity, cells were replated at  $2.5 \times 10^5$  cells/well in HMEM with 10% FBS in 6 well plates.

Various dilutions of ovIFN $\tau$  alone or in combination with peptides were added to achieve a final volume of 1 ml. Plates were incubated at 37° C. in 5% Co<sub>2</sub> for 12, 15, 18, 24, or 48 hours. Cells were treated with trypsin, collected by low speed centrifugation and washed. The cell pellet was blotted dry and 250  $\mu$ l of nuclear staining solution (5 mg propidium iodide, 0.3 ml NP40 and 0.1  $\mu$ m sodium citrate in 100 ml distilled H<sub>2</sub>O) was added to each tube. The tubes were incubated at room temperature. After 10 minutes, 250  $\mu$ l of RNase (500 units/ml in 1.12% sodium citrate) was added per tube and incubated an additional 20 minutes. Nuclei were filtered through 44  $\mu$ m mesh, and analyzed on a FACStar (Becton Dickinson, Mountain View, Calif.) using the DNA Star 2.0 software.

In the cellular growth assay using colony formation of both the bovine epithelial line, MDBK, and the human amniotic line, WISH, OvIFN $\tau$  inhibited both colony size and number. Ovine IFN $\tau$  was more effective than human IFN $\alpha$  on the human cell line; thus, it is very potent in cross-species activity. Its activity was dose-dependent, and inhibition of proliferation could be observed at concentrations as low as 1 unit/ml. Concentrations as high as 50,000 units/ml (units of antiviral activity/ml) stopped proliferation, while cell viability was not impaired.

Cell cycle analysis by flow cytometry with propidium iodide-stained WISH cells revealed an increased proportion of cells in G2/M after 48 hours of OvIFN $\tau$  treatment. IFN $\tau$ , therefore, appears to inhibit progress of cells through S phase. Ovine IFN $\tau$  antiproliferative effects can be observed as early as 12 hours after the initiation of culture and are maintained through 6 days.

The results presented above demonstrate both the antiproliferative effect of IFN $\tau$  as well as its low cytotoxicity.

EXAMPLE 14

Further Antiproliferative Effects of IFN $\tau$

The antiproliferative effects of OvIFN $\tau$  were studied for a rat cell line and a bovine cell line. The rate of <sup>3</sup>H-thymidine incorporation was used to assess the rate of cellular proliferation.

Rat (MtBr7 .c5) or bovine kidney (MDBK) cells were seeded in phenol red-free DME-F12 medium supplemented with 3% dextran-coated charcoal stripped Controlled Process Serum Replacement 2 (CPSR 2, Sigma) and 5% dextran-coated charcoal stripped fetal bovine serum (FBS). After attaching for approximately 15–18 hours, the cells were washed once with serum-free DME-F12 medium. The medium was replaced with phenol red-free DME-F12 medium supplemented with 3% stripped CPSR2, 1% stripped FBS (“3/1” medium) or 3/1 medium containing OvIFN $\tau$  at various units of antiviral activity as determined in the vesicular stomatitis virus challenge assay for interferons (Example 2). Media containing a similar dilution of buffer (undiluted buffer=10 mM Tris, 330 mM NaCl, [TS]), in which the OvIFN $\tau$  was dissolved was used for controls.

Cells were pulse labeled with <sup>3</sup>H-thymidine for 2 hours at approximately 48 hours post-treatment. The trichloroacetic acid (TCA) precipitable incorporated counts were determined by scintillation counting. Three replicates were included per treatment. Mean values for OvIFN $\tau$  treatments were compared to samples containing comparable dilutions of carrier TS buffer. Results of these experiments are shown in Table 7.



TABLE 7

<sup>3</sup> H-Thymidine Incorporation	
Treatment	% Reduction <sup>3</sup> H-Thymidine Incorporation
Experiment 1: MtBr7 .c5 (Rat)	
3/1	—
10 <sup>3</sup> u OvIFNτ/ml	0 (+12)
1:5000 TS	—
10 <sup>4</sup> u OvIFNτ/ml	24
1:500 TS	—
10 <sup>5</sup> OvIFNτ/ml	87
Experiment 2: MDBK	
3/1	—
10 <sup>3</sup> u OvIFNτ/ml	74
1:5000 TS	—
10 <sup>4</sup> u OvIFNτ/ml	83
1:500 TS	—
10 <sup>5</sup> u OvIFNτ/ml	83

As can be seen from Table 7, OvIFNτ drastically reduced the rate of cellular proliferation (based on thymidine incorporation) for each of the cell lines tested.

EXAMPLE 15

Antiproliferative Effects of IFNτ on Human Tumor Cell Lines

The antiproliferative activity of OvIFNτ on human tumor cell lines was evaluated by measuring the rate of <sup>3</sup>H-thymidine incorporation into cells which have been treated with OvIFNτ.

For experiments on tumor lines that grow in suspension, 1 ml of cells were plated at from 2.5–5×10<sup>5</sup> cells/well in 24-well plates. Triplicate wells received either the appropriate media, 100, 1,000 or 10,000 units/ml of OvIFNτ or equivalent antiviral concentrations of rHuIFNα2A (Lee Biomolecular). After 48 hours of incubation, cells were counted and viability assessed by trypan blue exclusion.

Adherent tumor lines were plated at 2.5×10<sup>5</sup> cells/well in 1 ml in 6-well plates. They received interferon treatments as just described, but were trypsinized prior to counting.

Significant differences between treatments were assessed by an analysis of variance followed by Scheffe's F-test. Cell cycle analysis was performed by flow cytometry using propidium iodide.

A. Breast Adenocarcinoma Cells.

Human MCF7 breast adenocarcinoma cells were seeded from logarithmically growing cultures in phenol red-free DME-F12 medium supplemented with 3% dextran-coated charcoal stripped CPSR and 5% dextran-coated FBS. After attaching for approximately 15–18 hours, the cells were washed once with serum-free DME-F12 medium. The medium was replaced with phenol red-free DME-F12 medium supplemented with 3% stripped CPSR2, 1% stripped FBS ("3/1" medium) or 3/1 medium containing OvIFNτ at the indicated number of units of antiviral activity as determined in the vesicular stomatitis virus challenge assay for interferons. Media containing a similar dilution of buffer (undiluted buffer=10 mM Tris, 330 mM NaCl [TS]) was used for controls. Cells were pulse labeled with <sup>3</sup>H-thymidine for 2 hours at approximately 48 hours post-treatment.

The trichloroacetic acid (TCA) precipitable incorporated counts were determined by scintillation counting. Three

replicates were included per treatment. Mean values for OvIFNτ treatments were compared to samples containing comparable dilutions of carrier TS buffer. The results of these analyses are shown in Table 8.

TABLE 8

<sup>3</sup> H-Thymidine Incorporation	
Treatment	% Reduction <sup>3</sup> H-Thymidine Incorporation
MCF7 Human	
3/1	—
10 <sup>3</sup> u OvIFNτ/ml	35
1:5000 TS	—
10 <sup>4</sup> u OvIFNτ/ml	53
1:500 TS	—
10 <sup>5</sup> u OvIFNτ/ml	70

As can be seen from the results shown in Table 8, OvIFNτ was able to substantially reduce the rate of <sup>3</sup>H-thymidine incorporation in the human carcinoma cell line. This demonstrates the efficacy of OvIFNτ in inhibiting tumor cell proliferation, in particular, mammary tumor cell proliferation.

B. Human Promyelocytic Leukemia.

A comparison of the antiproliferative effects of OvIFNτ and IFNα was conducted using HL-60 (human leukemia) cells (Foa, et al.; Todd, et al.) essentially as described above for MDBK cells. Both OvIFNτ and rHuIFNα inhibit HL-60 cell proliferation. Results of one of three replicate experiments are presented as mean % growth reduction ± SD in FIG. 4. FIG. 4 shows that both OvIFNτ and IFNα were able to drastically reduce growth of HL-60 cells. The growth reduction for each compound exceeded 60% for each concentration tested. At 10,000 units/ml, OvIFNτ caused an approximately 80% reduction in growth while IFNα caused a 100% reduction in growth.

However, the data presented in FIG. 4 reveal, that a substantial factor in the ability of IFNα to reduce growth was its toxic effect on the cells. At 10,000 units/ml, the toxicity of IFNα resulted in less than 25% of the cells remaining viable. By contrast, nearly 100% of the cells remained viable when OvIFNτ was applied at 10,000 units/ml.

FIG. 5 presents data demonstrating that rHuIFNα is cytotoxic. In the figure, results of one of three replicate experiments are presented as mean % viability ± SD.

C. Human Cutaneous T Cell Lymphoma.

The cutaneous T cell lymphoma, HUT 78, responded similarly to HL-60 when treated with IFNτ (FIG. 9). Both OvIFNτ and rHuIFNα reduce HUT 78 cell growth, but 10,000 units/ml of rHuIFNα decreased the cell number below that originally plated (533 10<sup>5</sup>). This is indicative of a reduction in cell viability to approximately 60%.

Cell cycle analysis (performed by cell flow cytometry) revealed an increased proportion of cells in G2/M phase of the cell cycle upon 48 hours of treatment with both interferons (Table 9). In Table 9 the results from one of three replicate experiments are presented as the percentage of cells in each phase of the cell cycle. 10,000 events were analyzed per sample.

This result is likely due to the slower progress of cells through the cell cycle. In the sample treated with 10,000 units/ml of rHuIFNα, a large percentage of events with low forward and high side scatter, identifying dead cells, were



present. This is consistent with the data obtained from proliferation experiments, where only OvIFN $\tau$  inhibited HUT 78 proliferation without toxicity.

TABLE 9

HUT 78 Cell Cycle Analysis.			
Treatment (units/ml)	G0/G1	S	G2/M
Media	44.43	49.95	5.61
100 OvIFN $\tau$	44.35	47.45	8.20
100 rHuIFN $\alpha$	40.01	57.53	2.45
1,000 OvIFN $\tau$	41.29	50.50	8.21
1,000 rHuIFN $\alpha$	41.73	44.91	13.36
10,000 OvIFN $\tau$	42.79	42.61	14.60
10,000 rHuIFN $\alpha$	18.01	71.31	10.67
			(cell death)

D. Human T Cell Lymphoma.

The T cell lymphoma cell line H9 was slightly less sensitive to the antiproliferative effects of the IFNs than the tumor cell lines described above. Results of one of three replicate experiments are presented in FIG. 10 as mean % growth reduction  $\pm$  SD. While rHuIFN $\alpha$  was not toxic to the H9 cells, it failed to inhibit cell division significantly at any of the concentrations examined. In contrast, OvIFN $\tau$  was observed to reduce H9 growth by approximately 60% (FIG. 10). Thus, only OvIFN $\tau$  is an effective growth inhibitor of this T cell lymphoma.

The results presented above demonstrate both the antiproliferative effect of IFN $\tau$  as well as its low cytotoxicity.

EXAMPLE 16

Preliminary In Vivo Treatment with OvIFN $\tau$

Three groups of 4 C57Bl/6 mice per group were given  $2.5 \times 10^4$  B16-F10 cells via the tail vein: B16-F10 is a syngeneic mouse transplantable tumor selected because of its high incidence of pulmonary metastases (Poste, et al., 1981). Interferon treatment was initiated 3 days after the introduction of the tumor cells. Each mouse received 100  $\mu$ l of either PBS alone, PBS containing  $1 \times 10^5$  units of OvIFN $\tau$ , or PBS containing  $1 \times 10^5$  units of recombinant murine IFN $\alpha$  (MuIFN $\alpha$ ), i.v. per day for 3 consecutive days.

Mice were sacrificed at 21 days and the lungs were preserved in 10% buffered formalin. The frequency of pulmonary metastases were compared between control mice (PBS), OvIFN $\tau$ -treated mice, and MuIFN $\alpha$ -treated mice. The results of these in vivo administrations demonstrated that OvIFN $\tau$  dramatically reduced B16-F10 pulmonary tumors. These results support the use of IFN $\tau$  as an efficacious antineoplastic agent in vivo.

EXAMPLE 17

Competitive Binding of IFN $\tau$  Peptide Fragments

A. The Ability of IFN $\tau$ -Based Peptides to Block IFN $\tau$  and IFN- $\alpha$  Antiviral Activity.

Overlapping synthetic peptides were synthesized corresponding to the entire IFN $\tau$  sequence (FIG. 6). Average hydropathicity values were calculated by taking the sum of the hydropathy values for each amino acid divided by the total number of amino acids in each sequence. Hydropathy values were taken from Kyte, et al. (1982).

These peptides were of approximately the same molecular weight but differed slightly in overall hydrophilicity. Despite

this difference, all peptides were antigenic as demonstrated by the production of rabbit antisera with titers greater than 1:3,000 as assessed by ELISA (Harlow, et al.).

The peptides were used to inhibit the antiviral activity (Example 2) of OvIFN $\tau$  and rBoIFN $\alpha$ . The results of this analysis are presented in FIG. 12: 1 mM N- and C-terminal peptides both effectively blocked the antiviral activity of OvIFN $\tau$  using MDBK cells. A third peptide, representing amino acids 62–92, also reduced IFN $\tau$  antiviral activity (70% inhibition). The peptide OvIFN $\tau$  (119–150) showed minimal inhibitory activity. The OvIFN $\tau$  (34–64) and (90–122) peptides had no apparent inhibitory activity.

Peptide inhibition of OvIFN $\tau$  antiviral activity was also examined as follows. Monolayers of Madin Darby bovine kidney cells were incubated with 40 units/ml OvIFN $\tau$  in the presence or absence of various concentrations of OvIFN $\tau$  peptides (see FIG. 13). Results in FIG. 13 are expressed as the percent of control antiviral activity: that is, in the absence of any competing peptide. Data presented are the means of 6 replicate experiments. The data demonstrate that inhibition by OvIFN $\tau$  (1–37), (62–92), (119–150), and (139–172) were significantly different than OvIFN $\tau$  (34–64) and (90–122) at  $10^{-3}$ M and  $3 \times 10^{-3}$ M. OvIFN $\tau$  (139–172) was significantly different than all other peptides at  $10^{-3}$ M. Significance was assessed by analysis of variance followed by Scheffe's F test at  $p < 0.05$ . Thus, OvIFN $\tau$  (1–37) (62–92), (119–150), and (139–172), in particular (139–172), may represent receptor binding regions for IFN $\tau$ .

The ability of the OvIFN $\tau$  peptides to inhibit bovine IFN $\alpha$  (BoIFN $\alpha$ ) antiviral activity was examined as follows. Monolayers of Madin Darby bovine kidney cells were incubated with 40 units/ml bovine IFN $\alpha$  in the presence or absence of various concentrations of OvIFN $\tau$  peptides. The results are presented in FIG. 14 and are expressed as the percent of control antiviral activity in the absence of OvIFN $\tau$  peptides. The data presented are the means of 4 replicate experiments. The results indicate that inhibition by OvIFN $\tau$  (62–92), (119–150), and (139–172) were significantly different from OvIFN $\tau$  (1–37), (34–64) and (90–122) at  $10^{-3}$ M. OvIFN $\tau$  (139–172) was significantly different than OvIFN $\tau$  (1–37), (34–64) and (90–122) at  $3 \times 10^{-3}$ M. Significance was assessed by analysis of variance followed by Scheffe's F test at  $p < 0.05$ . Thus, OvIFN $\tau$  (62–92), (119–150), and (139–172), in particular (139–172), may represent common receptor binding regions for IFN $\tau$  and bovine IFN $\alpha$ .

Peptide inhibition by OvIFN $\tau$  peptides of human IFN $\alpha$  antiviral activity was also examined. Monolayers of Madin Darby bovine kidney cells were incubated with 40 units/ml human IFN $\alpha$  in the presence or absence of various concentrations of OvIFN $\tau$  peptides. The results are expressed as the percent of control antiviral activity in the absence of OvIFN $\tau$  peptides. The data are presented in FIG. 15 and are the means of 3 replicate experiments. OvIFN $\tau$  (139–172) was significantly different from all other peptides at  $10^{-3}$ M. Significance was assessed by analysis of variance followed by Scheffe's F test at  $p < 0.05$ . Thus, OvIFN $\tau$  (139–172) may represent a common receptor binding region for IFN $\tau$  and various IFN $\alpha$ (s).

The OvIFN $\tau$  peptides described above appear to have no effect on the antiviral activity of IFN $\gamma$ . Peptide inhibition of bovine IFN $\gamma$  antiviral activity was evaluated as follows. Monolayers of Madin Darby bovine kidney cells were incubated with 40 units/ml bovine IFN gamma in the presence or absence of various concentrations of OvIFN $\tau$  peptides. Results are expressed as the percent of control



antiviral activity in the absence of OvIFN $\tau$  peptides. The data are presented in FIG. 16 and are the means of 3 replicate experiments. There were no significant differences among peptides as assessed by analysis of variance followed by Scheffe's F test at  $p<0.05$ .

The two synthetic peptides OvIFN $\tau$  (1–37) and OvIFN $\tau$  (139–172) also blocked OvIFN $\tau$  anti-FIV and anti-HIV activity. Reverse transcriptase (RT) activity (Examples 12 and 13) was monitored over a 14 day period in FIV-infected FET-1 cells ( $1\times 10^6$ /ml) and HIV-infected HPBL ( $1\times 10^6$ /ml). Control cultures received no OvIFN $\tau$ . OvIFN $\tau$  was used at 100 ng/ml, and peptides were used at 200  $\mu$ M. Data from a representative experiment are expressed as cpm/ml culture supernatant and are presented for FIV infected cells, FIG. 11A, and HIV infected cells, FIG. 11B. Both the N- and C-terminus of OvIFN $\tau$  appear to be involved in its anti-retroviral activity. While both peptides blocked FIV RT activity, only the C-terminal peptide, OvIFN $\tau$  (139–172), was an efficient inhibitor of vesicular stomatitis virus activity on the feline cell line, Fc9. Thus the C-terminal regions of type I IFNs may bind to common site on the type I IFN receptor, while the N-terminal region may be involved in the elicitation of unique functions.

B. Anti-Peptide Sera.

The ability of anti-peptide antisera to inhibit OvIFN $\tau$  antiviral activity was also determined. Antipeptide antisera inhibition of OvIFN $\tau$  antiviral activity was evaluated as follows. Monolayers of MDBK cells were incubated with 20 units/ml of OvIFN $\tau$  in the presence a 1:30 dilution of either preimmune sera or antisera to each of the OvIFN $\tau$  peptides described above. In FIG. 17 the data from duplicate experiments are presented as the mean percent inhibition of OvIFN $\tau$  antiviral activity produced by antipeptide antisera relative to the appropriate preimmune sera  $\pm$  standard error. Significant differences were assessed by analysis of variance followed by Scheffe's F test at  $p<0.05$ . Consistent with peptide inhibition of antiviral activities, sera containing antibodies immunoreactive to OvIFN $\tau$  (1–37), OvIFN $\tau$  (62–92), and OvIFN $\tau$  (139–172) were also the most effective inhibitors of OvIFN $\tau$  antiviral activity, with antibodies directed against the N-terminal and C-terminal peptides being the most efficacious.

The same sera were also used to examine their effect on the binding of IFN $\tau$  to its receptor.

The IFN $\tau$  binding assay was carried out as follows. Five  $\mu$ g of IFN $\tau$  was iodinated for 2 minutes with 500  $\mu$ Ci of Na<sup>126</sup>I (15 mCi/ $\mu$ g; Amersham corporation, Arlington Heights, Ill.) in 25  $\mu$ l of 0.5M potassium phosphate buffer,

pH 7.4, and 10  $\mu$ l of chloramine-T (5 mg/ml) (Griggs, et al., 1992). The specific activity of the iodinated protein was 137  $\mu$ Ci/ $\mu$ g. For binding assays, monolayers of MDBK cells were fixed with paraformaldehyde and blocked with 5% nonfat dry milk. Cells were incubated with 5 nM <sup>125</sup>I-IFN $\tau$  in phosphate buffered saline with 1% BSA for 2 hours at 4° C. in the presence or absence of a 1:30 dilution of sera containing antibodies raised against IFN $\tau$  peptides or the appropriate preimmune sera. Specific binding was assessed by incubation with a 100-fold molar excess of unlabeled IFN $\tau$ . Specific binding of 36% was determined by competition with 500 nM unlabeled IFN $\tau$ . For example, total counts bound were  $6850\pm 133$ , and a 100-fold molar excess of OvIFN $\tau$  produced  $4398\pm 158$  counts per minute. After incubation, the monolayers were washed three times, solubilized with 1% sodium dodecyl sulfate, and the radioactivity counted. Data from three replicate experiments are presented in FIG. 18 as the mean percent reduction of OvIFN $\tau$  specific binding produced by antipeptide antisera relative to the appropriate preimmune sera  $\pm$  standard deviation. Significant differences were assessed by analysis of variance followed by Scheffe's F test.

The same sera (containing antibodies immunoreactive to OvIFN $\tau$  (1–37), OvIFN $\tau$  (62–92), and OvIFN $\tau$  (139–172)) were the most effective inhibitors of <sup>125</sup>I-IFN $\tau$  binding to its receptor on MDBK cells. The lack of effect of sera immunoreactive with other IFN $\tau$ -derived peptides was not a function of titer against OvIFN $\tau$ , since each sera had equal or greater titer to their respective peptide relative to the three inhibiting sera: similar results were obtained when sera reactivity against the whole OvIFN $\tau$  molecule was assessed by ELISA for each sera.

These peptides, although apparently binding to the interferon receptor, did not in and of themselves elicit interferon-like effects in the cells.

C. Anti-Proliferative Activity.

Functionally important sites for the antiproliferative activity of IFN $\tau$  were also examined using synthetic peptides (Table 10). Cellular proliferation was assayed as described above using MDBK cells. MDBK cells were cultured at  $5\times 10^5$  cells/well in experiments 1 and 2 or  $10\times 10^5$  cells in experiment 3 and treated with medium alone, IFN $\tau$  at a concentration of 300 units/ml and peptides at 1 mM for 48 hours. Duplicate wells were counted in each of three replicate experiments. For statistical analysis, data were normalized based on medium alone and assessed by analysis of variance followed by Least Significant Difference multiplate comparison test ( $p>0.05$ ).

TABLE 10

Peptide Inhibition of IFN $\tau$ Antiproliferative Activity						
Treatment	Experiment 1		Experiment 2		Experiment 3	
	Cell Count	Via-bility	Cell Count	Via-bility	Cell Count	Via-bility
Medium alone	$9.8 \times 10^5$	99%	$13.0 \times 10^5$	96%	$27.3 \times 10^5$	97%
IFN $\tau$	$5.0 \times 10^5$	98%	$5.6 \times 10^5$	97%	$8.3 \times 10^5$	97%
IFN $\tau$ +IFN $\tau$ (1–37)	$6.3 \times 10^5$	100%	$10.6 \times 10^5$	98%	$13.4 \times 10^5$	100%
IFN $\tau$ +IFN $\tau$ (34–64)	$5.3 \times 10^5$	96%	$6.9 \times 10^5$	95%	$16.0 \times 10^5$	98%
IFN $\tau$ +IFN $\tau$ (62–82)	$6.5 \times 10^5$	97%	$9.2 \times 10^5$	93%	$8.9 \times 10^5$	96%
IFN $\tau$ +IFN $\tau$ (90–122)	$5.9 \times 10^5$	100%	$11.0 \times 10^5$	97%	$19.6 \times 10^5$	98%
IFN $\tau$ +IFN $\tau$ (119–150)	$8.4 \times 10^5$	100%	$13.2 \times 10^5$	96%	$31.8 \times 10^5$	90%
IFN $\tau$ +IFN $\tau$ (139–172)	$5.1 \times 10^5$	100%	$12.7 \times 10^5$	98%	$18.9 \times 10^5$	98%



When proliferation of MDBK cells was monitored over a two-day period, cell number increased roughly 2-fold with greater than 95% viability. Addition of 300 units/ml of OvIFN $\tau$  entirely eliminated cell proliferation without a decrease in cell viability. Ovine IFN $\tau$  (119–150) was the most effective inhibitor of IFN $\tau$  antiproliferative activity.

Antisera to IFN $\tau$  (119–150), which inhibited binding of OvIFN $\tau$  to receptor, also reversed the OvIFN $\tau$  antiproliferative effect. Several other peptides, notably IFN $\tau$  (139–172), reversed the OvIFN $\tau$  antiproliferative effect, but to a lesser extent.

EXAMPLE 18

Further Analysis of the Cellular and Anti-Viral Effects of IFN $\tau$

A. HIV Anti-Viral Effects.

The antiviral effects of IFN $\tau$  against HIV were evaluated by treating human PBMC cells with various amounts of either recombinant ovine IFN $\tau$  (r-OvIFN $\tau$ ) or recombinant human IFN $\alpha$ 2a at the time of infection with HIV. IFN $\tau$  was present throughout the experiment. At day 7 and day 14, p24 production was determined (by ELISA (Wang, et al., 1988, 1989) and compared to a zero drug control. The results of this analysis are presented in Table 11.

TABLE 11

Amounts of Drug Units/ml		% Inhibition	% Inhibition
IFN $\alpha$ 2a	IFN $\tau$	Day 7	Day 14
10		58%, 48%	91%, 91%
	26	48%, 45%	88%, 59%
100		68%, 74%	94%, 91%
	260	58%, 51%	82%, 70%
1,000		89%, 86%	97%, 93%
	2,600	65%, 68%	87%, 79%
10,000		90%, 86%	99%, 99%
	26,000	77%, 85%	77%, 96%
	260,000	85%, 84%	96%, 86%

The data from these experiments support the conclusion that, at relatively low concentrations, IFN $\alpha$ 2a and IFN $\tau$  are effective in reducing the replication of HIV in human lymphocytes.

B. In vitro Cytotoxicity Test in PBMC's

Human PBMC's were seeded at 5 $\times$ 10<sup>5</sup> cells/ml. Cells were stimulated at day 0 with 3  $\mu$ g/ml PHA. Cells were treated with recombinant human IFN $\alpha$ 2A (at concentrations of 10, 100, 1,000 and 10,000 units/ml) and IFN $\tau$  (at concentrations of 2.6, 26, 260, 2,600, 26,000, 260,000, and 2,600,000 units/ml) in 200  $\mu$ l/wells (4 replicates of each concentration using 96 well flat bottom plates). Control cultures were given no interferons. After 4 days of incubation, cells were pulsed for 9 hours using <sup>3</sup>H-thymidine at 1 uCi/well. The cells were harvested and the incorporation of labeled thymidine into DNA was determined (FIG. 8).

No cytotoxicity was observed by measuring the uptake of thymidine at any concentration of IFN $\tau$ . However, rHuIFN $\alpha$ 2 was toxic to cells at 1,000 units/ml.

In a second experiment, the same human PBMC's were treated with either IFN $\tau$  or human IFN $\alpha$ 2A at concentrations of 100 units/ml or 10,000 units/ml. After 3 days or 8 days of incubation, viable cells were counted by flow cytometry. The results of this analysis are presented in Table 12.

TABLE 12

Treatment (units/ml)	Number of Viable Cells $\times$ 10,000	
	Day 3	Day 8
No Treatment	735	840
IFN $\tau$ 100 units/ml	745	860
IFN $\tau$ 10,000 units/ml	695	910
IFN $\alpha$ 2a 100 units/ml	635	750
IFN $\alpha$ 2a 10,000 units/ml	680	495

No cytotoxicity was observed in the cells treated with IFN $\tau$ . However, there was 10% cell death in IFN $\alpha$ 2a treated cells at Day 3 and 49% cell death at Day 8.

C. Inhibition of Hepatitis B Virus DNA Replication in Hepatocytes

The cell line used, HepG2-T14, is a human cell that was derived from liver cells transfected with Hepatitis B Virus (HBV). The cell line semi-stably produces HBV virus: over time the cell line's production of HBV intracellular DNA and secreted virus decreases. In order to maximize production of HBV DNA and virus, the cells are pre-treated with deAZA-C (5-azacytidine; Miyoshi, et al.) to induce production of the virus. Treatment was for 2–3 days and the amount of induction was about a factor of two.

The cells were then treated with either the IFN $\alpha$  and IFN $\tau$  at levels of 0, 5,000, 10,000, 20,000 and 40,000 units per ml.

All levels of either IFN $\alpha$  or IFN $\tau$  reduced DNA production by about a factor of 2 compared to the no drug control.

D. Inhibition of Hepatospecific Messenger RNA Production in Hepatocytes

The hepatocyte cell line HepG2-T14 (described above) was examined for the effects of IFN $\alpha$  and IFN $\tau$  on hepatospecific mRNA production. Cells were incubated in concentrations of IFN $\alpha$  or IFN $\tau$  at 0, 5,000, 10,000, 20,000, and 40,000 units per ml. The messenger RNAs for the hepatocyte specific proteins Apo E and Apo A1 were detected by hybridization analysis (Sambrook, et al.; Maniatis, et al.) using probes specific for these two mRNA's (Shoulders, et al., and Wallis, et al.).

No reduction of mRNA production was seen for Apo E or Apo A1 mRNA production with up to 40,000 units of either IFN $\alpha$  or IFN $\tau$ . This result suggests that the reduction of viral DNA replication in previous experiments was not due to the effects of IFNs on cellular house-keeping activities; rather the reduction was likely due to specific inhibition of viral replication in the host cells.

E. In Vitro Toxicity of IFN $\beta$ , IFN $\gamma$  and IFN $\tau$ —L929 Cell Assay

The toxicity of IFN treatment was measured in vitro using the mouse L929 cell line. L929 cells were treated with 6000 U/ml to 200,000 U/ml of either OvIFN $\tau$  or MuIFN $\beta$ . The interferons were added at time zero and the cells were incubated for 72 hours and stained with crystal violet. The percentage of living cells was determined by measuring the absorbance at 405 nm.

Exemplary data are shown in FIG. 21. Values are presented as percent viability  $\pm$  standard error in which 100 percent is equal to the viability of L929 cells treated with media alone. At 6000 U/ml, IFN $\beta$ -treated cells exhibited a 77.0 $\pm$ 0.6% viability. Viability of L929 cells decreased as the concentrations of IFN $\beta$  increased in a dose-dependent manner. In contrast, L929 cells showed no decrease in viability at any of the IFN $\tau$  concentrations tested. These data indicate that, unlike IFN $\beta$ , IFN $\tau$  lacks toxicity at high concentrations in vitro.



Taken together, the results summarized above demonstrate that IFN $\tau$  is essentially non-toxic at concentrations at which IFN $\beta$  induces toxicity both in vitro and in vivo.

F. In Vivo Toxicity of IFN $\beta$ , IFN $\gamma$  and IFN $\tau$ —Cell Counts and Weight Changes

The effects of in vivo treatment with IFN $\tau$ , IFN $\beta$  and IFN $\alpha$  ( $10^5$  U/injection) on total white blood cell (WBC), total lymphocyte counts and weight measurements in NZW mice were assessed as follows. Interferons (OvIFN $\tau$ , MuIFN $\beta$ , and MuIFN $\alpha$ ) were injected intraperitoneally (i.p.) at a concentration of  $10^5$  U in a total volume of 0.2 ml in PBS into groups of New Zealand White (NZW) mice (Jackson Laboratories, Bar Harbor, Me.). Three to four animals were included in each group. White blood cell (WBC) counts were determined before injection and at selected timepoints thereafter (typically 12 and 24 hours) using a hemocytometer and standard techniques. Differential WBC counts were performed on Wright-Giemsa stained blood smears. The Before injection, the weights of the animals ranged from 20 to 23 grams. The results are summarized in Table 13, below.

TABLE 13

IN VIVO TOXICITY OF INTERFERONS AS MEASURED BY WHITE BLOOD CELL COUNTS AND PERCENT WEIGHT CHANGE						
Cell Count (Cell No. $\times 10^3$ )					% Lympho- cyte	% Weight Change 24
Before Injection		12 hr. after Injection		Depres- sion	Hours after Injection	
IFN	Total WBC	Lymphocytes	Total WBC	Lymphocytes		
none	7.3 $\pm$ 1.0	6.4 $\pm$ 0.7	8.0 $\pm$ 0.8	7.1 $\pm$ 0.7	0	+0.5 $\pm$ 0.7
$\tau$	6.7 $\pm$ 0.7	5.9 $\pm$ 0.6	6.7 $\pm$ 0.5	5.8 $\pm$ 0.4	1.7	+1.3 $\pm$ 0.5
$\beta$	7.0 $\pm$ 1.4	6.0 $\pm$ 0.5	6.8 $\pm$ 0.8	4.1 $\pm$ 0.3	31.7	-20.0 $\pm$ 1.0
$\alpha$	6.0 $\pm$ 0.8	5.2 $\pm$ 0.7	4.8 $\pm$ 0.5	2.3 $\pm$ 0.2	55.8	-8.5 $\pm$ 2.0

No significant differences in WBC counts, lymphocyte counts or weight change were observed between IFN $\tau$ -treated and untreated mice. In contrast, IFN $\beta$ -treated mice exhibited a 31.7% depression in lymphocyte counts 12 hours after injection, which continued for at least the next 12 hours. IFN $\alpha$ -treated mice exhibited a 55.8% lymphocyte depression and significant weight loss 12 hours after injection. These data indicate that, unlike IFN $\beta$  and IFN $\alpha$ , IFN $\tau$  lacks toxicity in vivo at the above concentrations as evidenced by peripheral blood cell counts and weight measurements.

EXAMPLE 19

Isolation of Interferon- $\tau$  Fusion Protein

Sepharose 4B beads conjugated with anti-beta galactosidase is purchased from Promega. The beads are packed in 2 ml column and washed successively with phosphate-buffered saline with 0.02% sodium azide and 10 ml TX buffer (10 mM Tris buffer, pH 7.4, 1% aprotinin).

The IFN $\tau$  coding sequence (e.g., SEQ ID NO:33, i.e., minus the nucleotides corresponding to the leader sequence) is cloned into the polylinker site of lambda gt11. The IFN $\tau$  coding sequence is placed in-frame with the amino terminal  $\beta$ -galactosidase coding sequences in lambda gt11. Lysogens infected with gt11/IFN $\tau$  are used to inoculate 500 ml of NZYDT broth. The culture is incubated at 32° C. with aeration to an O.D. of about 0.2 to 0.4, then brought to 43°

C. quickly in a 43° C. water bath for 15 minutes to induce gt11 peptide synthesis, and incubated further at 37° C. for 1 hour. The cells are pelleted by centrifugation, suspended in 10 ml of lysis buffer (10 mM Tris, pH 7.4 containing 2% “TRITON X-100” and 1% aprotinin added just before use.

The resuspended cells are frozen in liquid nitrogen then thawed, resulting in substantially complete cell lysis. The lysate is treated with DNaseI to digest bacterial and phage DNA, as evidenced by a gradual loss of viscosity in the lysate. Non-solubilized material is removed by centrifugation.

The clarified lysate material is loaded on the Sepharose column, the ends of the column closed, and the column placed on a rotary shaker for 2 hrs. at room temperature and 16 hours at 4° C. After the column settles, it is washed with 10 ml of TX buffer. The fused protein is eluted with 0.1M carbonate/bicarbonate buffer, pH10. Typically, 14 ml of the elution buffer is passed through the column, and the fusion protein is eluted in the first 4–6 ml of eluate.

The eluate containing the fusion protein is concentrated in “CENTRICON-30” cartridges (Amicon, Danvers, Mass.).

The final protein concentrate is resuspended in, for example, 400  $\mu$ l PBS buffer. Protein purity is analyzed by SDS-PAGE.

For polyclonal antibodies, the purified fused protein is injected subcutaneously in Freund’s adjuvant in a rabbit. Approximately 1 mg of fused protein is injected at days 0 and 21, and rabbit serum is typically collected at 6 and 8 weeks.

EXAMPLE 20

Preparation of Anti-IFN $\tau$  Antibody

A. Expression of Glutathione-S-Transferase Fusion Proteins.

The IFN $\tau$  coding sequence (e.g., SEQ ID NO:33) is cloned into the pGEX vector (Boyer, et al.; Frangioni, et al.; Guan, et al.; Hakes, et al.; Smith, et al., 1988). The pGEX vector (Smith, et al.) was modified by insertion of a thrombin cleavage sequence in-frame with the glutathione-S-transferase protein (GST—sj26 coding sequence). This vector is designated pGEXthr. The IFN $\tau$  coding sequence is placed in-frame with the sj26-thrombin coding sequences (Guan, et al.; Hakes, et al.). The IFN $\tau$  coding sequence insert can be generated by the polymerase chain reaction using PCR primers specific for the insert.

The IFN $\tau$  fragment is ligated to the linearized pGEXthr vector. The ligation mixture is transformed into *E. coli* and ampicillin resistant colonies are selected. Plasmids are iso-



lated from the ampicillin resistant colonies and analyzed by restriction enzyme digestion to identify clones containing the IFN $\tau$  insert (vector designated pGEXthr-IFN $\tau$ ).

*E. coli* strain XL-I Blue is transformed with pGEXthr-IFN $\tau$  and is grown at 37° C. overnight. DNA is prepared from randomly-picked colonies. The presence of the insert coding sequence is typically confirmed by (i) restriction digest mapping, (ii) hybridization screening using labelled IFN $\tau$  probes (i.e., Southern analysis), or (iii) direct DNA sequence analysis.

#### B. Partial Purification of Fusion Proteins.

ApGEXthr-IFN $\tau$  clone is grown overnight. The overnight culture is diluted 1:10 with LB medium containing ampicillin and grown for one hour at 37° C. Alternatively, the overnight culture is diluted 1:100 and grown to OD of 0.5–1.0 before addition of IPTG (isopropylthio- $\beta$ -galactoside). IPTG (GIBCO-BRL, Gaithersburg Md.) is added to a final concentration of 0.2–0.5 mM for the induction of protein expression and the incubation is typically continued for 2–5 hours, preferably 3.5 hours.

Bacterial cells are harvested by centrifugation and resuspended in 1/100 culture volume of MTPBS (150 mM NaCl, 16 mM Na<sub>2</sub>HPO<sub>4</sub>, 4 mM NaH<sub>2</sub>PO<sub>4</sub>). Cells are lysed by lysozyme, sonication or French press, and lysates cleared of cellular debris by centrifugation.

An aliquot of the supernatant obtained from IPTG-induced cultures of pGEXthr-IFN $\tau$ -containing cells and an aliquot of the supernatant obtained from IPTG-induced cultures of pGEXthr-vector alone are analyzed by SDS-polyacrylamide gel electrophoresis followed by Western blotting, as described below.

If necessary, the extracts can be concentrated by ultrafiltration using, for example, a “CENTRICON 10” filter.

Alternatively, the fusion proteins are partially purified over a glutathione agarose affinity column as described in detail by Smith, et al. In this method, 100 ml cultures are grown overnight. The cultures are diluted to 1 liter, and the cells grown another hour at 37° C. Expression of the fusion proteins is induced using IPTG. The induced cultures are grown at 37° C. for 3.5 hours. Cells are harvested and a sonicator used to lyse the cells. Cellular debris is pelleted and the clear lysate loaded onto a glutathione

“SEPHAROSE” column. The column is washed with several column volumes. The fusion protein is eluted from the affinity column with reduced glutathione and dialyzed. The IFN $\tau$  can be liberated from the hybrid protein by treatment with thrombin. The sj26 and IFN $\tau$  fragments of the hybrid protein can then be separated by size fractionation over columns or on gels.

Alternatively, the IFN $\tau$  portion of the hybrid protein is released from the column by treatment with thrombin (Guan, et al.; Hakes, et al.).

#### C. Antibodies Against the Fusion Protein.

The purified Sj26/IFN $\tau$  fused protein is injected subcutaneously in Freund’s adjuvant in a rabbit. Approximately 1 mg of fused protein is injected at days 0 and 21, and rabbit serum is typically collected at 6 and 8 weeks. A second rabbit is similarly immunized with purified Sj26 protein obtained from control bacterial lysate.

Minilysates from the following bacterial cultures are prepared: (1) KM392 cells infected with pGEXthr and pGEXthr containing the IFN $\tau$  insert; and (2) cells infected with lambda gt11 containing the IFN $\tau$  insert. The minilysates and a commercial source  $\beta$ -galactosidase are fractionated by SDS-PAGE, and the bands transferred to nitrocellulose filters for Western blotting (Sambrook, et al.; Ausubel, et al.).

Summarizing the expected results, serum from control (Sj26) rabbits is immunoreactive with each of the Sj26 and Sj26 fused protein antigens. Serum from the animal immunized with Sj26/IFN $\tau$  fused protein is reactive with all Sj-26 and beta-gal fusion proteins containing IFN $\tau$  coding sequences, indicating the presence of specific immunoreaction with the IFN $\tau$  antigen. None of the sera are expected to be immunoreactive with beta-galactosidase.

Anti-IFN $\tau$  antibody present in the sera from the animal immunized with the Sj26/IFN $\tau$  is purified by affinity chromatography (using immobilized recombinantly produced IFN $\tau$  as ligand, essentially as described above in Example 12 for the anti-beta-galactosidase antibody).

While the invention has been described with reference to specific methods and embodiments, it will be appreciated that various modifications and changes may be made without departing from the invention.

---

#### SEQUENCE LISTING

##### (1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 44

##### (2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 516 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: double  
 (D) TOPOLOGY: circular

(ii) MOLECULE TYPE: DNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
 (A) ORGANISM: *Ovis aries*  
 (B) STRAIN: Domestic



-continued

(D) DEVELOPMENTAL STAGE: Blastula (blastocyst)  
(F) TISSUE TYPE: Trophectoderm  
(G) CELL TYPE: Mononuclear trophectoderm cells

(vii) IMMEDIATE SOURCE:  
(B) CLONE: oTP-1a

(viii) POSITION IN GENOME:  
(C) UNITS: bp

(ix) FEATURE:  
(A) NAME/KEY: CDS  
(B) LOCATION: 1..516

(x) PUBLICATION INFORMATION:  
(A) AUTHORS: Ott, Troy L  
Van Heeke, Gino  
Johnson, Howard M  
Bazer, Fuller W  
(B) TITLE: Cloning and Expression in *Saccharomyces cerevisiae*  
of a Synthetic Gene for the Type I Trophoblast  
Interferon Ovine Trophoblast  
Protein-1:Purification and Antiviral Activity  
(C) JOURNAL: J. Interferon Res.  
(D) VOLUME: 11  
(F) PAGES: 357-364  
(G) DATE: 1991  
(K) RELEVANT RESIDUES IN SEQ ID NO:1: FROM 1 TO 516

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

TGC	TAC	CTG	TCG	CGA	AAA	CTG	ATG	CTG	GAC	GCT	CGA	GAA	AAT	TTA	AAA	48
Cys	Tyr	Leu	Ser	Arg	Lys	Leu	Met	Leu	Asp	Ala	Arg	Glu	Asn	Leu	Lys	
1				5					10					15		
CTG	CTG	GAC	CGT	ATG	AAT	CGA	TTG	TCT	CCG	CAC	AGC	TGC	CTG	CAA	GAC	96
Leu	Leu	Asp	Arg	Met	Asn	Arg	Leu	Ser	Pro	His	Ser	Cys	Leu	Gln	Asp	
			20					25					30			
CGG	AAA	GAC	TTC	GGT	CTG	CCG	CAG	GAA	ATG	GTT	GAA	GGT	GAC	CAA	CTG	144
Arg	Lys	Asp	Phe	Gly	Leu	Pro	Gln	Glu	Met	Val	Glu	Gly	Asp	Gln	Leu	
			35				40					45				
CAA	AAA	GAC	CAA	GCT	TTC	CCG	GTA	CTG	TAT	GAA	ATG	CTG	CAG	CAG	TCT	192
Gln	Lys	Asp	Gln	Ala	Phe	Pro	Val	Leu	Tyr	Glu	Met	Leu	Gln	Gln	Ser	
			50				55				60					
TTC	AAC	CTG	TTC	TAC	ACT	GAA	CAT	TCT	TCG	GCC	GCT	TGG	GAC	ACT	ACT	240
Phe	Asn	Leu	Phe	Tyr	Thr	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	
					70					75					80	
CTT	CTA	GAA	CAA	CTG	TGC	ACT	GGT	CTG	CAA	CAG	CAA	CTG	GAC	CAT	CTG	288
Leu	Leu	Glu	Gln	Leu	Cys	Thr	Gly	Leu	Gln	Gln	Gln	Leu	Asp	His	Leu	
					85				90					95		
GAC	ACT	TGC	CGT	GGC	CAG	GTT	ATG	GGT	GAA	GAA	GAC	TCT	GAA	CTG	GGT	336
Asp	Thr	Cys	Arg	Gly	Gln	Val	Met	Gly	Glu	Glu	Asp	Ser	Glu	Leu	Gly	
			100					105					110			
AAC	ATG	GAT	CCG	ATC	GTT	ACT	GTT	AAA	AAA	TAT	TTC	CAG	GGT	ATC	TAC	384
Asn	Met	Asp	Pro	Ile	Val	Thr	Val	Lys	Lys	Tyr	Phe	Gln	Gly	Ile	Tyr	
			115				120					125				
GAC	TAC	CTG	CAG	GAA	AAA	GGT	TAC	TCT	GAC	TGC	GCT	TGG	GAA	ATC	GTA	432
Asp	Tyr	Leu	Gln	Glu	Lys	Gly	Tyr	Ser	Asp	Cys	Ala	Trp	Glu	Ile	Val	
			130			135					140					
CGC	GTT	GAA	ATG	ATG	CGG	GCC	CTG	ACT	GTG	TCG	ACT	ACT	CTG	CAA	AAA	480
Arg	Val	Glu	Met	Met	Arg	Ala	Leu	Thr	Val	Ser	Thr	Thr	Leu	Gln	Lys	
			145			150				155				160		
CGG	TTA	ACT	AAA	ATG	GGT	GGT	GAC	CTG	AAT	TCT	CCG					516
Arg	Leu	Thr	Lys	Met	Gly	Gly	Asp	Leu	Asn	Ser	Pro					
				165				170								

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 172 amino acids

-continued

(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: amino acid sequence of a mature  
OvIFNtau protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Cys Tyr Leu Ser Arg Lys Leu Met Leu Asp Ala Arg Glu Asn Leu Lys  
1 5 10 15  
Leu Leu Asp Arg Met Asn Arg Leu Ser Pro His Ser Cys Leu Gln Asp  
20 25 30  
Arg Lys Asp Phe Gly Leu Pro Gln Glu Met Val Glu Gly Asp Gln Leu  
35 40 45  
Gln Lys Asp Gln Ala Phe Pro Val Leu Tyr Glu Met Leu Gln Gln Ser  
50 55 60  
Phe Asn Leu Phe Tyr Thr Glu His Ser Ser Ala Ala Trp Asp Thr Thr  
65 70 75 80  
Leu Leu Glu Gln Leu Cys Thr Gly Leu Gln Gln Gln Leu Asp His Leu  
85 90 95  
Asp Thr Cys Arg Gly Gln Val Met Gly Glu Glu Asp Ser Glu Leu Gly  
100 105 110  
Asn Met Asp Pro Ile Val Thr Val Lys Lys Tyr Phe Gln Gly Ile Tyr  
115 120 125  
Asp Tyr Leu Gln Glu Lys Gly Tyr Ser Asp Cys Ala Trp Glu Ile Val  
130 135 140  
Arg Val Glu Met Met Arg Ala Leu Thr Val Ser Thr Thr Leu Gln Lys  
145 150 155 160  
Arg Leu Thr Lys Met Gly Gly Asp Leu Asn Ser Pro  
165 170

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 516 base pairs  
(B) TYPE: nucleic acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: synthetic nucleotide sequence encoding  
a mature human interferon-tau protein, HuIFNtau1.

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TGTGACTTGT CTCAAAACCA CGTTTTGGTT GGTAGAAAGA ACTTAAGACT ACTAGACGAA 60  
ATGAGACGTC TATCTCCACG CTTCTGTCTA CAAGACAGAA AGGACTTCGC TTTGCCTCAG 120  
GAAATGGTTG AAGGTGGCCA ACTACAAGAA GCTCAAGCGA TATCTGTTTT GCACGAAATG 180  
TTGCAACAAA GCTTCAACTT GTTCCACACC GAACACTCTT CGGCCGCTTG GGACACCACC 240  
TTGTTGGAAC AGCTCAGAAC CGGTTTGCAC CAACAATTGG ACAACTTGGA TGCATGTTTG 300  
GGTCAAGTTA TGGGTGAAGA AGACTCTGCT CTCGGGAGAA CCGGTCCAAC GCTAGCTTTG 360  
AAGAGATACT TCCAAGGTAT CCACGTTTAC TTGAAGGAAA AGGGTTACTC TGA CTGTGCT 420  
TGGGAAACCG TCGTCTAGA AATCATGCGT AGCTTCTCTT CTTTGATCAG CTTGCAAGAA 480  
AGATTACGTA TGATGGACGG TGA CTGTGCTG AGCCCA 516



-continued

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 172 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: amino acid sequence for a mature HuIFNtau protein, HuIFNtau1.
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Cys Asp Leu Ser Gln Asn His Val Leu Val Gly Arg Lys Asn Leu Arg  
1 5 10 15  
Leu Leu Asp Glu Met Arg Arg Leu Ser Pro Arg Phe Cys Leu Gln Asp  
20 25 30  
Arg Lys Asp Phe Ala Leu Pro Gln Glu Met Val Glu Gly Gly Gln Leu  
35 40 45  
Gln Glu Ala Gln Ala Ile Ser Val Leu His Glu Met Leu Gln Gln Ser  
50 55 60  
Phe Asn Leu Phe His Thr Glu His Ser Ser Ala Ala Trp Asp Thr Thr  
65 70 75 80  
Leu Leu Glu Gln Leu Arg Thr Gly Leu His Gln Gln Leu Asp Asn Leu  
85 90 95  
Asp Ala Cys Leu Gly Gln Val Met Gly Glu Glu Asp Ser Ala Leu Gly  
100 105 110  
Arg Thr Gly Pro Thr Leu Ala Leu Lys Arg Tyr Phe Gln Gly Ile His  
115 120 125  
Val Tyr Leu Lys Glu Lys Gly Tyr Ser Asp Cys Ala Trp Glu Thr Val  
130 135 140  
Arg Leu Glu Ile Met Arg Ser Phe Ser Ser Leu Ile Ser Leu Gln Glu  
145 150 155 160  
Arg Leu Arg Met Met Asp Gly Asp Leu Ser Ser Pro  
165 170

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 37 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: amino acid sequence of fragment 1-37 of SEQ ID NO:2
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Cys Tyr Leu Ser Arg Lys Leu Met Leu Asp Ala Arg Glu Asn Leu Lys  
5 10 15  
Leu Leu Asp Arg Met Asn Arg Leu Ser Pro His Ser Cys Leu Gln Asp  
20 25 30  
Arg Lys Asp Phe Gly  
35

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 31 amino acids
  - (B) TYPE: amino acid

-continued

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: amino acid sequence of fragment 34-64  
of SEQ ID NO:2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Lys Asp Phe Gly Leu Pro Gln Glu Met Val Glu Gly Asp Gln Leu Gln  
5 10 15

Lys Asp Gln Ala Phe Pro Val Leu Tyr Glu Met Leu Gln Gln Ser  
20 25 30

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 31 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: amino acid sequence of fragment  
62-92 of SEQ ID NO:2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Gln Gln Ser Phe Asn Leu Phe Tyr Thr Glu His Ser Ser Ala Ala Trp  
5 10 15

Asp Thr Thr Leu Glu Gln Leu Cys Thr Gly Leu Gln Gln Gln  
20 25 30

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: amino acid sequence of fragment  
90-122 of SEQ ID NO:2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Gln Gln Gln Leu Asp His Leu Asp Thr Cys Arg Gly Gln Val Met Gly  
5 10 15

Glu Glu Asp Ser Glu Leu Gly Asn Met Asp Pro Ile Val Thr Val Lys  
20 25 30

Lys

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 32 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: amino acid sequence of fragment  
119-150 of SEQ ID NO:2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Thr Val Lys Lys Tyr Phe Gln Gly Ile Tyr Asp Tyr Leu Gln Glu Lys  
5 10 15



-continued

Gly Tyr Ser Asp Cys Ala Trp Glu Ile Val Arg Val Glu Met Met Arg  
20 25 30

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 34 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: amino acid sequence of fragment 139-172 of SEQ ID NO:2

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Cys Ala Trp Glu Ile Val Arg Val Glu Met Met Arg Ala Leu Thr Val  
5 10 15

Ser Thr Thr Leu Gln Lys Arg Leu Thr Lys Met Gly Gly Asp Leu Asn  
20 25 30

Ser Pro

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 588 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: HuIFNtau1 Human Interferon Tau coding sequence with a leader sequence.

- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..585

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

ATG GCC TTC GTG CTC TCT CTA CTC ATG GCC CTG GTG CTG GTC AGC TAC	48
Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr	
1 5 10 15	
GGC CCA GGA GGA TCC CTG GGT TGT GAC CTG TCT CAG AAC CAC GTG CTG	96
Gly Pro Gly Gly Ser Leu Gly Cys Asp Leu Ser Gln Asn His Val Leu	
20 25 30	
GTT GGC AGG AAG AAC CTC AGG CTC CTG GAC GAA ATG AGG AGA CTC TCC	144
Val Gly Arg Lys Asn Leu Arg Leu Leu Asp Glu Met Arg Arg Leu Ser	
35 40 45	
CCT CGC TTT TGT CTG CAG GAC AGA AAA GAC TTC GCT TTA CCC CAG GAA	192
Pro Arg Phe Cys Leu Gln Asp Arg Lys Asp Phe Ala Leu Pro Gln Glu	
50 55 60	
ATG GTG GAG GGC GGC CAG CTC CAG GAG GCC CAG GCC ATC TCT GTG CTC	240
Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile Ser Val Leu	
65 70 75 80	
CAT GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC CAC ACA GAG CAC TCC	288
His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr Glu His Ser	
85 90 95	
TCT GCT GCC TGG GAC ACC ACC CTC CTG GAG CAG CTC CGC ACT GGA CTC	336
Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr Gly Leu	
100 105 110	

-continued

CAT CAG CAG CTG GAC AAC CTG GAT GCC TGC CTG GGG CAG GTG ATG GGA His Gln Gln Leu Asp Asn Leu Asp Ala Cys Leu Gly Gln Val Met Gly 115 120 125	384
GAG GAA GAC TCT GCC CTG GGA AGG ACG GGC CCC ACC CTG GCT CTG AAG Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala Leu Lys 130 135 140	432
AGG TAC TTC CAG GGC ATC CAT GTC TAC CTG AAA GAG AAG GGA TAC AGC Arg Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys Gly Tyr Ser 145 150 155 160	480
GAC TGC GCC TGG GAA ACC GTC AGA CTG GAA ATC ATG AGA TCC TTC TCT Asp Cys Ala Trp Glu Thr Val Arg Leu Glu Ile Met Arg Ser Phe Ser 165 170 175	528
TCA TTA ATC AGC TTG CAA GAA AGG TTA AGA ATG ATG GAT GGA GAC CTG Ser Leu Ile Ser Leu Gln Glu Arg Leu Arg Met Met Asp Gly Asp Leu 180 185 190	576
AGC TCA CCT TGA Ser Ser Pro 195	588

- (2) INFORMATION FOR SEQ ID NO:12:
- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 195 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence  
of SEQ ID NO:11 (HuIFNtau1).
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr 1 5 10 15
Gly Pro Gly Gly Ser Leu Gly Cys Asp Leu Ser Gln Asn His Val Leu 20 25 30
Val Gly Arg Lys Asn Leu Arg Leu Leu Asp Glu Met Arg Arg Leu Ser 35 40 45
Pro Arg Phe Cys Leu Gln Asp Arg Lys Asp Phe Ala Leu Pro Gln Glu 50 55 60
Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile Ser Val Leu 65 70 75 80
His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr Glu His Ser 85 90 95
Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr Gly Leu 100 105 110
His Gln Gln Leu Asp Asn Leu Asp Ala Cys Leu Gly Gln Val Met Gly 115 120 125
Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala Leu Lys 130 135 140
Arg Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys Gly Tyr Ser 145 150 155 160
Asp Cys Ala Trp Glu Thr Val Arg Leu Glu Ile Met Arg Ser Phe Ser 165 170 175
Ser Leu Ile Ser Leu Gln Glu Arg Leu Arg Met Met Asp Gly Asp Leu 180 185 190
Ser Ser Pro 195



- (2) INFORMATION FOR SEQ ID NO:13:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 25 bases
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (synthetic)
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: 25-mer synthetic oligonucleotide
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CCTGTCTGCA GGACAGAAAA GACTT 25

- (2) INFORMATION FOR SEQ ID NO:14:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 25 bases
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: single
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (synthetic)
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: 25-mer synthetic oligonucleotide
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

TCTGAATTCT GACGATTTC CAGGC 25

- (2) INFORMATION FOR SEQ ID NO:15:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 37 amino acids
    - (B) TYPE: amino acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: peptide
  - (iii) HYPOTHETICAL: NO
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 1-37 of SEQ ID NO:4
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Cys Asp Leu Ser Gln Asn His Val Leu Val Gly Arg Lys Asn Leu Arg  
1 5 10 15  
Leu Leu Asp Glu Met Arg Arg Leu Ser Pro Arg Phe Cys Leu Gln Asp  
20 25 30  
Arg Lys Asp Phe Ala  
35

- (2) INFORMATION FOR SEQ ID NO:16:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 31 amino acids
    - (B) TYPE: amino acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: peptide
  - (iii) HYPOTHETICAL: NO
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 34-64 of SEQ ID NO:4
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

-continued

Lys Asp Phe Ala Leu Pro Gln Glu Met Val Glu Gly Gly Gln Leu Gln  
1 5 10 15  
Glu Ala Gln Ala Ile Ser Val Leu His Glu Met Leu Gln Gln Ser  
20 25 30

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 31 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment  
62-92 of SEQ ID NO:4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Gln Gln Ser Phe Asn Leu Phe His Thr Glu His Ser Ser Ala Ala Trp  
1 5 10 15  
Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr Gly Leu His Gln Gln  
20 25 30

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment  
90-122 of SEQ ID NO:4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

His Gln Gln Leu Asp Asn Leu Asp Ala Cys Leu Gly Gln Val Met Gly  
1 5 10 15  
Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala Leu Lys  
20 25 30

Arg

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 32 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment  
119-150 of SEQ ID NO:4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Ala Leu Lys Arg Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys  
1 5 10 15  
Gly Tyr Ser Asp Cys Ala Trp Glu Thr Val Arg Leu Glu Ile Met Arg  
20 25 30



(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 34 amino acids  
    (B) TYPE: amino acid  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(iii) HYPOTHETICAL: NO

(vi) ORIGINAL SOURCE:  
    (C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment  
        139-172 of SEQ ID NO:4

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Cys	Ala	Trp	Glu	Thr	Val	Arg	Leu	Glu	Ile	Met	Arg	Ser	Phe	Ser	Ser
1			5					10					15		
Leu	Ile	Ser	Leu	Gln	Glu	Arg	Leu	Arg	Met	Met	Asp	Gly	Asp	Leu	Ser
		20					25					30			
Ser	Pro														

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:  
    (A) LENGTH: 299 base pairs  
    (B) TYPE: nucleic acid  
    (C) STRANDEDNESS: double  
    (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA to mRNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(vi) ORIGINAL SOURCE:  
    (C) INDIVIDUAL ISOLATE: HuIFNtau6

(ix) FEATURE:  
    (A) NAME/KEY: CDS  
    (B) LOCATION: 2..298

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

C	CAG	GAG	ATG	GTG	GAG	GGC	GGC	CAG	CTC	CAG	GAG	GCC	CAG	GCC	ATC	46
	Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	Gln	Glu	Ala	Gln	Ala	Ile	
1				5				10					15			
TCT	GTG	CTC	CAC	AAG	ATG	CTC	CAG	CAG	AGC	TTC	AAC	CTC	TTC	CAC	ACA	94
Ser	Val	Leu	His	Lys	Met	Leu	Gln	Gln	Ser	Phe	Asn	Leu	Phe	His	Thr	
			20				25					30				
GAG	CGC	TCC	TCT	GCT	GCC	TGG	GAC	ACC	ACC	CTC	CTG	GAG	CAG	CTC	CGC	142
Glu	Arg	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	Leu	Leu	Glu	Gln	Leu	Arg	
			35				40					45				
ACT	GGA	CTC	CAT	CAG	CAG	CTG	GAT	GAC	CTG	GAC	GCC	TGC	CTG	GGG	CAG	190
Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	Asp	Ala	Cys	Leu	Gly	Gln	
		50				55					60					
GTG	ACG	GGA	GAG	GAA	GAC	TCT	GCC	CTG	GGA	AGG	ACG	GGC	CCC	ACC	CTG	238
Val	Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	Arg	Thr	Gly	Pro	Thr	Leu	
	65					70				75						
GCC	GTG	AAG	AGC	TAC	TTC	CAG	GGC	ATC	CAT	ATC	TAC	CTG	CAA	GAG	AAG	286
Ala	Val	Lys	Ser	Tyr	Phe	Gln	Gly	Ile	His	Ile	Tyr	Leu	Gln	Glu	Lys	
80					85			90						95		
GGA	TAC	AGC	GAC	T												299
Gly	Tyr	Ser	Asp													

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 99 amino acids	
(B) TYPE: amino acid	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: protein	
(vi) ORIGINAL SOURCE:	
(C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence of SEQ ID NO:21 (HuIFNtau6).	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:	
Gln Glu Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile Ser	
1 5 10 15	
Val Leu His Lys Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr Glu	
20 25 30	
Arg Ser Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr	
35 40 45	
Gly Leu His Gln Gln Leu Asp Asp Leu Asp Ala Cys Leu Gly Gln Val	
50 55 60	
Thr Gly Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala	
65 70 75 80	
Val Lys Ser Tyr Phe Gln Gly Ile His Ile Tyr Leu Gln Glu Lys Gly	
85 90 95	
Tyr Ser Asp	
(2) INFORMATION FOR SEQ ID NO:23:	
(i) SEQUENCE CHARACTERISTICS:	
(A) LENGTH: 288 base pairs	
(B) TYPE: nucleic acid	
(C) STRANDEDNESS: double	
(D) TOPOLOGY: linear	
(ii) MOLECULE TYPE: cDNA to mRNA	
(iii) HYPOTHETICAL: NO	
(iv) ANTI-SENSE: NO	
(vi) ORIGINAL SOURCE:	
(C) INDIVIDUAL ISOLATE: HuIFNtau7	
(ix) FEATURE:	
(A) NAME/KEY: CDS	
(B) LOCATION: 2..286	
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:	
C CAG GAG ATG GTG GAG GTC AGC CAG TTC CAG GAG GCC CAG GCC ATT	46
Gln Glu Met Val Glu Val Ser Gln Phe Gln Glu Ala Gln Ala Ile	
1 5 10 15	
TCT GTG CTC CAT GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC CAC AAA	94
Ser Val Leu His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Lys	
20 25 30	
GAG CGC TCC TCT GCT GCC TGG GAC ACT ACC CTC CTG GAG CAG CTC CTC	142
Glu Arg Ser Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Leu	
35 40 45	
ACT GGA CTC CAT CAG CAG CTG GAT GAC CTG GAT GCC TGT CTG GGG CAG	190
Thr Gly Leu His Gln Gln Leu Asp Asp Leu Asp Ala Cys Leu Gly Gln	
50 55 60	
TTG ACT GGA GAG GAA GAC TCT GCC CTG GGA AGG ACG GGC CCC ACC CTG	238
Leu Thr Gly Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu	
65 70 75	
GCC GTG AAG AGC TAC TTC CAG GGC ATC CAT GTC TAC CTG CAA GAG AAG	286
Ala Val Lys Ser Tyr Phe Gln Gly Ile His Val Tyr Leu Gln Glu Lys	



-continued

80	85	90	95
GG			288

- (2) INFORMATION FOR SEQ ID NO:24:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 95 amino acids
    - (B) TYPE: amino acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: protein
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence of SEQ ID NO:23 (HuIFNtau7).
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

Gln	Glu	Met	Val	Glu	Val	Ser	Gln	Phe	Gln	Glu	Ala	Gln	Ala	Ile	Ser
1				5				10						15	
Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	Phe	Asn	Leu	Phe	His	Lys	Glu
			20					25					30		
Arg	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	Leu	Leu	Glu	Gln	Leu	Leu	Thr
			35				40					45			
Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	Asp	Ala	Cys	Leu	Gly	Gln	Leu
	50					55				60					
Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	Arg	Thr	Gly	Pro	Thr	Leu	Ala
65				70				75						80	
Val	Lys	Ser	Tyr	Phe	Gln	Gly	Ile	His	Val	Tyr	Leu	Gln	Glu	Lys	
				85				90						95	

- (2) INFORMATION FOR SEQ ID NO:25:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 307 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: cDNA to mRNA
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: HuIFNtau4
  - (ix) FEATURE:
    - (A) NAME/KEY: CDS
    - (B) LOCATION: 2..307
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

C	CAG	GAG	ATG	GTG	GAG	GGT	GGC	CAG	CTC	CAG	GAG	GCC	CAG	GCC	ATC	46
	Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	Gln	Glu	Ala	Gln	Ala	Ile	
1				5				10						15		
TCT	GTG	CTC	CAC	GAG	ATG	CTC	CAG	CAG	AGC	TTC	AAC	CTC	TTC	CAC	ACA	94
	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	Phe	Asn	Leu	Phe	His	Thr
				20				25						30		
GAG	CAC	TCC	TCT	GCT	GCC	TGG	GAC	ACC	ACC	CTC	CTG	GAG	CAG	CTC	CGC	142
	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	Leu	Leu	Glu	Gln	Leu	Arg
				35				40					45			
ACT	GGA	CTC	CAT	CAG	CAG	CTG	GAT	GAC	CTG	GAT	GCC	TGC	CTG	GGG	CAG	190
	Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	Asp	Ala	Cys	Leu	Gly	Gln
				50			55					60				
GTG	ACG	GGA	GAG	GAA	GAC	TCT	GCC	CTG	GGA	AGG	ACG	GGC	CCC	ACC	CTG	238
	Val	Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	Arg	Thr	Gly	Pro	Thr	Leu

-continued

65	70	75	
GCC ATG AAG ACG TAT TTC CAG GGC ATC CAT GTC TAC CTG AAA GAG AAG			286
Ala Met Lys Thr Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys			
80	85	90	95
GGA TAT AGT GAC TGC GCC TGG			307
Gly Tyr Ser Asp Cys Ala Trp			
100			

- (2) INFORMATION FOR SEQ ID NO:26:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 102 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
- (C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence of SEQ ID NO:25 (HuIFNtau4).
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

Gln Glu Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile Ser	
1 5 10 15	
Val Leu His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr Glu	
20 25 30	
His Ser Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr	
35 40 45	
Gly Leu His Gln Gln Leu Asp Asp Leu Asp Ala Cys Leu Gly Gln Val	
50 55 60	
Thr Gly Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala	
65 70 75 80	
Met Lys Thr Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys Gly	
85 90 95	
Tyr Ser Asp Cys Ala Trp	
100	

- (2) INFORMATION FOR SEQ ID NO:27:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 294 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA to mRNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
- (C) INDIVIDUAL ISOLATE: HuIFNtau5
- (ix) FEATURE:
- (A) NAME/KEY: CDS
  - (B) LOCATION: 2..292
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

C CAG GAG ATG GTG GAG GGT GGC CAG CTC CAG GAG GCC CAG GCC ATC	46
Gln Glu Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile	
1 5 10 15	
TCT GTG CTC CAC GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC CAC ACA	94
Ser Val Leu His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr	
20 25 30	
GAG CAC TCC TCT GCT GCC TGG GAC ACC ACC CTC CTG GAG CAG CTC CGC	142



-continued

Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	Leu	Leu	Glu	Gln	Leu	Arg	
			35					40					45			
ACT	GGA	CTC	CAT	CAG	CAG	CTG	GAT	GAC	CTG	GAT	GCC	TGC	CTG	GGG	CAG	190
Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	Asp	Ala	Cys	Leu	Gly	Gln	
		50					55					60				
GTG	ACG	GGA	GAG	GAA	GAC	TCT	GCC	CTG	GGA	AGG	ACG	GGC	CCC	ACC	CTG	238
Val	Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	Arg	Thr	Gly	Pro	Thr	Leu	
	65					70				75						
GCC	ATG	AAG	ACG	TAT	TTC	CAG	GGC	ATC	CAT	GTC	TAC	CTG	AAA	GAG	AAG	286
Ala	Met	Lys	Thr	Tyr	Phe	Gln	Gly	Ile	His	Val	Tyr	Leu	Lys	Glu	Lys	
	80				85			90						95		
GGA	TAT	AG														294
Gly	Tyr															

- (2) INFORMATION FOR SEQ ID NO:28:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 97 amino acids
    - (B) TYPE: amino acid
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: protein
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence of SEQ ID NO:27 (HuIFNtau5).
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	Gln	Glu	Ala	Gln	Ala	Ile	Ser	
1				5				10						15		
Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	Phe	Asn	Leu	Phe	His	Thr	Glu	
		20						25					30			
His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	Leu	Leu	Glu	Gln	Leu	Arg	Thr	
		35				40						45				
Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	Asp	Ala	Cys	Leu	Gly	Gln	Val	
	50					55					60					
Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	Arg	Thr	Gly	Pro	Thr	Leu	Ala	
	65				70				75						80	
Met	Lys	Thr	Tyr	Phe	Gln	Gly	Ile	His	Val	Tyr	Leu	Lys	Glu	Lys	Gly	
				85				90						95		
Tyr																

- (2) INFORMATION FOR SEQ ID NO:29:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 516 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (genomic)
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: HuIFNtau2
  - (ix) FEATURE:
    - (A) NAME/KEY: CDS
    - (B) LOCATION: 1..516
    - (A) NAME/KEY: Modified-site
    - (B) LOCATION: 115-117
    - (D) OTHER INFORMATION: /note= "to allow expression of the encoded protein this site can be modified to encode an amino acid, e.g., Gln"

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:																
GAC	CTG	TCT	CAG	AAC	CAC	GTG	CTG	GTT	GGC	AGG	AAG	AAC	CTC	AGG	CTC	48
Asp	Leu	Ser	Gln	Asn	His	Val	Leu	Val	Gly	Arg	Lys	Asn	Leu	Arg	Leu	
1				5					10					15		
CTG	GAC	CAA	ATG	AGG	AGA	CTC	TCC	CCT	CGC	TTT	TGT	CTG	CAG	GAC	AGA	96
Leu	Asp	Gln	Met	Arg	Arg	Leu	Ser	Pro	Arg	Phe	Cys	Leu	Gln	Asp	Arg	
			20					25					30			
AAA	GAC	TTC	GCT	TTA	CCC	TAG	GAA	ATG	GTG	GAG	GGC	GGC	CAG	CTC	CAG	144
Lys	Asp	Phe	Ala	Leu	Pro		Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	Gln	
		35					40					45				
GAG	GCC	CAG	GCC	ATC	TCT	GTG	CTC	CAT	GAG	ATG	CTC	CAG	CAG	AGC	TTC	192
Glu	Ala	Gln	Ala	Ile	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	Phe	
	50					55					60					
AAC	CTC	TTC	CAC	ACA	GAG	CAC	TCC	TCT	GCT	GCC	TGG	GAC	ACC	ACC	CTC	240
Asn	Leu	Phe	His	Thr	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	Leu	
65					70				75						80	
CTG	GAG	CAG	CTC	CGC	ACT	GGA	CTC	CAT	CAG	CAG	CTG	GAC	AAC	CTG	GAT	288
Leu	Glu	Gln	Leu	Arg	Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asn	Leu	Asp	
				85					90					95		
GCC	TGC	CTG	GGG	CAG	GTG	ATG	GGA	GAG	GAA	GAC	TCT	GCC	CTG	GGA	AGG	336
Ala	Cys	Leu	Gly	Gln	Val	Met	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	Arg	
			100					105					110			
ACG	GGC	CCC	ACC	CTG	GCT	CTG	AAG	AGG	TAC	TTC	CAG	GGC	ATC	CAT	GTC	384
Thr	Gly	Pro	Thr	Leu	Ala	Leu	Lys	Arg	Tyr	Phe	Gln	Gly	Ile	His	Val	
		115					120					125				
TAC	CTG	AAA	GAG	AAG	GGA	TAC	AGC	GAC	TGC	GCC	TGG	GAA	ACC	GTC	AGA	432
Tyr	Leu	Lys	Glu	Lys	Gly	Tyr	Ser	Asp	Cys	Ala	Trp	Glu	Thr	Val	Arg	
	130					135					140					
GTG	GAA	ATC	ATG	AGA	TCC	TTC	TCT	TCA	TTA	ATC	AGC	TTG	CAA	GAA	AGG	480
Val	Glu	Ile	Met	Arg	Ser	Phe	Ser	Ser	Leu	Ile	Ser	Leu	Gln	Glu	Arg	
145					150					155					160	
TTA	AGA	ATG	ATG	GAT	GGA	GAC	CTG	AGC	TCA	CCT	TGA					516
Leu	Arg	Met	Met	Asp	Gly	Asp	Leu	Ser	Ser	Pro						
				165					170							

- (2) INFORMATION FOR SEQ ID NO:30:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 171 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
- (C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence of SEQ ID NO:29 (HuIFNtau2).
- (ix) FEATURE:
- (A) NAME/KEY: Modified-site
  - (B) LOCATION: 39
  - (D) OTHER INFORMATION: /note= "where Xaa a selected amino acid, for example, Gln"

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:																
Asp	Leu	Ser	Gln	Asn	His	Val	Leu	Val	Gly	Arg	Lys	Asn	Leu	Arg	Leu	
1				5					10					15		
Leu	Asp	Gln	Met	Arg	Arg	Leu	Ser	Pro	Arg	Phe	Cys	Leu	Gln	Asp	Arg	
			20					25					30			
Lys	Asp	Phe	Ala	Leu	Pro	Xaa	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	Gln	
		35					40					45				
Glu	Ala	Gln	Ala	Ile	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	Phe	
	50					55					60					



-continued

Asn Leu Phe His Thr Glu His Ser Ser Ala Ala Trp Asp Thr Thr Leu  
65 70 75 80  
Leu Glu Gln Leu Arg Thr Gly Leu His Gln Gln Leu Asp Asn Leu Asp  
85 90 95  
Ala Cys Leu Gly Gln Val Met Gly Glu Glu Asp Ser Ala Leu Gly Arg  
100 105 110  
Thr Gly Pro Thr Leu Ala Leu Lys Arg Tyr Phe Gln Gly Ile His Val  
115 120 125  
Tyr Leu Lys Glu Lys Gly Tyr Ser Asp Cys Ala Trp Glu Thr Val Arg  
130 135 140  
Val Glu Ile Met Arg Ser Phe Ser Ser Leu Ile Ser Leu Gln Glu Arg  
145 150 155 160  
Leu Arg Met Met Asp Gly Asp Leu Ser Ser Pro  
165 170

- (2) INFORMATION FOR SEQ ID NO:31:
- (i) SEQUENCE CHARACTERISTICS:
    - (A) LENGTH: 588 base pairs
    - (B) TYPE: nucleic acid
    - (C) STRANDEDNESS: double
    - (D) TOPOLOGY: linear
  - (ii) MOLECULE TYPE: DNA (genomic)
  - (iii) HYPOTHETICAL: NO
  - (iv) ANTI-SENSE: NO
  - (vi) ORIGINAL SOURCE:
    - (C) INDIVIDUAL ISOLATE: HuIFNtau3
  - (ix) FEATURE:
    - (A) NAME/KEY: CDS
    - (B) LOCATION: 1..588
  - (xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

ATG GCC TTC GTG CTC TCT CTA CTC ATG GCC CTG GTG CTG GTC AGC TAC 48  
Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr  
1 5 10 15  
GGC CCG GGA GGA TCC CTG CGG TGT GAC CTG TCT CAG AAC CAC GTG CTG 96  
Gly Pro Gly Gly Ser Leu Arg Cys Asp Leu Ser Gln Asn His Val Leu  
20 25 30  
GTT GGC AGC CAG AAC CTC AGG CTC CTG GGC CAA ATG AGG AGA CTC TCC 144  
Val Gly Ser Gln Asn Leu Arg Leu Leu Gly Gln Met Arg Arg Leu Ser  
35 40 45  
CTT CGC TTC TGT CTG CAG GAC AGA AAA GAC TTC GCT TTC CCC CAG GAG 192  
Leu Arg Phe Cys Leu Gln Asp Arg Lys Asp Phe Ala Phe Pro Gln Glu  
50 55 60  
ATG GTG GAG GGT GGC CAG CTC CAG GAG GCC CAG GCC ATC TCT GTG CTC 240  
Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile Ser Val Leu  
65 70 75 80  
CAC GAG ATG CTC CAG CAG AGC TTC AAC CTC TTC CAC ACA GAG CAC TCC 288  
His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr Glu His Ser  
85 90 95  
TCT GCT GCC TGG GAC ACC ACC CTC CTG GAG CAG CTC CGC ACT GGA CTC 336  
Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr Gly Leu  
100 105 110  
CAT CAG CAG CTG GAT GAC CTG GAT GCC TGC CTG GGG CAG GTG ACG GGA 384  
His Gln Gln Leu Asp Asp Leu Asp Ala Cys Leu Gly Gln Val Thr Gly  
115 120 125  
GAG GAA GAC TCT GCC CTG GGA AGA ACG GGC CCC ACC CTG GCC ATG AAG 432  
Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala Met Lys

-continued

130	135	140	
AGG TAT TTC CAG GGC ATC CAT GTC TAC CTG AAA GAG AAG GGA TAT AGT			480
Arg Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys Gly Tyr Ser			
145	150	155	160
GAC TGC GCC TGG GAA ATT GTC AGA CTG GAA ATC ATG AGA TCC TTG TCT			528
Asp Cys Ala Trp Glu Ile Val Arg Leu Glu Ile Met Arg Ser Leu Ser			
	165	170	175
TCA TCA ACC AGC TTG CAC AAA AGG TTA AGA ATG ATG GAT GGA GAC CTG			576
Ser Ser Thr Ser Leu His Lys Arg Leu Arg Met Met Asp Gly Asp Leu			
	180	185	190
AGC TCA CCT TGA			588
Ser Ser Pro			
	195		

- (2) INFORMATION FOR SEQ ID NO:32:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 195 amino acids
- (B) TYPE: amino acid
- (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
- (C) INDIVIDUAL ISOLATE: predicted amino acid coding sequence of SEQ ID NO:31 (HuIFNtau3)
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr	
1	15
Gly Pro Gly Gly Ser Leu Arg Cys Asp Leu Ser Gln Asn His Val Leu	
20	30
Val Gly Ser Gln Asn Leu Arg Leu Leu Gly Gln Met Arg Arg Leu Ser	
35	45
Leu Arg Phe Cys Leu Gln Asp Arg Lys Asp Phe Ala Phe Pro Gln Glu	
50	60
Met Val Glu Gly Gly Gln Leu Gln Glu Ala Gln Ala Ile Ser Val Leu	
65	80
His Glu Met Leu Gln Gln Ser Phe Asn Leu Phe His Thr Glu His Ser	
85	95
Ser Ala Ala Trp Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr Gly Leu	
100	110
His Gln Gln Leu Asp Asp Leu Asp Ala Cys Leu Gly Gln Val Thr Gly	
115	125
Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala Met Lys	
130	140
Arg Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys Gly Tyr Ser	
145	160
Asp Cys Ala Trp Glu Ile Val Arg Leu Glu Ile Met Arg Ser Leu Ser	
165	175
Ser Ser Thr Ser Leu His Lys Arg Leu Arg Met Met Asp Gly Asp Leu	
180	190
Ser Ser Pro	
195	

- (2) INFORMATION FOR SEQ ID NO:33:
- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 518 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: double



-continued

(D) TOPOLOGY: linear																
(ii) MOLECULE TYPE: DNA (genomic)																
(iii) HYPOTHETICAL: NO																
(iv) ANTI-SENSE: NO																
(vi) ORIGINAL SOURCE:																
(C) INDIVIDUAL ISOLATE: HuIFNtau3, mature no leader sequence																
(ix) FEATURE:																
(A) NAME/KEY: CDS																
(B) LOCATION: 1..518																
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:																
TGT	GAC	CTG	TCT	CAG	AAC	CAC	GTG	CTG	GTT	GGC	AGC	CAG	AAC	CTC	AGG	48
Cys	Asp	Leu	Ser	Gln	Asn	His	Val	Leu	Val	Gly	Ser	Gln	Asn	Leu	Arg	
1				5				10						15		
CTC	CTG	GGC	CAA	ATG	AGG	AGA	CTC	TCC	CTT	CGC	TTC	TGT	CTG	CAG	GAC	96
Leu	Leu	Gly	Gln	Met	Arg	Arg	Leu	Ser	Leu	Arg	Phe	Cys	Leu	Gln	Asp	
			20					25					30			
AGA	AAA	GAC	TTC	GCT	TTC	CCC	CAG	GAG	ATG	GTG	GAG	GGT	GGC	CAG	CTC	144
Arg	Lys	Asp	Phe	Ala	Phe	Pro	Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	
		35					40					45				
CAG	GAG	GCC	CAG	GCC	ATC	TCT	GTG	CTC	CAC	GAG	ATG	CTC	CAG	CAG	AGC	192
Gln	Glu	Ala	Gln	Ala	Ile	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	
	50					55				60						
TTC	AAC	CTC	TTC	CAC	ACA	GAG	CAC	TCC	TCT	GCT	GCC	TGG	GAC	ACC	ACC	240
Phe	Asn	Leu	Phe	His	Thr	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	
65				70				75						80		
CTC	CTG	GAG	CAG	CTC	CGC	ACT	GGA	CTC	CAT	CAG	CAG	CTG	GAT	GAC	CTG	288
Leu	Leu	Glu	Gln	Leu	Arg	Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	
			85					90					95			
GAT	GCC	TGC	CTG	GGG	CAG	GTG	ACG	GGA	GAG	GAA	GAC	TCT	GCC	CTG	GGA	336
Asp	Ala	Cys	Leu	Gly	Gln	Val	Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	
		100						105					110			
AGA	ACG	GGC	CCC	ACC	CTG	GCC	ATG	AAG	AGG	TAT	TTC	CAG	GGC	ATC	CAT	384
Arg	Thr	Gly	Pro	Thr	Leu	Ala	Met	Lys	Arg	Tyr	Phe	Gln	Gly	Ile	His	
		115					120					125				
GTC	TAC	CTG	AAA	GAG	AAG	GGA	TAT	AGT	GAC	TGC	GCC	TGG	GAA	ATT	GTC	432
Val	Tyr	Leu	Lys	Glu	Lys	Gly	Tyr	Ser	Asp	Cys	Ala	Trp	Glu	Ile	Val	
	130					135				140						
AGA	CTG	GAA	ATC	ATG	AGA	TCC	TTG	TCT	TCA	TCA	ACC	AGC	TTG	CAC	AAA	480
Arg	Leu	Glu	Ile	Met	Arg	Ser	Leu	Ser	Ser	Ser	Thr	Ser	Leu	His	Lys	
145				150				155						160		
AGG	TTA	AGA	ATG	ATG	GAT	GGA	GAC	CTG	AGC	TCA	CCT	TG				518
Arg	Leu	Arg	Met	Met	Asp	Gly	Asp	Leu	Ser	Ser	Pro					
			165					170								
(2) INFORMATION FOR SEQ ID NO:34:																
(i) SEQUENCE CHARACTERISTICS:																
(A) LENGTH: 172 amino acids																
(B) TYPE: amino acid																
(D) TOPOLOGY: linear																
(ii) MOLECULE TYPE: protein																
(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:																
Cys	Asp	Leu	Ser	Gln	Asn	His	Val	Leu	Val	Gly	Ser	Gln	Asn	Leu	Arg	
1				5				10						15		
Leu	Leu	Gly	Gln	Met	Arg	Arg	Leu	Ser	Leu	Arg	Phe	Cys	Leu	Gln	Asp	
			20					25					30			
Arg	Lys	Asp	Phe	Ala	Phe	Pro	Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	

-continued

35					40					45						
Gln	Glu	Ala	Gln	Ala	Ile	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	
50					55					60						
Phe	Asn	Leu	Phe	His	Thr	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	
65					70					75					80	
Leu	Leu	Glu	Gln	Leu	Arg	Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asp	Leu	
					85					90					95	
Asp	Ala	Cys	Leu	Gly	Gln	Val	Thr	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	
					100					105					110	
Arg	Thr	Gly	Pro	Thr	Leu	Ala	Met	Lys	Arg	Tyr	Phe	Gln	Gly	Ile	His	
					115					120					125	
Val	Tyr	Leu	Lys	Glu	Lys	Gly	Tyr	Ser	Asp	Cys	Ala	Trp	Glu	Ile	Val	
					130					135					140	
Arg	Leu	Glu	Ile	Met	Arg	Ser	Leu	Ser	Ser	Ser	Thr	Ser	Leu	His	Lys	
145					150					155					160	
Arg	Leu	Arg	Met	Met	Asp	Gly	Asp	Leu	Ser	Ser	Pro					
					165					170						

- (2) INFORMATION FOR SEQ ID NO:35:
- (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 37 amino acids

(B) TYPE: amino acid

(D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:

(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 1-37 of SEQ ID NO:33
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

Cys	Asp	Leu	Ser	Gln	Asn	His	Val	Leu	Val	Gly	Ser	Gln	Asn	Leu	Arg																			
1		5			10					15																								
Leu	Leu	Gly	Gln	Met	Arg	Arg	Leu	Ser	Leu	Arg	Phe	Cys	Leu	Gln	Asp																			
				20				25				30																						
Arg	Lys	Asp	Phe	Ala																														



-continued

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 62-92  
of SEQ ID NO:33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

Gln Gln Ser Phe Asn Leu Phe His Thr Glu His Ser Ser Ala Ala Trp  
1 5 10 15

Asp Thr Thr Leu Leu Glu Gln Leu Arg Thr Gly Leu His Gln Gln  
20 25 30

(2) INFORMATION FOR SEQ ID NO:38:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 33 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 90-122  
of SEQ ID NO:33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

His Gln Gln Leu Asp Asp Leu Asp Ala Cys Leu Gly Gln Val Thr Gly  
1 5 10 15

Glu Glu Asp Ser Ala Leu Gly Arg Thr Gly Pro Thr Leu Ala Met Lys  
20 25 30

Arg

(2) INFORMATION FOR SEQ ID NO:39:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 32 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment  
119-150 of SEQ ID NO:33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

Ala Met Lys Arg Tyr Phe Gln Gly Ile His Val Tyr Leu Lys Glu Lys  
1 5 10 15

Gly Tyr Ser Asp Cys Ala Trp Glu Ile Val Arg Leu Glu Ile Met Arg  
20 25 30

(2) INFORMATION FOR SEQ ID NO:40:

(i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 34 amino acids  
(B) TYPE: amino acid  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(vi) ORIGINAL SOURCE:  
(C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment  
139-172 of SEQ ID NO:33

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

Cys Ala Trp Glu Ile Val Arg Leu Glu Ile Met Arg Ser Leu Ser Ser  
1 5 10 15

-continued

Ser Thr Ser Leu His Lys Arg Leu Arg Met Met Asp Gly Asp Leu Ser  
20 25 30

Ser Pro

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 1-23 of SEQ ID NO:32
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr  
1 5 10 15

Gly Pro Gly Gly Ser Leu Arg  
20

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 23 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: protein
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: Amino acid sequence of fragment 1-23 of SEQ ID NO:11
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

Met Ala Phe Val Leu Ser Leu Leu Met Ala Leu Val Leu Val Ser Tyr  
1 5 10 15

Gly Pro Gly Gly Ser Leu Gly  
20

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 519 base pairs
  - (B) TYPE: nucleic acid
  - (C) STRANDEDNESS: double
  - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: DNA (genomic)
- (iii) HYPOTHETICAL: NO
- (iv) ANTI-SENSE: NO
- (vi) ORIGINAL SOURCE:
  - (C) INDIVIDUAL ISOLATE: HuIFNtau1 genomic-derived DNA coding sequence, without leader seq.
- (ix) FEATURE:
  - (A) NAME/KEY: CDS
  - (B) LOCATION: 1..519
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

TGT GAC CTG TCT CAG AAC CAC GTG CTG GTT GGC AGG AAG AAC CTC AGG 48  
Cys Asp Leu Ser Gln Asn His Val Leu Val Gly Arg Lys Asn Leu Arg  
1 5 10 15

CTC CTG GAC GAA ATG AGG AGA CTC TCC CCT CGC TTT TGT CTG CAG GAC 96  
Leu Leu Asp Glu Met Arg Arg Leu Ser Pro Arg Phe Cys Leu Gln Asp



-continued

20				25				30								
AGA	AAA	GAC	TTC	GCT	TTA	CCC	CAG	GAA	ATG	GTG	GAG	GGC	GGC	CAG	CTC	144
Arg	Lys	Asp	Phe	Ala	Leu	Pro	Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu	
35				40				45								
CAG	GAG	GCC	CAG	GCC	ATC	TCT	GTG	CTC	CAT	GAG	ATG	CTC	CAG	CAG	AGC	192
Gln	Glu	Ala	Gln	Ala	Ile	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser	
50				55				60								
TTC	AAC	CTC	TTC	CAC	ACA	GAG	CAC	TCC	TCT	GCT	GCC	TGG	GAC	ACC	ACC	240
Phe	Asn	Leu	Phe	His	Thr	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr	
65				70				75				80				
CTC	CTG	GAG	CAG	CTC	CGC	ACT	GGA	CTC	CAT	CAG	CAG	CTG	GAC	AAC	CTG	288
Leu	Leu	Glu	Gln	Leu	Arg	Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asn	Leu	
85				90				95								
GAT	GCC	TGC	CTG	GGG	CAG	GTG	ATG	GGA	GAG	GAA	GAC	TCT	GCC	CTG	GGA	336
Asp	Ala	Cys	Leu	Gly	Gln	Val	Met	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly	
100				105				110								
AGG	ACG	GGC	CCC	ACC	CTG	GCT	CTG	AAG	AGG	TAC	TTC	CAG	GGC	ATC	CAT	384
Arg	Thr	Gly	Pro	Thr	Leu	Ala	Leu	Lys	Arg	Tyr	Phe	Gln	Gly	Ile	His	
115				120				125								
GTC	TAC	CTG	AAA	GAG	AAG	GGA	TAC	AGC	GAC	TGC	GCC	TGG	GAA	ACC	GTC	432
Val	Tyr	Leu	Lys	Glu	Lys	Gly	Tyr	Ser	Asp	Cys	Ala	Trp	Glu	Thr	Val	
130				135				140								
AGA	CTG	GAA	ATC	ATG	AGA	TCC	TTC	TCT	TCA	TTA	ATC	AGC	TTG	CAA	GAA	480
Arg	Leu	Glu	Ile	Met	Arg	Ser	Phe	Ser	Ser	Leu	Ile	Ser	Leu	Gln	Glu	
145				150				155				160				
AGG	TTA	AGA	ATG	ATG	GAT	GGA	GAC	CTG	AGC	TCA	CCT	TGA				519
Arg	Leu	Arg	Met	Met	Asp	Gly	Asp	Leu	Ser	Ser	Pro					
165				170												

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 172 amino acids
  - (B) TYPE: amino acid
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

Cys	Asp	Leu	Ser	Gln	Asn	His	Val	Leu	Val	Gly	Arg	Lys	Asn	Leu	Arg
1				5					10					15	
Leu	Leu	Asp	Glu	Met	Arg	Arg	Leu	Ser	Pro	Arg	Phe	Cys	Leu	Gln	Asp
20				25				30							
Arg	Lys	Asp	Phe	Ala	Leu	Pro	Gln	Glu	Met	Val	Glu	Gly	Gly	Gln	Leu
35				40				45							
Gln	Glu	Ala	Gln	Ala	Ile	Ser	Val	Leu	His	Glu	Met	Leu	Gln	Gln	Ser
50				55				60							
Phe	Asn	Leu	Phe	His	Thr	Glu	His	Ser	Ser	Ala	Ala	Trp	Asp	Thr	Thr
65				70				75				80			
Leu	Leu	Glu	Gln	Leu	Arg	Thr	Gly	Leu	His	Gln	Gln	Leu	Asp	Asn	Leu
85				90				95							
Asp	Ala	Cys	Leu	Gly	Gln	Val	Met	Gly	Glu	Glu	Asp	Ser	Ala	Leu	Gly
100				105				110							
Arg	Thr	Gly	Pro	Thr	Leu	Ala	Leu	Lys	Arg	Tyr	Phe	Gln	Gly	Ile	His
115				120				125							
Val	Tyr	Leu	Lys	Glu	Lys	Gly	Tyr	Ser	Asp	Cys	Ala	Trp	Glu	Thr	Val
130				135				140							
Arg	Leu	Glu	Ile	Met	Arg	Ser	Phe	Ser	Ser	Leu	Ile	Ser	Leu	Gln	Glu
145				150				155				160			

