



US005998169A

United States Patent [19]

[11] Patent Number: **5,998,169**

Klein et al.

[45] Date of Patent: ***Dec. 7, 1999**

[54] **MULTIMERIC HYBRID GENE ENCODING A CHIMERIC PROTEIN WHICH CONFERS PROTECTION AGAINST PARAINFLUENZA VIRUS AND RESPIRATORY SYNCYTIAL VIRUS**

[75] Inventors: **Michel H. Klein**, Willowdale; **Run-Pan Du**, Thornhill; **Mary E. Ewasyshyn**, Willowdale, all of Canada

[73] Assignee: **Connaught Laboratories Limited**, North York, Canada

[*] Notice: This patent is subject to a terminal disclaimer.

[21] Appl. No.: **08/838,189**

[22] Filed: **Apr. 16, 1997**

Related U.S. Application Data

[63] Continuation of application No. 08/001,554, Jan. 6, 1993.

[30] Foreign Application Priority Data

Jan. 6, 1992 [GB] United Kingdom 9200117

[51] **Int. Cl.⁶** **C12N 21/06**; C12N 1/20; C12N 15/00; C07N 21/04

[52] **U.S. Cl.** **435/69.3**; 435/252.3; 435/252.32; 435/253.1; 536/23.72; 935/27; 935/32; 935/72

[58] **Field of Search** 435/69.3, 525.3, 435/252.32, 253.1, 320.1, 325; 536/23.72; 935/27, 32, 72

[56] References Cited

U.S. PATENT DOCUMENTS

- 4,578,269 3/1986 Morein .
- 4,722,898 2/1988 Paoletti et al. .
- 4,866,034 9/1989 Ribic .
- 4,879,213 11/1989 Fox et al. .
- 4,950,480 8/1990 Barber et al. .
- 5,098,998 3/1992 Mekalonos et al. .
- 5,110,587 5/1992 Paoletti et al. .

FOREIGN PATENT DOCUMENTS

- 0421626 4/1991 European Pat. Off. .
- WO 89/05823 7/1989 WIPO .
- WO 89/10405 11/1989 WIPO .
- WO 90/03437 4/1990 WIPO .
- WO 93/21310 10/1993 WIPO .

OTHER PUBLICATIONS

Murphy et al; Experimental Lung Research, 2:97-109 (1981).

Porter et al; J. of Virology, Jan. 1991, p. 103-111, vol.65, No. 1.

Kapikan et al Am. J. Epidemiology 89, 1969, p. 405-421.
 Roy et al, J. Inf. Diseases. 157(4): 648-659 (1988).
 Stover et al, Nature, 351:456-460 (1991).
 Olmstead et al, PNAS, 83: 7462-7466 (1986).
 Hall et al, Science 265:1393-1394 (1994).
 Bushe et al J. Inf. Dis. 145, 1982, P. 311-319.
 Ray et al, (1989) Virus Research, 12:169-180.
 Coelingh et al, (1987) Virology, 160: 465-472.
 Wathen et al, (1989) J. of Inf. Dis. 159:255-263.
 Spriggs et al, (1987) J. Virol. 61:3416-3423.
 Stott et al, (1987) J. Virol. 61:3855-3861.
 Wathen et al, (1989) J. Gen Virol. 70:2625-2635.
 Brideau et al, (1989) J. Gen. Virol. 70:2637-2644.
 Goebel et al, (1990) Virology 179:247-266.
 Perkus et al (1990) Virology 179:276-286.
 Goebel et al (1990) Virology 179:517-563.
 Conners et al, (1992) Vaccine 10:475-484.
 Targaglia et al (1992) Virology 188:217-232.
 Piccini et al, (1987) Methods in Enzymology, 153:545-563.
 Taylor et al, (1990) J. Virology, 64:1441-1450.
 Collins, P., (1990) J. Virol, 64: 4007-4012.
 Vijaya S., et al, (1988) Mol. Cell Biol. vol. 8, No. 4; 1709-1714.
 Perkus et al (1989) J. Virology 63:3829-3836.
 Bowie et al, Science 247: 1306-1310 (1990).
 Kumar et al, PNAS 87:1337-1347 (1990).
 Lazar et al, Molecular and Cellular Biology 8(3):1247-1252 (1988).
 Burges et al, Journal of Cell Biology 111:2129-2138 (1990).
 Chanock et al; Pediatrics vol. 90, No. 1, Jul. 1992 pp. 137-143.
 Kasel et al; J. of Virology, Dec. 1984, pp. 828-832, vol. 52, No. 3.
 Prince et al; Am. J. of Pathology, vol. 93, #3, 771-783, 1978.
 Prince et al; J. of Virology 57; 721-728; Mar. 1986.
 Tizard, et al "An Introduction to Veterinary Immunology", 2nd ed, published by Saunders (Philadelphia) pp. 123-124).
 Hendry et al (J. Gen. Virol. 64(8) pp. 3833-3843) Aug. 1990.
 Coelingh et al (Journal of Virology vol. 63(9) pp. 3755-3760) Sep. 1989.

Primary Examiner—Anthony C. Caputa
Assistant Examiner—Mark Navarro
Attorney, Agent, or Firm—Sim & McBurney

[57] ABSTRACT

Multimeric hybrid genes encoding the corresponding chimeric protein comprise a gene sequence coding for an antigenic region of a protein from a first pathogen linked to a gene sequence coding for an antigenic region of a protein from a second pathogen. The pathogens particularly are parainfluenza virus (PIV) and respiratory syncytial virus (RSV). A single recombinant immunogen is capable of protecting infants and similar susceptible individuals against diseases caused by both PIV and RSV.

18 Claims, 39 Drawing Sheets

FIG.1A. NUCLEOTIDE SEQUENCE OF THE PIV-3 F GENE (PCR-AMPLIFIED)

AAGTCAATACCAACA ACTATTAGCAGTCA TACGTGCA AGAACA AGAAGA AGAGATTCAA
 TTCAGTTATGGTTGTTGATAAATCGTCAGTATGCACGTTCTTCTTCTCTAAGTT 60
 10 20 30 40 50

AAAGCTAAATAAGAGAAATCAAAACAAGGTA TAGAACCCGAAACAACA AATCAAAA
 TTTTCGATTTATTTCTCTTTAGTTTTCCTTCCATATCTTGTGGGCTTGTGTTT TAGTTTT 120
 70 80 90 100 110

CATCCAATCCATTTTAAACAATAATCCAAAGAGACC GGCAACAACA AGCACCACAAC
 GTAGGTTAGGTAATAATTTGTTTAAAGGTTTTCCTCTGGCCGTTGTGTTGTTTCGTGTTTG 180
 130 140 150 160 170

MET PRO THR [LEU] ILE LEU ILE ILE THR THR MET ILE MET ALA [SER] SER CYS GLN
 ACAATGCCCAACTTTAATACTGCTAATTATACAAACAATGATTA TGGCATCTTCCTGCCCAA
 TGTTACGGTTGAAATTAATGACGATTAATAATGTTACTAATA CCGTAGAAGGACGGTT 240
 190 200 210 220 230

ILE ASP ILE THR LYS LEU GLN HIS VAL GLY VAL LEU VAL ASN SER PRO LYS GLY MET LYS
 ATACATATCACAAACTACAGCATGTAGGTGATTTGGTCAACAGTCCCA AAGGGATGAAG
 TATGTATAGTGTTTTGTGATGTCGTACATCCACATAAC CAGTTGTCAGGTTTCCCTACTTC 300
 250 260 270 280 290

ILE SER GLN ASN PHE GLU THR ARG TYR ILE LEU SER LEU ILE PRO LYS ILE GLU ASP
 ATATCACAAACTTCGAAACAAGATATCTAATTTGAGCCCTCATACCA AATAAGAGAC
 TATAGTGTTTTGAAGCTTTTGTCTATAGATTAAACTCGGAGTATGG TTTTATCTTCTG 360
 310 320 330 340 350

SER ASN SER CYS GLY ASP GLN ILE LYS GLN TYR LYS ARG LEU LEU ASP ARG LEU ILE
 TCTAACTCTTGTGGTGACCACAGATCAACAATACAGAGGTTATTGG ATAGACTGATC
 AGATTGAGAACACCACCTGGTTGTTGTTATGTTCTCCAAATAAC CTTATCTGACTAG 420
 370 380 390 400 410

FIG. 1B. ILE PRO LEU TYR ASP GLY LEU ARG LEU GLN LYS ASP VAL ILE VAL **THR** ASN GLN GLU SER
 ATCCCTCTATATGATGGATTAGATTACAGAAAGATGTGATAGTAACCAATCAAGAAATCC
 TAGGGAGATATACTACCTAATTCTAATGCTTTCTACACTATCATTTGGTTAGTTCTTAGG 480
 430 440 450 460 470
 ASN GLU ASN THR ASP PRO ARG THR **ARG** **ARG** **SER** PHE GLY GLY VAL ILE GLY THR ILE ALA
 AATGAAACAACACTGATCCCAAGAACAAAGACGATCCCTTTGGAGGGGTAATTGGAAACCAATTGCT
 TTACTTTTGTGACTAGGGTCTTGTCTGCTAGGAAACCTCCCCCAATTAACCTTGGTAACGA 540
 490 500 510 520 530
 F2-F1 CLEAVAGE SITE
 LEU GLY VAL ALA THR SER ALA GLN ILE THR ALA ALA VAL ALA LEU VAL GLU ALA LYS GLN
 CTGGGAGTAGCAACCTCAGCACAAATTACAGCGGCAGTTGCTCTGCTGTTGAGAGCCAGCAG
 GACCCTCATCGTTGGAGTCTGTTTAAATGTCGCCCGTCAACGAGACCAACTTCGGTTCGTC 600
 550 560 570 580 590
 ALA **LYS** SER ASP ILE GLU LYS LEU LYS GLU ALA ILE ARG ASP THR ASN LYS ALA VAL GLN
 GCAAAATCACACATCGAAAAACTCAAGAAAGCAATCAGGGACACAAACAAGCAGTGCCAG
 CGTTTTAGTGTTAGCTTTTGTGAGTTTCTTCCCTTAGTCCCTGTGTTTGGTTTCGTCACCGTC 660
 610 620 630 640 650
 SER VAL GLN SER SER ILE GLY ASN LEU ILE VAL ALA ILE LYS SER VAL GLN ASP TYR VAL
 TCAGTTCAGAGCTCTATAGGAATAATTAAATAGTAAGCAATTAATCAGTCCCAAGATTATGTC
 AGTCAAGTCTCGAGATAATCCCTTTAAATTAATTCATCGTTAATTTAGTCAAGGTTCTAATACAG 720
 670 680 690 700 710
 ASN **ASN** GLU ILE VAL PRO SER ILE ALA ARG LEU GLY CYS GLU ALA ALA GLY LEU GLN LEU
 AACAAAGAAATCGTGCCATCGATTGCTAGACTAGGTTGTGAAAGCAGCAGGACTTCAATTA
 TTGTTGCTTTAGCACGGTAGCTAACGATCTGATCCCAACACTTCGTCCTGAAAGTTAAT 780
 730 740 750 760 770
 GLY ILE ALA LEU THR GLN HIS TYR SER GLU LEU THR ASN ILE PHE GLY ASP ASN ILE GLY
 GGAATTGCATTAAACAACAGCATTACTCAGAAATTAACAACATAATTTGGTGATAACATAGGA
 CCTTAACGTAATTGTGTGTAATGCTTAAATTTGTTTGTATAAACCACTATTGTATCCT 840
 790 800 810 820 830 840

FIG.1C.

SER LEU GLN GLU LYS GLY ILE LYS LEU GLN GLY ILE ALA SER LEU TYR ARG THR ASN ILE
 TCGTTACAAGAAAGGAATAAAATTACAAGGTATAGCATTCATTATACCGCACAAATATC
 AGCAATGTTCTTTTCCCTTATTTAATGTTCCCATATCGTAGTAATAATGGCGTGTTTATAG
 850 860 870 880 890 900

THR GLU ILE PHE THR SER THR VAL ASP LYS TYR ASP ILE TYR ASP LEU LEU PHE THR
 ACAGAAATATTCACAACATCAACAGTTGATAAATATGATATCTATGATCTATTATTACCA
 TGTCTTTATAAGTGTGTGTCAACTATTTATACCTATAGATACTAGATAATAAAATGT
 910 920 930 940 950 960

GLU SER ILE LYS VAL ARG VAL ILE ASP VAL ASP LEU ASN ASP TYR SER ILE THR LEU GLN
 GAATCAATAAAGGTGAGAGTTATAGATGTTGATTTGAAATGATTACTCAATCACCCCTCCAA
 CTTAGTTATTTCCACTCTCAATAATCTACAACCTAACTTACTAATGAGTTAGTGGGAGGTT
 970 980 990 1000 1010 1020

VAL ARG LEU PRO LEU LEU THR ARG LEU LEU ASN THR GLN ILE TYR LYS VAL ASP SER ILE
 GTCAGACTCCCTTTATTAACTAGGCTGCTGAACTCAGATCTACAAGTAGATTCCATA
 CAGTCTGAGGGAATAAATTGATCCGACGACTTGTGAGTCTAGATGTTTTCATCTAAGGTAT
 1030 1040 1050 1060 1070 1080

SER TYR ASN ILE GLN ASN ARG GLU TRP TYR ILE PRO LEU PRO SER HIS ILE MET THR LYS
 TCATATAATATCCAAACAGAGAAATGGTATATCCCTCTTCCCAGCCATATCATGACGAAA
 AGTATATATAGGTTTGTCTCTTACCATAATAGGGAGAAAGGTCGGTATAGTACTGCTTT
 1090 1100 1110 1120 1130 1140

GLY ALA PKE LEU GLY GLY ALA ASP VAL LYS GLU CYS ILE GLU ALA PHE SER SER TYR ILE
 GGGCATTTCTAGGTGGAGCAGATGTC AAGGAATGTATAGAAAGCATTCAGCAGTTATATA
 CCCCATAAAGATCCACCTCGTCTACAGTTCCCTTACATATCTTCGTAAAGTCGTCAAATAT
 1150 1160 1170 1180 1190 1200

CYS PRO SER ASP PRO GLY PHE VAL LEU ASN HIS GLU KET GLU SER CYS LEU SER GLY ASN
 TGCCCTTCTGATCCAGGATTTGTACTAAACCATGAAATGGAGAGCTGCTTATCAGGAAAC
 ACGGGAAAGACTAGGTCCCTAAACATGATTTGGTACTTTACCTCTCGACCGAATAGTCCCTTG
 1210 1220 1230 1240 1250 1260

FIG. 1D. ILE SER GLN CYS PRO ARG THR **THR** VAL **THR** SER ASP ILE VAL PRO ARG TYR ALA PHE VAL
 A T A T C C C A A T G T C C A A G A A C C A C G G T C A C A T C A G A C A T T G T T C C A A G A T A T G C A T T C G T C
 T A T A G G G T T A C A G G T T C T T G G T G C C A G T G T A G T C T G T A A C A A G G T T C T A T A C G T A A G C A G
 1270 1280 1290 1300 1310

 ASN GLY GLY VAL VAL ALA ASN CYS ILE THR THR THR CYS THR CYS ASN GLY ILE **ASP** ASN
 A A T G G A G G A G T G G T T G C A A A C T G T A T A A C A A C C A C C T G T A C A T G C A A C G G A A T C G A C A A T
 T T A C C T C C T C A C C A A C G T T T G A C A T A T T G T T G G T G G A C A T G T A C G T T G C C T T A G C T G T T A
 1330 1340 1350 1360 1370 1380

 ARG ILE ASN GLN PRO PRO ASP GLN GLY VAL LYS ILE THR HIS LYS GLU CYS ASN THR
 A G A A T C A A T C A A C C A C C T G A T C A G G A G T A A A A T T A T A C A C A T A A A G A A T G T A A T A C A
 T C T T A G T T A G T T G G T G G A C T A G T T C C T C A T T T T A A T A T T G T G T A T T T C T T A C A T T A T G T
 1390 1400 1410 1420 1430 1440

 ILE GLY ILE ASN GLY MET LEU PHE ASN THR ASN LYS GLU GLY THR LEU ALA PHE TYR THR
 A T A G G T A T C A A C G G A A T G C T G T T C A A T A C A A A T A A A G A A G G A A C T C T T G C A T T C T A C A C A
 T A T C C A T A G T T G C C T T A C G A C A A G T T A T G T T A T T C T T C C T T G A G A A C G T A A G A T G T G T
 1450 1460 1470 1480 1490 1500

 PRO ASN ASP ILE THR LEU ASN SER VAL ALA LEU ASP PRO ILE ASP ILE SER ILE GLU
 C C A A A T G A T A T A C A C T A A A T A A T T C T G T T G C A C T T G A T C C C A A T T G A C A T A T C A A T C G A G
 G G T T T A C T A T A T T G T G A T T T A T T A A G A C A A C G T G A A C T A G G T T A A C T G T A T A G T T A G C T C
 1510 1520 1530 1540 1550 1560

 LEU ASN LYS ALA LYS SER ASP LEU GLU SER LYS GLU TRP ILE ARG ARG SER ASN GLN
 C T T A A C A A A G C C A A A T C A G A T C T A G A A G A A T C A A A G A A T G G A T A A G A A G G T C A A A T C A A
 G A A T T G T T T C G G T T T A G T C T A G A T C T T C T T A G T T T T C T T A C C C T A T T C T T C C A G T T T A G T T
 1570 1580 1590 1600 1610 1620

 LYS LEU ASP SER ILE GLY ASN TRP HIS GLN SER SER THR THR ILE ILE ILE **ILE** LEU ILE
 A A A C T A G A T T C T A T T G G A A A C T G G C A T C A A T C T A G C A C T A C A A T C A T A A T T A T T T A A T A
 T T T G A T C T A A G A T A A C C T T T G A C C C G T A G T T A G A T C G T G A T G T T A G T A T T A A A A T T A T
 1630 1640 1650 1660 1670 1680

FIG.1E

MET ILE ILE ILE LEU PHE ILE ILE ASN VAL THR ILE ILE **THR** ILE ALA **ILE** LYS TYR TYR
 ATGATCATTATATTGTTTATAATTAAATGTAACGATAATTACAAATTGCAATTAAAGTATTAC
 TACTAGTAATAATAACAATAATTAAATGCTATTGCTATTAAATGTTAACGTTAATTCAATAATG 1740
 1690 1700 1710 1720 1730

ARG ILE GLN LYS ARG ASN ARG VAL ASP GLN ASN ASP LYS PRO TYR VAL LEU THR ASN LYS
 AGAATTCAAGAAGAAATCGAGTGGATCAAAATGACAAAGCCATTATGTACTAACAAACAAA
 TCTTAAGTTTCTCTTAGCTCACCTAGTTTACTGTTCGGTATACATGATGTTGTTT 1800
 1750 1760 1770 1780 1790

TGACATATCTATAGATCATTAGATATTAAATTTATAAATAACTT
 ACTGTATAGATATCTAGTAACTCTAATAATTTAATAATTTTTTGAA 1840
 1810 1820 1830 1840

NUCLEOTIDE SEQUENCE OF THE PIV-3 F GENE. THE cDNA SEQUENCE IS SHOWN IN THE PLUS (MRNA) STRAND SENSE IN THE 5' TO 3' DIRECTION. THE SIGNAL PEPTIDE (SP) AND THE TRANSMEMBRANE (TM) ANCHOR DOMAIN ARE UNDERLINED. THE PREDICTED F2-F1 CLEAVAGE SITE IS INDICATED BY THE ARROW (↓). AMINO ACIDS DIFFERING FROM THE PUBLISHED PRIMARY SEQUENCE OF THE PROTEIN ENCODED BY THE PIV-3 F GENE ARE BOXED.

RESTRICTION MAP OF THE PIV-3 F GENE




FIG.2.

NUCLEOTIDE SEQUENCE OF THE PIV-3 HN GENE.

FIG.3A.

5' AGACAATCCAAATTTCGGAGATGGAAATACTGGAGAGCATACCAATCAACGGAAAGGATGCTGG
 TCTGTTTAGGTTTAAAGCTCTAAGCTTTCGTTAGTTCCTACGACC 60
 10 20 30 40 50 60
 MET GLU TYR TRP LYS HIS THR ASN HIS GLY LYS ASP ALA GLY
 ASN GLU LEU GLU THR SER MET ALA THR [ASN] GLY ASN LYS [LEU] THR ASN LYS ILE THR TYR
 CAATGAGCTGGAGACGTCCTACTAATGGCAACACAGCTCACCAATAAGATAACATA
 GTTACTCGACCCTCTGCAGGTACCGATGATTAACCGTTTGCAGTGGTTATTCTATTGTAT 120
 70 80 90 100 110 120
 ILE LEU TRP THR ILE ILE LEU VAL LEU SER ILE VAL PHE ILE ILE VAL LEU ILE ASN
 TATATGACCAATAATCCTGGTGTATTATTCATAAGTCTTCATCATAGTGCCTAATTAA
 ATATAACCTGTTATTAGGACCAATAATAAGTTATTCAGAGTAGTATCACGATTAA 180
 130 140 150 160 170 180
 SER ILE LYS SER GLU LYS ALA HIS GLU SER LEU LEU GLN ASP [ILE] ASN ASN GLU PHE MET
 TTCATAAAGTGAAGAAGGCTCATGAAATCATTGCTGCAAGACATAAATAATGAGTTTAT
 AAGGTAGTTTTCACCTTTCCGAGTACTTAGTAAACGACCGTTCCTGTTATTACTCAATA 240
 190 200 210 220 230 240
 GLU [ILE] THR GLU LYS ILE GLN MET ALA SER ASP ASN [THR] ASN ASP LEU ILE GLN SER GLY
 GGAATTACAGAAAGATCCAAATGGCATCGGATAAATACCAATGATCTAATAACAGTCAGG
 CCTTAAATGTCCTTCTAGGTTTACCCTATTAAGGTTAGCCTATTACTAGATTATGTCAGTCC 300
 250 260 270 280 290 300
 VAL ASN THR ARG LEU LEU THR ILE GLN SER HIS VAL GLN ASN TYR ILE PRO ILE SER LEU
 AGTGAATACAGGCTTCTTACCAATTCAGAGTCAATGATATAATACCAATAATAGTACT
 TCACCTTATGTTCCGAAGAAATGTTAAGTCTCAGGTACAGGTTAATAATGTTATAGTGA 360
 310 320 330 340 350 360

THR GLN GLN MET SER ASP LEU ARG LYS PHE ILE SER GLU ILE THR ILE ARG ASN ASP ASN
 GACACACAGATGTCAGATCTTAGGAAATTCATTAGTGAATAATACAAATTAAGAAATGATAA
 CTGTGTTGTCACAGTCTAGAAATCCCTTAAGTAATCACCTTTAATGTTAATCTTTACTATT
 370 380 390 400 410 420

[GLN] GLU VAL [LEU] PRO GLN ARG ILE THR HIS ASP [VAL] GLY ILE LYS PRO LEU ASN PRO ASP
 TCAAGAAAGTGTGCTGCCACAAGAAATAACACATGATGTGGGTATAAACAACCTTTAAATCCAGA
 AGTTCTTCAACGACGGTGTCTTATTTGTGTACTACCCCATATTTTGGAAATTTAGGTCT
 430 440 450 460 470 480

ASP PHE TRP ARG CYS THR SER GLY LEU PRO SER LEU MET LYS THR PRO LYS ILE ARG LEU
 TGATTTTGGAGATGCCACGTCTGGTCTTCCATCTTAAATGAACAACCTCCAAAATAAGGTT
 ACTAAAACCCTCTACGTGCGAGACCCAGAAAGGTAGAAATTTACTTTTGGAGGTTTATTCCCAA
 490 500 510 520 530 540

MET PRO GLY PRO GLY LEU LEU ALA MET PRO THR THR VAL ASP GLY CYS [ILE] ARG THR PRO
 AATGCCAGGGCCGGGATTAATTAGCTATGCCCAACGACTGTTGATGGCTGTATCAGAACTCC
 TTACGGTCCCGGCCCTAAATAATCGATAACGGTTGCTGACAACTACCGACATAGTCTTGAGG
 550 560 570 580 590 600

SER LEU VAL ILE ASN ASP LEU ILE TYR ALA TYR THR SER ASN LEU ILE THR ARG GLY CYS
 GTCCCTTAGTTATAAATGATCTGATTTATGCTTATACCTCAAAATCTAATTACTCGAGGTTG
 CAGGAATCAATATTACTAGACTAAATAACGAAATATGGAGTTTAGATTAATGAGCTCCAAC
 610 620 630 640 650 660

GLN ASP ILE GLY LYS SER TYR GLN VAL LEU GLN ILE ILE THR VAL ASN SER ASP
 TCAGGATATAGGAATAATCATATCAAGTCTTACAGATAGGGATAATAACTGTAAACTCAGAA
 AGTCCATAATCCCTTTAGTATAGTTTCAGAAATGTCCTATCCCTATTATTGACATTTGAGTCT
 670 680 690 700 710 720

LEU VAL PRO ASP LEU ASN PRO ARG ILE SER HIS THR PHE ASN ILE ASN ASP ASN ARG LYS
 CTGGTACCCTGACTTAATAATCCAGGATCTCTCATCTTAAACATAAATAAGCAATAGGAA
 GAACCATGGACTGAATTTAGGGTCCCTAGAGGATGAAATTTGTAATTTACTGTTATCCTT
 730 740 750 760 770 780

FIG.3B.

SRE CYS SER LEU ALA LEU LEU ASN THR ASP VAL TYR GLN LEU CYS SER THR PRO LYS VAL
 G T C A T G T T C T A G C A C T C C T A A A T A C A G A T G T A T A T C A A C T G T G T T C A A C T C C C A A A G T
 C A G T A C A A G A G A T C G T G A T T A T G T C T A C A T A T A G T T G A C A C A A G T T G A G G T T T C A
 790 800 810 820 830 840

ASP GLU ARG SER ASP TYR ALA SER SER GLY ILE GLU ASP ILE VAL LEU ASP ILE VAL ASN
 T G A T G A A A G A T C A G A T T A T G C A T C A G G C A T A G A A G A T A T T G T A C T T G A T A T T G T C A A
 A C T A C T T T C T A G T C T A A T A C G T A G T A G T C C G T A T C T T C T A T A C A T G A A C T A T A A C A G T T
 850 860 870 880 890 900

TYR ASP GLY SER ILE SER THR THR ARG PHE LYS ASN ASN ASN ILE SER PHE ASP GLN PRO
 T T A T G A T G G C T C A A T C T C A A C A A C A A G A T T T A A G A T A A T A A C A T A A G C T T T G A T C A A C C
 A A T A C T A C C G A G T T A G A T T G T T C T A A A T T C T T A T T A T T G T A T T C G A A A C T A G T T G G
 910 920 930 940 950 960

TYR ALA ALA LEU TYR PRO SER VAL GLY PRO GLY ILE TYR TYR LYS GLY LYS ILE ILE PHE
 T T A T G C T G C A C T A T A C C C A T C T G T T G G A C C A G G G A T A T A C T A C A A A G G C A A A A T A A T A T T
 A A T A C G A C C G T G A T A T G G G T A G A C A A C C T G G T C C C T A T A T G A T G T T C C G T T T A T T A T A A
 970 980 990 1000 1010 1020

LEU GLY TYR GLY GLY LEU GLU HIS PRO ILE ASN GLU ASN **VAL** ILE CYS ASN THR THR GLY
 T C T C G G G T A T G G A G G T C T T G A A C A T C C A A T A A A T G A G A A T G T A A T C T G C A A C A C A A C T G G
 A G A G C C C A T A C C T C C A G A A C T T G T A G G T T A T T A C T C T T A C A T T A G A C G T T G T G T T G A C C
 1030 1040 1050 1060 1070 1080

CYS PRO GLY LYS THR GLN ARG ASP CYS ASN GLN ALA SER HIS SER PRO TRP PHE SER ASP
 G T G T C C C G G A A A C A C A G A G A C T G C A A T C A G G C A T C T C A T A G T C C A T G G T T T C A G A
 C A C A G G G C C C T T T G T G T C T C T G A C G T T A G T C C G T A G A G T A T C A G G T A C C A A A G T C T
 1090 1100 1110 1120 1130 1140

ARG ARG MET VAL ASN SER ILE ILE VAL VAL ASP LYS GLY LEU ASN SER ILE PRO LYS LEU
 T A G G A G G A T G G T C A A C T C T A T C A T T G T T G A C A A A G G C T T A A A C T C A A T T C C A A A A T T
 A T C C T C C T A C C A G T T G A G A T A G T A A C A C A C T G T T C C G A A T T T G A G T T A A G G T T T A A
 1150 1160 1170 1180 1190 1200

FIG.3C.

LYS VAL TRP THR ILE SER MET ARG GLN ASN TYR TRP GLY SER GLU GLY ARG LEU LEU LEU
 G A G G T A T G G A C G A T A T C T A T G A G A C A G A A T T A C T G G G G T C A G A A G G A A G G T T A C T T C T
 C T T C C A T A C C T G C T A T A G A T A C T C T G T C T T A A T G A C C C C C A G T C T T C C T T C C A A T G A A G A
 1210 1220 1230 1240 1250 1260

LEU GLY ASN LYS ILE TYR ILE TYR THR ARG SER THR SER TRP HIS SER LYS LEU GLN LEU
 A C T A G G T A A C A A G A T C T A T A T A T A C A A G A T C C A C A A G T T G G C A T A G C A A G T T A C A A T T
 T G A T C C A T T G T T C T A G A T A T A T A T G T T C T A G G T G T T C A A C C G T A T C G T T C A A T G T T A A
 1270 1280 1290 1300 1310 1320

GLY ILE ILE ASP ILE THR ASP TYR SER ASP ILE ARG ILE LYS TRP THR TRP HIS ASN VAL
 A G G A A T A A T T G A T A T T A C T G A T T A C A G T G A T A T A A G G A T A A A A T G G A C A T G G C A T A A T G T
 T C C T T A T T A A C T A T A A T G A C T A A T G T C A C T A T A T T C C T A T T T A C C T G T A C C G T A T T A C A
 1330 1340 1350 1360 1370 1380

LEU SER ARG PRO GLY ASN ASN GLU CYS PRO TRP GLY HIS SER CYS PRO ASP GLY CYS ILE
 G C T A T C A A G A C C A G G A A A C A A T G A A T G T C C A T G G G G A C A T T C A T G T C C A G A T G G A T G T A T
 C G A T A G T T C T G G T C C T T T G T T A C T T A C A G G T A C C C C T G T A A G T A C A G G T C T A C C T A C A T A
 1390 1400 1410 1420 1430 1440

THR GLY VAL TYR THR ASP ALA TYR PRO LEU ASN PRO THR GLY SER ILE VAL SER SER VAL
 A A C A G G A G T A T A T A C T G A T G C A T A T C C A C A T C C A C A G G G A G C A T T G T G T C A T C T G T
 T T G T C C T C A T A T A T G A C T A C G T A T A G G T T A G G G T G T C C C T C G T A A C A C A G T A G A C A
 1450 1460 1470 1480 1490 1500

ILE LEU ASP SER GLN LYS SER ARG VAL ASN PRO VAL ILE THR TYR SER THR [ALA] THR GLU
 C A T A T T A G A T T C A C A A A A T C G A G A G T G A A C C C A G T C A T A A C T T A C T C A C A G C A A C C G A
 G T A T A A T C T A A G T G T T T T A G C T C T C A C T T G G G T C A G T A T T G A A T G A G T T G T C G T T G G C T
 1510 1520 1530 1540 1550 1560

ARG VAL ASN GLU LEU ALA ILE [ARG] ASN ARG THR LEU SER ALA GLY TYR THR THR THR SER
 A A G A G T A A C G A G C T G G C C A T C C G A A A C A G A A C A C T C T C A G C T G G A T A T A C A A C A A C A A G
 T T C T C A T T T G C T C G A C C G G T A G G C T T T G T C T T G T G A G T C G A C C T A T A T G T T G T T G T T C
 1570 1580 1590 1600 1610 1620

FIG. 3D.

```

CYS ILE THR HIS TYR ASN LYS GLY TYR CYS PHE HIS ILE VAL GLU ILE ASN GLN LYS SER
CTGCATCACACACTATAACAAAGGATATTGTTTTCATATAGTAAATAAATCAGAAAG
GACGTAGTGATATTGTTTCCCTATAACAAGTATAATCACTTATTAGTCTT
1630 1640 1650 1660 1670 1680

LEU ASN THR LEU GLN PRO MET LEU PHE LYS THR GLU VAL PRO LYS SER CYS SER ***
CTTAACAACACTTCAACCCATGTTGTTCAAGACAGAGGTTCCAAAGCTGCAGTTAATC
GAAATTTGTGTGAAGTTGGGTACAAAGTTCCTCCAAAGGTTTTCGACGTCAAATTAG
1690 1700 1710 1720 1730 1740

ATAATTAAACCGCAATAATGCATTAAACCTATCTATAATACAAGTATATGATAAGTAAATCAGC
TATTAAATTGGCGTTATAACGTAATTGGATAGATATTATGTTTCATATACCTATTAGTCCG
1750 1760 1770 1780 1790 1800

ATCAGACAATAGACAACAAGGAAATAAATAA
TTAGTCTGTTATCTGTTTCCCTTATAATTTT
1810 1820 1830

```

NUCLEOTIDE SEQUENCE OF THE PIV-3 HN GENE. THE cDNA SEQUENCE IS SHOWN IN THE PLUS (MRNA) STRAND SENSE IN THE 5' TO 3' DIRECTION. THE TRANSMEMBRANE (TM) ANCHOR DOMAIN IS UNDERLINED. AMINO ACIDS DIFFERING FROM THE PUBLISHED PRIMARY SEQUENCE OF THE PROTEIN ENCODED BY THE PIV-3 HN GENE ARE BOXED.

FIG.3E.

RESTRICTION MAP OF THE PIV-3 HN GENE




FIG.4.

FIG. 5A. NUCLEOTIDE SEQUENCE OF THE RSV F GENE.

5' MET GLU LEU **PRO** ILE LEU LYS ALA ASN ALA ILE THR THR ILE LEU ALA **ALA** VAL THR PHE
 ATGGAGTTTGCCATAAGCAATGCAATACCAATCCCTCGCTGCAGTCAATTT
 TACCCTCAACGGTTAGGAGTTTACGTTAATGGTGTAGGAGCGACGTCAGTGTAAA 60
 10 20 30 40 50
 CYS PHE ALA **SER** SER GLN ASN ILE THR GLU PHE TYR THR CYS SER ALA VAL
 TGC TTTGCTTCTAGTCAAAACATCACTGAGAAATTTATCAATCAACATGCAGTGCAGTT
 ACCGAAACGGAAGATCAGTTTGTAGTGACTTCTTAAATAGTTAGTTGTCACGTCACAA 120
 70 80 90 100 110
 SER LYS GLY TYR LEU SER ALA LEU ARG THR GLY TRP TYR THR SER VAL ILE THR ILE GLU
 AGCAAGGC TATCTTAGTGCCTCTAAGAACTGGTTGGTATACTAGTGTATAACTATAGAA
 TCGTTTCCGATAGAAATCACCGAGATTCTTGACCACCAATGATCAATAATGATATCTT 180
 130 140 150 160 170
 LEU SER ASN ILE LYS GLU ASN LYS CYS ASN GLY THR ASP ALA LYS VAL LYS LEU **MET** LYS
 TTAGTAAATATCAGGAATAAGTGTAATGGAAACAGATGCTAAGGTAAATGATGAA
 AATTCAATTAAGTTCCCTTTATTCACATTAACCAATTCCTACGATTTTAACTACT 240
 190 200 210 220 230
 GLN GLU LEU ASP LYS TYR LYS ASN ALA VAL THR GLU LEU MET GLN SER THR
 CAGAAATTAGATAAATAATAAATGCTGTAAACAGAAATGCAAGTTGCTCATGCAAGCAC
 GTTCTTAAATCTAATTAATTTTACGACATTTGCTTAAACGTCACGAGTACGTTTCGTGT 300
 250 260 270 280 290 300
 PRO **ALA** **ALA** ASN ASN ARG ALA ARG ARG GLU LEU PRO ARG PHE MET ASN TYR THR LEU ASN
 CCAAGCAACAATCGAGCCAGAGAACTACCAAGGTTTATGAAATATACACTCAAC
 GGTCCGTCGT TTAGCTTAGCTCCGGTCTTCTGATGGTTCCTTAAATAATGAGTTG 360
 310 320 330 340 350 360

ASN **THR** LYS LYS THR ASN VAL THR LEU SER LYS LYS ARG LYS ARG ARG PHE LEU GLY PHE
 AAT ACCCAA AATGTAACATTAAGCAAGAAAGAAAGAAAGATTTCTTGTTT
 TATGGTTT TTTGGTTACATTTGTAATTCCTTTCTTAAAGAACCAA
 370 380 390 400 410 420

F2-F1 CLEAVAGE SITE

LEU LEU GLY VAL GLY SER ALA ILE ALA SER GLY **ILE** ALA VAL SER LYS VAL LEU HIS LEU
 TTGTTAGGTGTTGGATCTGCAATCGCCAGTGGCATTGCTGTATCTAAGGTCCTGCACCTTA
 AACAAATCCACAACCTAGACGTTAGCGGTCAACCGTAAACGACATAGATTCCAGGACGTGAAT
 430 440 450 460 470 480

GLU GLY GLU VAL ASN LYS ILE LYS SER ALA LEU LEU SER THR ASN LYS ALA VAL VAL SER
 GAAGGAGAAAGTGAACAAGATCAAAAGTCTACTATCCACAACAAGCCGTCAGTCAAGT
 CTTCCCTTCCACTTGTTC TAGTTTTCACCGAGATGATAGGTTTGTTCCTCCGGCATCAGTCA
 490 500 510 520 530 540

LEU SER ASN GLY VAL SER VAL LEU THR SER LYS VAL LEU ASP LEU LYS ASN TYR ILE ASP
 TTATCAATGGAGTTAGTGTTTAAACCAGCAAGTGTAGACCTCAAAACTATATAGAT
 AATAGTTTACCCTCAATCACAGAAATTTGGTCTGAGTTTGTGATATATCTA
 550 560 570 580 590 600

LYS GLN LEU LEU PRO ILE VAL ASN LYS ARG SER CYS **ARG** ILE SER ASN ILE GLU THR VAL
 AACAAATTTGTTACCCTATTGTGAAATAAGCGAAGCTGCAAGAAATATAGAAACTGTG
 TTTGTTAACAAATGGATAACACTTATTCTGCTTCGACGTTTATAGTTTATATCTTTGACAC
 610 620 630 640 650 660

ILE GLU PHE GLN HIS LYS ASN ASN ARG LEU LEU GLU ILE THR ARG GLU SER VAL ASN
 ATAGAGTTCCAAACAAGAAACAACAGACTACTAGAGATTACCAGGGAATTAGTGTAAAT
 TATCTCAAGGTTGTGTTCTTGTGTTGATCTCTAATGGTCCCTTAAATCACAAATTA
 670 680 690 700 710 720

ALA GLY VAL THR THR PRO VAL SER THR TYR MET LEU THR ASN SER GLU LEU SER LEU
 GCAGGTGTAACCTACACCTGTAAAGCACTTACATGTTAATAAGTGAATTAATGTCATTA
 CGTCCACATTTGATGTGACATTCGTGAATGATTAATTAACAGTAAT
 730 740 750 760 770 780

FIG.5B.

```

ILE ASN ASP MET PRO ILE THR ASN ASP GLN LYS LYS LEU MET SER ASN ASN VAL GLN ILE
ATCAATGATATGCCCTATAACAATAATGATCAGAAAGTTAATGTCACAACAATGTTCAAAATA
TAGTTACTATACGGATATTGTTTACTAGTCTTTCATTAACAGGTTTACAGTTTACAGTTTAT
790 800 810 820 830
VAL ARG GLN GLN SER TYR SER ILE MET SER ILE ILE LYS GLU VAL LEU ALA TYR VAL
GTTAGACAGCAAGTTTACTCTATCATGTCCTAATAAAGAGGAGTCTTAGCATATGTATA
CAATCTGTCGTTTCAATGAGATAGTACAGGTAATTAATTTCTCCTTCAGAAATCGTATACAT
850 860 870 880 890 900
VAL GLN LEU PRO LEU TYR GLY VAL ILE ASP THR PRO CYS TRP LYS LEU HIS THR SER PRO
GTACAATTACCACCTAATATGGTGATAGATACACCTTGTGGAAATTAACAACATCCCT
CATGTTAATGGTGATATACCACCTATCTATGTGGACACAACCTTAAATGTTGTAGTGGGA
910 920 930 940 950 960
LEU CYS THR THR ASN THR LYS GLU GLY SER ASN ILE CYS LEU THR ARG THR ASP ARG GLY
CTATGTACAACCAACAACAAGAGGAGGTCACAACATCTGTTTACAACAAGACTGACAGAGGA
GATACATGTTGGTTGTGTTTCTTCCCAAGTGTGTAGACAACAATGTTCTTGACTGCTCTCCCT
970 980 990 1000 1010 1020
TRP TYR CYS ASP ASN ALA GLY SER VAL SER PHE PHE PRO GLN ALA GLU THR CYS LYS VAL
TGGTACTGTGACAATGCAAGGATCAAGTATCTTCTTCCACAAGCTGAACAATGTAAGTT
ACCATGACACTGTTACGTTCCCTAGTTCATAGACATAGAGAGGTTGTTCCGACTTGTACATTTCAA
1030 1040 1050 1060 1070 1080
GLN SER ASN ARG VAL PHE CYS ASP THR MET ASN SER LEU THR LEU PRO SER GLU VAL ASN
CAATCGAATCGAGTATTTTGTGACACAATGACAGTTTAAACATTAACAAGTGAAGTAAAT
GTTAGCTTAGCTCATAAACAACACTGTGTACTTGTCAAATTTGTAATGGTTTCACTTCAATTA
1090 1100 1110 1120 1130 1140
LEU CYS ASN VAL ASP ILE PHE ASN PRO LYS TYR ASP CYS LYS ILE MET THR SER LYS THR
CTCTGC AATGTTGACATAATTC AATCCCA AATGATGTA A AATTA TGA CTTC A A AACA
GAGACGTTACAACCTGTAT AAGTTAGGGTTTATACTAACAATTA AACTGAAGTTT 1150 1160 1170 1180 1190 1200

```

FIG. 5C.

ASP VAL SER SER SER VAL ILE THR SER LEU GLY ALA ILE VAL SER CYS TYR GLY LYS THR
 GATGTAAGCAGCTCCGTTATCACATCTCTAGGAGCCATTGTGTCAATGCTATGGCAAAACT
 CTACATTCGTCGAGGCAATAGTGTAGTCCCTCGGTAAACACAGTACGATACCGTTT 1260

LYS CYS THR ALA SER ASN LYS ASN ARG GLY ILE ILE LYS THR PHE SER ASN GLY CYS ASP
 AAATGTACAGCATCCCAATAAATCGTGGAAATCATAAAGACAATTTCTTAACGGGTGTGAT
 TTTACAATGTCGTAGGTTATTTTAGCACCTTAGTATTTCTGTAAAGATTGCCCACTA 1320

TYR VAL SER ASN LYS GLY VAL THR VAL SER VAL GLY ASN THR LEU TYR TYR VAL ASN
 TATGTATCAATAAAGGGGTGACACTGTGTCAGGTAAACACACATTAATAATGTAAT
 ATACATAGTTTATTTCCCACTGAGGTTGACACAGACATCCCAATGTTAATAATACAATTA 1380

LYS GLN GLU GLY LYS SER LEU TYR VAL LYS GLY GLU PRO ILE ILE ASN PHE TYR ASP PRO
 AAGCAAGAGGCAAAAGTCTCTATGTAAAGGTGAACCAATAATAAATTTCTATGACCCA
 TTCGTTCTTCCGTTTTCAGAGATACATTTTCCCACTTGGTTAATAAAGATACTGGGT 1440

LEU VAL PHE PRO SER ASP GLU PHE ASP ALA SER ILE SER GLN VAL ASN GLU LYS ILE ASN
 TTAGTATTCCCTCTGTGATGAAATTTGATGCATCAATAATCAAGTCAACGAGAGATTAAAC
 AATCATAAAGGGGAGACTACTTAACCTAACCTAGTTAATAGAGTTCAGTTGCTTCTAAT 1500

GLN SER LEU ALA PHE ILE ARG LYS SER ASP GLU LEU LEU HIS ASN VAL ASN ALA GLY LYS
 CAGAGTTTAGCATTATTCTGATGAAATTCAGTAAATCAATAATGTAATAATGCTGTTAA
 GTCACAATCGTAAATAAGCAATTTAGGCTACTTAATAATGTAATAATTAACGACCAAT 1560

SER THR THR ASN ILE MET ILE THR THR ILE ILE ILE GLU ILE VAL ILE LEU LEU SER
 TCAACCAATAATCATGATAACTACTATAATAGAGATTATAGTAATAATGTTATCA
 AGTTGGTGTATTAAGTACTATAATGATGATAATCAATTAACAATAGT 1620

FIG. 5D.

TM

LEU ILE ALA VAL GLY LEU LEU LEU TYR CYS LYS ALA ARG SER THR PRO VAL THR LEU SER
 TTAATTGCTGTGGACTGCTCCTATACCTGTAGGCAAGCAGCACACCAAGTCCACACTAAGC
 AATTAACGACAACCTGACCGAGGATATGACATTCCTGTTCCGGTCTTCCGGTCAAGTGTGATTCG
 1630 1640 1650 1660 1670 1680

LYS, ASP GLN LEU SER GLY ILE ASN ASN ILE ALA PHE SER ASN
 AAGGATCAACTGAGTGGTATAATAATAATTGCATTTAGTAACCTGAATAAATAAGCACCT
 TTCCTAGTTGACTCACCAATATTATTATAAACGTAATACTTGAATTAATTTTATTCGTGGA
 1690 1700 1710 1720 1730 1740

AATCATGTTCTTACAATGGTTTACTATCTGCTCATAGACAACCCATCTATCATTTGGATTT
 TTAGTACAAGAATGTTACCAATGATAGACGAGTATCTGTTGGGTAGATAGTAACCTAAA
 1750 1760 1770 1780 1790 1800

TCTTAAATCTGAACCTTCAATCGAAACTCTTATCTATAAACCATCTCACCTTACACTATTTA
 AGAATTTAGACTTGAAGTAGCTTTGAGAA TAGATATTGTTAGAGTGAATGTGATTAAT
 1810 1820 1830 1840 1850 1860

AGTAGATTCCCTAGTTTATAGTTATAT 3'
 TCATCTAAGGATCAATAATCAATAA
 1870 1880

NUCLEOTIDE SEQUENCE OF THE RSV F GENE. THE CDNA SEQUENCE IS SHOWN IN THE PLUS (MRNA)
 STRAND SENSE IN THE 5' TO 3' DIRECTION. THE SIGNAL PEPTIDE (SP) AND THE TRANSMEMBRANE (TM)
 ANCHOR DOMAIN ARE UNDERLINED. THE PREDICTED F2-F1 CLEAVAGE SITE IS INDICATED BY THE ARROW
 (↓). AMINO ACIDS DIFFERING FROM THE PUBLISHED PRIMARY SEQUENCE OF THE PROTEIN ENCODED BY
 THE RSV F GENE ARE BOXED.

FIG. 5E.

RESTRICTION MAP OF THE RSV F GENE




FIG.6.

FIG. 7A. NUCLEOTIDE SEQUENCE OF THE RSV G GENE

MET SER LYS ASN LYS ASP GLN ARG
 T G C A A A C A T G T C C A A A A A C A A G G A C C A A C G
 A C G T T T G T A C A G G T T T T T G T T C C T G G T T G C
 10 20 30
 THR ALA LYS THR LEU GLU **LYS** THR TRP ASP
 C A C C G C T A A G A C A C T A G A A A A G A C C T G G G A
 G T G G C G A T T C T G T G A T C T T T T C T G G A C C C T
 40 50 60
 THR LEU ASN HIS LEU LEU PHE ILE SER SER
 C A C T C T C A A T C A T T T A T T A T T C A T A T C A T C
 G T G A G A G T T A G T A A A T A A T A A G T A T A G T A G
 70 80 90
GLY LEU TYR LYS LEU ASN LEU LYS SER VAL
 G G G C T T A T A T A A G T T A A A T C T T A A A T C T G T
 C C C G A A T A T A T T C A A T T T A G A A T T T A G A C A
 100 110 120
 _____ TM _____
 ALA GLN ILE THR LEU SER ILE LEU ALA MET
 A G C A C A A A T C A C A T T A T C C A T T C T G G C A A T
 T C G T G T T T A G T G T A A T A G G T A A G A C C G T T A
 130 140 150

 ILE ILE SER THR SER LEU ILE ILE **THR** ALA
 G A T A A T C T C A A C T T C A C T T A T A A T T A C A G C
 C T A T T A G A G T T G A A G T G A A T A T T A A T G T C G
 160 170 180

 ILE ILE PHE ILE ALA SER ALA ASN HIS LYS
 C A T C A T A T T C A T A G C C T C G G C A A A C C A C A A
 G T A G T A T A A G T A T C G G A G C C G T T T G G T G T T
 190 200 210
 VAL THR **LEU** THR THR ALA ILE ILE GLN ASP
 A G T C A C A C T A A C A A C T G C A A T C A T A C A A G A
 T C A G T G T G A T T G T T G A C G T T A G T A T G T T C T
 220 230 240
 ALA THR SER GLN ILE LYS ASN THR THR PRO
 T G C A A C A A G C C A G A T C A A G A A C A C A A C C C C
 A C G T T G T T C G G T C T A G T T C T T G T G T T G G G G
 250 260 270
 THR TYR LEU THR GLN **ASP** PRO GLN LEU GLY
 A A C A T A C C T C A C T C A G G A T C C T C A G C T T G G
 T T G T A T G G A G T G A G T C C T A G G A G T C G A A C C
 280 290 300

FIG.7B.

ILE SER **PHE** SER ASN **LEU** SER GLU ILE THR
 A A T C A G C T T C T C C A A T C T G T C T G A A A T T A C
 T T A G T C G A A G A G G T T A G A C A G A C T T T A A T G
 310 320 330

SER GLN **THR** THR THR ILE LEU ALA SER THR
 A T C A C A A A C C A C C A C C A T A C T A G C T T C A A C
 T A G T G T T T G G T G G T G G T A T G A T C G A A G T T G
 340 350 360

THR PRO GLY VAL LYS SER **ASN** LEU GLN **PRO**
 A A C A C C A G G A G T C A A G T C A A A C C T G C A A C C
 T T G T G G T C C T C A G T T C A G T T T G G A C G T T G G
 370 380 390

THR THR VAL LYS THR LYS ASN THR THR THR
 C A C A A C A G T C A A G A C T A A A A A C A C A A C A A C
 G T G T T G T C A G T T C T G A T T T T T G T G T T G T T G
 400 410 420

THR GLN THR GLN PRO SER LYS PRO THR THR
 A A C C C A A A C A C A A C C C A G C A A G C C C A C T A C
 T T G G G T T T G T G T T G G G T C G T T C G G G T G A T G
 430 440 450

LYS GLN ARG GLN ASN LYS PRO PRO **ASN** LYS
 A A A A C A A C G C C A A A A C A A A C C A C C A A A C A A
 T T T T G T T G C G G T T T T G T T T G G T G G T T T G T T
 460 470 480

PRO ASN ASN ASP PHE HIS PHE GLU VAL PHE
 A C C C A A T A A T G A T T T T C A C T T C G A A G T G T T
 T G G G T T A T T A C T A A A A G T G A A G C T T C A C A A
 490 500 510

ASN PHE VAL PRO CYS SER ILE CYS SER ASN
 T A A C T T T G T A C C C T G C A G C A T A T G C A G C A A
 A T T G A A A C A T G G G A C G T C G T A T A C G T C G T T
 520 530 540

ASN PRO THR CYS TRP ALA ILE CYS LYS ARG
 C A A T C C A A C C T G C T G G G C T A T C T G C A A A A G
 G T T A G G T T G G A C G A C C C G A T A G A C G T T T T C
 550 560 570

ILE PRO ASN LYS LYS PRO GLY LYS LYS THR
 A A T A C C A A A C A A A A A A C C A G G A A A G A A A A C
 T T A T G G T T T G T T T T T G G T C C T T T C T T T T G
 580 590 600

FIG.7C.

THR THR LYS PRO THR LYS LYS PRO THR PHE
 C A C C A C C A A G C C T A C A A A A A A A C C A A C C T T
 G T G G T G G T T C G G A T G T T T T T T T G G T T G G A A
 610 620 630

LYS THR THR LYS LYS ASP **LEU** LYS PRO GLN
 C A A G A C A A C C A A A A A A G A T C T C A A A C C T C A
 G T T C T G T T G G T T T T T T C T A G A G T T T G G A G T
 640 650 660

THR THR LYS **PRO** LYS GLU VAL PRO THR THR
 A A C C A C T A A A C C A A A G G A A G T A C C C A C C A C
 T T G G T G A T T T G G T T T C C T T C A T G G G T G G T G
 670 680 690

LYS PRO THR GLU GLU PRO THR ILE ASN THR
 C A A G C C C A C A G A A G A G C C A A C C A T C A A C A C
 G T T C G G G T G T C T T C T C G G T T G G T A G T T G T G
 700 710 720

THR LYS THR ASN ILE **THR** THR THR LEU LEU
 C A C C A A A A C A A A C A T C A C A A C T A C A C T G C T
 G T G G T T T T G T T T G T A G T G T T G A T G T G A C G A
 730 740 750

THR **ASN** ASN THR THR GLY ASN PRO **LYS** LEU
 C A C C A A C A A C A C C A C A G G A A A T C C A A A A C T
 G T G G T T G T T G T G G T G T C C T T T A G G T T T T G A
 760 770 780

THR SER GLN MET GLU THR PHE HIS SER THR
 C A C A A G T C A A A T G G A A A C C T T C C A C T C A A C
 G T G T T C A G T T T A C C T T T G G A A G G T G A G T T G
 790 800 810

SER SER GLU GLY ASN **LEU** SER PRO SER GLN
 C T C C T C C G A A G G C A A T C T A A G C C C T T C T C A
 G A G G A G G C T T C C G T T A G A T T C G G G A A G A G T
 820 830 840

VAL SER THR THR SER GLU **HIS** PRO SER GLN
 A G T C T C C A C A A C A T C C G A G C A C C C A T C A C A
 T C A G A G G T G T T G T A G G C T C G T G G G T A G T G T
 850 860 870

PRO SER SER PRO PRO ASN THR **THR** ARG GLN
 A C C C T C A T C T C C A C C C A A C A C A A C A C G C C A
 T G G G A G T A G A G G T G G G T T G T G T T G T G C G G T
 880 890 900

G T A G T T A T T A A A A A A A A A A A
C A T C A A T A A T T T T T T T T T T T
 910 920

NUCLEOTIDE SEQUENCE OF THE RSV G GENE. THE cDNA SEQUENCE IS SHOWN IN THE PLUS (mRNA) STRAND SENSE IN THE 5' TO 3' DIRECTION. THE TRANSMEMBRANE (TM) ANCHOR DOMAIN IS UNDERLINED. AMINO ACIDS DIFFERING FROM THE PUBLISHED PRIMARY SEQUENCE OF THE PROTEIN ENCODED BY THE RSV G GENE ARE BOXED.

FIG.7D.

RESTRICTION MAP OF RSV G GENE




FIG.8.

Construction of a Bluescript-based expression vector containing the chimeric F_{PIV-3}-F_{RSV} gene with the 5' untranslated region of the PIV-3 F gene intact but lacking the nucleotide sequences coding for the hydrophobic anchor domains and cytoplasmic tails of both the PIV-3 and RSV F genes.

Step 1: Preparation of the plasmid containing the modified PIV-3 F gene






FIG.9A.








FIG.9D.

Construction of a Bluescript-based expression vector containing the PIV-3 F gene lacking the 5' untranslated sequence and transmembrane anchor and cytoplasmic tail coding regions.





FIG.10A.

FIG.10B.



Cut with EcoRI and BstBI

Retrieve: EcoRI-BstBI restricted vector


Ligate: EcoRI-BstBI restricted vector +

EcoRI

PpuMI

```

AATTCATGCCAACTTTAATACTGCTAATTATTACAACAATGATTATGG
CATCTTCCTGCCAAATAGATATCACAAAACACTACAGCAATGTAGGTGTA
TTGGTCAACAGTCCCAAAGGGATGAAGATATCACAAAACCT . . . 3'
. . . GTACGGTTGAAATTATGACGATTAATAATGTTGTTACTAATACC
GTAGAAGGACGGTTTATCTATAGTGTTTTGATGTCGTACATCCACATA
ACCAGTTGTCAGGGTTTCCCTACTTCTATAGTGTTTTGAAGCTT
    
```



Construction of the chimeric FPIV-3-FRSV gene consisting of the truncated PIV-3 F gene devoid of the 5' untranslated region linked to the truncated RSV F1 gene.





FIG.11.






FIG.12B.

FIG.13

IMMUNOBLOTS OF CELL LYSATES FROM Sf9 CELLS
INFECTED WITH RECOMBINANT BACULOVIRUSES




FIG 13 : Immunoblots of cell lysates from Sf9 cells infected with recombinant baculoviruses containing the truncated RSV F gene (Lane 1), the chimeric FPIV-3⁻FRSV gene (Lane 2) or infected with wild type virus (Lane 3) reacted with anti-F RSV Mab (panel A) and anti-F1 PIV-3 antiserum (panel B)

CONSTRUCTION OF THE BACULOVIRUS TRANSFER VECTOR pD2

FIG.14.




FIG.15A. CONSTRUCTION OF THE F_{RSV}-HNPIV₃ CHIMERIC GENE




FIG.15B.




FIG.16

SDS POLY ACRYLAMIDE GEL AND IMMUNOBLOTS OF
PURIFIED F_{RSV}-HN_{PIV-3} CHIMERIC PROTEIN




FIG 16 : A) Coomassie-stained SDS polyacrylamide gel of immunoaffinity-purified F_{RSV}-HN_{PIV-3}protein.

B) Immunoblots of F_{RSV}-HN_{PIV-3}protein reacted with an anti-F RSV Mab (lane 1) and anti-HN PIV-3 antiserum (lane 2)

FIG.17. MUTAGENESIS OF THE PIV-3 F GENE





FIG.18. CONSTRUCTION OF THE F_{PIV3}-G_{RSV} CHIMERIC GENE



**MULTIMERIC HYBRID GENE ENCODING A
CHIMERIC PROTEIN WHICH CONFERS
PROTECTION AGAINST PARAINFLUENZA
VIRUS AND RESPIRATORY SYNCYTIAL
VIRUS**

This application is a continuation of application Ser. No. 08/001,554 filed Jan. 6, 1993 which application is now pending.

FIELD OF INVENTION

The present invention relates to the engineering and expression of multimeric hybrid genes containing sequences from the gene coding for immunogenic proteins or protein fragments of numerous pathogens.

BACKGROUND TO THE INVENTION

The advantage of the approach taken by the present invention is to produce single immunogens containing protective antigens from a range of pathogens. Such chimeras greatly simplify the development of combination vaccines, in particular, with the view ultimately to produce single dose multivalent vaccines. Multivalent vaccines are currently made by separately producing pathogens and/or their pertinent antigens and combining them in various formulations. This is a labour intensive, costly and complex manufacturing procedure. In contrast, the availability of a single immunogen capable of protecting against a range of diseases would solve many of the problems of multivalent vaccine production. Several chimeric immunogens of the type provided herein may be combined to decrease the number of individual antigens required in a multivalent vaccine.

Human Parainfluenza virus types 1,2,3 and Respiratory syncytial virus types A and B are the major viral pathogens responsible for causing severe respiratory tract infections in infants and young children. It is estimated that, in the United States alone, approximately 1.6 million infants under one year of age will have a clinically significant RSV infection each year and an additional 1.4 million infants will be infected with PIV-3. Approximately 4000 infants less than one year of age in the United States die each year from complications arising from severe respiratory tract disease caused by infection with RSV and PIV-3. The WHO and NIALD vaccine advisory committees ranked RSV number two behind HIV for vaccine development while the preparation of an efficacious PIV-3 vaccine is ranked in the top ten vaccines considered a priority for vaccine development.

Safe and effective vaccines for protecting infants against these viral infections are not available and are urgently required. Clinical trials have shown that formaldehyde-inactivated and live-attenuated viral vaccines failed to adequately protect vaccinees against these infections. In fact, infants who received the formalin-inactivated RSV vaccine developed more serious lower respiratory tract disease during subsequent natural RSV infection than did the control group. [Am. J. Epidemiology 89, 1969, p.405-421; J. Inf. Dis. 145, 1982, p.311-319]. Furthermore, RSV glycoproteins purified by immunoaffinity chromatography using elution at acid pH induced immunopotential in cotton rats. [Vaccine, 10(7), 1992, p.475-484]. The development of efficacious PIV-3 and RSV vaccines which do not cause exacerbated pulmonary disease in vaccinees following injection with wild-type virus would have significant therapeutic implications. It is anticipated that the development of a single recombinant immunogen capable of simultaneously protecting infants against diseases caused by infection with

both Parainfluenza and Respiratory syncytial viruses could significantly reduce the morbidity and mortality caused by these viral infections.

It has been reported that a protective response against PIV-3 and RSV is contingent on the induction of neutralizing antibodies against the major viral surface glycoproteins. For PIV, these protective immunogens are the HN protein which has a molecular weight of 72 kDa and possesses both hemagglutination and neuraminidase activities and the fusion (F) protein, which has a molecular weight of 65 kDa and which is responsible for both fusion of the virus to the host cell membrane and cell-to-cell spread of the virus. For RSV, the two major immunogenic proteins are the 80 to 90 kDa G glycoprotein and the 70 kDa fusion (F) protein. The G and F proteins are thought to be functionally analogous to the PIV HN and F proteins, respectively. The PIV and RSV F glycoproteins are synthesized as inactive precursors (FO) which are proteolytically cleaved into N-terminal F2 and C-terminal F1 fragments which remain linked by disulphide bonds.

Recombinant surface glycoproteins from PIV and RSV have been individually expressed in insect cells using the baculovirus system [Ray et al., (1989), Virus Research, 12: 169-180; Coelingh et al., (1987), Virology, 160: 465-472; Wathen et al., (1989), J. of Inf. Dis. 159: 253-263] as well as in mammalian cells infected with recombinant poxviruses [Spriggs, et al., (1987), J. Virol. 61: 3416-3423; Stott et al., (1987), J. Virol. 61: 3855-3861]. Recombinant antigens produced in these systems were found to protect immunized cotton rats against live virus challenge. More recently, hybrid RSV F-G [Wathan et al., (1989), J. Gen Virol. 70: 2625-2635; Wathan, published International Patent application WO 89/05823] and PIV-3 F-HN [Wathan, published International Patent Application WO 89/10405], recombinant antigens have been engineered and produced in mammalian and insect cells. The RSV F-G hybrid antigen was shown to be protective in cotton rats [Wathan et al., (1989), J. Gen. Virol. 70: 2637-2644] although it elicited a poor anti-G antibody response [Connors et al., (1992), Vaccine 10: 475-484]. The protective ability of the PIV-3 F-HN protein was not reported in the published patent application. These antigens were engineered with the aim to protect against only the homologous virus, that is either RSV or PIV-3. However, it would be advantageous and economical to engineer and produce a single recombinant immunogen containing at least one protective antigen from each virus in order simultaneously to protect infants and young children against both PIV and RSV infections. The chimeric proteins provided herein for such purpose also may be administered to pregnant women or women of child bearing age to stimulate maternal antibodies to both PIV and RSV. In addition, the vaccine also may be administered to other susceptible individuals, such as the elderly.

SUMMARY OF INVENTION

In its broadest aspect, the present invention provides a multimeric hybrid gene, comprising a gene sequence coding for an immunogenic region of a protein from a first pathogen linked to a gene sequence coding for an immunogenic region of a protein from a second pathogen and to a chimeric protein encoded by such multimeric hybrid gene. Such chimeric protein comprises an immunogenic region of a protein from a first pathogen linked to an immunogenic region of a protein from a second pathogen.

The first and second pathogens are selected from bacterial and viral pathogens and, in one embodiment, may both be

viral pathogens. Preferably, the first and second pathogens are selected from those causing different respiratory tract diseases, which may be upper and lower respiratory tract diseases. In a preferred embodiment, the first pathogen is parainfluenza virus and the second pathogen is respiratory syncytial virus. The PIV protein particularly is selected from PIV-3 F and HN proteins and the RSV protein particularly is selected from RSV G and F proteins. Another aspect of the invention provides cells containing the multimeric hybrid gene for expression of a chimeric protein encoded by the gene. Such cells may be bacterial cells, mammalian cells, insect cells, yeast cells or fungal cells. Further, the present invention provides a live vector for antigen delivery containing the multimeric hybrid gene, which may be a viral vector or a bacterial vector, and a physiologically-acceptable carrier therefor. Such live vector may form the active component of a vaccine against diseases caused by multiple pathogenic infections. Such vaccine may be formulated to be administered in an injectable form, intranasally or orally.

In an additional aspect of the present invention, there is provided a process for the preparation of a chimeric protein, which comprises isolating a gene sequence coding for an immunogenic region of a protein from a first pathogen; isolating a gene sequence coding for an immunogenic region of a protein from a second pathogen; linking the gene sequences to form a multimeric hybrid gene; and expressing the multimeric hybrid gene in a cellular expression system. The first and second pathogens are selected from bacterial and viral pathogens. Such cellular expression system may be provided by bacterial cells, mammalian cells, insect cells, yeast cells or fungal cells. The chimeric protein product of gene expression may be separated from a culture of the cellular expression system and purified.

The present invention further includes a vaccine against diseases caused by multiple pathogen infections, comprising the chimeric protein encoded by the multimeric hybrid gene and a physiologically-acceptable carrier therefor. Such vaccine may be formulated to be administered in an injectable form, intranasally or orally.

The vaccines provided herein may be used to immunize a host against disease caused by multiple pathogenic infections, particularly those caused by a parainfluenza virus and respiratory syncytial virus, by administering an effective amount of the vaccine to the host. As noted above, for human PIV and RSV, the host may be infants and young children, pregnant women as well as those of a child-bearing age, and other susceptible persons, such as the elderly.

The chimeric protein provided herein also may be used as a diagnostic reagent for detecting infection by a plurality of different pathogens in a host, using a suitable assaying procedure.

It will be appreciated that, while the description of the present invention which follows focuses mainly on a chimeric molecule which is effective for immunization against diseases caused by infection by PIV and RSV, nevertheless the invention provided herein broadly extends to any chimeric protein which is effected for immunization against diseases caused by a plurality of pathogens, comprising an antigen from each of the pathogens linked in a single molecule, as well as to genes coding for such chimeric molecules.

In this application, by the term "multimeric hybrid genes" we mean genes encoding antigenic regions of proteins from different pathogens and by the term "chimeric proteins" we mean immunogens containing antigenic regions from proteins from different pathogens.

BRIEF DESCRIPTION OF DRAWINGS

FIG. 1 shows the nucleotide (SEQ ID No: 1) and amino acid (SEQ ID No: 2) sequence of a PCR-amplified PIV-3 F gene and F protein, respectively;

FIG. 2 shows the restriction map of the PIV-3 F gene;

FIG. 3 shows the nucleotide (SEQ ID No: 3) and amino acid (SEQ ID No: 4) sequences of the PIV-3 HN gene and HN protein, respectively;

FIG. 4 shows the restriction map of the PIV-3 HN gene;

FIG. 5 shows the nucleotide (SEQ ID No: 5) and amino acid (SEQ ID No: 6) sequences of the RSV F gene and RSV F protein, respectively;

FIG. 6 shows the restriction map of the RSV F gene;

FIG. 7 shows the nucleotide (SEQ ID No: 7) and amino acid (SEQ ID No: 8) sequences of the RSV G gene and RSV G protein, respectively;

FIG. 8 shows the restriction map of the RSV G gene;

FIG. 9 shows the steps involved in the construction of an expression vector containing a chimeric F_{PIV-3} - F_{RSV} gene;

FIG. 10 shows the steps involved in the construction of an expression vector containing a F_{PIV-3} gene lacking the 5'-untranslated sequence and transmembrane anchor and cytoplasmic tail coding regions;

FIG. 11 shows the steps involved in the construction of an expression vector containing a chimeric F_{PIV-3} - F_{RSV} gene containing a truncated PIV-3 F gene devoid of 5'-untranslated region linked to a truncated RSV F1 gene;

FIG. 12 shows the steps involved in construction of a modified pAC 610 baculovirus expression vector containing a chimeric F_{PIV-3} - F_{RSV} gene consisting of the PIV-3 F gene lacking both the 5'-untranslated sequence as well as transmembrane and cytoplasmic tail coding region linked to the truncated RSV F1 gene;

FIGS. 13A and 13B show immunoblots of cell lysates from Sf9 cells infected with recombinant baculoviruses;

FIG. 14 shows the steps involved in constructing a baculovirus transfer vector (pD2);

FIG. 15 shows the steps involved in construction of a chimeric F_{RSV} - HN_{PIV-3} gene;

FIGS. 16A and 16B show an SDS-PAGE gel and immunoblot of purified F_{RSV} - HN_{PIV-3} chimeric protein;

FIG. 17 illustrates mutagenesis of a PIV-3 F gene; and

FIG. 18 shows the steps involved in the construction of a chimeric F_{PIV-3} - G_{RSV} gene.

GENERAL DESCRIPTION OF INVENTION

In the present invention, a chimeric molecule protective against two different major childhood diseases is provided. The present invention specifically relates to the formulation of various recombinant Parainfluenza virus (PIV)/Respiratory syncytial virus (RSV) immunogens to produce safe and efficacious vaccines capable of protecting infants and young children, as well as other susceptible individuals, against diseases caused by infection with both PIV and RSV. However, as described above, the present invention extends to the construction of multimeric hybrid genes containing genes coding for protective antigens from many pathogens. Such vaccines may be administered in any desired manner, such as a readily-injectable vaccine, intranasally or orally.

In the present invention, the inventors have specifically engineered several model PIV/RSV chimeric genes containing relevant sequences from selected genes coding for PIV-3 and RSV surface glycoproteins linked in tandem. All genes

in the chimeric constructs described herein were obtained from recent clinical isolates of PIV-3 and RSV. The chimeric gene constructs may include gene sequences from either PIV-3 F or HN genes linked in tandem to either RSV F or G genes in all possible relative orientations and combinations.

The chimeric gene constructs provided herein may consist of either the entire gene sequences or gene segments coding for immunogenic and protective epitopes thereof. The natural nucleotide sequence of these genes may be modified by mutation while retaining antigenicity and such modifications may include the removal of putative pre-transcriptional terminators to optimize their expression in eukaryotic cells. The genes were designed to code for hybrid PIV-RSV surface glycoproteins linked in tandem in a single construct to produce gene products which elicit protective antibodies against both parainfluenza and respiratory syncytial viruses. Such multimeric hybrid genes consist of a gene sequence coding for a human PIV-3 F or HN protein or an immunogenic epitope-containing fragment thereof linked to a gene sequence coding for a human RSV G or F protein or an immunogenic epitope-containing fragment thereof. Specific gene constructs which may be employed include $F_{PIV-3}-F_{RSV}$, $F_{RSV}-HN_{PIV-3}$ and $F_{PIV-3}-G_{RSV}$ hybrid genes.

In addition, the present invention also extends to the construction of other multimeric genes, such as trimeric genes containing PIV and RSV genes or gene segments, linked in all possible relative orientations. For example:

$F_{PIV}-HN_{PIV}-F$ or G_{RSV}

$F_{PIV}-F_{RSV}-G_{RSV}$

$HN_{PIV}-F_{RSV}-G_{RSV}$

The multimeric genes provided herein also may comprise at least one gene encoding at least one immunogenic and/or immunostimulating molecule.

The multimeric hybrid genes provided herein may be sub-cloned into appropriate vectors for expression in cellular expression systems. Such cellular expression systems may include bacterial, mammalian, insect and fungal, such as yeast, cells.

The chimeric proteins provided herein also may be presented to the immune system by the use of a live vector, including live viral vectors, such as recombinant poxviruses, adenoviruses, retroviruses, Semliki Forest viruses, and live bacterial vectors, such as Salmonella and mycobacteria (e.g. BCG).

Chimeric proteins, such as a PIV/RSV chimera, present in either the supernatants or cell lysates of transfected, transformed or infected cells then can be purified in any convenient manner.

To evaluate the immunogenicity and protective ability of the chimeric proteins, suitable experimental animals are immunized with either varying doses of the purified chimeric proteins, such as the PIV/RSV chimera, and/or live recombinant vectors as described above. Such chimeric proteins may be presented to the immune system by either the use of physiologically-acceptable vehicles, such as aluminum phosphate, or by the use of delivery systems, such as ISCOMS and liposomes. The chimeras also may be formulated to be capable of eliciting a mucosal response, for example, by conjugation or association with immunotargeting vehicles, such as the cholera toxin B subunit, or by incorporation into microparticles. The vaccines may further comprise means for delivering the multimeric protein specifically to cells of the immune system, such as toxin molecules or antibodies. To further enhance the immunoprotective ability of the chimeric proteins, they may be

supplemented with other immunogenic and/or immunostimulating molecules. The chimeric PIV/RSV proteins specifically described herein may be formulated with an adjuvant, such as aluminum phosphate, to produce readily-injectable vaccines for protection against the diseases caused by both PIV-3 and RSV. The chimeric proteins also may be administered intranasally or orally. The chimeric proteins may be used in test kits for diagnosis of infection by PIV-3 and RSV.

The invention is not limited to the preparation of chimeric PIV-3 and RSV proteins, but is applicable to the production of chimeric immunogens composed of either the entire sequences or regions of the immunogenic proteins from at least two pathogens sequentially linked in a single molecule. Chimeric antigens also may be synthesized to contain the immunodominant epitopes of several proteins from different pathogens. These chimeric antigens may be useful as vaccines or as diagnostic reagents.

SEQUENCE IDENTIFICATION

Several nucleotide and amino acid sequences are referred to in the disclosure of this application. The following table identifies the sequences and the location of the sequence:

SEQ ID No.	Identification	Location
1	Nucleotide sequence for PCR-amplified PIV-3 F gene	FIG. 1, Example 1
2	Amino acid sequence for PCR-amplified PIV-F protein	FIG. 1, Example 1
3	Nucleotide sequence for PIV-3 HN gene	FIG. 3, Example 1
4	Amino acid sequence for PIV-3 HN protein	FIG. 3, Example 1
5	Nucleotide sequence for RSV F gene	FIG. 5, Example 1
6	Amino acid sequence for RSV F protein	FIG. 5, Example 1
7	Nucleotide sequence for RSV G gene	FIG. 7, Example 1
8	Amino acid sequence for RSV G protein	FIG. 7, Example 1
9	BsrI - BamHI oligonucleotide cassette	FIG. 9, Example 2
10	BspHI - BamHI oligonucleotide cassette	FIG. 9, Example 2
11	EcoRI - Ppu MI oligonucleotide cassette	FIG. 9, Example 2
12	BrsI - BamHI oligonucleotide cassette	FIG. 10, Example 3
13	EcoRI - Bsr BI oligonucleotide cassette	FIG. 10, Example 3
14	EcoRV - EcoRI oligonucleotide cassette	FIG. 11, Example 4
15	EcORV - BamHI oligonucleotide cassette	FIG. 14, Example 8
16	BspHI - BspHI oligonucleotide cassette	FIG. 15, Example 9
17	Nucleotide sequence for PIV-3 F gene	Example 15
18	Mutagenic oligonucleotide #2721	FIG. 17, Example 15
19	Nucleotide sequence for part of oligonucleotide #2721	Example 15
20	Oligonucleotide probe	Example 15

DEPOSIT INFORMATION

Certain plasmid DNAs described and referred to herein have been deposited with the American Type Culture Col-

lection (ATCC) located at 10801 University Boulevard, Manassas, Va. 20110-2209, USA, pursuant to the Budapest Treaty and prior to the filing of this application. The deposited purified plasmids will become available to the public and all restrictions imposed on access to the deposits will be irrevocable removed. The deposit will be replaced if viable samples cannot be dispensed by the depository upon grant of this U.S. patent application or upon publication of its corresponding European patent application, whichever first occurs. The invention described and claimed herein is not to be limited in scope by the plasmid DNAs of the constructs deposited, since the deposited embodiment is intended only as an illustration of the invention. The following purified plasmids were deposited at the ATCC with the noted accession numbers on Dec. 17, 1992:

Plasmid	Example No.	Accession No.
pAC DR7	5	75387
pD2RF-HN	9	75388
pD2F-G	16	75389

Any equivalent plasmids that can be used to produce equivalent antigens as described in this application are within the scope of the invention.

EXAMPLES

The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific Examples. These Examples are described solely for purposes of illustration and are not intended to limit the scope of the invention. Changes in form and substitution of equivalents are contemplated as circumstances may suggest or render expedient. Although specific terms have been employed herein, such terms are intended in a descriptive sense and not for purposes of limitations.

Methods for cloning and sequencing the PIV-3 and RSV genes as well as the procedures for sub-cloning the genes into appropriate vectors and expressing the gene constructs in mammalian and insect cells are not explicitly described in this disclosure but are well within the scope of those skilled in the art.

Example 1

This Example outlines the strategy used to clone and sequence the PIV-3 F, HN and RSV F, G genes (from a type A isolate). These genes were used in the construction of the $F_{PIV-3}-F_{RSV}$, $F_{RSV}-HN_{PIV-3}$, and $F_{PIV-3}-G_{RSV}$ chimeric genes detailed in Examples 2 to 4, 9 and 15, respectively.

Two PIV-3 F gene clones initially were obtained by PCR amplification of cDNA derived from viral RNA extracted from a recent clinical isolate of PIV-3. Two other PIV-3 F gene clones as well as the PIV-3 HN, RSV F and RSV G genes were cloned from a cDNA library prepared from mRNA isolated from MRC-5 cells infected with clinical isolates of either PIV-3 or RSV (type A isolate). The PIV-3 F (both PCR amplified and non-PCR amplified), PIV-3 HN, RSV F and RSV G gene clones were sequenced by the dideoxynucleotide chain termination procedure. Sequencing of both strands of the genes was performed by a combination of manual and automated sequencing.

The nucleotide (SEQ ID No: 1) and amino acid (SEQ ID No: 2) sequences of the PCR amplified PIV-3 F gene and F protein, respectively, are presented in FIG. 1 and the restriction map of the gene is shown in FIG. 2. Sequence analysis

of the 1844 nucleotides of two PCR amplified PIV-3 F gene clones confirmed that the clones were identical. Comparison of the coding sequence of the PCR-amplified PIV-3 F gene clone with that of the published PIV-3 F gene sequence revealed a 2.6% divergence in the coding sequence between the two genes resulting in fourteen amino acid substitutions.

The nucleotide sequence of the non-PCR amplified PIV-3 F gene clone differed from the PCR amplified gene clone in the following manner: the non-PCR amplified clone had ten additional nucleotides (AGGACAAAAG-SEQ ID NO: 38) at the 5' untranslated region of the gene and differed at four positions, 8 (T in PCR-amplified gene to C in non-PCR amplified gene), 512 (C in PCR-amplified gene to T in non-PCR amplified gene), 518 (G in PCR-amplified gene to A in non-PCR amplified gene) and 1376 (A in PCR-amplified gene to G in non-PCR amplified gene). These changes resulted in three changes in the amino acid sequence of the F protein encoded by the non-PCR amplified PIV-3 F gene. Serine (position 110), glycine (position 112), and aspartic acid (position 398) in the primary amino acid sequence of the F protein encoded by the PCR amplified PIV-3 F gene was changed to phenylalanine (position 110), glutamic acid (position 112) and glycine (position 398), respectively, in the primary amino acid sequence of the F protein encoded by the PCR amplified clone.

FIG. 3 shows the nucleotide (SEQ ID No: 3) and amino acid (SEQ ID No: 4) sequences of the PIV-3 HN gene and protein, respectively and the restriction map of the gene is presented in FIG. 4. Analysis of the 1833 nucleotide sequence from two HN clones confirmed that the sequences were identical. A 4.4% divergence in the coding sequence of the PIV-3 HN gene was noted when the S sequence was compared to the published PIV-3 HN coding sequence. This divergence resulted in seventeen amino acid substitutions in the amino acid sequence of the protein encoded by the PIV-3 HN gene.

The nucleotide (SEQ ID No: 5) and amino acid (SEQ ID No: 6) sequences of the RSV F gene and RSV F protein, respectively, are shown in FIG. 5 and the restriction map of the gene is shown in FIG. 6. Analysis of the 1886 nucleotide sequence from two RSV F clones verified complete sequence homology between the two clones. Comparison of this nucleotide sequence with that reported for the RSV F gene revealed approximately 1.8% divergence in the coding sequence resulting in eleven amino acid substitutions.

The nucleotide (SEQ ID No: 7) and amino acid (SEQ ID No: 8) sequences of the RSV G gene and RSV G protein, respectively, are presented in FIG. 7 while the restriction map of the gene is outlined in FIG. 8. Comparison of the 920 nucleotide sequence of the G gene clone with the published G sequence (type A isolate) revealed a 4.2% divergence in the nucleotide sequence and a 6.7% divergence in the amino acid sequence of the gene product. This divergence resulted in twenty amino acid substitutions.

The full-length PIV-3 F (non-PCR amplified), PIV-3 HN, RSV F and RSV G genes were cloned into λ gt11 and subcloned into the multiple cloning site of a Bluescript M13-SK vector, either by blunt end ligation or using appropriate linkers. The PCR-amplified PIV-3 F gene was directly cloned into the Bluescript vector. The cloning vectors containing the PIV-3 F-PCR amplified, PIV-3 F non-PCR amplified, PIV-3 HN, RSV F and RSV G genes were named pPI3F, pPI3Fc, pPIVHN, pRSVF and pRSVG, respectively.

Example 2

This Example illustrates the construction of a Bluescript-based expression vector (pMCR20) containing the chimeric

F_{PIV-3} - F_{RSV} gene. This chimeric gene construct contains the 5' untranslated region of the PIV-3 F gene but lacks the hydrophobic anchor and cytoplasmic tail coding regions of both the PIV-3 and RSV F genes. The steps involved in the construction of this plasmid are summarized in FIG. 9.

To prepare the PIV-3 portion of the chimeric gene (FIG. 9, step 1), the full length PIV-3 gene lacking the transmembrane region and cytoplasmic tail coding regions was retrieved from plasmid pPI3F by cutting the polylinker with BamHI, blunt-ending the linearized plasmid with Klenow polymerase and cutting the gene with BsrI. A BsrI-BamHI oligonucleotide cassette (SEQ ID No: 9) containing a PpuMI site and three successive translational stop codons were ligated to the truncated 1.6 Kb [BamHI]-BsrI PIV-3 F gene fragment and cloned into the EcoRV-BamHI sites of a Bluescript M13-SK expression vector containing the human methallothionin promoter and the poly A and IVS sequences of the SV40 genome (designated pMCR20), to generate plasmid pME1.

To engineer the RSV F gene component of the chimeric construct (FIG. 9, step 2), the RSV F gene lacking the transmembrane region and cytoplasmic tail coding regions was retrieved from plasmid pRSVF by cutting the polylinker with EcoRI and the gene with BspHI. A synthetic BspHI-BamHI oligonucleotide cassette (SEQ ID No: 10) containing three successive translational stop codons was ligated to the 1.6 Kb truncated RSV F gene and cloned into the EcoRI-BamHI sites of the Bluescript based expression vector, pMCR20 to produce plasmid pES13A. Plasmid pES13A then was cut with EcoRI and PpuMI to remove the leader and F2 coding sequences from the truncated RSV F gene. The leader sequence was reconstructed using an EcoRI-PpuMI oligocassette (SEQ ID No: 11) and ligated to the RSV F1 gene segment to generate plasmid pES23A.

To prepare the chimeric F_{PIV-3} - F_{RSV} gene (FIG. 9, step 3) containing the 5' untranslated region of the PIV-3 F gene linked to the truncated RSV F1 gene fragment, plasmid pME1 (containing the 1.6 Kb truncated PIV-3 F gene) first was cut with PpuMI and BamHI. The PpuMI-BamHI restricted pME1 vector was dephosphorylated with intestinal alkaline phosphatase. The 1.1 Kb RSV F1 gene fragment was retrieved from plasmid pES23A by cutting the plasmid with PpuMI and BamHI. The 1.1 Kb PpuMI-BamHI RSV F1 gene fragment was cloned into the PpuMI-BamHI sites of the dephosphorylated pME1 vector to generate plasmid pES29A. This chimeric gene construct contains the 5' untranslated region of the PIV-3 F gene but lacks the nucleotide sequences coding for the hydrophobic anchor domains and cytoplasmic tails of both the PIV-3 and RSV F proteins.

Example 3

This Example illustrates the construction of a Bluescript-based expression vector containing the PIV-3 F gene lacking both the 5' untranslated and transmembrane anchor and cytoplasmic tail coding regions. The steps involved in constructing this plasmid are outlined in FIG. 10.

Plasmid pPI3F containing the full length PIV-3 F gene was cut with BamHI, blunt ended with Klenow polymerase and then cut with BsrI to remove the transmembrane and cytoplasmic tail coding regions. The Bluescript-based expression vector, pMCR20, was cut with SmaI and BamHI. A synthetic BsrI-BamHI oligonucleotide cassette (SEQ ID No: 12) containing a translational stop codon was ligated with the 1.6 Kb blunt ended-BsrI PIV-3 F gene fragment to the SmaI-BamHI restricted pMCR20 vector to produce

plasmid PMpFB. The PIV-3 F gene of this construct lacked the DNA fragment coding for the transmembrane and cytoplasmic anchor domains but contained the 5' untranslated region. To engineer a plasmid containing the PIV-3 F gene devoid of both the 5' untranslated region and the DNA fragment coding for the hydrophobic anchor domain, plasmid pMpFB was cut with EcoRI and BstBI. An EcoRI-BstBI oligocassette (SEQ ID No: 13) containing the sequences to reconstruct the signal peptide and coding sequences removed by the EcoRI-BstBI cut was ligated to the EcoRI-BstBI restricted pMpFB vector to produce plasmid pMpFA.

Example 4

This Example illustrates the construction of the chimeric F_{PIV-3} - F_{RSV} gene composed of the truncated PIV-3 F gene devoid of the 5' untranslated region linked to the truncated RSV F1 gene. The steps involved in constructing this plasmid are summarized in FIG. 11.

To prepare this chimeric gene construct, plasmid pES29A (Example 2) was cut with BstBI and BamHI to release the 2.5 Kb BstBI-BamHI PIV-3 F-RSV F1 chimeric gene fragment. This BstBI-BamHI fragment was isolated from a low melting point agarose gel and cloned into the BstBI-BamHI sites of the dephosphorylated vector pMpFA to produce plasmid pES60A. This construct contained the PIV-3 F gene lacking both the 5' untranslated region and the hydrophobic anchor and cytoplasmic tail coding sequences linked to the F1 coding region of the truncated RSV F gene. This chimeric gene was subsequently subcloned into the baculovirus transfer vector (see Example 5).

Example 5

This Example illustrates the construction of the modified pAC 610 baculovirus transfer vector containing the native polyhedrin promoter and the chimeric F_{PIV-3} - F_{RSV} gene consisting of the PIV-3 F gene lacking both the 5' untranslated sequence and the nucleotide sequence coding for the hydrophobic anchor domain and cytoplasmic tail linked to the truncated RSV F1 gene. Construction of this plasmid is illustrated in FIG. 12.

The pAC 610 baculovirus expression vector was modified to contain the native polyhedrin promoter in the following manner. Vector pAC 610 was cut with EcoRV and BamHI. The 9.4 Kb baculovirus transfer vector lacking the EcoRV-BamHI DNA sequence was isolated from a low melting point agarose gel and treated with intestinal alkaline phosphatase. In a 3-way ligation, an EcoRV-EcoRI oligonucleotide cassette (SEQ ID No: 14) containing the nucleotides required to restore the native polyhedrin promoter was ligated with the 1.6 Kb EcoRI-BamHI truncated RSV F gene fragment isolated from construct pES13A (Example 2, step 2) and the EcoRV-BamHI restricted pAC 610 phosphatased vector to generate plasmid pES47A. To prepare the pAC 610 based expression vector containing the chimeric F_{PIV-3} - F_{RSV} gene, plasmid pES47A was first cut with EcoRI and BamHI to remove the 1.6 Kb truncated RSV F gene insert. The 2.8 Kb F_{PIV-3} - F_{RSV} chimeric gene was retrieved by cutting plasmid pES60A (Example 4) with EcoRI and BamHI. The 2.8 Kb EcoRI-BamHI chimeric gene was ligated to the EcoRI-BamHI restricted pES47A vector to generate plasmid pAC DR7 (ATCC 75387).

Example 6

This Example outlines the preparation of plaque-purified recombinant baculoviruses containing the chimeric F_{PIV-3} - F_{RSV} gene.

11

Spodoptera frugiperda (Sf9) cells were co-transfected with 1.0 μ g wild-type AcMNPV DNA and 2.5 μ g of F_{PIV-3}-FRSV plasmid DNA (plasmid pAC DR7-Example 5). Putative recombinant baculoviruses (purified once by serial dilution) containing the F_{PIV-3}-F_{RSV} chimeric gene were identified by dot-blot hybridization. Lysates of insect cells infected with the putative recombinant baculoviruses were probed with the ³²P-labelled F_{PIV-3}-F_{RSV} chimeric gene insert. Recombinant baculoviruses were plaque-purified twice before being used for expression studies. All procedures were carried out according to the protocols outlined by M. D. Summers and G. E. Smith in "A Manual of Methods for Baculovirus Vectors and Insect Cell Culture Procedures", Texas Agricultural Experiment Station, Bulletin 1555, 1987.

Example 7

This Example illustrates the presence of the chimeric F_{PIV-3}-F_{RSV} protein in supernatants and cell lysates of infected Sf9 cells.

Insect cells were infected with the plaque-purified recombinant baculoviruses prepared as described in Example 6 at a m.o.i. of 8. Concentrated supernatants from cells infected with the recombinant viruses were positive in a PIV-3 F specific ELISA. In addition, when lysates from ³⁵S-methioninelabelled infected cells were subjected to SDS-polyacrylamide gel electrophoresis and gels were analyzed by autoradiography, a strong band with apparent molecular weight of approximately 90 kDa was present in lysates of cells infected with the recombinant viruses but was absent in the lysates from wild-type infected cells. The presence of the chimeric F_{PIV-3}-F_{RSV} protein in the lysates of cells infected with the recombinant baculoviruses was confirmed further by Western blot analysis using monospecific anti-PIV-3 F and anti-RSV F antisera and/or monoclonal antibodies (Mabs). Lysates from cells infected with the recombinant baculoviruses reacted with both anti-PIV-3 and anti-RSV antisera in immunoblots. As shown in the immunoblot of FIG. 13, lysates from cells infected with either the RSV F or F_{PIV-3}-F_{RSV} recombinant baculoviruses reacted positively with the anti-F RSV Mab. As expected, lysates from cells infected with wild type virus did not react with this Mab. In addition, only lysates from cells infected with the chimeric F_{PIV-3}-F_{RSV} recombinant viruses reacted with the anti-PIV-3 F₁ antiserum.

Example 8

This Example illustrates modification of the baculovirus transfer vector pVL1392 (obtained from Invitrogen), wherein the polyhedrin ATG start codon was converted to ATT and the sequence CCG was present downstream of the polyhedrin gene at positions +4,5,6. Insertion of a structural gene several base pairs downstream from the ATT codon is known to enhance translation. The steps involved in constructing this modified baculovirus transfer vector are outlined in FIG. 14.

The baculovirus expression vector pVL1392 was cut with EcoRV and BamHI. The 9.5 kb restricted pVL1392 vector was ligated to an EcoRV-BamHI oligonucleotide cassette (SEQ ID No: 15) to produce the pD2 vector.

Example 9

This Example illustrates the construction of the pD2 baculovirus expression vector containing the chimeric F_{RSV}-HN_{PIV-3} gene consisting of the truncated RSV F and PIV-3 HN genes linked in tandem. The steps involved in constructing this plasmid are summarized in FIG. 15.

12

To engineer the F_{RSV}-HN_{PIV-3} gene, the RSV F gene lacking the nucleotide sequence coding for the transmembrane domain and cytoplasmic tail of the RSV F glycoprotein was retrieved from plasmid PRSVF (Example 1) by cutting the polylinker with EcoRI and the gene with BspHI. The PIV-3 HN gene devoid of the DNA fragment coding for the hydrophobic anchor domain was retrieved from plasmid pPIVHN (Example 1) by cutting the gene with BspHI and the polylinker with BamHI. The 1.6 Kb EcoRI-BspHI RSV F gene fragment and the 1.7 Kb BspHI-BamHI PIV-3 HN gene fragment were isolated from low melting point agarose gels. For cloning purposes, the two BspHI sites in the Bluescript based mammalian cell expression vector, pMCR20, were mutated. Mutations were introduced in the BspHI sites of the pMCR20 by cutting the expression vector with BspHI, treating both the BspHI restricted vector and the 1.1 Kb fragment released by the BspHI cut with Klenow polymerase and ligating the blunt-ended 1.1 Kb fragment to the blunt-ended Bluescript-based expression vector to generate plasmid pM'. Since insertion of the 1.1 Kb blunt-end fragment in the mammalian cell expression vector in the improper orientation would alter the Amp^r gene of the Bluescript-based expression vector, only colonies of HB101 cells transformed with the pM' plasmid DNA with the 1.1 Kb blunt-ended fragment in the proper orientation could survive in the presence of ampicillin. Plasmid DNA was purified from ampicillin-resistant colonies of HB101 cells transformed with plasmid pM' by equilibrium centrifugation in cesium chloride-ethidium bromide gradients. The 1.6 Kb EcoRI-BspHI RSV F and 1.7 Kb BspHI-BamHI PIV-3 HN gene fragments were directly cloned into the EcoRI-BamHI sites of vector pM' in a 3-way ligation to generate plasmid pM' RF-HN.

To restore specific coding sequences of the RSV F and PIV-3 HN genes removed by the BspHI cut, a BspHI-BspHI oligonucleotide cassette (SEQ ID No: 16) containing the pertinent RSV F and PIV-3 HN gene sequences was ligated via the BspHI site to the BspHI-restricted plasmid pM' RF-HN to produce plasmid pM RF-HN. Clones containing the BspHI-BspHI oligonucleotide cassette in the proper orientation were identified by sequence analysis of the oligonucleotide linker and its flanking regions.

To clone the chimeric F_{RSV}-HN_{PIV-3} gene into the baculovirus expression vector pD2 (Example 8), the F_{RSV}-HN_{PIV-3} truncated gene first was retrieved from plasmid pM RF-HN by cutting the plasmid with EcoRI. The 3.3 Kb F_{RSV}-HN_{PIV-3} gene then was cloned into the EcoRI site of the baculovirus transfer vector plasmid pD2 to generate plasmid pD2 RF-HN (ATCC 75388). Proper orientation of the 3.3 Kb EcoRI F_{RSV}-HN_{PIV-3} chimeric gene insert in plasmid pD2 RF-HN was confirmed by sequence analysis.

Example 10

This Example outlines the preparation of plaque-purified recombinant baculoviruses containing the chimeric F_{RSV}-HN_{PIV-3} gene.

Spodoptera frugiperda (Sf9) cells were co-transfected with 1 μ g wild-type AcNPV DNA and 2 μ g of F_{RSV}-HN_{PIV-3} plasmid DNA (plasmid pD2 RF-HN-Example 9). Putative recombinant baculoviruses (purified once by serial dilution) containing the F_{RSV}-HN_{PIV-3} chimeric gene were identified by dot-blot hybridization. Lysates of insect cells infected with the putative recombinant baculoviruses were probed with the ³²P-labelled RSV F or PIV-3 HN gene oligonucleotide probes. Recombinant baculoviruses were plaque-purified three times before being used for expression studies.

13

All procedures were carried out according to the protocols outlined by Summers and Smith (Example 6).

Example 11

This Example illustrates the presence of the chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein in supernatants of infected Sf9 and High 5 cells.

Insect cells (Sf9 and High 5), maintained in serum free medium EX401, were infected with the plaque purified recombinant baculoviruses of Example 10 at a m.o.i. of 5 to 10 pfu/cell. Supernatants from cells infected with the recombinant baculoviruses tested positive for expressed protein in both the RSV-F and PIV-3 HN specific ELISAS. In addition, supernatants from infected cells reacted positively with both an anti-F RSV monoclonal antibody and anti-HN peptide antisera on immunoblots. A distinct band of approximately 105 kDa was present in the immunoblots. These results confirm the secretion of the chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein into the supernatant of Sf9 and High 5 cells infected with the recombinant baculoviruses.

Example 12

This Example illustrates the purification of the chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein from the supernatants of infected High 5 cells.

High 5 cells, maintained in serum free medium, were infected with the plaque purified recombinant baculoviruses of Example 10 at a m.o.i of 5 pfu/cell. The supernatant from virus infected cells was harvested 2 days post-infection. The soluble $F_{RSV}\text{-HN}_{PIV-3}$ chimeric protein was purified from the supernatants of infected cells by immunoaffinity chromatography using an anti-HN PIV-3 monoclonal antibody. The anti-HN monoclonal antibody was coupled to CNBr-activated Sepharose 4B by conventional techniques. The immunoaffinity column was washed with 10 bed volumes of washing buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.02% v/v TRITON X-100 (Trademark for a non-ionic detergent) which is octadienyl phenol (ethylene glycol)₁₀ prior to use. After sample loading, the column was washed with 10 bed volumes of washing buffer followed by 3 bed volumes of high salt buffer (10 mM Tris-HCl pH 7.5, 500 mM NaCl, 0.02% v/v Triton-X 100). The chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein was eluted from the immunoaffinity column with 100 mM glycine, pH 2.5, in the presence of 0.02% TRITON X-100. Eluted protein was neutralized immediately with 1M Tris-HCl, pH 10.7.

Polyacrylamide gel electrophoretic analysis (FIG. 16, panel A) of the immunoaffinity-purified $F_{RSV}\text{-HN}_{PIV-3}$ protein revealed the presence of one major protein band with an apparent molecular weight of 105 kDa. The purified protein reacted with both an anti-RSV F monoclonal antibody and anti-HN peptide antisera on immunoblots (FIG. 16, panel B, lanes 1 and 2, respectively).

Example 13

This Example illustrates the immunogenicity of the $F_{RSV}\text{-HN}_{PIV-3}$ protein in guinea pigs.

Groups of four guinea pigs were injected intramuscularly with either 1.0 or 10.0 μg of the chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein purified as described in Example 12 and adjuvanted with aluminum phosphate. Groups of control animals were immunized with either placebo, or live PIV-3 or RSV (administered intranasally). Guinea pigs were bled 2 and 4 weeks after the primary injection and boosted at 4 weeks with an equivalent dose of the antigen formulation. Serum

14

samples also were taken 2 and 4 weeks after the booster dose. To assess the ability of the chimeric protein to elicit PIV-3 and RSV-specific antibody responses, sera samples were analyzed for the presence of PIV-3 specific hemagglutination inhibiting and neutralizing antibodies as well as RSV neutralizing antibodies. As summarized in Table 1 below (the Tables appear at the end of the disclosure), the sera of animals immunized with two 10 μg doses of the chimeric protein had titres of PIV-3 specific hemagglutination inhibition (HAI) and PIV-3/RSV neutralizing antibodies at the 6 and 8 week time points which were equivalent to the levels obtained following intranasal inoculation with either live PIV-3 or RSV. In addition, animals immunized with only two 1 μg doses of the chimeric protein elicited strong PIV-3 and RSV specific neutralizing antibodies. These results confirmed the immunogenicity of both the RSV and PIV-3 components of the chimeric protein and provided confirmatory evidence that a single recombinant immunogen can elicit neutralizing antibodies against both RSV and PIV-3.

Example 14

This Example illustrates the immunogenicity and protective ability of the $F_{RSV}\text{-HN}_{PIV-3}$ protein in cotton rats.

Groups of eight cotton rats were injected intramuscularly with either 1.0 or 10.0 μg of the chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein (prepared as described in Example 12) adjuvanted with aluminum phosphate. Groups of control animals were immunized with either placebo (PBS+aluminum phosphate) or live PIV-3 or RSV (administered intranasally). Cotton rats were bled 4 weeks after the primary injection and boosted at 4 weeks with an equivalent dose of the antigen formulation. Serum samples were also taken 1 week after the booster dose. As shown in Table 2 below, data from the 4-week bleed demonstrated that both a 1 and 10 μg dose of the chimeric protein was capable of inducing a strong primary response. Reciprocal mean \log_2 PIV-3 specific HAI and PIV-3/RSV neutralizing titers were equivalent to the titres obtained with live PIV-3 and RSV. Thus, a single inoculation of the chimeric protein was sufficient to elicit neutralizing antibodies against both PIV-3 and RSV. Strong neutralizing PIV-3 and RSV titres also were observed following the booster dose (5 week bleed). These results provide additional evidence that both the RSV and PIV-3 components of the chimeric protein are highly immunogenic.

To assess the ability of the chimeric immunogen to simultaneously protect animals against both RSV and PIV-3, four cotton rats from each group were challenged intranasally with 100 TCID₅₀ units of either PIV-3 or RSV. Animals were killed 4 days after virus challenge. Virus titers were determined in lung lavages. As shown in Table 3 below, animals immunized with either 1 or 10 μg of the chimeric $F_{RSV}\text{-HN}_{PIV-3}$ protein were completely protected against challenge with either PIV-3 or RSV. These results provide evidence that the chimeric protein is not only highly immunogenic but can also simultaneously protect cotton rats against disease caused by both PIV-3 and RSV infection.

Example 15

This Example illustrates the construction of a Bluescript M13-SK vector containing the chimeric $F_{PIV-3}\text{-G}_{RSV}$ gene. This chimeric gene construct contains the 5' untranslated region of a mutated PIV-3 F gene but lacks the nucleotide sequence coding for the hydrophobic anchor and cytoplasmic tail domains of both a mutated PIV-3 F and the native RSV G genes. The steps involved in constructing this plasmid are outlined in FIGS. 17 and 18.

The first step (FIG. 17) involved in preparing the PIV-3 F component of the chimeric F_{PIV-3} - G_{RSV} gene construct was to eliminate the putative pre-termination sites within the 18 nucleotide long sequence 5' CAAGAAAAGGAATAAAA 3' (SEQ ID No: 17) located between positions 857 and 874 of the non PCR-amplified PIV-3 F gene and positions 847 and 864 of the PCR-amplified PIV-3 F gene (see FIG. 1). To this end, the PIV-F cDNA of the non-PCR amplified PIV-3 F gene was cut at the BsaAI and EcoRI sites. The BsaAI-EcoRI PIV F gene fragment was cloned into the EcoRI site of a Bluescript M13-SK vector using an EcoRI-BsaAI linker. The 857–874 target region of the PIV-3 F gene (non-PCR amplified) then was mutated by oligonucleotide-mediated mutagenesis using the method of Morinaga et al. [1984, Biotechnology 2: 636–639]. Plasmid pPI3Fc (Example 1) was cut with ScaI in the Amp^r gene and dephosphorylated with alkaline phosphatase (plasmid #1). A second sample of plasmid pPI3Fc was cut with BstEII and NsiI to produce a 3.9 Kb restricted plasmid, lacking the 0.9 Kb BstEII-NsiI fragment of the PIV-3 F gene (plasmid #2). A mutagenic 78-mer synthetic oligonucleotide (#2721 shown in FIG. 17-SEQ ID No: 18)) containing the sequence 5' CAGGAGAAGGGTATCAAG 3' (SEQ ID No: 19) was synthesized to specifically mutate the 857–874 DNA segment without changing the F protein sequence. This oligonucleotide was added to plasmid DNAs #1 and #2, denatured at 100° C. for 3 min. and renatured by gradual cooling. The mixture then was incubated in the presence of DNA polymerase, dNTPs and T4 ligase and transformed into HB101 cells. Bacteria containing the 1.8 Kb mutated PIV-3 F gene were isolated on YT agar plates containing 100 µg/ml ampicillin. Hybridization with the oligonucleotide probe 5' AGGAGAAGGGTATCAAG 3' (SEQ ID No: 20) was used to confirm the presence of the mutated PIV-3 F gene. The mutated gene sequence was confirmed by DNA sequencing. The plasmid containing the mutated PIV-3 gene was designated pPI3Fm.

The second step (FIG. 18) in the engineering of the chimeric gene construct involved constructing a Bluescript based vector to contain the truncated PIV-3 Fm gene lacking the nucleotide sequence coding for the transmembrane anchor domain and cytoplasmic tail of the PIV-3 F protein linked in tandem with the RSV G gene lacking both the 5' leader sequence and the nucleotide sequence coding for the transmembrane anchor domain and cytoplasmic tail of the G glycoprotein.

To prepare this chimeric gene, the orientation of the mutated PIV-F gene in plasmid pPI3Fm first was reversed by EcoRI digestion and religation to generate plasmid pPI3Fmr. To prepare the PIV-3 F gene component of the chimeric gene, plasmid pPI3Fmr was cut with NotI and BsrI to release the 1.7 Kb truncated PIV-3 F gene. To prepare the RSV G component, the 0.95 Kb RSV-G gene lacking both the 5' leader sequence and the DNA segment encoding the G protein anchor domain and cytoplasmic tail was released from plasmid pRSVG (Example 1) by cutting the polylinker with EcoRI and the gene with BamHI. The 0.95 Kb EcoRI-BamHI RSV G gene fragment was subcloned into the EcoRI-BamHI sites of a restricted Bluescript vector, pM13-SK, to produce plasmid pRSVGt. The 0.95 Kb EcoRI-BamHI G gene fragment and the 1.5 Kb NotI-BsrI truncated PIV-3 F gene were linked via a BsrI-BamHI oligonucleotide cassette (SEQ ID No: 9) restoring the F and G gene coding sequences and cloned into the pRSVGt vector restricted with BamHI and NotI in a 3-way ligation. The plasmid thus generated was designated pFG.

Example 16

This Example outlines the construction of the pD2 baculovirus transfer vector (described in Example 8) containing

the chimeric F_{PIV-3} - G_{RSV} gene consisting of a mutated PIV-3 F gene lacking the hydrophobic anchor and cytoplasmic coding regions linked to the RSV G gene lacking both the 5' leader sequence and the nucleotide sequences encoding the transmembrane anchor domain and cytoplasmic tail of the G protein.

To prepare this construct, plasmid pFG (Example 15) was cut with EcoRI to release the 2.6 Kb F_{PIV-3} - G_{RSV} chimeric gene. The 2.6 Kb EcoRI restricted chimeric gene fragment then was sub-cloned into the EcoRI site of the dephosphorylated pD2 vector to generate the 12.1 Kb plasmid pD2F-G (ATCC 75389).

Example 17

This Example outlines the preparation of plaque-purified recombinant baculoviruses containing the chimeric F_{PIV-3} - G_{RSV} gene.

Spodoptera frugiperda (Sf9) cells were co-transfected with 2 µg of pD2F-G plasmid DNA (Example 16) and 1 µg of linear wild-type AcNPV DNA (obtained from Invitrogen). Recombinant baculoviruses containing the F_{PIV-3} - G_{RSV} gene were plaque-purified twice according to the procedure outlined in Example 10.

Example 18

This Example illustrates the presence of the chimeric F_{PIV-3} - G_{RSV} protein in the supernatant of Sf9 and High 5 cells infected with the recombinant baculoviruses.

Sf9 and High 5 cells were infected with recombinant baculoviruses containing the F_{PIV-3} - G_{RSV} gene (Example 16) at a m.o.i. of 5 to 10 pfu/cell. The supernatant of cells infected with the recombinant viruses tested positive for expressed protein in the PIV-3 F specific ELISA. Supernatants of infected cells reacted with both anti-F PIV-3 and anti-G RSV monoclonal antibodies in immunoblots. These results confirm the presence of the chimeric F_{PIV-3} - G_{RSV} protein in the supernatants of infected Sf9 and High 5 cells.

Example 19

This Example outlines the preparation of recombinant vaccinia viruses expressing the F_{PIV-3} - F_{RSV} and F_{RSV} - HN_{PIV-3} genes.

Vaccinia virus recombinant viruses expressing the F_{PIV-3} - F_{RSV} (designated vP1192) and F_{RSV} - HN_{PIV-3} (designated vP1195) genes were produced at Virogenetics Corporation (Troy, N.Y.) (an entity related to assignee hereof) using the COPAK host-range selection system. Insertion plasmids used in the COPAK host-range selection system contained the vaccinia K1L host-range gene [Perkus et al., (1990) Virology 179:276–286] and the modified vaccinia H6 promoter [Perkus et al. (1989), J. Virology 63:3829–3836]. In these insertion plasmids, the K1L gene, H6 promoter and polylinker region are situated between Copenhagen strain vaccinia flanking arms replacing the AT1 region [open reading frames (ORFs) A25L, A26L; Goebel et al., (1990), Virology 179: 247–266; 517–563]. COPAK insertion plasmids are designed for use in in vivo recombination using the rescue virus NYVAC (vP866) (Tartaglia et al., (1992) Virology 188: 217–232). Selection of recombinant viruses was done on rabbit kidney cells.

Recombinant viruses, vP1192 and vP1195 were generated using insertion plasmids pES229A-6 and PSD.RN, respectively. To prepare plasmid pES229A-6 containing the F_{PIV-3} - F_{RSV} gene, the COPAK-H6 insertion plasmid pSD555 was cut with SmaI and dephosphorylated with intestinal alkaline

phosphatase. The 2.6 Kb F_{PIV-3} - F_{RSV} gene was retrieved from plasmid pES60A (Example 4) by cutting the plasmid with EcoRI and BamHI. The 2.6 Kb EcoRI-BamHI F_{PIV-3} - F_{RSV} gene was blunt ended with Klenow polymerase, isolated from a low melting point agarose gel and cloned into the SmaI site of the COPAK-H6 insertion plasmid pSD555 to generate plasmid pES229A-6. This positioned the F_{PIV-3} - F_{RSV} ORF such that the 5' end is nearest the H6 promoter.

To prepare plasmid PSD.RN, the pSD555 vector first was cut with SmaI and BamHI. Plasmid pM RF-HN (Example 9) containing the truncated F_{RSV} -HN $_{PIV-3}$ gene was cut with ClaI, blunt ended with Klenow polymerase and then cut with BamHI. The 3.3 Kb F_{RSV} -HN $_{PIV-3}$ gene was cloned into the SmaI-BamHI sites of the pSD555 vector to generate plasmid PSD.RN. This positioned the F_{RSV} -HN $_{PIV-3}$ ORF such that the H6 5' end is nearest the H6 promoter.

Plasmids pES229A-6 and PSD.RN were used in in vitro recombination experiments in vero cells with NYVAC (vP866) as the rescuing virus. Recombinant progeny virus was selected on rabbit kidney (RK)-13 cells (ATCC #CCL37). Several plaques were passaged two times on RK-13 cells. Virus containing the chimeric genes were confirmed by standard in situ plaque hybridization [Piccini et al. (1987), Methods in Enzymology, 153:545–563] using radiolabeled probes specific for the PIV and RSV inserted DNA sequences. Plaque purified virus containing the F_{PIV-3} - F_{RSV} and F_{RSV} -HN $_{PIV-3}$ chimeric genes were designated vP1192 and vP1195, respectively.

Radioimmunoprecipitation was done to confirm the expression of the chimeric genes in vP1192 and vP1195 infected cells. These assays were performed with lysates prepared from infected Vero cells [according to the procedure of Taylor et al., (1990) J. Virology 64, 1441–1450] using guinea pig monospecific PIV-3 anti-HN and anti-F antiserum and rabbit anti-RSV F antiserum. Both the anti-PIV F and anti-RSV F antisera precipitated a protein with an apparent molecular weight of approximately 90 kDa from vP1192 infected Vero cells. Both anti-RSV F and guinea pig anti-PIV HN antisera precipitated a protein with an apparent molecular weight of approximately 100 kDa from vP1195 infected cells. These results confirmed the production of the F_{PIV-3} - F_{RSV} and F_{RSV} -HN $_{PIV-3}$ chimeric proteins in Vero cells infected with the recombinant poxviruses.

SUMMARY OF DISCLOSURE

In summary of the disclosure, the present invention provides multimeric hybrid genes which produce chimeric proteins capable of eliciting protection against infection by a plurality of pathogens, particularly PIV and RSV. Modifications are possible within the scope of this invention.

TABLE 1

Secondary antibody response of guinea pigs immunized with the chimeric F_{RSV} -HN $_{PIV-3}$ protein							
Antigen	Dose	HAI Titre ^a (log ₂ ± s.e.)		Neutralization Titre ^b (log ₂ ± s.e.)			
		PIV-3		PIV-3		RSV	
Formulation	(ug)	6 wk Bleed	8 wk Bleed	6 wk Bleed	8 wk Bleed	6 wk Bleed	8 wk Bleed
Buffer	—	<1.0 ± 0.0	<1.0 ± 0.0	<1.0 ± 0.0	<1.0 ± 0.0	<1.0 ± 0.0	<1.0 ± 0.0
F_{RSV} -HN $_{PIV-3}$	10.0	9.1 ± 0.3	9.1 ± 0.3	7.1 ± 0.3	7.1 ± 0.5	5.5 ± 0.9	4.5 ± 1.2
	1.0	7.0 ± 2.0	7.3 ± 2.2	5.0 ± 1.5	4.5 ± 1.4	4.5 ± 0.5	3.0 ± 1.0
Live PIV-3		8.6 ± 0.7	7.3 ± 0.6	7.0 ± 0.4	7.3 ± 0.6	N/A	N/A
Live RSV		N/A ^c	N/A	N/A	N/A	5.5 ± 1.5	5.0 ± 1.0

^aReciprocal mean log₂ serum dilution which inhibits erythrocyte agglutination by 4 hemagglutinating units of PIV-3

^bReciprocal mean log₂ serum dilution which blocks hemadsorption of 100 TCID₅₀ units of PIV-3 or RSV

^cN/A — not applicable

TABLE 2

Serum antibody response of cotton rate immunized with the chimeric F_{RSV} -HN $_{PIV-3}$ protein ^a							
Antigen	Dose	HAI Titre ^b (log ₂ ± s.d.)		Neutralization Titre ^c (log ₂ ± s.d.)			
		PIV-3		PIV-3		RSV	
Formulation	(ug)	4 wk Bleed	5 wk Bleed	4 wk Bleed	5 wk Bleed	4 wk Bleed	5 wk Bleed
Buffer	—	2.8 ± 0.5	<3.0 ± 0.0	<1.0 ± 0.0	<1.0 ± 0.0	1.8 ± 0.3	0.8 ± 0.7
F_{RSV} -HN $_{PIV-3}$	10.0	9.5 ± 1.3	10.5 ± 0.6	>9.0 ± 0.0	>9.0 ± 0.0	5.2 ± 1.1	5.8 ± 0.9
	1.0	9.3 ± 1.0	10.3 ± 0.5	>9.0 ± 0.0	>9.0 ± 0.0	5.0 ± 0.7	5.8 ± 1.2
Live PIV-3		7.0 ± 0.0	8.5 ± 0.7	>9.0 ± 0.0	9.2 ± 0.7	N/A	N/A
Live RSV		N/A ^d	N/A	N/A	N/A	5.5 ± 0.6	8.5 ± 0.6

^aEach value represents the mean titre of antisera from 8 animlas.

^bReciprocal mean log₂ serum dilution which inhibits erythrocyte agglutination by 4 hemagglutinating units of PIV-3

^cReciprocal mean log₂ serum dilution which blocks hemadsorption of 100 ICID₅₀ units of PIV-3 or RSV

^dN/A — not applicable

TABLE 3

Response of immunized cotton rats to PIV/RSV challenge ^a				5
Antigen	Dose	Mean virus lung titre		
		log ₁₀ /g lung ± s.d.		
Formulation	(ug)	RSV	PIV-3	
Buffer	—	3.7 ± 0.3	3.4 ± 0.3	10
F _{RSV} -HN _{PIV-3}	10.0	≅1.5 ± 0.0	≅1.5 ± 0.0	
F _{RSV} -HN _{PIV-3}	1.0	≅1.5 ± 0.0	≅1.5 ± 0.0	
Live RSV		≅1.5 ± 0.0	N.D.	
Live PIV-3		N.D.	≅1.5 ± 0.0	

^aAnimals were challenged intranasally with 100 TCID₅₀ units of PIV-3 or RSV and killed 4 days later. Each value represents the mean virus lung titre of 4 animals.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 38

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1844 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

```

AAGTCAATAC CAACAACAT TAGCAGTCAT ACGTGCAAGA ACAAGAAAGA AGAGATTCAA      60
AAAGCTAAAT AAGAGAAATC AAAACAAAAG GTATAGAACA CCCGAACAAC AAAATCAAAA      120
CATCCAATCC ATTTTAAACA AAAATTCCAA AAGAGACCGG CAACACAACA AGCACCAAAC      180
ACAATGCCAA CTTTAATACT GCTAATTATT ACAACAATGA TTATGGCATC TTCCTGCCAA      240
ATAGATATCA CAAAACCTACA GCATGTAGGT GTATTGGTCA ACAGTCCCAA AGGGATGAAG      300
ATATCACAAA ACTTCGAAAC AAGATATCTA ATTTTGAGCC TCATACCAA AATAGAAGAC      360
TCTAACTCTT GTGGTGACCA ACAGATCAAA CAATACAAGA GGTTATTGGA TAGACTGATC      420
ATCCCTCTAT ATGATGGATT AAGATTACAG AAAGATGTGA TAGTAACCAA TCAAGAATCC      480
AATGAAAACA CTGATCCCAG AACAAAGACGA TCCTTTGGAG GGGTAATTGG AACCATGCT      540
CTGGGAGTAG CAACCTCAGC ACAAATTACA GCGGCAGTTG CTCTGGTTGA AGCCAAGCAG      600
GCAAAATCAG ACATCGAAAA ACTCAAAGAA GCAATCAGGG ACACAAACAA AGCAGTGCAG      660
TCAGTTCAGA GCTCTATAGG AAATTTAATA GTAGCAATTA AATCAGTCCA AGATTATGTC      720
AACAACGAAA TGGTGCCATC GATTGCTAGA CTAGGTTGTG AAGCAGCAGG ACTTCAATTA      780
GGAATTGCAT TAACACAGCA TTAACAGAA TTAACAAACA TATTTGGTGA TAACATAGGA      840
TCGTTACAAG AAAAAGGAAT AAAATTACAA GGTATAGCAT CATTATACCG CACAAATATC      900
ACAGAAATAT TCACAACATC AACAGTTGAT AAATATGATA TCTATGATCT ATTATTTACA      960
GAATCAATAA AGGTGAGAGT TATAGATGTT GATTTGAATG ATTACTCAAT CACCCTCCAA     1020
GTCAGACTCC CTTTATTAAC TAGGCTGCTG AACACTCAGA TCTACAAAGT AGATTCCATA     1080
TCATATAATA TCCAAAACAG AGAATGGTAT ATCCCTCTTC CCAGCCATAT CATGACGAAA     1140

```

-continued

GGGGCATTTC TAGGTGGAGC AGATGTCAAG GAATGTATAG AAGCATTTCAG CAGTTATATA 1200
 TGCCCTTCTG ATCCAGGATT TGTACTAAAC CATGAAATGG AGAGCTGCTT ATCAGGAAAC 1260
 ATATCCCAAT GTCCAAGAAC CACGGTCACA TCAGACATTG TTCCAAGATA TGCATTTGTC 1320
 AATGGAGGAG TGGTTGCAAA CTGTATAACA ACCACCTGTA CATGCAACGG AATCGACAAT 1380
 AGAATCAATC AACCACCTGA TCAAGGAGTA AAAATTATAA CACATAAAGA ATGTAATACA 1440
 ATAGGTATCA ACGGAATGCT GTTCAATACA AATAAAGAAG GAACTCTTGC ATTCTACACA 1500
 CCAAATGATA TAACACTAAA TAATTCTGTT GCACTTGATC CAATTGACAT ATCAATCGAG 1560
 CTTAACAAAG CCAAATCAGA TCTAGAAGAA TCAAAAGAAT GGATAAGAAG GTCAAATCAA 1620
 AAAC TAGATT CTATTGAAA CTGGCATCAA TCTAGCACTA CAATCATAAT TATTTTAATA 1680
 ATGATCATT TATTGTTTAT AATTAATGTA ACGATAATTA CAATTGCAAT TAAGTATTAC 1740
 AGAATTCAAA AGAGAAATCG AGTGGATCAA AATGACAAGC CATATGTAAT ACAAACAAA 1800
 TGACATATCT ATAGATCATT AGATATTAATA ATTATAAAAA ACTT 1844

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 539 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Met Pro Thr Leu Ile Leu Leu Ile Ile Thr Thr Met Ile Met Ala Ser
 1 5 10 15
 Ser Cys Gln Ile Asp Ile Thr Lys Leu Gln His Val Gly Val Leu Val
 20 25 30
 Asn Ser Pro Lys Gly Met Lys Ile Ser Gln Asn Phe Glu Thr Arg Tyr
 35 40 45
 Leu Ile Leu Ser Leu Ile Pro Lys Ile Glu Asp Ser Asn Ser Cys Gly
 50 55 60
 Asp Gln Gln Ile Lys Gln Tyr Lys Arg Leu Leu Asp Arg Leu Ile Ile
 65 70 75 80
 Pro Leu Tyr Asp Gly Leu Arg Leu Gln Lys Asp Val Ile Val Thr Asn
 85 90 95
 Gln Glu Ser Asn Glu Asn Thr Asp Pro Arg Thr Arg Arg Ser Phe Gly
 100 105 110
 Gly Val Ile Gly Thr Ile Ala Leu Gly Val Ala Thr Ser Ala Gln Ile
 115 120 125
 Thr Ala Ala Val Ala Leu Val Glu Ala Lys Gln Ala Lys Ser Asp Ile
 130 135 140
 Glu Lys Leu Lys Glu Ala Ile Arg Asp Thr Asn Lys Ala Val Gln Ser
 145 150 155 160
 Val Gln Ser Ser Ile Gly Asn Leu Ile Val Ala Ile Lys Ser Val Gln
 165 170 175
 Asp Tyr Val Asn Asn Glu Ile Val Pro Ser Ile Ala Arg Leu Gly Cys
 180 185 190
 Glu Ala Ala Gly Leu Gln Leu Gly Ile Ala Leu Thr Gln His Tyr Ser
 195 200 205
 Glu Leu Thr Asn Ile Phe Gly Asp Asn Ile Gly Ser Leu Gln Glu Lys
 210 215 220
 Gly Ile Lys Leu Gln Gly Ile Ala Ser Leu Tyr Arg Thr Asn Ile Thr
 225 230 235 240

-continued

Glu Ile Phe Thr Thr Ser Thr Val Asp Lys Tyr Asp Ile Tyr Asp Leu
245 250 255

Leu Phe Thr Glu Ser Ile Lys Val Arg Val Ile Asp Val Asp Leu Asn
260 265 270

Asp Tyr Ser Ile Thr Leu Gln Val Arg Leu Pro Leu Leu Thr Arg Leu
275 280 285

Leu Asn Thr Gln Ile Tyr Lys Val Asp Ser Ile Ser Tyr Asn Ile Gln
290 295 300

Asn Arg Glu Trp Tyr Ile Pro Leu Pro Ser His Ile Met Thr Lys Gly
305 310 315 320

Ala Phe Leu Gly Gly Ala Asp Val Lys Glu Cys Ile Glu Ala Phe Ser
325 330 335

Ser Tyr Ile Cys Pro Ser Asp Pro Gly Phe Val Leu Asn His Glu Met
340 345 350

Glu Ser Cys Leu Ser Gly Asn Ile Ser Gln Cys Pro Arg Thr Thr Val
355 360 365

Thr Ser Asp Ile Val Pro Arg Tyr Ala Phe Val Asn Gly Gly Val Val
370 375 380

Ala Asn Cys Ile Thr Thr Thr Cys Thr Cys Asn Gly Ile Asp Asn Arg
385 390 395 400

Ile Asn Gln Pro Pro Asp Gln Gly Val Lys Ile Ile Thr His Lys Glu
405 410 415

Cys Asn Thr Ile Gly Ile Asn Gly Met Leu Phe Asn Thr Asn Lys Glu
420 425 430

Gly Thr Leu Ala Phe Tyr Thr Pro Asn Asp Ile Thr Leu Asn Asn Ser
435 440 445

Val Ala Leu Asp Pro Ile Asp Ile Ser Ile Glu Leu Asn Lys Ala Lys
450 455 460

Ser Asp Leu Glu Glu Ser Lys Glu Trp Ile Arg Arg Ser Asn Gln Lys
465 470 475 480

Leu Asp Ser Ile Gly Asn Trp His Gln Ser Ser Thr Thr Ile Ile Ile
485 490 495

Ile Leu Ile Met Ile Ile Ile Leu Phe Ile Ile Asn Val Thr Ile Ile
500 505 510

Thr Ile Ala Ile Lys Tyr Tyr Arg Ile Gln Lys Arg Asn Arg Val Asp
515 520 525

Gln Asn Asp Lys Pro Tyr Val Leu Thr Asn Lys
530 535

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1833 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

AGACAAATCC AAATTCGAGA TGAATACTG GAAGCATACC AATCACGGAA AGGATGCTGG 60

CAATGAGCTG GAGACGTCCA TGGCTACTAA TGGCAACAAG CTCACCAATA AGATAACATA 120

TATATTATGG ACAATAATCC TGGTGTATT ATCAATAGTC TTCATCATAG TGCTAATTAA 180

TTCCATCAAA AGTGAAAAGG CTCATGAATC ATTGCTGCAA GACATAAATA ATGAGTTTAT 240

GGAAATTACA GAAAAGATCC AAATGGCATC GGATAATACC AATGATCTAA TACAGTCAGG 300

AGTGAATACA AGGCTTCTTA CAATTCAGAG TCATGTCCAG AATTATATAC CAATATCACT 360

-continued

GACACAACAG ATGTCAGATC TTAGGAAATT CATTAGTGAA ATTACAATTA GAAATGATAA 420
TCAAGAAGTG CTGCCACAAA GAATAACACA TGATGTGGGT ATAAAACCTT TAAATCCAGA 480
TGATTTTTGG AGATGCACGT CTGGTCTTCC ATCTTTAATG AAAACTCCAA AAATAAGGTT 540
AATGCCAGGG CCGGGATTAT TAGCTATGCC AACGACTGTT GATGGCTGTA TCAGAACTCC 600
GTCCTTAGTT ATAAATGATC TGATTTATGC TTATACCTCA AATCTAATTA CTCGAGGTTG 660
TCAGGATATA GGAAAATCAT ATCAAGTCTT ACAGATAGGG ATAATAACTG TAAACTCAGA 720
CTTGGTACCT GACTTAAATC CCAGGATCTC TCATACTTTT AACATAAATG ACAATAGGAA 780
GTCATGTTCT CTAGCACTCC TAAATACAGA TGTATATCAA CTGTGTTCAA CTCCCAAAGT 840
TGATGAAAGA TCAGATTATG CATCATCAGG CATAGAAGAT ATTGTACTTG ATATTGTCAA 900
TTATGATGGC TCAATCTCAA CAACAAGATT TAAGAATAAT AACATAAGCT TTGATCAACC 960
TTATGCTGCA CTATACCCAT CTGTTGGACC AGGGATATAC TACAAAGGCA AAATAATATT 1020
TCTCGGGTAT GGAGGTCTTG AACATCCAAT AAATGAGAAT GTAATCTGCA ACACAACCTGG 1080
GTGTCCCGGG AAAACACAGA GAGACTGCAA TCAGGCATCT CATAGTCCAT GGTTTTCAGA 1140
TAGGAGGATG GTCAACTCTA TCATTGTTGT TGACAAAGGC TTAAACTCAA TTCCAAAATT 1200
GAAGGTATGG ACGATATCTA TGAGACAGAA TTACTGGGGG TCAGAAGGAA GGTACTTCT 1260
ACTAGGTAAC AAGATCTATA TATATACAAG ATCCACAAGT TGGCATAGCA AGTTACAATT 1320
AGGAATAATT GATATTACTG ATTACAGTGA TATAAGGATA AAATGGACAT GGCATAATGT 1380
GCTATCAAGA CCAGGAAACA ATGAATGTCC ATGGGGACAT TCATGTCCAG ATGGATGTAT 1440
AACAGGAGTA TATACTGATG CATATCCACT CAATCCCACA GGGAGCATTG TGTCATCTGT 1500
CATATTAGAT TCACAAAAAT CGAGAGTGAA CCCAGTCATA ACTTACTCAA CAGCAACCGA 1560
AAGAGTAAAC GAGCTGGCCA TCCGAAACAG AACACTCTCA GCTGGATATA CAACAACAAG 1620
CTGCATCACA CACTATAACA AAGGATATTG TTTTCATATA GTAGAAATAA ATCAGAAAAG 1680
CTTAAACACA CTTCAACCCA TGTGTTCAA GACAGAGGTT CCAAAAAGCT GCAGTTAATC 1740
ATAATTAACC GCAATATGCA TTAACCTATC TATAATACAA GTATATGATA AGTAATCAGC 1800
AATCAGACAA TAGACAAAAG GGAAATATAA AAA 1833

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 572 amino acids
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Met Glu Tyr Trp Lys His Thr Asn His Gly Lys Asp Ala Gly Asn Glu
1 5 10 15
Leu Glu Thr Ser Met Ala Thr Asn Gly Asn Lys Leu Thr Asn Lys Ile
20 25 30
Thr Tyr Ile Leu Trp Thr Ile Ile Leu Val Leu Leu Ser Ile Val Phe
35 40 45
Ile Ile Val Leu Ile Asn Ser Ile Lys Ser Glu Lys Ala His Glu Ser
50 55 60
Leu Leu Gln Asp Ile Asn Asn Glu Phe Met Glu Ile Thr Glu Lys Ile
65 70 75 80
Gln Met Ala Ser Asp Asn Thr Asn Asp Leu Ile Gln Ser Gly Val Asn
85 90 95

-continued

Thr	Arg	Leu	Leu	Thr	Ile	Gln	Ser	His	Val	Gln	Asn	Tyr	Ile	Pro	Ile
			100					105					110		
Ser	Leu	Thr	Gln	Gln	Met	Ser	Asp	Leu	Arg	Lys	Phe	Ile	Ser	Glu	Ile
		115					120					125			
Thr	Ile	Arg	Asn	Asp	Asn	Gln	Glu	Val	Leu	Pro	Gln	Arg	Ile	Thr	His
	130					135					140				
Asp	Val	Gly	Ile	Lys	Pro	Leu	Asn	Pro	Asp	Asp	Phe	Trp	Arg	Cys	Thr
145					150					155					160
Ser	Gly	Leu	Pro	Ser	Leu	Met	Lys	Thr	Pro	Lys	Ile	Arg	Leu	Met	Pro
				165					170					175	
Gly	Pro	Gly	Leu	Leu	Ala	Met	Pro	Thr	Thr	Val	Asp	Gly	Cys	Ile	Arg
			180					185					190		
Thr	Pro	Ser	Leu	Val	Ile	Asn	Asp	Leu	Ile	Tyr	Ala	Tyr	Thr	Ser	Asn
		195					200					205			
Leu	Ile	Thr	Arg	Gly	Cys	Gln	Asp	Ile	Gly	Lys	Ser	Tyr	Gln	Val	Leu
	210					215						220			
Gln	Ile	Gly	Ile	Ile	Thr	Val	Asn	Ser	Asp	Leu	Val	Pro	Asp	Leu	Asn
225					230					235					240
Pro	Arg	Ile	Ser	His	Thr	Phe	Asn	Ile	Asn	Asp	Asn	Arg	Lys	Ser	Cys
				245					250					255	
Ser	Leu	Ala	Leu	Leu	Asn	Thr	Asp	Val	Tyr	Gln	Leu	Cys	Ser	Thr	Pro
			260					265					270		
Lys	Val	Asp	Glu	Arg	Ser	Asp	Tyr	Ala	Ser	Ser	Gly	Ile	Glu	Asp	Ile
		275					280					285			
Val	Leu	Asp	Ile	Val	Asn	Tyr	Asp	Gly	Ser	Ile	Ser	Thr	Thr	Arg	Phe
	290					295					300				
Lys	Asn	Asn	Asn	Ile	Ser	Phe	Asp	Gln	Pro	Tyr	Ala	Ala	Leu	Tyr	Pro
305					310					315					320
Ser	Val	Gly	Pro	Gly	Ile	Tyr	Tyr	Lys	Gly	Lys	Ile	Ile	Phe	Leu	Gly
				325					330					335	
Tyr	Gly	Gly	Leu	Glu	His	Pro	Ile	Asn	Glu	Asn	Val	Ile	Cys	Asn	Thr
			340					345					350		
Thr	Gly	Cys	Pro	Gly	Lys	Thr	Gln	Arg	Asp	Cys	Asn	Gln	Ala	Ser	His
		355					360					365			
Ser	Pro	Trp	Phe	Ser	Asp	Arg	Arg	Met	Val	Asn	Ser	Ile	Ile	Val	Val
		370				375					380				
Asp	Lys	Gly	Leu	Asn	Ser	Ile	Pro	Lys	Leu	Lys	Val	Trp	Thr	Ile	Ser
385					390					395					400
Met	Arg	Gln	Asn	Tyr	Trp	Gly	Ser	Glu	Gly	Arg	Leu	Leu	Leu	Leu	Gly
			405						410					415	
Asn	Lys	Ile	Tyr	Ile	Tyr	Thr	Arg	Ser	Thr	Ser	Trp	His	Ser	Lys	Leu
			420					425					430		
Gln	Leu	Gly	Ile	Ile	Asp	Ile	Thr	Asp	Tyr	Ser	Asp	Ile	Arg	Ile	Lys
		435					440					445			
Trp	Thr	Trp	His	Asn	Val	Leu	Ser	Arg	Pro	Gly	Asn	Asn	Glu	Cys	Pro
	450					455					460				
Trp	Gly	His	Ser	Cys	Pro	Asp	Gly	Cys	Ile	Thr	Gly	Val	Tyr	Thr	Asp
465					470					475					480
Ala	Tyr	Pro	Leu	Asn	Pro	Thr	Gly	Ser	Ile	Val	Ser	Ser	Val	Ile	Leu
				485					490					495	
Asp	Ser	Gln	Lys	Ser	Arg	Val	Asn	Pro	Val	Ile	Thr	Tyr	Ser	Thr	Ala
			500					505					510		
Thr	Glu	Arg	Val	Asn	Glu	Leu	Ala	Ile	Arg	Asn	Arg	Thr	Leu	Ser	Ala
							520						525		

-continued

Gly Tyr Thr Thr Thr Ser Cys Ile Thr His Tyr Asn Lys Gly Tyr Cys
 530 535 540

Phe His Ile Val Glu Ile Asn Gln Lys Ser Leu Asn Thr Leu Gln Pro
 545 550 555 560

Met Leu Phe Lys Thr Glu Val Pro Lys Ser Cys Ser
 565 570

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1886 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

ATGGAGTTGC CAATCCTCAA AGCAAATGCA ATTACCACAA TCCTCGCTGC AGTCACATTT 60

TGCTTTGCTT CTAGTCAAAA CATCACTGAA GAATTTTATC AATCAACATG CAGTGCAGTT 120

AGCAAAGGCT ATCTTAGTGC TCTAAGAACT GGTGGGTATA CTAGTGTTAT AACTATAGAA 180

TTAAGTAATA TCAAGGAAAA TAAGTGTAAT GGAACAGATG CTAAGGTAAA ATTGATGAAA 240

CAAGAATTAG ATAAATATAA AAATGCTGTA ACAGAATTGC AGTTGCTCAT GCAAAGCACA 300

CCAGCAGCAA ACAATCGAGC CAGAAGAGAA CTACCAAGGT TTATGAATTA TACACTCAAC 360

AATACCAAAA AAACCAATGT AACATTAAGC AAGAAAAGGA AAAGAAGATT TCTTGTTTTT 420

TTGTTAGGTG TTGGATCTGC AATCGCCAGT GGCATTGCTG TATCTAAGGT CCTGCACTTA 480

GAAGGAGAAG TGAACAAGAT CAAAAGTGCT CTACTATCCA CAAACAAGGC CGTAGTCAGC 540

TTATCAAATG GAGTTAGTGT CTTAACCAGC AAAGTGTTAG ACCTCAAAAA CTATATAGAT 600

AAACAATTGT TACCTATTGT GAATAAGCGA AGCTGCAGAA TATCAAATAT AGAAACTGTG 660

ATAGAGTTCC AACAAAAGAA CAACAGACTA CTAGAGATTA CCAGGGAATT TAGTGTTAAT 720

GCAGGTGTAA CTACACCTGT AAGCACTTAC ATGTTAACTA ATAGTGAATT ATTGTCATTA 780

ATCAATGATA TGCCTATAAC AAATGATCAG AAAAAGTTAA TGTCCAACAA TGTTCAAATA 840

GTTAGACAGC AAAGTTACTC TATCATGTCC ATAATAAAAG AGGAAGTCTT AGCATATGTA 900

GTACAATTAC CACTATATGG TGTGATAGAT ACACCTTGTT GGAAATTACA CACATCCCCT 960

CTATGTACAA CCAACACAAA AGAAGGGTCA AACATCTGTT TAACAAGAAC TGACAGAGGA 1020

TGGTACTGTG ACAATGCAGG ATCAGTATCT TTCTTCCCAC AAGCTGAAAC ATGTAAAGTT 1080

CAATCGAATC GAGTATTTTG TGACACAATG AACAGTTTAA CATTACCAAG TGAAGTAAAT 1140

CTCTGCAATG TTGACATATT CAATCCCAAA TATGATTGTA AAATTATGAC TTCAAAAACA 1200

GATGTAAGCA GCTCCGTTAT CACATCTCTA GGAGCCATTG TGTCATGCTA TGGCAAAACT 1260

AAATGTACAG CATCCAATAA AAATCGTGGA ATCATAAAGA CATTTTCTAA CGGGTGTGAT 1320

TATGTATCAA ATAAAGGGGT GGACACTGTG TCTGTAGGTA ACACATTATA TTATGTAAAT 1380

AAGCAAGAAG GCAAAAAGTCT CTATGTAAAA GGTGAACCAA TAATAAATTT CTATGACCCA 1440

TTAGTATTCC CCTCTGATGA ATTTGATGCA TCAATATCTC AAGTCAATGA GAAGATTAAC 1500

CAGAGTTTAG CATTTATTCG TAAATCCGAT GAATTATTAC ATAATGTAAA TGCTGGTAAA 1560

TCAACCACAA ATATCATGAT AACTACTATA ATTATAGTGA TTATAGTAAT ATTGTTATCA 1620

TTAATTGCTG TTGGACTGCT CCTATACTGT AAGGCCAGAA GCACACCAGT CACACTAAGC 1680

AAGGATCAAC TGAGTGGTAT AAATAATATT GCATTTAGTA ACTGAATAAA AATAGCACCT 1740

AATCATGTTT TTACAATGGT TTAATATCTG CTCATAGACA ACCCATCTAT CATTGGATTT 1800

-continued

TCTTAAAATC TGAACCTTCAT CGAAACTCTT ATCTATAAAC CATCTCACTT ACACTATTTA 1860
 AGTAGATTCC TAGTTTATAG TTATAT 1886

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 594 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Met Glu Leu Pro Ile Leu Lys Ala Asn Ala Ile Thr Thr Ile Leu Ala
 1 5 10 15
 Ala Val Thr Phe Cys Phe Ala Ser Ser Gln Asn Ile Thr Glu Glu Phe
 20 25 30
 Tyr Gln Ser Thr Cys Ser Ala Val Ser Lys Gly Tyr Leu Ser Ala Leu
 35 40 45
 Arg Thr Gly Trp Tyr Thr Ser Val Ile Thr Ile Glu Leu Ser Asn Ile
 50 55 60
 Lys Glu Asn Lys Cys Asn Gly Thr Asp Ala Lys Val Lys Leu Met Lys
 65 70 75 80
 Gln Glu Leu Asp Lys Tyr Lys Asn Ala Val Thr Glu Leu Gln Leu Leu
 85 90 95
 Met Gln Ser Thr Pro Ala Ala Asn Asn Arg Ala Arg Arg Glu Leu Pro
 100 105 110
 Arg Phe Met Asn Tyr Thr Leu Asn Asn Thr Lys Lys Thr Asn Val Thr
 115 120 125
 Leu Ser Lys Lys Arg Lys Arg Arg Phe Leu Gly Phe Leu Leu Gly Val
 130 135 140
 Gly Ser Ala Ile Ala Ser Gly Ile Ala Val Ser Lys Val Leu His Leu
 145 150 155 160
 Glu Gly Glu Val Asn Lys Ile Lys Ser Ala Leu Leu Ser Thr Asn Lys
 165 170 175
 Ala Val Val Ser Leu Ser Asn Gly Val Ser Val Leu Thr Ser Lys Val
 180 185 190
 Leu Asp Leu Lys Asn Tyr Ile Asp Lys Gln Leu Leu Pro Ile Val Asn
 195 200 205
 Lys Arg Ser Cys Arg Ile Ser Asn Ile Glu Thr Val Ile Glu Phe Gln
 210 215 220
 His Lys Asn Asn Arg Leu Leu Glu Ile Thr Arg Glu Phe Ser Val Asn
 225 230 235 240
 Ala Gly Val Thr Thr Pro Val Ser Thr Tyr Met Leu Thr Asn Ser Glu
 245 250 255
 Leu Leu Ser Leu Ile Asn Asp Met Pro Ile Thr Asn Asp Gln Lys Lys
 260 265 270
 Leu Met Ser Asn Asn Val Gln Ile Val Arg Gln Gln Ser Tyr Ser Ile
 275 280 285
 Met Ser Ile Ile Lys Glu Glu Val Leu Ala Tyr Val Val Gln Leu Pro
 290 295 300
 Leu Tyr Gly Val Ile Asp Thr Pro Cys Trp Lys Leu His Thr Ser Pro
 305 310 315 320
 Leu Cys Thr Thr Asn Thr Lys Glu Gly Ser Asn Ile Cys Leu Thr Arg
 325 330 335
 Thr Asp Arg Gly Trp Tyr Cys Asp Asn Ala Gly Ser Val Ser Phe Phe

-continued

340	345	350
Pro Gln Ala Glu Thr Cys Lys Val Gln Ser Asn Arg Val Phe Cys Asp 355	360	365
Thr Met Asn Ser Leu Thr Leu Pro Ser Glu Val Asn Leu Cys Asn Val 370	375	380
Asp Ile Phe Asn Pro Lys Tyr Asp Cys Lys Ile Met Thr Ser Lys Thr 385	390	395
Asp Val Ser Ser Ser Val Ile Thr Ser Leu Gly Ala Ile Val Ser Cys 405	410	415
Tyr Gly Lys Thr Lys Cys Thr Ala Ser Asn Lys Asn Arg Gly Ile Ile 420	425	430
Lys Thr Phe Ser Asn Gly Cys Asp Tyr Val Ser Asn Lys Gly Val Asp 435	440	445
Thr Val Ser Val Gly Asn Thr Leu Tyr Tyr Val Asn Lys Gln Glu Gly 450	455	460
Lys Ser Leu Tyr Val Lys Gly Glu Pro Ile Ile Asn Phe Tyr Asp Pro 465	470	475
Leu Val Phe Pro Ser Asp Glu Phe Asp Ala Ser Ile Ser Gln Val Asn 485	490	495
Glu Lys Ile Asn Leu Val Phe Pro Ser Asp Glu Phe Asp Ala Ser Ile 500	505	510
Ser Gln Val Asn Glu Lys Ile Asn Gln Ser Leu Ala Phe Ile Arg Lys 515	520	525
Ser Asp Glu Leu Leu His Asn Val Asn Ala Gly Lys Ser Thr Thr Asn 530	535	540
Ile Met Ile Thr Thr Ile Ile Glu Ile Ile Val Ile Leu Leu Ser 545	550	555
Leu Ile Ala Val Gly Leu Leu Leu Tyr Cys Lys Ala Arg Ser Thr Pro 565	570	575
Val Thr Leu Ser Lys Asp Gln Leu Ser Gly Ile Asn Asn Ile Ala Phe 580	585	590
Ser Asn		

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 920 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

TGCAAACATG TCCAAAAACA AGGACCAACG CACCGCTAAG AACTAGAAA AGACCTGGGA	60
CACTCTCAAT CATTATTAT TCATATCATC GGGCTTATAT AAGTTAAATC TTAAATCTGT	120
AGCACAAATC ACATTATCCA TTCTGGCAAT GATAATCTCA ACTTCACTTA TAATTACAGC	180
CATCATATTC ATAGCCTCGG CAAACCACAA AGTCACACTA ACAACTGCAA TCATACAAGA	240
TGCAACAAGC CAGATCAAGA ACACAACCCC AACATACCTC ACTCAGGATC CTCAGCTTGG	300
AATCAGCTTC TCCAATCTGT CTGAAATTAC ATCACAAACC ACCACCATAC TAGCTTCAAC	360
AACACCAGGA GTCAAGTCAA ACCTGCAACC CACAACAGTC AAGACTAAAA ACACAACAAC	420
AACCCAAACA CAACCCAGCA AGCCCACTAC AAAACAACGC CAAAACAAAC CACCAAACAA	480
ACCCAATAAT GATTTTCACT TCGAAGTGTT TAACTTTGTA CCCTGCAGCA TATGCAGCAA	540
CAATCCAACC TGCTGGGCTA TCTGCAAAAAG AATACCAAAC AAAAAACCAG GAAAGAAAAC	600

-continued

CACCACCAAG CCTACAAAA AACCAACCTT CAAGACAACC AAAAAAGATC TCAAACCTCA 660
 AACCACTAAA CCAAAGGAAG TACCCACCAC CAAGCCCACA GAAGAGCCAA CCATCAACAC 720
 CACCAAAACA AACATCACAA CTACTGTGCT CACCAACAAC ACCACAGGAA ATCCAAAACCT 780
 CACAAGTCAA ATGGAAACCT TCCACTCAAC CTCCTCCGAA GGCAATCTAA GCCCTTCTCA 840
 AGTCTCCACA ACATCCGAGC ACCCATCACA ACCCTCATCT CCACCCAACA CAACAGCCA 900
 GTAGTTATTA AAAAAAAAAA 920

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 298 amino acids
 (B) TYPE: amino acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Met Ser Lys Asn Lys Asp Gln Arg Thr Ala Lys Thr Leu Glu Lys Thr
 1 5 10 15
 Trp Asp Thr Leu Asn His Leu Leu Phe Ile Ser Ser Gly Leu Tyr Lys
 20 25 30
 Leu Asn Leu Lys Ser Val Ala Gln Ile Thr Leu Ser Ile Leu Ala Met
 35 40 45
 Ile Ile Ser Thr Ser Leu Ile Ile Thr Ala Ile Ile Phe Ile Ala Ser
 50 55 60
 Ala Asn His Lys Val Thr Leu Thr Thr Ala Ile Ile Gln Asp Ala Thr
 65 70 75 80
 Ser Gln Ile Lys Asn Thr Thr Pro Thr Tyr Leu Thr Gln Asp Pro Gln
 85 90 95
 Leu Gly Ile Ser Phe Ser Asn Leu Ser Glu Ile Thr Ser Gln Thr Thr
 100 105 110
 Thr Ile Leu Ala Ser Thr Thr Pro Gly Val Lys Ser Asn Leu Gln Pro
 115 120 125
 Thr Thr Val Lys Thr Lys Asn Thr Thr Thr Thr Gln Thr Gln Pro Ser
 130 135 140
 Lys Pro Thr Thr Lys Gln Arg Gln Asn Lys Pro Pro Asn Lys Pro Asn
 145 150 155 160
 Asn Asp Phe His Phe Glu Val Phe Asn Phe Val Pro Cys Ser Ile Cys
 165 170 175
 Ser Asn Asn Pro Thr Cys Trp Ala Ile Cys Lys Arg Ile Pro Asn Lys
 180 185 190
 Lys Pro Gly Lys Lys Thr Thr Thr Lys Pro Thr Lys Lys Pro Thr Phe
 195 200 205
 Lys Thr Thr Lys Lys Asp Leu Lys Pro Gln Thr Thr Lys Pro Lys Glu
 210 215 220
 Val Pro Thr Thr Lys Pro Thr Glu Glu Pro Thr Ile Asn Thr Thr Lys
 225 230 235 240
 Thr Asn Ile Thr Thr Thr Leu Leu Thr Asn Asn Thr Thr Gly Asn Pro
 245 250 255
 Lys Leu Thr Ser Gln Met Glu Thr Phe His Ser Thr Ser Ser Glu Gly
 260 265 270
 Asn Leu Ser Pro Ser Gln Val Ser Thr Thr Ser Glu His Pro Ser Gln
 275 280 285
 Pro Ser Ser Pro Pro Asn Thr Thr Arg Gln
 290 295

-continued

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 26 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATCAATCAAA GGCCTGTGA TAATAG 26

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CATGACTGA TAATGAG 17

(2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 86 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

AATTCATGGA GTTGCTAATC CTCAAAGCAA ATGCAATTAC CACAATCCTC ACTGCAGTCA 60

CATTTTGT TT TGCTTCTGGT TCTAAG 86

(2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

ACTGGCATCA ATCTAGCACT ACATGAG 27

(2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

AATTCATGCC AACTTTAATA CTGCTAATTA TTACAACAAT GATTATGGCA TCTTCCTGCC 60

AAATAGATAT CACAAAATA CAGCATGTAG GTGTATTGGT CAACAGTCCC AAAGGGATGA 120

AGATATCACA AACTT 136

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 94 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

ATCATGGAGA TAATTAAAAT GATAACCATC TCGCAAATAA ATAAGTATTT TACTGTTTTC 60

GTAACAGTTT TGTAATAAAA AACCTATAA ATAG 94

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 141 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

ATCATGGAGA TAATTAAAAT GATAACCATC TCGCAAATAA ATAAGTATTT TACTGTTTTC 60

GTAACAGTTT TGTAATAAAA AACCTATAA ATATTCCGGA ATTCAGATCT GCAGCGGCCG 120

CTCCATCTAG AAGGTACCCG G 141

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 31 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CATGACTAAT TCCATCAAAA GTGAAAAGGC T 31

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CAAGAAAAG GAATAAAA 18

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 39 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

ATTTCTGTGA TATTTGTGCG GTATAATGAT GCTATACCT 39

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 18 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CAGGAGAAGG GTATCAAG 18

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 17 base pairs

-continued

(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:
AGGAGAAGGG TATCAAG 17

(2) INFORMATION FOR SEQ ID NO:21:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 94 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:
ATCATGGAGA TAATTAAAT GATAACCATC TCGCAAATAA ATAAGTATTT TACTGTTTTTC 60
GTAACAGTTT TGTAATAAAA AAACCTATAA ATAG 94

(2) INFORMATION FOR SEQ ID NO:22:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:
Gln Glu Lys Gly Ile Lys
1 5

(2) INFORMATION FOR SEQ ID NO:23:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 6 amino acids
(B) TYPE: amino acid
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:
Gln Glu Lys Gly Ile Lys
1 5

(2) INFORMATION FOR SEQ ID NO:24:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 20 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:
ATCAATCTAG CACTACACAG 20

(2) INFORMATION FOR SEQ ID NO:25:
(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 1617 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:
ATGCCAACTT TAATACTGCT AATTATTACA ACAATGATTA TGGCATCTTC CTGCCAAATA 60
GATATCACAA AACTACAGCA TGTAGGTGTA TTGGTCAACA GTCCCAAAGG GATGAAGATA 120
TCACAAAAC TCGAAACAAG ATATCTAATT TTGAGCCTCA TACCAAAAT AGAAGACTCT 180
AACTCTGTG GTGACCAACA GATCAAACA TACAAGAGGT TATTGGATAG ACTGATCATC 240

-continued

CCTCTATATG ATGGATTAAG ATTACAGAAA GATGTGATAG TAACCAATCA AGAATCCAAT	300
GAAAACACTG ATCCCAGAAC AAGACGATCC TTTGGAGGGG TAATTGGAAC CATTGCTCTG	360
GGAGTAGCAA CCTCAGCACA AATTACAGCG GCAGTTGCTC TGGTTGAAGC CAAGCAGGCA	420
AAATCAGACA TCGAAAAACT CAAAGAAGCA ATCAGGGACA CAAACAAAGC AGTGCAGTCA	480
GTTTCAGAGCT CTATAGGAAA TTTAATAGTA GCAATTAAAT CAGTCCAAGA TTATGTCAAC	540
AACGAAATGG TGCCATCGAT TGCTAGACTA GGTGTGAAG CAGCAGGACT TCAATTAGGA	600
ATTGCATTAA CACAGCATT A CTCAGAATTA ACAAACATAT TTGGTGATAA CATAGGATCG	660
TTACAAGAAA AAGGAATAAA ATTACAAGGT ATAGCATCAT TATACCGCAC AAATATCACA	720
GAAATATTCA CAACATCAAC AGTTGATAAAA TATGATATCT ATGATCTATT ATTTACAGAA	780
TCAATAAAGG TGAGAGTTAT AGATGTTGAT TTGAATGATT ACTCAATCAC CCTCCAAGTC	840
AGACTCCCTT TATTAAGTAG GCTGCTGAAC ACTCAGATCT ACAAAGTAGA TTCCATATCA	900
TATAATATCC AAAACAGAGA ATGGTATATC CCTCTTCCCA GCCATATCAT GACGAAAGGG	960
GCATTTCTAG GTGGAGCAGA TGTCAAGGAA TGTATAGAAG CATTTCAGCAG TTATATATGC	1020
CCTTCTGATC CAGGATTTGT ACTAAACCAT GAAATGGAGA GCTGCTTATC AGGAAACATA	1080
TCCCAATGTC CAAGAACCAC GGTACATCA GACATTGTTC CAAGATATGC ATTTGTCAAT	1140
GGAGGAGTGG TTGCAAACCTG TATAACAACC ACCTGTACAT GCAACGGAAT CGACAATAGA	1200
ATCAATCAAC CACCTGATCA AGGAGTAAAA ATTATAACAC ATAAAGAATG TAATACAATA	1260
GGTATCAACG GAATGCTGTT CAATACAAAT AAAGAAGGAA CTCTTGCATT CTACACACCA	1320
AATGATATAA CACTAAATAA TTCTGTTGCA CTTGATCCAA TTGACATATC AATCGAGCTT	1380
AACAAAGCCA AATCAGATCT AGAAGAATCA AAAGAATGGA TAAGAAGGTC AAATCAAAAA	1440
CTAGATTCTA TTGGAAACTG GCATCAATCT AGCACTACAA TCATAATTAT TTTAATAATG	1500
ATCATTATAT TGTTTATAAT TAATGTAACG ATAATTACAA TTGCAATTAA GTATTACAGA	1560
ATTCAAAAGA GAAATCGAGT GGATCAAAAT GACAAGCCAT ATGTACTAAC AAACAAA	1617

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1715 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

ATGGAATACT GGAAGCATA CAATCACGGA AAGGATGCTG GCAATGAGCT GGAGACGTCC	60
ATGGCTACTA ATGGCAACAA GCTCACCAAT AAGATAACAT ATATATTATG GACAATAATC	120
CTGGTGTAT TATCAATAGT CTTTCATCATA GTGCTAATTA ATTCCATCAA AAGTGAAAAG	180
GCTCATGAAT CATTGCTGCA AGACATAAAT AATGAGTTTA TGGAAATTAC AGAAAAGATC	240
CAAATGGCAT CGGATAATAC CAATGATCTA ATACAGTCAG GAGTGAATAC AAGGCTTCTT	300
ACAATTCAGA GTCATGTCCA GAATTATATA CCAATATCAC TGACACAACA GATGTCAGAT	360
CTTAGGAAAT TCATTAGTGA AATTACAATT AGAAATGATA ATCAAGAAGT GCTGCCACAA	420
AGAATAACAC ATGATGTGGG TATAAAACCT TTAAATCCAG ATGATTTTTG GAGATGCACG	480
TCTGGTCTTC CATCTTTAAT GAAAACCTCA AAAATAAGGT TAATGCCAGG GCCGGGATTA	540
TTAGCTATGC CAACGACTGT TGATGGCTGT ATCAGAACTC CGTCCTTAGT TATAAATGAT	600
CTGATTTATG CTTATACCTC AAATCTAATT ACTCGAGGTT GTCAGGATAT AGGAAAATCA	660

-continued

TATCAAGTCT	TACAGATAGG	GATAATAACT	GTAAACTCAG	ACTTGGTACC	TGACTTAAAT	720
CCCAGGATCT	CTCATACTTT	TAACATAAAT	GACAATAGGA	AGTCATGTTT	TCTAGCACTC	780
CTAAATACAG	ATGTATATCA	ACTGTGTTCA	ACTCCCAAAG	TTGATGAAAG	ATCAGATTAT	840
GCATCATCAG	GCATAGAAGA	TATTGTAATT	GATATTGTCA	ATTATGATGG	CTCAATCTCA	900
ACAACAAGAT	TTAAGAATAA	TAACATAAGC	TTTGATCAAC	CTTATGCTGC	ACTATACCCA	960
TCTGTTGGAC	CAGGGATATA	CTACAAAGGC	AAAATAATAT	TTCTCGGGTA	TGGAGGTCTT	1020
GAACATCCAA	TAAATGAGAA	TGTAATCTGC	AACACAACCTG	GGTGTCCCGG	GAAAACACAG	1080
AGAGACTGCA	ATCAGGCATC	TCATAGTCCA	TGGTTTTTTCAG	ATAGGAGGAT	GGTCAACTCT	1140
ATCATTGTTG	TTGACAAAGG	CTTAAACTCA	ATTCCAAAAT	TGAAGGTATG	GACGATATCT	1200
ATGAGACAGA	ATTACTGGGG	GTCAGAAGGA	AGGTTACTTC	TACTAGGTAA	CAAGATCTAT	1260
ATATATACAA	GATCCACAAG	TTGGCATAGC	AAGTTACAAT	TAGGAATAAT	TGATATTACT	1320
GATTACAGTG	ATATAAGGAT	AAAATGGACA	TGGCATAATG	TGCTATCAAG	ACCAGGAAAC	1380
AATGAATGTC	CATGGGGACA	TTCATGTCCA	GATGGATGTA	TAACAGGAGT	ATATACTGAT	1440
GCATATCCAC	TCAATCCCAC	AGGGAGCATT	GTGTCATCTG	TCATATTAGA	TTCAAAAAA	1500
TCGAGAGTGA	ACCCAGTCAT	AACTTACTCA	ACAGCAACCG	AAAGAGTAAA	CGAGCTGGCC	1560
ATCCGAAACA	GAACACTCTC	AGCTGGATAT	ACAACAACAA	GCTGCATCAC	ACACTATAAC	1620
AAAGGATATT	GTTTTCATAT	AGTAGAAATA	AATCAGAAAA	GCTTAAACAC	ACTTCAACCC	1680
ATGTTGTTCA	AGACAGAGGT	TCCAAAAAGC	TGCAG			1715

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1722 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

ATGGAGTTGC	CAATCCTCAA	AGCAAATGCA	ATTACCACAA	TCCTCGCTGC	AGTCACATTT	60
TGCTTTGCTT	CTAGTCAAAA	CATCACTGAA	GAATTTTATC	AATCAACATG	CAGTGCAGTT	120
AGCAAAGGCT	ATCTTAGTGC	TCTAAGAACT	GGTTGGTATA	CTAGTGTAT	AACTATAGAA	180
TTAAGTAATA	TCAAGGAAAA	TAAGTGTAAT	GGAACAGATG	CTAAGGTAAA	ATTGATGAAA	240
CAAGAATTAG	ATAAATATAA	AAATGCTGTA	ACAGAATTGC	AGTTGCTCAT	GCAAAGCACA	300
CCAGCAGCAA	ACAATCGAGC	CAGAAGAGAA	CTACCAAGGT	TTATGAATTA	TACACTCAAC	360
AATACCAAAA	AAACCAATGT	AACATTAAGC	AAGAAAAGGA	AAAGAAGATT	TCTTGGTTTT	420
TTGTTAGGTG	TTGGATCTGC	AATCGCCAGT	GGCATTGCTG	TATCTAAGGT	CCTGCACTTA	480
GAAGGAGAAG	TGAACAAGAT	CAAAAGTGCT	CTACTATCCA	CAAACAAGGC	CGTAGTCAGC	540
TTATCAAATG	GAGTTAGTGT	CTTAACCAGC	AAAGTGTTAG	ACCTCAAAAA	CTATATAGAT	600
AAACAATTGT	TACCTATTGT	GAATAAGCAA	AGCTGCAGAA	TATCAAATAT	AGAAACTGTG	660
ATAGAGTTCC	AACAAAAGAA	CAACAGACTA	CTAGAGATTA	CCAGGAATT	TAGTGTTAAT	720
GCAGGTGTAA	CTACACCTGT	AAGCACTTAC	ATGTTAACTA	ATAGTGAATT	ATTGTCATTA	780
ATCAATGATA	TGCCTATAAC	AAATGATCAG	AAAAAGTTAA	TGTCCAACAA	TGTTCAAATA	840
GTTAGACAGC	AAAGTTACTC	TATCATGTCC	ATAATAAAAG	AGGAAGTCTT	AGCATATGTA	900
GTACAATTAC	CACTATATGG	TGTGATAGAT	ACACCTTGTT	GGAAATTACA	CACATCCCCT	960
CTATGTACAA	CCAACACAAA	AGAAGGGTCA	AACATCTGTT	TAACAAGAAC	TGACAGAGGA	1020

-continued

TGGTACTGTG	ACAATGCAGG	ATCAGTATCT	TTCTTCCCAC	AAGCTGAAAC	ATGTAAAGTT	1080
CAATCGAATC	GAGTATTTTG	TGACACAATG	AACAGTTTAA	CATTACCAAG	TGAAGTAAAT	1140
CTCTGCAATG	TTGACATATT	CAATCCCCAAA	TATGATTGTA	AAATTATGAC	TTCAAAAACA	1200
GATGTAAGCA	GCTCCGTTAT	CACATCTCTA	GGAGCCATTG	TGTCATGCTA	TGGCAAAACT	1260
AAATGTACAG	CATCCAATAA	AAATCGTGGA	ATCATAAAGA	CATTTTCTAA	CGGGTGTGAT	1320
TATGTATCAA	ATAAAGGGGT	GGACACTGTG	TCTGTAGGTA	ACACATTATA	TTATGTAAAT	1380
AAGCAAGAAG	GCAAAAAGTCT	CTATGTAAAA	GGTGAACCAA	TAATAAATTT	CTATGACCCA	1440
TTAGTATTCC	CCTCTGATGA	ATTTGATGCA	TCAATATCTC	AAGTCAATGA	GAAGATTAAC	1500
CAGAGTTTAG	CATTTATTCG	TAAATCCGAT	GAATTATTAC	ATAATGTAAA	TGCTGGTAAA	1560
TCAACCACAA	ATATCATGAT	AACTACTATA	ATTATAGTGA	TTATAGTAAT	ATTGTTATCA	1620
TTAATTGCTG	TTGGACTGCT	CCTATACTGT	AAGGCCAGAA	GCACACCAGT	CACACTAAGC	1680
AAGGATCAAC	TGAGTGGTAT	AAATAATATT	GCATTTAGTA	AC		1722

(2) INFORMATION FOR SEQ ID NO:28:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 894 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

ATGTCCAAA	ACAAGGACCA	ACGCACCGCT	AAGACACTAG	AAAAGACCTG	GGACACTCTC	60
AATCATTAT	TATTCATATC	ATCGGGCTTA	TATAAGTTAA	ATCTTAAATC	TGTAGCACAA	120
ATCACATTAT	CCATTCTGGC	AATGATAATC	TCAACTTCAC	TTATAATTAC	AGCCATCATA	180
TTCATAGCCT	CGGCAAACCA	CAAAGTCACA	CTAACAACTG	CAATCATACA	AGATGCAACA	240
AGCCAGATCA	AGAACACAAC	CCCAACATAC	CTCACTCAGG	ATCCTCAGCT	TGGAATCAGC	300
TTCTCCAATC	TGTCTGAAAT	TACATCACAA	ACCACCACCA	TACTAGCTTC	AACAACACCA	360
GGAGTCAAGT	CAAACCTGCA	ACCCACAACA	GTCAAGACTA	AAAACACAAC	AACAACCCAA	420
ACACAACCCA	GCAAGCCCAC	TACAAAACAA	CGCCAAAACA	AACCACCAA	CAAACCCAAT	480
AATGATTTTC	ACTTCGAAGT	GTTTAACTTT	GTACCCTGCA	GCATATGCAG	CAACAATCCA	540
ACCTGCTGGG	CTATCTGCAA	AAGAATACCA	AACAAAAAAC	CAGGAAAGAA	AACCACCACC	600
AAGCCTACAA	AAAAACCAAC	CTTCAAGACA	ACCAAAAAAG	ATCTCAAACC	TCAAACCACT	660
AAACCAAAGG	AAGTACCCAC	CACCAAGCCC	ACAGAAGAGC	CAACCATCAA	CACCACCAA	720
ACAAACATCA	CAACTACACT	GCTCACCAAC	AACACCACAG	GAAATCCAAA	ACTCACAAGT	780
CAAATGGAAA	CCTTCCACTC	AACCTCCTCC	GAAGGCAATC	TAAGCCCTTC	TCAAGTCTCC	840
ACAACATCCG	AGCACCCATC	ACAACCTCA	TCTCCACCCA	ACACAACACG	CCAG	894

(2) INFORMATION FOR SEQ ID NO:29:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 32 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CGTAGTTAGT	TTCCAGGACA	CTATTATCCT	AG			32
------------	------------	------------	----	--	--	----

-continued

(2) INFORMATION FOR SEQ ID NO:30:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 17 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TGAACTATTA CTCCTAG 17

(2) INFORMATION FOR SEQ ID NO:31:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 85 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GTACCTCAAC GATTAGGAGT TTCGTTTACG TTAATGGTGT TAGGAGTGAC GTCAGTGTA 60

AACAAAACGA AGACCAAGAT TCCAG 85

(2) INFORMATION FOR SEQ ID NO:32:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 27 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

CGTAGTTAGA TCGTGATGTA CTCCTAG 27

(2) INFORMATION FOR SEQ ID NO:33:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 136 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GTACGGTTGA AATTATGACG ATTAATAATG TTGTTACTAA TACCGTAGAA GGACGGTTTA 60

TCTATAGTGT TTTGATGTCG TACATCCACA TAACCAGTTG TCAGGGTTTC CCTACTTCTA 120

TAGTGTTTTG AAGCTT 136

(2) INFORMATION FOR SEQ ID NO:34:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 98 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

TAGTACCTCT ATTAATTTTA CTATTGGTAG AGCGTTTATT TATTCATAAA ATGACAAAAG 60

CATTGTCAAA ACATTATTTT TTTGGATATT TATCTTAA 98

(2) INFORMATION FOR SEQ ID NO:35:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 145 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

-continued

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TAGTACCTCT ATTAATTTTA CTATTGGTAG AGCGTTTATT TATTCATAAA ATGACAAAAG 60
 CATTGTCAAA ACATTATTTT TTTGGATATT TATAAGGCCT TAAGTCTAGA CGTCGCCGGC 120
 GAGGTAGATC TTCCATGGGC CCTAG 145

(2) INFORMATION FOR SEQ ID NO:36:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 31 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

TGATTAAGGT AGTTTTCACT TTTCCGAGTA C 31

(2) INFORMATION FOR SEQ ID NO:37:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 39 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

TAAAGACACT ATAAACACGC CATATTACTA CGATATGGA 39

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 10 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

AGGACAAAAG 10

What is claim is:

1. A multimeric hybrid gene encoding a chimeric protein including a protein from parainfluenza virus (PIV) and a protein from respiratory syncytial virus (RSV), comprising a nucleotide sequence encoding a fragment of a PIV-3 F protein having fusion activity or a fragment of a PIV-3 HN protein having haemagglutinin-neuramidase activity linked to a nucleotide sequence coding for a fragment of a RSV G protein having attachment activity or a RSV F protein having fusion activity, wherein said nucleotide sequences code for the respective protein lacking the hydrophobic anchor and cytoplasmic tail domains of the respective protein.

2. A multimeric hybrid gene encoding a chimeric protein including a protein from parainfluenza virus (PIV) and a protein from respiratory syncytial virus (RSV), comprising a nucleotide sequence coding for a PIV F protein or an immunogenic fragment thereof having fusion activity or a nucleotide sequence coding for a PIV HN protein or an immunogenic fragment thereof having haemagglutinin-neuramidase activity linked to a nucleotide sequence coding for a RSV G protein or an immunogenic fragment thereof having attachment activity or a nucleotide sequence coding for a RSV F protein or an immunogenic fragment thereof having fusion activity and further comprising an upstream nucleotide sequence and a downstream nucleotide sequence

wherein said downstream nucleotide sequence lacks a 5'-untranslated region.

3. The hybrid gene of claim 2 wherein said upstream nucleotide sequence also lacks a 5'-untranslated region.

4. The hybrid gene of claim 2 wherein both said upstream and downstream nucleotide sequences lack a region coding for the transmembrane anchor and cytoplasmic tail of the respective protein.

5. The hybrid gene of claim 2 wherein said downstream sequence codes for a F1 subunit only of the RSV F protein.

6. The hybrid gene of claim 5 wherein said upstream nucleotide sequence codes for a PIV-3 F protein lacking its hydrophobic anchor and cytoplasmic tail domains and said downstream nucleotide sequence codes for a F1 subunit only of the RSV F protein which lacks its hydrophobic anchor and cytoplasmic tail domains.

7. The hybrid gene of claim 6 wherein said upstream nucleotide sequence lacks a 5'-untranslated region.

8. The hybrid gene of claim 2 wherein said downstream nucleotide sequence codes for a PIV-3 HN protein or a RSV G protein each lacking its hydrophobic anchor and cytoplasmic tail domains.

9. The hybrid gene of claim 8 wherein said upstream nucleotide sequence codes for a PIV-3 F or RSV F protein each lacking its hydrophobic anchor and cytoplasmic tail domains.

10. A process for the preparation of a chimeric protein including a protein from parainfluenza virus (PIV) and a protein from respiratory syncytial virus (RSV), which comprises:

isolating a first nucleotide sequence encoding a fragment
of a PIV-3 F protein having fusion activity or a first
nucleotide sequence encoding a fragment of a PIV-3
HN protein having haemagglutinin-neuramidase
activity,

isolating a second nucleotide sequence encoding a frag-
ment of a RSV G protein having attachment activity or
a second nucleotide sequence encoding a fragment of a
RSV F protein having fusion activity,

said first and second nucleotide sequences coding for the
respective protein lacking the hydrophobic anchor and
cytoplasmic tail domains of the respective protein,

linking said first and second nucleotide sequences to form
a multimeric hybrid gene, and

expressing the multimeric hybrid gene in a eukaryotic
cellular expression system.

11. A process for the preparation of a chimeric protein including a protein from parainfluenza virus (PIV) and a protein from respiratory syncytial virus (RSV), which comprises:

isolating a first nucleotide sequence coding for a PIV F
protein or an immunogenic fragment thereof having
fusion activity or a first nucleotide sequence coding for
a PIV HN protein or an immunogenic fragment thereof
having haemagglutinin-neuramidase activity,

isolating a second nucleotide sequence coding for a RSV
G protein or an immunogenic fragment thereof having
attachment activity or a second nucleotide sequence
coding for a RSV F protein or an immunogenic frag-
ment thereof having fusion activity,

wherein one of said first and second nucleotide sequences
comprises an upstream nucleotide sequence and the
other of said first and second nucleotide sequences
comprises a downstream nucleotide sequence wherein
said downstream nucleotide sequence lacks a
5'-untranslated region,

linking said first and second nucleotide sequences to form
a multimeric hybrid gene, and

expressing the multimeric hybrid gene in a eukaryotic
cellular expression system.

12. The process of claim **11** wherein said upstream
nucleotide sequence also lacks a 5'-untranslated region.

13. The process of claim **11** wherein both said upstream
and downstream nucleotide sequences lack a region coding
for the transmembrane anchor and cytoplasmic tail of the
respective protein.

14. The process of claim **11** wherein said downstream
sequence codes for a F1 subunit only of the RSV F protein.

15. The process of claim **14** wherein said upstream
nucleotide sequence codes for a PIV-3 F protein lacking its
hydrophobic anchor and cytoplasmic tail domains and said
downstream nucleotide sequence codes for a F1 subunit only
of the RSV F protein which lacks its hydrophobic anchor
and cytoplasmic tail domains.

16. The process of claim **15** wherein said upstream
nucleotide sequence lacks a 5'-untranslated region.

17. The process of claim **11** wherein said downstream
nucleotide sequence codes for a PIV-3 HN protein or a RSV
G protein each lacking its hydrophobic anchor and cyto-
plasmic tail domains.

18. The process of claim **17** wherein said upstream
nucleotide sequence codes for a PIV-3 F or RSV F protein
each lacking its hydrophobic anchor and cytoplasmic tail
domains.

* * * * *