



US006187576B1

(12) **United States Patent**  
**Svendsen et al.**

(10) **Patent No.:** **US 6,187,576 B1**  
(45) **Date of Patent:** **Feb. 13, 2001**

(54)  **$\alpha$ -AMYLASE MUTANTS**

(75) Inventors: **Allan Svendsen**, Birkerød; **Torben Vedel Borchert**, Jyllinge; **Henrik Bisgård-Frantzen**, Bagsværd, all of (DK)

(73) Assignee: **Novo Nordisk A/S**, Bagsvaerd (DK)

(\* ) Notice: Under 35 U.S.C. 154(b), the term of this patent shall be extended for 0 days.

(21) Appl. No.: **09/170,670**

(22) Filed: **Oct. 13, 1998**

**Related U.S. Application Data**

(60) Provisional application No. 60/063,306, filed on Oct. 28, 1997.

**Foreign Application Priority Data**

Oct. 13, 1997 (DK) ..... 1172/97

(51) **Int. Cl.**<sup>7</sup> ..... **C12N 9/00**; C12N 9/24; C12N 9/28; C11D 3/00; C11D 7/42

(52) **U.S. Cl.** ..... **435/202**; 435/183; 435/200; 510/226; 510/235; 510/320; 510/392

(58) **Field of Search** ..... 435/96, 98, 172.1, 435/252.3, 202, 220, 201, 204, 440; 536/23.2; 510/226, 235, 320, 392

(56) **References Cited**

**FOREIGN PATENT DOCUMENTS**

- WO 90/11352 10/1990 (WO) .
- WO 91/00353 1/1991 (WO) .
- WO 95/10603 4/1995 (WO) .
- WO 95/26397 10/1995 (WO) .

- WO 95/35382 12/1995 (WO) .
- WO 96/23873 8/1996 (WO) .
- WO 96/23874 8/1996 (WO) .

**OTHER PUBLICATIONS**

Gray G.L. et al. Structural genes encoding the thermophilic alpha-amylases of *Bacillus stearothermophilus* and *B.licheniformis*. J.Bacteriol., May 1986, vol. 166(2):635-643.\*

\* cited by examiner

*Primary Examiner*—Rebecca E. Prouty

*Assistant Examiner*—Manjunath Rao

(74) *Attorney, Agent, or Firm*—Elias J. Lambiris, Esq.; Reza Green, Esq.

(57) **ABSTRACT**

The invention relates to a variant of a parent Termamyl-like  $\alpha$ -amylase, comprising mutations in two, three, four, five or six regions/positions. The variants have increased thermostability at acidic pH and/or at low Ca<sup>2+</sup> concentrations (relative to the parent). The invention also relates to a DNA construct comprising a DNA sequence encoding an  $\alpha$ -amylase variant of the invention, a recombinant expression vector which carries a DNA construct of the invention, a cell which is transformed with a DNA construct of the invention, the use of an  $\alpha$ -amylase variant of the invention for washing and/or dishwashing, textile desizing, starch liquefaction, a detergent additive comprising an  $\alpha$ -amylase variant of the invention, a manual or automatic dishwashing detergent composition comprising an  $\alpha$ -amylase variant of the invention, a method for generating a variant of a parent Termamyl-like  $\alpha$ -amylase, which variant exhibits increased thermostability at acidic pH and/or at low Ca<sup>2+</sup> concentrations (relative to the parent).

**22 Claims, 3 Drawing Sheets**

1	HHNGTNGTMM	QYFEWHLPLND	GNHWNRLRDD	ASNLNRGKIT	AIWIPPAWKKG	50
2	..NGTNGTMM	QYFEWYLPND	GNHWNRLRSD	ASNLKDKGIS	AVWIPPAWKKG	
3	HHNGTNGTMM	QYFEWYLPND	GNHWNRLRDD	AANLKSCKGIT	AVWIPPAWKKG	
4	...VNGTLM	QYFEWYTPND	GQHWKRLQND	AEHLSDIGIT	AVWIPPAYKKG	
5	..ANLNGTLM	QYFEWYMPND	GQHWRRRLQND	SAYLAEHGKIT	AVWIPPAYKKG	
6	.AAPFNGTMM	QYFEWYLPDD	GTLWTKVANE	ANLSSLGIT	ALWLPPAYKKG	
51	TSQNDVGYGA	YDLYDLGEFN	QKGTVRTKYG	TRSQLESIAH	ALKNNGVQVY	100
1	ASQNDVGYGA	YDLYDLGEFN	QKGTIRTKYG	TRNQLQAAVN	ALKSNGIQVY	
2	TSQNDVGYGA	YDLYDLGEFN	QKGTVRTKYG	TRNQLQAAVT	SLKNNGIQVY	
3	LSQSDNGYGP	YDLYDLGEFQ	QKGTVRTKYG	TKSELQDAIG	SLHSRNVQVY	
4	TSQADVGYGA	YDLYDLGEFH	QKGTVRTKYG	TKGELQSAIK	SLHSRDINVY	
5	TSRSDVGYGV	YDLYDLGEFN	QKGTVRTKYG	TKAQYLQAIQ	AAHAAGMQVY	
6						
101	GDVVMNHKGG	ADATENVLAV	EVNPNRNQOE	ISGDYTI EAW	TKFDFPGRGN	150
1	GDVVMNHKGG	ADATEMVRVAV	EVNPNRNQOE	VSGEYTI EAW	TKFDFPGRGN	
2	GDVVMNHKGG	ADGTEIVNAV	EVNRSNRNQE	TSGEYAI EAW	TKFDFPGRGN	
3	GDVVLNHKAG	ADATEDVTAV	EVNPNRNQOE	TSEEYQIKAW	TDFRFPGRGN	
4	GDVVINHKGG	ADATEDVTAV	EVDPADRNRV	ISGEHLIKAW	THFHFPRGRS	
5	ADVVFEDHKGG	ADGTEWVDAV	EVNPSDRNQE	ISGTYQIQAW	TKFDFPGRGN	
6						
151	TYSDFKWRWY	HFDGVDWDQS	RQFQNR IYKF	RGDGKAWDWE	VDS ENGN YDY	200
1	THSNFKWRWY	HFDGVDWDQS	RKLNNR IYKF	RGDGKAWDWE	VDTE NGN YDY	
2	NHSSEKWRWY	HFDGTDWDQS	RQLQNK IYKF	RGTGKAWDWE	VDTE NGN YDY	
3	TYSDFKWHWY	HFDGADWDES	RKI.SRI FKF	RGEGKAWDWE	VSE NGN YDY	
4	TYSDFKWHWY	HFDGTDWDES	RKL.NRI YKF	.Q GKAWDWE	VSE NGN YDY	
5	TYSSEKWRWY	HFDGVDWDES	RKL.SRI YKF	RGIGKAWDWE	VDTE NGN YDY	
6						

Fig. 1a



201						250
1	LMYADVDMDH	PEVVNELRRW	GEWYNTLNL	DGERIDAVKH	DGERIDAVKH	IKYSFTRDWL
2	LMYADIDMDH	PEVVNELRNW	GVWYNTLGL	DGERIDAVKH	DGERIDAVKH	IKYSFTRDWS
3	LMYADVDMDH	PEVIHELNRW	GVWYNTLNL	DGERIDAVKH	DGERIDAVKH	IKYSFTRDWL
4	LMYADVVDYDH	PDVVAETKKW	GIWYANELSL	DGERIDAACKH	DGERIDAACKH	IKFSFLRDWV
5	LMYADIDYDH	PDVAAEIKRW	GTWYANELQL	DGERLDAVKH	DGERLDAVKH	IKFSFLRDWV
6	LMYADLDMDH	PEVVTELKNW	GKWYVNTTNI	DGERLDAVKH	DGERLDAVKH	IKFSFFPDWL
251						300
1	THVRNATGKE	MFAVAEFFWKN	DLGALENYLN	KTNWNHVSFD	KTNWNHVSFD	VPLHYNLYNA
2	IHVRSATGKN	MFAVAEFFWKN	DLGAIENYLN	KTNWNHVSFD	KTNWNHVSFD	VPLHYNFYNA
3	THVRNTTGKP	MFAVAEFFWKN	DLGAIENYLN	KTSWNHSAFD	KTSWNHSAFD	VPLHYNLYNA
4	QAVRQATGKE	MFTVAEYWQN	NAGKLENYLN	KTSFNQSVFD	KTSFNQSVFD	VPLHFNLQAA
5	NHVREKTGKE	MFTVAEYWQN	DLGALENYLN	KTNFNHVSFD	KTNFNHVSFD	VPLHYQFHAA
6	SYVRSQTGKP	LFTVGEYWSY	DINKLHNYIT	KTDGTMVSFD	KTDGTMVSFD	APLHNKFYTA
301						350
1	SNSGGNYDMA	KLLNGTVVQK	HPMHAUTFVD	NHDSQPGEAL	NHDSQPGEAL	ESFVQEWFKP
2	SKSGGNYDMR	QIFNGTVVQR	HPMHAUTFVD	NHDSQPGEAL	NHDSQPGEAL	ESFVEEWFKP
3	SNSGGYYDMR	NILNGSVVQK	HPTHAVTFVD	NHDSQPGEAL	NHDSQPGEAL	ESFVQQWFKP
4	SSQGGGYDMR	RLLDGTVVSR	HPEKAVTFVE	NHDTQPGQSL	NHDTQPGQSL	ESTVQTWFKP
5	STQGGGYDMR	KLLNGTVVSK	HPLKSVTFVD	NHDTQPGQSL	NHDTQPGQSL	ESTVQTWFKP
6	SKSGGAFDMR	TLMTNTLMKD	QPTLAVTFVD	NHDTQPGQAL	NHDTQPGQAL	QSWVDPWFKP
351						400
1	LAYALILTRE	QGYPSVIFYGD	YYGIPTHS..	VPAMKAKID	VPAMKAKID	PILEARQNFA
2	LAYALTLTRE	QGYPSVIFYGD	YYGIPTHG..	VPAMKSKID	VPAMKSKID	PILEARQKYA
3	LAYALVLTRE	QGYPSVIFYGD	YYGIPTHG..	VPAMKSKID	VPAMKSKID	PLLQARQTFA
4	LAYAFILTRE	SGYPQVIFYGD	MYGTKGTSPK	EIPSLKDNIE	EIPSLKDNIE	PILKARKEYA
5	LAYAFILTRE	SGYPQVIFYGD	MYGTKGDSQR	EIPALKHKIE	EIPALKHKIE	PILKARKQYA
6	LAYAFILTRQ	EGYPCVIFYGD	YYGIPQYN..	IPSLKSKID	IPSLKSKID	PLLIARRDYA

Figure 1 (continued)

Fig. 1b

401						450
1	YGTQHDYFDH	HNIIGWTREG	NTTHPNSGLA	TIMSDGPGGE	KWMYVGGQNK	
2	YGRQN.....	.....	.....	.....	.....	
3	YGTQHDYFDH	HDIIGWTREG	NSSHPNSGLA	TIMSDGPGGN	KWMYVGGKNA	
4	YGPQHDYIDH	PVIGWTREG	DSSAAKSGLA	ALITDGPGG	KRMYAGLKNA	
5	YGAQHDYFDH	HDIIGWTREG	DSSVANSGLA	ALITDGPGG	KRMYVGRQNA	
6	YGTQHDYLDH	SDIIGWTREG	GTEKPGSGLA	ALITDGPGG	KWMYVGGKQHA	
451						500
1	GQVWHDITGN	KPGTVTINAD	GWANFSVNGG	SVSIWVKR..	.....	
2	.....	.....	.....	.....	.....	
3	GQVWRDITGN	RTGTVTINAD	GWGNFSVNGG	SVSVWVKQ..	.....	
4	GETWYDITGN	RSDTVKIGSD	GWGEFHVNDG	SVSIYVQ...	.....	
5	GETWHDITGN	RSEPVINSE	GWGEFHVNGG	SVSIYVQR..	.....	
6	GKVFYDLTGN	RSDTVTINSD	GWGEFKVNGG	SVSVWVPRKT	TVSTIARPIIT	
501						519
1	.....	.....	.....	.....	.....	
2	.....	.....	.....	.....	.....	
3	.....	.....	.....	.....	.....	
4	.....	.....	.....	.....	.....	
5	.....	.....	.....	.....	.....	
6	TRPWTGEFVR	WTEPRLVAW				

Fig. 1c

Figure 1 (continued)



**α-AMYLASE MUTANTS****CROSS-REFERENCE TO RELATED APPLICATIONS**

The application claims priority under 35 U.S.C 119 of Danish application 1172/97 filed Oct. 13, 1997, and of U.S. provisional application 60/063,306 filed Oct. 28, 1997, the contents of which are fully incorporated herein by reference.

**FIELD OF THE INVENTION**

The present invention relates, inter alia, to novel variants (mutants) of parent Termamyl-like α-amylases, notably variants exhibiting increased thermostability at acidic pH and/or at low Ca<sup>2+</sup> concentrations (relative to the parent) which are advantageous with respect to applications of the variants in, industrial starch processing particularly (e.g. starch liquefaction or saccharification).

**BACKGROUND OF THE INVENTION**

α-Amylases (α-1,4-glucan-4-glucanohydrolases, EC 3.2.1.1) constitute a group of enzymes which catalyze hydrolysis of starch and other linear and branched 1,4-glucosidic oligo- and polysaccharides.

There is a very extensive body of patent and scientific literature relating to this industrially very important class of enzymes. A number of α-amylase such as Termamyl-like α-amylases variants are known from e.g. WO 90/11352, WO 95/10603, WO 95/26397, WO 96/23873 and WO 96/23874.

Among more recent disclosures relating to α-amylases, WO 96/23874 provides three-dimensional, X-ray crystal structural data for a Termamyl-like α-amylase which consists of the 300 N-terminal amino acid residues of the *B. amyloliquefaciens* α-amylase and amino acids 301–483 of the C-terminal end of the *B. licheniformis* α-amylase comprising the amino acid sequence (the latter being available commercially under the tradename Termamyl™), and which is thus closely related to the industrially important Bacillus α-amylases (which in the present context are embraced within the meaning of the term “Termamyl-like α-amylases”, and which include, inter alia, the *B. licheniformis*, *B. amyloliquefaciens* and *B. stearothermophilus* α-amylases). WO 96/23874 further describes methodology for designing, on the basis of an analysis of the structure of a parent Termamyl-like α-amylase, variants of the parent Termamyl-like α-amylase which exhibit altered properties relative to the parent.

WO 95/35382 (Gist Brocades B.V.) concerns amylolytic enzymes derived from *B. licheniformis* with improved properties allowing reduction of the Ca<sup>2+</sup> concentration under application without a loss of performance of the enzyme. The amylolytic enzyme comprises one or more amino acid changes at positions selected from the group of 104, 128, 187, 188 of the *B. licheniformis* α-amylase sequence.

WO 96/23873 (Novo Nordisk) discloses Termamyl-like α-amylase variants which have increased thermostability obtained by pairwise deletion in the region R181\*, G182\*, T183\* and G184\* of the sequence shown in SEQ ID NO: 1 herein.

**BRIEF DISCLOSURE OF THE INVENTION**

The present invention relates to novel α-amylolytic variants (mutants) of a Termamyl-like α-amylase, in particular variants exhibiting increased thermostability (relative to the parent) which are advantageous in connection with the

industrial processing of starch (starch liquefaction, saccharification and the like).

The inventors have surprisingly found out that in case of combining two, three, four, five or six mutations (will be described below), the thermostability of Termamyl-like α-amylases is increased at acidic pH and/or at low Ca<sup>2+</sup> concentration in comparison to single mutations, such as the mutation disclosed in WO 96/23873 (Novo Nordisk), i.e. pairwise deletion in the region R181\*, G182\*, T183\* and G184\* of the sequence shown in SEQ ID NO: 1 herein.

The invention further relates to DNA constructs encoding variants of the invention, to composition comprising variants of the invention, to methods for preparing variants of the invention, and to the use of variants and compositions of the invention, alone or in combination with other α-amylolytic enzymes, in various industrial processes, e.g., starch liquefaction.

**BRIEF DESCRIPTION OF THE DRAWING**

FIG. 1 is an alignment of the amino acid sequences of six parent Termamyl-like α-amylases in the context of the invention. The numbers on the Extreme left designate the respective amino acid sequences as follows:

- 1: SEQ ID NO: 2,
- 2: Kaoamyl,
- 3: SEQ ID NO: 1,
- 4: SEQ ID NO: 5,
- 5: SEQ ID NO: 4,
- 6: SEQ ID NO: 3.

**DETAILED DISCLOSURE OF THE INVENTION**

All patents, patent applications, and literature references referred to herein are hereby incorporated by reference in their entirety.

**The Termamyl-like α-amylase**

It is well known that a number of α-amylases produced by Bacillus spp. are highly homologous on the amino acid level. For instance, the *B. licheniformis* α-amylase comprising the amino acid sequence shown in SEQ ID NO: 4 (commercially available as Termamyl™) has been found to be about 89% homologous with the *B. amyloliquefaciens* α-amylase comprising the amino acid sequence shown in SEQ ID NO: 5 and about 79% homologous with the *B. stearothermophilus* α-amylase comprising the amino acid sequence shown in SEQ ID NO: 3. Further homologous α-amylases include an α-amylase derived from a strain of the Bacillus sp. NCIB 12289, NCIB 12512, NCIB 12513 or DSM 9375, all of which are described in detail in WO 95/26397, and the α-amylase described by Tsukamoto et al., Biochemical and Biophysical Research Communications, 151 (1988), pp. 25–31.

Still further homologous α-amylases include the α-amylase produced by the *B. licheniformis* strain described in EP 0252666 (ATCC 27811), and the α-amylases identified in WO 91/00353 and WO 94/18314. Other commercial Termamyl-like *B. licheniformis* α-amylases are Optitherm™ and Takathe™ (available from Solvay), Maxamyl™ (available from Gist-brocades/Genencor), Spezym AA™ and Spezyme Delta AA™ (available from Genencor), and Keistase™ (available from Daiwa).

Because of the substantial homology found between these α-amylases, they are considered to belong to the same class of α-amylases, namely the class of “Termamyl-like α-amylases”.

Accordingly, in the present context, the term “Termamyl-like α-amylase” is intended to indicate an α-amylase which,



at the amino acid level, exhibits a substantial homology to Termamyl™, i.e. the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 4 herein. In other words, a Termamyl-like  $\alpha$ -amylase is an  $\alpha$ -amylase which has the amino acid sequence shown in SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7 or 8 herein, and the amino acid sequence shown in SEQ ID NO: 1 of WO 95/26397 (the same as the amino acid sequence shown as SEQ ID NO: 7 herein) or in SEQ ID NO: 2 of WO 95/26397 (the same as the amino acid sequence shown as SEQ ID NO: 8 herein) or in Tsukamoto et al., 1988, (which amino acid sequence is shown in SEQ ID NO: 6 herein) or i) which displays at least 60%, preferred at least 70%, more preferred at least 75%, even more preferred at least 80%, especially at least 85%, especially preferred at least 90%, even especially more preferred at least 95% homology with at least one of said amino acid sequences shown in SEQ ID NOS 1 or 2 or 3 or 4 or 5 or 6 or 7 or 8 and/or ii) displays immunological cross-reactivity with an antibody raised against at least one of said  $\alpha$ -amylases, and/or iii) is encoded by a DNA sequence which hybridizes to the DNA sequences encoding the above-specified  $\alpha$ -amylases which are apparent from SEQ ID NOS: 9, 10, 11, or 12 of the present application (which encoding sequences encode the amino acid sequences shown in SEQ ID NOS: 1, 2, 3, 4 and 5 herein, respectively), from SEQ ID NO: 4 of WO 95/26397 (which DNA sequence, together with the stop codon TAA, is shown in SEQ ID NO: 13 herein and encodes the amino acid sequence shown in SEQ ID NO: 8 herein) and from SEQ ID NO: 5 of WO 95/26397 (shown in SEQ ID NO: 14 herein), respectively.

In connection with property i), the "homology" may be determined by use of any conventional algorithm, preferably by use of the GAP programme from the GCG package version 7.3 (June 1993) using default values for GAP penalties, which is a GAP creation penalty of 3.0 and GAP extension penalty of 0.1, (Genetic Computer Group (1991) Programme Manual for the GCG Package, version 7, 575 Science Drive, Madison, Wis., USA 53711).

A structural alignment between Termamyl and a Termamyl-like  $\alpha$ -amylase may be used to identify equivalent/corresponding positions in other Termamyl-like  $\alpha$ -amylases. One method of obtaining said structural alignment is to use the Pile Up programme from the GCG package using default values of gap penalties, i.e., a gap creation penalty of 3.0 and gap extension penalty of 0.1. Other structural alignment methods include the hydrophobic cluster analysis (Gaboriaud et al., (1987), FEBS LETTERS 224, pp. 149–155) and reverse threading (Huber, T; Torda, AE, PROTEIN SCIENCE Vol. 7, No. 1 pp. 142–149 (1998).

Property ii) of the  $\alpha$ -amylase, i.e. the immunological cross reactivity, may be assayed using an antibody raised against, or reactive with, at least one epitope of the relevant Termamyl-like  $\alpha$ -amylase. The antibody, which may either be monoclonal or polyclonal, may be produced by methods known in the art, e.g. as described by Hudson et al., Practical Immunology, Third edition (1989), Blackwell Scientific Publications. The immunological cross-reactivity may be determined using assays known in the art, examples of which are Western Blotting or radial immunodiffusion assay, e.g. as described by Hudson et al., 1989. In this respect, immunological cross-reactivity between the  $\alpha$ -amylases having the amino acid sequences SEQ ID NOS: 1, 2, 3, 4, 5, 6, 7, or 8 respectively, have been found.

The oligonucleotide probe used in the characterization of the Termamyl-like  $\alpha$ -amylase in accordance with property iii) above may suitably be prepared on the basis of the full or partial nucleotide or amino acid sequence of the  $\alpha$ -amylase in question.

Suitable conditions for testing hybridization involve pre-soaking in 5×SSC and prehybridizing for 1 hour at ~40° C. in a solution of 20% formamide, 5×Denhardt's solution, 50 mM sodium phosphate, pH 6.8, and 50 mg of denatured sonicated calf thymus DNA, followed by hybridization in the same solution supplemented with 100 mM ATP for 18 hours at ~40° C., followed by three times washing of the filter in 2×SSC, 0.2% SDS at 40° C. for 30 minutes to (low stringency), preferred at 50° C. (medium stringency), more preferably at 65° C. (high stringency), even more preferably at ~75° C. (very high stringency). More details about the hybridization method can be found in Sambrook et al., Molecular Cloning: A Laboratory Manual, 2nd Ed., Cold Spring Harbor, 1989.

In the present context, "derived from" is intended not only to indicate an  $\alpha$ -amylase produced or producible by a strain of the organism in question, but also an  $\alpha$ -amylase encoded by a DNA sequence isolated from such strain and produced in a host organism transformed with said DNA sequence. Finally, the term is intended to indicate an  $\alpha$ -amylase which is encoded by a DNA sequence of synthetic and/or cDNA origin and which has the identifying characteristics of the  $\alpha$ -amylase in question. The term is also intended to indicate that the parent  $\alpha$ -amylase may be a variant of a naturally occurring  $\alpha$ -amylase, i.e. a variant which is the result of a modification (insertion, substitution, deletion) of one or more amino acid residues of the naturally occurring  $\alpha$ -amylase.

#### Parent Hybrid $\alpha$ -amylases

The parent  $\alpha$ -amylase may be a hybrid  $\alpha$ -amylase, i.e. an  $\alpha$ -amylase which comprises a combination of partial amino acid sequences derived from at least two  $\alpha$ -amylases.

The parent hybrid  $\alpha$ -amylase may be one which on the basis of amino acid homology and/or immunological cross-reactivity and/or DNA hybridization (as defined above) can be determined to belong to the Termamyl-like  $\alpha$ -amylase family. In this case, the hybrid  $\alpha$ -amylase is typically composed of at least one part of a Termamyl-like  $\alpha$ -amylase and part(s) of one or more other  $\alpha$ -amylases selected from Termamyl-like  $\alpha$ -amylases or non-Termamyl-like  $\alpha$ -amylases of microbial (bacterial or fungal) and/or mammalian origin.

Thus, the parent hybrid  $\alpha$ -amylase may comprise a combination of partial amino acid sequences deriving from at least two Termamyl-like  $\alpha$ -amylases, or from at least one Termamyl-like and at least one non-Termamyl-like bacterial  $\alpha$ -amylase, or from at least one Termamyl-like and at least one fungal  $\alpha$ -amylase. The Termamyl-like  $\alpha$ -amylase from which a partial amino acid sequence derives may, e.g., be any of those specific Termamyl-like  $\alpha$ -amylases referred to herein.

For instance, the parent  $\alpha$ -amylase may comprise a C-terminal part of an  $\alpha$ -amylase derived from a strain of *B. licheniformis*, and a N-terminal part of an  $\alpha$ -amylase derived from a strain of *B. amyloliquefaciens* or from a strain of *B. stearotherophilus*. For instance, the parent  $\alpha$ -amylase may comprise at least 430 amino acid residues of the C-terminal part of the *B. licheniformis*  $\alpha$ -amylase, and may, e.g. comprise a) an amino acid segment corresponding to the 37 N-terminal amino acid residues of the *B. amyloliquefaciens*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 5 and an amino acid segment corresponding to the 445 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID No. 4, or b) an amino acid segment corresponding to the 68 N-terminal amino acid residues of the *B. stearotherophilus*  $\alpha$ -amylase having the amino acid sequence shown in



SEQ ID NO: 3 and an amino acid segment corresponding to the 415 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 4.

The non-Termamyl-like  $\alpha$ -amylase may, e.g., be a fungal  $\alpha$ -amylase, a mammalian or a plant  $\alpha$ -amylase or a bacterial  $\alpha$ -amylase (different from a Termamyl-like  $\alpha$ -amylase). Specific examples of such  $\alpha$ -amylases include the *Aspergillus oryzae* TAKA  $\alpha$ -amylase, the *A. niger* acid  $\alpha$ -amylase, the *Bacillus subtilis*  $\alpha$ -amylase, the porcine pancreatic  $\alpha$ -amylase and a barley  $\alpha$ -amylase. All of these  $\alpha$ -amylases have elucidated structures which are markedly different from the structure of a typical Termamyl-like  $\alpha$ -amylase as referred to herein.

The fungal  $\alpha$ -amylases mentioned above, i.e. derived from *A. niger* and *A. oryzae*, are highly homologous on the amino acid level and generally considered to belong to the same family of  $\alpha$ -amylases. The fungal  $\alpha$ -amylase derived from *Aspergillus oryzae* is commercially available under the tradename Fungamyl™.

Furthermore, when a particular variant of a Termamyl-like  $\alpha$ -amylase (variant of the invention) is referred to—in a conventional manner—by reference to modification (e.g. deletion or substitution) of specific amino acid residues in the amino acid sequence of a specific Termamyl-like  $\alpha$ -amylase, it is to be understood that variants of another Termamyl-like  $\alpha$ -amylase modified in the equivalent position(s) (as determined from the best possible amino acid sequence alignment between the respective amino acid sequences) are encompassed thereby.

A preferred embodiment of a variant of the invention is one derived from a *B. licheniformis*  $\alpha$ -amylase (as parent Termamyl-like  $\alpha$ -amylase), e.g. one of those referred to above, such as the *B. licheniformis*  $\alpha$ -amylase having the amino acid sequence shown in SEQ ID NO: 4.

#### Construction of Variants of the Invention

The construction of the variant of interest may be accomplished by cultivating a microorganism comprising a DNA sequence encoding the variant under conditions which are conducive for producing the variant. The variant may then subsequently be recovered from the resulting culture broth. This is described in detail further below.

#### Altered Properties of Variants of the Invention

The following discusses the relationship between mutations which may be present in variants of the invention, and desirable alterations in properties (relative to those a parent, Termamyl-like  $\alpha$ -amylase) which may result therefrom.

#### Increased Thermostability at Acidic pH and/or at Low $\text{Ca}^{2+}$ Concentration

Mutations of particular relevance in relation to obtaining variants according to the invention having increased thermostability at acidic pH and/or at low  $\text{Ca}^{2+}$  concentration include mutations at the following positions (relative to *B. licheniformis*  $\alpha$ -amylase, SEQ ID NO: 4): H156, N172, A181, N188, N190, H205, D207, A209, A210, E211, Q264, N265.

In the context of the invention the term “acidic pH” means a pH below 7.0, especially below the pH range, in which industrial starch liquefaction processes are normally performed, which is between pH 5.5 and 6.2.

In the context of the present invention the term “low Calcium concentration” means concentrations below the normal level used in industrial starch liquefaction. Normal concentrations vary depending of the concentration of free  $\text{Ca}^{2+}$  in the corn. Normally a dosage corresponding to 1 mM (40 ppm) is added which together with the level in corn gives between 40 and 60 ppm free  $\text{Ca}^{2+}$ .

In the context of the invention the term “high temperatures” means temperatures between 95° C. and 160° C., especially the temperature range in which industrial starch liquefaction processes are normally performed, which is between 95° C. and 105° C.

The inventors have now found that the thermostability at acidic pH and/or at low  $\text{Ca}^{2+}$  concentration may be increased even more by combining certain mutations including the above mentioned mutations and/or I201 with each other.

Said “certain” mutations are the following (relative to *B. licheniformis*  $\alpha$ -amylase, SEQ ID NO: 4): N190, D207, E211, Q264 and I201.

Said mutation may further be combined with deletions in one, preferably two or even three positions as described in WO 96/23873 (i.e. in positions R181, G182, T183, G184 in SEQ ID NO: 1 herein). According to the invention variants of a parent Termamyl-like  $\alpha$ -amylase with  $\alpha$ -amylase activity comprising mutations in two, three, four, five or six of the above positions are contemplated.

It should be emphasized that not only the Termamyl-like  $\alpha$ -amylases mentioned specifically below are contemplated. Also other commercial Termamyl-like  $\alpha$ -amylases are contemplated. An unexhaustive list of such  $\alpha$ -amylases is the following:  $\alpha$ -amylases produced by the *B. licheniformis* strain described in EP 0252666 (ATCC 27811), and the  $\alpha$ -amylases identified in WO 91/00353 and WO 94/18314. Other commercial Termamyl-like *B. licheniformis*  $\alpha$ -amylases are Optitherm™ and Takatherm™ (available from Solvay), Maxamyl™ (available from Gist-brocades/Genencor), Spezym AA™ Spezyme Delta AA™ (available from Genencor), and Keistase™ (available from Daiwa).

It may be mentioned here that amino acid residues, respectively, at positions corresponding to N190, I201, D207 and E211, respectively, in SEQ ID NO: 4 constitute amino acid residues which are conserved in numerous Termamyl-like  $\alpha$ -amylases. Thus, for example, the corresponding positions of these residues in the amino acid sequences of a number of Termamyl-like  $\alpha$ -amylases which have already been mentioned (vide supra) are as follows:

TABLE 1

Termamyl-like $\alpha$ -amylase	N	I	D	E	Q
<i>B. licheniformis</i> (SEQ ID NO: 4)	N190	I201	D207	E211	Q264
<i>B. amyloliquefaciens</i> (SEQ ID NO: 5)	N190	V201	D207	E211	Q264
<i>B. stearothermophilus</i> (SEQ ID NO: 3)	N193	L204	E210	E214	—
<i>Bacillus</i> WO 95/26397 (SEQ ID NO: 2)	N195	V206	E212	E216	—
<i>Bacillus</i> WO 95/26397 (SEQ ID NO: 1)	N195	V206	E212	E216	—
“ <i>Bacillus</i> sp. #707” (SEQ ID NO: 6)	N195	I206	E212	E216	—

Mutations of these conserved amino acid residues are very important in relation to improving thermostability at acidic pH and/or at low calcium concentration, and the following mutations are of particular interest in this connection (with reference to the numbering of the *B. licheniformis* amino acid sequence shown in SEQ ID NO: 4).

Pair-wise amino acid deletions at positions corresponding to R179-G182 in SEQ ID NO: 5 corresponding to a gap in Seq ID NO: 4. when aligned with a numerous Termamyl-like  $\alpha$ -amylases. Thus, for example, the corresponding positions of these residues in the amino acid sequences of a number of Termamyl-like  $\alpha$ -amylases which have already been mentioned (vide supra) are as follows:



TABLE 2

Termamyl-like $\alpha$ -amylase	Pair wise amino acid deletions among
<i>B. amyloliquefaciens</i> (SEQ ID No. 5)	R176, G177, E178, G179
<i>B. stearothermophilus</i> (SEQ ID No. 3)	R179, G180, I181, G182
Bacillus WO 95/26397 (SEQ ID No. 2)	R181, G182, T183, G184
Bacillus WO 95/26397 (SEQ ID No. 1)	R181, G182, D183, G184
"Bacillus sp. #707" (SEQ ID No. 6)	R181, G182, H183, G184

When using SEQ ID NO: 1 to SEQ ID NO: 6 as the backbone (i.e. as the parent Termamyl-like  $\alpha$ -amylase) two, three, four, five or six mutations may according to the invention be made in the following regions/positions to increase the thermostability at acidic pH and/or at low  $\text{Ca}^{2+}$  concentrations (relative to the parent):

(relative to Seq ID NO: 1 herein):

- 1: R181\*, G182\*, T183\*, G184\*
- 2: N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 3: V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- 4: E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5: E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 6: K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;

(relative to SEQ ID NO: 2 herein):

- 1: R181\*, G182\*, D183\*, G184\*
- 2: N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 3: V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- 4: E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5: E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 6: K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;

(Relative to SEQ ID NO: 3 herein):

- 1: R179\*, G180,I181\*, G182\*
- 2: N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 3: L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,Y,V;
- 4: E210A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5: E214A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 6: S267A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,Y,V

Relative to SEQ ID NO: 4 herein):

- 1: Q178\*, G179\*
- 2: N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 3: I201A,R,D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,Y,V;
- 4: D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5: E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 6: Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;

(relative to SEQ ID NO: 5 herein):

- 1: R176\*, G177\*, E178,G179\*
- 2: N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 3: V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- 4: D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5: E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 6: Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;

(relative to SEQ ID NO: 6 herein):

- 1: R181\*, G182\*, H183\*, G184\*
- 2: N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 3: I206A,R,D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,Y,V;
- 4: E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5: E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 6: K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V.

Comtemplated according to the present invention is combining three, four, five or six mutation.

Specific double mutations for backbone SEQ ID NO: 1 to SEQ ID NO: 6 are listed in the following.

Using SEQ ID NO: 1 as the backbone the following double mutantions resulting in the desired effect are contemplated according to the invention:

- R181\*/G182\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- G182\*/T183\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 5 -G183\*/G184\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- R181\*/G182\*/V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- G182\*/T183\*/V206A,R, D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- 10 -T183\*/G184\*/V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- R181\*/G182\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 15 -G182\*/T183\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- T183\*/G184\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- R181\*/G182\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 20 -G182\*/T183\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- T183\*/G184\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 25 -R181\*/G182\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- G182\*/T183\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- T183\*/G184\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- 30 -N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 35 -N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 40 -V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- 45 -E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;
- 50 Using SEQ ID NO: 2 as the backbone the following double mutantions resulting in the desired effect are contemplated according to the invention:
- R181\*/G182\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 55 -G182\*/D183\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- D183\*/G184\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;
- 60 -R181\*/G182\*/V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- G182\*/T183\*/V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- G183\*/G184\*/V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;
- 65 -R181\*/G182\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;



-G182\*/T183\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-T183\*/G184\*/E212A,R,D,N,C,Q,G,H,I,L,K,N,F,P,S,T,W,  
Y,V;  
-R181\*/G182\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G182\*/T183\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-T183\*/G184\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R181\*/G182\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,  
Y,V;  
-G182\*/T183\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,  
Y,V;  
-T183\*/G184\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,  
Y,V;  
-N195 A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/V206A,R,  
D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;  
-N195 A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E212A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N/95A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
-V216A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/E212A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-V206 A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E  
216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-V206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
-E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
-E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
Using SEQ ID NO. 3 as the backbone the following  
double mutations resulting in the desired effect are  
contemplated according to the invention:  
-R179\*/G180\*/N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-I180\*/I181\*/N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,  
V;  
-I181\*/G182\*/N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R179\*/G180\*/L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,  
Y,V;  
-G180\*/I181\*/L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,  
Y,V;  
-I181\*/G182\*/L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,  
Y,V;  
-R179\*/G180\*/E210A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G180\*/I181\*/E210A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-I181\*/G182\*/E210A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R179\*/G180\*/E214A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G180\*/I181\*/E214A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-I181\*/G182\*/E214A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R179\*/G180\*/S267A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,  
Y,V;  
-G180\*/I181\*/S267A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,  
Y,V;

-I181\*/I182\*/S267A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,Y,  
V;  
-N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/L204A,R,  
D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,Y,V;  
-N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E210A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E214A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/S267A,R,  
D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,Y,V;  
-L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,Y,V/E210A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,Y,V/E214A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-L204A,R,D,N,C,E,Q,G,H,I,K,M,F,P,S,T,W,Y,V/S267A,R,  
D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,Y,V;  
-E210A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-E210A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/S267A,R,  
D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,Y,V;  
-E214A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/S267A,R,  
D,N,C,E,Q,G,H,I,L,K,M,F,P,T,W,Y,V;  
Using SEQ ID NO. 4 as the backbone the following  
double mutations resulting in the desired effect are  
contemplated according to the invention:  
-Q178\*/G179\*/N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-Q178\*/G179\*/I201A,R,D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,  
Y,V;  
-Q178\*/G179\*/D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-Q178\*/G179\*/E201A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R179\*/G179\*/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-N190/I201A,R,D,N,C,E,Q,G,H,L,K,N,F,P,S,T,W,Y,V;  
-N190/D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N190/E201A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N190/Q264A,R,D,N,C,E,G,H,I,L,K,N,F,P,S,T,W,Y,V;  
-N201/D207A,R,N,C,Q,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-I201/E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-I201/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-D207/E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-D207/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-E211/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
Using SEQ ID NO: 5 as the backbone the following  
double mutations resulting in the desired effect are  
contemplated according to the invention:  
-R176\*/G177\*/N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G177\*/E178\*/N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-E178\*/G179\*/N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R176\*/G177\*/V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,  
W,Y;  
-G176\*/E178\*/V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,  
W,Y;  
-E178\*/G179\*/V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,  
W,Y;  
-R176\*/G177\*/D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G177\*/E178\*/D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-E178\*/G179\*/D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R176\*/G177\*/E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;



-G177\*/E178\*/E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-E178\*/G179\*/E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R176\*/G177\*/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G177\*/E178\*/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-E178\*/G179\*/Q264A,R,D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/V201A,R,  
D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y;  
-N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/D207A,R,  
N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E211A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/Q264A,R,  
D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/D207A,R,  
N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/E211A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-V201A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y/Q264A,R,  
D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E211A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-D207A,R,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/Q264A,R,  
D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-E211A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/Q264A,R,  
D,N,C,E,G,H,I,L,K,M,F,P,S,T,W,Y,V.  
Using SEQ ID NO: 6 as the backbone the following  
double mutations resulting in the desired effect are con-  
templated according to the invention:  
-R181\*/G182\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G182\*/H183\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-H183\*/G184\*/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R181\*/G182\*/I206A,R,D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,  
Y,V;  
-G182\*/H183\*/I206A,R,D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,  
Y,V;  
-H183\*/G184\*/I206A,R,D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,  
Y,V;  
-R181\*/G182\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G182\*/H183\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-H183\*/G184\*/E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R181\*/G182\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-G182\*/H183\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-H183\*/G184\*/E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,  
Y,V;  
-R181\*/G182\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,  
Y,V;  
-G182\*/H183\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,  
Y,V;  
-H183\*/G184\*/K269A,R,D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,  
Y,V;  
-N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/I206A,R,  
D,N,C,E,Q,G,H,L,K,M,F,P,S,T,W,Y,V;  
-N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E212A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;

/N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
-I206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E212A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-I206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-I206A,R,D,N,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,P,  
D,N,C,E,Q,C,H,I,L,M,F,P,S,T,W,Y,V;  
-E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/E216A,R,  
D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;  
-E212A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
-E216A,R,D,N,C,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V/K269A,R,  
D,N,C,E,Q,G,H,I,L,M,F,P,S,T,W,Y,V;  
All Termamyl-like  $\alpha$ -amylase defined above may suitably  
be used as backbone for preparing variants of the invention.  
However, in a preferred embodiment the variant com-  
prises the following mutations: N190F/Q264S in SEQ ID  
NO: 4 or in corresponding positions in another parent  
Termamyl-like  $\alpha$ -amylases.  
In another embodiment the variant of the invention com-  
prises the following mutations: I181\*/G182\*/N193F in SEQ  
ID NO: 3 (TVB146) or in corresponding positions in another  
parent Termamyl-like  $\alpha$ -amylases. Said variant may further  
comprise a substitution in position E214Q.  
In a preferred embodiment of the invention the parent  
Termamyl-like  $\alpha$ -amylase is a hybrid  $\alpha$ -amylase of SEQ ID  
NO: 4 and SEQ ID NO: 5. Specifically, the parent hybrid  
Termamyl-like  $\alpha$ -amylase may be a hybrid alpha-amylase  
comprising the 445 C-terminal amino acid residues of the *B.*  
*licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4 and the 37  
N-terminal amino acid residues of the  $\alpha$ -amylase derived  
from *B. amyloliquefaciens* shown in SEQ ID NO: 5, which  
may suitably further have the following mutations: H156Y+  
A181T+N190F+A209V+Q264S (using the numbering in  
SEQ ID NO: 4). The latter mentioned hybrid is used in the  
examples below and is referred to as LE174.  
General Mutations in Variants of the Invention  
It may be preferred that a variant of the invention com-  
prises one or more modifications in addition to those out-  
lined above. Thus, it may be advantageous that one or more  
proline residues present in the part of the  $\alpha$ -amylase variant  
which is modified is/are replaced with a non-proline residue  
which may be any of the possible, naturally occurring  
non-proline residues, and which preferably is an alanine,  
glycine, serine, threonine, valine or leucine.  
Analogously, it may be preferred that one or more cyste-  
ine residues present among the amino acid residues with  
which the parent  $\alpha$ -amylase is modified is/are replaced with  
a non-cysteine residue such as serine, alanine, threonine,  
glycine, valine or leucine.  
Furthermore, a variant of the invention may—either as the  
only modification or in combination with any of the above  
outlined modifications—be modified so that one or more  
Asp and/or Glu present in an amino acid fragment corre-  
sponding to the amino acid fragment 185–209 of SEQ ID  
NO: 4 is replaced by an Asn and/or Gln, respectively. Also  
of interest is the replacement, in the Termamyl-like  
 $\alpha$ -amylase, of one or more of the Lys residues present in an  
amino acid fragment corresponding to the amino acid frag-  
ment 185–209 of SEQ ID NO: 4 by an Arg.  
It will be understood that the present invention encom-  
passes variants incorporating two or more of the above  
outlined modifications.  
Furthermore, it may be advantageous to introduce point-  
mutations in any of the variants described herein.



### Methods for Preparing $\alpha$ -amylase Variants

Several methods for introducing mutations into genes are known in the art. After a brief discussion of the cloning of  $\alpha$ -amylase-encoding DNA sequences, methods for generating mutations at specific sites within the  $\alpha$ -amylase-encoding sequence will be discussed.

#### Cloning a DNA Sequence Encoding an $\alpha$ -amylase

The DNA sequence encoding a parent  $\alpha$ -amylase may be isolated from any cell or microorganism producing the  $\alpha$ -amylase in question, using various methods well known in the art. First, a genomic DNA and/or cDNA library should be constructed using chromosomal DNA or messenger RNA from the organism that produces the  $\alpha$ -amylase to be studied. Then, if the amino acid sequence of the  $\alpha$ -amylase is known, homologous, labelled oligonucleotide probes may be synthesized and used to identify  $\alpha$ -amylase-encoding clones from a genomic library prepared from the organism in question. Alternatively, a labelled oligonucleotide probe containing sequences homologous to a known  $\alpha$ -amylase gene could be used as a probe to identify  $\alpha$ -amylase-encoding clones, using hybridization and washing conditions of lower stringency.

Yet another method for identifying  $\alpha$ -amylase-encoding clones would involve inserting fragments of genomic DNA into an expression vector, such as a plasmid, transforming  $\alpha$ -amylase-negative bacteria with the resulting genomic DNA library, and then plating the transformed bacteria onto agar containing a substrate for  $\alpha$ -amylase, thereby allowing clones expressing the  $\alpha$ -amylase to be identified.

Alternatively, the DNA sequence encoding the enzyme may be prepared synthetically by established standard methods, e.g. the phosphoramidite method described by S. L. Beaucage and M. H. Caruthers (1981) or the method described by Matthes et al. (1984). In the phosphoramidite method, oligonucleotides are synthesized, e.g. in an automatic DNA synthesizer, purified, annealed, ligated and cloned in appropriate vectors.

Finally, the DNA sequence may be of mixed genomic and synthetic origin, mixed synthetic and cDNA origin or mixed genomic and cDNA origin, prepared by ligating fragments of synthetic, genomic or cDNA origin (as appropriate, the fragments corresponding to various parts of the entire DNA sequence), in accordance with standard techniques. The DNA sequence may also be prepared by polymerase chain reaction (PCR) using specific primers, for instance as described in U.S. Pat. No. 4,683,202 or R. K. Saiki et al. (1988).

#### Site-directed Mutagenesis

Once an  $\alpha$ -amylase-encoding DNA sequence has been isolated, and desirable sites for mutation identified, mutations may be introduced using synthetic oligonucleotides. These oligonucleotides contain nucleotide sequences flanking the desired mutation sites; mutant nucleotides are inserted during oligonucleotide synthesis. In a specific method, a single-stranded gap of DNA, bridging the  $\alpha$ -amylase-encoding sequence, is created in a vector carrying the  $\alpha$ -amylase gene. Then the synthetic nucleotide, bearing the desired mutation, is annealed to a homologous portion of the single-stranded DNA. The remaining gap is then filled in with DNA polymerase I (Klenow fragment) and the construct is ligated using T4 ligase. A specific example of this method is described in Morinaga et al. (1984). U.S. Pat. No. 4,760,025 discloses the introduction of oligonucleotides encoding multiple mutations by performing minor alterations of the cassette. However, an even greater variety of mutations can be introduced at any one time by the Morinaga method, because a multitude of

oligonucleotides, of various lengths, can be introduced. Another method for introducing mutations into  $\alpha$ -amylase-encoding DNA sequences is described in Nelson and Long (1989). It involves the 3-step generation of a PCR fragment containing the desired mutation introduced by using a chemically synthesized DNA strand as one of the primers in the PCR reactions. From the PCR-generated fragment, a DNA fragment carrying the mutation may be isolated by cleavage with restriction endonucleases and reinserted into an expression plasmid.

#### Random Mutagenesis

Random mutagenesis is suitably performed either as localised or region-specific random mutagenesis in at least three parts of the gene translating to the amino acid sequence shown in question, or within the whole gene.

The random mutagenesis of a DNA sequence encoding a parent  $\alpha$ -amylase may be conveniently performed by use of any method known in the art.

In relation to the above, a further aspect of the present invention relates to a method for generating a variant of a parent  $\alpha$ -amylase, e.g. wherein the variant exhibits altered or increased thermal stability relative to the parent, the method comprising:

- (a) subjecting a DNA sequence encoding the parent  $\alpha$ -amylase to random mutagenesis,
- (b) expressing the mutated DNA sequence obtained in step (a) in a host cell, and
- (c) screening for host cells expressing an  $\alpha$ -amylase variant which has an altered property (i.e. thermal stability) relative to the parent  $\alpha$ -amylase.

Step (a) of the above method of the invention is preferably performed using doped primers.

For instance, the random mutagenesis may be performed by use of a suitable physical or chemical mutagenizing agent, by use of a suitable oligonucleotide, or by subjecting the DNA sequence to PCR generated mutagenesis. Furthermore, the random mutagenesis may be performed by use of any combination of these mutagenizing agents. The mutagenizing agent may, e.g., be one which induces transitions, transversions, inversions, scrambling, deletions, and/or insertions.

Examples of a physical or chemical mutagenizing agent suitable for the present purpose include ultraviolet (UV) ir-radiation, hydroxylamine, N-methyl-N'-nitro-N-nitrosoguanidine (MNNG), O-methyl hydroxylamine, nitrous acid, ethyl methane sulphonate (EMS), sodium bisulphite, formic acid, and nucleotide analogues. When such agents are used, the mutagenesis is typically performed by incubating the DNA sequence encoding the parent enzyme to be mutagenized in the presence of the mutagenizing agent of choice under suitable conditions for the mutagenesis to take place, and selecting for mutated DNA having the desired properties.

When the mutagenesis is performed by the use of an oligonucleotide, the oligonucleotide may be doped or spiked with the three non-parent nucleotides during the synthesis of the oligonucleotide at the positions which are to be changed. The doping or spiking may be done so that codons for unwanted amino acids are avoided. The doped or spiked oligonucleotide can be incorporated into the DNA encoding the  $\alpha$ -amylase enzyme by any published technique, using e.g. PCR, LCR or any DNA polymerase and ligase as deemed appropriate.

Preferably, the doping is carried out using "constant random doping", in which the percentage of wild-type and mutation in each position is predefined. Furthermore, the doping may be directed toward a preference for the intro-



duction of certain nucleotides, and thereby a preference for the introduction of one or more specific amino acid residues. The doping may be made, e.g., so as to allow for the introduction of 90% wild type and 10% mutations in each position. An additional consideration in the choice of a doping scheme is based on genetic as well as protein-structural constraints. The doping scheme may be made by using the DOPE program which, inter alia, ensures that introduction of stop codons is avoided.

When PCR-generated mutagenesis is used, either a chemically treated or non-treated gene encoding a parent  $\alpha$ -amylase is subjected to PCR under conditions that increase the mis-incorporation of nucleotides (Deshler 1992; Leung et al., Technique, Vol. 1, 1989, pp. 11–15).

A mutator strain of *E. coli* (Fowler et al., Molec. Gen. Genet., 133, 1974, pp. 179–191), *S. cerevisiae* or any other microbial organism may be used for the random mutagenesis of the DNA encoding the  $\alpha$ -amylase by, e.g., transforming a plasmid containing the parent glycosylase into the mutator strain, growing the mutator strain with the plasmid and isolating the mutated plasmid from the mutator strain. The mutated plasmid may be subsequently transformed into the expression organism.

The DNA sequence to be mutagenized may be conveniently present in a genomic or cDNA library prepared from an organism expressing the parent  $\alpha$ -amylase. Alternatively, the DNA sequence may be present on a suitable vector such as a plasmid or a bacteriophage, which as such may be incubated with or otherwise exposed to the mutagenising agent. The DNA to be mutagenized may also be present in a host cell either by being integrated in the genome of said cell or by being present on a vector harboured in the cell. Finally, the DNA to be mutagenized may be in isolated form. It will be understood that the DNA sequence to be subjected to random mutagenesis is preferably a cDNA or a genomic DNA sequence.

In some cases it may be convenient to amplify the mutated DNA sequence prior to performing the expression step b) or the screening step c). Such amplification may be performed in accordance with methods known in the art, the presently preferred method being PCR-generated amplification using oligonucleotide primers prepared on the basis of the DNA or amino acid sequence of the parent enzyme.

Subsequent to the incubation with or exposure to the mutagenising agent, the mutated DNA is expressed by culturing a suitable host cell carrying the DNA sequence under conditions allowing expression to take place. The host cell used for this purpose may be one which has been transformed with the mutated DNA sequence, optionally present on a vector, or one which was carried the DNA sequence encoding the parent enzyme during the mutagenesis treatment. Examples of suitable host cells are the following: gram positive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, *Streptomyces lividans* or *Streptomyces murinus*; and gram-negative bacteria such as *E. coli*.

The mutated DNA sequence may further comprise a DNA sequence encoding functions permitting expression of the mutated DNA sequence.

#### Localized Random Mutagenesis

The random mutagenesis may be advantageously localized to a part of the parent  $\alpha$ -amylase in question. This may, e.g., be advantageous when certain regions of the enzyme have been identified to be of particular importance for a

given property of the enzyme, and when modified are expected to result in a variant having improved properties. Such regions may normally be identified when the tertiary structure of the parent enzyme has been elucidated and related to the function of the enzyme.

The localized, or region-specific, random mutagenesis is conveniently performed by use of PCR generated mutagenesis techniques as described above or any other suitable technique known in the art. Alternatively, the DNA sequence encoding the part of the DNA sequence to be modified may be isolated, e.g., by insertion into a suitable vector, and said part may be subsequently subjected to mutagenesis by use of any of the mutagenesis methods discussed above.

#### Alternative Methods of Providing $\alpha$ -amylase Variants

Alternative methods for providing variants of the invention include gene shuffling method known in the art including the methods e.g. described in WO 95/22625 (from Affymax Technologies N.V.) and WO 96/00343 (from Novo Nordisk A/S).

#### Expression of $\alpha$ -amylase Variants

According to the invention, a DNA sequence encoding the variant produced by methods described above, or by any alternative methods known in the art, can be expressed, in enzyme form, using an expression vector which typically includes control sequences encoding a promoter, operator, ribosome binding site, translation initiation signal, and, optionally, a repressor gene or various activator genes.

The recombinant expression vector carrying the DNA sequence encoding an  $\alpha$ -amylase variant of the invention may be any vector which may conveniently be subjected to recombinant DNA procedures, and the choice of vector will often depend on the host cell into which it is to be introduced. Thus, the vector may be an autonomously replicating vector, i.e. a vector which exists as an extrachromosomal entity, the replication of which is independent of chromosomal replication, e.g. a plasmid, a bacteriophage or an extrachromosomal element, minichromosome or an artificial chromosome. Alternatively, the vector may be one which, when introduced into a host cell, is integrated into the host cell genome and replicated together with the chromosome(s) into which it has been integrated.

In the vector, the DNA sequence should be operably connected to a suitable promoter sequence. The promoter may be any DNA sequence which shows transcriptional activity in the host cell of choice and may be derived from genes encoding proteins either homologous or heterologous to the host cell. Examples of suitable promoters for directing the transcription of the DNA sequence encoding an  $\alpha$ -amylase variant of the invention, especially in a bacterial host, are the promoter of the lac operon of *E. coli*, the *Streptomyces coelicolor* agarase gene *dagA* promoters, the promoters of the *Bacillus licheniformis*  $\alpha$ -amylase gene (*amyL*), the promoters of the *Bacillus stearothermophilus* maltogenic amylase gene (*amyM*), the promoters of the *Bacillus amyloliquefaciens*  $\alpha$ -amylase (*amyQ*), the promoters of the *Bacillus subtilis* *xylA* and *xylB* genes etc. For transcription in a fungal host, examples of useful promoters are those derived from the gene encoding *A. oryzae* TAKA amylase, *Rhizomucor miehei* aspartic proteinase, *A. niger* neutral  $\alpha$ -amylase, *A. niger* acid stable  $\alpha$ -amylase, *A. niger* glucoamylase, *Rhizomucor miehei* lipase, *A. oryzae* alkaline protease, *A. oryzae* triose phosphate isomerase or *A. nidulans* acetamidase.

The expression vector of the invention may also comprise a suitable transcription terminator and, in eukaryotes, polyadenylation sequences operably connected to the DNA sequence encoding the  $\alpha$ -amylase variant of the invention.



Termination and polyadenylation sequences may suitably be derived from the same sources as the promoter.

The vector may further comprise a DNA sequence enabling the vector to replicate in the host cell in question. Examples of such sequences are the origins of replication of plasmids pUC19, pACYC177, pUB110, pE194, pAMB1 and pIJ702.

The vector may also comprise a selectable marker, e.g. a gene the product of which complements a defect in the host cell, such as the *dal* genes from *B. subtilis* or *B. licheniformis*, or one which confers antibiotic resistance such as ampicillin, kanamycin, chloramphenicol or tetracycline resistance. Furthermore, the vector may comprise *Aspergillus* selection markers such as *amdS*, *argB*, *niaD* and *sC*, a marker giving rise to hygromycin resistance, or the selection may be accomplished by co-transformation, e.g. as described in WO 91/17243.

While intracellular expression may be advantageous in some respects, e.g. when using certain bacteria as host cells, it is generally preferred that the expression is extracellular. In general, the *Bacillus*  $\alpha$ -amylases mentioned herein comprise a preregion permitting secretion of the expressed protease into the culture medium. If desirable, this preregion may be replaced by a different preregion or signal sequence, conveniently accomplished by substitution of the DNA sequences encoding the respective preregions.

The procedures used to ligate the DNA construct of the invention encoding an  $\alpha$ -amylase variant, the promoter, terminator and other elements, respectively, and to insert them into suitable vectors containing the information necessary for replication, are well known to persons skilled in the art (cf., for instance, Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, 1989).

The cell of the invention, either comprising a DNA construct or an expression vector of the invention as defined above, is advantageously used as a host cell in the recombinant production of an  $\alpha$ -amylase variant of the invention. The cell may be transformed with the DNA construct of the invention encoding the variant, conveniently by integrating the DNA construct (in one or more copies) in the host chromosome. This integration is generally considered to be an advantage as the DNA sequence is more likely to be stably maintained in the cell. Integration of the DNA constructs into the host chromosome may be performed according to conventional methods, e.g. by homologous or heterologous recombination. Alternatively, the cell may be transformed with an expression vector as described above in connection with the different types of host cells.

The cell of the invention may be a cell of a higher organism such as a mammal or an insect, but is preferably a microbial cell, e.g. a bacterial or a fungal (including yeast) cell.

Examples of suitable bacteria are grampositive bacteria such as *Bacillus subtilis*, *Bacillus licheniformis*, *Bacillus lentus*, *Bacillus brevis*, *Bacillus stearothermophilus*, *Bacillus alkalophilus*, *Bacillus amyloliquefaciens*, *Bacillus coagulans*, *Bacillus circulans*, *Bacillus lautus*, *Bacillus megaterium*, *Bacillus thuringiensis*, or *Streptomyces lividans* or *Streptomyces murinus*, or gramnegative bacteria such as *E. coli*. The transformation of the bacteria may, for instance, be effected by protoplast transformation or by using competent cells in a manner known per se.

The yeast organism may favourably be selected from a species of *Saccharomyces* or *Schizosaccharomyces*, e.g. *Saccharomyces cerevisiae*. The filamentous fungus may advantageously belong to a species of *Aspergillus*, e.g.

*Aspergillus oryzae* or *Aspergillus niger*. Fungal cells may be transformed by a process involving protoplast formation and transformation of the protoplasts followed by regeneration of the cell wall in a manner known per se. A suitable procedure for transformation of *Aspergillus* host cells is described in EP 238 023.

In yet a further aspect, the present invention relates to a method of producing an  $\alpha$ -amylase variant of the invention, which method comprises cultivating a host cell as described above under conditions conducive to the production of the variant and recovering the variant from the cells and/or culture medium.

The medium used to cultivate the cells may be any conventional medium suitable for growing the host cell in question and obtaining expression of the  $\alpha$ -amylase variant of the invention. Suitable media are available from commercial suppliers or may be prepared according to published recipes (e.g. as described in catalogues of the American Type Culture Collection).

The  $\alpha$ -amylase variant secreted from the host cells may conveniently be recovered from the culture medium by well-known procedures, including separating the cells from the medium by centrifugation or filtration, and precipitating proteinaceous components of the medium by means of a salt such as ammonium sulphate, followed by the use of chromatographic procedures such as ion exchange chromatography, affinity chromatography, or the like.

#### Industrial Applications

The  $\alpha$ -amylase variants of this invention possess valuable properties allowing for a variety of industrial applications. In particular, enzyme variants of the invention are applicable as a component in washing, dishwashing and hard-surface cleaning detergent compositions. Numerous variants are particularly useful in the production of sweeteners and ethanol from starch, and/or for textile desizing. Conditions for conventional starch-conversion processes, including starch liquefaction and/or saccharification processes, are described in, e.g., U.S. Pat. No. 3,912,590 and in EP patent publications Nos. 252 730 and 63 909.

#### Production of Sweeteners from Starch:

A "traditional" process for conversion of starch to fructose syrups normally consists of three consecutive enzymatic processes, viz. a liquefaction process followed by a saccharification process and an isomerization process. During the liquefaction process, starch is degraded to dextrins by an  $\alpha$ -amylase (e.g. Termamyl™) at pH values between 5.5 and 6.2 and at temperatures of 95–160° C. for a period of approx. 2 hours. In order to ensure an optimal enzyme stability under these conditions, 1 mM of calcium is added (40 ppm free calcium ions).

After the liquefaction process the dextrins are converted into dextrose by addition of a glucoamylase (e.g. AMG™) and a debranching enzyme, such as an isoamylase or a pullulanase (e.g. Promozyme™). Before this step the pH is reduced to a value below 4.5, maintaining the high temperature (above 95° C.), and the liquefying  $\alpha$ -amylase activity is denatured. The temperature is lowered to 60° C., and glucoamylase and debranching enzyme are added. The saccharification process proceeds for 24–72 hours.

After the saccharification process the pH is increased to a value in the range of 6–8, preferably pH 7.5, and the calcium is removed by ion exchange. The dextrose syrup is then converted into high fructose syrup using, e.g., an immobilized glucoseisomerase (such as Sweetzyme™).

At least 1 enzymatic improvements of this process could be envisaged. Reduction of the calcium dependency of the liquefying  $\alpha$ -amylase. Addition of free calcium is required



to ensure adequately high stability of the  $\alpha$ -amylase, but free calcium strongly inhibits the activity of the glucosyltransferase and needs to be removed, by means of an expensive unit operation, to an extent which reduces the level of free calcium to below 3–5 ppm. Cost savings could be obtained if such an operation could be avoided and the liquefaction process could be performed without addition of free calcium ions.

To achieve that, a less calcium-dependent Termamyl-like  $\alpha$ -amylase which is stable and highly active at low concentrations of free calcium (<40 ppm) is required. Such a Termamyl-like  $\alpha$ -amylase should have a pH optimum at a pH in the range of 4.5–6.5, preferably in the range of 4.5–5.5.

#### Detergent Compositions

As mentioned above, variants of the invention may suitably be incorporated in detergent compositions. Increased thermostability at low calcium concentrations would be very beneficial for amylase performance in detergents, i.e. the alkaline region. Reference is made, for example, to WO 96/23874 and WO 97/07202 for further details concerning relevant ingredients of detergent compositions (such as laundry or dishwashing detergents), appropriate methods of formulating the variants in such detergent compositions, and for examples of relevant types of detergent compositions.

Detergent compositions comprising a variant of the invention may additionally comprise one or more other enzymes, such as a lipase, cutinase, protease, cellulase, peroxidase or laccase, and/or another  $\alpha$ -amylase.

$\alpha$ -amylase variants of the invention may be incorporated in detergents at conventionally employed concentrations. It is at present contemplated that a variant of the invention may be incorporated in an amount corresponding to 0.00001–1 mg (calculated as pure, active enzyme protein) of  $\alpha$ -amylase per liter of wash/dishwash liquor using conventional dosing levels of detergent.

The invention also relates to a composition comprising a mixture of one or more variants of the invention derived from (as the parent Termamyl-like  $\alpha$ -amylase) the *B. stearo-thermophilus*  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 3 and a Termamyl-like alpha-amylase derived from the *B. licheniformis*  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 4.

Further, the invention also relates to a composition comprising a mixture of one or more variants according to the invention derived from (as the parent Termamyl-like  $\alpha$ -amylase) the *B. stearo-thermophilus*  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 3 and a hybrid alpha-amylase comprising a part of the *B. amyloliquefaciens*  $\alpha$ -amylase shown in SEQ ID NO: 5 and a part of the *B. licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4. The latter mentioned hybrid Termamyl-like  $\alpha$ -amylase comprises the 445 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the  $\alpha$ -amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 5. Said latter mentioned hybrid  $\alpha$ -amylase may suitably comprise the following mutations: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4). In the examples below said hybrid parent Termamyl-like  $\alpha$ -amylase, is used in combination with variants of the invention, which variants may be used in compositions of the invention.

In a specific embodiment of the invention the composition comprises a mixture of TVB146 and LE174, e.g., in a ratio of 2:1 to 1:2, such as 1:1.

A  $\alpha$ -amylase variant of the invention or a composition of the invention may in an aspect of the invention be used for

washing and/or dishwashing; for textile desizing or for starch liquefaction.

#### MATERIALS AND METHODS

##### 5 Enzymes:

BSG alpha-amylase: *B. stearo-thermophilus* alpha-amylase depicted in SEQ ID NO: 3. TVB146 alpha-amylase variant: *B. stearo-thermophilus* alpha-amylase variant depicted in SEQ ID NO: 3 with the following mutations: with the deletion in positions I181-G182+N193F. LE174 hybrid alpha-amylase variant: LE174 is a hybrid Termamyl-like alpha-amylase being identical to the Termamyl sequence, i.e., the *Bacillus licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4, except that the N-terminal 35 amino acid residues (of the mature protein) has been replaced by the N-terminal 33 residues of BAN (mature protein), i.e., the *Bacillus amyloliquefaciens* alpha-amylase shown in SEQ ID NO: 5, which further have following mutations: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4). LE174 was constructed by SOE-PCR (Higuchi et al. 1988, Nucleic Acids Research 16:7351).

##### 15 Fermentation and Purification of $\alpha$ -amylase Variants

A *B. subtilis* strain harbouring the relevant expression plasmid is streaked on a LB-agar plate with 10  $\mu$ g/ml kanamycin from  $-80^{\circ}$  C. stock, and grown overnight at  $37^{\circ}$  C. The colonies are transferred to 100 ml BPX media supplemented with 10  $\mu$ g/ml kanamycin in a 500 ml shaking flask.

##### 25 Composition of BPX medium:

Potato starch	100 g/l
Barley flour	50 g/l
BAN 5000 SKB	0.1 g/l
Sodium caseinate	10 g/l
Soy Bean Meal	20 g/l
Na <sub>2</sub> HPO <sub>4</sub> · 12 H <sub>2</sub> O	9 g/l
Pluronic™	0.1 g/l

30 The culture is shaken at  $37^{\circ}$  C. at 270 rpm for 5 days.

Cells and cell debris are removed from the fermentation broth by centrifugation at 4500 rpm in 20–25 minutes. Afterwards the supernatant is filtered to obtain a completely clear solution. The filtrate is concentrated and washed on a UF-filter (10000 cut off membrane) and the buffer is changed to 20 mM Acetate pH 5.5. The UF-filtrate is applied on a S-sepharose F.F. and elution is carried out by step elution with 0.2M NaCl in the same buffer. The eluate is dialysed against 10 mM Tris, pH 9.0 and applied on a Q-sepharose F.F. and eluted with a linear gradient from 0–0.3M NaCl over 6 column volumes. The fractions which contain the activity (measured by the Phadebas assay) are pooled, pH was adjusted to pH 7.5 and remaining color was removed by a treatment with 0.5% W/vol. active coal in 5 minutes.

##### 55 Activity Determination—(KNU)

One Kilo alpha-amylase Unit (1 KNU) is the amount of enzyme which breaks down 5.26 g starch (Merck, Amylum Solubile, Erg. B 6, Batch 9947275) per hour in Novo Nordisk's standard method for determination of alpha-amylase based upon the following condition:

65 Substrate	soluble starch
Calcium content in solvent	0.0043M



-continued

Reaction time	7–20 minutes
Temperature	37° C.
pH	5.6

Detailed description of Novo Nordisk's analytical method (AF 9) is available on request.

### BS-amylase Activity Determination—KNU(S)

#### 1. Application Field

This method is used to determine  $\alpha$ -amylase activity in fermentation and recovery samples and formulated and granulated products.

#### 2. Principle

BS-amylase breaks down the substrate (4,6-ethylidene ( $G_7$ )-p-nitrophenyl ( $G_1$ )- $\alpha$ ,D-maltoheptaoside (written as ethylidene- $G_7$ -PNP) into, among other things,  $G_2$ -PNP and  $G_3$ -PNP, where G denoted glucose and PNP p-nitrophenol.

$G_2$ -PNP and  $G_3$ -PNP are broken down by  $\alpha$ -glucosidase, which is added in excess, into glucose and the yellow-coloured p-nitrophenol.

The colour reaction is monitored in situ and the change in absorbance over time calculated as an expression of the speed of the reaction and thus of the activity of the enzyme. See the Boehringer Mannheim 1442 309 guidelines for further details.

#### 2.1 Reaction conditions

<u>Reaction:</u>	
Temperature	37° C.
pH	7.1
Pre-incubation time	2 minutes
<u>Detection:</u>	
Wavelength	405 nm
Measurement time	3 minutes

#### 3. Definition of Units

*Bacillus stearothermophilus* alpha-amylase (BS-amylase) activity is determined relative to a standard of declared activity and stated in Kilo Novo Units (Stearothermophilus) or KNU(S)).

#### 4. Specificity and Sensitivity

Limit of determination: approx. 0.4 KNU(s)/g

#### 5. Apparatus

Cobas Fara analyser  
Diluted (e.g. Hamilton Microlab 1000)  
Analytical balance (e.g. Mettler AE 100)  
Stirrer plates

#### 6. Reagents/Substrates

A ready-made kit is used in this analysis to determine  $\alpha$ -amylase activity. Note that the reagents specified for the substrate and  $\alpha$ -glucosidase are not used as described in the Boehringer Mannheim guidelines. However, the designations "buffer", "glass 1", glass 1a" and Glass 2" are those referred to in those guidelines.

##### 6.1. Substrate

4,6-ethylidene( $G_7$ )-p-nitrophenyl( $G_1$ )- $\alpha$ ,D-maltoheptaoside (written as ethylidene- $G_7$ -PNP) e.g. Boehringer Mannheim 1442 309

### 6.2 $\alpha$ -glucosidase help reagent $\alpha$ -glucosidase, e.g. Boehringer Mannheim 1442 309

#### 6.3 BRIJ 35 solution

BRIJ 35 (30% W/V Sigma 430 AG-6)	1000 mL
Demineralized water	up to 2,000 mL

#### 6.4 Stabiliser

Brij 35 solution	33 mL
CaCl <sub>2</sub> *2H <sub>2</sub> O (Merck 2382)	882 g
Demineralized water	up to 2,000 mL

### 7. Samples and Standards

#### 7.1 Standard curve

Example: Preparation of BS-amylase standard curve

The relevant standard is diluted to 0.60 KNU(s)/mL as follows. A calculated quantity of standard is weighed out and added to 200 mL volumetric flask, which is filled to around the  $\frac{2}{3}$  mark with demineralized water. Stabiliser corresponding to 1% of the volume of the flask is added and the flask is filled to the mark with demineralized water.

A Hamilton Microlab 1000 is used to produce the dilutions shown below. Demineralized water with 1% stabiliser is used as the diluent.

Dilution No.	Enzyme stock solution	1% stabiliser	KNU(s)/mL
1	20 $\mu$ L	580 $\mu$ L	0.02
2	30 $\mu$ L	570 $\mu$ L	0.03
3	40 $\mu$ L	560 $\mu$ L	0.04
4	50 $\mu$ L	550 $\mu$ L	0.05
5	60 $\mu$ L	540 $\mu$ L	0.06

#### 7.2 Level control

A Novo Nordisk A/S BS amylase level control is included in all runs using the Cobas Fara. The control is diluted with 1% stabiliser so that the final dilution is within the range of the standard curve. All weights and dilutions are noted on the worklist

#### 7.3 Sample solutions

##### Single determination

Fermentation samples (not final samples) from production, all fermentation samples from pilot plants and storage stability samples are weighed out and analyzed once only.

##### Double determination over 1 run:

Process samples, final fermentation samples from production, samples from GLP studies and R&D samples are weighed out and analyzed twice.

##### Double determinations over 2 runs:

Finished product samples are weighed out and analyzed twice over two separate runs.

##### Maximum concentration of samples in powder form: 5%

Test samples are diluted with demineralized water with 1% stabiliser to approx. 0.037 KNU(S)/mL on the basis of their expected activity. The final dilution is made direct into the sample cup.

### 8. Procedure

#### 8.1 Cobas Menu Program

The Cobas Menu Program is used to suggest the weight/dilutions of samples and level control to be used.

The samples are entered into the program with a unique identification code and a worklist is printed out

The samples and control are weighed out and diluted as stated on the worklist with hand-written weight data is inserted into the BS-amylase analysis logbook



The results are computered automatically by the Cobas Fara as described in item 9 and printed out along with the standard curve.

Worklists and results printouts are inserted into the BS-amylase analysis logbook.

### 8.2 Cobas Fara Set-up

The samples are placed in the sample rack

The five standards are placed in the calibration rack at position 1 to 5 (strongest standard at position 5), and control placed in the same rack at position 10.

The substrate is transferred to a 30 mL reagent container and placed in that reagent rack at position 2 (holder 1).

The  $\alpha$ -glucosidase help reagent is transferred to a 50 mL reagent container and placed in the reagent rack at position 2 (holder C)

### 8.3 Cobas Fare Analysis

The main principles of the analysis are as follows: 20  $\mu$ L sample and 10  $\mu$ L rinse-water are pipetted into the cuvette along with 250  $\mu$ L  $\alpha$ -glucosidase help reagent. The cuvette rotates for 10 seconds and the reagents are thrown out into the horizontal cuvettes. 25  $\mu$ L substrate and 20  $\mu$ L rinse-water are pipetted off. After a 1 second wait to ensure that the temperature is 37° C., the cuvette rotates again and the substrate is mixed into the horizontal cuvettes. Absorbance is measured for the first time after 120 seconds and then every 5 seconds. Absorbance is measured a total of 37 times for each sample.

## 9. Calculations

The activity of the samples is calculated relative to Novo Nordisk A/S standard.

The standard curve is plotted by the analyzer. The curve is to be gently curved, rising steadily to an absorbance of around 0.25 for standard no. 5.

The activity of the samples in KNU(S)/mL is read off the standard curve by the analyzer.

The final calculations to allow for the weights/dilutions used employ the following formula:

$$\text{Activity in KNU(S)/g} = S \times V \times F / W$$

S=analysis result read off (KNU(S)/mL

V=volume of volumetric flask used in mL

F=dilution factor for second dilution

W=weight of enzyme sample in g

### 9.2 Calculation of Mean Values

Results are stated with 3 significant digits. However, for sample activity < 10 KNU(S)/g, only 2 significant digits are given.

The following rules apply on calculation of mean values:

1. Data which deviates more than 2 standard deviations from the mean value is not included in the calculation.

2. Single and double determination over one run:

The mean value is calculated on basis of results lying within the standard curve's activity area.

3. Double determinations over two runs: All values are included in the mean value. Outliers are omitted.

## 10. Accuracy and Precision

The coefficient of variation is 2.9% based on retrospective validation of analysis results for a number of finished products and the level control.

### Assay for $\alpha$ -Amylase Activity

$\alpha$ -Amylase activity is determined by a method employing Phadebas® tablets as substrate. Phadebas tablets (Phadebas® Amylase Test, supplied by Pharmacia Diagnostic) contain a cross-linked insoluble blue-coloured starch polymer which has been mixed with bovine serum albumin and a buffer substance and tabletted.

For every single measurement one tablet is suspended in a tube containing 5 ml 50 mM Britton-Robinson buffer (50

mM acetic acid, 50 mM phosphoric acid, 50 mM boric acid, 0.1 mM CaCl<sub>2</sub>, pH adjusted to the value of interest with NaOH). The test is performed in a water bath at the temperature of interest. The  $\alpha$ -amylase to be tested is diluted in x ml of 50 mM Britton-Robinson buffer. 1 ml of this  $\alpha$ -amylase solution is added to the 5 ml 50 mM Britton-Robinson buffer. The starch is hydrolysed by the  $\alpha$ -amylase giving soluble blue fragments. The absorbance of the resulting blue solution, measured spectrophotometrically at 620 nm, is a function of the  $\alpha$ -amylase activity.

It is important that the measured 620 nm absorbance after 10 or 15 minutes of incubation (testing time) is in the range of 0.2 to 2.0 absorbance units at 620 nm. In this absorbance range there is linearity between activity and absorbance (Lambert-Beer law). The dilution of the enzyme must therefore be adjusted to fit this criterion. Under a specified set of conditions (temp., pH, reaction time, buffer conditions) 1 mg of a given  $\alpha$ -amylase will hydrolyse a certain amount of substrate and a blue colour will be produced. The colour intensity is measured at 620 nm. The measured absorbance is directly proportional to the specific activity (activity/mg of pure  $\alpha$ -amylase protein) of the  $\alpha$ -amylase in question under the given set of conditions.

## EXAMPLES

### Example 1

#### Construction of Variants of BSG $\alpha$ -amylase (SEQ ID NO: 3)

The gene encoding BSG, amyS, is located in plasmid pPL1117. This plasmid contains also the gene conferring resistance towards kanamycin and an origin of replication, both obtained from plasmid pUB110 (Gryczan, T. J. et al (1978) J.Bact 134:318-329).

The DNA sequence of the mature part of amyS is shown as SEQ ID NO: 11 and the amino acid sequence of the mature protein is shown as SEQ ID NO: 3

BSG variant TVB145, which contains a deletion of 6 nucleotides corresponding to amino acids I181-G182 in the mature protein, is constructed as follows:

Polymerase Chain Reaction (PCR) is utilized to amplify the part of the amyS gene (from plasmid pPL1117), located between DNA primers BSG1 (SEQ ID NO: 15) and BSGM2 (SEQ ID NO: 18). BSG1 is identical to a part of the amyS gene whereas BSGM2 contains the 6 bp nucleotide deletion. A standard PCR reaction is carried out: 94° C. for 5 minutes, 25 cycles of (94° C. for 45 seconds, 50° C. for 45 seconds, 72° C. for 90 seconds), 72° C. for 7 minutes using the Pwo polymerase under conditions as recommended by the manufacturer, Boehringer Mannheim GmbH.

The resulting approximately 550 bp amplified band was used as a megaprimer (Barik, S and Galinski, MS (1991): Biotechniques 10: 489-490) together with primer BSG3 in a second PCR with pPL1117 as template resulting in a DNA fragment of approximately 1080 bp.

This DNA fragment is digested with restriction endonucleases Acc65I and Sall and the resulting approximately 550 bp fragment is ligated into plasmid pPL1117 digested with the same enzymes and transformed into the protease- and amylase-deleted *Bacillus subtilis* strain SHA273 (described in WO92/11357 and WO95/10603). Kanamycin resistant and starch degrading transformants were analysed for the presence of the desired mutations (restriction digest to verify the introduction of a HindIII site in the gene). The DNA sequence between restriction sites Acc65I and Sall was verified by DNA sequencing to ensure the presence of only the desired mutations.

BSG variant TVB146 which contains the same 6 nucleotide deletion as TVB145 and an additional substitution of



asparagine 193 for a phenylalanine, N193F, was constructed in a similar way as TVB145 utilizing primer BSGM3 (SEQ ID NO: 19) in the first PCR.

BSG variant TVB161, containing the deletion of I181-G182, N193F, and L204F, is constructed in a similar way as the two previous variants except that the template for the PCR reactions is plasmid pTVB146 (pPL1117 containing the TVB146-mutations within amyS and the mutagenic oligonucleotide for the first PCR is BSGM3.

BSG variant TVB162, containing the deletion of I181-G182, N193F, and E210H, is constructed in a similar way as TVB161 except that the mutagenic oligonucleotide is BSGM4 (SEQ ID NO: 20).

BSG variant TVB163, containing the deletion of I181-G182, N193F, and E214Q, is constructed in a similar way as TVB161 except that the mutagenic oligonucleotide is BSGM5 (SEQ ID NO: 21).

The above constructed BSG variants were then fermented and purified as described above in the "Material and Methods" section.

### Example 2

#### Measurement of the Calcium- and pH-dependent Stability

Normally, the industrial liquefaction process runs using pH 6.0–6.2 as liquefaction pH and an addition of 40 ppm free calcium in order to improve the stability at 95° C.–105° C. Some of the herein proposed substitutions have been made in order to improve the stability at

1. lower pH than pH 6.2 and/or
2. at free calcium levels lower than 40 ppm free calcium.

Two different methods have been used to measure the improvements in stability obtained by the different substitutions in the  $\alpha$ -amylase from *B.stearothermophilus*:

Method 1. One assay which measures the stability at reduced pH, pH 5.0, in the presence of 5 ppm free calcium. 10  $\mu$ g of the variant were incubated under the following conditions: A 0.1 M acetate solution, pH adjusted to pH 5.0, containing 5 ppm calcium and 5% w/w common corn starch (free of calcium). Incubation was made in a water bath at 95° C. for 30 minutes.

Method 2. One assay which measure the stability in the absence of free calcium and where the pH is maintained at pH 6.0. This assay measures the decrease in calcium sensitivity: 10  $\mu$ g of the variant were incubated under the following conditions: A 0.1 M acetate solution, pH adjusted to pH 6.0, containing 5% w/w common corn starch (free of calcium). Incubation was made in a water bath at 95° C. for 30 minutes.

#### Stability Determination

All the stability trials 1, 2 have been made using the same set up. The method was:

The enzyme was incubated under the relevant conditions (1–4). Samples were taken at 0, 5, 10, 15 and 30 minutes and diluted 25 times (same dilution for all taken samples) in assay buffer (0.1M 50 mM Britton buffer pH 7.3) and the activity was measured using the Phadebas assay (Pharmacia) under standard conditions pH 7.3, 37° C.

The activity measured before incubation (0 minutes) was used as reference (100%). The decline in percent was calculated as a function of the incubation time. The table shows the residual activity after 30 minutes of incubation.

#### Stability method 1. / Low pH stability improvement

MINUTES OF INCUBATION	WT. SEQ. ID. NO: 3 AMYLASE (BSG)	SEQ. ID NO: 3 VARIANT WITH DELETION IN POS. I181-G182 (TVB145)	SEQ. ID NO: 3 VARIANT WITH DELETION IN POS. I181-G182 + N193F (TVB146)	SEQ. ID NO: 3 VARIANT WITH DELETION IN POS. I181-G182 + N193F + E214Q (TVB163)
0	100	100	100	100
5	29	71	83	77
10	9	62	77	70
15	3	50	72	67
30	1	33	62	60

#### Stability Method 1./Low pH Stability Improvement

The temperature described in method 1 has been reduced from 95° C. to 70° C. since the amylases mentioned for SEQ ID NO: 1 and 2 have a lower thermostability than the one for SEQ ID NO: 3.

MINUTES OF INCUBATION	WT. SEQ. ID. NO: 2 AMYLASE	SEQ. ID NO: 2 VARIANT WITH DELETION IN POS. D183-G184	SEQ. ID NO: 1 AMYLASE	SEQ. ID NO: 1 VARIANT WITH DELETION IN POS. T183-G184
0	100	100	100	100
5	73	92	41	76
10	59	88	19	69
15	48	91	11	62
30	28	92	3	59

#### Stability method 2. / Low calcium sensitivity

MINUTES OF INCUBATION	WT. SEQ ID NO: 3 AMYLASE (BSG)	SEQ ID NO: 3 VARIANT WITH DELETION IN POS. I181-G182 (TVB145)	SEQ ID NO: 3 VARIANT WITH DELETION IN POS. I181-G182 + N193F (TVB146)	SEQ ID NO: 3 VARIANT WITH DELETION IN POS. I181-G182 + N193F + E214Q (TVB163)
0	100	100	100	100
5	60	82	81	82
10	42	76	80	83
15	31	77	81	79
30	15	67	78	79

#### Specific Activity Determination.

The specific activity was determined using the Phadebas assay (Pharmacia) as activity/mg enzyme. The activity was determined using the  $\alpha$ -amylase assay described in the Materials and Methods section herein.

The specific activity of the parent enzyme and a single and a double mutation was determined to:

BSG: SEQ ID NO:3 (Parent enzyme) 20000 NU/mg  
 TVB145: SEQ ID NO:3 with the deletion in positions I181-G182: (Single mutation) 34600 NU/mg



TVB146: SEQ ID NO:3 with the deletion in positions I181-G182+N193F: (Double mutation) 36600 NU/mg  
 TVB163: SEQ ID NO:3 with the deletion in positions I181-G182+N193F+E214Q: (Triple mutation) 36300 NU/mg

### Example 3

Pilot Plant Jet Cook and Liquefaction with Alpha-amylase Variant TVB146

Pilot plant liquefaction experiments were run in the mini-jet system using a dosage of 50 NU (S)/g DS at pH 5.5 with 5 ppm added Ca<sup>++</sup>, to compare the performance of formulated BSG alpha-amylase variant TVB146 (SEQ ID NO: 3 with deletion in positions I181-G182+N193F) with that of parent BSG alpha-amylase (SEQ ID NO: 3). The reaction was monitored by measuring the DE increase (Neocuproine method) as a function of time.

Corn starch slurries were prepared by suspending 11.8 kg Cerestar C\*Pharm GL 03406 (89 % starch) in deionized water and making up to 30 kg. The pH was adjusted to 5.5 at ambient temperature, after the addition of 0.55 g CaCl<sub>2</sub>·2H<sub>2</sub>O.

The following enzymes were used:

TVB146	108 KNU(S)/g, 146 KNU(SM9)/g
BSG amylase	101 KNU(S)/g, 98 KNU(SM9)/g

An amount of enzyme corresponding to 50 NU (SM9)/g DS was added, and the conductivity adjusted to 300 mS using NaCl. The standard conditions were as follows:

Substrate concentration	35% w/w (initial) 31.6–31.9% w/w (final)
Temperature	105° C., 5 min (Primary liquefaction) 95° C., 90 min (Secondary liquefaction)
pH (initial)	5.5

After jetting, the liquefied starch was collected and transported in sealed thermos-flasks from the pilot plant to the laboratory, where secondary liquefaction was continued at 95° C.

10 ml samples were taken at 15 minute intervals from 15–90 minutes. 2 drops of 1 N HCl were added to inactivate the enzyme. From these samples, 0.3–0.1 g (according to the expected DE) were weighed out and diluted to 100 ml. Reducing sugars were then determined according to the Neocuproine method (Determination of reducing sugar with improved precision. Dygert, Li, Florida and Thomas (1965). Anal. Biochem 13, 368) and DE values determined. The development of DE as a function of time is given in the following table:

Time (min.)	TVB146 DE (neocuproine)	BSG
15	2.80	2.32
30	4.88	3.56
45	6.58	4.98
60	8.17	6.00
75	9.91	7.40
90	11.23	8.03

As can be seen the alpha-amylase variant TVB146 performed significantly better under industrially relevant appli-

cation conditions at low levels of calcium than the parent BSG alpha-amylase.

### Example 4

5 Jet Cook and Liquefaction with a Combination of Alpha-amylase Variants (TVB146 and LE174)

Jet cook and liquefaction using a combination of the alpha-amylase variants, TVB146 and LE174 (ratio 1:1) were carried out at the following conditions:

10 Substrate A.E. Staley food grade powdered corn starch (100 lbs)

D.S. 35% using DI water

Free Ca<sup>2+</sup> 2.7 ppm at pH 5.3 (none added, from the starch only)

15 Initial pH 5.3

Dose AF9 units (AF9 is available on request) for each enzyme variant was 28 NU/g starch db for a total dose of 56 NU/g

Temperature in primary liquefaction 105° C.

20 Hold time in primary liquefaction 5 minutes

Temperature in secondary liquefaction 95° C.

At 15 minutes into secondary liquefaction 1.5 gms of hydrolyzate was added to a tared one liter volumetric containing 500 cc of DI water and 1 ml of one normal HCl and the 30 exact wt. added was recorded. This was repeated at 15 minute intervals out to 90 minutes with an additional point at 127 minutes. These were diluted to one liter and determined for dextrose equivalence via Neocuproine method as described by Dygert, Li, Florida and Thomas. 30 Determination of reducing sugar with improved precision (1965). Anal. Biochem 13, 368.

The results were as follows:

Time	DE
15	3.2
30	4.8
45	6.3
60	7.8
75	9.4
90	10.4
127	13.1

### REFERENCES CITED

- Klein, C., et al., *Biochemistry* 1992, 31, 8740–8746.  
 Mizuno, H., et al., *J. Mol. Biol.* (1993) 234, 1282–1283.  
 Chang, C., et al., *J. Mol. Biol.* (1993) 229, 235–238.  
 50 Larson, S. B., *J. Mol. Biol.* (1994) 235, 1560–1584.  
 Lawson, C. L., *J. Mol. Biol.* (1994) 236, 590–600.  
 Qian, M., et al., *J. Mol. Biol.* (1993) 231, 785–799.  
 Brady, R. L., et al., *Acta Crystallogr. sect. B*, 47, 527–535.  
 Swift, H. J., et al., *Acta Crystallogr. sect. B*, 47, 535–544.  
 55 A. Kadziola, Ph.D. Thesis: “An alpha-amylase from Barley and its Complex with a Substrate Analogue Inhibitor Studied by X-ray Crystallography”, Department of Chemistry University of Copenhagen 1993.  
 MacGregor, E. A., *Food Hydrocolloids*, 1987, Vol.1, No. 5–6.  
 60 B. Diderichsen and L. Christiansen, Cloning of a maltogenic α-amylase from *Bacillus stearothermophilus*, FEMS Microbiol. letters: 56: pp. 53–60 (1988).  
 Hudson et al., *Practical Immunology*, Third edition (1989), Blackwell Scientific Publications.  
 65 Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd Ed., Cold Spring Harbor, 1989.



- S. L. Beaucage and M. H. Caruthers, *Tetrahedron Letters* 22, 1981, pp. 1859–1869  
 Matthes et al., *The EMBO J.* 3, 1984, pp. 801–805.  
 R. K. Saiki et al., *Science* 239, 1988, pp. 487–491.  
 Morinaga et al., (1984, *Biotechnology* 2:646–639)  
 Nelson and Long, *Analytical Biochemistry* 180, 1989, pp. 147–151  
 Hunkapiller et al., 1984, *Nature* 310:105–111

- R. Higuchi, B. Krummel, and R. K. Saiki (1988). A general method of in vitro preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions. *Nucl. Acids Res.* 16:7351–7367.  
 5 Dubnau et al., 1971, *J. Mol. Biol.* 56, pp. 209–221.  
 Gryczan et al., 1978, *J. Bacteriol.* 134, pp. 318–329.  
 S. D. Erlich, 1977, *Proc. Natl. Acad. Sci.* 74, pp. 1680–1682.  
 Boel et al., 1990, *Biochemistry* 29, pp. 6244–6249.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 22

<210> SEQ ID NO 1

<211> LENGTH: 485

<212> TYPE: PRT

<213> ORGANISM: *Bacillus* sp.

<400> SEQUENCE: 1

```

His His Asn Gly Thr Asn Gly Thr Met Met Gln Tyr Phe Glu Trp Tyr
 1          5          10          15
Leu Pro Asn Asp Gly Asn His Trp Asn Arg Leu Arg Asp Asp Ala Ala
 20          25          30
Asn Leu Lys Ser Lys Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Trp
 35          40          45
Lys Gly Thr Ser Gln Asn Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr
 50          55          60
Asp Leu Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly
 65          70          75          80
Thr Arg Asn Gln Leu Gln Ala Ala Val Thr Ser Leu Lys Asn Asn Gly
 85          90          95
Ile Gln Val Tyr Gly Asp Val Val Met Asn His Lys Gly Gly Ala Asp
 100         105         110
Gly Thr Glu Ile Val Asn Ala Val Glu Val Asn Arg Ser Asn Arg Asn
 115         120         125
Gln Glu Thr Ser Gly Glu Tyr Ala Ile Glu Ala Trp Thr Lys Phe Asp
 130         135         140
Phe Pro Gly Arg Gly Asn Asn His Ser Ser Phe Lys Trp Arg Trp Tyr
 145         150         155         160
His Phe Asp Gly Thr Asp Trp Asp Gln Ser Arg Gln Leu Gln Asn Lys
 165         170         175
Ile Tyr Lys Phe Arg Gly Thr Gly Lys Ala Trp Asp Trp Glu Val Asp
 180         185         190
Thr Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Val Asp Met
 195         200         205
Asp His Pro Glu Val Ile His Glu Leu Arg Asn Trp Gly Val Trp Tyr
 210         215         220
Thr Asn Thr Leu Asn Leu Asp Gly Phe Arg Ile Asp Ala Val Lys His
 225         230         235         240
Ile Lys Tyr Ser Phe Thr Arg Asp Trp Leu Thr His Val Arg Asn Thr
 245         250         255
Thr Gly Lys Pro Met Phe Ala Val Ala Glu Phe Trp Lys Asn Asp Leu
 260         265         270
Gly Ala Ile Glu Asn Tyr Leu Asn Lys Thr Ser Trp Asn His Ser Val
 275         280         285
Phe Asp Val Pro Leu His Tyr Asn Leu Tyr Asn Ala Ser Asn Ser Gly
 290         295         300
Gly Tyr Tyr Asp Met Arg Asn Ile Leu Asn Gly Ser Val Val Gln Lys
 305         310         315         320
His Pro Thr His Ala Val Thr Phe Val Asp Asn His Asp Ser Gln Pro
 325         330         335
Gly Glu Ala Leu Glu Ser Phe Val Gln Gln Trp Phe Lys Pro Leu Ala
 340         345         350
Tyr Ala Leu Val Leu Thr Arg Glu Gln Gly Tyr Pro Ser Val Phe Tyr
 355         360         365
Gly Asp Tyr Tyr Gly Ile Pro Thr His Gly Val Pro Ala Met Lys Ser
 370         375         380
Lys Ile Asp Pro Leu Leu Gln Ala Arg Gln Thr Phe Ala Tyr Gly Thr
 385         390         395         400
Gln His Asp Tyr Phe Asp His His Asp Ile Ile Gly Trp Thr Arg Glu
 405         410         415
Gly Asn Ser Ser His Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asp
 420         425         430
Gly Pro Gly Gly Asn Lys Trp Met Tyr Val Gly Lys Asn Lys Ala Gly
 435         440         445

```



-continued

---

```

Gln Val Trp Arg Asp Ile Thr Gly Asn Arg Thr Gly Thr Val Thr Ile
  450                               455           460
Asn Ala Asp Gly Trp Gly Asn Phe Ser Val Asn Gly Gly Ser Val Ser
465                               470           475           480
Val Trp Val Lys Gln
                               485

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 485

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus sp.

&lt;400&gt; SEQUENCE: 2

```

His His Asn Gly Thr Asn Gly Thr Met Met Gln Tyr Phe Glu Trp His
  1                               5           10           15
Leu Pro Asn Asp Gly Asn His Trp Asn Arg Leu Arg Asp Asp Ala Ser
                               20           25           30
Asn Leu Arg Asn Arg Gly Ile Thr Ala Ile Trp Ile Pro Pro Ala Trp
  35                               40           45
Lys Gly Thr Ser Gln Asn Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr
  50                               55           60
Asp Leu Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly
  65                               70           75           80
Thr Arg Ser Gln Leu Glu Ser Ala Ile His Ala Leu Lys Asn Asn Gly
                               85           90           95
Val Gln Val Tyr Gly Asp Val Val Met Asn His Lys Gly Gly Ala Asp
                               100          105          110
Ala Thr Glu Asn Val Leu Ala Val Glu Val Asn Pro Asn Asn Arg Asn
                               115          120          125
Gln Glu Ile Ser Gly Asp Tyr Thr Ile Glu Ala Trp Thr Lys Phe Asp
  130                               135          140
Phe Pro Gly Arg Gly Asn Thr Tyr Ser Asp Phe Lys Trp Arg Trp Tyr
  145                               150          155          160
His Phe Asp Gly Val Asp Trp Asp Gln Ser Arg Gln Phe Gln Asn Arg
                               165          170          175
Ile Tyr Lys Phe Arg Gly Asp Gly Lys Ala Trp Asp Trp Glu Val Asp
                               180          185          190
Ser Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Val Asp Met
                               195          200          205
Asp His Pro Glu Val Val Asn Glu Leu Arg Arg Trp Gly Glu Trp Tyr
  210                               215          220
Thr Asn Thr Leu Asn Leu Asp Gly Phe Arg Ile Asp Ala Val Lys His
  225                               230          235          240
Ile Lys Tyr Ser Phe Thr Arg Asp Trp Leu Thr His Val Arg Asn Ala
                               245          250          255
Thr Gly Lys Glu Met Phe Ala Val Ala Glu Phe Trp Lys Asn Asp Leu
                               260          265          270
Gly Ala Leu Glu Asn Tyr Leu Asn Lys Thr Asn Trp Asn His Ser Val
  275                               280          285
Phe Asp Val Pro Leu His Tyr Asn Leu Tyr Asn Ala Ser Asn Ser Gly
  290                               295          300
Gly Asn Tyr Asp Met Ala Lys Leu Leu Asn Gly Thr Val Val Gln Lys
  305                               310          315          320
His Pro Met His Ala Val Thr Phe Val Asp Asn His Asp Ser Gln Pro
                               325          330          335
Gly Glu Ser Leu Glu Ser Phe Val Gln Glu Trp Phe Lys Pro Leu Ala
  340                               345          350
Tyr Ala Leu Ile Leu Thr Arg Glu Gln Gly Tyr Pro Ser Val Phe Tyr
  355                               360          365
Gly Asp Tyr Tyr Gly Ile Pro Thr His Ser Val Pro Ala Met Lys Ala
  370                               375          380
Lys Ile Asp Pro Ile Leu Glu Ala Arg Gln Asn Phe Ala Tyr Gly Thr
  385                               390          395          400
Gln His Asp Tyr Phe Asp His His Asn Ile Ile Gly Trp Thr Arg Glu
                               405          410          415
Gly Asn Thr Thr His Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asp
  420                               425          430
Gly Pro Gly Gly Glu Lys Trp Met Tyr Val Gly Gln Asn Lys Ala Gly
  435                               440          445
Gln Val Trp His Asp Ile Thr Gly Asn Lys Pro Gly Thr Val Thr Ile
  450                               455           460
Asn Ala Asp Gly Trp Ala Asn Phe Ser Val Asn Gly Gly Ser Val Ser
  465                               470           475           480
Ile Trp Val Lys Arg
                               485

```



-continued

<210> SEQ ID NO 3  
 <211> LENGTH: 514  
 <212> TYPE: PRT  
 <213> ORGANISM: *B. stearothermophilus*

<400> SEQUENCE: 3

```

Ala Ala Pro Phe Asn Gly Thr Met Met Gln Tyr Phe Glu Trp Tyr Leu
 1          5          10          15
Pro Asp Asp Gly Thr Leu Trp Thr Lys Val Ala Asn Glu Ala Asn Asn
 20          25          30
Leu Ser Ser Leu Gly Ile Thr Ala Leu Trp Leu Pro Pro Ala Tyr Lys
 35          40          45
Gly Thr Ser Arg Ser Asp Val Gly Tyr Gly Val Tyr Asp Leu Tyr Asp
 50          55          60
Leu Gly Glu Phe Asn Gln Lys Gly Ala Val Arg Thr Lys Tyr Gly Thr
 65          70          75          80
Lys Ala Gln Tyr Leu Gln Ala Ile Gln Ala Ala His Ala Ala Gly Met
 85          90          95
Gln Val Tyr Ala Asp Val Val Phe Asp His Lys Gly Gly Ala Asp Gly
 100         105         110
Thr Glu Trp Val Asp Ala Val Glu Val Asn Pro Ser Asp Arg Asn Gln
 115         120         125
Glu Ile Ser Gly Thr Tyr Gln Ile Gln Ala Trp Thr Lys Phe Asp Phe
 130         135         140
Pro Gly Arg Gly Asn Thr Tyr Ser Ser Phe Lys Trp Arg Trp Tyr His
 145         150         155         160
Phe Asp Gly Val Asp Trp Asp Glu Ser Arg Lys Leu Ser Arg Ile Tyr
 165         170         175
Lys Phe Arg Gly Ile Gly Lys Ala Trp Asp Trp Glu Val Asp Thr Glu
 180         185         190
Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Leu Asp Met Asp His
 195         200         205
Pro Glu Val Val Thr Glu Leu Lys Ser Trp Gly Lys Trp Tyr Val Asn
 210         215         220
Thr Thr Asn Ile Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys
 225         230         235         240
Phe Ser Phe Phe Pro Asp Trp Leu Ser Asp Val Arg Ser Gln Thr Gly
 245         250         255
Lys Pro Leu Phe Thr Val Gly Glu Tyr Trp Ser Tyr Asp Ile Asn Lys
 260         265         270
Leu His Asn Tyr Ile Met Lys Thr Asn Gly Thr Met Ser Leu Phe Asp
 275         280         285
Ala Pro Leu His Asn Lys Phe Tyr Thr Ala Ser Lys Ser Gly Gly Thr
 290         295         300
Phe Asp Met Arg Thr Leu Met Thr Asn Thr Leu Met Lys Asp Gln Pro
 305         310         315         320
Thr Leu Ala Val Thr Phe Val Asp Asn His Asp Thr Glu Pro Gly Gln
 325         330         335
Ala Leu Gln Ser Trp Val Asp Pro Trp Phe Lys Pro Leu Ala Tyr Ala
 340         345         350
Phe Ile Leu Thr Arg Gln Glu Gly Tyr Pro Cys Val Phe Tyr Gly Asp
 355         360         365
Tyr Tyr Gly Ile Pro Gln Tyr Asn Ile Pro Ser Leu Lys Ser Lys Ile
 370         375         380
Asp Pro Leu Leu Ile Ala Arg Arg Asp Tyr Ala Tyr Gly Thr Gln His
 385         390         395         400
Asp Tyr Leu Asp His Ser Asp Ile Ile Gly Trp Thr Arg Glu Gly Val
 405         410         415
Thr Glu Lys Pro Gly Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro
 420         425         430
Gly Gly Ser Lys Trp Met Tyr Val Gly Lys Gln His Ala Gly Lys Val
 435         440         445
Phe Tyr Asp Leu Thr Gly Asn Arg Ser Asp Thr Val Thr Ile Asn Ser
 450         455         460
Asp Gly Trp Gly Glu Phe Lys Val Asn Gly Gly Ser Val Ser Val Trp
 465         470         475         480
Val Pro Arg Lys Thr Thr Val Ser Thr Ile Ala Trp Ser Ile Thr Thr
 485         490         495
Arg Pro Trp Thr Asp Glu Phe Val Arg Trp Thr Glu Pro Arg Leu Val
 500         505         510
Ala Trp

```

<210> SEQ ID NO 4  
 <211> LENGTH: 483  
 <212> TYPE: PRT  
 <213> ORGANISM: *B. licheniformis*



-continued

&lt;400&gt; SEQUENCE: 4

```

Ala Asn Leu Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Met Pro
 1          5          10          15
Asn Asp Gly Gln His Trp Arg Arg Leu Gln Asn Asp Ser Ala Tyr Leu
 20          25          30
Ala Glu His Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly
 35          40          45
Thr Ser Gln Ala Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr Asp Leu
 50          55          60
Gly Glu Phe His Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly Thr Lys
 65          70          75          80
Gly Glu Leu Gln Ser Ala Ile Lys Ser Leu His Ser Arg Asp Ile Asn
 85          90          95
Val Tyr Gly Asp Val Val Ile Asn His Lys Gly Gly Ala Asp Ala Thr
100          105          110
Glu Asp Val Thr Ala Val Glu Val Asp Pro Ala Asp Arg Asn Arg Val
115          120          125
Ile Ser Gly Glu His Leu Ile Lys Ala Trp Thr His Phe His Phe Pro
130          135          140
Gly Arg Gly Ser Thr Tyr Ser Asp Phe Lys Trp His Trp Tyr His Phe
145          150          155          160
Asp Gly Thr Asp Trp Asp Glu Ser Arg Lys Leu Asn Arg Ile Tyr Lys
165          170          175
Phe Gln Gly Lys Ala Trp Asp Trp Glu Val Ser Asn Glu Asn Gly Asn
180          185          190
Tyr Asp Tyr Leu Met Tyr Ala Asp Ile Asp Tyr Asp His Pro Asp Val
195          200          205
Ala Ala Glu Ile Lys Arg Trp Gly Thr Trp Tyr Ala Asn Glu Leu Gln
210          215          220
Leu Asp Gly Phe Arg Leu Asp Ala Val Lys His Ile Lys Phe Ser Phe
225          230          235          240
Leu Arg Asp Trp Val Asn His Val Arg Glu Lys Thr Gly Lys Glu Met
245          250          255
Phe Thr Val Ala Glu Tyr Trp Gln Asn Asp Leu Gly Ala Leu Glu Asn
260          265          270
Tyr Leu Asn Lys Thr Asn Phe Asn His Ser Val Phe Asp Val Pro Leu
275          280          285
His Tyr Gln Phe His Ala Ala Ser Thr Gln Gly Gly Tyr Asp Met
290          295          300
Arg Lys Leu Leu Asn Gly Thr Val Val Ser Lys His Pro Leu Lys Ser
305          310          315          320
Val Thr Phe Val Asp Asn His Asp Thr Gln Pro Gly Gln Ser Leu Glu
325          330          335
Ser Thr Val Gln Thr Trp Phe Lys Pro Leu Ala Tyr Ala Phe Ile Leu
340          345          350
Thr Arg Glu Ser Gly Tyr Pro Gln Val Phe Tyr Gly Asp Met Tyr Gly
355          360          365
Thr Lys Gly Asp Ser Gln Arg Glu Ile Pro Ala Leu Lys His Lys Ile
370          375          380
Glu Pro Ile Leu Lys Ala Arg Lys Gln Tyr Ala Tyr Gly Ala Gln His
385          390          395          400
Asp Tyr Phe Asp His His Asp Ile Val Gly Trp Thr Arg Glu Gly Asp
405          410          415
Ser Ser Val Ala Asn Ser Gly Leu Ala Ala Leu Ile Thr Asp Gly Pro
420          425          430
Gly Gly Ala Lys Arg Met Tyr Val Gly Arg Gln Asn Ala Gly Glu Thr
435          440          445
Trp His Asp Ile Thr Gly Asn Arg Ser Glu Pro Val Val Ile Asn Ser
450          455          460
Glu Gly Trp Gly Glu Phe His Val Asn Gly Gly Ser Val Ser Ile Tyr
465          470          475          480
Val Gln Arg

```

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 480

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: B. amyloliquefaciens

&lt;400&gt; SEQUENCE: 5

```

Val Asn Gly Thr Leu Met Gln Tyr Phe Glu Trp Tyr Thr Pro Asn Asp
 1          5          10          15
Gly Gln His Trp Lys Arg Leu Gln Asn Asp Ala Glu His Leu Ser Asp
 20          25          30
Ile Gly Ile Thr Ala Val Trp Ile Pro Pro Ala Tyr Lys Gly Leu Ser
 35          40          45
Gln Ser Asp Asn Gly Tyr Gly Pro Tyr Asp Leu Tyr Asp Leu Gly Glu

```



-continued

50					55					60					
Phe	Gln	Gln	Lys	Gly	Thr	Val	Arg	Thr	Lys	Tyr	Gly	Thr	Lys	Ser	Glu
65					70					75					80
Leu	Gln	Asp	Ala	Ile	Gly	Ser	Leu	His	Ser	Arg	Asn	Val	Gln	Val	Tyr
				85					90					95	
Gly	Asp	Val	Val	Leu	Asn	His	Lys	Ala	Gly	Ala	Asp	Ala	Thr	Glu	Asp
			100					105					110		
Val	Thr	Ala	Val	Glu	Val	Asn	Pro	Ala	Asn	Arg	Asn	Gln	Glu	Thr	Ser
		115					120					125			
Glu	Glu	Tyr	Gln	Ile	Lys	Ala	Trp	Thr	Asp	Phe	Arg	Phe	Pro	Gly	Arg
		130				135					140				
Gly	Asn	Thr	Tyr	Ser	Asp	Phe	Lys	Trp	His	Trp	Tyr	His	Phe	Asp	Gly
145					150					155					160
Ala	Asp	Trp	Asp	Glu	Ser	Arg	Lys	Ile	Ser	Arg	Ile	Phe	Lys	Phe	Arg
				165					170					175	
Gly	Glu	Gly	Lys	Ala	Trp	Asp	Trp	Glu	Val	Ser	Ser	Glu	Asn	Gly	Asn
			180					185					190		
Tyr	Asp	Tyr	Leu	Met	Tyr	Ala	Asp	Val	Asp	Tyr	Asp	His	Pro	Asp	Val
		195					200					205			
Val	Ala	Glu	Thr	Lys	Lys	Trp	Gly	Ile	Trp	Tyr	Ala	Asn	Glu	Leu	Ser
		210				215					220				
Leu	Asp	Gly	Phe	Arg	Ile	Asp	Ala	Ala	Lys	His	Ile	Lys	Phe	Ser	Phe
225					230					235					240
Leu	Arg	Asp	Trp	Val	Gln	Ala	Val	Arg	Gln	Ala	Thr	Gly	Lys	Glu	Met
				245					250					255	
Phe	Thr	Val	Ala	Glu	Tyr	Trp	Gln	Asn	Asn	Ala	Gly	Lys	Leu	Glu	Asn
			260					265					270		
Tyr	Leu	Asn	Lys	Thr	Ser	Phe	Asn	Gln	Ser	Val	Phe	Asp	Val	Pro	Leu
		275					280					285			
His	Phe	Asn	Leu	Gln	Ala	Ala	Ser	Ser	Gln	Gly	Gly	Gly	Tyr	Asp	Met
		290				295					300				
Arg	Arg	Leu	Leu	Asp	Gly	Thr	Val	Val	Ser	Arg	His	Pro	Glu	Lys	Ala
305					310				315						320
Val	Thr	Phe	Val	Glu	Asn	His	Asp	Thr	Gln	Pro	Gly	Gln	Ser	Leu	Glu
				325					330					335	
Ser	Thr	Val	Gln	Thr	Trp	Phe	Lys	Pro	Leu	Ala	Tyr	Ala	Phe	Ile	Leu
			340					345					350		
Thr	Arg	Glu	Ser	Gly	Tyr	Pro	Gln	Val	Phe	Tyr	Gly	Asp	Met	Tyr	Gly
		355				360						365			
Thr	Lys	Gly	Thr	Ser	Pro	Lys	Glu	Ile	Pro	Ser	Leu	Lys	Asp	Asn	Ile
		370				375					380				
Glu	Pro	Ile	Leu	Lys	Ala	Arg	Lys	Glu	Tyr	Ala	Tyr	Gly	Pro	Gln	His
385					390					395					400
Asp	Tyr	Ile	Asp	His	Pro	Asp	Val	Ile	Gly	Trp	Thr	Arg	Glu	Gly	Asp
				405					410					415	
Ser	Ser	Ala	Ala	Lys	Ser	Gly	Leu	Ala	Ala	Leu	Ile	Thr	Asp	Gly	Pro
			420					425					430		
Gly	Gly	Ser	Lys	Arg	Met	Tyr	Ala	Gly	Leu	Lys	Asn	Ala	Gly	Glu	Thr
			435				440					445			
Trp	Tyr	Asp	Ile	Thr	Gly	Asn	Arg	Ser	Asp	Thr	Val	Lys	Ile	Gly	Ser
		450				455				460					
Asp	Gly	Trp	Gly	Glu	Phe	His	Val	Asn	Asp	Gly	Ser	Val	Ser	Ile	Tyr
465					470					475					480

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 485

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus sp.

&lt;400&gt; SEQUENCE: 6

His	His	Asn	Gly	Thr	Asn	Gly	Thr	Met	Met	Gln	Tyr	Phe	Glu	Trp	Tyr
1				5					10					15	
Leu	Pro	Asn	Asp	Gly	Asn	His	Trp	Asn	Arg	Leu	Asn	Ser	Asp	Ala	Ser
			20					25					30		
Asn	Leu	Lys	Ser	Lys	Gly	Ile	Thr	Ala	Val	Trp	Ile	Pro	Pro	Ala	Trp
		35					40					45			
Lys	Gly	Ala	Ser	Gln	Asn	Asp	Val	Gly	Tyr	Gly	Ala	Tyr	Asp	Leu	Tyr
		50				55					60				
Asp	Leu	Gly	Glu	Phe	Asn	Gln	Lys	Gly	Thr	Val	Arg	Thr	Lys	Tyr	Gly
65					70					75					80
Thr	Arg	Ser	Gln	Leu	Gln	Ala	Ala	Val	Thr	Ser	Leu	Lys	Asn	Asn	Gly
				85					90					95	
Ile	Gln	Val	Tyr	Gly	Asp	Val	Val	Met	Asn	His	Lys	Gly	Gly	Ala	Asp
			100					105					110		
Ala	Thr	Glu	Met	Val	Arg	Ala	Val	Glu	Val	Asn	Pro	Asn	Asn	Arg	Asn
			115				120					125			
Gln	Glu	Val	Thr	Gly	Glu	Tyr	Thr	Ile	Glu	Ala	Trp	Thr	Arg	Phe	Asp







-continued

```

195          200          205
Asp His Pro Glu Val Ile His Glu Leu Arg Asn Trp Gly Val Trp Tyr
210          215          220
Thr Asn Thr Leu Asn Leu Asp Gly Phe Arg Ile Asp Ala Val Lys His
225          230          235          240
Ile Lys Tyr Ser Phe Thr Arg Asp Trp Leu Thr His Val Arg Asn Thr
245          250          255
Thr Gly Lys Pro Met Phe Ala Val Ala Glu Phe Trp Lys Asn Asp Leu
260          265          270
Gly Ala Ile Glu Asn Tyr Leu Asn Lys Thr Ser Trp Asn His Ser Val
275          280          285
Phe Asp Val Pro Leu His Tyr Asn Leu Tyr Asn Ala Ser Asn Ser Gly
290          295          300
Gly Tyr Tyr Asp Met Arg Asn Ile Leu Asn Gly Ser Val Val Gln Lys
305          310          315          320
His Pro Thr His Ala Val Thr Phe Val Asp Asn His Asp Ser Gln Pro
325          330          335
Gly Glu Ala Leu Glu Ser Phe Val Gln Gln Trp Phe Lys Pro Leu Ala
340          345          350
Tyr Ala Leu Val Leu Thr Arg Glu Gln Gly Tyr Pro Ser Val Phe Tyr
355          360          365
Gly Asp Tyr Tyr Gly Ile Pro Thr His Gly Val Pro Ala Met Lys Ser
370          375          380
Lys Ile Asp Pro Leu Leu Gln Ala Arg Gln Thr Phe Ala Tyr Gly Thr
385          390          395          400
Gln His Asp Tyr Phe Asp His His Asp Ile Ile Gly Trp Thr Arg Glu
405          410          415
Gly Asn Ser Ser His Pro Asn Ser Gly Leu Ala Thr Ile Met Ser Asp
420          425          430
Gly Pro Gly Gly Asn Lys Trp Met Tyr Val Gly Lys Asn Lys Ala Gly
435          440          445
Gln Val Trp Arg Asp Ile Thr Gly Asn Arg Thr Gly Thr Val Thr Ile
450          455          460
Asn Ala Asp Gly Trp Gly Asn Phe Ser Val Asn Gly Gly Ser Val Ser
465          470          475          480
Val Trp Val Lys Gln
485

```

<210> SEQ ID NO 8  
 <211> LENGTH: 485  
 <212> TYPE: PRT  
 <213> ORGANISM: Bacillus sp.

<400> SEQUENCE: 8

```

His His Asn Gly Thr Asn Gly Thr Met Met Gln Tyr Phe Glu Trp His
1          5          10          15
Leu Pro Asn Asp Gly Asn His Trp Asn Arg Leu Arg Asp Asp Ala Ser
20          25          30
Asn Leu Arg Asn Arg Gly Ile Thr Ala Ile Trp Ile Pro Ala Trp
35          40          45
Lys Gly Thr Ser Gln Asn Asp Val Gly Tyr Gly Ala Tyr Asp Leu Tyr
50          55          60
Asp Leu Gly Glu Phe Asn Gln Lys Gly Thr Val Arg Thr Lys Tyr Gly
65          70          75          80
Thr Arg Ser Gln Leu Glu Ser Ala Ile His Ala Leu Lys Asn Asn Gly
85          90          95
Val Gln Val Tyr Gly Asp Val Val Met Asn His Lys Gly Gly Ala Asp
100          105          110
Ala Thr Glu Asn Val Leu Ala Val Glu Val Asn Pro Asn Asn Arg Asn
115          120          125
Gln Glu Ile Ser Gly Asp Tyr Thr Ile Glu Ala Trp Thr Lys Phe Asp
130          135          140
Phe Pro Gly Arg Gly Asn Thr Tyr Ser Asp Phe Lys Trp Arg Trp Tyr
145          150          155          160
His Phe Asp Gly Val Asp Trp Asp Gln Ser Arg Gln Phe Gln Asn Arg
165          170          175
Ile Tyr Lys Phe Arg Gly Asp Gly Lys Ala Trp Asp Trp Glu Val Asp
180          185          190
Ser Glu Asn Gly Asn Tyr Asp Tyr Leu Met Tyr Ala Asp Val Asp Met
195          200          205
Asp His Pro Glu Val Val Asn Glu Leu Arg Arg Trp Gly Glu Trp Tyr
210          215          220
Thr Asn Thr Leu Asn Leu Asp Gly Phe Arg Ile Asp Ala Val Lys His
225          230          235          240
Ile Lys Tyr Ser Phe Thr Arg Asp Trp Leu Thr His Val Arg Asn Ala
245          250          255
Thr Gly Lys Glu Met Phe Ala Val Ala Glu Phe Trp Lys Asn Asp Leu

```







-continued

actaagtttg	atthttccagg	gaggggtaat	acatactcag	actthtaaatg	gcgttggtat	480
catttcgatg	gtgtagattg	ggatcaatca	cgacaattcc	aaaatcgtat	ctacaaattc	540
cgaggtgatg	gtaaggcatg	ggattgggaa	gtagattcgg	aaaatggaaa	ttatgattat	600
ttaatgtatg	cagatgtaga	tatggatcat	ccggaggtag	taaagtagct	tagaagatgg	660
ggagaatggt	atacaaatc	attaaatctt	gatggattta	ggatcgtatc	ggtgaagcat	720
atthaaatata	gctttacacg	tgattgggtt	acccatgtaa	gaaacgcaac	gggaaaagaa	780
atgthttgctg	ttgctgaatt	ttggaaaaat	gatttaggtg	ccttgagaa	ctattthaaat	840
aaaacaaact	ggaatcattc	tgtctttgat	gtcccccttc	attataatct	ttataacgcg	900
tcaaatagtg	gaggcaacta	tgacatggca	aaacttctta	atggaacggt	tgttcaaaaag	960
catccaatgc	atgccgtaac	ttttgtggat	aatcacgatt	ctcaacctgg	ggaatcatta	1020
gaatcatttg	tacaagaatg	gtttaagcca	cttgcttatg	cgcttatttt	aacaagagaa	1080
caaggctatc	cctctgtctt	ctatgggtgac	tactatggaa	ttccaacaca	tagtgtccca	1140
gcaatgaaag	ccaagattga	tccaatctta	gaggcgcgtc	aaaattttgc	atatggaaca	1200
caacatgatt	atthttgacca	tcataatata	atcggatgga	cacgtgaagg	aaataaccacg	1260
catcccaatt	caggacttgc	gactatcatg	tcggatgggc	cagggggaga	gaaatggatg	1320
tacgtagggc	aaaataaagc	aggtcaagtt	tgcatgaca	taactggaaa	taaaccagga	1380
acagttacga	tcaatgcaga	tgatgggct	aatthtttcag	taaagtagg	atctgtttcc	1440
atthgggtga	aacga					1455

&lt;210&gt; SEQ ID NO 11

&lt;211&gt; LENGTH: 1548

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *B. stearothermophilus*

&lt;400&gt; SEQUENCE: 11

gccgcaccgt	ttaacggcac	catgatgcag	tatthttgaat	ggtacttgcc	ggatgatggc	60
acgttatgga	ccaaagtggc	caatgaagcc	aacaacttat	ccagccttgg	catcaccgct	120
ctthtgctgc	cgcccgctta	caaaggaaca	agccgcagcg	acgtagggtta	cggagtatac	180
gacttgtagt	acctcggcga	attcaatcaa	aaagggaccg	tccgcacaaa	atcgggaaca	240
aaagctcaat	atcttcaagc	cattcaagcc	gcccacgccc	ctggaatgca	agtgtacgcc	300
gatgtcgtgt	tcgaccataa	aggcggcgct	gacggcacgg	aatgggtgga	cgccgtcgaa	360
gtcaatccgt	ccgaccgcaa	ccaagaaatc	tcgggcacct	atcaaatcca	agcatggacg	420
aaatthtgatt	ttcccgggcg	gggcaacacc	tactccagct	ttaatggcg	ctggtaccat	480
thttgacggcg	ttgattggga	cgaaagccga	aaatthgacc	gcattthaca	attccgcggc	540
atcggcaaaag	cgtgggattg	ggaaagtagac	acggaaaacg	gaaactatga	ctacttaatg	600
tatgccgacc	ttgatatgga	tcatcccga	gtcgtgaccg	agctgaaaaa	ctgggggaaa	660
tggtatgtca	acacaacgaa	cattgatggg	ttccggcttg	atgccgtcaa	gcatattaag	720
ttcagthttt	ttcctgattg	gttgtcgtat	gtgcgttctc	agactggcaa	gccgctatth	780
accgtcgggg	aatattggag	ctatgacatc	aacaagthtg	acaattacat	tacgaaaaaca	840
gacggaacga	tgtctthgtt	tgatgccccg	ttacacaaca	aatthttatac	cgcttccaaa	900
tcagggggcg	catttgatat	gcgcacgtta	atgaccaata	ctctcatgaa	agatcaaccg	960
acattggccg	tcaccttcgt	tgataatcat	gacaccgaac	ccggccaagc	gctgcagtca	1020
tggttcgacc	catggttcaa	accgttggct	tacgcctthta	ttctaactcg	gcaggaagga	1080
taccctgctg	tctthttatg	tgactattat	ggcattccac	aatataacat	tccttcgctg	1140
aaaagcaaaa	tcgatccgct	cctcatcgcg	cgcagggatt	atgcttacgg	aacgcaacat	1200
gattatcttg	atcactccga	catcatcggg	tggaacaagg	aagggggcac	tgaaaaacca	1260
ggatccggac	tgcccgcaact	gatcaccgat	gggcccggag	gaagcaaatg	gatgtacgth	1320
ggcaaaacaac	acgttgga	agtgttctat	gaccttaccg	gcaaccggag	tgacaccgth	1380
accatcaaca	gtgatggatg	gggggaattc	aaagtcaatg	gcggttcggt	ttcggthttg	1440
gthtctagaa	aaacgaccgt	thtctaccatc	gctcggccga	tcacaaccg	accgtggact	1500
ggtgaattcg	tccgttgga	cgaaccacgg	ttggtggcat	ggccttga		1548

&lt;210&gt; SEQ ID NO 12

&lt;211&gt; LENGTH: 1920

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *B. licheniformis*

&lt;400&gt; SEQUENCE: 12

cggaagattg	gaagtacaaa	aataagcaaa	agattgtcaa	tcatgtcatg	agccatgcgg	60
gagacggaaa	aatcgtctta	atgcacgata	thttatgcaac	gthtcgagat	gctgctgaag	120
agattattaa	aaagctgaaa	gcaaaaggct	atcaattggt	aactgtatct	cagcttgaag	180
aagtgaagaa	gcagagaggc	tattgaataa	atgagtagaa	gcgcatatc	ggcgctthtc	240
thtttgaaga	aaatataggg	aaaatggtac	thgtthaaaa	thcggaatat	thatacaaca	300
tcatatgtht	cacattgaaa	ggggaggaga	atcatgaaac	aacaaaaacg	gctthtacgcc	360
cgattgctga	cgctgttatt	tgcgctcatc	thcttgctgc	ctcattctgc	agcagcggcg	420
gcaaatctta	atgggacgct	gatgcagtat	thttgaatggt	acatgccc	tgacggccaa	480
cattggaggc	gthttgcaaaa	cgactcggca	tatthtgctg	aacacggtat	tactgccgth	540
tgattcccc	cggcatataa	gggaacgagc	caagcggatg	tggtctacgg	tgcttacgac	600
ctthtatgatt	taggggagth	tcatcaaaaa	gggacggttc	ggacaaagta	cggcaca	660
ggagagctgc	aatctgcgat	caaaagtctt	cattcccgcg	acattaacgt	ttacggggat	720
gtggtcatca	accacaaagg	cggcgctgat	gcgaccgaag	atgtaaccgc	ggttgaagtc	780
gatcccgcgt	accgcaaccg	cgtaatthta	ggagaacacc	taatthaaagc	ctggacacat	840
thtcatthtc	cggggcgcgg	cagcacatac	agcgaatthta	aatggcattg	gtaccattht	900
gacggaaccg	atthggacga	gtcccgaag	ctgaccgca	tctataagth	tcaaggaaag	960
gctthggatt	gggaagthtc	caatgaaaac	ggcaactatg	atthattgatt	gtatgccgac	1020



-continued

atcgattatg	accatcctga	tgctgcagca	gaaattaaga	gatggggcac	ttggtatgcc	1080
aatgaactgc	aattggacgg	tttccgtctt	gatgctgtca	aacacattaa	atcttctttt	1140
ttgcgggatt	gggttaatca	tgctcagggaa	aaaacgggga	aggaaatggt	tacggtagct	1200
gaatattggc	agaatgactt	gggcgcgctg	gaaaactatt	tgaacaaaac	aaattttaat	1260
cattcagtgt	ttgacgtgcc	gcttcattat	cagttccatg	ctgcatcgac	acagggaggc	1320
ggctatgata	tgaggaaatt	gctgaacggg	acggtcgttt	ccaagcatcc	gttgaaatcg	1380
gttacatttg	tcgataacca	tgatacacag	cgggggcaat	cgcttgagtc	gactgtccaa	1440
acatggttta	agccgcttgc	ttacgctttt	attctcacia	gggaatctgg	ataccctcag	1500
gttttctacg	gggatatgta	cgggacgaaa	ggagactccc	agcgcgaaat	tcctgccttg	1560
aaacacaaaa	ttgaaccgat	cttaaaagcg	agaaaacagt	atgcgtacgg	agcacagcat	1620
gattatttctg	accaccatga	cattgtcggc	tggacaaggg	aaggcgacag	ctcggttgca	1680
aattcagggtt	tggcggcatt	aataacagac	ggaccgggtg	gggcaaagcg	aatgtatgtc	1740
ggccggcaaa	acgcgggtga	gacatggcat	gacattaccg	gaaaccgttc	ggagccgggtt	1800
gtcatcaatt	cggaaaggctg	gggagagttt	cacgtaaacg	gcgggtcggg	ttcaatttat	1860
gttcaaagat	agaagagcag	agaggacgga	tttctgaag	gaaatccggt	tttttatttt	1920

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 2084

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *B. amyloliquefaciens*

&lt;400&gt; SEQUENCE: 13

gccccgcaca	tacgaaaaga	ctggctgaaa	acattgagcc	tttgatgact	gatgatttgg	60
ctgaagaagt	ggatcgattg	tttgagaaaa	gaagaagacc	ataaaaatac	cttgtctgtc	120
atcagacagg	gtatttttta	tgctgtccag	actgtccgct	gtgtaaaaat	aaggaataaa	180
gggggggttg	tattatttta	ctgatatgta	aaatataatt	tgtataagaa	aatgagaggg	240
agaggaaaac	tgattcaaaa	acgaaagcgg	acagtttcgt	tcagacttgt	gcttatgtgc	300
acgctgttat	ttgtcagttt	gccgattaca	aaaacatcag	ccgtaaatgg	cacgctgatg	360
cagtattttg	aatggtatac	gccgaacgac	ggccagcatt	ggaaacgatt	gcagaatgat	420
gcggaacatt	tatcggatat	cggaaatcact	gccgtctgga	ttcctcccgc	atacaaagga	480
ttgagccaat	ccgataacgg	atacggacct	tatgatttgt	atgatttagg	agaattccag	540
caaaaaggga	cggtcagaac	gaaatacggc	acaaaatcag	agcttcaaga	tgcatcggc	600
tcactgcatt	cccggaacgt	ccaagtatac	ggagatgtgg	ttttgaaatc	taaggctggt	660
gctgatgcaa	cagaagatgt	aactgccgct	gaagtcaatc	cggccaatag	aatcaggaa	720
acttcggagg	aatatcaaat	caaagcgtgg	acggattttc	gttttcgggg	ccgtggaaac	780
acgtacagtg	attttaaatg	gcattgggat	catttcgacg	gagcggactg	ggatgaatcc	840
cggaagatca	gccgcatctt	taagtttcgt	ggggaaggaa	aagcgtggga	ttgggaagta	900
tcaagtgtaa	acggcaacta	tgactattta	atgtatgctg	atggtgacta	cgaccaccct	960
gatgtcgtgg	cagagacaaa	aaaatggggg	atctgggtat	cgaatgaact	gtcattagac	1020
ggcttccgta	ttgatgccgc	caaacatatt	aaattttcat	ttctgcgtga	ttgggttcag	1080
gcggtcagac	aggcgacggg	aaaagaaatg	tttacggttg	cggagtattg	gcagaataat	1140
gccgggaaac	tcgaaaacta	cttgaataaa	acaagcttta	atcaatccgt	gtttgatggt	1200
ccgcttcatt	tcaatttaca	ggcggttcc	tcacaaggag	gcggatatga	tatgagcgt	1260
ttgctggacg	gtaccgttgt	gtccaggcat	ccggaaaagg	cggttacatt	tgttgaaaat	1320
catgacacac	agccgggaca	gtcattggaa	tcgacagtcc	aaacttggtt	taaaccgctt	1380
gcatacgcct	ttattttgac	aagagaatcc	ggttatcctc	aggtgttcta	tggggatatg	1440
tacgggacaa	aaggacatc	gccaaaaggaa	attccctcac	tgaaagataa	tatagagccg	1500
attttaaaag	cgcgtaagga	gtacgcatac	gggccccagc	acgattatat	tgaccaccgg	1560
gatgtgatcg	gatggacgag	ggaagggtgac	agctccgccc	ccaaatcagg	tttggccgct	1620
ttaatcacgg	acggaccggg	cggatcaaaag	cggatgtatg	ccggcctgaa	aaatgccggc	1680
gagacatggt	atgacataac	gggcaaccgt	tcagatactg	taaaaatcgg	atctgacggc	1740
tggggagagt	ttcatgtaaa	cgatgggtcc	gtctccattt	atggttcagaa	ataaggtaat	1800
aaaaaaacac	ctccaagctg	agtgcgggta	tcagcttggg	ggtgcgttta	ttttttcagc	1860
cgtatgacaa	ggtcggcatc	agggtgtgaca	aatacgggat	gctggctgtc	ataggtgaca	1920
aatccgggtt	ttgcgccgtt	tggttttttc	acatgtctga	tttttgata	atcaacaggc	1980
acggagccgg	aatctttcgc	cttggaaaaa	taagcggcga	tcgtagctgc	ttccaatatg	2040
gattgttcat	cgggatcgct	gcttttaatc	acaacgtggg	atcc		2084

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 1455

&lt;212&gt; TYPE: DNA

<213> ORGANISM: *Bacillus* sp.

&lt;400&gt; SEQUENCE: 14

catcataatg	gaacaaatgg	tactatgatg	caatatttctg	aatggtatctt	gccaaatgac	60
gggaatcatt	ggaacaggtt	gagggatgac	gcagctaact	taaagagtaa	agggataaca	120
gctgtatgga	tcccacctgc	atggaagggg	acttcccaga	atgatgtagg	ttatggagcc	180
tatgatttat	atgatcttgg	agagttaaac	cagaagggga	cggttcgtac	aaaatatgga	240
acacgcaacc	agctacaggc	tgccggtgacc	tctttaaaaa	ataacggcat	tcaggatat	300
ggtgatgtcg	tcatgaatca	taaagggtgga	gcagatggta	cggaaattgt	aatgcccgtg	360
gaagtgaatc	ggagcaaccg	aaaccaggaa	acctcaggag	agtatgcaat	agaagcgtgg	420
acaaagtttg	atcttctctg	aagaggaaat	aaccattcca	gctttaagtg	gcgctggtat	480
cattttgatg	ggacagattg	ggatcagtca	cgccagcttc	aaaacaaaat	atataaattc	540
aggggaacag	gcaagcctg	ggactgggaa	gtcgatacag	agaatggcaa	ctatgactat	600
cttatgtatg	cagacgtgga	tatggatcac	ccagaagtaa	tacatgaact	tagaaactgg	660



-continued

ggagtgtggt	atacgaatac	actgaacctt	gatggattta	gaatagatgc	agtgaaacat	720
ataaaatata	gctttacgag	agattggctt	acacatgtgc	gtaacaccac	aggtaaacca	780
atgtttgag	tggctgagtt	ttggaaaaat	gaccttgggtg	caattgaaaa	ctatttgaat	840
aaaacaagtt	ggaatcactc	ggtgtttgat	gttcctctcc	actataattt	gtacaatgca	900
tctaatacg	gtggttatta	tgatatgaga	aatattttta	atggttctgt	ggtgcaaaaa	960
catccaacac	atgccgttac	ttttgttgat	aacctatgatt	ctcagcccgg	ggaagcattg	1020
gaatcctttg	ttcaacaatg	gtttaaacca	cttgcatatg	cattggttct	gacaagggaa	1080
caaggttatc	cttccgtatt	ttatggggat	tactacggta	tccaaccca	tggtgttccg	1140
gctatgaaat	ctaaaataga	ccctcttctg	caggcacgtc	aaacttttgc	ctatggtacg	1200
cagcatgatt	actttgatca	tcatgatatt	atcggttggga	caagagaggg	aaatagctcc	1260
catccaaatt	caggccttgc	caccattatg	tcagatggtc	caggtggtaa	caaatggatg	1320
tatgtgggga	aaaataaagc	gggacaagtt	tggagagata	ttaccgaaa	taggacaggc	1380
accgtcacia	ttaatgcaga	cggatggggg	aatttctctg	ttaatggagg	gtccgtttcg	1440
gtttgggtga	agcaa					1455

<210> SEQ ID NO 15  
 <211> LENGTH: 1455  
 <212> TYPE: DNA  
 <213> ORGANISM: Bacillus sp.

<400> SEQUENCE: 15

catcataatg	ggacaaatgg	gacgatgatg	caatactttg	aatggcactt	gcctaataatg	60
gggaatcact	ggaatagatt	aagagatgat	gctagtaatc	taagaaatag	aggataaacc	120
gctatttggg	ttccgcctgc	ctggaaaggg	acttcgcaa	atgatgtggg	gtatggagcc	180
tatgatcttt	atgatttagg	ggaatttaat	caaaagggga	cggttcgtac	taagtatggg	240
acacgtagtc	aattggagtc	tgccatccat	gctttaaaga	ataatggcgt	tcaagtttat	300
gggagtag	tgatgaacca	taaaggagga	gctgatgcta	cagaaaacgt	tcttgctgtc	360
gaggtagatc	caaataaccg	gaatcaagaa	atatctgggg	actacacaat	tgaggcttgg	420
actaagtttg	atthtccagg	gaggggtaat	acatactcag	actttaaatg	gcgttggtat	480
catttcgatg	gtgtagattg	ggatcaatca	cgacaattcc	aaaatcgtat	ctacaaattc	540
cgaggtagatg	gtaaggcatg	ggattgggaa	gtagattcgg	aaaatggaaa	ttatgattat	600
ttaatgtatg	cagatgtaga	tatggatcat	ccggaggtag	taaagagct	tagaagatgg	660
ggagaatggt	atacaatac	attaaatctt	gatggattta	ggatcgtatg	ggtgaagcat	720
attaaatata	gctttacacg	tgattgggtg	acccatgtaa	gaaacgcaac	gggaaaagaa	780
atgtttgctg	ttgctgaatt	ttggaaaaat	gatttaggtg	ccttgagaa	ctatttfaat	840
aaaacaaact	ggaatcattc	tgtctttgat	gtcccccttc	attataatct	ttataacgcg	900
tcaaatagtg	gaggcaacta	tgacatggca	aaacttctta	atggaacggt	tgttcaaaaag	960
catccaatgc	atgccgtaac	ttttgtggat	aatcacgatt	ctcaacctgg	ggaatcatta	1020
gaatcatttg	tacaagaatg	gtttaagcca	cttgcttatg	cgcttatttt	aacaagagaa	1080
caaggctatc	cctctgtctt	ctatgggtgac	tactatggaa	ttccaacaca	tagtgtccca	1140
gcaatgaaag	ccaagattga	tccaatctta	gaggcgcgtc	aaaattttgc	atatggaaca	1200
caacatgatt	atthtgacca	tcataatata	atcgatgga	cacgtgaagg	aaataccacg	1260
catcccaatt	caggacttgc	gactatcatg	tcggatgggc	cagggggaga	gaaatggatg	1320
tacgtagggc	aaaataaagc	aggtcaagtt	tggcatgaca	taactggaaa	taaaccagga	1380
acagttacga	tcaatgcaga	tggatgggct	aatttttcag	taaagggagg	atctgtttcc	1440
atthgggtga	aacga					1455

<210> SEQ ID NO 16  
 <211> LENGTH: 23  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer BSG1

<400> SEQUENCE: 16

ccatgatgca	gtatthtgaa	tgg	23
------------	------------	-----	----

<210> SEQ ID NO 17  
 <211> LENGTH: 22  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer BSG3

<400> SEQUENCE: 17

gtcaccataa	aagacgcacg	gg	22
------------	------------	----	----

<210> SEQ ID NO 18  
 <211> LENGTH: 68  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:



-continued

---

<223> OTHER INFORMATION: Primer BSGM1

<400> SEQUENCE: 18

gtcatagttt ccgaattccg tgtctacttc ccaatcccaa tcccaagctt tgccgcggaa	60
tttgtaaa	68

<210> SEQ ID NO 19

<211> LENGTH: 41

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer BSGM2

<400> SEQUENCE: 19

ctacttccca atcccaagct ttgccgcgga atttgtaaat g	41
---	----

<210> SEQ ID NO 20

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer BSGM3

<400> SEQUENCE: 20

ggatgatcca tgtcaaagtc ggcatac	27
-------------------------------	----

<210> SEQ ID NO 21

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer BSGM4

<400> SEQUENCE: 21

ctcggtcacc acgtggggat gatcc	25
-----------------------------	----

<210> SEQ ID NO 22

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Primer BSGM5

<400> SEQUENCE: 22

ccagtttttc agctgggtca cgac	24
----------------------------	----

---

What is claimed is:

1. A polypeptide having  $\alpha$ -amylase activity comprising a polypeptide having at least two alterations relative to a parent  $\alpha$ -amylase, wherein

- A) In SEQ ID NO:1, at least one of said alterations is selected from the group consisting of R181\*, G182\*, T183\*, and G184\*; and at least one of said alterations is selected from the group consisting of N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y, and V;
- B) In SEQ ID NO:2, at least one of said alterations is selected from the group consisting of R181\*, G182\*, D183\*, and G184\* and at least one of said alterations is selected from the group consisting of N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y, and V;
- C) In SEQ ID NO:3, at least one of said alterations is selected from the group consisting of R179\*, G180, I181\*, and G182\* and at least one of said alterations is selected from the group consisting of N193A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y,V;

- D) In SEQ ID NO:4, at least one of said alterations is selected from the group consisting of Q178\* and G179\* and at least one of said alterations is selected from the group consisting of N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y, and V;
- E) In SEQ ID NO:5, at least one of said alterations is selected from the group consisting of R176\*, G177\*, E178, G179\* and at least one of said alterations is selected from the group consisting of N190A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y, and V;
- E) In SEQ ID NO:6, at least one of said alterations is selected from the group consisting of R181\*, G182\*, H183\*, and G184\* and at least one of said alterations is selected from the group consisting of N195A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y, and V; and
- G) in an  $\alpha$ -amylase polypeptide having at least 60% homology to any of SEQ ID NO:s 1-6 or combinations thereof,



- (i) at least one of said alterations is a deletion of a residue corresponding to R181, G182, T183, or G184 of SEQ ID NO:1; R181, G182, T183, or G184 of SEQ ID NO:2; R179, G180, I181, or G182 of SEQ ID NO:3; Q178 or G179 in SEQ ID NO:4; R176, G177, E178, or G179 in SEQ ID NO:5; or R181, G182, H183, or G184 in SEQ ID NO:6 and
- (ii) at least one of said alterations is a substitution of a residue corresponding to N195 in SEQ ID Nos: 1, 2, or 6; N193 in SEQ ID NO:3; or N190 in SEQ ID NOs:4 or 5 with an amino acid selected from the group consisting of A,R,D,C,E,Q,G,H,I,L,K,M,F,P,S,T,W,Y, and V.
2. A polypeptide as defined in claim 1, wherein said alterations comprise I181\*, G182\*, and N193F in SEQ ID NO: 3 or in corresponding positions in another parent  $\alpha$ -amylase.
3. A polypeptide as defined in claim 2, wherein said alterations further comprise a substitution in position E214Q in SEQ ID NO: 3 or in a corresponding position in another parent  $\alpha$ -amylase.
4. A polypeptide as defined in claim 1, wherein the parent  $\alpha$ -amylase is a hybrid  $\alpha$ -amylase of SEQ ID NO: 4 and SEQ ID NO: 5.
5. A polypeptide as defined in claim 4, wherein the parent hybrid alpha-amylase comprises the 445 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the  $\alpha$ -amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 5.
6. A polypeptide as defined in claim 5, wherein the parent hybrid further comprises: H156Y+A181T+N190F+A209V+Q264S (using the numbering in SEQ ID NO: 4).
7. A polypeptide as defined in claim 1, wherein said at least one alteration results in increased stability at acidic pH and/or at low  $\text{Ca}^{2+}$  concentration relative to the parent  $\alpha$ -amylase.
8. A detergent additive comprising a polypeptide as defined in claim 1.
9. A detergent additive according to claim 8 comprising 0.02–200 mg of enzyme protein/g of the additive.
10. A detergent additive according to claim 8, further comprising an enzyme selected from the group consisting of a protease, a lipase, a peroxidase, another amyolytic enzyme, a cellulase, and combinations of any of the foregoing.
11. A detergent composition comprising a polypeptide as defined in claim 1.
12. The detergent composition according to claim 11 further comprising an enzyme selected from the group consisting of a protease, a lipase, a peroxidase, another amyolytic enzyme, a cellulase, and combinations of any of the foregoing.
13. A manual or automatic dishwashing detergent composition comprising a polypeptide as defined in claim 1.
14. A dishwashing detergent composition according to claim 13 further comprising an enzyme selected from the

group consisting of a protease, a lipase, a peroxidase, another amyolytic enzyme, a cellulase, and combinations of any of the foregoing.

15. A manual or automatic laundry washing composition comprising a polypeptide as defined in claim 1.
16. A laundry washing composition according to claim 15 further comprising an enzyme selected from the group consisting of a protease, a lipase, a peroxidase, an amyolytic enzyme, a cellulase, and combinations of any of the foregoing.
17. A composition comprising a mixture selected from the group consisting of:
- (i) a mixture of the  $\alpha$ -amylase from *B. licheniformis* having the sequence shown in SEQ ID NO: 4 with one or more polypeptides as defined in claim 1, wherein said polypeptides are derived from a parent  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 3;
- (ii) a mixture of the  $\alpha$ -amylase from *B. stearotherophilus* having the sequence shown in SEQ ID NO: 3 with one or more polypeptides as defined in claim 1, wherein said polypeptides are derived from one or more other parent Termamyl-like  $\alpha$ -amylases; and
- (iii) a mixture of one or more polypeptides as defined in claim 1 derived from a parent  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 3 with one or more polypeptides as defined in claim 1 derived from a different parent  $\alpha$ -amylase.
18. A composition comprising:
- a mixture of (i) one or more polypeptides as defined in claim 1 derived from a parent  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 3 and (ii) one or more polypeptides as defined in claim 1 derived from a parent  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 4.
19. A composition comprising:
- a mixture of (i) one or more polypeptides as defined in claim 1 derived from a parent  $\alpha$ -amylase having the sequence shown in SEQ ID NO: 3 and (ii) a hybrid  $\alpha$ -amylase comprising a part of the *B. amyloliquefaciens*  $\alpha$ -amylase shown in SEQ ID NO: 5 and a part of the *B. licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4.
20. The composition according to claim 19, wherein the hybrid  $\alpha$ -amylase is a hybrid  $\alpha$ -amylase comprising the 445 C-terminal amino acid residues of the *B. licheniformis*  $\alpha$ -amylase shown in SEQ ID NO: 4 and the 37 N-terminal amino acid residues of the  $\alpha$ -amylase derived from *B. amyloliquefaciens* shown in SEQ ID NO: 5.
21. The composition according to claim 20, wherein the hybrid  $\alpha$ -amylase further comprises the following alterations relative to SEQ ID NO:4: H156Y+A181T+N190F+A209V+Q264S.
22. The composition according to claim 19, comprising a mixture of TVB146 and LE174.