



US006255526B1

(12) **United States Patent**  
**Pesachovich et al.**

(10) **Patent No.: US 6,255,526 B1**  
(45) **Date of Patent: \*Jul. 3, 2001**

(54) **PREPARATION OF GABAPENTIN**

(75) Inventors: **Michael Pesachovich**, Givat Shmuel;  
**Claude Singer**, Kfar Saba; **Gideon Pilarski**, Holon, all of (IL)

(73) Assignee: **Teva Pharmaceutical Industries Ltd.**,  
Petah Tiqva (IL)

(\* ) Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 26 days.

(21) Appl. No.: **09/331,555**

(22) PCT Filed: **Dec. 24, 1997**

(86) PCT No.: **PCT/US97/23164**

§ 371 Date: **Jan. 14, 2000**

§ 102(e) Date: **Jan. 14, 2000**

(87) PCT Pub. No.: **WO98/28255**

PCT Pub. Date: **Jul. 2, 1998**

(30) **Foreign Application Priority Data**

Dec. 24, 1996 (IL) ..... 119890

(51) **Int. Cl.<sup>7</sup>** ..... **C07C 61/06**

(52) **U.S. Cl.** ..... **562/507**

(58) **Field of Search** ..... **562/507**

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

4,024,175	5/1977	Satzinger et al. .	
4,894,476	1/1990	Butler et al. ....	562/504
4,960,931	10/1990	Butler et al. .	
5,068,413	11/1991	Steiner et al. .	
5,095,148	3/1992	Mettler, et al. ....	562/507
5,132,451	7/1992	Jennings et al. ....	562/507

*Primary Examiner*—Michael L. Shippen

(74) *Attorney, Agent, or Firm*—Kenyon & Kenyon

(57) **ABSTRACT**

There is disclosed a method of converting gabapentin hydrochloride substantially free of inorganic salts to gabapentin form II. The method comprises the steps of: (1) obtaining gabapentin hydrochloride that is substantially free of inorganic salts; (2) mixing a solution of the gabapentin hydrochloride with an additional amine in a first solvent so as to obtain a precipitate comprising gabapentin; and then (3) recovering gabapentin form II from the precipitate. The precipitated gabapentin is a novel polymorphic form of gabapentin possessing a crystalline structure characterized by novel sets of peaks in the powder X-ray diffraction pattern and in the FTIR spectra. This newly disclosed polymorph of gabapentin is characterized herein as gabapentin form III. The recovery step may comprise, for example, one of two alternative methods, slurring the gabapentin form II in methanol, and then filtering the suspension to obtain gabapentin form II, or solubilizing the gabapentin form III in methanol with heating by reflux, and then cooling the solution to obtain gabapentin form II by crystallization.

**19 Claims, 7 Drawing Sheets**




FIG. 1





FIG. 2



2θ (DEGREES)

FIG. 3A




FIG. 3B




FIG. 4




FIG. 5




FIG. 6




## PREPARATION OF GABAPENTIN

### FIELD OF THE INVENTION

This invention relates to a new process for converting gabapentin hydrochloride salt to gabapentin via a novel polymorphic form of gabapentin.

### BACKGROUND OF THE INVENTION

Gabapentin is 1-(aminomethyl)-1-cyclohexaneacetic acid, having the chemical structure:



Gabapentin is used in the treatment of cerebral diseases such as epilepsy. The literature describes many ways of preparing gabapentin from a variety of starting materials. U.S. Pat. No. 4,024,175 describes at least three methods of preparing gabapentin from cyclohexyl-1,1-diacetic acid. Each of these methods results in the formation of gabapentin hydrochloride salt, which may be converted to 1-(aminomethyl)-1-cyclohexaneacetic acid by treatment with a basic ion exchanger and then crystallized from a solvent such as ethanol/ether.

U.S. Pat. No. 4,894,476 specifically discloses an improved method for converting the hydrochloride salt into the free amino acid. This involves pouring a deionized water solution of the salt over an ion exchange column, eluting with deionized water, producing a slurry from the eluate, adding an alcohol to the slurry, centrifuging and drying the slurry to obtain the free amino acid.

Alternative methods for preparing gabapentin have been described that do not proceed via the hydrochloride or any other mineral acid salt. Such methods include those described in U.S. Pat. Nos. 5,132,451, 5,095,148, 5,068,413. Each of these methods involve a cyanic intermediate which is hydrogenated under severe conditions to produce the free amino acid.

These methods are industrially impractical. Those methods comprising ion exchange columns require the use of large amounts of ion exchanger for lengthy periods of time to lower the level of chloride ions to the desired level. The alternative methods involve further more demanding steps.

Commercially available gabapentin is crystalline and exhibits an X-ray diffraction pattern with peaks of 2-theta values at 7.8, 13.3, 15.0, 17.0, 20.4, 21.3, 23.1, 23.6, 25.7, 27.0 and 28.2 degrees. Hereinafter, the commercially available polymorphic form of gabapentin is referred to as polymorph form "II".

### SUMMARY OF THE INVENTION

The present invention relates to an improved method for purifying gabapentin comprising converting gabapentin hydrochloride salt to gabapentin form II. The present invention avoids the disadvantages associated with prior art methods, by adding alternative steps and by proceeding via a novel polymorphic form of gabapentin.

Accordingly, the present invention relates to a method of converting gabapentin hydrochloride salt to gabapentin form II, comprising reacting a solution of gabapentin hydrochloride with an additional amine in a first solvent to produce a novel polymorphic form as a precipitate, thereafter, convert-

ing the novel polymorphic form to form II by forming a suspension and/or a solution of the precipitate in methanol, and then recovering gabapentin form II.

The present invention further relates to a novel polymorphic form of gabapentin designated as gabapentin form III. The polymorph may be identified by its unique X-ray diffraction pattern.

A further aspect of the present invention relates to the novel polymorphic form of gabapentin that is of use as an intermediate in the preparation of polymorphic form II.

Further objectives and advantages of the subject invention will be apparent to those skilled in the art from the detailed description of the disclosed invention.

### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is an x-ray diffraction pattern of a sample of gabapentin hydrate.

FIG. 2 is an x-ray diffraction pattern of gabapentin form II.

FIGS. 3a and 3b are x-ray diffraction pattern of gabapentin form III, possibly containing small amounts of gabapentin form II and/or gabapentin hydrate.

FIG. 4 is an FTIR (Fourier Transform Infra Red) spectrum of gabapentin hydrate.

FIG. 5 is an FTIR spectrum of gabapentin form II.

FIG. 6 is an FTIR spectrum of gabapentin form III, possibly containing small amounts of gabapentin form II and/or gabapentin hydrate.

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

The subject invention will now be described in detail for specific preferred embodiments of the invention, it being understood that these embodiments are intended only as illustrative examples and the invention is not to be limited thereto.

The method of the present invention comprises:

- (1) obtaining gabapentin hydrochloride that is substantially free of inorganic salts;
- (2) mixing a solution of the gabapentin hydrochloride with an additional amine in a first solvent so as to obtain a precipitate comprising gabapentin; and then
- (3) recovering gabapentin form II from the precipitate.

Preferably, the gabapentin hydrochloride used as a starting material in the process of the present invention is substantially free of other inorganic salts such as sodium chloride and sodium bromide, that is, such impurities are only present in trace amounts. Alternatively, gabapentin hydrochloride containing inorganic salts may be used subject to the addition of a further step for removing the inorganic salts prior to mixing gabapentin hydrochloride with the additional amine.

Thus, gabapentin hydrochloride containing inorganic salts may optionally be pre-treated to remove the inorganic salts by the steps of (a) dissolution in a solvent in which gabapentin hydrochloride is soluble, but the inorganic salts are not; (b) filtration of the inorganic salts and, optionally, (c) evaporating the solvent to recover gabapentin hydrochloride substantially free of inorganic salts. Appropriate solvents for the optional pre-treatment step include those selected from the group consisting of methanol, ethanol, n-propanol, isopropanol, butanol, t-butanol, n-butanol, ethyleneglycolmonomethylether, benzylalcohol or dimethylacetamide. Preferably the pre-treatment solvent is identical

to the first solvent. This pre-treatment solvent, which is identified in Table 1 and in the claims as the "second solvent", would be used prior to using the "first solvent", if, in fact, this optional pre-treatment step is included in the process.

The second step of the present invention comprises mixing a solution of the gabapentin hydrochloride in a solvent that contains an additional amine. The solvent may be any solvent in which the hydrochloride salt of the additional amine is soluble but in which gabapentin form III is insoluble, such that a precipitate of gabapentin form III is formed. Such solvents are preferably selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide. The precipitated gabapentin, which may be separated by filtration, is characterized herein as a novel polymorphic form of gabapentin, possessing a crystalline structure characterized by peaks in the powder X ray diffraction pattern with 2-theta values at 6.11, 12.22, 17.00, 18.20, 19.94, 20.81, 24.54, and 25.11 degrees, all  $\pm 0.2$  degrees. The x-ray diffraction pattern in the samples obtained appears to show a preferred orientation in which the peak at 6.11 degrees is larger than any other peak in the pattern, and the peaks at 12.22 and 24.54 degrees are larger than any of the remaining peaks in the pattern. This polymorph is referred to herein as gabapentin form III.

Two representative embodiments of the third step of the present invention, recovery of gabapentin form II from the precipitate, are slurring and crystallization. Slurring may be performed by suspending the precipitated gabapentin in methanol by mixing, stirring, and/or providing continuous agitation with some mechanical device so as to induce transformation into the gabapentin form II that is the commercially available polymorphic form of gabapentin. Gabapentin form II may then be filtered off and washed.

Alternatively, the precipitated gabapentin may be crystallized from methanol with heating by reflux until dissolved, cooling, optionally seeding with gabapentin, followed by further cooling, and then collecting and drying the crystals of gabapentin form II. Second and multiple crops may be obtained from the concentrated mother liqueurs.

Suitable amines for use in the present invention include triethylamine, tributylamine, tripropylamine, trihexylamine, diethylamine, ethanolamine and benzylamine. Preferably the amine is tributylamine.

The form II gabapentin obtained by the methods of the present invention may be crystallized using processes known in the art.

Certain specific representative embodiments of the invention are described in detail below, the materials, apparatus and process steps being understood as examples that are intended to be exemplary and illustrative only. In particular, the invention is not intended to be limited to the methods, materials, conditions, process parameters, apparatus and the like specifically recited herein.

### EXAMPLES

#### 1. Preparation of Gabapentin from Gabapentin Hydrochloride

##### A. Removal of Inorganic Salts

Gabapentin hydrochloride containing inorganic salts (NaCl, NaBr), which may be prepared, for example, as shown in U.S. Pat. No. 4,024,175, Example 4, Variant A,

was dissolved in 150 ml isopropanol and mixed at 25° C. Active carbon (0.9 g) was added and the suspension mixed for a further 2 hours. The inorganic salts were removed by filtering. The filter cake was washed twice with 15 ml isopropanol and the washings were added to the gabapentin hydrochloride solution.

##### B. Precipitation of Gabapentin Base; Formation of Gabapentin Form III

The gabapentin hydrochloride solution of part A was concentrated to dryness in vacuum while ensuring that the temperature of the heating bath did not exceed 35° C. 210 ml ethylacetate and 16.5 ml tributylamine were added and the solution was mixed for 2 hours at 25° C. The gabapentin precipitate was then separated by filtration, the filter cake being washed with 20 ml ethylacetate and then 20 ml methanol. The filter cake contained gabapentin form III which when dried displayed a characteristic X-ray diffraction pattern with 2-theta values at 6.1, 12.2, 17.0, 17.7, 18.3, 20.0, 20.8, 24.6 and 25.5 degrees.

##### C. Alternative Methods for Conversion of Precipitated Gabapentin to Gabapentin Form II

###### (1) Slurring

The still humid filter cake from step 2 was suspended in 52.5 ml methanol for about 14 hours at 25° C. Solid gabapentin was then separated from the suspension by filtration. The filter cake was washed with 20 ml methanol and then dried under vacuum at 35° C. 10.8 g crystalline gabapentin form II (yield 72%) was obtained.

###### (2) Crystallization

The still humid filter cake from step 2 was suspended in 210 ml of methanol and the suspension was heated to reflux (65° C.). If dissolution was incomplete, additional methanol was added.

After dissolution, the reactor content was cooled to 34° C. at which temperature crystallization was induced by seeding with pure Form II gabapentin base (0.1 g).

After maintaining the mixture at 34° C. for 60 minutes the reactor content was cooled to 25° C. and methanol was distilled by vacuum distillation. Approximately 160 ml of methanol was collected. Then the suspension was cooled to 0–10° C. and maintained at this temperature for 2 hours.

The crystalline gabapentin was separated by filtration from the suspension. The filter cake was washed with 20 ml methanol and then dried under vacuum at 35° C. 10.8 g of crystalline gabapentin form II (yield 72%) was thus obtained.

#### Examples 2–20

The method of Example 1 was followed using the slurring technique of step C(1) and employing the amines and solvents shown in Table 1 below. The percent yields are of purified product except where marked with an asterisk (\*), where the yield was measured at the stage prior to the slurring in methanol.

TABLE 1

Summary of the first solvent, second solvent, amine and yield of examples 2–20.				
Example No.	First Solvent	Second Solvent	Amine	Yield (%)
2	IPA	IPA	TBA	61.5
3	Et Ac	IPA	TBA	74.0
4	Et Ac	IPA	THA	66.2
5	Et Ac	IPA	TPA	63.0
6	Et Ac	IPA	THA	79.6*

TABLE 1-continued

Summary of the first solvent, second solvent, amine and yield of examples 2-20.				
Example No.	First Solvent	Second Solvent	Amine	Yield (%)
7	IPA Ac	IPA	TBA	64.9
8	ACN	IPA	TBA	67.8
9	DMC	IPA	TBA	57.9
10	DMA	DMA	TBA	65.7
11	BzOH	BzOH	TBA	43.0*
12	MEK	IPA	TBA	88.0*
13	t-BuOH	t-BuOH	TBA	79.4*
14	Acetone	IPA	TBA	73.1*
15	Et Ac	BuOH	TBA	69.8
16	MeOH	MeOH	TBA	67.4
17	EGMME	EGMME	TBA	66.8*
18	IPA	IPA	TEA	76.2*
19	IPA	IPA	BzA	56.0*
20	CH <sub>2</sub> Cl <sub>2</sub>	DMA	DEA	89.4*

IPA isopropanol  
 ACN acetonitrile  
 BzOH benzyl alcohol  
 MeOH methanol  
 CH<sub>2</sub>Cl<sub>2</sub> methylene chloride  
 THA trihexylamine  
 TPA tripropylamine  
 Et Ac ethyl acetate  
 MEK methyl ethyl ketone  
 EGMME ethyleneglycol monomethylether  
 TBA tributylamine  
 BzA benzylamine  
 IPA Ac isopropylacetate  
 DMC dimethylcarbonate  
 (t)-BuOH (tert)-butanol  
 TEA triethylamine

The gabapentin form III was characterized by comparing the x-ray diffraction pattern and the FTIR absorption spectra of gabapentin form III with gabapentin form II and gabapentin hydrate. The x-ray powder diffraction patterns of FIGS. 1, 2, and 3a were obtained using a Philips x-ray powder diffractometer with the following parameters:

Goniometer model 1050/70, Cu-tube, Curved graphite monochromator.

Sample holder:

Quartz monocrystal plate

Settings:

X-Ray tube:

KV-40

mA-28

Target-Cu

Divergence slit-1°

Receiving parallel slit-0.2 mm

Scatter slit-1°

Scintillation detector:

Voltage-832

Lower level-33.8%

Window-34.5%

Scan parameters:

Scanning speed:

2°/min

Paper speed:

2cm/min

Gain:

32

Calibration:

External calibration with silicon fine powder

Type of radiation:

copper K $\alpha$ .

FIG. 3b was obtained using a Siemens B5100 with a presample (K $\alpha$ <sub>1</sub> only) monochromator, step scan, Cu radiation, beam slits 0.3, receiving slit 0.05, standard Siemens rotating sample holder, start/stop/step angles in degrees were 4.0, 35, and 0.04, with a step duration of 15 seconds. Standard Siemens D-5000 software was used. The diffractometer was calibrated with NIST large d-spacing standard. No theta-compensating slits were employed.

The x-ray spectra of gabapentin hydrate, gabapentin form II and gabapentin form III are shown in FIGS. 1-3, respectively. The main diffraction peaks that characterize each material are listed in Table 2.

TABLE 2

X-ray Diffraction peaks of gabapentin hydrate, gabapentin form II and gabapentin form III.			
	Hydrate	Form II	Form III
	2-theta (°)	2-theta (°)	2-theta (°)
	6.1	7.9	6.11
	12.2	13.3	12.22
	16.0	15.0	17.00
	18.3	17.0	17.63
	19.1	19.5	18.20
	19.8	20.3	19.94
	20.7	21.3	20.81
	24.5	21.8	24.54
	26.4	23.0	25.11
	28.4	23.6	28.91
	30.7	25.7	30.20
	32.3	26.9	30.78
		28.2	31.46

The FTIR spectra for gabapentin hydrate, gabapentin form II and gabapentin form III are shown in FIGS. 4-6, respectively. The FTIR peaks are summarized in Table 3.

TABLE 3

FTIR peaks of gabapentin hydrate, gabapentin form II and gabapentin form III.			
	Form I	Form II	Form III
	cm <sup>-1</sup>	cm <sup>-1</sup>	cm <sup>-1</sup>
	648	709	708
	726	749	760
	880	890	885
	926	922,928	926
	968	976	974
	1154	1165	1160
	1175		1180
	1292	1300	1290
		1420	1420
		1476	1460
			1510
	1542	1546	1586
	1624	1615	1664
	1662		

Morphology of the three forms:

1. The hydrate form typically exists as large crystals with undefined shapes.
2. Form II typically exists as plate shaped crystals.
3. Form III typically exists as small rhomboidal crystals.

The melting point for gabapentin was not determined since gabapentin decomposes prior to melting.

What is claimed is:

1. A method of converting gabapentin hydrochloride to gabapentin form II comprising:

- (1) dissolving gabapentin hydrochloride in a first solvent in which gabapentin is relatively insoluble; and
- (2) adding an additional amine to the gabapentin hydrochloride solution so as to obtain a precipitate comprising gabapentin.

7

2. The method of claim 1, further comprising the step of recovering gabapentin form II from said precipitate.

3. The method of claim 2, the recovering step comprising slurring or crystallizing said precipitate in methanol.

4. The method of claim 3, wherein said first solvent is selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide.

5. The method of claim 1 wherein said additional amine is selected from the group consisting of triethylamine, tributylamine, tripropylamine, trihexylamine, diethylamine, ethanolamine and benzylamine.

6. The method of claim 5 wherein said additional amine is tributylamine.

7. The method of claim 1 wherein said first solvent is selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide.

8. The method of claim 7 wherein said first solvent is ethylacetate.

9. The method of claim 1 wherein said first solvent is one in which the hydrochloride salt of said additional amine is soluble.

10. The method of claim 1 further comprising pre-treating said gabapentin hydrochloride to remove inorganic salts, which pre-treatment comprises (a) dissolving gabapentin hydrochloride in a second solvent in which inorganic salts are insoluble, (b) filtering off the inorganic salts, and (c) recovering gabapentin hydrochloride from said second solvent.

11. The method of claim 10 wherein said second solvent is selected from the group consisting of methanol, ethanol, n-propanol, isopropanol, butanol, t-butanol, n-butanol, ethyleneglycolmonomethylether, benzylalcohol or dimethylacetamide.

12. The method of claim 11 wherein said second solvent is isopropanol.

8

13. The method of claim 1 or 10 wherein the precipitated gabapentin comprises gabapentin form III.

14. The method of claim 10 wherein the precipitated gabapentin comprises gabapentin form III, said second solvent is isopropanol, said first solvent is ethylacetate, and said additional amine is tributylamine.

15. The method of claim 14 wherein said precipitated gabapentin is slurried or crystallized from methanol to yield gabapentin form II.

16. Crystalline gabapentin form III characterized by powder X-ray diffraction peaks at  $6.11\pm 0.2$ ,  $12.22\pm 0.2$ ,  $17.00\pm 0.2$ ,  $17.63\pm 0.2$ ,  $18.20\pm 0.2$ ,  $19.94\pm 0.2$ ,  $20.81\pm 0.2$ ,  $24.54\pm 0.2$ ,  $25.11\pm 0.2$ ,  $28.91\pm 0.2$ ,  $30.20\pm 0.2$ ,  $30.78\pm 0.2$ , and  $31.46\pm 0.2$  degrees 2-theta.

17. Crystalline gabapentin form III characterized by powder X-ray diffraction peaks at  $6.11\pm 0.2$ ,  $12.22\pm 0.2$ ,  $17.00\pm 0.2$ ,  $17.63\pm 0.2$ ,  $18.20\pm 0.2$ ,  $19.94\pm 0.2$ ,  $20.81\pm 0.2$ ,  $24.54\pm 0.2$ ,  $25.11\pm 0.2$ ,  $28.91\pm 0.2$ ,  $30.20\pm 0.2$ ,  $30.78\pm 0.2$ , and  $31.46\pm 0.2$  degrees 2-theta, wherein the size of the peaks is in the order  $6.11 > 12.22$  and  $24.54 > 17.00$ ,  $17.63$ ,  $18.20$ ,  $19.94$ ,  $20.81$ ,  $25.11$ ,  $28.91$ ,  $30.20$ ,  $30.78$ , and  $31.46$ .

18. Crystalline gabapentin form III characterized by infrared absorptions having peaks at 708, 760, 885, 926, 974, 1160, 1180, 1290, 1420, 1460, 1510, 1586, and  $1664 \text{ cm}^{-1}$ .

19. Crystalline gabapentin form III produceable by a process comprising the steps of:

- (a) dissolving gabapentin hydrochloride in a solvent selected from the group consisting of ethyl acetate, dimethylcarbonate, ethanol, butanol, t-butanol, n-butanol, methanol, acetonitrile, toluene, isopropylacetate, isopropanol, methylethylketone, acetone, ethyleneglycolmonomethylether, methylene chloride, chloroform, benzylalcohol or dimethylacetamide;
- (b) adding an additional amine selected from the group consisting of triethylamine, tributylamine, tripropylamine, trihexylamine, diethylamine, ethanolamine and benzylamine; and
- (c) drying the precipitated gabapentin.

\* \* \* \* \*