



US006288034B1

(12) **United States Patent**  
**Murgita**

(10) **Patent No.:** **US 6,288,034 B1**  
(45) **Date of Patent:** **\*Sep. 11, 2001**

(54) **RECOMBINANT HUMAN ALPHA-FETOPROTEIN AS AN IMMUNOSUPPRESSIVE AGENT**

(75) Inventor: **Robert A. Murgita**, Montreal (CA)

(73) Assignee: **Martinx R & D Inc.** (CA)

(\* ) Notice: This patent issued on a continued prosecution application filed under 37 CFR 1.53(d), and is subject to the twenty year patent term provisions of 35 U.S.C. 154(a)(2).

Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **09/186,723**

(22) Filed: **Nov. 5, 1998**

**Related U.S. Application Data**

(63) Continuation of application No. 08/377,309, filed on Jan. 24, 1995, now Pat. No. 5,965,528.

(51) **Int. Cl.**<sup>7</sup> ..... **A61K 38/00**; C07K 14/47

(52) **U.S. Cl.** ..... **514/12**; 530/350

(58) **Field of Search** ..... 514/11, 12; 435/69.1; 424/88

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

|           |         |                |          |
|-----------|---------|----------------|----------|
| 4,692,332 | 9/1987  | McMichael      | 424/88   |
| 4,877,610 | 10/1989 | McMichael      | 424/88   |
| 4,966,753 | 10/1990 | McMichael      | 424/88   |
| 4,970,071 | 11/1990 | McMichael      | 424/88   |
| 4,996,193 | 2/1991  | Hewitt et al.  | 514/11   |
| 5,130,415 | 7/1992  | Tecce et al.   | 530/324  |
| 5,206,153 | 4/1993  | Tamaoki et al. | 435/69.7 |
| 5,365,948 | 11/1994 | McMichael      | 128/898  |
| 5,384,250 | 1/1995  | Murgita        | 435/69.1 |

**FOREIGN PATENT DOCUMENTS**

|             |        |        |
|-------------|--------|--------|
| 0 001 812   | 5/1979 | (EP) . |
| 20058666    | 1/1990 | (JP) . |
| WO 86/04241 | 7/1986 | (WO) . |
| WO 93/05774 | 4/1993 | (WO) . |
| WO 94/10199 | 5/1994 | (WO) . |

**OTHER PUBLICATIONS**

Abramsky et al., "Alpha-fetoprotein Suppresses Experimental Allergic Encephalomyelitis," *Journal of Neuroimmunology* 2:1-7 (1982).

Abramsky et al., *Annals New York Academy of Sciences*, pp. 108-115 (1983).

Aoyagi et al., "Differential Reactivity of  $\alpha$ -Fetoprotein with Lectins and Evaluation of Its Usefulness in the Diagnosis of Hepatocellular Carcinoma", *Gann* 75:809-815 (1984).

Biddle et al., "Specific Cytoplasmic Alpha-fetoprotein Binding Protein In MCF-7 Human Breast Cancer Cells and Primary Breast Cancer Tissue," *Breast Cancer Research and Treatment* 10:279-286 (1987).

Boismenu et al., "Expression of Domains of Mouse Alpha-Fetoprotein in *Escherichia coli*," *Life Sciences* 43:673-681 (1988).

Brenner et al., "Influence Of Alpha-fetoprotein On The In Vitro And In Vivo Immune Response To Acetylcholine Receptor," *Annals New York Academy of Sciences* 377:208-221 (1981).

Brenner et al., "Inhibitory Effect Of  $\alpha$ -fetoprotein On the Binding Of *Myasthenia gravis* Antibody To Acetylcholine Receptor," *Proc. Natl. Acad. Sci. USA* 77:3635-3639(1980).

Brenner et al., "Immunosuppression of Experimental Autoimmune *Myasthenia gravis* by Alpha-Fetoprotein Rich Formation," *Immunology Letters* 3:163-167 (1981).

Buamah et al., "Serum alpha fetoprotein heterogeneity as a means of differentiating between primary hepatocellular carcinoma and hepatic secondaries," *Clinica Chimica Acta*. 139:313-316 (1984).

Buschman et al., "Experimental *Myasthenia gravis* Induced in Mice by Passive Transfer of Human Myasthenic Immunoglobulin," *Journal of Neuroimmunology*, 13:315-330 (1987).

Cohen et al., "Suppression by Alpha-Fetoprotein of Murine Natural Killer Cell Activity Stimulated in Vitro and in Vivo by Interferon and Interleukin 2," *Scand. J. Immunol.* 23:211-223 (1986).

Dattwyler et al., "Binding Of  $\alpha$ -foetoprotein To Murine T Cells," *Nature* 256:656-657 (1975).

Fialova et al., "Serum Levels of Trophoblast-Specific Beta-1-Globulin (SP1) and Alpha-1-Fetoprotein (AFP) in Pregnant Women With Rheumatoid Arthritis," *Ceskoslovenska Gynekologie* 56:166-170 (1991) (Abstract only).

Gershwin et al., "The Influence of  $\alpha$ -Fetoprotein on Moloney Sarcoma Virus Oncogenesis: Evidence For Generation of Antigen Nonspecific Suppressor T Cells," *The Journal of Immunology* 121:2292-2297 (1978).

Giuliani et al., "Synthesis And Characterization Of A Recombinant Fragment Of Human  $\alpha$ -fetoprotein With Antigenic Selectivity Versus Albumin," *Protein Engineering* 2:605-610 (1989).

Glazier et al., "Graft-Versus-Host Disease in Cyclosporin A-Treated Rats After Syngeneic and Autologous Bone Marrow Reconstitution," *J. Exp. Med.* 158:1-8 (1983).

Goidl et al., "Studies On The Mechanisms of Alpha-Fetoprotein Induction of Immune Suppressive Activity," *Developmental Immunobiology* pp. 35-55 (1979).

(List continued on next page.)

*Primary Examiner*—Kenneth R. Horlick

*Assistant Examiner*—Teresa Strzelecka

(74) *Attorney, Agent, or Firm*—Clark & Elbing LLP

(57) **ABSTRACT**

Disclosed are methods of inhibiting autoreactive immune cell proliferation in a mammal, involving administering to the mammal a therapeutically effective amount of recombinant human alpha-fetoprotein or an immune cell anti-proliferative fragment or analog thereof.

**13 Claims, 7 Drawing Sheets**

## OTHER PUBLICATIONS

- Hamel et al., "Phenotype and Function of Bone Marrow-Derived T- and Non-T-Cells Activated in Vitro By Alpha-Fetoprotein," In: *Biological Activities of Alpha-Fetoprotein* (vol. 1), Mizejewski, G.J. and Jacobson, H.I. (eds.), CRC Press, Inc. (Boca Raton, FL), pp. 167-177 (1987).
- Heyward et al., "Early Detection Of Primary Hepatocellular Carcinoma By Screening For Alpha-Fetoprotein In High-Risk Families," *The Lancet* 2:1161-1162 (1983).
- Hooper et al., "Human AFP Inhibits Cell Proliferation and NK-Like Cytotoxic Activity Generated in Autologous, But Not In Allogeneic Mixed Lymphocyte Reactions," In: *Biological Activities of Alpha-Fetoprotein*, (vol. II) Mizejewski, G.J. and Jacobson, H.I. (eds.), CRC Press, Inc. (Boca Raton, FL), pp. 183-197 (1989).
- Hooper et al., "Selective Inhibition Of Murine T-Cell Proliferation And Lymphokine-Activated Natural Killer Cell Function By alpha-Fetoprotein," In: *Biological Activities of Alpha-Fetoprotein*, (vol. 1) Mizejewski, G.J. and Jacobson, H.I. (eds.), CRC Press, Inc. (Boca Raton, FL), pp. 153-165 (1987).
- Hooper et al., "Regulation of Murine T-Cell Responses to Autologous Antigens by  $\alpha$ -Fetoprotein," *Cellular Immunology* 63:417-425 (1981).
- Hooper et al., "Suppression Of Primary And Secondary Autologous Mixed Lymphocyte Reactions By Murine Alphafetoprotein," *Oncodevelopmental Biology and Medicine* 3:151-160 (1982).
- Hoskin et al., "Analysis of Pregnancy-Associated Immunoregulatory Pathways," In: *Alpha-Fetoprotein and Congenital Disorders*, Academic Press, Inc. (New York), pp. 59-78 (1985).
- Hoskin et al., "In Vitro Activation of Bone Marrow-Derived T- and Non-T-Cells Subsets by  $\alpha$ -Fetoprotein," *Cellular Immunology* 96:163-174 (1985).
- Hoskin et al., "Specific Maternal Anti-fetal Lymphoproliferative Responses And Their Regulation By Natural Immunosuppressive Factors," *Clin. Exp. Immunol.* 76:262-267 (1989).
- Innis et al., "Amplification of  $\alpha$ -Fetoprotein Complementary DNA by Insertion into a Bacterial Plasmid," *Archives of Biochemistry and Biophysics* 195:128-135 (1979).
- Ishiguro et al., "Serum Alpha-Fetoprotein Subfractions in Patients With Primary Hepatoma or Hepatic Metastasis of Gastric Cancer," *Cancer* 55:156-159 (1985).
- Jacobson et al., "Inhibition of Estrogen-dependent Breast Cancer Growth by a Reaction Product of  $\alpha$ -Fetoprotein and Estradiol," *Cancer Research* 50:415-420 (1990).
- Jiang et al., "Role of CD8<sup>+</sup>T Cells In Murine Experimental Allergic Encephalomyelitis," *Science* 256:1213-1215 (1992).
- Keller et al., "Immunosuppressive Properties of AFP: Role Of Estrogens," In: *Onco-Developmental Gene Expression*, Fishman, W.H. and Sell, S. (eds.), Academic Press, Inc. (New York) pp. 287-295 (1976).
- Kikutani et al., "The Murine Autoimmune Diabetes Model: NOD and Related Strains," *Advances in Immunology* 51:285-322 (1992).
- Line et al., "Medical Potential Of AFP As A Tumor Imaging Agent," In: *Biological Activities of Alpha-Fetoprotein* (vol. II), Mizejewski, G.J. and Jacobson, H.I. (eds.), CRC Press, Inc. (Boca Raton, FL), pp. 139-148 (1989).
- Lu et al., " $\alpha$ -Fetoprotein Inhibits Macrophage Expressions Of Ia Antigens," *The Journal of Immunology* 132:1722-1727 (1984).
- Masuda et al., "Selective Antitumor Effect of Thioether-Linked Immunotoxins Composed of Gelonin and Monoclonal Antibody to Alpha-Fetoprotein or its F(ab')<sub>2</sub> Fragment," *Tumor Biol.* 15:175-183 (1994).
- Mizejewski et al., "Alpha Fetoprotein Testing: Regulatory and Technical Considerations," *Laboratory Management* (1987).
- Morinaga et al., "Primary Structures Of Human  $\alpha$ -fetoprotein And Its mRNA," *Proc. Natl. Acad. Sci. USA* 80:4604-4608 (1983).
- Moro et al., "Monoclonal Antibodies Directed against a Widespread Oncofetal Antigen: The Alpha-Fetoprotein Receptor," *Tumor Biol.* 14:116-130 (1993).
- Murgita et al., "Selective Immunoregulatory Properties Of  $\alpha$ -Fetoprotein," *La Ricerca Clin. Lab.* 9:327-342 (1979).
- Murgita et al., "Regulation of Immune Functions in the Fetus and Newborn," *Progress Allergy* 29:54-132 (1981).
- Murgita et al., "Suppression Of The Immune Response By  $\alpha$ -Fetoprotein," *The Journal of Experimental Medicine* 141:440-452 (1975).
- Murgita et al., "Suppression Of The Immune Response By  $\alpha$ -Fetoprotein," *The Journal of Experimental Medicine* 141:269-286 (1975).
- Murgita et al., "Adult murine T cells activated in vitro by  $\alpha$ -fetoprotein and naturally occurring T cells in newborn mice: Identity in function and cell surface differentiation antigens," *Proc. Natl. Acad. Sci. USA* 75:2897-2901 (1978).
- Murgita et al., "Effects of human alpha-foetoprotein on human B and T lymphocyte proliferation in vitro," *Clin. Exp. Immunol.* 33:347-356 (1978).
- Murgita et al., "The Effects of Mouse Alpha-Fetoprotein on T-Cell-Dependent and T-Cell-Independent Immune Responses In Vitro," *Scand. J. Immunol.* 5:1215-1220 (1976).
- Murgita et al., "The Immunosuppressive Role of Alpha-Fetoprotein During Pregnancy," *Scand. J. Immunol.* 5:1003-1014 (1976).
- Murgita et al., " $\alpha$ -Fetoprotein induces suppressor T cells in vitro," *Nature* 267:257-258 (1977).
- Murgita et al., "Characterization Of Murine Newborn Inhibitory T Lymphocytes: Functional And Phenotype Comparison With An Adult T Subset Activated in vitro By Alpha-fetoprotein," *Eur. J. Immunol.* 11:957-964 (1981).
- Nelson et al., "Maternal-Fetal Disparity in HLA Class II Alloantigens And The Pregnancy-Induced Amelioration of Rheumatoid Arthritis," *The New England Journal of Medicine* 329:466-471 (1993).
- Nishi et al., "Expression of Rat  $\alpha$ -Fetoprotein cDNA in *Escherichia coli* and in Yeast," *J. Biochem.* 104:968-972 (1988).
- O'Neill et al., "Regulation Of Human Lymphocyte Activation By Alpha-Fetoprotein: Evidence For Selective Suppression Of Ia-Associated T-Cell Proliferation In Vitro," *Oncodevelopmental Biology and Medicine* 3:135-150 (1982).
- Peck et al., "Cellular And Genetic Restrictions In The Immunoregulatory Activity Of  $\alpha$ -Fetoprotein," *The Journal of Immunology* 128:1134-1140 (1982).
- Peck et al., "Cellular And Genetic Restrictions In The Immunoregulatory Activity Of Alpha-Fetoprotein," *The Journal of Experimental Medicine* 147:667-683 (1978).

Peck et al., "Cellular And Genetic Restrictions In The Immunoregulatory Activity Of Alpha-Fetoprotein," J. Exp. Med. 148:360-372 (1978)..

Sell, "Alphafetoprotein," In: Cancer Markers Diagnostic and Developmental Significance, Sell, S., (ed.), Humana Press, Clifton, NJ pp. 249-293 (1980).

Semeniuk et al., "Immunoregulation By Recombinant Alpha-Fetoproteins Produced In Eukaryotic And Prokaryotic Expression Systems," Abstracts 2799, Experimental Biology 94<sup>TM</sup>, Anaheim, CA (1994).

Soto et al., "Control Of Growth Of Estrogen-sensitive Cells: Role For  $\alpha$ -fetoprotein," Proc. Natl. Acad. Acad. Sci. USA 77:2084-2087 (1980).

van Oers et al., "Analytical- And Preparative-Scale Separation Of Molecular Variants of  $\alpha$ -Fetoprotein By Anion-Exchange Chromatography On Monobead<sup>TM</sup> Resins," Journal of Chromatography 525:59-69 (1990).

van Oers et al., "Isolation And Characterization Of A Distinct Immunoregulatory Isoform Of  $\alpha$ -Fetoprotein Produced By The Normal Fetus," J. Exp. Med. 170:811-825 (1989).

Villacampa et al., "Alpha-Fetoprotein Receptors In A Human Breast Cancer Cell Line," Biochemical and Biophysical Research Communications 122:1322-1327 (1984).




Fig. 1A




Fig. 1B




Fig. 1C

Fig. 1D




Fig. 2A




Fig. 2B




Fig. 3

AT (2)  
 -1  
 met lys trp val glu ser ile phe leu ile phe leu asn phe thr glu ser arg  
 -10  
 -19  
 ATG AAG TGG GTG GAA TCA ATT TTT TTA ATT TTC CTA AAT TTT ACT GAA TCC AGA (101)  
 ATTTGGCTTCCACCCTGCCAAATAACAATAACTAGCAACC  
 1  
 10  
 thr leu his arg asn glu tyr gly ile ala ser ile leu asp ser tyr gln cys thr ala glu ile ser leu ala asp leu ala thr ile  
 20  
 ACA CTG CAT AGA AAT GAA TAT GGA ATA GCT TCC TCA TTT GAT TCT TAC CAA TGT ACT GCA ATA AGT TTA GCT GAC CTG ACC ATA (191)  
 30  
 40  
 phe phe ala gln phe val gln glu ala thr tyr lys glu val ser lys met val lys asp ala leu thr ala ile glu lys pro thr gly  
 50  
 TTT TTT GCC CAG TTT GTT CAA GAA GCC ACT TAC AAG GAA GTA AGC AAA ATG GTG AAA GAT GCA TTG ACT GCA ATT GAG AAA CCC ACT GGA (281)  
 60  
 70  
 61  
 asp glu gln ser ser gly cys leu glu asn gln leu pro ala phe leu glu glu lys glu ile leu glu lys tyr gly  
 80  
 GAT GAA GAG TCT TCA GGG TGT TTA GAA AAC CAG CTA CCT GCC TTT CTG GAA GAA CTT TGC CAT GAG AAA GAA ATT TTG GAG AAG TAC GGA (371)  
 90  
 100  
 91  
 his ser asp cys cys ser gln ser glu glu gln arg his asn cys phe leu ala his lys pro thr pro ala ser ile pro leu phe  
 110  
 CAT TCA GAC TGC TGC AGC CAA AGT GAA GAG GGA AGA CAT AAC TGT TTT CTT GCA CAC AAA AAG CCC ACT CCA GCA TCG ATC CCA CTT TTC (461)  
 120  
 130  
 121  
 gln val pro glu pro val thr ser cys glu ala tyr glu glu asp arg glu thr phe met asn lys phe ile tyr glu ile ala arg arg  
 140  
 CAA GTT CCA GAA CCT GTC ACA AGC TGT GAA GCA TAT GAA GAC AGG GAG ACA TTC ATG AAC AAA TTC ATT TAT GAG ATA GCA AGA AGG (551)  
 150  
 160  
 151  
 his pro phe leu tyr ala pro thr ile leu trp ala ala arg tyr asp lys ile pro ser cys cys lys ala glu asn ala val  
 170  
 CAT CCC TTC CTG TAT GCA CCT ACA ATT CTT TGG GCT GCT CGC TAT GAC AAA ATA ATT CCA TCT TGC TGC AAA GCT GAA AAT GCA GTT (641)  
 180  
 190  
 181  
 glu cys phe gln thr lys ala ala thr val thr lys glu leu arg glu ser ser leu leu asn gln his ala cys ala val met lys asn  
 200  
 GAA TGC TTC CAA ACA AAG GCA GCA ACA GTT ACA AAA GAA TTA AGA GAA AGC AGC TTT TTA AAT CAA CAT GCA TGT GCA GTA ATG AAA AAT (731)  
 210

Fig. 4-1

211 phe gly thr arg thr phe gln ala ile thr val thr lys leu ser gln lys phe thr lys val asn phe thr glu ile gln lys leu val 240  
 TTT GGG ACC CGA ACT TTC CAA GCC ATA ACT GTT ACT GAA CTG AGT CAG AAG TTT ACC AAG GTT AAT TTT ACT GAA ATC CAG AAA CTA GTC (821)  
 220  
 241 leu asp val ala his val his glu his cys cys arg gly asp val leu asp cys leu gln asp gly glu lys ile met ser tyr ile cys 270  
 CTG GAT GTG GCC CAT GTA CAT GAG CAC TGT TGC AGA GGA GAT GTG CTG GAT TGT CTG CAG GAT GGG GAA AAA ATC ATG TCC TAC ATA TGT (911)  
 250  
 271 ser gln gln asp thr leu ser asn lys ile thr glu cys cys lys leu thr thr leu glu arg gly gln cys ile ile his ala glu asn 300  
 TCT CAA CAA GAC ACT CTG TCA AAC AAA ATA ACA GAA TGC TGC AAA CTG ACC ACC CTG GAA CGT GGT CAA TGT ATA ATT CAT GCA GAA AAT(1001)  
 280  
 301 glu lys pro glu gly leu ser pro asn leu ser pro asn leu gly asp arg phe asn gln phe ser ser gly glu lys asn ile 330  
 GAT GAA AAA CCT GAA GGT CTA TCT CCA AAT CTA AAC AGG TTT TTA GGA GAT AGA GAT TTT AAC CAA TTT TCT TCA GGG GAA AAA AAT ATC(1091)  
 310  
 331 leu ala ser phe val his glu tyr ser arg arg his pro gln leu ala val ser val ile leu arg val ala lys gly tyr gln glu 360  
 TTC TTG GCA AGT TTT GTT CAT GAA TAT TCA AGA AGA CAT CCT CAG CTT GCT GTC TCA GTA ATT CTA AGA GTT GCT AAA GGA TAC CAG GAG(1181)  
 340  
 361 leu glu lys cys phe gln thr glu asn pro leu glu cys gln asp lys gly glu leu gln lys tyr ile gln glu ser gln 390  
 TTA TTG GAG AAG TGT TTC CAG ACT GAA AAC CCT CTT GAA TGC CAA GAT AAA GGA GAA GAA TTA CAG AAA TAC ATC CAG GAG AGC CAA(1271)  
 370  
 391 ala leu ala lys arg ser cys gly leu phe gln lys leu gly glu tyr leu gln asn ala phe leu val ala tyr thr lys lys ala 420  
 GCA TTG GCA AAG CGA AGC TGC GGC CTC TTC CAG AAA CTA GGA GAA TAT TAC TTA CAA AAT GCG TTT CTC GTC ACA AAG AAA GCC(1361)  
 400  
 421 pro gln leu thr ser ser glu leu met ala ile thr arg lys met ala ala thr ala ala thr cys cys gln leu ser glu asp lys leu 450  
 CCC CAG CTG ACC TCG TCG GAG CTG ATG GCC ATC ACC AGA AAA ATG GCA GCC ACA GCA GCC ACT TGT TGC CAA CTC AGT GAG GAC AAA CTA(1451)  
 430

Fig. 4-2



451 leu ala cys gly glu gly ala ala asp ile ile ile gly his leu cys ile arg his glu met thr pro val asn pro gly val gly gln 480  
 TTG GCC TGT GGC GAG GGA GCG GCT GAC ATT ATT ATC GGA CAC TTA TGT ATC AGA CAT GAA ATG ACT CCA GTA AAC CCT GGT GTT GGC CAG (1541)  
 460  
 481 cys cys thr ser ser tyr ala asn arg arg pro cys phe ser leu val val asp glu thr tyr val pro pro ala phe ser asp asp 510  
 TGC TGC ACT TCT TCA TAT GCC AAC AGG AGG CCA TGC TTC AGC AGC TTTG GTG GAT GAA ACA TAT GTC CCT CCT GCA TTC TCT GAT GAC (1631)  
 490  
 511 lys phe ile phe his lys asp leu cys gln ala gln gly val ala leu gln thr met lys gln glu phe leu ile asn leu val lys gln 540  
 AAG TTC ATT TTC CAT AAG GAT AAG GAT CTG TGC CAA GCT CAG GGT GTA GCG CTG CAA ACG ATG AAG CAA GAG TTT CTC ATT AAC CTT GTG AAG CAA (1721)  
 520  
 541 lys pro gln ile thr glu glu glu leu glu ala val ile ala asp phe ser gly leu leu glu lys cys cys gln gly gln glu gln glu 570  
 AAG CCA CAA ATA ACA GAG GAA CAA CTT GAG GCT GTC ATT GCA GAT TTC TCA GGC CTG TGC AAA TGC TGC CAA GGC CAG GAA CAG GAA (1811)  
 530  
 550  
 571 val cys phe ala glu glu gly gln lys leu ile ser lys thr arg ala ala leu gly val ter 590  
 GTC TGC TTT GCT GAA GAG GGA CAA AAA CTG ATT TCA AAA ACT CGT GCT GCT TCG TTTG GGA GTT TAA ATTACTTCAGGGGAAGAGACAAAACGAGTCT (1908)  
 TTCATTCGGGTGAACCTTTCTCTTTAACTGATTTAAACACTTTTGTGAAATTAATGAAATGATAAAGACTTTTATGTGAGATTTCCCTTATCACAGAAATAAAAATATCTCCAAA (2027)

Fig. 4-3




Fig. 5

## RECOMBINANT HUMAN ALPHA-FETOPROTEIN AS AN IMMUNOSUPPRESSIVE AGENT

This application is a continuation of, and claims priority from, U.S. application Ser. No. 08/377,309, filed Jan. 24, 1995, U.S. Pat. No. 5,965,528.

### BACKGROUND OF THE INVENTION

This invention relates to methods for treating autoimmune diseases.

Responses of the immune system initiate the destruction and elimination of invading organisms and toxic molecules produced by them. Because these immune reactions are destructive, it is essential that they be made in response only to molecules that are foreign to the host and not to those of the host itself. The ability to distinguish foreign molecules from self molecules is a fundamental feature of the immune system. Occasionally the immune system fails to make this distinction and reacts destructively against the host's own molecules; such autoimmune diseases can be fatal. Thus, tolerance to self antigens breaks down, causing the components of the immune system such as T or B cells (or both) to react against their own tissue antigens. Multiple sclerosis, rheumatoid arthritis, myasthenia gravis, insulin-dependent diabetes mellitus, and systemic lupus erythematosus are a few examples of such autoimmune diseases.

### SUMMARY OF THE INVENTION

I have discovered that unglycosylated recombinant human alpha-fetoprotein made in a prokaryote (e.g., *E. coli*) is useful for inhibiting autoreactive immune cells derived from a mammal. Accordingly, the invention features a method of inhibiting autoreactive immune cell proliferation in a mammal (e.g., a human patient), involving administering to the mammal a therapeutically effective amount of recombinant human alpha-fetoprotein or an immune cell anti-proliferative fragment or analog thereof. Preferably, such immune cells include T cells or B cells; and the recombinant human alpha-fetoprotein used in such methods is produced in a prokaryotic cell (e.g., *E. coli*) and is unglycosylated.

In another aspect, the invention features a method of treating an autoimmune disease in a mammal (e.g., a human patient), involving administering to the mammal a therapeutically effective amount of recombinant human alpha-fetoprotein or an immune cell anti-proliferative fragment or analog thereof. Such an autoimmune disease is multiple sclerosis; is rheumatoid arthritis; is myasthenia gravis; is insulin-dependent diabetes mellitus; or is systemic lupus erythematosus. In yet other preferred embodiments the autoimmune disease is acquired immune deficiency syndrome or is a rejection of a transplanted organ, tissue or cell. Preferably, the recombinant human alpha-fetoprotein used in such methods is produced in a prokaryotic cell (e.g., *E. coli*) and is unglycosylated. In other preferred embodiments, such methods further involve administering to the mammal an immunosuppressive agent in an effective dose which is lower than the standard dose when the immunosuppressive agent is used by itself. Preferably, such an immunosuppressive agent is cyclosporine; is a steroid; is azathioprine; is FK-506; or is 15-deoxyspergualin. In yet another preferred embodiment, such a method involves administering to the mammal a tolerizing agent. Preferably, the recombinant human alpha-fetoprotein used in such methods is produced in a prokaryotic cell (e.g., *E. coli*) and is unglycosylated.

By "immune cell anti-proliferative" is meant capable of inhibiting the growth of an undesirable immune cell e.g., an autoreactive T cell as measured using the assays described herein).

By "therapeutically effective amount" is meant a dose of unglycosylated recombinant human alpha-fetoprotein (or a fragment or analog thereof) capable of inhibiting autoreactive immune cell proliferation.

By "recombinant human alpha-fetoprotein" is meant a polypeptide having substantially the same amino acid sequence as the protein encoded by the human alpha-fetoprotein gene as described by Morinaga et al., *Proc. Natl. Acad. Sci., USA* 80: 4604 (1983). The method of producing recombinant human alpha-fetoprotein in a prokaryotic cell is described in U.S. Ser. No. 08/133,773 issuing as U.S. Pat. No. 5,384,250.

According to the invention, administration of recombinant human alpha-fetoprotein ("rHuAFP") (or a fragment or analog thereof) can be an effective means of preventing or treating or ameliorating autoimmune diseases in a mammal. To illustrate this, I have shown that recombinant HuAFP produced in a prokaryotic expression system is effective in suppressing T cell proliferation in response to self antigens, despite the fact that such rHuAFP is not modified in the same fashion as naturally occurring HuAFP. The use of natural HuAFP has heretofore been limited by its unavailability, natural HuAFP is obtained by laborious purification from limited supplies of umbilical cords and umbilical cord serum. Because biologically rHuAPP can now be prepared in large quantities using the techniques of recombinant DNA, the use of rHuAFP for treating autoimmune diseases is now possible. The use of rHuAFP is especially advantageous since there are no known adverse side effects related to human alpha-fetoprotein and it is believed that relatively high doses can be safely administered.

Other features and advantages of the invention will be apparent from the following detailed description, and from the claims.

### DETAILED DESCRIPTION

The drawings will first be described.

#### Drawings

FIG. 1 is a series of panels (FIGS. 1A-1D) showing the purity and biochemical characteristics of rHuAFP using polyacrylamide gel electrophoresis and column chromatography. FIGS. 1A is a 10% non-denaturing alkaline polyacrylamide gel showing the purity of rHuAFP produced in *E. coli*. Mouse amniotic fluid proteins (transferrin, AFP and albumin) are shown in lane 1, natural HuAFP and rHuAFP are shown in lane 2 and lane 3, respectively. FIG. 1B is a 10% sodium dodecyl sulfate-polyacrylamide gel showing the purity of rHuAFP produced in *E. coli*. Molecular weight markers are shown in lane 1, natural HuAFP and rHuAFP are shown in lane 2 and lane 3, respectively. FIG. 1C is a series of FPLC chromatograms of natural HuAFP and rHuAFP eluted on a MonoQ anion exchange column. The superimposed chromatograms identify natural HuAFP (Chromatogram 1) and rHuAFP (Chromatogram 2). FIG. 1D is a series of HPLC chromatograms obtained by passing 50  $\mu$ g of natural and rHuAFP by passing through a reverse phase Delta Pak C18 column (Waters) and eluting with a gradient of 0-100% acetonitrile in 0.1% TFA. The superimposed chromatograms identify natural HuAFP (Chromatogram 1) and rHuAFP (Chromatogram 2).

FIG. 2 is a series of graphs showing the inhibitory effects of the rHuAFP on kinetics of T cell activation (FIG. 2A) and

the dose-response relationship of rHuAFP on autoproli-  
 erating T cells (FIG. 2B). FIG. 2A is a graph showing  
 proliferative responses over a 4 day time course of cells  
 cultured in the absence ( $\nabla$ ) and in the presence of 100  $\mu\text{g}/\text{ml}$   
 ( $\blacktriangle$ ) rHuAFP. ( $\bullet$ ) denotes the background proliferation of  
 the responder cell population cultured separately. Recombi-  
 nant HuAFP-mediated suppression on the AMLR over the  
 time course was significant ( $p < 0.01$ ). FIG. 2B is a graph  
 showing the inhibition of autoproli-  
 erating T cells at 144 hours with amounts of rHuAFP ranging from 6–100  $\mu\text{g}/\text{ml}$   
 ( $\blacktriangle$ ). ( $\nabla$ ) denotes the control response of the reaction in the  
 absence of protein. Inhibition of autoreactive T cells by  
 rHuAFP in the range of 12.5–100  $\mu\text{g}/\text{ml}$  is significant  
 ( $p < 0.005$ ).

FIG. 3 is a bar graph showing that monoclonal anti-  
 natural HuAFP antibodies block immunosuppression of the  
 AMLR by rHuAFP. Immunosuppression by rHuAFP was  
 significant ( $p < 0.002$ ) and blocking of rAFP-mediated immu-  
 nosuppression by the AMLR by monoclonal anti-natural  
 HuAFP antibodies was also significant ( $p < 0.03$ ).

FIG. 4 is the nucleotide sequence (SEQ ID NO: 1) and  
 deduced amino acid sequence (SEQ ID NO: 2) of the cDNA  
 encoding human alpha-fetoprotein.

FIG. 5 is the 10% SDS-PAGE analysis of rHuAFP Frag-  
 ment I or alpha-fetoprotein (Lane A, MW marker; Lane B,  
 natural human alpha-fetoprotein, Lane C, unpurified  
 rHuAFP and rHuAFP Fragment I, Lane D, purified rHuAFP  
 Fragment I, and Lane E, purified full-length rHuAFP).

#### PRODUCTION OF RECOMBINANT HUMAN ALPHA-FETOPROTEIN

As summarized above, the invention includes therapies  
 for the prevention and treatment of autoimmune diseases  
 involving recombinant human alpha-fetoprotein  
 ("rHuAFP") or fragments or analogs thereof. Methods for  
 producing such rHuAFP in a prokaryotic cell are described  
 in U.S. Ser. No. 08/133,773 and in U.S. Pat. No. 5,384,250,  
 issued Jan. 24, 1995.

##### Fragments and Analogs

The invention includes biologically active fragments of  
 rHuAFP. A biologically active fragment of rHuAFP is one  
 that possesses at least one of the following activities: (a)  
 directs a specific interaction with a target cell, e.g., binds to  
 a cell expressing a receptor which is recognized by rHuAFP  
 (e.g., the membrane of an autoreactive immune cell); or (b)  
 halts, reduces, or inhibits the growth of an autoreactive  
 immune cell (e.g., binds to a cell surface receptor and  
 imparts an anti-proliferative signal); or (c) blocks, inhibits,  
 or prevents an immunopathologic antibody reaction. The  
 ability of rHuAFP fragments or analogs to bind to a receptor  
 which is recognized by rHuAFP can be tested using any  
 standard binding assay known in the art. Methods for  
 assaying the biological activity of rHuAFP fragments and  
 analogs are also known in the art, e.g., those described  
 herein. Accordingly, a rHuAFP fragment, like the full-length  
 rHuAFP molecule, can be used to inhibit autoreactive immune  
 cell proliferation.

In general, fragments of rHuAFP are produced according  
 to the techniques of polypeptide expression and purification  
 described in U.S. Ser. No. 08/133,773 (U.S. Pat. No. 5,384,  
 250). For example, suitable fragments of rHuAFP can be  
 produced by transformation of a suitable host bacterial cell  
 with part of an HuAFP-encoding cDNA fragment (e.g., the  
 cDNA described above) in a suitable expression vehicle.  
 Alternatively, such fragments can be generated by standard  
 techniques of PCR and cloned into the expression vectors  
 (supra). Accordingly, once a fragment of rHuAFP is

expressed, it may be isolated by various chromatographic  
 and/or immunological methods known in the art. Lysis and  
 fractionation of rHuAFP-containing cells prior to affinity  
 chromatography may be performed by standard methods.  
 Once isolated, the recombinant protein can, if desired, be  
 further purified, e.g., by high performance liquid chroma-  
 tography (see, e.g., Fisher, *Laboratory Techniques In Bio-  
 chemistry And Molecular Biology*, Work and Burdon, eds.,  
 Elsevier, 1980).

A rHuAFP fragment may also be expressed as a fusion  
 protein with maltose binding protein produced in *E. coli*.  
 Using the maltose binding protein fusion and purification  
 system (New England Biolabs), the cloned human cDNA  
 sequence can be inserted downstream and in frame of the  
 gene encoding maltose binding protein (malE), and the malE  
 fusion protein can then be overexpressed. In the absence of  
 convenient restriction sites in the human cDNA sequence,  
 PCR can be used to introduce restriction sites compatible  
 with the vector at the 5' and 3' end of the cDNA fragment to  
 facilitate insertion of the cDNA fragment into the vector.

Following expression of the fusion protein, it can be  
 purified by affinity chromatography. For example, the fusion  
 protein can be purified by virtue of the ability of the maltose  
 binding protein portion of the fusion protein to bind to  
 amylose immobilized on a column.

To facilitate protein purification, the pMalE plasmid con-  
 tains a factor Xa cleavage site upstream of the site into  
 which the cDNA is inserted into the vector. Thus, the fusion  
 protein purified as described above can then be cleaved with  
 factor Xa to separate the maltose binding protein from a  
 fragment of the recombinant human cDNA gene product.  
 The cleavage products can be subjected to further chroma-  
 tography to purify rHuAFP from the maltose binding pro-  
 tein. Alternatively, a fragment of rHuAFP may be expressed  
 as a fusion protein containing a polyhistidine tag can be  
 produced. Such an alpha-fetoprotein fusion protein may then  
 be isolated by binding of the polyhistidine tag to an affinity  
 column having a nickel moiety which binds the polyhisti-  
 dine region with high affinity. The fusion protein may then  
 be eluted by shifting the pH within the affinity column. The  
 rHuAFP can be released from the polyhistidine sequences  
 present in the resultant fusion protein by cleavage of the  
 fusion protein with specific proteases.

Recombinant HuAFP fragment expression products (e.g.,  
 produced by any of the prokaryotic systems described in  
 U.S. Ser. No. 08/133,773) may be assayed by immunologi-  
 cal procedures, such as Western blot, immunoprecipitation  
 analysis of recombinant cell extracts, or immunofluores-  
 cence (using, e.g., the methods described in Ausubel et al.,  
*Current Protocols In Molecular Biology*, Greene Publishing  
 Associates and Wiley Interscience (John Wiley & Sons),  
 New York, 1994).

Once a fragment of rHuAFP is expressed, it is isolated  
 using the methods described supra. Once isolated, the frag-  
 ment of rHuAFP can, if desired, be further purified by using  
 the techniques described supra. Fragments can also be  
 produced by chemical synthesis (e.g., by the methods  
 described in *Solid Phase Peptide Synthesis*, 2nd ed., 1984,  
 The Pierce Chemical Co., Rockford, Ill.). The ability of a  
 candidate rHuAFP fragment to exhibit a biological activity  
 of alpha-fetoprotein is assessed by methods known to those  
 skilled in the art (e.g., those described herein).

The purified recombinant gene product or fragment  
 thereof can then be used to raise polyclonal or monoclonal  
 antibodies against the human recombinant alpha-fetoprotein  
 using well-known methods (see Coligan et al., eds., *Current  
 Protocols in Immunology*, 1992, Greene Publishing Associ-

ates and Wiley-Interscience). To generate monoclonal antibodies, a mouse can be immunized with the recombinant protein, and antibody-secreting B cells isolated and immortalized with a non-secretory myeloma cell fusion partner. Hybridomas are then screened for production of recombinant human alpha-fetoprotein (or a fragment or analog thereof)-specific antibodies and cloned to obtain a homogeneous cell population which produces monoclonal antibodies.

As used herein, the term "fragment," as applied to a rHuAFP polypeptide, is preferably at least 20 contiguous amino acids, preferably at least 50 contiguous amino acids, more preferably at least 100 contiguous amino acids, and most preferably at least 200 to 400 or more contiguous amino acids in length. Fragments of rHuAFP molecules can be generated by methods known to those skilled in the art, e.g., proteolytic cleavage or expression of recombinant peptides, or may result from normal protein processing (e.g., removal of amino acids from nascent polypeptide that are not required for biological activity).

Recombinant HuAFP fragments of interest include, but are not limited to, Domain I (amino acids 1 (Thr)-197 (Ser), see FIG. 4, SEQ ID NO: 3), Domain II (amino acids 198 (Ser)-389 (Ser), see FIG. 4, SEQ ID NO: 4), Domain III (amino acids 390 (Gln)-590 (Val), see FIG. 4, SEQ ID NO: 5), Domain I+II (amino acids 1 (Thr)-389 (Ser), see FIG. 4, SEQ ID NO: 6), Domain II+III (amino acids 198 (Ser)-590 (Val), see FIG. 4, SEQ ID NO: 7), and rHuAFP Fragment I (amino acids 266 (Met)-590 (Val), see FIG. 4, SEQ ID NO: 8). Activity of a fragment is evaluated experimentally using conventional techniques and assays, e.g., the assays described herein.

The invention further includes analogs of full-length rHuAFP or fragments thereof. Analogs can differ from rHuAFP by amino acid sequence differences, or by modifications (e.g., post-translational modifications) which do not affect sequence, or by both. Analogs of the invention will generally exhibit at least 80%, more preferably 85%, and most preferably 90% or even 99% amino acid identity with all or part of a rHuAFP amino acid sequence. Modifications (which do not normally alter primary sequence) include in vivo, or in vitro chemical derivatization of polypeptides, e.g., acetylation, or carboxylation; such modifications may occur during polypeptide synthesis or processing or following treatment with isolated modifying enzymes. Analogs can also differ from the naturally occurring rHuAFP by alterations in primary sequence, for example, substitution of one amino acid for another with similar characteristics (e.g., valine for glycine, arginine for lysine, etc.) or by one or more non-conservative amino acid substitutions, deletions, or insertions which do not abolish the polypeptide's biological activity. These include genetic variants, both natural and induced (for example, resulting from random mutagenesis by irradiation or exposure to ethanemethylsulfate or by site-specific mutagenesis as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd ed., Cold Spring Harbor Press, 1989, or Ausubel et al., supra). Also included are cyclized peptide molecules and analogs which contain residues other than L-amino acids, e.g., D-amino acids or non-naturally occurring or synthetic amino acids, e.g.,  $\beta$  or  $\gamma$  amino acids, or L-amino acids with non-natural side chains (see e.g., Noren et al., *Science* 244:182, 1989).

DomI25  
DomI3  
DomII5

5'-AAAAAAGGTACCACACTGCATAGAAATGAA-3' (SEQ ID NO:9)  
5'-AAAAAAGGATCCTTAGCTTTCTCTTAATTCCTT-3' (SEQ ID NO:10)  
5'-AAAAAATCGATATGAGCTTGTAAATCAACAT-3' (SEQ ID NO:11)

Methods for site-specific incorporation of non-natural amino acids into the protein backbone of proteins is described, e.g., in Ellman et al., *Science* 255:197, 1992. Also included are chemically synthesized polypeptides or peptides with modified peptide bonds (e.g., non-peptide bonds as described in U.S. Pat. No. 4,897,445 and U.S. Pat. No. 5,059,653) or modified side chains to obtain the desired pharmaceutical properties as described herein. Useful mutants and analogs are identified using conventional methods, e.g., those described herein.

The cloning, expression, isolation and characterization of exemplary rHuAFP fragments now follows. These examples are provided to illustrate, not limit, the invention.

## EXPERIMENTAL

### MATERIALS AND METHODS

#### Polymerase Chain Reaction (PCR) rHuAFP Fragments

Plasmid constructs encoding fragments of human alpha-fetoprotein were prepared using polymerase chain reaction (PCR) techniques known to those skilled in the art of molecular biology, using oligonucleotide primers designed to amplify specific portions of the human alpha-fetoprotein gene (see e.g., PCR Technology, H. A. Erlich, ed., Stockton Press, New York, 1989; *PCR Protocols: A Guide to Methods and Applications*, M. A. Innis, David H. Gelfand, John J. Sninsky, and Thomas J. White, eds., Academic Press, Inc., New York, 1990, and Ausubel et al., supra).

The following six rHuAFP fragments were prepared to evaluate their biological activity (e.g., according to the methods disclosed herein):

|                   |                                  |                       |
|-------------------|----------------------------------|-----------------------|
| Domain I          | Amino acids 1 (Thr)-197 (Ser),   | (FIG. 4, SEQ ID NO:3) |
| Domain II         | Amino acids 198 (Ser)-389 (Ser), | (FIG. 4, SEQ ID NO:4) |
| Domain III        | Amino acids 390 (Gln)-590 (Val), | (FIG. 4, SEQ ID NO:5) |
| Domain I + II     | Amino acids 1 (Thr)-389 (Ser),   | (FIG. 4, SEQ ID NO:6) |
| Domain II + III   | Amino acids 198 (Ser)-590 (Val), | (FIG. 4, SEQ ID NO:7) |
| rHuAFP Fragment I | Amino acids 266 (Met)-590 (Val), | (FIG. 4, SEQ ID NO:8) |

Amino acid sequences were deduced from those for human alpha-fetoprotein (1 (Thr)-590 (Val); SEQ ID NO: 2) shown in FIG. 4. Fragments of rHuAFP designated Domain I, Domain II, Domain III, Domain I+II, Domain II+III and rHuAFP Fragment I were synthesized using standard PCR reaction conditions in 100  $\mu$ L reactions containing 34  $\mu$ L H<sub>2</sub>O, 10  $\mu$ L 1X reaction buffer, 20  $\mu$ L 1 mM dNTP, 2  $\mu$ L DNA template (HuAFP cloned in pI18), appropriate 5' and 3' oligonucleotide primers (10  $\mu$ L 10 pmol/ $\mu$ L 5' primer, 10  $\mu$ L 10 pmol/ $\mu$ L 3' primer), 1  $\mu$ L glycerol, 10  $\mu$ L DMSO, and 1  $\mu$ L Pfu polymerase (Stratagene, LaJolla, Calif.). Primers used for PCR amplifications were:

## -continued

DomII3 5'-AAAAAAGGATCCTTAGCTCTCTGGATGTATTT-3' (SEQ ID NO:12)  
 DomIII5 5'-AAAAAATCGATATGCAAGCATTGGCAAAGCGA-3' (SEQ ID NO:13)  
 DomIII3 5'-AAAAAAGGATCCTTAAACTCCCAAAGCAGCACG-3' (SEQ ID NO:14)  
 5'rHuAFP Fragment I 5'-AAAAAATCGATATGTCTACATATGTTCTCAA-3' (SEQ ID NO:15)

Accordingly, primer pairs DomI25 and DomI3, DomII5 and DomII3, DomIII5 and DomIII3, 5'rHuAFP Fragment I and DomIII3, DomI25 and DomII3, and DomII5 and DomIII3 were used to isolate cDNA sequences of Domain I, Domain II, Domain III, rHuAFP Fragment I, Domain I+II, and Domain II+III, respectively, of rHuAFP. Annealing, extension, and denaturation temperatures were 50° C., 72° C., and 94° C., respectively, for 30 cycles. PCR products were purified according to standard methods. Purified PCR products encoding Domain I and Domain I+II were digested individually with KpnI and BamHI and cloned separately into KpnI/BamHI-treated pTrp4. Purified PCR products encoding Domain II, Domain III, Domain II+III, and rHuAFP Fragment I were digested individually with Bsp106I and BamHI and were cloned separately into Bsp106I/BamHI-treated pTrp4. Each plasmid construct was subsequently transformed into competent *E. coli* cells. Since the expression product will begin with the amino acid sequence encoded by the translation start signal methionine, it is expected that such signal will be removed, or in any event, not affect the bioactivity of the ultimate expression product.

## Autologous Mixed Lymphocyte Reactions (AMLR)

AMLR assays were performed as described below.

## RESULTS

## Expression and Purification

*E. coli* containing the expression plasmid encoding rHuAFP Fragment I was cultured and purified. FIG. 5 (lane D) shows the SDS-PAGE profile of the purified rHuAFP Fragment I. N-terminal amino acid sequence analysis showed that rHuAFP Fragment I possessed the amino acid sequence Ser<sub>267</sub>-Tyr-Ile-Cys-Ser-Gln-Gln-Asp-Thr<sub>275</sub> (SEQ ID NO: 16) which corresponds to the expected N-terminal amino acid sequence of rHuAFP Fragment I (see FIG. 4, SEQ ID NO: 2) where the initiating methionine is cleaved intracellularly.

## Inhibition of the Autologous Mixed Lymphocyte Reactions (AMLR)

The immunosuppressive activity of 100 µg/ml rHuAFP Fragment I was assessed by its ability to suppress human autologous mixed lymphocyte reactions (AMLR). As shown in Table I, rHuAFP Fragment I inhibited the proliferative response of autoreactive lymphocytes stimulated by autologous non-T cells at 144 hours.

TABLE I

| Reaction Setup                       | Thymidine Incorporation (CPM) |
|--------------------------------------|-------------------------------|
| T Cells                              | 7118 ± 964                    |
| AMLR                                 | 83103 ± 6480                  |
| AMLR + rHuAFP Fragment I (100 µg/ml) | 29692 ± 2963                  |

## Recombinant HuAFP As An Immunosuppressive Agent

Immunosuppressive attributes of rHuAFP (or a fragment or analog thereof) is evaluated by any standard assay for

analysis of immunoregulatory activity in vivo or in vitro. As discussed infra, the art provides a number of animal systems for in vivo testing of immunosuppressive characteristics of rHuAFP (or a fragment or analog thereof) on an autoimmune disease, e.g., the nonobese diabetic (NOD) mouse. Furthermore, a wide variety of in vitro systems are also available for testing immunosuppressive aspects of rHuAFP e.g., one such in vitro assay evaluates the inhibition of autoantigen-induced proliferation of T cells in an autologous mixed lymphocyte reaction (AMLR).

The following examples demonstrate that unglycosylated rHuAFP inhibits T cell autoprolieration in response to self antigens. These examples are provided to illustrate, not limit, the invention.

## EXPERIMENTAL

## MATERIALS AND METHODS

## Gel Electrophoresis, Immunoblotting and Purification

The purity and characterization of rHuAFP was evaluated by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) and nondenaturing alkaline PAGE (APAGE) according to standard methods. Gels were subsequently analyzed either by staining with Coomassie brilliant blue or by transferring electrophoretically separated polypeptides to Immobilon PVDF membranes (Millipore, Mississauga, ON) for immunoblotting analysis. Recombinant HuAFP-monospecific rabbit anti-natural HuAFP polyclonal antibody complexes were identified by alkaline-phosphatase-conjugated goat anti-rabbit IgG and the immunoreactive bands were detected with the BCIP/NBT color development solution (Bio-Rad-Laboratories, Mississauga, ON) according to the manufacturer's instructions.

Column chromatography was performed according to standard methods.

## Autologous Mixed Lymphocyte Reactions (AMLR)

Isolation of human peripheral blood mononuclear cells (PBMC), their fractionation into non-T cell populations, and the AMLR, were performed according to standard procedures. Responder T cells were isolated by passing  $1.5 \times 10^8$  PMBC over a commercial Ig-anti-Ig affinity column (Biotek Laboratories) and  $2 \times 10^5$  responder cells were subsequently cultured with  $2 \times 10^5$  autologous  $^{137}\text{Cs}$ -irradiated (2500 rads) non-T stimulator cells from a single donor. The medium employed consisted of RPMI-1640 supplemented with 20 mM HEPES (Gibco),  $5 \times 10^{-5}$  M 2-mercaptoethanol (BDH, Montreal, QC), 4 mM L-glutamine (Gibco), 100 U/ml penicillin (Gibco) and 100 µg/ml streptomycin sulfate, with the addition of 10% fresh human serum autologous to the responder T cell donor for the AMLR. Varying concentrations of purified rHuAFP, human serum albumin (HSA), anti-HuAFP monoclonal antibodies clone #164 (125 µg/ml final concentration in culture) (Leinco Technologies, St. Louis, Mo.) were added at the initiation of cultures. AMLR cultures were incubated for 4 to 7 days, at 37° C. in 95% air and 5% CO<sub>2</sub>. At the indicated intervals, DNA synthesis was

assayed by a 6 hour pulse with 1  $\mu$ Ci of  $^3$ H-thymidine (specific activity 56 to 80 Ci/mmmole, ICN). The cultures were harvested on a multiple sample harvester (Skatron, Sterling, Va.), and the incorporation of  $^3$ H-TdR was measured in a Packard 2500 TR liquid scintillation counter. Results are expressed as mean cpm  $\pm$  the standard error of the mean of triplicate or quadruplicate cultures.

## RESULTS

### Expression and Purification

Purity of isolated rHuAFP expressed in *E. coli* was verified as a single band on Coomassie stained APAGE and SDS-PAGE are shown in FIGS. 1A-1B, respectively. Soluble monomeric rHuAFP derived from *E. coli* was obtained by eluting a protein fraction containing rHuAFP employing Q-sepharose chromatography. Approximately 1 mg of pure rHuAFP per liter of bacterial culture was recovered as a single homogeneous peak by FPLC Mono-Q anion exchange with 220-230 mM NaCl and migrated at approximately 65 kD on SDS-PAGE (FIG. 1B). Recombinant HuAFP exhibits a lower molecular weight on SDS-PAGE than natural HuAFP, since prokaryotic expression systems lack the enzymatic machinery required for glycosylation of proteins. Rechromatographed samples of pure rHuAFP on FPLC and HPLC yielded a single peak as shown in FIG. 1C and FIG. 1D, confirming the purity of the rHuAFP preparation. In addition, N-terminal sequencing data correspond to the expected amino acid sequence at the N-terminus of rHuAFP.

### Inhibition of the Autologous Mixed Lymphocyte Reactions (AMLR)

The immunosuppressive activity of rHuAFP was assessed by its ability to suppress human autologous mixed lymphocyte reactions (AMLR). As shown in FIG. 2A, rHuAFP inhibited the proliferative response of autoreactive lymphocytes stimulated by autologous non-T cells, throughout the 4 to 7 day time course measuring autoproliiferation. Results from dose-response studies performed at the peak of T cell autoproliiferation, as shown in FIG. 2B, demonstrate that the addition of rHuAFP at the initiation of cultures suppressed the AMLR in a dose-dependent manner. Furthermore, parallel viability-studies established that the inhibitory activity of rHuAFP on human autoreactive T cells was not due to non-specific cytotoxic effects.

To further substantiate that rHuAFP was the agent responsible for the inhibition of autoproliiferating T cells, blocking of rHuAFP-mediated suppression of the AMLR was performed using commercial murine anti-human AFP monoclonal antibodies (MAb). As illustrated in FIG. 3, suppression of proliferating autoreactive T cells by 100  $\mu$ g/ml of rHuAFP was completely blocked by anti-HuAFP NAb. The addition of 100  $\mu$ g/ml of HSA did not diminish the AMLR response and the presence of MAb alone in the reaction culture was without any effect.

Recombinant polypeptides produced in prokaryotic expression systems are at risk for contamination with host cell lipopolysaccharide (LPS) during their isolation from bacteria. It has been demonstrated that small amounts of LPS can antagonize the biological activities of cytokines, thereby impairing the immune responsiveness of macrophages. Accordingly, the effect of endotoxin on various rHuAFP preparations was evaluated by performing AMLR experiments with recombinant protein depleted of endotoxin by passage over Detoxi-gel (Pierce) versus that of rHuAFP which was untreated. Results of these experiments showed

that both preparations had equivalent levels of immunosuppressive activity.

As shown in FIG. 2A and FIG. 2B, the results of this study also demonstrate that rHuAFP suppresses the proliferation of autoreaction T cells with a potency equivalent to glycosylated nHuAFP by eliciting inhibitory effects on autoproliiferating T cells throughout the in vitro reactions, with highly significant inhibition being achieved with rHuAFP concentrations ranging from 5  $\mu$ g/ml to 100  $\mu$ g/ml.

### Autoimmune Disease

As is discussed above, autoimmune diseases are characterized by a loss of tolerance to self antigens, causing cells of the immune systems, e.g., T or B cells (or both), to react against self tissue antigens. Autoimmune diseases may involve any organ system, although some are affected more commonly than others. Examples of tissues affected by autoimmune conditions include: the white matter of the brain and spinal cord in multiple sclerosis; the lining of the joints in rheumatoid arthritis; and the insulin secreting  $\beta$  islet cells of the pancreas in insulin-dependent diabetes mellitus. Other forms of autoimmune disease destroy the connections between nerve and muscle in myasthenia gravis or destroy the kidneys and other organs-in systemic lupus erythematosus. Examples of other autoimmune diseases include, without limitation, Addison's disease, Crohn's disease, Graves' disease, psoriasis, scleroderma, and ulcerative colitis.

The art provides a wide variety of experimental animal systems, transgenic and non-transgenic, for testing therapies for human illness involving autoimmune diseases (see e.g., Paul, W. E., *Fundamental Immunology*, 2nd ed., Raven Press, New York, 1989; and Kandel et al. *Principles of Neural Science*, 3rd ed., Appleton and Lange, Norwalk, Conn., 1991; and *Current Protocols In Immunology*, Coligan, J. E., Kruisbeek, A. M., Margulies, D. H., Shevach, E. M., and Strober, eds., Green Publishing Associates (John Wiley & Sons), New York, 1992). Based on the above-described experimental results showing immunosuppressive activity of unglycosylated rHuAFP, it is reasonable to believe that other autoimmune diseases can be treated by administration of such rHuAFP (or fragment or analog thereof) produced in a prokaryotic system. Accordingly, the invention provides the use of rHuAFP (or a fragment or analog thereof) for treatment (i.e., prevention or suppression or amelioration or promotion of remission) of any autoimmune disease.

There now follow examples of animal systems useful for evaluating the efficacy of recombinant human alpha-fetoprotein or an immune cell anti-proliiferative fragment or analog thereof in treating autoimmune diseases. These examples are provided for the purpose of illustrating, not limiting, the invention.

### Multiple Sclerosis

Multiple sclerosis (MS) is a demyelinating disease involving scattered areas of the white matter of the central nervous system. In MS, myelin basic protein and proteolipid protein are the major targets of an autoimmune response involving T lymphocytes, among other immune system components. Loss of the myelin sheath of nerve cells (demyelination) occurs, resulting in neurological symptoms that culminate in coma or paralysis.

Experimental autoimmune encephalomyelitis (EAE) is a primary model used in the art to examine and assess the effectiveness of therapeutic agents for treating MS. EAE is an inflammatory autoimmune demyelinating disease induced in laboratory animals by immunization with central nervous system tissue. When animals (e.g., mice, rats,

guinea pigs, rabbits, monkeys, etc.) are injected with adjuvant, e.g., complete Freund's adjuvant, plus myelin basic protein or proteolipid protein, EAE is induced, which is similar, pathologically to MS (see e.g., Alvord et al., *Experimental Allergic Encephalomyelitis-A Useful Model for Multiple Sclerosis*, Liss, New York, 1984; Swanborg, *Meth. Enzymol.* 162:413, 1988; and McCarron et al., *J. Immunol.*, 147: 3296, 1991.)

To evaluate rHuAFP or a fragment or analog thereof, EAE is induced in an appropriate laboratory animal, e.g., a mouse or rabbit, according to methods known in the art. To evaluate the compound's immunosuppressive effect on EAE, i.e., its ability to prevent or ameliorate EAE, the compound is administered according to standard methods, e.g., intravenously or intraperitoneal, at an appropriate dosage on a daily basis. Generally, administration is initiated prior to inducing EAE and/or after the clinical appearance of EAE. Control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP or related molecules. The effect of the test molecules on EAE is monitored according to any standard method. For example, weight loss and muscle paralysis in EAE-induced animals is monitored on a daily basis. If desired, histological inspection (e.g., by using any standard histochemical or immunohistochemical procedure, see e.g., Ausubel et al., *Current Protocols In Molecular Biology*, Greene Publishing Associates (John Wiley & Son), New York, 1994; Bancroft and Stevens, *Theory and Practice of Histochemical Techniques*, Churchill Livingstone, 1982) of brain and spinal cord tissues is performed and tissue samples examined microscopically for evidence of EAE, e.g., evidence of perivascular cellular infiltrates. Comparative studies between treated and control animals are used to determine the relative efficacy of the test molecules in preventing or ameliorating EAE. A molecule which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of EAE is considered useful in the invention.

#### Rheumatoid Arthritis

Rheumatoid arthritis (RA) is a common chronic illness in which the synovial membrane of multiple joints becomes inflamed, causing damage to cartilage and bone. RA is associated with human lymphocyte antigen (HLA)-DR4 and considered to be an autoimmune disorder involving T cells, see e.g., Sewell et al., *Lancet* 341: 283, 1993. RA results from a complex interaction of synovial cells with various cellular elements (and their soluble products) that infiltrate from the circulation into the synovial lining of joints. A series of biological events occur which ultimately lead to a lesion which invades and erodes collagen and the cartilage matrix of the joint.

A number of animal models of RA, e.g., the MRL-lpr/lpr mouse, are known in the art which develop a form of arthritis resembling the human disease (see e.g., *Fundamental Immunology*, supra). Alternatively, autoimmune collagen arthritis (ACA) and adjuvant arthritis (AA) can be induced in an appropriate animal according to standard methods.

To evaluate rHuAFP or a fragment or analog thereof on immunosuppressive on RA, i.e., the compound's ability to prevent or ameliorate RA, the test molecule is administered to a MRL-lpr/lpr mouse according to standard methods, e.g., intravenously or intraperitoneally, at an appropriate dosage on a daily basis. Generally, administration is initiated prior to the onset of RA and/or after the clinical appearance of RA. Control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP or related molecules. The effect of the test molecule on RA is monitored according to standard methods. For example, analysis

of the cellular component(s) of a synovial joint are monitored on a daily basis. If desired, histological inspection (e.g., by using any standard histochemical or immunohistochemical procedure, see e.g., Ausubel et al., supra; Bancroft and Stevens, supra) of the synovial joint is performed and tissue samples examined microscopically for evidence of RA, e.g., evidence of erosion of collagen and cartilage matrix in a joint. Comparative studies between treated and control animals are used to determine the relative efficacy of the test molecule in preventing or ameliorating RA. A test molecule which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of RA is considered useful in the invention.

#### Myasthenia Gravis

Myasthenia gravis (MG) is a disorder of neuromuscular transmission in which there are autoantibodies against acetylcholine receptors of neuromuscular junctions. Antibodies attack the junction, causing weakness and paralysis. Females are afflicted twice as often as males, typically during the third decade of life. Muscular weakness is the predominant feature of the disease. Clinical signs include drooping of the eyelids and double vision. There is an association between MG and hyperthyroidism.

Experimental autoimmune MG (EAMG) has been studied in a variety of animals including rabbits, monkeys, Lewis rats and inbred strains of mice (see e.g., *Principles of Neural Science*, supra), the symptoms of EAMG resemble the essential characteristics of the human disease. A single injection of acetylcholine receptor, e.g., purified from the electric organs of the eel *Torpedo californica*, along with adjuvants, causes an acute phase of weakness within 8 to 12 days and then chronic weakness after about 30 days. The response to the eel receptor is T cell dependent. The C57BL/6 strain (H-2B) is a high responder to Torpedo receptor and highly susceptible to EAMG.

To evaluate rHuAFP or a fragment or analog thereof, EAMG is induced in an appropriate laboratory animal, e.g., the C57BL/6 strain (H-2B) mouse, according to methods known in the art. To evaluate the compound's immunosuppressive effect on EAMG, i.e., its ability to prevent or ameliorate EAMG, the compound is administered according to standard methods, e.g., intravenously or intraperitoneally, at an appropriate dosage on a daily basis. Generally, administration is initiated prior to inducing EAMG and/or after the clinical appearance of EAMG. Control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP or related molecules. The effect of the test molecules on EAMG is monitored according to standard methods. For example, nerve stimulation in an electromyographic muscle assay (e.g., according to the methods of Pachner et al., *Ann. Neurol.* 11:48, 1982) in EAMG-induced animals can be assayed. If desired, histological inspection (e.g., by using any standard histochemical or immunohistochemical procedure, see e.g., Ausubel et al., supra; Bancroft and Stevens, supra) of tissue samples is performed and tissue samples examined microscopically for evidence of EAMG, e.g., evidence of monocyte infiltration and/or autoantibody localization at acetylcholine receptors of neuromuscular junctions. Comparative studies between treated and control animals are used to determine the relative efficacy of the test molecules in preventing or ameliorating EAMG. A molecule which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of EAMG is considered useful in the invention.

#### Insulin-Dependent Diabetes Mellitus

Diabetes is a disorder of glucose metabolism. Insulin-dependent diabetes mellitus (IDDM), also known as Type I



diabetes, is an autoimmune disease characterized by T-cell mediated destruction of pancreatic  $\beta$  cells in the islets of Langerhans, accompanied by an immune response to a diversity of self peptides leading to hyperglycemia, among other pathological events. IDDM patients depend on exogenous insulin to maintain normal glucose metabolism. Humans at risk for developing IDDM can be identified prior to onset of hyperglycemia by the abnormal occurrence of autoantibodies to insulin, islet cells, glutamic acid carboxylase, as well as other autologous proteins (see e.g., Baekkeskov et al., *J. Clin. Invest.* 79:926, 1987; Dean et al., *Diabetologia* 29: 339, 1986; Rossini et al., *Annu. Rev. Immunol.* 3:289, 1985; Srikanta et al., *N. Engl. J. Med.* 308:322, 1983). Autoantibody patterns, in general, are predictive for the eventual disease progression and/or risk for developing the disease (see e.g., Keller et al., *Lancet* 341:927, 1993).

Examples of animal models which spontaneously develop IDDM resembling the human disease include the Bio-Breeding (BB) rat and nonobese diabetic (NOD) mouse. Diabetes is also experimentally induced by streptozotocin.

The BB rat spontaneously develops a disease similar to IDDM, with insulinitis (infiltration of mononuclear cells into the pancreatic islets) and autoantibodies against self cells and insulin (see e.g., Baekkeskov et al., *J. Clin. Invest.* 79:926, 1987; Rossini et al, supra; Nakhooda et al., *Diabetes* 26: 100, 1977; Dean et al., *Clin. Exp. Immunol.* 69: 308, 1987).

NOD mice typically develop insulinitis between 5 and 8 weeks of age, and by 7 months 70% of the females and 40% of the males become diabetic. T cells transferred from diabetic mice to young nondiabetic NOD mice induce diabetes within 2 to 3 weeks (see e.g., Bendelac et al., *J. Exp. Med.* 166:823, 1987). NOD mice usually die within 1 to 2 months after the onset of diabetes unless they receive insulin therapy.

Chemically induced diabetes is accomplished using multiple injections of small doses of streptozotocin, a drug toxic for pancreatic  $\beta$  cells, which causes severe insulinitis and diabetes (see e.g., Kikutani et al., *Adv. Immunol.* 51:285, 1992).

Accordingly, the art provides a variety animal models resembling human IDDM which can be used to examine and assess approaches for the prevention or amelioration of diabetes involving rHuAFP (or a fragment or analog thereof).

To evaluate the immunosuppressive effect of rHuAFP or a fragment or analog thereof on the development of diabetes mouse, i.e., the compound's ability to treat or prevent insulinitis and diabetes, the test compound is administered to an appropriate test animal, e.g, a NOD mouse, according to standard methods, e.g., intravenously or intraperitoneally, at an appropriate dosage on a daily basis. Generally, administration is initiated prior to the onset of insulinitis and diabetes and/or after the clinical appearance of diabetic characteristics. control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP or related molecules. The effect of test molecules on insulinitis and diabetes is monitored according to standard methods. For example, weight loss, ketone body formation, and blood glucose concentration is monitored on a daily basis. If desired, histological inspection (e.g., by using any standard histochemical or immunohistochemical procedure, see e.g., Ausubel et al., supra; Bancroft and Stevens, supra) of pancreatic islet cells is performed and tissue samples examined microscopically for evidence of insulinitis and  $\beta$  cell destruction. Comparative studies between treated and con-

trol animals are used to determine the relative efficacy of the test molecules in preventing or ameliorating the diabetic condition. A molecule which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of diabetes, e.g., IDDM, is considered useful in the invention.

#### Systemic LuPus Erythematosus

Systemic lupus erythematosus (SLE) is a severe systemic autoimmune disease. About 90% of patients with this disease are young women. This marked preponderance of females is not seen before puberty or after menopause. The illness generally begins in young adulthood when a characteristic skin rash appears over cheekbones and forehead. Hair loss is common, as is severe kidney damage, arthritis, accumulation of fluid around the heart and inflammation of the lining of the lungs. In nearly half of the patients the blood vessels of the brain also become inflamed, leading to paralysis and convulsions. The activity of the disease, like other autoimmune diseases, can fluctuate: long quiescent periods of good health can terminate abruptly and inexplicably with the onset of a new attack. A large number of different autoantibodies are known to occur in SLE, e.g., autoantibodies against DNA, RNA and histones (see, e.g., *Fundamental Immunology*, supra)

A number of animal models of human SLE, e.g., inbred mouse strains including NZB mice-and their F<sub>1</sub> hybrids, MRL mice, and BXSB mice, are known in the art (see e.g., Bielschowsky et al. *Proc. Univ. Otago Med. Sch.* 37:9, 1959; Braverman et al., *J. Invest. Derm.* 50: 483, 1968; Howie et al. *Adv. Immunol.* 9:215, 1968; *Genetic Control of Autoimmune Disease*, Rose, M., Bigazzi, P. E., and Warner, N. L. eds., Elsevier, Amsterdam, 1979; and *Current Protocols In Immunology*, supra). For example, the NZBXNZW F<sub>1</sub> mouse is an excellent model of human SLE, female mice develop high levels of anti-double- and single-stranded DNA autoantibodies, other anti-nuclear antibodies, and renal disease; death usually occurs at approximately 8 months (see e.g., Theofilopoulos et al., *Adv. Immunol.* 37:269, 1985).

To evaluate the immunosuppressive effect of rHuAFP or a fragment or analog thereof on SLE, i.e., the compound's ability of rHuAFP to prevent or ameliorate SLE, test compounds are administered to an appropriate animal, e.g., the NZBXNZW F<sub>1</sub> mouse, according to standard methods, e.g., intravenously or intraperitoneally, at an appropriate dosage on a daily basis. Generally, administration is initiated prior to the onset of SLE and/or after the clinical appearance of SLE. Control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP or related molecules. The effect of the test compound on SLE is monitored according to standard methods. For example, analysis of autoantibodies, e.g., anti-DNA antibodies can be monitored. If desired, histological inspection (e.g., by using any standard histochemical or immunohistochemical procedure, see e.g., Ausubel et al., supra; Bancroft and Stevens, supra) of kidney tissue is performed and tissue samples examined microscopically for evidence of SLE, e.g., evidence of lupus nephritis. Comparative studies between treated and control animals are used to determine the relative efficacy of the test compounds in preventing or ameliorating SLE. A molecule which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of SLE is considered useful in the invention.

#### Therapeutic Administration

As demonstrated above, recombinant alpha-fetoprotein, e.g., rHuAFP (or a fragment or analog thereof) is effective

in inhibiting proliferation of autoimmune cells and accordingly is useful for the prevention or amelioration of autoimmune diseases including, but not limited to, multiple sclerosis, rheumatoid arthritis, diabetes mellitus, systemic lupus erythematosus, and myasthenia gravis. Accordingly, recombinant human alpha-fetoprotein (or a fragment or analog thereof) can be formulated according to known methods to prepare pharmaceutically useful compositions.

Recombinant alpha-fetoprotein, e.g., rHuAFP (or a fragment or analog thereof), is preferably administered to the patient in an amount which is effective in preventing or ameliorating the symptoms of an autoimmune disease. Generally, a dosage of 0.1 ng/kg to 10 g/kg body is adequate. If desired, administration is performed on a daily basis. Because there are no known adverse side effects related to recombinant human alpha-fetoprotein, it is believed that relatively high dosages can be safely administered. For example, treatment of human patients will be carried out using a therapeutically effective amount of rHuAFP (or a fragment or analog thereof) in a physiologically acceptable carrier. Suitable carriers and their formulation are described for example in Remington's *Pharmaceutical Sciences* by E. W. Martin. The amount of rHuAFP to be administered will vary depending upon the manner of administration, the age and body weight of the patient, and with the type of disease, and size of the patient predisposed to or suffering from the disease. Preferable routes of administration include, for example, subcutaneous, intravenous, intramuscular, or intradermal injections which provide continuous, sustained levels of the drug in the patient. In other preferred routes of administration, rHuAFP can be given to a patient by injection or implantation of a slow release preparation, for example, in a slowly dissociating polymeric or crystalline form; this sort of sustained administration can follow an initial delivery of the drug by more conventional routes (for example, those described above). Alternatively, rHuAFP can be administered using an infusion pump (e.g., an external or implantable infusion pump), thus allowing a precise degree of control over the rate of drug release, or through installation of rHuAFP in the nasal passages in a similar fashion to that used to promote absorption of insulin. As an alternative to nasal transmucosal absorption, rHuAFP can be delivered by aerosol deposition of the powder or solution into the lungs.

Furthermore, the method(s) of the invention can also employ combination therapy in which rHuAFP is administered either simultaneously or sequentially with a therapeutic agent such as a general or specific tolerizing agent (e.g., an anti-idiotypic agent (e.g., a monoclonal) or a therapeutic vaccine or an oral agent (e.g., insulin, collagen or myelin basic protein) or a cytokine (e.g., Il-15) or an interferon ( $\alpha$ -interferon) or an immunosuppressive agent. Preferably, an immunosuppressive agent is administered in an effective dose which is lower than the standard dose when the immunosuppressive agent is used by itself. Preferred immunosuppressive agents are cyclosporine, FK-506, steroids, azathioprine, or 15-deoxyspergualin.

Treatment is started generally with the diagnosis or suspicion of an autoimmune disease and is generally repeated on a daily basis. Protection or prevention from the development (or progression or exacerbation) of an autoimmune disease is also achieved by administration of rHuAFP prior to the onset of the disease. If desired, the efficacy of the treatment or protection regimens is assessed with the methods of monitoring or diagnosing patients for autoimmune disease.

The method(s) of the invention can also be used to treat non-human mammals, for example, domestic pets, or livestock.

## Other Embodiments

In other embodiments, the invention includes the use of rHuAFP (or fragment or analog thereof) for the prevention or treatment of acquired immunodeficiency syndrome (AIDS). To evaluate the immunosuppressive effect of rHuAFP or a fragment or analog thereof on AIDS, i.e., the compound's ability to prevent or ameliorate an autoimmune component of AIDS, test compounds are administered to an appropriate animal (e.g., a human patient), according to standard methods, e.g., intravenously or intraperitoneally, at an appropriate dosage on a daily basis as is discussed above. Generally, administration is initiated prior to the onset of AIDS and/or after the clinical appearance of AIDS. Control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP or related molecules. The effect of the test compound on AIDS is monitored according to standard methods. For example, analysis of the ability of the test compound to inhibit or prevent or ameliorate the destruction of helper T cells can be monitored. Comparative studies between treated and control animals are used to determine the relative efficacy of the test compounds in preventing or ameliorating AIDS. A molecule which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of AIDS is considered useful in the invention.

In the invention also includes the use of a therapeutically effective amount rHuAFP (or fragment or analog thereof) for inhibiting the rejection of a transplanted organ (e.g., the heart, the liver, the lung, the pancreas, and the kidney), tissue (e.g., skin, bone marrow, dura mater, bone, implanted collagen, an implanted bioreactor), or cell (e.g.,  $\beta$  islet cells of the pancreas, stem cells, hematopoietic cells, lymph cells, neuroendocrine or adrenal cells) in a mammal. Such transplanted organs, tissues, or cells may be derived from any source, e.g., such biological material can be allogenic, phenogenic, autologous, synthetic, artificial or genetically-engineered. For example, the method can also be used when the patient is the recipient of an allograft such a heart or kidney from another species.

In one working example, the immunosuppressive effect of rHuAFP on clinical transplantation, i.e., the ability of rHuAFP to prevent or ameliorate transplant rejection (e.g., hyperacute rejection, acute rejection and chronic rejection), is evaluated by administering rHuAFP to an NIH minipig according to standard methods, e.g., intravenously or intraperitoneally, at an appropriate dosage on a daily basis. Generally, administration of rHuAFP is initiated prior to the transplant, e.g., transplantation of a kidney and/or after the transplant procedure. Control animals receive a placebo, e.g., human serum albumin, similarly administered as for rHuAFP. The effect of rHuAFP on transplant rejection is monitored according to standard methods. One manifestation of the rejection process is diminished function of the transplanted organ, for example, analysis of urine output can be monitored. If desired, histological inspection (e.g., by using any standard histochemical or immunohistochemical procedure, see e.g., Ausubel et al., supra; Bancroft and Stevens, supra) of kidney tissue is performed and tissue samples obtained by biopsy are examined microscopically for evidence of transplant rejection, e.g., chronic interstitial fibrosis, vascular thrombosis, or the presence of abnormal lymphocytic infiltrates. Comparative studies between treated and control animals are used to determine the relative

efficacy of rHuAFP in preventing or ameliorating transplant rejection. Recombinant HuAFP (a fragment or analog thereof) which prevents or ameliorates (decreases or suppresses or relieves or promotes remission of) the symptoms of transplant rejection is considered useful in the invention.

All publications, manufacturer's instructions, patents, and patent applications mentioned in this specification are herein incorporated by reference to the same extent as if each individual publication or patent application was specifically and individually indicated to be incorporated by reference.

---

 SEQUENCE LISTING
 

---

<160> NUMBER OF SEQ ID NOS: 16

<210> SEQ ID NO 1

<211> LENGTH: 2027

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

atattgtgct tccaccactg ccaataacaa aataactagc aaccatgaag tgggtggaat    60
caatTTTTTT aattttccta ctaaatttta ctgaatccag aacctgcat agaatgaat    120
atggaatagc ttccatattg gattcttacc aatgtactgc agagataagt ttagctgacc    180
tggctaccat atTTTTTgcc cagtttgttc aagaagccac ttacaaggaa gtaagcaaaa    240
tggtgaaaga tgcattgact gcaattgaga aacctactgg agatgaacag tcttcagggt    300
gttttagaaaa ccagctacct gcctttctgg aagaactttg ccatgagaaa gaaatTTTgg    360
agaagtacgg acattcagac tgctgcagcc aaagtgaaga gggaagacat aactgTTTTc    420
ttgcacacaa aaagcccact gcagcatgga tcccactttt ccaagttcca gaacctgtca    480
caagctgtga agcatatgaa gaagacaggg agacattcat gaacaaattc atttatgaga    540
tagcaagaag gcatcccttc ctgtatgcac ctacaattct tctttcggct gctgggtatg    600
agaaaataat tccatcttgc tgcaaagctg aaaatgcagt tgaatgcttc caaacaagg    660
cagcaacagt taaaaagaa ttaagagaaa gcagcttggt aaatcaacat gcatgtccag    720
taatgaaaaa ttttgggacc cgaactttcc aagccataac tgttactaaa ctgagtcaga    780
agtttaccaa agttaatttt actgaaatcc agaaactagt cctggatgtg gcccatgtac    840
atgagcactg ttgcagagca gatgtgctgg attgtctgca ggatggggaa aaaatcatgt    900
cctacatatg ttctcaacaa gacactctgt caaacaaaat aacagaatgc tgcaaactga    960
ccacgctgga acgtggtcaa tgtataattc atgcagaaaa tgatgaaaaa cctgaaggtc   1020
tatctccaaa tctaaacagg tttttaggag atagagattt taaccaattt tcttcagggg   1080
aaaaaaaaat cttcttgcca agttttgttc atgaatattc aagaagacat cctcagcttg   1140
ctgtctcagt aattctaaga gttgctaaag gataccagga gttattggag aagtgtttcc   1200
agactgaaaa ccctcttgaa tgccaagata aaggagaaga agaattacag aaatacatcc   1260
aggagagcca agcattggca aagcgaagct gcggcctott ccagaaacta ggagaatatt   1320
acttacaaaa tgagtttctc gttgcttaca caaagaaagc ccccagctg acctcgtcgg   1380
agctgatggc catcaccaga aaaatggcag ccacagcagc cacttgttgc caactcagtg   1440
aggacaaact attggcctgt ggcgagggag cggtgacat tattatcgga cacttatgta   1500
tcagacatga aatgactcca gtaaaccctg gtgttgcca gtgctgact tcttcatatg   1560
ccaacaggag gccatgcttc agcagcttgg tggtgatga aacatatgtc cctcctgcat   1620
tctctgatga caagttcatt ttccataagg atctgtgcca agctcagggg gtagcgctgc   1680
aaaggatgaa gcaagagttt ctcatcacc ttgtgaagca aaagccacaa ataacagagg   1740
aacaacttga ggctctcatt gcagatttct caggcctggt ggagaaatgc tgccaaggcc   1800

```

-continued

---

```

aggAACAGGA agtctgcttt gctgaagagg gacaaaaact gatttcaaaa actggtgctg 1860
ctttgggagt ttaaattact tcaggggaag agaagacaaa acgagtcttt cattcggtgt 1920
gaacttttct ctttaatttt aactgattta acactttttg tgaattaatg aaatgataaa 1980
gacttttatg tgagatttcc ttatcacaga aataaaatat ctccaaa 2027

```

```

<210> SEQ ID NO 2
<211> LENGTH: 590
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 2

```

```

Thr Leu His Arg Asn Glu Tyr Gly Ile Ala Ser Ile Leu Asp Ser Tyr
 1           5           10           15
Gln Cys Thr Ala Glu Ile Ser Leu Ala Asp Leu Ala Thr Ile Phe Phe
          20           25           30
Ala Gln Phe Val Gln Glu Ala Thr Tyr Lys Glu Val Ser Lys Met Val
          35           40           45
Lys Asp Ala Leu Thr Ala Ile Glu Lys Pro Thr Gly Asp Glu Gln Ser
          50           55           60
Ser Gly Cys Leu Glu Asn Gln Leu Pro Ala Phe Leu Glu Glu Leu Cys
          65           70           75           80
His Glu Lys Glu Ile Leu Glu Lys Tyr Gly His Ser Asp Cys Cys Ser
          85           90           95
Gln Ser Glu Glu Gly Arg His Asn Cys Phe Leu Ala His Lys Lys Pro
          100          105          110
Thr Ala Ala Trp Ile Pro Leu Phe Gln Val Pro Glu Pro Val Thr Ser
          115          120          125
Cys Glu Ala Tyr Glu Glu Asp Arg Glu Thr Phe Met Asn Lys Phe Ile
          130          135          140
Tyr Glu Ile Ala Arg Arg His Pro Phe Leu Tyr Ala Pro Thr Ile Leu
          145          150          155          160
Leu Ser Ala Ala Gly Tyr Glu Lys Ile Ile Pro Ser Cys Cys Lys Ala
          165          170          175
Glu Asn Ala Val Glu Cys Phe Gln Thr Lys Ala Ala Thr Val Thr Lys
          180          185          190
Glu Leu Arg Glu Ser Ser Leu Leu Asn Gln His Ala Cys Pro Val Met
          195          200          205
Lys Asn Phe Gly Thr Arg Thr Phe Gln Ala Ile Thr Val Thr Lys Leu
          210          215          220
Ser Gln Lys Phe Thr Lys Val Asn Phe Thr Glu Ile Gln Lys Leu Val
          225          230          235          240
Leu Asp Val Ala His Val His Glu His Cys Cys Arg Ala Asp Val Leu
          245          250          255
Asp Cys Leu Gln Asp Gly Glu Lys Ile Met Ser Tyr Ile Cys Ser Gln
          260          265          270
Gln Asp Thr Leu Ser Asn Lys Ile Thr Glu Cys Cys Lys Leu Thr Thr
          275          280          285
Leu Glu Arg Gly Gln Cys Ile Ile His Ala Glu Asn Asp Glu Lys Pro
          290          295          300
Glu Gly Leu Ser Pro Asn Leu Asn Arg Phe Leu Gly Asp Arg Asp Phe
          305          310          315          320
Asn Gln Phe Ser Ser Gly Glu Lys Asn Ile Phe Leu Ala Ser Phe Val
          325          330          335

```

-continued

---

His Glu Tyr Ser Arg Arg His Pro Gln Leu Ala Val Ser Val Ile Leu  
                   340  345  350

Arg Val Ala Lys Gly Tyr Gln Glu Leu Leu Glu Lys Cys Phe Gln Thr  
                   355  360  365

Glu Asn Pro Leu Glu Cys Gln Asp Lys Gly Glu Glu Glu Leu Gln Lys  
                   370  375  380

Tyr Ile Gln Glu Ser Gln Ala Leu Ala Lys Arg Ser Cys Gly Leu Phe  
 385  390  395  400

Gln Lys Leu Gly Glu Tyr Tyr Leu Gln Asn Glu Phe Leu Val Ala Tyr  
                   405  410  415

Thr Lys Lys Ala Pro Gln Leu Thr Ser Ser Glu Leu Met Ala Ile Thr  
                   420  425  430

Arg Lys Met Ala Ala Thr Ala Ala Thr Cys Cys Gln Leu Ser Glu Asp  
                   435  440  445

Lys Leu Leu Ala Cys Gly Glu Gly Ala Ala Asp Ile Ile Ile Gly His  
                   450  455  460

Leu Cys Ile Arg His Glu Met Thr Pro Val Asn Pro Gly Val Gly Gln  
 465  470  475  480

Cys Cys Thr Ser Ser Tyr Ala Asn Arg Arg Pro Cys Phe Ser Ser Leu  
                   485  490  495

Val Val Asp Glu Thr Tyr Val Pro Pro Ala Phe Ser Asp Asp Lys Phe  
                   500  505  510

Ile Phe His Lys Asp Leu Cys Gln Ala Gln Gly Val Ala Leu Gln Arg  
                   515  520  525

Met Lys Gln Glu Phe Leu Ile Asn Leu Val Lys Gln Lys Pro Gln Ile  
                   530  535  540

Thr Glu Glu Gln Leu Glu Ala Leu Ile Ala Asp Phe Ser Gly Leu Leu  
 545  550  555  560

Glu Lys Cys Cys Gln Gly Gln Glu Gln Glu Val Cys Phe Ala Glu Glu  
                   565  570  575

Gly Gln Lys Leu Ile Ser Lys Thr Gly Ala Ala Leu Gly Val  
                   580  585  590

&lt;210&gt; SEQ ID NO 3

&lt;211&gt; LENGTH: 197

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Homo sapiens

&lt;400&gt; SEQUENCE: 3

Thr Leu His Arg Asn Glu Tyr Gly Ile Ala Ser Ile Leu Asp Ser Tyr  
 1                  5  10  15

Gln Cys Thr Ala Glu Ile Ser Leu Ala Asp Leu Ala Thr Ile Phe Phe  
                   20  25  30

Ala Gln Phe Val Gln Glu Ala Thr Tyr Lys Glu Val Ser Lys Met Val  
                   35  40  45

Lys Asp Ala Leu Thr Ala Ile Glu Lys Pro Thr Gly Asp Glu Gln Ser  
                   50  55  60

Ser Gly Cys Leu Glu Asn Gln Leu Pro Ala Phe Leu Glu Glu Leu Cys  
 65  70  75  80

His Glu Lys Glu Ile Leu Glu Lys Tyr Gly His Ser Asp Cys Cys Ser  
                   85  90  95

Gln Ser Glu Glu Gly Arg His Asn Cys Phe Leu Ala His Lys Lys Pro  
                   100  105  110

Thr Ala Ala Trp Ile Pro Leu Phe Gln Val Pro Glu Pro Val Thr Ser  
                   115  120  125

-continued

Cys Glu Ala Tyr Glu Glu Asp Arg Glu Thr Phe Met Asn Lys Phe Ile  
 130 135 140  
 Tyr Glu Ile Ala Arg Arg His Pro Phe Leu Tyr Ala Pro Thr Ile Leu  
 145 150 155 160  
 Leu Ser Ala Ala Gly Tyr Glu Lys Ile Ile Pro Ser Cys Cys Lys Ala  
 165 170 175  
 Glu Asn Ala Val Glu Cys Phe Gln Thr Lys Ala Ala Thr Val Thr Lys  
 180 185 190  
 Glu Leu Arg Glu Ser  
 195

<210> SEQ ID NO 4  
 <211> LENGTH: 192  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Ser Leu Leu Asn Gln His Ala Cys Pro Val Met Lys Asn Phe Gly Thr  
 1 5 10 15  
 Arg Thr Phe Gln Ala Ile Thr Val Thr Lys Leu Ser Gln Lys Phe Thr  
 20 25 30  
 Lys Val Asn Phe Thr Glu Ile Gln Lys Leu Val Leu Asp Val Ala His  
 35 40 45  
 Val His Glu His Cys Cys Arg Ala Asp Val Leu Asp Cys Leu Gln Asp  
 50 55 60  
 Gly Glu Lys Ile Met Ser Tyr Ile Cys Ser Gln Gln Asp Thr Leu Ser  
 65 70 75 80  
 Asn Lys Ile Thr Glu Cys Cys Lys Leu Thr Thr Leu Glu Arg Gly Gln  
 85 90 95  
 Cys Ile Ile His Ala Glu Asn Asp Glu Lys Pro Glu Gly Leu Ser Pro  
 100 105 110  
 Asn Leu Asn Arg Phe Leu Gly Asp Arg Asp Phe Asn Gln Phe Ser Ser  
 115 120 125  
 Gly Glu Lys Asn Ile Phe Leu Ala Ser Phe Val His Glu Tyr Ser Arg  
 130 135 140  
 Arg His Pro Gln Leu Ala Val Ser Val Ile Leu Arg Val Ala Lys Gly  
 145 150 155 160  
 Tyr Gln Glu Leu Leu Glu Lys Cys Phe Gln Thr Glu Asn Pro Leu Glu  
 165 170 175  
 Cys Gln Asp Lys Gly Glu Glu Glu Leu Gln Lys Tyr Ile Gln Glu Ser  
 180 185 190

<210> SEQ ID NO 5  
 <211> LENGTH: 201  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 5

Gln Ala Leu Ala Lys Arg Ser Cys Gly Leu Phe Gln Lys Leu Gly Glu  
 1 5 10 15  
 Tyr Tyr Leu Gln Asn Glu Phe Leu Val Ala Tyr Thr Lys Lys Ala Pro  
 20 25 30  
 Gln Leu Thr Ser Ser Glu Leu Met Ala Ile Thr Arg Lys Met Ala Ala  
 35 40 45  
 Thr Ala Ala Thr Cys Cys Gln Leu Ser Glu Asp Lys Leu Leu Ala Cys  
 50 55 60

-continued

Gly Glu Gly Ala Ala Asp Ile Ile Ile Gly His Leu Cys Ile Arg His  
 65 70 75 80  
 Glu Met Thr Pro Val Asn Pro Gly Val Gly Gln Cys Cys Thr Ser Ser  
 85 90 95  
 Tyr Ala Asn Arg Arg Pro Cys Phe Ser Ser Leu Val Val Asp Glu Thr  
 100 105 110  
 Tyr Val Pro Pro Ala Phe Ser Asp Asp Lys Phe Ile Phe His Lys Asp  
 115 120 125  
 Leu Cys Gln Ala Gln Gly Val Ala Leu Gln Arg Met Lys Gln Glu Phe  
 130 135 140  
 Leu Ile Asn Leu Val Lys Gln Lys Pro Gln Ile Thr Glu Glu Gln Leu  
 145 150 155 160  
 Glu Ala Leu Ile Ala Asp Phe Ser Gly Leu Leu Glu Lys Cys Cys Gln  
 165 170 175  
 Gly Gln Glu Gln Glu Val Cys Phe Ala Glu Glu Gly Gln Lys Leu Ile  
 180 185 190  
 Ser Lys Thr Gly Ala Ala Leu Gly Val  
 195 200

<210> SEQ ID NO 6  
 <211> LENGTH: 389  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 6

Thr Leu His Arg Asn Glu Tyr Gly Ile Ala Ser Ile Leu Asp Ser Tyr  
 1 5 10 15  
 Gln Cys Thr Ala Glu Ile Ser Leu Ala Asp Leu Ala Thr Ile Phe Phe  
 20 25 30  
 Ala Gln Phe Val Gln Glu Ala Thr Tyr Lys Glu Val Ser Lys Met Val  
 35 40 45  
 Lys Asp Ala Leu Thr Ala Ile Glu Lys Pro Thr Gly Asp Glu Gln Ser  
 50 55 60  
 Ser Gly Cys Leu Glu Asn Gln Leu Pro Ala Phe Leu Glu Glu Leu Cys  
 65 70 75 80  
 His Glu Lys Glu Ile Leu Glu Lys Tyr Gly His Ser Asp Cys Cys Ser  
 85 90 95  
 Gln Ser Glu Glu Gly Arg His Asn Cys Phe Leu Ala His Lys Lys Pro  
 100 105 110  
 Thr Ala Ala Trp Ile Pro Leu Phe Gln Val Pro Glu Pro Val Thr Ser  
 115 120 125  
 Cys Glu Ala Tyr Glu Glu Asp Arg Glu Thr Phe Met Asn Lys Phe Ile  
 130 135 140  
 Tyr Glu Ile Ala Arg Arg His Pro Phe Leu Tyr Ala Pro Thr Ile Leu  
 145 150 155 160  
 Leu Ser Ala Ala Gly Tyr Glu Lys Ile Ile Pro Ser Cys Cys Lys Ala  
 165 170 175  
 Glu Asn Ala Val Glu Cys Phe Gln Thr Lys Ala Ala Thr Val Thr Lys  
 180 185 190  
 Glu Leu Arg Glu Ser Ser Leu Leu Asn Gln His Ala Cys Pro Val Met  
 195 200 205  
 Lys Asn Phe Gly Thr Arg Thr Phe Gln Ala Ile Thr Val Thr Lys Leu  
 210 215 220  
 Ser Gln Lys Phe Thr Lys Val Asn Phe Thr Glu Ile Gln Lys Leu Val





-continued

---

Tyr Tyr Leu Gln Asn Glu Phe Leu Val Ala Tyr Thr Lys Lys Ala Pro  
 210 215 220  
 Gln Leu Thr Ser Ser Glu Leu Met Ala Ile Thr Arg Lys Met Ala Ala  
 225 230 235 240  
 Thr Ala Ala Thr Cys Cys Gln Leu Ser Glu Asp Lys Leu Leu Ala Cys  
 245 250 255  
 Gly Glu Gly Ala Ala Asp Ile Ile Ile Gly His Leu Cys Ile Arg His  
 260 265 270  
 Glu Met Thr Pro Val Asn Pro Gly Val Gly Gln Cys Cys Thr Ser Ser  
 275 280 285  
 Tyr Ala Asn Arg Arg Pro Cys Phe Ser Ser Leu Val Val Asp Glu Thr  
 290 295 300  
 Tyr Val Pro Pro Ala Phe Ser Asp Asp Lys Phe Ile Phe His Lys Asp  
 305 310 315 320  
 Leu Cys Gln Ala Gln Gly Val Ala Leu Gln Arg Met Lys Gln Glu Phe  
 325 330 335  
 Leu Ile Asn Leu Val Lys Gln Lys Pro Gln Ile Thr Glu Glu Gln Leu  
 340 345 350  
 Glu Ala Leu Ile Ala Asp Phe Ser Gly Leu Leu Glu Lys Cys Cys Gln  
 355 360 365  
 Gly Gln Glu Gln Glu Val Cys Phe Ala Glu Glu Gly Gln Lys Leu Ile  
 370 375 380  
 Ser Lys Thr Gly Ala Ala Leu Gly Val  
 385 390

<210> SEQ ID NO 8  
 <211> LENGTH: 325  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Met Ser Tyr Ile Cys Ser Gln Gln Asp Thr Leu Ser Asn Lys Ile Thr  
 1 5 10 15  
 Glu Cys Cys Lys Leu Thr Thr Leu Glu Arg Gly Gln Cys Ile Ile His  
 20 25 30  
 Ala Glu Asn Asp Glu Lys Pro Glu Gly Leu Ser Pro Asn Leu Asn Arg  
 35 40 45  
 Phe Leu Gly Asp Arg Asp Phe Asn Gln Phe Ser Ser Gly Glu Lys Asn  
 50 55 60  
 Ile Phe Leu Ala Ser Phe Val His Glu Tyr Ser Arg Arg His Pro Gln  
 65 70 75 80  
 Leu Ala Val Ser Val Ile Leu Arg Val Ala Lys Gly Tyr Gln Glu Leu  
 85 90 95  
 Leu Glu Lys Cys Phe Gln Thr Glu Asn Pro Leu Glu Cys Gln Asp Lys  
 100 105 110  
 Gly Glu Glu Glu Leu Gln Lys Tyr Ile Gln Glu Ser Gln Ala Leu Ala  
 115 120 125  
 Lys Arg Ser Cys Gly Leu Phe Gln Lys Leu Gly Glu Tyr Tyr Leu Gln  
 130 135 140  
 Asn Glu Phe Leu Val Ala Tyr Thr Lys Lys Ala Pro Gln Leu Thr Ser  
 145 150 155 160  
 Ser Glu Leu Met Ala Ile Thr Arg Lys Met Ala Ala Thr Ala Ala Thr  
 165 170 175  
 Cys Cys Gln Leu Ser Glu Asp Lys Leu Leu Ala Cys Gly Glu Gly Ala  
 180 185 190

-continued

Ala Asp Ile Ile Ile Gly His Leu Cys Ile Arg His Glu Met Thr Pro  
 195 200 205

Val Asn Pro Gly Val Gly Gln Cys Cys Thr Ser Ser Tyr Ala Asn Arg  
 210 215 220

Arg Pro Cys Phe Ser Ser Leu Val Val Asp Glu Thr Tyr Val Pro Pro  
 225 230 235 240

Ala Phe Ser Asp Asp Lys Phe Ile Phe His Lys Asp Leu Cys Gln Ala  
 245 250 255

Gln Gly Val Ala Leu Gln Arg Met Lys Gln Glu Phe Leu Ile Asn Leu  
 260 265 270

Val Lys Gln Lys Pro Gln Ile Thr Glu Glu Gln Leu Glu Ala Leu Ile  
 275 280 285

Ala Asp Phe Ser Gly Leu Leu Glu Lys Cys Cys Gln Gly Gln Glu Gln  
 290 295 300

Glu Val Cys Phe Ala Glu Glu Gly Gln Lys Leu Ile Ser Lys Thr Gly  
 305 310 315 320

Ala Ala Leu Gly Val  
 325

<210> SEQ ID NO 9  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

aaaaaaggta ccacactgca tagaaatgaa 30

<210> SEQ ID NO 10  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

aaaaaaggat ccttagcttt ctcttaattc ttt 33

<210> SEQ ID NO 11  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 11

aaaaaaatcg atatgagctt gttaaataca cat 33

<210> SEQ ID NO 12  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

aaaaaaggat ccttagctct cctggatgta ttt 33

<210> SEQ ID NO 13  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 13

aaaaaaatcg atatgcaagc attggcaaag cga 33

-continued

<210> SEQ ID NO 14  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

aaaaaaggat ccttaaactc ccaaagcagc acg

33

<210> SEQ ID NO 15  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

aaaaaatcg atatgtccta catatgttct caa

33

<210> SEQ ID NO 16  
 <211> LENGTH: 9  
 <212> TYPE: PRT  
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

Ser Tyr Ile Cys Ser Gln Gln Asp Thr  
 1 5

What is claimed is:

1. A method of inhibiting autoreactive immune cell proliferation in a mammal, said method comprising administering to said mammal a therapeutically effective amount of recombinant human alpha-fetoprotein or a fragment thereof comprising Domain I, Domain II, Domain III, Domain I+II, Domain II+III, or Fragment I.

2. The method of claim 1, wherein said immune cells include T cells.

3. The method of claim 1, wherein said mammal is a human patient.

4. A method of treating an autoimmune disease in a mammal, said method comprising administering to said mammal a therapeutically effective amount of recombinant human alpha-fetoprotein or a fragment thereof comprising Domain I, Domain II, Domain III, Domain I+II, Domain II+III, or Fragment I.

5. The method of claim 4, wherein said autoimmune disease is multiple sclerosis.

6. The method of claim 4, wherein said autoimmune disease is rheumatoid arthritis.

7. The method of claim 4, wherein said autoimmune disease is myasthenia gravis.

8. The method of claim 4, wherein said autoimmune disease is insulin-dependent diabetes mellitus.

9. The method of claim 4, wherein said autoimmune disease is systemic lupus erythematosus.

10. The method of claim 1, further comprising administering to said mammal an immunosuppressive agent in an effective dose which is lower than the standard dose when said immunosuppressive agent is used by itself.

11. The method of claim 1, further comprising administering to said mammal a tolerizing agent.

12. The method of claim 10, wherein said immunosuppressive agent is cyclosporine.

13. The method of claim 10, wherein said immunosuppressive agent is a steroid, azathioprine, FK-506, or 15-deoxyspergualin.

\* \* \* \* \*

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 6,288,034 B1  
DATED : September 11, 2001  
INVENTOR(S) : Robert A. Murgita

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title page,

OTHER PUBLICATIONS, replace the Brenner reference title "Inhibitory Effect Of  $\alpha$ -fetoprotein On the Binding Of *Myasthenia gravis* Antibody To Acetylcholine Receptor" with -- Inhibitory Effect Of  $\alpha$ -fetoprotein On the Binding Of *Myasthenia gravis* Antibody To Acetylcholine Receptor --;

OTHER PUBLICATIONS, replace the Murgita reference title "Adult murine T cells activated in vitro by  $\alpha$ -fetoprotein and naturally occurring T cells in newborn mice: Identity in function and cell surface differentiation antigens," with -- Adult murine T cells activated in vitro by  $\alpha$ -fetoprotein and naturally occurring T cells in newborn mice: Identity in function and cell surface differentiation antigens --;

Column 2,

Line 2, replace "cell e.g.," with -- cell (e.g., --;  
Line 29, replace "rHuAPP" with -- rHuAFP --;  
Line 36, replace "and" with -- and --;  
Line 47, replace "FIGS. 1A" with -- FIG. 1A --;

Column 3,

Line 5, replace "▲" with -- ▼ --;  
Line 11, replace "▲" with -- ▼ --;

Column 6,

Line 56, replace "1X reaction buffer," with -- 10X reaction buffer --;

Column 9,

Lines 6 through 8, put a blank line between "Results are expressed as mean cpm±the standard error of the mean of triplicate or quadruplicate cultures." and "RESULTS";  
Line 53, replace "NAb." with -- MAb --;

Column 10,

Line 23, replace "organs-in" with -- organs in --;

Column 12,

Line 29, replace "Torpedo californica" with -- *Torpedo californica* --;  
Line 33, replace "(H-2B)" with -- (H-2<sup>B</sup>) --;  
Line 37, replace "(H-2B)" with -- (H-2<sup>B</sup>) --;

UNITED STATES PATENT AND TRADEMARK OFFICE  
**CERTIFICATE OF CORRECTION**

PATENT NO. : 6,288,034 B1  
DATED : September 11, 2001  
INVENTOR(S) : Robert A. Murgita

Page 2 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 13,

Line 13, replace "*Enal.*" with -- *Engl.* --;

Line 56, replace "control animals" with -- Control animals --;

Column 14,

Line 7, replace "LuPus" with -- Lupus --; and

Line 26, replace "mice-and" with -- mice and --.

Signed and Sealed this

Sixteenth Day of April, 2002

Attest:



Attesting Officer

JAMES E. ROGAN  
Director of the United States Patent and Trademark Office