



US006346242B1

(12) **United States Patent**
Leone-Bay et al.

(10) **Patent No.: US 6,346,242 B1**
(45) **Date of Patent: Feb. 12, 2002**

(54) **COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS**

2,828,206 A 3/1958 Rosenberg 99/2
2,862,918 A 12/1958 Meyer et al. 260/123.5

(75) Inventors: **Andrea Leone-Bay**, Ridgefield, CT (US); **Koc-Kan Ho**, Monmouth Junction, NJ (US); **Donald J. Sarubbi**, Carmel; **Sam J. Milstein**, Larchmont, both of NY (US)

(List continued on next page.)

FOREIGN PATENT DOCUMENTS

CA 1077842 8/1976 A61K/9/50
DE 2 424 169 12/1974 A61K/9/00
DE 2 343 037 3/1975

(73) Assignee: **Emisphere Technologies, Inc.**, Tarrytown, NY (US)

(List continued on next page.)

OTHER PUBLICATIONS

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

Picciola, Sintesi Di Acidi Chinazolinonici . . . Il Farmaco—Ed. Sc. vol. 31, No. 9, pp. 655–664, 1976.*
Airaudo, C.B. et al. (1987) *Journal of Food Science*, vol. 52(6), pp. 1750–1752.
Andini, S. et al. (1975) *Origins of Life*, vol. 6, pp. 147–153.
Brooke, S. 1 et al. (1977) *BioSystems*, vol. 9, pp. 1–22.

(21) Appl. No.: **09/499,958**

(22) Filed: **Feb. 8, 2000**

(List continued on next page.)

Related U.S. Application Data

Primary Examiner—Jeffrey E. Russel

(63) Continuation of application No. 09/305,506, filed on May 5, 1999, which is a continuation of application No. 08/798,031, filed on Feb. 6, 1997, now Pat. No. 6,001,347, which is a continuation of application No. PCT/US96/04580, filed on Apr. 1, 1996, which is a continuation-in-part of application No. 08/414,654, filed on Mar. 31, 1995, now Pat. No. 5,650,386.

(74) *Attorney, Agent, or Firm*—Darby & Darby


(60) Provisional application No. 60/003,111, filed on Sep. 1, 1995, and provisional application No. 60/017,902, filed on Mar. 29, 1996.

(57) **ABSTRACT**

(51) **Int. Cl.**⁷ **A61K 31/715**; A61K 38/17; A61K 38/29; C07C 229/42

The present invention provides a compound having the formula

(52) **U.S. Cl.** **424/85.1**; 424/85.2; 424/85.4; 424/141.1; 424/184.1; 514/2; 514/3; 514/8; 514/11; 514/12; 514/21; 514/56; 514/563; 514/784; 562/457



(58) **Field of Search** 424/85.1, 85.2, 424/85.4, 141.1, 184.1; 514/2, 3, 8, 11, 12, 21, 56, 456, 563, 616, 784; 562/457


or a salt thereof which facilitates the delivery of active agents. Compositions and dosage unit forms comprising the compound of the present invention and at least one active agent, such as a peptide, mucopolysaccharide, carbohydrate, or a lipid, are also provided. Methods of administration and preparation of the compounds and compositions of the invention are provided as well.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2,671,451 A 3/1954 Bolger 128/260

27 Claims, 3 Drawing Sheets



U.S. PATENT DOCUMENTS			
2,868,740 A	1/1959	Luce	260/8
RE24,899 E	11/1960	Green	428/402.2
2,971,916 A	2/1961	Schleicher et al.	252/62.5
3,016,308 A	1/1962	Macaulay	177/37
3,052,655 A	9/1962	Fox et al.	260/78
3,057,344 A	10/1962	Abella et al.	128/2
3,076,790 A	2/1963	Fox et al.	260/78
3,170,802 A	2/1965	Fukushima	99/145
3,190,837 A	6/1965	Brynko et al.	252/316
3,474,777 A	10/1969	Figge et al.	128/2
3,491,093 A	1/1970	Pachter et al.	260/247.5
3,565,559 A	2/1971	Sato	424/37
3,567,650 A	3/1971	Bakan	252/316
3,574,832 A	4/1971	Engel et al.	424/183
3,576,758 A	4/1971	Emrick	252/316
3,687,926 A	8/1972	Arima et al.	260/112.5
3,725,113 A	4/1973	Chang	117/82
3,748,277 A	7/1973	Wagner	252/316
3,794,561 A	2/1974	Matsukawa et al.	195/29 R
3,795,739 A	3/1974	Birkmayer et al.	424/274
3,816,404 A	6/1974	Kablaoui et al.	260/239.3
3,822,348 A	7/1974	Higashi et al.	424/95
3,849,550 A	11/1974	Teitelbaum	424/78
3,933,873 A	1/1976	Love et al.	260/239.3
3,937,668 A	2/1976	Zolle	252/316
3,939,253 A	2/1976	Bodor et al.	424/309
3,956,172 A	5/1976	Saeki et al.	252/316
3,962,416 A	6/1976	Katzen	424/19
3,976,773 A	8/1976	Curran	424/250
4,035,507 A	7/1977	Bodor et al.	424/311
4,048,268 A	9/1977	Ludwig	264/15
4,061,466 A	12/1977	Sjoholm et al.	23/230 B
4,117,801 A	10/1978	Dannelly et al.	118/20
4,147,767 A	4/1979	Yapel	424/22
4,183,849 A	1/1980	Hansen	260/112.7
4,199,561 A	4/1980	Roth et al.	424/32
4,217,370 A	8/1980	Rawlings et al.	426/98
4,238,506 A *	12/1980	Stach et al.	562/442
4,239,635 A	12/1980	Rieder	252/34
4,239,754 A	12/1980	Sache et al.	424/183
4,272,506 A	6/1981	Schwarzberg	424/8
4,289,759 A	9/1981	Heavner et al.	424/177
4,345,588 A	8/1982	Widder et al.	128/1.3
4,348,384 A	9/1982	Horikoshi et al.	424/101
4,351,337 A	9/1982	Sidman	128/260
4,352,883 A	10/1982	Lim	435/178
4,357,259 A	11/1982	Senyei et al.	252/316
4,388,304 A	6/1983	Nyeki et al.	424/177
4,393,192 A	7/1983	Curatolo et al.	528/292
4,402,856 A	9/1983	Schnoring et al.	428/402.22
4,402,968 A	9/1983	Martin	424/273
4,405,598 A	9/1983	Brown	424/45
4,442,090 A	4/1984	Takeya et al.	424/178
4,446,138 A	5/1984	Pack	424/248.57
4,450,150 A	5/1984	Sidman	424/1.1
4,457,907 A	7/1984	Porter	424/7.1
4,460,563 A	7/1984	Calanchi	424/35
4,462,839 A	7/1984	McGinley et al.	106/198
4,462,991 A	7/1984	Higuchi et al.	424/177
4,473,620 A	9/1984	Wu et al.	428/402.24
4,483,807 A	11/1984	Asano	264/22
4,492,684 A	1/1985	Goosen et al.	424/19
4,518,433 A	5/1985	McGinley et al.	106/180
4,590,265 A	5/1986	Bogan et al.	536/63
4,608,278 A	8/1986	Frank	427/213.35
4,613,500 A	9/1986	Suzuki et al.	429/85
4,647,455 A	3/1987	De Bold	424/95
4,666,641 A	5/1987	Fickat et al.	264/4.3
4,671,954 A	6/1987	Goldberg	424/450
4,673,566 A	6/1987	Goosen et al.	424/19
4,683,092 A	7/1987	Tsang	264/4.3
4,690,786 A	9/1987	Ninomiya et al.	264/4.6
4,692,284 A	9/1987	Braden	264/4.3
4,692,433 A	9/1987	Hostetler et al.	514/12
4,703,042 A	10/1987	Bodor	514/56
4,708,952 A	11/1987	Salatinjants	514/158
4,745,161 A	5/1988	Saudek et al.	525/420
4,753,804 A	6/1988	Iaccheri et al.	424/491
4,757,007 A	7/1988	Satoh	435/69
4,757,024 A	7/1988	Roper	436/507
4,766,012 A	7/1988	Shiokari et al.	514/210
4,774,320 A	8/1988	Valenti	427/213.36
4,789,734 A	9/1988	Tagliabue et al.	530/328
4,835,312 A	12/1988	Pierschbacher	530/395
4,837,381 A	5/1989	Itoh et al.	564/205
4,844,904 A	6/1989	Steber et al.	424/502
4,873,087 A	7/1989	Hamaguchi et al.	424/450
4,878,942 A	10/1989	Morishita et al.	424/433
4,886,663 A	11/1989	Motegi et al.	71/109
4,895,725 A	12/1989	Houghten	424/88
4,897,444 A	1/1990	Kantor et al.	424/455
4,900,730 A	1/1990	Brynes et al.	525/54.1
4,908,233 A	2/1990	Miyauchi	514/12
4,919,939 A	3/1990	Takizawa et al.	427/213.35
4,925,673 A	4/1990	Baker	424/493
4,927,928 A	5/1990	Steiner	424/455
4,963,364 A	5/1990	Shroot et al.	544/154
4,976,968 A	10/1990	Fox et al.	424/455
4,983,402 A	12/1990	Steiner	424/491
4,996,292 A	1/1991	Steiner	424/491
5,019,400 A	2/1991	Fox et al.	528/328
5,023,374 A	5/1991	Gombotz et al.	424/497
5,039,481 A	6/1991	Simon	564/152
5,041,291 A	8/1991	Pacifici et al.	422/4
5,055,300 A	8/1991	Bader et al.	424/426
5,066,487 A	10/1991	Gupta	424/409
5,067,961 A	11/1991	Morelle et al.	424/68
5,069,936 A	11/1991	Kelman et al.	623/5
5,077,278 A	12/1991	Yen	427/213.33
5,100,669 A	12/1991	Hafner et al.	514/30
5,100,918 A	3/1992	Hyon et al.	424/426
5,122,367 A	3/1992	Sunshine et al.	514/577
5,126,147 A	6/1992	Ron et al.	424/80
5,137,892 A	6/1992	Silvestri et al.	424/497
5,186,947 A	8/1992	Chu et al.	514/278
5,204,099 A	2/1993	Goettsche et al.	424/638
5,206,384 A	4/1993	Barbier et al.	424/401
5,216,124 A	4/1993	Shibahara et al.	548/537
5,244,653 A	6/1993	Hansen, Jr. et al.	530/317
5,250,236 A	9/1993	Berke et al.	424/70
5,271,934 A	10/1993	Gasco	264/4.4
5,271,961 A	12/1993	Goldberg et al.	424/401
5,278,148 A	12/1993	Mathiowitz et al. ...	427/213.31
5,310,535 A	1/1994	Branca et al.	514/19
5,328,992 A	5/1994	Kruper, Jr. et al.	424/1.53
5,352,461 A	7/1994	Peter et al.	534/116
5,384,133 A	10/1994	Feldstein et al.	424/493
5,389,377 A	1/1995	Boyes et al.	424/501
5,389,379 A	2/1995	Chagnon et al.	424/450
5,401,516 A	2/1995	Dirix et al.	424/451
5,418,010 A	3/1995	Milstein et al.	424/491
5,439,686 A	5/1995	Janda et al.	427/213.31
5,443,841 A	8/1995	Desai et al.	424/451
5,447,728 A	8/1995	Milstein et al.	424/451
5,451,410 A	9/1995	Milstein et al.	424/490
5,474,997 A	9/1995	Milstein et al.	424/490
5,536,813 A	12/1995	Gray et al.	514/252
5,540,939 A	7/1996	Charpenel et al.	530/324
5,541,155 A	7/1996	Milstein et al.	424/491
5,578,323 A	7/1996	Leone-Bay et al.	514/2
	11/1996	Milstein et al.	424/499

5,601,846 A	2/1997	Milstein et al.	424/499	WO	WO 87/04076	7/1987	A61K/45/02
5,629,020 A	5/1997	Leone-Bay et al.	424/489	WO	WO 88/01213	2/1988	B23B/5/16
5,643,957 A *	7/1997	Leone-Bay et al.	514/563	WO	WO 92/19263	12/1992	A61K/39/00
5,650,386 A *	7/1997	Leone-Bay et al.	514/2	WO	WO 93/18754	9/1993	A61K/9/16
5,665,700 A	9/1997	Cho et al.	514/2	WO	WO 93/25583	12/1993	C07K/15/00
5,667,806 A	9/1997	Kantor	424/484	WO	WO 94/11015	5/1994	A61K/37/00
5,693,338 A	12/1997	Milstein	424/451	WO	WO 94/14420	7/1994	A61K/9/16
5,705,529 A	1/1998	Matyus et al.	514/541	WO	WO 94/18950	9/1994	A61K/9/127
5,709,861 A	1/1998	Santiago et al.	424/184.1	WO	WO 94/18997	9/1994	A61K/37/00
5,714,167 A	2/1998	Milstein et al.	424/490	WO	WO 94/21234	9/1994	A61K/7/00
5,750,147 A	5/1998	Kantor	424/491	WO	WO 94/23702	10/1994	A61K/9/16
5,766,633 A	6/1998	Milstein et al.	424/489	WO	WO 94/23767	10/1994	A61L/9/16
5,773,647 A	6/1998	Leone-Bay et al.	562/444	WO	WO 94/24291	10/1994	A61K/39/015
5,776,888 A	7/1998	Leone-Bay et al.	514/2	WO	WO 94/28878	12/1994	A61K/9/14
5,792,451 A	8/1998	Sarubbi et al.	424/85.4	WO	WO 95/11690	5/1995	A61K/37/00
5,804,688 A	9/1998	Leone-Bay et al.	562/444	WO	WO 85/02772	7/1995	A61K/49/00
5,811,127 A	9/1998	Milstein et al.	424/490	WO	WO 95/28838	11/1995	A01N/37/46
5,879,681 A	9/1998	Leone-Bay et al.	424/85.1	WO	WO 95/28920	11/1995	A61K/31/19
5,820,881 A	10/1998	Milstein	424/489	WO	WO 96/12473	5/1996	A61K/9/16
5,824,345 A	10/1998	Milstein	424/489	WO	WO 96/12474	5/1996	A61K/9/16
5,840,340 A	11/1998	Milstein et al.	424/499	WO	WO 96/12475	5/1996	A61K/9/16
5,863,944 A	1/1999	Leone-Bay et al.	514/559	WO	WO 96/21464	7/1996	A61K/39/00
5,866,536 A	2/1999	Leone-Bay et al.	514/2	WO	WO 96/30036	10/1996	A61K/38/00
5,876,710 A	3/1999	Leone-Bay et al.	424/85.1	WO	WO 96/33699	10/1996	A61K/9/16
6,001,347 A *	12/1999	Leone-Bay et al.	424/85.1	WO	WO 96/39835	12/1996	A01N/43/50
				WO	WO 96/40070	12/1996	A61K/9/14
				WO	WO 96/40076	12/1996	A61K/9/16
				WO	WO 97/10197	3/1997	C07C/51/10
				WO	WO 97/31938	9/1997	C07K/5/00
				WO	WO 97/36480	10/1997	A01N/37/12
				WO	WO 97/47270	12/1997	
				WO	WO 97/47288	12/1997	A61K/9/48
				WO	WO 98/49135	5/1998	C07C/231/00
				WO	WO 98/34632	8/1998	A61K/38/00
				WO	WO 98/50341	11/1998	C07C/229/00
				WO	WO 99/16427	4/1999	A61K/9/16

FOREIGN PATENT DOCUMENTS

DE	3 202 255	10/1982	C08L/89/00
DE	3 612 102.9	10/1986	C07K/15/00
EP	0 000 667 A1	2/1979	A61K/9/50
EP	0 036 145 A1	9/1981	A61K/31/62
EP	0 068 314	1/1983	A61K/31/16
EP	0 105 804	4/1984	C12N/15/00
EP	0 130 162 A2	1/1985	B01J/13/02
EP	0 170 540 A1	2/1986	A61K/9/52
EP	226223 A2	6/1987	C07C/103/46
EP	0 342 054 A2	11/1989	A61K/7/06
EP	0 342 056 A2	11/1989	A61K/7/06
EP	0 365 183	4/1990	A61K/31/18
EP	0 366 277	5/1990	A61K/9/107
EP	0 418 642	3/1991	A61K/37/30
EP	0 448 057	9/1991	C12P/21/08
EP	0 452 161	10/1991	A61K/7/48
EP	0 459 795	12/1991	A61K/37/02
EP	0 467 389	1/1992	A61K/9/52
EP	0 490 549 A1	6/1992	A61K/47/12
EP	0 517 211 A1	9/1992	A61K/47/12
EP	0 616 799 A1	9/1994	A61K/7/00
ES	369853	7/1969		
FR	1 351 358	3/1964		
FR	1 468 601	2/1967		
FR	2 133 926	12/1972	A61K/27/00
FR	2 326 934	5/1977	A61K/47/00
FR	2 565 102	12/1985	A61K/9/52
GB	929401	6/1963		
GB	1 075 952	8/1967		
GB	1 236 885	6/1971		
GB	1 567 763	5/1980	A61K/9/22
GB	2 095 994	10/1982	A61K/9/00
IL	71258/2	12/1987		
JP	48-24246	3/1973		
JP	54-66653	* 5/1979		
JP	56-68612	6/1981	A61K/31/19
JP	58-35111	3/1983	A61K/9/66
JP	06-107682	4/1994		
NL	280825	12/1964		
NL	280826	12/1964		
NO	B-146698	11/1982	A61K/37/26
WO	WO 85/00105	1/1985	A61K/9/52
WO	WO 85/00110	1/1985	A61K/47/00
WO	WO 85/00809	2/1985	C07D/233/64

OTHER PUBLICATIONS

- Chen et al. (1975) "Evidence for Hemiacetal Formation", *Biochemistry*, vol. 18, No. 5, pp. 921-925.
- Davis et al. (1983) "Leucinal Inhibits . . .", *Pharmacology Biochemistry Behavior*, vol. 19, pp. 791-794.
- Dose, K. (1974) *Origins of Life*, vol. 5, pp. 239-252.
- Fasman et al. (1964) *Biochemistry*, vol. 3, No. 11, pp. 1665-1674.
- Fox, S.W. et al. (1976) *BioSystems*, vol. 8, pp. 40-44.
- Fox, S.W. et al., *Molecular Evolution and the Origin of Life*, Maxel Decker, New York (1977).
- Fox, S.W. et al. (1968) *Biochim. Biophys. Acta*, vol. 160, pp. 246-249.
- Fox, S.W. (1976) *Origins of Life*, vol. 7, pp. 49-68.
- Fox, S.W. (1980) *Naturwissenschaften*, vol. 67, pp. 378-383.
- Fox, S.W. et al. (1960) *Archives of Biochemistry and Biophysics*, vol. 86, pp. 281-285.
- Fox, S.W. et al. (1974) *Origins of Life*, vol. 5, pp. 227-237.
- Fox, S.W. (1984) *Origins of Life*, vol. 14, pp. 485-488.
- Gol'dovskii, A.M. (1978) *Zhurnal Evolyutsionnoi Biokhimii i Fiziologii*, vol. 14(6), pp. 437-439.
- Gurrieri, S. et al. (1973) *Thermochimica Acta*, vol. 7, pp. 231-239.
- Harada, K. et al. (1979) *BioSystems*, vol. 11, pp. 47-53.
- Harada et al., (1960) *Archives of Biochemistry and Biophysics*, vol. 86, pp. 274-280.
- Hare (1970) *Etude Cenetique De La Polycondensation Thermique D'_x-Amino Acides*, vol. 45, pp. 330-339.
- Heinrich, M.R. et al. (1969) *Archives of Biochemistry and Biophysics*, vol. 130, pp. 441-448.

- Heinz, B. et al. (1981) *BioSystems*, vol. 14, pp. 33–40.
- Hennon, G. et al. (1975) *Biochimie*, vol. 57, pp. 1395–1396.
- Hsu, L.L. et al. (1976) *BioSystems*, vol. 8, pp. 89–101.
- Hsu, L.L. et al. (1971) *Currents in Modern Biology*, vol. 4, pp. 12–25.
- Ishima, Y. et al. (1981), *BioSystems*, vol. 14, pp. 243–251.
- Jackson et al. (1991) “Pharmacological . . .”, *J. Pharm. & Exp. Thera.*, vol. 261, No. 1, pp. 546–552.
- Jungck, J.R. et al. (1973) *Naturwissenschaften*, vol. 60, pp. 425–427.
- Kokufuta, E. et al. (1984) *BioSystems*, vol. 16, pp. 175–181.
- Krampitz, G. et al. (1967) *Naturwissenschaften*, pp. 516–517.
- Krampitz, G. et al. (1968) *Naturwissenschaften*, pp. 345 and 346.
- Krampitz, G. et al. (1966) *Naturwissenschaften*, pp. 7 and 8.
- Lacey, Jr., J.C. et al. (1979) *BioSystems*, vol. 11, pp. 9–17.
- Lacey, Jr., J.C. et al. (1979) *BioSystems*, vol. 11, pp. 1–7.
- Martinez Luque-Romero, M. et al. (1986) *BioSystems*, vol. 19, pp. 267–272.
- Masinovsky, Z. et al. (1989) *BioSystems*, vol. 22, pp. 305–310.
- Matsuno, K. (1982) *BioSystems*, vol. 15, pp. 1–11.
- Matsuno, K. (1984) *BioSystems*, vol. 17, pp. 11–14.
- Matsuno, K. (1981) *BioSystems*, vol. 14, pp. 163–170.
- McAlhaney, W.W. et al. (1976) *BioSystems*, vol. 8, pp. 45–50.
- Melius, P. et al. (1987) *BioSystems*, vol. 20, pp. 213–217.
- Melius, P. et al. (1975) *Bioorganic Chemistry*, vol. 4, pp. 385–391.
- Melius, P. (1979) *BioSystems*, vol. 11, pp. 125–132.
- Miquel, J. et al. (1971) *Currents in Modern Biology*, vol. 3, pp. 299–306.
- Nakashima, T. et al. (1980) *J. Mol. Evol.*, vol. 15, pp. 161–168.
- Nakashima, T. et al. (1981) *BioSystems*, vol. 14, pp. 151–161.
- Novak, V.J.A. (1984) *Origins of Life*, vol. 14, pp. 513–522.
- Olafsson, P.G. et al. (1971) *Polymer Letters*, vol. 9, pp. 521–528.
- Phillips, R.D. et al. (1974) *Int. J. Peptide Protein Res.*, vol. 6, pp. 309–319.
- Przybylski, A.T. et al. (1982) *Die Naturwissenschaften*, vol. 69, pp. 561–563.
- Przybylski, A.T. et al. (1984) *Applied Biochemistry and Biotechnology*, vol. 10, pp. 301–307.
- Przybylski, A.T. (1985) *BioSystems*, vol. 17, pp. 281–288.
- Rohlfing, D.L. (1975) *Origins of Life*, vol. 6, pp. 203–209.
- Rohlfing, D.L. (1970) *Science*, vol. 169, pp. 998–1000.
- Rohlfing, D.L. (1967) *Archives of Biochemistry and Biophysics*, vol. 118, pp. 468–474.
- Rohlfing, D.L. et al. *Catalytic Activities of Thermal Polyanhydro- α -Amino Acids*, pp. 373–418, 1969.
- Rohlfing, D.L. et al. (1976) *BioSystems*, vol. 8, pp. 139–145.
- Ryan, J.W. et al. (1973) *BioSystems*, vol. 5, pp. 115–118.
- Saunders, M.A. et al. (1974) *BioSystems*, vol. 6, pp. 81–92.
- Snyder, W.D. et al. (1975) *BioSystems*, vol. 7, pp. 222–229.
- Sokol, P.E. (1974) *Journal of the American Oil Chemists’ Society*, vol. 52, pp. 101–102.
- Tschager et al. (1988) *Milchwirtschaftliche Berichte*, vol. 95, pp. 79–83.
- Vaughan, G. et al. (1987) *BioSystems*, vol. 20, pp. 219–223.
- Vol’kenshtein, M.V. (1989) *Molekulyarnaya Biologiya*, vol. 23(1), pp. 23–37.
- Waehneltd, T.V. et al. (1968) *Biochim. Biophys. Acta*, vol. 160, pp. 239–245.
- Williams et al. (1991) *J. Biol. Chem.*, vol. 266, No. 8, pp. 5182–5190.
- Yuki, A. et al. (1969) *Biochemical and Biophysical Research Communications*, vol. 36(4), pp. 657–663.
- Zulaski et al. (1983) “New Carboxyalkyl Inhibitors of Brain Enkephalinase”, *J. Med. Chem.*, 26, pp. 60–65.
- (1985) *Chemical Abstracts*, vol. No. 105(1), Abstract No. 12027p.
- (1985) *Chemical Abstracts*, vol. No. 102(6), Abstract No. 50870d.
- Chemical Abstract*, vol. 80(9) Abst. No. 52392a (1974).
- Bergeron, Raymond J., et al. (1994) “Macromolecular Self-Assembly of Diketopiperazine Tetrapeptides”, *Journal of the American Chemical Society*, vol. 116, pp. 8479–8484.
- Bergeron, Raymond J., et al. (1993) “A Comparative Study of the Iron-Clearing Properties of Desferriothiocin Analogues With Desferrioxamine B in a Cebus Monkey Model”, *Blood*, vol. 81, No. 8, pp. 2166–2173.
- Bergeron, Raymond J., et al. (1992) “A Comparison of the Iron-Clearing Properties of 1,2-Dimethyl-3-Hydroxypyrid-4-One, 1,2-Diethyl-3-Hydroxypyrid-4-One, and Deferoxamine”, *Blood*, vol. 79, No. 7, pp. 1882–1890.
- Bergeron, Raymond J., et al. (1991) “Evaluation of Desferriothiocin and Its Synthetic Analogs as Orally Effective Iron Chelators”, *Journal of Medicinal Chemistry*, vol. 34, No. 7, pp. 2072–2078.
- Bergeron, Raymond et al., “A Comparative Evaluation of Iron Clearance Models”, *Annals New York Academy of Sciences*, pp. 278–393, Mar. 13–15, 1990.
- Andriuoli, G., et al. (1990), *Haemostasis* 20 (suppl. 1):154–158.
- Caramazza, I., et al. (1991), *Thrombosis Research* 62:785–789.
- Guarini, S., et al. (1983), *Experientia* 41:350–352.
- Guarini, S., et al. (1985), *Pharmacological Research Communications* 17(8):685–697.
- Dal Pozzo, A., et al. (1989), *Thrombosis Research* 56:119–124.
- Gelb, R., et al (1983), *Lite Sciences* 33(1):83–85.
- Watterberg et al. (1988), *Pediatric Research*, vol. 23, No. 4, part 2, p. 570A, col. 1, abstract No. 2209.
- Bernstein (1985), *Chest* 87(1):68S–73S.
- Damge et al. (1988), *Diabetes* 37:246–251.
- Chemical Abstracts*:83 184360k, (1975).
- Amino, Y., et al., *Chem. Pharm. Bull.* 36(11):4426–4434 (1988).
- Baughman, R.A. et al., *Proc. of the 6th Inter’l. Symp. on Recent Adv. in Drug Delivery Systems, Ctr. for Controlled Chem. Delivery*, University of Utah, Feb. 22–25, 1993, Salt Lake City, UT, pp. 179–180 “Method for Assessing The Stability of Proteinoid Microspheres”.
- Haas, S. et al., “Assessment Of Stability Of Proteinoid Microspheres”, *Proceed. Intern. Symp. Control. Rel. Bioact. Mater.*, 20 (1993), Controlled Release Society, Inc.
- X. Ma, et al., *Proceed. Intern. Symp. Control. Rel. Bioact. Mater.*, 20 (1993), Controlled Release Society, Inc. “In Vitro Mechanistic Investigation of the Proteinoid Microsphere Oral Delivery System”.

- Yen, H.-R H., et al., "Adsorption of Sulforhodamine 101 on Proteinoid Microspheres" *Proceed. Intern. Symp. Control. Rel. Bioact. Mater.*, 20 (1993), Controlled Release Society, Inc.
- Presented at "IBC Rational Drug Design Conference", San Diego, Calif.—Dec. 1994.
- Leone-Bay et al., Presented at "Winter Conference on Medicinal and Bioorganic Chemistry" Steamboat Springs, Colorado—Feb. 1995 "Microsphere Formation and Drug Delivery in a Series of Derivatized Amino Acids".
- Santiago et al., *Pharm. Res.* 11: 1994, p. S-298 "Oral Delivery of Heparin Microspheres made with Modified Amino Acids".
- Leone-Bay et al., *Pharm. Res.* 11: 1994, p. S-121 "Oral Delivery of Heparin using Acylated Amino Acids".
- Sarubbi et al., *Pharm. Res.* 11: 1994, p. S-299 "Oral Calcitonin Delivery using the PODDS Technology".
- Leipold et al., *Pharm. Res.* 11: 1994, p. S-298 "Oral Delivery of Interferon in Rats and Primates".
- Santiago et al., *Pharm. Res.* 11: 1994, p. S-298 "Evaluation in Rats of Vehicles for the Oral Delivery of Low Molecular Weight Heparin".
- X. Ma et al., PDD 7303 *Pharmaceutical Research* 9(10):S-244, 1992 (Oct. Supplement).
- Milstein et al., *Symposia Abstracts*. AAPS Annual Meeting, San Antonio, TX, Nov. 15-19, 1993.
- Santiago et al. "Initial Studies In The Assessment of Proteinoid Microsphere Activity" *Proceed. Intern. Symp. Control. Rel. Bioact. Mater.*, 20 (1993), Controlled Release Society, Inc.
- Santiago et al. "Oral Immunization of Rats with Influenza Virus M Protein (M1) Microspheres" *Proceed. Intern. Symp. Control. Rel. Bioact. Mater.*, 19 (1992), Controlled Release Society, Inc., p. 116-117.
- Santiago et al. "Proteinoid Microspheres For The Oral Delivery of Heparin" *Proceed. Intern. Symp. Control. Rel. Bioact. Mater.*, 19 (1992), Controlled Release Society, Inc. p. 514-515.
- Santiago et al. *American Society for Microbiology* 92nd General Meeting, Abstract of the General Meeting, p. 159, May 26-30, 1992.
- Milstein et al. "Preparation And In Vitro Characterization Of Proteinoid Microspheres" *Proceed Intern. Symp. Control. Rel. Bioact. Mater.*, 19 (1992), Controlled Release Society, Inc. p. 516-517.
- Doris K. Chiappetta, *Eastern Analytical Symposium*, Nov. 17, 1992 "Solutions for Problems in Bioanalysis".
- Elizabeth A. Harris. M.S., *Eastern Analytical Symposium*, Nov. 17, 1992 "Solutions for Problems in Bioanalysis".
- AAPS 6th Ann. Meeting and Expo.*, "Proteinoids—A Novel Drug Delivery System" Nov. 19, 1992, p. 33.
- Milstein et al., "Efficient Oral Delivery Of Monoclonal Antibodies By Proteinoid Encapsulation" *The 1993 Miami Bio/Technology Winter Symposium—Advances in Gene Technology: Protein Engineering and Beyond*, Jan. 17-22, 1993.
- Xinghang Ma, et al. "Stability Study of Durg-loaded Proteinoid Microsphere Formulations during Freeze-drying" *Journal of Drug Targeting*, 1994, vol. 2, pp 9-21.
- Baughman et al., "Screening Candidate Microsphere Formulations By Incubating In Simulated Digestive Fluids" *Proc. of the 6th Intern'l. Sympo. on Recent Advances in Drug Delivery Systems*, Ctr. for Controlled Chem. Delivery, University of Utah, Feb. 22-25, 1993, pp. 181-182.
- Robert O. Dillman, M.D., *Annals of Internal Medicine* 1989:111 pp. 592-600, "Monoclonal Antibodies for Treating Cancer".
- Brendan D. Curti, *Critical Reviews in Oncology/Hematology*, 1993: 14 pp. 29-39 "Physical barriers to drug delivery in tumors".
- V. Hird et al, *Genes and Cancer*, edited by Desmond Carney & Karol Sikora, pp. 183-189, Immunotherapy with Monoclonal Antibodies. (Not dated).
- Michael E. Osband et al., *Immunology Today*, vol. 11, No. 6 1990, pp. 193-195, "Problems in the investigational study and clinical use of cancer immunotherapy".
- William J. Harris, *Tibtech* Feb. 1993 vol. 11, pp. 42-44 "Therapeutic antibodies—the coming of age".
- Thomas A. Waldmann, *Science*, Jun. 21, 1991, 252:1657-1662, "Monoclonal Antibodies in Diagnosis and Therapy".
- Chemical Abstracts*, 76(14):72994u, (1971).
- Chemical Abstracts*, 84(7):44660d, (1975).
- Chemical Abstracts*, 86(16):107529g, (1976).
- Chemical Abstracts*, 112(15):134663h, (1989).
- Chemical Abstracts*, 114(22):214519x, (1990).
- J. Györe et al., *Thermal Analysis*, vol. 2—Proceeding Fourth ICTA Budapest 1974, p. 387-394.
- Chemical Abstracts*, 99(19) 158832b, (1982).
- Derwent Abstracts*, JP 67008622, (1967).
- Journal of Medicinal Chemistry*, vol. 38, No. 21, pp. 4257-4262, (1995), "Microsphere Formation in a Series of Derivatized α -Amino Acids: Properties, Molecular Modeling, and Oral Delivery of Salmon Calcitonin".
- Andrea Leone-Bay et al., *Journal of Medicinal Chemistry*, vol. 38, No. 21, pp. 4263-4269, (1995), "N-Acylated α -Amino Acids as Novel Oral Delivery Agents for Proteins".
- The Extra Pharmacopoeia*, Thirtieth Edition, pp. 325-326, (1993).
- Stephen J. Douglas et al., *Chemistry and Industry*, vol. 22:748-751, 1985.
- C.A. Finch, *Chemistry and Industry*, vol. 22:752-756, 1985.
- John A. Butera et al., *J. Med. Chem.*, vol. 34:3212-3228, 1990.
- Madeline G. Cimini et al., *Ann. Report in Med Chem.*, vol. 27:89-98., 1992.
- Bernadette Earley et al., *Brain Research*, vol. 546:282-286, 1991.
- John W. Ellingboe et al., *J. Med Chem.*, vol. 35:705-716, 1992.
- William C. Lumma et al., *J. Med Chem.*, vol. 30:758-763, 1987.
- Joseph J. Lynch et al., *J. of Pharm. and Exp. Therap.*, vol. 269:541-554, 1994.
- Kiyoshi Matsuno et al., *Brain Research*, vol. 575:315-319, 1992.
- Thomas K. Morgan et al., *J. Med. Chem.*, vol. 33:1091-1097, 1990.
- Hitoshi Oinuma et al., *J. Med Chem.*, vol. 33:903-905, 1990.
- Tadimeti S. Rao et al., *Molecular Pharmacology*, vol. 37:978-982, 1990.
- Asaji Kondo, *Microcapsule Processing and Technology*, pp. 154-165, 1979.
- G. Pastores et al., *J. Liquid Chromatography*, 18(15):3049-3059, 1995.

- D. Sinha et al., *J. Bio. Chem.*, 260(19):10714–10719, 1985.
E. Franssen et al., *J. Med. Chem.*, 35:1246–1259, 1992.
Chemical Abstracts, 99(23):191473h, Dec. 5, 1983.
R. Langer, *Science*, 249:1528, Sep. 28, 1990.
M. Alonso et al., *Vaccine*, 12:299, 1994.
A. Leone-Bay et al., *J. Med. Chem.*, 39:2571–2578, 1996.
R. Thompson, *Biochemistry*, 12:47–51, 1973.
S. Thompson, *J. Med. Chem.*, abstract, 86:174780, 1986.
G. Picciola, *Il Farmaco*, 31:655–664 (1976).
Tanaka et al., *Biophysical Chemistry*, vol. 50 (1994) 47–61.
Degrado et al., *Science*, vol. 243, (1989) 622–628.
Lynce Regan et al., *Science*, vol. 241 (1988), 976–978.
Matouschek et al., *Nature*, vol. 340, (1989) 122–126.
Parker et al., *Peptide Research* 347, vol. 4, No. 6 (1991).
Parker et al., *Peptide Research* 355, vol. 4, No. 6 (1991).
Fedorov et al., *J. Mol. Biol.* (1992) 225, 927–931.
Ptitsyn et al., *Biopolymers*, vol. 22, 15–25 (1983) 15–25.
Ptitsyn et al., *Protein Engineering* vol. 2, No. 6, 443–447, 1989.
J. M. Lehn, *Makromol. Chem., Macromol. Symp.* (1993) vol. 69, 1–17.
Paolo Scrimin, *Chemistry Chimicaoggi* (1989).
J.M. Lehn, *Angew. Chem. Int. Ed. Engl.* 27 (1988) 89–112.
Chemical Abstracts Registry No. 73548–12–6 (Apr. 1991).
Chemical Abstracts Registry No. 70204–54–5 (Apr. 1991).
* cited by examiner

FIG. 1

DRUG DOSE = 0.1mg/kg
DELIVERY AGENT DOSE = 10mg/kg




FIG. 2





FIG. 3

DELIVERY AGENT DOSE : 50 mg / kg
VEHICLE : 25% AQUEOUS PG
DOSE VOLUME : 1ml / kg
PH : 7.3-7.7



COMPOUNDS AND COMPOSITIONS FOR DELIVERING ACTIVE AGENTS

This is a continuation of application Ser. No. 09/305,506 filed May 5, 1999, which in turn is a continuation of application Ser. No. 08/798,031, filed Feb. 6, 1997, now U.S. Pat. No. 6,001,347, which in turn is a continuation of PCT/US96/04580, filed Apr. 1, 1996, published as WO 96/30036, which is a Continuation-In-Part of application Ser. No. 08/414,654, filed Mar. 31, 1995, now U.S. Pat. No. 5,650,386, and which claims benefit to application Ser. No. 60/003,111, filed Sep. 1, 1995, and application Ser. No. 60/017,902, filed Mar. 29, 1996. Each of these prior applications is hereby incorporated herein by reference, in its entirety.

FIELD OF THE INVENTION

The present invention relates to compounds for delivering active agents, and particularly biologically or chemically active agents such as, for example, bioactive peptides and the like. These compounds are used as carriers to facilitate the delivery of a cargo to a target. The carriers are modified amino acids and are well suited to form non-covalent mixtures with biologically-active agents for oral administration to animals. Methods for the preparation and for the administration of such compositions are also disclosed.

BACKGROUND OF THE INVENTION

Conventional means for delivering active agents are often severely limited by biological, chemical, and physical barriers. Typically, these barriers are imposed by the environment through which delivery occurs, the environment of the target for delivery, or the target itself.

Biologically or chemically active agents are particularly vulnerable to such barriers. For example in the delivery to animals of pharmacological and therapeutic agents, barriers are imposed by the body. Examples of physical barriers are the skin and various organ membranes that must be traversed before reaching a target. Chemical barriers include, but are not limited to, pH variations, lipid bi-layers, and degrading enzymes.

These barriers are of particular significance in the design of oral delivery systems. Oral delivery of many biologically or chemically active agents would be the route of choice for administration to animals if not for biological, chemical, and physical barriers such as varying pH in the gastro-intestinal (GI) tract, powerful digestive enzymes, and active agent impermeable gastrointestinal membranes. Among the

numerous agents which are not typically amenable to oral administration are biologically or chemically active peptides, such as calcitonin and insulin; polysaccharides, and in particular mucopolysaccharides including, but not limited to, heparin; heparinoids; antibiotics; and other organic substances. These agents are rapidly rendered ineffective or are destroyed in the gastro-intestinal tract by acid hydrolysis, enzymes, or the like.

Earlier methods for orally administering vulnerable pharmacological agents have relied on the co-administration of adjuvants (e.g., resorcinols and non-ionic surfactants such as polyoxyethylene oleyl ether and n-hexadecylpolyethylene ether) to increase artificially the permeability of the intestinal walls, as well as the co-administration of enzymatic inhibitors (e.g., pancreatic trypsin inhibitors, diisopropylfluorophosphate (DFF) and trasylol) to inhibit enzymatic degradation.

Liposomes have also been described as drug delivery systems for insulin and heparin. See, for example, U.S. Pat. No. 4,239,754; Patel et al. (1976), *FEBS Letters*, Vol. 62, pg. 60; and Hashimoto et al. (1979), *Endocrinology Japan*, Vol. 26, pg. 337.



However, broad spectrum use of such drug delivery systems is precluded because: (1) the systems require toxic amounts of adjuvants or inhibitors; (2) suitable low molecular weight cargos, i.e. active agents, are not available; (3) the systems exhibit poor stability and inadequate shelf life; (4) the systems are difficult to manufacture; (5) the systems fail to protect the active agent (cargo); (6) the systems adversely alter the active agent; or (7) the systems fail to allow or promote absorption of the active agent.

More recently, microspheres of artificial polymers of mixed amino acids (proteinoids) have been used to deliver pharmaceuticals. For example, U.S. Pat. No. 4,925,673 describes drug-containing proteinoid microsphere carriers as well as methods for their preparation and use. These proteinoid microspheres are useful for the delivery of a number of active agents.

There is still a need in the art for simple, inexpensive delivery systems which are easily prepared and which can deliver a broad range of active agents.

SUMMARY OF THE INVENTION









Compositions which are useful in the delivery of active agents are provided. These compositions include at least one active agent, and preferably a biologically or chemically active agent, and at least one of the following compounds I-CXXIII, or salts thereof.



I


II

-continued




-continued


XI




XII




XIII




XIV




XV




XVI



XVII




XVIII




-continued


XIX




XX




XXI




XXII




XXIII




XXIV




XXV




XXVI




-continued




XXVII




XXVIII




XXIX




XXX




XXXI



XXXII




XXXIII




XXXIV

-continued


XXXV



XXXVI




A



Compound	n	m	X
XXXVII	0	0	4-Cl
XXXVIII	3	0	H
XXXIX	3	1	4-CH ₃
XL	3	1	2-F
XLI	3	1	2-CH ₃
XLII	3	0	3-CF ₃
XLIII	3	4	H
XLIV	3	0	3-Cl
XLV	3	0	3-F
XLVI	3	0	3-CH ₃
XLVII	0	0	2-CF ₃
XLVIII	1	2	H
XLIX	3	2	2-F
L	3	0	3,4-OCH ₂ O—
LI	3	0	2-COOH
LII	1	0	2-OH
LIII	3	0	2,6-dihydroxy
LIV	2	0	2-OH
LV	0	0	2,4-difluoro
LVI	2	0	2,6-dihydroxy
LVII	0	0	4-CF ₃
LVIII	3	0	3-NMe ₂
LIX	2	0	3-NMe ₂
LX	3	0	2,6-dimethyl
LXI	3	0	2-NO ₂
LXII	3	0	2-CF ₃
LXIII	3	0	4-n-Pr
LXIV	3	0	2-NH ₂
LXV	3	0	2-OCH ₃
LXVI	3	0	3-NO ₂
LXVII	3	0	3-NH ₂
LXVIII	2	0	2-NO ₂
LXIX	2	0	2-NH ₂
LXX	3	0	2-OCF ₃
LXXI	2	0	2-OCH ₃
LXXII	2	0	2-OCF ₃

B




Compound	n	X
LXXIII	3	4-CF ₃
LXXIV	1	2-F
LXXV	1	4-CF ₃
LXXVI	3	3,4-dimethoxy

-continued


LXXVII	0	3-OCH ₃
LXXVIII	3	3-OCH ₃
LXXIX	3	2,6-difluoro
LXXX	3	4-CH ₃
LXXXI	1	4-OCH ₃
LXXXII	2	2-F
LXXXIII	0	2-F
LXXXIV	2	4-OCH ₃
LXXXV	0	2-OCH ₃
LXXXVI	2	2-OCH ₃
LXXXVII	0	4-CF ₃
LXXXVIII	3	3-F
LXXXIX	3	2-OCH ₃

C




Compound	n	m	X
XC	3	0	2-carboxycyclohexyl
XCI	3	3	cyclohexyl
XCII	3	0	2-adamantyl
XCIII	3	0	1-morpholino

D




Compound	m
XCIV	0
XCV	3

E










Compound	X
XCVI	OH
XCVII	=O

F




Compound	n
XCVIII	0
XCIX	2

-continued




-continued


CVII




CVIII




CIX



CX




G



Compound	n	m	X
CXI	6	0	2-OH
CXII	7	3	H
CXIII	7	0	2-I
CXIV	7	0	2-Br
CXV	7	0	3-NO ₂
CXVI	7	0	3-N(CH ₃) ₂
CXVII	7	0	2-NO ₂
CXVIII	7	0	4-NO ₂
CXIX	9	0	2-OH

H



Compound	X
CXX	1-morpholino
CXXI	O-t-Butyl
CXXII	CH(CH ₂ Ph)NC(O)O-t-Bu
CXXIII	2-hydroxyphenyl

It has been discovered that organic acid compounds, and their salts, having an aromatic amide group, having a hydroxy group substituted in the ortho position on the aromatic ring, and a lipophilic chain with from about 4 carbon atoms to about 20 atoms in the chain are useful as carriers for the delivery of active agents. In a preferred form the lipophilic chain can have from 5 to 20 carbon atoms.

Compositions comprising the carrier compounds discussed above and active agents have been shown effective in delivering active agents to selected biological systems. These compositions include at least one active agent which is preferably a biologically or chemically active agent, and at least one carrier compound having the formula



wherein Ar is a substituted or unsubstituted phenyl or naphthyl;

R⁷ is selected from the group consisting of C₄ to C₂₀ alkyl, C₄ to C₂₀ alkenyl, phenyl, naphthyl, (C₁ to C₁₀ alkyl) phenyl, (C₁ to C₁₀ alkenyl) phenyl, (C₁ to C₁₀ alkyl) naphthyl, (C₁ to C₁₀ alkenyl) naphthyl, phenyl (C₁ to C₁₀ alkyl), phenyl (C₁ to C₁₀ alkenyl), naphthyl (C₁ to C₁₀ alkyl), and naphthyl (C₁ to C₁₀ alkenyl);

R⁸ is selected from the group consisting of hydrogen, C₁ to C₄ alkyl, C₁ to C₄ alkenyl, hydroxy, and C₁ to C₄ alkoxy;

R⁷ is optionally substituted with C₁ to C₄ alkyl, C₁ to C₄ alkenyl, C₁ to C₄ alkoxy, —OH, —SH and —CO₂R⁹ or any combination thereof;


R⁹ is hydrogen, C₁ to C₄ alkyl or C₁ to C₄ alkenyl;

R⁷ is optionally interrupted by oxygen, nitrogen, sulfur or any combination thereof;

With the proviso that the compounds are not substituted with an amino group in the position alpha to the acid group, or salts thereof.

The preferred R⁶ groups are of C₄ to C₂₀ alkyl and C₄ to C₂₀ alkenyl. The most preferred R⁶ groups are C₅ to C₂₀ alkyl and C₅ to C₂₀ alkenyl.

A preferred carrier compound can have the formula




wherein R⁷ is defined above.

Further contemplated by the present invention are dosage unit forms that include these compositions.

Also contemplated is a method for preparing these compositions which comprises mixing at least one active agent with at least one compound as described above, and optionally, a dosing vehicle.

In an alternative embodiment, these non-toxic compounds are orally administered to animals as part of a delivery system by blending or mixing the compounds with an active agent prior to administration.

Further provided is a method for the preparation of a compound having the formula



Wherein Y is



or SO₂;


R¹ is C₃-C₂₄ alkyl, C₂-C₂₀ alkenyl, C₂-C₂₀ alkyne, cycloalkyl, or aromatic;

R² is hydrogen, C₁-C₄ alkyl, or C₂-C₄ alkenyl; and

R³ is C₁-C₇ alkyl, C₃-C₁₀ cycloalkyl, aryl, thienyl, pyrrolo, or pyridyl, where R³ is optionally substituted by one or more C₁-C₅ alkyl group, C₂-C₄ alkenyl group, F, Cl, OH, SO₂, COOH, or SO₃H;

said method comprising

(a) racking in water and the presence of a base, a compound having the formula



with a compound having the formula

R³-Y-X, wherein

Y, R¹, R², and R³ are as above and X is a leaving group.

DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graphic illustration of the results of subcutaneous injection of rhGH composition in rats.

FIG. 2 is a graphic illustration of the results of Sublingual (SL), intranasal (IN), and intracolonic (IC) dosing of rhGH in rats.

FIG. 3 is a graphic illustration of the results of intracolonic dosing of delivery of heparin with compound XXXI carrier.

DETAILED DESCRIPTION OF THE INVENTION

The specific compositions of the present invention include an active agent and a modified amino acid. These compositions may be used to deliver various active agents through various biological, chemical, and physical barriers and are particularly suited for delivering active agents which are subject to environmental degradation. These compositions of the subject invention are particularly useful for delivering or administering biologically or chemically active agents to any animals such as birds; mammals, such as primates and particularly humans; and insects.

Other advantages of the present invention include the use of easy to prepare, inexpensive raw materials. The compositions and the formulation methods of the present invention

are cost effective, simple to perform, and amenable to industrial scale up for commercial production.

Subcutaneous, sublingual, and intranasal coadministration of an active agent, such as recombinant human growth hormone (rhGH), and the delivery agents, and particularly proteins, described herein results in an increased bioavailability of the active agent compared to administration of the active agent alone. A similar result is obtained by coadministration of salmon calcitonin with the delivery agents, in rats. Data supporting these findings are presented in the examples.

Active Agents

Active agents suitable for use in the present invention include biologically or chemically active agents, chemically active agents, including, but not limited to, fragrances, as well as other active agents such as, for example, cosmetics.

Biologically or chemically active agents include, but are not limited to, pesticides, pharmacological agents, and therapeutic agents. For example, biologically or chemically active agents suitable for use in the present invention include, but are not limited to, peptides, and particularly small peptides; hormones, and particularly hormones which by themselves do not or only a fraction of the administered dose passes through the gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastrointestinal tract; polysaccharides, and particularly mixtures of muco-polysaccharides; carbohydrates; lipids; or any combination thereof. Further examples include, but are not limited to, human growth hormones; bovine growth hormones; growth releasing hormones; growth hormone-releasing hormones; interferons; interleukin-1; interleukin-II; insulin; heparin, and particularly low molecular weight heparin; calcitonin; erythropoietin; atrial natriuretic factor; antigens; monoclonal antibodies; somatostatin; adrenocorticotropin, gonadotropin releasing hormone; oxytocin; vasopressin; cromolyn sodium (sodium or disodium chromoglycate); vancomycin; desferrioxamine (DFO); parathyroid hormone anti-microbials, including, but not limited to anti-fungal agents; or any combination thereof.

Modified Amino Acids

The terms modified amino acid, modified poly amino acid, and modified peptide are meant to include amino acids which have been modified, or poly amino acids and peptides in which at least one amino acid has been modified, by acylating or sulfonating at least one free amine group with an acylating or sulfonating agent which reacts with at least one of the free amine groups present.

Amino acids, poly amino acids, and peptides, in modified form, may be used to deliver active agents including, but not limited to, biologically or chemically active agents such as for example, pharmacological and therapeutic agents.


An amino acid is any carboxylic acid having at least one free amine group and includes naturally occurring and synthetic amino acids.

Poly amino acids are either peptides or two or more amino acids linked by a bond formed by other groups which can be linked, e.g. an ester, anhydride, or an anhydride linkage.

Peptides are two or more amino acids joined by a peptide bond. Peptides can vary in length from dipeptides with two amino acids to poly peptides with several hundred amino acids. See *Chambers Biological Dictionary*, editor Peter M. B. Walker, Cambridge, England: Chambers Cambridge, 1989, page 215. Special mention is made of di-peptides, tri-peptides, tetra-peptides, and penta-peptides.

Although compounds I-CXXIII above have been found to act as carriers for the oral delivery of biologically or chemically active agents, special mention is made of compounds I-XXXI above.

Modified amino acids are typically prepared by modifying the amino acid or an ester thereof. Many of these compounds are prepared by acylation or sulfonation with agents having the formula



wherein: R^4 is the appropriate radical to yield the modification indicated in the final product,

Y is



or SO_2 , and X is a leaving group. Typical leaving groups include, but are not limited to, halogens such as, for example, chlorine, bromine, and iodine. Additionally, the corresponding anhydrides are modifying agents.

Many of the compounds of the present invention can be readily prepared from amino acids by methods within the skill of those in the art based upon the present disclosure. For example, compounds I-VII are derived from aminobutyric acid; Compounds VIII-X and XXXII-XXXV are derived from aminocaproic acid; and Compounds XI-XXVI and XXXVI are derived from aminocaprylic acid. For example, the modified amino acid compounds above may be prepared by reacting the single amino acid with the appropriate modifying agent which reacts with free amino moiety present in the amino acids to form amides. Protecting groups may be used to avoid unwanted side reactions as would be known to those skilled in the art.

The amino acid can be dissolved in aqueous alkaline solution of a metal hydroxide, e.g., sodium or potassium hydroxide, and heated at a temperature ranging between about 5°C . and about 70°C ., preferably between about 10°C . and about 40°C ., for a period ranging between about 1 hour and about 4 hours, preferably about 2.5 hours. The amount of alkali employed per equivalent of NH_2 groups in the amino acid generally ranges between about 1.25 and about 3 mmole, preferably between about 1.5 and about 2.25 mmole per equivalent of NH_2 . The pH of the solution generally ranges between about 8 and about 13, preferably ranging between about 10 and about 12.

Thereafter, the appropriate amino modifying agent is added to the amino acid solution while stirring. The temperature of the mixture is maintained at a temperature generally ranging between about 5°C . and about 70°C ., preferably between about 10°C . and about 40°C ., for a period ranging between about 1 and about 4 hours. The amount of amino modifying agent employed in relation to the quantity of amino acid is based on the moles of total free NH_2 in the amino acid. In general, the amino modifying agent is employed in an amount ranging between about 0.5 and about 2.5 mole equivalents, preferably between about 0.75 and about 1.25 equivalents, per molar equivalent of total NH_2 group in the amino acid.

The reaction is quenched by adjusting the pH of the mixture with a suitable acid, e.g., concentrated hydrochloric acid, until the pH reaches between about 2 and about 3. The mixture separates on standing at room temperature to form a transparent upper layer and a white or off-white precipitate. The upper layer is discarded, and the modified amino acid is collected from the lower layer by filtration or decantation. The crude modified amino acid is then dissolved in water at a pH ranging between about 9 and about 13, preferably between about 11 and about 13. Insoluble

23


materials are removed by filtration and the filtrate is dried in vacuo. The yield of modified amino acid generally ranges between about 30 and about 60%, and usually about 45%.

If desired, amino acid esters, such as, for example benzyl, methyl, or ethyl esters of amino acid compounds, may be used to prepare the modified amino acids of the invention. The amino acid ester, dissolved in a suitable organic solvent such as dimethylformamide, pyridine, or tetrahydrofuran is reacted with the appropriate amino modifying agent at a temperature ranging between about 5° C. and about 70° C., preferably about 25° C., for a period ranging between about 7 and about 24 hours. The amount of amino modifying agent used relative to the amino acid ester is the same as described above for amino acids. This reaction may be carried out with or without a base such as, for example, triethylamine or diisopropylethylamine.

Thereafter, the reaction solvent is removed under negative pressure and the ester functionality is removed by hydrolyzing the modified amino acid ester with a suitable alkaline solution, e.g. 1 N sodium hydroxide, at a temperature ranging between about 50° C. and about 80° C., preferably about 70° C., for a period of time sufficient to hydrolyze off the ester group and form the modified amino acid having a free carboxyl group. The hydrolysis mixture is then cooled to room temperature and acidified, e.g. aqueous 25% hydrochloric acid solution, to a pH ranging between about 2 and about 2.5. The modified amino acid precipitates out of solution and is recovered by conventional means such as filtration or decantation. Benzyl esters may be removed by hydrogenation in an organic solvent using a transition metal catalyst.

The modified amino acid may be purified by recrystallization or by fractionation on solid column supports. Suitable recrystallization solvent systems include acetonitrile, methanol and tetrahydrofuran. Fractionation may be performed on a suitable solid column supports such as alumina, using methanol-in-propanol mixtures as the mobile phase; reverse phase column supports using trifluoroacetic acid/acetonitrile mixtures as the mobile phase; and ion exchange chromatography using water as the mobile phase. When anion exchange chromatography is performed, preferably a subsequent 0–500 mM sodium chloride gradient is employed.

In an alternate method modified amino acids having the formula



CXXIV

wherein Y is



or SO₂;


R¹ is C₃–C₂₄ alkyl, C₂–C₂₀ alkenyl, C₂–C₂₀ alkyne, cycloalkyl, or aromatic;

R² is hydrogen, C₁–C₄ alkyl, or C₂–C₄ alkenyl; and

R³ is C₁–C₇ alkyl, C₃–C₁₀ cycloalkyl, aryl, thienyl, pyrrolo, or pyridyl, where R³ is optionally substituted by one or more C₁–C₅ alkyl group, C₂–C₄ alkenyl group, F, Cl, OH, SO₂, COOH or, SO₃H; may be prepared by


24

(a) reacting in water and the presence of a base a compound having the formula



CXXV

with a compound having the formula




Y, R¹, R², and R³ are as above and X is a leaving group.

Compound CXXV can be prepared, for example by the method described in Olah et al., *Synthesis*, 537–538 (1979).

Compound XXXI was prepared as described in Scheme I from 10-undecen-1-ol, 1, by a three step procedure in an overall yield of 31%. Alkylation of phthalimide with alkanol, 1, under Mitsunobu conditions, followed by reaction with hydrazine gave 1-aminoundec-10-ene, 2, in 66% yield. The amine was derivatized with O-acetyl'salicyloyl chloride and the resulting alkene, 3, was oxidized to the acid using potassium permanganate. Removal of the acetate, followed by acid precipitation provided compound XXXI in 47% yield based on amine 2.

Scheme I



Delivery Systems

The compositions of the present invention may include one or more active agents.

In one embodiment, compounds I–CXXIII or poly amino acids or peptides that include at least one of these compounds may be used directly as a delivery carrier by simply mixing one or more compound, poly amino acid or peptide with the active agent prior to administration.

In an alternative embodiment, the compounds, poly amino acids, or peptide may be used to form microspheres containing the active agent. These compounds, poly amino acids, or peptides are particularly useful for the oral administration of certain biologically-active agents, e.g., small peptide hormones, which, by themselves, do not pass or only a fraction of the administered dose passes through the

gastro-intestinal mucosa and/or are susceptible to chemical cleavage by acids and enzymes in the gastrointestinal tract.

If the modified amino acids, poly amino acids, or peptides are to be converted into microspheres, the mixture is optionally heated to a temperature ranging between about 20 and about 50° C., preferably about 40° C., until the modified amino acid(s) dissolve. The final solution contains between about 1 mg and to about 2000 mg of compound, poly amino acid, or peptide per mL of solution, preferably between about 1 and about 500 mg per mL. The concentration of active agent in the final solution varies and is dependent on the required dosage for treatment. When necessary, the exact concentration can be determined by, for example, reverse phase HPLC analysis.

When the compounds, poly amino acids, or peptides are used to prepare microspheres, another useful procedure is as follows: Compounds, poly amino acids, or peptides are dissolved in deionized water at a concentration ranging between about 75 and about 200 mg/ml, preferably about 100 mg/ml at a temperature between about 25° C. and about 60° C. preferably about 40° C. Particulate matter remaining in the solution may be removed by conventional means such as filtration.

Thereafter, the compound, poly amino acid, or peptide solution, maintained at a temperature of about 40° C., is mixed 1:1 (V/V) with an aqueous acid solution (also at about 40° C.) having an acid concentration ranging between about 0.05 N and about 2 N, preferably about 1.7 N. The resulting mixture is further incubated at 40° C. for a period of time effective for microsphere formation, as observed by light microscopy. In practicing this invention, the preferred order of addition is to add the compound, poly amino acid, or peptide solution to the aqueous acid solution.

Suitable acids for microsphere formation include any acid which does not

- (a) adversely effect the modified amino acids, poly amino acids, or peptides e.g., initiate or propagate chemical decomposition;
- (b) interfere with microsphere formation;
- (c) interfere with microsphere incorporation of the active agent cargo; and
- (d) adversely interact with the active agent cargo.

Preferred acids for use in this aspect include acetic acid, citric acid, hydrochloric acid, phosphoric acid, malic acid and maleic acid.

A microsphere stabilizing additive may be incorporated into the aqueous acid solution or into the compound or cargo solution prior to the microsphere formation process. With some active agents the presence of such additives promotes the stability and/or dispersibility of the microspheres in solution.

The stabilizing additives may be employed at a concentration ranging between about 0.1 and 5% (w/v), preferably about 0.50% (w/v). Suitable, but non-limiting, examples of microsphere stabilizing additives include gum acacia, gelatin, methyl cellulose, polyethylene glycol, polypropylene glycol, carboxylic acids and salts thereof, and polylysine. The preferred stabilizing additives are gum acacia, gelatin and methyl cellulose.

Under the above conditions, the compound molecules, poly amino acids, or peptides form hollow or solid matrix type microspheres wherein the cargo is distributed in a carrier matrix or capsule type microspheres encapsulating liquid or solid cargo. If the compound, poly amino acid, or peptide microspheres are formed in the presence of a soluble material, e.g., a pharmaceutical agent in the aforementioned aqueous acid solution, this material will be encapsulated

within the microspheres. In this way, one can encapsulate pharmacologically active materials such as peptides, proteins, and polysaccharides as well as charged organic molecules, e.g., antimicrobial agents, which normally have poor bioavailability by the oral route. The amount of pharmaceutical agent which may be incorporated by the microsphere is dependent on a number of factors which include the concentration of agent in the solution, as well as the affinity of the cargo for the carrier. The compound, poly amino acid, or peptide microspheres do not alter the physiological and biological properties of the active agent. Furthermore, the encapsulation process does not alter the pharmacological properties of the active agent. Any pharmacological agent can be incorporated within the microspheres. The system is particularly advantageous for delivering chemical or biological agents which otherwise would be destroyed or rendered less effective by conditions encountered within the body of the animal to which it is administered, before the microsphere reaches its target zone (i.e., the area in which the contents of the microsphere are to be released) and for delivering pharmacological agents which are poorly absorbed in the gastrointestinal tract. The target zones can vary depending upon the drug employed.

The particle size of the microsphere plays an important role in determining release of the active agent in the targeted area of the gastro-intestinal tract. The preferred microspheres have diameters between about ≤ 0.1 microns and about 10 microns, preferably between about 0.5 microns and about 5 microns. The microspheres are sufficiently small to release effectively the active agent at the targeted area within the gastrointestinal tract such as, for example, between the stomach and the jejunum. Small microspheres can also be administered parenterally by being suspended in an appropriate carrier fluid (e.g., isotonic saline) and injected directly into the circulatory system, intramuscularly or subcutaneously. The mode of administration selected will vary, of course, depending upon the requirement of the active agent being administered. Large amino acid microspheres (>50 microns) tend to be less effective as oral delivery systems.

The size of the microspheres formed by contacting compounds, poly amino acids, or peptides with water or an aqueous solution containing active agents can be controlled by manipulating a variety of physical or chemical parameters, such as the pH, osmolarity or ionic strength of the encapsulating solution, size of the ions in solution and by the choice of acid used in the encapsulating process.

The administration mixtures are prepared by mixing an aqueous solution of the carrier with an aqueous solution of the active ingredient, just prior to administration. Alternatively, the carrier and the biologically or chemically active ingredient can be admixed during the manufacturing process. The solutions may optionally contain additives such as phosphate buffer salts, citric acid, acetic acid, gelatin, and gum acacia.

Stabilizing additives may be incorporated into the carrier solution. With some drugs, the presence of such additives promotes the stability and dispersibility of the agent in solution.

The stabilizing additives may be employed at a concentration ranging between about 0.1 and 5% (W/V), preferably about 0.5% (W/V). Suitable, but non-limiting, examples of stabilizing additives include gum acacia, gelatin, methyl cellulose, polyethylene glycol, carboxylic acids and salts thereof, and polylysine. The preferred stabilizing additives are gum acacia, gelatin and methyl cellulose.

The amount of active agent is an amount effective to accomplish the purpose of the particular active agent. The

amount in the composition typically is a pharmacologically or biologically effective amount. However, the amount can be less than a pharmacologically or biologically effective amount when the composition is used in a dosage unit form, such as a capsule, a tablet or a liquid, because the dosage unit form may contain a multiplicity of carrier/biologically or chemically active agent compositions or may contain a divided pharmacologically or biologically effective amount. The total effective amounts can then be administered in cumulative units containing, in total, pharmacologically or biologically or chemically active amounts of biologically or pharmacologically active agent.



The total amount of active agent, and particularly biologically or chemically active agent, to be used can be determined by those skilled in the art. However, it has surprisingly been found that with some biologically or chemically active agents, the use of the presently disclosed carriers provides extremely efficient delivery, particularly in oral, intranasal, sublingual, intraduodenal, or subcutaneous systems. Therefore, lower amounts of biologically or chemically active agent than those used in prior dosage unit forms or delivery systems can be administered to the subject, while still achieving the same blood levels and therapeutic effects.

The amount of carrier in the present composition is a delivery effective amount and can be determined for any particular carrier or biologically or chemically active agent by methods known to those skilled in the art.


Dosage unit forms can also include any of excipients; diluents; disintegrants; lubricants; plasticizers; colorants; and dosing vehicles, including, but not limited to water, 1,2-propane diol, ethanol, olive oil, or any combination thereof.

Administration of the present compositions or dosage unit forms preferably is oral or by intraduodenal injection.

The delivery compositions of the present invention may also include one or more enzyme inhibitors. Such enzyme inhibitors include, but are not limited to, compounds such as actinonin or epiactinonin and derivatives thereof. These compounds have the formulas below:



Derivatives of these compounds are disclosed in U.S. Pat. No. 5,206,384. Actinonin derivatives have the formula:



15 wherein R⁵ is sulfoxymethyl or carboxyl or a substituted carboxy group selected from carboxamide, hydroxyaminocarbonyl and alkoxy carbonyl groups; and R⁶ is hydroxyl, alkoxy, hydroxyamino or sulfoxyamino group. Other enzyme inhibitors include, but are not limited to, aprotinin (Trasylol) and Bowman-Birk inhibitor.

20 The compounds and compositions of the subject invention are useful for administering biologically or chemically active agents to any animals such as birds; mammals, such as primates and particularly humans; and insects. The system is particularly advantageous for delivering chemically or biologically or chemically active agents which would otherwise be destroyed or rendered less effective by conditions encountered before the active agent its target zone (i.e. the area in which the active agent of the delivery composition are to be released) and within the body of the animal to which they are administered. Particularly, the compounds and composition of the present invention are useful in orally administering active agents, especially those which are not ordinarily orally deliverable.

35

DESCRIPTION OF THE PREFERRED EMBODIMENTS

The following examples illustrate the invention without limitation. All parts are given by weight unless otherwise indicated.

EXAMPLE 1

45 Compound XIX was prepared as follows:

A 3 L three-neck round bottom flask was fitted with an overhead mechanical stirrer and a thermometer, and the flask was cooled in an ice-bath. A solution of 8-aminocaprylic acid (100.0 g, 0.65 moles) in 2 M aqueous sodium hydroxide (1.4 L) was charged into the round bottom flask. The temperature of the solution was kept at about 5° C., and O-acetylsalicyloyl chloride (198.6 g, 0.76 moles, 1.2 equiv.) was added portionwise over 7 hours. The mixture was stirred at 5° C. for 12 hours to yield a yellow homogenous solution.

50 The solution was acidified with 1 M hydrochloric acid to pH 6.8 and was extracted with ethyl acetate (2×600 mL). The pH of the aqueous layer was readjusted to 6.3 and was further extracted with ethyl acetate (2×600 mL). The organic layers were combined, dried over anhydrous sodium sulfate, filtered, and evaporated under reduced pressure. The residue was redissolved in a minimum volume of 2 M aqueous sodium hydroxide, and the pH of the solution was between 9.5 and 10. The mixture was acidified with stirring with 1 M hydrochloric acid to pH of about 6.2, and a solid was formed. The solid was filtered, washed with water (3×300 mL), and recrystallized from 55% methanol/water (v/v) to yield Compound XVIII as an off-white solid (99.7 g, 57%).

65

Properties are listed below.

Mp 116–117° C. ¹H NMR (300 MHz, DMSO-d₆) δ: 12.70 (1H; br s), 11.95 (1H, br s) 8.81 (1H, t), 7.82 (1H, m), 7.38 (1H, m), 6.84 (2H, m), 2.36 (2H, q), 2.18 (2H, t), 1.50 (4H, br m), 1.28 (6H, m), Anal. Calcd for C₁₅H₂₁NO₄: C, 64.50; H, 7.58; N, 5.02. Found: C, 64.26; H, 7.81; N, 4.93.

Similar procedures were used to prepare Compounds I, II, III, IV, VI, IX, X, XI, XII, XIII, XIV, XX, XXI, XXIII, XXVII, XXVIII, XXXIII, and XXXIV.

Properties are listed below.

Compound I:	¹ H NMR (300MHz, D ₂ O): δ 1.5 (2H, m) 2.0 (2H, t) 2.3 (2H,t) 7.5 (2H, t) 7.6 (1H, m) 7.3 (2H, m)
Compound II:	¹ H NMR (300MHz, D ₂ O): δ 1.4 (8H, m) 1.7 (6H, m) 2.1 (2H,t) 1.25 (1H, m) 3.05 (2H, t)
Compound III:	¹ H NMR (300MHz, DMSO-d ₆): δ 0.7 (3H, m) 0.9 (2H, m) 1.1 (3H,q) 1.6.(5H, m) 1.75 (2H, q) 2.1 (2H, t) 3.0 (2h, q) 7.9 (1H, m)
Compound IV:	Anal. Calcd for C ₁₁ H ₁₃ NO ₄ : C, 59.9, H, 5.87, N, 6.27 Found: C, 58.89, H, 5.85, N, 6.07. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.8 (2H, m) 2.3 (2H, t) 3.1 (2H,q) 6.9 (2H, t) 7.4 (1H, t) 7.8 (1h, d) 8.85 (1H, t) 12.0 (1H, s) 12.15 (1H, s)
Compound VI:	¹ H NMR (300MHz, D ₂ O): δ 0.8 (2H, m) 1.1 (4H, m) 1.4 (2H,q) 1.6 (7H, m) 2.15 (4H, m) 3.1 (2H, t)
Compound IX:	¹ H NMR (300MHz, DMSO-d ₆): δ 0.9 (q, 3H), 1.2 (m, 7H), 1.3 (q, 2H), 1.5 (q, 3H), 1.9 (d, 2H), 2.0 (d, 1H), 2.2 (t, 2H), 3.0 (q, 3H), 7.7 (s, 1H)
Compound X:	¹ H NMR (300MHz, DMSO-d ₆): δ 0.7(d, 2H), 0.9 (dd, 1H), 1.2–1.3 (m, 7H), 1.5 (q, 3H), 1.6–1.8 (m, 5H), 2.15 (t, 2H), 3.0 (m, 3H), 7.5 (s, 1H), 12.0 (s, 1H)
Compound XI:	Anal. Calcd for C ₁₅ H ₂₀ NO ₃ Cl: C, 60.48, H, 6.78, N, 4.70 Found: C, 60.4, H, 6.68, N, 4.53. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.28 (m, 6H) 1.48 (m, 4H) 2.19 (t, 2H) 3.19 (qt, 2H), 7.323–7.48 (m, 4H), 8.39(t, H), 12.09 (s, 1H)
Compound XII:	Anal. Calcd for C ₁₇ H ₂₂ NO ₃ : C, 66.42, H, 7.23, N, 4.56 Found: C, 65.80, H, 7.17, N, 4.14. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.25 (m, 6H) 1.43–1.49 (m, 4H) 2.18 (t, 2H) 3.15 (qt, 2H), 6.72 (d, 1H), 7.21–7.26 (m, 2H), 7.39 (t, 1H), 7.48 (d, 1H), 7.65 (t, 1H), 8.21 (t, 1H)
Compound XIII:	Anal. Calcd for C ₁₅ H ₁₉ NO ₃ : C, 60.18, H, 6.41, N, 4.67 Found: C, 60.26, H, 6.53, N, 4.61. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.28 (m, 6H), 1.45–1.52 (m, 4H), 2.19 (t, 2H), 2.22 (qt, 2H), 7.13 (m, 2H), 7.43–7.53 (m, 1H), 8.67 (t, 1H)12.03 (s, 1H)
Compound XIV:	Anal. Calcd for C ₁₄ H ₂₀ N ₂ O ₃ : • 0.66 H ₂ O: C, 63.04, H, 7.91, N, 10.34 Found: C, 63.21, 7.59, 10.53 ¹ H NMR (300MHz, DMSO-d ₆): δ 1.22–12.8 (m, 6H), 1.48–1.50 (m, 4H), 2.18 (t, 2H), 3.24 (qt, 2H), 7.48 (m, 1H), 8.15 (d, 1H), 8.63–8.69 (m, 2H), 8.97 (d, 1H)
Compound XX:	Anal. Calcd for C ₁₅ H ₂₀ NO ₃ F: C, 60.09, H, 7.19, N, 4.98 Found: C, 63.82, H, 7.23, N, 4.94. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.28 (m, 6H) 1.49 (m, 4H) 2.19 (t, 2H) 3.23 (qt, 2H), 7.24–7.30 (m, 2H), 7.49–7.60 (m, 2H), 11.99 (s, 1H)
Compound XXI:	Anal. Calcd for C ₁₇ H ₂₃ NO ₄ : C, 66.85, H, 7.61, N, 4.58 Found: C, 66.81, R, 7.69, N, 4.37. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.26 (m, 6H) 1.42–1.50 (m, 4) 2.18 (t, 2H) 3.13 (qt, 2H), 6.63 (d, 1H), 6.80 (t, 1H), 6.86 (d, 1H), 7.15 (t, 1H)1 7.39 (d, 1H), 7.60 (d, 1H), 8.03 (t, 1H), 9.95 (s, 1H), 12.12 (s, 1H)
Compound XXIII:	Anal. Calcd for C ₁₅ H ₂₇ NO ₃ : C, 66.86, H, 10.22, N, 5.19 Found: C, 66.92, H, 10.72,

-continued

5	N, 5.14. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.56–1.34 (m, 13H) 1.46 (t, 2H) 1.60–1.68 (m, 5H), 2.04 (t, 1H), 2.17 (t, 2H), 2.97 (qt, 2H), 7.62 (t, 1H), 11.98 (s, 1H)
10	Compound XXVII: Anal. Calcd for C ₁₈ H ₂₇ NO ₄ : C, 67.25, H, 8.48, N, 4.36 Found: C, 67.23, H, 8.57, N, 4.20. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.22–1.26 (m, 12H) 1.45–1.51 (m, 4H) 2.16 (t, 2H) 3.25 (qt, 2H), 6.85 (t, 2H), 7.37 (t, 1H), 7.81 (d, 1H), 8.79 (t, 1H), 11.95 (s, 1H), 12.72 (s, 1H)
15	Compound XXVIII: ¹ H NMR (300MHz, DMSO-d ₆): δ 1.26 (8H, br m), 1.49 (4H, m), 2.17 (2H, t), 3.26 (2H, m), 6.86 (2H, m), 7.37 (1H, m), 7.83 (1H, m), 8.80 (1H, t), 11.95 (1H, s), 12.73 (1H, s).
20	Compound XXXIII: ¹ H NMR (300MHz, DMSO-d ₆): δ 1.2 (a, 2H), 1.3 (q, 2H), 1.3 (q, 2H), 1.5 (q, 2H), 2.2 (t, 2H), 3.0 (q, 2H), 3.5 (s, 2H), 7.3 (m, 5H), 8.0 (s, 1H)
25	Compound XXXIV: Anal. Calcd for C ₁₂ H ₁₇ NO ₄ : C, 62.23, H 6.83, N, 5.57 Found: C, 61.93, H, 6.80, N, 5.56. ¹ H NMR (300MHz, DMSO-d ₆): δ 1.24–1.34 (m, 2H) 1.49 (1.57 (m, 4H) 2.19 (t, 2H) 3.26 (qt, 2H), 6.68 (t, 2H), 7.37 (s, 1H), 7.83 (d, 1H) 8.81 (t, 1H), 12.08 (s, 1H), 12.72 (s, 1H)

EXAMPLE 1A

30 An alternate synthesis of compound XIX was as follows:

A 5 L three-neck round bottom flask was fitted with a heating mantle, an overhead mechanical stirrer, an addition funnel, and a thermometer. The reaction was performed under an argon atmosphere. Hydroxylamine-O-sulfonic acid (196.7 g, 1.74 moles, 1.10 equiv.) and formic acid (1 L) were charged into the round bottom flask and stirred to form a white slurry. A solution of cyclooctanone (200.0 g 1.58 moles, 1.0 equiv.) in formic acid (600 mL) was added dropwise to the white slurry via the addition funnel. After the addition, the addition funnel was replaced by a reflux condenser, and the reaction was heated to reflux (internal temperature about 105° C.) for 1 hour to give a brown solution. After the solution was cooled to room temperature, it was poured into a mixture of saturated aqueous ammonium chloride (1.5 L) and water (1.5 L). The aqueous mixture was extracted with chloroform (3×1200 mL). The combined chloroform layers were transferred into a breaker, and saturated sodium bicarbonate (2 L) was added slowly. The chloroform layer was then separated, dried over anhydrous sodium sulfate, and evaporated under reduced pressure to afford a brown oil. The oil was placed in a 500 mL round bottom flask with a magnetic stirrer. The round bottom flask was placed in a silicon oil bath and was fitted with a short path vacuum distillation head equipped with a thermometer. A Cow-type receiver was connected to three 250 mL flasks. 2-Azacyclononanone (145 g, 65%, mp 64–69° C.) was obtained by vacuum distillation (fraction with head temperature range from 80 to 120° C. at pressures between 3.0 and 3.4 mmHg).

60 A 5 L three-neck round bottom flask was fitted with a heating mantle, an overhead mechanical stirrer, a reflux condenser, and a 29 thermometer. A suspension of 2-azacyclononanone (83 g, 0.59 moles, 1.0 equiv.) in 5 M aqueous sodium hydroxide (650 mL, 3.23 moles, 5.5 equiv.) was charged into the round bottom flask. The mixture was heated to reflux (internal temperature about 110° C.) for 4 hours to yield a clear yellow solution. The heating mantle

and reflux condenser were removed. After the solution cooled to room temperature, it was diluted with water (650 mL) and cooled further in an ice bath. Finely ground O-acetylsalicyloyl chloride (114.7 g, 0.59 moles, 1.0 equiv.) was added portionwise to the solution with stirring and continued cooling over 1 hour. After an additional 30 minutes, the ice-bath was removed and stirring was continued at ambient temperature for 21 hours to give a brownish yellow solution. The stirred mixture was acidified with 2 M sulfuric acid (about 850 mL) to a pH of about 1, and a yellow solid was formed. The solid was collected by filtration and was dissolved in warm methanol (1.7 L). Activated charcoal (about 5 g) was added to the methanol, and the solution was stirred for 10 minutes. The activated charcoal was removed by filtration, and the charcoal residue was washed with additional 300 mL methanol. Water (2 L) was added to the combined filtrates (i.e. the 2 L methanol), and an off-white solid precipitated upon standing at 4° C. overnight. The crude product was filtered and was recrystallized from 65% methanol/water (v/v) to yield Compound XIX (69.1 g, 42%) as off-white solid.

Properties are listed below:

mp 116–117° C.; HPLC, ¹H NMR and Anal. Calcd for C₁₅H₂₁NO₄: C, 64.50; H, 7.58; N, 5.02. Found: C, 64.26; H, 7.81; N, 4.93.

EXAMPLE 2

Compound XXXI was prepared as follows:

1-Aminoundec-10-ene. A mixture of 10-undecene-1-ol (5.00 g, 29.36 mmol, 1 equiv), triphenylphosphine (7.70 g, 29.36 mmol, 1 equiv) and phthalimide (4.32 g, 29.36 mmol, 1 equiv) in dry tetrahydrofuran (THF, 30 mL) was stirred vigorously under argon. Diethyl azodicarboxylate (DEAD, 5.11 g, 29.36 mmol, 1 equiv) was diluted with THF (12 mL) and added dropwise by syringe. After the addition, the reaction was stirred at room temperature for 4 hours. The solvent was evaporated under vacuum and ether (30 mL) was added to precipitate the triphenylphosphine oxide and hydrazine dicarboxylate which were removed by filtration. The precipitate was rinsed with ether (2×30 mL) and the combined filtrates were evaporated to afford a yellow solid. The yellow solid was triturated with warm hexanes (3×50 mL) and filtered. The combined hexanes were evaporated to give 1-phthalimidylundec-10-ene as a yellow wax.

The yellow wax was dissolved in an ethanolic solution (38 mL) of hydrazine hydrate (1.47 g, 1 equiv, 29.36 mmol). The mixture was heated at reflux for 2 hours. After the mixture was cooled to room temperature, concentrated hydrochloric acid (30 mL) was added and the solid was filtered through a sintered glass filter. The residue was washed with water (50 mL) and the combined filtrates were evaporated to provide a yellow solid. The yellow solid was redissolved in 1M NaOH (100 mL) and extracted with ether (2×50 mL). The ether was dried and evaporated to provide a yellow oil. The oil was purified by Kugelrohr distillation (ca. 0.1 mmHg, 100° C.) to provide 1-aminoundec-10-ene (2) as a light yellow oil (3.29 g, 66%).

Properties are listed below.

¹H NMR (300 MHz, DMSO-d₆): δ 1.23 (14H, br m), 1.99 (2H, m), 2.48 (2H, m), 4.94 (2H, m), 5.77 (1H, m).

1-(O-Acetylsalicyloylamino)undec-10-ene. O-Acetylsalicyloyl chloride (3.82 g, 19.25 mmol, 1 equiv) in THF (30 mL) was cooled in an ice bath. Triethylamine (1.95 g, 19.25 mmol, 1 equiv), followed by 1-aminoundec-10-ene (3.26 g, 19.25 mmol, 1 equiv) in THF (10 mL) were added via syringe. The ice bath was removed and the reaction was

stirred at room temperature for 3.5 hours. After removal of the solvent, the residue was dissolved in EtOAc (50 mL) and washed with water (2×30 mL). The organic layer was dried and evaporated to afford 1-(O-acetylsalicyloylamino)undec-10-ene as a colorless oil, in a quantitative yield, 6.59 g.

Properties are listed below.

¹H NMR (300 MHz, DMSO-d₆): δ 1.26 (12H, br s), 1.47 (2H, m), 1.99 (2H, m), 2.19 (3H, s), 3.15 (2H, q), 4.95 (2H, m), 5.78 (1H, m), 7.15 (1H, m), 7.30 (1H, m), 7.50 (2H, m), 8.24 (1H, t).

Compound XXXI

1-(O-Acetylsalicyloylamino)undec-10-ene (6.59 g, 19.25 mmol, 1 equiv) in dichloromethane (108 mL) was added to a mixture of water (108 mL), sulfuric acid (9M, 13 mL), glacial acetic acid (2.16 mL) and methyltrialkyl(C₈–C₁₀) ammonium chloride (0.32 g) (Adogen® 464, available from Aldrich Chemical Co.). The mixture was stirred vigorously in an ice bath and potassium permanganate (9.13 g, 57.75 mmol, 3 equiv) was added in portions over 1.5 hours. After the addition, the ice bath was removed and the resultant purple solution was stirred at room temperature for 20 hours. The solution was cooled in an ice bath and sodium bisulfite (6.8 g) was added to dissipate the excess permanganate. The organic layer was separated and the aqueous layer was extracted with ethyl acetate (2×50 mL). The combined organic layers were washed with brine (50 mL), dried and evaporated. Sodium hydroxide (2M, 50 mL) was added to the residue and stirred for 30 min. The solution was diluted with water (50 mL), washed with ether (50 mL) and acidified to pH 1 with 2M hydrochloric acid. A solid formed and was collected by filtration. Recrystallization of the solid from 65 % MeOH/H₂O gave XXXI as a tan solid (2.78 g, 47% based on the amine).

Properties are listed below.

¹H NMR (300 MHz, DMSO-d₆): δ 1.24 (10H, br m), 1.51 (4H, m), 2.17 (2H, t), 3.27 (2H, m), 6.86 (2H, m), 7.37 (1H, m), 7.82 (1H, m), 8.80 (1H, t), 11.95 (1H, s), 12.72 (1H, s).

EXAMPLE 3

Compound LXXXVI was prepared as follows:

A one liter three-neck round bottom flask was fitted with a magnetic stirrer and a condenser. A solution of 3-(4-aminophenyl)propionic acid (30 g, 0.182 moles) in methylene chloride (300 mL) was charged to the flask and trimethylsilyl chloride (46.2 mL, 0.364 moles) was added in one portion. The reaction mixture was refluxed for 1.5 hours, allowed to cool to room temperature, and then immersed in an ice/water bath. Triethylamine (76.2 mL, 0.546 moles) was added, followed by 2-methoxycinnamoyl chloride (35.8 g, 0.182 moles). The reaction mixture was allowed to warm to room temperature and then stirred for 48 hours. The solvent was removed by rotary evaporation and saturated sodium bicarbonate solution and ethyl acetate were added to the residue. The layers were separated, the aqueous layer was acidified to pH 1.4 with 2N aqueous sulfuric acid and extracted with ethyl acetate (2×400 mL). The combined organic extracts were concentrated in vacuo and the residue recrystallized from 50% (v/v) aqueous methanol to provide the product as a tan solid (48.57 g, 82%).

Properties are listed below.

¹H NMR (300MHz, DMSO-d₆): δ 12.1 (1H, br), 7.8 (1H, dd), 7.6 (3H, m), 7.4 (1H, m), 7.3 (2H, m), 7.1 (1H, d), 7.0 (1H, t), 6.9 (1H, d), 3.9 (3H, s), 2.8 (2H, t), 2.5 (4H, m). Anal. Calcd for C₁₉H₁₉NO₄: C, 70.14; H, 5.88; N, 4.31. Found: C, 69.76; H, 5.91; N, 4.21.

EXAMPLE 4

Compound CXVII was prepared as follows:

A 3 L three-neck round bottom flask was fitted with an overhead mechanical stirrer and a thermometer. A solution of 8-aminocaprylic acid (10.0 g, 0.054 moles) in 2 M aqueous sodium hydroxide (1.4 L) was charged into the round bottom flask and O-nitrobenzoyl chloride (12.0 g, 0.065 moles, 1.2 equiv.) was added portionwise over 7 h. The mixture was stirred at 25 ° C. for 12 h to afford a yellow homogenous solution. The solution was acidified with 1 M hydrochloric acid to about pH 2, an oily residue separated and was decanted. The oil was dissolved in stirred water (300 mL) and cooled in and ice/water bath. The product precipitated as a white solid. The solid was filtered, washed with water (3×300 mL), and recrystallized from 55% acetonitrile/water (v/v) to provide Compound CXVII as an off-white solid (7.4 g, 47%). mp 89–92° C.

Properties are listed below.

¹H NMR (300 MHz, DMSO-d₆) δ: 12.0 (1H, s), 8.65 (1H, t), 8.0 (1H, dd), 7.8 (1H, m), 7.65 (1H, m), 7.5 (1H, m), 3.2 (2H, q), 2.2 (2H, t), 1.5 (4H, br m), 1.3 (6H, br m).
Anal. Calcd for C₁₅H₂₀N₂O₅: C, 58.41; H, 6.54; N, 9.09. Found: C, 58.50; H, 6.71; N, 9.14.

The other compounds of the invention can be readily prepared by following the procedures described in Examples 1–4.

EXAMPLES 5–15

In Vivo Evaluation of Recombinant Growth Hormone in Rats

Dosing compositions were prepared by mixing the modified amino acids and recombinant human growth hormone (rhGH) as listed in Table 1 below in a phosphate buffer solution at a pH of about 7–8.

Rats were administered the dosing composition by sublingual, oral gavage, intraduodenal administration, or colonic administration. Delivery was evaluated by using an ELISA assay for rhGH from Medix Biotech, Inc. For intracolonic administration, a sample was prepared and dosed to fasted rats at 25 mg/kg of carrier in a buffered solution containing propylene glycol (0–50%) and 1 mg/kg rhGH.

Results are illustrated in Table 1 below.

Comparative Example 5A

rhGH (6 mg/ml) was administered by oral gavage to a rat, and delivery was evaluated according to the procedure of Example 5.

Results are illustrated in Table 1 below.

TABLE 1

In Viva Delivery of rhGH					
Example	Carrier	Carrier Dose (mg/kg)	Drug Dose (mg/kg)	Method of Administration	Mean Peak Serum Levels of rhGH (ng/mL)
5	I	500	6	oral	26.6 +/- 43.83
5A	none	0	6	oral	<10 +/- 10
6	V	500	6	oral	3.22 +/- 7.2
7	VI	500	6	oral	19.34 +/- 18.73
8	VIII	500	6	oral	73.41 +/- 70.3
9	IX	500	6	oral	28.70 +/- 41.7
10	XIII	25	1	colonic	109.52 +/- 36.1
11	XIX	200	3	oral	60.92 +/- 26.3
12	XIX	25	1	colonic	111.52 +/- 16.4

TABLE 1-continued

In Viva Delivery of rhGH					
Example	Carrier	Carrier Dose (mg/kg)	Drug Dose (mg/kg)	Method of Administration	Mean Peak Serum Levels of rhGH (ng/mL)
13	XIX	100	3	sublingual	119.14 +/- 65.6
14	XIX	25	1	intranasal	92.7 +/- 73.2
15	XXVII	25	1	colonic	73.72 +/- 4.9

EXAMPLES 16–27

In Vivo Evaluation of Recombinant Growth Hormone in Rats

Preparation of Dosing solutions.

The delivery agents were reconstituted with distilled water and adjusted to pH 7.2–8.0 with either aqueous hydrochloric acid or aqueous sodium hydroxide. A stock solution of rhGH was prepared by mixing rhGH, D-mannitol and glycine and dissolving this mixture in 2% glycerol/water. The stock solution was then added to the delivery agent solution. Several delivery agent to active agent ratios were studied.

In vivo experiments.

Male Sprague-Dawley rats weighing 200–250 g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The rats were administered one of the dosing solutions described above by subcutaneous injection, intranasal instillation, or sublingual instillation. Blood samples were collected serially from the tail artery for serum calcium concentration determination or serum rhGH concentrations. The dose of rhGH administered in these experiments was 0.1 mg/kg.

Serum rhGH concentrations were quantified by an rhGH enzyme immunoassay test kit. The results are given in Table 2 and FIGS. 1 and 2.

In FIG. 2 the circles represent the response following SL dosing of an aqueous solution of compound CXXIII and rhGH. The squares represent the response following IN dosing of an aqueous solution of compound CXXIII and rhGH. The triangles represent the response following IC dosing of an aqueous solution of compound CXXIII and rhGH. The dose of compound CXXIII was 25 mg/kg and the dose of rhGH was 1 mg/kg.

Comparative Example 16A

rhGH (1 mg/kg) was administered by oral gavage to a rat, and delivery was evaluated according to the procedure of Example 16.

Results are illustrated in Table 2 below.

TABLE 2

Delivery Agent Enhancement of Recombinant Human Growth Hormone (rhGH) Bioavailability Administered by Subcutaneous Administration.			
Example	Deliver Agency	Delivery Agent Dose (mg/kg)	Peak Serum [rhGH] (ng/mL)
16	CXXIII	1.0	22 ± 3
16A	None	0.0	4 ± 2
17	CXXIII	2.5	25 ± 5
18	CXXIII	25	30 ± 6
19	CX	2.5	16 ± 2
20	LVIII	1.0	29 ± 10

TABLE 2-continued

Delivery Agent Enhancement of Recombinant Human Growth Hormone (rhGH) Bioavailability Administered by Subcutaneous Administration.			
Example	Deliver Ageny	Delivery Agent Dose (mg/kg)	Peak Serum [rhGH] (ng/mL)
21	LXXXVI	1.0	22 ± 7
22	LXXXVI	2.5	23 ± 5
23	LXI	2.5	26 ± 5
24	CX	1.0	15 ± 3
25	CXV	1.0	25 ± 3
26	LXVI	1.0	33 ± 5
27	CIX	1.0	16 ± 3

EXAMPLES 28-33

In Vivo Evaluation of Interferon in Rats

Dosing compositions were prepared by mixing the modified amino acid compounds and interferon α2b as listed in Table 3 below in a Trizma® hydrochloride buffer solution (Tris-HCl) at a pH of about 7-8. Propylene glycol (0-25%) was added as a solubilizing agent, if necessary.

Rats were administered the dosing composition by oral gavage, intraduodenal administration, or intracolonic administration. Delivery was evaluated by use of an ELISA assay for human interferon a from Biosource, Inc.

Results of intracolonic administration are illustrated in Table 3 below.

Comparative Example 28A

Interferon a2b (250 µg/kg) was administered intracolonicly to rats, and delivery was evaluated according to the procedure of Example 14.

Results are illustrated in Table 3 below.

TABLE 3

In Vivo Delivery of Interferon by Intracolonic Administration				
Example	Carrier	Carrier Dose (mg/kg)	Interferon Dose (µg/kg)	Mean Peak Serum Levels of Interferon (pg/mL)
28	VII	100	250	5241 +/- 2205
28A	none	0	250	0
29	XI	100	250	1189 +/- 1373
30	XII	100	250	6955 +/- 2163
31	XIX	100	250	11193 +/- 8559
32	XXI	100	250	4238 +/- 2789
33	XXXIV	100	250	4853 +/- 5231

Results are illustrated in Table 4 below.

EXAMPLES 34-37

In Vivo Evaluation of Salmon Calcitonin in Rats

Dosing compositions were prepared by mixing the modified amino acids and salmon calcitonin as listed in Table 4 below. 400 mg of carrier were added to 2.9 mL of 25% aqueous propylene glycol. The resultant solution was stirred, and the pH was adjusted to 7.2 with sodium hydroxide (1.0 N). Water was added to bring the total volume to 2.0 mL.

The sample had a final carrier concentration of 200 mg/mL. Calcitonin (10 µg) was added to the solution. The total calcitonin concentration was 2.5 µg/mL.

For each sample a group of fasted rats were anesthetized. The rats were administered the dosing composition by oral gavage, intracolonic instillation, or intraduodenal administration. Blood samples were collected serially from the tail artery. Serum calcium was determined by testing with a calcium Kit (Sigma Chemical Company, St. Louis, Mo., USA).

Results are illustrated in Table 4 below.

TABLE 4

In Vivo Delivery of Calcitonin					
Example	Carrier	Carrier Dose (mg/kg)	Drug Dose (µg/kg)	Method of Administration	Maximum Decrease in Serum Calcium (% below baseline)
34	I	400	10	oral	
35	V	400	10	oral	18.35 +/- 2.87
36	XIX	10	3	intracolonic	26.49 +/- 12.3
37	XIX	200	7.5	oral	25.48 +/- 4.7

EXAMPLES 38-43

In Vivo Evaluation of Salmon Calcitonin in Rats

Preparation of Dosing solution.

The delivery agents were reconstituted with distilled water and adjusted to pH 7.2-8.0 with either aqueous hydrochloric acid or aqueous sodium hydroxide. A stock solution of sCT was prepared by dissolving sCT in citric acid (0.085N). The stock solution was then added to the delivery agent solution. Several different delivery agent to active agent ratios were studied.

In vivo experiments.

Male Sprague-Dawley rats weighing 200-250 g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The rats were administered one of the dosing solutions described above by subcutaneous injection. Blood samples were collected serially from the tail artery for serum calcium concentration.

Serum calcium concentrations were quantified by the o-cresololphthalein complex one method (Sigma) using a UV/VIS spectrophotometer (Perkin Elmer). The results are given in Table 5.

EXAMPLES 38A

Salmon calcitonin was administered by oral gavage to rats, and delivery was evaluated according to the Procedure of Example 38. The results are given in Table 5 below.

TABLE 5

Delivery Agent Enhancement of Salmon Calcitonin (sCT, dosed at 0.2 µg/kg) Bioavailability Administered by Subcutaneous Administration.			
Example	Deliver Ageny	Delivery Agent Dose (µg/kg)	Percent Decrease in Serum Calcium
38	CXXIII	2	17 ± 3
38A	None	0	17 ± 2
39	CXXIII	20	25 ± 4
40	CXXIII	200	25 ± 5
41	CXXIII	2000	26 ± 5
42	CXI	20	21 ± 4
43	CXIV	20	20 ± 3

EXAMPLES 44-50

In Vivo Evaluation of HeDarin in Rats

Dosing compositions were prepared by mixing the modified amino acids and heparin as listed in Table 4. In a test tube, 900 mg of carrier were dissolved in 3 mL of propylene glycol, and 0.299 g of sodium heparin was dissolved in 3 mL of water. The solutions were mixed by vortex. Sodium hydroxide (10M) was added to the resulting mixture until a solution was obtained. The pH was then adjusted to 7.4 +/- 0.5 with concentrated hydrochloric acid, and the final solution was sonicated at 40° C. for 30 minutes.

A group of fasted, conscious rats were administered the dosing compositions by oral gavage. Blood samples were collected by cardiac puncture following the administration of ketamine (44 mg/kg)., Heparin activity was determined by utilizing the activated partial thromboplastin time (APTT) according to the method of Henry, J. B., *Clinical Diagnosis and Management by Laboratory Methods*; Philadelphia, Pa.; W B Saunders (1979).

Results are illustrated in table 6 below.

Comparative Example 44A

Heparin (100 mg/kg) was administered by oral gavage to rats, and heparin activity was determined according to the procedure of Example 44.

Results are illustrated in Table 6 below.

TABLE 6

In Vivo Delivery of Heparin by Oral Administration				
Example	Carrier	Carrier Dose (mg/kg)	Drug Dose (mg/kg)	Mean Peak APTT (sec)
44	II	300	100	25.45 +/- 2.8
44A	none	none	100	20.7 +/- 0.17
45	III	300	100	38.64 +/- 17
46	V	300	100	87.4 +/- 34.1
47	XII	300	100	49.53 +/- 17.1
48	XIX	300	100	119.99 +/- 56.3
49	XXXI	50	25	127.56 +/- 22.97
50	XXXI	50	10	50.85 +/- 9.1

EXAMPLE 51

The method of Example 44 was followed, substituting low molecular weight heparin for the heparin and varying the amounts of propylene glycol and water for solubilization as, necessary.

EXAMPLES 50-58

In vivo Evaluation of Parathyroid Hormone in Rats

Preparation of dosing solutions.

The delivery agents were reconstituted with distilled water and/or propylene glycol and adjusted to an apparent pH of 7.2-8.0 with either aqueous hydrochloric acid or aqueous sodium hydroxide. A stock solution of parathyroid hormone was prepared by dissolving parathyroid hormone in water. The parathyroid hormone solution was then added to the delivery agent solution. Several different delivery agent to active agent ratios, were studied.

In vivo experiments.

Male Sprague-Dawley rats weighing 200-250 g were fasted for 24 hours and administered ketamine (44 mg/kg) and chlorpromazine (1.5 mg/kg) 15 minutes prior to dosing. The rats were administered one of the dosing solutions described above by oral gavage or intracolonic instillation. Blood samples were collected serially from the tail artery for serum determination of parathyroid hormone concentration. Serum parathyroid hormone concentrations were quantified by a parathyroid hormone radioimmunoassay test kit.

In vivo Oral administration.

Oral administration of solutions containing parathyroid hormone (PTH) and the non-a-amino acid delivery agents was tested in vivo in rats. The result show a significant increase in the oral bioavailability of parathyroid hormone as compared to similar administration of the active agent alone. Data are presented in Table 7.

TABLE 7

Delivery Agent Enhancement of Parathyroid Hormone (PTH) Oral Bioavailability.						
Example	Carrier	Carrier Dose mg/kg	Method of Administration	Active Agent Dose (μg/kg)	Peak Serum [PTH] (pg/mL)	
51	CXXIII	100	intracolonic	25	130 ± 20	
52	CXXIII	250	oral	100	75 ± 25	
53	CXXIII	250	oral	25	20 ± 6	
54	CVIII	100	intracolonic	25	115 ± 20	
55	LXXXVI	100	intracolonic	25	40 ± 12	
56	LVIII	100	intracolonic	25	145 ± 25	
57	CXIV	100	intracolonic	25	65 ± 15	
58	LXXXIX	100	intracolonic	25	70 ± 15	

The above mentioned patents, applications, test methods, and publications are hereby incorporated by reference in their entirety.


Many variations of the present invention will suggest themselves to those skilled in the art in light of the above detailed description. All such obvious variations are within the full intended scope of the appended claims.

What is claimed is:

1. A composition comprising:

(A) at least one active agent; and

(B) a compound having the following formula



or a salt thereof.

2. A composition as defined in claim 1, wherein said active agent is selected from the group consisting of a biologically active agent, a chemically active agent, and a combination thereof.

3. A composition as defined in claim 2, wherein said biologically active agent comprises at least one peptide, hormone, polysaccharide, mucopolysaccharide, carbohydrate, or lipid.

4. The composition as defined in claim 3, wherein said biologically active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-1, interleukin-II, insulin, heparin, low molecular weight heparin, calcitonin, erythropoietin, atrial natriuretic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, parathyroid hormone, desferrioxamine, an anti-microbial agent, an anti-fungal agent, and any combination thereof.

5. A composition as defined in claim 4, wherein said biologically active agent comprises an interferon, interleukin-II, insulin, heparin, low molecular weight heparin, calcitonin, oxytocin, vasopressin, vancomycin, DFO, parathyroid hormone, or a combination thereof.

6. A composition as defined in claim 2, wherein said biologically active agent comprises parathyroid hormone.

7. A composition as defined in claim 2, wherein said biologically active agent comprises recombinant human growth hormone.

39

8. A composition as defined on claim 2, wherein said biologically active agent comprises interferon.

9. A composition as defined in claim 2, wherein said biologically active agent comprises heparin.

10. A composition as defined in claim 2, wherein said biologically active agent comprises low molecular weight heparin.

11. A dosage unit form comprising:

(A) a composition as defined in claim 1; and

(B) (a) an excipient

(b) a diluent,

(c) a disintegrant,

(d) a lubricant,

(e) a plasticizer,

(f) a colorant,

(g) a dosing vehicle, or

(h) any combination thereof.

12. The dosage unit form as defined in claim 11, wherein said active agent is selected from the group consisting of a biologically active agent, a chemically active agent, and a combination thereof.

13. The dosage unit form as defined in claim 12, wherein said biologically active agent comprises at least one peptide, hormone, polysaccharide, mucopolysaccharide, carbohydrate, or lipid.

14. The dosage unit form as defined in claim 13, wherein said biologically active agent is selected from the group consisting of human growth hormone, bovine growth hormone, growth hormone-releasing hormone, an interferon, interleukin-1, interleukin-II, insulin, heparin, low molecular weight heparin, calcitonin, erythropoietin, atrial natriuretic factor, an antigen, a monoclonal antibody, somatostatin, adrenocorticotropin, gonadotropin releasing hormone, oxytocin, vasopressin, cromolyn sodium, vancomycin, parathyroid hormone, desferrioxamine, an anti-microbial agent, an anti-fungal agent, and any combination thereof.

15. A dosage unit form as defined in claim 14, wherein said biologically active agent comprises an interferon, interleukin-II, insulin, heparin, low molecular weight heparin, calcitonin, oxytocin, vasopressin, vancomycin, DFO, parathyroid hormone, or a combination thereof.

16. A dosage unit form as defined in claim 12, wherein said biologically active agent comprises parathyroid hormone.

17. A dosage unit form as defined in claim 12, wherein said biologically active agent comprises recombinant human growth hormone.

18. A dosage unit form as defined on claim 12, wherein said biologically active agent comprises interferon.

40

19. A dosage unit form as defined in claim 12, wherein said biologically active agent comprises heparin.


20. A dosage unit form as defined in claim 12, wherein said biologically active agent comprises low molecular weight heparin.

21. A dosage unit form as defined in claim 11, comprising a tablet, a capsule, or a liquid.

22. A dosage unit form as defined in claim 21, wherein said dosing vehicle is selected from the group consisting of water, 1,2-propane diol, ethanol, and any combination thereof.

23. A method for administering a biologically-active agent to an animal in need of said agent, said method comprising administering orally to said animal a composition as defined in claim 2.

24. A compound having the formula



or a salt thereof.

25. A method for preparing a composition, said method comprising mixing:

(A) at least one active agent;


(B) at least one compound as defined in claim 24; and

(C) optionally, a dosing vehicle.

26. A composition comprising:

(A) at least one active agent; and

(B) a poly(amino acid) comprising a monomer having the formula



or a salt thereof.

27. A composition as defined in claim 26, wherein the poly(amino acid) comprises a peptide.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
Certificate

Patent No. 6,346,242 B1

Patented: February 12, 2002

On petition requesting issuance of a certificate for correction of inventorship pursuant to 35 U.S.C. 256, it has been found that the above identified patent, through error and without any deceptive intent, improperly sets forth the inventorship.

Accordingly, it is hereby certified that the correct inventorship of this patent is: Andrea Leone-Bay, Ridgefield, CT; Koc-Kan Ho, Monmouth Junction, NJ; Donald J. Sarubbi, Carmel, NY; Sam J. Milstein, Larchmont, NY; and Lise M. Genoble, Irvington, NY.

Signed and Sealed this Tenth Day of June 2003.

CHRISTOPHER LOW
Supervisory Patent Examiner
Art Unit 1653

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 6,346,242 B1
DATED : February 12, 2002
INVENTOR(S) : Andrea Leone-Bay et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Title page,

Item [73], Assignee, delete “**Emishpere Technologies, Inc.,**” and substitute
-- **Emisphere Technologies, Inc.,** --

Signed and Sealed this

Second Day of December, 2003

A handwritten signature in black ink, appearing to read "James E. Rogan", written over a horizontal line.

JAMES E. ROGAN
Director of the United States Patent and Trademark Office