



US006420541B1

(12) **United States Patent**  
**Behan et al.**

(10) **Patent No.:** **US 6,420,541 B1**  
(45) **Date of Patent:** **Jul. 16, 2002**

(54) **NON-ENDOGENOUS, CONSTITUTIVELY ACTIVATED HUMAN SEROTONIN RECEPTORS AND SMALL MOLECULE MODULATORS THEREOF**

(75) Inventors: **Dominic P. Behan**, San Diego; **Derek T. Chalmers**, Solana Beach; **Chen W. Liaw**; **Joseph F. Russo**, both of San Diego; **William J. Thomsen**, Del Mar, all of CA (US)

(73) Assignee: **Arena Pharmaceuticals, Inc.**, San Diego, CA (US)

(\* ) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/767,013**

(22) Filed: **Dec. 22, 2000**

#### Related U.S. Application Data

(60) Division of application No. 09/292,072, filed on Apr. 14, 1999, which is a continuation-in-part of application No. 09/060,188, filed on Apr. 14, 1998.

(60) Provisional application No. 60/090,783, filed on Jun. 29, 1998, provisional application No. 60/112,909, filed on Dec. 18, 1998, and provisional application No. 60/123,000, filed on Mar. 5, 1999.

(51) **Int. Cl.**<sup>7</sup> ..... **C07K 21/04**; C07K 14/00; C12N 15/00; C12N 15/09

(52) **U.S. Cl.** ..... **536/23.1**; 536/23.5; 435/320.1; 435/325; 435/252.3; 435/69.1; 435/172.3; 435/7.1; 530/300; 530/350

(58) **Field of Search** ..... 536/23.1, 23.5, 536/300, 350; 435/69.1, 320.1, 325, 172.3, 252.3, 7.1

#### (56) **References Cited**

##### U.S. PATENT DOCUMENTS

4,985,352 A 1/1991 Julius et al. .... 435/6  
5,661,024 A 8/1997 Kao et al. .... 435/240.2

##### FOREIGN PATENT DOCUMENTS

CA 2135253 5/1996  
WO WO 96/23783 8/1996

##### OTHER PUBLICATIONS

Barluenga, J. et al., "A New and Specific Method for the Monomethylation of Primary Amines," *J. Chem. Soc. Chem. Commun.*, 1984, 20, 1334-1335.

Batey, R.A. et al., "An Efficient New Protocol for the Formation of Unsymmetrical Tri- and Tetrasubstituted Ureas," *Tetra. Lett.*, 1988, 39, 6267-6270

Bernatowicz, M. et al., "A Comparison of Acid Labile Linkage Agents for the Synthesis of Peptide C-Terminal Amides," *Tetra. Lett.*, 1989, 30(35), 4645-4648.

Carter, H.E. et al., "Carbobenzoxy Chloride and Derivatives," *Org. Syn. Coll.*, 1955, vol. 3, 167-169.

Casey, C. et al., "Constitutively Active Mutant 5HT<sub>2A</sub> Serotonin Receptors: Inverse Agonist Activity of Classical 5HT<sub>2A</sub> Antagonists," *Society for Neuroscience Abstracts*, 1996, 22(3), Abstract No. 699.10.

Gutsche, C.D. et al., "2-Phenylcycloheptanone," *Org. Syn. Coll.*, 1963, vol. 4, 780-783.

Herrick-Davis, K. et al., "Activating Mutations of the Serotonin 5-HT<sub>2C</sub> Receptor," *J. Neurochem.*, 1997, 69(3), 1138-1144.

Herrick-Davis, K. et al., "Constitutively Active 5HT<sub>2C</sub> Serotonin Receptor Created by Site-Directed Mutagenesis," *Society for Neuroscience Abstracts*, 1996, 22(3), Abstract No. 699.18.

Konig, W. et al., "A New Method for Synthesis of Peptides: Activation of the Carboxyl Group with Dicyclohexylcarbodiimide Using 1-Hydroxybenzotriazoles as Additives," *Chem. Ber.*, 1970, 103, 788-798 (English Abstract included).

Marchini, P. et al., "Sodium Borohydride-Carboxylic Acid Systems. Useful Reagents for the Alkylation of Amines," *J. Org. Chem.*, 1975, 40(23), 3453-3456.

Sahgal, A. (ed.), "Practical behavioral neuroscience: problems, pitfalls and suggestions," in *Behavioral Neuroscience: A Practical Approach*, IRL Press, New York, 1993, vol. 1, 1-8.

Sheehan, J.C. et al., "1-Ethyl-3-(3-Dimethylamiono) Propylcarbodiimide Hydrochloride and Methiodide," *Org. Syn. Coll.*, 1973, vol. 5, 555-558.

Soresnon et al., "Characterization of the 5-HT<sub>2</sub> Receptor Antagonist MDL 100907 as Putative Atypical Antipsychotic: Behavioral, Electrophysiological and Neurochemical Studies," *J. Pharmacol. Exp. Ther.*, 1993, 266(2), 684-691.

White, E., "Deamination of Amines. 2-Phenylethyl Benzoate Via the Nitrosoamide Decomposition," *Org. Syn. Coll.*, 1973, vol. 5, 336-339.

*Primary Examiner*—Yvonne Eyler

*Assistant Examiner*—Nirmal S. Basi

(74) *Attorney, Agent, or Firm*—Woodcock Washburn LLP

#### (57) **ABSTRACT**

Disclosed herein are non-endogenous, constitutively activated forms of the human 5-HT<sub>2A</sub> and human 5-HT<sub>2C</sub> receptors and uses of such receptors to screen candidate compounds. Further disclosed herein are candidate compounds identified by the screening method which act at the 5HT<sub>2A</sub> receptors. Yet further disclosed is a new class of compounds which act at the 5HT<sub>2A</sub> receptors.

**1 Claim, 17 Drawing Sheets**




FIG. 1




FIG. 2

*FIG. 3A*

ATGGATATTCTTTGTGAAGAAAATACTTCTTTGAGCTCAACTACGAACTCCCTAATGCAATTA  
AATGATGACAACAGGCTCTACAGTAATGACTTTAACTCCGGAGAAGCTAACACTTCTGATGCA  
TTAACTGGACAGTCGACTCTGAAAATCGAACCAACCTTTCCTGTGAAGGGTGCCTCTCACCG  
TCGTGTCTCTCCTTACTTCATCTCCAGGAAAAAACTGGTCTGCTTTACTGACAGCCGTAGTGA  
TTATTCTAACTATTGCTGGAACATACTCGTCATCATGGCAGTGTCCCTAGAGAAAAAGCTGC  
AGAATGCCACCAACTATTTCCTGATGTCACTTGCCATAGCTGATATGCTGCTGGGTTTCCTTGT  
CATGCCCGTGTCCATGTTAACCATCCTGTATGGGTACCGGTGGCCTCTGCCGAGCAAGCTTTGT  
GCAGTCTGGATTTACCTGGACGTGCTTCTCCACGGCCTCCATCATGCACCTCTGCGCCATCT  
CGCTGGACCGCTACGTGCGCCATCCAGAATCCCATCCACCACAGCCGCTTCAACTCCAGAACTA  
AGGCATTTCTGAAAATCATTGCTGTTTGGACCATATCAGTAGGTATATCCATGCCAATACCAG  
TCTTTGGGCTACAGGACGATTCGAAGGTCTTTAAGGAGGGGAGTTGCTTACTCGCCGATGATA  
ACTTTGTCCTGATCGGCTCTTTTGTGTCATTTTTCATTCCTTAACCATCATGGTGATCACCTAC  
TTTCTAACTATCAAGTCACTCCAGAAAGAAGCTACTTTGTGTGTAAGTGATCTTGGCACACGG  
GCCAAATTAGCTTCTTTCAGCTTCCCTCCCTCAGAGTTCTTTGTCTTCAGAAAAGCTCTTCCAGC  
GGTCGATCCATAGGGAGCCAGGGTCTACACAGGCAGGAGGACTATGCAGTCCATCAGCAAT  
GAGCAAAAGGCATGCAAGGTGCTGGGCATCGTCTTCTTCCTGTTTGTGGTGATGTGGTGCCT  
TTCTTCATCACAAACATCATGGCCGTCATCTGCAAAGAGTCTGCAATGAGGATGTCATTGGG  
GCCCTGCTCAATGTGTTTGTGGATCGGTTATCTCTCTTCAGCAGTCAACCCACTAGTCTACA  
CACTGTTCAACAAGACCTATAGGTCAGCCTTTTCACGGTATATTCAGTGTGAGTACAAGGAAA  
ACAAAAACCATTGCAGTTAATTTTAGTGAACACAATAACCGGCTTTGGCCTACAAGTCTAGCC  
AACTTCAAATGGGACAAAAAAGAATTCAAAGCAAGATGCCAAGACAACAGATAATGACTGC  
TCAATGGTTGCTCTAGGAAAGCAGTATTCTGAAGAGGCTTCTAAAGACAATAGCGACGGAGT  
GAATGAAAAGGTGAGCTGTGTGTGA

*FIG. 3B*

MDILCEENTSLSSTTNSLMQLNDDNRLYSNDFNSGEANTSDAFNWTVDSERNTLSCEGCLSPSCL  
SLLHLQEKNEWSALLTAVVILTIAGNILVMAVSLEKKLQATNYFLMSLAIADMLLGFLVMPVSM  
LTILYGYRWPLPSKLCVWYLDVLFSTASIMHLCALSIDRYVAIQNPIHHSRFNSRTKAFLKIIAVW  
TISVGISMPIPVFGLQDDSKVFKEGSCLLADDNFVLIGSFVSFFIPLTMVITYFLTIKSLQKEATLCVS  
DLGTRAKLASFSFLPQSSLSSEKLFQRSIHREPGSYTGRRTMQSISNEQKACKVLGIVFFLVVMWC  
PFFITNIMAVICKESCNEVDIGALLNVFVWIGYLSSAVNPLVYTLFNKTYRSAFSRYIQCYKENKK  
PLQLLVNTIPALAYKSSQLQMGQKNSKQDAKTNDNDCSMVALGKQYSEEASKDNSDGVNEKV  
SCV

*FIG. 4B*

MVNLRNAVHSFLVHLIGLLVWQCDISVSPVAAIVTDIFNTSDGGRFKFPDGVQNWPAISIVIIIMTIGGN  
ILVMAVSMEKKLHNATNYFLMSLAIADMLVGLLVMPLSLLAILYDVWPLPRYLCPVWISLDVLFSTASI  
MHLCAISIDRYVAIRNPIEHSRFNSRTKAIMKIAVWALSIGVSVPIPVIGLRDEEKVFNNTTCVLNDPN  
FVLIGSFVAFFIPLTMVITYCLTIYVLRQALMLLHGHTTEPPGLSLDFLKCKRNTAEEENSANPNQDQ  
NARRRKKKERRPRGTMQAINNERKASKVLGIVFFVFLMWCPFFITNILSVLCEKSCNQKLMKLLNVFVW  
IGYVCSGINPLVYTLFNKTYRRAFSNYLRCNYKVEKKPPVRQIPVAATALS GRELVNVIYRHTNEPVIEK  
ASDNEPGIEMQVENLELPVNPSSVVSERISSV

*FIG. 4A*

ATGGTGAACCTGAGGAATGCGGTGCATTCATTCCTTGTGCACCTAATTGGCCTATTGGTTTGGC  
AATGTGATATTTCTGTGAGCCCAGTAGCAGCTATAGTAACTGACATTTTCAATACCTCCGATG  
GTGGACGCTTCAAATTCCCAGACGGGGTACAAAACCTGGCCAGCACTTTCAATCGTCATCATAA  
TAATCATGACAATAGGTGGCAACATCCTTGTGATCATGGCAGTAAGCATGGAAAAGAACTG  
CACAATGCCACCAATTACTTCTTAATGTCCCTAGCCATTGCTGATATGCTAGTGGGACTACTTG  
TCATGCCCTGTCTCTCCTGGCAATCCTTTATGATTATGTCTGGCCACTACCTAGATATTTGTG  
CCCCGTCTGGATTTCTTTAGATGTTTTATTTTCAACAGCGTCCATCATGCACCTCTGCGCTATAT  
CGCTGGATCGGTATGTAGCAATACGTAATCCTATTGAGCATAGCCGTTTCAATTCGCGGACTA  
AGGCCATCATGAAGATTGCTATTGTTTGGGCAATTTCTATAGGTGTATCAGTTCCTATCCCTGT  
GATTGGACTGAGGGACGAAGAAAAGGTGTTTCGTGAACAACACGACGTGCGTGCTCAACGACC  
CAAATTCGTTCTTATTGGGTCCCTTCGTAGCTTTCTTCATACCGCTGACGATTATGGTGATTAC  
GTATTGCCTGACCATCTACGTTCTGCGCCGACAAGCTTTGATGTTACTGCACGGCCACACCGA  
GGAACCGCCTGGACTAAGTCTGGATTTCTGAAAGTGCTGCAAGAGGAATACGGCCGAGGAAG  
AGAACTCTGCAAACCCTAACCAAGACCAGAACGCACGCCGAAGAAAGAAGAAGGAGAGACG  
TCCTAGGGGCACCATGCAGGCTATCAACAATGAAAGAAAAGCTTCGAAAGTCCTTGGGATTG  
TTTTCTTTGTGTTTCTGATCATGTGGTGCCCATTTTTTCATTACCAATATTCTGTCTGTTCTTTGTG  
AGAAGTCCTGTAACCAAAAAGCTCATGAAAAGCTTCTGAATGTTGTTTGGATTGGCTAG  
TTGTTCAGGAATCAATCCTCTGGTGTATACTCTGTTCAACAAAATTTACCGAAGGGCATTCTC  
CAACTATTTGCGTTGCAATTATAAGGTAGAGAAAAGCCTCCTGTCAGGCAGATTCCAAGAGT  
TGCCGCCACTGCTTTGTCTGGGAGGGAGCTTAATGTTAACATTTATCGGCATACCAATGAACC  
GGTGATCGAGAAAGCCAGTGACAATGAGCCCGGTATAGAGATGCAAGTTGAGAATTTAGAGT  
TACCAGTAAATCCCTCCAGTGTGGTTAGCGAAAGGATTAGCAGTGTGTGA

*FIG. 5A*

ATGGTGAACCTGAGGAATGCGGTGCATTCATTCCTTGTGCACCTAATTGGCCTATTGGTTTGGCAAT  
GTGATATTTCTGTGAGCCCAGTAGCAGCTATAGTAACTGACATTTCAATACCTCCGATGGTGGACG  
CTTCAAATCCCAGACGGGGTACAAAACCTGGCCAGCACTTTCAATCGTCATCATAATAATCATGAC  
AATAGGTCGCAACATCCTTGTGATCATGGCAGTAAGCATGGAAAAGAACTGCACAATGCCACCA  
ATTACTTCTTAATGTCCCTAGCCATTGCTGATATGCTAGTGGGACTACTTGTTCATGCCCTGTCTCTC  
CTGGCAATCCTTTATGATTATGTCTGGCCATCAACTAGATATTTGTGCCCGTCTGGATTTCTTTAGA  
TGTTTTATTTCAACAGCGTCCATCATGCACCTCTGCGCTATATCGCTGGATCGGTATGTAGCAATA  
CGTAATTTCTATTGAGCATAGCCGTTTCAATTCGCGGACTAAGGCCATCATGAAGATTGCTATTGTTT  
GGGCAATTTCTATAGGTGTATCAGTTCCTATCCCTGTGATTGGACTGAGGGACGAAGAAAAGGTGT  
TCGTGAACAACACGACGTGCGTGCTCAACGACCCAAATTTGCTTCTTATTGGGTCCTTCGTAGCTTT  
CTTCATACCGCTGACGATTATGGTGATTACGTATTGCCTGACCATCTACGTTCTGCGCCGACAAGCT  
TTGATGTTACTGCACGGCCACACCGAGGAACCGCCTGGACTAAGTCTGTATTTCTGAACTGCTGC  
AAGAGGAATACGGCCGAGGAAGAGA AACTCTGCAAACCTAACCAAGACCAGAACGCACGCCGAA  
GAAAGAAGAAGGAGAGACGTCTAGGGGCACCATGCAGGCTATCAACAATGAAAGAAAAGCTAA  
GAAAGTCCTTGGGATTGTTTTCTTTGTGTTTCTGATCATGTGGTGCCCATTTTTTCATTACCAATATTC  
TGTCTGTTCTTTGTGAGAAGTCCTGTAACCAAAAGCTCATGGAAAAGCTCTGAATGTGTTTGTGTTG  
GATTGGCTATGTTTGTTCAGGATTCAATCCTCTGGTGTATACTCTGTTCAACAAAATTTACCGAAGG  
GCATTTCTCCAAC TATTTGCGTTGCAATTATAAGGTAGAGAAAAGCCCTCCTGTCAGGCAGATTCCA  
AGAGTTGCCGCCACTGCTTTGTCTGGGAGGGAGCTTATTGTTAACATTTATCGGCATACCAATGAA  
CCGGTGATCGAGAAAGCCAGTGACAATGAGCCCGGTATAGAGATGCAAGTTGAGAATTTAGAGTT  
ACCAGTAAATCCCTCCAGTGTGGTTAGCGAAAGGATTAGCAGTGTGTGA


*FIG. 5B*

MVNLRNAVHSFLVHLIGLLVWQCDISVSPVAAIVTDIFNTSDGGRFKFPDG VQNWP ALSIVIIIIMTI  
 GGNILVIMAVSMEKKLHNATNYFLMSLAIADMLVGLLVMPLSLLAILDYVWPLPRYLCPVWISL  
 DVLFSTASIMHLC AISLD RYVAIRNPIEHSRFSR TKAIMKIAI VW AISIG VSVP IPIVIGLRDEEKV FV  
 NNTTCVLNDPNFV LIGSFVAFFIPLT MVTITYCLTIYV LRRQALMLLHGHT EEPGLSDFLKCKRN  
 TAE EENSANPNQDQ NARRRKKKERRPRGTMQAINNERKAKKVLGIVFFVFLIMWCPFFITNLSVL  
 CEKSCNQKLM EKLLNVFVWIGYVCSGINPLVYTLFNKIYRRAFSNYLRCNYKVEKKPPVRQIPRV  
 AAT ALSGRENLNVNTYRHTNEP VIEKASDNEPGIEMQVENLELPVNPSSVVSERISSV

*FIG. 6B*

MDILCEENTSL SSTTNSLMQLNDDNRLYSNDFNSGEANTS DAFNWTVDSENRTNLSCEGCLSPSCL  
 SLLHLQEK NWSALLTAVVILLTIAGN ILVIMAVSLEKKLQ NATNYFLMSLAIADMLLGFLVMPVSM  
 LTILYGYRWPLPSKLC AVWIYLDVLFSTASIMHLC AISLD RYVAIQNPIHHSRFSR TKAFKIAVW  
 TISVGISMPIPVFGLQDDSKVFKEGSCLLADDNFV LIGSFVSFFIPLT MVTITYFLT IKV LRRQALMLL  
 HGHT EEPGLSDFLKCKRNTAE EENSANPNQDQ NARRRKKKERRPRGTMQAINNERKAS  
 KVLGIVFFLFVVMWCPFFITNIMAVICKESCNE D VIGALLNVFVWIGYLSSAVNPLVYTLFNKIYR  
 RAFSNYLRCNYKVEKKPPVRQIPRVAAT ALSGRENLNVNTYRHTNEP VIEKASDNEPGIEMQVE  
 NLELPVNPSSVVSERISSV

*FIG. 6C*





*FIG. 6A*


ATGGATATTCTTTGTGAAGAAAATACTTCTTTGAGCTCAACTACGAACTCCCTAATGCAATTA  
AATGATGACAACAGGCTCTACAGTAATGACTTTAACTCCGGAGAAGCTAACACTTCTGATGCA  
TTAACTGGACAGTCGACTCTGAAAATCGAACCAACCTTTCCTGTGAAGGGTGCCCTCACCG  
TCGTGTCTCTCCTTACTTCATCTCAGGAAAAAACTGGTCTGCTTTACTGACAGCCGTAGTGA  
TTATTCTAACTATTGCTGGAAACATACTCGTCATCATGGCAGTGTCCCTAGAGAAAAAGCTGC  
AGAATGCCACCAACTATTTCTGATGTCACTTGCCATAGCTGATATGCTGCTGGGTTTCCTTGT  
CATGCCCGTGTCCATGTTAACCATCCTGTATGGGTACCGGTGGCCTCTGCCGAGCAAGCTTTGT  
GCAGTCTGGATTACCTGGACGTGCTTTCTCCACGGCCTCCATCATGCACCTCTGCGCCATCT  
CGCTGGACCGCTACGTCGCCATCCAGAATCCATCCACCACAGCCGCTTCAACTCCAGAACTA  
AGGCATTTCTGAAAATCATTGCTGTTGGACCATATCAGTAGGTATATCCATGCCAATACCAG  
TCTTTGGGCTACAGGACGATTCGAAGGTCTTTAAGGAGGGGAGTTGCTTACTCGCCGATGATA  
ACTTTGTCTGATCGGCTCTTTTGTGTCATTTTTCATTCCCTTAACCATCATGGTGATCACCTAC  
TTTCTAACTATCAAGGTTCTGCGCCCGACAAGCTTTGATGTTACTGCACGGCCACACCGAG  
GAACCGCCTGGACTAAGTCTGGATTTCCTGAAGTGCTGCAAGAGGAATACGGCCGAGGA  
AGAGAACTCTGCAAACCCTAACCAAGACCAGAACGCACGCCGAAGAAAGAAGAAGGAG  
AGACGTCTAGGGGCACCATGCAGGCTATCAACAATGAAAGAAAAGCTTCGAAGGTACT  
GGGCATCGTCTTCTTCTGTTTGTGGTGATGTGGTGCCCTTTCATCACAAACATCATGGCC  
GTCATCTGCAAAGAGTCTGCAATGAGGATGTCATTGGGGCCCTGCTCAATGTGTTTGTGG  
ATCGGTTATCTCTTTCAGCAGTCAACCCACTAGTCTATACTCTGTTCAACAAAATTTACCGA  
AGGGCATTCTCCAACCTATTGCGTTGCAATTATAAGGTAGAGAAAAGCCTCCTGTCAG  
GCAGATTCCAAGAGTTGCCGCCACTGCTTTGTCTGGGAGGGAGCTTAATGTTAACATTT  
ATCGGCATACCAATGAACCGGTGATCGAGAAAGCCAGTGACAATGAGCCCGGTATAGAG  
ATGCAAGTTGAGAATTTAGAGTTACAGTAAATCCCTCCAGTGTGGTTAGCGAAAGGAT  
TAGCAGTGTGTGA

*FIG. 7A*


ATGGATATTCTTTGTGAAGAAAATACTTCTTTGAGCTCAACTACGAACTCCCTAATGCAATTA  
AATGATGACAACAGGCTCTACAGTAATGACTTTAACTCCGGAGAAGCTAACACTTCTGATGCA  
TTAACTGGACAGTCGACTCTGAAAATCGAACCAACCTTTCTGTGAAGGGTGCCCTTCACCG  
TCGTGTCTCTCCTTACTTCATCTCCAGGAAAAAACTGGTCTGCTTACTGACAGCCGTAGTGA  
TTATTCTAACTATTGCTGGAAACATACTCGTCATCATGGCAGTGTCCCTAGAGAAAAGCTGC  
AGAATGCCACCAACTATTTCTGATGTCACTTGCCATAGCTGATATGCTGCTGGGTTTCCTTGT  
CATGCCCGTGTCCATGTTAACCATCCTGTATGGGTACCGGTGGCCTCTGCCGAGCAAGCTTTGT  
GCAGTCTGGATTTACCTGGACGTGCTCTTCTCCACGGCCTCCATCATGCACCTCTGCGCCATCT  
CGCTGGACCGCTACGTGCCATCCAGAATCCATCCACCACAGCCGCTTCAACTCCAGAACTA  
AGGCATTTCTGAAAATCATTGCTGTTTGGACCATATCAGTAGGTATATCCATGCCAATACCAG  
TCTTTGGGCTACAGGACGATTCGAAGGTCTTTAAGGAGGGGAGTTGCTTACTCGCCGATGATA  
ACTTTGTCTGATCGGCTCTTTTGTGTCATTTTTCATTTCCCTGACGATTATGGTGATTACGT  
ATTGCCTGACCATCTACGTTCTGCGCCGACAAGCTTTGATGTTACTGCACGGCCACACC  
GAGGAACCGCCTGGACTAAGTCTGGATTTCTGAAGTGCTGCAAGAGGAATACGGCCGA  
GGAAGAGAACTCTGCAAACCCTAACCAAGACCAGAACGCACGCCGAAGAAAGAAGAAG  
GAGAGACGTCCTAGGGGCACCATGCAGGCTATCAACAATGAAAGAAAAGCTAAGAAAGT  
CCTTGGGATTTGTTTCTTTGTGTTTCTGATCATGTGGTGCCCTTTCTTCATCACAAACATCA  
TGGCCGTCATCTGCAAAGAGTCTTGC AATGAGGATGTCATTTGGGGCCCTGCTCAATGTGTTT  
TTTGATCGGTTATCTCTTTCAGCAGTCAACCCACTAGTCTATACTCTGTTCAACAAAATTT  
ACCGAAGGGCATTCTCCA ACTATTTGCGTTGCAATTATAAGGTAGAGAAAAGCCTCCT  
GTCAGGCAGATTCCAAGAGTTGCCGCCACTGCTTTGTCTGGGAGGGAGCTTAATGTTAA  
CATTTATCGGCATACCAATGAACCGGTGATCGAGAAAGCCAGTGACAATGAGCCCGGTA  
TAGAGATGCAAGTTGAGAATTTAGAGTTACCAGTAAATCCCTCCAGTGTGGTTAGCGAA  
AGGATTAGCAGTGTGTGA

*FIG. 7B*

MDILCEENTSLSSTTNSLMQLNDDNRLYSNDFNSGEANTSDAFNWTVDSERNTNLSCEGCLSPSCL  
SLLHLQEKNSALLTAVVILTIAGNVLVMAVSLEKKLQATNYFLMSLAIADMLLGFLVMPVSM  
LTILYGYRWPLPSKLCVWIYLDVLFSTASIMHLCAISLDRYVAIQNPIHRSFNSRTKAFLKILAVW  
TISVGISMPIPVFGLQDDSKVFKEGSCLLADDNFVLIGSFVSFFIPLTMVTTYCLTIYVLRROALML  
LHGHTTEPPGLSLDFLKCKRNTAEENSANPNQDONARRRKKKERRRPRGTMQAINNERKA  
KKVLGIVFFVFLMWCFFITNMAVICKESCNEVDIGALLNVFVWIGYLSSAVNPLVYTLFNKIY  
RRAFSNYLRCNYKVEKKPPVRQIPRVAATASGRELVNVIYRHTNEPVIEKASDNEPGIEMQV  
ENLELPVNPSSVVSERISSV



*FIG. 7C*



\*Xho I (1312) to Sca I (3049) is identical to pRc/RSV Xho I (3045) to 4782.  
 \*Sca I (3049) to 4070 is identical to pCDM7 Amp Scal (2524) to 3545.  
 \*multiple cloning site includes Hind III to Sac I of pBluescript II  
 \*110 to 1312 is identical to pCMD7 Amp 76 to 1278.  
 \*Sac I and Spe I in MCS are not unique.

FIG. 8




FIG. 9




FIG. 10




FIG. 11A






FIG. 11B

*FIG. 12*



*FIG. 13*






FIG. 14





FIG. 15





*FIG. 16A*



*FIG. 16B*



*FIG. 16C*

FIG. 17A





FIG. 17B




FIG. 17C

**NON-ENDOGENOUS, CONSTITUTIVELY  
ACTIVATED HUMAN SEROTONIN  
RECEPTORS AND SMALL MOLECULE  
MODULATORS THEREOF**

This is a divisional of U.S. Ser. No. 09/292,072, filed Apr. 14, 1999, which in turn is a continuation-in-part of U.S. Ser. No. 09/060,188, filed Apr. 14, 1998 (owned by Arena Pharmaceuticals, Inc.), and claims priority to U.S. Provisional Application No. 60/090,783, filed Jun. 26, 1998 (owned by Arena Pharmaceuticals, Inc.), now abandoned, U.S. Provisional Application No. 60/112,909, filed Dec. 18, 1998, now abandoned and U.S. Provisional Application No. 60/123,000, filed Mar. 5, 1999, now abandoned. All of the foregoing patents and patent applications are incorporated in their entirety by reference.

**FIELD OF THE INVENTION**

The present invention relates to non-endogenous, constitutively active serotonin receptors and small molecule modulators thereof.

**BACKGROUND OF THE INVENTION**

**I. G Protein-coupled Receptors**

G protein-coupled receptors share a common structural motif. All these receptors have seven sequences of between 22 to 24 hydrophobic amino acids that form seven alpha helices, each of which spans the membrane. The transmembrane helices are joined by strands of amino acids having a larger loop between the fourth and fifth transmembrane helix on the extracellular side of the membrane. Another larger loop, composed primarily of hydrophilic amino acids, joins transmembrane helices five and six on the intracellular side of the membrane. The carboxy terminus of the receptor lies intracellularly with the amino terminus in the extracellular space. It is thought that the loop joining helices five and six, as well as, the carboxy terminus, interact with the G protein. Currently, Gq, Gs, Gi, and Go are G proteins that have been identified. The general structure of G protein-coupled receptors is shown in FIG. 1.

Under physiological conditions, G protein-coupled receptors exist in the cell membrane in equilibrium between two different states or conformations: an "inactive" state and an "active" state. As shown schematically in FIG. 2, a receptor in an inactive state is unable to link to the intracellular transduction pathway to produce a biological response. Changing the receptor conformation to the active state allows linkage to the transduction pathway and produces a biological response.

A receptor may be stabilized in an active state by an endogenous ligand or an exogenous agonist ligand. Recent discoveries such as, including but not exclusively limited to, modifications to the amino acid sequence of the receptor provide means other than ligands to stabilize the active state conformation. These means effectively stabilize the receptor in an active state by simulating the effect of a ligand binding to the receptor. Stabilization by such ligand-independent means is termed "constitutive receptor activation."

**II. Serotonin Receptors**

Receptors for serotonin (5-hydroxytryptamine, 5-HT) are an important class of G protein-coupled receptors. Serotonin is thought to play a role in processes related to learning and memory, sleep, thermoregulation, mood, motor activity, pain, sexual and aggressive behaviors, appetite, neurodegenerative regulation, and biological rhythms. Not surprisingly, serotonin is linked to pathophysiological conditions such as anxiety, depression, obsessive-compulsive

disorders, schizophrenia, suicide, autism, migraine, emesis, alcoholism and neurodegenerative disorders.

Serotonin receptors are divided into seven subfamilies, referred to as 5-HT1 through 5-HT7, inclusive. These subfamilies are further divided into subtypes. For example, the 5-HT2 subfamily- is divided into three receptor subtypes: 5-HT2A, 5-HT2B, and 5-HT2C. The human 5-HT2C receptor was first isolated and cloned in 1987, and the human 5-HT2A receptor was first isolated and cloned in 1990. These two receptors are thought to be the site of action of hallucinogenic drugs. Additionally, antagonists to the 5-HT2A and 5-HT2C receptors are believed to be useful in treating depression, anxiety, psychosis and eating disorders.

U.S. Pat. No. 4,985,352, describes the isolation, characterization, and expression of a functional cDNA clone encoding the entire human 5-HT1C receptor (now known as the 5HT2C receptor). U.S. Pat. No. 5,661,0124 describes the isolation, characterization, and expression of a functional cDNA clone encoding the entire human 5-HT2A receptor.

Mutations of the endogenous forms of the rat 5-HT2A and rat 5-HT2C receptors have been reported to lead to constitutive activation of these receptors (5-HT2A: Casey, C. et al. (1996) *Society for Neuroscience Abstracts*, 22:699.10, hereinafter "Casey"; 5-HT2C: Herrick-Davis, K., and Teitler, M. (1996) *Society for Neuroscience Abstracts*, 22:699.18, hereinafter "Herrick-Davis 1"; and Herrick-Davis, K. et al (1997) *J.Neurochemistry* 69(3): 1138, hereinafter "Herrick-Davis-2"). Casey describes a mutation of the cysteine residue at position 322 of the rat 5-HT2A receptor to lysine (C322K), glutamine (C322Q) and arginine (C322R) which reportedly led to constitutive activation. Herrick-Davis 1 and Herrick-Davis 2 describe mutations of the serine residue at position 312 of the rat 5-HT2C receptor to phenylalanine (S312F) and lysine (S312K), which reportedly led to constitutive activation.

**SUMMARY OF THE INVENTION**

The present invention relates to non-endogenous, constitutively activated forms of the human 5-HT2A and human 5-HT2C receptors and various uses of such receptors. Further disclosed are small molecule modulators of these receptors. Most preferably, these modulators have inverse agonist characteristics at the receptor.

More specifically, the present invention discloses nucleic acid molecules and the proteins for three non-endogenous, constitutively activated human serotonin receptors, referred to herein as, AP-1, AP-3, and AP4. The AP-1 receptor is a constitutively active form of the human 5-HT2C receptor created by an S310K point mutation. The AP-3 receptor is a constitutively active form of the human 5-HT2A receptor whereby the intracellular loop 3 (IC3) portion and the cytoplasmic-tail portion of the endogenous human 5-HT2A receptor have been replaced with the IC3 portion and the cytoplasmic-tail portion of the human 5-HT2C receptor. The AP-4 receptor is a constitutively active form of the human 5-HT2A receptor whereby (1) the region of the intracellular third loop between the proline of the transmembrane 5 region (TM5) and the proline of TM6 of the endogenous human 5-HT2A receptor has been replaced with the corresponding region of the human 5-HT2C receptor (including a S310K point mutation); and (2) the cytoplasmic-tail portion of the endogenous human 5-HT2A receptor has been replaced with the cytoplasmic-tail portion of the endogenous human 5-HT2C receptor.

The invention also provides assays that may be used to directly identify candidate compounds as agonists, partial agonists or inverse agonists to non-endogenous, constitu-

tively activated human serotonin receptors; such candidate compounds can then be utilized in pharmaceutical composition(s) for treatment of diseases and disorders which are related to the human 5-HT2A and/or human 5-HT2C receptors.

These and other aspects of the invention disclosed herein will be set forth in greater details as the patent disclosure proceeds.

The subject matter of the present application is related to the subject matter of U.S. Pat. No. 6,150,393, issued on Nov. 21, 2000; U.S. Pat. No. 6,140,509, issued on Oct. 31, 2000, and U.S. Pat. No. 6,107,324, issued Aug. 22, 2000. All of the foregoing patents and patent applications are incorporated in their entirety by reference

### BRIEF DESCRIPTION OF THE DRAWINGS

In the following figures, bold typeface indicates the location of the mutation in the non-endogenous, constitutively activated receptor relative to the corresponding endogenous receptor.

FIG. 1 shows a generalized structure of a G protein-coupled receptor with the numbers assigned to the transmembrane helices, the intracellular loops, and the extracellular loops.

FIG. 2 schematically shows the active and inactive states for a typical G protein-coupled receptor and the linkage of the active state to the second messenger transduction pathway.

FIG. 3a provides the nucleic acid sequence of the endogenous human 5-HT2A receptor (SEQ.ID.NO: 24).

FIG. 3b provides the corresponding amino acid sequence of the endogenous human 5-HT2A receptor (SEQ.ID.NO: 25).

FIG. 4a provides the nucleic acid sequence of the endogenous human 5-HT2C receptor (SEQ.ID.NO: 26).

FIG. 4b provides the corresponding amino acid sequence of the endogenous human 5-HT2C receptor (SEQ.ID.NO: 27).

FIG. 5a provides the nucleic acid sequence of a constitutively active form of the human 5-HT2C receptor ("AP-1 cDNA"—SEQ.ID.NO: 28).

FIG. 5b provides the corresponding amino acid sequence of the AP-1 cDNA ("AP-1"—SEQ.ID.NO: 29).

FIG. 6a provides the nucleic acid sequence of a constitutively active form of the human 5-HT2A receptor whereby the IC3 portion and the cytoplasmic-tail portion of the endogenous 5-HT2A receptor have been replaced with the IC3 portion and the cytoplasmic-tail portion of the human 5-HT2C receptor ("AP-3 cDNA"—SEQ.ID.NO: 30).

FIG. 6b provides the corresponding amino acid sequence of the AP-3 cDNA ("AP-3"—SEQ.ID.NO: 31).

FIG. 6c provides a schematic representation of AP-3, where the dashed-lines represent the portion obtained from the human 5-HT2C receptor.

FIG. 7a provides the nucleic acid sequence of a constitutively active form of the human 5-HT2A receptor whereby (1) the region of the between the proline of TM5 and the proline of TM6 of the endogenous human 5-HT2A receptor has been replaced with the corresponding region of the human 5-HT2C receptor (including a S310K point mutation); and (2) the cytoplasmic-tail portion of the endogenous 5-HT2A receptor has been replaced with the cytoplasmic-tail portion of the endogenous human 5-HT2C receptor ("AP-4 cDNA"—SEQ.ID.NO:32).

FIG. 7b provides the corresponding amino acid sequence of the AP4 cDNA ("AP-4"—SEQ.ID.NO: 33).

FIG. 7c provides a schematic representation of the mutated 5-HT2A receptor of FIG. 7b where the dashed-lines represent the portion obtained from the human 5-HT2C receptor.

FIG. 8 is a representation of the preferred vector, pCMV, used herein.

FIG. 9 is a diagram illustrating (1) enhanced [<sup>35</sup>S]GTPγS binding to membranes prepared from COS cells expressing the endogenous human 5-HT2C receptor in response to serotonin, and (2) inhibition by mianserin using wheatgerm agglutinin scintillation proximity beads. The concentration of [<sup>35</sup>S]GTPγS was held constant at 0.3 nM, and the concentration of GDP was held at 1 μM. The concentration of the membrane protein was 12.5 μg.

FIG. 10 is a diagram showing serotonin stimulation of [<sup>35</sup>S]GTPγS binding to membranes expressing AP-1 receptors in 293T cells and the inhibition by 30 μM mianserin on Wallac™ Scintistrips.

FIGS. 11A and 11B are diagrams showing the effects of protein concentration on [<sup>35</sup>S]GTPγS binding in membranes prepared from 293T cells transfected with the endogenous human 5-HT2C receptors and AP-1 receptors compared to cells transfected with the control vector (pCMV) alone in the absence (A) and presence (B) of 10 μM serotonin. The radiolabeled concentration of [<sup>35</sup>S]GTPγS was held constant at 0.3 nM, and the GDP concentration was held constant at 1 μM. The assay was performed on 96-well format on Wallac™ scintistrips.

FIG. 12 provides bar-graph comparisons of inositol triphosphate ("IP3") production between the endogenous human 5HT2A receptor and AP-2, a mutated form of the receptor.

FIG. 13 provides bar-graph comparisons of inositol triphosphate ("IP1") production between the endogenous human 5HT2A receptor and AP-4, a mutated form of the receptor.

FIG. 14 provides bar graph comparisons of IP3 production between the endogenous human 5-HT2A receptor and AP-3, a mutated form of the receptor.

FIG. 15 provides bar-graph comparisons of IP3 production between the endogenous human 5-HT2C receptor and AP-1.

FIGS. 16A–C provides representative autoradiograms showing displacement of I<sup>125</sup>-LSD from brain sections by spiperone and compound 116100.

FIGS. 17A–C shows in vivo response of animals to 116102 exposure.

### DEFINITIONS

The scientific literature that has evolved around receptors has adopted a number of terms to refer to ligands having various effects on receptors. For clarity and consistency, the following definitions will be used throughout this patent document. To the extent that these definitions conflict with other definitions for these terms, the following definitions shall control.

AGONISTS shall mean moieties that activate the intracellular response when they bind to the receptor, or enhance GTP binding to membranes.

AMINO ACID ABBREVIATIONS used herein are set out in Table 1:

TABLE 1

ALANINE	ALA	A
ARGININE	ARG	R
ASPARAGINE	ASN	N
ASPARTIC ACID	ASP	D
CYSTEINE	CYS	C
GLUTAMIC ACID	GLU	E
GLUTAMINE	GLN	Q
GLYCINE	GLY	G
HISTIDINE	HIS	H
ISOLEUCINE	ILE	I
LEUCINE	LEU	L
LYSINE	LYS	K
METHIONINE	MET	M
PHENYLALANINE	PHE	F
PROLINE	PRO	P
SERINE	SER	S
THREONINE	THR	T
TRYPTOPHAN	TRP	W
TYROSINE	TYR	Y
VALINE	VAL	V

PARTIAL AGONISTS shall mean moieties which activate the intracellular response when they bind to the receptor to a lesser degree/extent than do agonists, or enhance GTP binding to membranes to a lesser degree/extent than do agonists.

ANTAGONIST shall mean moieties that competitively bind to the receptor at the same site as the agonists but which do not activate the intracellular response initiated by the active form of the receptor, and can thereby inhibit the intracellular responses by agonists or partial agonists. ANTAGONISTS do not diminish the baseline intracellular response in the absence of an agonist or partial agonist.

CANDIDATE COMPOUND shall mean a molecule (for example, and not limitation, a chemical compound) which is amenable to a screening technique.

COMPOUND EFFICACY shall mean a measurement of the ability of a compound to inhibit or stimulate receptor functionality, as opposed to receptor binding affinity.

CONSTITUTIVELY ACTIVATED RECEPTOR shall mean a receptor subject to constitutive receptor activation.

CONSTITUTIVE RECEPTOR ACTIVATION shall mean stabilization of a receptor in the active state by means other than binding of the receptor with its endogenous ligand or a chemical equivalent thereof.

CONTACT or CONTACTING shall mean bringing at least two moieties together, whether in an in vitro system or an in vivo system.

ENDOGENOUS shall mean a material that a mammal naturally produces. ENDOGENOUS in reference to, for example and not limitation, the term "receptor" shall mean that which is naturally produced by a mammal (for example, and not limitation, a human) or a virus.

In contrast, the term NON-ENDOGENOUS in this context shall mean that which is not naturally produced by a mammal (for example, and not limitation, a human) or a virus. For example, and not limitation, a receptor which is not constitutively active in its endogenous form, but when manipulated becomes constitutively active, is most preferably referred to herein as a "non-endogenous, constitutively activated receptor." Both terms can be utilized to describe both "in vivo" and "in vitro" systems. For example, and not a limitation, in a screening approach, the endogenous or non-endogenous receptor may be in reference to an in vitro screening system. As a further example and not limitation, where the genome of a mammal has been manipulated to

include a non-endogenous constitutively activated receptor, screening of a candidate compound by means of an in vivo system is viable.

INHIBIT or INHIBITING, in relationship to the term "response" shall mean that a response is decreased or prevented in the presence of a compound as opposed to in the absence of the compound.

INVERSE AGONISTS shall mean moieties that bind the endogenous form of the receptor or to the constitutively activated form of the receptor, and which inhibit the baseline intracellular response initiated by the active form of the receptor below the normal base level of activity which is observed in the absence of agonists or partial agonists, or decrease GTP binding to membranes. Preferably, the baseline intracellular response is inhibited in the presence of the inverse agonist by at least 30%, more preferably by at least 50%, and most preferably by at least 75%, as compared with the baseline response in the absence of the inverse agonist.

LIGAND shall mean an endogenous, naturally occurring molecule specific for an endogenous, naturally occurring receptor.

PHARMACEUTICAL COMPOSITION shall mean a composition comprising at least one active ingredient, whereby the composition is amenable to investigation for a specified, efficacious outcome in a mammal (for example, and not limitation, a human). Those of ordinary skill in the art will understand and appreciate the techniques appropriate for determining whether an active ingredient has a desired efficacious outcome based upon the needs of the artisan.

STIMULATE or STIMULATING, in relationship to the term "response" shall mean that a response is increased in the presence of a compound as opposed to in the absence of the compound.

## DETAILED DESCRIPTION

### I. Particularly Preferred Mutations

For convenience, the sequence information regarding the non-endogenous, constitutively active human 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptors are referred to by identifiers as set forth in Table 2:

TABLE 2

IDENTIFIER	RECEPTOR	SEQ. ID. NO:	FIGURE
AP-1 cDNA	5-HT <sub>2C</sub>	28	5a
AP-1	5-HT <sub>2C</sub>	29	5b
AP-3 cDNA	5-HT <sub>2A</sub>	30	6a
AP-3	5-HT <sub>2A</sub>	31	6b
AP-4 cDNA	5-HT <sub>2A</sub>	32	7a
AP-4	5-HT <sub>2A</sub>	33	7b

As will be discussed in greater detail below, a mutation analogous to that reported by Casey (C322K) was utilized in the human 5-HT<sub>2A</sub> receptor and is referred to herein as AP-2. However, AP-2 did not lead to sufficient constitutive activation to allow for utilization in screening techniques.

### II. Introduction

While it is sometimes possible to make predictions as to the effect of nucleic acid manipulation from one species to another, this is not always the case. The results reported by Casey suggest that a point mutation in the rat 5-HT<sub>2A</sub> receptor evidences constitutive activation of the mutated receptor. Casey reports that the C322K mutation was approximately four fold more active than the native rat 5-HT<sub>2A</sub> receptor. However, for purposes of a most preferred use, i.e., screening of candidate compounds, this corresponding mutation in the human 5-HT<sub>2A</sub> receptor had little

discernable effect in evidencing constitutive activation of the human receptor. This, of course, creates the reasonable conclusion that the information reported in Herrick-Davis 1 or Herrick-Davis 2 is of limited predictive value relative to the manipulation of the human 5-HT<sub>2C</sub> receptor. Consequently, the ability to make reasonable predictions about the effects of mutations to the rat 5-HT receptors vis-à-vis the corresponding human receptors is not possible. Nonetheless, this unfortunate lack of reasonable predictability provides the opportunity for others to discover mutations to the human 5-HT receptors that provide evidence of constitutive activation.

Therefore, the present invention is based upon the desire of defining mutated sequences of the human serotonin receptors 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> whereby such mutated versions of the expressed receptor are constitutively active. These constitutively active receptors allow for, inter alia, screening candidate compounds.

What has been discovered and disclosed herein is that substantial activation of the human 5-HT<sub>2A</sub> receptor can be obtained by "domain swapping," i.e., by switching the third intracellular domain of the 5-HT<sub>2A</sub> receptor with the third intracellular domain of the 5-HT<sub>2C</sub> receptor. Additionally, swapping the cytoplasmic tail of the two receptors further increases the IP<sub>3</sub> response. Furthermore, mutation of the serine at position 310 to lysine (S310K) of the human 5-HT<sub>2C</sub> receptor leads to constitutive activation.

What follows is a most preferred approach to identification of candidate compounds; those in the art will readily appreciate that the particular order of screening approaches, and/or whether or not to utilize certain of these approaches, is a matter of choice. Thus, the order presented below, set for presentational efficiency and for indication of the most preferred approach utilized in screening candidate compounds, is not intended, nor is to be construed, as a limitation on the disclosure, or any claims to follow.

### III. Generic G Protein-Coupled Receptor Screening Assay Techniques

When a G protein receptor becomes constitutively active, it binds to a G protein (G<sub>q</sub>, G<sub>s</sub>, G<sub>i</sub>, G<sub>o</sub>) and stimulates the binding of GTP to the G protein. The G protein then acts as a GTPase and slowly hydrolyzes the GTP to GDP, whereby the receptor, under normal conditions, becomes deactivated. However, constitutively activated receptors continue to exchange GDP to GTP. A non-hydrolyzable analog of GTP, [<sup>35</sup>S]GTP $\gamma$ S, can be used to monitor enhanced binding to membranes which express constitutively activated receptors. It is reported that [<sup>35</sup>S]GTP $\gamma$ S can be used to monitor G protein coupling to membranes in the absence and presence of ligand. An example of this monitoring, among other examples well-known and available to those in the art, was reported by Traynor and Nahorski in 1995. The preferred use of this assay system is for initial screening of candidate compounds because the system is generically applicable to all G protein-coupled receptors regardless of the particular G protein that interacts with the intracellular domain of the receptor.

### IV. Confirmation of G Protein-Coupled Receptor Site Screening Assay Techniques

Once candidate compounds are identified using the "generic" G protein-coupled receptor assay (i.e. an assay to select compounds that are agonists, partial agonists, or inverse agonists), further screening to confirm that the compounds have interacted at the receptor site is preferred. For example, a compound identified by the "generic" assay may not bind to the receptor, but may instead merely "uncouple" the G protein from the intracellular domain.

Thus, by further screening those candidate compounds, which have been identified using a "generic" assay in an agonist and/or antagonist competitive binding assay, further refinement in the selection process is provided.

Lysergic acid diethylamide (LSD) is a well-known agonist of the 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptors, while mesulergine is a well-known antagonist to the 5-HT<sub>2C</sub> receptor. Accordingly, in most preferred embodiments, an agonist (LSD) and/or antagonist (mesulergine) competitive binding assay(s) is used to further screen those compounds selected from the "generic" assay for confirmation of serotonin receptor binding.

### V. Specified G Protein Assay Techniques

The art-accepted physiologically mediated pathway for the human 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptors is via G<sub>q</sub>. Intracellular accumulation of IP<sub>3</sub> can be used to confirm constitutive activation of these types of G<sub>q</sub> coupled receptors (see Herrick-Davis-1). As a result, "IP<sub>3</sub> accumulation" assays can be used to further screen those compounds selected from an agonist and/or antagonist competitive binding assay.

### VI. Pharmaceutical Compositions

Candidate compounds selected for further development can be formulated into pharmaceutical compositions using techniques well known to those in the art. Suitable pharmaceutically-acceptable carriers are available to those in the art; for example, see Remington's Pharmaceutical Sciences, 16<sup>th</sup> Edition, 1980, Mack Publishing Co., (Oslo et al., eds.)

## EXAMPLES

The following examples are presented for purposes of elucidation, and not limitation, of the present invention. While specific nucleic acid and amino acid sequences are disclosed herein, those of ordinary skill in the art are credited with the ability to make minor modifications to these sequences while achieving the same or substantially similar results reported below. It is intended that equivalent, non-endogenous, constitutively activated human serotonin receptor sequences having eighty-five percent (85%) homology, more preferably having ninety percent (90%) homology, and most preferably having ninety-five percent (95%) homology to the disclosed and claimed sequences all fall within the scope of any claims appended hereto.

### Example 1

#### Generation of Non-Endogenous, Constitutively Activated Human Serotonin Receptors 5-HT<sub>2C</sub> and 5-HT<sub>2A</sub>

##### A. Construction of Constitutively Active 5-HT<sub>2C</sub> Receptor cDNA

###### 1. Endogenous Human 5-HT<sub>2C</sub>

The cDNA encoding endogenous human 5-HT<sub>2C</sub> receptor was obtained from human brain poly-A<sup>+</sup> RNA by RT-PCR. The 5' and 3' primers were derived from the 5' and 3' untranslated regions and contained the following sequences:

5'-GACCTCGAGGTTGCTTAAGACTGAAGCA-3'  
(SEQ.ID.NO:1)

5'-ATTTCTAGACATATGTAGCTTGTACCGT-3'  
(SEQ.ID.NO:2)

PCR was performed using either TaqPlus<sup>TM</sup> precision polymerase (Stratagene) or rTth<sup>TM</sup> polymerase (Perkin Elmer) with the buffer systems provided by the manufacturers, 0.25  $\mu$ M of each primer, and 0.2 mM of each of the four (4) nucleotides. The cycle condition was 30 cycles of 94° C. for 1 minute, 57° C. for 1 minute and 72° C. for 2 minutes. The

1.5 kb PCR fragment was digested with Xho I and Xba I and subcloned into the Sal I-Xba I site of pBluescript.

The derived cDNA clones were fully sequenced and found to correspond to published sequences.

### 2. AP1 cDNA

The cDNA containing a S310K mutation (AP-1 cDNA) in the third intracellular loop of the human 5-HT<sub>2C</sub> receptor was constructed by replacing the Sty I restriction fragment containing amino acid 310 with synthetic double stranded oligonucleotides encoding the desired mutation. The sense strand sequence utilized had the following sequence:

5'-CTAGGGGCACCATGCAGGCTATCAACAATGAAA  
GAAAAGCTAAGAAAGTC-3' (SEQ. ID.NO: 3)

and the antisense strand sequence utilized had the following sequence:

5'-CAAGGACTTTCTTAGCTTTTCTTTCATTGTTGA  
TAGCCTGCATGGT GCCC-3' (SEQ. ID.NO: 4).

### B. Construction of Constitutively Active 5-HT<sub>2A</sub> Receptor cDNA

#### 1. Endogenous Human 5-HT<sub>2A</sub>

The cDNA encoding endogenous human 5-HT<sub>2A</sub> receptor was obtained by RT-PCR using human brain poly-A<sup>+</sup> RNA; a 5' primer from the 5' untranslated region with a Xho I restriction site:

5'-GACCTCGAGTCCTTCTACACCTCATC-3'  
(SEQ.ID.NO:5)

and a 3' primer from the 3' untranslated region containing an Xba I site:

5'-TGCTCTAGATTCCAGATAGGTGAAAAGTTG-3'  
(SEQ.ID.NO:6).

PCR was performed using either TaqPlus<sup>TM</sup> precision polymerase (Stratagene) or rTth<sup>TM</sup> polymerase (Perkin Elmer) with the buffer systems provided by the manufacturers, 0.25 μM of each primer, and 0.2 mM of each of the four (4) nucleotides. The cycle condition was 30 cycles of 94° C. for 1 minute, 57° C. for 1 minute and 72° C. for 2 minutes. The 1.5 kb PCR fragment was digested with Xba I and subcloned into the Eco RV-Xba I site of pBluescript.

The resulting cDNA clones were fully sequenced and found to encode two amino acid changes from the published sequences. The first change is a T25N mutation in the N-terminal extracellular domain and the second change is an H452Y mutation. These mutations are likely to represent sequence polymorphisms rather than PCR errors since the cDNA clones having the same two mutations were derived from two independent PCR procedures using Taq polymerase from two different commercial sources (TaqPlus<sup>TM</sup> Stratagene and rTth<sup>TM</sup> Perkin Elmer).

#### 2. Human 5-HT<sub>2A</sub> (C322K; AP-2)

The cDNA containing the point mutation C322K in the third intracellular loop was constructed by using the Sph I restriction enzyme site, which encompasses amino acid 322. For the PCR procedure, a primer containing the C322K mutation:

5'-CAAAGAAAGTACTGGGCATCGTCTTCTTCCT-3'  
(SEQ.ID.NO:7)

was used along with the primer from the 3' untranslated region set forth above as SEQ.ID.NO:6. The resulting PCR fragment was then used to replace the 3' end of the wild type 5-HT<sub>2A</sub> cDNA by the T4 polymerase blunted Sph I site. PCR was performed using pfu polymerase (Stratagene) with the buffer system provided by the manufacturer and 10% DMSO, 0.25 mM of each primer, 0.5 mM of each of the 4 nucleotides. The cycle conditions were 25 cycles of 94° C. for 1 minute, 60° C. for 1 minute and 72° C. for 1 minute.

#### 3. AP-3 cDNA

The human 5-HT<sub>2A</sub> cDNA with intracellular loop 3 (IC3) or IC3 and cytoplasmic tail replaced by the corresponding human 5-HT<sub>2C</sub> cDNA was constructed using PCR-based mutagenesis.

#### (a) Replacement of IC3 Loop

The IC3 loop of human 5-HT<sub>2A</sub> cDNA was first replaced with the corresponding human 5-HT<sub>2C</sub> cDNA. Two separate PCR procedures were performed to generate the two fragments, Fragment A and Fragment B, that fuse the 5-HT<sub>2C</sub> IC3 loop to the transmembrane 6 (TM6) of 5-HT<sub>2A</sub>. The 237 bp PCR fragment, Fragment A, containing 5-HT<sub>2C</sub> IC3 and the initial 13 bp of 5-HT<sub>2A</sub> TM6 was amplified by using the following primers:

5'-CCGCTCGAGTACTGCGCCGACAAGCTTTGAT-3'  
(SEQ.ID.NO:8)

5'-CGATGCCCAGCACTTTCGAAGCTTTTCTTTCAT  
TGTTG3'(SEQ.ID.NO:9)

The template used was human 5-HT<sub>2C</sub> cDNA.

The 529 bp PCR fragment, Fragment B, containing the C-terminal 13 bp of IC3 from 5-HT<sub>2C</sub> and the C-terminal of 5-HT<sub>2A</sub> starting at beginning of TM6, was amplified by using the following primers:

5'-AAAAGCTTCGAAAGTGCTGGGCATCGTCTTCTT  
CCT-3' (SEQ.ID.NO:10)

5'-TGCTCTAGATTCCAGATAGGTGAAAAGTTG-3'  
(SEQ.ID.NO: 11)

The template used was human 5-HT<sub>2A</sub> cDNA.

Second round PCR was performed using Fragment A and Fragment B as co-templates with SEQ.ID.NO:8 and SEQ.ID.NO:11 (it is noted that the sequences for SEQ.ID.NOS.: 6 and 11 are the same) as primers. The resulting 740 bp PCR fragment, Fragment C, contained the IC3 loop of human 5-HT<sub>2C</sub> fused to TM6 through the end of the cytoplasmic tail of human 5-HT<sub>2A</sub>. PCR was performed using pfu<sup>TM</sup> polymerase (Stratagene) with the buffer system provided by the manufacturer, and 10% DMSO, 0.25 mM of each primer, and 0.5 mM of each of the four (4) nucleotides. The cycle conditions were 25 cycles of 94° C. for 1 minute, 57° C. (1st round PCR) or 60° C. (2nd round PCR) for 1 minute. and 72° C. for 1 minute (1st round PCR) or 90 seconds. (2nd round PCR).

To generate a PCR fragment containing a fusion junction between the human 5-HT<sub>2A</sub> TM5 and the IC3 loop of 5-HT<sub>2C</sub>, four (4) primers were used. The two external primers, derived from human 5-HT<sub>2A</sub>, had the following sequences:

5'-CGTGTCTCTCCTTACTTCA-3' (SEQ.ID.NO:12)

The other primer used was SEQ.ID.NO.6 (see note above regarding SEQ.ID.NOS. 6 and 11). The first internal primer utilized was an antisense strand containing the initial 13 bp of IC3 of 5-HT<sub>2C</sub> followed by the terminal 23 bp derived from TM5 of 5-HT<sub>2A</sub>:

5'-TCGGCGCAGTACTTTGATAGTTAGAAAGTAGGT  
GAT-3' (SEQ.ID.NO:13)

The second internal primer was a sense strand containing the terminal 14 bp derived from TM5 of 5-HT<sub>2A</sub> followed by the initial 24 bp derived from IC3 of 5-HT<sub>2C</sub>:

5'-TTCTAACTATCAAAGTACTGCGCCGACAAGCT  
TTGATG-3' (SEQ.ID.NO:14).

PCR was performed using endogenous human 5-HT<sub>2A</sub> and a co-template, Fragment C, in a 50 ml reaction volume containing 1xpfu buffer, 10% DMSO, 0.5 mM of each of the four (4) nucleotides, 0.25 mM of each external primer (SEQ.ID.NOS. 11 and 12), 0.06 mM of each internal primer (SEQ.ID.NOS. 13 and 14) and 1.9 units of pfu polymerase (Stratagene). The cycle conditions were 25 cycles of 94° C. for 1 minute, 52° C. for 1 minute and 72° C. for 2 minutes and 10 seconds. The 1.3 kb PCR product was then gel purified and digested with Pst I and Eco RI. The resulting 1 kb PstI-Eco RI fragment was used to replace the corresponding fragment in the endogenous human 5-HT<sub>2A</sub> sequence to generate the mutant 5-HT<sub>2A</sub> sequence encoding the IC3 loop of 5-HT<sub>2C</sub>.

## (b) Replacement of the Cytoplasmic Tail

To replace the cytoplasmic tail of 5-HT2A with that of 5-HT2C, PCR was performed using a sense primer containing the C-terminal 22 bp of TM7 of endogenous human 5-HT2A followed by the initial 21 bp of the cytoplasmic tail of endogenous human 5-HT2C:

5'-TTCAGCAGTCAACCCACTAGTCTATACTCTGTTC AACAAAATT-3' (SEQ.ID.NO:15)

The antisense primer was derived from the 3' untranslated region of endogenous human 5-HT2C:

5'-ATTTCTAGACATATGTAGCTTGTACCGT-3' (SEQ.ID.NO:16).

The resulting PCR fragment, Fragment D, contained the last 22 bp of endogenous human 5-HT2A TM7 fused to the cytoplasmic tail of endogenous human 5-HT2C. Second round PCR was performed using Fragment D and the co-template was endogenous human 5-HT2A that was previously digested with Acc I to avoid undesired amplification. The antisense primer used was SEQ.ID.NO:16 (the sequences for SEQ.ID.NOS. 16 and 2 are the same) and the sense primer used was derived from endogenous human 5-HT2A:

5'-ATCACCTACTTTCTAACTA-3' (SEQ.ID.NO:17).

PCR conditions were as set forth in Example 1B3.(a) for the first round PCR, except that the annealing temperature was 48° C. and the extension time was 90 seconds. The resulting 710 bp PCR product was digested with Apa I and Xba I and used to replace the corresponding Apa I-Xba I fragment of either (a) endogenous human 5-HT2A, or (b) 5-HT2A with 2C IC3 to generate (a) endogenous human 5-HT2A with endogenous human 5-HT2C cytoplasmic tail and (b) AP-3, respectively.

## 4. AP-4 cDNA

This mutant was created by replacement of the region of endogenous human 5-HT2A from amino acid 247, the middle of TM5 right after Pro<sup>246</sup>, to amino acid 337, the middle of TM6 just before Pro<sup>338</sup>, by the corresponding region of AP-1 cDNA. For convenience, the junction in TM5 is referred to as the "2A-2C junction," and the junction in TM6 is referred to as the "2C-2A junction."

Three PCR fragments containing the desired hybrid junctions were generated. The 5' fragment of 561 bp containing the 2A-2C junction in TM5 was generated by PCR using endogenous human 5-HT2A as template, SEQ.ID.NO:12 as the sense primer, and the antisense primer was derived from 13 bp of 5-HT2C followed by 20 bp of 5-HT2A sequence: 5'-CCATAATCGTCAGGGGAATGAAAAATGACACAA-3' (SEQ.ID.NO:18)

The middle fragment of the 323 bp contains endogenous human 5-HT2C sequence derived from the middle of TM5 to the middle of TM6, flanked by 13 bp of 5-HT2A sequences from the 2A-2C junction and the 2C-2A junction. This middle fragment was generated by using AP-1 cDNA as a template, a sense primer containing 13 bp of 5-HT2A followed by 20 bp of 5-HT2C sequences across the 2A-2C junction and having the sequence:

5'-ATTTTTCATTCCCCTGACGATTATGGTGATTAC-3' (SEQ.ID.NO:19);

and an antisense primer containing 13 bp of 5-HT2A followed by 20 bp of 5-HT2C sequences across the 2C-2A junction and having the sequence:

5'-TGATGAAGAAAGGGCACCACATGATCAGAAACA-3' (SEQ.ID.NO:20).

The 3' fragment of 487 bp containing the 2C-2A junction was generated by PCR using endogenous human 5-HT2A as a template and a sense primer having the following sequence from the 2C-2A junction:

5'-GATCATGTGGTGCCCTTTCTTCATCACAAACAT-3' (SEQ.ID.NO:21)

and the antisense primer was SEQ.ID.NO:6 see note above regarding SEQ.ID.NOS. 6 and 11).

Two second round PCR reactions were performed separately to link the 5' and middle fragment (5'M PCR) and the middle and 3' fragment (M3' PCR). The 5'M PCR co-template used was the 5' and middle PCR fragment as described above, the sense primer was SEQ.ID.NO:12 and the antisense primer was SEQ.ID.NO:20. The 5'M PCR procedure resulted in an 857 bp PCR fragment.

The M3' PCR used the middle and M3' PCR fragment described above as the co-template, SEQ.ID.NO: 19 as the sense primer and SEQ.ID.NO:6 (see note above regarding SEQ.ID.NOS. 6 and 11) as the antisense primer, and generated a 784 bp amplification product. The final round of PCR was performed using the 857 bp and 784 bp fragments from the second round PCR as the co-template, and SEQ.ID.NO:12 and SEQ.ID.NO: 6 (see note above regarding SEQ.ID.NOS. 6 and 11) as the sense and the antisense primer, respectively. The 1.32 kb amplification product from the final round of PCR was digested with Pst I and Eco RI. Then resulting 1 kb Pst I-Eco RI fragment was used to replace the corresponding fragment of the endogenous human 5-HT2A to generate mutant 5-HT2A with 5-HT2C: C310K/IC3. The Apa I-Xba fragment of AP3 was used to replace the corresponding fragment in mutant 5-HT2A with 5-HT2C: C310K/IC3 to generate AP4.

## Example 2

## Receptor Expression

## A. pCMV

Although a variety of expression vectors are available to those in the art, for purposes of utilization for both the endogenous and non-endogenous receptors discussed herein, it is most preferred that the vector utilized be pCMV. This vector was deposited with the American Type Culture Collection (ATCC) on Oct. 13, 1998 (10801 University Blvd., Manassas, Va. 20110-2209 USA) under the provisions of the Budapest Treaty for the International Recognition of the Deposit of Microorganisms for the Purpose of Patent Procedure. The DNA was tested by the ATCC and determined to be viable. The ATCC has assigned the following deposit number to pCMV: ATCC #203351. See FIG. 8.

## B. Transfection Procedure

For the generic assay (<sup>35</sup>S]GTPγS; Example 3) and the antagonist binding assay (mesulergine; Example 4), transfection of COS-7 or 293T cells was accomplished using the following protocol.

On day one, 5×10<sup>6</sup> COS-7 cells or 1×10<sup>7</sup> 293T cells per 150 mm plate were plated out. On day two, two reaction tubes were prepared (the proportions to follow for each tube are per plate): tube A was prepared by mixing 20 μg DNA (e.g., pCMV vector; pCMV vector AP-1 cDNA, etc.) in 1.2 ml serum free DMEM (Irvine Scientific, Irvine, Calif.); tube B was prepared by mixing 120 μl lipofectamine (Gibco BRL) in 1.2 ml serum free DMEM. Tubes A and B were then admixed by inversions (several times), followed by incubation at room temperature for 30–45 min. The admixture is referred to as the "transfection mixture". Plated COS-7 cells were washed with 1×PBS, followed by addition of 10 ml serum free DMEM. 2.4 ml of the transfection mixture was then added to the cells, followed by incubation for 4 hrs at 37° C./5% CO<sub>2</sub>. The transfection mixture was then removed



by aspiration, followed by the addition of 25 ml of DMEM/10% Fetal Bovine Serum. Cells were then incubated at 37° C./5% CO<sub>2</sub>. After 72 hr incubation, cells were then harvested and utilized for analysis.

### Example 3

#### GTP Membrane Binding Scintillation Proximity Assay

The advantages of using [<sup>35</sup>S]GTPγS binding to measure constitutive activation are that: (a) [<sup>35</sup>S]GTPγS binding is generically applicable to all G protein-coupled receptors; and (b) [<sup>35</sup>S]GTPγS binding is proximal at the membrane surface, thereby making it less likely to pick-up molecules which affect the intracellular cascade. The assay utilizes the ability of G protein-coupled receptors to stimulate [<sup>35</sup>S]GTPγS binding to membranes expressing the relevant receptors. Therefore, the assay may be used to directly screen compounds at the disclosed serotonin receptors.

FIG. 9 demonstrates the utility of a scintillation proximity assay to monitor the binding of [<sup>35</sup>S]GTPγS to membranes expressing the endogenous human 5-HT<sub>2C</sub> receptor expressed in COS cells. In brief, the assay was incubated in 20 mM HEPES, pH 7.4, binding buffer with 0.3 nM [<sup>35</sup>S]GTPγS and 12.5 μg membrane protein and 1 μM GDP for 30 minutes. Wheatgerm agglutinin beads (25 μl; Amersham) were then added and the mixture was incubated for another 30 minutes at room temperature. The tubes were then centrifuged at 1500×g for 5 minutes at room temperature and then counted in a scintillation counter. As shown in FIG. 9, serotonin, which as the endogenous ligand activates the 5-HT<sub>2C</sub> receptor, stimulated [<sup>35</sup>S]GTPγS binding to the membranes in a concentration dependant manner. The stimulated binding was completely inhibited by 30 μM mianserin, a compound considered as a classical 5-HT<sub>2C</sub> antagonist, but also known as a 5-HT<sub>2C</sub> inverse agonist.

Although this assay measures agonist-induced binding of [<sup>35</sup>S]GTPγS to membranes and can be routinely used to measure constitutive activity of receptors, the present cost of wheatgerm agglutinin beads may be prohibitive. A less costly but equally applicable alternative also meets the needs of large-scale screening. Flash plates and Wallac™ scintistrips may be used to format a high throughput [<sup>35</sup>S]GTPγS binding assay. This technique allows one to monitor the tritiated ligand binding to the receptor while simultaneously monitoring the efficacy via [<sup>35</sup>S]GTPγS binding. This is possible because the Wallace beta counter can switch energy windows to analyze both tritium and <sup>35</sup>S-labeled probes.

Also, this assay may be used for detecting of other types of membrane activation events that result in receptor activation. For example, the assay may be used to monitor <sup>32</sup>P phosphorylation of a variety of receptors (including G protein-coupled and tyrosine kinase receptors). When the membranes are centrifuged to the bottom of the well, the bound [<sup>35</sup>S]GTPγS or the <sup>32</sup>P-phosphorylated receptor will activate the scintillant coated on the wells. Use of ScintiX strips (Wallac™) demonstrate this principle. Additionally, this assay may be used for measuring ligand binding to receptors using radiolabeled ligands. In a similar manner, the radiolabeled bound ligand is centrifuged to the bottom of the well and activates the scintillant. The [<sup>35</sup>S]GTPγS assay results parallel the results obtained in traditional second messenger assays of receptors.

As shown in FIG. 10, serotonin stimulates the binding of [<sup>35</sup>S]GTPγS to the endogenous human 5-HT<sub>2C</sub> receptor, while mianserin inhibits this response. Furthermore,

mianserin acts as a partial inverse agonist by inhibiting the basal constitutive binding of [<sup>35</sup>S]GTPγS to membranes expressing the endogenous human 5-HT<sub>2C</sub> receptor. As expected, there is no agonist response in the absence of GDP since there is no GDP present to exchange for [<sup>35</sup>S]GTPγS. Not only does this assay system demonstrate the response of the native 5-HT<sub>2C</sub> receptor, but it also measures the constitutive activation of other receptors.

FIGS. 11A and FIG. 11B demonstrate the enhanced binding of [<sup>35</sup>S]GTPγS to membranes prepared from 293T cells expressing the control vector alone, the native human 5-HT<sub>2C</sub> receptor or the AP-1 receptor. The total protein concentration used in the assay affects the total amount of [<sup>35</sup>S]GTPγS binding for each receptor. The c.p.m. differential between the CMV transfected and the constitutively active mutant receptor increased from approximately 1000 c.p.m. at 10 μg/well to approximately 6–8000 c.p.m. at 75 μg/well protein concentration, as shown in FIG. 11.

The AP-1 receptor showed the highest level of constitutive activation followed by the wild type receptor, which also showed enhanced [<sup>35</sup>S]GTPγS binding above basal. This is consistent with the ability of the endogenous human 5-HT<sub>2C</sub> receptor to accumulate intracellular IP<sub>3</sub>, in the absence of 5HT stimulation (Example 5) and is also consistent with published data claiming that the endogenous human 5-HT<sub>2C</sub> receptor has a high natural basal activity. Therefore, the AP-1 receptor demonstrates that constitutive activity may be measured by proximal [<sup>35</sup>S]GTPγS binding events at the membrane interface.

### Example 4

#### Serotonin Receptor Agonist/Antagonist Competitive Binding Assay

Membranes were prepared from transfected COS-7 cells (see Example 2) by homogenization in 20 mM HEPES and 10 mM EDTA, pH 7.4 and centrifuged at 49,000×g for 15 min. The pellet was resuspended in 20 mM HEPES and 0.1 mM EDTA, pH 7.4, homogenized for 10 sec. using polytron homogenizer (Brinkman) at 5000 rpm and centrifuged at 49,000×g for 15 min. The final pellet was resuspended in 20 mM HEPES and 10 mM MgCl<sub>2</sub>, pH 7.4, homogenized for 10 sec. using polytron homogenizer (Brinkman) at 5000 rpm.

Assays were performed in triplicate 200 μl volumes in 96 well plates. Assay buffer (20 mM HEPES and 10 mM MgCl<sub>2</sub>, pH 7.4) was used to dilute membranes, <sup>3</sup>H-LSD, <sup>3</sup>H-mesulergine, serotonin (used to define non-specific for LSD binding) and mianserin (used to define non-specific for mesulergine binding). Final assay concentrations consisted of 1 nM <sup>3</sup>H-LSD or 1nM <sup>3</sup>H-mesulergine, 50 μg membrane protein and 100 μM serotonin or mianserin. LSD assays were incubated for 1 hr at 37° C., while mesulergine assays were incubated for 1 hr at room temperature. Assays were terminated by rapid filtration onto Wallac Filtemat Type B with ice cold binding buffer using Skatron cell harvester. The radioactivity was determined in a Wallac 1205 BetaPlate counter.

### Example 5

#### Intracellular IP<sub>3</sub> Accumulation Assay

For the IP<sub>3</sub> accumulation assay, a transfection protocol different from the protocol set forth in Example 2 was utilized. In the following example, the protocols used for days 1–3 were slightly different for the data generated for

FIGS. 12 and 14 and for FIGS. 13 and 15; the protocol for day 4 was the same for all conditions.

#### A. COS-7 and 293 Cells

On day one, COS-7 cells or 293 cells were  $\mu$ lated onto 24 well plates, usually  $1 \times 10^5$  cells/well or  $2 \times 10^5$  cells/well, respectively. On day two, the cells were transfected by first mixing 0.25  $\mu$ g DNA (see Example 2) in 50  $\mu$ l serum-free DMEM/well and then 2  $\mu$ l lipofectamine in 50  $\mu$ l serum-free DMEM/well. The solutions ("transfection media") were gently mixed and incubated for 15–30 minutes at room temperature. The cells were washed with 0.5 ml PBS and then 400  $\mu$ l of serum free media was mixed with the transfection media and added to the cells. The cells were then incubated for 3–4 hours at 37° C./5%CO<sub>2</sub>. Then the transfection media was removed and replaced with 1 ml/well of regular growth media. On day 3, the media was removed and the cells were washed with 0.5 ml PBS. Then 0.5 ml inositol-free/serum-free media (GIBCO BRL) was added to each well with 0.25  $\mu$ Ci of <sup>3</sup>H-myo-inositol/well and the cells were incubated for 16–18 hours overnight at 37° C./5%CO<sub>2</sub>. Protocol A.

#### B. 293 Cells

On day one,  $1 \times 10^7$  293 cells per 150mm  $\mu$ late were plated out. On day two, two reaction tubes were prepared (the proportions to follow for each tube are per plate): tube A was prepared by mixing 20  $\mu$ g DNA (e.g., pCMV vector; pCMV vector AP-1 cDNA, etc.) in 1.2 ml serum free DMEM (Irvine Scientific, Irvine, Calif.); tube B was prepared by mixing 120  $\mu$ l lipofectamine (Gibco BRL) in 1.2 ml serum free DMEM. Tubes A and B were then admixed by inversions (several times), followed by incubation at room temperature for 30–45 min. The admixture is referred to as the "transfection mixture". Plated 293 cells were washed with 1xPBS, followed by addition of 10 ml serum free DMEM. 2.4 ml of the transfection mixture was then added to the cells followed by incubation for 4 hrs at 37° C./5% CO<sub>2</sub>. On day 3, cells were trypsinized and counted, followed by plating of  $1 \times 10^6$  cells/well (poly D-lysine treated 12-well plates). Cells were permitted to adhere to the wells, followed by one wash with 1xPBS. Thereafter, 0.5  $\mu$ Ci <sup>3</sup>H-inositol in 1 ml inositol-free DMEM was added per well. Protocol B.

On day 4, the cells were washed with 0.5 ml PBS and then 0.45 ml of assay medium was added containing inositol-free/serum free media, 10  $\mu$ M pargyline, 10 mM lithium chloride, or 0.4 ml of assay medium and 50  $\mu$ l of 10xketanserin (ket) to a final concentration of 10  $\mu$ M. The cells were then incubated for 30 minutes at 37° C. Then the cells were washed with 0.5 ml PBS and 200  $\mu$ l of fresh/icecold stop solution (1M KOH; 18 mM Na-borate; 3.8 mM EDTA) was added/well. The solution was kept on ice for 5–10 minutes or until the cells were lysed and then neutralized by 200  $\mu$ l of fresh/ice cold neutralization sol. (7.5% HCL). The lysate was then transferred into 1.5 ml microcentrifuge tubes and 1 ml of chloroform/methanol (1:2) was added/tube. The solution was vortexed for 15 seconds and the upper phase was applied to a Biorad AG1-X8 anion exchange resin (100–200 mesh). The resin was washed with water and 0.9 ml of the upper phase was loaded onto the column. The column was washed with 10 mls of 5 mM myo-inositol and 10 ml of 5 mM Na-borate/60 mM Na-formate. The inositol trisphosphates were eluted into scintillation vials containing 10 ml of scintillation cocktail with 2 ml of 0.1 M formic acid/ 1 M ammonium formate. The columns were regenerated by washing with 10 ml of 0.1 M formic acid/3M ammonium formate and rinsed twice with dd H<sub>2</sub>O and stored at room temperature in water. Results are discussed below.

FIG. 12 is an illustration of IP<sub>3</sub> production from the human 5-HT<sub>2A</sub> receptor which was mutated using the same point mutation as set forth in Casey, which rendered the rat receptor constitutively active. The results represented in FIG. 12, support the position that when the point mutation shown to activate the rat receptor is introduced into the human receptor, little activation of the receptor is obtained that would allow for appropriate screening of candidate compounds, with the response being only moderately above that of the endogenous human 5-HT<sub>2A</sub> receptor. Generally, a response of at least 2x above that of the endogenous response is preferred.

FIG. 13 provides an illustration comparing IP<sub>3</sub> production from endogenous 5-HT<sub>2A</sub> receptor and the AP4 mutation. The results illustrated in FIG. 13 support the position that when the novel mutation disclosed herein is utilized, a robust response of constitutive IP<sub>3</sub> accumulation is obtained (e.g., over 233 that of the endogenous receptor).

FIG. 14 provides an illustration of IP<sub>3</sub> production from AP3. The results illustrated in FIG. 14 support the position that when the novel mutation disclosed herein is utilized, a robust response of constitutive IP<sub>3</sub> accumulation is obtained.

FIG. 15 provides bar-graph comparisons of IP<sub>3</sub> accumulation between endogenous human 5-HT<sub>2C</sub> receptor and AP-1. Note that the endogenous receptor has a high degree of natural constitutive activity relative to the control CMV transfected cells (i.e., the endogenous receptor appears to be constitutively activated).

### Example 6

#### Screening of Compounds Known to Have 5HT<sub>2C</sub> Antagonist Activity Against Non-Endogenous, Constitutively Activated Human Serotonin Receptor: AP-1

A final concentration of 12.5  $\mu$ g membranes prepared from COS7 cells (see Example 2) transiently expressing constitutively active mutant human 5HT<sub>2C</sub> receptor AP-1 were incubated with binding buffer (20 mM HEPES, pH 7.4, 100 mM NaCl, 20 mM MgCl<sub>2</sub>.6H<sub>2</sub>O, 0.2% saponin, and 0.2 mM ascorbate), GDP(1  $\mu$ M) and compound in a 96-well plate format for a period of 60 minutes at ambient room temperature. Plates were then centrifuged at 4,000 rpm for 15 minutes followed by aspiration of the reaction mixture and counting for 1 minute in a Wallac™ MicroBeta plate scintillation counter. A series of compounds known to possess reported 5HT<sub>2C</sub> antagonist activity were determined to be active in the [<sup>35</sup>S]GTP $\gamma$ S binding assay using AP-1. IC<sub>50</sub> determinations were made for these commercially available compounds (RBI, Natick, Mass.). Results are summarized in Table 3. For each determination, eight concentrations of test compounds were tested in triplicate. The negative control in these experiments consisted of AP-1 receptor without test compound addition, and the positive control consisted of 12.5  $\mu$ g/well of COS7 cell membranes expressing the CMV promoter without expressed AP-1 receptor.

TABLE 3

Test Compound	Known Pharmacology	IC <sub>50</sub> (nM) in GTP- $\gamma$ -[ <sup>35</sup> S] Assay
Metergoline	5HT <sub>2/1C</sub> antagonist	32.0
Mesulergine	5HT <sub>2/1C</sub> antagonist	21.2
Methysergide	5HT <sub>2/1C</sub> antagonist	6.1

TABLE 3-continued

Test Compound	Known Pharmacology	IC <sub>50</sub> (nM) in GTP-γ-[ <sup>35</sup> S] Assay
Methiothepin	5HT <sub>1</sub> antagonist	20.4
Normethylclozapin	5HT <sub>2</sub> /1C antagonist	21.4
Fluoxetine	5HT reuptake inhibitor	114.0
Ritanserin	5HT <sub>2</sub> /1C antagonist	19.4

The IC<sub>50</sub> results confirm that the seven tested compounds showed antagonist activity at the AP-1 receptor.

#### Example 7

##### Screening of Candidate Compounds Against Non-Endogenous, Constitutively Activated Human Serotonin Receptors:AP-1

Approximately 5,500 candidate compounds (Tripos, Inc., St. Louis, Mo.) were screened using the assay protocol of Example 3 (with AP-1 mutant receptor) for identification as inverse agonists against the receptor; for this assay, an arbitrary cut-off of at least 50% inhibition was established for identification of inverse agonists. Approximately 120 of these compounds evidenced at least 50% inhibition of [<sup>35</sup>S]GTPγS binding at 10 μM candidate compound (data not shown).

#### Example 8

##### Screening of Selected Compounds to Confirm Receptor Binding: AP-1

The candidate compounds identified from Example 7 were then screened using the assay protocol of Example 4 (mesulergine), using the AP-1 mutant receptor. IC<sub>50</sub> (nM) values were determined; five of the nearly 120 compounds of Example 7 were determined to have potent binding affinity for the receptor. Results are summarized in Table 4.

TABLE 4

Candidate Compound	IC <sub>50</sub> (nM) in Mesulergine Assay
102461	205.0
102788	46.5
100341	209.0
100431	147.0
103487	1,810.0

#### Example 9a

##### General Screening Paradigm: Selection of Pre-Clinical Candidate Leads

The "primary" screen designed to directly identify human 5HT<sub>2A</sub>/5HT<sub>2C</sub> receptor inverse agonists consisted of a membrane-based GTPγS binding assay utilizing membranes prepared from COS7 cells transiently transfected with AP-1 human receptor. Candidate compounds (10 μM final assay concentration) directly identified as inhibiting receptor-mediated increases in GTPγS binding by greater than 50–75% (arbitrary cut-off value) were considered active "hits". Primary assay hits were then re-tested in the same assay to reconfirm their inverse agonist activity. If primary assay hits were reconfirmed active (50% or greater inhibition), and therefore directly identified as, e.g., an inverse agonist, one of two approaches were available: (a)

so-called "directed libraries" could be created, i.e., additional candidate compounds were synthesized based upon the structures of the reconfirmed hits (geared towards, e.g., improvement in the characteristics of the compounds) whereby the directed library compounds were then evaluated for the ability to compete for radioligand binding to both mutant 5HT<sub>2C</sub> (AP-1) and endogenous 5HT<sub>2A</sub> receptors, or (b) the reconfirmed hits were then evaluated for the ability to compete for radioligand binding to both mutant 5HT<sub>2C</sub> (AP-1) and endogenous 5HT<sub>2A</sub> receptors. Thus, when approach (a) was used, because these directed library candidate compounds were based upon the structures of compounds that were directly identified from the membrane-based GTPγS binding assay, the directed library compounds were not re-tested in the membrane-based GTPγS binding assay but rather were then confirmed via the radioligand binding analysis. The radioligand binding analysis tests were initially performed at 10 μM test compound in triplicate and if the compound inhibited radiolabeled binding by 50% or more, the analysis was followed by eight concentration competition curves to determine Ki values. The last step in secondary assay evaluation was to determine if test compounds were capable of inhibiting AP-3 receptor-mediated accumulation of inositol phosphates (e.g., IP<sub>3</sub>). This final assay confirms that the directly identified compounds retained inverse agonist properties.

#### Example 9b

##### Constitutively Activated Human 5HT<sub>2C</sub> Receptor (AP-1) Mediated Facilitation of GTPγS Binding to COS7 Membranes

This protocol is substantially the same as set forth above in Example 6.

Primary screening assays measuring GTPγS binding to membranes prepared from COS7 cells transiently transfected with human mutated 5HT<sub>2C</sub> receptor (AP-1) were used to directly identify inverse agonists in screening libraries (Tripos, Inc.). Candidate compound screens were performed in a total assay volume of 200 μl using scintillant-coated Wallac Scintistrip™ plates. The primary assay was comprised of the following chemicals (at indicated final assay concentrations): 20 mM HEPES, pH 7.4, 100 mM NaCl, 20 mM MgCl<sub>2</sub>, 0.2% saponin, 0.2 mM ascorbic acid, 1 μM GDP, 0.3 nM GTPγ<sup>35</sup>S, and 12.5 μg of the above defined membranes. Incubations were performed for 60 minutes at ambient room temperature. The binding assay incubation was terminated by centrifugation of assay plates at 4,000 rpm for 15 minutes, followed by rapid aspiration of the reaction mixture and counting in a Wallac MicroBeta™ scintillation counter.

Primary screening of candidate compounds initially involved testing of 72 test compounds per assay plate (96-well plates were utilized), at a final assay concentration of 10 μM candidate compound, in single replicates. A total of sixteen wells of each plate were dedicated for an eight concentration clozapine (a confirmed 5HT<sub>2C</sub>/2A inverse agonist) dose response curve (duplicate determinations at each concentration). Finally, a total of five assay wells of each plate were dedicated to define the negative control (AP-1 receptor expressing membranes without addition of candidate compounds) and three wells from each plate to define the positive control (membranes without AP-1 receptor).

Reconfirmation experiments involve re-testing candidate compounds in the same assay described above, except that

candidate compounds were evaluated in triplicate, thus allowing evaluation of 24 compounds per 96-well assay plate. Similar to the primary assay plates, an eight concentration clozapine dose response curve (duplicate determinations at each concentration) and the same negative and positive control wells were also included within each 96-well plate.

#### Example 9c(1)

##### Competition Studies Mutated Human 5HT2C Receptor (AP-1)

Radioligand binding competition experiments were performed in a total assay volume of 200  $\mu$ l using standard 96-well microtiter plates. The final assay ingredients consisted of assay buffer (20 mM HEPES and 10 mM MgCl<sub>2</sub>), 1 nM [<sup>3</sup>H]mesulergine, and 50  $\mu$ g of membranes (COS7 with AP-1 as defined above). Nonspecific [<sup>3</sup>H]mesulergine binding was defined in the presence of 100  $\mu$ M mianserin. Incubations were performed for 1 hour at 37° C. Receptor bound radioligand was resolved from free radioligand by rapid filtration of the assay mixture over a Wallac Filtermat™ Type B filter, followed by washing with ice-cold assay buffer using a Skatron™ cell harvester. Radioactivity was counted using a Wallac 1205 BetaPlate™ counter. Each assay plate contained five negative control wells (membranes expressing receptor and no candidate compound addition) and three positive control wells (each containing 100  $\mu$ M mianserin). For one concentration tests, candidate compounds were diluted into assay buffer and screened at a final concentration of 10  $\mu$ M, in triplicate. For IC<sub>50</sub> determinations, candidate compounds were diluted in assay buffer and eight different concentrations were evaluated, in triplicate. A total of 16 wells were designated for an eight concentration mianserin dose response curve evaluation for both assays.

#### Example 9c(2)

##### Competition Studies Wild Type Human 5HT2A Receptor

Radioligand binding competition experiments were performed in a total assay volume of 200  $\mu$ l using standard 96-well microtiter plates. The final assay ingredients comprised assay buffer (20 mM HEPES and 10 mM MgCl<sub>2</sub>), 1 nM [<sup>3</sup>H]LSD, and 50  $\mu$ g of the above-defined membranes (COS7 with AP-1). Nonspecific [<sup>3</sup>H]LSD binding was defined in the presence of 100  $\mu$ M serotonin. Incubations were performed for 1 hour at 37° C. Receptor bound radioligand was resolved from free radioligand by rapid filtration of the assay mixture over a Wallac Filtermat™ Type B filter, followed by washing with ice-cold assay buffer using a Skatron™ cell harvester. Radioactivity was counted using a Wallac 1205 Betaplate™ counter. Each assay plate contained five negative control wells (membranes expressing receptor and no candidate compound addition) and three positive control wells (containing 100  $\mu$ M mianserin). For one concentration tests, candidate compounds were diluted into assay buffer and screened at a final concentration of 10  $\mu$ M in triplicate. For IC<sub>50</sub> determinations, candidate compounds were diluted in assay buffer and eight different concentrations were evaluated in triplicate. A total of 16 wells were designated for an eight concentration serotonin dose response curve evaluation for both assays.

#### Example 9d

##### Receptor-Mediated Inositol Phosphate Accumulation

Candidate compound identified in the assays of Examples 9a–9c were then evaluated for inositol phosphate

accumulation, following the protocol of Example 5 (COS7 cells expressing human mutated 5HT2A receptor, AP-3), modified as follows: tube A was prepared by mixing 16  $\mu$ g DNA (e.g., pCMV vector; pCMV vector AP-1 cDNA, etc.) in 1.0 ml serum free DMEM (Irvine Scientific, Irvine, Calif.); tube B was prepared by mixing 60  $\mu$ l lipofectamine (Gibco BRL) in 1.0 ml serum free DMEM. Tubes A and B were then admixed by inversions (several times), followed by incubation at room temperature for 30 min. The admixture is referred to as the “transfection mixture”. Plated 293 cells were washed with 10 ml Serum Free DMEM, followed by addition of 11 ml Serum Free DMEM. 2.0 ml of the transfection mixture was then added to the cells, followed by incubation for 5 hrs at 37° C./5% CO<sub>2</sub>. On day 3, cells were trypsinized and counted, followed by plating of 1×10<sup>6</sup> cells/well (12-well plates). Cells were permitted to adhere to the wells for 8 hrs., followed by one wash with 1×PBS. Thereafter, 0.5  $\mu$ Ci <sup>3</sup>H-inositol in 1 ml inositol-free DMEM was added per well.

On day 4, the cells were washed with 1.5 ml PBS and then 0.9 ml of assay medium was added containing inositol-free/serum free media, 10  $\mu$ M pargyline, 10 mM lithium chloride, for 5 min in 37° C./5% CO<sub>2</sub> followed by 100  $\mu$ l addition of candidate compound diluted in the same material. The cells were then incubated for 120 minutes at 37° C. Then the cells were washed with 1.5 ml PBS and 200  $\mu$ l of fresh/icecold stop solution (1M KOH; 18 mM Na-borate; 3.8 mM EDTA) was added/well. The solution was kept on ice for 5–10 minutes or until the cells were lysed and then neutralized by 200  $\mu$ l of fresh/ice cold neutralization sol. (7.5% HCL). The lysate was then transferred into 1.5 ml micro-centrifuge tubes and 1 ml of chloroform/methanol (1:2) was added/tube. The solution was vortexed for 15 seconds and the upper phase was applied to a Biorad AG1-X8 anion exchange resin (100–200 mesh). The resin was washed with water and 0.9 ml of the upper phase was loaded onto the column. The column was washed with 10 mls of 5 mM myo-inositol and 10 ml of 5 mM Na-borate/60 mM Na-formate. The inositol trisphosphates were eluted into scintillation vials containing 10 ml of scintillation cocktail with 2 ml of 0.1 M formic acid/1 M ammonium formate. The columns were regenerated by washing with 10 ml of 0.1 M formic acid/3M ammonium formate and rinsed twice with dd H<sub>2</sub>O and stored at room temperature in water.

Following this round of assaying, candidate compounds having an IC<sub>50</sub> value of less than 10  $\mu$ M were considered as potential leads for the development of pharmaceutical compositions.

#### Screening Candidate Compounds

Following the protocols set forth above, one compound, 103487 (Example 8, supra) evidenced the following results:


Figure Number	GTP $\gamma$ S AP-1 Percent Inhibition Relative To Positive Control (Primary)	GTP $\gamma$ S AP-1 Percent Inhibition Relative To Positive Control (Reconfirm)	Competitive Binding AP-1 ([ <sup>3</sup> H]mesulergine) IC <sub>50</sub> Value (nM)	Competitive Binding WT 5HT2A ([ <sup>3</sup> H]LSD) IC <sub>50</sub> Value (nM)	Inositol Phosphate Accumulation AP-3 IC <sub>50</sub> Value (nM)
15A (103487)	-1%	31%	2100 850	46	52 90

Based upon these results, structure activity analysis of the 103487 compound suggested that a series of derivatives of 3-(4-bromo-1-methylpyrazole-3-yl)phenylamine would

## 21

exhibit similar 5-HT<sub>2A</sub> activity and selectivity. A series of derivatives of 3-(4-bromo-1-methylpyrazole-3-yl)phenylamine have now been synthesized. These "directed" library compounds (Tripos, Inc.) were then analyzed in accordance with the protocols of Examples 9c(1), 9c(2) and 9d.

This series of compounds exhibits highly selective 5-HT<sub>2A</sub> activity. Accordingly, in the first aspect of the invention, a series of compounds possessing 5-HT<sub>2A</sub> receptor activity that are useful as inverse agonists at such receptors is designated by the general formula (A):



Wherein:

W is lower alkyl (C<sub>1-6</sub>), or halogen;

V is lower alkyl (C<sub>1-6</sub>), or halogen;

X is either Oxygen or Sulfur;

Y is NR<sup>2</sup>R<sup>3</sup>, or (CH<sub>2</sub>)<sub>m</sub>R<sup>4</sup>, or O(CH<sub>2</sub>)<sub>n</sub>R<sup>4</sup>;

Z is lower alkyl (C<sub>1-6</sub>);

m=0-4

n=0-4

R<sup>1</sup> is H or lower alkyl (C<sub>1-4</sub>);

R<sup>2</sup> is H or lower alkyl (C<sub>1-4</sub>);

R<sup>3</sup> and R<sup>4</sup> are independently a C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>5</sup>R<sup>6</sup>, NR<sup>5</sup>R<sup>6</sup>, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, SO<sub>3</sub>R<sup>7</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, aryl, and aryloxy wherein each of the C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>5</sup>R<sup>6</sup>, NR<sup>5</sup>R<sup>6</sup>, NHCOCH<sub>3</sub>, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>3</sub>R<sup>7</sup>, SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, and aryl;

R<sup>5</sup> and R<sup>6</sup> are independently a H, or C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or aryl, or CH<sub>2</sub> aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>7</sup>R<sup>8</sup>, NR<sup>7</sup>R<sup>8</sup>, NHCOCH<sub>3</sub>, OCF<sub>3</sub>, SMe, COOR<sup>9</sup>, SO<sub>3</sub>R<sup>7</sup>, SO<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, and aryl wherein each of the C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>8</sup>R<sup>9</sup>, NR<sup>8</sup>R<sup>9</sup>, NHCOCH<sub>3</sub>, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, SO<sub>3</sub>R<sup>7</sup>, COMe, COEt,

## 22

CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, and aryl, or R<sup>5</sup> and R<sup>6</sup> may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, SO<sub>3</sub>R<sup>7</sup>, NHCOCH<sub>3</sub>, COEt, COMe, or halogen;

R<sup>7</sup> may be independently selected from H or C<sub>1-6</sub> alkyl;

R<sup>8</sup> and R<sup>9</sup> are independently a H, or C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or aryl, or CH<sub>2</sub>aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF<sub>3</sub>, OCF<sub>3</sub>, OEt, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, SMe, COMe, CN, COOR<sup>7</sup>, SO<sub>3</sub>R<sup>7</sup>, COEt, NHCOCH<sub>3</sub>, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic heterocyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle;

C<sub>1-6</sub> alkyl moieties can be straight chain or branched; optionally substituted C<sub>1-6</sub> alkyl moieties can be straight chain or branched;

C<sub>2-6</sub> alkenyl moieties can be straight chain or branched; and


optionally substituted C<sub>2-6</sub> alkenyl moieties can be straight chain or branched.

Examples of suitable C<sub>1-6</sub> alkyl groups include but are not limited to methyl, ethyl, n-propyl, i-propyl, n-butyl, and t-butyl.

Halogens are typically F, Cl, Br, and I.

Examples of 5 or 6 membered ring moieties include, but are not restricted to, phenyl, furanyl, thienyl, imidazolyl, pyridyl, pyrrolyl, oxazolyl, isoxazolyl, triazolyl, pyrazolyl, tetrazolyl, thiazolyl and isothiazolyl. Examples of polycycle moieties include, but are not restricted to, naphthyl, benzothiazolyl, benzofuranyl, benzimidazolyl, quinolyl, isoquinolyl, indolyl, quinoxalyl, quinazolyl and benzothienyl.

A more preferred series of compounds possessing 5-HT<sub>2A</sub> receptor activity that are useful as inverse agonists at such receptors is designated by the general formula (B):



Wherein:

W is Me, or Et, or halogen;

X is either Oxygen or Sulfur;

Y is NR<sup>2</sup>R<sup>3</sup>, or (CH<sub>2</sub>)<sub>m</sub>R<sup>4</sup>, or O(CH<sub>2</sub>)<sub>n</sub>R<sup>4</sup>;

Z is lower alkyl (C<sub>1-6</sub>);

m=0-4

n=0-4

R<sup>1</sup> is H or lower alkyl (C<sub>1-4</sub>);

R<sup>2</sup> is H or lower alkyl(C<sub>1-4</sub>);

R<sup>3</sup> and R<sup>4</sup> are independently a C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>5</sup>R<sup>6</sup>, NR<sup>5</sup>R<sup>6</sup>, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, SO<sub>3</sub>R<sup>7</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, aryl, and aryloxy wherein each of the C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>5</sup>R<sup>6</sup>, NR<sup>5</sup>R<sup>6</sup>, NHCOCH<sub>3</sub>, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>3</sub>R<sup>7</sup>, SO<sub>2</sub>NR<sup>5</sup>R<sup>6</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, and aryl;

R<sup>5</sup> and R<sup>6</sup> are independently a H, or C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or aryl, or CH<sub>2</sub> aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>7</sup>R<sup>8</sup>, NR<sup>7</sup>R<sup>8</sup>, NHCOCH<sub>3</sub>, OCF<sub>3</sub>, SMe, COOR<sup>9</sup>, SO<sub>3</sub>R<sup>7</sup>, SO<sub>2</sub>NR<sup>7</sup>R<sup>8</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, and aryl wherein each of the C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, CONR<sup>8</sup>R<sup>9</sup>, NR<sup>8</sup>R<sup>9</sup>, NHCOCH<sub>3</sub>, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, SO<sub>3</sub>R<sup>7</sup>, COMe, COEt, CO-lower alkyl, SCF<sub>3</sub>, CN, C<sub>2-6</sub> alkenyl, H, halogens, C<sub>1-4</sub> alkoxy, C<sub>3-6</sub> cycloalkyl, C<sub>1-6</sub> alkyl, and aryl,

or R<sup>5</sup> and R<sup>6</sup> may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from CF<sub>3</sub>, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, OEt, OCF<sub>3</sub>, SMe, COOR<sup>7</sup>, SO<sub>2</sub>NR<sup>8</sup>R<sup>9</sup>, SO<sub>3</sub>R<sup>7</sup>, NHCOCH<sub>3</sub>, COEt, COMe, or halogen;

R<sup>7</sup> may be independently selected from H or C<sub>1-6</sub> alkyl;

R<sup>8</sup> and R<sup>9</sup> are independently a H, or C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or aryl, or CH<sub>2</sub>aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen, CF<sub>3</sub>, OCF<sub>3</sub>, OEt, CCl<sub>3</sub>, Me, NO<sub>2</sub>, OH, OMe, SMe, COMe, CN, COOR<sup>7</sup>, SO<sub>3</sub>R<sup>7</sup>, COEt, NHCOCH<sub>3</sub>, or aryl;

an aryl moiety can be a 5 or 6 membered aromatic heterocyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle;

C<sub>1-6</sub> alkyl moieties can be straight chain or branched; optionally substituted C<sub>1-6</sub> alkyl moieties can be straight chain or branched;

C<sub>2-6</sub> alkenyl moieties can be straight chain or branched; and


optionally substituted C<sub>2-6</sub> alkenyl moieties can be straight chain or branched.

Examples of suitable C<sub>1-6</sub> alkyl groups include but are not limited to methyl, ethyl, n-propyl, i-propyl, n-butyl, and t-butyl.

Halogens are typically F, Cl, Br, and I.

Examples of 5 or 6 membered ring moieties include, but are not restricted to, phenyl, furanyl, thienyl, imidazolyl, pyridyl, pyrrolyl, oxazolyl, isoxazolyl, triazolyl, pyrazolyl, tetrazolyl, thiazolyl and isothiazolyl. Examples of polycyclic moieties include, but are not restricted to, naphthyl, benzothiazolyl, benzofuranyl, benzimidazolyl, quinolyl, isoquinolyl, indolyl, quinoxalyl, quinazolyl and benzothienyl.

A first series of compounds having 5-HT<sub>2A</sub> receptor activity is represented by a class (I) of compounds of formula (B) wherein Y=NR<sup>2</sup>R<sup>3</sup>:



(I)

Wherein:

Preferably R<sup>1</sup> and R<sup>2</sup> are H.

Preferably W is Br.

Preferably X is O.


Preferably Z is Me.

Preferably R<sup>3</sup> is 4-trifluoromethoxyphenyl or 4-trifluoromethoxybenzyl.

Preferred compounds are:

103487


N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl}amino]carboxamide



116115

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl)methyl}amino]carboxamide

25



These two compounds demonstrated the following activities using the assay protocols defined in the Examples above:

Compound Number	Competitive Binding AP-1 ([ <sup>3</sup> H]mesulergine) IC <sub>50</sub> Value (μM)	Competitive Binding WT 5HT <sub>2A</sub> ([ <sup>3</sup> H]LSD) IC <sub>50</sub> Value (μM)	Inositol Phosphate Accumulation AP-3 IC <sub>50</sub> Value (μM)
103487	2.1	.046	.052
116115	1.2	.45	.0171

26


Additional compounds of formula (B) wherein Y=NR<sup>2</sup>R<sup>3</sup> are set forth below. Inositol phosphate accumulation assays evidence the activity of test compounds. Both single concentration percentages of control values and IC<sub>50</sub> determinations indicate activity. In the tables below the column legends have the following meanings:

IP<sub>3</sub>% Control: The values in this column reflect an IP Accumulation Assay where the test compounds were evaluated at one concentration of 10 μM. For these assays, the compound was diluted into inositol-free Dulbecco's Eagle Media containing 10 μM pargyline and 10 mM LiCl and tested at a final assay concentration of 10 μM, in triplicate. The percent control value was calculated based on the control in which no test compound was added.

IP<sub>3</sub> AP-3 IC<sub>50</sub> nM: The values in this column reflect an IP accumulation assay in which the test compound was evaluated at several different concentrations whereby an IC<sub>50</sub> could be determined. This column corresponds to the column appearing in the tables above which is labeled: Inositol Phosphate Accumulation, AP-3, IC<sub>50</sub> Value (EM).

WT 5HT<sub>2A</sub> LSD IC<sub>50</sub> nM: The values in this column reflect a competitive binding assay using LSD. This column corresponds to the column appearing in the tables above which is labeled: Competitive Binding, WT 5HT<sub>2A</sub>, ([<sup>3</sup>H]LSD), IC<sub>50</sub> Value (μM).


Compounds listed in each of the following tables reference the structures immediately preceding the table. A "dash" in the table indicates that no value was determined.




Compound No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	X	U	IP <sub>3</sub> % of Control	IP <sub>3</sub> AP-3 IC <sub>50</sub> nM	WT 5HT <sub>2A</sub> LSD IC <sub>50</sub> nM
116079	SCH <sub>3</sub>	H	H	H	O	NH	16	17	4
116081	Cl	H	H	H	O	NH	10	3.2	11
116082	F	H	H	H	O	NH	11	—	7
116087	H	H	CF <sub>3</sub> O	H	O	NH	11	—	200
116089	H	H	NO <sub>2</sub>	H	O	NH	27	—	238
116091	MeO	H	H	H	O	NH	12	—	19
116092	H	H	Me	H	O	NH	32	—	131
116097	CF <sub>3</sub>	H	H	H	O	NH	11	—	65
116105	H	Cl	H	H	O	NH	11	—	39
116108	H	H	Cl	H	O	NH	6	—	249
116110	isopropyl	H	H	H	O	NH	7	—	338

-continued

116111	{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-methoxyphenyl)carboxamide	H	MeO	H	H	O	NH	7	—	106
116112	{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-methylphenyl)carboxamide	H	Me	H	H	O	NH	14	—	57
116113	[{3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino]-N-methyl-N-[4-(trifluoromethoxy)phenyl]carboxamide	CF <sub>3</sub> O	H	H	H	O	NCH <sub>3</sub>	—	193	2
116119	N-[4-(tert-butyl)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide	t-butyl	H	H	H	O	NH	17	—	476
116122	N-[4-(dimethylamino)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide	NMe <sub>2</sub>	H	H	H	O	NH	9	—	309
116138	N-(3,5-dichloro-4-methylphenyl){[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide	Me	Cl	H	Cl	O	NH	23	—	122
116139	{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide	CF <sub>3</sub> S	H	H	H	O	NH	12	—	56
116144	{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-fluorophenyl)carboxamide	H	H	F	H	O	NH	12	—	37
116145	2-({[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carbonylamino)benzamide	H	H	CONH <sub>2</sub>	H	O	NH	31	—	7473
116147	{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-cyanophenyl)carboxamide	CN	H	H	H	O	NH	12	—	2
116148	{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-cyanophenyl)carboxamide	H	H	CN	H	O	NH	30	—	348




Compound No.	N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]cyclohexylamino]carboxamide	IP <sub>3</sub> AP-3 IC <sub>50</sub> nM	WT 5HT <sub>2A</sub> LSD IC <sub>50</sub> nM
116141		114	81



Compound No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	IP <sub>3</sub> AP-3 IC <sub>50</sub> nM	WT 5HT <sub>2A</sub> LSD IC <sub>50</sub> nM
116143	H	H	H	H	H	120	47
116182	F	H	H	H	H	89	132
116183	OMe	OMe	H	H	H	—	1010
116184	OMe	OMe	H	OMe	H	—	2960
116185	H	H	Me	H	H	—	769
116189	OMe	H	H	H	H	—	102




-continued



Compound No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	R <sup>5</sup>	IP <sub>3</sub> AP-3 IC <sub>50</sub> nM	WT 5HT <sub>2A</sub> LSD IC <sub>50</sub> nM
N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-(4-methoxyphenyl)ethyl]amino]carboxamide 116194	OMe	H	H	H	H	32	61

A second series of compounds having 5-HT<sub>2A</sub> receptor activity is represented by a class (II) of compounds of formula (B) wherein Y=O(CH<sub>2</sub>)<sub>n</sub>R<sup>4</sup>:



Wherein:

Preferably R<sup>1</sup> is H.

Preferably W is Br.

Preferably X is O.


Preferably L is Me.

Preferably when n=0, R<sup>4</sup> is 4-methoxyphenyl or tertiary butyl.


Preferred compounds are:

116100

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide



116192  
(tert-butoxy)-N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]carboxamide



These two compounds demonstrated the following activity:

Compound No.	Competitive Binding AP-1 ([ <sup>3</sup> H]mesulergine) IC <sub>50</sub> Value (μM)	Competitive Binding WT 5HT <sub>2A</sub> ([ <sup>3</sup> H]LSD) IC <sub>50</sub> Value (μM)	Inositol Phosphate Accumulation AP-3 IC <sub>50</sub> Value (μM)
116100	1.8	<0.001	0.0003
116192	—	0.014	0.057

In addition to the assays discussed above, the specific activity of 116100 at the 5HT<sub>2A</sub> receptor was further confirmed by the following.

#### In Vitro Binding of 5HT<sub>2A</sub> Receptor

##### Animals

Animals (Sprague-Dawley rats) were sacrificed and brains were rapidly dissected and frozen in isopentane maintained at -42° C. Horizontal sections were prepared on a cryostat and maintained at -20° C.

##### LSD Displacement Protocol

Lysergic acid diethylamide (LSD) is a potent 5HT<sub>2A</sub> receptor and dopamine D<sub>2</sub> receptor ligand. An indication of the selectivity of compounds for either or both of these receptors involves displacement of radiolabeled-bound LSD from pre-treated brain sections. For these studies, radiola-

31

beled I<sup>125</sup>-LSD (NEN Life Sciences, Boston, Mass., Catalogue number NEX-199) was utilized; spiperone (RBI, Natick, Mass. Catalogue number s-128), a 5HT<sub>2A</sub> receptor and dopamine D2 receptor antagonist, was also utilized. Buffer consisted of 50 nanomolar TRIS-HCl, pH 7.4.


Brain sections were incubated in (a) Buffer plus 1 nanomolar I<sup>125</sup>-LSD; (b) Buffer plus 1 nanomolar I<sup>125</sup>-LSD and 1 micromolar spiperone; or Buffer plus 1 nanomolar I<sup>125</sup>-LSD and 1 micromolar 116100 for 30 minutes at room temperature. Sections were then washed 2×10 minutes at 4° C. in Buffer, followed by 20 seconds in distilled H<sub>2</sub>O. Slides were then air-dried.

After drying, sections were apposed to x-ray film (Kodak Hyperfilm) and exposed for 4 days.

#### Analysis

FIGS. 16A–C provide representative autoradiographic sections from this study. FIG. 16A evidences darker bands (derived from I<sup>125</sup>-LSD binding) primarily in both the fourth layer of the cerebral cortex (primarily 5HT<sub>2A</sub> receptors), and the caudate nucleus (primarily dopamine D2 receptors and some 5HT<sub>2A</sub> receptors). As can be seen from FIG. 16B, spiperone, which is a 5HT<sub>2A</sub> and dopamine D2 antagonist, displaces the I<sup>125</sup>-LSD from these receptors on both the cortex and the caudate. As can be further seen from FIG. 16C, 116100 appears to selectively displace the I<sup>125</sup>-LSD from the cortex (5HT<sub>2A</sub>) and not the caudate (dopamine D2).

A third series of compounds having 5-HT<sub>2A</sub> receptor activity is represented by a class (III) of compounds of formula (B) wherein Y=(CH<sub>2</sub>)<sub>m</sub>R<sup>4</sup>:



Wherein:

Preferably W is Br.

Preferably X is O.

Preferably Z is Me.

Preferably R<sup>1</sup> is H.

Preferably when m=0; R<sup>4</sup> is preferably 4-trifluoromethoxyphenyl, or thiophene, or 4-chlorophenyl.


Preferred compounds are:

116101

m=0, R<sup>1</sup>=H, R<sup>4</sup>=4-trifluoromethoxyphenyl

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide


32



116102

m=0, R<sup>1</sup>=H, R<sup>4</sup>=thiophene


N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide



116120

m=0, R<sup>1</sup>=H, R<sup>4</sup>=4-chlorophenyl

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-chlorophenyl]carboxamide



These three compounds demonstrated the following activities:

Compound Number	Competitive Binding AP-1 ([ <sup>3</sup> H]mesulergine) IC <sub>50</sub> Value (μM)	Competitive Binding WT 5HT <sub>2A</sub> ([ <sup>3</sup> H]LSD) IC <sub>50</sub> Value (μM)	Inositol Phosphate Accumulation AP-3 IC <sub>50</sub> Value (μM)
116101	6.1	.46	0.0213
116102	2.8	.17	0.080
116120	1.2	.21	0.0315

## In Vivo Analysis of Compound 116102

In addition to the in vitro assays shown in the above table, the in vivo response of animals to the 116102 compound is demonstrated by the following.

A 5HT<sub>2A</sub> receptor antagonist or inverse agonist is expected to decrease amphetamine-stimulated locomotion without affecting baseline locomotion. See, for example, Soresnon, et al, 266(2) J. Pharmacol. Exp. Ther. 684 (1993). Based upon the foregoing information, Compound 116102 is a potent inverse agonist at the human 5HT<sub>2A</sub> receptor. For the following study, the following parameters and protocol were utilized:

## Animals, Vehicle

Adult male Sprague-Dawley rats were utilized for these studies. Animals were housed in groups of 2–3 in hanging plastic cages with food and water available at all times. Animals were weighed and handled for at least one day prior to surgery and throughout the studies. For these studies, Vehicle consisted of 90% ethanol (100%) and 10% water.

## Amphetamine-stimulated Locomotor Activity: Assessment and Apparatus


A San Diego Instruments Flex Field apparatus was used to quantify baseline and amphetamine-stimulated locomotor activity. This apparatus consists of four 16"×16" clear plastic open fields. Photocell arrays (16 in each dimension) interfaced with a personal computer to automatically quantify activity. Several measures of activity can be assessed with the apparatus, including total photocell beam breaks. Animals (vehicle control and Compound treated) were injected s.c. 30 minutes prior to initiation of analysis. Following this 30 minute period, animals were placed individually into an open field and baseline activity was assessed for 30 minutes (habituation phase). Following baseline, animals were removed, injected with d-amphetamine sulfate (1.0 mg/kg) and immediately returned to the open field for 150 minutes, in order to follow the time course (10 minute intervals) of amphetamine-stimulated locomotor activity.

Vehicle Control	Compound 116102	Dose (mg/kg)
6 animals	6 animals	0.1
	6 animals	1.0
	6 animals	5.0
	6 animals	10.0

## Analysis

Results, based upon the number of recorded photobeam breaks (mean±sem), are presented in FIGS. 17A–C. As supported by FIGS. 17A,B and C, a general “inverted U” shaped pattern was observed (see, generally, Sahgal, A. “Practical behavioural neuroscience: problems, pitfalls and suggestions” pp 1–8, 5 in *Behavioral Neuroscience: A Practical Approach*, Volume 1 A. Sahgal (Ed.) 1993, IRL Press, New York). As FIG. 17 also indicates, with exception of the highest dose (10 mg/kg), in vivo, the tested doses of Compound 116102 evidenced a decrease in the amphetamine-stimulated locomotion, consistent with a 5HT<sub>2A</sub> receptor antagonist or inverse agonist.


Additional compounds of formula (B) wherein Y=(CH<sub>2</sub>)<sub>m</sub>R<sup>4</sup> are set forth below.



Compound No.	R <sup>1</sup>	R <sup>2</sup>	R <sup>3</sup>	R <sup>4</sup>	IP <sub>3</sub> IC <sub>50</sub> nM	LSD IC <sub>50</sub> nM
116137	OCF <sub>3</sub>	H	H	H	—	106
116174	H	F	H	H	153	318
116175	H	OMe	H	H	108	625
116176	H	H	F	H	129	662
116177	NO <sub>2</sub>	H	H	H	61	108
116178	H	H	OMe	H	165	2300

compound names not provided

Based upon the discovery of the specific inverse agonist activity of the above identified compounds at the 5HT<sub>2A</sub> receptor, a novel class of compounds has been identified which exhibits said activity. Accordingly, in the second aspect of the invention, there is provided a novel compound of formula (C):



Wherein:

W is Me, or Et, or halogen;

X is either Oxygen or Sulfur;

Y is NR<sup>2</sup>R<sup>3</sup>, or (CH<sub>2</sub>)<sub>m</sub>R<sup>4</sup>, or O(CH<sub>2</sub>)<sub>n</sub>R<sup>4</sup>;

Z is lower alkyl (C<sub>1-6</sub>);

m=0–4;

n=0–4;

R<sup>1</sup> is H or lower alkyl (C<sub>1-4</sub>);

R<sup>2</sup> is H or lower alkyl(C<sub>1-4</sub>);

R<sup>3</sup> is a C<sub>1-6</sub> alkyl, or C<sub>2-6</sub> alkenyl, or cycloalkyl, or (CH<sub>2</sub>)<sub>k</sub>aryl group (k=1–4), preferably k=1, and each

said group may be optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{CONR}^5\text{R}^6$ ,  $\text{NR}^5\text{R}^6$ ,  $\text{OCF}_3$ , SMe,  $\text{COOR}^7$ ,  $\text{SO}_2\text{NR}^5\text{R}^6$ ,  $\text{SO}_3\text{R}^7$ , COMe, COEt, CO-lower alkyl,  $\text{SCF}_3\text{CN}$ ,  $\text{C}_{2-6}$  alkenyl, H, halogens,  $\text{C}_{1-4}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, aryl, and aryloxy wherein each of the  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{CONR}^5\text{R}^6$ ,  $\text{NR}^5\text{R}^6$ ,  $\text{NHCOCH}_3$ ,  $\text{OCF}_3$ , SMe,  $\text{COOR}^7$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{SO}_2\text{NR}^5\text{R}^6$ , COMe, COEt, CO-lower alkyl,  $\text{SCF}_3$ , CN,  $\text{C}_{2-6}$  alkenyl, H, halogens,  $\text{C}_{1-4}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, and aryl;

$\text{R}^4$  is a  $\text{C}_{1-6}$  alkyl, or  $\text{C}_{2-6}$  alkenyl, or cycloalkyl, or aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{CONR}^5\text{R}^6$ ,  $\text{NR}^5\text{R}^6$ ,  $\text{NHCOCH}_3$ ,  $\text{OCF}_3$ , SMe,  $\text{COOR}^7$ ,  $\text{SO}_2\text{NR}^5\text{R}^6$ ,  $\text{SO}_3\text{R}^7$ , COMe, COEt, CO-lower alkyl,  $\text{SCF}_3\text{CN}$ ,  $\text{C}_{2-6}$  alkenyl, H, halogens,  $\text{C}_{1-4}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, aryl, and aryloxy wherein each of the  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, aryl, or aryloxy groups may be further optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{CONR}^5\text{R}^6$ ,  $\text{NR}^5\text{R}^6$ ,  $\text{NHCOCH}_3$ ,  $\text{OCF}_3$ , SMe,  $\text{COOR}^7$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{SO}_2\text{NR}^5\text{R}^6$ , COMe, COEt, CO-lower alkyl,  $\text{SCF}_3$ , CN,  $\text{C}_{2-6}$  alkenyl, H, halogens,  $\text{C}_{1-4}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, and aryl;

$\text{R}^5$  and  $\text{R}^6$  are independently a H, or  $\text{C}_{1-6}$  alkyl, or  $\text{C}_{2-6}$  alkenyl, or cycloalkyl, or aryl, or  $\text{CH}_2$  aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{CONR}^7\text{R}^8$ ,  $\text{NR}^7\text{R}^8$ ,  $\text{NHCOCH}_3$ ,  $\text{OCF}_3$ , SMe,  $\text{COOR}^9$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{SO}_2\text{NR}^7\text{R}^8$ , COMe, COEt, CO-lower alkyl,  $\text{SCF}_3$ , CN,  $\text{C}_{2-6}$  alkenyl, H, halogens,  $\text{C}_{1-4}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, and aryl wherein each of the  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, or aryl groups may be further optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{CONR}^8\text{R}^9$ ,  $\text{NR}^8\text{R}^9$ ,  $\text{NHCOCH}_3$ ,  $\text{OCF}_3$ , SMe,  $\text{COOR}^7$ ,  $\text{SO}_2\text{NR}^8\text{R}^9$ ,  $\text{SO}_3\text{R}^7$ , COMe, COEt, CO-lower alkyl,  $\text{SCF}_3$ , CN,  $\text{C}_{2-6}$  alkenyl, H, halogens,  $\text{C}_{1-4}$  alkoxy,  $\text{C}_{3-6}$  cycloalkyl,  $\text{C}_{1-6}$  alkyl, and aryl,

or  $\text{R}^5$  and  $\text{R}^6$  may form part of a 5, 6 or 7 membered cyclic structure which may be either saturated or unsaturated and that may contain up to four heteroatoms selected from O, N or S and said cyclic structure may be optionally substituted by up to four substituents in any position independently selected from  $\text{CF}_3$ ,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, OEt,  $\text{OCF}_3$ , SMe,  $\text{COOR}^7$ ,  $\text{SO}_2\text{NR}^8\text{R}^9$ ,  $\text{SO}_3\text{R}^7$ ,  $\text{NHCOCH}_3$ , COEt, COMe, or halogen;

$\text{R}^7$  may be independently selected from H or  $\text{C}_{1-6}$  alkyl;

$\text{R}^8$  and  $\text{R}^9$  are independently a H, or  $\text{C}_{1-6}$  alkyl, or  $\text{C}_{2-6}$  alkenyl, or cycloalkyl, or aryl, or  $\text{CH}_2$  aryl group and each said group may be optionally substituted by up to four substituents in any position independently selected from halogen,  $\text{CF}_3$ ,  $\text{OCF}_3$ , OEt,  $\text{CCl}_3$ , Me,  $\text{NO}_2$ , OH, OMe, SMe, COMe, CN,  $\text{COOR}^7$ ,  $\text{SO}_3\text{R}^7$ , COEt,  $\text{NHCOCH}_3$ , or aryl;

an aryl moiety can be a 5 or 6 membered aromatic heterocyclic ring (containing up to 4 hetero atoms independently selected from N, O, or S) or a 6 membered aromatic non-heterocyclic ring or a polycycle;

$\text{C}_{1-6}$  alkyl moieties can be straight chain or branched;

optionally substituted  $\text{C}_{1-6}$  alkyl moieties can be straight chain or branched;

$\text{C}_{2-6}$  alkenyl moieties can be straight chain or branched; and

optionally substituted  $\text{C}_{2-6}$  alkenyl moieties can be straight chain or branched;

with the proviso that said compound is not:

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][methylamino]carboxamide, or

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl}amino]carboxamide, or

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-chlorophenyl]carboxamide, or

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-chloro-3-pyridyl]carboxamide, or

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][trichloromethyl]carboxamide.

Examples of suitable  $\text{C}_{1-6}$  alkyl groups include but are not limited to methyl, ethyl, n-propyl, i-propyl, n-butyl, and t-butyl.


Halogens are typically F, Cl, Br, and I.

Examples of 5 or 6 membered ring moieties include, but are not restricted to, phenyl, furanyl, thienyl, imidazolyl, pyridyl, pyrrolyl, oxazolyl, isoxazolyl, triazolyl, pyrazolyl, tetrazolyl, thiazolyl and isothiazolyl. Examples of polycyclic moieties include, but are not restricted to, naphthyl, benzothiazolyl, benzofuranyl, benzimidazolyl, quinolyl, isoquinolyl, indolyl, quinoxalyl, quinazolinyl and benzothienyl.

### Synthetic Approaches


The compounds disclosed in this invention may be readily prepared according to a variety of synthetic manipulations, all of which would be familiar to one skilled in the art. In the general syntheses set forth below, the labeled substituents have the same identifications as set out in the definitions of the compounds above.

Compounds of general formula (I) can be obtained via a variety of synthetic routes all of which would be familiar to one skilled in the art. The reaction of isocyanates with amines is a commonly practised method for the formation of ureas (see Org. Syn. Coll. Vol. V, (1973), 555). Amine (IV), 3-(4-bromo-1-methylpyrazole-3-yl)phenylamine, commercially available from Maybridge Chemical Company, Catalog No. KM01978, CAS No. 175201-77-1] reacts readily with isocyanates (V) in inert solvents such as halocarbons to yield the desired ureas of general formula (I) wherein  $\text{R}^1=\text{R}^2=\text{H}$ :




37

-continued




Alternatively the amine (IV) can be converted to the corresponding isocyanate (VI) by the action of phosgene or a suitable phosgene equivalent, e.g. triphosgene, in an inert solvent such as a halocarbon in the presence of an organic base such as triethylamine or ethyldiisopropylamine. Isocyanate (VI) reacts with amines of general formula (VII), in an analogous fashion to that described above for the reaction of (IV) with (V), yielding the desired ureas of general formula (I) wherein  $R^1 = H$ :




38

Alternatively wherein the isocyanate of general formula (V) is not commercially available it can be prepared from the corresponding amine of general formula (VIII) in an analogous procedure to that described above for the preparation of (VI). Reaction of these isocyanates with (IV) would again yield the requisite ureas of general formula (I) wherein  $R^1 = R^2 = H$ :




39



40

Amines of general formula (VII) are also readily converted to activated isocyanate equivalents of general formula (IX) by the sequential action of carbonyldiimidazole and methyl iodide in tetrahydrofuran and acetonitrile respectively (R. A. Batey et al, *Tetrahedron Lett.*, (1998), 39, 6267–6270.) Reaction of (IX) with (IV) in an inert solvent such as a halocarbon would yield the requisite ureas of general formula (I) wherein  $R^1 = H$ :




55



Amine (IV) may be monomethylated according to the procedure of J. Barluenga et al, *J. Chem. Soc., Chem. Commun.*, (1984), 20, 1334–1335, or alkylated according to the procedure of P. Marchini et al, *J. Org. Chem.*, (1975), 40(23), 3453–3456, to yield compounds of general formula (X) wherein  $R^1 = \text{lower alkyl}$ . These materials may be reacted as above with reagents of general formula (V) and (IX) as depicted below:

60


41

(I)  $R^1 = \text{lower alkyl}, R^2 = \text{H}$ 


42

(X)  $R^1 = \text{lower alkyl}$ 


+



(IX)


(I)  $R^1 = \text{lower alkyl}$ 

Compounds of general formula (II) can similarly be obtained via a variety of synthetic manipulations, all of which would be familiar to one skilled in the art. The reaction of amine (IV) with chloroformates (see Org. Syn. Coll. Vol. IV, (1963), 780) of general formula (XI) in an inert solvent such as ether or halocarbon in the presence of a tertiary base such as triethylamine or ethyldiisopropylamine readily yields the requisite carbamates of general formula (II) wherein  $R^1 = \text{H}$ . Analogously amines of general formula (X) react similarly with chloroformates (XI) to yield the requisite carbamates of general formula (II) wherein  $R^1 = \text{lower alkyl}$ :




(IV)



+




(XI)





-continued

(II)  $R^1 = \text{H}$ (X)  $R^1 = \text{lower alkyl}$ 

+




(XI)

(II)  $R^1 = \text{lower alkyl}$ 


65

43

An alternative route employs the ready reaction of an alcohol with an isocyanate. Thus isocyanate (VI) described previously reacts readily with alcohols (XII) in an aprotic solvent such as ether or chlorocarbon to yield the desired carbamates of general formula (II) wherein  $R^1=H$ :




Chloroformates of general formula (XI) not commercially available may be readily prepared from the corresponding alcohol (XII) in an inert solvent such as toluene, chlorocarbon or ether by the action of excess phosgene (see *Org. Syn. Coll. Vol. III*, (1955), 167):



Compounds of general formula (III) can be obtained via a variety of synthetic manipulations, all of which would be familiar to one skilled in the art. The reaction of amine (IV) with acid chlorides (see *Org. Syn. Coll. Vol. V*, (1973), 336) wherein  $R^1=H$  is readily achieved in an inert solvent such



44

as chloroform or dichloromethane in the presence of an organic base such as triethylamine or ethyldiisopropylamine. In an identical fashion amines of general formula (X) would react with acid chlorides (XIII) to yield the desired amides (III) wherein  $R^1$ =lower alkyl:




Alternatively the corresponding acids of general formula (XIV) may be coupled with dicyclohexylcarbodiimide (DCC)/hydroxybenzotriazole (HOBT) (see W. König et al, *Chem. Ber.*, (1970), 103, 788) or hydroxybenzotriazole (HOBT)/2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (HBTU) (see M. Bematowicz et al., *Tetrahedron Lett.*, (1989), 30, 4645) as condensing agents in dimethylformamide or chloroform to amines (IV) and (X) respectively yielding products identical to those described in the previous scheme:





The acids of general formula (XIV) are readily converted to the corresponding acid chlorides (XIII) by the action of thionyl chloride or oxalyl chloride in the presence of catalytic dimethylformamide:



A third aspect of the present invention provides a compound of formula (A) or a solvate or physiologically functional derivative thereof for use as a therapeutic agent, specifically as a modifier of the activity of the serotonin 5-HT<sub>2A</sub> receptor. Modifiers of the activity of the serotonin 5-HT<sub>2A</sub> receptor are believed to be of potential use for the treatment or prophylaxis of CNS, gastrointestinal, cardiovascular, and inflammatory disorders. Compounds of the formula (A) may be administered by oral, sublingual, parenteral, rectal, or topical administration. In addition to

the neutral forms of compounds of formula (A) by appropriate addition of an ionizable substituent, which does not alter the receptor specificity of the compound, physiologically acceptable salts of the compounds may also be formed and used as therapeutic agents. Different amounts of the compounds of formula (A) will be required to achieve the desired biological effect. The amount will depend on factors such as the specific compound, the use for which it is intended, the means of administration, and the condition of the treated individual. A typical dose may be expected to fall in the range of 0.001 to 200 mg per kilogram of body weight of the treated individual. Unit does may contain from 1 to 200 mg of the compounds of formula (A) and may be administered one or more times a day, individually or in multiples. In the case of the salt or solvate of a compound of formulas (A), the dose is based on the cation (for salts) or the unsolvated compound.

A fourth aspect of the present invention provides pharmaceutical compositions comprising at least one compound of formula (A) and/or a pharmacologically acceptable salt or solvate thereof as an active ingredient combined with at least

one pharmaceutical carrier or excipient. Such pharmaceutical compositions may be used in the treatment of clinical conditions for which a modifier of the activity of the serotonin 5-HT<sub>2A</sub> receptor is indicated. At least one compound of formula (A) may be combined with the carrier in either solid or liquid form in a unit dose formulation. The pharmaceutical carrier must be compatible with the other ingredients in the composition and must be tolerated by the individual recipient. Other physiologically active ingredients may be incorporated into the pharmaceutical composition of the invention if desired, and if such ingredients are compatible with the other ingredients in the composition. Formulations may be prepared by any suitable method, typically by uniformly mixing the active compound(s) with liquids or finely divided solid carriers, or both, in the required proportions, and then, if necessary, forming the resulting mixture into a desired shape.

Conventional excipients, such as binding agents, fillers, acceptable wetting agents, tableting lubricants, and disintegrants may be used in tablets and capsules for oral administration. Liquid preparations for oral administration may be in the form of solutions, emulsions, aqueous or oily suspensions, and syrups. Alternatively, the oral preparations may be in the form of dry powder which can be reconstituted with water or another suitable liquid vehicle before use. Additional additives such as suspending or emulsifying agents, non-aqueous vehicles (including edible oils), preservatives, and flavorings and colorants may be added to the liquid preparations. Parenteral dosage forms may be prepared by dissolving the compound of the invention in a suitable liquid vehicle and filter sterilizing the solution before filling and sealing an appropriate vial or ampoule. These are just a few examples of the many appropriate methods well known in the art for preparing dosage forms.

The fifth aspect of the present invention provides for the use of a compound of formula (A) in the preparation of a medicament for the treatment of a medical condition for which a modifier of the activity of the serotonin 5-HT<sub>2A</sub> receptor is indicated.

A sixth aspect of the present invention provides for a method of treatment of a clinical condition of a mammal, such as a human, for which a modifier of the activity of the serotonin 5-HT<sub>2A</sub> receptor is indicated, which comprises the administration to the mammal of a therapeutically effective amount of a compound of formula (A) or a physiologically acceptable salt, solyate, or physiologically functional derivative thereof.

#### Experimental Data

Mass spectra were recorded on a Micromass Platform LC with Gilson HPLC. Infra-red spectra were recorded on a Nicolet Avatar 360 FT-IR. Melting points were recorded on a Electrothermal IA9200 apparatus and are uncorrected. Proton nuclear magnetic resonance spectra were recorded on a Bruker 300 MHz machine. Chemical shifts are given with respect to tetramethylsilane. In the text the following abbreviations are used; s (singlet), d (doublet), t (triplet), m (multiplet) or combinations thereof. Chemical shifts are quoted in parts per million (ppm) and with coupling constants in Hertz.

Thin layer chromatography was carried out using aluminium backed silica plates (250  $\mu$ L; GF<sub>254</sub>). HPLC was recorded either on a HP Chemstation 1100 HPLC using a Hichrom 3.5 C18 reverse phase column (50 mm $\times$ 2.1 mm i.d.). Linear gradient elution over 5 minutes—95% water (+0.1% TFA)/5% acetonitrile (+0.05% TFA) down to 5%

water/95% acetonitrile. Flow rate 0.8 mL/min [Method A]; or on a Hichrom 3.5 C18 reverse phase column (100 mm $\times$ 3.2 mm i.d.). Linear gradient elution over 11 minutes—95% water (+0.1% TFA)/5% acetonitrile (+0.05% TFA) down to 5% water/95% acetonitrile. Flow rate 1 mL/min [Method B]. Samples were routinely monitored at 254 nM unless otherwise stated.

All reagents were purchased from commercial sources.

#### Experiment 1

##### Preparation and Analysis of 103487

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl}amino]carboxamide

This compound is commercially available from Maybridge Chemical Company, Catalog No. KM04515.

#### Experiment 2

##### Preparation and Analysis of 116100

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-methoxyphenoxy]carboxamide

To 4-methoxyphenylchloroformate (19 mg, 0.10 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (25 mg, 0.10 mmol) and triethylamine (14  $\mu$ L, 0.10 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (40% EtOAc/hexane) gave the title compound as a colourless solid (21 mg, 52%), m.p. 140.3–141.8° C. (EtOAc/hexane).

IR:  $\nu_{max}$ =1748, 1592, 1504, 1412, 1190, 835, 764, 676 cm<sup>-1</sup>. MS (ES+): m/z (%)=404 (M+H<sup>81</sup>Br, 100), 402 (M+H<sup>79</sup>Br, 90).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD):  $\delta$ =3.80 (3H, s, CH<sub>3</sub>), 3.81 (3H, s, CH<sub>3</sub>), 6.91–6.98 (2H, m, ArH), 7.07–7.18 (3H, m, ArH), 7.42–7.53 (4H, m, ArH). HPLC: retention time 3.28 mins [Method A]. Tlc: Rf 0.4 (EtOAc/hexane).

#### Experiment 3

##### Preparation and Analysis of 116101

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-trifluoromethoxyphenyl]carboxamide

To 4-(trifluoromethoxy)benzoyl chloride (19  $\mu$ L, 0.12 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol) and triethylamine (17  $\mu$ L, 0.12 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL). The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (40 mg, 76%), m.p. 138.6–139.6° C. (EtOAc/hexane).

MS (ES+): m/z (%)=442 (M+H<sup>81</sup>Br, 93), 440 (M+H<sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>):  $\delta$ =3.79 (3H, s, CH<sub>3</sub>), 7.27 (1H, m, ArH), 7.45–7.60 (3H, m, ArH), 7.65 (1H, s, ArH), 7.87 (2H, m, ArH), 8.09 (2H, m, ArH), 10.51 (1H, s, NH).

HPLC: retention time 3.60 min [Method A]. TLC: Rf 0.40 (50% EtOAc/hexane).

#### Experiment 4

##### Preparation and Analysis of 116102

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][2-thienyl]carboxamide

To thiophene-2-carbonyl chloride (11  $\mu$ L, 0.09 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (25 mg, 0.09 mmol) and triethylamine (14  $\mu$ L, 0.09 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5

## 49

mL). The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (24 mg, 68%), m.p. 127.8–128.6° C. (EtOAc/hexane).

MS (ES+): m/z (%)=364 (M+H <sup>81</sup>Br, 96), 362 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ=3.81 (3H, s, CH<sub>3</sub>), 7.19 (2H, m, ArH), 7.48–7.58 (2H, m, ArH), 7.68–7.83 (3H, m, ArH), 7.93 (1H, dd, J=1.0, 3.8, ArH).

HPLC: retention time 3.12 min [Method A]. TLC: Rf 0.30 (30% EtOAc/hexane).

## Experiment 5

## Preparation and Analysis of 116115

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][{(4-trifluoromethoxy)phenyl)methyl}amino]carboxamide

To a stirred solution of triphosgene (12 mg, 0.04 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol) and triethylamine (33 μL, 0.24 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL). After 1 h, 4-(trifluoromethoxy)benzylamine (23 mg, 0.12 mmol) was added. The reaction mixture was stirred for 16 h and concentrated. Chromatography on flash silica (75%EtOAc/hexane) gave the title compound as a colourless solid (38 mg, 68%), m.p. 144.6–145.8° C. (EtOAc/hexane).

IR:  $\nu_{max}$ =1626, 1558, 1278, 1160, 969, 871, 789, 703 cm<sup>-1</sup>. MS (ES+): m/z (%)=471 (M+H <sup>81</sup>Br, 91), 469 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ=3.81 (3H, s, CH<sub>3</sub>), 4.42 (2H, s, CH<sub>2</sub>), 7.06 (1H, d, J=7.1, ArH), 7.24 (2H, d, J=8.4, ArH), 7.37–7.52 (6H, m, ArH). HPLC: retention time 3.06 mins [Method A]. Tlc: Rf 0.5 (EtOAc).

## Experiment 6

## Preparation and Analysis of 116120

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][4-chlorophenyl]carboxamide

To 4-chlorobenzoyl chloride (15 mg, 0.08 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (21 mg, 0.08 mmol) and triethylamine (12 μL, 0.08 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (0.5 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (50% EtOAc/hexane) gave the title compound as a colourless solid (23 mg, 72%), m.p. 184.4–184.8° C. (EtOAc/hexane).

MS (ES+): m/z (%)=394 (M+H <sup>81</sup>Br <sup>37</sup>Cl, 34), 392 (M+H <sup>79</sup>Br <sup>37</sup>Cl (<sup>81</sup>Br <sup>35</sup>Cl), 100), 390 (M+H <sup>79</sup>Br <sup>35</sup>Cl, 67).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.79 (3H, s, CH<sub>3</sub>), 7.25 (1H, d, J=7.9, ArH), 7.51–7.65 (3H, m, ArH), 7.69 (1H, s, ArH), 7.90 (2H, m, ArH), 8.00 (2H, m, ArH), 10.51 (1H, s, NH).

HPLC: retention time 3.40 min [Method A]. TLC: Rf 0.35 (50% EtOAc/hexane).

## Experiment 7

## Preparation and Analysis of 116137

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-[4-(trifluoromethoxy)phenyl]acetamide

A solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (35 mg, 0.14 mmol) and triethylamine (23 μL, 0.177 mmol) in DMF (0.5 mL) was added in one portion to a stirred solution of 4-trifluoromethoxyphenylacetic acid (31 mg, 0.14 mmol), HBTU (53 mg, 0.14 mmol) and HOBT (19 mg, 0.14 mmol) in DMF (1 mL). The mixture was heated at 70° C. for 24 h and then quenched with aqueous sodium bicarbonate solution. Ethyl acetate was added and the organic phase separated, washed with water (×3), brine, dried (MgSO<sub>4</sub>) and evaporated. Chromatography on flash

## 50

silica (50%EtOAc/hexane) gave the title compound as a colourless solid (43 mg, 68%), m.p. 141.2–142.5° C. (EtOAc/hexane).

IR:  $\nu_{max}$ =1684, 1592, 1510, 1253, 1217, 1157, 987, 798, 700 cm<sup>-1</sup>.

MS (ES+): m/z (%)=456 (M+H <sup>81</sup>Br, 100), 454 (M+H <sup>79</sup>Br, 94).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.72 (2H, s, CH<sub>2</sub>), 3.75 (3H, s, CH<sub>3</sub>), 7.17 (1H, d, J=7.7, ArH), 7.33 (2H, d, J=8.7, ArH), 7.38–7.51 (3H, m, ArH), 7.62–7.73 (3H, m, ArH), 10.44 (1H, s, NH).

HPLC: retention time 3.52 min [Method A].

## Experiment 8

## Preparation and Analysis of 116174

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(3-fluorophenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol), 3-fluorophenylacetic acid (18 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound 12 mg, 26%. Rf 0.41 (ethyl acetate-toluene, 1:1).

HPLC (Method B): retention time 7.07 min (100%).  $\delta_H$  (CDCl<sub>3</sub>) 3.77 (2H, s), 3.83 (3H, s), 7.02–7.20 (4H, m), 7.54 (1H, s), 7.60–7.63 (1H, m). MS (AP+): m/z (%)=390 (M+H <sup>81</sup>Br, 100), 388(M+H <sup>79</sup>Br, 100).

## Experiment 9

## Preparation and Analysis of 116175

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(3-methoxyphenyl)acetamide

A solution of 3-methoxyphenylacetyl chloride (0.02 ml, 0.12 mmol) in dichloromethane (0.75 ml) was added dropwise at 0° C. to a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol) and triethylamine (0.02 ml, 0.13 mmol) in dichloromethane (0.75 ml). The resulting mixture was stirred at room temperature for 16 h and then poured into brine. The organic layer was washed with more brine then dried over magnesium sulphate and concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (9 mg, 19%). Rf 0.30 (ethyl acetate-toluene, 1:1).

HPLC (Method B): retention time 8.62 min (97.09%).  $\Delta_H$  (CDCl<sub>3</sub>) 3.76 (2H, s), 3.82 (3H, s), 3.85 (3H, s), 6.84–6.90 (3H, m), 7.07–7.44 (5H, m), 7.53 (1H, s), 7.60 (1H, br s). MS (AP+): m/z (%)=402 (M+H <sup>81</sup>Br, 100), 400 (M+H <sup>79</sup>Br, 95).

## Experiment 10

## Preparation and Analysis of 116176

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(2-fluorophenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol), 2-fluorophenylacetic acid (18 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-

Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (15 mg, 32%). Rf 0.52 (ethyl acetate-toluene, 1:1).

HPLC (Method B): retention time 7.28 min (100%).  $\delta_H$  (CDCl<sub>3</sub>) 3.79 (2H, s), 3.83 (3H, s), 7.11–7.23 (3H, m), 7.30–7.55 (6H, m), 7.61–7.64 (1H, m). MS (AP+): m/z (%)=390 (M+H <sup>81</sup>Br, 100), 388 (M+H <sup>79</sup>Br, 100).

## Experiment 11

## Preparation and Analysis of 116177

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(4-nitrophenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol), 4-nitrophenylacetic acid (22 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropylethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (ethyl acetate-toluene, 1:1), giving the title compound (9 mg, 18%). Rf 0.19 (ethyl acetate-toluene, 1:1).

HPLC (Method B): retention time 7.22 min (94.30%).  $\delta_H$  (CDCl<sub>3</sub>) 3.83 (3H, s), 3.87 (2H, s), 7.18–7.23 (1H, m), 7.42–7.65 (7H, m), 8.22–8.30 (2H, m). MS (AP+): m/z (%)=417 (M+H <sup>81</sup>Br, 100), 415 (M+H <sup>79</sup>Br, 100).

## Experiment 12

## Preparation and Analysis of 116178

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]-2-(2-methoxyphenyl)acetamide

A mixture of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (30 mg, 0.12 mmol), 2-methoxyphenylacetic acid (20 mg, 0.12 mmol), 1-hydroxybenzotriazole hydrate (16 mg, 0.12 mmol) and 2-(1H-benzotriazole-1-yl)-1,1,3,3-tetramethyluronium hexafluoro-phosphate (46 mg, 0.12 mmol) were dissolved in chloroform (1.5 ml). N,N-Diisopropyl-ethylamine (0.02 ml, 0.13 mmol) was added and the mixture stirred at room temperature for 16 h. The reaction mixture was then poured into brine and the organic layer washed with further brine, dried over magnesium sulphate and then concentrated in vacuo. The crude product was purified by column chromatography (chloroform-methanol, 99:1), giving the title compound (18 mg, 38%) as a colourless solid. Rf 0.65 (chloroform-methanol, 98:2).

HPLC (Method B): retention time 7.16 min (100%).  $\delta_H$  (CDCl<sub>3</sub>) 3.76 (2H, s), 3.83 (3H, s), 3.98 (3H, s), 6.97–7.06 (2H, m), 7.11–7.16 (1H, m), 7.31–7.50 (4H, m), 7.53 (1H, s), 7.57–7.60 (1H, m), 7.91 (1H, br s). MS (AP-): m/z (%)=400 (M-H <sup>81</sup>Br, 90), 398 (M-H <sup>79</sup>Br, 100).

## Experiment 13

## Preparation and Analysis of 116192

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(1,1-dimethylethoxy)carboxamide

To di-tert-butyl dicarbonate (36 mg, 0.17 mmol) in methanol (1 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (42 mg, 0.17

mmol) in methanol (1 mL). The mixture was stirred for 16 h and concentrated. Chromatography on flash silica (40%EtOAc/hexane) gave the title compound as a colourless solid (29 mg, 49%) (EtOAc/hexane).

MS (CI-): m/z (%)=352 (M-H <sup>81</sup>Br, 100), 350 (M-H <sup>79</sup>Br, 96).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>):  $\delta$ =1.46 (9H, s, 3×CH<sub>3</sub>), 3.73 (3H, s, CH<sub>3</sub>), 7.07 (1H, m, ArH), 7.42 (1H, t, J=7.7, ArH), 7.53–7.60 (2H, m, ArH), 7.64 (1H, s, ArH), 9.57 (1 H, s, NH).

HPLC: retention time 7.15 min [Method B].

One or the other (as indicated) of the two following synthetic protocols was used to generate each of the compounds below:

## Protocol A

To an isocyanate (1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (1 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL). The mixture was stirred for 16 hours and concentrated. Chromatography on flash silica (20%–80% EtOAc/hexane) followed by recrystallisation gave the pure urea.

## Protocol B

To a stirred solution of triphosgene (0.33 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL) was added dropwise a solution of 3-(3-aminophenyl)-4-bromo-1-methylpyrazole (1 mmol) and triethylamine (2 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4 mL). After 1 hour, an aniline was added (1 mmol). The reaction mixture was stirred for 16 hours and concentrated. Chromatography on flash silica (20%–80%EtOAc/hexane) followed by recrystallisation gave the pure urea.

## Experiment 14

## Preparation and Analysis of 116079

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][(4-methylthiophenyl)amino]carboxamide

[Protocol A]—4-(methylthio)phenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+): m/z (%)=419 (M+H <sup>81</sup>Br, 100), 417 (M+H <sup>79</sup>Br, 94).

<sup>1</sup>H-NMR (MeOH d<sub>4</sub>):  $\delta$ =2.42 (3H, s, SCH<sub>3</sub>), 3.81 (3H, s, NCH<sub>3</sub>), 7.06 (1H, m, ArH), 7.22 (2H, m, ArH), 7.37 (2H, m, ArH), 7.42–7.61 (4H, m, ArH).

HPLC: retention time 3.35 min [Method A].

## Experiment 15

## Preparation and Analysis of 116081

N-[3-(4-bromo-1-methylpyrazol-3-yl)phenyl][(4-chlorophenyl)amino]carboxamide

[Protocol A]—4-chlorophenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+): m/z (%)=409 (M+H <sup>81</sup>Br <sup>37</sup>Cl, 19), 407 (M+H <sup>79</sup>Br <sup>37</sup>Cl (<sup>81</sup>Br <sup>35</sup>Cl), 100), 405 (M+H <sup>79</sup>Br <sup>35</sup>Cl, 81).

<sup>1</sup>H-NMR (MeOH d<sub>4</sub>):  $\delta$ =3.81 (3H, s, CH<sub>3</sub>), 7.07 (1H, m, ArH), 7.23 (2H, m, ArH), 7.36–7.60 (6H, m, ArH).

HPLC: retention time 3.42 min [Method A].

## Experiment 16

## Preparation and Analysis of 116082

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenyl)carboxamide

[Protocol A]—4-fluorophenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+): m/z (%)=391 (M+H <sup>81</sup>Br, 96), 389 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (MeOH d<sub>4</sub>):  $\delta$ =3.81 (3H, s, CH<sub>3</sub>), 6.93–7.11 (3H, m, ArH), 7.37–7.61 (6H, m, ArH).

HPLC: retention time 3.11 min.

## 53

## Experiment 17

## Preparation and Analysis of 116087

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[2-(trifluoromethoxy)phenyl]carboxamide  
[Protocol A]—2-(trifluoromethoxy)phenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=457 (M+H <sup>81</sup>Br, 100), 455 (M+H <sup>79</sup>Br, 95).

<sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>):  $\delta$ =3.79 (3H, s, CH<sub>3</sub>), 7.06–7.18 (2H, m, ArH), 7.38–7.49 (2H, m, ArH), 7.51–7.62 (2H, m, ArH), 7.65 (1H, m, ArH), 7.71 (1H, s, ArH), 8.24 (1H, dd, J=1.1, 8.2, ArH), 8.56 (1H, s, NH), 9.49 (1H, s, NH).

HPLC: retention time 3.40 min.

## Experiment 18

## Preparation and Analysis of 116089

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-nitrophenyl)carboxamide  
[Protocol A]—2-nitrophenyl isocyanate yellow solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=418 (M+H <sup>81</sup>Br, 98), 416 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>):  $\delta$ =<sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>):  $\square$ =3.79 (3H, s, NCH<sub>3</sub>), 7.14 (1H, m, ArH), 7.24 (1H, m, ArH), 7.50 (1H, t, J=7.7, ArH), 7.60 (2H, m, ArH), 7.67 (1H, s, ArH), 7.71 (1H, s, ArH), 8.10 (1H, m, ArH), 8.29 (1H, m, ArH), 9.65 (1H, s, NH), 10.09 (1H, s, NH).

HPLC: retention time 3.10 min [Method A].

## Experiment 19

## Preparation and Analysis of 116091

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenyl)carboxamide  
[Protocol A]—4-methoxyphenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=403 (M+H <sup>81</sup>Br, 100), 401 (M+H <sup>79</sup>Br, 96).

<sup>1</sup>H-NMR (DMSO *d*<sub>6</sub>):  $\delta$ =3.71 (3H, s, OCH<sub>3</sub>), 3.79 (3H, s, NCH<sub>3</sub>), 6.87 (2H, d, J=8.9, ArH), 7.96 (1H, d, J=7.5, ArH), 7.39 (2H, d, J=8.9, ArH), 7.45–7.61 (3H, m, ArH), 7.65 (1H, s, ArH), 8.52 (1H, s, NH), 8.84 (1H, s, NH).

HPLC: retention time 3.08 min.

## Experiment 20

## Preparation and Analysis of 116092

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenyl)carboxamide  
[Protocol A]—*o*-tolyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=387 (M+H <sup>81</sup>Br, 94), 385 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (MeOH *d*<sub>4</sub>):  $\delta$ =2.29 (3H, s, CH<sub>3</sub>), 3.81 (3H, s, NCH<sub>3</sub>), 7.03 (1H, dt, J=1.1, 7.5, ArH), 7.09 (1H, dt, J=1.1, 7.5, ArH), 7.13–7.22 (2H, m, ArH), 7.45 (1H, t, J=7.9, ArH), 7.49–7.57 (2H, m, ArH), 7.60–7.68 (2H, m, ArH).

HPLC: retention time 2.96 min.

## Experiment 21

## Preparation and Analysis of 116097

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethyl)phenyl]carboxamide  
[Protocol A]—4-(trifluoromethyl)phenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=441 (M+H <sup>81</sup>Br, 94), 439 (M+H <sup>79</sup>Br, 100).

## 54

H-NMR (MeOH *d*<sub>4</sub>):  $\delta$ =3.82 (3H, s, CH<sub>3</sub>), 7.04–7.16 (3H, m, ArH), 7.20–7.47 (6H, m, ArH).

HPLC: retention time 3.56 min.

## Experiment 22

## Preparation and Analysis of 116105

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-chlorophenyl)carboxamide  
[Protocol A]—3-chlorophenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=409 (M+H <sup>81</sup>Br <sup>37</sup>Cl, 26), 407 (M+H <sup>79</sup>Br <sup>37</sup>Cl (<sup>81</sup>Br <sup>35</sup>Cl), 100), 405 (M+H <sup>79</sup>Br <sup>35</sup>Cl, 70).

<sup>1</sup>H-NMR (MeOH *d*<sub>4</sub>):  $\delta$ =3.81 (3H, s, NCH<sub>3</sub>), 7.04 (1H, m, ArH), 7.10 (1H, m, ArH), 7.28 (2H, m, ArH), 7.47 (1H, t, J=7.8, ArH), 7.55 (1H, m, ArH), 7.63 (1H, m, ArH), 7.68 (1H, s, ArH), 7.73 (1H, m, ArH), 9.04 (2H, s, NH).

HPLC: retention time 3.20 min [Method A].

## Experiment 23

## Preparation and Analysis of 116108

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-chlorophenyl)carboxamide  
[Protocol A]—2-chlorophenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=409 (M+H <sup>81</sup>Br <sup>37</sup>Cl, 24), 407 (M+H <sup>79</sup>Br <sup>37</sup>Cl (<sup>81</sup>Br <sup>35</sup>Cl), 100), 405 (M+H <sup>79</sup>Br <sup>35</sup>Cl, 72).

<sup>1</sup>H-NMR (MeOH *d*<sub>4</sub>):  $\delta$ =3.81 (3H, s, NCH<sub>3</sub>), 7.03 (1H, m, ArH), 7.11 (1H, m, ArH), 7.28 (1H, m, ArH), 7.35–7.53 (3H, m, ArH), 7.55 (1H, s, ArH), 7.62 (1H, m, ArH), 8.11 (1H, m, ArH).

HPLC: retention time 3.13 min.

## Experiment 24

## Preparation and Analysis of 116110

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[4-(methylethyl)phenyl]carboxamide  
[Protocol A]—4-isopropylphenyl isocyanate colourless solid (THF/hexane)

MS (ES+):  $m/z$  (%)=415 (M+H <sup>81</sup>Br, 100), 413 (M+H <sup>79</sup>Br, 92).

<sup>1</sup>H-NMR (MeOH *d*<sub>4</sub>):  $\delta$ =1.23 (6H, d, J=6.8, 2×CH<sub>3</sub>), 2.86 (1H, septet, J=6.8, CH), 3.82 (3H, s, NCH<sub>3</sub>), 7.09 (1H, m, ArH), 7.16 (2H, d, J=7.6, ArH), 7.31 (2H, d, J=7.6, ArH), 7.42–7.51 (2H, m, ArH), 7.54 (1H, s, ArH), 7.59 (1H, m, ArH).

HPLC: retention time 3.66 min.

## Experiment 25

## Preparation and Analysis of 116111

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-methoxyphenyl)carboxamide  
[Protocol A]—3-methoxyphenyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=403 (M+H <sup>81</sup>Br, 100), 401 (M+H <sup>79</sup>Br, 96).

<sup>1</sup>H-NMR (MeOH *d*<sub>4</sub>):  $\delta$ =3.73 (3H, s, OCH<sub>3</sub>), 3.81 (3H, s, NCH<sub>3</sub>), 6.59 (1H, m, ArH), 6.91 (1H, in, ArH), 7.08 (1H, m, ArH), 7.14 (2H, m, ArH), 7.39–7.61 (4H, m, ArH).

HPLC: retention time 2.90 min.

## Experiment 26

## Preparation and Analysis of 116112

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3-methylphenyl)carboxamide  
[Protocol A]—*m*-tolyl isocyanate colourless solid (EtOAc/hexane)

MS (ES+):  $m/z$  (%)=387 (M+H <sup>81</sup>Br, 100), 385 (M+H <sup>79</sup>Br, 96).

## 55

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=2.26 (3H, s, CH<sub>3</sub>), 3.76 (3H, s, NCH<sub>3</sub>), 6.79 (1H, m, ArH), 7.06–7.22 (3H, m, ArH), 7.29 (1H, m, ArH), 7.43–7.62 (3H, m, ArH), 7.68 (1H, s, ArH), 8.65 (1H, s, NH), 8.89 (1H, s, NH).

HPLC: retention time 3.05 min [Method A].

## Experiment 27

## Preparation and Analysis of 116113

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-methyl-N-[4-(trifluoromethoxy)phenyl]carboxamide [Protocol B]-N-methyl-4-(trifluoromethoxy)aniline pale yellow solid (EtOAc/hexane)

MS (ES<sup>+</sup>): m/z (%)=471 (M+H <sup>81</sup>Br, 88), 469 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (MeOH d<sub>4</sub>): δ=3.35 (3H, s, NCH<sub>3</sub>), 3.81 (3H, s, NCH<sub>3</sub>), 7.09 (1H, m, ArH), 7.25–7.51 (8H, m, ArH).

HPLC: retention time 3.56 min [Method A].

## Experiment 28

## Preparation and Analysis of 116119

N-[4-(tert-butyl)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide [Protocol B]-4-tert-butylaniline colourless solid (OAc/hexane)

MS (ES<sup>+</sup>): m/z (%)=429 (M+H <sup>81</sup>Br, 98), 427 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=1.27 (9H, s, 3'CH<sub>3</sub>), 3.79 (3H, s, NCH<sub>3</sub>), 7.07 (1H, d, J=7.5, ArH), 7.29 (2H, d, J=8.7, ArH), 7.37 (2H, d, J=8.7, ArH), 7.45 (1H, t, J=7.5, ArH), 7.51–7.60 (2H, m, ArH), 7.66 (1H, s, ArH), 8.65 (1H, s, NH), 8.83 (1H, s, NH).

HPLC: retention time 3.77 min.

## Experiment 29

## Preparation and Analysis of 116122

N-[4-(dimethylamino)phenyl]{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide [Protocol B]-N,N-dimethyl-p-phenylenediamine colourless solid (EtOAc/hexane)

MS (ES<sup>+</sup>): m/z (%)=416 (M+H <sup>81</sup>Br, 96), 414 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=2.86 (6H, s, NCH<sub>3</sub>), 3.80 (3H, s, NCH<sub>3</sub>), 6.80 (2H, m, ArH), 7.09 (1H, d, J=7.7, ArH), 7.28 (2H, m, ArH), 7.42 (1H, t, J=7.8, ArH), 7.52 (1H, m, ArH), 7.59 (1H, s, ArH), 7.67 (1H, s, ArH), 8.45 (1H, s, NH), 8.75 (1H, s, NH).

HPLC: retention time 2.07 min [Method A].

## Experiment 30

## Preparation and Analysis of 116138

N-(3,5-dichloro-4-methylphenyl){[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carboxamide [Protocol B]-3,5-dichloro-4-methylphenylamine colourless solid (EtOAc/hexane)

MS (ES<sup>+</sup>): m/z (%)=457 (M+H, 35), 455 (M+H, 100), 453 (M+H, 65).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=2.32 (3H, s, CH<sub>3</sub>), 3.79 (3H, s, NCH<sub>3</sub>), 7.11 (1H, d, J=7.4, ArH), 7.46 (1H, t, J=7.8, ArH), 7.50–7.64 (4H, m, ArH), 7.68 (1H, s, ArH), 9.02 (1H, s, NH), 9.09 (1H, s, NH).

HPLC: retention time 3.66 min.

## Experiment 31

## Preparation and Analysis of 116139

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[4-(trifluoromethylthio)phenyl]carboxamide [Protocol B]-4-(trifluoromethylthio)aniline colourless solid (EtOAc/hexane)

MS (ES<sup>+</sup>): m/z (%)=473 (M+H <sup>81</sup>Br, 100), 471 (M+H <sup>79</sup>Br, 94).

## 56

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.81 (3H, s, NCH<sub>3</sub>), 7.11 (1H, d, J=7.5, ArH), 7.47 (1H, t, J=7.9, ArH), 7.51–7.63 (6H, m, ArH), 7.66 (1H, s, ArH), 9.03 (1H, s, NH), 9.16 (1H, s, NH).

HPLC: retention time 3.76 min.

## Experiment 32

## Preparation and Analysis of 116141

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(cyclohexyl)carboxamide

[Protocol B]-cyclohexylamine colourless solid, m.p. 155.5–156.3° C. (EtOAc/hexane).

MS (ES<sup>+</sup>): m/z (%)=379 (M+H <sup>81</sup>Br, 93), 377 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=1.07–1.34 (5H, m, 5xCH), 1.52 (1H, m, CH), 1.63 (2H, m, 2xCH), 1.76 (2H, m, 2xCH), 3.48 (1H, m, NCH), 3.74 (3H, s, CH<sub>3</sub>), 6.15 (1H, d, J=7.8, ArH), 6.98 (1H, d, J=7.5, ArH), 7.32–7.43 (2H, m, ArH), 7.51 (1H, m, NH), 7.62 (1H, s, ArH), 8.50 (1H, s, NH).

HPLC: retention time 3.16 min [Method A].

TLC: retention factor 0.35 (50% EtOAc/hexane).

## Experiment 33

## Preparation and Analysis of 116143

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(phenylmethyl)carboxamide

[Protocol B]-benzylamine colourless solid, m.p. 144.5–146.2° C. (EtOAc/hexane).

IR:  $\square_{max}$ =1622, 1565, 1467, 1374, 1239, 973, 802, 752, 695 cm<sup>-1</sup>.

MS (ES<sup>+</sup>): m/z (%)=387 (M+H <sup>81</sup>Br, 89), 385 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (CD<sub>3</sub>OD): δ=3.81 (3H, s, CH<sub>3</sub>), 4.40 (2H, s, CH<sub>2</sub>), 7.05 (1H, m, ArH), 7.19–7.51 (9H, m, ArH).

HPLC: retention time 3.06 min [Method A].a

## Experiment 34

## Preparation and Analysis of 116144

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-fluorophenyl)carboxamide

[Protocol A]-2-fluorophenyl isocyanate colourless solid (DCNM/hexane)

MS (ES<sup>+</sup>): m/z (%)=391 (M+H <sup>81</sup>Br, 100), 389 (M+H <sup>79</sup>Br, 90).

<sup>1</sup>H-NMR (MeOH d<sub>4</sub>): δ=3.79 (3H, s, NCH<sub>3</sub>), 7.00–7.11 (4H, m, ArH), 7.40–7.56 (3H, m, ArH), 7.61 (1H, m, ArH), 8.09 (1H, m, ArH).

HPLC: retention time 3.01 min.

## Experiment 35

## Preparation and Analysis of 116145

2-({[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}carbonylamino)benzamide

[Protocol B]-2-aminobenzamide colourless solid (EtOAc/hexane)

MS (ES<sup>+</sup>): m/z (%)=399 (M+H -17 <sup>81</sup>Br, 100), 397 (M+H -17 <sup>79</sup>Br, 94).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.79 (3H, s, NCH<sub>3</sub>), 6.93–7.10 (2H, m, ArH), 7.45 (2H, t, J=7.8, ArH), 7.59–7.72 (5H, m, ArH), 8.22 (2H, m), 9.92 (1H, s, NH), 10.69 (1H, s, NH).

HPLC: retention time 2.88 min.

57

## Experiment 36

## Preparation and Analysis of 116147

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-cyanophenyl)carboxamide  
[Protocol B]—4-aminobenzonitrile colourless solid (EtOAc/hexane)

MS (ES+): m/z (%)=398 (M+H <sup>81</sup>Br, 100), 396 (M+H <sup>79</sup>Br, 96).

<sup>1</sup>H-NMR (MeOH d<sub>4</sub>): δ=3.81 (3H, s, NCH<sub>3</sub>), 7.12 (1H, m, ArH), 7.46–7.57 (3H, m, ArH), 7.62–7.69 (5H, m, ArH).

HPLC: retention time 3.12 min.

## Experiment 37

## Preparation and Analysis of AR116148

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-cyanophenyl)carboxamide  
[Protocol B]—2-aminobenzonitrile colourless solid (EtOAc/hexane)

MS (ES+): m/z (%)=398 (M+H <sup>81</sup>Br, 95), 396 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (CDCl<sub>3</sub>): δ=3.79 (3H, s, CH<sub>3</sub>), 7.13–7.28 (2H, m, ArH), 7.49 (1H, t, J=7.8, ArH), 7.57 (1H, m, ArH), 7.62 (1H, m, ArH), 7.65–7.71 (2H, m, ArH), 7.78 (1H, m, ArH), 8.07 (1H, d, J=8.6, ArH), 8.83 (1H, s, NH), 9.62 (1H, s, NH).

HPLC: retention time 3.05 min [Method A].

## Experiment 38

## Preparation and Analysis of 116182

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-fluorophenylmethyl)carboxamide  
[Protocol B]—4-fluorobenzylamine colourless solid, m.p. 185.5–186.6° C. (EtOAc/hexane).

MS (ES+): m/z (%)=405 (M+H <sup>81</sup>Br, 97), 403 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.75 (3H, s, CH<sub>3</sub>), 4.28 (2H, d, J=6.0, CH<sub>2</sub>), 6.73 (1H, t, J=5.9, NH), 7.01 (1H, d, J=7.5, ArH), 7.10–7.18 (2H, m, ArH), 7.27–7.41 (4H, m, ArH), 7.56 (1H, s, ArH), 7.62 (1H, s, ArH), 8.82 (1H, s, NH).

HPLC: retention time 3.10 min [Method A].

TLC: retention factor 0.25 (50% EtOAc/hexane).

## Experiment 39

## Preparation and Analysis of 116183

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3,4-dimethoxyphenylmethyl)carboxamide  
[Protocol B]—3,4-dimethoxybenzylamine colourless solid, m.p. 174.9–175.5° C. (EtOAc/hexane).

MS (CI+): m/z (%)=447 (M+H <sup>81</sup>Br, 100), 445 (M+H <sup>79</sup>Br, 92).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.71 (3H, s, CH<sub>3</sub>), 3.73 (3H, s, CH<sub>3</sub>), 3.76 (3H, s, CH<sub>3</sub>), 4.22 (2H, d, J=5.8, CH<sub>2</sub>), 6.62 (1H, t, J=5.7, NH), 6.80 (1H, m, ArH), 6.89 (2H, m, ArH), 6.98 (1H, m, ArH), 7.36–7.51 (3H, m, ArH), 7.63 (1H, s, ArH), 8.76 (1H, s, NH).

HPLC: retention time 2.86 min [Method A].

TLC: retention factor 0.20 (50% EtOAc/hexane).

## Experiment 40

## Preparation and Analysis of 116184

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(3,4,5-trimethoxyphenylmethyl)carboxamide  
[Protocol B]—3,4,5-trimethoxybenzylamine colourless; solid (EtOAc/hexane).

MS (CI+): m/z (%)=477 (M+H <sup>81</sup>Br, 100), 475 (M+H <sup>79</sup>Br, 95).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.63 (3H, s, OCH<sub>3</sub>), 3.75 (9H, s, 3×CH<sub>3</sub>), 4.21 (1H, d, J=5.9, CH<sub>2</sub>), 6.61 (2H, s, ArH), 6.65

58

(1H, t, J=5.9, NH), 6.99 (1H, m, ArH), 7.40 (1H, t, J=7.7, ArH), 7.45 (1H, m, ArH), 7.56 (1H, m, ArH), 7.64 (1H, s, ArH), 8.77 (1H, s, NH).

HPLC: retention time 5.91 min [Method B].

TLC: retention factor 0.50 (50% EtOAc/hexane).

## Experiment 41

## Preparation and Analysis of 116185

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(2-methylphenylmethyl)carboxamide  
[Protocol B]—2-methylbenzylamine colourless solid (EtOAc/hexane).

MS (CI+): m/z (%)=401 (M-H <sup>81</sup>Br, 96), 399 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=2.28 (3H, s, CH<sub>3</sub>), 3.76 (3H, s, NCH<sub>3</sub>), 4.28 (1H, d, J=5.8, CH<sub>2</sub>), 6.60 (1H, t, J=5.8, NH), 7.01 (1H, m, ArH), 7.15 (3H, m, ArH), 7.24 (1H, m, ArH), 7.38–7.50 (2H, m, ArH), 7.57 (1H, m, ArH), 7.65 (1H, s, ArH), 8.77 (1H, s, NH).

HPLC: retention time 2.74 min [Method A].

TLC: retention factor 0.20 (50% EtOAc/hexane).

## Experiment 42

## Preparation and Analysis of 116189

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-(4-methoxyphenylmethyl)carboxamide  
[Protocol B]—4-methoxybenzylamine colourless solid (EtOAc/hexane).

MS (CI+): m/z (%)=417 (M+H <sup>81</sup>Br, 94), 415 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=3.72 (3H, s, CH<sub>3</sub>), 3.77 (3H, s, NCH<sub>3</sub>), 4.22 (1H, d, J=5.9, CH<sub>2</sub>), 6.62 (1H, t, J=5.9, NH), 6.90 (2H, d, J=8.8, ArH), 7.00 (1H, m, ArH), 7.23 (2H, d, J=8.8, ArH), 7.39 (1H, t, J=7.8, ArH), 7.43 (1H, m, ArH), 7.56 (1H, m, ArH), 7.64 (1H, s, ArH), 8.73 (1H, s, NH).

HPLC: retention time 6.41 min [Method B].

TLC: retention factor 0.25 (50% EtOAc/hexane).

## Experiment 43

## Preparation and Analysis of 116194

{[3-(4-bromo-1-methylpyrazol-3-yl)phenyl]amino}-N-[2-(4-methoxy)phenylethyl]carboxamide  
[Protocol B]—2-(4-methoxyphenyl)ethylamine colourless solid (EtOAc/hexane).

MS (ES+): m/z (%)=431 (M+H <sup>81</sup>Br, 95), 429 (M+H <sup>79</sup>Br, 100).

<sup>1</sup>H-NMR (DMSO d<sub>6</sub>): δ=2.68 (2H, t, J=7.1, CH<sub>2</sub>), 3.31 (2H, m, CH<sub>2</sub>), 3.71 (3H, s, CH<sub>3</sub>), 3.77 (3H, s, CH<sub>3</sub>), 6.16 (1H, t, J=5.8, NH), 6.87 (2H, d, J=8.6, ArH), 6.9 (1H, dt, J=1.4, 7.3, ArH), 7.16 (2H, d, J=8.6, ArH), 7.33–7.48 (2H, m, ArH), 7.52 (1H, m, ArH), 7.63 (1H, s, ArH), 8.71 (1H, s, NH).

HPLC: retention time 6.62 min [Method B].

An important point that can be derived from the foregoing data is that by using a constitutively activated form of the receptor in the direct identification of candidate compounds, the selectivity of the compounds is exceptional: as those in the art appreciate, the homology between the human 5HT<sub>2A</sub> and 5HT<sub>2C</sub> receptors is about 95%, and even with such homology, certain of the directly identified compounds evidence a 4-order-of-magnitude (10,000-fold) selectivity separation (116100). This is important for pharmaceutical compositions in that such selectivity can help to reduce side-effects associated with interaction of a drug with a non-target receptor.

Different embodiments of the invention will consist of different constitutively activated receptors, different expression systems, different assays, and different compounds.

Those skilled in the art will understand which receptors to use with which expression systems and assay methods. All are considered within the scope of the teaching of this invention. In addition, those skilled in the art will recognize

that various modifications, additions, substitutions, and variations to the illustrative examples set forth herein can be made without departing from the spirit of the invention and are, therefore, considered within the scope of the invention.

---

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 33

<210> SEQ ID NO 1  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 1

gacctcgagg ttgcttaaga ctgaagc 27

<210> SEQ ID NO 2  
 <211> LENGTH: 27  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 2

atttctagac atatgtagct tgtaccg 27

<210> SEQ ID NO 3  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 3

ctaggggcac catgcaggct atcaacaatg aaagaaaagc taagaaagtc 50

<210> SEQ ID NO 4  
 <211> LENGTH: 50  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 4

caaggacttt cttagctttt ctttcattgt tgatagcctg catggtgcc 50

<210> SEQ ID NO 5  
 <211> LENGTH: 26  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 5

gacctcgagt ccttctacac ctcac 26

<210> SEQ ID NO 6  
 <211> LENGTH: 30  
 <212> TYPE: DNA



-continued

---

<213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence  
  
 <400> SEQUENCE: 6  
  
 tgctctagat tccagatagg tgaaaacttg 30

<210> SEQ ID NO 7  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence  
  
 <400> SEQUENCE: 7  
  
 caaagaaagt actgggcatc gtcttcttcc t 31

<210> SEQ ID NO 8  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence  
  
 <400> SEQUENCE: 8  
  
 ccgctcgagt actgcgccga caagctttga t 31

<210> SEQ ID NO 9  
 <211> LENGTH: 38  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence  
  
 <400> SEQUENCE: 9  
  
 cgatgccag cacttctgaa gcttttcttt cattgttg 38

<210> SEQ ID NO 10  
 <211> LENGTH: 36  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence  
  
 <400> SEQUENCE: 10  
  
 aaaagcttcg aaagtgctgg gcatcgtctt cttcct 36

<210> SEQ ID NO 11  
 <211> LENGTH: 30  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence  
  
 <400> SEQUENCE: 11  
  
 tgctctagat tccagatagg tgaaaacttg 30

<210> SEQ ID NO 12  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence

<400> SEQUENCE: 12  
 cgtgtctctc cttacttca 19

<210> SEQ ID NO 13  
 <211> LENGTH: 36  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence

<400> SEQUENCE: 13  
 tcggcgcagt actttgatag ttagaaagta ggtgat 36

<210> SEQ ID NO 14  
 <211> LENGTH: 38  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence

<400> SEQUENCE: 14  
 ttctaactat caaagtactg cgccgacaag ctttgatg 38

<210> SEQ ID NO 15  
 <211> LENGTH: 43  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence

<400> SEQUENCE: 15  
 ttcagcagtc aaccactag tctatactct gttcaacaaa att 43

<210> SEQ ID NO 16  
 <211> LENGTH: 28  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence

<400> SEQUENCE: 16  
 atttctagac atatgtagct tgtaccgt 28

<210> SEQ ID NO 17  
 <211> LENGTH: 19  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel  
 Sequence

<400> SEQUENCE: 17  
 atcacctact ttctaacta 19

<210> SEQ ID NO 18  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:

-continued

---

<223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 18

ccataatcgt caggggaatg aaaaatgaca caa 33

<210> SEQ ID NO 19  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 19

atttttcatt cccctgacga ttatggtgat tac 33

<210> SEQ ID NO 20  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 20

tgatgaagaa agggcaccac atgatcagaa aca 33

<210> SEQ ID NO 21  
 <211> LENGTH: 33  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 21

gatcatgtgg tgccctttct tcatcacaaa cat 33

<210> SEQ ID NO 22  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 22

gagacatatt atctgccacg gagg 24

<210> SEQ ID NO 23  
 <211> LENGTH: 24  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 23

ttggcataga aaccggaccc aagg 24

<210> SEQ ID NO 24  
 <211> LENGTH: 1416  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel

-continued

## Sequence

&lt;400&gt; SEQUENCE: 24

atggatattc tttgtgaaga aaatacttct ttgagctcaa ctacgaactc cctaatagcaa 60  
ttaaatagatg acaacaggct ctacagtaat gactttaact ccggagaagc taacacttct 120  
gatgcattta actggacagt cgactctgaa aatcgaacca acctttcctg tgaaggggtgc 180  
ctctcaccgt cgtgtctctc cttacttcat ctccaggaaa aaaactggtc tgctttactg 240  
acagccgtag tgattattct aactattgct ggaaacatac tcgtcatcat ggcagtgctc 300  
ctagagaaaa agctgcagaa tgccaccaac tatttcctga tgtcacttgc catagctgat 360  
atgctgctgg gtttccttgt catgcccgtg tccatgttaa ccatcctgta tgggtaccgg 420  
tggcctctgc cgagcaagct ttgtgcagtc tggatttacc tggacgtgct cttctccacg 480  
gcctccatca tgcacctctg cgccatctcg ctggaccgct acgtcgccat ccagaatccc 540  
atccaccaca gccgcttcaa ctccagaact aaggcatttc tgaaaatcat tgctgtttgg 600  
accatatcag taggtatata catgccaaata ccagtctttg ggctacagga cgattcgaag 660  
gtctttaagg aggggagttg cttactcgcc gatgataact ttgtcctgat cggctctttt 720  
gtgtcatttt tcattccctt aaccatcatg gtgatcacct actttctaac tatcaagtca 780  
ctccagaaag aagctacttt gtgtgtaagt gatcttggca cacgggcaa attagcttct 840  
ttcagcttcc tcctcagag ttctttgtct tcagaaaagc tcttcacgag gtcgatccat 900  
aggagccag ggtcctacac aggcaggagg actatgcagt ccatcagcaa tgagcaaaag 960  
gcatgcaagg tgctgggcat cgtcttcttc ctgtttgtgg tgatgtgggt ccctttcttc 1020  
atcacaaca tcatggccgt catctgcaaa gagtctgca atgaggatgt cattggggcc 1080  
ctgctcaatg tgtttgttg gatcggttat ctctcttcag cagtcaacc actagtctac 1140  
acactgttca acaagaccta taggtcagcc ttttcacggt atattcagtg tcagtacaag 1200  
gaaaacaaaa aaccattgca gttaatttta gtgaacacaa taccggcttt ggcctacaag 1260  
tctagccaac ttcaaaggg acaaaaaaag aattcaaagc aagatgcaa gacaacagat 1320  
aatgactgct caatggtgc tctaggaaag cagtattctg aagaggcttc taaagacaat 1380  
agcgacggag tgaatgaaaa ggtgagctgt gtgtga 1416

&lt;210&gt; SEQ ID NO 25

&lt;211&gt; LENGTH: 470

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

&lt;400&gt; SEQUENCE: 25

Met Asp Ile Leu Cys Glu Glu Asn Thr Ser Leu Ser Ser Thr Thr Asn  
1 5 10 15  
Ser Leu Met Gln Leu Asn Asp Asp Asn Arg Leu Tyr Ser Asn Asp Phe  
20 25 30  
Asn Ser Gly Glu Ala Asn Thr Ser Asp Ala Phe Asn Trp Thr Val Asp  
35 40 45  
Ser Glu Asn Arg Thr Asn Leu Ser Cys Glu Gly Cys Leu Ser Pro Ser  
50 55 60  
Cys Ser Leu Leu His Leu Gln Glu Lys Asn Trp Ser Ala Leu Leu Thr  
65 70 75 80  
Ala Val Val Ile Ile Leu Thr Ile Ala Gly Asn Ile Leu Val Ile Met

-continued

85					90					95					
Ala	Val	Ser	Leu	Glu	Lys	Lys	Leu	Gln	Asn	Ala	Thr	Asn	Tyr	Phe	Leu
			100					105					110		
Met	Ser	Leu	Ala	Ile	Ala	Asp	Met	Leu	Leu	Gly	Phe	Leu	Val	Met	Pro
		115					120					125			
Val	Ser	Met	Leu	Thr	Ile	Leu	Tyr	Gly	Tyr	Arg	Trp	Pro	Leu	Pro	Ser
		130					135					140			
Lys	Leu	Cys	Ala	Val	Trp	Ile	Tyr	Leu	Asp	Val	Leu	Phe	Ser	Thr	Ala
145							150					155			160
Ser	Ile	Met	His	Leu	Cys	Ala	Ile	Ser	Leu	Asp	Arg	Tyr	Val	Ala	Ile
				165								170			175
Gln	Asn	Pro	Ile	His	His	Ser	Arg	Phe	Asn	Ser	Arg	Thr	Lys	Ala	Phe
			180						185					190	
Leu	Lys	Ile	Ile	Ala	Val	Trp	Thr	Ile	Ser	Val	Gly	Ile	Ser	Met	Pro
		195					200					205			
Ile	Pro	Val	Phe	Gly	Leu	Gln	Asp	Asp	Ser	Lys	Val	Phe	Lys	Glu	Gly
		210					215					220			
Ser	Cys	Leu	Leu	Ala	Asp	Asp	Asn	Phe	Val	Leu	Ile	Gly	Ser	Phe	Val
225							230					235			240
Ser	Phe	Phe	Ile	Pro	Leu	Thr	Ile	Met	Val	Ile	Thr	Tyr	Phe	Leu	Thr
				245					250					255	
Ile	Lys	Ser	Leu	Gln	Lys	Glu	Ala	Thr	Leu	Cys	Val	Ser	Asp	Leu	Gly
			260						265					270	
Thr	Arg	Ala	Lys	Leu	Ala	Ser	Phe	Ser	Phe	Leu	Pro	Gln	Ser	Ser	Leu
		275							280					285	
Ser	Ser	Glu	Lys	Leu	Phe	Gln	Arg	Ser	Ile	His	Arg	Glu	Pro	Gly	Ser
		290					295					300			
Tyr	Thr	Gly	Arg	Arg	Thr	Met	Gln	Ser	Ile	Ser	Asn	Glu	Gln	Lys	Ala
305							310					315			320
Cys	Lys	Val	Leu	Gly	Ile	Val	Phe	Phe	Leu	Phe	Val	Val	Met	Trp	Cys
				325					330					335	
Pro	Phe	Phe	Ile	Thr	Asn	Ile	Met	Ala	Val	Ile	Cys	Lys	Glu	Ser	Cys
			340						345					350	
Asn	Glu	Asp	Val	Ile	Gly	Ala	Leu	Leu	Asn	Val	Phe	Val	Trp	Ile	Gly
		355					360						365		
Tyr	Leu	Ser	Ser	Ala	Val	Asn	Pro	Leu	Val	Tyr	Thr	Leu	Phe	Asn	Lys
		370					375					380			
Thr	Tyr	Arg	Ser	Ala	Phe	Ser	Arg	Tyr	Ile	Gln	Cys	Gln	Tyr	Lys	Glu
385												395			400
Asn	Lys	Lys	Pro	Leu	Gln	Leu	Ile	Leu	Val	Asn	Thr	Ile	Pro	Ala	Leu
				405					410					415	
Ala	Tyr	Lys	Ser	Ser	Gln	Leu	Gln	Met	Gly	Gln	Lys	Lys	Asn	Ser	Lys
			420						425					430	
Gln	Asp	Ala	Lys	Thr	Thr	Asp	Asn	Asp	Cys	Ser	Met	Val	Ala	Leu	Gly
		435							440					445	
Lys	Gln	Tyr	Ser	Glu	Glu	Ala	Ser	Lys	Asp	Asn	Ser	Asp	Gly	Val	Asn
		450					455					460			
Glu	Lys	Val	Ser	Cys	Val										
465															

&lt;210&gt; SEQ ID NO 26

&lt;211&gt; LENGTH: 1377

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

-continued

---

<220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 26

```

atggtgaacc tgaggaatgc ggtgcattca ttccttgtgc acctaattgg cctattggtt    60
tggcaatgtg atatttctgt gagccagta gcagctatag taactgacat tttcaatacc    120
tccgatggtg gacgcttcaa attcccagac ggggtacaaa actggccagc actttcaatc    180
gtcatcataa taatcatgac aataggtggc aacatccttg tgatcatggc agtaagcatg    240
gaaaagaaac tgcacaatgc caccaattac ttcttaatgt ccctagccat tgctgatatg    300
ctagtgggac tacttgtcat gccctgtct ctcctggcaa tcctttatga ttatgtctgg    360
ccactaccta gatatttgtg ccccgctggg atttcttttag atgttttatt ttcaacagcg    420
tccatcatgc acctctgcgc tatatcgctg gatcgggatg tagcaatacg taatcctatt    480
gagcatagcc gtttcaattc gcggactaag gccatcatga agattgctat tgtttgggca    540
atctctatag gtgtatcagt tcctatccct gtgattggac tgagggacga agaaaagggtg    600
ttcgtgaaca acacgacgtg cgtgctcaac gacccaaatt tcgttcttat tgggtccttc    660
gtagctttct tcataccgct gacgattatg gtgattacgt attgcctgac catctacgtt    720
ctgcgccgac aagctttgat gttactgcac ggccacaccg aggaaccgcc tggactaagt    780
ctggatttcc tgaagtgctg caagaggaat acggccgagg aagagaactc tgcaaaccct    840
aaccaagacc agaacgcacg ccgaagaaag aagaaggaga gacgtcctag gggcaccatg    900
caggctatca acaatgaaag aaaagcttcg aaagtccttg ggattgtttt ctttgtgttt    960
ctgatcatgt ggtgccatt tttcattacc aatattctgt ctgttctttg tgagaagtcc   1020
tgtaaccaa agctcatgga aaagcttctg aatgtgtttg tttggattgg ctatgtttgt   1080
tcaggaatca atcctctggt gtatctctgt ttcaacaaaa tttaccgaag ggcattctcc   1140
aactatttgc gttgcaatta taaggtagag aaaaagcctc ctgtcaggca gattccaaga   1200
gttgccgcca ctgctttgtc tgggagggag cttaatgtta acatttatcg gcataccaat   1260
gaaccgggta tcgagaaagc cagtgacaat gagcccggta tagagatgca agttgagaat   1320
ttagagttac cagtaaattc ctccagtgtg gtttagcgaag ggattagcag tgtgtga    1377

```

<210> SEQ ID NO 27  
 <211> LENGTH: 458  
 <212> TYPE: PRT  
 <213> ORGANISM: Artificial Sequence  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

<400> SEQUENCE: 27

```

Met Val Asn Leu Arg Asn Ala Val His Ser Phe Leu Val His Leu Ile
  1           5           10          15
Gly Leu Leu Val Trp Gln Cys Asp Ile Ser Val Ser Pro Val Ala Ala
  20          25          30
Ile Val Thr Asp Ile Phe Asn Thr Ser Asp Gly Gly Arg Phe Lys Phe
  35          40          45
Pro Asp Gly Val Gln Asn Trp Pro Ala Leu Ser Ile Val Ile Ile Ile
  50          55          60
Ile Met Thr Ile Gly Gly Asn Ile Leu Val Ile Met Ala Val Ser Met
  65          70          75          80
Glu Lys Lys Leu His Asn Ala Thr Asn Tyr Phe Leu Met Ser Leu Ala

```

-continued

85					90					95					
Ile	Ala	Asp	Met	Leu	Val	Gly	Leu	Leu	Val	Met	Pro	Leu	Ser	Leu	Leu
			100						105				110		
Ala	Ile	Leu	Tyr	Asp	Tyr	Val	Trp	Pro	Leu	Pro	Arg	Tyr	Leu	Cys	Pro
		115					120					125			
Val	Trp	Ile	Ser	Leu	Asp	Val	Leu	Phe	Ser	Thr	Ala	Ser	Ile	Met	His
		130				135					140				
Leu	Cys	Ala	Ile	Ser	Leu	Asp	Arg	Tyr	Val	Ala	Ile	Arg	Asn	Pro	Ile
145					150					155					160
Glu	His	Ser	Arg	Phe	Asn	Ser	Arg	Thr	Lys	Ala	Ile	Met	Lys	Ile	Ala
				165					170					175	
Ile	Val	Trp	Ala	Ile	Ser	Ile	Gly	Val	Ser	Val	Pro	Ile	Pro	Val	Ile
			180					185					190		
Gly	Leu	Arg	Asp	Glu	Glu	Lys	Val	Phe	Val	Asn	Asn	Thr	Thr	Cys	Val
		195					200					205			
Leu	Asn	Asp	Pro	Asn	Phe	Val	Leu	Ile	Gly	Ser	Phe	Val	Ala	Phe	Phe
		210				215					220				
Ile	Pro	Leu	Thr	Ile	Met	Val	Ile	Thr	Tyr	Cys	Leu	Thr	Ile	Tyr	Val
225					230					235					240
Leu	Arg	Arg	Gln	Ala	Leu	Met	Leu	Leu	His	Gly	His	Thr	Glu	Glu	Pro
				245					250					255	
Pro	Gly	Leu	Ser	Leu	Asp	Phe	Leu	Lys	Cys	Cys	Lys	Arg	Asn	Thr	Ala
			260					265					270		
Glu	Glu	Glu	Asn	Ser	Ala	Asn	Pro	Asn	Gln	Asp	Gln	Asn	Ala	Arg	Arg
		275					280					285			
Arg	Lys	Lys	Lys	Glu	Arg	Arg	Pro	Arg	Gly	Thr	Met	Gln	Ala	Ile	Asn
		290				295					300				
Asn	Glu	Arg	Lys	Ala	Ser	Lys	Val	Leu	Gly	Ile	Val	Phe	Phe	Val	Phe
305					310					315					320
Leu	Ile	Met	Trp	Cys	Pro	Phe	Phe	Ile	Thr	Asn	Ile	Leu	Ser	Val	Leu
				325					330					335	
Cys	Glu	Lys	Ser	Cys	Asn	Gln	Lys	Leu	Met	Glu	Lys	Leu	Leu	Asn	Val
			340					345						350	
Phe	Val	Trp	Ile	Gly	Tyr	Val	Cys	Ser	Gly	Ile	Asn	Pro	Leu	Val	Tyr
		355					360					365			
Thr	Leu	Phe	Asn	Lys	Ile	Tyr	Arg	Arg	Ala	Phe	Ser	Asn	Tyr	Leu	Arg
				370		375					380				
Cys	Asn	Tyr	Lys	Val	Glu	Lys	Lys	Pro	Pro	Val	Arg	Gln	Ile	Pro	Arg
385					390					395					400
Val	Ala	Ala	Thr	Ala	Leu	Ser	Gly	Arg	Glu	Leu	Asn	Val	Asn	Ile	Tyr
				405					410					415	
Arg	His	Thr	Asn	Glu	Pro	Val	Ile	Glu	Lys	Ala	Ser	Asp	Asn	Glu	Pro
				420				425					430		
Gly	Ile	Glu	Met	Gln	Val	Glu	Asn	Leu	Glu	Leu	Pro	Val	Asn	Pro	Ser
		435					440					445			
Ser	Val	Val	Ser	Glu	Arg	Ile	Ser	Ser	Val						
		450				455									

&lt;210&gt; SEQ ID NO 28

&lt;211&gt; LENGTH: 1377

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

-continued

&lt;400&gt; SEQUENCE: 28

```

atggtgaacc tgaggaatgc ggtgcattca ttccttgtgc acctaattgg cctattggtt    60
tggcaatgtg atatttctgt gagcccagta gcagctatag taactgacat tttcaatacc    120
tccgatggtg gacgcttcaa attcccagac ggggtacaaa actggccagc actttcaatc    180
gtcatcataa taatcatgac aataggtggc aacatccttg tgatcatggc agtaagcatg    240
gaaaagaaac tgcacaatgc caccaattac ttcttaatgt ccctagccat tgctgatatg    300
ctagtgggac tacttgtcat gccctgtct ctcttgcaa tcctttatga ttatgtctgg    360
ccactaccta gatatttgtg ccccgctctg atttctttag atgttttatt ttcaacagcg    420
tccatcatgc acctctgctg tatatcgctg gatcgggatg tagcaatacg taatcctatt    480
gagcatagcc gtttcaattc gcggactaag gccatcatga agattgctat tgtttgggca    540
atctctatag gtgtatcagt tcctatccct gtgattggac tgagggacga agaaaagggtg    600
ttcgtgaaca acacgacgtg cgtgctcaac gacccaaatt tcgttcttat tgggtccttc    660
gtagctttct tcataccgct gacgattatg gtgattacgt attgcctgac catctacgtt    720
ctgcgccgac aagctttgat gttactgcac ggccacaccg aggaaccgcc tggactaagt    780
ctggatttcc tgaagtgctg caagaggaat acggccgagg aagagaactc tgcaaaccct    840
aaccaagacc agaacgcacg ccgaagaaaag aagaaggaga gacgtcctag gggcaccatg    900
caggctatca acaatgaaag aaaagctaag aaagtccttg ggattgtttt ctttgtgttt    960
ctgatcatgt ggtgccatt tttcattacc aatattctgt ctgttctttg tgagaagtcc   1020
tgtaaccaa agctcatgga aaagcttctg aatgtgtttg tttggattgg ctatgtttgt   1080
tcaggaatca atcctctggt gtatactctg ttcaacaaaa tttaccgaag gcattctcc   1140
aactatttgc gttgcaatta taaggtagag aaaaagcctc ctgtcaggca gattccaaga   1200
gttgccgcca ctgctttgtc tgggaggagg cttaatgtta acatttatcg gcataccaat   1260
gaaccggtga tcgagaaaag cagtgacaat gagcccggta tagagatgca agttgagaat   1320
ttagagttac cagtaaatcc ctccagtgtg gttagcgaaa ggattagcag tgtgtga    1377

```

&lt;210&gt; SEQ ID NO 29

&lt;211&gt; LENGTH: 458

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

&lt;400&gt; SEQUENCE: 29

```

Met Val Asn Leu Arg Asn Ala Val His Ser Phe Leu Val His Leu Ile
  1             5             10             15
Gly Leu Leu Val Trp Gln Cys Asp Ile Ser Val Ser Pro Val Ala Ala
          20             25             30
Ile Val Thr Asp Ile Phe Asn Thr Ser Asp Gly Gly Arg Phe Lys Phe
          35             40             45
Pro Asp Gly Val Gln Asn Trp Pro Ala Leu Ser Ile Val Ile Ile Ile
          50             55             60
Ile Met Thr Ile Gly Gly Asn Ile Leu Val Ile Met Ala Val Ser Met
          65             70             75             80
Glu Lys Lys Leu His Asn Ala Thr Asn Tyr Phe Leu Met Ser Leu Ala
          85             90             95
Ile Ala Asp Met Leu Val Gly Leu Leu Val Met Pro Leu Ser Leu Leu

```



-continued

100					105					110					
Ala	Ile	Leu	Tyr	Asp	Tyr	Val	Trp	Pro	Leu	Pro	Arg	Tyr	Leu	Cys	Pro
		115					120					125			
Val	Trp	Ile	Ser	Leu	Asp	Val	Leu	Phe	Ser	Thr	Ala	Ser	Ile	Met	His
	130				135						140				
Leu	Cys	Ala	Ile	Ser	Leu	Asp	Arg	Tyr	Val	Ala	Ile	Arg	Asn	Pro	Ile
145					150					155					160
Glu	His	Ser	Arg	Phe	Asn	Ser	Arg	Thr	Lys	Ala	Ile	Met	Lys	Ile	Ala
				165					170					175	
Ile	Val	Trp	Ala	Ile	Ser	Ile	Gly	Val	Ser	Val	Pro	Ile	Pro	Val	Ile
			180					185					190		
Gly	Leu	Arg	Asp	Glu	Glu	Lys	Val	Phe	Val	Asn	Asn	Thr	Thr	Cys	Val
		195					200					205			
Leu	Asn	Asp	Pro	Asn	Phe	Val	Leu	Ile	Gly	Ser	Phe	Val	Ala	Phe	Phe
		210				215					220				
Ile	Pro	Leu	Thr	Ile	Met	Val	Ile	Thr	Tyr	Cys	Leu	Thr	Ile	Tyr	Val
225					230					235					240
Leu	Arg	Arg	Gln	Ala	Leu	Met	Leu	Leu	His	Gly	His	Thr	Glu	Glu	Pro
				245					250					255	
Pro	Gly	Leu	Ser	Leu	Asp	Phe	Leu	Lys	Cys	Cys	Lys	Arg	Asn	Thr	Ala
			260					265					270		
Glu	Glu	Glu	Asn	Ser	Ala	Asn	Pro	Asn	Gln	Asp	Gln	Asn	Ala	Arg	Arg
			275				280					285			
Arg	Lys	Lys	Lys	Glu	Arg	Arg	Pro	Arg	Gly	Thr	Met	Gln	Ala	Ile	Asn
	290					295					300				
Asn	Glu	Arg	Lys	Ala	Lys	Lys	Val	Leu	Gly	Ile	Val	Phe	Phe	Val	Phe
305					310					315					320
Leu	Ile	Met	Trp	Cys	Pro	Phe	Phe	Ile	Thr	Asn	Ile	Leu	Ser	Val	Leu
				325					330					335	
Cys	Glu	Lys	Ser	Cys	Asn	Gln	Lys	Leu	Met	Glu	Lys	Leu	Leu	Asn	Val
			340					345					350		
Phe	Val	Trp	Ile	Gly	Tyr	Val	Cys	Ser	Gly	Ile	Asn	Pro	Leu	Val	Tyr
		355					360					365			
Thr	Leu	Phe	Asn	Lys	Ile	Tyr	Arg	Arg	Ala	Phe	Ser	Asn	Tyr	Leu	Arg
					370		375				380				
Cys	Asn	Tyr	Lys	Val	Glu	Lys	Lys	Pro	Pro	Val	Arg	Gln	Ile	Pro	Arg
385					390					395					400
Val	Ala	Ala	Thr	Ala	Leu	Ser	Gly	Arg	Glu	Leu	Asn	Val	Asn	Ile	Tyr
				405					410					415	
Arg	His	Thr	Asn	Glu	Pro	Val	Ile	Glu	Lys	Ala	Ser	Asp	Asn	Glu	Pro
			420					425					430		
Gly	Ile	Glu	Met	Gln	Val	Glu	Asn	Leu	Glu	Leu	Pro	Val	Asn	Pro	Ser
		435					440					445			
Ser	Val	Val	Ser	Glu	Arg	Ile	Ser	Ser	Val						
		450				455									

&lt;210&gt; SEQ ID NO 30

&lt;211&gt; LENGTH: 1437

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

&lt;400&gt; SEQUENCE: 30

-continued

---

```

atggatattc tttgtgaaga aaatacttct ttgagctcaa ctacgaactc cctaattgcaa 60
ttaaattgatg acaacaggct ctacagtaat gactttaact ccggagaagc taacacttct 120
gatgcattta actggacagt cgactctgaa aatcgaacca acctttcctg tgaagggtgc 180
ctctcaccgt cgtgtctctc cttacttcat ctccaggaaa aaaactggtc tgctttactg 240
acagccgtag tgattattct aactattgct ggaaacatac tcgtcatcat ggcagtgtcc 300
ctagagaaaa agctgcagaa tgccaccaac tatttcctga tgtcacttgc catagctgat 360
atgctgctgg gtttccttgt catgcccgtg tccatgttaa ccatcctgta tgggtaccgg 420
tggcctctgc cgagcaagct ttgtgcagtc tggatttacc tggacgtgct cttctccacg 480
gcctccatca tgcacctctg cgccatctcg ctggaccgct acgtcgccat ccagaatccc 540
atccaccaca gccgcttcaa ctccagaact aaggcatttc tgaaaatcat tgctgtttgg 600
accatatcag taggtatata catgccataa ccagtctttg ggctacagga cgattcgaag 660
gtctttaagg aggggagttg cttactcgcc gatgataact ttgtcctgat cggctctttt 720
gtgtcatttt tcattccctt aaccatcatg gtgatcacct actttctaac tatcaaggtt 780
ctgcgccgac aagctttgat gttactgcac ggccacaccg aggaaccgcc tggactaagt 840
ctggatttcc tgaagtgctg caagaggaat acggccgagg aagagaactc tgcaaaccct 900
aaccaagacc agaacgcacg ccgaagaaag aagaaggaga gacgtcctag gggcaccatg 960
caggctatca acaatgaaag aaaagcttcg aaggtagtgg gcatcgtctt cttcctgttt 1020
gtggtgatgt ggtgcccttt cttcatcaca aacatcatgg ccgtcatctg caaagagtcc 1080
tgcaatgagg atgtcattgg ggccctgctc aatgtgtttg tttggatcgg ttatctctct 1140
tcagcagtca acccactagt ctatactctg ttcaacaaaa tttaccgaag ggcattctcc 1200
aactatttgc gttgcaatta taaggtagag aaaaagcctc ctgtcaggca gattccaaga 1260
gttgccgcca ctgctttgtc tgggagggag cttaatgtta acatttatcg gcataccaat 1320
gaaccggtga tcgagaaaag cagtgacaat gagcccggta tagagatgca agttgagaat 1380
ttagagttac cagtaaatcc ctccagtgtg gttagcgaaa ggattagcag tgtgtga 1437

```

&lt;210&gt; SEQ ID NO 31

&lt;211&gt; LENGTH: 478

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

&lt;400&gt; SEQUENCE: 31

```

Met Asp Ile Leu Cys Glu Glu Asn Thr Ser Leu Ser Ser Thr Thr Asn
 1             5             10             15
Ser Leu Met Gln Leu Asn Asp Asp Asn Arg Leu Tyr Ser Asn Asp Phe
          20             25             30
Asn Ser Gly Glu Ala Asn Thr Ser Asp Ala Phe Asn Trp Thr Val Asp
          35             40             45
Ser Glu Asn Arg Thr Asn Leu Ser Cys Glu Gly Cys Leu Ser Pro Ser
          50             55             60
Cys Leu Ser Leu Leu His Leu Gln Glu Lys Asn Trp Ser Ala Leu Leu
          65             70             75             80
Thr Ala Val Val Ile Ile Leu Thr Ile Ala Gly Asn Ile Leu Val Ile
          85             90             95
Met Ala Val Ser Leu Glu Lys Lys Leu Gln Asn Ala Thr Asn Tyr Phe
          100            105            110

```

-continued

---

Leu Met Ser Leu Ala Ile Ala Asp Met Leu Leu Gly Phe Leu Val Met  
           115                                  120                                  125

Pro Val Ser Met Leu Thr Ile Leu Tyr Gly Tyr Arg Trp Pro Leu Pro  
           130                                  135                                  140

Ser Lys Leu Cys Ala Val Trp Ile Tyr Leu Asp Val Leu Phe Ser Thr  
 145                                  150                                  155                                  160

Ala Ser Ile Met His Leu Cys Ala Ile Ser Leu Asp Arg Tyr Val Ala  
                                   165                                  170                                  175

Ile Gln Asn Pro Ile His His Ser Arg Phe Asn Ser Arg Thr Lys Ala  
                                   180                                  185                                  190

Phe Leu Lys Ile Ile Ala Val Trp Thr Ile Ser Val Gly Ile Ser Met  
           195                                  200                                  205

Pro Ile Pro Val Phe Gly Leu Gln Asp Asp Ser Lys Val Phe Lys Glu  
           210                                  215                                  220

Gly Ser Cys Leu Leu Ala Asp Asp Asn Phe Val Leu Ile Gly Ser Phe  
 225                                  230                                  235                                  240

Val Ser Phe Phe Ile Pro Leu Thr Ile Met Val Ile Thr Tyr Phe Leu  
                                   245                                  250                                  255

Thr Ile Lys Val Leu Arg Arg Gln Ala Leu Met Leu Leu His Gly His  
           260                                  265                                  270

Thr Glu Glu Pro Pro Gly Leu Ser Leu Asp Phe Leu Lys Cys Cys Lys  
           275                                  280                                  285

Arg Asn Thr Ala Glu Glu Glu Asn Ser Ala Asn Pro Asn Gln Asp Gln  
           290                                  295                                  300

Asn Ala Arg Arg Arg Lys Lys Lys Glu Arg Arg Pro Arg Gly Thr Met  
 305                                  310                                  315                                  320

Gln Ala Ile Asn Asn Glu Arg Lys Ala Ser Lys Val Leu Gly Ile Val  
                                   325                                  330                                  335

Phe Phe Leu Phe Val Val Met Trp Cys Pro Phe Phe Ile Thr Asn Ile  
           340                                  345                                  350

Met Ala Val Ile Cys Lys Glu Ser Cys Asn Glu Asp Val Ile Gly Ala  
           355                                  360                                  365

Leu Leu Asn Val Phe Val Trp Ile Gly Tyr Leu Ser Ser Ala Val Asn  
           370                                  375                                  380

Pro Leu Val Tyr Thr Leu Phe Asn Lys Ile Tyr Arg Arg Ala Phe Ser  
 385                                  390                                  395                                  400

Asn Tyr Leu Arg Cys Asn Tyr Lys Val Glu Lys Lys Pro Pro Val Arg  
                                   405                                  410                                  415

Gln Ile Pro Arg Val Ala Ala Thr Ala Leu Ser Gly Arg Glu Leu Asn  
           420                                  425                                  430

Val Asn Ile Tyr Arg His Thr Asn Glu Pro Val Ile Glu Lys Ala Ser  
           435                                  440                                  445

Asp Asn Glu Pro Gly Ile Glu Met Gln Val Glu Asn Leu Glu Leu Pro  
 450                                  455                                  460

Val Asn Pro Ser Ser Val Val Ser Glu Arg Ile Ser Ser Val  
 465                                  470                                  475

&lt;210&gt; SEQ ID NO 32

&lt;211&gt; LENGTH: 1437

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

-continued

&lt;400&gt; SEQUENCE: 32

```

atggatattc tttgtgaaga aaatacttct ttgagctcaa ctacgaactc cctaattgcaa      60
ttaaattgatg acaacaggct ctacagtaat gactttaact ccggagaagc taacacttct      120
gatgcattta actggacagt cgactctgaa aatcgaacca acctttcctg tgaaggggtgc      180
ctctcaccgt cgtgtctctc cttacttcat ctccaggaaa aaaactggtc tgctttactg      240
acagccgtag tgattattct aactattgct ggaacatac tcgtcatcat ggcagtgtcc      300
ctagagaaaa agctgcagaa tgccaccaac tatttcctga tgtcacttgc catagctgat      360
atgctgctgg gtttccttgt catgcccgtg tccatgttaa ccatcctgta tgggtaccgg      420
tggcctctgc cgagcaagct ttgtgcagtc tggatttacc tggacgtgct cttctccacg      480
gcctccatca tgcacctctg cgccatctcg ctggaccgct acgtcgccat ccagaatccc      540
atccaccaca gccgcttcaa ctccagaact aaggcatttc tgaaaatcat tgctgtttgg      600
accatatcag taggtatata catgccaata ccagtctttg ggctacagga cgattcgaag      660
gtctttaagg aggggagttg cttactcgcc gatgataact ttgtcctgat cggctctttt      720
gtgtcatttt tcattcccct gacgattatg gtgattacgt attgcctgac catctacgtt      780
ctgcgccgac aagctttgat gttactgcac ggccacaccg aggaaccgcc tggactaagt      840
ctggatttcc tgaagtgctg caagaggaat acggccgagg aagagaactc tgcaaaccct      900
aaccaagacc agaacgcacg ccgaagaaag aagaaggaga gacgtcctag gggcaccatg      960
caggctatca acaatgaaag aaaagctaag aaagtccttg ggattgtttt ctttgtgttt     1020
ctgatcatgt ggtgcccttt cttcatcaca aacatcatgg ccgtcatctg caaagagtcc     1080
tgcaatgagg atgtcattgg ggccctgctc aatgtgtttg tttggatcgg ttatctctct     1140
tcagcagtca acccactagt ctatactctg ttcaacaaaa tttaccgaag ggcattctcc     1200
aactatttgc gttgcaatta taaggtagag aaaaagcctc ctgtcaggca gattccaaga     1260
gttgccgcca ctgctttgtc tgggaggag cttaatgtta acatttatcg gcataccaat     1320
gaaccggtga tcgagaaagc cagtgacaat gagcccggta tagagatgca agttgagaat     1380
ttagagttac cagtaaatcc ctccagtgtg gtttagcгаа ggattagcag tgtgtga      1437

```

&lt;210&gt; SEQ ID NO 33

&lt;211&gt; LENGTH: 478

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Artificial Sequence

&lt;220&gt; FEATURE:

&lt;223&gt; OTHER INFORMATION: Description of Artificial Sequence: Novel Sequence

&lt;400&gt; SEQUENCE: 33

```

Met Asp Ile Leu Cys Glu Glu Asn Thr Ser Leu Ser Ser Thr Thr Asn
 1             5             10             15
Ser Leu Met Gln Leu Asn Asp Asp Asn Arg Leu Tyr Ser Asn Asp Phe
 20             25             30
Asn Ser Gly Glu Ala Asn Thr Ser Asp Ala Phe Asn Trp Thr Val Asp
 35             40             45
Ser Glu Asn Arg Thr Asn Leu Ser Cys Glu Gly Cys Leu Ser Pro Ser
 50             55             60
Cys Leu Ser Leu Leu His Leu Gln Glu Lys Asn Trp Ser Ala Leu Leu
 65             70             75             80
Thr Ala Val Val Ile Ile Leu Thr Ile Ala Gly Asn Ile Leu Val Ile
 85             90             95

```

-continued

---

Met	Ala	Val	Ser	Leu	Glu	Lys	Lys	Leu	Gln	Asn	Ala	Thr	Asn	Tyr	Phe
			100					105					110		
Leu	Met	Ser	Leu	Ala	Ile	Ala	Asp	Met	Leu	Leu	Gly	Phe	Leu	Val	Met
		115					120					125			
Pro	Val	Ser	Met	Leu	Thr	Ile	Leu	Tyr	Gly	Tyr	Arg	Trp	Pro	Leu	Pro
	130					135					140				
Ser	Lys	Leu	Cys	Ala	Val	Trp	Ile	Tyr	Leu	Asp	Val	Leu	Phe	Ser	Thr
145					150					155					160
Ala	Ser	Ile	Met	His	Leu	Cys	Ala	Ile	Ser	Leu	Asp	Arg	Tyr	Val	Ala
				165					170					175	
Ile	Gln	Asn	Pro	Ile	His	His	Ser	Arg	Phe	Asn	Ser	Arg	Thr	Lys	Ala
			180					185					190		
Phe	Leu	Lys	Ile	Ile	Ala	Val	Trp	Thr	Ile	Ser	Val	Gly	Ile	Ser	Met
		195					200					205			
Pro	Ile	Pro	Val	Phe	Gly	Leu	Gln	Asp	Asp	Ser	Lys	Val	Phe	Lys	Glu
	210					215					220				
Gly	Ser	Cys	Leu	Leu	Ala	Asp	Asp	Asn	Phe	Val	Leu	Ile	Gly	Ser	Phe
225					230					235					240
Val	Ser	Phe	Phe	Ile	Pro	Leu	Thr	Ile	Met	Val	Ile	Thr	Tyr	Cys	Leu
				245					250					255	
Thr	Ile	Tyr	Val	Leu	Arg	Arg	Gln	Ala	Leu	Met	Leu	Leu	His	Gly	His
			260					265					270		
Thr	Glu	Glu	Pro	Pro	Gly	Leu	Ser	Leu	Asp	Phe	Leu	Lys	Cys	Cys	Lys
			275				280					285			
Arg	Asn	Thr	Ala	Glu	Glu	Glu	Asn	Ser	Ala	Asn	Pro	Asn	Gln	Asp	Gln
	290					295					300				
Asn	Ala	Arg	Arg	Arg	Lys	Lys	Lys	Glu	Arg	Arg	Pro	Arg	Gly	Thr	Met
305					310					315					320
Gln	Ala	Ile	Asn	Asn	Glu	Arg	Lys	Ala	Lys	Lys	Val	Leu	Gly	Ile	Val
				325					330					335	
Phe	Phe	Val	Phe	Leu	Ile	Met	Trp	Cys	Pro	Phe	Phe	Ile	Thr	Asn	Ile
			340					345					350		
Met	Ala	Val	Ile	Cys	Lys	Glu	Ser	Cys	Asn	Glu	Asp	Val	Ile	Gly	Ala
		355					360					365			
Leu	Leu	Asn	Val	Phe	Val	Trp	Ile	Gly	Tyr	Leu	Ser	Ser	Ala	Val	Asn
		370				375					380				
Pro	Leu	Val	Tyr	Thr	Leu	Phe	Asn	Lys	Ile	Tyr	Arg	Arg	Ala	Phe	Ser
385					390					395					400
Asn	Tyr	Leu	Arg	Cys	Asn	Tyr	Lys	Val	Glu	Lys	Lys	Pro	Pro	Val	Arg
				405					410					415	
Gln	Ile	Pro	Arg	Val	Ala	Ala	Thr	Ala	Leu	Ser	Gly	Arg	Glu	Leu	Asn
			420					425					430		
Val	Asn	Ile	Tyr	Arg	His	Thr	Asn	Glu	Pro	Val	Ile	Glu	Lys	Ala	Ser
		435					440					445			
Asp	Asn	Glu	Pro	Gly	Ile	Glu	Met	Gln	Val	Glu	Asn	Leu	Glu	Leu	Pro
	450					455					460				
Val	Asn	Pro	Ser	Ser	Val	Val	Ser	Glu	Arg	Ile	Ser	Ser	Val		
465					470					475					

---

We claim:

1. A constitutively active non-endogenous human 5HT<sub>2A</sub> serotonin receptor encoded by the cDNA of SEQ ID NO. 30 comprising SEQ ID NO. 31. 65

\* \* \* \* \*