



US006495317B1

(12) **United States Patent**
Loescher et al.

(10) **Patent No.:** **US 6,495,317 B1**
(45) **Date of Patent:** **Dec. 17, 2002**

(54) **DNA ENCODING MANNANOSE 6-PHOSPHATE
REDUCTASE AND RECOMBINANTS
PRODUCED THEREFROM**

(75) Inventors: **Wayne H. Loescher**, Okemos, MI
(US); **John D. Everard**, East Lansing,
MI (US); **Rebecca Grumet**, East
Lansing, MI (US)

(73) Assignee: **Board of Trustees of Michigan State
University**, East Lansing, MI (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **09/166,438**

(22) Filed: **Oct. 5, 1998**

Related U.S. Application Data

(62) Division of application No. 08/731,320, filed on Oct. 15,
1996, now Pat. No. 6,416,985.

(51) **Int. Cl.**⁷ **C12Q 1/68**

(52) **U.S. Cl.** **435/6; 536/23.2**

(58) **Field of Search** **435/6, 189, 190;**
536/23.2, 23.6

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,128,256 A 7/1992 Huse et al. 435/172.3
5,268,288 A 12/1993 Pharr 435/190
5,492,820 A 2/1996 Sonnewald et al. 435/69.1

OTHER PUBLICATIONS

Everard et al. "Cloning of mannose 6-phosphate reductase
from celery" *HortScience*, vol. 30 (4), p. 898, abstract number
829, Jul. 1995.*

Everard et al. "Molecular cloning of mannose-6-phosphate
reductase and its developmental expression in celery" *Plant
Physiol.* 113, 1427-1435, 1997.*

Bieleski, R.L., Sugar alcohols. In: F.A. Loewus and W.
Tanner, eds., *Plant Carbohydrates I. Intracellular Car. Encyc.*
Plant. Physiol. vol. 13A, New Series, Springer-Verlag, NY,
pp. 158-192 (1982).

Bohnert, Hans J., et al., *The Plant Cell*, 7:1099-1111 (1995).

Ahmad, I., et al., *New Phytol.* 82:671-679 (1979).

Hirai, M., *Plant Cell Physiol.* 24:925-931 (1983).

Everard et al., *Plant Physiol.* 102:345-356 (1993).

Stoop, J.M.H., et al., *Plant Physiol.* 106:503-511 (1994).

Kanayama, Y., et al., *Plant Physiology* 100:1607-1608
(1992).

Williamson, J.S., et al., *Proc. Natl. Acad. Sci* 92:7148-7152
(1995).

Tarczynski, M.C., et al., *Science* 259:508-510 (1993).

Thomas, J. C., et al., *Plant Cell and Environ.* 18:801-806
(1995).

Tao, R., et al., *Plant Cell Physiol* 36:525-532 (1995).

Hunkerpillar et al., *Methods in Enzymology* 91:227-236
(1983).

Pharr, D.M., et al., *Plant Physiol.* 180-194 (1995).

Loescher, W.H., et al., *Plant Physiol.* 170-178 (1995).

Bartels, D., et al., *EMBO J* 10:1037-1043 (1991).

Rumpho, M.E., et al., *Plant Physiol.* 73:869-873 (1983).

Harloff, H.J., et al., *J. Plant Physiol.* 141:513-520 (1993).

Loescher, W.H., et al., *Plant Physiol.* 98: 1396-1402 (1992).

Gilmour, S.J., et al., *Plant Physiol.* 87:745-750 (1988).

Hondred, D., et al., *Plant Mol. Biol.*, 9:259-275 (1987).

Ried, J.K., et al., *BioTechniques* 12:660-666 (1992).

Bradford, M.M., *Anal. Biochem.* 136:248-254 (1976).

Bohren et al., *J. Biol. Chem.* 264:9547-9551 (1989).

Borhani, D.W., et al., *J. Biol. Chem.* 267:24841-24847
(1992).

Morijana et al., *FASEB J* 46:1330 (Abstract) (1987).

Schade et al., *J.B.C.* 265(7) : 3628-3635 (1990).

Petrash et al., *JBC* 267 (34) : 24833-24840 (1992).

Anderson et al., *Planta* 196:118-124 (1995).

Laemmli, U.K., *Nature* vol. 227 680-685 (1970).

Loescher, W.H., *Physiol. Plantarum* 70:553-557 (1987).

Fox, T.C., et al *Plant Physiology* 82:307-311 (1986).

Flora and Madore Planta 189:484-490 (1993).

Loescher, W.H. and Everard, J.D., *Photoassimilate Distri-
bution in Plants and Crops*, 185-207 (1996).

Davis, J.M., et al., *Plant Physiol*, 86:129-133 (1988).

Everard, J.D., et al., *Plant Physiol.*, 106:281-292 (1994).

Escobar-Gutierrez, A., et al., *Plant Physiol. Supp* 105:
Abstract 575 (1994).

Quick, W.P. and Schaffer, A.A., Sucrose metabolism in
sources and sinks. In: *Photoassimilate distribution in plants
and crops: source-sink relationships*. Zamski, E., and Schaf-
fer A., (eds) Marcel Dekker, Inc.: pp. 115-156 (1996).

Preiss, J. and Sivak, M.N., Starch synthesis in sinks and
sources. In: *Photoassimilate distribution in plants and crops:
source-sink relationships*. Zamski, E., and Schaffer A.,
(eds), Marcel Dekker, Inc.: pp. 63-96 (1996).

* cited by examiner

Primary Examiner—Nashaat T. Nashed

(74) *Attorney, Agent, or Firm*—Ian C. McLeod

(57) **ABSTRACT**

DNA encoding mannose 6-phosphate reductase (M6PR) and
the use of the DNA in vectors and bacteria and in plants. The
enzyme enables the production of mannitol in plants which
increases stress tolerance, particularly to salt.

5 Claims, 7 Drawing Sheets




FIG. 1




FIG. 2A




FIG. 2B




FIG. 2C




FIG. 3A

TATGATGTTACTTGACCAGAGATATCTTGTTCATATCCCAAGATCAAGCCCTGCTGTAAATCAGATCGAGACGCCCTTACTTCCAAAGAGATTCTCTGATCAAAATCTGTCCAGAAATGATGGCATTGCTATCACAGCACA 700

M6PR ORF

Tyr Asp Val Tyr Leu Thr Arg Asp Ile Leu Ser Tyr Ser Lys Ile Lys Pro Ala Val Asn Gln Ile Glu Thr His Pro Tyr Phe Gln Arg Asp Ser Leu Ile Lys Phe Cys Gln Lys Tyr Gly Ile Ala Ile Thr Ala His

CACACCACTAGCGGGCCATTGGCTAACTAGCGGATTGGATCAGTTTGGTGCTTAGATGATCCAGTTCTTAAGAAAATTATCTGACAAACACAAAGTCACCAGCTCAGATTGTTCTCCGTTGGGGTGTGCAGCGCA 840

M6PR ORF

Thr Pro Leu Gly Ala Leu Ala Asn Thr Glu Arg Phe Gly Ser Val Ser Cys Leu Asp Asp Pro Val Leu Lys Lys Ser Asp Lys His Asn Lys Ser Pro Ala Gln Ile Val Leu Arg Trp Gly Val Gln Arg

ACACAAATTGTAATCCCAAGTCATCGAAAACATAAAGACTCGAGGAAAACATCAACATTTTTGACTTTGAGTTGAGCAAGGAAGATATGGAGCTCATCAAAACAATGGAGCGCAACCAAGGAGTAACACACCTGCTAAA 980

M6PR ORF

Aldoketo Reductase 3

Xho I

Asn Thr Ile Val Ile Pro Lys Ser Ser Lys Thr Lys Arg Leu Glu Glu Asn Ile Asn Ile Phe Asp Phe Glu Leu Ser Lys Glu Asp Met Glu Leu Ile Lys Thr Met Glu Arg Asn Thr Pro Ala Lys

GCTTGGGGAATAGATGTTTATGCTTGGCATAACACATCTTCACTGTATTTTTATCATTGTTATCCACAATTCAGAGTGGTTGCTATTTTACTTGTGTGGAGGGAATGTGTGTTGAGTTGTTGTAG 1120

M6PR ORF

Ala Trp Gly Ile Asp Val Tyr Ala Trp His Asn Thr Phe Thr Val Phe Leu Ser Leu Leu Phe His Asn Ser Glu Trp Leu Ser Phe Leu Leu Ala Ile Val Cys Gly Glu Cys Val Leu Ser Cys Cys Ser

TAATTGTACAAGGCATAAAGCCTTTAAATAACCCATCATATGTAAATGGGAAAATGCCATGATTTGGTCAAAAAAAAAAAAAAAAAAAAAA 1207

Asn Cys Thr Arg His Lys Ala Phe Lys Pro Ile Ile Cys Lys Trp Glu Met Pro Phe Gly Gln Lys Lys Lys Asn

FIG. 3B

M . . . V T L S S G . . . M P V L G L G V W R L D K . E L K . V Functional similarity

1 M S T V T L S S G Y E M P V I G L G L W R L E K D E L K E V A6PR protein aa

1 M A - I T L N S G F K M P V L G L G V W R M D R N E I K N L M6PR ORF 29 Dec 95

L L S A I . L G Y R H F D C A A . Y K S E A D V G E A L . E Functional similarity

31 I L N A I K I G Y R H F D C A A H Y K S E A D V G E A L A E A6PR protein aa

30 L L S A I N L G Y R H F D C A A D Y K N E L E V G E A F K E M6PR ORF 29 Dec 95

A F . T . L V K R E D L F I T T K L W N S D H G H V V E A C Functional similarity

61 A F K T G L V K R E E L F I T T K I W N S D H G H V V E A C A6PR protein aa

60 A F D T D L V K R E D L F I T T K L W N S D H G H V I E A C M6PR ORF 29 Dec 95

K N S L . K L Q L D Y L D L Y L V H . P M A S K H S . I G . Functional similarity

91 K N S L E K L Q I D Y L D L Y L V H Y P M P T K H N A I G K A6PR protein aa

90 K N S L K K L Q L E Y L D L Y L I H F P M A S K H S G I G T M6PR ORF 29 Dec 95

T . S L L . D D . V L D V D A T I S L . . T W . . M E K . V Functional similarity

121 T A S L L G E D K V L D I D V T I S L Q Q T W E G M E K T V A6PR protein aa

120 T R S I L D D E G V W E V D A T I S L E A T W H E M E K L V M6PR ORF 29 Dec 95

. L G L V R S I G L S N Y D V . L T R D . L . Y S K I K P A Functional similarity

151 S L G L V R S I G L S N Y E L F L T R D C L A Y S K I K P A A6PR protein aa

150 E M G L V R S I G I S N Y D V Y L T R D I L S Y S K I K P A M6PR ORF 29 Dec 95

V S Q I E T H P Y F Q R D S L V K F C . K . G V A I T A H T Functional similarity

181 V S Q F E T H P Y F Q R D S L V K F C M K H G V L P T A H T A6PR protein aa

180 V N Q I E T H P Y F Q R D S L I K F C Q K Y G I A I T A H T M6PR ORF 29 Dec 95

FIG. 4A

P L G G A A N . D . F G S V S . L D D P V L . . V . . K . Functional similarity

211 P L G G A A N K D M F G S V S P L D D P V L N D V A K K Y A6PR protein aa
210 P L G G A L A N T E R F G S V S C L D D P V L K K L S D K H M6PR ORF 29 Dec 95

G K S V A Q I . L R W G V Q R . T A V I P K S S K . . R L . Functional similarity

241 G K S V A Q I C L R W G I Q R K T A V I P K S S K I Q R L K A6PR protein aa
240 N K S P A Q I V L R W G V Q R N T I V I P K S S K T K R L E M6PR ORF 29 Dec 95

E N L . V L D F . L S . E D M . L I . S I D R . Q R S S . P Functional similarity

271 E N L E V L E F Q L S D E D M Q L I Y S I D R K Y R T S L P A6PR protein aa
270 E N I N I F D F E L S K E D M E L I K T M E R N Q R S N T P M6PR ORF 29 Dec 95

. K . W G L D V Y A . Functional similarity

301 S K T W G L D V Y A A6PR protein aa
300 A K A W G I D V Y A . M6PR ORF 29 Dec 95

Consensus 'Functional similarity': When all match the residue group of the
Consensus show the residue of the Consensus, otherwise show '.'. Residue
Groupings of Functional similarity are: a=(DE), b=(HKR), f=(AFILMPVW),
p=(CGNQSTY).

Decoration 'Decoration #1': Box residues that match M6PR ORF 29 Dec 95 exactly.

FIG. 4B




FIG. 5A




FIG. 5B

**DNA ENCODING MANNOSE 6-PHOSPHATE
REDUCTASE AND RECOMBINANTS
PRODUCED THEREFROM**

This application is a divisional of copending application Ser. No. 08/731,320 filed on Oct. 15, 1996, now U.S. Pat. No. 6,416,985.

GOVERNMENT RIGHTS

This invention was developed under U.S.D.A. Contract No. 93-37100-8907 and 94-01439. The U.S. Government has certain rights in this invention.

BACKGROUND OF THE INVENTION

(1) Field of the Invention

The present invention relates to a DNA encoding mannose 6-phosphate reductase (M6PR) which is part of the pathway which forms mannitol in plants. The DNA encoding M6PR was isolated from celery. When the DNA is transformed into a plant the resulting transformed plant can be more tolerant to environmental stresses in the form of dehydration, salinity and drought.

(2) Description of Related Art

In many plants, sucrose and starch are the primary products of photosynthetic carbon assimilation. In other species, however, acrylic polyols (e.g., sorbitol, mannitol) can also be primary products and account for between 15 and 60% of the assimilated carbon, depending on the species (Loescher, W. H., *Physiol. Plantarum* 70:553-557 (1987); Fox, T. C., et al *Plant Physiology* 82:307-311 (1986); Flora and Madore *Planta* 189:484-490 (1993); Loescher, W. H. and Everard, J. D., *Photoassimilate Distribution in Plants and Crops*, 185-207 (1996)), the stage of leaf development, (Davis, J. M., et al., *Plant Physiol*, 86:129-133 (1988) and environmental factors (e.g., salinity (Everard, J. D., et al., *Plant Physiol*, 106:281-292 (1994)) and water stress; Escobar-Gutierrez, A., et al., *Plant Physiol suppl* 105: abstract 575 (1994)). The influence of developmental and environmental factors suggest that the partitioning of photoassimilates between sugar alcohols, sucrose and starch is under strict metabolic control. This is consistent with the complexity and diversity of the control mechanisms known to govern sucrose and starch synthesis in species that do not synthesize sugar alcohols (Quick, W. P. and Schaffer, A. A., *Sucrose metabolism in sources and sinks*. In: *Photoassimilate distribution in plants and crops: source-sink relationships*. Zamski, E., and Schaffer A., (eds), Marcel Dekker, Inc.: pp 115-156 (1996); and Preiss, J. and Sivak, M. N., *Starch synthesis in sinks and sources*. In: *Photoassimilate distribution in plants and crops: source-sink relationships*. Zamski, E., and Schaffer A., (eds), Marcel Dekker, Inc.: pp 63-96 (1996)). There is, however, almost no equivalent information on the mechanisms by which polyol metabolism is regulated or integrated with these other pathways in sugar alcohol synthesizing species. Such mechanisms are of more than just esoteric interest since: (a) an estimated 30% of global annual carbon assimilation results in polyol production (Bialeski, R. L., *Sugar alcohols*. In: F. A. Loewus and W. Tanner, eds., *Plant Carbohydrates I. Intracellular Carbohydrates*, Encyc. *Plant Physiol*. Vol. 13A, New Series, Springer-Verlag, NY, pp. 158-192 (1982)), (b) many polyol producing species are of considerable economic value, and (c) substantial correlative evidence shows that polyols accumulate in higher plants subjected to stresses mediated at the cellular level by changes in water activity, suggesting a role in stress tolerance (Bohnert, Hans J., et al., *The Plant Cell*,

7:1099-1111 (1995); Ahmad, I., et al., *New Phytol* 82:671-679 (1979); Hirai, M., *Plant Cell Physiol*. 24:925-931 (1983); Everard, J. D., et al., *Plant Physiol* 106:281-292 (1994); Stoop, J. M. H., et al., *Plant Physiol*. 106:503-511 (1994)).

In recent years major advances in the understanding of sugar alcohol metabolism in higher plants have been facilitated by the detection, characterization, and purification of several key enzymes (see Loescher and Everard, *Photoassimilate Distribution in Plants and Crops*, 185-207 (1996) for a review). The successful cloning of genes for NADP-dependent sorbitol 6-phosphate reductase (Kanayama, Y., et al., *Plant Physiology* 100:1607-1608 (1992)), mannose 1-oxidoreductase (Williamson, J. D., et al., *Proc. Natl. Acad. Sci.* 92:7148-7152 (1995)) and the introduction of heterologous genes, which confer sugar alcohol synthesis to plants that normally do not produce them (Tarczynski, M. C., et al., *Science* 259:508-510 (1993); Thomas, J. C., et al., *Plant Cell and Environ* 18:801-806 (1995); Tao, R., et al., *Plant Cell Physiol* 36:525-532 (1995)), now provide powerful tools with which to study sugar alcohol biochemistry and physiology. For example, Tarczynski, M. C., et al. (*Science* 259:508-510 (1993)), and Thomas, J. C., et al. (*Plant Cell and Environ* 18:801-806 (1995)) have recently shown that transgenic tobacco and Arabidopsis plants expressing the bacterial mannitol dehydrogenase (mtlD) gene not only produce low levels of mannitol, but also exhibit enhanced sodium chloride tolerance. In another study the inhibitory effects of 300 mM NaCl on the growth of celery suspension cultures were substantially reduced when mannitol, rather than sucrose, was included as the sole carbon source (Pharr, D. M., et al., *Plant Physiology* 180-194 (1995)). Such studies convincingly demonstrate that polyols confer some stress protection but give little insight into the underlying mechanisms. Mechanisms are beginning to be elucidated however, mannitol accumulation in salt stressed celery plants has been associated with a down regulation (at both the mRNA and protein levels) of mannitol dehydrogenase (MTD), a key catabolic enzyme (Williamson, J. D., et al., *Proc. Natl. Acad. Sci.*, 92:7148-7152 (1995)), although, enhanced de novo synthesis is also undoubtedly involved in this acclimation response (Everard, J. D., *Plant Physiol*. 106:281-292 (1994); Loescher, W. H., et al., *Plant Physiology*, 170-178 (1995)). Other evidence showing a possible role for polyols in stress metabolism in higher plants includes the identification of an aldose reductase in *Craterostigma* leaves and barley embryos that accumulates when these tissues undergo desiccation (Bartels, D., et al., *EMBO J* 10:1037-1043 (1991)).

A pathway for mannitol synthesis in higher plants has been established in celery (Rumpho, M. E., et al., *Plant Physiol*. 73:869-873 (1983)) and appears to be present in other mannitol synthesizing species (Harloff, H. J., et al., *J. Plant Physiol* 141:513-520 (1993)). Biosynthesis involves three unique enzymatic steps consisting of an isomerization (F6P to mannose 6-P, mediated by mannose 6-P isomerase), a reduction (mannose 6-P to mannitol 1-P by mannose 6-P reductase (M6PR)) and a dephosphorylation (mannitol 1-P to mannitol, by mannitol 1-P phosphatase). Radiotracer studies and kinetic analyses indicate that M6PR plays a regulatory role in this pathway. This enzyme has been purified and partially characterized (Loescher, W. H., et al., *Plant Physiol*. 98:1396-1402 (1992)).

U.S. Pat. No. 5,268,288 to Pharr describes mannitol oxidoreductase protein. The enzyme converts mannitol to mannose in plants. It is thus different from the present invention which relates to mannitol production. The patent

describes various recombinant techniques useful in the present invention. U.S. Pat. No. 5,492,820 to Sonnewald et al describes plasmids (vectors) for producing recombinant plants with altered sugar expression. The disclosure of such vectors is incorporated into the disclosure of the present application as well.

OBJECTS

It is therefore an object of the present invention to provide DNA encoding mannose 6-phosphate reductase (M6PR) which is useful in producing plasmids and transgenic plants with increased tolerance to environmental stresses, particularly salinity. These and other objects will become increasingly apparent by reference to the following description and the drawings.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a photograph of an electrophoresis gel showing in vitro translation of poly(A)+RNA isolated from celery leaf tissue. Lanes 1 and 2 are Coomassie Brilliant blue R250 stained lanes from a 12.5% polyacrylamide gel showing Bio-Rad (Richmond, Calif.) molecular weight markers and purified celery leaf M6PR (0.1 μ g), respectively. Lanes 3 and 4 show an autoradiograph of a subsample of the in vitro translation products represented from leaves 5 and 7 after immunoprecipitation with M6PR-specific antisera; all lanes were from the same gel.

FIGS. 2A, 2B and 2C show verification of M6PR specific clones. FIG. 2A is a graph showing M6PR activities in sonicated extracts of one nonspecific and three putative M6PR clones with and without IPTG induction. FIG. 2B is a photograph of an SDS gel of the extracts (100 μ g/lane); the extreme right-hand lane contains 3 μ g of authentic celery leaf M6PR. FIG. 2C is a photograph of a Western blot; an SDS gel containing 5 μ g total protein per lane was blotted to PVDF membrane and probed with M6PR-specific antisera, as in Panel C the extreme left-hand lane contains authentic leaf enzyme.

FIGS. 3A and 3B are drawings showing the nucleic and translated amino acid sequence of celery cDNA clone D of SEQ ID NO:1 and SEQ ID NO:2, respectively. The open reading frame (M6PR ORF) coded for a peptide of 35.2 kD and had three domains which were identified, through computer data base searches, as being typical of the aldoketo reductase family. Also shown is the peptide resulting from tryptic digestion of authentic celery leaf M6PR and the Xho 1 restriction site within the coding region. Two other independent clones were sequenced on both strands and only differed from the displayed sequence in the lengths of their 3' and 5' non-coding regions.

FIGS. 4A and 4B are drawings showing an amino acid sequence comparison of M6PR of SEQ ID NO:2 and NADP-sorbitol-6-phosphate dehydrogenase from apple SEQ ID NO:4. Sequences were 64% identical (shown by enclosed areas) and showed 84% similarity, if the functional relatedness of the residues was considered. For the latter comparison the following groups were used: acidic (D,E), basic (H,K,R), hydrophobic (A,F,I,L,M,P,V,W) and polar (C,G,N,Q,S,T,Y); "." indicates lack of functional similarity.

FIG. 5A is a photograph of SDS-PAGE gel (12.5% acrylamide) of samples collected after the various purification steps (See Table 2). Lanes: 1 and 7, Bio-Rad molecular mass markers; 2, 27,500 g supernatant of disrupted cells (80 μ g protein); 3, 30 to 60% acetone fraction (120 μ g); 4, post gel-filtration chromatography on Sephacryl S-200 (30 μ g); 5, post affinity chromatography on Reactive Yellow 86

eluted with 0.1 mM NADPH (15 μ g); 6, product of a second pass over the RY86 column eluted with a linear gradient (0 to 0.2 mM) of NADPH (2 μ g). 8, authentic celery leaf M6PR (0.5 μ g).

FIG. 5B is a photograph of a Western blot of proteins from selected purification steps (see above), probed with M6PR-specific antisera. Lanes: 1, 6 and 7 authentic celery leaf M6PR; 2, crude supernatant; 3, post S200 gel filtration; 4 and 5, after 1 and 2 passes over RY86 affinity column, respectively.

DESCRIPTION OF PREFERRED EMBODIMENTS

The present invention relates to a DNA encoding mannose-6-phosphate reductase (M6PR) free of any other DNA as set forth in SEQ ID NO:1.

The present invention also relates to a DNA encoding mannose-6-phosphate reductase (M6PR) in a plasmid in an *Escherichia coli* as contained in a deposit identified as ATCC 98041.

The present invention also relates to a recombinant plasmid containing DNA encoding mannose-6-phosphate reductase as set forth in SEQ ID NO:1.

The present invention also relates to a recombinant plasmid containing DNA encoding mannose-6-phosphate reductase in a plasmid in *E. coli* as contained in a deposit identified as ATCC 98041.

The present invention also relates to a bacterium containing a recombinant plasmid containing DNA encoding mannose-6-phosphate as set forth in SEQ ID NO:1.

The present invention also relates to a bacterium containing a recombinant plasmid containing DNA encoding mannose 6-phosphate in the plasmid in an *E. coli* as contained in a deposit identified as ATCC 98041.

The present invention also relates to a method for detecting unknown DNA encoding mannose-6-phosphate reductase (M6PR) which comprises probing the unknown DNA with a probe DNA encoding a unique region of the M6PR as set forth in SEQ ID NO:1.

The present invention also relates to a method for detecting an unknown DNA encoding mannose-6-phosphate reductase (M6PR) which comprises probing the unknown DNA with a probe DNA encoding a unique region of the M6PR in a plasmid in *E. coli* as contained in a deposit identified as ATCC 98041.

The present invention also relates to a transgenic plant containing recombinant DNA encoding mannose-6-phosphate reductase (M6PR) based upon SEQ ID NO:1. The antisense of SEQ ID NO:1 is used.

The present invention also relates to a transgenic plant containing recombinant DNA encoding mannose-6-phosphate reductase (M6PR) in a plasmid in an *E. coli* contained in a deposit identified as ATCC 98041.

The present invention also relates to a method for detecting mannose 6-P reductase (M6PR) which comprises:

- (a) reacting an antibody selective for binding to the M6PR for screening an expression library suspected of encoding the M6PR; and
- (b) detecting the antibody binding to the M6PR.

The present invention also relates to a method for producing mannose 6-phosphate (M6PR) which comprises:

- (a) providing a bacterium containing a recombinant DNA encoding mannose-6-phosphate reductase (M6PR) free of any other DNA as set forth in SEQ ID NO:1 in a culture medium;

(b) expressing the M6PR in the culture medium; and

(c) isolating the M6PR.

The present invention also relates to a method for producing mannose 6-phosphate reductase (M6PR) which comprises:

(a) providing a bacterium containing a recombinant plasmid containing DNA encoding mannose 6-phosphate reductase in the plasmid in an *E. coli* as contained in a deposit identified as ATCC 98041 in a culture medium;

(b) expressing the M6PR in the culture medium; and

(c) isolating the M6PR.

The cloning of cDNAs encoding M6PR and the partial purification and characterization of active M6PR isolated from transformed *E. coli* is described.

The DNA of celery (*Apium graveolens*) encoding M6PR was deposited with the American Type Culture Collection on Apr. 30, 1996 in the BLUESCRIPT UNI-ZAP XR (Stratagene, LaJolla, Calif.) plasmid described and claimed in U.S. Pat. No. 5,128,256 contained in *Escherichia coli* SOLR and deposited as ATCC 98041. The plasmid contains a 1.3 kbp insert which is cleaved with Eco R1 and Kpn1 restriction enzymes. The culture is grown in the presence of 50 µg/ml ampicillin in LB media. No rights are granted to the DNA encoding M6PR except those accorded by the Budapest Treaty.

For M6PR, DNA encoding sequence is shown in SEQ ID NO:1. The encoded M6PR is shown in SEQ ID NO:2 and in FIGS. 3A and 3B. For aldose 6-phosphate reductase (S6PR), the DNA sequence is shown in SEQ ID NO:3 and the encoded S6PR is shown in SEQ ID NO:4. FIGS. 4A and 4B show the alignment of M6PR and S6PR as set forth in SEQ ID NO:2 and SEQ ID NO:4.

EXAMPLE 1

Isolation of M6PR Encoding DNA

Methods

RNA Isolation and Poly(A)+RNA Selection

Total RNA was extracted from approximately 10 g samples of the fifth and seventh leaves of mature celery (*Apium graveolens* c.v., Giant Pascal) plants, according to Gilmour, S. J., et al., *Plant Physiol.* 87:745–750 (1988). Slight modifications includes: (a), addition of the polyphenol oxidase inhibitors cupferron (1 mM) and 2-mercaptobenzothiazole (1 µg/ml) to the extraction buffer immediately to prior use; (b), inclusion of three phenol:chloroform:isoamyl alcohol (25:24:1 v:v:v) partitioning steps on the aqueous phase, followed by three chloroform/isoamyl alcohol (49:1 v:v) partitionings to remove residual phenol; (c), a single LiCl precipitation followed by four ethanol precipitation steps. Total RNA yields (as determined by OD₂₆₀) averaged 500±180 µg/gFwt. The absence of contaminating DNA was confirmed on an agarose gel after RNAase treatment.

Poly(A)+RNA was isolated by oligo-dT-cellulose chromatography essentially as described by Sambrook et al (Molecular cloning. Cold Spring Harbor Laboratory Press, NY (1989)) but using a modified loading buffer (0.12 M NaCl, 0.01 M Tris-HCl pH 7.5, 0.001 M EDTA). Poly(A)+RNA was eluted from the column in the above buffer with the NaCl omitted.

In vitro Translation and Immunoprecipitation

1 µg of poly(A)+RNA from each leaf was translated in vitro using rabbit reticulocyte lysates (Promega Corp., Madison, Wis.) and [³⁵S] methionine, according to the suppliers directions. Translation products, after 1 h at 30° C., were separated by SDS-PAGE (Laemmli, U. K., *Nature*

227:401–407 (1970)) either, before (total) or after, immunoprecipitation (Hondred, D., et al., *Plant Mol. Biol.*, 9:259–275 (1987)) with M6PR-specific antisera (Ried, J. L., et al., *BioTechniques* 12:660–666 (1992)).

cDNA Library Construction

A unidirectional cDNA expression library was constructed in UniZap™ XR vector (Stratagene, LaJolla, Calif.) using a mixture of poly(A)+RNA from leaves 5 and 7 (2.5 µg of each). After packaging the phage library was amplified once before screening.

Library Screening

Two hundred thousand plaque forming units (pfu's) were screened for M6PR expression at a density of 40,000 pfu's per 140 mm Petri dish. Once phage plaques became visible (after approximately 3 h at 42° C.), nitrocellulose disks (previously soaked in 10 mM IPTG and then air-dried) were laid onto the surface of the plates and the cultures were incubated for a further 3.5 h at 37° C. The membranes were replaced with fresh IPTG soaked membranes and the cultures were incubated for a further 3 h. Both sets of membranes were screened using M6PR-specific antisera (Ried et al., *BioTechniques* 12:660–666 (1992)) at a dilution of 1:10,000 (see Everard et al., *Plant Physiol* 102:345–356 (1993), for methods used). Over 100 plaque giving positive signals in the initial screening were recovered from the plates and ten of these were subjected to two additional rounds of screening. Twelve other recombinant (as determined by α-complementation; Sambrook et al., *Molecular cloning*. Cold Spring Harbor Laboratory Press, NY (1989)) plaques that did not give a positive reaction with M6PR antisera were also selected as non-specific control clones.

After selection through three rounds of screening, clones (both putative-M6PR and nonspecific) were in vivo excised to yield phagemid (plasmid) clones in *E. coli* strain SOLR (Stratagene, LaJolla, Calif.). It should be pointed out that although 10 individual M6PR-putative clones were selected for further study it is not certain that these represented 10 individual mRNA's isolated from the original population in the leaf material used. This is because the original library was amplified once before screening (see above).

Sequencing of M6PR Clones

Three putative-M6PR clones were sequenced on both strands with an Applied Biosystems 373A sequencer using dye-primer and dye-terminator methodologies. Sequences were obtained using T3, T7 and 20-mer primers corresponding to internal sequences. Consensus sequence was derived by matching the two strands of each individual clone and by comparison of the three independent clones.

Sequence analysis was performed using SeqEd (Applied Biosystems Inc., Foster City, Calif.) and Lasergene (DNASTAR Inc., Madison, Wis.). Sequence comparisons with other databases was performed through the National Center for Biotechnology Information via the BLAST server. Peptide comparisons were made through ExPASy-Prosite and PRINTS.

Clone Confirmation

Internal Peptide Sequencing

M6PR was found to be unsuitable for amino acid sequencing in the native state, presumably because of an N-terminus block. M6PR purified as described by Loescher et al., *Plant Physiol.* 98:1396–1402 (1992) was further purified by running approximately 200 mg on a preparative 10% polyacrylamide gel under denaturing conditions. After staining with Coomassie Blue R250 (0.05%, wt:v in acetic acid:methanol:H₂O; 10:40:50, v:v:v) for 2 minutes and destaining (in stain solvent alone) the band of gel containing the M6PR was excised and electroeluted (Hunkerpillar et al.,

Methods in Enzymology 91:227–236 (1983)). The eluted protein was dried in vacuo and taken up in 80% ethanol (to remove residual SDS). The precipitated protein was pelleted by centrifugation, and the pellet dissolved in 100 mM ammonium bicarbonate (pH 8.2) prior to digestion with trypsin at 37° C. for 16 h. Trypsin was added in two equal doses (2%, by weight at each addition), with the second dose added after 8 h digestion.

Digestion products were separated by reverse phase chromatography on a 1×25 mm column (Applied Biosystems) eluted in a 90 minute linear gradient of TFA (0.1% v:v in H₂O) and acetonitrile (90:9.91.5:0.085; acetonitrile:H₂O:TFA v:v:v), at a flow rate of 830 nl/sec. Prominent peptides (as detected by OD_{212 nm}) were collected, dried down in vacuo, and subjected to amino acid sequencing on an Applied Biosystems 477A sequencer.

Test for M6PR Activity in Putative Clones

Three putative M6PR and one non-positive clone (as determined by antibody screening) were tested for M6PR activity. Duplicate 10 ml cultures of each clone were grown in LB+ampicillin (amp; 50 mg/ml). Once an average OD₆₀₀ of 0.5 had been attained one culture of each pair was induced by the addition of isopropyl-β-D-thiogalactopyranoside (IPTG; final concentration 10 mM). Control (uninduced) cultures had an equal volume of sterile water added. Cultures were maintained at 30° C. for a further 3 h and were harvested at an average OD₆₀₀ of 1.29±0.02. Cells were pelleted by centrifugation (2000×g for 5 min) and washed twice by resuspending the pellets in 5 ml Tris (100 mM, pH 7.5) containing 250 mM PMSF, with a centrifugation step between each wash. After the final wash pelleted cells were maintained on ice. Just prior to disruption the cells were resuspended in 4 ml of extraction buffer (100 mM Tris-HCl pH 7.5, 250 mM PMSF, 10 mM DTT and 0.1% Triton X-100) and transferred to blood dilution vials (American Scientific Products, McGaw Park, Ill.). Cells were ruptured by suspending the vial containing the cells 0.5 cm above the cuphorn probe of a Heat Systems W-385 sonicator (Misonics Farmingdale, N.Y.) and subjecting them to 2 minutes sonication at full power. Coolant was circulated around the vial to maintain the temperature at approximately 4° C. After sonication, 1.5 ml aliquots of the homogenate were centrifuged at 13,000×g for 5 minutes at 4° C. Supernatants were assayed for M6PR activity (Loescher et al., Plant Physiol. 98:1396–1402 (1992)); with at least two different aliquot volumes from each extract to test for linearity. Supernatant aliquots (containing approximately 100 mg of total protein) were precipitated by adding a 7 times volume of acetone. After standing overnight at -21° C. the samples were prepared by SDS-PAGE and Western blotting as described in Everard et al., Plant Physiol 102:345–356 (1993).

Large Scale Preparations and Purification

One liter LB/amp cultures were initiated by the addition of 50 ml of an overnight culture of clone D (initial OD₆₀₀, approx., 0.15). At an OD₆₀₀ of approximately 0.25 IPTG was added to a final concentration of 1 mM and the culture was incubated by 30° C. until OD₆₀₀ was between 0.8 and 1.0 (after approximately 6 hours; a preliminary experiment showing that M6PR specific activity began to level off when OD₆₀₀>1.0). Cells were harvested by centrifugation and, after washing as described above, were frozen at -80° C. Prior to extraction, cells were thawed slowly on ice and ruptured by either: 1. Sonication: cells were suspended in 20 ml extraction buffer (see previous section for composition) in a blood dilution vial and sonicated for 5 min at full power. The homogenate was centrifuged (20,000×g for 10 min), the

supernatant collected, and the pellet resuspended in 10 ml extraction buffer and sonicated for a further 5 min, this cycle was repeated three times: or 2. Decompression/shearing: Cells suspended in 20 ml extraction buffer were ruptured by three passes through a French Press followed by centrifugation (20,000×g for 20 min).

Subsequent purification steps were essentially as described in Loescher et al., Plant Physiol. 98:1396–1402 (1992) except for Reactive Yellow 86 (RY86) chromatography. Here the active fraction from the gel filtration step was split in two, and each fraction run separately on the RY86 column as described in Loescher et al., Plant Physiol. 98:1396–1402 (1992). The active fractions were pooled and either diluted by the addition of 0.5 v of column buffer or desalted using centrifugal concentrating devices (30 kD cut-off membranes) and washed with 2 volumes of column buffer. This step was performed to dilute or remove NADPH. The sample was then loaded back onto the RY86 column and activity eluted in a linear gradient between 0 and 0.2 mM NADPH in column buffer. The purified M6PR was desalted and concentrated using centrifugal filtration and was either used immediately for kinetic characterization or stored at -21° C. after adding glycerol (1:1, v:v).

Protein Determinations

Protein content was determined by the method of Bradford, M. M., Anal Biochem 136:248–254 (1976) using Bovine Serum Albumen (BSA) as a standard.

Results

Characteristics of M6PR-specific Immunoprecipitation Products

FIG. 1 shows that immunoprecipitation (with M6PR-specific antisera) of the in vitro translation products synthesized from poly(A)+RNA isolated from leaves 5 and 7 yielded a single dominant peptide (molecular mass, 35.1 kD) from each leaf. The molecular mass of authentic celery M6PR run on the same gel (but not immunoprecipitated) was 35.1 kD; immunoprecipitation prior to SDS-PAGE had no effect on the relative mobility of M6PR (data not shown). Five and seven percent of the total TCA precipitable radioactivity was recovered in the immunoprecipitation products from leaves 5 and 7, respectively.

Characteristics of the cDNA Library

The primary library consisted of >1.5×10⁶ plaque forming units (pfu's) of which 0.33% were non-recombinant, as estimated by α-complementation. Phagemids from 12 randomly selected recombinant clones were in vivo excised and used to transform *E. coli* (strain SOLR). The average insert size (after digestion with Eco R1 and Xho 1) was 1.7 kb, with a size range between 1.0–2.3 kb.

Library Screening

Of the 200,000 pfu's screened an estimated 0.15±0.04% gave a positive signal with M6PR-specific antisera and were thus identified as putative M6PR clones. Ten of these were subjected to two more rounds of screening and three of these were characterized and their authenticity as M6PR clones was confirmed as described below.

Expression of M6PR Enzyme Activity

FIG. 2A shows the activity of M6PR, in one non-specific (2) and the three putative M6PR clones (B, C and D), with and without induction with IPTG. Under IPTG induction, the putative M6PR-specific clone expressed M6PR activity, whereas little or no activity was observed in the non-specific clone (2), with or without IPTG induction, or in the uninduced cultures of the putative specific clones. The presence and absence of a peptide with identical molecular mass to authentic celery M6PR (FIG. 2B), which also reacted with M6PR-specific antisera (FIG. 2C), correlated well with the

measured enzyme activities. Trace amounts of M6PR peptide were detected in uninduced cultures of the specific clones (FIG. 2C) but the levels were below that detectable in the enzyme assay (FIG. 2A). This experiment was repeated twice with identical results.

Sequencing M6PR Clones

EcoR1-Kpn1 restriction digestion of the three putative M6PR clones yielded inserts of approximately 1.3 kb. The sequence of one of these inserts is shown in FIGS. 3A and 3B. The sequence of the two other putative clones was also determined (data not show). Each of the sequenced inserts coded for a 927 bp open reading frame; the clones differed from each other in the length of the 5' non-coding region and the poly-A tail, suggesting that they represent independent cloning events rather than duplications arising during library amplification. Translation of the open reading frame yielded a polypeptide with a molecular mass of 35.2 kD (FIGS. 3A and 3B). This value was consistent with the previously determined value for authentic celery M6PR (determined by SDS-PAGE) of 34.5 kD (Loescher et al., Plant Physiol. 98:1396-2401 (1992)) and with MALDITOFMS determined values for the recombinant and authentic M6PR protein (see below, Table 3). The predicted translation product was also consistent with an internal peptide sequence obtained after tryptic digestion of purified celery M6PR. The predicted translation product was identical to the internal amino acid sequence R, S, I, L, D, D, E, G (Arg Ser Ile Leu Asp Asp Glu Gly) (FIGS. 3A and 3B). A search of the non-redundant data base at NCBI gave only one entry showing homology (*Mycoplasma pneumoniae* M129B18 cytoadherence-accessory) indicating the rarity of this specific sequence.

Sequence Comparisons

Sequence similarities resulting from a search of the NCBI non-redundant data base resulted in 61 entries showing greater than 55% sequence homology with either the whole length of defined regions of the M6PR ORF. This information and an analysis of the amino acid sequence of M6PR through two motif analysis programs (see Materials and Methods) showed M6PR to be a member of the aldoketo reductase family. A detailed description of the features and members of this group can be gained from Bohren et al., J. Biol. Chem. 264:9547-9551 (1989). In brief, the group is typified by the three conserved domains marked in FIGS. 3A and 3B.

Five sequences of plant origin were obtained from the nucleotide sequence comparison (listed in Table 1).

TABLE 1

Sequence similarity between M6PR and entries accessible through the National Institute of Biotechnological Information (NCBI) data base.			
Accession #	Clone Identity	Comparable sequence	% Identity
D11080	Apple S6PDH NADP-dependent mRNA	Full mRNA	66
Z48383	<i>Arabidopsis thaliana</i> 315 bp EST	bases 14-325 of M6PR ORF	70
D41273	<i>Oryza sativa</i> (Rice) 462 bp EST	bases 3-291 of M6PR ORF	67
D48175	<i>Oryza sativa</i> 430 bp EST	bases 118-331 of M6PR ORF	73
X57526	<i>Hordeum vulgare</i> (Barley) aldose reductase	Bases 413-635	53

A direct sequence comparison with NADP-dependent sorbitol-6-phosphate dehydrogenase, which showed the greatest degree of homology over the entire M6PR sequence, is given in FIGS. 4A and 4B.

Purification and Characterization of Recombinant M6PR

Table 2 shows the results of a typical purification of M6PR from clone D.

TABLE 2

Summary of a typical purification of recombinant M6PR from a 1 liter, IPTG induced, culture of clone D.

STEP	vol (ml)	protein (mg)	Total Activity (mU)	Specific Activity (mU/mg prot)	Yield (%)	Purification (X)
Crude	22.5	178	13433	75	100	1
30-60% acetone	10	79	14155	179	105	2.4
S200 gel filt.	67	77	12663	164	94	2.18
RY 86, 2 passes	4	0.5	2216	4432	16	59

1 mU = 1 nmol NADPH oxidized per minute. RY 86 = Reactive Yellow 86 affinity chromatography. On the first pass M6PR was eluted with 0.1 mM NADPH, on the second pass M6PR was eluted with a linear gradient of 0 to 0.2 mM NADPH. At the end of the purification the enzyme was concentrated and NADPH was removed by ultrafiltration. Buffers used during the purification and the assay conditions were as described in Loescher et al., 1992.

With RY86 purification and desalting, a 50 to 60 fold purification was achieved. The average specific activity of the purified recombinant enzyme, 3926 ± 833 mU/mg protein (average of three preparations; final two preparations 4441 ± 13 mU/mg), was comparable to that of purified celery M6PR (3756 mU/mg protein, Loescher et al., Plant Physiol. 98:1396-1402 (1992)). FIG. 5A shows an SDS-PAGE of the various purification steps. After the second pass over the RY86 column the dominant peptide, which had a molecular mass identical to authentic celery M6PR (FIG. 5A) and cross reacted with M6PR-specific antisera (FIG. 5B) represented $88 \pm 4\%$ of the protein present (estimated by scanning densitometry; mean of 2 independent preparations).

Characterization of Recombinant M6PR

Table 3 shows some characteristics of M6PR purified from the two sources.

TABLE 3

A comparison of the characteristics of purified leaf and recombinant M6PR.		
	Authentic leaf M6PR	Recombinant M6PR
SDSPAGE determined Molecular mass. (kD)	34.8 ± 0.4 (2;3)	34.3 ± 0.4 (2;3)
MALDITOFMS determined Molecular mass. (kD)	35.21 (1;1)	35.3 ± 0.01 (1;3)
V_{max} ; mannose 6-P ($\mu\text{mol mg prot}^{-1} \text{min}^{-1}$)	6.8 ± 1.3 (2;2)	8.0 ± 1.9 (4;6)
k_m ; mannose 6-P (mM)	13.6 ± 2.7 (2;2)	10.1 ± 1.4 (4;6)
V_{max} ; NADPH ($\mu\text{mol mg prot}^{-1} \text{min}^{-1}$)	3.7 ± 0.7 (2;2)	6.0 ± 2.3 (2;3)

TABLE 3-continued

A comparison of the characteristics of purified leaf and recombinant M6PR.		
	Authentic leaf M6PR	Recombinant M6PR
k_m ; NADPH (μM)	2.1 \pm 1.1 (2;2)	6.2 \pm 2.4 (2;3)

SDSPAGE determined molecular masses of purified authentic and recombinant M6PR were determined by running them in adjacent wells on 12.5% acrylamide gels. MALDITOFMS = Matrix Assisted Laser Desorption Ionization Time Of Flight Mass Spectrometry; the matrix was sinapinic acid. The values within parentheses represent the number of individual enzyme preparations and the number of independent determinations used to calculate the mean of each parameter, respectively. Kinetic parameters were determined at 30° C. in 33 mM Tris-HCl buffer (pH 7.5) containing 3 mM DTT (see Loescher et al. (1992) for other details concerning the assays). Mannose 6-phosphate (M6P) kinetics were determined at 12 concentrations of M6P ranging from 1–50 mM under saturating NADPH concentrations (200 μM). NADPH kinetics were determined at 9 NADPH concentrations ranging from 1–50 μM . Mannose 6-phosphate concentrations in these assays were 12 mM. Best line fits on double reciprocal plots were calculated by linear regression. At least 4 data points were used for the regression analyses (in most cases more than 8); r^2 values were >0.99 except for one recombinant NADPH determination

The molecular masses of the authentic and the recombinant proteins determined by either SDS-PAGE or MALDITOF were almost identical and both were consistent with the value of 35.2 kD determined by translation of the coding region (see above). The V_{max} and k_m values determined for mannose 6-phosphate were the same for both enzymes (Table 3) and those determined for NADPH were not significantly different (by t-test).

In this Example 1 shows the successful cloning of a full length transcript coding for mannose 6-phosphate reductase from celery. This represents the first report of the cloning of the purification of competent recombinant enzyme from transformed *E. coli*. The purified recombinant protein had physical and kinetic properties indistinguishable from the purified plant enzyme.

The authenticity of the clones was confirmed by the following criteria. 1). Only putative clones displayed M6PR activity when induced with IPTG. The activity correlated with the presence of a peptide, of identical molecular mass to authentic celery leaf M6PR (Table 3). This peptide cross reacted with M6PR-specific antisera (FIG. 2C). No activity or immuno reactive peptide was observed in non-specific clones. 2). A tryptic digestion product from authentic celery M6PR had 100% homology to a peptide present within the open reading frame of the putative clone (FIGS. 3A and 3B). 3). Database comparisons showed that the clones had a high degree of homology to the aldo-keto reductase family. The greatest degree of homology (spanning the whole ORF with 67% sequence and 64% amino acid identity; 84% similarity when amino acids with the same functional properties were considered (FIGS. 4A and 4B)) was with NADP-dependent D-sorbitol 6-phosphate dehydrogenase (Kanayama et al., Plant Physiology 10:1607–1608 (1992)), a key enzyme in sorbitol biosynthesis in woody Rosaceae species. The sequence and amino acid similarity was very low (<22%) on an amino acid and sequence comparison basis) between M6PR and mannitol dehydrogenase (MTD; Williamson et al., Proc Natl. Acad. Sci. 92:7148–7152 (1995)), the only other mannitol metabolizing enzyme cloned to date from higher plants. However, four other similar plant sequences were obtained from the databases. Three of these were for partial sequences from arabidopsis and rice which showed strong homology with the 5' end of M6PR (Table 1). Although there are not known to be any reports of sugar

alcohols in arabidopsis and rice, the recent discovery of a homologue of mannitol dehydrogenase in arabidopsis (Williamson et al., Proc. Natl. Acad. Sci. 92:7148–7152 (1995)) may indirectly confirm Bieleski's admonition that the presence of sugar alcohols should not be discounted until proven absent (Bieleski, R. L., Sugar Alcohols. In: F. A. Loewus and W. Tanner, eds., Plant carbohydrates I. Intracellular Carbohydrates, Encyc. Plant Physiol., Vol. 13A, New Series. Springer-Verlag, NY, pp. 158–192 (1982)). The fifth plant enzyme sequence with similarity to M6PR was an aldose reductase from barley (see below). As mentioned, there is almost no information as to how sugar alcohol metabolism is regulated and integrated with the other products of primary production (sucrose, starch and nitrogen metabolism) in higher plants. In contrast, there is a relatively large body of work on the kinetics and regulation of animal aldose reductase, driven by their putative role in the pathology of diabetes mellitus (Borhani, D. W., et al., J. Biol. Chem. 267:24841–24847 (1992)). The sequence homologies suggest that insights into regulatory sites and mechanisms in the plant enzymes may be gained from the animal literature. For example, several sites are highly conserved between the animal and plant enzymes (marked as motifs 1, 2 and 3 in FIG. 3B) including the peptide IPKS (within motif 3, towards the 3' end of the coding region). The lysine (K) within this motif has been shown, by chemical modification studies in animals, to be the likely NADPH binding site (Morijana et al., FASEB J 46:1330 (abstract) (1987); Bohren et al., J. Biol. Chem. 264:9547–9551 (1989)). This motif is completely conserved in terms of amino acids and position in M6PR (IPKS, amino acids 260–264), NADP-dependent D-sorbitol 6-phosphate dehydrogenase (IPKS, amino acids 260–264) and in aldose reductase (IPKS, amino acids 257–260) cloned from desiccated barley embryo's (Bartels et al., EMBO J 10:1037–1043 (1991)). This latter aldose reductase has been associated with tissues subjected to desiccation and is inducible with ABA which is of interest given the role that mannitol synthesis and M6PR play in salinity stressed celery (Everard et al., Plant Physiol. 106:281–292 (1994); Loescher et al., Plant Physiology 170–178 (1995)). Another similarity between the animal aldose reductases and M6PR is the apparent lack of post-translational modification. In vitro translation of celery poly(A)+RNA resulted in a peptide immunoprecipitation product of identical molecular mass to authentic leaf M6PR (FIG. 1B) indicating that post-translational modification is unlikely to occur in vivo, a conclusion also drawn for bovine lens aldose reductase (the model for study of this class of enzymes prior to the availability of recombinant enzymes, Schade et al., J.B.C. 265(7):3628–3635 (1990). Finally, aldose reductases in animals are sensitive to oxidation (Petrasch et al., JBC 267(34) :24833–24840 (1992)), as is the M6PR (Loescher et al., Plant Physiol. 98:1396–1402 (1992)), and the current thinking is that redox activation/inactivation may play an important role in the in vivo regulation of M6PR. The importance of redox activation of extraplasmidic plant enzymes has grown in recent years and an increasing number of cytosolic (Anderson et al., Planta 196:118–124 (1995)) and mitochondrial enzymes are being reported to be regulated in part by this mechanism. The recombinant enzyme has a specific activity similar to the plant enzyme which suggests that if redox activation is a factor then the *E. coli* thioredoxin system is competent. Such information should lead to approaches to look for regulatory mechanisms in the plant enzymes, studies that will be simpler with the recombinant enzyme which appears fully competent and kinetically indistinguishable from the

authentic enzyme. Ultimately, mechanisms may be explored at the enzyme level using site directed mutagenesis and at the whole plant level by studies into message level regulation, such as that recently reported for NADP-dependent D-sorbitol 6-phosphate dehydrogenase (Kanayama et al., Plant Physiology 100:1607-1608 (1995)), and by pathway suppression using antisense and cosuppression techniques. Such studies should give insight into the roles of polyols in primary carbon metabolism and hence plant productivity as well as in stress tolerance.

DNA is incorporated into plants in a manner known to those skilled in the art as represented by U.S. Pat. No. 5,492,820 to Sonnewald et al. Various well known vectors are used.

EXAMPLE 2

A large number of techniques are available for inserting M6PR DNA into a plant host cell. Those techniques often include transformation with T-DNA using *Agrobacterium tumefaciens* or *A. rhizogenes* containing the Ti or Ri plasmids (respectively) as transformation agents. Some of the other methods used include fusion, biolistic or conventional injection, or electroporation. If *Agrobacterium* related methods are used, the DNA is cloned into a special plasmid, either an intermediate vector or into a binary vector. Intermediate vectors are integrated into the Ti or Ri plasmid by homologous recombination resulting from sequences that are homologous to sequences in the T-DNA. The Ti or Ri plasmid also comprises the vir region necessary for transfer of the T-DNA. The intermediate vector are transferred into *Agrobacterium* by means of a helper plasmid (via conjugation). Binary vectors replicate themselves both in *E. coli* and *Agrobacterium*. These vectors include a selection marker gene and a linker or polylinker that are framed by the right and left T-DNA border regions. These are transformed directly into *Agrobacterium* and the *Agrobacterium* is then used as a host cell for the plasmid carrying the vir region. The vir region is also necessary for the transfer of the

T-DNA into the plant cell. Additional T-DNA may be contained. The transformed bacteria are used for transformation of plant cells. Plant explants (e.g., sections of leaves, stems and roots, segments of petioles, flowers, and flower parts) are cultivated with *Agrobacterium tumefaciens* or *A. rhizogenes* for the transfer of the DNA into the plant cell. Whole plants are then regenerated (from the infected plant material, or from protoplasts or suspension-cultivated cells) in a suitable medium which can contain antibiotics or biocides (e.g., kanamycin, bleomycin, hygromycin, chloramphenicol, among others) for selection of transformed plant cells. Unlike *Agrobacterium*-mediated insertion of M6PR DNA, no special demands are necessary for construction for plasmids used for particle bombardment, fusion, injection, or electroporation. It is possible to use ordinary plasmids, e.g., pUC derivatives, although selection markers are usually included. However obtained, whole plants are then tested for the presence of the inserted DNA. The transformed cells grow normally in the plant and eventually give rise to reproductive organs, i.e., flowers, that can be used in an ordinary breeding program. The resulting hybrids have the appropriate phenotypic properties.

Expression of the M6PR DNA requires a promoter associated with the cloned gene. Among the many examples available are viral promoters such as the cauliflower mosaic virus 35 S promoter, heat shock protein promoters such as the HSP 70 promoter, light induced promoters such as the ST-Ls1 or the rubisco small subunit (SSU) promoters, stress response proteins such as the PR protein promoter, the *Agrobacterium tumefaciens* nos promoter, and various organ, root, tuber (e.g., the class I patatin), and leaf specific promoters. A termination signal is also used in these constructs, e.g., the 3'-end of the poly-A side of the octopine synthase gene.

It is intended that the foregoing description be only illustrative of the present invention and that the present invention be limited only by the hereinafter appended claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(iii) NUMBER OF SEQUENCES: 4

(2) INFORMATION FOR SEQ ID NO:1:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1207
- (B) TYPE: Nucleic Acid
- (C) STRANDEDNESS: Single
- (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE:

- (A) DESCRIPTION: Synthetic DNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:

- (A) ORGANISM: Celery

(vii) IMMEDIATE SOURCE:

- (A) LIBRARY:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

-continued

TAGAGAAAGA AGGAGGATAG TTTTTTAGGC TACACAACAC	40
AGTTCTAAAA ATCTTTTATC GTTTGTTAGG TTTGACAATG	80
GCAATAACTC TTAACAGCGG CTTTAAAATG CCCGTTCTGG	120
GTCTCGGCGT CTGGCGTATG GACCGTAATG AAATCAAGAA	160
TCTCCTCCTT TCCGCGATTA ACCTTGGTTA TCGTCACTTT	200
GACTGTGCTG CTGACTACAA GAATGAGTTA GAAGTAGGGG	240
AGGCATTTAA AGAGGCTTTT GATACTGATC TTGTCAAGAG	280
GGAGGATCTG TTTATTACTA CCAAGCTCTG GAACTCAGAC	320
CATGGACATG TAATTGAGGC ATGCAAAAAC AGTCTCAAGA	360
AGCTTCAGCT AGAATATCTT GATCTTTACC TCATTCACCT	400
CCCAATGGCT TCTAAACATT CCGGAATTGG TACTACTCGA	440
AGTATCTTGG ATGATGAAGG TGTTTGGGAG GTTGATGCAA	480
CCATTTCACT GGAAGCTACA TGGCATGAGA TGGAGAAGCT	520
GGTTGAAATG GGCTTAGTCC GTAGCATAGG AATCAGCAAC	560
TATGATGTTT ACTTGACCAG AGATATCTTG TCATATTCCA	600
AGATCAAGCC TGCTGTAAAT CAGATCGAGA CGCACCTTA	640
CTTCCAAAGA GATTCTCTGA TCAAATCTG TCAGAAGTAT	680
GGCATTGCTA TCACAGCACA CACACCACTA GGCGGCGCAT	720
TGGCTAATAC TGAGCGATTT GGATCAGTTT CGTGCTTAGA	760
TGATCCAGTT CTTAAGAAAT TATCTGACAA ACACAACAAG	800
TCACCAGCTC AGATTGTTCT CCGTTGGGGT GTGCAGCGCA	840
ACACAATTGT AATTCCCAAG TCATCGAAAA CTAAAAGACT	880
CGAGGAAAAC ATCAACATTT TTGACTTTGA GTTGAGCAAG	920
GAAGATATGG AGCTCATCAA AACAATGGAG CGCAACCAAA	960
GGAGTAACAC ACCTGCTAAA GCTTGGGGAA TAGATGTTTA	1000
TGCTTGATGG CATAACACAT TCTTCACTGT ATTTTTATCA	1040
TTGTTATTCC ACAATTCAGA GTGGTTGTCA TTTTACTTGG	1080
CTATTGTGTG TGGAGGGGAA TGTGTGTTGA GTTGTGTAG	1120
TAATTGTACA AGGCATAAAG CCTTTAAATA ACCCATCATA	1160
TGTAAATGGG AAATGCCATG ATTTGGTCAA AAAAAAAAAA	1200
AAAAAAA	1207

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 309
 (B) TYPE: Amino Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

- (ii) MOLECULE TYPE:
 (A) DESCRIPTION:

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

-continued

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: celery

(vii) IMMEDIATE SOURCE:
 (A) LIBRARY:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

Met Ala Ile Thr Leu Asn Ser Gly Phe Lys Met Pro Val Leu Gly
 5 10 15

Leu Gly Val Trp Arg Met Asp Arg Asn Glu Ile Lys Asn Leu Leu
 20 25 30

Leu Ser Ala Ile Asn Leu Gly Tyr Arg His Phe Asp Cys Ala Ala
 35 40 45

Asp Tyr Lys Asn Glu Leu Glu Val Gly Glu Ala Phe Lys Glu Ala
 50 55 60

Phe Asp Thr Asp Leu Val Lys Arg Glu Asp Leu Phe Ile Thr Thr
 65 70 75

Lys Leu Trp Asn Ser Asp His Gly His Val Ile Glu Ala Cys Lys
 80 85 90

Asn Ser Leu Lys Lys Leu Gln Leu Glu Tyr Leu Asp Leu Tyr Leu
 95 100 105

Ile His Phe Pro Met Ala Ser Lys His Ser Gly Ile Gly Thr Thr
 110 115 120

Arg Ser Ile Leu Asp Asp Glu Gly Val Trp Glu Val Asp Ala Thr
 125 130 135

Ile Ser Leu Glu Ala Thr Trp His Glu Met Glu Lys Leu Val Glu
 140 145 150

Met Gly Leu Val Arg Ser Ile Gly Ile Ser Asn Tyr Asp Val Tyr
 155 160 165

Leu Thr Arg Asp Ile Leu Ser Tyr Ser Lys Ile Lys Pro Ala Val
 170 175 180

Asn Gln Ile Glu Thr His Pro Tyr Phe Gln Arg Asp Ser Leu Ile
 185 190 195

Lys Phe Cys Gln Lys Tyr Gly Ile Ala Ile Thr Ala His Thr Pro
 200 205 210

Leu Gly Gly Ala Leu Ala Asn Thr Glu Arg Phe Gly Ser Val Ser
 215 220 225

Cys Leu Asp Asp Pro Val Leu Lys Lys Leu Ser Asp Lys His Asn
 230 235 240

Lys Ser Pro Ala Gln Ile Val Leu Arg Trp Gly Val Gln Arg Asn
 245 250 255

Thr Ile Val Ile Pro Lys Ser Ser Lys Thr Lys Arg Leu Glu Glu
 260 265 270

Asn Ile Asn Ile Phe Asp Phe Glu Leu Ser Lys Glu Asp Met Glu
 275 280 285

Leu Ile Lys Thr Met Glu Arg Asn Gln Arg Ser Asn Thr Pro Ala
 290 295 300

Lys Ala Trp Gly Ile Asp Val Tyr Ala
 305

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1259
 (B) TYPE: Nucleic Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

-continued

(ii) MOLECULE TYPE:
 (A) DESCRIPTION: Synthetic DNA

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: apple

(vii) IMMEDIATE SOURCE:
 (A) LIBRARY: N/A

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

CCGCTCTAGA ACTAGTGGAC GAGAGAAAAA CAGAAACAGG	40
CTGCAGCAGT CGCTGAGAGA GTTTGGAGAG TGAGAAAAACA	80
TGTCCACCGT CACCCTGAGC AGTGGCTACG AGATGCCGGT	120
CATCGGTCTC GGCCTTTGGC GTCTGGAGAA GGACGAGCTT	160
AAAGAAGTCA TCTTAAATGC TATTAAGATT GGCTATCGCC	200
ATTTTACTG TGCTGCTCAT TACAAGAGTG AAGCAGACGT	240
TGGAGAAGCA CTTGCAGAAG CATTTAAGAC TGGACTTGTT	280
AAGAGGGAAG AACTTTTCAT TACCACCAAG ATTTGGAATT	320
CAGACCATGG GCATGTGGTG GAGGCCTGTA AGAACAGCCT	360
CGAGAAGCTT CAGATAGATT ATCTGGATCT CTACCTGGTT	400
CACTACCCAA TGCCCACAAA GCACAATGCA ATTGGTAAAA	440
CTGCCAGTCT TTTGGGCGAG GATAAGGTGT TGGACATCGA	480
TGTAACAATT TCCCTTCAAC AAACCTGGGA GGGCATGGAA	520
AAGACCGTCT CTTTGGGCTT AGTTCGCAGC ATTGGTCTCA	560
GCAACTATGA GCTCTTTCTA ACTAGAGATT GCTTGGCTTA	600
CTCCAAAATA AAGCCTGCTG TGAGCCAATT TGAAACCCAC	640
CCCTATTTCC AGCGCGACTC TCTCGTCAAA TTCTGTATGA	680
AACACGGCGT TCTTCCCACA GCTCACACCC CTCTCGGAGG	720
TGCTGCTGCC AACAAGGATA TGTTTGGTTC TGTTTCACCT	760
TTGGATGATC CAGTTCTCAA TGATGTGGCT AAGAAATACG	800
GAAAGAGCGT GGCACAAATC TGTCTGAGGT GGGGAATTCA	840
GAGGAAAACA GCAGTGATTC CAAAATCATC GAAAATTCAG	880
CGATTGAAAG AGAATTTGGA GGTTC TTGAA TTCCAGCTGA	920
GCGATGAAGA CATGCAGCTC ATCTACAGTA TCGACAGGAA	960
GTATCGTACC AGTCTACCTT CCAAGACTTG GGGCTTAGAC	1000
GTGTATGCAT AAGCGTGCCA TTCAAAAACC TTCGAATTGC	1040
TGCCTCCGCA ACTTCTTCCA AGGCTGTTCA ACGGAAGCGA	1080
AATGGAAACT ATCGTGAATC TTA CT TACAA TAAACTGAGC	1120
TTCATATAAT TTTCCAGAAG CTCATCTATC TGCTAGTTTG	1160
AAA ACTTCAT TATTCGCCCT TTGCATTAGG CCTTGCAAAG	1200
GAAAATATAA TAAACGGCCC TTGTATTTTT TTTGGTACTT	1240
AATAAATGAG TTATTTAAAG	1259

-continued

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 310
 (B) TYPE: Amino Acid
 (C) STRANDEDNESS: Single
 (D) TOPOLOGY: Linear

(ii) MOLECULE TYPE:
 (A) DESCRIPTION:

(iii) HYPOTHETICAL: No

(iv) ANTI-SENSE: No

(vi) ORIGINAL SOURCE:
 (A) ORGANISM: apple

(vii) IMMEDIATE SOURCE:
 (A) LIBRARY: N/A

(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

```

Met Ser Thr Val Thr Leu Ser Ser Gly Tyr Glu Met Pro Val Ile
                5                10                15
Gly Leu Gly Leu Trp Arg Leu Glu Lys Asp Glu Leu Lys Glu Val
                20                25                30
Ile Leu Asn Ala Ile Lys Ile Gly Tyr Arg His Phe Asp Cys Ala
                35                40                45
Ala His Tyr Lys Ser Glu Ala Asp Val Gly Glu Ala Leu Ala Glu
                50                55                60
Ala Phe Lys Thr Gly Leu Val Lys Arg Glu Glu Leu Phe Ile Thr
                65                70                75
Thr Lys Ile Trp Asn Ser Asp His Gly His Val Val Glu Ala Cys
                80                85                90
Lys Asn Ser Leu Glu Lys Leu Gln Ile Asp Tyr Leu Asp Leu Tyr
                95                100               105
Leu Val His Tyr Pro Met Pro Thr Lys His Asn Ala Ile Gly Lys
                110               115               120
Thr Ala Ser Leu Leu Gly Glu Asp Lys Val Leu Asp Ile Asp Val
                125               130               135
Thr Ile Ser Leu Gln Gln Thr Trp Glu Gly Met Glu Lys Thr Val
                140               145               150
Ser Leu Gly Leu Val Arg Ser Ile Gly Leu Ser Asn Tyr Glu Leu
                155               160               165
Phe Leu Thr Arg Asp Cys Leu Ala Tyr Ser Lys Ile Lys Pro Ala
                170               175               180
Val Ser Gln Phe Glu Thr His Pro Tyr Phe Gln Arg Asp Ser Leu
                185               190               195
Val Lys Phe Cys Met Lys His Gly Val Leu Pro Thr Ala His Thr
                200               205               210
Pro Leu Gly Gly Ala Ala Ala Asn Lys Asp Met Phe Gly Ser Val
                215               220               225
Ser Pro Leu Asp Asp Pro Val Leu Asn Asp Val Ala Lys Lys Tyr
                230               235               240
Gly Lys Ser Val Ala Gln Ile Cys Leu Arg Trp Gly Ile Gln Arg
                245               250               255
Lys Thr Ala Val Ile Pro Lys Ser Ser Lys Ile Gln Arg Leu Lys
                260               265               270

```

-continued

Glu	Asn	Leu	Glu	Val	Leu	Glu	Phe	Gln	Leu	Ser	Asp	Glu	Asp	Met
				275					280					285
Gln	Leu	Ile	Tyr	Ser	Ile	Asp	Arg	Lys	Tyr	Arg	Thr	Ser	Leu	Pro
				290					295					300
Ser	Lys	Thr	Trp	Gly	Leu	Asp	Val	Tyr	Ala					
				305					310					

We claim:

1. A method for detecting unknown DNA encoding mannose-6-phosphate reductase (M6PR) which comprises probing the unknown DNA with a probe DNA comprising a sequence or a portion of the sequence as set forth in SEQ ID NO:1.
2. A method for detecting unknown DNA encoding mannose-6-phosphate reductase (M6PR) which comprises probing the unknown DNA with a DNA probe comprising DNA encoding M6PR or a portion of the M6PR, the DNA probe including a region which is noncoding, in a plasmid in *E. Coli* as contained in a deposit identified as ATCC 98041.
3. A method for detecting whether DNA from a plant encodes mannose-6-phosphate reductase (M6PR) comprising:

- (a) isolating the DNA from the plant; and
- (b) probing the DNA with a probe DNA encoding the M6PR or a portion of the M6PR DNA probe which has a sequence as set forth in SEQ ID NO:1, including a region in the DNA probe which is noncoding, wherein the probe hybridizes to the DNA which encodes the M6PR.
4. The method of claim 3 wherein the probe is the M6PR or the portion thereof in a plasmid in *Escherichia Coli* as contained in a deposit identified as ATCC 98041.
5. The method of claim 3 wherein the DNA is a cDNA of RNA from the plant.

* * * * *