



US006939692B2

(12) **United States Patent**  
**Bathe et al.**

(10) **Patent No.:** **US 6,939,692 B2**  
(45) **Date of Patent:** **Sep. 6, 2005**

(54) **NUCLEOTIDE SEQUENCES CODING FOR THE PKNB GENE**

(75) Inventors: **Brigitte Bathe**, Saltzkotten (DE); **Stephan Hans**, Osnabrueck (DE); **Mike Farwick**, Bielefeld (DE); **Thomas Hermann**, Bielefeld (DE)

(73) Assignee: **Degussa AG**, Duesseldorf (DE)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 234 days.

(21) Appl. No.: **09/949,970**

(22) Filed: **Sep. 12, 2001**

(65) **Prior Publication Data**

US 2002/0042105 A1 Apr. 11, 2002

**Related U.S. Application Data**

(60) Provisional application No. 60/297,250, filed on Jun. 12, 2001.

(30) **Foreign Application Priority Data**

Sep. 12, 2000 (DE) ..... 100 44 912  
Apr. 25, 2001 (DE) ..... 101 20 095

(51) **Int. Cl.**<sup>7</sup> ..... **C12P 13/04**

(52) **U.S. Cl.** ..... **435/106**; 435/183; 435/193;  
435/194; 435/252.3; 435/252.32; 435/320.1;  
536/23.2

(58) **Field of Search** ..... 435/106, 183,  
435/193, 194, 252.3, 252.32, 320.1; 536/23.2

(56) **References Cited**

**FOREIGN PATENT DOCUMENTS**

EP 1 029 919 8/2000  
EP 1 108 790 6/2001

**OTHER PUBLICATIONS**

Cocito et al. Accession AAZ38383. Feb. 22, 2000 (Alignment No. 1).\*

Cocito et al. Accession AAZ38383. Feb. 22, 2000 (Alignment No. 2).\*

Zhao et al. Accession AQ391796. Mar. 6, 1999.\*

Attwood et al. Which craft is best in bioinformatics? Comput. Chem. 2001, vol. 25(4), pp. 329–339.\*

Ponting, C.P. Issues in predicting protein function from sequence. Brief. Bioinform. Mar. 2001, vol. 2(1), pp. 19–29.\*

Database Swail “Online”, EBI, “M. Leprae PknB”, XP 002185694, Acc. No. P54744, Oct. 1, 1996.

Josef Cremer, et al., Applied and Environmental Microbiology, vol. 57, No. 6, pp. 1746–1752, “Control of the Lysine Biosynthesis Sequence in Corynebacterium Glutamicum as Analyzed by Overexpression of the Individual Corresponding Genes”, Jun. 1, 1991.

\* cited by examiner

*Primary Examiner*—Ponnathapu Achutamurthy

*Assistant Examiner*—Christian L. Fronda

(74) *Attorney, Agent, or Firm*—Oblon, Spivak, McClelland, Maier & Neustadt, P.C.

(57) **ABSTRACT**

An isolated polynucleotide which contains a polynucleotide sequence selected from the group comprising:

- (a) a polynucleotide which is at least 70% identical to a polynucleotide coding for a polypeptide containing the amino acid sequence of SEQ ID No. 2,
- (b) a polynucleotide coding for a polypeptide containing an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID No. 2,
- (c) a polynucleotide which is complementary to the polynucleotides of (a) or (b), and
- (d) a polynucleotide containing at least 15 consecutive nucleotides of the polynucleotide sequence of (a), (b) or (c),

and a fermentation process for the preparation of L-amino acids using corynebacteria in which at least the pknB gene is amplified, and to the use, as hybridization probes, of polynucleotides containing the sequences according to the invention.

**38 Claims, No Drawings**

## NUCLEOTIDE SEQUENCES CODING FOR THE PKNB GENE

### CROSS-REFERENCES TO RELATED APPLICATIONS

This application claims benefit to U.S. Provisional Application Ser. No. 60/297,250, filed on Jun. 12, 2001, and incorporated herein by reference.

### BACKGROUND OF THE INVENTION

#### 1. Field of the Invention

The present invention relates nucleotide sequences from corynebacteria coding for the *pknB* gene and a fermentation process for the preparation of amino acids using bacteria in which the endogenous *pknB* gene is amplified.

#### 2. State of the Art

L-Amino acids, especially L-lysine, are used in human medicine, in the pharmaceutical industry, in the food industry, and, in particular, in animal nutrition. It is known that amino acids are prepared by the fermentation of strains of corynebacteria, especially *Corynebacterium glutamicum*. Because of their great importance, attempts are constantly being made to improve these preparative processes. Improvements to the processes may relate to measures involving the fermentation technology, e.g. stirring and oxygen supply, or the composition of the nutrient media, e.g. the sugar concentration during fermentation, or the work-up to the product form, e.g. by ion exchange chromatography, or the intrinsic productivity characteristics of the microorganism itself.

The productivity characteristics of these microorganisms are improved by using methods of mutagenesis, selection and mutant choice to give strains which are resistant to antimetabolites or auxotrophic for metabolites important in regulation, and produce amino acids.

Methods of recombinant DNA technology have also been used for some years to improve L-amino acid-producing strains of *Corynebacterium* by amplifying individual amino acid biosynthesis genes and studying the effect on amino acid production. However, there remains a need for improved methods of producing L-amino acids.

### SUMMARY OF THE INVENTION

It is an object of the present invention to provide novel measures for improving the preparation of amino acids by fermentation.

It is another object of the present invention to provide nucleic acid sequences which are useful for the production of amino acids.

Accordingly, the present invention provides an isolated polynucleotide from corynebacteria which contains a polynucleotide sequence coding for the *pknB* gene and is selected from the following group:

(a) a polynucleotide which is at least 70% identical to a polynucleotide coding for a polypeptide containing the amino acid sequence of SEQ ID No. 2,

(b) a polynucleotide coding for a polypeptide containing an amino acid sequence which is at least 70% identical to the amino acid sequence of SEQ ID No. 2,

(c) a polynucleotide which is complementary to the polynucleotides of (a) or (b), and

(d) a polynucleotide containing at least 15 consecutive nucleotides of the polynucleotide sequence of (a), (b) or (c), where the polypeptide preferably has the activity of protein kinase B.

The present invention also provides the above-mentioned polynucleotide, which is preferably a replicatable DNA containing:

- (i) the nucleotide sequence shown in SEQ ID No. 1, or
- (ii) at least one sequence corresponding to sequence (i) within the degeneracy of the genetic code, or
- (iii) at least one sequence which hybridizes with the sequence complementary to sequence (i) or (ii), and optionally
- (iv) neutral sense mutations in (i).

The present invention also provides:

a replicatable polynucleotide, especially DNA, containing the nucleotide sequence as shown in SEQ ID No. 1, a polynucleotide coding for a polypeptide containing the amino acid sequence as shown in SEQ ID No. 2, a vector containing the polynucleotide according to the invention, especially a shuttle vector or plasmid vector, and

corynebacteria which contain the vector or in which the endogenous *pknB* gene is amplified.

The present invention also provides a fermentation process for the preparation of an L-amino acid, comprising:

(a) fermenting corynebacteria which produce the L-amino acid in a medium, wherein at least the *pknB* gene or nucleotide sequences coding therefor are amplified in the corynebacteria,

(b) enriching the L-amino acid in the medium or in the cells of the corynebacteria, and

(c) isolating the L-amino acid.

The present invention additionally provides polynucleotides consisting substantially of a polynucleotide sequence which are obtainable by screening, by means of hybridization, of an appropriate gene library of a corynebacterium, containing the complete gene or parts thereof, with a probe containing the sequence of the polynucleotide of the invention according to SEQ ID No. 1 or a fragment thereof, and by isolation of the polynucleotide sequence.

A more complete appreciation of the invention and many of the attendant advantages thereof will be readily obtained as the same becomes better understood by reference to the following detailed description.

### DETAILED DESCRIPTION OF THE INVENTION

When L-amino acids or amino acids are mentioned hereafter, this is understood to refer to one or more amino acids, including their salts, selected from the group comprising L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine. L-Lysine is particularly preferred.

When L-lysine or lysine is mentioned hereafter, this is understood to refer to not only the bases but also the salts, e.g. lysine monohydrochloride or lysine sulfate.

As hybridization probes for RNA, cDNA and DNA, polynucleotides containing the sequences according to the invention are suitable for isolating the full length of nucleic acids, or polynucleotides or genes, coding for protein kinase B, or for isolating nucleic acids, or polynucleotides or genes, whose sequence exhibits a high degree of similarity to the sequence of the *pknB* gene. They are also suitable for incorporation into so-called arrays, micro-arrays or DNA chips for detecting and determining the corresponding polynucleotides.

Polynucleotides containing the sequences according to the invention are further suitable as primers for the preparation, by the polymerase chain reaction (PCR), of DNA of genes coding for protein kinase B.

Such oligonucleotides serving as probes or primers contain at least 25, 26, 27, 28, 29 or 30, preferably at least 20, 21, 22, 23 or 24 and very particularly preferably at least 15, 16, 17, 18 or 19 consecutive nucleotides. Oligonucleotides with a length of at least 31, 32, 33, 34, 35, 36, 37, 38, 39 or 40 or at least 41, 42, 43, 44, 45, 46, 47, 48, 49 or 50 nucleotides are also suitable. Oligonucleotides with a length of at least 100, 150, 200, 250 or 300 nucleotides are also suitable.

As used herein, the term "isolated" refers to a material, e.g., a nucleic acid sequence, separated from its natural environment.

"Polynucleotide" refers in general to polyribonucleotides and polydeoxyribonucleotides, it being possible for the RNAs or DNAs in question to be unmodified or modified.

The polynucleotides according to the invention include a polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom, as well as polynucleotides which are in particular at least 70% to 80%, preferably at least 81% to 85%, particularly preferably at least 86% to 90% and very particularly preferably at least 91%, 93%, 95%, 97% or 99% identical to the polynucleotide according to SEQ ID No. 1 or a fragment prepared therefrom.

"Polypeptides" are understood as meaning peptides or proteins containing two or more amino acids bonded via peptide links.

The polypeptides according to the invention include a polypeptide according to SEQ ID No. 2, especially those with the biological activity of protein kinase B and also those which are at least 70% to 80%, preferably at least 81% to 85%, particularly preferably at least 86% to 90% and very particularly preferably at least 91%, 93%, 95%, 97% or 99% identical to the polypeptide according to SEQ ID No. 2, and having the activity of protein kinase B.

The invention further relates to a fermentation process for the preparation of amino acids selected from the group comprising L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-lysine, L-tryptophan and L-arginine, using corynebacteria which, in particular, already produce amino acids and in which the nucleotide sequences coding for the *pknB* gene are amplified and, in particular, overexpressed.

In this context the term "enhancement" describes the increase in the intracellular activity, in a microorganism, of one or more enzymes which are coded for by the appropriate DNA, for example by increasing the copy number of the gene(s) or allele(s), using a strong promoter or using a gene or allele coding for an appropriate enzyme with a high activity, and optionally combining these measures.

By enhancement measures, in particular over-expression, the activity or concentration of the corresponding protein is in general increased by at least 10%, 25%, 50%, 75%, 100%, 150%, 200%, 300%, 400% or 500%, up to a maximum of 1000% or 2000%, based on the starting microorganism.

The microorganisms provided by the present invention can produce L-amino acids from glucose, sucrose, lactose, fructose, maltose, molasses, starch or cellulose or from glycerol and ethanol. The microorganisms can be representatives of corynebacteria, especially of the genus *Coryne-*

*bacterium*. The species *Corynebacterium glutamicum* may be mentioned in particular in the genus *Corynebacterium*, which is known to those skilled in the art for its ability to produce L-amino acids.

The following known wild-type strains:

*Corynebacterium glutamicum* ATCC13032

*Corynebacterium acetoglutamicum* ATCC 15806

*Corynebacterium acetoacidophilum* ATCC 13870

*Corynebacterium thermoaminogenes* FERM BP-1539

*Corynebacterium melassecola* ATCC17965

*Brevibacterium flavum* ATCC14067

*Brevibacterium lactofermentum* ATCC13869 and

*Brevibacterium divaricatum* ATCC14020

and L-amino acid-producing mutants or strains prepared therefrom, are particularly suitable strains of the genus *Corynebacterium*, especially of the species *Corynebacterium glutamicum* (*C. glutamicum*).

The novel *pknB* gene of *C. glutamicum* coding for the enzyme protein kinase B (EC 2.7.1.37) has been isolated using the procedures described herein.

The first step in isolating the *pknB* gene or other genes of *C. glutamicum* is to construct a gene library of this microorganism in *Escherichia coli* (*E. coli*). The construction of gene libraries is documented in generally well-known textbooks and manuals. Examples which may be mentioned are the textbook by Winnacker entitled *From Genes to Clones, Introduction to Gene Technology* (Verlag Chemie, Weinheim, Germany, 1990) or the manual by Sambrook et al. entitled *Molecular Cloning, A Laboratory Manual* (Cold Spring Harbor Laboratory Press, 1989). A very well-known gene library is that of the *E. coli* K-12 strain W3110, which was constructed by Kohara et al. (*Cell* 50, 495–508 (1987)) in  $\lambda$  vectors. Bathe et al. (*Molecular and General Genetics* 252, 255–265, 1996) describe a gene library of *C. glutamicum* ATCC13032, which was constructed using cosmid vector SuperCos I (Wahl et al., 1987, *Proceedings of the National Academy of Sciences USA* 84, 2160–2164) in the *E. coli* K-12 strain NM554 (Raleigh et al., 1988, *Nucleic Acids Research* 16, 1563–1575). Börmann et al. (*Molecular Microbiology* 6(3), 317–326 (1992)) in turn describe a gene library of *C. glutamicum* ATCC13032 using cosmid pHC79 (Hohn and Collins, *Gene* 11, 291–298 (1980)).

A gene library of *C. glutamicum* in *E. coli* can also be constructed using plasmids like pBR322 (Bolivar, *Life Sciences* 25, 807–818 (1979)) or pUC9 (Vieira et al., 1982, *Gene* 19, 259–268). Restriction- and recombination-defective *E. coli* strains are particularly suitable as hosts, an example being the strain DH5 $\alpha$ mc $r$ , which has been described by Grant et al. (*Proceedings of the National Academy of Sciences USA* 87 (1990) 4645–4649). The long DNA fragments cloned with the aid of cosmids can then in turn be subcloned into common vectors suitable for sequencing, and subsequently sequenced, e.g. as described by Sanger et al. (*Proceedings of the National Academy of Sciences of the United States of America* 74, 5463–5467, 1977).

The DNA sequences obtained can then be examined with known algorithms or sequence analysis programs, e.g. that of Staden (*Nucleic Acids Research* 14, 217–232 (1986)), that of Marck (*Nucleic Acids Research* 16, 1829–1836 (1988)) or the GCG program of Butler (*Methods of Biochemical Analysis* 39, 74–97 (1998)).

The novel DNA sequence of *C. glutamicum* coding for the *pknB* gene was found and, as SEQ ID No. 1, forms part of the present invention. Furthermore, the amino acid sequence of the corresponding protein was derived from the DNA

sequence by the methods described above. The resulting amino acid sequence of the *pknB* gene product is shown in SEQ ID No. 2.

Coding DNA sequences which result from SEQ ID No. 1 due to the degeneracy of the genetic code also within the scope of the present invention. Likewise, DNA sequences which hybridize with SEQ ID No. 1 or portions of SEQ ID No. 1 form part of the invention. Furthermore, conservative amino acid exchanges, e.g. the exchange of glycine for alanine or of aspartic acid for glutamic acid in proteins, are known to those skilled in the art as "sense mutations", which do not cause a fundamental change in the activity of the protein, i.e. they are neutral. It is also known that changes at the N- and/or C-terminus of a protein do not substantially impair its function or may even stabilize it. Those skilled in the art will find information on this subject in Ben-Bassat et al. (Journal of Bacteriology 169, 751-757 (1987)), O'Regan et al. (Gene 77, 237-251 (1989)), Sahin-Toth et al. (Protein Sciences 3, 240-247 (1994)) and Hochuli et al. (Bio/Technology 6, 1321-1325 (1988)), inter alia, and in well-known textbooks on genetics and molecular biology. Amino acid sequences which correspondingly result from SEQ ID No. 2 also form part of the invention.

Likewise, DNA sequences which hybridize with SEQ ID No. 1 or portions of SEQ ID No. 1 are within the scope of the present invention. Finally, DNA sequences which are prepared by the polymerase chain reaction (PCR) using primers resulting from SEQ ID No. 1 form part of the invention. Such oligonucleotides typically have a length of at least 15 nucleotides.

Those skilled in the art can find instructions on the identification of DNA sequences by means of hybridization in the manual entitled "The DIG System User's Guide for Filter Hybridization" from Boehringer Mannheim GmbH (Mannheim, Germany, 1993) and in Liebl et al. (International Journal of Systematic Bacteriology (1991) 41, 255-260), inter alia. Hybridization takes place under stringent conditions; in other words, only hybrids for which the probe and the target sequence, i.e. the polynucleotides treated with the probe, are at least 70% identical are formed. It is known that the stringency of hybridization, including the washing steps, is influenced or determined by varying the buffer composition, the temperature and the salt concentration. The hybridization reaction is preferably carried out under relatively low stringency compared with the washing steps (Hybaid Hybridisation Guide, Hybaid Limited, Teddington, UK, 1996).

The hybridization reaction can be carried out for example using a 5× SSC buffer at a temperature of approx. 50° C.-68° C., it also being possible for probes to hybridize with polynucleotides which are less than 70% identical to the sequence of the probe. Such hybrids are less stable and are removed by washing under stringent conditions. This can be achieved for example by lowering the salt concentration to 2× SSC and subsequently to 0.5× SSC if necessary (The DIG System User's Guide for Filter Hybridization, Boehringer Mannheim, Mannheim, Germany, 1995), the temperature being adjusted to approx. 50° C.-68° C. It is possible to lower the salt concentration to 0.1× SSC if necessary. By raising the hybridization temperature in approx. 1-2° C. steps from 50° C. to 68° C., it is possible to isolate polynucleotide fragments which are e.g. at least 70%, at least 80% or at least 90% to 95% identical to the sequence of the probe used. Further instructions on hybridization are commercially available in the form of kits (e.g. DIG Easy Hyb from Roche Diagnostics GmbH, Mannheim, Germany, Catalog No. 1603558).

Those skilled in the art can find instructions on the amplification of DNA sequences with the aid of the polymerase chain reaction (PCR) in the manual by Gait entitled Oligonucleotide Synthesis: A Practical Approach (IRL Press, Oxford, UK, 1984) and in Newton and Graham: PCR (Spektrum Akademischer Verlag, Heidelberg, Germany, 1994), inter alia.

It has been found that, after overexpression of the *pknB* gene, the production of amino acids by corynebacteria is improved.

Overexpression can be achieved by increasing the copy number of the appropriate genes or mutating the promoter and regulatory region or the ribosome binding site located upstream from the structural gene. Expression cassettes incorporated upstream from the structural gene work in the same way. Inducible promoters additionally make it possible to increase the expression in the course of the production of amino acid by fermentation. Measures for prolonging the life of the mRNA also improve the expression. Furthermore, the enzyme activity is also enhanced by preventing the degradation of the enzyme protein. The genes or gene constructs can either be located in plasmids of variable copy number or integrated and amplified in the chromosome. Alternatively, it is also possible to achieve overexpression of the genes in question by changing the composition of the media and the culture technique.

Those skilled in the art can find relevant instructions in Martin et al. (Bio/Technology 5, 137-146 (1987)), Guerrero et al. (Gene 138, 35-41 (1994)), Tsuchiya and Morinaga (Bio/Technology 6, 428-430 (1988)), Eikmanns et al. (Gene 102, 93-98 (1991)), EP 0 472 869, U.S. Pat. No. 4,601,893, Schwarzer and Puhler (Bio/Technology 9, 84-87 (1991)), Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)), LaBarre et al. (Journal of Bacteriology 175, 1001-1007 (1993)), WO 96/15246, Malumbres et al. (Gene 134, 15-24 (1993)), JP-A-10-229891, Jensen and Hammer (Biotechnology and Bioengineering 58, 191-195 (1998)) and Makrides (Microbiological Reviews 60, 512-538 (1996)), inter alia, and in well-known textbooks on genetics and molecular biology.

For amplification, the *pknB* gene according to the invention has been overexpressed for example with the aid of episomal plasmids. Suitable plasmids are those which are replicated in corynebacteria. Numerous known plasmid vectors, e.g. pZ1 (Menkel et al., Applied and Environmental Microbiology (1989) 64, 549-554), pEKEx1 (Eikmanns et al., Gene 102, 93-98 (1991)) or pHS2-1 (Sonnen et al., Gene 107, 69-74 (1991)), are based on cryptic plasmids pHM1519, pBL1 or pGA1. Other plasmid vectors, e.g. those based on pCG4 (U.S. Pat. No. 4,489,160), pNG2 (Serwold-Davis et al., FEMS Microbiology Letters 66, 119-124 (1990)) or pAG1 (U.S. Pat. No. 5,158,891), can be used in the same way.

Other suitable plasmid vectors are those which make it possible to use the gene amplification process by integration into the chromosome, as described for example by Reinscheid et al. (Applied and Environmental Microbiology 60, 126-132 (1994)) for the duplication or amplification of the *hom-thrB* operon. In this method the complete gene is cloned into a plasmid vector which can replicate in a host (typically *E. coli*), but not in *C. glutamicum*. Examples of suitable vectors are pSUP301 (Simon et al., Bio/Technology 1, 784-791 (1983)), pK18mob or pK19mob (Schäfer et al., Gene 145, 69-73 (1994)), pGEM-T (Promega Corporation, Madison, Wis., USA), pCR2.1-TOPO (Shuman (1994), Journal of Biological Chemistry 269, 32678-84; U.S. Pat. No. 5,487,993), pCR@Blunt (Invitrogen, Groningen, The

Netherlands; Bernard et al., *Journal of Molecular Biology* 234, 534–541 (1993)), pEM1 (Schrumpf et al., 1991, *Journal of Bacteriology* 173, 4510–4516) or pBGS8 (Spratt et al., 1986, *Gene* 41, 337–342). The plasmid vector containing the gene to be amplified is then transferred to the desired strain of *C. glutamicum* by conjugation or transformation. The method of conjugation is described for example in Schäfer et al. (*Applied and Environmental Microbiology* 60, 756–759 (1994)). Methods of transformation are described for example in Thierbach et al. (*Applied Microbiology and Biotechnology* 29, 356–362 (1988)), Dunican and Shivnan (*Bio/Technology* 7, 1067–1070 (1989)) and Tauch et al. (*FEMS Microbiological Letters* 123, 343–347 (1994)). After homologous recombination by means of a crossover event, the resulting strain contains at least two copies of the gene in question.

It has also been found that amino acid exchanges in the section between position 581 and position 587 of the amino acid sequence of protein kinase B, shown in SEQ ID No. 2, improve the production of amino acids, especially lysine, by corynebacteria. Preferably, L-proline in position 584 is exchanged for any other proteogenic amino acid except L-proline, preferably for L-serine or L-threonine and very particularly preferably for L-serine.

SEQ ID No.3 shows the base sequence of the *pknB*-1547 allele contained in the strain DM1547. The *pknB*-1547 allele codes for a protein whose amino acid sequence is shown in SEQ ID No. 4. The protein contains L-serine in position 584. The DNA sequence of the *pknB*-1547 allele (SEQ ID No. 3) contains the base thymine in place of the base cytosine contained in the *pknB* wild-type gene (SEQ ID No. 1) in position 2343.

Mutagenesis can be carried out by conventional methods using mutagenic substances such as N-methyl-N'-nitro-N-nitrosoguanidine or ultraviolet light. Mutagenesis can also be carried out using in vitro methods such as treatment with hydroxylamine (Miller, J. H.: *A Short Course in Bacterial Genetics. A Laboratory Manual and Handbook for Escherichia coli and Related Bacteria*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, 1992) or mutagenic oligonucleotides (T. A. Brown: *Gentechnologie für Einsteiger (Gene Technology for Beginners)*, Spektrum Akademischer Verlag, Heidelberg, 1993), or the polymerase chain reaction (PCR) as described in the manual by Newton and Graham (PCR, Spektrum Akademischer Verlag, Heidelberg, 1994).

The corresponding alleles or mutations are sequenced and introduced into the chromosome by the method of gene replacement, for example as described in Peters-Wendisch et al. (*Microbiology* 144, 915–927 (1998)) for the *pyc* gene of *C. glutamicum*, in Schäfer et al. (*Gene* 145, 69–73 (1994)) for the *hom-thrB* gene region of *C. glutamicum* or in Schäfer et al. (*Journal of Bacteriology* 176, 7309–7319 (1994)) for the *cg1* gene region of *C. glutamicum*. The corresponding alleles or the associated proteins can optionally be amplified in turn.

In addition it can be advantageous for the production of L-amino acids to amplify and, in particular, overexpress not only the *pknB* gene but also one or more enzymes of the particular biosynthetic pathway, the glycolysis, the anaplerosis, the citric acid cycle, the pentose phosphate cycle or the amino acid export, and optionally regulatory proteins.

Thus, for the production of L-amino acids, one or more genes selected from the following group can be amplified and, in particular, overexpressed in addition to amplification of the endogene *pknB* gene: the *dapA* gene coding for dihydrodipicolinate synthase (EP-B-0 197 335),

the *gap* gene coding for glyceraldehyde 3-phosphate dehydrogenase (Eikmanns (1992), *Journal of Bacteriology* 174, 6076–6086),

the *tpi* gene coding for triose phosphate isomerase (Eikmanns (1992), *Journal of Bacteriology* 174, 6076–6086),

the *pgk* gene coding for 3-phosphoglycerate kinase (Eikmanns (1992), *Journal of Bacteriology* 174, 6076–6086),

the *zwf* gene coding for glucose-6-phosphate dehydrogenase (JP-A-09224661),

the *pyc* gene coding for pyruvate carboxylase (DE-A-198 31 609),

the *lysC* gene coding for a feedback-resistant aspartate kinase (Accession no. P26512; EP-B-0387527; EP-A-0699759),

the *lysE* gene coding for lysine export (DE-A-195 48 222),

the *hom* gene coding for homoserine dehydrogenase (EP-A-0131171),

the *ilvA* gene coding for threonine dehydratase (Möckel et al., *Journal of Bacteriology* (1992) 8065–8072) or the *ilvA(Fbr)* allele coding for a feedback-resistant threonine dehydratase (Mockel et al. (1994), *Molecular Microbiology* 13, 833–842),

the *ilvBN* gene coding for acetohydroxy acid synthase (EP-B-0356739),

the *ilvD* gene coding for dihydroxy acid dehydratase (Sahm and Eggeling (1999), *Applied and Environmental Microbiology* 65, 1973–1979), or

the *zwa1* gene coding for the Zwa1 protein (DE 199 59 328.0, DSM13115).

In addition to amplification of the *pknB* gene, it can also be advantageous for the production of L-amino acids to attenuate one or more genes selected from the following group:

the *pck* gene coding for phosphoenol pyruvate carboxykinase (DE 199 50 409.1, DSM13047),

the *pgi* gene coding for glucose-6-phosphate isomerase (U.S. Ser. No. 09/396,478, DSM12969),

the *poxB* gene coding for pyruvate oxidase (DE 199 51 975.7, DSM13114), or

the *zwa2* gene coding for the Zwa2 protein (DE 199 59 327.2, DSM13113),

and, in particular, to reduce the expression.

In this context the term “attenuation” describes the reduction or switching-off of the intracellular activity, in a microorganism, of one or more enzymes (proteins) which are coded for by the appropriate DNA, for example by using a weak promoter or using a gene or allele coding for an appropriate enzyme with a low activity, or inactivating the appropriate gene or enzyme (protein), and optionally combining these measures.

By attenuation measures, the activity or concentration of the corresponding protein is in general reduced to 0 to 50%, 0 to 25%, 0 to 10% or 0 to 5% of the activity or concentration of the wild-type protein.

It can also be advantageous for the production of amino acids not only to overexpress the *pknB* gene but also to switch off unwanted secondary reactions (Nakayama: “Breeding of Amino Acid Producing Micro-organisms”, in: *Overproduction of Microbial Products*, Krumphanzl, Sikyta, Vanek (eds.), Academic Press, London, UK, 1982).

The microorganisms prepared according to the invention are also provided by the invention and can be cultivated for

the production of amino acids continuously or discontinuously by the batch process, the fed batch process or the repeated fed batch process. A summary of known cultivation methods is described in the textbook by Chmiel (Bioprozesstechnik 1. Ein uung in die Bioverfahrenstechnik (Bioprocess Technology 1. Introduction to Bioengineering) (Gustav Fischer Verlag, Stuttgart, 1991)) or in the textbook by Storhas (Bioreaktoren und periphere Einrichtungen (Bioreactors and Peripheral Equipment) (Vieweg Verlag, Brunswick/Wiesbaden, 1994)).

The culture medium to be used must appropriately meet the demands of the particular strains. Descriptions of culture media for various microorganisms can be found in "Manual of Methods for General Bacteriology" of the American Society for Bacteriology (Washington D.C., USA, 1981).

Carbon sources which can be used are sugars and carbohydrates, e.g. glucose, sucrose, lactose, fructose, maltose, molasses, starch and cellulose, oils and fats, e.g. soybean oil, sunflower oil, groundnut oil and coconut fat, fatty acids, e.g. palmitic acid, stearic acid and linoleic acid, alcohols, e.g. glycerol and ethanol, and organic acids, e.g. acetic acid. These substances can be used individually or as a mixture.

Nitrogen sources which can be used are organic nitrogen-containing compounds such as peptones, yeast extract, meat extract, malt extract, corn steep liquor, soybean flour and urea, or inorganic compounds such as ammonium sulfate, ammonium chloride, ammonium phosphate, ammonium carbonate and ammonium nitrate. The nitrogen sources can be used individually or as a mixture.

Phosphorus sources which can be used are phosphoric acid, potassium dihydrogenphosphate or dipotassium hydrogenphosphate or the corresponding sodium salts. The culture medium must also contain metal salts, e.g. magnesium sulfate or iron sulfate, which are necessary for growth. Finally, essential growth-promoting substances such as amino acids and vitamins can be used in addition to the substances mentioned above. Suitable precursors can also be added to the culture medium. The feed materials can be added to the culture all at once or fed in appropriately during cultivation.

The pH of the culture is controlled by the appropriate use of basic compounds such as sodium hydroxide, potassium hydroxide, ammonia or aqueous ammonia, or acidic compounds such as phosphoric acid or sulfuric acid. Foaming can be controlled using antifoams such as fatty acid polyglycol esters. The stability of plasmids can be maintained by adding suitable selectively acting substances, e.g. antibiotics, to the medium. Aerobic conditions are maintained by introducing oxygen or oxygen-containing gaseous mixtures, e.g. air, into the culture. The temperature of the culture is normally 20° C. to 45° C. and preferably 25° C. to 40° C. The culture is continued until the formation of the desired product has reached a maximum. This objective is normally achieved within 10 hours to 160 hours.

Methods of determining L-amino acids are well-known to those skilled in the art. They can be analyzed for example by ion exchange chromatography with subsequent ninhydrin derivation, as described by Spackman et al. (Analytical Chemistry 30 (1958) 1190), or by reversed phase HPLC, as described by Lindroth et al. (Analytical Chemistry (1979) 51, 1167–1174).

A pure culture of the *Corynebacterium glutamicum* strain DM1547 was deposited as DSM13994 in the Deutsche Sammlung für Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell Cultures (DSMZ), Brunswick, Germany) on Jan. 16, 2001 under the terms of the Budapest Treaty.

The fermentation process according to the invention is used for the preparation of amino acids.

Another aspect of the present invention is a method of identifying nucleic acids which code for protein kinase B or have a high degree of similarity to the sequence of the pknB gene, comprising:

contacting a sample with the polynucleotide of claim 1 under conditions suitable for the polynucleotide to hybridize to another nucleic acid which codes for protein kinase B or have a high degree of similarity to the sequence of the pknB gene. In one embodiment, the another nucleic acid referred to above is RNA, cDNA, or DNA. In another embodiment, the method is conducted on an array, micro-array, or DNA chip.

## EXAMPLES

Having generally described this invention, a further understanding can be obtained by reference to certain specific examples which are provided herein for purposes of illustration only and are not intended to be limiting unless otherwise specified.

The isolation of plasmid DNA from *Escherichia coli* and all the techniques of restriction, Klenow treatment and alkaline phosphatase treatment were carried out according to Sambrook et al. (Molecular Cloning. A Laboratory Manual (1989), Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., USA). Methods of transforming *Escherichia coli* are also described in this manual.

The composition of common nutrient media, such as LB or TY medium, can also be found in the manual by Sambrook et al.

### Example 1

#### Preparation of a Genomic Cosmid Gene Library from *Corynebacterium glutamicum* ATCC13032

Chromosomal DNA from *Corynebacterium glutamicum* ATCC13032 was isolated as described in Tauch et al. (1995, Plasmid 33, 168–179) and partially cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, product description Sau3AI, code no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, product description SAP, code no. 1758250). The DNA of cosmid vector SuperCos1 (Wahl et al. (1987), Proceedings of the National Academy of Sciences USA 84, 2160–2164), obtained from Stratagene (La Jolla, USA, product description SuperCos1 Cosmid Vector Kit, code no. 251301), was cleaved with the restriction enzyme XbaI (Amersham Pharmacia, Freiburg, Germany, product description XbaI, code no. 27-0948-02) and also dephosphorylated with shrimp alkaline phosphatase.

The cosmid DNA was then cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, product description BamHI, code no. 27-0868-04). The cosmid DNA treated in this way was mixed with the treated ATCC13032 DNA and the mixture was treated with T4 DNA ligase (Amersham Pharmacia, Freiburg, Germany, product description T4 DNA ligase, code no. 27-0870-04). The ligation mixture was then packaged into phages using Gigapack II XL Packing Extract (Stratagene, La Jolla, USA, product description Gigapack II XL Packing Extract, code no. 200217).

For infection of the *E. coli* strain NM554 (Raleigh et al., 1988, Nucleic Acid Research 16, 1563–1575), the cells were taken up in 10 mM MgSO<sub>4</sub> and mixed with an aliquot of the

## 11

phage suspension. Infection and titering of the cosmid library were carried out as described in Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the cells being plated on LB agar (Lennox, 1955, Virology 1, 190) containing 100 mg/l of ampicillin. After incubation overnight at 37° C., recombinant single clones were selected.

## Example 2

## Isolation and Sequencing of the pknB Gene

The cosmid DNA of a single colony was isolated with the Qiaprep Spin Miniprep Kit (product no. 27106, Qiagen, Hilden, Germany) in accordance with the manufacturer's instructions and partially cleaved with the restriction enzyme Sau3AI (Amersham Pharmacia, Freiburg, Germany, product description Sau3AI, product no. 27-0913-02). The DNA fragments were dephosphorylated with shrimp alkaline phosphatase (Roche Diagnostics GmbH, Mannheim, Germany, product description SAP, product no. 1758250). After separation by gel electrophoresis, the cosmid fragments in the size range from 1500 to 2000 bp were isolated with the QiaExII Gel Extraction Kit (product no. 20021, Qiagen, Hilden, Germany).

The DNA of sequencing vector pZero-1, obtained from Invitrogen (Groningen, The Netherlands, product description Zero Background Cloning Kit, product no. K2500-01), was cleaved with the restriction enzyme BamHI (Amersham Pharmacia, Freiburg, Germany, product description BamHI, product no. 27-0868-04). Ligation of the cosmid fragments into sequencing vector pZero-1 was carried out as described in Sambrook et al. (1989, Molecular Cloning: A Laboratory Manual, Cold Spring Harbor), the DNA mixture being incubated overnight with T4 ligase (Pharmacia Biotech, Freiburg, Germany). This ligation mixture was then introduced into the *E. coli* strain DH5aMCR (Grant, 1990, Proceedings of the National Academy of Sciences USA 87, 4645-4649) by electroporation (Tauch et al. 1994, FEMS Microbiol. Letters 123, 343-7) and plated on LB agar (Lennox, 1955, Virology 1, 190) containing 50 mg/l of zeocin.

## 12

Plasmid preparation of the recombinant clones was carried out with Biorobot 9600 (product no. 900200, Qiagen, Hilden, Germany). Sequencing was carried out by the dideoxy chain termination method of Sanger et al. (1977, Proceedings of the National Academy of Sciences USA 74, 5463-5467) with modifications by Zimmermann et al. (1990, Nucleic Acids Research 18, 1067). The "RR dRhodamin Terminator Cycle Sequencing Kit" from PE Applied Biosystems (product no. 403044, Weiterstadt, Germany) was used. Separation by gel electrophoresis and analysis of the sequencing reaction were carried out in a "Rotiphorese NF acrylamide/bisacrylamide" gel (29:1) (product no. A124.1, Roth, Karlsruhe, Germany) with the "ABI Prism 377" sequencer from PE Applied Biosystems (Weiterstadt, Germany).

The raw sequence data obtained were then processed using the Staden programming package (1986, Nucleic Acids Research 14, 217-231), version 97-0. The individual sequences of the pZero-1 derivatives were assembled into a cohesive contig. Computer-assisted coding region analysis was performed with the XNIP program (Staden, 1986, Nucleic Acids Research 14, 217-231).

The nucleotide sequence obtained is shown in SEQ ID No. 1. Analysis of the nucleotide sequence gave an open reading frame of 1884 base pairs, which was called the pknB gene. The pknB gene codes for a protein of 627 amino acids.

Obviously, numerous modifications and variations of the present invention are possible in light of the above teachings. It is therefore to be understood that within the scope of the appended claims, the invention may be practiced otherwise than as specifically described herein.

All of the publications cited above are incorporated herein by reference.

This application is based, inter alia, on German Patent Application Serial No.100 44 912.3, filed on Sep. 12, 2000; and German Patent Application Serial No. 101 20 095.1, filed on Apr. 25, 2001, both of which are incorporated herein by reference.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 4

<210> SEQ ID NO 1  
 <211> LENGTH: 2875  
 <212> TYPE: DNA  
 <213> ORGANISM: Corynebacterium glutamicum  
 <220> FEATURE:  
 <221> NAME/KEY: CDS  
 <222> LOCATION: (594)..(2474)  
 <223> OTHER INFORMATION:

<400> SEQUENCE: 1

```

cgcacgagcg cgatgatggc gcaggcggag gcgccgtcgc caagcgaatc aacggcgatg      60
ctgggcaggg tggcccggcc tgcaacaatc acccaagaag cggccccgaa acgcggttcc      120
ggcattggca ttggtctgtt catcgcagct ttgcttgccg tgattattgg cgcggtgatc      180
tatgcgggca ccaccggaat tttgttcaac gacactccgg aagaaaccac cacacctgaa      240
accattacgg aaacatacac cccaaccgtg gaggaaacca cctctcagtg ggtaccgcca      300
acgcctcaa caggtcaac attcaccgaa cctgaaacaa cttcacaccg tccgacgaca      360

```

-continued

agtgaagaga gcacatccga ggaaccaacc acggaagctc caacaagtag ccgaactgtg	420
cctcaaatcc ctacctctac acctaggacg agtgctagcg ttccagttga gactaatgca	480
ccggctgatg atttaatcga cgccgtaaat ggcctattgg atgtaggagg agcgcagtga	540
ccttcgtgat cgctgatcgc tatgaactgg atgccgtcat cggctccggt ggc atg	596
	Met
	1
agc gag gtg ttc gcg gcc acc gac acg ctc att ggt cgg gag gtc gcg	644
Ser Glu Val Phe Ala Ala Thr Asp Thr Leu Ile Gly Arg Glu Val Ala	
	5 10 15
gta aag atg ctg cgc atc gac ctt gcg aaa gat ccc aat ttc cga gaa	692
Val Lys Met Leu Arg Ile Asp Leu Ala Lys Asp Pro Asn Phe Arg Glu	
	20 25 30
cgc ttc cgc agg gaa gcc caa aac tcc gga agg ttg agc cac tct tcg	740
Arg Phe Arg Arg Glu Ala Gln Asn Ser Gly Arg Leu Ser His Ser Ser	
	35 40 45
atc gtc gct gtt ttt gac acc ggc gaa gta gac aaa gac ggc acc tct	788
Ile Val Ala Val Phe Asp Thr Gly Glu Val Asp Lys Asp Gly Thr Ser	
	50 55 60 65
gtt ccc tac att gtg atg gaa cgc gtg cag ggt cga aac ctg cgc gaa	836
Val Pro Tyr Ile Val Met Glu Arg Val Gln Gly Arg Asn Leu Arg Glu	
	70 75 80
gtt gtc acc gaa gac ggc gta ttc acc cca gtt gag gca gcc aac atc	884
Val Val Thr Glu Asp Gly Val Phe Thr Pro Val Glu Ala Ala Asn Ile	
	85 90 95
ctc atc cct gtg tgt gaa gcg ctg cag gca tcc cat gac gcc ggc att	932
Leu Ile Pro Val Cys Glu Ala Leu Gln Ala Ser His Asp Ala Gly Ile	
	100 105 110
att cac cgc gat gtg aaa ccc gcc aac atc atg atc acc aac acc ggt	980
Ile His Arg Asp Val Lys Pro Ala Asn Ile Met Ile Thr Asn Thr Gly	
	115 120 125
ggc gtg aaa gtc atg gac ttc ggc atc gcc cgc gcg gtc aac gat tcc	1028
Gly Val Lys Val Met Asp Phe Gly Ile Ala Arg Ala Val Asn Asp Ser	
	130 135 140 145
acc tcc gcc atg act caa acc tcc gca gtc atc ggc acc gcc cag tac	1076
Thr Ser Ala Met Thr Gln Thr Ser Ala Val Ile Gly Thr Ala Gln Tyr	
	150 155 160
ctc tcc cct gag cag gcc cgc ggc aaa ccc gcc gat gcg cgt tcc gat	1124
Leu Ser Pro Glu Gln Ala Arg Gly Lys Pro Ala Asp Ala Arg Ser Asp	
	165 170 175
att tac gcc acc ggc tgc gtc atg tac gaa tta gtc acc ggt aag cca	1172
Ile Tyr Ala Thr Gly Cys Val Met Tyr Glu Leu Val Thr Gly Lys Pro	
	180 185 190
cct ttt gaa ggc gag tcc cct ttc gcc gtg gcc tac caa cac gtc cag	1220
Pro Phe Glu Gly Glu Ser Pro Phe Ala Val Ala Tyr Gln His Val Gln	
	195 200 205
gaa gac ccc acc cct cct tcg gat ttc atc gcg gac ctc acc ccg acc	1268
Glu Asp Pro Thr Pro Pro Ser Asp Phe Ile Ala Asp Leu Thr Pro Thr	
	210 215 220 225
tct gct gtc aac gtg gat gcc gtg gta ctc acc gcc atg gca aaa cac	1316
Ser Ala Val Asn Val Asp Ala Val Val Leu Thr Ala Met Ala Lys His	
	230 235 240
ccc gcc gac cgc tac caa aca gcc tcc gaa atg gcc gct gac ctg ggc	1364
Pro Ala Asp Arg Tyr Gln Thr Ala Ser Glu Met Ala Ala Asp Leu Gly	
	245 250 255
cgg cta tcc cgc aat gca gtc tcc cat gcc gca cgc gcg cat gta gaa	1412
Arg Leu Ser Arg Asn Ala Val Ser His Ala Ala Arg Ala His Val Glu	
	260 265 270
aca gaa gaa acc cca gaa gag ccc gaa act cgc ttc tcg acg cgc acc	1460



-continued

Thr	Glu	Glu	Thr	Pro	Glu	Glu	Pro	Glu	Thr	Arg	Phe	Ser	Thr	Arg	Thr		
	275					280					285						
tcc	acc	caa	gtg	gcc	ccc	gcc	gca	ggc	gtg	gct	gcg	gcc	agt	acg	ggg	1508	
Ser	Thr	Gln	Val	Ala	Pro	Ala	Ala	Gly	Val	Ala	Ala	Ala	Ser	Thr	Gly		
	290				295				300						305		
tca	ggg	tct	tct	tcg	cgt	aaa	cgt	gga	tcc	aga	ggc	ctc	acc	gcc	ctg	1556	
Ser	Gly	Ser	Ser	Ser	Arg	Lys	Arg	Gly	Ser	Arg	Gly	Leu	Thr	Ala	Leu		
				310					315					320			
gcc	atc	gtg	tta	tcc	cta	ggt	gtc	gtc	ggc	ggt	gcc	ggt	gcc	ttc	acc	1604	
Ala	Ile	Val	Leu	Ser	Leu	Gly	Val	Val	Gly	Val	Ala	Gly	Ala	Phe	Thr		
			325					330					335				
tac	gac	tac	ttt	gcc	aac	agc	tcc	tcc	act	gca	acc	agc	gcg	atc	ccc	1652	
Tyr	Asp	Tyr	Phe	Ala	Asn	Ser	Ser	Ser	Thr	Ala	Thr	Ser	Ala	Ile	Pro		
		340					345						350				
aat	gtg	gaa	ggc	ctc	ccg	cag	caa	gaa	gct	ctc	aca	gaa	ctt	caa	gca	1700	
Asn	Val	Glu	Gly	Leu	Pro	Gln	Gln	Glu	Ala	Leu	Thr	Glu	Leu	Gln	Ala		
	355					360					365						
gca	gga	ttt	ggt	gtc	aac	atc	gtc	gaa	gaa	gcc	agc	gcc	gac	gtc	gcc	1748	
Ala	Gly	Phe	Val	Val	Asn	Ile	Val	Glu	Glu	Ala	Ser	Ala	Asp	Val	Ala		
	370				375					380					385		
gaa	ggc	ctc	gtc	atc	cga	gca	aac	cca	agc	ggt	gga	tcc	gaa	atc	cgc	1796	
Glu	Gly	Leu	Val	Ile	Arg	Ala	Asn	Pro	Ser	Val	Gly	Ser	Glu	Ile	Arg		
			390						395					400			
caa	ggg	gcc	acc	gtc	acc	atc	acc	gtg	tcc	acc	ggc	cga	gaa	atg	atc	1844	
Gln	Gly	Ala	Thr	Val	Thr	Ile	Thr	Val	Ser	Thr	Gly	Arg	Glu	Met	Ile		
			405					410						415			
aac	atc	cca	gac	gtc	tcc	ggc	atg	aca	ctt	gag	gac	gcc	gcc	cgc	gcc	1892	
Asn	Ile	Pro	Asp	Val	Ser	Gly	Met	Thr	Leu	Glu	Asp	Ala	Ala	Arg	Ala		
		420					425					430					
ctc	gaa	gac	ggt	ggt	ctc	ata	ctc	aac	caa	aac	ggt	cgg	gaa	gaa	acc	1940	
Leu	Glu	Asp	Val	Gly	Leu	Ile	Leu	Asn	Gln	Asn	Val	Arg	Glu	Glu	Thr		
	435					440					445						
tcc	gac	gac	gtc	gaa	tct	ggc	ctc	gtc	atc	gac	caa	aac	ccc	gaa	gcc	1988	
Ser	Asp	Asp	Val	Glu	Ser	Gly	Leu	Val	Ile	Asp	Gln	Asn	Pro	Glu	Ala		
	450				455					460					465		
ggc	caa	gaa	gta	gtc	gtg	ggt	tcc	tct	gta	tct	cta	acc	atg	tct	tca	2036	
Gly	Gln	Glu	Val	Val	Val	Gly	Ser	Ser	Val	Ser	Leu	Thr	Met	Ser	Ser		
			470						475					480			
ggc	acc	gag	agc	atc	cga	gtg	ccc	aac	ctc	acc	ggc	atg	aac	tgg	tca	2084	
Gly	Thr	Glu	Ser	Ile	Arg	Val	Pro	Asn	Leu	Thr	Gly	Met	Asn	Trp	Ser		
		485					490						495				
caa	gca	gaa	caa	aac	ctc	atc	tcc	atg	ggc	ttt	aac	ccc	aca	gct	tcc	2132	
Gln	Ala	Glu	Gln	Asn	Leu	Ile	Ser	Met	Gly	Phe	Asn	Pro	Thr	Ala	Ser		
		500					505					510					
tac	tta	gac	agc	agc	gaa	cca	gaa	ggc	gaa	gtc	ctc	tca	ggt	tcc	agc	2180	
Tyr	Leu	Asp	Ser	Ser	Glu	Pro	Glu	Gly	Glu	Val	Leu	Ser	Val	Ser	Ser		
	515					520						525					
caa	gga	act	gaa	cta	ccc	aag	ggt	tca	tcc	atc	aca	gtg	gaa	gtc	tcc	2228	
Gln	Gly	Thr	Glu	Leu	Pro	Lys	Gly	Ser	Ser	Ile	Thr	Val	Glu	Val	Ser		
	530				535					540				545			
aac	ggc	atg	ctc	atc	caa	gcc	ccc	gat	ctc	gcc	cgc	atg	tcc	acc	gaa	2276	
Asn	Gly	Met	Leu	Ile	Gln	Ala	Pro	Asp	Leu	Ala	Arg	Met	Ser	Thr	Glu		
			550						555					560			
cag	gcc	atc	agt	gcc	ctc	cgc	gct	gct	ggc	tgg	acc	gcc	cca	gat	caa	2324	
Gln	Ala	Ile	Ser	Ala	Leu	Arg	Ala	Ala	Gly	Trp	Thr	Ala	Pro	Asp	Gln		
			565					570					575				
tcc	ctg	atc	gtc	ggc	gac	ccc	atc	cac	acc	gca	gcc	ctc	gtg	gat	caa	2372	
Ser	Leu	Ile	Val	Gly	Asp	Pro	Ile	His	Thr	Ala	Ala	Leu	Val	Asp	Gln		
		580					585						590				



-continued

---

Gly Arg Leu Ser Arg Asn Ala Val Ser His Ala Ala Arg Ala His Val  
                   260                                  265                                  270  
  
 Glu Thr Glu Glu Thr Pro Glu Glu Pro Glu Thr Arg Phe Ser Thr Arg  
                   275                                  280                                  285  
  
 Thr Ser Thr Gln Val Ala Pro Ala Ala Gly Val Ala Ala Ala Ser Thr  
           290                                  295                                  300  
  
 Gly Ser Gly Ser Ser Ser Arg Lys Arg Gly Ser Arg Gly Leu Thr Ala  
 305                                  310                                  315                                  320  
  
 Leu Ala Ile Val Leu Ser Leu Gly Val Val Gly Val Ala Gly Ala Phe  
                   325                                  330                                  335  
  
 Thr Tyr Asp Tyr Phe Ala Asn Ser Ser Ser Thr Ala Thr Ser Ala Ile  
                   340                                  345                                  350  
  
 Pro Asn Val Glu Gly Leu Pro Gln Gln Glu Ala Leu Thr Glu Leu Gln  
                   355                                  360                                  365  
  
 Ala Ala Gly Phe Val Val Asn Ile Val Glu Glu Ala Ser Ala Asp Val  
           370                                  375                                  380  
  
 Ala Glu Gly Leu Val Ile Arg Ala Asn Pro Ser Val Gly Ser Glu Ile  
 385                                  390                                  395                                  400  
  
 Arg Gln Gly Ala Thr Val Thr Ile Thr Val Ser Thr Gly Arg Glu Met  
                   405                                  410                                  415  
  
 Ile Asn Ile Pro Asp Val Ser Gly Met Thr Leu Glu Asp Ala Ala Arg  
                   420                                  425                                  430  
  
 Ala Leu Glu Asp Val Gly Leu Ile Leu Asn Gln Asn Val Arg Glu Glu  
           435                                  440                                  445  
  
 Thr Ser Asp Asp Val Glu Ser Gly Leu Val Ile Asp Gln Asn Pro Glu  
           450                                  455                                  460  
  
 Ala Gly Gln Glu Val Val Val Gly Ser Ser Val Ser Leu Thr Met Ser  
 465                                  470                                  475                                  480  
  
 Ser Gly Thr Glu Ser Ile Arg Val Pro Asn Leu Thr Gly Met Asn Trp  
                   485                                  490                                  495  
  
 Ser Gln Ala Glu Gln Asn Leu Ile Ser Met Gly Phe Asn Pro Thr Ala  
                   500                                  505                                  510  
  
 Ser Tyr Leu Asp Ser Ser Glu Pro Glu Gly Glu Val Leu Ser Val Ser  
           515                                  520                                  525  
  
 Ser Gln Gly Thr Glu Leu Pro Lys Gly Ser Ser Ile Thr Val Glu Val  
           530                                  535                                  540  
  
 Ser Asn Gly Met Leu Ile Gln Ala Pro Asp Leu Ala Arg Met Ser Thr  
 545                                  550                                  555                                  560  
  
 Glu Gln Ala Ile Ser Ala Leu Arg Ala Ala Gly Trp Thr Ala Pro Asp  
                   565                                  570                                  575  
  
 Gln Ser Leu Ile Val Gly Asp Pro Ile His Thr Ala Ala Leu Val Asp  
           580                                  585                                  590  
  
 Gln Asn Lys Ile Gly Phe Gln Ser Pro Thr Pro Ala Thr Leu Phe Arg  
           595                                  600                                  605  
  
 Lys Asp Ala Gln Val Gln Val Arg Leu Phe Glu Phe Asp Leu Ala Ala  
 610                                  615                                  620  
  
 Leu Val Gln  
 625

<210> SEQ ID NO 3  
 <211> LENGTH: 2875  
 <212> TYPE: DNA  
 <213> ORGANISM: Corynebacterium glutamicum  
 <220> FEATURE:  
 <221> NAME/KEY: CDS

-continued

<222> LOCATION: (594)..(2474)  
 <223> OTHER INFORMATION:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (2343)..(2343)  
 <223> OTHER INFORMATION: C-T transition  
  
 <400> SEQUENCE: 3

cgcacgagcg cgatgatggc gcaggcggag gcgccgtcgc caagcgaatc aacggcgatg	60
ctgggcaggg tggcccggcc tgcaacaatc acccaagaag cggccccgaa acgcggttcc	120
ggcattggca ttggtctgtt catcgcagct ttgcttgccg tgattattgg cgcggtgatc	180
tatgcgggca ccaccggaat tttgttcaac gacactccgg aagaaaccac cacacctgaa	240
accattacgg aacatacac cccaaccgtg gagaaacca cctctcagtg ggtaccgcca	300
acgcctccaa cacggtcaac attcaccgaa cctgaaacaa cttcacaccg tccgacgaca	360
agtgaagaga gcacatccga ggaaccaacc acggaagctc caacaagtag ccgaactgtg	420
cctcaaatcc ctacctctac acctaggacg agtgctagcg ttccagttga gactaatgca	480
ccggctgatg atttaatcga cgccgtaaat ggcctattgg atgtaggagg agcgcagtga	540
ccttcgtgat cgctgatcgc tatgaactgg atgccgtcat cggctccggt ggc atg	596
	Met
	1
agc gag gtg ttc gcg gcc acc gac acg ctc att ggt cgg gag gtc gcg	644
Ser Glu Val Phe Ala Ala Thr Asp Thr Leu Ile Gly Arg Glu Val Ala	
	5 10 15
gta aag atg ctg cgc atc gac ctt gcg aaa gat ccc aat ttc cga gaa	692
Val Lys Met Leu Arg Ile Asp Leu Ala Lys Asp Pro Asn Phe Arg Glu	
	20 25 30
cgc ttc cgc agg gaa gcc caa aac tcc gga agg ttg agc cac tct tcg	740
Arg Phe Arg Arg Glu Ala Gln Asn Ser Gly Arg Leu Ser His Ser Ser	
	35 40 45
atc gtc gct gtt ttt gac acc ggc gaa gta gac aaa gac ggc acc tct	788
Ile Val Ala Val Phe Asp Thr Gly Glu Val Asp Lys Asp Gly Thr Ser	
	50 55 60 65
gtt ccc tac att gtg atg gaa cgc gtg cag ggt cga aac ctg cgc gaa	836
Val Pro Tyr Ile Val Met Glu Arg Val Gln Gly Arg Asn Leu Arg Glu	
	70 75 80
gtt gtc acc gaa gac ggc gta ttc acc cca gtt gag gca gcc aac atc	884
Val Val Thr Glu Asp Gly Val Phe Thr Pro Val Glu Ala Ala Asn Ile	
	85 90 95
ctc atc cct gtg tgt gaa gcg ctg cag gca tcc cat gac gcc ggc att	932
Leu Ile Pro Val Cys Glu Ala Leu Gln Ala Ser His Asp Ala Gly Ile	
	100 105 110
att cac cgc gat gtg aaa ccc gcc aac atc atg atc acc aac acc ggt	980
Ile His Arg Asp Val Lys Pro Ala Asn Ile Met Ile Thr Asn Thr Gly	
	115 120 125
ggc gtg aaa gtc atg gac ttc ggc atc gcc cgc gcg gtc aac gat tcc	1028
Gly Val Lys Val Met Asp Phe Gly Ile Ala Arg Ala Val Asn Asp Ser	
	130 135 140 145
acc tcc gcc atg act caa acc tcc gca gtc atc ggc acc gcc cag tac	1076
Thr Ser Ala Met Thr Gln Thr Ser Ala Val Ile Gly Thr Ala Gln Tyr	
	150 155 160
ctc tcc cct gag cag gcc cgc ggc aaa ccc gcc gat gcg cgt tcc gat	1124
Leu Ser Pro Glu Gln Ala Arg Gly Lys Pro Ala Asp Ala Arg Ser Asp	
	165 170 175
att tac gcc acc ggc tgc gtc atg tac gaa tta gtc acc ggt aag cca	1172
Ile Tyr Ala Thr Gly Cys Val Met Tyr Glu Leu Val Thr Gly Lys Pro	
	180 185 190
cct ttt gaa ggc gag tcc cct ttc gcc gtg gcc tac caa cac gtc cag	1220

-continued

Pro	Phe	Glu	Gly	Glu	Ser	Pro	Phe	Ala	Val	Ala	Tyr	Gln	His	Val	Gln	
	195					200					205					
gaa	gac	ccc	acc	cct	cct	tcg	gat	ttc	atc	gcg	gac	ctc	acc	ccg	acc	1268
Glu	Asp	Pro	Thr	Pro	Pro	Ser	Asp	Phe	Ile	Ala	Asp	Leu	Thr	Pro	Thr	
210					215					220					225	
tct	gct	gtc	aac	gtg	gat	gcc	gtg	gta	ctc	acc	gcc	atg	gca	aaa	cac	1316
Ser	Ala	Val	Asn	Val	Asp	Ala	Val	Val	Leu	Thr	Ala	Met	Ala	Lys	His	
				230					235					240		
ccc	gcc	gac	cgc	tac	caa	aca	gcc	tcc	gaa	atg	gcc	gct	gac	ctg	ggc	1364
Pro	Ala	Asp	Arg	Tyr	Gln	Thr	Ala	Ser	Glu	Met	Ala	Ala	Asp	Leu	Gly	
			245					250					255			
cgg	cta	tcc	cgc	aat	gca	gtc	tcc	cat	gcc	gca	cgc	gcg	cat	gta	gaa	1412
Arg	Leu	Ser	Arg	Asn	Ala	Val	Ser	His	Ala	Ala	Arg	Ala	His	Val	Glu	
		260					265					270				
aca	gaa	gaa	acc	cca	gaa	gag	ccc	gaa	act	cgc	ttc	tcg	acg	cgc	acc	1460
Thr	Glu	Glu	Thr	Pro	Glu	Glu	Pro	Glu	Thr	Arg	Phe	Ser	Thr	Arg	Thr	
		275					280				285					
tcc	acc	caa	gtg	gcc	ccc	gcc	gca	ggc	gtg	gct	gcg	gcc	agt	acg	ggg	1508
Ser	Thr	Gln	Val	Ala	Pro	Ala	Ala	Gly	Val	Ala	Ala	Ala	Ser	Thr	Gly	
290					295					300					305	
tca	ggg	tct	tct	tcg	cgt	aaa	cgt	gga	tcc	aga	ggc	ctc	acc	gcc	ctg	1556
Ser	Gly	Ser	Ser	Ser	Arg	Lys	Arg	Gly	Ser	Arg	Gly	Leu	Thr	Ala	Leu	
				310					315					320		
gcc	atc	gtg	tta	tcc	cta	ggt	gtc	gtc	ggc	ggt	gcc	ggt	gcc	ttc	acc	1604
Ala	Ile	Val	Leu	Ser	Leu	Gly	Val	Val	Gly	Val	Ala	Gly	Ala	Phe	Thr	
			325					330					335			
tac	gac	tac	ttt	gcc	aac	agc	tcc	tcc	act	gca	acc	agc	gcg	atc	ccc	1652
Tyr	Asp	Tyr	Phe	Ala	Asn	Ser	Ser	Ser	Thr	Ala	Thr	Ser	Ala	Ile	Pro	
		340					345					350				
aat	gtg	gaa	ggc	ctc	ccg	cag	caa	gaa	gct	ctc	aca	gaa	ctt	caa	gca	1700
Asn	Val	Glu	Gly	Leu	Pro	Gln	Gln	Glu	Ala	Leu	Thr	Glu	Leu	Gln	Ala	
		355					360				365					
gca	gga	ttt	ggt	gtc	aac	atc	gtc	gaa	gaa	gcc	agc	gcc	gac	gtc	gcc	1748
Ala	Gly	Phe	Val	Val	Asn	Ile	Val	Glu	Glu	Ala	Ser	Ala	Asp	Val	Ala	
370					375					380				385		
gaa	ggc	ctc	gtc	atc	cga	gca	aac	cca	agc	ggt	gga	tcc	gaa	atc	cgc	1796
Glu	Gly	Leu	Val	Ile	Arg	Ala	Asn	Pro	Ser	Val	Gly	Ser	Glu	Ile	Arg	
				390					395					400		
caa	ggg	gcc	acc	gtc	acc	atc	acc	gtg	tcc	acc	ggc	cga	gaa	atg	atc	1844
Gln	Gly	Ala	Thr	Val	Thr	Ile	Thr	Val	Ser	Thr	Gly	Arg	Glu	Met	Ile	
			405					410					415			
aac	atc	cca	gac	gtc	tcc	ggc	atg	aca	ctt	gag	gac	gcc	gcc	cgc	gcc	1892
Asn	Ile	Pro	Asp	Val	Ser	Gly	Met	Thr	Leu	Glu	Asp	Ala	Ala	Arg	Ala	
		420					425					430				
ctc	gaa	gac	ggt	ggt	ctc	ata	ctc	aac	caa	aac	ggt	cgg	gaa	gaa	acc	1940
Leu	Glu	Asp	Val	Gly	Leu	Ile	Leu	Asn	Gln	Asn	Val	Arg	Glu	Glu	Thr	
			435			440					445					
tcc	gac	gac	gtc	gaa	tct	ggc	ctc	gtc	atc	gac	caa	aac	ccc	gaa	gcc	1988
Ser	Asp	Asp	Val	Glu	Ser	Gly	Leu	Val	Ile	Asp	Gln	Asn	Pro	Glu	Ala	
450					455					460				465		
ggc	caa	gaa	gta	gtc	gtg	ggt	tcc	tct	gta	tct	cta	acc	atg	tct	tca	2036
Gly	Gln	Glu	Val	Val	Val	Gly	Ser	Ser	Val	Ser	Leu	Thr	Met	Ser	Ser	
				470					475					480		
ggc	acc	gag	agc	atc	cga	gtg	ccc	aac	ctc	acc	ggc	atg	aac	tgg	tca	2084
Gly	Thr	Glu	Ser	Ile	Arg	Val	Pro	Asn	Leu	Thr	Gly	Met	Asn	Trp	Ser	
			485				490						495			
caa	gca	gaa	caa	aac	ctc	atc	tcc	atg	ggc	ttt	aac	ccc	aca	gct	tcc	2132
Gln	Ala	Glu	Gln	Asn	Leu	Ile	Ser	Met	Gly	Phe	Asn	Pro	Thr	Ala	Ser	
		500					505					510				

-continued

tac tta gac agc agc gaa cca gaa ggc gaa gtc ctc tca gtt tcc agc	2180
Tyr Leu Asp Ser Ser Glu Pro Glu Gly Glu Val Leu Ser Val Ser Ser	
515 520 525	
caa gga act gaa cta ccc aag ggt tca tcc atc aca gtg gaa gtc tcc	2228
Gln Gly Thr Glu Leu Pro Lys Gly Ser Ser Ile Thr Val Glu Val Ser	
530 535 540 545	
aac ggc atg ctc atc caa gcc ccc gat ctc gcc cgc atg tcc acc gaa	2276
Asn Gly Met Leu Ile Gln Ala Pro Asp Leu Ala Arg Met Ser Thr Glu	
550 555 560	
cag gcc atc agt gcc ctc cgc gct gct ggc tgg acc gcc cca gat caa	2324
Gln Ala Ile Ser Ala Leu Arg Ala Ala Gly Trp Thr Ala Pro Asp Gln	
565 570 575	
tcc ctg atc gtc ggc gac tcc atc cac acc gca gcc ctc gtg gat caa	2372
Ser Leu Ile Val Gly Asp Ser Ile His Thr Ala Ala Leu Val Asp Gln	
580 585 590	
aac aaa atc gga ttc caa tcc cca acc cct gca acc ctc ttc cgc aaa	2420
Asn Lys Ile Gly Phe Gln Ser Pro Thr Pro Ala Thr Leu Phe Arg Lys	
595 600 605	
gac gcc caa gtg caa gtg cga ctc ttc gaa ttc gat ctc gct gca ctc	2468
Asp Ala Gln Val Gln Val Arg Leu Phe Glu Phe Asp Leu Ala Ala Leu	
610 615 620 625	
gtg caa tagccaacaa ggaaccgctc aaggtagctg gcccggaac tgatagctta	2524
Val Gln	
agctcaaaca agataagtac cagttgctgg ggtttttcca agacaataaa ttatgaaggt	2584
gtgaacaatg ccaaaggcaa gagtaactaa aaacgagacc gcaccggttt caagcaaccc	2644
aagcgcacaaac cgcaccccgg ttaagatcaa ttccgccgga accccaatgt ggtacaaggt	2704
catcatgttt gccttcatga tcgtcggcct agcctggttg atcattaact acctcgtggg	2764
cccacagatc ccattcatgg ctgatcttgg tgcatggaac tatggcatcg gcttcggtct	2824
gatgatcatc ggccactca tgaccatggg ttggcggtta tccttcaaaa a	2875

<210> SEQ ID NO 4  
 <211> LENGTH: 627  
 <212> TYPE: PRT  
 <213> ORGANISM: Corynebacterium glutamicum  
 <220> FEATURE:  
 <221> NAME/KEY: misc\_feature  
 <222> LOCATION: (2343)..(2343)  
 <223> OTHER INFORMATION: C-T transition

<400> SEQUENCE: 4

Met Ser Glu Val Phe Ala Ala Thr Asp Thr Leu Ile Gly Arg Glu Val	
1 5 10 15	
Ala Val Lys Met Leu Arg Ile Asp Leu Ala Lys Asp Pro Asn Phe Arg	
20 25 30	
Glu Arg Phe Arg Arg Glu Ala Gln Asn Ser Gly Arg Leu Ser His Ser	
35 40 45	
Ser Ile Val Ala Val Phe Asp Thr Gly Glu Val Asp Lys Asp Gly Thr	
50 55 60	
Ser Val Pro Tyr Ile Val Met Glu Arg Val Gln Gly Arg Asn Leu Arg	
65 70 75 80	
Glu Val Val Thr Glu Asp Gly Val Phe Thr Pro Val Glu Ala Ala Asn	
85 90 95	
Ile Leu Ile Pro Val Cys Glu Ala Leu Gln Ala Ser His Asp Ala Gly	
100 105 110	
Ile Ile His Arg Asp Val Lys Pro Ala Asn Ile Met Ile Thr Asn Thr	
115 120 125	

-continued

Gly	Gly	Val	Lys	Val	Met	Asp	Phe	Gly	Ile	Ala	Arg	Ala	Val	Asn	Asp
130						135					140				
Ser	Thr	Ser	Ala	Met	Thr	Gln	Thr	Ser	Ala	Val	Ile	Gly	Thr	Ala	Gln
145					150					155					160
Tyr	Leu	Ser	Pro	Glu	Gln	Ala	Arg	Gly	Lys	Pro	Ala	Asp	Ala	Arg	Ser
				165					170					175	
Asp	Ile	Tyr	Ala	Thr	Gly	Cys	Val	Met	Tyr	Glu	Leu	Val	Thr	Gly	Lys
			180					185					190		
Pro	Pro	Phe	Glu	Gly	Glu	Ser	Pro	Phe	Ala	Val	Ala	Tyr	Gln	His	Val
		195					200					205			
Gln	Glu	Asp	Pro	Thr	Pro	Pro	Ser	Asp	Phe	Ile	Ala	Asp	Leu	Thr	Pro
	210					215					220				
Thr	Ser	Ala	Val	Asn	Val	Asp	Ala	Val	Val	Leu	Thr	Ala	Met	Ala	Lys
225					230					235					240
His	Pro	Ala	Asp	Arg	Tyr	Gln	Thr	Ala	Ser	Glu	Met	Ala	Ala	Asp	Leu
				245					250					255	
Gly	Arg	Leu	Ser	Arg	Asn	Ala	Val	Ser	His	Ala	Ala	Arg	Ala	His	Val
			260					265					270		
Glu	Thr	Glu	Glu	Thr	Pro	Glu	Glu	Pro	Glu	Thr	Arg	Phe	Ser	Thr	Arg
		275					280					285			
Thr	Ser	Thr	Gln	Val	Ala	Pro	Ala	Ala	Gly	Val	Ala	Ala	Ala	Ser	Thr
	290					295					300				
Gly	Ser	Gly	Ser	Ser	Ser	Arg	Lys	Arg	Gly	Ser	Arg	Gly	Leu	Thr	Ala
305					310					315					320
Leu	Ala	Ile	Val	Leu	Ser	Leu	Gly	Val	Val	Gly	Val	Ala	Gly	Ala	Phe
				325					330					335	
Thr	Tyr	Asp	Tyr	Phe	Ala	Asn	Ser	Ser	Ser	Thr	Ala	Thr	Ser	Ala	Ile
			340					345					350		
Pro	Asn	Val	Glu	Gly	Leu	Pro	Gln	Gln	Glu	Ala	Leu	Thr	Glu	Leu	Gln
		355					360					365			
Ala	Ala	Gly	Phe	Val	Val	Asn	Ile	Val	Glu	Glu	Ala	Ser	Ala	Asp	Val
	370					375					380				
Ala	Glu	Gly	Leu	Val	Ile	Arg	Ala	Asn	Pro	Ser	Val	Gly	Ser	Glu	Ile
385					390					395					400
Arg	Gln	Gly	Ala	Thr	Val	Thr	Ile	Thr	Val	Ser	Thr	Gly	Arg	Glu	Met
				405					410					415	
Ile	Asn	Ile	Pro	Asp	Val	Ser	Gly	Met	Thr	Leu	Glu	Asp	Ala	Ala	Arg
			420					425					430		
Ala	Leu	Glu	Asp	Val	Gly	Leu	Ile	Leu	Asn	Gln	Asn	Val	Arg	Glu	Glu
		435					440					445			
Thr	Ser	Asp	Asp	Val	Glu	Ser	Gly	Leu	Val	Ile	Asp	Gln	Asn	Pro	Glu
	450					455					460				
Ala	Gly	Gln	Glu	Val	Val	Val	Gly	Ser	Ser	Val	Ser	Leu	Thr	Met	Ser
465					470					475					480
Ser	Gly	Thr	Glu	Ser	Ile	Arg	Val	Pro	Asn	Leu	Thr	Gly	Met	Asn	Trp
				485					490					495	
Ser	Gln	Ala	Glu	Gln	Asn	Leu	Ile	Ser	Met	Gly	Phe	Asn	Pro	Thr	Ala
			500					505					510		
Ser	Tyr	Leu	Asp	Ser	Ser	Glu	Pro	Glu	Gly	Glu	Val	Leu	Ser	Val	Ser
		515					520					525			
Ser	Gln	Gly	Thr	Glu	Leu	Pro	Lys	Gly	Ser	Ser	Ile	Thr	Val	Glu	Val
	530					535					540				
Ser	Asn	Gly	Met	Leu	Ile	Gln	Ala	Pro	Asp	Leu	Ala	Arg	Met	Ser	Thr

-continued

545	550	555	560
Glu Gln Ala Ile Ser Ala Leu Arg Ala Ala Gly Trp Thr Ala Pro Asp	565	570	575
Gln Ser Leu Ile Val Gly Asp Ser Ile His Thr Ala Ala Leu Val Asp	580	585	590
Gln Asn Lys Ile Gly Phe Gln Ser Pro Thr Pro Ala Thr Leu Phe Arg	595	600	605
Lys Asp Ala Gln Val Gln Val Arg Leu Phe Glu Phe Asp Leu Ala Ala	610	615	620
Leu Val Gln			
625			

What is claimed is:

1. An isolated polynucleotide from corynebacteria comprising a polynucleotide sequence selected from the group consisting of:

- (a) a polynucleotide which is at least 90% identical to the polynucleotide of SEQ ID No. 1 and encodes a polypeptide having protein kinase B activity,
- (b) a polynucleotide coding for a polypeptide containing an amino acid sequence which is at least 90% identical to the amino acid sequence of SEQ ID No. 2, wherein the polypeptide has protein kinase B activity, or
- (c) a polynucleotide which is complementary to the polynucleotide (a) or (b).

2. The polynucleotide of claim 1, which is a DNA.

3. The polynucleotide of claim 1, which is an RNA.

4. The isolated polynucleotide of claim 1 which comprises the nucleic acid sequence as shown in SEQ ID NO:1.

5. The polynucleotide that is DNA according to claim 2 comprising:

- (i) the nucleotide sequence shown in SEQ ID NO: 1, or
- (ii) a nucleotide sequence encoding a polypeptide comprising the amino acid sequence of SEQ ID NO: 2.

6. An isolated polynucleotide isolated from the species *Corynebacterium glutamicum* that hybridizes to the complement of SEQ ID NO: 1, wherein the isolated polynucleotide encodes a polypeptide having the enzymatic activity of protein kinase B and the hybridization is carried out under a stringency 0.5× SSC at a temperature of at 68° C.

7. The isolated polynucleotide from corynebacteria of claim 1, which encodes a polypeptide comprising the amino acids sequence shown in SEQ ID NO: 2.

8. The polynucleotide of claim 1, which is (a).

9. The polynucleotide of claim 1, which is (b).

10. The polynucleotide of claim 1, which is (c).

11. Recombinant Corynebacteria in which the isolated polynucleotide of claim 1 or claim 6 is amplified.

12. The recombinant Corynebacteria of claim 11, in which the isolated polynucleotide is overexpressed.

13. A fermentation process for the preparation of an L-amino acid, comprising:

- (a) fermenting in a medium recombinant corynebacteria which produce the L-amino acid, wherein at least the isolated polynucleotide of claim 1 or claim 6 is amplified in the recombinant corynebacteria,
- (b) enriching the L-amino acid in the medium or in the cells of the corynebacteria, and
- (c) isolating the L-amino acid.

14. The process of claim 13, wherein the L-amino acid is L-lysine.

15. The process of claim 13, wherein the L-amino acid is selected from the group consisting of L-asparagine, L-threonine, L-serine, L-glutamate, L-glycine, L-alanine, L-cysteine, L-valine, L-methionine, L-isoleucine, L-leucine, L-tyrosine, L-phenylalanine, L-histidine, L-tryptophan, and L-arginine.

16. The process of claim 13, wherein at least the isolated polynucleotide is overexpressed.

17. The process of claim 13, wherein said recombinant corynebacteria is transformed with a plasmid vector, and the said plasmid vector comprises the isolated polynucleotide of claim 1 or claim 6.

18. The process of claim 13, wherein one or more genes selected from the following group are simultaneously amplified or overexpressed in the corynebacteria:

- the dapA gene coding for dihydrodipicolinate synthase,
- the gap gene coding for glyceraldehyde 3-phosphate dehydrogenase,
- the tpi gene coding for triose phosphate isomerase,
- the pgk gene coding for 3-phosphoglycerate kinase,
- the zwf gene coding for glucose-6-phosphate dehydrogenase,
- the pyc gene coding for pyruvate carboxylase,
- the lysC gene coding for a feedback-resistant aspartate kinase,
- the lysE gene coding for lysine export,
- the hom gene coding for homoserine dehydrogenase,
- the ilvA gene coding for threonine dehydratase or the ilvA(Fbr) allele coding for a feedback-resistant threonine dehydratase,
- the ilvBN gene coding for acetohydroxy acid synthase,
- the ilvD gene coding for dihydroxy acid dehydratase, and
- the zwa1 gene coding for the Zwa1 protein.

19. The process of claim 13, wherein one or more genes selected from the following group are simultaneously attenuated in the corynebacteria:

- the pck gene coding for phosphoenol pyruvate carboxykinase,
- the pgi gene coding for glucose-6-phosphate isomerase,
- the poxB gene coding for pyruvate oxidase, and
- the zwa2 gene coding for the Zwa2 protein.

20. The process of claim 13, wherein the corynebacteria are *Corynebacterium glutamicum*.

21. The process of claim 20, wherein the *Corynebacterium glutamicum* is strain DSM 13994.

22. Recombinant corynebacteria which contain a vector comprises an isolated polynucleotide as claimed in claim 1 or claim 6.



## 31

**23.** A method of identifying a nucleic acid which code for SEQ ID NO:2 or a polynucleotide which is at least 90% identical to the polynucleotide of SEQ ID NO:1 and encodes a polypeptide having protein kinase B activity, comprising:

contacting a sample with the polynucleotide of claim 1 <sup>5</sup>  
under conditions suitable for the nucleic acid to hybridize to the polynucleotide; and

determining whether the nucleic acid codes for SEQ ID NO:2 or a polynucleotide which is at least 90% identical to the polynucleotide of SEQ ID NO:1 and encodes a polypeptide having protein kinase B activity. <sup>10</sup>

**24.** The method of claim **23**, wherein said nucleic acid is RNA, cDNA, or DNA.

**25.** The method of claim **23**, which is conducted on an array, micro-array, or DNA chip. <sup>15</sup>

**26.** An isolated DNA originating from corynebacteria and coding for protein kinase B, wherein the corresponding amino acid sequences contain any other proteogenic amino acid except L-proline in position 584 in SEQ ID No. 2.

**27.** An isolated DNA originating from corynebacteria and coding for protein kinase B, wherein the corresponding amino acid sequences contain L-serine or L-threonine in position 584 in SEQ ID No. 2. <sup>20</sup>

**28.** An isolated DNA originating from corynebacteria and coding for protein kinase B, wherein the corresponding amino acid sequence contains L-serine in position 584, shown in SEQ ID No. 4. <sup>25</sup>

**29.** The DNA of claim **28**, which contains the nucleobase thymine in position 2343, shown in SEQ ID No. 3.

## 32

**30.** Recombinant Corynebacteria which comprises a DNA as claimed in claim **26**.

**31.** Recombinant Corynebacteria which comprises a DNA as claimed in claim **27**.

**32.** Recombinant Corynebacteria which comprises a DNA as claimed in claim **28**.

**33.** *Corynebacterium glutamicum* DM1547 deposited as DSM13994 in the Deutsche Sammlung für Mikroorganismen und Zellkulturen (German Collection of Microorganisms and Cell Cultures), Brunswick, Germany.

**34.** A member of the Coryneform group of bacteria transformed by introduction of the polynucleotide according to any one of claim **1**, **2-7** and **26-28**.

**35.** The bacteria of claim **34**, which are of the genus *Corynebacterium*. <sup>15</sup>

**36.** A vector comprising the polynucleotide of any one of claims **1**, **2-7** and **26-28**.

**37.** A host cell of the *Corynebacterium* group of bacteria transformed with the vector of claim **36**. <sup>20</sup>

**38.** An isolated polynucleotide consisting of at least 26 consecutive nucleotides consisting of SEQ ID NO: 1 or of the full complement of SEQ ID NO: 1, wherein said isolated polynucleotide is a probe or primer for a polynucleotide sequence encoding a polypeptide having protein kinase B activity. <sup>25</sup>

\* \* \* \* \*