



US007148050B2

(12) **United States Patent**  
**Smolyar**

(10) **Patent No.: US 7,148,050 B2**  
(45) **Date of Patent: Dec. 12, 2006**

(54) **REGULATION OF HUMAN PROTEIN  
KINASE-LIKE PROTEIN**

2004/0053394 A1\* 3/2004 Gururajan et al. .... 435/252.3

FOREIGN PATENT DOCUMENTS

(75) Inventor: **Alex Smolyar**, Brookline, MA (US)

WO	WO 01 29564	4/2001
WO	WO 01 66594	9/2001
WO	WO 02 33099	4/2002

(73) Assignee: **Bayer Healthcare AG**, Leverkusen  
(DE)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 177 days.

OTHER PUBLICATIONS

(21) Appl. No.: **10/471,762**

Hanks et al.: "Protein kinases 6. The eukaryotic protein kinase  
superfamily: kinase (catalytic) domain structure and classification";  
FASEB Journal; May 1995; 9 (8); 576-96.

(22) PCT Filed: **Mar. 15, 2002**

Hanks et al.: "Protein Kinase Catalytic Domain Sequence Database:  
Identification of Conserved Features of Primary Structure and  
Classification of Family Members"; Methods in Enzymology; vol.  
200; 1991; pp. 38-62.

(86) PCT No.: **PCT/EP02/02887**

Database EMBL Online!; Sep. 29, 2000; Sugano et al.: "Homo  
sapiens cDNA: FLJ23119 is clone LNG07978"; Database acces-  
sion No. Ak026772; XP002222894.

§ 371 (c)(1),  
(2), (4) Date: **Apr. 13, 2004**

Nagase et al.: "Prediction of the coding sequences of unidentified  
human genes XX. The complete sequences of 100 new cDNA  
clones from brain which code for large proteins in vitro"; DNA  
Research; 8 (2); Apr. 27, 2001; 85-95; XP002943489.

(87) PCT Pub. No.: **WO02/081704**

PCT Pub. Date: **Oct. 17, 2002**

Dorow et al.: "Identification of a new family of human epithelial  
protein kinases containing two leucine/isoleucine-zipper domains";  
European Journal of Biochemistry; 213 (2); 1993; 701-710;  
XP001084152.

(65) **Prior Publication Data**

US 2004/0171539 A1 Sep. 2, 2004

Hanes et al.: "Characterization by cDNA cloning of two new human  
protein kinases. Evidence by sequence comparison of a new family  
of mammalian protein kinases"; Journal of Molecular Biology; vol.  
224; 1994; 665-672; XP002064038.

**Related U.S. Application Data**

(60) Provisional application No. 60/337,124, filed on Dec.  
10, 2001, provisional application No. 60/326,458,  
filed on Oct. 3, 2001, provisional application No.  
60/324,053, filed on Sep. 24, 2001, provisional appli-  
cation No. 60/276,055, filed on Mar. 16, 2001.

Taylor et al.: "How do protein kinases discriminate between serine/  
threonine and tyrosine? Structural insights from the insulin receptor  
protein-tyrosine kinase"; FASEB Journal; 9(131); 1995; 1255-1266;  
XP001109316.

(51) **Int. Cl.**

**C12N 9/12** (2006.01)

**C12N 15/00** (2006.01)

**C12N 1/20** (2006.01)

**C07H 21/04** (2006.01)

**C07K 1/00** (2006.01)

**C12Q 1/68** (2006.01)

Strausberg et al.: "Generation and initial analysis of more than  
15,000 full-length human and mouse cDNA sequences"; PROC  
NATL ACAD SCI USA; 2002; 99 (26); 16899-903 [PubMed/  
12477932].

\* cited by examiner

*Primary Examiner*—Maryam Monshipouri

(74) *Attorney, Agent, or Firm*—Banner & Witcoff, Ltd.

(52) **U.S. Cl.** ..... **435/194**; 530/350; 435/320.1;  
435/252.3; 435/6; 536/243.2

(57) **ABSTRACT**

(58) **Field of Classification Search** ..... 435/194,  
435/6, 320.1, 325, 252.3; 536/23.2; 530/356  
See application file for complete search history.

Reagents that regulate human protein kinase-like protein and  
reagents which bind to human protein kinase-like gene  
products can play a role in preventing, ameliorating, or  
correcting dysfunctions or diseases including, but not lim-  
ited to, cancer, CNS disorders, COPD, obesity, diabetes, and  
cardiovascular disorders.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

5,635,177 A 6/1997 Goeddel et al.

**18 Claims, 14 Drawing Sheets**

**Fig. 1**

atgaccgact	tcccggccag	gctcttcctg	gagaacagca
agctggagca	cagcgaggac	gagggcagcg	tcctggggcca
gggcggcagt	ggcaccgtca	tctaccgggc	ccgggtaccag
ggccagccctg	tggccgtcaa	gcgcttccac	atcaaaaaat
tcaagaactt	tgctaacgta	ccggcagaca	ccatgctgag
gcacctgcgg	gccaccgatg	ccatgaagaa	cttctccgag
ttccgggcagg	aggccagcat	gctgcacgcg	ctgcagcacc
cctgcatcgt	ggcgctcatc	ggcatcagca	tccacccgct
ctgcttcgcc	ctggagctcg	cgccgctcag	cagcctcaac
accgtgctgt	ccgagaacgc	cagagattct	tcctttatac
ccctgggaca	catgctcacc	caaaaaatag	cctaccagat
cgccctcgggc	ctggcctacc	tgcacaagaa	aaacatcatc
ttctgtgacc	tgaagtcgga	caacattctg	gtgtggtccc
ttgacgtcaa	ggagcacatc	aacatcaagc	tatctgacta
cgggattttcg	aggcagtcac	tccatgaggg	cgccctaggc
gtggaggggca	ctcctggcta	ccaggcccca	gagatcaggc
ctcgcatctgt	atatgatgag	aaggtagata	tgttctccta
tggaatggtg	ctctacgagt	tgctgtcagg	acagcgccct
gcactgggcc	accaccagct	ccagattgcc	aagaagctgt
ccaagggcat	ccgcccgggt	ctggggcagc	cggaggaagt
gcagttccgg	cgactgcagg	cgctcatgat	ggagtgctgg
gacactaagc	cagagaag		

**Fig. 2**

MTDFPARLFL	ENSKLEHSED	EGSVLGQGGG	GTVIYRARYQ
GQPVAVKRFH	IKKFKNFANV	PADTMLRHLR	ATDAMKNFSE
FRQEASMLHA	LQHPCIVALI	GISIHPLCFA	LELAPLSSLN
TVLSENARDS	SFIPLGHMLT	OKIAYQIASG	LAYLHKKNII
FCDLKSDNIL	VWSLDVKEHI	NIKLSDYGIS	RQSFHEGALG
VEGTPGYQAP	EIRPRIVYDE	KVDMFSYGMV	LYELLSGQRP
ALGHHQLQIA	KKLSKGIRPV	LGQPEEVQFR	RLQALMMECW DTKPEK

**Fig. 3**

AELTLEEIIGIGGGFGKVYRAFWIGDEVAVKAARHDPDEDISQTIENVRQEAKL  
FAMLKHPNIIALRGVCLKEPNLCLVMEFARGGPLNRVLSGKRIPPDILVNWAV  
QIARGMNYLHDEAIVPIIHRDLKSSNILILOKVENGDLNKKILKITDFGLARE  
WHRTTKMSAAGTYAWMAPEVIRASMFSKGSVDWSYGVLLWELLTGEVPFRGID  
GLRVAYGVAMNKLALPIPSTCPEPFAKLMEDCWNPDPHSRPSFTNILDQLTTI  
EESGFFEMPKDSFHCLQDNWKHEIQEMFDQLRAKEKELRTWEEELTRAALQOK  
NQEELLRRREQELAEREIDILEREELNIIHQLCQEKPRVKKRKGKFRKSRLAQ  
PVLPPHGHGSRCPGGTGSSWGGQ

**Fig. 4**

CACGAGGCTGGTCCCTGAACTGTTTCATGACCGACTTCCCGGCCAGGCTCTTCC  
TGGAGAACAGCAAGCTGGAGCACAGCGAGGACGAGGGCAGCGTCCTGGGCCAG  
GGCGGCAGTGGCACCGTCATCTACCGGGCCCGGTACCAGGGCCAGCCTGTGGC  
CGTCAAGCGCTTCCACATCAAAAAATTCAAGAACTTTGCTAACGTACCGGCAG  
ACACCATGCTGAGGCACCTGCGGGGCCACCGATGCCATGAAGAACTTCTCCGAG  
TTCCGGCAGGAGGCCAGCATGCTGCACGCGCTGCAGCACCCCTGCATCGTGGC  
GCTCATCGGCATCAGCATCCACCCGCTCTGCTTCGCCCTGGAGCTCGCGCCGC  
TCAGCAGCCTCAACACCGTGCTGTCCGAGAACGCCAGAGATTCTTCCTTTATA  
CCCCTGGGACACATGCTCACCCAAAAAATAGCCTACCAGATCGCCTCGG

**Fig. 5**

CACGAGGCTGGTCCCTGAACTGTTTCATGACCGACTTCCCGGCCAGGCTCTTCC  
TGGAGAACAGCAAGCTGGAGCACAGCGAGGACGAGGGCAGCGTCCTGGGCCAG  
GGCGGCAGTGGCACCGTCATCTACCGGGCCCGGTACCAGGGCCAGCCTGTGGC  
CGTCAAGCGCTTCCACATCAAAAAATTCAAGAACTTTGCTAACGTACCGGCAG  
ACACCATGCTGAGGCACCTGCGGGGCCACCGATGCCATGAAGAACTTCTCCGAG  
TTCCGGCAGGAGGCCAGCATGCTGCACGCGCTGCAGCACCCCTGCATCGTGGC  
GCTCATCGGCATCAGCATCCACCCGCTCTGCTTCGCCCTGGAGCTCGCGCCGC  
TCAGCAGCCTCAACACCGTGCTGTCCGAGAACGCCAGAGATTCTTCCTTTATA  
CCCCTGGGACACATGCTCACCCAAAAAATA

**Fig. 6**

CTCTTCCTGTGAGAACAGCAAGCTGGAGCACAGCGAGGACGAGGGCAGCGTCC  
TGGGCCACGGCGGCAGTGGCACCGTCATCTACCGGGCCCGGTACCAGGGCCAG  
CCTGTGGCCGTCAAGCGCTTCCACATCAAAAAATTCAAGAACTTTGCTAACGT  
ACCGGCAGACACCATGCTGAGGCACCTGCGGGGCCACCGATGCCATGAAGAACT  
TCTCCGAGTTCCGGCAGGAGGCCAGCATGCTGCACGCGCTGCAGCACCCCTGC  
ATCGTGGCGCTCATCGGCATCAGCATCCACCCGACTCTGCTTCGCCCTGGAGC  
TCGCGCCGCTCAGCAGCCTCAACACCGTGCTGTCCGAGAACGCCAGAGATTCT  
TCCTTTATACCCCTGGGACACATGCTCACCCAAAAAATAGCCTACCAGATCGC  
CTCGGGCCTGGCCTACCTGCACAAGAAAAACATCATCTTCTGTGACCTGAAGT  
CGGACAACATTCTGGTGTGGTCCCTTGACGTCAAGGAGCACATCAACATCAAG  
CTATCTGACTACGGGATTTTCGAGGCAGTCATTCCATGAGGGCGCCCTAGGCGT  
GGAGGGCACTCCTGGCTACCAGGGCCCCAGAGATCAGGCCTCGCATTGTATATG  
ATGAGAAGGTAGCATATGTCTCCTATGGAATGGTGCTCTACGAGTTGCTGTCA  
GGACAGGGGCCTGCATTGGGCCACCAACAGCTTCCGATTGCCAAGAGGTGTCC  
AAGGCATCCGCCCCGGTACTGAGGACAGCCGGAAGGAAGTGAATCCGGCGATG  
CATGGCTCATTATTGAGATGCTGGGAACTATAATCAAGCGAACGGAGCACTG  
TGGCCCTATACGTAGGTAGCCCCAATTAGACGACCCTCACATTATGAACCAAT  
CGGTAATAAACGTGTACGTGCGGGACTAGCAGCCACCTATAACACACAGCCAC  
GGCCACCTGGGCTGTGATGCATACAAGACGCTCAAACACGCGGGTCACAGAAA  
GCCTCATCGAGTCGCATCCACCCGAGAGAGTAGCGCCGCGGAAGTAGTATAGA  
CGGTAT

**Fig. 7**

CACGAGGAAAAAATTCAAGAACTTTGCTAACGTACCGGCAGACACCATGCTGA  
GGCACCTGCGGGCCACCGATGCCATGAAGAACTTCTCCGAGTTCCGGCAGGAG  
GCCAGCATGCTGCACGCGCTGCAGCACCCCTGCATCGTGGCGCTCATCGGCAT  
CAGCATCCACCCGCTCTGCTTCGCCCTGGAGCTCGCGCCGCTCAGCAGCCTCA  
ACACCGTGCTGTCCGAGAACGCCAGAGATTCTTCCTTTATACCCCTGGGACAC  
ATGCTCACCCAAAAAATAGCCTACCAGATCGCCTCGGGCCTGGCCTACCTGCA  
CAAGAAAAACATCATCTTCTGTGACCTGAAGTCGGACAACATTCTGGTGTGGT  
CCCTTGACGTCAAGGAGCACATCAACATCAAGCTATCTGACTACGGGATTTTCG  
AGGCAGTCATTCCATGAGGGCGCCCTANGCGTGGAGGGCACTCCTGG

**Fig. 8**

CCCCTTAATGATTTTGCACAGGCCCCCAAAGCAATGCTGTCTTACCTGCGTGCT  
CAGCTGCGGAAAGCGGAAAAGTGCAAGCTGATGAAGATGATCATCGTGGGTCC  
cccgcgccaggggcaagtccaccctcctggagatcttacagacggggagggccc  
cccaggtggtgcatggagaggccaccatcaggaccaccaagtgggagctccag  
aggccggctggctcaagagccaaggttgagtccgtggagtccaacgtctggga  
catcgggggacccggccagcatggccactgtcaaccagtgttcttcacggaca  
aggccctgtacgtggtggtctggaacctggcgctgggggaggaggccgtggcc  
aacctccagtctctggctgctcaacatcgaggccaaggccccaacgccgtggt  
gctggtggtcgggacgcacctggatttaattgaagccaagtccgtgtggaaa  
ggattgcaacgctgcgtgcctatgtgctggcactctgccgctccccctccggc  
tccagggccacaggcttcccagacatcaccttcaaacacttacatgagatttc  
ctgcaagagcctggaagggtcaggaagggtgcgacagctgattttccacgtca  
cgtgcagcatgaaggacgtgggcagcaccatcggctgccagcgactggcaggg  
cggctgatccccaggagctacctgagcctgcaggaggccgtgctggcagagca  
gcagcgcgcagccgggacgcagcagctgcagtacctgacggacaggcagctgg  
agcagctggtggagcagacgcccagacaacgacatcaaggactacgaggacctg  
cagtcagccaatcagcttccctcatagaaaccggcaccctgctccatttcccggg  
caccagccacggcctgaggaacctctacttccctcgacctatttggctctccg  
aatgtctgcagaggaatctttaatatgaagggtctcggctcagtggccaagaat  
ggggtgatcagagcagaagacctcaggatgctgctggtggggactggcttcac  
gcagcagacggaagagcagtagcttccagttcctggccaagtgttgagatcgccc  
tgcccgtcgccaatgacagctacctcctgccccatctccttccatctaaacct  
ggcctggacacccacgggtatgcggcacccccacagccaacaccattcagagggt  
atttaagatgagcttcgttcccgttggcttctggcaaagggttatagcacgga  
tgctgatcagcctggcggagatggacctgcagctttttgaaaacaagaagaat  
actaaaagcaggaacaggaaagtccacatttacagttttacaggaaaccagag  
aatcgctgtagcacatcagagtgaaaagaaatcagaccatctattggcagg  
aagggtcctggtcacttttgatgggggctacctcagtggtggaatcttccgac  
gtgaactggaaaaagaagaaaagcggagggaatgaaaattgtttgccaatcaga  
agtgagggaacttctcagccatggctttcatcacggaccacgtcaattccttga  
ttgatcagtggtttcccggccctgacggccacagagagcgacgggacgccactc  
atggagcagtagctgccctgcccggctctgcgagacagcctggggcccagcacac  
ggaccccagtgagaaatcagaggatgtgtagtagtacttcgacatggaagactgtg  
tcctgacggccatcgagcgggacttcatctcctgccccagacacccggacctc  
cccgtgccgctgcaggagctgggtccctgaactgttcatgaccgacttcccggc  
caggctcttccctggagaacagcaagctggagcacagcgaggacgagggcagcg  
tcctggggccagggcggcagtgggcaccgtcatctaccggggcccggtaccagggc  
cagcctgtggccgtcaagcgcctccacatcaaaaaaattcaagaactttgctaa  
cgtaccgggcagacaccatgctgaggcacctgcggggccaccgatgccatgaaga  
acttctccgagttccggcaggaggccagcatgctgcacgcgctgcagcaccctc  
tgcattcgtggcgctcatcggcacatccacccgctctgcttcgcccctgga  
gctcgcgcgcgtcagcagcctcaacaccgtgctgtccgagaacgccagagatt  
cttcctttataccccctgggacacatgctcacccaaaaaatagcctaccagatc  
gcctcgggcctggcctacctgcacaagaaaaaacatcatcttctgtgacctgaa  
gtcggacaacaattctggtgtggtcccttgacgtcaaggagcacatcaacatca  
agctatctgactacgggatttccgaggcagtcattccatgagggcgccctaggc  
gtggaggggcactcctggctaccaggccccagagatcaggccctcgcatgtgata  
tgatgagaaggtagatatgttctcctatggaaatgggtgctctacgagttgctgt  
caggacagcgccctgcactggggccaccaccagctccagattgccaagaagctg  
tccaagggcacccgcccgggttctggggcagccggagggaagtgcagttccggcg  
actgcaggcgctcatgatggagtgctggggacactaagccagagaagcgaccgc  
tgggccctgtcggtggtgagccagatgaaggacccgacttttgccaccttcatg  
tatgaactgtgctgtgggaagcagacagccttcttctcatcccagggccagga

**Fig. 8 (continued)**

gtacaccgtggtgttttgggatggaaaagaggagtccaggaactacacgggtgg  
tgaacacagagaagggcctcatggaggtgcagaggatgtgctgccctgggatg  
aaggtgagctgccagctccaggtccagagatccctgtggacagccaccgagga  
ccagaaaatctacatctacaccctcaagggcatgtgccccttaaacacacccc  
aacaggccttggatactccagctgtcgtcacctgcttcttggccgtgcctgtt  
attaaaaagaattcctacctggtcttagcgggcctcgccgatgggcttgtggc  
tgtgtttcccgtggtgcggggcaccaccaaggacagctgctcctacctgtgct  
cacacacagccaacagggtccaagtccagcatcgccgatgaagacgcacggcag  
aaccctacctcagtgaaaggccatggaggtggtcaacagcggctctgaggtctg  
gtacagcaatgggcccgggcctccttgtcatcgactgtgctccttgagatct  
gcaggcggctggagccctacatggccccctccatggttacgtcagtcgtgtgc  
agctctgagggcagaggggaggaggtcgtctggtgcttgatgacaaggccaa  
ctccttgggtgatgtaccactccaccacctaccagctgtgtgcccgggtacttct  
gcgggggtccccagccccctcagggacatgtttcccgtgcggcccttggacacg  
gaacccccggcagccagccacacggccaaccccaaagggtgcctgagggggactc  
catcgccgacgtgagcatcatgtacagtgaggagctgggcacgcagatcctga  
tccaccaggaatcactcactgactactgctccaatgtcctcctactcctcatcc  
ccaccccgccaggctgccagggtccccctcaagcctccccagctccccagcaag  
ttcttccagtgtgcttcttccaccgactgcgaggactcagacatgctacata  
cgcccgggtgctgcttccgacagggtctgagcatgacctgacccccatggacggg  
gagaccttcagccagcacctgcaggccgtgaagatcctcgccgtcagagacct  
catttgggtccccaggcgcgggtggagatgttatcgctcattggcctggagaagg  
attctgaagcccagcggggccgagtcattggcgtcttaaaaagcccagagctg  
actccgcatgggggtgctggtggatgctgccgtgggtggcaaggacactgttgt  
gtgcacctttgaaaatgaaaacacagagtgggtgcctggccgtctggaggggct  
ggggcgccaggaggttcgacattttctaccagtcctacgaggagctggggccgg  
ctggaggcttgcactcgcaagagaaggtaattcctgtggaatgactgtcacac  
atcagagctggctggccccggggctgcagcctgacccctctgccatcggcctct  
agtcttccaaggacctagaagacagatggagtcttccccctgaactccttgctg  
ctaagaagtgtgagaagttactcgccctggcgggtggctccagggttctctggt  
tctctggagcagagttctctgaatacccccatcccccaactgctgattttacag  
ccccagggaagacagtggtafcagggtgggagcggcctcctctggcctcccc  
atcagtttgcaggagcaggggtgcaggatcctgttctgagctgggtcaaaca  
agcagggccgggccttccctgccatccccagggtctcagatggaattacactaga  
ggccctccgctgggaagcacttgaggtagggcaggaggggggctgtgacccct  
gccctttccccgcccagagacctcgggctctcagcacattccacaggctcctga  
gtccccgaggcctggggcagcttgggcaagccaagatcagatgtctctgtgtt  
cgggaaggctctccgtgtgggaaagcccttgggggatcccgggtgaggagtggt  
gccccatccagagaatgaatgagttcctttaagtgccaccgcccagcaagccca  
gaggcacacagtcagagtgaccccgcttagcctttacattcctctccaccgac  
aaaagggaaggggaaactcaatcagcaggacttcagaaagggccttgtgtttat  
agctttgtcaagttaaatttggaacgcagctggagcacaggccctgtttgtttgc  
acataataatcttgtttatcactttaaaaaattcagtaatatctcagcagtc  
ggcttctggttgtgaaatcacattgtatgggattttataccaaattatgtattt  
gctaaacattcactgcacacgtgtacagcggagtacgaaaagggaacggtgtcc  
acaggggatatttatggatacaacagcaaacattttataaactatgcacatgcat  
tacacacatgcacacacatatgcacacacatgtgcaaacatagccactttttt  
gtcaagagttaccctttggggctccttaaacagaaatgggagtttgaaagaga  
gatcafactccagctgaagttgttgacccttttctaaaattaaaaagatcaa  
atttagtatttgctggatatgcaggggagatgagactcttttaattctcaaaata  
aacagattctttcaag

**Fig. 9**

MLSYLRAQLRKA EKCKLMKMIIVGPPRQ GKSTLLEILQTGRAPQVVHGEATIR  
TTKWELQRPAGSRAKVESVEFNVWDIGGPASMATVNQCFFTDKALYVVVWNLA  
LGEEAVANLQFWLLNIEAKAPNAVVLVVGTHLDLIEAKFRVERIATLRAYVLA  
LCRSPSGSRATGFPDITFKHLHEISCKSLEGQEGRLQLIFHVTC SMKDVGSTI  
GCQRLAGRLIPRSYLSLQEAVLAEQQRRSRDDDVQYLTD RQLEQLVEQTPDND  
IKDYEDLQSAISFLIETGTL LHFPDTSHGLRNLYFLDPIWLSECLQRI FNIKG  
SRSVAKNGVIRAEDLRMLLVGTGFTQQTEE QYFQFLAKFEIALPVANDSYLLP  
HLLPSKPGLDTHGMRHPTANTIQRVFKMSFVPVGF WQRFIARMLISLAEMDLQ  
LFENKKNTKSRNRKVTIYSFTGNQRNRCSTFRVKRNQTIYWQEGLLVTFDGGY  
LSVESSDVNWKKKKSGGMKIVCQSEVRDFSAMAFITDHVNSLIDQWFPALTAT  
ESDGTPLMEQYVPCPVCETAWAQHTDPSEKSE DVQYFDMEDCVLTAIERDFIS  
CPRHPDLPVPLQELVPELFMTDFPARLFLENSKLEHSEDEGSVLGQGGSGTVI  
YRARYQGQPVAVKRFHIKKFKNFANVPADTMLRHLRATDAMKNFSEFRQEASM  
LHALQHPCIVALIGISIHPLCFALELAPLSSLNTVLSENARDSSFIPLGHMLT  
QKIAYQIASGLAYLHKKNIIFCDLKSDNILVWSLDVKEHINIKLS DYGISRQS  
FHEGALGVEGTPGYQAPEIRPRIVYDEKVDMFSYGMVLYELL SGQRPALGHHQ  
LQIAKKLSKGIRPVLGQP EEVQFRRLQALMMECWDTKPEKRPLALSVVSQMKD  
PTFATFMYELCCGKQTAFSSQGEYTVVFWDGKEESRNYTVVNTEKGLMEVQ  
RMCCPGMKVSCQLQVQRSLWTATEDQKIYIYTLKGMCP LNTPQQALDTPAVVT  
CFLAVPVIKKNSYLVLAGLADGLVAVFPVVRGTPK DSCSYLCSHTANRSKFST  
ADEDARQNPYPVKAMEVVNSGSEVWYSNGPGLLVIDCASLEICRRLEPYMAPS  
MVTSVVCSSEGRGEEVWCLDDKANSLVMYHSTTYQLCARYFCGVPSPLRDMF  
PVRPLDTEPPAASHTANPKVPEGDSIADV SIMYSEELGTQILIHQESLTDYCS  
MSSYSSSPPRQAARSPSSLPSSPASSSSVPFSTDCEDSDMLHTPGAASDRSEH  
DLTPMDGETFSQHLQAVKILAVRDLI WVPRRGGDVIVIGLEKDSEAQRGRVIA  
VLKARELTPHGVLVDAAVVAKDTVVCTFENENTEWCLAVWRGWGAREFDIFYQ  
SYEELGRLEACTRKRR

Fig. 10

CAAAGCAATGCTGTCTTACCTGCGTGCTCAGCTGCGGAAAGCGGAAAAGTGCA  
AGCTGATGAAGATGATCATCGTGGGTCCCCCGCGCCAGGGCAAGTCCACCCTC  
CTGGAGATCTTACAGACGGGGAGGGCCCCCAGGTGGTGCATGGAGAGGCCAC  
CATCAGGACCACCAAGTGGGAGCTCCAGAGGCCGGCTGGCTCAAGAGCCAAGG  
TTGAGTCCGTGGAGTTCAACGTCTGGGACATCGGGGGACCGGCCAGCATGGCC  
ACTGTCAACCAGTGCTTCTTCACGGACAAGGCCCTGTACGTGGTGGTCTGGAA  
CCTGGCGCTGGGGGAGGAGGCCGTGGCCAACCTCCAGTTCTGGCTGCTCAACA  
TCGAGGCCAAGGCCCAAACGCCGTGGTGCTGGTGGTTCGGGACGCACCTGGAT  
TTAATTGAAGCCAAGTTCCTGTGGAAAGGATTGCAACGCTGCGTGCCCTATGT  
GCTGGCACTCTGCCGCTCCCCCTCCGGCTCCAGGGCCACAGGCTTCCAGACA  
TCACCTTCAAACACTTACATGAGATTTCTGCAAGAGCCTGGAAGGTCAGGAA  
GGGCTGCGACAGCTGATTTTCCACGTACGTGCAGCATGAAGGACGTGGGCAG  
CACCATCGGCTGCCAGCGACTGGCAGGGCGGCTGATCCCCAGGAGCTACCTGA  
GCCTGCAGGAGGCCGTGCTGGCAGAGCAGCAGCGCCGCAGCCGGGACGACGAC  
GTGCAGTACCTGACGGACAGGCAGCTGGAGCAGCTGGTGGAGCAGACGCCCGA  
CAACGACATCAAGGACTACGAGGACCTGCAGTCAGCCATCAGCTTCCTCATAG  
AAACCGGCACCCTGCTCCATTTCCCGGACACCAGCCACGGCCTGAGGAACCTC  
TACTTCCTCGACCCTATTTGGCTCTCCGAATGTCTGCAGAGGATCTTTAATAT  
TAAGGGCTCTCGGTGAGTGGCCAAGAATGGGGTGATCAGAGCAGAAGACCTCA  
GGATGCTGCTGGTGGGGACTGGCTTCACGCAGCAGACGGAAGAGCAGTACTTC  
CAGTTCCTGGCCAAGTTTGAGATCGCCCTGCCCGTCCGCAATGACAGCTACCT  
CCTGCCCCATCTCCTTCCATCTAAACCTGGCCTGGACACCCACGGTATGCGGC  
ACCCACAGCCAACACCATTTCAGAGGGTATTTAAGATGAGCTTCGTTCCCGTT  
GGCTTCTGGCAAAGGTTTATAGCACGGATGCTGATCAGCCTGGCGGAGATGGA  
CCTGCAGCTTTTTGAAAACAAGAAGAATACTAAAAGCAGGAACAGGAAAGTCA  
CCATTTACAGTTTTTACAGGAAACCAGAGAAATCGCTGTAGCACATTCAGAGTG  
AAAAGAAATCAGACCATCTATTGGCAGGAAGGGCTCCTGGTCACTTTTGATGG  
GGGCTACCTCAGTGTGGAATCTTCCGACGTGAACTGGAAAAAGAAGAAAAGCG  
GAGGAATGAAAATTGTTTGCCAATCAGAAGTGAGGGACTTCTCAGCCATGGCT  
TTCATCACGGACCACGTCAATTCCTTGATTGATCAGTGGTTTCCCGCCCTGAC  
AGCCACAGAGAGCGACGGGACGCCACTCATGGAGCAGTACGTGCCCTGCCCGG  
TCTGCGAGACAGCCTGGGCCCAGCACACGGACCCAGTGAGAAATCAGAGGAT  
GTGCAGTACTTCGACATGGAAGACTGTGTCTGACGGCCATCGAGCGGGACTT  
CATCTCCTGCCCCAGACACCCGGACCTCCCCGTGCCGCTGCAGGAGCTGGTCC  
CTGAACTGTTTCATGACCGACTTCCCGGCCAGGCTCTTCTGAGAACAGCAAG  
CTGGAGCACAGCGAGGACGAGGGCAGCGTCTGGGCCAGGGCGGCAGTGGCAC  
CGTCATCTACCGGGCCCGGTACCAGGGCCAGCCTGTGGCCGTCAAGCGCTTCC  
ACATCAAAAAAATTCAAGAACTTTGCTAACGTACCGGCAGACACCATGCTGAGG  
CACCTGCGGGCCACCGATGCCATGAAGAACTTCTCCGAGTTCCGGCAGGAGGC  
CAGCATGCTGCACGCGCTGCAGCACCCCTGCATCGTGGCGCTCATCGGCATCA  
GCATCCACCCGCTCTGCTTCGCCCTGGAGCTCGCGCCGCTCAGCAGCCTCAAC  
ACCGTGCTGTCCGAGAACGCCAGAGATTCTTCTTTTATACCCCTGGGACACAT  
GCTCACCCAAAAAATAGCCTACCAGATCGCCTCGGGCCTGGCCTACCTGCACA  
AGAAAAACATCATCTTCTGTGACCTGAAGTCGGACAACATTCTGGTGTGGTCC  
CTTGACGTCAAGGAGCACATCAACATCAAGCTATCTGACTACGGGATTTTCGAG  
GCAGTCATTCCATGAGGGCGCCCTAGGCGTGGAGGGCACTCCTGGCTACCAGG  
CCCCAGAGATCAGGCCTCGCATTGTATATGATGAGAAGGTAGATATGTTCTCC  
TATGGAATGGTGCTCTACGAGTTGCTGTGAGGACAGCGCCCTGCACTGGGCCA  
CCACCAGCTCCAGATTGCCAAGAAGCTGTCCAAGGGCATCCGCCCGGTTCTGG  
GGCAGCCGGAGGAAGTGCAAGTTCGGGCGACTGCAGGCGCTCATGATGGAGTGC  
TGGGACACTAAGCCAGAGAAGCGACCGCTGGCCCTGTCCGTGGTGGAGCCAGAT  
GAAGGACCCGACTTTTGCCACCTTCATGTATGAACTGTGCTGTGGGAAGCAGA  
CAGCCTTCTTCTCATCCCAGGGCCAGGAGTACACCGTGGTGTTTTGGGATGGA

**Fig. 10 (continued)**

AAAGAGGAGTCCAGGAACTACACGGTGGTGAACACAGAGAAGGGCCTCATGGA  
GGTGCAGAGGATGTGCTGCCCTGGGATGAAGGTGAGCTGCCAGCTCCAGGTCC  
AGAGATCCCTGTGGACAGCCACCGAGGACCAGAAAATCTACATCTACACCCTC  
AAGGGCATGTGCCCCCTTAAACACACCCCCAACAGGCCTTGGATACTCCAGCTGT  
CGTCACCTGCTTCTTGGCCGTGCCTGTTATTAAAAAGAATTCTTACCTGGTCT  
TAGCGGGCCTCGCCGATGGGCTTGTGGCTGTGTTTCCCGTGGTGCGGGGCACC  
CCAAAGGACAGCTGCTCCTACCTGTGCTCACACACAGCCAACAGGTCCAAGTT  
CAGCATCGCGGATGAAGACGCACGGCAGAACCCCTACCCAGTGAAGGCCATGG  
AGGTGGTCAACAGCGGCTCTGAGGTCTGGTACAGCAATGGGCGGGCCTCCTT  
GTCATCGACTGTGCCTCCCTGGAGATCTGCAGGCGGCTGGAGCCCTACATGGC  
CCCCTCCATGGTTACGTCAGTCGTGTGCAGCTCTGAGGGCAGAGGGGAGGAGG  
TCGTCTGGTGCCTGGATGACAAGGCCAACTCCTTGGTGATGTACCACTCCACC  
ACCTACCAGCTGTGTGCCCGGTACTTCTGCGGGGTCCCCAGCCCCCTCAGGGA  
CATGTTTCCCGTGCGGCCCTTGGACACGGAACCCCCCGGCAGCCAGCCACACGG  
CCAACCCAAAGGTGCCTGAGGGGGGACTCCATCGCGGACGTGAGCATCATGTAC  
AGTGAGGAGCTGGGCACGCAGATCCTGATCCACCAGGAATCACTCACTGACTA  
CTGCTCCATGTCTCTACTCCTCATCCCCACCCCGCCAGGCTGCCAGGTCCC  
CCTCAAGCCTCCCCAGCTCCCCAGCAAGTTCTTCCAGTGTGCCTTTCTCCACC  
GACTGCGAGGACTCAGACATGCTACATACGCCCGGTGCTGCCTCCGACAGGTC  
TGAGCATGACCTGACCCCCATGGACGGGGGAGACCTTCAGCCAGCACCTGCAGG  
CCGTGAAGATCCTCGCCGTCAGAGACCTCATTTGGGTCCCCAGGCGCGGTGGA  
GATGTTATCGTCATTGGCCTGGAGAAGGATTCTGAAGCCCAGCGGGGCCGAGT  
CATTGCCGTCTTAAAAGCCCGAGAGCTGACTCCGCATGGGGTGCTGGTGGATG  
CTGCCGTGGTGGCAAAGGACACTGTTGTGTGCACCTTTGAAAATGAAAACACA  
GAGTGGTGCCTGGCCGTCTGGAGGGGCTGGGGCGCCAGGGAGTTCGACATTTT  
CTACCAGTCCTACGAGGAGCTGGGCCGGCTGGAGGCTTGCACTCGCAAGAGAA  
GGTAATTCCTGTGGAATGACTGTCACACA

**BLASTP - alignment of 423 against swiss P80192 M3K9 HUMAN**

```

This hit is scoring at : 5e-23 (expectation value)
Alignment length (overlap) : 287
Identities : 31 %
Scoring matrix : BLOSUM62 (used to infer consensus pattern)
Database searched : nrdb_1;

```

```

Q: 24 VLQGGSGTVIYRARYQGQPVAVKRFHIKKFKNFANVPADTMLRHLRATDAMKNFSEFRQ
      :G GG G.V YRA : G.V VAVK.
H: 8 IIGIGFGKV-YRAFWIGDEVAVKA-----ARHDPDEDISQTIENTVRQ

      bind ATP
      active site residue
      EA.:. L:HP I:AL G:.. LC.:E.A ..LN.VLS .. IP .:..
      EAKLFAMLKHPNIIALRGVCLKEPNLCLVMEFARGGPLNRVLSGKR-----IPPDILVNW
      KIAYQIASGLAYLHKKNI---IFCDLKSDNIIVWSL---DVKEHINIKLSDYGISRQSF
      A.QIA.G: YLH.: I I.DLKS.NIL: . D:..I :K:D:G::R: .
      --AVQIARGMNYLHDEAIVPIIHRDLKSSNIILOKVENGDLNKI-LKITDFGLAREWH

      HEGALGVEGTPGYQAEIRPRIVYDEKVDMSYGMVLYELLSGQRPALGHHQLQIA-----
      :.:.:.GT :.APE: .:.. D::SYG::L:ELL:G: P G . L::A
      RTTKMSAAGTYAWMAPEVIRASMFSGSDVWSYGVLLWELLTGEVFPFRGIDGLRVAYGVA

      -KKLSKGIRPVLGQPPEEVQFRRLQALMMECWDTKPEKRP 284
      .KL: I :..:P ..LM :CW:..P..RP
      MNKLALPIPSTCPEP-----FAKLMEDCWNPDPHSRP 249

```

Fig. 12

HMMPFAM - alignment of 423 against pfam|hmm|pkinese

Eukaryotic protein kinase domain

This hit is scoring at : 161.2 E=2.6e-42  
Scoring matrix : BLOSUM62 (used to infer consensus pattern)

Q: 19 EDEGSVLGQGSGTViYRARYQ-GQPVAVKRFHIKKFKnfanvpadtmlrlratdamkn  
: LG:G. G.V Y:A::: G: VAVK ..:..  
H: 1 yelleklGeGsfGkv.ykakhktgkiVAvKilkkesl.....  
  
fsefRQEASMLHALQHPCIVALIGISI---HPLCFALELAPLSSLNTVLSenardssfiP  
:E:::L..L.HP IV.L:G: L::E...:L::L..N. P  
....lrEiqilkrIshPNIvrllgvfedtdhhlylvmEymegGdLfdylrrng.....p  
  
LGHMLTQKIAyQIASGLAYLHKKNIIFCDLKSdNIlVWSLdvkehINIKLSDYGISRQsf  
L...:KIA.QI..GL.YLH...I::DLK::NIL: :K::D:G::R  
lsekeakkialQilrGleYlHsngivHRDLKpenIllden.....gtvKiadFGLArll.  
  
hEGALGVEGTPGYQ-APEI--RPrIVYDEKVDmFSYGMVLyELLSG-----  
E..:GTP Y. APE: . Y..KVD::S.G::LYELL:G L: L:::  
.eklttfvGTPwYmmAPEvileg.rgysskvDvWSIGviLyElltggplfpgadlpaftg  
  
-----QRPALGH-----HQLQIAKKLSKG-IRPVLGQPPEEvqfrLQALMME L: L:::  
gdevdqliifvklpfsdelpktridpleelfrikrrlplpsncSee.....lkdlLlkk  
  
CWDTKPEK 286  
C :..P.K  
cLnkDPsk 262

Fig. 13

Relative expression of HUMAN PROTEIN KINASE-LIKE PROTEIN mRNA :  
Cancer/NAT pairs




Fig. 14




Fig. 15





Fig. 16



## REGULATION OF HUMAN PROTEIN KINASE-LIKE PROTEIN

This application is a National Stage application of co-  
pending PCT application PCT/EP02/02887 filed Mar. 15, 2002, which was published in English under PCT Article 21(2) on Oct. 17, 2002, which claims the benefit of U.S. provisional application Ser. No. 60/276,055 filed Mar. 16, 2001, Ser. No. 60/324,053 filed Sep. 24, 2001, Ser. No. 60/326,458 filed Oct. 3, 2001, and Ser. No. 60/337,124 filed Dec. 10, 2001. These applications are incorporated herein by reference in their entireties.

### TECHNICAL FIELD OF THE INVENTION

The invention relates to the regulation of a human protein kinase-like protein.

### BACKGROUND OF THE INVENTION

Protein kinases play critical roles in the regulation of biochemical and morphological changes associated with cellular growth and division. D'Urso et al., *Science* 250, 786-91, 1990; Birchmeier et al., *Bioessays* 15, 185-89, 1993; U.S. Pat. No. 6,200,770. They serve as growth factor receptors and signal transducers and have been implicated in cellular transformation and malignancy. Hunter et al., *Cell* 70, 375-87, 1992; Posada et al., *Mol. Biol. Cell* 3, 583-92, 1992; Hunter et al., *Cell* 79, 573-82, 1994. For example, protein kinases have been shown to participate in the transmission of signals from growth-factor receptors (Sturgill et al., *Nature* 344, 715-18, 1988; Gomez et al., *Nature* 353, 170-73, 1991), control of entry of cells into mitosis (Nurse et al., *Nature* 344, 503-08, 1990; Maller, *Curr. Opin. Cell Biol.* 3, 269-75, 1991), and regulation of actin bundling (Husain-Chishti et al., *Nature* 334, 718-21, 1988).

Protein kinases can be divided into two main groups based on either amino acid sequence similarity or specificity for either serine/threonine or tyrosine residues. A small number of dual-specificity kinases are structurally like the serine/threonine-specific group. Within the broad classification, kinases can be further sub-divided into families whose members share a higher degree of catalytic domain amino acid sequence identity and also have similar biochemical properties. Most protein kinase family members also share structural features outside the kinase domain that reflect their particular cellular roles. These include regulatory domains that control kinase activity or interaction with other proteins. Hanks et al., *Science* 241, 42-52, 1988).

There is a need in the art to identify protein kinase-like proteins, which can be regulated to provide therapeutic effects.

### SUMMARY OF THE INVENTION

It is an object of the invention to provide reagents and methods of regulating a human protein kinase-like protein. This and other objects of the invention are provided by one or more of the embodiments described below.

One embodiment of the invention is a proteine kinase-like protein polypeptide comprising an amino acid sequence selected from the group consisting of:

amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 2;  
the amino acid sequence shown in SEQ ID NO: 2;  
amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 9 and;  
the amino acid sequence shown in SEQ ID NO:9.

Yet another embodiment of the invention is a method of screening for agents which decrease extracellular matrix degradation. A test compound is contacted with a proteine kinase-like protein polypeptide comprising an amino acid sequence selected from the group consisting of:

amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 2;  
the amino acid sequence shown in SEQ ID NO: 2;  
amino acid sequences which are at least about 50% identical to the amino acid sequence shown in SEQ ID NO: 9 and;  
the amino acid sequence shown in SEQ ID NO:9.

Binding between the test compound and the proteine kinase-like protein polypeptide is detected. A test compound which binds to the proteine kinase-like protein polypeptide is thereby identified as a potential agent for decreasing extracellular matrix degradation. The agent can work by decreasing the activity of the proteine kinase-like protein.

Another embodiment of the invention is a method of screening for agents which decrease extracellular matrix degradation. A test compound is contacted with a polynucleotide encoding a proteine kinase-like protein polypeptide, wherein the polynucleotide comprises a nucleotide sequence selected from the group consisting of:

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 1;  
the nucleotide sequence shown in SEQ ID NO: 1;  
nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 8;  
the nucleotide sequence shown in SEQ ID NO: 8;  
nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 10; and  
the nucleotide sequence shown in SEQ ID NO: 10.

Binding of the test compound to the polynucleotide is detected. A test compound which binds to the polynucleotide is identified as a potential agent for decreasing extracellular matrix degradation. The agent can work by decreasing the amount of the proteine kinase-like protein through interacting with the proteine kinase-like protein mRNA.

Another embodiment of the invention is a method of screening for agents which regulate extracellular matrix degradation. A test compound is contacted with a proteine kinase-like protein polypeptide comprising an amino acid sequence selected from the group consisting of:

amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 2;  
the amino acid sequence shown in SEQ ID NO: 2;  
amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 9 and;  
the amino acid sequence shown in SEQ ID NO:9.

A proteine kinase-like protein activity of the polypeptide is detected. A test compound which increases proteine kinase-like protein activity of the polypeptide relative to proteine kinase-like protein activity in the absence of the test compound is thereby identified as a potential agent for increasing extracellular matrix degradation. A test compound which decreases proteine kinase-like protein activity of the polypeptide relative to proteine kinase-like protein activity in the absence of the test compound is thereby identified as a potential agent for decreasing extracellular matrix degradation.

Even another embodiment of the invention is a method of screening for agents which decrease extracellular matrix degradation. A test compound is contacted with a proteine kinase-like protein product of a polynucleotide which comprises a nucleotide sequence selected from the group consisting of:

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 1; the nucleotide sequence shown in SEQ ID NO: 1; nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 8; the nucleotide sequence shown in SEQ ID NO: 8; nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 10; and the nucleotide sequence shown in SEQ ID NO:10.

Binding of the test compound to the protein kinase-like protein product is detected. A test compound which binds to the protein kinase-like protein product is thereby identified as a potential agent for decreasing extracellular matrix degradation.

Still another embodiment of the invention is a method of reducing extracellular matrix degradation. A cell is contacted with a reagent which specifically binds to a polynucleotide encoding a protein kinase-like protein polypeptide or the product encoded by the polynucleotide, wherein the polynucleotide comprises a nucleotide sequence selected from the group consisting of:

nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 1; the nucleotide sequence shown in SEQ ID NO: 1; nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 8; the nucleotide sequence shown in SEQ ID NO: 8; nucleotide sequences which are at least about 50% identical to the nucleotide sequence shown in SEQ ID NO: 10; and the nucleotide sequence shown in SEQ ID NO: 10.

Protein kinase-like protein activity in the cell is thereby decreased.

The invention thus provides a human protein kinase-like protein that can be used to identify test compounds that may act, for example, as activators or inhibitors at the enzyme's active site. Human protein kinase-like protein and fragments thereof also are useful in raising specific antibodies that can block the enzyme and effectively reduce its activity.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:1).

FIG. 2 shows the amino acid sequence deduced from the DNA-sequence of FIG. 1 (SEQ ID NO:2).

FIG. 3 shows the amino acid sequence of the protein identified by swissP80192M3K9\_HUMAN (SEQ ID NO:3).

FIG. 4 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:4).

FIG. 5 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:5).

FIG. 6 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:6).

FIG. 7 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:7).

FIG. 8 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:8).

FIG. 9 shows the amino acid sequence deduced from the DNA-sequence of FIG. 8 (SEQ ID NO:9).

FIG. 10 shows the DNA-sequence encoding a protein kinase-like protein Polypeptide (SEQ ID NO:10).

FIG. 11 shows the BLASTP—alignment of 423 (SEQ ID NO:2) against swissP80192M3K9\_HUMAN (SEQ ID NO:3).

FIG. 12 shows the HMMPFAM—alignment of 423 (SEQ ID NO:2) against pfamhmmmpkinase.

FIG. 13 shows the relative expression of human protein kinase-like protein mRNA in cancer cells and tissue.

FIG. 14 shows the relative expression of protein kinase-like protein mRNA in tissue relevant for obesity and diabetes.

FIG. 15 shows the copy number of protein kinase-like protein mRNA in tissue relevant for obesity and diabetes.

FIG. 16 shows the relative expression profile of protein kinase-like protein mRNA in various human tissues (body-map).

#### DETAILED DESCRIPTION OF THE INVENTION

The invention relates to an isolated polynucleotide from the group consisting of:

- a) a polynucleotide encoding a protein kinase-like protein polypeptide comprising an amino acid sequence selected from the group consisting of:
  - amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 2;
  - the amino acid sequence shown in SEQ ID NO: 2;
  - amino acid sequences which are at least about 35% identical to the amino acid sequence shown in SEQ ID NO: 9 and;
  - the amino acid sequence shown in SEQ ID NO:9.
- b) a polynucleotide comprising the sequence of SEQ ID NOS: 1, 8 or 10;
- c) a polynucleotide which hybridizes under stringent conditions to a polynucleotide specified in (a) and (b) and encodes a protein kinase-like protein polypeptide;
- d) a polynucleotide the sequence of which deviates from the polynucleotide sequences specified in (a) to (c) due to the degeneration of the genetic code and encodes a protein kinase-like protein polypeptide; and
- e) a polynucleotide which represents a fragment, derivative or allelic variation of a polynucleotide sequence specified in (a) to (d) and encodes a protein kinase-like protein polypeptide.

Furthermore, it has been discovered by the present applicant that a novel protein kinase-like protein, particularly a human protein kinase-like protein, can be used in therapeutic methods to treat cancer, a CNS disorder, COPD, obesity, diabetes, or a cardiovascular disorder.

Human protein kinase-like protein comprises the amino acid sequence shown in SEQ ID NOS:2 and 9. A coding sequence for human protein kinase-like protein is shown in SEQ ID NOS: 1, 8 and 10. This sequence is located on chromosome 15. Related ESTs (SEQ ID NOS:4–7) are expressed in lymphoma and germinal center B cells.

Human protein kinase-like protein is 31% identical over 287 amino acids to swissP80192M3K9\_HUMAN (SEQ ID NO:3) (FIG. 1). Pfam homology search results support the identification of this protein as a eukaryotic protein kinase. Its likely function as a tyrosine kinase class m protein also is supported by the results of a BLOCKS database search. Consensus patterns are underlined in FIG. 1; the active site Asp residue and the ATP-binding Lys residue are highlighted in bold.

Human protein kinase-like protein of the invention is expected to be useful for the same purposes as previously identified protein kinase-like protein enzymes. Human protein kinase-like protein is believed to be useful in therapeutic methods to treat disorders such as cancer, CNS disorders, COPD, obesity, diabetes, and cardiovascular disorders.

Human protein kinase-like protein also can be used to screen for human protein kinase-like protein activators and inhibitors.

#### Polypeptides

Human protein kinase-like polypeptides according to the invention comprise at least 6, 10, 15, 20, 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, or 286 contiguous amino acids selected from the amino acid sequence shown in SEQ ID NO:2 or a biologically active variant thereof, as defined below. Human protein kinase-like polypeptides according to the invention comprise at least 6, 10, 15, 20, 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1025, 1050, 1075, 1100, 1125, 1150, 1175, 1200, 1225, 1250, 1275, 1300, 1325, 1350, 1375 or 1394 contiguous amino acids selected from the amino acid sequence shown in SEQ ID NO:9 or a biologically active variant thereof, as defined below. A protein kinase-like polypeptide of the invention therefore can be a portion of a protein kinase-like protein, a full-length protein kinase-like protein, or a fusion protein comprising all or a portion of a protein kinase-like protein.

#### Biologically Active Variants

Human protein kinase-like polypeptide variants which are biologically active, e.g., retain enzymatic activity, also are human protein kinase-like polypeptides. Preferably, naturally or non-naturally occurring human protein kinase-like polypeptide variants have amino acid sequences which are at least about 32, 35, 40, 45, 50, 55, 60, 65, or 70, preferably about 75, 80, 85, 90, 96, 98, or 99% identical to the amino acid sequence shown in SEQ ID NOS:2 and 9 or a fragment thereof. Percent identity between a putative human protein kinase-like polypeptide variant and an amino acid sequence of SEQ ID NO:2 or 9 is determined by conventional methods. See, for example, Altschul et al., *Bull. Math. Bio.* 48:603 (1986), and Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1992). Briefly, two amino acid sequences are aligned to optimize the alignment scores using a gap opening penalty of 10, a gap extension penalty of 1, and the "BLOSUM62" scoring matrix of Henikoff & Henikoff, 1992.

Those skilled in the art appreciate that there are many established algorithms available to align two amino acid sequences. The "FASTA" similarity search algorithm of Pearson & Lipman is a suitable protein alignment method for examining the level of identity shared by an amino acid sequence disclosed herein and the amino acid sequence of a putative variant. The FASTA algorithm is described by Pearson & Lipman, *Proc. Nat'l Acad. Sci. USA* 85:2444 (1988), and by Pearson, *Meth. Enzymol.* 183:63 (1990). Briefly, FASTA first characterizes sequence similarity by identifying regions shared by the query sequence (e.g., SEQ ID NO: 2 or 9) and a test sequence that have either the highest density of identities (if the ktup variable is 1) or pairs of identities (if ktup=2), without considering conservative amino acid substitutions, insertions, or deletions. The ten regions with the highest density of identities are then rescored by comparing the similarity of all paired amino acids using an amino acid substitution matrix, and the ends of the regions are "trimmed" to include only those residues that contribute to the highest score. If there are several regions with scores greater than the "cutoff" value (calculated by a predetermined formula based upon the length of the sequence the ktup value), then the trimmed initial regions are examined to determine whether the regions can be joined

to form an approximate alignment with gaps. Finally, the highest scoring regions of the two amino acid sequences are aligned using a modification of the Needleman-Wunsch-Sellers algorithm (Needleman & Wunsch, *J. Mol. Biol.* 48:444 (1970); Sellers, *SIAM J. Appl. Math.* 26:787 (1974)), which allows for amino acid insertions and deletions. Preferred parameters for FASTA analysis are: ktup=1, gap opening penalty=10, gap extension penalty=1, and substitution matrix=BLOSUM62. These parameters can be introduced into a FASTA program by modifying the scoring matrix file ("SMATRIX"), as explained in Appendix 2 of Pearson, *Meth. Enzymol.* 183:63 (1990).

FASTA can also be used to determine the sequence identity of nucleic acid molecules using a ratio as disclosed above. For nucleotide sequence comparisons, the ktup value can range between one to six, preferably from three to six, most preferably three, with other parameters set as default.

Variations in percent identity can be due, for example, to amino acid substitutions, insertions, or deletions. Amino acid substitutions are defined as one for one amino acid replacements. They are conservative in nature when the substituted amino acid has similar structural and/or chemical properties. Examples of conservative replacements are substitution of a leucine with an isoleucine or valine, an aspartate with a glutamate, or a threonine with a serine.

Amino acid insertions or deletions are changes to or within an amino acid sequence. They typically fall in the range of about 1 to 5 amino acids. Guidance in determining which amino acid residues can be substituted, inserted, or deleted without abolishing biological or immunological activity of a human protein kinase-like polypeptide can be found using computer programs well known in the art, such as DNASTAR software.

The invention additionally, encompasses protein kinase-like polypeptides that are differentially modified during or after translation, e.g., by glycosylation, acetylation, phosphorylation, amidation, derivatization by known protecting/blocking groups, proteolytic cleavage, linkage to an antibody molecule or other cellular ligand, etc. Any of numerous chemical modifications can be carried out by known techniques including, but not limited, to specific chemical cleavage by cyanogen bromide, trypsin, chymotrypsin, papain, V8 protease, NaBH<sub>4</sub>, acetylation, formylation, oxidation, reduction, metabolic synthesis in the presence of tunicamycin, etc.

Additional post-translational modifications encompassed by the invention include, for example, e.g., N-linked or O-linked carbohydrate chains, processing of N-terminal or C-terminal ends), attachment of chemical moieties to the amino acid backbone, chemical modifications of N-linked or O-linked carbohydrate chains, and addition or deletion of an N-terminal methionine residue as a result of prokaryotic host cell expression. The protein kinase-like polypeptides may also be modified with a detectable label, such as an enzymatic, fluorescent, isotopic or affinity label to allow for detection and isolation of the protein.

The invention also provides chemically modified derivatives of protein kinase-like polypeptides that may provide additional advantages such as increased solubility, stability and circulating time of the polypeptide, or decreased immunogenicity (see U.S. Pat. No. 4,179,337). The chemical moieties for derivitization can be selected from water soluble polymers such as polyethylene glycol, ethylene glycol/propylene glycol copolymers, carboxymethylcellulose, dextran, polyvinyl alcohol, and the like. The polypep-

tides can be modified at random or predetermined positions within the molecule and can include one, two, three, or more attached chemical moieties.

Whether an amino acid change or a polypeptide modification results in a biologically active protein kinase-like polypeptide can readily be determined by assaying for enzymatic activity, as described for example, in U.S. Pat. No. 6,194,186.

#### Fusion Proteins

Fusion proteins are useful for generating antibodies against protein kinase-like polypeptide amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins that interact with portions of a protein kinase-like polypeptide. Protein affinity chromatography or library-based assays for protein-protein interactions, such as the yeast two-hybrid or phage display systems, can be used for this purpose. Such methods are well known in the art and also can be used as drug screens.

A protein kinase-like polypeptide fusion protein comprises two polypeptide segments fused together by means of a peptide bond. The first polypeptide segment comprises at least 6, 10, 15, 20, 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, or 286 contiguous amino acids of SEQ D NOS:2 or of a biologically active variant, such as those described above. The first polypeptide segment comprises at least 6, 10, 15, 20, 25, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, 550, 575, 600, 625, 650, 675, 700, 725, 750, 775, 800, 825, 850, 875, 900, 925, 950, 975, 1000, 1025, 1050, 1075, 1100, 1125, 1150, 1175, 1200, 1225, 1250, 1275, 1300, 1325, 1350, 1375 or 1394 contiguous amino acids of SEQ ID NOS:2 or of a biologically active variant, such as those described above. The first polypeptide segment also can comprise full-length protein kinase-like protein.

The second polypeptide segment can be a full-length protein or a protein fragment. Proteins commonly used in fusion protein construction include 13-galactosidase, P-glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Additionally, epitope tags are used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex a DNA binding domain (DBD) fusions, GALA DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions. A fusion protein also can be engineered to contain a cleavage site located between the protein kinase-like polypeptide-encoding sequence and the heterologous protein sequence, so that the protein kinase-like polypeptide can be cleaved and purified away from the heterologous moiety.

A fusion protein can be synthesized chemically, as is known in the art. Preferably, a fusion protein is produced by covalently linking two polypeptide segments or by standard procedures in the art of molecular biology. Recombinant DNA methods can be used to prepare fusion proteins, for example, by making a DNA construct which comprises coding sequences selected from SEQ ID NOS: 1, 8 and 10 in proper reading frame with nucleotides encoding the second polypeptide segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies

such as Promega Corporation (Madison, Wis.), Stratagene (La Jolla, Calif.), CLONTECH (Mountain View, Calif.), Santa Cruz Biotechnology (Santa Cruz, Calif.), MBL International Corporation (MIC; Watertown, Mass.), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

#### Identification of Species Homologs

Species homologs of human protein kinase-like polypeptide can be obtained using protein kinase-like polypeptide polynucleotides (described below) to make suitable probes or primers for screening cDNA expression libraries from other species, such as mice, monkeys, or yeast, identifying cDNAs which encode homologs of protein kinase-like polypeptide, and expressing the cDNAs as is known in the art.

#### Polynucleotides

A protein kinase-like polynucleotide can be single- or double-stranded and comprises a coding sequence or the complement of a coding sequence for a protein kinase-like polypeptide. A coding sequence for human protein kinase-like protein is shown in SEQ ID NOS: 1, 8 and 10.

Degenerate nucleotide sequences encoding human protein kinase-like polypeptides, as well as homologous nucleotide sequences which are at least about 50, 55, 60, 65, 70, preferably about 75, 90, 96, 98, or 99% identical to the nucleotide sequence shown in SEQ ID NOS: 1, 8 and 10 or its complement also are protein kinase-like polynucleotides. Percent sequence identity between the sequences of two polynucleotides is determined using computer programs such as ALIGN which employ the FASTA algorithm, using an affine gap search with a gap open penalty of -12 and a gap extension penalty of -2. Complementary DNA (cDNA) molecules, species homologs, and variants of protein kinase-like polynucleotides that encode biologically active protein kinase-like polypeptides also are protein kinase-like polynucleotides. Polynucleotide fragments comprising at least 8, 9, 10, 11, 12, 15, 20, or 25 contiguous nucleotides of SEQ ID NOS: 1, 8 and 10 or its complement also are protein kinase-like polynucleotides. These fragments can be used, for example, as hybridization probes or as antisense oligonucleotides.

#### Identification of Polynucleotide Variants and Homologs

Variants and homologs of the protein kinase-like polynucleotides described above also are protein kinase-like polynucleotides. Typically, homologous protein kinase-like polynucleotide sequences can be identified by hybridization of candidate polynucleotides to known protein kinase-like polynucleotides under stringent conditions, as is known in the art. For example, using the following wash conditions—2×SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room temperature twice, 30 minutes each; then 2×SSC, 0.1% SDS, 50° C. once, 30 minutes; then 2×SSC, room temperature twice, 10 minutes each—homologous sequences can be identified which contain at most about 25–30% basepair mismatches. More preferably, homologous nucleic acid strands contain 15–25% basepair mismatches, even more preferably 5–15% basepair mismatches.

Species homologs of the protein kinase-like polynucleotides disclosed herein also can be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, or yeast. Human variants of protein kinase-like polynucleotides can be identified, for example, by screening human cDNA expression libraries. It is well known that the  $T_m$  of a double-stranded DNA decreases by 1–1.5° C. with every 1%

decrease in homology (Bonner et al., *J. Mol. Biol.* 81, 123 (1973). Variants of human protein kinase-like polynucleotides or protein kinase-like polynucleotides of other species can therefore be identified by hybridizing a putative homologous protein kinase-like polynucleotide with a polynucleotide having a nucleotide sequence of SEQ ID NOS: 1, 8 and 10 or the complement thereof to form a test hybrid. The melting temperature of the test hybrid is compared with the melting temperature of a hybrid comprising polynucleotides having perfectly complementary nucleotide sequences, and the number or percent of basepair mismatches within the test hybrid is calculated.

Nucleotide sequences which hybridize to protein kinase-like polynucleotides or their complements following stringent hybridization and/or wash conditions also are protein kinase-like polynucleotides. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, 2d ed., 1989, at pages 9.50–9.51.

Typically, for stringent hybridization conditions a combination of temperature and salt concentration should be chosen that is approximately 12–20° C. below the calculated  $T_m$  of the hybrid under study. The  $T_m$  of a hybrid between a protein kinase-like polynucleotide having a nucleotide sequence shown in SEQ ID NOS: 1, 8 and 10 or the complement thereof and a polynucleotide sequence which is at least about 50, preferably about 75, 90, 96, or 98% identical to one of those nucleotide sequences can be calculated, for example, using the equation of Bolton and McCarthy, *Proc. Natl. Acad. Sci. U.S.A.* 48, 1390 (1962):

$$T_m = 81.5^\circ \text{ C.} - 16.6(\log_{10}[\text{Na}^+]) - 0.41(\% \text{ G+C}) - 0.63(\% \text{ formamide}) - 600/l,$$

where  $l$  = the length of the hybrid in basepairs.

Stringent wash conditions include, for example, 4×SSC at 65° C., or 50% formamide, 4×SSC at 42° C., or 0.5×SSC, 0.1% SDS at 65° C. Highly stringent wash conditions include, for example, 0.2×SSC at 65° C.

#### Preparation of Polynucleotides

A protein kinase-like polynucleotide can be isolated free of other cellular components such as membrane components, proteins, and lipids. Polynucleotides can be made by a cell and isolated using standard nucleic acid purification techniques, or synthesized using an amplification technique, such as the polymerase chain reaction (PCR), or by using an automatic synthesizer. Methods for isolating polynucleotides are routine and are known in the art. Any such technique for obtaining a polynucleotide can be used to obtain isolated protein kinase-like polynucleotides. For example, restriction enzymes and probes can be used to isolate polynucleotide fragments, which comprise protein kinase-like protein nucleotide sequences. Isolated polynucleotides are in preparations that are free or at least 70, 80, or 90% free of other molecules.

Human protein kinase-like cDNA molecules can be made with standard molecular biology techniques, using protein kinase-like mRNA as a template. Human protein kinase-like cDNA molecules can thereafter be replicated using molecular biology techniques known in the art and disclosed in manuals such as Sambrook et al. (1989). An amplification technique, such as PCR, can be used to obtain additional copies of polynucleotides of the invention, using either human genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize protein kinase-like polynucleotides. The degeneracy of the genetic code allows alternate nucleotide

sequences to be synthesized which will encode a protein kinase-like polypeptide having, for example, an amino acid sequence shown in SEQ ID NOS: 2 and 9 or a biologically active variant thereof.

#### Extending Polynucleotides

The partial sequence disclosed herein can be used to identify the corresponding full length gene from which it was derived. The partial sequence can be nick-translated or end-labeled with  $^{32}\text{P}$  using polynucleotide kinase using labeling methods known to those with skill in the art (*BASIC METHODS IN MOLECULAR BIOLOGY*, Davis et al., eds., Elsevier Press, N.Y., 1986). A lambda library prepared from human tissue can be directly screened with the labeled sequences of interest or the library can be converted en masse to pBluescript (Stratagene Cloning Systems, La Jolla, Calif. 92037) to facilitate bacterial colony screening (see Sambrook et al., *MOLECULAR CLONING: A LABORATORY MANUAL*, Cold Spring Harbor Laboratory Press (1989, pg. 1.20).

Both methods are well known in the art. Briefly, filters with bacterial colonies containing the library in pBluescript or bacterial lawns containing lambda plaques are denatured, and the DNA is fixed to the filters. The filters are hybridized with the labeled probe using hybridization conditions described by Davis et al., 1986. The partial sequences, cloned into lambda or pBluescript, can be used as positive controls to assess background binding and to adjust the hybridization and washing stringencies necessary for accurate clone identification. The resulting autoradiograms are compared to duplicate plates of colonies or plaques; each exposed spot corresponds to a positive colony or plaque. The colonies or plaques are selected, expanded and the DNA is isolated from the colonies for further analysis and sequencing.

Positive cDNA clones are analyzed to determine the amount of additional sequence they contain using PCR with one primer from the partial sequence and the other primer from the vector. Clones with a larger vector-insert PCR product than the original partial sequence are analyzed by restriction digestion and DNA sequencing to determine whether they contain an insert of the same size or similar as the mRNA size determined from Northern blot Analysis.

Once one or more overlapping cDNA clones are identified, the complete sequence of the clones can be determined, for example after exonuclease III digestion (McCombie et al., *Methods* 3, 33–40, 1991). A series of deletion clones are generated, each of which is sequenced. The resulting overlapping sequences are assembled into a single contiguous sequence of high redundancy (usually three to five overlapping sequences at each nucleotide position), resulting in a highly accurate final sequence.

Various PCR-based methods can be used to extend the nucleic acid sequences disclosed herein to detect upstream sequences such as promoters and regulatory elements. For example, restriction-site PCR uses universal primers to retrieve unknown sequence adjacent to a known locus (Sarkar, *PCR Methods Applic.* 2, 318–322, 1993). Genomic DNA is first amplified in the presence of a primer to a linker sequence and a primer specific to the known region. The amplified sequences are then subjected to a second round of PCR with the same linker primer and another specific primer internal to the first one. Products of each round of PCR are transcribed with an appropriate RNA polymerase and sequenced using reverse transcriptase.

Inverse PCR also can be used to amplify or extend sequences using divergent primers based on a known region (Triglia et al., *Nucleic Acids Res.* 16, 8186, 1988). Primers

can be designed using commercially available software, such as OLIGO 4.06 Primer Analysis software (National Biosciences Inc., Plymouth, Minn.), to be 22–30 nucleotides in length, to have a GC content of 50% or more, and to anneal to the target sequence at temperatures about 68–72° C. The method uses several restriction enzymes to generate a suitable fragment in the known region of a gene. The fragment is then circularized by intramolecular ligation and used as a PCR template.

Another method which can be used is capture PCR, which involves PCR amplification of DNA fragments adjacent to a known sequence in human and yeast artificial chromosome DNA (Lagerstrom et al., *PCR Methods Applic.* 1, 111–119, 1991). In this method, multiple restriction enzyme digestions and ligations also can be used to place an engineered double-stranded sequence into an unknown fragment of the DNA molecule before performing PCR.

Another method which can be used to retrieve unknown sequences is that of Parker et al., *Nucleic Acids Res.* 19, 3055–3060, 1991). Additionally, PCR, nested primers, and PROMOTERFINDER libraries (CLONTECH, Palo Alto, Calif) can be used to walk genomic DNA (CLONTECH, Palo Alto, Calif.). This process avoids the need to screen libraries and is useful in finding intron/exon junctions.

When screening for full-length cDNAs, it is preferable to use libraries that have been size-selected to include larger cDNAs. Randomly-primed libraries are preferable, in that they will contain more sequences which contain the 5' regions of genes. Use of a randomly primed library may be especially preferable for situations in which an oligo d(T) library does not yield a full-length cDNA. Genomic libraries can be useful for extension of sequence into 5' non-transcribed regulatory regions.

Commercially available capillary electrophoresis systems can be used to analyze the size or confirm the nucleotide sequence of PCR or sequencing products. For example, capillary sequencing can employ flowable polymers for electrophoretic separation, four different fluorescent dyes (one for each nucleotide) that are laser activated, and detection of the emitted wavelengths by a charge coupled device camera. Output/light intensity can be converted to electrical signal using appropriate software (e.g. GENOTYPER and Sequence NAVIGATOR, Perkin Elmer), and the entire process from loading of samples to computer analysis and electronic data display can be computer controlled. Capillary electrophoresis is especially preferable for the sequencing of small pieces of DNA that might be present in limited amounts in a particular sample.

#### Obtaining Polypeptides

Human protein kinase-like polypeptides can be obtained, for example, by purification from human or other cells, by expression of protein kinase-like polynucleotides, or by direct chemical synthesis.

#### Protein Purification

Human protein kinase-like polypeptides can be purified from any cell that expresses the polypeptide, including host cells that have been transfected with protein kinase-like protein expression constructs. A purified protein kinase-like polypeptide is separated from other compounds that normally associate with the protein kinase-like polypeptide in the cell, such as certain proteins, carbohydrates, or lipids, using methods well-known in the art. Such methods include, but are not limited to, size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity chromatography, and preparative gel electrophoresis. A preparation of purified protein kinase-like

polypeptides is at least 80% pure; preferably, the preparations are 90%, 95%, or 99% pure. Purity of the preparations can be assessed by any means known in the art, such as SDS-polyacrylamide gel electrophoresis.

#### Expression of Polynucleotides

To express a protein kinase-like polynucleotide, the polynucleotide can be inserted into an expression vector that contains the necessary elements for the transcription and translation of the inserted coding sequence. Methods that are well known to those skilled in the art can be used to construct expression vectors containing sequences encoding protein kinase-like polypeptides and appropriate transcriptional and translational control elements. These methods include in vitro recombinant DNA techniques, synthetic techniques, and in vivo genetic recombination. Such techniques are described, for example, in Sambrook et al. (1989) and in Ausubel et al., *CURRENT PROTOCOLS IN MOLECULAR BIOLOGY*, John Wiley & Sons, New York, N.Y., 1989.

A variety of expression vector/host systems can be utilized to contain and express sequences encoding a protein kinase-like polypeptide. These include, but are not limited to, microorganisms, such as bacteria transformed with recombinant bacteriophage, plasmid, or cosmid DNA expression vectors; yeast transformed with yeast expression vectors, insect cell systems infected with virus expression vectors (e.g., baculovirus), plant cell systems transformed with virus expression vectors (e.g., cauliflower mosaic virus, CaMV; tobacco mosaic virus, TMV) or with bacterial expression vectors (e.g., Ti or pBR322 plasmids), or animal cell systems.

The control elements or regulatory sequences are those non-translated regions of the vector—enhancers, promoters, 5' and 3' untranslated regions—which interact with host cellular proteins to carry out transcription and translation. Such elements can vary in their strength and specificity. Depending on the vector system and host utilized, any number of suitable transcription and translation elements, including constitutive and inducible promoters, can be used. For example, when cloning in bacterial systems, inducible promoters such as the hybrid lacZ promoter of the BLUE-SCRIPT phagemid (Stratagene, LaJolla, Calif.) or pSPORT1 plasmid (Life Technologies) and the like can be used. The baculovirus polyhedrin promoter can be used in insect cells. Promoters or enhancers derived from the genomes of plant cells (e.g., heat shock, RUBISCO, and storage genes) or from plant viruses (e.g., viral promoters or leader sequences) can be cloned into the vector. In mammalian cell systems, promoters from mammalian genes or from mammalian viruses are preferable. If it is necessary to generate a cell line that contains multiple copies of a nucleotide sequence encoding a protein kinase-like polypeptide, vectors based on SV40 or EBV can be used with an appropriate selectable marker.

#### Bacterial and Yeast Expression Systems

In bacterial systems, a number of expression vectors can be selected depending upon the use intended for the protein kinase-like polypeptide. For example, when a large quantity of a protein kinase-like polypeptide is needed for the induction of antibodies, vectors which direct high level expression of fusion proteins that are readily purified can be used. Such vectors include, but are not limited to, multifunctional *E. coli* cloning and expression vectors such as BLUESCRIPT (Stratagene). In a BLUESCRIPT vector, a sequence encoding the protein kinase-like polypeptide can be ligated into the vector in frame with sequences for the amino-terminal Met and the subsequent 7 residues of  $\beta$ -galactosidase so that

a hybrid protein is produced. pIN vectors (Van Heeke & Schuster, *J. Biol. Chem.* 264, 5503–5509, 1989) or pGEX vectors (Promega, Madison, Wis.) also can be used to express foreign polypeptides as fusion proteins with glutathione S-transferase (GST). In general, such fusion proteins are soluble and can easily be purified from lysed cells by adsorption to glutathione-agarose beads followed by elution in the presence of free glutathione. Proteins made in such systems can be designed to include heparin, thrombin, or factor Xa protease cleavage sites so that the cloned polypeptide of interest can be released from the GST moiety at will.

In the yeast *Saccharomyces cerevisiae*, a number of vectors containing constitutive or inducible promoters such as alpha factor, alcohol oxidase, and PGH can be used. For reviews, see Ausubel et al. (1989) and Grant et al., *Methods Enzymol.* 153, 516–544, 1987.

#### Plant and Insect Expression Systems

If plant expression vectors are used, the expression of sequences encoding protein kinase-like polypeptides can be driven by any of a number of promoters. For example, viral promoters such as the 35S and 19S promoters of CaMV can be used alone or in combination with the omega leader sequence from TMV (Takamatsu, *EMBO J.* 6, 307–311, 1987). Alternatively, plant promoters such as the small subunit of RUBISCO or heat shock promoters can be used (Coruzzi et al., *EMBO J.* 3, 1671–1680, 1984; Broglie et al., *Science* 224, 838–843, 1984; Winter et al., *Results Probl. Cell Differ.* 17, 85–105, 1991). These constructs can be introduced into plant cells by direct DNA transformation or by pathogen-mediated transfection. Such techniques are described in a number of generally available reviews (e.g., Hobbs or Murray, in McGRAW HILL YEARBOOK OF SCIENCE AND TECHNOLOGY, McGraw Hill, New York, N.Y., pp. 191–196, 1992).

An insect system also can be used to express a protein kinase-like polypeptide. For example, in one such system *Autographa californica* nuclear polyhedrosis virus (AcNPV) is used as a vector to express foreign genes in *Spodoptera frugiperda* cells or in *Trichoplusia* larvae. Sequences encoding protein kinase-like polypeptides can be cloned into a non-essential region of the virus, such as the polyhedrin gene, and placed under control of the polyhedrin promoter. Successful insertion of protein kinase-like polypeptides will render the polyhedrin gene inactive and produce recombinant virus lacking coat protein. The recombinant viruses can then be used to infect *S. frugiperda* cells or *Trichoplusia* larvae in which protein kinase-like polypeptides can be expressed (Engelhard et al., *Proc. Nat. Acad. Sci.* 91, 3224–3227, 1994).

#### Mammalian Expression Systems

A number of viral-based expression systems can be used to express protein kinase-like polypeptides in mammalian host cells. For example, if an adenovirus is used as an expression vector, sequences encoding protein kinase-like polypeptides can be ligated into an adenovirus transcription/translation complex comprising the late promoter and tripartite leader sequence. Insertion in a non-essential E1 or E3 region of the viral genome can be used to obtain a viable virus that is capable of expressing a protein kinase-like polypeptide in infected host cells (Logan & Shenk, *Proc. Natl. Acad. Sci.* 81, 3655–3659, 1984). If desired, transcription enhancers, such as the Rous sarcoma virus (RSV) enhancer, can be used to increase expression in mammalian host cells.

Human artificial chromosomes (HACs) also can be used to deliver larger fragments of DNA than can be contained and expressed in a plasmid. HACs of 6M to 10M are constructed and delivered to cells via conventional delivery methods (e.g., liposomes, polycationic amino polymers, or vesicles).

Specific initiation signals also can be used to achieve more efficient translation of sequences encoding protein kinase-like polypeptides. Such signals include the ATG initiation codon and adjacent sequences. In cases where sequences encoding a protein kinase-like polypeptide, its initiation codon, and upstream sequences are inserted into the appropriate expression vector, no additional transcriptional or translational control signals may be needed. However, in cases where only coding sequence, or a fragment thereof, is inserted, exogenous translational control signals (including the ATG initiation codon) should be provided. The initiation codon should be in the correct reading frame to ensure translation of the entire insert. Exogenous translational elements and initiation codons can be of various origins, both natural and synthetic. The efficiency of expression can be enhanced by the inclusion of enhancers which are appropriate for the particular cell system which is used (see Scharf et al., *Results Probl. Cell Differ.* 20, 125–162, 1994).

#### Host Cells

A host cell strain can be chosen for its ability to modulate the expression of the inserted sequences or to process the expressed protein kinase-like polypeptide in the desired fashion. Such modifications of the polypeptide include, but are not limited to, acetylation, carboxylation, glycosylation, phosphorylation, lipidation, and acylation. Post-translational processing which cleaves a “prepro” form of the polypeptide also can be used to facilitate correct insertion, folding and/or function. Different host cells that have specific cellular machinery and characteristic mechanisms for post-translational activities (e.g., CHO, HeLa, MDCK, HEK293, and WI38), are available from the American Type Culture Collection (ATCC; 10801 University Boulevard, Manassas, Va. 20110-2209) and can be chosen to ensure the correct modification and processing of the foreign protein.

Stable expression is preferred for long-term, high-yield production of recombinant proteins. For example, cell lines which stably express protein kinase-like polypeptides can be transformed using expression vectors which can contain viral origins of replication and/or endogenous expression elements and a selectable marker gene on the same or on a separate vector. Following the introduction of the vector, cells can be allowed to grow for 1–2 days in an enriched medium before they are switched to a selective medium. The purpose of the selectable marker is to confer resistance to selection, and its presence allows growth and recovery of cells which successfully express the introduced protein kinase-like protein sequences. Resistant clones of stably transformed cells can be proliferated using tissue culture techniques appropriate to the cell type. See, for example, ANIMAL CELL CULTURE, R. I. Freshney, ed., 1986.

Any number of selection systems can be used to recover transformed cell lines.

These include, but are not limited to, the herpes simplex virus thymidine kinase (Wigler et al., *Cell* 11, 223–32, 1977) and adenine phosphoribosyltransferase (Lowy et al., *Cell* 22, 817–23, 1980) genes which can be employed in tk<sup>-</sup> or aprt<sup>-</sup> cells, respectively. Also, antimetabolite, antibiotic, or herbicide resistance can be used as the basis for selection. For example, dhfr confers resistance to methotrexate

(Wigler et al., *Proc. Natl. Acad. Sci.* 77, 3567–70, 1980), npt confers resistance to the aminoglycosides, neomycin and G-418 (Colbere-Garapin et al., *J. Mol. Biol.* 150, 1–14, 1981), and als and pat confer resistance to chlorsulfuron and phosphinotricin acetyltransferase, respectively (Murray, 1992, *supra*). Additional selectable genes have been described. For example, trpB allows cells to utilize indole in place of tryptophan, or hisD, which allows cells to utilize histinol in place of histidine (Hartman & Mulligan, *Proc. Natl. Acad. Sci.* 85, 8047–51, 1988). Visible markers such as anthocyanins,  $\beta$ -glucuronidase and its substrate GUS, and luciferase and its substrate luciferin, can be used to identify transformants and to quantify the amount of transient or stable protein expression attributable to a specific vector system (Rhodes et al., *Methods Mol. Biol.* 55, 121–131, 1995).

#### Detecting Expression

Although the presence of marker gene expression suggests that the protein kinase-like polynucleotide is also present, its presence and expression may need to be confirmed. For example, if a sequence encoding a protein kinase-like polypeptide is inserted within a marker gene sequence, transformed cells containing sequences that encode a protein kinase-like polypeptide can be identified by the absence of marker gene function. Alternatively, a marker gene can be placed in tandem with a sequence encoding a protein kinase-like polypeptide under the control of a single promoter. Expression of the marker gene in response to induction or selection usually indicates expression of the protein kinase-like polynucleotide.

Alternatively, host cells which contain a protein kinase-like polynucleotide and which express a protein kinase-like polypeptide can be identified by a variety of procedures known to those of skill in the art. These procedures include, but are not limited to, DNA-DNA or DNA-RNA hybridizations and protein bioassay or immunoassay techniques that include membrane, solution, or chip-based technologies for the detection and/or quantification of nucleic acid or protein. For example, the presence of a polynucleotide sequence encoding a protein kinase-like polypeptide can be detected by DNA-DNA or DNA-RNA hybridization or amplification using probes or fragments or fragments of polynucleotides encoding a protein kinase-like polypeptide. Nucleic acid amplification-based assays involve the use of oligonucleotides selected from sequences encoding a protein kinase-like polypeptide to detect transformants that contain a protein kinase-like polynucleotide.

A variety of protocols for detecting and measuring the expression of a protein kinase-like polypeptide, using either polyclonal or monoclonal antibodies specific for the polypeptide, are known in the art. Examples include enzyme-linked immunosorbent assay (ELISA), radioimmunoassay (RIA), and fluorescence activated cell sorting (FACS). A two-site, monoclonal-based immunoassay using monoclonal antibodies reactive to two non-interfering epitopes on a protein kinase-like polypeptide can be used, or a competitive binding assay can be employed. These and other assays are described in Hampton et al., *SEROLOGICAL METHODS: A LABORATORY MANUAL*, APS Press, St. Paul, Minn., 1990) and Maddox et al., *J. Exp. Med.* 158, 1211–1216, 1983).

A wide variety of labels and conjugation techniques are known by those skilled in the art and can be used in various nucleic acid and amino acid assays. Means for producing labeled hybridization or PCR probes for detecting sequences related to polynucleotides encoding protein kinase-like

polypeptides include oligolabeling, nick translation, end-labeling, or PCR amplification using a labeled nucleotide. Alternatively, sequences encoding a protein kinase-like polypeptide can be cloned into a vector for the production of an mRNA probe. Such vectors are known in the art, are commercially available, and can be used to synthesize RNA probes in vitro by addition of labeled nucleotides and an appropriate RNA polymerase such as T7, T3, or SP6. These procedures can be conducted using a variety of commercially available kits (Amersham Pharmacia Biotech, Promega, and US Biochemical). Suitable reporter molecules or labels which can be used for ease of detection include radionuclides, enzymes, and fluorescent, chemiluminescent, or chromogenic agents, as well as substrates, cofactors, inhibitors, magnetic particles, and the like.

#### Expression and Purification of Polypeptides

Host cells transformed with nucleotide sequences encoding a protein kinase-like polypeptide can be cultured under conditions suitable for the expression and recovery of the protein from cell culture. The polypeptide produced by a transformed cell can be secreted or contained intracellularly depending on the sequence and/or the vector used. As will be understood by those of skill in the art, expression vectors containing polynucleotides which encode protein kinase-like polypeptides can be designed to contain signal sequences which direct secretion of soluble protein kinase-like polypeptides through a prokaryotic or eukaryotic cell membrane or which direct the membrane insertion of membrane-bound protein kinase-like polypeptide.

As discussed above, other constructions can be used to join a sequence encoding a protein kinase-like polypeptide to a nucleotide sequence encoding a polypeptide domain which will facilitate purification of soluble proteins. Such purification facilitating domains include, but are not limited to, metal chelating peptides such as histidine-tryptophan modules that allow purification on immobilized metals, protein A domains that allow purification on immobilized immunoglobulin, and the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp., Seattle, Wash.). Inclusion of cleavable linker sequences such as those specific for Factor Xa or enterokinase (Invitrogen, San Diego, Calif.) between the purification domain and the protein kinase-like polypeptide also can be used to facilitate purification. One such expression vector provides for expression of a fusion protein containing a protein kinase-like polypeptide and 6 histidine residues preceding a thioredoxin or an enterokinase cleavage site. The histidine residues facilitate purification by IMAC (immobilized metal ion affinity chromatography, as described in Porath et al., *Prot. Exp. Purif.* 3, 263–281, 1992), while the enterokinase cleavage site provides a means for purifying the protein kinase-like polypeptide from the fusion protein. Vectors that contain fusion proteins are disclosed in Kroll et al., *DNA Cell Biol.* 12, 441–453, 1993.

#### Chemical Synthesis

Sequences encoding a protein kinase-like polypeptide can be synthesized, in whole or in part, using chemical methods well known in the art (see Caruthers et al., *Nucl. Acids Res. Symp. Ser.* 215–223, 1980; Horn et al. *Nucl. Acids Res. Symp. Ser.* 225–232, 1980). Alternatively, a protein kinase-like polypeptide itself can be produced using chemical methods to synthesize its amino acid sequence, such as by direct peptide synthesis using solid-phase techniques (Merrifield, *J. Am. Chem. Soc.* 85, 2149–2154, 1963; Roberge et al., *Science* 269, 202–204, 1995). Protein synthesis can be performed using manual techniques or by automation. Auto-

mated synthesis can be achieved, for example, using Applied Biosystems 431A Peptide Synthesizer (Perkin Elmer). Optionally, fragments of protein kinase-like polypeptides can be separately synthesized and combined using chemical methods to produce a full-length molecule.

The newly synthesized peptide can be substantially purified by preparative high performance liquid chromatography (e.g., Creighton, *PROTEINS: STRUCTURES AND MOLECULAR PRINCIPLES*, WH Freeman and Co., New York, N.Y., 1983). The composition of a synthetic protein kinase-like polypeptide can be confirmed by amino acid analysis or sequencing (e.g., the Edman degradation procedure; see Creighton, *supra*). Additionally, any portion of the amino acid sequence of the protein kinase-like polypeptide can be altered during direct synthesis and/or combined using chemical methods with sequences from other proteins to produce a variant polypeptide or a fusion protein.

#### Production of Altered Polypeptides

As will be understood by those of skill in the art, it may be advantageous to produce protein kinase-like polypeptide-encoding nucleotide sequences possessing non-naturally occurring codons. For example, codons preferred by a particular prokaryotic or eukaryotic host can be selected to increase the rate of protein expression or to produce an RNA transcript having desirable properties, such as a half-life that is longer than that of a transcript generated from the naturally occurring sequence.

The nucleotide sequences disclosed herein can be engineered using methods generally known in the art to alter protein kinase-like polypeptide-encoding sequences for a variety of reasons, including but not limited to, alterations which modify the cloning, processing, and/or expression of the polypeptide or mRNA product. DNA shuffling by random fragmentation and PCR reassembly of gene fragments and synthetic oligonucleotides can be used to engineer the nucleotide sequences. For example, site-directed mutagenesis can be used to insert new restriction sites, alter glycosylation patterns, change codon preference, produce splice variants, introduce mutations, and so forth.

#### Antibodies

Any type of antibody known in the art can be generated to bind specifically to an epitope of a protein kinase-like polypeptide. "Antibody" as used herein includes intact immunoglobulin molecules, as well as fragments thereof, such as Fab, F(ab')<sub>2</sub>, and Fv, which are capable of binding an epitope of a protein kinase-like polypeptide.

Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, e.g., at least 15, 25, or 50 amino acids.

An antibody which specifically binds to an epitope of a protein kinase-like polypeptide can be used therapeutically, as well as in immunochemical assays, such as Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other immunochemical assays known in the art. Various immunoassays can be used to identify antibodies having the desired specificity. Numerous protocols for competitive binding or immunoradiometric assays are well known in the art. Such immunoassays typically involve the measurement of complex formation between an immunogen and an antibody that specifically binds to the immunogen.

Typically, an antibody which specifically binds to a protein kinase-like polypeptide provides a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in an immu-

nochemical assay. Preferably, antibodies which specifically bind to protein kinase-like polypeptides do not detect other proteins in immunochemical assays and can immunoprecipitate a protein kinase-like polypeptide from solution.

Human protein kinase-like polypeptides can be used to immunize a mammal, such as a mouse, rat, rabbit, guinea pig, monkey, or human, to produce polyclonal antibodies. If desired, a protein kinase-like polypeptide can be conjugated to a carrier protein, such as bovine serum albumin, thyroglobulin, and keyhole limpet hemocyanin. Depending on the host species, various adjuvants can be used to increase the immunological response. Such adjuvants include, but are not limited to, Freund's adjuvant, mineral gels (e.g., aluminum hydroxide), and surface active substances (e.g. lysolecithin, pluronic polyols, polyanions, peptides, oil emulsions, keyhole limpet hemocyanin, and dinitrophenol). Among adjuvants used in humans, BCG (bacilli Calmette-Guerin) and *Corynebacterium parvum* are especially useful.

Monoclonal antibodies that specifically bind to a protein kinase-like polypeptide can be prepared using any technique which provides for the production of antibody molecules by continuous cell lines in culture. These techniques include, but are not limited to, the hybridoma technique, the human B-cell hybridoma technique, and the EBV-hybridoma technique (Kohler et al., *Nature* 256, 495-497, 1985; Kozbor et al., *J. Immunol. Methods* 81, 31-42, 1985; Cote et al., *Proc. Natl. Acad. Sci.* 80, 2026-2030, 1983; Cole et al., *Mol. Cell Biol.* 62, 109-120, 1984).

In addition, techniques developed for the production of "chimeric antibodies," the splicing of mouse antibody genes to human antibody genes to obtain a molecule with appropriate antigen specificity and biological activity, can be used (Morrison et al., *Proc Natl. Acad. Sci.* 81, 6851-6855, 1984; Neuberger et al., *Nature* 312, 604-608, 1984; Takeda et al., *Nature* 314, 452-454, 1985). Monoclonal and other antibodies also can be "humanized" to prevent a patient from mounting an immune response against the antibody when it is used therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences by site directed mutagenesis of individual residues or by grating of entire complementarity determining regions. Alternatively, humanized antibodies can be produced using recombinant methods, as described in GB2188638B. Antibodies that specifically bind to a protein kinase-like polypeptide can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. Pat. No. 5,565,332.

Alternatively, techniques described for the production of single chain antibodies can be adapted using methods known in the art to produce single chain antibodies that specifically bind to protein kinase-like polypeptides. Antibodies with related specificity, but of distinct idiotype composition, can be generated by chain shuffling from random combinatorial immunoglobulin libraries (Burton, *Proc. Natl. Acad. Sci.* 88, 11120-23, 1991).

Single-chain antibodies also can be constructed using a DNA amplification method, such as PCR, using hybridoma cDNA as a template (Thirion et al., 1996, *Eur. J. Cancer Prev.* 5, 507-11). Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma & Morrison, 1997, *Nat. Biotechnol.* 15,

159–63. Construction of bivalent, bispecific single-chain antibodies is taught in Mallender & Voss, 1994, *J. Biol. Chem.* 269, 199–206.

A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology (Verhaar et al., 1995, *Int. J. Cancer* 61, 497–501; Nicholls et al., 1993, *J. Immunol. Meth.* 165, 81–91).

Antibodies which specifically bind to protein kinase-like polypeptides also can be produced by inducing in vivo production in the lymphocyte population or by screening immunoglobulin libraries or panels of highly specific binding reagents as disclosed in the literature (Orlandi et al., *Proc. Natl. Acad. Sci.* 86, 3833–3837, 1989; Winter et al., *Nature* 349, 293–299, 1991).

Other types of antibodies can be constructed and used therapeutically in methods of the invention. For example, chimeric antibodies can be constructed as disclosed in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the “diabodies” described in WO 94/13804, also can be prepared.

Antibodies according to the invention can be purified by methods well known in the art. For example, antibodies can be affinity purified by passage over a column to which a protein kinase-like polypeptide is bound. The bound antibodies can then be eluted from the column using a buffer with a high salt concentration.

#### Antisense Oligonucleotides

Antisense oligonucleotides are nucleotide sequences that are complementary to a specific DNA or RNA sequence. Once introduced into a cell, the complementary nucleotides combine with natural sequences produced by the cell to form complexes and block either transcription or translation. Preferably, an antisense oligonucleotide is at least 11 nucleotides in length, but can be at least 12, 15, 20, 25, 30, 35, 40, 45, or 50 or more nucleotides long. Longer sequences also can be used. Antisense oligonucleotide molecules can be provided in a DNA construct and introduced into a cell as described above to decrease the level of protein kinase-like gene products in the cell.

Antisense oligonucleotides can be deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkyl-phosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters. See Brown, *Meth. Mol. Biol.* 20, 1–8, 1994; Sonveaux, *Meth. Mol. Biol.* 26, 1–72, 1994; Uhlmann et al., *Chem. Rev.* 90, 543–583, 1990.

Modifications of protein kinase-like gene expression can be obtained by designing antisense oligonucleotides that will form duplexes to the control, 5', or regulatory regions of the protein kinase-like gene. Oligonucleotides derived from the transcription initiation site, e.g., between positions –10 and +10 from the start site, are preferred. Similarly, inhibition can be achieved using “triple helix” base-pairing methodology. Triple helix pairing is useful because it causes inhibition of the ability of the double helix to open suffi-

ciently for the binding of polymerases, transcription factors, or chaperons. Therapeutic advances using triplex DNA have been described in the literature (e.g., Gee et al., in Huber & Carr, *MOLECULAR AND IMMUNOLOGIC APPROACHES*, Futura Publishing Co., Mt. Kisco, N.Y., 1994). An antisense oligonucleotide also can be designed to block translation of mRNA by preventing the transcript from binding to ribosomes.

Precise complementarity is not required for successful complex formation between an antisense oligonucleotide and the complementary sequence of a protein kinase-like polynucleotide. Antisense oligonucleotides which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a protein kinase-like polynucleotide, each separated by a stretch of contiguous nucleotides which are not complementary to adjacent protein kinase-like protein nucleotides, can provide sufficient targeting specificity for protein kinase-like mRNA. Preferably, each stretch of complementary contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular protein kinase-like polynucleotide sequence.

Antisense oligonucleotides can be modified without affecting their ability to hybridize to a protein kinase-like polynucleotide. These modifications can be internal or at one or both ends of the antisense molecule. For example, internucleoside phosphate linkages can be modified by adding cholesteryl or diamine moieties with varying numbers of carbon residues between the amino groups and terminal ribose. Modified bases and/or sugars, such as arabinose instead of ribose, or a 3',5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, also can be employed in a modified antisense oligonucleotide. These modified oligonucleotides can be prepared by methods well known in the art. See, e.g., Agrawal et al., *Trends Biotechnol.* 10, 152–158, 1992; Uhlmann et al., *Chem. Rev.* 90, 543–584, 1990; Uhlmann et al., *Tetrahedron. Lett.* 215, 3539–3542, 1987.

#### Ribozymes

Ribozymes are RNA molecules with catalytic activity. See, e.g., Cech, *Science* 236, 1532–1539, 1987; Cech, *Ann. Rev. Biochem.* 59, 543–568, 1990; Cech, *Curr. Opin. Struct. Biol.* 2, 605–609, 1992; Couture & Stinchcomb, *Trends Genet.* 12, 510–515, 1996. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (e.g., Haseloff et al., U.S. Pat. No. 5,641,673). The mechanism of ribozyme action involves sequence-specific hybridization of the ribozyme molecule to complementary target RNA, followed by endonucleolytic cleavage. Examples include engineered hammerhead motif ribozyme molecules that can specifically and efficiently catalyze endonucleolytic cleavage of specific nucleotide sequences.

The coding sequence of a protein kinase-like polynucleotide can be used to generate ribozymes that will specifically bind to mRNA transcribed from the protein kinase-like polynucleotide. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art (see Haseloff et al. *Nature* 334, 585–591, 1988). For example, the cleavage activity of ribozymes can be targeted to specific RNAs by engineering

a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target (see, for example, Gerlach et al., EP 321,201).

Specific ribozyme cleavage sites within a protein kinase-like protein RNA target can be identified by scanning the target molecule for ribozyme cleavage sites which include the following sequences: GUA, GUU, and GUC. Once identified, short RNA sequences of between 15 and 20 ribonucleotides corresponding to the region of the target RNA containing the cleavage site can be evaluated for secondary structural features which may render the target inoperable. Suitability of candidate protein kinase-like protein RNA targets also can be evaluated by testing accessibility to hybridization with complementary oligonucleotides using ribonuclease protection assays. Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the ribozyme can be integrally related such that upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

Ribozymes can be introduced into cells as part of a DNA construct. Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce a ribozyme-containing DNA construct into cells in which it is desired to decrease protein kinase-like protein expression. Alternatively, if it is desired that the cells stably retain the DNA construct, the construct can be supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art. A ribozyme-encoding DNA construct can include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator signal, for controlling transcription of ribozymes in the cells.

As taught in Haseloff et al., U.S. Pat. No. 5,641,673, ribozymes can be engineered so that ribozyme expression will occur in response to factors that induce expression of a target gene. Ribozymes also can be engineered to provide an additional level of regulation, so that destruction of mRNA occurs only when both a ribozyme and a target gene are induced in the cells.

#### Differentially Expressed Genes

Described herein are methods for the identification of genes whose products interact with human protein kinase-like protein. Such genes may represent genes that are differentially expressed in disorders including, but not limited to, cancer, CNS disorders, COPD, obesity, diabetes, and cardiovascular disorders

Further, such genes may represent genes that are differentially regulated in response to manipulations relevant to the progression or treatment of such diseases. Additionally, such genes may have a temporally modulated expression, increased or decreased at different stages of tissue or organism development. A differentially expressed gene may also have its expression modulated under control versus experimental conditions. In addition, the human protein kinase-like gene or gene product may itself be tested for differential expression.

The degree to which expression differs in a normal versus a diseased state need only be large enough to be visualized via standard characterization techniques such as differential display techniques. Other such standard characterization techniques by which expression differences may be visual-

ized include but are not limited to, quantitative RT (reverse transcriptase), PCR, and Northern analysis.

#### Identification of Differentially Expressed Genes

To identify differentially expressed genes total RNA or, preferably, mRNA is isolated from tissues of interest. For example, RNA samples are obtained from tissues of experimental subjects and from corresponding tissues of control subjects. Any RNA isolation technique that does not select against the isolation of mRNA may be utilized for the purification of such RNA samples. See, for example, Ausubel et al., ed., CURRENT PROTOCOLS IN MOLECULAR BIOLOGY, John Wiley & Sons, Inc. New York, 1987-1993. Large numbers of tissue samples may readily be processed using techniques well known to those of skill in the art, such as, for example, the single-step RNA isolation process of Chomczynski, U.S. Pat. No. 4,843,155.

Transcripts within the collected RNA samples that represent RNA produced by differentially expressed genes are identified by methods well known to those of skill in the art. They include, for example, differential screening (Tedder et al., *Proc. Natl. Acad. Sci. U.S.A.* 85, 208-12, 1988), subtractive hybridization (Hedrick et al., *Nature* 308, 149-53; Lee et al., *Proc. Natl. Acad. Sci. U.S.A.* 88, 2825, 1984), and, preferably, differential display (Liang & Pardee, *Science* 257, 967-71, 1992; U.S. Pat. No. 5,262,311).

The differential expression information may itself suggest relevant methods for the treatment of disorders involving the human protein kinase-like protein. For example, treatment may include a modulation of expression of the differentially expressed genes and/or the gene encoding the human protein kinase-like protein. The differential expression information may indicate whether the expression or activity of the differentially expressed gene or gene product or the human protein kinase-like gene or gene product are up-regulated or down-regulated.

#### Screening Methods

The invention provides assays for screening test compounds that bind to or modulate the activity of a protein kinase-like polypeptide or a protein kinase-like polynucleotide. A test compound preferably binds to a protein kinase-like polypeptide or polynucleotide. More preferably, a test compound decreases or increases enzyme activity by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the test compound.

#### Test Compounds

Test compounds can be pharmacologic agents already known in the art or can be compounds previously unknown to have any pharmacological activity. The compounds can be naturally occurring or designed in the laboratory. They can be isolated from microorganisms, animals, or plants, and can be produced recombinantly, or synthesized by chemical methods known in the art. If desired, test compounds can be obtained using any of the numerous combinatorial library methods known in the art, including but not limited to, biological libraries, spatially addressable parallel solid phase or solution phase libraries, synthetic library methods requiring deconvolution, the "one-bead one-compound" library method, and synthetic library methods using affinity chromatography selection. The biological library approach is limited to polypeptide libraries, while the other four approaches are applicable to polypeptide, non-peptide oligomer, or small molecule libraries of compounds. See Lam, *Anticancer Drug Des.* 12, 145, 1997.

Methods for the synthesis of molecular libraries are well known in the art (see, for example, DeWitt et al., *Proc. Natl.*

*Acad. Sci. U.S.A.* 90, 6909, 1993; Erb et al. *Proc. Natl. Acad. Sci. U.S.A.* 91, 11422, 1994; Zuckermann et al., *J. Med. Chem.* 37, 2678, 1994; Cho et al., *Science* 261, 1303, 1993; Carell et al., *Angew. Chem. Int. Ed. Engl.* 33, 2059, 1994; Carell et al., *Angew. Chem. Int. Ed. Engl.* 33, 2061, 1994; Gallop et al., *J. Med. Chem.* 37, 1233, 1994). Libraries of compounds can be presented in solution (see, e.g., Houghten, *BioTechniques* 13, 412–421, 1992), or on beads (Lam, *Nature* 354, 82–84, 1991), chips (Fodor, *Nature* 364, 555–556, 1993), bacteria or spores (Ladner, U.S. Pat. No. 5,223,409), plasmids (Cull et al., *Proc. Natl. Acad. Sci. U.S.A.* 89, 1865–1869, 1992), or phage (Scott & Smith, *Science* 249, 386–390, 1990; Devlin, *Science* 249, 404–406, 1990); Cwirla et al., *Proc. Natl. Acad. Sci.* 97, 6378–6382, 1990; Felici, *J. Mol. Biol.* 222, 301–310, 1991; and Ladner, U.S. Pat. No. 5,223,409).

#### High Throughput Screening

Test compounds can be screened for the ability to bind to protein kinase-like polypeptides or polynucleotides or to affect protein kinase-like protein activity or protein kinase-like gene expression using high throughput screening. Using high throughput screening, many discrete compounds can be tested in parallel so that large numbers of test compounds can be quickly screened. The most widely established techniques utilize 96-well microtiter plates. The wells of the microtiter plates typically require assay volumes that range from 50 to 500  $\mu$ l. In addition to the plates, many instruments, materials, pipettors, robotics, plate washers, and plate readers are commercially available to fit the 96-well format.

Alternatively, “free format assays,” or assays that have no physical barrier between samples, can be used. For example, an assay using pigment cells (melanocytes) in a simple homogeneous assay for combinatorial peptide libraries is described by Jayawickreme et al., *Proc. Natl. Acad. Sci. U.S.A.* 19, 1614–18 (1994). The cells are placed under agarose in petri dishes, then beads that carry combinatorial compounds are placed on the surface of the agarose. The combinatorial compounds are partially released the compounds from the beads. Active compounds can be visualized as dark pigment areas because, as the compounds diffuse locally into the gel matrix, the active compounds cause the cells to change colors.

Another example of a free format assay is described by Chelsky, “Strategies for Screening Combinatorial Libraries: Novel and Traditional Approaches,” reported at the First Annual Conference of The Society for Biomolecular Screening in Philadelphia, Pa. (Nov. 7–10, 1995). Chelsky placed a simple homogenous enzyme assay for carbonic anhydrase inside an agarose gel such that the enzyme in the gel would cause a color change throughout the gel. Thereafter, beads carrying combinatorial compounds via a photolinker were placed inside the gel and the compounds were partially released by UV-light. Compounds that inhibited the enzyme were observed as local zones of inhibition having less color change.

Yet another example is described by Salmon et al., *Molecular Diversity* 2, 57–63 (1996). In this example, combinatorial libraries were screened for compounds that had cytotoxic effects on cancer cells growing in agar.

Another high throughput screening method is described in Beutel et al., U.S. Pat. No. 5,976,813. In this method, test samples are placed in a porous matrix. One or more assay components are then placed within, on top of, or at the bottom of a matrix such as a gel, a plastic sheet, a filter, or other form of easily manipulated solid support. When samples are introduced to the porous matrix they diffuse

sufficiently slowly, such that the assays can be performed without the test samples running together.

#### Binding Assays

For binding assays, the test compound is preferably a small molecule that binds to and occupies, for example, the active site of the protein kinase-like polypeptide, such that normal biological activity is prevented. Examples of such small molecules include, but are not limited to, small peptides or peptide-like molecules.

In binding assays, either the test compound or the protein kinase-like polypeptide can comprise a detectable label, such as a fluorescent, radioisotopic, chemiluminescent, or enzymatic label, such as horseradish peroxidase, alkaline phosphatase, or luciferase. Detection of a test compound that is bound to the protein kinase-like polypeptide can then be accomplished, for example, by direct counting of radio-emission, by scintillation counting, or by determining conversion of an appropriate substrate to a detectable product.

Alternatively, binding of a test compound to a protein kinase-like polypeptide can be determined without labeling either of the interactants. For example, a microphysiometer can be used to detect binding of a test compound with a protein kinase-like polypeptide. A microphysiometer (e.g., Cytosensor™) is an analytical instrument that measures the rate at which a cell acidifies its environment using a light-addressable potentiometric sensor (LAPS). Changes in this acidification rate can be used as an indicator of the interaction between a test compound and a protein kinase-like polypeptide (McConnell et al., *Science* 257, 1906–1912, 1992).

Determining the ability of a test compound to bind to a protein kinase-like polypeptide also can be accomplished using a technology such as real-time Bimolecular Interaction Analysis (BIA) (Sjolander & Urbaniczky, *Anal. Chem.* 63, 2338–2345, 1991, and Szabo et al., *Curr. Opin. Struct. Biol.* 5, 699–705, 1995). BIA is a technology for studying biospecific interactions in real time, without labeling any of the interactants (e.g., BIAcore™). Changes in the optical phenomenon surface plasmon resonance (SPR) can be used as an indication of real-time reactions between biological molecules.

In yet another aspect of the invention, a protein kinase-like polypeptide can be used as a “bait protein” in a two-hybrid assay or three-hybrid assay (see, e.g., U.S. Pat. No. 5,283,317; Zervos et al., *Cell* 72, 223–232, 1993; Madura et al., *J. Biol. Chem.* 268, 12046–12054, 1993; Bartel et al., *BioTechniques* 14, 920–924, 1993; Iwabuchi et al., *Oncogene* 8, 1693–1696, 1993; and Brent WO94/10300), to identify other proteins which bind to or interact with the protein kinase-like polypeptide and modulate its activity.

The two-hybrid system is based on the modular nature of most transcription factors, which consist of separable DNA-binding and activation domains. Briefly, the assay utilizes two different DNA constructs. For example, in one construct, polynucleotide encoding a protein kinase-like polypeptide can be fused to a polynucleotide encoding the DNA binding domain of a known transcription factor (e.g., GAL-4). In the other construct a DNA sequence that encodes an unidentified protein (“prey” or “sample”) can be fused to a polynucleotide that codes for the activation domain of the known transcription factor. If the “bait” and the “prey” proteins are able to interact in vivo to form an protein-dependent complex, the DNA-binding and activation domains of the transcription factor are brought into close

25

proximity. This proximity allows transcription of a reporter gene (e.g., LacZ), which is operably linked to a transcriptional regulatory site responsive to the transcription factor. Expression of the reporter gene can be detected, and cell colonies containing the functional transcription factor can be isolated and used to obtain the DNA sequence encoding the protein that interacts with the protein kinase-like polypeptide.

It may be desirable to immobilize either the protein kinase-like polypeptide (or polynucleotide) or the test compound to facilitate separation of bound from unbound forms of one or both of the interactants, as well as to accommodate automation of the assay. Thus, either the protein kinase-like polypeptide (or polynucleotide) or the test compound can be bound to a solid support. Suitable solid supports include, but are not limited to, glass or plastic slides, tissue culture plates, microtiter wells, tubes, silicon chips, or particles such as beads (including, but not limited to, latex, polystyrene, or glass beads). Any method known in the art can be used to attach the enzyme polypeptide (or polynucleotide) or test compound to a solid support, including use of covalent and non-covalent linkages, passive absorption, or pairs of binding moieties attached respectively to the polypeptide (or polynucleotide) or test compound and the solid support. Test compounds are preferably bound to the solid support in an array, so that the location of individual test compounds can be tracked. Binding of a test compound to a protein kinase-like polypeptide (or polynucleotide) can be accomplished in any vessel suitable for containing the reactants. Examples of such vessels include microtiter plates, test tubes, and microcentrifuge tubes.

In one embodiment, the protein kinase-like polypeptide is a fusion protein comprising a domain that allows the protein kinase-like polypeptide to be bound to a solid support. For example, glutathione-S-transferase fusion proteins can be adsorbed onto glutathione sepharose beads (Sigma Chemical, St. Louis, Mo.) or glutathione derivatized microtiter plates, which are then combined with the test compound or the test compound and the non-adsorbed protein kinase-like polypeptide; the mixture is then incubated under conditions conducive to complex formation (e.g., at physiological conditions for salt and pH). Following incubation, the beads or microtiter plate wells are washed to remove any unbound components. Binding of the interactants can be determined either directly or indirectly, as described above. Alternatively, the complexes can be dissociated from the solid support before binding is determined.

Other techniques for immobilizing proteins or polynucleotides on a solid support also can be used in the screening assays of the invention. For example, either a protein kinase-like polypeptide (or polynucleotide) or a test compound can be immobilized utilizing conjugation of biotin and streptavidin. Biotinylated protein kinase-like polypeptides (or polynucleotides) or test compounds can be prepared from biotin-NHS(N-hydroxysuccinimide) using techniques well known in the art (e.g., biotinylation kit, Pierce Chemicals, Rockford, Ill.) and immobilized in the wells of streptavidin-coated 96 well plates (Pierce Chemical). Alternatively, antibodies which specifically bind to a protein kinase-like polypeptide, polynucleotide, or a test compound, but which do not interfere with a desired binding site, such as the active site of the protein kinase-like polypeptide, can be derivatized to the wells of the plate. Unbound target or protein can be trapped in the wells by antibody conjugation.

Methods for detecting such complexes, in addition to those described above for the GST-immobilized complexes, include immunodetection of complexes using antibodies

26

which specifically bind to the protein kinase-like polypeptide or test compound, enzyme-linked assays which rely on detecting an activity of the protein kinase-like polypeptide, and SDS gel electrophoresis under non-reducing conditions.

Screening for test compounds which bind to a protein kinase-like polypeptide or polynucleotide also can be carried out in an intact cell. Any cell which comprises a protein kinase-like polypeptide or polynucleotide can be used in a cell-based assay system. A protein kinase-like polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Binding of the test compound to a protein kinase-like polypeptide or polynucleotide is determined as described above.

#### Enzyme Assays

Test compounds can be tested for the ability to increase or decrease the protein kinase activity of a human protein kinase-like polypeptide. Protein kinase activity can be measured, for example, as described in U.S. Pat. No. 6,194,186

Enzyme assays can be carried out after contacting either a purified protein kinase-like polypeptide, a cell membrane preparation, or an intact cell with a test compound. A test compound that decreases a protein kinase activity of a protein kinase-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential therapeutic agent for decreasing protein kinase-like protein activity. A test compound which increases a protein kinase activity of a human protein kinase-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% is identified as a potential therapeutic agent for increasing human protein kinase-like protein activity.

#### Gene Expression

In another embodiment, test compounds that increase or decrease protein kinase-like gene expression are identified. A protein kinase-like polynucleotide is contacted with a test compound, and the expression of an RNA or polypeptide product of the protein kinase-like polynucleotide is determined. The level of expression of appropriate mRNA or polypeptide in the presence of the test compound is compared to the level of expression of mRNA or polypeptide in the absence of the test compound. The test compound can then be identified as a modulator of expression based on this comparison. For example, when expression of mRNA or polypeptide is greater in the presence of the test compound than in its absence, the test compound is identified as a stimulator or enhancer of the mRNA or polypeptide expression. Alternatively, when expression of the mRNA or polypeptide is less in the presence of the test compound than in its absence, the test compound is identified as an inhibitor of the mRNA or polypeptide expression.

The level of protein kinase-like mRNA or polypeptide expression in the cells can be determined by methods well known in the art for detecting mRNA or polypeptide. Either qualitative or quantitative methods can be used. The presence of polypeptide products of a protein kinase-like polynucleotide can be determined, for example, using a variety of techniques known in the art, including immunochemical methods such as radioimmunoassay, Western blotting, and immunohistochemistry. Alternatively, polypeptide synthesis can be determined in vivo, in a cell culture, or in an in vitro translation system by detecting incorporation of labeled amino acids into a protein kinase-like polypeptide.

Such screening can be carried out either in a cell-free assay system or in an intact cell. Any cell that expresses a protein kinase-like polynucleotide can be used in a cell-

based assay system. The protein kinase-like polynucleotide can be naturally occurring in the cell or can be introduced using techniques such as those described above. Either a primary culture or an established cell line, such as CHO or human embryonic kidney 293 cells, can be used.

#### Pharmaceutical Compositions

The invention also provides pharmaceutical compositions that can be administered to a patient to achieve a therapeutic effect. Pharmaceutical compositions of the invention can comprise, for example, a protein kinase-like polypeptide, protein kinase-like polynucleotide, ribozymes or antisense oligonucleotides, antibodies which specifically bind to a protein kinase-like polypeptide, or mimetics, activators, or inhibitors of a protein kinase-like polypeptide activity. The compositions can be administered alone or in combination with at least one other agent, such as stabilizing compound, which can be administered in any sterile, biocompatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions can be administered to a patient alone, or in combination with other agents, drugs or hormones.

In addition to the active ingredients, these pharmaceutical compositions can contain suitable pharmaceutically-acceptable carriers comprising excipients and auxiliaries that facilitate processing of the active compounds into preparations which can be used pharmaceutically. Pharmaceutical compositions of the invention can be administered by any number of routes including, but not limited to, oral, intravenous, intramuscular, intra-arterial, intramedullary, intrathecal, intraventricular, transdermal, subcutaneous, intraperitoneal, intranasal, parenteral, topical, sublingual, or rectal means. Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees, capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents can be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores can be used in conjunction with suitable coatings, such as concentrated sugar solutions, which also can contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments can be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound, i.e., dosage.

Pharmaceutical preparations that can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium stearate, and,

optionally, stabilizers. In soft capsules, the active compounds can be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration can be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks' solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions can contain substances that increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Additionally, suspensions of the active compounds can be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Non-lipid polycationic amino polymers also can be used for delivery. Optionally, the suspension also can contain suitable stabilizers or agents that increase the solubility of the compounds to allow for the preparation of highly concentrated solutions. For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention can be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes. The pharmaceutical composition can be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation can be a lyophilized powder which can contain any or all of the following: 1–50 mM histidine, 0.1%–2% sucrose, and 2–7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

Further details on techniques for formulation and administration can be found in the latest edition of REMINGTON'S PHARMACEUTICAL SCIENCES (Maack Publishing Co., Easton, Pa.). After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

#### Therapeutic Indications and Methods

Human protein kinase-like protein can be regulated to treat cancer, CNS disorders, COPD, obesity, diabetes, and cardiovascular disorders.

#### Cancer

Cancer is a disease fundamentally caused by oncogenic cellular transformation. There are several hallmarks of transformed cells that distinguish them from their normal counterparts and underlie the pathophysiology of cancer. These include uncontrolled cellular proliferation, unresponsiveness to normal death-inducing signals (immortalization), increased cellular motility and invasiveness, increased ability to recruit blood supply through induction of new blood vessel formation (angiogenesis), genetic instability, and dysregulated gene expression. Various combinations of these aberrant physiologies, along with the acquisition of drug-resistance frequently lead to an intractable disease state in which organ failure and patient death ultimately ensue.

Most standard cancer therapies target cellular proliferation and rely on the differential proliferative capacities

between transformed and normal cells for their efficacy. This approach is hindered by the facts that several important normal cell types are also highly proliferative and that cancer cells frequently become resistant to these agents. Thus, the therapeutic indices for traditional anti-cancer therapies rarely exceed 2.0.

The advent of genomics-driven molecular target identification has opened up the possibility of identifying new cancer-specific targets for therapeutic intervention that will provide safer, more effective treatments for cancer patients. Thus, newly discovered tumor-associated genes and their products can be tested for their role(s) in disease and used as tools to discover and develop innovative therapies. Genes playing important roles in any of the physiological processes outlined above can be characterized as cancer targets.

Genes or gene fragments identified through genomics can readily be expressed in one or more heterologous expression systems to produce functional recombinant proteins. These proteins are characterized in vitro for their biochemical properties and then used as tools in high-throughput molecular screening programs to identify chemical modulators of their biochemical activities. Activators and/or inhibitors of target protein activity can be identified in this manner and subsequently tested in cellular and in vivo disease models for anti-cancer activity. Optimization of lead compounds with iterative testing in biological models and detailed pharmacokinetic and toxicological analyses form the basis for drug development and subsequent testing in humans.

Protein phosphorylation is an essential component in intracellular signaling, with diverse and crucial functions including mediation of cell proliferation, survival, apoptosis, differentiation, migration and attachment. It is regulated by the balance between the opposing activities of protein kinases and protein phosphatases. Protein phosphorylation is mainly mediated by two types of protein kinases—protein tyrosine kinases and protein serine/threonine kinases. A number of protein tyrosine kinases are encoded by proto-oncogenes or viral oncogenes, and are thus strongly implicated in cancer. Protein serine/threonine kinases are known to play a role in intracellular signal transduction mediated by growth factors, cytokines, etc. inducing either cell proliferation, apoptosis or differentiation. Inhibitors of protein kinases are expected to provide efficacious therapeutic agents for the treatment of cancer.

#### CNS Disorders

Central and peripheral nervous system disorders also can be treated, such as primary and secondary disorders after brain injury, disorders of mood, anxiety disorders, disorders of thought and volition, disorders of sleep and wakefulness, diseases of the motor unit, such as neurogenic and myopathic disorders, neurodegenerative disorders such as Alzheimer's and Parkinson's disease, and processes of peripheral and chronic pain.

Pain that is associated with CNS disorders also can be treated by regulating the activity of human protein kinase-like protein. Pain which can be treated includes that associated with central nervous system disorders, such as multiple sclerosis, spinal cord injury, sciatica, failed back surgery syndrome, traumatic brain injury, epilepsy, Parkinson's disease, post-stroke, and vascular lesions in the brain and spinal cord (e.g., infarct, hemorrhage, vascular malformation). Non-central neuropathic pain includes that associated with post mastectomy pain, reflex sympathetic dystrophy (RSD), trigeminal neuralgia, radioculopathy, post-surgical pain, HIV/AIDS related pain, cancer pain, metabolic neuropathies (e.g., diabetic neuropathy, vasculitic

neuropathy secondary to connective tissue disease), paraneoplastic polyneuropathy associated, for example, with carcinoma of lung, or leukemia, or lymphoma, or carcinoma of prostate, colon or stomach, trigeminal neuralgia, cranial neuralgias, and post-herpetic neuralgia. Pain associated with cancer and cancer treatment also can be treated, as can headache pain (for example, migraine with aura, migraine without aura, and other migraine disorders), episodic and chronic tension-type headache, tension-type like headache, cluster headache, and chronic paroxysmal hemicrania.

#### COPD

Chronic obstructive pulmonary (or airways) disease (COPD) is a condition defined physiologically as airflow obstruction that generally results from a mixture of emphysema and peripheral airway obstruction due to chronic bronchitis (Senior & Shapiro, *Pulmonary Diseases and Disorders*, 3d ed., New York, McGraw-Hill, 1998, pp. 659–681, 1998; Barnes, *Chest* 117, 10S–14S, 2000). Emphysema is characterized by destruction of alveolar walls leading to abnormal enlargement of the air spaces of the lung. Chronic bronchitis is defined clinically as the presence of chronic productive cough for three months in each of two successive years. In COPD, airflow obstruction is usually progressive and is only partially reversible. By far the most important risk factor for development of COPD is cigarette smoking, although the disease does occur in non-smokers.

Chronic inflammation of the airways is a key pathological feature of COPD (Senior & Shapiro, 1998). The inflammatory cell population comprises increased numbers of macrophages, neutrophils, and CD8<sup>+</sup> lymphocytes. Inhaled irritants, such as cigarette smoke, activate macrophages which are resident in the respiratory tract, as well as epithelial cells leading to release of chemokines (e.g., interleukin-8) and other chemotactic factors. These chemotactic factors act to increase the neutrophil/-monocyte trafficking from the blood into the lung tissue and airways. Neutrophils and monocytes recruited into the airways can release a variety of potentially damaging mediators such as proteolytic enzymes and reactive oxygen species. Matrix degradation and emphysema, along with airway wall thickening, surfactant dysfunction, and mucus hypersecretion, all are potential sequelae of this inflammatory response that lead to impaired airflow and gas exchange.

Obesity and overweight are defined as an excess of body fat relative to lean body mass. An increase in caloric intake or a decrease in energy expenditure or both can bring about this imbalance leading to surplus energy being stored as fat. Obesity is associated with important medical morbidities and an increase in mortality. The causes of obesity are poorly understood and may be due to genetic factors, environmental factors or a combination of the two to cause a positive energy balance. In contrast, anorexia and cachexia are characterized by an imbalance in energy intake versus energy expenditure leading to a negative energy balance and weight loss. Agents that either increase energy expenditure and/or decrease energy intake, absorption or storage would be useful for treating obesity, overweight, and associated comorbidities. Agents that either increase energy intake and/or decrease energy expenditure or increase the amount of lean tissue would be useful for treating cachexia, anorexia and wasting disorders.

This gene, translated proteins and agents which modulate this gene or portions of the gene or its products are useful for treating obesity, overweight, anorexia, cachexia, wasting disorders, appetite suppression, appetite enhancement, increases or decreases in satiety, modulation of body weight,

and/or other eating disorders such as bulimia. Also this gene, translated proteins and agents which modulate this gene or portions of the gene or its products are useful for treating obesity/overweight-associated comorbidities including hypertension, type 2 diabetes, coronary artery disease, hyperlipidemia, stroke, gallbladder disease, gout, osteoarthritis, sleep apnea and respiratory problems, some types of cancer including endometrial, breast, prostate, and colon cancer, thrombotic disease, polycystic ovarian syndrome, reduced fertility, complications of pregnancy, menstrual irregularities, hirsutism, stress incontinence, and depression.

#### Diabetes

Diabetes mellitus is a common metabolic disorder characterized by an abnormal elevation in blood glucose, alterations in lipids and abnormalities (complications) in the cardiovascular system, eye, kidney and nervous system. Diabetes is divided into two separate diseases: type 1 diabetes (juvenile onset), which results from a loss of cells which make and secrete insulin, and type 2 diabetes (adult onset), which is caused by a defect in insulin secretion and a defect in insulin action.

Type 1 diabetes is initiated by an autoimmune reaction that attacks the insulin secreting cells (beta cells) in the pancreatic islets. Agents that prevent this reaction from occurring or that stop the reaction before destruction of the beta cells has been accomplished are potential therapies for this disease. Other agents that induce beta cell proliferation and regeneration also are potential therapies.

Type II diabetes is the most common of the two diabetic conditions (6% of the population). The defect in insulin secretion is an important cause of the diabetic condition and results from an inability of the beta cell to properly detect and respond to rises in blood glucose levels with insulin release. Therapies that increase the response by the beta cell to glucose would offer an important new treatment for this disease.

The defect in insulin action in Type II diabetic subjects is another target for therapeutic intervention. Agents that increase the activity of the insulin receptor in muscle, liver, and fat will cause a decrease in blood glucose and a normalization of plasma lipids. The receptor activity can be increased by agents that directly stimulate the receptor or that increase the intracellular signals from the receptor. Other therapies can directly activate the cellular end process, i.e. glucose transport or various enzyme systems, to generate an insulin-like effect and therefore a produce beneficial outcome. Because overweight subjects have a greater susceptibility to Type II diabetes, any agent that reduces body weight is a possible therapy.

Both Type I and Type diabetes can be treated with agents that mimic insulin action or that treat diabetic complications by reducing blood glucose levels. Likewise, agents that reduces new blood vessel growth can be used to treat the eye complications that develop in both diseases.

#### Cardiovascular Diseases

Cardiovascular diseases include the following disorders of the heart and the vascular system: congestive heart failure, myocardial infarction, ischemic diseases of the heart, all kinds of atrial and ventricular arrhythmias, hypertensive vascular diseases, and peripheral vascular diseases.

Heart failure is defined as a pathophysiologic state in which an abnormality of cardiac function is responsible for the failure of the heart to pump blood at a rate commensurate with the requirement of the metabolizing tissue. It includes all forms of pumping failure, such as high-output and

low-output, acute and chronic, right-sided or left-sided, systolic or diastolic, independent of the underlying cause.

Myocardial infarction (MI) is generally caused by an abrupt decrease in coronary blood flow that follows a thrombotic occlusion of a coronary artery previously narrowed by arteriosclerosis. MI prophylaxis (primary and secondary prevention) is included, as well as the acute treatment of MI and the prevention of complications.

Ischemic diseases are conditions in which the coronary flow is restricted resulting in a perfusion which is inadequate to meet the myocardial requirement for oxygen. This group of diseases includes stable angina, unstable angina, and asymptomatic ischemia.

Arrhythmias include all forms of atrial and ventricular tachyarrhythmias (atrial tachycardia, atrial flutter, atrial fibrillation, atrio-ventricular reentrant tachycardia, preexcitation syndrome, ventricular tachycardia, ventricular flutter, and ventricular fibrillation), as well as bradycardic forms of arrhythmias.

Vascular diseases include primary as well as all kinds of secondary arterial hypertension (renal, endocrine, neurogenic, others). The disclosed gene and its product may be used as drug targets for the treatment of hypertension as well as for the prevention of all complications. Peripheral vascular diseases are defined as vascular diseases in which arterial and/or venous flow is reduced resulting in an imbalance between blood supply and tissue oxygen demand. It includes chronic peripheral arterial occlusive disease (PAOD), acute arterial thrombosis and embolism, inflammatory vascular disorders, Raynaud's phenomenon, and venous disorders.

This invention further pertains to the use of novel agents identified by the screening assays described above. Accordingly, it is within the scope of this invention to use a test compound identified as described herein in an appropriate animal model. For example, an agent identified as described herein (e.g., a modulating agent, an antisense nucleic acid molecule, a specific antibody, ribozyme, or a protein kinase-like polypeptide binding molecule) can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent identified as described herein can be used in an animal model to determine the mechanism of action of such an agent. Furthermore, this invention pertains to uses of novel agents identified by the above-described screening assays for treatments as described herein.

A reagent which affects protein kinase-like protein activity can be administered to a human cell, either in vitro or in vivo, to reduce protein kinase-like protein activity. The reagent preferably binds to an expression product of a human protein kinase-like gene. If the expression product is a protein, the reagent is preferably an antibody. For treatment of human cells ex vivo, an antibody can be added to a preparation of stem cells that have been removed from the body. The cells can then be replaced in the same or another human body, with or without clonal propagation, as is known in the art.

In one embodiment, the reagent is delivered using a liposome. Preferably, the liposome is stable in the animal into which it has been administered for at least about 30 minutes, more preferably for at least about 1 hour, and even more preferably for at least about 24 hours. A liposome comprises a lipid composition that is capable of targeting a reagent, particularly a polynucleotide, to a particular site in an animal, such as a human. Preferably, the lipid composition of the liposome is capable of targeting to a specific

organ of an animal, such as the lung, liver, spleen, heart brain, lymph nodes, and skin.

A liposome useful in the present invention comprises a lipid composition that is capable of fusing with the plasma membrane of the targeted cell to deliver its contents to the cell. Preferably, the transfection efficiency of a liposome is about 0.5  $\mu\text{g}$  of DNA per 16 nmole of liposome delivered to about  $10^6$  cells, more preferably about 1.0  $\mu\text{g}$  of DNA per 16 nmole of liposome delivered to about  $10^6$  cells, and even more preferably about 2.0  $\mu\text{g}$  of DNA per 16 nmol of liposome delivered to about  $10^6$  cells. Preferably, a liposome is between about 100 and 500 nm, more preferably between about 150 and 450 nm, and even more preferably between about 200 and 400 nm in diameter.

Suitable liposomes for use in the present invention include those liposomes standardly used in, for example, gene delivery methods known to those of skill in the art. More preferred liposomes include liposomes having a polycationic lipid composition and/or liposomes having a cholesterol backbone conjugated to polyethylene glycol. Optionally, a liposome comprises a compound capable of targeting the liposome to a particular cell type, such as a cell-specific ligand exposed on the outer surface of the liposome.

Complexing a liposome with a reagent such as an antisense oligonucleotide or ribozyme can be achieved using methods that are standard in the art (see, for example, U.S. Pat. No. 5,705,151). Preferably, from about 0.1  $\mu\text{g}$  to about 10  $\mu\text{g}$  of polynucleotide is combined with about 8 nmol of liposomes, more preferably from about 0.5  $\mu\text{g}$  to about 5  $\mu\text{g}$  of polynucleotides are combined with about 8 nmol liposomes, and even more preferably about 1.0  $\mu\text{g}$  of polynucleotides is combined with about 8 mol liposomes.

In another embodiment, antibodies can be delivered to specific tissues in vivo using receptor-mediated targeted delivery. Receptor-mediated DNA delivery techniques are taught in, for example, Findeis et al. *Trends in Biotechnol.* 11, 202–05 (1993); Chiou et al., *GENE THERAPEUTICS: METHODS AND APPLICATIONS OF DIRECT GENE TRANSFER* (J. A. Wolff, ed.) (1994); Wu & Wu, *J. Biol. Chem.* 263, 621–24 (1988); Wu et al., *J. Biol. Chem.* 269, 542–46 (1994); Zenke et al., *Proc. Natl. Acad. Sci. U.S.A.* 87, 3655–59 (1990); Wu et al., *J. Biol. Chem.* 266, 338–42 (1991).

#### Determination of a Therapeutically Effective Dose

The determination of a therapeutically effective dose is well within the capability of those skilled in the art. A therapeutically effective dose refers to that amount of active ingredient which increases or decreases protein kinase-like protein activity relative to the protein kinase-like protein activity which occurs in the absence of the therapeutically effective dose.

For any compound, the therapeutically effective dose can be estimated initially either in cell culture assays or in animal models, usually mice, rabbits, dogs, or pigs. The animal model also can be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

Therapeutic efficacy and toxicity, e.g.,  $\text{ED}_{50}$  (the dose therapeutically effective in 50% of the population) and  $\text{LD}_{50}$  (the dose lethal to 50% of the population), can be determined by standard pharmaceutical procedures in cell cultures or experimental animals. The dose ratio of toxic to therapeutic effects is the therapeutic index, and it can be expressed as the ratio,  $\text{LD}_{50}/\text{ED}_{50}$ .

Pharmaceutical compositions that exhibit large therapeutic indices are preferred. The data obtained from cell culture assays and animal studies is used in formulating a range of dosage for human use. The dosage contained in such compositions is preferably within a range of circulating concentrations that include the  $\text{ED}_{50}$  with little or no toxicity. The dosage varies within this range depending upon the dosage form employed, sensitivity of the patient, and the route of administration.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active ingredient or to maintain the desired effect. Factors that can be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and tolerance/response to therapy. Long-acting pharmaceutical compositions can be administered every 3 to 4 days, every week, or once every two weeks depending on the half-life and clearance rate of the particular formulation.

Normal dosage amounts can vary from 0.1 to 100,000 micrograms, up to a total dose of about 1 g, depending upon the route of administration. Guidance as to particular dosages and methods of delivery is provided in the literature and generally available to practitioners in the art. Those skilled in the art will employ different formulations for nucleotides than for proteins or their inhibitors. Similarly, delivery of polynucleotides or polypeptides will be specific to particular cells, conditions, locations, etc.

If the reagent is a single-chain antibody, polynucleotides encoding the antibody can be constructed and introduced into a cell either ex vivo or in vivo using well-established techniques including, but not limited to, transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, protoplast fusion, viral infection, electroporation, “gene gun,” and DEAE- or calcium phosphate-mediated transfection.

Effective in vivo dosages of an antibody are in the range of about 5  $\mu\text{g}$  to about 50  $\mu\text{g}/\text{kg}$ , about 50  $\mu\text{g}$  to about 5 mg/kg, about 100  $\mu\text{g}$  to about 500  $\mu\text{g}/\text{kg}$  of patient body weight, and about 200 to about 250  $\mu\text{g}/\text{kg}$  of patient body weight. For administration of polynucleotides encoding single-chain antibodies, effective in vivo dosages are in the range of about 100 ng to about 200 ng, 500 ng to about 50 mg, about 1  $\mu\text{g}$  to about 2 mg, about 5  $\mu\text{g}$  to about 500  $\mu\text{g}$ , and about 20  $\mu\text{g}$  to about 100  $\mu\text{g}$  of DNA.

If the expression product is mRNA, the reagent is preferably an antisense oligo-nucleotide or a ribozyme. Polynucleotides that express antisense oligonucleotides or ribozymes can be introduced into cells by a variety of methods, as described above.

Preferably, a reagent reduces expression of a protein kinase-like gene or the activity of a protein kinase-like polypeptide by at least about 10, preferably about 50, more preferably about 75, 90, or 100% relative to the absence of the reagent. The effectiveness of the mechanism chosen to decrease the level of expression of a protein kinase-like gene or the activity of a protein kinase-like polypeptide can be assessed using methods well known in the art, such as hybridization of nucleotide probes to protein kinase-like protein-specific mRNA, quantitative RT-PCR, immunologic detection of a protein kinase-like polypeptide, or measurement of protein kinase-like protein activity.

In any of the embodiments described above, any of the pharmaceutical compositions of the invention can be administered in combination with other appropriate therapeutic agents. Selection of the appropriate agents for use in combination therapy can be made by one of ordinary skill in the art, according to conventional pharmaceutical principles. The combination of therapeutic agents can act synergistically to effect the treatment or prevention of the various disorders described above. Using this approach, one may be able to achieve therapeutic efficacy with lower dosages of each agent, thus reducing the potential for adverse side effects.

Any of the therapeutic methods described above can be applied to any subject in need of such therapy, including, for example, mammals such as dogs, cats, cows, horses, rabbits, monkeys, and most preferably, humans.

#### Diagnostic Methods

Human protein kinase-like protein also can be used in diagnostic assays for detecting diseases and abnormalities or susceptibility to diseases and abnormalities related to the presence of mutations in the nucleic acid sequences that encode the enzyme. For example, differences can be determined between the cDNA or genomic sequence encoding protein kinase-like protein in individuals afflicted with a disease and in normal individuals. If a mutation is observed in some or all of the afflicted individuals but not in normal individuals, then the mutation is likely to be the causative agent of the disease.

Sequence differences between a reference gene and a gene having mutations can be revealed by the direct DNA sequencing method. In addition, cloned DNA segments can be employed as probes to detect specific DNA segments. The sensitivity of this method is greatly enhanced when combined with PCR. For example, a sequencing primer can be used with a double-stranded PCR product or a single-stranded template molecule generated by a modified PCR. The sequence determination is performed by conventional procedures using radiolabeled nucleotides or by automatic sequencing procedures using fluorescent tags.

Genetic testing based on DNA sequence differences can be carried out by detection of alteration in electrophoretic mobility of DNA fragments in gels with or without denaturing agents. Small sequence deletions and insertions can be visualized, for example, by high resolution gel electrophoresis. DNA fragments of different sequences can be distinguished on denaturing formamide gradient gels in which the mobilities of different DNA fragments are retarded in the gel at different positions according to their specific melting or partial melting temperatures (see, e.g., Myers et al., *Science* 230, 1242, 1985). Sequence changes at specific locations can also be revealed by nuclease protection assays, such as RNase and S1 protection or the chemical cleavage method (e.g., Cotton et al., *Proc. Natl. Acad. Sci. USA* 85, 4397-4401, 1985). Thus, the detection of a specific DNA sequence can be performed by methods such as hybridization, RNase protection, chemical cleavage, direct DNA sequencing or the use of restriction enzymes and Southern blotting of genomic DNA. In addition to direct methods such as gel-electrophoresis and DNA sequencing, mutations can also be detected by in situ analysis.

Altered levels of protein kinase-like protein also can be detected in various tissues. Assays used to detect levels of the receptor polypeptides in a body sample, such as blood or a tissue biopsy, derived from a host are well known to those

of skill in the art and include radioimmunoassays, competitive binding assays, Western blot analysis, and ELISA assays.

All patents, patent applications, and references cited in this disclosure are expressly incorporated herein by reference in their entireties. The above disclosure generally describes the present invention. A more complete understanding can be obtained by reference to the following specific examples, which are provided for purposes of illustration only and are not intended to limit the scope of the invention.

#### EXAMPLE 1

##### Detection of Proteine Kinase-like Protein Activity

The polynucleotide of SEQ ID NO: 1 is inserted into the expression vector pCEV4 and the expression vector pCEV4-proteine kinase-like protein polypeptide obtained is transfected into human embryonic kidney 293 cells. From these cells extracts are obtained and proteine kinase-like protein activity is assayed by measuring the incorporation of  $^{32}\text{P}$  into calf thymus H1 histone from  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ . Cell extracts are incubated in a reaction mixture containing 20 mM Tris HCl at pH 7.5, 10 mM  $\text{MgCl}_2$ , 20  $\mu\text{M}$  ATP, 15-50 kBq of  $[\gamma\text{-}^{32}\text{P}]\text{ATP}$ , and 200  $\mu\text{g/ml}$  H1 istone. The incubation is carried out for 5 min at 30° C., and the phosphorylated proteins are seperated by SDS/PAGE and visualized and quantitated by measuring the intensity of photostimulated luminescence (PSL) using a Bio-Imaging Analyzer. It is shown that the polypeptide of SEQ ID NO: 2 has a proteine kinase-like protein activity.

#### EXAMPLE 2

##### Expression of Recombinant Human Protein Kinase-like Protein

The *Pichia pastoris* expression vector pPICZB (Invitrogen, San Diego, Calif.) is used to produce large quantities of recombinant human protein kinase-like polypeptides in yeast. The protein kinase-like protein-encoding DNA sequence is derived from SEQ ID NOS: 1, 8 and 10. Before insertion into vector pPICZB, the DNA sequence is modified by well known methods in such a way that it contains at its 5'-end an initiation codon and at its 3'-end an enterokinase cleavage site, a His6 reporter tag and a termination codon. Moreover, at both termini recognition sequences for restriction endonucleases are added and after digestion of the multiple cloning site of pPICZ B with the corresponding restriction enzymes the modified DNA sequence is ligated into pPICZB. This expression vector is designed for inducible expression in *Pichia pastoris*, driven by a yeast promoter. The resulting pPICZ/md-His6 vector is used to transform the yeast.

The yeast is cultivated under usual conditions in 5 liter shake flasks and the recombinantly produced protein isolated from the culture by affinity chromatography (Ni-NTA-Resin) in the presence of 8 M urea. The bound polypeptide is eluted with buffer, pH 3.5, and neutralized. Separation of the polypeptide from the His6 reporter tag is accomplished by site-specific proteolysis using enterokinase (Invitrogen, San Diego, Calif.) according to manufacturer's instructions. Purified human protein kinase-like polypeptide is obtained.

#### Identification of Test Compounds that Bind to Protein Kinase-like Polypeptides

Purified protein kinase-like polypeptides comprising a glutathione-S-transferase protein and absorbed onto glutathione-derivatized wells of 96-well microtiter plates are contacted with test compounds from a small molecule library at pH 7.0 in a physiological buffer solution. Human protein kinase-like polypeptides comprise the amino acid sequence shown in SEQ ID NOS:2 and 9. The test compounds comprise a fluorescent tag. The samples are incubated for 5 minutes to one hour. Control samples are incubated in the absence of a test compound.

The buffer solution containing the test compounds is washed from the wells. Binding of a test compound to a protein kinase-like polypeptide is detected by fluorescence measurements of the contents of the wells. A test compound that increases the fluorescence in a well by at least 15% relative to fluorescence of a well in which a test compound is not incubated is identified as a compound which binds to a protein kinase-like polypeptide.

#### EXAMPLE 4

#### Identification of a Test Compound which Decreases Protein Kinase-like Gene Expression

A test compound is administered to a culture of human cells transfected with a protein kinase-like protein expression construct and incubated at 37° C. for 10 to 45 minutes. A culture of the same type of cells that have not been transfected is incubated for the same time without the test compound to provide a negative control.

RNA is isolated from the two cultures as described in Chirgwin et al., *Biochem.* 18, 5294–99, 1979). Northern blots are prepared using 20 to 30 µg total RNA and hybridized with a <sup>32</sup>P-labeled protein kinase-like protein-specific probe at 65° C. in Express-hyb (CLONTECH). The probe comprises at least 11 contiguous nucleotides selected from the complement of SEQ ID NOS: 1, 8 and 10. A test compound that decreases the protein kinase-like protein-specific signal relative to the signal obtained in the absence of the test compound is identified as an inhibitor of protein kinase-like gene expression.

#### EXAMPLE 5

#### Identification of a Test Compound which Decreases Protein Kinase-like Protein Activity

A test compound is administered to a culture of human cells transfected with a protein kinase-like protein expression construct and incubated at 37° C. for 10 to 45 minutes. A culture of the same type of cells that have not been transfected is incubated for the same time without the test compound to provide a negative control. protein kinase-like protein activity is measured using the method of U.S. Pat. No. 6,194,186.

A test compound which decreases the protein kinase-like protein activity of the protein kinase-like protein relative to the protein kinase-like protein activity in the absence of the test compound is identified as an inhibitor of protein kinase-like protein activity.

#### Tissue-Specific Expression of Protein Kinase-like Protein

The qualitative expression pattern of protein kinase-like protein in various tissues is determined by Reverse Transcription-Polymerase Chain Reaction (RT-PCR).

To demonstrate that protein kinase-like protein is involved in cancer, expression is determined in the following tissues: adrenal gland, bone marrow, brain, cerebellum, colon, fetal brain, fetal liver, heart, kidney, liver, lung, mammary gland, pancreas, placenta, prostate, salivary gland, skeletal muscle, small intestine, spinal cord, spleen, stomach, testis, thymus, thyroid, trachea, uterus, and peripheral blood lymphocytes. Expression in the following cancer cell lines also is determined: DU-145 (prostate), NCI-H125 (lung), HT-29 (colon), COLO-205 (colon), A-549 (lung), NCI-H460 (lung), HT-116 (colon), DLD-1 (colon), MDA-MD-231 (breast), LS 174T (colon), ZF-75 (breast), MDA-MN-435 (breast), HT-1080, MCF-7 (breast), and U87. Matched pairs of malignant and normal tissue from the same patient also are tested.

To demonstrate that protein kinase-like protein is involved in CNS disorders, the following tissues are screened: fetal and adult brain, muscle, heart, lung, kidney, liver, thymus, testis, colon, placenta, trachea, pancreas, kidney, gastric mucosa, colon, liver, cerebellum, skin, cortex (Alzheimer's and normal), hypothalamus, cortex, amygdala, cerebellum, hippocampus, choroid, plexus, thalamus, and spinal cord.

To demonstrate that protein kinase-like protein is involved in the disease process of COPD, the initial expression panel consists of RNA samples from respiratory tissues and inflammatory cells relevant to COPD: lung (adult and fetal), trachea, freshly isolated alveolar type II cells, cultured human bronchial epithelial cells, cultured small airway epithelial cells, cultured bronchial smooth muscle cells, cultured H441 cells (Clara-like), freshly isolated neutrophils and monocytes, and cultured monocytes (macrophage-like). Body map profiling also is carried out, using total RNA panels purchased from Clontech. The tissues are adrenal gland, bone marrow, brain, colon, heart, kidney, liver, lung, mammary gland, pancreas, prostate, salivary gland, skeletal muscle, small intestine, spleen, stomach, testis, thymus, trachea, thyroid, and uterus.

To demonstrate that protein kinase-like protein is involved in the disease process of obesity, expression is determined in the following tissues: subcutaneous adipose tissue, mesenteric adipose tissue, adrenal gland, bone marrow, brain (cerebellum, spinal cord, cerebral cortex, caudate, medulla, substantia nigra, and putamen), colon, fetal brain, heart, kidney, liver, lung, mammary gland, pancreas, placenta, prostate, salivary gland, skeletal muscle small intestine, spleen, stomach, testes, thymus, thyroid trachea, and uterus. Neuroblastoma cell lines SK-Nr-Be (2), Hr, Sk-N-As, HTB-10, IMR-32, SNSY-5Y, T3, SK-N-D2, D283, DAOY, CHP-2, U87MG, BE(2)C, T986, KANTS, M059K, CHP234, C6 (rat), SK-N-F1, SK-PU-DW, PFSK-1, BE(2) M17, and MCIXC also are tested for protein kinase-like protein expression. As a final step, the expression of protein kinase-like protein in cells derived from normal individuals with the expression of cells derived from obese individuals is compared.

To demonstrate that protein kinase-like protein is involved in the disease process of diabetes, the following whole body panel is screened to show predominant or relatively high expression: subcutaneous and mesenteric

adipose tissue, adrenal gland, bone marrow, brain, colon, fetal brain, heart, hypothalamus, kidney, liver, lung, mammary gland, pancreas, placenta, prostate, salivary gland, skeletal muscle, small intestine, spleen, stomach, testis, thymus, thyroid, trachea, and uterus. Human islet cells and an islet cell library also are tested. As a final step, the expression of protein kinase-like protein in cells derived from normal individuals with the expression of cells derived from diabetic individuals is compared.

Quantitative expression profiling. Quantitative expression profiling is performed by the form of quantitative PCR analysis called "kinetic analysis" firstly described in Higuchi et al., *BioTechnology* 10, 413-17, 1992, and Higuchi et al., *BioTechnology* 11, 1026-30, 1993. The principle is that at any given cycle within the exponential phase of PCR, the amount of product is proportional to the initial number of template copies.

If the amplification is performed in the presence of an internally quenched fluorescent oligonucleotide (TaqMan probe) complementary to the target sequence, the probe is cleaved by the 5'-3' endonuclease activity of Taq DNA polymerase and a fluorescent dye released in the medium (Holland et al., *Proc. Natl. Acad. Sci. U.S.A.* 88, 7276-80, 1991). Because the fluorescence emission will increase in direct proportion to the amount of the specific amplified product, the exponential growth phase of PCR product can be detected and used to determine the initial template concentration (Heid et al., *Genome Res.* 6, 986-94, 1996, and Gibson et al., *Genome Res.* 6, 995-1001, 1996).

The amplification of an endogenous control can be performed to standardize the amount of sample RNA added to a reaction. In this kind of experiment, the control of choice is the 18S ribosomal RNA. Because reporter dyes with differing emission spectra are available, the target and the endogenous control can be independently quantified in the same tube if probes labeled with different dyes are used.

All "real time PCR" measurements of fluorescence are made in the ABI Prism 7700.

RNA extraction and cDNA preparation. Total RNA from the tissues listed above are used for expression quantification. RNAs labeled "from autopsy" were extracted from autaptic tissues with the TRIzol reagent (Life Technologies, MD) according to the manufacturer's protocol.

Fifty µg of each RNA were treated with DNase I for 1 hour at 37° C. in the following reaction mix: 0.2 U/µl RNase-free DNase I (Roche Diagnostics, Germany); 0.4 U/µl RNase inhibitor (PE Applied Biosystems, CA); 10 mM Tris-HCl pH 7.9; 10 mM MgCl<sub>2</sub>; 50 mM NaCl; and 1 mM DTT.

After incubation, RNA is extracted once with 1 volume of phenol:chloroform:—isoamyl alcohol (24:24:1) and once with chloroform, and precipitated with 1/10 volume of 3 M NaAcetate, pH5.2, and 2 volumes of ethanol.

Fifty µg of each RNA from the autaptic tissues are DNase treated with the DNA-free kit purchased from Ambion (Ambion, Tex.). After resuspension and spectrophotometric quantification, each sample is reverse transcribed with the TaqMan Reverse Transcription Reagents (PE Applied Biosystems, CA) according to the manufacturer's protocol. The final concentration of RNA in the reaction mix is 200 ng/µL. Reverse transcription is carried out with 2.5 µM of random hexamer primers.

TaqMan quantitative analysis. Specific primers and probe are designed according to the recommendations of PE Applied Biosystems; the probe can be labeled at the 5' end with FAM (6-carboxy-fluorescein) and at the 3' end with TAMRA (6-carboxy-tetramethyl-rhodamine). Quantification experiments are performed on 10 ng of reverse transcribed RNA from each sample. Each determination is done in triplicate.

Total cDNA content is normalized with the simultaneous quantification (multiplex PCR) of the 18S ribosomal RNA using the Pre-Developed TaqMan Assay Reagents (PDAR) Control Kit (PE Applied Biosystems, CA).

The assay reaction mix is as follows: 1× final TaqMan Universal PCR Master Mix (from 2× stock) (PE Applied Biosystems, CA); 1× PDAR control —18S RNA (from 20× stock); 300 nM forward primer; 900 nM reverse primer; 200 nM probe; 10 ng cDNA; and water to 25 µl.

Each of the following steps are carried out once: pre PCR, 2 minutes at 50° C., and 10 minutes at 95° C. The following steps are carried out 40 times: denaturation, 15 seconds at 95° C., annealing/extension, 1 minute at 60° C.

The experiment is performed on an ABI Prism 7700 Sequence Detector (PE Applied Biosystems, CA). At the end of the run, fluorescence data acquired during PCR are processed as described in the ABI Prism 7700 user's manual in order to achieve better background subtraction as well as signal linearity with the starting target quantity.

## EXAMPLE 7

Proliferation Inhibition Assay: Antisense Oligonucleotides Suppress the Growth of Cancer Cell Lines

The cell line used for testing is the human colon cancer cell line HCT116. Cells are cultured in RPMI-1640 with 10-15% fetal calf serum at a concentration of 10,000 cells per milliliter in a volume of 0.5 ml and kept at 37° C. in a 95% air/5% CO<sub>2</sub> atmosphere.

Phosphorothioate oligoribonucleotides are synthesized on an Applied Biosystems Model 380B DNA synthesizer using phosphoroamidite chemistry. A sequence of 24 bases complementary to the nucleotides at position 1 to 24 of SEQ ID NOS: 1, 8 and 10 is used as the test oligonucleotide. As a control, another (random) sequence is used: 5'-TCA ACT GAC TAG ATG TAC ATG GAC-3'. Following assembly and deprotection, oligonucleotides are ethanol-precipitated twice, dried, and suspended in phosphate buffered saline at the desired concentration. Purity of the oligonucleotides is tested by capillary gel electrophoresis and ion exchange HPLC. The purified oligonucleotides are added to the culture medium at a concentration of 10 µM once per day for seven days.

The addition of the test oligonucleotide for seven days results in significantly reduced expression of human protein kinase-like protein as determined by Western blotting. This effect is not observed with the control oligonucleotide. After 3 to 7 days, the number of cells in the cultures is counted using an automatic cell counter. The number of cells in cultures treated with the test oligonucleotide (expressed as 100%) is compared with the number of cells in cultures treated with the control oligonucleotide. The number of cells in cultures treated with the test oligonucleotide is not more than 30% of control, indicating that the inhibition of human protein kinase-like protein has an anti-proliferative effect on cancer cells.

## EXAMPLE 8

In Vivo Testing of Compounds/Target Validation

### 1. Acute Mechanistic Assays

#### 1.1. Reduction in Mitogenic Plasma Hormone Levels

This non-tumor assay measures the ability of a compound to reduce either the endogenous level of a circulating hormone or the level of hormone produced in response to a biologic stimulus. Rodents are administered test compound (p.o., i.p., i.v., i.m., or s.c.). At a predetermined time after

administration of test compound, blood plasma is collected. Plasma is assayed for levels of the hormone of interest. If the normal circulating levels of the hormone are too low and/or variable to provide consistent results, the level of the hormone may be elevated by a pre-treatment with a biologic stimulus (i.e., LHRH may be injected i.m. into mice at a dosage of 30 ng/mouse to induce a burst of testosterone synthesis). The timing of plasma collection would be adjusted to coincide with the peak of the induced hormone response. Compound effects are compared to a vehicle-treated control group. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test. Significance is  $p \leq 0.05$  compared to the vehicle control group.

### 1.2. Hollow Fiber Mechanism of Action Assay

Hollow fibers are prepared with desired cell line(s) and implanted intraperitoneally and/or subcutaneously in rodents. Compounds are administered p.o., i.p., i.v., i.m., or s.c. Fibers are harvested in accordance with specific readout assay protocol, these may include assays for gene expression (bDNA, PCR, or Taqman), or a specific biochemical activity (i.e., cAMP levels. Results are analyzed by Student's t-test or Rank Sum test after the variance between groups is compared by an F-test, with significance at  $p < 0.05$  as compared to the vehicle control group.

## 2. Subacute Functional In Vivo Assays

### 2.1. Reduction in Mass of Hormone Dependent Tissues

This is another non-tumor assay that measures the ability of a compound to reduce the mass of a hormone dependent tissue (i.e., seminal vesicles in males and uteri in females). Rodents are administered test compound (p.o., i.p., i.v., i.m., or s.c.) according to a predetermined schedule and for a predetermined duration (i.e., 1 week). At termination of the study, animals are weighed, the target organ is excised, any fluid is expressed, and the weight of the organ is recorded. Blood plasma may also be collected. Plasma may be assayed for levels of a hormone of interest or for levels of test agent. Organ weights may be directly compared or they may be normalized for the body weight of the animal. Compound effects are compared to a vehicle-treated control group. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test. Significance is  $p \leq 0.05$  compared to the vehicle control group.

### 2.2. Hollow Fiber Proliferation Assay

Hollow fibers are prepared with desired cell line(s) and implanted intraperitoneally and/or subcutaneously in rodents. Compounds are administered p.o., i.p., i.v., i.m., or s.c. Fibers are harvested in accordance with specific readout assay protocol. Cell proliferation is determined by measuring a marker of cell number (i.e., MTT or LDH). The cell number and change in cell number from the starting inoculum are analyzed by Student's t-test or Rank Sum test after the variance between groups is compared by an F-test, with significance at  $p \leq 0.05$  as compared to the vehicle control group.

## 2.3 Anti-angiogenesis Models

### 2.3.1. Corneal Angiogenesis

Hydron pellets with or without growth factors or cells are implanted into a micropocket surgically created in the rodent cornea. Compound administration may be systemic or local (compound mixed with growth factors in the hydron pellet). Corneas are harvested at 7 days post implantation immediately following intracardiac infusion of colloidal carbon and are fixed in 10% formalin. Readout is qualitative scoring

and/or image analysis. Qualitative scores are compared by Rank Sum test. Image analysis data is evaluated by measuring the area of neovascularization (in pixels) and group averages are compared by Student's t-test (2 tail). Significance is  $p < 0.05$  as compared to the growth factor or cells only group.

### 2.3.2. Matrigel Angiogenesis

Matrigel, containing cells or growth factors, is injected subcutaneously. Compounds are administered p.o., i.p., i.v., i.m., or s.c. Matrigel plugs are harvested at predetermined time point(s) and prepared for readout. Readout is an ELISA-based assay for hemoglobin concentration and/or histological examination (i.e. vessel count, special staining for endothelial surface markers: CD31, factor-8). Readouts are analyzed by Student's t-test, after the variance between groups is compared by an F-test, with significance determined at  $p \leq 0.05$  as compared to the vehicle control group.

## 3. Primary Antitumor Efficacy

### 3.1. Early Therapy Models

#### 3.1.1. Subcutaneous Tumor

Tumor cells or fragments are implanted subcutaneously on Day 0. Vehicle and/or compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule starting at a time, usually on Day 1, prior to the ability to measure the tumor burden. Body weights and tumor measurements are recorded 2–3 times weekly. Mean net body and tumor weights are calculated for each data collection day. Anti-tumor efficacy may be initially determined by comparing the size of treated (T) and control (C) tumors on a given day by a Student's t-test, after the variance between groups is compared by an F-test, with significance determined at  $p \leq 0.05$ . The experiment may also be continued past the end of dosing in which case tumor measurements would continue to be recorded to monitor tumor growth delay. Tumor growth delays are expressed as the difference in the median time for the treated and control groups to attain a predetermined size divided by the median time for the control group to attain that size. Growth delays are compared by generating Kaplan-Meier curves from the times for individual tumors to attain the evaluation size. Significance is  $p < 0.05$ .

#### 3.1.2. Intraperitoneal/Intracranial Tumor Models

Tumor cells are injected intraperitoneally or intracranially on Day 0. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule starting on Day 1. Observations of morbidity and/or mortality are recorded twice daily. Body weights are measured and recorded twice weekly. Morbidity/mortality data is expressed in terms of the median time of survival and the number of long-term survivors is indicated separately. Survival times are used to generate Kaplan-Meier curves. Significance is  $p \leq 0.05$  by a log-rank test compared to the control group in the experiment.

### 3.2. Established Disease Model

Tumor cells or fragments are implanted subcutaneously and grown to the desired size for treatment to begin. Once at the predetermined size range, mice are randomized into treatment groups. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule. Tumor and body weights are measured and recorded 2–3 times weekly. Mean tumor weights of all groups over days post inoculation are graphed for comparison. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test to compare tumor sizes in the

treated and control groups at the end of treatment. Significance is  $p \leq 0.05$  as compared to the control group. Tumor measurements may be recorded after dosing has stopped to monitor tumor growth delay. Tumor growth delays are expressed as the difference in the median time for the treated and control groups to attain a predetermined size divided by the median time for the control group to attain that size. Growth delays are compared by generating Kaplan-Meier curves from the times for individual tumors to attain the evaluation size. Significance is  $p \text{ value} \leq 0.05$  compared to the vehicle control group.

### 3.3. Orthotopic Disease Models

#### 3.3.1. Mammary Fat Pad Assay

Tumor cells or fragments, of mammary adenocarcinoma origin, are implanted directly into a surgically exposed and reflected mammary fat pad in rodents. The fat pad is placed back in its original position and the surgical site is closed. Hormones may also be administered to the rodents to support the growth of the tumors. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule. Tumor and body weights are measured and recorded 2–3 times weekly. Mean tumor weights of all groups over days post inoculation are graphed for comparison. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test to compare tumor sizes in the treated and control groups at the end of treatment. Significance is  $p \leq 0.05$  as compared to the control group.

Tumor measurements may be recorded after dosing has stopped to monitor tumor growth delay. Tumor growth delays are expressed as the difference in the median time for the treated and control groups to attain a predetermined size divided by the median time for the control group to attain that size. Growth delays are compared by generating Kaplan-Meier curves from the times for individual tumors to attain the evaluation size. Significance is  $p \text{ value} \leq 0.05$  compared to the vehicle control group. In addition, this model provides an opportunity to increase the rate of spontaneous metastasis of this type of tumor. Metastasis can be assessed at termination of the study by counting the number of visible foci per target organ, or measuring the target organ weight. The means of these endpoints are compared by Student's t-test after conducting an F-test, with significance determined at  $p \leq 0.05$  compared to the control group in the experiment.

#### 3.3.2. Intraprostatic Assay

Tumor cells or fragments, of prostatic adenocarcinoma origin, are implanted directly into a surgically exposed dorsal lobe of the prostate in rodents. The prostate is externalized through an abdominal incision so that the tumor can be implanted specifically in the dorsal lobe while verifying that the implant does not enter the seminal vesicles. The successfully inoculated prostate is replaced in the abdomen and the incisions through the abdomen and skin are closed. Hormones may also be administered to the rodents to support the growth of the tumors. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule. Body weights are measured and recorded 2–3 times weekly. At a predetermined time, the experiment is terminated and the animal is dissected. The size of the primary tumor is measured in three dimensions using either a caliper or an ocular micrometer attached to a dissecting scope. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test to compare tumor sizes in the treated and control groups at the

end of treatment. Significance is  $p \leq 0.05$  as compared to the control group. This model provides an opportunity to increase the rate of spontaneous metastasis of this type of tumor. Metastasis can be assessed at termination of the study by counting the number of visible foci per target organ (i.e., the lungs), or measuring the target organ weight (i.e., the regional lymph nodes). The means of these endpoints are compared by Student's t-test after conducting an F-test, with significance determined at  $p \leq 0.05$  compared to the control group in the experiment.

#### 3.3.3. Intrabronchial Assay

Tumor cells of pulmonary origin may be implanted intrabronchially by making an incision through the skin and exposing the trachea. The trachea is pierced with the beveled end of a 25 gauge needle and the tumor cells are inoculated into the main bronchus using a flat-ended 27 gauge needle with a 90° bend. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule. Body weights are measured and recorded 2–3 times weekly. At a predetermined time, the experiment is terminated and the animal is dissected. The size of the primary tumor is measured in three dimensions using either a caliper or an ocular micrometer attached to a dissecting scope. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test to compare tumor sizes in the treated and control groups at the end of treatment. Significance is  $p \leq 0.05$  as compared to the control group. This model provides an opportunity to increase the rate of spontaneous metastasis of this type of tumor. Metastasis can be assessed at termination of the study by counting the number of visible foci per target organ (i.e., the contralateral lung), or measuring the target organ weight. The means of these endpoints are compared by Student's t-test after conducting an F-test, with significance determined at  $p < 0.05$  compared to the control group in the experiment.

#### 3.3.4. Intracecal Assay

Tumor cells of gastrointestinal origin may be implanted intracecally by making an abdominal incision through the skin and externalizing the intestine. Tumor cells are inoculated into the cecal wall without penetrating the lumen of the intestine using a 27 or 30 gauge needle. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule. Body weights are measured and recorded 2–3 times weekly. At a predetermined time, the experiment is terminated and the animal is dissected. The size of the primary tumor is measured in three dimensions using either a caliper or an ocular micrometer attached to a dissecting scope. An F-test is preformed to determine if the variance is equal or unequal followed by a Student's t-test to compare tumor sizes in the treated and control groups at the end of treatment. Significance is  $p \leq 0.05$  as compared to the control group. This model provides an opportunity to increase the rate of spontaneous metastasis of this type of tumor. Metastasis can be assessed at termination of the study by counting the number of visible foci per target organ (i.e., the liver), or measuring the target organ weight. The means of these endpoints are compared by Student's t-test after conducting an F-test, with significance determined at  $p \leq 0.05$  compared to the control group in the experiment.

### 4. Secondary (Metastatic) Antitumor Efficacy

#### 4.1. Spontaneous Metastasis

Tumor cells are inoculated s.c. and the tumors allowed to grow to a predetermined range for spontaneous metastasis studies to the lung or liver. These primary tumors are then excised. Compounds are administered p.o., i.p., i.v., i.m., or

s.c. according to a predetermined schedule which may include the period leading up to the excision of the primary tumor to evaluate therapies directed at inhibiting the early stages of tumor metastasis. Observations of morbidity and/or mortality are recorded daily. Body weights are measured and recorded twice weekly. Potential endpoints include survival time, numbers of visible foci per target organ, or target organ weight. When survival time is used as the endpoint the other values are not determined. Survival data is used to generate Kaplan-Meier curves. Significance is  $p \leq 0.05$  by a log-rank test compared to the control group in the experiment. The mean number of visible tumor foci, as determined under a dissecting microscope, and the mean target organ weights are compared by Student's t-test after conducting an F-test, with significance determined at  $p \leq 0.05$  compared to the control group in the experiment for both of these endpoints.

#### 4.2. Forced Metastasis

Tumor cells are injected into the tail vein, portal vein, or the left ventricle of the heart in experimental (forced) lung, liver, and bone metastasis studies, respectively. Compounds are administered p.o., i.p., i.v., i.m., or s.c. according to a predetermined schedule. Observations of morbidity and/or mortality are recorded daily. Body weights are measured and recorded twice weekly. Potential endpoints include survival time, numbers of visible foci per target organ, or target organ weight. When survival time is used as the endpoint the other values are not determined. Survival data is used to generate Kaplan-Meier curves. Significance is  $p \leq 0.05$  by a log-rank test compared to the control group in the experiment. The mean number of visible tumor foci, as determined under a dissecting microscope, and the mean target organ weights are compared by Student's t-test after conducting an F-test, with significance at  $p \leq 0.05$  compared to the vehicle control group in the experiment for both endpoints.

### EXAMPLE 9

#### In Vivo Testing of Compounds/Target Validation

##### 1. Pain

###### Acute Pain

Acute pain is measured on a hot plate mainly in rats. Two variants of hot plate testing are used: In the classical variant animals are put on a hot surface (52 to 56° C.) and the latency time is measured until the animals show nocifensive behavior, such as stepping or foot licking. The other variant is an increasing temperature hot plate where the experimental animals are put on a surface of neutral temperature. Subsequently this surface is slowly but constantly heated until the animals begin to lick a hind paw. The temperature which is reached when hind paw licking begins is a measure for pain threshold.

Compounds are tested against a vehicle treated control group. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

###### Persistent Pain

Persistent pain is measured with the formalin or capsaicin test, mainly in rats. A solution of 1 to 5% formalin or 10 to 100 µg capsaicin is injected into one hind paw of the experimental animal. After formalin or capsaicin application the animals show nocifensive reactions like flinching, licking and biting of the affected paw. The number of nocifensive reactions within a time frame of up to 90 minutes is a measure for intensity of pain.

Compounds are tested against a vehicle treated control group. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to formalin or capsaicin administration.

###### Neuropathic Pain

Neuropathic pain is induced by different variants of unilateral sciatic nerve injury mainly in rats. The operation is performed under anesthesia. The first variant of sciatic nerve injury is produced by placing loosely constrictive ligatures around the common sciatic nerve. The second variant is the tight ligation of about the half of the diameter of the common sciatic nerve. In the next variant, a group of models is used in which tight ligations or transections are made of either the L5 and L6 spinal nerves, or the L % spinal nerve only. The fourth variant involves an axotomy of two of the three terminal branches of the sciatic nerve (tibial and common peroneal nerves) leaving the remaining sural nerve intact whereas the last variant comprises the axotomy of only the tibial branch leaving the sural and common nerves uninjured. Control animals are treated with a sham operation.

Postoperatively, the nerve injured animals develop a chronic mechanical allodynia, cold allodynia, as well as a thermal hyperalgesia. Mechanical allodynia is measured by means of a pressure transducer (electronic von Frey Anesthesiometer, IITC Inc.-Life Science Instruments, Woodland Hills, SA, USA; Electronic von Frey System, Somedic Sales AB, Hörby, Sweden). Thermal hyperalgesia is measured by means of a radiant heat source (Plantar Test, Ugo Basile, Comerio, Italy), or by means of a cold plate of 5 to 10° C. where the nocifensive reactions of the affected hind paw are counted as a measure of pain intensity. A further test for cold induced pain is the counting of nocifensive reactions, or duration of nocifensive responses after plantar administration of acetone to the affected hind limb. Chronic pain in general is assessed by registering the circadian rhythms in activity (Suijo and Arndt, Universität zu Köln, Cologne, Germany), and by scoring differences in gait (foot print patterns; FOOTPRINTS program, Klapdor et al., 1997. A low cost method to analyze footprint patterns. J. Neurosci. Methods 75, 49-54).

Compounds are tested against sham operated and vehicle treated control groups. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

###### Inflammatory Pain

Inflammatory pain is induced mainly in rats by injection of 0.75 mg carrageenan or complete Freund's adjuvant into one hind paw. The animal develop an edema with mechanical allodynia as well as thermal hyperalgesia.

Mechanical allodynia is measured by means of a pressure transducer (electronic von Frey Anesthesiometer, IITC Inc.-Life Science Instruments, Woodland Hills, SA, USA). Thermal hyperalgesia is measured by means of a radiant heat source (Plantar Test, Ugo Basile, Comerio, Italy, Paw thermal stimulator, G. Ozaki, University of California, USA). For edema measurement two methods are being used. In the first method, the animals are sacrificed and the affected hindpaws sectioned and weighed. The second method comprises differences in paw volume by measuring water displacement in a plethysmometer (Ugo Basile, Comerio, Italy).

Compounds are tested against uninflamed as well as vehicle treated control groups. Substance application is

performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

#### Diabetic Neuropathic Pain

Rats treated with a single intraperitoneal injection of 50 to 80 mg/kg streptozotocin develop a profound hyperglycemia and mechanical allodynia within 1 to 3 weeks. Mechanical allodynia is measured by means of a pressure transducer (electronic von Frey Anesthesiometer, IITC Inc.-Life Science Instruments, Woodland Hills, SA, USA).

Compounds are tested against diabetic and non-diabetic vehicle treated control groups. Substance application is performed at different time points via different application routes (i.v., i.p., p.o., i.t., i.c.v., s.c., intradermal, transdermal) prior to pain testing.

## 2. Parkinson's Disease

### 6-Hydroxydopamine (6-OH-DA) Lesion

Degeneration of the dopaminergic nigrostriatal and striatopallidal pathways is the central pathological event in Parkinson's disease. This disorder has been mimicked experimentally in rats using single/sequential unilateral stereotaxic injections of 6-OH-DA into the medium forebrain bundle (MFB).

Male Wistar rats (Harlan Winkelmann, Germany), weighing  $200 \pm 250$  g at the beginning of the experiment, are used. The rats are maintained in a temperature- and humidity-controlled environment under a 12 h light/dark cycle with free access to food and water when not in experimental sessions. The following in vivo protocols are approved by the governmental authorities. All efforts are made to minimize animal suffering, to reduce the number of animals used, and to utilize alternatives to in vivo techniques.

Animals are administered pargyline on the day of surgery (Sigma, St. Louis, Mo., USA; 50 mg/kg i.p.) in order to inhibit metabolism of 6-OHDA by monoamine oxidase and desmethylimipramine HCl (Sigma; 25 mg/kg i.p.) in order to prevent uptake of 6-OHDA by noradrenergic terminals. Thirty minutes later the rats are anesthetized with sodium pentobarbital (50 mg/kg) and placed in a stereotaxic frame. In order to lesion the DA nigrostriatal pathway 4  $\mu$ l of 0.01% ascorbic acid-saline containing 8  $\mu$ g of 6-OHDA HBr (Sigma) are injected into the left medial fore-brain bundle at a rate of 1  $\mu$ l/min (2.4 mm anterior, 1.49 mm lateral, -2.7 mm ventral to Bregma and the skull surface). The needle is left in place an additional 5 min to allow diffusion to occur.

#### Stepping Test

Forelimb akinesia is assessed three weeks following lesion placement using a modified stepping test protocol. In brief, the animals are held by the experimenter with one hand fixing the hindlimbs and slightly raising the hind part above the surface. One paw is touching the table, and is then moved slowly sideways (5 s for 1 m), first in the forehand and then in the backhand direction. The number of adjusting steps is counted for both paws in the backhand and forehand direction of movement. The sequence of testing is right paw forehand and backhand adjusting stepping, followed by left paw forehand and backhand directions. The test is repeated three times on three consecutive days, after an initial training period of three days prior to the first testing. Forehand adjusted stepping reveals no consistent differences between lesioned and healthy control animals. Analysis is therefore restricted to backhand adjusted stepping.

#### Balance Test

Balance adjustments following postural challenge are also measured during the stepping test sessions. The rats are held in the same position as described in the stepping test and,

instead of being moved sideways, tilted by the experimenter towards the side of the paw touching the table. This maneuver results in loss of balance and the ability of the rats to regain balance by forelimb movements is scored on a scale ranging from 0 to 3. Score 0 is given for a normal forelimb placement. When the forelimb movement is delayed but recovery of postural balance detected, score 1 is given. Score 2 represents a clear, yet insufficient, forelimb reaction, as evidenced by muscle contraction, but lack of success in recovering balance, and score 3 is given for no reaction of movement. The test is repeated three times a day on each side for three consecutive days after an initial training period of three days prior to the first testing.

#### Staircase Test (Paw Reaching)

A modified version of the staircase test is used for evaluation of paw reaching behavior three weeks following primary and secondary lesion placement. Plexiglass test boxes with a central platform and a removable staircase on each side are used. The apparatus is designed such that only the paw on the same side at each staircase can be used, thus providing a measure of independent forelimb use. For each test the animals are left in the test boxes for 15 min. The double staircase is filled with 7x3 chow pellets (Precision food pellets, formula: P, purified rodent diet, size 45 mg; Sandown Scientific) on each side. After each test the number of pellets eaten (successfully retrieved pellets) and the number of pellets taken (touched but dropped) for each paw and the success rate (pellets eaten/pellets taken) are counted separately. After three days of food deprivation (12 g per animal per day) the animals are tested for 11 days. Full analysis is conducted only for the last five days.

#### MPTP Treatment

The neurotoxin 1-methyl-4-phenyl-1,2,3,6-tetrahydro-pyridine (MPTP) causes degeneration of mesencephalic dopaminergic (DAergic) neurons in rodents, non-human primates, and humans and, in so doing, reproduces many of the symptoms of Parkinson's disease. MPTP leads to a marked decrease in the levels of dopamine and its metabolites, and in the number of dopaminergic terminals in the striatum as well as severe loss of the tyrosine hydroxylase (TH)-immunoreactive cell bodies in the substantia nigra, pars compacta.

In order to obtain severe and long-lasting lesions, and to reduce mortality, animals receive single injections of MPTP, and are then tested for severity of lesion 7-10 days later. Successive MPTP injections are administered on days 1, 2 and 3. Animals receive application of 4 mg/kg MPTP hydrochloride (Sigma) in saline once daily. All injections are intraperitoneal (i.p.) and the MPTP stock solution is frozen between injections. Animals are decapitated on day 11.

#### Immunohistology

At the completion of behavioral experiments, all animals are anaesthetized with 3 ml thiopental (1 g/40 ml i.p., Tyrol Pharma). The mice are perfused transcardially with 0.01 M PBS (pH 7.4) for 2 min, followed by 4% paraformaldehyde (Merck) in PBS for 15 min. The brains are removed and placed in 4% paraformaldehyde for 24 h at 4° C. For dehydration they are then transferred to a 20% sucrose (Merck) solution in 0.1 M PBS at 4° C. until they sink. The brains are frozen in methylbutan at -20° C. for 2 min and stored at -70° C. Using a sledge microtome (mod. 3800-Frigocut, Leica), 25  $\mu$ m sections are taken from the genu of the corpus callosum (AP 1.7 mm) to the hippocampus (AP 21.8 mm) and from AP 24.16 to AP 26.72. Forty-six sections are cut and stored in sorters in 0.25 M Tris buffer (pH 7.4) for immunohistochemistry.

A series of sections is processed for free-floating tyrosine hydroxylase (TH) immunohistochemistry. Following three rinses in 0.1 M PBS, endogenous peroxidase activity is quenched for 10 min in 0.3% H<sub>2</sub>O<sub>2</sub>+PBS. After rinsing in PBS, sections are preincubated in 10% normal bovine serum (Sigma) for 5 min as blocking agent and transferred to either primary anti-rat TH rabbit antiserum (dilution 1:2000).

Following overnight incubation at room temperature, sections for TH immunoreactivity are rinsed in PBS (2×10 min) and incubated in biotinylated anti-rabbit immunoglobulin G raised in goat (dilution 1:200) (Vector) for 90 min, rinsed repeatedly and transferred to Vectastain ABC (Vector) solution for 1 h. 3,3'-Diaminobenzidine tetrahydrochloride (DAB; Sigma) in 0.1 M PBS, supplemented with 0.005% H<sub>2</sub>O<sub>2</sub>, serves as chromogen in the subsequent visualization reaction. Sections are mounted on to gelatin-coated slides, left to dry overnight, counter-stained with hematoxylin dehydrated in ascending alcohol concentrations and cleared in butylacetate. Coverslips are mounted on entellan.

#### Rotarod Test

We use a modification of the procedure described by Rozas and Labandeira-Garcia (1997), with a CR-1 Rotamex system (Columbus Instruments, Columbus, Ohio) comprising an IBM-compatible personal computer, a CIO-24 data acquisition card, a control unit, and a four-lane rotarod unit. The rotarod unit consists of a rotating spindle (diameter 7.3 cm) and individual compartments for each mouse. The system software allows preprogramming of session protocols with varying rotational speeds (0–80 rpm). Infrared beams are used to detect when a mouse has fallen onto the base grid beneath the rotarod. The system logs the fall as the end of the experiment for that mouse, and the total time on the rotarod, as well as the time of the fall and all the set-up parameters, are recorded. The system also allows a weak current to be passed through the base grid, to aid training.

### 3. Dementia

#### The Object Recognition Task

The object recognition task has been designed to assess the effects of experimental manipulations on the cognitive performance of rodents. A rat is placed in an open field, in which two identical objects are present. The rats inspects both objects during the first trial of the object recognition task. In a second trial, after a retention interval of for example 24 hours, one of the two objects used in the first trial, the 'familiar' object, and a novel object are placed in the open field. The inspection time at each of the objects is registered. The basic measures in the OR task is the time spent by a rat exploring the two object the second trial. Good retention is reflected by higher exploration times towards the novel than the 'familiar' object.

Administration of the putative cognition enhancer prior to the first trial predominantly allows assessment of the effects on acquisition, and eventually on consolidation processes. Administration of the testing compound after the first trial allows to assess the effects on consolidation processes, whereas administration before the second trial allows to measure effects on retrieval processes.

#### The Passive Avoidance Task

The passive avoidance task assesses memory performance in rats and mice. The inhibitory avoidance apparatus consists of a two-compartment box with a light compartment and a dark compartment. The two compartments are separated by a guillotine door that can be operated by the experimenter. A threshold of 2 cm separates the two com-

partments when the guillotine door is raised. When the door is open, the illumination in the dark compartment is about 2 lux. The light intensity is about 500 lux at the center of the floor of the light compartment.

Two habituation sessions, one shock session, and a retention session are given, separated by inter-session intervals of 24 hours. In the habituation sessions and the retention session the rat is allowed to explore the apparatus for 300 sec. The rat is placed in the light compartment, facing the wall opposite to the guillotine door. After an accommodation period of 15 sec. the guillotine door is opened so that all parts of the apparatus can be visited freely. Rats normally avoid brightly lit areas and will enter the dark compartment within a few seconds.

In the shock session the guillotine door between the compartments is lowered as soon as the rat has entered the dark compartment with its four paws, and a scrambled 1 mA footshock is administered for 2 sec. The rat is removed from the apparatus and put back into its home cage. The procedure during the retention session is identical to that of the habituation sessions.

The step-through latency, that is the first latency of entering the dark compartment (in sec.) during the retention session is an index of the memory performance of the animal; the longer the latency to enter the dark compartment, the better the retention is. A testing compound is given half an hour before the shock session, together with 1 mg\*kg<sup>-1</sup> scopolamine. Scopolamine impairs the memory performance during the retention session 24 hours later. If the test compound increases the enter latency compared with the scopolamine-treated controls, is likely to possess cognition enhancing potential.

#### The Morris Water Escape Task

The Morris water escape task measures spatial orientation learning in rodents. It is a test system that has extensively been used to investigate the effects of putative therapeutic on the cognitive functions of rats and mice. The performance of an animal is assessed in a circular water tank with an escape platform that is submerged about 1 cm below the surface of the water. The escape platform is not visible for an animal swimming in the water tank. Abundant extra-maze cues are provided by the furniture in the room, including desks, computer equipment, a second water tank, the presence of the experimenter, and by a radio on a shelf that is playing softly.

The animals receive four trials during five daily acquisition sessions. A trial is started by placing an animal into the pool, facing the wall of the tank. Each of four starting positions in the quadrants north, east, south, and west is used once in a series of four trials; their order is randomized. The escape platform is always in the same position. A trial is terminated as soon as the animal had climbs onto the escape platform or when 90 seconds have elapsed, whichever event occurs first. The animal is allowed to stay on the platform for 30 seconds. Then it is taken from the platform and the next trial is started. If an animal did not find the platform within 90 seconds it is put on the platform by the experimenter and is allowed to stay there for 30 seconds. After the fourth trial of the fifth daily session, an additional trial is given as a probe trial: the platform is removed, and the time the animal spends in the four quadrants is measured for 30 or 60 seconds. In the probe trial, all animals start from the same start position, opposite to the quadrant where the escape platform had been positioned during acquisition.

Four different measures are taken to evaluate the performance of an animal during acquisition training: escape latency, traveled distance, distance to platform, and swim-

51

ming speed. The following measures are evaluated for the probe trial: time (s) in quadrants and traveled distance (cm) in the four quadrants. The probe trial provides additional information about how well an animal learned the position of the escape platform. If an animal spends more time and swims a longer distance in the quadrant where the platform had been positioned during the acquisition sessions than in any other quadrant, one concludes that the platform position has been learned well.

In order to assess the effects of putative cognition enhancing compounds, rats or mice with specific brain lesions which impair cognitive functions, or animals treated with compounds such as scopolamine or MK-801, which interfere with normal learning, or aged animals which suffer from cognitive deficits, are used.

The T-Maze Spontaneous Alternation Task

The T-maze spontaneous alternation task (TeMCAT) assesses the spatial memory performance in mice. The start arm and the two goal arms of the T-maze are provided with guillotine doors which can be operated manually by the experimenter. A mouse is put into the start arm at the beginning of training. The guillotine door is closed. In the first trial, the ‘forced trial’, either the left or right goal arm is blocked by lowering the guillotine door. After the mouse has been released from the start arm, it will negotiate the maze, eventually enter the open goal arm, and return to the start position, where it will be confined for 5 seconds, by lowering the guillotine door. Then, the animal can choose freely between the left and right goal arm (all guillotine-doors opened) during 14 ‘free choice’ trials. As soon a the mouse has entered one goal arm, the other one is closed. The mouse eventually returns to the start arm and is free to visit whichever go alarm it wants after having been confined to the start arm for 5 seconds. After completion of 14 free choice trials in one session, the animal is removed from the maze. During training, the animal is never handled.

The percent alternations out of 14 trials is calculated. This percentage and the total time needed to complete the first forced trial and the subsequent 14 free choice trials (in s) is analyzed. Cognitive deficits are usually induced by an injection of scopolamine, 30 min before the start of the training session. Scopolamine reduced the per-cent alternations to chance level, or below. A cognition enhancer, which is always administered before the training session, will at least partially, antagonize the scopolamine-induced reduction in the spontaneous alternation rate.

EXAMPLE 10

Expression of Human Protein Kinase-like Protein

Total RNA used for Taqman quantitative analysis were either purchased (Clontech,CA) or extracted from tissues using TRIzol reagent (Life Technologies, MD) according to a modified vendor protocol which utilizes the Rneasy protocol (Qiagen, CA).

One hundred µg of each RNA were treated with DNase I using RNase free- DNase (Qiagen, CA) for use with RNeasy or QiaAmp columns.

After elution and quantitation with Ribogreen (Molecular Probes Inc., OR), each sample was reverse transcribed using the GibcoBRL Superscript II First Strand Synthesis System for RT-PCR according to vendor protocol (Life Technologies, MD). The final concentration of RNA in the reaction mix was 50 ng/µL. Reverse transcription was performed with 50 ng of random hexamers.

52

Specific primers and probe were designed according to PE Applied Biosystems’ Primer Express program recommendations and are listed below:

forward primer: 5'-(CTGTGACCTGAAGTCGGACAAC)-3'

reverse primer: 5'-(GGAATGACTGCCTCGAAATCC)-3'

probe: SYBR Green

Quantitation experiments were performed on 25 ng of reverse transcribed RNA from each sample. 18S ribosomal RNA was measured as a control using the Pre-Developed TaqMan Assay Reagents (PDAR)(PE Applied Biosystems, CA). The assay reaction mix was as follows:

	final
TaqMan SYBR Green PCR Master Mix (2x)	1x
(PE Applied Biosystems, CA)	
Forward primer	300 nM
Reverse primer	300 nM
cDNA	25 ng
Water to	25 uL

PCR Conditions:

Once:	2' minutes at 50° C. 10 minutes at 95° C.
40cycles:	15 sec. at 95° C. 1 minute at 60° C.

The experiment was performed on an ABI Prism 7700 Sequence Detector (PE Applied Biosystems, CA). At the end of the run, fluorescence data acquired during PCR were processed as described in the ABI Prism 7700 user’s manual. Fold change was calculated using the delta-delta CT method with normalization to the 18S values. Relative expression was calculated by normalizing to 18s (D Ct), then making the highest expressing tissue 100 and everything else relative to it. Copy number conversion was performed without normalization using the formula Cn=10(Ct-40.007)/-3.623.

The results are shown in FIGS. 13, 14, 15 and 16.

Protein kinase-like protein expression in adipose and skeletal muscle could be regulated to increase insulin sensitivity.

References

1. Protein kinases 6. The eukaryotic protein kinase superfamily: kinase (catalytic) domain structure and classification. FASEB J 1995 May;9(8):576-96

2. Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. Methods Enzymol 1991;200:38-62

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 12

<210> SEQ ID NO 1  
<211> LENGTH: 858  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

atgaccgact tcccggccag gctcttcctg gagaacagca agctggagca cagcgaggac	60
gagggcagcg tcctgggcca gggcggcagt ggcaccgtca tctaccgggc ccggtaccag	120
ggccagcctg tggccgtcaa gcgcttcac atcaaaaaat tcaagaactt tgctaacgta	180
ccggcagaca ccatgctgag gcacctgcgg gccaccgatg ccatgaagaa cttctccgag	240
ttccggcagg aggccagcat gctgcacgcg ctgcagcacc cctgcatcgt ggcgctcatc	300
ggcatcagca tccacccgct ctgcttcgcc ctggagctcg cgccgctcag cagcctcaac	360
accgtgctgt ccgagaacgc cagagattct tcctttatac ccctgggaca catgctcacc	420
caaaaaatag cctaccagat cgccctcggg ctggcctacc tgcacaagaa aaacatcatc	480
ttctgtgacc tgaagtcgga caacattctg gtgtgggtccc ttgacgtcaa ggagcacatc	540
aacatcaagc tatctgacta cgggatttcg aggcagtcac tccatgaggg cgccctaggg	600
gtggagggca ctcttggtta ccaggcccca gagatcaggg ctgcatttgt atatgatgag	660
aaggtagata tgttctccta tggaatggtg ctctacgagt tgctgtcagg acagcgccct	720
gcactggggc accaccagct ccagattgcc aagaagctgt ccaagggcat ccgcccgggt	780
ctggggcagc cggaggaagt gcagttccgg cgactgcagg cgctcatgat ggagtgtctg	840
gacactaagc cagagaag	858

<210> SEQ ID NO 2  
<211> LENGTH: 286  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Thr Asp Phe Pro Ala Arg Leu Phe Leu Glu Asn Ser Lys Leu Glu	
1 5 10 15	
His Ser Glu Asp Glu Gly Ser Val Leu Gly Gln Gly Gly Ser Gly Thr	
20 25 30	
Val Ile Tyr Arg Ala Arg Tyr Gln Gly Gln Pro Val Ala Val Lys Arg	
35 40 45	
Phe His Ile Lys Lys Phe Lys Asn Phe Ala Asn Val Pro Ala Asp Thr	
50 55 60	
Met Leu Arg His Leu Arg Ala Thr Asp Ala Met Lys Asn Phe Ser Glu	
65 70 75 80	
Phe Arg Gln Glu Ala Ser Met Leu His Ala Leu Gln His Pro Cys Ile	
85 90 95	
Val Ala Leu Ile Gly Ile Ser Ile His Pro Leu Cys Phe Ala Leu Glu	
100 105 110	
Leu Ala Pro Leu Ser Ser Leu Asn Thr Val Leu Ser Glu Asn Ala Arg	
115 120 125	
Asp Ser Ser Phe Ile Pro Leu Gly His Met Leu Thr Gln Lys Ile Ala	
130 135 140	
Tyr Gln Ile Ala Ser Gly Leu Ala Tyr Leu His Lys Lys Asn Ile Ile	
145 150 155 160	

-continued

Phe	Cys	Asp	Leu	Lys	Ser	Asp	Asn	Ile	Leu	Val	Trp	Ser	Leu	Asp	Val
				165					170					175	
Lys	Glu	His	Ile	Asn	Ile	Lys	Leu	Ser	Asp	Tyr	Gly	Ile	Ser	Arg	Gln
			180					185					190		
Ser	Phe	His	Glu	Gly	Ala	Leu	Gly	Val	Glu	Gly	Thr	Pro	Gly	Tyr	Gln
		195					200					205			
Ala	Pro	Glu	Ile	Arg	Pro	Arg	Ile	Val	Tyr	Asp	Glu	Lys	Val	Asp	Met
	210					215					220				
Phe	Ser	Tyr	Gly	Met	Val	Leu	Tyr	Glu	Leu	Leu	Ser	Gly	Gln	Arg	Pro
225					230				235						240
Ala	Leu	Gly	His	His	Gln	Leu	Gln	Ile	Ala	Lys	Lys	Leu	Ser	Lys	Gly
			245						250					255	
Ile	Arg	Pro	Val	Leu	Gly	Gln	Pro	Glu	Glu	Val	Gln	Phe	Arg	Arg	Leu
			260					265					270		
Gln	Ala	Leu	Met	Met	Glu	Cys	Trp	Asp	Thr	Lys	Pro	Glu	Lys		
		275					280					285			
<210> SEQ ID NO 3															
<211> LENGTH: 394															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 3															
Ala	Glu	Leu	Thr	Leu	Glu	Glu	Ile	Ile	Gly	Ile	Gly	Gly	Phe	Gly	Lys
1				5					10					15	
Val	Tyr	Arg	Ala	Phe	Trp	Ile	Gly	Asp	Glu	Val	Ala	Val	Lys	Ala	Ala
			20					25					30		
Arg	His	Asp	Pro	Asp	Glu	Asp	Ile	Ser	Gln	Thr	Ile	Glu	Asn	Val	Arg
		35					40					45			
Gln	Glu	Ala	Lys	Leu	Phe	Ala	Met	Leu	Lys	His	Pro	Asn	Ile	Ile	Ala
	50					55					60				
Leu	Arg	Gly	Val	Cys	Leu	Lys	Glu	Pro	Asn	Leu	Cys	Leu	Val	Met	Glu
65				70					75						80
Phe	Ala	Arg	Gly	Gly	Pro	Leu	Asn	Arg	Val	Leu	Ser	Gly	Lys	Arg	Ile
			85					90						95	
Pro	Pro	Asp	Ile	Leu	Val	Asn	Trp	Ala	Val	Gln	Ile	Ala	Arg	Gly	Met
			100					105					110		
Asn	Tyr	Leu	His	Asp	Glu	Ala	Ile	Val	Pro	Ile	Ile	His	Arg	Asp	Leu
		115					120					125			
Lys	Ser	Ser	Asn	Ile	Leu	Ile	Leu	Gln	Lys	Val	Glu	Asn	Gly	Asp	Leu
	130					135					140				
Ser	Asn	Lys	Ile	Leu	Lys	Ile	Thr	Asp	Phe	Gly	Leu	Ala	Arg	Glu	Trp
145					150					155					160
His	Arg	Thr	Thr	Lys	Met	Ser	Ala	Ala	Gly	Thr	Tyr	Ala	Trp	Met	Ala
				165					170					175	
Pro	Glu	Val	Ile	Arg	Ala	Ser	Met	Phe	Ser	Lys	Gly	Ser	Asp	Val	Trp
			180					185					190		
Ser	Tyr	Gly	Val	Leu	Leu	Trp	Glu	Leu	Leu	Thr	Gly	Glu	Val	Pro	Phe
		195					200					205			
Arg	Gly	Ile	Asp	Gly	Leu	Arg	Val	Ala	Tyr	Gly	Val	Ala	Met	Asn	Lys
	210					215					220				
Leu	Ala	Leu	Pro	Ile	Pro	Ser	Thr	Cys	Pro	Glu	Pro	Phe	Ala	Lys	Leu
225					230					235					240
Met	Glu	Asp	Cys	Trp	Asn	Pro	Asp	Pro	His	Ser	Arg	Pro	Ser	Phe	Thr

	245	250	255
Asn Ile Leu Asp Gln Leu Thr Thr Ile Glu Glu Ser Gly Phe Phe Glu			
	260	265	270
Met Pro Lys Asp Ser Phe His Cys Leu Gln Asp Asn Trp Lys His Glu			
	275	280	285
Ile Gln Glu Met Phe Asp Gln Leu Arg Ala Lys Glu Lys Glu Leu Arg			
	290	295	300
Thr Trp Glu Glu Glu Leu Thr Arg Ala Ala Leu Gln Gln Lys Asn Gln			
305	310	315	320
Glu Glu Leu Leu Arg Arg Arg Glu Gln Glu Leu Ala Glu Arg Glu Ile			
	325	330	335
Asp Ile Leu Glu Arg Glu Leu Asn Ile Ile Ile His Gln Leu Cys Gln			
	340	345	350
Glu Lys Pro Arg Val Lys Lys Arg Lys Gly Lys Phe Arg Lys Ser Arg			
	355	360	365
Leu Ala Gln Pro Val Leu Pro Phe Pro His Gly His Ser Arg Cys Pro			
	370	375	380
Gly Gly Thr Gly Ser Ser Trp Gly Gly Gln			
385	390		
<210> SEQ ID NO 4			
<211> LENGTH: 473			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 4			
cacgaggctg gtcctgaac tgttcacgac cgacttcccg gccaggctct tcctggagaa	60		
cagcaagctg gagcacagcg aggacgaggg cagcgtcctg ggccagggcg gcagtggcac	120		
cgtcatctac cgggcccgtt accagggcca gcctgtggcc gtcaagcgct tccacatcaa	180		
aaaattcaag aactttgcta acgtaccggc agacaccatg ctgaggcacc tgcggggccac	240		
cgatgccatg aagaacttct ccgagttccg gcaggaggcc agcatgctgc acgcgctgca	300		
gcaccctgc atcgtggcgc tcatcggcat cagcatccac ccgctctgct tcgccctgga	360		
gctcgcgccg ctacgcagcc tcaacaccgt gctgtccgag aacgccagag attcttcctt	420		
tatacccttg ggacacatgc tcacccaaaa aatagcctac cagatcgctt cgg	473		
<210> SEQ ID NO 5			
<211> LENGTH: 454			
<212> TYPE: DNA			
<213> ORGANISM: Homo sapiens			
<400> SEQUENCE: 5			
cacgaggctg gtcctgaac tgttcacgac cgacttcccg gccaggctct tcctggagaa	60		
cagcaagctg gagcacagcg aggacgaggg cagcgtcctg ggccagggcg gcagtggcac	120		
cgtcatctac cgggcccgtt accagggcca gcctgtggcc gtcaagcgct tccacatcaa	180		
aaaattcaag aactttgcta acgtaccggc agacaccatg ctgaggcacc tgcggggccac	240		
cgatgccatg aagaacttct ccgagttccg gcaggaggcc agcatgctgc acgcgctgca	300		
gcaccctgc atcgtggcgc tcatcggcat cagcatccac ccgctctgct tcgccctgga	360		
gctcgcgccg ctacgcagcc tcaacaccgt gctgtccgag aacgccagag attcttcctt	420		
tatacccttg ggacacatgc tcacccaaaa aata	454		
<210> SEQ ID NO 6			

-continued

<211> LENGTH: 1066		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
<400> SEQUENCE: 6		
ctcttctctgt gagaacagca agctggagca cagcgaggac gagggcagcg tcctggggcca	60	
cggcggcagt ggcaccgtca tctaccgggc ccggtaccag ggccagcctg tggccgtcaa	120	
gcgcttccac atcaaaaaat tcaagaactt tgctaacgta ccggcagaca ccatgctgag	180	
gcacctgcgg gccaccgatg ccatgaagaa cttctccgag ttccggcagg aggccagcat	240	
gctgcacgcg ctgcagcacc cctgcatcgt ggcgctcacc ggcatcagca tccacccgac	300	
tctgcttcgc cctggagctc gcgccgctca gcagcctcaa caccgtgctg tccgagaacg	360	
ccagagattc ttcctttata cccctgggac acatgctcac ccaaaaaata gcctaccaga	420	
tcgcctcggg cctggcctac ctgcacaaga aaaacatcat cttctgtgac ctgaagtcgg	480	
acaacattct ggtgtggtcc cttgacgtca aggagcacat caacatcaag ctatctgact	540	
acgggatttc gaggcagtca ttccatgagg gcgccctagg cgtggagggc actcctggct	600	
accaggcccc agagatcagg cctcgcattg tatatgatga gaaggtagca tatgtctcct	660	
atggaatggt gctctacgag ttgctgtcag gacaggggcc tgcattgggc caccaacagc	720	
ttccgattgc caagaggtgt ccaaggcatc cgcccggtag tgaggacagc cggaaggaac	780	
tgcaatccgg cgatgcatgg ctcatatttg agatgctggg aaactataat caagcgaacg	840	
gagcactgtg gccctatacg taggtagccc caattagacg accctcacat tatgaaccaa	900	
tcggtaataa acgtgtacgt gcgggactag cagccaccta taacacacag ccacggccac	960	
ctgggctgtg atgcatacaa gacgctcaaa cacgcgggtc acagaaagcc tcatcgagtc	1020	
gcatccaccc gagagagtag cgccgccgaa ctagtataga cggtat	1066	
<210> SEQ ID NO 7		
<211> LENGTH: 471		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
<220> FEATURE:		
<221> NAME/KEY: misc_feature		
<222> LOCATION: (452)..(452)		
<223> OTHER INFORMATION: n=a, c, g or t		
<400> SEQUENCE: 7		
cacgaggaaa aaattcaaga actttgctaa cgtaccggca gacaccatgc tgaggcacct	60	
gcggggccacc gatgccatga agaacttctc cgagttccgg caggaggcca gcatgctgca	120	
cgcgctgcag caccctgca tcgtggcgct catcggcacc agcatccacc cgctctgctt	180	
cgccctggag ctgcgcccgc tcagcagcct caacaccgtg ctgtccgaga acgccagaga	240	
ttcttctctt atacccttg gacacatgct cacccaaaaa atagcctacc agatcgctc	300	
gggcctggcc tacctgcaca agaaaaacat catcttctgt gacctgaagt cggacaacat	360	
tctggtgtgg tcccttgacg tcaaggagca catcaacatc aagctatctg actacgggat	420	
ttcgaggcag tcattccatg agggcgccct angcgtggag ggcactcctg g	471	
<210> SEQ ID NO 8		
<211> LENGTH: 5475		
<212> TYPE: DNA		
<213> ORGANISM: Homo sapiens		
<400> SEQUENCE: 8		
ccccttaatg attttgaca ggccccaaag caatgctgtc ttacctgcgt gctcagctgc	60	

-continued

ggaaagcgga	aaagtgcaag	ctgatgaaga	tgatcatcgt	gggtcccccg	cgccagggca	120
agtccaccct	cctggagatc	ttacagacgg	ggagggcccc	ccaggtggtg	catggagagg	180
ccaccatcag	gaccaccaag	tgggagctcc	agaggccggc	tggctcaaga	gccaaaggtt	240
agtccgtgga	gttcaacgtc	tgggacatcg	ggggaccggc	cagcatggcc	actgtcaacc	300
agtgccttct	cacggacaag	gccctgtacg	tgggtggtctg	gaacctggcg	ctgggggagg	360
aggccgtggc	caacctccag	ttctggctgc	tcaacatcga	ggccaaggcc	ccaaacgccg	420
tggtgctggt	ggtcgggacg	cacctggatt	taattgaagc	caagttccgt	gtggaaagga	480
ttgcaacgct	gcgtgcctat	gtgctggcac	tctgccgctc	cccctccggc	tccagggcca	540
caggcttccc	agacatcacc	ttcaaacact	tacatgagat	ttcctgcaag	agcctggaag	600
gtcaggaagg	gctgcgacag	ctgatttttc	acgtcacgtg	cagcatgaag	gacgtgggca	660
gcaccatcgg	ctgccagcga	ctggcagggc	ggctgatccc	caggagctac	ctgagcctgc	720
aggaggccgt	gctggcagag	cagcagcgcc	gcagccggga	cgacgacgtg	cagtacctga	780
cggacaggca	gctggagcag	ctggtggagc	agacgcccga	caacgacatc	aaggactacg	840
aggacctgca	gtcagccatc	agcttcctca	tagaaaccgg	cacctgctc	catttcccgg	900
acaccagcca	cggcctgagg	aacctctact	tcctcgaccc	tatttggtct	tccgaatgtc	960
tgcaagaggat	ctttaatat	aagggtctct	ggtcagtggc	caagaatggg	gtgatcagag	1020
cagaagacct	caggatgctg	ctggtgggga	ctggcttcac	gcagcagacg	gaagagcagt	1080
acttccagtt	cctggccaag	tttgagatcg	ccctgcccgt	cgccaatgac	agctacctcc	1140
tgccccatct	ccttccatct	aaacctggcc	tggacacca	cggtatgcgg	cacccacag	1200
ccaacaccat	tcagagggtg	tttaagatga	gcttcgttcc	cgttggcttc	tggcaaaggt	1260
ttatagcacg	gatgctgac	agcctggcgg	agatggacct	gcagcttttt	gaaaacaaga	1320
agaatactaa	aagcaggaac	aggaaagtca	ccatttacag	ttttacagga	aaccagagaa	1380
atcgctgtag	cacattcaga	gtgaaaagaa	atcagacct	ctattggcag	gaagggtcc	1440
tggtcacttt	tgatgggggc	tacctcagt	tggaatcttc	cgacgtgaac	tggaaaaaga	1500
agaaaagcgg	aggaatgaaa	attgtttgcc	aatcagaagt	gagggacttc	tcagccatgg	1560
ctttcatcac	ggaccacgtc	aattccttga	ttgatcagt	gtttcccggc	ctgacggcca	1620
cagagagcga	cgggacgcca	ctcatggagc	agtacgtgcc	ctgcccggtc	tgcgagacag	1680
cctgggcccc	gcacacggac	cccagtgaga	aatcagagga	tgtgcagtac	ttcgacatgg	1740
aagactgtgt	cctgacggcc	atcgagcggg	acttcatctc	ctgccccaga	cacccggacc	1800
tccccgtgcc	gctgcaggag	ctggtccctg	aactgttcat	gaccgacttc	ccggccaggc	1860
tcttcctgga	gaacagcaag	ctggagcaca	gcgaggacga	gggcagcgtc	ctggggccagg	1920
gcggcagtg	caccgtcatc	taccggggcc	ggtaccagg	ccagcctgtg	gccgtcaagc	1980
gcttccacat	caaaaaattc	aagaactttg	ctaactgacc	ggcagacacc	atgctgaggc	2040
acctgcgggc	caccgatgcc	atgaagaact	tctccgagtt	ccggcaggag	gccagcatgc	2100
tgacgcgct	gcagaccccc	tgatcgtgg	cgctcatcgg	catcagcatc	caccgctct	2160
gcttcgccct	ggagctcgcg	ccgctcagca	gcctcaacac	cgtgctgtcc	gagaacgcca	2220
gagattcttc	ctttataccc	ctgggacaca	tgctcaccca	aaaaatagcc	taccagatcg	2280
cctcgggcct	ggcctacctg	cacaagaaaa	acatcatctt	ctgtgacctg	aagtcggaca	2340
acattctggt	gtggtccctt	gacgtcaagg	agcacatcaa	catcaagcta	tctgactacg	2400

-continued

ggatttgcgag	gcagtcattc	catgagggcg	ccctagggcg	ggagggcact	cctggctacc	2460
aggccccaga	gatcaggcct	cgcattgtat	atgatgagaa	ggtagatatg	ttctcctatg	2520
gaatgggtgct	ctacgagttg	ctgtcaggac	agcgccctgc	actggggccac	caccagctcc	2580
agattgccaa	gaagctgtcc	aagggcatcc	gcccggttct	ggggcagccg	gaggaagtgc	2640
agttccggcg	actgcaggcg	ctcatgatgg	agtgctggga	cactaagcca	gagaagcgac	2700
cgttgccct	gtcggtggtg	agccagatga	aggacccgac	ttttgccacc	ttcatgtatg	2760
aactgtgctg	tgggaagcag	acagccttct	tctcatccca	gggccaggag	tacaccgtgg	2820
tgttttggga	tggaaaagag	gagtccagga	actacacggg	ggtgaacaca	gagaagggcc	2880
tcatggaggt	gcagaggatg	tgctgccctg	ggatgaaggt	gagctgccag	ctccaggtcc	2940
agagatccct	gtggacagcc	accgaggacc	agaaaatcta	catctacacc	ctcaagggca	3000
tgtgccccct	aaacacaccc	caacaggcct	tggatactcc	agctgtcgtc	acctgcttct	3060
tggccgtgcc	tgttattaaa	aagaattcct	acctgggtct	agcgggcctc	gccgatgggc	3120
ttgtggctgt	gtttcccggtg	gtgcggggca	ccccaaagga	cagctgctcc	tacctgtgct	3180
cacacacagc	caacaggtcc	aagttcagca	tcgcggatga	agacgcacgg	cagaaccctt	3240
accagtgaa	ggccatggag	gtggtcaaca	gcggctctga	ggtctggtac	agcaatgggc	3300
cgggcctcct	tgtcatcgac	tgtgcctccc	tggagatctg	caggcggctg	gagccctaca	3360
tggccccctc	catggttacg	tcagtcgtgt	gcagctctga	gggcagaggg	gaggaggtcg	3420
tctggtgcct	ggatgacaag	gccaaactcct	tgggtgatgta	ccactccacc	acctaccagc	3480
tgtgtgcccg	gtacttctgc	ggggtcccca	gccccctcag	ggacatgttt	cccgtgcggc	3540
ccttggaac	ggaacccccg	gcagccagcc	acacggccaa	cccaaaggtg	cctgaggggg	3600
actccatcgc	ggacgtgagc	atcatgtaca	gtgaggagct	gggcacgcag	atcctgatcc	3660
accaggaatc	actcactgac	tactgctcca	tgtcctccta	ctcctcatcc	ccaccccgcc	3720
aggctgccag	gtccccctca	agcctcccca	gctccccagc	aagttcttcc	agtgtgcctt	3780
tctccaccga	ctgcgaggac	tcagacatgc	tacatacgcc	cgggtgctgcc	tccgacaggt	3840
ctgagcatga	cctgaccccc	atggacgggg	agaccttcag	ccagcacctg	caggccgtga	3900
agatcctcgc	cgtcagagac	ctcatttggg	tccccaggcg	cggtgagat	gttatcgtca	3960
ttggcctgga	gaaggattct	gaagcccagc	ggggccgagt	cattgccgtc	ttaaaagccc	4020
gagagctgac	tccgcatggg	gtgctgggtg	atgctgccgt	ggtggcaaag	gacactgttg	4080
tgtgcacctt	tgaaaatgaa	aacacagagt	ggtgcctggc	cgtctggagg	ggctggggcg	4140
ccaggggagt	cgacattttc	taccagtcct	acgaggagct	gggccggctg	gaggcttgca	4200
ctcgcaagag	aaggtaatc	ctgtggaatg	actgtcacac	atcagagctg	gctggcccgg	4260
ggctgcagcc	tgaccctct	gccatcggcc	tctagtctc	caaggacctc	gaagacagat	4320
ggagttctcc	cctgaactcc	ttgctgctaa	gaagtgtgta	gaagttactc	gcctggcggt	4380
ggctccaggg	ttctctggtt	ctctggagca	gagttctctg	aatacccat	cccccaactg	4440
ctgattttac	agccccaggg	aagacagtgg	tatcaggctg	ggagcggcct	cctctggcct	4500
cccccatcag	tttgaggag	caggggtgca	ggatcctgtt	ctgagctggg	tcaaacaag	4560
cagggccggg	ccttctctgc	atccccagg	ctcagatgga	attacactag	aggccctccg	4620
ctgggaagca	cttgaggtag	ggcaggaggg	gggctgtgac	ccctgccctt	tccccgccag	4680
agacctcggg	ctctcagcac	attccacagg	ctcctgagtc	cccaggcct	gggccagctt	4740
gggcaagcca	agatcagatg	tctctgtgtt	cgggaaggtc	tccgtgtggg	aaagcccttg	4800

-continued

ggggatcccg ggtgaggagt gttgccccat ccagagaatg aatgagttcc ttttaagtgcc 4860  
accgccagca agcccagagg cacacagtcc gagtgcaccc gcttagcctt tacattcctc 4920  
tccaccgaca aaaggaaggg gaaactcaat cagcaggact tcagaaaggg ccttgtgttt 4980  
atagctttgt caagtaaatt tggacgcagc tggagcacag gccctgtttg tttgcacata 5040  
ataatcttgt ttatcacttt aaaaaattca gtaatatctc agcagtcagg cttctggttg 5100  
tgaaatcaca ttgtatggga tttataccaa attatgtatt tgctaaacat tcaactgcaca 5160  
cgtgtacagc ggagtacgaa aaggaacgtt gtccacaggg gatttatgga tacaacagca 5220  
aacattttat aaactatgca catgcattac acacatgcac acacatatgc acacacatgt 5280  
gcaaacatag ccactttttt gtcaagagtt accctttggg gctccttaaa ccagaatggg 5340  
agtttgaaag agagatcata ctccagctga agtttggtga cccttttcta aaattaaaaa 5400  
gatcaaattt agtatttgct ggatatgcag ggagatgaga ctcttttaat ctcaaaataa 5460  
acagattcctt tcaag 5475

<210> SEQ ID NO 9  
<211> LENGTH: 1394  
<212> TYPE: PRT  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 9

Met Leu Ser Tyr Leu Arg Ala Gln Leu Arg Lys Ala Glu Lys Cys Lys  
1 5 10 15  
Leu Met Lys Met Ile Ile Val Gly Pro Pro Arg Gln Gly Lys Ser Thr  
20 25 30  
Leu Leu Glu Ile Leu Gln Thr Gly Arg Ala Pro Gln Val Val His Gly  
35 40 45  
Glu Ala Thr Ile Arg Thr Thr Lys Trp Glu Leu Gln Arg Pro Ala Gly  
50 55 60  
Ser Arg Ala Lys Val Glu Ser Val Glu Phe Asn Val Trp Asp Ile Gly  
65 70 75 80  
Gly Pro Ala Ser Met Ala Thr Val Asn Gln Cys Phe Phe Thr Asp Lys  
85 90 95  
Ala Leu Tyr Val Val Val Trp Asn Leu Ala Leu Gly Glu Glu Ala Val  
100 105 110  
Ala Asn Leu Gln Phe Trp Leu Leu Asn Ile Glu Ala Lys Ala Pro Asn  
115 120 125  
Ala Val Val Leu Val Val Gly Thr His Leu Asp Leu Ile Glu Ala Lys  
130 135 140  
Phe Arg Val Glu Arg Ile Ala Thr Leu Arg Ala Tyr Val Leu Ala Leu  
145 150 155 160  
Cys Arg Ser Pro Ser Gly Ser Arg Ala Thr Gly Phe Pro Asp Ile Thr  
165 170 175  
Phe Lys His Leu His Glu Ile Ser Cys Lys Ser Leu Glu Gly Gln Glu  
180 185 190  
Gly Leu Arg Gln Leu Ile Phe His Val Thr Cys Ser Met Lys Asp Val  
195 200 205  
Gly Ser Thr Ile Gly Cys Gln Arg Leu Ala Gly Arg Leu Ile Pro Arg  
210 215 220  
Ser Tyr Leu Ser Leu Gln Glu Ala Val Leu Ala Glu Gln Gln Arg Arg  
225 230 235 240  
Ser Arg Asp Asp Asp Val Gln Tyr Leu Thr Asp Arg Gln Leu Glu Gln

-continued

245					250					255					
Leu	Val	Glu	Gln	Thr	Pro	Asp	Asn	Asp	Ile	Lys	Asp	Tyr	Glu	Asp	Leu
		260						265					270		
Gln	Ser	Ala	Ile	Ser	Phe	Leu	Ile	Glu	Thr	Gly	Thr	Leu	Leu	His	Phe
		275					280					285			
Pro	Asp	Thr	Ser	His	Gly	Leu	Arg	Asn	Leu	Tyr	Phe	Leu	Asp	Pro	Ile
	290					295					300				
Trp	Leu	Ser	Glu	Cys	Leu	Gln	Arg	Ile	Phe	Asn	Ile	Lys	Gly	Ser	Arg
305					310					315					320
Ser	Val	Ala	Lys	Asn	Gly	Val	Ile	Arg	Ala	Glu	Asp	Leu	Arg	Met	Leu
				325					330					335	
Leu	Val	Gly	Thr	Gly	Phe	Thr	Gln	Gln	Thr	Glu	Glu	Gln	Tyr	Phe	Gln
		340						345					350		
Phe	Leu	Ala	Lys	Phe	Glu	Ile	Ala	Leu	Pro	Val	Ala	Asn	Asp	Ser	Tyr
		355					360					365			
Leu	Leu	Pro	His	Leu	Leu	Pro	Ser	Lys	Pro	Gly	Leu	Asp	Thr	His	Gly
	370					375					380				
Met	Arg	His	Pro	Thr	Ala	Asn	Thr	Ile	Gln	Arg	Val	Phe	Lys	Met	Ser
385					390					395					400
Phe	Val	Pro	Val	Gly	Phe	Trp	Gln	Arg	Phe	Ile	Ala	Arg	Met	Leu	Ile
				405					410					415	
Ser	Leu	Ala	Glu	Met	Asp	Leu	Gln	Leu	Phe	Glu	Asn	Lys	Lys	Asn	Thr
			420					425					430		
Lys	Ser	Arg	Asn	Arg	Lys	Val	Thr	Ile	Tyr	Ser	Phe	Thr	Gly	Asn	Gln
		435					440					445			
Arg	Asn	Arg	Cys	Ser	Thr	Phe	Arg	Val	Lys	Arg	Asn	Gln	Thr	Ile	Tyr
	450					455					460				
Trp	Gln	Glu	Gly	Leu	Leu	Val	Thr	Phe	Asp	Gly	Gly	Tyr	Leu	Ser	Val
465					470					475					480
Glu	Ser	Ser	Asp	Val	Asn	Trp	Lys	Lys	Lys	Lys	Ser	Gly	Gly	Met	Lys
				485					490					495	
Ile	Val	Cys	Gln	Ser	Glu	Val	Arg	Asp	Phe	Ser	Ala	Met	Ala	Phe	Ile
		500						505					510		
Thr	Asp	His	Val	Asn	Ser	Leu	Ile	Asp	Gln	Trp	Phe	Pro	Ala	Leu	Thr
		515					520					525			
Ala	Thr	Glu	Ser	Asp	Gly	Thr	Pro	Leu	Met	Glu	Gln	Tyr	Val	Pro	Cys
	530					535					540				
Pro	Val	Cys	Glu	Thr	Ala	Trp	Ala	Gln	His	Thr	Asp	Pro	Ser	Glu	Lys
545					550					555					560
Ser	Glu	Asp	Val	Gln	Tyr	Phe	Asp	Met	Glu	Asp	Cys	Val	Leu	Thr	Ala
				565					570					575	
Ile	Glu	Arg	Asp	Phe	Ile	Ser	Cys	Pro	Arg	His	Pro	Asp	Leu	Pro	Val
			580					585					590		
Pro	Leu	Gln	Glu	Leu	Val	Pro	Glu	Leu	Phe	Met	Thr	Asp	Phe	Pro	Ala
		595					600					605			
Arg	Leu	Phe	Leu	Glu	Asn	Ser	Lys	Leu	Glu	His	Ser	Glu	Asp	Glu	Gly
	610					615					620				
Ser	Val	Leu	Gly	Gln	Gly	Gly	Ser	Gly	Thr	Val	Ile	Tyr	Arg	Ala	Arg
625					630					635					640
Tyr	Gln	Gly	Gln	Pro	Val	Ala	Val	Lys	Arg	Phe	His	Ile	Lys	Lys	Phe
				645					650					655	
Lys	Asn	Phe	Ala	Asn	Val	Pro	Ala	Asp	Thr	Met	Leu	Arg	His	Leu	Arg
			660					665						670	

-continued

Ala	Thr	Asp	Ala	Met	Lys	Asn	Phe	Ser	Glu	Phe	Arg	Gln	Glu	Ala	Ser	
		675					680					685				
Met	Leu	His	Ala	Leu	Gln	His	Pro	Cys	Ile	Val	Ala	Leu	Ile	Gly	Ile	
	690					695					700					
Ser	Ile	His	Pro	Leu	Cys	Phe	Ala	Leu	Glu	Leu	Ala	Pro	Leu	Ser	Ser	
705					710					715					720	
Leu	Asn	Thr	Val	Leu	Ser	Glu	Asn	Ala	Arg	Asp	Ser	Ser	Phe	Ile	Pro	
				725					730					735		
Leu	Gly	His	Met	Leu	Thr	Gln	Lys	Ile	Ala	Tyr	Gln	Ile	Ala	Ser	Gly	
			740					745					750			
Leu	Ala	Tyr	Leu	His	Lys	Lys	Asn	Ile	Ile	Phe	Cys	Asp	Leu	Lys	Ser	
		755					760					765				
Asp	Asn	Ile	Leu	Val	Trp	Ser	Leu	Asp	Val	Lys	Glu	His	Ile	Asn	Ile	
	770					775					780					
Lys	Leu	Ser	Asp	Tyr	Gly	Ile	Ser	Arg	Gln	Ser	Phe	His	Glu	Gly	Ala	
785					790					795					800	
Leu	Gly	Val	Glu	Gly	Thr	Pro	Gly	Tyr	Gln	Ala	Pro	Glu	Ile	Arg	Pro	
				805					810					815		
Arg	Ile	Val	Tyr	Asp	Glu	Lys	Val	Asp	Met	Phe	Ser	Tyr	Gly	Met	Val	
			820					825					830			
Leu	Tyr	Glu	Leu	Leu	Ser	Gly	Gln	Arg	Pro	Ala	Leu	Gly	His	His	Gln	
		835					840					845				
Leu	Gln	Ile	Ala	Lys	Lys	Leu	Ser	Lys	Gly	Ile	Arg	Pro	Val	Leu	Gly	
	850					855					860					
Gln	Pro	Glu	Glu	Val	Gln	Phe	Arg	Arg	Leu	Gln	Ala	Leu	Met	Met	Glu	
865					870					875					880	
Cys	Trp	Asp	Thr	Lys	Pro	Glu	Lys	Arg	Pro	Leu	Ala	Leu	Ser	Val	Val	
				885					890					895		
Ser	Gln	Met	Lys	Asp	Pro	Thr	Phe	Ala	Thr	Phe	Met	Tyr	Glu	Leu	Cys	
			900					905					910			
Cys	Gly	Lys	Gln	Thr	Ala	Phe	Phe	Ser	Ser	Gln	Gly	Gln	Glu	Tyr	Thr	
		915					920					925				
Val	Val	Phe	Trp	Asp	Gly	Lys	Glu	Glu	Ser	Arg	Asn	Tyr	Thr	Val	Val	
						935					940					
Asn	Thr	Glu	Lys	Gly	Leu	Met	Glu	Val	Gln	Arg	Met	Cys	Cys	Pro	Gly	
945					950					955					960	
Met	Lys	Val	Ser	Cys	Gln	Leu	Gln	Val	Gln	Arg	Ser	Leu	Trp	Thr	Ala	
				965					970					975		
Thr	Glu	Asp	Gln	Lys	Ile	Tyr	Ile	Tyr	Thr	Leu	Lys	Gly	Met	Cys	Pro	
			980					985					990			
Leu	Asn	Thr	Pro	Gln	Gln	Ala	Leu	Asp	Thr	Pro	Ala	Val	Val	Thr	Cys	
			995				1000					1005				
Phe	Leu	Ala	Val	Pro	Val	Ile	Lys	Lys	Asn	Ser	Tyr	Leu	Val	Leu		
	1010					1015					1020					
Ala	Gly	Leu	Ala	Asp	Gly	Leu	Val	Ala	Val	Phe	Pro	Val	Val	Arg		
	1025					1030					1035					
Gly	Thr	Pro	Lys	Asp	Ser	Cys	Ser	Tyr	Leu	Cys	Ser	His	Thr	Ala		
	1040					1045					1050					
Asn	Arg	Ser	Lys	Phe	Ser	Ile	Ala	Asp	Glu	Asp	Ala	Arg	Gln	Asn		
	1055					1060					1065					
Pro	Tyr	Pro	Val	Lys	Ala	Met	Glu	Val	Val	Asn	Ser	Gly	Ser	Glu		
	1070					1075					1080					

-continued

Val	Trp	Tyr	Ser	Asn	Gly	Pro	Gly	Leu	Leu	Val	Ile	Asp	Cys	Ala
1085						1090					1095			
Ser	Leu	Glu	Ile	Cys	Arg	Arg	Leu	Glu	Pro	Tyr	Met	Ala	Pro	Ser
1100						1105					1110			
Met	Val	Thr	Ser	Val	Val	Cys	Ser	Ser	Glu	Gly	Arg	Gly	Glu	Glu
1115						1120					1125			
Val	Val	Trp	Cys	Leu	Asp	Asp	Lys	Ala	Asn	Ser	Leu	Val	Met	Tyr
1130						1135					1140			
His	Ser	Thr	Thr	Tyr	Gln	Leu	Cys	Ala	Arg	Tyr	Phe	Cys	Gly	Val
1145						1150					1155			
Pro	Ser	Pro	Leu	Arg	Asp	Met	Phe	Pro	Val	Arg	Pro	Leu	Asp	Thr
1160						1165					1170			
Glu	Pro	Pro	Ala	Ala	Ser	His	Thr	Ala	Asn	Pro	Lys	Val	Pro	Glu
1175						1180					1185			
Gly	Asp	Ser	Ile	Ala	Asp	Val	Ser	Ile	Met	Tyr	Ser	Glu	Glu	Leu
1190						1195					1200			
Gly	Thr	Gln	Ile	Leu	Ile	His	Gln	Glu	Ser	Leu	Thr	Asp	Tyr	Cys
1205						1210					1215			
Ser	Met	Ser	Ser	Tyr	Ser	Ser	Ser	Pro	Pro	Arg	Gln	Ala	Ala	Arg
1220						1225					1230			
Ser	Pro	Ser	Ser	Leu	Pro	Ser	Ser	Pro	Ala	Ser	Ser	Ser	Ser	Val
1235						1240					1245			
Pro	Phe	Ser	Thr	Asp	Cys	Glu	Asp	Ser	Asp	Met	Leu	His	Thr	Pro
1250						1255					1260			
Gly	Ala	Ala	Ser	Asp	Arg	Ser	Glu	His	Asp	Leu	Thr	Pro	Met	Asp
1265						1270					1275			
Gly	Glu	Thr	Phe	Ser	Gln	His	Leu	Gln	Ala	Val	Lys	Ile	Leu	Ala
1280						1285					1290			
Val	Arg	Asp	Leu	Ile	Trp	Val	Pro	Arg	Arg	Gly	Gly	Asp	Val	Ile
1295						1300					1305			
Val	Ile	Gly	Leu	Glu	Lys	Asp	Ser	Glu	Ala	Gln	Arg	Gly	Arg	Val
1310						1315					1320			
Ile	Ala	Val	Leu	Lys	Ala	Arg	Glu	Leu	Thr	Pro	His	Gly	Val	Leu
1325						1330					1335			
Val	Asp	Ala	Ala	Val	Val	Ala	Lys	Asp	Thr	Val	Val	Cys	Thr	Phe
1340						1345					1350			
Glu	Asn	Glu	Asn	Thr	Glu	Trp	Cys	Leu	Ala	Val	Trp	Arg	Gly	Trp
1355						1360					1365			
Gly	Ala	Arg	Glu	Phe	Asp	Ile	Phe	Tyr	Gln	Ser	Tyr	Glu	Glu	Leu
1370						1375					1380			
Gly	Arg	Leu	Glu	Ala	Cys	Thr	Arg	Lys	Arg	Arg				
1385						1390								

<210> SEQ ID NO 10  
<211> LENGTH: 4216  
<212> TYPE: DNA  
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 10

caaagcaatg ctgtcttacc tgcgtgctca gctgcggaaa gcggaaaagt gcaagctgat	60
gaagatgatc atcgtgggtc ccccgcgcca gggcaagtcc accctcctgg agatcttaca	120
gacggggagg gccccccagg tgggtgatgg agaggccacc atcaggacca ccaagtggga	180
gctccagagg ccggctggct caagagccaa ggttgagtcc gtggagttca acgtctggga	240

-continued

catcggggga	cgggccagca	tggccactgt	caaccagtgc	ttcttcacgg	acaaggccct	300
gtacgtggtg	gtctggaacc	tggcgctggg	ggaggaggcc	gtggccaacc	tccagttctg	360
gctgctcaac	atcgaggcca	aggcccaaaa	cgccgtggtg	ctggtggtcg	ggacgcacct	420
ggatttaatt	gaagccaagt	tccgtgtgga	aaggattgca	acgctgcgtg	cctatgtgct	480
ggcactctgc	cgctccccct	cgggctccag	ggccacaggc	ttcccagaca	tcaccttcaa	540
acacttacat	gagatttcct	gcaagagcct	ggaaggtcag	gaagggctgc	gacagctgat	600
tttccacgtc	acgtgcagca	tgaaggacgt	gggcagcacc	atcggctgcc	agcgactggc	660
agggcggtg	atccccagga	gctacctgag	cctgcaggag	gccgtgctgg	cagagcagca	720
gcgccgcagc	cgggacgacg	acgtgcagta	cctgacggac	aggcagctgg	agcagctggt	780
ggagcagacg	cccgacaacg	acatcaagga	ctacgaggac	ctgcagtcag	ccatcagctt	840
cctcatagaa	accggcaccc	tgctccattt	cccggacacc	agccacggcc	tgaggaacct	900
ctacttctc	gacctattt	ggctctccga	atgtctgcag	aggatcttta	atattaaggg	960
ctctcggtca	gtggccaaga	atggggtgat	cagagcagaa	gacctcagga	tgctgctggt	1020
ggggactggc	ttcacgcagc	agacggaaga	gcagtacttc	cagttcctgg	ccaagtttga	1080
gatcgccctg	cccgtcgcca	atgacagcta	cctcctgccc	catctccttc	catctaaacc	1140
tggcctggac	accacggta	tgcggcaccc	cacagccaac	accattcaga	gggtatttaa	1200
gatgagcttc	gttcccgttg	gcttctggca	aaggtttata	gcacggatgc	tgatcagcct	1260
ggcggagatg	gacctgcagc	tttttgaaaa	caagaagaat	actaaaagca	ggaacaggaa	1320
agtcaccatt	tacagtttta	caggaaacca	gagaaaatgc	tgtagcacat	tcagagtga	1380
aagaaatcag	accatctatt	ggcaggaagg	gctcctggtc	acttttgatg	ggggctacct	1440
cagtgtggaa	tcttcgcagc	tgaactggaa	aaagaagaaa	agcggaggaa	tgaaaattgt	1500
ttgccaatca	gaagtgaggg	acttctcagc	catggctttc	atcacggacc	acgtcaattc	1560
cttgattgat	cagtggtttc	cgccttgac	agccacagag	agcgacggga	cgccactcat	1620
ggagcagtac	gtgccctgcc	cggctctgca	gacagcctgg	gccagcaca	cggaccccag	1680
tgagaaatca	gaggatgtgc	agtacttcga	catggaagac	tgtgtcctga	cggccatcga	1740
gcgggacttc	atctcctgcc	ccagacaccc	ggacctcccc	gtgccgtgc	aggagctggt	1800
ccctgaactg	ttcatgaccg	acttcccggc	caggctcttc	ctggagaaca	gcaagctgga	1860
gcacagcgag	gacgagggca	gcgtcctggg	ccagggcggc	agtggcaccg	tcatctaccg	1920
ggcccggtag	cagggccagc	ctgtggccgt	caagcgcttc	cacatcaaaa	aattcaagaa	1980
ctttgctaac	gtaccggcag	acaccatgct	gaggcacctg	cgggccaccg	atgccatgaa	2040
gaacttctcc	gagttccggc	aggaggccag	catgctgcac	gcgctgcagc	accttgcat	2100
cgtggcgctc	atcggcatca	gcatccaccc	gctctgcttc	gccctggagc	tcgcgccgct	2160
cagcagcctc	aacaccgtgc	tgtccgagaa	cgccagagat	tcttccttta	tacctctggg	2220
acacatgctc	acccaaaaaa	tagcctacca	gatcgccctg	ggcctggcct	acctgcacaa	2280
gaaaaacatc	atcttctgtg	acctgaagtc	ggacaacatt	ctggtgtggt	cccttgacgt	2340
caaggagcac	atcaacatca	agctatctga	ctacgggatt	tcgaggcagt	cattccatga	2400
gggcgcctta	ggcgtggagg	gcactcctgg	ctaccaggcc	ccagagatca	ggcctcgcat	2460
tgtatatgat	gagaaggtag	atatgttctc	ctatggaatg	gtgctctacg	agttgctgtc	2520
aggacagcgc	cctgcactgg	gccaccacca	gctccagatt	gccaagaagc	tgtccaaggg	2580
catccgcccg	gttctggggc	agccggagga	agtgcagttc	cggcgactgc	aggcgctcat	2640

-continued

gatggagtgc tgggacacta agccagagaa gcgaccgctg gccctgtcgg tggtgagcca	2700
gatgaaggac ccgacttttg ccaccttcat gtatgaactg tgctgtggga agcagacagc	2760
cttcttctca tcccagggcc aggagtacac cgtgggtgtt tgggatggaa aagaggagtc	2820
caggaactac acggtggtga acacagagaa gggcctcatg gaggtgcaga ggatgtgctg	2880
ccctgggatg aaggtgagct gccagctcca ggtccagaga tccctgtgga cagccaccga	2940
ggaccagaaa atctacatct acaccctcaa gggcatgtgc cccttaaaca caccccaaca	3000
ggccttggat actccagctg tcgtcacctg cttcttggcc gtgcctgtta ttaaaaagaa	3060
ttcctacctg gtcttagcgg gcctcgccga tgggcttggt gctgtgtttc ccgtgggtgcg	3120
gggcacccca aaggacagct gtcctacct gtgctcacac acagccaaca ggtccaagtt	3180
cagcatcgcg gatgaagacg cacggcagaa cccctaccca gtgaaggcca tggagggtgt	3240
caacagcggc tctgaggtct ggtacagcaa tgggccgggc ctcttgtca tcgactgtgc	3300
ctccctggag atctgcaggc ggctggagcc ctacatggcc ccctccatgg ttacgtcagt	3360
cgtgtgcagc tctgagggca gaggggagga ggtcgtctgg tgcttgatg acaaggccaa	3420
ctccttggtg atgtaccact ccaccacct ccagctgtgt gcccggtact tctgcggggt	3480
ccccagcccc ctcagggaca tgtttcccg tgcggccctg gacacggaac ccccggcagc	3540
cagccacacg gccaacccaa aggtgcctga gggggactcc atcgcgagc tgagcatcat	3600
gtacagtgag gagctgggca cgcagatcct gatccaccag gaatcactca ctgactactg	3660
ctccatgtcc tcctactcct catccccacc ccgccaggct gccaggctcc cctcaagcct	3720
ccccagctcc ccagcaagtt cttccagtgt gcctttctcc accgactgcg aggactcaga	3780
catgctacat acgcccgtg ctgcctccga caggctctgag catgacctga ccccatgga	3840
cggggagacc ttcagccagc acctgcaggc cgtgaagatc ctgcgcgtca gagacctcat	3900
ttgggtcccc aggcgcggtg gagatgttat cgtcattggc ctggagaagg attctgaagc	3960
ccagcggggc cgagtcattg ccgtcttaaa agcccagag ctgactccgc atgggggtgct	4020
ggtggatgct gccgtggtgg caaaggacac tgttgtgtgc accttgaaa atgaaaacac	4080
agagtgggtg ctggccgtct ggaggggctg gggcgccagg gagttcgaca ttttctacca	4140
gtcctacgag gagctgggcc ggctggaggc ttgcactcgc aagagaaggt aattcctgtg	4200
gaatgactgt cacaca	4216
<210> SEQ ID NO 11	
<211> LENGTH: 22	
<212> TYPE: DNA	
<213> ORGANISM: Artificial	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<223> OTHER INFORMATION: forward primer	
<400> SEQUENCE: 11	
ctgtgacctg aagtcggaca ac	22
<210> SEQ ID NO 12	
<211> LENGTH: 21	
<212> TYPE: DNA	
<213> ORGANISM: Homo sapiens	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	

-continued

<223> OTHER INFORMATION: reverse primer	
<400> SEQUENCE: 12	
ggaatgactg cctcgaaatc c	21

- The invention claimed is:
1. An isolated and purified protein consisting of the amino acid sequence shown in SEQ ID NO:2.
  2. An isolated and purified protein consisting an amino acid sequence which is at least 90% identical to the amino acid sequence shown in SEQ ID NO:2 and which has a kinase activity.
  3. An isolated and purified polynucleotide comprising at least one strand which consists of a coding sequence for the amino acid sequence shown in SEQ ID NO:2.
  4. The polynucleotide of claim 3 wherein the coding sequence is the nucleotide sequence shown in SEQ ID NO: 1.
  5. The polynucleotide of claim 3 which is a cDNA.
  6. An isolated and purified single-stranded polynucleotide consisting of at least 8 contiguous nucleotides of the nucleotide sequence shown in SEQ ID NO: 1 or its complement.
  7. An expression construct, comprising;  
the polynucleotide comprising at least one strand which consists of a coding sequence for the amino acid sequence shown in SEQ ID NO:2; and  
a promoter which is located upstream from the coding sequence and which controls expression of the coding sequence.
  8. The expression construct of claim 7 wherein the coding sequence consists of the nucleotide sequence shown in SEQ ID NO: 1.
  9. A host cell comprising an expression construct, wherein the expression construct comprises:  
a polynucleotide comprising at least one strand which consists of a coding sequence for the amino acid sequence shown in SEQ ID NO:2; and  
a promoter which is located upstream from the coding sequence and which controls expression of the coding sequence.
  10. The host cell of claim 9 which is prokaryotic.
  11. The host cell of claim 9 which is eukaryotic.
  12. A method of producing a protein, comprising the steps of:  
culturing a host cell in a culture medium, wherein the host cell comprises an expression construct comprising (a) a polynucleotide comprising at least one strand which consists of a coding sequence for the amino acid sequence shown in SEQ ID NO:2 and (b) a promoter which is located upstream from the coding sequence and which controls expression of the coding sequence, wherein the step of culturing is carried out under conditions whereby the protein is expressed; and  
recovering the protein.
  13. A composition comprising:  
a protein consisting of the amino acid sequence shown in SEQ ID NO:2; and  
a pharmaceutically acceptable carrier.
  14. A composition comprising:  
a polynucleotide comprising at least one strand which consists of a coding sequence for the amino acid sequence shown in SEQ ID NO:2; and  
a pharmaceutically acceptable carrier.
  15. The composition of claim 14 wherein the polynucleotide consists of the nucleotide sequence shown in SEQ ID NO: 1.
  16. The isolated and purified protein of claim 2 which is at least 96% identical.
  17. The isolated and purified protein of claim 2 which is at least 98% identical.
  18. The isolated and purified protein of claim 2 which is at least 99% identical.
- \* \* \* \* \*