



US007230080B2

(12) **United States Patent**
Matz et al.

(10) **Patent No.:** **US 7,230,080 B2**
(45) **Date of Patent:** **Jun. 12, 2007**

(54) **FLUORESCENT AND COLORED PROTEINS,
AND POLYNUCLEOTIDES THAT ENCODE
THESE PROTEINS**

(75) Inventors: **Mikhail Vladimirovitch Matz**, Palm
Coast, FL (US); **Naila Omar Khayyam**
Alieva, Palm Coast, FL (US); **Karen**
Ann Konzen, St. Augustine, FL (US);
Steven Field, Daytona Beach, FL (US);
Anyah Salih, Waterloo (AU)

(73) Assignees: **University of Florida Research**
Foundation, Inc., Gainesville, FL (US);
University of Sydney, Sydney (AU)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **11/058,952**

(22) Filed: **Feb. 16, 2005**

(65) **Prior Publication Data**
US 2006/0183133 A1 Aug. 17, 2006

(51) **Int. Cl.**
C07K 1/00 (2006.01)
C12N 15/00 (2006.01)
C12N 9/12 (2006.01)
C12N 1/20 (2006.01)

(52) **U.S. Cl.** **530/350**; 435/69.1; 435/252.3;
435/320.1

(58) **Field of Classification Search** 530/350;
435/69.1, 252.3, 320.1
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,268,463 A 12/1993 Jefferson

5,491,084 A	2/1996	Chalfie et al.
5,639,663 A	6/1997	Crosby et al.
5,804,387 A	9/1998	Cormack et al.
5,874,304 A	2/1999	Zolotukhin et al.
5,919,445 A	7/1999	Chao
5,958,713 A	9/1999	Thastrup et al.
5,968,738 A	10/1999	Anderson et al.
5,968,750 A	10/1999	Zolotukhin et al.
5,976,796 A	11/1999	Szalay et al.
5,985,577 A	11/1999	Bulinski
6,020,192 A	2/2000	Muzyczka et al.
6,066,476 A	5/2000	Tsien et al.
6,455,759 B1	9/2002	Vierstra et al.

OTHER PUBLICATIONS

Anderluh G. et al., "Cloning, Sequencing, and Expression of
Equinatoxin II," *Biochemical and Biophysical Research Communi-*
cations, 1996, 220:437-442.

Ando, R. et al., "An Optical Marker Based on the UV-Induced
Green-to-Red Photoconversion of a Fluorescent Protein," *Proceed-*
ings of the National Academy of Sciences (2002), 99(20):12651-
12656.

Chudakov D. M. et al., "Kindling Fluorescent Proteins for Precise
in vivo Photolabeling," *Nature Biotechnology*, 2003, 21:191-194.

(Continued)


Primary Examiner—Maryam Monshipouri

(74) *Attorney, Agent, or Firm*—Saliwanchik, Lloyd &
Saliwanchik

(57) **ABSTRACT**

The subject invention provides new fluorescent and/or col-
ored proteins, and polynucleotide sequences that encode
these proteins. The subject invention further provides mate-
rials and methods useful for expressing these detectable
proteins in biological systems.

2 Claims, 21 Drawing Sheets



OTHER PUBLICATIONS

- Eichinger L. et al., "Dictyostelium as Model System for Studies of the Actin Cytoskeleton by Molecular Genetics," *Microscopy Research and Technique*, 1999, 47:124-134.
- Falk M. M. et al., "High Resolution, Fluorescent Deconvolution Microscopy and Tagging With the Autofluorescent Tracers CFP, GFP, and YFP to Study the Structural Composition of Gap Junctions in Living Cells," *Microscopy Research and Technique*, 2001, 52:251-262.
- Fradkov A. F. et al., "Novel Fluorescent Protein From Discosoma Coral and Its Mutants Possesses a Unique Far-Red Fluorescent," *FEBS Letters*, 2000, 479:127-130.
- Gurskaya N. G. et al., "GFP-like Chromosomes as a Source of Far-Red Fluorescent Proteins," *FEBS Letters*, 2001, 507:16-20.
- Gurskaya N. G. et al., "Color Transitions in Coral's Fluorescent Proteins by Site-Directed Mutagenesis," *BMC Biochemistry*, 2001, 2:6.
- Hanson M. R. et al., "GFP Imaging: Methodology and Application to Investigate Cellular Compartmentation in plants," *Journal of Experimental Botany*, 2001, 52:529-539.
- Hillisch A. et al., "Recent Advances in FRET: Distance Determination in Protein-DNA Complexes," *Current Opinion in Structural Biology*, 2001, 11:201-207.
- Houtsmuller A. B. et al., "Macromolecular Dynamics in Living Cell Nuclei Revealed by Fluorescent Redistribution After Photobleaching," *Histochem Cell Biol*, 2001, 115:13-21.
- Kallal L. et al., "Using Green Fluorescent Proteins to Study G-Protein-Coupled Receptor Localization Trafficking," *Trends Pharmacol Sci*, 21:175-180, 2000.
- Labas Y. A. et al., "Diversity and Evolution of the Green Fluorescent Protein Family," *Proc Natl Acad Sci USA*, 2002, 99:4256-4261.
- Laird D. W. et al., "Comparative Analysis and Application of Fluorescent Protein-Tagged Connexins," *Microscopy and Research Technique*, 2001, 52:263-272.
- Lukyanov K. A. et al., "Natural Animal Coloration Can Be Determined by a Nonfluorescent Green Fluorescent Protein Homolog," *J Biol Chemistry*, 2000, 275(34):25879-25882.
- Macek, P. et al., "Intrinsic Tryptophan Fluorescence of Equinatoxin II, a Pore-Forming Polypeptide From the Sea Anemone *Actinia equina* L, Monitors Its Interaction With Lipid Membranes," *European Journal of Biochemistry* (1995), 234:329-335.
- Martynov V. I. et al., "Alternative Cyclization in GFP-like Proteins Family," *J Biol Chem*, 2001, 276:21012-6.
- Matz M. V. et al., "Family of the Green Fluorescent Protein: Journey to the End of the Rainbow," *Bioessays*, 2002, 24:953-959.
- Matz M. V. et al., "Fluorescent Proteins From Nonbioluminescent Anthozoa Species," *Nature Biotechnol*, 1999, 17:969-973.
- Patterson G. H. et al., "A Photoactivatable GFP for Selective Photolabeling of Proteins and Cells," *Science*, 2002, 297:1873-1877.
- Pollok B. A. and Roger Heim, "Using GFP in FRET-based Applications," *Cell Biology*, 1999, 9:57-60.
- Reits E. et al., "From Fixed to FRAP: Measuring Protein Mobility and Activity in Living Cells," *Nature Cell Biology*, 2001, 3:E145-147.
- Terskikh A. et al., "Fluorescent Timer: Protein That Changes Color With Time," *Science*, 2000, 290:1585-8.
- Tsien R. Y., "The Green Fluorescent Protein," *Annu Rev Biochem*, 1998, 67:509-544.
- Tsien R. Y., "Rosy Dawn for Fluorescent Proteins," *Nat Biotech*, 1999, 17:956-957.
- Verkhusha V. V., et al., "An Enhanced Mutant of Red Fluorescent Protein DsRed for Developmental Timer of Neural Fiber Bundle Formation," *Journal of Biological Chemistry*, 2001, 276:29621-29624.
- Ward W. W. et al., "An Energy Transfer Protein in Coelenterate Bioluminescence," *J Biol Chem*, 1979, 254:781-788.
- Yanushevich Y. G. et al., "A Strategy for the Generation of Non-Aggregating Mutants of Anthozoa Fluorescent Proteins," *FEBS Letters*, 2002, 511:11-14.
- Yarbrough D. et al., "Refined Crystal Structure of DsRed, a Red Fluorescent Protein From Coral, at 2.0-A Resolution," *Proc Natl Acad Sci USA*, 2001, 98:462-7.




Fig. 1




Fig. 2




Fig. 3




Fig. 4




Fig. 5




Fig. 6




Fig. 7




Fig. 8





Fig. 9

**Fig. 10**




Fig. 11




Fig. 12




Fig. 13




Fig. 14




Fig. 15




Fig. 16




Fig. 17





Fig. 18

**Fig. 19**




Fig. 20




Fig. 21

FLUORESCENT AND COLORED PROTEINS, AND POLYNUCLEOTIDES THAT ENCODE THESE PROTEINS

GOVERNMENT SUPPORT

The subject matter of this application has been supported in part by U.S. Government Support under NIH RO1 GM066243-01. Accordingly, the U.S. Government has certain rights in this invention.

FIELD OF THE INVENTION

The present invention relates to novel fluorescent and colored proteins, and their use. These materials and methods are particularly advantageous for labeling and detection technology. Specifically, exemplified are novel colored and/or fluorescent proteins, and mutants thereof, isolated from marine organisms. These new proteins offer a wider array of colors and biochemical features compared to existing wild-type green fluorescent protein (GFP) or its modified variants utilized in current labeling and detection technology.

BACKGROUND OF THE INVENTION

Genetic markers are important for monitoring gene expression and tracking movement of proteins in cells. Markers have been extensively used for monitoring biological activity of genetic elements such as promoters, enhancers and terminators, and other aspects of gene regulation in numerous biological systems. Over the years numerous marker genes have been developed and utilized widely in molecular and genetic studies aimed at the identification, isolation and characterization of genetic regulatory elements and genes, and the development of gene transfer techniques.

In general, markers can be grouped into selectable markers and reporter markers. Selectable markers are typically enzymes with catalytic capability to convert chemical substrates usually harmful to host cells into non-toxic products, thus providing transformed host cells a conditionally selectable growth advantage under selective environment and allowing the recovery of stable transformants after transformation. A number of commonly used selectable markers include those that confer resistance characteristics to antibiotics (Gritz and Davies 1983; Bevan et al., 1983) and herbicides (De Block et al., 1987), and those with enzymatic activity to detoxify metabolic compounds that can adversely affect cell growth (Joersbo and Okkels 1996).

Reporter markers are compounds that provide biochemically assayable or identifiable activities. Reporter markers have been widely used in studies to reveal biological functions and modes of action of genetic elements such as promoters, enhancers, terminators, and regulatory proteins including signal peptides, transcription factors and related gene products. Over the years, several reporter markers have been developed for use in both prokaryotic and eukaryotic systems, including β -galactosidase (LacZ) (Stanley and Luzio 1984), β -glucuronidase (GUS) (Jefferson et al., 1987; U.S. Pat. No. 5,268,463), chloramphenicol acetyltransferase (CAT) (Gorman et al., 1982), green fluorescent protein (GFP) (Prasher et al., 1992; U.S. Pat. No. 5,491,084) and luciferase (Luc) (Ow et al., 1986).

Among reporter markers, GUS offers a sensitive and versatile reporting capability for gene expression in plants. β -glucuronidase or GUS, encoded by the uidA gene from *Escherichia coli*, catalyzes the conversion of several colorogenic and fluorogenic glucorogenic substrates such as p-ni-

trophenyl β -D-glucuronide and 4-methylumbelliferyl β -D-glucuronide into easily detectable products. GUS activity can be measured by highly sensitive colorimetric and fluorimetric methods (Jefferson et al., 1987). However, the GUS assay often requires total destruction of the sample tissues or exposure of sample tissues to phytotoxic chemical substrates. This prevents repeated use of the same sample tissue for continuous expression analysis and precludes the recovery of transformants from analyzed materials.

Recently, GFP isolated from the Pacific Northwest jellyfish (*Aequorea victoria*) has become an important reporter marker for non-destructive analysis of gene expression. GFP fluoresces in vivo by receiving light energy without the involvement of any chemical substrates. Thus, GFP is especially suitable for real time and continuous monitoring of temporal and spatial control of gene expression and protein activities without any physical damage to assay samples.

The gene for GFP has been cloned and used as a reporter gene, which can be expressed as a functional transgene in living organisms, marking the organisms with fluorescent color and thus allowing detection of those organisms. Accordingly, GFP has become a versatile fluorescent marker for monitoring a variety of physiological processes, visualizing protein localization and detecting the expression of transferred genes in various living systems, including bacteria, fungi, and mammalian tissues.

This in vivo labeling and detection technology was originally based on a single fluorescent protein: the green fluorescent protein from *Aequorea victoria*. Numerous modifications have been made to alter the spectral properties of GFP to provide for significant enhancement in fluorescence intensity (Prasher et al., 1992; Cubitt et al., 1995; Heim et al., 1994, 1995; Cormack et al., 1996; U.S. Pat. No. 5,804,387). In addition, GFP genes have been modified to contain more silent base mutations that correspond to codon-usage preferences in order to improve its expression efficacy, making it a reporter gene in both animal and plant systems (U.S. Pat. Nos. 5,874,304; 5,968,750; and 6,020,192).

In addition to GFP, there are now a number of other fluorescent proteins, substantially different from GFP, which are being developed into biotechnology tools. Most prominent of these proteins is the red fluorescent protein DsRed. See, for example, Labas, Y. A., N. G. Gurskaya, Y. G. Yanushevich, A. F. Fradkov, K. A. Lukyanov, S. A. Lukyanov and M. V. Matz. (2002) "Diversity and evolution of the green fluorescent protein family" *Proc Natl Acad Sci USA* 99:4256-4261 and Matz, M. V., K. A. Lukyanov and S. A. Lukyanov (2002) "Family of the green fluorescent protein: journey to the end of the rainbow" *Bioessays* 24: 953-959.

Labeling technologies based on GFP and related proteins have become indispensable in such areas as basic biomedical research, cell and molecular biology, transgenic research and drug discovery. The number of PubMed records containing the phrase "green fluorescent protein" exceeds 5500 only within the last three years. Demand for labeling and detection based on the fluorescent protein technology is large and steady.

Currently, there are very few known natural pigments essentially encoded by a single gene, wherein both the substrate for pigment biosynthesis and the necessary catalytic moieties are provided within a single polypeptide chain. The limited availability of fluorescent marker proteins makes the current technology based on fluorescent proteins very expensive, rendering it unaffordable and inaccessible to many mid-size (or smaller) companies that are interested in

using the technology. Therefore, there is a need for less expensive, readily available fluorescent and/or colored materials.

BRIEF SUMMARY OF THE INVENTION

The subject invention provides new fluorescent and/or colored proteins, and polynucleotide sequences that encode these proteins. The subject invention further provides materials and methods useful for expressing these detectable proteins in biological systems.

In specific embodiments, the subject invention provides advantageous fluorescent proteins. The invention also includes proteins substantially similar to, or mutants or variants of, the exemplified proteins.

Another aspect of the subject invention pertains to polynucleotide sequences that encode the detectable proteins of the present invention. In one embodiment, the present invention provides polynucleotide constructs comprising cDNA encoding novel colored and/or fluorescent proteins and mutants thereof.

In one embodiment, the invention provides nucleotide sequences of the inserts in pGEM-T vector (Promega), the conceptual translations of these inserts, and special properties of purified protein products.

The proteins and polynucleotides of the present invention can be used as described herein as colored and/or fluorescent (detectable) labels in a variety of ways, including but not limited to, as reporter genes for monitoring gene expression in living organisms, as protein tags for tracing the location of proteins within living cells and organisms, as reporter molecules for engineering various protein-based biosensors, and as genetically encoded pigments for modifying color and/or fluorescence of living organisms or their parts.

In a specific embodiment, the proteins of the subject invention can be used in molecular fluorescent tagging whereby the coding region of a protein of interest is fused with the coding region for a fluorescent protein of the subject invention. The product of such a gene shows the functional characteristics of the protein of interest, but bears the fluorescent label allowing tracing its movements.

Advantageously, the present invention provides proteins and polynucleotides to improve on the current technology of labeling and detection by offering a wider choice of colors and biochemical features never before provided by GFP and its modified variants.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows the design of bacterial expression constructs for the proteins of interests of the present invention.

FIG. 2 shows the excitation and emission spectrum of *Montipora millepora*, gbr1.

FIG. 3 shows the excitation and emission spectrum of *Echinophyllia echinata*, gbr3.

FIG. 4 shows the excitation and emission spectra of *Mycedium elephantotus*, gbr4.

FIG. 5 shows the excitation and emission spectra of *Mycedium elephantotus*, gbr5.

FIG. 6 shows the excitation and emission spectra of *Echinophyllia echinata*, gbr6.

FIG. 7 shows the excitation and emission spectra of *Echinophyllia echinata*, gbr7.

FIG. 8 shows the excitation and emission spectra of *Echinophyllia echinata*, gbr8.

FIG. 9 shows the excitation and emission spectra of *Fungia scutaria*, gbr9.

FIG. 10 shows the excitation and emission spectra of *Galaxea fascicularis*, gbr10.

FIG. 11 shows the excitation and emission spectra of *Montipora efflorescens*, gbr14.

FIG. 12 shows the excitation and emission spectra of *Porites porites*, gbr15.

FIG. 13 shows the excitation and emission spectra of *Porites porites*, gbr17.

FIG. 14 shows the excitation and emission spectra of *Stylocoeniella* sp., gbr18.

FIG. 15 shows the excitation and emission spectra of *Montipora efflorescens*, gbr24.

FIG. 16 shows the excitation and emission spectra of *Fungia cf danai*, gbr19.

FIG. 17 shows the excitation and emission spectra of *Fungia cf danai*, gbr20.

FIG. 18 shows the excitation and emission spectra of *Galaxea fascicularis*, gbr11.

FIG. 19 shows the excitation and emission spectra of *Goniopora djiboutiensis*, gbr21.

FIG. 20 shows the excitation and emission spectra of *Montipora efflorescens*, gbr22.

FIG. 21 shows the excitation and emission spectra of *Stylocoeriella* sp., gbr23.

BRIEF DESCRIPTION OF THE SEQUENCES

SEQ ID NO:1 is the 5' heel of an upstream primer used according to the subject invention.

SEQ ID NO:2 is the 5' heel of a downstream primer used according to the subject invention.

SEQ ID NO:3 is the open reading frame of the cDNA encoding the gbr1 protein of interest from *Montipora millepora*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:4 is the open reading frame of the cDNA encoding the gbr3 protein of interest from *Echinophyllia echinata*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:5 is the open reading frame of the cDNA encoding the gbr4 protein of interest from *Mycedium elephantotus*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:6 is the open reading frame of the cDNA encoding the gbr5 protein of interest from *Mycedium elephantotus*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:7 is the open reading frame of the cDNA encoding the gbr6 protein of interest from *Echinophyllia echinata*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature. A sequence identified in another misc_feature is derived from the cloning vector pGEM-T; it is included since in this particular construct it becomes translated during protein expression.

SEQ ID NO:8 is the open reading frame of the cDNA encoding the gbr7 protein of interest from *Echinophyllia echinata*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:9 is the open reading frame of the cDNA encoding the gbr8 protein of interest from *Echinophyllia echinata*. Parts of the sequence that have been artificially

added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:10 is the open reading frame of the cDNA encoding the gbr9 protein of interest from *Fungia scutaria*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:11 is the open reading frame of the cDNA encoding the gbr10 protein of interest from *Galaxea fascicularis*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:12 is the open reading frame of the cDNA encoding the gbr11 protein of interest from *Galaxea fascicularis*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:13 is the open reading frame of the cDNA encoding the gbr14 protein of interest from *Montipora efflorescens*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:14 is the open reading frame of the cDNA encoding the gbr15 protein of interest from *Porites porites*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:15 is the open reading frame of the cDNA encoding the gbr17 protein of interest from *Porites porites*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:16 is the open reading frame of the cDNA encoding the gbr18 protein of interest from *Stylocoeniella* sp. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:17 is the open reading frame of the cDNA encoding the gbr19 protein of interest from *Fungia cf danai*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:18 is the open reading frame of the cDNA encoding the gbr20 protein of interest from *Fungia cf danai*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature. SEQ ID NO:19 is the open reading frame of the cDNA encoding the gbr21 protein of interest from *Goniopora djiboutiensis*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:20 is the open reading frame of the cDNA encoding the gbr22 protein of interest from *Montipora efflorescens*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:21 is the open reading frame of the cDNA encoding the gbr23 protein of interest from *Stylocoeniella* sp. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:22 is the open reading frame of the cDNA encoding the gbr24 protein of interest from *Montipora efflorescens*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:23 is the open reading frame of the cDNA encoding the gbr25 protein of interest from *Montipora efflorescens*. Parts of the sequence that have been artificially added during the cloning process to facilitate gene expression in *E. coli* are identified in misc_feature.

SEQ ID NO:24 is the amino acid sequence encoded by SEQ ID NO:3.

SEQ ID NO:25 is the amino acid sequence encoded by SEQ ID NO:4.

SEQ ID NO:26 is the amino acid sequence encoded by SEQ ID NO:5.

SEQ ID NO:27 is the amino acid sequence encoded by SEQ ID NO:6.

SEQ ID NO:28 is the amino acid sequence encoded by SEQ ID NO:7.

SEQ ID NO:29 is the amino acid sequence encoded by SEQ ID NO:8.

SEQ ID NO:30 is the amino acid sequence encoded by SEQ ID NO:9.

SEQ ID NO:31 is the amino acid sequence encoded by SEQ ID NO:10.

SEQ ID NO:32 is the amino acid sequence encoded by SEQ ID NO:11.

SEQ ID NO:33 is the amino acid sequence encoded by SEQ ID NO:12.

SEQ ID NO:34 is the amino acid sequence encoded by SEQ ID NO:13.

SEQ ID NO:35 is the amino acid sequence encoded by SEQ ID NO:14.

SEQ ID NO:36 is the amino acid sequence encoded by SEQ ID NO:15.

SEQ ID NO:37 is the amino acid sequence encoded by SEQ ID NO:16.

SEQ ID NO:38 is the amino acid sequence encoded by SEQ ID NO:17.

SEQ ID NO:39 is the amino acid sequence encoded by SEQ ID NO:18.

SEQ ID NO:40 is the amino acid sequence encoded by SEQ ID NO:19.

SEQ ID NO:41 is the amino acid sequence encoded by SEQ ID NO:20.

SEQ ID NO:42 is the amino acid sequence encoded by SEQ ID NO:21.

SEQ ID NO:43 is the amino acid sequence encoded by SEQ ID NO:22.

SEQ ID NO:44 is the amino acid sequence encoded by SEQ ID NO:23.

DETAILED DESCRIPTION OF THE INVENTION

The present invention provides novel fluorescent and colored proteins isolated from marine organisms. In a particularly preferred embodiment, these proteins are fluorescent proteins. Specifically, exemplified herein are novel fluorescent proteins.

The subject invention further provides polynucleotide sequences encoding these proteins. These polynucleotide sequences include open reading frames encoding the specific exemplified detectable proteins, as well as expression constructs for expressing these proteins, for example, in bacterial hosts.

The proteins of the present invention can be readily, expressed by any one of the recombinant technology methods known to those skilled in the art having the benefit of the instant disclosure. The preferred method will vary depending upon many factors and considerations, including the

host, and the cost and availability of materials and other economic considerations. The optimum production procedure for a given situation will be apparent to those skilled in the art having the benefit of the current disclosure.

The subject invention also concerns cells transformed with a polynucleotide of the present invention comprising a nucleotide sequences encoding a novel detectable protein. These cells may be prokaryotic or eukaryotic, plant or animal. In one embodiment, animals, such as fish, are transformed to provide them with a unique color or ability to fluoresce. Polynucleotides providing the markers of the present invention are stable in a diverse range of hosts, including prokaryotic and eukaryotic organisms, and the translation products are fully functional and capable of providing assayable characteristics.

In another embodiment, the present invention provides methods to synthesize colored and fluorescent proteins in a recombinant cell.

In a specific embodiment, the proteins of the subject invention can be used in molecular fluorescent tagging whereby the coding region of a protein of interest is fused with the coding region for a fluorescent protein of the subject invention. The product of such a gene shows the functional characteristics of the protein of interest, but bears the fluorescent label allowing tracing its movements. See, for example, Eichinger, L., S. S. Lee and M. Schleicher (1999) "Dictyostelium as model system for studies of the actin cytoskeleton by molecular genetics" *Microsc Res Tech* 47:124-134; Falk, M. M. and U. Lauf (2001) "High resolution, fluorescence deconvolution microscopy and tagging with the autofluorescent tracers CFP, GFP, and YFP to study the structural composition of gap junctions in living cells" *Microsc Res Tech* 52:251-262; Kallal, L. and J. L. Benovic (2000) "Using green fluorescent proteins to study G-protein-coupled receptor localization and trafficking" *Trends Pharmacol Sci* 21:175-180; and Laird, D. W., K. Jordan, T. Thomas, H. Qin, P. Fistouris and Q. Shao (2001) "Comparative analysis and application of fluorescent protein-tagged connexins" *Microsc Res Tech* 52:263-272.

In a further embodiment, the subject invention concerns polynucleotides comprising an in-frame fusion of nucleotide sequences encoding multiple genetic markers. In one embodiment, the polynucleotides encode the genetic markers GUS, and a detectable protein of the subject invention.

The subject invention helps to provide a more abundant and diverse collection of proteins, which can be used in place of a GFP protein, such that new proteins are readily available for commercial exploitation by small companies that cannot take advantage of the current technology for financial reasons.

Definitions

As used herein, the terms "nucleic acid" and "polynucleotide" refer to a deoxyribonucleotide, ribonucleotide, or a mixed deoxyribonucleotide and ribonucleotide polymer in either single- or double-stranded form, and unless otherwise limited, would encompass known analogs of natural nucleotides that can function in a similar manner as naturally-occurring nucleotides.

As used herein, "a vector" is a DNA sequence having the elements necessary for the transcription/translation of a gene. Such elements would include, for example, promoters. Various classes of promoters are well known in the art and can be obtained commercially or assembled from the sequences and methods, which are also well known in the art. A number of vectors are available for expression and/or

cloning, and include, but are not limited to, pBR322, pUC series, M13 series, and pBLUESCRIPT vectors (Stratagene, La Jolla, Calif.).

As used herein, the term "expression construct" refers to a combination of nucleic acid sequences that provides for transcription of an operably linked nucleic acid sequence. As used herein, the term "operably linked" refers to a juxtaposition of the components described wherein the components are in a relationship that permits them to function in their intended manner. In general, operably linked components are in contiguous relation.

Detectable Proteins

The subject invention provides novel fluorescent and/or colored proteins. The novel colored and fluorescent proteins of the present invention can be detected using standard long-wave UV light sources or, preferably, optical designs appropriate for detecting agents with the excitation/emission characteristics of the proteins exemplified herein (see, for example, FIGS. 2-21). These proteins are referred to herein as "detectable proteins" or "marker proteins." The interaction of two or more residues of the protein and external agents such as molecular oxygen give rise to the colored and/or fluorescent feature of the proteins.

Advantageously, the use of these proteins facilitate real-time detection in vivo, a substrate is not required, and the relatively small size make the proteins very advantageous.

Substitution of amino acids other than those specifically exemplified or naturally present in the genetic marker proteins of the invention are also contemplated within the scope of the present invention. Such substitutions will create "variant proteins" within the scope of the subject invention. Variants and fragments preferably have emission and excitation maxima within 10 nm of the values shown in FIGS. 2-21. For example, non-natural amino acids can be substituted for the amino acids of the marker proteins, so long as a marker protein having the substituted amino acids retains its ability to be detected through fluorescence and/or color. Examples of non-natural amino acids include, but are not limited to, ornithine, citrulline, hydroxyproline, homoserine, phenylglycine, taurine, iodotyrosine, 2,4-diaminobutyric acid, α -amino isobutyric acid, 4-aminobutyric acid, 2-amino butyric acid, γ -amino butyric acid, ϵ -amino hexanoic acid, 6-amino hexanoic acid, 2-amino isobutyric acid, 3-amino propionic acid, norleucine, norvaline, sarcosine, homocitrulline, cysteic acid, τ -butylglycine, τ -butylalanine, phenylglycine, cyclohexylalanine, β -alanine, fluoro-amino acids, designer amino acids such as β -methyl amino acids, C-methyl amino acids, N-methyl amino acids, and amino acid analogues in general. Non-natural amino acids also include amino acids having derivatized side groups. Furthermore, any of the amino acids in the protein can be of the D (dextrorotary) form or L (levorotary) form. Allelic variants of a protein sequence of a detectable protein used in the present invention are also encompassed within the scope of the invention.

Amino acids can be generally categorized in the following classes: non-polar, uncharged polar, basic, and acidic. Conservative substitutions whereby a marker protein having an amino acid of one class is replaced with another amino acid of the same class fall within the scope of the subject invention so long as a marker protein having the substitution still is detectable Table 1 below provides a listing of examples of amino acids belonging to each class.

TABLE 1

Class of Amino Acid	Examples of Amino Acids
Nonpolar	Ala, Val, Leu, Ile, Pro, Met, Phe, Trp
Uncharged Polar	Gly, Ser, Thr, Cys, Tyr, Asn, Gln
Acidic	Asp, Glu
Basic	Lys, Arg, His

Polynucleotides

cDNA sequences encoding the proteins of the present invention are provided. Polynucleotides of the present invention can be composed of either RNA or DNA. Preferably, the polynucleotides are composed of DNA. The subject invention also encompasses those polynucleotides that are complementary in sequence to the polynucleotides disclosed herein.

Specifically exemplified are DNA sequences that encode novel fluorescent proteins. These DNA sequences are set forth in SEQ. ID NOS. 3–23.

Sequences of the subject invention may utilize codons preferred for expression by the selected host strains. These sequences may also have sites for cleavage by restriction enzymes, and/or initial, terminal, or intermediate DNA sequences which facilitate construction of readily expressed vectors.

Because of the degeneracy of the genetic code, a variety of different polynucleotide sequences can encode the detectable proteins of the present invention. In addition, it is well within the skill of a person trained in the art to create alternative polynucleotide sequences encoding the same, or essentially the same, detectable proteins of the subject invention. These variant or alternative polynucleotide sequences are within the scope of the subject invention. As used herein, references to “essentially the same” sequence refers to sequences which encode amino acid substitutions, deletions, additions, or insertions which do not eliminate the detectability of the polypeptide encoded by the polynucleotides of the present invention. Allelic variants of the nucleotide sequences encoding a genetic marker protein of the invention are also encompassed within the scope of the invention.

The subject invention also concerns variants of the polynucleotides of the present invention that encode detectable proteins. Variant sequences include those sequences wherein one or more nucleotides of the sequence have been substituted, deleted, and/or inserted. The nucleotides that can be substituted for natural nucleotides of DNA have a base moiety that can include, but is not limited to, inosine, 5-fluorouracil, 5-bromouracil, hypoxanthine, 1-methylguanine, 5-methylcytosine, and tritylated bases. The sugar moiety of the nucleotide in a sequence can also be modified and includes, but is not limited to, arabinose, xylulose, and hexose. In addition, the adenine, cytosine, guanine, thymine, and uracil bases of the nucleotides can be modified with acetyl, methyl, and/or thio groups. Sequences containing nucleotide substitutions, deletions, and/or insertions can be prepared and tested using standard techniques known in the art.

Polynucleotides and polypeptides of the subject invention can also be defined in terms of more particular identity and/or similarity ranges with those exemplified herein. The sequence identity will typically be greater than 60%, preferably greater than 75%, more preferably greater than 80%, even more preferably greater than 90%, and can be greater than 95%. The identity and/or similarity of a sequence can

be 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, or 99% as compared to a sequence exemplified herein. Unless otherwise specified, as used herein percent sequence identity and/or similarity of two sequences can be determined using the algorithm of Karlin and Altschul (1990), modified as in Karlin and Altschul (1993). Such an algorithm is incorporated into the NBLAST and XBLAST programs of Altschul et al. (1990). BLAST searches can be performed with the NBLAST program, score=100, wordlength=12, to obtain sequences with the desired percent sequence identity. To obtain gapped alignments for comparison purposes, Gapped BLAST can be used as described in Altschul et al. (1997). When utilizing BLAST and Gapped BLAST programs, the default parameters of the respective programs (NBLAST and XBLAST) can be used.

The subject invention also contemplates those polynucleotide molecules having sequences that are sufficiently homologous with the polynucleotide sequences exemplified herein so as to permit hybridization with that sequence under standard stringent conditions and standard methods (Maniatis et al. 1982). As used herein, “stringent” conditions for hybridization refers to conditions wherein hybridization is typically carried out overnight at 20–25 C below the melting temperature (T_m) of the DNA hybrid in 6× SSPE, 5× Denhardt’s solution, 0.1% SDS, 0.1 mg/ml denatured DNA. The melting temperature, T_m, is described by the following formula (Beltz et al., 1983):

$$T_m = 81.5 \text{ } ^\circ\text{C} + 16.6 \text{ Log } [\text{Na}^+] + 0.41(\% \text{ G+C}) - 0.61(\% \text{ formamide}) - 600/\text{length of duplex in base pairs.}$$

Washes are typically carried out as follows:

(1) Twice at room temperature for 15 minutes in 1× SSPE, 0.1% SDS (low stringency wash).

(2) Once at T_m–20 C for 15 minutes in 0.2× SSPE, 0.1% SDS (moderate stringency wash).

The polynucleotide sequences include the DNA strand sequence that is transcribed into RNA and the strand sequence that is complementary to the DNA strand that is transcribed. The polynucleotide sequences also include both full-length sequences as well as shorter sequences derived from the full-length sequences. The polynucleotide sequence includes both the sense and antisense strands either as individual strands or in the duplex.

Recombinant Hosts

Polynucleotide molecules containing DNA sequences encoding the colored and/or fluorescent proteins of the present invention can be introduced into a variety of host cells including bacterial cells, yeast cells, fungal cells, plant cells and animal cells. Methods by which the exogenous genetic material can be introduced into such host cells are well known in the art.

In one embodiment, the invention provides a bacteria cell capable of expressing the novel colored and fluorescent proteins.

Plants, plant tissues, and plant cells bred to contain, or transformed with, a polynucleotide of the invention are also contemplated by the present invention. In one embodiment, the polynucleotide encodes a detectable polypeptide shown in SEQ ID NOS. 7–10, or a functional fragment or variant thereof. Plants within the scope of the present invention include monocotyledonous plants, such as rice, wheat, barley, oats, sorghum, maize, sugarcane, pineapple, onion, bananas, coconut, lilies, grasses, and millet; and dicotyledonous plants, such as peas, alfalfa, tomato, melon, chick-

pea, chicory, clover, kale, lentil, soybean, tobacco, potato, sweet potato, radish, cabbage, rape, apple trees, grape, cotton, sunflower, and lettuce; and conifers. Techniques for transforming plant cells with a gene are known in the art and include, for example, *Agrobacterium* infection, biolistic methods, electroporation, calcium chloride treatment, etc. Transformed cells can be selected, redifferentiated, and grown into plants using standard methods known in the art. The progeny of any transformed plant cells or plants are also included within the scope of the present invention.

The subject invention also concerns non-human transgenic animals which have incorporated into the host cell genome a polynucleotide of the invention. Methods for producing transgenic animals, including mice, rats, pigs, sheep, cows, fish, and the like are well known in the art.

The subject invention also concerns methods for isolating transformants expressing a transgene. In one embodiment, an expression construct of the present invention comprising a transgene of interest operably linked to a nucleotide sequence encoding a detectable marker of the present invention is used to transform a cell. Methods for transforming cells are well known in the art. Transformed cells expressing the transgene are selected by identifying those cells expressing a genetic marker of the invention.

Expression Constructs

An expression construct of the invention typically comprises a structural gene sequence (encoding a protein), an antisense sequence, or other polynucleotide sequences, or a site for insertion of such sequences, operably linked to a polynucleotide of the present invention encoding a marker. The structural gene can be a gene encoding a protein from a prokaryotic or eukaryotic organism, for example, a human, mammal, insect, plant, bacteria, or virus. Proteins that can be encoded by a gene sequence include, but are not limited to, enzymes, hormones, cytokines, interleukins, receptors, growth factors, immunoglobulins, transcription factors, and *Bacillus thuringiensis* (B.t.) crystal toxin proteins. Sequences encoding B.t. proteins which have codon usage for preferential expression in plants are described in U.S. Pat. Nos. 5,380,831; 5,567,862; 5,567,600; 6,013,523; and 6,015,891. An antisense sequence is a sequence wherein the RNA transcribed from the antisense sequence is at least partially complementary to RNA transcribed from a gene encoding a protein.

Expression constructs of the invention will also generally include regulatory elements that are functional in the intended host cell in which the expression construct is to be expressed. Thus, a person of ordinary skill in the art can select regulatory elements for use in, for example, bacterial host cells, yeast host cells, plant host cells, insect host cells, mammalian host cells, and human host cells. Regulatory elements include promoters, transcription termination sequences, translation termination sequences, enhancers, and polyadenylation elements.

An expression construct of the invention can comprise a promoter sequence operably linked to a polynucleotide sequence encoding a marker of the invention. Promoters can be incorporated into a polynucleotide using standard techniques known in the art. Multiple copies of promoters or multiple promoters can be used in an expression construct of the invention. In a preferred embodiment, a promoter can be positioned about the same distance from the transcription start site as it is from the transcription start site in its natural genetic environment. Some variation in this distance is

permitted without substantial decrease in promoter activity. A transcription start site is typically included in the expression construct.

For expression in prokaryotic systems, an expression construct of the invention can comprise promoters such as, for example, alkaline phosphatase promoter, tryptophan (trp) promoter, lambda P_L promoter, β -lactamase promoter, lactose promoter, phoA promoter, T3 promoter, T7 promoter, or tac promoter (de Boer et al., 1983).

If the expression construct is to be provided in a plant cell, plant viral promoters, such as, for example, the cauliflower mosaic virus (CaMV) 35S (including the enhanced CaMV 35S promoter (see, for example U.S. Pat. No. 5,106,739)) or 19S promoter can be used. Plant promoters such as prolifera promoter, Ap3 promoter, heat shock promoters, T-DNA 1'- or 2'-promoter of *A. tumefaciens*, polygalacturonase promoter, chalcone synthase A (CHS-A) promoter from petunia, tobacco PR-1a promoter, ubiquitin promoter, actin promoter, alcA gene promoter, pin2 promoter (Xu et al., 1993), maize WipI promoter, maize trpA gene promoter (U.S. Pat. No. 5,625,136), maize CDPK gene promoter, and RUBISCO SSU promoter (U.S. Pat. No. 5,034,322) can also be used. Seed-specific promoters such as the promoter from a β -phaseolin gene (of kidney bean) or a glycinin gene (of soybean), and others, can also be used. Constitutive promoters (such as the CaMV, ubiquitin, actin, or NOS promoter), tissue-specific promoters (such as the E8 promoter from tomato), developmentally-regulated promoters, and inducible promoters (such as those promoters that can be induced by heat, light, hormones, or chemicals) are contemplated for use with the polynucleotides of the invention.

For expression in animal cells, an expression construct of the invention can comprise suitable promoters that can drive transcription of the polynucleotide sequence. If the cells are mammalian cells, then promoters such as, for example, actin promoter, metallothionein promoter, NF-kappaB promoter, EGR promoter, SRE promoter, IL-2 promoter, NFAT promoter, osteocalcin promoter, SV40 early promoter and SV40 late promoter, Lck promoter, BMP5 promoter, TRP-1 promoter, murine mammary tumor virus long terminal repeat promoter, STAT promoter, or an immunoglobulin promoter can be used in the expression construct. The baculovirus polyhedrin promoter can be used with an expression construct of the invention for expression in insect cells. Promoters suitable for use with an expression construct of the invention in yeast cells include, but are not limited to, 3-phosphoglycerate kinase promoter, glyceraldehyde-3-phosphate dehydrogenase promoter, metallothionein promoter, alcohol dehydrogenase-2 promoter, and hexokinase promoter.

Expression constructs of the invention may optionally contain a transcription termination sequence, a translation termination sequence, signal peptide sequence, and/or enhancer elements. Transcription termination regions can typically be obtained from the 3' untranslated region of a eukaryotic or viral gene sequence. Transcription termination sequences can be positioned downstream of a coding sequence to provide for efficient termination. Signal peptides are a group of short amino terminal sequences that encode information responsible for the relocation of an operably linked mature polypeptide to a wide range of post-translational cellular destinations, ranging from a specific organelle compartment to sites of protein action and the extracellular environment. Targeting marker gene products to an intended cellular and/or extracellular destination through the use of operably linked signal peptide sequence is contemplated for use with the polypeptides of the inven-

tion. Enhancers are cis-acting elements that increase activity of a promoter and can also be included in the expression construct. Enhancer elements are known in the art, and include, but are not limited to, the CaMV 35S enhancer element, maize shrunken-1 enhancer element, cytomegalovirus (CMV) early promoter enhancer element, and the SV40 enhancer element.

DNA sequences which direct polyadenylation of the mRNA encoded by the structural gene can also be included in the expression construct. The expression constructs of the invention can also include a polynucleotide sequence that directs transposition of other genes, i.e., a transposon.

Applications

There are many ways in which the novel proteins of the subject invention can be used. In one embodiment, the proteins can be used to identify cells. In these methods the proteins can be used to express fluorescence in a cell. One use for this method is in pre-labeling isolated cells or a population of similar cells prior to exposing the cells to an environment in which different cell types are present. Detection of fluorescence in only the original cells allows the location of such cells to be determined and compared with the total population.

A second group of methods concerns the identification of cells that have been transformed with exogenous DNA of interest. Identifying cells transformed with exogenous DNA is required in many in vitro procedures as well as in in vivo applications such as gene therapy.

In one embodiment of the subject invention, a polynucleotide sequence encoding a protein of the subject invention is fused to a DNA sequence encoding a selected protein in order to directly label the encoded protein. Expressing such a fluorescent and/or colored protein in a cell results in the production of labeled proteins that can be readily detected. This is useful in confirming that a protein is being produced by a chosen host cell. It also allows the location of the selected protein to be determined.

Cells that have been transformed with exogenous DNA can also be identified without creating a fusion protein. Here, the method relies on the identification of cells that have received a plasmid or vector that comprises at least two transcriptional or translational units. A first unit encodes and directs expression of the desired protein, while the second unit encodes and directs expression of the detectable protein. Co-expression of the detectable protein from the second transcriptional or translational unit ensures that cells containing the vector are detected and differentiated from cells that do not contain the vector.

In methods to produce fluorescent molecular weight markers, a gene sequence is generally fused to one or more DNA sequences that encode proteins having defined amino acid sequences and the fusion proteins are expressed from an expression vector. Expression results in the production of fluorescent proteins of defined molecular weight or weights that may be used as markers (following calculation of the size of the complete amino acid sequence).

Amino acid replacements that produce different color forms permit simultaneous use of multiple reporter genes. Different colored proteins can be used to identify multiple cell populations in a mixed cell culture or to track multiple cell types, enabling differences in cell movement or migration to be visualized in real time without the need to add additional agents or fix or kill the cells.

Other options include tracking and determining the ultimate location of multiple proteins within a single cell, tissue or organism; differential promoter analysis in which gene

expression from two different promoters is determined in the same cell, tissue or organism; and FACS sorting of mixed cell populations.

The techniques that can be used with spectrally separable proteins are exemplified by confocal microscopy, flow cytometry, and fluorescence activated cell sorting (FACS) using modular flow, dual excitation techniques.

In one embodiment, the subject invention concerns polynucleotides comprising an in-frame fusion of nucleotide sequences encoding multiple genetic markers. For example, a polynucleotide of the invention may comprise a first nucleotide sequence that is operably linked in-frame to a second nucleotide sequence. The polynucleotide encodes the amino acid sequences of the detectable protein and another genetic marker such that the genetic markers are in direct contact with one another, i.e., where the last amino acid of the fluorescent genetic marker is immediately contiguous with the first amino acid of the other genetic marker, or they can be separated by a peptide linker sequence, for example, as described in U.S. Pat. No. 5,891,680 and Li et al., 2001, that do not substantially alter functional activity of the genetic markers.

The subject invention also concerns kits comprising in one or more containers and a polynucleotide and/or protein of the present invention.

Additional useful applications of the technology described herein include, but are not limited to, the following:

FRET—Fluorescence Resonant Energy Transfer: This technique allows observation and quantification of molecular interactions. It requires at least two fluorescent proteins of different colors. Currently the most widely used pair is CFP and YFP (mutated variants of GFP); the proteins of the subject invention may be substituted for either or both of them.

REFERENCES

1. Hanson, M. R. and R. H. Kohler. 2001. GFP imaging: methodology and application to investigate cellular compartmentation in plants. *J Exp Bot* 52: 529–539.
2. Pollok, B. A. and R. Heim. 1999. Using GFP in FRET-based applications. *Trends Cell Biol* 9: 57–60.
3. Schuttrigkeit, T. A., U. Zachariae, T. von Feilitzsch, J. Wiehler, J. von Hummel, B. Steipe and M. E. Michel-Beyerle. 2001. Picosecond time-resolved FRET in the fluorescent protein from *Discosoma Red* (wt-DsRed). *Chemphyschem* 2: 325–328.
4. Hillisch, A., M. Lorenz and S. Diekmann. 2001. Recent advances in FRET: distance determination in protein-DNA complexes. *Curr Opin Struct Biol* 11: 201–207.
- FRAP—Fluorescence Redistribution After Photobleaching: This technique quantifies the dynamics of tagged molecules or the reporter molecules themselves. It involves in photobleaching (burning out) of all the fluorescent molecules within a small area by intense excitation light and monitoring the process of fluorescence recovery within this area (due to migration of tagged molecules from adjacent areas).

REFERENCES

1. Reits, E. A. and J. J. Neefjes. 2001. From fixed to FRAP: measuring protein mobility and activity in living cells. *Nat Cell Biol* 3: E145–147.

15

2. Houtsmuller, A. B. and W. Vermeulen. 2001. Macromolecular dynamics in living cell nuclei revealed by fluorescence redistribution after photobleaching. *Histochem Cell Biol* 115:13–21.

“Fluorescent timer” applications: one of the proteins exemplified herein—scubRFP—due to its natural spectroscopic properties, can be used as a reporter that changes color with time. Such reporters make it possible to estimate the time elapsed since the reporter protein was synthesized by quantifying its color. In addition, since the maturation speed (the rate of conversion from green to red) in scubRFP can be increased by UV-A light, it is possible to adjust its timing scale: experiments that need timing in shorter intervals may use appropriate background UV illumination to speed up the green-to-red conversion.

REFERENCES

1. Terskikh, A. V., A. Fradkov, A. Zaiskiy, A. V. Kajava, M. Matz, S. Kim, I. Weissman and P. Siebert. 2000. “Fluorescent timer”: Protein that changes color over time. *Molecular Biology of the Cell* 11: 648.
2. Verkhusha, V. V., H. Otsuna, T. Awasaki, H. Oda, S. Tsukita and K. Ito. 2001. An enhanced mutant of red fluorescent protein DsRed for double labeling and developmental timer of neural fiber bundle formation. *Journal of Biological Chemistry* 276: 29621–29624.

“Light-inducible fluorescence”: since the red fluorescence of scubRFP can be induced by exposure to UV-A light, it is possible to use this protein as a light-inducible reporter. Such a reporter can be used for studying molecular dynamics, in a way that is analogous to FRAP (see above). A small area can be irradiated by the fluorescence-inducing light, after which the process of redistribution of active fluorescent molecules from the irradiated spot can be followed.

REFERENCES

1. Ando, R., H. Hama, M. Yamamoto-Hino, H. Mizuno and A. Miyawaki. 2002. An optical marker based on the UV-induced green-to-red photoconversion of a fluorescent protein. *Proceedings of the National Academy of Sciences of the United States of America* 99: 12651–12656.
2. Patterson, G. H. and J. Lippincott-Schwartz. 2002. A photoactivatable GFP for selective photolabeling of proteins and cells. *Science* 297: 1873–1877.
3. Chudakov, D. M., V. V. Belousov, A. G. Zaiskiy, V. V. Novoselov, D. B. Staroverov, D. B. Zorov, S. Lukyanov and K. A. Lukyanov. 2003. Kindling fluorescent proteins for precise in vivo photolabeling (vol 21, pg 191, 2003). *Nature Biotechnology* 21: 452–452.

Coloring of biological objects for decorative and other non-scientific purposes. Examples: producing decorative fish for aquariums; coloring of fur, wool and milk by means of genetic modifications of appropriate animals; and coloring of decorative plants. Such uses can be implemented by a person skilled in the art having the benefit of the teachings of the current disclosure.

All patents, patent applications, provisional applications, and publications referred to or cited herein are incorporated by reference in their entirety, including all figures and tables, to the extent they are not inconsistent with the explicit teachings of this specification.

16

Following are examples which illustrate procedures for practicing the invention. These examples should not be construed as limiting. All percentages are by weight and all solvent mixture proportions are by volume unless otherwise noted.

EXAMPLE 1

Bacterial Expression Construct

As illustrated in FIG. 1, to prepare a bacterial expression construct, the ORF of the target detectable protein can be amplified by means of polymerase chain reaction (PCR), using primers corresponding to the beginning and end of the protein’s ORF. The upstream primer can carry a 5'-heel ttgattgattgaaggagaaatcATG (SEQ ID NO:1), which encodes three termination codons in three frames (bold), followed by the ribosome binding site (underlined), 6 spacer bases and initiation ATG codon.

The downstream primer can encode a 6×His tag in place of the original termination codon (the heel sequence can be 5'-tta tta gtg atg gtg atg gtg atg (SEQ ID NO:2)), to facilitate protein purification by means of metal-affinity chromatography.

The products of amplification can be cloned into pGEM-T vector (Promega) using manufacturer-provided reagents and protocol. The expressing clones can be identified after overnight growth of the colonies by their fluorescent appearance.

EXAMPLE 2

Excitation and Emission Spectra of the Detectable Proteins

The excitation spectra were measured from the proteins purified after bacterial expression. The spectra are shown in FIGS. 2–21. Emission spectra (dotted lines) were measured using USB2000 uv-vis spectrometer (Ocean Optics), excitation spectra (solid lines)—using spectrofluorometer LS-50B (Perkin Elmer). The indicated positions of excitation and emission maxima are accurate within 5 nm.

EXAMPLE 3

Multiple Marker Constructs

There are several advantages associated with the use of fusion markers, including: 1) achievement of combined functionalities in a single transcription unit, 2) reduced usage of genetic elements, such as promoters and terminators, for expressing multiple marker genes, 3) reduced overall length of insertion sequences that may lead to increased transformation efficiency, and most importantly 4) elimination of molecular interactions between adjacent genetic elements. Such unwanted interactions are frequently encountered when multiple expression units associated with different marker genes are used simultaneously and often complicate the interpretation of expression results.

In an effort to improve marker functionality and versatility, several translational fusions between two genetic markers have been developed. Datla et al. (1991; U.S. Pat. No. 5,639,663) created a bifunctional fusion between GUS and neomycin phosphotransferase (NPTII) to provide a biochemically assayable reporter activity and a conditionally selectable growth advantage for use in plant transformation. Another bifunctional fusion, between GUS and GFP, was also developed to provide both indicative and assayable reporter activities for monitoring transient and stable trans-

gene expression in plant cells (Quaedvlieg et al., 1998). More recently, Li et al. (2001) constructed a bifunctional fusion between GFP and NPTII and successfully used this marker for continuous analysis of promoter activity and transgene expression in transgenic grape plants throughout the entire process of plant development.

Small portions of a protein that provide unique functions such as protein/DNA/substrate binding activity can be inserted into another heterologous protein to create a hybrid fusion with enhanced functionality and utility. In other cases, an entire gene or protein of interest has been fused in-frame to another heterologous gene or protein to form a double fusion to provide combined functionalities. Production of multiple proteins using fusion constructs composed of two genes from transgenic plants has been demonstrated previously (U.S. Pat. No. 6,455,759).

In one embodiment, the subject invention provides cells transformed with a polynucleotide of the present invention comprising an in-frame fusion of nucleotide sequences encoding multiple markers. Preferably, the polynucleotide sequence is provided in an expression construct of the invention. The transformed cell can be a prokaryotic cell, for example, a bacterial cell such as *E. coli* or *B. subtilis*, or the transformed cell can be a eukaryotic cell, for example, a plant or animal cell. Animal cells include human cells, mammalian cells, avian cells, fish cells and insect cells. Mammalian cells include, but are not limited to, COS, 3T3, and CHO cells.

Genetic markers that can be used in conjunction with the detectable proteins of the present invention are known in the art and include, for example, polynucleotides encoding proteins that confer a conditionally selective growth advantage, such as antibiotic resistance and herbicide-resistance; polynucleotides encoding proteins that confer a biochemically assayable reporter activity; and polynucleotides encoding proteins that confer an indicative reporter activity. Examples of polynucleotides encoding proteins providing antibiotic resistance include those that can provide for resistance to one or more of the following antibiotics: hygromycin, kanamycin, bleomycin, G418, streptomycin, paromomycin, and spectinomycin. Kanamycin resistance can be provided by neomycin phosphotransferase (NPTII). Examples of polynucleotides encoding proteins providing herbicide resistance include those that can provide for resistance to phosphinothricin acetyltransferase or glyphosate. Examples of genetic markers that confer assayable or indicative reporters activity that can be used in the present invention include, but are not limited to, polynucleotides encoding β -glucuronidase (GUS), β -galactosidase, chloramphenicol acetyltransferase (CAT), luciferase, nopaline synthase (NOS), and green fluorescence protein (GFP).

It should be understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 44

<210> SEQ ID NO 1
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' heel of upstream primer used according to subject invention

<400> SEQUENCE: 1

ttgattgatt gaaggagaaa tatcatg 27

<210> SEQ ID NO 2
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: 5' heel of a downstream primer used according to the subject invention

<400> SEQUENCE: 2

ttattagtga tggatgatgt gatg 24

<210> SEQ ID NO 3
<211> LENGTH: 720
<212> TYPE: DNA
<213> ORGANISM: Montipora millepora
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in *E. coli*.
<220> FEATURE:
<221> NAME/KEY: misc_feature

-continued

<222> LOCATION: (712)..(720)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 3	
ttgattgatt gaaggagaaa tatcatggct ctaccaagc agatgaaact gacataccac	60
atggagggga ctgttaatgg gcatttcttt ataatcaagg gcgaaggcgg tggagagcct	120
tacgaaggaa cacatactat taagctgcaa gtggttgaag gaagtccact gccattctcc	180
cctgacatat tgtcgactgt gtttcaatac gaaacaggt gcttcactaa atatcccccc	240
aacatagttg actatttcaa gaactcatgt tctggtggcg gatatacatt tggaaggtct	300
tttctctatg aagatggagc agtttgaca gccagtggag atataacatt gagctctgat	360
aagagtagct ttgaacacaa atccaagttt cttggagtca actttcctgc tgatggacct	420
gtgatgaaaa aggagacgac taattgggag ccacctgcg agaaaatgac acctaattggg	480
atgacattga taggggatgt cactgagttc cttctgaaga aagatggtaa acgttacaag	540
tgccagttcc acacatttca cgatgcaaag gagaagtcga gaaacatgcc aatgccagac	600
ttccacttcg tgcaacatga gatagaaagg aaagacctac ccggtcctat gcagacatgg	660
caactgacag aacatgctgc tgcatgtaaa aatgtttcac catcaccatc acatcactaa	720
<210> SEQ ID NO 4	
<211> LENGTH: 733	
<212> TYPE: DNA	
<213> ORGANISM: Echinophyllia echinata	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(24)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (712)..(733)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 4	
ttgattgatt gaaggagaaa tatcatgagt gtgtttaatc cagatatgaa gatcaagctg	60
tatatggaag gcgctgtaaa cgggcacaag ttcgagatta aaggagaagg aaacgggaag	120
ccttttgagg gaaaacagac catggacctg gcagtcgtag acggcggacc tctgcctttt	180
gctttcgata tcttgacaac ttcattcaat tacggcaaca gggatttcac caaataccca	240
gatactatag tagactatth caagccgtcg tttcctgagg ggtattcctg ggaacgaagc	300
atgacttacg aagatggagg catttgcatc gccacaaatg acataacact gctgaaagat	360
accgacgact cgaactatth ctactataaa attcgatttg atggtgtgaa ctttgctgcc	420
aatggtccag ttatgcagaa gaagaccgcg aaatgggagc catccactga gaaaatgtat	480
gtgcgtgatg gagtgctgaa ggggtgaagtt aacatggctc tgttgcttga aggaggtggc	540
cattaccgat gtgactttaa aactacctat aaagctaaga aggttgctcg gttgccaaagc	600
tatcactttg tggaccaccg tatagagatt ttaagccaca gcaaagatta caaccaagtt	660
aggctgcatg agcatgctga agctcattcc gggctgccga gacaagccaa gcatcaccat	720
caccatcact aaa	733
<210> SEQ ID NO 5	

-continued

<211> LENGTH: 726	
<212> TYPE: DNA	
<213> ORGANISM: Mycedium elephantotus	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(24)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (706)..(726)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 5	
ttgattgatt gaaggagaaa tatcatgagt gtgattaagc cagacatgag gataaggctg	60
caaatgcaag gcgcggtaaa cgggcacccg ttcgtgatta caggagaagg agagggcaag	120
ccttacgaag gaaaacacac tataaacctt acagtccaag acggtggacc tctccctttc	180
gctttcgata tcttaacgac agcattccag tacggcaaca gggatttcac caaataccca	240
aaagacatcc cagactattt caagcagtcg tttcccgcgg ggtattcctg ggagcgatgc	300
atgacgttcg aagacggagg cctttgcacc gtgtcgagcc acataaaaaat tgaaggtgac	360
tattttacct acgacattcg atttcatggt gtgaactttc cagccggtgg tccagtcatg	420
cagaagaaga cgctgagatg ggagccatcc actgagaata tgtatgtgcg tgatggagtg	480
ctggtggggg aggtagagag gactctgttg cttgaaggaa ataagcatca ccgatgtaac	540
ttcagaacta cttacaaagc taagaaagaa gtggtgttac cagaatatca ctttgtggat	600
caccgaatag agatattagg ccatgacaaa gattacaaca acgtggtggt gtatgagaat	660
gcggttgccc gccagcaggc ttctactctg ccaagcaagg ccaagcatca ccatcaccat	720
cactaa	726
<210> SEQ ID NO 6	
<211> LENGTH: 727	
<212> TYPE: DNA	
<213> ORGANISM: mycedium elephantotus	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(24)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (703)..(727)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 6	
ttgattgatt gaaggagaaa tatcatgaat gtgattaaac cagacatgaa gctcaagctg	60
cgcatggaag gcacggtaaa tgggcactac ttcgtgattg aaggagatgg taaaggcagg	120
ccttttgagg gaaaacagag tatggactta gatgtaaaag agggcggacc actgcctttc	180
gcctatgata tcttaacaac agcattccat tatggcaaca gggttttcgc agaataccca	240
gatcatatac cagactattt caaacagtca tttcctggag ggtattcctg ggaacgaagc	300
ctcacgtttg aagacggggg catttgcatc gccagaaacg acataaaaaat ggtaggcgac	360
actttctata atacagttcg atttgatggt gttaactttc cccccaatgg tccagtgatg	420
caaaggagga cccagaaatg ggagccatcc accgagaaaa tatatgtgcg tgatggagtg	480

-continued

ttgacgggtg acattacat ggctctgttg cttgaaggag gtgtccatta ccgatgtgac 540
ttcagaacta cttacaaagc taaggagaag ggcgtccagt tgccaggcta tcactttgta 600
gatcactgta tagaaatfff aagtcacgac aaagattata acaaggttaa actgtacgag 660
catgccgtag ctcattctgg attgccggac aacaaacggc aacatcacca tcaccatcac 720
taataaa 727

<210> SEQ ID NO 7
<211> LENGTH: 768
<212> TYPE: DNA
<213> ORGANISM: Echinophyllia echinata
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (697)..(712)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (713)..(768)
<223> OTHER INFORMATION: Sequence derived from the cloning vector pGEM-T; in this particular construct it becomes translated during protein expression.

<400> SEQUENCE: 7
ttgattgatt gaaggagaaa tatcatgaat gtgattaaac cagacatgaa gattaagctg 60
cgtatggaag gcgctgtaaa cgggcacaag tttgctattg aagggaagg aaacggccag 120
cccttcgagg gaaaacagac tatgaacctg aaagtcaaag aaggtggacc tctgcctttt 180
gcttacgata tcttgacaac aatattcaat tacggcaaca gggatattgt caaatacca 240
gatgatatag tagactatft caagcagtcg tttcccgagg gctattcctg ggaacgcagc 300
atgatttatg aagacggagg catttgcac gccacaaacg acataactft ggaaggtgat 360
tgtttcgtct ataaaattcg atttgatggg gtaaacfttc ccgccaaaag tccagftttg 420
cagaagatga cgaaaaaatg ggagccatcc actgagaaat tgtatgtacg tgatggagtg 480
ctgaagggtg atgttaacat ggctctgttg cttgaaggag gtggccactt ccggtgtgac 540
tttaaaacta cttacaaagc taaaaaggft gttcaactac cagattatca ctttgtggat 600
caccgcattg aaattatgag ccacgacaaa gattacaaca acgttaagct atgtgagcat 660
gccgaagctc attccgggct gccagggcag gcgaagcac accatcacca taatcccgcg 720
gcatggcgg ccgggagcat gcgacgtcgg gcccaattcg ccctatag 768

<210> SEQ ID NO 8
<211> LENGTH: 736
<212> TYPE: DNA
<213> ORGANISM: Echinophyllia echinata
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(30)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (709)..(709)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in

-continued

E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (715)..(736)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 8	
ttagtgatga ttgattgaag gagaaatatc atgaatgtga ttaaaccaga catgaagatc	60
aggctgcgta tggaaggcgc tgtaaacggg cacaagttcg taattatcgg aaaaggagat	120
ggcaagcctt acgagggaac tcagactatg gaccttgaag tcatagaggg cggacctctg	180
ccctttgctt ttgatatctt gacaacagta ttcaaatacg gcaacagggc tttcgttaaa	240
tatccaacgg atatagcaga ctatttcaag caatcgtttc ctgaagggtt ttcttgggag	300
cgaagcatga cttacgaaga cggaggaatt tgcatcgcca caaatgacat aacactaagt	360
aaagacatcg ccaactgctt tgattataac attcgatttg atgggtgtgaa ctttcccccg	420
aatagtccgg ttttgcagaa gacaacaata aagtgggagc cttccactga aaacatgtat	480
gtgcgtgatg gagttctgaa aggcgacatt aacatgtctc tgttgcttga aggagggtgca	540
ggccattacc ggtgtgactt caaaactact tacaaaagta agaaggctgt caagttgcc	600
gactatcact ttgtggacca ccgcattaca attgtaagcc acgacaagga ttacaacaaa	660
gtgaagctgc gtgagcatgc cgaagctcat tccgggctgc agatggagcc caagcatcac	720
catcaccatc actaaa	736
<210> SEQ ID NO 9	
<211> LENGTH: 710	
<212> TYPE: DNA	
<213> ORGANISM: Echinophyllia echinata	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(13)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (689)..(710)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 9	
aaggagaaat atcatgagtc tgattaaacc agaaatgaag atcaagctgc ttatggaagg	60
caatgtaaac gggcacccgt ttgttattga gggagatgga aaaggccatc cttttgaggg	120
aaaacagagt atggaccttg tagtcaaaga aggcgcacct ctcccttttg cctacgatat	180
cttgacaaca gcattccatt acggcaacag ggtttttgct aaatacccag accatatacc	240
agactacttc aagcagtcgt ttcccaacgg gttttcttgg gagcgaagcc tgatgttcga	300
ggacgggggc gtttgcacgc ccacaaatga cataaacactg gaaggagaca ctttctttaa	360
caaagttcga ttttatggtg taaactttcc cccaaatggt cctgttatgc agaagaagac	420
gctgaaatgg gaggcatcca ctgagaaaat gtatttgcgt gatggagtgt tgacgggcga	480
tattaccatg gctctgctgc ttaaaggaga tgtccattac cgatgtgact tcagaactac	540
ttacaaatct aggcaggagg gtgtcaagtt gccaggctat cactttgtcg atcactgcat	600
cagcattgtg agccatgaca aagactacac gaaggttaag ctgtatgagc atgctgttgc	660
ccatttggga ttgccgaaa acgtcaagca tcaccatcac catcactaaa	710

-continued

<210> SEQ ID NO 10
<211> LENGTH: 711
<212> TYPE: DNA
<213> ORGANISM: *Fungia scutaria*
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(12)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in *E. coli*.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (684)..(684)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in *E. coli*.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (691)..(711)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in *E. coli*.

<400> SEQUENCE: 10

aggagaaata tcatgagtgt gattgtaaag gaaatgatga ctaagctaca catggaaggt 60
actgttaacg ggcacgcctt tacaattgaa ggcaaaggaa aaggcgatcc ttacaatgga 120
gtgcagtcta tgaaccttga cgtcaaaggc ggtgcgcctt tgccgttctc tttcgatcct 180
ttgacgccag cattcatgta cggcaacaga gtgttcacga agtatccaga agacatacca 240
gactttttca agcaggtgtt tcctgaaggg taccactggg aaagaagtat tacctttgaa 300
gatcaggccg tttgtacggc aaccagccac ataaggctgg accagaaaga gatgtgtttt 360
atctatgacg tccgttttca cgggtgtgaac tttcccgcca atggcccaat catgcagaag 420
aagatactgg gatgggagcc atccactgag aaaatgtatg cacgtgatgg ggtgctgaag 480
ggtgatgtta atatgactct tcgtgttgaa ggaggtggcc attaccgagc tgacttcagg 540
actacttaca aagcaaagaa gccagtcaac ctgccaggct atcacttcat agaccaccgc 600
attgagatta ccaagcacag caaagattac accaatgttg ctttgtatga ggcagcagtt 660
gctcgtcatt ctccgtgcc taaagttgct catcaccatc accatcacta a 711

<210> SEQ ID NO 11
<211> LENGTH: 720
<212> TYPE: DNA
<213> ORGANISM: *Galaxea fascicularis*
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in *E. coli*.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (700)..(720)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in *E. coli*.

<400> SEQUENCE: 11

ttgattgatt gaaggagaaa tatcatgaat gtgattaaac cagacatgaa gatcaagctg 60
tgtatgagag gcactataaa cgggcataat ttcgtgattg aaggagaagg aaaaggaaac 120
ccttacgagg gaacgcagat tttagacctg aacgtcactg aaggcgcacc tctgcctttc 180
gcttacgata tcttgacaac agtgttccag tacggcaaca gggcattcac caagtaccca 240

-continued

ttgattgatt gaaggagaaa tatcatggct ctttcaaaga acggtgtcaa agacagaatg 60
aagctgaaat tccatatgga ggggagtgtc aacgggcatg aatttacaat caagggcgaa 120
ggcactgggc aaccttacga agggacacag tctattcaac tgcgtgtgga aaaagggggt 180
ccgttgccat tctccgtaga catattgtcg gctgtgtttc tgtacggaaa caggggtcttt 240
actaaatata ctcaagacct tgttgactat ttcaagaact cgtgtcctgc tggatataca 300
tggcaaaggt cttttctctt tgaagatggg gcagtttgca cagccagtgc agatataaca 360
gtgagtgttg aggagaactg cttttatcac gagtccaagt ttcatggagt gaactttcct 420
gctgatggac ctgtgatgaa aaagatgaca actaactggg aaccatcctg cgagaaaatc 480
acaccaatac ctaatgaggg gatattgaaa ggagatgtca ccatgttcct ccttctgaag 540
gatggtgggc gttaccggtg ccagttcgat acagtttaca aagcaaagtc tgacccaaaa 600
acgatcatga tgccggactg gcacttcata caacataagc tcaaccgcga agaccgcagc 660
gatgctaagc accagaaatg gcgactggta gaaaatgcta ttgcataaccg atccacatta 720
tcccatcacc atcaccatca ctaa 744

<210> SEQ ID NO 14
<211> LENGTH: 738
<212> TYPE: DNA
<213> ORGANISM: Porites porites
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (718)..(738)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.

<400> SEQUENCE: 14

ttgattgatt gaaggagaaa tatcatggct ctttcaaagc aaagtggggt caaagatgta 60
atgaacaccg agcttcatat ggacgggatc gtcaatggac acccctttga gataaaagga 120
aaaggaaagg gaaacccgta caaggggtgtg cagaccatga agcttacagt cattaagggt 180
gcgcctttgc ctttttctat tgacattttg ctgcctcaac acatgtatgg aagcaagcca 240
tttattaagt atcctgagag tatcccagac tacatcaagt tgtcatttcc cgaggggaatc 300
acatgggaaa ggtccatgac ctttgaagat ggtgcagtgt gcactgtctc taacgactcc 360
aggctcgatg gcgactcttt catctacgaa gtcaggtttc ttggcgtgaa ctttccccga 420
gatggacctg ttatgcagaa gaagacgcga ggctgggacc cgtccacaga gagactgtat 480
gagtgtggtg ggtggcagag aggagacgtc cacatggcct tgaagttgga gaacgggtggc 540
cattatacgt gcgacttcaa aactacttac aaatcaaaga agggcttgaa ggtgccaccg 600
tatcacttcg ttgaccacaa actagattta ctgagccaca acaccgatgg tgctaccttt 660
gaagagtttg aacaacgaga aattgcacat gcacatcttt ctaacttacc ggtagcccat 720
caccatcacc atcactaa 738

<210> SEQ ID NO 15
<211> LENGTH: 689
<212> TYPE: DNA
<213> ORGANISM: Porites porites
<220> FEATURE:

-continued

<221> NAME/KEY: misc_feature		
<222> LOCATION: (1)..(14)		
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.		
<220> FEATURE:		
<221> NAME/KEY: misc_feature		
<222> LOCATION: (675)..(689)		
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.		
<400> SEQUENCE: 15		
taaggagaaa tatcatggct ctttcaaacc aagtgacaat gaaataccac atggatggca	60	
gattcgagga caaggagttt acaatcgagg gtgaaggcac agggaagccg tacgagggga	120	
agcaaaccgt gacactgtgg gtaaccaagg gtgcacccct cccattctcc tttgacatat	180	
tgtcggctgt gtttctctat ggtaacagag ccttcactga ttatcctaaa ggaatcgttg	240	
actattttcaa gccatctttt cctgaaggat attcatttga aagaactctt gaatttgagg	300	
atggcgggata ttgcacagcc agtgcgata taagtcttga cagtgcaagc aactgcttca	360	
tccacaagtc cagtttcaag ggcgtcaagt ttcctgacaa tggaccagtg aagcaaaaga	420	
agacaactaa ctgggagccg tccatcgaga aaatgactgt gcgtgacggg atattgaagg	480	
gtgatgttac catgttcctg tcgctgacag atggaggaaa tcatcgttgc cagttcagca	540	
ctttatacaa agcaaagaag gctgtcaagt tgccaacgga aagccactat gtggagcacc	600	
gcctggtgag gactgacctt cctaattgaa aagttcagtt ggaagagcat gctgctgcac	660	
gtttaaacac cgtgcatcac catcaataa	689	
<210> SEQ ID NO 16		
<211> LENGTH: 717		
<212> TYPE: DNA		
<213> ORGANISM: Stylocoeniella sp		
<220> FEATURE:		
<221> NAME/KEY: misc_feature		
<222> LOCATION: (1)..(24)		
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.		
<220> FEATURE:		
<221> NAME/KEY: misc_feature		
<222> LOCATION: (697)..(717)		
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.		
<400> SEQUENCE: 16		
ttgattgatt gaaggagaaa tatcatggct cttacaaagc agtgtatcgc aaacgaaatg	60	
acgatgactt tccacatgga tggctgcgtc aatggccatt actttactat cgaaggagaa	120	
ggctccggga ggccatacga ggggaagcag atgtcaaagt ttaaagtcac caaggggtggg	180	
ccccttccat tctcctttga catactatcg tcagcattca aatatggaaa tcgatgcttc	240	
actgcgtatc ctgccggcat gcacgactac ttcaaacaag catttcctga gggaatgtca	300	
tatgaaagga catttacctt tgaagatgga ggagttgcta cagcgagtgg ggacataagc	360	
cttaaaggta actgctttgt ccacaaatcc atgtttcacg gagtgaactt tcctgctgat	420	
ggacctgtga tgaaaaagaa gacaactggg tgggaccggt cctttgagaa aatgactgtg	480	
tgcaatggaa tattgaaggg cgatgttacc atgttcctca tgctcgaaga tggtaaaaat	540	
tacaaatgcc aattccacac ttcttacaag acaaagaaac cggttacgct gccatcaaac	600	
catgtcgtgg aacatcgcat tgtgaggacc aaccttgata aagctggcaa ccatgttcaa	660	

-continued

ctggatgagc atgctgttgc acatgttaat cctttgcac accatcacca tcaactaa 717

<210> SEQ ID NO 17
<211> LENGTH: 684
<212> TYPE: DNA
<213> ORGANISM: Fungia cf danai
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (664)..(684)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.

<400> SEQUENCE: 17

atgsctcttt caamrcaagy gattggaaaa gacatgaraa ttaactatth tatggatggc 60
agtgtgaacg ggcacgagtt tactgttaaa ggtgaaggca taggcaaacc ttacgagggg 120
caccatgaga tgacactacg cgtcactatg gctaagggcg ggccattgcc tttctcgtht 180
gacttattgt cacacacgth ctgttatggc aatagacctt tcactaaata ccctgaagag 240
atacccgact atttcaaaca agcatttcct gaaggcctgt catgggaaag gtcattgcag 300
ttcgaagatg gtgggtttgc tgcagtcaac gcgaacataa gccttaaagg agactgcttc 360
gagcacaatt ccaaatttgt tggcgthaac tttcccgcg agggtcctgt gatgcaaaac 420
aaaagtctgg attgggagcc atctaccgag aaaattactg tctccgacgg agtgctgaag 480
ggtgatgttc cgatgttcct aaagctcgtg ggaggtggta atcataaatg ccaattcacg 540
actacttaca aagcggccaa aaaggttctt gacatgcctc aaagccattt catcttccat 600
cgcctagtca ggaaaaccga aggcaacatt accaagctgg tagaggatgt agaagctcat 660
aaccatcacc atcaccatca ctaa 684

<210> SEQ ID NO 18
<211> LENGTH: 738
<212> TYPE: DNA
<213> ORGANISM: Fungia cf danai
<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (1)..(24)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.

<220> FEATURE:
<221> NAME/KEY: misc_feature
<222> LOCATION: (718)..(738)
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.

<400> SEQUENCE: 18

ttgattgatt gaaggagaaa tatcatgtct tattcaaagc agggcatcgt acaagaaatg 60
aagacgaaat accgtatgga aggcagtgtc aatggccatg aattcacgat cgaagggtga 120
ggaactgggt acccttacga agggaaacag atgtccgaat tagtgatcat caagcctaag 180
ggaaagcccc ttccattctc ctttgacata ctgtcatcag tctttcaata tggaaacagg 240
tgcttcacaa agtaccctgc agacatgcct gactatttca agcaagcatt cccagatgga 300
atgtcatatg aaaggtcatt tctatttgag gatggagcag ttgctacagc cagctggaac 360
attcgtctcg aaggaaattg cttcatccac aattccatct ttcattggcg aaactttccc 420
gccgatggac ccgtaatgaa aaagaagaca attggctggg ataagtcctt cgaaaaaatg 480
actgtgtcta aagaggtgtt aacaggtgat gtgactatgt ttcttatgct cgaaggaggt 540

-continued

ggttaccaca gatgccagtt tcaactccact tacaaaaacag agaagccggt cgaactgccc	600
ccgaatcatg tcgtagaaca tcaaattgtg aggaccgacc ttggccaaag tgcaaaaggc	660
ttcacagtca agctggaagc acatgctgcg gtcctatgta accctttgaa ggttcaacat	720
caccatcacc atcactaa	738
 <210> SEQ ID NO 19 <211> LENGTH: 708 <212> TYPE: DNA <213> ORGANISM: Goniopora djiboutiensis <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (1)..(24) <223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli. <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (688)..(708) <223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli. <400> SEQUENCE: 19	
ttgattgatt gaaggagaaa tatcatgagt gtgatcgcta aacaaatgac ctacaaggtt	60
tatatgtcag gcacggtcaa tggacactac tttgaggtcc aaggcgatgg aaaaggaaag	120
ccttacgagg gggagcagac agtaaagctg actgtcacca aggggtggacc tctgccattt	180
gcttgggata ttttatcacc acaggctcag tacggaagca taccattcac caagtaccct	240
gaagacatcc ctgactatgt aaagcagtca ttccctgagg gatatacatg ggagaggatc	300
atgaactttg aagatgggtgc agtgtgtact gtcagcaatg attccagcat ccaaggcaac	360
tgtttcatct acaatgtcaa gttctctggt ttgaactttc ctcccagtgg accagtcatg	420
cagaagaaga cacagggctg ggaacccaac actgagcgtc tccttgcacg agatggaatg	480
ctgataggaa acaactttat ggctctgaag ttggaaggag gtggtcacta tttgtgtgaa	540
ttcaaatacta cctacaaggc aaagaagcct gtgaagatgc cagggtatca ctttgttgac	600
cgcaaactgg atgtaaccaa tcacaaccag gattacactt ccgttgagca gtgtgaaatt	660
tccattgcac gcaaacctgt ggtcgcccat caccatcacc atcactaa	708
 <210> SEQ ID NO 20 <211> LENGTH: 708 <212> TYPE: DNA <213> ORGANISM: Montipora efflorescens <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (1)..(24) <223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli. <220> FEATURE: <221> NAME/KEY: misc_feature <222> LOCATION: (688)..(708) <223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli. <400> SEQUENCE: 20	
ttgattgatt gaaggagaaa tatcatgagt gtgatcgcta aacaaatgac ctacaaggtt	60
tatatgtcag gcacggtcaa tggacactac tttgaggtcg aaggcgatgg aaaaggaaag	120
ccttacgagg gggagcagac ggtaaagctc actgtcacca aggggtggacc tctgccattt	180
gcttgggata ttttatcacc actgtctcaa tacggaagca taccattcac caagtaccct	240

-continued

gaagacatcc ctgattatgt aaagcagtca ttccctgagg gatatacatg ggagaggatc	300
atgaactttg aagatgggtgc agtgtgtact gtcagcaatg attccagcat ccaaggcaac	360
tgtttcatct acaatgtcaa aatctctggg gtgaactttc ctccaatgg acctgttatg	420
cagaagaaga cacagggctg ggaacccaac actgagcgtc tctttgcacg agatggaatg	480
ctgataggaa acaactttat ggctctgaag ttggaaggag gtggttacta tttgtgtgaa	540
ttcaaatacta cttacaaggc aaagaagcct gtgaggatgc cagggtatca ctatgttgac	600
cgcaaactgg atgtaaccag tcacaacaag gattacacat ttgttgagca gtgtgaaata	660
tccattgcac gccactcttt gctcgggtcat caccatcacc atcactaa	708
 <210> SEQ ID NO 21	
<211> LENGTH: 708	
<212> TYPE: DNA	
<213> ORGANISM: Stylocoeniella sp	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(24)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (688)..(708)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
 <400> SEQUENCE: 21	
ttgattgatt gaaggagaaa tatcatgagt gtgatcgcta aacaaatgac ctacaaggtt	60
tatatgtcag gcacggtcaa tggacactac tttgaggtcc aaggcgatgg aaaaggaaag	120
ccttacgagg gggagcagac agtaaggctc actgtcacca aggggtgggcc tctgccattt	180
gcatgggata ttttatcacc actgtctcag tacggaagca tacctttcac caagtaccct	240
gaagacatcc ctgattatgt aaagcagtca ttccctgagg gatatacatg ggagaggatc	300
atgaactttg aagatgggtgc agtgtgtact gtcagcaayg attccagcat ccaaggcaac	360
tgtttcatct acartgtcaa aatctctggg ttgaactttc ctccaatgg acctgttatg	420
cagaagaaga cacagggctg ggaacccaac actgagcgtc tctttgcacg agatggaatg	480
ctgataggaa acaactttat ggctctgaag ttggaaggag gtggtcacta tttgtgtgaa	540
ttcaaatacta cttacaaggc aaagaagcct gtgaggatgc cagggtatca ctatgttgac	600
cgcaaactgg atgtaaccag tcacaacagg gattacacat ctgttgagca gtgtgaaata	660
tccatagcac gccactcttt gctcgggtcat caccatcacc atcactaa	708
 <210> SEQ ID NO 22	
<211> LENGTH: 741	
<212> TYPE: DNA	
<213> ORGANISM: Montipora efflorescens	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(24)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (721)..(741)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	

-continued

<400> SEQUENCE: 22	
ttgattgatt gaaggagaaa tatcatggct ctttcaaagc aaagtctacc cagcgacatg	60
aaactgatat accacatgga tgggaatggt aatgggcatt cctttgtaat caagggcgaa	120
ggcgagggaa agccttacga agggacacat actattaagc tgcaagtggg tgaaggaagt	180
ccactgccat tctccgctga catattgtcg actgtgtttc aatacggaaa cagggtgcttc	240
acaaaatatc cccccaacat agttgactat ttcaagaact catgttctgg tggcggatat	300
aaatttggaa ggtcttttct ctatgaagat ggagcagttt gcacagccag tggagatata	360
acattgagcg ctgataagaa aagctttgaa cacaaatcca agtttcttgg agtcaacttt	420
cctgctgatg gacctgtgat gaaaaaggag acgactaatt gggagccatc ctgcgagaaa	480
atgacaccta atgggatgac attgataggg gatgtcactg ggttccttct gaaggaagat	540
ggtaaacggt acaagtgcc gttccacaca tttcacgatg caaaggataa gtcgaaaaag	600
atgccaatgc cagacttcca cttcgtgcaa cataagatag aaaggaaaga cctaccgggt	660
tctatgcaga catggcgact gacagaacat gctgctgcat gtaaacctg tttcactgag	720
catcaccatc accatcacta a	741
<210> SEQ ID NO 23	
<211> LENGTH: 747	
<212> TYPE: DNA	
<213> ORGANISM: Montipora efflorescens	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (1)..(24)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<220> FEATURE:	
<221> NAME/KEY: misc_feature	
<222> LOCATION: (727)..(747)	
<223> OTHER INFORMATION: Part of the sequence that has been artificially added during the cloning process to facilitate gene expression in E. coli.	
<400> SEQUENCE: 23	
ttgattgatt gaaggagaaa tatcatggct ctttcaaaga acggtctaac aaagaacatg	60
acgacgaaat accgcatgga aggggtgtgtc gatgggcata aatttgtaat cacgggcgac	120
ggcattggag atcctttcga ggggaaacag actagtattg atctgtgtgt ggttgaaggg	180
ggaccactgc cattctccga agatatattg tctgctgtgt ttgactacgg aaacagggtc	240
tttactaaat atcctcaaga ccttgttgac tatttcaaga actcgtgtcc tgctggatat	300
acatggcaaa ggtcttttct ctttgaagat ggtgcagttt gcacagccag tgcagatata	360
acagtgagtg ttgaggagaa ctgcttttat cacgagtcca agtttcatgg agtgaacttt	420
cctgctgatg gacctgtgat gaaaaagatg acaactaact gggaaccatc ctgcgagaaa	480
atcacaccaa tacctaata ggggatattg aaaggagatg tcaccatgtt cctccttctg	540
aaggatgggt ggcgttaccg gtgccagttc gatacagttt acaaagcaaa gtctgaccca	600
aaaacgatca tgatgccga ctggcacttc atccaacata agctcaaccg cgaagaccgc	660
agcgatgcta agcaccagaa atggcgactg gtagaaaatg ctattgcata ccgatccaca	720
ttatcccatc accatcacca tcaactaa	747
<210> SEQ ID NO 24	
<211> LENGTH: 231	
<212> TYPE: PRT	
<213> ORGANISM: Montipora millepora	

-continued

<400> SEQUENCE: 24

Met Ala Leu Pro Lys Gln Met Lys Leu Thr Tyr His Met Glu Gly Thr
1 5 10 15

Val Asn Gly His Phe Phe Ile Ile Lys Gly Glu Gly Gly Gly Glu Pro
20 25 30

Tyr Glu Gly Thr His Thr Ile Lys Leu Gln Val Val Glu Gly Ser Pro
35 40 45

Leu Pro Phe Ser Pro Asp Ile Leu Ser Thr Val Phe Gln Tyr Gly Asn
50 55 60

Arg Cys Phe Thr Lys Tyr Pro Pro Asn Ile Val Asp Tyr Phe Lys Asn
65 70 75 80

Ser Cys Ser Gly Gly Gly Tyr Thr Phe Gly Arg Ser Phe Leu Tyr Glu
85 90 95

Asp Gly Ala Val Cys Thr Ala Ser Gly Asp Ile Thr Leu Ser Ser Asp
100 105 110

Lys Ser Ser Phe Glu His Lys Ser Lys Phe Leu Gly Val Asn Phe Pro
115 120 125

Ala Asp Gly Pro Val Met Lys Lys Glu Thr Thr Asn Trp Glu Pro Ser
130 135 140

Cys Glu Lys Met Thr Pro Asn Gly Met Thr Leu Ile Gly Asp Val Thr
145 150 155 160

Glu Phe Leu Leu Lys Lys Asp Gly Lys Arg Tyr Lys Cys Gln Phe His
165 170 175

Thr Phe His Asp Ala Lys Glu Lys Ser Arg Asn Met Pro Met Pro Asp
180 185 190

Phe His Phe Val Gln His Glu Ile Glu Arg Lys Asp Leu Pro Gly Pro
195 200 205

Met Gln Thr Trp Gln Leu Thr Glu His Ala Ala Ala Cys Lys Asn Val
210 215 220

Ser Pro Ser Pro Ser His His
225 230

<210> SEQ ID NO 25
<211> LENGTH: 235
<212> TYPE: PRT
<213> ORGANISM: Echinophyllia echinata

<400> SEQUENCE: 25

Met Ser Val Phe Asn Pro Asp Met Lys Ile Lys Leu Tyr Met Glu Gly
1 5 10 15

Ala Val Asn Gly His Lys Phe Glu Ile Lys Gly Glu Gly Asn Gly Lys
20 25 30

Pro Phe Glu Gly Lys Gln Thr Met Asp Leu Ala Val Val Asp Gly Gly
35 40 45

Pro Leu Pro Phe Ala Phe Asp Ile Leu Thr Thr Ser Phe Asn Tyr Gly
50 55 60

Asn Arg Val Phe Thr Lys Tyr Pro Asp Thr Ile Val Asp Tyr Phe Lys
65 70 75 80

Pro Ser Phe Pro Glu Gly Tyr Ser Trp Glu Arg Ser Met Thr Tyr Glu
85 90 95

Asp Gly Gly Ile Cys Ile Ala Thr Asn Asp Ile Thr Leu Leu Lys Asp
100 105 110

Thr Asp Asp Ser Asn Tyr Phe Tyr Tyr Lys Ile Arg Phe Asp Gly Val
115 120 125

Asn	Phe	Ala	Ala	Asn	Gly	Pro	Val	Met	Gln	Lys	Lys	Thr	Ala	Lys	Trp
130						135			140						
Glu	Pro	Ser	Thr	Glu	Lys	Met	Tyr	Val	Arg	Asp	Gly	Val	Leu	Lys	Gly
145			150						155			160			
Glu	Val	Asn	Met	Ala	Leu	Leu	Leu	Glu	Gly	Gly	Gly	His	Tyr	Arg	Cys
			165						170			175			
Asp	Phe	Lys	Thr	Thr	Tyr	Lys	Ala	Lys	Lys	Val	Val	Arg	Leu	Pro	Ser
			180			185						190			
Tyr	His	Phe	Val	Asp	His	Arg	Ile	Glu	Ile	Leu	Ser	His	Ser	Lys	Asp
195						200						205			
Tyr	Asn	Gln	Val	Arg	Leu	His	Glu	His	Ala	Glu	Ala	His	Ser	Gly	Leu
210						215			220						
Pro	Arg	Gln	Ala	Lys	His	His	His	His	His	His					
225			230						235						

<400> SEQUENCE: 26

Met 1	Ser	Val	Ile 5	Lys	Pro	Asp	Met	Arg	Ile 10	Arg	Leu	Gln	Met	Gln 15	Gly
Ala	Val	Asn 20	Gly	His	Pro	Phe	Val	Ile 25	Thr	Gly	Glu	Gly 30	Glu	Gly	Lys
Pro	Tyr 35	Glu	Gly	Lys	His	Thr	Ile 40	Asn	Leu	Thr	Val	Gln 45	Asp	Gly	Gly
Pro 50	Leu	Pro	Phe	Ala	Phe	Asp 55	Ile	Leu	Thr	Thr	Ala 60	Phe	Gln	Tyr	Gly
Asn 65	Arg	Val	Phe	Thr 70	Lys	Tyr	Pro	Lys	Asp 75	Ile	Pro	Asp	Tyr	Phe	Lys 80
Gln	Ser	Phe	Pro 85	Ala	Gly	Tyr	Ser	Trp	Glu 90	Arg	Cys	Met	Thr 95	Phe	Glu
Asp	Gly	Gly 100	Leu	Cys	Thr	Val	Ser	Ser 105	His	Ile	Lys	Ile 110	Glu	Gly	Asp
Tyr	Phe 115	Thr	Tyr	Asp	Ile	Arg	Phe 120	His	Gly	Val	Asn 125	Phe	Pro	Ala	Gly
Gly 130	Pro	Val	Met	Gln	Lys 135	Lys	Thr	Leu	Arg	Trp	Glu 140	Pro	Ser	Thr	Glu
Asn 145	Met	Tyr	Val	Arg 150	Asp	Gly	Val	Leu	Val	Gly 155	Glu	Val	Glu	Arg	Thr 160
Leu	Leu	Leu	Glu 165	Gly	Asn	Lys	His	His	Arg 170	Cys	Asn	Phe	Arg	Thr 175	Thr
Tyr	Lys	Ala 180	Lys	Lys	Glu	Val	Val 185	Leu	Pro	Glu	Tyr	His 190	Phe	Val	Asp
His	Arg 195	Ile	Glu	Ile	Leu	Gly 200	His	Asp	Lys	Asp	Tyr	Asn 205	Asn	Val	Val
Val	Tyr 210	Glu	Asn	Ala	Val	Ala 215	Arg	Gln	Gln	Ala	Ser 220	Thr	Leu	Pro	Ser
Lys 225	Ala	Lys	His	His 230	His	His	His	His							

```
<210> SEQ ID NO 27
<211> LENGTH: 232
<212> TYPE: PRT
```

-continued

<213> ORGANISM: Mycedium elephantotus																							
<400> SEQUENCE: 27																							
Met	Asn	Val	Ile	Lys	Pro	Asp	Met	Lys	Leu	Lys	Leu	Arg	Met	Glu	Gly								
1				5					10					15									
Thr	Val	Asn	Gly	His	Tyr	Phe	Val	Ile	Glu	Gly	Asp	Gly	Lys	Gly	Arg								
			20					25					30										
Pro	Phe	Glu	Gly	Lys	Gln	Ser	Met	Asp	Leu	Asp	Val	Lys	Glu	Gly	Gly								
		35					40					45											
Pro	Leu	Pro	Phe	Ala	Tyr	Asp	Ile	Leu	Thr	Thr	Ala	Phe	His	Tyr	Gly								
	50					55					60												
Asn	Arg	Val	Phe	Ala	Glu	Tyr	Pro	Asp	His	Ile	Pro	Asp	Tyr	Phe	Lys								
65					70				75					80									
Gln	Ser	Phe	Pro	Gly	Gly	Tyr	Ser	Trp	Glu	Arg	Ser	Leu	Thr	Phe	Glu								
				85					90					95									
Asp	Gly	Gly	Ile	Cys	Ile	Ala	Arg	Asn	Asp	Ile	Lys	Met	Val	Gly	Asp								
			100					105					110										
Thr	Phe	Tyr	Asn	Thr	Val	Arg	Phe	Asp	Gly	Val	Asn	Phe	Pro	Pro	Asn								
		115					120					125											
Gly	Pro	Val	Met	Gln	Arg	Arg	Thr	Gln	Lys	Trp	Glu	Pro	Ser	Thr	Glu								
	130					135					140												
Lys	Ile	Tyr	Val	Arg	Asp	Gly	Val	Leu	Thr	Gly	Asp	Ile	Thr	Met	Ala								
145					150					155				160									
Leu	Leu	Leu	Glu	Gly	Gly	Val	His	Tyr	Arg	Cys	Asp	Phe	Arg	Thr	Thr								
			165						170					175									
Tyr	Lys	Ala	Lys	Glu	Lys	Gly	Val	Gln	Leu	Pro	Gly	Tyr	His	Phe	Val								
			180					185					190										
Asp	His	Cys	Ile	Glu	Ile	Leu	Ser	His	Asp	Lys	Asp	Tyr	Asn	Lys	Val								
		195					200					205											
Lys	Leu	Tyr	Glu	His	Ala	Val	Ala	His	Ser	Gly	Leu	Pro	Asp	Asn	Lys								
	210					215					220												
Arg	Gln	His	His	His	His	His	His																
225					230																		
<210> SEQ ID NO 28																							
<211> LENGTH: 247																							
<212> TYPE: PRT																							
<213> ORGANISM: Echinophyllia echinata																							
<400> SEQUENCE: 28																							
Met	Asn	Val	Ile	Lys	Pro	Asp	Met	Lys	Ile	Lys	Leu	Arg	Met	Glu	Gly								
1				5					10					15									
Ala	Val	Asn	Gly	His	Lys	Phe	Ala	Ile	Glu	Gly	Glu	Gly	Asn	Gly	Gln								
			20					25					30										
Pro	Phe	Glu	Gly	Lys	Gln	Thr	Met	Asn	Leu	Lys	Val	Lys	Glu	Gly	Gly								
		35					40					45											
Pro	Leu	Pro	Phe	Ala	Tyr	Asp	Ile	Leu	Thr	Thr	Ile	Phe	Asn	Tyr	Gly								
	50					55					60												
Asn	Arg	Val	Phe	Val	Lys	Tyr	Pro	Asp	Asp	Ile	Val	Asp	Tyr	Phe	Lys								
65					70				75					80									
Gln	Ser	Phe	Pro	Glu	Gly	Tyr	Ser	Trp	Glu	Arg	Ser	Met	Ile	Tyr	Glu								
				85					90					95									
Asp	Gly	Gly	Ile	Cys	Ile	Ala	Thr	Asn	Asp	Ile	Thr	Leu	Glu	Gly	Asp								
			100					105				110											
Cys	Phe	Val	Tyr	Lys	Ile	Arg	Phe	Asp	Gly	Val	Asn	Phe	Pro	Ala	Lys								

-continued

115					120					125						
Ser	Pro	Val	Leu	Gln	Lys	Met	Thr	Lys	Lys	Trp	Glu	Pro	Ser	Thr	Glu	
130					135					140						
Lys	Leu	Tyr	Val	Arg	Asp	Gly	Val	Leu	Lys	Gly	Asp	Val	Asn	Met	Ala	
145					150					155					160	
Leu	Leu	Leu	Glu	Gly	Gly	Gly	His	Phe	Arg	Cys	Asp	Phe	Lys	Thr	Thr	
					165					170					175	
Tyr	Lys	Ala	Lys	Lys	Val	Val	Gln	Leu	Pro	Asp	Tyr	His	Phe	Val	Asp	
					180					185					190	
His	Arg	Ile	Glu	Ile	Met	Ser	His	Asp	Lys	Asp	Tyr	Asn	Asn	Val	Lys	
					195					200					205	
Leu	Cys	Glu	His	Ala	Glu	Ala	His	Ser	Gly	Leu	Pro	Gly	Gln	Ala	Lys	
					210					215					220	
His	His	His	His	His	Asn	Pro	Ala	Ala	Met	Ala	Ala	Gly	Ser	Met	Arg	
225					230					235					240	
Arg	Arg	Ala	Gln	Phe	Ala	Leu										
					245											
<210> SEQ ID NO 29																
<211> LENGTH: 234																
<212> TYPE: PRT																
<213> ORGANISM: Echinophyllia echinata																
<400> SEQUENCE: 29																
Met	Asn	Val	Ile	Lys	Pro	Asp	Met	Lys	Ile	Arg	Leu	Arg	Met	Glu	Gly	
1					5					10					15	
Ala	Val	Asn	Gly	His	Lys	Phe	Val	Ile	Ile	Gly	Lys	Gly	Asp	Gly	Lys	
				20					25					30		
Pro	Tyr	Glu	Gly	Thr	Gln	Thr	Met	Asp	Leu	Glu	Val	Ile	Glu	Gly	Gly	
				35					40					45		
Pro	Leu	Pro	Phe	Ala	Phe	Asp	Ile	Leu	Thr	Thr	Val	Phe	Lys	Tyr	Gly	
50				55				60								
Asn	Arg	Ala	Phe	Val	Lys	Tyr	Pro	Thr	Asp	Ile	Ala	Asp	Tyr	Phe	Lys	
65				70				75				80				
Gln	Ser	Phe	Pro	Glu	Gly	Phe	Ser	Trp	Glu	Arg	Ser	Met	Thr	Tyr	Glu	
				85					90					95		
Asp	Gly	Gly	Ile	Cys	Ile	Ala	Thr	Asn	Asp	Ile	Thr	Leu	Ser	Lys	Asp	
				100					105					110		
Ile	Ala	Asn	Cys	Phe	Asp	Tyr	Asn	Ile	Arg	Phe	Asp	Gly	Val	Asn	Phe	
115								120					125			
Pro	Pro	Asn	Ser	Pro	Val	Leu	Gln	Lys	Thr	Thr	Ile	Lys	Trp	Glu	Pro	
130				135				140								
Ser	Thr	Glu	Asn	Met	Tyr	Val	Arg	Asp	Gly	Val	Leu	Lys	Gly	Asp	Ile	
145				150				155				160				
Asn	Met	Ser	Leu	Leu	Leu	Glu	Gly	Gly	Ala	Gly	His	Tyr	Arg	Cys	Asp	
				165				170				175				
Phe	Lys	Thr	Thr	Tyr	Lys	Ala	Lys	Lys	Ala	Val	Lys	Leu	Pro	Asp	Tyr	
				180				185				190				
His	Phe	Val	Asp	His	Arg	Ile	Thr	Ile	Val	Ser	His	Asp	Lys	Asp	Tyr	
195								200				205				
Asn	Lys	Val	Lys	Leu	Arg	Glu	His	Ala	Glu	Ala	His	Ser	Gly	Leu	Gln	
210				215				220								
Met	Glu	Pro	Lys	His	His	His	His	His	His							
225					230											

-continued

<210> SEQ ID NO 30
<211> LENGTH: 231
<212> TYPE: PRT
<213> ORGANISM: Echinophyllia echinata

<400> SEQUENCE: 30

Met Ser Leu Ile Lys Pro Glu Met Lys Ile Lys Leu Leu Met Glu Gly
1 5 10 15

Asn Val Asn Gly His Pro Phe Val Ile Glu Gly Asp Gly Lys Gly His
20 25 30

Pro Phe Glu Gly Lys Gln Ser Met Asp Leu Val Val Lys Glu Gly Ala
35 40 45

Pro Leu Pro Phe Ala Tyr Asp Ile Leu Thr Thr Ala Phe His Tyr Gly
50 55 60

Asn Arg Val Phe Ala Lys Tyr Pro Asp His Ile Pro Asp Tyr Phe Lys
65 70 75 80

Gln Ser Phe Pro Asn Gly Phe Ser Trp Glu Arg Ser Leu Met Phe Glu
85 90 95

Asp Gly Gly Val Cys Ile Ala Thr Asn Asp Ile Thr Leu Glu Gly Asp
100 105 110

Thr Phe Phe Asn Lys Val Arg Phe Tyr Gly Val Asn Phe Pro Pro Asn
115 120 125

Gly Pro Val Met Gln Lys Lys Thr Leu Lys Trp Glu Ala Ser Thr Glu
130 135 140

Lys Met Tyr Leu Arg Asp Gly Val Leu Thr Gly Asp Ile Thr Met Ala
145 150 155 160

Leu Leu Leu Lys Gly Asp Val His Tyr Arg Cys Asp Phe Arg Thr Thr
165 170 175

Tyr Lys Ser Arg Gln Glu Gly Val Lys Leu Pro Gly Tyr His Phe Val
180 185 190

Asp His Cys Ile Ser Ile Val Ser His Asp Lys Asp Tyr Thr Lys Val
195 200 205

Lys Leu Tyr Glu His Ala Val Ala His Leu Gly Leu Pro Glu Asn Val
210 215 220

Lys His His His His His His
225 230

<210> SEQ ID NO 31
<211> LENGTH: 232
<212> TYPE: PRT
<213> ORGANISM: Fungia scutaria

<400> SEQUENCE: 31

Met Ser Val Ile Val Lys Glu Met Met Thr Lys Leu His Met Glu Gly
1 5 10 15

Thr Val Asn Gly His Ala Phe Thr Ile Glu Gly Lys Gly Lys Gly Asp
20 25 30

Pro Tyr Asn Gly Val Gln Ser Met Asn Leu Asp Val Lys Gly Gly Ala
35 40 45

Pro Leu Pro Phe Ser Phe Asp Leu Leu Thr Pro Ala Phe Met Tyr Gly
50 55 60

Asn Arg Val Phe Thr Lys Tyr Pro Glu Asp Ile Pro Asp Phe Phe Lys
65 70 75 80

Gln Val Phe Pro Glu Gly Tyr His Trp Glu Arg Ser Ile Thr Phe Glu
85 90 95

-continued

<210> SEQ ID NO 33
<211> LENGTH: 227
<212> TYPE: PRT
<213> ORGANISM: Galaxea fascicularis

<400> SEQUENCE: 33
Met Ser Val Ile Ala Lys Gln Met Thr Tyr Lys Val Tyr Met Ser Gly
1 5 10 15
Thr Val Asn Gly His Tyr Phe Glu Val Glu Gly Asp Gly Lys Gly Lys
20 25 30
Pro Tyr Glu Gly Glu Gln Thr Val Lys Leu Thr Val Thr Lys Gly Gly
35 40 45
Pro Leu Pro Phe Ala Trp Asp Ile Leu Ser Pro Gln Ser Gln Tyr Gly
50 55 60
Ser Ile Pro Phe Thr Lys Tyr Pro Glu Asp Ile Pro Asp Tyr Val Lys
65 70 75 80
Gln Ser Phe Pro Glu Gly Tyr Thr Trp Glu Arg Ile Met Asn Phe Glu
85 90 95
Asp Gly Ala Val Cys Thr Val Ser Asn Asp Ser Ser Ile Gln Gly Asn
100 105 110
Cys Phe Ile Tyr His Val Lys Phe Ser Gly Leu Asn Phe Pro Pro Asn
115 120 125
Gly Pro Val Met Gln Lys Lys Thr Gln Gly Trp Glu Pro Asn Thr Glu
130 135 140
Arg Leu Phe Ala Arg Asp Gly Met Leu Ile Gly Asn Asn Phe Met Ala
145 150 155 160
Leu Lys Leu Glu Gly Gly Gly His Tyr Leu Cys Glu Phe Lys Ser Thr
165 170 175
Tyr Lys Ala Lys Lys Pro Val Lys Met Pro Gly Tyr His Tyr Val Asp
180 185 190
Arg Lys Leu Asp Val Thr Asn His Asn Lys Asp Tyr Thr Ser Val Glu
195 200 205
Gln Cys Glu Ile Ser Ile Ala Arg Lys Ser Val Val Ala His His His
210 215 220
His His His
225

<210> SEQ ID NO 34
<211> LENGTH: 239
<212> TYPE: PRT
<213> ORGANISM: Montipora efflorescens

<400> SEQUENCE: 34
Met Ala Leu Ser Lys Asn Gly Val Lys Asp Arg Met Lys Leu Lys Phe
1 5 10 15
His Met Glu Gly Ser Val Asn Gly His Glu Phe Thr Ile Lys Gly Glu
20 25 30
Gly Thr Gly Gln Pro Tyr Glu Gly Thr Gln Ser Ile Gln Leu Arg Val
35 40 45
Glu Lys Gly Gly Pro Leu Pro Phe Ser Val Asp Ile Leu Ser Ala Val
50 55 60
Phe Leu Tyr Gly Asn Arg Val Phe Thr Lys Tyr Pro Gln Asp Leu Val
65 70 75 80
Asp Tyr Phe Lys Asn Ser Cys Pro Ala Gly Tyr Thr Trp Gln Arg Ser
85 90 95

-continued

Phe Leu Phe Glu Asp Gly Ala Val Cys Thr Ala Ser Ala Asp Ile Thr
100 105 110

Val Ser Val Glu Glu Asn Cys Phe Tyr His Glu Ser Lys Phe His Gly
115 120 125

Val Asn Phe Pro Ala Asp Gly Pro Val Met Lys Lys Met Thr Thr Asn
130 135 140

Trp Glu Pro Ser Cys Glu Lys Ile Thr Pro Ile Pro Asn Glu Gly Ile
145 150 155 160

Leu Lys Gly Asp Val Thr Met Phe Leu Leu Leu Lys Asp Gly Gly Arg
165 170 175

Tyr Arg Cys Gln Phe Asp Thr Val Tyr Lys Ala Lys Ser Asp Pro Lys
180 185 190

Thr Ile Met Met Pro Asp Trp His Phe Ile Gln His Lys Leu Asn Arg
195 200 205

Glu Asp Arg Ser Asp Ala Lys His Gln Lys Trp Arg Leu Val Glu Asn
210 215 220

Ala Ile Ala Tyr Arg Ser Thr Leu Ser His His His His His His
225 230 235

<210> SEQ ID NO 35
<211> LENGTH: 237
<212> TYPE: PRT
<213> ORGANISM: Porites porites

<400> SEQUENCE: 35

Met Ala Leu Ser Lys Gln Ser Gly Val Lys Asp Val Met Asn Thr Glu
1 5 10 15

Leu His Met Asp Gly Ile Val Asn Gly His Pro Phe Glu Ile Lys Gly
20 25 30

Lys Gly Lys Gly Asn Pro Tyr Lys Gly Val Gln Thr Met Lys Leu Thr
35 40 45

Val Ile Lys Gly Ala Pro Leu Pro Phe Ser Ile Asp Ile Leu Leu Pro
50 55 60

Gln His Met Tyr Gly Ser Lys Pro Phe Ile Lys Tyr Pro Glu Ser Ile
65 70 75 80

Pro Asp Tyr Ile Lys Leu Ser Phe Pro Glu Gly Ile Thr Trp Glu Arg
85 90 95

Ser Met Thr Phe Glu Asp Gly Ala Val Cys Thr Val Ser Asn Asp Ser
100 105 110

Arg Leu Asp Gly Asp Ser Phe Ile Tyr Glu Val Arg Phe Leu Gly Val
115 120 125

Asn Phe Pro Arg Asp Gly Pro Val Met Gln Lys Lys Thr Arg Gly Trp
130 135 140

Asp Pro Ser Thr Glu Arg Leu Tyr Glu Cys Gly Gly Trp Gln Arg Gly
145 150 155 160

Asp Val His Met Ala Leu Lys Leu Glu Asn Gly Gly His Tyr Thr Cys
165 170 175

Asp Phe Lys Thr Thr Tyr Lys Ser Lys Lys Gly Leu Lys Val Pro Pro
180 185 190

Tyr His Phe Val Asp His Lys Leu Asp Leu Leu Ser His Asn Thr Asp
195 200 205

Gly Ala Thr Phe Glu Glu Phe Glu Gln Arg Glu Ile Ala His Ala His
210 215 220

Leu Ser Asn Leu Pro Val Ala His His His His His His

-continued

225	230	235
<div><210> SEQ ID NO 36 <211> LENGTH: 224 <212> TYPE: PRT <213> ORGANISM: Porites porites <400> SEQUENCE: 36</div>		
Met Ala Leu Ser	Asn Gln Val Thr	Met Lys Tyr His Met Asp Gly Arg
1	5	10 15
Phe Glu Asp Lys	Glu Phe Thr Ile	Glu Gly Glu Gly Thr Gly Lys Pro
	20	25 30
Tyr Glu Gly Lys	Gln Thr Val Thr	Leu Trp Val Thr Lys Gly Ala Pro
	35	40 45
Leu Pro Phe Ser	Phe Asp Ile Leu Ser	Ala Val Phe Leu Tyr Gly Asn
	50	55 60
Arg Ala Phe Thr	Asp Tyr Pro Lys Gly	Ile Val Asp Tyr Phe Lys Pro
65	70	75 80
Ser Phe Pro Glu	Gly Tyr Ser Phe Glu	Arg Thr Leu Glu Phe Glu Asp
	85	90 95
Gly Gly Tyr Cys	Thr Ala Ser Ala	Asp Ile Ser Leu Asp Ser Ala Ser
	100	105 110
Asn Cys Phe Ile	His Lys Ser Ser	Phe Lys Gly Val Lys Phe Pro Asp
	115	120 125
Asn Gly Pro Val	Lys Gln Lys Lys	Thr Thr Asn Trp Glu Pro Ser Ile
	130	135 140
Glu Lys Met Thr	Val Arg Asp Gly	Ile Leu Lys Gly Asp Val Thr Met
145	150	155 160
Phe Leu Ser Leu	Thr Asp Gly Gly	Asn His Arg Cys Gln Phe Ser Thr
	165	170 175
Leu Tyr Lys Ala	Lys Lys Ala Val	Lys Leu Pro Thr Glu Ser His Tyr
	180	185 190
Val Glu His Arg	Leu Val Arg Thr	Asp Leu Pro Asn Gly Lys Val Gln
	195	200 205
Leu Glu Glu His	Ala Ala Ala Arg	Leu Asn Thr Val His His His Gln
	210	215 220
<div><210> SEQ ID NO 37 <211> LENGTH: 230 <212> TYPE: PRT <213> ORGANISM: Stylocoeniella sp <400> SEQUENCE: 37</div>		
Met Ala Leu Thr	Lys Gln Cys Ile	Ala Asn Glu Met Thr Met Thr Phe
1	5	10 15
His Met Asp Gly	Cys Val Asn Gly	His Tyr Phe Thr Ile Glu Gly Glu
	20	25 30
Gly Ser Gly Arg	Pro Tyr Glu Gly	Lys Gln Met Ser Lys Phe Lys Val
	35	40 45
Thr Lys Gly Gly	Pro Leu Pro Phe	Ser Phe Asp Ile Leu Ser Ser Ala
	50	55 60
Phe Lys Tyr Gly	Asn Arg Cys Phe	Thr Ala Tyr Pro Ala Gly Met His
65	70	75 80
Asp Tyr Phe Lys	Gln Ala Phe Pro	Glu Gly Met Ser Tyr Glu Arg Thr
	85	90 95
Phe Thr Phe Glu	Asp Gly Gly Val	Ala Thr Ala Ser Gly Asp Ile Ser

-continued

100					105					110						
Leu	Lys	Gly	Asn	Cys	Phe	Val	His	Lys	Ser	Met	Phe	His	Gly	Val	Asn	
115					120					125						
Phe	Pro	Ala	Asp	Gly	Pro	Val	Met	Lys	Lys	Lys	Thr	Thr	Gly	Trp	Asp	
130					135					140						
Pro	Ser	Phe	Glu	Lys	Met	Thr	Val	Cys	Asn	Gly	Ile	Leu	Lys	Gly	Asp	
145					150					155					160	
Val	Thr	Met	Phe	Leu	Met	Leu	Glu	Asp	Gly	Lys	Asn	Tyr	Lys	Cys	Gln	
165					170					175						
Phe	His	Thr	Ser	Tyr	Lys	Thr	Lys	Lys	Pro	Val	Thr	Leu	Pro	Ser	Asn	
180					185					190						
His	Val	Val	Glu	His	Arg	Ile	Val	Arg	Thr	Asn	Leu	Asp	Lys	Ala	Gly	
195					200					205						
Asn	His	Val	Gln	Leu	Asp	Glu	His	Ala	Val	Ala	His	Val	Asn	Pro	Leu	
210					215					220						
His	His	His	His	His	His											
225					230											

<210> SEQ ID NO 38

<211> LENGTH: 227

<212> TYPE: PRT

<213> ORGANISM: Fungia cf danai

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (2)..(2)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (5)..(5)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

<220> FEATURE:

<221> NAME/KEY: misc_feature

<222> LOCATION: (13)..(13)

<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid

Met	Xaa	Leu	Ser	Xaa	Gln	Xaa	Ile	Gly	Lys	Asp	Met	Xaa	Ile	Asn	Tyr
1				5					10					15	
Phe	Met	Asp	Gly	Ser	Val	Asn	Gly	His	Glu	Phe	Thr	Val	Lys	Gly	Glu
			20					25					30		
Gly	Ile	Gly	Lys	Pro	Tyr	Glu	Gly	His	His	Glu	Met	Thr	Leu	Arg	Val
			35					40					45		
Thr	Met	Ala	Lys	Gly	Gly	Pro	Leu	Pro	Phe	Ser	Phe	Asp	Leu	Leu	Ser
			50				55					60			
His	Thr	Phe	Cys	Tyr	Gly	Asn	Arg	Pro	Phe	Thr	Lys	Tyr	Pro	Glu	Glu
65					70					75					80
Ile	Pro	Asp	Tyr	Phe	Lys	Gln	Ala	Phe	Pro	Glu	Gly	Leu	Ser	Trp	Glu
				85					90					95	
Arg	Ser	Leu	Gln	Phe	Glu	Asp	Gly	Gly	Phe	Ala	Ala	Val	Asn	Ala	Asn
			100					105					110		
Ile	Ser	Leu	Lys	Gly	Asp	Cys	Phe	Glu	His	Asn	Ser	Lys	Phe	Val	Gly
			115					120					125		
Val	Asn	Phe	Pro	Ala	Glu	Gly	Pro	Val	Met	Gln	Asn	Lys	Ser	Leu	Asp
			130				135					140			
Trp	Glu	Pro	Ser	Thr	Glu	Lys	Ile	Thr	Val	Ser	Asp	Gly	Val	Leu	Lys
145					150					155					160

-continued

Gly Asp Val Pro Met Phe Leu Lys Leu Val Gly Gly Gly Asn His Lys
165 170 175
Cys Gln Phe Thr Thr Thr Tyr Lys Ala Ala Lys Lys Val Leu Asp Met
180 185 190
Pro Gln Ser His Phe Ile Phe His Arg Leu Val Arg Lys Thr Glu Gly
195 200 205
Asn Ile Thr Lys Leu Val Glu Asp Val Glu Ala His Asn His His His
210 215 220
His His His
225

<210> SEQ ID NO 39
<211> LENGTH: 237
<212> TYPE: PRT
<213> ORGANISM: Fungia cf danai

<400> SEQUENCE: 39

Met Ser Tyr Ser Lys Gln Gly Ile Val Gln Glu Met Lys Thr Lys Tyr
1 5 10 15
Arg Met Glu Gly Ser Val Asn Gly His Glu Phe Thr Ile Glu Gly Val
20 25 30
Gly Thr Gly Tyr Pro Tyr Glu Gly Lys Gln Met Ser Glu Leu Val Ile
35 40 45
Ile Lys Pro Lys Gly Lys Pro Leu Pro Phe Ser Phe Asp Ile Leu Ser
50 55 60
Ser Val Phe Gln Tyr Gly Asn Arg Cys Phe Thr Lys Tyr Pro Ala Asp
65 70 75 80
Met Pro Asp Tyr Phe Lys Gln Ala Phe Pro Asp Gly Met Ser Tyr Glu
85 90 95
Arg Ser Phe Leu Phe Glu Asp Gly Ala Val Ala Thr Ala Ser Trp Asn
100 105 110
Ile Arg Leu Glu Gly Asn Cys Phe Ile His Asn Ser Ile Phe His Gly
115 120 125
Val Asn Phe Pro Ala Asp Gly Pro Val Met Lys Lys Lys Thr Ile Gly
130 135 140
Trp Asp Lys Ser Phe Glu Lys Met Thr Val Ser Lys Glu Val Leu Thr
145 150 155 160
Gly Asp Val Thr Met Phe Leu Met Leu Glu Gly Gly Gly Tyr His Arg
165 170 175
Cys Gln Phe His Ser Thr Tyr Lys Thr Glu Lys Pro Val Glu Leu Pro
180 185 190
Pro Asn His Val Val Glu His Gln Ile Val Arg Thr Asp Leu Gly Gln
195 200 205
Ser Ala Lys Gly Phe Thr Val Lys Leu Glu Ala His Ala Ala Ala His
210 215 220
Val Asn Pro Leu Lys Val Gln His His His His His His
225 230 235

<210> SEQ ID NO 40
<211> LENGTH: 227
<212> TYPE: PRT
<213> ORGANISM: Goniopora djiboutiensis

<400> SEQUENCE: 40

Met Ser Val Ile Ala Lys Gln Met Thr Tyr Lys Val Tyr Met Ser Gly
1 5 10 15

-continued

Thr	Val	Asn	Gly	His	Tyr	Phe	Glu	Val	Gln	Gly	Asp	Gly	Lys	Gly	Lys	
		20						25					30			
Pro	Tyr	Glu	Gly	Glu	Gln	Thr	Val	Lys	Leu	Thr	Val	Thr	Lys	Gly	Gly	
		35					40					45				
Pro	Leu	Pro	Phe	Ala	Trp	Asp	Ile	Leu	Ser	Pro	Gln	Ala	Gln	Tyr	Gly	
		50				55					60					
Ser	Ile	Pro	Phe	Thr	Lys	Tyr	Pro	Glu	Asp	Ile	Pro	Asp	Tyr	Val	Lys	
65					70					75					80	
Gln	Ser	Phe	Pro	Glu	Gly	Tyr	Thr	Trp	Glu	Arg	Ile	Met	Asn	Phe	Glu	
				85					90					95		
Asp	Gly	Ala	Val	Cys	Thr	Val	Ser	Asn	Asp	Ser	Ser	Ile	Gln	Gly	Asn	
			100					105					110			
Cys	Phe	Ile	Tyr	Asn	Val	Lys	Phe	Ser	Gly	Leu	Asn	Phe	Pro	Pro	Ser	
		115					120					125				
Gly	Pro	Val	Met	Gln	Lys	Lys	Thr	Gln	Gly	Trp	Glu	Pro	Asn	Thr	Glu	
	130					135					140					
Arg	Leu	Leu	Ala	Arg	Asp	Gly	Met	Leu	Ile	Gly	Asn	Asn	Phe	Met	Ala	
145					150					155					160	
Leu	Lys	Leu	Glu	Gly	Gly	Gly	His	Tyr	Leu	Cys	Glu	Phe	Lys	Ser	Thr	
				165					170					175		
Tyr	Lys	Ala	Lys	Lys	Pro	Val	Lys	Met	Pro	Gly	Tyr	His	Phe	Val	Asp	
			180					185					190			
Arg	Lys	Leu	Asp	Val	Thr	Asn	His	Asn	Gln	Asp	Tyr	Thr	Ser	Val	Glu	
		195					200					205				
Gln	Cys	Glu	Ile	Ser	Ile	Ala	Arg	Lys	Pro	Val	Val	Ala	His	His	His	
	210					215					220					
His	His	His														
225																
<210> SEQ ID NO 41																
<211> LENGTH: 227																
<212> TYPE: PRT																
<213> ORGANISM: Montipora efflorescens																
<400> SEQUENCE: 41																
Met	Ser	Val	Ile	Ala	Lys	Gln	Met	Thr	Tyr	Lys	Val	Tyr	Met	Ser	Gly	
1				5					10					15		
Thr	Val	Asn	Gly	His	Tyr	Phe	Glu	Val	Glu	Gly	Asp	Gly	Lys	Gly	Lys	
		20						25					30			
Pro	Tyr	Glu	Gly	Glu	Gln	Thr	Val	Lys	Leu	Thr	Val	Thr	Lys	Gly	Gly	
		35					40					45				
Pro	Leu	Pro	Phe	Ala	Trp	Asp	Ile	Leu	Ser	Pro	Leu	Ser	Gln	Tyr	Gly	
		50				55					60					
Ser	Ile	Pro	Phe	Thr	Lys	Tyr	Pro	Glu	Asp	Ile	Pro	Asp	Tyr	Val	Lys	
65					70					75					80	
Gln	Ser	Phe	Pro	Glu	Gly	Tyr	Thr	Trp	Glu	Arg	Ile	Met	Asn	Phe	Glu	
				85					90					95		
Asp	Gly	Ala	Val	Cys	Thr	Val	Ser	Asn	Asp	Ser	Ser	Ile	Gln	Gly	Asn	
			100					105					110			
Cys	Phe	Ile	Tyr	Asn	Val	Lys	Ile	Ser	Gly	Val	Asn	Phe	Pro	Pro	Asn	
		115					120					125				
Gly	Pro	Val	Met	Gln	Lys	Lys	Thr	Gln	Gly	Trp	Glu	Pro	Asn	Thr	Glu	
	130					135					140					
Arg	Leu	Phe	Ala	Arg	Asp	Gly	Met	Leu	Ile	Gly	Asn	Asn	Phe	Met	Ala	

-continued

145				150						155					160
Leu	Lys	Leu	Glu	Gly	Gly	Gly	Tyr	Tyr	Leu	Cys	Glu	Phe	Lys	Ser	Thr
				165					170					175	
Tyr	Lys	Ala	Lys	Lys	Pro	Val	Arg	Met	Pro	Gly	Tyr	His	Tyr	Val	Asp
			180					185					190		
Arg	Lys	Leu	Asp	Val	Thr	Ser	His	Asn	Lys	Asp	Tyr	Thr	Phe	Val	Glu
		195					200					205			
Gln	Cys	Glu	Ile	Ser	Ile	Ala	Arg	His	Ser	Leu	Leu	Gly	His	His	His
	210					215					220				
His	His	His													
225															
<210> SEQ ID NO 42															
<211> LENGTH: 227															
<212> TYPE: PRT															
<213> ORGANISM: Stylocoeniella															
<400> SEQUENCE: 42															
Met	Ser	Val	Ile	Ala	Lys	Gln	Met	Thr	Tyr	Lys	Val	Tyr	Met	Ser	Gly
1				5					10					15	
Thr	Val	Asn	Gly	His	Tyr	Phe	Glu	Val	Gln	Gly	Asp	Gly	Lys	Gly	Lys
			20					25					30		
Pro	Tyr	Glu	Gly	Glu	Gln	Thr	Val	Arg	Leu	Thr	Val	Thr	Lys	Gly	Gly
		35					40					45			
Pro	Leu	Pro	Phe	Ala	Trp	Asp	Ile	Leu	Ser	Pro	Leu	Ser	Gln	Tyr	Gly
	50					55					60				
Ser	Ile	Pro	Phe	Thr	Lys	Tyr	Pro	Glu	Asp	Ile	Pro	Asp	Tyr	Val	Lys
65					70				75						80
Gln	Ser	Phe	Pro	Glu	Gly	Tyr	Thr	Trp	Glu	Arg	Ile	Met	Asn	Phe	Glu
				85					90					95	
Asp	Gly	Ala	Val	Cys	Thr	Val	Ser	Asn	Asp	Ser	Ser	Ile	Gln	Gly	Asn
			100					105					110		
Cys	Phe	Ile	Tyr	Asn	Val	Lys	Ile	Ser	Gly	Leu	Asn	Phe	Pro	Pro	Asn
		115					120					125			
Gly	Pro	Val	Met	Gln	Lys	Lys	Thr	Gln	Gly	Trp	Glu	Pro	Asn	Thr	Glu
	130					135					140				
Arg	Leu	Phe	Ala	Arg	Asp	Gly	Met	Leu	Ile	Gly	Asn	Asn	Phe	Met	Ala
145					150					155					160
Leu	Lys	Leu	Glu	Gly	Gly	Gly	His	Tyr	Leu	Cys	Glu	Phe	Lys	Ser	Thr
				165					170					175	
Tyr	Lys	Ala	Lys	Lys	Pro	Val	Arg	Met	Pro	Gly	Tyr	His	Tyr	Val	Asp
			180					185					190		
Arg	Lys	Leu	Asp	Val	Thr	Ser	His	Asn	Arg	Asp	Tyr	Thr	Ser	Val	Glu
		195					200					205			
Gln	Cys	Glu	Ile	Ser	Ile	Ala	Arg	His	Ser	Leu	Leu	Gly	His	His	His
	210					215					220				
His	His	His													
225															
<210> SEQ ID NO 43															
<211> LENGTH: 238															
<212> TYPE: PRT															
<213> ORGANISM: Montipora efflorescens															
<400> SEQUENCE: 43															
Met	Ala	Leu	Ser	Lys	Gln	Ser	Leu	Pro	Ser	Asp	Met	Lys	Leu	Ile	Tyr

-continued

1	5	10	15
His Met Asp Gly	Asn Val Asn Gly	His Ser Phe Val	Ile Lys Gly Glu
20	25	30	
Gly Glu Gly Lys	Pro Tyr Glu Gly	Thr His Thr Ile	Lys Leu Gln Val
35	40	45	
Val Glu Gly Ser	Pro Leu Pro Phe	Ser Ala Asp Ile	Leu Ser Thr Val
50	55	60	
Phe Gln Tyr Gly	Asn Arg Cys Phe	Thr Lys Tyr Pro	Pro Asn Ile Val
65	70	75	80
Asp Tyr Phe Lys	Asn Ser Cys Ser	Gly Gly Gly Tyr	Lys Phe Gly Arg
85	90	95	
Ser Phe Leu Tyr	Glu Asp Gly Ala	Val Cys Thr Ala	Ser Gly Asp Ile
100	105	110	
Thr Leu Ser Ala	Asp Lys Lys Ser	Phe Glu His Lys	Ser Lys Phe Leu
115	120	125	
Gly Val Asn Phe	Pro Ala Asp Gly	Pro Val Met Lys	Lys Glu Thr Thr
130	135	140	
Asn Trp Glu Pro	Ser Cys Glu Lys	Met Thr Pro Asn	Gly Met Thr Leu
145	150	155	160
Ile Gly Asp Val	Thr Gly Phe Leu	Leu Lys Glu Asp	Gly Lys Arg Tyr
165	170	175	
Lys Cys Gln Phe	His Thr Phe His	Asp Ala Lys Asp	Lys Ser Lys Lys
180	185	190	
Met Pro Met Pro	Asp Phe His Phe	Val Gln His Lys	Ile Glu Arg Lys
195	200	205	
Asp Leu Pro Gly	Ser Met Gln Thr	Trp Arg Leu Thr	Glu His Ala Ala
210	215	220	
Ala Cys Lys Thr	Cys Phe Thr Glu	His His His His	His His His His
225	230	235	
<210> SEQ ID NO 44			
<211> LENGTH: 240			
<212> TYPE: PRT			
<213> ORGANISM: Montipora efflorescens			
<400> SEQUENCE: 44			
Met Ala Leu Ser	Lys Asn Gly Leu	Thr Lys Asn Met	Thr Thr Lys Tyr
1	5	10	15
Arg Met Glu Gly	Cys Val Asp Gly	His Lys Phe Val	Ile Thr Gly Asp
20	25	30	
Gly Ile Gly Asp	Pro Phe Glu Gly	Lys Gln Thr Ser	Ile Asp Leu Cys
35	40	45	
Val Val Glu Gly	Gly Pro Leu Pro	Phe Ser Glu Asp	Ile Leu Ser Ala
50	55	60	
Val Phe Asp Tyr	Gly Asn Arg Val	Phe Thr Lys Tyr	Pro Gln Asp Leu
65	70	75	80
Val Asp Tyr Phe	Lys Asn Ser Cys	Pro Ala Gly Tyr	Thr Trp Gln Arg
85	90	95	
Ser Phe Leu Phe	Glu Asp Gly Ala	Val Cys Thr Ala	Ser Ala Asp Ile
100	105	110	
Thr Val Ser Val	Glu Glu Asn Cys	Phe Tyr His Glu	Ser Lys Phe His
115	120	125	
Gly Val Asn Phe	Pro Ala Asp Gly	Pro Val Met Lys	Lys Met Thr Thr
130	135	140	

-continued

Asn	Trp	Glu	Pro	Ser	Cys	Glu	Lys	Ile	Thr	Pro	Ile	Pro	Asn	Glu	Gly
145					150					155					160
Ile	Leu	Lys	Gly	Asp	Val	Thr	Met	Phe	Leu	Leu	Leu	Lys	Asp	Gly	Gly
				165					170					175	
Arg	Tyr	Arg	Cys	Gln	Phe	Asp	Thr	Val	Tyr	Lys	Ala	Lys	Ser	Asp	Pro
			180					185					190		
Lys	Thr	Ile	Met	Met	Pro	Asp	Trp	His	Phe	Ile	Gln	His	Lys	Leu	Asn
		195					200					205			
Arg	Glu	Asp	Arg	Ser	Asp	Ala	Lys	His	Gln	Lys	Trp	Arg	Leu	Val	Glu
	210					215					220				
Asn	Ala	Ile	Ala	Tyr	Arg	Ser	Thr	Leu	Ser	His	His	His	His	His	His
225					230					235					240

We claim:

1. An isolated protein comprising an amino sequence set forth as SEQ ID NO:35 and a variant of said sequence, wherein the amino acid sequence of the variant is at least 95% identical to SEQ ID NO:35 and wherein said variant

20 has emission and excitation maxima that are within +10 nm of the emission and excitation maxima for SEQ ID NO:35.

2. The protein, according to claim 1, wherein said protein has an amino acid sequence set forth as SEQ ID NO:35.

* * * * *