



US007241880B2

(12) **United States Patent**
Adler et al.

(10) **Patent No.:** **US 7,241,880 B2**
(45) **Date of Patent:** **Jul. 10, 2007**

(54) **T1R TASTE RECEPTORS AND GENES
ENCODING SAME**

(75) Inventors: **Jon Elliot Adler**, San Diego, CA (US);
Xiaodong Li, San Diego, CA (US);
Lena Staszewski, San Diego, CA (US);
Shawn O'Connell, Encinitas, CA (US);
Sergey Zozulya, San Diego, CA (US)

(73) Assignee: **Senomyx, Inc.**, San Diego, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 365 days.

(21) Appl. No.: **10/035,045**

(22) Filed: **Jan. 3, 2002**

(65) **Prior Publication Data**

US 2003/0054448 A1 Mar. 20, 2003

Related U.S. Application Data

(60) Provisional application No. 60/284,547, filed on Apr.
19, 2001, provisional application No. 60/259,227,
filed on Jan. 3, 2001.

(51) **Int. Cl.**

C07H 21/04 (2006.01)
C12N 15/63 (2006.01)
C12N 15/12 (2006.01)
C12N 1/15 (2006.01)
G01N 33/567 (2006.01)

(52) **U.S. Cl.** **536/23.5**; 435/6; 435/7.21;
435/69.1; 435/252.3; 435/325; 436/501; 530/300;
530/350; 514/2

(58) **Field of Classification Search** 536/23.5;
435/6, 7.21, 69.5, 252.3, 320.1; 530/300,
530/350, 388.22; 436/501
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

2003/0036089 A1* 2/2003 Wei et al.

FOREIGN PATENT DOCUMENTS

WO WO 95/08627 3/1995
WO WO 00/06593 A 2/2000
WO WO 2002/064631 A3 8/2002

OTHER PUBLICATIONS

Hoon et al., Cell 96(541-551), Feb. 19, 1999.*
Krautwurst et al. Cell 95(917-926)1998.*

Alexander et al., Proc. Natl. Acad. Sci. 89(3352-3356)1992.*
Bowie et al., 1990, Science 247:1306-1310.*

Mark A. Hoon, et al., "Putative Mammalian Taste Receptors: A
Class of Taste-Specific GPCRs with Distinct Topographic Selectiv-
ity", Cell, vol. 96, pp. 541-551, Feb. 19, 1999—XP-000922524.
Xiaodong Li, et al. "Human Receptors for Sweet and Umami Taste",
Proceedings of the National Academy of Sciences of USA, National
Academy of Science, vol. 99, No. 7, Apr. 2, 2002, pp. 4692-
4696—XP002254830.

R.H. Waterston, "Direct Submission", Genome Sequencing Center,
Apr. 21, 2000—XP-002262036.

Elliot Adler, et al. "A Novel Family of Mammalian Taste Recep-
tors", Cell, vol. 100, pp. 693-702, Mar. 17, 2000—XP-000982304.
Montmayeur, J-P, et al., "A Candidate Taste Receptor Gene Near a
Sweet Taste Locus," Nature Neuroscience, May 2001, pp. 492-498,
vol. 4, No. 5, Nature America, Inc., US.

Kitagawa M., et al., "Molecular Genetic Identification of a Candi-
date Receptor Gene for Sweet Taste," Biochemical and Biophysical
Research Communications, Apr. 27, 2001, pp. 236-242, vol. 283,
No. 1, Academic Press, San Diego, California, US.

Max Marianna, et al., "Tas1r3, Encoding a New Candidate Taste
Receptor, is Allelic to the Sweet Responsiveness Locus Sac,"
Nature Genetics, May 2001, pp. 58-63, vol. 28, No. 1, Nature
America, New York, US.

Sainz E., et al., "Identification of a Novel Member of the T1R
Family of Putative Taste Receptors," Journal of Neurochemistry,
May 2001, pp. 896-903, vol. 77, No. 3, New York, NY, US.

Johnson C., et al., "The Effect of the Sweetness Inhibitor 2(4-
methoxyphenoxy) Propanoic Acid (Sodium Salt) (Na-PMP) on the
Taste of Bitter-sweet Stimuli," Chemical Sense, Aug. 1994, pp.
349-358, vol. 19, No. 4, Oxford, UK.

* cited by examiner

Primary Examiner—Janet L. Andres

Assistant Examiner—Michael Brannock

(74) *Attorney, Agent, or Firm*—Hunton & Williams LLP

(57) **ABSTRACT**

Newly identified mammalian taste-cell-specific G protein-
coupled receptors, and the genes and cDNA encoding said
receptors are described. Specifically, T1R G protein-coupled
receptors active in taste signaling, and the genes and cDNA
encoding the same, are described, along with methods for
isolating such genes and for isolating and expressing such
receptors. Methods for representing taste perception of a
particular taste stimulus in a mammal are also described, as
are methods for generating novel molecules or combinations
of molecules that elicit a predetermined taste perception in
a mammal, and methods for simulating one or more tastes.
Further, methods for stimulating or blocking taste perception
in a mammal are also disclosed.

29 Claims, 5 Drawing Sheets






Figure 1




Figure 2





Figure 3

A.



H2-IT-Myc



H2-PDZP-IT-Myc

B.




Figure 4




Figure 5




Figure 6

T1R TASTE RECEPTORS AND GENES ENCODING SAME

RELATED APPLICATIONS

This application claims benefit of priority to U.S. Ser. No. 60/259,227 filed on Jan. 3, 2001 and U.S. Ser. No. 60/284,547, filed on Apr. 19, 2001, both of which are incorporated by reference in their entirety herein.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The invention relates to newly identified mammalian chemosensory G protein-coupled receptors, to family of such receptors, and to the genes and cDNA encoding said receptors. More particularly, the invention relates to newly identified mammalian chemosensory G protein-coupled receptors active in taste signaling, to a family of such receptors, to the genes and cDNA encoding said receptors, and to methods of using such receptors, genes, and cDNA in the analysis and discovery of taste modulators. The invention provides in particular a DNA sequence encoding a novel human taste receptor identified infra as T1R2 and the corresponding receptor polypeptide.

2. Description of the Related Art

The taste system provides sensory information about the chemical composition of the external world. Mammals are believed to have at least five basic taste modalities: sweet, bitter, sour, salty, and umami. See, e.g., Kawamura et al., *Introduction to Umami: A Basic Taste* (1987); Kinnamon et al., *Ann. Rev. Physiol.*, 54:715–31 (1992); Lindemann, *Physiol. Rev.*, 76:718–66 (1996); Stewart et al., *Am. J. Physiol.*, 272:1–26(1997). Each taste modality is thought to be mediated by a distinct protein receptor or receptors that are expressed in taste receptor cells on the surface of the tongue (Lindemann, *Physiol. Rev.* 76:718–716 (1996)). The taste receptors that recognize bitter, sweet, and umami taste stimuli belong to the G-protein-coupled receptor (GPCR) superfamily (Hoon et al., *Cell* 96:451 (1999); Adler et al., *Cell* 100:693 (2000)). (Other taste modalities are believed to be mediated by ion channels.)

G protein-coupled receptors mediate many other physiological functions, such as endocrine function, exocrine function, heart rate, lipolysis, and carbohydrate metabolism. The biochemical analysis and molecular cloning of a number of such receptors has revealed many basic principles regarding the function of these receptors. For example, U.S. Pat. No. 5,691,188 describes how upon a ligand binding to a GPCR, the receptor undergoes a conformational change leading to activation of a heterotrimeric G protein by promoting the displacement of bound GDP by GTP on the surface of the Ga subunit and subsequent dissociation of the Ga subunit from the Gb and Gg subunits. The free Ga subunits and Gbg complexes activate downstream elements of a variety of signal transduction pathways.

Complete or partial sequences of numerous human and other eukaryotic chemosensory receptors are currently known. See, e.g., Pilpel, Y. and Lancet, D., *Protein Science*, 8:969–977 (1999); Mombaerts, P., *Annu. Rev. Neurosci.*, 22:487–50 (1999). See also, EP0867508A2, U.S. Pat. No. 5,874,243, WO 92/17585, WO 95/18140, WO 97/17444, WO 99/67282. Because of the complexity of ligand-receptor interactions, and more particularly taste stimulus-receptor interactions, information about ligand-receptor recognition is lacking.

The identification and characterization of the GPCRs that function as sweet and umami taste receptors could allow for new methods of discovery of new taste stimuli. For example, the availability of receptors could permit the screening for receptor modulators. Such compounds would modulate taste and could be useful in the food industry to improve the taste of a variety of consumer products; e.g., improving the palatability of low-calorie beverages through the development of new artificial sweeteners.

In part, the present invention addresses the need for better understanding of the interactions between chemosensory receptors and chemical stimuli. The present invention also provides, among other things, novel chemosensory receptors, and methods for utilizing such receptors, and the genes a cDNAs encoding such receptors, to identify molecules that can be used to modulate chemosensory transduction, such as taste sensation.

SUMMARY OF THE INVENTION

The invention relates to a new family of G protein-coupled receptors, and to the genes and cDNAs encoding said receptors. The receptors are thought to be primarily involved in sweet taste transduction, but can be involved in transducing signals from other taste modalities as well.

The invention provides methods for representing the perception of taste and/or for predicting the perception of taste in a mammal, including in a human. Preferably, such methods may be performed by using the receptors and genes encoding said receptors disclosed herein.

Toward that end, it is an object of the invention to provide a new family of mammalian G protein-coupled receptors, herein referred to as T1Rs, active in taste perception. It is another object of the invention to provide fragments and variants of such T1Rs that retain taste stimulus-binding activity.

It is yet another object of the invention to provide nucleic acid sequences or molecules that encode such T1Rs, fragments, or variants thereof.

It is still another object of the invention to provide expression vectors which include nucleic acid sequences that encode such T1Rs, or fragments or variants thereof, which are operably linked to at least one regulatory sequence such as a promoter, enhancer, or other sequence involved in positive or negative gene transcription and/or translation.

It is still another object of the invention to provide human or non-human cells that functionally express at least one of such T1Rs, or fragments or variants thereof.

It is still another object of the invention to provide T1R fusion proteins or polypeptides which include at least a fragment of at least one of such T1Rs.

It is another object of the invention to provide an isolated nucleic acid molecule encoding a T1R polypeptide comprising a nucleic acid sequence that is at least 50%, preferably 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to a nucleic acid sequence selected from the group consisting of: SEQ ID NOS: 1, 2, 3, 9, 11, 13, 15, 16, 20, and conservatively modified variants thereof.

It is a further object of the invention to provide an isolated nucleic acid molecule comprising a nucleic acid sequence that encodes a polypeptide having an amino acid sequence at least 35 to 50%, and preferably 60%, 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to an amino acid sequence selected from the group consisting of: SEQ ID NOS: 4, 10, 12, 14, 17, 21, and conservatively modified variants thereof, wherein the fragment is at least 20, pref-

erably 40, 60, 80, 100, 150, 200, or 250 amino acids in length. Optionally, the fragment can be an antigenic fragment which binds to an anti-T1R antibody.

It is still a further object of the invention to provide an isolated polypeptide comprising a variant of said fragment, wherein there is a variation in at most 10, preferably 5, 4, 3, 2, or 1 amino acid residues.

It is still another object of the invention to provide agonists or antagonists of such T1Rs, or fragments or variants thereof.

It is yet another object of the invention to provide methods for representing the perception of taste and/or for predicting the perception of taste in a mammal, including in a human. Preferably, such methods may be performed by using the T1Rs, or fragments or variants thereof, and genes encoding such T1Rs, or fragments or variants thereof, disclosed herein.

It is yet another object of the invention to provide novel molecules or combinations of molecules which elicit a predetermined taste perception in a mammal. Such molecules or compositions can be generated by determining a value of taste perception in a mammal for a known molecule or combinations of molecules; determining a value of taste perception in a mammal for one or more unknown molecules or combinations of molecules; comparing the value of taste perception in a mammal for one or more unknown compositions to the value of taste perception in a mammal for one or more known compositions; selecting a molecule or combination of molecules that elicits a predetermined taste perception in a mammal; and combining two or more unknown molecules or combinations of molecules to form a molecule or combination of molecules that elicits a predetermined taste perception in a mammal. The combining step yields a single molecule or a combination of molecules that elicits a predetermined taste perception in a mammal.

It is still a further object of the invention to provide a method of screening one or more compounds for the presence of a taste detectable by a mammal, comprising: a step of contacting said one or more compounds with at least one of the disclosed T1Rs, fragments or variants thereof, preferably wherein the mammal is a human.

It is another object of the invention to provided a method for simulating a taste, comprising the steps of: for each of a plurality of T1Rs, or fragments of variants thereof disclosed herein, preferably human T1Rs, ascertaining the extent to which the T1R interacts with the taste stimulus; and combining a plurality of compounds, each having a previously ascertained interaction with one or more of the T1Rs, in amounts that together provide a receptor-stimulation profile that mimics the profile for the taste. Interaction of a taste stimulus with a T1R can be determined using any of the binding or reporter assays described herein. The plurality of compounds may then be combined to form a mixture. If desired, one or more of the plurality of the compounds can be combined covalently. The combined compounds substantially stimulate at least 50%, 60%, 70%, 75%, 80% or 90% or all of the receptors that are substantially stimulated by the taste stimulus.

In yet another aspect of the invention, a method is provided wherein a plurality of standard compounds are tested against a plurality of T1Rs, or fragments or variants thereof, to ascertain the extent to which the T1Rs each interact with each standard compound, thereby generating a receptor stimulation profile for each standard compound. These receptor stimulation profiles may then be stored in a relational database on a data storage medium. The method may further comprise providing a desired receptor-stimula-

tion profile for a taste; comparing the desired receptor stimulation profile to the relational database; and ascertaining one or more combinations of standard compounds that most closely match the desired receptor-stimulation profile.

The method may further comprise combining standard compounds in one or more of the ascertained combinations to simulate the taste.

It is a further object of the invention to provide a method for representing taste perception of a particular taste stimulus in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n T1Rs of said vertebrate, where n is greater than or equal to 2; and generating from said values a quantitative representation of taste perception. The T1Rs may be a taste receptor disclosed herein, or fragments or variants thereof, the representation may constitute a point or a volume in n -dimensional space, may constitute a graph or a spectrum, and may constitute a matrix of quantitative representations. Also, the providing step may comprise contacting a plurality of recombinantly-produced T1Rs, or fragments or variants thereof, with a test composition and quantitatively measuring the interaction of said composition with said receptors.

It is yet another object of the invention to provide a method for predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n T1Rs of said vertebrate, where n is greater than or equal to 2; for one or more molecules or combinations of molecules yielding known taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal, providing values X_1 to X_n representative of the quantitative stimulation of each of n T1Rs of said vertebrate, where n is greater than or equal to 2; for one or more molecules or combinations of molecules yielding unknown taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, and predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal by comparing the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal to the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal. The T1Rs used in this method may include a taste receptor, or fragment or variant thereof, disclosed herein.

DETAILED DESCRIPTION OF THE FIGURES

FIGS. 1a–1c present functional data for the human T1R2/T1R3 receptor. Intracellular calcium responses of HEK cells stably expressing $G\alpha_{15}$ that are transiently transfected with human T1R2, T1R3, and T1R2/T1R3 to various concentrations of sucrose are shown in FIG. 1a. Each panel corresponds to approximately 1000 confluent, transfected, and calcium-dye-loaded cells. Inhibition of T1R2/T1R3 activity by the sweet-taste inhibitor gurmarin is shown in FIG. 1b. T1R2/T1R3 dose responses to four sweeteners and correlated psychophysical detection thresholds (X-axis circles) are shown in FIG. 1c.

FIG. 2 presents functional data for the rat T1R2/T1R3 receptor. Human T1R2/T1R3 and rat T1R2/T1R3 (as well as mixed rat/human receptors) responses to 350 mM sucrose, 25 mM tryptophan, 15 mM aspartame, and 0.05% monellin are shown. Rat T1R2/T1R3 does not respond to aspartame or monellin, which are not palatable to rodents.

FIGS. 3a–3c present functional data for the human T1R2/T1R3 receptor. Intracellular calcium responses of HEK cells stably expressing G α 15 that are transiently transfected with human T1R1, T1R3, and T1R1/T1R3 to various concentrations of L-glutamate are shown in FIG. 3a. Potentiation of the T1R1/T1R3 response by IMP is shown in FIG. 3b. T1R1/T1R3 dose responses to L-glutamate and L-glutamate plus 0.2 mM IMP and correlated psychophysical detection thresholds (X-axis circles) are shown in FIG. 3c.

FIGS. 4a–4b present immunofluorescence and FACS experiments that demonstrate that fusing the PDZIP peptide (SEQ ID No: 1) to human T1R2 enhanced its expression on the surface of HEK cells.

FIG. 5 presents automated fluorescence imaging data for cell lines that stably express G α 15 and human T1R1/T1R3. L-glutamate dose responses were determined in the presence of 0.5 mM IMP.

FIG. 6 presents automated fluorescence imaging data for cell lines that stably express G α 15 and human T1R2/T1R3. Sucrose, D-tryptophan, saccharin, and aspartame dose responses are shown for on stable cell line.

DETAILED DESCRIPTION OF THE INVENTION

The invention thus provides isolated nucleic acid molecules encoding taste-cell-specific G protein-coupled receptors (“GPCR”), and the polypeptides they encode. These nucleic acid molecules and the polypeptides that they encode are members of the T1R family of taste-cell-specific GPCRs. Members of the T1R family of taste-cell-specific GPCRs are identified in Hoon et al., *Cell*, 96:541–551 (1999), WO 00/06592, and WO 00/06593, all of which are incorporated herein by reference in their entireties.

More particularly, the invention provides nucleic acids encoding a novel family of taste-cell-specific GPCRs. These nucleic acids and the receptors that they encode are referred to as members of the “T1R” family of taste-cell-specific GPCRs. In particular embodiments of the invention, the T1R family members include human T1R1, T1R2, and T1R3. As described supra, different T1R combinations likely mediate sweet and umami taste.

Further, it is believed that T1R family members may act in combination with other T1R family members, other taste-cell-specific GPCRs, or a combination thereof, to thereby effect chemosensory taste transduction. For instance, it is believed that T1R1 and T1R3 maybe coexpressed within the same taste receptor cell type, and the two receptors may physically interact to form a heterodimeric taste receptor. Alternatively, T1R1 and T1R3 may both independently bind to the same type of ligand, and their combined binding may result in a specific perceived taste sensation.

These nucleic acids provide valuable probes for the identification of taste cells, as the nucleic acids are specifically expressed in taste cells. For example, probes for T1R polypeptides and proteins can be used to identify taste cells present in foliate, circumvallate, and fungiform papillae, as well as taste cells present in the geschmackstreifen, oral cavity, gastrointestinal epithelium, and epiglottis. They may also serve as tools for the generation of taste topographic

maps that elucidate the relationship between the taste cells of the tongue and taste sensory neurons leading to taste centers in the brain. In particular, methods of detecting T1Rs can be used to identify taste cells sensitive to sweet taste stimuli or other specific modalities of taste stimuli. Furthermore, the nucleic acids and the proteins they encode can be used as probes to dissect taste-induced behaviors. Also, chromosome localization of the genes encoding human T1Rs can be used to identify diseases, mutations, and traits caused by and associated with T1R family members.

The nucleic acids encoding the T1R proteins and polypeptides of the invention can be isolated from a variety of sources, genetically engineered, amplified, synthesized, and/or expressed recombinantly according to the methods disclosed in WO 00/035374, which is herein incorporated by reference in its entirety.

The invention also provides methods of screening for modulators, e.g., activators, inhibitors, stimulators, enhancers, agonists, and antagonists, of these novel taste-cell-specific GPCRs. Such modulators of taste transduction are useful for pharmacological, chemical, and genetic modulation of taste signaling pathways. These methods of screening can be used to identify high affinity agonists and antagonists of taste cell activity. These modulatory compounds can then be used in the food and pharmaceutical industries to customize taste, e.g., to modulate the sweet tastes of foods or drugs.

Thus, the invention provides assays for detecting and characterizing taste modulation, wherein T1R family members act as direct or indirect reporter molecules of the effect of modulators on taste transduction. GPCRs can be used in assays to, e.g., measure changes in ligand binding, ion concentration, membrane potential, current flow, ion flux, transcription, signal transduction, receptor-ligand interactions, second messenger concentrations, in vitro, in vivo, and ex vivo. In one embodiment, members of the T1R family can be used as indirect reporters via attachment to a second reporter molecule such as green fluorescent protein (see, e.g., Mistili & Spector, *Nature Biotechnology*, 15:961–964 (1997)). In another embodiment, T1R family members may be recombinantly expressed in cells, and modulation of taste transduction via GPCR activity may be assayed by measuring changes in Ca²⁺ levels and other intracellular messages such as cAMP, cGMP, or IP3.

In certain embodiments, a domain of a T1R polypeptide, e.g., an extracellular, transmembrane, or intracellular domain, is fused to a heterologous polypeptide, thereby forming a chimeric polypeptide, e.g., a chimeric polypeptide with GPCR activity. Such chimeric polypeptides are useful, e.g., in assays to identify ligands, agonists, antagonists, or other modulators of a T1R polypeptide. In addition, such chimeric polypeptides are useful to create novel taste receptors with novel ligand binding specificity, modes of regulation, signal transduction pathways, or other such properties, or to create novel taste receptors with novel combinations of ligand binding specificity, modes of regulation, signal transduction pathways, etc.

In one embodiment, a T1R polypeptide is expressed in a eukaryotic cell as a chimeric receptor with a heterologous, chaperone sequence that facilitates plasma membrane trafficking, or maturation and targeting through the secretory pathway. The optional heterologous sequence may be a rhodopsin sequence, such as an N-terminal fragment of a rhodopsin. Such chimeric T1R receptors can be expressed in any eukaryotic cell, such as HEK-293 cells. Preferably, the cells comprise a G protein, e.g., G α 15 or G α 16 or another type of promiscuous G protein capable of pairing a wide range of

chemosensory GPCRs to an intracellular signaling pathway or to a signaling protein such as phospholipase C. Alternatively, the cells may express a chimeric or variant G protein that is selected based on its ability to couple with T1Rs to produce a functional T1R taste receptor. Examples of variant G proteins which are especially preferred include the G protein variants disclosed in U.S. Ser. No. 09/984,292, filed on Oct. 29, 2001, incorporated by reference herein in its entirety and the chimeric $G_{\alpha 15}$ variants disclosed in U.S. Provisional Application No. 60/243,770 also incorporated by reference in its entirety. These applications disclose G protein variants that have been shown to couple better with T1Rs than $G_{\alpha 15}$, a well known promiscuous G protein. Activation of such chimeric receptors in such cells can be detected using any standard method, such as by detecting changes in intracellular calcium by detecting FURA-2 dependent fluorescence in the cell. If preferred host cells do not express an appropriate G protein, they may be transfected with a gene encoding a promiscuous G protein such as those described in U.S. Application Ser. No. 60/243,770, which is herein incorporated by reference in its entirety.

Methods of assaying for modulators of taste transduction include in vitro ligand-binding assays using: T1R polypeptides, portions thereof, i.e., the extracellular domain, transmembrane region, or combinations thereof, or chimeric proteins comprising one or more domains of a T1R family member; oocyte or tissue culture cells expressing T1R polypeptides, fragments, or fusion proteins; phosphorylation and dephosphorylation of T1R family members; G protein binding to GPCRs; ligand-binding assays; voltage, membrane potential and conductance changes; ion flux assays; changes in intracellular second messengers such as cGMP, CAMP and inositol triphosphate; changes in intracellular calcium levels; and neurotransmitter release.

Further, the invention provides methods of detecting T1R nucleic acid and protein expression, allowing investigation of taste transduction regulation and specific identification of taste receptor cells. T1R family members also provide useful nucleic acid probes for paternity and forensic investigations. T1R genes are also useful as a nucleic acid probes for identifying taste receptor cells, such as foliate, fungiform, circumvallate, geschmackstreifen, and epiglottis taste receptor cells. T1R receptors can also be used to generate monoclonal and polyclonal antibodies useful for identifying taste receptor cells. Taste receptor cells can be identified using techniques such as reverse transcription and amplification of mRNA, isolation of total RNA or poly A+ RNA, northern blotting, dot blotting, in situ hybridization, RNase protection, S1 digestion, probing DNA microchip arrays, western blots, and the like.

Functionally, the T1R polypeptides comprise a family of related seven transmembrane G protein-coupled receptors, which are believed to be involved in taste transduction and may interact with a G protein to mediate taste signal transduction (see, e.g., Fong, *Cell Signal*, 8:217 (1996); Baldwin, *Curr. Opin. Cell Biol.*, 6:180 (1994)). Structurally, the nucleotide sequences of T1R family members may encode related polypeptides comprising an extracellular domain, seven transmembrane domains, and a cytoplasmic domain. Related T1R family genes from other species share at least about 50%, and optionally 60%, 70%, 80%, or 90%, nucleotide sequence identity over a region of at least about 50 nucleotides in length, optionally 100, 200, 500, or more nucleotides in length to SEQ ID NOS: 1, 2, 3, 9, 11, 13, 15, 16, 20, or conservatively modified variants thereof, or encode polypeptides sharing at least about 35 to 50%, and optionally 60%, 70%, 80%, or 90%, amino acid sequence

identity over an amino acid region at least about 25 amino acids in length, optionally 50 to 100 amino acids in length to SEQ ID NOS: 4, 10, 12, 14, 17, 21, or conservatively modified variants thereof.

Several consensus amino acid sequences or domains have also been identified that are characteristic of T1R family members. For example, T1R family members typically comprise a sequence having at least about 50%, optionally 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95–99%, or higher, identity to T1R consensus sequences 1 and 2 (SEQ ID NOS 18 and 19, respectively). These conserved domains thus can be used to identify members of the T1R family, by identity, specific hybridization or amplification, or specific binding by antibodies raised against a domain. Such T1R consensus sequences have the following amino acid sequences:

T1R Family Consensus Sequence 1: (SEQ ID NO: 18)
(TR)C(FL)(RQP)R(RT)(SPV)(VERKT)FL(AE)(WL)
(RHG)E

T1R Family Consensus Sequence 2: (SEQ ID NO: 19)
(LQP)(EGT)(NRC)YN(RE)A(RK)(CGF)(VLI)T(FL)
(AS)(ML)

These consensus sequences are inclusive of those found in the T1R polypeptides described herein, but T1R family members from other organisms may be expected to comprise consensus sequences having about 75% identity or more to the inclusive consensus sequences described specifically herein.

Specific regions of the T1R nucleotide and amino acid sequences may be used to identify polymorphic variants, interspecies homologs, and alleles of T1R family members. This identification can be made in vitro, e.g., under stringent hybridization conditions or PCR (e.g., using primers encoding the T1R consensus sequences identified above), or by using the sequence information in a computer system for comparison with other nucleotide sequences. Different alleles of T1R genes within a single species population will also be useful in determining whether differences in allelic sequences correlate to differences in taste perception between members of the population. Classical PCR-type amplification and cloning techniques are useful for isolating orthologs, for example, where degenerate primers are sufficient for detecting related genes across species, which typically have a higher level of relative identity than paralogous members of the T1R family within a single species.

For instance, degenerate primers SAP077 (SEQ. ID NO. 5) and SAP0079 (SEQ. ID NO. 6) can be used to amplify and clone T1R3 genes from different mammalian genomes. In contrast, genes within a single species that are related to T1R3 are best identified using sequence pattern recognition software to look for related sequences. Typically, identification of polymorphic variants and alleles of T1R family members can be made by comparing an amino acid sequence of about 25 amino acids or more, e.g., 50–100 amino acids. Amino acid identity of approximately at least 35 to 50%, and optionally 60%, 70%, 75%, 80%, 85%, 90%, 95–99%, or above typically demonstrates that a protein is a polymorphic variant, interspecies homolog, or allele of a T1R family member. Sequence comparison can be performed using any of the sequence comparison algorithms discussed below. Antibodies that bind specifically to T1R polypeptides or a conserved region thereof can also be used to identify alleles, interspecies homologs, and polymorphic variants.

Polymorphic variants, interspecies homologs, and alleles of T1R genes can be confirmed by examining taste-cell-specific expression of the putative T1R polypeptide. Typi-

cally, T1R polypeptides having an amino acid sequence disclosed herein can be used as a positive control in comparison to the putative T1R polypeptide to demonstrate the identification of a polymorphic variant or allele of the T1R family member. The polymorphic variants, alleles, and inter-species homologs are expected to retain the seven trans-membrane structure of a G protein-coupled receptor. For further detail, see WO 00/06592, which discloses related T1R family members, GPCR-B3s, the contents of which are herein incorporated by reference in a manner consistent with this disclosure. GPCR-B3 receptors are referred to herein as rT1R1 and mT1R1. Additionally, see WO 00/06593, which also discloses related T1R family members, GPCR-B4s, the contents of which are herein incorporated by reference in a manner consistent with this disclosure. GPCR-B4 receptors are referred to herein as rT1R2 and mT1R2.

Nucleotide and amino acid sequence information for T1R family members may also be used to construct models of taste-cell-specific polypeptides in a computer system. These models can be subsequently used to identify compounds that can activate or inhibit T1R receptor proteins. Such compounds that modulate the activity of T1R family members can then be used to investigate the role of T1R genes and receptors in taste transduction.

The present invention also provides assays, preferably high throughput assays, to identify molecules that interact with and/or modulate a T1R polypeptide. In numerous assays, a particular domain of a T1R family member is used, e.g., an extracellular, transmembrane, or intracellular domain or region. In numerous embodiments, an extracellular domain, transmembrane region or combination thereof may be bound to a solid substrate, and used, e.g., to isolate ligands, agonists, antagonists, or any other molecules that can bind to and/or modulate the activity of a T1R polypeptide.

In one aspect of the invention, a new human GPCR gene of the T1R family, termed hT1R3, is provided. The hT1R3 gene was identified from the human genome sequence database including the HTGS division of GenBank. The nucleotide and conceptually translated amino acid sequence for hT1R3 are provided in SEQ. ID NOS 1–4. The hT1R3 receptor was identified in the partially sequenced BAC genomic clone RP5-89003 (database accession number AL139287) by virtue of its sequence similarity to the candidate rat taste receptor rT1R1 (accession number AF127389). By reference, the pairwise identity between the predicted hT1R3 and rT1R1 protein sequences is approximately 34%. Sequence comparisons with additional members of the GPCR Family C (which includes the calcium-sensing receptors, putative V2R pheromone receptors, GABA-B receptors, fish taste receptors, and metabotropic glutamate receptors) indicate that hT1R3 is likely to belong to the Family C subgroup defined by T1R1 and a second rat candidate taste receptor (rT1R2, accession number AF127390).

The invention also provides the human ortholog, termed hT1R1, of a rat taste receptor, designated rT1R1. The gene products of rT1R1 and hT1R1 are approximately 74% identical. The mouse gene, mT1R1 has been reported, see Hoon et al., *Cell*, 96:541–551 (2000), and maps to a chromosomal interval homologous to the interval containing hT1R1. The nucleotide and conceptually-translated hT1R1 sequences are described herein as SEQ. ID NOS 15 and 16, respectively.

While not wishing to be bound to any particular theory, the T1R family of receptors is predicted to be involved in sweet taste transduction by virtue of the linkage of mT1R3

to the Sac locus, a locus on the distal end of chromosome four that influences sweet taste. Human T1R3 has also been reported to localize to 1p36.2–1p36.33, a region that displays conserved synteny with the mouse interval containing Sac and T1R1. However, T1R type receptors may mediate other taste modalities, such as bitter, umami, sour and salty.

Various conservative mutations and substitutions are envisioned to be within the scope of the invention. For instance, it would be within the level of skill in the art to perform amino acid substitutions using known protocols of recombinant gene technology including PCR, gene cloning, site-directed mutagenesis of cDNA, transfection of host cells, and in-vitro transcription. The variants could then be screened for taste-cell-specific GPCR functional activity.

A. Identification and Characterization of T1R Polypeptides

The amino acid sequences of the T1R proteins and polypeptides of the invention can be identified by putative translation of the coding nucleic acid sequences. These various amino acid sequences and the coding nucleic acid sequences may be compared to one another or to other sequences according to a number of methods.

For example, in sequence comparison, typically one sequence acts as a reference sequence, to which test sequences are compared. When using a sequence comparison algorithm, test and reference sequences are entered into a computer, subsequence coordinates are designated, if necessary, and sequence algorithm program parameters are designated. Default program parameters can be used, as described below for the BLASTN and BLASTP programs, or alternative parameters can be designated. The sequence comparison algorithm then calculates the percent sequence identities for the test sequences relative to the reference sequence, based on the program parameters.

A “comparison window,” as used herein, includes reference to a segment of any one of the number of contiguous positions selected from the group consisting of from 20 to 600, usually about 50 to about 200, more usually about 100 to about 150 in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Methods of alignment of sequences for comparison are well known in the art. Optimal alignment of sequences for comparison can be conducted, e.g., by the local homology algorithm of Smith & Waterman, *Adv. Appl. Math.* 2:482 (1981), by the homology alignment algorithm of Needleman & Wunsch, *J Mol. Biol.* 48:443 (1970), by the search for similarity method of Pearson & Lipman, *Proc. Natl. Acad. Sc. USA* 85:2444 (1988), by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by manual alignment and visual inspection (see, e.g., *Current Protocols in Molecular Biology* (Ausubel et al., eds. 1995 supplement)).

A preferred example of an algorithm that is suitable for determining percent sequence identity and sequence similarity are the BLAST and BLAST 2.0 algorithms, which are described in Altschul et al., *Nuc. Acids Res.* 25:3389–3402 (1997) and Altschul et al., *J Mol. Biol.* 215:403–410 (1990), respectively. Software for performing BLAST analyses is publicly available through the National Center for Biotechnology Information. This algorithm involves first identifying high scoring sequence pairs (HSPs) by identifying short words of length W in the query sequence, which either match or satisfy some positive-valued threshold score T when aligned with a word of the same length in a database

sequence. T is referred to as the neighborhood word score threshold (Altschul et al. Altschul et al., *Nuc. Acids Res.* 25:3389–3402 (1977) and Altschul et al, *J Mol Biol.* 215: 403–410 (1990)). These initial neighborhood word hits act as seeds for initiating searches to find longer HSPs containing them. The word hits are extended in both directions along each sequence for as far as the cumulative alignment score can be increased. Cumulative scores are calculated using, for nucleotide sequences, the parameters M (reward score for a pair of matching residues; always>0) and N (penalty score for mismatching residues; always>0). For amino acid sequences, a scoring matrix is used to calculate the cumulative score. Extension of the word hits in each direction are halted when: the cumulative alignment score falls off by the quantity X from its maximum achieved value; the cumulative score goes to zero or below, due to the accumulation of one or more negative-scoring residue alignments; or the end of either sequence is reached. The BLAST algorithm parameters W, T, and X determine the sensitivity and speed of the alignment. The BLASTN program (for nucleotide sequences) uses as defaults a wordlength (XV) of 11, an expectation (E) of 10, M=5, N=-4 and a comparison of both strands. For amino acid sequences, the BLASTP program uses as defaults a wordlength of 3, and expectation (E) of 10, and the BLOSUM62 scoring matrix (see Henikoff & Henikoff, *Proc. Natl. Acad. Sci. USA* 89:10915 (1989)) alignments (B) of 50, expectation (E) of 10, M=5, N=-4, and a comparison of both strands.

Another example of a useful algorithm is PILEUP. PILEUP creates a multiple sequence alignment from a group of related sequences using progressive, pairwise alignments to show relationship and percent sequence identity. It also plots a so-called “tree” or “dendrogram” showing the clustering relationships used to create the alignment (see, e.g., FIG. 2). PILEUP uses a simplification of the progressive alignment method of Feng & Doolittle, *J Mol. Evol.* 35:351–360 (1987). The method used is similar to the method described by Higgins & Sharp, *CABIOS* 5:151–153 (1989). The program can align up to 300 sequences, each of a maximum length of 5,000 nucleotides or amino acids. The multiple alignment procedure begins with the pairwise alignment of the two most similar sequences, producing a cluster of two aligned sequences. This cluster is then aligned to the next most related sequence or cluster of aligned sequences. Two clusters of sequences are aligned by a simple extension of the pairwise alignment of two individual sequences. The final alignment is achieved by a series of progressive, pairwise alignments. The program is run by designating specific sequences and their amino acid or nucleotide coordinates for regions of sequence comparison and by designating the program parameters. Using PILEUP, a reference sequence is compared to other test sequences to determine the percent sequence identity relationship using the following parameters: default gap weight (3.00), default gap length weight (0.10), and weighted end gaps. PILEUP can be obtained from the GCG sequence analysis software package, e.g., version 7. 0 (Devereaux et al., *Nuc. Acids Res.* 12:387–395 (1984) encoded by the genes were derived by conceptual translation of the corresponding open reading frames. Comparison of these protein sequences to all known proteins in the public sequence databases using BLASTP algorithm revealed their strong homology to the members of the T1R family, each of the T1R family sequences having at least about 35 to 50%, and preferably at least 55%, at least 60%, at least 65%, and most preferably at least 70%, amino acid identity to at least one known member of the family.

B. Definitions

As used herein, the following terms have the meanings ascribed to them unless specified otherwise.

“Taste cells” include neuroepithelial cells that are organized into groups to form taste buds of the tongue, e.g., foliate, fungiform, and circumvallate cells (see, e.g., Roper et al., *Ann. Rev. Neurosci.* 12:329–353 (1989)). Taste cells are also found in the palate and other tissues, such as the esophagus and the stomach.

“T1R” refers to one or more members of a family of G protein-coupled receptors that are expressed in taste cells such as foliate, fungiform, and circumvallate cells, as well as cells of the palate, and esophagus (see, e.g., Hoon et al., *Cell*, 96:541–551 (1999), herein incorporated by reference in its entirety). Members of this family are also referred to as GPCR-B3 and TR1 in WO 00/06592 as well as GPCR-B4 and TR2 in WO 00/06593. GPCR-B3 is also herein referred to as rT1R1, and GPCR-B4 is referred to as rT1R2. Taste receptor cells can also be identified on the basis of morphology (see, e.g., Roper, supra), or by the expression of proteins specifically expressed in taste cells. T1R family members may have the ability to act as receptors for sweet taste transduction, or to distinguish between various other taste modalities.

“T1R” nucleic acids encode a family of GPCRs with seven transmembrane regions that have “G protein-coupled receptor activity,” e.g., they may bind to G proteins in response to extracellular stimuli and promote production of second messengers such as IP3, cAMP, cGMP, and Ca²⁺ via stimulation of enzymes such as phospholipase C and adenylyl cyclase (for a description of the structure and function of GPCRs, see, e.g., Fong, supra, and Baldwin, supra). A single taste cell may contain many distinct T1R polypeptides.

The term “T1R” family therefore refers to polymorphic variants, alleles, mutants, and interspecies homologs that: (1) have at least about 35 to 50% amino acid sequence identity, optionally about 60, 75, 80, 85, 90, 95, 96, 97, 98, or 99% amino acid sequence identity to SEQ ID NOS: 4, 10, 12, 14, 17, or 21 over a window of about 25 amino acids, optionally 50–100 amino acids; (2) specifically bind to antibodies raised against an immunogen comprising an amino acid sequence selected from the group consisting of SEQ ID NOS: 4, 10, 12, 14, 17, 21 and conservatively modified variants thereof; (3) are encoded by a nucleic acid molecule which specifically hybridize (with a size of at least about 100, optionally at least about 500–1000 nucleotides) under stringent hybridization conditions to a sequence selected from the group consisting of SEQ ID NOS: 1, 2, 3, 9, 11, 13, 15, 16, 20, and conservatively modified variants thereof; (4) comprise a sequence at least about 35 to 50% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS: 4, 10, 12, 14, 17, or 21; or (5) are amplified by primers that specifically hybridize under stringent hybridization conditions to the same sequence as degenerate primer sets encoding SEQ ID NOS: 7, 8, and conservatively modified variants thereof.

Topologically, certain chemosensory GPCRs have an “N-terminal domain;” “extracellular domains;” “transmembrane domains” comprising seven transmembrane regions, and corresponding cytoplasmic, and extracellular loops; “cytoplasmic domains,” and a “C-terminal domain” (see, e.g., Hoon et al., *Cell*, 96:541–551 (1999); Buck & Axel, *Cell*, 65:175–187 (1991)). These domains can be structurally identified using methods known to those of skill in the art, such as sequence analysis programs that identify hydrophobic and hydrophilic domains (see, e.g., Stryer, *Biochemistry*,

(3rd ed. 1988); see also any of a number of Internet based sequence analysis programs, such as those found at dot.im-gen.bcm.tmc.edu). Such domains are useful for making chimeric proteins and for in vitro assays of the invention, e.g., ligand-binding assays.

“Extracellular domains” therefore refers to the domains of T1R polypeptides that protrude from the cellular membrane and are exposed to the extracellular face of the cell. Such domains generally include the “N terminal domain” that is exposed to the extracellular face of the cell, and optionally can include portions of the extracellular loops of the transmembrane domain that are exposed to the extracellular face of the cell, i.e., the loops between transmembrane regions 2 and 3, between transmembrane regions 4 and 5, and between transmembrane regions 6 and 7.

The “N terminal domain” region starts at the N-terminus and extends to a region close to the start of the transmembrane domain. More particularly, in one embodiment of the invention, this domain starts at the N-terminus and ends approximately at the conserved glutamic acid at amino acid position 563 plus or minus approximately 20 amino acid. These extracellular domains are useful for in vitro ligand-binding assays, both soluble and solid phase. In addition, transmembrane regions, described below, can also bind ligand either in combination with the extracellular domain, and are therefore also useful for in vitro ligand-binding assays.

“Transmembrane domain,” which comprises the seven “transmembrane regions,” refers to the domain of T1R polypeptides that lies within the plasma membrane, and may also include the corresponding cytoplasmic (intracellular) and extracellular loops. In one embodiment, this region corresponds to the domain of T1R family members which starts approximately at the conserved glutamic acid residue at amino acid position 563 plus or minus 20 amino acids and ends approximately at the conserved tyrosine amino acid residue at position 812 plus or minus approximately 10 amino acids. The seven transmembrane regions and extracellular and cytoplasmic loops can be identified using standard methods, as described in Kyte & Doolittle, *J. Mol. Biol.*, 157:105–32 (1982), or in Stryer, *supra*.

“Cytoplasmic domains” refers to the domains of T1R polypeptides that face the inside of the cell, e.g., the “C terminal domain” and the intracellular loops of the transmembrane domain, e.g., the intracellular loop between transmembrane regions 1 and 2, the intracellular loop between transmembrane regions 3 and 4, and the intracellular loop between transmembrane regions 5 and 6. “C terminal domain” refers to the region that spans the end of the last transmembrane domain and the C-terminus of the protein, and which is normally located within the cytoplasm. In one embodiment, this region starts at the conserved tyrosine amino acid residue at position 812 plus or minus approximately 10 amino acids and continues to the C-terminus of the polypeptide.

The term “ligand-binding region” or “ligand-binding domain” refers to sequences derived from a chemosensory receptor, particularly a taste receptor, that substantially incorporates at least the extracellular domain of the receptor. In one embodiment, the extracellular domain of the ligand-binding region may include the N-terminal domain and, optionally, portions of the transmembrane domain, such as the extracellular loops of the transmembrane domain. The ligand-binding region may be capable of binding a ligand, and more particularly, a taste stimulus.

The phrase “functional effects” in the context of assays for testing compounds that modulate T1R family member medi-

ated taste transduction includes the determination of any parameter that is indirectly or directly under the influence of the receptor, e.g., functional, physical and chemical effects. It includes ligand binding, changes in ion flux, membrane potential, current flow, transcription, G protein binding, GPCR phosphorylation or dephosphorylation, signal transduction, receptor-ligand interactions, second messenger concentrations (e.g., cAMP, cGMP, IP3, or intracellular Ca²⁺), in vitro, in vivo, and ex vivo and also includes other physiologic effects such increases or decreases of neurotransmitter or hormone release.

By “determining the functional effect” in the context of assays is meant assays for a compound that increases or decreases a parameter that is indirectly or directly under the influence of a T1R family member, e.g., functional, physical and chemical effects. Such functional effects can be measured by any means known to those skilled in the art, e.g., changes in spectroscopic characteristics (e.g., fluorescence, absorbance, refractive index), hydrodynamic (e.g., shape), chromatographic, or solubility properties, patch clamping, voltage-sensitive dyes, whole cell currents, radioisotope efflux, inducible markers, oocyte T1R gene expression; tissue culture cell T1R expression; transcriptional activation of T1R genes; ligand-binding assays; voltage, membrane potential and conductance changes; ion flux assays; changes in intracellular second messengers such as cAMP, cGMP, and inositol triphosphate (IP3); changes in intracellular calcium levels; neurotransmitter release, and the like.

“Inhibitors,” “activators,” and “modulators” of T1R genes or proteins are used interchangeably to refer to inhibitory, activating, or modulating molecules identified using in vitro and in vivo assays for taste transduction, e.g., ligands, agonists, antagonists, and their homologs and mimetics. Inhibitors are compounds that, e.g., bind to, partially or totally block stimulation, decrease, prevent, delay activation, inactivate, desensitize, or down regulate taste transduction, e.g., antagonists. Activators are compounds that, e.g., bind to, stimulate, increase, open, activate, facilitate, enhance activation, sensitize, or up regulate taste transduction, e.g., agonists. Modulators include compounds that, e.g., alter the interaction of a receptor with: extracellular proteins that bind activators or inhibitor (e.g., ebnerin and other members of the hydrophobic carrier family); G proteins; kinases (e.g., homologs of rhodopsin kinase and beta adrenergic receptor kinases that are involved in deactivation and desensitization of a receptor); and arrestins, which also deactivate and desensitize receptors. Modulators can include genetically modified versions of T1R family members, e.g., with altered activity, as well as naturally occurring and synthetic ligands, antagonists, agonists, small chemical molecules and the like. Such assays for inhibitors and activators include, e.g., expressing T1R family members in cells or cell membranes, applying putative modulator compounds, in the presence or absence of taste stimuli, e.g., sweet taste stimuli, and then determining the functional effects on taste transduction, as described above. Samples or assays comprising T1R family members that are treated with a potential activator, inhibitor, or modulator are compared to control samples without the inhibitor, activator, or modulator to examine the extent of modulation. Control samples (untreated with modulators) are assigned a relative T1R activity value of 100%. Inhibition of a T1R is achieved when the T1R activity value relative to the control is about 80%, optionally 50% or 25–0%. Activation of a T1R is achieved when the T1R activity value relative to the control is 110%, optionally 150%, optionally 200–500%, or 1000–3000% higher.

The terms “purified,” “substantially purified,” and “isolated” as used herein refer to the state of being free of other, dissimilar compounds with which the compound of the invention is normally associated in its natural state, so that the “purified,” “substantially purified,” and “isolated” subject comprises at least 0.5%, 1%, 5%, 10%, or 20%, and most preferably at least 50% or 75% of the mass, by weight, of a given sample. In one preferred embodiment, these terms refer to the compound of the invention comprising at least 95% of the mass, by weight, of a given sample. As used herein, the terms “purified,” “substantially purified,” and “isolated” “isolated,” when referring to a nucleic acid or protein, of nucleic acids or proteins, also refers to a state of purification or concentration different than that which occurs naturally in the mammalian, especially human, body. Any degree of purification or concentration greater than that which occurs naturally in the mammalian, especially human, body, including (1) the purification from other associated structures or compounds or (2) the association with structures or compounds to which it is not normally associated in the mammalian, especially human, body, are within the meaning of “isolated.” The nucleic acid or protein or classes of nucleic acids or proteins, described herein, may be isolated, or otherwise associated with structures or compounds to which they are not normally associated in nature, according to a variety of methods and processes known to those of skill in the art.

As used herein, the term “isolated,” when referring to a nucleic acid or polypeptide refers to a state of purification or concentration different than that which occurs naturally in the mammalian, especially human, body. Any degree of purification or concentration greater than that which occurs naturally in the body, including (1) the purification from other naturally-occurring associated structures or compounds, or (2) the association with structures or compounds to which it is not normally associated in the body are within the meaning of “isolated” as used herein. The nucleic acids or polypeptides described herein may be isolated or otherwise associated with structures or compounds to which they are not normally associated in nature, according to a variety of methods and processed known to those of skill in the art.

As used herein, the terms “amplifying” and “amplification” refer to the use of any suitable amplification methodology for generating or detecting recombinant or naturally expressed nucleic acid, as described in detail, below. For example, the invention provides methods and reagents (e.g., specific degenerate oligonucleotide primer pairs) for amplifying (e.g., by polymerase chain reaction, PCR) naturally expressed (e.g., genomic or mRNA) or recombinant (e.g., cDNA) nucleic acids of the invention (e.g., taste stimulus-binding sequences of the invention) in vivo or in vitro.

The term “7-transmembrane receptor” means a polypeptide belonging to a superfamily of transmembrane proteins that have seven domains that span the plasma membrane seven times (thus, the seven domains are called “transmembrane” or “TM” domains TM I to TM VII). The families of olfactory and certain taste receptors each belong to this super-family. 7-transmembrane receptor polypeptides have similar and characteristic primary, secondary and tertiary structures, as discussed in further detail below.

The term “library” means a preparation that is a mixture of different nucleic acid or polypeptide molecules, such as the library of recombinantly generated chemosensory, particularly taste receptor ligand-binding domains generated by amplification of nucleic acid with degenerate primer pairs, or an isolated collection of vectors that incorporate the

amplified ligand-binding domains, or a mixture of cells each randomly transfected with at least one vector encoding an taste receptor.

The term “nucleic acid” or “nucleic acid sequence” refers to a deoxyribonucleotide or ribonucleotide oligonucleotide in either single- or double-stranded form. The term encompasses nucleic acids, i.e., oligonucleotides, containing known analogs of natural nucleotides. The term also encompasses nucleic-acid-like structures with synthetic backbones (see e.g., *Oligonucleotides and Analogues, a Practical Approach*, ed. F. Eckstein, Oxford Univ. Press (1991); *Antisense Strategies, Annals of the N.Y. Academy of Sciences*, Vol. 600, Eds. Baserga et al. (NYAS 1992); Milligan *J. Med. Chem.* 36:1923–1937 (1993); *Antisense Research and Applications* (1993, CRC Press), WO 97/03211; WO 96/39154; Mata, *Toxicol. Appl. Pharmacol.* 144:189–197 (1997); Strauss-Soukup, *Biochemistry* 36:8692–8698 (1997); Samstag, *Antisense Nucleic Acid Drug Dev.* 6:153–156 (1996)).

Unless otherwise indicated, a particular nucleic acid sequence also implicitly encompasses conservatively modified variants thereof (e.g., degenerate codon substitutions) and complementary sequences, as well as the sequence explicitly indicated. Specifically, degenerate codon substitutions may be achieved by generating, e.g., sequences in which the third position of one or more selected codons is substituted with mixed-base and/or deoxyinosine residues (Batzer et al., *Nucleic Acid Res.*, 19:5081 (1991); Ohtsuka et al., *J. Biol. Chem.*, 260:2605–2608 (1985); Rossolini et al., *Mol. Cell. Probes*, 8:91–98 (1994)). The term nucleic acid is used interchangeably with gene, cDNA, mRNA, oligonucleotide, and polynucleotide.

The terms “polypeptide,” “peptide” and “protein” are used interchangeably herein to refer to a polymer of amino acid residues. The terms apply to amino acid polymers in which one or more amino acid residue is an artificial chemical mimetic of a corresponding naturally occurring amino acid, as well as to naturally occurring amino acid polymers and non-naturally occurring amino acid polymer.

The term “plasma membrane translocation domain” or simply “translocation domain” means a polypeptide domain that, when incorporated into the amino terminus of a polypeptide coding sequence, can with great efficiency “chaperone” or “translocate” the hybrid (“fusion”) protein to the cell plasma membrane. For instance, a “translocation domain” may be derived from the amino terminus of the bovine rhodopsin receptor polypeptide, a 7-transmembrane receptor. However, rhodopsin from any mammal may be used, as can other translocation facilitating sequences. Thus, the translocation domain is particularly efficient in translocating 7-transmembrane fusion proteins to the plasma membrane, and a protein (e.g., a taste receptor polypeptide) comprising an amino terminal translocating domain will be transported to the plasma membrane more efficiently than without the domain. However, if the N-terminal domain of the polypeptide is active in binding, the use of other translocation domains may be preferred.

The “translocation domain,” “ligand-binding domain”, and chimeric receptors compositions described herein also include “analogs,” or “conservative variants” and “mimetics” (“peptidomimetics”) with structures and activity that substantially correspond to the exemplary sequences. Thus, the terms “conservative variant” or “analog” or “mimetic” refer to a polypeptide which has a modified amino acid sequence, such that the change(s) do not substantially alter the polypeptide’s (the conservative variant’s) structure and/or activity, as defined herein. These include conservatively

modified variations of an amino acid sequence, i.e., amino acid substitutions, additions or deletions of those residues that are not critical for protein activity, or substitution of amino acids with residues having similar properties (e.g., acidic, basic, positively or negatively charged, polar or non-polar, etc.) such that the substitutions of even critical amino acids does not substantially alter structure and/or activity.

More particularly, “conservatively modified variants” applies to both amino acid and nucleic acid sequences. With respect to particular nucleic acid sequences, conservatively modified variants refers to those nucleic acids which encode identical or essentially identical amino acid sequences, or where the nucleic acid does not encode an amino acid sequence, to essentially identical sequences. Because of the degeneracy of the genetic code, a large number of functionally identical nucleic acids encode any given protein.

For instance, the codons GCA, GCC, GCG and GCU all encode the amino acid alanine. Thus, at every position where an alanine is specified by a codon, the codon can be altered to any of the corresponding codons described without altering the encoded polypeptide.

Such nucleic acid variations are “silent variations,” which are one species of conservatively modified variations. Every nucleic acid sequence herein which encodes a polypeptide also describes every possible silent variation of the nucleic acid. One of skill will recognize that each codon in a nucleic acid (except AUG, which is ordinarily the only codon for methionine, and TGG, which is ordinarily the only codon for tryptophan) can be modified to yield a functionally identical molecule. Accordingly, each silent variation of a nucleic acid which encodes a polypeptide is implicit in each described sequence.

Conservative substitution tables providing functionally similar amino acids are well known in the art. For example, one exemplary guideline to select conservative substitutions includes (original residue followed by exemplary substitution): ala/gly or ser; arg/lys; asn/gln or his; asp/glu; cys/ser; gln/asn; gly/asp; gly/ala or pro; his/asn or gln; ile/leu or val; leu/ile or val; lys/arg or gln or glu; met/leu or tyr or ile; phe/met or leu or tyr; ser/thr; thr/ser; trp/tyr; tyr/trp or phe; val/ile or leu. An alternative exemplary guideline uses the following six groups, each containing amino acids that are conservative substitutions for one another: 1) Alanine (A), Serine (S), Threonine (T); 2) Aspartic acid (D), Glutamic acid (E); 3) Asparagine (N), Glutamine (Q); 4) Arginine (R), Lysine (I); 5) Isoleucine (I), Leucine (L), Methionine (M), Valine (V); and 6) Phenylalanine (F), Tyrosine (Y), Tryptophan (W); (see also, e.g., Creighton, *Proteins*, W.H. Freeman and Company (1984); Schultz and Schimer, *Principles of Protein Structure*, Springer-Verlag (1979)). One of skill in the art will appreciate that the above-identified substitutions are not the only possible conservative substitutions. For example, for some purposes, one may regard all charged amino acids as conservative substitutions for each other whether they are positive or negative. In addition, individual substitutions, deletions or additions that alter, add or delete a single amino acid or a small percentage of amino acids in an encoded sequence can also be considered “conservatively modified variations.”

The terms “mimetic” and “peptidomimetic” refer to a synthetic chemical compound that has substantially the same structural and/or functional characteristics of the polypeptides, e.g., translocation domains, ligand-binding domains, or chimeric receptors of the invention. The mimetic can be either entirely composed of synthetic, non-natural analogs of amino acids, or may be a chimeric molecule of partly natural

peptide amino acids and partly non-natural analogs of amino acids. The mimetic can also incorporate any amount of natural amino acid conservative substitutions as long as such substitutions also do not substantially alter the mimetic’s structure and/or activity.

As with polypeptides of the invention which are conservative variants, routine experimentation will determine whether a mimetic is within the scope of the invention, i.e., that its structure and/or function is not substantially altered. Polypeptide mimetic compositions can contain any combination of non-natural structural components, which are typically from three structural groups: a) residue linkage groups other than the natural amide bond (“peptide bond”) linkages; b) non-natural residues in place of naturally occurring amino acid residues; or c) residues which induce secondary structural mimicry, i.e., to induce or stabilize a secondary structure, e.g., a beta turn, gamma turn, beta sheet, alpha helix conformation, and the like. A polypeptide can be characterized as a mimetic when all or some of its residues are joined by chemical means other than natural peptide bonds. Individual peptidomimetic residues can be joined by peptide bonds, other chemical bonds or coupling means, such as, e.g., glutaraldehyde, N-hydroxysuccinimide esters, bifunctional maleimides, N,N'-dicyclohexylcarbodiimide (DCC) or N,N'-diisopropylcarbodiimide (DIC). Linking groups that can be an alternative to the traditional amide bond (“peptide bond”) linkages include, e.g., ketomethylene (e.g., $-\text{C}(=\text{O})-\text{CH}_2-$ for $-\text{C}(=\text{O})-\text{NH}-$), aminomethylene (CH_2-NH), ethylene, olefin ($\text{CH}=\text{CH}$), ether (CH_2-O), thioether (CH_2-S), tetrazole (CN_4), thiazole, retroamide, thioamide, or ester (see, e.g., Spatola, *Chemistry and Biochemistry of Amino Acids, Peptides and Proteins*, Vol. 7, pp 267–357, “Peptide Backbone Modifications,” Marcell Dekker, NY (1983)). A polypeptide can also be characterized as a mimetic by containing all or some non-natural residues in place of naturally occurring amino acid residues; non-natural residues are well described in the scientific and patent literature.

A “label” or a “detectable moiety” is a composition detectable by spectroscopic, photochemical, biochemical, immunochemical, or chemical means. For example, useful labels include ^{32}P , fluorescent dyes, electron-dense reagents, enzymes (e.g., as commonly used in an ELISA), biotin, digoxigenin, or haptens and proteins which can be made detectable, e.g., by incorporating a radiolabel into the peptide or used to detect antibodies specifically reactive with the peptide.

A “labeled nucleic acid probe or oligonucleotide” is one that is bound, either covalently, through a linker or a chemical bond, or noncovalently, through ionic, van der Waals, electrostatic, or hydrogen bonds to a label such that the presence of the probe may be detected by detecting the presence of the label bound to the probe.

As used herein a “nucleic acid probe or oligonucleotide” is defined as a nucleic acid capable of binding to a target nucleic acid of complementary sequence through one or more types of chemical bonds, usually through complementary base pairing, usually through hydrogen bond formation. As used herein, a probe may include natural (i.e., A, G, C, or T) or modified bases (7-deazaguanosine, inosine, etc.). In addition, the bases in a probe may be joined by a linkage other than a phosphodiester bond, so long as it does not interfere with hybridization. Thus, for example, probes may be peptide nucleic acids in which the constituent bases are joined by peptide bonds rather than phosphodiester linkages. It will be understood by one of skill in the art that probes may bind target sequences lacking complete complementa-

rity with the probe sequence depending upon the stringency of the hybridization conditions. The probes are optionally directly labeled as with isotopes, chromophores, lumiphores, chromogens, or indirectly labeled such as with biotin to which a streptavidin complex may later bind. By assaying for the presence or absence of the probe, one can detect the presence or absence of the select sequence or subsequence.

The term "heterologous" when used with reference to portions of a nucleic acid indicates that the nucleic acid comprises two or more subsequences that are not found in the same relationship to each other in nature. For instance, the nucleic acid is typically recombinantly produced, having two or more sequences from unrelated genes arranged to make a new functional nucleic acid, e.g., a promoter from one source and a coding region from another source. Similarly, a heterologous protein indicates that the protein comprises two or more subsequences that are not found in the same relationship to each other in nature (e.g., a fusion protein).

A "promoter" is defined as an array of nucleic acid sequences that direct transcription of a nucleic acid. As used herein, a promoter includes necessary nucleic acid sequences near the start site of transcription, such as, in the case of a polymerase type promoter, a TATA element. A promoter also optionally includes distal enhancer or repressor elements, which can be located as much as several thousand base pairs from the start site of transcription. A "constitutive" promoter is a promoter that is active under most environmental and developmental conditions. An "inducible" promoter is a promoter that is active under environmental or developmental regulation. The term "operably linked" refers to a functional linkage between a nucleic acid expression control sequence (such as a promoter, or array of transcription factor binding sites) and a second nucleic acid sequence, wherein the expression control sequence directs transcription of the nucleic acid corresponding to the second sequence.

As used herein, "recombinant" refers to a polynucleotide synthesized or otherwise manipulated in vitro (e.g., "recombinant polynucleotide"), to methods of using recombinant polynucleotides to produce gene products in cells or other biological systems, or to a polypeptide ("recombinant protein") encoded by a recombinant polynucleotide. "Recombinant means" also encompass the ligation of nucleic acids having various coding regions or domains or promoter sequences from different sources into an expression cassette or vector for expression of, e.g., inducible or constitutive expression of a fusion protein comprising a translocation domain of the invention and a nucleic acid sequence amplified using a primer of the invention.

The phrase "selectively (or specifically) hybridizes to" refers to the binding, duplexing, or hybridizing of a molecule only to a particular nucleotide sequence under stringent hybridization conditions when that sequence is present in a complex mixture (e.g., total cellular or library DNA or RNA).

The phrase "stringent hybridization conditions" refers to conditions under which a probe will hybridize to its target subsequence, typically in a complex mixture of nucleic acid, but to no other sequences. Stringent conditions are sequence-dependent and will be different in different circumstances. Longer sequences hybridize specifically at higher temperatures. An extensive guide to the hybridization of nucleic acids is found in Tijssen, *Techniques in Biochemistry and Molecular Biology—Hybridisation with Nucleic Probes*, "Overview of principles of hybridization and the strategy of nucleic acid assays" (1993). Generally, stringent

conditions are selected to be about 5–10° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic concentration) at which 50% of the probes complementary to the target hybridize to the target sequence at equilibrium (as the target sequences are present in excess, at T_m, 50% of the probes are occupied at equilibrium). Stringent conditions will be those in which the salt concentration is less than about 1.0 M sodium ion, typically about 0.01 to 1.0 M sodium ion concentration (or other salts) at pH 7.0 to 8.3 and the temperature is at least about 30° C. for short probes (e.g., 10 to 50 nucleotides) and at least about 60° C. for long probes (e.g., greater than 50 nucleotides). Stringent conditions may also be achieved with the addition of destabilizing agents such as formamide. For selective or specific hybridization, a positive signal is at least two times background, optionally 10 times background hybridization. Exemplary stringent hybridization conditions can be as following: 50% formamide, S×SSC, and 1% SDS, incubating at 42° C., or, S×SSC, 1% SDS, incubating at 65° C., with wash in 0.2×SSC, and 0.1% SDS at 65° C. Such hybridizations and wash steps can be carried out for, e.g., 1, 2, 5, 10, 15, 30, 60; or more minutes.

Nucleic acids that do not hybridize to each other under stringent conditions are still substantially related if the polypeptides which they encode are substantially related. This occurs, for example, when a copy of a nucleic acid is created using the maximum codon degeneracy permitted by the genetic code. In such cases, the nucleic acids typically hybridize under moderately stringent hybridization conditions. Exemplary "moderately stringent hybridization conditions" include a hybridization in a buffer of 40% formamide, 1 M NaCl, 1% SDS at 37° C., and a wash in 1×SSC at 45° C. Such hybridizations and wash steps can be carried out for, e.g., 1, 2, 5, 10, 15, 30, 60, or more minutes. A positive hybridization is at least twice background. Those of ordinary skill will readily recognize that alternative hybridization and wash conditions can be utilized to provide conditions of similar stringency.

"Antibody" refers to a polypeptide comprising a framework region from an immunoglobulin gene or fragments thereof that specifically binds and recognizes an antigen. The recognized immunoglobulin genes include the kappa, lambda, alpha, gamma, delta, epsilon, and mu constant region genes, as well as the myriad immunoglobulin variable region genes. Light chains are classified as either kappa or lambda. Heavy chains are classified as gamma, mu, alpha, delta, or epsilon, which in turn define the immunoglobulin classes, IgG, IgM, IgA, IgD and IgE, respectively.

An exemplary immunoglobulin (antibody) structural unit comprises a tetramer. Each tetramer is composed of two identical pairs of polypeptide chains, each pair having one "light" (about 25 kDa) and one "heavy" chain (about 50–70 kDa). The N-terminus of each chain defines a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The terms variable light chain (VL) and variable heavy chain (VH) refer to these light and heavy chains respectively.

A "chimeric antibody" is an antibody molecule in which (a) the constant region, or a portion thereof, is altered, replaced or exchanged so that the antigen binding site (variable region) is linked to a constant region of a different or altered class, effector function and/or species, or an entirely different molecule which confers new properties to the chimeric antibody, e.g., an enzyme, toxin, hormone, growth factor, drug, etc.; or (b) the variable region, or a

portion thereof, is altered, replaced or exchanged with a variable region having a different or altered antigen specificity.

An “anti-T1R” antibody is an antibody or antibody fragment that specifically binds a polypeptide encoded by a T1R gene, cDNA, or a subsequence thereof.

The term “immunoassay” is an assay that uses an antibody to specifically bind an antigen. The immunoassay is characterized by the use of specific binding properties of a particular antibody to isolate, target, and/or quantify the antigen.

The phrase “specifically (or selectively) binds” to an antibody or, “specifically (or selectively) immunoreactive with,” when referring to a protein or peptide, refers to a binding reaction that is determinative of the presence of the protein in a heterogeneous population of proteins and other biologics. Thus, under designated immunoassay conditions, the specified antibodies bind to a particular protein at least two times the background and do not substantially bind in a significant amount to other proteins present in the sample. Specific binding to an antibody under such conditions may require an antibody that is selected for its specificity for a particular protein. For example, polyclonal antibodies raised to a T1R family member from specific species such as rat, mouse, or human can be selected to obtain only those polyclonal antibodies that are specifically immunoreactive with the T1R polypeptide or an immunogenic portion thereof and not with other proteins, except for orthologs or polymorphic variants and alleles of the T1R polypeptide. This selection may be achieved by subtracting out antibodies that cross-react with T1R molecules from other species or other T1R molecules. Antibodies can also be selected that recognize only T1R GPCR family members but not GPCRs from other families. A variety of immunoassay formats may be used to select antibodies specifically immunoreactive with a particular protein. For example, solid-phase ELISA immunoassays are routinely used to select antibodies specifically immunoreactive with a protein (see, e.g., Harlow & Lane, *Antibodies, A Laboratory Manual*, (1988), for a description of immunoassay formats and conditions that can be used to determine specific immunoreactivity). Typically a specific or selective reaction will be at least twice background signal or noise and more typically more than 10 to 100 times background.

The phrase “selectively associates with” refers to the ability of a nucleic acid to “selectively hybridize” with another as defined above, or the ability of an antibody to “selectively (or specifically) bind to a protein, as defined above.

The term “expression vector” refers to any recombinant expression system for the purpose of expressing a nucleic acid sequence of the invention in vitro or in vivo, constitutively or inducibly, in any cell, including prokaryotic, yeast, fungal, plant, insect or mammalian cell. The term includes linear or circular expression systems. The term includes expression systems that remain episomal or integrate into the host cell genome. The expression systems can have the ability to self-replicate or not, i.e., drive only transient expression in a cell. The term includes recombinant expression “cassettes which contain only the minimum elements needed for transcription of the recombinant nucleic acid.

By “host cell” is meant a cell that contains an expression vector and supports the replication or expression of the expression vector. Host cells may be prokaryotic cells such as *E. coli*, or eukaryotic cells such as yeast, insect, amphibian, or mammalian cells such as CHO, HeLa, HEK-293, and the like, e.g., cultured cells, explants, and cells in vivo.

A. Isolation and Expression of T1R Polypeptides

Isolation and expression of the T1Rs, or fragments or variants thereof, of the invention can be performed as described below. PCR primers can be used for the amplification of nucleic acids encoding taste receptor ligand-binding regions, and libraries of these nucleic acids can optionally be generated. Individual expression vectors or libraries of expression vectors can then be used to infect or transfect host cells for the functional expression of these nucleic acids or libraries. These genes and vectors can be made and expressed in vitro or in vivo. One of skill will recognize that desired phenotypes for altering and controlling nucleic acid expression can be obtained by modulating the expression or activity of the genes and nucleic acids (e.g., promoters, enhancers and the like) within the vectors of the invention. Any of the known methods described for increasing or decreasing expression or activity can be used. The invention can be practiced in conjunction with any method or protocol known in the art, which are well described in the scientific and patent literature.

The nucleic acid sequences of the invention and other nucleic acids used to practice this invention, whether RNA, cDNA, genomic DNA, vectors, viruses or hybrids thereof, may be isolated from a variety of sources, genetically engineered, amplified, and/or expressed recombinantly. Any recombinant expression system can be used, including, in addition to mammalian cells, e.g., bacterial, yeast, insect, or plant systems.

Alternatively, these nucleic acids can be synthesized in vitro by well-known chemical synthesis techniques, as described in, e.g., Carruthers, *Cold Spring Harbor Symp. Quant. Biol.* 47:411–418 (1982); Adams, *Am. Chem. Soc.* 105:661 (1983); Belousov, *Nucleic Acids Res.* 25:3440–3444 (1997); Frenkel, *Free Radic. Biol. Med.* 19:373–380 (1995); Blommers, *Biochemistry* 33:7886–7896 (1994); Narang, *Meth. Enzymol.* 68:90 (1979); Brown, *Meth. Enzymol.* 68:109 (1979); Beaucage, *Tetra. Lett.* 22:1859 (1981); U.S. Pat. No. 4,458,066. Double-stranded DNA fragments may then be obtained either by synthesizing the complementary strand and annealing the strands together under appropriate conditions, or by adding the complementary strand using DNA polymerase with an appropriate primer sequence.

Techniques for the manipulation of nucleic acids, such as, for example, for generating mutations in sequences, subcloning, labeling probes, sequencing, hybridization and the like are well described in the scientific and patent literature. See, e.g., Sambrook, ed., *Molecular Cloning: a Laboratory manual* (2nd ed.), Vols. 1–3, Cold Spring Harbor Laboratory (1989); *Current Protocols in Molecular Biology*, Ausubel, ed. John Wiley & Sons, Inc., New York (1997); *Laboratory Techniques in Biochemistry and Molecular Biology: Hybridization With Nucleic Acid Probes, Part I, Theory and Nucleic Acid Preparation*, Tijssen, ed. Elsevier, N.Y. (1993).

Nucleic acids, vectors, capsids, polypeptides, and the like can be analyzed and quantified by any of a number of general means well known to those of skill in the art. These include, e.g., analytical biochemical methods such as NMR, spectrophotometry, radiography, electrophoresis, capillary electrophoresis, high performance liquid chromatography (HPLC), thin layer chromatography (TLC), and hyperdiffusion chromatography, various immunological methods, e.g., fluid or gel precipitin reactions, immunodiffusion, immunoelectrophoresis, radioimmunoassays (RIAs), enzyme-linked immunosorbent assays (ELISAs), immuno-fluorescent assays, Southern analysis, Northern analysis, dot-blot analysis, gel electrophoresis (e.g., SDS-PAGE), RT-PCR, quan-

titative PCR, other nucleic acid or target or signal amplification methods, radiolabeling, scintillation counting, and affinity chromatography.

Oligonucleotide primers may be used to amplify nucleic acid fragments encoding taste receptor ligand-binding regions. The nucleic acids described herein can also be cloned or measured quantitatively using amplification techniques. Amplification methods are also well known in the art, and include, e.g., polymerase chain reaction, PCR (*PCR Protocols, a Guide to Methods and Applications*, ed. Innis, Academic Press, N.Y. (1990) and *PCR Strategies*, ed. Innis, Academic Press, Inc., N.Y. (1995), ligase chain reaction (LCR) (see, e.g., Wu, *Genomics* 4:560 (1989); Landegren, *Science* 241:1077,(1988); Barringer, *Gene* 89:117 (1990)); transcription amplification (see, e.g., Kwoh, *Proc. Natl. Acad. Sci. USA* 86:1173 (1989)); and, self-sustained sequence replication (see, e.g., Guatelli, *Proc. Natl. Acad. Sci. USA* 87:1874 (1990)); Q Beta replicase amplification (see, e.g., Smith, *J. Clin. Microbiol.* 35:1477–1491 (1997)); automated Q-beta replicase amplification assay (see, e.g., Burg, *Mol. Cell. Probes* 10:257–271 (1996)) and other RNA polymerase mediated techniques (e.g., NASBA, Cangene, Mississauga, Ontario); see also Berger, *Methods Enzymol.* 152:307–316 (1987); Sambrook; Ausubel; U.S. Pat. Nos. 4,683,195 and 4,683,202; Sooknanan, *Biotechnology* 13:563–564 (1995). The primers can be designed to retain the original sequence of the “donor” 7-membrane receptor. Alternatively, the primers can encode amino acid residues that are conservative substitutions (e.g., hydrophobic for hydrophobic residue, see above discussion) or functionally benign substitutions (e.g., do not prevent plasma membrane insertion, cause cleavage by peptidase, cause abnormal folding of receptor, and the like). Once amplified, the nucleic acids, either individually or as libraries, may be cloned according to methods known in the art, if desired, into any of a variety of vectors using routine molecular biological methods; methods for cloning in vitro amplified nucleic acids are described, e.g., U.S. Pat. No. 5,426,039.

The primer pairs may be designed to selectively amplify ligand-binding regions of the T1R family members. These regions may vary for different ligands or taste stimuli. Thus, what may be a minimal binding region for one taste stimulus, may be too limiting for a second taste stimulus. Accordingly, ligand-binding regions of different sizes comprising different extracellular domain structures may be amplified.

Paradigms to design degenerate primer pairs are well known in the art. For example, a Consensus-DEgenerate Hybrid Oligonucleotide Primer (CODEHOP) strategy computer program is accessible as <http://blocks.fhcrc.org/codehop.html>, and is directly linked from the BlockMaker multiple sequence alignment site for hybrid primer prediction beginning with a set of related protein sequences, as known taste receptor ligand-binding regions (see, e.g., Rose, *Nucleic Acids Res.* 26:1628–1635 (1998); Singh, *Biotechniques* 24:318–319 (1998)).

Means to synthesize oligonucleotide primer pairs are well known in the art. “Natural” base pairs or synthetic base pairs can be used. For example, use of artificial nucleobases offers a versatile approach to manipulate primer sequence and generate a more complex mixture of amplification products. Various families of artificial nucleobases are capable of assuming multiple hydrogen bonding orientations through internal bond rotations to provide a means for degenerate molecular recognition. Incorporation of these analogs into a single position of a PCR primer allows for generation of a complex library of amplification products. See, e.g., Hoops, *Nucleic Acids Res.* 25:4866–4871 (1997). Nonpolar mol-

ecules can also be used to mimic the shape of natural DNA bases. A non-hydrogen-bonding shape mimic for adenine can replicate efficiently and selectively against a nonpolar shape mimic for thymine (see, e.g., Morales, *Nat. Struct. Biol.* 5:950–954 (1998)). For example, two degenerate bases can be the pyrimidine base 6H, 8H-3,4-dihydropyrimido[4, 5-c][1,2]oxazin-7-one or the purine base N6-methoxy-2,6-diaminopurine (see, e.g., Hill, *Proc. Natl. Acad. Sci. USA* 95:4258–4263 (1998)). Exemplary degenerate primers of the invention incorporate the nucleobase analog

5'-(Dimethoxytrityl)-N-benzoyl-2'-deoxy-Cytidine,3'-[(2-cyanoethyl)-(N,N-diisopropyl)]-phosphoramidite (the term “P” in the sequences, see above). This pyrimidine analog hydrogen bonds with purines, including A and G residues. Polymorphic variants, alleles, and interspecies homologs that are substantially identical to a taste receptor disclosed herein can be isolated using the nucleic acid probes described above. Alternatively, expression libraries can be used to clone T1R polypeptides and polymorphic variants, alleles, and interspecies homologs thereof, by detecting expressed homologs immunologically with antisera or purified antibodies made against a T1R polypeptide, which also recognize and selectively bind to the T1R homolog.

Nucleic acids that encode ligand-binding regions of taste receptors may be generated by amplification (e.g., PCR) of appropriate nucleic acid sequences using degenerate primer pairs. The amplified nucleic acid can be genomic DNA from any cell or tissue or mRNA or cDNA derived from taste receptor-expressing cells.

In one embodiment, hybrid protein-coding sequences comprising nucleic acids encoding T1Rs fused to a translocation sequences may be constructed. Also provided are hybrid T1Rs comprising the translocation motifs and taste stimulus-binding domains of other families of chemosensory receptors, particularly taste receptors. These nucleic acid sequences can be operably linked to transcriptional or translational control elements, e.g., transcription and translation initiation sequences, promoters and enhancers, transcription and translation terminators, polyadenylation sequences, and other sequences useful for transcribing DNA into RNA. In construction of recombinant expression cassettes, vectors, and transgenics, a promoter fragment can be employed to direct expression of the desired nucleic acid in all desired cells or tissues.

In another embodiment, fusion proteins may include C-terminal or N-terminal translocation sequences. Further, fusion proteins can comprise additional elements, e.g., for protein detection, purification, or other applications. Detection and purification facilitating domains include, e.g., metal chelating peptides such as polyhistidine tracts, histidine-tryptophan modules, or other domains that allow purification on immobilized metals; maltose binding protein; protein A domains that allow purification on immobilized immunoglobulin; or the domain utilized in the FLAGS extension/affinity purification system (Immunex Corp, Seattle Wash.).

The inclusion of a cleavable linker sequences such as Factor Xa (see, e.g., Ottavi, *Biochimie* 80:289–293 (1998)), subtilisin protease recognition motif (see, e.g., Polyak, *Protein Eng.* 10:615–619 (1997)); enterokinase (Invitrogen, San Diego, Calif.), and the like, between the translocation domain (for efficient plasma membrane expression) and the rest of the newly translated polypeptide may be useful to facilitate purification. For example, one construct can include a polypeptide encoding a nucleic acid sequence linked to six histidine residues followed by a thioredoxin, an enterokinase cleavage site (see, e.g., Williams, *Biochemistry* 34:1787–1797 (1995)), and an C-terminal translocation

domain. The histidine residues facilitate detection and purification while the enterokinase cleavage site provides a means for purifying the desired protein(s) from the remainder of the fusion protein. Technology pertaining to vectors encoding fusion proteins and application of fusion proteins are well described in the scientific and patent literature, see, e.g., Kroll, *DNA Cell. Biol.* 12:441–53 (1993).

Expression vectors, either as individual expression vectors or as libraries of expression vectors, comprising the ligand-binding domain encoding sequences may be introduced into a genome or into the cytoplasm or a nucleus of a cell and expressed by a variety of conventional techniques, well described in the scientific and patent literature. See, e.g., Roberts, *Nature* 328:731 (1987); Berger supra; Schneider, *Protein Expr. Purif.* 6435:10 (1995); Sambrook; Tijssen; Ausubel. Product information from manufacturers of biological reagents and experimental equipment also provide information regarding known biological methods. The vectors can be isolated from natural sources, obtained from such sources as ATCC or GenBank libraries, or prepared by synthetic or recombinant methods.

The nucleic acids can be expressed in expression cassettes, vectors or viruses which are stably or transiently expressed in cells (e.g., episomal expression systems). Selection markers can be incorporated into expression cassettes and vectors to confer a selectable phenotype on transformed cells and sequences. For example, selection markers can code for episomal maintenance and replication such that integration into the host genome is not required. For example, the marker may encode antibiotic resistance (e.g., chloramphenicol, kanamycin, G418, bleomycin, hygromycin) or herbicide resistance (e.g., chlorosulfuron or Basta) to permit selection of those cells transformed with the desired DNA sequences (see, e.g., Blondelet-Rouault, *Gene* 190:315–317 (1997); Aubrecht, *J. Pharmacol. Exp. Ther.* 281:992–997 (1997)). Because selectable marker genes conferring resistance to substrates like neomycin or hygromycin can only be utilized in tissue culture, chemoresistance genes are also used as selectable markers in vitro and in vivo.

A chimeric nucleic acid sequence may encode a T1R ligand-binding domain within any 7-transmembrane polypeptide. Because 7-transmembrane receptor polypeptides have similar primary sequences and secondary and tertiary structures, structural domains (e.g., extracellular domain, TM domains, cytoplasmic domain, etc.) can be readily identified by sequence analysis. For example, homology modeling, Fourier analysis and helical periodicity detection can identify and characterize the seven domains with a 7-transmembrane receptor sequence. Fast Fourier Transform (FFT) algorithms can be used to assess the dominant periods that characterize profiles of the hydrophobicity and variability of analyzed sequences. Periodicity detection enhancement and alpha helical periodicity index can be done as by, e.g., Donnelly, *Protein Sci.* 2:55–70 (1993). Other alignment and modeling algorithms are well known in the art, see, e.g., Peitsch, *Receptors Channels* 4:161–164 (1996); Kyte & Doolittle, *J. Md. Bio.*, 157:105–132 (1982); Cronet, *Protein Eng.* 6:59–64 (1993) (homology and “discover modeling”).

The present invention also includes not only the DNA and proteins having the specified nucleic and amino acid sequences, but also DNA fragments, particularly fragments of, e.g., 40, 60, 80, 100, 150, 200, or 250 nucleotides, or more, as well as protein fragments of, e.g., 10, 20, 30, 50, 70, 100, or 150 amino acids, or more. Optionally, the nucleic acid fragments can encode an antigenic polypeptide which is capable of binding to an antibody raised against a T1R family member. Further, a protein fragment of the invention

can optionally be an antigenic fragment which is capable of binding to an antibody raised against a T1R family member.

Also contemplated are chimeric proteins, comprising at least 10, 20, 30, 50, 70, 100, or 150 amino acids, or more, of one of at least one of the T1R polypeptides described herein, coupled to additional amino acids representing all or part of another GPCR, preferably a member of the 7 transmembrane superfamily. These chimeras can be made from the instant receptors and another GPCR, or they can be made by combining two or more of the present receptors. In one embodiment, one portion of the chimera corresponds to or is derived from the extracellular domain of a T1R polypeptide of the invention. In another embodiment, one portion of the chimera corresponds to, or is derived from the extracellular domain and one or more of the transmembrane domains of a T1R polypeptide described herein, and the remaining portion or portions can come from another GPCR. Chimeric receptors are well known in the art, and the techniques for creating them and the selection and boundaries of domains or fragments of G protein-coupled receptors for incorporation therein are also well known. Thus, this knowledge of those skilled in the art can readily be used to create such chimeric receptors. The use of such chimeric receptors can provide, for example, a taste selectivity characteristic of one of the receptors specifically disclosed herein, coupled with the signal transduction characteristics of another receptor, such as a well known receptor used in prior art assay systems.

For example, a domain such as a ligand-binding domain, an extracellular domain, a transmembrane domain, a transmembrane domain, a cytoplasmic domain, an N-terminal domain, a C-terminal domain, or any combination thereof, can be covalently linked to a heterologous protein. For instance, an T1R extracellular domain can be linked to a heterologous GPCR transmembrane domain, or a heterologous GPCR extracellular domain can be linked to a T1R transmembrane domain. Other heterologous proteins of choice can include, e.g., green fluorescent protein, β -gal, glutamate receptor, and the rhodopsin presequence.

Also within the scope of the invention are host cells for expressing the T1Rs, fragments, or variants of the invention. To obtain high levels of expression of a cloned gene or nucleic acid, such as cDNAs encoding the T1Rs, fragments, or variants of the invention, one of skill typically subclones the nucleic acid sequence of interest into an expression vector that contains a strong promoter to direct transcription, a transcription/translation terminator, and if for a nucleic acid encoding a protein, a ribosome binding site for translational initiation. Suitable bacterial promoters are well known in the art and described, e.g., in Sambrook et al. However, bacterial or eukaryotic expression systems can be used.

Any of the well known procedures for introducing foreign nucleotide sequences into host cells may be used. These include the use of calcium phosphate transfection, polybrene, protoplast fusion, electroporation, liposomes, microinjection, plasma vectors, viral vectors and any of the other well known methods for introducing cloned genomic DNA, cDNA, synthetic DNA or other foreign genetic material into a host cell (see, e.g., Sambrook et al.) It is only necessary that the particular genetic engineering procedure used be capable of successfully introducing at least one nucleic acid molecule into the host cell capable of expressing the T1R, fragment, or variant of interest.

After the expression vector is introduced into the cells, the transfected cells are cultured under conditions favoring expression of the receptor, fragment, or variant of interest,

which is then recovered from the culture using standard techniques. Examples of such techniques are well known in the art. See, e.g., WO 00/06593, which is incorporated by reference in a manner consistent with this disclosure.

B. Detection of T1R Polypeptides

In addition to the detection of T1R genes and gene expression using nucleic acid hybridization technology, one can also use immunoassays to detect T1Rs, e.g., to identify taste receptor cells, and variants of T1R family members. Immunoassays can be used to qualitatively or quantitatively analyze the T1Rs. A general overview of the applicable technology can be found in Harlow & Lane, *Antibodies: A Laboratory Manual* (1988).

1. Antibodies to T1R Family Members

Methods of producing polyclonal and monoclonal antibodies that react specifically with a T1R family member are known to those of skill in the art (see, e.g., Coligan, *Current Protocols in Immunology* (1991); Harlow & Lane, *supra*; Goding, *Monoclonal Antibodies: Principles and Practice* (2d ed. 1986); and Kohler & Milstein, *Nature*, 256:495–497 (1975)). Such techniques include antibody preparation by selection of antibodies from libraries of recombinant antibodies in phage or similar vectors, as well as preparation of polyclonal and monoclonal antibodies by immunizing rabbits or mice (see, e.g., Huse et al., *Science*, 246:1275–1281 (1989); Ward et al., *Nature*, 341:544–546 (1989)).

A number of T1R-comprising immunogens may be used to produce antibodies specifically reactive with a T1R family member. For example, a recombinant T1R polypeptide, or an antigenic fragment thereof, can be isolated as described herein. Suitable antigenic regions include, e.g., the consensus sequences that are used to identify members of the T1R family. Recombinant proteins can be expressed in eukaryotic or prokaryotic cells as described above, and purified as generally described above. Recombinant protein is the preferred immunogen for the production of monoclonal or polyclonal antibodies. Alternatively, a synthetic peptide derived from the sequences disclosed herein and conjugated to a carrier protein can be used as an immunogen. Naturally occurring protein may also be used either in pure or impure form. The product is then injected into an animal capable of producing antibodies. Either monoclonal or polyclonal antibodies may be generated, for subsequent use in immunoassays to measure the protein.

Methods of production of polyclonal antibodies are known to those of skill in the art. For example, an inbred strain of mice (e.g., BALB/C mice) or rabbits is immunized with the protein using a standard adjuvant, such as Freund's adjuvant, and a standard immunization protocol. The animal's immune response to the immunogen preparation is monitored by taking test bleeds and determining the titer of reactivity to the T1R. When appropriately high titers of antibody to the immunogen are obtained, blood is collected from the animal and antisera are prepared. Further fractionation of the antisera to enrich for antibodies reactive to the protein can be done if desired (see Harlow & Lane, *supra*).

Monoclonal antibodies may be obtained by various techniques familiar to those skilled in the art. Briefly, spleen cells from an animal immunized with a desired antigen may be immortalized, commonly by fusion with a myeloma cell (see Kohler & Milstein, *Eur. J. Immunol.*, 6:511–519 (1976)). Alternative methods of immortalization include transformation with Epstein Barr Virus, oncogenes, or retroviruses, or other methods well known in the art. Colonies arising from single immortalized cells are screened for production of antibodies of the desired specificity and affinity for the

antigen, and yield of the monoclonal antibodies produced by such cells may be enhanced by various techniques, including injection into the peritoneal cavity of a vertebrate host. Alternatively, one may isolate DNA sequences which encode a monoclonal antibody or a binding fragment thereof by screening a DNA library from human B cells according to the general protocol outlined by Huse et al., *Science*, 246:1275–1281 (1989).

Monoclonal antibodies and polyclonal sera are collected and titered against the immunogen protein in an immunoassay, for example, a solid phase immunoassay with the immunogen immobilized on a solid support. Typically, polyclonal antisera with a titer of 10⁴ or greater are selected and tested for their cross reactivity against non-T1R polypeptides, or even other T1R family members or other related proteins from other organisms, using a competitive binding immunoassay. Specific polyclonal antisera and monoclonal antibodies will usually bind with a K_d of at least about 0.1 mM, more usually at least about 1 pM, optionally at least about 0.1 pM or better, and optionally 0.01 pM or better.

Once T1R family member specific antibodies are available, individual T1R proteins and protein fragments can be detected by a variety of immunoassay methods. For a review of immunological and immunoassay procedures, see *Basic and Clinical Immunology* (Stites & Terr eds., 7th ed. 1991). Moreover, the immunoassays of the present invention can be performed in any of several configurations, which are reviewed extensively in *Enzyme Immunoassay* (Maggio, ed., 1980); and Harlow & Lane, *supra*.

2. Immunological Binding Assays

T1R proteins, fragments, and variants can be detected and/or quantified using any of a number of well recognized immunological binding assays (see, e.g., U.S. Pat. Nos. 4,366,241; 4,376,110; 4,517,288; and 4,837,168). For a review of the general immunoassays, see also *Methods in Cell Biology: Antibodies in Cell Biology*, volume 37 (Asai, ed. 1993); *Basic and Clinical Immunology* (Stites & Terr, eds., 7th ed. 1991). Immunological binding assays (or immunoassays) typically use an antibody that specifically binds to a protein or antigen of choice (in this case a T1R family member or an antigenic subsequence thereof). The antibody (e.g., anti-T1R) may be produced by any of a number of means well known to those of skill in the art and as described above.

Immunoassays also often use a labeling agent to specifically bind to and label the complex formed by the antibody and antigen. The labeling agent may itself be one of the moieties comprising the antibody/antigen complex. Thus, the labeling agent may be a labeled T1R polypeptide or a labeled anti-T1R antibody. Alternatively, the labeling agent may be a third moiety, such as a secondary antibody, that specifically binds to the antibody/T1R complex (a secondary antibody is typically specific to antibodies of the species from which the first antibody is derived). Other proteins capable of specifically binding immunoglobulin constant regions, such as protein A or protein G may also be used as the label agent. These proteins exhibit a strong non-immunogenic reactivity with immunoglobulin constant regions from a variety of species (see, e.g., Kronval et al., *J. Immunol.*, 111:1401–1406 (1973); Akerstrom et al., *J. Immunol.*, 135:2589–2542 (1985)). The labeling agent can be modified with a detectable moiety, such as biotin, to which another molecule can specifically bind, such as streptavidin. A variety of detectable moieties are well known to those skilled in the art.

Throughout the assays, incubation and/or washing steps may be required after each combination of reagents. Incu-

bation steps can vary from about 5 seconds to several hours, optionally from about 5 minutes to about 24 hours. However, the incubation time will depend upon the assay format, antigen, volume of solution, concentrations, and the like. Usually, the assays will be carried out at ambient temperature, although they can be conducted over a range of temperatures, such as 10° C. to 40° C.

a. Non-competitive Assay Formats

Immunoassays for detecting a T1R polypeptide in a sample may be either competitive or noncompetitive. Non-competitive immunoassays are assays in which the amount of antigen is directly measured. In one preferred "sandwich" assay, for example, the anti-T1R antibodies can be bound directly to a solid substrate on which they are immobilized. These immobilized antibodies then capture the T1R polypeptide present in the test sample. The T1R polypeptide is thus immobilized is then bound by a labeling agent, such as a second T1R antibody bearing a label. Alternatively, the second antibody may lack a label, but it may, in turn, be bound by a labeled third antibody specific to antibodies of the species from which the second antibody is derived. The second or third antibody is typically modified with a detectable moiety, such as biotin, to which another molecule specifically binds, e.g., streptavidin, to provide a detectable moiety.

b. Competitive Assay Formats

In competitive assays, the amount of T1R polypeptide present in the sample is measured indirectly by measuring the amount of a known, added (exogenous) T1R polypeptide displaced (competed away) from an anti-T1R antibody by the unknown T1R polypeptide present in a sample. In one competitive assay, a known amount of T1R polypeptide is added to a sample and the sample is then contacted with an antibody that specifically binds to the T1R. The amount of exogenous T1R polypeptide bound to the antibody is inversely proportional to the concentration of T1R polypeptide present in the sample. In a particularly preferred embodiment, the antibody is immobilized on a solid substrate. The amount of T1R polypeptide bound to the antibody may be determined either by measuring the amount of T1R polypeptide present in a T1R/antibody complex, or alternatively by measuring the amount of remaining uncomplexed protein. The amount of T1R polypeptide may be detected by providing a labeled T1R molecule.

A hapten inhibition assay is another preferred competitive assay. In this assay the known T1R polypeptide is immobilized on a solid substrate. A known amount of anti-T1R antibody is added to the sample, and the sample is then contacted with the immobilized T1R. The amount of anti-T1R antibody bound to the known immobilized T1R polypeptide is inversely proportional to the amount of T1R polypeptide present in the sample. Again, the amount of immobilized antibody may be detected by detecting either the immobilized fraction of antibody or the fraction of the antibody that remains in solution. Detection may be direct where the antibody is labeled or indirect by the subsequent addition of a labeled moiety that specifically binds to the antibody as described above.

c. Cross-reactivity Determinations

Immunoassays in the competitive binding format can also be used for cross-reactivity determinations. For example, a protein at least partially encoded by the nucleic acid sequences disclosed herein can be immobilized to a solid support. Proteins (e.g., T1R polypeptides and homologs) are added to the assay that compete for binding of the antisera to the immobilized antigen. The ability of the added proteins to compete for binding of the antisera to the immobilized

protein is compared to the ability of the T1R polypeptide encoded by the nucleic acid sequences disclosed herein to compete with itself. The percent cross-reactivity for the above proteins is calculated, using standard calculations. Those antisera with less than 10% cross-reactivity with each of the added proteins listed above are selected and pooled. The cross-reacting antibodies are optionally removed from the pooled antisera by immunoabsorption with the added considered proteins, e.g., distantly related homologs. In addition, peptides comprising amino acid sequences representing conserved motifs that are used to identify members of the T1R family can be used in cross-reactivity determinations.

The immunoabsorbed and pooled antisera are then used in a competitive binding immunoassay as described above to compare a second protein, thought to be perhaps an allele or polymorphic variant of a T1R family member, to the immunogen protein (i.e., T1R polypeptide encoded by the nucleic acid sequences disclosed herein). In order to make this comparison, the two proteins are each assayed at a wide range of concentrations and the amount of each protein required to inhibit 50% of the binding of the antisera to the immobilized protein is determined. If the amount of the second protein required to inhibit 50% of binding is less than 10 times the amount of the protein encoded by nucleic acid sequences disclosed herein required to inhibit 50% of binding, then the second protein is said to specifically bind to the polyclonal antibodies generated to a T1R immunogen.

Antibodies raised against T1R conserved motifs can also be used to prepare antibodies that specifically bind only to GPCRs of the T1R family, but not to GPCRs from other families.

Polyclonal antibodies that specifically bind to a particular member of the T1R family can be made by subtracting out cross-reactive antibodies using other T1R family members. Species-specific polyclonal antibodies can be made in a similar way. For example, antibodies specific to human T1R1 can be made by, subtracting out antibodies that are cross-reactive with orthologous sequences, e.g., rat T1R1 or mouse T1R1.

d. Other Assay Formats

Western blot (immunoblot) analysis is used to detect and quantify the presence of T1R polypeptide in the sample. The technique generally comprises separating sample proteins by gel electrophoresis on the basis of molecular weight, transferring the separated proteins to a suitable solid support, (such as a nitrocellulose filter, a nylon filter, or derivatized nylon filter), and incubating the sample with the antibodies that specifically bind the T1R polypeptide. The anti-T1R polypeptide antibodies specifically bind to the T1R polypeptide on the solid support. These antibodies may be directly labeled or alternatively may be subsequently detected using labeled antibodies (e.g., labeled sheep anti-mouse antibodies) that specifically bind to the anti-T1R antibodies.

Other, assay formats include liposome immunoassays (LIA), which use liposomes designed to bind specific molecules (e.g., antibodies) and release encapsulated reagents or markers. The released chemicals are then detected according to standard techniques (see Monroe et al., *Amer. Clin. Prod. Rev.*, 5:34-41 (1986)).

e. Reduction of Non-specific Binding

One of skill in the art will appreciate that it is often desirable to minimize non-specific binding in immunoassays. Particularly, where the assay involves an antigen or antibody immobilized on a solid substrate it is desirable to minimize the amount of non-specific binding to the sub-

strate. Means of reducing such non-specific binding are well known to those of skill in the art. Typically, this technique involves coating the substrate with a proteinaceous composition. In particular, protein compositions such as bovine serum albumin (BSA), nonfat powdered milk, and gelatin are widely used with powdered milk being most preferred.

f. Labels

The particular label or detectable group used in the assay is not a critical aspect of the invention, as long as it does not significantly interfere with the specific binding of the antibody used in the assay. The detectable group can be any material having a detectable physical or chemical property. Such detectable labels have been well developed in the field of immunoassays and, in general, most any label useful in such methods can be applied to the present invention. Thus, a label is any composition detectable by spectroscopic, photochemical, biochemical, immunochemical, electrical, optical, or chemical means. Useful labels in the present invention include magnetic beads (e.g., DYNABEADSTM), fluorescent dyes (e.g., fluorescein isothiocyanate, Texas red, rhodamine, and the like), radiolabels (e.g., ³H, ¹²⁵I, ³S, ¹⁴C, or ³²P), enzymes (e.g., horseradish peroxidase, alkaline phosphatase and others commonly used in an ELISA), and colorimetric labels such as colloidal gold or colored glass or plastic beads (e.g., polystyrene, polypropylene, latex, etc.).

The label may be coupled directly or indirectly to the desired component of the assay according to methods well known in the art. As indicated above, a wide variety of labels may be used, with the choice of label depending on sensitivity required, ease of conjugation with the compound, stability requirements, available instrumentation, and disposal provisions.

Non-radioactive labels are often attached by indirect means. Generally, a ligand molecule (e.g., biotin) is covalently bound to the molecule. The ligand then binds to another molecules (e.g., streptavidin) molecule, which is either inherently detectable or covalently bound to a signal system, such as a detectable enzyme, a fluorescent compound, or a chemiluminescent compound. The ligands and their targets can be used in any suitable combination with antibodies that recognize a T1R polypeptide, or secondary antibodies that recognize anti-T1R.

The molecules can also be conjugated directly to signal generating compounds, e.g., by conjugation with an enzyme or fluorophore. Enzymes of interest as labels will primarily be hydrolases, particularly phosphatases, esterases and glycosidases, or oxidotases, particularly peroxidases. Fluorescent compounds include fluorescein and its derivatives, rhodamine and its derivatives, dansyl, umbelliferone, etc. Chemiluminescent compounds include luciferin, and 2,3-dihydrophthalazinediones, e.g., luminol. For a review of various labeling or signal producing systems that may be used, see U.S. Pat. No. 4,391,904.

Means of detecting labels are well known to those of skill in the art. Thus, for example, where the label is a radioactive label, means for detection include a scintillation counter or photographic film as in autoradiography. Where the label is a fluorescent label, it may be detected by exciting the fluorochrome with the appropriate wavelength of light and detecting the resulting fluorescence. The fluorescence may be detected visually, by means of photographic film, by the use of electronic detectors such as charge-coupled devices (CCDs) or photomultipliers and the like. Similarly, enzymatic labels may be detected by providing the appropriate substrates for the enzyme and detecting the resulting reaction product. Finally simple calorimetric labels may be detected simply by observing the color associated with the

label. Thus, in various dipstick assays, conjugated gold often appears pink, while various conjugated beads appear the color of the bead.

Some assay formats do not require the use of labeled components. For instance, agglutination assays can be used to detect the presence of the target antibodies. In this case, antigen-coated particles are agglutinated by samples comprising the target antibodies. In this format, none of the components need be labeled and the presence of the target antibody is detected by simple visual inspection.

C. Detection of Modulators

Compositions and methods for determining whether a test compound specifically binds to a chemosensory receptor of the invention, both in vitro and in vivo, are described below. Many aspects of cell physiology can be monitored to assess the effect of ligand binding to a T1R polypeptide of the invention. These assays may be performed on intact cells expressing a chemosensory receptor, on permeabilized cells, or on membrane fractions produced by standard methods.

Taste receptors bind taste stimuli and initiate the transduction of chemical stimuli into electrical signals. An activated or inhibited G protein will in turn alter the properties of target enzymes, channels, and other effector proteins. Some examples are the activation of cGMP phosphodiesterase by transducin in the visual system, adenylate cyclase by the stimulatory G protein, phospholipase C by Gq and other cognate G proteins, and modulation of diverse channels by Gi and other G proteins. Downstream consequences can also be examined such as generation of diacyl glycerol and IP3 by phospholipase C, and in turn, for calcium mobilization by IP3.

The T1R proteins or polypeptides of the assay will typically be selected from a polypeptide having a sequence of SEQ ID NOS: 4, 10, 12, 14, 17, 21 or fragments or conservatively modified variants thereof. Optionally, the fragments and variants can be antigenic fragments and variants which bind to an anti-T1R antibody.

Alternatively, the T1R proteins or polypeptides of the assay can be derived from a eukaryote host cell and can include an amino acid subsequence having amino acid sequence identity to SEQ ID NOS: 4, 10, 12, 14, 17, 21, or fragments or conservatively modified variants thereof. Generally, the amino acid sequence identity will be at least 35 to 50%, or optionally 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99%. Optionally, the T1R proteins or polypeptides of the assays can comprise a domain of a T1R protein, such as an extracellular domain, transmembrane region, transmembrane domain, cytoplasmic domain, ligand-binding domain, and the like. Further, as described above, the T1R protein or a domain thereof can be covalently linked to a heterologous protein to create a chimeric protein used in the assays described herein.

Modulators of T1R receptor activity are tested using T1R proteins or polypeptides as described above, either recombinant or naturally occurring. The T1R proteins or polypeptides can be isolated, expressed in a cell, expressed in a membrane derived from a cell, expressed in tissue or in an animal, either recombinant or naturally occurring. For example, tongue slices, dissociated cells from a tongue, transformed cells, or membranes can be used. Modulation can be tested using one of the in vitro or in vivo assays described herein.

1. In Vitro Binding Assays

Taste transduction can also be examined in vitro with soluble or solid state reactions, using a T1R polypeptide of the invention. In a particular embodiment, a T1R ligand-

binding domain can be used in vitro in soluble or solid state reactions to assay for ligand binding.

For instance, the T1R N-terminal domain is predicted to be involved in ligand binding. More particularly, the T1Rs belong to a GPCR sub-family that is characterized by large, approximately 600 amino acid, extracellular N-terminal segments. These N-terminal segments are thought to form, at least in part, the ligand-binding domains, and are therefore useful in biochemical assays to identify T1R agonists and antagonists. The ligand-binding domain may also contain additional portions of the extracellular domain, such as the extracellular loops of the transmembrane domain. Similar assays have been used with other GPCRs that are related to the T1Rs, such as the metabotropic glutamate receptors (see, e.g., Han and Hampson, *J. Biol. Chem.* 274:10008–10013 (1999)). These assays might involve displacing a radioactively or fluorescently labeled ligand, measuring changes in intrinsic fluorescence or changes in proteolytic susceptibility, etc.

Ligand binding to a T1R polypeptide of the invention can be tested in solution, in a bilayer membrane, optionally attached to a solid phase, in a lipid monolayer, or in vesicles. Binding of a modulator can be tested using, e.g., changes in spectroscopic characteristics (e.g., fluorescence, absorbance, refractive index) hydrodynamic (e.g., shape), chromatographic, or solubility properties. Preferred binding assays of the invention are biochemical binding assays that use recombinant soluble N-terminal T1R domains.

Receptor-G protein interactions can also be examined. For example, binding of the G protein to the receptor, or its release from the receptor can be examined. More particularly, in the absence of GTP, an activator will lead to the formation of a tight complex of a G protein (all three subunits) with the receptor. This complex can be detected in a variety of ways, as noted above. Such an assay can be modified to search for inhibitors, e.g., by adding an activator to the receptor and G protein in the absence of GTP, which form a tight complex, and then screen for inhibitors by looking at dissociation of the receptor-G protein complex. In the presence of GTP, release of the alpha subunit of the G protein from the other two G protein subunits serves as a criterion of activation. An activated or inhibited G protein will in turn alter the properties of target enzymes, channels, and other effector proteins.

In another embodiment of the invention, a GTP γ S assay may be used. As described above, upon activation of a GPCR, the G α subunit of the G protein complex is stimulated to exchange bound GDP for GTP. Ligand-mediated stimulation of G protein exchange activity can be measured in a biochemical assay measuring the binding of added radioactively-labeled GTP γ ³⁵S to the G protein in the presence of a putative ligand. Typically, membranes containing the chemosensory receptor of interest are mixed with a complex of G proteins. Potential inhibitors and/or activators and GTP γ S are added to the assay, and binding of GTP γ S to the G protein is measured. Binding can be measured by liquid scintillation counting or by any other means known in the art, including scintillation proximity assays (SPA). In other assays formats, fluorescently-labeled GTP γ S can be utilized.

2. Fluorescence Polarization Assays

In another embodiment, Fluorescence Polarization (“FP”) based assays may be used to detect and monitor ligand binding. Fluorescence polarization is a versatile laboratory technique for measuring equilibrium binding, nucleic acid hybridization, and enzymatic activity. Fluorescence polarization assays are homogeneous in that they do not require

a separation step such as centrifugation, filtration, chromatography, precipitation, or electrophoresis. These assays are done in real time, directly in solution and do not require an immobilized phase. Polarization values can be measured repeatedly and after the addition of reagents since measuring the polarization is rapid and does not destroy the sample. Generally, this technique can be used to measure polarization values of fluorophores from low picomolar to micromolar levels. This section describes how fluorescence polarization can be used in a simple and quantitative way to measure the binding of ligands to the T1R polypeptides of the invention.

When a fluorescently labeled molecule is excited with plane polarized light, it emits light that has a degree of polarization that is inversely proportional to its molecular rotation. Large fluorescently labeled molecules remain relatively stationary during the excited state (4 nanoseconds in the case of fluorescein) and the polarization of the light remains relatively constant between excitation and emission. Small fluorescently labeled molecules rotate rapidly during the excited state and the polarization changes significantly between excitation and emission. Therefore, small molecules have low polarization values and large molecules have high polarization values. For example, a single-stranded fluorescein-labeled oligonucleotide has a relatively low polarization value but when it is hybridized to a complementary strand, it has a higher polarization value. When using FP to detect and monitor taste stimulus-binding which may activate or inhibit the chemosensory receptors of the invention, fluorescence-labeled taste stimuli or auto-fluorescent taste stimuli may be used.

Fluorescence polarization (P) is defined as:

$$P = \frac{Int_{\parallel} - Int_{\perp}}{Int_{\parallel} + Int_{\perp}}$$

Where \parallel is the intensity of the emission light parallel to the excitation light plane and \perp is the intensity of the emission light perpendicular to the excitation light plane. P, being a ratio of light intensities, is a dimensionless number. For example, the Beacon® and Beacon 2000™ System may be used in connection with these assays. Such systems typically express polarization in millipolarization units (1 Polarization Unit=1000 mP Units).

The relationship between molecular rotation and size is described by the Perrin equation and the reader is referred to Jolley, M. E. (1991) in *Journal of Analytical Toxicology*, pp. 236–240, which gives a thorough explanation of this equation. Summarily, the Perrin equation states that polarization is directly proportional to the rotational relaxation time, the time that it takes a molecule to rotate through an angle of approximately 68.5°. Rotational relaxation time is related to viscosity (η), absolute temperature (T), molecular volume (V), and the gas constant (R) by the following equation:

$$RotationalRelaxationTime = \frac{3\eta V}{RT}$$

The rotational relaxation time is small (\approx 1 nanosecond) for small molecules (e.g. fluorescein) and large (\approx 100 nanoseconds) for large molecules (e.g. immunoglobulins). If viscosity and temperature are held constant, rotational relaxation time, and therefore polarization, is directly related to

the molecular volume. Changes in molecular volume may be due to interactions with other molecules, dissociation, polymerization, degradation, hybridization, or conformational changes of the fluorescently labeled molecule. For example, fluorescence polarization has been used to measure enzymatic cleavage of large fluorescein labeled polymers by proteases, DNases, and RNases. It also has been used to measure equilibrium binding for protein/protein interactions, antibody/antigen binding, and protein/DNA binding.

3. Solid State and Soluble High Throughput Assays

In yet another embodiment, the invention provides soluble assays using a T1R polypeptide; or a cell or tissue expressing an T1R polypeptide. In another embodiment, the invention provides solid phase based in vitro assays in a high throughput format, where the T1R polypeptide, or cell or tissue expressing the T1R polypeptide is attached to a solid phase substrate.

In the high throughput assays of the invention, it is possible to screen up to several thousand different modulators or ligands in a single day. In particular, each well of a microtiter plate can be used to run a separate assay against a selected potential modulator, or, if concentration or incubation time effects are to be observed, every 5–10 wells can test a single modulator. Thus, a single standard microtiter plate can assay about 100 (e.g., 96) modulators. If 1536 well plates are used, then a single plate can easily assay from about 1000 to about 1500 different compounds. It is also possible to assay multiple compounds in each plate well. It is possible to assay several different plates per day; assay screens for up to about 6,000–20,000 different compounds is possible using the integrated systems of the invention. More recently, microfluidic approaches to reagent manipulation have been developed.

The molecule of interest can be bound to the solid state component, directly or indirectly, via covalent or non-covalent linkage, e.g., via a tag. The tag can be any of a variety of components. In general, a molecule which binds the tag (a tag binder) is fixed to a solid support, and the tagged molecule of interest (e.g., the taste transduction molecule of interest) is attached to the solid support by interaction of the tag and the tag binder.

A number of tags and tag binders can be used, based upon known molecular interactions well described in the literature. For example, where a tag has a natural binder, for example, biotin, protein A, or protein G, it can be used in conjunction with appropriate tag binders (avidin, streptavidin, neutravidin, the Fc region of an immunoglobulin, etc.). Antibodies to molecules with natural binders such as biotin are also widely available and appropriate tag binders (see, SIGMA Immunochemicals 1998 catalogue SIGMA, St. Louis Mo.).

Similarly, any haptenic or antigenic compound can be used in combination with an appropriate antibody to form a tag/tag binder pair. Thousands of specific antibodies are commercially available and many additional antibodies are described in the literature. For example, in one common configuration, the tag is a first antibody and the tag binder is a second antibody which recognizes the first antibody. In addition to antibody-antigen interactions, receptor-ligand interactions are also appropriate as tag and tag-binder pairs. For example, agonists and antagonists of cell membrane receptors (e.g., cell receptor-ligand interactions such as transferrin, c-kit, viral receptor ligands, cytokine receptors, chemokine receptors, interleukin receptors, immunoglobulin receptors and antibodies, the cadherin family, the integrin family, the selectin family, and the like; see, e.g., Pigott & Power, *The Adhesion Molecule Facts Book I* (1993)).

Similarly, toxins and venoms, viral epitopes, hormones (e.g., opiates, steroids, etc.), intracellular receptors (e.g., which mediate the effects of various small ligands, including steroids, thyroid hormone, retinoids and vitamin D; peptides), drugs, lectins, sugars, nucleic acids (both linear and cyclic polymer configurations), oligosaccharides, proteins, phospholipids and antibodies can all interact with various cell receptors.

Synthetic polymers, such as polyurethanes, polyesters, polycarbonates, polyureas, polyamides, polyethyleneimines, polyarylene sulfides, polysiloxanes, polyimides, and polyacetates can also form an appropriate tag or tag binder. Many other tag/tag binder pairs are also useful in assay systems described herein, as would be apparent to one of skill upon review of this disclosure.

Common linkers such as peptides, polyethers, and the like can also serve as tags, and include polypeptide sequences, such as poly gly sequences of between about 5 and 200 amino acids. Such flexible linkers are known to persons of skill in the art. For example, poly(ethylene glycol) linkers are available from Shearwater Polymers, Inc. Huntsville, Alabama. These linkers optionally have amide linkages, sulfhydryl linkages, or heterofunctional linkages.

Tag binders are fixed to solid substrates using any of a variety of methods currently available. Solid substrates are commonly derivatized or functionalized by exposing all or a portion of the substrate to a chemical reagent which fixes a chemical group to the surface which is reactive with a portion of the tag binder. For example, groups which are suitable for attachment to a longer chain portion would include amines, hydroxyl, thiol, and carboxyl groups. Aminoalkylsilanes and hydroxyalkylsilanes can be used to functionalize a variety of surfaces, such as glass surfaces. The construction of such solid phase biopolymer arrays is well described in the literature. See, e.g., Merrifield, *J. Am. Chem. Soc.*, 85:2149–2154 (1963) (describing solid phase synthesis of, e.g., peptides); Geysen et al., *J. Immun. Meth.*, 102:259–274 (1987) (describing synthesis of solid phase components on pins); Frank & Doring, *Tetrahedron*, 44:60316040 (1988) (describing synthesis of various peptide sequences on cellulose disks); Fodor et al., *Science*, 251:767–777 (1991); Sheldon et al., *Clinical Chemistry*, 39(4):718–719 (1993); and Kozal et al., *Nature Medicine*, 2(7):753759 (1996) (all describing arrays of biopolymers fixed to solid substrates). Non-chemical approaches for fixing tag binders to substrates include other common methods, such as heat, cross-linking by UV radiation, and the like.

4. Computer-based Assays

Yet another assay for compounds that modulate T1R polypeptide activity involves computer assisted compound design, in which a computer system is used to generate a three-dimensional structure of an T1R polypeptide based on the structural information encoded by its amino acid sequence. The input amino acid sequence interacts directly and actively with a preestablished algorithm in a computer program to yield secondary, tertiary, and quaternary structural models of the protein. The models of the protein structure are then examined to identify regions of the structure that have the ability to bind, e.g., ligands. These regions are then used to identify ligands that bind to the protein.

The three-dimensional structural model of the protein is generated by entering protein amino acid sequences of at least 10 amino acid residues or corresponding nucleic acid sequences encoding a T1R polypeptide into the computer system. The nucleotide sequence encoding the T1R polypep-

tion, or the amino acid sequence thereof, can be any sequence disclosed herein, and conservatively modified versions thereof.

The amino acid sequence represents the primary sequence or subsequence of the protein, which encodes the structural information of the protein. At least 10 residues of the amino acid sequence (or a nucleotide sequence encoding 10 amino acids) are entered into the computer system from computer keyboards, computer readable substrates that include, but are not limited to, electronic storage media (e.g., magnetic diskettes, tapes, cartridges, and chips), optical media (e.g., CD ROM), information distributed by internet sites, and by RAM. The three-dimensional structural model of the protein is then generated by the interaction of the amino acid sequence and the computer system, using software known to those of skill in the art.

The amino acid sequence represents a primary structure that encodes the information necessary to form the secondary, tertiary and quaternary structure of the protein of interest. The software looks at certain parameters encoded by the primary sequence to generate the structural model. These parameters are referred to as "energy terms," and primarily include electrostatic potentials, hydrophobic potentials, solvent accessible surfaces, and hydrogen bonding. Secondary energy terms include van der Waals potentials. Biological molecules form the structures that minimize the energy terms in a cumulative fashion. The computer program is therefore using these terms encoded by the primary structure or amino acid sequence to create the secondary structural model.

The tertiary structure of the protein encoded by the secondary structure is then formed on the basis of the energy terms of the secondary structure. The user at this point can enter additional variables such as whether the protein is membrane bound or soluble, its location in the body, and its cellular location, e.g., cytoplasmic, surface, or nuclear. These variables along with the energy terms of the secondary structure are used to form the model of the tertiary structure. In modeling the tertiary structure, the computer program matches hydrophobic faces of secondary structure with like, and hydrophilic faces of secondary structure with like.

Once the structure has been generated, potential ligand-binding regions are identified by the computer system. Three-dimensional structures for potential ligands are generated by entering amino acid or nucleotide sequences or chemical formulas of compounds, as described above. The three-dimensional structure of the potential ligand is then compared to that of the T1R polypeptide to identify ligands that bind to the protein. Binding affinity between the protein and ligands is determined using energy terms to determine which ligands have an enhanced probability of binding to the protein.

Computer systems are also used to screen for mutations, polymorphic variants, alleles, and interspecies homologs of T1R genes. Such mutations can be associated with disease states or genetic traits. As described above, GeneChip™ and related technology can also be used to screen for mutations, polymorphic variants, alleles, and interspecies homologs. Once the variants are identified, diagnostic assays can be used to identify patients having such mutated genes. Identification of the mutated T1R genes involves receiving input of a first nucleic acid or amino acid sequence of a T1R gene, or conservatively modified versions thereof. The sequence is entered into the computer system as described above. The first nucleic acid or amino acid sequence is then compared to a second nucleic acid or amino acid sequence that has

substantial identity to the first sequence. The second sequence is entered into the computer system in the manner described above. Once the first and second sequences are compared, nucleotide or amino acid differences between the sequences are identified. Such sequences can represent allelic differences in various T1R genes, and mutations associated with disease states and genetic traits.

5. Cell-based Binding Assays

In one embodiment, a T1R protein or polypeptide is expressed in a eukaryotic cell as a chimeric receptor with a heterologous, chaperone sequence that facilitates its maturation and targeting through the secretory pathway. Such chimeric T1R polypeptides can be expressed in any eukaryotic cell, such as HEK-293 cells. Preferably, the cells comprise a functional G protein, e.g., G α I5, that is capable of coupling the chimeric receptor to an intracellular signaling pathway or to a signaling protein such as phospholipase C. Activation of such chimeric receptors in such cells can be detected using any standard method, such as by detecting changes in intracellular calcium by detecting FURA-2 dependent fluorescence in the cell.

Activated GPCR receptors become substrates for kinases that phosphorylate the C-terminal tail of the receptor (and possibly other sites as well). Thus, activators will promote the transfer of 32P from gamma-labeled GTP to the receptor, which can be assayed with a scintillation counter. The phosphorylation of the C-terminal tail will promote the binding of arrestin-like proteins and will interfere with the binding of G proteins. The kinase/arrestin pathway plays a key role in the desensitization of many GPCR receptors. For example, compounds that modulate the duration a taste receptor stays active would be useful as a means of prolonging a desired taste or cuffing off an unpleasant one. For a general review of GPCR signal transduction and methods of assaying signal transduction, see, e.g., *Methods in Enzymology*, vols. 237 and 238 (1994) and volume 96 (1983); Bourne et al., *Nature*, 10:349:117-27 (1991); Bourne et al., *Nature*, 348:125-32 (1990); Pitcher et al., *Annu. Rev. Biochem.*, 67:653-92 (1998).

T1R modulation may be assayed by comparing the response of a T1R polypeptide treated with a putative T1R modulator to the response of an untreated control sample. Such putative T1R modulators can include taste stimuli that either inhibit or activate T1R polypeptide activity. In one embodiment, control samples (untreated with activators or inhibitors) are assigned a relative T1R activity value of 100. Inhibition of a T1R polypeptide is achieved when the T1R activity value relative to the control is about 90%, optionally 50%, optionally 25-0%. Activation of a T1R polypeptide is achieved when the T1R activity value relative to the control is 110%, optionally 150%, 200-500%, or 1000-2000%.

Changes in ion flux may be assessed by determining changes in ionic polarization (i.e., electrical potential) of the cell or membrane expressing a T1R polypeptide. One means to determine changes in cellular polarization is by measuring changes in current (thereby measuring changes in polarization) with voltage-clamp and patch-clamp techniques (see, e.g., the "cell-attached" mode, the "inside-out" mode, and the "whole cell" mode, e.g., Ackerman et al., *New Engl. J. Med.*, 336:1575-1595 (1997)). Whole cell currents are conveniently determined using the standard. Other known assays include: radiolabeled ion flux assays and fluorescence assays using voltage-sensitive dyes (see, e.g., Vestergaard-Bogind et al., *J. Membrane Biol.*, 88:67-75 (1988); Gonzalez & Tsien, *Chem. Biol.*, 4:269-277 (1997); Daniel et al., *J. Pharmacol. Meth.*, 25:185-193 (1991); Holevinsky et al., *J.*

Membrane Biology, 137:59–70 (1994)). Generally, the compounds to be tested are present in the range from 1 pM to 100 mM.

The effects of the test compounds upon the function of the polypeptides can be measured by examining any of the parameters described above. Any suitable physiological change that affects GPCR activity can be used to assess the influence of a test compound on the polypeptides of this invention. When the functional consequences are determined using intact cells or animals, one can also measure a variety of effects such as transmitter release, hormone release, transcriptional changes to both known and uncharacterized genetic markers (e.g., northern blots), changes in cell metabolism such as cell growth or pH changes, and changes in intracellular second messengers such as Ca^{2+} , IP₃, cGMP, or cAMP.

Preferred assays for GPCRs include cells that are loaded with ion or voltage sensitive dyes to report receptor activity. Assays for determining activity of such receptors can also use known agonists and antagonists for other G protein-coupled receptors as negative or positive controls to assess activity of tested compounds. In assays for identifying modulatory compounds (e.g., agonists, antagonists), changes in the level of ions in the cytoplasm or membrane voltage will be monitored using an ion sensitive or membrane voltage fluorescent indicator, respectively. Among the ion-sensitive indicators and voltage probes that may be employed are those disclosed in the Molecular Probes 1997 Catalog. For G protein-coupled receptors, promiscuous G proteins such as G α 15 and G α 16 can be used in the assay of choice (Wilkie et al., *Proc. Nat'l Acad. Sci.*, 88:10049–10053 (1991)). Such promiscuous G proteins allow coupling of a wide range of receptors.

Receptor activation typically initiates subsequent intracellular events, e.g., increases in second messengers such as IP₃, which releases intracellular stores of calcium ions. Activation of some G protein-coupled receptors stimulates the formation of inositol triphosphate (IP₃) through phospholipase C-mediated hydrolysis of phosphatidylinositol (Berridge & Irvine, *Nature*, 312:315–21 (1984)). IP₃ in turn stimulates the release of intracellular calcium ion stores. Thus, a change in cytoplasmic calcium ion levels, or a change in second messenger levels such as IP₃ can be used to assess G protein-coupled receptor function. Cells expressing such G protein-coupled receptors may exhibit increased cytoplasmic calcium levels as a result of contribution from both intracellular stores and via activation of ion channels, in which case it may be desirable although not necessary to conduct such assays in calcium-free buffer, optionally supplemented with a chelating agent such as EGTA, to distinguish fluorescence response resulting from calcium release from internal stores.

Other assays can involve determining the activity of receptors which, when activated, result in a change in the level of intracellular cyclic nucleotides, e.g., cAMP or cGMP, by activating or inhibiting enzymes such as adenylyl cyclase. There are cyclic nucleotide-gated ion channels, e.g., rod photoreceptor cell channels and olfactory neuron channels that are permeable to cations upon activation by binding of cAMP or cGMP (see, e.g., Altenhofen et al., *Proc. Nat'l Acad. Sci.*, 88:9868–9872 (1991) and Dhallan et al., *Nature*, 347:184–187 (1990)). In cases where activation of the receptor results in a decrease in cyclic nucleotide levels, it may be preferable to expose the cells to agents that increase intracellular cyclic nucleotide levels, e.g., forskolin, prior to adding a receptor-activating compound to the cells in the assay. Cells for this type of assay can be made by

co-transfection of a host cell with DNA encoding a cyclic nucleotide-gated ion channel, GPCR phosphatase and DNA encoding a receptor (e.g., certain glutamate receptors, muscarinic acetylcholine receptors, dopamine receptors, serotonin receptors, and the like), which, when activated, causes a change in cyclic nucleotide levels in the cytoplasm.

In a preferred embodiment, T1R polypeptide activity is measured by expressing a T1R gene in a heterologous cell with a promiscuous G protein that links the receptor to a phospholipase C signal transduction pathway (see Offermanns & Simon, *J. Biol. Chem.*, 270:15175–15180 (1995)). Optionally the cell line is HEK-293 (which does not naturally express T1R genes) and the promiscuous G protein is G α 15 (Offermanns & Simon, *supra*). Modulation of taste transduction is assayed by measuring changes in intracellular Ca^{2+} levels, which change in response to modulation of the T1R signal transduction pathway via administration of a molecule that associates with a T1R polypeptide. Changes in Ca^{2+} levels are optionally measured using fluorescent Ca^{2+} indicator dyes and fluorometric imaging.

In one embodiment, the changes in intracellular cAMP or cGMP can be measured using immunoassays. The method described in Offermanns & Simon, *J. Bio. Chem.*, 270:15175–15180 (1995), may be used to determine the level of cAMP. Also, the method described in Felley-Bosco et al., *Am. J. Resp. Cell and Mol. Biol.*, 11:159–164 (1994), may be used to determine the level of cGMP. Further, an assay kit for measuring cAMP and/or cGMP is described in U.S. Pat. No. 4,115,538, herein incorporated by reference.

In another embodiment, phosphatidyl inositol (PI) hydrolysis can be analyzed according to U.S. Pat. No. 5,436,128, herein incorporated by reference. Briefly, the assay involves labeling of cells with 3H-myoinositol for 48 or more hrs. The labeled cells are treated with a test compound for one hour. The treated cells are lysed and extracted in chloroform-methanol-water after which the inositol phosphates were separated by ion exchange chromatography and quantified by scintillation counting. Fold stimulation is determined by calculating the ratio of cpm in the presence of agonist, to cpm in the presence of buffer control. Likewise, fold inhibition is determined by calculating the ratio of cpm in the presence of antagonist, to cpm in the presence of buffer control (which may or may not contain an agonist).

In another embodiment, transcription levels can be measured to assess the effects of a test compound on signal transduction. A host cell containing a T1R polypeptide of interest is contacted with a test compound for a sufficient time to effect any interactions, and then the level of gene expression is measured. The amount of time to effect such interactions may be empirically determined, such as by running a time course and measuring the level of transcription as a function of time. The amount of transcription may be measured by using any method known to those of skill in the art to be suitable. For example, mRNA expression of the protein of interest may be detected using northern blots or their polypeptide products may be identified using immunoassays. Alternatively, transcription based assays using reporter gene may be used as described in U.S. Pat. 5,436,128, herein incorporated by reference. The reporter genes can be, e.g., chloramphenicol acetyltransferase, luciferase, β -galactosidase and alkaline phosphatase. Furthermore, the protein of interest can be used as an indirect reporter via attachment to a second reporter such as green fluorescent protein (see, e.g., Mistili & Spector, *Nature Biotechnology*, 15:961–964 (1997)).

The amount of transcription is then compared to the amount of transcription in either the same cell in the absence of the test compound, or it may be compared with the amount of transcription in a substantially identical cell that lacks the T1R polypeptide of interest. A substantially identical cell may be derived from the same cells from which the recombinant cell was prepared but which had not been modified by introduction of heterologous DNA. Any difference in the amount of transcription indicates that the test compound has in some manner altered the activity of the T1R polypeptide of interest.

6. Transgenic Non-human Animals Expressing Chemosensory Receptors

Non-human animals expressing one or more chemosensory receptor sequences of the invention, can also be used for receptor assays. Such expression can be used to determine whether a test compound specifically binds to a mammalian taste transmembrane receptor polypeptide in vivo by contacting a non-human animal stably or transiently transfected with a nucleic acid encoding a chemosensory receptor or ligand-binding region thereof with a test compound and determining whether the animal reacts to the test compound by specifically binding to the receptor polypeptide.

Animals transfected or infected with the vectors of the invention are particularly useful for assays to identify and characterize taste stimuli/ligands that can bind to a specific or sets of receptors. Such vector-infected animals expressing human chemosensory receptor sequences can be used for in vivo screening of taste stimuli and their effect on, e.g., cell physiology (e.g., on taste neurons), on the CNS, or behavior.

Means to infect/express the nucleic acids and vectors, either individually or as libraries, are well known in the art. A variety of individual cell, organ, or whole animal parameters can be measured by a variety of means. The T1R sequences of the invention can be for example expressed in animal taste tissues by delivery with an infecting agent, e.g., adenovirus expression vector.

The endogenous chemosensory receptor genes can remain functional and wild-type (native) activity can still be present. In other situations, where it is desirable that all chemosensory receptor activity is by the introduced exogenous hybrid receptor, use of a knockout line is preferred. Methods for the construction of non-human transgenic animals, particularly transgenic mice, and the selection and preparation of recombinant constructs for generating transformed cells are well known in the art.

Construction of a "knockout" cell and animal is based on the premise that the level of expression of a particular gene in a mammalian cell can be decreased or completely abrogated by introducing into the genome a new DNA sequence that serves to interrupt some portion of the DNA sequence of the gene to be suppressed. Also, "gene trap insertion" can be used to disrupt a host gene, and mouse embryonic stem (ES) cells can be used to produce knockout transgenic animals (see, e.g., Holzschu, *Transgenic Res* 6:97-106 (1997)). The insertion of the exogenous is typically by homologous recombination between complementary nucleic acid sequences. The exogenous sequence is some portion of the target gene to be modified, such as exonic, intronic or transcriptional regulatory sequences, or any genomic sequence which is able to affect the level of the target gene's expression; or a combination thereof. Gene targeting via homologous recombination in pluripotential embryonic stem cells allows one to modify precisely the genomic sequence of interest. Any technique can be used to create, screen for, propagate, a knockout animal, e.g., see Bijvoet,

Hum. Mol. Genet. 7:53-62 (1998); Moreadith, *J. Mol. Med.* 75:208-216 (1997); Tojo, *Cytotechnology* 19:161-165 (1995); Mudgett, *Methods Mol. Biol.* 48:167-184 (1995); Longo, *Transgenic Res.* 6:321-328 (1997); U.S. Pat. Nos. 5,616,491; 5,464,764; 5,631,153; 5,487,992; 5,627,059; 5,272,071; WO 91/09955; WO93/09222; WO 96/29411; WO 95/31560; WO 91/12650.

The nucleic acids of the invention can also be used as reagents to produce "knockout" human cells and their progeny. Likewise, the nucleic acids of the invention can also be used as reagents to produce "knock-ins" in mice. The human or rat T1R gene sequences can replace the orthologous T1R in the mouse genome. In this way, a mouse expressing a human or rat T1R is produced. This mouse can then be used to analyze the function of human or rat T1Rs, and to identify ligands for such T1Rs.

D. Modulators

The compounds tested as modulators of a T1R family member can be any small chemical compound, or a biological entity, such as a protein, sugar, nucleic acid or lipid. Alternatively, modulators can be genetically altered versions of a T1R gene. Typically, test compounds will be small chemical molecules and peptides. Essentially any chemical compound can be used as a potential modulator or ligand in the assays of the invention, although most often compounds can be dissolved in aqueous or organic (especially DMSO-based) solutions are used. The assays are designed to screen large chemical libraries by automating the assay steps and providing compounds from any convenient source to assays, which are typically run in parallel (e.g., in microtiter formats on microtiter plates in robotic assays). It will be appreciated that there are many suppliers of chemical compounds, including Sigma (St. Louis, Mo.), Aldrich (St. Louis, Mo.), Sigma-Aldrich (St. Louis, Mo.), Fluka Chemika-Biochemica Analytika (Buchs, Switzerland) and the like.

In one preferred embodiment, high throughput screening methods involve providing a combinatorial chemical or peptide library containing a large number of potential therapeutic compounds (potential modulator or ligand compounds). Such "combinatorial chemical libraries" or "ligand libraries" are then screened in one or more assays, as described herein, to identify those library members (particular chemical species or subclasses) that display a desired characteristic activity. The compounds thus identified can serve as conventional "lead compounds" or can themselves be used as potential or actual therapeutics.

A combinatorial chemical library is a collection of diverse chemical compounds generated by either chemical synthesis or biological synthesis, by combining a number of chemical "building blocks" such as reagents. For example, a linear combinatorial chemical library such as a polypeptide library is formed by combining a set of chemical building blocks (amino acids) in every possible way for a given compound length (i.e., the number of amino acids in a polypeptide compound). Millions of chemical compounds can be synthesized through such combinatorial mixing of chemical building blocks.

Preparation and screening of combinatorial chemical libraries is well known to those of skill in the art. Such combinatorial chemical libraries include, but are not limited to, peptide libraries (see, e.g., U.S. Pat. No. 5,010,175, Furka, *Int. J. Pept. Prot. Res.*, 37:487-493 (1991) and Houghton et al., *Nature*, 354:84-88 (1991)). Other chemistries for generating chemical diversity libraries can also be used. Such chemistries include, but are not limited to: peptoids (e.g., PCT Publication No. WO 91/19735), encoded

peptides (e.g., PCT Publication WO 93/20242), random bio-oligomers (e.g., PCT Publication No. WO 92/00091), benzodiazepines (e.g., U.S. Pat. No. 5,288,514), diversomers such as hydantoins, benzodiazepines and dipeptides (Hobbs et al., *Proc. Nat. Acad. Sci.*, 90:6909–6913 (1993)), vinylogous polypeptides (Hagihara et al., *J. Amer. Chem. Soc.*, 114:6568 (1992)), nonpeptidal peptidomimetics with glucose scaffolding (Hirschmann et al., *J. Amer. Chem. Soc.*, 114:9217–9218 (1992)), analogous organic syntheses of small compound libraries (Chen et al., *J. Amer. Chem. Soc.*, 116:2661 (1994)), oligocarbamates (Cho et al., *Science*, 261:1303 (1993)), peptidyl phosphonates (Campbell et al., *J. Org. Chem.*, 59:658 (1994)), nucleic acid libraries (Ausubel, Berger and Sambrook, all supra), peptide nucleic acid libraries (U.S. Pat. No. 5,539,083), antibody libraries (Vaughn et al., *Nature Biotechnology*, 14(3):309–314 (1996) and PCT/US96/10287), carbohydrate libraries (Liang et al., *Science*, 274:1520–1522 (1996) and U.S. Pat. No. 5,593,853), small organic molecule libraries (benzodiazepines, Baum, *C&EN*, January 18, page 33 (1993); thiazolidinones and metathiazanones, U.S. Pat. No. 5,549,974; pynrolidines, U.S. Pat. Nos. 5,525,735 and 5,519,134; morpholino compounds, U.S. Pat. No. 5,506,337; benzodiazepines, U.S. Pat. No. 5,288,514, and the like).

Devices for the preparation of combinatorial libraries are commercially available (see, e.g., 357 MPS, 390 MPS (Advanced Chem Tech, Louisville Ky.), Symphony (Rainin, Woburn, Mass.), 433A (Applied Biosystems, Foster City, Calif.), 9050 Plus (Millipore, Bedford, Mass.)). In addition, numerous combinatorial libraries are themselves commercially available (see, e.g., ComGenex, Princeton, N.J.; Tripos, Inc., St. Louis, Mo.; 3D Pharmaceuticals, Exton, Pa.; Martek Biosciences; Columbia, Md.; etc.).

In one aspect of the invention, the T1R modulators can be used in any food product, confectionery, pharmaceutical composition, or ingredient thereof to thereby modulate the taste of the product, composition, or ingredient in a desired manner. For instance, T1R modulators which enhance sweet taste sensation can be added to sweeten a product or composition, while T1R modulators which block undesirable taste sensations can be added to improve the taste of a product or composition.

E. Methods for Representing and Predicting the Perception of Taste

The invention also preferably provides methods for representing the perception of taste and/or for predicting the perception of taste in a mammal, including in a human. Preferably, such methods may be performed by using the receptors and genes encoding said T1R polypeptides disclosed herein.

Also contemplated as within the invention, is a method of screening one or more compounds for the presence of a taste detectable by a mammal, comprising: contacting said one or more compounds with the disclosed receptors, preferably wherein the mammal is a human. Also contemplated as within the invention, is a method for representing taste perception of a particular taste in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n chemosensory receptors of said vertebrate, where n is greater than or equal to 2; and generating from said values a quantitative representation of taste perception. The chemosensory receptors may be a chemosensory receptor disclosed herein, the representation may constitute a point or a volume in n -dimensional space, may constitute a graph or a spectrum, and may constitute a matrix of quantitative representations. Also, the providing

step may comprise contacting a plurality of recombinantly-produced chemosensory receptors with a test composition and quantitatively measuring the interaction of said composition with said receptors.

Also contemplated as within the invention, is a method for predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, comprising the steps of: providing values X_1 to X_n representative of the quantitative stimulation of each of n chemosensory receptors of said vertebrate, where n is greater than or equal to 2, for one or more molecules or combinations of molecules yielding known taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal, providing values X_1 to X_n representative of the quantitative stimulation of each of n chemosensory receptors of said vertebrate, where n is greater than or equal to 2, for one or more molecules or combinations of molecules yielding unknown taste perception in a mammal; and generating from said values a quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal, and predicting the taste perception in a mammal generated by one or more molecules or combinations of molecules yielding unknown taste perception in a mammal by comparing the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding unknown taste perception in a mammal to the quantitative representation of taste perception in a mammal for the one or more molecules or combinations of molecules yielding known taste perception in a mammal. The chemosensory receptors used in this method may include a chemosensory receptor disclosed herein.

In another embodiment, novel molecules or combinations of molecules are generated which elicit a predetermined taste perception in a mammal by determining a value of taste perception in a mammal for a known molecule or combinations of molecules as described above; determining a value of taste perception in a mammal for one or more unknown molecules or combinations of molecules as described above; comparing the value of taste perception in a mammal for one or more unknown compositions to the value of taste perception in a mammal for one or more known compositions; selecting a molecule or combination of molecules that elicits a predetermined taste perception in a mammal; and combining two or more unknown molecules or combinations of molecules to form a molecule or combination of molecules that elicits a predetermined taste perception in a mammal. The combining step yields a single molecule or a combination of molecules that elicits a predetermined taste perception in a mammal.

In another embodiment of the invention, there is provided a method for simulating a taste, comprising the steps of: for each of a plurality of cloned chemosensory receptors, preferably human receptors, ascertaining the extent to which the receptor interacts with the taste stimulus; and combining a plurality of compounds, each having a previously-ascertained interaction with one or more of the receptors, in amounts that together provide a receptor-stimulation profile that mimics the profile for the taste stimulus. Interaction of a taste stimulus with a chemosensory receptor can be determined using any of the binding or reporter assays described herein. The plurality of compounds may then be combined to form a mixture. If desired, one or more of the

plurality of the compounds can be combined covalently. The combined compounds substantially stimulate at least 75%, 80%, or 90% of the receptors that are substantially stimulated by the taste stimulus.

In another preferred embodiment of the invention, a plurality of standard compounds are tested against a plurality of chemosensory receptors to ascertain the extent to which the receptors each interact with each standard compound, thereby generating a receptor stimulation profile for each standard compound. These receptor stimulation profiles may then be stored in a relational database on a data storage medium. The method may further comprise providing a desired receptor-stimulation profile for a taste; comparing the desired receptor stimulation profile to the relational database; and ascertaining one or more combinations of standard compounds that most closely match the desired receptor-stimulation profile. The method may further comprise combining standard compounds in one or more of the ascertained combinations to simulate the taste.

F. Kits

T1R genes and their homologs are useful tools for identifying chemosensory receptor cells, for forensics and paternity determinations, and for examining taste transduction. T1R family member-specific reagents that specifically hybridize to T1R nucleic acids, such as T1R probes and primers, and T1R specific reagents that specifically bind to a T1R polypeptide, e.g., T1R antibodies are used to examine taste cell expression and taste transduction regulation.

Nucleic acid assays for the presence of DNA and RNA for a T1R family member in a sample include numerous techniques are known to those skilled in the art, such as southern analysis, northern analysis, dot blots, RNase protection, S1 analysis, amplification techniques such as PCR, and in situ hybridization. In in situ hybridization, for example, the target nucleic acid is liberated from its cellular surroundings in such as to be available for hybridization within the cell while preserving the cellular morphology for subsequent

interpretation and analysis. The following articles provide an overview of the art of in situ hybridization: Singer et al., *Biotechniques*, 4:230250 (1986); Haase et al., *Methods in Virology*, vol. VII, pp. 189–226 (1984); and *Nucleic Acid Hybridization: A Practical Approach* (Names et al., eds. 1987). In addition, a T1R polypeptide can be detected with the various immunoassay techniques described above. The test sample is typically compared to both a positive control (e.g., a sample expressing a recombinant T1R polypeptide) and a negative control.

The present invention also provides for kits for screening for modulators of T1R family members. Such kits can be prepared from readily available materials and reagents. For example, such kits can comprise any one or more of the following materials: T1R nucleic acids or proteins, reaction tubes, and instructions for testing T1R activity. Optionally, the kit contains a biologically active T1R receptor. A wide variety of kits and components can be prepared according to the present invention, depending upon the intended user of the kit and the particular needs of the user.

EXAMPLES

In the protein sequences presented herein, the one-letter code X or Xaa refers to any of the twenty common amino acid residues. In the DNA sequences presented herein, the one letter codes N or n refers to any of the of the four common nucleotide bases, A, T, C, or G.

Example 1—hT1R3

The hT1R3 genomic DNA is provided below as SEQ ID NO 1 and SEQ ID NO 2 with predicted coding sequences (cds) shown in boldface. The break between the 5' and 3' contigs is shown as ellipses ('. . .'). The hT1R3 predicted cds are described in SEQ ID NO 3. Finally, a preferred, predicted hT1R3 amino acid sequence is provided as SEQ ID NO 4, using the one-letter code for the amino acids.

hT1R3 genomic DNA—5' contig (SEQ ID NO 1)

```

AGCCTGGCAGTGGCCTCAGGCAGAGTCTGACGCGCACAACTTTTCAGGCC (SEQ ID NO:1)
AGGAAGCGAGGACACCCTGGGGCCCCAGGGTGTGGCAAGTGAGGATGGC
AAGGGTTTTGCTAAACAAATCCTCTGCCCGCTCCCCGCCCGGGCTCACTCC
ATGTGAGGCCCCAGTCGGGGCAGCCACCTGCCGTGCGTGTGGAAGTTGCC
TCTGCCATGCTGGGCCCTGCTGTCTGGGCCCTCAGCCTCTGGGCTCTCCTG
CACCCTGGGACGGGGGCCCATTTGTGCCTGTACAGCAACTTAGGATGAAG
GGGACTACGTGCTGGGGGGCTGTTCCCCTGGGCGAGGCCGAGGAGGC
TGGCCTCCGCAGCCGGACACGGCCAGCAGCCCTGTGTGCACCAGGTACA
GAGGTGGGACGGCCTGGGTCGGGGTCAGGGTGACCAGGTCTGGGGTGCTC
CTGAGCTGGGGCCGAGGTGGCCATCTGCGGTTCTGTGTGGCCCCAGGTTCT
CCTCAAACGGCTGCTCTGGGCACTGGCCATGAAAATGGCCGTGGAGGAGA
TCAACAACAAGTCGGATCTGCTGCCCGGGCTGCGCCTGGGCTACGACCTCT
TTGATACGTGCTCGGAGCCTGTGGTGGCCATGAAGCCCAGCCTCATGTTCT
GGCCAAGGCAGGCAGCCGGACATCGCCGCTACTGCAACTACACGCAGTA
CCAGCCCCGTGTGCTGGCTGTTCATCGGGCCCCACTCGTCAGAGCTCGCCAT

```

-continued

GGTCACCGGCAAGTTCTTCAGCTTCTTCTCATGCCCCAGTGGGGCGCCCC
CCACCATCACCCACCCCAACCAACCCCTGCCCGTGGGAGCCCCTTGTGT
CAGGAGAATGC

hTIR3 genomic DNA-3' contig (SEQ ID NO 2)

.....TACATGCACCCACCCAGCCCTGCCCTGGGAGCCCTGTGTCAG (SEQ ID NO 2)
AAGATGCTCTTGGCCTTGCAGGTCAGCTACGGTGCTAGCATGGAGCTGCTG
AGGCCCGGGAGACCTTCCCTCCTTCTTCCGCACCGTGCCAGCGACCGT
GTGCAGCTGACGGCCGCCGCGGAGCTGCTGCAGGAGTTCGGCTGGAAGT
GGTGGCCGCCCTGGGCAGCGACGAGTACGGCCGGCAGGGCCTGAGC
ATCTTCTCGGCCCTGGCCGCGCACGCGGCATCTGCATCGCGCACGAGGG
CCTGGTGCCGCTGCCCGTGGCGATGACTCGCGGCTGGGGAAGGTGCAGG
ACGTCCTGCACCAGGTGAACCAGAGCAGCGTGCAGGTGGTGCTGCTGTTTCG
CCTCCGTGCACGCCGCCACGCCCTTCAACTACAGCATCAGCAGCAGGC
TCTCGCCCAAGGTGTGGGTGGCCAGCGAGGCTGGCTGACCTCTGACCTGG
TCATGGGGCTGCCCGCATGGCCAGATGGGCACGGTGCTTGGCTTCTCTC
AGAGGGGTGCCAGCTGCACGAGTTCCTCCAGTACGTGAAGACGCACCTG
GCCCTGGCCACCGACCCGGCCTTCTGCTCTGCCCTGGGCGAGAGGGAGCA
GGTCTGGAGGAGGACGTGGTGGCCAGCGCTGCCCGCAGTGTGACTGCA
TCACGCTGCAGAACGTGAGCGCAGGGCTAAATCACCACCAGACGTTCTCTG
TCTACGAGCTGTGTATAGCGTGGCCAGGCCCTGCACAACACTCTTCAGTG
CAACGCCTCAGGCTGCCCGCGCAGGACCCCGTGAAGCCCTGGCAGGTGA
GCCCCGGAGATGGGGGTGTGCTGTCTCTGCATGTGCCAGGCCACCAGG
CACGGCCACCACGCTGAGCTGGAGGTGGCTGGCGGCTCAGCCCCGTCCC
CCGCCCGCAGCTCCTGGAGAACATGTACAACCTGACCTTCCACGTGGGCGG
GCTGCCGCTGCGGTTGACAGCAGCGGAAACGTGGACATGGAGTACGACCT
GAAGCTGTGGGTGTGGCAGGGCTCAGTGCCAGGCTCCACGACGTGGGCA
GGTTCAACGGCAGCCTCAGGACAGAGCGCTGAAGATCCGCTGGCACACGT
CTGACAACCAGGTGAGGTGAGGGTGGGTGTGCCAGGCGTGCCCGTGGTAG
CCCCCGGGCAGGGCGCAGCCTGGGGTGGGGCCGTTCCAGTCTCCCCT
GGGCATGCCAGCCGAGCAGAGCCAGACCCAGGCTGTGCGCAGAAGCC
CGTGTCCCGGTGCTCGCGGAGTGCCAGGAGGGCCAGGTGCGCCGGGTCA
AGGGGTTCCACTCCTGCTGCTACGACTGTGTGACTGCGAGGCGGGCAGCT
ACCGGCAAAACCAGGTGAGCCGCTTCCCGGAGGCGGGGGTGGGAACG
CAGCAGGGGAGGGTCTGCCAAGTCTGACTCTGAGACCAGAGCCACAGG
GTACAAGACGAACACCAGCGCCCTTCTCTCTCTCACAGACGACATCGCCT
GCACCTTTTGTGGCCAGGATGAGTGGTCCCCGGAGCGAAGCACACGCTGCT
TCCGCCGAGGTTCTCGGTTCTGGCATGGGGCGAGCCGGCTGTGCTGCTGC
TGCTCCTGCTGCTGAGCCTGGCGCTGGGCTTGTGCTGGCTGCTTTGGGGC
TGTTTCGTTACCATCGGGACAGCCACTGGTTCAGGCCTCGGGGGGGCCCC
TGGCCTGCTTTGGCCTGGTGTGCTGGGCTGGTCTGCCTCAGCGTCTCC

-continued

TGTTCCCTGGCCAGCCAGCCCTGCCCGATGCCCTGGCCCAGCAGCCCTTGT
 CCCACCTCCCGCTCACGGGCTGCCTGAGCACACTCTTCTGCAGGGCGGCCG
 AGATCTTCGTGGAGTCAGAACTGCCTCTGAGCTGGGCAGACCGGCTGAGTG
 GCTGCCTGCGGGGGCCCTGGGCCTGGCTGGTGGTGTCTGCTGGCCATGCTG
 GTGGAGGTGCACTGTGCACCTGGTACCTGGTGGCCTTCCCGCCGAGGTG
 GTGACGGACTGGCACATGTGCCACGGAGGCGCTGGTGCCTGCCGCAC
 ACGCTCCTGGGTGAGCTTCGGCCTAGCGCACGCCACCAATGCCACGCTGGC
 CTTTCTCTGCTTCCCTGGGCACTTTCTCTGGTGGGAGCCAGCCGGGCTGCTAC
 AACCGTGCCCGTGGCCTCACCTTTGCCATGCTGGCCTACTTCATCACCTGGG
 TCTCCTTTGTGCCCTCCTGGCCAATGTGCAGGTGGTCTCAGGCCCGCCGT
 GCAGATGGGGCCCTCCTGCTCTGTGTCTGGGCATCCTGGCTGCCTTCCA
 CCTGCCCAGGTGTTACCTGCTCATGCGGCAGCCAGGGCTCAACACCCCGA
 GTTCTTCTGGGAGGGGGCCCTGGGGATGCCCAAGGCCAGAATGACGGGA
 ACACAGGAAATCAGGGGAAACATGAGTGACCCAACCCTGTGATCTCAGCCC
 CGGTGAACCCAGACTTAGCTGCATCCCCCAAGCCAGCAATGACCCGTG
 TCTCGCTACAGAGACCTCCCGCTCTAGGTTCTGACCCAGGTTGTCTCCTG
 ACCCTGACCCACAGTGAGCCCTAGGCCTGGAGCACGTGGACACCCCTGTG
 ACCATC

hT1R3 full-length genomic DNA (SEQ ID NO 20)

AGCCTGGCAGTGGCCTCAGGCAGAGTCTGACGCGCACAACTTTCAGGCC
 AGGAAGCGAGGACAGCACTGGGGCCCCAGGGTGTGGCAAGTGAGGATGGC
 AAGGGTTTTGCTAAACAAATCCTCTGCCCGCTCCCCGCCCCGGGCTCACTCC
 ATGTGAGGCCCCAGTCGGGGCAGCCACCTGCCGTGCCTGTTGGAAGTTGCC
 TCTGCCATGCTGGGCCCTGCTGTCTGGCCCTCAGCCTCTGGGCTCTCCTG
 CACCCTGGGACGGGGGCCCATTTGTGCCTGTCACAGCAACTTAGGATGAAG
 GGGACTACGTGCTGGGGGGCTGTTCCCCCTGGGCGAGGCCGAGGAGGC
 TGGCCTCCGAGCCGGACACGGCCAGCAGCCCTGTGTGCACCAGGTACA
 GAGGTGGGACGGCCTGGGTGCGGGTCAGGGTGACCAGGTCTGGGGTGCTC
 CTGAGCTGGGGCCGAGGTGGCCATCTGCGGTTCTGTGTGGCCCCAGGTTCT
 CCTCAAACGGCCTGCTCTGGGCACTGGCCATGAAAATGGCCGTGGAGGAGA
 TCAACAACAAGTCGGATCTGCTGCCCGGGCTGCGCCTGGGCTACGACCTCT
 TTGATACGTGCTCGGAGCCTGTGGTGGCCATGAAGCCAGCCTCATGTTCTT
 GGCCAAGGCAGGCAGCCGCGACATCGCCGCTACTGCAACTACACGCAGTA
 CCAGCCCCGTGTGCTGGCTGTCATCGGGCCCCACTCGTCAGAGCTCGCCAT
 GGTACCCGGCAAGTTCTTCAGCTTCTTCTCATGCCCCAGTGGGGCGCCCC
 CCACCATCACCCACCCCAACCAACCCTGCCCGTGGGAGCCCTTGTGT
 CAGGAGAATGCTACATGCACCCACCCAGCCCTGCCCTGGGAGCCCTGTGT
 CAGAAGATGCTCTTGGCCTTGCAGGTCAGCTACGGTGTCTAGCATGGAGCTG
 CTGAGCGCCCGGAGACCTTCCCTCCTTCTCCGCACCGTGCCAGCGAC
 CGTGTGCAGCTGACGGCCGCGGGAGCTGCTGCAGGAGTTCGGCTGGAA

(SEQ ID NO
 20)

-continued

CTGGGTGGCCGCCCTGGGCAGCGACGACGAGTACGGCCGGCAGGGCCTGA
GCATCTTCTCGGCCCTGGCCGGCAGCGGCATCTGCATCGCGCACGAG
GGCCTGGTGCCGCTGCCCGTGCCGATGACTCGCGGCTGGGGAAGGTGCA
GGACGTCTGCACCAGGTGAACCAGAGCAGCGTGCAGGTGGTGCTGCTGTT
CGCCTCCGTGCACGCCGCCACGCCCTCTTCAACTACAGCATCAGCAGCAG
GCTCTCGCCCAAGGTGTGGGTGGCCAGCGAGGCCCTGGCTGACCTCTGACCT
GGTCATGGGGCTGCCCGCATGGCCAGATGGGCACGGTGCTTGGCTTCCT
CCAGAGGGGTGCCAGCTGCACGAGTCCCCAGTACGTGAAGACGCACCT
GGCCCTGGCCACCGACCCGGCCTTCTGCTCTGCCCTGGGCGAGAGGGAGC
AGGGTCTGGAGGAGGACGTGGTGGCCAGCGCTGCCCGCAGTGTGACTGC
ATCACGCTGCAGAACGTGAGCGCAGGGCTAAATCACCACCAGACGTTCTCT
GTCTACGCAGCTGTGTATAGCGTGGCCAGGCCCTGCACAACCTCTTCACT
GCAACGCCTCAGGCTGCCCGCGCAGGACCCCGTGAAGCCCTGGCAGGTG
AGCCCGGGAGATGGGGTGTGCTGTCTCTGCATGTGCCAGGCCACCAG
GCACGGCCACCACGCTGAGCTGGAGGTGGTGGCGGCTCAGCCCCGTCC
CCCGCCCGCAGCTCCTGGAGAACATGTACAACCTGACCTTCCACGTGGGCG
GGCTGCCGCTGCGGTTCGACAGCAGCGGAAACGTGGACATGGAGTACGAC
CTGAAGCTGTGGGTGTGGCAGGGCTCAGTGCCAGGCTCCACGACGTGGG
CAGGTTCAACGGCAGCTCAGGACAGAGCGCCTGAAGATCCGCTGGCACAC
GTCTGACAACCAGGTGAGGTGAGGGTGGGTGTGCCAGGCGTGCCCGTGGT
AGCCCCCGGGCAGGGCGCAGCCTGGGGGTGGGGGCCGTTCCAGTCTCCC
GTGGGCATGCCAGCCGAGCAGAGCCAGACCCAGGCTGTGCGCAGAAG
CCCGTGTCCCGGTGCTCGCGGCAGTGCCAGGAGGGCCAGGTGCGCCGGGT
CAAGGGGTTCACCTCCTGCTGCTACGACTGTGTGGACTGCGAGGGGGCAG
CTACCGGCAAAACCCAGGTGAGCCGCTTCCCGGCAGGCGGGGGTGGGAA
CGCAGCAGGGGAGGGTCTGCAAGTCTGACTCTGAGACCAGAGCCCACA
GGGTACAAGACGAACACCAGCGCCCTTCTCTCTCACAGACGACATCG
CCTGCACCTTTTGTGGCCAGGATGAGTGGTCCCCGGAGCGAAGCACACGCT
GCTTCCGCCCGCAGGTCTCGGTTCTTGGCATGGGGCGAGCCGGCTGTGCTGC
TGCTGCTCCTGCTGCTGAGCCTGGCGCTGGGCCTGTGCTGGCTGCTTTGG
GGCTGTTGTTTACCATCGGGACAGCCACTGGTTCAGGCCCTCGGGGGGGC
CCCTGGCCTGCTTTGGCCTGGTGTGCTGGCCTGGTCTGCTCAGCGTCC
TCCTGTTCCCTGGCCAGCCAGCCCTGCCGATGCCCTGGCCAGCAGCCCT
TGTTCCACCTCCCGCTCACGGGCTGCCTGAGCACACTCTTCTGTCAGGCGG
CCGAGATCTTCTGGAGTCAGAACTGCCTCTGAGCTGGGCAGACCGGCTGA
GTGGCTGCCTGCGGGGGCCCTGGGCCTGGCTGGTGGTGTGCTGGCCATG
CTGGTGGAGGTGCACTGTGCACCTGGTACCTGGTGGCTTCCCGCCGGAG
GTGGTGACGGACTGGCACATGCTGCCACGGAGGCGCTGGTGCACCTGCCG
CACACGCTCCTGGGTGAGCTTCCGGCTAGCGCACGCCACCAATGCCAGCT
GGCCTTCTCTGCTTCTGGCACTTCTGTTGGGAGCCAGCCGGGCTG

-continued

CTACAACCGTGCCCGTGGCCTCACCTTTGCCATGCTGGCCTACTTCATCACC
 TGGGTCTCCTTTGTGCCCTCCTGGCCAATGTGCAGGTGGTCTCAGGCC
 GCCGTGCAGATGGGCGCCCTCCTGCTCTGTGTCTGGGCATCCTGGCTGCC
 TTCCACCTGCCAGGTGTTACCTGCTCATGCGGCAGCCAGGGCTCAACACC
 CCCGAGTTCTTCTGGGAGGGGGCCCTGGGGATGCCCAAGGCCAGAATGA
 CGGGAACACAGGAAATCAGGGGAAACATGAGTGAACCAACCTGTGATCTC
 AGCCCCGGTGAACCCAGACTTAGCTGCGATCCCCCAAGCCAGCAATGAC
 CCGTGTCTCGCTACAGAGACCCTCCCGCTCTAGGTTCTGACCCAGGTTGTC
 TCCTGACCCTGACCCACAGTGAGCCCTAGGCCTGGAGCACGTGGACACC
 CTGTGACCATC

hTIR3 predicted cds (SEQ ID NO 3)

ATGCTGGGCCCTGCTGTCTGGGCCTCAGCCTCTGGGCTCTCCTGCACCCT
 GGGACGGGGGCCCCATTGTGCCTGTCACAGCAACTTAGGATGAAGGGGGAC
 TACGTGCTGGGGGGGCTGTCCCCCTGGGCGAGGCCGAGGAGGCTGGCCT
 CCGCAGCCGGACACGGCCCAGCAGCCCTGTGTGCACCAGGTTCTCCTCAA
 CGCCTGCTCTGGGCAC TGCCATGAAAATGGCCGTGGAGGAGATCAACAA
 CAAGTCGGATCTGTGCCGGGCTGCGCCTGGGCTACGACCTCTTTGATAC
 GTGCTCGGAGCCTGTGGTGGCCATGAAGCCCAGCCTCATGTTCTGGCCAA
 GGCAGGCAGCCGACATCGCCGCTACTGCAACTACACGCAGTACCAGCC
 CCGTGTGCTGGCTGTTCATCGGGCCCCACTCGTCAGAGCTCGCCATGGTCAC
 CGGCAAGTTCTTTCAGCTTCTTCCTCATGCCCCAGGTCAGCTACGGTGCTAGC
 ATGGAGCTGCTGAGCGCCGGGAGACCTTCCCCCTCTTCTTCCGCACCGTG
 CCCAGCGACCGTGTGCAGCTGACGGCCGCGGAGCTGCTGCAGGAGTT
 CGGCTGGAAC TGGTGGCCGCCCTGGGCAGCGACGACGAGTACGGCCGGC
 AGGCCTGAGCATCTTCTCGGCCCTGGCCGCGGCACGCGGCATCTGCATCG
 CGCACGAGGGCCTGGTGCCGCTGCCCCGTGCCGATGACTCGGGCTGGGG
 AAGGTGCAGGACGTCTGCACCAGGTGAACCAGAGCAGCGTGCAGGTGGTG
 CTGCTGTTTCGCTCCGTGCACGCCGCCACGCCCTTCAACTACAGCATCA
 GCAGCAGGCTCTGCCCAAGGTGTGGGTGGCCAGCGAGGCCCTGGCTGACC
 TGTGACCTGGTCATGGGGCTGCCCGGCATGGCCAGATGGGCACGGTGCTT
 GGCTTCTCCAGAGGGGTGCCAGCTGCACGAGTCCCCCAGTAGGTGAAG
 ACGCACCTGGCCCTGGCCACCGACCCGGCCTTCTGCTCTGCCCTGGGCGAG
 AGGGAGCAGGGTCTGGAGGAGACGTGGTGGGCCAGCGCTGCCCGCAGTG
 TGA CTGCATCACGCTGGAGAACGTGAGCGCAGGGCTAAATCACCACCAGAC
 GTTCTCTGTCTACGGAGCTGTGTATAGCGTGGCCCAGGCCCTGCACAACACT
 CTTCACTGCAACGCCCTCAGGCTGCCCCGCGCAGGACCCCGTGAAGCCCTGG
 CAGCTCCTGGAGAACATGTACAACCTGACCTTCCACGTGGGCGGGCTGCCG
 CTGCGGTTTCGACAGCAGCGAAACGTGGACATGGAGTACGACCTGAAGGTG
 TGGGTGTGGCAGGGCTCAGTGCCAGGCTCCACGACGTGGGCAGGTTCAAC
 GGCAGCCTCAGGACAGAGCGCCTGAAGATCCGCTGGCACACGTCTGACAAC

(SEQ ID NO 3)

-continued

CAGAAAGCCCGTGTCCCGGTGCTCGCGGCAGTGCCAGGAGGGCCAGGTGCG
 CCGGGTCAAGGGGTCCACTCCTGCTGCTACGACTGTGTGGACTGCGAGGC
 GGGCAGCTACCGGCAAACCCAGACGACATCGCCTGCACCTTTTGTGGCCA
 GGATGAGTGGTCCCCGGAGCGAAGCACAGGGTGCCTCCGCCGAGGTCTCG
 GTTCTGGCATGGGGCGAGCCGGCTGTGCTGCTGCTGCTCCTGCTGCTGAG
 CCTGGCGCTGGGCCTTGTGCTGGCTGCTTTGGGGCTGTTCTTACCATCG
 GGACAGCCCACTGGTTCAGGCCTCGGGGGGGCCCTGGCCTGCTTTGGCC
 TGGTGTGCTGGGCCTGGTCTGCCTCAGCGTCCTCCTGTTCCCTGGCCAGC
 CCAGCCCTGCCCAGTGCCTGGCCCAGCAGCCCTTGTCCACCTCCCGCTCA
 CGGGCTGCCTGAGCACACTCTTCTGTCAGGCGGCCGAGATCTTCGTGGAGT
 CAGAACTGCCTGTGAGCTGGGCAGACCGGCTGAGTGGCTGCCTGCGGGGG
 CCCTGGGCCTGGCTGGTGGTGTGCTGCTGGCCATGCTGGTGGAGGTGCGACT
 GTGCACCTGGTACCTGGTGGCCTTCCCGCCGGAGGTGGTGACGGACTGGCA
 CATGCTGCCACGGAGGCGCTGGTGCCTGCGCACACGCTCCTGGGTGAG
 CTTCCGGCTAGCGCACGCCACCAATGCCACGCTGGCCTTTCTCTGCTTCTG
 GGCACTTTCTGGTGGGAGGCAGCCGGGCTGCTACAACCGTGCCCGTGGC
 CTCACCTTTGCCATGCTGGCCTACTTCATCACCTGGGTCTCCTTTGTGCCCC
 TCCTGGCCAATGTGCAGGTGGTCTCAGGCCCGCCGTGCAGATGGGCGCCC
 TCCTGCTCTGTGCTCCTGGGCATCCTGGCTGCCTTCCAGCTGCCAGGTGTTA
 CCTGCTCATGCGGCAGCCAGGGCTCAACACCCCGAGTCTTCTGTTGGGAGG
 GGGCCCTGGGGATGCGCAAGGCCAGAATGACGGGAACACAGGAAATCAGG

GGAAACATGAGTGA

hTIR3 conceptual translation (SEQ ID NO 4)

MLGPAVLGLSLWALLHPGTGAPLCLSQQLRMKGDVVLGGLFPLGEAEELRLS
 RTRPSSPVCTRFSSNGLLWALAMKMAVEEINNKSDDLPLGLRLGYDLFDTCSEPV
 VAMKPSLMFLAKAGSRDIAAYCNYTQYQPRVLAVIGPHSSELAMVTGKFFSFFLM
 PQVSYGASMELLSARETFPSFFRTVPSDRVQLTAAAEELQEFGWNVAAALGSD
 DEYGRQLSIFALAAARGICIAHEGLVPLPRADDSRLGKVQDVLHQVNQSSVQ
 VVLLFASVHAAHALFNYSISSRLSPKVWVASEAWLTSDLVMGLPGMAQMGTVLG
 FLQGAQLHEFPQYVKTHLALATDPAFCSALGEREQGLEEDVVGQRCPQDCIT
 LQNVSAGLNHHQTFSVYAAVYSVAQALHNTLQCNASGCPAQDPVKPWQLLENM
 YNLTFHVGGPLRFDSSGNVMEYDLKLWVWQGSVPRLHDVGRFNGSLRTERL
 KIRWHTSDNQKPVSRCSRQCQEGQVRRVKGFHSCCYDCVDCEAGSYRQNPDD
 IACTFCGQDEWSPERSTRCFRRRSRFLAWGEPVLLLLLLLLSLALGLVLAALGLF
 VHHRDSPLVQASGGPLACFLVCLGLVCLSVLLFPQPSPARCLAQQPLSHLPL
 TGCLSTLFLQAAEIFVESELPLSWADRLSGCLRGPWAWLVLLAMLVEVALCTW
 YLVAFPPEWTDWHMLPTEALVHCRTRSWSVFGLAHATNATLAFLCFLGTFLVR
 SQPGCYNRARGLTFAMLAYFITWVSFVPLLANVQVLRPAVQMGALLLCVLGILA
 AFHLPRCYLLMRQPGLNTPFEFFLGGGPGDAQQNDGNTGNQKHE

(SEQ ID NO 4)

Example 2—rT1R3 and mT1R3

Segments of the rat and mouse T1R3 genes were isolated by PCR amplification from genomic DNA using degenerate primers based on the human T1R3 sequence. The degenerate primers SAP077 (5'-CGNTTYTNGCNTGGGGN-GARCC-3'; SEQ ID NO 5) and SAP079 (5'-CGNGC-
5 NCGRTTRTARCANCCNGG-3'; SEQ ID NO 6) are complementary to human T1R3 residues RFLAWGEPA (corresponding to SEQ ID NO 7) and PGCYNRAR (corresponding to SEQ ID NO 8), respectively. The PCR products

were cloned and sequenced. Plasmid SAV115 carries a cloned segment of the mouse T1R3 gene, and SAV118 carries a segment of the rat gene. These sequences, shown below, clearly represent the rodent counterparts of human T1R3, since the mouse segment is 74% identical to the corresponding segment of human T1R3, and the rat segment is 80% identical to the corresponding segment of human T1R3. The mouse and rat segments are 88% identical. No other database sequences are more than 40% identical to these T1R3 segments.

SAV115 mouse T1R3 segment in sense orientation (sequence corresponding to degenerate primer removed) (SEQ ID NO 9)

GTGCTGTCACTCCTCTGCTGCTTTGCCTGGTGTGGGTCTAGCACTGGCTG (SEQ ID NO 9)
CTCTGGGGCTCTCTGTCCACCACTGGGACAGCCCTGTTGTCCAGGCCTCAG
GCGGCTCACAGTTCTGCTTTGGCCTGATCTGCCTAGGCCTCTTCTGCCTCAG
TGTCTTTCTGTTCCCAGGACGGCCAAGCTCTGCCAGCTGCCTTGACAACAA
CCAATGGCTCACCTCCCTCTCACAGGCTGCCTGAGCACACTCTTCTGCAAG
CAGCTGAGACCTTTGTGGAGTCTGAGCTGCCACTGAGCTGGGCAAACCTGGC
TATGCAGCTACCTTCGGGACTCTGGCCTGCTAGTGGTACTGTTGGCACTTT
TGTGGAGGCAGCACTATGTGCCTGGTATTTGACCGCTTACCAGAAGTGGTG
ACAGACTGGTCAGTGTGCCACAGAGGTACTGGAGCACTGCCACGTGCGT
TCCTGGGTCAACCTGGGCTTGGTGCACATCACC AATGCAATGGTAGCTTTTC
TCTGCTTTCTGGGCACTTTCTCTGGTACAAGACCAG

mT1R3 segment, conceptual translation (SEQ ID NO 10)

VLSLLLLLCLVLGLLALGLSVHHWDSPLVQASGGSQFCFLICLGLFCLSVLLF (SEQ ID NO 10)
PGRPSASCLAQQPMAHLPLTGCLSTLFLQAAETFVESELPLSWANWLCSYL RD
SGLLVVLLATFVEAALCAWYLTASPEVVTDWSVLPTEVLEHCHVRSWVNLGLVH
ITNAMVAFLCFLGTFLVQDQ

SAV118 rat T1R3 segment in sense orientation (sequence corresponding to degenerate primer removed) (SEQ ID NO 11)

GTGCTGTCACTTCTCCTGCTGCTTTGCCTGGTGTGGCCTGACACTGGCTG (SEQ ID NO 11)
CCCTGGGGCTCTTTGTCCACTACTGGGACAGCCCTCTTGTTCAGGCCTCAGG
TGGGTCACTGTTCTGCTTTGGCCTGATCTGCCTAGGCCTCTTCTGCCTCAGT
GTCCTTCTGTTCCCAGGACGACCAGCTCTGCCAGCTGCCTTGCCCAACAAC
CAATGGCTCACCTCCCTCTCACAGGCTGCCTGAGCACACTCTTCTGCAAGC
AGCCGAGATCTTTGTGGAGTGTGAGCTGCCACTGAGTTGGGCAAACCTGGCT
CTGCAGCTACCTTCGGGGCCCTGGGCTTGGCTGGTGGTACTGCTGGCCAC
TCTTGTGGAGGCTGCACTATGTGCCTGGTACTTGTGCTTTCCCTCCAGAG
GTGGTGACAGATTGGCAGGTGCTGCCACGGAGGTACTGGAACACTGCCGC
ATGCGTTCCTGGGTGAGCCTGGGCTTGGTGCACATCACC AATGCAGGGGTA
GCTTTCCTCTGCTTTCTGGGCACTTTCTCTGGTACAAAGCCAG

rT1R3 segment, conceptual translation (SEQ ID NO 12)

VLSLLLLLCLVLGLTLAALGLFVHYWDSPLVQASGGSFLFCFLICLGLFCLSVLLFP (SEQ ID NO 12)
GRPRSASCLAQQPMAHLPLTGCLSTLFLQAAEIFVESELPLSWANWLCSYL RGP

-continued

WAWLVVLLATLVEAALCAWYLMAFPPEVVTDWQVLPTEVLEHCRMRSWVSLGL

VHITNAGVAFLCFLGTFVLVQSQ

Example 3—Cloning of rT1R3

The mT1R3 and rT1R3 fragments identified above as SEQ ID NOs 9 and 11 were used to screen a rat taste tissue-derived cDNA library. One positive clone was sequenced and found to contain the full-length rT1R3 sequence presented below as SEQ ID NO 13. Sequence

comparison to the mT1R3 and rT1R3 partial sequences and to the full-length hT1R3 sequence established that this cDNA represents the rat counterpart to hT1R3. For example, the pairwise amino acid identity between rT1R3 and hT1R3 is approximately 72%, whereas the most related annotated sequence in public DNA sequence data banks is only approximately 33% identical to rT1R3.

rT1R3 predicted cds (SEQ. ID NO. 13)

(SEQ ID NO 13)

ATGCCGGTTTGGCTATCTTGGCCTCAGTCTGGCTGCTTTCCTGGAGCTTG
GGATGGGGTCCCTCTTTGTGTCTGTACAGCAATTCAAGGCACAAGGGGACTA
TATATTGGGTGGACTATTTCCCCTGGGCACAACGAGGAGGCCACTCTCAAC
CAGAGAACACAGCCCAACGGCATCCTATGTACCAGGTTCTCGCCCTTGGTT
TGTTCCCTGGCCATGGCTATGAAGATGGCTGTAGAGGAGATCAACAATGGATC
TGCCTTGCTCCCTGGGCTGCGACTGGGCTATGACCTGTTTGACACATGCTCA
GAGCCAGTGGTCACCATGAAGCCCAGCCTCATGTTTATGGCCAAGGTGGGA
AGTCAAAGCATTTGCTGCCTACTGCAACTACACACAGTACCAACCCCGTGTGC
TGGCTGTCAATTGGTCCCCACTCATCAGAGCTTGCCCTCATTACAGGCAAGTT
CTTCAGCTTCTTCCCTCATGCCACAGGTCAGCTATAGTGCCAGCATGGATCGG
CTAAGTGACCGGAAACATTTCCATCCTTCTTCCGCACAGTGCCAGTGACC
GGGTGCAGCTGCAGGCCGTTGTGACACTGTTGCAGAATTTAGCTGGAAC
GGGTGGCTGCCTTAGGTAGTGATGACTATGGCCGGAAGGTCTGAGCA
TCTTTTCTGGTCTGGCCAACACAGGATATCTGCATTGCACACGAGGGCCT
GGTGCCACAACATGACACTAGTGGCCAACAATTGGGCAAGGTGGTGGATGT
GCTACGCCAAGTGAACCAAAGCAAAGTACAGGTGGTGGTGTGTTTGCATCT
GCCCGTGTGTCTACTCCCTTTTGTAGCTACAGCATCCTTCATGACCTCTCACC
CAAGGTATGGGTGGCCAGTGAGTCTGGCTGACCTCTGACCTGGTCATGAC
ACTTCCCAATATTGCCCGTGTGGGCACTGTTCTTGGGTTTCTGCAGCGCGGT
GCCCTACTGCCTGAATTTTCCATTATGTGGAGACTCGCCTTGCCCTAGCTG
CTGACCGAACATTCTGTGCCTCCGTGAAAGGTGAGTTGGATCTGGAGGAGCG
CGTGATGGGGCCACGCTGTTACAATGTGACTACATCATGCTACAGAACCTG
TCATCTGGGCTGATGCAGAACCATCAGCTGGGCAGTTGCACCACCAATAT
TTGCAACCTATGCAGCTGTGTACAGTGTGGCTCAGGCCCTTCACAACACCCT
GCAGTGCAATGTCTCACATTGCCACACATCAGAGCCTGTTCAACCCTGGCAG
CTCCTGGAGAACATGTACAATATGAGTTTCCGTGCTCGAGACTTGACACTGC
AGTTTGATGCCAAAGGGAGTGTAGACATGGAATATGACCTGAAGATGTGGGT
GTGGCAGAGCCCTACACCTGTACTACATACTGTAGGCACCTTCAACGGCACC
CTTCAGCTGCAGCACTCGAAAATGTATTGGCCAGGCAACCAGGTGCCAGTCT
CCCAGTGTCCCGGCAGTGCAAAGATGGCCAGGTGCGCAGAGTAAAGGGCT

-continued

TTCATTCTGCTGCTATGACTGTGTGGACTGCAAGGCAGGGAGCTACCGGAA
GCATCCAGATGACTTCACCTGTACTCCATGTGGCAAGGATCAGTGGTCCCCA
GAAAAAGCACAACTGCTTACCTCGCAGGCCCAAGTTTCTGGCTTGGGG
GAGCCAGCTGTGCTGTCACTTCTCCTGCTGCTTTGCCTGGTGTGGGCTGA
CACTGGCTGCCCTGGGGCTCTTTGTCCACTACTGGGACAGCCCTCTTGTTC
GGCTCAGGTGGGTCACTGTCTGCTTTGGCTGATCTGCCTAGGCCTCTTC
TGCTCAGTGTCTTCTGTTCAGGACGACCACGCTCTGCCAGCTGCCTTG
CCCAACAACCAATGGCTCACCTCCCTCTCACAGGCTGCCTGAGCACACTCTT
CCTGCAAGCAGCCGAGATCTTTGTGGAGTCTGAGCTGCCACTGAGTTGGGC
AAACTGGCTCTGCAGCTACCTTCGGGGCCCCCTGGGCTTGGCTGGTGGTACT
GCTGGCCACTCTTGTGGAGGCTGCACTATGTGCCCTGGTACTTGATGGCTTTC
CCTCCAGAGGTGGTGACAGATTGGCAGGTGCTGCCACGGAGGTACTGGAA
CACTGCCGCATGCGTTCCTGGGTGAGCCTGGGCTTGGTGCACATACCAAT
GCAGTGTAGCTTTCCTCTGCTTTCTGGGCACCTTCTGGTACAGAGCCAGC
CTGGTCGCTATAACCGTGCCTGGCCTGACCTTCGCCATGCTAGCTTATTT
CATCATCTGGGTCTCTTTTGTGCCCTCCTGGCTAATGTGCAGGTGGCTTAC
CAGCCAGCTGTGCAGATGGGTGCTATCTTATTCTGTGCCCTGGGCATCCTGG
CCACCTTCCACCTGCCCAAATGCTATGTACTTCTGTGGCTGCCAGAGCTCAA
CAGCCAGGAGTTCTTCTGGGAAGGAGCCCCAAGGAAGCATCAGATGGGAA
TAGTGGTAGTAGTGAGGCAACTCGGGGACACAGTGAATGA

rT1R3 conceptual translation (SEQ. ID NO. 14)

MPGLAILGLSLAAFLELGMSSLSLQFKAQGDYIILGLFPLGTTEEATLNQRT (SEQ ID NO 14)
QPNGILCTRFSPLGLFLAMAMKMAVEE INNGSALLPGLRLGYDLFDTCSEPVTM
KPSLMFMAKVGVSQSIAYCNYTQYQPRVLAVIGPHSSELALITGKFFSFFLMPQV
SYSASMDRLSDRETFFSFFRTVPSDRVQLQAVVTLQNFSWNWVAALGSDDDY
GREGLSIFSGLANSRGICIAHEGLVPQHDTSGQQLGKWDVLRQVNQSKVQVVV
LFASARAVYSLFSYSILHDLSPKVWVASESWLTSDLVMTLPNIARVGTVLGFLQR
GALLPEFSHYVETRLALADPTFCASLKAELDLEERVMPGRCSQCDYIMLQNLSS
GLMQNLSAGQLHHQIFATYAAVYSVAQALHNTLQCNVSHCHTSEPVQPWQLE
NMYNMSFRARDLTLQFDAQGSVDMEYDLKMWWQSPTPVLHTVGTFNGLTLQL
QHSKMYWPGNQVPVSQCSRQCKDQVRRVKGFHSCCYDCVDCKAGSYRKHP
DDFTCTPCGKDQWSPEKSTTCLPRRPKFLAWGEPAVLSLLLLLCLVLGLTLAAL
GLFVHYWDSPLVQASGSLFCFLICLGLFCLSVLLFPGRPRSASCLAQQPMAH
LPLTGCLSTLFLQAEIFVESELPLSWANWLC SYLRGPWAWLVLLATLVEAALC
AWYLMAFPPEVVTDWQVLPTEVLEHCRMRSWVSLGLVHITNAVLAFLCFLGTFL
VQSQPGRYNRARGLTFAMLAYFI IWVSFVPLL ANVQVAYQPAVQMGAILFCALGI
LATFHLPKCYVLLWLPENLTQEFFLGRSPKEASDGNSSSEATRHSSE

Example 4—Expression of mT1R3

The above described mouse T1R3 fragment contained in SAV115 was PCR amplified using M13 forward and M13 reverse primers and then gel purified. The T1R3 DNA

template was placed into an in vitro transcription labeling reaction where Digoxigenin labeled UTP was incorporated into an antisense cRNA probe. This probe was hybridized to adult mouse taste tissue containing circumvallate papillae. The T1R3 in situ hybridization and detection were per-

-continued

TAAACAACCTCCACGGCCCTGCTGCCCAACATCACCCCTGGGGTACCAGCTGT
ATGATGTGTGTTCTGACTCTGCCAATGTGTATGCCACGCTGAGAGTGCTCTC
CCTGCCAGGGCAACACCACATAGAGCTCCAAGGAGACCTTCTCCACTATTCC
CCTACGGTGCTGGCAGTGATTGGGCCTGACAGCACCAACCGTGCTGCCACC
ACAGCCGCCCTGCTGAGCCCTTCTCTGGTGCCCATGGTAAGCTGGAGCCTC
AGACCTTTGCCCATCTCCCTTCAGGCAAGTCTGGGNNNNNNNNNNNNNNNN
NNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNNGCCACCATGCCCGGCTA
ATTTTTTTGTATTTTTAGTAGAGACGGGGTTTCACCGTGTAGCCAGGCTGGT
CGCAAACCTCCTAACCTCGTGATCCACCCACCTCGGCCTCCCAATGTGCTGGG
ATTACAGGTGTGAGCCACTGCACCCGGCCATAATGTATTAATATAATAAAATA
ATTATACAACCTCACATAATGTAGAATCAGTGGGAGCCCTGAGCTTGTTTTCC
TACAACCTAGATGGTCCCATCTGGGGGTGATGGGAGACAGTGACAGATCATCA
GACATTAGATTCTCATAAGTAGCGTGCAACCCAGATCCCTCGCATGTGCAGT
TCACAGTAGGGTTCAAGCTCCTACAAGAATCTGATGCTGCTGATCTGAC
AGGAGGGGAGCAGCTGTAAATACAGATGAAGCTTCGCTTACTCACCAGCTGC
TCACCTCCTCCTGTGAGGCCGGTTCTAACAGGCCACTGACCTAACTTCTG
CCCTGACCTACACATGCTTCTTCTTCTTCTTGCAAACCTGCCTCCAGTGAAGT
CCCTGAAGGTCCCCAACACACGGGACTATTTCACTCCTATGCAGTTTTGT
CTCCTTTGCTTGGAATGCATCCCCCTCACCCCTTGTCCTCCAGGCAGATTCCCA
CCCCCTCCCCAGAACCTGCCCCAGTGGAGCCTTCGCAGGTGATTTGTGAGTT
TCACAGGCTGAGGGGTGCTCTCCTGGTCTCCCCGGCTCCCTGTATCCCCAC
ACCCAGCACAGGGCCAGGCACTGGGGGGCCTTCAGTGGAGACTGAAATG
GCTGAACGGGACCTCCCATAGATTAGCTATGCGGCCAGCAGCGAGACGCTC
AGCGTGAAGCGGCAGTATCCCTCTTCTGCGCACCATCCCCAATGACAAGT
ACCAGGTGGAGACCATGGTGTCTGCTGCTGCAGAAGTTCGGGTGGACCTGGA
TCTCTCTGGTTGGCAGCAGTGACGACTATGGGCAGCTAGGGGTGCAGGCAC
TGGAGAACCAGGCCACTGGTCAGGGGATCTGCATGCTTTCAAGGACATCAT
GCCCTTCTCTGCCAGGTGGGCGATGAGAGGATGCAGTGCTCATGCGCCA
CCTGGCCCAGGCCGGGGCCACCGTCGTGGTTGTTTTTCCAGCCGGCAGTT
GGCCAGGGTGTTTTTCGAGTCCGTGGTGCTGACCAACCTGACTGGCAAGGT
GTGGGTGCCTCAGAAGCCTGGGCCCTCTCCAGGCACATCACTGGGGTGCC
CGGGATCCAGCGCATTGGGATGGTGCTGGGCGTGCCATCCAGAAGAGGG
CTGTCCCTGGCCTGAAGCGTTTGAAGAAGCCTATGCCCGGGCAGACAAGA
AGGCCCTAGGCCTTGCCACAAGGGCTCCTGGTGCAGCAGCAATCAGCTCT
GCAGAGAATGCCAAGCTTTCATGGCACACACGATGCCCAAGCTCAAAGCCTT
CTCCATGAGTTCTGCCTACAACGCATACCGGGCTGTGTATGCGGTGGCCAT
GGCCTCCACCAGCTCCTGGGCTGTGCCTCTGGAGCTTGTTCAGGGGCCGA
GTCTACCCCTGGCAGGTAAGAGAGCCACCCAGCACCTCCTGTCAGGGAG
AACAGCCAATCCTGAGATGAGCAGAGTGGGCACTCTCCGGTCACTCTAAATG
CCAAGGGGGATAAATGCCACTAACTTGAGGTTTTTTGTTTTGTTTTGTTTTGT

-continued

TTTTTGAGACAGTCTGGCTCTGTCACCCAGGCTGCAGTGTAGTGATGCGATC
TCGGCTCTCTGCAACTTCCACCTCCTGGGTTCAAGTGATTCTCTTGCCTCGG
CCTCCTGAGTAGCTGGGATTACAGGCACCCACCACCATGCCTGGATAATTTT
TCTTTTCTTTTTTTTTTTTTTTTGGAGATAGAGTCTCGCTCTGTTGCCAGGCTGGA
ATGCAGTGGTGCATCTTGGCTCACTGTGAGCTCCGCCTCCAGGTTCACTC
CATTTCCCTGCCTCAGCTCCCAAGTAGGTGGGACTACGGGCGCCCGCCAC
CACGCCAGCTAATTTTTTTTGTATTTTGTAGTAGAGACGGGGTTTACCATGT
TAGCCAGGATGGTCTCAATCTCCTGACCTTGTCTATCCGCCACCTCGTCCTC
CCAAAGTGCTGGGATTACAGGCGTGAGCCACCGCACCCGGCCTAATTTTTTGT
ATTTTGTAGTAGAGATGGGGTTTACCATGTTGGCCAGGCTGGTCTCGAACTC
CTGGCATCAAGTGATCCCTCCTGCTTTCGGCCTCCCAAAGTGCTGGGATTACAG
GCATTAGCTCTCTTCTCTTAGACAGATCTTCTCTCTGATCCTTGCCTTCTCTC
ACCCACTGTGTCTTGGAAAGTGCAAGTGATAAGATCCAGGGCTAAAAGTGT
TGTAAGGAGTGTGTTGTTAGAGGCTCCTCTCAGGAGGTTGGTGGGAAGAT
TGAGGGGCTTCTAAGAAGGAAGGACGAGACCTTCTGATGGGCTGAAAC
CACCAGGACGGAACCCAGGAAGGCCCCAGGCCCTTGCTTCTGGGACCATG
TGGGTCTGTGCTGTCTGTGGTGGCTTCATGATACGCGTTTCTTTTCTAGCTTTTG
GAGCAGATCCACAAGGTGCATTTCTTCTACACAAGGACACTGTGGCGTTTA
ATGACAACAGAGATCCCTCAGTAGCTATAACATAATTGCCTGGGACTGGAA
TGGACCCAAGTGGACCTTCCAGGTCCTCGGTTCCCTCCACATGGTCTCCAGTT
CAGCTAAACATAAATGAGACCAAAATCCAGTGGCACGGAAAGGACAACCAG
GTAATGGGGATGTGGCTACTCACCATGTAAGTGGCTTATGGGCAACCTAGAG
CCTGGGGGTGATGCTGACACAGTGTACAGGGAGCAGGAGGGGGGCCCCAG
GGTCCAGCTGCCACCACTCTACCCATCCTGGCCAGGGAAGCAGGGAAGAC
ACTCCGTAGGCGAGTGTGCAGATGCCCTGGGGCGGAAGTTCACACGACCAG
GGGCCCTGCCCTGGGAGTGAGCCCTGAGGGCAGATGCACAGAGATTCTGTT
TTCTGTTCCACATGTGAGCTGTCTTTGACTTGGGCCCTACGTGTGGCCCC
TCTGGCTTCTTACAGGTGCCTAAGTCTGTGTGTTCCAGCGACTGTCTTGAAG
GGCACCAGCGAGTGGTTACGGGTTTCCATCACTGCTGCTTTGAGTGTGTGCC
CTGTGGGGCTGGGACCTTCCCTCAACAAGAGTGGTGAGTGGGCAATGGAGCA
GGCGAGCTACCCAGCACTCCCGGGGGCTGCACGGTGGAGGGAGGGCCTCC
CTTGGGCCCCATGTGCCCTGCCCCAGAACCAAGGCCAGTCACTGGGCTGC
CAGTTAGCTTACAGGTTGGAGGACACCTGCTACCAGACAGAATTCTGATCAAG
AGAATCAGCCACTGGGTGCGGTGGCTCATGCCGTGTAATCCAGCACTTTGG
GAGGCTGAGGCGGGTGGATCACTTGAGGTGGGAGTTCGAGACCAGCCTG
GCCAACATGGTGAACCCCATCTCTACCAAAAATATAAAAATTAGCTGGGTG
TGGTGGCGCGTGCCTGTAATCCAGCTACTCGGGAGGCTGAGGCAGGAGAA
TCACTTGAACCCAGGAGCGGAGGTTGCAGTGAGCCAAGATGCATTCCAGC
CTGGACCACAAAGCGAGAATTCGTCCCCCAAAAAAGAAAGGAGGCCGGG
CGCGGTGGCTCACACCTGTAATCCAGCACTTTGGGAGGCCGAGGTGGGTG

-continued

GATCACCTGAGGTCAGGAGTTCGAGACCAGCCTGACCAACATGGTGAAACC
 CCATCTCTACTAAAAATACAAAAAAGTTAGCCGGGCGTTGTGGCGTGTGCC
 TGTAAATCCAGCTACTCGGGAGGCTGAGGCAGGAGAATTGCTTGAACCCGG
 GAGGCGGAGGTTGCAGTGAGCCAAGATTGCACCATTGCACTCCAGCCTGGG
 CGACAAGAGAAAACTCTGTCTCAAAAAAAGAAAGAAAGAATTAGCC
 AACTGAAAGCCTTAGACTGAGGTGTGTCTCTGTTAGAGAGCTGTCATCACA
 ACTCCTACAAAAGCAGTCGTATCCTGAATTCAACCTCTTTCTCTAAATGAATAT
 AGCTATTGTTCCCTTTGTGCCCTCTTGTCTACTGTCCCTTCTGTTGCCCATG
 CCAAAGACAGCTAGCTCCTTGAACAGCTTGGCCTGAATACAGATACTAGCGT
 GTCTGCAGCAGAGAAAAACAGCATTCCCCATCCAGAAATGCAAGGTCAAG
 AACAGAGAGCAAATTAGGTAGCTAAGGACTCAGGTCTTAGTTGGTGTCCAG
 GGGCCACATTCTTTCCCTTACCATCTCTGTAGGGACAGGAATACTTCCCTTC
 TGTCTCAGAGGGTCCAGGACTCAGAGAAACCACAGAGCAGCAGCTCAGGAA
 AGTGGTTCATGAAATGCTGGCAAGAGAGAGGGGTTACAATGCCCTCCCTTG
 GGAGCAGGCTGCTCCCATCAGATCGTAACCTCTCTGGTATGTGGGCAGAGC
 TACCAGGTTAAGTCCCTCCCTAGGGTTTGCAAACCTCATGGGATCATGAG
 CCATACAGAACCGACCTGTGTGTCTCCAGAGTCTGTAATTAACACAGGCATTT
 TGAGGAAATGCGTGGCTCAGGCCCACTCCCGGCTACCCCATCCACTA
 TGCTTAGTATAGTCTAGCTGCCCTGGTACAATTCTCCAGTATCTTGCAGGC
 CCCTATTTCTTACTTCTGCTCATCTGGCTCTCAGGAACCTTCTTGGCC
 TTCCCTTTCAGACCTCTACAGATGCCAGCCTTGTGGGAAAGAAGAGTGGGCA
 CCTGAGGGAAGCCAGACCTGCTTCCCGCGCACTGTGGTGTGTTTTGGCTTTGC
 GTGAGCACACCTCTTGGTGTCTGCTGGCAGCTAACACGCTGCTGCTGCTGC
 TGTGCTTGGGACTGCTGGCCTGTTGCTGGCACCTAGACACCCCTGTGGT
 GAGGTCAGCAGGGGGCCGCTGTGCTTTCTTATGCTGGGCTCCCTGGCAGC
 AGGTAGTGGCAGCCTCTATGGCTTCTTTGGGGAACCCACAAGGCCTGCGTG
 CTTGCTACGCCAGGCCCTCTTTGCCCTTGGTTTACCATCTTCTGTCTGCTGCC
 TGACAGTTTCGCTCATTCCAATAATCATCATCTTCAAGTTTTCCACCAAGGTA
 CCTACATTCTACCAGCCTGGGTCCAAAACCACGGTGTGGCCTGTTTGTGA
 TGATCAGCTCAGCGGCCAGCTGCTTATCTGTCTAACTTGGCTGGTGGTGTG
 GACCCCACTGCCTGCTAGGGAATACCAGCGCTTCCCCATCTGGTGATGCTT
 GAGTGCACAGAGACCAACTCCCTGGGCTTCATACTGGCCTTCTCTACAATG
 GCCTCCTCTCCATCAGTGCCTTTGCTGCTGCTACCTGGGTAAGGACTTGCC
 AGAGAACTACAACGAGGCCAAATGTGTACCTTCAGCCTGCTCTTCAACTTC
 GTGTCTGGATCGCCTTCTTACCACGGCCAGCGTCTACGACGGCAAGTAC
 CTGCCTGCGGCCAACATGATGGCTGGGCTGAGCAGCCTGAGCAGCGGCTTC
 GGTGGGTATTTTCTGCCTAAGTGTACGTGATCCTCTGCCGCCAGACCTCA
 ACAGCACAGAGCACTTCCAGCCTCCATTACAGACTACACGAGGCGCTGCG
 GCTCCACCTGACCAGTGGGTCAGCAGGCACGGCTGGCAGCCTTCTCTGCC
 TGAGGGTCGAAGTTCGAGCAGGCCGGGGTGTCCGGGAGGTCTTTGGGCA

-continued

TCGCGGTCTGGGGTTGGGACGTGTAAGCGCCTGGGAGAGCCTAGACCAGG
 CTCCGGGCTGCCAATAAAGAAGTGAAATGCGTATCTGGTCTCCTGTCTGTTGGG
 AGAGTGTGAGGTGTAACGGATTCAAGTCTGAACCCAGAGCCTGGAAAAGGC
 TGACCGCCAGATTGACGTTGCTAGGCAACTCCGGAGGCGGGCCAGCGCC
 AAAAGAACAGGGCGAGGCGTCTCCCCGCATCCCATTTGGCCGTTCTCTGCG
 GGGCCCCGCCCTCGGGGGCCGGAGCTAGAAGCTCTACGCTTCCGAGGCGC
 ACCTCCTGGCCTGCACGCTTTGACGT

hT1R1 predicted cds (SEQ ID NO 16)

ATGCTGCTCTGCACGGCTCGCCTGGTCGGCCTGCAGCTTCTCATTTCCTGCT (SEQ ID NO 16)
 GCTGGGCCTTTGCCTGCCATAGCACGGAGTCTTCTCCTGACTTCACCCCTCCC
 CGGAGATTACCTCCTGGCAGGCTGTTCCCTCTCCATTCTGGCTGTCTGCAG
 GTGAGGCACAGACCCGAGGTGACCCCTGTGTGACAGGTCTTGTAGCTTCAAT
 GAGCATGGCTACCACCTCTTCCAGGCTATGCGGCTTGGGGTTGAGGAGATA
 AACAACTCCACGGCCCTGCTGCCAACATCACCCCTGGGGTACCAGCTGTATG
 ATGTGTGTTCTGACTCTGCCAATGTGTATGCCACGCTGAGAGTGTCTCCCT
 GCCAGGGCAACACCACATAGAGCTCCAAGGAGACCTTCTCCACTATTCCCCT
 ACGGTGCTGGCAGTGATTGGGCCTGACAGCACCAACCGTGTGCCACCACA
 GCCGCCCTGCTGAGCCCTTCTCCTGGTGCCATGATTAGCTATGCGGCCAGC
 AGCGAGACGCTCAGCGTGAAGCGGCAGTATCCCTCTTCTGCGCACCATC
 CCCAATGACAAGTACCAGGTGGAGACCATGGTGTGCTGCTGCAGAAGTTC
 GGGTGGACCTGGATCTCTCTGGTTGGCAGCAGTGACGACTATGGGCAGCTA
 GGGGTGCAGGCACTGGAGAACCAGGCCACTGGTCAGGGGATCTGCATTGCT
 TTCAAGGACATCATGCCCTTCTCTGCCAGGTGGGCGATGAGAGGATGCAG
 TGCCTCATGCGCCACCTGGCCAGGCCGGGGCCACCCTCGTGGTTGTTTTT
 TCCAGCCGGCAGTTGGCCAGGTGTTTTTCGAGTCCGTGGTGTGACCAAC
 CTGACTGGCAAGGTGTGGGTCGCCTCAGAAGCCTGGGCCCTCTCCAGGCAC
 ATCACTGGGGTGCCCGGATCCAGCGCATTGGGATGGTGTGCTGGGCGTGGC
 CATCCAGAAGAGGGCTGTCCCTGGCCTGAAGGCGTTTGAAGAAGCCTATGC
 CCGGGCAGACAAGAAGGCCCTAGGCCCTGGCCACAAGGGCTCCTGGTGCAG
 CAGCAATCAGCTCTGCAGAGAATGCCAAGCTTTCATGGCACACACGATGCC
 AAGCTCAAAGCCTTCTCCATGAGTTCTGCCTACAACGCATACCGGGCTGTGT
 ATGCGGTGGCCATGGCCTCCACCAGCTCCTGGGCTGTGCCTCTGGAGCTT
 GTTCCAGGGGCGAGTCTACCCCTGGCAGCTTTTGGAGCAGATCCACAAGG
 TGCAATTCCTTCTACACAAGGACACTGTGGCGTTAATGACAACAGAGATCCC
 CTCAGTAGCTATAACATAATTGCCTGGGACTGGAATGGACCCAAGTGGACCT
 TCACGGTCCCTCGGTTCTCCACATGGTCTCCAGTTCAGCTAAACATAAATGA
 GACCAAAATCCAGTGGCACGGAAAGGACAACCAGGTGCCTAAGTCTGTGTG
 TTCCAGGACTGTCTTGAAGGGCACAGCGAGTGGTTACGGGTTTCCATCAC
 TGCTGCTTTGAGTGTGTGCCCTGTGGGGCTGGGACCTTCTCAACAAGAGT
 GACCTCTACAGATGCCAGCCTTGTGGGAAAGAAGAGTGGGCACCTGAGGGA

-continued

AGCCAGACCTGCTTCCCGCGCACTGTGGTGTGTTTTGGCTTTGCGTGAGCACA
 CCTCTTGGGTGCTGCTGGCAGCTAACACGCTGCTGCTGCTGCTGCTGCTG
 GGACTGCTGGCCTGTTTGCCTGGCACCTAGACACCCCTGTGGTGAGGTCAG
 CAGGGGGCCGCTGTGCTTCTTATGCTGGGCTCCCTGGCAGCAGGTAGTG
 GCAGCCTCTATGGCTTCTTTGGGGAACCCACAAGGCCTGCGTGCTTGCTAC
 GCCAGGCCCTCTTTGCCCTTGGTTTACCATCTTCTGCTGCTGCTGACAGT
 TCGCTCATTCCAACATAATCATCATCTTCAAGTTTTCCACCAAGGTACCTACATT
 CTACCACGCCTGGGTCCAAAACCACGGTGTGGCCTGTTTGTGATGATCAGC
 TCAGCGGCCAGCTGCTTATCTGTCTAACTTGGCTGGTGGTGTGGACCCAC
 TGCTGCTAGGGAATACCAGCGCTTCCCCATCTGGTGATGCTTGAGTGACAC
 AGAGACCAACTCCCTGGGCTTCATACTGGCCTTCTCTACAATGGCCTCCTC
 TCCATCAGTGCCTTTGCCTGCAGCTACCTGGGTAAGGACTTGCCAGAGAACT
 ACAACGAGGCCAAATGTGTACCTTACAGCGTGTCTTCAACTTCGTGTCCTG
 GATCGCCTTCTTACCACGGCCAGCGTCTACGACGGCAAGTACCTGCCTGC
 GGCCAACATGATGGCTGGGCTGAGCAGCCTGAGCAGCGGCTTCGGTGGGTA
 TTTTCTGCCTAAGTGCTACGTGATCCTCTGCCGCCAGACCTCAACAGCACA
 GAGCACTTCCAGGCCTCCATTCAGGACTACACGAGGCGCTGCGGCTCCACC
 TGA

hT1R1 conceptual translation (SEQ ID NO 17)

MLLCTARLVOLQLLISCCWAFACHSTESSPDFTLPGDYLLAGLFLPHSGCLQVRH (SEQ ID NO 17)
 RPEVTLCDRSCSFNEHGYYHLFQAMRLGVVEINNSTALLPNITLGYQLYDVCSDSA
 NVYATLRVLSLPGQHHEIQGDLHYSPTVLAVIGPDSTNRAATTAALLSPFLVPM
 ISYAASSETLSVKRQYPSFLRTIPNDKYQVETMVLQKFGWTWISLVGSSDDYG
 QLGVALENQATGQICIAFKDIMPFSQVGDERMQLMRHLAQAGATVVVVS
 SRQLARVFFESVVLTNLTGKVVVASEAWALSRHITGVPIQRIGMVLGVAIQKRA
 VPGLKAFEEAYARADKKAPRCHKGSWCSSNQLCRECQAFMAHTMPKPKAFS
 MSSAYNAYRAVYAVAHGLHQLLGCASGACSRGRVYPWQLLEQIHKVHFLHDK
 TVAFNDNRDPLSSYNI IAWDWNPKWTFVTLGSSVSPVQLNINETKIQWHGK
 DNQVPKSVCSDDCLEGHQRVVTGFHHCFCVPCGAGTFLNKSDLYRCQPCG
 KEWAPEGSQTCFPRTWFLALREHTSWVLLAANTLLLLLLGTAGLFAWHLDT
 PVVRSAGGRLCFLMLGSLAAGSGSLYGFGEPTRPACLLRQALFALGFTIFLSCL
 TVRSFQLIIIFKFSTKVPTFYHAWVQNHGAGLFVMISSAAQLLICLTWLVVWTPLP
 AREYQRFPHLVMLECTETNSLGFILAFLYNGLLSISAFACSYLGKDLPENYNEAKC
 VTFSLLFNFVSWIAFFTTASVYDGKYLPAANMMAGLSSLSGFGGYFLPKCYVIL
 CRPDLNSTEHFQASIQDYTRRCGST

Example 6—hT1R2

The predicted cds of the human ortholog of a rat taste receptor, designated rT1R2, is provided below as SEQ ID

NO 20. The conceptually-translated hT1R2 sequences is also described herein as SEQ ID NO 21. According to the present invention, the first two coding exons of hT1R2 were identified within a PAC by Southern blot. Exon 1 was isolated within a BamHI/BglII fragment that was identified

in the Southern experiment, and exon 2 was isolated within a PCR product that spans exon 1 to exon 3. Comparison of the first two coding exons to the rT1R2 sequence established that the two exons encode the N-terminus of the human counterpart to rT1R2. For example, the pairwise amino acid

identity between the hT1R2 N-terminal sequence coded by the two exons and corresponding regions of rT1R2 is approximately 72%, whereas the most related annotated sequence in public DNA sequence data banks is only approximately 48% identical to hT1R2.

hT1R2 predicted cds (SEQ ID NO 20)

ATGGGGCCCAGGGCAAAGACCATCTGCTCCCTGTTCTTCCTCCATGGGTCC (SEQ ID NO. 20)
 TGGCTGAGCCGGCTGAGAACTCGGACTTCTACCTGCCTGGGGATTACCTCC
 TGGGTGGCCTCTTCTCCCTCCATGCCAACATGAAGGGCATGTTCACCTTAA
 CTTCTGCAGGTGCCCATGTGCAAGGAGTATGAAGTGAAGGTGATAGGCTAC
 AACCTCATGCAGGCCATGCGCTTCGCGGTGGAGGAGATCAACAATGACAGC
 AGCCTGCTGCCTGGTGTGCTGCTGGGCTATGAGATCGTGGATGTGTGCTAC
 ATCTCCAACAATGTCCAGCCGGTGTCTACTTCTGGCACACGAGGACAACC
 TCCTTCCCATCCAAGAGGACTACAGTAACTACATTTCCCGTGTGGTGGCTGT
 CATTGGCCCTGACAACTCCGAGTCTGTGACTGTGGCCAACTTCTCTCC
 CTATTTCTCCTTCCACAGATCACCTACAGCGCCATCAGCGATGAGCTGCGAG
 ACAAGGTGCGCTTCCCGGCTTTGCTGCGTACCACACCAGCGCCGACCACC
 ACGTCGAGGCCATGGTGCAGCTGATGCTGCACTTCCGCTGGAAGTGGATCA
 TTGTGCTGGTGTGAGCAGCGACACCTATGGCCGCGACAATGGCCAGCTGCTTG
 GCGAGCGCTGGCCCGCGGACATCTGCATCGCCTTCCAGGAGACGCTG
 CCCACACTGCAGCCGAAGCAGAACATGACGTGAGAGGAGCGCCAGCGCCTG
 GTGACCATTTGTGGACAAGCTGCAGCAGAGCACAGCGCGCTCGTGGTCTGTG
 TTCTCGCCCGACCTGACCCTGTACCACTTCTTCAATGAGGTGCTGCGCCAGA
 ACTTCACGGGCGCCGTGTGGATCGCCTCCGAGTCTGGGCCATCGACCCGG
 TCCTGCACAACCTCACGGAGGTGGGCCACTTGGGCACCTTCTGGGCATCA
 CCATCCAGAGCGTGGCCATCCCGGCTTCAGTGAGTTCGCGAGTGGGGCC
 CACAGGCTGGGCGCCACCCCTCAGCAGGACCAGCCAGAGCTATACCTGCA
 ACCAGGAGTGCACAACCTGCCTGAACGCCACCTTGTCTTCAACACCATTCT
 CAGGCTCTGGGGAGCGTGTCTTACAGCGTGTACTCTGCGGTCTATGC
 TGTGGCCATGCCCTGCACAGCCTCCTCGGCTGTGACAAAAGCACCTGCAC
 CAAGAGGGTGGTCTACCCCTGGCAGCTGCTTGAGGAGATCTGGAAGGTCAA
 CTTCACTCTCCTGGACCACCAAATCTTCTTCGACCCGCAAGGGGACGTGGCT
 CTGCACTTGGAGATTGTCCAGTGGCAATGGGACCGGAGCCAGAATCCCTTC
 CAGAGCGTCGCCTCTACTACCCCTGCAGCGACAGCTGAAGAACATCCAA
 GACATCTCCTGGCACACCGTCAACAACACGATCCCTATGTCCATGTGTTCCA
 AGAGGTGCCAGTCAGGGCAAAGAAGAAGCCTGTGGGCATCCACGTCTGCT
 GCTTCGAGTGCATCGACTGCCTTCCCGGCACCTTCTCAACCACACTGAAGA
 TGAATATGAATGCCAGGCTGCCCGAATAACGAGTGGTCTTACCAGAGTGAG
 ACCTCCTGCTTCAAGCGGCAGCTGGTCTTCTGGAATGGCATGAGGCACCC
 ACCATCGCTGTGGCCCTGCTGGCCGCCCTGGGCTTCTCAGCACCCCTGGCC
 ATCCTGGTGATATTCTGGAGGCACTTCCAGACACCCATAGTTCGCTCGGCTG
 GGGCCCCATGTGCTTCTGATGCTGACACTGCTGCTGGTGGCATAACATGG

-continued

TGGTCCCGGTGTACGTGGGGCCGCCAAGGTCTCCACCTGCCTCTGCCGCC
 AGGCCCTCTTTCCCTCTGCTTACAATTTGCATCTCCTGTATCGCCGTGCGT
 TCTTTCCAGATCGTCTGCGCCTTCAAGATGGCCAGCCGCTTCCCACGCGCCT
 ACAGCTACTGGGTCCGCTACCAGGGGCCCTACGTCTCTATGGCATTATCAC
 GGTACTCAAAATGGTCATTGTGGTAATTGGCATGCTGGCCACGGGCCTCAGT
 CCCACCACCCGTACTGACCCCGATGACCCCAAGATCACAATTTGCTCCTGTA
 ACCCAACTACCGCAACAGCCTGCTGTTCAACACCAGCCTGGACCTGCTGCT
 CTCAGTGGTGGGTTTCAGCTTCGCCTACATGGGCAAAGAGCTGCCACCAAC
 TACAACGAGGCCAAGTTCATCACCCCTCAGCATGACCTTCTATTTACCTCATC
 CGTCTCCCTCTGCACCTTCATGTCTGCCTACAGCGGGGTGCTGGTCACCATC
 GTGGACCTCTTGGTCACTGTGCTCAACCTCCTGGCCATCAGCCTGGGCTACT
 TCGGCCCCAAGTGCTACATGATCCTCTTCTACCCGGAGCGCAACACGCCCCG
 CCTACTTCAACAGCATGATCCAGGGCTACACCATGAGGAGGGACTAG

hT1R2 conceptual translation (SEQ ID NO 21)

MGPRAKTICSLFLLWVLAEPANSDFYLPDYLGLGFLSHANMKGIVHLNFLQ (SEQ ID NO. 21)

VPMCKEYEVKVIYNLMQAMRFAVEEINNDSSLLPGVLLGYEIVDVCYISNNVQP
 VLYFLAHEDNLLPIQEDYSNYISRVAVIGPDNSEVMTVANFLSLFLLPQITYSAI
 SDELRDKVRFALLRTPSADHHVEAMVQLMLHFRWNWIIIVLVSSDTYGRDNGQ
 LLGERVARRDICIAFQETLPTLQPNQNTSEERQRLVTIVDKLQQSTARVVVVS
 PDLTLYHFFNEVLRQNFAGVWIASWAI DPVLHNLTELGHGTLGTLGITIQSVPIP
 GFSEFREWGPQAGPPPLSRTSQSYTCNQECDNCLNATLSFNITLRLSGERVVYS
 VYSAVYAVAHALHSLGCDKSTCTKRVPVWPQLLEE IWKVNFTLLDHQIFFDPPQG
 DVALHLEIVQWQDRSQNPFQSVASYPLQRQLKNIQDISWHTVNNTIPMSMC
 SKRCQSGQKKKPVGIHVCCFECIDCLPGTFLNHTEDEYEQACPNNEWSYQSE
 TSCFKRQLVFLWHEAPTIAVALLAALGFLSTLAILVIFWRHFQTPIVRSAGGPMC
 FLMLTLLLVAYMVVPVYVGGPKVSTCLCRQALFPLCFTICISCIAVRSFQIVCAFKM
 ASRFPRAYSYWVRYQGPYVSMAFITVLKMIIVIGMLATGLSPTTRTDPDDPKITI
 VSCNPNYRNSLLFNTSLDLLSVVGFSAFYMKGELPTNYNEAKFITLSMTFYFTSS
 VSLCTFMSAYSGLVLTIVDLLVTVLNLLAISLGYFGPKCYMILFYPERNTPAYFNS
 MIQGYTMRRD

Example 7

Methods for Heterologous Expression of T1Rs in Heterologous Cells

An HEK-293 derivative (Chandrashekar et al., Cell 100 (6): 703–11 (2000)), which stably expresses Gα15, was grown and maintained at 37° C. in Dulbecco's Modified Eagle Medium (DMEM, Gibco BRL) supplemented with 10% FBS, MEM non-essential amino acids (Gibco BRL), and 3 μg/ml blasticidin. For calcium-imaging experiments, cells were first seeded onto 24-well tissue-culture plates (approximately 0.1 million cells per well), and transfected by lipofection with Mirus Transit-293 (PanVera). To minimize glutamate-induced and glucose-induced desensitization, supplemented DMEM was replaced with low-glucose

DMEM/GlutaMAX (Gibco BRL) approximately 24 hours after transfection. 24 hours later, cells were loaded with the calcium dye Fluo-4 (Molecular Probes), 31 μM in Dulbecco's PBS buffer (DPBS, GibcoBRL), for 1.5 hours at room temperature. After replacement with 250 μl DPBS, stimulation was performed at room temperature by addition of 200 μl DPIBS supplemented with taste stimuli. Calcium mobilization was monitored on a Axiovert S100 TV microscope (Zeiss) using Imaging Workbench 4.0 software (Axon). T1R1/T1R3 and T1R2/T1R3 responses were strikingly transient—calcium increases rarely persisted longer than 15 seconds—and asynchronous. The number of responding cells was thus relatively constant over time; therefore, cell responses were quantitated by manually counting the number of responding cells at a fixed time point, typically 30 seconds after stimulus addition.

Example 8

Human T1R2/T1R3 Functions as a Sweet Taste Receptor

HEK cells stably expressing $G_{\alpha 15}$ were transiently transfected with human T1R2, T1R3 and T1R2/T1R3, and assayed for increases in intracellular calcium in response to increasing concentrations of sucrose (FIG. 1(a)). Also, T1R2/T1R3 dose responses were determined for several sweet taste stimuli (FIG. 1(b)). The maximal percentage of responding cells was different for different sweeteners, ranging from 10–30%. For clarity, dose responses were normalized to the maximal percentage of responding cells. The values in FIG. 1 represent the mean \pm s.e. of four independent responses. X-axis circles mark psychophysical detection thresholds determined by taste testing. Gurmarin (50-full dilution of a filtered 10 g/l *Gymnema sylvestre* aqueous extract) inhibited the response of T1R2/T1R3 to 250 mM sucrose, but not the response of endogenous β 2-adrenergic receptor to 20 μ M isoproterenol (FIG. 1(b)). FIG. 1(c) contains the normalized response of T1R2/T1R3 co-expressing cell lines to different sweeteners (sucrose, aspartame, tryptophan and saccharin)

Example 9

Rat T1R2/T1R3 Also Functions as a Sweet Taste Receptor

HEK cells stably expressing $G_{\alpha 15}$ were transiently transfected with hT1R2/hT1R3, rT1R2/rT1R3, hT1R2/rT1R3, and rT1R2/hT1R3. These transfected cells were then assayed for increased intracellular calcium in response to 350 mM sucrose, 25 mM tryptophan, 15 mM aspartame, and 0.05 of monellin. The results with sucrose and aspartame are contained in FIG. 2 and indicate that rT1R2/rT1R3 also functions as a sweet taste receptor. Also, these results suggest that T1R2 may control T1R2/T1R3 ligand specificity.

Example 10

Human T1R1/T1R3 Functions as Umami Taste Receptors

HEK cells stably expressing $G_{\alpha 15}$ were transiently transfected with human T1R1, T1R3 and T1R1/T1R3 and assayed for increases in intracellular calcium in response to increasing concentrations of glutamate (FIG. 3(a) and 0.5 mM glutamate), 0.2 mM IMP, and 0.5 mM glutamate plus 0.2 mM IMP (FIG. 3(b)). Human T1R1/T1R3 dose responses were determined for glutamate in the presence and absence of 0.2 mM IMP (FIG. 3(c)). The maximal percentages of responding cells was approximately 5% for glutamate and approximately 10% for glutamate plus IMP. For clarity, does responses are normalized to the maximal percentage of responding cells. The values represent the mean \pm s.e. of four independent responses. X-axis circles mark taste detection thresholds determined by taste testing.

Example 11

PDZIP as an Export Sequence

The six residue PDZIP sequence (SVSTW (SEQ ID NO:22)) was fused to the C-terminus of hT1R2 and the

chimeric receptor (i.e. hT1R2-PDZIP) was transfected into an HEK-293 host cell. The surface expression of hT1R2 was then monitored using immunofluorescence and FACS scanning data. As shown in FIGS. 6A and 6B, the inclusion of the PDZIP sequence increased the surface expression of hT1R2-PDZIP relative to hT1R2.

PKZIP Sequence

SVSTW (SEQ ID NO:22)

More specifically, FIG. 4A shows an immunofluorescence staining of myc-tagged hT1R2 demonstrating that PDZIP significantly increases the amount of hT1R2 protein on the plasma membrane. FIG. 4B shows FACS analysis data demonstrating the same result.—Cells expressing myc-tagged hT1R2 are indicated by the dotted line and cells expressing myc-tagged hT1R2-PDZIP are indicated by the solid line.

Example 12

Generation of Cell Lines that Stably Co-Express T1R1/T1R3 or T1R2/T1R3

Human cell lines that stably co-express human T1R2/T1R3 or human T1R1/T1R3 were generated by transfecting linearized PEAK10-derived (Edge Biosystems) vectors containing pCDNA 3.1/ZEO-derived (Invitrogen) vectors respectively containing hT1R1 or hT1R2 expression construct (plasmid SAV2485 for T1R1, SAV2486 for T1R2) and hT1R3 (plasmid SXV550 for T1R3) into a $G_{\alpha 15}$ expressing cell line. Specifically, T1R2/T1R3 stable cell lines were produced by co-transfecting linearized SAV2486 and SXV550 into Aurora Bioscience's HEK-293 cell line that stably expresses $G_{\alpha 15}$. T1R1/T1R3 stable cell lines were produced by co-transfecting linearized SAV2485 and SXV550 into the same HEK-293 cell line that stably expresses $G_{\alpha 15}$. Following SAV2485/SCV550 and SAV2486/SXV550 transfections, puromycin-resistant and zeocin-resistant colonies were selected, expanded, and tested by calcium imaging for responses to sweet or umami taste stimuli. Cells were selected in 0.0005 mg/ml puromycin (CALBIOCHEM) and 0.1 mg/ml zeocin (Invitrogen) at 37° C. in low-glucose DMEM supplemented with GlutaMAX, 10% dialyzed FBS, and 0.003 mg/ml blasticidin. Resistant colonies were expanded, and their responses to sweet taste stimuli evaluated by Fluorescence microscopy. For automated fluorimetric imaging on VIPR-II instrumentation (Aurora Biosciences), T1R2/T1R3 stable cells were first seeded onto 96-well plates (approximately 15,000 cells per well). Twenty-four hours later, cells were loaded with the calcium dye fluo-3-AM (Molecular Probes), 0.005 mM in PBS, for one hour at room temperature. After replacement with 70 ml PBS, stimulation was performed at room temperature by addition of 70 ml PBS supplemented with taste stimuli. Fluorescence (480 nm excitation and 535 nm emission) responses from 20 to 30 seconds following compound addition were averaged, corrected for background fluorescence measured prior to compound addition, and normalized to the response to 0.001 mM ionomycin (CALBIOCHEM), a calcium ionophore.

It was then observed that when these cell lines were contacted with sweet or umami, that for active clones typically 80–100% of cells responded to taste stimuli. Unexpectedly, the magnitude of individual cell responses was markedly larger than that of transiently transfected cells.

Based on this observation, the inventors tested the activity of T1R stable cell lines by automated fluorescence imaging

using Aurora Bioscience's VIPR instrumentation as described above. The responses of two T1R1/T1R3 and one T1R2/T1R3 cell line are shown in FIG. 5 and FIG. 6 respectively.

Remarkably, the combination of increased numbers of responding cells and increased response magnitudes resulted in a greater than 10-fold increase in activity relative to transiently transfected cells. (By way of comparison, the percent ionomycin response for cells transiently transfected with T1R2/T1R3 was approximately 5% under optimal conditions.) Moreover, dose responses obtained for stably expressed human T1R2/T1R3 and T1R1/T1R3 correlated

with human taste detection thresholds. The robust T1R activity of these stable cell lines suggests that they are well suited for use in high-throughput screening of chemical libraries in order to identify compounds, e.g. small molecules, that modulate the sweet or umami taste receptor and which therefore modulate, enhance, block or mimic sweet or umami taste.

While the foregoing detailed description has described several embodiments of the present invention, it is to be understood that the above description is illustrative only and not limiting of the disclosed invention. The invention is to be limited only by the claims which follow.

 SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 24

<210> SEQ ID NO 1

<211> LENGTH: 876

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

agcctggcag tggcctcagg cagagtctga cgcgcacaaa ctttcaggcc caggaagcga      60
ggacaccact ggggccccag ggtgtggcaa gtgaggatgg caagggtttt gctaaacaaa      120
tcctctgccc gctccccgcc ccgggctcac tccatgtgag gccccagtcg gggcagccac      180
ctgccgtgcc tgttggaagt tgccctgcc atgctgggcc ctgctgtcct gggcctcagc      240
ctctgggctc tcctgcacc tgggacgggg gccccattgt gcctgtcaca gcaacttagg      300
atgaaggggg actacgtgct gggggggctg ttccccctgg gcgaggccga ggaggctggc      360
ctccgcagcc ggacacggcc cagcagccct gtgtgcacca ggtacagagg tgggacggcc      420
tgggtcgggg tcaggtgac caggtctggg gtgtcctga gctggggccg aggtggccat      480
ctgcggttct gtgtggcccc aggttctcct caaacggcct gctctgggca ctggccatga      540
aatggccgt ggaggagatc aacaacaagt cggatctgct gcccggtg cgccctgggct      600
acgacctctt tgatacgtgc tcggagcctg tgggtggccat gaagcccagc ctcatgttcc      660
tggccaaggg aggcagccgc gacatcgccg cctactgcaa ctacacgcag taccagcccc      720
gtgtgctggc tgtcatcggg cccactcgt cagagctcgc catggtcacc ggcaagttct      780
tcagcttctt cctcatgccc cagtggggcg cccccacca tcaccaccc ccaaccaacc      840
cctgccccgt gggagcccct tgtgtcagga gaatgc      876
  
```

<210> SEQ ID NO 2

<211> LENGTH: 2687

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

```

tacatgcacc ccaccagcc ctgccctggg agccctgtgt cagaagatgc tcttggcctt      60
gcaggtcagc tacggtgcta gcatggagct gctgagcgcc cgggagacct tccccctctt      120
cttccgcacc gtgccagcg accgtgtgca gctgacggcc gccgcggagc tgctgcagga      180
gttcggctgg aactgggtgg ccgccctggg cagcgcagac gagtacggcc ggcagggcct      240
gagcatcttc tcggccctgg ccgcggcacg cggcatctgc atcgcgcacg agggcctggg      300
gccgctgccc cgtgccgatg actcgcggct ggggaaggtg caggacgtcc tgcaccaggt      360
gaaccagagc agcgtgcagg tgggtgctgct gttcgctcc gtgcacgccg cccacgcctt      420
  
```

-continued

cttcaactac agcatcagca gcaggctctc gcccaagggtg tgggtggcca gcgaggcctg	480
gctgacctct gacctggtca tggggctgcc cggcatggcc cagatgggca cggtgcttgg	540
cttcctccag aggggtgcc agctgcacga gttccccag tacgtgaaga cgcacctggc	600
cctggccacc gaccggcct tctgctctgc cctgggcgag agggagcagg gtctggagga	660
ggacgtggtg ggccagcgt gcccgcagtg tgactgcac acgctgcaga acgtgagcgc	720
agggctaaat caccaccaga cgttctctgt ctacgcagct gtgtatagcg tggcccaggc	780
cctgcacaac actcttcagt gcaacgcctc aggtgcccc gcgcaggacc ccgtgaagcc	840
ctggcagggtg agcccgggag atgggggtgt gctgtcctct gcatgtgccc aggccaccag	900
gcacggccac cacgcctgag ctggagggtg ctggcggctc agccccgtcc cccgcccga	960
gctcctggag aacatgtaca acctgacctt ccacgtgggc gggctgccgc tgcggttcga	1020
cagcagcggg aacgtggaca tggagtacga cctgaagctg tgggtgtggc agggctcagt	1080
gcccaggctc cacgacgtgg gcaggttaa cggcagcctc aggacagagc gcctgaagat	1140
ccgctggcac acgtctgaca accaggtgag gtgaggggtg gtgtgccagg cgtgcccgtg	1200
gtagcccccg cggcagggcg cagcctgggg gtgggggccc ttccagtctc ccgtgggcat	1260
gccagccga gcagagccag accccaggcc tgtgcgcaga agcccgtgc ccggtgctcg	1320
cggcagtgcc aggagggcca ggtgcgccgg gtcaaggggt tccactcctg ctgctacgac	1380
tgtgtggact gcgagggcgg cagctaccgg caaaaccag gtgagccgcc ttcccggcag	1440
gcgggggtgg gaacgcagca ggggagggtc ctgccaagtc ctgactctga gaccagagcc	1500
cacaggttac aagacgaaca cccagcgcctt ttctcctctc tcacagacga catgcctgc	1560
accttttgtg gccaggatga gtggtccccg gagcgaagca cacgctgctt ccgccgagc	1620
tctcggttcc tggcatgggg cgagccggct gtgctgctgc tgctcctgct gctgagcctg	1680
gcgctgggccc ttgtgctggc tgctttgggg ctgttcgttc accatcggga cagcccactg	1740
gttcaggcct cgggggggccc cctggcctgc tttggcctgg tgtgcctggg cctggtctgc	1800
ctcagcgtcc tcctgttccc tggccagccc agccctgccc gatgcctggc ccagcagccc	1860
ttgtcccacc tcccgtcac gggctgcctg agcacactct tcctgcaggc ggccgagatc	1920
ttcgtggagt cagaactgcc tctgagctgg gcagaccggc tgagtggctg cctgcggggg	1980
ccctgggcct ggctggtggt gctgctggcc atgctggtgg aggtcgcact gtgcacctgg	2040
tacctggtgg ccttcccgcc ggaggtggtg acggactggc acatgctgcc cacggaggcg	2100
ctggtgcact gccgcacacg ctctgggtc agcttcggcc tagcgcacgc caccaatgcc	2160
acgctggcct ttctctgctt cctgggcact ttctggtgc ggagccagcc gggctgctac	2220
aaccgtgccc gtggcctcac ctttgccatg ctggcctact tcatcacctg ggtctccttt	2280
gtgcccctcc tggccaatgt gcaggtggtc ctcaggcccc cgtgcagat gggcgccctc	2340
ctgctctgtg tcctgggcat cctggctgcc ttccacctgc ccaggtgta cctgctcatg	2400
cggcagccag ggctcaacac ccccagttc ttctgggag ggggccctgg ggatgccaa	2460
ggccagaatg acgggaacac aggaaatcag gggaaacatg agtgaccaa ccctgtgatc	2520
tcagccccgg tgaaccaga cttagctgcg atcccccca agccagcaat gaccctgtc	2580
tcgctacaga gaccctccc ctctaggttc tgaccccagg ttgtctcctg acctgacct	2640
cacagtgagc cctaggcctg gagcacgtgg acaccctgt gaccatc	2687

-continued

<211> LENGTH: 2559

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 3

atgctgggcc ctgctgtcct gggcctcagc ctctgggctc tcctgcaccc tgggacgggg 60
gccccattgt gcctgtcaca gcaacttagg atgaaggggg actacgtgct gggggggctg 120
ttccccctgg gcgaggccga ggaggctggc ctccgcagcc ggacacggcc cagcagccct 180
gtgtgcacca ggttctcctc aaacggcctg ctctgggcac tggccatgaa aatggccgtg 240
gaggagatca acaacaagtc ggatctgctg cccgggctgc gcctgggcta cgacctcttt 300
gatacgtgct cggagcctgt ggtggccatg aagcccagcc tcatgttcct ggccaaggca 360
ggcagccgcg acatcgccgc ctactgcaac tacacgcagt accagccccg tgtgctggct 420
gtcatcgggc cccactcgtc agagctcgcc atggtcaccg gcaagttctt cagcttcttc 480
ctcatgcccc aggtcagcta cggtgctagc atggagctgc tgagcgcccg ggagaccttc 540
ccctccttct tccgcaccgt gcccagcgac cgtgtgcagc tgacggccgc cggggagctg 600
ctgcaggagt tcggctggaa ctgggtggcc gccctgggca gcgacgacga gtacggccgg 660
cagggcctga gcatcttctc ggccctggcc gcggcaccg gcacctgcat cgcgcacgag 720
ggcctggtgc cgctgccccg tgccgatgac tcgcggctgg ggaaggtgca ggacgtcctg 780
caccaggatga accagagcag cgtgcaggtg gtgctgctgt tcgcctccgt gcacgcccgc 840
cacgccctct tcaactacag catcagcagc aggctctcgc ccaaggtgtg ggtggccagc 900
gaggcctggc tgacctctga cctggatcatg gggctgcccg gcatggccca gatgggcacg 960
gtgcttggct tcctccagag gggtgcccag ctgcacgagt tccccagta cgtgaagacg 1020
cacctggccc tggccaccga cccggccttc tgctctgccc tgggcgagag ggagcagggg 1080
ctggaggagg acgtggtggg ccagcgctgc ccgcagtggt actgcatcac gctgcagaac 1140
gtgagcgagc ggctaaatca ccaccagacg ttctctgtct acgcagctgt gtatagcgtg 1200
gcccaggccc tgcacaacac tcttcagtgc aacgcctcag gctgccccgc gcaggacccc 1260
gtgaagccct ggcagctcct ggagaacatg tacaacctga ccttccacgt gggcgggctg 1320
ccgctgcggg tcgacagcag cggaaacgtg gacatggagt acgacctgaa gctgtgggtg 1380
tggcagggct cagtgccag gctccacgac gtgggcagggt tcaacggcag cctcaggaca 1440
gagcgcctga agatccgctg gcacacgtct gacaaccaga agcccgtgtc ccggtgctcg 1500
cggcagtgcc aggagggcca ggtgcgcccg gtcaaggggt tccactcctg ctgctacgac 1560
tgtgtggact gcgaggcggg cagctaccgg caaaaccag acgacatcgc ctgcaccttt 1620
tgtggccagg atgagtgtc cccggagcga agcacacgct gcttccgccg caggtctcgg 1680
ttcctggcat ggggcgagcc ggctgtgctg ctgctgctcc tgctgctgag cctggcgctg 1740
ggccttgtgc tggctgcttt ggggctgttc gttcaccatc gggacagccc actggttcag 1800
gcctcggggg ggcccctggc ctgctttggc ctggtgtgcc tgggcctggt ctgcctcagc 1860
gtcctcctgt tccctggcca gcccagccct gcccgatgcc tggcccagca gcccttgtcc 1920
cacctcccgc tcacgggctg cctgagcaca ctcttctcgc aggcggccga gatcttcgtg 1980
gagtcagaac tgccctctgag ctgggcagac cggctgagtg gctgcctgcg ggggccctgg 2040
gcctggctgg tggctgctgct ggccatgctg gtggaggtcg cactgtgcac ctggtacctg 2100
gtggccttcc cgccggagggt ggtgacggac tggcacatgc tgcccacgga ggcgctgggtg 2160
cactgcccga cacgctcctg ggtcagcttc ggcttagcgc acgccaccaa tgccacgctg 2220

-continued

```

gcctttctct gcttctctggg cactttctctg gtgcggagcc agccgggctg ctacaaccgt 2280
gcccgtaggc tcaccttgc catgctggcc tacttcatca cctgggtctc ctttgtgccc 2340
ctcctggcca atgtgcaggt ggctctcagg cccgccgtgc agatgggggc cctcctgctc 2400
tgtgtcctgg gcatcctggc tgccttccac ctgcccaggt gttacctgct catgcccag 2460
ccagggtcca acacccccga gttcttctctg ggagggggcc ctggggatgc ccaaggccag 2520
aatgacggga acacaggaaa tcaggggaaa catgagtga 2559

```

```

<210> SEQ ID NO 4
<211> LENGTH: 852
<212> TYPE: PRT
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 4

```

```

Met Leu Gly Pro Ala Val Leu Gly Leu Ser Leu Trp Ala Leu Leu His
 1                    5                    10                    15
Pro Gly Thr Gly Ala Pro Leu Cys Leu Ser Gln Gln Leu Arg Met Lys
                20                    25                    30
Gly Asp Tyr Val Leu Gly Gly Leu Phe Pro Leu Gly Glu Ala Glu Glu
          35                    40                    45
Ala Gly Leu Arg Ser Arg Thr Arg Pro Ser Ser Pro Val Cys Thr Arg
          50                    55                    60
Phe Ser Ser Asn Gly Leu Leu Trp Ala Leu Ala Met Lys Met Ala Val
          65                    70                    75                    80
Glu Glu Ile Asn Asn Lys Ser Asp Leu Leu Pro Gly Leu Arg Leu Gly
                85                    90                    95
Tyr Asp Leu Phe Asp Thr Cys Ser Glu Pro Val Val Ala Met Lys Pro
          100                    105                    110
Ser Leu Met Phe Leu Ala Lys Ala Gly Ser Arg Asp Ile Ala Ala Tyr
          115                    120                    125
Cys Asn Tyr Thr Gln Tyr Gln Pro Arg Val Leu Ala Val Ile Gly Pro
          130                    135                    140
His Ser Ser Glu Leu Ala Met Val Thr Gly Lys Phe Phe Ser Phe Phe
          145                    150                    155                    160
Leu Met Pro Gln Val Ser Tyr Gly Ala Ser Met Glu Leu Leu Ser Ala
                165                    170                    175
Arg Glu Thr Phe Pro Ser Phe Phe Arg Thr Val Pro Ser Asp Arg Val
          180                    185                    190
Gln Leu Thr Ala Ala Ala Glu Leu Leu Gln Glu Phe Gly Trp Asn Trp
          195                    200                    205
Val Ala Ala Leu Gly Ser Asp Asp Glu Tyr Gly Arg Gln Gly Leu Ser
          210                    215                    220
Ile Phe Ser Ala Leu Ala Ala Ala Arg Gly Ile Cys Ile Ala His Glu
          225                    230                    235                    240
Gly Leu Val Pro Leu Pro Arg Ala Asp Asp Ser Arg Leu Gly Lys Val
                245                    250                    255
Gln Asp Val Leu His Gln Val Asn Gln Ser Ser Val Gln Val Val Leu
          260                    265                    270
Leu Phe Ala Ser Val His Ala Ala His Ala Leu Phe Asn Tyr Ser Ile
          275                    280                    285
Ser Ser Arg Leu Ser Pro Lys Val Trp Val Ala Ser Glu Ala Trp Leu
          290                    295                    300
Thr Ser Asp Leu Val Met Gly Leu Pro Gly Met Ala Gln Met Gly Thr

```

-continued

305	310	315	320
Val Leu Gly Phe Leu Gln Arg Gly Ala Gln Leu His Glu Phe Pro Gln 325 330 335			
Tyr Val Lys Thr His Leu Ala Leu Ala Thr Asp Pro Ala Phe Cys Ser 340 345 350			
Ala Leu Gly Glu Arg Glu Gln Gly Leu Glu Glu Asp Val Val Gly Gln 355 360 365			
Arg Cys Pro Gln Cys Asp Cys Ile Thr Leu Gln Asn Val Ser Ala Gly 370 375 380			
Leu Asn His His Gln Thr Phe Ser Val Tyr Ala Ala Val Tyr Ser Val 385 390 395 400			
Ala Gln Ala Leu His Asn Thr Leu Gln Cys Asn Ala Ser Gly Cys Pro 405 410 415			
Ala Gln Asp Pro Val Lys Pro Trp Gln Leu Leu Glu Asn Met Tyr Asn 420 425 430			
Leu Thr Phe His Val Gly Gly Leu Pro Leu Arg Phe Asp Ser Ser Gly 435 440 445			
Asn Val Asp Met Glu Tyr Asp Leu Lys Leu Trp Val Trp Gln Gly Ser 450 455 460			
Val Pro Arg Leu His Asp Val Gly Arg Phe Asn Gly Ser Leu Arg Thr 465 470 475 480			
Glu Arg Leu Lys Ile Arg Trp His Thr Ser Asp Asn Gln Lys Pro Val 485 490 495			
Ser Arg Cys Ser Arg Gln Cys Gln Glu Gly Gln Val Arg Arg Val Lys 500 505 510			
Gly Phe His Ser Cys Cys Tyr Asp Cys Val Asp Cys Glu Ala Gly Ser 515 520 525			
Tyr Arg Gln Asn Pro Asp Asp Ile Ala Cys Thr Phe Cys Gly Gln Asp 530 535 540			
Glu Trp Ser Pro Glu Arg Ser Thr Arg Cys Phe Arg Arg Arg Ser Arg 545 550 555 560			
Phe Leu Ala Trp Gly Glu Pro Ala Val Leu Leu Leu Leu Leu Leu Leu 565 570 575			
Ser Leu Ala Leu Gly Leu Val Leu Ala Ala Leu Gly Leu Phe Val His 580 585 590			
His Arg Asp Ser Pro Leu Val Gln Ala Ser Gly Gly Pro Leu Ala Cys 595 600 605			
Phe Gly Leu Val Cys Leu Gly Leu Val Cys Leu Ser Val Leu Leu Phe 610 615 620			
Pro Gly Gln Pro Ser Pro Ala Arg Cys Leu Ala Gln Gln Pro Leu Ser 625 630 635 640			
His Leu Pro Leu Thr Gly Cys Leu Ser Thr Leu Phe Leu Gln Ala Ala 645 650 655			
Glu Ile Phe Val Glu Ser Glu Leu Pro Leu Ser Trp Ala Asp Arg Leu 660 665 670			
Ser Gly Cys Leu Arg Gly Pro Trp Ala Trp Leu Val Val Leu Leu Ala 675 680 685			
Met Leu Val Glu Val Ala Leu Cys Thr Trp Tyr Leu Val Ala Phe Pro 690 695 700			
Pro Glu Val Val Thr Asp Trp His Met Leu Pro Thr Glu Ala Leu Val 705 710 715 720			
His Cys Arg Thr Arg Ser Trp Val Ser Phe Gly Leu Ala His Ala Thr 725 730 735			

-continued

Asn Ala Thr Leu Ala Phe Leu Cys Phe Leu Gly Thr Phe Leu Val Arg
740 745 750

Ser Gln Pro Gly Cys Tyr Asn Arg Ala Arg Gly Leu Thr Phe Ala Met
755 760 765

Leu Ala Tyr Phe Ile Thr Trp Val Ser Phe Val Pro Leu Leu Ala Asn
770 775 780

Val Gln Val Val Leu Arg Pro Ala Val Gln Met Gly Ala Leu Leu Leu
785 790 795 800

Cys Val Leu Gly Ile Leu Ala Ala Phe His Leu Pro Arg Cys Tyr Leu
805 810 815

Leu Met Arg Gln Pro Gly Leu Asn Thr Pro Glu Phe Phe Leu Gly Gly
820 825 830

Gly Pro Gly Asp Ala Gln Gly Gln Asn Asp Gly Asn Thr Gly Asn Gln
835 840 845

Gly Lys His Glu
850

<210> SEQ ID NO 5
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (3)
<223> OTHER INFORMATION: a, t, c, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (9)
<223> OTHER INFORMATION: a, t, c, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (12)
<223> OTHER INFORMATION: a, t, c, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (18)
<223> OTHER INFORMATION: a, t, c, g, other or unknown

<400> SEQUENCE: 5

cgnttyytng cntgggnga rcc

23

<210> SEQ ID NO 6
<211> LENGTH: 23
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Description of Artificial Sequence: Primer
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (3)
<223> OTHER INFORMATION: a, t, c, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (6)
<223> OTHER INFORMATION: a, t, c, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (18)
<223> OTHER INFORMATION: a, t, c, g, other or unknown
<220> FEATURE:
<221> NAME/KEY: modified_base
<222> LOCATION: (21)
<223> OTHER INFORMATION: a, t, c, g, other or unknown

<400> SEQUENCE: 6

-continued

cgngcncgrt trtarcanc cc ngg

23

<210> SEQ ID NO 7
 <211> LENGTH: 9
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 7

Arg Phe Leu Ala Trp Gly Glu Pro Ala
 1 5

<210> SEQ ID NO 8
 <211> LENGTH: 8
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 8

Pro Gly Cys Tyr Asn Arg Ala Arg
 1 5

<210> SEQ ID NO 9
 <211> LENGTH: 552
 <212> TYPE: DNA
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 9

gtgctgtcac tcctcctgct gctttgacct gtgctgggtc tagcactggc tgctctgggg 60
 ctctctgtcc accactggga cagccctctt gtccaggcct caggcggctc acagttctgc 120
 tttggcctga tctgcctagg cctcttctgc ctcaagtgtc ttctgttccc aggacggcca 180
 agctctgcca gctgccttgc acaacaacca atggctcacc tccctctcac aggctgcctg 240
 agcacactct tcctgcaagc agctgagacc tttgtggagt ctgagctgcc actgagctgg 300
 gcaaactggc tatgcagcta ccttcgggac tctggcctgc tagtggtact gttggccact 360
 tttgtggagg cagcactatg tgcttggtat ttgaccgctt caccagaagt ggtgacagac 420
 tggtcagtgc tgcccacaga ggtactggag cactgccacg tgcgttcctg ggtcaacctg 480
 ggcttggtgc acatcaccaa tgcaatggta gcttttctct gctttctggg cactttcctg 540
 gtacaagacc ag 552

<210> SEQ ID NO 10
 <211> LENGTH: 184
 <212> TYPE: PRT
 <213> ORGANISM: Mus sp.

<400> SEQUENCE: 10

Val Leu Ser Leu Leu Leu Leu Cys Leu Val Leu Gly Leu Ala Leu
 1 5 10 15
 Ala Ala Leu Gly Leu Ser Val His His Trp Asp Ser Pro Leu Val Gln
 20 25 30
 Ala Ser Gly Gly Ser Gln Phe Cys Phe Gly Leu Ile Cys Leu Gly Leu
 35 40 45
 Phe Cys Leu Ser Val Leu Leu Phe Pro Gly Arg Pro Ser Ser Ala Ser
 50 55 60
 Cys Leu Ala Gln Gln Pro Met Ala His Leu Pro Leu Thr Gly Cys Leu
 65 70 75 80
 Ser Thr Leu Phe Leu Gln Ala Ala Glu Thr Phe Val Glu Ser Glu Leu
 85 90 95
 Pro Leu Ser Trp Ala Asn Trp Leu Cys Ser Tyr Leu Arg Asp Ser Gly

-continued

100	105	110
Leu Leu Val Val Leu Leu Ala Thr Phe Val Glu Ala Ala Leu Cys Ala		
115	120	125
Trp Tyr Leu Thr Ala Ser Pro Glu Val Val Thr Asp Trp Ser Val Leu		
130	135	140
Pro Thr Glu Val Leu Glu His Cys His Val Arg Ser Trp Val Asn Leu		
145	150	155
Gly Leu Val His Ile Thr Asn Ala Met Val Ala Phe Leu Cys Phe Leu		
165	170	175
Gly Thr Phe Leu Val Gln Asp Gln		
180		

<210> SEQ ID NO 11
 <211> LENGTH: 558
 <212> TYPE: DNA
 <213> ORGANISM: Rattus sp.

<400> SEQUENCE: 11

```

gtgctgtcac ttctcctgct gctttgcctg gtgctgggcc tgacactggc tgccctgggg      60
ctctttgtcc actactggga cagccctctt gttcaggcct cagggtgggtc actgttctgc    120
tttggcctga tctgcctagg cctcttctgc ctcagtgtcc ttctgttccc aggacgacca    180
cgctctgcca gctgccttgc ccaacaacca atggctcacc tccctctcac aggctgcctg    240
agcacactct tcctgcaagc agccgagatc tttgtggagt ctgagctgcc actgagttgg    300
gcaaactggc tctgcagcta ccttcggggc ccctgggctt ggctgggtgt actgctggcc    360
actcttgggg aggctgcact atgtgcctgg tacttggatg ctttcctcc agaggtgggtg    420
acagattggc aggtgctgcc cacggaggta ctggaacact gccgcatgcg ttctggggtc    480
agcctgggct tgggtgcacat caccaatgca ggggtagctt tcctctgctt tctgggcact    540
ttcctggtac aaagccag                                     558
    
```

<210> SEQ ID NO 12
 <211> LENGTH: 186
 <212> TYPE: PRT
 <213> ORGANISM: Rattus sp.

<400> SEQUENCE: 12

Val Leu Ser Leu Leu Leu Leu Cys Leu Val Leu Gly Leu Thr Leu		
1	5	10
Ala Ala Leu Gly Leu Phe Val His Tyr Trp Asp Ser Pro Leu Val Gln		
20	25	30
Ala Ser Gly Gly Ser Leu Phe Cys Phe Gly Leu Ile Cys Leu Gly Leu		
35	40	45
Phe Cys Leu Ser Val Leu Leu Phe Pro Gly Arg Pro Arg Ser Ala Ser		
50	55	60
Cys Leu Ala Gln Gln Pro Met Ala His Leu Pro Leu Thr Gly Cys Leu		
65	70	75
Ser Thr Leu Phe Leu Gln Ala Ala Glu Ile Phe Val Glu Ser Glu Leu		
85	90	95
Pro Leu Ser Trp Ala Asn Trp Leu Cys Ser Tyr Leu Arg Gly Pro Trp		
100	105	110
Ala Trp Leu Val Val Leu Leu Ala Thr Leu Val Glu Ala Ala Leu Cys		
115	120	125
Ala Trp Tyr Leu Met Ala Phe Pro Pro Glu Val Val Thr Asp Trp Gln		
130	135	140

-continued

Val Leu Pro Thr Glu Val Leu Glu His Cys Arg Met Arg Ser Trp Val
 145 150 155 160

Ser Leu Gly Leu Val His Ile Thr Asn Ala Gly Val Ala Phe Leu Cys
 165 170 175

Phe Leu Gly Thr Phe Leu Val Gln Ser Gln
 180 185

<210> SEQ ID NO 13
 <211> LENGTH: 2577
 <212> TYPE: DNA
 <213> ORGANISM: Rattus sp.

<400> SEQUENCE: 13

atgccggggtt tggctatcctt gggcctcagt ctggctgctt tcctggagct tgggatgggg 60

tcctctttgt gtctgtcaca gcaattcaag gcacaagggg actatatatt gggatggacta 120

ttccccctgg gcacaactga ggaggccact ctcaaccaga gaacacagcc caacggcatc 180

ctatgtacca ggttctcgcc ccttggtttg ttcctggcca tggctatgaa gatggctgta 240

gaggagatca acaatggatc tgccttgctc cctgggctgc gactgggcta tgacctgttt 300

gacacatgct cagagccagt ggtcaccatg aagcccagcc tcatgttcat ggccaagggtg 360

ggaagtcaaa gcattgctgc ctactgcaac tacacacagt accaaccctg tgtgctggct 420

gtcattggtc cccactcatc agagcttgcc ctattacag gcaagttctt cagcttcttc 480

ctcatgccac aggtcagcta tagtgccagc atggatcggc taagtgaccg ggaaacattt 540

ccatccttct tccgcacagt gccagtgac cgggtgcagc tgcaggccgt tgtgacactg 600

ttgcagaatt tcagctggaa ctgggtggct gccttaggta gtgatgatga ctatggccgg 660

gaaggctctga gcatcttttc tggctctggc aactcacgag gtatctgcat tgcacacgag 720

ggcctggtgc cacaacatga cactagtggc caacaattgg gcaagggtgt ggatgtgcta 780

cgccaagtga accaaagcaa agtacaggtg gtggtgctgt ttgcatctgc ccgtgctgtc 840

tactcccttt ttagctacag catccttcat gacctctcac ccaaggtatg ggtggccagt 900

gagtccctggc tgacctctga cctggctcatg aacttccca atattgccc tgtgggact 960

gttcttgggt ttctgcagcg cgggtgccca ctgcctgaat tttcccatta tgtggagact 1020

cgccttgccc tagctgctga cccaacattc tgtgcctccc tgaaagctga gttggatctg 1080

gaggagcgcg tgatggggcc acgctgttca caatgtgact acatcatgct acagaacctg 1140

tcactctggc tgatgcagaa cctatcagct gggcagttgc accaccaa atttgcaacc 1200

tatgcagctg tgtacagtgt ggctcaggcc cttcacaaca ccctgcagtg caatgtctca 1260

cattgccaca catcagagcc tgttcaacc tggcagctcc tggagaacat gtacaatatg 1320

agtttccgtg ctcgagactt gacactgcag tttgatgcca aaggagtgat agacatggaa 1380

tatgacctga agatgtgggt gtggcagagc cctacacctg tactacatac tgtaggcacc 1440

ttcaacggca ccctcagct gcagcactcg aaaatgtatt ggccaggcaa ccagggtgcca 1500

gtctcccagt gctcccggca gtgcaaagat ggccagggtg gcagagtaaa gggctttcat 1560

tcctgctgct atgactgtgt ggactgcaag gcagggagct accggaagca tccagatgac 1620

ttcacctgta ctccatgtg caaggatcag tggctcccag aaaaaagcac aacctgctta 1680

cctcgcaggc ccaagtttct ggcttggggg gagccagctg tgctgtcact tctcctgctg 1740

ctttgcctgg tgctgggctt gacactggct gcctggggc tctttgtcca ctactgggac 1800

agccctcttg ttcaggcctc aggtgggtca ctgttctgct ttggcctgat ctgcctaggc 1860

-continued

```

ctcttctgcc tcagtgtcct tctgttccca ggacgaccac gctctgccag ctgccttgcc 1920
caacaaccaa tggctcacct cctctcaca ggctgcctga gcacactctt cctgcaagca 1980
gccgagatct ttgtggagtc tgagctgccca ctgagttggg caaactggct ctgcagctac 2040
cttcggggcc cctgggcttg gctgggtgga ctgctggcca ctcttggtga ggctgcacta 2100
tgtgcctggt acttgatggc tttccctcca gaggtgggta cagattggca ggtgctgccc 2160
acggaggtag tggaacactg ccgcatgcgt tcctgggtca gcctgggctt ggtgcacatc 2220
accaatgcag tgtagcttt cctctgcttt ctgggcactt tcctggtaca gagccagcct 2280
ggtcgctata accgtgcccg tggcctcacc ttcgccatgc tagcttattt catcatctgg 2340
gtctcttttg tgcccctcct ggctaagtgc caggtggcct accagccagc tgtgcagatg 2400
ggtgctatct tattctgtgc cctgggcatc ctggccacct tccacctgcc caaatgctat 2460
gtacttctgt ggctgccaga gctcaacacc caggagtctt tcctgggaag gagccccaag 2520
gaagcatcag atgggaatag tggtagtagt gaggcaactc ggggacacag tgaatga 2577

```

<210> SEQ ID NO 14

<211> LENGTH: 858

<212> TYPE: PRT

<213> ORGANISM: Rattus sp.

<400> SEQUENCE: 14

```

Met Pro Gly Leu Ala Ile Leu Gly Leu Ser Leu Ala Ala Phe Leu Glu
 1           5           10          15
Leu Gly Met Gly Ser Ser Leu Cys Leu Ser Gln Gln Phe Lys Ala Gln
          20          25          30
Gly Asp Tyr Ile Leu Gly Gly Leu Phe Pro Leu Gly Thr Thr Glu Glu
          35          40          45
Ala Thr Leu Asn Gln Arg Thr Gln Pro Asn Gly Ile Leu Cys Thr Arg
          50          55          60
Phe Ser Pro Leu Gly Leu Phe Leu Ala Met Ala Met Lys Met Ala Val
          65          70          75          80
Glu Glu Ile Asn Asn Gly Ser Ala Leu Leu Pro Gly Leu Arg Leu Gly
          85          90          95
Tyr Asp Leu Phe Asp Thr Cys Ser Glu Pro Val Val Thr Met Lys Pro
          100         105         110
Ser Leu Met Phe Met Ala Lys Val Gly Ser Gln Ser Ile Ala Ala Tyr
          115         120         125
Cys Asn Tyr Thr Gln Tyr Gln Pro Arg Val Leu Ala Val Ile Gly Pro
          130         135         140
His Ser Ser Glu Leu Ala Leu Ile Thr Gly Lys Phe Phe Ser Phe Phe
          145         150         155         160
Leu Met Pro Gln Val Ser Tyr Ser Ala Ser Met Asp Arg Leu Ser Asp
          165         170         175
Arg Glu Thr Phe Pro Ser Phe Phe Arg Thr Val Pro Ser Asp Arg Val
          180         185         190
Gln Leu Gln Ala Val Val Thr Leu Leu Gln Asn Phe Ser Trp Asn Trp
          195         200         205
Val Ala Ala Leu Gly Ser Asp Asp Asp Tyr Gly Arg Glu Gly Leu Ser
          210         215         220
Ile Phe Ser Gly Leu Ala Asn Ser Arg Gly Ile Cys Ile Ala His Glu
          225         230         235         240
Gly Leu Val Pro Gln His Asp Thr Ser Gly Gln Gln Leu Gly Lys Val

```


-continued

245					250					255					
Val	Asp	Val	Leu	Arg	Gln	Val	Asn	Gln	Ser	Lys	Val	Gln	Val	Val	Val
			260					265					270		
Leu	Phe	Ala	Ser	Ala	Arg	Ala	Val	Tyr	Ser	Leu	Phe	Ser	Tyr	Ser	Ile
		275					280					285			
Leu	His	Asp	Leu	Ser	Pro	Lys	Val	Trp	Val	Ala	Ser	Glu	Ser	Trp	Leu
	290					295					300				
Thr	Ser	Asp	Leu	Val	Met	Thr	Leu	Pro	Asn	Ile	Ala	Arg	Val	Gly	Thr
305					310					315					320
Val	Leu	Gly	Phe	Leu	Gln	Arg	Gly	Ala	Leu	Leu	Pro	Glu	Phe	Ser	His
				325					330					335	
Tyr	Val	Glu	Thr	Arg	Leu	Ala	Leu	Ala	Ala	Asp	Pro	Thr	Phe	Cys	Ala
			340					345						350	
Ser	Leu	Lys	Ala	Glu	Leu	Asp	Leu	Glu	Glu	Arg	Val	Met	Gly	Pro	Arg
		355					360						365		
Cys	Ser	Gln	Cys	Asp	Tyr	Ile	Met	Leu	Gln	Asn	Leu	Ser	Ser	Gly	Leu
	370					375					380				
Met	Gln	Asn	Leu	Ser	Ala	Gly	Gln	Leu	His	His	Gln	Ile	Phe	Ala	Thr
385					390					395					400
Tyr	Ala	Ala	Val	Tyr	Ser	Val	Ala	Gln	Ala	Leu	His	Asn	Thr	Leu	Gln
				405					410					415	
Cys	Asn	Val	Ser	His	Cys	His	Thr	Ser	Glu	Pro	Val	Gln	Pro	Trp	Gln
		420						425					430		
Leu	Leu	Glu	Asn	Met	Tyr	Asn	Met	Ser	Phe	Arg	Ala	Arg	Asp	Leu	Thr
		435					440						445		
Leu	Gln	Phe	Asp	Ala	Lys	Gly	Ser	Val	Asp	Met	Glu	Tyr	Asp	Leu	Lys
	450					455					460				
Met	Trp	Val	Trp	Gln	Ser	Pro	Thr	Pro	Val	Leu	His	Thr	Val	Gly	Thr
465					470					475					480
Phe	Asn	Gly	Thr	Leu	Gln	Leu	Gln	His	Ser	Lys	Met	Tyr	Trp	Pro	Gly
			485						490					495	
Asn	Gln	Val	Pro	Val	Ser	Gln	Cys	Ser	Arg	Gln	Cys	Lys	Asp	Gly	Gln
		500						505					510		
Val	Arg	Arg	Val	Lys	Gly	Phe	His	Ser	Cys	Cys	Tyr	Asp	Cys	Val	Asp
		515					520					525			
Cys	Lys	Ala	Gly	Ser	Tyr	Arg	Lys	His	Pro	Asp	Asp	Phe	Thr	Cys	Thr
	530					535					540				
Pro	Cys	Gly	Lys	Asp	Gln	Trp	Ser	Pro	Glu	Lys	Ser	Thr	Thr	Cys	Leu
545					550					555					560
Pro	Arg	Arg	Pro	Lys	Phe	Leu	Ala	Trp	Gly	Glu	Pro	Ala	Val	Leu	Ser
			565						570					575	
Leu	Leu	Leu	Leu	Leu	Cys	Leu	Val	Leu	Gly	Leu	Thr	Leu	Ala	Ala	Leu
			580					585					590		
Gly	Leu	Phe	Val	His	Tyr	Trp	Asp	Ser	Pro	Leu	Val	Gln	Ala	Ser	Gly
		595					600					605			
Gly	Ser	Leu	Phe	Cys	Phe	Gly	Leu	Ile	Cys	Leu	Gly	Leu	Phe	Cys	Leu
	610					615					620				
Ser	Val	Leu	Leu	Phe	Pro	Gly	Arg	Pro	Arg	Ser	Ala	Ser	Cys	Leu	Ala
625					630					635					640
Gln	Gln	Pro	Met	Ala	His	Leu	Pro	Leu	Thr	Gly	Cys	Leu	Ser	Thr	Leu
			645						650					655	
Phe	Leu	Gln	Ala	Ala	Glu	Ile	Phe	Val	Glu	Ser	Glu	Leu	Pro	Leu	Ser
			660					665					670		

-continued

Trp Ala Asn Trp Leu Cys Ser Tyr Leu Arg Gly Pro Trp Ala Trp Leu
 675 680 685
 Val Val Leu Leu Ala Thr Leu Val Glu Ala Ala Leu Cys Ala Trp Tyr
 690 695 700
 Leu Met Ala Phe Pro Pro Glu Val Val Thr Asp Trp Gln Val Leu Pro
 705 710 715 720
 Thr Glu Val Leu Glu His Cys Arg Met Arg Ser Trp Val Ser Leu Gly
 725 730 735
 Leu Val His Ile Thr Asn Ala Val Leu Ala Phe Leu Cys Phe Leu Gly
 740 745 750
 Thr Phe Leu Val Gln Ser Gln Pro Gly Arg Tyr Asn Arg Ala Arg Gly
 755 760 765
 Leu Thr Phe Ala Met Leu Ala Tyr Phe Ile Ile Trp Val Ser Phe Val
 770 775 780
 Pro Leu Leu Ala Asn Val Gln Val Ala Tyr Gln Pro Ala Val Gln Met
 785 790 795 800
 Gly Ala Ile Leu Phe Cys Ala Leu Gly Ile Leu Ala Thr Phe His Leu
 805 810 815
 Pro Lys Cys Tyr Val Leu Leu Trp Leu Pro Glu Leu Asn Thr Gln Glu
 820 825 830
 Phe Phe Leu Gly Arg Ser Pro Lys Glu Ala Ser Asp Gly Asn Ser Gly
 835 840 845
 Ser Ser Glu Ala Thr Arg Gly His Ser Glu
 850 855

<210> SEQ ID NO 15
 <211> LENGTH: 8194
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1251)..(1300)
 <223> OTHER INFORMATION: a, t, c, g, other or unknown
 <220> FEATURE:
 <221> NAME/KEY: modified_base
 <222> LOCATION: (1951)..(2000)
 <223> OTHER INFORMATION: a, t, c, g, other or unknown

<400> SEQUENCE: 15

gagaatctcg cgagatcccg tcggtccgcc ccgctgccct cccagctgcc gaaaagaggg 60
 gcctccgagc cgccggcgcc ctctgccggc aaacctccgga agcacactag gaggttccag 120
 ccgatctggt cgaggggctc cacggaggac tccatctacg ttacgcaaat tccctacccc 180
 agccggccgg agagagaaaag ccagaaacct cgcgaccagc catggggccac ctctccggaa 240
 aaacaccggg atattttttt tctcctgcag aaaaagcttt aggattggca gtttaaaaaa 300
 aacatgtcta ttgcatacc ttcggtttgc atgcatttgt ttcgaagtga gcaaccctgg 360
 gtaacaaggc gaaagtatat gacaatttgc tcagaatctt aatgtcagaa aactggagac 420
 tggggcaggg ggggtgctgac tcaaagctgt gtctcattta gtaaactgag gccaggtaa 480
 aaagtctga aacctcgcaa caccgggaga aattgtgttc cagcctcca cctcgcccca 540
 aatgccaga gctccttttc taagccaggt gaagtcacag agcgtggaca gaaccacaaa 600
 ccgtccagag gaagggtcac tgggtgccac ctggtttgca tctgtgcctt cgtcctgccc 660
 agttcctgag tgggaccgca ggcccgaat gtcaaggcaa acagtcctgc ttcagccact 720
 gggctccagt cccaccctt ttgggggcct gaagtttaga agcatccggc agctgccttc 780

-continued

tatttaagca	actggcctcc	ttagaggcca	ctccttggcc	atgccaggcg	cgggcatctg	840
gccagcatgc	tgctctgcac	ggctcgcctg	gtcggcctgc	agcttctcat	ttcctgctgc	900
tgggcctttg	cctgccatag	cacggagtct	tctcctgact	tcaccctccc	cggagattac	960
ctcctggcag	gcctgttccc	tctccattct	ggctgtctgc	aggtgaggca	cagacccgag	1020
gtgaccctgt	gtgacaggtg	agtgaggggc	cagcagagcc	acacttagtg	ggaccctgg	1080
ctatagggcc	cctctggctg	ccatcctcca	aacaggacct	tgctctgcc	tttgcctctt	1140
gaactgtccc	caggccttgt	tcatcaatcc	acttgccacc	taagtgctgg	ctagaccttc	1200
ctagacactt	cggccagttt	ccaattatth	cacccttgct	gttagaatgt	nnnnnnnnnn	1260
nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	aattccttaa	actaaatttc	1320
tcactttctc	tctctctctg	gaaaacactg	actaatgtag	caggtttctc	tgctccagga	1380
cttcaggacc	ttttcgatgc	taataagttt	ctccatcagg	gccagcttgt	tcctcctact	1440
gagcttgaga	gcccttgttg	aagttgtgg	ttgggggact	ggaccgatga	cctcaaaggt	1500
tccctttgct	cccaagcctc	agagtctagg	aggccagagg	gtctcagcag	gcctttgtcc	1560
ttctcagctg	tctcttactg	gctttctcca	caggcttctg	agcttcaatg	agcatggcta	1620
ccacctcttc	caggctatgc	ggcttgggg	tgaggagata	aacaactcca	cggcctgct	1680
gccaacatc	accctgggg	accagctgta	tgatgtgtgt	tctgactctg	ccaatgtgta	1740
tgccacgctg	agagtgtct	ccctgccagg	gcaacaccac	atagagctcc	aaggagacct	1800
tctccactat	tcccctacgg	tgctggcagt	gattgggcct	gacagcacca	accgtgctgc	1860
caccacagcc	gccctgctga	gccctttcct	ggtgccatg	gtaagctgga	gcctcagacc	1920
tttgcctatc	tcccttcagg	caagtctggg	nnnnnnnnnn	nnnnnnnnnn	nnnnnnnnnn	1980
nnnnnnnnnn	nnnnnnnnnn	gccaccatgc	ccggctaatt	tttttgatt	tttagtagag	2040
acggggtttc	accgtgttag	ccaggctgg	cgaaactcc	taacctctg	atccaccac	2100
ctcggcctcc	caatgtgctg	ggattacagg	tgtgagccac	tgacccggc	cataatgtat	2160
taataataa	aaataattat	acaactcacc	ataatgtaga	atcagtggga	gccctgagct	2220
tgttttccta	caactagatg	gtcccatctg	gggtgatgg	gagacagtga	cagatcatca	2280
gacattagat	tctcataagt	agcgtgcaac	ccagatccct	cgcatgtgca	gttcacagta	2340
gggttcaagc	tcctacaaga	atctgatgct	gctgctgatc	tgacaggagg	ggagcagctg	2400
taaatacaga	tgaagcttcg	cttactcacc	agctgtcac	ctcctcctgt	gaggcccgg	2460
tcctaacagg	ccactgacct	aacttctgcc	ctgacctaca	catgcttctc	ttcttccttg	2520
caaactgcct	ccagtggaag	tccctgaagg	tcccaaaaca	cacgggacta	tttcaactcct	2580
atgcaggttt	tgtctccttt	gcttggaatg	catcccctca	ccccttgctc	ccaggcagat	2640
tcccaccctc	cccccagaac	ctgccccagt	ggagccttcg	cagggtgatt	gtcagtttca	2700
caggctgagg	ggtgctctcc	tggtctcccc	ggctccctgt	atccccacac	ccagcacagg	2760
gccaggcact	gggggggcct	tcagtggaga	ctgaaatggc	tgaacgggac	ctccataga	2820
ttagctatgc	ggccagcagc	gagacgctca	gctggaagcg	gcagtatccc	tctttcctgc	2880
gcaccatccc	caatgacaag	taccaggtgg	agaccatggt	gctgctgctg	cagaagttcg	2940
ggtggacctg	gatctctctg	gttggcagca	gtgacgacta	tgggcagcta	ggggtgcagg	3000
cactggagaa	ccaggccact	ggtcagggga	tctgcattgc	tttcaaggac	atcatgccct	3060
tctctgccc	ggtggcgat	gagaggatgc	agtgcctcat	gcgccacctg	gcccaggccg	3120
gggccaccgt	cgtggttgtt	ttttccagcc	ggcagttggc	cagggtgttt	ttcagatccg	3180

-continued

tggtgctgac	caacctgact	ggcaagggtgt	gggtcgccctc	agaagcctgg	gccctctcca	3240
ggcacatcac	tggggtgccc	gggatccagc	gcattgggat	ggtgctgggc	gtggccatcc	3300
agaagagggc	tgtccctggc	ctgaaggcgt	ttgaagaagc	ctatgcccg	gcagacaaga	3360
aggcccctag	gccttgccac	aagggtcct	ggtgcagcag	caatcagctc	tgagagaaat	3420
gccaagcttt	catggcacac	acgatgccc	agctcaaagc	cttctccatg	agttctgcct	3480
acaacgcata	ccgggctgtg	tatgcgggtg	cccatggcct	ccaccagctc	ctgggctgtg	3540
cctctggagc	ttgttccagg	ggccgagtct	accctggca	ggtaagagag	cccaccccag	3600
cacctcctgt	cagggagaac	agccaatcct	gagatgagca	gagtgggcac	tctccggcca	3660
ctctaaatgc	caaggggat	aatgcccact	aacttgaggt	tttttgttt	gttttgttt	3720
gttttttgag	acagtctggc	tctgtcacc	aggtgcag	gtagtgatgc	gatctcggt	3780
ctctgcaact	tccacctcct	gggttcaagt	gattctcttg	cctcggcctc	ctgagtagct	3840
gggattacag	gcaccacca	ccatgcctgg	ataatttttc	ttttctttt	ttttttttg	3900
agatagagtc	tcgctctgtt	gccagggctg	gaatgcagtg	gtgcgatctt	ggctcactgt	3960
gagctccgcc	tcccaggtc	actccattcc	cctgcctcag	cctcccaagt	aggtgggact	4020
acgggcgccc	gccaccacgc	ccagctaatt	ttttttgtat	tttgagtaga	gacgggggtt	4080
caccatgtta	gccaggatgg	tctcaatctc	ctgacctgtg	catccgcca	cctcgtcctc	4140
ccaaagtgct	gggattacag	gcgtgagcca	ccgcaccgg	cctaattttt	gtatttttag	4200
tagagatggg	gtttcaccat	gttggccagg	ctggtctoga	actcctggca	tcaagtgatc	4260
ctcctgcttc	ggcctcccaa	agtgctggga	ttacaggcat	tagctctctt	ctcttagaca	4320
gatctttctc	tctgatcctt	gccttctctc	accactgtg	tcttggaggt	gtcaagtgat	4380
aagatccagg	gctaaaactg	tctgtaaagg	agtgtttgtt	agaggcctcc	tctcaggagg	4440
ttggtgggga	agattgaggg	gcttcctaag	aaggaaggga	cgagacctc	ctgatgggct	4500
gaaaccacca	ggacggaaac	ccaggaaggc	cccaggccct	tgcttctggg	accatgtggg	4560
tctgtgctgt	ctgtgggtgg	ttcatgatac	gcgtttcttt	cagcttttgg	agcagatcca	4620
caaggtgcat	ttccttctac	acaaggacac	tgtggcggtt	aatgacaaca	gagatcccct	4680
cagtagctat	aacataattg	cctgggactg	gaatggaccc	aagtggacct	tcacggctcct	4740
cggttctctc	acatggtctc	cagttcagct	aaacataaat	gagaccaaaa	tccagtggca	4800
cggaaggac	aaccaggtaa	tggggatgtg	gctactcacc	atgtaactgg	cttatgggca	4860
acctagagcc	tgggggtgat	gctgacacag	tgtacaggga	gcaggagggg	ggccccaggg	4920
gtccagctgc	caccactcta	cccctcctgg	ccagggaagc	agggaagaca	ctccgtaggc	4980
gagtgtgcag	atgccctggg	gcggaagttc	acacgaccag	gggcccctgc	ctgggagtga	5040
gccctgaggg	cagatgcaca	gagattctgt	tttctgttcc	acatgtgagc	tgtcctttga	5100
cttgggcccc	tacgtgtggc	ccctctggct	tcttacaggt	gcctaagtct	gtgtgttcca	5160
gcgactgtct	tgaagggcac	cagcgagtgg	ttacgggttt	ccatcactgc	tgctttgagt	5220
gtgtgccttg	tggggctggg	accttctca	acaagagtgg	tgagtgggca	atggagcagg	5280
cgactaccc	agcactccc	ggggctgcac	ggtggaggga	gggcccctcc	tgggccccat	5340
gtgccctgcc	ccagaaccaa	ggcccagtca	ctgggctgcc	agttagcttc	aggttggagg	5400
acacctgcta	ccagacagaa	ttctgatcaa	gagaatcagc	cactgggtgc	ggtggctcat	5460
gcctgtaatc	ccagcacttt	gggaggctga	ggcgggtgga	tcacttgagg	tcgggagttc	5520

-continued

gagaccagcc	tggccaacat	ggtgaaaccc	catctctacc	aaaaatataa	aaaattagct	5580
gggtgtggtg	gcgcgctgcct	gtaatcccag	ctactcggga	ggctgaggca	ggagaatcac	5640
ttgaacccag	gaggcggagg	ttgcagtgag	ccaagatgca	ttccagcctg	gaccacaaag	5700
cgagaattcg	tcccccaaaa	aaaagaaagg	aggccggggc	cggtggctca	cacctgtaat	5760
cccagcactt	tgggaggccg	agggtgggtg	atcacctgag	gtcaggagtt	cgagaccagc	5820
ctgaccaaca	tgggtgaaacc	ccatctctac	taaaaataca	aaaaaagtta	gccgggctgt	5880
gtggcgtgtg	cctgtaattc	cagctactcg	ggaggctgag	gcaggagaat	tgcttgaacc	5940
cgggaggcgg	aggttgcagt	gagccaagat	tgcaccattg	cactccagcc	tgggcgacaa	6000
gagaaaaact	ctgtctcaaa	aaaaagaaa	gaaagaaaga	attagccaac	tgaaagcctt	6060
agactgaggt	gtgtcctctg	ttagagagct	gtcatcacia	ctcctacaaa	agcagtcgta	6120
tcctgaattc	aacctctttc	tctaaatgaa	tatagctatt	gttccctttg	tgccctcttg	6180
tcctactgtc	ccttctgttg	cccatgccaa	agacagctag	ctccttgaac	agcttggcct	6240
gaatacagat	actagcgtgt	ctgcagcaga	gaaaaaaca	gcattcccca	tccagaaatg	6300
caaggtcaag	aacagagagc	aaattaggta	gctaaggact	caggtcctta	gttgggtgtcc	6360
agggggccaca	ttctttcctt	tcaccatctc	tgtagggaca	ggaatacttc	ccttctgtcc	6420
tcagagggtc	aggactcaga	gaaaccacag	agcagcagct	caggaaagtg	gttcatggaa	6480
atgctggcaa	gagagagggg	ttacaatgcc	ctcccttggg	agcaggctgc	tcccatcaga	6540
tcgtaacctc	tctggtatgt	gggcagagct	accaggttaa	ggcctccctc	agggtttgca	6600
aaaccctcat	gggatcatga	gccatacaga	accgacctgt	gtgtctccag	agtctgtaat	6660
taacacaggg	atthttgagga	aatgcgtggc	ctcaggcccc	actcccggct	acccccatcc	6720
cactatgcct	agtatagtct	agctgcctctg	gtacaattct	cccagtatct	tgacggcccc	6780
tatttctctat	tcctactctg	ctcatctggc	tctcaggaac	cttcttggcc	ttccctttca	6840
gacctctaca	gatgccagcc	ttgtgggaaa	gaagagtggg	cacctgaggg	aagccagacc	6900
tgcttcccgc	gcactgtggt	gtttttggct	ttgcgtgagc	acacctcttg	ggtgctgctg	6960
gcagctaaca	cgctgctgct	gctgctgctg	cttgggactg	ctggcctggt	tgccctggcac	7020
ctagacaccc	ctgtggtgag	gtcagcaggg	ggccgcctgt	gctttcttat	gctgggctcc	7080
ctggcagcag	gtagtggcag	cctctatggc	ttctttgggg	aaccacaag	gcctgcgtgc	7140
ttgctacgcc	aggccctctt	tgcccttggg	ttaccatct	tcctgtcctg	cctgacagtt	7200
cgctcattcc	aactaatcat	catcttcaag	ttttccacca	aggtacctac	attctaccac	7260
gcctgggtcc	aaaaccacgg	tgctggcctg	tttgtgatga	tcagctcagc	ggcccagctg	7320
cttatctgtc	taacttggct	ggtggtgtgg	accccactgc	ctgctaggga	ataccagcgc	7380
ttcccccatc	tggatgatgct	tgagtgcaca	gagaccaact	ccctgggctt	catactggcc	7440
ttcctctaca	atggcctcct	ctccatcagt	gcctttgcct	gcagctacct	gggtaaggac	7500
ttgccagaga	actacaacga	ggccaaatgt	gtcaccttca	gcctgctctt	caacttcgtg	7560
tcctggatcg	ccttcttcac	cacggccagc	gtctacgagc	gcaagtacct	gcctgcggcc	7620
aacatgatgg	ctgggctgag	cagcctgagc	agcggcttcg	gtgggtattt	tctgcctaag	7680
tgtactgtga	tcctctgccg	cccagacctc	aacagcacag	agcacttcca	ggcctccatt	7740
caggactaca	cgaggcgtcg	cggctccacc	tgaccagtgg	gtcagcaggc	acggctggca	7800
gccttctctg	ccctgagggg	cgaaggctca	gcaggccggg	ggtgtccggg	aggctcttgg	7860
gcatcgcggt	ctggggttgg	gacgtgtaag	cgctgggag	agcctagacc	aggctccggg	7920

-continued

```

ctgccaataa agaagtgaaa tgcgtatctg gtctcctgtc gtgggagagt gtgaggtgta 7980
acggattcaa gtctgaacct agagcctgga aaaggctgac cgcccagatt gacgttgcta 8040
ggcaactccg gaggcgggcc cagcgccaaa agaacagggc gaggcgtcgt ccccgcatcc 8100
cattggccgt tctctgcggg gccccgccct cgggggccgg agctagaagc tctacgcttc 8160
cgaggcgcac ctctggcct gcacgctttg acgt 8194

```

<210> SEQ ID NO 16

<211> LENGTH: 2526

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

```

atgctgctct gcacggctcg cctggctggc ctgcagcttc tcatttcctg ctgctgggcc 60
tttgctgcc atagcacgga gtcttctcct gacttcacct tccccggaga ttacctcctg 120
gcaggcctgt tccctctcca ttctggctgt ctgcaggtga ggcacagacc cgaggtgacc 180
ctgtgtgaca ggtctttag cttcaatgag catggctacc acctcttcca ggctatgcgg 240
cttgggggtg aggagataaa caactccacg gccctgctgc ccaacatcac cctgggggtac 300
cagctgtatg atgtgtgttc tgactctgcc aatgtgtatg ccacgctgag agtgctctcc 360
ctgccagggc aacaccacat agagctccaa ggagaccttc tccactattc ccctacggtg 420
ctggcagtga ttggcctga cagcaccaac cgtgctgcca ccacagccgc cctgctgagc 480
cctttcctgg tgcccatgat tagctatgag gccagcagcg agacgctcag cgtgaagcgg 540
cagtatccct ctttctgctg caccatcccc aatgacaagt accaggtgga gaccatggtg 600
ctgctgctgc agaagttcgg gtggacctgg atctctctgg ttggcagcag tgacgactat 660
gggcagctag ggggtgcaggc actggagaac caggccactg gtcaggggat ctgcattgct 720
ttcaaggaca tcatgccctt ctctgcccag gtgggcgatg agaggatgca gtgcctcatg 780
cgccacctgg cccaggccgg ggccaccgtc gtggtgtgtt tttccagccg gcagttggcc 840
agggtgtttt tcgagtcctg ggtgctgacc aacctgactg gcaaggtgtg ggtgcctca 900
gaagcctggg ccctctccag gcacatcact ggggtgcccc ggatccagcg cattgggatg 960
gtgctgggcg tggccatcca gaagagggtc gtccctggcc tgaaggcgtt tgaagaagcc 1020
tatgcccggg cagacaagaa ggcccctagg ccttgccaca agggctcctg gtgcagcagc 1080
aatcagctct gcagagaatg ccaagctttc atggcacaca cgatgcccga gctcaaagcc 1140
ttctccatga gttctgccta caacgcatac cgggctgtgt atgcggtggc ccatggcctc 1200
caccagctcc tgggctgtgc ctctggagct tgttccaggg gccgagtcta cccctggcag 1260
cttttgagc agatccacaa ggtgcatttc cttctacaca aggacactgt ggcgtttaat 1320
gacaacagag atcccctcag tagctataac ataattgctt gggactggaa tggacccaag 1380
tggaccttca cggctctcgg ttcctccaca tggctctcag ttcagctaaa cataaatgag 1440
acaaaaatcc agtggcacgg aaaggacaac caggtgccta agtctgtgtg ttccagcgac 1500
tgtcttgaag ggcaccagcg agtggttacg gtttccatc actgctgctt tgagtgtgtg 1560
ccctgtgggg ctgggacctt cctcaacaag agtgacctct acagatgcca gccttgtggg 1620
aaagaagagt gggcacctga ggaagccag acctgcttcc cgcgactgt ggtgtttttg 1680
gctttgcgtg agcacacctc ttgggtgctg ctggcagcta acacgctgct gctgctgctg 1740
ctgcttggga ctgctggcct gtttgcctgg cacctagaca cccctgtggt gaggtcagca 1800

```

-continued

```

gggggcccgc tgtgctttct tatgctgggc tccctggcag caggtagtgg cagcctctat 1860
ggcttctttg ggggaaccac aaggcctgcg tgcttgctac gccaggccct ctttgccctt 1920
ggtttcacca tcttctgtc ctgcctgaca gttcgtcat tccaactaat catcatcttc 1980
aagttttcca ccaaggtacc tacattctac cacgcctggg tccaaaacca cggtgctggc 2040
ctgtttgtga tgatcagctc agcggcccag ctgcttatct gtctaacttg gctggtggtg 2100
tggacccccc tgctgctag ggaataccag cgcttcccc atctggtgat gcttgagtgc 2160
acagagacca actccctggg cttcatactg gccttctct acaatggcct cctctccatc 2220
agtgcctttg cctgcagcta cctgggtaag gacttgccag agaactaaa cgaggccaaa 2280
tgtgtcacct tcagcctgct cttcaacttc gtgtcctgga tcgccttctt caccacggcc 2340
agcgtctacg acggcaagta cctgcctgcg gccaacatga tggctgggct gagcagcctg 2400
agcagcggct tcggtgggta ttttctgcct aagtgtacg tgatcctctg ccgccagac 2460
ctcaacagca cagagcactt ccaggcctcc attcaggact acacgaggcg ctgcggtcc 2520
acctga 2526

```

<210> SEQ ID NO 17

<211> LENGTH: 841

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

```

Met Leu Leu Cys Thr Ala Arg Leu Val Gly Leu Gln Leu Leu Ile Ser
 1           5           10          15
Cys Cys Trp Ala Phe Ala Cys His Ser Thr Glu Ser Ser Pro Asp Phe
          20          25          30
Thr Leu Pro Gly Asp Tyr Leu Leu Ala Gly Leu Phe Pro Leu His Ser
          35          40          45
Gly Cys Leu Gln Val Arg His Arg Pro Glu Val Thr Leu Cys Asp Arg
          50          55          60
Ser Cys Ser Phe Asn Glu His Gly Tyr His Leu Phe Gln Ala Met Arg
          65          70          75          80
Leu Gly Val Glu Glu Ile Asn Asn Ser Thr Ala Leu Leu Pro Asn Ile
          85          90          95
Thr Leu Gly Tyr Gln Leu Tyr Asp Val Cys Ser Asp Ser Ala Asn Val
          100         105         110
Tyr Ala Thr Leu Arg Val Leu Ser Leu Pro Gly Gln His His Ile Glu
          115         120         125
Leu Gln Gly Asp Leu Leu His Tyr Ser Pro Thr Val Leu Ala Val Ile
          130         135         140
Gly Pro Asp Ser Thr Asn Arg Ala Ala Thr Thr Ala Ala Leu Leu Ser
          145         150         155         160
Pro Phe Leu Val Pro Met Ile Ser Tyr Ala Ala Ser Ser Glu Thr Leu
          165         170         175
Ser Val Lys Arg Gln Tyr Pro Ser Phe Leu Arg Thr Ile Pro Asn Asp
          180         185         190
Lys Tyr Gln Val Glu Thr Met Val Leu Leu Leu Gln Lys Phe Gly Trp
          195         200         205
Thr Trp Ile Ser Leu Val Gly Ser Ser Asp Asp Tyr Gly Gln Leu Gly
          210         215         220
Val Gln Ala Leu Glu Asn Gln Ala Thr Gly Gln Gly Ile Cys Ile Ala
          225         230         235         240

```

-continued

Phe	Lys	Asp	Ile	Met	Pro	Phe	Ser	Ala	Gln	Val	Gly	Asp	Glu	Arg	Met	245	250	255	
Gln	Cys	Leu	Met	Arg	His	Leu	Ala	Gln	Ala	Gly	Ala	Thr	Val	Val	Val	260	265	270	
Val	Phe	Ser	Ser	Arg	Gln	Leu	Ala	Arg	Val	Phe	Phe	Glu	Ser	Val	Val	275	280	285	
Leu	Thr	Asn	Leu	Thr	Gly	Lys	Val	Trp	Val	Ala	Ser	Glu	Ala	Trp	Ala	290	295	300	
Leu	Ser	Arg	His	Ile	Thr	Gly	Val	Pro	Gly	Ile	Gln	Arg	Ile	Gly	Met	305	310	315	320
Val	Leu	Gly	Val	Ala	Ile	Gln	Lys	Arg	Ala	Val	Pro	Gly	Leu	Lys	Ala	325	330	335	
Phe	Glu	Glu	Ala	Tyr	Ala	Arg	Ala	Asp	Lys	Lys	Ala	Pro	Arg	Pro	Cys	340	345	350	
His	Lys	Gly	Ser	Trp	Cys	Ser	Ser	Asn	Gln	Leu	Cys	Arg	Glu	Cys	Gln	355	360	365	
Ala	Phe	Met	Ala	His	Thr	Met	Pro	Lys	Leu	Lys	Ala	Phe	Ser	Met	Ser	370	375	380	
Ser	Ala	Tyr	Asn	Ala	Tyr	Arg	Ala	Val	Tyr	Ala	Val	Ala	His	Gly	Leu	385	390	395	400
His	Gln	Leu	Leu	Gly	Cys	Ala	Ser	Gly	Ala	Cys	Ser	Arg	Gly	Arg	Val	405	410	415	
Tyr	Pro	Trp	Gln	Leu	Leu	Glu	Gln	Ile	His	Lys	Val	His	Phe	Leu	Leu	420	425	430	
His	Lys	Asp	Thr	Val	Ala	Phe	Asn	Asp	Asn	Arg	Asp	Pro	Leu	Ser	Ser	435	440	445	
Tyr	Asn	Ile	Ile	Ala	Trp	Asp	Trp	Asn	Gly	Pro	Lys	Trp	Thr	Phe	Thr	450	455	460	
Val	Leu	Gly	Ser	Ser	Thr	Trp	Ser	Pro	Val	Gln	Leu	Asn	Ile	Asn	Glu	465	470	475	480
Thr	Lys	Ile	Gln	Trp	His	Gly	Lys	Asp	Asn	Gln	Val	Pro	Lys	Ser	Val	485	490	495	
Cys	Ser	Ser	Asp	Cys	Leu	Glu	Gly	His	Gln	Arg	Val	Val	Thr	Gly	Phe	500	505	510	
His	His	Cys	Cys	Phe	Glu	Cys	Val	Pro	Cys	Gly	Ala	Gly	Thr	Phe	Leu	515	520	525	
Asn	Lys	Ser	Asp	Leu	Tyr	Arg	Cys	Gln	Pro	Cys	Gly	Lys	Glu	Glu	Trp	530	535	540	
Ala	Pro	Glu	Gly	Ser	Gln	Thr	Cys	Phe	Pro	Arg	Thr	Val	Val	Phe	Leu	545	550	555	560
Ala	Leu	Arg	Glu	His	Thr	Ser	Trp	Val	Leu	Leu	Ala	Ala	Asn	Thr	Leu	565	570	575	
Leu	Leu	Leu	Leu	Leu	Leu	Gly	Thr	Ala	Gly	Leu	Phe	Ala	Trp	His	Leu	580	585	590	
Asp	Thr	Pro	Val	Val	Arg	Ser	Ala	Gly	Gly	Arg	Leu	Cys	Phe	Leu	Met	595	600	605	
Leu	Gly	Ser	Leu	Ala	Ala	Gly	Ser	Gly	Ser	Leu	Tyr	Gly	Phe	Phe	Gly	610	615	620	
Glu	Pro	Thr	Arg	Pro	Ala	Cys	Leu	Leu	Arg	Gln	Ala	Leu	Phe	Ala	Leu	625	630	635	640
Gly	Phe	Thr	Ile	Phe	Leu	Ser	Cys	Leu	Thr	Val	Arg	Ser	Phe	Gln	Leu	645	650	655	
Ile	Ile	Ile	Phe	Lys	Phe	Ser	Thr	Lys	Val	Pro	Thr	Phe	Tyr	His	Ala				

-continued

660				665				670							
Trp	Val	Gln	Asn	His	Gly	Ala	Gly	Leu	Phe	Val	Met	Ile	Ser	Ser	Ala
		675					680					685			
Ala	Gln	Leu	Leu	Ile	Cys	Leu	Thr	Trp	Leu	Val	Val	Trp	Thr	Pro	Leu
	690					695					700				
Pro	Ala	Arg	Glu	Tyr	Gln	Arg	Phe	Pro	His	Leu	Val	Met	Leu	Glu	Cys
705					710					715					720
Thr	Glu	Thr	Asn	Ser	Leu	Gly	Phe	Ile	Leu	Ala	Phe	Leu	Tyr	Asn	Gly
					725				730					735	
Leu	Leu	Ser	Ile	Ser	Ala	Phe	Ala	Cys	Ser	Tyr	Leu	Gly	Lys	Asp	Leu
			740					745					750		
Pro	Glu	Asn	Tyr	Asn	Glu	Ala	Lys	Cys	Val	Thr	Phe	Ser	Leu	Leu	Phe
		755					760						765		
Asn	Phe	Val	Ser	Trp	Ile	Ala	Phe	Phe	Thr	Thr	Ala	Ser	Val	Tyr	Asp
	770					775					780				
Gly	Lys	Tyr	Leu	Pro	Ala	Ala	Asn	Met	Met	Ala	Gly	Leu	Ser	Ser	Leu
785					790					795					800
Ser	Ser	Gly	Phe	Gly	Gly	Tyr	Phe	Leu	Pro	Lys	Cys	Tyr	Val	Ile	Leu
				805					810					815	
Cys	Arg	Pro	Asp	Leu	Asn	Ser	Thr	Glu	His	Phe	Gln	Ala	Ser	Ile	Gln
			820					825					830		
Asp	Tyr	Thr	Arg	Arg	Cys	Gly	Ser	Thr							
		835					840								

<210> SEQ ID NO 18
 <211> LENGTH: 14
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Consensus
 sequence
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (1)
 <223> OTHER INFORMATION: Thr or Arg
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (3)
 <223> OTHER INFORMATION: Phe or Leu
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (4)
 <223> OTHER INFORMATION: Arg, Gln or Pro
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (6)
 <223> OTHER INFORMATION: Arg or Thr
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (7)
 <223> OTHER INFORMATION: Ser, Pro or Val
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (8)
 <223> OTHER INFORMATION: Val, Glu, Arg, Lys or Thr
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (11)
 <223> OTHER INFORMATION: Ala or Glu
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (12)
 <223> OTHER INFORMATION: Trp or Leu
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (13)
 <223> OTHER INFORMATION: Arg, His or Gly

-continued

<400> SEQUENCE: 18

Xaa Cys Xaa Xaa Arg Xaa Xaa Xaa Phe Leu Xaa Xaa Xaa Glu
 1 5 10

<210> SEQ ID NO 19

<211> LENGTH: 15

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Consensus
 sequence

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (1)

<223> OTHER INFORMATION: Leu or Gln

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (3)

<223> OTHER INFORMATION: Glu, Gly or Thr

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (4)

<223> OTHER INFORMATION: Asn, Arg or Cys

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (7)

<223> OTHER INFORMATION: Arg or Glu

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (9)

<223> OTHER INFORMATION: Arg or Lys

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (10)

<223> OTHER INFORMATION: Cys, Gly or Phe

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (11)

<223> OTHER INFORMATION: Val, Leu or Ile

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (13)

<223> OTHER INFORMATION: Phe or Leu

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (14)

<223> OTHER INFORMATION: Ala or Ser

<220> FEATURE:

<221> NAME/KEY: MOD_RES

<222> LOCATION: (15)

<223> OTHER INFORMATION: Met or Leu

<400> SEQUENCE: 19

Xaa Pro Xaa Xaa Tyr Asn Xaa Ala Xaa Xaa Xaa Thr Xaa Xaa Xaa
 1 5 10 15

<210> SEQ ID NO 20

<211> LENGTH: 3563

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

agcctggcag tggcctcagg cagagtctga cgcgcacaaa ctttcaggcc caggaagcga 60
 ggacaccact ggggccccag ggtgtggcaa gtgaggatgg caagggtttt gctaaacaaa 120
 tcctctgccc gctccccgcc ccgggctcac tccatgtgag gccccagtcg gggcagccac 180
 ctgccgtgcc tgttggaagt tgcctctgcc atgctgggcc ctgctgtcct gggcctcagc 240
 ctctgggctc tcctgcacc tgggacgggg gccccattgt gcctgtcaca gcaacttagg 300
 atgaaggggg actacgtgct gggggggctg ttccccctgg gcgaggccga ggaggctggc 360

-continued

ctccgcagcc	ggacacggcc	cagcagccct	gtgtgcacca	ggtacagagg	tgggacggcc	420
tgggtcgggg	tcagggtgac	caggtctggg	gtgctcctga	gctggggccg	aggtggccat	480
ctgcggttct	gtgtggcccc	aggttctcct	caaacggcct	gctctgggca	ctggccatga	540
aaatggccgt	ggaggagatc	aacaacaagt	cggatctgct	gcccgggctg	cgccctgggct	600
acgacctctt	tgatacgtgc	toggagcctg	tggtagccat	gaagcccagc	ctcatgttcc	660
tggccaaggc	aggcagccgc	gacatcgccg	cctactgcaa	ctacacgcag	taccagcccc	720
gtgtgctggc	tgtcatcggg	cccactcgt	cagagctcgc	catggtcacc	ggcaagttct	780
tcagcttctt	cctcatgccc	cagtggggcg	ccccccacca	tcaccacccc	ccaaccaacc	840
cctgccccgt	gggagcccct	tgtgtcagga	gaatgctaca	tgcaccccac	ccagccctgc	900
cctgggagcc	ctgtgtcaga	agatgctctt	ggcctgcag	gtcagctacg	gtgctagcat	960
ggagctgctg	agcggccggg	agaccttccc	ctccttcttc	cgccaccgtg	ccagcgaccg	1020
tgtgcagctg	acggccggcg	cggagctgct	gcaggagttc	ggctggaact	gggtggccgc	1080
cctgggcagc	gacgacgagt	acggccggca	ggcctgagc	atcttctcgg	ccctggccgc	1140
ggcacgcggc	atctgcatcg	cgccagggg	cctggtgccg	ctgccccgtg	ccgatgactc	1200
gcggctgggg	aaggtgcagg	acgtcctgca	ccaggtgaac	cagagcagcg	tgcaggtggt	1260
gctgctgttc	gcctccgtgc	acgccgccc	cgccctcttc	aactacagca	tcagcagcag	1320
gctctcgccc	aaggtgtggg	tggccagcga	ggcctggctg	acctctgacc	tggatcatggg	1380
gctgcccggc	atggcccaga	tgggcacggg	gcttggcttc	ctccagaggg	gtgcccagct	1440
gcacgagttc	ccccagtacg	tgaagacgca	cctggcccctg	gccaccgacc	cgcccttctg	1500
ctctgccctg	ggcgagaggg	agcagggctc	ggaggaggac	gtggtgggcc	agcgtgccc	1560
gcagtgtgac	tgcatacgc	tgcagaacgt	gagcgcaggg	ctaaatcacc	accagacggt	1620
ctctgtctac	gcagctgtgt	atagcgtggc	ccaggcccctg	cacaacactc	ttcagtgcaa	1680
cgccctcaggc	tgccccgcgc	aggaccccgt	gaagcccctg	caggtgagcc	cgggagatgg	1740
gggtgtgctg	tcctctgcat	gtgcccaggc	caccaggcac	ggccaccacg	cctgagctgg	1800
aggtggctgg	cggtcagcc	ccgtcccccg	cccgcagctc	ctggagaaca	tgtacaacct	1860
gaccttccac	gtgggcgggc	tgccgctgcg	gttcgacagc	agcggaaacg	tggacatgga	1920
gtacgacctg	aagctgtggg	tgtggcaggg	ctcagtgccc	aggctccacg	acgtgggag	1980
gttcaacggc	agcctcagga	cagagcgcct	gaagatccgc	tggcacacgt	ctgacaacca	2040
ggtgaggtga	gggtgggtgt	gccaggcgtg	cccgtggtag	cccccgcggc	agggcgagc	2100
ctgggggtgg	gggccgttcc	agtctcccgt	ggcatgccc	agccgagcag	agccagaccc	2160
caggcctgtg	cgcagaagcc	cgtgtcccgg	tgctcgcggc	agtgccagga	gggcccaggtg	2220
cgccgggtca	aggggttcca	ctcctgctgc	tacgactgtg	tggactgcga	ggcgggcagc	2280
taccggcaaa	accaggtga	gccgccttcc	cggcagggcg	gggtgggaac	gcagcagggg	2340
aggtcctgc	caagtcctga	ctctgagacc	agagcccaca	gggtacaaga	cgaacaccca	2400
gcgccttct	cctctctcac	agacgacatc	gcctgcacct	tttgtggcca	ggatgagtgg	2460
tccccggagc	gaagcacacg	ctgcttccgc	cgcaggtctc	ggttcctggc	atggggcgag	2520
ccggctgtgc	tgctgctgct	cctgctgctg	agcctggcgc	tgggccttgt	gctggctgct	2580
ttggggctgt	tcgttcacca	togggacagc	ccactggttc	aggcctcggg	ggggcccctg	2640
gcctgctttg	gcctggtgtg	cctgggcctg	gtctgcctca	gcgtcctcct	gttccctggc	2700

-continued

```

cagcccagcc ctgcccgatg cctggcccag cagcccttgt cccacctccc gctcacgggc 2760
tgcttgagca cactcttctt gcaggcggcc gagatcttctg tggagtacaga actgcctctg 2820
agctgggcag accggctgag tggctgcctg cgggggcctt gggcctggct ggtggtgctg 2880
ctggccatgc tgggtggaggt cgcactgtgc acctggtacc tgggtggcctt cccgccggag 2940
gtggtgacgg actggcacat gctgcccacg gaggcgctgg tgcactgccg cacacgctcc 3000
tgggtcagct tcggcctagc gcacgccacc aatgccacgc tggcctttct ctgcttctg 3060
ggcactttcc tgggtcggag ccagccgggc tgctacaacc gtgcccgtgg cctcaccttt 3120
gccatgctgg cctacttcat cacctgggtc tcctttgtgc ccctcctggc caatgtgcag 3180
gtggtcctca ggcccggct gcagatgggc gccctcctgc tctgtgtcct gggcatcctg 3240
gctgccttcc acctgccag gtgttacctg ctcatgcggc agccagggt caacaccccc 3300
gagttcttcc tgggagggg ccctggggat gcccaaggcc agaatgacgg gaacacagga 3360
aatcagggga aacatgagtg acccaaccct gtgatctcag ccccggtgaa cccagactta 3420
gctgcgatcc cccccaagcc agcaatgacc cgtgtctcgc tacagagacc ctcccgtctt 3480
aggttctgac cccaggttgt ctctgacct tgaccccaca gtgagcccta ggctggagc 3540
acgtggacac ccctgtgacc atc 3563

```

<210> SEQ ID NO 21

<211> LENGTH: 839

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

```

Met Gly Pro Arg Ala Lys Thr Ile Cys Ser Leu Phe Phe Leu Leu Trp
 1           5           10           15

Val Leu Ala Glu Pro Ala Glu Asn Ser Asp Phe Tyr Leu Pro Gly Asp
          20           25           30

Tyr Leu Leu Gly Gly Leu Phe Ser Leu His Ala Asn Met Lys Gly Ile
          35           40           45

Val His Leu Asn Phe Leu Gln Val Pro Met Cys Lys Glu Tyr Glu Val
          50           55           60

Lys Val Ile Gly Tyr Asn Leu Met Gln Ala Met Arg Phe Ala Val Glu
          65           70           75           80

Glu Ile Asn Asn Asp Ser Ser Leu Leu Pro Gly Val Leu Leu Gly Tyr
          85           90           95

Glu Ile Val Asp Val Cys Tyr Ile Ser Asn Asn Val Gln Pro Val Leu
          100          105          110

Tyr Phe Leu Ala His Glu Asp Asn Leu Leu Pro Ile Gln Glu Asp Tyr
          115          120          125

Ser Asn Tyr Ile Ser Arg Val Val Ala Val Ile Gly Pro Asp Asn Ser
          130          135          140

Glu Ser Val Met Thr Val Ala Asn Phe Leu Ser Leu Phe Leu Leu Pro
          145          150          155          160

Gln Ile Thr Tyr Ser Ala Ile Ser Asp Glu Leu Arg Asp Lys Val Arg
          165          170          175

Phe Pro Ala Leu Leu Arg Thr Thr Pro Ser Ala Asp His His Val Glu
          180          185          190

Ala Met Val Gln Leu Met Leu His Phe Arg Trp Asn Trp Ile Ile Val
          195          200          205

Leu Val Ser Ser Asp Thr Tyr Gly Arg Asp Asn Gly Gln Leu Leu Gly
          210          215          220

```

-continued

Glu Arg Val Ala Arg Arg Asp Ile Cys Ile Ala Phe Gln Glu Thr Leu
 225 230 235 240
 Pro Thr Leu Gln Pro Asn Gln Asn Met Thr Ser Glu Glu Arg Gln Arg
 245 250 255
 Leu Val Thr Ile Val Asp Lys Leu Gln Gln Ser Thr Ala Arg Val Val
 260 265 270
 Val Val Phe Ser Pro Asp Leu Thr Leu Tyr His Phe Phe Asn Glu Val
 275 280 285
 Leu Arg Gln Asn Phe Thr Gly Ala Val Trp Ile Ala Ser Glu Ser Trp
 290 295 300
 Ala Ile Asp Pro Val Leu His Asn Leu Thr Glu Leu Gly His Leu Gly
 305 310 315 320
 Thr Phe Leu Gly Ile Thr Ile Gln Ser Val Pro Ile Pro Gly Phe Ser
 325 330 335
 Glu Phe Arg Glu Trp Gly Pro Gln Ala Gly Pro Pro Pro Leu Ser Arg
 340 345 350
 Thr Ser Gln Ser Tyr Thr Cys Asn Gln Glu Cys Asp Asn Cys Leu Asn
 355 360 365
 Ala Thr Leu Ser Phe Asn Thr Ile Leu Arg Leu Ser Gly Glu Arg Val
 370 375 380
 Val Tyr Ser Val Tyr Ser Ala Val Tyr Ala Val Ala His Ala Leu His
 385 390 395 400
 Ser Leu Leu Gly Cys Asp Lys Ser Thr Cys Thr Lys Arg Val Val Tyr
 405 410 415
 Pro Trp Gln Leu Leu Glu Glu Ile Trp Lys Val Asn Phe Thr Leu Leu
 420 425 430
 Asp His Gln Ile Phe Phe Asp Pro Gln Gly Asp Val Ala Leu His Leu
 435 440 445
 Glu Ile Val Gln Trp Gln Trp Asp Arg Ser Gln Asn Pro Phe Gln Ser
 450 455 460
 Val Ala Ser Tyr Tyr Pro Leu Gln Arg Gln Leu Lys Asn Ile Gln Asp
 465 470 475 480
 Ile Ser Trp His Thr Val Asn Asn Thr Ile Pro Met Ser Met Cys Ser
 485 490 495
 Lys Arg Cys Gln Ser Gly Gln Lys Lys Lys Pro Val Gly Ile His Val
 500 505 510
 Cys Cys Phe Glu Cys Ile Asp Cys Leu Pro Gly Thr Phe Leu Asn His
 515 520 525
 Thr Glu Asp Glu Tyr Glu Cys Gln Ala Cys Pro Asn Asn Glu Trp Ser
 530 535 540
 Tyr Gln Ser Glu Thr Ser Cys Phe Lys Arg Gln Leu Val Phe Leu Glu
 545 550 555 560
 Trp His Glu Ala Pro Thr Ile Ala Val Ala Leu Leu Ala Ala Leu Gly
 565 570 575
 Phe Leu Ser Thr Leu Ala Ile Leu Val Ile Phe Trp Arg His Phe Gln
 580 585 590
 Thr Pro Ile Val Arg Ser Ala Gly Gly Pro Met Cys Phe Leu Met Leu
 595 600 605
 Thr Leu Leu Leu Val Ala Tyr Met Val Val Pro Val Tyr Val Gly Pro
 610 615 620
 Pro Lys Val Ser Thr Cys Leu Cys Arg Gln Ala Leu Phe Pro Leu Cys
 625 630 635 640

-continued

Phe Thr Ile Cys Ile Ser Cys Ile Ala Val Arg Ser Phe Gln Ile Val
 645 650 655
 Cys Ala Phe Lys Met Ala Ser Arg Phe Pro Arg Ala Tyr Ser Tyr Trp
 660 665 670
 Val Arg Tyr Gln Gly Pro Tyr Val Ser Met Ala Phe Ile Thr Val Leu
 675 680 685
 Lys Met Val Ile Val Val Ile Gly Met Leu Ala Thr Gly Leu Ser Pro
 690 695 700
 Thr Thr Arg Thr Asp Pro Asp Asp Pro Lys Ile Thr Ile Val Ser Cys
 705 710 715 720
 Asn Pro Asn Tyr Arg Asn Ser Leu Leu Phe Asn Thr Ser Leu Asp Leu
 725 730 735
 Leu Leu Ser Val Val Gly Phe Ser Phe Ala Tyr Met Gly Lys Glu Leu
 740 745 750
 Pro Thr Asn Tyr Asn Glu Ala Lys Phe Ile Thr Leu Ser Met Thr Phe
 755 760 765
 Tyr Phe Thr Ser Ser Val Ser Leu Cys Thr Phe Met Ser Ala Tyr Ser
 770 775 780
 Gly Val Leu Val Thr Ile Val Asp Leu Leu Val Thr Val Leu Asn Leu
 785 790 795 800
 Leu Ala Ile Ser Leu Gly Tyr Phe Gly Pro Lys Cys Tyr Met Ile Leu
 805 810 815
 Phe Tyr Pro Glu Arg Asn Thr Pro Ala Tyr Phe Asn Ser Met Ile Gln
 820 825 830
 Gly Tyr Thr Met Arg Arg Asp
 835

<210> SEQ ID NO 22
 <211> LENGTH: 6
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Synthetic
 PDZIP peptide sequence

<400> SEQUENCE: 22

Ser Val Ser Thr Val Val
 1 5

<210> SEQ ID NO 23
 <211> LENGTH: 2520
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 23

atggggccca gggcaaagac catctgctcc ctgttcttcc tcctatgggt cctggctgag 60
 ccggctgaga actcggactt ctacctgcct ggggattacc tcctgggtgg cctcttctcc 120
 ctccatgcca acatgaaggc cattgttcac cttacttcc tgcaggtgcc catgtgcaag 180
 gagtatgaag tgaagtgat aggctacaac ctcatgcagg ccatgcgctt cgcggtggag 240
 gagatcaaca atgacagcag cctgctgcct ggtgtgctgc tgggctatga gatcgtggat 300
 gtgtgctaca tctccaacaa tgtccagccg gtgctctact tcctggcaca cgaggacaac 360
 ctccttccca tccaagagga ctacagtaac tacatttccc gtgtggtggc tgcattggc 420
 cctgacaact ccgagtctgt catgactgtg gccaaactcc tctccctatt tctccttcca 480
 cagatcacct acagcgccat cagcgatgag ctgcgagaca aggtgcgctt cccggctttg 540

-continued

```

ctgcgtacca caccagcgc cgaccaccac gtcgaggcca tgggtgcagct gatgctgcac 600
ttccgctgga actggatcat tgtgctggtg agcagcgaca cctatggccg cgacaatggc 660
cagctgcttg gcgagcgcgt ggcccggcgc gacatctgca tcgccttcca ggagacgctg 720
cccacactgc agcccaacca gaacatgacg tcagaggagc gccagcgcct ggtgaccatt 780
gtggacaagc tgcagcagag cacagcgcgc gtcgtggtcg tgttctcgcc cgacctgacc 840
ctgtaccact tcttcaatga ggtgctgctg cagaacttca cgggcgccgt gtggatcgcc 900
tccgagtccct gggccatcga cccggctctg cacaacctca cggagctggg ccacttgggc 960
accttcctgg gcatcacat ccagagcgtg cccatcccgg gcttcagtga gttccgcgag 1020
tggggcccac aggctgggccc gccaccctc agcaggacca gccagagcta tacctgcaac 1080
caggagtgcg acaactgcct gaacgccacc ttgtccttca acaccattct caggctctct 1140
ggggagcgtg tcgtctacag cgtgtactct gcggtctatg ctgtggccca tgccctgcac 1200
agcctcctcg gctgtgacaa aagcacctgc accaagaggg tggcttacc ctggcagctg 1260
cttgaggaga tctggaaggt caacttact ctctggacc accaaatctt cttcgaccgg 1320
caaggggacg tggctctgca cttggagatt gtccagtggc aatgggaccg gagccagaat 1380
cccttcacaga gcgtcgctc ctactacccc ctgcagcgac agctgaagaa catccaagac 1440
atctcctggc acaccgtcaa caacacgatc cctatgtcca tgtgttcaa gaggtgccag 1500
tcagggcaaa agaagaagcc tgtgggcatc cacgtctgct gcttcgagtg catcgactgc 1560
cttcccggca ccttctcaa ccacactgaa gatgaatatg aatgccaggc ctgcccgaat 1620
aacgagtggc cctaccagag tgagacctcc tgcttcaagc ggcagctggt cttcctggaa 1680
tggcatgagg caccaccat cgctgtggcc ctgctggccg ccctgggctt cctcagcacc 1740
ctggccatcc tggatgatatt ctggaggcac ttccagacac ccatagttcg ctggctggg 1800
ggccccatgt gcttctgat gctgacactg ctgctggtgg catacatggt ggtcccgggtg 1860
tacgtggggc cgcccaaggt ctccacctgc ctctgccgcc aggcctctt tcccctctgc 1920
ttcacaattt gcatctcctg tatcgccgtg cgttctttcc agatcgtctg cgcttcaag 1980
atggccagcc gcttcccacg cgcctacagc tactgggtcc gctaccaggg gccctacgtc 2040
tctatggcat ttatcacggt actcaaatg gtcatgtggt taattggcat gctggccacg 2100
ggcctcagtc ccaccaccg tactgacccc gatgaccca agatcacaat tgtctcctgt 2160
aaccccaact accgcaacag cctgctgttc aacaccagcc tggacctgct gctctcagtg 2220
gtgggtttca gcttcgcta catgggcaaa gagctgcca ccaactaaa cgaggccaag 2280
ttcatcacc tcagcatgac cttctatttc acctcatccg tctccctctg caccttcag 2340
tctgcctaca gcggggtgct ggtcaccatc gtggacctct tggctactgt gctcaacctc 2400
ctggccatca gcctgggcta cttcggcccc aagtgtaca tgatcctctt ctaccggag 2460
cgcaacacgc ccgcctactt caacagcatg atccagggt acaccatgag gagggactag 2520

```

<210> SEQ ID NO 24

<211> LENGTH: 6

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Six-His tag

<400> SEQUENCE: 24

His His His His His His

1

5

What is claimed:

1. An isolated nucleic acid sequence which encodes a human hT1R2 taste receptor that specifically responds to sweet taste stimuli wherein said hT1R2 nucleic acid sequence is selected from the following:

- (i) a nucleic acid sequence which encodes the human T1R2 polypeptide comprising SEQ ID NO: 21;
- (ii) a nucleic acid sequence which encodes a polypeptide having at least 95% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21;
- (iii) a nucleic acid sequence which hybridizes to the human T1R2 encoding nucleic acid sequence contained in SEQ ID NO: 20 under stringent hybridization conditions which consist of hybridization in 50% formamide, 5×SSC and 1% SDS, incubating at 42° C., with wash in 0.2×SSC and 0.1% SDS at 65° C., wherein the hybridization and wash steps are each carried out for at least 1 minute; and
- (iv) the nucleic sequence contained in SEQ ID NO: 20 encoding a human T1R2 polypeptide.

2. A human T1R2 nucleic acid sequence which encodes the human T1R2 polypeptide contained in SEQ ID NO: 21.

3. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses at least 95% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

4. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses greater than 95% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

5. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses at least 96% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

6. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses at least 97% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

7. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses at least 98% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

8. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses at least 99% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

9. The human T1R2 nucleic acid sequence of claim 1 which encodes a human T1R2 polypeptide which possesses greater than 99% sequence identity to the human T1R2 polypeptide contained in SEQ ID NO: 21.

10. The human T1R2 nucleic acid sequence of claim 1 which is a genomic sequence.

11. The human T1R2 nucleic acid sequence of claim 1 which is a cDNA.

12. The human T1R2 sequence of claims 1 which is an isolated mRNA.

13. The human T1R2 nucleic acid sequence of claim 1 which is operably linked to a nucleic acid sequence that facilitates the transcription of said human T1R2 nucleic acid sequence.

14. The human T1R2 nucleic acid sequence of claim 1 which is operably linked to a sequence that facilitates the surface expression of human T1R2 polypeptide by a host cell containing said nucleic acid sequence.

15. The human T1R2 nucleic acid sequence of claim 14 wherein said sequence that facilitates surface expression is from a mammalian rhodopsin gene.

16. The human T1R2 nucleic acid sequence of claims 1 which is contained on a nucleic acid sequence that further comprises a nucleic acid sequence which encodes a detectable marker.

17. The human T1R2 nucleic acid sequence of claim 16 wherein said detectable marker is a green fluorescent protein.

18. The human T1R2 nucleic acid sequence of claim 1 which is operably linked to a constitutive promoter.

19. The human T1R2 nucleic acid sequence of claim 1 which is operably linked to a regulatable promoter.

20. A human T1R2 nucleic acid sequence according to claim 1 which is contained in a nucleic acid construct.

21. The human T1R2 nucleic acid sequence containing construct of claim 20 which is a plasmid.

22. The human T1R2 nucleic acid sequence containing construct of claim 20 which is a viral vector.

23. The human T1R2 nucleic acid sequence containing construct of claim 20 which comprises a gene encoding a detectable marker.

24. The human T1R2 nucleic acid sequence containing construct of claim 20 wherein said human T1R2 nucleic acid sequence is operably linked to a promoter.

25. The human T1R2 nucleic acid sequence containing construct of claim 24 wherein said promoter is constitutive.

26. The human T1R2 nucleic acid sequence containing construct of claim 24 wherein said promoter is regulatable.

27. The human T1R2 nucleic acid sequence containing construct of claim 20 wherein said human T1R2 nucleic acid sequence is operably linked to a nucleic acid sequence that encodes a polypeptide that facilitates the surface expression of said human T1R2 polypeptide in a host cell containing the vector.

28. The human nucleic acid sequence containing construct of claim 27 wherein said polypeptide that facilitates surface expression is a mammalian rhodopsin polypeptide.

29. The human T1R2 nucleic acid sequence containing construct of claim 28 wherein said mammalian rhodopsin is bovine rhodopsin.

* * * * *