

US007351803B2

(12) **United States Patent**
Johnson et al.

(10) **Patent No.:** **US 7,351,803 B2**
(45) **Date of Patent:** **Apr. 1, 2008**

(54) **CD16A BINDING PROTEINS AND USE FOR THE TREATMENT OF IMMUNE DISORDERS**

EP 0 343 950 B1 10/2000
WO WO 00/42072 7/2000
WO WO 02/061090 8/2002

(75) Inventors: **Leslie S. Johnson**, Darnestown, MD (US); **Ling Huang**, Gaithersburg, MD (US); **Hua Li**, North Potomac, MD (US); **Nadine Tuailon**, Sykesville, MD (US)

(73) Assignee: **Macrogenics, Inc.**, Rockville, MD (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 353 days.

(21) Appl. No.: **10/449,566**

(22) Filed: **May 29, 2003**

(65) **Prior Publication Data**

US 2004/0010124 A1 Jan. 15, 2004

Related U.S. Application Data

(60) Provisional application No. 60/439,320, filed on Jan. 10, 2003, provisional application No. 60/384,689, filed on May 30, 2002.

(51) **Int. Cl.**
C07K 16/00 (2006.01)
C12P 21/08 (2006.01)
C07K 17/00 (2006.01)

(52) **U.S. Cl.** **530/387.1; 530/387.3; 530/388.1; 530/388.22**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,816,567 A	3/1989	Cabilly et al.
4,946,778 A	8/1990	Ladner et al.
5,116,964 A	5/1992	Capon et al.
5,219,728 A	6/1993	Khayat et al.
5,399,493 A	3/1995	Emerson et al.
5,428,130 A	6/1995	Capon et al.
5,437,994 A	8/1995	Emerson et al.
5,455,030 A	10/1995	Ladner et al.
5,545,806 A	8/1996	Lonberg et al.
5,569,825 A	10/1996	Lonberg et al.
5,624,821 A	4/1997	Winter et al.
5,641,863 A	6/1997	Schreiber et al.
5,834,597 A *	11/1998	Tso et al. 530/387.3
5,837,821 A	11/1998	Wu
5,976,831 A	11/1999	Peltz et al.
5,985,599 A	11/1999	McKenzie et al.
6,025,158 A	2/2000	Gonzalez et al.
6,103,889 A	8/2000	Whitlow et al.
6,194,551 B1	2/2001	Idusogie et al.
6,207,804 B1	3/2001	Huston et al.
6,277,375 B1	8/2001	Ward
6,528,624 B1	3/2003	Idusogie et al.

FOREIGN PATENT DOCUMENTS

EP 0 753 065 1/1997

OTHER PUBLICATIONS

Vaswani et al. *Annals of Allergy, Asthma, & Immunology*. 1998, 81:105-116, 119 (pp. 117-118 do not exit).*

Brummell et al. *Biochemistry*. 1993. 32;4:1180-1187.*

Rudikoff et al., *PNAS* 1982 vol. 79, pp. 1979-1983.*

Amit et al. *Science* 1986. 233:747-753.*

Alegre, M-L et al. (Jun. 1, 1992). "Effect of a Single Amino Acid Mutation on the Activating and Immunosuppressive Properties of a 'Humanized' OKT3 Monoclonal Antibody," *J. Immunol.* 148(11):3461-3468.

Altschul, S. F. et al. (1990). "Basic Local Alignment Search Tool," *J. Mol. Biol.* 215:403-410.

Anonymous (Dec. 20, 2000). "PerCP Mouse Anti-Human Monoclonal Antibody," PharMingen Technical Data Sheet, CD3/CD16/CD45, 2 pages.

Anonymous (Jun. 6, 2001). "R-Phycoerythrin (R-PE)-Conjugated Mouse Anti-Human Monoclonal Antibody," BD PharMingen Technical Data Sheet., CD16, one page.

Barbas, III, C. F. et al. (Sep. 1991). "Assembly of Combinatorial Antibody Libraries on Phage Surfaces: The Gene III Site," *PNAS* 88:7978-7982.

Bedzyk, W. D. et al. (Jan. 25, 1989). "Comparison of Variable Region Primary Structures Within Anti-Fluorescein Idiotype Family," *J. Biol. Chem.* 264(3):1565-1569.

Bird, R. E. et al. (Oct. 21, 1988). "Single-Chain Antigen-Binding Proteins," *Science* 242:423-426.

Boder, E. T. et al. (Sep. 26, 2000). "Directed Evolution of Antibody Fragments with Monovalent Femtomolar Antigen-Binding Affinity," *PNAS*. 97(20):10701-10705.

Boder, E. T. et al. (Jun. 1997). "Yeast Surface Display for Screening Combinatorial Polypeptide Libraries," *Nature Biotechnology* 15:553-557.

Bolland, S. et al. (1999). "Inhibitory Pathways Triggered by ITIM-Containing Receptors," *Adv. In Immunol.* 72:149-177.

Bussel, J. et al. (1988). "Infusion of a Monoclonal Anti-FcR III in Patients with Refractory ITP," *Neo-Adjuvant Chemotherapy* 169:883-887.

Canfield, S. M. et al. (1991). "The Binding Affinity of Human IgG for Its High Affinity Fc Receptor is Determined by Multiple Amino Acids in the C_H2 Domain and Is Modulated by Hinge Region," *J. Exp. Med.* 173:1483-1491.

Carter, P. et al. (1992). "High Level *Escherichia coli* Expression and Production of a Bivalent Humanized Antibody Fragment," *Bio/Technology* 10:163-167.

(Continued)

Primary Examiner—Maher M. Haddad

Assistant Examiner—Chun Crowder

(74) *Attorney, Agent, or Firm*—King & Spalding LLP; Margaret B. Brivanlou

(57) **ABSTRACT**

CD16A binding proteins useful for the reduction of a deleterious immune response are described. In one aspect, humanized anti-CD16A antibodies, optionally lacking effector function, are used for treatment of immune disorders such as idiopathic thrombocytopenic purpura and autoimmune hemolytic anemia.

18 Claims, 12 Drawing Sheets

OTHER PUBLICATIONS

- Casali, P. et al. (1986). "Human Monoclonals from Antigen-Specific Selection of B Lymphocytes and Transformation by EBV," *Science* 234:476-552.
- Cheung, R. C. et al. (1990). "Epitope-Specific Antibody Response to the Surface Antigen of Duck Hepatitis B Virus in Infected Ducks," *Virology* 176:546-552.
- Chothia, C. et al. (Dec. 1989). "Conformation of Immunoglobulin Hypervariable Regions," *Nature* 342:877-883.
- Clarkson, S. B. et al. (Aug. 1986). "Blockade of Clearance of Immune Complexes by an Anti-Fc γ Receptor Monoclonal Antibody," *J. Exp. Med.* 164:474-489.
- Clarkson, S. B. et al. (1986). "Treatment of Refractory Immune Thrombocytopenic Purpura with an Anti-Fc γ -Receptor Antibody," *N. Engl. J. Med.* 314(19):1236-1239.
- Clynes, R. A. et al. (Apr. 2000). "Inhibitory Fc Receptors Modulate In Vivo Cytotoxicity Against Tumor Targets," *Nat. Med.* 6(4):443-446.
- Co, M. S. et al. (Apr. 1991). "Humanized Antibodies for Antiviral Therapy," *PNAS* 88:2869-2873.
- Daugherty, P. S. et al. (1998). "Antibody Affinity Maturation Using Bacterial Surface Display," *Protein Enginee.* 11(9):825-832.
- Ding, Y.-H. et al. (Apr. 1998). "Two Human T Cell Receptors Bind in a Similar Diagonal Mode to the HLA-A2/Tax Peptide Complex Using Different TCR Amino Acids," *Immunity* 8:403-411.
- Duncan, A. R. et al. (1988). "Localization of the Binding Site for the Human High-Affinity Fc Receptor on IgG," *Nature* 332:563-564.
- Edge, J. C. et al. (1997). "Cell Type-Specific Glycoforms of Fc γ RIIIa (CD16): Differential Ligand Binding," *J. Immunol.* 159(8):3849-3857.
- Fleit, H. B. et al. (May 1982). "Human Neutrophil Fc γ Receptor Distribution and Structure," *PNAS* 79:3275-3279.
- Foote, J. et al. (1992). "Antibody Framework Residues Affecting the Conformation of the Hypervariable Loops," *J. Mol. Biol.* 224:487-499.
- Friend, P.J. et al. (1999). "Phase I Study of an Engineered Aglycosylated Humanized CD3 Antibody in Renal Transplant Rejection," *Transplantation* 68(11):1632-1637.
- Hoogenboom, H. R. and Winter, G. (1991). "By-Passing Immunisation Human Antibodies from Synthetic Repertoires of Germline VH Gene Segments Rearranged in Vitro," *J. Mol. Biol.* 227:381-388.
- Hudson, P. J. et al. (Jan. 2003). "Engineered Antibodies," *Nature Medicine* 9(1):129-139.
- Huse, W. D. et al. (Dec. 1989). "Generation of a Large Combinatorial Library of the Immunoglobulin Repertoire in Phage Lambda," *Science* 246:1275-1281.
- Jefferis, R. et al. (1995). "Recognition Sites on Human IgG for Fc γ Receptors: the Role of Glycosylation," *Immunology Letters* 44:111-117.
- Jefferis, R. et al. (1997). "Glycosylation of Antibody Molecules: Structural and Functional Significance," *Chem. Immunol.* 65:111-128.
- Jones, P. T. et al. (1986). "Replacing the Complementarity Determining Regions in a Human Antibody with Those from Mouse," *Nature* 321:522-525.
- Kirkland, T. N. et al. (1986). "Analysis of the Fine Specificity and Cross-Reactivity of Monoclonal Anti-Lipid A Antibodies," *J. Immunol* 137(11):3614-3619.
- Koumemis, I. L. et al. (2000). "Modulating Pharmacokinetics of An Anti-Interleukin-8 F(ab')₂ by Amine-Specific PEGylation with Preserved Bioactivity," *Int. J. Pharm.* 198:83-95.
- Krebs, B. et al. (2001). "High-Throughput Generation and Engineering of Recombinant Human Antibodies," *J. Imm. Methods* 254:67-84.
- Lanzavecchia, A. et al. (1987). "The Use of Hybridoma Target Human Cytotoxic T Lymphocytes," *Eur. J. Immunol.* 17:105-111.
- Larrick, J. W. et al. (1991). "Recombinant Antibodies," *Hum Antibodies Hybridomas* 2:172-189.
- Leong, S. R. et al. (2001). "Adapting Pharmacokinetic Properties of a Humanized Anti-Interleukin-8 Antibody for Therapeutic Applications Using Site-Specific Pegylation," *Cytokine* 16(3):106-119.
- Li, M. et al. (Mar. 1996). "Reconstitution of Human Fc γ RIII Cell Type Specificity in Transgenic Mice," *J. Exp. Med.* 183:1259-1263.
- Lucas, B. K. et al. (1996). "High-Level Production of Recombinant Proteins in CHO Cells Using A Dicistronic DHFR Intron Expression Vector," *Nucleic Acids Res.* 24(9):1774-1779.
- Malmqvist, M. et al. (1997). "Biomolecular Interaction Analysis: Affinity Biosensor Technologies for Functional Analysis of Proteins," *Curr. Opinion Chem. Biol.* 1:378-383.
- McCrae, K. R. et al. (2001). "Platelets: An Update on Diagnosis and Management of Thrombocytopenic Disorders," *Hematology* pp. 282-305.
- Mizutani, H. et al. (Aug. 1, 1993). "Development and Characterization of Monoclonal Antiplatelet Autoantibodies from Autoimmune Thrombocytopenic Purpura-Prone (NZW x BXSB)F₁ Mice," *Blood* 82(3):837-844.
- Moldenhauer, G. et al. (1990). "Identity of HML-1 Antigen on Intestinal Intraepithelial T Cells and of B-ly7 Antigen on Hairy Cell Leukaemia," *Scand. J. Immunol.* 32:77-82.
- Morel, G. A. et al. (1988). "Monoclonal Antibodies to Bovine Serum Albumin: Affinity and Specificity Determination," *Molec. Immunol.* 25(1):7-15.
- Needleman, S. B. et al. (1970). "A General Method Applicable to the Search for Similarities in the Amino Acid Sequence of Two Proteins," *J. Mol. Biol.* 48:443-453.
- O'Connell, D. et al. (2002). "Phage versus Phagemid Libraries for Generation of Human Monoclonal Antibodies," *J. Mol. Biol.* 321:49-56.
- Orlandi, R. et al. (May 1989). "Cloning Immunoglobulin Variable Domains for Expression by the Polymerase Chain Reaction," *PNAS* 86:3833-3837.
- Oyaizu, N. et al. (1988). "NZW x BXSB)F₁ Mouse: A New Animal Model of Idiopathic Thrombocytopenic Purpura," *J. Exp. Med.* 167:2017-2022.
- Padlan, E. A. (1991). "A Possible Procedure for Reducing the Immunogenicity of Antibody Variable Domains While Preserving Their Ligand-Binding Properties," *Mol. Immunol* 28(4/5):489-498.
- Pearson, W. R. et al. (1988). "Improved Tools for Biological Sequence Comparison," *PNAS* 85:2444-2448.
- Queen, C. et al. (1989). "A Humanized Antibody that Binds to the Interleukin 2 Receptor," *PNAS USA* 86:10029-10033.
- Radaev, S. et al. (May 11, 2001). "Recognition of IgG by Fc γ Receptor," *J. of Biological Chemistry* 276(19):16478-16483.
- Ravetch, J. V. et al. (Aug. 1989). "Alternative Membrane Forms of Fc γ RIII(CD16) on Human Natural Killer Cells and Neutrophils," *J. Exp. Med.* 170:481-497.
- Ravetch, J. V. et al. (2001). "I_GG Fc Receptors," *Annual Rev. Immunol.* 19:275-290.
- Ravtech, J. V. et al. (1991). "Fc Receptors," *Annual Rev. Immunol.* 9:457-492.
- Riechmann, L. et al. (1988). "Reshaping Human Antibodies for Therapy," *Nature* 332:323-327.
- Samuelsson, A. et al. (Jan. 19, 2001). "Anti-Inflammatory Activity of IVIG Mediated Through the Inhibitory Fc Receptor," *Science* 291:484-486.
- Scallon, B. J. et al. (1989). "A Human Immunoglobulin G Receptor Exists in Both Polypeptide-Anchored and Phosphatidylinositol-Glycan Anchored Forms," *PNAS* 86:5079-5093.
- Schier, R. et al. (1996). "Isolation of Picomolar Affinity Anti-c-erbB-2 Single-Chain Fv by Molecular Evolution of the Complementarity Determining Regions in the Center of the Antibody Binding Site," *J. Mol. Biol* 263:551-567.
- Schmidt, R. E. et al. (1989). "N2 Cluster Report: CD 16," In *Leukocyte Typing IV White Cell Differentiation Antigens*, Knapp, W. et al., eds., Oxford University Press, pp. 574-597.
- Shields, R. L. et al. (2001). "High Resolution Mapping of the Binding Site on Human IgG1 for Fc γ RI, Fc γ RIII, and FcRN and Design of IgG1 Variants with Improved Binding to the Fc γ R*," *J. of Biological Chemistry* 276(9):6591-6604.
- Smith, T. F. et al. (1981). "Comparison of Biosequence," *Adv. Appl. Math.* 2:482-489.

- Soubrane, C. et al. (Jan. 1, 1993). "Biologic Response Anti-CD16 Monoclonal Antibodies Therapy in a Human Immunodeficiency Virus-Related Immune Thrombocytopenic Purpura Patient," *Blood* 81(1):15-19.
- Stähli, C. et al. (1983). "Distinction of Epitopes by Monoclonal Antibodies," *Methods in Enzymology*, 92:242-253.
- Tao, M.-H. (Oct. 15, 1989). "Studies of Aglycosylated Chimeric Mouse-Human IgG: Role of Carbohydrate in the Structure and Effector Functions Mediated by the Human IgG Constant Region," *J. of Immunology* 143(8):2595-2601.
- Woof, J. et al. (1986). "Localisation of the Monocyte-Binding Region on Human Immunoglobulin G," *Mol. Imm.* 23(3):319-330.
- Xu, D. et al. (2000). "In Vitro Characterization of Five Humanized OKT3 Effector Function Variant Antibodies," *Cell Immunology* 200:16-26.
- Zu Putlitz, J. et al. (Jul. 1990) "Antibody Prosecution in Baculovirus Infected Insect Cells," *Bio/Technology* 8:651-654.
- International Search Report mailed on Oct. 20, 2003, for PCT application No. PCT/US03/17111 filed May 29, 2003, 6 pages.
- Tamm, A. et al. (1996). "The Binding Epitopes of Human CD 16 (FcγRIII) Monoclonal Antibodies," *Journal of Immunology* 157:1576-1581.
- Zhu, X. et al. (1998). "Intracellular Expression of FcγRIII (CD16) and Its Mobilization by Chemoattractants in Human Eosinophils¹," *Journal of Immunology* 161:2574-2579.
- Cartron et al., "Therapeutic activity of humanized anti-CD20 monoclonal antibody and polymorphism in IgG Fc receptor FcγRIIIa gene," *Blood* Feb. 1, 2002; 99(3):754-8.
- Clynes et al., "Uncoupling of immune complex formation and kidney damage in autoimmune glomerulonephritis," *Science* Feb. 13, 1998; 279(5353):1052-1054.
- Edberg JC, Kimberly, RP., "Cell type-specific glycoforms of Fc gamma RIIIa (CD16): differential ligand binding," *J. Immunol.* Oct. 15, 1997; 159(8):3849-3857.
- Farag, S.F. et al., "FcγRIIIa and FcγRIIIa Polymorphisms Do Not Predict Response to Rituximab in B-Cell Chronic Lymphocytic Leukemia," *Blood* Feb. 15, 2004; 103(4):1472-4; pre-published online Oct. 16, 2003.
- Fijen et al., "The role of Fcγreceptor polymorphisms and C3 in the immune defence against *Neisseria meningitidis* in complement-deficient individuals," *Clin Exp Immunol.* May 2000; 120(2):338-45.
- Frank et al., "Defective reticuloendothelial system Fc-receptor function in systemic lupus erythematosus," *N Engl J Med.* Mar. 8, 1979; 300(10):518-523.
- Fujiwara et al., "Determination of granulocyte-specific antigens on neutrophil FcA receptor IIIb by PCR-preferential homoduplex formation assay, and gene frequencies in the Japanese population," *Vox Sang.* 1999; 77(4):218-22.
- Hibbs et al., "Membrane-proximal Ig-like domain of Fc gamma RIII (CD16) contains residues critical for ligand binding," *J Immunol.* May 1, 1994; 152(9):4466-4474.
- Hosea et al., "Opsonic requirements for intravascular clearance after splenectomy," *N Engl J Med.* Jan. 29, 1981; 304(5):245-250.
- Imbach et al., "High-dose intravenous gammaglobulin for idiopathic thrombocytopenic purpura in childhood," *Lancet.* Jun. 6, 1981; 1(8232):1228-1231.
- Jones et al., "Two distinct classes of IgG Fc receptors on a human monocyte line (U937) defined by differences in binding of murine IgG subclasses at low ionic strength," *J Immunol.* Nov. 1985; 135(5):3348-53.
- Khayat et al., "Soluble circulating Fc gamma receptors in human serum. A new ELISA assay for specific and quantitative detection," *J Immunol Methods.* Jun. 26, 1987; 100(1-2):235-241.
- Kim et al., "Analysis of FcγRIII and IgG polymorphism reveals functional and evolutionary implications of protein-protein interaction," *J Mol Evol.* Jul. 2001; 53(1):1-9.
- Kurlander et al., "The blockade of Fc receptor-mediated clearance of immune complexes in vivo by a monoclonal antibody (2.4G2) directed against Fc receptors on murine leukocytes," *J Immunol.* Aug. 1984; 133(2):855-862.
- McDougal et al., "Binding of immunoglobulin G aggregates and immune complexes in human sera to Staphylococci containing protein A," *J Clin Invest.* Apr. 1979; 63(4):627-636.
- Nagarajan et al., "Ligand binding and phagocytosis by CD16 (Fc gamma receptor III) isoforms. Phagocytic signaling by associated zeta and gamma subunits in Chinese hamster ovary cells," *J Biol Chem.* Oct. 27, 1995; 270(43):25762-70.
- Olsson et al., "Serum from patients with chronic idiopathic thrombocytopenic purpura frequently affect the platelet function," *Thromb Res.* Aug. 15, 2002; 107(3-4):135-139.
- Parker CW., "Spectrofluorometric Methods." Chapter 14 of *Handbook of Experimental Immunology*. Weir ed. Blackwell Scientific Publications: pp. 14.1-14.25.
- Tebo et al., "Fcγ receptor-mediated phagocytosis of *Plasmodium falciparum*-infected erythrocytes in vitro," *Clin Exp Immunol.* Nov. 2002; 130(2):300-6.
- Todorovska et al., "Design and application of diabodies, triabodies and tetrabodies for cancer targeting," *J Immunol Methods.* Feb. 1, 2001; 248(1-2):47-66. Review.
- Treon, S.P. et al., "Polymorphisms in Fcγ RIIIa(CD16) Receptor Expression Are Associated with Clinical Response to Rituximab in Waldenström's Macroglobulinemia," *Biologic Therapy of Lymphomas: Laboratory Investigations.* 537a.
- Van Sorge et al., "FcγR polymorphisms: Implications for function, disease susceptibility and immunotherapy," *Tissue Antigens.* 2003; 61:189-202.
- Warmerdam et al., "Molecular basis for a polymorphism of human Fc gamma receptor II (CD32)," *J Exp Med.* Jul. 1, 1990; 172(1):19-25.
- West et al., "Natural cytotoxic reactivity of human lymphocytes against a myeloid cell line: characterization of effector cells," *J Immunol.* Jan. 1977; 118(1):355-361.
- Bolt et al., 1993, "The generation of a humanized, non-mitogenic CD3 monoclonal antibody which retains in vitro immunosuppressive properties," *Eur. J. Immunol.* 23(2):403-11.
- Bussel et al., 2000, "Fc receptor blockade and immune thrombocytopenic purpura," *Semin. Hematol.* 37(3):261-6.
- Cassatella et al., 1989, "Fc gamma R(CD16) interaction with ligand induces Ca²⁺ mobilization and phosphoinositide turnover in human natural killer cells. Role of Ca²⁺ in Fc gamma R(CD16)-induced transcription and expression of lymphokine genes," *J. Exp. Med.* 169(2):549-67.
- Kimberly et al., 1989, "In vivo handling of soluble complement fixing Ab/dsDNA immune complexes in chimpanzees," *J. Clin. Invest.* 84(3):962-70.

* cited by examiner

FIGURE 1


Binding of Mabs to sFcRIIIa

FIGURE 2




FIGURE 3

Competitive Cell Binding Assay




FIGURE 4




FIGURE 5




FIGURE 6




FIGURE 7




FIGURE 8




FIGURE 9




FIGURE 10




FIGURE 11




FIGURE 12




FIGURE 13(A)




FIGURE 13(B)




FIGURE 14




FIGURE 15(A)




FIGURE 15(B)

5 Hr Dose Response




FIGURE 16(A)




FIGURE 16(B)





FIGURE 17



1

CD16A BINDING PROTEINS AND USE FOR THE TREATMENT OF IMMUNE DISORDERS

CROSS-REFERENCE TO RELATED APPLICATIONS

This application claims benefit of provisional patent application No. 60/384,689, filed May 30, 2002, and provisional patent application No. 60/439,320, filed Jan. 10, 2003, the entire contents of which are incorporated herein by reference for all purposes.

FIELD OF THE INVENTION

The invention relates to CD16A binding proteins and methods for treatment of immune disorders. The invention finds application in the fields of biomedicine and immunology.

BACKGROUND

Fcγ receptors (FcγR) are cell surface receptors that bind the Fc region of immunoglobulin G (IgG) molecules. Among other functions, these receptors couple the formation of antibody-antigen complexes to effector cell responses. For example, cross-linking of activating Fcγ receptors by immune complexes can result in the phagocytosis of pathogens, killing of foreign and transformed cells by direct cytotoxicity, the clearance of toxic substances, and the initiation of an inflammatory response. Notably, the Fcγ receptors play a key role in autoimmunity. Autoantibody binding to activating Fc receptors triggers the pathogenic sequelae of autoimmune diseases such as idiopathic thrombocytopenic purpura, arthritis, systemic lupus erythematosus, autoimmune hemolytic anemia, and others.

In humans and rodents there are three classes of Fcγ receptors, designated FcγRI, FcγRII, and FcγRIII (see, Ravetch and Bolland, 2001 *Annual Rev. Immunol.* 19:275-90; and Ravetch and Kinet, 1991, *Annual Rev. Immunol.* 9:457-92). FcγRI sites are generally occupied by monomeric IgG, while RII and RIII receptors are generally unoccupied and available to interact with immune complexes. FcγRI, also called CD64, binds monomeric IgG with high affinity, and is present on monocytes and macrophages. FcγRII, also called CD32, binds to multimeric IgG (immune complexes or aggregated IgG) with moderate affinity, and is present on a variety of cell types, including B cells, platelets, neutrophils, macrophages and monocytes. FcγRIII, also called CD16, binds to multimeric IgG with moderate affinity and is the predominant activating FcγR on myeloid cells. FcγRIII is found in two forms. FcγRIIIA (CD16A), a transmembrane signaling form (50-65 kDa), is expressed by NK cells, monocytes, macrophages, and certain T cells. FcγRIIIB (CD16B), a glycosyl-phosphatidyl-inositol anchored form (48 kDa) form, is expressed by human neutrophils. See, e.g., Scallon et al., 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:5079-83 and Ravetch et al., 1989, *J. Exp. Med.* 170:481-97. Protein and nucleic acid sequences for CD16A are reported in Genbank as accession numbers P08637 (protein) and X52645 (nucleic acid) and in SWISS-PROT as accession number CAA36870. Protein and nucleic acid sequences for CD16B are reported in Genbank as accession numbers O75015 (protein) and X16863 (nucleic acid) and in SWISS-PROT as CAA34753.

2

SUMMARY OF THE INVENTION

In one aspect, the invention provides a CD16A binding protein that may be used for treatment of an individual with an autoimmune disease. CD16A binding proteins of the invention are other than mouse antibodies, and include chimeric, human and humanized anti-CD16A monoclonal antibodies, fragments thereof, single chain antibodies, and other binding proteins comprising a V_H domain and/or a V_L domain.

In one aspect the CD16A binding protein comprises a Fc region derived from a human IgG heavy chain (e.g., a Fc region derived from human IgG₁) where the Fc region lacks effector function and/or is modified to reduce binding to a Fc effector ligand. In one embodiment, the CD16A binding protein is not glycosylated, for example, due to a substitution at residue 297 of the Fc region.

In one aspect, the CD16A binding protein is a humanized 3G8 antibody with a V_H domain comprising three complementarity determining regions (CDRs) derived from the V_H domain of mouse monoclonal antibody 3G8. In one embodiment, the V_H domain has the sequence of the V_H domain of Hu3G8V_H-1. In one embodiment, the CDRs of the binding protein have the sequence of the mouse CDRs. In some versions, the V_H domain CDRs differ from those of 3G8 at least by one or more of the following substitutions: Val at position 34 in CDR1, Leu at position 50 in CDR2, Phe at position 52 in CDR2, Asn at position 54 in CDR2, Ser at position 60 in CDR2, Ser at position 62 in CDR2, Tyr at position 99 in CDR3, and Asp at position 101 of CDR3. In one embodiment, the V_H domain has the sequence of the V_H domain of Hu3G8V_H-22. In one embodiment V_H domain comprises an FR3 domain having the sequence of SEQ ID NO:51. The V_H domain may be linked to an antibody heavy chain constant domain, for example the human Cγ1 constant domain.

In some versions the CD16A binding protein has a V_H domain having a sequence set forth in Table 3. In some versions the CD16A binding protein has a V_H domain that differs from the sequence of Hu3G8VH-1 by one or more of the substitutions shown in Table 1.

In one aspect, the CD16A binding protein is a humanized 3G8 antibody with a V_L domain comprising three complementarity determining regions (CDRs) derived from the V_L domain of mouse monoclonal antibody 3G8. In one embodiment, the CDRs of the binding protein have the sequence of the mouse CDRs. In some versions, the V_L domain CDRs differ from those of 3G8 at least by one or more of the following substitutions: Arg at position 24 in CDR1; Ser at position 25 in CDR1; Tyr at position 32 in CDR1; Leu at position 33 in CDR1; Ala at position 34 in CDR1; Asp, Trp or Ser at position 50 in CDR2; Ala at position 51 in CDR2; Ser at position 53 in CDR2; Ala or Gln at position 55 in CDR2; Thr at position 56 in CDR2; Tyr at position 92 in CDR3; Ser at position 93 in CDR3; and Thr at position 94 in CDR3. In one embodiment, the V_L domain has the sequence of the V_L domain of Hu3G8VL-1, Hu3G8VL-22 or Hu3G8VL-43. The V_L domain may be linked to an antibody light chain constant domain, for example the human Cκ constant region.

In some versions the CD16A binding protein has a V_L domain having a sequence set forth in Table 4. In some versions the CD16A binding protein has a V_L domain that differs from the sequence of Hu3G8VL-1 by one or more of the substitutions shown in Table 2.

In one aspect, the CD16A binding protein comprises both a V_H domain and a V_L domain, as described above (which

may be prepared by coexpression of polynucleotides encoding heavy and light chains). Optionally the humanized heavy chain variable region comprises a sequence set forth in Table 3 and/or the humanized light chain variable region comprises a sequence set forth in Table 4. For example, in exemplary embodiments, the binding protein has a heavy chain variable region having the sequence of SEQ ID NO:113 and a light chain variable region having the sequence of SEQ ID NO:96, 100 or 118. In another exemplary embodiment, the binding protein has a heavy chain variable region having the sequence of SEQ ID NO:109 and light chain variable regions having the sequence of SEQ ID NO:96. In another exemplary embodiment, the binding protein has a heavy chain variable region having the sequence of SEQ ID NO:104 and a light chain variable region having the sequence of SEQ ID NO:96.

In an embodiment, the CD16A binding protein is tetrameric antibody comprising two light chains and two heavy chains, said light chains comprising a V_L domain and a light chain constant domain and said heavy chains comprising a V_H domain and a heavy chain constant domain. In an embodiment, the light chain constant domain is human C κ and/or the heavy chain constant region is C γ 1.

In one embodiment of the invention, the CD16A binding protein comprises an antigen binding site that binds CD16A or sCD16A with a binding constant of less than 5 nM.

In one embodiment, the CD16A binding protein comprises an aglycosyl Fc region that has reduced binding to at least one Fc effector ligand compared to a reference CD16A binding protein that comprises an unmodified Fc region (e.g., a human IgG₁ Fc domain glycosylated at position 297). The Fc effector ligand can be Fc γ RIII or the C1q component of complement.

In one embodiment, the invention provides a CD16A binding protein that is humanized antibody that binds to CD16A and inhibits the binding of Fc to CD16.

In an aspect, the invention provides a pharmaceutical composition comprising of CD16A binding protein described herein and a pharmaceutically acceptable excipient.

In an aspect, the invention provides an isolated polynucleotide, optionally an expression vector, encoding a V_H domain of a CD16A binding protein described herein. In an aspect, the invention provides an isolated nucleic acid, optionally an expression vector, encoding a V_L domain of a CD16A binding protein described herein. In an aspect, the invention provides a cell, optionally a mammalian cell, comprising a polynucleotide described herein. In an aspect, the invention a cell line, optionally a mammalian cell line, expressing a CD16A binding protein described herein.

The invention further provides a method of reducing an deleterious immune response (or undesired immune response) in a mammal comprising administering to a mammal a CD16A binding protein described herein. In an embodiment, reducing the deleterious immune response comprises protecting against antibody-mediated platelet depletion.

In one aspect, the invention provides a method of treating an deleterious immune response in a mammal without inducing neutropenia in the mammal (e.g., severe neutropenia or moderate neutropenia), where the method comprises administering to the mammal a CD16A binding protein having an Fc region derived from human IgG, and where the amino acid at position 297 of the Fc region is aglycosyl.

In embodiments of the above-described methods, the deleterious immune response is an inflammatory response, for example, an inflammatory response caused by an

autoimmune disease. In an embodiment, the inflammatory response is caused by idiopathic thrombocytopenic purpura (ITP), rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), autoimmune hemolytic anemia (AHA), scleroderma, autoantibody triggered urticaria, pemphigus, vasculitic syndromes, systemic vasculitis, Goodpasture's syndrome, multiple sclerosis (MS), psoriatic arthritis, ankylosing spondylitis, Sjögren's syndrome, Reiter's syndrome, Kawasaki's disease, polymyositis and dermatomyositis.

Other examples of diseases or conditions that can be treated according to the invention also include any diseases susceptible to treatment with intravenous immunoglobulin (IVIg) therapy (e.g., allergic asthma). The invention provides CD16A binding proteins that both protect against autoimmune diseases and do not result in significant neutrophil diminution in a mammal. In an embodiment, the CD16A binding proteins are anti-CD16A antibodies. These CD16A binding proteins are particularly advantageous for use as human therapeutics. In one aspect, the invention provides a method of treating an autoimmune disease in a mammal without neutrophil diminution or neutropenia in the mammal, by administering a CD16A binding protein having an Fc region derived from human IgG and an aglycosyl amino acid at position 297 of each of the C_H2 domains of the Fc region.

In yet another aspect, the invention provides a method of inhibiting the binding of IgG antibodies to Fc γ RIII on a cell by contacting the cell with a CD16A binding protein under conditions in which the CD16A binding protein binds the Fc γ RIII on the cell.

In one aspect, the invention provides a method of making a CD16A binding protein with improved therapeutic efficacy in treating an deleterious immune response, comprising the following steps: i) obtaining a first CD16A binding protein, where the first CD16A binding protein comprises an Fc region derived from IgG; and ii) modifying the Fc region of the first CD16A binding protein to produce a second CD16A binding protein that is aglycosylated at position 297 of the Fc region, where the second CD16A binding protein is more effective in treating the deleterious immune response when administered to a mammal than the first CD16A binding protein.

In one aspect, the invention provides a method of making a CD16A binding protein with improved therapeutic efficacy in treating an deleterious immune response, comprising the following steps: i) obtaining a first CD16A binding protein, wherein the first CD16A binding protein comprises an Fc region derived from IgG; and ii) modifying the Fc region of the first CD16A binding protein to produce a second CD16A binding protein that has reduced binding to an Fc effector ligand compared to the unmodified Fc region of the first CD16A binding protein, where the second CD16A binding protein is more effective in treating the deleterious immune response when administered to a mammal than the first CD16A binding protein. In one embodiment, the Fc effector ligand is Fc γ RIII or the C1q component of complement.

In one aspect the method involves administering a CD16A binding protein to reduce a deleterious immune response in a subject without eliciting one or more significant deleterious effects that result from 3G8 administration, or eliciting significantly lower levels of such effects than does administration of murine 3G8.

In one embodiment of the invention, the improved therapeutic efficacy in treating a deleterious immune response comprises improved effectiveness at protecting against antibody-mediated platelet depletion. The deleterious immune response is optionally due to idiopathic thrombocytopenic

5

purpura (ITP) or the administration of murine monoclonal antibody 6A6 to a muFcγRIII^{-/-}, huFcγRIIIA transgenic mouse.

The invention provides the use of a CD16A binding protein comprising an Fc region derived from a human IgG heavy chain, wherein the Fc region lacks effector function, for treatment of an immune disorder or for preparation of a medicament for treatment of an immune disorder.

BRIEF DESCRIPTION OF THE FIGURES

FIG. 1 shows results from an ELISA for binding of sCD16A by CD16A binding proteins. Hu3G8-24.43 is an antibody with the heavy chain Hu3G8VH-24, and the light chain Hu3G8VL-43. Hu3G8-5.1 is an antibody with the heavy chain Hu3G8VH-5, and the light chain Hu3G8VL-1. Ch3G8 is the chimeric 3G8 antibody. Hu1gG1 is an irrelevant immunoglobulin.

FIG. 2 shows results of an assay for binding of humanized and chimeric antibodies to CHO-K1 cells expressing the extracellular domain of CD16A. Hu3G8-22.1 is an antibody with the heavy chain Hu3G8VH-22, and the light chain Hu3G8VL-1. Hu3G8-5.1 is an antibody with the heavy chain Hu3G8VH-5, and the light chain Hu3G8VL-1. Hu3G8-22.43 is an antibody with the heavy chain Hu3G8VH-22, and the light chain Hu3G8VL-43. N297Q indicates the antibody is aglycosylated.

FIG. 3 shows results of a cell based competition assay. The aglycosylated humanized antibodies shown compete with aglycosylated chimeric antibody for binding to CHO-K1 cells expressing the extracellular domain of CD16A.

FIG. 4 shows inhibition of binding of sCD16A to immune complexes. Hu3G8-1.1 is an antibody with the heavy chain Hu3G8VH-1, and the light chain Hu3G8VL-1.

FIG. 5 shows ITP protection in mice injected i.v. with mAb 3G8 (0.5 μg/g) or human IVIG (1 mg/g) one hour before ch6A6 i.p. injection.

FIG. 6 shows ITP protection in mice injected i.v. with mAb 3G8 (0.5 μg/g) or human IVIG (1 mg/g) one hour before ch6A6 i.v. injection.

FIG. 7 shows the absence of ITP protection in mice injected i.v. with ch3G8 (0.5 μg/g) one hour before 6A6 i.p. injection.

FIG. 8 shows protection from ITP in mice injected i.v. with ch3G8 N297Q one hour before ch6A6 i.p. injection.

FIG. 9 shows protection from ITP in mice injected i.v. with ch3G8 N297Q one hour before ch6A6 i.v. injection.

FIG. 10 shows the results of FACS scans of neutrophils following administration of CD16A binding protein or controls. The x-axis shows labeling with antibody to CD16, and the y-axis shows labeling with antibody to the Gr-1 antigen. The upper right quadrant shows neutrophils; the upper left quadrant shows other granulocytes and neutrophils that no longer stain with 3G8-FITC.

FIG. 11 shows prevention of AIHA with a humanized anti-CD16 antibody.

FIG. 12 shows inhibition of ch4D5 mediated ADCC by humanized 3G8 antibodies.

FIGS. 13A-B show inhibition of ch4-4-20 mediated ADCC by mouse 3G8 (FIG. 13A) and humanized 3G8 antibodies (FIG. 13B).

FIG. 14 shows protection of FcγRIII^{-/-}, hCD16A, hCD32A mice against ITP by administration of hu3G8-5.1.

FIGS. 15A-B show protection of FcγRIII^{-/-}, hCD16A mice against ITP by administration of hu3G8-5.1 N297Q.

6

FIG. 15(A) shows data points for each dose at indicated times. FIG. 15(B) shows dose response at the 5 hour time point.

FIGS. 16A-B show the therapeutic effect of administration of aglycosylated humanized antibody subsequent to mice in which thrombocytopenia has been induced. FIG. 16(A) shows administration of Hu3G8-5.1-N297Q. FIG. 16(B) shows administration of Hu3G8-22.1-N297Q and Hu3G8-22.43-N297Q.

FIG. 17 shows the therapeutic effect of a humanized anti-CD16A antibody in treatment of autoimmune hemolytic anemia.

DETAILED DESCRIPTION

1. Definitions

Unless otherwise defined, all terms of art, notations and other scientific terms or terminology used herein are intended to have the meanings commonly understood by those of skill in the art to which this invention pertains. In some cases, terms with commonly understood meanings are defined herein for clarity and/or for ready reference, and the inclusion of such definitions herein should not necessarily be construed to represent a substantial difference over what is generally understood in the art. The practice of the present invention will employ, unless otherwise indicated, conventional techniques of molecular biology (including recombinant techniques), microbiology, cell biology, biochemistry, nucleic acid chemistry, and immunology, which are within the skill of the art. Such techniques are explained fully in the literature, such as, *Current Protocols in Immunology* (J. E. Coligan et al., eds., 1999, including supplements through 2001); *Current Protocols in Molecular Biology* (F. M. Ausubel et al., eds., 1987, including supplements through 2001); *Molecular Cloning: A Laboratory Manual*, third edition (Sambrook and Russet, 2001); *PCR: The Polymerase Chain Reaction*, (Mullis et al., eds., 1994); *The Immunoassay Handbook* (D. Wild, ed., Stockton Press NY, 1994); *Bioconjugate Techniques* (Greg T. Hermanson, ed., Academic Press, 1996); *Methods of Immunological Analysis* (R. Masseyeff, W. H. Albert, and N. A. Staines, eds., Weinheim: VCH Verlags gesellschaft mbH, 1993); Harlow and Lane, *Using Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y., 1999; and Beaucage et al. eds., *Current Protocols in Nucleic Acid Chemistry* John Wiley & Sons, Inc., New York, 2000).

The terms "heavy chain," "light chain," "variable region," "framework region," "constant domain," and the like, have their ordinary meaning in the immunology art and refer to domains in naturally occurring immunoglobulins and the corresponding domains of synthetic (e.g., recombinant) binding proteins (e.g., humanized antibodies). The basic structural unit of naturally occurring immunoglobulins (e.g., IgG) is a tetramer having two light chains and two heavy chains. Usually naturally occurring immunoglobulin is expressed as a glycoprotein of about 150,000 daltons, although, as described below, IgG can also be produced in a nonglycosylated form. The amino-terminal ("N") portion of each chain includes a variable region of about 100 to 110 or more amino acids primarily responsible for antigen recognition. The carboxy-terminal ("C") portion of each chain defines a constant region, with light chains having a single constant domain and heavy chains usually having three constant domains and a hinge region. Thus, the structure of the light chains of an IgG molecule is N-V_L-C_L-C and the structure of IgG heavy chains is N-V_H-C_{H1}-H-C_{H2}-C_{H3}-C

(where H is the hinge region). The variable regions of an IgG molecule consist of the complementarity determining regions (CDRs), which contain the residues in contact with antigen and non-CDR segments, referred to as framework segments, which maintain the structure and determine the positioning of the CDR loops. Thus, the V_L and V_H domains have the structure N-FR1, CDR1, FR2, CDR2, FR3, CDR3, FR4-C.

As used herein, the terms “CD16A binding protein,” “CD16A antibody,” and “anti-CD16A antibody,” are used interchangeably and refer to a variety of immunoglobulin-like or immunoglobulin-derived proteins. “CD16A binding proteins” bind CD16A via an interaction with V_L and/or V_H domains (as distinct from Fc-mediated binding). Examples of CD16A binding proteins include chimeric, humanized and human antibodies (e.g., comprising 2 heavy and 2 light chains), fragments thereof (e.g., Fab, Fab', F(ab')₂, and Fv fragments), bifunctional or multifunctional antibodies (see, e.g., Lanzavecchia et al., 1987, *Eur. J. Immunol.* 17:105), single chain antibodies (see, e.g., Bird et al., 1988, *Science* 242:423-26), fusion proteins (e.g., phage display fusion proteins), “minibodies” (see, e.g., U.S. Pat. No. 5,837,821) and other antigen binding proteins comprising a V_L and/or V_H domain or fragment thereof. In one aspect, the CD16A binding protein is a “tetrameric antibody,” i.e., having generally the structure of a naturally occurring IgG and comprising both variable and constant domains, (i.e., two light chains comprising a V_L domain and a light chain constant domain, such as human C κ and two heavy chains comprising a V_H domain and a heavy chain hinge and constant domains, such as human C γ 1). Except as expressly noted, the mouse antibody 3G8 is specifically excluded from the definition of CD16A binding protein.

When referring to binding proteins or antibodies (as broadly defined herein) the assignment of amino acids to each domain is in accordance with the definitions of Kabat, SEQUENCES OF PROTEINS OF IMMUNOLOGICAL INTEREST (National Institutes of Health, Bethesda, Md., 1987 and 1991). Amino acids from the variable regions of the mature heavy and light chains of immunoglobulins are designated by the position of an amino acid in the chain. Kabat described numerous amino acid sequences for antibodies, identified an amino acid consensus sequence for each subgroup, and assigned a residue number to each amino acid. Kabat's numbering scheme is extendible to antibodies not included in his compendium by aligning the antibody in question with one of the consensus sequences in Kabat by reference to conserved amino acids. This method for assigning residue numbers has become standard in the field and readily identifies amino acids at equivalent positions in different antibodies, including chimeric or humanized variants. For example, an amino acid at position 50 of a human antibody light chain occupies the equivalent position to an amino acid at position 50 of a mouse antibody light chain. Thus, as used herein in the context of chimeric or humanized antibodies, a reference such as “at position 297 of the Fc region” refers to the amino acid position in an immunoglobulin chain, region of an immunoglobulin chain, or region of a polypeptide derived from an immunoglobulin chain, that corresponds to position 297 of the corresponding human immunoglobulin.

The “Fc region” of immunoglobulins refers to the C-terminal region of an immunoglobulin heavy chain. Although the boundaries of the Fc region may vary somewhat, usually the Fc region is from about position 226-230 extending to the carboxy terminus of the polypeptide (and encompassing

the C_H2 and C_H3 domains). Sequences of human Fc regions are found in Kabat, supra. In addition, a variety of allotypic variants are known to exist.

An “Fc effector ligand” is a ligand that binds to the Fc region of an IgG antibody, thereby activating effector mechanisms resulting in the clearance and destruction of pathogens. Fc effector ligands include three cellular Fc receptors types—FcR γ I, FcR γ II, and FcR γ III. The multiple isoforms of each of the three Fc receptor types are also included. Accordingly, the term “Fc effector ligand” includes both FcR γ IIIA (CD16A) and FcR γ IIIB (CD16B). The term “Fc effector ligand” also includes the neonatal Fc receptor (Fc γ n) and the C1q component of complement. Binding of IgG to the Fc receptors triggers a variety of biological processes including antibody-dependent cell-mediated cytotoxicity (ADCC), release of inflammatory mediators, control of antibody production, clearance of immune complexes and destruction of antibody-coated particles. Binding of the C1q component of complement to IgG activates the complement system. Activation of complement plays important roles in opsonization, lysis of cell pathogens, and inflammatory responses.

As used herein, an Fc region that “lacks effector function” does not bind the Fc receptor and/or does not bind the C1q component of complement and trigger the biological responses characteristic of such binding.

The term “glycosylation site” refers to an amino acid residue that is recognized by a mammalian cell as a location for the attachment of sugar residues. Amino acid residues to which carbohydrates, such as oligosaccharides, are attached are usually asparagine (N-linkage), serine (O-linkage), and threonine (O-linkage) residues. The specific sites of attachment usually have a characteristic sequence of amino acids, referred to as a “glycosylation site sequence.” The glycosylation site sequence for N-linked glycosylation is: -Asn-X-Ser- or -Asn-X-Thr-, where X can be any of the conventional amino acids, other than proline. The Fc region of human IgG has two glycosylation sites, one in each of the C_H2 domains. The glycosylation that occurs at the glycosylation site in the C_H2 domain of human IgG is N-linked glycosylation at the asparagine at position 297 (Asn 297).

The term “chimeric,” when referring to antibodies, has the ordinary meaning in the art and refers to an antibody in which a portion of a heavy and/or light chain is identical to or homologous with an antibody from one species (e.g., mouse) while the remaining portion is identical to or homologous with an antibody of another species (e.g., human).

As used herein, the term “humanized” has its usual meaning in the art. In general terms, humanization of a non-human antibody involves substituting the CDR sequences from non-human immunoglobulin V_L and V_H regions into human framework regions. Further, as used herein, “humanized” antibodies may comprise additional substitutions and mutations in the CDR and/or framework regions introduced to increase affinity or for other purposes. For example, substitution of nonhuman framework residues in the human sequence can increase affinity. See, e.g., Jones et al., 1986, *Nature* 321:522-25; Queen et al., 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:10029-33; Foote and Winter, 1992, *J. Mol. Biol.* 224:487-99; Chothia et al., 1989, *Nature* 342:877-83; Riechmann et al., 1988, *Nature* 332:323-27; Co et al., 1991, *Proc. Natl. Acad. Sci. U.S.A.* 88:2869-73; Padlan, 1991, *Mol. Immunol.* 28:489-98. The resulting variable domains have non-human CDR sequences and framework sequences derived from human antibody framework sequence(s) or a human consensus sequence (e.g., as dis-

closed in Kabat, supra). A variety of different human framework regions may be used singly or in combination as a basis for the humanized monoclonal antibodies of the present invention. The framework sequences of a humanized antibody are "substantially human," by which is meant that at least about 70% of the human antibody sequence, usually at least about 80% human, and most often at least about 90% of the framework sequence is from human antibody sequence. In some embodiments, the substantially human framework comprises a serine at position 113 of the V_H FR4 domain (e.g., SEQ ID NO: 64). As used herein, a "humanized antibody" includes, in addition to tetrameric antibodies, single chain antibodies, antibody fragments and the like that comprise CDRs derived from a non-human antibody and framework sequences derived from human framework regions.

As used herein, "mammals" include humans, non-human primates, rodents, such as, mice and rats, and other mammals.

As used herein, "neutropenia" has its ordinary meaning, and refers to a state in which the number of neutrophils circulating in the blood is abnormally low. The normal level of neutrophils in human blood varies slightly by age and race. The average adult level is about 1500 cells/mm³ of blood. Neutrophil counts less than 500 cells/mm³ result in great risk of severe infection. Generally, in humans, severe neutropenia is defined by a blood neutrophil count less than about 500 cells/mm³, and moderate neutropenia is characterized by a blood neutrophil count from about 500-1000 cells/mm³.

As used herein, "treatment" refers to clinical intervention in an attempt to alter the disease course of the individual or cell being treated, and can be performed either for prophylaxis or during the course of clinical pathology. Therapeutic effects of treatment include without limitation, preventing occurrence or recurrence of disease, alleviation of symptoms, diminishment of any direct or indirect pathological consequences of the disease, decreasing the rate of disease progression, amelioration or palliation of the disease state, and remission or improved prognosis.

An "effective amount" is an amount sufficient to effect a beneficial or desired clinical result upon treatment. An effective amount can be administered to a patient in one or more doses. A "therapeutically effective amount" is an amount that is sufficient to palliate, ameliorate, stabilize, reverse or slow the progression of the disease, or otherwise reduce the pathological consequences of the disease, or reduce the symptoms of the disease. The amelioration or reduction need not be, and usually is not, permanent, but may be for a period of time ranging from at least one hour, at least one day, or at least one week or more. The effective amount is generally determined by the physician on a case-by-case basis and is within the skill of one in the art. Several factors are typically taken into account when determining an appropriate dosage to achieve an effective amount. These factors include age, sex and weight of the patient, the condition being treated, the severity of the condition and the form and effective concentration of the binding protein administered. An "inflammation reducing amount" is an amount that reduces inflammation in a subject. A reduction in inflammation can be assessed by art known criteria, including decreased C-reactive protein levels, decreased consumption of complement, reduced immune complex deposition at sites of inflammation (e.g., joints in subjects with RA, kidney in subjects with lupus, myelin sheath, etc.), reduced cytokine release, migration of macrophages and neutrophils, and the like.

"Substantial sequence identity," as used herein, refers to two or more sequences or subsequences (e.g., domains) that have at least about 80% amino acid residue identity, preferably at least about 90%, or at least about 95% identity when compared and aligned for maximum correspondence. Sequence identity between two similar sequences (e.g., antibody variable regions) can be measured by (1) aligning the sequences to maximize the total number of identities across the entire length of the sequences, or across the entire length of the shorter of the two sequences, if of different lengths (and where the length of the aligned sequences or shorter of the aligned sequences is "L" residues); (2) counting the number of positions (not including the number "E" residues designated as excluded from the comparison) at which there is an amino acid identity, where the number of identities is designated "N"; (3) and dividing the N by the "L" minus "E." For example, in a comparison of two sequences each of length 80 residues, in which 6 specific residues are excluded from the comparison and for which there are 65 identities in the remaining 74 positions, the sequence identity would be N/(L-E) or 65/(80-6) or 87.8%. (Residues might be specified as "excluded" from the calculation when, for illustration but not limitation, they are in a non-antibody domain of fusion protein.) Alternatively, optimal alignment and sequence identity can be calculated by computerized implementations of algorithms described in Smith & Waterman, 1981, *Adv. Appl. Math.* 2:482 [local homology algorithm], Needleman & Wunsch, 1970, *J. Mol. Biol.* 48:443 [homology alignment algorithm], Pearson & Lipman, 1988, *Proc. Natl. Acad. Sci. USA* 85:2444 [search for similarity method], or Altschul et al., 1990, *J. Mol. Biol.* 215:403-10 [BLAST algorithm]. See Ausubel et al., supra and GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package, Genetics Computer Group, 575 Science Dr., Madison, Wis.). When using any of the aforementioned algorithms, the default parameters (for Window length, gap penalty, etc.) are used. An amino acid or nucleic acid sequence is "substantially similar to" a second sequence when the degree of sequence identity is at least about 70% identical, preferably at least about 80%, or at least about 90%, or even at least about 95%, identical. Sequences that are substantially identical are also substantially similar.

As used herein, a polypeptide, polypeptide domain or region, or amino acid sequence is "derived from" another when the two sequences are identical or substantially similar and have a similar biological function. For example, in a humanized mouse monoclonal antibody the complementary determining regions (CDRs) are "derived from" the corresponding CDRs of the mouse monoclonal antibody, and the variable domain framework regions can be "derived from" framework sequences of the corresponding human antibody. It will be apparent that one domain, etc., can be derived from a parental domain, etc., even though the two differ in sequence due to, for example, the introduction of mutations that affect, or alternatively do not change, binding affinity or other properties of the protein in which the domain, etc., is contained, such as those described herein. It will also be understood that normally a domain, etc., "derived from" a parental domain, etc., is made, produced or designed using materials (e.g. genetic material) or information (e.g., nucleotide or amino acid sequence) from the parental molecule.

Standard abbreviations are used for amino acids: alanine, Ala (A); serine, Ser (S); threonine, Thr (T); aspartic acid, Asp (D); glutamic acid, Glu (E); asparagine, Asn (N); glutamine, Gln (Q); arginine, Arg (R); lysine, Lys (K); isoleucine, Ile (I); leucine, Leu (L); methionine, Met (M); valine, Val (V); phenylalanine, Phe (F); tyrosine, Tyr (Y);

tryptophan, Trp (W); glycine, Gly (G); histidine, His (H); proline, Pro (P); and cysteine, Cys (C).

2. Introduction

The FcγRIIIA receptor, CD16A, plays a role in coupling cytotoxic and immune complex antibodies to effector responses. It is believed that the interaction of the FcγRIIIA receptor and immunoglobulin aggregates (e.g. immune complexes) present in autoimmune diseases and other pathogenic conditions results in a deleterious inflammatory response in subjects. Without intending to be bound by a specific mechanism, it is believed that reducing the interaction of the FcγRIIIA receptor (generally referred to herein as “CD16A” or “the CD16A receptor”) and immunoglobulin aggregates will alleviate this inflammatory response. Also without intending to be bound by a specific mechanism, it is believed that one method for reducing the interaction of CD16A and immunoglobulin aggregates is by use of anti-CD16A antibodies, or other CD16A binding proteins, to block the interaction.

Monoclonal antibody 3G8 (“mAb 3G8”) is a mouse monoclonal antibody that binds the Fc-binding domain of human CD16A and B with a K_a of $1 \times 10^9 \text{ M}^{-1}$ (Fleit et al., 1982, *Proc. Natl. Acad. Sci. U.S.A* 79:3275-79). 3G8 blocks the binding of human IgG₁ immune complexes to isolated human NK cells, monocytes and neutrophils, as well as to CD16A-transfected 293 cells. Experiments in which mAb 3G8 has been administered to human patients for treatment of idiopathic thrombocytopenic purpura (ITP) have been conducted (Clarkson et al., 1986, *N. Engl. J Med.* 314:1236-39; Soubrane, et al., 1993, *Blood* 81:15-19). Administration of the 3G8 antibody was reported to result in increased platelet levels and was accompanied by one or more significant side effects, including a HAMA response, cytokine release syndrome, and/or pronounced neutropenia.

The present invention provides novel CD16A binding proteins, including humanized and/or aglycosylated monoclonal antibodies, and methods for reducing a deleterious immune response in a subject by administering the proteins. Administration of these binding proteins is shown to be protective in well established models for two distinct autoimmune diseases: autoimmune hemolytic anemia (AHA) and idiopathic thrombocytopenic purpura. These results are indicative of efficacy of this treatment for other autoimmune diseases as well. Moreover, the inventors have discovered that, unexpectedly, administration of anti-CD16A antibodies with altered effector function (e.g., aglycosylated antibodies) protects against the deleterious immune responses characteristic of autoimmune disorders without inducing acute severe neutropenia. Thus, the invention provides new reagents and methods for antibody-mediated effected treatment of autoimmune conditions without pronounced side-effects observed using alternative treatments.

3. CD16A Binding Proteins

A variety of CD16A binding proteins may be used in the methods of the invention. Suitable CD16A binding proteins include human or humanized monoclonal antibodies as well as CD16A binding antibody fragments (e.g., scFv or single chain antibodies, Fab fragments, minibodies) and another antibody-like proteins that bind to CD16A via an interaction with a light chain variable region domain, a heavy chain variable region domain, or both.

In some embodiments, the CD16A binding protein for use according to the invention comprises a V_L and/or V_H domain that has one or more CDRs with sequences derived from a non-human anti-CD16A antibody, such as a mouse anti-

CD16A antibody, and one or more framework regions derived from framework sequences of one or more human immunoglobulins. A number of non-human anti-CD16A monoclonal antibodies, from which CDR and other sequences may be obtained, are known (see, e.g., Tamm and Schmidt, 1996, *J. Imm.* 157:1576-81; Fleit et al., 1989, p. 159; LEUKOCYTE TYPING II: HUMAN MYELOID AND HEMATOPOIETIC CELLS, Reinherz et al., eds. New York: Springer-Verlag; 1986; LEUCOCYTE TYPING III: WHITE CELL DIFFERENTIATION ANTIGENS, McMichael A J, ed., Oxford: Oxford University Press, 1986); LEUKOCYTE TYPING IV: WHITE CELL DIFFERENTIATION ANTIGENS, Kapp et al., eds. Oxford Univ. Press, Oxford; LEUKOCYTE TYPING V: WHITE CELL DIFFERENTIATION ANTIGENS, Schlossman et al., eds. Oxford Univ. Press, Oxford; LEUKOCYTE TYPING VI: WHITE CELL DIFFERENTIATION ANTIGENS, Kishimoto, ed. Taylor & Francis. In addition, as shown in the Examples, new CD16A binding proteins that recognize human CD16A expressed on cells can be obtained using well known methods for production and selection of monoclonal antibodies or related binding proteins (e.g., hybridoma technology, phage display, and the like). See, for example, O’Connel et al., 2002, *J. Mol. Biol.* 321:49-56; Hoogenboom and Chames, 2000, *Imm. Today* 21:371078; Krebs et al., 2001, *J. Imm. Methods* 254:67-84; and other references cited herein. Monoclonal antibodies from a non-human species can be chimerized or humanized using techniques of antibody humanization known in the art.

Alternatively, fully human antibodies against CD16A can be produced using transgenic animals having elements of a human immune system (see, e.g., U.S. Pat. Nos. 5,569,825 and 5,545,806), using human peripheral blood cells (Casali et al., 1986, *Science* 234:476), by screening a DNA library from human B cells according to the general protocol outlined by Huse et al., 1989, *Science* 246:1275, and by other methods.

It is contemplated that, for some purposes, it may be advantageous to use CD16A binding proteins that bind the CD16A receptor at the same epitope bound by 3G8, or at least sufficiently close to this epitope to block binding by 3G8. Methods for epitope mapping and competitive binding experiments to identify binding proteins with the desired binding properties are well known to those skilled in the art of experimental immunology. See, for example, Harlow and Lane, cited supra; Stahl et al., 1983, *Methods in Enzymology* 9:242-53; Kirkland et al., 1986, *J. Immunol.* 137:3614-19; Morel et al., 1988, *Molec. Immunol.* 25:7-15; Cheung et al., 1990, *Virology* 176:546-52; and Moldenhauer et al., 1990, *Scand. J. Immunol.* 32:77-82. Also see Examples and §3G (i), infra. For instance, it is possible to determine if two antibodies bind to the same site by using one of the antibodies to capture the antigen on an ELISA plate and then measuring the ability of the second antibody to bind to the captured antigen. Epitope comparison can also be achieved by labeling a first antibody, directly or indirectly, with an enzyme, radionuclide or fluorophore, and measuring the ability of an unlabeled second antibody to inhibit the binding of the first antibody to the antigen on cells, in solution, or on a solid phase.

It is also possible to measure the ability of antibodies to block the binding of the CD16A receptor to immune complexes formed on ELISA plates. Such immune complexes are formed by first coating the plate with an antigen such as fluorescein, then applying a specific anti-fluorescein antibody to the plate. This immune complex then serves as the ligand for soluble Fc receptors such as sFcRIIIa. Alternatively a soluble immune complex may be formed and labeled, directly or indirectly, with an enzyme radionuclide

or fluorophore. The ability of antibodies to inhibit the binding of these labeled immune complexes to Fc receptors on cells, in solution or on a solid phase can then be measured.

CD16A binding proteins of the invention may or may not comprise a human immunoglobulin Fc region. Fc regions are not present, for example, in scFv binding proteins. Fc regions are present, for example, in human or humanized tetrameric monoclonal IgG antibodies. As described in detail below, in some embodiments of the present invention, the CD16A binding protein includes an Fc region that has an altered effector function, e.g., reduced affinity for an effector ligand such as an Fc receptor or C1 component of complement compared to the unaltered Fc region (e.g., Fc of naturally occurring IgG₁, proteins). In one embodiment the Fc region is not glycosylated at the Fc region amino acid corresponding to position 297. Such antibodies lack Fc effector function.

Thus, in some embodiments of the invention, the CD16A binding protein does not exhibit Fc-mediated binding to an effector ligand such as an Fc receptor or the C1 component of complement due to the absence of the Fc domain in the binding protein while, in other cases, the lack of binding or effector function is due to an alteration in the constant region of the antibody.

CD16A Binding Proteins Comprising CDR Sequences Similar to mAb 3G8 CDR Sequences

CD16A binding proteins that can be used in the practice of the invention include proteins comprising a CDR sequence derived from (i.e., having a sequence the same as or similar to) the CDRs of the mouse monoclonal antibody 3G8. Complementary cDNAs encoding the heavy chain and light chain variable regions of the mouse 3G8 monoclonal antibody, including the CDR encoding sequences, were cloned and sequenced as described in the Examples. The nucleic acid and protein sequences of 3G8 are provided below and are designated SEQ ID NO:1 and 2 (V_H) and SEQ ID NO:3 and 4 (V_L). Using the mouse variable region and CDR sequences, a large number of chimeric and humanized monoclonal antibodies, comprising complementary determining regions derived from 3G8 CDRs were produced and their properties analyzed. To identify humanized antibodies that bind CD16A with high affinity and have other desirable properties, antibody heavy chains comprising a V_H region with CDRs derived from 3G8 were produced and combined (by coexpression) with antibody light chains comprising a V_L region with CDRs derived from 3G8 to produce a tetrameric antibody for analysis. Properties of the resulting tetrameric antibodies were determined as described below. As described below, CD16A binding proteins comprising 3G8 CDRs, such as the humanized antibody proteins described hereinbelow, may be used according to the invention to reduce an deleterious immune response.

A. V_H Region

In one aspect, the CD16A binding protein of the invention may comprise a heavy chain variable domain in which at least one CDR (and usually three CDRS) have the sequence of a CDR (and more typically all three CDRS) of the mouse monoclonal antibody 3G8 heavy chain and for which the remaining portions of the binding protein are substantially human (derived from and substantially similar to, the heavy chain variable region of a human antibody or antibodies).

In an aspect, the invention provides a humanized 3G8 antibody or antibody fragment containing CDRs derived from the 3G8 antibody in a substantially human framework, but in which at least one of the CDRs of the heavy chain

variable domain differs in sequence from the corresponding mouse antibody 3G8 heavy chain CDR. For example, in one embodiment, the CDR(S) differs from the 3G8 CDR sequence at least by having one or more CDR substitutions shown in Table 1 (e.g., valine at position 34 in CDR1, leucine at position 50 in CDR2, phenylalanine at position 52 in CDR2, tyrosine at position 52 in CDR2, aspartic acid at position 52 in CDR2, asparagine at position 54 in CDR2, serine at position 60 in CDR2, serine at position 62 in CDR2, tyrosine at position 99 in CDR3, and/or aspartic acid at position 101 of CDR3). Suitable CD16 binding proteins may comprise 0, 1, 2, 3, or 4, or more of these substitutions (and often have from 1 to 4 of these substitutions) and optionally can have additional substitutions as well.

In one embodiment, a CD16A binding protein may comprise a heavy chain variable domain sequence that is the same as, or similar to, the V_H domain of the Hu3G8VH-1 construct, the sequence of which is provided in Table 3. For example, the invention provides a CD16A binding protein comprising a V_H domain with a sequence that (1) differs from the V_H domain of Hu3G8VH-1 by zero, one, or more than one of the CDR substitutions set forth in Table 1; (2) differs from the V_H domain of Hu3G8VH-1 by zero, one or more than one of the framework substitutions set forth in Table 1; and (3) is at least about 80% identical, often at least about 90%, and sometimes at least about 95% identical, or even at least about 98% identical to the Hu3G8VH-1 V_H sequence at the remaining positions.

Exemplary V_H domains of CD16 binding proteins of the invention have the sequence of Hu3G8VH-5 and Hu3G8VH-22, as shown in Tables 3 and 6.

The V_H domain may have a sequence that differs from that of Hu3G8VH-1 (Table 3) by at least one, at least two, at least three, at least four, at least five, or at least six of the substitutions shown in Table 1. These substitutions are believed to result in increased affinity for CD16A and/or reduce the immunogenicity of a CD16A binding protein when administered to humans. In certain embodiments, the degree of sequence identity with the Hu3G8VH-1 V_H domain at the remaining positions is at least about 80%, at least about 90%, at least about 95% or at least about 98%.

TABLE 1

V _H Domain Substitutions			
No.	Kabat Position	Region	Substitutions
1	2	FR1	Ile
2	5	FR1	Lys
3	10	FR1	Thr
4	30	FR1	Arg
5	34	CDR1	Val
6	50	CDR2	Leu
7	52	CDR2	Phe or Tyr or Asp
8	54	CDR2	Asn
9	60	CDR2	Ser
10	62	CDR2	Ser
11	70	FR3	Thr
12	94	FR3	Gln or Lys or Ala or His
13	99	CDR3	Tyr
14	101	CDR3	Asp

For illustration and not limitation, the sequences of a number of CD16A building protein V_H domains are shown

in Table 3. As described in the Examples, *infra*, heavy chains comprising these sequences fused to a human C γ 1 constant region were coexpressed with the hu3G8VL-1 light chain (described below) to form tetrameric antibodies, and binding of the antibodies to CD16A was measured to assess the effect of amino acid substitutions compared to the hu3G8VH-1 V_H domain. Constructs in which the V_H domain has a sequence of hu3G8VH-1, 2, 3, 4, 5, 8, 12, 14, 16, 17, 18, 19, 20, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 42, 43, 44 and 45 showed high affinity binding, with hu3G8VH-6 and -40 V_H domains showing intermediate binding. CD16A binding proteins comprising the V_H domains of hu3G8VH-5 and hu3G8VH-22 are considered to have particularly favorable binding properties.

B. V_L Region

Similar studies were conducted to identify light chain variable domain sequences with favorable binding properties. In one aspect, the invention provides a CD16A binding protein containing a light chain variable domain in which at least one CDR (and usually three CDRs) has the sequence of a CDR (and more typically all three CDRs) of the mouse monoclonal antibody 3G8 light chain and for which the remaining portions of the binding protein are substantially human (derived from and substantially similar to, the heavy chain variable region of a human antibody or antibodies).

In one aspect, the invention provides a humanized 3G8 antibody or antibody fragment containing CDRs derived from the 3G8 antibody in a substantially human framework, but in which at least one of the CDRs of the light chain variable domain differs in sequence from the mouse monoclonal antibody 3G8 light chain CDR. In one embodiment, the CDR(s) differs from the 3G8 sequence at least by having one or more amino acid substitutions in a CDR, such as, one or more substitutions shown in Table 2 (e.g., arginine at position 24 in CDR1; serine at position 25 in CDR1; tyrosine at position 32 in CDR1; leucine at position 33 in CDR1; aspartic acid, tryptophan or serine at position 50 in CDR2; serine at position 53 in CDR2; alanine or glutamine at position 55 in CDR2; threonine at position 56 in CDR2; serine at position 93 in CDR3; and/or threonine at position 94 in CDR3). In various embodiments, the variable domain can have 0, 1, 2, 3, 4, 5, or more of these substitutions (and often have from 1 to 4 of these substitutions) and optionally, can have additional substitutions as well.

In one embodiment, a suitable CD16A binding protein may comprise a light chain variable domain sequence that is the same as, or similar to, the V_L domain of the Hu3G8VL-1 construct, the sequence of which is provided in Table 4. For example, the invention provides a CD16A binding protein comprising a V_L domain with a sequence that (1) differs from the V_L domain of Hu3G8VL-1 by zero, one, or more of the CDR substitutions set forth in Table 2; (2) differs from the V_L domain of Hu3G8VL-1 by zero, one or more of the framework substitutions set forth in Table 2; and (3) is at least about 80% identical, often at least about 90%, and sometimes at least about 95% identical, or even at least about 98% identical to the Hu3G8VL-1 V_L sequence at the remaining positions.

Exemplary V_L domains of CD16 binding proteins of the invention have the sequence of Hu3G8VL-1 or Hu3G8VL-43, as shown in Tables 4 and 6.

The V_L domain may have a sequence that differs from that of Hu3G8VL-1 (Table 4) by zero, one, at least two, at least 3, at least 4, at least 5, at least 6, at least 7, at least 8, or at least 9 of the substitutions shown in Table 2. These substitutions are believed to result in increased affinity for CD16A

and/or reduce the immunogenicity of a CD16A binding protein when administered to humans. In certain embodiments, the degree of sequence identity at the remaining positions is at least about 80%, at least about 90% at least about 95% or at least about 98%.

TABLE 2

V _L Domain Substitutions			
No.	Kabat Position	Region	Substitutions
1	24	CDR1	Arg
2	25	CDR1	Ser
3	32	CDR1	Tyr
4	33	CDR1	Leu
5	50	CDR2	Asp or Trp or Ser
6	51	CDR2	Ala
7	53	CDR2	Ser
8	55	CDR2	Ala or Gln
9	56	CDR2	Thr
10	93	CDR3	Ser
11	94	CDR3	Thr

For illustration and not limitation, the sequences of a number of CD16A binding proteins V_L domains are shown in Table 4. As described in the Examples, *infra*, light chains comprising these sequences fused to a human CK constant domain were coexpressed with the Hu3G8VH-1 heavy chain (described above) to form tetrameric antibodies, and the binding of the antibodies to CD16A was measured to assess the effect of amino acid substitutions compared to the Hu3G8VL-1 V_L domain. Constructs in which the V_L domain has a sequence of hu3G8VL-1, 2, 3, 4, 5, 10, 16, 18, 19, 21, 22, 24, 27, 28, 32, 33, 34, 35, 36, 37, and 42 showed high affinity binding and hu3G8VL-15, 17, 20, 23, 25, 26, 29, 30, 31, 38, 39, 40 and 41 showed intermediate binding. CD16A binding proteins comprising the V_L domains of hu3G8VL-1, hu3G8VL-22, and hu3G8VL-43 are considered to have particularly favorable binding properties.

C. Combinations of V_L and/or V_H Domains

As is known in the art and described elsewhere herein, immunoglobulin light and heavy chains can be recombinantly expressed under conditions in which they associate to produce a tetrameric antibody, or can be so combined *in vitro*. Similarly, combinations of V_L and/or V_H domains can be expressed in the form of single chain antibodies, and still other CD16A binding proteins that comprise a V_L and/or V_H domain can be expressed by known methods. It will thus be appreciated that a 3G8-derived V_L-domain described herein can be combined with a 3G8-derived V_H-domain described herein to produce a CD16A binding protein, and all such combinations are contemplated.

For illustration and not for limitation, examples of useful CD16A binding proteins are those comprising at least one V_H domain and at least one V_L domain, where the V_H domain is from hu3G8VH-1, hu3G8VH-22 or hu3G8VH-5 and the V_L domain is from hu3G8VL-1, hu3G8VL-22 or hu3G8VL-43. In particular, humanized antibodies that comprise hu3G8VH-22 and either hu3G8VL-1, hu3G8VL-22 or hu3G8VL-43, or hu3G8VH-5 and hu3G8VL-1 have favorable properties.

It will be appreciated by those of skill that the sequences of V_L and V_H domains described here can be further modified by art-known methods such as affinity maturation (see Schier et al., 1996, *J. Mol. Biol.* 263:551-67; Daugherty et

al., 1998, *Protein Eng.* 11:825-32; Boder et al., 1997, *Nat. Biotechnol.* 15:553-57; Boder et al., 2000, *Proc. Natl. Acad. Sci. U.S.A* 97:10701-705; Hudson and Souriau, 2003, *Nature Medicine* 9:129-39). For example, the CD16A binding proteins can be modified using affinity maturation techniques to identify proteins with increased affinity for CD16A and/or decreased affinity for CD16B.

D. Constant Domains and Fc Region

As noted above, the CD16A binding protein of the invention may contain light chain and/or heavy chain constant regions (including the hinge region connecting the C_{H1} and C_{H2} domains in IgG molecules). It is contemplated that a constant domain from any type (e.g., IgM, IgG, IgD, IgA and IgE) of immunoglobulin may be used. The constant domain for the light chain can be lambda or kappa. The constant domain for the heavy chain can be any isotype (e.g., IgG₁, IgG₂, IgG₃ and IgG₄). Chimeric constant domains, portions of constant domains, and variants of naturally occurring human antibody constant domains (containing deletions, insertions or substitutions of amino acid residues) may be used. For instance, a change in the amino acid sequence of a constant domain can be modified to provide additional or different properties, such as altered immunogenicity or half-life of the resultant polypeptide. The changes range from insertion, deletion or substitution of a small number (e.g., less than ten, e.g., one, two, three or more) amino acid residues to substantial modifications of the constant region domain. Changes contemplated include those that affect the interaction with membrane receptors, complement fixation, persistence in circulation, and other effector functions. For example, the hinge or other regions can be modified as described in U.S. Pat. No. 6,277,375. In particular, it will often be advantageous to delete or alter amino acids of the Fc region. For example, the Fc region can be modified to reduce or eliminate binding to Fc effector ligands such as FcγRIII and the C1q component of complement, such that the antibodies lack (or have substantially reduced) effector function. Antibodies having such modified Fc regions induce little or no antibody dependent cellular cytotoxicity (ADCC) and/or complement mediated lysis when administered to a mammal, compared to unmodified antibodies. Assays to identify antibodies lacking effector function are known in the art. See, e.g., U.S. Pat. Nos. 6,194,551; 6,528,624; and 5,624,821, European Pat. No. EP 0 753 065 B1, and PCT publication WO 00/42072.

The CD16A binding protein of the invention may include a human IgG₁ Fc domain comprising one or more amino acid substitutions or deletions (relative to the parental naturally occurring IgG₁) that result in a reduced interaction between the Fc domain of the binding protein and FcγRIIA and/or FcγRIIIA (e.g., to minimize potential activation of macrophages and/or minimize neutrophil diminution) and/or increased binding of the Fc region to FcγRIIB (e.g., to increase FcγRIIB-mediated inhibition of effector cell activation; see Bolland and Ravetch, 1999, *Adv. in Immunol.* 72:149). Specific mutations effecting the desired changes in binding can be identified by selection using display of mutant Fc libraries expressed on the surface of microorganisms, viruses or mammalian cells, and screening such libraries for mutant Fc variants having the desired property or properties. In addition, the literature reports that particular residues or regions of the Fc are involved in Fcγ interactions such that deletion or mutation of these residues would be expected to result in reduced FcR binding. The binding site on human antibodies for FcγR was reported to be the residues 233-239 (Canfield et al., 1991, *J Exp Med* 173:

1483-91; Woof et al., 1986, *Mol. Imm.* 23:319-30; Duncan et al., 1988, *Nature* 332:563). The crystal structure of FcγRIII complexed with human IgG1 Fc revealed potential contacts between the receptor and its ligand and also revealed that a single FcγRIII monomer binds to both subunits of the Fc homodimer in an asymmetric fashion. Alanine-scanning mutagenesis of the Fc region confirmed the importance of most of the predicted contact residues (Shields et al., 2001, *J Biol. Chem.* 276:6591-6604).

Exemplary Fc region mutations include, for example, L235E, L234A, L235A, and D265A, which have been shown to have low affinity for all FcR, into Cγ-1 (Clynes et al., 2000, *Nat. Med.* 6:443-46; Alegre et al., 1992, *J Immunol* 148:3461-68; Xu et al., 2000, *Cell Immunol* 200:16-26). Additional Fc region modifications purported to affect FcR binding are described in WO 00/42072 (e.g., "class 4" Fc region variants) and WO 02/061090.

Fc binding to FcγRIIA and FcγRIIIA or other proteins can be measured by any of a number of methods, including ELISA to measure binding to isolated recombinant FcγR and RIA or FACS to measure binding to cells. Immune complexes and heat aggregated or chemically crosslinked Fc or IgG can be used to test affinity for FcRs in such assays. In one embodiment, immune complexes are produced by expressing an Fc in the context of an Fab with affinity for an antigen (such as fluorescein) and mixing the antibody and antigen to form an immune complex.

E. Fc Regions with Reduced Binding to Fc Effector Ligands Due to Aglycosylation or Changes in Glycosylation

As discussed above, in CD16A binding proteins of the invention that comprise Fc domains (e.g., anti-CD16A monoclonal antibodies) the Fc domain can be modified to achieve desired properties. In a particular aspect, the invention provides a CD16A binding protein, such as a human or humanized anti-CD16A monoclonal antibody, comprising an Fc region that is not glycosylated. As demonstrated in Example 10, infra, the inventors have discovered that, unexpectedly, administration of anti-CD16A antibodies with altered effector function (aglycosylated antibodies) protects against autoimmune disorders without inducing acute severe neutropenia. On the basis of this discovery, therapeutic anti-CD16A antibodies can be designed to protect against autoimmune diseases without inducing dangerous side effects.

In one embodiment, the invention provides a CD16A binding protein comprising an Fc region derived from human IgG₁, where the amino acids corresponding to position 297 of the C_{H2} domains of the Fc region are aglycosyl. The terms "aglycosyl" or "aglycosylated," when referring to an Fc region in its entirety, or a specific amino acid residue in the Fc region, mean that no carbohydrate residues are attached to the specified region or residue.

Human IgG antibodies that are aglycosylated show decreased binding to Fc effector ligands such as Fc receptors and C1q (see, e.g., Jefferis et al., 1995, *Immunology Letters* 44:111-17; Tao, 1989, *J. of Immunology*, 143:2595-2601; Friend et al., 1999, *Transplantation* 68:1632-37; Radaev and Sun, 2001, *J. of Biological Chemistry* 276:16478-83; Shields et al., 2001, *J. of Biological Chemistry* 276:6591-6604, and U.S. Pat. No. 5,624,821). Without intending to be bound by a particular mechanism, it is believed that the aglycosylation of the amino acid at position 297 of the Fc domains of CD16A binding proteins described herein results in reduced binding to CD16A and the C1q component of complement. Such aglycosylated antibodies lack effector function.

In human IgG₁ constant regions, the residue at position 297 is asparagine. In one embodiment of the present invention, the residue at, or corresponding to, position 297 of the Fc region of the CD16A binding protein is other than asparagine. Substitution of another amino acid residue in the place of asparagine eliminates the N-glycosylation site at position 297. Substitution of any amino acid residues which will not result in glycosylation upon expression of the CD16A binding protein in a mammalian cell is appropriate for this embodiment. For instance, in some embodiments of the invention, the amino acid residue at position 297 is glutamine or alanine. In some embodiments, the amino acid residue at position 297 is cysteine, which is optionally linked to PEG.

In other embodiments of the invention, the residue at position 297 may or may not be asparagine, but is not glycosylated. This can be accomplished in a variety of ways. For example, amino acid residues other than the asparagine at position 297 are known to be important for N-linked glycosylation at position 297 (see Jefferis and Lund, 1997, *Chem. Immunol.* 65:111-28), and the substitution of residues at positions other than position 297 of the C_H2 domain can result in a CD16A binding protein aglycosylated at residue 297. For illustration and not limitation, a residue at position 299 in the C_H2 domain that is other than threonine or serine will result in an antibody that is aglycosylated at position 297. Similarly, substitution of the amino acid at position 298 with proline will produce an antibody with an aglycosylated amino acid at position 297. In other embodiments of the invention, Fc domains of IgG₂ or IgG₄ are used rather than IgG₁ domains.

Modification of the amino acid residues of CD16A binding proteins is well within the ability of the ordinarily skilled practitioner, and can be achieved by mutation of a polynucleotide encoding the binding protein or portion thereof. The CD16A binding protein comprising an IgG-derived Fc region need not necessarily be mutated at the amino acid level to be aglycosylated. Binding proteins aglycosylated at position 297 of the IgG-derived Fc region can be produced by expressing the CD16A binding protein in certain cells (e.g., *E. coli*; see PCT publication WO 02061090A2), cell lines or under certain cell culture growth conditions where glycosylation at Asn 297 does not take place. Alternatively, carbohydrate groups may be removed from a CD16A binding protein following expression of the protein, e.g., enzymatically. Methods for removing or modifying carbohydrate groups on proteins are known and include use of endoglycosidases and peptide:N-glycosidases.

It will be apparent that a variety of methods can be used to modify the Fc region of a CD16A binding protein to change its properties. Accordingly, unless otherwise specified, as used herein the term "modifying" in the context of modifying the Fc region of a CD16A binding protein includes modifying the protein itself directly, modifying the polynucleotide that encodes the protein and/or modifying or selecting a suitable expression system for production of the protein.

In addition to CD16A binding proteins that are aglycosylated at the position corresponding to arginine 297, variants with reduced binding to Fc effector ligands due to only partial removal, or modification, of the carbohydrate at that position may be used in the present invention. For example, the Fc region can be modified to include a non-naturally occurring carbohydrate that does not bestow binding protein with effector function. As used herein, a "modified Fc

region" is an Fc region that has been derived from a parent Fc region, but which differs in glycosylation pattern from the parent Fc region.

F. Production of CD16A Binding Proteins

CD16A binding proteins of the invention can be produced using a variety of methods well known in the art, including de novo protein synthesis and recombinant expression of nucleic acids encoding the binding proteins. The desired nucleic acid sequences can be produced by recombinant methods (e.g., PCR mutagenesis of an earlier prepared variant of the desired polynucleotide) or by solid-phase DNA synthesis. Usually recombinant expression methods are used. In one aspect, the invention provides a polynucleotide that comprises a sequence encoding a CD16A binding protein disclosed herein or a CD16A binding fragment thereof, for example a sequence encoding a V_L or V_H described herein, or antibody heavy chain or light chain described herein. Because of the degeneracy of the genetic code, a variety of nucleic acid sequences encode each immunoglobulin amino acid sequence, and the present invention includes all nucleic acids encoding the binding proteins described herein.

Recombinant expression of antibodies is well known in the art and can be carried out, for example, by inserting nucleic acids encoding light and heavy chain variable regions, optionally linked to constant regions, into expression vectors. Expression vectors typically include control sequences such as a promoter, an enhancer, and a transcription termination sequence to which DNA segments encoding polypeptides (e.g., immunoglobulin chains) are operably linked to ensure the expression of immunoglobulin polypeptides. Expression vectors are typically replicable in the host organisms either as episomes or as an integral part of the host chromosomal DNA. The light and heavy chains can be cloned in the same or different expression vectors.

Immunoglobulin light and heavy chains are expressed using standard methods. A multiple polypeptide chain antibody or antibody fragment species can be made in a single host cell expression system wherein the host cell produces each chain of the antibody or antibody fragment and assembles the polypeptide chains into a multimeric structure to form the antibody or antibody fragment in vivo. See e.g., Lucas et al., 1996, *Nucleic Acids Res.*, 24:1774-79. When heavy and light chains are cloned on separate expression vectors, the vectors are co-transfected to obtain expression and assembly of intact immunoglobulins. Alternatively, recombinant production of antibody heavy and light chains in separate expression hosts followed by assembly of antibody from separate heavy and light chains in vitro is known. See, e.g., U.S. Pat. No. 4,816,567 and Carter et al., 1992, *Bio/Technology* 10:163-67.

The CD16A binding proteins are conveniently expressed in prokaryotic or eukaryotic cells. Useful hosts for antibody expression include bacteria (see, e.g., PCT publication WO 02/061090), yeast (e.g., *Saccharomyces*), insect cell culture (Putlitz et al., 1990, *Bio/Technology* 8:651-54), plants and plant cell cultures (Larrick and Fry, 1991, *Hum. Antibodies Hybridomas* 2:172-89), and mammalian cells. Methods for expression are well known in the art. For example, in *E. coli*, vectors using the lac promoter to drive expression of heavy and light chains fused to various prokaryotic secretion signal sequences such as pelB have resulted in successful secretion of scFv and Fab fragments into the periplasmic space or into the culture medium (Barbas et al., 1991, *Proc. Natl. Acad.*

Sci. U.S.A. 88:7978-82). A vector derived from pET25b in which the lac promoter has been inserted in place of the T7 promoter may be used.

Mammalian cells are especially useful for producing CD16A binding proteins, including tetrameric antibodies or fragments thereof. A number of suitable host cell lines capable of secreting intact heterologous proteins are known, and include CHO cell lines, COS cell lines, HeLa cells, L cells and myeloma cell lines. Expression vectors for mammalian cells can include expression control sequences, such as an origin of replication, a promoter, an enhancer, ribosome binding sites, RNA splice sites, polyadenylation sites, and transcriptional terminator sequences. Examples of expression control sequences are promoters derived from endogenous genes, cytomegalovirus, SV40, adenovirus, bovine papillomavirus, and the like. In one embodiment, binding proteins are expressed using the CMV immediate early enhancer/promoter in the vector pCDNA3.1 or a similar vector. To facilitate secretion, the genes can be fused to a gene cassette containing the signal sequence of a mouse VH gene described by Orlandi et al., 1989, *Proc. Natl. Acad. Sci. U.S.A.* 86:3833-37, which has been widely used for high-level secretion of immunoglobulins.

The vectors containing the DNA segments encoding the polypeptides of interest can be transferred into the host cell using routine methods, depending on the type of cellular host. For example, calcium chloride transfection is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment, electroporation, lipofection, biolistics or viral-based transfection may be used for other cellular hosts. Other methods used to transform mammalian cells include the use of polybrene, protoplast fusion, liposomes, electroporation, and microinjection (see generally, Sambrook et al., supra). For transient expression, cells, e.g., HEK293, can be co-transfected with separate heavy and light chain expression vectors using a cationic lipid (e.g., LIPO-FECTAMINE™ 2000, Invitrogen). This method can achieve expression levels of 10-20 mg/l of IgG in conditioned medium after 3 days. The cells can then be re-fed and similar quantities harvested after 3 more days. It will be appreciated that, for some uses, the cells expressing CD16A binding proteins can be maintained in medium containing FBS screened for very low levels of bovine IgG, or, alternatively, in serum-free medium.

In addition to expression of tetrameric antibodies, single chain antibodies, antibody fragments, and other CD16A binding proteins can be prepared. For example, immunoglobulin fragments can be prepared by proteolytic digestion of tetrameric antibodies, or more often, by recombinant expression of truncated antibody constructs. Usually, single chain V region ("scFv") constructs are made by linking V_L and/or V_H domain using a short linking peptide (see, e.g., Bird et al., 1988, *Science* 242:423-26; U.S. Pat. Nos. 4,946,778; 5,455,030; 6,103,889; and 6,207,804).

Once expressed, the binding proteins can be purified using procedures well known in the art, including ammonium sulfate precipitation, affinity chromatography, gel electrophoresis and the like (see, generally, Harris and Angal, 1990, *PROTEIN PURIFICATION APPLICATIONS, A PRACTICAL APPROACH* Oxford University Press, Oxford, UK; and Coligan et al., supra). In one embodiment, purification is accomplished by capturing the antibody using a high flow rate protein A resin such as Poros A (Perseptive Biosystems, Inc), and elution at low pH, followed by size exclusion chromatography to remove any traces of aggregate present. Since FcγRIIIA binds preferentially to aggregated IgG, removal of aggregates will be desirable for certain applications. The binding

proteins can be purified to substantial purity if desired, e.g., at least about 80% pure, often at least about 90% pure, more often least about 95%, or at least about 98% pure. In this context, the percent purity is calculated as a weight percent of the total protein content of the preparation, and does not include constituents which are deliberately added to the composition after the binding protein is purified.

CD16A binding proteins can be modified after expression. For example, derivation of antibodies with polyethylene glycol ("pegylation") is reported to increase residence time (half-life and stability) and reduce immunogenicity in vivo without alteration of biological activity. See, e.g., Leong et al., 2001, *Cytokine* 16:106-19; Koumenis et al., 2000, *Int J Pharm* 198:83-95; U.S. Pat. No. 6,025,158. CD16A binding proteins can be conjugated to a detectable label or ligand (e.g., a radioisotope or biotin). Other modifications are well known in the art and are also contemplated.

G. Properties of CD16A Binding Proteins

In certain embodiments, CD16A binding proteins having properties as described below are used in the methods of the invention.

i) Binding Affinity

CD16A binding proteins can be described by reference to their binding properties and biological activity. In various embodiments, the binding constant for the interaction of a CD16A binding protein of the invention and CD16A is between 0.1 and 5 nM, less than about 2.5 nM, less than about 1 nM, or less than about 0.5 nM. Usually the binding protein binds CD16A with an affinity that is within 4-fold, optionally within 2-fold, of the binding affinity exhibited under similar conditions by 3G8 or the chimeric antibody comprising the heavy chain Ch3G8VH and the light chain Ch3G8VL as described herein below. In an embodiment, the binding affinity for CD16A is greater than that of 3G8. In an alternative embodiment, the binding affinity for CD16B is no greater than, and preferably less than, 3G8 or the chimeric antibody Ch3G8.

Binding can be measured using a variety of methods, including ELISA, biosensor (kinetic analysis), and radioimmunoassay (RIA). ELISA is well known (see, Harlow and Lane, supra, and Ausubel et al., supra) and can be carried out using conditioned medium containing binding proteins or, alternatively, with purified antibodies. The concentration of antibody that results in 50% apparent maximal binding provides an estimate of antibody K_d .

Binding can also be detected using a biosensor assay, which also provides information on the kinetic and equilibrium properties of antibody binding to FcγRIIIA. An exemplary biosensor assay uses the BIAcore system (Malmqvist et al., 1997, *Curr. Opin. Chem. Biol.* 1:378-83). The BIAcore system relies on passing analyte over a sensor chip onto which the ligand (e.g., CD16A) is immobilized. The binding of the analyte can be measured by following surface plasmon resonance (SPR) signal, which changes in direct proportion to the mass bound to the chip. A fixed concentration of analyte is passed over the chip for a specific amount of time, allowing for the measurement of the association rate, k_{on} . Following this phase, buffer alone is passed over the chip and the rate at which the analyte dissociates from the surface, k_{off} can be measured. The equilibrium dissociation constant can be calculated from the ratio of the kinetic constants; $K_d = k_{on}/k_{off}$.

A radioimmunoassay (RIA) can be used to measure the affinity of antibodies for FcγRIII-bearing cells, and to measure inhibition of IgG complexes to cells by these antibodies. In an exemplary assay, ^{125}I labeled binding protein is

prepared and specific radioactivity of the protein determined. Labeled binding protein and cells are mixed for several hours, the cells and bound material are separated from the unbound material by centrifugation, and the radioactivity in both compartments is determined. A direct binding format is used to determine the K_d of, and the number of binding sites for, iodinated binding protein using Scatchard analysis of the binding data. Controls containing an excess of cold (unlabeled) binding protein competitor can be included to ensure the results reflect specific interactions. Examples of suitable cells include (1) NK cells or macrophages derived from normal human peripheral blood lymphocytes; (2) Cells obtained from huCD16A transgenic mice (Li, 1996 *J. Exp. Med.* 183:1259-63); (3) mammalian cell lines expressing the extracellular portion of CD16A fused to the transmembrane and intracellular domain of RII or another receptor (such as CD8 or LFA-3); (4) mammalian cell lines (e.g., CHO, HEK-293, COS) transfected transiently or stably with CD16A expression vectors (and optionally coexpressing gamma chain for optimal expression receptor expression).

Examples of expression vectors useful for expression of CD16A and other polypeptides for use in binding assays include mammalian expression vectors (e.g., pCDNA 3.1 or pCI-neo) that contain a strong promoter/enhancer sequence (e.g., CMV immediate early) and a polyadenylation/transcription termination site flanking a polylinker region into which the CD16A gene is introduced. Usually the vector includes a selectable marker such as a neomycin resistance gene.

In one embodiment, the CD16A expressed for use in assays has the sequence:

(SEQ ID NO:116)

MWQLLLPTALLLLVSAGMRTE³⁵DLPKAVVFLEPQWYRVLEKDSVTLKCQGA
 YSPEDNSTQWFHNESLISSQASSYFIDAATVDDSGEYRCQTNLSTLSDPV
 QLEVHIGWLLLQAPRWVFKEEDPIHLRCHSWKNTALHKV⁴⁰TYLQNGKGRKY
 FHHNSDFYIPKATLKDSGSYFCRGLFGSKNVSETV⁴⁵NITITQGLAVSTIS
 SFFPPGYQVSFCLVMVLLFAVDTGLYFSVKTNIRSSSTRDWDHKFKWRKD
 PQDK.

CD16A with the sequence:

(SEQ ID NO:117)

MWQLLLPTALLLLVSAGMRTE⁵⁰DLPKAVVFLEPQWYRVLEKDSVTLKCQGA
 YSPEDNSTQWFHNESLISSQASSYFIDAATVDDSGEYRCQTNLSTLSDPV
 QLEVHIGWLLLQAPRWVFKEEDPIHLRCHSWKNTALHKV⁵⁵TYLQNGKGRKY
 FHHNSDFYIPKATLKDSGSYFCRGLVGSKNVSETV⁶⁰NITITQGLAVSTIS
 SFFPPGYQVSFCLVMVLLFAVDTGLYFSVKTNIRSSSTRDWDHKFKWRKD
 PQDK

can also be used. Additional CD16A variants and substitutes will be known to, or readily discernible from the scientific literature by, the ordinarily skilled reader.

Competitive assay formats can be used to measure the ability of a CD16A binding protein to inhibit binding of another molecule to the receptor. For example, in one competitive assay format a fixed amount of labeled 3G8 is mixed with varying amounts of either unlabeled 3G8,

CD16A binding protein or an irrelevant IgG (control) and added to FcγRIIIA expressing cells. After incubation and separation of the cell-bound material from the material free in solution, the amount of bound labeled 3G8 (and/or optionally also the unbound labeled 3G8) is determined. The concentration of unlabeled mAb which results in a 50% decrease in the binding of labeled 3G8 (IC_{50}) is then determined from this data.

ii. Blocking Immune Complex Binding to FcγRIIIA

Another characteristic of the CD16A binding proteins of the invention is the ability to inhibit binding of immune complexes to CD16A ("IC Blocking Activity"). Usually the binding protein has IC Blocking Activity that is within 4-fold, preferably within 2-fold, of the activity exhibited under similar conditions by 3G8 or the chimeric antibody, Ch3G8, described herein.

Assays for measuring ability of an antibody to block binding of complexed IgG to CD16 are known. See, e.g., Knapp et al., 1989, *LEUKOCYTE TYPING IV*, Oxford University Press, Oxford, p. 574-97; and Edberg and Kimberly, 1997, *J Immunol* 159:3849-57. One suitable assay is an RIA assay with the format described above for the competitive assay, but substituting ¹²⁵I-labeled aggregated irrelevant human IgG for the ¹²⁵I-labeled 3G8 used in the competitive assay described above.

The invention provides a method of inhibiting the binding of IgG antibodies to CD16 on a cell by contacting the cell with a CD16A binding protein under conditions in which the CD16A binding protein binds the FcγRIII on the cell. The contacting can be in vivo (e.g., by administering the binding protein in a mammal) or in vitro (e.g., by addition of antibodies to cultured cells expressing the FcγRIII). IgG antibodies that are inhibited from binding the FcγRIII can be administered to the animal or added to a culture medium before or after addition or administration of the binding protein, or may be present in an animal normally or in response to a disease state. In one embodiment, the CD16 on the surface of the cell is CD16A.

iii. In Vivo Protection Against Platelet Depletion

The ability of the CD16A binding proteins of the invention to reduce deleterious immune responses can be assessed in a variety of animal models. An exemplary model system is a mouse model for idiopathic thrombocytopenic purpura (ITP) (see, Oyaizu et al., 1988, *J Exp. Med.* 167:2017-22; Mizutani et al., 1993, *Blood* 82:837-44). See Example 9, *infra*. Other suitable models are known in the art. Other animal models include rodent models of inflammatory diseases described in, for example, *Current Protocols in Immunology* (in some cases modified by using animals transgenic for human CD16A). Transgenic mice can be made using routine methods or can be purchased from commercial sources (e.g., Taconic Inc., German Town, N.Y.).

An example of a procedure suitable for assessing the ability of a CD16A binding protein to provide protection from platelet depletion in a mouse model is described in Example 8, *infra*. CD16A binding proteins can be administered to muFcγRIII^{-/-}, huFcγRIIIA transgenic mice at a variety of concentrations, and ITP subsequently induced in the mice (e.g., by administering the 6A6 or chimeric 6A6 antibody) to the mice. At timed intervals after the administration of 6A6/ch6A6, the mice are bled and the platelet counts are determined. Optionally, the IC_{50} for each molecule is then determined at the time point where maximal platelet depletion is observed in the negative control group. Based on the results of Example 8 and on prior studies, maximum depletion occurred 2-6 hr after 6A6 administration. IC_{50} s are determined graphically, using a curve-fitting program such as the four-parameter fit provided in the SigmaPlot program.

Statistically significant inhibition of depletion of platelets after administration of 6A6 in the treatment group compared to the untreated group and a group administered an identical formulation of an irrelevant, isotype matched mAb is indicative of the desired biological activity.

Experiments in which protection by CD16A binding proteins was assayed are described in the Examples, *infra*. Preparations of recombinant mouse 3G8 produced in HEK-293 cells, chimeric 3G8 with human IgG1 or IgG2 constant domains (ch3G8- γ 1 produced in HEK-293 and CHO-K1 cells, and ch3G8- γ 2 produced in HEK-293 cells), and a ch3G8- γ 1 variant (ch3G8- γ 1 D265A) did not provide significant protection. Murine 3G8, produced from the hybridoma, and a chimeric version of 3G8 containing an aglycosylated human G1 constant region (Ch3G8-G1 N297Q), produced in HEK-293 cells, were able to protect animals from platelet depletion in the mouse model. As shown in Example 10, 11 and 15-17, *infra*, Ch3G8 N297Q and aglycosylated humanized antibodies protected against platelet depletion in the ITP mouse model. Although not intending to be bound by a particular theory, one possibility is that since ch3G8 N297Q is largely devoid of effector function, it is more efficient than ch3G8 in protecting mice against ITP. Thus, these data suggest that anti-CD16A antibodies without effector function (e.g., aglycosylated antibodies) have advantages compared to some glycosylated (e.g., glycosylated recombinant) antibodies. Further, as described in the examples, administration of aglycosylated anti-CD16A antibody to muFc γ RIII $^{-/-}$, huFc γ RIIB transgenic mice did not result in neutrophil depletion in the blood, spleen, and bone marrow. Without intending to be bound by a particular theory, there are several possible explanations for these unexpected results. Protein glycosylation is known to vary in different cell lines, especially those from different species. A difference in the nature of the carbohydrate attached to the antibody constant region as a consequence of expression in different cell types may be responsible for the difference in activity, i.e., if the lack of activity results in part from effector cell activation caused by ch3G8 binding to Fc receptors (or complement) via the antibody Fc region in a glycosylation-dependent manner. Alternatively, recombinant murine and ch3G8 may contain other post-translational modifications that affect activity and which can be eliminated by using different cell lines to express the CD16A binding proteins. It is possible that a combination of isotype and/or isotype containing mutations to eliminate effector function may provide similar protective effects as elimination of the carbohydrate on the Fc.

5. Methods of Treatment

A number of diseases and conditions characterized by a deleterious immune response can be treated using the binding proteins of the invention, i.e., a CD16A binding protein as described herein (e.g., comprising a V_L and/or V_H sequence as disclosed herein and, optionally, a Fc region modified as disclosed herein to have a reduced effector function). In one embodiment, the binding protein is administered to a subject with an autoimmune disease (i.e., a disease characterized by the production of autoantibodies). It is believed that pathogenic IgG antibodies observed in autoimmune diseases are either the pathogenic triggers for these diseases or contribute to disease progression and mediate disease through the inappropriate activation of cellular Fc receptors. Aggregated autoantibodies and/or autoantibodies complexed with self antigens (immune complexes) bind to activating FcRs, thereby triggering the

pathogenic sequelae of numerous autoimmune diseases (which occur in part because of immunologically mediated inflammation against self tissues). Without intending to be bound by a particular mechanism of action, the CD16A binding proteins described herein interfere with and reduce the interaction of the autoimmune antibodies and Fc γ RIII receptors.

Examples of autoimmune diseases that can be treated include, without limitation, idiopathic thrombocytopenic purpura (ITP), rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), autoimmune hemolytic anemia (AHA), scleroderma, autoantibody triggered urticaria, pemphigus, vasculitic syndromes, systemic vasculitis, Goodpasture's syndrome, multiple sclerosis (MS), psoriatic arthritis, ankylosing spondylitis, Sjögren's syndrome, Reiter's syndrome, Kawasaki's disease, polymyositis and dermatomyositis. Other examples of diseases or conditions that can be treated according to the invention also include any diseases susceptible to treatment with intravenous immunoglobulin (IVIG) therapy (e.g., allergic asthma). Thus, the treatment of autoimmune diseases heretofore treated by IVIG therapy (in one embodiment, a condition other than ITP) is contemplated. While detailed understanding of the mechanism of action of IVIG has not been established, it is proposed that modulating the activity of cellular Fc γ Rs plays a role in its *in vivo* efficacy. The protective activity of IVIG may rely on the small percentage of dimeric or polymeric IgG present in the preparation. The specificity of the Fc γ RIII pathway in coupling cytotoxic and immune complex antibodies to effector responses and the ability to directly block this pathway with a mAb strongly suggests that an anti-Fc γ RIII antibody will have enhanced activity relative to IVIG.

A reduction in a deleterious immune response can be detected as a reduction in inflammation. Alternatively, a reduction in a deleterious immune response can be detected as a reduction in symptoms characteristic of the condition being treated (e.g., a reduction in symptoms exhibited by a subject suffering from an autoimmune condition), or by other criteria that will be easily recognized by physicians and experimentalists in the field of autoimmunity. It will be apparent that, in many cases, specific indicia of reduction will depend on the specific condition being treated. For example, for illustration and not limitation, a reduction in a deleterious immune response in a subject with ITP can be detected as a rise in platelet levels in the subject. Similarly, a reduction in a deleterious immune response in a subject with anemia can be detected as a rise in RBC levels in the subject. A clinician will recognize significant changes in platelet or RBC levels, or other responses following treatment.

The deleterious immune response is optionally due to idiopathic thrombocytopenic purpura resulting from the administration of an antiplatelet antibody, optionally murine monoclonal antibody 6A6, to a muFc γ RIII $^{-/-}$, huFc γ RIIA transgenic mouse.

In one aspect, the invention provides a method for treating an autoimmune disease, such as ITP, by administering a CD16A binding protein that is largely devoid of effector function. In an embodiment, the CD16A binding protein comprises Fc regions derived from human IgG. In an embodiment, the Fc regions are aglycosyl. In an embodiment, position 297 of each of the C_H2 domains is a residue other than asparagine or proline. In one aspect, the binding protein comprises a variable region sequence as described elsewhere herein. However, as discussed herein, the compositions and treatment methods of the invention are not limited to specific CD16A binding proteins derived from

murine mAb 3G8, but are applicable to CD16A binding proteins in general. In an embodiment, the CD16A binding protein is a tetrameric antibody protein having two light chains and two heavy chains.

In a related aspect, the invention provides methods of reducing a deleterious immune response in a mammal without significantly reducing neutrophil levels or inducing neutropenia (e.g., severe neutropenia or moderate neutropenia) by administering to the mammal a therapeutically effective amount of a pharmaceutical composition comprising a CD16A binding protein described herein. In an embodiment, the mammal is human. In an embodiment, the mammal is a nonhuman mammal (e.g., mouse) comprising one or more human transgenes.

For therapeutic applications, the binding proteins of the invention are formulated with a pharmaceutically acceptable excipient or carrier, e.g., an aqueous carrier such as water, buffered water, 0.4% saline, 0.3% glycine and the like, optionally including other substances to increase stability, shelf-life or to approximate physiological conditions (sodium acetate, sodium chloride, potassium chloride, calcium chloride, sodium lactate, histidine and arginine). For administration to an individual, the composition is preferably sterile, and free of pyrogens and other contaminants. The concentration of binding protein can vary widely, e.g., from less than about 0.01%, usually at least about 0.1% to as much as 5% by weight. Methods for preparing parentally administerable compositions are known or apparent to those skilled in the art and are described in more detail in, for example, Remington, *THE SCIENCE OF PRACTICE AND PHARMACY*, 20th Edition Mack Publishing Company, Easton, Pa., 2001). The pharmaceutical compositions of the invention are typically administered by a parenteral route, most typically intravenous, subcutaneous, intramuscular, but other routes of administration can be used (e.g., mucosal, epidermal, intraperitoneal, oral, intranasal, and intrapulmonary). Although not required, pharmaceutical compositions are preferably supplied in unit dosage form suitable for administration of a precise amount. In one embodiment, CD16A binding proteins can be administered in a form, formulation or apparatus for sustained release (e.g., release over a period of several weeks or months).

In one embodiment, polynucleotides encoding CD16A binding proteins (e.g., CD16A binding protein expression vectors) are administered to a patient. Following administration, the CD16A binding protein is expressed in the patient. Vectors useful in administration of CD16A binding proteins can be viral (e.g., derived from adenovirus) or nonviral. Usually the vector will comprise a promoter and, optionally, an enhancer that serve to drive transcription of a protein or proteins. Such therapeutic vectors can be introduced into cells or tissues in vivo, in vitro or ex vivo. For ex vivo therapy, vectors may be introduced into cells, e.g., stem cells, taken from the patient and clonally propagated for autologous transplant back into the same patient (see, e.g., U.S. Pat. Nos. 5,399,493 and 5,437,994).

The compositions can be administered for prophylactic and/or therapeutic treatments. In prophylactic applications, compositions are administered to a patient prior to an expected or potential deleterious immune response. For example, idiopathic thrombocytopenic purpura and systemic lupus erythematosus are conditions in which a deleterious immune response can be exacerbated by administration of certain medications. The CD16A binding compositions of the invention can be administered in anticipation of such medication-induced responses to reduce the magnitude of the response. In therapeutic applications, compositions are

administered to a patient already suffering from a deleterious immune response in an amount sufficient to at least partially ameliorate the condition and its complications. An amount adequate to accomplish this may be a "therapeutically effective amount" or "therapeutically effective dose." Amounts effective for these uses depend upon the severity of the condition and the general state of the patient's own immune system, but generally range from about 0.01 to about 100 mg of antibody per dose, with dosages of from 0.1 to 50 mg and 1 to 10 mg per patient being more commonly used. An "inflammation reducing amount" of the binding protein can also be administered to a mammal to reduce a deleterious immune response.

The administration of the CD16A binding proteins can be administered according to the judgement of the treating physician, e.g., daily, weekly, biweekly or at any other suitable interval, depending upon such factors, for example, as the nature of the ailment, the condition of the patient and half-life of the binding protein.

CD16A binding proteins can be administered in combination with other treatments directed to alleviation of the deleterious immune response or its symptoms or sequelae. Thus, the binding proteins can be administered as part of a therapeutic regimen that includes co-administration of another agent or agents, e.g., a chemotherapeutic agent such as a non-steroidal anti-inflammatory drug (e.g., aspirin, ibuprofen), steroids (e.g., a corticosteroid, prednisone), immunosuppressants (e.g., cyclosporin A, methotrexate, cytoxan), and antibodies (e.g., in conjunction with IVIG).

6. Increasing the Therapeutic Efficacy of a CD16A Binding Protein

In a related aspect, the invention provides a method for increasing the therapeutic efficacy of a CD16A binding protein comprising one or more Fc domains (e.g., anti-CD16A antibodies comprising two Fc domains) by modifying the protein so that it has Fc region(s) with reduced binding to at least one Fc effector ligand compared to the original (i.e., unmodified) Fc region. For example, the Fc region can be modified so that the Fc region is not glycosylated. As described above, modification of the Fc region can be accomplished in several ways (e.g., by genetic mutation, by choice of expression system to change the Fc glycosylation pattern, and the like). In one embodiment, the Fc effector ligand is FcγRIII. In one embodiment, the Fc effector ligand is the C1q component of complement. As used in this context, a subject CD16A binding protein has increased "therapeutic efficacy" compared to a reference binding protein that induces neutropenia when administered if the subject CD16A binding protein does not induce neutropenia (or results in less severe neutropenia). For example, a CD16A binding protein that reduces the severity of a deleterious immune response (e.g., ITP or experimentally induced ITP in a mammal) and reduces neutrophil levels in the animal by x % has greater "therapeutic efficacy" than a CD16A binding protein that reduces the severity of a deleterious immune response and reduces neutrophil levels in the animal by y %, if y is greater than x, e.g. two-fold greater. In one embodiment, the protein is modified by mutation such that the modified protein is aglycosylated.

For example, the invention provides methods for producing a modified CD16 binding protein comprising a modified immunoglobulin heavy chain, the modified CD16 binding protein having greater therapeutic efficacy than a parent CD16 binding protein comprising a parent immunoglobulin heavy chain, by (i) introducing at least one mutation into a parent polynucleotide that encodes the parent immunoglo-

bulin heavy chain to produce a modified polynucleotide that encodes the modified immunoglobulin heavy chain, the mutation introducing into the modified immunoglobulin heavy chain an amino acid substitution that changes, reduces or eliminates glycosylation in the C_H2 domain of the parent immunoglobulin heavy chain; and (ii) expressing the modified polynucleotide in a cell as the modified immunoglobulin heavy chain so as to produce the modified CD16 binding protein heavy chain. Optionally, the heavy chain is produced under conditions of co-expression with a light chain to produce a tetrameric antibody.

7. EXAMPLES

Example 1

Mouse 3G8 VH and VL and Chimeric Molecules
Generated therefrom

A) Mouse 3G8 VH and VL

The cDNA encoding the mouse 3G8 antibody light chain was cloned. The sequence of the 3G8 antibody heavy chain was provided by Dr. Jeffry Ravetch. The amino acid sequences of the 3G8 V_H and V_L are provided in Tables 1 and 3, infra. Nucleic acid sequences encoding the variable regions are:

{3G8VH}
CAGGTTACTCTGAAAGAGTCTGGCCCTGGGATATTGCAGCCCTCCAGAC
CCTCAGTCTGACTTGTCTTTCTCTGGGTTTTCACTGAGGACTTCTGGTA
TGGGTGTAGGCTGGATTTCGTCAGCCTTCAGGGAAGGGTCTAGAGTGGCTG
GCACACATTTGGTGGGATGATGACAAGCGCTATAATCCAGCCCTGAAGAG

-continued

CCGACTGACAATCTCCAAGGATACCTCCAGCAACCAGGTATTCCTCAAAA
5 TCGCCAGTGTGGACACTGCAGATACTGCCACATACTACTGTGCTCAAATA
AACCCCGCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTCTC
TGCA
10 {3G8VL}
GACACTGTGCTGACCCAATCTCCAGCTTCTTTGGCTGTGTCTCTAGGGCA
GAGGGCCACCATCTCCTGCAAGGCCAGCCAAAGTGTTGATTTTGATGGTG
15 ATAGTTTTATGAACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTC
CTCATCTATACTACATCCAATCTAGAATCTGGGATCCCAGCCAGGTTTAG
TGCCAGTGGGTCTGGGACAGACTTCACCCTCAACATCCATCCTGTGGAGG
20 AGGAGGATACTGCAACCTATTACTGTGTCAGCAAAGTAATGAGGATCCGTAC
ACGTTTCGGAGGGGGGACCAAGCTGGAAATAAAA

B) Chimeric Heavy Chain

To create a chimeric gene coding for expression of the mouse 3G8 VH fused to a human constant domain, the nucleic acid encoding the 3G8 V_H was fused to sequences encoding a signal peptide (see Orlandi et al., 1989, *Proc. Natl. Acad. Sci. U.S.A* 86:3833-37; in lower case underline
25 below) and a human C_γ1 constant region (in lower case below) using standard techniques (including overlapping PCR amplification). To facilitate cloning, a SacI site was introduced, resulting in a single residue change in VH FR4 (ala→ser). This change in FR4 does not affect binding to
35 CD16. The resulting nucleic acid had the sequence shown below. The region encoding the V_H domain is in upper case.

{ch3G8VH}
gctagcgttttaaacttaagcttggtgactagtgagatcacagttctctctacagtta SEQ ID NO:5
ctgagcacacaggacctcaccatgggatggagctgtatcatcctcttcttggtagca
acagctacaggtaaggggctcacagtagcaggcttgaggctctggacatatatatggg
tgacaatgacatccactttgcctttctctccacaggtgtccactccCAGGTTACCT
GAAAGAGTCTGGCCCTGGGATATTGCAGCCCTCCAGACCTCAGTCTGACTTGTTT
TTTCTCTGGGTTTTCACTGAGGACTTCTGGTATGGGTGTAGGCTGGATTTCGTCAGCC
TTCAGGGAAGGGTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTA
TAATCCAGCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCAGCAACCAGGT
ATTCTCAAAATCGCCACTGTGGACACTGCAGATACTGCCACATACTACTGTGCTCA
AATAAACCCCGCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTGAGCTC
Agcctocaccaagggcccatcggtcttccccctggcaccctcctccaagagcacctc
tgggggcacagcgccctgggctgctggtcaaggactacttccccgaaccggtgac
gggtgctgtggaactcaggcgccctgaccagcggtgcacaccttccccggctgtcct
acagtccctcaggactctactccctcagcagcgtggtgaccgtgcctccagcagctt
gggcacccagacctacatctgcaacgtgaatcacaagcccagcaacaccaaggtgga
caagagagttgagcccaaatcttggtgacaaaactcacacatgccaccgtgcccagc
acctgaactcctggggggaccgtcagtccttctcttccccccaaaaccaaggacac

-continued

cctcatgatctcccgaccctgaggtcacatgctggtggtggacgtgagccacga
agaccctgaggtcaagttcaactggtacgtggacggcgtggaggtgcataaatgccaa
gacaaagccgcgggaggagcagtacaacagcacgtaccgtgtggtcagcgtcctcac
cgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaa
agccctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgaga
accacaggtgtacaccctgcccccatcccggtgagctgaccaagaaccaggtcag
cctgacctgctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagag
caatgggcagccggagaacaactacaagaccacgcctcccgctgctggactccgacgg
ctccttcttctctacagcaagctcacgtggacaagagcaggtggcagcaggggaa
cgtcttctcatgctccgtgatgcatgaggtctgcacaaccactacacgcagaagag
cctctccctgtctccgggtaaatgagtgcggccgcgaattc

This construct was inserted into the pCI-Neo (Promega Biotech) at the NheI-EcoRI sites of the polylinker for use for expression of the chimeric heavy chain in cells.

-continued

C) Chimeric Light Chain

To create a chimeric gene coding for the mouse 3G8 V_L fused to a human constant domain, this 3G8 V_L segment was fused to a signal sequence (as for the V_H above; (lower case underlined) and a human Ck constant region (lower case) cDNA using standard techniques, resulting in a nucleic acid with the sequence shown below:

ch3G8VL

SEQ ID NO: 6

gctagctgagatcacagttctctctacagttactgagcacacaggacctc
accatgggatggagctgtatcatcctcttcttggtagcaacagctacagg
taaggggctcacagtagcaggcttgaggtctggacatatatatgggtgac
aatgacatccactttgcctttctctccacaggtgtccactccGACACTGT
GCTGACCCAATCTCCAGCTTCTTTGGCTGTGTCTCTAGGGCAGAGGGCCA
CCATCTCCTGCAAGGCCAGCCAAAGTGTTGATTTTGATGGTGATAGTTTT
ATGAACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTCCTCATCTA
TACTACATCCAATCTAGAATCTGGGATCCCAGCCAGGTTTAGTGCCAGTG
GGTCTGGGACAGACTTCACCTCAACATCCATCCTGTGGAGGAGGAGGAT
ACTGCAACCTATTACTGTCAGCAAAGTAATGAGGATCCGTACACGTTCCG
AGGGGGGACCAAGCTTGAGATCAAAcgaactgtggctgcaccatcggtct
tcatcttcccgccatctgatgagcagttgaaatctggaactgcctctgtt
gtgtgcctgctgaataacttctatcccagagaggccaaagtacagtggaa

25 ggtggataacgcctccaatcgggtaactcccaggagagtgtcacagagc
aggacagcaaggacagcacctacagcctcagcagcaccctgacgtgagc
30 aaagcagactacgagaaacacaaagtctacgcctgcgaagtca
gggcctgagctcgcccgtcacaaagagcttcaacaggggagagtgttagt
35 tctagagtcgaactctagaggatccccgggtaccgagctcgaattc

This construct was inserted into pCI-Neo (Promega Biotech) at the NheI-EcoRI sites of the polylinker for use for expression of the chimeric light chain in cells.

D) Expression


The ch3G8VH and ch3G8VL chimeric proteins described above can be co-expressed to form a chimeric antibody, referred to as ch3G8. The chimeric antibody ch3G8 can be expressed either in a myeloma or in other mammalian cells (e.g., CHO, HEK-293). An example of a procedure for expression of CD16A binding proteins such as ch3G8 and variants is provided in Example 4, infra.

Example 2

Humanized Anti-CD16A Binding Proteins

A) Humanized Heavy Chain

CDR encoding sequences from the mouse 3G8 V_H clone were fused to framework sequences derived from the human germline VH sequence VH2-70 to create a polynucleotide encoding a V_H designated Hu3G8VH. The polynucleotide was generated by an overlapping PCR procedure. In a first step, using the primers and strategy shown below and the mouse 3G8 V_H polynucleotide (SEQ ID NO: 1) as template.



The resulting fragment was digested with EcoRI and SacI and cloned into pUC18. After sequencing, one plasmid was selected for a final round of overlapping PCR to correct a deletion which occurred during the second PCR step. The resulting polynucleotide had the sequence:

{hu3G8VH}

SEQ ID NO:16
CAGGTTACCCTGAGAGAGTCTGGCCCTGCGCTGGTGAAGCCACACAGAC
CCTCACACTGACTTGTACCTTCTCTGGGTTTTCACTGAGCACTTCTGGTA
TGGGTGTAGGCTGGATTCTGTCAGCCTCCCGGAAGGCTCTAGAGTGGCTG
GCACACATTTGGTGGGATGATGACAAGCGCTATAATCCAGCCCTGAAGAG
CCGACTGACAATCTCCAAGGATACCTCCAAAACCAGGTAGTCCTCACAA
TGACCAACATGGACCCTGTGGATACTGCCACATACTACTGTGCTCGGATA
AACCCCGCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTGAG
CTCA

The Hu3G8VH sequence was then combined with segments coding for a secretion signal sequence (as described above; lower case underline) and cDNA for the human Cyl

{hu3G8VH-1}

SEQ ID NO:17
gctagcgtttaaacttaagcttggtgactagtgagatcacagttctctct
acagttactgagcacacaggacctcaccatgggatggagctgtatcatcc
tcttcttggttagcaacagctacaggtaaggggctcacagtagcaggcttg
aggctctggacatatatatgggtgacaatgacatccactttgctttctct
ccacaggtgtccactccCAGGTTACCTGAGAGAGTCTGGCCCTGCGCTG
GTGAAGCCACACAGACCCTCACACTGACTTGTACCTTCTCTGCGTTTTTC
ACTGAGCACTTCTGGTATGGGTGTAGGCTGGATTCTGTCAGCCTCCCGGA

-continued


AGGCTCTAGAGTGGCTGGCACACATTTCTGGGATGATGACAAGCGCTAT
AATCCAGCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCAAAAA
CCAGGTAGTCCTCACAAATGACCAACATGGACCCTGTGGATACTGCCACAT
ACTACTGTGCTCGGATAAACCCCGCCTGGTTTGCTTACTGGGGCCAAGGG
ACTCTGGTCACTGTGAGCTCAgcctccaccaaggggccatcggtcttccc
cctggcaccctcctccaagagcacctctgggggcacagcgccctgggct
gcctggtcaaggactacttccccgaaccggtgacggtgtcgtggaactca
ggcgccctgaccagcggtgcacaccttccgggtgtcctacagtctctc
aggactctactccctcagcagcgtggtgacggtgccttccagcagcttgg
gcaccagacctacatctgcaacgtgaatcacaagcccagcaacaccaag
gtggacaagagagttgagcccaaactctgtgacaaaactcacacatgccc
accgtgcccagcacctgaactcctgggggaccgtcagtccttctcttcc
ccccaaaacccaaggacaccctcatgatctcccggaaccctgaggtcaca
tgcggtggtggtgacgtgagccacgaagaccctgaggtcaagttcaactg
gtacgtggacggcgtggaggtgcataatgccaagacaaagccgcgggagg
agcagtaaacagcacgtaccgtgtggtcagcgtcctcaccgtcctgcac
caggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagc
cctcccagcccccatcgagaaaaccatctccaaagccaaagggcagcccc
gagaaccacaggtgtacaccctgcccccatcccggtatgagctgaccaag
aaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacat
cgccgtggagtgggagagcaatgggcagccggagaaactacaagacca
cgctcccgtgctggactccgacggctccttcttctctacagcaagctc
accgtggacaagagcaggtggcagcaggggaacgtcctctcatgctccgt

-continued
gatgcatgaggctctgcacaaccactacacgcagaagagcctctccctgt
ctccgggtaaatgagtgcggccgcgaattc

For expression in mammalian cells (HEK-293), the Hu3G8VH-1 sequence was cloned into the pCI-Neo polylinker at the NheI-EcoRI sites, following intervening cloning into pUC and pCDNA3.1.

B) Humanized Light Chain

CDR encoding sequences from the mouse 3G8 V_L clone were fused to framework sequences derived from the human B3 germline V-κ gene. The polynucleotide was generated by an overlapping PCR procedure using the primers and strategy shown below and the mouse 3G8 V_L polynucleotide (SEQ ID NO: 2) as template.



The resulting polynucleotide had the sequence

{hu3G8VL}
SEQ ID NO:25
GACACTGTGCTGACCCAATCTCCAGCTTCTTTGGCTGTGTCTCTAGGGCA
GAGGGCCACCATCTCCTGCAAGGCCAGCCAAAGTGTGATTTTGATGGTC
ATAGTTTTATGAACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTC
CTCATCTATACTACATCCAATCTAGAATCTGGGATCCCAGCCAGCTTTAG
TGCCAGTGGGTCTGGGACAGACTTCACCCTCAACATCCATCCTGTGGAGG
AGGAGGATACTGCAACCTATTACTGTCTAGCAAAGTAATGAGGATCCGTAC
ACGTTCCGGAGGGGGGACCAAGCTTGAGATCAAA

The Hu3G8 VL gene segment was combined with a signal sequence (as described above, lower case, underline) and a human C-κ constant region (lower case) cDNA using standard techniques resulting in a product with the sequence below:

{hu3G8VL-1}
SEQ ID NO:26
gctagctgagatcacagttctctctacagttactgagcacacaggacctc
accatgggatggagctgtatcatcctcttcttggtagcaacagctacagg

-continued
taaggggctcacagtagcaggcttgaggctctggacatatatatgggtgac
5 aatgacatccactttgcctttctctccacaggtgtccactccGACACTGT
GCTGACCCAATCTCCAGCTTCTTTGGCTGTGTCTCTAGGCCAGAGGGCCA
CCATCTCCTGCAAGGCCAGCCAAAGTGTGATTTTGATGCTGATAGTTTT
10 ATGAACTGGTACCAACAGAAACCAGGACAGCCACCCAAACTCCTCATCTA
TACTACATCCAATCTAGAATCTGGGATCCCAGCCAGGTTTAGTGCCAGTG
GGTCTGGGACAGACTTCACCCTCAACATCCATCCTGTGGAGCAGGAGGAT
15 ACTGCAACCTATTACTGTCTAGCAAAGTAATGAGGATCCGTACACGTTCCG

-continued
AGGGGGGACCAAGCTTGAGATCAAAcgaactgtggctgcaccatcggtct
tcatcttcccgccatctgatgagcagttgaaatctggaactgcctctgtt
45 gtgtgcctgctgaataaacttctatcccagagaggccaaagtacagtggaa
gggtggataacgccctccaatcgggtaactcccaggagagtgtcacagagc
aggacagcaaggacagcacctacagcctcagcagcacctgacgctgagc
50 aaagcagactacgagaaacacaaagtctacgcctgccaagtaccccatca
gggcctgagctcgcccgtcacaaagagcttcaacaggggagagtgttagt
tctagagtcgactctagaggatccccgggtaccgagctcgaattc

55 This construct was inserted into pCI-Neo for expression in mammalian cells.

Example 3

60 Variant CD16A Binding Proteins

Additional expression constructs were made in which sequence changes were introduced in the V_L or V_H domains by site directed mutagenesis. A typical mutagenesis reaction
65 contained 10 ng plasmid DNA (isolated from a methylation competent strain of *E. coli*), 125 ng each of a forward and reverse primer, each containing the mutation of interest,

37

reaction buffer, and dNTPs in 0.05 ml volume. 2.5 units of PfuTURBO® DNA polymerase (Stratagene) was added and the reaction was subjected to 15 cycles of 95°, 30 sec; 55°, 1 min; 68°, 12 min. The product of the PCR was then digested with DpnI endonuclease and the restricted DNA was used to transform *E. coli* strain XL-10 GOLD® ultra-competent cells. Because DpnI only digests methylated DNA it will digest the parental, non-mutated, plasmid leaving the newly synthesized non-methylated product, containing the mutation of interest, as the predominant species.

The sequences of the variant V_H domains are shown in Table 3. The sequences of the variant V_L domains are shown in Table 4.

Example 4

Expression in Mammalian Cells

Various combinations of heavy and light chain expression plasmids (e.g., comprising the chimeric, humanized and variant V_L and V_H domains fused to human C γ 1 and C κ constant domains as described above) were co-transfected into HEK-293 cells for transient expression of recombinant tetrameric antibodies (i.e., comprising 2 heavy chains and 2 light chains), sometimes referred to herein as “recombinant antibodies.” Transfection was carried out using LIPO-FECTAMINE® 2000 (Invitrogen) in 6 well plates according to the manufacturer’s instructions.

Recombinant antibodies were prepared by cotransfection of a heavy chain expression plasmid (i.e., encoding a heavy chain comprising a V_H and constant domains) and light chain expression plasmids (i.e., encoding a light chain comprising a V_L and constant domains) together into HEK-293 cells for transient expression of recombinant antibodies.

Hu3G8VH variants listed in Table 3 were coexpressed with the hu3G8VL-1 light chain. For reference, most assays included (i) recombinant antibodies produced by coexpression of ch3G8VH and ch3G8VL (“ch3G8VH/ch3G8VL”) and (ii) recombinant antibodies produced by coexpression of hu3G8VH-1 and hu3G8VL-1 (“hu3G8VH-1/hu3G8VL-1”).

Hu3G8VL variants listed in Table 4 were coexpressed with the ch3G8VH heavy chain. For reference, most assays included (i) recombinant antibodies produced by coexpression of ch3G8VH and ch3G8VL (“ch3G8VH/ch3G8VL”) and (ii) recombinant antibodies produced by coexpression of ch3G8VH and hu3G8VL-1 (“ch3G8VH/hu3G8VL-1”).

After three days, the levels of recombinant antibodies in the conditioned media were determined by ELISA, and the recombinant antibodies were analyzed by ELISA for binding to captured sCD16A as described in Examples 5. Selected antibodies were assayed for cell binding to cells expressing the extracellular domain of CD16A, as shown in Example 6.

Example 5

ELISA Determination of Binding to CD16A

Sandwich ELISA was performed to detect binding of antibodies to a soluble form of CD16A.

Soluble Human CD16A

A soluble form of human CD16A was expressed from HEK-293 cells using a pcDNA3.1-derived expression vector containing the CD16A gene truncated just prior to the transmembrane region. To create the vector, cDNA encoding CD16A was amplified using the primers 3A_{left} [gttgatcctc-
caactgctctgctacttctagttt] (SEQ ID NO:27) and 3A_{right} [gaaaagcttaagaatgatgagatggtgacact] (SEQ ID NO:28)

38

digested with BamHI and HindIII, and cloned into the vector pcDNA3.1 (Novagen) at the Bam/HindIII site of the polylinker. The construct was used to transiently transfect HEK-293 cells. For some assays, the secreted product was purified from conditioned medium using affinity chromatography on a human IgG Sepharose column. In some assays, the amount of sCD16A in conditioned medium was quantitated and unpurified sCD16A was used. Purification was not required since the ELISA capture antibody (LNK16 mAb) specifically bound the antigen, allowing removal of contaminants in washing steps.

The amino acid sequence of the sCD16 construct is shown below. (The signal sequence, underlined, is cleaved off during expression. Note the last seven residues are derived from the vector pcDNA3.1 rather than from the CD16A gene):

(SEQ ID NO:29)

MWOLLPTALLLLVSAGMRTEDLPKAVVFLEPQWYRVLEKDSVTLKCQGA
YSPEDNSTQWFHNESLISSQASSYFIDAATVDDSGEYRCQTNLSTLSDPV
QLEVHIGWLLLQAPRWVFKEEDPIHLRCHSWKNTALHKVITYLQNGKGRKY
FHHNSDFYIPKATLKDSGSYFCRGLFGSKNVSSSETVNITITQGLAVSTIS
SFFKLAAARV

ELISA Format

Plates were first coated with 100 ng/well of the anti-CD16 mAb LNK-16 (Advanced ImmunoChemical, Long Beach Calif.; see 5th Human Lymphocyte Differentiation Antigens Workshop) in carbonate buffer at room temperature for 2 hrs. Any anti-sCD16A antibody that does not block binding by 3G8 can be used. After blocking for 30 minutes with PBS-T-BSA, sCD16A conditioned medium was added at a dilution of 1/10 and incubated at room temperature for 16 hrs. Alternatively, when purified sCD16 was used, it was diluted to a concentration of 50 ng/ml in PBS-T-BSA. 0.05 ml was added to each well and incubated for at least 2 hrs at room temperature.

The plate was washed and dilutions of recombinant antibodies starting at 0.5 μ g/ml in PBS-T-BSA were then added and incubated for 1 hr at room temp. Binding of recombinant antibodies to the captured sCD16A was then measured using an anti-human IgG-HRP conjugate and TMB substrate. After stopping color development using dilute sulfuric acid, the plate was read at 450 nM.

Results of Binding Assays

This example shows that the binding properties of humanized anti-CD16A antibodies for binding to CD16A are the same or similar to the properties of the chimeric 3G8 antibody.

Based on the comparative binding studies, the recombinant antibodies were classified as binding with high, intermediate, or low affinity. Antibodies with high and intermediate binding affinity are discussed above in section 4. The recombinant antibodies with a V_H domain of hu3G8VH-9, 10, 11, 13, 15, 21, 38, 39, or 41 showed little or no binding to sCD16A. From these data it appears certain substitutions (or combinations of substitutions) are generally detrimental to binding. For example, substitution of tyrosine or aspartic acid at V_H position 52 (i.e., 52Y and 52D) or threonine at position 94 (94T) are detrimental to binding. Similarly, the combination of leucine at position 50 with aspartic acid at position 54 (50L+54N) is detrimental to binding, as is the combination of arginine at 94 and aspartic acid at 101 (94R+101D). However, aspartic acid at 101 is tolerated

when position 94 is glutamine, lysine, histidine or alanine (but not arginine). Further 34V+94R+101D has intermediate activity. This indicates a relationship between positions 34, 94 and 101 in maintaining high affinity binding, and suggests that 34V may be an especially important residue. Likewise, recombinant antibodies with a V_L domain of hu3G8VL-6, 7, 8, 9, 11, 12, 13, and 14 showed little or no binding to sCD16A. From these data it appears certain substitutions (or combinations of substitutions) are generally detrimental to binding. For example, substitution of alanine at position 34 (34A) or tyrosine at position 92 (92Y) is generally detrimental to binding.

Results of an exemplary binding assay are shown in FIG. 1.

Example 6

Antibody Binding to Cells Expressing CD16A

The ability of selected humanized antibodies to bind to CD16A expressed by CHO-K1 cells as assayed by direct binding competition assays.

CHO-K1 cells expressing extracellular domain of FcRIIIa fused to the transmembrane and intracellular domain of FcRIIb were used for cell binding assays. Cells were plated at 40,000 cells per well in 96 well flat bottom tissue culture plates (FALCON® MICROTEST™ Tissue Culture plate, 96 well) and incubated at 37° C. CO₂ incubator for approximately 24 hr. The plate was then gently washed three times with 25 mM Hepes, 75 uM EDTA, 11.5 mM KCl, 115 mM NaCl, 6 mM MgSO₄, 1.8 mM CaCl₂, 0.25% BSA (binding buffer).

For indirect binding assays, 100 µl of a serial dilution of anti-CD16 Mab (final concentration: 1 ug/ml, 0.5, 0.25, 0.125, 0.0625, 0.03, 0.015, 0 ug/ml) was then added to wells in binding buffer. The plate was then incubated at 23° C. for 1 hr and washed three times with binding buffer. 50 µl/well of Europium (EU)-labeled-antihuman-IgG (100 ng/ml) was then added to each well and the plate was incubated at 23° C. for 30 minutes then washed three times with binding buffer. Finally, 100 µl DELFIA® enhancement solution, an acidic chelating detergent solution intended for use in the quantitative determination of Eu³⁺/Sm³⁺ (PerkinElmer/Wallac) was added. After incubating with shaking for 15 minutes, the plate was read for time resolved fluorescence (excitation 340 nm; emission 615 nm) in a VICTOR² instrument (PerkinElmer/Wallac). The results of the assay are shown in FIG. 2.

The CHO-K1 cells described above were also used in competition assays. After washing with binding buffer as described above, varying amounts of purified unlabeled Mab (1.2-75 nM final concentration) were mixed with a fixed concentration of Eu-Ch3G8-N297Q (final concentration 2.5 nM). The plate was then incubated at 23° C. for 1 hr and washed three times with binding buffer. 100 µl DELFIA® enhancement solution (PerkinElmer/Wallac) was added and after incubating with shaking for 15 minutes, the plate was read for time resolved fluorescence (excitation 340 nm; emission 615 nm) in a VICTOR² instrument (PerkinElmer/Wallac). The results of the assay are shown in FIG. 3.

These assays demonstrate that the humanized anti CD16A monoclonal antibodies bind with high affinity to CD16A on the surface of transfected cells. Hu3G8-22.1-N297Q binds to CD16A bearing cells with higher affinity than Ch3G8-N297Q.

Example 7

Inhibition of Binding of sCD16A to Immune Complexes

Assay of 4-4-20 Binding to FITC-BSA

The binding of ch4-4-20 or ch4-4-20 (D265A) to FITC-BSA was assessed by ELISA. (Ch4-4-20 is identical to Ch3G8 except that it contains the respective VH and VL regions of 4-4-20 instead of those of 3G8. Thus it retains high affinity and specificity for the hapten fluorescein. 4-4-20 is described in Bedzyk et al., 1989, *J Biol Chem* 264:1565-9.) FITC-BSA (1 ug/ml-50 ng/well) was coated onto Nunc maxisorb immunoplates in carbonate buffer and allowed to bind for approximately 16 hr. Following blocking with BSA, dilutions of ch4-4-20 were added to the wells and allowed to bind for 1 hr at RT. After washing out unbound Mab, HRP-conjugated goat anti-human Ig secondary was added. One hour later the secondary antibody was removed, washed and developed with TMB substrate. Following addition of an acidic stop solution the plate was read at 450 nm. Both ch4-4-20 and ch4-4-20(D265A) bound to the FITC-BSA with high affinity (data not shown).

Assay of sFcR Binding to ch4-4-20/FITC-BSA Immune Complexes

The same format was used to assay binding of sFcRs to immune complexes (IC) formed on the ELISA plate between ch4-4-20 and FITC-BSA. In this case we have used either biotinylated sFcR or biotinylated anti-human G2 Mab as a secondary reagent, followed by streptavidin-HRP detection.

Inhibition of sFcR Binding to IC by Murine, Chimeric and Humanized 3G8

The concentrations of ch4-4-20 and sFcR were fixed to give approximately 90 percent maximal signal in the assay. sCD16A was premixed with serial dilutions of murine, chimeric or humanized 3G8 and incubated for one hour prior to adding to the plate containing the immune complex. Serial dilutions of humanized or chimeric 3G8 were incubated with sCD16A-G2-biotin for one hour. The mixtures were then added to ELISA wells containing an immune complex between a human IgG1 chimeric form of the anti-fluorescein Mab 4-4-20 and FITC-BSA. After one hour, binding of the soluble receptor to the IC was detected using streptavidin-HRP conjugate and TMB development. The results are shown in FIG. 4. This assay indicates that humanized anti-CD16A antibodies are potent inhibitors of CD16A binding to IgG in immune complexes.

Example 8

Analysis of Anti-CD16A Monoclonal Antibody Panel

A panel of hybridomas was generated following immunizing and boosting mice with sCD16A using standard methods. Eight 96-well plates were screened by ELISA for binding activity on plates coated directly with sCD16A. Ninety-three of these gave a positive signal and were expanded further. Of these, 37 were positive for binding to human blood cells by FACS. These supernatants were then analyzed for their ability to block the interaction of CD16A with immune complexes and for the similarity of the binding site (epitope) to that of 3G8. Assays included capture ELISA using chimeric 3G8 down and inhibition of immune complex binding to sRIIIa-Ig. Based on these assays antibodies

with binding and inhibitory properties similar to 3G8 were isolated, as well as Mabs with binding and/or inhibitory properties distinct from 3G8.

DJ130c (DAKO) and 3G8 were used as controls in the assays. Mab DJ130c is a commercially available Mab which binds CD16 at an epitope distinct from 3G8 (Tamm and Schmidt). This Mab does not block FcRIIIa-immune complex binding (Tamm and Schmidt). In an ELISA-based inhibition assay, DJ130c enhances rather than inhibits binding.

The data indicate that the panel contains antibodies which bind to the same epitope as Ch3G8 and block sCD16A binding to immune complexes. The panel of Mabs also contains antibodies which do not bind to the same epitope as Ch3G8. Most of these latter antibodies do not block the interaction of sCD16a with IgG in immune complexes.

TABLE

Effect on sCD16a Binding to Immune Complexes				
Assay	Result	Inhibition	Enhancement	No Effect
Binding to sCD16	Positive	2	5 (+DJ-130c)	17
Captured by Ch3G8	Negative	11 (+3G8)	0	2

Example 9

Induction of Platelet Depletion In Vivo

The in vivo activity of a CD16A binding protein for blocking human Fc-FcγRIII interactions induced by autoantibodies can be evaluated using animal models of autoimmune diseases. One suitable model is the "passive mouse model" of ITP and the anti-platelet mAb 6A6 (see, Oyaizu et al., 1988, *J Exp. Med.* 167:2017-22; Mizutani et al., 1993, *Blood* 82:837-44). 6A6 is an IgG2a isotype mAb derived from a NZW×BSXB F1 individual. Administration of 6A6 depletes platelets in muFcγRIII^{-/-}, huFcγRIIIA transgenic mice but not in muFcγRIII^{-/-} mice without the human transgene. See Samuelsson et al., 2001, *Science* 291:484-86. Other anti-platelet monoclonal antibodies can be used in place of 6A6 in the model. Alternatively, a polyclonal anti-platelet antibody can be used.

CD16A binding proteins that confer the greatest degree of protection from platelet depletion can be identified by administering CD16A binding proteins to a muFcγRIII^{-/-}, huFcγRIIIA transgenic mouse and measuring any reduction in mAb 6A6 induced platelet depletion.

A related assay can be carried out using a chimeric human IgG₁κ chimeric derivative of 6A6 in place of the mouse mAb in the protocol provided above, so that the depleting mAb had a human isotype. To conduct this assay, a chimeric 6A6 monoclonal antibody (ch6A6) was prepared by fusing the cDNA segments encoding the murine anti-platelet monoclonal antibody 6A6 V_H and V_L regions to the human Cγ1 and Cκ cDNA segments, respectively. The resulting genes were co-expressed in 293 cells and chimeric 6A6 was purified by protein A affinity chromatography followed by size exclusion chromatography.

To demonstrate that the chimeric 6A6 antibody induces platelet depletion, ch6A6 was administered to muFcγRIII^{-/-}, huFcγRIIIA transgenic mice. The ch6A6 was administered to each animal either i.v. or intraperitoneally (i.p.) (0.1 μg/g).

Animals were bled 2 hrs, 5 hrs, 24 hrs and 48 hrs after administration of ch6A6, and plasma platelet counts were determined using a Z2™ COULTER COUNTER® particle counter and size analyzer equipped with a 70 μm aperture. Particles between 1.5 and 4 μm in size (corresponding to platelets) were counted and the data were analyzed by plotting the platelet count versus time for each concentration.

Two hours after injection of 0.1 μg/g ch6A6 i.p., approximately 75% of the platelets were depleted. The number of platelets remained low for 5 hours after ch6A6 injection then progressively increased to return to normal 72 hours after ch6A6 injection.

Two hours after injection of 0.1 μg/g ch6A6 i.v., approximately 60% of the platelets were depleted. The number of platelets remained low for 6 hours after ch6A6 injection then progressively increased to return to normal 48 hours after ch6A6 injection.

Example 10

Analysis of the Ability of CD16 Binding Antibodies to Protect Mice from Platelet Depletion

The ability of CD16A binding proteins to reduce platelet depletion in experimental ITP can be assayed as described below. CD16A binding proteins were administered intravenously (i.v.) to groups of muFcγRIII^{-/-}, huFcγRIIIA transgenic mice at concentrations of 0.5, 1, 2 or 5 μg/g in phosphate buffered saline (PBS). Controls were PBS alone or an irrelevant human IgG1 (negative control) or human intravenous immunoglobulin (IVIG; positive control). One hour after administration of the CD16A binding protein or control, ITP was induced by administering 0.1 μg/g ch6A6 to each animal either intravenously or intraperitoneally. Animals were bled 2 hrs, 5 hrs, 24 hrs and 48 hrs after administration of ch6A6. Plasma platelet counts were determined using the Z2™ COULTER COUNTER® particle counter and size analyzer as described above and the data were analyzed by plotting the platelet count versus time for each concentration of administered binding protein.

When muFcγRIII^{-/-}, huFcγRIIIA transgenic mice were injected with murine 3G8 (0.5 μg/g) one hour before i.p. injection of ch6A6, 33% of the platelets were depleted at the 2 hours time point (FIG. 5). The number of platelets then progressively increased to return to normal 24 hours after ch6A6 injection. When muFcγRIII^{-/-}, huFcγRIIIA transgenic mice were injected with murine 3G8 (0.5 μg/g) one hour before i.v. injection of ch6A6, 30% of the platelets were depleted at the 2 hours time point (FIG. 6). The number of platelets then rapidly increased to return to normal 5 hours after ch6A6 injection.

These results were similar to the protection seen when human IVIG is administered. When muFcγRIII^{-/-}, huFcγRIIIA transgenic mice were injected with human IVIG (1 mg/g) one hour before i.p. injection of ch6A6, 33% of the platelets were depleted at the 2 hours time point (FIG. 5). The number of platelets then progressively increased to return to normal 24 hours after ch6A6 injection. When muFcγRIII^{-/-}, huFcγRIIIA transgenic mice were injected with human IVIG (1 mg/g) one hour before i.v. injection of ch6A6, 20% of the platelets were depleted at the 2 hours time point (FIG. 6). The number of platelets then rapidly increased to return to normal 5 hours after ch6A6 injection.

The results shown in FIGS. 5 and 6 show that m3G8 protects mice from ch6A6-mediated platelet depletion, and that the level of protection was similar to the protection conferred by IVIG.

Preparations of recombinant mouse 3G8 produced in HEK-293 cells, chimeric 3G8 with human IgG1 or IgG2 constant domains (ch3G8- γ 1 produced in HEK-293 and CHO-K1 cells, and ch3G8- γ 2 produced in HEK-293 cells), and a ch3G8- γ 1 variant (ch3G8- γ 1 D265A) did not provide significant protection in this experiment. When muFc γ RIII^{-/-}, huFc γ RIIIA transgenic mice were injected with ch3G8 γ 1 or γ 2 (0.5 μ g/g) one hour before i.p. injection of 6A6, approximately 60% of the platelets were depleted at the 5 hour time point (FIG. 7). The number of platelets then progressively returned to normal. Although depletion was not as severe as in mice that received no anti-CD16A binding protein, these chimeric antibodies provided significantly less protection, if any, than murine 3G8. A ch3G8 variant in which aspartic acid 265 was changed to alanine showed similar results. Interestingly, as is shown in Example 11, modification of the ch3G8 to produce an aglycosylated variant increased the protective effect of the antibody.

Example 11

Ch3G8 N297Q Protects Mice from ch6A6-Mediated Platelet Depletion

An aglycosylated version of ch3G8- γ 1 was prepared by mutating the expression polynucleotide encoding ch3G8- γ 1 so that residue 297 was changed from asparagine (N) to glutamine acid (Q), and expressing the encoded antibody. Residue 297 lies in an N-linked glycosylation site, and this mutation prevents glycosylation of the Fc domain at this site. This aglycosylated antibody, ch3G8 N297Q, was produced in HEK-293 cells as described for ch3G8- γ 1 (see Example 4, supra). The ability of ch3G8-N297Q to protect against ch6A6-mediated platelet depletion was tested using the protocol described above.

When muFc γ RIII^{-/-}, huFc γ RIIIA transgenic mice were injected with 1 μ g/g of the aglycosyl form of ch3G8 (ch3G8 N297Q) one hour before i.p. injection of ch6A6, approximately 75% of the platelets were depleted at the 2-hour time point (FIG. 8). Platelet levels increased faster than in the absence of ch3G8 N297Q, and returned to normal by 24 hours after ch6A6 injection.

When muFc γ RIII^{-/-}, huFc γ RIIIA transgenic mice were injected with 1 μ g/g ch3G8 N297Q one hour before i.v. injection of ch6A6, approximately 60% of the platelets were depleted at the 2 hours time point (FIG. 9). Platelet levels increased faster than in the absence of ch3G8 N297Q, and returned to normal by 48 hours after ch6A6 injection.

When muFc γ RIII^{-/-}, huFc γ RIIIA transgenic mice were injected with ch3G8 N297Q (2 μ g/g) one hour before i.v. injection of ch6A6, only 40% of the platelets were depleted at the 2 hours time point (FIG. 9). Platelet levels increased faster than in the absence of ch3G8 N297Q, and returned to normal by 5 hours after ch6A6 injection.

Thus, ch3G8-N297Q was consistently able to significantly improve platelet counts. Binding of 3G8 to human CD16 on effector cells blocks the ability of CD16 to interact with immune complexes and trigger effector functions such as ADCC or phagocytosis. Chimeric and mouse 3G8 molecules have similar ability to bind CD16 and are similar in their ability to inhibit the binding of sCD16 to immune complexes in vitro. Without intending to be bound by a particular mechanism, the binding (and thus) the blocking activity of the mAb is thought to be confined to the Fab portion of the antibody and blocking of huCD16 is believed to be the mechanism of protection in the transgenic mouse ITP model. The data above suggest that the glycosylation

state of the Fc domain can affect the in vivo protective capacity of anti-CD16A antibodies. Ablation of Fc domain glycosylation (e.g., with D265A or N297Q mutations, or by using a human gamma2 Fc domain) reduces or eliminates Fc binding to FcR. In the case of the aglycosyl (N297Q) variant, complement fixation is also abolished.

Example 12

Neutrophil Levels Following Administration of Aglycosyl CD16A Binding Proteins

The effect of an aglycosylated CD16A binding protein on neutrophil levels was tested and compared to that of glycosylated CD16A binding proteins. CD16A binding proteins, or the controls such as irrelevant human IgG1 (negative control) or murine RB6-8C5 (positive control), were administered to groups of muFc γ RIII^{-/-}, huFc γ RIIIB transgenic mice at a concentration of 5 μ g/g in phosphate buffered saline (PBS). Another negative control was administered PBS alone. Twenty four hours later, mice were euthanized and blood, spleen and bone marrow were collected. Neutrophils were analyzed by FACS. Staining experiments were performed in RPMI containing 3% FCS. Murine cells were stained using FITC-conjugated 3G8 (PharMingen) and R-PE-conjugated RB6-8C5 (PharMingen). Samples were analyzed by flow-cytometry using a FACSCALIBUR™ (Becton Dickinson).

Intraperitoneal injection of 5 μ g/g ch3G8 (prepared as described above) resulted in murine neutrophil depletion in the blood and spleen (FIG. 10; upper right quadrant). Similar results were seen following administration of murine 3G8 (results not shown). In the bone marrow of ch3G8 treated animals, neutrophils stained weakly for CD16, which could indicate receptor occupancy by the chimeric antibody or shedding (FIG. 10; see shift from the upper right quadrant to the upper left quadrant). In contrast, intraperitoneal injection of 5 μ g/g ch3G8 N297Q did not result in murine neutrophil depletion in the blood, spleen or bone marrow (FIG. 10). In additional experiments, humanized glycosylated 3G8 antibodies showed substantially more depletion of circulating blood neutrophils compared to aglycosylated forms of the same antibodies.

Example 13

Autoimmune Hemolytic Anemia Model

This example demonstrates that administration of CD16A binding protein prevents red blood cell depletion in a model of autoimmune hemolytic anemia.

The ability of the Hu3G8-5.1-N297Q monoclonal antibody to prevent antibody-dependent red blood cell depletion in muFcRIII^{-/-}, huFcRIIIa+ mice was evaluated. Hu3G8-5.1-N297Q is an aglycosyl antibody with the heavy chain Hu3G8VH-5 and the light chain Hu3G8VH-1 and the indicated substitution of asparagine 297. Mice were bled on day 0 and RBC levels were determined using a Z2™ COULTER COUNTER® particle analyzer. The next day groups of 3 animals each were then injected intravenously with either 0.5 mg/kg Hu3G8-5.1-N297Q or PBS. One group of mice did not receive any compound. One hour later, RBC depletion was induced in the first two groups by administering mouse anti-RBC IgG2a Mab 34-3C to each animal intraperitoneally (i.p.) (2.5 mg/kg). Animals were bled 2 hrs, 5 hrs, 24 hrs and 48 hrs after administration of 34-3C and RBC counts were determined. Data was analyzed by plotting

45

RBC count versus. The data, depicted in FIG. 11, demonstrate the ability of Hu3G8-5.1-N297Q to prevent RBC depletion in this model.

Example 14

Inhibition of Antibody-Dependent Cellular Cytotoxicity (ADCC)

This example demonstrates that humanized 3G8 variants inhibit ADCC in vitro and with an activity similar to that of mouse 3G8.

Methods: The protocol for assessment of antibody dependent cellular cytotoxicity (ADCC) is similar to that previously described in (Ding et al., 1998, *Immunity* 8:403-11). Briefly, target cells from the HER2-overexpressing breast cancer cell line SK-BR-3 were labeled with the europium chelate bis(acetoxymethyl)2,2':6',2''-terpyridine-6,6''-dicarboxylate (DELFLIA® BATDA Reagent, Perkin Elmer/Wallac). The labeled target cells were then opsonized (coated) with either chimeric anti-HER2 (ch4D5, 100 ng/ml) or chimeric anti-fluorescein (ch4-4-20, 1 ug/ml) antibodies. In the case of the anti-fluorescein antibody, SK-BR-3 cells were coated with the fluorescein hapten prior to antibody opsonization. Peripheral blood mononuclear cells (PBMC), isolated by FICOLL-PAQUE™ (Amersham Pharmacia) gradient centrifugation, were used as effector cells (Effector:Target ratio: ch4D5=(37.5:1) and ch4-4-20=(75:1)). Following a 3.5 hour incubation at 37° C., 5% CO₂, cell supernatants were harvested and added to an acidic europium solution (DELFLIA® Europium Solution, Perkin Elmer/Wallac). The fluorescence of the Europium-TDA chelates formed was quantitated in a time-resolved fluorometer (VICTOR² 1420, Perkin Elmer/Wallac). Maximal release (MR) and spontaneous release (SR) were determined by incubation of target cells with 2% TX-100 and media alone, respectively. Antibody independent cellular cytotoxicity (AICC) was measured by incubation of target and effector cells in the absence of antibody. Each assay is performed in triplicate. The mean percentage specific lysis was calculated as: (ADCC-AICC)/(MR-SR)×100.

Results: Addition of anti-CD16 variants inhibited ADCC mediated through antibodies directed against the HER2/neu protein (ch4D5) (FIG. 12), or the hapten, fluorescein (ch4-4-20) (FIG. 13). Inhibition of the ch4D5 mediated ADCC was greater than 50% at 300 ng/ml for all 3G8 variants tested while isotype control antibodies had no effect in the assay. In the case of the anti-fluorescein antibody, inhibition was approximately 50% at concentrations above 1 ug/ml for murine 3G8 (FIG. 13A) and humanized 3G8 variants (FIG. 13B), while isotype control antibodies and chimeric 3G8 had little effect.

Example 15

Administration of Hu3G8-5.1-N297Q Prevents Immune Thrombocytopenia (ITP) in huFcRIIIa+, huFcRIIIa-Mice

This example shows that that administration of anti-CD16A antibodies protects against ITP mediated by CD32A. As in FcγRIII-/-, hCD16A mice, administration of the ch6A6 antibody induces ITP in FcγRIII-/-, hCD32A transgenic mice. Five hours after injection of 0.1 μg/g ch6A6 i.p., approximately 80% of the platelets are depleted (not shown). The number of platelets remained low for 24 hours after ch6A6 injection, and then progressively increased to

46

return to normal 48 hours after ch6A6 injection. As expected, the i.v. injection of hu3G8-5.1 (0.5 μg/g) one hour prior to ch6A6 injection did not protect FcγRIII-/-, hCD32A mice against ITP (not shown).

As in single transgenic mice, ch6A6 induces ITP in FcγRIII-/-, hCD16A, hCD32A double transgenic mice. Five hours after injection of 0.1 μg/g ch6A6 i.p., approximately 80% of the platelets were depleted (FIG. 14). The number of platelets remained low for 24 hours after ch6A6 injection, and then progressively increased to return to normal 48 hours after ch6A6 injection.

In contrast to FcγRIII-/-, hCD32A mice, FcγRIII-/-, hCD16A, hCD32A mice were protected against ITP by administration of hu3G8-5.1. Complete protection was observed when 1 μg/g h3G8 5.1 is injected one hour prior to ch6A6 i.p. injection; and partial protection resulted from administration of 0.75 μg/g or 0.5 μg/g of h3G8-5.1. (FIG. 14). Thus, the data indicate that although CD32A can mediate ITP, the injection of 1 μg/g of h3G8-5.1 completely and unexpectedly protects mice against platelet depletion.

Example 16

Prevention of Platelet Depletion Using Hu3G8-5.1-N297Q Produced in CHO-S Cell Line

Hu3G8-5.1-N297Q was produced in a CHO-S cell line. The ability of this antibody to protect against ITP in FcγRIII-/-, hCD16A single transgenic mice was determined using the procedure described in Example 13. As is shown in FIG. 15, administration of 0.5 mg/kg or more Hu3G8-5.1-N297Q produced in CHO-S cells one hour prior to ch6A6 i.p. injection completely protects mice against ITP.

Example 17

Therapeutic Effect of Aglycosylated Humanized Antibodies

ITP was induced in mice as described above, by i.p. injection of 0.1 ug/g ch6A6 at time 0. Two hours later, the number of platelets in the plasma was determined to confirm the presence of ITP. Three hours after i.p. injection of ch6A6, mice were injected i.v. with hu3G8-5.1-N297Q at different concentration (arrow). The results (FIG. 16A) indicate that the number of platelets rapidly returns to normal after Hu3G8-5.1-N297Q injection whereas the number of platelets remains low in non-treated mice. These results demonstrate that administration of the hu3G8-5.1-N297Q antibody can be used to cure ITP in the mouse model.

In this experiment, ITP was induced by i.p. injection of 0.1 ug/g ch6A6 at time 0. Two hours later, the number of platelets in the plasma was determined to confirm the presence of ITP. Three hours after i.p. injection of ch6A6, mice were injected i.v. with hu3G8-22.1-N297Q or hu3G8-22.43-N297Q at 0.5 ug/g (arrow). The results indicate that the number of platelets rapidly returns to normal after Hu3G8-22.1-N297Q injection whereas the number of platelets remains low in non-treated mice and in mice treated with Hu3G8-22.43-N297Q (FIG. 16B). These data indicate that hu3G8-22.1-N297Q can be used to cure ITP in the mouse model.

Therapeutic Effect of Hu3G8-22.1-N297Q in AHA
in muFcγRIII-/-, huFcγRIIIA Transgenic Mice

In this experiment, AHA was induced by i.p. injection of 50 ug mouse anti-RBC IgG2a Mab 34-3C at day 0. On day 1, the number of RBC in the blood was determined to confirm the presence of AHA. Two hours later, mice were injected i.v. with Hu3G8-22.1-N297Q at various concentrations (arrow). The results indicate that the number of RBC remained stable after Hu3G8-22.1-N297Q injection whereas the number of RBC continued to drop in non-treated mice (FIG. 17). The optimal concentration of Hu3G8-22.1-N297Q is 0.5 ug/g. The number of RBC returned to normal in all mice at day 7. Control mice were bled every day but not injected in order to determine the effect of repeated bleedings on the number of RBC. These results in the mouse model indicate that Hu3G8-22.1-N297Q can be used to cure AHA. Hu3G8-22.1-N297Q prevents further RBC depletion by autoantibodies and therefore protects mice against anemia.

TABLE 3A*

V _H SEQUENCES							
	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
3G8VH	A	A	A	A	A	A	A
Ch3G8VH	A	A	A	A	A	A	B
HxC	B	A	B	A	A	A	B
CxH	A	A	A	A	B	A	B
Hu3G8VH-1	B	A	B	A	B	A	B
Hu3G8VH-2	C	A	B	A	B	A	B
Hu3G8VH-3	D	A	B	A	B	A	B
Hu3G8VH-4	B	A	B	A	C	B	B
Hu3G8VH-5	B	A	B	A	C	A	B
Hu3G8VH-6	B	B	B	A	B	B	B
Hu3G8VH-7	B	B	B	A	B	A	B
Hu3G8VH-8	B	A	B	A	B	C	B
Hu3G8VH-9	B	A	B	B	B	B	B
Hu3G8VH-10	B	A	B	A	B	B	B
Hu3G8VH-11	B	A	B	B	B	A	B
Hu3G8VH-12	B	A	B	C	B	A	B
Hu3G8VH-13	B	A	B	D	B	A	B
Hu3G8VH-14	B	A	B	E	B	A	B
Hu3G8VH-15	B	A	B	A	D	A	B
Hu3G8VH-16	B	A	B	A	E	A	B
Hu3G8VH-17	B	A	B	A	F	A	B
Hu3G8VH-18	B	A	B	A	G	A	B
Hu3G8VH-19	B	A	B	A	C	C	B
Hu3G8VH-20	B	B	B	C	B	A	B
Hu3G8VH-21	B	A	B	A	D	B	B
Hu3G8VH-22	B	B	B	C	B	C	B
Hu3G8VH-23	B	B	B	C	E	C	B
Hu3G8VH-24	B	B	B	C	F	C	B
Hu3G8VH-25	B	B	B	C	G	C	B
Hu3G8VH-26	B	B	B	C	C	C	B
Hu3G8VH-27	B	B	B	C	E	D	B
Hu3G8VH-28	B	B	B	C	F	D	B
Hu3G8VH-29	B	B	B	C	G	D	B
Hu3G8VH-30	B	B	B	C	C	D	B
Hu3G8VH-31	E	B	B	C	B	A	B
Hu3G8VH-32	E	B	B	H	B	A	B
Hu3G8VH-33	E	B	B	H	B	A	B
Hu3G8VH-34	E	B	B	C	B	C	B
Hu3G8VH-35	E	B	B	C	C	C	B
Hu3G8VH-36	E	B	B	H	C	D	B
Hu3G8VH-37	E	B	B	H	E	C	B
Hu3G8VH-38	E	B	B	F	B	A	B
Hu3G8VH-39	E	B	B	I	B	A	B
Hu3G8VH-40	E	B	B	G	B	A	B
Hu3G8VH-41	E	B	B	J	B	A	B
Hu3G8VH-42	E	B	B	C	H	A	B
Hu3G8VH-43	E	B	B	C	H	C	B

TABLE 3A*-continued

V _H SEQUENCES							
	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
Hu3G8VH-44	E	B	B	C	I	D	B
Hu3G8VH-45	E	B	B	C	J	D	B

*Letters in Table 3A refer to sequences in Tables 3B-H.

TABLE 3B

FR1					
A	B	C	D	E	RESIDUE
Q	Q	Q	Q	Q	1
V	V	V	V	I	2
T	T	T	T	T	3
L	L	L	L	L	4
K	R	K	R	K	5
E	E	E	E	E	6
S	S	S	S	S	7
G	G	G	G	G	8
P	P	P	P	P	9
G	A	A	A	T	10
I	L	L	L	L	11
L	V	V	V	V	12
Q	K	K	K	K	13
P	P	P	P	P	14
S	T	T	T	T	15
Q	Q	Q	Q	Q	16
T	T	T	T	T	17
L	L	L	L	L	18
S	T	T	T	T	19
L	L	L	L	L	20
T	T	T	T	T	21
C	C	C	C	C	22
S	T	T	T	T	23
F	F	F	F	F	24
S	S	S	S	S	25
G	G	G	G	G	26
F	F	F	F	F	27
S	S	S	S	S	28
L	L	L	L	L	29
R	S	S	R	S	30
30	31	32	33	34	Seq ID No

TABLE 3C

CDR1		
A	B	RESIDUE
T	T	31
S	S	32
G	G	33
M	V	34
G	G	35
V	V	35A
G	G	35B
35	36	Seq ID No

TABLE 3D

FR2		
A	B	RESIDUE
W	W	36
I	I	37
R	R	38
Q	Q	39
P	P	40

49

TABLE 3D-continued

FR2		
A	B	RESIDUE
S	P	41
G	G	42
K	K	43
G	A	44
L	L	45
E	E	46
W	W	47
L	L	48
A	A	49
37	38	Seq ID No.

TABLE 3E

CDR2										RESIDUE
A	B	C	D	E	F	G	H	I	J	
H	H	H	H	H	L	H	L	H	L	50
I	I	I	I	I	I	I	I	I	I	51
W	Y	W	Y	W	D	F	W	D	W	52
W	W	W	W	W	W	W	W	W	W	53
D	N	D	D	N	D	D	D	D	N	54
D	D	D	D	D	D	D	D	D	D	55
D	D	D	D	D	D	D	D	D	D	56
K	K	K	K	K	K	K	K	K	K	57
R	R	R	R	R	R	R	R	R	R	58
Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	59
N	N	S	N	N	S	S	S	S	S	60
P	P	P	P	P	P	P	P	P	P	61
A	A	S	A	A	S	S	S	S	S	62
L	L	L	L	L	L	L	L	L	L	63
K	K	K	K	K	K	K	K	K	K	64
S	S	S	S	S	S	S	S	S	S	65
39	40	41	42	43	44	45	46	47	48	Seq ID No

TABLE 3F

FR3										RESIDUE
A	B	C	D	E	F	G	H	I	J	
R	R	R	R	R	R	R	R	R	R	66
L	L	L	L	L	L	L	L	L	L	67
T	T	T	T	T	T	T	T	T	T	68
I	I	I	I	I	I	I	I	I	I	69
S	S	S	S	S	S	S	T	T	T	70
K	K	K	K	K	K	K	K	K	K	71
D	D	D	D	D	D	D	D	D	D	72
T	T	T	T	T	T	T	T	T	T	73
S	S	S	S	S	S	S	S	S	S	74
S	K	K	K	K	K	K	K	K	K	75
N	N	N	N	N	N	N	N	N	N	76
Q	Q	Q	Q	Q	Q	Q	Q	Q	Q	77
V	V	V	V	V	V	V	V	V	V	78
F	V	V	V	V	V	V	V	V	V	79
L	L	L	L	L	L	L	L	L	L	80
K	T	T	T	T	T	T	T	T	T	81
I	M	M	M	M	M	M	M	M	M	82
A	T	T	T	T	T	T	T	T	T	82A
S	N	N	N	N	N	N	N	N	N	82B
V	M	M	M	M	M	M	M	M	M	82C
D	D	D	D	D	D	D	D	D	D	83
T	P	P	P	P	P	P	P	P	P	84
A	V	V	V	V	V	V	V	V	V	85
D	D	D	D	D	D	D	D	D	D	86
T	T	T	T	T	T	T	T	T	T	87
A	A	A	A	A	A	A	A	A	A	88
T	T	T	T	T	T	T	T	T	T	89
Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	90
Y	Y	Y	Y	Y	Y	Y	Y	Y	Y	91

50

TABLE 3F-continued

FR3										RESIDUE
A	B	C	D	E	F	G	H	I	J	
C	C	C	C	C	C	C	C	C	C	92
A	A	A	A	A	A	A	A	A	A	93
Q	R	Q	T	K	A	H	R	H	Q	94
49	50	51	52	53	54	55	56	57	58	Seq ID No

TABLE 3G

CDR3					RESIDUE
A	B	C	D		
I	I	I	I		95
N	N	N	N		96
P	P	P	P		97
A	A	A	A		98
W	W	Y	Y		99
F	F	F	F		100
A	D	A	D		101
Y	Y	Y	Y		102
59	60	61	62		Seq ID No

TABLE 3H

FR4			RESIDUE
A	B		
W	W		103
G	G		104
Q	Q		105
G	G		106
T	T		107
L	L		108
V	V		109
T	T		110
V	V		111
S	S		112
A	S		113
63	64		Seq ID No

TABLE 4A*

V _L SEQUENCES							
	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
3G8VL	A	A	A	A	A	A	A
Ch3G8VL	A	A	A	A	A	A	A
Hu3G8VL-1	B	A	A	A	B	A	B
Hu3G8VL-2	B	B	A	A	B	A	B
Hu3G8VL-3	B	C	A	A	B	A	B
Hu3G8VL-4	B	D	A	A	B	A	B
Hu3G8VL-5	B	E	A	A	B	A	B
Hu3G8VL-6	B	F	A	A	B	A	B
Hu3G8VL-7	B	G	A	A	B	A	B
Hu3G8VL-8	B	A	A	B	B	A	B
Hu3G8VL-9	B	A	A	C	B	A	B
Hu3G8VL-10	B	A	A	D	B	A	B
Hu3G8VL-11	B	A	A	E	B	A	B
Hu3G8VL-12	B	A	A	F	B	A	B
Hu3G8VL-13	B	A	A	G	B	A	B
Hu3G8VL-14	B	A	A	A	B	B	B
Hu3G8VL-15	B	A	A	A	B	C	B
Hu3G8VL-16	B	A	A	A	B	D	B
Hu3G8VL-17	B	A	A	A	B	E	B
Hu3G8VL-18	B	B	A	D	B	A	B
Hu3G8VL-19	B	B	A	D	B	D	B
Hu3G8VL-20	B	B	A	D	B	E	B

TABLE 4A*-continued

<u>V_L SEQUENCES</u>							
	FR1	CDR1	FR2	CDR2	FR3	CDR3	FR4
Hu3G8VL-21	B	C	A	D	B	A	B
Hu3G8VL-22	B	C	A	D	B	D	B
Hu3G8VL-23	B	C	A	D	B	E	B
Hu3G8VL-24	B	D	A	D	B	A	B
Hu3G8VL-25	B	D	A	D	B	D	B
Hu3G8VL-26	B	D	A	D	B	E	B
Hu3G8VL-27	B	E	A	D	B	A	B
Hu3G8VL-28	B	E	A	D	B	D	B
Hu3G8VL-29	B	E	A	D	B	E	B
Hu3G8VL-30	B	A	A	D	B	D	B
Hu3G8VL-31	B	A	A	D	B	E	B
Hu3G8VL-32	B	A	A	H	B	A	B
Hu3G8VL-33	B	A	A	I	B	A	B
Hu3G8VL-34	B	A	A	J	B	A	B
Hu3G8VL-35	B	B	A	H	B	D	B
Hu3G8VL-36	B	C	A	H	B	D	B
Hu3G8VL-37	B	E	A	H	B	D	B
Hu3G8VL-38	B	B	A	I	B	D	B
Hu3G8VL-39	B	C	A	I	B	D	B
Hu3G8VL-40	B	E	A	I	B	D	B
Hu3G8VL-41	B	B	A	J	B	D	B
Hu3G8VL-42	B	C	A	J	B	D	B
Hu3G8VL-43	B	E	A	J	B	D	B
Hu3G8VL-44	B	A	A	K	B	A	B

*Letters in Table 4A refer to sequences in Tables 4B-H.

TABLE 4B

<u>FR1</u>		
A	B	RESIDUE
D	D	1
T	I	2
V	V	3
L	M	4
T	T	5
Q	Q	6
S	S	7
P	P	8
A	D	9
S	S	10
L	L	11
A	A	12
V	V	13
S	S	14
L	L	15
G	G	16
Q	E	17
R	R	18
A	A	19
T	T	20
I	I	21
S	N	22
C	C	23
65	66	Seq ID No

TABLE 4C

<u>CDR1</u>							
A	B	C	D	E	F	G	RESIDUE
K	R	K	K	K	K	K	24
A	A	S	A	A	A	A	25
S	S	S	S	S	S	S	26
Q	Q	Q	Q	Q	Q	Q	27
S	S	S	S	S	S	S	27A
V	V	V	V	V	V	V	27B
D	D	D	D	D	D	D	27C

TABLE 4C-continued

<u>CDR1</u>							
A	B	C	D	E	F	G	RESIDUE
F	F	F	F	F	F	F	27D
D	D	D	D	D	D	D	28
G	G	G	G	G	G	G	29
D	D	D	D	D	D	D	30
S	S	S	S	S	S	S	31
F	F	F	Y	F	F	Y	32
M	M	M	M	L	M	L	33
N	N	N	N	N	A	A	34
67	68	69	70	71	72	73	Seq ID No

TABLE 4D

<u>FR2</u>	
A	RESIDUE
W	35
Y	36
Q	37
Q	38
K	39
P	40
G	41
Q	42
P	43
P	44
K	45
L	46
L	47
I	48
Y	49
74	Seq ID No

TABLE 4E

<u>CDR2</u>											
A	B	C	D	E	F	G	H	I	J	K	RESIDUE
T	D	W	T	D	D	S	S	S	T	T	50
T	A	A	T	A	A	A	T	T	T	T	51
S	S	S	S	S	S	S	S	S	S	S	52
N	N	N	N	N	N	N	N	N	N	S	53
L	L	L	L	L	L	L	L	L	L	L	54
E	E	E	E	E	A	Q	E	Q	Q	Q	55
S	S	S	T	T	T	S	S	S	S	S	56
75	76	77	78	79	80	81	82	83	84	85	Seq ID No

TABLE 4F

<u>FR3</u>		
A	B	RESIDUE
G	G	57
I	V	58
P	P	59
A	D	60
R	R	61
F	F	62
S	S	63
A	G	64
S	S	65
G	G	66
S	S	67
G	G	68
T	T	69
D	D	70

TABLE 4F-continued					
FR3			5		
A	B	RESIDUE			
F	F	71	10		
T	T	72			
L	L	73			
N	T	74			
I	I	75			
H	S	76			
P	S	77			
V	L	78			
E	Q	79			
E	A	80			
E	E	81	15		
D	D	82			
T	V	83			
A	A	84			
T	V	85			
Y	Y	86			
Y	Y	87			
C	C	88			
86	87	Seq ID No			

TABLE 4G					
<u>CDR3</u>					
A	B	C	D	E	RESIDUE
Q	Q	Q	Q	Q	89
Q	Q	Q	Q	Q	90
S	S	S	S	S	91

TABLE 4G-continued					
CDR3					
A	B	C	D	E	RESIDUE
N	Y	Y	N	N	92
E	S	E	S	E	93
D	T	D	D	T	94
P	P	P	P	P	95
Y	Y	Y	Y	Y	96
T	T	T	T	T	97
88	89	90	91	92	Seq ID No

TABLE 4H		
<u>FR4</u>		
A	B	RESIDUE
F	F	98
G	G	99
G	Q	100
G	G	101
T	T	102
K	K	103
L	L	104
E	E	105
I	I	106
K	K	107
93	94	Seq ID No

TABLE 5

Hu3G8VL-1	
(SEQ ID NO:95)	
CGAGCTAGCTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCCTC TTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATGGGTGACAATGACA TCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCGACATCGTGATGACCCAATCTCCAGACTCTTTGGCTGTGTCTCTA GGGGAGAGGGCCACCATCAACTGCAAGGCCAGCCAAAGTGTTGATTTTGATGGTGATAGTTTTATGAACTGGTACCAACA GAAACCAGGACAGCCACCCAAACTCCTCATCTATACTACATCCAATCTAGAATCTGGGGTCCCAGACAGGTTTAGTGCCA GTGGGTCTGGGACAGACTTCACCCCTCACCATCAGCAGCCTGCAGGCTGAGGATGTGGCAGTTTATTACTGTCAGCAAAGT AATGAGGATCCGTACACGTTCCGGACAGGGGACCAAGCTTGAgATcAAACGaACTGTGGCTGCACCATCGGTCTTCATCTT AATGAGGATCCGTACACGTTCCGGACAGGGGACCAAGCTTGAgATcAAA	
Hu3G8VL-1	
(SEQ ID NO:96)	
DIVMTQSPDSLAVSLGERATINCKASQSVDFDGDSEFMNWWYQQKPGQPPKLLIYTTSNLESGVPDRFSGSGSGTDFTLTIS SLQAEDVAVYYCQQSNEDPYTFGQGTKLEIK	
Hu3G8VL-1K	
(SEQ ID NO:97)	
CGAGCTAGCTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCCTC TTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATGGGTGACAATGACA TCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCGACATCGTGATGACCCAATCTCCAGACTCTTTGGCTGTGTCTCTA GGGGAGAGGGCCACCATCAACTGCAAGGCCAGCCAAAGTGTTGATTTTGATGGTGATAGTTTTATGAACTGGTACCAACA GAAACCAGGACAGCCACCCAAACTCCTCATCTATACTACATCCAATCTAGAATCTGGGGTCCCAGACAGGTTTAGTGCCA GTGGGTCTGGGACAGACTTCACCCCTCACCATCAGCAGCCTGCAGGCTGAGGATGTGGCAGTTTATTACTGTCAGCAAAGT AATGAGGATCCGTACACGTTCCGGACAGGGGACCAAGCTTGAgATcAAACGaACTGTGGCTGCACCATCGGTCTTCATCTT CCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAAATTCTATCCCAGAGAGGCCA AAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAAGTCCCAGGAGAGTGTCACAGAGCAGGACAGCAAGGACAGC ACCTACAGCCTCAGCAGCACCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCA TCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTTAGTTCAGAGTCGACTCTAGAGGATCCCCG GGTACCGAGCTCGAATTC	
Hu3GSVL-1K	
(SEQ ID NO:98)	
DIVMTQSPDSLAVSLGERATINCKASQSVDFDGDSEFMNWWYQQKPGQPPKLLIYTTSNLESGVPDRFSGSGSGTDFTLTIS SLQAEDVAVYYCQQSNEDPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKDSSTLSSTLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC	
Hu3G8VL-43	
(SEQ ID NO:99)	
CGAGCTAGCTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCCTC TTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATGGGTGACAATGACA TCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCGACATCGTGATGACCCAATCTCCAGACTCTTTGGCTGTGTCTCTA	

TABLE 5-continued

GGGGAGAGGGGCCACCATCAACTGCAAGtCCAGCCAAAGTGTGATTTTGGATGGTGATAGTTTTATGAACTGGTACCAACA GAAACCAGGACAGCCACCCAAACTCCTCATCTATACTACATCCAgTCTAGAATCTGGGGTCCCAGACAGGTTTAGTGCCA GTGGGTCTGGGACAGACTTCACCTCACCATCAGCAGCCTGCAGGCTGAGGATGTGGCAGTTTATTACTGTCAGCAAAGT AATtcGGATCCGTACACGTTCTGGACAGGGGACCAAGCTTGAgATcAAA Hu3G8VL-43	(SEQ ID NO:100)
DIVMTQSPDSLAVSLGERATINCKSSQSVDFDGDSEFMNWWYQQKPGQPPKLLIYTTSSLESQVDFRSGSGSGTDFTLTIS SLQAEDVAVYYCQQSNSDPYTFGQGTKLEIK Hu3G8VL-43 + Kappa	(SEQ ID NO:101)
CGAGCTAGCTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCCTC TTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATGGGTGACAATGACA TCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCGACATCGTGATGACCAATCTCCAGACTCTTTGGCTGTGTCTCTA GGGGAGAGGGGCCACCATCAACTGCAAGtCCAGCCAAAGTGTGATTTTGGATGGTGATAGTTTTATGAACTGGTACCAACA GAAACCAGGACAGCCACCCAAACTCCTCATCTATACTACATCCAgTCTAGAATCTGGGGTCCCAGACAGGTTTAGTGCCA GTGGGTCTGGGACAGACTTCACCTCACCATCAGCAGCCTGCAGGCTGAGGATGTGGCAGTTTATTACTGTCAGCAAAGT AATtcGGATCCGTACACGTTCTGGACAGGGGACCAAGCTTGAgATcAAACGaACTGTGGCTGCACCATCGGTCTTCATCTT CCCGCCATCTGATGAGCAGTTGAAATCTGGAAGTGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCAGAGAGGCCA AAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAACTCCCAGGAGAGTGTACAGAGCAGGACAGCAAGGACAGC ACCTACAGCCTCAGCAGCACCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCA TCAGGGCCTGAGCTCGCCCGTCACAAAGAGCTTCAACAGGGGAGAGTGTAGTTCTAGAGTCGACTCTAGAGGATCCCCG GGTACCGAGCTCGAATTC Hu3G8VL-43K	(SEQ ID NO:102)
DIVMTQSPDSLAVSLGERATINCKSSQSVDFDGDSEFMNWWYQQKPGQPPKLLIYTTSSLESQVDFRSGSGSGTDFTLTIS SLQAEDVAVYYCQQSNSDPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS GNSQESVTEQDSKDYSLSSLTLTLKADYEKHKVYACEVTHQGLSSPVTKSPNRGEC Hu3G8VH-1	(SEQ ID NO:103)
GCTAGCgtttaaacttaagcttGTTGACTAGTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCAT GGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGAC ATATATATGGGTGACAATGACATCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTTACCTTGAGAGAGTCTGG CCCTGCGCTGGTGAAGCCCACACAGACCCTCACACTGACTTGTACCTTCTCTGGGTTTTACTGAGCACTTCTGGTATGG GTGTAGGCTGGATTCTGTCAGCCTCCCGGAAGGCTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTAT AATCCAGCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCTCAAAAACAGGTAGTCCTCACAATGACCAACATGGA CCCTGTGGATACTGCCACATACTACTGTGCTCGGATAAACCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCA CTGTGAGCTCA Hu3G8VH-1	(SEQ ID NO:104)
QVTLRESGPALVKPTQTTLTLCTFSGFSLSTSGMGVGVWIRQPPGKALEWLAHIWWDDDKRYNPALKSRLTISKDTSKNQV VLTMTNMDPVDATYYCARINPAWFAYWGQGLTVTVSS Hu3G8VH-1G1	(SEQ ID NO:105)
GCTAGCgtttaaacttaagcttGTTGACTAGTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCAT GGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGAC ATATATATGGGTGACAATGACATCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTTACCTTGAGAGAGTCTGG CCCTGCGCTGGTGAAGCCCACACAGACCCTCACACTGACTTGTACCTTCTCTGGGTTTTACTGAGCACTTCTGGTATGG GTGTAGGCTGGATTCTGTCAGCCTCCCGGAAGGCTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTAT AATCCAGCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCTCAAAAACAGGTAGTCCTCACAATGACCAACATGGA CCCTGTGGATACTGCCACATACTACTGTGCTCGGATAAACCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCA CTGTGAGCTCAgcctccaccaagggcccatcggtcttccccctggcaccctcctccaagagcacctctgggggacagcg gccctgggctgcctggtcaaggactacttccccgaaccgggtgacgggtgtcgtggaactcaggcgccctgaccagcggtg gcacacctccccggctgtcctacagtcctcaggactctactccctcagcagcggtggtgaccgtgccctccagcagcttgg gcacccagacatcacatctgcaacgtgaatcacagcccagcaacaccaaggtggacaagagagttgagcccaaatcttgt gacaaaactcacacatgccacccgtgcccagcacctgaactcctggggggaccgtcagtccttctcttccccccaaaacc caaggacaccctcatgatctccccgacccctgaggtcacatgcgtggtggtggacgtgagccacgaagaccctgaggtca agttcaactggtacgtggacggcggtggaggtgcataatgccaaagacaaagccgaggaggagcagtaaacagcacgtac cgtgtggtcagcgctcctcaccgtcctgcaccaggactggctgaatggcaaggagtaaacgtgcaaggtctccaacaaagc cctcccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaaccacaggtgtacaccctgcccccat cccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggag tgggagagcaatgggcagccggagacaactacaagaccagcctcccgtgctggactccgacggctccttcttccctcta cagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctctgcaca accactacacgcagaagcctctccctgtctccgggtaaatgagtgcgccgcGAATTC Hu3GSVH-1G1	(SEQ ID NO:107)
QVTLRESGPALVKPTQTTLTLCTFSGFSLSTSGMGVGVWIRQPPGKALEWLAHIWWDDDKRYNPALKSRLTISKDTSKNQV VLTMTNMDPVDATYYCARINPAWFAYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN SGALTSGVHTFPAVLQSSGLYSLSSVVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPS VFLFPPKPKDTLMISRTPEVTCVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYNSTYRVVSVLTVLHQDNLNGKEY KCKVSNKALPAPIEKTIKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTTPVLD SDGSFFLYSKLTVDKSRWQQGNVFPSCSVMEALHNHYTQKSLSLSPGK Hu3G8VH-5	(SEQ ID NO:108)
GCTAGCgtttaaacttaagcttGTTGACTAGTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCAT GGGATGGAGCTGTATCATCCTCTTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGAC ATATATATGGGTGACAATGACATCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTTACCTTGAGAGAGTCTGG CCCTGCGCTGGTGAAGCCCACACAGACCCTCACACTGACTTGTACCTTCTCTGGGTTTTACTGAGCACTTCTGGTATGG GTGTAGGCTGGATTCTGTCAGCCTCCCGGAAGGCTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTAT AATCCAGCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCTCAAAAACAGGTAGTCCTCACAATGACCAACATGGA	

TABLE 5-continued

CCCTGTGGATACTGCCACATACTACTGTGCTCaaATAAAACCCCGCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTGAGCTCAHu3G8VH- 5	(SEQ ID NO:109)
QVTLRESGPALVKPTQTLTLTCTFSGFSLSTSGMGVGWIRQPPGKALEWLAHIWDDDKRYNPALKSRLTISKDTSKNQV VLTMTNMDPVDATATYYCAQINPAWFAYWGQGLTVTVSSHu3G8VH- 5G1Ag	(SEQ ID NO:110)
GCTAGCgtttaaacttaagcttGTTGACTAGTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCTCTTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATATGGGTGACAATGACATCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTTACCCTGAGAGAGTCTGGCCCTGCGCTGGTGAAGCCCACACAGACCCTCACACTGACTTGTACCTTCTCTGGGTTTTTACTGAGCACTTCTGGTATGGGTGTAGGCTGGATTTCGTACAGCTCCCGGAAGGCTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTATAATCCAGCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCAAAACCCAGGTAGTCCTCACAATGACCAACATGGA	
CCCTGTGGATACTGCCACATACTACTGTGCTCaaATAAAACCCCGCCTGGTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTGAGCTCAgacctccaccaagggcccatcggtcttccccctggcacctcctccaagagcacctctgggggacagcggccctgggctgcctggtcaaggactacttccccgaaccgggtgacgggtgtcgtggaactcaggcgccctgaccagcggcgtgcacaccttccccggctgtcctacagtccctcaggactctactccctcagcagcgtggtgacctgacctccagcagcttgggcacccagacctacatctgcaacgtgaatcacaaagcccagcaacaccaaggtggacaagagagttgagcccaaatcttgtgacaaaaactcacacatgcccaccgtgcccagcacctgaactcctggggggaccgtcagttcttctcttccccccaaaacc	
caaggacacctcatgatctcccgacccctgaggtcacatgcgtggtggtggacgtgagccacgaagacctgaggtcaagttcaactggtagctggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtagCaGagcacgtac	
cgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagacaactacaagaccacgcctcccggtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctctgcaca	
accactacacgcagaagagcctctccctgtctccgggtaaatgagtgcgccgcGAATTC	
Hu3G8VH- 5G1Ag	(SEQ ID NO:111)
QVTLRESGPALVKPTQTLTLTCTFSGFSLSTSGMGVGWIRQPPGKALEWLAHIWDDDKRYNPALKSRLTISKDTSKNQV VLTMTNMDPVDATATYYCAQINPAWFAYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN	
SGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI	
SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD	
SDGSFFLYSKLTVDKSRWQQGNVFPSCSVMHEALHNHYTQKSLSLSPGKHu3G8VH- 22	
GCTAGCgtttaaacttaagcttGTTGACTAGTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCTCTTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATATGGGTGACAATGACATCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTTACCCTGAGAGAGTCTGGCCCTGCGCTGGTGAAGCCCACACAGACCCTCACACTGACTTGTACCTTCTCTGGGTTTTTACTGAGCACTTCTGGTgTGGGTGTAGGCTGGATTTCGTACAGCTCCCGGAAGGCTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTAT	
tctCCAtCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCAAAACCCAGGTAGTCCTCACAATGACCAACATGGA	
CCCTGTGGATACTGCCACATACTACTGTGCTCGGATAAAACCCCGCCTacTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTGAGCTCAHu3G8VH- 22	
QVTLRESGPALVKPTQTLTLTCTFSGFSLSTSGVGVGWIRQPPGKALEWLAHIWDDDKRYSPSLKSRLTISKDTSKNQV VLTMTNMDPVDATATYYCARINPAYFAYWGQGLTVTVSHu3G8VH- 22G1Ag	(SEQ ID NO:113)
GCTAGCgtttaaacttaagcttGTTGACTAGTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCTCTTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCTGGACATATATATATGGGTGACAATGACATCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCCAGGTTACCCTGAGAGAGTCTGGCCCTGCGCTGGTGAAGCCCACACAGACCCTCACACTGACTTGTACCTTCTCTGGGTTTTTACTGAGCACTTCTGGTgTGGGTGTAGGCTGGATTTCGTACAGCTCCCGGAAGGCTCTAGAGTGGCTGGCACACATTTGGTGGGATGATGACAAGCGCTAT	
tctCCAtCCCTGAAGAGCCGACTGACAATCTCCAAGGATACCTCCAAAACCCAGGTAGTCCTCACAATGACCAACATGGA	
CCCTGTGGATACTGCCACATACTACTGTGCTCGGATAAAACCCCGCCTacTTTGCTTACTGGGGCCAAGGGACTCTGGTCACTGTGAGCTCAgacctccaccaagggcccatcggtcttccccctggcacctcctccaagagcacctctgggggacagcggccctgggctgcctggtcaaggactacttccccgaaccgggtgacgggtgtcgtggaactcaggcgccctgaccagcggcgtgcacaccttccccggctgtcctacagtcctcaggactctactccctcagcagcgtggtgacctgacctccagcagcttgggcacccagacctacatctgcaacgtgaatcacaaagcccagcaacaccaaggtggacaagagagttgagcccaaatcttgtgacaaaaactcacacatgcccaccgtgcccagcacctgaactcctggggggaccgtcagttcttctcttccccccaaaacc	
caaggacacctcatgatctcccgacccctgaggtcacatgcgtggtggtggacgtgagccacgaagacctgaggtcaagttcaactggtagctggacggcgtggaggtgcataatgccaaagacaaagccgcgggaggagcagtagCaGagcacgtac	
cgtgtggtcagcgtcctcaccgtcctgcaccaggactggctgaatggcaaggagtacaagtgaaggtctccaacaaagccctccagcccccatcgagaaaaccatctccaaagccaaagggcagccccgagaaccacaggtgtacacctgcccccatcccgggatgagctgaccaagaaccaggtcagcctgacctgcctggtcaaaggcttctatcccagcgacatcgccgtggagtgggagagcaatgggcagccggagacaactacaagaccacgcctcccggtgctggactccgacggctccttcttctctacagcaagctcaccgtggacaagagcaggtggcagcaggggaacgtcttctcatgctccgtgatgcatgaggtctctgcaca	
accactacacagcagaagagcctctccctgtctccgggtaaatgagtgcgccgcGAATTC	
Hu3G8VH- 22G1Ag	(SEQ ID NO:114)
QVTLRESGPALVKPTQTLTLTCTFSGFSLSTSGVGVGWIRQPPGKALEWLAHIWDDDKRYSPSLKSRLTISKDTSKNQV VLTMTNMDPVDATATYYCARINPAYFAYWGQGLTVTVSSASTKGPSVFPLAPSSKSTSGGTAALGCLVKDYFPEPVTVSWN	
SGALTSGVHTFPAVLQSSGLYSLSSVTVPSSSLGTQTYICNVNHKPSNTKVDKRVEPKSCDKTHTCPPCPAPELLGGPSVFLFPPKPKDTLMI	
SRTPEVTCVVVDVSHEDPEVKFNWYVDGVEVHNAKTKPREEQYQSTYRVVSVLTVLHQDWLNGKEYKCKVSNKALPAPIEKTISKAKGQPREPQVYTLPPSRDELTKNQVSLTCLVKGFYPSDIAVEWESNGQPENNYKTTPPVLD	

TABLE 5-continued

SDGSFFLYSKLTVDKSRWQQGNVFSQSVMHENHNTQKSLSLSPGK	
Hu3G8VL-22	(SEQ ID NO:118)
DIVMTQSPDSLAVSLGERATINCKSSQSVDFDGDGSMNWFYQKPGQPPKLLIYTTSNLETGVPDRFSGSGSGTDFTLTIS	
SLQAEDVAVYYCQQSNSDPYTFGQGTKLEIK	
Hu3G8VL-22K	(SEQ ID NO:119)
DIVMTQSPDSLAVSLGERATINCKSSQSVDFDGDGSMNWFYQKPGQPPKLLIYTTSNLETGVPDRFSGSGSGTDFTLTIS	
SLQAEDVAVYYCQQSNSDPYTFGQGTKLEIKRTVAAPSVFIFPPSDEQLKSGTASVVCLLNNFYPREAKVQWKVDNALQS	
GNSQESVTEQDSKDSYSLSSLTLSKADYEKHKVYACEVTHQGLSSPVTKSFNRGEC	
Hu3G8VL-22	(SEQ ID NO:106)
CGAGCTAGCTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCCTC	
TTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCGGACATATATATGGGTGACAATGACA	
TCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCGACATCGTGATGACCCAATCTCCAGACTCTTTGGCTGTGTCTCTA	
GGGGAGAGGGCCACCATCAACTGCAAGTCCAGCCAAAGTGTTGATTTTGATGGTGATAGTTTATGAACTGGTACCAACA	
GAAACCAGGACAGCCACCCAACTCCTCATCTATACTACATCCAATCTAGAACTGGGGTCCCAGACAGGTTTAGTGCCA	
GTGGGTCTGGGACAGACTTCACCCCTCACCATCAGCAGCCTGCAGGCTGAGGATGTGGCAGTTTATTACTGTCAGCAAAGT	
AATTCGGATCCGTACACGTTCCGGACAGGGGACCAAGCTTGAGATcAAA	
Hu3G8VL-22K	(SEQ ID NO:24)
CGAGCTAGCTGAGATCACAGTTCTCTCTACAGTTACTGAGCACACAGGACCTCACCATGGGATGGAGCTGTATCATCCTC	
TTCTTGGTAGCAACAGCTACAGGTAAGGGGCTCACAGTAGCAGGCTTGAGGTCGGACATATATATGGGTGACAATGACA	
TCCACTTTGCCTTTCTCTCCACAGGTGTCCACTCCGACATCGTGATGACCCAATCTCCAGACTCTTTGGCTGTGTCTCTA	
GGGGAGAGGGCCACCATCAACTGCAAGTCCAGCCAAAGTGTTGATTTTGATGGTGATAGTTTATGAACTGGTACCAACA	
GAAACCAGGACAGCCACCCAACTCCTCATCTATACTACATCCAATCTAGAACTGGGGTCCCAGACAGGTTTAGTGCCA	
GTGGGTCTGGGACAGACTTCACCCCTCACCATCAGCAGCCTGCAGGCTGAGGATGTGGCAGTTTATTACTGTCAGCAAAGT	
AATTCGGATCCGTACACGTTCCGGACAGGGGACCAAGCTTGAGATcAAACGaACTGTGGCTGCACCATCGGTCTTCATCTT	
CCCGCCATCTGATGAGCAGTTGAAATCTGGAATGCCTCTGTTGTGTGCCTGCTGAATAACTTCTATCCCAGAGAGGCCA	
AAGTACAGTGGAAGGTGGATAACGCCCTCCAATCGGGTAACCTCCAGGAGAGTGTCACAGAGCAGGACAGCAAGGACAGC	
ACCTACAGCCTCAGCAGCACCCCTGACGCTGAGCAAAGCAGACTACGAGAAACACAAAGTCTACGCCTGCGAAGTCACCCA	
TCAGGGCCTGAGCTCGCCCCTCACAAGAGCTTCAACAGGGGAGAGTGTTAGTTCTAGAGTCGACTCTAGAGGATCCCCG	
GGTACCGAGCTCGAATTC	

It is understood that the examples and embodiments described herein are for illustrative purposes only and that various modifications or changes in light thereof will be suggested to persons skilled in the art and are to be included within the spirit and purview of this application and scope of the appended claims. All publications (including sequence

accession numbers and corresponding annotations), patents and patent applications cited herein are hereby incorporated by reference in their entirety for all purposes to the same extent as if each individual publication, patent or patent application were specifically and individually indicated to be so incorporated by reference.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 119	
<210> SEQ ID NO 1	
<211> LENGTH: 354	
<212> TYPE: DNA	
<213> ORGANISM: mouse	
<400> SEQUENCE: 1	
caggttactc tgaagagtc tggccctggg atattgcagc cctcccagac cctcagtctg	60
acttgttctt tctctgggtt ttcactgagg acttctggta tgggtgtagg ctggattcgt	120
cagccttcag ggaaggtct agagtggctg gcacacattt ggtgggatga tgacaagcgc	180
tataatccag ccctgaagag ccgactgaca atctccaagg atacctccag caaccaggta	240
ttcctcaaaa tcgccagtgt ggacactgca gatactgcca catactactg tgctcaaata	300
aaccccgccct ggtttgctta ctgggggcaa gggactctgg tcactgtctc tgca	354
<210> SEQ ID NO 2	
<211> LENGTH: 354	
<212> TYPE: PRT	
<213> ORGANISM: mouse	

<400> SEQUENCE: 2

<400> SEQUENCE: 3

gacactgtgc tgaccaatc tccagcttct ttggctgtgt ctctagggca gagggccacc 60

-continued

atctcctgca aggccagcca aagtgttgat tttgatggtg atagttttat gaactggtac 120
caacagaaac caggacagcc acccaaactc ctcatctata ctacatccaa tctagaatct 180
gggatcccag ccaggtttag tgccagtggg tctgggacag acttcaccct caacatccat 240
cctgtggagg aggaggatac tgcaacctat tactgtcagc aaagtaatga ggatccgtac 300
acgttcggag gggggaccaa gctggaaata aaa 333

<210> SEQ ID NO 4
<211> LENGTH: 333
<212> TYPE: PRT
<213> ORGANISM: mouse

<400> SEQUENCE: 4

Gly Ala Cys Ala Cys Thr Gly Thr Gly Cys Thr Gly Ala Cys Cys Cys
1 5 10 15
Ala Ala Thr Cys Thr Cys Cys Ala Gly Cys Thr Thr Cys Thr Thr Thr
20 25 30
Gly Gly Cys Thr Gly Thr Gly Thr Cys Thr Cys Thr Ala Gly Gly Gly
35 40 45
Cys Ala Gly Ala Gly Gly Gly Cys Cys Ala Cys Cys Ala Thr Cys Thr
50 55 60
Cys Cys Thr Gly Cys Ala Ala Gly Gly Cys Cys Ala Gly Cys Cys Ala
65 70 75 80
Ala Ala Gly Thr Gly Thr Thr Gly Ala Thr Thr Thr Thr Gly Ala Thr
85 90 95
Gly Gly Thr Gly Ala Thr Ala Gly Thr Thr Thr Thr Ala Thr Gly Ala
100 105 110
Ala Cys Thr Gly Gly Thr Ala Cys Cys Ala Ala Cys Ala Gly Ala Ala
115 120 125
Ala Cys Cys Ala Gly Gly Ala Cys Ala Gly Cys Cys Ala Cys Cys Cys
130 135 140
Ala Ala Ala Cys Thr Cys Cys Thr Cys Ala Thr Cys Thr Ala Thr Ala
145 150 155 160
Cys Thr Ala Cys Ala Thr Cys Cys Ala Ala Thr Cys Thr Ala Gly Ala
165 170 175
Ala Thr Cys Thr Gly Gly Gly Ala Thr Cys Cys Cys Ala Gly Cys Cys
180 185 190
Ala Gly Gly Thr Thr Thr Ala Gly Thr Gly Cys Cys Ala Gly Thr Gly
195 200 205
Gly Gly Thr Cys Thr Gly Gly Gly Ala Cys Ala Gly Ala Cys Thr Thr
210 215 220
Cys Ala Cys Cys Cys Thr Cys Ala Ala Cys Ala Thr Cys Cys Ala Thr
225 230 235 240
Cys Cys Thr Gly Thr Gly Gly Ala Gly Gly Ala Gly Gly Ala Gly Gly
245 250 255
Ala Thr Ala Cys Thr Gly Cys Ala Ala Cys Cys Thr Ala Thr Thr Ala
260 265 270
Cys Thr Gly Thr Cys Ala Gly Cys Ala Ala Ala Gly Thr Ala Ala Thr
275 280 285
Gly Ala Gly Gly Ala Thr Cys Cys Gly Thr Ala Cys Ala Cys Gly Thr
290 295 300
Thr Cys Gly Gly Ala Gly Gly Gly Gly Gly Gly Ala Cys Cys Ala Ala
305 310 315 320

-continued

Gly Cys Thr Gly Gly Ala Ala Ala Thr Ala Ala Ala Ala	
325	330
<div><210> SEQ ID NO 5</div> <div><211> LENGTH: 1580</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Synthetic construct</div> <div><400> SEQUENCE: 5</div>	
gctagcgttt aaacttaagc ttgttgacta gtgagatcac agttctctct acagttactg	60
agcacacagg acctcaccat gggatggagc tgtatcatcc tcttcttggt agcaacagct	120
acaggtaagg ggctcacagt agcaggcttg aggtctggac atatatatgg gtgacaatga	180
catccacttt gcctttctct ccacagggtgt ccactcccag gttaccctga aagagtctgg	240
ccctgggata ttgcagccct ccagaccct cagtctgact tgttctttct ctgggttttc	300
actgaggact tctggtatgg gtgtaggctg gattcgtcag ccttcaggga agggctctaga	360
gtggctggca cacatttggg gggatgatga caagcgctat aatccagccc tgaagagccg	420
actgacaatc tccaaggata cctccagcaa ccaggtattc ctcaaaatcg ccagtgtgga	480
cactgcagat actgccacat actactgtgc tcaaataaac cccgcctggg ttgcttactg	540
gggccaaggg actctggtea ctgtgagctc agcctccacc aagggcccat cggctctccc	600
cctggcacc cctccaaga gcacctctgg gggcacagcg gccctgggct gcctgggtcaa	660
ggactacttc cccgaaccgg tgacgggtgc gtggaactca ggcgcctga ccagcggcgt	720
gcacaccttc ccggctgtcc tacagtctc aggactctac tccctcagca gcgtgggtgac	780
cgtgccctcc agcagcttgg gcacccagac ctacatctgc aacgtgaatc acaagcccag	840
caacaccaag gtggacaaga gagttgagcc caaatcttgt gacaaaactc acacatgccc	900
accgtgccc gcacctgaac tcctgggggg accgtcagtc ttcctcttc ccccaaaacc	960
caaggacacc ctcatgatct cccggacccc tgaggtcaca tgcgtgggtg tggacgtgag	1020
ccacgaagac cctgaggtca agttcaactg gtacgtggac ggcgtggagg tgcataatgc	1080
caagacaaag ccgcgggagg agcagtacaa cagcacgtac cgtgtggtca gcgtcctcac	1140
cgtcctgcac caggactggc tgaatggcaa ggagtacaag tgcaaggtct ccaacaaagc	1200
cctcccagcc cccatcgaga aaaccatctc caaagccaaa gggcagcccc gagaaccaca	1260
ggtgtacacc ctgcccccat cccgggatga gctgaccaag aaccagggtca gcctgacctg	1320
cctgggtcaaa ggcttctatc ccagcgacat cgccgtggag tgggagagca atgggcagcc	1380
ggagaacaac tacaagacca cgctcccgt gctggactcc gacggctcct tcttctctta	1440
cagcaagctc accgtggaca agagcagggt gcagcagggg aacgtcttct catgctccgt	1500
gatgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctccgggtaa	1560
atgagtgcgg ccgcgaattc	1580
<div><210> SEQ ID NO 6</div> <div><211> LENGTH: 895</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Synthetic construct</div> <div><400> SEQUENCE: 6</div>	
gctagctgag atcacagttc tctctacagt tactgagcac acaggacctc accatgggat	60

-continued

ggagctgtat catcctcttc ttggtagcaa cagctacagg taaggggctc acagtagcag	120
gcttgaggtc tggacatata tatgggtgac aatgacatcc actttgcctt tctctccaca	180
ggtgtccact ccgacactgt gctgacccaa tctccagctt ctttggtgt gtctctaggg	240
cagagggcca ccatctcctg caaggccagc caaagtgttg attttgatgg tgatagtttt	300
atgaactggg accaacagaa accaggacag ccacccaaac tcctcatcta tactacatcc	360
aatctagaat ctgggatccc agccaggttt agtgccagtg ggtctgggac agacttcacc	420
ctcaacatcc atcctgtgga ggaggaggat actgcaacct attactgtca gcaaagtaat	480
gaggatccgt acacgttcgg aggggggacc aagcttgaga tcaaacgaac tgtggctgca	540
ccatcggtct tcctctccc gccatctgat gaggcgtga aatctggaac tgcctctgtt	600
gtgtgcctgc tgaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac	660
gccctccaat cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc	720
tacagcctca gcagcacct gacgctgagc aaagcagact acgagaaaca caaagtctac	780
gcctgcgaag tcacccatca gggcctgagc tcgcccgtca caaagagctt caacagggga	840
gagtgttagt tctagagtcg actctagagg atccccgggt accgagctcg aattc	895
<210> SEQ ID NO 7	
<211> LENGTH: 62	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 7	
ccgcgaattc tggccagggt accctgagag agtctggccc tgcgctgggt aagccacac	60
ag	62
<210> SEQ ID NO 8	
<211> LENGTH: 80	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 8	
gcgctggtga agccacaca gaccctcaca ctgacttgta ccttctctgg gttttcactg	60
agcacttctg gtatgggtgt	80
<210> SEQ ID NO 9	
<211> LENGTH: 42	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 9	
tggattcgtc agcctcccgg gaaggctcta gagggtgg ca	42
<210> SEQ ID NO 10	
<211> LENGTH: 42	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 10	
tgccagccac tctagagcct tcccgggagg ctgacgaatc ca	42

-continued

<210> SEQ ID NO 11
<211> LENGTH: 72
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 11

gtcctcaca tgaccaacat ggaccctgtg gatactgccatactactg tgetcggata 60

aacccccgcct gg 72

<210> SEQ ID NO 12
<211> LENGTH: 51
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 12

catgttggtc attgtgagga ctacctggtt tttggaggta tccttgga t 51

<210> SEQ ID NO 13
<211> LENGTH: 37
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 13

ggctgagctc acagtgacca gagtcccttg gccccag 37

<210> SEQ ID NO 14
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 14

gtgtaggctg gattcgtcag cctcccg 27

<210> SEQ ID NO 15
<211> LENGTH: 33
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Primer

<400> SEQUENCE: 15

gacgaatcca gcctacacc ataccagaag tgc 33

<210> SEQ ID NO 16
<211> LENGTH: 354
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 16

caggttaccc tgagagagtc tggccctgctg ctggtgaagc ccacacagac cctcacactg 60

acttgtaacct tctctgggtt ttcactgagc acttctggta tgggtgtagg ctggattcgt 120

cagcctcccg ggaaggctct agagtggctg gcacacattt ggtgggatga tgacaagcgc 180

-continued

tataatccag ccctgaagag ccgactgaca atctccaagg atacctccaa aaaccaggta	240
gtcctcacia tgaccaacat ggaccctgtg gatactgcca catactactg tgetcggata	300
aaccccgccct ggtttgctta ctggggccaa gggactctgg tcactgtgag ctca	354
<210> SEQ ID NO 17	
<211> LENGTH: 1580	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic construct	
<400> SEQUENCE: 17	
gctagcgttt aaacttaagc ttgttgacta gtgagatcac agttctctct acagttactg	60
agcacacagg acctcaccat gggatggagc tgtatcatcc tcttcttggt agcaacagct	120
acaggtaagg ggctcacagt agcaggcttg aggtctggac atatatatgg gtgacaatga	180
catccacttt gcctttctct ccacagggtg ccactcccag gttaccctga gagagtctgg	240
ccctgcgctg gtgaagccca cacagaccct cacactgact tgtaccttct ctgggttttc	300
actgagcact tctggtatgg gtgtaggctg gattcgtcag cctcccggga aggctctaga	360
gtggctggca cacatttggt gggatgatga caagcgctat aatccagccc tgaagagccg	420
actgacaatc tccaaggata cctccaaaaa ccaggtagtc ctcacaatga ccaacatgga	480
ccctgtggat actgccacat actactgtgc tcggataaac cccgcctggg ttgettactg	540
gggccaaggg actctggcca ctgtgagctc agcctccacc aaggggccat cggctctccc	600
cctggcacc cctccaaga gcacctctgg gggcacagcg gccctgggct gcctgggtcaa	660
ggactacttc cccgaaccgg tgacgggtgc gtggaactca ggcgcctga ccagcggcgt	720
gcacaccttc ccggctgtcc tacagtctc aggactctac tccctcagca gcgtgggtgac	780
cgtgccctcc agcagcttgg gcacccagac ctacatctgc aacgtgaatc acaagcccag	840
caacaccaag gtggacaaga gagttgagcc caaatcttgt gacaaaactc acacatgccc	900
accgtgccca gcacctgaac tctgggggg accgtcagtc ttctcttcc ccccaaaacc	960
caaggacacc ctcatgatct cccggacccc tgaggtcaca tgcgtgggtg tggacgtgag	1020
ccacgaagac cctgagggtca agttcaactg gtacgtggac ggcgtggagg tgcataatgc	1080
caagacaaag ccgcgggagg agcagtacaa cagcacgtac cgtgtgggtca gcgtcctcac	1140
cgtcctgcac caggactggc tgaatggcaa ggagtacaag tgcaagggtc ccaacaaagc	1200
cctcccagcc cccatcgaga aaaccatctc caaagccaaa gggcagcccc gagaaccaca	1260
ggtgtacacc ctgcccccat cccgggatga gctgaccaag aaccagggtca gcctgacctg	1320
cctgggtcaaa ggcttctatc ccagcgacat cgcctgggag tgggagagca atgggcagcc	1380
ggagaacaac tacaagacca cgctcccgt gctggactcc gacggctcct tcttctctta	1440
cagcaagctc accgtggaca agagcagggtg gcagcagggg aacgtcttct catgctccgt	1500
gatgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctccgggtaa	1560
atgagtgccg ccgcgaattc	1580
<210> SEQ ID NO 18	
<211> LENGTH: 63	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Primer	
<400> SEQUENCE: 18	

-continued

actcttttggc tgtgtctcta ggggagaggg ccaccatcaa ctgcaaggcc agccaaagtg	60
ttg	63
<div><210> SEQ ID NO 19</div> <div><211> LENGTH: 66</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Primer</div>	
<400> SEQUENCE: 19	
ctctccacag gtgtccactc cgacatcgtg atgacccaat ctccagactc tttggctgtg	60
tctcta	66
<div><210> SEQ ID NO 20</div> <div><211> LENGTH: 71</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Primer</div>	
<400> SEQUENCE: 20	
ggtgaggggtg aagtctgtcc cagacccact gccactaaac ctgtctggga ccccagattc	60
tagattggat g	71
<div><210> SEQ ID NO 21</div> <div><211> LENGTH: 67</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Plasmid</div>	
<400> SEQUENCE: 21	
tgacagtaat aaactgccac atcctcagcc tgcaggctgc tgatggtgag ggtgaagtct	60
gtcccag	67
<div><210> SEQ ID NO 22</div> <div><211> LENGTH: 71</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Plasmid</div>	
<400> SEQUENCE: 22	
gcggcaagct tgggtccctg tccgaacgtg tacggatcct cattactttg ctgacagtaa	60
taaactgcc a	71
<div><210> SEQ ID NO 23</div> <div><211> LENGTH: 30</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Plasmid</div>	
<400> SEQUENCE: 23	
cgagctagct gagatcacag ttctctctac	30
<div><210> SEQ ID NO 24</div> <div><211> LENGTH: 898</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div>	

-continued

<223> OTHER INFORMATION: Synthetic Construct		
<400> SEQUENCE: 24		
cgagctagct gagatcacag ttctctctac agttactgag cacacaggac ctcaccatgg	60	
gatggagctg tatcatcctc ttcttggttag caacagctac aggtaagggg ctcacagtag	120	
caggcttgag gtctggacat atatatgggt gacaatgaca tccactttgc ctttctctcc	180	
acaggtgtcc actccgacat cgtgatgacc caatctccag actctttggc tgtgtctcta	240	
ggggagaggg ccaccatcaa ctgcaagtcc agccaaagtg ttgattttga tggatgatgt	300	
tttatgaact ggtaccaaca gaaaccagga cagccacca aactcctcat ctatactaca	360	
tccaatctag aaactggggc cccagacagg tttagtggca gtgggtctgg gacagacttc	420	
accctcacca tcagcagcct gcaggctgag gatgtggcag tttattactg tcagcaaagt	480	
aattcggatc cgtacacgtt cggacagggg accaagcttg agatcaaacg aactgtggct	540	
gcaccatcgg tcttcatctt cccgccatct gatgagcagt tgaaatctgg aactgcctct	600	
gttgtgtgcc tgctgaataa cttctatccc agagaggcca aagtacagtg gaaggtggat	660	
aacgccctcc aatcgggtaa ctcccaggag agtgtcacag agcaggacag caaggacagc	720	
acctacagcc tcagcagcac cctgacgtg agcaaagcag actacgagaa acacaaagtc	780	
tacgcctgcg aagtcacca tcagggcctg agctcgcccg tcacaaagag cttcaacagg	840	
ggagagtgtt agttctagag tcgactctag aggatccccg ggtaccgagc tcgaattc	898	
<210> SEQ ID NO 25		
<211> LENGTH: 333		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic Construct		
<400> SEQUENCE: 25		
gacactgtgc tgacccaatc tocagcttct ttggctgtgt ctctagggca gagggccacc	60	
atctcctgca aggccagcca aagtgttgat tttgatggtg atagttttat gaactggtac	120	
caacagaaac caggacagcc acccaaactc ctcatctata ctacatcaa tctagaatct	180	
gggatcccag ccaggtttag tgccagtggg tctgggacag acttcaccct caacatccat	240	
cctgtggagg aggaggatac tgcaacctat tactgtcagc aaagtaatga ggatccgtac	300	
acgttcggag gggggaccaa gcttgagatc aaa	333	
<210> SEQ ID NO 26		
<211> LENGTH: 895		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic Construct		
<400> SEQUENCE: 26		
gctagctgag atcacagttc tctctacagt tactgagcac acaggacctc accatgggat	60	
ggagctgtat catcctcttc ttggtagcaa cagctacagg taaggggctc acagtagcag	120	
gcttgaggtc tggacatata tatgggtgac aatgacatcc actttgcctt tctctccaca	180	
ggtgtccact ccgacactgt gctgaccaa tctccagctt ctttggtgt gtctctaggg	240	
cagagggcca ccatctcctg caaggccagc caaagtgttg attttgatgg tgatagtttt	300	
atgaactggg accaacagaa accaggacag ccacccaaac tcctcatcta tactacatcc	360	
aatctagaat ctgggatccc agccaggttt agtgccagtg ggtctgggac agacttcacc	420	

-continued

ctcaacatcc atcctgtgga ggaggaggat actgcaacct attactgtca gcaaagtaat 480
gaggatccgt acacgttcgg aggggggacc aagcttgaga tcaaacgaac tgtggctgca 540
ccatcggtct tcattctccc gccatctgat gagcagttga aatctggaac tgctctgtt 600
gtgtgcctgc tgaataactt ctatcccaga gaggccaaag tacagtggaa ggtggataac 660
gccctccaat cgggtaactc ccaggagagt gtcacagagc aggacagcaa ggacagcacc 720
tacagcctca gcagcaccct gacgctgagc aaagcagact acgagaaaca caaagtctac 780
gcctgcgaag tcacccatca gggcctgagc tcgcccgtca caaagagctt caacagggga 840
gagtgttagt tctagagtcg actctagagg atccccgggt accgagctcg aattc 895

<210> SEQ ID NO 27
<211> LENGTH: 35
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 27

gttggatcct ccaactgctc tgctacttct agttt 35

<210> SEQ ID NO 28
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 28

gaaaagctta aagaatgatg agatggttga cact 34

<210> SEQ ID NO 29
<211> LENGTH: 210
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 29

Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Val Ser Ala
1 5 10 15
Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro
20 25 30
Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr Leu Lys Cys Gln
35 40 45
Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp Phe His Asn Glu
50 55 60
Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile Asp Ala Ala Thr
65 70 75 80
Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn Leu Ser Thr Leu
85 90 95
Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln
100 105 110
Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile His Leu Arg Cys
115 120 125
His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr Tyr Leu Gln Asn
130 135 140
Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp Phe Tyr Ile Pro

-continued

145	150	155	160
Lys Ala Thr Leu	Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Phe		
	165	170	175
Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln			
	180	185	190
Gly Leu Ala Val Ser Thr Ile Ser Ser Phe Phe Lys Leu Ala Ala Ala			
	195	200	205
Arg Val			
210			
<210> SEQ ID NO 30			
<211> LENGTH: 30			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic Construct			
<400> SEQUENCE: 30			
Gln Val Thr Leu Lys Glu Ser Gly Pro Gly Ile Leu Gln Pro Ser Gln			
1	5	10	15
Thr Leu Ser Leu Thr Cys Ser Phe Ser Gly Phe Ser Leu Arg			
	20	25	30
<210> SEQ ID NO 31			
<211> LENGTH: 30			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic Construct			
<400> SEQUENCE: 31			
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln			
1	5	10	15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser			
	20	25	30
<210> SEQ ID NO 32			
<211> LENGTH: 30			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic Construct			
<400> SEQUENCE: 32			
Gln Val Thr Leu Lys Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln			
1	5	10	15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser			
	20	25	30
<210> SEQ ID NO 33			
<211> LENGTH: 30			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic Construct			
<400> SEQUENCE: 33			
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln			
1	5	10	15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Arg			
	20	25	30

-continued

```
<210> SEQ ID NO 34
<211> LENGTH: 30
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 34
Gln Ile Thr Leu Lys Glu Ser Gly Pro Thr Leu Val Lys Pro Thr Gln
 1             5             10            15
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser
      20             25             30

<210> SEQ ID NO 35
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 35
Thr Ser Gly Met Gly Val Gly
 1             5

<210> SEQ ID NO 36
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 36
Thr Ser Gly Val Gly Val Gly
 1             5

<210> SEQ ID NO 37
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 37
Trp Ile Arg Gln Pro Ser Gly Lys Gly Leu Glu Trp Leu Ala
 1             5             10

<210> SEQ ID NO 38
<211> LENGTH: 14
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 38
Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu Trp Leu Ala
 1             5             10

<210> SEQ ID NO 39
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 39
His Ile Trp Trp Asp Asp Lys Arg Tyr Asn Pro Ala Leu Lys Ser
 1             5             10            15
```

-continued

<210> SEQ ID NO 40
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 40

His Ile Tyr Trp Asn Asp Asp Lys Arg Tyr Asn Pro Ala Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 41
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 41

His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 42
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 42

His Ile Tyr Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 43
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 43

His Ile Trp Trp Asn Asp Asp Lys Arg Tyr Asn Pro Ala Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 44
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 44

Leu Ile Asp Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 45
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 45

His Ile Phe Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
1 5 10 15

-continued

<210> SEQ ID NO 46
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 46

Leu Ile Trp Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 47
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 47

His Ile Asp Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 48
<211> LENGTH: 16
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 48

Leu Ile Trp Trp Asn Asp Asp Lys Arg Tyr Ser Pro Ser Leu Lys Ser
1 5 10 15

<210> SEQ ID NO 49
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 49

Arg Leu Thr Ile Ser Lys Asp Thr Ser Ser Asn Gln Val Phe Leu Lys
1 5 10 15

Ile Ala Ser Val Asp Thr Ala Asp Thr Ala Thr Tyr Tyr Cys Ala Gln
20 25 30

<210> SEQ ID NO 50
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 50

Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val Val Leu Thr
1 5 10 15

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Arg
20 25 30

<210> SEQ ID NO 51
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 51

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Gln
20 25 30

```
<210> SEQ ID NO 52
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 52

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Thr
20 25 30

```
<210> SEQ ID NO 53
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 53

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Lys
20 25 30

```
<210> SEQ ID NO 54
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 54

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Ala
20 25 30

```
<210> SEQ ID NO 55
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 55

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala His
20 25 30

```
<210> SEQ ID NO 56
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```


-continued

<400> SEQUENCE: 56

Arg Leu Thr Ile Thr Lys Asp Thr Ser Lys Asn Gln Val Val Leu Thr
1 5 10 15

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Arg
20 25 30

<210> SEQ ID NO 57
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 57

Arg Leu Thr Ile Thr Lys Asp Thr Ser Lys Asn Gln Val Val Leu Thr
1 5 10 15

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala His
20 25 30

<210> SEQ ID NO 58
<211> LENGTH: 32
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 58

Arg Leu Thr Ile Thr Lys Asp Thr Ser Lys Asn Gln Val Val Leu Thr
1 5 10 15

Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr Cys Ala Gln
20 25 30

<210> SEQ ID NO 59
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 59

Ile Asn Pro Ala Trp Phe Ala Tyr
1 5

<210> SEQ ID NO 60
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 60

Ile Asn Pro Ala Trp Phe Asp Tyr
1 5

<210> SEQ ID NO 61
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 61

Ile Asn Pro Ala Tyr Phe Ala Tyr
1 5

-continued

<210> SEQ ID NO 62
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 62

Ile Asn Pro Ala Tyr Phe Asp Tyr
1 5

<210> SEQ ID NO 63
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 63

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ala
1 5 10

<210> SEQ ID NO 64
<211> LENGTH: 11
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 64

Trp Gly Gln Gly Thr Leu Val Thr Val Ser Ser
1 5 10

<210> SEQ ID NO 65
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 65

Asp Thr Val Leu Thr Gln Ser Pro Ala Ser Leu Ala Val Ser Leu Gly
1 5 10 15
Gln Arg Ala Thr Ile Ser Cys
20

<210> SEQ ID NO 66
<211> LENGTH: 23
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 66

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15
Glu Arg Ala Thr Ile Asn Cys
20

<210> SEQ ID NO 67
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

-continued

<400> SEQUENCE: 67

Lys Ala Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Phe Met Asn
1 5 10 15

<210> SEQ ID NO 68
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 68

Arg Ala Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Phe Met Asn
1 5 10 15

<210> SEQ ID NO 69
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 69

Lys Ser Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Phe Met Asn
1 5 10 15

<210> SEQ ID NO 70
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 70

Lys Ala Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Tyr Met Asn
1 5 10 15

<210> SEQ ID NO 71
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 71

Lys Ala Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Phe Leu Asn
1 5 10 15

<210> SEQ ID NO 72
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 72

Lys Ala Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Phe Met Ala
1 5 10 15

<210> SEQ ID NO 73
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 73

Lys Ala Ser Gln Ser Val Asp Phe Asp Gly Asp Ser Tyr Leu Ala
1 5 10 15

```
<210> SEQ ID NO 74
<211> LENGTH: 15
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 74

Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro Lys Leu Leu Ile Tyr
1 5 10 15

```
<210> SEQ ID NO 75
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 75

Thr Thr Ser Asn Leu Glu Ser
1 5

```
<210> SEQ ID NO 76
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 76

Asp Ala Ser Asn Leu Glu Ser
1 5

```
<210> SEQ ID NO 77
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 77

Trp Ala Ser Asn Leu Glu Ser
1 5

```
<210> SEQ ID NO 78
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 78

Thr Thr Ser Asn Leu Glu Thr
1 5

```
<210> SEQ ID NO 79
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 79

-continued

Asp Ala Ser Asn Leu Glu Thr
1 5

<210> SEQ ID NO 80
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 80

Asp Ala Ser Asn Leu Ala Thr
1 5

<210> SEQ ID NO 81
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 81

Ser Ala Ser Asn Leu Gln Ser
1 5

<210> SEQ ID NO 82
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 82

Ser Thr Ser Asn Leu Glu Ser
1 5

<210> SEQ ID NO 83
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 83

Ser Thr Ser Asn Leu Gln Ser
1 5

<210> SEQ ID NO 84
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 84

Thr Thr Ser Asn Leu Gln Ser
1 5

<210> SEQ ID NO 85
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 85

Thr Thr Ser Ser Leu Gln Ser
1 5

Gly Ile Pro Ala Arg Phe Ser Ala Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15

Leu Asn Ile His Pro Val Glu Glu Asp Thr Ala Thr Tyr Tyr Cys
20 25 30

Gly Val Pro Asp Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr
1 5 10 15
Leu Thr Ile Ser Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys
20 25 30

Gln Gln Ser Asn Glu Asp Pro Tyr Thr
1 5

Gln Gln Ser Tyr Ser Thr Pro Tyr Thr
1 5

Gln Gln Ser Tyr Glu Asp Pro Tyr Thr
1 5

```
<210> SEQ ID NO 91
<211> LENGTH: 9
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
```


-continued

<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
<400> SEQUENCE: 91	
Gln Gln Ser Asn Ser Asp Pro Tyr Thr	
1 5	
<210> SEQ ID NO 92	
<211> LENGTH: 9	
<212> TYPE: PRT	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
<400> SEQUENCE: 92	
Gln Gln Ser Asn Glu Thr Pro Tyr Thr	
1 5	
<210> SEQ ID NO 93	
<211> LENGTH: 10	
<212> TYPE: PRT	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
<400> SEQUENCE: 93	
Phe Gly Gly Gly Thr Lys Leu Glu Ile Lys	
1 5 10	
<210> SEQ ID NO 94	
<211> LENGTH: 10	
<212> TYPE: PRT	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
<400> SEQUENCE: 94	
Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys	
1 5 10	
<210> SEQ ID NO 95	
<211> LENGTH: 528	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
<400> SEQUENCE: 95	
cgagctagct gagatcacag ttctctctac agttactgag cacacaggac ctcaccatgg	60
gatggagctg tatcatcctc ttcttggtag caacagctac aggtaagggg ctcacagtag	120
caggcttgag gtctggacat atatatgggt gacaatgaca tccactttgc ctttctctcc	180
acaggtgtcc actccgacat cgtgatgacc caatctccag actctttggc tgtgtctcta	240
ggggagaggg ccaccatcaa ctgcaaggcc agccaaagtg ttgattttga tggatgatgt	300
tttatgaact ggtaccaaca gaaaccagga cagccacca aactcctcat ctatactaca	360
tccaatctag aatctggggt cccagacagg tttagtggca gtgggtctgg gacagacttc	420
accctcacca tcagcagcct gcaggctgag gatgtggcag tttattactg tcagcaaagt	480
aatgaggatc cgtacacgtt cggacagggg accaagcttg agatcaaa	528
<210> SEQ ID NO 96	
<211> LENGTH: 111	
<212> TYPE: PRT	

-continued

<213> ORGANISM: Artificial Sequence															
<220> FEATURE:															
<223> OTHER INFORMATION: Synthetic Construct															
<400> SEQUENCE: 96															
Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	Leu	Gly
1				5					10					15	
Glu	Arg	Ala	Thr	Ile	Asn	Cys	Lys	Ala	Ser	Gln	Ser	Val	Asp	Phe	Asp
			20					25					30		
Gly	Asp	Ser	Phe	Met	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro
		35					40					45			
Lys	Leu	Leu	Ile	Tyr	Thr	Thr	Ser	Asn	Leu	Glu	Ser	Gly	Val	Pro	Asp
	50					55					60				
Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser
65					70					75					80
Ser	Leu	Gln	Ala	Glu	Asp	Val	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Ser	Asn
				85					90					95	
Glu	Asp	Pro	Tyr	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	
			100					105					110		
<210> SEQ ID NO 97															
<211> LENGTH: 897															
<212> TYPE: DNA															
<213> ORGANISM: Artificial Sequence															
<220> FEATURE:															
<223> OTHER INFORMATION: Synthetic Construct															
<400> SEQUENCE: 97															
cgagctagct	gagatcacag	ttctctctac	agttactgag	cacacaggac	ctcaccatgg										60
gatggagctg	tatcatcctc	ttcttggtag	caacagctac	aggtaagggg	ctcacagtag										120
caggcttgag	gtctggacat	atatatgggt	gacaatgaca	tccactttgc	ctttctctcc										180
acaggtgtcc	actccgacat	cgtgatgacc	caatctccag	actctttggc	tgtgtctcta										240
ggggagaggg	ccaccatcaa	ctgcaaggcc	agccaaagtg	ttgattttga	tggtgatagt										300
tttatgaact	ggtaccaaca	gaaaccagga	cagccacca	aactcctcat	ctatactaca										360
tccaatctag	aatctgggg	cccagacagg	tttagtggca	gtgggtctgg	gacagacttc										420
accctcacca	tcagcagcct	gcaggctgag	gatgtggcag	tttattactg	tcagcaaagt										480
aatgaggatc	cgtacacgtt	cggacagggg	accaagcttg	agatcaaacg	aactgtggct										540
gcaccatcgg	tcttcatctt	cccgccatct	gatgagcagt	tgaaatctgg	aactgcctct										600
gttgtgtgcc	tgctgaataa	cttctatccc	agagaggcca	aagtacagtg	gaagggtggat										660
aacgccctcc	aatcgggtaa	ctcccaggag	agtgtcacag	agcaggacag	caaggacagc										720
acctacagcc	tcagcagcac	cctgacgctg	agcaaagcag	actacgagaa	acacaaagtc										780
tacgtgcga	agtcacccat	cagggcctga	gtcgcgccgt	cacaaagagc	ttcaacaggg										840
gagagtgtta	gttctagagt	cgactctaga	ggatccccgg	gtaccgagct	cgaattc										897
<210> SEQ ID NO 98															
<211> LENGTH: 218															
<212> TYPE: PRT															
<213> ORGANISM: Artificial Sequence															
<220> FEATURE:															
<223> OTHER INFORMATION: Synthetic Construct															
<400> SEQUENCE: 98															
Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	Leu	Gly
1				5					10					15	

-continued

Glu Arg Ala Thr Ile Asn Cys Lys Ala Ser Gln Ser Val Asp Phe Asp
20 25 30

Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45

Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Ser Gly Val Pro Asp
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
85 90 95

Glu Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
145 150 155 160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
165 170 175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
180 185 190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
195 200 205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

<210> SEQ ID NO 99
<211> LENGTH: 528
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 99

cgagctagct gagatcacag ttctctctac agttactgag cacacaggac ctcaccatgg 60

gatggagctg tatcatcctc ttcttggttag caacagctac aggtaagggg ctcacagtag 120

caggcttgag gtctggacat atatatgggt gacaatgaca tccactttgc ctttctctcc 180

acaggtgtcc actccgacat cgtgatgacc caatctccag actctttggc tgtgtctcta 240

ggggagaggg ccaccatcaa ctgcaagtec agccaaagtg ttgattttga tggatgatag 300

tttatgaact ggtaccaaca gaaaccagga cagccacca aactcctcat ctatactaca 360

tccagtctag aatctggggg cccagacagg tttagtggca gtgggtctgg gacagacttc 420

accctcacca tcagcagcct gcaggctgag gatgtggcag tttattactg tcagcaaagt 480

aattcggatc cgtacacgtt cggacagggg accaagcttg agatcaaa 528

<210> SEQ ID NO 100
<211> LENGTH: 111
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 100

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly

-continued

1	5	10	15
Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Asp Phe Asp			
20	25	30	
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro			
35	40	45	
Lys Leu Leu Ile Tyr Thr Thr Ser Ser Leu Glu Ser Gly Val Pro Asp			
50	55	60	
Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser			
65	70	75	80
Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn			
85	90	95	
Ser Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys			
100	105	110	
<210> SEQ ID NO 101			
<211> LENGTH: 898			
<212> TYPE: DNA			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic Construct			
<400> SEQUENCE: 101			
cgagctagct gagatcacag ttctctctac agttactgag cacacaggac ctcaccatgg	60		
gatggagctg tatcatcctc ttcttggttag caacagctac aggtaagggg ctcacagtag	120		
caggcttgag gtctggacat atatatgggt gacaatgaca tccactttgc ctttctctcc	180		
acagggtgcc actccgacat cgtgatgacc caatctccag actctttggc tgtgtctcta	240		
ggggagaggg ccaccatcaa ctgcaagtcc agccaaagtg ttgattttga tggatgatagt	300		
tttatgaact ggtaccaaca gaaaccagga cagccacca aactcctcat ctatactaca	360		
tccagtctag aatctggggg cccagacagg tttagtggca gtgggtctgg gacagacttc	420		
accctcacca tcagcagcct gcaggctgag gatgtggcag tttattactg tcagcaaagt	480		
aattcggatc cgtacacgtt cggacagggg accaagcttg agatcaaacg aactgtggct	540		
gcaccatcgg tcttcatctt cccgccatct gatgagcagt tgaaatctgg aactgcctct	600		
gttgtgtgcc tgctgaataa cttctatccc agagaggcca agtacagtg gaaggtggat	660		
aacgccctcc aatcgggtaa ctcccaggag agtgtcacag agcaggacag caaggacagc	720		
acctacagcc tcagcagcac cctgacgctg agcaaagcag actacgagaa acacaaagtc	780		
tacgcctgcg aagtcacca tcagggcctg agctcgcccg tcacaaagag cttcaacagg	840		
ggagagtgtt agttctagag togactctag aggatccccg ggtaccgagc togaattc	898		
<210> SEQ ID NO 102			
<211> LENGTH: 218			
<212> TYPE: PRT			
<213> ORGANISM: Artificial Sequence			
<220> FEATURE:			
<223> OTHER INFORMATION: Synthetic Construct			
<400> SEQUENCE: 102			
Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly			
1	5	10	15
Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Asp Phe Asp			
20	25	30	
Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro			
35	40	45	

-continued

Lys	Leu	Leu	Ile	Tyr	Thr	Thr	Ser	Ser	Leu	Glu	Ser	Gly	Val	Pro	Asp	
50						55					60					
Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	
65					70					75					80	
Ser	Leu	Gln	Ala	Glu	Asp	Val	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Ser	Asn	
				85					90					95		
Ser	Asp	Pro	Tyr	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys	Arg	
			100					105					110			
Thr	Val	Ala	Ala	Pro	Ser	Val	Phe	Ile	Phe	Pro	Pro	Ser	Asp	Glu	Gln	
		115					120					125				
Leu	Lys	Ser	Gly	Thr	Ala	Ser	Val	Val	Cys	Leu	Leu	Asn	Asn	Phe	Tyr	
130						135					140					
Pro	Arg	Glu	Ala	Lys	Val	Gln	Trp	Lys	Val	Asp	Asn	Ala	Leu	Gln	Ser	
145					150					155					160	
Gly	Asn	Ser	Gln	Glu	Ser	Val	Thr	Glu	Gln	Asp	Ser	Lys	Asp	Ser	Thr	
			165					170						175		
Tyr	Ser	Leu	Ser	Ser	Thr	Leu	Thr	Leu	Ser	Lys	Ala	Asp	Tyr	Glu	Lys	
		180						185					190			
His	Lys	Val	Tyr	Ala	Cys	Glu	Val	Thr	His	Gln	Gly	Leu	Ser	Ser	Pro	
	195						200					205				
Val	Thr	Lys	Ser	Phe	Asn	Arg	Gly	Glu	Cys							
210						215										
<210> SEQ ID NO 103																
<211> LENGTH: 571																
<212> TYPE: DNA																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Synthetic Construct																
<400> SEQUENCE: 103																
gctagcgttt aaacttaagc ttgttgacta gtgagatcac agttctctct acagttactg															60	
agcacacagg acctcaccat gggatggagc tgtatcatcc tcttcttggt agcaacagct															120	
acaggtaagg ggctcacagt agcaggcttg aggtctggac atatatatgg gtgacaatga															180	
catccacttt gcctttctct ccacaggtgt ccactcccag gttaccctga gagagtctgg															240	
ccctgcgctg gtgaagccca cacagaccct cacactgact tgtaccttct ctgggttttc															300	
actgagcact tctggtatgg gtgtaggctg gattcgtcag cctcccggga aggctctaga															360	
gtggctggca cacatttggt gggatgatga caagcgctat aatccagccc tgaagagccg															420	
actgacaatc tccaaggata cctccaaaaa ccaggtagtc ctcacaatga ccaacatgga															480	
ccctgtggat actgccacat actactgtgc tcggataaac cccgcctggt ttgcttactg															540	
gggccaaggg actctgggtca ctgtgagctc a															571	
<210> SEQ ID NO 104																
<211> LENGTH: 118																
<212> TYPE: PRT																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Synthetic Construct																
<400> SEQUENCE: 104																
Gln	Val	Thr	Leu	Arg	Glu	Ser	Gly	Pro	Ala	Leu	Val	Lys	Pro	Thr	Gln	
1				5					10					15		
Thr	Leu	Thr	Leu	Thr	Cys	Thr	Phe	Ser	Gly	Phe	Ser	Leu	Ser	Thr	Ser	
		20					25					30				

-continued

[illegible]

```
<210> SEQ ID NO 105
<211> LENGTH: 1580
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct
```

<400> SEQUENCE: 105

gctagcgcttt	aaacttaagc	ttgttgacta	gtgagatcac	agttctctct	acagttactg	60
agcacacagg	acctcaccat	gggatggagc	tgtatcatcc	tcttcttggt	agcaacagct	120
acaggtaaagg	ggctcacagt	agcaggcttg	aggctctggac	atatatatgg	gtgacaatga	180
catccacttt	gcctttctct	ccacagggtg	ccactoccag	gttaccctga	gagagtctgg	240
ccctgcgctg	gtgaagccca	cacagaccct	cacactgact	tgtaccttct	ctggggtttc	300
actgagcact	tctggtatgg	gtgtaggctg	gattcgctcag	cctcccggga	aggctctaga	360
gtggctggca	cacatttggt	gggatgatga	caagcgctat	aatccagccc	tgaagagccg	420
actgacaatc	tccaaggata	cctccaaaaa	ccaggtagtc	ctcacaatga	ccaacatgga	480
ccctgtggat	actgccacat	actactgtgc	tgggataaac	cccgcctggg	ttgcttactg	540
gggccaaagg	actctggtca	ctgtgagctc	agcctccacc	aaggggcccat	cggctcttccc	600
cctggcaccc	tcctccaaga	gcacctctgg	gggcacagcg	gccttgggct	gcctgggtcaa	660
ggactacttc	ccgaaccgg	tgacgggtgc	gtggaactca	ggcgccctga	ccagcggcgt	720
gcacaccttc	ccggctgtcc	tacagtcctc	aggactctac	tccttcagca	gcgtgggtgac	780
cgtgccctcc	agcagcttgg	gcaccagac	ctacatctgc	aacgtgaatc	acaagcccag	840
caacaccaag	gtggacaaga	gagttgagcc	caaatcttgt	gacaaaactc	acacatgccc	900
accgtgccc	gcacctgaac	tcctgggggg	accgtcagtc	ttctcttcc	ccccaaaacc	960
caaggacacc	ctcatgatct	cccgacccc	tgaggtcaca	tgcgtgggtg	tggacgtgag	1020
ccacgaagac	cctgagggtca	agttcaactg	gtacgtggac	ggcgtggagg	tgcataatgc	1080
caagacaaa	ccgcggggag	agcagtacaa	cagcacgtac	cgtgtgggtca	gcgtcctcac	1140
cgtcctgcac	caggactggc	tgaatggcaa	ggagtacaag	tgcaagggtct	ccaacaaagc	1200
cctcccagcc	cccacgaga	aaaccatctc	caaagccaaa	gggcagcccc	gagaaccaca	1260
ggtgtacacc	ctgcccccat	ccggggatga	gctgaccaag	aaccagggtca	gcctgacctg	1320
cctgggtcaaa	ggcttctatc	ccagcgacat	cgcctgggag	tgggagagca	atgggcagcc	1380
ggagaacaac	tacaagacca	cgctcccgt	gctggactcc	gacggctcct	tcttctctta	1440
cagcaagctc	accgtggaca	agagcagggtg	gcagcagggg	aacgtcttct	catgctccgt	1500
gatgcatgag	gctctgcaca	accactacac	gcagaagagc	ctctccctgt	ctccgggttaa	1560

-continued

atgagtgcgg ccgcgaattc	1580
<div><210> SEQ ID NO 106</div> <div><211> LENGTH: 528</div> <div><212> TYPE: DNA</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Synthetic Construct</div> <div><400> SEQUENCE: 106</div>	
cgagctagct gagatcacag ttctctctac agttactgag cacacaggac ctcaccatgg	60
gatggagctg tatcatcctc ttcttggttag caacagctac aggtaagggg ctcacagtag	120
caggcttgag gtctggacat atatatgggt gacaatgaca tccactttgc ctttctctcc	180
acaggtgtcc actccgacat cgtgatgacc caatctccag actctttggc tgtgtctcta	240
ggggagaggg ccaccatcaa ctgcaagtcc agccaaagtg ttgattttga tggatgatagt	300
tttatgaact ggtaccaaca gaaaccagga cagccacca aactcctcat ctatactaca	360
tccaatctag aaactggggt cccagacagg tttagtggca gtgggtctgg gacagacttc	420
accctcacca tcagcagcct gcaggctgag gatgtggcag tttattactg tcagcaaagt	480
aattcggatc cgtacacgtt cggacagggg accaagcttg agatcaaa	528
<div><210> SEQ ID NO 107</div> <div><211> LENGTH: 448</div> <div><212> TYPE: PRT</div> <div><213> ORGANISM: Artificial Sequence</div> <div><220> FEATURE:</div> <div><223> OTHER INFORMATION: Synthetic Construct</div> <div><400> SEQUENCE: 107</div>	
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln	
1 5 10 15	
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser	
20 25 30	
Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu	
35 40 45	
Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala	
50 55 60	
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val	
65 70 75 80	
Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr	
85 90 95	
Cys Ala Arg Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr	
100 105 110	
Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro	
115 120 125	
Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly	
130 135 140	
Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn	
145 150 155 160	
Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln	
165 170 175	
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser	
180 185 190	
Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser	
195 200 205	

Asn	Thr	Lys	Val	Asp	Lys	Arg	Val	Glu	Pro	Lys	Ser	Cys	Asp	Lys	Thr
210						215					220				
His	Thr	Cys	Pro	Pro	Cys	Pro	Ala	Pro	Glu	Leu	Leu	Gly	Gly	Pro	Ser
225					230					235					240
Val	Phe	Leu	Phe	Pro	Pro	Lys	Pro	Lys	Asp	Thr	Leu	Met	Ile	Ser	Arg
				245					250					255	
Thr	Pro	Glu	Val	Thr	Cys	Val	Val	Val	Asp	Val	Ser	His	Glu	Asp	Pro
			260					265					270		
Glu	Val	Lys	Phe	Asn	Trp	Tyr	Val	Asp	Gly	Val	Glu	Val	His	Asn	Ala
		275					280					285			
Lys	Thr	Lys	Pro	Arg	Glu	Glu	Gln	Tyr	Asn	Ser	Thr	Tyr	Arg	Val	Val
290						295					300				
Ser	Val	Leu	Thr	Val	Leu	His	Gln	Asp	Trp	Leu	Asn	Gly	Lys	Glu	Tyr
305					310					315					320
Lys	Cys	Lys	Val	Ser	Asn	Lys	Ala	Leu	Pro	Ala	Pro	Ile	Glu	Lys	Thr
				325					330					335	
Ile	Ser	Lys	Ala	Lys	Gly	Gln	Pro	Arg	Glu	Pro	Gln	Val	Tyr	Thr	Leu
			340					345					350		
Pro	Pro	Ser	Arg	Asp	Glu	Leu	Thr	Lys	Asn	Gln	Val	Ser	Leu	Thr	Cys
		355					360					365			
Leu	Val	Lys	Gly	Phe	Tyr	Pro	Ser	Asp	Ile	Ala	Val	Glu	Trp	Glu	Ser
370						375					380				
Asn	Gly	Gln	Pro	Glu	Asn	Asn	Tyr	Lys	Thr	Thr	Pro	Pro	Val	Leu	Asp
385					390					395					400
Ser	Asp	Gly	Ser	Phe	Phe	Leu	Tyr	Ser	Lys	Leu	Thr	Val	Asp	Lys	Ser
				405					410					415	
Arg	Trp	Gln	Gln	Gly	Asn	Val	Phe	Ser	Cys	Ser	Val	Met	His	Glu	Ala
			420					425					430		
Leu	His	Asn	His	Tyr	Thr	Gln	Lys	Ser	Leu	Ser	Leu	Ser	Pro	Gly	Lys
		435					440					445			

<400> SEQUENCE: 108

gctagcgttt	aaacttaagc	ttgttgacta	gtgagatcac	agttctctct	acagttactg	60
agcacacagg	acctcaccat	gggatggagc	tgtatcatcc	tcttcttggt	agcaacagct	120
acaggtaagg	ggctcacagt	agcaggcttg	aggctcggac	atatatatgg	gtgacaatga	180
catccacttt	gcctttctct	ccacagggtg	ccactcccag	gttacccctga	gagagtctgg	240
ccctgcgctg	gtgaagccca	cacagaccct	cacactgact	tgtaccttct	ctggggtttc	300
actgagcact	tctgggatgg	gtgtaggctg	gattcgctcag	cctcccgga	aggctctaga	360
gtggctggca	cacatttggt	gggatgatga	caagcgctat	aatccagccc	tgaagagccg	420
actgacaatc	tccaaggata	cctccaaaaa	ccaggtagtc	ctcacaatga	ccaacatgga	480
ccctgtggat	actgccacat	actactgtgc	tcaaataaac	cccgcctgg	ttgcttactg	540
gggcccaagg	actctggtca	ctgtgaqctc	a			571

```
<210> SEQ ID NO 109
<211> LENGTH: 118
```


-continued

cgctcctgcac caggactggc tgaatggcaa ggagtacaag tgcaaggtct ccaacaaagc	1200
cctcccagcc ccccatcgaga aaaccatctc caaagccaaa gggcagcccc gagaaccaca	1260
ggtgtacacc ctgcccccat cccgggatga gctgaccaag aaccaggtca gcctgacctg	1320
cctggtcaaa ggcttctatc ccagcgacat cgccgtggag tgggagagca atgggcagcc	1380
ggagaacaac tacaagacca cgctcccgt gctggactcc gacggctcct tcttctcta	1440
cagcaagctc accgtggaca agagcaggtg gcagcagggg aacgtcttct catgctccgt	1500
gatgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctccgggtaa	1560
atgagtgcgg ccgcgaattc	1580
<210> SEQ ID NO 111	
<211> LENGTH: 448	
<212> TYPE: PRT	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic Construct	
<400> SEQUENCE: 111	
Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln	
1 5 10 15	
Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser	
20 25 30	
Gly Met Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu	
35 40 45	
Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Asn Pro Ala	
50 55 60	
Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val	
65 70 75 80	
Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr	
85 90 95	
Cys Ala Gln Ile Asn Pro Ala Trp Phe Ala Tyr Trp Gly Gln Gly Thr	
100 105 110	
Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro	
115 120 125	
Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly	
130 135 140	
Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn	
145 150 155 160	
Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln	
165 170 175	
Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser	
180 185 190	
Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser	
195 200 205	
Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr	
210 215 220	
His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser	
225 230 235 240	
Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg	
245 250 255	
Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro	
260 265 270	
Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala	
275 280 285	

-continued

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Gln Ser Thr Tyr Arg Val Val
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
340 345 350

Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys
355 360 365

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
435 440 445

<210> SEQ ID NO 112
<211> LENGTH: 571
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 112

gctagcgttt aaacttaagc ttgttgacta gtgagatcac agttctctct acagttactg 60

agcacacagg acctcaccat gggatggagc tgtatcatcc tcttcttggt agcaacagct 120

acaggtaagg ggctcacagt agcaggcttg aggtctggac atatatatgg gtgacaatga 180

catccacttt gcctttctct ccacaggtgt cactcccag gttaccctga gagagtctgg 240

ccctgcgctg gtgaagccca cacagaccct cacactgact tgtaccttct ctggggttttc 300

actgagcact tctgggtgtg gtgtaggctg gattcgtcag cctcccggga aggcctctaga 360

gtggctggca cacatttggt gggatgatga caagcgctat tctccatccc tgaagagccg 420

actgacaatc tccaaggata cctccaaaaa ccaggtagtc ctcaaatga ccaacatgga 480

ccctgtggat actgccacat actactgtgc tcggataaac cccgcctact ttgcttactg 540

gggccaaggg actctggtca ctgtgagctc a 571

<210> SEQ ID NO 113
<211> LENGTH: 117
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 113

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

Gly Val Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

-continued

Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Ile Asn Pro Ala Tyr Phe Ala Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser
115

<210> SEQ ID NO 114
<211> LENGTH: 1580
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 114

gctagcgttt aaacttaagc ttgttgacta gtgagatcac agttctctct acagttactg 60

agcacacagg acctcaccat gggatggagc tgtatcatcc tcttcttggt agcaacagct 120

acaggtaagg ggctcacagt agcaggcttg aggtctggac atatatatgg gtgacaatga 180

catccacttt gcctttctct ccacaggtgt ccactcccag gttaccctga gagagtctgg 240

ccctgcgctg gtgaagccca cacagaccct cactgact tgtaccttct ctgggttttc 300

actgagcact tctggtgtgg gtgtaggctg gattcgtcag cctcccggga aggcctctaga 360

gtggctggca cacatttggg gggatgatga caagcgctat tctccatccc tgaagagccg 420

actgacaatc tccaaggata cctccaaaaa ccaggtagtc ctcaaatga ccaacatgga 480

ccctgtggat actgccacat actactgtgc tcggataaac cccgcctact ttgcttactg 540

gggccaaggg actctggtca ctgtgagctc agcctccacc aaggggcccat cggctctccc 600

cctggcacc cctccaaga gcacctctgg gggcacagcg gccctgggct gcctgggtcaa 660

ggactacttc cccgaaccgg tgacgggtgc gtggaactca ggcgcctga ccagcggcgt 720

gcacaccttc ccggctgtcc tacagtcttc aggactctac tccctcagca gcgtggtgac 780

cgtgccctcc agcagcttgg gcacccagac ctacatctgc aacgtgaatc acaagcccag 840

caacaccaag gtggacaaga gagttgagcc caaatcttgt gacaaaactc acacatgccc 900

accgtgccc gcacctgaac tcctgggggg accgtcagtc ttcctcttcc ccccaaaacc 960

caaggacacc ctcatgatct cccggacccc tgaggtcaca tgcgtggtgg tggacgtgag 1020

ccacgaagac cctgaggcca agttcaactg gtacgtggac ggcgtggagg tgcataatgc 1080

caagacaaaag ccgcgggagg agcagtacca gagcacgtac cgtgtggtca gcgtcctcac 1140

cgtcctgcac caggactggc tgaatggcaa ggagtacaag tgcaaggtct ccaacaaagc 1200

cctcccagcc cccatcgaga aaaccatctc caaagccaaa gggcagcccc gagaaccaca 1260

ggtgtacacc ctgcccccat cccgggatga gctgaccaag aaccaggcca gcctgacctg 1320

cctggtcaaa ggcttctatc ccagcgacat cgcctggagg tgggagagca atgggcagcc 1380

ggagaacaac tacaagacca cgctcccgt gctggactcc gacggctcct tcttctctta 1440

cagcaagctc accgtggaca agagcagggt gcagcagggg aacgtcttct catgctccgt 1500

gatgcatgag gctctgcaca accactacac gcagaagagc ctctccctgt ctccgggtaa 1560

atgagtgcgg ccgcgaattc 1580

-continued

<210> SEQ ID NO 115
<211> LENGTH: 448
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 115

Gln Val Thr Leu Arg Glu Ser Gly Pro Ala Leu Val Lys Pro Thr Gln
1 5 10 15

Thr Leu Thr Leu Thr Cys Thr Phe Ser Gly Phe Ser Leu Ser Thr Ser
20 25 30

Gly Val Gly Val Gly Trp Ile Arg Gln Pro Pro Gly Lys Ala Leu Glu
35 40 45

Trp Leu Ala His Ile Trp Trp Asp Asp Asp Lys Arg Tyr Ser Pro Ser
50 55 60

Leu Lys Ser Arg Leu Thr Ile Ser Lys Asp Thr Ser Lys Asn Gln Val
65 70 75 80

Val Leu Thr Met Thr Asn Met Asp Pro Val Asp Thr Ala Thr Tyr Tyr
85 90 95

Cys Ala Arg Ile Asn Pro Ala Tyr Phe Ala Tyr Trp Gly Gln Gly Thr
100 105 110

Leu Val Thr Val Ser Ser Ala Ser Thr Lys Gly Pro Ser Val Phe Pro
115 120 125

Leu Ala Pro Ser Ser Lys Ser Thr Ser Gly Gly Thr Ala Ala Leu Gly
130 135 140

Cys Leu Val Lys Asp Tyr Phe Pro Glu Pro Val Thr Val Ser Trp Asn
145 150 155 160

Ser Gly Ala Leu Thr Ser Gly Val His Thr Phe Pro Ala Val Leu Gln
165 170 175

Ser Ser Gly Leu Tyr Ser Leu Ser Ser Val Val Thr Val Pro Ser Ser
180 185 190

Ser Leu Gly Thr Gln Thr Tyr Ile Cys Asn Val Asn His Lys Pro Ser
195 200 205

Asn Thr Lys Val Asp Lys Arg Val Glu Pro Lys Ser Cys Asp Lys Thr
210 215 220

His Thr Cys Pro Pro Cys Pro Ala Pro Glu Leu Leu Gly Gly Pro Ser
225 230 235 240

Val Phe Leu Phe Pro Pro Lys Pro Lys Asp Thr Leu Met Ile Ser Arg
245 250 255

Thr Pro Glu Val Thr Cys Val Val Val Asp Val Ser His Glu Asp Pro
260 265 270

Glu Val Lys Phe Asn Trp Tyr Val Asp Gly Val Glu Val His Asn Ala
275 280 285

Lys Thr Lys Pro Arg Glu Glu Gln Tyr Gln Ser Thr Tyr Arg Val Val
290 295 300

Ser Val Leu Thr Val Leu His Gln Asp Trp Leu Asn Gly Lys Glu Tyr
305 310 315 320

Lys Cys Lys Val Ser Asn Lys Ala Leu Pro Ala Pro Ile Glu Lys Thr
325 330 335

Ile Ser Lys Ala Lys Gly Gln Pro Arg Glu Pro Gln Val Tyr Thr Leu
340 345 350

Pro Pro Ser Arg Asp Glu Leu Thr Lys Asn Gln Val Ser Leu Thr Cys
355 360 365

-continued

Leu Val Lys Gly Phe Tyr Pro Ser Asp Ile Ala Val Glu Trp Glu Ser
370 375 380

Asn Gly Gln Pro Glu Asn Asn Tyr Lys Thr Thr Pro Pro Val Leu Asp
385 390 395 400

Ser Asp Gly Ser Phe Phe Leu Tyr Ser Lys Leu Thr Val Asp Lys Ser
405 410 415

Arg Trp Gln Gln Gly Asn Val Phe Ser Cys Ser Val Met His Glu Ala
420 425 430

Leu His Asn His Tyr Thr Gln Lys Ser Leu Ser Leu Ser Pro Gly Lys
435 440 445

<210> SEQ ID NO 116
<211> LENGTH: 254
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 116

Met Trp Gln Leu Leu Leu Pro Thr Ala Leu Leu Leu Leu Val Ser Ala
1 5 10 15

Gly Met Arg Thr Glu Asp Leu Pro Lys Ala Val Val Phe Leu Glu Pro
20 25 30

Gln Trp Tyr Arg Val Leu Glu Lys Asp Ser Val Thr Leu Lys Cys Gln
35 40 45

Gly Ala Tyr Ser Pro Glu Asp Asn Ser Thr Gln Trp Phe His Asn Glu
50 55 60

Ser Leu Ile Ser Ser Gln Ala Ser Ser Tyr Phe Ile Asp Ala Ala Thr
65 70 75 80

Val Asp Asp Ser Gly Glu Tyr Arg Cys Gln Thr Asn Leu Ser Thr Leu
85 90 95

Ser Asp Pro Val Gln Leu Glu Val His Ile Gly Trp Leu Leu Leu Gln
100 105 110

Ala Pro Arg Trp Val Phe Lys Glu Glu Asp Pro Ile His Leu Arg Cys
115 120 125

His Ser Trp Lys Asn Thr Ala Leu His Lys Val Thr Tyr Leu Gln Asn
130 135 140

Gly Lys Gly Arg Lys Tyr Phe His His Asn Ser Asp Phe Tyr Ile Pro
145 150 155 160

Lys Ala Thr Leu Lys Asp Ser Gly Ser Tyr Phe Cys Arg Gly Leu Phe
165 170 175

Gly Ser Lys Asn Val Ser Ser Glu Thr Val Asn Ile Thr Ile Thr Gln
180 185 190

Gly Leu Ala Val Ser Thr Ile Ser Ser Phe Phe Pro Pro Gly Tyr Gln
195 200 205

Val Ser Phe Cys Leu Val Met Val Leu Leu Phe Ala Val Asp Thr Gly
210 215 220

Leu Tyr Phe Ser Val Lys Thr Asn Ile Arg Ser Ser Thr Arg Asp Trp
225 230 235 240

Lys Asp His Lys Phe Lys Trp Arg Lys Asp Pro Gln Asp Lys
245 250

<210> SEQ ID NO 117
<211> LENGTH: 254
<212> TYPE: PRT
<213> ORGANISM: Homo Sapien

<400> SEQUENCE: 117

-continued

Met	Trp	Gln	Leu	Leu	Leu	Pro	Thr	Ala	Leu	Leu	Leu	Leu	Val	Ser	Ala	
1				5					10					15		
Gly	Met	Arg	Thr	Glu	Asp	Leu	Pro	Lys	Ala	Val	Val	Phe	Leu	Glu	Pro	
			20					25					30			
Gln	Trp	Tyr	Arg	Val	Leu	Glu	Lys	Asp	Ser	Val	Thr	Leu	Lys	Cys	Gln	
		35					40					45				
Gly	Ala	Tyr	Ser	Pro	Glu	Asp	Asn	Ser	Thr	Gln	Trp	Phe	His	Asn	Glu	
	50					55					60					
Ser	Leu	Ile	Ser	Ser	Gln	Ala	Ser	Ser	Tyr	Phe	Ile	Asp	Ala	Ala	Thr	
65					70					75					80	
Val	Asp	Asp	Ser	Gly	Glu	Tyr	Arg	Cys	Gln	Thr	Asn	Leu	Ser	Thr	Leu	
				85					90					95		
Ser	Asp	Pro	Val	Gln	Leu	Glu	Val	His	Ile	Gly	Trp	Leu	Leu	Leu	Gln	
			100					105					110			
Ala	Pro	Arg	Trp	Val	Phe	Lys	Glu	Glu	Asp	Pro	Ile	His	Leu	Arg	Cys	
		115					120					125				
His	Ser	Trp	Lys	Asn	Thr	Ala	Leu	His	Lys	Val	Thr	Tyr	Leu	Gln	Asn	
						135					140					
Gly	Lys	Gly	Arg	Lys	Tyr	Phe	His	His	Asn	Ser	Asp	Phe	Tyr	Ile	Pro	
145					150					155					160	
Lys	Ala	Thr	Leu	Lys	Asp	Ser	Gly	Ser	Tyr	Phe	Cys	Arg	Gly	Leu	Val	
				165					170					175		
Gly	Ser	Lys	Asn	Val	Ser	Ser	Glu	Thr	Val	Asn	Ile	Thr	Ile	Thr	Gln	
			180					185					190			
Gly	Leu	Ala	Val	Ser	Thr	Ile	Ser	Ser	Phe	Phe	Pro	Pro	Gly	Tyr	Gln	
		195					200					205				
Val	Ser	Phe	Cys	Leu	Val	Met	Val	Leu	Leu	Phe	Ala	Val	Asp	Thr	Gly	
		210				215					220					
Leu	Tyr	Phe	Ser	Val	Lys	Thr	Asn	Ile	Arg	Ser	Ser	Thr	Arg	Asp	Trp	
225					230					235					240	
Lys	Asp	His	Lys	Phe	Lys	Trp	Arg	Lys	Asp	Pro	Gln	Asp	Lys			
				245					250							
<210> SEQ ID NO 118																
<211> LENGTH: 111																
<212> TYPE: PRT																
<213> ORGANISM: Artificial Sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Synthetic Construct																
<400> SEQUENCE: 118																
Asp	Ile	Val	Met	Thr	Gln	Ser	Pro	Asp	Ser	Leu	Ala	Val	Ser	Leu	Gly	
1				5					10					15		
Glu	Arg	Ala	Thr	Ile	Asn	Cys	Lys	Ser	Ser	Gln	Ser	Val	Asp	Phe	Asp	
			20					25					30			
Gly	Asp	Ser	Phe	Met	Asn	Trp	Tyr	Gln	Gln	Lys	Pro	Gly	Gln	Pro	Pro	
		35					40					45				
Lys	Leu	Leu	Ile	Tyr	Thr	Thr	Ser	Asn	Leu	Glu	Thr	Gly	Val	Pro	Asp	
	50					55					60					
Arg	Phe	Ser	Gly	Ser	Gly	Ser	Gly	Thr	Asp	Phe	Thr	Leu	Thr	Ile	Ser	
65					70					75					80	
Ser	Leu	Gln	Ala	Glu	Asp	Val	Ala	Val	Tyr	Tyr	Cys	Gln	Gln	Ser	Asn	
			85						90					95		
Ser	Asp	Pro	Tyr	Thr	Phe	Gly	Gln	Gly	Thr	Lys	Leu	Glu	Ile	Lys		
		100						105					110			

-continued

<210> SEQ ID NO 119
<211> LENGTH: 218
<212> TYPE: PRT
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic Construct

<400> SEQUENCE: 119

Asp Ile Val Met Thr Gln Ser Pro Asp Ser Leu Ala Val Ser Leu Gly
1 5 10 15

Glu Arg Ala Thr Ile Asn Cys Lys Ser Ser Gln Ser Val Asp Phe Asp
20 25 30

Gly Asp Ser Phe Met Asn Trp Tyr Gln Gln Lys Pro Gly Gln Pro Pro
35 40 45

Lys Leu Leu Ile Tyr Thr Thr Ser Asn Leu Glu Thr Gly Val Pro Asp
50 55 60

Arg Phe Ser Gly Ser Gly Ser Gly Thr Asp Phe Thr Leu Thr Ile Ser
65 70 75 80

Ser Leu Gln Ala Glu Asp Val Ala Val Tyr Tyr Cys Gln Gln Ser Asn
85 90 95

Ser Asp Pro Tyr Thr Phe Gly Gln Gly Thr Lys Leu Glu Ile Lys Arg
100 105 110

Thr Val Ala Ala Pro Ser Val Phe Ile Phe Pro Pro Ser Asp Glu Gln
115 120 125

Leu Lys Ser Gly Thr Ala Ser Val Val Cys Leu Leu Asn Asn Phe Tyr
130 135 140

Pro Arg Glu Ala Lys Val Gln Trp Lys Val Asp Asn Ala Leu Gln Ser
145 150 155 160

Gly Asn Ser Gln Glu Ser Val Thr Glu Gln Asp Ser Lys Asp Ser Thr
165 170 175

Tyr Ser Leu Ser Ser Thr Leu Thr Leu Ser Lys Ala Asp Tyr Glu Lys
180 185 190

His Lys Val Tyr Ala Cys Glu Val Thr His Gln Gly Leu Ser Ser Pro
195 200 205

Val Thr Lys Ser Phe Asn Arg Gly Glu Cys
210 215

What is claimed is:

1. An isolated anti-CD16A antibody comprising a V_H domain comprising a CDR1 having the amino acid sequence of SEQ ID NO:35, a CDR2 having the amino acid sequence of SEQ ID NO:39, and a CDR3 having the amino acid sequence of SEQ ID NO:59 and a V_L domain comprising a CDR1 having the amino acid sequence of SEQ ID NO:67, a CDR2 having the amino acid sequence of SEQ ID NO:75, and a CDR3 having the amino acid sequence of SEQ ID NO:88, wherein said CDRs have an amino acid substitution selected from the group consisting of, in the V_H domain, M34V in CDR1; H50L in CDR2, W52F in CDR2, D54N in CDR2, N60S in CDR2, A62S in CDR2; W99Y in CDR3, A101D in CDR3, and in the V_L domain, K24R in CDR1, A25S in CDR1, F32Y in CDR1, M33L in CDR1, N34A in CDR1, T50D, T50W or T50S in CDR2, T51A in CDR2, N53S in CDR2, E55A or E55Q in CDR2, S56T in CDR2; N92Y in CDR3, N93S in CDR3, and D94T in CDR3, which positions are according to the Kabat numbering scheme.

- 2. The isolated antibody of claim 1 that lacks an Fc region or comprises an Fc region that lacks effector function.
- 3. The isolated antibody of claim 2 comprising an Fc region that lacks effector function, which Fc region is a human IgG₁ Fc region.
- 4. The isolated antibody of claim 3 wherein the amino acid corresponding to residue 297 of the Fc region according to the Kabat numbering scheme is not asparagine.
- 5. The isolated antibody of claim 2 that is a single chain antibody.
- 6. The isolated antibody of claim 1, wherein said antibody comprises two light chains and two heavy chains.
- 7. An isolated humanized anti-CD16A antibody comprising a heavy chain variable region having the sequence of SEQ ID NO:113 and a light chain variable region having the sequence of SEQ ID NO:96.
- 8. An isolated humanized anti-CD16A antibody comprising a V_H CDR1 having the sequence of SEQ ID NO:35, a V_H CDR2 having the sequence of SEQ ID NO:39, a V_H CDR3

133

having the sequence of SEQ ID NO:59, and a V_L domain having the amino acid sequence of SEQ ID NO:96.

9. An isolated humanized anti-CD16A antibody that comprises a V_L domain having the amino acid sequence SEQ ID NO:96.

10. The isolated antibody of claim 9 that comprises a V_H domain having the amino acid sequence of SEQ ID NO: 109 or SEQ ID NO: 104.

11. The isolated antibody of any of claims 7 and 8-10, wherein said antibody lacks an Fc region or comprises an Fc region that lacks effector function.

12. The isolated antibody of claim 11 comprising an Fc region that lacks effector function, which Fc region is a human IgG₁ Fc region.

13. The isolated antibody of claim 11 comprising an Fc region that lacks effector function, wherein amino acid sequence corresponding to residue 297 of the Fc region according to the Kabat numbering scheme is not glycosylated.

134

14. The isolated antibody of claim 13, wherein the amino acid corresponding to residue 297 of the Fc region according to the Kabat numbering scheme is not asparagine.

15. The isolated antibody of claim 3, wherein the amino acid corresponding to residue 297 of the Fc region according to the Kabat numbering scheme is not glycosylated.

16. An isolated anti-CD16A antibody comprising a V_H CDR1 having the sequence of SEQ ID NO:35, a V_H CDR2 having the sequence of SEQ ID NO:39, a V_H CDR3 having the sequence of SEQ ID NO:59, and a light chain having the amino acid sequence of SEQ ID NO:98.

17. The isolated antibody claim 9, wherein said antibody comprises a heavy chain having the amino acid sequence of SEQ ID NO:111.

18. An isolated humanized anti-CD16A antibody comprising a light chain having the amino acid sequence of SEQ ID NO:98 and a heavy chain having the amino acid sequence of SEQ ID NO:111.

* * * * *