

US007560278B2

(12) **United States Patent**
Liu et al.

(10) **Patent No.:** **US 7,560,278 B2**
(45) **Date of Patent:** ***Jul. 14, 2009**

(54) **METHODS TO CULTURE CIRCOVIRUS**

(75) Inventors: **Qiang Liu**, Saskatoon (CA); **Suresh K. Tikoo**, Saskatoon (CA); **Philip Willson**, Saskatoon (CA); **Lorne A. Babiuk**, Saskatoon (CA)

(73) Assignee: **University of Saskatchewan**, Saskatoon (CA)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

| | | | |
|----|--------------|----|---------|
| WO | WO-99/53047 | A2 | 10/1999 |
| WO | WO-99/53047 | A3 | 10/1999 |
| WO | WO-00/01409 | A1 | 1/2000 |
| WO | WO-00/01409 | A3 | 1/2000 |
| WO | WO-00/26395 | A2 | 5/2000 |
| WO | WO-00/26395 | A3 | 5/2000 |
| WO | WO-00/47756 | A1 | 8/2000 |
| WO | WO-00/77216 | A2 | 12/2000 |
| WO | WO-00/77216 | A3 | 12/2000 |
| WO | WO-01/16330 | A2 | 3/2001 |
| WO | WO-01/16330 | A3 | 3/2001 |
| WO | WO-01/83737 | A2 | 11/2001 |
| WO | WO-01/83737 | A3 | 11/2001 |
| WO | WO-02/077210 | A2 | 10/2002 |
| WO | WO-02.077210 | A3 | 10/2002 |

(21) Appl. No.: **11/654,869**

(22) Filed: **Jan. 17, 2007**

(65) **Prior Publication Data**

US 2007/0128700 A1 Jun. 7, 2007

Related U.S. Application Data

(60) Continuation of application No. 10/840,879, filed on May 7, 2004, now Pat. No. 7,172,899, which is a division of application No. 10/112,540, filed on Mar. 27, 2002, now Pat. No. 6,794,163.

(60) Provisional application No. 60/279,173, filed on Mar. 27, 2001.

(51) **Int. Cl.**

C12N 5/00 (2006.01)

A61K 39/12 (2006.01)

(52) **U.S. Cl.** **435/325**; 435/69.1; 435/70.1; 424/233.1

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

| | | | |
|--------------|----|---------|-----------------|
| 6,217,883 | B1 | 4/2001 | Allan et al. |
| 6,287,856 | B1 | 9/2001 | Poet et al. |
| 6,368,601 | B1 | 4/2002 | Allan et al. |
| 6,391,314 | B1 | 5/2002 | Allan et al. |
| 6,492,343 | B1 | 12/2002 | Reddy et al. |
| 6,660,272 | B2 | 12/2003 | Allan et al. |
| 6,703,023 | B1 | 3/2004 | Jestin et al. |
| 6,794,163 | B2 | 9/2004 | Liu et al. |
| 6,943,152 | B1 | 9/2005 | Audonnet et al. |
| 7,172,899 | B2 | 2/2007 | Liu et al. |
| 2002/0177216 | A1 | 11/2002 | Liu et al. |
| 2003/0130187 | A1 | 7/2003 | Reddy et al. |
| 2004/0132178 | A1 | 7/2004 | Haines et al. |

FOREIGN PATENT DOCUMENTS

| | | | |
|----|-------------|----|--------|
| AU | 756554 | A | 4/1999 |
| WO | WO-99/08706 | A1 | 2/1999 |
| WO | WO-99/18214 | A1 | 4/1999 |
| WO | WO-99/29871 | A2 | 6/1999 |
| WO | WO-99/29871 | A3 | 6/1999 |
| WO | WO-99/45956 | A1 | 9/1999 |

OTHER PUBLICATIONS

Allan, G.M. et al. (1994). "Some Biological and Physico-Chemical Properties of Porcine Circovirus," *J. Vet Med. B.* 41:17-26.

Allan, G.M. et al. (1995). "Pathogenesis of Porcine Circovirus; Experimental Infections of Colostrum Deprived Piglets and Examination of Pig Foetal Material," *Veterinary Microbiology* 44:49-64.

Allan, G.M. et al. (1998). "Isolation of Porcine Circovirus-Like Viruses from Pigs with a Wasting Disease in the USA and Europe," *J. Vet. Diagn. Invest.* 10:3-10.

Allan, G.M. et al. (2000). "Porcine Circoviruses: A Review," *J. Vet. Diagn. Invest.* 12:3-14.

Certified English translation of PCT publication No. WO 00/47756, published Aug. 17, 2000, entitled "Recombined Porcine Adenovirus Based Viral Vaccines and Vectors," 60 pages.

Chartier, C. et al. (Jul. 1996). "Efficient Generation of Recombinant Adenovirus Vectors by Homologous Recombination in *Escherichia coli*," *J. Virol.* 70(7):4805-4810.

Chroboczek, J. et al. (1992). "The Sequence of the Genome of Adenovirus Type 5 and Its Comparison with the Genome of Adenovirus Type 2," *Virology* 186:280-285.

Ellis, J. et al. (Jan. 1998). "Isolation of Circovirus from Lesions of Pigs with Postweaning Multisystemic Wasting Syndrome," *Can. Vet. J.* 39:44-51.

Ellis, J. et al. (1999). "Reproduction of Lesions of Postweaning Multisystemic Wasting Syndrome in Gnotobiotic Piglets," *J. Vet. Diagn. Invest.* 11:3-14.

Flint, S.J. (1986). "Regulation of Adenovirus mRNA Formation," *Advances in Virus Research* 31:169-228.

Flint, S.J. et al. (1982). "Expression of Adenoviral Genetic Information in Productively Infected Cells," *Biochem. Biophys. Acta* 651:175-208.

Grand, R.J.A. (1987). "The Structure and Functions of the Adenovirus Early Region 1 Proteins," *Biochem. J.* 241:25-38.

(Continued)

Primary Examiner—Stacy B Chen

(74) *Attorney, Agent, or Firm*—Morrison & Foerster LLP

(57) **ABSTRACT**

The present invention relates to methods for culturing circovirus and in particular, porcine circovirus. The present invention provides compositions and methods for culturing porcine circovirus in mammalian cells expressing mammalian adenovirus E1 function.

8 Claims, 4 Drawing Sheets

OTHER PUBLICATIONS

- Hamel, A.L. et al. (May 1998). Genbank Accession No. AF027217. Porcine Circovirus . . . [gi:2689645]. "Nucleotide Sequence of Porcine Circovirus Associated with Postweaning Multisystemic Wasting Syndrome in Pigs," total pages 4.
- Hamel, A.L. et al. (Jun. 1998). "Nucleotide Sequence of Porcine Circovirus Associated with Postweaning Multisystemic Wasting Syndrome in Pigs," *J. Virol.* 72(6):5262-5267.
- Hirt, B. (1967). "Selective Extraction of Polyma DNA from Infected Mouse Cell Cultures," *J. Mol. Biol.* 26:365-369.
- International Search Report mailed Jun. 6, 2003, for PCT Application No. PCT/CA02/00413, filed Mar. 27, 2003, five pages (with International Search Report for PCT Application No. PCT/CA01/00598 inadvertently included with above report, two pages.)
- Kennedy, S. et al. (2000). "Reproduction of Lesions of Postweaning Multisystemic Wasting Syndrome by Infection of Conventional Pigs with Porcine Circovirus Type 2 Alone or in Combination with Porcine Parvovirus," *Journal of Comparative Pathology* 122(1):9-24.
- Kleiboeker, S.B. (1995). "Identification and Sequence Analysis of the E1 Genomic Region of a Porcine Adenovirus," *Virus Res.* 36:259-268.
- Krakowka, S. et al. (2000). "Viral Wasting Syndrome of Swine: Experimental Reproduction of Postweaning Multisystemic Wasting Syndrome in Gnotobiotic Swine by Coinfection with Porcine Circovirus 2 and Porcine Parvovirus," *Vet. Pathol.* 37:254-263.
- Liu, Q. et al. (Sep. 2000). "Quantitative Competitive PCR Analysis of Porcine Circovirus DNA in Serum from Pigs with Postweaning Multisystemic Wasting Syndrome," *J. Clin. Microbiol.* 38(9):3474-3477.
- Liu, Q. et al. (2001). "Bacterial Expression of an Immunologically Reactive PCV2 ORF2 Fusion Protein," *Protein Express and Purif.* 21:115-120.
- Liu, Q. et al. (2001). "Nuclear Localization of the ORF2 Protein Encoded by Porcine Circovirus Type 2," *Virology* 285(1):91-99.
- Lukert, P. D. et al. (1995). "Circoviridae", In *Virus Taxonomy: Classification and Nomenclature of Viruses*. F. A. Murphy, et al. (eds.) Sixth Report of the International Committee on Taxonomy of Viruses, Virology Division International Union of Microbiological Societies, Springer-Verlag Wien New York. pp. 166-168.
- Mahé, D. et al. (2000). "Differential Recognition of ORF2 Protein from Type 1 and Type 2 Porcine Circoviruses and Identification of Immunorelevant Epitopes," *Journal of General Virology* 81:1815-1824.
- Mankertz, A. et al. (Mar. 1997). "Mapping and Characterization of the Origin of DNA Replication of Porcine Circovirus," *J. Virol.* 71(3):2562-2566.
- Mankertz, A. et al. (1998). "Identification of a Protein Essential for Replication of Porcine Circovirus," *Journal of General Virology* 79:381-384.
- Mankertz, A. et al. (2000). "Characterisation of PCV-2 Isolates from Spain, Germany, and France," *Virus Res.* 66:65-77.
- Meehan, B. M. et al. (1997). "Sequence of Porcine Circovirus DNA: Affinities with Plant Circoviruses," *Journal of General Virology* 78:221-227.
- Meehan, B. M. et al. (1998). "Characterization of Novel Circovirus DNAs Associated with Wasting Syndromes in Pigs," *Journal of General Virology* 79:2171-2179.
- Morozov, I. et al. (Sep. 1998). "Detection of a Novel Strain of Porcine Circovirus in Pigs with Postweaning Multisystemic Wasting Syndrome," *J. Clin. Microbiol.* 36(9):2535-2541.
- Murphy, F.A. et al. eds. (1999). "Circoviridae," Chapter 22 in *Veterinary Virology*. 3rd ed. Academic Press, p. 357-361.
- Nagy, M. et al. (2000). "Porcine Adenoviruses: An Update On Genome Analysis and Vector Development," *Acta Veterinaria Hungarica* 48(4):491-499.
- Nagy, M. et al. (2001). "The Complete Nucleotide Sequence of Porcine Adenovirus Serotype 5," *Journal of General Virology* 82(3):525-529.
- Nawagitgul, P. et al. (2000). "Open Reading Frame 2 of Porcine Circovirus Type 2 Encodes a Major Capsid Protein," *Journal of General Virology* 81:2281-2287.
- Non-Final Office Action issued Sep. 28, 2005, for U.S. Appl. No. 10/840,879, six pages.
- Non-Final Office Action issued Apr. 18, 2006, for U.S. Appl. No. 10/840,879, four pages.
- Pogranichnyy, R. M. et al. (2000). "Characterization of Immune Response of Young Pigs to Porcine Circovirus Type 2 Infection," *Viral. Immunol.* 13(2):143-153.
- Reddy, P.A. et al. (1999). "Replication-Defective Bovine Adenovirus Type 3 as an Expression Vector," *Journal of Virology* 73(11):9137-9144.
- Reddy, P.S. et al. (1995). "Comparison of the Inverted Terminal Repetition Sequences from Five Porcine Adenovirus Serotypes," *Virology* 212(1):237-239.
- Reddy, P.S. et al. (Feb. 1998). "Nucleotide Sequence, Genome Organization, and Transcription Map of Bovine Adenovirus Type 3," *Journal of Virology* 72(2):1394-1402.
- Reddy, P.S. et al. (1999). "Porcine Adenovirus-3 As A Helper-Dependent Expression Vector," *J. Gen. Virol.* 80:2909-2916.
- Sambrook, J. et al. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press, p. 7.52 entitled "Hybridization and Autoradiography".
- Shenk, T. (1996). "Adenoviridae: The Viruses and Their Replications," in *Fields Virology*. 3rd ed., Fields, B.N. et al. eds., Lippincott-Raven Publishers: Philadelphia, PA, pp. 2111-2148.
- Shenk, T. (1996). "Adenoviridae: The Viruses and Their Replications," Chapter 30 in *Fields Virology*, 3rd ed., Fields, B.N. et al. eds., Lippincott-Raven Publishers: Philadelphia, PA, pp. 979-1016.
- Tischer, I. et al. (1986). "Studies on Epidemiology and Pathogenicity of Porcine Circovirus," *Arch. Virol.* 91:271-276.
- Tischer, I. et al. (1987). "Replication of Porcine Circovirus: Induction by Glucosamine and Cell Cycle Dependence," *Arch. Virol.* 96:39-57.
- Tischer, I.D. et al. (1995). "Occurrence and Role of an Early Antigen and Evidence for Transforming Ability of Porcine Circovirus," *Arch. Virol.* 140:1799-1816.
- Tuboly, T. et al. (1993). "Potential Viral Vectors for the Stimulation of Mucosal Antibody Responses Against Enteric Viral Antigens in Pigs," *Research In Veterinary Science* 54(3):345-350.
- Tuboly, T. et al. (1995). "Restriction Endonuclease Analysis and Physical Mapping of the Genome of Porcine Adenovirus Type 5," *Virus Research* 37(1):49-54.
- Tuboly, T. et al. (2000). "Characterization of Early Region 4 of Porcine Adenovirus Serotype 5," *Virus Genes* 20(3):217-219.
- Tuboly, T. et al. (2000). "Sequence Analysis and Deletion of Porcine Adenovirus Serotype 5 E3 Region," *Virus Research* 68(2):109-117.
- Tuboly, T. et al. (2001). "Construction and Characterization of Recombinant Porcine Adenovirus Serotype 5 Expressing the Transmissible Gastroenteritis Virus Spike Gene," *Journal of General Virology* 82(1):183-190.
- Wang, L. et al. (Sep. 1998). Genbank Accession No. AF086834. Porcine Circovirus . . . [gi:3661515]. "Porcine Circovirus Strain B9, Complete Genome," total pages 4.
- Wang, L. et al. (Sep. 1998). Genbank Accession No. AF086835. Porcine Circovirus . . . [gi:3661518]. "Porcine Circovirus Strain 9741, Complete Genome," total pages 3.
- Wang, L. et al. (Sep. 1998). Genbank Accession No. AF086836. Porcine Circovirus . . . [gi:3661521]. "Porcine Circovirus Strain M226, Complete Genome," total pages 3.
- English translation of PCT Publication No. WO 99/29871, entitled "Circovirus Sequences Associated With Piglet Weight Loss Disease (PWD)", created on Jun. 25, 2003, 118 pages.




Fig. 1A-1B




Fig. 2A-2B

1 accagcgcac ttcggcagcg gcagcacctc ggcaacacct cagcagcaac atgccagca
61 agaagaatgg aagaagcgga cccaaccac ataaaagggtg ggtgttcacg ctgaataatc
121 cttccgaaga cgagcgcaag aaaatacggg agtcccaat ctcctattt gattatttta
181 ttgttggcga ggagggtaat gaggaaggac gaacacctca cctccagggg ttcgctaatt
241 ttgtgaagaa gcaaactttt aataaagtga agtgggtattt gggtgcccgc tgccacatcg
301 agaaagccaa aggaactgat cagcagaata aagaatattg tagtaaagaa ggcaacttac
361 ttattgaatg tggagctcct cgatctcaag gacaacggag tgacctgtct actgctgtga
421 gtaccttgtt ggagagcggg attctggtga ccgttgcaa gcagcacctt gtaacgtttg
481 tcaaaaattt ccgcgggctg gctgaacttt tgaaagtgag cgggaaaatg caaaagcgtg
541 attggaaaac caatgtacac ttcattgtgg ggccacctgg gtgtggtaaa agcaaatggg
601 ctgctaattt tgcaaaccgg gaaaccacat actggaaacc acctaaaaac aagtgggtgg
661 atggttacca tggtgaaaaa gtggttggtt ttgatgactt ttatggctgg ctgccgtggg
721 atgatctact gagactgtgt gatcgatata cattgactgt aaaaactaaa ggtggaactg
781 tacctttttt ggcccgcagt attctgatta ccagcaatca aaccccgttg gaatggtact
841 cctcaactgc tgtcccagct gtagaagctc tctatcggag gattacttcc ttggtatttt
901 ggaagaatgt tacagaacaa tccacggagg aagggggcca gtttgtcacc ctttcccccc
961 catgccctga atttccatat gaaataaatt actgagtctt tttatcact tcgtaatggt
1021 ttttattatt catttagggt ttaagtgggg ggtctttaag attaaattct ctgaattgta
1081 catacatggt tacacggata ttgtagctct ggtcgtattt actgttttcg aacgcagtgc
1141 cgaggcctac gtgggtccaca tttctagagg ttgtagcct cagccaaagc tgattccttt
1201 tgttatttgg ttggaagtaa tcaatagtgg agtcaagaac aggtttgggt gtgaagtaac
1261 gggagtggta ggagaagggt tgggggattg tatggcggga ggagtgttt acatatgggt
1321 cataggttag ggctgtggcc tttgttacia agttatcatc tagaataaca gcagtggagc
1381 ccactcccct atcaccctgg gtgatggggg agcagggcca gaattcaacc ttaaccttcc
1441 ttattctgta gtattcaaag ggtatagaga ttttgttggg cccccctccc gggggaacaa
1501 agtcgtcaat attaaatctc atcatgtcca ccgccagga gggcgttgtg actgtggtag
1561 ccttgacagt atatccgaag gtgcgggaga ggcgggtgtt gaagatgcca ttttctctc
1621 tccaacggta gcggtggcgg ggggtggacga gccagggggc gcggcggagg atctggccaa
1681 gatggctgcg ggggcggtgt cttctctctg ggtaacgcct ccttggatac gtcatactg
1741 aaaacgaaag aagtgcgctg taagtatt

FIG. 3A

MPSKKNRSGPQPHKRWFVTLNPNSEDERKKIRELPISLFDYFI
VGEEGNEEGRTPHLQGFANFVKKQTFNKVKWYLGARCHIEKAKGTDQONKEYCSKEGN
LLIECGAPRSQGQRSDLSTAVSTLLESGILVTVAKQHPVTFVKNFRGLAELLKVS GKM
QKRDWKTNVHFIVGPPGCGKSKWAANFANPETTYWKPPKNKWWDGYHGEKVVVIDDFY
GWLPWDDLRLCDRYPLTVKTKGGTVPFLARSILITSNQTPLEWYSSTAVPAVEALYR
RITSLVFWKNVTEQSTEEGGQFVTLSPPCPEFPYEINY

FIG. 3B

MTYPRRRYRRRRHRPRSHLGQILRRRPWLHVHPRHRYRWRKNGI
FNTRLSRTFGYTVKATTVTTPSWAVDMMRFNIDDFVPPGGGTNKISIPFEYYRIRKVK
VEFWPCSPITQGDRGVGSTAVILDDNFVTKATALTYDPYVNYSSRHTIPQPFSYHSRY
FTPKPVL DSTIDYFQPNNKRNQLWLR LQTSRNV DHVGLGTAFENSKYDQDYNIRVTMY
VQFREFNLKDPPLKP

FIG. 3C

METHODS TO CULTURE CIRCOVIRUS**CROSS-REFERENCE TO RELATED APPLICATIONS**

This application is a continuation application of U.S. patent application Ser. No. 10/840,879, filed May 7, 2004, now U.S. Pat. No. 7,172,899, which is a divisional of U.S. patent application Ser. No. 10/112,540, filed Mar. 27, 2002, issued on Sep. 21, 2004 as U.S. Pat. No. 6,794,163, which claims the benefit of U.S. Provisional Application Ser. No. 60/279,173, filed Mar. 27, 2001, all of which are hereby incorporated by reference herein in their entirety.

TECHNICAL FIELD

The present invention relates to the field of circovirus and provides compositions and methods for culturing circovirus, in particular porcine circovirus. In particular, the present invention relates to methods for culturing porcine circovirus in mammalian cells expressing a mammalian adenovirus E1 gene function.

BACKGROUND ART

A family of viruses, named Circoviridae, found in a range of plant and animal species and commonly referred to as circoviruses, are characterized as round, non-enveloped virions with mean diameters from 17 to 23.5 nm containing circular, single-stranded deoxyribonucleic acid (ssDNA). The ssDNA genome of the circoviruses represent the smallest viral DNA replicons known. As disclosed in WO 99/45956, at least six viruses have been identified as members of the family according to The Sixth Report of the International Committee for the Taxonomy of Viruses (Lukert, P. D. et al. 1995, *The Circoviridae*, pp. 166-168. In F. A. Murphy, et al. (eds.) *Virus Taxonomy, Sixth Report of the International Committee on Taxonomy of Viruses*, Arch. Virol. 10 Suppl.).

Animal viruses included in the family are chicken anemia virus (CAV); beak and feather disease virus (BFDV); porcine circovirus (PCV); and pigeon circovirus. PCV was originally isolated in porcine kidney cell cultures. PCV replicates in the cell nucleus and produces large intranuclear inclusion bodies. See Murphy et al. (1999, *Circoviridae* p. 357-361, *Veterinary Virology*, 3rd ed. Academic Press, San Diego). There are currently two recognized types of PCV, PCV type 1 (PCV1) and PCV type 2 (PCV2). PCV1, isolated as a persistent contaminant of the continuous porcine kidney cell line PK-15 (ATCC CCL31), does not cause detectable cytopathic effects in cell culture and fails to produce clinical disease in pigs after experimental infection (see Allan G., 1995, *Vet. Microbiol.* 44: 49-64; Tischer, I. et al., 1982, *Nature* 295:64-66; and Tischer, I. et al., 1986, *Arch. Virol.* 91:271-276). PCV2, in contrast to PCV1, is closely associated with post weaning multisystemic wasting syndrome (PMWS) in weanling pigs (see Allan G. et al., 1998, *Europe. J. Vet. Diagn. Investig.* 10:3-10; Ellis, J. et al., 1998, *Can. Vet. J* 39:44-51 and Morozov, I. et al., 1998, *J Clin. Microbiol.* 36:2535-2541). The nucleotide sequences for PCV1 are disclosed in Mankertz, A., et al. (1997, *J. Virol.* 71:2562-2566) and Meehan, B. M., et al. (1997, *J. Gen. Virol.* 78:221-227) and the nucleotide sequences for PCV2 are disclosed in Hamel, A. L. et al. (1998, *J Virol* 72:5262-5267); Mankertz, A. et al. (2000, *Virus Res.* 66:65-77) and Meehan, B. M. et al. (1998, *J. Gen. Virol.* 79:2171-2179). Strains of PCV2 are disclosed in WO 00/01409 and have been deposited at the European Collection of Cell Cultures, Centre for Applied Microbiology &

Research, Porton Down, Salisbury, Wiltshire SP4 OJG, United Kingdom and include: accession No. V97100219; accession No. V9700218; accession No. V97100217; accession No. V98011608; and accession No. V98011609. WO 00/77216 also discloses PCV2.

Published studies to date on PCV2 used either tissue homogenate or cultured virus derived from field isolates. Tischer et al. (1987, *Arch Virol.* 96:39-57) report that porcine kidney cells are stimulated to entry to the S phase in the cell cycle by D-glucosamine treatment. However, the treatment must be performed with caution because D-glucosamine is toxic for cell culture (see, Allan et al., (2000). *J. Vet. Diagn. Investigation.* 12:3-14). There remains a need for methods for culturing circovirus, such as for example, PCV1 and PCV2, and other circoviruses, such that pure circovirus is obtained. Such methods would be advantageous, in particular for preparation of PCV2 antigens as vaccines directed against PMWS. The present invention addresses that need.

All patents and publications are hereby incorporated herein in their entirety.

DISCLOSURE OF THE INVENTION

The present invention provides methods for culturing mammalian circovirus comprising: a) obtaining mammalian cells expressing a mammalian adenovirus E1 function, wherein said cells are permissive for mammalian circovirus replication; b) introducing said mammalian circovirus genome, or a portion thereof capable of replication, into said mammalian cells; and c) culturing said mammalian cells under conditions suitable for replication of said mammalian circovirus. In some embodiments, the method further comprises recovering said circovirus from said cultured cells.

In some embodiments, the mammalian circovirus is porcine circovirus, such as for example, porcine circovirus 1 (PCV1) or porcine circovirus 2 (PCV2). In yet additional embodiments, the porcine circovirus comprises a chimeric nucleotide sequence. In other embodiments, the mammalian cells are of porcine origin. In yet other embodiments, the mammalian cells are porcine retina cells.

In other embodiments, the mammalian adenovirus E1 function is human adenovirus E1 function. In yet other embodiments, the mammalian adenovirus E1 function is porcine adenovirus E1 function. In further embodiments, the E1 function is E1A and/or E1B function. In yet further embodiments, the mammalian cell expressing the mammalian E1 function is stably transformed with mammalian E1 gene sequences. In other embodiments, the mammalian E1 gene sequence is heterologous to said mammalian cell.

The present invention also provides recombinant mammalian cells that express a mammalian adenovirus E1 function and comprise a mammalian circovirus genome, or a portion thereof capable of replication, and wherein said cells are permissive for the replication of said mammalian circovirus. In some embodiments, the mammalian circovirus is porcine circovirus, such as for example, porcine circovirus 1 (PCV1) or porcine circovirus 2 (PCV2). In yet additional embodiments, the porcine circovirus comprises a chimeric nucleotide sequence. In some embodiments, the adenovirus E1 function is human adenovirus E1 function. In other embodiments, the E1 function is porcine adenovirus E1 function. In other embodiments, the mammalian cell is of porcine origin. In further embodiments, the mammalian cell is a porcine retinal cell. In yet further embodiments, the mammalian cell expressing the mammalian E1 function is stably transformed

with mammalian adenovirus E1 gene sequences. In other embodiments, the mammalian E1 gene sequence is heterologous to said mammalian cell.

The present invention also provides methods of preparing a recombinant mammalian cell expressing a mammalian adenovirus E1 function and comprising a mammalian circovirus genome comprising the steps of, a) obtaining a mammalian cell expressing a mammalian adenovirus E1 function; and b) introducing said mammalian circovirus genome, or a portion thereof capable of replication, into said mammalian cell. In additional embodiments, the method comprises the additional step of culturing the recombinant mammalian cell under conditions suitable for the replication of said mammalian circovirus. In further embodiments, the method comprises recovering said circovirus from said cultured cells. In some embodiments, the mammalian circovirus is porcine circovirus, such as for example, porcine circovirus 1 (PCV1) or porcine circovirus 2 (PCV2). In yet additional embodiments, the porcine circovirus comprises a chimeric nucleotide sequence. In further embodiments, the mammalian cells are of porcine origin. In yet further embodiments, the mammalian cells are porcine retina cells. In additional embodiments, the adenovirus E1 function is human adenovirus E1 function or porcine adenovirus E1 function. In yet further embodiments, the mammalian cell expressing the mammalian adenovirus E1 function is stably transformed with mammalian adenovirus E1 gene sequences. In other embodiments, the mammalian E1 gene sequence is heterologous to said mammalian cell.

BRIEF DESCRIPTION OF THE DRAWINGS

FIGS. 1A-1B provide the characterization and titration of PCV2 virus generated by DNA transfection and extraction from infected VIDO R1 cells by Hirt's method. (A) PCR using PCV2-specific primers and DNA from PCV 2-infected (lane 1) and mock-infected (lane 2) cells. A plasmid containing PCV2 genome was used as a control (lane 3). The 1-kb DNA ladder from GIBCO BRL was loaded in lane M. (B) Viral DNA from PCV2-infected (lanes 1, 3, and 5) and mock-infected (lanes 2, 4, and 6) cells were digested with NcoI and StuI (lanes 1 and 2), EcoRI and StuI (lanes 3 and 4), and EcoRI and EcoRV (lanes 5 and 6). The 1-kb-plus DNA ladder from GIBCO BRL was loaded in lane M.

FIGS. 2A-2B depict titration of PCV2 by immunoperoxidase staining. At 72 h.p.i., mock-(A) or PCV2-(B) infected VIDO R1 cells were incubated with rabbit anti-ORF2 polyclonal antibody and biotinylated secondary antibody. After application of an avidin and biotinylated horseradish peroxidase complex, the monolayer was developed by diaminobenzidine tetrahydrochloride (DAB). One dark cell resulted from one virus particle infection.

FIGS. 3A-3C show the nucleotide sequence (SEQ ID NO:1) (A) and amino acid sequence for ORF 1 (SEQ ID NO:2) (B) and ORF 2 (SEQ ID NO:3) (C) of porcine circovirus 2 (PCV2) as described in Genbank accession number AF086834.

BEST MODE FOR CARRYING OUT THE INVENTION

The present invention relates to compositions and methods for culturing mammalian circovirus, in particular porcine circovirus. The present invention is based on the finding that a porcine cell expressing human E1 function was able to be transfected with a PCV2 virus genome and generated PCV2 virus with a high virus titer. VIDO R1 cell line, deposited with

the ATCC and having ATCC accession number PTA-155, is a porcine retina cell line transformed with human adenovirus-5 (HAV-5) E1, that has been shown to induce the S phase of the cell cycle and transactivate transcription. See, Shenk, T. (1996). "Adenoviridae: the viruses and their replication" In *Fields Virology*. 3rd ed. B. N. Fields, D. M. Knipe and P. M. Howley (ed.) Lippincott-Raven Publishers, Philadelphia, New York, pp. 2111-2148. As described herein in Example 3, VIDO R1 cells, were transfected with a PCV2 genome and generated virus at 2×10^7 IU/ml.

The practice of the present invention employs, unless otherwise indicated, conventional microbiology, immunology, virology, molecular biology, and recombinant DNA techniques which are within the skill of the art. These techniques are fully explained in the literature. See, e.g., Maniatis et al., *Molecular Cloning: A Laboratory Manual* (1982); *DNA Cloning: A Practical Approach*, vols. I & II (D. Glover, ed.); *Oligonucleotide Synthesis* (N. Gait, ed. (1984)); *Nucleic Acid Hybridization* (B. Hames & S. Higgins, eds. (1985)); *Transcription and Translation* (B. Hames & S. Higgins, eds. (1984)); *Animal Cell Culture* (R. Freshney, ed. (1986)); Perbal, *A Practical Guide to Molecular Cloning* (1984); Ausubel, et al., *Current Protocols In Molecular Biology*, John Wiley & Sons (1987, 1988, 1989, 1990, 1991, 1992, 1993, 1994, 1995, 1996); and Sambrook et al., *Molecular Cloning: A Laboratory Manual* (2nd Edition); vols. I, II & III (1989).

Circoviridae, a family of viruses having round, non-enveloped virions with mean diameters from 17 to 23.5 nm containing circular, single-stranded DNA (ssDNA), are described in The Sixth Report of the International Committee for the Taxonomy of Viruses, supra. Members of the group include the porcine circoviruses, PCV1 and PCV2. Some of the PCVs are known to be pathogenic, such as PCV2, associated with PMWS.

Nucleotide sequences for PCV1 are provided in Mankertz, A., et al., 1997, *J. Virol.* 71:2562-2566 and Meehan, B. M. et al., 1997, *J. Gen. Virol.* 78:221-227. Nucleotide sequences for PCV2 are provided in Hamel, A. L. et al., 1998, *J. Virol.* 72:5262-5267; Mankertz, A. et al., 2000, *Virus Res.* 66:65-77 and Meehan, B. M. et al., 1998, *J. Gen. Virol.* 79:2171-2179. Representative strains of PCV2 have been deposited with the European Collection of Cell Cultures, Centre for Applied Microbiology & Research, Porton Down, Salisbury, Wiltshire SP4 OJG, United Kingdom and include accession No. V97100219; accession No. V9700218; accession No. V97100217; accession No. V98011608; and accession No. V98011609. WO 00/77216 also discloses PCV2. PCV2 nucleotide sequences have also been published in Hamel et al., (1998), *J. Virol.* vol. 72, 6:5262-5267 (GenBank AF027217) and in Morozov et al., (1998), *J. Clinical Microb.* vol. 36, 9:2535-2541, as well as GenBank AF086834; AF086835; and AF086836. Comparison of the published nucleotide sequences for PCV1 and PCV2 reveals a <80% identity, although the genomic organization is similar, especially in the arrangement of the two largest open reading frames (ORFs) with a putative origin of DNA replication.

The present invention encompasses methods of culturing mammalian circovirus and in particular, porcine circovirus (PCV). The present invention encompasses methods of culturing PCV comprising the PCV nucleotide sequences disclosed herein or known in the art, or ORFs thereof, or portions thereof that are capable of replication. The present invention also encompasses methods of culturing PCV having PCV nucleotide sequences differing through the degeneracy of the genetic code to those disclosed herein or known in the art, or ORFs thereof, or portions thereof capable of replication. The present invention further encompasses methods of culturing

PCV comprising PCV nucleotide sequence variations which do not change the functionality or strain specificity of the nucleotide sequence, or ORFs thereof, or portions thereof capable of replication. The present invention also encompasses methods of culturing PCV comprising PCV nucleotide sequences capable of hybridizing to those sequences disclosed herein under conditions of intermediate to high stringency, and methods of culturing PCV comprising mutations of the PCV nucleotide sequence disclosed herein or known in the art, such as deletions or point mutations, or ORFs thereof, or portions thereof, capable of replication. The present invention also encompasses methods of culturing PCV comprising heterologous nucleotide sequences. The present invention encompasses methods of culturing PCV that comprise chimeric circovirus nucleotide sequences, such as, for example, nucleotide sequences from porcine circovirus in fusion with nucleotide sequences from other pathogenic viruses, such as a pathogenic porcine virus, including parvovirus.

As used herein, a heterologous nucleotide sequence, with respect to a circovirus or mammalian cell, is one which is not normally associated with the circovirus sequences as part of the circovirus genome or one which is not normally associated with the mammalian cell, respectively. Heterologous nucleotide sequences include synthetic sequences. Hybridization reactions can be performed under conditions of different "stringency". Conditions that increase stringency of a hybridization reaction are widely known and published in the art. See, for example, Sambrook et al. (1989) at page 7.52. Examples of relevant conditions include (in order of increasing stringency): incubation temperatures of 25° C., 37° C., 50° C. and 68° C.; buffer concentrations of 10×SSC, 6×SSC, 1×SSC, 0.1×SSC (where SSC is 0.15 M NaCl and 15 mM citrate buffer) and their equivalents using other buffer systems; formamide concentrations of 0%, 25%, 50%, and 75%; incubation times from 5 minutes to 24 hours; 1, 2, or more washing steps; wash incubation times of 1, 2, or 15 minutes; and wash solutions of 6×SSC, 1×SSC, 0.1×SSC, or deionized water. An exemplary set of stringent hybridization conditions is 68° C. and 0.1×SSC.

The PCV genomes encode several polypeptide sequences, ranging in approximate size from 8 to 35 kD. It is deemed routine to determine open reading frames (ORFs) for porcine circoviruses using standard software such as for example, MacVector® (Oxford Molecular Group Inc., MD 21030). The largest ORF, ORF1, of the two types of PCV shows only minor variation with an identity of 85% (as measured by the clustal program) and has been demonstrated to be the Rep protein in PCV1 (Mankertz, A., et al., 1998, *J. Gen. Virol.* 79:381-384). Without wanting to be bound by theory, a higher rate of variation displayed in the ORF2 sequences of PCV1 and PCV2 (identity about 65%) would suggest that type-specific features of PCV might be determined by the respective ORF2 protein. Several PCV type-specific epitopes have been mapped on PCV2 ORF2 sequences. See Mahe, D. et al., 2000, *J. Gen. Virol.* 81:1815-24. In another recent study, PCV2 ORF2 has been identified as a major structural protein that can form viral capsid-like particles in insect cells infected with ORF2 expressing recombinant baculovirus. See Nawagitgul, P. et al., 2000, *J. Gen. Virol.* 81:2281-2287.

In some illustrative embodiments of the invention disclosed herein, a recombinant vector comprising a PCV genome or an ORF thereof, or a portion thereof, such as an antigenic region, is constructed by in vitro recombination between a plasmid and a PCV genome. In some embodiments, the PCV genome is a PCV2 genome. In other embodiments, a recombinant vector comprising a PCV genome or an ORF thereof, or a portion thereof, such as an antigenic region,

is constructed by in vivo recombination. Methods for in vivo recombination are known in the art and include, for example, the methods disclosed in Chartier, et al. (1996, *J. Virol.* 70:48054810). Vectors for constructing circovirus genomes include for example, bacterial plasmids which allow multiple copies of the cloned circovirus nucleotide sequence to be produced. In some embodiments, the plasmid is co-transfected into a suitable host cell for recombination. Suitable host cells for recombination include any cell that will support recombination between a PCV genome and a plasmid containing PCV sequences, or between two or more plasmids, each containing PCV sequences. Recombination is generally performed in prokaryotic cells, such as *E. coli* for example, while generation of circovirus is preferably performed in mammalian cells permissive for PCV replication, such as for example porcine cells and in particular, porcine cells capable of expressing mammalian adenovirus E1 function.

The present invention encompasses the use of any mammalian host cell permissive for circovirus replication, and in particular, permissive for replication of PCV, such as PCV1 and PCV2. Allan et al. (1995, *Veterinary Microbiology* 44:49-64) report that PCV replicate in porcine and bovine monocyte/macrophage cultures. Tischer et al. (1987, *Arch. Virol.* 96:39-57) report that PCV is known to require actively dividing cells for replication in cell culture. Examples of cells or cell lines useful for replication of PCV include mammalian cells comprising E1 function and permissive for PCV replication, including porcine cells, such as porcine monocyte/macrophage cells and porcine retinal cells, expressing adenovirus E1 function. In an illustrative embodiment disclosed herein, porcine retina cells expressing human adenovirus E1 function are shown to be permissive for replication of PCV2 and shown to generate virus at 2×10^7 IU/ml. Porcine cell lines are available from public sources such as for example, the American Type Tissue Collection (ATCC). The growth of bacterial cell cultures, as well as culture and maintenance of eukaryotic cells and mammalian cell lines are procedures well-known to those of skill in the art.

The present invention encompasses methods of culturing mammalian circovirus, in particular, porcine circovirus, in mammalian host cells transfected with mammalian adenovirus E1 gene sequences. In some embodiments, the mammalian cell is stably transformed with adenovirus E1 gene sequences. In some embodiments, the E1 gene sequences are integrated into the genome of the mammalian cell. In other embodiments, the E1 gene sequences are present on a replicating plasmid. In yet other embodiments, the E1 gene sequence is heterologous to the mammalian cell. In an illustrative embodiment disclosed herein a porcine mammalian cell is transformed with a human adenovirus E1 gene sequence. The present invention encompasses the use of any mammalian cell or mammalian cell line expressing E1 function as long as the mammalian cell or cell line expressing E1 function is permissive for the replication of circovirus, in particular porcine circovirus, such as for example, porcine circovirus 1 or porcine circovirus 2. In preferred embodiments, the mammalian cell is a porcine cell or cell line. The present invention encompasses the use of any mammalian E1 function as long as the mammalian host cell expressing the mammalian E1 function is permissive for replication of circovirus, in particular, PCV, such as for example, porcine circovirus 1 or porcine circovirus 2. Mammalian adenovirus genomes are known in the art and are disclosed in, for example, Reddy et al. (1998, *Journal of Virology*, 72:1394) which discloses nucleotide sequence, genome organization, and transcription map of bovine adenovirus 3 (BAV3); and Kleiboeker (1995, *Virus Res.* 36:259-268), which discloses

the E1 region of PAV-4. The present invention encompasses E1 function from any of the various serotypes of human adenovirus, such as Ad2, Ad5, Ad12, and Ad40. In an illustrative embodiment disclosed herein in Example 1, E1 function is human Ad5 E1 function. The human E1A gene is expressed immediately after viral infection (0-2 hours) and before any other viral genes. Flint (1982) *Biochem. Biophys. Acta* 651:175-208; Flint (1986) *Advances Virus Research* 31:169-228; Grand (1987) *Biochem. J.* 241:25-38. The transcription start site of Ad5 E1A is at nucleotide 498 and the ATG start site of the E1A protein is at nucleotide 560 in the virus genome. The E1B protein functions in trans and is necessary for transport of late mRNA from the nucleus to the cytoplasm. The E1B promoter of Ad5 consists of a single high-affinity recognition site for Sp1 and a TATA box. In particular, human adenovirus 5 E1A and E1B gene sequences are located at nucleotides 505-4034 of the nucleotide sequence provided in Chroboezek, J. et al. (1992, *Virology*. 186:280-285). In an illustrative embodiment disclosed herein in the Examples, the mammalian host cell is a porcine host cell transfected with human adenovirus 5 E1 gene sequences.

The PCV genome can be isolated from PCV virions, or can comprise a PCV genome that has been inserted into a plasmid, using standard techniques of molecular biology and biotechnology. Cloning of the full-length PCV2 genome into vector pbluescript II KS(+) from Stratagene by PCR is described in Liu, et al. (2000, *J. Clin. Microbiol.* vol 38:3474-3477). The full-length PCV2 genome DNA can be released from the resulting plasmid upon SacII digestion.

Introduction of circovirus nucleotide sequences into permissive mammalian host cells can be achieved by any method known in the art, including, but not limited to, transfection and transformation including, but not limited to, microinjection, electroporation, CaPO₄ precipitation, DEAE-dextran, liposomes, particle bombardment, etc. An illustrative method for transfecting PCV2 nucleotide sequences into VIDO R1 cells is described herein in Example 3.

Methods for culturing procaryotic cells, such as bacterial cells, and eukaryotic cells, such as mammalian host cells expressing adenovirus E1 function are deemed routine to those of skill in the art.

The following examples are provided to illustrate but not limit the invention. All references and patent publications disclosed herein are hereby incorporated in their entirety by reference.

EXAMPLES

Example 1

Preparation of Porcine Retinal Cells Transfected with Human Adenovirus E1 Gene Sequences (VIDO R1 Cells)

Primary cultures of porcine embryonic retina cells were transfected with 10 µg of plasmid pTG 4671 (Transgene, Strasbourg, France) by the calcium phosphate technique. The pTG 4671 plasmid contains the entire E1A and E1 B sequences (nts 505-4034) of HAV-5, along with the puromycin acetyltransferase gene as a selectable marker. In this plasmid, the E1 region is under the control of the constitutive promoter from the mouse phosphoglycerate kinase gene, and the puromycin acetyltransferase gene is controlled by the constitutive SV40 early promoter. Transformed cells were selected by three passages in medium containing 7 µg/ml puromycin, identified based on change in their morphology from single foci (i.e., loss of contact inhibition), and sub-

jected to single cell cloning. The established cell line was first tested for its ability to support the growth of E1 deletion mutants of HAV-5. Subsequently the cell line was further investigated for the presence of E1 sequences in the genome by PCR, expression of the E1A and E1B proteins by Western blot, and doubling time under cell culture conditions. E1 sequences were detected, and production of E1A and E1B proteins was demonstrated by immunoprecipitation. Doubling time was shorter, when compared to that of the parent cell line.

To assess the stability of E1 expression, VIDO R1 cells were cultured through more than 50 passages (split 1:3 twice weekly) and tested for their ability to support the replication of E1-deleted HAV-5. Expression of the E1A and E1B proteins at regular intervals was also monitored by Western blot. The results indicated that the VIDO R1 line retained the ability to support the growth of E1-deleted virus and expressed similar levels of E1 proteins during more than 50 passages in culture. Therefore, VIDO R1 can be considered to be an established cell line. VIDO R1 cell line has been deposited with the American Type Culture collection (ATCC) and has ATCC accession number PTA-155.

Example 2

Example 2 provides a description of the molecular cloning of full-length PCV2 genome.

Initially, PCV2 DNA was amplified by PCR from total DNA extracted from a piglet with PMWS. The cloning of the full-length PCV2 genome DNA into vector pBluescript II KS(+) (Stratagene) by polymerase chain reaction (PCR) was described in Liu et al. (2000). *J. Clin. Microbiol.* 38:3474-3477). The PCV2 sequence was submitted to GenBank (Accession no. AF086834). The full-length PCV2 genome DNA is released from the resulting plasmid upon SacII digestion.

Example 3

Example 3 describes the transfection of VIDO R1 cells, as described in Example 1, with a plasmid containing the PCV2 genome as constructed in Example 2.

Material and Methods

Cell Culture

Fetal porcine retina cell line, VIDO R1, as described in Example 1 and Vero cells (ATCC) were maintained at 37° C. with 5% CO₂ in Eagles based MEM media supplemented with 10% or 5% heat-inactivated fetal bovine serum (FBS), respectively.

Transfection and Infection

Monolayers of VIDO R1 cells grown in a six-well dish were transfected with cloned PCV2 DNA using Lipofectin according to the manufacturer's recommendations (GIBCO BRL). Prior to transfection, PCV2 full-length genome was released from the plasmid by digestion with SacII (Liu, Q., et al., 2000, *J Clin. Microbiol.* 38:3474-3477). For infection, the transfected VIDO R1 cells were subjected to three cycles of freezing (-70° C.) and thawing (37° C.). The lysate was then clarified by centrifugation and used to infect fresh VIDO R1 cells. In published reports, a PCV1-free porcine kidney cell line is used to culture PCV2 virus. To stimulate the entry to the S phase in the cell cycle, the porcine kidney cells are always treated by D-glucosamine (see Tischer et al., (1987). *Arch Virol.* 96:39-57.). However, the treatment must be performed with caution because D-glucosamine is toxic for cell culture (see, Allan et al., (2000). *J. Vet. Diagn. Investigation.* 12:3-14). In contrast, since the VIDO R1 cell line used in this study

has been transformed by HAV5-E1 that can induce the S phase, the D-glucosamine treatment was not necessary.

Example 4

Virus Purification and Titration

For the purification of PCV2 virus, PCV2-infected VIDO R1 cells were incubated with 0.5% Triton X-114 in phosphate-buffered saline (PBS) at 37° C. for 45 min followed by Freon 113 (1,1,2-trichloro-trifluoroethane) extraction. The cell debris and membranes were clarified by centrifugation at 2000 g for 15 min. The viruses in the supernatant were pelleted at 35000 g for 3 h through a 20% sucrose cushion. The virus pellet was suspended in PBS and stored at -70° C. Virus titers were determined as infectious units (IU) by quantitative ORF2 protein immuno-peroxidase staining. For this purpose, the cell monolayers in 12-well dishes were infected with serial dilutions of virus. After adsorption of virus for 1 h, the cells were washed and overlaid with MEM containing 2% FBS and 0.7% agarose. On day 3 post infection (p.i.), the agarose overlay was removed and the cells were fixed and permeabilized with methanol/acetone (1:1 in volume) for 20 min at -20° C. After blocking with 1% bovine serum albumin for 1 h at room temperature, the cells were incubated with rabbit anti-ORF2 serum (Liu et al., 2001, Protein Expression and Purification. 21:115-120). After 2 h incubation, the plates were washed with PBS and then processed using

VECTASTAIN Elite ABC kit (Vector Laboratories). The reaction was developed with 3,3'diaminobenzidine (DAB) tetrahydrochloride and observed under a microscope. By counting the positively stained cells, the virus titer was expressed as IU where 1 IU was defined as one positively stained cell/foci at 3 d.p.i.

Viral DNA Extraction and Characterization

Viral DNA was extracted from PCV2-infected VIDO RI cell monolayers by the method of Hirt (1967, *J. Mol. Biol.* 26:365-369). The viral DNA was then characterized by restriction analysis and polymerase chain reaction (PCR) as described in Liu et al., (2000). *J. Clin. Microbiol.* 38:3474-3477).

PCR using DNA extracted from the infected cells as template and PCV-2-specific primers amplified a product of specific size, while no DNA was amplified from control, uninfected cells. Consistent with the expected restriction patterns, digestions of viral DNA with NcoI and StuI resulted in two fragments of 1291 bp and 477 bp in size, respectively, digestion with EcoRI and StuI produced two fragments of 1492 bp and 276 bp in size, respectively; and digestion with EcoRI and EcoRV generated two fragments of 1094 bp and 674 bp in size, respectively. The data indicate that PCV2 virus was obtained. Using an immunostaining assay and by counting the positive stained cells, the virus titer of this preparation was determined to be 2×10^7 IU/ml.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 3

<210> SEQ ID NO 1

<211> LENGTH: 1768

<212> TYPE: DNA

<213> ORGANISM: Porcine circovirus

<400> SEQUENCE: 1

```

accagcgcac ttcggcagcg gcagcacctc ggcaacacct cagcagcaac atgccagca      60
agaagaatgg aagaagcggg cccaaccac ataaaagggtg ggtgttcacg ctgaataatc     120
cttccgaaga cgagcgcaag aaaatacggg agctcccaat ctccctatct gattatctta     180
ttgttgccga ggagggtaat gaggaaggac gaacacctca cctccagggg ttcgtaatt      240
ttgtgaagaa gcaaactttt aataaagtga agtggtatct ggggtgccgc tgccacatcg     300
agaaagccaa aggaactgat cagcagaata aagaatattg tagtaagaa ggcaacttac     360
ttattgaatg tggagctcct cgatctcaag gacaacggag tgacctgtct actgctgtga     420
gtacctgtgt ggagagcggg attctggtga ccgttgcaaa gcagcacctc gtaacgcttg     480
tcaaaaattt ccgcccggctg gctgaacttt tgaaagtggc cgggaaaatg caaaagcgtg     540
attgaaaac caatgtacac ttcattgtgg ggccacctgg gtgtggtaaa agcaaatggg     600
ctgctaattt tgcaaacccg gaaaccacat actggaaacc acctaaaaac aagtgggtggg     660
atggttacca tggtgaaaaa gtggttggtt ttgatgactt ttatggctgg ctgccgtggg     720
atgatctact gagactgtgt gatcgatctc cattgactgt aaaaactaaa ggtggaactg     780
tacctttttt ggcccgcagt attctgatta ccagcaatca aaccccgttg gaatggtact     840
cctcaactgc tgtcccagct gtagaagctc tctatcggag gattacttcc ttgggtatctt     900
ggaagaatgt tacagaacaa tccacggagg aagggggcca gtttgtcacc ctttcccccc     960
catgccctga atttccatat gaaataaatt actgagtctt tttatcact tcgtaatggt    1020

```

-continued

```

ttttattatt catttagggt ttaagtgggg ggtctttaag attaaattct ctgaattgta 1080
catacatggt tacacggata ttgtagtcct ggtcgtatct actgttttcg aacgcagtgc 1140
cgaggcctac gtggccaca tttctagagg tttgtagcct cagccaaagc tgattccttt 1200
tgttatttgg ttggaagtaa tcaatagtgg agtcaagaac aggtttgggt gtgaagtaac 1260
gggagtggta ggagaagggt tgggggattg tatggcggga ggagtagttt acatatgggt 1320
cataggttag ggctgtggcc tttgttaca agttatcatc tagaataaca gcagtggagc 1380
ccactcccct atcacctgg gtgatggggg agcaggcca gaattcaacc ttaaccttc 1440
ttattctgta gtattcaaag ggtatagaga tttgttgggt cccccctccc gggggaacaa 1500
agtcgtcaat attaatctc atcatgtcca ccgccagga gggcgttggt actgtggtag 1560
ccttgacagt atatccgaag gtgctgggga ggcgggtgtt gaagatgcca ttttctctc 1620
tccaacggta gcggtggcgg ggggtggacga gccaggggcg gcggcggagg atctggccaa 1680
gatggctgcg ggggctgtgt cttcttctgc ggtaacgct ccttgatac gtcatactg 1740
aaaacgaaag aagtgcgctg taagtatt 1768

```

<210> SEQ ID NO 2

<211> LENGTH: 314

<212> TYPE: PRT

<213> ORGANISM: Porcine circovirus

<400> SEQUENCE: 2

```

Met Pro Ser Lys Lys Asn Gly Arg Ser Gly Pro Gln Pro His Lys Arg
 1          5          10          15
Trp Val Phe Thr Leu Asn Asn Pro Ser Glu Asp Glu Arg Lys Lys Ile
          20          25          30
Arg Glu Leu Pro Ile Ser Leu Phe Asp Tyr Phe Ile Val Gly Glu Glu
          35          40          45
Gly Asn Glu Glu Gly Arg Thr Pro His Leu Gln Gly Phe Ala Asn Phe
          50          55          60
Val Lys Lys Gln Thr Phe Asn Lys Val Lys Trp Tyr Leu Gly Ala Arg
          65          70          75          80
Cys His Ile Glu Lys Ala Lys Gly Thr Asp Gln Gln Asn Lys Glu Tyr
          85          90          95
Cys Ser Lys Glu Gly Asn Leu Leu Ile Glu Cys Gly Ala Pro Arg Ser
          100          105          110
Gln Gly Gln Arg Ser Asp Leu Ser Thr Ala Val Ser Thr Leu Leu Glu
          115          120          125
Ser Gly Ile Leu Val Thr Val Ala Lys Gln His Pro Val Thr Phe Val
          130          135          140
Lys Asn Phe Arg Gly Leu Ala Glu Leu Leu Lys Val Ser Gly Lys Met
          145          150          155          160
Gln Lys Arg Asp Trp Lys Thr Asn Val His Phe Ile Val Gly Pro Pro
          165          170          175
Gly Cys Gly Lys Ser Lys Trp Ala Ala Asn Phe Ala Asn Pro Glu Thr
          180          185          190
Thr Tyr Trp Lys Pro Pro Lys Asn Lys Trp Trp Asp Gly Tyr His Gly
          195          200          205
Glu Lys Val Val Val Ile Asp Asp Phe Tyr Gly Trp Leu Pro Trp Asp
          210          215          220
Asp Leu Leu Arg Leu Cys Asp Arg Tyr Pro Leu Thr Val Lys Thr Lys
          225          230          235          240

```

-continued

Gly Gly Thr Val Pro Phe Leu Ala Arg Ser Ile Leu Ile Thr Ser Asn
 245 250 255
 Gln Thr Pro Leu Glu Trp Tyr Ser Ser Thr Ala Val Pro Ala Val Glu
 260 265 270
 Ala Leu Tyr Arg Arg Ile Thr Ser Leu Val Phe Trp Lys Asn Val Thr
 275 280 285
 Glu Gln Ser Thr Glu Glu Gly Gly Gln Phe Val Thr Leu Ser Pro Pro
 290 295 300
 Cys Pro Glu Phe Pro Tyr Glu Ile Asn Tyr
 305 310

<210> SEQ ID NO 3
 <211> LENGTH: 233
 <212> TYPE: PRT
 <213> ORGANISM: Porcine circovirus

<400> SEQUENCE: 3

Met Thr Tyr Pro Arg Arg Arg Tyr Arg Arg Arg Arg His Arg Pro Arg
 1 5 10 15
 Ser His Leu Gly Gln Ile Leu Arg Arg Arg Pro Trp Leu Val His Pro
 20 25 30
 Arg His Arg Tyr Arg Trp Arg Arg Lys Asn Gly Ile Phe Asn Thr Arg
 35 40 45
 Leu Ser Arg Thr Phe Gly Tyr Thr Val Lys Ala Thr Thr Val Thr Thr
 50 55 60
 Pro Ser Trp Ala Val Asp Met Met Arg Phe Asn Ile Asp Asp Phe Val
 65 70 75 80
 Pro Pro Gly Gly Gly Thr Asn Lys Ile Ser Ile Pro Phe Glu Tyr Tyr
 85 90 95
 Arg Ile Arg Lys Val Lys Val Glu Phe Trp Pro Cys Ser Pro Ile Thr
 100 105 110
 Gln Gly Asp Arg Gly Val Gly Ser Thr Ala Val Ile Leu Asp Asp Asn
 115 120 125
 Phe Val Thr Lys Ala Thr Ala Leu Thr Tyr Asp Pro Tyr Val Asn Tyr
 130 135 140
 Ser Ser Arg His Thr Ile Pro Gln Pro Phe Ser Tyr His Ser Arg Tyr
 145 150 155 160
 Phe Thr Pro Lys Pro Val Leu Asp Ser Thr Ile Asp Tyr Phe Gln Pro
 165 170 175
 Asn Asn Lys Arg Asn Gln Leu Trp Leu Arg Leu Gln Thr Ser Arg Asn
 180 185 190
 Val Asp His Val Gly Leu Gly Thr Ala Phe Glu Asn Ser Lys Tyr Asp
 195 200 205
 Gln Asp Tyr Asn Ile Arg Val Thr Met Tyr Val Gln Phe Arg Glu Phe
 210 215 220
 Asn Leu Lys Asp Pro Pro Leu Lys Pro
 225 230

We claim:

1. A recombinant mammalian cell that expresses a mammalian adenovirus E1 functional protein and comprises a mammalian circovirus genome, or a portion thereof capable of replication, and wherein said cell is permissive for the replication of said circovirus.

60 2. The recombinant mammalian cell of claim 1 wherein Said adenovirus E1 functional protein is human adenovirus E1 function.

65 3. The recombinant mammalian cell of claim 1 wherein The mammalian cell expressing mammalian adenovirus E1 functional protein is stably transformed with a mammalian adenovirus E1 gene sequence.

15

4. The recombinant mammalian cell of claim 3 wherein said E1 gene sequence is a human adenovirus E1 gene sequence.

5. The recombinant mammalian cell of claim 3 wherein said mammalian adenovirus E1 gene sequence is heterolo- 5 gous to said mammalian cell.

6. The recombinant cell of claim 1 wherein said circovirus comprises a heterologous nucleotide sequence.

16

7. The recombinant cell of claim 4 wherein the human adenovirus is human adenovirus 5.

8. A composition comprising the recombinant mammalian cell of claim 1 or 2.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,560,278 B2
APPLICATION NO. : 11/654869
DATED : July 14, 2009
INVENTOR(S) : Qiang Liu et al.

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Cover: item (56),

- In the Foreign Patent Documents, please replace “WO-99/29871 A2 6/1999” with --WO-99/29871 A2 6/1999 English Translation--.
- In the Foreign Patent Documents, please replace “WO-99/29871 A3 6/1999” with --WO-99/29871 A3 6/1999 English Translation--.

In the Specification:

- In column 1, line 26, please replace “Circoviridae” with --*Circoviridae*--.
- In column 1, line 57, please replace “*Can. Vet. J*” with --*Can. Vet. J.*--.
- In column 4, line 5, please replace “Adenoviridae” with --*Adenoviridae*--.
- In column 4, line 6, please replace “Fields Virology” with --Fields Virology--.
- In column 4, line 15, please replace “e.g.” with --*e.g.*--.
- In column 4, line 27, please replace “Circoviridae” with --*Circoviridae*--.
- In column 4, line 31, please replace “supra” with --*supra*--.
- In column 4, line 49, please replace “*vol*” with --*vol*--.
- In column 6, line 1, please replace “in vivo” with --*in vivo*--.
- In column 6, line 4, please replace “70:48054810” with --70:4805-4810--.
- In column 6, line 50, please replace “adenovirus” with --adenovirus 5--.
- In column 7, line 26, please replace “pbluescript” with --pBluescript--.
- In column 7, line 29, please replace “SacII” with --*SacII*--.
- In column 7, line 67, please replace “i.e.” with --*i.e.*--.
- In column 8, line 36, please replace “SacII” with --*SacII*--.


- In column 8, line 43, please replace “Material and Methods” with --Material and Methods--.
- In column 8, line 55, please replace “SacII” with --*SacII*--.
- In column 9, line 6, please replace “Virus Purification and Titration” with --Virus Purification and Titration--.
- In column 10, line 7, please replace “Viral DNA Extraction and Characterization” with --Viral DNA Extraction and Characterization--.

In the Claims:

- In claim 2, line 62, please replace “Said” with --said--.

Signed and Sealed this

Twentieth Day of April, 2010

A handwritten signature in black ink that reads "David J. Kappos". The signature is written in a cursive, flowing style.

David J. Kappos
Director of the United States Patent and Trademark Office

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 7,560,278 B2
APPLICATION NO. : 11/654869
DATED : July 14, 2009
INVENTOR(S) : Qiang Liu et al.

Page 1 of 2

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the Cover: item (56),

- In the Foreign Patent Documents, please replace “WO-99/29871 A2 6/1999” with --WO-99/29871 A2 6/1999 English Translation--.
- In the Foreign Patent Documents, please replace “WO-99/29871 A3 6/1999” with --WO-99/29871 A3 6/1999 English Translation--.


In the Specification:

- In column 1, line 26, please replace “Circoviridae” with --*Circoviridae*--.
- In column 1, line 57, please replace “*Can. Vet. J*” with --*Can. Vet. J.*--.
- In column 4, line 5, please replace “Adenoviridae” with --*Adenoviridae*--.
- In column 4, line 6, please replace “Fields Virology” with --Fields Virology--.
- In column 4, line 15, please replace “e.g.” with --*e.g.*--.
- In column 4, line 27, please replace “Circoviridae” with --*Circoviridae*--.
- In column 4, line 31, please replace “supra” with --*supra*--.
- In column 4, line 49, please replace “*vol*” with --*vol*--.
- In column 6, line 1, please replace “in vivo” with --*in vivo*--.
- In column 6, line 4, please replace “70:48054810” with --70:4805-4810--.

This certificate supersedes the Certificate of Correction issued April 20, 2010.

Signed and Sealed this

Eleventh Day of May, 2010



David J. Kappos
Director of the United States Patent and Trademark Office

- In column 6, line 50, please replace “adenovirus” with --adenovirus 5--.
- In column 7, line 26, please replace “pbluescript” with --pBluescript--.
- In column 7, line 29, please replace “SacII” with --*SacII*--.
- In column 7, line 67, please replace “i.e.” with --*i.e.*--.
- In column 8, line 36, please replace “SacII” with --*SacII*--.
- In column 8, line 43, please replace “Material and Methods” with --Material and Methods--.
- In column 8, line 55, please replace “SacII” with --*SacII*--.
- In column 9, line 6, please replace “Virus Purification and Titration” with --Virus Purification and Titration--.
- In column 10, line 7, please replace “Viral DNA Extraction and Characterization” with --Viral DNA Extraction and Characterization--.

In the Claims:

- In column 14, claim 2, line 62, please replace “Said” with --said--.