

US007745174B2

(12) **United States Patent**
Kim et al.(10) **Patent No.:** **US 7,745,174 B2**
(45) **Date of Patent:** ***Jun. 29, 2010**(54) **HYBRID HEPATOCYTE GROWTH FACTOR GENE HAVING HIGH EXPRESSION EFFICIENCY OF TWO HETEROTYPES OF HEPATOCYTE GROWTH FACTOR**(75) Inventors: **Jong-Mook Kim**, Seoul (KR); **Woong Hahn**, Goyang (KR); **Eun-Jin Park**, Seoul (KR)(73) Assignee: **Viomed Co., Ltd.**, Seoul (KR)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **12/650,860**(22) Filed: **Dec. 31, 2009**(65) **Prior Publication Data**

US 2010/0105878 A1 Apr. 29, 2010

Related U.S. Application Data

(63) Continuation of application No. 10/944,277, filed on Sep. 20, 2004, which is a continuation of application No. PCT/KR03/00548, filed on Mar. 20, 2003.

(30) **Foreign Application Priority Data**

Mar. 20, 2002 (KR) 10-2002-0015074

(51) **Int. Cl.****C12P 21/06** (2006.01)**C12P 21/04** (2006.01)**C12N 15/09** (2006.01)(52) **U.S. Cl.** **435/69.1**; 435/69.4; 435/70.1; 435/71.1(58) **Field of Classification Search** None
See application file for complete search history.(56) **References Cited**

U.S. PATENT DOCUMENTS

5,328,836 A 7/1994 Shima et al.
5,500,354 A 3/1996 Kitamura et al.
5,580,859 A 12/1996 Felgner et al.
5,587,359 A 12/1996 Higashio et al.
5,652,225 A 7/1997 Isner
5,693,622 A 12/1997 Wolff et al.
6,013,624 A 1/2000 Goldberg et al.
6,121,246 A 9/2000 Isner
6,248,722 B1 6/2001 Morishita et al.
6,258,787 B1 7/2001 Isner
6,316,419 B1 11/2001 Leiden et al.
6,413,942 B1 7/2002 Felgner et al.
6,498,144 B1 12/2002 Goldberg et al.
6,706,694 B1 3/2004 Wolff et al.
6,887,477 B1 5/2005 Nagano et al.
7,285,540 B2 10/2007 Morishita et al.
2002/0172663 A1 11/2002 Palasis
2003/0148968 A1 8/2003 Hammond et al.
2003/0171287 A1 9/2003 Morishita et al.2004/0105882 A1 6/2004 Morishita et al.
2004/0228834 A1 11/2004 Isner et al.
2005/0079581 A1 4/2005 Kim et al.
2006/0286072 A1 12/2006 Giordano et al.
2007/0059288 A1 3/2007 Dinsmore et al.
2008/0268030 A1 10/2008 Morishita et al.
2009/0004260 A1 1/2009 Morishita et al.
2009/0082293 A1 3/2009 Giordano et al.
2009/0131350 A1 5/2009 Kim et al.
2009/0202606 A1 8/2009 Kim et al.

FOREIGN PATENT DOCUMENTS

JP 11-246433 9/1999
WO WO 98/50079 A2 11/1998
WO WO 99/45775 A1 9/1999
WO WO 01/34208 A1 5/2001
WO WO 02/089856 A1 11/2002

OTHER PUBLICATIONS

Deng, et al., "Secretory Expression of the Deleted Variant of Human Hepatocyte Growth Factor (hdHGF) in *Pichia pastoris*," *Chinese Journal of Biochemistry and Molecular Biology*, 2001, 17:590-594, China Academic Journal Electronic Publishing House, Beijing, China.Kisselev, L., "Polypeptide Release Factors in Prokaryotes and Eukaryotes: Same Function, Different Structure," *Structure*, 2002, 10, pp. 8-9, Elsevier Science Ltd., Cambridge, Massachusetts, USA.
Liu, Y., "The human hepatocyte growth factor receptor gene: complete structural organization and promoter characterization," *Gene* 215:159-169, Elsevier/North-Holland (1998).Miyazawa, K., et al., "Molecular Cloning and Sequence Analysis of cDNA for Human Hepatocyte Growth Factor," *Biochem. Biophys. Res. Commun.* 163:967-973, Academic Press (1989).Nakamura, T., et al., "Molecular cloning and expression of human hepatocyte growth factor," *Nature* 342:440-443, Nature Publishing Group (1989).NCBI Entrez, GenBank Database, Accession No. AC004960, "*Homo sapiens* PAC clone RP5-1098B1 from 7q11.23-q21, complete sequence," 51 pages (first available 1998).Ngo et al., In *The Protein Folding Problem and Tertiary Structure Prediction*, 1994, Merz et al. (ed.), Birkhauser, Boston, MA, pp. 433 and 492-495.Potrykus. Gene transfer to cereals: an assessment, *Biotechnology*, 1990, 8(6):535-542.Rubin, et al., "A broad-spectrum human lung fibroblast-derived mitogen is a variant of hepatocyte growth factor," *Proc. Natl. Acad. Sci. USA*, 1991, 88:415-419, Proceedings of the National Academy of Sciences of the United States of America, 500 5th St., Washington, DC 20001.

(Continued)

Primary Examiner—Suzanne M. Noakes*Assistant Examiner*—Jae W Lee(74) *Attorney, Agent, or Firm*—Sterne, Kessler, Goldstein & Fox P.L.L.C.(57) **ABSTRACT**

The present invention relates to a hybrid Hepatocyte Growth Factor (HGF) gene which is prepared by inserting an inherent or foreign intron between exons 4 and 5 in HGF cDNA, which has a base sequence of SEQ ID NO: 2. The gene has high expression efficiency and simultaneously expresses two heterotypes of HGF and dHGF (deleted variant HGF). Further the gene may be used for treating or preventing ischemic or liver diseases.

21 Claims, 12 Drawing Sheets

OTHER PUBLICATIONS

Seki, T., et al., "Isolation and Expression of cDNA for Different Forms of Hepatocyte Growth Factor from Human Leukocyte," *Biochem. Biophys. Res. Commun.* 172:321-327, Academic Press (1990).

Seki, T., et al., "Organization of the human hepatocyte growth factor-encoding gene," *Gene* 102:213-219, Elsevier/North-Holland (1991).

Shima, N., et al., "Hepatocyte Growth Factor and its Variant with a Deletion of Five Amino Acids are Distinguishable in their Biological Activity and Tertiary Structure," *Biochem. Biophys. Res. Commun.* 200:808-815, Academic Press (1994).

Warnecke C. et al., "Efficient transcription of the human angiotensin II type 2 receptor gene requires intronic sequence elements," *Biochemical journal*, 1999, 340 (1), pp. 17-24, Portland Press, Colchester, Great Britain.

Wishart et al., "A Single Mutation Converts a Novel Phosphotyrosine Binding into a Dual-Specificity Phosphatase," *Journal of Biological Chemistry*, 1995, 270 (45), pp. 26782-26785, American Society for Biochemistry and Molecular Biology, Bethesda, MD, USA.

Witkowski et al., "Conversion of a A-Ketoacyl Synthase to a Malonyl Decarboxylase by Replacement of the Active-Site Cysteine with Glutamine," *Biochemistry*, 1999, 38 (36), pp. 11643-11650, American Chemical Society, Washington, DC, USA.

European Search Report for European Application No. EP 03 74 4561, mailed Apr. 18, 2006, European Patent Office, Munich, Germany.

Office Action for Co-pending U.S. Appl. No. 10/944,277, mailed Jan. 9, 2008.

Office Action for Co-Pending U.S. Appl. No. 10/944,277, mailed Feb. 13, 2009.

Notice of Allowance and Fees Due for Co-Pending U.S. Appl. No. 10/944,277, mailed May 29, 2009.

Notice of Allowance and Fees Due for Co-Pending U.S. Appl. No. 10/944,277, mailed Oct. 23, 2009.

Office Action for Co-pending U.S. Appl. No. 11/957,170, mailed Jan. 28, 2010.

Esp@cenet Database, English language abstract of JP 11-246433 A, published Sep. 14, 1999 (listed as document FP5 on the accompanying form PTO/SB/08A).




FIG. 2






FIG. 3








FIG. 5




FIG. 6




FIG. 7




FIG. 8





FIG. 9



FIG. 10






FIG. 11

**HYBRID HEPATOCYTE GROWTH FACTOR
GENE HAVING HIGH EXPRESSION
EFFICIENCY OF TWO HETEROTYPES OF
HEPATOCYTE GROWTH FACTOR**

This application is a continuation of U.S. application Ser. No. 10/944,277, filed Sep. 20, 2004, which is a continuation of International Application No. PCT/KR03/00548, filed Mar. 20, 2003, which claims priority benefit to Korean Appl. No. 10-2002-0015074, filed Mar. 20, 2002, each of which are herein incorporated by reference in their entirety.

REFERENCE TO A SEQUENCE LISTING
SUBMITTED ELECTRONICALLY

The content of the electronically submitted sequence listing ("sequencelisting.ascii.txt", 29,879 bytes, created on Dec. 30, 2009) filed with the application is incorporated herein by reference in its entirety.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to a highly efficient hybrid Hepatocyte Growth Factor (HGF) gene which simultaneously expresses two heterotypes of HGF.

2. Related Art

The present invention relates to a hybrid HGF gene prepared by inserting an inherent or foreign intron between exons 4 and 5 in HGF cDNA, which has higher expression efficiency than HGF cDNA and simultaneously expresses two heterotypes of HGF and dHGF (deleted variant HGF).

HGF is a heparin binding glycoprotein called a scatter factor. A gene encoding HGF is located at chromosome 721.1 and comprises 18 exons and 17 introns, having the nucleotide sequence of SEQ ID NO: 1 (Seki T., et al., *Gene* 102:213-219 (1991)). A transcript of about 6 kb is transcribed from the HGF gene, and then, a polypeptide HGF precursor consisting of 728 amino acids is synthesized therefrom. Simultaneously, a polypeptide of dHGF precursor consisting of 723 amino acids is also synthesized by an alternative splicing of the HGF gene. The biologically inactive precursors may be converted into active forms of disulfide-linked heterodimer by protease in serum. In the heterodimers, the alpha chain having a high molecular weight forms four kringle domains and an N-terminal hairpin loop like a preactivated peptide region of plasminogen. The kringle domains of a triple disulfide-bonded loop structure consisting of about 80 amino acids may play an important role in protein-protein interaction. The low molecular weight beta chain forms an inactive serine protease-like domain. dHGF consisting 723 amino acids is a polypeptide with deletion of five amino acids in the 1st kringle domain of the alpha chain, i.e., F, L, P, S and S.

It has been recently reported that both of HGF and dHGF have several biological functions, e.g., promoting the growth and morphogenesis of epithelial cell, melanocyte and endothelial cell. However, they are different in terms of immunological or biological properties.

For example, HGF shows about 20-fold, 10-fold and 2-fold higher activities than dHGF in promoting DNA synthesis in human umbilical cord venous endothelial cell, arterial smooth muscle cell and NSF-60 (murine myeloblast cell), respectively. dHGF shows about 3-fold and 2-fold higher activities than HGF in promoting DNA synthesis of LLC-PK1 (pig kidney epithelial cell), and OK (American opossum kidney epithelial cell) and mouse interstitial cell, respectively. HGF has a 70-fold higher solubility in PBS than dHGF.

Several anti-dHGF monoclonal antibodies recognize only dHGF, but not HGF or a reduced form of dHGF, which implies structures of HGF and dHGF are different. Accordingly, the simultaneous synthesis of HGF and dHGF in vivo suggests that they biologically interact with each other (Shima, N. et al., *Biochemical and Biophysical Research Communications* 200:808-815 (1994)).

HGF secreted from mesoderm-derived cells has various biological functions, e.g., 1) inducing epithelial cells into a tubular structure; 2) stimulating vascularization from endothelial cells in vitro and in vivo; 3) regeneration of liver and kidney, owing to its anti-apoptosis activity; 4) organogenesis of kidney, ovary and testis; 5) controlling osteogenesis; 6) stimulating the growth and differentiation of erythroid hematopoietic precursor cells; and 7) axon sprouting of neurons (Stella, M. C. and Comoglio, P. M., *The International Journal of Biochemistry & Cell Biology* 31:1357-1362 (1999)). Based on these various functions, HGF or a gene encoding HGF may be developed as a therapeutic agent for treating ischemic or liver diseases. Actually, in vivo, the HGF may exist as either HGF or dHGF, and therefore, the coexpression of HGF and dHGF is important for maximizing the therapeutic effect. Accordingly, the present inventors have endeavored to develop a hybrid HGF gene which can simultaneously express HGF and dHGF with a high efficiency for gene therapy.

SUMMARY OF THE INVENTION

Accordingly, it is a primary object of the present invention to provide a hybrid HGF gene which simultaneously expresses two heterotypes of HGF.

In accordance with one aspect of the present invention, there is provide the hybrid HGF gene having an inherent or foreign intron is inserted between exons 4 and 5 of HGF cDNA.

It is a another object of the present invention to provide a recombinant vector comprising the hybrid HGF gene and a microorganism transformed with the above vector.

It is a still further object of the present invention to provide a pharmaceutical composition for treating or preventing ischemic or liver diseases, which comprises the HGF gene.

BRIEF DESCRIPTION OF THE DRAWINGS

The above and other objects and features of the present invention will become apparent from the following description of the invention, when taken in conjunction with the accompanying drawings which respectively show:

FIG. 1: a schematic diagram of HGF-X prototype illustrating the positions of the gene fragments,

FIG. 2: a process for cloning gene fragments from HepG2 genomic DNA,

FIG. 3: a process for cloning gene fragments from human placenta cDNA,

FIGS. 4A and 4B: processes for preparing expression vectors pCK-HGF-X,

FIG. 5: a process for preparing expression vectors pCK-cHGF and pCK-dHGF,

FIG. 6: a process for preparing expression vectors pCP-HGF-X family,

FIG. 7: a process for preparing expression vectors pCP-cHGF and pCP-dHGF,

FIG. 8: gene expression levels of pCP-cHGF, pCP-dHGF and pCP-HGF-X.

FIG. 9: gene expression patterns of pCP-cHGF, pCP-dHGF and pCP-HGF-X observed by electrophoresis on 12% polyacrylamide gel,

FIG. 10: gene expression levels of pCP-cHGF, pCP-dHGF and pCP-HGF-X7, in vivo,

FIG. 11: cerebral angiogenesis of two groups of rabbits which were subject to administering pCP and pCP-HGF-X7, respectively,

DETAILED DESCRIPTION OF THE INVENTION

The hybrid Hepatocyte Growth Factor (HGF) gene of the present invention comprises cDNA corresponding to the exons 1 to 18, and an inherent or foreign intron inserted between exons 4 and 5 of the cDNA. The intron comprises a fragment of the inherent intro or a recombinant sequence.

An embodiment of the hybrid HGF gene of the present invention comprising the inherent intron is 7113 bp long and has the nucleotide sequence of SEQ ID NO: 2. The hybrid HGF gene simultaneously expresses both HGF and dHGF, and has higher expression efficiency than HGF cDNA.

Codon degeneracy enables the hybrid HGF gene of the present invention to be modified or changed in the coding and/or non-coding region without altering the amino acid sequence of the protein and the expression of the gene. Accordingly, polynucleotides which is substantially identical to the hybrid HGF gene of SEQ ID NO:2, and the fragments thereof fall within the scope of the invention. "Substantially identical" means that the sequence homology is not less than 80%, preferably not less than 90%, and more preferably not less than 95%.

A hybrid HGF gene may comprise a fragment of inherent intron optionally having a small recombinant sequence inserted thereinto between exons 4 and 5 of HGF cDNA. Herein, such a hybrid HGF gene comprising a fragment of inherent intron designates "HGF-X". HGF-X6, HGF-X7 and HGF-X8 having the nucleotide sequence of SEQ ID Nos: 19 to 21, respectively, are preferred.

The hybrid HGF gene of the present invention is synthesized and inserted into an expression vector, according to the known genetic engineering methods. Then, the vector can be introduced into an appropriate host cells such as *E. coli* and yeast. For example, *Escherichia coli* Top10F' may be transfected with HGF-X7 gene of the present invention. *Escherichia coli* Top10F' pCK-HGFX7 and *Escherichia coli* Top10F' pCP-HGFX7 then obtained were deposited as the accession numbers KCCM-10361 and KCCM-10362, respectively, on Mar. 12, 2002.

By using the transformed cells, the gene of the present invention and the protein encoded thereby may be produced on a large scale.

The vector of the present invention may selectively comprise sequence(s) for regulating gene expression such as promoter or terminator, self-replication sequence and secretory signal, depending on host cells.

Further, the present invention comprises a pharmaceutical composition for treating or preventing ischemic and liver diseases, which comprises the hybrid HGF gene or the vector comprising the gene as an active ingredient. Preferably, the composition is formulated for injection.

The composition of the present invention may further comprise pharmaceutically acceptable carriers. Any of the conventional procedures in the pharmaceutical field may be used to prepare oral formulations such as tablets, capsules, pills, granules, suspensions and solutions; rejection formulations such as solutions, suspensions, or dried powders that may be

mixed with distilled water before injection; locally-applicable formulations such as ointments, creams and lotions; and other formulations.

Carriers generally used in the pharmaceutical field may be employed in the composition of the present invention. For example, orally-administered formulations may include binders, emulsifiers, disintegrating agents, excipients, solubilizing agents, dispersing agents, stabilizing agents, suspending agents, coloring agents or spicery. Injection formulations may comprise preservatives, unagonizing agents, solubilizing agents or stabilizing agents. Preparation for local administration may contain bases, excipients, lubricants or preservatives. Any of the suitable formulations known in the art (Remington's Pharmaceutical Science [the new edition], Mack Publishing Company, Eaton Pa.) may be used in the present invention.

The inventive composition can be clinically administered as various oral and parenteral formulations. A suitable formulation may be prepared using such excipients as additives, enhancers, binders, wetting agents, disintegrating agents and surfactants, or diluents. Solid formulations for oral administration include pills, tablets, dusting powder, granules and capsules. Those solid formulations may be prepared by mixing one or more excipients, e.g. starch, calcium carbonate, sucrose, lactose and gelatin with dibenzylbutyllacton lignan derivatives. Also, lubricants such as magnesium stearate and talc may be included in the present formulation. Liquid formulations for oral administration include suspension, solution, emulsion and syrup. Those formulations may contain wetting agents, sweeteners, aromatics and preservatives, in addition to general simple diluents such as water and liquid paraffin. Formulations for parenteral administration include sterilized aqueous solution, suspension, emulsion, freeze-dried alternative treatment and suppositories. Water-insoluble excipients and suspending agents comprise vegetable fats such as propylene glycol, polyethylene glycol and olive oil, and injectable esters such as ethyl oleate. Witepsol®, Macrogol®, Tween® 61, cacao fats, laurin fats and glycerogelatin may be used as bases of suppositories.

The inventive composition may be administered orally or via parenteral routes such as intravenous, intramuscular, subcutaneous, intraabdominal, sternal and arterial injection or infusion, or topically through rectal, intranasal, inhalational or intraocular administration.

It should be understood that the typical daily dose of composition of the present invention ought to be determined in light of various relevant factors including the conditions to be treated, the chosen route of administration, the age, sex and body weight of the individual patient, and the severity of the patient's symptom, and can be administered in a single dose or in divided dose. Therefore, the daily dose should not be construed as a limitation to the scope of the invention in any way.

The following Examples are given for the purpose of illustration only, and are not intended to limit the scope of the invention.

EXAMPLE 1

Preparation of Hybrid Gene Constructs Encoding Human HGF

(1) Cloning of HGF Gene Fragments Obtained from Genomic DNA

Human HepG2 cells (ATCC Accession NO: HB-8065) were suspended in TES buffer (10 mM Tris-HCl; 1 mM

EDTA; 0.7% SDS) and treated with 400 µg/ml of proteinase K at 50° C. for 1 hour. Subsequently, genomic DNA was extracted from the cell suspension by phenol/chloroform extraction and ethanol precipitation according to the conventional method in the art.

In the PCR amplification, the extracted genomic DNA was employed as a template DNA. As primer pairs, the synthetic nucleotides of SEQ ID NOs: 3 and 4 were employed to obtain DNA fragments containing: HGF gene fragment 2 (HGF-F2), SEQ ID NOs: 3 and 5; HGF-F3, SEQ ID NOs: 6 and 7; HGF-F5, SEQ ID NOs: 8 and 7; HGF-F7, SEQ ID NOs: 9 and 7; HGF-F8, SEQ ID NOs: 10 and 7; HGF-F6, respectively (FIG. 1). The PCR amplification mixture was prepared by mixing 1 µl of template DNA, 1 µl each of primer (10 pmol/µl), 10 µl of dNTP (10 mM), 3.5 unit of Expand High Fidelity enzyme (Gibco BRL, USA) and 10 µl of enzyme buffer solution and adjusted to a final volume of 100 µl with distilled water. 30 cycles of the PCR amplification was carried out, each cycle consisting of 1 min at 94° C., 1 min at 55° C. and 30 sec at 72° C. The primers used herein and the amplified gene fragments obtained therefrom are shown in Table 1.

TABLE 1

5' primer	3' primer	Amplified fragment
gHGF3 (SEQ ID NO: 3)	gHGF4 (SEQ ID NO: 4)	HGF gene fragment 2 (HGF-F2)
gHGF3 (SEQ ID NO: 3)	gHGF10 (SEQ ID NO: 5)	HGF gene fragment 3 (HGF-F3)
gHGF5 (SEQ ID NO: 6)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 5 (HGF-F5)
gHGF12 (SEQ ID NO: 8)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 7 (HGF-F7)
gHGF13 (SEQ ID NO: 9)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 8 (HGF-F8)
gHGF6 (SEQ ID NO: 10)	gHGF7 (SEQ ID NO: 7)	HGF gene fragment 6 (HGF-F6)

The amplified HGF-F2 comprised the sequence ranging from 392 to 2247 of human HGF cDNA prototype (HGF-X1; composed of exons 1 to 4-intron 4-exons 5 to 18) of SEQ ID NO: 2; HGF-F3, the sequence ranging from 392 to 727; HGF-F5, the sequence ranging from 2229 to 5471; HGF-F6, the sequence ranging from 5117 to 5471; HGF-F7, the sequence ranging from 3168 to 5471; and HGF-F8, the sequence ranging from 4168 to 5471.

The amplified HGF gene fragments were each inserted into the multiple cloning site of pGEM-T easy vector (Promega, WI, USA) to obtain pGEM-T easy-HGF-F2, pGEM-T easy-HGF-F3, pGEM-T easy-HGF-F5, pGEM-T easy-HGF-F6, pGEM-T easy-HGF-F7 and pGEM-T easy-HGF-F8, respectively (FIG. 2). The nucleotide sequences of the amplified HGF gene fragments were confirmed by a sequence analysis.

(2) Cloning of HGF Gene Fragments Obtained from cDNA

In the PCR amplification, human placenta cDNA (Clontech, CA, USA) was employed as a template DNA under the same condition as described in Example 1. As primer pairs, the synthetic oligonucleotides of SEQ ID NOs: 11 and 12, and SEQ ID NOs: 13 and 14 were employed to obtain DNA fragments containing HGF-F1 and HGF-F4, respectively. Further, DNA fragments containing cDNAs of HGF gene (cHGF) and deleted HGF gene (dHGF) were amplified by PCR using synthetic oligonucleotides of SEQ ID NOs: 15 and 16 as a primer pair, respectively. dHGF is a HGF gene with deletion of 5 base sequences.

The primers used herein and the amplified gene fragments obtained therefrom are shown in Table 2.

TABLE 2

5' primer	3' primer	Amplified fragment
gHGF1 (SEQ ID NO: 11)	gHGF2 (SEQ ID NO: 12)	HGF gene fragment 1 (HGF-F1)
gHGF8 (SEQ ID NO: 13)	gHGF9 (SEQ ID NO: 14)	HGF gene fragment 4 (HGF-F4)
cHGF5 (SEQ ID NO: 15)	cHGF3 (SEQ ID NO: 16)	HGF gene cDNA (cHGF)
		dHGF gene cDNA (dHGF)

The amplified HGF-F1 and HGF-F4 comprised the nucleotide sequences ranging from 1 to 402 and from 6533 to 7113 of SEQ ID NO: 2 of human HGF cDNA prototype, respectively. HGF gene cDNA comprised the nucleotide sequence ranging from 1 to 2184 of SEQ ID NO: 1 of human HGF gene, and dHGF gene cDNA has the same sequence as HGF gene cDNA except for the deletion of the sequence ranging from 483 to 495.

The amplified fragments of HGF gene were each inserted into the multiple cloning site of pGEM-T easy vector (Promega, WI, USA) to obtain pGEM-T easy-HGF-F1, pGEM-T easy-HGF-F4, pGEM-T easy-cHGF and pGEM-T easy-dHGF, respectively (FIG. 3). The nucleotide sequences of the human HGF gene fragments, HGF gene cDNA and dHGF gene cDNA were confirmed by sequence analyses.

(3) Preparation of Hybrid HGF Gene Constructs

Hybrid HGF gene constructs of genomic DNA and cDNA were prepared by combining the fragments of HGF gene cloned in steps (1) and (2) as follows (FIGS. 4A and 4B).

Plasmid pGEM-T-easy-HGF-F1 was treated with HindIII/BamHI to obtain HGF-F1. Plasmid pCK (see PCT International Publication NO: WO/0040737) was treated with HindIII/BamHI, and HGF-F1 was inserted therein to obtain pCK-F1. And then, plasmids pGEM-T-easy-HGF-F2 and pGEM-T-easy-HGF-F3 were treated with MluI/BamHI to obtain HGF-F2 and HGF-F3, respectively. pCK-1 was treated with MluI/BamHI, and then HGF-F2 and HGF-F3 were inserted therein to obtain pCK-F12M and pCK-F13M. The MluI restriction site of vectors pCK-F12M and pCK-F13M was substituted with an HgaI restriction site by employing a site-directed mutagenesis kit (Stratagene, CA., USA) to obtain pCK-F12 and pCK-F13, respectively.

Further, plasmid pGEM-T-easy-HGF-F4 was treated with BamHI/XbaI to obtain HGF-F4. pCK-F12 and pCK-F13 were treated with BamHI/XbaI, and HGF-F4 was inserted therein to obtain pCK-F124 and pCK-F134, respectively. And then, plasmids pGEM-T-easy-HGF-F5, pGEM-T-easy-HGF-F6, pGEM-T-easy-HGF-F7 and pGEM-T-easy-HGF-F8 were treated with BamHI/XhoI to obtain HGF-F5, HGF-F6, HGF-F7 and HGF-F8, respectively. pCK-F124 and pCK-F134 were treated with BamHI/XhoI, and then HGF-F5, HGF-F6, HGF-F7 and HGF-F8 were inserted therein to obtain pCK-F1254 and pCK-F1264, pCK-F1274, pCK-F1284, pCK-F1354, pCK-F1364, pCK-F1374 and pCK-F1384, respectively.

And then, pGEM-T easy-cHGF was treated with XhoI to obtain cDNA fragment of HGF gene (HGF-XhoI) of about 1100 bp. Then, HGF-XhoI was inserted into pCK-F1254, pCK-F1264, pCK-F1274, pCK-F1284, pCK-F1354, pCK-F1364, pCK-F1374 and pCK-F1384 to obtain pCK-HGF-X1, pCK-HGF-X2, pCK-HGF-X3, pCK-HGF-X4, pCK-HGF-X5, pCK-HGF-X6, pCK-HGF-X7 and pCK-HGF-X8, respectively. On the other hand, pGEM-T easy-cHGF and pGEM-T easy-dHGF were treated with BamHI to obtain HGF gene cDNA and dHGF gene cDNA. Then, HGF gene

cDNA and dHGF gene cDNA were inserted into the BamHI restriction site of pCK to obtain pCK-cHGF and pCK-dHGF, respectively (FIG. 5).

(4) Preparation of an Expression Vector Containing a Hybrid HGF Gene Construct

Plasmid pcDNA3.1 (Invitrogen, USA) was digested with NdeI, treated with the Klenow fragment to build blunt ends, and then digested with NheI to obtain a DNA fragment containing human cytomegalovirus promoter. Plasmids pCK-HGF-X1, pCK-HGF-X2, pCK-HGF-X3, pCK-HGF-X4, pCK-HGF-X5, pCK-HGF-X6, pCK-HGF-X7 and pCK-HGF-X8 were digested with SnaBI, treated with the Klenow fragment to make blunt ends and digested with NheI, and then the above DNA fragment containing human cytomegalovirus promoter was inserted thereto to obtain pCP-HGF-X1, pCP-HGF-X2, pCP-HGF-X3, pCP-HGF-X4, pCP-HGF-X5, pCP-HGF-X6, pCP-HGF-X7 and pCP-HGF-X8, respectively (FIG. 6).

Plasmid pcDNA3.1 (Invitrogen, USA) was digested with NheI, treated with the Klenow fragment to make blunt ends and digested with NdeI to obtain the DNA fragment containing human cytomegalovirus promoter. pCK-cHGF and pCK-dHGF were digested with MluI, treated with the Klenow fragment to make blunt ends and digested with NdeI, and then the above DNA fragment containing human cytomegalovirus promoter was inserted thereto to obtain pCP-cHGF and pCP-dHGF, respectively (FIG. 7).

EXAMPLE 2

Examination of the Expression Efficiency of Hybrid HGF Gene Construct and the Coexpression of HGF/dHGF

Studies was conducted to examine whether the hybrid HGF gene constructs (HGF-X1 to HGF-X8) obtained in Example 1 can simultaneously express HGF and dHGF and whether there is any difference in the gene expression level between hybrid HGF gene constructs and HGF cDNA.

(1) Gene Expression Efficiency

First, 5 μ g of pCP-HGF-X2, pCP-HGF-X3, pCP-HGF-X6, pCP-HGF-X7 and pCP-HGF-X8 were transfected into 5×10^6 cells of 293 cell (ATCC CRL 1573) together with 0.5 μ g of DONAI-LacZ (TAKARA SHUZO, Japan) DNA using FuGENE6 (Gibco BRL, MD, USA), according to the manufacturer's instructions. At this time, 5 μ g each of pCP-cHGF and pCP-dHGF were used as controls, and DONAI-LacZ DNA was used to calibrate the infection efficiency. 3 hours after transfection, cells were re-fed with a fresh medium and further cultured for 48 hours. The culture solution thus obtained was divided into two parts. One part of the 293 cell culture solution was subjected to RNA extraction, and the other, to measurement of LacZ activity. The LacZ activity was measured using an activity measuring kit (Stratagene, CA, USA) according to the manufacturer's instructions.

In order to compare the gene expression levels, the amount of HGF in the cell culture was measured by an enzyme-linked immunosorbent assay kit (ELISA, R&D System, MN, USA). After calibrating the infection efficiency by the measured LacZ activity, the expression level of HGF-X gene was found to be from 20 to 150-fold higher than those of HGF cDNA and dHGF cDNA (FIG. 8). HGF-X7, in particular, showed the highest gene expression level.

(2) Coexpression of HGF and dHGF

In order to examine coexpression of HGF and dHGF from hybrid HGF gene constructs, total cellular RNAs were extracted from the transfected 293 cells using the Trizol method (Trizol; Gibco BRL, USA) and subjected to RT-PCR to obtain cDNA. Then, using cDNA as a template DNA, PCR amplification was carried out using synthetic oligonucleotides of SEQ ID NOs: 17 and 18 as a primer pair. The PCR amplification mixture was prepared by mixing 1 μ l of the template DNA, 1 μ l each of the primer (10 pmol/ μ l), 10 μ l of dNTP (10 mM), 3.5 unit of Taq polymerase (TAKARA SHUZO, Japan) and 10 μ l of enzyme buffer solution and adjusted to a final volume of 100 μ l with distilled water. 30 cycles of PCR amplification was conducted, each cycle consisting of 1 min at 94° C., 1 min at 55° C., and 90 sec at 72° C.

The amplified PCR products corresponded to the boundary region between exons 4 and 5 of HGF gene; HGF gene cDNA of 142 by and dHGF gene cDNA of 127 bp, respectively. With no splicing, the PCR product of at least 1 kb in length was amplified; and if alternative splicing occurred, HGF gene cDNA of 142 bp and dHGF gene cDNA of 127 by were simultaneously synthesized and amplified. The amplified PCR products were distinguished by electrophoresis on a 12% polyacrylamide gel.

As shown in FIG. 9, while the bands of 142 by and 127 by were detected in the lanes loading HGF gene cDNA and dHGF gene cDNA, respectively, both bands of 142 by and 127 by were detected in the lanes loading HGF-X. The above results suggest that HGF and dHGF are simultaneously expressed from hybrid HGF-X gene constructs of the present invention.

EXAMPLE 3

Comparison of Expression Levels of HGF-X7, HGF Gene cDNA and dHGF Gene cDNA (In Vivo)

100 μ g each of pCP-HGF-X7, pCP-cHGF and pCP-dHGF were injected into the anterior tibial muscle of the hind limb of mice with an insulin syringe. After 5 days, the mice were sacrificed and the muscles around the injection spot were removed and smashed in a protein extraction buffer (25 mM Tris-HCl (pH 7.4), 50 mM NaCl, 0.5% Na-deoxycholate, 2% NP-40, 0.2% SDS) to separate total proteins. The amount of the total proteins was measured with a DC protein analysis kit (Bio-Rad Laboratories, CA, USA) and the amount of expressed HGF was determined with an ELISA kit (R&D System) according to the manufacturer's instruction.

As can be seen from the result shown in FIG. 10, the amount of HGF expressed from HGF-X7 is 250-fold higher than that from HGF gene cDNA or dHGF gene cDNA.

Together with the result of the experiment of Example 2 (in vivo), this result demonstrates that the expression efficiency of HGF-X gene is much superior to those of HGF gene cDNA or dHGF gene cDNA.

EXAMPLE 4

Gene Therapy Employing HGF-X7 in a Rabbit Ischemic Hind Limb Model

In order to examine whether HGF-X7 gene is effective in the treatment of ischemic hind limb disease, gene therapy was carried out on a rabbit ischemic hind limb model as follows.

A rabbit ischemic hind limb model, which is a standard animal model for the ischemic limb disease, was prepared by the method described by Takeshita et al., *Journal of Clinical*

Investigation 93:662 (1994). At the day before operation (day 0), each of 30 white rabbits from New Zealand (male, from 3.8 to 4.2 kg) was intramuscularly injected with 5 mg/kg of xylazine and, then, anesthetized by an intramuscular injection of 50 mg/kg of ketamine. Subsequently, the left femoral region of the rabbit was incised and all branches of the femoral artery were separated and tied. The region from the proximal part to the branching point of the saphenous and popliteal arteries was incised to prepare the model. After the operation, 15 mg/kg/day of cefazolin was injected intramuscularly for 5 days and 0.3 mg/day of morphine, for 10 days. 10 days after the operation (day 10), angiography was carried out for the left hind limb where the ischemia was induced, and the degree of arteriogenesis was recorded as a basal level. The rabbits were randomly divided into two groups and injected at four sites in the femoral muscle with 500 µg of plasmid pCP-HGF-

X7 (experimental group) or 500 µg of plasmid pCP (control), respectively. 40 days after the operation (day 40), angiography was carried out again for the left hind limb and the degree of arteriogenesis at the arteriole level was determined and compared to that of day 10.

As can be seen from the result in FIG. 11, the degree of angiogenesis was significantly enhanced in the experimental group administered with pCP-HGF-X7 as compared with the pCP-administered control group.

This result demonstrates that HGF-X7 gene can be effectively used in the gene therapy of an ischemic disease.

While the invention has been described with respect to the above specific embodiments, it should be recognized that various modifications and changes may be made to the invention by those skilled in the art which also fall within the scope of the invention as defined by the appended claims.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 21

<210> SEQ ID NO 1

<211> LENGTH: 2187

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 1

```

atgtgggtga ccaaactcct gccagccctg ctgtcgcagc atgtcctcct gcatctcctc      60
ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat      120
gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa      180
acaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt      240
ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc      300
ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa      360
aacaagact acattagaaa ctgcatcatt ggtaaaggac gcagctaaa gggaacagta      420
tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac      480
agctttttgc cttcgagcta tcggggtaaa gacctacagg aaaactactg tcgaaatcct      540
cgaggggaag aagggggacc ctgggtgttc acaagcaatc cagaggtacg ctacgaagtc      600
tgtgacattc ctcagtgttc agaagttgaa tgcatgacct gcaatgggga gagttatcga      660
ggtctcatgg atcatacaga atcaggcaag atttgtcagc gctgggatca tcagacacca      720
caccggcaca aattcttgcc tgaaagatat cccgacaagg gctttgatga taattattgc      780
cgcaatcccg atggccagcc gagggcatgg tgctatactc ttgaccctca caccgctgg      840
gagtactgtg caattaaaac atgcgctgac aatactatga atgacactga tgttcctttg      900
gaaacaactg aatgcatcca aggtcaagga gaaggctaca ggggactgt caataccatt      960
tggaatggaa ttccatgtca gcggtgggat tctcagtatc ctcacgagca tgacatgact     1020
cctgaaaatt tcaagtgcaa ggacctacga gaaaattact gccgaaatcc agatgggtct     1080
gaatcacccct ggtgttttac cactgatcca aacatccgag ttggctactg ctcccaaatt     1140
ccaaactgtg atatgtcaca tggacaagat tgttatcgtg ggaatggcaa aaattatag     1200
ggcaacttat cccaacaag atctggacta acatgttcaa tgtgggaaa gaacatggaa     1260
gacttacatc gtcatatctt ctgggaacca gatgcaagta agctgaatga gaattactgc     1320
cgaaatccag atgatgatgc tcatggacct tgggtctaca cgggaaatcc actcattcct     1380

```

-continued

tgggattatt gccctatttc tcgttgtgaa ggtgatacca cacctacaat agtcaattta	1440
gaccatcccc taatatcttg tgccaaaacg aaacaattgc gagttgtaaa tgggattcca	1500
acacgaacaa acataggatg gatggttagt ttgagataca gaaataaaca tatctgcgga	1560
ggatcattga taaaggagag ttgggttctt actgcacgac agtgtttccc ttctcgagac	1620
ttgaaagatt atgaagcttg gcttgggaatt catgatgtcc acggaagagg agatgagaaa	1680
tgcaaacagg ttctcaatgt ttcccagctg gtatatggcc ctgaaggatc agatctgggt	1740
ttaatgaagc ttgccaggcc tgctgtcctg gatgattttg ttagtacgat tgatttacct	1800
aattatggat gcacaattcc tgaaaagacc agttgcagtg tttatggctg gggctacact	1860
ggattgatca actatgatgg cctattacga gtggcacatc tctatataat gggaaatgag	1920
aatgcagcc agcatcatcg agggaagggt actctgaatg agtctgaaat atgtgctggg	1980
gctgaaaaga ttggatcagg accatgtgag ggggattatg gtggcccact tgtttgtgag	2040
caacataaaa tgagaatggt tcttgggtgc attgttctctg gtcgtggatg tgccattcca	2100
aatcgtcctg gtatttttgt ccgagtagca tattatgcaa aatggatata caaaattatt	2160
ttaacatata aggtaccaca gtcatag	2187

<210> SEQ ID NO 2

<211> LENGTH: 7113

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc	60
ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat	120
gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa	180
acaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt	240
ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc	300
ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa	360
aacaaagact acattagaaa ctgcatcatt ggtaaaggac gcagctacaa gggaacagta	420
tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac	480
aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacct gttaacagtc	540
tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat	600
tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata	660
tgtaataaaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg	720
tatttgtgaa ataataacgt aaacttcggt taaaaggatt cttcttctctg tctttgagaa	780
agtacggcac tgtgcagggg gagagggtga ttgtgaaaaa tcagaggtag atgagaatct	840
tactgagggc tgagggttct ttaaccttgg tggatctcaa cattgggttc acattaaaat	900
cacctgctgc aagcccttga cgaatcttac ttagaagatg acaacacaga acaattaaat	960
cagaatctct ggggagaata gggcaccagt attttttgag ctcccacat gattccaaag	1020
tgagccaaa tttgagaacc actgctaaaa gctcaagctt cagattgacc agcttttcca	1080
tctcacctat cgctaaaga ccaaattgga taaatgtggt cattacgaca gatgggtact	1140
atttaaagat gagtaaacac aatatactta ggctcgtcag actgagagtt ttaatcatca	1200
ctgaggaaaa acatagatat ctaatactga ctggagtatt agtcaaggct tatttcacac	1260
acaattttat cagaaaccaa agtagtttaa aacagctctc cccttattag taatgcattg	1320

-continued

gagggtttac	tttaccatgt	accttgctga	gcaactgtacc	ttgttaatct	catttacttg	1380
taatgagaac	cacacagcgg	gtagttttat	tggttctatt	ttacctacat	gacaaaactg	1440
aagcataaaa	acacttagta	agttttcagt	gtcatgcaca	actaggaagt	gacatggcca	1500
gaatataagc	ccagtcacca	tcaactctata	acctgcgctt	ttaacaactt	cagggcatga	1560
cacatttggc	cggtcagtag	aacccatgct	gtgatttggt	tttgcagtg	tggtgatgac	1620
tgcttgttg	aatccacttt	ttattctatt	ccattttggg	gacacaattc	tgcaagatga	1680
ttcttcatta	ggaaacagag	atgagttatt	gaccaacaca	gaaagaaaa	gagtttgttg	1740
ctccacactg	ggattaaacc	tatgatcttg	gcctaattaa	cactagctag	taagtgtcca	1800
agctgatcat	ctctacaaca	tttcaataac	agaaaacaac	aattttcaa	attagttact	1860
tacaattatg	tagaaatgcc	tctaaaacac	agtattttcc	ttatattaca	aaaacaaaa	1920
ttataattgg	ttttgtctc	ttttgagagt	ttgcatggtg	ttactcctg	catagtgaag	1980
aaaacatttt	atttaagtag	atggatctaa	gtttttcatg	aacaaaggaa	tgacatttga	2040
aatcaatcct	accctagtcc	aggagaatgc	attagattaa	cctagtagag	gtcttatttc	2100
accctgagtt	ttctatgatc	gtgattctct	gctggaggag	taattgtgaa	atagatctct	2160
ctgggaactg	gcttcctagt	ccaatcagct	cttttacc	tgaacacttc	cttgtgatat	2220
agatgtttat	ggccgagagg	atccagtata	ttaataaaat	ccctttttgt	attcaatgag	2280
ggaaacacat	aattttcatc	aattagcagc	ttattggaat	atctgcatga	tggtttaaca	2340
cttttaagtg	ttgactaaag	attaatttta	cagaaaatag	aaaaagaaat	atgtttctgt	2400
ctggaggaat	gatttattgt	tgaccctaa	attgaaatat	tttactagt	gcttaatgga	2460
aagatgatga	aagatgatga	aattaatgta	gaagcttaac	tagaaaatca	ggtgacctga	2520
tatctacatc	tgtatccttc	attggccacc	cagcattcat	taatgaatca	gatgatggaa	2580
tagatcaagt	ttcctaggaa	cacagtgaat	attaaaagaa	aacaaaggga	gcctagcacc	2640
tagaagacct	agtttatatt	tcaaagtata	tttggatgta	acceatttt	aaacatttcc	2700
tcacttgtct	ctcttaaagc	cttgccaaca	gcaaggacag	agaacaaaa	atagtgtata	2760
tatgaataaa	tgcttattac	agaatctgct	gactggcaca	tgctttgtgt	gtaatgggtt	2820
ctcataaaca	cttgttgaat	gaacacacat	aagtgaaaga	gcatggctag	gcttcatccc	2880
ttggtcaaat	atggggtgct	aaagaaaagc	aggggaaata	cattgggaca	ctaacaaaa	2940
aaaacagtta	atthaggtaa	aagataaaat	acaccacaga	atgaagaaaa	gagatgaccc	3000
agactgctct	ttaaccttca	tgtcctagag	aggtttttga	tatgaattgc	attcagaatt	3060
gtggaaagga	gcccactttt	tctcttcatt	ttgattttat	taactccaat	gggggaattt	3120
tattcgtggt	ttggccatat	ctacttttga	tttctacatt	attctctctt	cctttctacc	3180
tgtatttgtc	ctaataaatt	gttgacttat	taattcacta	cttctcaca	gctttttttt	3240
ggctttacaa	atccactgga	aaggtatatg	ggtgtatcac	tttgtgtatt	tcggtgtgca	3300
tgtgtagagg	ggacaaaaat	cctctctcaa	actataaata	ttgagtattt	gtgtattgaa	3360
catttgctat	aactactagg	tttcttaaat	aatcttaata	tataaaatga	tatagaaaa	3420
gggaaattat	agttcgtatt	attcatctaa	gtgaagagat	taaaaccag	ggagtaaata	3480
aattgtctaa	ggactaaggt	tgtatactat	ttaggtgata	gatatggggc	aaccgtatgg	3540
gttttatgat	taacaaataa	acttctcacc	actctacat	atcaactttt	ccataaaaga	3600
gagctatagt	attctttgct	taaataaatt	tgattagtgc	atgacttctt	gaaaacatat	3660

-continued

aaagcaaaag	tcacatttga	ttctatcaga	aaagtgagta	agccatggcc	caaacaaaag	3720
atgcattaa	atattctgga	atgatggagc	taaaagtaag	aaaaatgact	ttttaaaaaa	3780
gtttactggt	aggaatttg	aaattatgct	gaatttttagt	tgcatataa	tttttgtcag	3840
tcatacggtc	tgacaacctg	tcttatttct	atttcccat	atgaggaatg	ctagttaagt	3900
atggatatta	actattacta	cttagatgca	ttgaagttgc	ataatatgga	taatacttca	3960
ctggttccct	gaaaatgttt	agttagtaat	aagtctctta	cactatttgt	tttgtccaat	4020
aatttatatt	ttctgaagac	ttaactctag	aatacactca	tgtcaaatg	aaagaatttc	4080
attgcaaaat	attgcttgg	acatgacgca	tacctgtatt	tgttttgtgt	cacaacatga	4140
aaaatgatgg	tttattagaa	gtttcattgg	gtaggaaaca	catttgaatg	gtatttacta	4200
agatactaaa	atccttggac	ttcactctaa	ttttagtgcc	atttagaact	caaggtctca	4260
gtaaaagtag	aaataaagcc	tgtaacaaa	acacaagctg	aatattaata	atgtaactgg	4320
attttcaaag	aatgtttac	tggtattacc	tgtagatgta	tattctttat	tatgatcttt	4380
tgtgtaaagt	ctggcagaca	aatgcaatat	ctaattgttg	agtccaatat	cacaagcagt	4440
acaaaagtat	aaaaaagact	tggccttttc	taatgtgta	aaatacttta	tgctggtaat	4500
aacactaaga	gtagggcact	agaaatttta	agtgaagata	atgtgttgca	gttactgcac	4560
tcaatggctt	actattataa	acaaaaactg	ggatcactaa	gctccagtca	gtcaaatga	4620
tcaaaattat	tgaagagaat	aagcaattct	gttctttatt	aggacacagt	agatacagac	4680
tacaaagtgg	agtgtgctta	ataagaggta	gcatttgta	agtgtcaatt	actctattat	4740
cccttggagc	ttctcaaat	aacatataa	ggtgtaagat	gttaaagggt	atggttacac	4800
tcagtgcaca	ggtaagctaa	taggctgaga	gaagctaaat	tacttactgg	ggtctcacag	4860
taagaaagtg	agctgaagtt	tcagcccaga	ttaactgga	ttctgggctc	tttattcatg	4920
ttacttcatg	aatctgtttc	tcaattgtgc	agaaaaagg	gggctattta	taagaaaagc	4980
aataaaciaa	caagtaatga	tctcaataa	gtaatgcaag	aaatagtgag	atctcaaat	5040
cagtggcagc	gatttctcag	ttctgtccta	agtggccttg	ctcaatcacc	tgctatcttt	5100
tagtggagct	ttgaaattat	gtttcagaca	acttcgattc	agttctagaa	tgtttgactc	5160
agcaaatca	caggctcatc	tttctaactt	gatgggtgaat	atggaaattc	agctaaatgg	5220
atgtaataa	aattcaaacy	ttttaaggac	agatgaaaat	gacagaattt	taaggtaaaa	5280
tatatgaagg	aatataagat	aaaggatttt	tctaccttca	gcaaaaacat	accactaat	5340
tagtaaaatt	aataggcaaa	aaaaagttgc	atgctcttat	actgtaatga	ttatcatttt	5400
aaaactagct	ttttgccttc	gagctatcgg	ggtaaagacc	tacaggaaaa	ctactgtcga	5460
aatcctcgag	gggaagaagg	gggaccctgg	tgtttcacaa	gcaatccaga	ggtacgctac	5520
gaagtctgtg	acattcctca	gtgttcagaa	gtgaaatgca	tgacctgcaa	tgaggagagt	5580
tatcgaggtc	tcatggatca	tacagaatca	ggcaagattt	gtcagcgctg	ggatcatcag	5640
acaccacacc	ggcaciaaatt	cttgctgaa	agatatcccg	acaagggtt	tgatgataat	5700
tattgcccga	atcccgatgg	ccagccgagg	ccatgggtgt	atactcttga	ccctcacacc	5760
cgctgggagt	actgtgcaat	taaaacatgc	gctgacaata	ctatgaatga	cactgatggt	5820
cctttggaaa	caactgaatg	catccaaggt	caaggagaag	gctacagggg	cactgtcaat	5880
accatttggg	atggaattcc	atgtcagcgt	tgggattctc	agtatcctca	cgagcatgac	5940
atgactcctg	aaaatttcaa	gtgcaaggac	ctacgagaaa	attactgccg	aaatccagat	6000
gggtctgaat	caccctggtg	ttttaccact	gatccaaaca	tccgagttgg	ctactgctcc	6060

-continued

```

caaattccaa actgtgatat gtcacatgga caagattggt atcgtgggaa tggcaaaaat 6120
tatatgggca acttatcca aacaagatct ggactaacat gttcaatgtg ggacaagaac 6180
atggaagact tacatcgtca tatcttctgg gaaccagatg caagtaagct gaatgagaat 6240
tactgccgaa atccagatga tgatgctcat ggaccctggt gctacacggg aaatccactc 6300
attccttggg attattgcc tatttctcgt tgtgaagggt ataccacacc tacaatagtc 6360
aatttagacc atcccgtaat atcttgtgcc aaaacgaaac aattgctgagtg tgtaaattggg 6420
attccaacac gaacaaacat aggatggatg gttagtttga gatacagaaa taaacatatac 6480
tgcggaggat cattgataaa ggagagttgg gttcttactg cacgacagtg tttcccttct 6540
cgagacttga aagattatga agcttggctt ggaattcatg atgtccacgg aagaggagat 6600
gagaaatgca aacaggttct caatgtttcc cagctgggat atggccctga aggatcagat 6660
ctggttttaa tgaagcttgc caggcctgct gtctggatg attttgtag tacgattgat 6720
ttacctaat atggatgac aattcctgaa aagaccagtt gcagtgttta tggctggggc 6780
tacctggat tgatcaacta tgatggccta ttacgagtgg cacatctcta tataatggga 6840
aatgagaaat gcagccagca tcatcgaggg aagtgactc tgaatgagtc tgaaatatgt 6900
gctggggctg aaaagattgg atcaggacca tgtgaggggg attatggtgg cccacttggt 6960
tgtgagcaac ataaaatgag aatggttctt ggtgtcattg ttctggtcg tggatgtgcc 7020
attccaaatc gtctgggat ttttgtccga gtagcatatt atgcaaaatg gatacacaaa 7080
attattttaa catataaggt accacagtca tag 7113

```

```

<210> SEQ ID NO 3
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: gHGF3 primer

```

```

<400> SEQUENCE: 3

```

```

gtaaaggacg cgtctacaag ggaacagtat ctat 34

```

```

<210> SEQ ID NO 4
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: gHGF4 primer

```

```

<400> SEQUENCE: 4

```

```

actggatcct ctggccata aacatct 27

```

```

<210> SEQ ID NO 5
<211> LENGTH: 34
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:
<223> OTHER INFORMATION: gHGF10 primer

```

```

<400> SEQUENCE: 5

```

```

gaagcttagc accatgtggg tgaccaaact cctg 34

```

```

<210> SEQ ID NO 6
<211> LENGTH: 27
<212> TYPE: DNA
<213> ORGANISM: Artificial Sequence
<220> FEATURE:

```

-continued

<223> OTHER INFORMATION: gHGF5 primer

<400> SEQUENCE: 6

tggccgagag gatccagtat attaata

27

<210> SEQ ID NO 7

<211> LENGTH: 27

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: gHGF7 primer

<400> SEQUENCE: 7

cccctcgagg atttcgacag tagtttt

27

<210> SEQ ID NO 8

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: gHGF12 primer

<400> SEQUENCE: 8

gggatccctt cctttctacc tgtatttg

28

<210> SEQ ID NO 9

<211> LENGTH: 25

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: gHGF13 primer

<400> SEQUENCE: 9

gggatcctgg gtaaacacat ttgaa

25

<210> SEQ ID NO 10

<211> LENGTH: 28

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: gHGF6 primer

<400> SEQUENCE: 10

gggatcctta tgtttcagac aacttcga

28

<210> SEQ ID NO 11

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: gHGF1 primer

<400> SEQUENCE: 11

gaagcttgcc accatgtggg tgaccaaact cctg

34

<210> SEQ ID NO 12

<211> LENGTH: 34

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: gHGF2 primer

<400> SEQUENCE: 12

gggatccaga acgcgtcctt taccgatgat gcag

34

-continued

<210> SEQ ID NO 13
 <211> LENGTH: 35
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: gHGF8 primer

 <400> SEQUENCE: 13

 gggatccctt ctcgagactt gaaagattat gaagc 35

<210> SEQ ID NO 14
 <211> LENGTH: 32
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: gHGF9 primer

 <400> SEQUENCE: 14

 gtctagagcg gccgctatga ctgtggtacc tt 32

<210> SEQ ID NO 15
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: cHGF5 primer

 <400> SEQUENCE: 15

 ggatccacgc gtagcagcac catgtgggtg accaaa 36

<210> SEQ ID NO 16
 <211> LENGTH: 36
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: cHGF3 primer

 <400> SEQUENCE: 16

 ggatcctcta gattacttca gctatgactg tggtag 36

<210> SEQ ID NO 17
 <211> LENGTH: 26
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: GHGF5' primer

 <400> SEQUENCE: 17

 caaatgtcag ccctggagtt ccatga 26

<210> SEQ ID NO 18
 <211> LENGTH: 23
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: GHGF3' primer

 <400> SEQUENCE: 18

 ctggattgct tgtgaaacag ggt 23

<210> SEQ ID NO 19
 <211> LENGTH: 4679
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: HGF-X6 gene

-continued

<400> SEQUENCE: 19

```

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcactcctc 60
ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120
gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa 180
acaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240
ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc 300
ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa 360
aacaagact acattagaaa ctgcatcatc ggtaaaggac gcagctaca gggaacagta 420
tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac 480
aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacct gttaacagtc 540
tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat 600
tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata 660
tgttaataaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg 720
tatttgtgga tcccttcctt tctacctgta tttgtcctaa taaattgttg acttattaat 780
tcactacttc ctcacagctt ttttttggct ttacaaatcc actggaaagg tatatgggtg 840
tactactttg tgtatttcgg tgtgcatgtg tagaggggac aaaaatcctc tctcaacta 900
taaattattg gtatttgtgt attgaacatt tgcataact actaggtttc ttaaataatc 960
ttaatatata aatgatata gaaaaagga aattatagtt cgtattattc atctaagtga 1020
agagattaaa acccaggagag taaataaatt gtctaaggac taaggttgta tactatttag 1080
gtgatagata tggggcaacc gtatgggttt tatgattaac aaataaactt ctcaccactc 1140
taccatatca acttttccat aaaagagagc tatagtattc tttgcttaa taaatttgat 1200
tagtgcatac cttcttgaaa acatataaag caaaagtcac atttgattct atcagaaaag 1260
tgagtaagcc atggcccaaa caaaagatgc attaaaatat tctggaatga tggagctaaa 1320
agtaagaaaa atgacttttt aaaaaagttt actgttagga attgtgaaat tatgctgaat 1380
tttagttgca ttataatttt tgtcagtcac acggtctgac aacctgtctt atttctattt 1440
ccccatatga ggaatgctag ttaagtatgg atattaacta ttactactta gatgcattga 1500
agttgcataa tatggataat acttcactgg ttccctgaaa atgttttagt agtaataagt 1560
ctcttacact atttgttttg tccaataatt tatattttct gaagacttaa ctctagaata 1620
cactcatgtc aaaatgaaag aatttcattg caaaatattg cttggtacat gacgcatacc 1680
tgtatttggt ttgtgtcaca acatgaaaaa tgatggttta ttagaagttt cattgggtag 1740
gaaacacatt tgaatggtat ttactaagat actaaaatcc ttggacttca ctctaatttt 1800
agtgccattt agaactcaag gtctcagtaa aagtagaaat aaagcctgtt aacaaaacac 1860
aagctgaata ttaaaaatgt aactggattt tcaaagaaat gtttactggt attacctgta 1920
gatgtatatt ctttattatg atcttttgtg taaagtctgg cagacaaatg caatatctaa 1980
ttgttgagtc caatatcaca agcagtaca aagtataaaa aagacttggc cttttctaat 2040
gtgttaaaat actttatgct ggtaataaca ctaagagtag ggcactagaa attttaagtg 2100
aagataatgt gttgcagtta ctgcactcaa tggttacta ttataacca aaactgggat 2160
cactaagctc cagtcagtca aatgatcaa aattattgaa gagaataagc aattctgttc 2220
tttattagga cacagtagat acagactaca aagtggagtg tgcttaataa gaggtagcat 2280
ttgttaagtg tcaattactc tattatccct tggagcttct caaataacc atataagtg 2340

```

-continued

taagatgta	aaggttatgg	ttacactcag	tgcacaggta	agctaatagg	ctgagagaag	2400
ctaaattact	tactggggtc	tcacagtaag	aaagtgagct	gaagtttcag	cccagattta	2460
actggattct	gggctcttta	ttcatgttac	ttcatgaatc	tgtttctcaa	ttgtgcagaa	2520
aaaagggggc	tatttataag	aaaagcaata	aacaaacaag	taatgatctc	aaataagtaa	2580
tgcaagaaat	agtgagattt	caaaatcagt	ggcagcgatt	tctcagttct	gtcctaagtg	2640
gccttgctca	atcacctgct	atcttttagt	ggagctttga	aattatgttt	cagacaactt	2700
cgattcagtt	ctagaatggt	tgactcagca	aattcacagg	ctcatctttc	taacttgatg	2760
gtgaatatgg	aaattcagct	aatggatgt	taataaaatt	caaacgtttt	aaggacagat	2820
gaaaatgaca	gaattttaag	gtaaaatata	tgaaggaata	taagataaag	gatttttcta	2880
ccttcagcaa	aacataccc	actaattagt	aaaattaata	ggcaaaaaaa	agttgcatgc	2940
tcttatactg	taatgattat	cattttaaaa	ctagcttttt	gccttcgagc	tatcggggta	3000
aagacctaca	ggaaaactac	tgtcgaaatc	ctcaggggga	agaaggggga	ccctgggtgtt	3060
tcacaagcaa	tccagaggta	cgctacgaag	tctgtgacat	tcctcagtgt	tcagaagttg	3120
aatgcatgac	ctgcaatggg	gagagttatc	gaggtctcat	ggatcataca	gaatcaggca	3180
agatttgca	gcgctgggat	catcagacac	cacaccggca	caaattcttg	cctgaaagat	3240
atcccgacaa	gggctttgat	gataattatt	gccgcaatcc	cgatggccag	ccgaggccat	3300
ggtgctatac	tcttgaccct	cacacccgct	gggagtactg	tgcaattaaa	acatgctgctg	3360
acaatactat	gaatgacact	gatgttcctt	tggaaacaac	tgaatgcatc	caagggtcaag	3420
gagaaggcta	caggggcact	gtcaatacca	tttggaatgg	aattccatgt	cagcgttggg	3480
attctcagta	tcctcacgag	catgacatga	ctcctgaaaa	tttcaagtgc	aaggacctac	3540
gagaaaatta	ctgccgaaat	ccagatgggt	ctgaatcacc	ctgggtgttt	accactgatc	3600
caaacatccg	agttggctac	tgctcccaaa	ttccaaactg	tgatatgtca	catggacaag	3660
attgttatcg	tgggaatggc	aaaaattata	tgggcaactt	atcccaaca	agatctggac	3720
taacatgttc	aatgtgggac	aagaacatgg	aagacttaca	tcgtcatatc	ttctgggaac	3780
cagatgcaag	taagctgaat	gagaattact	gccgaaatcc	agatgatgat	gctcatggac	3840
cctggtgcta	cacgggaaat	ccactcattc	cttgggatta	ttgccctatt	tctcgttgtg	3900
aagggtgatac	cacacctaca	atagtcaatt	tagaccatcc	cgtaatatct	tgtgccaaaa	3960
cgaaacaatt	gcgagttgta	aatgggatcc	caacacgaac	aaacatagga	tggatggtta	4020
gtttgagata	cagaaataaa	catatctgcg	gaggatcatt	gataaaggag	agttgggttc	4080
ttactgcacg	acagtgtttc	ccttctcgag	acttgaaaga	ttatgaagct	tggcttggaa	4140
ttcatgatgt	ccacggaaga	ggagatgaga	aatgcaaca	ggttctcaat	gtttcccagc	4200
tggatatatgg	ccctgaagga	tcagatctgg	ttttaatgaa	gcttgccagg	cctgctgtcc	4260
tggatgattt	tgtagtacg	attgatttac	ctaattatgg	atgcacaatt	cctgaaaaga	4320
ccagttgcag	tgtttatggc	tggggctaca	ctggattgat	caactatgat	ggcctattac	4380
gagtggcaca	tctctatata	atgggaaatg	agaaatgcag	ccagcatcat	cgaggggaagg	4440
tgactctgaa	tgagtctgaa	atatgtgctg	gggtgaaaa	gattggatca	ggacctgtg	4500
agggggatta	tgggtggcca	cttgtttgtg	agcaacataa	aatgagaatg	gttcttgggtg	4560
tcattgttcc	tggctgtgga	tgtgccattc	caaatcgtec	tggatatttt	gtccgagtag	4620
catattatgc	aaaatggata	cacaaaatta	ttttaacata	taaggtacca	cagtcatag	4679

-continued

<210> SEQ ID NO 20
 <211> LENGTH: 3679
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: HGF-X7 gene

<400> SEQUENCE: 20

```

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcatctcctc   60
ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat   120
gaattcaaaa aatcagcaaa gactacccta atcaaaatag atccagcact gaagataaaa   180
acaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt   240
ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc   300
ttcaatagca tgtcaagtgg agtgaaaaaa gaatttggcc atgaatttga cctctatgaa   360
aacaagact acattagaaa ctgcatcatc ggtaaaggac gcagctacaa gggaacagta   420
tctatcacta agagtggcat caaatgtcag ccctggagtt ccatgatacc acacgaacac   480
aggtaagaac agtatgaaga aaagagatga agcctctgtc ttttttacct gttaacagtc   540
tcatattagt ccttcagaat aattctacaa tcctaaaata acttagccaa cttgctgaat   600
tgtattacgg caaggtttat atgaattcat gactgatatt tagcaaatga ttaattaata   660
tgttaataaa atgtagccaa aacaatatct taccttaatg cctcaatttg tagatctcgg   720
tatttgtgga tcctgggtag gaaacacatt tgaatggtat ttactaagat actaaaatcc   780
ttggacttca ctctaatttt agtgccattt agaactcaag gtctcagtaa aagtagaaat   840
aaagcctggt aacaaaacac aagctgaata ttaaaaatgt aactggattt tcaaagaaat   900
gtttactggg attacctgta gatgtatatt ctttattatg atcttttgtg taaagtctgg   960
cagacaaatg caatatctaa ttgttgagtc caatatcaca agcagtacaa aagtataaaa  1020
aagacttggc cttttcta atgtgtaaaat actttatgct ggtaataaca ctaagagtag  1080
ggcactagaa attttaagtg aagataatgt gttgcagtta ctgcactcaa tggcttacta  1140
ttataacca aaactgggat cactaagctc cagtcagtca aatgatcaa aattattgaa  1200
gagaataagc aattctgttc tttattagga cacagtagat acagactaca aagtggagtg  1260
tgcttaataa gaggtagcat ttgttaagtg tcaattactc tattatccct tggagcttct  1320
caaaaataacc atataaggtg taagatgtaa aaggttatgg ttacactcag tgcacaggta  1380
agctaatagg ctgagagaag ctaaattact tactggggtc tcacagtaag aaagtgagct  1440
gaagtttcag cccagattta actggattct gggctcttta ttcattgtac ttcattgaatc  1500
tgtttctcaa ttgtgcagaa aaaagggggc tatttataag aaaagcaata acaaaacaag  1560
taatgatctc aaataagtaa tgcaagaaat agtgagattt caaaatcagt ggcagcgatt  1620
tctcagttct gtcctaagtg gccttgctca atcacctgct atcttttagt ggagctttga  1680
aattatgttt cagacaactt cgattcagtt ctagaatggt tgactcagca aattcacagg  1740
ctcatctttc taacttgatg gtgaatatgg aaattcagct aaatggatgt taataaaatt  1800
caaacgtttt aaggacagat gaaaatgaca gaattttaag gtaaaatata tgaaggaata  1860
taagataaag gatttttcta ccttcagcaa aaacataccc actaattagt aaaattaata  1920
ggcaaaaaaa agttgcatgc tcttatactg taatgattat cattttaaaa ctagcttttt  1980
gccttcgagc tateggggta aagacctaca ggaaaactac tgtcgaaatc ctcgagggga  2040
agaaggggga ccctggtggt tcacaagcaa tccagaggta cgctacgaag tctgtgacat  2100

```

-continued

```

tcctcagtgt tcagaagttg aatgcatgac ctgcaatggg gagagttatc gaggtctcat 2160
ggatcataca gaatcaggca agatttgtca gcgctgggat catcagacac cacaccggca 2220
caaattcttg cctgaaagat atccccgaaa gggctttgat gataattatt gccgcaatcc 2280
cgatggccag ccgaggccat ggtgctatac tcttgaccct cacaccgct gggagtactg 2340
tgcaattaa acatgcgctg acaatactat gaatgacact gatgttcctt tggaaacaac 2400
tgaatgcac caaggtcaag gagaaggcta caggggact gtcaatacca tttggaatgg 2460
aattccatgt cagcgttggg attctcagta tctcaccgag catgacatga ctctgaaaa 2520
tttcaagtgc aaggacctac gagaaaatta ctgccgaaat ccagatgggt ctgaatcacc 2580
ctggtgtttt accactgatc caaacatccg agttggctac tgctccaaa ttccaaactg 2640
tgatatgtca catggacaag attgttatcg tgggaatggc aaaaattata tgggcaactt 2700
atcccaaaca agatctggac taacatgttc aatgtgggac aagaacatgg aagacttaca 2760
tcgtcatatc ttctgggaac cagatgcaag taagctgaat gagaattact gccgaaatcc 2820
agatgatgat gctcatggac cctggtgcta cacgggaaat ccactcattc cttgggatta 2880
ttgccctatt tctcgttggg aaggtgatac cacacctaca atagtcaatt tagaccatcc 2940
cgtaatatct tgtgccaaa cgaacaatt gcgagttgta aatgggattc caacacgaac 3000
aaacatagga tggatggtta gtttgagata cagaaataaa catatctgcg gaggatcatt 3060
gataaaggag agttgggttc ttactgcacg acagtgttcc ccttctcgag acttgaaaga 3120
ttatgaagct tggcttgaa tcatgatgt ccacggaaga ggagatgaga aatgcaaca 3180
ggttctcaat gtttcccagc tggatatgg ccctgaagga tcagatctgg ttttaatgaa 3240
gcttgccagg cctgctgtcc tggatgattt tgtagtacg attgatttac ctaattatgg 3300
atgcacaatt cctgaaaaga ccagttgcag tgtttatggc tggggctaca ctggattgat 3360
caactatgat ggcctattac gagtggcaca tctctatata atgggaaatg agaaatgcag 3420
ccagcatcat cgagggaag tgactctgaa tgagtctgaa atatgtgctg gggctgaaaa 3480
gattggatca ggaccatgtg agggggatta tgggtggcca cttgtttgat agcaacataa 3540
aatgagaatg gttcttggg tcattgttcc tggctgtgga tgtgccattc caaatcgtcc 3600
tggtatTTTT gtccgagtag catattatgc aaaatggata cacaaaatta ttttaacata 3660
taaggtacca cagtcatag 3679

```

<210> SEQ ID NO 21

<211> LENGTH: 2729

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: HGF-X8 gene

<400> SEQUENCE: 21

```

atgtgggtga ccaaactcct gccagccctg ctgctgcagc atgtcctcct gcactcctc 60
ctgctcccca tcgccatccc ctatgcagag ggacaaagga aaagaagaaa tacaattcat 120
gaattcaaaa aatcagcaaa gactacccta atcaaatag atccagcact gaagataaaa 180
acaaaaaag tgaatactgc agaccaatgt gctaatagat gtactaggaa taaaggactt 240
ccattcactt gcaaggcttt tgtttttgat aaagcaagaa aacaatgcct ctggttcccc 300
ttcaatagca tgtcaagtgg agtgaaaaaa gaatttgccc atgaatttga cctctatgaa 360
aacaagact acattagaaa ctgcatcatc ggtaaaggac gcagctacaa gggaacagta 420

```

-continued

tctatcacta	agagtggcat	caaatgtcag	ccttgagtt	ccatgatacc	acacgaacac	480
aggtaagaac	agtatgaaga	aaagagatga	agcctctgtc	ttttttacat	gttaacagtc	540
tcatattagt	ccttcagaat	aattctacaa	tctaaaata	acttagccaa	cttgctgaat	600
tgtattacgg	caaggtttat	atgaattcat	gactgatatt	tagcaaatga	ttaattaata	660
tgtaataaaa	atgtagccaa	aacaatatct	taccttaatg	cctcaatttg	tagatctcgg	720
tatttggtga	tccttatgtt	tcagacaact	tcgattcagt	tctagaatgt	ttgactcagc	780
aaattcacag	gctcatcttt	ctaacttgat	ggtgaatatg	gaaattcagc	taaatggatg	840
ttaataaaaat	tcaaactgtt	taaggacaga	tgaaaatgac	agaattttaa	ggtaaaaat	900
atgaaggaat	ataagataaa	ggatttttct	accttcagca	aaaacatacc	cactaattag	960
taaaattaat	aggcaaaaaa	aagttgcatg	ctcttatact	gtaatgatta	tcattttaaa	1020
actagctttt	tgcttcgag	ctatcggggt	aaagacctac	aggaaaacta	ctgtcgaaat	1080
cctcgagggg	aagaagggg	accctgggtg	ttcacaagca	atccagaggt	acgctacgaa	1140
gtctgtgaca	ttctcagtg	ttcagaagtt	gaatgcatga	cctgcaatgg	ggagagttat	1200
cgaggtctca	tggatcatac	agaatcaggc	aagatttgtc	agcgctggga	tcatacagaca	1260
ccacaccggc	acaaattctt	gctgaaaga	tatcccagca	agggctttga	tgataattat	1320
tgccgcaatc	ccgatggcca	gccgaggcca	tggtgctata	ctcttgacc	tcacaccgc	1380
tgggagtact	gtgcaattaa	aacatgcgct	gacaatacta	tgaatgacac	tgatgttctt	1440
ttgaaacaa	ctgaatgcat	ccaaggtcaa	ggagaaggct	acaggggcac	tgtcaatacc	1500
atttggaatg	gaattccatg	tcagcgttgg	gattctcagt	atcctcacga	gcatgacatg	1560
actcctgaaa	atttcaagtg	caaggaccta	cgagaaaatt	actgccgaaa	tccagatggt	1620
ctgaatcacc	ctgggtgttt	accactgatc	caaacatccg	agttggctac	tgctcccaaa	1680
ttccaaactg	tgatatgtca	catggacaag	attgttatcg	tgggaatggc	aaaaattata	1740
tgggcaactt	atcccaaaca	agatctggac	taacatgttc	aatgtgggac	aagaacatgg	1800
aagacttaca	tcgtcatatc	ttctgggaac	cgatgcaag	taagctgaat	gagaattact	1860
gccgaaatcc	agatgatgat	gctcatggac	cctggtgcta	cacgggaaat	ccactcattc	1920
cttgggatta	ttgcctatt	tctcgttggtg	aaggtgatac	cacacctaca	atagtcaatt	1980
tagaccatcc	cgtaatatct	tgtgccaaaa	cgaaacaatt	gcgagttgta	aatgggattc	2040
caacacgaac	aaacatagga	tggatggtta	gtttgagata	cagaaataaa	catatctgcg	2100
gaggatcatt	gataaaggag	agttgggttc	ttactgcacg	acagtgtttc	ccttctcgag	2160
acttgaaga	ttatgaagct	tggcttgga	ttcatgatgt	ccacggaaga	ggagatgaga	2220
aatgcaaca	ggttctcaat	gtttcccagc	tggtatatgg	ccctgaagga	tcagatctgg	2280
ttttaatgaa	gcttgccag	cctgctgtcc	tggatgattt	tgtagtacg	attgatttac	2340
ctaattatgg	atgcacaatt	cctgaaaaga	ccagttgcag	tgtttatggc	tggggctaca	2400
ctggattgat	caactatgat	ggcctattac	gagtggcaca	tctctatata	atgggaaatg	2460
agaaatgcag	ccagcatcat	cgaggggaag	tgactctgaa	tgagtctgaa	atatgtgctg	2520
gggctgaaaa	gattggatca	ggaccatgtg	agggggatta	tgggtggcca	cttgtttggtg	2580
agcaacataa	aatgagaatg	gttcttggtg	tcattgttcc	tggctgtgga	tgtgccattc	2640
caaatcgtcc	tggatatttt	gtccgagtag	catattatgc	aaaatggata	cacaaaatta	2700
ttttaacata	taagtagca	cagtcatag				2729

What is claimed is:

1. A method for co-expressing two heterotypes of Hepatocyte Growth Factor (HGF) comprising transforming or transfecting to a cell a DNA construct comprising:

- (a) a promoter,
 - (b) a first cDNA which has the same sequence as exons 1-4 of the human HGF gene wherein said exons 1-4 are arranged in sequential order without an intron therebetween, or degenerates thereof which do not alter the amino acid sequence encoded by said first cDNA,
 - (c) a polynucleotide that has the same sequence as intron 4 of the HGF gene or a functional fragment thereof, and
 - (d) a second cDNA which has the same sequence as exons 5-18 of the human HGF gene wherein said exons 5-18 are arranged in sequential order without an intron therebetween, or degenerates thereof which do not alter the amino acid sequence encoded by said second cDNA;
- wherein (c) is located between (b) and (d); and the HGF construct simultaneously encodes two heterotypes of human HGF.

2. The method of claim **1**, wherein said intron has the same sequence as a fragment of intron 4 of the HGF gene.

3. The method of claim **2**, wherein the construct comprises a nucleotide sequence not less than 90% identical to SEQ ID NO: 19.

4. The method of claim **3**, wherein the construct comprises a nucleotide sequence not less than 95% identical to SEQ ID NO: 19.

5. The method of claim **4**, wherein the construct comprises the sequence of SEQ ID NO: 19.

6. The method of claim **2**, wherein the construct comprises a nucleotide sequence not less than 90% identical to SEQ ID NO: 20.

7. The method of claim **6**, wherein the construct comprises a nucleotide sequence not less than 95% identical to SEQ ID NO: 20.

8. The method of claim **7**, wherein the construct comprises the sequence of SEQ ID NO: 20.

9. The method of claim **2**, wherein the construct comprises a nucleotide sequence not less than 90% identical to SEQ ID NO: 21.

10. The method of claim **9**, wherein the construct comprises a nucleotide sequence not less than 95% identical to SEQ ID NO: 21.

11. The method of claim **10**, wherein the construct comprises the sequence of SEQ ID NO: 21.

12. The method of claim **1**, wherein the one intron has the same sequence as the full intron 4 of the HGF gene.

13. The method of claim **1**, wherein the construct comprises a nucleotide sequence not less than 90% identical to SEQ ID NO: 2.

14. The method of claim **13**, wherein the construct comprises a nucleotide sequence not less than 95% identical to SEQ ID NO: 2.

15. The method of claim **14**, wherein the construct comprises the sequence of SEQ ID NO: 2.

16. The method of claim **1**, wherein the construct further comprises a terminator sequence, a self-replication sequence, or a secretory signal.

17. The method of claim **1**, wherein the expression efficiency of the construct is higher than the expression efficiency of HGF cDNA or deleted variant HGF (dHGF) cDNA.

18. The method of claim **1**, wherein the expression level of the construct is about 20- to 100-fold higher than the expression level of the HGF cDNA or dHGF cDNA.

19. The method of claim **1**, wherein the cell is a mammalian cell, a bacterial cell or a yeast cell.

20. The method of claim **19**, wherein the cell is a mammalian cell.

21. The method of claim **20**, wherein the transformation of said mammalian cell is in vivo.

* * * * *