

US007790381B2

(12) **United States Patent**
Arnold et al.

(10) **Patent No.:** **US 7,790,381 B2**
(45) **Date of Patent:** ***Sep. 7, 2010**

(54) **METHOD FOR CREATING
POLYNUCLEOTIDE AND POLYPEPTIDE
SEQUENCES**

(75) Inventors: **Frances Arnold**, Pasadena, CA (US);
Zhixin Shao, Penzberg (DE); **Alexander
Volkov**, South Pasadena, CA (US)

(73) Assignee: **California Institute of Technology**,
Pasadena, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 595 days.

This patent is subject to a terminal dis-
claimer.

(21) Appl. No.: **11/636,421**

(22) Filed: **Dec. 7, 2006**

(65) **Prior Publication Data**

US 2008/0268505 A1 Oct. 30, 2008

Related U.S. Application Data

(63) Continuation of application No. 10/371,168, filed on
Feb. 19, 2003, now Pat. No. 7,166,466, which is a
continuation of application No. 09/205,448, filed on
Dec. 4, 1998, now Pat. No. 6,537,746.

(60) Provisional application No. 60/067,908, filed on Dec.
8, 1997.

(51) **Int. Cl.**
C12Q 1/68 (2006.01)
C12N 15/00 (2006.01)
C07H 21/00 (2006.01)
C07H 21/02 (2006.01)

(52) **U.S. Cl.** **435/6; 435/440; 536/22.1;**
536/23.1

(58) **Field of Classification Search** **435/6,**
435/440; 536/22.1, 23.1, 24.3
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,683,202 A 7/1987 Mullis
4,800,159 A 1/1989 Mullis et al.
4,816,567 A 3/1989 Cabilly et al.
4,959,312 A 9/1990 Sirotkin
4,965,188 A 10/1990 Mullis et al.
4,994,368 A 2/1991 Goodman et al.
4,994,379 A 2/1991 Chang
5,023,171 A 6/1991 Ho et al.
5,043,272 A 8/1991 Hartley
5,066,584 A 11/1991 Gyllensten et al.
5,093,257 A 3/1992 Gray
5,106,727 A 4/1992 Hartley et al.
5,169,764 A 12/1992 Shooter et al.
5,176,995 A 1/1993 Sninsky et al.
5,187,083 A 2/1993 Mullis
5,223,408 A 6/1993 Goeddel et al.
5,234,824 A 8/1993 Mullis

5,264,563 A 11/1993 Huse
5,279,952 A 1/1994 Wu
5,314,809 A 5/1994 Erlich et al.
5,316,935 A 5/1994 Arnold et al.
5,324,830 A * 6/1994 Resnick et al. 536/23.2
5,356,801 A 10/1994 Rambosek et al.
5,360,728 A 11/1994 Prasher
5,418,149 A 5/1995 Gelfand et al.
5,422,266 A 6/1995 Cormier et al.
5,470,725 A 11/1995 Borriss et al.
5,489,523 A 2/1996 Mathur
5,502,167 A 3/1996 Waldmann et al.
5,512,463 A 4/1996 Stemmer
5,514,568 A 5/1996 Stemmer
5,521,077 A 5/1996 Khosla et al.
5,523,388 A 6/1996 Huse
5,541,309 A 7/1996 Prasher
5,556,750 A 9/1996 Modrich
5,556,772 A 9/1996 Sorge et al.
5,571,708 A 11/1996 Yang et al.
5,574,205 A 11/1996 Kucherlapati et al.
5,605,793 A 2/1997 Stemmer
5,629,179 A 5/1997 Mierendorf et al.
5,652,116 A 7/1997 Grandi et al.
5,679,522 A 10/1997 Modrich
5,698,426 A 12/1997 Huse
5,714,316 A 2/1998 Weiner et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 746786 5/2002

(Continued)

OTHER PUBLICATIONS

Barnes W.M. PCR amplification of up to 35-kb DNA with high
fidelity and high yield from lambda bacteriophage template. PNAS
91 : 2216-2220 (1991).*

(Continued)

Primary Examiner—Ethan Whisenant
(74) *Attorney, Agent, or Firm*—Townsend and Townsend and
Crew LLP

(57) **ABSTRACT**

The invention provides methods for evolving a polynucle-
otide toward acquisition of a desired property. Such methods
entail incubating a population of parental polynucleotide
variants under conditions to generate annealed polynucle-
otides comprising heteroduplexes. The heteroduplexes are
then exposed to a cellular DNA repair system to convert the
heteroduplexes to parental polynucleotide variants or recom-
bined polynucleotide variants. The resulting polynucleotides
are then screened or selected for the desired property.

34 Claims, 24 Drawing Sheets

U.S. PATENT DOCUMENTS

5,723,323	A	3/1998	Kauffman et al.
5,756,316	A	5/1998	Schellenberger
5,763,192	A	6/1998	Kauffman et al.
5,770,434	A	6/1998	Huse
5,773,267	A	6/1998	Jacobs et al.
5,783,431	A	7/1998	Peterson et al.
5,795,747	A	8/1998	Henco et al.
5,808,022	A	9/1998	Huse
5,811,238	A	9/1998	Stemmer et al.
5,814,476	A	9/1998	Kauffman et al.
5,817,483	A	10/1998	Kauffman et al.
5,824,469	A	10/1998	Horwitz et al.
5,824,485	A	10/1998	Thompson et al.
5,824,514	A	10/1998	Kauffman et al.
5,830,696	A	11/1998	Short
5,830,721	A	11/1998	Stemmer et al.
5,834,252	A	11/1998	Stemmer et al.
5,837,458	A	11/1998	Minshull et al.
5,843,643	A	12/1998	Ratner
5,851,813	A	12/1998	Desrosiers
5,858,725	A	1/1999	Crowe et al.
5,866,363	A	2/1999	Pieczzenik
5,871,974	A	2/1999	Huse
5,877,402	A	3/1999	Maliga et al.
5,925,749	A	7/1999	Mathur et al.
5,928,905	A	7/1999	Stemmer et al.
5,939,250	A	8/1999	Short
5,955,358	A	9/1999	Huse
5,958,672	A	9/1999	Short
5,962,258	A	10/1999	Mathur et al.
5,965,408	A	10/1999	Short
5,965,415	A	10/1999	Radman
5,976,862	A	11/1999	Kauffman et al.
6,001,574	A	12/1999	Short et al.
6,004,788	A	12/1999	Short
6,030,779	A	2/2000	Short
6,051,409	A	4/2000	Hansen et al.
6,054,267	A	4/2000	Short
6,057,103	A	5/2000	Short
6,071,889	A	6/2000	Weiss et al.
6,074,853	A	6/2000	Pati et al.
6,087,177	A	7/2000	Wohlstadter
6,087,341	A	7/2000	Khavari
6,093,873	A	7/2000	Chambon et al.
6,096,548	A	8/2000	Stemmer
6,103,463	A	8/2000	Chetverin et al.
6,117,679	A	9/2000	Stemmer
6,132,970	A	10/2000	Stemmer
6,165,793	A	12/2000	Stemmer
6,168,919	B1	1/2001	Short
6,171,820	B1	1/2001	Short
6,174,673	B1	1/2001	Short
6,180,406	B1	1/2001	Stemmer
6,277,638	B1	8/2001	Stemmer
6,287,861	B1	9/2001	Stemmer et al.
6,291,242	B1	9/2001	Stemmer
6,297,053	B1	10/2001	Stemmer
6,323,030	B1	11/2001	Stemmer
6,352,842	B1	3/2002	Short et al.
6,358,709	B1	3/2002	Short et al.
6,361,974	B1	3/2002	Short et al.
6,365,408	B1	4/2002	Stemmer
6,379,964	B1	4/2002	del Cardayre et al.
6,583,336	B1 *	6/2003	Reiss et al. 800/278

FOREIGN PATENT DOCUMENTS

EP	252 666	12/1998
EP	544 809	12/1998
EP	563 296	3/1999
EP	552 266	1/2000

EP	876 509	9/2001
EP	911 396	9/2001
EP	1 138 763	10/2001
EP	934 999	1/2002
EP	752 008	4/2002
JP	2-303489	12/1990
WO	WO 90/07576	7/1990
WO	WO 90/14424	11/1990
WO	WO 90/14430	11/1990
WO	WO 91/01087	2/1991
WO	WO 91/06570	5/1991
WO	WO 91/06643	5/1991
WO	WO 91/06645	5/1991
WO	WO 91/07506	5/1991
WO	WO 91/15581	10/1991
WO	WO 91/16427	10/1991
WO	WO 92/06176	4/1992
WO	WO 92/07075	4/1992
WO	WO 92/18645	10/1992
WO	WO 93/01282	1/1993
WO	WO 93/02191	2/1993
WO	WO 93/06213	4/1993
WO	WO 93/11237	6/1993
WO	WO 93/12228	6/1993
WO	WO 93/15208	8/1993
WO	WO 93/16192	8/1993
WO	WO 93/18141	9/1993
WO	WO 93/19172	9/1993
WO	WO 93/25237	12/1993
WO	WO 94/03596	2/1994
WO	WO 94/09817	5/1994
WO	WO 94/11496	5/1994
WO	WO 94/13804	6/1994
WO	WO 95/17413	6/1995
WO	WO 95/22625	8/1995
WO	WO 96/17056	6/1996
WO	WO 96/33207	10/1996
WO	WO 97/07205	2/1997
WO	WO 97/20078	6/1997
WO	WO 97/25410	7/1997
WO	WO 97/35957	10/1997
WO	WO 97/35966	10/1997
WO	WO 98/01581	1/1998
WO	WO 98/27230	6/1998
WO	WO 98/28416	7/1998
WO	WO 98/41622	9/1998
WO	WO 98/41623	9/1998
WO	WO 98/41653	9/1998
WO	WO 98/42832	10/1998
WO	WO 99/29902	6/1999
WO	WO 00/04190	1/2000
WO	WO 00/06718	2/2000
WO	WO 00/09727	2/2000
WO	WO 00/18906	4/2000

OTHER PUBLICATIONS

Abastado et al., "Processing of complex heteroduplexes in *Escherichia coli* and *Cos-1* monkey cells," *PNAS* 81(18):5792-5796 (1984).

Adey et al., "Preparation of second-generation phage libraries," *Phage Disp. Pept. Proteins*, eds. Kay et al., pp. 277-291 (1996).

Amended Statement of Particulars re: opposition 703264 in Australia (Jan. 25, 2001).

Andersson et al., "Muller's ratchet decreases fitness of a DNA-based microbe", *PNAS*, 93: 906-907 (Jan. 1996).

Arkin et al., "An Algorithm for Protein Engineering: Simulations of Recursive Ensemble Mutagenesis" *Proc. Natl. Acad. Sci. USA*, 89(16):7811-7815 (1992).

Atreya et al., "Construction of in-frame chimeric plant genes by simplified PCR strategies," *Plant Mol. Biol.*, 19:517-522 (1992).

Bailey, "Toward a Science of Metabolic Engineering", *Science*, 252: 1668-1680 (1991).

- Balint et al., "Antibody Engineering By Parsimonious Mutagenesis", *Gene*, 137(1):109-118 (1993).
- Barrett et al., "Genotypic analysis of multiple loci in somatic cells by whole genome amplification", *Nuc. Acids Res.*, 23(17): 3488-3492 (1995).
- Bartel et al., "Isolation of New Ribozymes From a Large Pool of Random Sequences", *Science*, 261:1411-1418 (1993).
- Beaudry et al., "Directed Evolution of an RNA Enzyme," *Science*, 257:635-641 (1992).
- Berger et al., "Phoenix Mutagenesis: One-Step Reassembly of Multiply Cleaved Plasmids With Mixtures of Mutant and Wild-Type Fragments," *Anal. Biochem.*, 214:571-579 (1993).
- Berkhout et al., "In Vivo Selection of Randomly Mutated Retroviral Genomes," *Nucleic Acids Research*, 21(22):5020-5024 (1993).
- Biotransformations, Pathogenesis, and Evolving Biotechnology, Program and Abstracts, Pseudomonas '89, American Society for Microbiology and the University of Illinois, Jul. 9-13, 1989, abstracts 11-21 to 11-25.
- Blattman, A., Letter of Feb. 27, 2002 to Commissioner for Patents re: Australian Application 16241/99.
- Bock et al., "Selection of single-stranded DNA molecules that bind and inhibit human thrombin," *Nature*, 355:564-566 (Feb. 2, 1992).
- Cadwell et al., "Randomization of Genes by PCR Mutagenesis," *PCR Methods and Applications*, 2:28-33 (1992).
- Calogero et al., "In Vivo Recombination and the Production of Hybrid Genes," *Microbiology Letters*, 76:41-44 (1992).
- Cameron et al., "Cellular and Metabolic Engineering An Overview", *Applied Biochem. Biotech.*, 38: 105-140 (1993).
- Caren et al., "Efficient Sampling of Protein Sequence Space for Multiple Mutants," *Biotechnology*, 12(5):517-520 (1994).
- Carter, P., "Improved Oligonucleotide-Directed Mutagenesis Using M13 Vectors," *Methods in Enzymology*, 154:382-383 (1985).
- U.S. Appl. No. 60/067,908 filed Dec. 8, 1997.
- Chakrabarty, "Microbial Degradation of Toxic Chemicals: Evolutionary Insights and Practical Considerations", *ASM News*, 62(3): 130-137 (1996).
- Chang et al., "Recombination following transformation of *Escherichia coli* by heteroduplex plasmid DNA molecules," *Gene*, 29:255-261 (1984).
- Chang, et al., "Recombination following transformation of *Escherichia coli* by heteroduplex plasmid DNA molecules", *Gene*, 29:255-261 (1984).
- Chater, "The Improving Prospects for Yield Increase by Genetic Engineering in Antibiotic-Producing Streptomycetes", *Biotechnology*, 8: 115-121 (Feb. 1990).
- Chen et al., "Tuning the activity of an enzyme for unusual environments: Sequential random mutagenesis of subtilisin E for catalysis in dimethylformamide", *PNAS*, 90: 5618-5622 (Jun. 1993).
- Clackson et al., "Making antibody fragments using phage display libraries," *Nature*, 352:624-628 (Aug. 15, 1991).
- Claverys et al., "Heteroduplex Deoxyribonucleic Acid Base Mismatch Repair in Bacteria," *Microbiological Reviews*, 50(2): 133-165 (1986).
- Collet et al., "A Binary plasmid System for shuffling combinatorial antibody Libraries," *PNAS*, 89(21):10026-10030 (1992).
- Cramer et al., "10(20)-Fold aptamer library amplification without gel purification," *Nuc. Acids Res.*, 21(18):4410 (1993).
- Cramer et al., "Combinatorial Multiple Cassette Mutagenesis Creates All The Permutations Of Mutant And Wild-Type Sequences", *Biotechniques*, 18(2):194-196 (1995).
- Cramer et al., "Construction And Evolution Of Antibody-Phage Libraries By DNA Shuffling", *Nat. Med.*, 2(1):100-102 (1996).
- Cramer et al., "DNA Shuffling Of A Family Of Genes From Diverse Species Accelerates Directed Evolution", *Nature*, 391(3664):288-291 (1998).
- Cramer et al., "Improved Green Fluorescent Protein By Molecular Evolution Using DNA Shuffling", *Nat. Biotechnol.*, 14(3):315-319 (1996).
- Cramer et al., "Molecular Evolution Of An Arsenate Detoxification Pathway By DNA Shuffling", *Nat. Biotechnol.*, 15(5):436-438 (1997).
- Cull et al., "Screening for receptor ligands using large libraries of peptides linked to the C terminus of the *lac* repressor," *PNAS*, 89:1865-1869 (Mar. 1992).
- Cwirla et al., "Peptides on phage: A vast library of peptides for identifying ligands," *PNAS*, 87:6378-6382 (Aug. 1990).
- Daugherty et al., "Polymerase chain reaction facilitates the cloning, CDR-grafting, and rapid expression of a murine monoclonal antibody directed against the CD18 component of leukocyte integrins," *Nuc. Acids Res.*, 19(9):2471-2476 (1991).
- Declaration of Gerald F. Joyce in Opposition to Australian Patent Application 746786 (Aug. 8, 2004).
- Delagrave et al., "Recursive Ensemble Mutagenesis," *Protein Engineering*, 6(3):327-331 (1993).
- Delagrave et al., "Searching Sequence Space to Engineer Proteins: Exponential Ensemble Mutagenesis," *Biotechnology*, 11:1548-1552 (Dec. 1993).
- Dieffenbach et al., *PCR Primer, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, pp. 583-589, 591-601, 603-612, and 613-621 (1995).
- Dohet et al., "Methyl-directed repair of frameshift mutations in heteroduplex DNA," *PNAS*, 83(10):3395-3397 (1986).
- Dube et al., "Artificial mutants Generated by the Insertion of Random Oligonucleotides into the Putative Nucleoside Binding Site of the HSV-1 Thymidine Kinase Gene," *Biochemistry*, 30(51):11760-11767 (1991).
- Evnin et al., "Substrate specificity of trypsin investigated by using a genetic selection", *PNAS*, 87: 6659-6663 (Sep. 1990).
- Fang et al., "Methyl-directed Repair of Mismatched Small Heterologous Sequences in Cell Extracts from *Escherichia coli*," *J. Biological Chemistry*, 272(36):22714-22720 (1997).
- Fang et al., Human Strand-specific Mismatch Repair Occurs by a Bidirectional Mechanism Similar to.
- Feinberg et al., "A Technique for Radiolabeling DNA Restriction Endonuclease Fragments to High Specific Activity," *Anal. Biochem.*, 132:6-13 (1983).
- Fisch et al., "A Strategy Of Exon Shuffling For Making Large Peptide Repertoires Displayed On Filamentous Bacteriophage", *Proc Natl Acad Sci USA*, 93(15):7761-7766 (1996).
- Folger et al., "Efficient Correction of Mismatched Bases in Plasmid Heteroduplexes Injected into Cultured Mammalian Cell Nuclei," *Molecular and Cellular Biology*, 5(1):70-74 (1985).
- Fullen et al., "Genetic Algorithms and Recursive Ensemble Mutagenesis in Protein Engineering," *Complexity Int'l 1994 I*, printed from website <http://www.csu.edu.au/ci/vol1/fullen/REM.html> on Dec. 7, 1999.
- Gates et al., "Affinity Selective Isolation Of Ligands From Peptide Libraries Through Display On A Lac Repressor 'Headpiece Dimer'", *J. Mol. Biol.*, 255(3):373-386 (1996).
- Ghosh et al., "Arginine-395 Is Required for Efficient in Vivo and in Vitro Aminoacylation of tRNAs by *Escherichia coli* Methionyl-tRNA Synthetase," *Biochemistry*, 30:11767-11774 (1991).
- Goldman et al., "An Algorithmically Optimized Combinatorial Library Screened by digital Imaging Spectroscopy," *Biotechnology*, 10:1557-1561 (Dec. 1992).
- Graf et al., "Random circular permutation of genes and expressed polypeptide chains: Application of the method to the catalytic chains of aspartate transcarbamoylase," *PNAS*, 93:11591-11596 (1996).
- Gram et al., "In Vitro Selection and Affinity Maturation of Antibodies From a Naive Combinatorial Immunoglobulin Library", *Proc. Natl. Acad. Sci. USA*, 89:3576-3580 (1992).
- Greener et al., "An Efficient Random Mutagenesis Technique Using An *E. coli* Mutator Strain", *Methods in Molecular Biology*, 57:375-385 (1995).
- Harlow et al., "Construction of Linker-Scanning Mutations using the Polymerase Chain Reaction," *Methods in Mol. Biol.*, 31:87-96 (1994).
- Heda et al., "A simple in vitro site directed mutagenesis of concatamerized cDNA by inverse polymerase chain reaction," *Nuc. Acids Res.*, 20(19):5241-5242 (1992).
- Heim et al., "Wavelength Mutations And Posttranslational Autoxidation Of Green Fluorescent Protein" *Proc. Natl. Acad. Sci. USA*, 91(26):12501-12504 (1994).

- Hermes et al., "Searching Sequence Space by Definably Random Mutagenesis: Improving the Catalytic Potency of an Enzyme," *Proc. Natl. Acad. Sci. USA*, 87(2):696-700 (1990).
- Higuchi et al., "A general method of in vitro preparation and specific mutagenesis of DNA fragments: study of protein and DNA interactions," *Nuc. Acids Res.*, 16(15):7351-7367 (1988).
- Ho et al., "DNA and Protein Engineering Using the Polymerase Chain Reaction: Splicing by Overlap Extension," *DNA and Protein Eng. Techniques*, 2(2):50-55 (1990).
- Ho et al., "Site-Directed Mutagenesis by Overlap Extension Using the Polymerase Chain Reaction" *Gene*, 77:51-59 (1989).
- Hodgson, "The Whys and Wherefores of DNA Amplification," *Biotechnology*, 11:940-942 (Aug. 1993).
- Horton et al., "Engineering Hybrid Genes Without the Use of Restriction Enzymes: Gene Splicing by Overlap Extension," *Gene*, 77:61-68 (1989).
- Horton et al., "Gene Splicing by Overlap Extension," *Methods in Enzymology*, 217:270-279 (1993).
- Horton et al., "Gene Splicing by Overlap Extension: Tailor-Made Genes Using the Polymerase chain Reaction," *BioTechniques*, 8(5):528-535 (May 1990).
- Ippolito et al., "Structure assisted redesign of a protein-zinc-binding site with femtomolar affinity", *PNAS*, 92: 5017-5021 (May 1995).
- Janczewski et al., "Molecular phylogenetic inference from saber-toothed cat fossils of Rancho La Brea," *PNAS*, 89:9769-9773 (1992).
- Jayaraman et al., "Polymerase chain reaction-mediated gene synthesis: Synthesis of a gene coding for isozyme c of horseradish peroxidase," *PNAS*, 88:4084-4088 (May 1991).
- Jones et al., "A Rapid Method for Recombination and Site-Specific Mutagenesis by Placing Homologous ends on DNA Using Polymerase Chain Reaction," *BioTechniques*, 10(1): 62-66 (1991).
- Jones et al., "Recombinant Circle PCR and Recombination PCR for Site-Specific Mutagenesis Without PCR Product Purification" *Biotechniques* 12(4):528-534 (1992).
- Joyce, G. F., "Directed Molecular Evolution," *Scientific American*, (Dec. 1992).
- Kang et al., "Antibody redesign by chain shuffling from random combinatorial immunoglobulin libraries," *PNAS*, 88(24):11120-11123 (1991).
- Kellogg et al., "Plasmid-Assisted Molecular Breeding: New Technique for Enhanced Biodegradation of Persistent Toxic Chemicals", *Science*, 214: 1133-1135 (Dec. 4, 1981).
- Kim et al., "Cloning and Nucleotide Sequence of the Collb Shufflon," *Plasmid*, 22:180-184 (1989).
- Kim et al., "Human Immunodeficiency Virus Reverse Transcriptase," *The Journal of Biological Chemistry*, 271(9):4872-4878 (1996).
- Klug et al., "Creating chimeric molecules by PCR directed homologous DNA recombination," *Nuc. Acids Res.*, 19(10):2793 (1991).
- Komano et al., "Distribution of Shufflon among Incl Plasmids," *J. Bacteriology*, 169(11):5317-5319 (1987).
- Komano et al., "Physical and Genetic Analyses of Incl2 Plasmid R721: Evidence for the Presence of Shufflon," *Plasmid*, 23:248-251 (1990).
- Kramer et al., "Oligonucleotide-directed construction of mutations via gapped duplex DNA," *Methods in Enzymology*, 154:350-367 (1987).
- Krishnan et al., "Direct and crossover PCR amplification to facilitate Tn5supF-based sequencing of I phage clones," *Nuc. Acids Res.*, 19(22):6177-6182 (1991).
- Kunkel et al., "Rapid and efficient site-specific mutagenesis without phenotypic selection," *Methods in Enzymology*, 154:367-382 (1987).
- Kunkel, "Rapid and efficient site-specific mutagenesis without phenotypic selection", *PNAS*, 82: 488-493 (Jan. 1985).
- Lahue et al., "DNA Mismatch Correction in a Defined System," *Science*, 245:160-164 (1989).
- Leung et al., "A Method For Random Mutagenesis of a Defined DNA Segment Using a Modified Polymerase Chain Reaction," *Techniques*, 1:11-15 (1989).
- Levichkin et al., "A New Approach to Construction of Hybrid Genes: Homolog Recombination Method", *Mol. Biology*, 29(5) part 1: 572-577 (1995).
- Lewis et al., "Efficient site directed in vitro mutagenesis using ampicillin selection", *Nuc. Acids Res.*, 18(12): 3439-3443 (1990).
- List of Publications authored by Gerald F. Joyce, 2008.
- Lorberboum-Calski et al., "Cytotoxic activity of an interleukin 2-*Pseudomonas* exotoxin chimeric protein produced in *Escherichia coli*," *PNAS*, 85:1922-1926 (1988).
- Lowman, H.B. et al., "Affinity Maturation of Human Growth Hormone by Monovalent Phage Display," *J. Mol. Biol.*, 234:564-578 (1993).
- Majumder, K., "Ligation-free gene synthesis by PCR: synthesis and mutagenesis at multiple loci of a chimeric gene encoding OmpA signal peptide and hirudin" *Gene*, 110:89-94 (1992).
- Marks et al., "By-passing Immunization, Human Antibodies from V-gene Libraries Displayed on Phage," *J. Mol. Biol.*, 222:581-597 (1991).
- Marks et al., "By-Passing Immunization: Building High Affinity Human Antibodies by Chain Shuffling," *Bio/Technology*, 10:779-783 (1992).
- Marton et al., "DNA Nicking Favors PCR Recombination", *Nucleic Acids Res.*, 19(9):2423-2426 (1991).
- Maryon et al., "Characterization of recombination intermediates from DNA injected into *Xenopus laevis* oocytes: evidence for a nonconservative mechanism of homologous recombination," *Mol. Cell Biol.*, 11(6):3278-3287 (1991).
- McCafferty et al., "Phage antibodies: filamentous phage displaying antibody variable domains," *Nature*, 348:552-554 (Dec. 6, 1990).
- Meyerhans et al., "DNA Recombination During PCR," *Nucleic Acids Research*, 18(7):1687-1691 (1990).
- Michael, S.F., "Thermostable Ligase-Mediated Incorporation of Mutagenic Oligonucleotides During PCR Amplification," chapter 19 from *Methods in Molecular Biology, PCR Cloning Protocols from Molecular Cloning to Genetic Engineering*, eds. B. White, Humana Press, totowa, New Jersey, pp. 189-195 (1997).
- Moore et al., "Directed evolution of a *para*-nitrobenzyl esterase for aqueous-organic solvents", *Nature Biotech.*, 14: 458-467 (Apr. 1996).
- Morl et al., "Group II intron RNA-catalyzed recombination of RNA in vitro," *Nuc. Acids Res.*, 18(22):6545-6551 (1990).
- Mullis et al., "Specific Enzymatic Amplification of DNA In Vitro: The Polymerase Chain Reaction," Cold Spring Harbor Symposia on Quantitative Biology, 51:263-273 (1986).
- Mullis et al., "Specific Synthesis of DNA in Vitro via a Polymerase-Catalyzed Chain Reaction," *Methods in Enzymology*, 155:335-351 (1987).
- Near, "Gene Conversion Of Immunoglobulin Variable Regions In Mutagenesis Cassettes By Replacement PCR Mutagenesis", *Biotechniques*, 12(1):88-97 (1992).
- Ner et al., "Laboratory Methods: A Simple and Efficient Procedure for Generating Random Point Mutations and for Codon Replacements Using Mixed Oligodeoxynucleotides," *DNA*, 7(2):127-134 (1988).
- Nissim et al., "Antibody fragments from a 'single pot' display library as immunochemical reagents," *EMBO Journal*, 13(3):692-698 (1994).
- Oliphant et al., "Cloning of Random-Sequence Oligodeoxynucleotides," *Gene*, 44(2-3):177-183 (1986).
- Olsen et al., "Hybrid *Bacillus* (1-3,1-4)-beta-glucanases: engineering thermostable enzymes by construction of hybrid genes," *Mol. Gen. Genet.*, 225(2):177-185 (1991).
- Omura, "Philosophy of New Drug Discovery", *Microbiol. Rev.*, 50(3): 259-279 (Sep. 1986).
- Opposition Statement in matter of Australian Patent Application 703264 (Affymax Technologies NV), filed by Diversa Corporation on Sep. 23, 1999.
- Osuna et al., "Combinatorial mutagenesis of three major groove-contacting residues of *Eco* RI: single and double amino acid replacements retaining methyltransferase-sensitive activities," *Gene*, 106:7-12 (1991).
- Paabo et al., "DNA Damage Promotes Jumping between Templates during Enzymatic Amplification," *J. Biol. Chem.*, 265(8):4718-4721 (Mar. 15, 1990).

- Perlak, "Single Step Large Scale Site-Directed In Vitro Mutagenesis Using Multiple Oligonucleotides", *Nucleic Acids Res.*, 18(24):7457-7458 (1990).
- Pharmacia Catalog* pp. 70-71 (1993 Edition).
- Piepersberg, "Pathway Engineering in Secondary Metabolite-Producing Actinomycetes", *Crit. Rev. Biotech.*, 14(3):251-285 (1994).
- Pompon et al., "Protein Engineering by cDNA Recombination in Yeasts: Shuffling of Mammalian Cytochrome P-450 Functions," *Gene*, 83(1):15-24 (1989).
- Prasher, "Using GFP to see the light", *TIG*, 11(8) (Aug. 1995).
- Prodromou et al., "Protocol, Recursive PCR: a novel technique for total gene synthesis," *Protein Engineering*, 5(8):827-829 (1992).
- Rao et al., "Recombination and Polymerase Error Facilitate Restoration of Infectivity in Brome Mosaic Virus," *Journal of Virology*, 67(2):969-979 (1993).
- Rapley, "Enhancing PCR Amplification And Sequencing Using DNA-Binding Proteins", *Mol. Biotechnol.*, 2(3):295-298 (1994).
- Reidhaar-Olson et al., "Combinatorial Cassette Mutagenesis as a Probe of the Informational Content of Protein Sequences," *Science*, 241:53-57 (1988).
- Request for leave to amend the Statement of Grounds and Particulars re: opposition 703264 in Australia (Jan. 25, 2001).
- Rice et al., "Random PCR mutagenesis screening of secreted proteins by direct expression in mammalian cells", *PNAS*, 89: 5467-5471 (Jun. 1992).
- Robles et al., "Hydropathy and Molar Volume Constraints on Combinatorial mutants of the Photosynthetic Reaction Center," *J. Mol. Biol.*, 232:242-252 (1993).
- Rouwendal et al., "Simultaneous Mutagenesis of Multiple Sites: Application of the Ligase Chain Reaction Using PCR Products Instead of Oligonucleotides," *BoiTechniques*, 15(1):68-70, 72-74, 76 (1993).
- Saiki et al., "analysis of enzymatically amplified b-globin and HLA-DQa DNA with allele-specific oligonucleotide probes," *Nature*, 324:163-166 (Nov. 13, 1986).
- Saiki et al., "Diagnosis of sickle Cell Anemia and b-Thalassemia with Enzymatically Amplified DNA and Nonradioactive Allele-Specific Oligonucleotide Probes," *New England J. of Medicine*, 319(9):537-541 (Sep. 1, 1988).
- Saiki et al., "Enzymatic Amplification of b-Globin Genomic Sequences and Restriction Site analysis for Diagnosis of Sickle Cell Anemia," *Science*, 230:1350-1354 (Dec. 20, 1985).
- Saiki et al., "Primer-Directed Enzymatic Amplification of DNA with a Thermostable DNA Polymerase," *Science*, 239:487-491 (Jan. 20, 1988).
- Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd edition, Cold Spring Harbor Laboratory Press, pp. 14.2, 14.34, and 14.35 (1989).
- Sambrook et al., *Molecular Cloning A Laboratory Manual*, 2nd Edition, Cold Spring Harbor Laboratory Press, table of contents and pp. 5.15-5.27, 15.1-15.113, 16.1-16.81, and 17.1-17.44 (1989).
- Sambrook et al., *Molecular Cloning, A Laboratory Manual*, Cold Spring Laboratory Press, Cold Spring Harbor, New York (1989).
- Sandhu et al., "Dual Asymmetric PCR: One-Step Construction of Synthetic Genes," *BioTechniques*, 12(1):14-16 (1992).
- Scharf et al., "Direct Cloning and Sequence Analysis of Enzymatically Amplified Genomic Sequences," *Science*, 233:1076-1078 (Sep. 1986).
- Scott et al., "Searching for Peptide Ligands with an Epitope Library," *Science*, 249:386-390 (Jul. 20, 1990).
- Shao et al., "Random-priming in vitro recombination: an effective tool for directed evolution," *Nuc. Acids Res.*, 26(2):681-683 (1998).
- Shi et al., "Rapid PCR Construction of a Gene Containing Lym-1 Antibody Variable Regions," *PCR Methods and Applications*, 3:46-53 (1993).
- Shuldiner et al., "Hybrid DNA artifact from PCR of closely related target sequences," *Nuc. Acids Res.*, 17(11):4409 (1989).
- Sikorski et al., "In Vitro Mutagenesis and Planned Shuffling: From Cloned Gene to Mutant Yeast," *Methods in Enzymology*, 194:302-318 (1991).
- Simpson et al., "Two paradigms of metabolic engineering applied to amino acid biosynthesis", *Biochem. Soc. Transactions*, vol. 23 (1995).
- Smith et al., "Localized sex in bacteria," *Nature*, 349:29-31 (1991).
- Smith et al., "Unwanted Mutations in PCR Mutagenesis: Avoiding the Predictable," *PCR Methods and Applications*, 2(3):253-257 (Feb. 1993).
- Statutory Declaration of Dr. Gerald Joyce in Australian Opposition against application 703264, 2008.
- Statutory Declaration of Mae Li Gan in Australian Opposition against application 703264, 2008.
- Statutory Declaration of Ngarie Petit-Young in Australian Opposition against application 703264, 2008.
- Statutory Declaration of Ruth Bird in Australian Opposition against application 703264, 2008.
- Steele et al., "Techniques for Selection of Industrially Important Microorganisms", *Ann. Rev. Microbiol.*, 45: 89-106 (1991).
- Stemmer et al., "Increased Antibody Expression from *Escherichia-coli* Through Wobble-Base Library Muatagenesis by Enzymatic Inverse PCR," *Gene*, 123(1):1-7 (1993).
- Stemmer et al., "A 20-Minute Ethidium Bromide High-slat Extraction Protocol for Plasmid DNA," *Biotechniques*, 10(6):726 (1991).
- Stemmer et al., "Enzymatic Inverse PCR—A Restriction Site Independent, single-Fragment Method for High-Efficiency, Site-Directed Mutagenesis," *Biotechniques*, 13(2):214 (1992).
- Stemmer et al., "Expression of Antibody FV Fragments Specific for a Heavy Metal Chelate Indium Edta in *Escherichia-coli*," *J. Cell Biochem.*, Suppl. 0(15 part G), p. 217 (1991).
- Stemmer et al., "Selection of an Active Single Chain FV Antibody from a Protein Linker Library Prepared by Enzymatic Inverse PCR," *Biotechniques*, 14(2):256-265 (1992).
- Stemmer et al., "Single-Step Assembly Of A Gene And Entire Plasmid From Large Numbers Of Oligodeoxyribonucleotides", *Gene*, 164(1):49-53 (1995).
- Stemmer, "DNA Shuffling by Random Fragmentation and Reassembly: In Vitro Recombination for Molecular Evolution" *Proc. Natl. Acad. Sci. USA*, 91(22):10747-10751 (1994).
- Stemmer, "Rapid Evolution of a Protein in Vitro by DNA Shuffling," *Nature*, 370:389-391 (1994).
- Stemmer, "Searching Sequence Space", *Biotechnology*, 13:549-553 (1995).
- Stemmer, "Sexual PCR and Assembly PCR" *Encyclopedia Mol. Biol.*, VCH Publishers, New York, pp. 447-457 (1996).
- Stemmer, "The Evolution of Molecular Computation", *Science*, 270(5241):1510 (1995).
- Stephanopoulos et al., "Metabolic engineering—methodologies and future prospects", *Trends Biotech.* 11: 392-396 (1993).
- Stephanopoulos, "Metabolic engineering", *Curr. Opin. Biotech.*, 5: 196-200 (1994).
- Strickberger, M. W., cover page, preface, table of contents, and pp. 844-846 from *Genetics, Second Edition*, MacMillan Publishing Co., Inc., 1976.
- That of the Bacterial Reaction, *J. Biol. Chem.*, 268(16): 11838-11844 (Jun. 5, 1993).
- Umar et al., "DNA Loop Repair by Human Cell Extracts," *Science*, 266:814-816 (1994).
- Vassilyev et al., "DNA-repair enzymes," *Current Opinion in Structural Biology*, 7:103-109 (1997).
- Villarreal et al., "A General Method of Polymerase-Chain-Reaction-Enabled Protein Domain Mutagenesis: Construction of a Human Protein S-Osteonectin Gene," *Analytical Biochem.*, 197:362-367 (1991).
- Volkov et al., "Random Chimeragenesis by heteroduplex recombination," *Methods in Enzymology*, 328:456-463 (2000).
- Volkov et al., "Recombination and chimeragenesis by in vitro heteroduplex formation and in vitro repair," *Nuc. Acids. Research*, 27(18):i-vi (1999).
- Wang et al., "Identification Of Glial Filament Protein And Vimentin In The Same Intermediate Filament System In Human Glioma Cells", *Proc. Natl. Acad. Sci. USA*, 81(7):2102-2106 (1984).
- Weber et al., "Formation of Genes Coding for Hybrid Proteins by Recombination Between Related, Cloned Genes in *E. Coli*," *Nucleic Acids Research*, 11(16):5661-5669 (1983).
- Wehmeier, "New multifunctional *Escherichia coli-Streptomyces* shuttle vectors allowing blue-white screening on XGal plates", *Gene*, 165: 149-150 (1995).

- Weissenhorn et al., "Chimerization of antibodies by isolation of rearranged genomic variable regions by the polymerase chain reaction," *Gene*, 106:273-277 (1991).
- Winter et al., "Making Antibodies By Phage Display Technology", *Ann. Rev. Immunol.*, 12:433-455 (1994).
- Wu et al., "Allele-specific enzymatic amplification of beta-globin genomic for diagnosis of sickle cell anemia," *PNAS*, 86(6):2757-2760 (1989).
- Yao et al., "Site-directed Mutagenesis of Herpesvirus Glycoprotein Phosphorylation Sites by Recombination Polymerase Chain Reaction," *PCR Methods and Applications*, 1(3):205-207 (Feb. 1992).
- Yolov et al., "Constructing DNA by polymerase recombination," *Nuc. Acids Res.*, 18(13):3983-3986 (1990).
- Yon et al., "Precise gene fusion by PCR," *Nuc. Acids Res.*, 17(12):4895 (1989).
- Youvan et al., "Recursive Ensemble Mutagenesis: A Combinatorial Optimization Technique for Protein Engineering," from *Parallel Problem Solving from Nature*, 2, Manner eds., pp. 401-410 (1992).
- Zarucki-Schulz et al., "Point Mutagenesis of the Ovalbumin Gene Promoter Sequence and Its Effect on in Vitro Transcription," *J Biol. Chem.*, 257(18):11070-11077 (1982).
- Zhang et al., "Directed Evolution Of A Fucosidase From A Galactosidase By DNA Shuffling And Screening", *Proc. Natl. Acad. Sci. USA*, 94(9):4504-4509 (1997).
- Zhao et al., "Molecular Evolution by Staggered Extension Process (StEP) In Vitro Recombination", *Nature Biotech.*, 16:258-261 (1998).
- Zoller et al., "Oligonucleotide-directed mutagenesis: a simple method using two oligonucleotide primers and a single-stranded DNA template," *Methods in Enzymology*, 154:329-350 (1987).
- Zoller, M.J., "New recombinant DNA methodology for protein engineering," *Curr. Opin. Biotech.*, 3:348-354 (1992).

* cited by examiner




FIG. 1




FIG. 2




FIG. 3




FIG. 4

1 MASVLTNINAMSALQTLRSISSNMEDTQSRISSGMRVGSASDNAAWYSIATTMRSDNASLSAVQDAIGLG R. lupini Flaa
|.:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:
1 MTSILTNNSAMAAALSTLRSISSMEDTQSRISSGLRVGSASDNAAWYSIATTMRSDNQALSAVQDALGLG R. meliоти Flaa

71 AAKVDTASAGMDAVIDVVKQIKNKLVTAEQESSADKTKIQGEVVKQLQEQKGIVDASFSNGENWLKGDLS⁷¹ R. lupini Flaa
|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:
71 AAKVDTAYS⁷¹GMESAIEVVKEIKAKLVAAATEDGV⁷¹DKAKIQEEITQLKDLT⁷¹SIAEAA⁷¹SFSGENWLQADLSG R. meliotti Flaa

141 TT.TKSVVGSFVRE.GGT¹⁴¹VSVK¹⁴¹TIDYALNASKVLVDTRA¹⁴¹TGK¹⁴¹TGILDTAY¹⁴¹TGLNANTV¹⁴¹TDINKGGVIT R. lupini Flaa
|.:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:
141 GPVTKSVVGGFVRDSSGAVSVK¹⁴¹VDYSLN¹⁴¹DTVLFD...TTGNTGILDK¹⁴¹VY.NV¹⁴¹SQASVTL¹⁴¹PNVNGTTS R. meliotti Flaa

209 QASVRAYSTDEMLSLGAKVDGANSNVAVGGSASSRSTAAGLRVASTLRPPSPHQHQLASLPPLTPFLK R. lupini Flaa
|.:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:
207 EYTVGAYNVDDLIDASATFDGDYANVGAGALAGDYVKVQGSWVKAVDVAATGQE R. meliotti Flaa

279 LVLQLLPVT²⁷⁹SPSSTK²⁷⁹P²⁷⁹TAA²⁷⁹P.VQVNLT²⁷⁹QSVLT²⁷⁹MDVS.SM²⁷⁹SSTD²⁷⁹VGSYL²⁷⁹TGVEKAL²⁷⁹TSL²⁷⁹SAGAE²⁷⁹LSIKQ R. lupini Flaa
|.:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:
277 VVYDDGTTK²⁷⁷WGVDTT²⁷⁷VTGAPAT²⁷⁷NVAA²⁷⁷PASIA²⁷⁷TIDITIAAQAGNLDAL²⁷⁷IAGVDEAL²⁷⁷TDMTSA²⁷⁷AAASLS²⁷⁷SISS R. meliotti Flaa

347 RIDLQVDFASKLGDALAKGIGRLVDADMNEESTK³⁴⁷KLQ³⁴⁷TQ³⁴⁷QLAIQ³⁴⁷LSIANS³⁴⁷SDSQN³⁴⁷I³⁴⁷LSLPR 410 R. lupini Flaa
|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:.|:
331 RIDLQSD³³¹FV³³¹NKLS³³¹DS³³¹IDS³³¹GV³³¹RLVD³³¹ADMNEEST³³¹RL³³¹KAL³³¹Q³³¹TQ³³¹QLAIQ³³¹ALS³³¹ANS³³¹SDSQN³³¹V³³¹LSLFR 394 R. meliotti Flaa

FIG. 5




FIG. 6A




FIG. 6B

SCS01 ATGGCAAGCGTTCTCAAAACATTAACGCAATGTCTGCTCTTCAGACGGTTCGATTTCTTCCAAACATGGAGACACCCAGAGCCGGTATTTCCAGCG
SCS01 GCATGCGCGTTGGTTCGGCTTCCGACAAACGCCGCTTATTGGTCTATCGGACCAACCATGCGCTCGGACAAATGCCCTCGCTTTCCGGTTCAGGATGCAAT
SCS01 TGGCCTCGGTGCGCCCAAGGTCGATACCGCTTCGGCGGGTATGGATGCGGTTATCGATGTTGTAAGCAGATCAAGAACAAACTGGTCACTGCCACCCGAA
SCS01 GACGGCTCGACAAAGCCAAAGATCCAAAGAAGAAATCACTCAGCTCAAGGACCAAGCTGACGAGCATCGCCGCTTCCCTTCCGGTGAATAACTGGC
SCS01 TCAAGGGCGATCTTTCCACGACGAC...AACCAATCAGTGGTTCCTTCGTTTCGTTGA...AGGCGGTACCGTATCGGTCAAGACCATCGATTAC
SCS01 GCTCTGAATGCTTCCAAAGTTCGGTGGATACCCGCAACGGGCAACAGACCGGCAATTCGGACAAGGTCTACAAACGCTTCGCAGGCAAGCGTCAAGC
SCS01 TGACGG...TCAACACCAACGGGTCGAAATCCAGGCTCCGTCGGCCCTATTCGCTGGA...GTCCCTCACCCGAAGCCGGTCCGGAGTT
SCS01 CCAGGCAACTATGCTCTTCAGGGGGTAAACAGCTACGTCGAAACCGTCTGGGT...TCGA.GCTGAG...ACCGCATCA
SCS01 ACACCAAGTCGCTGGCAAGTTTGCCCGCCTTACACCCCGCTGAAGCTGGTACTGCAGCTGCTGCCGGTGACGCCATCATCGTCGACGAAACCAACAGCG
SCS01 GCGCCGGTGCAGGTAAACCTCACCCAGTCGGTCCCTGACCAATGGATGTCAGCTCGATGAGCTCGACGGATGTCGGCAGCTACCTCACGGCGGTGGAAGAAG
SCS01 CTCTCACAGCCTGACGAGCGCTGGCGCTGAACCTCGGCTCTATCAAACAGCCATCGATCTGCAGGTTGATTTGCTTCCAAGCTGGCGACGCTCTCGC
SCS01 AAAAGGTAATTGGCCGCTCCTCGTTGATGCTGACATGAATGAAGTCCACTAAGCTTAAGGCTCTTCAGACGCGCAGCAGCTGGCTATCCAGTCGCTCTCC
SCS01 ATCGCAACACGCGACTCGCAGAACATTCCTGCTCGCTGTTCCGGTTAA

FIG. 7A

SCS02 ATGACGAGCATTTCTACCCAACTCCGGCAATGGCCCGCCTTTCCGGAGTGCCGCTCGATCTCTCCAGCATGGAAAGACACGCCAGAGCCGCAATCTCCTCCG
SCS02 GCCTTCGCGTCCGGTTCGGCTCCGACACCGCCCTACTGGTCCGATTCGGACACCAATGCCCTCCGACAAACAGGCCCTTTTCGGCCGTCAGGACGCCCT
SCS02 CGCCCTCGCCCGCCAAAGTTGATACCGCCTATTCCGGTATGGAATCGGCGATCGAAGTCGTTAAGGAAATCAAGAAACAACACTGGTCACTGCTCAGGAA
SCS02 TCTTCTGCCGACAAACGAGATTCAAGGCGAAGTCAAGCAGCTTCAGGAGCAGTTGAAGGGCaTCGTTGATTCCGCTTCCCTTCTCCGGTGAGAACTGGC
SCS02 TGCAGGCGGACC..TCAGCGCGCGCCGTCACCAAGAGCGTCCGTCGGCTCGTTCCGTGACGGAAAGCGGTTCCCGTAGCCGTCAGAAAGGTCGATTAC
SCS02 GCTCTGAATGCTTCCAAAGTTCTGGTGGATACCCCGGCAACGGGCACCAAGACCGGCATTCTCGATACTGCTTATACCGGCCCTTAACGCCA...ACACGG
SCS02 TGACGGTTGATATCAACAAGGGCGGTGATCACCCAGGCCCTCCGTCCGCCCTATCCACgGACGAAATGCTCTCCCTCGGGCGCAAGGTCGATGGCGC
SCS02 AAACAGCAACGTTGCTGTTGGCGGGG.CTCCGCTTCGTeAAGTTCGACGGCAGCTGGGTTAAGGGTAGCCTCGACgCTGGGCCCTCCATCACCCGATCA
SCS02 ACCGGCCCAACCGGTCAGAAATCGCCGC...CACCAACGACGGCAGCTGGTACCATCACTGCAGACAGCTGGGTCGTCCGATGTGGCAACGCTCCCTGCCC
SCS02 CCAACGTTTCGG...CCGGCCAGTCCGGTCGCGAACATCAACATCGTCGGAATGGGCTCGACCGGATGTCCGCGAGCTACCTCACGGCGTGGAAAGG
SCS02 CTCTCACCCAGCATGACCAAGCGCTGCCCGCTCGCTCGCATCTCCTCGCGCATCGACCTGCAGAGCGGAATTCTGTCAACRAAGCTCTCGGACTCGATCGA
SCS02 GTCGGCGTCCGGCCGTCCTCGTCGACGGGACATGAACGAGGAGTCCGACCCCGCTCAAGGCCCTGCAGACCCAGCAGCAGCTCCGCCATCCAGGCCCTGTCCG
SCS02 ATCGCCCAACTCGGACTCGCAGAACGTTCTGTCTCCGGCTAA

FIG. 7B

ES01 ATGACGAGCATTTCCACCACAACCTCCGCAATGGCCGCGCTTCCGGAGTGGCTCGATCTTCCAGCATGGAAAGACACGAGAGCCGCATCTCCTCCG
ES01 GCCTTCGGCTCGGTTCCGGCTCCGACAAACGCCCTACTGGTGGATTGGACCAACCATGGCTCCGACAAACCAGGCCCTTTCCGGCCGTCCAGGACGCCCT
ES01 CGGCCTCGGCCCGCAAGGTTGATACCCCTATTCGGTATGGAATCGGATCGAATCGAATCAAGGCCAAGCTCGTAGCTGCCACCCGAA
ES01 GACGGCTCGACAAAGCCCAAGATCCAAGAAGAAATCACTCAGCTCAAGGACCCAGCTGACGAGCATCGCCGACCGGCTTCTCCGGTGAGAACTGGC
ES01 TGCAGGGCGGACC..TCAGCGGGCGCCGTCACCAAGAGCTCGTCCGGCTCGTTCGGTCCGTGACGGAAAGCGGTTCCGTAGCCGTCAAGACCATCGATTAC
ES01 GCTCTGAATGCTTCCAAAGTTCTGGTGGATAACCCGGACACCGGTCCGGCATTCGGACAAGGTCTACAACGTCGCAAGGCAAGCGTCAAGCGTCAAGC
ES01 TGACGG.....TCAACACCACCGGCTCGAATCGCATACGGTTGCTGCCTATTGCTGGA.....GTCCCTCACCGAAGCCGGTCCGGAGTT
ES01 CCAGGGCAACTATGCTCTTCAGGGCGGTAACAGCTACGTCAAAGGTCGACGGCAGCTGGGTTAAGGGTAGCGTGCAGCCTGCGGCCCTCCATCACCCGCATCA
ES01 ACACCAGTCCGTGGCAAGTTGCCCGCCGCTTACACCCCGCTGAAGCTGGTACTGCAGCTGCTGCCGGTGCACCTCATCGTCGACGAAACCAACAGCG
ES01 GCGCCGGTGCAGGTAAACCTCACCCAGTCCGTTCTGACCATGGATGTCAGCTCGATGTCGGCAGCTACCTCACGGGCGTGGAAAAGG
ES01 CTCTCACCCAGCCTGACGAGCCGTGGCGCTGAACCTCCGCTCCATCTCCGCGCATCGACCTGCAGAGCGAATTCGTCAACAAGCTCTCGGACTCGATCGA
ES01 GTCGGCGTCCGGCCGTCCTCGTCGACCGGGACATGAACGAGGAGTCCGACCCGCTCAAGGCCCTGCAGACCCAGCAGCTCGCCATCCAGGCCCTGTGTCG
ES01 ATCGCCAACTCGGACTCGCAGAAGCTCTGTCTCCGCTCTTCCGGCTAA

FIG. 7C

ES02 ATGACGAGCATTCTCACCACAACTCCGCAATGGCCCGCGCTTCCGGAGTGCGCTCCTCAGCATGGAAGACACGAGCCGCAATCTCCCTCCG
 ES02 GCCTTCGCGTGGTTCGGCTCCGACACAGCCCGCTACTGGTCGATGCGACCAACCAGGCCCTTTCGGCCGTCCAGGACGCCCT
 ES02 CGCCTCGGCCCGCCAGGTTGATACCGCTATTCCGGTATGGAATCGGCCATCGAAGTCGTTAAGGAAATCAAGCCAGCTCGTAGCTGCCACCCGAA
 ES02 GACGGCGTCGACAAAGCCCAAGATCCAAGAAGAAATCACTAGCTCAAGGACCAAGCTGACGAGCATCGCCGACGGCTTCTTCCGGTGAGAACTGGC
 ES02 TGCAGGGCGACC..TCAGCGGCGGCGCTCACCAGAGCGTCCGTCGGTCGTCAGCGGAGCGGTTCCGTAGCCGTCGTAAGACCATCGATTAC
 ES02 GCTCTGAATGCTTCCAAGGTTCTGGTGGATACCCCGCAACGGCAACAGCCGCAATCTCGATACTGCTTATACCGGCTTAACCGA...ACACGG
 ES02 TGACGGTTGATATCAACAAGGGCGGTGATCACCCAGGCTCCGGTCCGAAATGCTCTCCTCACCGAAGCCGGTCCGGAGTT
 ES02 CCAGGGCAACTATGCTCTTCAGGGCGGTAACAGCTACGTCAGGTCGAAAACGTCTGGGT.....TCGA.GCTGAG.....ACCGCTGCA
 ES02 ACCGGCGCCACCGGTCAAGAAATCGCCG..CACCAAGACGGCAGCTGGTACCATCACTGCAGACAGCTGGGTCGTCGATGTCGGCAACGCTCCTGCCG
 ES02 CCAACGTTTCGG.....CCGGCCAGTCCGGTCGCGAACATCAACATCGTCCGGAATGGGTGCGAGCTGGCTGATCAGCGGTGTCGACGCCG
 ES02 CTTTGACAGACATGACAGCGGCTGCCGCTCCATCTCCTCGCCATCGACCTGCAGAGCGAATTCGTCACAAGCTCTCGGACTCGATCGA
 ES02 GTCGGGCGTCGGCCGTCTCGTCGACGGGACATGAACGAGGAGTCCGACCCGCTCAAGGCCCTGCAGACCCAGCAGCTCGCCATCCAGGCCCTGTCTG
 ES02 ATCGCCAACTCGGACTCGCAGAACGTCTCTCCGCTAA

FIG. 7D




FIG. 8




FIG. 9A




FIG. 9B




FIG. 10




FIG. 11




FIG. 12

1 CTGCAGCGTGCCAGCTGTTTCGTGGTGGTGATCGCGGCCGCGCTGGCCGCCGTCGCGGTC
 61 GCCGCCGCCGGGCCGATCGAGTTCGTGCGCTTCGTGCGTCCGCGAGATCGCCCTGCGGCTC
 121 TCGGGCGGCAGCCGGCCGCCCTGCTCGCCTCGGGCGATGCTCGGGCGGCTGCTGGTGGTC
 181 GGCGCCGACCTGGTTCGCTCAGATCGTGGTGGCGCCGAAGGAGCTGCCGGTCCGGCTGCTC
 241 ACCGCGATGATCGGCACCCCGTACCTGCTCTGGCTCCTGCTTCGGCGATCAAGAAAGGTG
 301 AGCGGATGAACGCCCGCCTGCGTGGCGAGGGCCTGCACCTCGCGTACGGGGACCTGACCG
 361 TGATCGACGGCCTCGACGTCGACGTGCACGACGGGCTGGTACCACCATCATCGGGCCCA
 421 ACGGGTGCGGCAAGTCGACGCTGCTCAAGGCGCTCGGCCGGCTGCTGCGCCCGACCGGCG
 481 GGCAGGTGCTGCTGGACGGCCCGCATCGACCGGACCCCAACCCGTGACGTGGCCCGGG
 541 TGCTCGGGCGTGTGCCGAGTCGCCACCGCGCCCGAAGGGCTCACCGTCGCCGACCTGG
 601 TGATGCGCGGCCGCCACCCGCACCAGACCTGGTTCGGCAGTGGTCCGCGGACGACGAGG
 661 ACCAGGTCCCGACGCGCTGCGCTGGACCGACATGCTGGCGTACGCGGACCGCCCGGTGG
 721 ACGCCCTCTCCGGCGGTTCAGCGCCAGCGCGCCTGGATCAGCATGGCGCTGGCCCAAGGCA
 781 CCGACCTGCTGCTGCTGGACGAGCCGACCACCTTCCTCGACCTGGCCCAACAGATCGACG
 841 TGCTGGACCTGGTCCGCCGGCTGCACGCCGAGATGGGCCGGACCGTGGTGGTGGTGC
 901 ACGACCTGAGCCTGGCCGCCCGGTACGCCGACCGGCTGATCGCGATGAAGGACGGCCGGA
 961 TCGTGGCGAGCGGGCGCCGGACGAGGTGCTCACCCCGCGCTGCTGGAGTCGGTCTTCG
 1021 GGCTGCGCGCGATGGTGGTGGCCGACCCGGCGACCGGCACCCCGCTGGTGGTATCCCCCTGC
 1081 CGCGCCCCGCCACCTCGGTGCGGGCCTGAAATCGATGAGCGTGGTTGCTTCATCGGCCTG
 1141 CCGAGCGATGAGAGTATGTGGGCGGTAGAGCGAGTCTCGAGGGGGAGATGCCGCCGTGAC

V T

1201 GTCCTCGTACATGCGCCTGAAAGCAGCAGCGATCGCCTTCGGTGTGATCGTGGCGACCGC
 3 S S Y M R L K A A A I A F G V I V A T A
 1261 AGCCGTGCCGTCACCCGCTTCGGCAGGGAACATGACGGCGGCTATGCGGCCCTGATCCG
 23 A V P S P A S G R E H D G G Y A A L I R
 1321 CCGGGCCTCGTACGGCGTCCCGCACATCACCGCCGACGACTTCGGGAGCCTCGGTTTCGG
 43 R A S Y G V P H I T A D D F G S L G F G
 1381 CGTCGGGTACGTGCAGGCCGAGGACAACATCTGCGTCATCGCCGAGAGCGTAGTGACGGC
 63 V G Y V Q A E D N I C V I A E S V V T A
 1441 CAACGGTGAGCGGTTCGCGGTGGTTCGGTTCGACCGGGCCGGACGACCGCGATGTGCGCAG
 83 N G E R S R W F G A T G P D D A D V R S

FIG. 13A

1501 CGACCTCTTCCACCGCAAGGCGATCGACGACCGCGTCGCCGAGCGGCTCCTCGAAGGGCC
103 D L F H R K A I D D R V A E R L L E G P

1561 CCGCGACGGCGTGCGGGCGCCGTCGGACGACGTCCGGGACCAGATGCGCGGCTTCGTCCG
123 R D G V R A P S D D V R D Q M R G F V A

1621 CGGCTACAACCACTTCCTACGCCGCACCGGCGTGCACCGCCTGACCGACCCGGCGTGCCG
143 G Y N H F L R R T G V H R L T D P A C R

1681 CGGCAAGGCCTGGGTGCGCCCGCTCTCCGAGATCGATCTCTGGCGTACGTTCGTGGGACAG
163 G K A W V R P L S E I D L W R T S W D S

1741 CATGGTCCGGGCCGGTTCGGGGCGCTGCTCGACGGCATCGTCGCCGCGACGCCACCTAC
183 M V R A G S G A L L D G I V A A T P P T

1801 AGCCGCCGGGCCCGCGTCAGCCCCGGAGGCACCCGACGCCCGCGATCGCCGCCGCCCT
203 A A G P A S A P E A P D A A A I A A A L

1861 CGACGGGACGAGCGCGGGCATCGGCAGCAACGCGTACGGCCTCGGCGCGCAGGCCACCGT
223 D G T S A G I G S N A Y G L G A Q A T V

1921 GAACGGCAGCGGGATGGTGCTGGCCAACCCGCACTTCCCCTGGCAGGGCGCCGCACGCTT
243 N G S G M V L A N P H F P W Q G A A R F

1981 CTACCGGATGCACCTCAAGGTGCCCGGCCGCTACGACGTTCGAGGGCGCGGCGCTGATCGG
263 Y R M H L K V P G R Y D V E G A A L I G

2041 CGACCCGATCATCGGGATCGGGCACAACCGCACGGTTCGCTGGAGCCACACCGTCTCCAC
283 D P I I G I G H N R T V A W S H T V S T

2101 CGCCCGCCGGTTCGTGTGGCACCGCCTGAGCCTCGTGCCCGGGCAGCCACCTCCTATTA
303 A R R F V W H R L S L V P G D P T S Y Y

2161 CGTCGACGGCCGGCCCCGAGCGGATGCGCGCCCGCACGGTTCACGGTCCAGACCGGCAGCGG
323 V D G R P E R M R A R T V T V Q T G S G

FIG. 13B

2221 CCCGGTCAGCCGCACCTTCCACGACACCCGCTACGGCCCGGTGGCCGTGATGCCGGGCAC
343 P V S R T F H D T R Y G P V A V M P G T

2281 CTTCGACTGGACGCCGGCCACCGGTACGCCATCACCGACGTCAACGCGGGCAACAACCG
363 F D W T P A T A Y A I T D V N A G N N R

2341 CGCCTTCGACGGGTGGCTGCGGATGGGCCAGGCCAAGGACGTCCGGGCGCTCAAGGCGGT
383 A F D G W L R M G Q A K D V R A L K A V

2401 CCTCGACCGGCACCAGTTCCTGCCCTGGGTCAACGTGATCGCCCGGACGCGCGGGGCGA
403 L D R H Q F L P W V N V I A A D A R G E

2461 GGCCCTCTACGGCGATCATTCGGTTCGTCGCCCCGGGTGACCGGCGCGCTCGCTGCCGCCTG
423 A L Y G D H S V V P R V T G A L A A A C

2521 CATCCCGGCGCCGTTCCAGCCGCTCTACGCCTCCAGCGGCCAGGCGGTCTTGGACGGTTC
443 I P A P F Q P L Y A S S G Q A V L D G S

2581 CCGGTCGGAÇTGCGCGCTCGGGCGCCGACCCCGACGCGCGGTCCCGGGCATTCTCGGCCC
463 R S D C A L G A D P D A A V P G I L G P

2641 GGCGAGCCTGCCGGTGCGGTTCGCGGACGACTACGTCACCAACTCCAACGACAGTCACTG
483 A S L P V R F R D D Y V T N S N D S H W

2701 GCTGGCCAGCCCGGCCGCCCCGCTGGAAGGCTTCCCGCGGATCCTCGGCAACGAACGCAC
503 L A S P A A P L E G F P R I L G N E R T

2761 CCCGCGCAGCCTGCGCACC CGGCTCGGGCTGGACCAGATCCAGCAGCGCCTCGCCGGCAC
523 P R S L R T R L G L D Q I Q Q R L A G T

2821 GGACGGTCTGCCCGGCAAGGGCTTCACCACCGCCCGGCTCTGGCAGGTTCATGTTCCGGCAA
543 D G L P G K G F T T A R L W Q V M F G N

2881 CCGGATGCACGGCGCCGAACCTCGCCCGGACGACCTGGTTCGCGCTCTGCCCGCCGCCAGCC
563 R M H G A E L A R D D L V A L C R R Q P

FIG. 13C

2941 GACCGCGACCGCCTCGAACGGCGCGATCGTCGACCTCACCGCGGCCTGCACGGCGCTGTC
583 T A T A S N G A I V D L T A A C T A L S

3001 CCGCTTCGATGAGCGTGCCGACCTGGACAGCCGGGGCGCGCACCTGTTTCACCGAGTTCGC
603 R F D E R A D L D S R G A H L F T E F A

3061 CCTCGCGGGCGGAATCAGGTTCGCCGACACCTTCGAGGTGACCGATCCGGTACGCACCCC
623 L A G G I R F A D T F E V T D P V R T P

3121 GCGCCGTCTGAACACCACGGATCCGCGGGTACGGACGGCGCTCGCCGACGCCGTGCAACG
643 R R L N T T D P R V R T A L A D A V Q R

3181 GCTCGCCGGCATCCCCCTCGACGCGAAGCTGGGAGACATCCACACCGACAGCCGCGGCGA
663 L A G I P L D A K L G D I H T D S R G E

3241 ACGGCGCATCCCCATCCACGGTGGCCGCGGGGAAGCAGGCACCTTCAACGTGATCACCAA
683 R R I P I H G G R G E A G T F N V I T N

3301 CCCGCTCGTGCCGGGCGTGGGATACCCGACGGTCCGTCACGGAACATCGTTCGTGATGGC
703 P L V P G V G Y P Q V V H G T S F V M A

3361 CGTCGAACTCGGCCCCGACGGCCCCGTCGGGACGGCAGATCCTCACCTATGCGCAGTCGAC
723 V E L G P H G P S G R Q I L T Y A Q S T

3421 GAACCCGAACTCACCTGGTACGCCGACCAGACCGTGCTCTACTCGCGGAAGGGCTGGGA
743 N P N S P W Y A D Q T V L Y S R K G W D

3481 CACCATCAAGTACACCGAGGCGCAGATCGCGGCCGACCCGAACCTGCGCGTCTACCGGGT
763 T I K Y T E A Q I A A D P N L R V Y R V

3541 GGCACAGCGGGGACGCTGACCCACGTCACGCCGGCTCGGCCCGTGCGGGGGCGCAGGGCG
783 A Q R G R *

FIG. 13D

3601 CCGATCGTCTCTGCATCGCCGGTCAGCCGGGGCCTGCGTCGACCGGCGGCGGCCGGTTCGA
3661 CGCCCGCGTCCCGGCGCAGCGACTGGCTGAAGCGCCAGGCGTCGGCGGCCCGGGCAGGT
3721 TGTTGAACATCACGTACGCCGGGCCGCGTTCGAGGATGCCGGCGAGGTGTGCCAGCTCGG
3781 CATCCGTGTACACATGCCGGGCGCCGGTGATGCCGTGCAGCCGGTAATAGGCCATCGGCC
3841 TCAGACTGCGGCGCAGGAACGGGTCGGCGGCGTGGGTCAGGTCCAGCTCCTGGCACAAAGC
3901 CCTCGACCACCTCGTCCGGCCACGGGCCGCGCGGCTCCCACAACAGCCGGACACCGGCCG
3961 GCCGGCGCGCTCGGGCGCAGAACTCACGCAGTCGCGCGATGGCGGGTTCGGTTCGGCCGGA
4021 AACTCGCCGGGCACTGCAG

FIG. 13E




FIG. 14




FIG. 15




FIG. 16

1

**METHOD FOR CREATING
POLYNUCLEOTIDE AND POLYPEPTIDE
SEQUENCES**

CROSS-REFERENCES TO RELATED
APPLICATIONS

The present application is a continuation of U.S. Ser. No. 10/371,168 filed Feb. 19, 2003, which is a continuation of U.S. Ser. No. 09/205,448 filed Dec. 4, 1998, which is a non-provisional of 60/607,908 filed Dec. 8, 1997, all of which are incorporated by reference for all purposes.

This application derives priority from U.S. Ser. No. 60/067,908, filed Dec. 8, 1997, which is incorporated by reference in its entirety for all purposes.

TECHNICAL FIELD

The invention resides in the technical field of genetics, and more specifically, forced molecular evolution of polynucleotides to acquire desired properties.

BACKGROUND

A variety of approaches, including rational design and directed evolution, have been used to optimize protein functions (1, 2). The choice of approach for a given optimization problem depends, in part, on the degree of understanding of the relationships between sequence, structure and function. Rational redesign typically requires extensive knowledge of a structure-function relationship. Directed evolution requires little or no specific knowledge about structure-function relationship; rather, the essential features is a means to evaluate the function to be optimized. Directed evolution involves the generation of libraries of mutant molecules followed by selection or screening for the desired function. Gene products which show improvement with respect to the desired property or set of properties are identified by selection or screening. The gene(s) encoding those products can be subjected to further cycles of the process in order to accumulate beneficial mutations. This evolution can involve few or many generations, depending on how far one wishes to progress and the effects of mutations typically observed in each generation. Such approaches have been used to create novel functional nucleic acids (3, 4), peptides and other small molecules (3), antibodies (3), as well as enzymes and other proteins (5, 6, 7). These procedures are fairly tolerant to inaccuracies and noise in the function evaluation (7).

Several publications have discussed the role of gene recombination in directed evolution (see WO 97/07205, WO 98/42727, U.S. Pat. Nos. 5,807,723, 5,721,367, 5,776,744 and WO 98/41645 U.S. Pat. No. 5,811,238, WO 98/41622, WO 98/41623, and U.S. Pat. No. 5,093,257).

A PCR-based group of recombination methods consists of DNA shuffling [5,6], staggered extension process [89, 90] and random-priming recombination [87]. Such methods typically involve synthesis of significant amounts of DNA during assembly/recombination step and subsequent amplification of the final products and the efficiency of amplification decreases with gene size increase.

Yeast cells, which possess an active system for homologous recombination, have been used for in vivo recombination. Cells transformed with a vector and partially overlapping inserts efficiently join the inserts together in the regions of homology and restore a functional, covalently-closed plasmid [91]. This method does not require PCR amplification at any stage of recombination and therefore is free from the size

2

considerations inherent in this method. However, the number of crossovers introduced in one recombination event is limited by the efficiency of transformation of one cell with multiple inserts. Other in vivo recombination methods entail recombination between two parental genes cloned on the same plasmid in a tandem orientation. One method relies on homologous recombination machinery of bacterial cells to produce chimeric genes [92]. A first gene in the tandem provides the N-terminal part of the target protein, and a second provides the C-terminal part. However, only one crossover can be generated by this approach. Another in vivo recombination method uses the same tandem organization of substrates in a vector [93]. Before transformation into *E. coli* cells, plasmids are linearized by endonuclease digestion between the parental sequences. Recombination is performed in vivo by the enzymes responsible for double-strand break repair. The ends of linear molecules are degraded by a 5'->3' exonuclease activity, followed by annealing of complementary single-strand 3' ends and restoration of the double-strand plasmid [94]. This method has similar advantages and disadvantages of tandem recombination on circular plasmid.

SUMMARY OF THE INVENTION

The invention provides methods for evolving a polynucleotide toward acquisition of a desired property. Such methods entail incubating a population of parental polynucleotide variants under conditions to generate annealed polynucleotides comprises heteroduplexes. The heteroduplexes are then exposed to a cellular DNA repair system to convert the heteroduplexes to parental polynucleotide variants or recombined polynucleotide variants. The resulting polynucleotides are then screened or selected for the desired property.

In some methods, the heteroduplexes are exposed to a DNA repair system in vitro. A suitable repair system can be prepared in the form of cellular extracts.

In other methods, the products of annealing including heteroduplexes are introduced into host cells. The heteroduplexes are thus exposed to the host cells' DNA repair system in vivo.

In several methods, the introduction of annealed products into host cells selects for heteroduplexes relative to transformed cells comprising homoduplexes. Such can be achieved, for example, by providing a first polynucleotide variant as a component of a first vector, and a second polynucleotide variant is provided as a component of a second vector. The first and second vectors are converted to linearized forms in which the first and second polynucleotide variants occur at opposite ends. In the incubating step, single-stranded forms of the first linearized vector reanneal with each other to form linear first vector, single-stranded forms of the second linearized vector reanneal with each other to form linear second vector, and single-stranded linearized forms of the first and second vectors anneal with each to form a circular heteroduplex bearing a nick in each strand. Introduction of the products into cells thus selects for circular heteroduplexes relative to the linear first and second vector. Optionally, in the above methods, the first and second vectors can be converted to linearized forms by PCR. Alternatively, the first and second vectors can be converted to linearized forms by digestion with first and second restriction enzymes.

In some methods, polynucleotide variants are provided in double stranded form and are converted to single stranded form before the annealing step. Optionally, such conversion is by conducting asymmetric amplification of the first and second double stranded polynucleotide variants to amplify a first strand of the first polynucleotide variant, and a second strand

of the second polynucleotide variant. The first and second strands anneal in the incubating step to form a heteroduplex.

In some methods, a population of polynucleotides comprising first and second polynucleotides is provided in double stranded form, and the method further comprises incorporating the first and second polynucleotides as components of first and second vectors, whereby the first and second polynucleotides occupy opposite ends of the first and second vectors. In the incubating step single-stranded forms of the first linearized vector reanneal with each other to form linear first vector, single-stranded forms of the second linearized vector reanneal with each other to form linear second vector, and single-stranded linearized forms of the first and second vectors anneal with each to form a circular heteroduplex bearing a nick in each strand. In the introducing step selects for transformed cells comprises the circular heteroduplexes relative to the linear first and second vector.

In some methods, the first and second polynucleotides are obtained from chromosomal DNA. In some methods, the polynucleotide variants encode variants of a polypeptide. In some methods, the population of polynucleotide variants comprises at least 20 variants. In some methods, the population of polynucleotide variants are at least 10 kb in length.

In some methods, the polynucleotide variants comprises natural variants. In other methods, the polynucleotide variants comprise variants generated by mutagenic PCR or cassette mutagenesis. In some methods, the host cells into which heteroduplexes are introduced are bacterial cells. In some methods, the population of variant polynucleotide variants comprises at least 5 polynucleotides having at least 90% sequence identity with one another.

Some methods further comprise a step of at least partially demethylating variant polynucleotides. Demethylation can be achieved by PCR amplification or by passaging variants through methylation-deficient host cells.

Some methods include a further step of sealing one or more nicks in heteroduplex molecules before exposing the heteroduplexes to a DNA repair system. Nicks can be sealed by treatment with DNA ligase.

Some methods further comprise a step of isolating a screened recombinant polynucleotide variant. In some methods, the polynucleotide variant is screened to produce a recombinant protein or a secondary metabolite whose production is catalyzed thereby.

In some methods, the recombinant protein or secondary metabolite is formulated with a carrier to form a pharmaceutical composition.

In some methods, the polynucleotide variants encode enzymes selected from the group consisting of proteases, lipases, amylases, cutinases, cellulases, amylases, oxidases, peroxidases and phytases. In other methods, the polynucleotide variants encode a polypeptide selected from the group consisting of insulin, ACTH, glucagon, somatostatin, somatotropin, thymosin, parathyroid hormone, pigmentary hormones, somatomedin, erythropoietin, luteinizing hormone, chorionic gonadotropin, hyperthymic releasing factors, antidiuretic hormones, thyroid stimulating hormone, relaxin, interferon, thrombopoietic (TPO), and prolactin.

In some methods, each polynucleotide in the population of variant polynucleotides encodes a plurality of enzymes forming a metabolic pathway.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 illustrates the process of heteroduplex formation using polymerase chain reaction (PCR) with one set of primers for each different sequence to amplify the target sequence and vector.

FIG. 2 illustrates the process of heteroduplex formation using restriction enzymes to linearize the target sequences and vector.

FIG. 3 illustrates a process of heteroduplex formation using asymmetric or single primer polymerase chain reaction (PCR) with one set of primers for each different sequence to amplify the target sequence and vector.

FIG. 4 illustrates heteroduplex recombination using unique restriction enzymes (X and Y) to remove the homoduplexes.

FIG. 5 shows the amino acid sequences of the FlaA from *R. lupini* (SEQ ID NO: 1) and *R. meliloti* (SEQ ID NO:2).

FIGS. 6A and 6B show the locations of the unique restriction sites utilized to linearize pRL20 and pRM40.

FIGS. 7A, B, C and D show the DNA sequences of four mosaic flaA genes created by in vitro heteroduplex formation followed by in vivo repair ((a) is SEQ ID NO:3, (b) is SEQ ID NO:4, (c) is SEQ ID NO:5 and (d) is SEQ ID NO:6).

FIG. 8 illustrates how the heteroduplex repair process created mosaic flaA genes containing sequence information from both parent genes.

FIG. 9 shows the physical maps of *Actinoplanes utahensis* ECB deacylase mutants with enhanced specific activity ((a) is pM7-2 for Mutant 7-2, and (b) is pM16 for Mutant 16).

FIG. 10 illustrates the process used for Example 2 to recombine mutations in Mutant 7-2 and Mutant 16 to yield ECB deacylase recombinant with more enhanced specific activity.

FIG. 11 shows specific activities of wild-type ECB deacylase and improved mutants Mutant 7-2, Mutant 16 and recombinant Mutant 15.

FIG. 12 shows positions of DNA base changes and amino acid substitutions in recombinant ECB deacylase Mutant 15 with respect to parental sequences of Mutant 7-2 and Mutant 16.

FIGS. 13 A, B, C, D and E show the DNA sequence of *A. utahensis* ECB deacylase gene mutant M-15 genes created by in vitro heteroduplex formation followed by in vivo repair (SEQ ID NO:7).

FIG. 14 illustrates the process used for Example 3 to recombine mutations in RC1 and RC2 to yield thermostable subtilisin E.

FIG. 15 illustrates the sequences of RC1 and RC2 and the ten clones picked randomly from the transformants of the reaction products of duplex formation as described in Example 3. The x's correspond to base positions that differ between RC1 and RC2. The mutation at 995 corresponds to amino acid substitution at 181, while that at 1107 corresponds to an amino acid substitution at 218 in the subtilisin protein sequence.

FIG. 16 shows the results of screening 400 clones from the library created by heteroduplex formation and repair for initial activity (A_i) and residual activity (A_r). The ratio A_i/A_r was used to estimate the enzymes' thermostability. Data from active variants are sorted and plotted in descending order.

Approximately 12.9% of the clones exhibit a phenotype corresponding to the double mutant containing both the N181D and the N218S mutations.

DEFINITIONS

Screening is, in general, a two-step process in which one first physically separates the cells and then determines which cells do and do not possess a desired property. Selection is a form of screening in which identification and physical separation are achieved simultaneously by expression of a selection marker, which, in some genetic circumstances, allows cells expressing the marker to survive while other cells die (or vice versa). Exemplary screening members include luciferase, β galactosidase and green fluorescent protein. Selection markers include drug and toxin resistance genes. Although spontaneous selection can and does occur in the course of natural evolution, in the present methods selection is performed by man.

An exogenous DNA segment is one foreign (or heterologous) to the cell or homologous to the cell but in a position within the host cell nucleic acid in which the element is not ordinarily found. Exogenous DNA segments are expressed to yield exogenous polypeptides.

The term gene is used broadly to refer to any segment of DNA associated with a biological function. Thus, genes include coding sequences and/or the regulatory sequences required for their expression. Genes also include non-expressed DNA segments that, for example, form recognition sequences for other proteins.

The term "wild-type" means that the nucleic acid fragment does not comprise any mutations. A "wild-type" protein means that the protein will be active at a level of activity found in nature and typically will comprise the amino acid sequence found in nature. In an aspect, the term "wild type" or "parental sequence" can indicate a starting or reference sequence prior to a manipulation of the invention.

"Substantially pure" means an object species is the predominant species present (i.e., on a molar basis it is more abundant than any other individual macromolecular species in the composition), and preferably a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present. Generally, a substantially pure composition will comprise more than about 80 to 90 percent of all macromolecular species present in the composition. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species. Solvent species, small molecules (<500 Daltons), and elemental ion species are not considered macromolecular species.

Percentage sequence identity is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison. Optimal alignment of sequences for aligning a comparison window can be conducted by computerized implementations of algorithms GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, Wis.

The term naturally-occurring is used to describe an object that can be found in nature as distinct from being artificially

produced by man. For example, a polypeptide or polynucleotide sequence that is present in an organism (including viruses) that can be isolated from a source in nature and which has not been intentionally modified by man in the laboratory is naturally-occurring. Generally, the term naturally-occurring refers to an object as present in a non-pathological (undiseased) individual, such as would be typical for the species.

A nucleic acid is operably linked when it is placed into a functional relationship with another nucleic acid sequence. For instance, a promoter or enhancer is operably linked to a coding sequence if it increases the transcription of the coding sequence. Operably linked means that the DNA sequences being linked are typically contiguous and, where necessary to join two protein coding regions, contiguous and in reading frame. However, since enhancers generally function when separated from the promoter by several kilobases and intronic sequences may be of variable lengths, some polynucleotide elements may be operably linked but not contiguous.

A specific binding affinity between, for example, a ligand and a receptor, means a binding affinity of at least $1 \times 10^6 \text{ M}^{-1}$.

The term "cognate" as used herein refers to a gene sequence that is evolutionarily and functionally related between species. For example but not limitation, in the human genome, the human CD4 gene is the cognate gene to the mouse CD4 gene, since the sequences and structures of these two genes indicate that they are highly homologous and both genes encode a protein which functions in signaling T cell activation through MHC class II-restricted antigen recognition.

The term "heteroduplex" refers to hybrid DNA generated by base pairing between complementary single strands derived from the different parental duplex molecules, whereas the term "homoduplex" refers to double-stranded DNA generated by base pairing between complementary single strands derived from the same parental duplex molecules.

The term "nick" in duplex DNA refers to the absence of a phosphodiester bond between two adjacent nucleotides on one strand. The term "gap" in duplex DNA refers to an absence of one or more nucleotides in one strand of the duplex. The term "loop" in duplex DNA refers to one or more unpaired nucleotides in one strand.

A mutant or variant sequence is a sequence showing substantial variation from a wild type or reference sequence that differs from the wild type or reference sequence at one or more positions.

DETAILED DESCRIPTION

I. General

The invention provides methods of evolving a polynucleotide toward acquisition of a desired property. The substrates for the method are a population of at least two polynucleotide variant sequences that contain regions of similarity with each other but, which also have point(s) or regions of divergence. The substrates are annealed in vitro at the regions of similarity. Annealing can regenerate initial substrates or can form heteroduplexes, in which component strands originate from different parents. The products of annealing are exposed to enzymes of a DNA repair, and optionally a replication system, that repairs unmatched pairings. Exposure can be in vivo as when annealed products are transformed into host cells and exposed to the hosts DNA repair system. Alternatively, exposure can be in vitro, as when annealed products are exposed to cellular extracts containing functional DNA repair systems. Exposure of heteroduplexes to a DNA repair system results in

DNA repair at bulges in the heteroduplexes due to DNA mismatching. The repair process differs from homologous recombination in promoting nonreciprocal exchange of diversity between strands. The DNA repair process is typically effected on both component strands of a heteroduplex molecule and at any particular mismatch is typically random as to which strand is repaired. The resulting population can thus contain recombinant polynucleotides encompassing an essentially random reassortment of points of divergence between parental strands. The population of recombinant polynucleotides is then screened for acquisition of a desired property. The property can be a property of the polynucleotide per se, such as capacity of a DNA molecule to bind to a protein or can be a property of an expression product thereof, such as mRNA or a protein.

II. Substrates for Shuffling

The substrates for shuffling are variants of a reference polynucleotide that show some region(s) of similarity with the reference and other region(s) or point(s) of divergence. Regions of similarity should be sufficient to support annealing of polynucleotides such that stable heteroduplexes can be formed. Variants forms often show substantial sequence identity with each other (e.g., at least 50%, 75%, 90% or 99%). There should be at least sufficient diversity between substrates that recombination can generate more diverse products than there are starting materials. Thus, there must be at least two substrates differing in at least two positions. The degree of diversity depends on the length of the substrate being recombined and the extent of the functional change to be evolved. Diversity at between 0.1-25% of positions is typical. Recombination of mutations from very closely related genes or even whole sections of sequences from more distantly related genes or sets of genes can enhance the rate of evolution and the acquisition of desirable new properties. Recombination to create chimeric or mosaic genes can be useful in order to combine desirable features of two or more parents into a single gene or set of genes, or to create novel functional features not found in the parents. The number of different substrates to be combined can vary widely in size from two to 10, 100, 1000, to more than 10^5 , 10^7 , or 10^9 members.

The initial small population of the specific nucleic acid sequences having mutations may be created by a number of different methods. Mutations may be created by error-prone PCR. Error-prone PCR uses low-fidelity polymerization conditions to introduce a low level of point mutations randomly over a long sequence. Alternatively, mutations can be introduced into the template polynucleotide by oligonucleotide-directed mutagenesis. In oligonucleotide-directed mutagenesis, a short sequence of the polynucleotide is removed from the polynucleotide using restriction enzyme digestion and is replaced with a synthetic polynucleotide in which various bases have been altered from the original sequence. The polynucleotide sequence can also be altered by chemical mutagenesis. Chemical mutagens include, for example, sodium bisulfite, nitrous acid, hydroxylamine, hydrazine or formic acid. Other agents which are analogues of nucleotide precursors include nitrosoguanidine, 5-bromouracil, 2-aminopurine, or acridine. Generally, these agents are added to the PCR reaction in place of the nucleotide precursor thereby mutating the sequence. Intercalating agents such as proflavine, acriflavine, quinacrine and the like can also be used. Random mutagenesis of the polynucleotide sequence can also be achieved by irradiation with X-rays or ultraviolet light. Generally, plasmid DNA or DNA fragments so mutagenized are introduced into *E. coli* and propagated as a pool or library of mutant plasmids.

Alternatively the small mixed population of specific nucleic acids can be found in nature in the form of different alleles of the same gene or the same gene from different related species (i.e., cognate genes). Alternatively, substrates can be related but nonallelic genes, such as the immunoglobulin genes. Diversity can also be the result of previous recombination or shuffling. Diversity can also result from resynthesizing genes encoding natural proteins with alternative codon usage.

The starting substrates encode variant forms of sequences to be evolved. In some methods, the substrates encode variant forms of a protein for which evolution of a new or modified property is desired. In other methods, the substrates can encode variant forms of a plurality of genes constituting a multigene pathway. In such methods, variation can occur in one or any number of the component genes. In other methods, substrates can contain variants segments to be evolved as DNA or RNA binding sequences. In methods, in which starting substrates containing coding sequences, any essential regulatory sequences, such as a promoter and polyadenylation sequence, required for expression may also be present as a component of the substrate. Alternatively, such regulatory sequences can be provided as components of vectors used for cloning the substrates.

The starting substrates can vary in length from about 50, 250, 1000, 10,000, 100,000, 10^6 or more bases. The starting substrates can be provided in double- or single-stranded form. The starting substrates can be DNA or RNA and analogs thereof. If DNA, the starting substrates can be genomic or cDNA. If the substrates are RNA, the substrates are typically reverse-transcribed to cDNA before heteroduplex formation. Substrates can be provided as cloned fragments, chemically synthesized fragments or PCR amplification products. Substrates can derive from chromosomal, plasmid or viral sources. In some methods, substrates are provided in concatenated form.

III. Procedures for Generating Heteroduplexes

Heteroduplexes are generated from double stranded DNA substrates, by denaturing the DNA substrates and incubating under annealing conditions. Hybridization conditions for heteroduplex formation are sequence-dependent and are different in different circumstances. Longer sequences hybridize specifically at higher temperatures. Generally, hybridization conditions are selected to be about 25° C. lower than the thermal melting point (T_m) for the specific sequence at a defined ionic strength and pH. The T_m is the temperature (under defined ionic strength, pH, and nucleic acid concentration) at which 50% of the probes complementary to the target sequence hybridize to the target sequence at equilibrium.

Exemplary conditions for denaturation and renaturation of double stranded substrates are as follows. Equimolar concentrations (~1.0-5.0 nM) of the substrates are mixed in 1×SSPE buffer (180 mM NaCl, 1.0 mM EDTA, 10 mM NaH_2PO_4 , pH 7.4). After heating at 96° C. for 10 minutes, the reaction mixture is immediately cooled at 0° C. for 5 minutes; The mixture is then incubated at 68° C. for 2-6 hr. Denaturation and reannealing can also be carried out by the addition and removal of a denaturant such as NaOH. The process is the same for single stranded DNA substrates, except that the denaturing step may be omitted for short sequences.

By appropriate design of substrates for heteroduplex formation, it is possible to achieve selection for heteroduplexes relative to reformed parental homoduplexes. Homoduplexes merely reconstruct parental substrates and effectively dilute recombinant products in subsequent screening steps. In gen-

eral, selection is achieved by designing substrates such that heteroduplexes are formed in open-circles, whereas homoduplexes are formed as linear molecules. A subsequent transformation step results in substantial enrichment (e.g., 100-fold) for the circular heteroduplexes.

FIG. 1 shows a method in which two substrate sequences in separate vectors are PCR-amplified using two different sets of primers (P1, P2 and P3, P4). Typically, first and second substrates are inserted into separate copies of the same vector. The two different pairs of primers initiate amplification at different points on the two vectors. FIG. 1 shows an arrangement in which the P1/P2 primer pairs initiates amplification at one of the two boundaries of the vector with the substrate and the P1/P2 primer pair initiates replication at the other boundary in a second vector. The two primers in each primer pair prime amplification in opposite directions around a circular plasmid. The amplification products generated by this amplification are double-stranded linearized vector molecules in which the first and second substrates occur at opposite ends of the vector. The amplification products are mixed, denatured and annealed. Mixing and denaturation can be performed in either order. Reannealing generates two linear homoduplexes, and an open circular heteroduplex containing one nick in each strand, at the initiation point of PCR amplification. Introduction of the amplification products into host cells selects for the heteroduplexes relative to the homoduplexes because the former transform much more efficiently than the latter.

It is not essential in the above scheme that amplification is initiated at the interface between substrate and the rest of the vector. Rather, amplification can be initiated at any points on two vectors bearing substrates provided that the amplification is initiated at different points between the vectors. In the general case, such amplification generates two linearized vectors in which the first and second substrates respectively occupy different positions relative to the remainder of the vector. Denaturation and reannealing generator heteroduplexes similar to that shown in FIG. 1, except that the nicks occur within the vector component rather than at the interface between plasmid and substrate. Initiation of amplification outside the substrate component of a vector has the advantage that it is not necessary to design primers specific for the substrate borne by the vector.

Although FIG. 1 is exemplified for two substrates, the above scheme can be extended to any number of substrates. For example, an initial population of vector bearing substrates can be divided into two pools. One pool is PCR-amplified from one set of primers, and the other pool from another. The amplification products are denatured and annealed as before. Heteroduplexes can form containing one strand from any substrate in the first pool and one strand from any substrate in the second pool. Alternatively, three or more substrates cloned into multiple copies of a vector can be subjected to amplification with amplification in each vector starting at a different point. For each substrate, this process generates amplification products varying in how flanking vector DNA is divided on the two sides of the substrate. For example, one amplification product might have most of the vector on one side of the substrate, another amplification product might have most of the vector on the other side of the substrate, and a further amplification product might have an equal division of vector sequence flanking the substrate. In the subsequent annealing step, a strand of substrate can form a circular heteroduplex with a strand of any other substrate, but strands of the same substrate can only reanneal with each other to form a linear homoduplex. In a still further variation, multiple substrates can be performed by performing multiple

iterations of the scheme in FIG. 1. After the first iteration, recombinant polynucleotides in a vector, undergo heteroduplex formation with a third substrate incorporated into a further copy of the vector. The vector bearing the recombinant polynucleotides and the vector bearing the third substrate are separately PCR amplified from different primer pairs. The amplification products are then denatured and annealed. The process can be repeated further times to allow recombination with further substrates.

An alternative scheme for heteroduplex formation is shown in FIG. 2. Here, first and second substrates are incorporated into separate copies of a vector. The two copies are then respectively digested with different restriction enzymes. FIG. 2 shows an arrangement in which, the restriction enzymes cut at opposite boundaries between substrates and vector, but all that is necessary is to use two different restriction enzymes that cut at different places. Digestion generates linearized first and second vector bearing first and second substrates, the first and second substrates occupying different positions relative to the remaining vector sequences. Denaturation and reannealing generates open circular heteroduplexes and linear homoduplexes. The scheme can be extended to recombination between more than two substrates using analogous strategies to those described with respect to FIG. 1. In one variation, two pools of substrates are formed, and each is separately cloned into vector. The two pools are then cut with different enzymes, and annealing proceeds as for two substrates. In another variation, three or more substrates can be cloned into three or more copies of vector, and the three or more result molecules cut with three or more enzymes, cutting at three or more sites. This generates three different linearized vector forms differing in the division of vector sequences flanking the substrate moiety in the vectors. Alternatively, any number of substrates can be recombined pairwise in an iterative fashion with products of one round of recombination annealing with a fresh substrate in each round.

In a further variation, heteroduplexes can be formed from substrates molecules in vector-free form, and the heteroduplexes subsequently cloned into vectors. Such can be achieved by asymmetric amplification of first and second substrates as shown in FIG. 3. Asymmetric or single primer PCR amplifies only one strand of a duplex. By appropriate selection of primers, opposite strands can be amplified from two different substrates. On reannealing amplification products, heteroduplexes are formed from opposite strands of the two substrates. Because only one strand is amplified from each substrate, reannealing does not reform homoduplexes (other than for small quantities of unamplified substrate). The process can be extended to allow recombination of any number of substrates using analogous strategies to those described with respect to FIG. 1. For example, substrates can be divided into two pools, and each pool subject to the same asymmetric amplification, such that amplification products of one pool can only anneal with amplification products of the other pool, and not with each other. Alternatively, shuffling can proceed pairwise in an iterative manner, in which recombinants formed from heteroduplexes of first and second substrates, are subsequently subjected to heteroduplex formation with a third substrate. Point mutations can also be introduced at a desired level during PCR amplification.

FIG. 4 shows another approach of selecting for heteroduplexes relative to homoduplexes. First and second substrates are isolated by PCR amplification from separate vectors. The substrates are denatured and allowed to anneal forming both heteroduplexes and reconstructed homoduplexes. The products of annealing are digested with restriction enzymes X and Y. X has a site in the first substrate but not the second sub-

strate, and vice versa for Y. Enzyme X cuts reconstructed homoduplex from the first substrate and enzyme Y cuts reconstructed homoduplex from the second substrate. Neither enzyme cuts heteroduplexes. Heteroduplexes can effectively be separated from restriction fragments of homoduplexes by further cleavage with enzymes A and B having sites proximate to the ends of both the first and second substrates, and ligation of the products into vector having cohesive ends compatible with ends resulting from digestion with A and B. Only heteroduplexes cut with A and B can ligate with the vector. Alternatively, heteroduplexes can be separated from restriction fragments of homoduplexes by size selection on gels. The above process can be generalized to N substrates by cleaving the mixture of heteroduplexes and homoduplexes with N enzymes, each one of which cuts a different substrate and no other substrate. Heteroduplexes can be formed by directional cloning. Two substrates for heteroduplex formation can be obtained by PCR amplification of chromosomal DNA and joined to opposite ends of a linear vector. Directional cloning can be achieved by digesting the vector with two different enzymes, and digesting or adapting first and second substrates to be respectively compatible with cohesive ends of only of the two enzymes used to cut the vector. The first and second substrates can thus be ligated at opposite ends of a linearized vector fragment. This scheme can be extended to any number of substrates by using principles analogous to those described for FIG. 1. For example, substrates can be divided into two pools before ligation to the vector. Alternatively, recombinant products formed by heteroduplex formation of first and second substrates, can subsequently undergo heteroduplex formation with a third substrate.

IV. Vectors and Transformation

In general, substrates are incorporated into vectors either before or after the heteroduplex formation step. A variety of cloning vectors typically used in genetic engineering are suitable.

The vectors containing the DNA segments of interest can be transferred into the host cell by standard methods, depending on the type of cellular host. For example, calcium chloride transformation is commonly utilized for prokaryotic cells, whereas calcium phosphate treatment, lipofection, or electroporation may be used for other cellular hosts. Other methods used to transform mammalian cells include the use of Polybrene, protoplast fusion, liposomes, electroporation, and microinjection, and biolistics (see, generally, Sambrook et al., supra). Viral vectors can also be packaged in vitro and introduced by infection. The choice of vector depends on the host cells. In general, a suitable vector has an origin of replication recognized in the desired host cell, a selection marker capable of being expressed in the intended host cells and/or regulatory sequences to support expression of genes within substrates being shuffled.

V. Types of Host Cells

In general any type of cells supporting DNA repair and replication of heteroduplexes introduced into the cells can be used. Cells of particular interest are the standard cell types commonly used in genetic engineering, such as bacteria, particularly, *E. coli* (16, 17). Suitable *E. coli* strains include *E. coli* mutS, mutL, dam⁻, and/or recA⁺, *E. coli* XL-10-Gold ([Tet^rΔ(mcrA)183 Δ(mcrCB-hsdSMR-mrr)173 endA1 supE44 thi-1 recA1 gyrA96 relA1 lac Hte] [F' proAB lacI^gZAM15 Tn10 (Tet^r) Amy Cam^r]), *E. coli* ES1301 mutS [Genotype: lacZ53, mutS201::Tn5, thyA36, rha-5, metBI, deoC, IN(rrnD-rrnE)] (20, 24, 28-42). Preferred *E. coli* strains are *E. coli* SCS110 [Genotype: rpsI, (Str^r), thr, leu, endA, thi-1, lacy, galk gait, ara tona, tsx, dam, dcm, supE44,

Δ(lac-proAB), [F, traD36, proA⁺B⁺IacI^gZAM15], which have normal cellular mismatch repair systems (17). This strain type repairs mismatches and unmatches in the heteroduplex with little strand-specific preference. Further, because this strain is dam⁻ and dcm⁻, plasmid isolated from the strain is unmethylated and therefore particularly amenable for further rounds of DNA duplex formation/mismatch repair (see below). Other suitable bacterial cells include gram-negative and gram-positive, such as *Bacillus*, *Pseudomonas*, and *Salmonella*.

Eukaryotic organisms are also able to carry out mismatch repair (43-48). Mismatch repair systems in both prokaryotes and eukaryotes are thought to play an important role in the maintenance of genetic fidelity during DNA replication. Some of the genes that play important roles in mismatch repair in prokaryotes, particularly mutS and mutL, have homologs in eukaryotes. In the outcome of genetic recombinations, and in genome stability. Wild-type or mutant *S. cerevisiae* has been shown to carry out mismatch repair of heteroduplexes (49-56), as have COS-1 monkey cells (57). Preferred strains of yeast are *Picchia* and *Saccharomyces*. Mammalian cells have been shown to have the capacity to repair G-T to G-C base pairs by a short-patch mechanism (38, 58-63). Mammalian cells (e.g., mouse, hamster, primate, human), both cell lines and primary cultures can also be used. Such cells include stem cells, including embryonic stem cells, zygotes, fibroblasts, lymphocytes, Chinese hamster ovary (CHO), mouse fibroblasts (NIH3T3), kidney, liver, muscle, and skin cells. Other eucaryotic cells of interest include plant cells, such as maize, rice, wheat, cotton, soybean, sugarcane, tobacco, and *arabidopsis*; fish, algae, fungi (*aspergillus*, *podospora*, *neurospora*), insect (e.g., baculo lepidoptera) (see, Winnacker, "From Genes to Clones," VCH Publishers, N.Y., (1987), which is incorporated herein by reference).

In vivo repair occurs in a wide variety of prokaryotic and eukaryotic cells. Use of mammalian cells is advantage in certain application in which substrates encode polypeptides that are expressed only in mammalian cells or which are intended for use in mammalian cells. However, bacterial and yeast cells are advantageous for screening large libraries due to the higher transformation frequencies attainable in these strains.

V. In Vitro DNA Repair Systems

As an alternative to introducing annealed products into host cells, annealed products can be exposed a DNA repair system in vitro. The DNA repair system can be obtained as extracts from repair-competent *E. coli*, yeast or any other cells (64-67). Repair-competent cells are lysed in appropriate buffer and supplemented with nucleotides. DNA is incubated in this cell extract and transformed into competent cells for replication.

VI. Screening and Selection

After introduction of annealed products into host cells, the host cells are typically cultured to allow repair and replication to occur and optionally, for genes encoded by polynucleotides to be expressed. The recombinant polynucleotides can be subject to further rounds of recombination using the heteroduplex procedures described above, or other shuffling methods described below. However, whether after one cycle of recombination or several, recombinant polynucleotides are subjected to screening or selection for a desired property. In some instances, screening or selection is performed in the same host cells that are used for DNA repair. In other instances, recombinant polynucleotides, their expression products or secondary metabolites produced by the expression products are isolated from such cells and screened in

vitro. In other instances, recombinant polynucleotides are isolated from the host cells in which recombination occurs and are screened or selected in other host cells. For example, in some methods, it is advantageous to allow DNA repair to occur in a bacterial host strain, but to screen an expression product of recombinant polynucleotides in eucaryotic cells. The recombinant polynucleotides surviving screening or selection are sometimes useful products in themselves. In other instances, such recombinant polynucleotides are subjected to further recombination with each other or other substrates. Such recombination can be effected by the heteroduplex methods described above or any other shuffling methods. Further round(s) of recombination are followed by further rounds of screening or selection on an iterative basis. Optionally, the stringency of selection can be increased at each round.

The nature of screening or selection depends on the desired property sought to be acquired. Desirable properties of enzymes include high catalytic activity, capacity to confer resistance to drugs, high stability, the ability to accept a wider (or narrower) range of substrates, or the ability to function in nonnatural environments such as organic solvents. Other desirable properties of proteins include capacity to bind a selected target, secretion capacity, capacity to generate an immune response to a given target, lack of immunogenicity and toxicity to pathogenic microorganisms. Desirable properties of DNA or RNA polynucleotides sequences include capacity to specifically bind a given protein target, and capacity to regulate expression of operably linked coding sequences. Some of the above properties, such as drug resistance, can be selected by plating cells on the drug. Other properties, such as the influence of a regulatory sequence on expression, can be screened by detecting appearance of the expression product of a reporter gene linked to the regulatory sequence. Other properties, such as capacity of an expressed protein to be secreted, can be screened by FACSTM, using a labelled antibody to the protein. Other properties, such as immunogenicity or lack thereof, can be screened by isolating protein from individual cells or pools of cells, and analyzing the protein in vitro or in a laboratory animal.

VII. Variations

1. Demethylation

Most cell types methylate DNA in some manner, with the pattern of methylation differing between cells types. Sites of methylation include 5-methylcytosine (m⁵C), N4-methylcytosine (m⁴C) and N⁶-methyladenine (m⁶A), 5-hydroxymethylcytosine (hm⁵C) and 5-hydroxymethyluracil (hm⁵U). In *E. coli*, methylation is effected by Dam and Dcm enzymes. The methylase specified by the dam gene methylates the N6-position of the adenine residue in the sequence GATC, and the methylase specified by the dcm gene methylates the C5-position of the internal cytosine residue in the sequence CCWGG. DNA from plants and mammal is often subject to CG methylation meaning that CG or CNG sequences are methylated. Possible effects of methylated on cellular repair are discussed by references 18-20.

In some methods, DNA substrates for heteroduplex formation are at least partially demethylated on one or both strands, preferably the latter. Demethylation of substrate DNA promotes efficient and random repair of the heteroduplexes. In heteroduplexes formed with one strand dam-methylated and one strand unmethylated, repair is biased to the unmethylated strand, with the methylated strand serving as the template for correction. If neither strand is methylated, mismatch repair occurs, but shows insignificant strand preference (23, 24).

Demethylation can be performed in a variety of ways. In some methods, substrate DNA is demethylated by PCR-amplification. In some instances, DNA demethylation is accomplished in one of the PCR steps in the heteroduplex formation procedures described above. In other methods, an additional PCR step is performed to effect demethylation. In other methods, demethylation is effected by passaging substrate DNA through methylation deficient host cells (e.g. an *E. coli* dam⁻dcm⁻ strain). In other methods, substrate DNA is demethylated in vitro using a demethylating enzymes. Demethylated DNA is used for heteroduplex formation using the same procedures described above. Heteroduplexes are subsequently introduced into DNA-repair-proficient but restriction-enzyme-defective cells to prevent degradation of the unmethylated heteroduplexes.

2. Sealing Nicks

Several of the methods for heteroduplex formation described above result in circular heteroduplexes bearing nicks in each strand. These nicks can be sealed before introducing heteroduplexes into host cells. Sealing can be effected by treatment with DNA ligase under standard ligating conditions. Ligation forms a phosphodiester bond to link two adjacent bases separated by a nick in one strand of double helix of DNA. Sealing of nicks increases the frequency of recombination after introduction of heteroduplexes into host cells.

3. Error Prone PCR Attendant To Amplification

Several of the formats described above include a PCR amplification step. Optionally, such a step can be performed under mutagenic conditions to induce additional diversity between substrates.

VIII. Other Shuffling Methods

The methods of heteroduplex formation described above can be used in conjunction with other shuffling methods. For example, one can perform one cycle of heteroduplex shuffling, screening or selection, followed by a cycle of shuffling by another method, followed by a further cycle of screening or selection. Other shuffling formats are described by WO 95/22625; U.S. Pat. Nos. 5,605,793; 5,811,238; WO 96/19256; Stemmer, *Science* 270, 1510 (1995); Stemmer et al., *Gene*, 164, 49-53 (1995); Stemmer, *Bio/Technology*, 13, 549-553 (1995); Stemmer, *Proc. Natl. Acad. Sci. USA* 91, 10747-10751 (1994); Stemmer, *Nature* 370, 389-391 (1994); Crameri et al., *Nature Medicine*, 2(1):1-3, (1996); Crameri et al., *Nature Biotechnology* 14, 315-319 (1996); WO 98/42727; WO 98/41622; WO 98/05764 and WO 98/42728, WO 98/27230 (each of which is incorporated by reference in its entirety for all purposes).

IX. Protein Analogs

Proteins isolated by the methods also serve as lead compounds for the development of derivative compounds. The derivative compounds can include chemical modifications of amino acids or replace amino acids with chemical structures. The analogs should have a stabilized electronic configuration and molecular conformation that allows key functional groups to be presented in substantially the same way as a lead protein. In particular, the non-peptic compounds have spatial electronic properties which are comparable to the polypeptide binding region, but will typically be much smaller molecules than the polypeptides, frequently having a molecular weight below about 2 CHD and preferably below about 1 CHD. Identification of such non-peptic compounds can be performed through several standard methods such as self-consistent field (CSF) analysis, configuration interaction (CHI) analysis, and normal mode dynamics analysis. Computer programs for implementing these techniques are readily

available. See Rein et al., *Computer-Assisted Modeling of Receptor-Ligand Interactions* (Alan Liss, New York, 1989).

IX. Pharmaceutical Compositions

Polynucleotides, their expression products, and secondary metabolites whose formation is catalyzed by expression products, generated by the above methods are optionally formulated as pharmaceutical compositions. Such a composition comprises one or more active agents, and a pharmaceutically acceptable carrier. A variety of aqueous carriers can be used, e.g., water, buffered water, phosphate-buffered saline (PBS), 0.4% saline, 0.3% glycine, human albumin solution and the like. These solutions are sterile and generally free of particulate matter. The compositions may contain pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions such as pH adjusting and buffering agents, toxicity adjusting agents and the like, for example, sodium acetate, sodium chloride, potassium chloride, calcium chloride and sodium is selected primarily based on fluid volumes, viscosities, and so forth, in accordance with the particular mode of administration selected.

EXAMPLES

Example 1

Novel *Rhizobium* FlaA Genes from Recombination of *Rhizobium Lupini* FlaA and *Rhizobium Meliloti* FlaA

Bacterial flagella have a helical filament, a proximal hook and a basal body with the flagellar motor (68). This basic design has been extensively examined in *E. coli* and *S. typhimurium* and is broadly applicable to many other bacteria as well as some archaea. The long helical filaments are polymers assembled from flagellin subunits, whose molecular weights range between 20,000 and 65,000, depending on the bacterial species (69). Two types of flagellar filaments, named plain and complex, have been distinguished by their electron microscopically determined surface structures (70). Plain filaments have a smooth surface with faint helical lines, whereas complex filaments exhibit a conspicuous helical pattern of alternating ridges and grooves. These characteristics of complex flagellar filaments are considered to be responsible for the brittle and (by implication) rigid structure that enables them to propel bacteria efficiently in viscous media (71-73). Whereas flagella with plain filaments can alternate between clockwise and counter clockwise rotation (68), all known flagella with complex filaments rotate only clockwise with intermittent stops (74). Since this latter navigation pattern is found throughout bacteria and archaea, it has been suggested that complex flagella may reflect the common background of an ancient, basic motility design (69).

Differing from plain bacterial flagella in the fine structure of their filaments dominated by conspicuous helical bands and in their fragility, the filaments are also resistant against heat decomposition (72). Schmitt et al. (75) showed that bacteriophage 7-7-1 specifically adsorbs to the complex flagella of *R. lupini* H13-3 and requires motility for a productive infection of its host. Though the flagellins from *R. meliloti* and *R. lupini* are quite similar, bacteriophage 7-7-1 does not infect *R. meliloti*. Until now complex flagella have been observed in only three species of soil bacteria: *Pseudomonas rhodos* (73), *R. meliloti* (76), and *R. lupini* H13-3 (70, 72). Cells of *R. lupini* H13-3 possess 5 to 10 peritrichously inserted complex flagella, which were first isolated and analyzed by high resolution electron microscopy and by optical diffraction (70).

Maruyama et al. (77) further found that a higher content of hydrophobic amino acid residues in the complex filament may be one of the main reasons for the unusual properties of complex flagella. By measuring mass per unit length and obtaining three-dimensional reconstruction from electron micrographs, Trachtenberg et al. (73, 78) suggested that the complex filaments of *R. lupini* are composed of functional dimers. FIG. 6 shows the comparison between the deduced amino acid sequence of the *R. lupini* H13-3 FlaA and the deduced amino acid sequence of the *R. meliloti* FlaA. Perfect matches are indicated by vertical lines, and conservative exchanges are indicated by colons. The overall identity is 56%. The *R. lupini* flaA and *R. meliloti* flaA were subjected to in vitro heteroduplex formation followed by in vivo repair in order to create novel FlaA molecules and structures.

A. Methods

pRL20 containing *R. lupini* H13-3flaA gene and pRM40 containing *R. meliloti* flaA gene are shown in FIGS. 6A and 6B. These plasmids were isolated from *E. coli* SCS110 (free from dam- and dcm-type methylation). About 3.0 pg. of unmethylated pRL20 and pRM40 DNA were digested with Bam HI and Eco RI, respectively, at 37° C. for 1 hour. After agarose gel separation, the linearized DNA was purified with Wizard PCR Prep kit (Promega, Wis., USA). Equimolar concentrations (2.5 nM) of the linearized unmethylated pRL20 and pRM40 were mixed in 1×SSPE buffer (180 mM NaCl, 1 mM EDTA, 10 mM NaH₂PO₄, pH 7.4). After heating at 96° C. for 10 minutes, the reaction mixture was immediately cooled at 0° C. for 5 minutes. The mixture was incubated at 68° C. for 2 hour for heteroduplexes to form.

One microliter of the reaction mixture was used to transform 50111 of *E. coli* ES 1301 mutS, *E. coli* SCS110 and *E. coli* JM109 competent cells. The transformation efficiency with *E. coli* JM109 competent cells was about seven times higher than that of *E. coli* SCS110 and ten times higher than that of *E. coli* ES1301 mutS, although the overall transformation efficiencies were 10-200 times lower than those of control transformations with the close, covalent and circular pUC19 plasmid.

Two clones were selected at random from the *E. coli* SCS10 transformants and two from *E. coli* ES1301 mutS transformants, and plasmid DNA was isolated from these four clones for further DNA sequencing analysis.

B. Results

FIG. 7 shows (a) the sequence of SCS01 (clone#1 from *E. coli* SCS110 transformant library), (b) the sequence of SCS02 (clone #2 from *E. coli* SCS110 transformant library), (c) the sequence of ES01 (clone #1 from *E. coli* ES1301 transformant library), and (d) the sequence of ES02 (clone #2 from *E. coli* ES1301 transformant library). All four sequences were different from wild-type *R. lupini* flaA and *R. meliloti* flaA sequences. Clones SCS02, ES01 and ES02 all contain a complete open-reading frame, but SCS01 was truncated. FIG. 8 shows that recombination mainly occurred in the loop regions (unmatched regions). The flaA mutant library generated from *R. meliloti* flaA and *R. lupini* flaA can be transformed into *E. coli* SCS110, ES1301, XL10-Gold and JM109, and transformants screened for functional FlaA recombinants.

Example 2

Directed Evolution of ECB Deacylase for Variants with Enhanced Specific Activity

Streptomyces are among the most important industrial microorganisms due to their ability to produce numerous important secondary metabolites (including many antibiot-

ics) as well as large amounts of enzymes. The approach described here can be used with little modification for directed evolution of native *Streptomyces* enzymes, some or all of the genes in a metabolic pathways, as well as other heterologous enzymes expressed in *Streptomyces*.

New antifungal agents are critically needed by the large and growing numbers of immune-compromised AIDS, organ transplant and cancer chemotherapy patients who suffer opportunistic infections. Echinocandin B (ECB), a lipopeptide produced by some species of *Aspergillus*, has been studied extensively as a potential antifungal. Various antifungal agents with significantly reduced toxicity have been generated by replacing the linoleic acid side chain of *A. nidulans* echinocandin B with different aryl side chains (79-83). The cyclic hexapeptide ECB nucleus precursor for the chemical acylation is obtained by enzymatic hydrolysis of ECB using *Actinoplanes utahensis* ECB deacylase. To maximize the conversion of ECB into intact nucleus, this reaction is carried out at pH 5.5 with a small amount of miscible organic solvent to solubilize the ECB substrate. The product cyclic hexapeptide nucleus is unstable at pH above 5.5 during the long incubation required to fully deacylate ECB (84). The pH optimum of ECB deacylase, however, is 8.0-8.5 and its activity is reduced at pH 5.5 and in the presence of more than 2.5% ethanol (84). To improve production of ECB nucleus it is necessary to increase the activity of the ECB deacylase under these process-relevant conditions.

Relatively little is known about ECB deacylase. The enzyme is a heterodimer whose two subunits are derived by processing of a single precursor protein (83). The 19.9 kD α -subunit is separated from the 60.4 kD β -subunit by a 15-amino acid spacer peptide that is removed along with a signal peptide and another spacer peptide in the native organism. The polypeptide is also expressed and processed into functional enzyme in *Streptomyces lividans*, the organism used for large-scale conversion of ECB by recombinant ECB deacylase. The three-dimensional structure of the enzyme has not been determined, and its sequence shows so little similarity to other possibly related enzymes such as penicillin acylase that a structural model reliable enough to guide a rational effort to engineer the ECB deacylase will be difficult to build. We therefore decided to use directed evolution (85) to improve this important activity.

Protocols suitable for mutagenic PCR and random-priming recombination of the 2.4 kb ECB deacylase gene (73% G+C) have been described recently (86). Here, we further describe the use of heteroduplex recombination to generate new ECB deacylase with enhanced specific activity.

In this case, two *Actinoplanes utahensis* ECB deacylase mutants, M7-2 and M16, which show higher specific activity at pH 5.5 and in the presence of 10% MeOH were recombined using technique of the in vitro heteroduplex formation and in vivo mismatch repair.

FIG. 12 shows the physical maps of plasmids pM7-2 and pM16 which contain the genes for the M7-2 and M16 ECB deacylase mutants. Mutant M7-2 was obtained through mutagenic PCR performed directly on whole *Streptomyces lividans* cells containing wild-type ECB deacylase gene, expressed from plasmid pSHP150-2*. *Streptomyces* with pM7-2 show 1.5 times the specific activity of cells expressing the wild-type ECB deacylase (86). Clone pM16 was obtained using the random-priming recombination technique as described (86, 87). It shows 2.4 times specific activity of the wild-type ECB deacylase clone.

A. Methods:

M7-2 and M16 plasmid DNA (pM7-2 and pM16) (FIG. 9) were purified from *E. coli* SCS110 (in separate reactions).

About 5.0 μ g of unmethylated M7-2 and M16 DNA were digested with Xho I and Psh AI, respectively, at 37° C. for 1 hour (FIG. 10). After agarose gel separation, the linearized DNA was purified using a Wizard PCR Prep Kit (Promega, Wis., USA).

Equimolar concentrations (2.0 nM) of the linearized unmethylated pM7-2 and pM16 DNA were mixed in 1 \times SSPE buffer (1 \times SSPE: 180 mM NaCl, 1.0 mM EDTA, 10 mM NaH₂PO₄, pH 7.4). After heating at 96° C. for 10 minutes, the reaction mixture is immediately cooled at 0° C. for 5 minutes. The mixture was incubated at 68° C. for 3 hours to promote formation of heteroduplexes.

One microliter of the reaction mixture was used to transform 50 μ l of *E. coli* ES1301 mutS, SCS110 and JM109 competent cells. All transformants from *E. coli* ES1301 mutS were pooled and *E. coli* SCS110 were pooled. A plasmid pool was isolated from each pooled library, and this pool was used to transform *S. lividans* TK23 protoplasts to form a mutant library for deacylase activity screening. Transformants from the *S. lividans* TK23 libraries were screened for ECB deacylase activity with an in situ plate assay. Transformed protoplasts were allowed to regenerate on R2YE agar plates for 24 hr at 30° C. and to develop in the presence of thiostrepton for 48 hours. When the colonies grew to the proper size, 6 ml of 0.7% agarose solution containing 0.5 mg/ml ECB in 0.1 M sodium acetate buffer (pH 5.5) was poured on top of each R2YE-agar plate and allowed to develop for 18-24 hr at 30° C. Colonies surrounded by a clearing zone larger than that of a control colony containing wild-type plasmid pSHP150-2*, were selected for further characterization.

Selected transformants were inoculated into 20 ml medium containing thiostrepton and grown aerobically at 30° C. for 48 hours, at which point they were analyzed for ECB deacylase activity using HPLC. 100 μ l of whole broth was used for a reaction at 30° C. for 30 minutes in 0.1 M NaAc buffer (pH 5.5) containing 10% (v/v) MeOH and 200 μ g/ml of ECB substrate. The reactions were stopped by adding 2.5 volumes of methanol, and 20 μ l of each sample were analyzed by HPLC on a 100 \times 4.6 mm polyhydroxyethyl aspartamide column (PolyLC Inc., Columbia, Md., USA) at room temperature using a linear acetonitrile gradient starting with 50:50 of A:B (A=93% acetonitrile, 0.1% phosphoric acid; B=70% acetonitrile, 0.1% phosphoric acid) and ending with 30:70 of A:B in 22 min at a flow rate of 2.2 ml/min. The areas of the ECB and ECB nucleus peaks were calculated and subtracted from the areas of the corresponding peaks from a sample culture of *S. lividans* containing pIJ702* in order to estimate the ECB deacylase activity.

2.0 ml pre-cultures of positive mutants were used to inoculate 50-ml medium and allowed to grow at 30° C. for 96 hr. The supernatants were further concentrated to 1/30 their original volume using an Amicon filtration unit (Beverly, Mass., USA) with molecular weight cutoff of 10 kD. The resulting enzyme samples were diluted with an equal volume of 50 mM KH₂PO₄ (pH 6.0) buffer and were applied to Hi-Trap ion exchange column (Pharmacia Biotech, Piscataway, N.J., USA). The binding buffer was 50 mM KH₂PO₄ (pH 6.0), and the elution buffer was 50 mM KH₂PO₄ (pH 6.0) containing 1.0 M NaCl. A linear gradient from 0 to 1.0 M NaCl was applied in 8 column volumes with a flow rate of 2.7 mL/min. The ECB deacylase fraction eluting at 0.3 M NaCl was concentrated and the buffer was exchanged for 50 mM KH₂PO₄ (pH 6.0) using Centricon-10 units. Enzyme purity was verified by SDS-PAGE using Coomassie Blue stain, and the concentration was determined using the Bio-Rad Protein Assay Reagent (Hercules, Calif., USA).

A modified HPLC assay was used to determine the activities of the ECB deacylase mutants on ECB substrate (84). Four μg of each purified ECB deacylase mutant was used for activity assay reaction at 30° C. for 30 minutes in 0.1 M NaAc buffer (pH 5.5) containing 10% (v/v) MeOH and different concentrations of ECB substrate. Assays were performed in duplicate. The reactions were stopped by adding 2.5 volumes of methanol, and the HPLC assays were carried out as described above. The absorbance values were recorded, and the initial rates were calculated by least-squares regression of the time progress curves from which the K_m and the k_{cat} were calculated.

Activities as a function of pH were measured for the purified ECB deacylases at 30° C. at different pH values: 5, 5.5 and 6 (0.1 M acetate buffer); 7, 7.5, 8 and 8.5 (0.1 M phosphate buffer); 9 and 10 (0.1 M carbonate buffer) using the HPLC assay. Stabilities of purified ECB deacylases were determined at 30° C. in 0.1 M NaAc buffer (pH 5.5) containing 10% methanol. Samples were withdrawn at different time intervals, and the residual activity was measured in the same buffer with the HPLC assay described above.

B. Results

FIG. 11 shows that after one round of applying this heteroduplex repair technique on the mutant M7-2 and M16 genes, one mutant (M15) from about 500 original transformants was found to possess 3.1 times the specific activity of wild-type. Wild type and evolved M15 ECB deacylases were purified and their kinetic parameters for deacylation of ECB were determined by HPLC. The evolved deacylases M15 has an increased catalytic rate constant, k_{cat} , by 205%. The catalytic efficiency (k_{cat}/K_m) of M20 is enhanced by a factor of 2.9 over the wild-type enzyme.

Initial rates of deacylation with the wild type and M15 at different pH values from 5 to 10 were determined at 200 $\mu\text{g}/\text{ml}$ of ECB. The recombined M15 is more active than wild type at pH 5-8. Although the pH dependence of the enzyme activity in this assay is not strong, there is a definite shift of 1.0-1.5 units in the optimum to lower pH, as compared to wild type.

The time courses of deactivation of the purified ECB deacylase mutant M15 was measured in 0.1 M NaAc (pH 5.5) at 30° C. No significant difference in stability was observed between wild type and mutant M15.

The DNA mutations with respect to the wild type ECB deacylase sequence and the positions of the amino acid substitutions in the evolved variants M7-2, M-16 and M15 are summarized in FIG. 12.

The heteroduplex recombination technique can recombine parent sequences to create novel progeny. Recombination of the M7-2 and M16 genes yielded M15, whose activity is higher than any of its parents (Fig. 13). Of the six base substitutions in M15, five (at positions $\alpha 50$, $\alpha 171$, $\beta 57$, $\beta 129$ and $\beta 340$) were inherited from M7-2, and the other one ($\beta 30$) came from M16.

This approach provides an alternative to existing methods of DNA recombination and is particularly useful in recombining large genes or entire operons. This method can be used to create recombinant proteins to improve their properties or to study structure-function relationship.

Example 3

Novel Thermostable *Bacillus Subtilis* Subtilisin E Variants

This example demonstrates the use in vitro heteroduplex formation followed by in vivo repair for combining sequence

information from two different sequences in order to improve the thermostability of *Bacillus subtilis* subtilisin E.

Genes RC1 and RC2 encode thermostable *B. subtilis* subtilisin E variants (88). The mutations at base positions 1107 in RC1 and 995 in RC2 (FIG. 14), giving rise to amino acid substitutions Asn218/Ser (N218S) and Asn181/Asp (N181 ID), lead to improvements in subtilisin E thermostability; the remaining mutations, both synonymous and nonsynonymous, have no detectable effects on thermostability. At 65° C., the single variants N181D and N218S have approximately 3-fold and 2-fold longer half-lives, respectively, than wild subtilisin E, and variants containing both mutations have half-lives that are 8-fold longer (88). The different half-lives in a population of subtilisin E variants can therefore be used to estimate the efficiency by which sequence information is combined. In particular, recombination between these two mutations (in the absence of point mutations affecting thermostability) should generate a library in which 25% of the population exhibits the thermos/ability of the double mutant. Similarly, 25% of the population should exhibit wild-type like stability, as N181D and N218S are eliminated at equal frequency. We used the fractions of the recombined population as a diagnostic

A. Methods

The strategy underlying this example is shown in FIG. 15.

Subtilisin E thermostable mutant genes RC1 and RC2 (FIG. 14) are 986-bp fragments including 45 nt of subtilisin E prosequence, the entire mature sequence and 113 nt after the stop codon. The genes were cloned between Bam HI and Nde I in *E. coli/B. subtilis* shuttle vector pBE3, resulting in pBE3-1 and pBE3-2, respectively. Plasmid DNA pBE3-1 and pBE3-2 was isolated from *E. coli* SCS110.

About 5.0 μg of unmethylated pBE3-1 and pBE3-2 DNA were digested with Bam HI and Nde I, respectively, at 37° C. for 1 hour. After agarose gel separation, equimolar concentrations (2.0 nM) of the linearized unmethylated pBE3-1 and pBE3-2 were mixed in 1 \times SSPE buffer (180 mM NaCl, 1.0 mM EDTA, 10 mM NaH_2PO_4 , pH 7.4). After heating at 96° C. for 10 minutes, the reaction mixture was immediately cooled at 0° C. for 5 min. The mixture was incubated at 68° C. for 2 hr for heteroduplexes to form.

One microliter of the reaction mixture was used to transform 50 μl of *E. coli* ES 1301 mutS, *E. coli* SCS110 and *E. coli* HB101 competent cells.

The transformation efficiency with *E. coli* HB101 competent cells was about ten times higher than that of *E. coli* SCS110 and 15 times higher than that of *E. coli* ES1301 mutS. But in all these cases, the transformation efficiencies were 10-250 times lower than that of the transformation with closed, covalent and circular control pUC19 plasmids.

Five clones from *E. coli* SCS110 mutant library and five from *E. coli* ES1301 mutS library were randomly chosen, and plasmid DNA was isolated using a QIAprep spin plasmid miniprep kit for further DNA sequencing analysis.

About 2,000 random clones from *E. coli* HB101 mutant library were pooled and total plasmid DNA was isolated using a QIAGEN-100 column. 0.5-4.0 μg of the isolated plasmid was used to transform *Bacillus subtilis* DB428 as described previously (88).

About 400 transformants from the *Bacillus subtilis* DB428 library were subjected to screening. Screening was performed using the assay described previously (88), on succinyl-Ala-Ala-Pro-Phe-p-nitroanilide. *B. subtilis* DB428 containing the plasmid library were grown on LB plates containing kanamycin (20 $\mu\text{g}/\text{ml}$) plates. After 18 hours at 37° C. single colonies were picked into 96-well plates containing 200 μl SG/kanamycin medium per well. These plates were incu-

bated with shaking at 37° C. for 24 hours to let the cells to grow to saturation. The cells were spun down, and the supernatants were sampled for the thermostability assay.

Two replicates of 96-well assay plates were prepared for each growth plate by transferring 10 µl of supernatant into the replica plates. The subtilisin activities were then measured by adding 100 µl of activity assay solution (0.2 mM succinyl-Ala-Ala-Pro-Phe-p-nitroanilide, 100 mM Tris-HCl, 10 mM CaCl₂, pH 8.0, 37° C.). Reaction velocities were measured at 405 nm to over 1.0 min in a ThermoMax microplate reader (Molecular Devices, Sunnyvale Calif.). Activity measured at room temperature was used to calculate the fraction of active clones (clones with activity less than 10% of that of wild type were scored as inactive). Initial activity (A_i) was measured after incubating one assay plate at 65° C. for 10 minutes by immediately adding 100 µl of prewarmed (37° C.) assay solution (0.2 mM succinyl-Ala-Ala-Pro-Phe-p-nitroanilide, 100 mM Tris-HCl, pH 8.0, 10 mM CaCl₂, pH 8.0) into each well. Residual activity (A_r) was measured after 40 minute incubation.

B. Results

In vitro heteroduplex formation and in vivo repair was carried out as described above. Five clones from *E. coli* SCS110 mutant library and five from *E. coli* ES1301 mutS libraries were selected at random and sequenced. FIG. 14 shows that four out of the ten clones were different from the parent genes. The frequency of occurrence of a particular point mutation from parent RC1 or RC2 in the resulting genes ranged from 0% to 50%, and the ten point mutations in the heteroduplex have been repaired without strong strand-specific preference.

Since none of the ten mutations locates within the dcm site, the mismatch repair appears generally done via the *E. coli* long-patch mismatch repair systems. The system repairs different mismatches in a strand-specific manner using the state of N6-methylation of adenine in GATC sequences as the major mechanism for determining the strand to be repaired. With heteroduplexes methylated at GATC sequences on only one DNA strand, repair was shown to be highly biased to the unmethylated strand, with the methylated strand serving as the template for correction. If neither strand was methylated, mismatch repair occurred, but showed little strand preference (23, 24). These results shows that it is preferable to demethylate the DNA to be recombined to promote efficient and random repair of the heteroduplexes.

The rates of subtilisin E thermo-inactivation at 65° C. were estimated by analyzing the 400 random clones from the *Bacillus subtilis* DB428 library. The thermostabilities obtained from one 96-well plate are shown in FIG. 16, plotted in descending order. About 12.9% of the clones exhibited thermostability comparable to the mutant with the N181D and N218S double mutations. Since this rate is only half of that expected for random recombination of these two markers, it indicates that the two mismatches at positions 995 and 1107 within the heteroduplexes have been repaired with lower position randomness.

Sequence analysis of the clone exhibiting the highest thermostability among the screened 400 transformants from the *E. coli* SCS110 heteroduplex library confirmed the presence of both N181D and N218S mutations. Among the 400 transformants from the *B. subtilis* DB428 library that were screened, approximately 91% of the clones expressed N181D- and/or N218S-type enzyme stabilities, while about 8.0% of the transformants showed only wild-type subtilisin E stability.

Less than 1.0% inactive clone was found, indicating that few new point mutations were introduced in the recombina-

tion process. This is consistent with the fact that no new point mutations were identified in the ten sequenced genes (FIG. 14). While point mutations may provide useful diversity for some in vitro evolution applications, they can also be problematic for recombination of beneficial mutations, especially when the mutation rate is high.

Example 4

Optimizing Conditions for the Heteroduplex Recombination

We have found that the efficiency of heteroduplex recombination can differ considerably from gene to gene [17,57]. In this example, we investigate and optimize a variety of parameters that improve recombination efficiency. DNA substrates used in this example were site-directed mutants of green fluorescent protein from *Aequorea victoria*. The GFP mutants had a stop codon(s) introduced at different locations along the sequence that abolished their fluorescence. Fluorescent wild type protein could be only restored by recombination between two or more mutations. Fraction of fluorescent colonies was used as a measure of recombination efficiency.

A. Methods

About 2-4 µg of each parent plasmid was used in one recombination experiment. One parent plasmid was digested with Pst I endonuclease another parent with EcoRI. Linearized plasmids were mixed together and 20×SSPE buffer was added to the final concentration 1× (180 mM NaCl, 1 mM EDTA, 10 mM NaH₂PO₄, pH 7.4). The reaction mixture was heated at 96° C. for 4 minutes, immediately transferred on ice for 4 minutes and the incubation was continued for 2 hours at 68° C.

Target genes were amplified in a PCR reaction with primers corresponding to the vector sequence of pGFP plasmid. Forward primer: 5'-CCGACTGGAAAGCGGGCAGTG-3', reverse primer 5'-CGGGGCTGGCTTAACTATGCGG-3'. PCR products were mixed together and purified using Qiagen PCR purification kit. Purified products were mixed with 20×SSPE buffer and hybridized as described above. Annealed products were precipitated with ethanol or purified on Qiagen columns and digested with EcoRI and PstI enzymes. Digested products were ligated into PstI and EcoRI digested pGFP vector.

dUTP was added into PCR reaction at final concentrations 200 µM, 40 µM, 8 µM, 1.6 µM, 0.32 µM. PCR reaction and subsequent cloning procedures were performed as described above.

Recombinant plasmids were transformed into XL10 *E. coli* strain by a modified chemical transformation method. Cells were plated on ampicillin containing LB agar plates and grown overnight at 37° C., followed by incubation at room temperature or at 4° C. until fluorescence developed.

B. Results.

1. Effect of Ligation on Recombination Efficiency.

Two experiments have been performed to test the effect of breaks in the DNA heteroduplex on the efficiency of recombination. In one experiment heteroduplex plasmid was treated with DNA ligase to close all existing single-strand breaks and was transformed in identical conditions as an unligated sample (see Table 1). The ligated samples show up to 7-fold improvement in recombination efficiency over unligated samples.

In another experiment, dUTP was added into PCR reaction to introduce additional breaks into DNA upon repair by uracyl N-glycosylase in the host cells. Table 2 shows that dUMP

incorporation significantly suppressed recombination, the extent of suppression increasing with increased dUTP concentration.

2. Effect of Plasmid Size on the Efficiency of Heteroduplex Formation.

Plasmid size was a significant factor affecting recombination efficiency. Two plasmids pGFP (3.3 kb) and a *Bacillus* shuttle vector pCT1 (about 9 kb) were used in preparing circular heteroduplex-like plasmids following traditional heteroduplex protocol. For the purpose of this experiment (to study the effect of plasmid size on duplex formation), both parents had the same sequences. While pGFP formed about 30-40% of circular plasmid, the shuttle vector yielded less than 10% of this form.

Increase in plasmid size decreases concentration of the ends in the vicinity of each and makes annealing of very long (>0.8 kb) ends that are single-stranded more difficult. This difficulty is avoided by the procedure shown in FIG. 3, in which heteroduplex formation occurs between substrates in vector-free form, and, heteroduplexes are subsequently inserted into a vector.

3. Efficiency of Recombination vs. Distance Between Mutations

A series of GFP variants was recombined pairwise to study the effect of distance between mutations on the efficiency of recombination. Parental genes were amplified by PCR, annealed and ligated back into pGFP vector. Heteroduplexes were transformed into XL10 *E. coli* strain.

The first three columns in Table 3 show the results of three independent experiments and demonstrate the dependence of recombination efficiency on the distance between mutations. As expected recombination becomes less and less efficient for very close mutations.

However, it is still remarkable that long-patch repair has been able to recombine mutations separated by only 27 bp.

The last line in Table 3 represents recombination between one single and one double mutants. Wild type GFP could only be restored in the event of double crossover with each individual crossover occurring in the distance of 99 bp only, demonstrating the ability of this method to recombine multiple, closely-spaced mutations.

4. Elimination of the Parental Double Strands from Heteroduplex Preparations.

Annealing of substrates in vector-free form offers size-advantages relative to annealing of substrates as components of vectors, but does not allow selection for heteroduplexes relative to homoduplexes simply by transformation into host. Asymmetric PCR reactions with only one primer for each parent seeded with appropriate amount of previously amplified and purified gene fragment were run for 100 cycles, ensuring a 100-fold excess of one strand over another. Products of these asymmetrical reactions were mixed and annealed together producing only a minor amount of non-recombinant duplexes. The last column in Table 3 shows the recombination efficiency obtained from these enriched heteroduplexes. Comparison of the first three columns with the fourth one demonstrates the improvement achieved by asymmetric synthesis of the parental strands.

While the foregoing invention has been described in some detail for purposes of clarity and understanding, it will be clear to one skilled in the art from a reading of this disclosure that various changes in form and detail can be made without departing from the true scope of the invention. All publications and patent documents cited in this application are incorporated by reference in their entirety for all purposes to the

same extent as if each individual publication or patent document were so individually denoted.

REFERENCES

1. Shao, Z. and Arnold, F. H. 1996. Engineering new functions and altering existing functions. *Curr. Opin. Struct. Biol.* 6:513-518.
2. Kuchner, O and Arnold, F. H. 1997. Directed evolution of enzyme catalysts. *Trends in Biotechnol.* 15:523-530.
3. Abelson, J. N. (ed.) 1996. Combinatorial chemistry. *Methods in Enzymol.* 267, Academic Press, Inc. San Diego.
4. Joyce, G. F. 1992. Directed molecular evolution. *Scientific American* 267:90-97.
5. Stemmer, W. P. C. 1994a. Rapid evolution of a protein in vitro by DNA shuffling. *Nature* 370:389-391.
6. Stemmer, W. P. C. 1994b. DNA shuffling by random fragmentation and reassembly: in vitro recombination for molecular evolution. *Proc. Natl. Acad. Sci. USA* 91:10747-10751.
7. Moore, J. C. and Arnold, F. H. 1996. Directed evolution of a para-nitrobenzyl esterase for aqueous-organic solvents. *Nature Biotech.* 14:458-467.
8. Holland, J. H. 1975. Adaptation in natural and artificial systems. The University Press, Ann Arbor.
9. Goldberg, D. E. 1989. Genetic algorithms in search, optimization and machine learning. Addison-Wesley. Reading.
10. Eigen, M. 1971. Self-organization of matter and the evolution of biological macromolecules. *Naturwissenschaften* 58:465-523.
11. Rechenberg, L. 1973. Evolutions strategic: Optimierung technischer Systeme nach Prinzipien der biologischen Evolution. Fronimann-Holzboog, Stuttgart.
12. Brady, R. M. 1985. Optimization strategies gleaned from biological evolution. *Nature* 317:804-806.
13. Muhlenbein, H. 1991. The parallel genetic algorithm as function optimizer. *Parallel Computing* 17:619-632.
14. Pal, K. F. 1993. Genetic algorithms for the traveling salesman problem-based on a heuristic crossover operation. *Bio. Cybern.* 69:539-546.
15. Pal, K. F. 1995. Genetic algorithm with local optimization. *Bio. Cybern.* 73:335-341.
16. Cami, B., P. Chambon, P. Kourlsky. 1984. Correction of complex heteroduplexes made of mouse H-2 gene sequences in *E. coli* K-12. *Proc. Natl. Acad. Sci. USA* 81:503-507.
17. Westmoreland, J, G. Porter, M. Radman and M. A. Resnick. 1997. Highly mismatched molecules resembling recombination intermediates efficiently transform mismatch repair proficient *E. coli*. *Genetics* 145:29-38.
18. Kramer, B., W. Kramer and H.-J. Fritz. 1984. Different base/base mismatches are corrected with different efficiencies by the methyl-directed DNA mismatch-repair system of *E. coli*. *Cell* 38:879-887.
19. Lu, A.-L., S. Clark and P. Modrich. 1983. Methyl-affected repair of DNA base pair mismatches in vitro. *Proc. Natl. Acad. Sci. USA* 80:4639-4643.
20. Carraway, M. and Marinus, M. G. 1993. Repair of heteroduplex DNA molecules with multibase loops in *Escherichia coli*. *Bacteriol.* 175:3972-3980.
21. Cooper, D. L., Lahue, R. S, and Modrich, P. 1993. Methyl-directed mismatch repair is bi-directional. *J. Biol. Chem.* 268:11823-11829.
22. Au, K. G., Welsh, K. and Modrich, P. 1992. Initiation of methyl-directed mismatch repair. *J. Biol. Chem.* 267:12142-12148.

23. Meselson, M. 1988. Methyl-directed repair of DNA mismatches, p. 91-113. In K. B. Low (ed.), *Recombination of the Genetic Material*. Academic Press, Inc., San Diego, Calif.
24. Fishel, R. A., Siegel, E. C. and Kolodner, R. 1986. Gene conversion in *Escherichia coli*. Resolution of heteroallelic mismatched nucleotides by co-repair. *J. Mol. Biol.* 188: 147-157.
25. Pukkila, P. J., J. Peterson, G. Herman, P. Modrich, and M. Meselson. 1983. Effects of high levels of DNA adenine methylation on methyl-directed mismatch repair in *Escherichia coli*. *Genetics* 104:571-582.
26. Radman, M., R. E. Wagner, B. W. Glickman, and M. Meselson. 1980. DNA methylation, mismatch correction and genetic stability, p. 121-130. In M. Alacevic (ed.) *Process in Environmental Mutagenesis*. Elsevier/North-Holland Biochemical Press, Amsterdam.
27. Sambrook, J., Fritsch, E. F. and Maniatis, T. 1989. *Molecular cloning: A Laboratory Manual*. 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.
28. Allen, D. J., Makhov, A., Grilley, M., Taylor, J., Thresher, R., Modrich, P. and Griffith, J. D. MutS mediates heteroduplex loop formation by a translocation mechanism. 1997. *EMBO J.* 16: 4467-4476.
30. Tsai-Wu, J. J. and Lu, A. L. 1994. *Escherichia coli* mutY-dependent mismatch repair involves DNA polymerase I and a short repair tract. *Mol. Gen. Genet.* 244:444-450.
31. Worth, L. Jr., Clark, S., Radman, M. and Modrich, P. 1994. Mismatch repair proteins MutS and MutL inhibit RecA-catalyzed strand transfer between diverged DNAs. *Proc. Natl. Acad. Sci. USA* 91:3238-3241.
32. Fox, M. S., Radicella, J. P. and Yamamoto, K. 1994. Some features of base pair mismatch repair and its role in the formation of genetic recombinants. *Experientia* 50:253-260.
33. Radicella, J. P., Clark, E. A., Chen, S. and Fox, M. S. 1993. Patch length of localized repair events: role of DNA polymerase I in mutY-dependent mismatch repair. *J. Bacteriol.* 175: 7732-7736.
34. Kraczkiewicz-Dowjat, A. and Fishel, R. 1990. RecB-recC-dependent processing of heteroduplex DNA stimulates recombination of an adjacent gene in *Escherichia coli*. *J. Bacteriol.* 172:172-178.
35. Radman, M. 1989. Mismatch repair and the fidelity of genetic recombination. *Genome* 31: 68-73.
36. Raposa, S. and Fox, M. S. 1987. Some features of base pair mismatch and heterology repair in *Escherichia coli*. *Genetics* 117:381-390.
37. Jones, M., Wagner, R. and Radman, M. 1987. Mismatch repair and recombination in *E. coli*. *Cell* 50:621-626.
38. Langle-Rouault, F., Maenhaut-Michel, G. and Radman, M. 1987. GATC sequences, DNA nicks and the MutH function in *Escherichia coli* mismatch repair. *EMBO J.* 6:1121-1127
39. Glazer, P. M., Sarkar, S. N., Chisholm, G. E. and Summers, W. C. 1987. DNA mismatch repair detected in human cell extracts. *Mol. Cell. Biol.* 7:218-224
40. Laengle-Rouault, F., Maenhaut-Michel, G. and Radman M. 1986. GATC sequence and mismatch repair in *Escherichia coli*. *EMBO J.* 5:2009-2013.
41. Bauer, J., Krammer, G. and Knippers, R. 1981. Asymmetric repair of bacteriophage T7 heteroduplex DNA. *Mol. Gen. Genet.* 181:541-547.
42. Wildenberg, J. and Meselson, M. 1975. Mismatch repair in heteroduplex DNA. *Proc. Natl. Acad. Sci. USA* 72:2202-2206.

43. Kirkpatrick, D. T. and Petes, T. D. 1997. Repair of DNA loops involves DNA-mismatch and nucleotide-excision repair proteins. *Nature* 387: 929-31.
44. Leung, W., Malkova, A. and Haber, J. E. 1997. Gene targeting by linear duplex DNA frequently occurs by assimilation of a single strand that is subject to preferential mismatch correction. *Proc. Natl. Acad. Sci. USA* 94: 6851-6856.
45. Hunter, N. and Borts, R. H. 1997. Mlh1 is unique among mismatch repair proteins in its ability to promote crossing-over during meiosis. *Genes Dev.* 11:0890-9369.
46. Alani, E., Lee, S., Kane, M. F., Griffith, J. and Kolodner, R. D. 1997. *Saccharomyces cerevisiae* MSH2, a mispaired base recognition protein, also recognizes Holliday junctions in DNA. *J. Mol. Biol.* 265:289-301.
47. Varlet, I., Canard, B., Brooks, P., Cerovic, G. and Radman, M. 1996. Mismatch repair in *Xenopus* egg extracts: DNA strand breaks act as signals rather than excision points. *Proc. Natl. Acad. Sci. USA* 93:10156-10161.
48. Nicolas, A. and Petes, T. D. 1994. Polarity of meiotic gene conversion in fungi: contrasting views. *Experientia* 50:242-52.
49. Bishop, D. K., J. Andersen, and R. D. Kolodner. 1989. Specificity of mismatch repair following transformation of *Saccharomyces cerevisiae* with heteroduplex plasmid DNA. *Proc. Natl. Acad. Sci. USA* 86:3713-3717.
50. Kramer, B., W. Kramer, M. S. Williamson, and S. Fogel. 1989. Heteroduplex DNA correction in *Saccharomyces cerevisiae* is mismatch specific and requires functional PMS genes *Mol. Cell. Biol.* 9:4432-4440.
51. Baynton, K., Bresson-Roy, A. and Fuchs, R. P. 1998. Analysis of damage tolerance pathways in *Saccharomyces cerevisiae*: a requirement for Rev3 DNA polymerase in translation synthesis. *Mol. Cell. Biol.* 18: 960-966.
52. Alani, E., Reenan, R. A. and Kolodner, R. D. 1994. Interaction between mismatch repair and genetic recombination in *Saccharomyces cerevisiae*. *Genetics* 137:19-39.
54. Bishop, D. K., Williamson, M. S., Fogel, S. and Kolodner, R. D. 1987. The role of heteroduplex correction in gene conversion in *Saccharomyces cerevisiae*. *Nature* 328:362-364.
55. Bishop, D. K. and Kolodner, R. D. 1986. Repair of heteroduplex plasmid DNA after transformation into *Saccharomyces cerevisiae*. *Mol. Cell. Biol.* 6:3401-3409.
56. White, J. H., Lusnak, K. and Fogel, S. 1985. Mismatch-specific post-meiotic segregation frequency in yeast suggests a heteroduplex recombination intermediate. *Nature* 315: 350-352.
57. Abastado, J.-P., B. Cami, T. H. Dinh, J. Igoler and P. Kourilsky. 1984. Processing of complex heteroduplexes in *E. coli* and Cos-1 monkey cells. *Proc. Natl. Acad. Sci. USA* 81:5792-5796.
58. Brown, T. C. and J. Jiricny. 1987. A specific mismatch repair event protects mammalian cells from loss of 5-methylcytosine. *Cell* 50:945-950.
59. Sibghat-Ullah, and R-S, Day. 1993. DNA-substrate sequence specificity of human G:T mismatch repair activity. *Nucleic Acids Res.* 21:1281-1287.
60. Miller, E. M., Hough, H. L., Cho, J. W. and Nickoloff, J. A. 1997. Mismatch repair by efficient nick-directed, and less efficient mismatch-specific, mechanisms in homologous recombination intermediates in Chinese hamster ovary cells. *Genetics* 147: 743-753.

61. Deng, W. P. and Nickoloff, J. A. 1994. Mismatch repair of heteroduplex DNA intermediates of extrachromosomal recombination in mammalian cells. *Mol. Cell. Biol.* 14:400-406.
62. Thomas, D. C., Roberts, J. D. and Kunkel, T. A. 1991. Heteroduplex repair in extracts of human HeLa cells. *J. Biol. Chem.* 266:3744-51.
63. Folger, K. R., Thomas, K. and Capecchi, M. R. 1985. Efficient correction of mismatched bases in plasmid heteroduplexes injected into cultured mammalian cell nuclei. *Mol. Cell. Biol.* 5:70-74.
64. Fang, W., Wu, J. Y. and Su, M. J. 1997. Methyl-directed repair of mismatched small heterologous sequences in cell extracts from *Escherichia coli*. *J. Biol. Chem.* 272: 22714-22720.
65. Smith, J. and Modrich, P. 1997. Removal of polymerase-produced mutant sequences from PCR products. *Proc. Natl. Acad. Sci. USA* 94: 6847-50.
66. Su, S. S., Grilley, M., Thresher, R., Griffith, J. and Modrich, P. 1989. Gap formation is associated with methyl-directed mismatch correction under conditions of restricted DNA synthesis. *Genome* 31:104-11.
67. Muster-Nassal, C. and Kolodner, R. 1986. Mismatch correction catalyzed by cell-free extracts of *Saccharomyces cerevisiae*. *Proc. Natl. Acad. Sci. USA* 83:7618-7622.
68. Macnab, R. M. 1992. Genetic and biogenesis of bacterial flagella. *Annul Rev. Genet.* 26:131-158.
69. Wilson, D. R. and Beveridge, T. J. 1993. Bacterial flagellar filaments and their component flagellins. *Can. J. Microbiol.* 39:451-472.
70. Schmitt, R., Raskal, A. and Mayer, F. 1974. Plain and complex flagella of *Pseudomonas rhodos*: analysis of fine structure and composition. *J. Bacteriol.* 117:844-857.
71. Gotz, R., Limmer, N., Ober, K. and Schmitt, R. 1982. Motility and chemotaxis in two strains of *Rhizobium* with complex flagella. *J. Gen. Microbiol.* 128:789-798.
72. Schmitt, R., Bambergerl., Acker G. and Mayer, F. 1974. Fine structure analysis of the complex flagella of *Rhizobium lupini* H13-3. *Arch. Microbiol.* 100:145-162.
73. Trachtenberg, S., DeRosier, D. J. and Macnab, R. M. 1987. Three-dimensional structure of the complex flagellar filament of *Rhizobium lupini* and its relation to the structure of the plain filaments. *J. Mol. Biol.* 195:603-620,
74. Gotz, R. and Schmitt, R. 1987. *Rhizobium meliloti* swims by unidirectional intermittent rotation of right-handed flagellar helices. *J. Bacteriol.* 169:3146-3150.
75. Lotz, W., Acker, G. and Schmitt, R. 1977. Bacteriophage 7-7-1 adsorbs to the complex flagella of *Rhizobium lupini* H13-3. *J. Gen. Virol.* 34:9-17.
76. Krupski, G., Gotz, F., Ober, K., Pleicr, E. and Schmitt, R. 1985. Structure of complex flagellar filaments in *Rhizobium meliloti*. *J. Bacteriol.* 162:361-366.
77. Maruyama, M., Lodderstaedt, G. and Schmitt, R. 1978. Purification and, biochemical properties of complex flagella isolated from *Rhizobium lupini* H13-3. *Biochem. Biophys. Acta* 535:110-124.
78. Trachtenberg, S., DeRosier, D. J., Aizawa, S.-I. and Macnab, R. M. 1986. Pairwise perturbation of flagellin subunits. The structural basis for the differences between plain and complex bacterial flagellar filaments. *J. Mol. Biol.* 190:569-576.

79. Gordee, R. S., Zeckner, D. J., Ellis, L. F., Thakkar, A. L. and Howard, L. C. 1984. In vitro and in vivo anti-Candida activity and toxicity of LY121019. *J. Antibiotics* 37:1054-1065.
- 5 80. Debono, M., Willard, K. E., Kirst, H. A., Wind, J. A., Crouse, G. D., Tao, E. V., Vicenzi, J. T., Counter, F. T., Ott, J. L., Ose, E. E. and Omura, S. 1989. Synthesis of new analogs of echinocandin B by enzymatic deacylation and chemical reacylation of the echinocandin B peptide: synthesis of the antifungal agent cilofungin (LY121019). *J. Antibiotics* 42(3): 389-397.
- 10 81. Debono, M. and Gordee, R. S. 1994. Antibiotics that inhibit fungal cell-wall development. *Annu. Rev. Microbiol.* 48: 471-497.
- 15 82. Debono, M., Turner, W. W., Lagrandeur, L., Burkhardt, F. J., Nissen, J. S., Nichols, K. K., Rodriguez, M. J., Zweifel, M. J., Zeckner, D. J., Gordee, R. S., Tang, J. and Parr, T. R. 1995. Semisynthetic chemical modification of the antifungal lipopeptide echinocandin B (ECB): structure-activity studies of the lipophilic and geometric parameters of polyarylated acyl analogs of ECB. *J. Med. Chem.* 38(17): 3271-3281.
- 20 83. Yeh, W. K. 1997. Evolving enzyme technology for pharmaceutical applications: case studies. *J. Ind. Microbiol. Biotechnol.* 19(5-6): 334-343.
- 25 84. Boeck, L. D., Fukuda, D., Abbott, B. J. and M. Debono. 1989. Deacylation of echinocandin B by *Actinoplanes utahensis*. *J. Antibiotics* 42(3): 382-388
- 30 85. Arnold, F. H. 1998. Design by directed evolution. *Accts. Chem. Res.* 31:125-131.
- 35 86. Shao, Z., Callahan, M. and Arnold, F. H. 1998. Directed enzyme evolution of *Actinoplane utahensis* ECB deacylase in *Streptomyces lividans* for enhanced specific activity. Manuscript submitted.
- 40 87. Shao, Z., Zhao, H., Giver, L. and Arnold, F. H. 1998. Random-priming in vitro recombination: an effective tool for directed evolution. *Nucleic Acids Res.* 26 (2): 681-683.
- 45 88. Zhao, H. and Arnold, F. H. 1997. Functional and nonfunctional mutations distinguished by random recombination of homologous genes. *Proc. Natl. Acad. Sci. USA* 94:7997-8000.
- 50 89. Zhao, H., Giver, L., Shao, Z., Affholter, J. A., and Arnold, F. H. 1998. Molecular evolution by staggered extension process (StEP) in vitro recombination. *Nat. Biotechnol.* 16: 258-261.
- 55 90. Judo, M. S. B., Wedel, A. B. and Wilson, C. 1998. Stimulation and suppression of PCR-mediated recombination. *Nucleic Acids Res.* 26: 1819-1825.
91. Okkels, J. S. 1997. Method for preparing polypeptide variants. PCT application WO 97/07205.
92. Gray, G. L. 1992. Hybrid prokaryotic polypeptides produced by in vivo homologous recombination. U.S. Pat. No. 5,093,257.
93. Weber, H. and Weissmann, C. 1983. Formation of genes coding for hybrid proteins by recombination between related, cloned genes in *E. coli*. *Nucl. Acids Res.* 11:5661-5669.
- 65 94. Maryon, E. and Carroll, D. 1991. Characterization of recombination intermediates from DNA injected into *Xenopus laevis* oocytes: evidence for a nonconservative mechanism of homologous recombination. *Mol. Cell. Biol.* 11:3278-3287.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 11

<210> SEQ ID NO 1

<211> LENGTH: 410

<212> TYPE: PRT

<213> ORGANISM: Rhizobium lupini

<220> FEATURE:

<223> OTHER INFORMATION: flagellin A (FlaA)

<400> SEQUENCE: 1

```

Met Ala Ser Val Leu Thr Asn Ile Asn Ala Met Ser Ala Leu Gln Thr
 1           5           10           15
Leu Arg Ser Ile Ser Ser Asn Met Glu Asp Thr Gln Ser Arg Ile Ser
          20           25           30
Ser Gly Met Arg Val Gly Ser Ala Ser Asp Asn Ala Ala Tyr Trp Ser
          35           40           45
Ile Ala Thr Thr Met Arg Ser Asp Asn Ala Ser Leu Ser Ala Val Gln
          50           55           60
Asp Ala Ile Gly Leu Gly Ala Ala Lys Val Asp Thr Ala Ser Ala Gly
          65           70           75           80
Met Asp Ala Val Ile Asp Val Val Lys Gln Ile Lys Asn Lys Leu Val
          85           90           95
Thr Ala Gln Glu Ser Ser Ala Asp Lys Thr Lys Ile Gln Gly Glu Val
          100          105          110
Lys Gln Leu Gln Glu Gln Leu Lys Gly Ile Val Asp Ser Ala Ser Phe
          115          120          125
Ser Gly Glu Asn Trp Leu Lys Gly Asp Leu Ser Thr Thr Thr Thr Lys
          130          135          140
Ser Val Val Gly Ser Phe Val Arg Glu Gly Gly Thr Val Ser Val Lys
          145          150          155          160
Thr Ile Asp Tyr Ala Leu Asn Ala Ser Lys Val Leu Val Asp Thr Arg
          165          170          175
Ala Thr Gly Thr Lys Thr Gly Ile Leu Asp Thr Ala Tyr Thr Gly Leu
          180          185          190
Asn Ala Asn Thr Val Thr Val Asp Ile Asn Lys Gly Gly Val Ile Thr
          195          200          205
Gln Ala Ser Val Arg Ala Tyr Ser Thr Asp Glu Met Leu Ser Leu Gly
          210          215          220
Ala Lys Val Asp Gly Ala Asn Ser Asn Val Ala Val Gly Gly Gly Ser
          225          230          235          240
Ala Ser Ser Arg Ser Thr Ala Ala Gly Leu Arg Val Ala Ser Thr Leu
          245          250          255
Arg Pro Pro Ser Pro His Gln His Gln Ser Leu Ala Ser Leu Pro Pro
          260          265          270
Leu Thr Pro Pro Leu Lys Leu Val Leu Gln Leu Leu Pro Val Thr Pro
          275          280          285
Ser Ser Ser Thr Lys Pro Thr Ala Ala Pro Val Gln Val Asn Leu Thr
          290          295          300
Gln Ser Val Leu Thr Met Asp Val Ser Ser Met Ser Ser Thr Asp Val
          305          310          315          320
Gly Ser Tyr Leu Thr Gly Val Glu Lys Ala Leu Thr Ser Leu Thr Ser
          325          330          335
Ala Gly Ala Glu Leu Gly Ser Ile Lys Gln Arg Ile Asp Leu Gln Val
          340          345          350

```

-continued

Asp Phe Ala Ser Lys Leu Gly Asp Ala Leu Ala Lys Gly Ile Gly Arg
 355 360 365

Leu Val Asp Ala Asp Met Asn Glu Glu Ser Thr Lys Leu Lys Ala Leu
 370 375 380

Gln Thr Gln Gln Gln Leu Ala Ile Gln Ser Leu Ser Ile Ala Asn Ser
 385 390 395 400

Asp Ser Gln Asn Ile Leu Ser Leu Phe Arg
 405 410

<210> SEQ ID NO 2
 <211> LENGTH: 394
 <212> TYPE: PRT
 <213> ORGANISM: Rhizobium meliloti
 <220> FEATURE:
 <223> OTHER INFORMATION: flagellin A (FlaA)

<400> SEQUENCE: 2

Met Thr Ser Ile Leu Thr Asn Asn Ser Ala Met Ala Ala Leu Ser Thr
 1 5 10 15

Leu Arg Ser Ile Ser Ser Ser Met Glu Asp Thr Gln Ser Arg Ile Ser
 20 25 30

Ser Gly Leu Arg Val Gly Ser Ala Ser Asp Asn Ala Ala Tyr Trp Ser
 35 40 45

Ile Ala Thr Thr Met Arg Ser Asp Asn Gln Ala Leu Ser Ala Val Gln
 50 55 60

Asp Ala Leu Gly Leu Gly Ala Ala Lys Val Asp Thr Ala Tyr Ser Gly
 65 70 75 80

Met Glu Ser Ala Ile Glu Val Val Lys Glu Ile Lys Ala Lys Leu Val
 85 90 95

Ala Ala Thr Glu Asp Gly Val Asp Lys Ala Lys Ile Gln Glu Glu Ile
 100 105 110

Thr Gln Leu Lys Asp Gln Leu Thr Ser Ile Ala Glu Ala Ala Ser Phe
 115 120 125

Ser Gly Glu Asn Trp Leu Gln Ala Asp Leu Ser Gly Gly Pro Val Thr
 130 135 140

Lys Ser Val Val Gly Gly Phe Val Arg Asp Ser Ser Gly Ala Val Ser
 145 150 155 160

Val Lys Lys Val Asp Tyr Ser Leu Asn Thr Asp Thr Val Leu Phe Asp
 165 170 175

Thr Thr Gly Asn Thr Gly Ile Leu Asp Lys Val Tyr Asn Val Ser Gln
 180 185 190

Ala Ser Val Thr Leu Pro Val Asn Val Asn Gly Thr Thr Ser Glu Tyr
 195 200 205

Thr Val Gly Ala Tyr Asn Val Asp Asp Leu Ile Asp Ala Ser Ala Thr
 210 215 220

Phe Asp Gly Asp Tyr Ala Asn Val Gly Ala Gly Ala Leu Ala Gly Asp
 225 230 235 240

Tyr Val Lys Val Gln Gly Ser Trp Val Lys Ala Val Asp Val Ala Ala
 245 250 255

Thr Gly Gln Glu Val Val Tyr Asp Asp Gly Thr Thr Lys Trp Gly Val
 260 265 270

Asp Thr Thr Val Thr Gly Ala Pro Ala Thr Asn Val Ala Ala Pro Ala
 275 280 285

Ser Ile Ala Thr Ile Asp Ile Thr Ile Ala Ala Gln Ala Gly Asn Leu
 290 295 300

-continued

Asp Ala Leu Ile Ala Gly Val Asp Glu Ala Leu Thr Asp Met Thr Ser
 305 310 315 320
 Ala Ala Ala Ser Leu Gly Ser Ile Ser Ser Arg Ile Asp Leu Gln Ser
 325 330 335
 Asp Phe Val Asn Lys Leu Ser Asp Ser Ile Asp Ser Gly Val Gly Arg
 340 345 350
 Leu Val Asp Ala Asp Met Asn Glu Glu Ser Thr Arg Leu Lys Ala Leu
 355 360 365
 Gln Thr Gln Gln Gln Leu Ala Ile Gln Ala Leu Ser Ile Ala Asn Ser
 370 375 380
 Asp Ser Gln Asn Val Leu Ser Leu Phe Arg
 385 390

<210> SEQ ID NO 3
 <211> LENGTH: 1201
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:SCS01 mosaic
 flaA gene created by in vitro heteroduplex
 formation followed by in vivo repair

<400> SEQUENCE: 3

atggcaagcg ttctcacaaa cattaacgca atgtctgctc ttcagacgct gcgttcgatt 60
 tcttccaaca tggaagacac ccagagccgt atttccagcg gcatgcgctg tggttcggct 120
 tccgacaacg ccgcttattg gtctatcgcg accaccatgc gctcggacaa tgectcgctt 180
 tccgctgttc aggatgcaat tggcctcggt gccgccaagg tcgataccgc ttcggcgggt 240
 atggatgctg ttatcgatgt tgtaaagcag atcaagaaca aactggctac tgccaccgaa 300
 gacggcgctg acaaggccaa gatccaagaa gaaatcactc agctcaagga ccagctgacg 360
 agcatcgccg acgcggtctc cttctccggt gaaaactggc tcaagggcga tctttccacg 420
 acgacaacca aatcagtggg tggtccttc gttcgtgaag gcggtaccgt atcgggtcaag 480
 accatcgatt acgctctgaa tgcttccaag gttctgggtg ataccgctg aacgggcacc 540
 aagaccggca ttctggacaa ggtctacaac gtctcgcagg caagcgtcac gctgacggtc 600
 aacaccaacg gcgtcgaatc ccaggcctcc gtccgcgctc attcgtgga gtcctcacc 660
 gaagccgggtg cggagtcca gggcaactat gctcttcagg gcggtaacag ctacgtcaag 720
 gtcgaaaacg tctgggttcg agctgagacc gcatcaaacac cagtcgctgg caagtttgcc 780
 gccgcttaca ccgccgctga agctgggtact gcagctgctg ccggtgacgc catcatcgtc 840
 gacgaaacca acagcggcgc cgggtgcagg aaacctcacc cagtcggtcc tgaccatgga 900
 tgtcagctcg atgagctcga cggatgtcgg cagctacctc acgggcgtgg aaaaggctct 960
 caccagcctg acgagcgtg gcgctgaact cggtctatc aaacagcgc tcatctgca 1020
 ggttgatttt gcttccaagc tgggacgacg tctcgaaaaa ggtattggcc gtctcgttga 1080
 tgctgacatg aatgaagagt ccactaagct taaggctctt cagacgcagc agcagctggc 1140
 tatccagtcg ctctccatcg caaacagcga ctgcgagaac attctgtcgc tgttccgtta 1200
 a 1201

<210> SEQ ID NO 4
 <211> LENGTH: 1229
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:

-continued

<223> OTHER INFORMATION: Description of Artificial Sequence:SCS02 mosaic
flaA gene created by in vitro heteroduplex
formation followed by in vivo repair

<400> SEQUENCE: 4

```

atgacgagca ttctcaccaa caactccgca atggccgcgc tttccggagt gcgctcgatc    60
tcttccagca tggaagacac gcagagccgc atctcctccg gccttcgcgt cggttcggcc    120
tccgacaacg ccgctactg gtcgattgcg accaccatgc gctccgacaa ccaggccctt    180
tcggccgtcc aggacgccct cggcctcggc gccgccaagg ttgataccgc ctattccggg    240
atggaatcgg cgatcgaagt cgtaaggaa atcaagaaca aactggtcac tgctcaggaa    300
tcttctgccg acaaaacgaa gattcagggc gaagtcaagc agcttcagga gcagttgaag    360
ggcatcgttg attccgcttc cttctccggg gagaactggc tgcaggcgga cctcagcggc    420
ggcgccgtca ccaagagcgt cgtcggctcg ttcgtccgtg acggaagcgg ttccgtagcc    480
gtcaagaagg tcgattacgc tctgaatgct tccaaggttc tggggtgatac ccgcgcaacg    540
ggcaccaaga ccggcattct cgatactgct tataccggcc ttaacgcgaa cacggtgacg    600
gttgatatca acaagggcgg cgtgatcacc caggcctccg tccgcgccta ttccacggac    660
gaaatgctct ccctcggcgc aaaggtcgat ggcgcaaaaca gcaacgttgc tgttggcggc    720
ggctccgctt cgtcaaggtc gacggcagct gggtaaggg tagcgtcgac gctgcggcct    780
ccatcacccg atcaaccggc gccaccggtc aagaaatcgc cgccaccacg acggcagctg    840
gtaccatcac tgacagacgc tgggtcgtcg atgtcggcaa cgctcctgcc gccaacgttt    900
cggccggcca gtcggtcgcg aacatcaaca tcgtcggaat gggctcgacg gatgtcggca    960
gctacctcac gggcgtgaa aaggctctca ccagcatgac cagcgtgcc gcctcgtcgc    1020
gtccatctc ctcgcgcac gacctgcaga gcgaattcgt caacaagctc tcggactcga    1080
tcgagtcggg cgtcggccgt ctcgtcgacg cggacatgaa cgaggagtcg acccgcctca    1140
aggccctgca gaccagcag cagctcgcca tccaggccct gtcgatcgcc aactcggact    1200
cgcagaacgt cctgtcgtc ttccgctaa    1229

```

<210> SEQ ID NO 5

<211> LENGTH: 1228

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence:ES01 mosaic
flaA gene created by in vitro heteroduplex
formation followed by in vivo repair

<400> SEQUENCE: 5

```

atgacgagca ttctcaccaa caactccgca atggccgcgc tttccggagt gcgctcgatc    60
tcttccagca tggaagacac gcagagccgc atctcctccg gccttcgcgt cggttcggcc    120
tccgacaacg ccgctactg gtcgattgcg accaccatgc gctccgacaa ccaggccctt    180
tcggccgtcc aggacgccct cggcctcggc gccgccaagg ttgataccgc ctattccggg    240
atggaatcgg cgatcgaagt cgtaaggaa atcaaggcca agctcgtagc tgccaccgaa    300
gacggcgtcg acaaggcca gatccaagaa gaaatcactc agctcaagga ccagctgacg    360
agcatcgccg acgcggttc cttctccggg gagaactggc tgcaggcgga cctcagcggc    420
ggcgccgtca ccaagagcgt cgtcggctcg ttcgtccgtg acggaagcgg ttccgtagcc    480
gtcaagacca tcgattacgc tctgaatgct tccaaggttc tggggtgatac ccgcgacacg    540
gtcggcgata ccggcattct ggacaaggtc tacaacgtct cgcaggcaag cgtcacgctg    600

```


-continued

```

acggtcaaca ccaacggcgt cgaatcgag catacggttg ctgcctattc gctggagtcc 660
ctcaccgaag ccggtgcgga gttccagggc aactatgctc ttcagggcgg taacagctac 720
gtcaaggctg acggcagctg ggtaagggt agcgtcgacg ctgcggcctc catcaccgca 780
tcaacaccag tcgctggcaa gtttgccgcc gcttacaccg ccgctgaagc tggtagtca 840
gctgctgccg gtgacgccat catcgctgac gaaaccaaca gcggcgccgg tgcaggtaaa 900
cctcaccag tcggtcctga ccatggatgt cagctcgatg agctcgacgg atgtcggcag 960
ctacctcag gccgtgaaa aggctctcac cagcctgacg agcgtggcg ctgaactcgg 1020
ctccatctcc tcgcgcatcg acctgcagag cgaattcgtc aacaagctct cggactcgat 1080
cgagtcgggc gtcggccgtc tcgctgacgc ggacatgaac gaggagtca cccgcctcaa 1140
ggccctgcag acccagcagc agctcgccat ccaggccctg tcgactgcca actcggactc 1200
gcagaacgtc ctgctgctct tccgctaa 1228

```

<210> SEQ ID NO 6

<211> LENGTH: 1209

<212> TYPE: DNA

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

```

<223> OTHER INFORMATION: Description of Artificial Sequence:ES02 mosaic
flaA gene created by in vitro heteroduplex
formation followed by in vivo repair

```

<400> SEQUENCE: 6

```

atgacgagca ttctaccaa caactccgca atggccgcgc tttccggagt gcgctcgatc 60
tcttccagca tggaagacac gcagagccgc atctcctccg gccttcgctg cggttcggcc 120
tccgacaacg ccgcctactg gtcgattgcg accaccatgc gctccgaca ccaggccctt 180
tcggccgtcc aggacgccct cggcctcggc gccgccaaagg ttgataccgc ctattccggg 240
atggaatcgg cgatcgaagt cgttaaggaa atcaaggcca agctcgtagc tgccaccgaa 300
gacggcgtcg acaaggcca gatccaagaa gaaatcactc agctcaagga ccagctgacg 360
agcatcgccg acgcggttc cttctccggg gagaactggc tgcaggcgga cctcagcggc 420
ggcgccgtca ccaagagcgt cgtcggctcg ttcgtccgtg acggaagcgg ttccgtagcc 480
gtcaagacca tcgattacgc tctgaatgct tccaaggttc tggtaggata ccgcgcaacg 540
ggaccaaga ccggcattct cgatactgct tataccggcc ttaacgcgaa cacggtgacg 600
gttgatatca acaagggcgg cgtgatcacc caggcctccg tccgcgccta ttccacggac 660
gaaatgctct ccctaccga agccgggtgc gagttccagg gcaactatgc tcttcagggc 720
ggtaacagct acgtcaaggt cgaaaacgtc tgggttcgag ctgagaccgc tgcaaccggc 780
gccaccggtc aagaaatcgc cgcaccacg acggcagctg gtaccatcac tgcagacagc 840
tgggtcgtcg atgtcggcaa cgtcctgcc gccaacggtt cggccggcca gtcggctcgcg 900
aacatcaaca tcgctggaat ggggtcagct gcgctcgatg ccctgatcag cgggtgctgac 960
gccgctttga cagacatgac cagcgtgcc gcctcgtcgc gctccatctc ctccgcatc 1020
gacctgcaga gcgaattcgt caacaagctc tcggactcga tcgagtcggg cgtcggccgt 1080
ctcgtcgacg cggacatgaa cgaggagtgc acccgcctca aggcctgca gaccacgacg 1140
cagctcgcca tccaggccct gtcgatcgcc aactcggact cgcagaacgt cctgctgctc 1200
ttccgctaa 1209

```

<210> SEQ ID NO 7

-continued

<211> LENGTH: 4039
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Actinoplanes
 utahensis echinocandin B (ECB) deacylase gene
 mutant M-15 created by in vitro heteroduplex
 formation followed by in vivo repair
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (1196)..(3559)

<400> SEQUENCE: 7

```

ctgcagcgtg cccagctggt cgtggtggtg atcgcggccg cgctggccgc cgctcgcggtc   60
gccgcccgcg ggccgatcga gttcgtcgcc ttcgtcgtgc cgcagatcgc cctgcccgtc   120
tgcggcggca gccggccgcc cctgctcgcc tcggcgatgc tcggcgcgct gctggtggtc   180
ggcggcggcc tggctcgctca gatcgtggtg gcgccgaagg agctgccggt cggcctgctc   240
accgcatgga tcggcaccac gtacctgctc tggctcctgc ttcggcgatc aagaaagggtg   300
agcggatgaa cggccgcctg cgtggcgagg gcctgcacct cgcgtacggg gacctgaccg   360
tgatcgacgg cctcgacgtc gacgtgcacg acgggctggt caccaccatc atcgggcccga   420
acgggtgctg caagtgcgac ctgctcaagg cgtcggccgc gctgctgcgc ccgaccggcg   480
ggcagggtgt gctggacggc cggccgatcg accggacccc caccctgac gtggcccggg   540
tgctcggcgt gctgcccagc tcggccaccg cggccgaagg gctcaccgtc gccgacctgg   600
tgatgcgcgg ccggcaccac caccagacct ggttccggca gtggtcgcgc gacgacgagg   660
accaggctgc cgacgcgctg cgctggaccg acatgctggc gtacgcggac cggccggtgg   720
acgccctctc cggcggctcag cggcagcgcg cctggatcag catggcgctg gccaggggca   780
ccgacctgct gctgctggac gagccgacca ccttctcoga cctggcccac cagatcgacg   840
tgctggacct ggtccgcccg ctgcacgccg agatgggccc gaccgtggtg atggtgctgc   900
acgacctgag cctggcccgc cggtagcccg accggctgat cgcgatgaag gacggcccga   960
tcgtggcgag cggggcgccc gacgaggtgc tcaccccggc gctgctggag tcggtcttcg  1020
ggctgcgcgc gatggtggtg cccgaccgag cggccggcac cccgctggtg atccccctgc  1080
cgcgccccgc cacctcgggt cgggctgaa atcgatgagc gtggttgett catcggcctg  1140
ccgagcgatg agagtatgtg ggcggtagag cgagtctcga ggggggatg ccgcc  gtg  1198
                               Val
                               1

acg tcc tcg tac atg cgc ctg aaa gca gca gcg atc gcc ttc ggt gtg   1246
Thr Ser Ser Tyr Met Arg Leu Lys Ala Ala Ala Ile Ala Phe Gly Val
      5                10                15

atc gtg gcg acc gca gcc gtg ccg tca ccc gct tcc ggc agg gaa cat   1294
Ile Val Ala Thr Ala Ala Val Pro Ser Pro Ala Ser Gly Arg Glu His
      20                25                30

gac ggc ggc tat gcg gcc ctg atc cgc cgg gcc tcg tac ggc gtc ccg   1342
Asp Gly Gly Tyr Ala Ala Leu Ile Arg Arg Ala Ser Tyr Gly Val Pro
      35                40                45

cac atc acc gcc gac gac ttc ggg agc ctc ggt ttc ggc gtc ggg tac   1390
His Ile Thr Ala Asp Asp Phe Gly Ser Leu Gly Phe Gly Val Gly Tyr
      50                55                60                65

gtg cag gcc gag gac aac atc tgc gtc atc gcc gag agc gta gtg acg   1438
Val Gln Ala Glu Asp Asn Ile Cys Val Ile Ala Glu Ser Val Val Thr
      70                75                80

gcc aac ggt gag cgg tcg cgg tgg ttc ggt gcg acc ggg ccg gac gac   1486
Ala Asn Gly Glu Arg Ser Arg Trp Phe Gly Ala Thr Gly Pro Asp Asp
      85                90                95

```

-continued

gcc gat gtg cgc agc gac ctc ttc cac cgc aag gcg atc gac gac cgc	1534
Ala Asp Val Arg Ser Asp Leu Phe His Arg Lys Ala Ile Asp Asp Arg	
100 105 110	
gtc gcc gag cgg ctc ctc gaa ggg ccc cgc gac ggc gtg cgg gcg ccg	1582
Val Ala Glu Arg Leu Leu Glu Gly Pro Arg Asp Gly Val Arg Ala Pro	
115 120 125	
tcg gac gac gtc cgg gac cag atg cgc ggc ttc gtc gcc ggc tac aac	1630
Ser Asp Asp Val Arg Asp Gln Met Arg Gly Phe Val Ala Gly Tyr Asn	
130 135 140 145	
cac ttc cta cgc cgc acc ggc gtg cac cgc ctg acc gac ccg gcg tgc	1678
His Phe Leu Arg Arg Thr Gly Val His Arg Leu Thr Asp Pro Ala Cys	
150 155 160	
cgc ggc aag gcc tgg gtg cgc ccg ctc tcc gag atc gat ctc tgg cgt	1726
Arg Gly Lys Ala Trp Val Arg Pro Leu Ser Glu Ile Asp Leu Trp Arg	
165 170 175	
acg tcg tgg gac agc atg gtc cgg gcc ggt tcc ggg gcg ctg ctc gac	1774
Thr Ser Trp Asp Ser Met Val Arg Ala Gly Ser Gly Ala Leu Leu Asp	
180 185 190	
ggc atc gtc gcc gcg acg cca cct aca gcc gcc ggg ccc gcg tca gcc	1822
Gly Ile Val Ala Ala Thr Pro Pro Thr Ala Ala Gly Pro Ala Ser Ala	
195 200 205	
ccg gag gca ccc gac gcc gcc gcg atc gcc gcc gcc ctc gac ggg acg	1870
Pro Glu Ala Pro Asp Ala Ala Ala Ile Ala Ala Ala Leu Asp Gly Thr	
210 215 220 225	
agc gcg ggc atc ggc agc aac gcg tac ggc ctc ggc gcg cag gcc acc	1918
Ser Ala Gly Ile Gly Ser Asn Ala Tyr Gly Leu Gly Ala Gln Ala Thr	
230 235 240	
gtg aac ggc agc ggg atg gtg ctg gcc aac ccg cac ttc ccg tgg cag	1966
Val Asn Gly Ser Gly Met Val Leu Ala Asn Pro His Phe Pro Trp Gln	
245 250 255	
ggc gcc gca cgc ttc tac cgg atg cac ctc aag gtg ccc ggc cgc tac	2014
Gly Ala Ala Arg Phe Tyr Arg Met His Leu Lys Val Pro Gly Arg Tyr	
260 265 270	
gac gtc gag ggc gcg gcg ctg atc ggc gac ccg atc atc ggg atc ggg	2062
Asp Val Glu Gly Ala Ala Leu Ile Gly Asp Pro Ile Ile Gly Ile Gly	
275 280 285	
cac aac cgc acg gtc gcc tgg agc cac acc gtc tcc acc gcc cgc cgg	2110
His Asn Arg Thr Val Ala Trp Ser His Thr Val Ser Thr Ala Arg Arg	
290 295 300 305	
ttc gtg tgg cac cgc ctg agc ctc gtg ccc ggc gac ccc acc tcc tat	2158
Phe Val Trp His Arg Leu Ser Leu Val Pro Gly Asp Pro Thr Ser Tyr	
310 315 320	
tac gtc gac ggc cgg ccc gag cgg atg cgc gcc cgc acg gtc acg gtc	2206
Tyr Val Asp Gly Arg Pro Glu Arg Met Arg Ala Arg Thr Val Thr Val	
325 330 335	
cag acc ggc agc ggc ccg gtc agc cgc acc ttc cac gac acc cgc tac	2254
Gln Thr Gly Ser Gly Pro Val Ser Arg Thr Phe His Asp Thr Arg Tyr	
340 345 350	
ggc ccg gtg gcc gtg atg ccg ggc acc ttc gac tgg acg ccg gcc acc	2302
Gly Pro Val Ala Val Met Pro Gly Thr Phe Asp Trp Thr Pro Ala Thr	
355 360 365	
gcg tac gcc atc acc gac gtc aac gcg ggc aac aac cgc gcc ttc gac	2350
Ala Tyr Ala Ile Thr Asp Val Asn Ala Gly Asn Asn Arg Ala Phe Asp	
370 375 380 385	
ggg tgg ctg cgg atg ggc cag gcc aag gac gtc cgg gcg ctc aag gcg	2398
Gly Trp Leu Arg Met Gly Gln Ala Lys Asp Val Arg Ala Leu Lys Ala	
390 395 400	
gtc ctc gac cgg cac cag ttc ctg ccc tgg gtc aac gtg atc gcc gcc	2446
Val Leu Asp Arg His Gln Phe Leu Pro Trp Val Asn Val Ile Ala Ala	

-continued

405	410	415	
gac gcg cgg ggc gag gcc ctc tac ggc gat cat tcg gtc gtc ccc cgg Asp Ala Arg Gly Glu Ala Leu Tyr Gly Asp His Ser Val Val Pro Arg 420 425 430			2494
gtg acc ggc gcg ctc gct gcc gcc tgc atc ccg gcg ccg ttc cag ccg Val Thr Gly Ala Leu Ala Ala Ala Cys Ile Pro Ala Pro Phe Gln Pro 435 440 445			2542
ctc tac gcc tcc agc ggc cag gcg gtc ctg gac ggt tcc cgg tcg gac Leu Tyr Ala Ser Ser Gly Gln Ala Val Leu Asp Gly Ser Arg Ser Asp 450 455 460 465			2590
tgc gcg ctc ggc gcc gac ccc gac gcc gcg gtc ccg ggc att ctc ggc Cys Ala Leu Gly Ala Asp Pro Asp Ala Ala Val Pro Gly Ile Leu Gly 470 475 480			2638
ccg gcg agc ctg ccg gtg cgg ttc cgc gac gac tac gtc acc aac tcc Pro Ala Ser Leu Pro Val Arg Phe Arg Asp Asp Tyr Val Thr Asn Ser 485 490 495			2686
aac gac agt cac tgg ctg gcc agc ccg gcc gcc ccg ctg gaa ggc ttc Asn Asp Ser His Trp Leu Ala Ser Pro Ala Ala Pro Leu Glu Gly Phe 500 505 510			2734
ccg cgg atc ctc ggc aac gaa cgc acc ccg cgc agc ctg cgc acc cgg Pro Arg Ile Leu Gly Asn Glu Arg Thr Pro Arg Ser Leu Arg Thr Arg 515 520 525			2782
ctc ggg ctg gac cag atc cag cag cgc ctc gcc ggc acg gac ggt ctg Leu Gly Leu Asp Gln Ile Gln Gln Arg Leu Ala Gly Thr Asp Gly Leu 530 535 540 545			2830
ccc ggc aag ggc ttc acc acc gcc cgg ctc tgg cag gtc atg ttc ggc Pro Gly Lys Gly Phe Thr Thr Ala Arg Leu Trp Gln Val Met Phe Gly 550 555 560			2878
aac cgg atg cac ggc gcc gaa ctc gcc cgc gac gac ctg gtc gcg ctc Asn Arg Met His Gly Ala Glu Leu Ala Arg Asp Asp Leu Val Ala Leu 565 570 575			2926
tgc cgc cgc cag ccg acc gcg acc gcc tcg aac ggc gcg atc gtc gac Cys Arg Arg Gln Pro Thr Ala Thr Ala Ser Asn Gly Ala Ile Val Asp 580 585 590			2974
ctc acc gcg gcc tgc acg gcg ctg tcc cgc ttc gat gag cgt gcc gac Leu Thr Ala Ala Cys Thr Ala Leu Ser Arg Phe Asp Glu Arg Ala Asp 595 600 605			3022
ctg gac agc cgg ggc gcg cac ctg ttc acc gag ttc gcc ctc gcg ggc Leu Asp Ser Arg Gly Ala His Leu Phe Thr Glu Phe Ala Leu Ala Gly 610 615 620 625			3070
gga atc agg ttc gcc gac acc ttc gag gtg acc gat ccg gta cgc acc Gly Ile Arg Phe Ala Asp Thr Phe Glu Val Thr Asp Pro Val Arg Thr 630 635 640			3118
ccg cgc cgt ctg aac acc acg gat ccg cgg gta cgg acg gcg ctc gcc Pro Arg Arg Leu Asn Thr Thr Asp Pro Arg Val Arg Thr Ala Leu Ala 645 650 655			3166
gac gcc gtg caa cgg ctc gcc ggc atc ccc ctc gac gcg aag ctg gga Asp Ala Val Gln Arg Leu Ala Gly Ile Pro Leu Asp Ala Lys Leu Gly 660 665 670			3214
gac atc cac acc gac agc cgc ggc gaa cgg cgc atc ccc atc cac ggt Asp Ile His Thr Asp Ser Arg Gly Glu Arg Arg Ile Pro Ile His Gly 675 680 685			3262
ggc cgc ggg gaa gca ggc acc ttc aac gtg atc acc aac ccg ctc gtg Gly Arg Gly Glu Ala Gly Thr Phe Asn Val Ile Thr Asn Pro Leu Val 690 695 700 705			3310
ccg ggc gtg gga tac ccg cag gtc gtc cac gga aca tcg ttc gtg atg Pro Gly Val Gly Tyr Pro Gln Val Val His Gly Thr Ser Phe Val Met 710 715 720			3358
gcc gtc gaa ctc ggc ccg cac ggc ccg tcg gga cgg cag atc ctc acc			3406

-continued

Ala	Val	Glu	Leu	Gly	Pro	His	Gly	Pro	Ser	Gly	Arg	Gln	Ile	Leu	Thr	
			725					730					735			
tat	gcg	cag	tcg	acg	aac	ccg	aac	tca	ccc	tgg	tac	gcc	gac	cag	acc	3454
Tyr	Ala	Gln	Ser	Thr	Asn	Pro	Asn	Ser	Pro	Trp	Tyr	Ala	Asp	Gln	Thr	
		740					745					750				
gtg	ctc	tac	tcg	cgg	aag	ggc	tgg	gac	acc	atc	aag	tac	acc	gag	gcg	3502
Val	Leu	Tyr	Ser	Arg	Lys	Gly	Trp	Asp	Thr	Ile	Lys	Tyr	Thr	Glu	Ala	
		755				760					765					
cag	atc	gcg	gcc	gac	ccg	aac	ctg	cgc	gtc	tac	cgg	gtg	gca	cag	cgg	3550
Gln	Ile	Ala	Ala	Asp	Pro	Asn	Leu	Arg	Val	Tyr	Arg	Val	Ala	Gln	Arg	
770					775				780					785		
gga	cgc	tgaccacgt	cacgccggct	cggcccgtgc	gggggcgcag	ggcgcgatc										3606
Gly	Arg															
gtctctgcat	cgccggtcag	cgggggctg	cgctgaccgg	cgccggccgg	tcgacgccc											3666
cgctccggcg	cagcgactgg	ctgaagcgcc	aggcgtcggc	ggcccggggc	aggttgttga											3726
acatcacgta	cgccgggccc	ccgtcgagga	tgccggcgag	gtgtgccagc	tcggcatccg											3786
tgtacacatg	ccgggcccgg	gtgatgccgt	gcagccggta	ataggccatc	ggcgtcagac											3846
tgccggcgag	gaacgggtcg	gcccgcgtggg	tcaggtccag	ctcctggcac	aagccctcga											3906
ccacctcgtc	cggccacggg	ccgcgcggct	cccacaacag	ccggacaccg	gccggcccggc											3966
gcgctcgggc	gcagaactca	cgcagtcgcg	cgatggcggg	ttcggtcggc	cggaaactcg											4026
ccgggcactg	cag															4039

<210> SEQ ID NO 8

<211> LENGTH: 787

<212> TYPE: PRT

<213> ORGANISM: Artificial Sequence

<220> FEATURE:

<223> OTHER INFORMATION: Description of Artificial Sequence: Actinoplanes utahensis echinocandin B (ECB) deacylase protein mutant M-15 transcribed from gene created by in vitro heteroduplex formation followed by in vivo repair

<400> SEQUENCE: 8

Val	Thr	Ser	Ser	Tyr	Met	Arg	Leu	Lys	Ala	Ala	Ala	Ile	Ala	Phe	Gly	
1				5					10					15		
Val	Ile	Val	Ala	Thr	Ala	Ala	Val	Pro	Ser	Pro	Ala	Ser	Gly	Arg	Glu	
			20					25					30			
His	Asp	Gly	Gly	Tyr	Ala	Ala	Leu	Ile	Arg	Arg	Ala	Ser	Tyr	Gly	Val	
		35					40					45				
Pro	His	Ile	Thr	Ala	Asp	Asp	Phe	Gly	Ser	Leu	Gly	Phe	Gly	Val	Gly	
		50				55					60					
Tyr	Val	Gln	Ala	Glu	Asp	Asn	Ile	Cys	Val	Ile	Ala	Glu	Ser	Val	Val	
65					70					75					80	
Thr	Ala	Asn	Gly	Glu	Arg	Ser	Arg	Trp	Phe	Gly	Ala	Thr	Gly	Pro	Asp	
			85					90						95		
Asp	Ala	Asp	Val	Arg	Ser	Asp	Leu	Phe	His	Arg	Lys	Ala	Ile	Asp	Asp	
			100					105					110			
Arg	Val	Ala	Glu	Arg	Leu	Leu	Glu	Gly	Pro	Arg	Asp	Gly	Val	Arg	Ala	
		115					120					125				
Pro	Ser	Asp	Asp	Val	Arg	Asp	Gln	Met	Arg	Gly	Phe	Val	Ala	Gly	Tyr	
		130				135					140					
Asn	His	Phe	Leu	Arg	Arg	Thr	Gly	Val	His	Arg	Leu	Thr	Asp	Pro	Ala	
145					150					155					160	
Cys	Arg	Gly	Lys	Ala	Trp	Val	Arg	Pro	Leu	Ser	Glu	Ile	Asp	Leu	Trp	
			165						170					175		

-continued

Arg Thr Ser Trp Asp Ser Met Val Arg Ala Gly Ser Gly Ala Leu Leu
 180 185 190

Asp Gly Ile Val Ala Ala Thr Pro Pro Thr Ala Ala Gly Pro Ala Ser
 195 200 205

Ala Pro Glu Ala Pro Asp Ala Ala Ala Ile Ala Ala Ala Leu Asp Gly
 210 215 220

Thr Ser Ala Gly Ile Gly Ser Asn Ala Tyr Gly Leu Gly Ala Gln Ala
 225 230 235 240

Thr Val Asn Gly Ser Gly Met Val Leu Ala Asn Pro His Phe Pro Trp
 245 250 255

Gln Gly Ala Ala Arg Phe Tyr Arg Met His Leu Lys Val Pro Gly Arg
 260 265 270

Tyr Asp Val Glu Gly Ala Ala Leu Ile Gly Asp Pro Ile Ile Gly Ile
 275 280 285

Gly His Asn Arg Thr Val Ala Trp Ser His Thr Val Ser Thr Ala Arg
 290 295 300

Arg Phe Val Trp His Arg Leu Ser Leu Val Pro Gly Asp Pro Thr Ser
 305 310 315 320

Tyr Tyr Val Asp Gly Arg Pro Glu Arg Met Arg Ala Arg Thr Val Thr
 325 330 335

Val Gln Thr Gly Ser Gly Pro Val Ser Arg Thr Phe His Asp Thr Arg
 340 345 350

Tyr Gly Pro Val Ala Val Met Pro Gly Thr Phe Asp Trp Thr Pro Ala
 355 360 365

Thr Ala Tyr Ala Ile Thr Asp Val Asn Ala Gly Asn Asn Arg Ala Phe
 370 375 380

Asp Gly Trp Leu Arg Met Gly Gln Ala Lys Asp Val Arg Ala Leu Lys
 385 390 395 400

Ala Val Leu Asp Arg His Gln Phe Leu Pro Trp Val Asn Val Ile Ala
 405 410 415

Ala Asp Ala Arg Gly Glu Ala Leu Tyr Gly Asp His Ser Val Val Pro
 420 425 430

Arg Val Thr Gly Ala Leu Ala Ala Ala Cys Ile Pro Ala Pro Phe Gln
 435 440 445

Pro Leu Tyr Ala Ser Ser Gly Gln Ala Val Leu Asp Gly Ser Arg Ser
 450 455 460

Asp Cys Ala Leu Gly Ala Asp Pro Asp Ala Ala Val Pro Gly Ile Leu
 465 470 475 480

Gly Pro Ala Ser Leu Pro Val Arg Phe Arg Asp Asp Tyr Val Thr Asn
 485 490 495

Ser Asn Asp Ser His Trp Leu Ala Ser Pro Ala Ala Pro Leu Glu Gly
 500 505 510

Phe Pro Arg Ile Leu Gly Asn Glu Arg Thr Pro Arg Ser Leu Arg Thr
 515 520 525

Arg Leu Gly Leu Asp Gln Ile Gln Gln Arg Leu Ala Gly Thr Asp Gly
 530 535 540

Leu Pro Gly Lys Gly Phe Thr Thr Ala Arg Leu Trp Gln Val Met Phe
 545 550 555 560

Gly Asn Arg Met His Gly Ala Glu Leu Ala Arg Asp Asp Leu Val Ala
 565 570 575

Leu Cys Arg Arg Gln Pro Thr Ala Thr Ala Ser Asn Gly Ala Ile Val
 580 585 590

-continued

Asp Leu Thr Ala Ala Cys Thr Ala Leu Ser Arg Phe Asp Glu Arg Ala
 595 600 605
 Asp Leu Asp Ser Arg Gly Ala His Leu Phe Thr Glu Phe Ala Leu Ala
 610 615 620
 Gly Gly Ile Arg Phe Ala Asp Thr Phe Glu Val Thr Asp Pro Val Arg
 625 630 635 640
 Thr Pro Arg Arg Leu Asn Thr Thr Asp Pro Arg Val Arg Thr Ala Leu
 645 650 655
 Ala Asp Ala Val Gln Arg Leu Ala Gly Ile Pro Leu Asp Ala Lys Leu
 660 665 670
 Gly Asp Ile His Thr Asp Ser Arg Gly Glu Arg Arg Ile Pro Ile His
 675 680 685
 Gly Gly Arg Gly Glu Ala Gly Thr Phe Asn Val Ile Thr Asn Pro Leu
 690 695 700
 Val Pro Gly Val Gly Tyr Pro Gln Val Val His Gly Thr Ser Phe Val
 705 710 715 720
 Met Ala Val Glu Leu Gly Pro His Gly Pro Ser Gly Arg Gln Ile Leu
 725 730 735
 Thr Tyr Ala Gln Ser Thr Asn Pro Asn Ser Pro Trp Tyr Ala Asp Gln
 740 745 750
 Thr Val Leu Tyr Ser Arg Lys Gly Trp Asp Thr Ile Lys Tyr Thr Glu
 755 760 765
 Ala Gln Ile Ala Ala Asp Pro Asn Leu Arg Val Tyr Arg Val Ala Gln
 770 775 780
 Arg Gly Arg
 785

<210> SEQ ID NO 9
 <211> LENGTH: 21
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:forward
 primer corresponding to the vector sequence of pGFP
 plasmid (Aequorea victoria green fluorescent
 protein)

<400> SEQUENCE: 9

ccgactggaa agcgggcagt g

21

<210> SEQ ID NO 10
 <211> LENGTH: 22
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence:reverse
 primer corresponding to the vector sequence of pGFP
 plasmid (Aequorea victoria green fluorescent
 protein)

<400> SEQUENCE: 10

cggggctggc ttaactatgc gg

22

<210> SEQ ID NO 11
 <211> LENGTH: 4
 <212> TYPE: PRT
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <221> NAME/KEY: MOD_RES
 <222> LOCATION: (1)
 <223> OTHER INFORMATION: Xaa = succinyl-Ala
 <220> FEATURE:

-continued

<221> NAME/KEY: MOD_RES
 <222> LOCATION: (4)
 <223> OTHER INFORMATION: Xaa = Phe-p-nitroanilide
 <220> FEATURE:
 <223> OTHER INFORMATION: Description of Artificial Sequence: Bacillus
 subtilis subtilisin E thermostability assay
 substrate

<400> SEQUENCE: 11

Xaa Ala Pro Xaa
 1

What is claimed is:

1. A method for evolving a polynucleotide toward acquisition of a desired property, comprising

(a) incubating a population of parental polynucleotide variants comprising allelic or species variants of a gene, or mutants created by error-prone PCR, under conditions to generate annealed polynucleotides comprising heteroduplexes;

(b) exposing the heteroduplexes to enzymes of a DNA repair system to convert the heteroduplexes to parental polynucleotide variants or recombined polynucleotide variants;

(c) screening or selecting the recombined polynucleotide variants for the desired property.

2. The method of claim **1**, wherein the heteroduplexes are exposed to the enzymes of the DNA repair system *in vitro*.

3. The method of claim **2**, wherein the enzymes of the DNA repair system are provided as cellular extracts.

4. The method of claim **1**, further comprising introducing the heteroduplexes into cells, whereby the heteroduplexes are exposed to the enzymes of the DNA repair system of the cells *in vivo*.

5. The method of claim **4**, wherein the annealed polynucleotides further comprise homoduplexes and the introducing step selects for transformed cells comprising the heteroduplexes relative to transformed cells comprising homoduplexes.

6. The method of claim **4**, wherein a first polynucleotide variant is provided as a component of a first vector, and a second polynucleotide variant is provided as a component of a second vector, and the method further comprises converting the first and second vectors to linearized forms in which the first and second polynucleotide variants occur at opposite ends, whereby in the incubating step single-stranded forms of the first linearized vector reanneal with each other to form linear first vector, single-stranded forms of the second linearized vector reanneal with each other to form linear second vector, and single-stranded linearized forms of the first and second vectors anneal with each to form a circular heteroduplex bearing a nick in each strand, and the introducing step selects for transformed cells comprising the circular heteroduplexes relative to the linear first and second vector.

7. The method of claim **6**, wherein the first and second vectors are converted to linearized forms by PCR.

8. The method of claim **6**, wherein the first and second vectors are converted to linearized forms by digestion with first and second restriction enzymes.

9. The method of claim **1-3** wherein the population of polynucleotide variants are provided in double stranded form, and the method further comprising converting the double stranded polynucleotides to single stranded polynucleotides before the annealing step.

10. The method of claim **9**, wherein the converting step comprises:

conducting asymmetric amplification of the first and second double stranded polynucleotide variants to amplify a strand of the first polynucleotide variant, and a strand of the second polynucleotide variant, whereby the strands anneal in the incubating step to form a heteroduplex.

11. The method of claim **10**, wherein the first and second double-stranded polynucleotides variants are provided in vector-free form, and the method further comprises incorporating the heteroduplex into a vector.

12. The method of claim **4** wherein the population of polynucleotides comprises first and second polynucleotides provided in double stranded form, and the method further comprises incorporating the first and second polynucleotides as components of first and second vectors, whereby the first and second polynucleotides occupy opposite ends of the first and second vectors, whereby in the incubating step single-stranded forms of the first linearized vector reanneal with each other to form linear first vector, single-stranded forms of the second linearized vector reanneal with each other to form linear second vector, and single-stranded linearized forms of the first and second vectors anneal with each to form a circular heteroduplex bearing a nick in each strand, and the introducing step selects for transformed cells comprising the circular heteroduplexes relative to the linear first and second vector.

13. The method of claim **4**, further comprising sealing nicks in the heteroduplexes to form covalently-closed circular heteroduplexes before the introducing step.

14. The method of claim **11**, wherein the first and second polynucleotides are obtained from chromosomal DNA.

15. The method of claim **1**, further comprising repeating steps (a)-(c) whereby the incubating step in a subsequent cycle is performed on recombinant variants from a previous cycle.

16. The method of claim **1**, wherein the polynucleotide variants encode a polypeptide.

17. The method of claim **1**, wherein the population of polynucleotide variants comprises at least 20 variants.

18. The method of claim **1**, wherein the population of polynucleotide variants are at least 10 kb in length.

19. The method of claim **1**, wherein the population of polynucleotide variants comprises natural variants.

20. The method of claim **1**, wherein the population of polynucleotides comprises variants generated by mutagenic PCR.

21. The method of claim **4**, wherein the cells are bacterial cells.

22. The method of claim **1**, further comprising at least partially demethylating the population of variant polynucleotides.

23. The method of claim 22, whether the at least partially demethylating step is performed by PCR amplification of the population of variant polynucleotides.

24. The method of claim 22, wherein the at least partially demethylating step is performed by amplification of the population of variant polynucleotides in host cells.

25. The method of claim 24, wherein the host cells are defective in a gene encoding a methylase enzyme.

26. The method of claim 1 wherein the population of variant polynucleotide variants comprises at least 5 polynucleotides having at least 90% sequence identity with one another.

27. The method of claim 1, further comprising isolating a screened recombinant variant.

28. The method of claim 27, further comprising expressing a screened recombinant variant to produce a recombinant protein.

29. The method of claim 28 further comprising formulating the recombinant protein with a carrier to form a pharmaceutical composition.

30. The method of claim 1, wherein the polynucleotide variants encode enzymes selected from the group consisting of proteases, lipases, amylases, cutinases, cellulases, amylases, oxidases, peroxidases and phytases.

31. The method of claim 1, wherein the polynucleotide variants encode a polypeptide selected from the group consisting of insulin, ACTH, glucagon, somatostatin, somatotro-

pin, thymosin, parathyroid hormone, pigmentary hormones, somatomedin, erythropoietin, luteinizing hormone chorionic gonadotropin, hyperthymic releasing factors, antidiuretic hormones, thyroid stimulating hormone, relaxin, interferon, thrombopoietin (TPO), and prolactin.

32. The method of claim 1, wherein the polynucleotide variants encode a plurality of enzymes forming a metabolic pathway.

33. The method of claim 1, wherein the polynucleotide variants are in concatemeric form.

34. A method for evolving a polynucleotide toward acquisition of a desired property, comprising

- (a) incubating a population of parental polynucleotide variants, under conditions to generate annealed polynucleotides comprising heteroduplexes;
- (b) exposing the heteroduplexes to enzymes of a DNA repair system to convert the heteroduplexes to parental polynucleotide variants or recombined polynucleotide variants;
- (c) screening or selecting the recombined polynucleotide variants for the desired property; and
- (d) recombining screened or selected recombined polynucleotides with each other or other substrates, followed by screening or selection.

* * * * *