

US007906638B2

(12) **United States Patent**
Kipps et al.

(10) **Patent No.:** **US 7,906,638 B2**
(45) **Date of Patent:** **Mar. 15, 2011**

(54) **CHIMERIC NUCLEIC ACIDS ENCODING
POLYPEPTIDES COMPRISING CD70 AND
FAS LIGAND DOMAINS**

(75) Inventors: **Thomas J. Kipps**, Solana Beach, CA
(US); **Sanjai Sharma**, La Jolla, CA
(US); **Mark Cantwell**, La Jolla, CA
(US)

(73) Assignee: **Regents of the University of California**,
Oakland, CA (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 217 days.

(21) Appl. No.: **12/371,188**

(22) Filed: **Feb. 13, 2009**

(65) **Prior Publication Data**

US 2010/0267129 A1 Oct. 21, 2010

Related U.S. Application Data

(63) Continuation of application No. 11/015,117, filed on
Dec. 17, 2004, now Pat. No. 7,524,944, which is a
continuation of application No. 08/982,272, filed on
Dec. 1, 1997, now Pat. No. 7,070,771.

(60) Provisional application No. 60/032,145, filed on Dec.
9, 1996, now abandoned.

(51) **Int. Cl.**
C12N 15/62 (2006.01)
C12N 15/12 (2006.01)
C12N 15/00 (2006.01)
C12N 15/63 (2006.01)

(52) **U.S. Cl.** **536/23.4**; 536/23.1; 536/23.5;
435/252.3; 435/320.1; 435/455

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,990,455	A	2/1991	Yamagishi et al.
5,422,104	A	6/1995	Fiers et al.
5,480,981	A	1/1996	Goodwin et al.
5,486,463	A	1/1996	Lesslauer et al.
5,519,119	A	5/1996	Yamada et al.
5,540,926	A	7/1996	Aruffo et al.
5,565,321	A	10/1996	Spriggs et al.
5,573,924	A	11/1996	Beckmann et al.
5,606,023	A	2/1997	Chen et al.
5,817,516	A	10/1998	Kehry et al.
5,861,310	A	1/1999	Freeman et al.
6,017,527	A	1/2000	Maraskovsky et al.
6,106,832	A	8/2000	Spriggs et al.
7,070,771	B1 *	7/2006	Kipps et al. 424/93.21
7,495,090	B2 *	2/2009	Prussak et al. 536/23.4
7,524,944	B2 *	4/2009	Kipps et al. 536/23.4
7,786,282	B2 *	8/2010	Prussak et al. 536/23.4
2004/0209295	A1	10/2004	Schwabe et al.

FOREIGN PATENT DOCUMENTS

EP	0 317 641	A1	5/1989
EP	0 585 943	A2	3/1993
EP	0 675 200	A1	10/1995
EP	1 016 721	A1	7/2000
WO	WO 91/02540	A1	3/1991
WO	WO 93/08207	A1	4/1993
WO	WO 93/19777		10/1993
WO	WO 93/24135		12/1993
WO	WO 94/04570		3/1994
WO	WO 94/04680	A1	3/1994
WO	WO 94/17196	A1	8/1994
WO	WO 95/14487	A1	6/1995
WO	WO 95/17202		6/1995
WO	WO 95/18819	A1	7/1995
WO	WO 95/32627	A1	12/1995
WO	WO 96/03141		2/1996
WO	WO 96/14876	A1	5/1996
WO	WO 96/18413	A1	6/1996
WO	WO 96/22370	A1	7/1996
WO	WO 98/21232		5/1998

OTHER PUBLICATIONS

Addison, Christina L. et al., "Intratumoral injection of an adenovirus expressing interleukin 2 induces regression and immunity in a murine breast cancer model," *Proc. Natl. Acad. Sci. U.S.A.*, 92:8522-8526 (1995).

Alderson et al., "CD40 expression by human monocytes: Regulation by cytokines and activation of monocytes by the ligand for CD40", *J. Exp. Med.*, 178: 669-674 (1993).

Ali, Munaf et al., "The use of DNA viruses as vectors for gene therapy", *Gene Therapy*, 1:367-384 (1994).

Ali, Stuart Alvaro et al., "PCR-Ligation-PCR Mutagenesis: A Protocol for Creating Gene Fusions and Mutations", *BioTechniques*, 18:746-750 (1995).

Armitage, Richard J. et al., "Molecular and biological characterization of a murine ligand for CD40", *Nature*, 357:80-82 (1992).

Banchereau, Jacques et al., "Long-Term Human B Cell Lines Dependent on Interleukin-4 and Antibody to CD40," *Science*, 251:70-72 (1991).

Berman, Joan W. et al., "Gene transfer in lymphoid cells: Expression of the Thy-1.2 antigen by Thy-1.1 BW5147 lymphoma cells transfected with unfractionated cellular DNA", *Proc. Natl. Acad. Sci. USA*, 81:7176-7179 (1984).

Blieden, Timothy M. et al., "Class-I MHC Expression in the Mouse Lung Carcinoma, Line 1: A Model for Class-I Inducible Tumors", *Int. J. Cancer Supp.*, 6:82-89 (1991).

Boles, Eckhard et al., "A rapid and highly efficient method for PCR-based site-directed mutagenesis using only one new primer", *Curr. Genet.*, 28:197-198 (1995).

(Continued)

Primary Examiner — Phillip Gambel

(74) *Attorney, Agent, or Firm* — DLA Piper LLP (US)

(57) **ABSTRACT**

This invention relates to genes which encode accessory molecule ligands and their use for immunomodulation, vaccination and treatments of various human diseases, including malignancies and autoimmune diseases. This invention also describes the use of accessory molecule ligands which are made up of various domains and subdomain portions of molecules derived from the tumor necrosis factor family. The chimeric molecules of this invention contain unique properties which lead to the stabilization of their activities and thus greater usefulness in the treatment of diseases. Vectors for expressing genes which encode the accessory molecule ligands of this invention are also disclosed.


24 Claims, 30 Drawing Sheets

OTHER PUBLICATIONS

- Boris-Lawrie, Kathleen A. et al., "Recent advances in retrovirus vector technology", *Current Opinion in Genetics and Development*, 3:102-109 (1993).
- Brody, Steven L. et al., "Adenovirus-mediated in Vivo Gene Transfer", *Ann. N. Y. Acad. Sci.*, 716:90-103 (1994).
- Cadwell, R. Craig et al., "Randomization of Genes by PCR Mutagenesis", *PCR Methods and Applications*, 2:28-33 (1992).
- Cantwell, M.J. et al., "CD95 and FAS-ligand expression and apoptosis in rheumatoid arthritis", *Arthritis and Rheumatism*, vol. 39, No. 9, Suppl., Sep. 1996, p. 287.
- Cantwell, Mark et al., "Acquired CD40-ligand deficiency in chronic lymphocytic leukemia", *Nature Medicine*, 3:984-989 (1997).
- Cantwell, Mark J. et al., "Adenovirus Vector Infection of Chronic Lymphocytic Leukemia B Cells", *Blood*, 88:4676-4683 (1996).
- Carter, Barrie J., "Adeno-associated virus vectors", *Current Opinion in Biotechnology*, 3:533-539 (1992).
- Clark, Edward A. et al., "Activation of human B cells mediated through two distinct cell surface differentiation antigens, Bp35 and Bp50", *Proc. Natl. Acad. Sci. U.S.A.*, 83:4494-4498 (1986).
- Cooper, Mark J., "Noninfectious Gene Transfer and Expression Systems for Cancer Gene Therapy", *Seminars in Oncology*, 23:172-187 (1996).
- Cosman, David et al., "Cloning, sequence and expression of human interleukin-2 receptor", *Nature*, 312:768-771 (1984).
- Danko, Istvan et al., "Direct gene transfer into muscle", *Vaccine*, 12:1499-1502 (1994).
- Davis, Heather L. et al., "Direct Gene Transfer into Skeletal Muscle In Vivo: Factors Affecting Efficiency of Transfer and Stability of Expression", *Human Gene Therapy*, 4:151-159 (1993).
- Deans, Robert J. et al., "Expression of an immunoglobulin heavy chain gene transfected into lymphocytes", *Proc. Natl. Acad. Sci. USA*, 81:1292-1296 (1984).
- DeMatteo, Ronald P. et al., "Gene Transfer to the Thymus", *Annals of Surgery*, 222:229-242 (1995).
- Dilloo, Dagmar et al., "CD40 Ligand Induces an Antileukemia Immune Response In Vivo", *Blood*, 90:1927-1933 (1997).
- Evan, Christopher et al., "Clinical Trial to Assess the Safety, Feasibility, and Efficacy of Transferring a Potentially Anti-Arthritic Cytokine Gene to Human Joints with Rheumatoid Arthritis", *Human Gene Therapy*, 7:1261-1280 (1996).
- Fanslow, William C. et al., "Structural characteristics of CD40 ligand that determine biological function", *Seminars in Immunology*, 6:267-278 (1994).
- Felgner, Philip L. et al., "Improved Cationic Lipid Formulations for In Vivo Gene Therapy", *Ann. N. Y. Acad. Sci.*, 772:126-139 (1995).
- Fisher, Lisa J. et al., "In vivo and ex vivo gene transfer to the brain", *Current Opinion in Neurobiology*, 4:735-741 (1994).
- Flotte, T. R. et al., "Adeno-associated virus vectors for gene therapy", *Gene Therapy*, 2:357-362 (1995).
- Furth, Priscilla A. et al., "Gene Transfer into Mammalian Cells by Jet Injection", *Hybridoma*, 14:149-152 (1995).
- Galle, Peter R. et al., "Involvement of the CD95 (APO-1/Fas) Receptor and Ligand in Liver Damage", *J. Exp. Med.*, 182:1223-1230 (1995).
- Glorioso, J. C. et al., "Development and application of herpes simplex virus vectors for human gene therapy", *Annu. Rev. Microbiol.*, 49:675-710 (1995).
- Goldspiel, Barry R. et al., "Human gene therapy", *Clinical Pharmacology*, 12:488-505 (1993).
- Graham, Frank L. et al., "Manipulation of Adenovirus Vectors", *Methods in Molecular Biology*, 7(11):109-128 (1991).
- Hengge, Ulrich R. et al., "Expression of Naked DNA in Human, Pig, and Mouse Skin", *Journal of Clinical Investigation*, 97:2911-2916 (1996).
- Henkel, Thomas et al., "Functional Analysis of Mutated cDNA Clones by Direct Use of PCR Products in in Vitro Transcription/Translation Reactions", *Analytical Biochemistry*, 214:351-352 (1993).
- Hollenbaugh, Diane et al., "The human T cell antigen gp39, a member of the TNF gene family, is a ligand for the CD40 receptor: expression of a soluble form of gp39 with B cell co-stimulatory activity", *The EMBO Journal*, 11:4313-4321 (1992).
- Horton, Robert M., "PCR-mediated Recombination and Mutagenesis", *Molecular Biotechnology*, 3:93-99 (1995).
- International Search Report for PCT Application No. PCT/US97/22740.
- Jolly, Douglas, "Viral vector systems for gene therapy", *Cancer Gene Therapy*, 1:51-64 (1994).
- Kass-Eisler, Alyson et al., "Prospects for Gene Therapy with Direct Injection of Polynucleotides", *Ann. N. Y. Acad. Sci.*, 772:232-240 (1995).
- Kato et al., Adenovirus-mediated gene transfer of CD40-ligand induces autologous immune recognition of chronic lymphocytic leukemia B cells, *Blood*, 90: Abstract No. 1157, 1997.
- Kipps, Thomas J. et al., "New developments in flow cytometric analyses of lymphocyte markers", *Laboratory Immunology II*, 12:237-275 (1992).
- Koc, Omer N. et al., "Transfer of Drug Resistance Genes Into Hematopoietic Progenitors to Improve Chemotherapy Tolerance", *Seminars in Oncology*, 23:46-65 (1996).
- Kohn, Donald B., "The current status of gene therapy using hematopoietic stem cell", *Current Opinion in Pediatrics*, 7:56-63 (1995).
- Kouskoff, Valerie et al., "Organ-Specific Disease Provoked by Systemic Autoimmunity", *Cell*, 87:811-822 (1997).
- Kunkel, Thomas A. et al., "Rapid and Efficient Site-Specific Mutagenesis without Phenotypic Selection", *Methods in Enzymology*, 154:367-382 (1987).
- Kunkel, Thomas A., "Rapid and efficient site-specific mutagenesis without phenotypic selection", *Proc. Natl. Acad. Sci. USA*, 82:488-492 (1985).
- Lederman et al., "T-BAM/CD40-L on helper T lymphocytes augments lymphokine-induced B cell Ig isotype switch recombination and rescues B cells from programmed cell death", *Journal of Immunology*, 152:2163-2171, 1994.
- Lu, Li et al., "Stem cells from bone marrow, umbilical cord blood and peripheral blood for clinical application: current status and future application", *Critical Reviews in Oncology/Hematology*, 22:61-78 (1996).
- Majumder, Kumud et al., "Background-minimized Cassette Mutagenesis by PCR Using Cassette-specific Selection Markers: A Useful General Approach for Studying Structure-Function Relationships of Multisubstrate Enzymes", *PCR Methods and Applications*, 4:212-218 (1995).
- Morrison, Hilary G. et al., "A PCR-Based Strategy for Extensive Mutagenesis of a Target DNA Sequence", *BioTechniques*, 14:454-457 (1993).
- Nadler, Lee M., "The Malignant Lymphomas", *Harrison's Principles of Internal Medicine*, Wilson et al., eds., McGraw-Hill, New York, Chapter 302, pp. 1599-1612.
- Nagase, H. et al., "Human Matrix Metalloproteinase Specificity Studies Using Collagen Sequence-Based Synthetic Peptides", *Biopolymers (Peptide Science)*, 40:399-416 (1996).
- Okayama, Hiroto and Paul Berg, "A cDNA Cloning Vector That Permits Expression of cDNA Inserts in Mammalian Cells", *Molecular and Cellular Biology*, 3:280-289 (1983).
- Peitsch, Manuel C. et al., "A 3-D model for the CD40 ligand predicts that it is a compact trimer similar to the tumor necrosis factors", *International Immunology*, 5:233-238 (1993).
- Prentice, Howard et al., "Ischemic/Reperfused Myocardium Can Express Recombinant Protein Following Direct DNA or Retroviral Injection", *J. Mol. Cell Cardiol.*, 28:133-140 (1996).
- Randrianarison-Jewtoukoff, Voahangy et al., "Recombinant Adenoviruses as Vaccines", *Biologicals*, 23:145-157 (1995).
- Ranheim, Eric A. et al., "Activated T Cells Induce Expression of B7/BB1 on Normal or Leukemic B Cells through a CD40-dependent Signal", *J. Exp. Med.*, 177:925-935 (1993).
- Raper, Steven E. et al., "Safety and Feasibility of Liver-Directed Ex Vivo Gene Therapy for Homozygous Familial Hypercholesterolemia", *Annals of Surgery*, 223:116-126 (1996).
- Rassenti, Laura Z. et al., "Lack of Allelic Exclusion in B Cell Chronic Lymphocytic Leukemia", *J. Exp. Med.*, 185:1435-1445 (1992).
- Raz, Eyal et al., "Systemic immunological effects of cytokine genes injected into skeletal muscle", *Proc. Natl. Acad. Sci. U.S.A.*, 90:4523-4527 (1993).

- Russell, S. J., "Replicating Vectors for Gene Therapy of Cancer: Risks, Limitations and Prospects", *European Journal of Cancer*, 30A:1165-1171 (1994).
- Russell, Stephen J., "Replicating vectors for cancer therapy: a question of strategy", *Seminars in Cancer Biology*, 5:437-443 (1994).
- Sambrook, J. et al., "Standard Protocol for Calcium Phosphate-mediated Transfection of Adherent Cells", *Molecular Cloning. A Laboratory Manual*, 2d edition, Chapter 16:33-37 (1989).
- Sato, Ken et al., "An aggressive nasal lymphoma accompanied by high levels of soluble Fas ligand", *British Journal of Haematology*, 94:379-382 (1996).
- Schultze, Joachim L. et al., "Autologous Tumor Infiltrating T Cells Cytotoxic for Follicular Lymphoma Cells Can Be Expanded In Vitro", *Blood*, 89:3806-3816 (1997).
- Shaughnessy, Elizabeth et al., "Parvoviral Vectors for the Gene Therapy of Cancer", *Seminars in Oncology*, 23:159-171 (1996).
- Skolnick and Fetrow, "From genes to protein structure and function: novel applications of computational approaches in the genomic era", *Trends in Biotechnology*, 18:34-39 (2000).
- Smith, K. T. et al., "Gene delivery systems for use in gene therapy: an overview of quality assurance and safety issues", *Gene Therapy*, 3:190-200 (1996).
- Smith, Matthew M. et al., "Rapid Identification of Highly Active and Selective Substrates for Stromelysin and Matrilysin Using Bacteriophage Peptide Display Libraries", *The Journal of Biological Chemistry*, 270:6440-6449 (1995).
- Soubrane, C. et al., "Direct Gene Transfer of a Plasmid Carrying the Herpes Simplex Virus-Thymidine Kinase Gene (HSV-TK) in Transplanted Murine Melanoma: In Vivo Study", *European Journal of Cancer*, 32A:691-695 (1996).
- Spessot, Robert, "Cloning of the Herpes Simplex Virus ICP4 Gene in an Adenovirus Vector: Effects on Adenovirus Gene Expression and Replication", *Virology*, 168:378-387 (1989).
- Srivastava, Arun, "Parvovirus-Based Vectors for Human Gene Therapy", *Blood Cells*, 20:531-538 (1994).
- Stappert, Jorg et al., "A PCR method for introducing mutations into cloned DNA by joining an internal primer to a tagged flanking primer", *Nucleic Acids Research*, 20:624 (1992).
- Sugaya, Susumu et al., "Inhibition of Tumor Growth by Direct Intratumoral Gene Transfer of Herpes Simplex Virus Thymidine Kinase Gene with DNA-Liposome Complexes", *Human Gene Therapy*, 7:223-230 (1996).
- Tesselaar, Kiki et al., "Characterization of Murine CD70, the Ligand of the TNF Receptor Family Member CD27", *The Journal of Immunology*, 159:4959-4965 (1997).
- Tessier, Daniel C. et al., "PCR-Assisted Large Insertion/Deletion Mutagenesis", *BioTechniques*, 15:498-501 (1993).
- Thomas, J. Alero et al., "Epstein-Barr Virus-Associated Lymphoproliferative Disorders in Immunocompromised Individuals", *Advances in Cancer Research*, Woude et al., eds., Academic Press, Inc., 57:329-380 (1991).
- Tolstoshev, Paul, "Gene therapy, concepts, current trials and future directions", *Annu. Rev. Pharmacol. Toxicol.*, 33:573-596 (1993).
- Tracey, Kevin J. et al., "Tumor Necrosis Factor: A Pleiotropic Cytokine and Therapeutic [sic] Target", *Annu. Rev. Med.*, 45:491-503 (1994).
- Vallejo, Abbe N. et al., "In Vitro Synthesis of Novel Genes: Mutagenesis and Recombination by PCR", *PCR Methods and Applications*, 4:S123-S130 (1994).
- van Oers, M. H. J. et al., "Expression and Release of CD27 in Human B-Cell Malignancies", *Blood*, 82:3430-3436 (1993).
- Villardaga, J. P. et al., "Improved PCR Method for High-Efficiency Site-Directed Mutagenesis Using Class 2S Restriction Enzymes", *BioTechniques*, 18:604-606 (1995).
- Vile, R. G. et al., "Retroviruses as vectors", *British Medical Bulletin*, 51:12-30 (1995).
- Vile, R.G. et al., "Targeting of cytokine gene expression to malignant melanoma cells using tissue specific promoter sequences", *Annals of Oncology*, 5 Suppl. 4:S59-S65 (1994).
- Wierda et al., "Infection of B-cell lymphoma with adenovirus vector encoding CD40-ligand (CD154) induces phenotypic changes that allow for autologous immune recognition", *Blood*, 90: Abstract No. 2280, 1997.
- Wiley, James A. et al., "Exogenous CD40 Ligand Induces a Pulmonary Inflammation Response", *Journal of Immunology*, 158:2932-2938 (1997).
- Woll, P. J. et al., "Gene therapy for lung cancer", *Annals of Oncology*, 6 Suppl. 1:S73-S77 (1995).
- Yee, Jiing-Kuan et al., "Generation of High-Tier Pseudotyped Retroviral Vectors with Very Broad Host Range", *Methods in Cell Biology*, Chapter 5, 43:99-112 (1994).
- Yellin et al., "T lymphocyte T Cell-B Cell-activating molecule/CD40-L molecules induce normal B Cells or chronic lymphocytic leukemia B cells to express CD80 (B7/BB-1) and enhance their costimulatory activity", *J. Immun.*, 153:666-674 (1994).
- Yovandich, Jason et al., "Gene Transfer to Synovial Cells by Intra-Articular Administration of Plasmid DNA", *Human Gene Therapy*, 6:603-610 (1995).
- Zhang, Haidi et al., "Amelioration of Collagen-induced Arthritis by CD95 (Apo-1/Fas)-ligand Gene Transfer", *J. Clin. Invest.*, 100:1951-1957 (1997).
- Ngo, Thomas J. et al., "Computational Complexity, Protein Structure Prediction, and the Levinthal Paradox," *The Protein Folding Problem and Tertiary Structure Prediction*, Mertz et al. Editors, Birkhäuser Boston, pp. 491-495 (1994).

* cited by examiner



DOMAINS: I - Cytoplasmic Domain; II - Transmembrane Domain; III - Proximal Extracellular Domain; IV - Distal Extracellular Domain (putative soluble form)

FIG. 1






FIG. 2





Allogenic T cell response to CLL cells transfected with adeno-mCD40L




FIG. 5

Production of IFN γ by allogenic T lymphocytes
stimulated with CLL B cells




FIG. 6




FIG. 7




FIG. 8




FIG. 9




FIG. 10




FIG. 11A






FIG. 11B












FIG. 14C






FIG. 16A




FIG. 16B




FIG. 17A




FIG. 17B




FIG. 18




FIG. 19




FIG. 20A




FIG. 20B




FIG. 21B

SK-Mes-1 Squamous Cell (CD40++)




FIG. 21D




FIG. 21A

NCI 460 Large Cell CA (CD40+)




FIG. 21C




FIG. 22A




FIG. 22B


RA SYNOVIAL FLUID AND PLASMA INHIBITION
OF FAS-LIGAND

FIG. 23

Gene Therapy of Leukemia




FIG. 24

1 MQQPFNYPYPQIYWVDSSASSPWAPPGTVLPCPTSVPRRPGQRRPPPPPP 50
|||||
1 MQQPFNYPYPQIYWVDSSASSPWAPPGTVLPCPTSVPRRPGQRRPPPPPP 50
51 PPPLPPPPPPPPPLPPLPLPPLKKRGNHSTGLCLLMFFMVLVALVGLGLG 100
|||||
51 PPPLPPPPPPPPPLPPLPLPPLKKRGNHSTGLCLLMFFMVLVALVGLGLG 100
101 MFQLFHLOKELAELRESTSOMHTASSLEKQIGHPSPPPEKKELRKVAHLT 150
||||| |. | .-||:|||||
101 MFQLFHLOKELAELREFTNQSLKVSSFEKQIGHPSPPPEKKELRKVAHLT 150
151 GKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSKVYFRGQ 200
|||||
151 GKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSKVYFRGQ 200
201 SCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGAVERN 250
|||||
201 SCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGAVERN 250
251 LTSADHLYVNVSELSLVNFEESQTFFGLYKL 281
|||||
251 LTSADHLYVNVSELSLVNFEESQTFFGLYKL 281

FIG. 25

1 MQQPFNYPYPQIYWVDSSASSPWAPP GTVLP CPTSVPRRPGQRRPPPPPP 50
|||||
1 MQQPFNYPYPQIYWVDSSASSPWAPP GTVLP CPTSVPRRPGQRRPPPPPP 50
51 PPPLPPPPPPPPPLPPLPLPPLKKRGNHSTGLCLLMFFMVLVALVGLGLG 100
|||||
51 PPPLPPPPPPPPPLPPLPLPPLKKRGNHSTGLCLLMFFMVLVALVGLGLG 100
101 MFQLFHLOKELAELRESTSOMHTASSLEKQIGHPSPPPEKKELRKVAHLT 150
|||||:
101 MFQLFR.....FAQAIGHPSPPPEKKELRKVAHLT 130
151 GKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSKVYFRGQ 200
|||||
131 GKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSKVYFRGQ 180
201 SCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGA VFN 250
|||||
181 SCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGA VFN 230
251 LTSADHLYVNVSELSLVNFEESQTFFGLYKL 281
|||||
231 LTSADHLYVNVSELSLVNFEESQTFFGLYKL 261

FIG. 26

1 MQQPFNYPYPQIYWVDSSASSPWAPPGTVLPCPTSVPRRPGQRRPPPPPP 50
|||||
1 MQQPFNYPYPQIYWVDSSASSPWAPPGTVLPCPTSVPRRPGQRRPPPPPP 50
51 PPPLPPPPPPPPPLPPLPLPPLKKRGNHSTGLCLLMFFMVLVALVGLGLG 100
|||||
51 PPPLPPPPPPPPPLPPLPLPPLKKRGNHSTGLCLLMFFMVLVALVGLGLG 100
101 MFQLFHLOKELAELRESTSOMHTASSLEKQOIGHPSPPPEKKELRKVAHLT 150
||||| : : | : : | : : : : . |||||
101 MFQLF.....MPEEGSGCSVRRRPYGCVLRIHPSPPPEKKELRKVAHLT 145
151 GKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSKVYFRGQ 200
|||||
146 GKSNSRSMPLEWEDTYGIVLLSGVKYKKGGLVINETGLYFVYSKVYFRGQ 195
201 SCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGAVFN 250
|||||
196 SCNNLPLSHKVYMRNSKYPQDLVMMEGKMMSYCTTGQMWARSSYLGAVFN 245
251 LTSADHLYVNVSELVLNFEESQTFFGLYKL 281
|||||
246 LTSADHLYVNVSELVLNFEESQTFFGLYKL 276

FIG. 27

Matrix Metalloproteinase Cleavage Sites

Cleavage ⇓									
P ₄	P ₃	P ₂	P ₁	P' ₁	P' ₂	P' ₃	P' ₄		
Collagenases									
MMP-1 Interstitial Collagenase									
SEQ ID NO: 50	P ₄ Ala	P ₃ Pro	P ₂ Leu	P ₁ Gly	P' ₁ Met	P' ₂ Arg	P' ₃ Met/Ala	P' ₄ Arg	
SEQ ID NO: 51	Gly/Leu	Leu	Met/Tyr	His	Leu	Leu	Gly	Lys	
SEQ ID NO: 52	Met	Ala	Val/Gly	Glu	Ile	Phe	Val	Gln	
SEQ ID NO: 53	Glu	Asp	Ile	Tyr	Gln	Trp	Ser	Ile	
SEQ ID NO: 54	Pro	Ser	Gln/Arg	Ala	Pro	Glu	Glu	Gly	
SEQ ID NO: 55	Tyr	Glu	Asp	Phe	Phe	Ala	Phe	Ser	
SEQ ID NO: 56	Ile	Gly	Glu	Gln	Ala	Val/Gly	Arg	Glu	
SEQ ID NO: 57	Thr	Arg	Ala	Asn	Tyr/Val	Ser	Pro	Ala	
SEQ ID NO: 58	Arg				[not K,E,W]	Asn			
MMP-8 Neutrophil Collagenase									
SEQ ID NO: 59	P ₄ Ala	P ₃ Pro	P ₂ Leu	P ₁ Glu	P' ₁ Tyr	P' ₂ Ala	P' ₃ Gly	P' ₄ Arg	
SEQ ID NO: 60	Gly/Leu	Leu	Gln	Gly/His	Ile	Leu	Met	Gln	
SEQ ID NO: 61	Met			Ala	Val	Trp	Ala		
SEQ ID NO: 62	Glu								
SEQ ID NO: 63	Pro				Phe				
SEQ ID NO: 64	Tyr/Ile/Thr/Arg			Ala					
(otherwise same as MMP-1)									

FIG. 28-1

Gelatinases																
MMP-2		Gelatinase A														
SEQ ID NO: 65	P ₄	Gly	P ₃	Pro	P ₂	Arg	P ₁	Gly	P ₁	Leu	P ₁	Ala/Leu	P ₃	Gly/Ala	P ₄	Gln
SEQ ID NO: 66		Ile		Ala		Gln	Asn.	Ile/Phe		Ile/Phe		Phe/Trp		Leu		Arg
SEQ ID NO: 67		Pro		Arg		Leu	Ala	Val/Met		Val/Met		Gly		Ser		His
SEQ ID NO: 68		Arg				Ala	His			Ala		Arg/Gln		Pro		Pro
SEQ ID NO: 69		Leu				Lys	Leu			Glu		His				
SEQ ID NO: 70						Ile	Tyr			Gln/Asn		Val				
SEQ ID NO: 71						His				Ser						
MMP-9		Gelatinase B														
SEQ ID NO: 72	P ₄		P ₃	Pro	P ₂	Arg	P ₁	Gly	P ₁	Leu	P ₁	Ala/Leu/Phe	P ₃	Ala	P ₄	Thr
SEQ ID NO: 73						Gln		Ile/Phe		Ile/Phe		Glu		Leu		
SEQ ID NO: 74		Gln/Arg								Val/Met		Trp/Gly		Ser		
SEQ ID NO: 75										Ala				Gly		
SEQ ID NO: 76						Leu										
Stromelysins																
MMP-3		Stromelysin 1														
SEQ ID NO: 77	P ₄	Asp	P ₃	Pro	P ₂	Phe	P ₁	Glu	P ₁	Leu	P ₁	Arg	P ₃	Ala	P ₄	Thr
SEQ ID NO: 78		Gly		Ala		Leu/Met		Ala		Phe		Leu/Phe		Arg/Met		
SEQ ID NO: 79		Gln/Arg														
SEQ ID NO: 80		Leu		Val		Tyr	Gln/Phe	Trp/Tyr		Trp/Tyr		Trp		Gly		Pro
SEQ ID NO: 81		Ile		Leu		Pro/Gly/Glu	Asn	Ile		Ile		Val		Val/Ile		
SEQ ID NO: 82		Glu/Val														
SEQ ID NO: 83		Leu		Thr		Ile	His	Val		Val		Gln		Ser/Asn		Ala
SEQ ID NO: 84		Lys		Phe		Ala	Gly	Met		Met		His/Met		Glu/Thr		
SEQ ID NO: 85		Gly/Asp														
SEQ ID NO: 86		Arg		Arg		Ser	Leu/Pro	Glu		Glu		Glu/Ser/Thr		Leu		
SEQ ID NO: 87		Ser/Lys/Phe														
SEQ ID NO: 88		Pro/Met		Ser/Gly			Lys/Tyr/Arg									
SEQ ID NO: 89		Ala/Phe/Gln														

FIG. 28-2

1

CHIMERIC NUCLEIC ACIDS ENCODING POLYPEPTIDES COMPRISING CD70 AND FAS LIGAND DOMAINS

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation application of U.S. application Ser. No. 11/015,117 filed Dec. 17, 2004, now U.S. Pat. No. 7,524,944 which is a continuation application of U.S. application Ser. No. 08/982,272 filed Dec. 1, 1997, now issued as U.S. Pat. No. 7,070,771; which claims the benefit under 35 USC §119(e) to U.S. Application Ser. No. 60/032,145 filed Dec. 9, 1996, now abandoned. The disclosure of each of the prior applications is considered part of and is incorporated by reference in the disclosure of this application.

GRANT INFORMATION

This invention was made with government support under Grant No. CA49870 awarded by National Institutes of Health. The government has certain rights in the invention.

BACKGROUND OF THE INVENTION

1. Field of the Invention

The present invention relates to novel expression vectors containing genes which encode an accessory molecule ligand and the use of those vectors for immunomodulation, improved vaccination protocols and the treatment of malignancies and autoimmune diseases. More particularly, this invention provides expression vectors and methods for treating various neoplastic or malignant cells, and expression vectors and methods for treating autoimmune Disease. This invention also contemplates the production and expression of accessory molecule ligands with greater stability and enhanced function.

2. Background Information

Leukemias, lymphomas, carcinomas and other malignancies are well known and described in, e.g., *Harrison's Principles of Internal Medicine*, Wilson et al., eds., McGraw-Hill, New York, pp. 1599-1612. These malignancies appear to have somehow escaped the immune system surveillance mechanisms that eliminate rapidly and continuously proliferating cells. The exact mechanism by which these malignancies escape the immune system surveillance is not known.

Some of these malignant immune system cells are malignant antigen presenting cells which do not function properly within the immune cascade. For example, neoplastic B cells cannot induce even weak allogeneic or autologous mixed lymphocyte reactions in vitro. Further evidence that malignancies survive due to the failure of the immune surveillance mechanism includes the increased frequency of such malignancies in immunocompromised individuals, such as allograft recipients and those receiving long-term immunosuppressant therapy. Further, the frequency of these malignancies is increased in patients having Acquired Immune Deficiency Syndrome (AIDS) and patients with primary immune deficiency syndromes, such as X-linked lymphoproliferative syndrome or Wiscott-Aldrich Syndrome (Thomas et al., *Adv. Cancer Res.* 57:329, 1991).

The immune system normally functions to eliminate malignant cells by recognizing the malignant cells as foreign cells and clearing those cells from the body. An immune reaction depends on both the immune system's antibody response and on the cellular immune response within a patient. More specifically, the cellular immune response

2

which acts to recognize the malignant cells as foreign requires a number of different cells of the immune system and the interaction between those cells. An immune reaction begins with a T lymphocyte (T cell) which has on its cell surface the T cell receptor. The T cell also has the ability to express on its surface various accessory molecules which interact with accessory molecules on the B lymphocyte (B cell). When the T cell receptor of the T cell specifically binds to a foreign antigen, such as a malignant cell, it becomes activated and expresses the accessory molecule ligand, CD40 ligand on its cell surface. The accessory cell molecule ligand is only present on the activated T cells for a short period of time and is rapidly removed from the cell surface. After the accessory cells molecule ligand is removed from the surface of the activated T cell, its ability to bind to B cells via the accessory molecule ligand is destroyed.

When present on the surface of an activated T cell, the accessory cell ligand can specifically bind to the accessory cell molecule present on the B cell. This specific T-B cell interaction causes the B and T cell to express costimulatory surface accessory molecule and cytokines which result in an immune activation which lead to cytolytic T cells which specifically kill and remove the malignant cell from the body.

The interaction with an activated T cell is not solely limited to B cells but rather can be carried out by any cell which is able to present antigen to the T cell (an antigen presenting cell). These cells include B lymphocyte, macrophages, dendritic cells, monocytes, Langerhans cells, interdigitating cells, follicular dendritic cells or Kupffer cells. These cells all are known to have various accessory molecules on the cell surface which allow them to interact with other cells of the immune system. For example, these antigen presenting cells all have the accessory molecule CD40 on their cell surface. The presence of these accessory molecules allows these antigen presenting cells to specifically bind to complimentary accessory molecule ligand and thus directly interact with other immune cells.

A large number of accessory molecule ligands are members of the tumor necrosis factor superfamily. (Fanslow et al., *Sem. Immun.*, 6:267-268 (1994). The genes for a number of these accessory molecule ligands have been cloned and identified. These accessory molecule ligand genes encode accessory molecules which all have the configuration of Type II membrane proteins and exhibit varying degrees of homology with other accessory molecule ligand genes. For example, the accessory molecule ligand genes encoding both murine CD40 ligand and human CD40 ligand have been isolated. See, Armitage et al., *Nature*, 357:80-82 (1992) and Hollenbaugh et al., *EMBO J.*, 11:4313-4321 (1992).

CD40 and its ligand, CD40 ligand are critical components of a normal immune response. CD40 mediated signals induce immune lymphocytes to proliferate and differentiate and become potent antigen presenting cells. Malignant or neoplastic B cells are poor antigen presenter cells and are unable to stimulate a vigorous allogeneic mixed lymphocyte reaction. Successful cross linking of CD40 molecules on immune cells results in a strong allogeneic mixed lymphocyte reaction suggesting a strong immune reaction. Various soluble CD40 ligands or antibodies specific for CD40 have been used to potentially cross link CD40. These soluble CD40 ligands and CD40-specific antibodies are not optimal for cross linking the CD40 molecules on antigen presenting cells and do not work as effectively as CD40 ligand expressed on a cell membrane to produce strong stimulation of antigen presenting cells. These methods are also difficult to implement because large amounts of CD40 ligand constructs or antibodies must be isolated which is difficult and time-consuming work. Other

strategies to utilize CD40 ligand in solution or as a membrane bound molecule including transformation of fibroblasts with CD40 ligand to produce cultured cells which are then used to present antigen are not amenable to in vivo human clinical protocols.

CD95 (Fas) interaction with its ligand (Fas-ligand, or FasL) functions to limit the duration of the immune response and/or life-span of activated lymphocytes. Apoptosis induced by Fas-FasL binding serves to clear activated self-reactive lymphocytes. Problems caused by altering this pathway have been demonstrated in animals with defects in Fas<->Fas-ligand interactions. Mice having mutations, which inactivate CD95 or FasL, develop numerous disorders including autoimmune pathology resembling that seen in patients with rheumatoid arthritis (RA) or systemic lupus. Zhang, et al., in *J. Clin. Invest.* 100:1951-1957 (1997) show that injection of FasL-expressing virus, into the joints of mice with collagen-induced-arthritis, results in apoptosis of synovial cells and relief of arthritis symptoms. Expression of Fas ligand allows clearance of activated cells which play a role in the pathogenesis of autoimmune disease. Therefore, a gene therapy strategy for introducing FasL into the joints of rheumatoid arthritis patients could function to improve disease pathology by leading to destruction of the infiltrating mononuclear cells.

Administration of soluble accessory molecules and accessory molecule ligands has been shown to trigger or to be associated with adverse physiological effects. For example, treatment of mice, having wild-type CD40-receptor expression, with soluble CD40L-CD8 fusion protein resulted in a pulmonary inflammatory response. This was not observed in mice in which the gene for the CD40 receptor had been knocked out. These experiments, described in Wiley, J. A. et al., *Journal of Immunology* 158:2932-2938 (1997), support in vitro data which suggest that CD40 ligation can result in inflammatory responses.

Direct administration of purified recombinant soluble Tumor Necrosis Factor (either a or P) results in shock and tissue injury, as described in Tracey, K. J., and A. Cerami, *Annu. Rev. Med.* 45:491-503 (1994). Within minutes after acute intravenous or intra-arterial administration of TNF, a syndrome of shock, tissue injury, capillary leakage syndrome, hypoxia, pulmonary edema, and multiple organ failure associated with a high mortality ensues. Chronic low dose of TNF causes anorexia, weight loss, dehydration and depletion of whole-body protein and lipid.

Soluble Fas ligand and receptor have also been shown to be associated with tissue damage and other adverse effects. CD95, the Fas receptor, is a mediator of apoptosis. Fas ligand induces apoptosis by binding to Fas receptor. As shown in Galle, P. R., et al., *J. Exp. Med.* 182:1223-1230 (1995) administering an agonistic anti-Fas antibody resulted in liver damage to mice. Mice injected intraperitoneally with the agonistic antibody died within several hours, and analyses revealed that severe liver damage by apoptosis was the most likely cause of death.

The role of soluble Fas ligand (FasL), in the pathogenesis of systemic tissue injury in aggressive lymphoma is described in Sato, K. et al., *British Journal of Haematology*, 94:379-382 (1996). The findings presented in this report indicate that soluble FasL is directly associated with the pathogenesis of liver injury and pancytopenia.

CD27, the receptor for the accessory molecule ligand, CD70, was shown, in a report written by van Oers, et al., in *Blood* 82:3430-3436 (1993), to be associated with B cell malignancies.

The above findings all contraindicate the administration of soluble accessory molecule ligands, highlighting the need for

therapies that increase the levels of these molecules without resulting in an elevation of their soluble forms.

Despite the wealth of information regarding accessory molecule ligand genes and their expression on the surface of various immune cells, the exact mechanism by which the accessory molecule ligand genes are regulated on antigen presenting cells is not yet known. Without specific knowledge of the regulation of expression of accessory molecule ligand genes on these antigen presenting cells, altering the immune response by varying expression of an accessory molecule ligand gene has to date not been possible. Without any specific knowledge as to how to regulate the expression of an accessory molecule ligand gene on an antigen presenting cell, it is not possible to alter the immune response towards malignant cells. Thus, there was a need for a method of increasing the expression of an accessory molecule ligand gene on normal and malignant cells including antigen presenting cells.

Further, without the ability to regulate the expression of accessory molecule ligands, it is not possible to alter the immune clearance of these cells.

SUMMARY OF THE INVENTION

The present invention fills these needs by providing novel expression vectors containing accessory molecule ligand genes and methods for introducing those genes into normal and malignant antigen presenting cells thereby allowing the alteration of an immune response, the treatment of autoimmune diseases and the treatment of various neoplasias. This invention provides vectors, including gene therapy vectors which contain accessory molecule ligand genes. These vectors also contain the additional genetic elements, such as promoters, enhancers, polyadenylation signals (3' ends), which allow that vector to be successfully placed within the cell and to direct the expression of the accessory molecule ligand gene in a cell. Such gene therapy vectors are capable of transforming animal cells directly and thereby introducing the accessory molecule ligand gene into the cells of that animal in a form which can be utilized to produce accessory molecule ligands within that cell.

In other aspects of the present invention, the function of an accessory molecule ligand is modified by altering the half life of the molecule on the cell surface or by changing the level of expression of that molecule on the cell surface. In preferred embodiments, the present invention provides accessory molecule ligands which are modified to improve the stability of such accessory molecule ligands on the cell surface. Such increased stability may be accomplished using any of the disclosed methods of molecules described in this application, including chimeric molecules and molecules into which mutations have been introduced at least one location. The present invention also contemplates increasing the expression of such a molecule.

The present invention also provides gene therapy vectors containing the accessory molecule ligand genes which are chimeric in that portions of the gene are derived from two separate accessory molecule ligands which may or may not be from different species. The accessory molecule ligand genes of the present invention include genes which encode molecules of the tumor necrosis factor (TNF) family. The molecules which make up the TNF family include TNF α , TNF β , CD40 ligand, Fas ligand, CD70, CD30 ligand, 41BB ligand (4-1BBL), nerve growth factor and TNF-related apoptosis inducing ligand (TRAIL). In some embodiments of the present invention, the chimeric accessory molecule ligand genes of the present invention contain at least a portion of a murine accessory molecule ligand gene together with por-

5

tions of accessory molecule ligand genes derived from, either mouse, humans or other species. Some preferred embodiments of the present invention utilize murine CD40 ligand genes and chimeric CD40 ligand genes containing at least a segment of the murine CD40 ligand gene together with at least a segment of the human CD40 ligand gene. The present invention contemplates chimeric accessory molecule ligand genes wherein segments from the accessory molecule ligand gene of one species have been interchanged with segments from a second accessory molecule ligand gene which may optionally be from a different species. For example, in one preferred embodiment, the murine CD40 ligand gene transmembrane and cytoplasmic domains have been attached to the extracellular domains of human CD40 ligand gene.

The present invention contemplates gene therapy vectors which are capable of directly infecting the human, mammal, insect, or other cell. The use of such gene therapy vectors greatly simplifies inserting an accessory molecule ligand gene into those cells. The contemplated gene therapy vectors may be used in vivo or in vitro to infect the desired cell and are particularly useful for infecting malignant cells to effect sustained high-level expression of a physiologic ligand.

The present invention also contemplates animal, mammal, and human cells containing a gene therapy vector which includes an accessory molecule ligand gene and sufficient genetic information to express that accessory molecule ligand within that cell. In preferred embodiments, the present invention also contemplates human neoplastic antigen presenting cells which contain the gene therapy vectors of the present invention or contain an accessory molecule ligand gene together with a promoter and 3' end region.

The present invention also contemplates human cells and human neoplastic cells containing a gene therapy vector which includes a chimeric accessory molecule ligand gene. The present invention also contemplates bacterial cells or animal cells containing accessory molecule ligand genes, chimeric accessory molecule ligand genes, murine accessory molecule ligand genes, human accessory molecule ligand genes, the gene therapy vectors of the present invention, the vectors of the present invention, and a chimeric accessory molecule ligand gene together with a heterologous promoter, enhancer or polyadenylation sequence.

The present invention also contemplates methods of altering immune response within a human patient or the immunoreactivity of human cells in vivo by introducing a gene which encodes an accessory molecule ligand gene into the human cells so that that accessory molecule ligand is expressed on the surface of those human cells. This method includes the introduction of the accessory molecule ligand gene as part of a gene therapy vector or in association with a heterologous or native promoter, enhancer or polyadenylation signal. Some preferred embodiments of the present invention utilize introduction of Fas ligand genes and chimeric Fas ligand genes, constructed as contemplated above for CD40, into human cells to alter their immunoreactivity. The present invention also includes methods in which such accessory molecule ligand genes are inserted into cells which have the accessory molecule to which the accessory molecule ligand binds on the surface of the cell into which the accessory molecule ligand gene.

The present methods of altering immunoreactivity are applicable to all types of human, animal, and murine cells including human neoplastic cells such as human lymphomas, leukemias and other malignancies. In preferred embodiments, this method is used to introduce the gene encoding the accessory molecule ligand into potential antigen presenting cells of a human patient or cell which can stimulate bystand-

6

ing antigen presenting cells. Such antigen presenting cells include monocytes, macrophages, B cells, Langerhans cells, interdigitating cells, follicular dendritic cells, Kupffer cells, and the like. The various antigen presenting cells may be present as part of a known malignancy in a human patient such as leukemias, lymphomas, acute monocytic leukemia (AML)/chronic lymphocytic leukemia (CLL), acute myelomonocytic leukemia (AMML), chronic myelogenous or chronic myelomonocytic leukemia (CMML) and thus would include all tumors of any cell capable of presenting antigen to the human or animal immune system or are capable of stimulating bystander antigen presenting cells. The present invention also contemplates modulating the immune system by introducing genes encoding an accessory molecule ligand gene of the present invention into any number of different cells found in a patient, including muscle cells, skin cells, stromal cells, connective tissue cells, fibroblasts and the like.

The present invention also contemplates methods of treating neoplasias in either a human patient or an animal patient. In one preferred embodiment, the method comprises isolating the neoplastic cells from the human or animal patient and inserting into those isolated cells the gene which encodes the chimeric accessory molecule ligand or the accessory molecule ligand so that that molecule is expressed on the cell surface of those neoplastic cells or other somatic cells. The neoplastic cells are then infused back into the human or animal patient and may then participate in an enhanced immune response.

The present invention also contemplates the co-infection or co-introduction of the accessory molecule ligand gene together with a gene which encodes a tumor or carcinoma specific antigen. This combination of molecules are then expressed on the surface of the neoplastic cells and when those cells are introduced into the patient lead to the rapid immune response resulting in the destruction of those cells.

The present methods also include directly introducing the gene therapy vector or other vector carrying the accessory molecule ligand gene directly into the tumor or tumor bed of a patient. Upon entering the tumor bed of the patient, the gene therapy vector or other vector enter the cells present in the tumor or tumor bed and then express the accessory molecule ligand gene on the surface of those cells. These cells then are able to participate fully in the human immune or animal immune response.

The present invention also contemplates methods of augmenting an immune response to a vaccine. The present method of vaccinating an animal against a predetermined organism or antigen by administering to that animal a vaccine which has a genetic vector containing an accessory molecule ligand gene. Other embodiments of the present invention include vaccinating an animal by administering two separate genetic vectors, one containing the antigens from the organism to which immunity is desired by isolating the cells of the target animal and contacting with those cells a vector encoding at least one antigen from a predetermined organism so that the antigen is expressed by the cells and also contacting those cells with a different vector which expresses the accessory molecule ligand gene on the surface of the animal's antigen presenting cells. Together these two separate vectors produce a vaccination which is much stronger and of longer duration than is vaccination with antigen alone.

The present methods of vaccination are applicable to vaccinations designed to produce immunity against a virus, a cell, a bacteria, any protein or a fungus. The present methods are also applicable to immunization against various carcinomas and neoplasias. In these embodiments, the tumor antigen

against which immunity is desired is introduced into the animal together with the genetic vector containing the accessory molecule ligand gene.

The present invention also contemplates methods of treating arthritis utilizing a gene therapy vector encoding an accessory molecule ligand. Of particular interest for use with arthritis is the Fas ligand molecule in which the expression of Fas ligand activity has been increased in the joint and/or the stability of the Fas ligand activity on cells within the joint enhanced. In other embodiments, the present invention contemplated methods of treating arthritis utilizing chimeric accessory molecule ligands and chimeric accessory molecule ligand genes. The present invention also contemplates both ex vivo therapy and in vivo therapy of arthritis utilizing the expression vectors of the present invention together with the Fas ligand and modified versions of that molecule including chimeric molecules.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a diagram showing a number of accessory molecule ligand genes and Domains I-IV of those genes as deduced from sequence data.

FIG. 2 is a diagram showing example chimeric accessory molecule ligand genes. The domains derived from the murine accessory module are shown shaded.

FIGS. 3A through 3D show the amount of either mouse or human CD40 ligand found on the surface of Hela or CLL cells infected with gene therapy vectors containing the genes encoding these molecules. FIG. 3A shows uninfected Hela cells (shaded) and Hela cells infected with a gene therapy vector encoding murine CD40 ligand. FIG. 3B shows uninfected Hela cells (shaded) and Hela cells infected with a gene therapy vector encoding human CD40 ligand. FIG. 3C shows uninfected CLL cells (shaded) and CLL cells infected with a gene therapy vector encoding murine CD40 ligand. FIG. 3D shows uninfected CLL cells (shaded) and CLL cells infected with a gene therapy vector encoding human CD40 ligand.

FIGS. 4A through 4D show histograms of the increased expression of CD54 (FIG. 4B) and CD80 (FIG. 4D) on CLL cells into which a gene therapy vector containing the accessory molecule ligand gene (murine CD40 ligand gene) has been introduced. The shaded graph indicates control stain in FACS analysis and the open graph indicates staining with monoclonal antibodies immunospecific for either CD54 (FIGS. 4A and 4B) or CD80 (FIGS. 4C and 4D).

FIG. 5 shows the cell proliferation as measured by ^3H -TdR incorporation of allogeneic T cells in response to various stimulation regimes. The CLL cells containing a gene therapy vector expressing an accessory molecule ligand gene (the murine CD40 ligand gene) were introduced, stimulating allogeneic T cells to proliferate.

FIG. 6 shows the production of gamma interferon ($\text{IFN}\gamma$) by allogeneic T cells stimulated with CLL cells containing an accessory molecule ligand gene.

FIG. 7 shows the treatment of a neoplasia in an animal using a gene therapy vector containing an accessory molecule ligand gene of the present invention. The open squares show mice immunized with neoplastic cell not expressing an accessory molecule ligand of the present invention. Mice immunized with neoplastic cells expressing an accessory molecule ligand of the present invention are shown as the horizontal line at the top of the Figure and show no morbidity.

FIG. 8 shows the production levels and stabilities of CD40 ligand and CD40 ligand transcript in CLL (upper graph) and normal blood mononuclear cells (lower graph).

FIG. 9 shows the time course of transgene expression in CLL B cells infected with the accessory molecule ligand (CD40 ligand). The MFIR (mean fluorescence intensity ratio), comparing the fluorescence intensity of $\text{CD}19^+$ CLL cells stained with PE-labeled CD40 ligand versus the same stained with a PE-labeled isotype control mAb at each time point, are represented by the closed circles connected by solid lines according to the scale provided on the left-hand ordinate.

FIG. 10 shows changes in surface antigen phenotype of CLL B cells infected with a gene therapy vector containing an accessory molecule ligand, CD40 ligand. Shaded histograms represent staining of uninfected CLL cells (thin lines) stained with nonspecific control antibody, open histograms drawn with thin lines represent uninfected CLL cells stained with FITC-conjugated specific mAb, and open histograms drawn with thick lines (labeled CD154-CLL) represent CLL cells infected with the accessory molecule ligand gene therapy vector and stained with FITC-conjugated specific mAb.

FIGS. 11A and 11B show levels of CD27 produced in CLL cells infected with a gene therapy vector containing an accessory molecule ligand. FIG. 11A shows that CD40L-infected CLL (CD154-CLL) cells express reduced levels of surface CD27. Open histograms represent staining of non-infected CLL cells (thin lines) or infected CLL (thick lines) with FITC-conjugated $\alpha\text{CD}27$ mAb, respectively. FIG. 11B shows production of soluble form of CD27 by CLL B cells.

FIGS. 12A through 12D show allogeneic T cell responses induced by CLL cells infected with a gene therapy vector containing an accessory molecule ligand (CD40 ligand, also called CD154). FIG. 12A indicates the concentration of $\text{IFN}\gamma$ in the supernatants after stimulation of allogeneic T cells with CLL cells containing the accessory molecule ligand. FIG. 12B shows cell proliferation, as assessed by incorporation of ^3H -thymidine. FIGS. 12C and 12D show secondary allogeneic T cell responses induced by CLL containing the accessory molecule ligand.

FIGS. 13A through 13C depict autologous T cell responses induced by CLL B cells containing the accessory molecule ligand, CD40 ligand or CD154, and controls. FIG. 13A shows incorporation of ^3H -thymidine by autologous T cells co-cultured with the CLL cells. FIG. 13B shows the levels of human $\text{IFN}\gamma$ produced by autologous T cells co-cultured with the CLL cells. In FIG. 13C, the CTL activities of autologous T cells induced by CLL B cells containing the accessory molecule ligand are graphed.

FIGS. 14A through 14C show specificity of CTL for autologous CLL B cells. $\text{IFN}\gamma$ concentration was assessed in the supernatants after 48 h of culture (FIG. 14A), and cytolytic activity was assessed at 3 h of culture (FIG. 14B). In FIG. 14C, mAb were added to the autologous leukemia target cells prior to the CTL assay.

FIGS. 15A through 15D show that intercellular stimulation plays a role in production of the phenotypic changes observed in CLL cells expressing the accessory molecule ligand. In FIG. 15A, the effect of culture density on the induced expression of CD54 and CD80 following infection with a gene therapy vector containing the accessory molecule ligand (CD40 ligand, CD154) is shown. Shaded histograms represent staining of leukemia B cells with a FITC-conjugated isotype control mAb. Open histograms represent CD154-CLL B cells, cultured at high or low density (indicated by arrows), and stained with a FITC-conjugated mAb specific for CD54 or CD80. FIG. 15B shows inhibition of CD154-CLL cell activation by anti-CD154 mAb. FIGS. 15C and 15D depict expression of immune accessory molecules on bystander non-infected CLL B cells induced by CLL cells

expressing the accessory molecule ligand. Shaded histograms represent staining with PE-conjugated isotype control mAb.

FIGS. 16A and 16B show that the vector encoding an accessory molecule ligand enhances immunization against β -gal in mice. FIG. 16A shows that mice that received intramuscular injections of the pCD40L vector produced significantly more antibodies to β -gal than did mice injected with either the non-modified pcDNA3 vector or pCD40L. FIG. 16B, ELISA analyses of serial dilutions of sera collected at d28, shows that mice co-injected with placZ and pCD40L had an eight-fold higher mean titer of anti- β -gal antibodies at d28 than mice treated with placZ+pcDNA3.

FIGS. 17A and 17B show analysis of the IgG₁ and IgG_{2a} immune responses to intramuscular plasmid DNA immunizations with and without a vector, pCD40L, encoding an accessory molecule ligand. IgG_{2a} anti- β -gal antibodies predominated over IgG₁ subclass antibodies in the sera of mice injected with either placZ and pcDNA3 or placZ and pCD40L. In contrast, BALB/c mice injected with β -gal protein developed predominantly IgG₁ anti- β -gal antibodies, and no detectable IgG_{2a} anti- β -gal antibodies.

FIG. 18 shows the comparison between injection of mice with a vector, pCD40L, encoding an accessory molecule ligand, at the same and different sites as placZ. Adjuvant effect of pCD40L requires co-injection with placZ at the same site.

FIG. 19 shows that co-injection into dermis of a vector encoding an accessory molecule ligand, pCD40L, with placZ enhances the IgG anti- β -gal response in BALB/c mice.

FIGS. 20A and 20B show that a vector encoding an accessory molecule ligand, pCD40L, enhances the ability of placZ to induce CTL specific for syngeneic β -gal-expressing target cells. Splenocyte effector cells, taken from mice which had received injections of placZ and pCD40L, specifically lysed significantly more cells than did splenocytes from mice that received control injections.

FIGS. 21A through 21D show downmodulation of human CD40L, but not murine CD40L, in lung tumor cell lines that express CD40.

FIG. 22A shows that CD40 binding induces enhanced expression of the tumor cell surface markers CD95 (Fas), CD54 (ICAM-1), and MHC-I, in lung tumor cell lines. FIG. 22B shows downmodulation of human CD40L by CD40-positive tumor cells.

FIG. 23 shows the inhibition of Fas ligand expression by lymphocytes in the presence of RA synovial fluid.

FIG. 24 shows an outline for a clinical trial of an accessory molecule ligand (CD40L) gene therapy treatment for B cell CLL.

FIG. 25 shows a sequence line-up of human Fas ligand with human Fas ligand in which Domain III is replaced by Domain III of murine Fas ligand. The top protein sequence is native human Fas ligand. Domain III is underlined with the dotted line. The double underline indicates a putative MMP cleavage site. The bottom protein sequence is that of chimeric human-mouse Fas ligand. Domain III of the mouse Fas ligand (underlined with dotted line) is substituted for Domain III of human Fas ligand. The numbers correspond to the amino acid sequence number using 1 for the start of the polypeptide sequence. The number of the first nucleotide base for the codon encoding the amino acid is $1+3 \times (n-1)$, where n is the amino acid sequence number.

FIG. 26 shows a sequence line-up of human Fas ligand with human Fas ligand in which Domain III has been replaced with Domain III of human CD70. The top protein sequence is native human Fas ligand, and the bottom sequence is that of

chimeric Fas ligand, in which Domain III of human CD70 has been substituted for Fas Domain III. Other markings are used similarly as in FIG. 25.

FIG. 27 shows a sequence line-up of human Fas ligand with human Fas ligand in which Domain I has been replaced with Domain III of human CD70. The top protein is native human Fas ligand, and the bottom protein sequence is that of chimeric Fas ligand, in which Domain III has been replaced with Domain I of human CD70. Other markings are used similarly as in FIG. 25.

FIG. 28 shows the amino acids around and at known matrix metalloproteinase (MMP) cleavage sites, as described in Smith, M. M. et al., Journal of Biol. Chem. 270:5440-6449 (95) and Nagase, H., and G. B. Fields, Biopolymers (Peptide Science) 40:399-416 (96). The cleavage site is indicated with an arrow.

DETAILED DESCRIPTION OF THE INVENTION

All references cited herein are hereby incorporated in their entirety by reference.

I. Definitions

An "accessory molecule ligand gene" is a gene which encodes all or part of an accessory molecule ligand. The gene comprises at least the nucleotide sequence required to encode the functional portion of an accessory molecule ligand. The gene may optionally include such genetic elements as promoters, enhancers and 3' ends. The accessory molecule ligand gene is derived from a ligand which is a member of the tumor necrosis factor (TNF) family, including CD40 ligand, Fas ligand, CD70, TNF α , TNF β , CD30 ligand, 4-1BB ligand (4-1BBL), nerve growth factor and TNF-related apoptosis inducing ligand (TRAIL). As used herein, the term "accessory molecule ligand gene" includes chimeric accessory molecule ligand genes as defined below.

As used herein, the term "malignant cells or neoplastic cells," is defined to mean malignant or cancerous cells which are found in a human patient or an animal. Preferred types of malignant or neoplastic cells include any malignant antigen-presenting cell. In some preferred embodiments, these malignant antigen presenting cells have at least low levels of CD40 present on the cell surface.

As used herein, the term "neoplastic human cells" is defined to mean human cells which are neoplastic including but not limited to antigen presenting cells, any neoplastic cell which may function as an antigen presenting cell or function to facilitate antigen presentation, neoplastic monocytes, neoplastic macrophages, neoplastic B cells, neoplastic dendritic cells, neoplastic Langerhans cells, neoplastic interdigitating cells, neoplastic follicular dendritic cells, or neoplastic Kupffer cells and the like. The definition of neoplastic human cells includes those cells which are associated with neoplastic cells in the tumor bed of human patients. Typically, the neoplastic human cells are either leukemias, lymphomas, AML, ALL, AMML, CML, CMML, CLL other tumors of antigen presenting cells or breast, ovarian or lung neoplastic cells. It is also contemplated that the accessory molecule ligand genes or chimeric accessory molecule ligand genes of the present invention may be inserted into somatic cells. These somatic cells can be created by a genetic engineering process which has introduced into those cells genes which encode molecules which render those cells capable of presenting antigen to the immune system.

As used herein, the term "chimeric gene" is defined to mean a gene in which part of the gene is derived from a second

different gene and combined with the first gene so that at least a portion of each gene is present in the resulting chimeric gene. A gene may be chimeric if any portion of the sequence which encodes the resulting protein is derived from a second and different gene. Typical chimeric genes include genes in which specific functional domains from one gene have been transferred to a second gene and replace the analogous domains of that second gene. For example, the resulting chimeric gene may have one domain derived from a murine gene and several domains derived from a human gene. These domains may range in size from 5 amino acids to several hundred amino acids. Other examples of chimeric accessory molecule ligand genes include genes which contain nucleotides encoding amino acids not found in any naturally occurring accessory molecule ligand gene. Examples of chimeric genes and potential various combinations of domains are numerous and one of skill in the art will understand that no limit is placed on the amount of one gene that must be present in a second gene to render it chimeric.

As used herein, the term "murine CD40 ligand gene" is defined to mean an accessory molecule ligand gene which is derived from a murine CD40 ligand gene. Examples of such murine CD40 ligand genes include the gene isolated by Armitage et al., *Nature*, 357:80-82 (1992) and other genes derived from murine origin which hybridize to the gene described by Armitage et al. under low stringency hybridization conditions.

As used herein, the term "vector or genetic vector" is defined to mean a nucleic acid which is capable of replicating itself within, an organism such as a bacterium or animal cell. Typical genetic vectors include the plasmids commonly used in recombinant DNA technology and various viruses capable of replicating within bacterial or animal cells. Preferred types of genetic vectors includes plasmids, phages, viruses, retroviruses, and the like.

As used herein, the term "gene therapy vector" is defined to mean a genetic vector which is capable of directly infecting cells within an animal, such as a human patient. A number of gene therapy vectors have been described in the literature, and include, the gene therapy vector described in Cantwell et al., *Blood*, In Press (1996) entitled "Adenovirus Vector Infection of Chronic Lymphocytic Leukemia B Cells." Such vectors have been described for example by Woll, P. J. and I. R. Hart, *Ann. Oncol.*, 6 Suppl 1:73 (1995); Smith, K. T., A. J. Shepherd, J. E. Boyd, and G. M. Lees, *Gene Ther.*, 3:190 (1996); Cooper, M. J., *Semin. Oncol.*, 23:172 (1996); Shaughnessy, E., D. Lu, S. Chatterjee, and K. K. Wong, *Semin. Oncol.*, 23:159 (1996); Glorioso, J. C., N. A. DeLuca, and D. J. Fink, *Annu. Rev. Microbiol.*, 49:675 (1995); Flotte, T. R. and B. J. Carter, *Gene Ther.*, 2:357 (1995); Randrianarison-Jewtoulkoff, V. and M. Perricaudet, *Biologicals.*, 23:145 (1995); Kohn, D. B., *Curr. Opin. Pediatr.*, 7:56 (1995); Vile, R. G. and S. J. Russell, *Br. Med. Bull.*, 51:12 (1995); Russell, S. J., *Semin. Cancer Biol.*, 5:437 (1994); and Ali, M., N. R. Lemoine, and C. J. Ring, *Gene Ther.*, 1:367 (1994). All references cited herein are hereby incorporated by reference.

II. Genetic Vectors and Constructs Containing an Accessory Molecule Ligand Gene

A. Accessory Molecule Ligand Genes

In one embodiment of the present invention, preferred gene therapy vectors contain an accessory molecule ligand gene. This accessory molecule ligand gene may be derived from any source and may include molecules which are man-made and do not appear in nature. The present invention contemplates accessory molecule ligand genes which are derived

from the genes encoding molecules within the tumor necrosis family (TNF) which includes the genes encoding: murine CD40 ligand, human CD40 ligand, Fas ligand, TNF α , TNF β , CD30 ligand, 4-1BB ligand, nerve growth factor, CD70, TNF-related apoptosis inducing ligand (TRAIL) and chimeric accessory molecule ligands. The nucleotide sequence of one accessory molecule ligand, the sequence of at least one form of the murine CD40 ligand gene, has been determined and is listed as SEQ ID NO: 1. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to the sequence present in SEQ ID NO: 1, and thus hybridizes to this sequence at low stringency hybridization conditions. One of skill in the art will understand that accessory molecule ligand genes, including murine CD40 ligand gene, useful in the present invention may be isolated from various different murine strains.

The nucleotide sequence of a human CD40 ligand gene has been determined and is shown as SEQ ID NO: 2. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to SEQ ID NO: 2, and thus hybridizes to this sequence at low stringency conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the human CD40 ligand genes, useful in the present invention, may vary depending on the individual from which the gene is isolated and such variations may prove useful in producing unique accessory molecule ligand genes. The present invention contemplates the use of the domains, sub-domains, amino acid or nucleotide sequence of the human CD40 ligand and/or human CD40 ligand gene as part of a chimeric accessory molecule ligand or chimeric accessory molecule ligand gene.

The nucleotide sequence of a bovine CD40 ligand gene has been determined and is shown as SEQ ID NO: 8. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to SEQ ID NO: 8, and thus hybridizes to the sequence at low stringency conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the bovine CD40 ligand genes, may vary depending on the individual animal from which the gene is isolated and that such variations may prove useful in producing unique accessory molecule ligand genes.

The nucleotide sequence of human TNF α and human TNF β have been determined and are shown as SEQ ID NOS: 9 and 10, respectively. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to either human TNF α or human TNF β (SEQ ID NOS: 9 and 10, respectively), and thus hybridizes to these sequences at low stringency conditions. The accessory molecule ligand genes useful in the present invention, including the human TNF α and TNF β genes, may vary depending on the particular individual from which the gene has been isolated and these variations may prove useful in producing unique accessory molecule genes.

The nucleotide sequence of porcine TNF α and TNF β have been determined and are shown as SEQ ID NO: 11. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to either SEQ ID NO: 11, and thus would hybridize to these sequences at low stringency conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the porcine TNF α and TNF β genes, may vary depending on the particular animal from which the gene is isolated and that such variation may prove useful in producing unique accessory molecule genes.

The nucleotide sequence of a murine TNF α gene has been determined and is shown as SEQ ID NO: 12. The present invention contemplates the use of any accessory molecule

ligand gene which is homologous to SEQ ID NO: 12, and thus hybridizes to the sequence at low stringency conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the murine TNF α gene may vary depending on the individual from which the gene is isolated and that these variations may prove useful in producing unique accessory molecule genes.

The nucleotide sequence of human Fas ligand and murine (C57BL/6) Fas ligand have been determined and are shown as SEQ ID NOS: 13 and 14, respectively. The nucleotide sequence of murine Balb/c Fas ligand is shown as SEQ ID NO: 31. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to any of SEQ ID NOS: 13, 14, and 31, and thus hybridizes to the sequences at low stringency conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the human Fas ligand or murine Fas ligand genes may vary depending on the particular individual or animal from which the gene is isolated and that such variations may prove useful in producing any accessory molecule genes.

The nucleotide sequence of a human CD70 gene has been determined and is shown as SEQ ID NO: 15. The murine CD70 gene sequence has also been determined, and is shown as SEQ ID NO: 36 and was described by Tesselaar et. al, *J. Immunol.* 159:4959-65(1997). The present invention contemplates the use of any accessory molecule ligand gene which is homologous to SEQ ID NO: 15 or 36, and thus hybridizes to this sequence at low stringency-conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the human CD70 gene may vary depending on the individual from which the gene is isolated and that these variations may prove useful in producing unique accessory molecule ligand genes.

The nucleotide sequence of human CD30 ligand gene has been determined and is shown as SEQ ID NO: 16. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to SEQ ID NO: 16, and thus hybridizes to this sequence at low stringency conditions. One of ordinary skill in the art will understand that the accessory molecule ligand genes, including the human CD30 ligand gene, may vary depending on the individual from which the gene is isolated and that such variations may prove useful in producing unique accessory molecule ligand genes.

The present invention also contemplates variations and variants of the nucleotide sequences of the accessory molecule ligand genes provided herein which are caused by alternative splicing of the messenger RNA. This alternative splicing of the messenger RNA inserts additional nucleotide sequences which may encode one or more optional amino acid segments which in turn allows the accessory molecule ligand encoded to have additional properties or functions.

The nucleotide sequence of a human and mouse 4-1BBL have been determined and are shown as SEQ ID NOS: 17 and 18, respectively. The present invention contemplates the use of any accessory molecule ligand gene which is homologous to either SEQ ID NOS: 17 or 18, and thus hybridizes to these sequences at low stringency conditions. One of ordinary skill in the art will understand that accessory molecule ligand genes, including the human 4-1BBL gene may vary depending on the individual from which it is isolated and that such variations may prove useful in producing unique accessory molecule ligand genes.

The present invention also contemplates chimeric accessory molecules containing any domain, sub-domain portion, or amino acid sequence encoded by the following genes: bovine TNF- α (SEQ ID NO: 21), murine CD40 ligand (SEQ

ID NO: 22), human nerve growth factor- β (SEQ ID NO: 23), murine nerve growth factor (SEQ ID NO: 24), rat Fas ligand (SEQ ID NO: 25), human TNF-related apoptosis inducing ligand (TRAIL) (SEQ ID NO: 41, Genbank accession number U37518), murine TNF-related apoptosis inducing ligand (TRAIL) (SEQ ID NO: 42, Genbank accession number U37522), murine CD30-Ligand (SEQ ID NO: 43), human 4-1BBL (SEQ ID NO: 17), an murine 4-1BBL (SEQ ID NOS: 44 and 18). The present invention also contemplates chimeric accessory molecules which utilize genes encoding amino acid sequences homologous to these sequences.

The present invention contemplates chimeric accessory molecule ligand genes which are comprised of a nucleotide segment derived from one accessory molecule ligand gene operatively linked to a nucleotide sequence derived from a different accessory molecule ligand gene or other gene.

For example, chimeric accessory molecule ligand genes are contemplated which are comprised of a segment of the murine CD40 ligand gene which has been operatively linked to at least one other additional gene segment derived from a different accessory molecule ligand gene. The size of the particular segment derived from the different accessory molecule ligand gene may vary from a nucleotide sequence encoding a few amino acids, a sub-domain of the accessory molecule ligand, a domain of the accessory molecule ligand or more than a domain of an accessory molecule ligand. Other chimeric accessory molecules of the present invention are comprised of an accessory molecule ligand gene into which nucleotides encoding an amino acid segment which is not found as part of a naturally occurring accessory molecule ligand have been inserted. This amino acid segment may be artificially created or derived from a protein found in nature. The chimeric accessory molecule ligand gene encodes a chimeric amino acid sequence and thus a chimeric accessory molecule ligand encoded may possess unique properties in addition to the properties found on the individual segments derived from the different accessory molecule ligand genes. The chimeric accessory molecule ligand gene may encode an accessory molecule ligand which has properties derived from the accessory molecule ligand used to construct the chimeric gene.

Each of the accessory molecule ligand genes which are a member of the tumor necrosis factor family have a similar secondary structure consisting of a number of domains. This domain structure includes a first domain which is encoded by the 5' region of the accessory molecule ligand gene. The second domain (Domain II) is the domain which contains the amino acids which span the cell membrane and is thus called the transmembrane domain. The third domain (Domain III) is the proximal extracellular domain and these amino acids are the amino acids which are found proximal to the cellular membrane. The fourth domain (Domain IV), is encoded by the 3' end of the accessory molecule ligand gene and has been called the distal extracellular domain. The distal extracellular domain (Domain IV) generally makes up the soluble form of the tumor necrosis factor family molecule. Based on the x-ray crystal structure of human TNF, the predicted secondary structure of the accessory molecule, CD40 ligand has been deduced together with the domain structure of these molecules by M. Peitsch and C. Jongeneel, *International Immunology*, 5:233-238 (1993). The secondary structures of the other members of the tumor necrosis factor family were deduced using computer analysis together with comparison to the human TNF and CD40 ligand domain structure. In Table I, the domain boundaries of a number of accessory molecule ligand genes is shown. A diagram of these domains for a number of these accessory cell molecule ligands is

shown in FIG. 1. The assignments of the domain boundaries are approximate and one of ordinary skill in the art will understand that these boundaries may vary and yet still provide useful identification of domains.

TABLE I

DOMAIN STRUCTURE OF TUMOR NECROSIS FACTOR FAMILY MOLECULES*				
	Domain I (Cytoplasmic)	Domain II (Transmembrane)	Domain III (Proximal Extracellular)	Domain IV (Distal Extracellular)
Human CD40 Ligand	1-42	43-135	136-330	331-786
Murine CD40 Ligand	1-42	43-135	136-327	328-783
Bovine CD40 Ligand	1-42	43-135	136-330	331-786
Human TNF- α	1-87	88-168	169-228	229-699
Murine TNF- α	1-87	88-168	169-237	238-705
Porcine TNF- α	1-87	88-168	169-228	229-696
Human TNF- β	1-39	40-129	130-153	154-615
Porcine TNF- β	1-39	40-126	127-150	151-612
Human Fas Ligand	1-237	238-315	316-390	391-843
Murine Fas Ligand	1-237	238-309	310-384	385-837
Human CD70	1-61	62-117	118-132	133-579
Murine CD70	1-73	74-123	124-138	139-585
Human CD30 Ligand	1-117	118-186	187-240	241-702
Murine CD30 Ligand	1-135	136-201	202-255	256-717
Human 4-1BBL	1-69	70-174	175-210	211-762
Murine 4-1BBL	1-237	238-333	334-369	370-927
Human TRAIL	1-39	40-117	118-375	376-843
Murine TRAIL	1-51	52-111	112-387	388-873

*The Domains above are identified by the nucleotide boundaries of each domain using the first nucleotide of the initial methionine of the cDNA as nucleotide number 1.

35

One of ordinary skill in the art will understand that typical chimeric accessory molecule genes would include genes produced by exchanging domains or sub-domain segments between, for example, a mouse CD40 ligand gene and a human CD40 ligand gene. For example, chimeric accessory molecule gene may be constructed by operatively linking Domain I of the human CD40 ligand gene to Domains II-IV of the murine CD40 ligand gene. One of ordinary skill in the art will understand the variety of chimeric accessory molecule ligand genes which may be produced using the accessory molecules identified in Table I. The present invention also contemplates chimeric accessory molecules which are not shown in Table I but which are shown to have a similar domain structure. Other chimeric genes are also contemplated in which smaller segments (sub-domain segments) are exchanged between, for example, a murine CD40 ligand gene and a human CD40 ligand gene or a second murine CD40 ligand gene. One of skill in the art will understand that genes encoding accessory molecules will have at least gene segments which correspond to various functional segments of an accessory molecule ligand such as the murine CD40 ligand encoded by the murine CD40 ligand gene (SEQ ID NO: 1). It will also be apparent to one of skill in the art that the nucleotide boundaries identified in Table I may vary considerably from those identified for the murine CD40 ligand gene (SEQ ID NO: 1) and still define domains which are useful in the present invention.

In one preferred embodiment, the chimeric accessory molecule ligand gene is comprised of the nucleotides encoding extracellular domains (Domains III and IV) of human CD40 ligand operatively linked to the nucleotides encoding trans-

membrane (Domain II) and the nucleotides encoding cytoplasmic domain (Domain I) of the murine CD40 ligand gene. Examples of such preferred chimeric accessory molecules are shown in FIG. 2. An exemplary nucleotide sequence for such

a gene is SEQ ID NO: 7. In other chimeric accessory molecule ligand genes of the present invention, the nucleotides encoding the extracellular domains (Domains III and IV) of the murine CD40 ligand gene may be operatively linked to nucleotides encoding the transmembrane (Domain II) and cytoplasmic domain (Domain I) of the human CD40 ligand gene. An exemplary nucleotide sequence for such a gene is SEQ ID NO: 3. In other preferred chimeric accessory molecule ligand genes of the present invention, the nucleotides encoding the extracellular domains (Domains III and IV) and transmembrane domain (Domain II) of human CD40 ligand are coupled to the nucleotides encoding cytoplasmic domain (Domain I) of murine CD40 ligand gene. An exemplary nucleotide sequence for such a gene is SEQ ID NO: 6. Other chimeric accessory molecule genes contemplated by the present invention comprise the nucleotides encoding the extracellular domains (Domains III and IV) and transmembrane domain (Domain I) of the murine CD40 ligand gene operatively linked to the nucleotides encoding cytoplasmic domain of the human CD40 ligand gene. An exemplary nucleotide sequence for such a gene is SEQ ID NO: 5. Other chimeric accessory molecule ligand genes are contemplated by the present invention in which the human CD40 ligand gene extracellular domains (Domain III and IV) is operatively linked to the murine CD40 ligand gene transmembrane domain (Domain I) which is operatively linked to the human CD40 ligand gene cytoplasmic domain (Domain I). An exemplary nucleotide sequence for such a gene is SEQ ID NO: 4.

One of ordinary skill in the art will understand that many more combinations which utilize domains or other selected segments of any of the accessory molecule ligand genes

including the human CD40 ligand genes and the mouse CD40 ligand genes are possible. Such additional chimeric accessory molecule genes would include the following genes: chimeric accessory molecule genes in which the nucleotides encoding Domain I are selected from a particular accessory molecule ligand gene and operatively linked, either directly or by an additional nucleotide sequence to the nucleotides encoding Domain II from a particular accessory molecule ligand gene. These domains then would be operatively linked either directly or by an additional nucleotide sequence to the nucleotides encoding Domain III from a particular, accessory molecule ligand gene. This molecule would then be operatively linked either directly or by an additional nucleotide sequence to the nucleotides encoding Domain IV of a particular accessory molecule ligand gene. The chimeric accessory molecule ligand gene constructed in this manner may have additional nucleotides on either end or between domains which are useful to provide different amino acids in these positions. One of ordinary skill in the art will understand that these particular combinations are merely illustrations and that numerous other combinations could be contemplated in which gene segments comprising nucleotides encoding less than the entire domain of an accessory molecule are exchanged between different accessory molecules.

The present invention also contemplates chimeric accessory molecule ligand genes which are comprised of gene segments of mouse or human CD40 ligand in combination with gene segments derived from Fas ligand, TNF_α , TNF_β , CD70, CD30L, 4-1BBL, nerve growth factor or TNF-related apoptosis inducing ligand (TRAIL). Particularly useful chimeric accessory molecule ligand genes comprise at least one gene segment which is derived from a murine CD40 ligand gene together with gene segments or a gene segments derived from a different accessory molecule ligand gene.

The present invention also contemplates chimeric accessory molecule ligand genes in which the accessory molecules produced have been modified to remove amino acids within the chimeric accessory molecule that are used by post-translational mechanisms to regulate the level of expression of the accessory molecule or accessory molecule protein on a particular cell. The sites removed from the chimeric accessory molecules or chimeric molecule may include amino acids or sites which make up protease cleavage sites including metallothionine proteases, serine proteases and other proteases that recognize an amino acid sequence either specifically or non-specifically. In particular preferred embodiments, amino acids in Domain III which make up potential or actual recognition site(s) used by post-translational regulatory mechanisms have been modified or removed.

The present invention also contemplates chimeric accessory molecule ligand genes in which the domains, subdomain fragments or other amino acid residues have been taken from one accessory molecule ligand gene and moved into a second accessory molecule ligand gene from the same species. For example, in this particular embodiment, the human Domain I, and the human Domain II from the CD40 ligand molecule may be operatively linked to the nucleotides encoding the human Domain III from, for example, the CD70 molecule which is in turn operatively linked to human Domain IV for the CD40 ligand molecule. This chimeric accessory molecule therefore contains human CD40L Domains I, II and IV and human CD70 Domain III. An exemplary nucleotide sequence for such a gene is SEQ ID NO: 19. One of ordinary skill in the art will understand that a number of such combinations using domains from the same species from different accessory mol-

ecule ligand genes may create a number of chimeric accessory molecule genes which may all have specific activities and properties.

The present invention contemplates chimeric accessory molecule ligand genes in which the Domain III of a particular accessory molecule ligand gene has been replaced with a Domain III from a different accessory molecule ligand gene. In one particularly preferred embodiment, the mouse Domain III has been used to replace the human Domain III in the CD40 ligand molecule. This chimeric accessory molecule therefore contains the human CD40L Domain I, the human CD40L Domain II, mouse CD40L Domain III, and human CD40L Domain IV. An exemplary nucleotide sequence for such a gene is SEQ ID NO: 20.

The present invention also contemplates the use of chimeric accessory molecules that contain man-made amino acid sequences inserted into or in place of a portion of a domain or other amino acid sequence of an accessory molecule gene. These man-made amino acid segments may be created by selecting any amino acid sequence that may be used to give the accessory molecule a particular function or to remove another undesired function. These man-made amino acid segments are produced by inserting into the accessory molecule ligand gene or chimeric accessory molecule ligand gene the nucleotide sequences required to encode those particular man-made amino acid segments in the desired positions. Further, the chimeric accessory molecule ligand genes may contain nucleotide segments which comprise sub-domain segments of other molecules or small segments in which amino acids have been changed for a desired purpose. The use of sub-domain nucleotide segments allows the introduction of short amino acid sequences derived from other molecules into chimeric accessory molecules of the present invention. The incorporation of such short sub-domain segments or amino acid changes into the accessory-molecule ligand allows the introduction of desired or the removal of undesired features of that molecule.

The identification of domain structures within accessory cell molecules is well known in the art and generally requires the identification of cysteine residues within the accessory molecules and the subsequent mapping of disulfide bonds between various cysteine residues. The mapping of various sub-domain segments of an accessory molecule is well known in the art and involves analysis of the amino acid sequence of the accessory molecules and generally involves a comparison of the crystal structure of tissue necrosis factor with the use of predictive algorithms thereby producing a predicted structure of a chimeric accessory molecule or an accessory molecule. This predicted structure of these molecules can then be used to select various sub-domain portions of the molecule to be used to construct further chimeric accessory molecules. Examples of such mapping studies include the studies by M. Pitsch and C. V. Jongeneel, *International Immunology*, 5:233-238 (1993) and the analysis shown in FIG. 1.

The present invention also contemplates accessory molecule ligand genes and chimeric accessory molecule ligand genes which are truncated and encode less than the full length of the amino acid sequence found in the native accessory molecule ligand. These truncations may alter the properties of the accessory molecule ligand gene but some identified activity is maintained. Such truncations may be made by removing a gene segment or gene segments from the accessory molecule gene and typically would be performed by removing nucleotides encoding domains which are not directly involved in the binding of the accessory molecule ligand with its accessory molecule. These truncated accessory molecule

ligand genes or chimeric truncated accessory molecule ligand genes may contain further gene segments which encode amino acid segments or domains which replace the domains removed from that truncated accessory molecule gene. However, such replacement of the portions of the accessory molecule removed by truncation is not necessary.

The chimeric accessory molecule genes of the present invention may be constructed using standard genetic engineering methods to operatively link a particular nucleotide sequence from one accessory molecule ligand gene to a different nucleotide sequence derived from the same or different accessory molecule ligand gene. In addition, standard genetic engineering methods may be used to insert man-made nucleotide sequences or sub-domain nucleotide sequences into the chimeric accessory molecule ligand gene. One of ordinary skill in the art will understand that various methods may be utilized to produce such chimeric accessory molecule genes. For example, a gene conversion method known as "SOEN" may be used to produce a chimeric accessory molecule gene which contains nucleotide segments derived from different chimeric accessory molecules. The methods for using this gene conversion method are well known in the art and have been described for example in Horton, R. M., *Mol. Biotechnol.*, 3:93 (1995); Ali, S. A. and A. Steinkasserer, *Biotechniques*, 18:746 (1995); Vilardaga, J. P., E. Di Paolo, and A. Bollen, *Biotechniques*, 18:604 (1995); Majumder, K., F. A. Fattah, A. Selvapandian, and R. K. Bhatnagar, *PCR. Methods Appl.*, 4:212 (1995); Boles, E. and T. Miosga, *Curr. Genet.* 28:197 (1995); Vallejo, A. N., R. J. Pogulis, and L. R. Pease, *PCR. Methods Appl.*, 4:S123 (1994); Henkel, T. and P. A. Baeuerle, *Anal. Biochem.*, 214:351 (1993); Tessier, D. C. and D. Y. Thomas, *Biotechniques*, 15:498 (1993); Morrison, H. G. and R. C. Desrosiers, *Biotechniques*, 14:454 (1993); Cadwell, R. C. and G. F. Joyce, *PCR. Methods Appl.*, 2:28 (1992); and, Stappert, J., J. Wirsching, and R. Kemler, *Nucleic Acids Res.*, 20:624 (1992). Alternatively, one of ordinary skill in the art will understand that site-directed mutagenesis may be used to introduce changes into a particular nucleotide sequence to directly produce or indirectly be used to produce a chimeric accessory molecule gene of the present invention. For example, the mutagen kit provided by BioRad Laboratories may be used together with the methods and protocols described within that kit to produce the desired changes in the nucleotide sequence. These methods were originally described by Kunkel, *Proc. Natl. Acad. Sci. USA*, 82:488-492 (1985) and Kunkel et al., *Meth. Enzol. Mol.*, 154:367-382 (1987). By using the site directed mutagenesis protocols described herein and known within the art, a skilled investigator may induce individual nucleotide changes which result in an altered amino acid sequence or which preserve an amino acid sequence but introduce a desired restriction enzyme recognition sequence into the gene. This new restriction endonuclease recognition site may then be used to cut the gene at that particular point and use it to a gene or segment of another accessory molecule ligand gene. In addition to these methods, one of ordinary skill in the art will understand that an entire chimeric accessory molecule ligand gene may be synthesized using synthetic methods known in the art. This methodology only requires that the skilled artisan generating nucleotide sequence of a chimeric accessory molecule ligand gene and provide that sequence to a company which is capable of synthesizing such a gene.

B. Genetic Constructs

The present invention contemplates the use of accessory molecule ligand genes or chimeric accessory molecule ligand genes which are present in various types of genetic vectors. A genetic vector refers to a DNA molecule capable of auton-

mous replication in a cell into which another DNA segment can be inserted to cause the additional DNA segments to replicate. Vectors capable of expressing genes contained in that vector are referred to as "expression vectors." Thus, the genetic vectors and expression vectors of the present invention are recombinant DNA molecules which comprise at least two nucleotide sequences not normally found together in nature.

The genetic vectors useful in the present invention contain an accessory molecule ligand gene which encodes an accessory molecule ligand which is optionally operatively linked to a suitable transcriptional or translational regulatory nucleotide sequence, such as one derived from a mammalian, microbial, viral, or insect gene. Such regulatory sequences include sequences having a regulatory role in gene expression, such as a transcriptional promoter or enhancer, an operator sequence to control transcription, a sequence encoding a ribosomal binding site within the messenger RNA and appropriate sequences which control transcription, translation initiation or transcription termination.

Particularly useful regulatory sequences include the promoter regions from various mammalian, viral, microbial, and insect genes. The promoter region directs an initiation of transcription of the gene and causes transcription of DNA through and including the accessory molecule ligand gene. Useful promoter regions include the promoter found in the Rous Sarcoma Virus (RSV)-long terminal repeat (LTR), human cytomegalovirus (HCMV) enhancer/promoter region lac promoters, and promoters isolated from adenovirus, and any other promoter known by one of ordinary skill in the art would understand to be useful for gene expression in eukaryotes, prokaryotes, viruses, or microbial cells. Other promoters that are particularly useful for expressing genes and proteins within eukaryotic cells include mammalian cell promoter sequences and enhancer sequences such as those derived from polyoma virus, adenovirus, simian virus 40 (SV40), and the human cytomegalovirus. Particularly useful are the viral early and late promoters which are typically found adjacent to the viral origin of replication in viruses such as the SV40. Examples of various promoters which have been used in expression vectors have been described by Okama and Berg (*Mol. Cell. Biol.* 3:280, 1983), the pMLSVN SV40 described by Kossman et al., *Nature* 312:768 (1984). One of ordinary skill in the art will understand that the selection of a particular useful promoter depends on the exact cell lines and the other various parameters of the genetic construct to be used to express the accessory molecule ligand gene or the chimeric accessory molecule ligand gene within a particular cell line. In addition, one of ordinary skill in the art will select a promoter which is known to express genes in the target cell at a sufficiently high level to be useful in the present invention.

The genetic vectors and expression vectors of the present invention optionally contain various additional regulatory sequences including ribosome binding sites which allow the efficient translation of the messenger RNA produced from an expression vector into proteins, the DNA sequence encoding various signals peptides which may be operatively linked to the accessory molecule ligand gene or the chimeric accessory molecule ligand gene. The signal peptide, if present, is expressed as a precursor amino acid which enables improved extracellular secretion of translation fusion polypeptide.

The genetic constructs contemplated by the present invention therefore include various forms of accessory molecule ligand genes described above which are operatively linked to either a promoter sequence or a promoter and enhancer sequence and also operatively linked to a polyadenylation sequence which directs the termination and polyadenylation

of messenger RNA. It is also contemplated that the genetic constructs of the present invention will contain other genetic sequences which allow for the efficient replication and expression of that construct within the desired cells. Such sequence may include introns which are derived from native accessory molecule ligand genes or, for example, from a virus gene.

The present invention also contemplates gene therapy vectors which are able to directly infect mammalian cells so as to introduce the desired accessory molecule ligand gene or chimeric accessory molecule ligand gene into that cell. These gene therapy vectors are useful for directly infecting cells which have been isolated from an animal or patient, or can be directly introduced into an animal or patient and thereby directly infect the desired cell within that animal or patient.

Many types of gene therapy vectors which are able to successfully transfer genes and cause the expression of desired foreign DNA sequences have been developed and described in the literature. For example, the article entitled "Gene Transfer Vectors for Mammalian Cells" in *Current Comm. Mol. Biol.*, Cold Springs Harbor Laboratory, New York (1987). Further, naked DNA can be physically introduced into eukaryotic cells including human cells by transfection using any number of techniques including calcium phosphate transfection (Berman et al., *Proc. Natl. Acad. Sci. USA*, 81:7176 (1984)), DEAE-Dextran Transfection, protoplast fusion (Deans et al., *Proc. Natl. Acad. Sci. USA*, 81:1292 (1984)), electroporation, liposome fusion, polybrene transfection and direct gene transfer by laser micropuncture of the cell membrane. In addition, one of ordinary skill in the art will understand that any technique which is able to successfully introduce the DNA into a cell in such a manner as to allow it to integrate into the genome of a cell and allow the expression of the desired gene would be useful in the present invention.

Specifically, gene therapy vectors which utilize recombinant infectious virus particles for gene delivery have been widely described. See, for example, Brody, S. L. and R. G. Crystal, *Ann. N. Y. Acad. Sci.*, 716:90 (1994); Srivastava, A., *Blood Cells*, 20:531 (1994); Jolly, D., *Cancer Gene Ther.*, 1:51 (1994); Russell, S. J., *Eur. J. Cancer*, 30A:1165 (1994); Yee, J. K., T. Friedmann, and J. C. Burns, *Methods Cell Biol.*, 43 Pt A:99 (1994); Boris-Lawrie, K. A. and H. M. Temin, *Curr. Opin. Genet. Dev.*, 3:102 (1993); Tolstoshev, P., *Annu. Rev. Pharmacol. Toxicol.*, 33:573 (1993); and, Carter, B. J., *Curr. Opin. Biotechnol.*, 3:533 (1992). The present invention contemplates the use of gene therapy vectors to carry out the desired methodology of the present invention by introducing a gene encoding an accessory molecule ligand gene or a chimeric accessory molecule ligand gene into the cell. Many viral vectors have been defined and used as gene therapy vectors and include virus vectors derived from simian virus 40 (SV40), adenoviruses, adeno-associated viruses, and retroviruses. One of ordinary skill in the art will understand that useful gene therapy vectors are vectors which are able to directly introduce into the target cells the DNA which encodes the accessory molecule ligand and allow that DNA to persist in the cell so as to express the accessory molecule ligand in the desired manner within the cell.

The gene therapy vectors of the present invention are useful for introducing accessory molecule ligand genes into a variety of mammalian cells including human cells. The particular cells infected by the gene therapy vector will, depend on the various specifics of the vector and such vectors can be used to introduce the accessory molecule ligand genes of the present invention into hematopoietic or lymphoid stem cells, antigen presenting cells, embryonic stem cells, and other cells which are capable of presenting antigen within the immune system

including cells which have CD40 on their surface. Further, such gene therapy vectors are able to introduce a gene encoding an accessory molecule ligand gene into a human neoplastic cell such as a lymphoma, leukemia, AML, CLL, CML, AMML, CMML, breast cancer, lung cancer, ovarian cancer or any tumor capable of acting as antigen presenting cells or cells which can stimulate bystander antigen presenting cells. Further, the contemplated gene therapy vectors may be used to introduce the accessory molecule ligand genes of the present invention into cells which have been engineered to make those cells capable of presenting antigen to the immune system.

III. Cells Containing Genetic Constructs Encoding an Accessory Molecule Ligand or Chimeric Accessory Molecule Ligand

The present invention also contemplates various cells which contain the genetic constructs of the present invention. These cells contain the constructs which encode the accessory molecule ligand gene and thus contain the various genetic elements described in Section II.B. above. These cells may be microbial cells, eukaryotic cells, insect cells, and various mammalian cells including human cells. In preferred embodiments of the present invention, these cells include various neoplastic cells including human neoplastic cells. These neoplastic cells may be of any cell type and include cells of the immune system, and other blood cells. Particularly preferred are any neoplastic-cells which may function as an antigen presenting cells within the immune system or which may stimulate bystander antigen presenting cells by expression of a transgenic accessory cell molecule of the present invention. Typically these neoplastic which are able to function to present antigen to the immune system have or have had an accessory molecule, such as the CD40 molecule, on the cell surface. Generally, these cells are naturally capable of presenting antigen to the immune system, but the present invention also contemplates the introduction of accessory molecule ligand genes into a cell which is not naturally able to present antigen to the immune system but which has been genetically engineered to make that cell capable of presenting antigen to the immune system. Typically, these cells include various known cell types such as monocytes, macrophages, B cells, Langerhans cells, interdigitating cells, follicular dendritic cells or Kupffer cells and the like which have become neoplastic. In addition, the present invention also contemplates cells from various carcinomas, breast, ovarian and lung cancers which contain the genetic constructs described herein. In other preferred embodiments, an accessory molecule ligand gene of the present invention is placed into cells which may be injected into a treatment site such as a tumor bed or joint. For example, the accessory molecule ligand gene of the present invention may be inserted into a fibroblast cell and the accessory molecule ligand expressed on the surface of that cell. The fibroblasts are then injected into the treatment site and cause the desired immuno effect due to the presence of the accessory molecule ligand on the surface of those cells. These cells stimulate other immune cells present in that treatment site (bystander cells). This process then results in the desired effect on the immune system.

IV. Methods Utilizing Genetic Vectors and Constructs Containing an Accessory Molecule Ligand Gene

The present invention contemplates methods of altering the immunoreactivity of human cells using a method which

includes introducing a gene encoding an accessory molecule ligand gene into the human cells so that the accessory molecule ligand encoded by that gene is expressed on the surface of those cells. The present invention is useful for any human cells which participate in an immune reaction either as a target for the immune system or as part of the immune system which responds to the foreign target. A large variety of methods are contemplated in which the final result is that the accessory molecule ligand gene is introduced into the desired cells. These methods include ex vivo methods, in vivo methods and various other methods which involve injection of DNA, genetic vectors or gene therapy vectors into the animal or human, including injection directly into the tumor bed present in any animal or human.

Ex vivo methods are contemplated wherein the cells into which the accessory molecule ligand gene is to be introduced are isolated from the animal or patient and then the gene is introduced into those isolated cells using suitable methods. Examples of useful ex vivo methods have been described for example by Raper, S. E., M. Grossman, D. J. Rader, J. G. Thoene, B. J. Clark, D. M. Kolansky, D. W. Muller, and J. M. Wilson, *Ann. Surg.*, 223:116 (1996); Lu, L., R. N. Shen, and H. E. Broxmeyer, *Crit. Rev. Oncol. Hematol.*, 22:61 (1996); Koc, O. N., J. A. Allay, K. Lee, B. M. Davis, J. S. Reese, and S. L. Gerson, *Semin. Oncol.*, 23:46 (1996); Fisher, L. J. and J. Ray, *Curr. Opin. Neurobiol.*, 4:735 (1994); and, Goldspiel, B. R., L. Green, and K. A. Calis, *Clin. Pharm.*, 12:488 (1993). D. Dilloo et al., in *Blood* 90:1927-1933 (1997), describe a method, using CD40L-activated cells, for treating B-acute lymphoblastic leukemia (ALL). They cocultured leukemia cells with fibroblasts infected with a retroviral vector encoding CD40L, then injected the cell mix into mice. Such an approach, if taken in humans, would differ from that contemplated here in that the therapeutic cells are stimulated in vitro, by another cell line expressing the accessory molecule ligand. Schultze, J. L. et al., in *Blood* 89:3806-3816 (1997), describe a method for stimulating T-TILs (tumor-infiltrating T cells) cytotoxic for follicular lymphoma (FL) cells by exposing them, in vitro, to FL B cells which were previously cultured with CD40L-expressing fibroblasts. They propose an adoptive immunotherapy in which T-TILs stimulated in this manner are transfused into patients. This method also requires in vitro stimulation, of the cells to be transfused, with another cell line expressing an accessory molecule.

Following the introduction of the gene, including any optional steps to assure that the accessory molecule ligand gene has been successfully introduced into those isolated cells, the isolated cells are introduced into the patient either at a specific site or directly into the circulation of the patient. In preferred embodiments of the present invention, cell surface markers, including molecules such tumor markers or antigens identify the cells are used to specifically isolate these molecules from the patient. One of ordinary skill in the art will understand that such isolation methods are well known and include such methodologies as fluorescence activated cell sorting (FACS), immunoselection involving a variety of formats including panning, columns and other similar methods.

The present invention also contemplates introducing the accessory molecule ligand gene into the desired cells within the body of an animal or human patient without first removing those cells from the patient. Methods for introducing genes into specific cells in vivo, or within the patient's body are well known and include use of gene therapy vectors and direct injection of various genetic constructs into the animal or patient. Examples of useful methods have been described by Danko, I. and J. A. Wolff, *Vaccine*, 12:1499 (1994); Raz, E., A. Watanabe, S. M. Baird, R. A. Eisenberg, T. B. Parr, M.

Lotz, T. J. Kipps, and D. A. Carson, *Proc. Natl. Acad. Sci. U.S.A.*, 90:4523 (1993); Davis, H. L., R. G. Whalen, and B. A. Demeneix, *Hum. Gene Ther.*, 4:151 (1993); Sugaya, S., K. Fujita, A. Kikuchi, H. Ueda, K. Takakuwa, S. Kodama, and K. Tanaka, *Hum. Gene Ther.*, 7:223 (1996); Prentice, H., R. A. Kloner, Y. Li, L. Newman, and L. Kedes, *J. Mol. Cell Cardiol.*, 28:133 (1996); Soubrane, C, R. Mouawad, O. Rixe, V. Calvez, A. Ghoumari, O. Verola, M. Weil, and D. Khayat, *Eur. J. Cancer*, 32A:691 (1996); Kass-Eisler, A., K. Li, and L. A. Leinwand, *Ann. N. Y. Acad. Sci.*, 772:232 (1995); DeMatteo, R. P., S. E. Raper, M. Ahn, K. J. Fisher, C. Burke, A. Radu, G. Widera, B. R. Claytor, C. F. Barker, and J. F. Markmann, *Ann. Surg.*, 222:229 (1995); Addison, C. L., T. Braciak, R. Ralston, W. J. Muller, J. Gaudie, and F. L. Graham, *Proc. Natl. Acad. Sci. U.S.A.*, 92:8522 (1995); Hengge, U. R., P. S. Walker, and J. C. Vogel, *J. Clin. Invest.*, 97:2911 (1996); Feigner, P. L., Y. J. Tsai, L. Sukhu, C. J. Wheeler, M. Manthorpe, J. Marshall, and S. H. Cheng, *Ann. N. Y. Acad. Sci.*, 772:126 (1995); and, Furth, P. A., A. Shamay, and L. Hennighausen, *Hybridoma*, 14:149 (1995). In a typical application, a gene therapy vector containing an accessory molecule ligand gene is introduced into the circulation or at a localized site of the patient to allow the gene therapy vector to specifically infect the desired cells. In other preferred embodiments the gene therapy vector is injected directly into the tumor bed present in an animal which contains at least some of the cells into which the accessory molecule ligand gene is to be introduced.

The present invention also contemplates the direct injection of DNA from a genetic construct which has a promoter and accessory molecule ligand gene followed by a polyadenylation sequence into a patient or animal. Examples of such useful methods have been described by Vile, R. G. and I. R. Hart, *Ann. Oncol.*, 5 Suppl 4:59 (1994). The genetic construct DNA is directly injected into the muscle or other sites of the animal or patient or directly into the tumor bed of the animal or patient. Alternatively, DNA from a genetic construct containing at least an accessory molecule ligand gene is used and directly injected into the animal.

In preferred embodiments of the present invention, the immune reaction or response of a human patient or animal is altered by introducing the accessory molecule ligand gene into cells, including human cells which have an accessory molecule present on the cell surface. Such cells include human cells, human antigen presenting cells and optionally these cells may be neoplastic antigen presenting cells which have the capacity to express the accessory molecule on the surface of the cell or cells which are capable of stimulating. In some embodiments, the amount of accessory molecule present on the surface of the cells into which the accessory molecule ligand gene is to be introduced is very small and such small amounts of the accessory molecule may result from down-regulation of that accessory molecule on the surface of such cells. In some embodiments, the cells into which the accessory molecule ligand gene is introduced have at least low levels of the CD40 molecule present on the cell surface or are derived from cells, which did express the CD40 ligand molecule on the cell surface but have reduced or eliminated that expression.

The preferred methods of altering the immunoreactivity of a particular cell are applicable to mammalian cells including human cells. These human cells may include neoplastic human cells such as human lymphomas, leukemias, and other malignancies including breast, lung and ovarian cancers. In some preferred embodiments the cells are normal antigen presenting cells of a human patient such as monocytes, macrophages, B cells, Langerhans cells, interdigitating cells, follicular dendritic cells, Kupffer cells, and other similar cells. In

25

preferred embodiments, the cells are lymphocytes which acquire altered immunoreactivity when the accessory molecules of the present invention are introduced into those cells. In other preferred embodiments, the cells may be neoplastic or normal cells which are capable of stimulating bystander antigen presenting cells when the accessory molecule ligand genes of the present invention are introduced into these cells. The present invention also contemplates that cells which are not naturally capable of presenting antigen to the immune system may be genetically engineered to introduce the genes encoding the molecules required for antigen presentation, including genes encoding an accessory molecule, and thus allow these cells to act as artificial antigen presenting cells. The accessory molecule ligand gene may then be introduced into these artificial antigen presenting cells. Various tests are well known in the literature to determine whether a particular cell is able to function as an antigen presenting cell, such as cell proliferation or the production of lymphokines and therefore this aspect of the present invention may be easily determined.

In addition to the above normal human cells, the present invention also contemplates introducing the accessory molecule ligand gene into various neoplastic or malignant cells which optionally are antigen presenting cells. Such human neoplastic cells which are contemplated include leukemias, lymphomas, AML, AMML, or CMML, CML, CLL and any neoplastic cell which is capable of stimulating bystander antigen presenting cells when an accessory molecule ligand is introduced into that cell. Also contemplated are neoplastic cells such as a breast, ovarian or lung cancer cell which is capable of or is engineered to act as an antigen presenting cell. However, the present immunomodulation also applicable to other malignancies not specifically identified and thus would include any tumor of any cell capable of presenting antigen within the animal or human immune system or any cell which is capable of acting as an antigen presenting cell or capable of stimulating bystander antigen presenting cells after an accessory molecule ligand gene has been introduced into those cells. Generally these antigen presenting cells have accessory molecules on the surface of the cells.

The present methods of altering the immunoreactivity of a human or animal cell contemplate the introduction of an accessory molecule ligand gene into the cells for which altered immunoreactivity is desired. The genes useful in the present invention include the wide range of accessory molecule ligand genes and chimeric accessory molecule ligand genes identified above and in preferred embodiments include at least a portion of the murine CD40 ligand gene. In particularly preferred embodiments, the accessory molecule ligand gene introduced into the cells using the methods of the present invention is selected to correspond to the accessory molecule present on the surface of the cells for which altered immunoreactivity is desired. In one particular application of the present invention, the immunoreactivity of a cell which expresses the CD40 molecule on the cell surface would be accomplished by introducing the gene which encodes the CD40 ligand molecule and more preferably the murine CD40 ligand molecule.

The present invention also contemplates altering the immunoreactivity of human or animal cells by introducing an accessory molecule ligand gene which is a chimeric accessory molecule ligand gene into the cell. The various useful chimeric accessory molecule ligand genes were identified above and could include a wide variety of molecules and allow the unique properties of those chimeric accessory molecule ligand genes to be utilized to alter the immunoreactivity of the target cells. In preferred embodiments, useful chimeric

26

accessory molecule ligand genes are genes which encode at least a portion of the accessory molecule ligand which is capable of binding the accessory molecule present on the surface of the cells for which altered immunoreactivity is desired.

The methods of the present invention for altering the immunoreactivity contemplate the use of genetic vectors and genetic constructs including gene therapy vectors which encode an accessory molecule ligand and therefore contain an accessory molecule ligand gene. Typically, the genetic vectors and genetic constructs including the gene therapy vectors of the present invention have a promoter which is operatively linked to the accessory molecule ligand gene followed by a polyadenylation sequence. In other embodiments, the only requirement is that the genetic vectors, genetic constructs, and gene therapy vectors of the present invention contain the accessory molecule ligand gene or the chimeric accessory molecule ligand gene.

V. Methods of Treating Neoplasia

The present invention also contemplates methods of treating human neoplasia comprising inserting into a human neoplastic cell a gene which encodes an accessory molecule ligand so that the accessory molecule ligand is expressed on the surface of the neoplastic cells. The present invention contemplates treating human neoplasia both in vivo, ex vivo and by directly injecting various DNA molecules containing a gene which encodes an accessory molecule ligand into the patient. However, at a minimum, the present methods for treating human neoplasia involve inserting the gene encoding the accessory molecule ligand into the neoplastic cells in such a way as to allow those neoplastic cells to express the accessory molecule ligand on the cell surface. The expression of the accessory molecule ligand gene in these neoplastic cells modulates the immune system to cause the neoplasia to be reduced or eliminated.

In a preferred method of treating human neoplasia, the method further comprises the steps of first obtaining the human neoplastic cells from a human patient and then inserting into the isolated human neoplastic cells a gene which encodes an accessory molecule ligand so that the accessory molecule ligand is expressed on the surface of the neoplastic cells. The human neoplastic cells having the accessory molecule ligand on the surface of that cell are then infused back into the human patient. One of ordinary skill in the art will understand that numerous methods are applicable for infusing the altered human neoplastic cells containing the gene encoding the accessory molecule ligand back into the patient and that these methods are well known in the art.

The contemplated methods of treating human neoplasia are applicable to a wide variety of human neoplasias including lymphomas, leukemias, and other malignancies. In preferred embodiments the human neoplasia is a neoplasia which involves the antigen presenting cells of the human immune system and includes monocytes, macrophages, B cells, Langerhans cells, interdigitating cells, follicular dendritic cells, Kupffer cells, and the like. In other preferred embodiments, the human neoplasia is a leukemia, a lymphoma, AML, AMML, CMML, CML or CLL, lung cancer, breast cancer, ovarian cancer and other similar neoplasias.

The genetic vectors, genetic constructs and gene therapy vectors useful in the methods of treating human neoplasia of the present invention have been disclosed above and include constructs in which a promoter is operatively linked to the accessory molecule ligand gene or the chimeric accessory molecule ligand gene which is in turn operatively linked to a

polyadenylation sequence. The methods of treating human neoplasia contemplate the use of genetic constructs, genetic vectors and gene therapy vectors as described in this specification. In addition, the present invention contemplates the use of DNA which contains at least a gene encoding an accessory molecule ligand gene. This gene may or may not contain a promoter and other regulatory sequences.

In preferred embodiments of the present invention, the cells comprising the human neoplasia are located in at least one defined site termed a tumor bed within the human patient. This tumor bed typically contains the tumor or neoplastic cell together with a number of other cells which are associated with the tumor or neoplastic cells. The present invention contemplates methods of treating such human neoplasia present in a tumor bed by injecting into the tumor bed of the patient, a gene which encodes an accessory molecule ligand so that the accessory molecule ligand is expressed on the surface of the tumor cells thereby causing the cells to participate in an immune reaction. The gene which encodes the accessory molecule ligand may be present as part of a gene therapy vector, genetic construct or genetic vector.

In preferred embodiments, the accessory molecule ligand gene is a chimeric accessory molecule ligand gene which has at least a portion of the murine CD40 ligand gene is used. In other preferred embodiments, the accessory molecule ligand encoded is capable of binding an accessory molecule present on the human neoplasia to be treated.

The various gene therapy vectors used in the treatment methods of the present invention include vectors which are capable of directly infecting human cells. Such vectors have been described in the literature and are readily adaptable to the methods described in the present invention.

The present invention contemplates the use of any type of gene therapy including the methods of Raper, S. E. et al., *Ann. Surg.*, 223:116 (1996); Lu, L. et al., *Crit. Rev. Oncol. Hematol.*, 22:61 (1996); Koc, O. N. et al., *Semin. Oncol.* 23:46 (1996); Fisher, L. J. et al., *Curr. Opin. Neurobiol.*, 4:735 (1994); Goldspiel, B. R. et al., *Clin. Pharm.*, 12:488 (1993); Danko, I. et al., *Vaccine*, 12:1499 (1994); Raz, E. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 90:4523 (1993); Davis, H. L. et al., *Hum. Gene Ther.*, 4:151 (1993); Sugaya, S. et al., *Hum. Gene Ther.*, 7:223 (1996); Prentice, H. et al., *J. Mol. Cell Cardiol.*, 28:133 (1996); Soubrane, C. et al., *Eur. J. Cancer*, 32A:691 (1996); Kass-Eisler, A. et al., *Ann. N. Y. Acad. Sci.*, 772:232 (1995); DeMatteo, R. P. et al., *Ann. Surg.*, 222:229 (1995); Addison, C. L. et al., *Proc. Natl. Acad. Sci. U.S.A.*, 92:8522 (1995); Hengge, U. R. et al., *J. Clin. Invest.*, 97:2911 (1996); Feigner, P. L. et al., *Ann. N. Y. Acad. Sci.*, 772:126 (1995); Furth, P. A., *Hybridoma*, 14:149 (1995); Yovandich, J. et al., *Hum. Gene Ther.*, 6:603 (1995); Evans, C. H. et al., *Hum. Gene Ther.*, 7:1261.

VI. Methods of Vaccination

The present invention contemplates methods of vaccinating an animal against a predetermined organism comprising administering to that animal a vaccine containing immunogenic animal antigens capable of causing an immune response in that animal against the desired organism together with a vector containing a gene encoding an accessory molecule ligand. The present invention also contemplates methods of vaccinating an animal which include administering the genes which encode the immunogenic antigen capable of causing a desired immune response or altering the immune response to a particular antigen together with a vector containing a gene including the accessory molecule ligand gene. In this particular embodiment, the vector or vectors intro-

duced encode the immunogenic antigens desired and the desired accessory molecule ligand. The present invention also contemplates that the gene or genes encoding the immunogenic peptide or peptides may be present on the same vector as is the gene or genes encoding the accessory molecule ligand.

The vaccination methods of the present invention are general in that they may be used to produce a vaccination against any predetermined organism, such as a virus, a bacteria, a fungus or other organism. In addition, the present vaccination methods may be used to produce an immune response against a neoplastic cell.

In other preferred embodiments, the vaccination methods of the present invention utilize a genetic vector, a genetic construct or a gene therapy vector which contains an accessory molecule ligand gene which is a chimeric accessory molecule ligand gene. That chimeric accessory molecule ligand gene preferably contains at least a portion of the murine CD40 ligand gene. In other preferred embodiments, the vaccination method utilizes a DNA molecule which encodes at the minimum the accessory molecule ligand gene or a chimeric accessory molecule ligand gene. This particular DNA may or may not include a promoter sequence which directs the expression of the accessory molecule ligand gene.

The present invention also contemplates that the vaccination method may utilize a genetic vector which is capable of expressing an accessory molecule ligand within a particular cell or organism together with a vector which is capable of expressing at least a single polypeptide from an andovirus. This andovirus polypeptide may be expressed from the same or different vector which expresses the accessory molecule ligand in that cell. In this particular embodiment, the andovirus polypeptide is also expressed in at least one cell type within the organism and serves to modulate the immune response found in response to this vaccination protocol.

The present invention also contemplates the introduction of an accessory molecule ligand gene into cells which are present in the joints of patients with rheumatoid arthritis. In preferred embodiments, the accessory molecule ligand gene introduced comprises at least a portion of the Fas ligand gene and upon expression the accessory ligand induces the cell death of cells expressing Fas on the cell surface. This process leads to the reduction of the destructive inflammatory process.

The following examples are provided to illustrate various aspects of the present invention and do not limit the scope of that invention.

VII. Methods of Treating Arthritis

The present invention also contemplates methods of treating arthritis comprising inserting into a joint, cells which have been transformed with an accessory molecule, such as the Fas ligand. In preferred embodiments, the expression of that accessory molecule ligand or the stability of that molecule on the surface of the cells has been altered. In these preferred embodiments, the accessory molecule ligand functions in an enhanced manner to aid in the treatment of arthritis within the joint. The present invention contemplates treating human arthritis both in vivo, ex vivo, and by directly injecting various DNA molecules containing genes which encode the useful accessory molecule ligand into the patients. Various useful protocols may be designed to rheumatoid arthritis including those described in the example section below.

The present invention contemplates the treatment of arthritis utilizing accessory molecule ligand genes which may be chimeric accessory molecule ligand genes comprised of por-

tions of that gene being derived from two different accessory molecule ligand genes. In other embodiments, the chimeric accessory molecule ligands may be produced by utilizing domains from the same accessory molecule ligand gene. The resulting chimeric accessory molecule ligands have an altered stability on the surface of cells upon which they are expressed. This altered stability modulates the function of the immune system in the local environment around the cells in which these chimeric accessory molecule ligands are expressed. For example, in certain preferred embodiments, Fas ligand stability is altered on the surface of cells within a joint of a patient suffering from arthritis. This altered stability modulates the immune system and causes the cells to be targeted for apoptosis and thus reducing the immune response within the inflamed joint. In other embodiments, the accessory molecule ligand genes described within are altered such that the resulting accessory molecule ligand has an altered stability and causes an immunomodulatory effect which can be useful in the treatment of arthritis.

The present invention contemplates in preferred embodiments that chimeric accessory molecule ligands genes be utilized in the treatment of arthritis. These chimeric accessory molecule ligand genes preferably contain at least a portion of the Fas ligand gene Domain IV, which carries the effect or function for Fas ligand. In preferred embodiments, at least in the portion of that domain, is present which allows Fas ligand to have its biologic effects. In other preferred chimeric accessory molecule ligands, those ligands contain domains from other accessory molecule ligand genes of the present invention or from a different domain of the same accessory molecule ligand. Particularly preferred are Fas chimeric accessory molecule ligand genes made up on Domain IV of the human Fas ligand operatively linked with Domain III of the mouse Fas ligand. This particular combination results in more stable Fas ligand and thus, by replacing Domain III of human Fas ligand with Domain III of the mouse ligand, the activity of the human Fas ligand gene is altered.

Alternatively, in other preferred embodiments, the murine Fas ligand gene is used to encode the murine Fas ligand on the surface of cells in place of the human Fas ligand. The murine Fas ligand is more stable than the human Fas ligand and thus, alters the Fas ligand activity in the joint. The resulting altered Fas ligand activity is useful in the treatment of rheumatoid arthritis.

Further preferred embodiments include embodiments in which the effect or function present on Domain IV of the human Fas ligand is combined with other domains from other accessory molecule ligands. For example, CD70 Domain III is more stable than Domain III of the human Fas ligand and thus the chimeric accessory-molecule ligand made up of Domain III from the human CD70 and Domain IV of the Fas ligand together with other supporting domains would be more stable. The increased stability leads to increase Fas ligand activity. In other preferred embodiments, Domain III of the Fas ligand is replaced with multiple copies of a domain or domains. Such multiple copies of domains include domains made up of two or more copies of other domains such as Domains III or I of the CD70 molecule.

In other preferred embodiments, the present invention contemplates accessory molecule ligand genes, such as Fas ligand genes, in which a cleavage site for matrix-metalloproteinase (MMP), have been removed from the accessory molecule ligand. MMP cleavage and recognition sites, charted in FIG. 28, are discussed in Smith, M. M. et al., *Journal of Biol. Chem.*, 270:6440-6449 (95) and Nagase, H., and G. B. Fields, *Biopolymers (Peptide Science)*, 40:399-416 (96). In preferred embodiments, at least one MMP site has been removed from

at least Domain III of the Fas ligand gene. The removal of the MMP site from the Fas ligand gene makes the Fas ligand more stable and thus, more effective in the treatment of arthritis.

In other preferred embodiments, chimeric accessory molecule ligand genes are comprised of portions of the human Fas ligand gene with other domains from other human accessory molecule ligands or domains from accessory molecules derived from other species. For example, the present invention contemplates the use of domains from CD40 ligand, CD70 ligand, CD30 ligand, TNF-related apoptosis inducing ligand (TRAIL), TNF- α as well as mutants of human Fas ligand and murine Fas ligand. Production of such chimeric accessory molecule ligands is easily accomplished by manipulating and producing accessory molecule ligand genes which are chimeric and thus has portions derived from at least two different accessory molecule ligand genes.

Examples

1. Expression of Human and Mouse Accessory Molecule Ligand in Human CLL Cells

a. Construction of a Genetic Construct and Gene Therapy Vector Containing a Human and Mouse Accessory Molecule Ligand Gene

Either the human accessory molecule ligand gene (human CD40 ligand) or the murine accessory molecule ligand gene (murine CD40 ligand) was constructed utilizing the respective human and murine genes. Each of these genes was cloned in the following manner.

i. Murine CD40-L Cloning

Total RNA was isolated using the RNA STAT-60 kit (Tel-Test "B" Inc., Friendswood, Tex.) from 1×10^7 B6 mouse splenocytes that were previously activated for 8 hours with immobilized CD3-specific mAb. cDNA was then synthesized with the Superscript cDNA synthesis kit (Gibco BRL, Grand Island, N.Y.) using oligo-dT primers. The murine CD40 ligand (mCD40-L) gene was then amplified from the cDNA by PCR using the following mCD40-L specific primers. 5'-GTTAAGCTTTTCAGTCAGCATGATAGAA (SEQ ID NO: 26), 5'-GTTTCTAGATCAGAGTTTGAGTAAGCC (SEQ ID NO: 27). The amplified mCD40-L PCR product was subcloned into the HindIII and XbaI sites of the eukaryotic expression vector pcDNA3 (Invitrogen, San Diego, Calif.). A DNA fragment encompassing the CMV promoter, mCD40-L gene, and polyadenylation signal was released from this plasmid construct after restriction digestion with BglII and XhoI enzymes. This DNA fragment was then subcloned into the shuttle plasmid MCS(SK) pXCX2 (Spessot R, 1989, *Virology* 168:378) that was designated mCD40-L pXCX2. This plasmid was used for adenovirus production as described below.

ii. Human CD40-L Cloning

A plasmid containing the gene for human CD40-L was used to produce the human CD40-L gene used herein. The sequence of this gene is available and thus this source of the gene was used merely for convenience. See GenBank accession no. X67878. This plasmid was used for PCR amplification of the human CD40-L gene using the specific primers, sense primer 5'CCAAGACTAGTTAACACAGCATGATCGAAA 3' (SEQ ID NO: 28) and antisense primer 5' CCAATGCGGCCGCACTCAGAATTCAACCTG 3' (SEQ ID NO: 29).

These primers contain flanking restriction enzyme sites for subcloning into the eukaryotic expression plasmid pRc/CMV (Invitrogen). The PCR amplified CD40-L fragment was subcloned into the SpeI and NotI sites of pRc/CMV and designated hCD40-L pRc/CMV. A BglII and XhoI fragment

encompassing the CMV promoter, hCD40-L gene, and polyadenylation signal was then released from this plasmid and subcloned into the shuttle plasmid MCS(SK)pXCX2 as described above. This plasmid was designated hCD40-L pXCX2. This plasmid was used for adenovirus production as described below.

iii. Adenovirus Synthesis

Either mCD40-L pXCX2 or hCD40-L pXCX2 plasmids were co-transfected with pJM17 (Graham and Prevec, 1991, *Methods in Molecular Biology*, Vol 7) into 293 cells (American Type Culture Collection, Rockville, Md.) using the calcium phosphate method (Sambrook, Fritsch, and Maniatis, 1989, *Molecular Cloning, A Laboratory Manual*, 2nd edition, chapter 16:33-34). Isolated adenovirus plaques were picked and expanded by again infecting 293 cells. High titer adenovirus preparations were obtained as described (Graham and Prevec, 1991, *Methods in Molecular Biology*, Vol 7), except for the following modifications. The cesium chloride gradient used for concentrating viral particles was a step gradient, with densities of 1.45 g/cm³ and 1.2 g/cm³. The samples were spun in a SW41 rotor (Beckman, Brea, Calif.) at 25,000 rpm at 4° C. The viral band was desalted using a Sephadex G25 DNA grade column (Pharmacia, Piscataway, N.J.). The isolated virus was stored at 70° C. in phosphate buffered saline with 10% glycerol. The virus titer was determined by infecting 293 cells with serial dilutions of the purified adenovirus and counting the number of plaques formed. Viral titers typically ranged from 10¹⁰ to 10¹² plaque forming units/ml (PFU/ml).

b. Introduction of a Murine and Human Accessory Molecule Ligand Gene into CLL Cells and HeLa Cells

For adenovirus infection, 106 freshly thawed and washed CLL cells or HeLa cells were suspended in 0.5 to 1 mL of culture medium for culture at 37° C. in a 5% CO₂-in-air incubator. Adenovirus was added to the cells at varying multiplicity of infection (MOI), and the infected cells were cultured for 48 hours, unless otherwise stated, before being analyzed for transgene expression.

c. Expression of an Accessory Molecule Ligand Gene in CLL Cells and HeLa Cells

The CLL and HeLa cells which were infected with the adenovirus vector containing either mouse or human CD40 ligand genes prepared in Example 1b. were then stained with commercially available monoclonal antibodies immunospecific for either human or mouse CD40 ligand (Pharmingen, San Diego, Calif.) using the manufacturer's directions. The CLL and HeLa cells were washed in staining media (SM) consisting of RPMI-1640, 3% fetal calf serum and 0.05% sodium azide and containing propidium iodide and then analyzed on a FACScan (Becton Dickinson, San Jose, Calif.). Dead cells and debris were excluded from analysis by characteristic forward and side light scatter profiles and propidium iodide staining. Surface antigen expression was measured as the mean fluorescence intensity ratio (MFIR). MFIR equals the mean fluorescence intensity (MFI) of cells stained with a specific FITC-conjugated MoAb, divided by the MFI of cells stained with a control IgG-FITC. This method controls for the nonspecific increases in auto-fluorescence seen in larger, more activated cells.

The histograms, generated for the CLL cells and HeLa cells containing either a genetic vector containing the human CD40 ligand gene or the murine CD40 ligand gene and the appropriate controls, are shown in FIG. 3A-3D. The expression of both the murine and human accessory molecule ligand gene (CD40 ligand) in HeLa cells is shown in FIGS. 3A and 3B, respectively. The expression of the murine and human accessory molecule ligand in CLL cells is shown in FIGS. 3C and 3D. The expression of an accessory molecule ligand gene

in CLL cells and the expression of murine CD40 ligand on the surface of the CLL cells is shown in FIG. 3C. The failure of the human accessory molecule ligand to be expressed on the surface of the CLL cells is shown in FIG. 3D.

FIG. 8 shows data from an experiment done to examine whether the CD4⁺ T cells of CLL patients could be induced to express the accessory molecule ligand mRNA after CD3 ligation. An ELISA-based quantitative competitive RT-PCR was used to measure CD40 ligand transcript levels. In this experiment, CD40 ligand and RNA transcribed from the CD40 ligand gene in CLL cells are compared with levels of CD40 ligand and RNA made in normal donor cells, after induction by CD3 ligation. For CD3 activation, plate coats of CD3 mAb were made and incubated with plated CLL or normal donor mononuclear cells for the indicated amount of time, after which cells were analyzed for expression of surface antigens or CD154 RNA message levels. CLL or normal donor serum was added to the cells at the beginning of the activation assay for examination of modulation of CD40 ligand surface expression.

For quantitative CD154 RT-PCR ELISA, total RNA was extracted and competitor RNA was generated from the insert containing CD40 ligand (CD154) cDNA. Varying amounts of competitor RNA were added to separate wells of isolated total RNA that subsequently were converted into cDNA. CD3 activation, ELISAs and PCR reactions were performed as described in Cantwell, M. et al., *Nature Medicine* 3:984-989 (1997). Biotinylated PCR products were captured onto microtiter plates (Becton Dickinson, Oxnard, Calif.) coated with streptavidin (Sigma), and incubated. The plate was treated with NaOH to remove the sense strands and subsequently washed. The DNA was then hybridized with either wild-type gene-specific or competitor-specific oligonucleotides. Using terminal transferase, each probe was labeled with a molecule of digoxigenin-11-dideoxyUTP (Boehringer Mannheim). The plate was incubated and washed with HYBE buffer and blocking buffer, then peroxidase-conjugated anti-digoxigenin antibody (150 U/ml; Boehringer Mannheim) in blocking buffer was added. TMB (tetramethylbenzidine) and peroxidase (Kirkegaard and Perry Laboratories, Gaithersburg, Md.) were added for color development, and optical densities were measured at 450 nm and Deltasoft II (Biometallics, Princeton, N.J.) was used for data analysis.

Standard curves plotting the moles of RNA product versus the optical density were made for the standard cDNA reactions. The equations describing these standard curves were then used to calculate the moles of wild-type or competitor DNA present in the unknown PCR reactions based on the optical densities obtained in the ELISA readings. The ratio of the quantity of wild-type DNA to the amount of competitor DNA was then plotted against the known quantity of competitor RNA added in the initial samples. The ratio of 1 was taken for the extrapolation of the amount of unknown moles of target RNA in the sample (a ratio of 1 means the amount of target RNA versus competitor RNA are equal). The molecules of target RNA per CD4 cell was then calculated based on the following formula: [(moles target CD154 RNA) × (6 × 10²³ molecules/mole) × (dilution factor of test RNA)] / (% of CD4 T cells in total cell population).

The upper graph in FIG. 8 shows that T cells of patients with CLL do not express detectable CD40 ligand after CD3 ligation. CD40 ligand RNA is produced, but it is not stable. Although both CD40 ligand and CD40 ligand RNA are expressed in normal donor T cells (lower graph), the levels of neither the protein or RNA are stably maintained.

FIG. 9 shows a time course for surface expression of CD40 ligand. Expression reached a peak level at 48 hours after

infection and persisted at high levels for at least 6 days thereafter. In this experiment, CLL B cells were infected with a gene therapy vector containing an accessory molecule ligand, at a MOI of 1000 at time zero, and then assessed by flow cytometry at various times thereafter. At each time point listed on the abscissa, the proportions of viable CLL B cells that expressed detectable CD154 are indicated by the vertical bars corresponding to the percentage scale depicted on the right-hand ordinate.

d. Function of the Human and Murine Accessory Molecule Ligands

i. Induction of CD80 and CD54 on Cells Containing a Gene Therapy Vector Encoding an Accessory Molecule

The CLL cells infected with the murine accessory molecule ligand gene prepared in Example 1b. were then cultured in tissue culture plates. The CLL cells were then analyzed using multiparameter FACS analysis to detect induction of CD80 and CD54 expression using fluorescein isothiocyanate-conjugated monoclonal antibodies immunospecific for each of these respective surface antigens. Non-infected CLL cells were used as a control. The cells were subjected to the appropriate FACS analysis and histograms were generated. CD80 mAb was obtained from Dr. Edward Clark and CD54 mAb was purchased from CALTAG Inc. The CD80 was conjugated using standard methods which have been described in Kipps et al., *Laboratory Immunology II*, 12:237-275 (1992).

The results of this analysis are shown in FIG. 4A-4D. FIGS. 4A-4B compare the amount of CD54 expression in CLL cells which have not been transfected (FIG. 4A) or CLL cells into which a gene therapy-vector containing the murine CD40 ligand gene was introduced (FIG. 4B). The shaded graph indicates the isotype control for FACS staining and the open graph indicates the cells stained with the anti-CD54 antibody. These results show that the level of expression of CD54 is increased in CLL cells into which the gene therapy vector containing the murine CD40 ligand was introduced.

FIGS. 4C and 4D compare the amount of CD80 expression in CLL cells which have not been transfected (FIG. 4C) or CLL cells into which a gene therapy vector containing the murine CD40 ligand gene was introduced (FIG. 4D). The shaded graph indicates the isotype control for FACS staining and the open graph indicates the cells stained with the anti-CD80 antibody. These results show that the level of expression of CD80 is increased in the CLL cells into which the gene therapy vector containing the murine CD40 ligand was introduced.

In an additional experiment, CLL cells infected with a gene therapy vector containing the murine accessory molecule ligand gene were evaluated by flow cytometry for induced expression of not only CD54 and CD80, but also CD86, CD58, CD70 and CD95. Fluorescein-conjugated mAb specific for human CD54 and CD70 were purchased from CALTAG. Fluorescein-conjugated mAb specific for human CD27, CD58, CD80, CD86, or CD95, and phycoerythrin-conjugated mAb specific for human or mouse CD40 ligand, were obtained from PharMingen. Shaded histograms represent staining of CLL B cells with FITC-conjugated isotype nonspecific mAb. In contrast to uninfected CLL cells (FIG. 10, thin-lined histograms), or Ad-lacZ-infected CLL cells (data similar to that obtained with uninfected cells, but not shown), CLL cells infected with the adenovirus vector encoding the CD40 ligand (CD154) expressed high levels of CD54 (FIG. 10, top left), CD80 (FIG. 10, top middle), CD86 (FIG. 10, top right), CD58 (FIG. 10, bottom left), CD70 (FIG. 10, bottom middle), and CD95 (FIG. 10, bottom right). On the other hand, CD40 ligand-CLL (CD154 CLL) expressed significantly lower levels of both surface membrane CD27 (FIG.

11A, thick-lined histogram) and soluble CD27 (FIG. 11B) than uninfected (FIG. 11A, thin-lined histogram) ($P < 0.01$, Bonferroni t-test) or Ad-lacZ-infected CLL cells (data similar to that obtained with uninfected cells, but not shown). In the experiment shown in FIG. 11A, the CLL B cells were examined for expression of CD27 via flow cytometry, three days after infection. Shaded histograms represent staining of CLL B cells with FITC-conjugated isotype control mAb. In FIG. 11B, cell-free supernatants were collected, after the infection or stimulation of CLL B cells, for 72 hours and tested for the concentration of human CD27 by ELISA. The reduced expression of CD27 (FIG. 11B) is similar to that noted for leukemia B cells stimulated via CD40 cross-linking with mAb G28-5 presented by CD32-expressing L cells, as described in Rassenti, L. Z. and T. J. Kipps, *J. Exp. Med.* 185:1435-1445.

ii. Allogeneic T Cell Responses to CLL Cells into which a Genetic Therapy Vector Containing a Murine CD40 Ligand Gene has been Introduced

The ability of CLL cells which have been infected with a gene therapy vector containing the murine CD40 ligand gene to stimulate allogeneic T cells (i.e., from another individual) was analyzed using cell proliferation assays. Briefly, the test cells were co-cultured with the genetic therapy vector containing the lac-Z gene or the murine CD40 ligand gene at a multiplicity of infection of 1,000 in the presence of IL-4 at a concentration of 10 ng/ml. In other samples, the CLL cells were stimulated with MOPC21 (a control IgG) or G28-5 (an anti-CD40 monoclonal antibody) or were preincubated on CD32-L cells and at the same time treated with IL-4. The preincubation with the CD32-L cells together with IL-4 treatment have been shown to be an efficient form of cross-linking the CD40 molecule other than direct gene transfection.

After three days of culture at 37° C., these cells were treated with mitomycin C to prevent their proliferation and then used to stimulate allogeneic T cells. Prior to this co-culture, the different aliquots of CLL cells had either been treated with the anti-CD40 monoclonal antibody or had been infected with the gene therapy vector containing either the lac-Z or murine CD40 ligand gene at a stimulator ratio of 1:10. After two days of culture at 37° C., interferon gamma (IFN γ) production was measured by ELISA assay. After five days of co-culture at 37° C., the incorporation of ³H-thymidine into replicating cells was measured after an eight hour pulse label. The results of this assay are shown in Table II below and in FIG. 5.

In another experiment, CLL B cells infected with the gene therapy vector containing the CD40 ligand gene were evaluated for their ability to act as stimulator cells in an allogeneic mixed lymphocyte T cell reaction (MLTR). In parallel, the stimulatory capacity of control lac-Z-vector-infected CLL cells and CLL B cells that had been cultured with CD32-L cells and an anti-CD40 mAb (G28-5) or an isotype control Ig, was also examined as described in Ranheim, E. A. and T. J. Kipps, *J. Exp. Med.*, 177:925-935 (1993), Clark, E. A. and J. A. Ledbetter, *Proc. Natl. Acad. Sci. USA*, 83:4494-4498 (1986), and Banchereau, J. et al., *Science* 251:70-72 (1991). Effector T cells from a non-related donor were co-cultured with the CLL stimulator cells at an effector to target ratio of 4:1. After 18 h culture at 37° C., over 30% of the allogeneic CD3⁺ cells were found to express the activation-associated antigen CD69 when cultured with CD154-CLL cells (data not shown). In contrast, less than 4% of the T cells expressed CD69 when co-cultured with uninfected or Ad-lacZ-infected CLL cells (data not shown).

Two days after the initiation of the MLTR, the concentrations of IFN γ in the culture supernatants were assayed by

ELISA. The supernatants of the MLTR stimulated with CLL cells infected with the accessory molecule ligand CD40L (FIG. 12A, CD154-CLL) contained significantly higher levels of IFN γ (306 ± 5 ng/ml, $m \pm SE$, $n=3$) than that of MLTR cultures stimulated with the anti-CD40 mAb (FIG. 12A, α CD40-CLL) (23 ± 3 ng/ml) ($P < 0.05$, Bonferroni t-test). The latter was not significantly different from that of MLTR cultures stimulated with control Ad-lacZ-infected CLL cells (FIG. 12A, lacZ-CLL) (43 ± 10 ng/ml) ($P > 0.1$, Bonferroni t-test). The supernatants of effector cells alone, or of MLTR cultures stimulated with uninfected CLL cells (FIG. 12A, CLL) or control Ig treated CLL cells (FIG. 12A, MOPC-CLL), did not contain detectable amounts of IFN γ (< 2 ng/ml). Similarly, none of the leukemia B cell populations produced detectable amounts of IFN γ when cultured alone, without added effector T cells (data not shown).

After 5 days, cell proliferation was assessed by incorporation of ^{3} H-thymidine. Cultures with isotype control IgG-treated (FIG. 12B, MOPC-CLL) or uninfected (FIG. 12B, CLL) stimulator cells did not incorporate more ^{3} H-thymidine than cultures without added leukemia-stimulator cells (FIG. 12B, None). Ad-lacZ-infected CLL B cells (FIG. 12B, lacZ-CLL) also were unable to stimulate allogeneic T cells to incorporate amounts of ^{3} H-thymidine that were much greater than that of control cultures. In contrast, anti-CD40-stimulated leukemia cells or CD154-CLL cells each induced significant effector cell proliferation (FIG. 12B, α CD40-CLL or CD154-CLL) ($P < 0.05$, Bonferroni t-test). Moreover, the amount of ^{3} H-thymidine incorporated by cultures stimulated with CD154-CLL cells ($41,004 \pm 761$ cpm ($m \pm SE$), $n=3$) was significantly greater than that of cultures stimulated with equal numbers of α CD40-CLL cells ($22,935 \pm 1,892$ cpm, $n=3$) ($P < 0.05$, Bonferroni t test). However, neither of these mitomycin-C-treated leukemia cell populations incorporated ^{3} H-thymidine when cultured without effector T cells (data not shown). Also, as described for the MLTR between allogeneic T cells and CD40-stimulated CLL cells {6549, 7167, 7168}, allogeneic T cell proliferation in response to CD154-CLL could be inhibited by CTLA-4-Ig or CD11a mAb when added at the initiation of the MLTR, indicating that respective interactions between CD80/CD86 and CD28, or CD54 and CD11a/CD18, contribute to the noted allogeneic T cell reaction (data not shown).

TABLE II

Allogeneic T cell responses to CLL cells infected with mCD40-L adenovirus				
Stimulators	% positive cells		Allogeneic response (mean $\pm$ SEM)	
	mCD40-L	Human CD80	3H-TdR uptake (cpm)	IFN γ production (ng/ml)
None (t cells only)	—	—	3577 $\pm$ 821	n.d.*
CLL with:				
No activation	0	1.4	4577 $\pm$ 1097	n.d.
MOPC21	0	1.0	5259 $\pm$ 1788	n.d.
G28-5	0	26.7	22935 $\pm$ 1892	22.3 $\pm$ 1.6
lac-Z adeno	0	4.8	9037 $\pm$ 1781	43.2 $\pm$ 10.5
mCD40-L adeno	17.5	19.7	41004 $\pm$ 761	305.7 $\pm$ 4.5

*n.d.—not detectable

iii. Stimulation of Gamma Interferon by CLL Cells Containing an Accessory Molecule Ligand Gene

The function of CLL cells containing an accessory molecule ligand gene (mouse CD40 ligand) was analyzed by determining the ability of those cells to activate T lymphocytes. The procedure was performed as follows: allogeneic T lymphocytes from a healthy donor (greater than 90% CD3 $^{+}$) were purified using magnetic beads and monoclonal antibodies specific for the CD14 and CD19 antigen. These allogeneic T lymphocytes then were cultured together with MMC-treated CLL cells which were infected with the accessory molecule ligand gene (murine CD40 ligand) or the lac-Z gene. This co-culture was performed in RPMI-1640 medium containing 10% fetal calf serum. After culture for 24 hours, the cells were collected and analyzed to determine the expression of the antigen CD69 on the T lymphocytes using a standard FACS sorting protocol. The cell culture supernatants were collected after two days in culture and tested to determine the concentration of human interferon gamma using an ELISA assay. A portion of the CLL cells containing an accessory molecule ligand gene (murine CD40 ligand) and a portion of the cells containing the adenovirus expressing the lac-Z were cultured in the presence of human interleukin 4 IL-4 (5 ng/mL). The production of interferon gamma by allogeneic T lymphocytes in the presence of this amount of human interleukin 4 was also analyzed. The results from these analyses are shown in FIG. 6.

As can be seen, the human CLL cells containing the accessory molecule ligand gene (murine CD40) produced substantially higher concentrations of interferon gamma in the cell culture supernatant when compared to CLL cells which contained the lac-Z gene. The increased production of interferon gamma (IFN γ) by T lymphocytes exposed to CLL cells containing the accessory molecule ligand gene indicates that these CLL cells containing the accessory molecule ligand genes were effective in producing an enhanced immune response.

iv. Stimulation of Allogeneic T Cells Pre-Exposed to Non-Modified CLL B Cells Containing an Accessory Molecule Ligand Gene

Prior studies indicated that antigen presentation to T cells, in the absence of the signals derived from costimulatory molecules such as CD28, can lead to specific T cell clonal anergy. For this reason, allogeneic T cells that had previously been cultured, with non-modified CLL B cells lacking expression of CD80 and other immune accessory molecules, were tested for their ability to respond to CLL cells containing the CD40 ligand gene. Allogeneic effector cells did not incorporate more ^{3} H-thymidine in response to non-modified CLL cells (FIG. 12C, CLL), or control CLL cells infected with Ad-lacZ (FIG. 12C, lacZ-CLL), than when they were cultured alone (FIG. 12C, None). In contrast, even after prior co-culture with non-modified CLL B cells, allogeneic effector cells could still be induced to proliferate (FIG. 12C, CD154-CLL) or to produce IFN γ (FIG. 12D, CD154 CLL) in response to cells expressing an accessory molecule ligand. Although modest amounts of IFN γ were detected in the supernatants of such secondary cultures when Ad-lacZ-infected leukemia cells were used as stimulator cells (FIG. 12D, lacZ-CLL), this level was significantly lower than that noted for secondary cultures with Ad-CD40-ligand-infected CLL cells (FIG. 12D, CD154-CLL) ($P < 0.05$, Bonferroni t-test). Similarly, the supernatants of the leukemia cells alone (data not shown), and the effector cells alone (FIG. 12D, None), of the MLTR cultures stimulated with uninfected CLL cells (FIG. 12D, CLL), contained negligible amounts of IFN γ (< 2 ng/ml). These results indicate that allogeneic effector cells cultured

with nonmodified CLL B cells are not precluded from responding to CLL B cells infected with a gene therapy vector containing the accessory molecule ligand gene.

v. Autologous T Cell Responses to CLL Cells into which a Gene Therapy Vector Encoding a Murine Accessory Molecule Ligand Gene has been Introduced

T cells isolated from the blood of CLL patients were examined for their ability to respond in vitro to autologous CLL B cells containing a gene therapy vector which encodes the murine accessory molecule, CD40 ligand. T cells were isolated to >95% purity, and then co-cultured with mitomycin-C-treated autologous leukemia cells in serum-free AIM-V medium supplemented with exogenous interleukin-2 at 25 U/ml. Modest ³H-thymidine incorporation ($\leq 10,000$ cpm) was detected in cultures without added stimulator cells, secondary in part to the exogenous IL-2 (FIG. 13A, and data not shown). The level of T cell proliferation, however, did not increase in response to uninfected CLL cells (FIG. 13A, CLL) or Ad-lacZ-infected CLL cells (FIG. 13A, lacZ-CLL). In contrast, CLL cells infected with a gene therapy vector containing the accessory molecule ligand (FIG. 13A, CD154-CLL) induced autologous T cells to incorporate significantly more ³H-thymidine ($17,368 \pm 1,093$ cpm, $n=3$) than any of the control cultures ($P<0.05$, Bonferroni t-test). Furthermore, the MLTR stimulated with CLL cells infected with a vector encoding an accessory molecule ligand (CD40L) also generated significantly more IFN γ (165 ± 3 ng/ml, $n=3$) than any of the other cultures (FIG. 13B) ($P<0.05$, Bonferroni t-test).

The T cells were harvested after 5 days from the autologous MLTR and assessed for CTL activity against autologous CLL B cells. T cells co-cultured with autologous CD40-ligand-CLL cells developed CTL activity for non-modified CLL B cells, effecting 40.1% lysis ($\pm 2.3\%$) at an E:T ratio of 2:1 (FIG. 13C, CD154). However, such T cells did not develop detectable CTL activity for the same target cells in the control reactions, when co-cultured with uninfected or Ad-lacZ-infected CLL cells (FIG. 13C).

vi. Specificity of CTL Stimulated by Autologous CD40-Ligand-CLL B Cells for Allogeneic CLL B Cells

Effector cells stimulated with autologous CD40-ligand-CLL were evaluated for their ability to secrete IFN γ or manifest CTL activity against allogeneic CLL B cells (FIG. 14). After 5 days of autologous MLTR with CD154-CLL or lacZ-CLL, T cells were isolated by Ficoll density gradient centrifugation, washed extensively, and then cultured in media for 24 h. Washed T cells were mixed with autologous ("Auto CLL", solid bar) or allogeneic ("Allo-1 CLL" or "Allo-2 CLL", shaded or hatched bars) target CLL B cells. T cells stimulated in the autologous MLTR with CD40-ligand-CLL cells, but not with lacZ-CLL cells, produced significantly more IFN γ in response to secondary culture with non-modified autologous CLL B cells than with allogeneic CLL B cells (FIG. 14A) ($P<0.05$, Bonferroni t-test). Furthermore, T cells stimulated with CD40-ligand-CLL cells, but not with lacZ-CLL cells, were cytotoxic for autologous CLL cells, but not allogeneic CLL cells (FIG. 14B). Similar results were obtained with the autologous MLTR-activated T cells of the allogeneic donor, again demonstrating specific cytotoxicity for autologous CLL B cells (data not shown). Finally, W6/32, a mAb to class I major histocompatibility complex (MHC I) antigens could significantly inhibit the cytotoxicity of T cells stimulated with CD40-ligand-CLL cells for autologous CLL B cells (FIG. 14C, α HLA-class I) ($P<0.05$, Bonferroni t-test). Such inhibition was not observed with mAb specific for MHC class II antigen (FIG. 14C, α HLA-DP), mAb specific for the Fas-ligand (FIG. 14C, α FasL), or an isotype control mAb of irrelevant specificity (FIG. 14C, MOPC-21). Collectively,

these studies indicate that Ad-CD40-ligand-infected CLL cells can induce an autologous anti-leukemia cellular immune response in vitro, leading to the generation of MHC-class I-restricted CTL specific for autologous non-modified leukemia B cells.

e. Transactivation of Non-Infected Bystander Leukemia B Cells by Ad-CD40L CLL Cells

To address whether the changes in tumor marker expression (described in section 1di.) resulted from intracellular versus intercellular stimulation, the effect of culture density on the induced expression of CD54 and CD80 following infection with adenovirus gene therapy vector encoding the accessory molecule ligand (CD40L, or CD154) was examined. After infection, CLL cells were cultured at standard high density (e.g. 1×10^6 cells/ml) or low density (e.g. 2×10^5 cells/ml) for 3 days at 37° C. Cells plated at high density contained homotypic aggregates, whereas cells plated at low density remained evenly dispersed and without substantial cell-cell contact (data not shown). Despite expressing similar levels of heterologous CD154, CD154-CLL B cells cultured at high density were induced to express higher levels of CD54 and CD80 than CD154-CLL cells cultured at low density (FIG. 15A). The stimulation achieved at high density could be inhibited by culturing the cells with a hamster anti-mouse CD154 mAb capable of blocking CD40 $\leftrightarrow$ CD154 interactions (FIG. 15B, α CD154 Ab). Collectively, these studies indicate that CD154-CLL cells can activate each other in trans and that surface expression of CD154 is necessary for optimal leukemia cell stimulation.

In addition, Ad-CD154-infected, uninfected, Ad-lacZ-infected, or G28-5-stimulated CLL cells were labeled with a green-fluorescence dye to examine whether CD154-CLL could stimulate non-infected bystander leukemia cells. Dye-labeled cells were used as stimulator cells for equal numbers of non-labeled syngeneic CLL B cells. After 2 days' culture, stimulator cells cultured by themselves retained the green-fluorescence dye, allowing such cells to be distinguished from non-labeled CLL cells by flow cytometry. Bystander (green-fluorescence-negative) CD19⁺ CLL B cells were induced to express CD54 (FIG. 15C, right histogram) or CD86 (FIG. 15D, right histogram) when co-cultured with Ad-CD154-infected leukemia B cells, but not with mock infected CLL cells (FIGS. 15C and 15D, left histograms), G28-5-stimulated CLL cells, or Ad-lacZ-infected CLL cells (data not shown). As expected, these bystander (green-fluorescence-negative) CLL cells also were negative for heterologous CD154.

f. Treatment of Leukemia with Gene Therapy Vectors Encoding an Accessory Molecule Ligand

FIG. 24 shows an outline for a clinical trial for testing treatment of B cell CLL with adenovirus gene therapy vectors encoding modified CD40 ligand. Leukemia cells harvested by pheresis are infected with replication-defective vectors that encode the modified CD40 ligand. Following expression of this protein, the cells will be administered back to the patient for the purpose of stimulating a host anti-leukemia-cell immune response. This strategy is far superior to one that uses gene therapy to affect expression of only one immune stimulatory molecule on the leukemia cell surface. Indeed, this strategy results in the leukemia cells expressing an array of immune-stimulatory accessory molecules and cytokines, as well as a molecule that can affect the same changes in leukemia cells of the patient that were never harvested.

2. Expression of Chimeric Accessory Molecule Ligand Genes

The chimeric accessory molecule ligand genes described below are prepared using standard techniques as described herein.

a. Preparation of Chimeric Accessory Molecule Ligand Genes Utilizing Domains from Two Different Accessory Molecule Genes

The human CD40 ligand gene was isolated from RNA prepared from T cells which had been activated by an anti-CD3 monoclonal antibody using 5' and 3' primers together with well known PCR methods. Chimeric accessory molecule genes of human CD40 ligand and murine CD40 ligand are constructed from the newly cloned human CD40 ligand gene and mouse CD40 ligand gene described herein as SEQ ID NO: 2. The transmembrane and cytoplasmic domains of human CD40 ligand genes are exchanged with those of the murine CD40 ligand gene and designated H(Ex)-M(Tm-Cy) CD40 ligand. These chimeric accessory molecule ligand genes are produced using the gene conversion technique described as SOEN which has been previously described by Horton, *Mol. Biotechnol.*, 3:93 (1995). A diagram depicting the chimeric accessory molecule ligand genes which are produced is shown in FIG. 4. The nucleotide sequences of each of these respective chimeric accessory molecule ligand genes is designated SEQ ID NOS: 3-7 as indicated in the Table below.

TABLE III

Chimeric Accessory Molecule Ligand Gene	SEQ ID NO:
HuIC/HuTM/MuEX CD40-Ligand	SEQ ID NO: 3
HuIC/MuTM/HuEX CD40-Ligand	SEQ ID NO: 4
HuIC/MuTM/MuEX CD40-Ligand	SEQ ID NO: 5
MuIC/HuTM/HuEX CD40-Ligand	SEQ ID NO: 6
MuIC/MuTM/HuEX CD40-Ligand	SEQ ID NO: 7

Adenovirus vectors encoding each of the chimeric accessory molecules shown in FIG. 2 are constructed using the methods described in Example 1. Each of these constructs are then transfected into either HeLa cells or CLL cells according to the methods of Example 1.

b. Expression of Chimeric Accessory Molecule Ligands on CLL and HeLa Cells

The expression of each of the chimeric accessory molecule ligand genes constructed above is analyzed by using FACS analysis as specified in Example 1. The appropriate monoclonal antibody immunospecific for the external domain of either human or mouse CD40 ligand is selected and used to determine the level of expression of the chimeric accessory molecules on the surface of these cells. After appropriate analysis and preparation of appropriate histograms, the expression of chimeric accessory molecules containing at least a portion of the murine CD40 ligand gene is confirmed.

c. Function of Chimeric Accessory Molecule Ligands

CLL cells are infected with various MOI of the mCD40L adenovirus and then cultured in 48 or 24 well tissue culture plates for various times after infection (48, 72, and 96 hours). The CD19⁺ B cells are then analyzed by multiparameter FACS analysis for induction of CD80 and CD54 expression using fluorescein isothiocyanate-conjugated mAb specific for each respective surface antigen as described in Example 1. Increased amounts of CD54 and CD80 are found on cells which have the chimeric accessory molecules containing the domain or domains derived from the mouse CD40 ligand gene.

Further analysis of the cells containing the chimeric accessory molecule genes is carried out according to Example 1(d). The cells containing the chimeric accessory molecule genes which contain the domains derived from the murine CD40 ligand gene are able to stimulate the production of gamma interferon and T cell proliferation.

d. Expression of Chimeric Accessory Molecule Genes Which Contain Proximal Extracellular Domains from Two Different Accessory Molecules from the Same Species

A chimeric accessory molecule ligand gene is prepared which contains the proximal extracellular domain from the human CD70 gene (Domain III) with the remainder of the domains derived from the human CD40 ligand gene. This gene is prepared using standard biologic techniques as previously described herein. This chimeric accessory molecule ligand gene has the DNA sequence shown as SEQ ID NO: 19. A different chimeric accessory molecule ligand gene is prepared which contains the proximal extracellular domain from the murine CD40 ligand gene with the remainder of the domains derived from the human CD40 ligand gene. This gene is prepared using standard techniques as previously described herein. This chimeric accessory molecule ligand gene has the DNA sequence shown as SEQ ID NO: 20.

The chimeric accessory molecule genes shown as SEQ ID NOS: 19 and 20 are inserted into the appropriate vectors as described in Example 1 and introduced into human neoplastic cells. The expression of that chimeric accessory molecule gene in the cells is determined as was described in Example 1.

The chimeric accessory molecule encoded by each of these chimeric accessory molecule genes is found on the surface of the human neoplastic cells using the FACS analysis described in Example 1. Increased amounts of CD54 and CD80 are found on the cells containing the chimeric accessory molecule genes using the techniques described in Example 1. The cells containing the chimeric accessory molecule gene are able to stimulate the production of gamma interferon and T cell proliferation as described and assayed according to Example 1.

3. Augmentation of Vaccination Using Vectors Encoding Accessory Molecules

The following procedures were used to demonstrate the augmentation of a vaccination protocol using a gene therapy vector encoding an accessory molecule.

a. Augmentation of the Antibody Response in Mice Co-Injected with an Accessory Molecule Gene Therapy Vector and placZ

Three different gene therapy constructs were prepared using standard techniques including those techniques described herein. The first was a control gene therapy vector, pcDNA3, which did not contain any gene. The second, placZ, contained the Lac-Z gene which encoded β -galactosidase (β -gal). The third, p-mCD40L, contained the murine CD40 ligand gene described in Example 1.

Prior to any immunizations, serum was isolated from 6-8 week old BALB/c-mice to determine the amount of any initial antibodies to β -galactosidase. Each animal was injected i.m. with 100 micrograms of plasmid DNA per injection. Four separate injections were given at one week intervals.

Prior to the third injection, the animals were bled to monitor the early antibody response to β -gal. One week after the final injection of plasmid DNA, the animals were bled to monitor the late antibody response to beta-galactosidase. To test the sensitivity of the assay, known amounts of anti- β -gal antibodies isolated from an anti- β -gal antiserum were tested in parallel.

Serum dilutions of 1:40, 1:200, or 1:1000 were tested in an ELISA for anti- β -gal antibodies. For this, polystyrene microtiter ELISA plates were coated with β -gal at 10 microgram/ml in phosphate buffered saline. The plates were washed thrice with blocking buffer containing 1% bovine serum albumin (BSA), 0.2% Tween 20 in borate buffered saline (BBS) (0.1M

borate, 0.2M NaCl, pH 8.2). 50 microliters of diluted serum were added to separate wells. After at least 1 hour at room temperature, the plates were washed thrice with blocking buffer and then allowed to react with alkaline phosphatase-conjugated goat anti-mouse IgG antibody. One hour later, the plates again were washed four times with blocking buffer and incubated with 25 ml of TMB peroxidase substrate (Kirkegaard & Perry, Gaithersburg, Md.). The absorbance at 405 nm of each well was measured using a microplate reader (Molecular devices, Menlo Park, Calif.). The higher the O.D. reading, the greater the amount of specific antibody in the sample.

The data for each of two experiments are provided in Tables IV and V which follow on separate sheets. The results are summarized in Tables VI and VII collating the data from the two experiments is provided as well. On the summary page *n* stands for the number of animals in each of the four groups. S.D. stands for standard deviation and Avg. is the average O.D. reading for all the animals in a particular group.

The results of Group 4 demonstrate that the use of a gene therapy vector encoding an accessory molecule ligand (CD40L) enhances the immunization against β -gal encoded by a genetic or gene therapy vector. The average O.D. reading of the 1:40 dilution of the sera from animals of this group is significantly higher than that of groups 1, 2, and 3 ($P < 0.05$, Bonferroni *t* tests, see Table VII).

Data from an additional experiment further reinforce the finding that the gene therapy vector encoding an accessory molecule ligand enhances immunization against β -gal (FIG. 16). Here, pCD40L and placZ were co-injected into skeletal muscle, to test for enhancement of the immune response to placZ, a pcDNA3-based vector encoding *E. coli* β -galactosidase. The relative anti- β -gal Ab activities were determined via ELISA. As expected, mice injected with either the non-modified pcDNA3 vector or pCD40L alone did not produce detectable antibodies to β -gal (FIG. 16A). Mice were injected with either 100 μ g pcDNA3 (checkered bar), 50 μ g pcDNA3+50 μ g pCD40L (lined bar), 50 μ g pcDNA3+50 μ g placZ (striped bar), or 50 μ g pCD40L+50 μ g placZ (solid bar). On the other hand, mice that received placZ and pcDNA3 developed detectable anti- β -gal antibodies one week after the fourth and final injection, at d28. Mice that received placZ and pCD40L developed higher titers of anti- β -gal antibodies than mice injected with placZ and pcDNA3. FIG. 16B, ELISA analyses of serial dilutions of sera collected at d28, shows that mice co-injected with placZ and pCD40L had an eight-fold higher mean titer of anti- β -gal antibodies at d28 than mice treated with placZ+pcDNA3.

i. Immunoglobulin Subclass Production Stimulated by Accessory Molecule Vector Co-Injection

Despite enhancing the titer of the anti- β -gal antibody response, the subclass of anti- β -gal IgG induced by injection of placZ was not altered by the co-injection of pCD40L. IgG_{2a} anti- β -gal antibodies predominated over IgG₁ subclass antibodies in the sera of mice injected with either placZ and pcDNA3 or placZ and pCD40L (FIG. 17). Also depicted are the ELISA O.D. measurements of anti- β -gal IgG₁ and anti- β -gal IgG_{2a} present in the pre-immune sera (striped bar) or post-immune sera (solid bar), collected at d28) of each group of mice, injected as indicated on the abscissa. In contrast, BALB/c mice injected with β -gal protein developed predominantly IgG₁ anti- β -gal antibodies, and no detectable IgG_{2a} anti- β -gal antibodies.

ii. Augmentation of Vaccination by Accessory-Molecule Vector Requires Co-Injection with placZ at the Same Site

The adjuvant effect of the pCD40L plasmid on the anti- β -gal antibody response was noted only when it was injected

into the same site as placZ (FIG. 18). Groups of BALB/c mice (*n*=4) received intramuscular injections of placZ and pCD40L together at the same site, or as simultaneous separate injections at distal sites (right and left hind leg quadriceps). A control group received intramuscular injections of placZ and pcDNA3 at the same site. Animals were bled at d28 and the sera tested for anti- β -gal Ab at different dilutions, as indicated on the abscissa. The graph illustrates a representative experiment depicting the mean O.D. at 405 nm of replicate wells of each of the serum samples for each group, at a 1:40, 1:200, or 1:1000 dilution. Animals injected simultaneously with placZ and pCD40L, but at different sites, did not develop detectable anti- β -gal antibodies until d28. Moreover, the anti- β -gal antibody titers of the sera from such animals at d28 were similar to that of mice that received placZ and pcDNA3, and significantly less than that of animals that received placZ and pCD40L together at the same site.

iii. Augmentation of Vaccination when Accessory Molecule Vector and placZ are Co-Injected into Dermis

The pCD40L plasmid also enhanced the anti- β -gal antibody response to placZ when injected into the dermis. In the experiment shown in FIG. 19, mice received intradermal injections, near the base of the tail, with either 50 μ g pcDNA3 (checkered bar), 25 μ g pcDNA3+25 μ g pCD40L (lined bar), 25 μ g pcDNA3+25 μ g placZ (striped bar), or 25 μ g pCD40L+25 μ g placZ (solid bar). Injections, bleeds and ELISA analyses were performed as in FIG. 16A. The checkered bar and lined bar groups each consisted of 8 mice while the striped bar and solid bar groups each consisted of 12 mice. The height of each bar represents the mean O.D. of sera at a 1:40 dilution of each group $\pm$ S.E. A statistical analysis of the data indicated that the striped bar and solid bar groups are independent ($P < 0.05$). As observed with intramuscular injection, mice co-injected with placZ and pCD40L developed detectable serum anti- β -gal antibodies one week following the second injection (d14), and two weeks earlier than mice injected with placZ and pcDNA3. Moreover, these animals also had an eight-fold higher mean titer of anti- β -gal antibodies than mice of the placZ-injected group at d28. Mice injected with either the non-modified pcDNA3 vector or pCD40L alone did not produce detectable antibodies to β -gal.

b. Augmentation of the CTL Response in Mice Co-Injected with an Accessory Molecule Gene Therapy Vector and placZ

The ability of pCD40L to enhance induction, by placZ, of CTL specific for syngeneic β -gal-expressing target cells was tested. BALB/c mice co-injected with pCD40L and placZ into skeletal muscle (FIG. 20A) or dermis (FIG. 20B) generated greater numbers of CTL specific for P13.2, a placZ transfected P815 cell line, than mice co-injected with placZ and pcDNA3. At a 5:1 effector:target ratio, the splenocyte effector cells from mice that received intramuscular injections of placZ and pCD40L achieved greater than 20% specific lysis of P13.2. In contrast, when splenocytes of mice that received the control injection with placZ and pcDNA3 were used, a 9-fold greater ratio of effector to target cells was required to achieve this level of specific lysis. Similarly, the splenocyte effector cells from mice that received intradermal injections of placZ and pCD40L killed more than 50% of the P13.2 cells at effector:target ratios of 4:1. To achieve comparable levels of specific lysis required eight-fold higher effector:target ratios using splenocytes from mice that received intradermal injections of placZ and pcDNA3. Nevertheless, the splenocytes of mice co-injected with pCD40L and placZ did not have greater non-specific CTL activity for P815 cells than that of mice that received placZ along with pcDNA3 (FIG. 20). As expected, the splenocytes from mice that

received injections of pcDNA3 alone, or pcDNA3 and pCD40L, did not mediate specific lysis of P13.2 or P815 cells.

TABLE IV

Experiment #1 Injections of plasmid DNA i.m.: Apr. 3, 1996; Apr. 10, 1996; Apr. 17, 1996; Apr. 24, 1996						
ELISA for anti-beta galactosidase		Dilution of Pre-Bleed (4/3)			Dilution of Bleed (4/17)	
antibodies:	Animal	1/140	1/200	1/1000	1/140	1/200
Group	1	0.09	0.11	0.09	0.06	0.06
pcDNA3	2	0.11	0.09	0.09	0.07	0.07
(p-control, 100 mcg)	3	0.12	0.11	0.10	0.09	0.09
(Control vector)	4	0.11	0.10	0.10	0.08	0.11
	Avg.	0.11	0.11	0.11	0.11	0.11
	S.D.	0.01	0.01	0.01	0.01	0.02
p-lacZ (50 mcg) +	5	0.13	0.10	0.10	0.07	0.11
p-Control (50 mcg)	6	0.10	0.11	0.10	0.07	0.06
	7	0.19	0.10	0.18	0.07	0.07
	8	0.10	0.09	0.10	0.08	0.07
	Avg.	0.13	0.10	0.12	0.07	0.08
	S.D.	0.04	0.01	0.04	0.01	0.02
p-lacZ (50 mcg) +	27	0.06	0.06	0.06	0.13	0.11
pReCMV-mCD40L	18	0.06	0.06	0.06	0.27	0.13
(p-mCD40L, 50 mcg)	19	0.06	0.06	0.06	0.23	0.19
	20	0.06	0.06	0.06	0.23	0.19
	Avg.	0.06	0.06	0.06	0.74	0.47
	S.D.	0.00	0.00	0.00	1.06	0.66

TABLE V

Experiment #2 Injections of plasmid DNA i.m.: Jun. 5, 1996; Jun. 12, 1996; Jun. 19, 1996; Jun. 26, 1996						
Dilutions of sera for anti-beta galactosidase		Dilution of pre-Bleed (6/5)			Dilution of Bleed (7/1)	
antibodies:	Animal	1/140	1/200	1/1000	1/140	1/200
Group	9	0.02	0.02	0.06	0.04	0.01
p-Control (50 mcg) +	10	0.06	0.02	0.10	0.02	0.02
p-mCD40L (50 mcg)	11	0.02	0.02	0.07	0.03	0.01
	12	0.06	0.03	0.05	0.18	0.04
	Avg.	0.04	0.04	0.04	0.04	0.04
	S.D.	0.02	0.01	0.02	0.07	0.01
p-lacZ (50 mcg) +	5	0.02	0.03	0.02	0.06	0.04
p-Control (50 mcg)	6	0.03	0.02	0.03	0.14	0.03
	7	0.56	0.13	0.06	0.29	0.06
	8	0.01	0.02	0.05	0.06	0.02
	Avg.	0.15	0.05	0.04	0.13	0.04
	S.D.	0.27	0.05	0.02	0.11	0.02
p-lacZ (50 mcg) +	13	0.23	0.06	0.05	0.28	0.07
p-mCD40L (50 mcg)	14	0.02	0.02	0.03	0.04	0.02
	15	0.02	0.02	0.02	0.89	0.21
	16	0.05	0.04	0.02	0.11	0.04
	Avg.	0.08	0.04	0.03	0.33	0.08
	S.D.	0.10	0.02	0.02	0.39	0.09

TABLE VI

Summary						
		Pre-Immune @ beta-gal			Early @ beta-gal	
		1/140	1/200	1/1000	1/140	1/200
1) p-Control (n = 4)	Avg.	0.11	0.11	0.11	0.11	0.11
	S.D.	0.01	0.01	0.01	0.01	0.02
2) p-mCD40L +	Avg.	0.04	0.04	0.04	0.04	0.04
p-Control (n = 4)	S.D.	0.02	0.01	0.02	0.07	0.01
3) p-lacZ +	Avg.	0.11	0.04	0.04	0.11	0.03
p-Control (n = 8)	S.D.	0.22	0.04	0.01	0.09	0.02
4) p-lacZ +	Avg.	0.11	0.04	0.03	0.25	0.06

TABLE VI-continued

Summary						
		Pre-Immune @ beta-gal			Early @ beta-gal	
		1/140	1/200	1/1000	1/140	1/200
p-mCD40L (n = 8)	S.D.	0.10	0.02	0.01	0.32	0.07
Anti-beta-	67 ng	22 ng	7.4 ng	2.5 ng	.82 ng	.27 ng
galactosidase						
standard:						
O.D.	3.01	2.98	2.05	1.10	0.52	0.26
	3.14	3.14	2.25	1.20	0.56	0.26

TABLE VII

BONFERRONI t-TESTS				
Comparison	Difference of means	t	P < .05	
4 vs 2:	2.06 – 0.04 = 2.02	3.782	Yes	
4 vs 1:	2.06 – 0.11 = 1.95	3.651	Yes	
4 vs 3:	2.06 – 0.61 = 1.45	3.325	Yes	
3 vs 2:	0.61 – 0.04 = 0.57	1.067	No	
3 vs 1:	0.61 – 0.11 = 0.50		Do not test	
1 vs 2:	0.11 – 0.04 = 0.07		Do not test	
Degrees of freedom: 20				
ONE WAY ANALYSIS OF VARIANCE				
Group	N	Mean	Std Dev	SEM
1	4	0.11	0.01	0.00
2	4	0.04	0.04	0.02
3	8	0.61	1.11	0.39
4	8	2.06	0.97	0.34
5	4	1.51	0.77	0.38
6	4	1.14	0.53	0.26
7	4	0.83	0.43	0.22
ONE WAY ANALYSIS OF VARIANCE				
Source of Variation	SS	DF	Variance Est (MS)	
Between Groups	18.29	6	3.05	
Within Groups	18.39	29	0.63	
Total	36.69	35		
$F = \frac{s2_bet}{s2_wit} = \frac{MSbet}{Mswit} = \frac{3.05}{0.63} = 4.81$			P = 0.002	

4. Treatment of Neoplasia Using a Gene Therapy-Vector Containing an Accessory Molecule Gene or Chimeric Accessory Molecule Gene

a. Treatment of Neoplasia in Mice

The treatment of a neoplasia in a mouse model system has been demonstrated using the genes encoding accessory molecule ligands of the present invention. Gene therapy vectors containing an accessory molecule ligand gene (murine CD40 ligand) were prepared as has been previously described in the above examples. These gene therapy vectors were used to introduce that accessory molecule ligand gene into neoplastic cells, Linel cells, from a tumor which originated in BALB/c mice. The accessory molecules were introduced into the neoplastic cells according to the above examples. The expression of the accessory molecule ligand on the surface of these neoplastic cells was confirmed using flow cytometry as has been described in the above examples.

The effectiveness of the accessory molecule ligand genes for treating neoplasia was shown as follows. Female BALB/c

mice (6-8 weeks old) were injected i.p. with 1.0×10^5 irradiated Linel neoplastic cells. The neoplastic Linel cells are derived from a spontaneous lung adenocarcinoma in a BALB/c mouse. This neoplastic cell has been described by Blieden et al., *Int. J. Cancer Supp.*, 6:82 (1991). Other female BALB/c mice were injected i.p. with 1.0×10^5 irradiated Linel tumor cells that had previously been transduced with the gene therapy vector encoding the accessory molecule ligand gene (murine CD40) as described above.

Each group of mice was allowed to generate an immune response for 10 days. After 10 days each mouse was challenged with 1.0×10^4 live, non-irradiated Linel neoplastic cells. These mice were then monitored for the formation of tumors and then sacrificed when the tumors grew to 2.0 cm because of morbidity. The results of this monitoring are shown in FIG. 7. As can be seen by FIG. 7, the mice immunized with the neoplastic cell expressing the accessory molecule ligands of the present invention on the cell surface remained free of tumor throughout the experiment. Mice immunized with the neoplastic cells not having the accessory molecule ligand genes of the present invention succumbed to tumor 50 days after challenge with the neoplastic cells.

FIG. 21 demonstrates downmodulation of human CD40L, but not murine CD40L, in lung tumor cell lines that express CD40. Human cell lines HeLa (CD40-negative cervical carcinoma, FIG. 21A), A427 (CD40-negative lung carcinoma, FIG. 21B), NCI 460 (weakly CD40-positive lung large cell carcinoma, FIG. 21C), and SK-Mes-1 (strongly CD40-positive lung squamous cell tumor, FIG. 21D) were infected with adenovirus encoding lac-Z (Ad-LacZ), murine CD40L (Ad-mCD40L), and human CD40L (Ad-hCD40L) at an MOI of 0 (Blank), 1, and 10. 48 hours after infection, murine CD40L and human CD40L surface expression was determined. The percentage of cells that express ligand are plotted on the Y-axis. Human and mouse CD40L are expressed at equal levels in CD40-negative cell lines. However, only murine CD40L expression is stable on cell lines that express CD40. In contrast to mCD40L, human CD40L is downmodulated on CD40-positive tumors.

The data graphed in FIG. 22A show that CD40 binding induces expression of tumor surface markers. Treating CD40-expressing lung cancer cell lines with α CD40 mAb resulted in enhanced expression of the tumor cell surface markers CD95 (Fas), CD54 (ICAM-1) and class I major histocompatibility antigens (MHC I). NCI 460, a weakly CD40-positive lung large cell carcinoma, was incubated with a CD40-specific monoclonal antibody (thick line), or MOPC21, an isotype control mAb (thin line), on CD32-expressing mouse fibroblasts for 48 hours. Following the 48 hr incubation, the lung tumor cells were analyzed for CD95, CD54, and MHC-I expression by FACS.

FIG. 22B again shows downmodulation of human CD40L by CD40-positive tumor cells. HeLa (CD40-negative), CLL (CD40-positive), and SK-MES-1 (CD40-positive) tumor cells were cocultured for 24 hours with CD3-activated normal donor T cells at a tumor cell:T cell ration of 2.5:1. Following coculture, CD2-expressing T cells were analyzed for CD40L surface expression by FACS. Thin lines represent T cells stained with FITC-labeled isotype control antibody (MOPC21) and thick lines represent activated T cells stained with FITC-labeled α CD40L antibody (α CD154 antibody). The CD40-positive tumor cell lines, SK-MES-1, and CLL, do not express CD40 ligand on their surfaces.

5. Expression of the Human and Mouse Accessory Molecule Ligand, Fas Ligand, in Human Blood Lymphocytes

a. Construction of a Genetic Construct and Gene Therapy Vector Containing the Human and Mouse Fas Ligand Gene

Either the human accessory molecule ligand gene (human Fas ligand) or the murine accessory molecule ligand gene (murine Fas ligand) was constructed utilizing the respective human and murine genes. An altered accessory cell molecule, in which a putative MMP-cleavage site was removed, was made and designated AFasL-pcDNA3. The nucleotide sequence of AFasL-pcDNA3 is listed as SEQ ID NO: 40. Human Fas ligand nucleotides 325 to 342, encoding six amino acids, are missing from AFasL. The design of AFasL was based on reasoning that Domain III contains sites most accessible to MMPs, and could thus be the target on the molecule for cleavage from the surface of the cell. Sequences of the human Fas ligand gene have been determined and are listed as SEQ ID NOS: 13 and 30 (Genbank accession U11821). Sequences of mouse Fas ligand genes have been determined and are listed as SEQ ID NOS: 14 (C57BL/6, Genbank accession U10984) and 31 (Balb/c, Genbank accession U58995). The sequence of the rat Fas ligand gene has been determined and is listed as SEQ ID NO: 25 (Genbank accession U03470). Chimeric constructs are made, as described in Example 2 for CD40 ligand chimeric constructs, in which Domain III of human Fas ligand is replaced with Domains of other proteins, particularly proteins of the TNF family. Chimeric constructs include, but are not limited to, human Fas ligand with Domain III replaced by Domain III of murine Fas ligand (chimeric sequence listed as SEQ ID NO: 37, sequence line-up shown in FIG. 37), or replaced by Domain III of human CD70 (chimeric sequence listed as SEQ ID NO: 38, sequence line-up shown in FIG. 38), or replaced with Domain I of human CD70 (chimeric sequence listed as SEQ ID NO: 39, sequence line-up shown in FIG. 39). Chimeric constructs in which multiple domains, for example, two copies of human CD70 Domain III, are inserted into human Fas ligand in place of Domain III, are also made using methods described in Example I. Chimeric constructs in which synthetic sequences are used to replace Domain III of human Fas ligand are also made.

i. Human Fas Ligand Cloning

The cDNA encoding human Fas-ligand was subcloned in the eukaryotic expression vector pcDNA3. Normal donor blood lymphocytes were activated for 4 hours with 1 ng/ml PMA plus 0.5 μ M ionomycin. Total RNA was isolated with the Qiagen Rneasy kit. cDNA was then synthesized from poly-A RNA with oligo-dT primers using the Gibco-BRL Superscript cDNA synthesis kit. The gene encoding human Fas-ligand was then PCR amplified with the Fas-ligand-specific primers (sense primer, SEQ ID NO: 32, antisense primer, SEQ ID NO: 33). The Fas-ligand PCR product was then subcloned into pcDNA3 using standard molecular biology techniques. RT-PCR products, subcloned into pcDNA3, are designated hFasL-pcDNA3.

ii. Murine Fas Ligand Cloning

The murine Fas-ligand genes from Balb/c and C57/BL6 strains of mice were also amplified following activation of mouse splenocytes with PMA plus ionomycin as described above, and amplified from poly-A synthesized cDNA as described above (sense primer, SEQ ID NO: 34, antisense primer, SEQ ID NO: 35). These genes were subcloned in the pTARGET expression vector (Promega, Madison, Wis.). RT-PCR products, subcloned into pcDNA3, are designated mFasL-pcDNA3.

iii. Adenovirus Vector Construction

For construction of adenovirus vectors encoding human Fas-ligand, murine Fas-ligand or Δ Fas-ligand, the cloned cDNA insert is subcloned into the plasmid pRc/RSV (Invitrogen, San Diego, Calif.) at the HindIII-XbaI site. A BglII-XhoI fragment with the RSV promoter-enhancer and the

bovine growth hormone poly-A signal sequence was sub-cloned into the BamHI-XhoI site of plasmid MCS(SK) pXCX2. The plasmid MCS(SK)pXCX2 is a modification of the plasmid pXCX2, in which the pBluescript polylinker sequence was cloned into the E1 region. The resulting plasmid then is co-transfected along with pJM17 into 293 cells using the calcium phosphate method. Isolated plaques of adenovirus vectors are picked and expanded by infecting 293 cells. High titer adenovirus preparations are obtained, as described above which uses a cesium chloride gradient for concentrating virus particles via a step gradient, with the densities of 1.45 g/cm³ and 1.20 g/cm³, in which samples are centrifuged for 2 hours in an SW41 rotor (Beckman, Brea, Calif.) at 25,000 rpm at 4° C. The virus band is desalted using a Sephadex G-25 DNA grade column (Pharmacia, Piscataway, N.J.), and the isolated virus is stored at -70° C. in phosphate-buffered saline with 10% glycerol. The titer of the virus is determined by infecting permissive 293 cells at various dilutions and counting the number of plaques. Titers typically range from 10¹⁰ to 10¹² plaque forming units/ml. The adenovirus constructs are designated Ad-hFasL, Ad-mFasL and Ad-AFasL.

b. Introduction of the Murine and Human Fas Ligand Genes into Human Cells

The constructs hFasL-pcDNA3, mFasL-pcDNA3 and ΔFasL-pcDNA3 are transfected into 293 via electroporation. The transfected cells are selected in medium containing G418. Fas-ligand transfectants are screened for expression of the transgene using anti-Fas-ligand antibody and flow cytometry. The methods used are similar to those described for transfection of CD40L into CLL cells.

For FasL-adenovirus infection, 10⁶ freshly thawed and washed CLL cells or HeLa cells are suspended in 0.5 to 1 mL of culture medium for culture at 37° C. in a 5% CO₂-in-air incubator. Adenovirus are added to the cells at varying multiplicity of infection (MOI), and the infected cells are cultured for 48 hours, unless otherwise stated, before being analyzed for transgene expression.

c. Expression of the Fas Ligand Genes in Human Cells

Mice with the lymphoproliferative or generalized lymphoproliferative disorder are unable to delete activated self-reactive cells outside of the thymus. This is related to the fact that, in these mice, interactions between the Fas receptor and an accessory molecule ligand, Fas ligand, are defective. These animals develop numerous disorders, including lymphadenopathy, splenomegaly, nephritis, and systemic autoimmune pathology which resembles that seen in patients with systemic lupus erythematosus or rheumatoid arthritis (RA). It is conceivable that the normal interactions between the Fas receptor and the accessory molecule ligand that are responsible for clearance of activated lymphocytes from joints may be impaired in RA patients.

RA synovial lymphocytes express the Fas receptor at a higher proportion than that of matched RA blood lymphocytes to matched normal donor blood lymphocytes. On the other hand, RA synovial lymphocytes express little or no accessory molecule ligand. Since the RA synovial lymphocytes are sensitive to Fas-induced apoptosis, it is feasible that local expression of Fas ligand in the RA joint could serve to eliminate the synovial mononuclear cells that potentially mediate RA autoimmune pathology.

FIG. 23 shows that Fas-ligand expression in lymphocytes is inhibited by exposure to RA synovial fluid. Normal donor blood T cells were activated for 5 hours with 1 ng/ml PMA plus 0.5 μM ionomycin. Cells were incubated in the presence of rheumatoid arthritis blood plasma (circles), RA synovial fluid (diamonds), or neither (squares). In addition, cells were

incubated with increasing concentrations of the MMP inhibitor BB94. Following activation, cells were analyzed for Fas-ligand surface expression by FACS. The percentage of cells expressing Fas ligand are plotted in FIG. 23. This experiment demonstrates that there is a factor(s) present in RA synovial fluid and serum that prevents surface expression of Fas-ligand.

d. Function of Human, Murine and Chimeric Accessory Molecule Ligand, Fas Ligand

To determine the capacity of the AFasL constructs, the above-mentioned transfected cells are mixed with the Fas-ligand sensitive human T cell line, JURKAT. Following 4 hours coculture, the nonadherent JURKAT cells are collected and evaluated for apoptosis. The fluorescent compound 3,3'-dihexyloxacarbocyanine iodide (DiOC₆) is used to evaluate for apoptosis using a modification of a previously described protocol. For this, the cells are washed once at room temperature in phosphate buffered saline (PBS, pH 7.2). Cells are placed into separate wells of a 96 well U-bottom plastic microtiter plate at 10⁵-5×10⁵ cells/well in 50 ml total volume. If indicated, saturating amounts of PE-conjugated antibodies are added followed by addition of DiOC₆ and propidium iodide (PI). DiOC₆ and PI are used at 40 nM and 10 ng/ml final concentrations, respectively. The cells are then incubated 15 minutes in a 37° C., 5% CO₂ tissue culture incubator. The stained cells are then washed twice in ice cold PBS and ultimately suspended in 200 ml SM and analyzed by FACS. Dead cells and debris with characteristic forward and light scatter profiles and PI staining are excluded from analysis.

The ability of cells expressing ΔFasL-pcDNA3 to direct Fas-mediated apoptosis of cells expressing CD95 is compared with that of cells expressing FasL-pcDNA3. Relative stability of the protein products encoded by ΔFasL-pcDNA3 or FasL-pcDNA3 pre- and post-culture with RA synovial fluid, and with or without the metalloproteinase inhibitors, are assessed via flow cytometry of cells expressing either ligand.

6. Treatment of Arthritis with Gene Therapy Vectors Encoding an Accessory Molecule Ligand, Fas Ligand

The heterologous Fas-ligand constructs, made as described above, that show the highest stability of expression in combination with the greatest ability to mediate Fas-induced apoptosis, are used in gene therapy for RA. Potential therapeutic constructs are tested in well-characterized mouse models of arthritis to assess efficacy and function in vivo.

a. Gene Therapy Treatment of Arthritis in Mice

i. Mouse Models for Arthritis

One mouse arthritis model is collagen-induced arthritis. It is known that injecting DBA/1 mice with type II collagen in complete Freund's adjuvant (CFA) induces an arthritis with synovitis and erosions that histologically resemble RA. For our studies, male DBA/1 mice are immunized with bovine type II collagen in complete Freund's adjuvant on day 0 and boosted intraperitoneally (i.p.) on day 21. On day 28, animals are given an additional i.p. injection with lipopolysaccharide (LPS) and/or the same type collagen, or an injection of acetic acid alone. Swelling and/or redness of a fore or hind paw in animals immunized with collagen typically is detected the third or fourth week following the second injection. The vertebrae are only rarely affected, and then only weeks after the initial peripheral joint swelling. Affected joints display initial histologic changes of synovial edema, followed by synovial hyperplasia.

Another animal model, recently described by Kouskoff, V. et al., in *Cell* 87:811-822 (1997) was generated fortuitously, by crossing a T cell receptor (TCR) transgenic mouse line with the non-obese-diabetic (NOD) strain to produce the KRN×NOD mouse model of RA. The offspring of such a mating universally develop a joint disease that is highly similar to that of patients with RA. Moreover, the disease in these animals has an early and reproducible time of onset and a highly reproducible course. The arthritis apparently is induced by chance recognition of an NOD-derived major histocompatibility complex (MHC) class II molecule by the transgenic TCR, leading to breakdown in the general mechanisms of self-tolerance and systemic self-reactivity.

ii. Relief of Arthritis Symptoms in Mice Treated with a Gene Therapy Vector Encoding an Accessory Molecule Ligand

We have adapted and modified a protocol originally described by Sawchuk and colleagues for micro-injecting adenovirus vectors into mouse joints. Using this procedure we can reproducibly inject a 5 μ l volume into the articular space of the mouse knee. In this procedure, the mice are anesthetized with metofane. A small incision of approximately 2-3 mm is made with a #11 scalpel blade in the skin over the lateral aspect of the knee to visualize the patello-tibial ligament. We can inject up to 5 μ l of fluid using a micro-100 μ l-Hamilton syringe and a 30-gauge needle. After the injection, the knee incision is closed with Nexabond (Veterinary Products Laboratory). Our adenovirus titers typically exceed 10^{10} plaque forming units (pfu) per ml, making it possible to deliver at least 5×10^8 pfu of virus in 5 ml into the knee joints, as outlined above. Control animals are injected with control Ad-lacZ vector, a replication-defective adenovirus vector lacking a transgene, or with the buffer used to suspend the virus (10 mM Tris, 1 mM $MgCl_2$, 10% glycerol).

In another method, splenocytes will be harvested from mice that are syngeneic to the host animal intended for adoptive transfer of transduced cells. Cell proliferation will be induced with exogenous IL-12 (100 units/ml) for 48 h. Cells are counted and then re-plated at densities of 5×10^5 or 1×10^6 cells per ml in a 12-well dish with 1 ml complete culture medium per well. Virus and ConA are added together at the time of re-plating in the presence of polybrene (8 μ g/ml). The medium is changed 24 hours after infection with complete medium containing 100 units of recombinant IL-2 per ml. Aliquots of the transduced cells are examined, for Fas-ligand expression, at 48 hours after infection via flow cytometry.

Animals will receive standardized numbers of cytokine-producing cells or control mock-transfected cells intraperitoneally. Concentrated cell suspensions are injected directly into the mouse synovium, as described in section 4A above. In parallel, aliquots of the transferred cell populations are maintained in tissue culture supplemented with exogenous IL-2.

Mice are monitored in a blinded fashion for signs of arthritis. The date of disease onset is recorded and clinical severity of each joint or group of joints (toes, tarsus, ankle, wrist, knee) are graded as follows: 0 (normal), 1 (erythema), 2 (swelling), 3 (deformity), 4 (necrosis). The scores are summed to yield the arthritic score. The severity of arthritis is expressed both as the mean score observed on a given day, and as the mean of the maximal arthritic score reached by each mouse during the clinical course of the disease. At the time of death, hind paws are dissected free and processed for histologic examination or for RT-PCR. The histologic severity of the arthritis is scored on a scale of 0-3 for synovial proliferation and inflammatory cell infiltration, where a score of 0=normal and 3=severe.

For mice receiving intra-synovial injection of control of test adenovirus vector, the level of arthritis observed between contralateral sites is compared. In addition, the overall joint score minus that of the injected joint for the entire animal is compared with that observed in the joint injected with the control or test adenovirus vector.

Local administration of Fas-ligand adenovirus expression vectors will result in clearance of activated cells, as assessed by measuring the relative levels of CD80 mRNA by quantitative RT-PCR. This treatment also will lead to an enhanced level. Also, whether such level of apoptosis identified in affected mouse synovial tissue is assessed by the TUNEL assay ("Terminal deoxynucleotidyl transferase (TdT)-mediated dUTP Nick End Labeling"). TUNEL is performed by immersing the sections in TdT buffer (30 mM Tris-HCl, pH 7.2, 140 mM sodium cacodylate, 1 mM cobalt chloride), and then adding TdT (GIBCO BRL, Grand Island, N.Y.) and biotinylated dUTP (Boehringer Mannheim, Indianapolis, Ind.). The reaction is terminated by immersing the sections in TB buffer (300 mM sodium chloride, 30 mM sodium citrate). Subsequently, the samples are treated with peroxidase-labeled streptavidin and then visualized using the VECTASTAIN ABC kit (Vector Laboratories Inc., Burlingame, Calif.). For immunohistochemistry, the sections are blocked with 4% skim milk for 30 minutes at room temperature, then incubated with biotinylated mAbs specific for mouse CD3, B220, CD80, or CD95 (Fas). These antibodies are available from Pharmingen (San Diego, Calif.).

b. Treatment of Rheumatoid Arthritis Patients with a Gene Therapy Vector Encoding an Accessory Molecule Ligand, Fas Ligand

Candidate Fas-ligand constructs identified as having potential therapeutic benefit are used in human protocols to treat RA. Human protocols encompass either in vivo or ex vivo methods to deliver the Fas-ligand constructs. Furthermore, the Fas-ligand constructs are potentially delivered by either viral or non-viral methods. Outlines of therapeutic strategies are described below.

An ex vivo therapy is similar to a protocol described for intra-articular transplantation of autologous synoviocytes retrovirally transduced to synthesize interleukin-1 receptor antagonist (Evan, Christopher et. al., Clinical Trial to Assess the Safety, Feasibility, and Efficacy of Transferring a Potentially Anti-Arthritic Cytokine Gene to Human Joints with Rheumatoid arthritis, *Human Gene Therapy*, 7:1261-1280). In this procedure, after clinical diagnosis of RA, the synovium is harvested during total joint replacement. The synoviocytes re-isolated and expanded, then transduced or transfected with heterologous Fas-ligand into synoviocytes (via retrovirus, adenovirus, naked DNA, etc.). The gene-modified synoviocytes are then reinjected into the patient, who is monitored and tested for amelioration of RA-associated symptoms, and for expression and function of the Fas-ligand in modified synoviocytes.

In another ex vivo protocol, an allogeneic immortalized cell line that stably expresses the heterologous Fas-ligand is administered to the RA patient. In this protocol, a stable immortalized cell line expressing Fas-ligand (introduced by transfection of the gene into the cell by nonviral methods, such as electroporation), or by viral transduction of the gene into the cell) is constructed. The modified cell line is injected into the patient, who is monitored and tested for amelioration of RA associated symptoms, and for expression and function of the hFas-ligand in modified synoviocytes.

An in vivo based therapy will be similar in concept to the amelioration of collagen-induced-arthritis using a murine

Fas-ligand adenovirus gene therapy vector, described in Zhang, et al., *J. Clin. Invest.* 100:1951-1957 (1997). In our use of such an approach, delivery of the hFas-ligand construct or chimeric ΔfasL directly to the joints of RA patients is performed using either viral or non-viral methods. In this

procedure, the Fas-ligand construct (e.g. hFas-ligand adenovirus) is directly injected into the synovium. Patients are monitored and tested for amelioration of RA-associated symptoms as well as biological testing for expression and function of the hFas-ligand in modified synoviocytes.

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 95

<210> SEQ ID NO 1
<211> LENGTH: 786
<212> TYPE: DNA
<213> ORGANISM: Mus sp.

<400> SEQUENCE: 1

atgatcgaaa catacaacca aacttctccc cgatctgceg ccactggact gcccattcagc	60
atgaaaattt ttatgtattt acttactgtt tttcttatca cccagatgat tgggtcagca	120
ctttttgctg tgtatcttca tagaagggtg gacaagatag aagatgaaag gaatcttcat	180
gaagattttg tattcatgaa aacgatacag agatgcaaca caggagaaag atccttatcc	240
ttactgaact gtgaggagat taaaagccag tttgaaggct ttgtgaagga tataatgtta	300
aacaaagagg agacgaagaa agaaaacagc tttgaaatgc aaaaagggtga tcagaatcct	360
caaattgceg cacatgtcat aagtgaggcc agcagtaaaa caacatctgt gttacagtgg	420
gctgaaaaag gatactacac catgagcaac aacttggtaa ccctggaaaa tgggaaacag	480
ctgaccgtta aaagacaagg actctattat atctatgccc aagtcacctt ctgttccaat	540
cgggaagctt cgagtcaagc tccatttata gccagcctct gcctaaagtc ccccggtaga	600
ttcgagagaa tcttactcag agctgcaaat acccacagtt ccgccaaacc ttgcgggcaa	660
caatccattc acttgggagg agtatttgaa ttgcaaccag gtgcttcggt gtttgtcaat	720
gtgactgatc caagccaagt gagccatggc actggcttca cgtccttttg cttactcaaa	780
ctctga	786

<210> SEQ ID NO 2
<211> LENGTH: 783
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

atgatatgaaa catacagcca accttcccc agatccgtgg caactggact tccagcgagc	60
atgaagattt ttatgtattt acttactgtt ttccttatca cccaaatgat tggatctgtg	120
ctttttgctg tgtatcttca tagaagattg gataaggctg aagaggaagt aaaccttcat	180
gaagattttg tattcataaa aaagctaaag agatgcaaca aaggagaagg atctttatcc	240
ttgctgaact gtgaggagat gagaaggcaa tttgaagacc ttgtcaagga tataacgtta	300
aacaaagaag agaaaaaaga aaacagcttt gaaatgcaa gaggtgatga ggatcctcaa	360
attgcagcac acgttgtaag cgaagccaac agtaatgcag catccgttct acagtgggcc	420
aagaaaggat attataccat gaaaagcaac ttggtaatgc ttgaaaatgg gaaacagctg	480
acgggttaaaa gagaaggact ctattatgtc tacactcaag tcaccttctg ctctaactcg	540
gagccttcga gtcaacgcc attcatcgtc ggctctggc tgaagcccag cattggatct	600
gagagaatct tactcaaggc ggcaaatacc cacagttcct cccagctttg cgagcagcag	660
tctgttcaact tgggcggagt gtttgaatta caagctggtg cttctgtgtt tgtcaacgtg	720
actgaagcaa gccaaagtat ccacagagtt ggcttctcat cttttggctt actcaaactc	780

-continued

tga	783
<210> SEQ ID NO 3 <211> LENGTH: 783 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic chimeric construct <400> SEQUENCE: 3	
atgatcgaaa catacaacca aacttctccc cgatctgcgg ccactggact gcccacagc	60
atgaaaattt ttatgtattt acttactgtt tttcttatca cccagatgat tgggtcagca	120
ctttttgctg tgtatcttca tagaagattg gataaggctg aagaggaagt aaaccttcac	180
gaagattttg tattcataaa aaagctaaag agatgcaaca aaggagaagg atctttatcc	240
ttgctgaact gtgaggagat gagaaggcaa tttgaagacc ttgtcaagga tataacgtta	300
aacaaagaag agaaaaaaga aaacagcttt gaaatgcaaa gaggtgatga ggatcctcaa	360
attgcagcac acgttgtaag cgaagccaac agtaatgcag catccgttct acagtgggcc	420
aagaaaggat attataccat gaaaagcaac ttggtaatgc ttgaaaatgg gaaacagctg	480
acggttaaaa gagaaggact ctattatgtc tacactcaag tcaccttctg ctctaactcg	540
gagccttcga gtcaacgcc attcatcgtc ggctctggc tgaagcccag cattggatct	600
gagagaatct tactcaaggc ggcaaatacc cacagttcct cccagctttg cgagcagcag	660
tctgttcaact tgggcggagt gtttgaatta caagctggtg cttctgtgtt tgtcaacgtg	720
actgaagcaa gccaagtgat ccacagagtt ggcttctcat cttttggctt actcaaactc	780
tga	783
<210> SEQ ID NO 4 <211> LENGTH: 786 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic chimeric construct <400> SEQUENCE: 4	
atgatcgaaa catacaacca aacttctccc cgatctgcgg ccactggact gcccacagc	60
atgaaaattt ttatgtattt acttactgtt ttccttatca cccaaatgat tggatctgtg	120
ctttttgctg tgtatcttca tagaagggtg gacaagatag aagatgaaag gaatcttcac	180
gaagattttg tattcatgaa aacgatacag agatgcaaca caggagaaag atccttatcc	240
ttactgaact gtgaggagat taaaagccag tttgaaggct ttgtgaagga tataatgtta	300
aacaaagagg agacgaagaa agaaaacagc tttgaaatgc aaaaagggtga tcagaatcct	360
caaattgcgg cacatgtcat aagtgaggcc agcagtaaaa caacatctgt gttacagtgg	420
gctgaaaaag gatactacac catgagcaac aacttggtta ccctggaaaa tgggaaacag	480
ctgaccgtta aaagacaagg actctattat atctatgccc aagtcacctt ctgttccaat	540
cgggaagctt cgagtcaagc tccatttata gccagcctct gcctaaagtc ccccggtaga	600
ttcgagagaa tcttactcag agctgcaaat acccacagtt ccgccaaacc ttgcgggcaa	660
caatccattc acttgggagg agtatttgaa ttgcaaccag gtgcttcggt gtttgtcaat	720
gtgactgac caagccaagt gagccatggc actggcttca cgtccttttg cttactcaaa	780
ctctga	786

-continued

<210> SEQ ID NO 5		
<211> LENGTH: 783		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic chimeric construct		
<400> SEQUENCE: 5		
atgatcgaaa catacaacca aacttctccc cgatctgcgg ccactggact gcccacgagc	60	
atgaaaattt ttatgtattt acttactgtt ttccttatca cccaaatgat tggatctgtg	120	
ctttttgctg tgtatcttca tagaagattg gataaggtcg aagaggaagt aaaccttcac	180	
gaagattttg tattcataaa aaagctaaag agatgcaaca aaggagaagg atctttatcc	240	
ttgctgaact gtgaggagat gagaaggcaa tttgaagacc ttgtcaagga tataacgtta	300	
aacaaagaag agaaaaaaga aaacagcttt gaaatgcaaa gaggtgatga ggatcctcaa	360	
attgcagcac acgttgtaag cgaagccaac agtaatgcag catccgttct acagtgggcc	420	
aagaaaggat attataccat gaaaagcaac ttggtaatgc ttgaaaatgg gaaacagctg	480	
acggttaaaa gagaaggact ctattatgtc tacactcaag tcaccttctg ctctaacgg	540	
gagccttcga gtcaacgccc attcatcgtc ggctctcggc tgaagcccag cattggatct	600	
gagagaatct tactcaaggc ggcaaatacc cacagttcct ccagctttg cgagcagcag	660	
tctgttcaact tgggcggagt gtttgaatta caagctggtg cttctgtgtt tgtcaacgtg	720	
actgaagcaa gccaaagtgat ccacagagtt ggcttctcat cttttggctt actcaaactc	780	
tga	783	
<210> SEQ ID NO 6		
<211> LENGTH: 786		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic chimeric construct		
<400> SEQUENCE: 6		
atgatatgaaa catacagcca accttcccc agatccgtgg caactggact tccagcgagc	60	
atgaagattt ttatgtattt acttactgtt tttcttatca ccagatgat tgggtcagca	120	
ctttttgctg tgtatcttca tagaagggtg gacaagatag aagatgaaag gaatcttcat	180	
gaagattttg tattcatgaa aacgatacag agatgcaaca caggagaaaag atccttatcc	240	
ttactgaact gtgaggagat taaaagccag tttgaaggct ttgtgaagga tataatgtta	300	
aacaaagagg agacgaagaa agaaaacagc tttgaaatgc aaaaagggtga tcagaatcct	360	
caaattgctg cacatgtcat aagtgaggcc agcagtaaaa caacatctgt gttacagtgg	420	
gctgaaaaag gatactacac catgagcaac aacttggtta ccctggaaaa tgggaaacag	480	
ctgaccgtta aaagacaagg actctattat atctatgcc aagtcacctt ctgttccaat	540	
cgggaagctt cgagtcaagc tccatttata gccagcctct gcctaaagtc ccccggtaga	600	
ttcgagagaa tcttactcag agctgcaaat acccacagtt ccgccaaacc ttgcgggcaa	660	
caatccattc acttgggagg agtatttgaa ttgcaaccag gtgcttcggt gtttgtcaat	720	
gtgactgata caagccaagt gagccatggc actggcttca cgtccttttg ctactcaaa	780	
ctctga	786	
<210> SEQ ID NO 7		
<211> LENGTH: 786		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		

-continued

<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic chimeric construct	
<400> SEQUENCE: 7	
atgatagaaa catacagcca accttcccc agatccgtgg caactggact tccagcgagc	60
atgaagattt ttatgtattt acttactgtt ttccttatca cccaaatgat tggatctgtg	120
ctttttgctg tgtatcttca tagaagggtg gacaagatag aagatgaaag gaatcttcat	180
gaagattttg tattcatgaa aacgatacag agatgcaaca caggagaaag atccttatcc	240
ttactgaact gtgaggagat taaaagccag tttgaaggct ttgtgaagga tataatgtta	300
aacaaagagg agacgaagaa agaaaacagc tttgaaatgc aaaaagggtga tcagaatcct	360
caaattgcgg cacatgtcat aagtgaggcc agcagtaaaa caacatctgt gttacagtgg	420
gctgaaaaag gatactacac catgagcaac aacttggtaa ccctggaaaa tgggaaacag	480
ctgaccgtta aaagacaagg actctattat atctatgccc aagtcacctt ctgttccaat	540
cggaagcgtt cgagtcaagc tccatttata gccagcctct gcctaaagtc ccccggtaga	600
ttcgagagaa tcttactcag agctgcaaat acccacagtt ccgccaaacc ttgcgggcaa	660
caatccattc acttgggagg agtat ttgaa ttgcaaccag gtgcttcggt gtttgtcaat	720
gtgactgata caagccaagt gagccatggc actgggttca cgtccttttg cttactcaaa	780
ctctga	786
<210> SEQ ID NO 8	
<211> LENGTH: 864	
<212> TYPE: DNA	
<213> ORGANISM: Bos sp.	
<400> SEQUENCE: 8	
aactctaacg cagcatgata gaaacataca gtcaaccttc tccccgctcc gtggccactg	60
gaccacctgt cagtatgaaa atttttatgt atttacttac agtttttctt atcaccaga	120
tgattgggtc agcgcttttt gctgtgtatc ttcacagacg attggacaag atagaagacg	180
aaaggaatct tcatgaagat tttgtgttca tgaaaacgat acagagatgc aataaaggag	240
aggggtcctt atccttactg aactgtgagg aaattagaag ccggtttgaa gacttggtca	300
aggatataat gcaaaacaaa gaagtaaaga agaaagaaaa aaactttgaa atgcacaagg	360
gtgatcagga gcctcagata gcggcacatg tcatcagtga ggccagtagt aaaacaacct	420
ctgttctcca gtgggcccc aaaggatact acaccctaag caacaacctg gtaaccctcg	480
aaaacgggaa acagctggcc gtgaaaagac aaggattcta ttacatctac acccaagtca	540
ccttctgttc caatcgggaa actttgagtc aagctccatt tatagccagc ctctgcctga	600
agtccccaag tggatcagag agaattctac tgagagctgc aaacaccac agttcttcca	660
aaccatgcgg gcagcaatcc attcacttag gaggagtctt tgaattgcaa tcgggtgctt	720
cggtgtttgt caatgtgact gatccaagtc aagtgagcca cgggacgggc ttcacatcat	780
ttggcttact caaactctga acggtgtaag ccagcaggct gcggctgggc tgatgctggt	840
ggtcttcaca atccaggaaa gcag	864
<210> SEQ ID NO 9	
<211> LENGTH: 3634	
<212> TYPE: DNA	
<213> ORGANISM: Homo sapiens	
<400> SEQUENCE: 9	
gaattccggg tgatttcact cccggctgtc caggcttgtc ctgctacccc acccagcctt	60

-continued

tcttgaggcc	tcaagcctgc	caccaagccc	ccagctcctt	ctccccgcag	gacccaaaca	120
caggcctcag	gactcaacac	agctttttccc	tccaacccgt	tttctctccc	tcaacggact	180
cagctttctg	aagccccctc	cagttctagt	tctatctttt	tcttgcattc	tgtctggaag	240
ttagaaggaa	acagaccaca	gacctgggtc	ccaaaagaaa	tggaggcaat	aggttttgag	300
gggcatgggg	acgggggttca	gcctccaggg	tcctacacac	aaatcagtca	gtggcccaga	360
agacccccct	cggaatcgga	gcagggagga	tggggagtgt	gaggggtatc	cttgatgctt	420
gtgtgtcccc	aactttccaa	atccccgccc	ccgcgatgga	gaagaaaccg	agacagaagg	480
tgcagggccc	actaccgctt	cctccagatg	agctcatggg	tttctccacc	aaggaagttt	540
tccgctgggt	gaatgattct	ttccccgccc	tcctctcgcc	ccagggacat	ataaaggcag	600
ttgttggcac	accagccag	cagacgctcc	ctcagcaagg	acagcagagg	accagctaag	660
agggagagaa	gcaactacag	acccccctg	aaaacaaccc	tcagacgcca	catccccga	720
caagctgcca	ggcaggttct	cttctctca	catactgacc	cacggcttca	ccctctctcc	780
cctggaaagg	acaccatgag	cactgaaagc	atgatccggg	acgtggagct	ggccgaggag	840
gcgctcccca	agaagacagg	ggggccccag	ggctccaggc	ggtgcttggt	cctcagcctc	900
ttctccttcc	tgatcgtagc	aggcgccacc	acgtctttct	gcctgctgca	ctttggagtg	960
atcgggcccc	agaggaaga	ggtgagtgcc	tggccagcct	tcattccactc	tcccacccaa	1020
ggggaaatga	gagacgcaag	agagggagag	agatgggatg	ggtgaaagat	gtgcgctgat	1080
agggagggat	gagagagaaa	aaaacatgga	gaaagacggg	gatgcagaaa	gagatgtggc	1140
aagagatggg	gaagagagag	agagaaagat	ggagagacag	gatgtctggc	acatggaagg	1200
tgctcactaa	gtgtgtatgg	agtgaatgaa	tgaatgaatg	aatgaacaag	cagatatata	1260
aataagatat	ggagacagat	gtgggggtgtg	agaagagaga	tgggggaaga	aacaagtgat	1320
atgaataaag	atggtgagac	agaaagagcg	ggaaatatga	cagctaagga	gagagatggg	1380
ggagataagg	agagaagaag	ataggggtgc	tggcacacag	aagacactca	gggaaagagc	1440
tgttgaatgc	tggaaggtga	atacacagat	gaatggagag	agaaaaccag	acacctcagg	1500
gctaagagcg	caggccagac	aggcagccag	ctgttcctcc	tttaagggtg	actccctcga	1560
tgtaaacat	tctccttctc	cccaacagtt	cccaggggac	ctctctctaa	tcagccctct	1620
ggcccaggca	gtcagtaagt	gtctccaaac	ctctttccta	attctgggtt	tgggtttggg	1680
ggtagggtta	gtaccggtat	ggaagcagtg	ggggaaatth	aaagtthtgg	tcttggggga	1740
ggatggatgg	aggatgaaat	aggggggtat	tttctaggaa	gtttaagggt	ctcagcttht	1800
tctthtctct	ctcctcttca	ggatcatctt	ctcgaacccc	gagtgcacag	cctgtagccc	1860
atgttgtagg	taagagctct	gaggatgtgt	cttggaactt	ggagggttag	gatttgggga	1920
ttgaagcccc	gctgatggta	ggcagaactt	ggagacaatg	tgagaaggac	tcgctgagct	1980
caaggggaagg	gtggaggaac	agcacaggcc	ttagtgggat	actcagaacg	tcattggccag	2040
gtgggatgtg	ggatgacaga	cagagaggac	aggaaccgga	tgtgggtgtg	gcagagctcg	2100
agggccagga	tgtggagagt	gaaccgacat	ggccacactg	actctcctct	ccctctctcc	2160
ctccctccag	caaaccctca	agctgagggg	cagctccagt	ggctgaaccg	ccgggccaat	2220
gccctcctgg	ccaatggcgt	ggagctgaga	gataaccagc	tgggtgtgtc	atcagagggc	2280
ctgtacctca	tctactccca	ggtcctcttc	aagggccaaag	gctgccccctc	cacctatgtg	2340
ctcctcacc	acaccatcag	ccgcatcgcc	gtctcctacc	agaccaaggt	caacctctct	2400
tctgccatca	agagcccctg	ccagagggag	accccagagg	gggctgaggc	caagccctgg	2460

-continued

tatgagccca	tctatctggg	aggggtcttc	cagctggaga	agggtgaccg	actcagcgct	2520
gagatcaatc	ggcccgacta	tctcgacttt	gccgagtctg	ggcaggtcta	ctttgggata	2580
attgccctgt	gaggaggacg	aacatccaac	cttcccaaac	gcctcccctg	ccccaatccc	2640
tttattaccc	cctccttcag	acaccctcaa	cctcttcttg	ctcaaaaaga	gaattggggg	2700
cttagggtcg	gaacccaagc	ttagaacttt	aagcaacaag	accaccactt	cgaaacctgg	2760
gattcaggaa	tgtgtggcct	gcacagtga	gtgtggcaa	ccactaagaa	ttcaaactgg	2820
ggcctccaga	actcactggg	gcctacagct	ttgatccctg	acatctggaa	tctggagacc	2880
agggagcctt	tggttctggc	cagaatgctg	caggacttga	gaagacctca	cctagaaatt	2940
gacacaagtg	gaccttaggc	cttcctctct	ccagatgttt	ccagacttcc	ttgagacacg	3000
gagcccagcc	ctcccatgg	agccagctcc	ctctatttat	gtttgcactt	gtgattattt	3060
attatttatt	tattatttat	ttatttacag	atgaatgtat	ttatttggga	gaccggggta	3120
tcctggggga	cccaatgtag	gagctgcctt	ggctcagaca	tgttttccgt	gaaaacggag	3180
ctgaacaata	ggctgttccc	atgtagcccc	ctggcctctg	tgccttcttt	tgattatgtt	3240
ttttaaaata	tttatctgat	taagttgtct	aaacaatgct	gatttgggtga	ccaactgtca	3300
ctcattgctg	agcctctgct	ccccagggga	gttgtgtctg	taatcgccct	actattcagt	3360
ggcgagaaat	aaagtttgct	tagaaaagaa	acatgggtctc	cttcttggaa	ttaattctgc	3420
atctgcctct	tcttgtgggt	gggaagaagc	tcctaagtc	ctctctccac	aggetttaag	3480
atccctcgga	cccagtccca	tccttagact	cctagggccc	tggagaccct	acataaacia	3540
agcccaacag	aatattcccc	atcccccagg	aaacaagagc	ctgaacctaa	ttacctctcc	3600
ctcagggcat	gggaatttcc	aactctggga	attc			3634
<210> SEQ ID NO 10						
<211> LENGTH: 1997						
<212> TYPE: DNA						
<213> ORGANISM: Homo sapiens						
<400> SEQUENCE: 10						
gagacagagt	cttgctctgt	ccccaggct	ggaatacagt	ggtgcgatct	tgactcactg	60
cagcctccgc	ctcccagggt	caaataattc	tccagcctca	gcctcccag	tagctgggac	120
tgcagatgcg	caccagcacg	cctggcta	ttttgtat	attatagaga	tggggtttca	180
ccatgttggc	cagctgggtct	caaactcctg	acctcaagta	atccgcccac	ctcagactcc	240
caaagtgcc	ggattacagg	tgtgagccac	tgcaccaggc	ctggaacaat	tttaaaataa	300
tgtattggct	ctgcaaatgc	agcttcagaa	caagtccctt	agctgtcccc	acccaccct	360
aagtcaccac	ccttaagcct	cacccatgtg	gaattctgaa	acttcctttg	tagaaaactt	420
tggaagggtg	ctgccacatt	gatectggaa	tgtgtgttta	tttgggggta	tataaatctg	480
ttctgtggaa	gccacctgaa	gtcaggaaga	gatggagggc	atccttcagg	agtgagatga	540
gacctcatca	tacttgactg	tccagcatca	tctctgagta	aggggaccaa	aaaatttatc	600
ttccaaacta	ggacactttc	aagagtggaa	gggggatcca	ttaatatttt	cacctggaca	660
agaggcaaac	accagaatgt	ccccgatgaa	ggggatatat	aatggacctt	cttgatgtga	720
aacctgccag	atgggctgga	aagtccgtat	actgggacaa	gtatgatttg	agttgtttgg	780
gacaaggaca	ggggtacaag	agaaggaaat	gggcaaagag	agaagcctgt	actcagccaa	840
gggtgcagag	atgttatata	tgattgctct	tcagggaacc	gggcctccag	ctcacacccc	900
agctgctcaa	ccacctctc	tctgaattga	ctgtcccttc	tttggaaactc	taggcctgac	960

-continued

cccaactccct	ggccctccca	gcccacgatt	cccctgaccc	gactcccttt	cccagaactc	1020
agtcgcctga	acccccagcc	tgtggttctc	tcctaggcct	cagcctttcc	tgcctttgac	1080
tgaaacagca	gtatcttcta	agccctgggg	gcttccccgg	gccccagccc	cgacctagaa	1140
cccgcccgct	gcctgccacg	ctgccactgc	cgcttcctct	ataaagggac	ctgagcgtcc	1200
gggcccaggg	gctccgcaca	gcaggtagagg	ctctcctgcc	ccatctcctt	gggctgcccg	1260
tgcttcgtgc	tttgactac	cgcccagcag	tgtcctgccc	tctgcctggg	cctcggtccc	1320
tcctgcacct	gctgcctgga	tccccggcct	gcctgggcct	gggcttggtg	ggtttggttt	1380
tggtttcctt	ctctgtctct	gactctccat	ctgtcagtct	cattgtctct	gtcacacatt	1440
ctctgtttct	gccatgattc	ctctctgttc	ccttcctgtc	tctctctgtc	tcctctgct	1500
caccttgggg	tttctctgac	tgcattctgt	ccccttctct	gtcgatctct	ctctcggggg	1560
tcgggggggtg	ctctctccca	gggcgggagg	tctgtcttcc	gccgcgtgcc	ccgccccgct	1620
cactgtctct	ctctctctct	ctctttctct	gcaggttctc	cccatgacac	cacctgaacg	1680
tctcttctct	ccaagggtgt	gtggcaccac	cctacacctc	ctccttctgg	ggctgctgct	1740
ggttctgctg	cctggggccc	aggtagagga	gcaggagaat	gggggctgct	ggggtggctc	1800
agccaaacct	tgagccctag	agccccctc	aactctgttc	tcccctaggg	gtccctgggt	1860
gttggcctca	caccttcagc	tgccagact	gcccgtcagc	acccaagat	gcattctgcc	1920
cacagcacc	tcaaacctgc	tgtcacctc	attggtaaac	atccacctga	cctcccagac	1980
atgtccccac	cagctct					1997
<210> SEQ ID NO 11						
<211> LENGTH: 10240						
<212> TYPE: DNA						
<213> ORGANISM: Sus sp.						
<400> SEQUENCE: 11						
gaattccccg	gatcaaagtc	agcattaaat	cccagtttag	gttttgaggc	taagttcaag	60
tttgagtcta	atgtcatttc	agccttgttt	ggaggactca	gagatttcac	tagtttctcc	120
gcagagacca	ctgtagaaac	tgcatttccc	tgagttttgg	gcacaagact	ccagtcatca	180
cccctcccac	acagggaaa	ccccaaacca	actgctggcc	tcctcaagaa	agaaaccgaa	240
tttcacacaa	cctccgaaac	taagattgaa	accaagattg	gcccattctc	aggcgcgtcc	300
tccagcacat	tgagaatgtc	gctgatggag	cctcggccca	gctctcgagc	ttccttcctt	360
tctgtctctc	atgtcttctc	atcactcctt	ctcaccttcc	cgtttttgtc	ctgcaatgcc	420
cccttcttcc	tctcttcctg	gggtttttcc	ctttattttct	cactgtacca	ttttatatatt	480
taataaagcc	gaggtctcct	agtccatcag	ctcctactgt	tggagaggag	gcagaaagaa	540
acagcaggac	ggcaaaggga	ctccagagaa	agagactcag	aggaaaggca	agaaacaggg	600
accaagagag	aggccaacag	tgacacaaga	cacagtgagg	ttaaaagaaa	taagatgagg	660
ccaagataga	gaccaagcta	tttaaaagag	ccatctgtgg	ctacccttct	tcgccatcg	720
catctggtca	gccaccaaga	ttttgcctag	aaacgttccct	cctctccatt	ctcctgctgc	780
tgtgtgtgct	gctgtgtgtg	ctgtgtgtgc	tgtgtgtgct	gctgtgtgtg	ctgccttaat	840
acgaatgcag	gctcttggtca	tctccttgct	gggttggtgc	aaaatcctcc	taactgggtct	900
ccacacttct	catttcccct	ccagcccccc	atcttccata	cttccattta	tttattttgg	960
ccatgcccac	ggcatgtggc	agttccaggg	gccagggatc	aaacctgtgc	caatgcagtg	1020
accgtgtcag	atccttaacc	cactgcacac	aaggcaacgc	ccctcgagtc	attctcattt	1080

-continued

tttaaatata	ccaatttgag	ggggtccttc	tttcacttaa	aaattttggc	agctccctat	1140
catgatgaga	aggaattcca	aaccattttt	cttgtgtgca	aacccttcag	catgtgtcct	1200
cagcttactt	cccaagcctc	atccctgctc	cttctacgtg	taccatgtg	tacatctcca	1260
cacaccatat	actctttttt	acctcccatc	tttgacctt	ctgttccttc	tctctgcccc	1320
tcaccatctt	ttttgctttg	atacttaatg	cctctccctc	aggccagggt	caatggcttt	1380
tctgtgggct	gctttaagcc	cactgtcatg	gaacttatca	cattttattt	tatttgactt	1440
tctttttagg	gccgcaccca	gcatatggag	attcccaggc	tagggatcta	atcggagctg	1500
tatctgccag	cctgcgctgg	agccacagca	acgtgggata	cgagcctgag	gggttttgat	1560
gtcctgtggc	acagaagtta	cattcaggct	gtgcatgaac	tatttctcct	gttctcctcc	1620
ccctgcttga	ggccctgcag	ctttgcctct	catgccttgc	tgctctgacc	tatgacttct	1680
ttttgtttgc	attccatctc	tttagttttc	tctctgttcc	acaaacattt	actgagcatc	1740
tacatgaggc	attgaggata	cggatgggaa	agacagtccc	ctgacctctg	ggacctcaaa	1800
gaccaattgt	ggaagactgg	ttggttatca	gataattaca	atgaagtgtg	ggagtccctg	1860
tcatgggtca	gcaggtaatg	aaccagtaa	acgatccatg	aggatgcaga	ttcaatccct	1920
ggccttgctc	agcgggttaa	ggatccagcg	ttcccacaag	ctgtggtgta	ggtcgcagat	1980
gcgactcaga	tcttgcatth	ctgtggctgt	ggtgtaggct	ggtggctacc	cctagcctgg	2040
gaacctccat	atgcctcagg	tgccggccct	aaagacaaaa	aaaaaaaaa	gagaaacttt	2100
tctttttctt	aatgtgtaac	ctacaagcta	agtgaaaact	ggctcctatt	ccataacgth	2160
tgtatcattt	ttcatactag	ccaaatacta	gaaacaggga	gttcccgtcg	tggtgcagca	2220
gaaacaaatt	cgactaggaa	ccatgagggt	gcgggttcga	tccttgccct	tgctcagtgg	2280
gttaaggata	cggcgttgcc	gtgagctgtg	gtgtaggctg	cagatgtggc	tcggatctag	2340
tgttgctgtg	gctctggtgt	aggccggcag	caacagctct	gattagactc	ctagcctgag	2400
aacctccata	agctgtggct	gcggccctat	aaagacaaaa	aaaaaaaaa	ggccaaatac	2460
tagaaacaaa	ccaaatgcc	atcaacagaa	gaatagataa	gttaattggg	gtatatgcac	2520
acaatagcat	cacacaataa	catgcacaca	ataacatcac	aatgaaataa	aaattactac	2580
tgacagacac	aaccatatag	atgaatttca	caaacacaac	agcgagaata	aaagccaagc	2640
acagatgagt	tgtctgtgtg	gattcatttc	tatgaagttc	aagcgcagga	agaacttaat	2700
ctatagtga	agaggtcaga	gagcagttgg	ttgtctttgg	caggtatgaa	ctgggagtgg	2760
gcatgagaga	actttctgga	gacctaaaaa	tatattggac	tggtatgggt	caacatggct	2820
acaagaagat	ggaaaagtth	ctcaggctgt	ccacttgga	gacgggcttc	tcacgggacc	2880
taagttctgc	atcagcagag	ggggaaatcc	ttaatgattt	gacaattaca	aagtgtattg	2940
gctttaccga	tgtattttca	acacaatccc	tctgctgtcc	ccacccacc	ctaggtcacc	3000
acccttaagc	tcacactgtg	tggaattctg	aagcctcccc	tgtagagaac	tttagcagth	3060
gccacgttct	tttgatgcag	gaacgtgttg	tctagagtta	gacacatctg	atctgtgggg	3120
cccacccaag	gttgggacat	ggtggggggc	ggccttctgc	agtgagatga	aacctcattg	3180
taggtgattt	cgtggcctca	tcctgagtc	agatcttcca	aatgaggaca	ctttggagag	3240
caaaaggggg	ctccctgaag	atttcctcca	ggacagcagg	aacaaaccag	gatgtcccag	3300
gcaggagggg	atagaaggga	acttgttgat	atgaaatcag	ccagatgacc	tggaaaatac	3360
acagactggg	acaagtgtga	cttgagcctc	ttgggcccag	gacaggggta	cagaggagga	3420
aacgtgcaca	gagagaagcc	cgtaatcagc	caaggctgca	gaggtgttat	acataatcgc	3480

-continued

tcttcacgca	accgggcaag	cagcccacgc	cccagctgca	ctccatctcc	tcctctgaac	3540
tcaccgtccc	ttctctggaa	ctcctaagcc	tgaccccgct	ccctggccct	cccagcccac	3600
ggttcccctg	accccactcc	ctttcccaga	actcagtcac	ctgagccccc	agcctgcgtt	3660
ctctcctagg	cctcagcctt	tcctgccttc	gcgtgaaaca	gcagcatctt	ctaagccctg	3720
ggcttcccca	ggccccagcc	cgggcctaga	acccgcccag	ccgacctgcc	cacgctgcca	3780
ctgccggctt	cctctataaa	gggacccagg	gcgccagaa	aggggcccac	aggggtcccg	3840
cacagcaggt	gagactctcc	caccccatct	cctagggctg	tccgggtgct	ggactcccc	3900
ctcaactcgg	tcctccgcc	cgtccctgg	ccttcctgcc	cctcctgcat	cttcaccccg	3960
gcctgggcct	tgggtgggtt	ggttttgggt	tgttctctct	gattctttat	ctgtcaggct	4020
ctttctagct	ctcacacact	ctgatccctc	tctgttccct	tcccatctct	gtttctctct	4080
gggtctcccc	ctgctcacct	cgggatttcc	ctgagtgcct	ctgggtccct	tctctgtctg	4140
gcgccccgtc	tcttgtctct	cgggggtggc	gtctccgagg	gcaggaggcc	ttcttccgca	4200
gggtccccgc	cccgtccact	gtctctctcc	ccccacaggt	tttcccatg	acaccacctg	4260
gacgcctcta	cctccggagg	gtgtgcagca	ccccatcct	cctcctcctg	gggtgctgc	4320
tggccctgcc	gcccgaggcc	caggtgaggc	agcaggagag	cgggcccgtg	gggcagcctt	4380
cgccaacctt	gggcctcaga	gcctctctga	cgtctttctc	ccctaggggc	tcctggcgt	4440
cggcctccca	ccctcagctg	cacagcctgc	ccatcagcac	ccccaaaagc	acttgccag	4500
aggcacctc	aaacctgccg	ctcacctcgt	tggtaaacad	ccacctggcc	tcccagacct	4560
gtagccccca	gtcctcctcc	tatgccctg	cttcaggagc	tgaagcatcc	ctccccccca	4620
tctcccccca	ccccctaaat	ggaggcatcc	cactcccgac	tccctcccaa	ccatccccca	4680
ggaactcagt	ccagcacctg	cttcctcagg	gattgagacc	tccgaccccc	aggtccttga	4740
ctcccacccc	ctctggctct	tcctaggaga	cccagcacc	ccggactcac	tgcgtggag	4800
agcgaacacg	gategtgcct	tcctccgcca	tggcttcttg	ctgagcaaca	actccctgct	4860
ggccccacc	agtggcctct	actttgtcta	ctccaggctc	gtcttctccg	gggaaggctg	4920
cttccccaa	gccaccccca	cccctctcta	cctggcccac	gaggtccagc	tcttctcctc	4980
ccagtacccc	ttccacgtgc	cgtcctcag	cgtcagaag	tccgtgtgcc	ccgggccaca	5040
gggaccttgg	gtgcgtctct	tgtaccagg	ggctgtgttc	ctgctcacc	aggagatca	5100
gctgtccaca	cacacagacg	gcacccccca	cctgctcctc	agccccagta	gcgtcttctt	5160
tggagccttc	gctctataga	agaatccaga	aagaaaaaaa	ttggtttcaa	ggccttctcc	5220
ccttttcacc	tcccttatga	ccacttcgga	ggcacccg	cctctcctct	gacaatttcc	5280
aacagtctca	tcttccccca	cgtcagcac	ctggagcttc	tgtagaagga	attctaggca	5340
cctcggggga	actggaacca	ccccgatgc	tctgctgagg	atctgaatgc	ccgcctggag	5400
cccttcccct	gtcctgccc	tctagggg	ctcgtccagg	acgtggaagg	gaagctgacc	5460
catgagggg	tttgaacgga	tgaccggagc	gggtgtgggg	ggttatttat	gaaggggaaa	5520
attaaattat	ttatttatgg	aggatggaga	gaagggaatc	acagagggat	gtcagaagag	5580
tgtgacacat	gtgccaaga	gataaagtga	cagaaggcat	gggctccaga	tgaccgggcc	5640
agagagggga	aagtggctca	ggaaggggct	gcttgactgg	aggctcatga	ggagacggct	5700
gaccctcgat	gaaacccaat	aaagctcttt	tctctgaaat	gctgtctgct	cgtatctgtc	5760
actcgggagg	ggagaattct	ccagatgtct	ctaaggagtg	gagggaggac	aggaatcaga	5820
ggggacggga	gctgtgggtg	tgtgatgagg	cctaaggggc	tcaggtgaga	gatggcggcc	5880

-continued

tcagggtgag	ggcagccaga	ccccgcagg	agaagcagat	ggttcctctg	agaagacaaa	5940
ggaagagatg	cagggccaag	gtcttgagaa	ccgaggtcgg	gggtcgcttg	gcagatatgg	6000
ccacaggtag	agggaacagag	gaataggggt	gacaggaggc	ttcccgggag	aagggaacac	6060
actgaggggt	gttcgggatt	ctgagggagg	agcacgggga	cgccctggga	gacatgccgt	6120
ccagggccat	gaggagtggg	agagcctctg	aggctagcgg	ctggagatac	agggaacattt	6180
gaggagacac	ggtcatggcc	aggagccgcg	agggcctgga	cagtctctag	gaatctcgaa	6240
gaagcaggaa	ttctttgagg	atacgtggcc	acacaaaggg	aggctgaggt	gtggggactt	6300
catgcagaag	tcagggcctc	acattccctt	ggaagccgag	actgaaacca	gcagcagagt	6360
tttggtgagt	tcctgtcaga	gtgaaaggag	aaggcccggc	atggtggggt	tgtgaattcc	6420
cagcctggct	tcctctccct	ctggggctgt	cccaggcctg	ttcctgccgt	cctccccag	6480
cccgtgtagg	gcctccagct	gcccttctcc	cagctcctct	tcctccagg	agacgaaaca	6540
tgggtctcag	caccagcgc	gggtgtgtct	aagttttctc	tccattaaga	actcagcttt	6600
ctgaagctcc	tcccattcct	agttctaccc	ctacctgagc	cctgttcgga	aatcagagag	6660
aaatagaagt	catcccccaa	agaaaaggaa	tttgtcccc	aaagaaacag	aacttgtccc	6720
ccaaagaaat	ggaaacaatg	ggaaatggga	ggcagggggg	acctggggtc	cagcctccag	6780
ggtcctacac	acagagcagt	aactggccca	gcaagcccac	ctcaggatcc	gggcagggag	6840
ggtaggaagt	atccctgatg	cctgggtgtc	cccaactttc	caaaccgccg	cccccgctat	6900
ggagatgaaa	ctaagacaga	aggtgcaggg	cccgtaccg	cttcctccag	atgagctcat	6960
gggtttctcc	accaaggaag	ttttccgctg	gttgaaagag	agcctctccc	cgccctcttc	7020
tcaccagag	cgtataaatg	cagctgtttg	cacaccagc	cagcagaagc	tcccagagt	7080
aggacaccag	gggaccagcc	aggagagaga	caagccatct	ccaggacccc	ctagaaataa	7140
cctctcagaa	gacacacccc	cgaacaggca	gccggacgac	tctctccctc	tcacacgctg	7200
ccccggggcg	ccaccatctc	ccagctggac	ctgagcccct	ctgaaaaaga	caccatgagc	7260
actgagagca	tgatccgaga	cgtggagctg	gcggaggagg	cgctcgccaa	gaaggccggg	7320
ggcccccagg	gctccaggag	gtgcctgtgc	ctcagcctct	tctccttctc	cctggtcgca	7380
ggagccacca	cgctcttctg	cctactgcac	ttcgaggtta	tcggccccca	gaaggaagag	7440
gtgagcgcct	ggccagcctt	ggctcattct	cccacccgga	gagaaatggg	gaagaaagag	7500
ggccagagac	gagctggggg	aaagaagtgt	gctgatgggg	agtgtgggga	ggaaatcatg	7560
gagaaagatg	gggaggcaga	aggagacgtg	gagagagatg	gggggagaga	gagaaggatg	7620
gagagaaatc	cgggtggccc	gcccttgga	atgctctcta	aatatttggt	gcacgaatga	7680
gtgagtaagc	agggaacccg	atataaagag	agatgagtag	acagacaagg	gggtgtggtag	7740
aaagataggg	aaaaaacaag	tgatctggat	aaagatagtg	agacaggaag	aggtagagga	7800
gataggaaag	agagataagg	agagaagaag	gaagcgtggg	tgtctggcac	gtggaaggca	7860
ctcaatgaag	gagttgttga	atggatgggt	ggatgagaaa	atggatgagt	ggagagaaaa	7920
aactagacat	cagggcagag	agtacaagct	agagaagcag	gtggctgttt	tccttcaga	7980
ggggacttat	tcaaacttaa	ttaatccttc	ttcttctccc	caacagtttc	cagctggccc	8040
cttgagcatc	aaccctctgg	cccaaggact	cagtaagtat	ctctaaaacc	tgtctctcag	8100
ttctgagctt	ggacaggggt	ggggtagtg	ctgggtgga	aggaagaagg	gaaatttagg	8160
gtctgggttt	ggcgggggga	atgcaggcca	aagtagtgag	atattttctg	ggaagtctga	8220
gggtctcatc	tttttctttc	ctctttcctc	ctcaggatca	tcgtctcaaa	cctcagataa	8280

-continued

gcccgtcgcc	cacgttgtag	gtaagagttc	tgaggatgtg	tctgggggat	gaagaaatag	8340
gcaggacaga	gagggatagg	atttgggggc	tgaagccagg	ctgagggtag	ccagagcttg	8400
gagatagtat	gaggaggact	cgctgagctc	caggggagga	tgggggatac	tcagaacttg	8460
aggaggatac	tcggaacctc	atggacagat	gggatgtggg	aagacagacc	gaggggacag	8520
gaaccggatg	tggggggcgg	gcagaactcg	agggccagga	tgtggagagt	ggaactgaca	8580
gggtcacact	gactaccccc	tccctctttg	tctcctccct	ccagccaatg	tcaaagccga	8640
gggacagctc	caatggcaga	gtgggtatgc	caatgccctc	ctggccaacg	gcgtgaagct	8700
gaaagacaac	cagctggtgg	tgccgacaga	tgggctgtac	ctcatctact	cccaggctct	8760
cttcaggggc	caaggctgcc	cttccaccaa	cgttttcttc	actcacacca	tcagccgcat	8820
cgccgtctcc	taccagacca	aggtcaacct	cctctctgcc	atcaagagcc	cttgccagag	8880
ggagaccccc	gagggggcgg	aggccaagcc	ctggtacgaa	cccatctacc	tgggaggggt	8940
cttccagctg	gagaaggatg	atcgactcag	tgccgagatc	aacctgcccg	actatctgga	9000
ctttgctgaa	tctgggcagg	tctattttgg	gatcattgcc	ctgtgagggg	gcaggacatc	9060
cgttccctcc	cctgtccatc	cctttattat	tttactcctt	cagaccccct	cacgtccttc	9120
tggttttagaa	agagaatgag	gggctgggga	ctgggctcca	agcttaaaac	tttaaacaac	9180
aacagcaaca	cttagaaatc	agggattcag	ggatgtgtgg	cctggacaac	caggcactga	9240
ccaccaccaa	gaattggaac	tggggcttcc	agactcgctg	gggtccttgg	gtttggattc	9300
ctggatgcaa	cctgggacat	ctggaatgtg	gctgccaggg	aagcttgggg	tccaatcgga	9360
atacttcaga	acatttcctg	agaagatttc	acctcaatct	tgatgacttt	ttaggcttcc	9420
ctttcttcca	attttccaga	cttccctggg	atggggagcc	cagccccaaa	ccccacaggc	9480
cagctccctc	ttatttatat	ttgcacttgg	cattattatt	tatttattta	tttattattt	9540
atttactagt	gaatgtattt	attcaggagg	gcgaggtgtc	ctgggagacc	cagcataagg	9600
gctgccttgg	ttcagatgtg	ttttctgtga	aaacggagct	gaactgtagg	ttgctccac	9660
ctggcctcct	agcctctgtg	cctccttttg	cttatgtttt	taaaaacaaa	tatttatctg	9720
atcgagttgt	ctaaataatg	ctgatttggg	gactaacttg	tcgctacatc	gctgaacctc	9780
tgctccccag	gggagttgtg	tctgtaaccg	ccctactggg	cagtggcgag	aaataaaaagc	9840
gtgcttagaa	aagaaatctg	gcctctttct	gcgactgaat	tctgcatctc	cttggggggg	9900
tgaggctgct	cccaaaaatt	ctttctccac	cgggcttagg	attccctggg	cttcaactct	9960
gagcttggac	tgcttggtc	aggagcctct	gcaagaaaca	aagcccagcc	aaacagggtcc	10020
ctccccctaa	aaaggaacct	gaaggtaatt	acctctccct	cagggtgtgg	gaatttccaa	10080
gtctgggaat	tcctatccag	ctgggggaagt	ctgcagtgca	ggtgagactt	ccggctgaaa	10140
gagccaggga	gcggccagat	gctcaggtac	ctgaaccaga	gccaagggac	ttccagacag	10200
tgaggcaact	gggctccaaa	taacctgatc	cggggaattc			10240
<210> SEQ ID NO 12						
<211> LENGTH: 1644						
<212> TYPE: DNA						
<213> ORGANISM: Mus sp.						
<400> SEQUENCE: 12						
cctcagcgag	gacagcaagg	gactagccag	gagggagaac	agaaactcca	gaacatcctg	60
gaaatagctc	ccagaaaagc	aagcagccaa	ccaggcaggt	tctgtccctt	tactcactg	120
gccaaggcgg	ccacatctcc	ctccagaaaa	gacaccatga	gcacagaaag	catgatccgc	180

-continued

gacgtggaac tggcagaaga ggcactcccc caaaagatgg ggggcttcca gaactccagg	240
cgggtgcctat gtctcagcct cttctcattc ctgcttgtgg cagggggccac cacgctcttc	300
tgtctactga acttcggggg gatcgggtccc caaagggatg agaagttccc aaatggcctc	360
cctctcatca gttctatggc ccagaccctc aactcagat catcttctca aaattcgagt	420
gacaagcctg tagcccacgt cgtagcaaac caccaagtgg aggagcagct ggagtggctg	480
agccagcgcg ccaacgccct cctggccaac ggcatggatc tcaaagacaa ccaactagt	540
gtgccagccg atgggttgta ccttgtctac tcccaggttc tcttcaaggg acaaggctgc	600
cccgactacg tgctcctcac ccacaccgtc agccgatttg ctatctcata ccaggagaaa	660
gtcaacctcc tctctgccgt caagagcccc tgccccaagg acaccctga gggggctgag	720
ctcaaaccct ggtatgagcc catatacctg ggaggagtct tccagctgga gaagggggac	780
caactcagcg ctgaggtcaa tctgccaag tacttagact ttgcggagtc cgggcaggtc	840
tactttggag tcattgctct gtgaaggga tgggtgttca tccattctct acccagcccc	900
cactctgacc cctttactct gacccttta ttgtctactc ctgagagccc ccagtctgtg	960
tccttctaac ttagaaaggg gattatggct cagagtccaa ctctgtgctc agagctttca	1020
acaactactc agaaacacaa gatgctggga cagtgcctg gactgtgggc ctctcatgca	1080
ccaccatcaa ggactcaaat gggctttccg aattcactgg agcctcgaat gtccattcct	1140
gagttctgca aagggagagt ggtcaggttg cctctgtctc agaatgaggc tggataagat	1200
ctcaggcctt cctaccttca gacctttcca gactcttccc tgaggtgcaa tgcacagcct	1260
tcctcacaga gccagcccc ctctatttat atttgcactt attatttatt atttatttat	1320
tatttattta tttgcttatg aatgtattta tttggaaggc cgggggtgtcc tggaggacct	1380
agtgtgggaa gctgtcttca gacagacatg tttctgtga aaacggagct gagctgtccc	1440
cacctggcct ctctaccttg ttgcctctc ttttgcctat gtttaaaaca aaatatttat	1500
ctaaccat atgtcttaata acgtgattt ggtgaccagg ctgtcgctac atcactgaac	1560
ctctgtctcc cacgggagcc gtgactgtaa ttgccctaca gtcaattgag agaaataaag	1620
atcgcttaaa ataaaaaacc cccc	1644
<210> SEQ ID NO 13	
<211> LENGTH: 1890	
<212> TYPE: DNA	
<213> ORGANISM: Homo sapiens	
<400> SEQUENCE: 13	
aaacagagag agatagagaa agagaaagac agaggtgttt cccttagcta tggaaactct	60
ataagagaga tccagcttgc ctctcttga gcagtcagca acagggctcc gtccttgaca	120
cctcagcctc tacaggactg agaagaagta aaaccgtttg ctggggctgg cctgactcac	180
cagctgccat gcagcagccc ttcaattacc catatcccca gatctactgg gtggacagca	240
gtgccagctc tccttgggcc cctccaggca cagttcttcc ctgtccaacc tctgtgcca	300
gaaggcctgg tcaaaggagg ccaccaccac caccgccacc gccaccacta ccacctccgc	360
cgccgcgcgc accactgcct ccactaccgc tgccaccct gaagaagaga gggaaccaca	420
gcacaggcct gtgtctctt gtgatgtttt tcatggttct ggttgcttg gtaggattgg	480
gcctggggat gtttcagctc ttccacctac agaaggagct ggcagaactc cgagagtcta	540
ccagccagat gcacacagca tcattcttgg agaagcaaat aggccacccc agtccacccc	600
ctgaaaaaaa ggagctgagg aaagtggccc atttaacagg caagtccaac tcaagggtcca	660

-continued

tgctcttgga atgggaagac acctatggaa ttgtcctgct ttctggagtg aagtataaga	720
agggtggcct tgtgatcaat gaaactgggc tgtactttgt atattccaaa gtatacttcc	780
ggggtcaatc ttgcaacaac ctgcccctga gccacaaggt ctacatgagg aactctaagt	840
atccccagga tctggatgat atggagggga agatgatgag ctactgcact actgggcaga	900
tgtgggcccg cagcagctac ctgggggcag tgttcaatct taccagtgct gatcatttat	960
atgtcaacgt atctgagctc tctctgggtca attttgagga atctcagacg tttttcggct	1020
tatataagct ctaagagaag cactttggga ttctttccat tatgattctt tggtacaggc	1080
accgagaatg ttgtattcag tgaggggtctt cttacatgca ttgaggtca agtaagaaga	1140
catgaaccaa gtggaccttg agaccacagg gttcaaaatg tctgtagctc ctcaactcac	1200
ctaattgttta tgagccagac aaatggagga atatgacgga agaacataga actctgggct	1260
gccatgtgaa gagggagaag catgaaaaag cagctaccca ggtgttctac actcatctta	1320
gtgcctgaga gtatttaggc agattgaaaa ggacaccttt taactcacct ctcaagggtg	1380
gccttgctac ctcaaggggg actgtctttc agatacatgg ttgtgacctg aggatttaag	1440
ggatggaaaa ggaagactag aggcttgcat aataagctaa agaggctgaa agaggccaat	1500
gccccactgg cagcatcttc acttctaaat gcatatcctg agccatcggg gaaactaaca	1560
gataagcaag agagatgttt tggggactca tttcattcct aacacagcat gtgtatttcc	1620
agtgccaatt gtaggggtgt gtgtgtgtgt gtgtgtgtgt gtgtatgact aaagagagaa	1680
tgtagatatt gtgaagtaca tattaggaaa atatgggttg catttggtca agattttgaa	1740
tgttctctga caatcaactc taatagtgtc taaaaatcat tgattgtcag ctactaatga	1800
tgttttctta taatataata aatatttatg tagatgtgca tttttgtgaa atgaaaacat	1860
gtaataaaaa gtatatgtta ggatacaaat	1890
<210> SEQ ID NO 14	
<211> LENGTH: 1541	
<212> TYPE: DNA	
<213> ORGANISM: Mus sp.	
<400> SEQUENCE: 14	
gggtgtctca cagagaagca aagagaagag aacaggagaa atggtgtttc ccttgactgc	60
ggaaacttta taaagaaaac ttagcttctc tggagcagtc agcgtcagag ttctgtcctt	120
gacacctgag tctcctccac aaggctgtga gaaggaaacc ctttctctggg gctgggtgcc	180
atgcagcagc ccatgaatta cccatgtccc cagatcttct gggtagacag cagtgccact	240
tcatcttggg ctctccagg gtcagttttt ccctgtccat cttgtgggcc tagagggccg	300
gaccaaagga gaccgccacc tccaccacca cctgtgtcac cactaccacc gccatcacia	360
ccactcccac tgccgccact gaccctcta aagaagaagg accacaacac aaatctgtgg	420
ctaccggtgg tatttttcat ggttctggtg gctctggttg gaatgggatt aggaatgtat	480
cagctcttcc acctgcagaa ggaactggca gaactccgtg agttcaccaa ccaaagcctt	540
aaagtatcat cttttgaaaa gcaaatagcc aaccccagta caccctctga aaaaaagag	600
ccgaggagtg tggcccattt aacagggaac cccactcaa ggtccatccc tctggaatgg	660
gaagacacat atggaaccgc tctgatctct ggagtgaagt ataagaaagg tggccttgtg	720
atcaacgaaa ctgggttgta cttcgtgtat tccaaagtat acttccgggg tcagtcttgc	780
aacaaccagc ccctaaacca caaggtctat atgaggaact ctaagtatcc tgaggatctg	840
gtgctaattg aggagaagag gttgaactac tgcactactg gccagatatg ggcccacagc	900

-continued

agctacctgg	gggcagtatt	caatcttacc	agtgctgacc	atztatatgt	caacatatct	960
caactctctc	tgatcaatth	tgaggaatct	aagaccttht	tcggcttgta	taagctthta	1020
aagaaaaagc	atthtaaaat	gatctactat	tctthtatcat	gggcaccagg	aatattgtct	1080
tgaatgagag	tcttcttaag	acctattgag	attaattaag	actacatgag	ccacaaagac	1140
ctcatgaccg	caagggtcaa	caggtcagct	atccttcatt	ttctcgaggt	ccatggagtg	1200
gtccttaatg	cctgcatcat	gagccagatg	gaaggagggtc	tgtgactgag	ggacataaag	1260
ctthgggctg	ctgtgtagca	atgcagaggc	acagagaaaag	aactgtctga	tgttaaatgg	1320
ccaagagaa	thtaaccatt	gaagaagaca	cctthtacct	cacttccagg	gtgggtctac	1380
ttactacctc	acagaggccg	ththtgagac	atagttgtgg	tatgaatata	caagggtgag	1440
aaaggaggct	catttgactg	ataagctaga	gactgaaaaa	aagacagtgt	ctcattggca	1500
ccatctthac	tgttacctga	tgtthtctga	gccgacctth	g		1541
<210> SEQ ID NO 15						
<211> LENGTH: 888						
<212> TYPE: DNA						
<213> ORGANISM: Homo sapiens						
<400> SEQUENCE: 15						
ggctgggtccc	ctgacaggth	gaagcaagta	gacgcccagg	agccccggga	gggggctgca	60
gtthcctthc	thcctthctg	gcagcgctcc	gcgcccccat	cgccctcct	gcgctagcgg	120
aggatgacgc	cgcggcgatg	cggaggagg	gttcgggctg	ctcggtgcg	cgcaggccct	180
atgggtgcgt	cctgcgggct	gctthggthc	cattggthgc	gggcttggtg	atctgcctcg	240
tggtgtgcat	ccagcgcttc	gcacaggctc	agcagcagct	gccgctcgag	tcacttgggt	300
gggacgtagc	tgagctgcag	ctgaatcaca	caggacctca	gcaggacccc	aggtatact	360
ggcagggggg	cccagcactg	ggccgctcct	tcctgcatgg	accagagctg	gacaaggggc	420
agctacgtat	ccatcgatg	ggcatctaca	tggtagacat	ccaggtagc	ctggccatct	480
gtcctccac	gacggcctcc	aggcaccacc	ccaccacct	ggcgtggga	atctgctctc	540
ccgcctccc	tagcatcagc	ctgctgcgtc	tcagctthca	ccaaggthgt	accattgcct	600
cccagcgct	gacgcccctg	gcccgagggg	acacactctg	caccaacctc	actgggacac	660
ththgccttc	ccgaaacact	gatgagacct	tctthggagt	gcagtgggtg	cgccctgac	720
cactgctgct	gattagggth	ththaaatth	tattthattth	tattthaagth	caagagaaaa	780
agtgtacaca	cagggggccac	cgggggttg	ggtgggagtg	tggtgggggg	tagtgggtggc	840
aggacaagag	aaggcattga	gctthtctth	tcattthcct	attaaaaa		888
<210> SEQ ID NO 16						
<211> LENGTH: 1906						
<212> TYPE: DNA						
<213> ORGANISM: Homo sapiens						
<400> SEQUENCE: 16						
ccaagtcaca	tgattcagga	ttcaggggga	gaatcctthc	tggaacagag	atgggcccag	60
aactgaatca	gatgaagaga	gataaggtht	gatgtgggga	agactatata	aagaatggac	120
ccagggtgc	agcaagcact	caacggaatg	gcccctcctg	gagacacagc	catgcatgtg	180
ccggcgggct	ccgtggccag	ccacctgggg	accacgagcc	gcagctattth	ctatttgacc	240
acagccactc	tggtctgtg	ccttgtcttc	acggtggcca	ctattatggt	gttggtcgth	300
cagaggacgg	actccattcc	caactcacct	gacaacgtcc	ccctcaaagg	aggaaattgc	360

-continued

tcagaagacc	tcttatgtat	cctgaaaaga	gtccattca	agaagtcag	ggcctacctc	420
caagtggcaa	agcatctaaa	caaaaccaag	ttgtcttgga	acaaagatgg	cattctccat	480
ggagtcagat	atcaggatgg	gaatctgggtg	atccaattcc	ctggtttgta	cttcatcatt	540
tgccaactgc	agtttcttgt	acaatgccc	aataattctg	tcgatctgaa	gttggagctt	600
ctcatcaaca	agcatatcaa	aaaacaggcc	ctggtgacag	tgtgtgagtc	tggaatgcaa	660
acgaaacacg	tataccagaa	tctctctcaa	ttcttgctgg	attacctgca	ggtcaacacc	720
accatatcag	tcaatgtgga	tacattccag	tacatagata	caagcacctt	tcctcttgag	780
aatgtgttgt	ccatcttctt	atacagtaat	tcagactgaa	cagtttctct	tggccttcag	840
gaagaaagcg	cctctctacc	atacagtatt	tcacccctcc	aaacacttgg	gcaaaaagaa	900
aactttagac	caagacaaac	tacacagggt	attaaatagt	atacttctcc	ttctgtctct	960
tggaaagata	cagctccagg	gttaaaaaga	gagtttttag	tgaagtatct	ttcagatagc	1020
aggcagggaa	gcaatgtagt	gtgggtgggca	gagccccaca	cagaatcaga	agggatgaat	1080
ggatgtccca	gcccaccac	taattcactg	tatggctctg	atctatttct	tctgttttga	1140
gagcctccag	ttaaatggg	gcttcagtac	cagagcagct	agcaactctg	ccctaattggg	1200
aaatgaaggg	gagctgggtg	tgagtgttta	cactgtgccc	ttcacgggat	acttctttta	1260
tctgcagatg	gcctaattgct	tagttgtcca	agtcgcatc	aaggactctc	tcacacagga	1320
aacttccta	tactggcaga	tacacttggt	actgaacat	gcccagttta	tgcctgtctg	1380
actgtcactc	tggcactagg	aggtgatct	tgtactccat	atgacccac	ccctaggaac	1440
ccccagggaa	aaccaggctc	ggacagcccc	ctgttctga	gatggaaagc	acaaatttaa	1500
tacaccacca	caatggaaaa	caagttcaaa	gacttttact	tacagatcct	ggacagaaag	1560
ggcataatga	gtctgaaggg	cagtcctcct	tctccagggt	acatgaggca	ggaataagaa	1620
gtcagacaga	gacagcaaga	cagttaacaa	cgtaggtaaa	gaaatagggt	gtggtcactc	1680
tcaattcact	ggcaaatgcc	tgaatgggtc	gtctgaagga	agcaacagag	aagtggggaa	1740
tccagtctgc	taggcaggaa	agatgcctct	aagtcttgt	ctctggccag	aggtgtggta	1800
tagaaccaga	aaccatatac	aagggtgact	aagcccggt	tccggtatga	gaaattaaac	1860
ttgtatacaa	aatggttgcc	aaggcaacat	aaaattataa	gaattc		1906
<210> SEQ ID NO 17						
<211> LENGTH: 1619						
<212> TYPE: DNA						
<213> ORGANISM: Homo sapiens						
<400> SEQUENCE: 17						
gtcatggaat	acgcctctga	cgttccactg	gacccgaag	ccccgtggcc	tcccgcccc	60
cgcgtctcg	cctgccggt	actgccttg	gccctggctg	cggtgctgct	gtgctgctg	120
ctgctcgctg	ccgcctgct	cgtcttctc	gcctgcccct	gggcccgtgc	cggtgctcgc	180
gcctcgcccg	gctccgccc	cagcccaga	ctccgcagg	gtcccgagct	ttcgccgac	240
gatcccgccg	gcctcttga	cctgcggcag	ggcatgttg	cgcagctgg	ggccaaaat	300
gttctgctga	tcgatggg	cctgagctgg	tacagtgac	caggcctgg	aggcgtgtcc	360
ctgacggggg	gcctgagcta	caaagaggac	acgaaggag	tggtgggtgg	caaggctgga	420
gtctactatg	tcttcttca	actagagctg	cgccgctgg	tgccggcga	gggctcaggc	480
tccgtttcac	ttgcgtgca	cctgcagcca	ctgcgtctg	ctgctggggc	cgcgcctg	540
gctttgaccg	tggacctgcc	acccgcctcc	tccgaggtc	ggaactcgg	cttcggtttc	600

-continued

cagggcgctg	tgctgcacct	gagtgcgggc	cagcgctgg	gcgtccatct	tcacactgag	660
gccagggcac	gccatgcctg	gcagcttacc	cagggcgcca	cagtcttggg	actcttccgg	720
gtgacccccg	aaatcccagc	cggactccct	tcaccgaggt	cggaataacg	cccagcctgg	780
gtgcagccca	cctggacaga	gtccgaatcc	tactccatcc	ttcatggaga	cccctgggtgc	840
tgggtccctg	ctgctttctc	tacctcaagg	ggcttggcag	gggtccctgc	tgttgacctc	900
cccttgagga	ccctcctcac	ccactccttc	cccaagttgg	accttgatat	ttattctgag	960
cctgagctca	gataatatat	tatatatatt	atatatatat	atataattct	atttaaagag	1020
gatcctgagt	ttgtgaatgg	acttttttag	aggagttggt	ttgggggggg	ggctcttcgac	1080
attgccgagg	ctggtcttga	actcctggac	ttagacgata	ctcctgcctc	agcctcccaa	1140
gcaactggga	ttcatccttt	ctattaattc	attgtactta	tttgccctatt	tgtgtgtatt	1200
gagcatctgt	aatgtgccag	cattgtgccc	aggctagggg	gctatagaaa	catctagaaa	1260
tagactgaaa	gaaaatctga	gttatggtaa	tacgtgagga	atttaaagac	tcacccccag	1320
cctccacctc	ctgtgtgata	cttgggggct	agcttttttc	tttctttctt	ttttttgaga	1380
tggctcttgt	ctgtcaacca	ggctagaatg	cagcggtgca	atcatgagtc	aatgcagcct	1440
ccagcctcga	cctcccgagg	ctcaggtgat	cctcccatct	cagcctctcg	agtagctggg	1500
accacagttg	tgtgccacca	cacttggtca	actttttaat	ttttttgcgg	agacgggtatt	1560
gctatgttgc	caaggttggt	tacatgccag	tacaatttat	aataaacact	catttttcc	1619
<210> SEQ ID NO 18						
<211> LENGTH: 1239						
<212> TYPE: DNA						
<213> ORGANISM: Mus sp.						
<400> SEQUENCE: 18						
agcctataaa	gcacgggcac	tggcgggaga	cgtgcactga	ccgaccgtgg	taatggacca	60
gcacacactt	gatgtggagg	ataccgcgga	tgcagacat	ccagcaggta	cttcgtgccc	120
ctcggatgcg	gcgctcctca	gagataccgg	gtcctcgcg	gacgctgcgc	tcctctcaga	180
tactgtgcgc	cccacaaatg	cgcgcctccc	cacggatgct	gcctaccctg	cggttaatgt	240
tcgggatcgc	gaggccgcgt	ggcgcctgc	actgaacttc	tgttcccgc	acccaaagct	300
ctatggccta	gtcgctttgg	ttttgctgct	tctgacgcc	gcctgtgttc	ctatcttcac	360
ccgcaccgag	cctcggccag	cgtcacaaat	caccacctcg	cccaacctgg	gtacccgaga	420
gaataatgca	gaccaggtca	cccctgtttc	ccacattggc	tgccccaaca	ctacacaaca	480
gggctctcct	gtgttcgcca	agctactggc	taaaaacca	gcctcgttgt	gcaatacaac	540
tctgaactgg	cacagccaag	atggagctgg	gagctcatac	ctatctcaag	gtctgaggta	600
cgaagaagac	aaaaaggagt	tgggtgtaga	cagtcccggg	ctctactacg	tatttttgga	660
actgaagctc	agtccaacat	tcacaaacac	aggccacaag	gtgcagggct	gggtctctct	720
tgttttgcaa	gcaaagcctc	aggtagatga	ctttgacaac	ttggccctga	cagtggaact	780
gttcccttgc	tccatggaga	acaagttagt	ggaccgttcc	tggagtcaac	tgttgctcct	840
gaaggctggc	caccgcctca	gtgtgggtct	gagggttat	ctgcatggag	cccaggatgc	900
atacagagac	tgggagctgt	cttatcccaa	caccaccagc	tttgactct	ttcttgtaaa	960
acccgacaac	ccatgggaat	gagaactatc	cttcttggtg	ctcctagttg	ctaagtcctc	1020
aagctgctat	gttttatggg	gtctgagcag	gggtcccttc	catgactttc	tcttgctttt	1080
aactggactt	ggtatttatt	ctgagcatag	ctcagacaag	actttatata	attcactaga	1140

-continued

tagcattagt aaactgctgg gcagctgcta gataaaaaaa aatttctaaa tcaaagttta	1200
tatttatatt aatatataaa aataaatgtg tttgtaaat	1239
<210> SEQ ID NO 19 <211> LENGTH: 606 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic chimeric construct	
<400> SEQUENCE: 19	
atgatcgaaa catacaacca aacttctccc cgatctgcgg ccactggact gcccacagc	60
atgaaaattt ttatgtattt acttactgtt tttcttatca cccagatgat tgggtcagca	120
ctttttgctg tgtatcgctt cgcacaggct tttgaaatgc aaaaagggtga tcagaatcct	180
caaattgcgg cacatgtcat aagtgaggcc agcagtaaaa caacatctgt gttacagtgg	240
gctgaaaaag gatactacac catgagcaac aacttggtaa ccctggaaaa tgggaaacag	300
ctgaccgtta aaagacaagg actctattat atctatgccc aagtcacctt ctgttccaat	360
cggggaagctt cgagtcaagc tccatttata gccagcctct gcctaaagtc ccccggtaga	420
ttcgagagaa tcttactcag agctgcaaat acccacagtt ccgccaaacc ttgcggggcaa	480
caatccattc acttgggagg agtat ttgaaaccag gtgcttcggt gtttgtcaat	540
gtgactgac caagccaagt gagccatggc actggcttca cgtcctttgg cttactcaaa	600
ctctga	606
<210> SEQ ID NO 20 <211> LENGTH: 783 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic chimeric construct	
<400> SEQUENCE: 20	
atgatcgaaa catacaacca aacttctccc cgatctgcgg ccactggact gcccacagc	60
atgaaaattt ttatgtattt acttactgtt tttcttatca cccagatgat tgggtcagca	120
ctttttgctg tgtatcttca tagaagattg gataaggctg aagaggaagt aaaccttcat	180
gaagattttg tattcataaa aaagctaaag agatgcaaca aaggagaagg atctttatcc	240
ttgctgaact gtgaggagat gagaaggcaa tttgaagacc ttgtcaagga tataacgtta	300
aacaaagaag agaaaaaaga aaacagcttt gaaatgcaaa aaggatgatca gaatcctcaa	360
attgcggcac atgtcataag tgaggccagc agtaaaacaa catctgtgtt acagtgggct	420
gaaaaaggat actacaccat gagcaacaac ttggtaacct tggaaaatgg gaaacagctg	480
accgttaaaa gacaaggact ctattatata tatgcccagg tcaccttctg ttccaatcgg	540
gaagcttcga gtcaagctcc atttatagcc agcctctgcc taaagtcccc cggtagattc	600
gagagaatct tactcagagc tgcaaatacc cacagttccg ccaaaccttg cgggcaacaa	660
tccattcact tgggaggagt atttgaattg caaccagggt cttcgggtgt tgtcaatgtg	720
actgatccaa gccaaagtgag ccatggcact ggcttcacgt cctttggctt actcaaaactc	780
tga	783
<210> SEQ ID NO 21 <211> LENGTH: 558 <212> TYPE: DNA <213> ORGANISM: Bos sp.	

-continued

<400> SEQUENCE: 21		
ctgctgcact tcggggtaat cggccccag agggaagagc agtccccagg tggccccctcc	60	
atcaacagcc ctctggttca aacactcagg tcctcttctc aagcctcaag taacaagccg	120	
gtagcccacg ttgtagccga catcaactct ccggggcagc tccggtggtg ggactcgtat	180	
gccaatgccc tcatggccaa cgggtgtgaag ctggaagaca accagctggt ggtgcctgct	240	
gacgggcttt acctcatcta ctcacaggtc ctcttcaggg gccaaggctg cccttccacc	300	
cccttgttcc tcaccacac catcagccgc attgcagtct cctaccagac caaggccaac	360	
atcctgtctg ccatcaagag cccttgccac agggagaccc cagagtgggc tgaggccaag	420	
ccctggtacg aaccatcta ccagggagga gtcttccagc tggagaaggg agatcgctc	480	
agtgtgaga tcaacctgcc ggactacctg gactatgccg agtccgggca ggtctacttt	540	
gggatcattg ccctgtga	558	
<210> SEQ ID NO 22		
<211> LENGTH: 1783		
<212> TYPE: DNA		
<213> ORGANISM: Mus sp.		
<400> SEQUENCE: 22		
caagtcacat gatccaggat gcaggggaaa atccttcttg gaacagagct gggtagagaa	60	
ccgaatcaga tgaggagaga taagggtgtga tgtgggacag actatataaa gcatggagcc	120	
agggtgcaa caagcaggca gctgtggggc tccttccct gaccagcca tgcaggtgca	180	
gcccggctcg gtagccagcc cctggagaag cagcaggccc tggagaagca caagtcgcag	240	
ctacttctac ctcagacca ccgcactggt gtgccttggt gtggcagtgg cgatcattct	300	
ggtactggta gtccagaaaa aggactccac tccaaataca actgagaagg ccccccttaa	360	
aggaggaaat tgctcagagg atctcttctg taccctgaaa agtactccat ccaagaagtc	420	
atgggcctac ctccaagtgt caaagcatct caacaatacc aaactgtcat ggaacgaaga	480	
tggcaccatc cacggactca tataccagga cgggaacctg atagtccaat tccttggtt	540	
gtacttcacg gtttgccaac tgcagttcct cgtgcagtgc tcaaatcatt ctgtggacct	600	
gacattgcag ctctcatca attccaagat caaaaagcag acgttggtaa cagtgtgtga	660	
gtctggagtt cagagtaaga acatctacca gaatctctct cagtttttgc tgcattactt	720	
acaggccaac tctaccatat cagtcagggt ggataatttc cagtatgtgg atacaaacac	780	
tttccctctt gataatgtgc tatccgtctt cttatatagt agctcagact gaatagttgt	840	
tcttaacctt tatgaaaatg ctgtctacca tacagtactt catctgtcca aacatgggcc	900	
aaagaaaata ttaggacaac tcaaactaag catgtgagtt agtgcacttc tctttctgtc	960	
ctttggaaaa atacaaaccc aggatttaga aagtggagtc tccttcagat gcacaaacag	1020	
gaaagaatgt gatatgtgca cagagacctc cttgggcact agaaggggtg tgagttgtcc	1080	
cagtataacc actaattcac tgaccttgag ccatttttcc tccccctgg aacttggggg	1140	
ctgaatctgg aaaagtagga gatgagattt acatttcccc aatattttct tcaactcaga	1200	
agacgagact gtggagctga gctccctaca cagatgaagg cctcccatgg catgaggaaa	1260	
atgatggtac cagtaatgtc tgtctgactg tcatctcagc aagtcctaag gacttccatg	1320	
ctgccttggt gaaagatact ctaacctctt gtaatgggca aagtgatcct gtctctcact	1380	
gaggggagta gctgctgcca tctcctgaga catacatgga gacattttct gccc aaattc	1440	
cattctgtgt gcagttttta agtattcccc caaaagttct tgacaatgag aactttgaat	1500	

-continued

gtgggaagag cttctggaca gcaaacatta acagctttctc ctgaccagag agaccatgca	1560
agcttggtct tagacccatc aagcttgagg tttctacatt gtgggagaca gacttttgac	1620
aaaccatttg agttgatgtc tgggcccttg ggagttctcc ttcagtaagg agagcaagcc	1680
gttctagtgc tgtgtcagag gatggagtaa aatagacact tttctgaagg aaaggagaac	1740
aaagtccag aaaaaggcta gaaaatgttt aaaaagaaaa aaa	1783
<210> SEQ ID NO 23	
<211> LENGTH: 1047	
<212> TYPE: DNA	
<213> ORGANISM: Homo sapiens	
<400> SEQUENCE: 23	
agagagcgct gggagccgga ggggagcgca gcgagttttg gccagtggtc gtgcagtcca	60
aggggctgga tggcatgctg gacccaagct cagctcagcg tccggacca ataacagttt	120
taccaaggga gcagctttct atcctggcca cactgaggty catagcgtaa tgtccatgtt	180
gttctacact ctgatcacag cttttctgat cggcatacag gcggaaccac actcagagag	240
caatgtccct gcaggacaca ccatcccca agtccactgg actaaacttc agcattccct	300
tgacactgcc cttcgcagag ccgcagcgc cccggcagcg gcgatatctg cacgcgtggc	360
ggggcagacc cgcaacatta ctgtggaccc caggctgttt aaaaagcggc gactccgttc	420
accccggtgtg ctgttttagca cccagcctcc ccgtaagct gcagacactc aggatctgga	480
cttcgaggtc ggtggtgctg ccccttcaa caggactcac aggagcaagc ggtcatcatc	540
ccatcccata ttcacaggg gcgaattctc ggtgtgtgac agtgtcagcg tgtgggttgg	600
ggataagacc accgccacag acatcaagg caaggaggty atggtgttgg gagaggtgaa	660
cattaacaac agtgtattca aacagtactt ttttgagacc aagtgccggg acccaaatcc	720
cgttgacagc gggtgccggg gcattgactc aaagcactgg aactcatatt gtaccacgac	780
tcacaccttt gtcaaggcgc tgaccatgga tggcaagcag gctgcctggc ggtttatccg	840
gatagatacg gcctgtgtgt gtgtgctcag caggaaggct gtgagaagag cctgacctgc	900
cgacacgctc cctccccctg ccccttctac actctcctgg gcccctccct acctcaacct	960
gtaaattatt ttaaattata aggactgcat ggtaatttat agtttataca gttttaaaga	1020
atcattattt attaaatfff tggaagc	1047
<210> SEQ ID NO 24	
<211> LENGTH: 1176	
<212> TYPE: DNA	
<213> ORGANISM: Mus sp.	
<400> SEQUENCE: 24	
gagcgctgg agccggagg gagcgcatcg agtgactttg gagctggcct tatatttgga	60
tctcccgggc agcttttttg aaactcctag tgaacatgct gtgcctcaag ccagtgaaat	120
taggctccct ggagggtgga cacgggcagc atggtggagt tttggcctgt ggtcgtgcag	180
tccagggggc tggatggcat gctggacca agctcacctc agtgtctggg cccaataaag	240
gttttgccaa ggacgcagct ttctatactg gccgcagtga ggtgcatagc gtaatgtcca	300
tgttgttcta cactctgatc actgcgtttt tgatcggcgt acaggcagaa ccgtacacag	360
atagcaatgt cccagaagga gactctgtcc ctgaagccca ctggactaaa cttcagcatt	420
cccttgacac agccctccgc agagcccgca gtgccctac tgcaccaata gctgcccgag	480
tgacagggca gaccgcaac atcactgtag accccagact gtttaagaaa cggagactcc	540

-continued

actcaccocg	tgtgctgttc	agcaccacgc	ctccaccac	ctcttcagac	actctggatc	600
tagactttcca	ggcccatggt	acaatccctt	tcaacaggac	tcaccggagc	aagcgctcat	660
ccaccacccc	agtcttcac	atgggggagt	tctcagtgtg	tgacagtgtc	agtgtgtggg	720
ttggagataa	gaccacagcc	acagacatca	agggaagga	ggtgacagtg	ctggccgagg	780
tgaacattaa	caacagtgtg	ttcagacagt	acttttttga	gaccaagtgc	cgagcctcca	840
atcctgttga	gagtgggtgc	cggggcatcg	actccaaaca	ctggaactca	tactgcacca	900
cgactcacac	cttcgtcaag	gcgttgacaa	cagatgagaa	gcaggctgcc	tggagggttca	960
tccggataga	cacagcctgt	gtgtgtgtgc	tcagcaggaa	ggctacaaga	agaggctgac	1020
ttgcctgcag	cccccttccc	cacctgcccc	ctccacactc	tcttggggccc	ctccctacct	1080
cagcctgtaa	attattttta	attataagga	ctgcatgata	atttatcggt	tatacaattt	1140
taaagacatt	atttattaaa	ttttcaaagc	atcctg			1176
<210> SEQ ID NO 25						
<211> LENGTH: 1623						
<212> TYPE: DNA						
<213> ORGANISM: Rattus sp.						
<400> SEQUENCE: 25						
tcagagtcc	gtccctgaca	cttcagtctc	cacaagactg	agaggaggaa	accctttcc	60
ggggctgggt	gccatgcagc	agcccgtaga	ttacccatgt	cccagatct	actgggtaga	120
cagcagtgcc	acttctcctt	gggtccctcc	agggtcagtt	ttttcttgtc	catactctgg	180
gcctagaggg	ccaggacaaa	ggagaccacc	gcctccacca	ccacctccat	caccactacc	240
accgccttcc	caaccacccc	cgtgcctcc	actaagccct	ctaaagaaga	aggacaacat	300
agagctgtgg	ctaccgggtga	tatttttcat	ggtgctgggtg	gctctgggtg	gaatgggggt	360
aggaatgtat	caactctttc	atctacagaa	ggaactggca	gaactccgtg	agttcaccaa	420
ccacagcctt	agagtatcat	cttttgaaaa	gcaaatagcc	aacccagca	cacctctga	480
aacaaaaag	ccaaggagtg	tggcccactt	aacagggaac	ccccgctcaa	ggtccatccc	540
tctggaatgg	gaagacacat	atggaactgc	ttgatctct	ggagtgaagt	ataagaaagg	600
cggccttgtg	atcaatgagg	ctgggttgta	cttcgtatat	tccaaagtat	acttccgggg	660
tcagtcttgc	aacagccagc	ccctaagcca	caagggtctat	atgaggaact	ttaagtatcc	720
tggggatctg	gtgctaattg	aggagaagaa	gttgaattac	tgcactactg	gccagatatg	780
ggcccacagc	agctacctag	gggcagtatt	taatcttacc	gttgctgacc	atttatatgt	840
caacatatct	caactctctc	tgatcaattt	tgaggaatct	aagacctttt	ttggcttata	900
taagctttta	aggaaaaagc	attttagaat	gatctattat	tctttatcat	ggatgccagg	960
aatattgtct	tcaatgagag	tcttcttaag	accaattgag	ccacaaagac	cacaagggtcc	1020
aacaggctcag	ctacccttca	ttttctagag	gtccatggag	tggtccttaa	tgctgtcatc	1080
atgagccaga	tgggaagaag	actgttcctg	aggaacataa	agttttgggc	tgctgtgtgg	1140
caatgcagag	gcaaagagaa	ggaactgtct	gatgttaa	ggccaagagc	attttagcca	1200
ttgaagaaaa	aaaaaacctt	taaactcacc	ttccagggtg	ggtctacttg	ctacctcaca	1260
ggaggccgct	ttttagacac	atggttgtgg	tatgactata	caagggtgag	aaaggatgct	1320
aggtttcatg	gataagctag	agactgaaaa	aagccagtgt	cccattggca	tcatctttat	1380
ttttaactga	tgttttctga	gcccaccttt	gatgctaaca	gagaaataag	aggggtgttt	1440
gaggcacaag	tcattctcta	catagcatgt	gtacctccag	tgcaatgatg	tctgtgtgtg	1500

-continued

tttttatgta tgagagtaga gcgattctaa agagtcacat gagtacaacg cgtacattac	1560
ggagtacata ttagaaacgt atgtgttaca tttgatgcta gaatatctga atgtttcttg	1620
cta	1623
<210> SEQ ID NO 26 <211> LENGTH: 28 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 26	
gttaagcttt tcagtcagca tgatagaa	28
<210> SEQ ID NO 27 <211> LENGTH: 27 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 27	
gtttctagat cagagtttga gtaagcc	27
<210> SEQ ID NO 28 <211> LENGTH: 30 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 28	
ccaagactag ttaacacagc atgatcgaaa	30
<210> SEQ ID NO 29 <211> LENGTH: 30 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 29	
ccaatgcggc cgcactcaga attcaacctg	30
<210> SEQ ID NO 30 <211> LENGTH: 972 <212> TYPE: DNA <213> ORGANISM: Homo sapiens	
<400> SEQUENCE: 30	
tctagactca ggactgagaa gaagtaaaac cgtttgctgg ggctggcctg actcaccagc	60
tgccatgcag cagcccttca attacccata tcccagatc tactgggttg acagcagtgc	120
cagctctccc tgggccctc caggcacagt tcttccctgt ccaacctctg tgcccagaag	180
gcctggtcaa aggaggccac caccaccacc gccaccgcca ccactaccac ctccgccgcc	240
gccgccacca ctgcctccac taccgctgcc acccctgaag aagagaggga accacagcac	300
aggcctgtgt ctcttgtga tgtttttcat ggttctggtt gccttggtag gattgggcct	360
ggggatgttt cagctcttcc acctacagaa ggagctggca gaactccgag agtctaccag	420
ccagatgcac acagcatcat ctttggagaa gcaaataggc caccacagtc caccacctga	480
aaaaaaggag ctgaggaaag tggcccatth aacaggcaag tccaactcaa ggtccatgcc	540

-continued

tctggaatgg gaagacacct atggaattgt cctgctttct ggagtgaagt ataagaaggg	600
tggccttggtg atcaatgaaa ctgggctgta ctttgtatat tccaaagtat acttccgggg	660
tcaatcttgc aacaacctgc ccctgagcca caaggtctac atgaggaact ctaagtatcc	720
ccaggatctg gtgatgatgg aggggaagat gatgagctac tgcactactg ggcagatgtg	780
ggcccgcagc agctacctgg gggcagtggt caatcttacc agtgctgac atttatatgt	840
caacgtatct gagctctctc tgggtcaattt tgaggaatct cagacgtttt tcggcttata	900
taagctctaa gagaagcact ttgggattct ttccattatg attctttgtt acaggcacccg	960
agatgttcta ga	972
<210> SEQ ID NO 31	
<211> LENGTH: 885	
<212> TYPE: DNA	
<213> ORGANISM: Mus sp.	
<400> SEQUENCE: 31	
atgcagcagc ccatgaatta cccatgtccc cagatcttct gggtagacag cagtgccact	60
tcatcttggg ctctccagg gtcagttttt ccctgtccat cttgtgggcc tagagggccg	120
gaccaaagga gaccgccacc tccaccacca cctgtgtcac cactaccacc gccatcacia	180
ccactcccac tgccgccact gacccctcta aagaagaagg accacaacac aaatctgtgg	240
ctaccggtgg tatttttcat ggttctggtg gctctggtg gaatgggatt aggaatgtat	300
cagctcttcc acctgcagaa ggaactggca gaactccgtg agttcaccaa ccaaagcctt	360
aaagtatcat cttttgaaaa gcaaatagcc aaccccagta caccctctga aaaaaagag	420
ccgaggagtg tggcccattht aacagggaac cccactcaa ggtccatccc tctggaatgg	480
gaagacacat atggaaccgc tctgatctct ggagtgaagt ataagaaagg tggccttggtg	540
atcaacgaag ctgggttgta cttcgtatat tccaaagtat acttccgggg tcagtcttgc	600
aacaaccagc ccctaaacca caaggtctat atgaggaact ctaagtatcc tggggatctg	660
gtgctaattg aggagaagag gttgaactac tgcactactg gacagatatg ggcccacagc	720
agctacctgg gggcagtatt caatcttacc agtgctgacc atttatatgt caacatatct	780
caactctctc tgatcaattt tgaggaatct aagacctttt tcggcttgta taagctttaa	840
aagaaaaagc attttaaaat gatctactat tctttatcat gggca	885
<210> SEQ ID NO 32	
<211> LENGTH: 29	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 32	
cttaagcttc tacaggactg agaagaagt	29
<210> SEQ ID NO 33	
<211> LENGTH: 30	
<212> TYPE: DNA	
<213> ORGANISM: Artificial Sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic primer	
<400> SEQUENCE: 33	
cttgaattcc aacattctcg gtgcctgtaa	30

-continued

<210> SEQ ID NO 34		
<211> LENGTH: 27		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic primer		
<400> SEQUENCE: 34		
tcaggatcca caaggctgtg agaagga	27	
<210> SEQ ID NO 35		
<211> LENGTH: 26		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic primer		
<400> SEQUENCE: 35		
cttgtctaga cctggtgccc atgata	26	
<210> SEQ ID NO 36		
<211> LENGTH: 680		
<212> TYPE: DNA		
<213> ORGANISM: Mus sp.		
<400> SEQUENCE: 36		
atgccggagg aaggtcgccc ttgccctcgg gttcgctgga gcgggaccgc gttccagcgc	60	
caatggccat ggctgctgct ggtggtgttt attactgtgt tttgctgttg gtttcattgt	120	
agcggactac tcagtaagca gcaacagagg ctgctggagc accctgagcc gcacacagct	180	
gagttacagc tgaatctcac agttcctcgg aaggacccca cactgcgctg gggagcaggc	240	
ccagccttgg gaaggtcctt cacacacgga ccagagctgg aggagggccca tctgcgtatc	300	
catcaagatg gcctctacag gctgcatatc caggtgacac tggccaactg ctcttcccca	360	
ggcagcacc ctcagcacag ggccaccctg gctgtgggca tctgctcccc cgctgcgcac	420	
ggcatcagct tgctgcgtgg gcgctttgga caggactgta cagtggcatt acagcgctg	480	
acatacctgg tccacggaga tgtcctctgt accaacctca ccctgcctct gctgccgtcc	540	
cgcaacgctg atgagacctt ctttggagtt cagtggatat gcccttgacc acaactccag	600	
gatgacttgt gaatattttt tttcttttca agttctacgt atttataaat gtatatagta	660	
cacataaaaa aaaaaaaaaa	680	
<210> SEQ ID NO 37		
<211> LENGTH: 846		
<212> TYPE: DNA		
<213> ORGANISM: Artificial Sequence		
<220> FEATURE:		
<223> OTHER INFORMATION: Synthetic chimeric construct		
<400> SEQUENCE: 37		
atgcagcagc ctttcaatta cccatatccc cagatctact gggtggaagc cagtgccagc	60	
tctccctggg cccctccagg cacagttctt ccctgtccaa cctctgtgcc cagaaggcct	120	
ggtcaaagga ggccaccacc accaccgcca ccgccaccac taccacctcc gccgcgcgcg	180	
ccaccactgc ctccactacc gctgccaccc ctgaagaaga gagggaacca cagcacaggc	240	
ctgtgtctcc ttgtgatgtt tttcatggtt ctggttgcc tggtaggatt gggcctgggg	300	
atgtttcagc tcttccacct gcagaaggaa ctggcagaac tccgtgagtt caccaaccaa	360	
agccttaaag tatcatcttt tgaaaagcaa ataggccacc ccagtccacc ccctgaaaaa	420	
aaggagctga ggaaagtggc ccatttaaca ggcaagtcca actcaaggtc catgcctctg	480	

-continued

gaatgggaag acacctatgg aattgtcctg ctttctggag tgaagtataa gaaggggtggc	540
cttgtgatca atgaaactgg gctgtacttt gtatattcca aagtatactt ccgggggtcaa	600
tcttgcaaca acctgcccct gagccacaag gtctacatga ggaactctaa gtatccccag	660
gatctggtga tgatggaggg gaagatgatg agctactgca ctactgggca gatgtggggc	720
cgcagcagct acctgggggc agtgttcaat cttaccagtg ctgatcattt atatgtcaac	780
gtatctgagc tctctctggt caattttgag gaatctcaga cgtttttcgg cttatataag	840
ctctaa	846
 <210> SEQ ID NO 38 <211> LENGTH: 786 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic chimeric construct <400> SEQUENCE: 38	
atgcagcagc ccttcaatta cccatatccc cagatctact ggggtggacag cagtgccagc	60
tctccctggg cccctccagg cacagttctt ccctgtccaa cctctgtgcc cagaaggcct	120
ggtcaaagga ggccaccacc accaccgcca ccgccaccac taccacctcc gccgcgcgcg	180
ccaccactgc ctccactacc gctgccaccc ctgaagaaga gagggaacca cagcacaggc	240
ctgtgtctcc ttgtgatgtt tttcatgggt ctggttgccct tggtaggatt gggcctgggg	300
atgtttcagc tcttccgctt cgcacaggct ataggccacc ccagtccacc ccctgaaaaa	360
aaggagctga ggaaagtggc ccatttaaca ggcaagtcca actcaaggtc catgcctctg	420
gaatgggaag acacctatgg aattgtcctg ctttctggag tgaagtataa gaaggggtggc	480
cttgtgatca atgaaactgg gctgtacttt gtatattcca aagtatactt ccgggggtcaa	540
tcttgcaaca acctgcccct gagccacaag gtctacatga ggaactctaa gtatccccag	600
gatctggtga tgatggaggg gaagatgatg agctactgca ctactgggca gatgtggggc	660
cgcagcagct acctgggggc agtgttcaat cttaccagtg ctgatcattt atatgtcaac	720
gtatctgagc tctctctggt caattttgag gaatctcaga cgtttttcgg cttatataag	780
ctctaa	786
 <210> SEQ ID NO 39 <211> LENGTH: 864 <212> TYPE: DNA <213> ORGANISM: Artificial Sequence <220> FEATURE: <223> OTHER INFORMATION: Synthetic chimeric construct <400> SEQUENCE: 39	
atgcagcagc ccttcaatta cccatatccc cagatctact ggggtggacag cagtgccagc	60
tctccctggg cccctccagg cacagttctt ccctgtccaa cctctgtgcc cagaaggcct	120
ggtcaaagga ggccaccacc accaccgcca ccgccaccac taccacctcc gccgcgcgcg	180
ccaccactgc ctccactacc gctgccaccc ctgaagaaga gagggaacca cagcacaggc	240
ctgtgtctcc ttgtgatgtt tttcatgggt ctggttgccct tggtaggatt gggcctgggg	300
atgtttcagc tcttccaatc ctccatcctc ccctatgccg gaggaggggt cgggctgctc	360
ggtgcggcgc aggccttatg ggtgcgtcct gcggccatcc tcaatcctat aggccacccc	420
agtccacccc ctgaaaaaaaa ggagctgagg aaagtggccc atttaacagg caagtccaac	480
tcaagggtcca tgcctctgga atgggaagac acctatggaa ttgtcctgct ttctggagtg	540

-continued

aagtataaga aggggtggcct tgtgatcaat gaaactgggc tgtactttgt atattccaaa	600
gtatacttcc ggggtcaatc ttgcaacaac ctgcccctga gccacaaggt ctacatgagg	660
aactctaagt atccccagga tctgggtgatg atggagggga agatgatgag ctactgcact	720
actgggcaga tgtgggcccg cagcagctac ctgggggcag tgttcaatct taccagtgct	780
gatcatttat atgtcaacgt atctgagctc tctctgggtca attttgagga atctcagacg	840
tttttcggct tatataagct ctaa	864

<210> SEQ ID NO 40
 <211> LENGTH: 828
 <212> TYPE: DNA
 <213> ORGANISM: Artificial Sequence
 <220> FEATURE:
 <223> OTHER INFORMATION: Synthetic delta-FasL-pcDNA3 sequence

<400> SEQUENCE: 40

atgcagcagc ccttcaatta cccatatccc cagatctact ggggtggacag cagtgccagc	60
tctccctggg cccctccagg cacagttctt ccctgtccaa cctctgtgcc cagaaggcct	120
ggtcaaagga ggccaccacc accaccgcca ccgccaccac taccacctcc gccgcgcgcg	180
ccaccactgc ctccactacc gctgccaccc ctgaagaaga gagggaacca cagcacaggc	240
ctgtgtctcc ttgtgatgtt tttcatgggt ctggttgccct tggtaggatt gggcctgggg	300
atgtttcagc tcttccacct acagcgagag tctaccagcc agatgcacac agcatcatct	360
ttggagaagc aaataggcca cccaggtcca cccctgaaa aaaaggagct gaggaaagtg	420
gcccatttaa caggcaagtc caactcaagg tccatgcctc tggaatggga agacacctat	480
ggaattgtcc tgctttctgg agtgaagtat aagaaggggtg gccttgtgat caatgaaact	540
gggctgtact ttgtatatc caaagtatac ttccgggggc aatcttgcaa caacctgccc	600
ctgagccaca aggtctacat gaggaactct aagtatcccc aggatctggt gatgatggag	660
gggaagatga tgagctactg cactactggg cagatgtggg cccgcagcag ctacctgggg	720
gcagtgttca atcttaccag tgctgatcat ttatatgtca acgtatctga gctctctctg	780
gtcaattttg aggaatctca gacgtttttc ggcttatata agctctaa	828

<210> SEQ ID NO 41
 <211> LENGTH: 846
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 41

atggctatga tggaggtcca ggggggaccc agcctgggac agacctgcgt gctgatcgtg	60
atcttcacag tgctcctgca gtctctctgt gtggctgtaa cttacgtgta ctttaccaac	120
gagctgaagc agatgcagga caagtactcc aaaagtggca ttgcttgttt cttaaaagaa	180
gatgacagtt attgggaccc caatgacgaa gagagtatga acagcccctg ctggcaagtc	240
aagtggcaac tccgtcagct cgttagaaag atgattttga gaacctctga ggaaaccatt	300
tctacagttc aagaaaagca acaaaatatt tctcccctag tgagagaaag aggtcctcag	360
agagtagcag ctacataaac tgggaccaga ggaagaagca acacattgtc ttctccaaac	420
tccaagaatg aaaaggctct gggccgcaaa ataaactcct gggaatcatc aaggagtggg	480
cattcattcc tgagcaactt gcacttgagg aatggtgaac tggatcatca tgaaaaaggg	540
ttttactaca tctattccca aacatacttt cgatttcagg aggaaataaa agaaaacaca	600
aagaacgaca aacaaatggc ccaatatatt taaaaataca caagttatcc tgaccctata	660

-continued

ttgttgatga aaagtgctag aaatagttgt tggctctaaag atgcagaata tggactctat	720
tccatctatc aagggggaat atttgagctt aaggaaaatg acagaatttt tgtttctgta	780
acaaatgagc acttgataga catggaccat gaagccagtt ttttcggggc ctttttagtt	840
ggctaa	846
 <210> SEQ ID NO 42 <211> LENGTH: 876 <212> TYPE: DNA <213> ORGANISM: Mus sp. <400> SEQUENCE: 42	
atgccttcct caggggccct gaaggacctc agcttcagtc agcacttcag gatgatggtg	60
atttgcatag tgctcctgca ggtgctcctg caggctgtgt ctgtggctgt gacttacatg	120
tacttcacca acgagatgaa gcagctgcag gacaattact ccaaaattgg actagcttgc	180
ttctcaaaga cggatgagga tttctgggac tccactgatg gagagatctt gaacagacct	240
tgcttgacag ttaagaggca actgtatcag ctcatgaag aggtgacttt gagaaccttt	300
caggacacca tttctacagt tccagaaaag cagctaagta ctctccctt gccagaggt	360
ggaagacctc agaaagtggc agctcacatt actgggatca ctcggaaga caactcagct	420
ttaattccaa tctccaagga tggaaagacc ttaggccaga agattgaatc ctgggagtcc	480
tctcggaag ggcattcatt tctcaaccac gtgctcttta ggaatggaga gctggtcac	540
gagcaggagg gcctgtatta catctattcc caaacatact tccgatttca ggaagctgaa	600
gacgcttcca agatggtctc aaaggacaag gtgagaacca aacagctggt gcagtacac	660
tacaagtaca ccagctatcc ggatcccata gtgctcatga agagcgccag aaacagctgt	720
tggtcagag atgccgagta cggactgtac tccatctatc agggaggatt gttcgagcta	780
aaaaaaaaatg acaggatttt tgtttctgtg acaaatgaac atttgatgga cctggatcaa	840
gaagccagct tctttggagc ctttttaatt aactaa	876
 <210> SEQ ID NO 43 <211> LENGTH: 720 <212> TYPE: DNA <213> ORGANISM: Mus sp. <400> SEQUENCE: 43	
atggagccag ggctgcaaca agcaggcagc tgtggggctc cttccctga cccagccatg	60
cagggtgcagc cgggctcgtt agccagcccc tggagaagca cgaggccctg gagaagcaca	120
agtcgcagct acttctacct cagcaccacc gcactggtgt gccttgttgt ggcagtggcg	180
atcattctgg tactggtagt ccagaaaaag gactccactc caaatacaac tgagaaggcc	240
ccccttaaag gaggaattg ctcagaggat ctcttctgta ccctgaaaag tactccatcc	300
aagaagtcat gggcctacct ccaagtgtca aagcatctca acaataccaa actgtcatgg	360
aacgaagatg gcaccatcca cggactcata taccaggacg ggaacctgat agtccaattc	420
cctggcttgt acttcatcgt ttgccaaactg cagtccctcg tgcagtgtc aaatcattct	480
gtggacctga cattgcagct cctcatcaat tccaagatca aaaagcagac gttggtaaca	540
gtgtgtgagt ctggagttca gagtaagaac atctaccaga atctctctca gtttttgctg	600
cattacttac aggtcaactc taccatatca gtcagggtgg ataatttcca gtatgtggat	660
acaaacactt tccctcttga taatgtgcta tccgtcttct tatatagtag ctcagactga	720

-continued

<210> SEQ ID NO 44															
<211> LENGTH: 930															
<212> TYPE: DNA															
<213> ORGANISM: Mus sp.															
<400> SEQUENCE: 44															
atggaccagc	acacacttga	tgtggaggat	accgcggatg	ccagacatcc	agcagggtact										60
tcgtgccccct	cggatgcggc	gctcctcaga	gataccgggc	tcctcgcgga	cgtcgcgtc										120
ctctcagata	ctgtgcgccc	cacaaatgcc	gcgctcccca	cggatgctgc	ctaccctgcg										180
gttaatgttc	gggatcgga	ggccgcgtgg	ccgcctgcac	tgaacttctg	ttcccggcac										240
ccaaagctct	atggcctagt	cgttttggtt	ttgctgcttc	tgatcgccgc	ctgtgttctt										300
atcttcaccc	gcaccgagcc	tcggccagcg	ctcacatca	ccacctcgcc	caacctgggt										360
acccgagaga	ataatgcaga	ccaggtcacc	cctgtttccc	acattggctg	ccccaacact										420
acacaacagg	gctctcctgt	gttcgccaag	ctactggcta	aaaaccaagc	atcgttgtgc										480
aatacaactc	tgaactggca	cagccaagat	ggagctggga	gtcatacct	atctcaaggt										540
ctgaggtacg	aagaagacaa	aaaggagttg	gtggtagaca	gtcccgggct	ctactacgta										600
tttttggaac	tgaagctcag	tccaacattc	acaaacacag	gccacaaggt	gcagggtctg										660
gtctctcttg	ttttgcaagc	aaagcctcag	gtagatgact	ttgacaactt	ggccctgaca										720
gtggaactgt	tcccttgctc	catggagaac	aagttagtgg	accgttcctg	gagtcaactg										780
ttgctcctga	aggctggcca	ccgcctcagt	gtgggtctga	gggcttatct	gcatggagcc										840
caggatgcat	acagagactg	ggagctgtct	tatcccaaca	ccaccagctt	tggactcttt										900
cttgtgaaac	ccgacaaccc	atgggaatga												930	

<210> SEQ ID NO 45															
<211> LENGTH: 281															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 45															
Met	Gln	Gln	Pro	Phe	Asn	Tyr	Pro	Tyr	Pro	Gln	Ile	Tyr	Trp	Val	Asp
1				5					10					15	
Ser	Ser	Ala	Ser	Ser	Pro	Trp	Ala	Pro	Pro	Gly	Thr	Val	Leu	Pro	Cys
			20					25					30		
Pro	Thr	Ser	Val	Pro	Arg	Arg	Pro	Gly	Gln	Arg	Arg	Pro	Pro	Pro	Pro
		35					40					45			
Pro	Pro	Pro	Pro	Pro	Leu	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Leu	Pro
		50				55				60					
Pro	Leu	Pro	Leu	Pro	Pro	Leu	Lys	Lys	Arg	Gly	Asn	His	Ser	Thr	Gly
65					70				75					80	
Leu	Cys	Leu	Leu	Val	Met	Phe	Phe	Met	Val	Leu	Val	Ala	Leu	Val	Gly
				85					90					95	
Leu	Gly	Leu	Gly	Met	Phe	Gln	Leu	Phe	His	Leu	Gln	Lys	Glu	Leu	Ala
		100						105					110		
Glu	Leu	Arg	Glu	Ser	Thr	Ser	Gln	Met	His	Thr	Ala	Ser	Ser	Leu	Glu
		115					120					125			
Lys	Gln	Ile	Gly	His	Pro	Ser	Pro	Pro	Pro	Glu	Lys	Lys	Glu	Leu	Arg
		130				135					140				
Lys	Val	Ala	His	Leu	Thr	Gly	Lys	Ser	Asn	Ser	Arg	Ser	Met	Pro	Leu
145					150					155					160
Glu	Trp	Glu	Asp	Thr	Tyr	Gly	Ile	Val	Leu	Leu	Ser	Gly	Val	Lys	Tyr
				165					170					175	

-continued

Lys	Lys	Gly	Gly	Leu	Val	Ile	Asn	Glu	Thr	Gly	Leu	Tyr	Phe	Val	Tyr	
			180					185					190			
Ser	Lys	Val	Tyr	Phe	Arg	Gly	Gln	Ser	Cys	Asn	Asn	Leu	Pro	Leu	Ser	
		195					200					205				
His	Lys	Val	Tyr	Met	Arg	Asn	Ser	Lys	Tyr	Pro	Gln	Asp	Leu	Val	Met	
	210					215					220					
Met	Glu	Gly	Lys	Met	Met	Ser	Tyr	Cys	Thr	Thr	Gly	Gln	Met	Trp	Ala	
225					230					235					240	
Arg	Ser	Ser	Tyr	Leu	Gly	Ala	Val	Phe	Asn	Leu	Thr	Ser	Ala	Asp	His	
				245					250					255		
Leu	Tyr	Val	Asn	Val	Ser	Glu	Leu	Ser	Leu	Val	Asn	Phe	Glu	Glu	Ser	
			260					265					270			
Gln	Thr	Phe	Phe	Gly	Leu	Tyr	Lys	Leu								
		275					280									
<210> SEQ ID NO 46																
<211> LENGTH: 281																
<212> TYPE: PRT																
<213> ORGANISM: Artificial sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Synthetic chimeric construct																
<400> SEQUENCE: 46																
Met	Gln	Gln	Pro	Phe	Asn	Tyr	Pro	Tyr	Pro	Gln	Ile	Tyr	Trp	Val	Asp	
1				5					10					15		
Ser	Ser	Ala	Ser	Ser	Pro	Trp	Ala	Pro	Pro	Gly	Thr	Val	Leu	Pro	Cys	
			20					25					30			
Pro	Thr	Ser	Val	Pro	Arg	Arg	Pro	Gly	Gln	Arg	Arg	Pro	Pro	Pro	Pro	
		35					40					45				
Pro	Pro	Pro	Pro	Pro	Leu	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Leu	Pro	
	50					55				60						
Pro	Leu	Pro	Leu	Pro	Pro	Leu	Lys	Lys	Arg	Gly	Asn	His	Ser	Thr	Gly	
65					70				75						80	
Leu	Cys	Leu	Leu	Val	Met	Phe	Phe	Met	Val	Leu	Val	Ala	Leu	Val	Gly	
				85					90					95		
Leu	Gly	Leu	Gly	Met	Phe	Gln	Leu	Phe	His	Leu	Gln	Lys	Glu	Leu	Ala	
			100					105					110			
Glu	Leu	Arg	Glu	Phe	Thr	Asn	Gln	Ser	Leu	Lys	Val	Ser	Ser	Phe	Glu	
		115					120					125				
Lys	Gln	Ile	Gly	His	Pro	Ser	Pro	Pro	Pro	Glu	Lys	Lys	Glu	Leu	Arg	
	130					135					140					
Lys	Val	Ala	His	Leu	Thr	Gly	Lys	Ser	Asn	Ser	Arg	Ser	Met	Pro	Leu	
145					150					155					160	
Glu	Trp	Glu	Asp	Thr	Tyr	Gly	Ile	Val	Leu	Leu	Ser	Gly	Val	Lys	Tyr	
				165					170					175		
Lys	Lys	Gly	Gly	Leu	Val	Ile	Asn	Glu	Thr	Gly	Leu	Tyr	Phe	Val	Tyr	
			180					185					190			
Ser	Lys	Val	Tyr	Phe	Arg	Gly	Gln	Ser	Cys	Asn	Asn	Leu	Pro	Leu	Ser	
		195					200					205				
His	Lys	Val	Tyr	Met	Arg	Asn	Ser	Lys	Tyr	Pro	Gln	Asp	Leu	Val	Met	
	210					215					220					
Met	Glu	Gly	Lys	Met	Met	Ser	Tyr	Cys	Thr	Thr	Gly	Gln	Met	Trp	Ala	
225					230					235					240	
Arg	Ser	Ser	Tyr	Leu	Gly	Ala	Val	Phe	Asn	Leu	Thr	Ser	Ala	Asp	His	
				245					250					255		
Leu	Tyr	Val	Asn	Val	Ser	Glu	Leu	Ser	Leu	Val	Asn	Phe	Glu	Glu	Ser	

-continued

260					265					270					
Gln	Thr	Phe	Phe	Gly	Leu	Tyr	Lys	Leu							
		275					280								
<210> SEQ ID NO 47															
<211> LENGTH: 261															
<212> TYPE: PRT															
<213> ORGANISM: Artificial sequence															
<220> FEATURE:															
<223> OTHER INFORMATION: Synthetic chimeric construct															
<400> SEQUENCE: 47															
Met	Gln	Gln	Pro	Phe	Asn	Tyr	Pro	Tyr	Pro	Gln	Ile	Tyr	Trp	Val	Asp
1				5					10					15	
Ser	Ser	Ala	Ser	Ser	Pro	Trp	Ala	Pro	Pro	Gly	Thr	Val	Leu	Pro	Cys
			20					25					30		
Pro	Thr	Ser	Val	Pro	Arg	Arg	Pro	Gly	Gln	Arg	Arg	Pro	Pro	Pro	Pro
			35				40					45			
Pro	Pro	Pro	Pro	Pro	Leu	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Leu	Pro
			50			55					60				
Pro	Leu	Pro	Leu	Pro	Pro	Leu	Lys	Lys	Arg	Gly	Asn	His	Ser	Thr	Gly
65					70					75					80
Leu	Cys	Leu	Leu	Val	Met	Phe	Phe	Met	Val	Leu	Val	Ala	Leu	Val	Gly
				85					90					95	
Leu	Gly	Leu	Gly	Met	Phe	Gln	Leu	Phe	Arg	Phe	Ala	Gln	Ala	Ile	Gly
			100					105					110		
His	Pro	Ser	Pro	Pro	Pro	Glu	Lys	Lys	Glu	Leu	Arg	Lys	Val	Ala	His
			115				120					125			
Leu	Thr	Gly	Lys	Ser	Asn	Ser	Arg	Ser	Met	Pro	Leu	Glu	Trp	Glu	Asp
			130				135				140				
Thr	Tyr	Gly	Ile	Val	Leu	Leu	Ser	Gly	Val	Lys	Tyr	Lys	Lys	Gly	Gly
145					150					155					160
Leu	Val	Ile	Asn	Glu	Thr	Gly	Leu	Tyr	Phe	Val	Tyr	Ser	Lys	Val	Tyr
				165					170					175	
Phe	Arg	Gly	Gln	Ser	Cys	Asn	Asn	Leu	Pro	Leu	Ser	His	Lys	Val	Tyr
			180					185					190		
Met	Arg	Asn	Ser	Lys	Tyr	Pro	Gln	Asp	Leu	Val	Met	Met	Glu	Gly	Lys
			195				200					205			
Met	Met	Ser	Tyr	Cys	Thr	Thr	Gly	Gln	Met	Trp	Ala	Arg	Ser	Ser	Tyr
			210				215				220				
Leu	Gly	Ala	Val	Phe	Asn	Leu	Thr	Ser	Ala	Asp	His	Leu	Tyr	Val	Asn
225					230					235					240
Val	Ser	Glu	Leu	Ser	Leu	Val	Asn	Phe	Glu	Glu	Ser	Gln	Thr	Phe	Phe
				245					250					255	
Gly	Leu	Tyr	Lys	Leu											
			260												
<210> SEQ ID NO 48															
<211> LENGTH: 282															
<212> TYPE: PRT															
<213> ORGANISM: Homo sapiens															
<400> SEQUENCE: 48															
Met	Gln	Gln	Pro	Phe	Asn	Tyr	Pro	Tyr	Pro	Gln	Ile	Tyr	Trp	Val	Asp
1				5					10					15	
Ser	Ser	Ala	Ser	Ser	Pro	Trp	Ala	Pro	Pro	Gly	Thr	Val	Leu	Pro	Cys
			20					25					30		

-continued

Pro	Thr	Ser	Val	Pro	Arg	Arg	Pro	Gly	Gln	Arg	Arg	Pro	Pro	Pro	Pro	
		35					40					45				
Pro	Pro	Pro	Pro	Pro	Leu	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Leu	Pro	
		50				55				60						
Pro	Leu	Pro	Leu	Pro	Pro	Leu	Lys	Lys	Arg	Gly	Asn	His	Ser	Thr	Gly	
65					70					75					80	
Leu	Cys	Leu	Leu	Val	Met	Phe	Phe	Met	Val	Leu	Val	Ala	Leu	Val	Gly	
				85					90					95		
Leu	Gly	Leu	Gly	Met	Phe	Gln	Leu	Phe	His	Leu	Gln	Lys	Glu	Leu	Ala	
			100					105					110			
Glu	Leu	Arg	Glu	Ser	Thr	Ser	Gln	Met	His	Thr	Ala	Ser	Ser	Leu	Glu	
		115					120					125				
Lys	Gln	Gln	Ile	Gly	His	Pro	Ser	Pro	Pro	Pro	Glu	Lys	Lys	Glu	Leu	
		130				135					140					
Arg	Lys	Val	Ala	His	Leu	Thr	Gly	Lys	Ser	Asn	Ser	Arg	Ser	Met	Pro	
145					150					155					160	
Leu	Glu	Trp	Glu	Asp	Thr	Tyr	Gly	Ile	Val	Leu	Leu	Ser	Gly	Val	Lys	
				165					170					175		
Tyr	Lys	Lys	Gly	Gly	Leu	Val	Ile	Asn	Glu	Thr	Gly	Leu	Tyr	Phe	Val	
			180					185					190			
Tyr	Ser	Lys	Val	Tyr	Phe	Arg	Gly	Gln	Ser	Cys	Asn	Asn	Leu	Pro	Leu	
		195					200					205				
Ser	His	Lys	Val	Tyr	Met	Arg	Asn	Ser	Lys	Tyr	Pro	Gln	Asp	Leu	Val	
		210				215					220					
Met	Met	Glu	Gly	Lys	Met	Met	Ser	Tyr	Cys	Thr	Thr	Gly	Gln	Met	Trp	
225					230					235					240	
Ala	Arg	Ser	Ser	Tyr	Leu	Gly	Ala	Val	Phe	Asn	Leu	Thr	Ser	Ala	Asp	
				245					250					255		
His	Leu	Tyr	Val	Asn	Val	Ser	Glu	Leu	Ser	Leu	Val	Asn	Phe	Glu	Glu	
			260					265					270			
Ser	Gln	Thr	Phe	Phe	Gly	Leu	Tyr	Lys	Leu							
		275					280									
<210> SEQ ID NO 49																
<211> LENGTH: 276																
<212> TYPE: PRT																
<213> ORGANISM: Artificial sequence																
<220> FEATURE:																
<223> OTHER INFORMATION: Synthetic chimeric construct																
<400> SEQUENCE: 49																
Met	Gln	Gln	Pro	Phe	Asn	Tyr	Pro	Tyr	Pro	Gln	Ile	Tyr	Trp	Val	Asp	
1				5					10					15		
Ser	Ser	Ala	Ser	Ser	Pro	Trp	Ala	Pro	Pro	Gly	Thr	Val	Leu	Pro	Cys	
			20					25					30			
Pro	Thr	Ser	Val	Pro	Arg	Arg	Pro	Gly	Gln	Arg	Arg	Pro	Pro	Pro	Pro	
		35					40					45				
Pro	Pro	Pro	Pro	Pro	Leu	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Pro	Leu	Pro	
		50				55				60						
Pro	Leu	Pro	Leu	Pro	Pro	Leu	Lys	Lys	Arg	Gly	Asn	His	Ser	Thr	Gly	
65					70					75					80	
Leu	Cys	Leu	Leu	Val	Met	Phe	Phe	Met	Val	Leu	Val	Ala	Leu	Val	Gly	
				85					90					95		
Leu	Gly	Leu	Gly	Met	Phe	Gln	Leu	Phe	Met	Pro	Glu	Glu	Gly	Ser	Gly	
			100					105					110			
Cys	Ser	Val	Arg	Arg	Arg	Pro	Tyr	Gly	Cys	Val	Leu	Arg	Ile	Gly	His	

```

115          120          125
Pro Ser Pro Pro Pro Glu Lys Lys Glu Leu Arg Lys Val Ala His Leu
130          135          140

Thr Gly Lys Ser Asn Ser Arg Ser Met Pro Leu Glu Trp Glu Asp Thr
145          150          155          160

Tyr Gly Ile Val Leu Leu Ser Gly Val Lys Tyr Lys Lys Gly Gly Leu
165          170          175

Val Ile Asn Glu Thr Gly Leu Tyr Phe Val Tyr Ser Lys Val Tyr Phe
180          185          190

Arg Gly Gln Ser Cys Asn Asn Leu Pro Leu Ser His Lys Val Tyr Met
195          200          205

Arg Asn Ser Lys Tyr Pro Gln Asp Leu Val Met Met Glu Gly Lys Met
210          215          220

Met Ser Tyr Cys Thr Thr Gly Gln Met Trp Ala Arg Ser Ser Tyr Leu
225          230          235          240

Gly Ala Val Phe Asn Leu Thr Ser Ala Asp His Leu Tyr Val Asn Val
245          250          255

Ser Glu Leu Ser Leu Val Asn Phe Glu Glu Ser Gln Thr Phe Phe Gly
260          265          270

Leu Tyr Lys Leu
275

<210> SEQ ID NO 50
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Met or Ala

<400> SEQUENCE: 50

Ala Pro Leu Gly Met Arg Xaa Arg
1          5

<210> SEQ ID NO 51
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Gly or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is Met or Tyr

<400> SEQUENCE: 51

Xaa Leu Xaa His Leu Leu Gly Lys
1          5

<210> SEQ ID NO 52
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)

```

-continued

<223> OTHER INFORMATION: Xaa is Val or Gly

<400> SEQUENCE: 52

Met Ala Xaa Glu Ile Phe Val Gln
1 5

<210> SEQ ID NO 53
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 53

Glu Asp Ile Tyr Gln Trp Ser Ile
1 5

<210> SEQ ID NO 54
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is Gln or Arg

<400> SEQUENCE: 54

Pro Ser Xaa Ala Pro Glu Glu Gly
1 5

<210> SEQ ID NO 55
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 55

Tyr Glu Asp Phe Phe Ala Phe Ser
1 5

<210> SEQ ID NO 56
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is Val or Gly

<400> SEQUENCE: 56

Ile Gly Glu Gln Ala Xaa Arg Glu
1 5

<210> SEQ ID NO 57
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Tyr or Val

<400> SEQUENCE: 57

-continued

Thr Arg Ala Asn Xaa Ser Pro Ala
1 5

<210> SEQ ID NO 58
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa can be any naturally occurring amino acid
except Lys, Glu, Trp
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Met or Ala

<400> SEQUENCE: 58

Arg Pro Leu Gly Xaa Asn Xaa Arg
1 5

<210> SEQ ID NO 59
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 59

Ala Pro Leu Glu Tyr Ala Gly Arg
1 5

<210> SEQ ID NO 60
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Gly or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Gly or His

<400> SEQUENCE: 60

Xaa Leu Gln Xaa Ile Leu Met Gln
1 5

<210> SEQ ID NO 61
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 61

Met Pro Leu Ala Leu Trp Ala Arg
1 5

<210> SEQ ID NO 62
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

-continued

<400> SEQUENCE: 62

Glu Pro Leu Glu Val Ala Gly Arg
1 5

<210> SEQ ID NO 63
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 63

Pro Pro Leu Glu Phe Ala Gly Arg
1 5

<210> SEQ ID NO 64
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Tyr, Ile, Thr or Arg

<400> SEQUENCE: 64

Xaa Pro Leu Ala Tyr Ala Gly Arg
1 5

<210> SEQ ID NO 65
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is Ala or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Gly or Ala

<400> SEQUENCE: 65

Gly Pro Arg Gly Leu Xaa Xaa Gln
1 5

<210> SEQ ID NO 66
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Ile or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is Thr or Phe

<400> SEQUENCE: 66

Ile Ala Gln Asn Xaa Xaa Leu Arg
1 5

<210> SEQ ID NO 67

-continued

<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Val or Met

<400> SEQUENCE: 67

Pro Arg Leu Ala Xaa Gly Ser His
1 5

<210> SEQ ID NO 68
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is Arg or Gln

<400> SEQUENCE: 68

Arg Pro Ala His Ala Xaa Pro Pro
1 5

<210> SEQ ID NO 69
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Gly or Ala

<400> SEQUENCE: 69

Leu Pro Lys Leu Glu His Xaa Gln
1 5

<210> SEQ ID NO 70
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Gln or Asn
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Gly or Ala

<400> SEQUENCE: 70

Gly Pro Ile Tyr Xaa Val Xaa Gln
1 5

<210> SEQ ID NO 71
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)

-continued

<223> OTHER INFORMATION: Xaa is Ala or Leu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Cly or Ala

<400> SEQUENCE: 71

Gly Pro His Gly Ser Xaa Xaa Gln
1 5

<210> SEQ ID NO 72
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 72

Pro Arg Gly Leu Glu Ala Thr
1 5

<210> SEQ ID NO 73
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Ile or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Ala, Leu or Phe

<400> SEQUENCE: 73

Pro Gln Gly Xaa Xaa Leu Thr
1 5

<210> SEQ ID NO 74
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Gln or Arg

<400> SEQUENCE: 74

Xaa Pro Arg Gly Leu Glu Ala Thr
1 5

<210> SEQ ID NO 75
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Val or Met
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Trp or Gly

<400> SEQUENCE: 75

-continued

Pro Leu Gly Xaa Xaa Ser Thr
1 5

<210> SEQ ID NO 76
<211> LENGTH: 7
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 76

Pro Arg Gly Ala Glu Gly Thr
1 5

<210> SEQ ID NO 77
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 77

Asp Pro Phe Glu Leu Arg Ala Thr
1 5

<210> SEQ ID NO 78
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is Leu or Met
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is Leu or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Arg or Met

<400> SEQUENCE: 78

Gly Ala Xaa Ala Phe Xaa Xaa Thr
1 5

<210> SEQ ID NO 79
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Gln or Arg

<400> SEQUENCE: 79

Xaa Pro Phe Glu Leu Arg Ala Thr
1 5

<210> SEQ ID NO 80
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE

-continued

<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Gln or Phe
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (5)..(5)
<223> OTHER INFORMATION: Xaa is Trp or Tyr

<400> SEQUENCE: 80

Leu Val Tyr Xaa Xaa Trp Gly Pro
1 5

<210> SEQ ID NO 81
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (3)..(3)
<223> OTHER INFORMATION: Xaa is Pro, Gly or Glu
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Val or Ile

<400> SEQUENCE: 81

Ile Leu Xaa Asn Ile Val Xaa Thr
1 5

<210> SEQ ID NO 82
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Glu or Val

<400> SEQUENCE: 82

Xaa Pro Phe Glu Leu Arg Ala Thr
1 5

<210> SEQ ID NO 83
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Ser or Asn

<400> SEQUENCE: 83

Leu Thr Ile His Val Gln Xaa Ala
1 5

<210> SEQ ID NO 84
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is His or Met
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE

-continued

<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Glu or Thr

<400> SEQUENCE: 84

Lys Phe Ala Gly Met Xaa Xaa Thr
1 5

<210> SEQ ID NO 85
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Gly or Asp

<400> SEQUENCE: 85

Xaa Pro Phe Glu Leu Arg Ala Thr
1 5

<210> SEQ ID NO 86
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Leu or Pro
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (6)..(6)
<223> OTHER INFORMATION: Xaa is Glu, Ser or Thr

<400> SEQUENCE: 86

Arg Arg Ser Xaa Glu Xaa Leu Thr
1 5

<210> SEQ ID NO 87
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Ser, Lys or Phe

<400> SEQUENCE: 87

Xaa Pro Phe Glu Leu Arg Ala Thr
1 5

<210> SEQ ID NO 88
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Pro or Met
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (2)..(2)
<223> OTHER INFORMATION: Xaa is Ser or Gly
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE

-continued

<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Lys, Tyr or Arg

<400> SEQUENCE: 88

Xaa Xaa Phe Xaa Leu Arg Ala Thr
1 5

<210> SEQ ID NO 89
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (1)..(1)
<223> OTHER INFORMATION: Xaa is Ala, Phe or Gln

<400> SEQUENCE: 89

Xaa Pro Phe Glu Leu Arg Ala Thr
1 5

<210> SEQ ID NO 90
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 90

Arg Ala Ile His Ile Gln Ala Glu
1 5

<210> SEQ ID NO 91
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 91

Gly Pro His Leu Leu Val Glu Ala
1 5

<210> SEQ ID NO 92
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct

<400> SEQUENCE: 92

Ile Pro Leu Glu Leu Arg Ala Gln
1 5

<210> SEQ ID NO 93
<211> LENGTH: 8
<212> TYPE: PRT
<213> ORGANISM: Artificial sequence
<220> FEATURE:
<223> OTHER INFORMATION: Synthetic construct
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (4)..(4)
<223> OTHER INFORMATION: Xaa is Met or Ala
<220> FEATURE:
<221> NAME/KEY: MISC_FEATURE
<222> LOCATION: (7)..(7)
<223> OTHER INFORMATION: Xaa is Val, Arg or Met

-continued

<400> SEQUENCE: 93	
Gly Leu Gln Xaa Ile Met Xaa Gln	
1 5	
<210> SEQ ID NO 94	
<211> LENGTH: 8	
<212> TYPE: PRT	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic construct	
<220> FEATURE:	
<221> NAME/KEY: MISC_FEATURE	
<222> LOCATION: (4)..(4)	
<223> OTHER INFORMATION: Xaa is Pro or Gln	
<400> SEQUENCE: 94	
Pro Pro Val Xaa Met Gln Gly Gln	
1 5	
<210> SEQ ID NO 95	
<211> LENGTH: 8	
<212> TYPE: PRT	
<213> ORGANISM: Artificial sequence	
<220> FEATURE:	
<223> OTHER INFORMATION: Synthetic construct	
<400> SEQUENCE: 95	
Ile Pro Leu Gly Leu Arg Ala Gln	
1 5	

- What is claimed is:
1. A chimeric nucleic acid molecule comprising a first polynucleotide encoding CD70 Domain III and a second polynucleotide encoding Fas ligand (FasL) Domain IV.
 2. The chimeric nucleic acid molecule according to claim 1, wherein the CD70 Domain III lacks a matrix metalloproteinase cleavage site, and the FasL Domain IV contains a binding site for CD95.
 3. The chimeric nucleic acid molecule according to claim 1, wherein the first polynucleotide further encodes Domain II from CD70 or FasL.
 4. The chimeric nucleic acid molecule according to claim 3, wherein the first polynucleotide further encodes Domain I from CD70 or FasL.
 5. An expression vector comprising the chimeric nucleic acid molecule according to claim 1.
 6. A genetic construct comprising the chimeric nucleic acid molecule according to claim 1.
 7. A host cell comprising the expression vector according to claim 5.
 8. A host cell comprising the genetic construct according to claim 6.
 9. A chimeric nucleic acid molecule comprising a first polynucleotide which encodes human Fas ligand (FasL) Domain III and a second polynucleotide which encodes non-human FasL Domain IV.
 10. The chimeric nucleic acid molecule according to claim 9, wherein the human FasL Domain III lacks a matrix metalloproteinase cleavage site, and the non-human FasL Domain IV contains a binding site for CD95.
 11. The chimeric nucleic acid molecule according to claim 7, wherein the first polynucleotide further encodes Domain II from human or non-human FasL.
 12. The chimeric nucleic acid molecule according to claim 7, wherein the first polynucleotide further encodes Domain I from human or non-human FasL.
 13. A genetic construct comprising the chimeric nucleic acid molecule according to claim 9.
 14. An expression vector comprising the chimeric nucleic acid molecule according to claim 9.
 15. A host cell comprising the genetic construct according to claim 13.
 16. A host cell comprising the expression vector according to claim 14.
 17. A chimeric nucleic acid molecule comprising a first polynucleotide which encodes non-human Fas ligand (FasL) Domain III and a second polynucleotide which encodes human FasL Domain IV.
 18. The chimeric nucleic acid molecule according to claim 17, wherein the non-human FasL Domain III lacks a matrix metalloproteinase cleavage site, and the human FasL Domain IV contains a binding site for CD95.
 19. The chimeric nucleic acid molecule according to claim 17, wherein the first polynucleotide further encodes Domain II from non-human or human FasL.
 20. The chimeric nucleic acid molecule according to claim 19, wherein the first polynucleotide further encodes Domain I from non-human or human FasL.
 21. A genetic construct comprising the chimeric nucleic acid molecule according to claim 17.
 22. An expression vector comprising the chimeric nucleic acid molecule according to claim 17.
 23. A host cell comprising the genetic construct according to claim 21.
 24. A host cell comprising the expression vector according to claim 22.

* * * * *