



US007989189B2

(12) **United States Patent**  
**Ebert et al.**

(10) **Patent No.:** **US 7,989,189 B2**  
(45) **Date of Patent:** **Aug. 2, 2011**

(54) **MUTANT GTP CYCLOHYDROLASE II ENZYMES**

(75) Inventors: **Sybille Ebert**, Biberach (DE);  
**Hans-Peter Hohmann**, Loerrach (DE);  
**Martin Lehmann**, Grenzach-Wyhlen (DE);  
**Nigel John Mouncey**, Binningen (CH);  
**Markus Wyss**, Liestal (CH)

(73) Assignee: **DSM IP Assets B.V.**, Te Heerlen (NL)

(\*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 143 days.

(21) Appl. No.: **11/631,542**

(22) PCT Filed: **Jul. 7, 2005**

(86) PCT No.: **PCT/EP2005/007320**  
§ 371 (c)(1),  
(2), (4) Date: **Nov. 13, 2007**

(87) PCT Pub. No.: **WO2006/003015**  
PCT Pub. Date: **Jan. 12, 2006**

(65) **Prior Publication Data**  
US 2008/0193998 A1 Aug. 14, 2008

(30) **Foreign Application Priority Data**  
Jul. 7, 2004 (EP) ..... 04015584

(51) **Int. Cl.**  
**C12P 17/00** (2006.01)  
**C12P 21/04** (2006.01)  
**C07H 21/04** (2006.01)  
**C12N 9/14** (2006.01)  
**C12N 15/00** (2006.01)  
**C12N 1/20** (2006.01)  
**C12Q 1/00** (2006.01)  
**C12Q 1/68** (2006.01)

(52) **U.S. Cl.** ..... **435/117**; 435/4; 435/6; 435/69.1;  
435/71.1; 435/252.3; 435/195; 435/440; 435/320.1;  
536/23.2

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,821,090 A 10/1998 Revuelta Doval et al.  
6,322,995 B1 11/2001 Hohmann et al.

OTHER PUBLICATIONS

Branden et al. Introduction to Protein Structure, Garland Publishing Inc., New York, p. 247, 1991.\*  
Gusarov et al. UniProtKB—Accession T50543—2000.\*  
Protein Purification Handbook—2001.\*  
Gusarov et al. Riboflavin biosynthetic genes in *Bacillus amyloliquefaciens*: primary structure, organization and regulation of activity. Mol Biol (Mosk). May-Jun. 1997;31(3):446-53.\*  
Richter et al., "Biosynthesis of Riboflavin: Cloning, Sequencing, Mapping, and Expression of the Gene Coding for GTP Cyclohydrolase II in *Escherichia coli*," J. Bact., vol. 175, No. 13, pp. 4045-4051, XP002095462 ISSN: 0021-9193 (1993).  
Herz et al., "Biosynthesis of riboflavin in plants. The ribA gene of *Arabidopsis thaliana* specifies a bifunctional GTP cyclohydrolase II/3,4-di-hydroxy-2-butanone 4-phosphate synthase," Phytochem., vol. 53, No. 7, pp. 723-731, XP004291367 ISSN: 0031-9422 (2000).  
Huembelin et al., "GTP cyclohydrolase II and 3,4-dihydroxy-2-butanone 4-phosphate synthase are rate-limiting enzymes in riboflavin synthesis of an industrial *Bacillus subtilis* strain used for riboflavin production," J. Ind. Microbiol. & Tech., vol. 22, No. 1, pp. 1-7, XP002348036 ISSN 1367-5435 (1999).  
Kaiser et al., "Biosynthesis of vitamin B2. An essential zinc ion at the catalytic site of GTP cyclohydrolase II," Eur. J. Biochem., vol. 269, No. 21, pp. 5264-5270, XP002348037 ISSN: 0014-2956 (2002).  
Bacher et al., "Biosynthesis of Riboflavin: GTP Cyclohydrolase II, Deaminase, And Reductase," Methods in Enzymology, vol. 280, pp. 382-389, XP009030498 ISSN: 0076-6879 (1997).  
Ritz et al., "Biosynthesis of riboflavin: Studies on the mechanism of GTP cyclohydrolase II," J. Biol. Chem., vol. 276, No. 25, pp. 22273-22277, XP002233121 ISSN: 0021-9258 (2001).

\* cited by examiner

*Primary Examiner* — Yong D Pak  
(74) *Attorney, Agent, or Firm* — Hoxie & Associates LLC

(57) **ABSTRACT**

The present invention relates to modified GTP cyclohydrolase II enzymes that display increased specific activity, and to polynucleotides encoding them. The invention further pertains to vectors comprising these polynucleotides and host cells containing such vectors. The invention provides a method for producing the modified enzyme and a method for producing riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof.

**4 Claims, No Drawings**



## MUTANT GTP CYCLOHYDROLASE II ENZYMES

This application is the National Stage of International Application No. PCT/EP2005/007320, filed Jul. 7, 2005.

The present invention provides modified enzymes with higher GTP cyclohydrolase II activity than the respective wild-type enzymes. The modified enzymes and polynucleotides encoding the same can be used for the production of riboflavin, riboflavin precursors, flavin mononucleotide (FMN), flavin adenine dinucleotide (FAD), and derivatives thereof.

Riboflavin (vitamin B2) is synthesized by all plants and many microorganisms but is not produced by higher animals. Because it is a precursor to coenzymes such as flavin adenine dinucleotide and flavin mononucleotide that are required in the enzymatic oxidation of carbohydrates, riboflavin is essential to basic metabolism. In higher animals, insufficient riboflavin can cause loss of hair, inflammation of the skin, vision deterioration, and growth failure.

Engineering of riboflavin production strains with increased rates and yields of riboflavin has been achieved in the past in a number of different ways. For instance, (1) classical mutagenesis was used to create variants with random mutations in the genome of the organism of choice, followed by selection for higher resistance to purine analogs and/or by screening for increased production of riboflavin. (2) Alternatively, the terminal enzymes of riboflavin biosynthesis, i.e., the enzymes catalyzing the conversion of guanosine triphosphate (GTP) and ribulose-5-phosphate to riboflavin, were overexpressed, resulting also in a higher flux towards the target product. However, in this latter approach, strong overexpression of the riboflavin biosynthesis proteins imposes an additional metabolic burden on the host cells which may, in turn, induce stress response reactions and other undesirable negative effects on the cells' physiology.

The enzymes required catalyzing the biosynthesis of riboflavin from guanosine triphosphate (GTP) and ribulose-5-phosphate are encoded by four genes (ribG, ribB, ribA, and ribH) in *B. subtilis*. These genes are located in an operon, the gene order of which differs from the order of the enzymatic reactions catalyzed by the enzymes. For example, GTP cyclohydrolase II, which catalyzes the first step in riboflavin biosynthesis, is encoded by the third gene in the operon, ribA. The ribA gene also encodes a second enzymatic activity, i.e., 3,4-dihydroxy-2-butanone 4-phosphate synthase (DHBPS), which catalyzes the conversion of ribulose-5-phosphate to the four-carbon unit 3,4-dihydroxy-2-butanone 4-phosphate (DHBP). Deaminase and reductase are encoded by the first gene of the operon, ribG. The penultimate step in riboflavin biosynthesis is catalyzed by lumazine synthase, the product of the last rib gene, ribH. Riboflavin synthase, which controls the last step of the pathway, is encoded by the second gene of the operon, ribB. The function of the gene located at the 3' end of the rib operon is, at present, unclear; however, its gene product is not required for riboflavin synthesis.

Transcription of the riboflavin operon from the ribP1 promoter is controlled by an attenuation mechanism involving a regulatory leader region located between ribP1 and ribG. The ribO mutations within this leader region result in deregulated expression of the riboflavin operon. Deregulated expression is also observed in strains containing missense mutations in the ribC gene. The ribC gene has been shown to encode the flavin kinase/FAD synthase of *B. subtilis* (Mack, M., et al., J. Bacteriol., 180:950-955, 1998). Deregulating mutations reduce the flavokinase activity of the ribC gene product

resulting in reduced intracellular concentrations of flavin mononucleotide (FMN), the effector molecule of the riboflavin regulatory system.

Recently, *Bacillus subtilis* was genetically engineered to produce high yields of riboflavin during a short fermentation cycle (U.S. Pat. No. 5,837,528). This approach combined classical genetic mutant selection and fermentation improvement with genetic engineering of the riboflavin biosynthetic genes by deregulating and increasing the level of gene expression. In this system, the expression of the rib genes was increased by mutating the flavokinase encoding ribC gene, by linking the rib genes to strong, constitutive promoters, and by increasing the copy number of the rib genes.

As already discussed above, overexpression of the rib genes poses an additional burden on the production strains which may, potentially, have a negative impact on the production of riboflavin precursors, riboflavin, FMN, FAD, or their derivatives. In order to circumvent this shortcoming, it is a subject of the present invention to describe GTP cyclohydrolase II mutants with increased specific activity. Use of such mutant enzymes in production strains, either alone or combined with improved mutants of the other Rib proteins, will allow higher flux rates with less or no additional burden on the cells' metabolism.

As used herein, the term "GTP cyclohydrolase II" may include any enzyme that is capable of catalyzing the conversion of GTP to 2,5-diamino-6-ribosylamino-4 (3H)-pyrimidinone-5'-phosphate (DRAPP). It is irrelevant whether this enzyme is capable of catalyzing further reactions, as for example the conversion of ribulose-5-phosphate to DHBP. A "GTP cyclohydrolase II" may be homologous to one or more of the enzymes the amino acid sequences of which are shown in Table 4. "Homologous" refers to a GTP cyclohydrolase II that is at least about 50% identical, preferably at least about 60% identical, more preferably at least about 70% identical, even more preferably at least about 80% identical, even more preferably at least about 85% identical, even more preferably at least about 90% or 95% identical, and most preferably at least about 98% identical to one or more of the amino acid sequences as shown in Table 4.

The term "% identity", as known in the art, means the degree of relatedness between polypeptide or polynucleotide sequences, as the case may be, as determined by the match between strings of such sequences. "Identity" can be readily determined by known methods, e.g., with the program BEST-FIT (GCG Wisconsin Package, version 10.2, Accelrys Inc., 9685 Scranton Road, San Diego, Calif. 92121-3752, USA) using the following parameters: gap creation penalty 8, gap extension penalty 2 (default parameters).

"Wild-type enzyme" or "wild-type GTP cyclohydrolase II" may include any GTP cyclohydrolase II homologous to any one of the enzymes shown in Table 4 that is used as starting point for designing mutants with increased activity according to the present invention. "Wild-type" in the context of the present invention may include both GTP cyclohydrolase II sequences derivable from nature as well as variants of synthetic GTP cyclohydrolase II enzymes (as long as they are homologous to any one of the sequences shown in Table 4), if they can be made more active by any of the teachings of the present invention. The terms "wild-type GTP cyclohydrolase II" and "non-modified GTP cyclohydrolase II" are used interchangeably herein.

A "mutant", "mutant enzyme", or "mutant GTP cyclohydrolase II" may include any variant derivable from a given wild-type enzyme/GTP cyclohydrolase II (according to the above definition) according to the teachings of the present invention and being more active than the respective wild-type



enzyme. For the scope of the present invention, it is not relevant how the mutant(s) are obtained; such mutants may be obtained, e.g., by site-directed mutagenesis, saturation mutagenesis, random mutagenesis/directed evolution, chemical or UV mutagenesis of entire cells/organisms, and other methods which are known in the art. These mutants may also be generated, e.g., by designing synthetic genes, and/or produced by in vitro (cell-free) translation. For testing of specific activity, mutants may be (over-) expressed by methods known to those skilled in the art. The terms “mutant GTP cyclohydrolase II” and “modified GTP cyclohydrolase II” are used interchangeably herein. This also applies to the terms “mutant enzyme” and “modified enzyme”.

“Riboflavin precursor” and “derivatives of riboflavin, FMN or FAD” in the context of this patent application may include any and all metabolite(s) requiring GTP cyclohydrolase II as an intermediate enzyme in their (bio-) synthesis. In the context of this patent application, it is irrelevant whether such (bio-) synthesis pathways are natural or non-natural (i.e., pathways not occurring in nature, but engineered biotechnologically). Preferably, the synthesis pathways are biochemical in nature. Riboflavin precursors and derivatives of riboflavin, FMN or FAD include but are not limited to: DRAPP; 5-amino-6-ribosylamino-2,4 (1H,3H)-pyrimidinedione-5'-phosphate; 2,5-diamino-6-ribitylamino-4 (3H)-pyrimidinedione-5'-phosphate; 5-amino-6-ribitylamino-2,4 (1H,3H)-pyrimidinedione-5'-phosphate; 5-amino-6-ribitylamino-2,4 (1H,3H)-pyrimidinedione; 6,7-dimethyl-8-ribityllumazine (DMRL); and flavoproteins. The term “riboflavin” also includes derivatives of riboflavin, such as e.g. riboflavin-5-phosphate and salts thereof, such as e.g. sodium riboflavin-5-phosphate.

It is in general an object of the present invention to provide an enzyme having GTP cyclohydrolase II activity, said enzyme being modified in a way that its catalytic properties are more favorable (i.e., showing higher specific activity) than those of the non-modified GTP cyclohydrolase II enzymes.

The invention relates to a modified GTP cyclohydrolase II which exhibits higher (specific) activity in comparison to the corresponding non-modified GTP cyclohydrolase II wherein

(i) the amino acid sequence of the modified GTP cyclohydrolase II contains at least one mutation when compared with the amino acid sequence of the corresponding non-modified GTP cyclohydrolase II, and

(ii) the at least one mutation is at one or more amino acid positions selected from the group consisting of amino acid positions corresponding to positions 261, 270, 276, 279, 308 and 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO:2.

Thus, it is an object of the present invention to provide a modified GTP cyclohydrolase II, wherein

(i) the specific activity of the modified enzyme is increased in comparison to the corresponding non-modified enzyme, and

(ii) the amino acid sequence of the modified enzyme comprises one or more mutation(s) including 1, 2, 3, 4, 5, or 6 mutation(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308 and/or 347 of SEQ ID NO:2.

The term “at least one mutation” means one or more mutation on a position as defined above leading to a modified GTP cyclohydrolase II having an increased specific activity compared to the non-modified enzyme. A modified enzyme as described above may consist of only 1, 2, 3, 4, 5 or 6 mutation(s) on a position as defined above leading to an increased specific activity compared to the non-modified enzyme, but may also include further amino acid mutations on other posi-

tions, as long as the resulting modified enzyme has an increased specific activity. Thus, the modified enzyme comprises one or more mutation(s) including 1, 2, 3, 4, 5, or 6 mutation(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308 and/or 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO:2. Examples of such mutations on positions other than the ones defined above are amino acid mutation(s) on a position corresponding to amino acid position 196, 282, and/or 325 of SEQ ID NO:2.

As used herein, the term “specific activity” denotes the reaction rate of the wild-type and mutant GTP cyclohydrolase II enzymes under properly defined reaction conditions as described e.g. in Ritz et al. (J. Biol. Chem. 276, 22273-22277, 2001), Koh et al. (Mol. Gen. Genet. 251, 591-598, 1996), or Schramek et al. (J. Biol. Chem. 276, 44157-44162, 2001) or as described in detail in Example 2. The “specific activity” defines the amount of substrate consumed and/or product produced in a given time period and per defined amount of protein at a defined temperature. Typically, “specific activity” is expressed in  $\mu\text{mol}$  substrate consumed or product formed per min per mg of protein. Typically,  $\mu\text{mol}/\text{min}$  is abbreviated by U (=unit). Therefore, the unit definitions for specific activity of  $\mu\text{mol}/\text{min}/(\text{mg of protein})$  or  $\text{U}/(\text{mg of protein})$  are used interchangeably throughout this document. It is understood that in the context of the present invention, specific activity must be compared on the basis of a similar, or preferably identical, length of the polypeptide chain. The invention shall not be circumvented by increasing the size of a given wild-type enzyme through, e.g., formation of a fusion protein, thereby reducing the apparent specific activity of the overall enzyme.

According to the present invention the modified GTP cyclohydrolase II exhibits a specific activity that is higher than that of the corresponding non-modified enzyme. Preferably, the specific activity of the modified GTP cyclohydrolase II of the invention is increased by at least about 5, 10, 25, 40, 60, 70, 80, 85, 90%, more preferably at least about 70% in comparison to the corresponding non-modified GTP cyclohydrolase II (for measurement of specific activity, see below). Preferably, increases in specific activity refer to the experimental conditions described in Example 1 of this application. Approx. 0.004-0.02 U/ml (corresponding to approx. 40  $\mu\text{g}/\text{ml}$  of *Bacillus subtilis* GTP cyclohydrolase II or 20  $\mu\text{g}/\text{ml}$  for the best mutants described here), preferably approx. 0.004 U/ml of GTP cyclohydrolase II activity, were present in the assay mixture, and the reaction was carried out at 37° C.

The amino acid sequence of the modified GTP cyclohydrolase II of the invention contains at least one mutation as defined above when compared with the amino acid sequence of the corresponding non-modified GTP cyclohydrolase II. Said mutation may be one or more addition, deletion and/or substitution, preferably one or more amino acid substitution wherein a given amino acid present in the amino acid sequence of the non-modified GTP cyclohydrolase II is replaced with a different amino acid in the amino acid sequence of the modified GTP cyclohydrolase II of the invention. The amino acid sequence of the modified GTP cyclohydrolase II may contain at least one amino acid substitution when compared with the amino acid sequence of the corresponding non-modified GTP cyclohydrolase II, i.e. may comprise one or more mutation(s) including 1, 2, 3, 4, 5, or 6 amino acid substitution(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308 and/or 347 of SEQ ID NO:2, preferably 2, 3, 4 or 5 amino acid substitutions. Thus, the modified enzyme preferably contains at least 2, at least 3, at least 4 or at least 5 substitutions when compared



## 5

with the amino acid sequence of the corresponding non-modified GTP cyclohydrolase II.

In one embodiment, a modified GTP cyclohydrolase II obtainable from *Bacillus*, preferably *Bacillus subtilis*, is provided, wherein

(i) the specific activity of the modified enzyme is increased in comparison to the corresponding non-modified enzyme, and

(ii) the amino acid sequence of the modified enzyme comprises one or more mutation(s) including 1, 2, 3, 4, 5, or 6 mutation(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308 and/or 347 of SEQ ID NO:2.

In one embodiment the non-modified enzyme corresponds to *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO:2. Thus, the modified enzyme having an increased specific activity in comparison to the wild type enzyme comprises one or more mutation(s) including 1, 2, 3, 4, 5, or 6 mutation(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308 and/or 347 of SEQ ID NO:2. In a further embodiment the modified enzyme having increased specific activity as defined above contains amino acid mutation(s) beside the amino acid positions as above, said further mutation(s) being on a position selected from the group consisting of position 196, 282, 235, and any combination thereof, preferably amino acid substitutions, more preferably the substitutions are Y196C (replacement of tyrosine by cysteine), A282T (replacement of alanine by threonine) or F325Y (replacement of phenylalanine by tyrosine).

A non-modified GTP cyclohydrolase II may be any GTP cyclohydrolase II for which increasing the specific activity is desirable. Non-modified GTP cyclohydrolase II enzymes include but are not limited to GTP cyclohydrolase II enzymes derivable from nature, such as enzymes of eukaryotic or prokaryotic origin, preferably fungal or bacterial origin. More preferably the non-modified enzyme is selected from the ones shown in Table 4 or which is homologous to any of the amino acid sequences as shown in Table 4, in particular selected from the group consisting of *Ashbya*, *Saccharomyces*, *Eremothecium*, *Candida*, *Neurospora*, *Schizosaccharomyces*, *Archeoglobus*, *Streptomyces*, *Helicobacter*, *Escherichia*, *Corynebacterium*, *Thermotoga*, *Arabidopsis*, *Lycopersicum*, *Oryza*, *Alcaligenes*, *Pseudomonas*, *Dinococcus*, *Lactobacillus*, *Photobacterium* and *Bacillus* and preferably selected from the group consisting of *Candida guilliermondii*, *Ashbya gossypii* (*Eremothecium ashbyii*) (SEQ ID NO:33), *Saccharomyces cerevisiae*, *Neurospora crassa*, *Schizosaccharomyces pombe*, *Archeoglobus fulgidus*, *Streptomyces coelicolor*, *Helicobacter pylori* J99, *Escherichia coli* (SEQ ID NO:35), *Corynebacterium glutamicum* (SEQ ID NO:37), *Bacillus amyloliquefaciens* (SEQ ID NO:39), *Bacillus cereus* (SEQ ID NO:41), *Bacillus halodurans* (SEQ ID NO:43), *Thermotoga maritima*, *Arabidopsis thaliana*, *Lycopersicum esculentum*, *Oryza sativum*, *Alcaligenes eutrophus*, *Pseudomonas putida* strain KT2440, *Corynebacterium efficiens*, *Deinococcus radiodurans*, *Lactobacillus plantarum*, *Photobacterium phosphoreum*, *Pseudomonas putida* strain KT2440 (second gene) and *Bacillus subtilis* (SEQ ID NO:2). Most preferably the non-modified enzyme is obtainable from *Bacillus subtilis*.

The modified GTP cyclohydrolase II of the invention may be obtained by mutating the corresponding non-modified GTP cyclohydrolase II. In one embodiment, the non-modified enzyme corresponds to the *B. subtilis* GTP cyclohydrolase II shown in SEQ ID NO:2 and the modified enzyme comprises one or more amino acid mutation(s) including 1, 2, 3, 4, 5, or 6 mutation(s) on amino acid position(s) 261, 270, 276, 279, 308 and/or 347 of SEQ ID NO:2, wherein the

## 6

specific activity of said modified enzyme is increased compared to the non-modified enzyme.

Preferably, the at least one mutation is at one or more amino acid positions selected from the group consisting of amino acid positions corresponding to positions 261, 279, 308 and 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO: 2. Thus, in one embodiment the modified GTP cyclohydrolase II comprises one or more mutation(s) including 1, 2, 3 or 4 mutation(s) on amino acid position(s) corresponding to positions 261, 279, 308, and/or 347 of SEQ ID NO:2. In a preferred embodiment, the modified enzyme is obtainable from *B. subtilis* and comprises mutated amino acid positions 261, 279, 308, and/or 347 as shown in SEQ ID NO:2, corresponding to amino acids V261, Q279, K308, and M374, respectively.

In another preferred embodiment the at least one mutation is at one or more amino acid positions selected from the group consisting of amino acid positions corresponding to positions 270, 279, 308 and 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO: 2. Thus, in one embodiment the modified GTP cyclohydrolase II comprises one or more mutation(s) including 1, 2, 3 or 4 mutation(s) on amino acid position(s) corresponding to positions 270, 279, 308, and/or 347 of SEQ ID NO:2. Preferably, the modified enzyme is obtainable from *B. subtilis* and comprises mutated amino acid positions 270, 279, 308, and/or 347 as shown in SEQ ID NO:2, corresponding to amino acids G270, Q279, K308, and M374, respectively.

In a further preferred embodiment the at least one mutation is at one or more amino acid positions selected from the group consisting of amino acid positions corresponding to positions 276, 279, 308 and 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO: 2. Thus, in a further embodiment the modified GTP cyclohydrolase II comprises one or more mutation(s) including 1, 2, 3 or 4 mutation(s) on amino acid position(s) corresponding to positions 276, 279, 308, and/or 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO:2. Preferably, the modified enzyme is obtainable from *B. subtilis* and comprises mutated amino acid positions 276, 279, 308, and/or 347 as shown in SEQ ID NO:2, corresponding to amino acids A276, Q279, K308, and M374, respectively.

Preferably, the one or more amino acid mutation(s) of the modified GTP cyclohydrolase II is one or more amino acid substitution(s).

A modified GTP cyclohydrolase II may comprise one or more mutation(s) including only one mutation on an amino acid position as defined above, such mutation, particularly an amino acid substitution, may include one mutation on an amino acid position corresponding to position 261, 270, 276, 279, 308, or 347 of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II as shown in SEQ ID NO:2. The amino acid present in the non-modified GTP cyclohydrolase II corresponding to position 261 may be valine, the amino acid present in the non-modified GTP cyclohydrolase II corresponding to position 270 may be glycine, the amino acid present in the non-modified GTP cyclohydrolase II corresponding to position 276 may be alanine, the amino acid present in the non-modified GTP cyclohydrolase II corresponding to position 279 may be glutamine, the amino acid present in the non-modified GTP cyclohydrolase II corresponding to position 308 may be lysine, and the amino acid present in the non-modified GTP cyclohydrolase II corresponding to position 347 may be methionine.

The amino acid in the sequence of the non-modified GTP cyclohydrolase II maybe changed such that the amino acid



corresponding to position 261 may be changed to alanine (e.g. V261A), the amino acid corresponding to position 270 may be changed to alanine or arginine (e.g. G270A and G270R), the amino acid corresponding to position 276 may be changed to threonine (e.g. A276T), the amino acid corresponding to position 279 may be changed to arginine (e.g. Q279A), the amino acid corresponding to position 308 may be changed to arginine (e.g. K308R), and the amino acid corresponding to position 347 may be changed to isoleucine (e.g. M374I). In one embodiment, the modified enzyme is obtainable from *B. subtilis* comprising an amino acid substitution in a position of SEQ ID NO:2 which is selected from the group consisting of position 261, 270, 276, 279, 308, and 347. Preferably, the substitution is V261A, G270A, G270R, A276T, Q279R, K308R or M347I.

A modified GTP cyclohydrolase II may comprise one or more mutation(s) including two mutations on amino acid positions as defined above, such mutations, particularly amino acid substitutions, may include mutations on amino acid positions corresponding to two of the positions as defined above, e.g. combinations of positions corresponding to positions 261/270, 261/276, 261/279, 261/308, 261/347, 270/276, 270/279, 270/308, 270/347, 276/279, 276/308, 276/347, 279/308, 279/347, or 308/347 as shown in SEQ ID NO:2. Preferred are amino acid substitutions such as V261A/A276T, V261A/Q279R, V261A/K308R, V261A/M347I, G270A/Q279R, G270A/K308R, G270A/M347I, A276T/Q279R, A276T/K308R, or A276T/M347I, wherein the positions correspond to the amino acid positions of SEQ ID NO:2. In one embodiment, such preferred substitutions are comprised in a modified GTP cyclohydrolase II obtainable from *B. subtilis* wherein the non-modified enzyme corresponds to SEQ ID NO:2. Preferably, the modified *B. subtilis* GTP cyclohydrolase II as of SEQ ID NO:2 comprises substitutions V261A/A276T or A276T/M347I.

A modified GTP cyclohydrolase II may comprise one or more mutation(s) including three mutations on amino acid positions as defined above, such mutations, particularly amino acid substitutions, may include mutations on amino acid positions corresponding to three of the positions as defined above, in particular combinations of positions corresponding to positions 261/279/308, 261/279/347, 261/308/347, 270/279/308, 270/279/347, 270/308/347, 276/279/308, 276/308/347, or 276/279/347 as shown in SEQ ID NO:2. Preferred are amino acid substitutions such as V261A/Q279R/K308R, V261A/K308R/M347I, V261A/Q279R/M347I, G270A/Q279R/K308R, G270A/K308R/M347I, G270A/Q279R/M347I, A276T/Q279R/K308R, A276T/K308R/M347I, or A276T/Q279R/M347I, wherein the positions correspond to the amino acid positions of SEQ ID NO:2. In one embodiment, such preferred substitutions are comprised in a modified GTP cyclohydrolase II obtainable from *B. subtilis* wherein the non-modified enzyme corresponds to SEQ ID NO:2. Preferably, the modified *B. subtilis* GTP cyclohydrolase II as of SEQ ID NO:2 comprises substitutions A276T/Q279R/M347I.

A modified GTP cyclohydrolase II may comprise one or more mutation(s) including four mutations on amino acid positions as defined above, such mutations, particularly amino acid substitutions, may include mutations in amino acid positions corresponding to four of the positions as defined above, in particular combinations of positions corresponding to positions 261/279/308/347, 270/279/308/347, or 276/279/308/347 as shown in SEQ ID NO:2. Preferred are amino acid substitutions such as V261A/Q279R/K308R/M347I, G270A/Q279R/K308R/M347I or A276T/Q279R/K308R/M347I, wherein the positions correspond to the

amino acid positions of SEQ ID NO:2. In one embodiment, such preferred substitutions are comprised in a modified GTP cyclohydrolase II obtainable from *B. subtilis* wherein the non-modified enzyme corresponds to SEQ ID NO:2. Preferably, the modified *B. subtilis* GTP cyclohydrolase II as of SEQ ID NO:2 comprises substitutions A276T/Q279R/K308R/M347I.

Most preferred are the combinations of mutations disclosed in Table 1 or 2 (see *infra*). The amino acid positions identified in these examples may be transferred to GTP cyclohydrolase II enzymes of different origin, as e.g. shown in Table 4.

The modified GTP cyclohydrolase II of the invention may comprise foreign amino acids, preferably at its N- or C-terminus. "Foreign amino acids" mean amino acids which are not present in a native (occurring in nature) GTP cyclohydrolase II, preferably a stretch of at least about 3, at least about 5 or at least about 7 contiguous amino acids which are not present in a native GTP cyclohydrolase II. Preferred stretches of foreign amino acids include but are not limited to "tags" that facilitate purification of the recombinantly produced modified GTP cyclohydrolase II. Examples of such tags include but are not limited to a "His<sub>6</sub>" tag, a FLAG tag, a myc tag, and the like. For calculation of specific activity, the values need to be corrected for these additional amino acids (see also above).

In another embodiment the modified GTP cyclohydrolase II may contain one or more, e.g. two, deletions when compared with the amino acid sequence of the corresponding non-modified GTP cyclohydrolase II. Preferably, the deletions affect N- or C-terminal amino acids of the corresponding non-modified GTP cyclohydrolase II and do not significantly reduce the functional properties, e.g., the specific activity, of the enzyme.

The polypeptides and polynucleotides of the present invention, including modified GTP cyclohydrolase II enzymes, may be provided in an isolated form, and preferably are purified to homogeneity. As used herein, the term "isolated" means that the material is removed from its original environment (e.g., the natural environment if it is naturally occurring). For example, a naturally-occurring polynucleotide or polypeptide present in a living microorganism is not isolated, but the same polynucleotide or polypeptide, separated from some or all of the coexisting materials in the natural system, is isolated. Such polynucleotides could be part of a vector and/or such polynucleotides or polypeptides may be part of a composition and still be isolated in that such vector or composition is not part of its natural environment. An isolated polypeptide is preferably greater than 80% pure, more preferably greater than 90% pure, even more preferably greater than 95% pure, most preferably greater than 99% pure. Purity may be determined according to methods known in the art, e.g., by SDS-PAGE and subsequent protein staining. Protein bands can then be quantified by densitometry. Further methods for determining the purity are within the level of ordinary skill.

The invention further relates to a polynucleotide comprising a nucleotide sequence which codes for a modified GTP cyclohydrolase II according to the invention. "Polynucleotide" as used herein refers to a polyribonucleotide or polydeoxyribonucleotide that may be unmodified RNA or DNA or modified RNA or DNA. Polynucleotides include but are not limited to single- and double-stranded DNA, DNA that is a mixture of single- and double-stranded regions, single- and double-stranded RNA, and RNA that is a mixture of single- and double-stranded regions, hybrid molecules comprising DNA and RNA that may be single-stranded or,



more typically, double-stranded or a mixture of single- and double-stranded regions. The term "polynucleotide" includes DNA or RNA that comprises one or more unusual bases, e.g., inosine, or one or more modified bases, e.g., tritylated bases.

The polynucleotide of the invention can easily be obtained by modifying a polynucleotide sequence which codes for a non-modified GTP cyclohydrolase II. Examples of such polynucleotide sequences encoding non-modified GTP cyclohydrolase II enzymes include but are not limited to the amino acid sequences in Table 4, in particular to SEQ ID NOs:2, 33, 35, 37, 39, 41, and 43. Non-limiting examples of polynucleotides encoding modified GTP cyclohydrolase II enzymes according to the invention are shown in SEQ ID NOs:6, 8, 10, 12, 14, 16, 18, 20, 22, 24 and 26.

Methods for introducing mutations, e.g., additions, deletions and/or substitutions into the nucleotide sequence coding for the non-modified GTP cyclohydrolase II include but are not limited to site-directed mutagenesis and PCR-based methods.

DNA sequences of the present invention can be constructed starting from genomic or cDNA sequences coding for GTP cyclohydrolase II enzymes known in the state of the art, as are available from, e.g., Genbank (Intelligenetics, California, USA), European Bioinformatics Institute (Hinton Hall, Cambridge, GB), NBRF (Georgetown University, Medical Centre, Washington D.C., USA) and Vecbase (University of Wisconsin, Biotechnology Centre, Madison, Wis., USA) or from the sequence information disclosed in Table 4 by methods of in vitro mutagenesis (see e.g. Sambrook et al., Molecular Cloning, Cold Spring Harbor Laboratory Press, New York). Another possibility of mutating a given DNA sequence which is also preferred for the practice of the present invention is mutagenesis by using the polymerase chain reaction (PCR). DNA as starting material can be isolated by methods known in the art and described, e.g., in Sambrook et al. (Molecular Cloning) from the respective strains/organisms. It is, however, understood that DNA encoding a GTP cyclohydrolase II to be constructed/mutated in accordance with the present invention can also be prepared on the basis of a known DNA sequence, e.g. by construction of a synthetic gene by methods known in the art (as described, e.g., in EP 747483).

The polynucleotide of the invention may be an isolated polynucleotide, i.e. a polynucleotide that is substantially free from other nucleic acid sequences such as but not limited to other chromosomal and extrachromosomal DNA and RNA. Conventional nucleic acid purification methods known to people skilled in the art may be used to obtain isolated polynucleotides. The term also embraces recombinant polynucleotides and chemically synthesized polynucleotides.

In yet another embodiment the invention pertains to a functional polynucleotide in which a promoter, a ribosome-binding site, if necessary as in the case of bacterial cells, and a terminator are operably linked with a polynucleotide according to the invention. In yet a further embodiment the invention pertains to a vector or plasmid comprising such a polynucleotide. The vector or plasmid preferably comprises at least one marker gene. The term "operably linked" as used herein refers to the association of nucleic acid sequences on a single nucleic acid fragment so that the function of one is affected by the other. For example, a promoter is operably linked with a coding sequence when it is capable of affecting the expression of that coding sequence, i.e., the coding sequence is under the transcriptional control of the promoter. Coding sequences may be operably linked to regulatory sequences in sense or anti-sense orientation. The term "expression" denotes the transcription of a DNA sequence into mRNA and/or the translation of mRNA into an amino acid sequence.

The term "over-expression" means the production of a gene product in a modified organism (e.g., modified by transformation or transfection) that exceeds levels of production in the corresponding non-modified organism by deregulating the expression of the gene and/or by multiplying the gene itself inside of the organism.

Once complete DNA sequences of the present invention have been obtained, they may be integrated into vectors or directly introduced into the genome of a host organism by methods known in the art and described in, e.g., Sambrook et al. (s.a.) to (over-) express the encoded polypeptide in appropriate host systems. However, a man skilled in the art knows that also the DNA sequences themselves can be used to transform the suitable host systems of the invention to get (over-) expression of the encoded polypeptide.

Suitable host cells may be eukaryotic or prokaryotic cells. Examples of suitable host cells include but are not limited to bacterial cells such as cells of cyanobacteria, streptococci, staphylococci, enterococci, e.g., *Bacillus* as, e.g., *Bacillus subtilis*, or *Streptomyces*, as, e.g. *Streptomyces lividans* or *Streptococcus pneumoniae*, *E. coli* as, e.g., *E. coli* K12 strains, e.g. M15 or HB 101. The host cells may be a fungal cell including yeast cells, such as cells of *Aspergilli*, e.g. *Aspergillus niger* or *Aspergillus oryzae*, *Trichoderma*, e.g. *Trichoderma reesei*, *Ashbya*, e.g. *Ashbya gossypii*, *Eremothecium*, e.g. *Eremothecium ashbyii*, *Saccharomyces*, e.g. *Saccharomyces cerevisiae*, *Candida*, e.g. *Candida flareri*, *Pichia*, e.g. *Pichia pastoris*, *Hansenula polymorpha*, e.g. *H. polymorpha* (DSM 5215), and *Kluyveromyces*. A suitable host cell may further be selected from animal cells, including mammalian cells, such as for instance CHO, COS, HeLa, 3T3, BHK, 293, CV-1 and insect cells like *Drosophila* S2 and *Spodoptera* Sf9 cells; and plant cells such as cells of a gymnosperm or angiosperm.

Vectors which may be used for expression in fungi are known in the art and described e.g. in EP 420358, or by Cullen et al. (Bio/Technology 5, 369-376, 1987), Ward (in Molecular Industrial Mycology, Systems and Applications for Filamentous Fungi, Marcel Dekker, New York, 1991), Upshall et al. (Bio/Technology 5, 1301-1304, 1987), Gwynne et al. (Bio/Technology 5, 71-79, 1987), or Punt et al. (J. Biotechnol. 17, 19-34, 1991), and for yeast by Sreekrishna et al. (J. Basic Microbiol. 28, 265-278, 1988; Biochemistry 28, 4117-4125, 1989), Hitzemann et al. (Nature 293, 717-722, 1981) or in EP 183070, EP 183071, EP 248227, or EP 263311. Suitable vectors which may be used for expression in *E. coli* are known in the art as described by Sambrook et al. (s.a.). Vectors which may be used for expression in *Bacilli* are known in the art and described, e.g. in EP 207459 or EP 405370, by Yansura and Henner in Proc. Natl. Acad. Sci. USA 81, 439-443 (1984), or by Henner, Le Grice and Nagarajan in Meth. Enzymol. 185, 199-228, 1990. Vectors which maybe used for expression in *H. polymorpha* are known in the art as described, e.g. in Gellissen et al., Biotechnology 9, 291-295, 1991.

Either such vectors already carry regulatory elements, e.g. promoters, or the polynucleotides of the present invention may be engineered to contain such elements. Suitable promoter elements which may be used are known in the art and are, e.g., for *Trichoderma reesei* the *cbh1*- or the *pki1*-promoter, for *Aspergillus oryzae* the *amy*-promoter, and for *Aspergillus niger* the *glaA*-, *alcA*-, *aphA*-, *tpiA*-, *gpdA*- and the *pkiA*-promoter. Suitable promoter elements which may be used for expression in yeast are known in the art and are, e.g., the *pho5*- or the *gap*-promoter for expression in *Saccharomyces cerevisiae*, and e.g. the *aox1*-promoter for *Pichia pastoris* or the *FMD*- or *MOX* promoter for *H. polymorpha*.

Suitable promoters and vectors for bacterial expression include, e.g., a synthetic promoter described by Giacomini et



al. (Gene 144, 17-24, 1994), the *vegI* promoter from *Bacillus subtilis* or the strong bacteriophage T5 promoter. Appropriate teachings for expression of the claimed (mutant) GTP cyclohydrolase II enzymes in bacteria, either by appropriate plasmids or through integration of GTP cyclohydrolase II-encoding DNA sequences into the chromosomal DNA, may be found in many places, e.g., U.S. Pat. No. 6,322,995.

Accordingly, vectors comprising a polynucleotide of the present invention, preferably for the expression of said polynucleotides in bacterial, fungal, animal or plant hosts, and such transformed bacteria or fungal, animal or plant hosts are also an object of the present invention.

The invention further relates to a method for producing riboflavin, a riboflavin precursor, FMN, FAD, or one or more derivatives thereof, comprising:

(a) culturing the host cell of the invention in a suitable medium under conditions that allow expression of the modified GTP cyclohydrolase II in said host cell; and

(b) optionally separating the product (riboflavin, a riboflavin precursor, FMN, FAD, or one or more derivatives thereof) from the medium.

Such a method can be used for the biotechnological production of either one or more of the following products: riboflavin, a riboflavin precursor, FMN, FAD, or one or more derivatives thereof. Such derivatives may include flavoproteins.

Methods of genetic and metabolic engineering of suitable host cells according to the present invention are known to the man skilled in the art. Similarly, (potentially) suitable purification methods for riboflavin, a riboflavin precursor, FMN, FAD, or one or more derivatives thereof are well known in the area of fine chemical biosynthesis and production.

It is understood that methods for biotechnological production of riboflavin, a riboflavin precursor, FMN, FAD, or one or more derivatives thereof according to the present invention are not limited to whole-cellular fermentation processes as described above, but may also use, e.g., permeabilized host cells, crude cell extracts, cell extracts clarified from cell remnants by, e.g., centrifugation or filtration, or even reconstituted reaction pathways with isolated enzymes. Also combinations of such processes are in the scope of the present invention. In the case of cell-free biosynthesis (such as with reconstituted reaction pathways), it is irrelevant whether the isolated enzymes have been prepared by and isolated from a host cell, by *in vitro* transcription/translation, or by still other means.

The invention further relates to a method for producing a modified GTP cyclohydrolase II of the invention comprising:

(a) culturing a host cell of the invention under conditions that allow expression of the modified GTP cyclohydrolase II of the invention; and

(b) recovering the modified GTP cyclohydrolase II from the cells or from the media.

The modified GTP cyclohydrolase II enzymes of the invention may be prepared from genetically engineered host cells comprising appropriate expression systems.

For recombinant production of the polypeptides of the invention, host cells may be genetically engineered to incorporate polynucleotides or vectors or plasmids of the invention. Introduction of a polynucleotide or vector into the host cell may be effected by standard methods known in the art such as calcium phosphate transfection, DEAE-dextran mediated transfection, microinjection, cationic lipid-mediated transfection, electroporation, transduction, ballistic introduction and infection.

A great variety of expression systems may be used to produce the modified GTP cyclohydrolase II enzymes of the

invention. Such vectors include, among others, those described supra. Generally, any system or vector suitable to maintain, propagate or express polynucleotides and/or to express a polypeptide in a host may be used for expression in this regard.

In recombinant expression systems in eukaryotes, for secretion of a translated protein into the lumen of the endoplasmic reticulum, into the periplasmic space or into the extracellular environment, appropriate secretion signals may be incorporated into the expressed polypeptide. These signals may be endogenous to the polypeptide or they may be heterologous signals.

Polypeptides of the invention may be recovered and purified from recombinant cell cultures by well-known methods including ammonium sulfate or ethanol precipitation, acid extraction, anion or cation exchange chromatography, phosphocellulose chromatography, hydrophobic interaction chromatography, affinity chromatography, hydroxyapatite chromatography, and high performance liquid chromatography. Well-known techniques for protein refolding may be employed to regenerate active conformation when the polypeptide is denatured during isolation and/or purification.

GTP cyclohydrolase II enzymes of the present invention may also be expressed in plants according to methods as described, e.g., by Pen et al. in *Bio/Technology* 11, 811-814, 1994 or in EP 449375, preferably in seeds as described, e.g., in EP 449376. Some suitable examples of promoters and terminators include those from nopaline synthase (*nos*), octopine synthase (*ocs*) and cauliflower mosaic virus (*CaMV*) genes. One type of efficient plant promoter that may be used is a high-level plant promoter. Such promoters, in operable linkage with the genetic sequences of the present invention should be capable of promoting expression of a gene product of the present invention. High-level plant promoters that may be used in this invention include the promoter of the small subunit (*ss*) of the ribulose-1,5-bisphosphate carboxylase, for example from soybean, and the promoter of the chlorophyll *a/b* binding protein.

Where commercial production of the instant proteins is desired, a variety of culture methodologies may be applied. For example, large-scale production of a specific gene product, overexpressed from a recombinant microbial host may be achieved by both batch or continuous culture methodologies. Batch and fed-batch culturing methods are common and well known in the art, and examples have been described by Thomas D. Brock in *Biotechnology: A Textbook of Industrial Microbiology*, Second Edition (1989), Sinauer Associates, Inc., Sunderland, Mass., or Deshpande, *Appl. Biochem. Biotechnol.* 36, 227-234, 1992. Methods of modulating nutrients and growth factors for continuous culture processes as well as techniques for maximizing the rate of product formation are well known in the art of industrial microbiology, and a variety of methods are detailed by Brock, supra.

Fermentation media may further contain suitable carbon substrates. Suitable substrates may include but are not limited to monosaccharides such as glucose and fructose, oligosaccharides such as lactose or sucrose, polysaccharides such as starch or cellulose or mixtures thereof and unpurified mixtures from renewable feedstocks. It is contemplated that the source of carbon utilized in the present invention may encompass a wide variety of carbon containing substrates and will only be limited by the choice of organism.

The invention further relates to a method for the preparation of a GTP cyclohydrolase II having increased specific activity, comprising the following steps:



(a) providing a polynucleotide encoding a first GTP cyclohydrolase II with a specific activity that, desirably, should be increased;

(b) introducing one or more mutation(s) into the polynucleotide sequence such that the mutated polynucleotide sequence encodes a modified GTP cyclohydrolase II comprising one or more mutation(s) when compared to the first GTP cyclohydrolase II wherein the one or more mutation(s) include 1, 2, 3, 4, 5, or 6 mutation(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308 and/or 347 of SEQ ID NO:2;

(c) optionally inserting the mutated polynucleotide in a vector or plasmid;

(d) introducing the polynucleotide or the vector or plasmid into a suitable host cell; and

(e) culturing the host cell under conditions that allow expression of the modified GTP cyclohydrolase II.

The present invention includes further the provision of a method for the preparation of a GTP cyclohydrolase II having increased specific activity, comprising the following steps:

(a) providing a polynucleotide encoding a first GTP cyclohydrolase II with a specific activity that, desirably, should be increased;

(b) providing the positions that have an effect on the specific activity;

(c) defining the optimal amino acid for replacement of a given amino acid of the wild-type GTP cyclohydrolase II as defined in (b) and introducing one or more mutations into the polynucleotide sequence of (a) at the positions defined in (b) such that the mutated polynucleotide sequence encodes a new GTP cyclohydrolase II;

(d) optionally inserting the mutated polynucleotide in a vector or plasmid;

(d) introducing the polynucleotide or the vector or plasmid into a suitable host cell; and

(e) culturing the host cell under conditions that allow expression of the modified GTP cyclohydrolase II.

In one embodiment, step (c) or the method described above is performed via saturated mutagenesis. However, it is understood that this may be not the only way to define the amino acid which should replace an amino acid at a given position of the wild-type GTP cyclohydrolase II in order to obtain a modified GTP cyclohydrolase II with increased specific activity.

The preparation of a modified GTP cyclohydrolase II having increased specific activity from a non-modified GTP cyclohydrolase II as described above, e.g., via saturated mutagenesis, includes, but is not limited to, the preparation of mutated GTP cyclohydrolase II proteins from non-modified proteins as in Table 4, in particular those identified by SEQ ID NOs:2, 33, 35, 37, 39, 41, and 43, such as for example non-modified GTP cyclohydrolase II proteins of *Bacillus subtilis* or *Ashbya gossypii*. The primers used for the PCR reaction are such that one primer, e.g., the sense primer, may contain a mutated nucleotide sequence and the other primer, e.g., the anti-sense primer, may contain the wild-type nucleotide sequence. PCR with these primer pairs and genomic DNA of the wild-type ribA may result in a PCR product carrying the particular mutation at a given position, depending on the mutated nucleotide sequence of the primer used. After purification of the resulting PCR products using standard methods like, e.g., the QIAquick PCR purification kit (Qiagen), the DNA may be cut with restriction enzymes such as BamHI and EcoRI, ligated into a suitable vector, e.g., pQE60, and transformed into a strain which is negative for GTP cyclohydrolase II. An example for such a strain is the *E. coli* strain Rib7 (Richter et al., J. Bacteriol. 175, 4045-4051, 1993) containing

the plasmid pREP4. After confirmation of the correct sequence by DNA sequencing, the mutated RibA may be purified and characterized as described above. If *Ashbya gossypii* is used for the generation of a GTP cyclohydrolase II having increased specific activity, saturated mutagenesis has to be performed at amino acid residues/positions T126, G135, A141, L144, N182 and I221 corresponding to the respective residues V261, G270, A276, Q279, K308 and M347 of *Bacillus subtilis* GTP cyclohydrolase II as of SEQ ID NO:2 that were shown to have an impact on the specific activity of the latter enzyme (see Table 4).

The preferred embodiments of this method correspond to the preferred embodiments of the modified GTP cyclohydrolase II, the polynucleotides encoding them, the vectors and plasmids, the host cells, and the methods described herein. The first and second GTP cyclohydrolase II correspond to the non-modified and modified GTP cyclohydrolase II, respectively (see supra).

It is an object of the present invention to provide a polynucleotide comprising a nucleic acid sequence coding for a modified GTP cyclohydrolase II as described above, a vector, preferably an expression vector, comprising such a polynucleotide, a host cell which has been transformed by such a polynucleotide or vector, a process for the preparation of a GTP cyclohydrolase II of the present invention wherein the host cell as described before is cultured under suitable culture conditions and the GTP cyclohydrolase II is isolated from such host cell or the culture medium by methods known in the art, and a process for the biotechnological production of riboflavin, a riboflavin precursor, FMN, FAD, or one or more derivatives thereof based on a host cell which has been transformed by such a polynucleotide or vector, and/or which may have stably integrated such a polynucleotide into its chromosome(s).

It is also an object of the present invention to provide (i) a DNA sequence which codes for a GTP cyclohydrolase II carrying at least one of the specific mutations of the present invention and which hybridizes under standard conditions with any of the DNA sequences of the specific modified GTP cyclohydrolase II enzymes of the present invention, or (ii) a DNA sequence which codes for a GTP cyclohydrolase II carrying at least one of the specific mutations of the present invention but, because of the degeneracy of the genetic code, does not hybridize but which codes for a polypeptide with exactly the same amino acid sequence as a DNA sequence which hybridizes under standard conditions with any of the DNA sequences of the specific modified GTP cyclohydrolase II enzymes of the present invention, or (iii) a DNA sequence which is a fragment of such DNA sequences which maintains the activity properties of the polypeptide of which it is a fragment.

“Standard conditions” for hybridization mean in the context of the present invention the conditions which are generally used by a man skilled in the art to detect specific hybridization signals and which are described, e.g. by Sambrook et al., “Molecular Cloning”, second edition, Cold Spring Harbor Laboratory Press 1989, New York, or preferably so-called stringent hybridization and non-stringent washing conditions or more preferably so-called stringent hybridization and stringent washing conditions a man skilled in the art is familiar with and which are described, e.g., in Sambrook et al. (s.a.). A specific example of stringent hybridization conditions is overnight incubation (e.g., 15 hours) at 42° C. in a solution comprising: 50% formamide, 5×SSC (150 mM NaCl, 15 mM trisodium citrate), 50 mM sodium phosphate (pH 7.6), 5× Denhardt’s solution, 10% dextran sulfate, and 20 µg/ml of



denatured, sheared salmon sperm DNA, followed by washing the hybridization support in 0.1×SSC at about 65° C.

It is furthermore an object of the present invention to provide a DNA sequence which can be obtained by the so-called polymerase chain reaction method (“PCR”) by PCR primers designed on the basis of the specifically described DNA sequences of the present invention. It is understood that the so obtained DNA sequences code for GTP cyclohydrolase II enzymes with at least the same mutation as the ones from which they are designed and show comparable activity properties.

The various embodiments of the invention described herein may be cross-combined.

Multiple sequence alignment was calculated by the program PILEUP of the GCG program package of 92 GTP cyclohydrolase II sequences found by the program BLASTN using standard databases as SWISS-PROT and TrEMBL (*Candida guilliermondii*, *Ashbya gossypii*, *Saccharomyces cerevisiae*, *Neurospora crassa*, *Schizosaccharomyces pombe*, *Archaeoglobus fulgidus*, *Streptomyces coelicolor*, *Helicobacter pylori* J99, *Helicobacter pylori*, *Pyrococcus furiosus*, *Thermotoga maritima*, *Chlamydia muridarum*, *Chlamydia trachomatis*, *Chlamydia caviae* GPIC, *Arabidopsis thaliana*, *Lycopersicon esculentum*, *Oryza sativa*, *Alcaligenes eutrophus*, *Neisseria meningitidis* (serogroup A), *Neisseria meningitidis* (serogroup B, two GTP cyclohydrolase II enzymes), *Pseudomonas putida* (two GTP cyclohydrolase II enzymes), *Pseudomonas syringae* (two GTP cyclohydrolase II enzymes), *Actinobacillus actinomycetemcomitans* (*Haemophilus actinomycetemcomitans*), *Haemophilus influenzae*, *Pasteurella multocida*, *Escherichia coli*, *Escherichia coli* O6, *Salmonella typhimurium*, *Yersinia pestis*, *Buchnera aphidicola* (subsp. *Acyrtosiphon pisum*) (*Acyrtosiphon pisum* symbiotic bacterium), *Buchnera aphidicola* (subsp. *Schizaphis graminum*), *Wigglesworthia glossinidia brevipalpis*, *Buchnera aphidicola* (subsp. *Baizongia pistaciae*), *Mycobacterium leprae*, *Mycobacterium tuberculosis*, *Corynebacterium efficiens*, *Corynebacterium glutamicum*, *Corynebacterium ammoniagenes* (*Brevibacterium ammoniagenes*), *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Actinobacillus pleuropneumoniae*, *Lactococcus lactis* (*Streptococcus lactis*), *Streptococcus agalactiae*, *Streptococcus pneumoniae*, *Clostridium acetobutylicum*, *Fusobacterium nucleatum*, *Anabaena spec.*, *Synechocystis spec.*, *Synechococcus elongatus* (*Thermosynechococcus elongatus*), *Bacillus amyloliquefaciens*, *Bacillus subtilis*, *Bacillus cereus*, *Bacillus halodurans*, *Clostridium perfringens*, *Clostridium tetani*, *Chlorobium tepidum*, *Aquifex aeolicus*, *Leptospira ininterrogans*, *Deiniococcus radiodurans*, *Bacteroides thetaiotaomicron*, *Caulobacter crescentus*, *Coxiella burnetii*, *Rhizobium etli*, *Lactobacillus plantarum*, *Pseudomonas glumae*, *Streptomyces avermitilis*, *Photobacterium phosphoreum*, *Azospirillum brasilense*, *Agrobacterium tumefaciens*, *Rhizobium meliloti* (*Sinorhizobium meliloti*), *Brucella melitensis*, *Brucella suis*, *Rhizobium loti* (*Mesorhizobium loti*), *Nitrosomonas europaea*, *Ralstonia solanacearum* (*Pseudomonas solanacearum*), *Xanthomonas axonopodis*, *Xanthomonas campestris*, *Vibrio parahaemolyticus*, *Vibrio vulnificus*, *Vibrio cholerae*, *Vibrio fischeri*, *Shewanella oneidensis*, *Photobacterium phosphoreum*, *Photobacterium leiognathi*, *Pseudomonas aeruginosa*, *Dehalospirillum multivorans*, *Xylella fastidiosa*). Numbering relates to the alignment made. Some of the amino acid sequences code for an enzyme that has just GTP cyclohydrolase II activity like the enzymes from *Ashbya gossypii*, *Streptomyces coelicolor*, *Helicobacter pylori* J99, *Helicobacter pylori*, *Arabidopsis thaliana*, *Alcaligenes eutrophus*, *Neisseria meningitidis* (serogroup A), *Neisseria meningitidis* (serogroup B), *Pseudomonas putida*, *Pseudomonas syringae*, *Actinobacillus actinomycetemcomitans* (*Haemophilus actinomycetemcomitans*), *Haemophilus*

*influenzae*, *Pasteurella multocida*, *Escherichia coli*, *Escherichia coli* O6, *Salmonella typhimurium*, *Yersinia pestis*, *Buchnera aphidicola* (subsp. *Acyrtosiphon pisum*) (*Acyrtosiphon pisum* symbiotic bacterium), *Buchnera aphidicola* (subsp. *Schizaphis graminum*), *Wigglesworthia glossinidia brevipalpis*, *Buchnera aphidicola* (subsp. *Baizongia pistaciae*), *Pseudomonas glumae*, *Streptomyces avermitilis*, or *Photobacterium phosphoreum*. Other enzymes like the RibA enzyme from *B. subtilis* contain, in addition, a domain having DHBP synthase activity. Positions that are homologous/equivalent to the amino acid residues found to have a positive effect on specific activity (amino acid residues 261, 270, 276, 279, 308, 347) and on protease sensitivity (196) of RibA from *B. subtilis* are discussed in one of the following examples. The numbering used for those positions is done according to the *B. subtilis* wild-type amino acid sequence.

The following non-limiting examples further illustrate the invention.

#### EXAMPLE 1

##### Measurement of GTP Cyclohydrolase II Activity and Determination of Specific Activity

The enzymatic assay used for measuring GTP cyclohydrolase II activity was adapted from Ritz et al. (J. Biol. Chem. 276, 22273-22277, 2001). The final assay buffer contained 50 mM Tris/HCl, pH 8.5, 10 mM MgCl<sub>2</sub>, 7.5 mM mercaptoethanol, 2.5 mM GTP and 0.1 mg/ml bovine serum albumin. After purification (see Example 5), the enzyme was kept in a buffer containing 50 mM Tris/HCl, pH 8.5, 10 mM MgCl<sub>2</sub>, 7.5 mM mercaptoethanol, and 10% glycerol. Substrate was added to the enzyme and the absorption at 310 nm, at which GTP shows no absorption, was followed over 20-30 min. The final reaction mixture contained between 0.02 and 0.04 mg/ml GTP cyclohydrolase II from *B. subtilis* or one of the mutants as shown in Table 1 or 2. An absorption coefficient of 6.28 [mM<sup>-1</sup> cm<sup>-1</sup>] for DRAPP was used for the calculation of the activity. Protein determination was done with the Protein Assay from Bio-Rad (Cat. No. 500-0002, Bio-Rad Laboratories AG, Nenzlingerweg 2, CH-4153 Reinach, Switzerland).

According to the definition of “specific activity” given above, one unit is the amount of RibA that catalyzes the formation of 1 μmol DRAPP per minute under the conditions as described above. The specific activity is the amount of DRAPP that is formed by 1 mg of RibA per minute under the conditions as described above. Using the aforementioned definitions, the specific GTP cyclohydrolase II activity of the His<sub>6</sub>-tagged RibA protein of *B. subtilis* was 0.115 U/mg.

#### EXAMPLE 2

##### Testing of the Quality of the Enzymatic Assay

An optimal assay should fulfill a number of requirements, such as linearity with enzyme concentration and linearity with time. Using the conditions described in Example 1 and 22 μg enzyme, the increase in the absorption at 310 nm was followed over 25 min. To test in which range the assay is linear with the enzyme concentration, the dependence of the assay on increasing enzyme concentration (0-40 μg His<sub>6</sub>-tagged RibA) was tested. The assay proved to be linear over 25 min and between 0 and 40 μg His<sub>6</sub>-tagged RibA from *B. subtilis*.

After this the dependence of the GTP cyclohydrolase II activity of His<sub>6</sub>-tagged RibA from *B. subtilis* on GTP concentration was tested. The conditions as described in Example 1 were used. However, the GTP concentration was varied from 0.05 to 2.5 mM final concentration. The data



## 17

indicate a  $K_m$  value for GTP of 0.07 mM and a specific activity of around 115 mU/mg protein at 37° C. for the GTP cyclohydrolase II activity of the His<sub>6</sub>-tagged RibA enzyme from *B. subtilis*. The experiments of this example showed that the GTP cyclohydrolase II activity assay, in fact, is linear with time and enzyme (GTP cyclohydrolase II) concentration, and that under the given conditions for *Bacillus subtilis* GTP cyclohydrolase II, a GTP concentration of 2.5 mM may be optimal to allow reliable measurements of the specific activity of the enzyme.

## EXAMPLE 3

Isolation of Genomic DNA from *Bacillus Subtilis*

*B. subtilis* was grown at 30° C. in Veal Infusion Broth (Becton Dickinson, Sparks, Md. 21152, USA) overnight. 1.5 ml culture was transferred into a 1.5 ml tube and centrifuged.

The cell pellet was resuspended in 0.5 ml suspension buffer (50 mM Tris/HCl, pH 7.5, 50 mM Na<sub>2</sub>EDTA, 15% sucrose and 1 mg/ml freshly added lysozyme). After 10 min incubation at room temperature 1 µl diethyloxidiformate was added. Then 10 µl of 10% SDS solution was added and the tube was inverted several times. The tube was incubated for 5 min at 70° C. to release the bacterial DNA. 50 µl 5 M potassium acetate was added, the tube was cooled on ice and left there for 45 min. After this, the sample was centrifuged for 30 min at 4° C. The supernatant was transferred into a new 1.5-ml tube, which was filled (up to 1.5 ml) with ethanol at room temperature. After 5 min centrifugation, the supernatant was discarded and the DNA pellet was dried. Then the DNA was washed with 70% and 96% ethanol and dissolved in 10 mM Tris/HCl, pH 7.5, 1 mM EDTA, and 10 µg/ml RNase A.

## EXAMPLE 4

Construction of the Expression Plasmids for Expressing RibA Coding for GTP Cyclohydrolase II and DHBP Synthase from *B. Subtilis* and its Mutants

Cloning of the ribA gene (SEQ ID NO:1) of *B. subtilis* that codes for the GTP cyclohydrolase II and the DHBP synthase was done by PCR. Genomic DNA of *B. subtilis* was isolated according to Example 3. 100 ng of this DNA or of a template coding for a mutated form of the ribA gene were used for a PCR using primers RibA 1S (SEQ ID NO:27) and RibA 1AS (SEQ ID NO:28). The following PCR conditions were used: 2 µM of each primer, 0.2 mM of each nucleotide, 2.5 U of a proof-reading DNA polymerase (Stratagene, Gebouw Calif., 1101 CB Amsterdam Zuidoost, The Netherlands), and 100 ng genomic DNA in the appropriate buffer as supplied together with the DNA polymerase.

Temperature regulation was as follows:

Step 1: 3 min at 95° C.

Step 2: 30 sec at 95° C.

Step 3: 30 sec at 52° C.

Step 4: 60 sec at 72° C.

Steps 2 to 4 were repeated 30-times.

The PCR product of 1.3 kb was used as template for PCR 2, in which primer RibA 1S was replaced by primer RibA 2S (SEQ ID NO:29). The PCR product of this reaction (SEQ ID NO:3), encoding an N-terminally His<sub>6</sub>-tagged version of *B. subtilis* RibA (SEQ ID NO:4), was separated by agarose gel electrophoresis, eluted from the gel, digested with EcoRI and BamHI, and ligated into the EcoRI and BamHI digested vector pQE60 (Qiagen AG, Hilden, Germany). The plasmid was called pQE60ribANhis.

## 18

## EXAMPLE 5

## Characterization of the Wild-Type and the Mutant Enzymes

The generation of mutated enzymes was performed using methods described above and which are known to the skilled person. Mutants of RibA from *B. subtilis* that were further investigated are depicted in Table 1. All mutant genes were cloned into a pQE60 vector as described in Example 4. All final constructs contained an N-terminal His<sub>6</sub>-tag.

TABLE 1

Rib A mutants as defined by the amino acid exchanges compared to the wild-type RibA protein of <i>B. subtilis</i> (the numbers define the respective amino acid positions in SEQ ID NO: 2)		
Mutant	SEQ ID NO (DNA)	SEQ ID NO (protein)
RibA M347I	9	10
RibA G270R	23	24
RibA K220E, G270A	19	20
RibA Y196C, A276T, A282T (PCR III)	11	12
RibA Y196C, A276T	5	6
RibA Y196C, V261A	7	8
RibA Y196C, V261A, A276T	25	26
RibA Y196C, A276T, A282T, M347I	13	14
RibA Y196C, A276T, Q279R, A282T, M347I	15	16
RibA Y196C, A276T, Q279R, A282T, K308R, M347I (construct C)	17	18
RibA Y196C, A276T, Q279R, A282T, K308R, F325Y, M347I (construct E)	21	22

The RibA mutant enzymes were expressed from the plasmids of Example 4 and purified as described in "The QiaExpressionist", Qiagen, Hilden, Germany, March 2001, edition 5. The enzymatic properties of the purified enzymes (RibA mutants) were analyzed as described in Examples 1 and 2. Table 2 compares the specific GTP cyclohydrolase II activities of the RibA mutants (see Table 1) to that of the GTP cyclohydrolase II of the wild-type RibA of *B. subtilis*. The activity was measured using the N-terminally His<sub>6</sub>-tagged enzyme versions of RibA as described in Example 4. The numbers define the respective amino acid positions in SEQ ID NO:2.

TABLE 2

Comparison of the specific GTP cyclohydrolase II activities of mutated and wild-type (WT) <i>B. subtilis</i> RibAs (all N-terminally His <sub>6</sub> -tagged)	
Mutations	specific activity (in % relative to wild-type RibA)
WT	100
M347I	120
G270R	140
G270A, (K220E)	160
(Y196C), A276T, (A282T)	160
(Y196C), A276T	160
(Y196C), V261A	160
(Y196C), V261A, A276T	160
(Y196C), A276T, (A282T), M347I	180
(Y196C), A276T, Q279R, (A282T), M347I	185
(Y196C), A276T, Q279R, (A282T), K308R, M347I	201
(Y196C), A276T, Q279R, (A282T), K308R, (F325Y), M347I	200



## 19

The amino acid replacements in parentheses have most probably no effect on GTP cyclohydrolase II activity of the mutants. Amino acid exchange Y196C reduces the protease sensitivity of RibA.

## EXAMPLE 6

Construction of Recombinant *B. Subtilis* Strains Over-Expressing RibA Mutants that Show a Higher Specific GTP Cyclohydrolase II Activity

In the following example, the mutated ribA polynucleotide sequences of RibA Y196C,A276T,282T (PCR III), RibA Y196C,A276T,Q279R,A282T,K308R,M347I (construct C), and RibA Y196C,A276T,Q279R,A282T,K308R,F325Y, M347I (construct E) were first introduced into a vector containing the strong constitutive promoter  $P_{vegI}$ , and then further manipulated in *E. coli*. Transformation of a natural competent *B. subtilis* microorganism with the polynucleotide sequence and flanking vector sequences resulted in a *B. subtilis* strain over-expressing the mutated ribA. Standard recombinant DNA techniques were used for the construction of the polynucleotide sequence and the *B. subtilis* strains. See, for example, Sambrook et al., Molecular Cloning. A Laboratory Manual (2<sup>nd</sup> Ed.), Cold Spring Harbor Laboratory Press (1989), and Harwood and Cutting, Molecular Biology Methods for *Bacillus*, John Wiley and Sons (1990).

To amplify the mutated ribA, a 1.2-kb DNA fragment containing the entire ribA coding sequence was amplified by PCR using DNA from a plasmid containing mutants PCR III, construct C or construct E, and RibANde+1 (SEQ ID NO:30) and RibA4AS (SEQ ID NO:31) as primers.

The reaction conditions for the PCR reaction consisted of 30 cycles of denaturation at 95° C. for 1 min, annealing at 52° C. for 1 min, and extension at 72° C. for 2 min. The Pfu Turbo DNA polymerase (Stratagene, Gebouw Calif., 1101 CB Amsterdam Zuidoost, The Netherlands) was used to minimize PCR-generated errors. The PCR products were purified using the QIAquick PCR purification kit (Qiagen), and doubly digested using NdeI and BamHI. The digested PCR products were cloned into the pXI16 vector (Huembelin et al., J. Ind. Microbiol. Biotechnol. 22, 1-7, 1999), which consists of suitable restriction sites for the cloning of polynucleotide sequences immediately downstream of the strong constitutive  $P_{vegI}$  promoter from *B. subtilis*. The pXI16 vector also contains the cryT transcriptional terminator from *B. thuringiensis*, the sacB flanking sequences for homologous recombination into the *B. subtilis* genome by a double-crossover event, and an erythromycin-resistance marker. That each plasmid contained the mutated ribA was confirmed by DNA sequencing.

Each plasmid was digested with ApaI to remove the spacer region from the  $P_{vegI}$  promoter, re-ligated and digested again with FspI, and transformed into natural competent *B. subtilis* 1012 cells. Transformants were selected on TBAB plates (Tryptose Blood Agar Base, Becton Dickinson, Sparks, Md. 21152, USA) containing erythromycin to a final concentration of 2 µg/ml. DNA sequencing verified that the mutated ribA polynucleotide sequences were correct in these strains. Overproduction of riboflavin was tested according to Example 7.

The mutated ribA polynucleotide sequences driven by the  $P_{vegI}$  promoter were introduced into riboflavin over-producing *B. subtilis* strain RB50::(pRF69)<sub>n</sub>::(pRF93)<sub>m</sub>, which has been described in Perkins et al., J. Ind. Microbiol. Biotechnol. 22:8-18 (1999), by generalized transduction. Standard techniques using bacteriophage PBS1 were employed according

## 20

to Harwood and Cutting, Molecular Biology Methods for *Bacillus*, John Wiley and Sons (1990). Transductants were selected for on TBAB plates containing erythromycin to a final concentration of 2 µg/ml. Transformants were checked by PCR analysis and DNA sequencing to verify correct insertion of the mutated ribA polynucleotide sequence.

## EXAMPLE 7

Improved Production of Riboflavin Using a GTP Cyclohydrolase II with Increased Specific Activity

To test the in vivo effect of mutations affecting the specific activity of GTP cyclohydrolase II, the *Bacillus subtilis* GTP cyclohydrolase II (RibA) mutants PCR III, construct C, or construct E were introduced into riboflavin over-producing *B. subtilis* strains, such as strain RB50::(pRF69)<sub>n</sub>::(pRF93)<sub>m</sub> (Perkins et al, J. Ind. Microbiol. Biotechnol. 22, 8-18, 1999), e.g. in the sacB locus. The production of riboflavin was compared directly in two recombinant strains of *B. subtilis* that differ only by the presence or absence of the mutations in the ribA gene. Culturing of the *Bacillus* strains was done as described in Example 8.

## EXAMPLE 8

Culture Conditions for Evaluating Riboflavin Production

Riboflavin production was tested in fed-batch cultivations of riboflavin-overproducing *B. subtilis* strain RB50::(pRF69)<sub>n</sub>::(pRF93)<sub>m</sub> in which the GTP cyclohydrolase II mutants PCR III, construct C, or construct E driven by the  $P_{vegI}$  promoter were integrated in the sacB locus (see Example 6). Fermentation of the strains was done as described in EP 405370.

## EXAMPLE 9

Analytical Methods for Determination of Riboflavin

For determination of riboflavin, the following analytical method can be used (Bretzel et al, J. Ind. Microbiol. Biotechnol. 22, 19-26, 1999).

The chromatographic system was a Hewlett-Packard 1100 System equipped with a binary pump, a column thermostat and a cooled autosampler. Both a diode array detector and a fluorescence detector were used in line. Two signals were recorded, UV at 280 nm and a fluorescence trace at excitation 446 nm, emission 520 nm.

A stainless-steel Supelcosil LC-8-DB column (150×4.6 mm, 3 µm particle size) was used, together with a guard cartridge. The mobile phases were 100 mM acetic acid (A) and methanol (B). A gradient elution according to the following scheme was used:

Time [min]	% A	% B
0	98	2
6	98	2
15	50	50
25	50	50

The column temperature was set to 20° C., and the flow rate was 1.0 ml/min. The run time was 25 min.



## 21

Fermentation samples were diluted, filtered and analyzed without further treatment. Riboflavin was quantitated by comparison with an external standard. The calculations were based on the UV signal at 280 nm. Riboflavin purchased from Fluka (9471 Buchs, Switzerland) was used as standard material (purity  $\geq 99.0\%$ ).

## EXAMPLE 10

Identification of Corresponding Residues in GTP  
Cyclohydrolase II Enzymes that are Homologous to  
*Bacillus subtilis* GTP Cyclohydrolase II

A multiple amino acid sequence alignment of 92 different GTP cyclohydrolase II enzymes found by the program BLASTN using standard databases such as SWISS-PROT and TrEMBL was calculated with the program "PILEUP" (GCG Wisconsin Package, version 10.2, Accelrys Inc., 9685 Scranton Road, San Diego, Calif. 92121-3752, USA) using the following parameters: gap creation penalty 8, gap extension penalty 2, and blosum62.cmp matrix (default parameters).

A homologous GTP cyclohydrolase II in the context of the present invention may show sequence similarity with any of the GTP cyclohydrolase II amino acid sequences shown in Table 4, which serves as an example and is not meant to be a complete collection of all known GTP cyclohydrolase II enzymes. Homologous residues, i.e. residues of the different GTP cyclohydrolase II enzymes that are located at the same position in the amino acid sequence alignment, are expected to be similarly positioned in the 3D structure of each protein and to fulfill in each protein a comparable function structure-wise and function-wise. Amino acid residues homologous to the amino acid residues of the GTP cyclohydrolase II from *B. subtilis* are discussed in the Examples.

Amino acid residues of 92 different organisms corresponding to specific amino acid positions, i.e. positions that are homologous/equivalent to the amino acid residues found to have a positive effect on specific activity (amino acid residues 261, 270, 276, 279, 308, 347) of the amino acid sequence of *Bacillus subtilis* GTP cyclohydrolase II (SEQ ID NO:2) are summarized in Table 4, wherein the number in the left column defines the organism, starting with the name of the sequence used, the database accession number, and in parenthesis the source organism of the sequence:

- (1) gch2\_bacsu: SWISS-PROT: gch2\_bacsu (*Bacillus subtilis*)
- (2) gch2\_cangu: geneseqp:aay69776 (*Candida guilliermondii*)
- (3) gch2\_ashgo: TrEMBL: CAA02912: (*Ashbya gossypii* (*Eremothecium gossypii*))
- (4) gch2\_yeast: SWISS-PROT: gch2\_yeast (*Saccharomyces cerevisiae*)
- (5) gch2\_neucr: TrEMBL: Q871B3 (*Neurospora crassa*)
- (6) gch2\_schpo: TrEMBL: Q9P7M9 (*Schizosaccharomyces pombe*)
- (7) gch2\_arcfu: SWISS-PROT: gch2\_arcfu (*Archaeoglobus fulgidus*)
- (8) gch2\_strco: SWISS-PROT: gch2\_strco (*Streptomyces coelicolor*)
- (9) gch2\_helpj: SWISS-PROT: gch2\_helpj (*Helicobacter pylori* J99)
- (10) gch2\_helpy: SWISS-PROT: gch2\_helpy (*Heliobacter pylori*)
- (11) gch2\_pyrfu: TrEMBL: Q8U4L7 (*Pyrococcus furiosus*)

## 22

- (12) gch2\_thema: SWISS-PROT: gch2\_thema (*Thermotoga maritima*)
- (13) gch2\_chlmu: SWISS-PROT: gch2\_chlmu (*Chlamydia muridarum*)
- (14) gch2\_chltr: SWISS-PROT: gch2\_chltr (*Chlamydia trachomatis*)
- (15) gch2\_chlca: TrEMBL: AAP05635 (*Chlamydia caviae* GPIC)
- (16) gch2\_chlpn: SWISS-PROT: gch2\_chlpn (*Chlamydia pneumoniae*)
- (17) gch2\_arath: SWISS-PROT: gch2\_arath (*Arabidopsis thaliana*)
- (18) gch2\_lyces: TrEMBL: CAC09119 (*Lycopersicon esculentum*)
- (19) gch2\_orysa: TrEMBL: AAO72560 (*Oryza sativum*)
- (20) gch2\_alceu: TrEMBL: Q9F184 (*Alcaligenes eutrophus*)
- (21) gch2\_neima: SWISS-PROT: gch2\_neima (*Neisseria meningitidis* (serogroup A))
- (22) gch2\_neimb: SWISS-PROT: gch2\_neimb (*Neisseria meningitidis* (serogroup B))
- (23) gch2\_psepik: SWISS-PROT: gch2\_psepik (*Pseudomonas putida* (strain KT2440))
- (24) gch2\_psesm: SWISS-PROT: gch2\_psesm (*Pseudomonas syringae* (pv. tomato))
- (25) gch2\_actac: TrEMBL: Q9JRR0 (*Actinobacillus actinomycetemcomitans* (*Haemophilus actinomycetemcomitans*))
- (26) gch2\_haein: SWISS-PROT: gch2\_pasmu gch2\_haein (*Haemophilus influenzae*)
- (27) gch2\_pasmu: SWISS-PROT: (*Pasteurella multocida*)
- (28) gch2\_ecO6: TrEMBL: Q8FHU5 (*Escherichia coli* O6)
- (29) gch2\_ecoli: SWISS-PROT: gch2\_ecoli (*Escherichia coli*)
- (30) gch2\_salty: TrEMBL: Q8XFY7 (*Salmonella typhimurium*)
- (31) gch2\_yerpe: TrEMBL: Q8ZEF0 (*Yersinia pestis*)
- (32) gch2\_bucal: SWISS-PROT: gch2\_bucal (*Buchnera aphidicola* (subsp. *Acyrtosiphon pisum*) (*Acyrtosiphon pisum* symbiotic bacterium))
- (33) gch2\_bucap: SWISS-PROT: gch2\_bucap (*Buchnera aphidicola* (subsp. *Schizaphis graminum*))
- (34) gch2\_wigbr: SWISS-PROT: gch2\_wigbr (*Wigglesworthia glossinidia brevipalpis*)
- (35) gch2\_bucbp: SWISS-PROT: gch2\_wigbr (*Buchnera aphidicola* (subsp. *Baizongia pistaciae*))
- (36) gch2\_mytle: TrEMBL: Q9CCP4 (*Mycobacterium leprae*)
- (37) gch2\_myctu: SWISS-PROT: gch2\_myctu (*Mycobacterium tuberculosis*)
- (38) gch2\_coref: TrEMBL: Q8FT57 (*Corynebacterium efficiens*)
- (39) gch2\_corgl: GENESEQP: AAB79913 (*Corynebacterium glutamicum*)
- (40) gch2\_coram: SWISS-PROT: gch2\_coram (*Corynebacterium ammoniagenes* (*Brevibacterium ammoniagenes*))
- (41) gch2\_staau: TrEMBL: Q8NW14 (*Staphylococcus aureus* (strain MW2))
- (42) gch2\_staep: GENESEQP: ABP40248 (*Staphylococcus epidermidis*)
- (43) gch2\_actpl: SWISS-PROT: gch2\_actpl (*Actinobacillus pleuropneumoniae*)
- (44) gch2\_lacla: TrEMBL: Q9CGU7 (*Lactococcus lactis* (subsp. *lactis*) (*Streptococcus lactis*))
- (45) gch2\_stcag: TrEMBL: Q8E658 (*Streptococcus agalactiae* (serotype III))



## 23

- (46) gch2\_stcpn: TrEMBL: Q8DRF1 (*Streptococcus pneumoniae* (strain ATCC BAA-255/R6))
- (47) gch2\_cloac: TrEMBL: Q97LG9 (*Clostridium acetobutylicum*)
- (48) gch2\_fusnu: TrEMBL: Q8RIR1 (*Fusobacterium nucleatum* (subsp. *nucleatum*))
- (49) gch2\_anasp: TrEMBL: Q8RIR1 (*Anabaena* sp. (strain PCC 7120))
- (50) gch2\_syny3: SWISS-PROT: gch2\_syny3 (*Synechocystis* sp. (strain PCC 6803))
- (51) gch2\_synel: TrEMBL: Q8DI64 *Synechococcus elongatus* (*Thermosynechococcus elongatus*)
- (52) gch2\_bacam: SWISS-PROT: gch2\_bacam (*Bacillus amyloliquefaciens*)
- (53) gch2\_bacce: TrEMBL: AAP11030 (*Bacillus cereus* ATCC 14579)
- (54) gch2\_bacha: TrEMBL: Q9KCL5 (*Bacillus halodurans*)
- (55) gch2\_clope: TrEMBL: Q8XMX0 (*Clostridium perfringens*)
- (56) gch2\_clote: TrEMBL: Q897Q8 (*Clostridium tetani*)
- (57) gch2\_chlte: TrEMBL: Q8KC35 (*Chlorobium tepidum*)
- (58) gch2\_aquae: SWISS-PROT: gch2\_aquae (*Aquifex aeolicus*)
- (59) gch2\_lepin: TrEMBL: Q8F701 (*Leptospira interrogans*)
- (60) gch2\_deira: TrEMBL: Q9RXZ9 (*Deinococcus radiodurans*)
- (61) gch2\_bacth: TrEMBL: Q8A528 (*Bacteroides thetaiotaomicron*)
- (62) gch2\_caucr: TrEMBL: Q9A9S5 (*Caulobacter crescentus*)
- (63) gch2\_coxbu: TrEMBL: AAO90191 (*Coxiella burnetii* RSA 493)
- (64) gch2\_rhiet: TrEMBL: Q8KL38 (*Rhizobium etli*)
- (65) gch2\_lacpl: TrEMBL: Q88X17 (*Lactobacillus plantarum*)
- (66) gch2\_psegl: TrEMBL: Q8RS38 (*Pseudomonas glumae*)
- (67) gch2\_strav: TrEMBL: BAC71833 (*Streptomyces avermitilis*)
- (68) gch2\_phopo: SWISS-PROT: gch2\_phopo (*Photobacterium phosphoreum*)
- (69) gch2\_azobr: SWISS-PROT: gch2\_azobr (*Azospirillum brasilense*)
- (70) gch2\_agrtu: TrEMBL: Q8UHC9 (*Agrobacterium tumefaciens* (strain C58/ATCC 33970))
- (71) gch2\_rhime: TrEMBL: Q92RH2 (*Rhizobium meliloti* (*Sinorhizobium meliloti*))
- (72) gch2\_brume: TrEMBL: Q8YFL5 (*Brucella melitensis*)
- (73) gch2\_brusu: TrEMBL: Q8G298 (*Brucella suis*)
- (74) gch2\_rhilo: TrEMBL: Q985Z3 (*Rhizobium loti* (*Mesorhizobium loti*))
- (75) gch2\_braja: TrEMBL: Q89RZ7 (*Bradyrhizobium japonicum*)
- (76) gch2\_niteu: TrEMBL: CAD86468 (*Nitrosomonas europaea* ATCC 19718)
- (77) gch2\_ralso: TrEMBL: Q8Y1H7 (*Ralstonia solanacearum* (*Pseudomonas solanacearum*))
- (78) gch2\_neime: TrEMBL: Q9JZ77 (*Neisseria meningitidis* (serogroup B, second enzyme found))
- (79) gch2\_xanax: TrEMBL: Q8PPD7 (*Xanthomonas axonopodis* (pv. *citri*))
- (80) gch2\_xanca: TrEMBL: Q8PCM8 (*Xanthomonas campestris* (pv. *campestris*))

## 24

- (81) gch2\_vibpa: TrEMBL: Q87RU5 (*Vibrio parahaemolyticus*)
- (82) gch2\_vibvu: TrEMBL: Q8DF98 (*Vibrio vulnificus*)
- (83) gch2\_vibch: TrEMBL: Q9KPU3 (*Vibrio cholerae*)
- (84) gch2\_vibfi: TrEMBL: Q8G9G5 (*Vibrio fischeri*)
- (85) gch2\_sheon: TrEMBL: Q8EBP2 (*Shewanella oneidensis*)
- (86) gch2\_phoph: TrEMBL: Q8G9H7 (*Photobacterium phosphoreum*)
- (87) ribb\_phole: TrEMBL: Q93E93 (*Photobacterium leiognathi*)
- (88) gch2\_psepu: TrEMBL: Q88GB1 (*Pseudomonas putida* (strain KT2440, second enzyme found))
- (89) gch2\_psesy: TrEMBL: Q882G0 (*Pseudomonas syringae* (pv. *Tomato*, second enzyme found))
- (90) gch2\_pseae: TrEMBL: Q9HWX4 (*Pseudomonas aeruginosa*)
- (91) ribb\_dehmu: SWISS-PROT: ribb\_dehmu (*Dehalospirillum multivorans*)
- (92) gch2\_xylfa: TrEMBL: Q87D69 (*Xylella fastidiosa* (strain Temecula1/ATCC 700964))

TABLE 4

Positions/amino acid residues corresponding to positions V261, G270, A276, Q279, K308, and M347 of RibA of *B. subtilis* as of SEQ ID NO: 2. The numbers in the left column refer to the different organisms (see above).

							SEQ ID NO DNA; protein
	(1) V261	G270	A276	Q279	K308	M347	1; 2
	(2) T181	G190	A196	L199	N228	I267	
	(3) T126	G135	A141	L144	N182	I221	32; 33
	(4) N153	G162	A168	L171	N211	V250	
	(5) T255	G264	A270	L273	N305	I344	
	(6) T176	G185	A191	L194	N223	I262	
	(7) V139	G248	A254	M257	K287	I326	
	(8) A73	G82	A88	Q91	A121	V160	
	(9) A60	G69	A75	R78	A106	M145	
	(10) A60	G69	A75	R78	A106	M145	
	(11) T151	G160	A166	T169	E297	I336	
	(12) V245	G254	F260	Y263	S291	V330	
	(13) V267	G276	A282	Y285	A315	V354	
	(14) I267	G276	A282	Y285	A315	V354	
	(15) I272	G281	A287	Y290	A320	I359	
	(16) I272	G281	A287	Y290	A320	I359	
	(17) I91	G100	S106	Q109	N139	M178	
	(18) I299	G308	A314	Q317	N347	M386	
	(19) I295	G304	A310	L313	N343	M382	
	(20) V61	G70	A76	K79	R109	V148	
	(21) A60	G69	A75	A78	H107	V146	
	(22) A60	G69	A75	A78	H107	V146	
	(23) A59	G68	A74	A77	E106	L145	
	(24) A59	G68	A74	A77	E106	L145	
	(25) A61	G70	A76	A79	S108	I147	
	(26) A61	G70	A76	A79	S108	V147	
	(27) A61	G70	A76	A79	S108	V147	
	(28) A80	G89	A95	Q98	A127	V166	
	(29) A59	G68	A74	Q77	A106	V145	34; 35
	(30) A59	G68	A74	H77	A106	V145	
	(31) A59	G68	A74	R77	A106	V145	
	(32) A57	G66	S72	R75	A104	I143	
	(33) A59	G68	A74	R77	S106	I145	
	(34) A59	G68	A74	H77	S106	I145	
	(35) A60	G69	A75	E78	A107	I146	
	(36) V279	G288	A294	M297	Q327	M366	
	(37) V269	G278	A284	M287	Q317	M356	
	(38) V281	G290	S296	M299	Q329	I368	
	(39) V273	G282	S288	L291	Q321	L360	36; 37
	(40) V274	G283	S289	I292	S322	A361	
	(41) I259	G268	S274	Y277	E305	I344	
	(42) I263	G272	S276	Y279	E309	I348	
	(43) A264	G273	A279	Q282	E311	I350	
	(44) A262	G271	A277	K280	S309	L348	



TABLE 4-continued

Positions/amino acid residues corresponding to positions V261, G270, A276, Q279, K308, and M347 of RibA of <i>B. subtilis</i> as of SEQ ID NO: 2. The numbers in the left column refer to the different organisms (see above).						SEQ ID NO DNA; protein
(45)	V261	G270	A276	Q279	H308	I347
(46)	V261	G270	A276	M279	H308	L347
(47)	V264	G273	A279	A282	M311	V350
(48)	I261	G270	A276	R279	N308	I347
(49)	A301	R310	A316	M319	S348	I387
(50)	A265	R274	A280	M283	S312	I351
(51)	A267	R276	A282	M285	S315	I353
(52)	V261	G270	A276	Q279	R308	M347 38; 39
(53)	V260	G269	A275	Q278	K307	L346 40; 41
(54)	V264	G273	A279	Q282	K311	M350 42; 43
(55)	V244	G253	A259	K262	K291	I330
(56)	A265	G274	A280	A283	N312	I351
(57)	T267	G276	A282	M285	N314	M353
(58)	V270	R279	A285	M288	E317	M356
(59)	I263	G272	A278	M281	N310	M349
(60)	A269	G278	A284	A287	A316	L355
(61)	I265	G274	A281	M283	K314	M351
(62)	A260	G269	S276	Q278	A307	V346
(63)	V265	G274	A280	E283	A311	I350
(64)	A283	G292	A298	A301	S330	V369
(65)	V262	G271	A277	K280	A309	V348
(66)	V70	G79	S85	L88	R117	V156
(67)	V66	G75	A81	R84	A112	L151
(68)	G61	G70	T76	I79	K108	I147
(69)	L250	G259	A265	E268	R297	V336
(70)	V264	—	Y274	K277	K304	I340
(71)	V264	—	I274	R277	K304	I340

TABLE 4-continued

Positions/amino acid residues corresponding to positions V261, G270, A276, Q279, K308, and M347 of RibA of <i>B. subtilis</i> as of SEQ ID NO: 2. The numbers in the left column refer to the different organisms (see above).						SEQ ID NO DNA; protein
(72)	V264	—	I275	R278	R305	I345
(73)	V264	—	I275	R278	R305	I345
(74)	V264	—	I274	A277	—	I340
(75)	I261	—	V271	H274	—	I330
(76)	L262	S271	A277	V280	D303	M338
(77)	L285	S294	A300	A303	Q327	M362
(78)	F263	S271	A277	H280	D303	L336
(79)	L262	G271	A277	A280	R304	L339
(80)	L262	G271	A277	A280	R304	L339
(81)	V261	S271	A277	R280	R303	M343
(82)	L261	S271	A277	R280	R303	M343
(83)	L261	S271	A277	R280	R303	M343
(84)	I261	S271	A277	R280	R303	M343
(85)	L260	S270	A276	R279	K302	M343
(86)	I262	—	A275	R278	K301	M341
(87)	I262	—	A275	R278	Q301	I339
(88)	L262	N272	A278	K281	R305	L342
(89)	L262	N272	A278	R281	R305	L342
(90)	L262	R270	A276	K279	H303	M339
(91)	L269	Y276	A282	Y285	—	I324
(92)	L230	G239	S245	T248	G277	I316

The examples shown in Table 4 serve as illustration of the principle. Corresponding residues can be determined for all other GTP cyclohydrolase II amino acid sequences that are homologous to any one of the sequences shown in Table 4.

## SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 43

<210> SEQ ID NO 1

<211> LENGTH: 1197

<212> TYPE: DNA

<213> ORGANISM: *Bacillus subtilis*

<400> SEQUENCE: 1

```

atgtttcatc cgatagaaga agcactggac gctttaaaaa aaggcgaagt catcatcggt      60
gtagatgatg aagacagaga aaatgaagga gactttgtgg ctcttgccga gcatgcaacg     120
ccggaagtca ttaactttat ggcgacacat gggagaggac tgatctgcac gccgctcagt     180
gaggaaatcg cagacaggct tgatcttcac cctatggttg agcataatac agactctcac     240
cactactgcat ttaccgtaag catagacatc cgtgaaacga agacaggtat cagcgtctcaa     300
gaaagatctt ttaccgttca agcattgctg gacagcaaat ccgtgccatc tgattttcag     360
cgtccggggc acatttttcc actgattgcg aaaaaaggag gtgtcctgaa aagagcgggc     420
catacagaag ctgctgttga tcttgctgaa gcttggtgat ctccaggagc cggcgtcatt     480
tgtgaaatta tgaatgaaga cggaacgatg gcgagagtgc ctgagctcat tgaaattgctg     540
aaaaagcatc aattaaat gatcaccatt aaggatttga ttcaataccg ttacaatctg       600
acaacacttg tcgagcgtga agttgacatt acgctgccta ctgattttgg gacatttaag     660
gtttatggat acacaaatga ggtagatgga aaagagcatg tcgcatttgt gatgggagat     720
gtgccgttcg gagaagaacc ggtattggtc cgggtgcatt cagaatgtct cacaggtgac     780
gtgtttggct ctcatcgctg tgattgcgga ccgagctgc acgccgcgt gaaccaaatt      840

```



-continued

---

```

gccgcagaag gccgtggagt gctcctgtac ttgcgccaag aaggacgagg catcggttta 900
atcaataaat taaaagctta taagcttcag gaacaaggct atgacaccgt agaagccaat 960
gaggcgcttg gattcttgcc ggatcttcgc aactatggca tcggagcaca aatcttacgc 1020
gacctcgggtg tccggaatat gaagcttttg acgaataatc cgcgaaaaat cgcaggcctt 1080
gaaggctacg gactcagtat ttcagaaaga gtgccgcttc aaatggaggc gaaagaacac 1140
aataaaaaat atttgcaaac caaaatgaac aagctaggtc atttacttca tttctaa 1197

```

&lt;210&gt; SEQ ID NO 2

&lt;211&gt; LENGTH: 398

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 2

```

Met Phe His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu
1           5           10           15
Val Ile Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe
20           25           30
Val Ala Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala
35           40           45
Thr His Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala
50           55           60
Asp Arg Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His
65           70           75           80
His Thr Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly
85           90           95
Ile Ser Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser
100          105          110
Lys Ser Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu
115          120          125
Ile Ala Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala
130          135          140
Ala Val Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile
145          150          155          160
Cys Glu Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu
165          170          175
Ile Glu Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp
180          185          190
Leu Ile Gln Tyr Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val
195          200          205
Asp Ile Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr
210          215          220
Thr Asn Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp
225          230          235          240
Val Pro Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys
245          250          255
Leu Thr Gly Asp Val Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln
260          265          270
Leu His Ala Ala Leu Asn Gln Ile Ala Ala Glu Gly Arg Gly Val Leu
275          280          285
Leu Tyr Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu
290          295          300
Lys Ala Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn
305          310          315          320

```



-continued

Glu Ala Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala  
 325 330 335  
 Gln Ile Leu Arg Asp Leu Gly Val Arg Asn Met Lys Leu Leu Thr Asn  
 340 345 350  
 Asn Pro Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser  
 355 360 365  
 Glu Arg Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr  
 370 375 380  
 Leu Gln Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
 385 390 395

<210> SEQ ID NO 3  
 <211> LENGTH: 1239  
 <212> TYPE: DNA  
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 3

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa 60  
 gaagcactgg acgctttaa aaaaggcgaa gtcacatcgt ttgtagatga tgaagacaga 120  
 gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt 180  
 atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg 240  
 cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta 300  
 agcatagacc atcgtgaaac gaagacaggt atcagcgtc aagaaagatc ttttaccggt 360  
 caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt 420  
 cactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt 480  
 gatcttgctg aagcttggtg atctccagga gccggcgtca tttgtgaaat tatgaatgaa 540  
 gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa 600  
 atgatcacca ttaaggattt gattcaatac cgttacaatc tgacaacact tgtcagcgt 660  
 gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720  
 gagtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccgtt cggagaagaa 780  
 ccggtattgg tccgggtgca ttcagaatgt ctcacagggt acgtgtttgg ctctcatcgc 840  
 tgtgattgcg gaccgcagct gcacgccgcg ctgaaccaa ttgccgcaga aggccgtgga 900  
 gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct 960  
 tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgtc tggattcttg 1020  
 ccggatcttc gcaactatgg catcggagca caaatcttac gcgacctcgg tgtccggaat 1080  
 atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140  
 atttcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200  
 accaaaatga acaagctagg tcatttactt catttctaa 1239

<210> SEQ ID NO 4  
 <211> LENGTH: 412  
 <212> TYPE: PRT  
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 4

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe  
 1 5 10 15  
 His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile  
 20 25 30



-continued

Ile	Val	Val	Asp	Asp	Glu	Asp	Arg	Glu	Asn	Glu	Gly	Asp	Phe	Val	Ala
		35					40					45			
Leu	Ala	Glu	His	Ala	Thr	Pro	Glu	Val	Ile	Asn	Phe	Met	Ala	Thr	His
	50					55					60				
Gly	Arg	Gly	Leu	Ile	Cys	Thr	Pro	Leu	Ser	Glu	Glu	Ile	Ala	Asp	Arg
65					70					75					80
Leu	Asp	Leu	His	Pro	Met	Val	Glu	His	Asn	Thr	Asp	Ser	His	His	Thr
				85					90					95	
Ala	Phe	Thr	Val	Ser	Ile	Asp	His	Arg	Glu	Thr	Lys	Thr	Gly	Ile	Ser
			100					105					110		
Ala	Gln	Glu	Arg	Ser	Phe	Thr	Val	Gln	Ala	Leu	Leu	Asp	Ser	Lys	Ser
		115					120					125			
Val	Pro	Ser	Asp	Phe	Gln	Arg	Pro	Gly	His	Ile	Phe	Pro	Leu	Ile	Ala
	130					135					140				
Lys	Lys	Gly	Gly	Val	Leu	Lys	Arg	Ala	Gly	His	Thr	Glu	Ala	Ala	Val
145					150					155					160
Asp	Leu	Ala	Glu	Ala	Cys	Gly	Ser	Pro	Gly	Ala	Gly	Val	Ile	Cys	Glu
				165					170					175	
Ile	Met	Asn	Glu	Asp	Gly	Thr	Met	Ala	Arg	Val	Pro	Glu	Leu	Ile	Glu
			180					185					190		
Ile	Ala	Lys	Lys	His	Gln	Leu	Lys	Met	Ile	Thr	Ile	Lys	Asp	Leu	Ile
		195					200					205			
Gln	Tyr	Arg	Tyr	Asn	Leu	Thr	Thr	Leu	Val	Glu	Arg	Glu	Val	Asp	Ile
	210					215					220				
Thr	Leu	Pro	Thr	Asp	Phe	Gly	Thr	Phe	Lys	Val	Tyr	Gly	Tyr	Thr	Asn
225					230					235					240
Glu	Val	Asp	Gly	Lys	Glu	His	Val	Ala	Phe	Val	Met	Gly	Asp	Val	Pro
				245					250					255	
Phe	Gly	Glu	Glu	Pro	Val	Leu	Val	Arg	Val	His	Ser	Glu	Cys	Leu	Thr
			260					265					270		
Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys	Gly	Pro	Gln	Leu	His
		275					280					285			
Ala	Ala	Leu	Asn	Gln	Ile	Ala	Ala	Glu	Gly	Arg	Gly	Val	Leu	Leu	Tyr
	290					295					300				
Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Ile	Asn	Lys	Leu	Lys	Ala
305					310					315					320
Tyr	Lys	Leu	Gln	Glu	Gln	Gly	Tyr	Asp	Thr	Val	Glu	Ala	Asn	Glu	Ala
			325						330					335	
Leu	Gly	Phe	Leu	Pro	Asp	Leu	Arg	Asn	Tyr	Gly	Ile	Gly	Ala	Gln	Ile
			340					345					350		
Leu	Arg	Asp	Leu	Gly	Val	Arg	Asn	Met	Lys	Leu	Leu	Thr	Asn	Asn	Pro
	355						360					365			
Arg	Lys	Ile	Ala	Gly	Leu	Glu	Gly	Tyr	Gly	Leu	Ser	Ile	Ser	Glu	Arg
	370					375					380				
Val	Pro	Leu	Gln	Met	Glu	Ala	Lys	Glu	His	Asn	Lys	Lys	Tyr	Leu	Gln
385					390					395					400
Thr	Lys	Met	Asn	Lys	Leu	Gly	His	Leu	Leu	His	Phe				
				405						410					

&lt;210&gt; SEQ ID NO 5

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis



-continued

&lt;400&gt; SEQUENCE: 5

```

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa    60
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga    120
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt    180
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg    240
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta    300
agcatagacc atcgtgaaac gaagacaggt atcagcgtc aagaaagatc ttttaccgtt    360
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt    420
ccactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt    480
gatcttgctg aagcttgagg atctccagga gccggcgtea tttgtgaaat tatgaatgaa    540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa    600
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgtcagcgt    660
gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat    720
gaggtagatg gaaaagagca tgtcgcatth gtgatgggag atgtgccgtt cggagaagaa    780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgtttgg ctctcatcgc    840
tgtgattgcg gaccgcagct gcacgccacg ctgaaccaa ttgccgcaga aggccgtgga    900
gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct    960
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg   1020
ccggatcttc gcaactatgg catcggagca caaatthtac gcgacctcgg tgtccggaat   1080
atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt   1140
atctcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa   1200
acaaaaatga acaagctagg tcatttactt catttctaa                            1239

```

&lt;210&gt; SEQ ID NO 6

&lt;211&gt; LENGTH: 412

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 6

```

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1          5          10          15

His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
          20          25          30

Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
          35          40          45

Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
          50          55          60

Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80

Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
          85          90          95

Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
          100          105          110

Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
          115          120          125

Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
          130          135          140

```



-continued

Lys	Lys	Gly	Gly	Val	Leu	Lys	Arg	Ala	Gly	His	Thr	Glu	Ala	Ala	Val
145					150					155					160
Asp	Leu	Ala	Glu	Ala	Cys	Gly	Ser	Pro	Gly	Ala	Gly	Val	Ile	Cys	Glu
			165						170					175	
Ile	Met	Asn	Glu	Asp	Gly	Thr	Met	Ala	Arg	Val	Pro	Glu	Leu	Ile	Glu
		180						185					190		
Ile	Ala	Lys	Lys	His	Gln	Leu	Lys	Met	Ile	Thr	Ile	Lys	Asp	Leu	Ile
		195					200					205			
Gln	Cys	Arg	Tyr	Asn	Leu	Thr	Thr	Leu	Val	Glu	Arg	Glu	Val	Asp	Ile
	210					215					220				
Thr	Leu	Pro	Thr	Asp	Phe	Gly	Thr	Phe	Lys	Val	Tyr	Gly	Tyr	Thr	Asn
225					230					235					240
Glu	Val	Asp	Gly	Lys	Glu	His	Val	Ala	Phe	Val	Met	Gly	Asp	Val	Pro
				245					250					255	
Phe	Gly	Glu	Glu	Pro	Val	Leu	Val	Arg	Val	His	Ser	Glu	Cys	Leu	Thr
			260					265					270		
Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys	Gly	Pro	Gln	Leu	His
		275					280					285			
Ala	Thr	Leu	Asn	Gln	Ile	Ala	Ala	Glu	Gly	Arg	Gly	Val	Leu	Leu	Tyr
		290				295					300				
Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Ile	Asn	Lys	Leu	Lys	Ala
305					310					315					320
Tyr	Lys	Leu	Gln	Glu	Gln	Gly	Tyr	Asp	Thr	Val	Glu	Ala	Asn	Glu	Ala
				325					330					335	
Leu	Gly	Phe	Leu	Pro	Asp	Leu	Arg	Asn	Tyr	Gly	Ile	Gly	Ala	Gln	Ile
			340					345					350		
Leu	Arg	Asp	Leu	Gly	Val	Arg	Asn	Met	Lys	Leu	Leu	Thr	Asn	Asn	Pro
		355					360					365			
Arg	Lys	Ile	Ala	Gly	Leu	Glu	Gly	Tyr	Gly	Leu	Ser	Ile	Ser	Glu	Arg
	370					375					380				
Val	Pro	Leu	Gln	Met	Glu	Ala	Lys	Glu	His	Asn	Lys	Lys	Tyr	Leu	Gln
385					390					395					400
Thr	Lys	Met	Asn	Lys	Leu	Gly	His	Leu	Leu	His	Phe				
			405						410						

&lt;210&gt; SEQ ID NO 7

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 7

atgagaggat ctcacatca ccatcacat gggatcgatc atatgtttca tccgatagaa	60
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga	120
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt	180
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg	240
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta	300
agcatagacc atcgtgaaac gaagacaggt atcagcgctc aagaaagatc ttttaccggt	360
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt	420
ccactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgtctgt	480
gatcttgctg aagcttgctg atctccagga gccggcgctca tttgtgaaat tatgaatgaa	540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa	600
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgctcgagcgt	660



-continued

---

```

gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720
gaggtagatg gaaaagagca tgtcgcatth gtgatgggag atgtgccggt cggagaagaa 780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgcgtttg ctctcatcgc 840
tgtgattgcg gaccgcagct gcacgccgcg ctgaacccaa ttgccgcaga aggccgtgga 900
gtgctcctgt acttgcgcca agaaggacga ggcacgggtt taatcaataa attaaaagct 960
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgcct tggattcttg 1020
ccggatcttc gcaactatgg catcggagca caaatthtac gcgacctcgg tgtccggaat 1080
atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140
atttcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200
acccaaatga acaagctagg tcatttactt catttctaa 1239

```

```

<210> SEQ ID NO 8
<211> LENGTH: 412
<212> TYPE: PRT
<213> ORGANISM: Bacillus subtilis

```

```

<400> SEQUENCE: 8

```

```

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1          5          10          15
His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
20          25          30
Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
35          40          45
Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
50          55          60
Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80
Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
85          90          95
Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
100         105         110
Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
115         120         125
Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
130         135         140
Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val
145         150         155         160
Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu
165         170         175
Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu
180         185         190
Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile
195         200         205
Gln Cys Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile
210         215         220
Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn
225         230         235         240
Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro
245         250         255
Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr
260         265         270

```



-continued

---

Gly Asp Ala Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln Leu His  
 275 280 285

Ala Ala Leu Asn Gln Ile Ala Ala Glu Gly Arg Gly Val Leu Leu Tyr  
 290 295 300

Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu Lys Ala  
 305 310 315 320

Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn Glu Ala  
 325 330 335

Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala Gln Ile  
 340 345 350

Leu Arg Asp Leu Gly Val Arg Asn Met Lys Leu Leu Thr Asn Asn Pro  
 355 360 365

Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser Glu Arg  
 370 375 380

Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr Leu Gln  
 385 390 395 400

Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
 405 410

<210> SEQ ID NO 9  
 <211> LENGTH: 1239  
 <212> TYPE: DNA  
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 9

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa 60  
 gaagcactgg acgctttaa aaaaggcgaa gtcatcatcg ttgtagatga tgaagacaga 120  
 gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt 180  
 atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg 240  
 cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta 300  
 agcatagacc atcgtgaaac gaagacaggt atcagcgcgc aagaaagatc ttttaccggt 360  
 caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt 420  
 cactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt 480  
 gatcttgctg aagcttggtg atctccagga gccggcgtca tttgtgaaat tatgaatgaa 540  
 gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa 600  
 atgatcacca ttaaggattt gattcaatac cgttacaatc tgacaacact tgtcgagcgt 660  
 gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720  
 gaggtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccgtt cggagaagaa 780  
 ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgttttg ctctcatcgc 840  
 tgtgattgcg gaccgcagct gcacgccgcg ctgaacccaa ttgccgcaga aggccgtgga 900  
 gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct 960  
 tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg 1020  
 ccggatcttc gcaactatgg catcggagca caaatcttac gcgacctcgg tgtccggaat 1080  
 ataaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140  
 atttcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200  
 accaaaatga acaagctagg tcatttactt catttctaa 1239



-continued

---

```

<210> SEQ ID NO 10
<211> LENGTH: 412
<212> TYPE: PRT
<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 10

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1          5          10          15
His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
20          25          30
Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
35          40          45
Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
50          55          60
Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80
Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
85          90          95
Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
100         105         110
Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
115         120         125
Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
130         135         140
Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val
145         150         155         160
Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu
165         170         175
Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu
180         185         190
Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile
195         200         205
Gln Tyr Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile
210         215         220
Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn
225         230         235         240
Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro
245         250         255
Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr
260         265         270
Gly Asp Val Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln Leu His
275         280         285
Ala Ala Leu Asn Gln Ile Ala Ala Glu Gly Arg Gly Val Leu Leu Tyr
290         295         300
Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu Lys Ala
305         310         315         320
Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn Glu Ala
325         330         335
Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala Gln Ile
340         345         350
Leu Arg Asp Leu Gly Val Arg Asn Ile Lys Leu Leu Thr Asn Asn Pro
355         360         365
Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser Glu Arg
370         375         380

```



-continued

Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr Leu Gln  
385 390 395 400

Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
405 410

<210> SEQ ID NO 11

<211> LENGTH: 1239

<212> TYPE: DNA

<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 11

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa 60  
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga 120  
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt 180  
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg 240  
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta 300  
agcatagacc atcgtgaaac gaagacaggt atcagcgctc aagaaagatc ttttaccggt 360  
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt 420  
ccactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt 480  
gatcttgctg aagcttggtg atctccagga gccggcgtca tttgtgaaat tatgaatgaa 540  
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa 600  
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgtcagcgt 660  
gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720  
gagtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccgtt cggagaagaa 780  
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgtttgg ctctcatcgc 840  
tgtgattgcg gaccgcagct gcacgccacg ctgaaccaa ttgccacaga aggccgtgga 900  
gtgctcctgt acttgcgcca agaaggacga ggcacgggt taatcaataa attaaaagct 960  
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg 1020  
ccggatcttc gcaactatgg catcggagca caaattttac gcgacctcgg tgtccggaat 1080  
atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140  
atctcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200  
acaaaaatga acaagctagg tcatttactt catttctaa 1239

<210> SEQ ID NO 12

<211> LENGTH: 412

<212> TYPE: PRT

<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 12

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe  
1 5 10 15

His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile  
20 25 30

Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala  
35 40 45

Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His  
50 55 60

Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg  
65 70 75 80



-continued

---

Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr  
85 90 95

Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser  
100 105 110

Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser  
115 120 125

Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala  
130 135 140

Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val  
145 150 155 160

Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu  
165 170 175

Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu  
180 185 190

Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile  
195 200 205

Gln Cys Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile  
210 215 220

Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn  
225 230 235 240

Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro  
245 250 255

Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr  
260 265 270

Gly Asp Val Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln Leu His  
275 280 285

Ala Thr Leu Asn Gln Ile Ala Thr Glu Gly Arg Gly Val Leu Leu Tyr  
290 295 300

Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu Lys Ala  
305 310 315 320

Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn Glu Ala  
325 330 335

Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala Gln Ile  
340 345 350

Leu Arg Asp Leu Gly Val Arg Asn Met Lys Leu Leu Thr Asn Asn Pro  
355 360 365

Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser Glu Arg  
370 375 380

Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr Leu Gln  
385 390 395 400

Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
405 410

&lt;210&gt; SEQ ID NO 13

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 13

atgagaggat ctcacatca ccatcaccat gggatcgatc atatgtttca tccgatagaa 60

gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga 120

gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt 180

atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg 240

cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta 300



-continued

---

```

agcatagacc atcgtgaaac gaagacaggt atcagcgcctc aagaaagatc ttttaccggt 360
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt 420
cactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt 480
gatcttgctg aagcttgtgg atctccagga gccggcgtca tttgtgaaat tatgaatgaa 540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa 600
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgtcgagcgt 660
gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720
gaggtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccgtt cggagaagaa 780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgtttgg ctctcatcgc 840
tgtgattgcg gaccgcagct gcacgccacg ctgaacccaa ttgccacaga aggccgtgga 900
gtgctcctgt acttgcgcca agaaggacga ggcacgggtt taatcaataa attaaaagct 960
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgtt tggattcttg 1020
ccggatcttc gcaactatgg catcggagca caaattttac gcgacctcgg tgtccggaat 1080
ataaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140
atctcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200
accaaaatga acaagctagg tcatttactt catttctaa 1239

```

&lt;210&gt; SEQ ID NO 14

&lt;211&gt; LENGTH: 412

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 14

```

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1           5           10          15

His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
20          25          30

Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
35          40          45

Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
50          55          60

Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80

Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
85          90          95

Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
100         105         110

Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
115         120         125

Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
130         135         140

Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val
145         150         155         160

Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu
165         170         175

Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu
180         185         190

Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile
195         200         205

```



-continued

Gln	Cys	Arg	Tyr	Asn	Leu	Thr	Thr	Leu	Val	Glu	Arg	Glu	Val	Asp	Ile
210						215					220				
Thr	Leu	Pro	Thr	Asp	Phe	Gly	Thr	Phe	Lys	Val	Tyr	Gly	Tyr	Thr	Asn
225					230					235					240
Glu	Val	Asp	Gly	Lys	Glu	His	Val	Ala	Phe	Val	Met	Gly	Asp	Val	Pro
				245					250					255	
Phe	Gly	Glu	Glu	Pro	Val	Leu	Val	Arg	Val	His	Ser	Glu	Cys	Leu	Thr
			260					265					270		
Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys	Gly	Pro	Gln	Leu	His
		275					280					285			
Ala	Thr	Leu	Asn	Gln	Ile	Ala	Thr	Glu	Gly	Arg	Gly	Val	Leu	Leu	Tyr
		290				295					300				
Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Ile	Asn	Lys	Leu	Lys	Ala
305					310					315					320
Tyr	Lys	Leu	Gln	Glu	Gln	Gly	Tyr	Asp	Thr	Val	Glu	Ala	Asn	Glu	Ala
			325						330					335	
Leu	Gly	Phe	Leu	Pro	Asp	Leu	Arg	Asn	Tyr	Gly	Ile	Gly	Ala	Gln	Ile
			340					345					350		
Leu	Arg	Asp	Leu	Gly	Val	Arg	Asn	Ile	Lys	Leu	Leu	Thr	Asn	Asn	Pro
		355					360						365		
Arg	Lys	Ile	Ala	Gly	Leu	Glu	Gly	Tyr	Gly	Leu	Ser	Ile	Ser	Glu	Arg
		370				375					380				
Val	Pro	Leu	Gln	Met	Glu	Ala	Lys	Glu	His	Asn	Lys	Lys	Tyr	Leu	Gln
385					390					395					400
Thr	Lys	Met	Asn	Lys	Leu	Gly	His	Leu	Leu	His	Phe				
			405						410						

&lt;210&gt; SEQ ID NO 15

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 15

```

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa      60
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga      120
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt      180
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg      240
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta      300
agcatagacc atcgtgaaac gaagacaggt atcagcgctc aagaaagatc ttttaccggt      360
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt      420
cactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt      480
gatcttgctg aagcttggtg atctccagga gccggcgctc tttgtgaaat tatgaatgaa      540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa      600
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgtcgagcgt      660
gaagttgaca ttacgtgcc tactgatttt gggacattta aggtttatgg atacacaaat      720
gaggtagatg gaaaagagca tgtcgcatth gtgatgggag atgtgccgtt cggagaagaa      780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgtttgg ctctcatcgc      840
tgtgattgcg gaccgcagct gcacgccacg ctgaaccgaa ttgccacaga aggccgtgga      900
gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct      960
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg     1020

```



-continued

---

```

ccggatcttc gcaactatgg catcggagca caaatTTTtac gcgacctcgg tgtccggaat 1080
ataaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140
atttcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200
accaaaatga acaagctagg tcatttactt catttctaa 1239

```

&lt;210&gt; SEQ ID NO 16

&lt;211&gt; LENGTH: 412

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 16

```

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1          5          10          15
His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
20          25          30
Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
35          40          45
Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
50          55          60
Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80
Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
85          90          95
Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
100         105         110
Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
115         120         125
Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
130         135         140
Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val
145         150         155         160
Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu
165         170         175
Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu
180         185         190
Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile
195         200         205
Gln Cys Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile
210         215         220
Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn
225         230         235         240
Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro
245         250         255
Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr
260         265         270
Gly Asp Val Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln Leu His
275         280         285
Ala Thr Leu Asn Arg Ile Ala Thr Glu Gly Arg Gly Val Leu Leu Tyr
290         295         300
Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu Lys Ala
305         310         315         320
Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn Glu Ala
325         330         335

```



-continued

Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala Gln Ile  
 340 345 350  
 Leu Arg Asp Leu Gly Val Arg Asn Ile Lys Leu Leu Thr Asn Asn Pro  
 355 360 365  
 Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser Glu Arg  
 370 375 380  
 Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr Leu Gln  
 385 390 395 400  
 Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
 405 410

<210> SEQ ID NO 17  
 <211> LENGTH: 1239  
 <212> TYPE: DNA  
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 17

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa 60  
 gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga 120  
 gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt 180  
 atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg 240  
 cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta 300  
 agcatagacc atcgtgaaac gaagacaggt atcagcgctc aagaaagatc ttttaccgtt 360  
 caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt 420  
 ccaactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt 480  
 gatcttgctg aagcttggtg atctccagga gccggcgctca tttgtgaaat tatgaatgaa 540  
 gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa 600  
 atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgtcagcgt 660  
 gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720  
 gaggtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccggt cggagaagaa 780  
 ccggatttgg tccgggtgca ttcagaatgt ctcacagggt acgtgtttgg ctctcatcgc 840  
 tgtgattgcg gaccgcagct gcacgccact ctgaaccgaa ttgccacaga aggccgtgga 900  
 gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct 960  
 tataggcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg 1020  
 ccggatcttc gcaactatgg catcggagca caaattttac gcgacctcgg tgtccggaat 1080  
 ataaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140  
 atttcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200  
 accaaaaatga acaagctagg tcatttactt cattttctaa 1239

<210> SEQ ID NO 18  
 <211> LENGTH: 412  
 <212> TYPE: PRT  
 <213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 18

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe  
 1 5 10 15  
 His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile  
 20 25 30

-continued

Ile	Val	Val	Asp	Asp	Glu	Asp	Arg	Glu	Asn	Glu	Gly	Asp	Phe	Val	Ala
		35					40					45			
Leu	Ala	Glu	His	Ala	Thr	Pro	Glu	Val	Ile	Asn	Phe	Met	Ala	Thr	His
	50					55					60				
Gly	Arg	Gly	Leu	Ile	Cys	Thr	Pro	Leu	Ser	Glu	Glu	Ile	Ala	Asp	Arg
65					70					75					80
Leu	Asp	Leu	His	Pro	Met	Val	Glu	His	Asn	Thr	Asp	Ser	His	His	Thr
				85					90					95	
Ala	Phe	Thr	Val	Ser	Ile	Asp	His	Arg	Glu	Thr	Lys	Thr	Gly	Ile	Ser
			100					105					110		
Ala	Gln	Glu	Arg	Ser	Phe	Thr	Val	Gln	Ala	Leu	Leu	Asp	Ser	Lys	Ser
		115					120					125			
Val	Pro	Ser	Asp	Phe	Gln	Arg	Pro	Gly	His	Ile	Phe	Pro	Leu	Ile	Ala
	130					135					140				
Lys	Lys	Gly	Gly	Val	Leu	Lys	Arg	Ala	Gly	His	Thr	Glu	Ala	Ala	Val
145					150					155					160
Asp	Leu	Ala	Glu	Ala	Cys	Gly	Ser	Pro	Gly	Ala	Gly	Val	Ile	Cys	Glu
				165					170					175	
Ile	Met	Asn	Glu	Asp	Gly	Thr	Met	Ala	Arg	Val	Pro	Glu	Leu	Ile	Glu
			180					185					190		
Ile	Ala	Lys	Lys	His	Gln	Leu	Lys	Met	Ile	Thr	Ile	Lys	Asp	Leu	Ile
		195					200					205			
Gln	Cys	Arg	Tyr	Asn	Leu	Thr	Thr	Leu	Val	Glu	Arg	Glu	Val	Asp	Ile
	210					215					220				
Thr	Leu	Pro	Thr	Asp	Phe	Gly	Thr	Phe	Lys	Val	Tyr	Gly	Tyr	Thr	Asn
225					230					235					240
Glu	Val	Asp	Gly	Lys	Glu	His	Val	Ala	Phe	Val	Met	Gly	Asp	Val	Pro
				245					250					255	
Phe	Gly	Glu	Glu	Pro	Val	Leu	Val	Arg	Val	His	Ser	Glu	Cys	Leu	Thr
				260				265					270		
Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys	Gly	Pro	Gln	Leu	His
		275					280					285			
Ala	Thr	Leu	Asn	Arg	Ile	Ala	Thr	Glu	Gly	Arg	Gly	Val	Leu	Leu	Tyr
	290					295					300				
Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Ile	Asn	Lys	Leu	Lys	Ala
305					310					315					320
Tyr	Arg	Leu	Gln	Glu	Gln	Gly	Tyr	Asp	Thr	Val	Glu	Ala	Asn	Glu	Ala
				325					330					335	
Leu	Gly	Phe	Leu	Pro	Asp	Leu	Arg	Asn	Tyr	Gly	Ile	Gly	Ala	Gln	Ile
			340					345					350		
Leu	Arg	Asp	Leu	Gly	Val	Arg	Asn	Ile	Lys	Leu	Leu	Thr	Asn	Asn	Pro
		355					360					365			
Arg	Lys	Ile	Ala	Gly	Leu	Glu	Gly	Tyr	Gly	Leu	Ser	Ile	Ser	Glu	Arg
	370					375					380				
Val	Pro	Leu	Gln	Met	Glu	Ala	Lys	Glu	His	Asn	Lys	Lys	Tyr	Leu	Gln
385					390					395					400
Thr	Lys	Met	Asn	Lys	Leu	Gly	His	Leu	Leu	His	Phe				
				405						410					

&lt;210&gt; SEQ ID NO 19

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis



-continued

&lt;400&gt; SEQUENCE: 19

```

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa    60
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga    120
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt    180
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg    240
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta    300
agcatagacc atcgtgaaac gaagacaggt atcagcgtc aagaaagatc ttttaccggt    360
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt    420
ccactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt    480
gatcttgctg aagcttgagg atctccagga gccggcgtea tttgtgaaat tatgaatgaa    540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa    600
atgatcacca ttaaggattt gattcaatac cgttacaatc tgacaacact tgtcgagcgt    660
gaagttgaca ttacgctgcc tactgatttt gggacatttg aggtttatgg atacacaaat    720
gaggtagatg gaaaagagca tgtcgcatth gtgatgggag atgtgccgtt cggagaagaa    780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgtttgg ctctcatcgc    840
tgtgattgcg caccgcagct gcacgccgcg ctgaaccaa ttgccgcaga aggccgtgga    900
gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct    960
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg   1020
ccggatcttc gcaactatgg catcggagca caaatthtac gcgacctcgg tgtccggaat   1080
atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt   1140
atctcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa   1200
acaaaaatga acaagctagg tcatttactt catttctaa                               1239

```

&lt;210&gt; SEQ ID NO 20

&lt;211&gt; LENGTH: 412

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 20

```

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1           5           10           15

His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
          20           25           30

Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
          35           40           45

Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
          50           55           60

Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65           70           75           80

Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
          85           90           95

Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
          100          105          110

Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
          115          120          125

Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
          130          135          140

```

-continued

Lys	Lys	Gly	Gly	Val	Leu	Lys	Arg	Ala	Gly	His	Thr	Glu	Ala	Ala	Val
145					150					155					160
Asp	Leu	Ala	Glu	Ala	Cys	Gly	Ser	Pro	Gly	Ala	Gly	Val	Ile	Cys	Glu
			165						170					175	
Ile	Met	Asn	Glu	Asp	Gly	Thr	Met	Ala	Arg	Val	Pro	Glu	Leu	Ile	Glu
		180						185					190		
Ile	Ala	Lys	Lys	His	Gln	Leu	Lys	Met	Ile	Thr	Ile	Lys	Asp	Leu	Ile
		195					200					205			
Gln	Tyr	Arg	Tyr	Asn	Leu	Thr	Thr	Leu	Val	Glu	Arg	Glu	Val	Asp	Ile
	210					215					220				
Thr	Leu	Pro	Thr	Asp	Phe	Gly	Thr	Phe	Glu	Val	Tyr	Gly	Tyr	Thr	Asn
225					230					235					240
Glu	Val	Asp	Gly	Lys	Glu	His	Val	Ala	Phe	Val	Met	Gly	Asp	Val	Pro
				245					250					255	
Phe	Gly	Glu	Glu	Pro	Val	Leu	Val	Arg	Val	His	Ser	Glu	Cys	Leu	Thr
			260					265					270		
Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys	Ala	Pro	Gln	Leu	His
		275					280					285			
Ala	Ala	Leu	Asn	Gln	Ile	Ala	Ala	Glu	Gly	Arg	Gly	Val	Leu	Leu	Tyr
	290					295					300				
Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Ile	Asn	Lys	Leu	Lys	Ala
305					310					315					320
Tyr	Lys	Leu	Gln	Glu	Gln	Gly	Tyr	Asp	Thr	Val	Glu	Ala	Asn	Glu	Ala
				325					330					335	
Leu	Gly	Phe	Leu	Pro	Asp	Leu	Arg	Asn	Tyr	Gly	Ile	Gly	Ala	Gln	Ile
			340					345					350		
Leu	Arg	Asp	Leu	Gly	Val	Arg	Asn	Met	Lys	Leu	Leu	Thr	Asn	Asn	Pro
		355					360					365			
Arg	Lys	Ile	Ala	Gly	Leu	Glu	Gly	Tyr	Gly	Leu	Ser	Ile	Ser	Glu	Arg
	370					375					380				
Val	Pro	Leu	Gln	Met	Glu	Ala	Lys	Glu	His	Asn	Lys	Lys	Tyr	Leu	Gln
385					390					395					400
Thr	Lys	Met	Asn	Lys	Leu	Gly	His	Leu	Leu	His	Phe				
				405						410					

&lt;210&gt; SEQ ID NO 21

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 21

atgagaggat ctcacatca ccatcacat gggatcgatc atatgtttca tccgatagaa	60
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga	120
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt	180
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg	240
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta	300
agcatagacc atcgtgaaac gaagacaggt atcagcgctc aagaaagatc ttttaccggt	360
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt	420
ccactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgtctgt	480
gatcttgctg aagcttgctg atctccagga gccggcgctca tttgtgaaat tatgaatgaa	540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa	600
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgctcgagcgt	660



-continued

---

```

gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720
gaggtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccggt cggagaagaa 780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgtttg ctctcatcgc 840
tgtgattgcg gaccgcagct gcacgccacg ctgaaccgaa ttgccacaga aggccgtgga 900
gtgctcctgt acttgcgcca agaaggacga ggcacgggtt taatcaataa attaaaagct 960
tataggcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggataactg 1020
ccggatcttc gcaactatgg catcggagca caaatTTTtac gcgacctcgg tgtccggaat 1080
ataaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140
atttcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200
acccaaatga acaagctagg tcatttactt catttctaa 1239

```

&lt;210&gt; SEQ ID NO 22

&lt;211&gt; LENGTH: 412

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 22

```

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1          5          10          15
His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
20          25          30
Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
35          40          45
Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
50          55          60
Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80
Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
85          90          95
Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
100         105         110
Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
115         120         125
Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
130         135         140
Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val
145         150         155         160
Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu
165         170         175
Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu
180         185         190
Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile
195         200         205
Gln Cys Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile
210         215         220
Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn
225         230         235         240
Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro
245         250         255
Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr
260         265         270

```

-continued

Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys	Gly	Pro	Gln	Leu	His
		275					280					285			
Ala	Thr	Leu	Asn	Arg	Ile	Ala	Thr	Glu	Gly	Arg	Gly	Val	Leu	Leu	Tyr
		290				295					300				
Leu	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Ile	Asn	Lys	Leu	Lys	Ala
305					310					315					320
Tyr	Arg	Leu	Gln	Glu	Gln	Gly	Tyr	Asp	Thr	Val	Glu	Ala	Asn	Glu	Ala
				325					330					335	
Leu	Gly	Tyr	Leu	Pro	Asp	Leu	Arg	Asn	Tyr	Gly	Ile	Gly	Ala	Gln	Ile
			340					345					350		
Leu	Arg	Asp	Leu	Gly	Val	Arg	Asn	Ile	Lys	Leu	Leu	Thr	Asn	Asn	Pro
		355					360					365			
Arg	Lys	Ile	Ala	Gly	Leu	Glu	Gly	Tyr	Gly	Leu	Ser	Ile	Ser	Glu	Arg
		370				375					380				
Val	Pro	Leu	Gln	Met	Glu	Ala	Lys	Glu	His	Asn	Lys	Lys	Tyr	Leu	Gln
385					390					395					400
Thr	Lys	Met	Asn	Lys	Leu	Gly	His	Leu	Leu	His	Phe				
			405						410						

&lt;210&gt; SEQ ID NO 23

&lt;211&gt; LENGTH: 1239

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus subtilis

&lt;400&gt; SEQUENCE: 23

```

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa    60
gaagcactgg acgctttaa aaaaggcgaa gtcatcatcg ttgtagatga tgaagacaga    120
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt    180
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg    240
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta    300
agcatagacc atcgtgaaac gaagacaggt atcagcgcgc aagaaagatc ttttaccgtt    360
caagcattgc tggacagcaa atccgtgcc tctgattttc agcgtccggg gcacattttt    420
cactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt    480
gatcttgctg aagcttgtgg atctccagga gccggcgctc tttgtgaaat tatgaatgaa    540
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa    600
atgatcacca ttaaggattt gattcaatac cgttacaatc tgacaacact tgtcgagcgt    660
gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat    720
gaggtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccgtt cggagaagaa    780
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgtgttttg ctctcatcgc    840
tgtgattgca gaccgcagct gcacgccgcg ctgaacccaa ttgccgcaga aggccgtgga    900
gtgctcctgt acttgcgcca agaaggacga ggcacggtt taatcaataa attaaaagct    960
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgtt tggattcttg   1020
ccggatcttc gcaactatgg catcggagca caaatcttac gcgacctcgg tgtccggaat   1080
atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt   1140
atctcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa   1200
acaaaaatga acaagctagg tcatttactt catttctaa                               1239

```



-continued

---

```

<210> SEQ ID NO 24
<211> LENGTH: 412
<212> TYPE: PRT
<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 24

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe
1          5          10          15
His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile
20          25          30
Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala
35          40          45
Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His
50          55          60
Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg
65          70          75          80
Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr
85          90          95
Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser
100         105         110
Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser
115         120         125
Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala
130         135         140
Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val
145         150         155         160
Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu
165         170         175
Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu
180         185         190
Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile
195         200         205
Gln Tyr Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile
210         215         220
Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn
225         230         235         240
Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro
245         250         255
Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr
260         265         270
Gly Asp Val Phe Gly Ser His Arg Cys Asp Cys Arg Pro Gln Leu His
275         280         285
Ala Ala Leu Asn Gln Ile Ala Ala Glu Gly Arg Gly Val Leu Leu Tyr
290         295         300
Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu Lys Ala
305         310         315         320
Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn Glu Ala
325         330         335
Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala Gln Ile
340         345         350
Leu Arg Asp Leu Gly Val Arg Asn Met Lys Leu Leu Thr Asn Asn Pro
355         360         365
Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser Glu Arg
370         375         380

```

-continued

---

Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr Leu Gln  
385 390 395 400

Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
405 410

<210> SEQ ID NO 25

<211> LENGTH: 1239

<212> TYPE: DNA

<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 25

atgagaggat ctcaccatca ccatcaccat gggatcgatc atatgtttca tccgatagaa 60  
gaagcactgg acgctttaa aaaaggcgaa gtcacatcg ttgtagatga tgaagacaga 120  
gaaaatgaag gagactttgt ggctcttgcc gagcatgcaa cgccggaagt cattaacttt 180  
atggcgacac atgggagagg actgatctgc acgccgctca gtgaggaaat cgcagacagg 240  
cttgatcttc accctatggt tgagcataat acagactctc accacactgc atttaccgta 300  
agcatagacc atcgtgaaac gaagacaggt atcagcgctc aagaaagatc ttttaccggt 360  
caagcattgc tggacagcaa atccgtgcca tctgattttc agcgtccggg gcacattttt 420  
ccactgattg cgaaaaaagg aggtgtcctg aaaagagcgg gccatacaga agctgctggt 480  
gatcttgctg aagcttggtg atctccagga gccggcgtca tttgtgaaat tatgaatgaa 540  
gacggaacga tggcgagagt gcctgagctc attgaaattg cgaaaaagca tcaattaaaa 600  
atgatcacca ttaaggattt gattcaatgc cgttacaatc tgacaacact tgtcagcgt 660  
gaagttgaca ttacgctgcc tactgatttt gggacattta aggtttatgg atacacaaat 720  
gaggtagatg gaaaagagca tgtcgcattt gtgatgggag atgtgccgtt cggagaagaa 780  
ccggtattgg tccgggtgca ttcagaatgt ctcacaggtg acgcgtttgg ctctcatcgc 840  
tgtgattgcg gaccgcagct gcacgccacg ctgaaccaa ttgccgcaga aggcctgga 900  
gtgctcctgt acttgcgcca agaaggacga ggcacgggtt taatcaataa attaaaagct 960  
tataagcttc aggaacaagg ctatgacacc gtagaagcca atgaggcgct tggattcttg 1020  
ccggatcttc gcaactatgg catcggagca caaattttac gcgacctcgg tgtccggaat 1080  
atgaagcttt tgacgaataa tccgcgaaaa atcgcaggcc ttgaaggcta cggactcagt 1140  
atctcagaaa gagtgccgct tcaaatggag gcgaaagaac acaataaaaa atatttgcaa 1200  
acaaaaatga acaagctagg tcatttactt catttctaa 1239

<210> SEQ ID NO 26

<211> LENGTH: 412

<212> TYPE: PRT

<213> ORGANISM: Bacillus subtilis

<400> SEQUENCE: 26

Met Arg Gly Ser His His His His His His Gly Ile Asp His Met Phe  
1 5 10 15

His Pro Ile Glu Glu Ala Leu Asp Ala Leu Lys Lys Gly Glu Val Ile  
20 25 30

Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe Val Ala  
35 40 45

Leu Ala Glu His Ala Thr Pro Glu Val Ile Asn Phe Met Ala Thr His  
50 55 60

Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Glu Ile Ala Asp Arg  
65 70 75 80



-continued

---

Leu Asp Leu His Pro Met Val Glu His Asn Thr Asp Ser His His Thr  
 85 90 95

Ala Phe Thr Val Ser Ile Asp His Arg Glu Thr Lys Thr Gly Ile Ser  
 100 105 110

Ala Gln Glu Arg Ser Phe Thr Val Gln Ala Leu Leu Asp Ser Lys Ser  
 115 120 125

Val Pro Ser Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu Ile Ala  
 130 135 140

Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala Ala Val  
 145 150 155 160

Asp Leu Ala Glu Ala Cys Gly Ser Pro Gly Ala Gly Val Ile Cys Glu  
 165 170 175

Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Leu Ile Glu  
 180 185 190

Ile Ala Lys Lys His Gln Leu Lys Met Ile Thr Ile Lys Asp Leu Ile  
 195 200 205

Gln Cys Arg Tyr Asn Leu Thr Thr Leu Val Glu Arg Glu Val Asp Ile  
 210 215 220

Thr Leu Pro Thr Asp Phe Gly Thr Phe Lys Val Tyr Gly Tyr Thr Asn  
 225 230 235 240

Glu Val Asp Gly Lys Glu His Val Ala Phe Val Met Gly Asp Val Pro  
 245 250 255

Phe Gly Glu Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu Thr  
 260 265 270

Gly Asp Ala Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln Leu His  
 275 280 285

Ala Thr Leu Asn Gln Ile Ala Ala Glu Gly Arg Gly Val Leu Leu Tyr  
 290 295 300

Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu Lys Ala  
 305 310 315 320

Tyr Lys Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn Glu Ala  
 325 330 335

Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala Gln Ile  
 340 345 350

Leu Arg Asp Leu Gly Val Arg Asn Met Lys Leu Leu Thr Asn Asn Pro  
 355 360 365

Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser Glu Arg  
 370 375 380

Val Pro Leu Gln Met Glu Ala Lys Glu His Asn Lys Lys Tyr Leu Gln  
 385 390 395 400

Thr Lys Met Asn Lys Leu Gly His Leu Leu His Phe  
 405 410

<210> SEQ ID NO 27  
 <211> LENGTH: 61  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer RibA 1S

<400> SEQUENCE: 27

atgagaggat ctcaccatca ccataccat gggatcgatc atatgtttca tccgatagaa 60

g 61

-continued

---

<210> SEQ ID NO 28  
 <211> LENGTH: 38  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer RibA 1AS  
  
 <400> SEQUENCE: 28  
  
 tataattgga tccttagaaa tgaagtaa at gacctagc 38

<210> SEQ ID NO 29  
 <211> LENGTH: 54  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer RibA 2S  
  
 <400> SEQUENCE: 29  
  
 attaatgaat tcattaaaga ggagaaatta actatgagag gatctcacca tcac 54

<210> SEQ ID NO 30  
 <211> LENGTH: 31  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer RibANde+1  
  
 <400> SEQUENCE: 30  
  
 ggagggtttc atatgtttca tccgatagaa g 31

<210> SEQ ID NO 31  
 <211> LENGTH: 21  
 <212> TYPE: DNA  
 <213> ORGANISM: Artificial  
 <220> FEATURE:  
 <223> OTHER INFORMATION: Primer RibA4AS  
  
 <400> SEQUENCE: 31  
  
 taattaagct tggatcctta g 21

<210> SEQ ID NO 32  
 <211> LENGTH: 906  
 <212> TYPE: DNA  
 <213> ORGANISM: Ashbya gossypii  
  
 <400> SEQUENCE: 32  
  
 atgactgaat acacagtgcc agaagtgagg tgtgtcgcac gcgcgcgcat accgacggta 60  
 cagggcaccg atgtcttctt ccatctatac cacaactoga tcgacagcaa ggaacaccta 120  
 gcgattgtct tggcgagaa catacgctcg cggagtctgt tccggtagcg gaaagacgac 180  
 acgcagcagg cgcgatggt gcgggcgcc tacgtgggac agctgtacc cgggaggacc 240  
 gaggcagacg cggatcggcg tcagggcctg gagctgcggt ttgatgagac agggcagctg 300  
 gtggtggagc gggcgacgac gtggaccagg gagccgacac tgggtcggct gcaactcggag 360  
 tgttacacgg gcgagacggc gtggagcgcg cgggtgcgact gcggggagca gttcgaccag 420  
 gcgggtaagc tgatggctgc ggcgacagag ggcgaggtgg ttggcggctg ggggcacggc 480  
 gtgatcgtgt acctgcggca ggagggccgc ggcacgggc taggcgagaa gctgaaggcg 540  
 tacaacctgc aggacctggg cgcggacacg gtgcaggcga acgagctgct caaccacct 600  
 gcggacgcgc ggcacttctc gttggggcgc gcaatcctac tggacctcg tatcgaggac 660  
 atccggttgc tcacgaataa ccccgacaag gtgcagcagg tgcactgtcc gccggcgcta 720  
 cgctgcatcg agcgggtgcc catggtgccg ctttcatgga ctgagccac acagggcgtg 780



-continued

---

```

cgctcgcgcg agctggacgg ctacctgccc gccaaaggctg agcgcatggg gcacatgctg 840
cagcgccgcg tgggtgctgca cacgtctgcg gcggccgagc tccccgcgc caacacacac 900
atataa 906

```

```

<210> SEQ ID NO 33
<211> LENGTH: 301
<212> TYPE: PRT
<213> ORGANISM: Ashbya gossypii

```

```

<400> SEQUENCE: 33

```

```

Met Thr Glu Tyr Thr Val Pro Glu Val Arg Cys Val Ala Arg Ala Arg
1           5           10           15

Ile Pro Thr Val Gln Gly Thr Asp Val Phe Leu His Leu Tyr His Asn
          20           25           30

Ser Ile Asp Ser Lys Glu His Leu Ala Ile Val Phe Gly Glu Asn Ile
          35           40           45

Arg Ser Arg Ser Leu Phe Arg Tyr Arg Lys Asp Asp Thr Gln Gln Ala
          50           55           60

Arg Met Val Arg Gly Ala Tyr Val Gly Gln Leu Tyr Pro Gly Arg Thr
65           70           75           80

Glu Ala Asp Ala Asp Arg Arg Gln Gly Leu Glu Leu Arg Phe Asp Glu
          85           90           95

Thr Gly Gln Leu Val Val Glu Arg Ala Thr Thr Trp Thr Arg Glu Pro
          100          105          110

Thr Leu Val Arg Leu His Ser Glu Cys Tyr Thr Gly Glu Thr Ala Trp
          115          120          125

Ser Ala Arg Cys Asp Cys Gly Glu Gln Phe Asp Gln Ala Gly Lys Leu
          130          135          140

Met Ala Ala Ala Thr Glu Gly Glu Val Val Gly Gly Ala Gly His Gly
          145          150          155          160

Val Ile Val Tyr Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Gly Glu
          165          170          175

Lys Leu Lys Ala Tyr Asn Leu Gln Asp Leu Gly Ala Asp Thr Val Gln
          180          185          190

Ala Asn Glu Leu Leu Asn His Pro Ala Asp Ala Arg Asp Phe Ser Leu
          195          200          205

Gly Arg Ala Ile Leu Leu Asp Leu Gly Ile Glu Asp Ile Arg Leu Leu
          210          215          220

Thr Asn Asn Pro Asp Lys Val Gln Gln Val His Cys Pro Pro Ala Leu
          225          230          235          240

Arg Cys Ile Glu Arg Val Pro Met Val Pro Leu Ser Trp Thr Gln Pro
          245          250          255

Thr Gln Gly Val Arg Ser Arg Glu Leu Asp Gly Tyr Leu Arg Ala Lys
          260          265          270

Val Glu Arg Met Gly His Met Leu Gln Arg Pro Leu Val Leu His Thr
          275          280          285

Ser Ala Ala Ala Glu Leu Pro Arg Ala Asn Thr His Ile
          290          295          300

```

```

<210> SEQ ID NO 34
<211> LENGTH: 591
<212> TYPE: DNA
<213> ORGANISM: Escherichia coli

```

-continued

&lt;400&gt; SEQUENCE: 34

```

atgcagctta aacgtgtggc agaagccaaa ctgccaaacc catggggcga tttcctgatg    60
gtgggatttg aagaactggc aaccggacac gatcatgtcg cgctagtcta tggcgatatt    120
tccgggcata ccccggtact tgcgcgcgtc cattccgaat gtctgaccgg tgacgccttg    180
ttcagcttgc gctgcgattg tggttccag ctccaagcgg cattgacgca aattgccgag    240
gaaggccttg gtattttgct gtatcacctg caggaaggtc gtaacattgg tctgctgaat    300
aaaatccgcg cttacgcact gcaggatcaa ggttacgata ccgtagaggc taaccaccag    360
ttaggcttcg ccgctgatga gcgcgacttc actctttgcy ctgatatgtt caaactcctt    420
ggcgtcaatg aagtcgctt gttaaccaat aaccgaaaa aagtcgaaat tctgaccgaa    480
gcagggatta atattgttga acgcgtacca ttgattgtag gtcgtaacc caataacgaa    540
cattatctcg ataccaaagc cgagaaaatg ggccatttgc tgaacaaata a          591

```

&lt;210&gt; SEQ ID NO 35

&lt;211&gt; LENGTH: 196

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Escherichia coli

&lt;400&gt; SEQUENCE: 35

```

Met Gln Leu Lys Arg Val Ala Glu Ala Lys Leu Pro Thr Pro Trp Gly
1           5           10           15
Asp Phe Leu Met Val Gly Phe Glu Glu Leu Ala Thr Gly His Asp His
          20           25           30
Val Ala Leu Val Tyr Gly Asp Ile Ser Gly His Thr Pro Val Leu Ala
          35           40           45
Arg Val His Ser Glu Cys Leu Thr Gly Asp Ala Leu Phe Ser Leu Arg
          50           55           60
Cys Asp Cys Gly Phe Gln Leu Glu Ala Ala Leu Thr Gln Ile Ala Glu
          65           70           75           80
Glu Gly Arg Gly Ile Leu Leu Tyr His Arg Gln Glu Gly Arg Asn Ile
          85           90           95
Gly Leu Leu Asn Lys Ile Arg Ala Tyr Ala Leu Gln Asp Gln Gly Tyr
          100          105          110
Asp Thr Val Glu Ala Asn His Gln Leu Gly Phe Ala Ala Asp Glu Arg
          115          120          125
Asp Phe Thr Leu Cys Ala Asp Met Phe Lys Leu Leu Gly Val Asn Glu
          130          135          140
Val Arg Leu Leu Thr Asn Asn Pro Lys Lys Val Glu Ile Leu Thr Glu
          145          150          155          160
Ala Gly Ile Asn Ile Val Glu Arg Val Pro Leu Ile Val Gly Arg Asn
          165          170          175
Pro Asn Asn Glu His Tyr Leu Asp Thr Lys Ala Glu Lys Met Gly His
          180          185          190
Leu Leu Asn Lys
          195

```

&lt;210&gt; SEQ ID NO 36

&lt;211&gt; LENGTH: 1269

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Corynebacterium glutamicum

&lt;400&gt; SEQUENCE: 36

```

gtgagtgaac atgagcaggc acacagccaa ttagattctg ttgaagaggc catcgctgac    60
atcgctgagg gtaaagccgt cgtggtgta gatgatgaag atcgtgaaaa tgaaggcgac    120

```



-continued

```

atcatctttg cgcgcaatt agccactcca gaattagtcg ctttcatggt gcgttattcc 180
tcgggataca tctgtgcgcc attaaccgca aaggatgcag atcgtettga tctgcctccg 240
atgaccgcgc acaatcagga tgcccgcggc accgcttaca ccgtgaccgt tgatgccaac 300
accggcacca caggcatttc tgcaacagac cgcgcccaca ctttgcgctt gcttgctgat 360
ccagaagccg accgcacgga tttcacccgt cccggacacg ttgtgccact gcgtgctcgt 420
gaaggtggcg tcttggtgcg cgctggacac accgaagcag ctgtcgattt ggctcgcgct 480
gcaggcctgc gcccagcagg tgttatctgc gaagtggcca gtgaagagga ccccaccggc 540
atggctcggg ttctgagct gcgccccttc tgcgatgagc acgatctgaa gctgatctct 600
attgagcagc tcattgagtg gcgtcgcaag aatgaaattt tggaggagcg ccaggaggaa 660
actgtgctgc ctaccgattt cggcacgttc aaggctggtg gttaccgttc catcatcgat 720
ggcaccgagc ttggtgccat tgttgccggc gacgtggcat ccgacggtgg cgaaaacgtc 780
ctggttcgag tccactctga gtgcttgact ggtgatggtt ttggatcccg gcgctgcgac 840
tgtggacagc agctgcacga gtctttgccc ctgatccagg aagctggtcg gggagtagtg 900
gtgtacatgc gtgggcatga gggacgaggc attggtctgc tcgccaagct acgcgcctac 960
caactccagg atgaaggtgc cgacaccgtc gatgccaaac tcgcaactgg tcttccagcc 1020
gatgcccgcg aatttggcac cagcgcaccg attctctacg acttgggtgt gcgctcgtcc 1080
aacttgatca gcaacaacc agccaagaag gtgggacttg aaggccacgg catttccatt 1140
gccagccgaa ccccatccc tgttgctggt catgaagaca atgttcgata cctgaaaacc 1200
aagcgtgacc gcatgggaca tgacctcca gatgtcgcac tgtgggaaca agagcaccca 1260
gaaaactaa 1269

```

&lt;210&gt; SEQ ID NO 37

&lt;211&gt; LENGTH: 422

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Corynebacterium glutamicum

&lt;400&gt; SEQUENCE: 37

```

Val Ser Glu His Glu Gln Ala His Ser Gln Leu Asp Ser Val Glu Glu
1          5          10          15
Ala Ile Ala Asp Ile Ala Ala Gly Lys Ala Val Val Val Val Asp Asp
20          25          30
Glu Asp Arg Glu Asn Glu Gly Asp Ile Ile Phe Ala Ala Glu Leu Ala
35          40          45
Thr Pro Glu Leu Val Ala Phe Met Val Arg Tyr Ser Ser Gly Tyr Ile
50          55          60
Cys Ala Pro Leu Thr Ala Lys Asp Ala Asp Arg Leu Asp Leu Pro Pro
65          70          75          80
Met Thr Ala His Asn Gln Asp Ala Arg Gly Thr Ala Tyr Thr Val Thr
85          90          95
Val Asp Ala Asn Thr Gly Thr Thr Gly Ile Ser Ala Thr Asp Arg Ala
100         105         110
His Thr Leu Arg Leu Leu Ala Asp Pro Glu Ala Asp Arg Thr Asp Phe
115        120        125
Thr Arg Pro Gly His Val Val Pro Leu Arg Ala Arg Glu Gly Gly Val
130        135        140
Leu Val Arg Ala Gly His Thr Glu Ala Ala Val Asp Leu Ala Arg Ala
145        150        155        160
Ala Gly Leu Arg Pro Ala Gly Val Ile Cys Glu Val Val Ser Glu Glu
165        170        175

```

-continued

Asp	Pro	Thr	Gly	Met	Ala	Arg	Val	Pro	Glu	Leu	Arg	Arg	Phe	Cys	Asp
			180					185					190		
Glu	His	Asp	Leu	Lys	Leu	Ile	Ser	Ile	Glu	Gln	Leu	Ile	Glu	Trp	Arg
		195					200					205			
Arg	Lys	Asn	Glu	Ile	Leu	Val	Glu	Arg	Gln	Val	Glu	Thr	Val	Leu	Pro
	210					215					220				
Thr	Asp	Phe	Gly	Thr	Phe	Lys	Ala	Val	Gly	Tyr	Arg	Ser	Ile	Ile	Asp
225					230					235					240
Gly	Thr	Glu	Leu	Val	Ala	Ile	Val	Ala	Gly	Asp	Val	Ala	Ser	Asp	Gly
				245					250					255	
Gly	Glu	Asn	Val	Leu	Val	Arg	Val	His	Ser	Glu	Cys	Leu	Thr	Gly	Asp
		260						265					270		
Val	Phe	Gly	Ser	Arg	Arg	Cys	Asp	Cys	Gly	Gln	Gln	Leu	His	Glu	Ser
		275					280					285			
Leu	Arg	Leu	Ile	Gln	Glu	Ala	Gly	Arg	Gly	Val	Val	Val	Tyr	Met	Arg
	290					295						300			
Gly	His	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Leu	Ala	Lys	Leu	Arg	Ala	Tyr
305					310					315					320
Gln	Leu	Gln	Asp	Glu	Gly	Ala	Asp	Thr	Val	Asp	Ala	Asn	Leu	Ala	Leu
			325						330					335	
Gly	Leu	Pro	Ala	Asp	Ala	Arg	Glu	Phe	Gly	Thr	Ser	Ala	Gln	Ile	Leu
			340					345					350		
Tyr	Asp	Leu	Gly	Val	Arg	Ser	Leu	Asn	Leu	Ile	Ser	Asn	Asn	Pro	Ala
	355						360					365			
Lys	Lys	Val	Gly	Leu	Glu	Gly	His	Gly	Ile	Ser	Ile	Ala	Ser	Arg	Thr
	370					375					380				
Pro	Ile	Pro	Val	Ala	Val	His	Glu	Asp	Asn	Val	Arg	Tyr	Leu	Lys	Thr
385					390					395					400
Lys	Arg	Asp	Arg	Met	Gly	His	Asp	Leu	Pro	Asp	Val	Ala	Leu	Trp	Glu
			405						410					415	
Gln	Glu	His	Pro	Glu	Asn										
			420												

&lt;210&gt; SEQ ID NO 38

&lt;211&gt; LENGTH: 1197

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus amyloliquefaciens

&lt;400&gt; SEQUENCE: 38

atgtttcatc cgatagaaga ggcattagaa gcgctgaaaa aaggtgaagt catcatcggt	60
gtc gatgatg aagacagaga aaacgaagga gatttcgtag cgctcgctga gcatgctacg	120
cctgaagtgg tgaattttat ggcgaccac gggagaggcc tgatctgcac gccgctttct	180
gaagacatcg ccggccggct ggatcttcat ccaatggctg atcataatac agactcgcat	240
gagaccgcgt ttacagtcag cattgaccac aagctgacaa aaacgggaat cagcgctcag	300
gaacgttctt ttacgattca ggcgcttttg gacgaagaat ctgtgectgg cgattttcag	360
cgccggggtc atatttttcc cttaatagca aaaaaaggag gcgtcctgaa gcgggagggc	420
cacacggaag cagccgttga cctggcaaaa gcatgcggtt ctcaaggagc ggacgtcatt	480
tgtgaaatta tgaatgaaga cggcacaatg gcgagagtgc ctgagattag cgagattgcg	540
aaaagccacc agctgaaaat gattacgata aaagacttaa tagaataccg ctacaacatt	600
acaacacttg tgaacagaga agttgacatt acgctgccga ctgacttcgg cacgttccgg	660
gtttacggat atacaaacga ggtggacgga aaagaacatc tcgcctttgt catgggcat	720



-continued

---

```

gtcccgttta acagcggacc cgttcttgtc agagtgcact cagaatgcct gaccggcgat 780
gtgtttgcat cccaccgctg tgattgctgg cctcagcttc atgccgcggt gcgccaaatt 840
gccgaagaag gccgcggcgt tctattgtat ttgcgtcagg aaggcagagg aatcgggtctc 900
atcaataagc tgaaagcgta tcgattgcag gaacaagggt acgacacggt tgaagcgaac 960
gaagcgctcg gctttctgcc tgacttgctc aactatggca tcggcgccca gattctccgc 1020
gatttagggg ttcagcatat gaaactttta accaataacc cccggaaaat cgccggcctt 1080
gaagggtagc gactaagcat ttcagatcgg gtgccgcttc aaatggaagc gagtgagcac 1140
aacaagcagt atttacaac caaatgaaa aaactcggac acttgcttca tttctaa 1197

```

&lt;210&gt; SEQ ID NO 39

&lt;211&gt; LENGTH: 398

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus amyloliquefaciens

&lt;400&gt; SEQUENCE: 39

```

Met Phe His Pro Ile Glu Glu Ala Leu Glu Ala Leu Lys Lys Gly Glu
1           5           10          15
Val Ile Ile Val Val Asp Asp Glu Asp Arg Glu Asn Glu Gly Asp Phe
          20          25          30
Val Ala Leu Ala Glu His Ala Thr Pro Glu Val Val Asn Phe Met Ala
          35          40          45
Thr His Gly Arg Gly Leu Ile Cys Thr Pro Leu Ser Glu Asp Ile Ala
          50          55          60
Gly Arg Leu Asp Leu His Pro Met Val Asp His Asn Thr Asp Ser His
65          70          75          80
Glu Thr Ala Phe Thr Val Ser Ile Asp His Lys Leu Thr Lys Thr Gly
          85          90          95
Ile Ser Ala Gln Glu Arg Ser Phe Thr Ile Gln Ala Leu Leu Asp Glu
          100         105         110
Glu Ser Val Pro Gly Asp Phe Gln Arg Pro Gly His Ile Phe Pro Leu
          115         120         125
Ile Ala Lys Lys Gly Gly Val Leu Lys Arg Ala Gly His Thr Glu Ala
130         135         140
Ala Val Asp Leu Ala Lys Ala Cys Gly Ser Gln Gly Ala Asp Val Ile
145         150         155         160
Cys Glu Ile Met Asn Glu Asp Gly Thr Met Ala Arg Val Pro Glu Ile
          165         170         175
Ser Glu Ile Ala Lys Ser His Gln Leu Lys Met Ile Thr Ile Lys Asp
          180         185         190
Leu Ile Glu Tyr Arg Tyr Asn Ile Thr Thr Leu Val Asn Arg Glu Val
          195         200         205
Asp Ile Thr Leu Pro Thr Asp Phe Gly Thr Phe Arg Val Tyr Gly Tyr
210         215         220
Thr Asn Glu Val Asp Gly Lys Glu His Leu Ala Phe Val Met Gly Asp
225         230         235         240
Val Pro Phe Asn Ser Gly Pro Val Leu Val Arg Val His Ser Glu Cys
          245         250         255
Leu Thr Gly Asp Val Phe Ala Ser His Arg Cys Asp Cys Gly Pro Gln
          260         265         270
Leu His Ala Ala Leu Arg Gln Ile Ala Glu Glu Gly Arg Gly Val Leu
          275         280         285
Leu Tyr Leu Arg Gln Glu Gly Arg Gly Ile Gly Leu Ile Asn Lys Leu
          290         295         300

```

-continued

---

Lys Ala Tyr Arg Leu Gln Glu Gln Gly Tyr Asp Thr Val Glu Ala Asn  
 305 310 315 320

Glu Ala Leu Gly Phe Leu Pro Asp Leu Arg Asn Tyr Gly Ile Gly Ala  
 325 330 335

Gln Ile Leu Arg Asp Leu Gly Val Gln His Met Lys Leu Leu Thr Asn  
 340 345 350

Asn Pro Arg Lys Ile Ala Gly Leu Glu Gly Tyr Gly Leu Ser Ile Ser  
 355 360 365

Asp Arg Val Pro Leu Gln Met Glu Ala Ser Glu His Asn Lys Gln Tyr  
 370 375 380

Leu Gln Thr Lys Met Lys Lys Leu Gly His Leu Leu His Phe  
 385 390 395

<210> SEQ ID NO 40  
 <211> LENGTH: 1194  
 <212> TYPE: DNA  
 <213> ORGANISM: Bacillus cereus

<400> SEQUENCE: 40

```

atgtttcatc gtattgaaga agctctagaa gatttaaaaa aaggtaaagt cgttatcgta    60
tgtgatgatg aaaaccgaga aaatgaaggc gattttattg ctttagcaga gtacattaca    120
ccagaaacaa taaattttat gattacacat ggccgtggtc tcgtttgtgt accgattacg    180
gaaggatcag cagaacgtct acaattagaa ccaatgggat ctcataatac agattcacat    240
catactgcgt ttacagtgag cattgacat gtctctacaa caacagggat tagcgctcac    300
gaacgtgcaa ctacgatata agaattgtta aaccccgcat caaaagggtgc tgatttcaat    360
cgacctggac atatctttcc attaattgag aaagaaggcg gtgtcctgcg tcgtgcaggt    420
catacagaag ctgctgttga tttagcaaag ctatgcggtg ccgaaccagc tggagttatt    480
tgcgagatta taaatgagga cggcacgatg gcacgtgtac ctgatttaat agaatgcgca    540
aaacaatttg atataaaaat gattacaata gaagatttaa ttgcttaccg ccgccatcat    600
gaaacacttg tgacgagaga agcggaaatt acattaccta cagatttcgg tactttccac    660
gcaattggct attctaactc attagatcag aaagaacata tcgcacttgt aaaagggtgat    720
atctcaacag gtgaaccggt acttgtagct gttcattctg aatgcttaac aggagatgta    780
ttcggttcac atcgctgca ttgcggacca caactccatg cagcacttgc tcaaattgag    840
cgtgaaggaa aagggtgttct tctttatatg aggcaagaag gaagaggcat tgggcttctt    900
aataagcttc gtgcttataa attacaagaa gaaggattcg atactgtaga agcaaatgaa    960
aaactcggct tcctgctga tcttcgtgat tacggtatcg gtgctcaaat attaaaagat   1020
ttaggtttac agagtttacg attattaacg aataacccaa gaaaaattgc tggcttacia   1080
ggttacgatt tagaagtagt cgagcgtgta ccgttgcaaa tgccagcaaa agaagagaat   1140
aatcgtatt tacaacgaa agtaaacaaa ttaggacact tactaaactt ataa           1194

```

<210> SEQ ID NO 41  
 <211> LENGTH: 397  
 <212> TYPE: PRT  
 <213> ORGANISM: Bacillus cereus

<400> SEQUENCE: 41

Met Phe His Arg Ile Glu Glu Ala Leu Glu Asp Leu Lys Lys Gly Lys  
 1 5 10 15

Val Val Ile Val Cys Asp Asp Glu Asn Arg Glu Asn Glu Gly Asp Phe  
 20 25 30



-continued

Ile Ala Leu Ala Glu Tyr Ile Thr Pro Glu Thr Ile Asn Phe Met Ile  
 35 40 45  
 Thr His Gly Arg Gly Leu Val Cys Val Pro Ile Thr Glu Gly Tyr Ala  
 50 55 60  
 Glu Arg Leu Gln Leu Glu Pro Met Val Ser His Asn Thr Asp Ser His  
 65 70 75 80  
 His Thr Ala Phe Thr Val Ser Ile Asp His Val Ser Thr Thr Thr Gly  
 85 90 95  
 Ile Ser Ala His Glu Arg Ala Thr Thr Ile Gln Glu Leu Leu Asn Pro  
 100 105 110  
 Ala Ser Lys Gly Ala Asp Phe Asn Arg Pro Gly His Ile Phe Pro Leu  
 115 120 125  
 Ile Ala Lys Glu Gly Gly Val Leu Arg Arg Ala Gly His Thr Glu Ala  
 130 135 140  
 Ala Val Asp Leu Ala Lys Leu Cys Gly Ala Glu Pro Ala Gly Val Ile  
 145 150 155 160  
 Cys Glu Ile Ile Asn Glu Asp Gly Thr Met Ala Arg Val Pro Asp Leu  
 165 170 175  
 Ile Glu Cys Ala Lys Gln Phe Asp Ile Lys Met Ile Thr Ile Glu Asp  
 180 185 190  
 Leu Ile Ala Tyr Arg Arg His His Glu Thr Leu Val Thr Arg Glu Ala  
 195 200 205  
 Glu Ile Thr Leu Pro Thr Asp Phe Gly Thr Phe His Ala Ile Gly Tyr  
 210 215 220  
 Ser Asn Ser Leu Asp Thr Lys Glu His Ile Ala Leu Val Lys Gly Asp  
 225 230 235 240  
 Ile Ser Thr Gly Glu Pro Val Leu Val Arg Val His Ser Glu Cys Leu  
 245 250 255  
 Thr Gly Asp Val Phe Gly Ser His Arg Cys Asp Cys Gly Pro Gln Leu  
 260 265 270  
 His Ala Ala Leu Ala Gln Ile Glu Arg Glu Gly Lys Gly Val Leu Leu  
 275 280 285  
 Tyr Met Arg Gln Glu Gly Arg Gly Ile Gly Leu Leu Asn Lys Leu Arg  
 290 295 300  
 Ala Tyr Lys Leu Gln Glu Glu Gly Phe Asp Thr Val Glu Ala Asn Glu  
 305 310 315 320  
 Lys Leu Gly Phe Pro Ala Asp Leu Arg Asp Tyr Gly Ile Gly Ala Gln  
 325 330 335  
 Ile Leu Lys Asp Leu Gly Leu Gln Ser Leu Arg Leu Leu Thr Asn Asn  
 340 345 350  
 Pro Arg Lys Ile Ala Gly Leu Gln Gly Tyr Asp Leu Glu Val Val Glu  
 355 360 365  
 Arg Val Pro Leu Gln Met Pro Ala Lys Glu Glu Asn Lys Ser Tyr Leu  
 370 375 380  
 Gln Thr Lys Val Asn Lys Leu Gly His Leu Leu Asn Leu  
 385 390 395

&lt;210&gt; SEQ ID NO 42

&lt;211&gt; LENGTH: 1215

&lt;212&gt; TYPE: DNA

&lt;213&gt; ORGANISM: Bacillus halodurans

&lt;400&gt; SEQUENCE: 42

atggacaaaa agctatttga tccgattgaa gaagcaatat atgaattaat gcaaggtcga 60

gtcgtgatcg tttgtgatga tgaggatcgg gaaaacgaag gggattttgt agcccttgct 120

-continued

---

```

gaaaaagcaa caccagaagt gattaacttc atgatcacgc atggccgtgg tctcgtttgc 180
acgccaatca cggaagagcg ggcaaaggaa ttagatcttg tccccatggt ggaccataat 240
accgatcccc atggtacggc gtttaccgtc agcattgatc atcaaatgac gaccacagga 300
atctctgccc atgaacgggc tatgacgatt caggcgtaa ttgataagaa aacgaaaaag 360
caccacttca aacgaccagg tcacattttc ccctaataag cgaaaaacgg aggagtactc 420
cgacggggcg gtcatacaga agcggccggt gatctagctc gtttgtcagg cgctgagccg 480
gcaggggtta tttgtgaaat cattaagaa gatggttcaa tggcacgagt tcttgatttg 540
cgaaaaatcg ccgatcagtt tgaactgaag atgatcacia ttaaagattt aatcgaatat 600
cgtcaccgta aagacaagct tgtcaagcgt gaagtagata tttccttacc gacggatttc 660
ggctcattcc gtgcaatcgg ttatacagat gtcattgatg gaaaagagag tgctcgctta 720
gtgaaaggac agattgttga aggtgaacca aactcgttc gtgttcactc cgaatgttta 780
acaggtgatg tgctcggttc tcaccgttgc gattgtggcc cacaactcca ggcagctctc 840
acacaaatcg agcaacaagg caaaggata ctctttata tgcgtcaaga gggctcgtgg 900
atcggcttca tgaataagtt gaaggcatac aagcttcaag aagaaggcta tgatactgta 960
gaagcaaatg agaaattagg ctttctctgct gatcttcggg actatggaat gggcgcgcaa 1020
atcttacgcg acttaggtgt gtcaaaaatg cgctcctta caaacaatcc gcgaaaaatt 1080
acgggcttga aagggtatgg ccttgaagtg gttgaacggg tgccgctcca attacctcat 1140
aacaagata atgagcgcta tttgaaaaca aagcacgaaa agttaggaca tctgctaaat 1200
tttactcatt cgtaa 1215

```

&lt;210&gt; SEQ ID NO 43

&lt;211&gt; LENGTH: 404

&lt;212&gt; TYPE: PRT

&lt;213&gt; ORGANISM: Bacillus halodurans

&lt;400&gt; SEQUENCE: 43

```

Met Asp Lys Lys Leu Phe Asp Pro Ile Glu Glu Ala Ile Tyr Glu Leu
1           5           10           15
Met Gln Gly Arg Val Val Ile Val Cys Asp Asp Glu Asp Arg Glu Asn
          20           25           30
Glu Gly Asp Phe Val Ala Leu Ala Glu Lys Ala Thr Pro Glu Val Ile
          35           40           45
Asn Phe Met Ile Thr His Gly Arg Gly Leu Val Cys Thr Pro Ile Thr
          50           55           60
Glu Glu Arg Ala Lys Glu Leu Asp Leu Val Pro Met Val Asp His Asn
65           70           75           80
Thr Asp Pro His Gly Thr Ala Phe Thr Val Ser Ile Asp His Gln Met
          85           90           95
Thr Thr Thr Gly Ile Ser Ala His Glu Arg Ala Met Thr Ile Gln Ala
          100          105          110
Leu Ile Asp Lys Lys Thr Lys Lys His His Phe Lys Arg Pro Gly His
          115          120          125
Ile Phe Pro Leu Ile Ala Lys Asn Gly Gly Val Leu Arg Arg Ala Gly
          130          135          140
His Thr Glu Ala Ala Val Asp Leu Ala Arg Leu Ser Gly Ala Glu Pro
145          150          155          160
Ala Gly Val Ile Cys Glu Ile Ile Lys Glu Asp Gly Ser Met Ala Arg
          165          170          175

```



-continued

---

Val	Pro	Asp	Leu	Arg	Lys	Ile	Ala	Asp	Gln	Phe	Glu	Leu	Lys	Met	Ile
			180					185					190		
Thr	Ile	Lys	Asp	Leu	Ile	Glu	Tyr	Arg	His	Arg	Lys	Asp	Lys	Leu	Val
		195					200					205			
Lys	Arg	Glu	Val	Asp	Ile	Ser	Leu	Pro	Thr	Asp	Phe	Gly	Ser	Phe	Arg
	210					215					220				
Ala	Ile	Gly	Tyr	Thr	Asp	Val	Ile	Asp	Gly	Lys	Glu	Ser	Val	Ala	Leu
225					230					235					240
Val	Lys	Gly	Gln	Ile	Val	Glu	Gly	Glu	Pro	Thr	Leu	Val	Arg	Val	His
				245					250					255	
Ser	Glu	Cys	Leu	Thr	Gly	Asp	Val	Phe	Gly	Ser	His	Arg	Cys	Asp	Cys
			260					265					270		
Gly	Pro	Gln	Leu	Gln	Ala	Ala	Leu	Thr	Gln	Ile	Glu	Gln	Gln	Gly	Lys
		275					280					285			
Gly	Ile	Leu	Leu	Tyr	Met	Arg	Gln	Glu	Gly	Arg	Gly	Ile	Gly	Leu	Met
	290					295					300				
Asn	Lys	Leu	Lys	Ala	Tyr	Lys	Leu	Gln	Glu	Glu	Gly	Tyr	Asp	Thr	Val
305					310					315					320
Glu	Ala	Asn	Glu	Lys	Leu	Gly	Phe	Pro	Ala	Asp	Leu	Arg	Asp	Tyr	Gly
				325					330					335	
Met	Gly	Ala	Gln	Ile	Leu	Arg	Asp	Leu	Gly	Val	Ser	Lys	Met	Arg	Leu
			340					345						350	
Leu	Thr	Asn	Asn	Pro	Arg	Lys	Ile	Thr	Gly	Leu	Lys	Gly	Tyr	Gly	Leu
		355				360						365			
Glu	Val	Val	Glu	Arg	Val	Pro	Leu	Gln	Leu	Pro	His	Asn	Lys	Asp	Asn
	370					375					380				
Glu	Arg	Tyr	Leu	Lys	Thr	Lys	His	Glu	Lys	Leu	Gly	His	Leu	Leu	Asn
385					390					395					400
Phe	Thr	His	Ser												

---

The invention claimed is:

1. A method for producing riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof comprising:

culturing a host cell in a suitable medium, the host cell comprising a polynucleotide comprising a nucleotide sequence which codes for a modified GTP cyclohydrolase II from *Bacillus subtilis*, wherein (i) the specific activity of the modified enzyme is increased in comparison to the corresponding non-modified enzyme and (ii) the amino acid sequence of the modified enzyme comprises one or more mutations(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308, and/or 347 of SEQ ID NO: 2 and combinations of the one or more mutations(s) thereof such that said modified enzyme includes from 1 to 6 position substitutions selected from: Alanine at an amino acid position corresponding to the position 261, Alanine or Arginine at an amino acid position corresponding to the position 270, Threonine at an amino acid position corresponding to the position 276, Arginine at an amino acid position corresponding to the position 279, Arginine at an amino acid position corresponding to the position 308, and Isoleucine at an amino acid position corresponding to the position 347; said modified enzyme exhibiting at least 98% identity to SEQ ID NO: 2, and

optionally separating riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof from the medium.

2. A method for increasing the production of riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof by

introducing into a host cell a modified GTP cyclohydrolase II from *Bacillus subtilis*, wherein

(i) the specific activity of the modified enzyme is increased in comparison to the corresponding non-modified enzyme, and

(ii) the amino acid sequence of the modified enzyme comprises one or more mutations(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308, and/or 347 of SEQ ID NO: 2 and combinations of the one or more mutation(s) thereof such that said modified enzyme includes from 1 to 6 position substitutions selected from: Alanine at an amino acid position corresponding to the position 261, Alanine or Arginine at an amino acid position corresponding to the position 270, Threonine at an amino acid position corresponding to the position 276, Arginine at an amino acid position corresponding to the position 279, Arginine at an amino acid position corresponding to the position 308, and a sixth mutation comprises an Isoleucine at an amino acid position corresponding to the position 347, said modified enzyme exhibiting at least 98% identity to SEQ ID NO: 2.

3. A method for increasing the production of riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof by introducing into a host cell a polynucleotide comprising a nucleotide sequence which codes for a modified GTP cyclohydrolase II from *Bacillus subtilis*, wherein

- (i) the specific activity of the modified enzyme is increased in comparison to the corresponding non-modified enzyme, and
- (ii) the amino acid sequence of the modified enzyme comprises one or more mutations(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308, and/or 347 of SEQ ID NO: 2 and combinations of the one or more mutation(s) thereof such that said modified enzyme includes from 1 to 6 position substitutions selected from: Alanine at an amino acid position corresponding to the position 261, Alanine or Arginine at an amino acid position corresponding to the position 270, Threonine at an amino acid position corresponding to the position 276, Arginine at an amino acid position corresponding to the position 279, Arginine at an amino acid position corresponding to the position 308, and a sixth mutation comprises an Isoleucine at an amino acid position corresponding to the position 347, said modified enzyme exhibiting at least 98% identity to SEQ ID NO: 2.
4. A method for producing riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof comprising: culturing a host cell in a suitable medium, the host cell comprising a polynucleotide comprising a nucleotide

sequence which codes for a modified GTP cyclohydrolase II from *Bacillus subtilis*, wherein (i) the specific activity of the modified enzyme is increased in comparison to the corresponding non-modified enzyme and (ii) the amino acid sequence of the modified enzyme consisting of one or more mutations(s) on amino acid position(s) corresponding to positions 261, 270, 276, 279, 308, and/or 347 of SEQ ID NO: 2 and combinations of the one or more mutations(s) thereof such that said modified enzyme includes from 1 to 6 position substitutions selected from: Alanine at an amino acid position corresponding to the position 261; Alanine or Arginine at an amino acid position corresponding to the position 270; Threonine at an amino acid position corresponding to the position 276; Arginine at an amino acid position corresponding to the position 279; Arginine at an amino acid position corresponding to the position 308, and Isoleucine at an amino acid position corresponding to the position 347; said modified enzyme exhibiting at least 98% identity to SEQ ID NO: 2, and optionally separating riboflavin, a riboflavin precursor, FMN, FAD, or a derivative thereof from the medium.

\* \* \* \* \*