



US008022028B2

(12) **United States Patent**
Szewczyk et al.

(10) **Patent No.:** **US 8,022,028 B2**
(45) **Date of Patent:** **Sep. 20, 2011**

(54) **LIGHT DUTY LIQUID CLEANING
COMPOSITIONS AND METHODS OF
MANUFACTURE AND USE THEREOF
COMPRISING ORGANIC ACIDS**

(75) Inventors: **Gregory Szewczyk**, Flemington, NJ
(US); **Caryn R. Westphal**, East
Brunswick, NJ (US); **Karen
Wisniewski**, Bridgewater, NJ (US);
Cynthia McCullar Murphy, Belle
Meade, NJ (US); **David Frank Suriano**,
Edison, NJ (US)

(73) Assignee: **Colgate-Palmolive Company**, New
York, NY (US)

(*) Notice: Subject to any disclaimer, the term of this
patent is extended or adjusted under 35
U.S.C. 154(b) by 0 days.

(21) Appl. No.: **12/140,754**

(22) Filed: **Jun. 17, 2008**

(65) **Prior Publication Data**

US 2009/0312225 A1 Dec. 17, 2009

(51) **Int. Cl.**
C11D 17/00 (2006.01)

(52) **U.S. Cl.** **510/424; 510/425; 510/426; 510/428;**
510/490; 510/506

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,658,985	A	4/1972	Olson, Jr.
3,676,340	A	7/1972	Berg et al.
3,769,398	A	10/1973	Hewitt
3,935,129	A	1/1976	Jabalee
3,982,892	A	9/1976	Gray
4,013,787	A	3/1977	Varlerberghe et al.
4,129,515	A	12/1978	Foster
4,154,706	A	5/1979	Kenkare et al.
4,224,195	A	9/1980	Kawasaki et al.
4,259,204	A	3/1981	Homma
4,329,334	A	5/1982	Su et al.
4,329,335	A	5/1982	Su et al.
4,329,336	A	5/1982	Su et al.
4,450,091	A	5/1984	Schmolka
4,595,526	A	6/1986	Lai
5,130,056	A	7/1992	Jakobson et al.
5,192,460	A	3/1993	Thomas et al.
5,294,364	A	3/1994	Thomas et al.
5,437,809	A	8/1995	Chaudhuri
5,531,938	A	7/1996	Erilli
5,554,320	A	9/1996	Yianakopoulos
5,741,769	A	4/1998	Erilli
5,914,300	A	6/1999	Fujiwara et al.
5,962,388	A	10/1999	Sherry et al.
5,968,539	A	10/1999	Beerse et al.
6,010,992	A	1/2000	Arvanitidou et al.
6,046,146	A	4/2000	Erilli
6,046,150	A	4/2000	Choy et al.
6,048,834	A	4/2000	Drapier et al.
6,051,542	A	4/2000	Pollack et al.
6,071,866	A	6/2000	Fujiwara et al.
6,121,228	A	9/2000	Drapier et al.
6,136,769	A	10/2000	Asano et al.

6,147,039	A	11/2000	Jacques et al.
6,156,296	A	12/2000	Riedel et al.
6,159,925	A	12/2000	Blandiaux
6,251,844	B1	6/2001	Leonard et al.
6,255,269	B1	7/2001	Leonard et al.
6,262,003	B1	7/2001	Leonard et al.
6,268,330	B1	7/2001	Leonard et al.
6,291,419	B1	9/2001	D'Ambrogio et al.
6,306,817	B1	10/2001	Kott et al.
6,313,084	B1	11/2001	D'Ambrogio et al.
6,339,055	B1	1/2002	Brooker et al.
6,361,768	B1	3/2002	Galleguillos et al.
6,441,037	B1	8/2002	Arvanitidou et al.
6,444,636	B1	9/2002	Toussaint et al.
6,475,967	B1	11/2002	Arvanitidou et al.
6,495,507	B1	12/2002	Arvanitidou
6,534,472	B1	3/2003	Arvanitidou et al.
6,537,952	B2	3/2003	Hayward et al.
6,537,955	B1	3/2003	Raso et al.
6,541,436	B1	4/2003	Arvanitidou et al.
6,551,980	B1	4/2003	Wisniewski et al.
6,583,178	B2	6/2003	D'Ambrogio et al.
6,586,014	B2	7/2003	Arvanitidou et al.
6,593,284	B2	7/2003	Arvanitidou et al.
6,605,579	B1	8/2003	Arvanitidou et al.
6,617,296	B1	9/2003	Connors et al.
6,627,589	B1	9/2003	Arvanitidou
6,696,399	B1	2/2004	Chernin et al.
6,733,986	B1	5/2004	Armstrong et al.
6,815,406	B1	11/2004	Szewczyk
6,821,939	B1	11/2004	Szewczyk et al.
6,884,764	B2	4/2005	Szewczyk et al.
6,914,038	B2	7/2005	Terazaki et al.
7,033,986	B2	4/2006	Szewczyk
7,470,653	B2	12/2008	Gambogi et al.
2003/0144218	A1	7/2003	Arvanitidou et al.
2003/0147822	A1	8/2003	Doi et al.
2003/0170197	A1	9/2003	Terazaki et al.
2004/0101504	A1	5/2004	Kinscherf et al.
2004/0204331	A1	10/2004	Connors et al.
2004/0229763	A1	11/2004	Hutton, III et al.
2004/0229766	A1	11/2004	Hutton, III et al.
2004/0229767	A1	11/2004	Ford et al.
2004/0229963	A1	11/2004	Culeron et al.
2004/0254253	A1	12/2004	Culeron et al.
2005/0003990	A1*	1/2005	Smith et al. 510/421
2005/0019161	A1	1/2005	Ichikawa et al.
2005/0031570	A1	2/2005	Grit et al.
2005/0049161	A1	3/2005	Szewczyk
2005/0049170	A1	3/2005	Szewczyk
2005/0192196	A1	9/2005	Hutton, III et al.
2006/0115440	A1	6/2006	Arata et al.
2006/0198807	A1	9/2006	Morioka

FOREIGN PATENT DOCUMENTS

EP 0 250 181 A2 6/1987

(Continued)

OTHER PUBLICATIONS

PAIC Anti-bacterial washing up liquid <http://www.gnpd.com>.

(Continued)

Primary Examiner — Necholus Ogden, Jr.

(74) *Attorney, Agent, or Firm* — Michael F. Morgan

(57) **ABSTRACT**

The invention encompasses liquid cleaning compositions, for example, dish washing liquids, and methods of their manufacture and use, which possess enhanced cleaning ability. The cleaning compositions of the invention include acidic light duty liquid cleaning compositions with low toxicity and anti-bacterial efficacy on surfaces, for example, hard surfaces.

13 Claims, No Drawings

FOREIGN PATENT DOCUMENTS

WO 94/11476 A1 5/1994
WO 00/66079 A1 11/2000
WO 01/79404 A2 10/2001
WO 02/092743 A1 11/2002
WO 03/097779 A1 11/2003

OTHER PUBLICATIONS

File History U.S. Appl. No. 12/140,795, filed Jun. 17, 2008.
File History U.S. Appl. No. 12/140,754, filed Jun. 17, 2008.
File History U.S. Appl. No. 12/140,806, filed Jun. 17, 2008.

File History PCT Serial No. PCT/US2008/067219, Jun. 17, 2008.
File History PCT Serial No. PCT/US2008/067220, Jun. 17, 2008.
File History PCT Serial No. PCT/US2008/067228, Jun. 17, 2008.
International Search Report for PCT/US2009/047595, Jun. 17, 2009.
International Search Report for PCT/US2009/047604, Jun. 17, 2009.
International Search Report for PCT/US2009/047601, Jun. 17, 2009.
International Search Report for PCT/US2008/067219, Jun. 17, 2008.
International Search Report for PCT/US2008/067220, Jun. 17, 2008.
International Search Report for PCT/US2008/067228, Jun. 17, 2008.
File history of U.S. Appl. No. 12/808,376, filed Jun. 16, 2010.

* cited by examiner

1

**LIGHT DUTY LIQUID CLEANING
COMPOSITIONS AND METHODS OF
MANUFACTURE AND USE THEREOF
COMPRISING ORGANIC ACIDS**

FIELD OF THE INVENTION

The invention encompasses liquid cleaning compositions, for example, dish washing liquids, and methods of their manufacture and use, which possess enhanced cleaning ability. The cleaning compositions of the invention include acidic light duty liquid cleaning compositions with low toxicity and antibacterial efficacy on surfaces, for example, hard surfaces.

BACKGROUND OF THE INVENTION

Light duty liquid cleaning compositions should be designed with acceptable foaming and cleaning properties. Such cleaning compositions should maintain acceptable cleaning performance, have ease of rinsing, and contain a low level of dye mix that yields a near colorless visual appearance. Light duty liquid cleaning compositions should include an ingredient mix that increases utilization of naturally derived ingredients, results in a cleaned surface with minimal spotting and is both mild and hypoallergenic. Light duty liquid cleaning compositions should also be designed to be biodegradable and not to leave any harmful residue on surfaces.

Accordingly, the inventors of the invention have developed light duty liquid cleaning compositions, which are suitable for disinfecting all types of surfaces including animate surfaces (e.g., human skin and/or mouth when used as an oral preparation or toothpaste) and inanimate surfaces. This technology is suitable for use on delicate surfaces including those surfaces in contact with food in a safe manner. Moreover, the light duty liquid cleaning compositions according to the invention reduce the amount of chemical residues left on a surface disinfected therewith. Thus, it may be not necessary to rinse, for example, a surface after the compositions of the invention have been applied thereto in diluted conditions. The inventors have developed compositions and methods that include cleaning compositions with enhanced cleaning possessing antibacterial efficacy and low toxicity.

BRIEF SUMMARY OF THE INVENTION

The invention encompasses acidic liquid cleaning compositions designed for cleaning surfaces including hard surfaces, which deliver acceptable cleaning and foaming performance and exhibit ease of rinsing while leaving low amounts residue.

The inventors have surprisingly found that cleaning compositions including a combination of anionic surfactants, a zwitterionic surfactant and an acid in specific amounts have antibacterial activity while at the same time having low toxicity. In certain embodiments, the cleaning composition is a colorless liquid.

In one embodiment the invention encompasses cleaning compositions including an acidic formulation that exhibits ease of rinsing, which assists with the removal of residue while exhibiting antibacterial efficacy.

In other embodiments, the invention encompasses cleaning compositions including a surfactant based cleaning composition comprising at least one anionic surfactant, at least one zwitterionic surfactant, and at least one organic acid or salt thereof, wherein the composition has a \log_{10} reduction in microbes of at least about 3 when a surface containing bacteria is contacted with the composition for about 30 seconds

2

at 25° C., wherein the composition is stable for at least about 1 year at room temperature, and wherein the composition has a low toxicity.

In certain embodiments, the invention encompasses a cleaning composition including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 3 wt. % to about 20 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 2 wt. % to about 20 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 1 wt. % to about 8 wt. % by weight of the total composition, and lactic acid or a salt thereof, wherein the lactic acid or salt thereof is present in an amount of about 1 wt. % to about 2.5 wt. % by weight of the total composition.

In certain embodiments, the invention encompasses compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 20 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 18 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 2 wt. % to about 8 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition.

In certain embodiments, the invention encompasses compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 12 wt. % to about 18 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 14 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 2 wt. % to about 6 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In certain embodiments, the invention encompasses compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 9 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 9 wt. % to about 20 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 1 wt. % to about 7 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In certain embodiments, the invention encompasses compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 2 wt. % to about 5 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 8 wt. % by weight of the total composition, at least one zwitterionic

3

surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In certain embodiments, the invention encompasses an acidic liquid cleaning composition designed for cleaning hard surfaces as well as glass surfaces and effective in removing grease soil and/or other soil.

In other embodiments, the invention encompasses cleaning compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 12.5 wt. % by weight of the total composition a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 13.5 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 5.5 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 to about 2.5 wt. % by weight of the total composition.

In other embodiments, the invention encompasses cleaning compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 5 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 18 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 2.5 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In other embodiments, the invention encompasses cleaning compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 8.5 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 18 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 5.5 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In another embodiment, the invention encompasses cleaning compositions including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 5.7 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 19.5 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 6.5 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

Another embodiment of the invention encompasses cleaning compositions including dodecyl benzene sulfonate or salt thereof is present in an amount of about 5 wt. % to about 20 wt. % by weight of the total composition, lauryl ether sulfate with about two EO units or salt thereof is present in an amount

4

of about 5 wt. % to about 15 wt. % by weight of the total composition, a betaine or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 3 wt. % to about 8 wt. % by weight of the total composition, and lactic acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition.

Another embodiment of the invention encompasses a method of making a liquid cleaning composition designed for cleaning surfaces including hard surfaces and effective in removing soil, which includes combining a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 20 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 15 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 3 wt. % to about 8 wt. % by weight of the total composition, and at least one acid or a salt thereof wherein the acid or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition.

Another embodiment of the invention encompasses a method of removing soil and bacteria, which includes contacting the surface with a composition including a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 3 wt. % to about 20 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 5 wt. % to about 18 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 2 wt. % to about 8 wt. % by weight of the total composition, and at least one acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition.

To achieve the foregoing and other embodiments and in accordance with the purpose of the invention, as embodied and broadly described herein the light duty liquid detergent of this invention includes at least one first anionic surfactant, at least one second anionic surfactant, at least one zwitterionic surfactant, and at least one acidic component, which has both good disinfecting properties on hard surfaces and good food soil and/or other soil removal and leaves surfaces with a shiny appearance.

The compositions have utility in a broad range of applications including, for example, in consumer product fluids such as surface cleaners, cleansers and the like.

DETAILED DESCRIPTION OF THE INVENTION

As used throughout, ranges are used as a shorthand for describing each and every value that is within the range. Any value within the range can be selected as the terminus of the range. In addition, all references cited herein are hereby incorporated by reference in their entireties. In the event of a conflict in a definition in the present disclosure and that of a cited reference, the present disclosure controls.

General Description

The cleaning compositions of the invention are useful as ultra and regular density dish liquid formulas designed for several key formula characteristics including, but not limited to, antibacterial efficacy from a naturally-derived organic acid, at minimum about a 3-log reduction in about 30 seconds

for both Gram-positive (e.g., *Staphylococcus aureus*) and Gram-negative (e.g., *Salmoicella enterica*, *E. coli*) on surfaces, minimal toxicity of inert (non-antibacterial) cleaning materials in the formulation, minimize corrosivity to processing equipment, competitive or superior foaming/cleaning performance with existing commercial cleaning products, competitive or superior rinsing and/or shine performance with existing products, and delivery of both active and aesthetic product stability performance over product lifetime.

Accordingly, the invention encompasses cleaning compositions including a surfactant based cleaning composition comprising at least one anionic surfactant, at least one zwitterionic surfactant, and at least one organic acid or salt thereof, wherein the composition has a \log_{10} reduction in bacteria of at least about 3 when a surface containing bacteria is contacted with the composition for about 30 seconds at 25 °C., wherein the composition is stable for at least about 1 year at room temperature, and wherein the composition has a low toxicity.

In certain embodiments, the organic acid is lactic acid.

In certain embodiments, the at least one anionic surfactant is present in an amount of about 3 wt. % to about 20 wt. % by weight of the total composition.

In certain embodiments, the at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 1 wt. % to about 8 wt. % by weight of the total composition.

In certain embodiments, the at least one organic acid or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition.

In certain embodiments, the anionic surfactant is a C_{10} - C_{14} LA sulfonate.

In certain embodiments, the anionic surfactant is sodium lauryl ether sulfate with about two ethylene oxide units.

In certain embodiments, the zwitterionic surfactant is laurylamidopropyl betaine.

In certain embodiments, the anionic surfactants and/or the zwitterionic surfactants are derived from a natural source and biodegradable surfactants.

In other embodiments, the invention encompasses a cleaning composition comprising a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 8 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 9 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 1.5 wt. % by weight of the total composition, and lactic acid or a salt thereof, wherein the lactic acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In other embodiments, the invention encompasses a cleaning composition comprising a first anionic surfactant or a salt thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 4 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 12 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 3 wt. % by weight of the total composition, and lactic acid or a salt thereof, wherein the lactic acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In other embodiments, the invention encompasses a cleaning composition comprising a first anionic surfactant or a salt

thereof wherein the first anionic surfactant or salt thereof is present in an amount of about 8 wt. % by weight of the total composition, a second anionic surfactant or a salt thereof, wherein the second anionic surfactant or salt thereof is present in an amount of about 18 wt. % by weight of the total composition, at least one zwitterionic surfactant or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 5.5 wt. % by weight of the total composition, and lactic acid or a salt thereof, wherein the lactic acid or salt thereof is present in an amount of about 2 wt. % by weight of the total composition.

In other embodiments, the invention encompasses a cleaning composition comprising dodecyl benzene sulfonate or salt thereof is present in an amount of about 3 wt. % to about 20 wt. % by weight of the total composition, lauryl ether sulfate with about two EO units or salt thereof is present in an amount of about 3 wt. % to about 20 wt. % by weight of the total composition, laurylamidopropyl betaine or a salt thereof, wherein the zwitterionic surfactant or salt thereof is present in an amount of about 1 wt. % to about 8 wt. % by weight of the total composition, and lactic acid or a salt thereof, wherein the acid or salt thereof is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition.

The invention also encompasses methods of cleaning a surface including contacting the surface with a composition of the invention, diluted or undiluted. The cleaning compositions possess antibacterial efficacy from an acid, for example lactic acid or a salt thereof. In certain embodiments, the acid is a naturally-derived, weak-organic acid manufactured from renewable plant resources via microbial fermentation. In other embodiments, the acid is natural and readily biodegradable, non-toxic to the environment, and a natural product. In other embodiments, the surfactant is natural and readily biodegradable, non-toxic to the environment, and a natural product.

As used herein the phrase “from a natural source” refers to surfactants that have a natural origin and are derived from, for example, crops, animal fats and/or trees. These are also referred to in the art as oleochemical surfactants and are derived from sources including but not limited to plant oils such as palm, palm kernel or coconut oil, or from animal fats such as tallow, lard or fish oil. This is in contrast to petroleum or petrochemical surfactants derived from, for example, crude oil.

As used herein, the term “biodegradable surfactants” refers to surfactant-based cleaning ingredients that are designed to be used with water and disposed of down the drain. There they combine with other wastes for treatment in either a municipal treatment plant or a household septic tank system. During treatment, microorganisms biodegrade surfactants and other organic materials, ultimately breaking them down into carbon dioxide, water and minerals. Any small amount of surfactants that remain after treatment continue to biodegrade in the environment. In certain embodiments, the surfactants of the invention biodegrade quickly and thoroughly and do not present a risk to organisms living in the environment.

The cleaning compositions of the invention, diluted or undiluted, result in a minimal 3-log reduction in about 30 seconds or about one minute of both Gram-positive (e.g., *Staphylococcus aureus*) and Gram-negative (e.g., *Salmonella enterica*, *E. coli*) bacteria, or run-off solutions. Without being limited by theory, the inventors believe that the cleaning compositions of the invention, which include an acid result in the acid crossing the bacterial cell membrane in its protonated or charge-neutral form. Lactic acid with a pKa of about 3.8 (the point at which half of the molecules are protonated and

half are not protonated) is effective at a pH below 3.5. In certain embodiments, the recommended pH for the cleaning compositions of the invention for maximal efficacy balanced against safety is about 3.25. Without being limited by theory, the mechanism of action for lactic acid is thought to be two-fold: (1) as protonated molecules cross the bacterial membrane they become deprotonated at the internal pH of the cell and progressively lower the internal bacterial cell pH that can lead to protein deformation and halt critical cellular processes, but (2) this change in internal pH can act to collapse the delta psi gradients critical to microbial nutrient and energy transport systems in the bacterial cell membrane—also leading to a cut-off of critical nutrients and energy sources.

Anionic Surfactants

In certain embodiments, the compositions of the invention include one or more anionic surfactants. The anionic surfactants, which may be used in the compositions of the invention include water soluble anionic sulfonate surfactants and include, but are not limited to, sodium, potassium, ammonium, magnesium and ethanolanionium salts of linear C_8 - C_{16} alkyl benzene sulfonates; C_{10} - C_{20} paraffin sulfonates, alpha olefin sulfonates containing about 10 to about 24 carbon atoms and C_8 - C_{18} alkyl sulfates and mixtures thereof.

The anionic surfactant may be any of the anionic surfactants known or previously used in the art of aqueous surfactant compositions. Suitable anionic surfactants include, but are not limited to, alkyl sulfates, alkyl ether sulfates, alkaryl sulfonates, alkyl succinates, alkyl sulfosuccinates, N-alkoyl sarcosinates, alkyl phosphates, alkyl ether phosphates, alkyl ether carboxylates, alkylamino acids, alkyl peptides, alkoyl taurates, carboxylic acids, acyl and alkyl glutamates, alkyl isethionates, and alpha-olefin sulfonates, especially their sodium, potassium, magnesium, ammonium and mono-, di- and triethanolamine salts. The alkyl groups generally contain about 8 to about 18 carbon atoms and may be unsaturated.

In certain embodiments, suitable anionic surfactants include sodium lauryl ether sulfate, ammonium lauryl ether sulfate, sodium lauryl sulfate, ammonium lauryl sulfate, triethanolamine lauryl sulfate, disodium laureth sulfosuccinate, sodium cocoyl isethionate, sodium C_{12} - C_{14} olefin sulfonate, sodium laureth-6 carboxylate, sodium C_{12} - C_{15} pareth sulfate, sodium methyl cocoyl taurate, sodium dodecylbenzene sulfonate, sodium cocoyl sarcosinate, triethanolamine monolauryl phosphate, and fatty acid soaps.

In certain illustrative embodiments, examples of suitable sulfonated anionic surfactants include, but are not limited to, alkyl mononuclear aromatic sulfonates, such as the higher alkylbenzene sulfonates containing in one embodiment 8 to 18 carbon atoms, in another embodiment 11 to 16 carbon atoms, and in another embodiment 14 or 15 carbon atoms, the higher alkyl group in a straight or branched chain, or C_8 - C_{15} alkyl toluene sulfonates and C_8 - C_{15} alkyl phenol sulfonates. In another embodiment, the alkylbenzene sulfonate is a linear alkylbenzene sulfonate having a higher content of 3-phenyl (or higher) isomers and a correspondingly lower content (well below 50%) of 2-phenyl (or lower) isomers, such as those sulfonates wherein the benzene ring is attached mostly at the 3 or higher (for example 4, 5, 6 or 7) position of the alkyl group and the content of the isomers in which the benzene ring is attached in the 2 or 1 position is correspondingly low. Illustrative materials are described in U.S. Pat. No. 3,320,174.

In another embodiment, examples of suitable sulfonated anionic surfactants include, but are not limited to, those surface-active or detergent compounds, which contain an organic hydrophobic group containing generally about 8 to about 26 carbon atoms or 10 to 18 carbon atoms in their molecular structure and at least one water-solubilizing group

including, but not limited to, sulfonate, sulfate and carboxylate so as to form a water-soluble detergent. Usually, the hydrophobic group will include a C_8 - C_{22} alkyl, alkyl or acyl group. Such surfactants are employed in the form of water-soluble salts and the salt-forming cation is sodium, potassium, ammonium, magnesium and mono-, di- or tri- C_2 - C_3 alkanolammonium. In an illustrative embodiment the cations are sodium, magnesium or ammonium cations.

Other suitable anionic surfactants encompassed within the scope of the invention include, but are not limited to, the olefin sulfonates, including long-chain alkene sulfonates, long-chain hydroxyalkane sulfonates or mixtures of alkene sulfonates and hydroxyalkane sulfonates. These olefin sulfonate detergents may be prepared in a known manner by the reaction of sulfur trioxide (SO_3) with long-chain olefins containing 8 to 25, or 12 to 21 carbon atoms and having the formula $RCH=CHR_1$ where R is a higher alkyl group of 6 to 23 carbons and R_1 is an alkyl group of 1 to 17 carbons or hydrogen to form a mixture of sulfones and alkene sulfonic acids which is then treated to convert the sulfones to sulfonates. In other embodiments olefin sulfonates contain about 14 to about 16 carbon atoms in the R alkyl group and are obtained by sulfonating an alpha-olefin.


Other examples of suitable anionic sulfonate surfactants encompassed within the scope of the invention include the paraffin sulfonates containing about 10 to about 20, or about 13 to about 17 carbon atoms. Primary paraffin sulfonates are made by reacting long-chain alpha olefins and bisulfites and paraffin sulfonates having the sulfonate group distributed along the paraffin chain are shown in U.S. Pat. Nos. 2,503,280; 2,507,088; 3,260,744; 3,372,188; and German Patent 735,096.

The compositions of the invention may also include alkyl ethoxylated ether sulfates. Another surfactant utilized in the instant composition at a concentration of about 2 to about 15% by weight in one embodiment or about 4 to about 14% by weight in another embodiment is a metal salt of a C_8 - C_{18} alkyl ethoxylated ether sulfate. The ethoxylated alkyl ether sulfate (AEOS.xEO) is depicted by the Formula I:



In one embodiment, x is 1 to 22; in another embodiment x is 1 to 10. In certain embodiments, R^1 is an alkyl group having 10 to 16 carbon atoms; in other embodiments R_3 is an alkyl group having 12 to 15 carbon atoms. In other embodiments, R_3 is C_{12} - C_{14} , C_{12} - C_{13} or C_{12} - C_{15} and M is an alkali metal cation such as, for example, lithium, potassium and sodium or an alkali earth metal cation such as magnesium.

Other examples of anionic ethoxylated sulfates are the C_8 - C_{18} ethoxylated alkyl ether sulfate salts having the Formula II:



where R^1 is defined above.

In another embodiment, the anionic surfactant is present in an amount of about 3 wt. % to about 20 wt. %. In another embodiment, the anionic surfactant is present in an amount of about 5 wt. % to about 15 wt. %. In another embodiment, the anionic surfactant is present in an amount of about 8 wt. % to about 13 wt. %. In another embodiment, the anionic surfactant is present in an amount of about 12 wt. % to about 13 wt. %.

In certain embodiments, the compositions include a first surfactant and a second surfactant. In certain embodiments, the first anionic surfactant is present in an amount of about 8 wt. % to about 18 wt. % based on the weight of the total composition. In certain embodiments, the first anionic sur-

factant is present in an amount of about 8 wt. % to about 13 wt. % based on the weight of the total composition. In certain embodiments, the first anionic surfactant is present in an amount of about 8.5 wt. % based on the weight of the total composition. In certain embodiments, the first anionic surfactant is present in an amount of about 12.5 wt. % based on the weight of the total composition. In certain embodiments, the second anionic surfactant is present in an amount of about 5 wt. % to about 20 wt. % based on the weight of the total composition. In certain embodiments, the second anionic surfactant is present in an amount of about 13 wt. % to about 20 wt. % based on the weight of the total composition. In certain embodiments, the second anionic surfactant is present in an amount of about 13.5 wt. % based on the weight of the total composition. In certain embodiments, the second anionic surfactant is present in an amount of about 18 wt. % based on the weight of the total composition.

Zwitterionic Surfactants

The compositions of the invention also include one or more zwitterionic surfactants. In certain embodiments, the zwitterionic surfactant is also an amphoteric surfactant. Amphoteric and zwitterionic surfactants are those compounds that have the capacity of behaving either as an acid or a base. Suitable zwitterionic or amphoteric surfactants include, but are not limited to, alkyl betaines, alkyl amidopropyl betaines, alkyl sulphobetaines, alkyl glycinate, alkyl carboxyglycinate, alkyl amphopropionates, alkyl amidopropyl hydroxysulfates, acyl taurates and acyl glutamates wherein the alkyl and acyl groups have about 8 to about 18 carbon atoms. Examples include cocamidopropyl betaine, sodium cocoamphoacetate, cocamidopropyl hydroxysulfate, and sodium cocamphopropionate.

In another embodiment, suitable zwitterionic surfactants for use herein contain both a cationic hydrophilic group (i.e., a quaternary ammonium group) and anionic hydrophilic group on the same molecule at a relatively wide range of pH's. The typical anionic hydrophilic groups are carboxylates and sulfonates, although other groups like sulfates, phosphonates, and the like can be used.

In certain embodiments, the zwitterionic surfactants also include hydrophobic groups including aliphatic or aromatic, saturated or unsaturated, substituted or unsubstituted hydrocarbon chains that can contain linking groups such as amido groups, ester groups. In another embodiment the hydrophobic group is an alkyl group containing about 1 to about 24 carbon atoms, in another embodiment about 8 to about 18, and in another embodiment about 10 to about 16. In certain embodiments, simple alkyl groups are utilized for cost and stability reasons.

Some common examples of betaine/sulphobetaine are described in U.S. Pat. Nos. 2,082,275, 2,702,279 and 2,255,082.

Examples of suitable alkyldimethyl betaines include, but are not limited. cocodimethyl betaine, lauryl dimethyl betaine, decyl dimethyl betaine, 2-(N-decyl-N,N-dimethylammonia)acetate, 2-(N-coco N,N-dimethylammonio)acetate, myristyl dimethyl betaine, palmityl dimethyl betaine, cetyl dimethyl betaine, stearyl dimethyl betaine. For example Coconut dimethyl betaine is commercially available from Seppic under the trade name of Amony 265®. Lauryl betaine is commercially available from Albright & Wilson under the trade name Empigen BB/L®.

Examples of amidobetaines include cocoamidoethylbetaine, cocoamido-propyl betaine or C₁₀-C₁₄ fatty acylamidopropylene(hydropropylene)-sulfobetaine. For example C₁₀-C₁₄ fatty acylamidopropylene(hydropropylene)-sulfobetaine is commercially available from Sherex Company

under the trade name "Varion CAS® sulfobetaine." A further example of betaine is Lauryl-imino-dipropionate. Laurylamido propylbetaine is commercially available from Stepan Chemical under tradename AmphoSol LB.

In certain embodiments, the zwitterionic surfactant is present in an amount of about 2 wt. % to about 7 wt. % based on the weight of the total composition. In certain embodiments, the zwitterionic surfactant is present in an amount of about 2.5 wt. % to about 6.5 wt. % based on the weight of the total composition. In certain embodiments, the zwitterionic surfactant is present in an amount of about 5.5 wt. % to about 6.5 wt. % based on the weight of the total composition.

Acids of the Invention

The cleaning compositions of the invention also include an acid constituent, which can be a water soluble inorganic acid or a water soluble organic acid. By way of non-limiting example inorganic acids include, but are not limited to hydrochloric acid, phosphoric acid, boric acid, and sulfuric acid. The organic acids of the invention generally include at least one carbon atom, and include at least one carboxyl group (—COOH) in its structure. In certain embodiments, water soluble organic acids contain from 1 to about 6 carbon atoms and at least one carboxyl group.

In certain embodiments, organic acids include, but are not limited to, formic acid, citric acid, sorbic acid, acetic acid, glycolic acid, propanoic acid, propionic acid, oxalic acid, maleic acid, tartaric acid, adipic acid, lactic acid, malic acid, malonic acid, glycolic acid, and mixtures thereof.

In certain embodiments, the acid constituent is lactic acid, for example. D- and/or L-lactic acid or mixtures thereof.

The compositions are acidic in nature (pH<7.0). Accordingly, there should be sufficient acid present in the composition such that the pH of the composition in various embodiments is less than about 6, or about 2 to about 3.5, or about 2.8 to about 3.3, or about 3.0 to about 3.3. The pH of the composition incorporating a selected acid is preferred to be within 10% of the pka of the selected acid. Mixtures of two or more acids may be used. and the acid constituent may be present in any effective amount. The pH of the composition after the aging period remains less than about 6.5. or about 2.1 to about 3.6. or about 2.9 to about 3.4, or about 3.1 to about 3.4. The pH of the composition after the aging period incorporating a selected acid is preferred to be within 10% of the pka of the selected acid. The aging period should be at least about two months, or about 6 months, or about 1 year or about 2 years.

The acid is present in an amount of less than about of 5% wt. based on the total weight of the compositions. In other embodiments, the acid is present in an amount of about 0.05 to about 4% wt., from about 1 to about 3% wt., and in an amount of about 2% wt. to about 2.5% wt. The amount of acid present after the aging period should not differ substantially from the level of acid in the initial composition.

Sequestering/Chelating Agents of the Invention

In certain embodiments, the cleaning compositions of the invention can also contain an organic or inorganic sequestrant or mixtures of sequestrants. Organic sequestrants such as citric acid, the alkali metal salts of nitrilotriacetic acid (NTA), EDTA or salts thereof, alkali metal gluconates, polyelectrolytes such as a polyacrylic acid, and the like can be used herein. In certain embodiments, sequestrants are organic sequestrants such as sodium gluconate due to the compatibility of the sequestrant with the formulation base.

The sequestering agent of the invention also includes an effective amount of a water-soluble organic phosphonic acid, which has sequestering properties. In certain embodiments, phosphonic acids include low molecular weight compounds containing at least two anion-forming groups, at least one of

which is a phosphonic acid group. Such useful phosphonic acids include mono-, di-, tri- and tetra-phosphonic acids which can also contain groups capable of forming anions under alkaline conditions such as carboxy, hydroxy, thio and the like.

The phosphonic acid may also include a low molecular weight phosphonopolycarboxylic acid such as one having about 2-4 carboxylic acid moieties and about 1-3 phosphonic acid groups. Such acids include 1-phosphono-1-methylsuccinic acid, phosphosuccinic acid and 2-phosphonobutane-1,2,4-tricarboxylic acid.

Other organic phosphonic acids include 1-hydroxyethylidene-1,1-diphosphonic acid ($\text{CH}_3\text{C}(\text{PO}_3\text{H}_2)_2\text{OH}$), available from Monsanto Industrial Chemicals Co., St. Louis, Mo. as Dequest® 2010, a 58-62% aqueous solution; amino [tri(methylenephosphonic acid)] ($\text{N}[\text{CH}_2\text{PO}_3\text{H}_2]_3$), available from Monsanto as Dequest® 2000, a 50% aqueous solution; ethylenediamine [tetra(methylene-phosphonic acid)] available from Monsanto as Dequest® 2041, a 90% solid acid product; and 2-phosphonobutane-1,2,4-tricarboxylic acid available from Mobay Chemical Corporation, Inorganic Chemicals Division, Pittsburgh, Pa. as Bayhibit AM, a 45-50% aqueous solution. It will be appreciated that, the above-mentioned phosphonic acids can also be used in the form of water-soluble acid salts, particularly the alkali metal salts, such as sodium or potassium; the ammonium salts or the alkylol amine salts where the alkylol has 2 to 3 carbon atoms, such as mono-, di-, or tri-ethanolamine salts. If desired, mixtures of the individual phosphonic acids or their acid salts can also be used. Further useful phosphonic acids are disclosed in U.S. Pat. No. 4,051,058, the disclosure of which is incorporated by reference herein. In certain embodiments, phosphonic acids useful in the present invention do not contain amino groups since they produce substantially less degradation of the active chlorine source than do phosphonic acids including amino groups.

Sequestrants of the invention also include materials such as, complex phosphate sequestrants, including sodium tripolyphosphate, sodium hexametaphosphate, and the like, as well as mixtures thereof. Phosphates, the sodium condensed phosphate hardness sequestering agent component functions as a water softener, a cleaner, and a detergent builder. Alkali metal (M) linear and cyclic condensed phosphates commonly have a $\text{M}_2\text{O}:\text{P}_2\text{O}_5$ mole ratio of about 1:1 to 2:1 and greater. Typical polyphosphates of this kind are sodium tripolyphosphate, sodium hexametaphosphate, sodium metaphosphate as well as corresponding potassium salts of these phosphates and mixtures thereof. The particle size of the phosphate is not critical, and any finely divided or granular commercially available product can be employed.

In certain embodiments, sodium tripolyphosphate is an inorganic hardness sequestering agent for reasons of its ease of availability, low cost, and high cleaning power. Sodium tripolyphosphate acts to sequester calcium and/or magnesium cations, providing water softening properties. It contributes to the removal of soil from hard surfaces and keeps soil in suspension. It has little corrosive action on common surface materials and is low in cost compared to other water conditioners. Sodium tripolyphosphate has relatively low solubility in water (about 14 wt-%) and its concentration must be increased using means other than solubility. Typical examples of such phosphates being alkaline condensed phosphates (i.e. polyphosphates) such as sodium or potassium pyrophosphate, sodium or potassium tripolyphosphate, sodium or potassium hexametaphosphate.; carbonates such as sodium or potassium carbonate; borates, such as sodium borate.

If utilized, the sequestering or chelating agent(s) will generally include about 0.00015% to about 15% by weight of the cleaning compositions herein. In other various embodiments, if utilized, the sequestering or chelating agent(s) will include about 0.0003% to about 3.0% by weight of such compositions or about 0.003% to about 1.0% by weight of such compositions or about 0.03% to about 0.1% by weight of such compositions.

Fragrance Agents

The compositions and methods of the invention can also include one or more fragrance agents. Fragrance agents useful in the compositions and methods include a wide variety of natural and synthetic chemical ingredients, including, but not limited to, aldehydes, ketones, esters, and the like. Also included are various natural extracts and essences, which can include complex mixtures of ingredients, such as orange oil, lemon oil, rose extract, lavender, musk, patchouli, balsamic essence, sandalwood oil, pine oil, cedar, and the like. Finished fragrance agents can include extremely complex mixtures of such ingredients. Finished fragrance agents typically include about 0.01% to about 2%, by weight, of the detergent compositions herein, and individual fragrance agents can include about 0.0001% to about 90% of a finished perfume composition.

In a certain embodiment of the invention, the composition includes a blooming perfume. A blooming perfume ingredient is characterized by its boiling point (B.P.) and its octanol/water partition coefficient (P). The octanol/water partition coefficient of a perfume ingredient is the ratio between its equilibrium concentrations in octanol and in water. The fragrance agents of the invention have a B.P., determined at the normal, standard pressure of about 760 mm Hg, of about 260° C. or lower, less than about 255° C.; and less than about 250° C., and an octanol/water partition coefficient P of about 1,000 or higher. Since the partition coefficients of the fragrance agents of the invention have high values, they are more conveniently given in the form of their logarithm to the base 10, log P. Thus the fragrance agents have log P of about 3 or higher, or more than about 3. 1, or more than about 3.2.

In certain embodiments, the compositions can include a combination of fragrance agents. In certain embodiments, the composition includes a first perfume ingredient having boiling point of 250° C. or less and C log P of 3.0 or less; and a second perfume ingredient having boiling point of 250° C. or less and C log P of 3.0 or more.

Hydrotropes

The compositions of the invention can also include one or more hydrotrope(s). Without being limited by theory it is believed that the hydrotrope contributes to the physical and chemical stability of the compositions.

Suitable hydrotropes include sulphonated hydrotropes. Any sulphonated hydrotropes known to those skilled in the art are suitable for use herein. In certain embodiments, alkyl aryl sulphonates or alkyl aryl sulphonic acids are used. In other embodiments alkyl aryl sulphonates include sodium, potassium, calcium and ammonium xylene sulphonates, sodium, potassium, calcium and ammonium toluene sulphonates, sodium, potassium, calcium and ammonium cumene sulphonates, sodium, potassium, calcium and ammonium substituted or unsubstituted naphthalene sulphonates and mixtures thereof. In other embodiments alkyl aryl sulphonic acids include xylenesulphonic acid, toluenesulphonic acid, cumenesulphonic acid, substituted or unsubstituted naphthalenesulphonic acid and mixtures thereof. In other embodiments, xylenesulphonic acid or p-toluene sulphonate or mixtures thereof are used.

In various embodiments, the compositions may include hydrotropes in amounts of about 0.01 wt. % to 20 wt. %, about wt. 0.05% to 10 wt. % or about 0.1 wt. % to 5 wt. % or about 3 wt. % by weight of the total composition.

Solvents of the Invention

The invention in certain embodiments can also include one or more solvents. Typical solvents used in the composition are aqueous soluble, miscible or immiscible. Solvents can include aliphatic and aromatic hydrocarbons, chlorinated hydrocarbons, alcohols, ether compounds, fluorocarbon compounds, and other similar low molecular weight generally volatile liquid materials. In various embodiments, the compositions may include solvents in amounts of up to about 6 wt. %, preferably at least about wt. 0.1% by weight of the total composition.

In certain embodiments, water is not a solvent but when used acts as a diluent or as a dispersing medium for the active materials. In other embodiments, water is a solvent.

These materials can be used in solution or as a miscible mixture or as a dispersion of the solvent in the aqueous liquid. A solvent or cosolvent can be used to enhance certain soil removal properties of this invention. Cosolvents include alcohols and the mono and di-alkyl ethers of alkylene glycols, dialkylene glycols trialkylene glycols, etc. Alcohols which are useful as cosolvents in this invention include methanol, ethanol, propanol and isopropanol. Particularly useful in this invention are the mono and dialkyl ethers of ethylene glycol and diethylene glycol, which have acquired trivial names such as polyglymes, cellosolves, and carbitols. Representative examples of this class of cosolvent include methyl cellosolves, butyl carbitol, dibutyl carbitol, diglyme, triglyme. Nonaqueous liquid solvents can be used for varying compositions of the present invention. These include the higher glycols, polyglycols, polyoxides and glycol ethers.

Suitable substances are propylene glycol, polyethylene glycol, polypropylene glycol, diethylene glycol monoethyl ether, diethylene glycol monopropyl ether, diethylene glycol monobutyl ether, tripropylene glycol methyl ether, propylene glycol methyl ether (PM), dipropylene glycol methyl ether (DPM), propylene glycol methyl ether acetate (PMA), dipropylene glycol methyl ether acetate (CPMA), propylene glycol n-butyl ether, dipropylene glycol monobutyl ether, ethylene glycol n-butyl ether and ethylene glycol n-propyl ether, and combinations thereof. In certain embodiments, the glycol solvent is propylene glycol n-butyl ether. In certain embodiments, the glycol solvent is dipropylene glycol monobutyl ether.

Other useful solvents are ethylene oxide/propylene oxide, liquid random copolymer such as Synalox® solvent series from Dow Chemical (e.g., Synalox® 50-50B). Other suitable solvents are propylene glycol ethers such as PnB, DPnB and TPnB (propylene glycol mono n-butyl ether, dipropylene glycol and tripropylene glycol mono n-butyl ethers sold by Dow Chemical under the trade name Dowanol.®). Also tripropylene glycol mono methyl ether "Dowanol TPM®" from Dow Chemical is suitable.

The final ingredient in the inventive cleaning compositions is water. The proportion of water in the compositions generally is in the range of about 35% to about 90% or about 50% to 85% by weight of the cleaning composition.

Thickening Agents

In certain embodiments, the compositions of the invention also include a thickening agent. Suitable thickeners may be organic or inorganic in nature. The thickener may thicken the composition by either thickening the aqueous portions of the composition, or by thickening the non-aqueous portions of the composition. In certain embodiments, the thickening

agent is a water soluble polymer. In other embodiments, the thickening agent is a cationic water soluble polymer.

In certain embodiments, the polymeric thickener may be added to the composition to achieve two objectives (i) to increase the "flow" viscosity and (ii) to neutralize the anionic detergent carry-over guaranteeing a high level of softness in washing conditions where the carry-over is high. The flow viscosity corresponds to the viscosity measured with a flowmeter. The flowability of the tested composition is expressed as time needed to a fixed amount of product flowing through a small tube.

Thickeners can be divided into organic and inorganic thickeners. Of the organic thickeners there are (1) cellulosic thickeners and their derivatives, (2) natural gums, (3) acrylates, (4) starches, (5) stearates, (6) fatty acid alcohols and inorganic thickeners including (7) clays, and (8) salts. Some non-limiting examples of cellulosic thickeners include carboxymethyl hydroxyethylcellulose, cellulose, hydroxybutyl methylcellulose, hydroxyethylcellulose hydroxypropylcellulose, hydroxypropyl methyl cellulose, methylcellulose, microcrystalline cellulose, sodium cellulose sulfate, and the like. Some non-limiting examples of natural gums include acacia, calcium carrageenan, guar, gelatin, guar gum, hydroxypropyl guar, karaya gum, kelp, locust bean gum, pectin, sodium carrageenan, gellan gum, tragacanth gum, xanthan gum, and the like. Some non-limiting examples of acrylates include potassium aluminum polyacrylate, sodium acrylate/vinyl alcohol copolymer, sodium polymethacrylate, and the like. Some non-limiting examples of starches include oat flour, potato starch, wheat flour, wheat starch, and the like. Some non-limiting examples of stearates include methoxy PEG-22/dodecyl glycol copolymer, PEG-2M, PEG-5M, and the like. Some non-limiting examples of fatty acid alcohols include caprylic alcohol, cetearyl alcohol, lauryl alcohol, oleyl alcohol, palm kernel alcohol, and the like. Some non-limiting examples of clays include bentonite, magnesium aluminum silicate, magnesium trisilicate, stearylaluminum bentonite, tromethamine magnesium aluminum silicate, and the like. Some non-limiting examples of salts include calcium chloride, sodium chloride, sodium sulfate, ammonium chloride, and the like.

Some non-limiting examples of thickeners that thicken the non-aqueous portions of the composition include waxes such as candelilla wax, carnauba wax, beeswax, and the like, oils, vegetable oils and animal oils, and the like.

The composition may contain one thickener or a mixture of two or more thickeners. In certain embodiments the thickeners do not adversely react with the other components or compounds of the invention or otherwise render the composition of the invention ineffective. It is understood that a person skilled in the art will know how to select an appropriate thickener and control any adverse reactions through formulating.

The amount of thickener present in the composition depends on the desired viscosity of the composition. The composition may have a viscosity of about 100 to about 15,000 centipoise, of about 150 to about 10,000 centipoise, and of about 200 to about 5,000 centipoise as determined using a Brookfield DV-II+rotational viscometer using spindle # 21 @ 20 rpm @ 70° F. Accordingly, to achieve the desired viscosities, the thickener may be present in the composition in an amount about 0.001 wt. % to about 5 wt. % of the total composition, about 0.01 wt. % to about 3 wt. %, and about 0.05 wt. % to about 2 wt. % of the total composition.

Thickeners from said classes of substance are commercially broadly available and are obtainable, for example, under the trade names Acusol® 820 (methacrylic acid

(stearyl alcohol-20 EO) ester-acrylic acid copolymer, 30% strength in water. Rohm & Haas). Dapral®-GT-282-S (alkyl polyglycol ether, Akzo), Deuterol® polymer-11 (dicarboxylic acid copolymer, Schoner GmbH), Deuteron® XG (anionic heteropolysaccharide based on beta-D-glucose, D-manose, D-glucuronic acid, Schoner GmbH), Deuteron®-XN (nonionogenic polysaccharide, Schoner GmbH), Dicrylan® thickener-O (ethylene oxide adduct, 50% strength in water/isopropanol, Pfersse Chemie), EMA®-81 and EMA®-91 (ethylene-maleic anhydride copolymer, Monsanto), thickener-QR-1001 (polyurethane emulsion, 19 21% strength in water/diglycol ether, Rohm & Haas), Mirox®-AM (anionic acrylic acid-acrylic ester copolymer dispersion, 25% strength in water, Stockhausen), SER-AD-FX-1100 (hydrophobic urethane polymer, Servo Delden), Shellflo®-S (high molecular weight polysaccharide, stabilized with formaldehyde, Shell) and Shellflo®-XA (xanthan biopolymer, stabilized with formaldehyde, Shell).

The inventors have discovered that xanthan gum is useful as a thickening agent for suspending fragrance molecules in a hard surface cleaner. In certain embodiments, the thickening agent is xanthan gum. In other embodiments, the thickening agent is xanthan gum present in at least about 0.2 weight %.

Additional Optional Ingredients

Examples of additional optional components include, but are not limited to, hydrotropes, fluorescent whitening agents, photobleaches, fiber lubricants, reducing agents, enzymes, enzyme stabilizing agents, powder finishing agents, builders, bleaches, bleach catalysts, soil release agents, dye transfer inhibitors, buffers, colorants, fragrances, pro-fragrances, rheology modifiers, anti-ashing polymers, preservatives, soil repellents, water-resistance agents, suspending agents, aesthetic agents, structuring agents, sanitizers, solvents, fabric finishing agents, dye fixatives, fabric conditioning agents and deodorizers.

The instant cleaning compositions may contain optionally about 0.25 wt. % to about 10 wt. %, about 1 wt. % to about 8 wt. %, of at least one solubilizing agent. The solubilizing agents include, but are not limited to, C₁-C₅ mono, dihydroxy or polyhydroxy alkanols such as ethanol, isopropanol, alkylene glycols such as hexylene glycol, glycerol ethylene glycol, diethylene glycol and propylene glycol and mixtures thereof and alkali metal cumene or xylene sulfonates such as sodium cumene sulfonate and sodium xylene sulfonate. The solubilizing agents are included in order to control low temperature cloud clear properties. Urea can be optionally employed in the instant composition as a supplemental solubilizing agent at a concentration of 0 to about 10 wt. %, about 0.5 wt. % to about 8 wt. %.

Other surfactants which can be utilized in the present invention are set forth in more detail in WO 99/21530, U.S. Pat. No. 3,929,678; U.S. Pat. No. 4,565,647; U.S. Pat. No. 5,720,964; and U.S. Pat. No. 5,858,948. Other suitable surfactants are described in McCutcheon's Emulsifiers and Detergents (North American and International Editions, by Schwartz, Perry and Berch), which is hereby fully incorporated by reference.

In addition to the previously mentioned constituents of the composition, one may also employ normal and conventional adjuvants, provided they do not adversely affect the properties of the detergent. Thus there may be used a cationic antibacterial agent, coloring agents and perfumes; polyethylene glycol, ultraviolet light absorbers such as the Uvinuls, which are products of GAF Corporation; pH modifiers; etc. The proportion of such adjuvant materials, in total will normally not exceed 15% by weight of the detergent composition, and the percentages of illustrative examples of such individual

components will be about 5% by weight. Sodium formate or formalin or Quaternium 15 (Dowicil 75) can be included in the formula as a preservative at a concentration of about 0.1 to about 4.0 wt. %.

5 Process of Manufacture

The compositions are readily made by simple mixing methods from readily available components which, on storage, do not adversely affect the entire composition. Solubilizing agent such as ethanol, hexylene glycol, sodium chloride and/or sodium xylene or sodium xylene sulfonate are used to assist in solubilizing the surfactants. The viscosity of the light duty liquid composition desirably will be at least 100 centipoises (cps) at room temperature, but may be up to 1,000 centipoises. The viscosity of the light duty liquid composition and the light duty liquid composition itself remain stable on storage for lengthy periods of time, without color changes or settling out of any insoluble materials. In certain embodiments, the pH of the composition is about 3.5.

20 Methods of Use

The invention encompasses cleaning compositions useful for cleaning a surface. The compositions surprisingly possess antibacterial efficacy and low toxicity.

By surfaces, it is meant herein any kind of surfaces typically found in houses like kitchens, bathrooms, or the exterior surfaces of a vehicle, for example, floors, walls, tiles, windows, sinks, showers, shower plastified curtains, wash basins, WCs, dishes and other food contact surfaces, fixtures and fittings and the like made of different materials like ceramic, vinyl, no-wax vinyl, linoleum, melamine, glass, any plastics, plastified wood, metal, especially steel and chrome metal or any painted or varnished or sealed surface and the like. Surfaces also include household appliances including, but not limited to, refrigerators, garbage cans, freezers, washing machines, automatic dryers, ovens, microwave ovens, dishwashers and so on. The present composition is especially efficacious in the cleaning of ceramic, steel, plastic, glass and the exterior painted or otherwise finished surface of a vehicle, for example, a car. The cleaning compositions are also safe on the skin.

The cleaning composition is applied to the surface, undiluted or diluted, optionally after a pre-rinse step. The composition can be applied using a cloth or sponge onto which the composition has been applied or by pouring the composition over the surface. Alternatively the composition may be applied by spraying the composition onto the surface using a spraying device as described above. The cleaning compositions of the invention can be left to sit on a surface or be wiped or scrubbed on or from the surface.

Once the composition has been applied to the surface, the surface can then be optionally rinsed, usually with water and left to dry naturally. Optionally the user can wait in between application of the composition and rinsing in order to allow the composition maximum working time. A particular benefit of the composition is that the surface can be cleaned as described above and the surface left to dry naturally with minimal formation of water spots, streaks, and/or with reduced or no corrosion.

The following examples illustrate compositions of the invention. Unless otherwise specified, all percentages are by weight. The exemplified compositions are illustrative only and do not limit the scope of the invention. Unless otherwise specified, the proportions in the examples and elsewhere in the specification are by active weight. The active weight of a material is the weight of the material itself excluding water or other materials that may be present in the supplied form of the material.

17
EXAMPLES

The following examples illustrate liquid cleaning compositions of the invention. Unless otherwise specified, all percentages are by weight. The exemplified compositions are illustrative only and do not limit the scope of the invention. It will be understood by those of skill in the art that numerous and various modifications can be made without departing from the spirit of the present invention. Therefore, it should be clearly understood that the forms of the present invention described herein are illustrative only and are not intended to limit the scope of the invention.

Example 1

Tables 1a-1d illustrate several non-limiting illustrative embodiments of the invention illustrating regular and ultra dishwashing liquids.

TABLE 1a

Ingredient	Reg 2a (wt. %)	Reg 2b (wt. %)	Reg 1a (wt. %)	Reg 1b (wt. %)	Ultra 2a (wt. %)	Ultra 2b (wt. %)	Ultra 1a (wt. %)	Ultra 1b (wt. %)
Na DBS ¹	3.7	3.7	8.4	8.4	5.8	5.8	8.5	8.5
SLES (2EO) C ₁₂ -C ₁₄	12	12	9.2	9.2	19.6	19.6	17.9	17.9
Laurylamido-propyl betaine	3.4	3.4	1.5	1.5	6.6	6.6	5.4	5.4
Ethanol	3.5	3.5	2.1	2.1	6	6	4.3	4.3
Lactic Acid	2	2	2	2	2	2	2	2
Mg ₂ SO ₄	0	0	3.5	3.5	0	0	0.9	0.9
SXS	0.8	0.8	0.8	0.8	2.2	2.2	3	3
NaCl	0	0	0.7	0.7	0	0	0	0
Na ₄ EDTA	0.1	0.1	0.1	0.1	0.1	0.1	0.1	0.1
Color solution	0.2	0.2	0.2	0.2	0.2	0.2	0.2	0.2
Fragrance option A	0.26	0	0.26	0	0.4	0	0.4	0
Fragrance option B	0	0.26	0	0.26	0	0.45	0	0.45
Water	qs	qs	qs	qs	qs	qs	qs	qs
Total Surfactant %	19	19	19	19	32	32	32	32
Calculated LD ₅₀	4100	4100	3895	3895	4157	4157	3721	3721

¹Sodium dodecyl benzene sulfonate

TABLE 1b

Ingredient	Reg. 1	Reg. 2	Ultra 1	Ultra 2
NaLAS	8.4	3.7	8.5	5.75
SLES	9.3	12.1	17.9	19.6
Betaine	1.5	3.4	5.4	6.6
Lactic	2	2	2	2

TABLE 1c

Ingredient	Reg. 1	Reg. 2
NaLAS	5.3	2.3
SLES	5.8	7.6
Betaine	0.1	2.2
Lactic	2	2

TABLE 1d

Ingredient	(wt. %)
Na DBS ¹	12.7
SLES (2EO) C ₁₂ -C ₁₄	13.7
Laurylamido propyl Betaine	5.6
Ethanol	4
Lactic Acid	2
SXS	2.5
Color Solution	0.3

18

TABLE 1d-continued

Ingredient	(wt. %)
Fragrance	0.35
Water	qs

Example 2

Tables 2a-2d illustrate several non-limiting illustrative embodiments of the invention. Amounts added are based on the percent raw amount of ingredient added.

In certain illustrative embodiments of the invention, the EPA mandated antibacterial efficacy of the cleaning compositions has been validated for independent lots of dish liquid manufactured under GMP conditions. The lots were tested on 10 carriers (replicate surfaces) against *Staphylococcus aureus*

ATCC 6538 and, separately on 10 carriers (replicate surfaces) against *Salmonella enterica* ATCC 10708 as mandated by the EPA. A confirmatory test, on independent lots, was also tested on 10 carriers (replicate surfaces) against *Escherichia coli* O157 H7 ATCC 43895 for additional on-pack claims against this specific and relevant food pathogen. The tests were conducted on 1:20 use-dilutions with a 30-second exposure time. In all cases a minimum 3-log reduction or 99.9% kill rate was attained for both the surfaces and the run-off counts, as prescribed for EPA acceptance.

TABLE 2a

Organism	Lot Number	Dilution	Replicate	CFU/carrier	Average log ₁₀ reduction
<i>S. aureus</i>	1a	1:20	1	5.4 × 10 ²	3.41
			2	1.8 × 10 ²	3.89
			3	3.0 × 10 ¹	4.67
		1:50	1	8.0 × 10 ¹	4.54
			2	3.7 × 10 ²	3.58
			3	1.9 × 10 ²	4.15
	2a	1:20	1	4.7 × 10 ²	3.47
			2	5.9 × 10 ²	3.38
			3	7.3 × 10 ²	3.28
		1:50	1	8.8 × 10 ²	3.20
			2	6.1 × 10 ²	3.36
			3	5.5 × 10 ²	3.41
3a	1:20	1	1.0 × 10 ¹	5.51	
		2	5.0 × 10 ¹	4.45	

19

TABLE 2a-continued

Organism	Lot Number	Dilution	Replicate	CFU/carrier	Average log ₁₀ reduction	
	4a	1:50	3	2.0 × 10 ¹	4.85	
			1	1.0 × 10 ¹	5.51	
			2	5.0 × 10 ¹	4.45	
		1:20	3	5.0 × 10 ¹	4.45	
			1	3.4 × 10 ²	3.61	
			2	3.6 × 10 ²	3.59	
	1:50	3	1.2 × 10 ²	4.07		
		1	5.1 × 10 ²	3.44		
		2	3.0 × 10 ²	3.67		
				3	2.4 × 10 ²	3.77

TABLE 2b

Organism	Lot Number	Dilution	Replicate	CFU/carrier	Average log ₁₀ reduction	
<i>S. aureus</i>	1b	1:20	1	4.5 × 10 ²	3.99	
			2	4.7 × 10 ²	3.97	
			3	3.7 × 10 ²	4.08	
		1:50	1	9.3 × 10 ²	3.67	
			2	8.1 × 10 ²	3.73	
			3	7.6 × 10 ²	3.76	
		2b	1:20	1	1.5 × 10 ³	3.47
				2	1.1 × 10 ³	3.60
				3	1.4 × 10 ³	3.50
	1:50		1	1.7 × 10 ³	3.41	
			2	2.4 × 10 ³	3.26	
			3	2.1 × 10 ³	3.32	
	3b	1:20	1	1.2 × 10 ²	4.56	
			2	1.1 × 10 ²	4.60	
			3	4.2 × 10 ²	4.02	
		1:50	1	5.0 × 10 ¹	4.94	
			2	9.0 × 10 ¹	4.69	
			3	7.0 × 10 ¹	4.80	
	4b	1:20	1	3.2 × 10 ²	4.14	
			2	4.9 × 10 ²	3.95	
			3	4.4 × 10 ²	4.00	
		1:50	1	1.1 × 10 ³	3.60	
			2	1.5 × 10 ³	3.47	
			3	6.7 × 10 ²	3.82	

TABLE 2c

Organism	Lot Number	Dilution	Replicate	CFU/carrier	Average log ₁₀ reduction	
<i>E. coli</i>	1c	1:20	1	1.9 × 10 ²	4.17	
			2	2.5 × 10 ²	4.04	
			3	8.3 × 10 ²	3.53	
		1:50	1	3.3 × 10 ²	3.93	
			2	2.0 × 10 ²	4.15	
			3	1.5 × 10 ²	4.27	
		2c	1:20	1	6.3 × 10 ²	3.65
				2	5.0 × 10 ²	3.75
				3	5.1 × 10 ²	3.74
	1:50		1	1.6 × 10 ²	4.24	
			2	2.2 × 10 ²	4.10	
			3	3.0 × 10 ¹	4.97	
	3c	1:20	1	6.2 × 10 ⁴	1.65	
			2	6.6 × 10 ³	2.63	
			3	6.0 × 10 ⁴	1.67	
		1:50	1	6.3 × 10 ⁴	1.65	
			2	6.8 × 10 ³	2.61	
			3	6.2 × 10 ³	2.65	
	3d	1:20	1	4.0 × 10 ²	3.85	
			2	7.3 × 10 ²	3.58	
			3	1.4 × 10 ²	4.30	
		1:50	1	2.0 × 10 ¹	5.15	

20

TABLE 2c-continued

Organism	Lot Number	Dilution	Replicate	CFU/carrier	Average log ₁₀ reduction			
			2	2.0 × 10 ¹	5.15			
			3	2.8 × 10 ²	4.00			
			TABLE 2d					
			<i>E. coli</i>	1d	1:20	1	9.0 × 10 ¹	4.74
						2	<1.0 × 10 ¹	>5.70
						3	<1.0 × 10 ¹	>5.70
1:50	1	6.0 × 10 ²			3.92			
	2	4.4 × 10 ²			4.06			
	3	2.4 × 10 ²			4.32			
2d	1:20	1		1.0 × 10 ¹	5.70			
		2		<1.0 × 10 ¹	>5.70			
		3		4.0 × 10 ²	4.10			
	1:50	1		7.0 × 10 ¹	4.85			
		2		5.0 × 10 ¹	5.00			
		3		2.6 × 10 ²	4.28			
3d	1:20	1	1.8 × 10 ⁴	2.44				
		2	4.4 × 10 ³	3.06				
		3	1.1 × 10 ⁴	2.66				
	1:50	1	4.2 × 10 ³	3.08				
		2	1.7 × 10 ³	3.47				
		3	1.7 × 10 ⁴	2.47				
4d	1:20	1	2.0 × 10 ¹	5.40				
		2	9.0 × 10 ¹	4.74				
		3	2.0 × 10 ¹	5.40				
	1:50	1	1.0 × 10 ²	4.70				
		2	<1.0 × 10 ¹	>5.70				
		3	3.8 × 10 ²	4.12				

Example 3

In certain embodiments, the cleaning compositions of the invention include inert ingredients. The inert ingredients include the surfactants that provide surface cleaning benefits, viscosity modifiers, salts, hydrotropes, chelants that deliver conventional and consumer parameters such as dispensing and clarity, and color/fragrance to provide a consumer-delightful product use experience. The inerts are shown in Table 3 with its status on the EPA Inert List.

TABLE 3

Ingredient	EPA Inert List
Na DBS ¹	3
SLES (2EO) C ₁₂ -C ₁₄	4B
Laurylamidopropyl Betaine	3
Ethanol	4B
Lactic Acid	4B
Mg ₂ SO ₄	4A
SXS	3
NaCl	4A
Na ₄ EDTA	4B
Gellan gum	4A
Water	n/a

¹Sodium dodecyl benzene sulfonate

Example 4

The cleaning compositions of the invention were designed for minimal corrosivity for processing equipment. In Tables 4a, 4b and 4c, high salt, Sample 1, and low salt, Sample 2,

versions of formulas were tested by both short-term, accelerated electrochemical polarization tests at 100° F., and longer-term (6 weeks) immersion tests at 100° F. and at 140° F. with creviced-corrosion coupons made of varying grades of stainless steel.

TABLE 4a

Ingredient	Ingredient % Weight in Formulation			
	Sample 1	Sample 2	Surfactant Control	3% NaCl solution
Na DBS ¹	8.5	8.5	8.2	0
SLES (2EO) C ₁₂ -C ₁₄	17.9	17.9	0	0
NH ₄ AEOS (1.3 EO)	0	0	11.3	0
Laurylamidopropyl betaine	5.4	5.4	0	0
Amine Oxide	0	0	3.5	0
Lactic Acid	2.0	2.0	0	0
MgSO ₄	0.9	0.9	1.31	0
Alcohol	3.0	4.3	1.4	0
NaCl (added)	3.3	0	0.8	3.0
SXS	3.0	3.0	0.65	0
Chealant	0.1	0.1	0.1	0
Color Solution	0.2	0.2	0.1	0
Fragrance A	0.4	0.4	0	0
Fragrance B	0	0	0.3	0

¹Sodium dodecyl benzene sulfonate

TABLE 4b

Formula Description	pH	Cond. ¹ (1st)	Cond. (2nd)	Cl ⁻ (ppm)	SO ₄ ⁻ (ppm)	AcO ⁻ (ppm)	LPR 1 (mpy)	LPR 2 (mpy)
Sample 1	3.04	39.3 □S	41.7 mS	23.980	7.999	14.138	0.248	0.223
Sample 2	2.89	25.6 mS	25.4 mS	6.190	10.496	15.726	0.081	0.068
Surfactant Control	6.60	28.2 mS	28.5 mS	4.494	11.648	BDL	0.059	0.050
3% NaCl Solution	3.00		40.6 mS				0.083	0.122

¹A first conductivity reading was taken and then a second conductivity reading was taken. Electrochemical testing was done at 40° C.

TABLE 4c

Formula	Alloy					
	40° C.			60° C.		
	Surfactant Control	Sample 2	Sample 1	Surfactant Control	Sample 2	Sample 1
Attribute	High pH	Low pH	Low pH	High pH	Low pH	Low pH
Description	Low Cl ⁻	Low Cl ⁻	High Cl ⁻	Low Cl ⁻	Low Cl ⁻	High Cl ⁻
pH	6.6	3.0	3.0	6.6	3.0	3.0
Chloride (ppm)	4500	6200	24000	4500	6200	24000
Stainless Steel ¹	No Attack	No Attack	Crevice Attack	No Attack	Crevice Attack	Crevice Attack
Stainless Steel	Not Tested	Not Tested	Not Tested	No Attack	No Attack	Crevice Attack

¹2 Types of Stainless steel were tested (316L and AL6XN)

Example 5

The cleaning compositions of the invention provide competitive foaming/cleaning performance with existing commercial products. Traditional performance tests were completed to assess the flash foam profile with and without soil (shake-foam), the foam mileage (miniplate), and typical/dynamic (Baumgartner) grease soil removal for both the ultra and regular density dishliquids. The resulting performance profiles against in-market products are shown in the tables

below. These results indicate an unexpectedly higher performance profile that should be more acceptable to the consumer.

(1) Baumgartner Grease Removal

5 The Baumgartner test measures grease removal in every day cleaning situations. Plastic tubes covered with solidified lard, tallow, or mixed greasy soil are dipped in a warm LDL solution 100 times; the concentration of the solution is 0.0667%. The total dipping time is approximately 1 minute. 10 The tubes are weighed before and after grease is applied. After the tubes dry, the % grease removal is calculated.

(2) Shake-foam Test

15 100 ml of a diluted (0.033%) test solution in 150 ppm hardness water at RT is filled into a 500 ml graduated cylinder with a stopper. The stoppered cylinder is placed on an agitating machine, which rotates the cylinder for 40 cycles at 30 rpm. The height of the foam in the cylinder is observed. A milk soil is then introduced (about 175 μL) into the cylinder. 20 The cylinder is then inserted 40 times more, and the height after soil addition is recorded.

25 The number of miniplates is measured using an automated miniplate test. The procedure is described in detail in U.S. Pat. No. 4,556,509, which is incorporated herein by reference. The test is used to determine the number of theoretical plates that can be washed in a cleaning solution until the foam disappears. This test is used to demonstrate the improvement

in cleaning efficiency as gauged by foam volume and foam stability. Foam is generated in a detergent solution by the action of an agitating brush. The foam is electronically measured by reflectance of the solution surface (with an added dye) as a mixed soil (potato, milk, olive oil, crisco) is added to the detergent solution at a steady rate. The disappearance of the foam determines the endpoint of the test and the number of m-niplates is then calculated based on foam duration and the rate of soil addition. For these, tests the detergent solution

was an illustrative cleaning composition of the invention at 3.3 wt. % with 150 ppm Mg/CaCO₃ hardness and was initially heated to 47° C. at the start of soil addition.

TABLE 5a

Ingredient	Ingredient % Weight in Formulation				In-market Control
	Sample 1	Sample 2	Sample 3	Sample 4	
Na DBS ¹	8.4	4	4	4	0
Mg DBS ²	0	0	0	0	12.2
SLES (2EO) C ₁₂ -C ₁₄	9.3	13	13	13	0
NH ₄ AEOS (1.3 EO)	0	0	0	0	7
Laurylamidopropyl betaine	1.5	3.7	3.7	3.7	0
Amine Oxide	0	0	0	0	1
MgSO ₄	0.9	0.9	0	0	0.5
NaCl (added)	1.85	3.25	2.5	0	0
Antibacterial	Lactic acid	Lactic acid	Lactic acid	Lactic acid	Triclosan
Total % Surfactant	19.2	20.7	20.7	20.7	20.2

¹Sodium dodecyl benzene sulfonate

²Magnesium dodecyl benzene sulfonate

TABLE 5b

Sample	Total % Surfactant	Baumgartner	SFI Mean	SFS Mean	Miniplate
1	19.2%	75.83	391.67	128.33	14
2	20.7%	81.13	391.67	130.00	20
3	20.7%	76.17	391.67	136.37	19.5
4	20.7%	80.27	408.33	138.33	23
In-market Control	20.2%	88.32	383.33	123.33	17

Example 6

25 The cleaning compositions of the invention provide superior rinsing and/or shine performance with existing products. Studies showed that the acidic formula can deliver advantages on rinsing attributes versus in-market formulas. This is likely to be especially noticeable in hard water environments.

30 The rinsing benefits of the compositions of the invention were demonstrated by actual in lab rinsing measurements. This method involves applying an illustrative cleaning composition of the invention to a plate and recording the time it

TABLE 5c

Ingredient	Ingredient % Weight in Formulation				In-market Control
	Sample 1	Sample 2	Sample 3	Sample 4	
Na DBS ¹	5.7	5.7	8.5	8.5	0.9
Mg DBS ²	0	0	0	0	13.1
SLES (2EO) C ₁₂ -C ₁₄	18.9	18.9	17.9	17.9	0
NH ₄ AEOS (1.3 EO)	0	0	0	0	16.2
Laurylamidopropyl betaine	5.4	5.4	5.4	5.4	0
Amine Oxide	0	0	0	0	5.9
MgSO ₄	0	0	0.85	0.85	0
NaCl (added)	3.5	0	3.0	0	0.4
Antibacterial	Lactic acid	Lactic acid	Lactic acid	Lactic acid	Triclosan
Total % Surfactant	30	30	31.8	31.8	35.6

¹Sodium dodecyl benzene sulfonate

²Magnesium dodecyl benzene sulfonate

TABLE 5d

Sample	Total % Surfactant	Baumgartner	SFI Mean	SFS Mean	Miniplate
1	30%	68.267	390	141.67	20.4
2	30%	67.75	398.33	146.67	19
3	31.8%	71.12	403.33	145.00	21.4
4	31.8%	75.85	395.67	163.33	21.4
In-Market Control	35.6%	84.03	380	161.67	23.4

55 takes for full rinsing of the product. The illustrative cleaning compositions of the invention were nearly twice as fast to rinse.

TABLE 6

	Composition of the Invention	Control
60		
65	Rinsing Time	6.5 sec
		11.8 sec

25

Example 7

Tables 7a-7e illustrate properties including good Foam Volume, good Grease Redeposition, and good rinsibility of illustrative embodiments of the invention.

TABLE 7a

Ingredient	Sample 1	Sample 2	Sample 3	Sample 4 In-market Control
Na DBS ¹	10	10	6.4	0
Mg DBS ²	0	0	0	14
SLES (2EO) C ₁₂ -C ₁₄	21	21	24.6	0
NH ₄ AEOS (1.3 EO)	0	0	0	16.1
Laurylamidopropyl betaine	3.6	3.6	3.6	0
Amine Oxide	0	0	0	5.9
MgSO ₄	0	1.7	1.6	0
NaCl (added)	0	0.25	1.3	0.4
Antibacterial	Lactic acid	Lactic acid	Lactic acid	Triclosan
Total % Surfactant	34.7	34.6	34.6	36

¹Sodium dodecyl benzene sulfonate

²Magnesium dodecyl benzene sulfonate

TABLE 7b

Product	ini	soil	re-en	Neat ¹ Foam1	Neat ² Foam2
Sample 1	0.42 ³	0.55	0.51	0.70	0.73
Sample 2	6.5 ⁴	4.0	4.1	7.4	3.7
Sample 3	6.7	4.5	4.7	7.0	3.6
Sample 4	6.6	4.5	4.7	7.1	3.6
Sample 4	6.7	4.5	4.8	5.7	2.6

¹Neat foam without soils - 15 squeezes.

²Neat foam with soils added - 10 additional squeezes.

³Minimum significant difference.

⁴Means having the same letter are not significantly different (alpha = 0.10).

TABLE 7c

Product	Gr on glass	Gr on plate	Gr on plastic	Gr on knife	tub
Sample 1	0.32	0.17	0.29	0.22	0.40
Sample 2	1.1	0.3	0.8	0.4	1.7
Sample 3	0.9	0.3	0.8	0.5	1.3
Sample 4	0.9	0.4	0.8	0.4	1.4
Sample 4	0.7	0.2	0.6	0.3	1.0

TABLE 7d

Product	DLRIN (Ease of rinse) ¹ Glass	Ease of Rinse Plate (sec) ²	Amount of water film (%) ³	Neat ⁴ Glass	Neat ⁴ Plate
Sample 1	0.49 ⁵	1.04	10.73	0.74	0.70
Sample 2	6.2 ⁶	8.3	46.9	1.9	1.2
Sample 3	6.2	10.0	42.4	2.1	1.9
Sample 4	6.3	9.7	50.2	2.6	1.7
Sample 4	6.8	13.0	48.0	2.6	1.8

¹Number of rinses till no foam.

²Seconds needed to rinse detergent off plate.

³Percent water film on plate.

⁴Detergent residue remaining on dishes.

⁵Minimum significant difference.

⁶Means having the same letter are not significantly different (alpha = 0.10).

26

TABLE 7e

Product	Fat soils remaining
Sample 1	2.4
Sample 2	2.5
Sample 3	2.3
Sample 4	1.58

Example 8

The compositions exhibit stability at reduced and increased temperatures. More specifically, such compositions remain clear and stable in the range of about 0° C. to about 50° C.

Creep Yield Stress Test (static test)—This rheological test was conducted on the TA Instruments ARG2 rheometer. It uses the high surface area vane geometry that is very sensitive and can measure very low yield stresses. The test is run in a 50-gram water-jacketed sample holder at a constant temperature of 25° C. The test runs a creep test (strain vs. stress) at stresses ranging from 0.01 Pa to 0.6 Pa. Custom software then calculates yield stress from the family of curves generated for each sample. A yield stress above 0.5 dyn/cm² is ideal for supporting particulates, but a yield stress above 0.15 dyn/cm² is sufficient to justify product positioning as a gel or dish gel.

Brookfield Yield Stress Test (dynamic test)—This test was developed to approximate the creep analysis above, but with much more rapid output to provide rapid feedback when processing formulas at manufacturing conditions. It also uses the high surface area vane geometry that is very sensitive and can measure very low yield stresses. The test is run in a 400-milliliter glass beaker. The test runs a torque sweep at decreasing RPMs, or revolutions per minute, ranging from 50 to 0.3 rpm. Once the torques are recorded, custom software then calculates yield stress for each sample. A yield stress above 0.5 dyn/cm² is ideal for supporting particulates, but a yield stress above 0.2 dyn/cm² is sufficient to justify product positioning as a gel or dish gel.

ARG2 Viscosity Test—This rheological test was conducted on the ARG2 rheometer. It simply measures viscosity at a constant shear rate of 21 s⁻¹ with a constant temperature of 25° C. This test simulates the shear rate of the product coming out of the bottle when the consumer dispenses the product under normal conditions. If the value is above 2000 cP, the cap orifice may need to be modified to assure consumer-friendly dispensing.

TABLE 8a

Ingredient	Ex. Ultra acidic LDL gel #1 (wt. %)	Ex. Ultra acidic LDL gel #2 (wt. %)	Ex. Ultra acidic LDL gel #3 (wt. %)
Na DBS ¹	5.75	5.75	5.75
SLES (2EO) C ₁₂ -C ₁₄	19.6	19.6	19.6
Laurylamidopropylbetaine	6.6	6.6	6.6
Alcohol	4	4	4
Lactic Acid	2	2	2
SXS	2.5	2.5	2.5
Gellan Gum	0.075	0.094	0.125
Na ₄ EDTA	0.83	0.83	0.83
Water	qs	qs	qs

¹Sodium dodecyl benzene sulfonate

TABLE 8b

Ultra Acidic LDL Gel	Brookfield Viscosity (cP)	ARG2 Viscosity (cP)	ARG2 Creep Yield Stress (dyn/cm ²)	Brookfield Yield Stress (dyn/cm ²)
#1	773	1472	0.16	0.231
#2	768	1388	0.40	0.859
#3	905	1778	— ¹	1.046

¹Surface skinning over testing creep test timing cycle yielded invalidated data

The liquid compositions are readily pourable and exhibit a viscosity in the range of 6 to 300 milliPascal second (mPas or mps) as measured at 25° C. with a Brookfield RVTDV-II Viscometer using a #21 spindle rotating at 20 RPM. In certain embodiments, the viscosity is maintained in the range of 10 to 200 mPas.

Example 9

The compositions of the invention are nearly colorless. The relative amount of a coloring agent to deliver near-colorless aesthetics is mainly dependent on the color of the dodecyl benzene sulfonate being used. Color is measured on a Klett scale where the higher the Klett the more yellow a particular material used. The next table gives an approximate amount of color needed to deliver the aesthetic according to Klett of NaLAS. The formulation uses a mixture of violet and pink dyes to yield the final color aesthetic. The colors are chosen based on the color wheel. Violet dye is added to offset the light yellow color present in the base. Because it may be an imperfect match a slight green color can be generated which is accounted for with a pink colorant. The net result is a product that has an appearance of a colorless material. The composition has a color on a Klett scale of about 0.1 to about 20.

TABLE 9

Dodecyl Benzene Sulfonate Color	Wt. % Coloring Agent Added
0-5	<0.0035
6-10	0.0035
10-15	0.006
15-20	0.008
20-25	0.01

Example 10

Toxicity Testing was conducted using animal studies and alternative tests. Animal studies were completed using Table 1a formula Ultra 2a/b (with or without fragrance).

The alternate test methodologies (human testing) were done for the Inventive formula as listed in Table 1d. Studies were completed with 4 different formula/fragrance options, but the base formula is 1d. The Toxicity Testing was conducted using the following test protocols:

(1) Acute Oral Toxicity—Exposure is via a single, limit dose of dish liquid at the maximum required upper limit dose of 5000 mg/kg. The Acute Oral Toxicity was conducted using OPPTS Guideline Study 870.1100, EPA Publication #98-190. The LD₅₀ was not reached and was greater than 5000 mg/kg.

(2) Acute Dermal Toxicity—Exposure is via a single, limit dose of dish liquid at the upper limit dose of 5000 mg/kg. The Acute Dermal Toxicity was conducted using OPPTS Guideline Study 870.1200, EPA Publication # 98-192. The LD₅₀ was not reached and was greater than 5000 mg/kg.

(3) Acute Eye Irritation—Exposure is via a single 10 μl dose with scoring for irritation at fixed intervals after exposure. The Acute Eye Irritation was conducted using a lower volume of test material placed directly on the eye. The amount of test material used in the LVET is 1/10th of that used in the Draize eye irritation test. There was some initial irritation that fully reversed within the 7 day scoring endpoint, and no corneal opacity.

(4) Acute Dermal Irritation—Exposure is via repeated, occluded, prolonged exposure to concentrated (undiluted) dish liquid. The Acute Dermal Irritation was conducted using OPPTS Guideline Study 870.2500, EPA Publication #98-196. Results showed irritation within the 72 hour period that fully reversed with seven days. A more relevant measure skin irritation uses the standard 21 -day Cumulative Irritation study methodology (applied commonly in the cosmetic industry) on humans. This method is a semi-occluded exposure at a relevant product use-dilution, although it is still clearly an extreme/maximal exposure scenario. This method shows no significant irritation for lactic acid based formulations.

(5) Skin Sensitization—Exposure is via three weekly induction doses and then a challenge dose (following a 2 week intervening rest period). The Skin Sensitization was conducted using OPPTS Guideline Study 870.2600, EPA Publication #98-197. A naïve control group is used as a comparison for the group receiving the challenge dose. Result is that the formula is a non-sensitizer. A more relevant measure of sensitization potential uses the Human Repeat Insult Patch Test methodology (applied commonly in the cosmetic industry) on humans. This is an occlusive patch exposure method at a relevant product use-dilution. This method also documents no skin sensitization for the sample.

All of the references cited and appended hereto, including patents, patent applications, literature publications, and the like, are hereby incorporated in their entireties by reference.

What is claimed is:

1. A colorless surfactant based cleaning composition comprising a surfactant system and at least one organic acid or salt thereof, and a combination of violet and pink dyes such that the composition has a color on a Klett scale of about 0.1 to about 20.

2. The composition of claim 1, wherein the cleaning composition comprises a viscosity of about 10 to about 1,000 cps at 25° C. using a #21 spindle at 20 rpm measured on a Brookfield RVTDV-II viscometer.

3. The composition of claim 1, wherein the composition has a Baumgartner grease removal of at least about 30, wherein the composition has about 50 ppm to about 150 ppm hardness water.

4. The composition of claim 1, wherein the composition has a log₁₀ reduction in microbes of about 3 when a surface containing microbes is contacted with the composition for about 30 seconds at 25° C.

5. The surfactant based composition of claim 1, wherein the organic acid is lactic acid.

6. The surfactant based cleaning composition of claim 1, wherein the surfactant system includes an anionic surfactant and the anionic surfactant is a C₁₀-C₁₄ LA sulfonate, sodium lauryl ether sulfate with about two ethylene oxide units, or combinations thereof.

7. The surfactant based cleaning composition of claim 1, wherein the surfactant system includes a zwitterionic surfactant that is laurylamidopropyl betaine.

8. The surfactant based cleaning composition of claim 1, wherein one or more of the surfactants are derived from a natural source and biodegradable surfactants.

29

9. The composition of claim 1, wherein

A) the surfactants are a combination comprising

(i) sodium dodecyl benzene sulfonate, which is present in an amount of about 3 wt. % to about 20 wt. % or about 2 wt. % to about 9 wt. % by weight of the total composition, 5

(ii) sodium lauryl ether sulfate with about two EO units, which is present in an amount of about 3 wt. % to about 20 wt. % by weight of the total composition,

(iii) laurylamidopropyl betaine, which is present in an amount of about 1 wt. % to about 8 wt. % by weight of the total composition, 10

B) the at least one organic acid is at least one acid chosen from lactic acid, formic acid, citric acid, sorbic acid, acetic acid, glycolic acid, propanoic acid, propionic acid, oxalic acid, maleic acid, tartaric acid, adipic acid, malic acid, malonic acid, and glycolic acid, and is present in an amount of about 1 wt. % to about 3 wt. % by weight of the total composition,

C) optionally comprising one or more of each of the following: thickener, solvent, suspending agent, viscosity

30

modifier, hydrotrope, fragrance agent, preservative, diluent, chelating agent, coloring, water, fluorescent whitening agent, photobleach, fiber lubricant, reducing agent, enzyme, enzyme stabilizing agent, powder finishing agent, builder, bleach, bleach catalyst, soil release agent, dye transfer inhibitor, buffer, pro-fragrance, anti-ashing polymer, soil repellent, water-resistance agent, aesthetic agent, structuring agent, sanitizer, fabric finishing agent, dye fixative, fabric conditioning agent, deodorizer or combination thereof

10. The composition of claim 9, wherein the solvent is present.

11. The composition of claim 9, wherein the thickener is present.

12. The composition of claim 9, wherein the suspending agent is present. 15

13. A method of cleaning a surface comprising contacting the surface with a composition of claim 1.

* * * * *