

US008143488B2

(12) **United States Patent**
Burns et al.

(10) **Patent No.:** **US 8,143,488 B2**
(45) **Date of Patent:** **Mar. 27, 2012**

(54) **PLANTS AND SEEDS OF SPRING CANOLA VARIETY SCV470336**

(75) Inventors: **Dale Burns**, Lethbridge (CA); **Scott McClinchey**, Orton (CA); **Zenon Lisieczko**, Guelph (CA)

(73) Assignee: **Monsanto Technology LLC**, St. Louis, MO (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 224 days.

(21) Appl. No.: **12/713,594**

(22) Filed: **Feb. 26, 2010**

(65) **Prior Publication Data**

US 2011/0212248 A1 Sep. 1, 2011

(51) **Int. Cl.**

A01H 5/00 (2006.01)

A01H 5/10 (2006.01)

A01H 1/00 (2006.01)

C12N 15/82 (2006.01)

(52) **U.S. Cl.** **800/306; 800/260; 435/430**

(58) **Field of Classification Search** None
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

3,710,511 A	1/1973	Patterson
3,861,709 A	1/1975	Mulcahy et al.
4,535,060 A	8/1985	Comai
4,654,465 A	3/1987	Brar et al.
4,658,085 A	4/1987	Beversdorf et al.
4,727,219 A	2/1988	Brar et al.
4,769,061 A	9/1988	Comai
4,810,648 A	3/1989	Stalker
4,940,835 A	7/1990	Shah et al.
4,971,908 A	11/1990	Kishore et al.
4,975,374 A	12/1990	Goodman et al.
5,015,580 A	5/1991	Christou et al.
5,034,323 A	7/1991	Jorgensen et al.
5,107,065 A	4/1992	Shewmaker et al.
5,145,783 A	9/1992	Kishore et al.
5,188,642 A	2/1993	Shah et al.
5,266,317 A	11/1993	Tomalski et al.
5,310,667 A	5/1994	Eischholtz et al.
5,312,910 A	5/1994	Kishore et al.
5,322,783 A	6/1994	Tomes et al.
5,432,068 A	7/1995	Albertsen et al.
5,453,566 A	9/1995	Shewmaker et al.
5,463,175 A	10/1995	Barry et al.
5,478,369 A	12/1995	Albertsen et al.
5,491,288 A	2/1996	Chaubet et al.
5,494,813 A	2/1996	Hepher et al.
5,510,471 A	4/1996	Lebrun et al.
5,559,223 A	9/1996	Falco et al.
5,563,055 A	10/1996	Townsend et al.
5,607,914 A	3/1997	Rao et al.
5,627,061 A	5/1997	Barry et al.
5,633,435 A	5/1997	Barry et al.
5,633,436 A	5/1997	Wandelt
5,633,448 A	5/1997	Lebrun et al.
5,759,829 A	6/1998	Shewmaker et al.
5,767,373 A	6/1998	Ward et al.

5,776,760 A	7/1998	Barry et al.
5,789,566 A	8/1998	Bonhomme et al.
5,792,931 A	8/1998	Duvick et al.
5,804,425 A	9/1998	Barry et al.
5,824,524 A	10/1998	Albertsen et al.
5,850,014 A	12/1998	Albertsen et al.
5,850,016 A	12/1998	Jung et al.
5,859,341 A	1/1999	Albertsen et al.
5,866,775 A	2/1999	Eichholtz et al.
5,885,801 A	3/1999	Rao
5,885,802 A	3/1999	Rao
5,891,859 A	4/1999	Thomashow et al.
5,892,009 A	4/1999	Thomashow et al.
5,912,414 A	6/1999	Falco et al.
5,929,305 A	7/1999	Thomashow et al.
5,939,599 A	8/1999	Chui et al.

(Continued)

FOREIGN PATENT DOCUMENTS

EP 0333033 A1 9/1989

(Continued)

OTHER PUBLICATIONS

Grahame E. Pratt, et al., Identification of an Allatostatin from Adult *Diptera Punctata*, Received Jul. 18, 1989, Biochemical and Biophysical Research Communications, vol. 163, No. 3, Sep. 19, 1989, pp. 1243-1247. 0006-291X/89 \$.50. Copyright 1989 by Academic Press, Inc. All rights of reproduction in any form reserved.

Elisabeth Przbilla, et al., Site-Specific Mutagenesis of the D1 Subunit of Photosystem II in Wild-Type *Chlamydomonas*, accepted Nov. 16, 1990, *The Plant Cell*, vol. 3, 169-174, Feb. 1991. Copyright 1991 American Society of Plant Physiologists.

V. Raboy, et al., A survey of maize kernel mutants for variation in phytic acid, *Maydica* 35 (1990): 383-390.

John C. Sanford, et al., Delivery of Substances into Cells and Tissues Using a Particle Bombardment Process, *Particulate Science and Technology* 5:27-37, Copyright 1987 by Hemisphere Publishing Corporation.

John C. Sanford, Biolistic plant transformation, revised Oct. 18, 1989, *Physiologia Plantarum* 79: 206-209. Copenhagen 1990.

(Continued)

Primary Examiner — Elizabeth McElwain

(74) *Attorney, Agent, or Firm* — Nelson Mullins Riley & Scarborough, LLP

(57) **ABSTRACT**

The invention relates to a novel canola line designated as SCV470336. The invention also relates to the seeds, the plants, and the plant parts of canola line SCV470336 as well as to methods for producing a canola plant produced by crossing canola line SCV470336 with itself or with another canola line. The invention also relates to methods for producing a canola plant containing in its genetic material one or more transgenes and to the transgenic canola plants and plant parts produced by those methods. The invention further relates to canola lines or breeding lines and plant parts derived from canola line SCV470336, to methods for producing other canola lines or plant parts derived from canola line SCV470336 and to the canola plants, varieties, and their parts derived from use of those methods. The invention additionally relates to hybrid canola seeds, plants, and plant parts produced by crossing the line SCV470336 with another canola line.

26 Claims, No Drawings

U.S. PATENT DOCUMENTS

5,959,185	A	9/1999	Streit et al.
5,965,705	A	10/1999	Thomashow et al.
5,973,233	A	10/1999	Burns et al.
5,973,234	A	10/1999	Mueller et al.
5,977,445	A	11/1999	Soper et al.
5,990,389	A	11/1999	Rao et al.
RE36,449	E	12/1999	Lebrun et al.
6,040,497	A	3/2000	Spencer et al.
6,063,947	A	5/2000	DeBonte et al.
6,080,913	A	6/2000	Tarczynski et al.
6,084,153	A	7/2000	Good et al.
6,107,547	A	8/2000	Coruzzi et al.
6,118,055	A	9/2000	Livesey
6,127,600	A	10/2000	Beach et al.
6,130,366	A	10/2000	Herrera-Estrella et al.
6,162,967	A	12/2000	Webb
6,177,275	B1	1/2001	Coruzzi et al.
6,194,638	B1	2/2001	Dhugga et al.
6,197,561	B1	3/2001	Martino-Catt et al.
6,225,114	B1	5/2001	Eichholtz et al.
6,229,072	B1	5/2001	Burns et al.
6,232,529	B1	5/2001	Singletary et al.
6,248,876	B1	6/2001	Barry et al.
RE37,287	E	7/2001	Lebrun et al.
6,265,640	B1	7/2001	Albertsen et al.
6,282,837	B1	9/2001	Ward et al.
6,288,306	B1	9/2001	Ward et al.
6,291,224	B1	9/2001	Martin-Catt et al.
6,297,426	B1	10/2001	Albertsen et al.
6,307,126	B1	10/2001	Harberd et al.
6,323,392	B1	11/2001	Charne
6,338,961	B1	1/2002	DeRose et al.
6,346,403	B1	2/2002	Rafalski et al.
6,391,348	B1	5/2002	Stilborn et al.
6,399,859	B1	6/2002	Nichols et al.
6,417,428	B1	7/2002	Thomashow et al.
6,423,886	B1	7/2002	Singletary et al.
6,441,274	B1	8/2002	Cahoon et al.
6,459,019	B1	10/2002	Falco et al.
6,563,020	B1	5/2003	Simmons et al.
6,566,587	B1	5/2003	Lebrun et al.
6,573,430	B1	6/2003	Bradley et al.
6,664,445	B1	12/2003	Falco et al.
6,664,446	B2	12/2003	Heard et al.
6,706,866	B1	3/2004	Thomashow et al.
6,713,663	B2	3/2004	Weigel et al.
6,717,034	B2	4/2004	Jiang
6,787,683	B1	9/2004	Penna et al.
6,794,560	B2	9/2004	Harberd et al.
6,801,104	B2	10/2004	Zhu et al.
6,803,498	B2	10/2004	Dhugga et al.
6,825,397	B1	11/2004	Lowe et al.
6,858,778	B1	2/2005	Jung et al.
6,875,907	B2	4/2005	Simmons et al.
6,911,577	B2	6/2005	Simmons et al.
6,930,225	B2	8/2005	Dhugga et al.
6,992,237	B1	1/2006	Habben et al.
7,067,720	B2	6/2006	Shi et al.
7,087,810	B2	8/2006	Muller et al.
7,098,381	B2	8/2006	Singletary et al.
7,135,615	B2	11/2006	Kato
7,145,060	B2	12/2006	Muller et al.
7,154,029	B2	12/2006	Cahoon et al.
7,157,621	B2	1/2007	Allen et al.
7,179,955	B2	2/2007	Dhugga et al.
7,205,453	B2	4/2007	Altier et al.
7,288,386	B2	10/2007	Lightfoot et al.
7,462,481	B2	12/2008	Castle et al.
7,531,723	B2	5/2009	Habben et al.
7,723,582	B2	5/2010	Chungu et al.
7,947,877	B2	5/2011	Lisieczko
7,964,744	B2	6/2011	Nara et al.
8,071,848	B2 *	12/2011	Lisieczko 800/306
2001/0037515	A1	11/2001	Patel
2003/0079247	A1	4/2003	Shi et al.
2003/0166197	A1	9/2003	Ecker et al.
2004/0068767	A1	4/2004	Dhugga et al.
2004/0078852	A1	4/2004	Thomashow et al.

2004/0098764	A1	5/2004	Heard et al.
2004/0128719	A1	7/2004	Klee et al.
2004/0148654	A1	7/2004	Helentjaris
2005/0160488	A1	7/2005	Jung et al.
2005/0204418	A1	9/2005	Jung et al.
2005/0235383	A1	10/2005	Patel
2005/0278812	A1	12/2005	Patel
2006/0075516	A1	4/2006	Kubik
2006/0075517	A1	4/2006	Kubik
2006/0225146	A1	10/2006	Kubik
2006/0225156	A1	10/2006	Jonsson
2006/0225157	A1	10/2006	Jonsson
2006/0225158	A1	10/2006	Jonsson
2006/0225159	A1	10/2006	Kubik
2007/0089181	A1	4/2007	Patel
2008/0256653	A1	10/2008	Chungu et al.
2008/0256654	A1	10/2008	Chungu et al.
2008/0256655	A1	10/2008	Chungu et al.
2008/0263714	A1	10/2008	Chungu et al.
2008/0263715	A1	10/2008	Chungu et al.
2008/0263716	A1	10/2008	Chungu et al.
2008/0263717	A1	10/2008	Chungu et al.
2009/0064359	A1	3/2009	Lisieczko
2009/0285967	A1	11/2009	Lisieczko
2009/0285968	A1	11/2009	Lisieczko
2009/0288184	A1	11/2009	Lisieczko
2009/0288185	A1	11/2009	Lisieczko et al.

FOREIGN PATENT DOCUMENTS

EP	0242236	A1	10/1990
EP	0599042	A1	10/1993
EP	1173580	A1	1/2002
EP	1173581	A1	1/2002
EP	1173582	A1	1/2002
JP	2002281975	A	10/2002
WO	92/05251	A1	4/1992
WO	92/13956	A1	8/1992
WO	92/13957	A1	8/1992
WO	93/02197	A1	2/1993
WO	93/11245	A1	6/1993
WO	94/00992	A1	1/1994
WO	95/15392	A1	6/1995
WO	96/38560	A2	6/1995
WO	95/18855	A2	7/1995
WO	96/01905	A1	1/1996
WO	96/14414	A1	5/1996
WO	96/30530	A1	10/1996
WO	97/02737	A1	1/1997
WO	97/10339	A1	3/1997
WO	97/29123	A2	8/1997
WO	97/49811	A1	12/1997
WO	98/09521	A1	3/1998
WO	98/20133	A3	5/1998
WO	98/22604	A1	5/1998
WO	98/27806	A1	7/1998
WO	98/42831	A2	10/1998
WO	98/45448	A1	10/1998
WO	98/45458	A1	10/1998
WO	98/53083	A1	11/1998
WO	98/56918	A1	12/1998
WO	98/56935	A2	12/1998
WO	99/05298	A1	2/1999
WO	99/09174	A1	2/1999
WO	99/10498	A2	3/1999
WO	99/25821	A1	5/1999
WO	99/25853	A1	5/1999
WO	99/29882	A1	6/1999
WO	99/38977	A2	8/1999
WO	99/40209	A1	8/1999
WO	99/49064	A2	9/1999
WO	99/53050	A1	10/1999
WO	99/55882	A1	11/1999
WO	00/09706	A2	2/2000
WO	00/31964	A1	6/2000
WO	00/32761	A1	6/2000
WO	00/42219	A1	7/2000
WO	00/44918	A1	8/2000
WO	00/46358	A2	8/2000
WO	00/60089	A1	10/2000

WO	00/68393	A1	11/2000
WO	00/73475	A1	12/2000
WO	01/04147	A2	1/2001
WO	01/12800	AS	2/2001
WO	01/12825	A1	2/2001
WO	01/21822	A1	3/2001
WO	01/26459	A2	4/2001
WO	01/29237	A2	4/2001
WO	01/34726	A2	5/2001
WO	01/35725	A1	5/2001
WO	01/35727	A1	5/2001
WO	01/36444	A1	5/2001
WO	01/36596	A2	5/2001
WO	01/36598	A1	5/2001
WO	01/52620	A2	7/2001
WO	01/64898	A2	9/2001
WO	01/66704	A2	9/2001
WO	01/79516	A2	10/2001
WO	02/15675	A1	2/2002
WO	02/17430	A1	2/2002
WO	02/42424	A2	5/2002
WO	02/057439	A2	7/2002
WO	02/059324	A2	8/2002
WO	02/02776	A1	10/2002
WO	02/077185	A2	10/2002
WO	02/079403	A2	10/2002
WO	03/000863	A2	1/2003
WO	03/011015	A2	2/2003
WO	03/013227	A2	2/2003
WO	03/013228	A2	2/2003
WO	03/027243	A2	4/2003
WO	03/048345	A1	6/2003
WO	03/052063	A2	6/2003
WO	03/076574	A2	9/2003
WO	03/082899	A2	10/2003
WO	2004/031349	A2	4/2004
WO	2004/076638	A2	9/2004
WO	2004/009143	A2	10/2004

OTHER PUBLICATIONS

John C. Sanford, Reviews, The biolistic process, TIBTECH, Dec. 1988 [vol. 6], pp. 299-302, Copyright 1988, Elsevier Science Publishers Ltd. (UK) 0167-9430/88/\$02.00.

Mark Schena, et al., A steroid-inducible gene expression system for plant cells, Contributed by Ronald W. Davis, Aug. 8, 1991, Proc. Natl. Acad. Sci. USA, vol. 88, pp. 10421-10425, Dec. 1991, Genetics.

Champa Sengupta-Gopalan, et al., Developmentally regulated expression of the bean B-phaseolin gene in tobacco seed, Communicated by Arthur Kelman, Jan. 14, 1985, Proc. Natl. Acad. Sci. USA, vol. 82, pp. 3320-3324, May 1985, Developmental Biology.

Dilip M. Shah, et al., Engineering Herbicide Tolerance in Transgenic Plants, Science, vol. 233, pp. 478-481, Dec. 12, 1985; accepted May 6, 1986.

Phillip A. Sharp, RNAi and double-strand RNA, Genes Dev. 1999 13: 139-141 Cold Spring Harbor Laboratory Press. To subscribe to Genes & Development go to: <http://genesdev.cshlp.org/subscriptions>.

Raymond E. Sheehy, et al., Reduction of polygalacturonase activity in tomato fruit by antisense RNA, Communicated by E. Peter Geiduschek, Aug. 4, 1988, Proc. Natl., Acad. Sci. USA, vol. 85, pp. 8805-8809, Dec. 1988, Biochemistry.

Teruaki Shiroza, et al., Sequence Analysis of the Streptococcus mutans Fructosyltransferase Gene and Flanking Regions, Accepted Nov. 19, 1987, Journal of Bacteriology, Feb. 1988, p. 810-816. 0021-9193/88/020810-07\$02.00/0. Copyright 1988, American Society for Microbiology.

June Simpson, et al., Light-inducible and tissue-specific expression of a chimaeric gene under control of the 5' flanking sequence of a pea chlorophyll a/b-binding protein gene, The EMBO Journal, vol. 4, No. 11, pp. 2723-2729, 1985. Copyright IRL Press Limited, Oxford, England.

Neil A. Smith, et al., Gene expression, Total silencing by intron-spliced hairpin RNAs, Nature, vol. 407, Sep. 21, 2000, pp. 319-320, www.nature.com Copyright Macmillan Magazines Ltd.

Morten Sogaardt, et al., Site-directed Mutagenesis of Histidine 93, Aspartic Acid 180, Glutamic Acid 205, Histidine 290, and Aspartic

Acid 291 at the Active Site and Tryptophan 279 at the Raw Starch Binding in Barley α -Amylase I, The Journal of Biological Chemistry, vol. 268, No. 30, Issue of Oct. 25, pp. 22480-22484, 1993, Copyright 1993 by The American Society for Biochemistry and Molecular Biology, Inc. Printed in U.S.A.

Imre E. Somssich, Previews, Closing Another Gap in the Plant SAR Puzzle, Cell, 2003, pp. 815-816.

Cheryl Montain Laursen, et al., Production of fertile transgenic maize by electroporation of suspension culture cells, accepted in revised form Sep. 17, 1994, Plant Molecular Biology 24: 51-61, Copyright 1994 Kluwer Academic Publishers. Printed in Belgium.

David M. Stalker, et al., Herbicide Resistance in Transgenic Plants Expressing a Bacterial Detoxification Gene, Apr. 29, 1988; accepted Aug. 9, 1988, Reports, Oct. 21, 1988, pp. 419-423.

Virginia Stiefel, et al., Expression of a Maize Cell Wall Hydroxyproline-Rich Glycoprotein Gene in Early Leaf and Root Vascular Differentiation, revised Jun. 8, 1990, The Plant Cell, vol. 2, 785-793, Aug. 1990. Copyright 1990 American Society of Plant Physiologists.

Peter Steinecke, et al., Expression of a chimeric ribozyme gene results in endonucleolytic cleavage of target mRNA and a concomitant reduction of gene expression in vivo, revised on Dec. 20, 1991, The EMBO Journal, vol. 11, No. 4, pp. 1525-1530, 1992. Copyright Oxford University Press.

Michel Steinmetz, et al., The DNA sequence of the gene for the secreted *Bacillus subtilis* enzyme levansucrase and its genetic control sites, Mol. Gen. Genet (1985) 200: 220-228. Copyright Springer-Verlag 1985.

Jun-Ichi Sumitani, et al., Molecular Cloning and Expression of Proteinaceous β -Amylase Inhibitor Gene from *Streptomyces nitrosporeus* in *Escherichia coli*, Biosci. Biotech. Biochem., 57 (8), 1243-1248, 1993.

Zora Svab, et al., Aminoglycoside-3" -denyltransferase confers resistance to spectinomycin and streptomycin in *Nicotiana tabacum*, accepted Sep. 19, 1989, Plant Molecular Biology 14: 197-205, Copyright 1990 Kluwer Academic Publishers. Printed in Belgium.

Paraskevi Taviadoraki, et al., Transgenic plants expressing a functional single-chain Fv antibody are specifically protected from virus attack, Letters to Nature, Nature, vol. 366, Dec. 2, 1993, pp. 469-472. Copyright 1993 Nature Publishing Group.

Crispin B. Taylor, In this issue: Comprehending Cosuppression, The Plant Cell, vol. 9, 1245-1249, Aug. 1997.

T.H. Teeri, et al., Gene fusions to lacZ reveal new expression patterns of chimeric genes in transgenic plants, revised on Dec. 2, 1988, The EMBO Journal, vol. 8, No. 2, pp. 343-350, 1989. Copyright IRL Press.

M. P. Timko, et al., Letters to Nature, Light regulation of plant gene expression by an upstream enhancer-like element, Received Jun. 13; accepted Oct. 9, 1985, Nature, vol. 318, Dec. 12, 1985, pp. 579-582. Copyright 1985 Nature Publishing Group.

Patrick Toubart, et al., Cloning and characterization of the gene encoding the endopolygalacturonase-inhibiting protein (PGIP) of *Phaseolus vulgaris* L., revised Jan. 2, 1992, The Plant Journal (1992) 2 (3), 367-373.

Kazuhiro Toyoda, et al., Review: Resistance and susceptibility of plants to fungal pathogens, Transgenic Research 11: 567-582, 2002. Copyright 2002 Kluwer Academic Publishers. Printed in the Netherlands.

David Twell, et al., Isolation and expression of an anther-specific gene from tomato, Received Nov. 20, 1988, Mol. Gen. Genet (1989) 217: 240-245. Copyright Springer-Verlag 1989.

David Twell, et al., Activation and developmental regulation of an Arabidopsis anther-specific promoter in microspores and pollen of *Nicotiana tabacum*, Sexual Plant Reproduction (1993), 6: 217-224. Copyright Springer-Verlag 1993.

Els J. M. Van Damme, et al., Short Communication: Molecular cloning of mannose-binding lectins from *Clivia miniata*, accepted in revised form Jan. 24, 1994, Plant Molecular Biology 24: 825-830, 1994. Copyright 1994 Kluwer Academic Publishers. Printed in Belgium.

- Wim Van Hartingsveldt, et al., Gene 07019, Cloning, characterization and overexpression of the phytase-encoding gene (phyA) of *Aspergillus niger*, Received by J. R. Kinghorn: Oct. 20, 1992; Revised/Accepted: Dec. 10/Dec. 12, 1992; received at publishers: Jan. 6, 1993, Gene, 127 (1993) 87-94. Copyright 1993 Elsevier Science Publishers B.V. All rights reserved. 0378-1119/93/\$06.00.
- Peter J. M. Van Den Elzen, et al., Short Communication: A chimaeric hygromycin resistance gene as a selectable marker in plant cells, in revised form Jul. 17, 1985; accepted Jul. 27, 1985, Plant Molecular Biology 5: 299-302, 1985. Copyright Martinus Nijhoff Publishers, Dordrecht—Printed in The Netherlands.
- L. Verdoodt, et al., Use of the multi-allelic self-incompatibility gene in apple to assess homozygosity in shoots obtained through haploid induction, Accepted: Sep. 10, 1997, Theor Appl Genet (1998) 96: 294-300. Copyright Springer-Verlag 1998.
- Vicki Chandler, Chapter 118, Overview of Cloning Genes Using Transposon Tagging, The Maize Handbook—M Freeling, V. Walbott, eds., pp. 647-652. Copyright 1994 Springer-Verlag, New York, Inc. Supplied by the British Library—"The world's knowledge".
- Y. Wan, et al., Efficient production of doubled haploid plants through colchicine treatment of anther-derived maize callus, Accepted Nov. 28, 1988, Theor Appl Genet (1989) 77: 889-892, Copyright Springer-Verlag 1989.
- David G. Wang, et al., Reports, Large-Scale Identification, Mapping, and Genotyping of Single-Nucleotide Polymorphisms in the Human Genome, Feb. 5, 1998; accepted Mar. 31, 1998, Science, vol. 280, May 15, 1998, pp. 1077-1082. www.sciencemag.org.
- E. R. Ward, et al., Update section, Mini review, Chemical regulation of transgene expression in plants, Plant Molecular Biology 22: 361-366, 1993. Copyright 1993 Kluwer Academic Publishers. Printed in Belgium.
- Phillip D. Zamore, et al., RNAi: Double-Stranded RNA Directs the ATP-Dependent Cleavage of mRNA at 21 to 223 Nucleotide Intervals, Cell, vol. 101, 25-33, Mar. 31, 2000. Copyright 2000 by Cell Press.
- Li-Jian Zhang, et al. Efficient Transformation of Tobacco by Ultrasonication, accepted Jun. 14, 1991, Copyright 1991 Nature Publishing Group. <http://www.nature.com/naturebiotechnology>.
- Final Office Action, mailed Jan. 26, 2011, from United States Patent and Trademark Office, U.S. Appl. No. 12/120,297, filed May 14, 2008, Zenon Lisieczko.
- Response to Final Office Action, filed Feb. 4, 2011, with the United States Patent and Trademark Office, in response to the Final Office Action, mailed Jan. 26, 2011, U.S. Appl. No. 12/120.297, filed May 14, 2008, Zenon Lisieczko.
- Roger N. Beachy, et al., Coat Protein-Mediated Resistance Against Virus Infection, Annu. Rev. Phytopathol. 1990, 28:451-74. Copyright 1990 by Annual Reviews Inc. 0066-4286/90/0901-0451\$02.00.
- Thomas W. Becker, et al., The cab-m7 gene: a light-inducible, mesophyll-specific gene of maize, accepted in revised form Apr. 16, 1992, Plant Molecular Biology 20: 49-60, 1992. Copyright 1992 Kluwer Academic Publishers. Printed in Belgium.
- R. Bernardo, et al., North American study of essential derivation in maize: inbreds developed without and with selection from F2 populations, Accepted: Jul. 28, 2000, Theor Appl Genet (2001) 102: 986-992. Copyright Springer-Verlag 2001.
- Jose R. Botella, et al., Differential expression of two calmodulin genes in response to physical and chemical stimuli, accepted in revised form Jan. 12, 1994. Plant Molecular Biology 24: 757-766, 1994. Copyright 1994 Kluwer Academic Publishers. Printed in Belgium.
- Steven P. Briggs, Plant Disease Resistance, Grand unification theory in sight, Molecular characterization of the components of signalling pathways that mediate disease resistance is at least providing a unified picture of how plants fight disease. Copyright Current Biology 1995, vol. 5, No. 2., pp. 128-131.
- Rachel A. Burton, et al., Virus-Induced Silencing of a Plant Cellulose Synthase Gene, The Plant Cell, vol. 12, 691-705, May 2000. www.plantcell.org Copyright 2000 American Society of Plant Physiologists. Copyright American Society of Plant Biologists. Advancing the Science of Plant Biology.
- W. R. Busnell, et al., Genetic engineering of disease resistance in cereals, , The Canadian Phytopathological Society, vol. 20(2):137-220 Jun. 1998, ISSN 0706-0661. Accepted for publication Mar. 30, 1998. This paper was presented at a symposium on bioengineering plants for disease resistance, held at the annual meeting of the Canadian Phytopathological Society, Jul. 6-9, 1997. Canadian Journal of Plant Pathology, Online publication date: Dec. 22, 2009. To link to this Article: DOI: 10.1080/07060669809500419. Publisher Taylor & Francis.
- Martin Chalfie, et al., Green Fluorescent Protein as a Marker for Gene Expression, Source: Science, New Series, vol. 263, No. 5148 (Feb. 11, 1994), pp. 802-803. Published by: American Association for the Advancement of Science. Stable URL: <http://www.jstor.org/stable/2882924> Accessed: Apr. 24, 2008 19:27.
- S. T. Chalyk, et al., Transgressive segregation in the progeny of a cross between two inducers of maize maternal haploids, Maize Genetics Cooperation News Letter (Journal Article), Std. No. 10904573, Issue 68, p. 47, Pub Date: 1994.
- Pierre J. Charest, et al., In vitro study of transgenic tobacco expressing *Arabidopsis* wild type and mutant acetohydroxyacid synthase genes. Received Jun. 28, 1989/Revised version received Sep. 15, 1989—Communicated by F. Constabel. Plant Cell Reports (1990) 8: 643-646. Copyright Springer-Verlag 1990.
- Cho, Un-Haing, et al., Detection of Genetic Variation and Gene Introgression in Potato Dihaploids Using Randomly Amplified Polymorphic DNA (RAPD) Markers, J. Plant Biol. 39(3): 185-188 (1996). Copyright 1996 by Botanical Society of Korea, Seoul.
- Alan H. Christensen, et al., Sequence analysis and transcriptional regulation by heat shock of polyubiquitin transcripts from maize. accepted in revised form Feb. 7, 1989. Plant Molecular Biology 12: 619-632, 1989. Copyright 1989 Kluwer Academic Publishers. Printed in Belgium.
- Alan H. Christensen, et al., Maize polyubiquitin genes: structure, thermal perturbation of expression and transcript splicing, and promoter activity following transfer to protoplasts by electroporation. accepted in revised form Oct. 25, 1991. Plant Molecular Biology 18: 675-689, 1992. Copyright 1992 Kluwer Academic Publishers. Printed in Belgium.
- Paul Christou, et al., Stable transformation of soybean by electroporation and root formation from transformed callus. Communicated by Oliver E. Nelson, Jr., Feb. 25, 1987. Proc. Natl. Acad. Sci. USA, vol. 84, pp. 3962-3966, Jun. 1987. Applied Biology.
- Phan V. Chuong, et al., A simple culture method for Brassica hypototyl protoplasts. Revised version received Nov. 27, 1984—Communicated by F. Constabel. Plant Cell Reports (1985) 4:4-6.
- L. Comai, et al., Expression in plants of a mutant *aroA* gene from *Salmonella typhimurium* confers tolerance to glyphosate. Letters to Nature, Nature vol. 317 Oct. 24, 1985, pp. 741-744. Copyright 1985 Nature Publishing Group.
- Ben J. C. Cornelissen, et al., Update on Biotechnology, Strategies for Control of Fungal Diseases with Transgenic Plants, Plant Physiol. (1993) 101: 709-712.
- Gary Creissen, et al., Short Communication, Molecular characterization of glutathione reductase cDNAs from pea (*Pisum sativum* L.). The Plant Journal (1991) 2(1), 129-131.
- Swapna K. Datta, et al., Herbicide-resistant Indica rice plants from IIRI breeding line IR72 after PEG-mediated transformation of protoplasts. accepted in revised form Jun. 8, 1992. Plant Molecular Biology 20: 619-629, 1992. Copyright 1992 Kluwer Academic Publishers. Printed in Belgium.
- Marc De Block, et al., Expression of foreign genes in regenerated plants and in their progeny. The EMBO Journal vol. 3 No. 8 pp. 1681-1689, 1984. Copyright IRL Press Limited, Oxford, England.
- Willy De Greef, et al., Evaluation of Herbicide Resistance in Transgenic Crops Under Field Conditions, BIO/Technology vol. 7 Jan. 1989, pp. 61-64. Copyright 1989 Nature Publishing Group <http://www.nature.com/naturebiotechnology>.
- Alain Deshayes, et al., Liposome-mediated transformation of tobacco mesophyll protoplasts by an *Escherichia coli* plasmid, The EMBO Journal vol. 4 No. 11 pp. 2731-2737, 1985. Copyright IRL Press Limited, Oxford, England.

- Kathleen D'Halluin, et al., Transgenic Maize Plants by Tissue Electroporation, *The Plant Cell*, vol. 4, 1495-1505, Dec. 1992. Copyright 1992 American Society of Plant Physiologists.
- J. Draper, et al., Ti Plasmid Homologous Sequences Present in Tissues from *Agrobacterium* Plasmid-transformed *Petunia* Protoplasts, *Plant & Cell Physiol.* 23(3): 451-458 (1982).
- David A. Eichholtz, et al., Expression of Mouse Dihydrofolate Reductase Gene Confers Methotrexate Resistance in Transgenic *Petunia* Plants. Final Oct. 6, 1986. *Somatic Cell and Molecular Genetics*, vol. 13, No. 1, 1987, pp. 67-76. 0740-7750/8710100/006\$05.00/0 Copyright 1987 Plenum Publishing Corporation.
- Kathryn J. Elliott, et al., Isolation and characterization of fruit vacuolar invertase genes from two tomato species and temporal differences in mRNA levels during fruit ripening. accepted in revised form Oct. 20, 1992. *Plant Molecular Biology* 21: 515-524, 1993. Copyright 1993 Kluwer Academic Publishers. Printed in Belgium.
- Jean Finnegan, et al., Transgene Inactivation: Plants Fight Back!, Review, *Bio/Technology* vol. 12 Sep. 1994, pp. 883-888. Copyright 1994 Nature Publishing Group <http://www.nature.com/naturebiotechnology>.
- Dane K. Fisher, et al., Plant Gene Register, Starch Branching Enzyme II from Maize Endosperm, *Plant Physiol.* (1993) 102: 1045-1046.
- R. B. Flavell, Review, Inactivation of gene expression in plants as a consequence of specific sequence duplication, *Proc. Natl. Acad. Sci. USA*, vol. 91, pp. 3490-3496, Apr. 1994.
- Elizabeth B. P. Fontes, et al., Characterization of an Immunoglobulin Binding Protein Homolog in the Maize flourly-2 Endosperm Mutant, *The Plant Cell*. vol. 3, 483-496, May 1991. Copyright 1991 American Society of Plant Physiologists.
- Robert T. Fraley, et al., Expression of bacterial genes in plant cells, *Proc. Natl. Acad. Sci. USA*, vol. 80, pp. 4803-4807, Aug. 1983. *Genetics*.
- Christiane Gatz, et al., Regulation of a modified CaMV 35S promoter by the Tn10-encoded Tet repressor in transgenic tobacco, *Mol. Gen. Genet* (1991) 227:229-237. 002689259100168M.
- Frank T. Roder, et al., Efficiency of the tetracycline-dependent gene expression system: complete suppression and efficient induction of the rolB phenotype in transgenic plants. Accepted: Oct. 11, 1993. *Mol. Gen. Genet.* (1994) 243:32-38. Copyright Springer-Verlag 1994.
- William J. Gordon-Kamm, et al., Transformation of Maize Cells and Regeneration of Fertile Transgenic Plants. *The Plant Cell*, vol. 2, 603-618, Jul. 1990. Copyright 1990 American Society of Plant Physiologists.
- Stephen J. Gould, et al., A Conserved Tripeptide Sorts Proteins to Peroxisomes, Published May 1, 1989, *The Journal of Cell Biology*, vol. 108, May 1989 1657-1664. Copyright The Rockefeller University Press, 0021-9525/89/05/1657/8 \$2.00.
- Eike A. Griess, et al., Plant Gene Register, *Plant Physiol.* (1994) 104: 1467-1468.
- Margaret Y. Gruber, et al., Chapter 7, Vectors for Plant Transformation, *Methods in Plant Molecular Biology and Biotechnology*, Std. No. 9780849351648, pp. 89-119, Published 1993.
- Bernard R. Glick, et al., References, *Methods in plant molecular biology and biotechnology*, Std. No. 9780849351648, pp. 284-285, Published 1993.
- Felix D. Guerrero, et al., Promoter sequences from a maize pollen-specific gene direct tissue-specific transcription in tobacco, *Mol. Gen. Genet.* (1990) 24: 161-168. Copyright Springer-Verlag 1990.
- R. Hain, et al. Uptake, integration, expression and genetic transmission of a selectable chimaeric gene by plant protoplasts, *Mol. Gen. Genet.* (1985) 199: 161-168. Copyright Springer-Verlag 1985.
- Bruce D. Hammock, et al., Expression and effects of the juvenile hormone esterase in a baculovirus vector, *Letters to Nature*, *Nature*, vol. 344, pp. 485-461, Mar. 29, 1990.
- Jim Haseloff, et al., Article, Simple RNA enzymes with new and highly specific endoribonuclease activities, *Nature* vol. 334, pp. 585-591, Aug. 18, 1988. Copyright 1988 Nature Publishing Group.
- Jiro Hattori, et al., Original Paper, An acetohydroxy acid synthase mutant reveals a single site involved in multiple herbicide resistance, Accepted Aug. 18, 1994, *Mol. Gen. Genet* (1995) 246: 419-425. Copyright Springer-Verlag 1995.
- John D. Hayes, et al., Molecular cloning and heterologous expression of a cDNA encoding a mouse glutathione S-transferase Yc subunit possessing high catalytic activity for aflatoxin B1-8,9-epoxide, *Biochem. J.* (1992) 285, 173-180 (Printed in Great Britain).
- Maria B. Hayford, et al., Development of a Plant Transformation Selection System Based on Expression of Genes Encoding Gentamicin Acetyltransferases, Received for publication Nov. 6, 1987 and in revised form Jan. 8, 1988. *Plant Physiol.* 86, 1216-1222. 0032-0889/88/86/126/07/\$01.00/0.
- Gayle Heney, et al., The Purification of Avidin and Its Derivatives on 2-Iminobiotin-6aminohexyl-Sepharose 4B, *Analytical Biochemistry* 114, 92-96 (1981). 0003-2697/81/090092-05\$02.00/0. Copyright 1981 by Academic Press, Inc.
- Christiane Gatz, et al., Regulation of a modified CaMV 35S promoter by the TN10-encoded Tet repressor in transgenic tobacco, *Mol. Gen. Genet* (1991) 227: 229-237. 00268925100168M.
- Jacques Hille, et al., Bleomycin resistance: a new dominant selectable marker for plant cell transformation, *Plant Molecular Biology* 7: 171-176 (1986). Copyright Martinus Nijhoff Publishers, Dordrecht—Printed in Netherlands.
- U.S. Patent and Trademark Office, Non-Final Office Action, mailed Jul. 1, 2011, U.S. Appl. No. 12/486,161, filed Jun. 17, 2009, Confirmation No. 4613, Zenon Lisieczko.
- R. B. Horsch, et al., A Simple and General Method for Transferring Genes into Plants, Mar. 8, 1985, pp. 1229-1231.
- Huib J.M. Linthorst, et al., Tobacco proteinase inhibitor I genes are locally, but not systemically induced by stress, accepted in revised form Dec. 20, 1992. *Plant Molecular Biology* 21: 985-992, 1993. Copyright 1993 Kluwer Academic Publishers. Printed in Belgium.
- Jesse M. Jaynes, et al., Expression of a Cecropin B lytic peptide analog in transgenic tobacco confers enhanced resistance to bacterial wilt caused by *Pseudomonas solanacearum*, revision received Nov. 4, 1992; accepted Dec. 7, 1992). 0168-9452/93\$06.00 Copyright 1993 Elsevier Scientific Publishers Ireland Ltd. Printed and Published in Ireland.
- Richard A. Jefferson, *Experimental Protocols, Assaying Chimeric Genes in Plants: The GUS Gene Fusion System*, *Plant Molecular Biology Reporter*, vol. 5, No. 1, 1987, pp. 387-405.
- Jonathan D.G. Jones, et al., A dominant nuclear streptomycin resistance marker for plant cell transformation, *Mol. Gen. Genet.* (1987) 210: 86-91. Copyright Springer-Verlag 1987.
- David A. Jones, et al., Reports, Isolation of the Tomato Cf-9 Gene for Resistance to *Cladosporium fulvum* by Transposon Tagging, *Science*, vol. 266, pp. 789-793, Nov. 4, 1994.
- Richard Jorgensen, Undercurrents, Altered gene expression in plants due to trans interactions between homologous genes, TIBTECH—Dec. 1990 (vol. 8) pp. 340-344. Copyright 1990, Elsevier Science Publishers Ltd. (UK) 0167-940/90/\$2.00.
- Clarence I. Kado, Ph.D., Molecular Mechanisms of Crown Gall Tumorigenesis, *Critical Reviews in Plant Sciences*, 10(1):1-32 (1991). 0735-2689/91/\$.50 Copyright 1991 by CRC Press, Inc.
- Daniel Calderon, et al., A Short Amino Acid Sequence Able to Specify Nuclear Location, *Cell*, vol. 39, 499-509, Dec. 1984 (Part 2), Copyright 1984 by MT. 0092-8674/84/130499-11 \$02.00/0.
- K. K. Kartha, et al., In vitro Plant Formation from Stem Explants of Rape (*Brassica napus* cv. Zephyr), *Physiol. Plant.* 31: 217-220, 1974.
- Petra Kawalleck, et al., Polyubiquitin gene expression and structural properties of the ubi4-2 gene in *Petroselinum crispum*, accepted in revised form Nov. 22, 1992, *Plant Molecular Biology* 21: 673-684, 1993. Copyright 1993 Kluwer Academic Publishers. Printed in Belgium.
- T. M. Klein, Factors Influencing Gene Delivery Into *Zea mays* Cells by High-Velocity Microprojectiles, *Bio/Technology* vol. 6, May 1988, pp. 559-563. Copyright 1988 Nature Publishing Group <http://www.nature.com/naturebiotechnology>.
- Cheryl A. P. Knox, et al., Structure and organization of two divergent α -amylase genes from barley, in revised form Mar. 2, 1987; accepted Mar. 3, 1987, *Plant Molecular Biology* 9: 3-17 (1987). Copyright Martinus Nijhoff Publishers, Dordrecht—Printed in the Netherlands.

- Deborah S. Knutzon, et al., Modification of *Brassica* seed oil by antisense expression of a stearyl-acyl carrier protein desaturase gene, Communicated by Donald R. Helinski, Dec. 23, 1991, Proc. Natl. Acad. Sci. USA, vol. 89, pp. 2624-2628, Apr. 1992. Plant Biology.
- Mitsue Kobayashi, et al. Haploid induction and its genetic mechanism in alloplasmic common wheat, The Journal of Heredity 71: 9-14, 1980.
- Csaba Koncz, et al., Expression and assembly of functional bacterial luciferase in plants, Contributed by Jeff Schell, Sep. 10, 1986, Proc. Natl. Acad. Sci. USA, vol. 84, pp. 131-135, Jan. 1987, Cell Biology.
- Karl J. Kramer, et al., Sequence of a cDNA and Expression of the Gene Encoding Epidermal and Gut Chitinases of *Manduca sexta*, revised and accepted Jan. 13, 1993. Insect Biochem. Molec. Biol. vol. 23, No. 6, pp. 691-701, 1993. Printed in Great Britain. 0965-1748/93 \$6.00+0.00, Pergamon Press Ltd.
- Christopher J. Lamb, et al., Review/Emerging Strategies for Enhancing Crop Resistance to Microbial Pathogens, Biotechnology vol. 10, Nov. 1992, pp. 1436-1445. Copyright 1992 Nature Publishing Group <http://www.nature.com/naturebiotechnology>.
- Corne MJ Pieterse, et al., NPR1: the spider in the web of induced resistance signaling pathways, Current Opinion in Plant Biology 2004, 7:456-464. Elsevier. Full text provided by www.sciencedirect.com.
- Joan T. Odell, et al., Letters to Nature, Identification of DNA sequences required for activity of the cauliflower mosaic virus 35S promoter, Nature vol. 313, Feb. 28, 1985. Copyright 1985 Nature Publishing Group.
- S. B. Narasimhulu, et al., Species specific shoot regeneration response of cotyledonary explants of *Brassicaceae*,—Communicated by E. Earle, Plant Cell Reports (1988) 7: 104-106. Plant Cell Reports Copyright Springer-Verlag 1988.
- Wyatt Paul, et al., The isolation and characterization of the tapetum-specific *Arabidopsis thaliana* A9 gene, accepted in revised form Mar. 19, 1992, Plant Molecular Biology 19: 611-622, 1992. Copyright 1992 Kluwer Academic Publishers. Printed in Belgium.
- Jan Van Parijs, et al., Hevein: an antifungal protein from rubber-tree (*Hevea brasiliensis*) latex, accepted Jul. 25, 1990, Planta (1991) 183: 258-264. Planta Copyright Springer-Verlag 1991.
- F. Neuberger, et al., Susceptibility of transgene loci to homology-dependent gene silencing, Accepted: Feb. 7, 1994, Mol. Gen. Genet (1994) 244: 230-241. Copyright Springer-Verlag 1994.
- D. I. Last, et al., pEmu: an improved promoter for gene expression in cereal cells, Accepted Oct. 16, 1990, Theor. Appl. Genet (1991) 81: 581-588.
- Kathleen Y. Lee, et al., The molecular basis of sulfonylurea herbicide resistance in tobacco, The EMBO Journal vol. 7, No. 5, pp. 1241-1248, 1988.
- Marc Lepetit, et al., A plant histone gene promoter can direct both replication-dependent and -independent gene expression in transgenic plants, Mol. Gen. Genet (1992) 231: 276-285, MGG Copyright Springer-Verlag 1992.
- David R. Lerner, et al., Cloning and Characterization of Root-Specific Barley Lectin, Received for publication Mar. 13, 1989 and in revised form May 3, 1989, Plant Physiol. (1989) 91, 124-129, 0032-0889/89/91/0124/06/\$01.00/0.
- J. Logemann, et al., Expression of a Barley Ribosome-Inactivating Protein Leads to Increased Fungal Protection in Transgenic Tobacco Plants, Bio/Technology, vol. 10, Mar. 1992, pp. 305-308. Copyright 1992 Nature Publishing Group <http://www.nature.com/naturebiotechnology>.
- L.A. Lyznik, et al., Site-specific recombination for genetic engineering in plants, Received: Dec. 20, 2002 / Accepted: Feb. 24, 2003 / Published online: Apr. 26, 2003. Plant Cell Rep (2003) 21: 925-932. DOI 10.1007/s00299-003-0616-7. Copyright Springer-Verlag 2003.
- L. C. Marshall, et al., Allelic mutations in acetyl-coenzyme A carboxylase confer herbicide tolerance in maize, Accepted Jul. 9, 1991, Theor. Appl. Genet (1992) 83: 435-442. Copyright Springer-Verlag 1992.
- Gregory B. Martin, et al., Reports, Map-Based Cloning of a Protein Kinase Gene Conferring Disease Resistance in Tomato, Science, vol. 262, pp. 1432-1436, Nov. 26, 1993.
- J. Velten, et al., Isolation of a dual plant promoter fragment from the Ti plasmid of *Agrobacterium tumefaciens*, The EMBO Journal, vol. 3, No. 12, pp. 2723-2730, 1984. Copyright IRL Press Limited, Oxford, England.
- Ken Matsuoka, et al., Propeptide of a precursor to a plant vacuolar protein required for vacuolar targeting, Communicated by Shang Fa Yang, Oct. 15, 1990 (received for review Sep. 4, 1990), Proc. Natl. Acad. Sci. USA, vol. 88, pp. 834-838, Feb. 1991, Cell Biology.
- John M. McDowell, et al., Review, Plant disease resistance genes: recent insights and potential applications, Trends in Biotechnology, vol. 21, No. 4, pp. 178-183, Apr. 2003. <http://tibtec.trends.com/0167-7799/0111>—see front matter. Copyright 2003 Elsevier Science Ltd. All rights reserved. doi:10.1016/S0167-7799(03)00053-2.
- David McElroy, et al., Isolation of an Efficient Actin Promoter for Use in Rice Transformation, The Plant Cell, vol. 2, 163-171, Feb. 1990. Copyright 1990 American Society of Plant Physiologists.
- Vadim L. Mett, et al., Copper-controllable gene expression system for whole plants, Communicated by Eric E. Conn, Jan. 26, 1993, Proc. Natl. Acad. Sci. USA, vol. 90, pp. 4567-4571, May 1993, Plant Biology.
- B. L. Miki, et al., Methods in Plant Molecular Biology and Biotechnology, Chapter 6, Procedures for Introducing Foreign DNA into Plants, pp. 67-88. 0-8493-5164-2/1993/\$0.00+\$0.50. Copyright 1993 by CRC Press, Inc.
- B. L. Miki, et al., Transformation of *Brassica napus* canola cultivars with *Arabidopsis thaliana* acetohydroxyacid synthase genes and analysis of herbicide resistance, Accepted Apr. 11, 1990, Theor. Appl. Genet (1990) 80: 449-458. Copyright Springer-Verlag 1990.
- Michael Mindrinos, et al., The *A. thaliana* Disease Resistance Gene RPS2 Encodes a Protein Containing a Nucleotide-Binding Site and Leucine-Rich Repeats, *Arabidopsis* Disease Resistance Gene, Cell, vol. 78, 1089-1099, Sep. 23, 1994, Copyright 1994 by Cell Press.
- Maurice M. Moloney, et al., High efficiency transformation of *Brassica napus* using *Agrobacterium* vectors,—Communicated by F. Constable, Plant Cell Reports (1989) 8:238-242. Copyright Springer-Verlag 1989.
- Mary K. Montgomery, et al., RNA as a target of double-stranded RNA-mediated genetic interference in *Caenorhabditis elegans*, Communicated by Joseph G. Gall, . . . , Nov. 2, 1998 (received for review Oct. 6, 1998), Proc. Natl. Acad. Sci. USA, vol. 95, pp. 15502-15507, Dec. 1998. Genetics. Copyright 1998 by the National Academy of Sciences 0027-8424/98/9515502-6\$2.00/0 PNAS is available online at www.pnas.org.
- Norimoto Murai, et al., Research Article, Phaseolin Gene from Bean Is Expressed After Transfer to Sunflower Via Tumor-Inducing Plasmid Vectors, Science, vol. 222, pp. 476-482. accepted Sep. 14, 1983.
- Carolyn Napoli, et al., Introduction of a Chimeric Chalcone Synthase Gene into Petunia Results in Reversible Co-Suppression of Homologous Genes in trans, The Plant Cell, vol. 2, 279-289, Apr. 1990, Copyright 1990 American Society of Plant Physiologists.
- S.J. Openshaw, et al., Marker-assisted Selection in Backcross Breeding, Analysis of Molecular Marker Data, Joint Plant Breeding Symposia Series, Aug. 5-6, 1994, Corvallis, Oregon, American Society for Horticultural Science Crop Science Society of America, pp. 41-43. Supplied by the British Library—"The world's knowledge".
- Sheng-Zhi Pang, et al., Expression of a gene encoding a scorpion insectotoxin peptide in yeast, bacteria and plants, Gene. 116 (1992) 165-172, Copyright 1992 Elsevier Science Publishers B.V. All rights reserved 0378-1199/92/ \$05.00.
- Jan Pen, et al., Research, Production of Active *Bacillus licheniformis* Alpha-Amylase in Tobacco and Its Application in Starch Liquefaction, Bio/Technology, vol. 10, pp. 292-296, Mar. 1992. Copyright 1992 Nature Publishing Group <http://www.nature.com/naturebiotechnology>.
- Rhonda Perriman, et al., A Ribozyme That Enhances Gene Suppression in Tobacco Protoplasts, accepted for publication Jun. 2, 1993, Antisense Research and Development, 3:253-263 (1993) Mary Ann Liebert, Inc., Publishers.

J. Plieske, et al., Microsatellite markers for genome analysis in *Brassica*. I. development in *Brassica napus* and abundance in *Brassicaceae* species, *Theor Appl. Genet* (2001) 102: 689-694. Copyright Springer-Verlag 2001.

M. Pollacsek, Plant improvement, Management of the *ig* gene for haploid induction in maize, accepted Jan. 16, 1992), *Agronomie* (1991) 12, 247-251. Copyright Elsevier/INRA.

United States Patent and Trademark Office, Non-Final Office Action, mailed Dec. 29, 2011, U.S. Appl. No. 12/713,411, filed Feb. 26, 2010, Jun Liu.

United States Patent and Trademark Office, Non-Final Office Action, mailed Dec. 29, 2011, U.S. Appl. No. 12/713,479, filed Feb. 26, 2010, Jun Liu.

United States Patent and Trademark Office, Non-Final Office Action, mailed Dec. 29, 2011, U.S. Appl. No. 12/713,637, filed Mar. 11, 2010, Chunren Wu.

* cited by examiner

**PLANTS AND SEEDS OF SPRING CANOLA
VARIETY SCV470336**

BACKGROUND OF THE INVENTION

Field of the Invention

The present invention relates to a new and distinctive canola line, designated SCV470336.

SUMMARY OF THE INVENTION

The present invention is directed, in an embodiment, to a plant, plant part, or seed of canola line SCV470336, representative sample of seed of which was deposited under ATCC Accession No. PTA-10628. The invention is also directed, in an embodiment, to a method for producing a canola seed comprising crossing two canola plants and harvesting the resultant canola seed, wherein at least one of the two canola plants is of canola line SCV470336. In another embodiment, the invention is also directed to a method for producing a canola plant with a particular trait, such as male sterility, herbicide or insect resistance or tolerance, or modified fatty acid metabolism or modified carbohydrate metabolism, wherein the method comprises transforming the canola plant of canola line SCV470336 with a nucleic acid molecule that confers that trait.

DEFINITIONS

Allele. Allele is any of one or more alternative forms of a gene which relate to one trait or characteristic. In a diploid cell or organism, the two alleles of a given gene occupy corresponding loci on a pair of homologous chromosomes.

Alter. The utilization of up-regulation, down-regulation, or gene silencing.

Anther arrangement. The orientation of the anthers in fully opened flowers; anther arrangement can be useful as an identifying trait. Anther arrangements can range from introse (facing inward toward pistil), erect (neither inward not outward), or extrose (facing outward away from pistil).

Anther dotting. The presence/absence of anther dotting (colored spots on the tips of anthers); if present, the percentage of anther dotting on the tips of anthers in newly opened flowers is a distinguishing trait for varieties.

Anther fertility. A measure of the amount of pollen produced on the anthers of a flower. Anther fertility can range from sterile (such as in female parents used for hybrid seed production) to fertile (all anthers shedding).

Backcrossing. A process in which a breeder repeatedly crosses hybrid progeny back to one of the parents; for example, a first generation hybrid F_1 may be crossed with one of the parental genotypes of the F_1 hybrid.

Blackleg (*Leptosphaeria maculans*). A fungal canker or dry rot disease of the actively growing crop that causes stem girdling and lodging. In heavily infested crops, up to 100% of the stems may be infected, resulting in major yield loss. For purposes of this application, resistance to blackleg is measured using ratings of "R" (resistant), "MR" (moderately resistant), "MS" (moderately susceptible) or "S" (susceptible).

Cell. As used herein, the term cell includes a plant cell, whether isolated, in tissue culture, or incorporated in a plant or plant part.

Cotyledon width. The cotyledons are leaf structures that form in the developing seeds of canola that make up the majority of the mature seed of these species. When the seed germi-

nates, the cotyledons are pushed out of the soil by the growing hypocotyls (segment of the seedling stem below the cotyledons and above the root) and they unfold as the first photosynthetic leaves of the plant. The width of the cotyledons varies by variety and can be classified as narrow, medium, or wide.

Elite canola line. A canola line which has been sold commercially.

Elite canola parent line. A canola line which is the parent line of a canola line which has been commercially sold.

Embryo. The embryo is the small plant contained within a mature seed. Essentially all of the physiological and morphological characteristics. This phrase refers to a plant having essentially all of the physiological and morphological characteristics of the recurrent parent, except for the characteristics derived from any converted trait.

Fatty Acid Methyl Ester ("FAME") analysis. A method that allows for accurate quantification of the fatty acids which make up complex lipid classes.

Flower bud location. The location of the unopened flower buds relative to the adjacent opened flowers is useful in distinguishing between the canola species. The unopened buds are held above the most recently opened flowers in *B. napus* and they are positioned below the most recently opened flower buds in *B. rapa*.

Flowering date. This is measured by the number of days from planting to the stage when 50% of the plants in a population have one or more open flowers. This varies from variety to variety.

***Fusarium* Wilt.** *Fusarium* wilt, largely caused by *Fusarium oxysporum*, is a disease of canola that causes part or all of a plant to wilt, reducing yield by 30% or more on badly affected fields. For purposes of this application, resistance to *Fusarium* wilt is measured using ratings of "R" (resistant), "MR" (moderately resistant), "MS" (moderately susceptible) or "S" (susceptible).

Gene silencing. Gene silencing means the interruption or suppression of the expression of a gene at the level of transcription or translation.

Genotype. This term refers to the genetic constitution of a cell or organism.

Glucosinolates. A secondary metabolite of Brassicales that is organic and contains sulfur and nitrogen; derived from glucose and an amino acid. Glucosinolates are measured in micromoles (μm) of total aliphatic glucosinolates per gram of air-dried oil-free meal. The level of glucosinolates is somewhat influenced by the sulfur fertility of the soil, but is also controlled by the genetic makeup of each variety and thus can be useful in characterizing varieties.

Growth habit. At the end of flowering, the angle relative to the ground surface of the outermost fully expanded leaf petioles; a variety-specific trait. This trait can range from erect (very upright along the stem) to prostrate (almost horizontal and parallel with the ground surface).

Leaf attachment to the stem. The manner in which the base of the leaf blade clasps the stem. This trait is especially useful for distinguishing between the two canola species. The base of the leaf blade of the upper stem leaves of *B. rapa* completely clasp the stem whereas those of the *B. napus* only partially clasp the stem. Those of the mustard species do not clasp the stem at all.

Leaf blade color. The color of the leaf blades is variety-specific and can range from light to medium dark green to blue-green.

Leaf development of lobes. The varying degrees of development of lobes on leaves on the upper portion of the stem which are disconnected from one another along the petiole

3

of the leaf. The degree of lobing is variety-specific and can range from absent (no lobes)/weak to very strong (abundant lobes).

Leaf glaucosity. This refers to the waxiness of the leaves and is characteristic of specific varieties although environment can have some effect on the degree of waxiness. This trait can range from absent (no waxiness)/weak to very strong. The degree of waxiness can be best determined by rubbing the leaf surface and noting the degree of wax present.

Leaf indentation of margin. The varying degrees of serration along the leaf margins on leaves on the upper portion of the stem. The degree of serration or indentation of the leaf margins can vary from absent (smooth margin)/weak to strong (heavy saw-tooth like margin).

Leaf pubescence. The leaf pubescence is the degree of hairiness of the leaf surface and is especially useful for distinguishing between the canola species. There are two main classes of pubescence, glabrous (smooth/not hairy) and pubescent (hairy), which mainly differentiate between the *B. napus* and *B. rapa* species, respectively.

Leaf surface. A measure of the surface texture of a leaf; the leaf surface can be used to distinguish between varieties. The surface can be smooth or rugose (lumpy) with varying degrees between the two extremes.

Linkage. This term refers to a phenomenon wherein alleles on the same chromosome tend to segregate together more often than expected if their transmission was independent.

Linkage disequilibrium. A phenomenon wherein alleles tend to remain together in linkage groups when segregating from parents to offspring, with a greater frequency than expected from their individual frequencies.

Locus. A position on a genomic sequence that is usually found by a point of reference, for example, the position of a DNA sequence that is a gene, or part of a gene or intergenic region. A locus confers one or more traits such as, for example, male sterility, herbicide tolerance, insect resistance, disease resistance, modified fatty acid metabolism, modified phytic acid metabolism, modified carbohydrate metabolism and modified protein metabolism. The trait may be, for example, conferred by a naturally occurring gene introduced into the genome of the variety by backcrossing, a natural or induced mutation, or a transgene introduced through genetic transformation techniques. A locus may comprise one or more alleles integrated at a single chromosomal location.

Lodging resistance. This term refers to a variety's ability to remain erect. Lodging is rated on a scale of 1 to 5, wherein a score of 1 indicates erect plants and a score of 5 indicates plants that are lying on the ground.

Maturity. The maturity of a variety is measured as the number of days between planting and physiological maturity. This trait is useful in distinguishing varieties relative to one another and when used in this context it is referred to as "Relative Maturity".

Oil content. Oil content is measured as a percent of the whole dried seed and is variety-specific. It can be determined using various analytical techniques such as nuclear magnetic resonance (NMR) spectroscopy, near-infrared (NIR) spectroscopy, and Soxhlet extraction.

Percent linolenic acid. This refers to the percent of total oil in the seed that is linolenic acid.

Percent oleic acid (OLE). OLE refers to the percent of total oil in the seed that is oleic acid.

Percentage of total fatty acids. This may be determined by extracting a sample of oil from seed, producing the methyl esters of fatty acids present in that oil sample and analyzing the proportions of the various fatty acids in the sample

4

using gas chromatography. The fatty acid composition can also be a distinguishing characteristic of a variety.

Petal color. The petal color on the first day a flower opens can be a distinguishing characteristic for a variety. It can be white, varying shades of yellow, or orange.

Plant. As used herein, the term "plant" includes plant cells, plant protoplasts, plant cell tissue cultures from which canola plants can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants, such as embryos, pollen, ovules, flowers, pods, leaves, roots, root tips, anthers, cotyledons, hypocotyls, meristematic cells, stems, pistils, petiole, and an immature or mature whole plant, including a plant from which seed or grain or anthers have been removed. Seed or embryo that will produce the plant is also considered to be the plant.

Plant height. This is the height of the plant at the end of flowering if the floral branches are extended upright (i.e., not lodged). This varies from variety to variety and although it can be influenced by environment, relative comparisons between varieties grown side by side are useful for variety identification.

Plant parts. As used herein, the term "plant parts" (or "canola plant, or a part thereof" or similar phrasings) includes protoplasts, leaves, stems, roots, root tips, anthers, pistils, seed, grain, embryo, pollen, ovules, cotyledon, hypocotyl, pod, flower, shoot, tissue, petiole, cells, meristematic cells and the like.

Protein content. Protein content is measured as a percent of whole dried seed and may vary from variety to variety. This can be determined using various analytical techniques such as NIR and Kjeldahl.

Quantitative trait loci (QTL). This term refers to genetic loci that control to some degree numerically-representable traits that are usually continuously distributed.

Regeneration. Regeneration refers to the development of a plant from tissue culture.

Resistance to lodging. This term refers to the ability of a variety to stand up in the field under high yield conditions and severe environmental factors. A variety can have very good, good (remains upright), fair, or poor (falls over) resistance to lodging. The degree of resistance to lodging is not expressed under all conditions but is most meaningful when there is some degree of lodging in a field trial.

Seed coat color. The color of the seed coat can be variety-specific and can range from black to brown to yellow. Color can also be mixed for some varieties.

Seed coat mucilage. This refers to a gel-like substance on the seed coat that draws water to itself. It can be useful for differentiating between the two species of canola with *B. rapa* varieties having mucilage present in their seed coats whereas *B. napus* varieties do not have this present. It is detected by imbibing seeds with water and monitoring the mucilage that is exuded by the seed.

Seedling growth habit. This term refers to the rosette, which consists of the first 2-8 true leaves; a variety can be characterized as having a strong rosette (closely packed leaves) or a weak rosette (loosely arranged leaves).

Silique (pod) habit. This term refers to the orientation of the pods along the racemes (flowering stems) and is variety-specific. This trait can range from erect (pods angled close to racemes) to horizontal (pods perpendicular to racemes) to arching (pods show distinct arching habit).

Silique (pod) length of beak. The beak is the segment at the end of the pod which does not contain seed (it is a remnant of the stigma and style for the flower). The length of the beak can be variety-specific and can range from short to medium to long.

Silique (pod) length of pedicel. The pedicel is the stem that attaches the pod to the raceme or flowering shoot. The length of the pedicel can be variety-specific and can vary from short to medium to long.

Silique (pod) length. This is the length of the fully developed pods and can range from short to medium to long. In characterizing a variety, it is best used by making comparisons relative to reference varieties.

Silique (pod) type. The type of pod; this is typically a bilateral single pod for both species of canola and is, therefore, not very useful for variety identification within these species.

Silique (pod) width. This is the width of a fully developed pod and can range from narrow to medium to wide. In characterizing a variety, it is best used by making comparisons relative to reference varieties.

Single gene converted (conversion). This term refers to plants that are developed using a plant breeding technique known as backcrossing, or via genetic engineering, wherein essentially all of the desired morphological and physiological characteristics of a variety are recovered in addition to the single gene transferred into the variety via the backcrossing technique or via genetic engineering.

Stem intensity of anthocyanin coloration. The varying degrees of purple coloration on stems and other organs of canola plants, which is due to the presence of anthocyanin (purple) pigments. The degree of coloration is somewhat subject to growing conditions, but varieties typically show varying degrees of coloration ranging from: absent (no purple) to very weak to very strong (deep purple coloration).

Total saturated (TOTSAT). Measured as a percent of the total oil of the seed, this refers to the amount of saturated fats in the oil including C12:0, C14:0, C16:0, C18:0, C20:0, C22:0 and C24:0.

DETAILED DESCRIPTION OF THE INVENTION

Reference now will be made in detail to the embodiments of the invention, one or more examples of which are set forth below. Each example is provided by way of explanation of the invention, not a limitation of the invention. In fact, it will be apparent to those skilled in the art that various modifications and variations can be made in the present invention without departing from the scope or spirit of the invention. For instance, features illustrated or described as part of one embodiment, can be used on another embodiment to yield a still further embodiment.

Thus, it is intended that the present invention covers such modifications and variations as come within the scope of the appended claims and their equivalents. Other objects, features and aspects of the present invention are disclosed in or are obvious from the following detailed description. It is to be understood by one of ordinary skill in the art that the present discussion is a description of exemplary embodiments only, and is not intended as limiting the broader aspects of the present invention.

According to the invention, there is provided a new canola line designated SCV470336. This invention relates to the seeds, plants, and plant parts of canola SCV470336 and to methods for producing a canola plant produced by crossing the canola SCV470336 with itself or another canola genotype, and to the creation of variants by mutagenesis or transformation of canola SCV470336.

Thus, any such methods using the canola line SCV470336 are part of this invention, including but not limited to selfing, backcrossing, hybrid production, and crosses to populations. All plants produced using canola line SCV470336 as a parent

are within the scope of this invention. In an embodiment, the canola line could be used in crosses with other different canola plants to produce first generation (F₁) canola hybrid seeds and plants with superior characteristics.

In another embodiment, the present invention provides for single or multiple gene-converted plants of SCV470336. The transferred gene(s) may be a dominant or recessive allele. The transferred gene(s) may confer such traits as herbicide tolerance, insect resistance, resistance for bacterial, fungal, or viral disease, male fertility, male sterility, enhanced nutritional quality, modified fatty acid metabolism, modified carbohydrate metabolism, modified seed yield, modified oil percent, modified protein percent, modified lodging resistance, modified glucosinolate content, modified chlorophyll content and industrial usage. The gene may be a naturally occurring canola gene or a transgene introduced through genetic engineering techniques.

In another embodiment, the present invention provides regenerable cells for use in tissue culture of canola plant SCV470336. The tissue culture may be capable of regenerating plants having essentially all of the physiological and morphological characteristics of the foregoing canola plant, and capable of regenerating plants having substantially the same genotype as the foregoing canola plant. The regenerable cells in such tissue cultures may be embryos, protoplasts, meristematic cells, callus, pollen, leaves, anthers, pistils, cotyledons, roots, root tips, flowers, seeds, pods or stems. Still further, the present invention, in an embodiment, provides canola plants regenerated from the tissue cultures of the invention.

In a particular embodiment, the invention comprises a plant cell or a descendant of a plant cell of a *Brassica* plant designated variety SCV470336. In some embodiments, the descendant has the same desirable traits as the plant designated variety SCV470336.

In another aspect, the present invention provides a method of introducing a desired trait into canola line SCV470336, wherein the method comprises crossing a SCV470336 plant with a plant of another canola genotype that comprises a desired trait to produce F₁ progeny plants, wherein the desired trait is selected from the group consisting of male sterility, herbicide tolerance, insect resistance, modified fatty acid metabolism, modified carbohydrate metabolism, modified seed yield, modified oil percent, modified protein percent, modified lodging resistance, and resistance to bacterial disease, fungal disease, or viral disease; selecting one or more progeny plants that have the desired trait to produce selected progeny plants; crossing the selected progeny plants with the SCV470336 plants to produce backcross progeny plants; selecting for backcross progeny plants that have the desired trait and essentially all of the physiological and morphological characteristics of canola line SCV470336 to produce selected backcross progeny plants; and repeating these steps three or more times to produce selected fourth or higher backcross progeny plants that comprise the desired trait and essentially all of the physiological and morphological characteristics of canola line SCV470336 as listed in Table 1. Included in this aspect of the invention is the plant produced by the method wherein the plant has the desired trait and essentially all of the physiological and morphological characteristics of canola line SCV470336 as listed in Table 1.

A. Origin and Breeding History

Spring canola variety SCV470336 is a spring canola inbred line developed from the cross of SCV378221 and SCV475099 (proprietary spring canola inbred lines of Monsanto Technology LLC). Progeny from this cross were self-pollinated and the pedigree system of plant breeding was then

used to develop SCV470336. Some of the criteria used for selection in various generations include: fertility, standability, disease tolerance, combining ability, oil content, maturity and total saturated fats.

SCV470336 is a conventional (non transgenic), pollen-fertile canola restorer inbred line (R-line) used for producing hybrids with resistance to blackleg and *Fusarium* wilt. Canola line SCV470336 is stable, uniform and no off-type plants have been exhibited in evaluation. The line has shown uniformity and stability, as described in the following variety description information. It has been self-pollinated a sufficient number of generations with careful attention to uniformity of plant type. The line has been increased with continued observation for uniformity.

B. Phenotypic Description

In accordance with another aspect of the present invention, there is provided a canola plant having the physiological and morphological characteristics of canola variety SCV470336. A description of various physiological and morphological characteristics of canola variety SCV470336 is presented in Table 1.

TABLE 1

Physiological and Morphological Characteristics of SCV470336			
CHARACTERISTIC	VALUE		
	SCV470336	SCV378221*	SCV204738*
1. PLANT			
Plant Height	132 cm	125 cm	130 cm
Days to 50%	50 days	49 days	50 days
Flowering			
Maturity	Medium	Early	Early
Resistance To	Very Good	Good	Very Good
Lodging			
Blackleg Resistance	R	MR	R
<i>Fusarium</i> Wilt Resistance	R	S	R
Rf Restorer Gene Source	From "old" source (INRA)	From "old" source (INRA)	From "old" source (INRA)
2. SEED			
Coat Color	Black	Black	Black
Oil Content (% of whole seed @ 8.5% moisture)**	45.39%	45.82%	45.68%
Protein Content (% defatted, dry meal)**	44.16%	46.87%	45.86%
Erucic Acid Content	Trace	Trace	Trace
Glucosinolate Content**	16.94%	16.23%	16.38%

*SCV378221 and SCV204738 are proprietary canola inbred lines of Monsanto Technology LLC.

**These are average values. Values may vary due to environment.

In an embodiment, the invention is directed to methods for producing a canola plant by crossing a first parent canola plant with a second parent canola plant, wherein the first or second canola plant is the canola plant from the line SCV470336. Further, both first and second parent canola plants may be from the line SCV470336. Any methods using the line SCV470336 are part of this invention: selfing, backcrosses, hybrid breeding, and crosses to populations. Any plants produced using line SCV470336 as a parent are within the scope of the invention.

Additional methods of the present invention include, but are not limited to, expression vectors introduced into plant tissues using a direct gene transfer method such as microprojectile-mediated delivery, DNA injection, and electroporation. Expression vectors may be introduced into plant tissues by using either microprojectile-mediated delivery with a biolistic device or by using *Agrobacterium*-mediated trans-

formation. Transformant plants obtained with the protoplasm of the invention are intended to be within the scope of this invention.

The advent of new molecular biological techniques has allowed the isolation and characterization of genetic elements with specific functions, such as encoding specific protein products. Scientists in the field of plant biology developed a strong interest in engineering the genome of plants to contain and express foreign genetic elements, or additional, or modified versions of native or endogenous genetic elements in order to alter the traits of a plant in a specific manner. Any DNA sequences, whether from a different species or from the same species which are inserted into the genome using transformation, are referred to herein collectively as "transgenes". In some embodiments of the invention, a transgenic variant of SCV470336 may contain at least one transgene but could contain at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 transgenes and/or no more than 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, 5, 4, 3, or 2 transgenes. Over the last fifteen to twenty years several methods for producing transgenic plants have been developed, and the present invention also relates to transgenic variants of the claimed canola line SCV470336.

One embodiment of the invention is a process for producing canola line SCV470336 further comprising a desired trait, said process comprising transforming a canola plant of line SCV470336 with a transgene that confers a desired trait. Another embodiment of the invention comprises a canola plant produced by this process. In one embodiment, the desired trait may be one or more of herbicide tolerance, insect resistance, disease resistance, modified seed yield, modified oil percent, modified protein percent, modified lodging resistance, or modified fatty acid or carbohydrate metabolism. The specific gene may be any gene known in the art or listed herein, including but not limited to a polynucleotide conferring resistance or tolerance to imidazolinone, sulfonylurea, glyphosate, glufosinate, 2,4-D, Dicamba, L-phosphinothricin, triazine, hydroxyphenylpyruvate dioxygenase inhibitor, protoporphyrinogen oxidase inhibitor, phenoxy propionic acid, cyclohexone, and benzonitrile; a polynucleotide encoding a *Bacillus thuringiensis* polypeptide, a polynucleotide encoding phytase, fatty acid desaturase (FAD)-2, FAD-3, galactinol synthase or a raffinose synthetic enzyme; or a polynucleotide conferring resistance to blackleg, white rust or other common canola diseases.

Numerous methods for plant transformation have been developed and are included as part of the invention, including biological and physical plant transformation protocols. See, for example, Miki, et al., "Procedures for Introducing Foreign DNA into Plants" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B. R. and Thompson, J. E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 67-88 and "Genetic Transformation for the Improvement of Canola", in *World Conference on Biotechnology for the Fats and Oils Industry*, 1988. In addition, expression vectors and in vitro culture methods for plant cell or tissue transformation and regeneration of plants are available and are included within the invention. See e.g. Gruber, et al., "Vectors for Plant Transformation" in *Methods in Plant Molecular Biology and Biotechnology*, Glick, B. R. and Thompson, J. E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 89-119.

As an embodiment of the invention, a genetic trait which has been engineered into the genome of a particular canola plant may then be moved into the genome of another variety using traditional breeding techniques that are well-known in the plant breeding arts. For example, a backcrossing approach may be used to move a transgene from a transformed canola

variety into an already-developed canola variety, and the resulting backcross conversion plant would then comprise the transgene(s).

As an embodiment of the invention, various genetic elements can be introduced into the plant genome using transformation techniques. These elements include, but are not limited to genes, coding sequences, inducible, constitutive, and tissue specific promoters, enhancing sequences, and signal and targeting sequences. For example, see the traits, genes and transformation methods listed in U.S. Pat. No. 6,118,055, incorporated herein by reference in its entirety.

Plant transformation involves the construction of an expression vector which will function in plant cells. Such a vector comprises DNA comprising a gene under control of or operatively linked to a regulatory element (for example, a promoter). The expression vector may contain one or more such operably linked gene/regulatory element combinations. The vector(s) may be in the form of a plasmid, and can be used alone or in combination with other plasmids, to provide transformed canola plants, using transformation methods as described below to incorporate transgenes into the genetic material of the canola plant(s).

Expression Vectors for Canola Transformation: Marker Genes

Expression vectors include at least one genetic marker operably linked to a regulatory element (a promoter, for example) that allows transformed cells containing the marker to be either recovered by negative selection, i.e., inhibiting growth of cells that do not contain the selectable marker gene, or by positive selection, i.e., screening for the product encoded by the genetic marker. Any selectable marker known in the art and/or discussed herein may be used in the present invention. Many commonly used selectable marker genes for plant transformation are well-known in the transformation arts, and include, for example, genes that code for enzymes that metabolically detoxify a selective chemical agent which may be an antibiotic or an herbicide, or genes that encode an altered target which is insensitive to the inhibitor. A few positive selection methods are also known in the art.

One commonly used selectable marker gene for plant transformation is the neomycin phosphotransferase II (nptII) gene which, when under the control of plant regulatory signals, confers resistance to kanamycin. Fraley, et al., *Proc. Natl. Acad. Sci. U.S.A.*, 80:4803 (1983). Another commonly used selectable marker gene is the hygromycin phosphotransferase gene which confers resistance to the antibiotic hygromycin. Vanden Elzen, et al., *Plant Mol. Biol.*, 5:299 (1985).

Additional selectable marker genes of bacterial origin that confer resistance to antibiotics include gentamycin acetyl transferase, streptomycin phosphotransferase, aminoglycoside-3'-adenyl transferase and the bleomycin resistance determinant. Hayford, et al., *Plant Physiol.* 86:1216 (1988), Jones, et al., *Mol. Gen. Genet.*, 210:86 (1987), Svab, et al., *Plant Mol. Biol.* 14:197 (1990) Hille, et al., *Plant Mol. Biol.* 7:171 (1986). Other selectable marker genes confer resistance or tolerance to herbicides such as glyphosate, glufosinate or bromoxynil. Comai, et al., *Nature* 317:741-744 (1985), Gordon-Kamm, et al., *Plant Cell* 2:603-618 (1990) and Stalker, et al., *Science* 242:419-423 (1988).

Selectable marker genes for plant transformation not of bacterial origin include, for example, mouse dihydrofolate reductase, plant 5-enolpyruvylshikimate-3-phosphate synthase and plant acetolactate synthase. Eichholtz, et al., *Somatic Cell Mol. Genet.* 13:67 (1987), Shah, et al., *Science* 233:478 (1986), Charest, et al., *Plant Cell Rep.* 8:643 (1990).

Another class of marker genes for plant transformation requires screening of presumptively transformed plant cells rather than direct genetic selection of transformed cells for resistance to a toxic substance such as an antibiotic. These genes are particularly useful to quantify or visualize the spatial pattern of expression of a gene in specific tissues and are frequently referred to as reporter genes because they can be fused to a gene or gene regulatory sequence for the investigation of gene expression. Commonly used genes for screening presumptively transformed cells include β -glucuronidase (GUS), β -galactosidase, luciferase and chloramphenicol acetyltransferase. Jefferson, R. A., *Plant Mol. Biol. Rep.* 5:387 (1987), Teeri, et al., *EMBO J.* 8:343 (1989), Koncz, et al., *Proc. Natl. Acad. Sci. U.S.A.* 84:131 (1987), DeBlock, et al., *EMBO J.* 3:1681 (1984).

In vivo methods for visualizing GUS activity that do not require destruction of plant tissue are available. Molecular Probes publication 2908, IMAGENE GREEN J., p. 1-4 (1993) and Naleway, et al., *J. Cell Biol.* 115:151a (1991). However, these in vivo methods for visualizing GUS activity have not proven useful for recovery of transformed cells because of low sensitivity, high fluorescent backgrounds and limitations associated with the use of luciferase genes as selectable markers.

More recently, a gene encoding Green Fluorescent Protein (GFP) has been utilized as a marker for gene expression in prokaryotic and eukaryotic cells. Chalfie, et al., *Science* 263:802 (1994). GFP and mutants of GFP may be used as screenable markers.

Expression Vectors for Canola Transformation: Promoters

Genes included in expression vectors must be driven by a nucleotide sequence comprising a regulatory element, for example, a promoter. Any promoter known in the art and/or discussed herein may be used in the present invention. Several types of promoters are now well-known in the transformation arts, as are other regulatory elements that can be used alone or in combination with promoters.

As used herein, "promoter" includes reference to a region of DNA upstream from the start of transcription and involved in recognition and binding of RNA polymerase and other proteins to initiate transcription. A "plant promoter" is a promoter capable of initiating transcription in plant cells. Examples of promoters under developmental control include promoters that preferentially initiate transcription in certain tissues, such as leaves, roots, seeds, fibers, xylem vessels, tracheids, or sclerenchyma. Such promoters are referred to as "tissue-preferred". Promoters which initiate transcription only in certain tissues are referred to as "tissue-specific". A "cell type" specific promoter primarily drives expression in certain cell types in one or more organs, for example, vascular cells in roots or leaves. An "inducible" promoter is a promoter which is under environmental control. Examples of environmental conditions that may effect transcription by inducible promoters include anaerobic conditions or the presence of light. Tissue-specific, tissue-preferred, cell type specific, and inducible promoters constitute the class of "non-constitutive" promoters. A "constitutive" promoter is a promoter which is active under most environmental conditions.

A. Inducible Promoters—An inducible promoter is operably linked to a gene for expression in canola. Optionally, the inducible promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a gene for expression in canola.

With an inducible promoter the rate of transcription increases in response to an inducing agent.

Any inducible promoter can be used in the instant invention. See Ward, et al., *Plant Mol. Biol.* 22:361-366 (1993). Exemplary inducible promoters include, but are not limited to, that from the activating copper-metallothionein expression I (ACEI) system which responds to copper (Mett, et al., *PNAS* 90:4567-4571 (1993)); In2 gene from maize which responds to benzenesulfonamide herbicide safeners (Hershey, et al., *Mol. Gen. Genetics* 227:229-237 (1991) and Gatz, et al., *Mol. Gen. Genetics* 243:32-38 (1994)) or Tet repressor from Tn10 (Gatz, et al., *Mol. Gen. Genetics* 227:229-237 (1991)). A particular inducible promoter responds to an inducing agent to which plants do not normally respond. An exemplary inducible promoter is the inducible promoter from a steroid hormone gene, the transcriptional activity of which is induced by a glucocorticosteroid hormone. Schena, et al., *Proc. Natl. Acad. Sci. U.S.A.* 88:0421 (1991).

B. Constitutive Promoters—A constitutive promoter is operably linked to a gene for expression in canola or the constitutive promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a gene for expression in canola.

Many different constitutive promoters can be utilized in the instant invention. Exemplary constitutive promoters include, but are not limited to, the promoters from plant viruses such as the 35S promoter from CaMV (Odell, et al., *Nature* 313:810-812 (1985)) and the promoters from such genes as rice actin (McElroy, et al., *Plant Cell* 2:163-171 (1990)); ubiquitin (Christensen, et al., *Plant Mol. Biol.* 12:619-632 (1989) and Christensen, et al., *Plant Mol. Biol.* 18:675-689 (1992)); pEMU (Last, et al., *Theor. Appl. Genet.* 81:581-588 (1991)); MAS (Velten, et al., *EMBO J.* 3:2723-2730 (1984)) and maize H3 histone (Lepetit, et al., *Mol. Gen. Genetics* 231:276-285 (1992) and Atanassova, et al., *Plant Journal* 2 (3): 291-300 (1992)). The ALS promoter, Xba1/Nco1 fragment 5' to the *Brassica napus* ALS3 structural gene (or a nucleotide sequence similarity to said Xba1/Nco1 fragment), represents a particularly useful constitutive promoter. See PCT Application No. WO 96/30530, incorporated herein by reference in its entirety.

C. Tissue-Specific or Tissue-Preferred Promoters—A tissue-specific promoter is operably linked to a gene for expression in canola. Optionally, the tissue-specific promoter is operably linked to a nucleotide sequence encoding a signal sequence which is operably linked to a gene for expression in canola. Plants transformed with a gene of interest operably linked to a tissue-specific promoter produce the protein product of the transgene exclusively, or preferentially, in a specific tissue.

Any tissue-specific or tissue-preferred promoter can be utilized in the instant invention. Exemplary tissue-specific or tissue-preferred promoters include, but are not limited to, a root-preferred promoter such as that from the phaseolin gene (Murai, et al., *Science* 23:476-482 (1983) and Sengupta-Gopalan, et al., *Proc. Natl. Acad. Sci. U.S.A.* 82:3320-3324 (1985)); a leaf-specific and light-induced promoter such as that from cab or rubisco (Simpson, et al., *EMBO J.* 4(11):2723-2729 (1985) and Timko, et al., *Nature* 318:579-582 (1985)); an anther-specific promoter such as that from LAT52 (Twell, et al., *Mol. Gen. Genetics* 217:240-245 (1989)); a pollen-specific promoter such as that from Zm13 (Guerrero, et al., *Mol. Gen. Genetics* 244:161-168 (1993)) or a microspore-preferred promoter such as that from apg (Twell, et al., *Sex. Plant Reprod.* 6:217-224 (1993)).

Signal Sequences for Targeting Proteins to Subcellular Compartments

Transport of protein produced by transgenes to a subcellular compartment such as the chloroplast, vacuole, peroxisome, glyoxysome, cell wall or mitochondrion or for secretion into the apoplast, is accomplished by means of operably linking the nucleotide sequence encoding a signal sequence to the 5' and/or 3' region of a gene encoding the protein of interest. Targeting sequences at the 5' and/or 3' end of the structural gene may determine, during protein synthesis and processing, where the encoded protein is ultimately compartmentalized.

The presence of a signal sequence directs a polypeptide to either an intracellular organelle or subcellular compartment or for secretion to the apoplast. Many signal sequences are known in the art. See e.g. Becker, et al., *Plant Mol. Biol.* 20:49 (1992), Knox, C., et al., *Plant Mol. Biol.* 9:3-17 (1987), Lerner, et al., *Plant Physiol.* 91:124-129 (1989), Fontes, et al., *Plant Cell* 3:483-496 (1991), Matsuoka, et al., *Proc. Natl. Acad. Sci.* 88:834 (1991), Gould, et al., *J. Cell. Biol.* 108:1657 (1989), Creissen, et al., *Plant J.* 2:129 (1991), Kalderon, et al., *Cell* 39:499-509 (1984), Steifel, et al., *Plant Cell* 2:785-793 (1990). Any signal sequence known in the art and/or discussed herein may be used in the present invention.

Foreign Protein Genes and Agronomic Genes

With transgenic plants according to the present invention, a foreign protein can be produced in commercial quantities. Thus, techniques for the selection and propagation of transformed plants, which are well understood in the art, yield a plurality of transgenic plants which are harvested in a conventional manner, and a foreign protein then can be extracted from a tissue of interest or from total biomass. Protein extraction from plant biomass can be accomplished by known methods which are discussed, for example, by Heney and Orr, *Anal. Biochem.* 114:92-6 (1981).

According to an embodiment, the transgenic plant provided for commercial production of foreign protein is a canola plant. In another embodiment, the biomass of interest is seed. For the relatively small number of transgenic plants that show higher levels of expression, a genetic map can be generated, primarily via conventional restriction fragment length polymorphism (RFLP), polymerase chain reaction (PCR), short sequence repeats (SSR) analysis, which identifies the approximate chromosomal location of the integrated DNA molecule. For exemplary methodologies in this regard, see Glick and Thompson, *Methods in Plant Molecular Biology and Biotechnology* CRC Press, Boca Raton 269:284 (1993). Map information concerning chromosomal location is useful for proprietary protection of a subject transgenic plant. If unauthorized propagation is undertaken and crosses made with other germplasm, the map of the integration region can be compared to similar maps for suspect plants, to determine if the latter have a common parentage with the subject plant. Map comparisons would involve hybridizations, RFLP, PCR, SSR and sequencing, all of which are conventional techniques.

Wang, et al. discuss "Large Scale Identification, Mapping and Genotyping of Single-Nucleotide Polymorphisms in the Human Genome", *Science*, 280:1077-1082, 1998, and similar capabilities are becoming increasingly available for the canola genome. Map information concerning chromosomal location is useful for proprietary protection of a subject transgenic plant. If unauthorized propagation is undertaken and crosses made with other germplasm, the map of the integra-

tion region can be compared to similar maps for suspect plants to determine if the latter have a common parentage with the subject plant. Map comparisons would involve hybridizations, RFLP, PCR, SSR and sequencing, all of which are conventional techniques. Single nucleotide polymorphism (SNP)s may also be used alone or in combination with other techniques.

In an embodiment of the present invention, plants can be genetically engineered to express various phenotypes of agronomic interest. Through the transformation of canola the expression of genes can be altered to enhance disease resistance, insect resistance, herbicide tolerance, agronomic, grain quality and other traits. Transformation can also be used to insert DNA sequences which control or help control male-sterility. DNA sequences native to canola as well as non-native DNA sequences can be transformed into canola and used to alter levels of native or non-native proteins. Various promoters, targeting sequences, enhancing sequences, and other DNA sequences can be inserted into the genome for the purpose of altering the expression of proteins. Reduction of the activity of specific genes (also known as gene silencing, or gene suppression) may also be desirable for several aspects of genetic engineering in plants.

Many techniques for gene silencing are well-known to one of skill in the art, including but not limited to knock-outs (such as by insertion of a transposable element such as mu (Vicki Chandler, *The Maize Handbook* ch. 118, Springer-Verlag, 1994) or other genetic elements such as a FRT, Lox or other site specific integration site, antisense technology (see, e.g., Sheehy, et al. (1988) *PNAS USA* 85:8805-8809; and U.S. Pat. Nos. 5,107,065; 5,453,566; and 5,759,829, incorporated herein by reference in their entirety); co-suppression (e.g., Taylor (1997) *Plant Cell* 9:1245; Jorgensen (1990) *Trends Biotech.* 8(12):340-344; Flavell (1994) *PNAS USA* 91:3490-3496; Finnegan, et al. (1994) *Bio/Technology* 12: 883-888; and Neuhuber, et al. (1994) *Mol. Gen. Genet.* 244:230-241); RNA interference (Napoli, et al. (1990) *Plant Cell* 2:279-289; U.S. Pat. No. 5,034,323, incorporated herein by reference in its entirety; Sharp (1999) *Genes Dev.* 13:139-141; Zamore, et al. (2000) *Cell* 101:25-33; and Montgomery, et al. (1998) *PNAS USA* 95:15502-15507), virus-induced gene silencing (Burton, et al. (2000) *Plant Cell* 12:691-705; and Baulcombe (1999) *Curr. Op. Plant Bio.* 2:109-113); target-RNA-specific ribozymes (Haseloff, et al. (1988) *Nature* 334: 585-591); hairpin structures (Smith, et al. (2000) *Nature* 407:319-320; WO 99/53050; and WO 98/53083), incorporated herein by reference in their entirety; MicroRNA (Aukerman & Sakai (2003) *Plant Cell* 15:2730-2741); ribozymes (Steinecke, et al. (1992) *EMBO J.* 11:1525; and Perriman, et al. (1993) *Antisense Res. Dev.* 3:253); oligonucleotide mediated targeted modification (e.g., WO 03/076574 and WO 99/25853, incorporated herein by reference in their entirety); Zn-finger targeted molecules (e.g., WO 01/52620; WO 03/048345; and WO 00/42219, incorporated herein by reference in their entirety); and other methods or combinations of the above methods known to those of skill in the art.

Likewise, by means of the present invention, agronomic genes can be expressed in transformed plants. More particularly, plants can be genetically engineered to express various phenotypes of agronomic interest. Exemplary genes implicated in this regard include, but are not limited to, those categorized below:

1. Genes that Confer Resistance to Pests or Disease and that Encode:

A. Plant disease resistance genes. Plant defenses are often activated by specific interaction between the product of a disease resistance gene (R) in the plant and the product

of a corresponding avirulence (Avr) gene in the pathogen. A plant variety can be transformed with cloned resistance genes to engineer plants that are resistant to specific pathogen strains. See, for example Jones, et al., *Science* 266:789 (1994) (cloning of the tomato Cf-9 gene for resistance to *Cladosporium fulvum*); Martin, et al., *Science* 262:1432 (1993) (tomato Pto gene for resistance to *Pseudomonas syringae* pv. tomato encodes a protein kinase); Mindrinos, et al., *Cell* 78:1089 (1994) (Arabidopsis RSP2 gene for resistance to *Pseudomonas syringae*), McDowell & Woffenden, (2003) *Trends Biotechnol.* 21(4): 178-83 and Toyoda, et al., (2002) *Transgenic Res.* 11 (6):567-82.

B. A gene conferring resistance to fungal pathogens, such as oxalate oxidase or oxalate decarboxylase (Zhou, et al., (1998) *Pl. Physiol.* 117:33-41).

C. A *Bacillus thuringiensis* protein, a derivative thereof or a synthetic polypeptide modeled thereon. See, for example, Geiser, et al., *Gene* 48:109 (1986), who disclose the cloning and nucleotide sequence of a Bt δ -endotoxin gene. Moreover, DNA molecules encoding δ -endotoxin genes can be purchased from American Type Culture Collection, Manassas, Va., for example, under ATCC Accession Nos. 40098, 67136, 31995 and 31998.

D. A lectin. See, for example, the disclosure by Van Damme, et al., *Plant Molec. Biol.* 24:25 (1994), who disclose the nucleotide sequences of several *Clivia miniata* mannose-binding lectin genes.

E. A vitamin-binding protein such as avidin. See PCT Application No. WO 93/06487, incorporated herein by reference in its entirety. The application teaches the use of avidin and avidin homologues as larvicides against insect pests.

F. An enzyme inhibitor, for example, a protease or proteinase inhibitor or an amylase inhibitor. See, for example, Abe, et al., *J. Biol. Chem.* 262:16793 (1987) (nucleotide sequence of rice cysteine proteinase inhibitor), Huub, et al., *Plant Molec. Biol.* 21:985 (1993) (nucleotide sequence of cDNA encoding tobacco proteinase inhibitor I), Sumitani, et al., *Biosci. Biotech. Biochem.* 57:1243 (1993) (nucleotide sequence of *Streptomyces nitrosporeus* α -amylase inhibitor) and U.S. Pat. No. 5,494,813 (Hepher and Atkinson, issued Feb. 27, 1996), incorporated herein by reference in its entirety.

G. An insect-specific hormone or pheromone such as an ecdysteroid or juvenile hormone, a variant thereof, a mimetic based thereon, or an antagonist or agonist thereof. See, for example, the disclosure by Hammock, et al., *Nature* 344:458 (1990), of baculovirus expression of cloned juvenile hormone esterase, an inactivator of juvenile hormone.

H. An insect-specific peptide or neuropeptide which, upon expression, disrupts the physiology of the affected pest. For example, see the disclosures of Regan, *J. Biol. Chem.* 269:9 (1994) (expression cloning yields DNA coding for insect diuretic hormone receptor), and Pratt, et al., *Biochem. Biophys. Res. Comm.* 163:1243 (1989) (an allostatin is identified in *Diploptera punctata*). See also U.S. Pat. No. 5,266,317, incorporated herein by reference in its entirety, to Tomalski, et al., who disclose genes encoding insect-specific, paralytic neurotoxins.

I. An insect-specific venom produced in nature by a snake, a wasp, etc. For example, see Pang, et al., *Gene* 116:165 (1992), for disclosure of heterologous expression in plants of a gene coding for a scorpion insectotoxic peptide.

- J. An enzyme responsible for a hyperaccumulation of a monoterpene, a sesquiterpene, a steroid, hydroxamic acid, a phenylpropanoid derivative or another non-protein molecule with insecticidal activity.
- K. An enzyme involved in the modification, including the post-translational modification, of a biologically active molecule; for example, a glycolytic enzyme, a proteolytic enzyme, a lipolytic enzyme, a nuclease, a cyclase, a transaminase, an esterase, a hydrolase, a phosphatase, a kinase, a phosphorylase, a polymerase, an elastase, a chitinase and a glucanase, whether natural or synthetic. See PCT Application No. WO 93/02197, incorporated herein by reference in its entirety, in the name of Scott, et al., which discloses the nucleotide sequence of a callase gene. DNA molecules which contain chitinase-encoding sequences can be obtained, for example, from the ATCC under Accession Nos. 39637 and 67152. See also Kramer, et al., *Insect Biochem. Molec. Biol.* 23:691 (1993), who teach the nucleotide sequence of a cDNA encoding tobacco hornworm chitinase, and Kawalleck, et al., *Plant Molec. Biol.* 21:673 (1993), who provide the nucleotide sequence of the parsley ubi4-2 polyubiquitin gene, U.S. Pat. Nos. 7,145,060, 7,087,810 and 6,563,020, incorporated herein by reference in their entirety.
- L. A molecule that stimulates signal transduction. For example, see the disclosure by Botella, et al., *Plant Molec. Biol.* 24:757 (1994), of nucleotide sequences for mung bean calmodulin cDNA clones, and Griess, et al., *Plant Physiol.* 104:1467 (1994), who provide the nucleotide sequence of a maize calmodulin cDNA clone.
- M. A hydrophobic moment peptide. See PCT Application No. WO 95/16776 (disclosure of peptide derivatives of Tachyplesin which inhibit fungal plant pathogens) and PCT application WO 95/18855 and U.S. Pat. No. 5,607,914 which teaches synthetic antimicrobial peptides that confer disease resistance, each of which is incorporated herein by reference in its entirety.
- N. A membrane permease, a channel former or a channel blocker. For example, see the disclosure of Jaynes, et al., *Plant Sci* 89:43 (1993), of heterologous expression of a cecropin- β , lytic peptide analog to render transgenic tobacco plants resistant to *Pseudomonas solanacearum*.
- O. A viral-invasive protein or a complex toxin derived therefrom. For example, the accumulation of viral coat proteins in transformed plant cells imparts resistance to viral infection and/or disease development effected by the virus from which the coat protein gene is derived, as well as by related viruses. See Beachy, et al., *Ann. rev. Phytopathol.* 28:451 (1990). Coat protein-mediated resistance has been conferred upon transformed plants against alfalfa mosaic virus, cucumber mosaic virus, tobacco streak virus, potato virus X, potato virus Y, tobacco etch virus, tobacco rattle virus and tobacco mosaic virus. Id.
- P. An insect-specific antibody or an immunotoxin derived therefrom. Thus, an antibody targeted to a critical metabolic function in the insect gut would inactivate an affected enzyme, killing the insect. Cf. Taylor, et al., Abstract #497, Seventh Int'l Symposium on Molecular Plant-Microbe Interactions (Edinburgh, Scotland) (1994) (enzymatic inactivation in transgenic tobacco via production of single-chain antibody fragments).
- Q. A virus-specific antibody. See, for example, Tavladoraki, et al., *Nature* 366:469 (1993), who show that transgenic plants expressing recombinant antibody genes are protected from virus attack.

- R. A developmental-arrestive protein produced in nature by a pathogen or a parasite. Thus, fungal endo- α -1,4-D-polygalacturonases facilitate fungal colonization and plant nutrient release by solubilizing plant cell wall homo- α -1,4-D-galacturonase. See Lamb, et al., *Bio/Technology* 10:1436 (1992). The cloning and characterization of a gene which encodes a bean endopolygalacturonase-inhibiting protein is described by Toubart, et al., *Plant J.* 2:367 (1992).
- S. A developmental-arrestive protein produced in nature by a plant. For example, Logemann, et al., *Bio/Technology* 10:305 (1992), have shown that transgenic plants expressing the barley ribosome-inactivating gene have an increased resistance to fungal disease.
- T. Genes involved in the Systemic Acquired Resistance (SAR) Response and/or the pathogenesis-related genes. Briggs, S., *Current Biology*, 5 (2) (1995); Pieterse & Van Loon (2004) *Curr. Opin. Plant Bio.* 7(4):456-64 and Somssich (2003) *Cell* 113(7):815-6.
- U. Antifungal genes. See Cornelissen and Meichers, *Plant Physiol.*, 101:709-712 (1993); Parijs, et al., *Planta* 183:258-264 (1991) and Bushnell, et al., *Can. J of Plant Path.* 20(2):137-149 (1998); see also U.S. Pat. No. 6,875,907, incorporated herein by reference in its entirety.
- V. Detoxification genes, such as for fumonisin, beauvericin, moniliformin and zearalenone and their structurally related derivatives. For example, see U.S. Pat. No. 5,792,931, incorporated herein by reference in its entirety.
- W. Cystatin and cysteine proteinase inhibitors. See U.S. Pat. No. 7,205,453, incorporated herein by reference in its entirety.
- X. Defensin genes. See WO 03/000863 and U.S. Pat. No. 6,911,577, incorporated herein by reference in their entirety.
- Y. Genes that confer resistance to *Phytophthora* root rot, such as the *Brassica* equivalents of the Rps 1, Rps 1-a, Rps 1-b, Rps 1-c, Rps 1-d, Rps 1-e, Rps 1-k, Rps 2, Rps 3-a, Rps 3-b, Rps 3-c, Rps 4, Rps 5, Rps 6, Rps 7 and other Rps genes. See, for example, Shoemaker, et al., *Phytophthora* Root Rot Resistance Gene Mapping in Soybean, Plant Genome IV Conference, San Diego, Calif. (1995).
2. Genes that Confer Resistance or Tolerance to an Herbicide:
- A. An herbicide that inhibits the growing point or meristem, such as an imidazolinone or a sulfonylurea. Exemplary genes in this category code for mutant ALS and AHAS enzyme as described, for example, by Lee, et al., *EMBO J.* 7:1241 (1988), and Miki, et al., *Theor. Appl. Genet.* 80:449 (1990), respectively.
- B. Glyphosate (resistance or tolerance conferred by mutant 5-enolpyruvylshikimate-3-phosphate synthase (EPSP) and aroA genes, respectively) and other phosphono compounds such as glufosinate (phosphinothricin acetyl transferase (PAT) and *Streptomyces hygrosopicus* PAT bar genes), and pyridinoyl or phenoxy propionic acids and cyclohexones (ACCase inhibitor-encoding genes). See, for example, U.S. Pat. No. 4,940,835 to Shah, et al., incorporated herein by reference in its entirety, which discloses the nucleotide sequence of a form of EPSP which can confer glyphosate resistance or tolerance. U.S. Pat. No. 5,627,061 to Barry, et al., incorporated herein by reference in its entirety, also describes genes encoding EPSPS enzymes. See also U.S. Pat. Nos. 6,566,587; 6,338,961; 6,248,876 B1; 6,040,497; 5,804,

425; 5,633,435; 5,145,783; 4,971,908; 5,312,910; 5,188,642; 4,940,835; 5,866,775; 6,225,114 B1; 6,130,366; 5,310,667; 4,535,060; 4,769,061; 5,633,448; 5,510,471; Re. 36,449; RE 37,287 E; and 5,491,288; and international publications EP1173580; WO 01/66704; EP1173581 and EP1173582, which are each incorporated herein by reference in their entireties. Glyphosate resistance or tolerance is also imparted to plants that express a gene that encodes a glyphosate oxido-reductase enzyme as described more fully in U.S. Pat. Nos. 5,776,760 and 5,463,175, which are incorporated herein by reference in their entireties. In addition glyphosate resistance or tolerance can be imparted to plants by the overexpression of genes encoding glyphosate N-acetyltransferase. See, for example, U.S. Pat. No. 7,462,481, incorporated herein by reference in its entirety. A DNA molecule encoding a mutant *aroA* gene can be obtained under ATCC accession number 39256, and the nucleotide sequence of the mutant gene is disclosed in U.S. Pat. No. 4,769,061 to Comai, incorporated herein by reference in its entirety. European patent application No. 0 333 033 to Kumada, et al., and U.S. Pat. No. 4,975,374 to Goodman, et al., incorporated herein by reference in their entireties, disclose nucleotide sequences of glutamine synthetase genes which confer resistance or tolerance to herbicides such as L-phosphinothricin.

The nucleotide sequence of a PAT gene is provided in European application No. 0 242 246 to Leemans, et al., incorporated herein by reference in its entirety, DeGreef, et al., *Bio/Technology* 7:61 (1989), describe the production of transgenic plants that express chimeric bar genes coding for PAT activity. Exemplary of genes conferring resistance or tolerance to phenoxy propionic acids and cyclohexones, such as sethoxydim and haloxyfop are the Acc1-S1, Acc1-S2 and Acc1-S3 genes described by Marshall, et al., *Theor. Appl. Genet.* 83:435 (1992).

C. An herbicide that inhibits photosynthesis, such as a triazine (*psbA* and *gs+* genes) and a benzonitrile (nitrilase gene). Przibila, et al., *Plant Cell* 3:169 (1991), describe the transformation of *Chlamydomonas* with plasmids encoding mutant *psbA* genes. Nucleotide sequences for nitrilase genes are disclosed in U.S. Pat. No. 4,810,648 to Stalker, incorporated herein by reference in its entirety, and DNA molecules containing these genes are available under ATCC Accession Nos. 53435, 67441, and 67442. Cloning and expression of DNA coding for a glutathione S-transferase is described by Hayes, et al., *Biochem. J.* 285:173 (1992).

D. Acetohydroxy acid synthase, which has been found to make plants that express this enzyme resistant to multiple types of herbicides, has been introduced into a variety of plants. See Hattori, et al., *Mol. Gen. Genet.* 246:419, 1995. Other genes that confer tolerance to herbicides include a gene encoding a chimeric protein of rat cytochrome P4507A1 and yeast NADPH-cytochrome P450 oxidoreductase (Shiota, et al., *Plant Physiol.*, 106: 17, 1994), genes for glutathione reductase and superoxide dismutase (Aono, et al., *Plant Cell Physiol.* 36:1687, 1995), and genes for various phosphotransferases (Datta, et al., *Plant Mol. Biol.* 20:619, 1992).

E. Protoporphyrinogen oxidase (*protox*) is necessary for the production of chlorophyll, which is necessary for all plant survival. The *protox* enzyme serves as the target for a variety of herbicidal compounds. These herbicides also inhibit growth of all the different species of plants present, causing their total destruction. The development of plants containing altered *protox* activity which

are resistant to these herbicides are described in U.S. Pat. Nos. 6,288,306; 6,282,837; 5,767,373; and international publication WO 01/12825, each of which is incorporated herein by reference in its entirety.

3. Genes that Confer or Contribute to a Value-Added Trait:

A. Modified fatty acid metabolism, for example, by transforming a plant with an antisense gene of stearyl-ACP desaturase to increase stearic acid content of the plant. See Knultzon, et al., *Proc. Natl. Acad. Sci. U.S.A.* 89:2624 (1992).

B. Decreased phytate content: 1) Introduction of a phytase-encoding gene would enhance breakdown of phytate, adding more free phosphate to the transformed plant. For example, see Van Hartingsveldt, et al., *Gene* 127:87 (1993), for a disclosure of the nucleotide sequence of an *Aspergillus niger* phytase gene. 2) A gene could be introduced that reduced phytate content. In maize for example, this could be accomplished by cloning and then reintroducing DNA associated with the single allele which is responsible for maize mutants characterized by low levels of phytic acid. See Raboy, et al., *Maydica* 35:383 (1990) and/or by altering inositol kinase activity as in WO 02/059324, U.S. Pat. No. 7,067,720, WO 03/027243, US2003/0079247, WO 99/05298, U.S. Pat. Nos. 6,197,561, 6,291,224, 6,391,348, WO 2002/059324, WO 98/45448, WO 99/55882, WO 01/04147, each of which is incorporated herein by reference in its entirety.

C. Modified carbohydrate composition effected, for example, by transforming plants with a gene coding for an enzyme that alters the branching pattern of starch, or, a gene altering thioredoxin such as NTR and/or TRX (See U.S. Pat. No. 6,531,648, which is incorporated herein by reference in its entirety) and/or a gamma zein knock out or mutant such as *cs27* or *TUSC27* or *en27* (See U.S. Pat. No. 6,858,778 and U.S. Patent Application Nos. 2005/0160488, 2005/0204418; which are incorporated herein by reference in their entireties). See Shiroza, et al., *J. Bacteriol.* 170: 810 (1988) (nucleotide sequence of *Streptococcus mutans* fructosyltransferase gene), Steinmetz, et al., *Mol. Gen. Genet.* 200: 220 (1985) (nucleotide sequence of *Bacillus subtilis* levansucrase gene), Pen, et al., *Bio/Technology* 10: 292 (1992) (production of transgenic plants that express *Bacillus licheniformis* alpha-amylase), Elliot, et al., *Plant Molec. Biol.* 21: 515 (1993) (nucleotide sequences of tomato invertase genes), Sogaard, et al., *J. Biol. Chem.* 268: 22480 (1993) (site-directed mutagenesis of barley alpha-amylase gene), and Fisher, et al., *Plant Physiol.* 102: 1045 (1993) (maize endosperm starch branching enzyme II), WO 99/10498 (improved digestibility and/or starch extraction through modification of UDP-D-xylose 4-epimerase, Fragile 1 and 2, 1, HCHL, C4H), U.S. Pat. No. 6,232,529 (method of producing high oil seed by modification of starch levels (AGP)), each of which is incorporated herein by reference in its entirety. The fatty acid modification genes mentioned above may also be used to affect starch content and/or composition through the interrelationship of the starch and oil pathways.

D. Elevated oleic acid via FAD-2 gene modification and/or decreased linolenic acid via FAD-3 gene modification. See U.S. Pat. Nos. 6,063,947; 6,323,392; and international publication WO 93/11245, each of which is incorporated herein by reference in its entirety.

E. Altering conjugated linolenic or linoleic acid content, such as in WO 01/12800, incorporated herein by refer-

ence in its entirety. Altering LEC1, AGP, Dek1, Supera11, mi 1 ps, various Ipa genes such as Ipa1, Ipa3, hpt or hgg. For example, see WO 02/42424, WO 98/22604, WO 03/011015, U.S. Pat. Nos. 6,423,886, 6,197,561, 6,825,397, 7,157,621, U.S. Patent Application No. 2003/0079247, International publications WO 02/057439 and WO 03/011015 and Rivera-Madrid, R., et al. *Proc. Natl. Acad. Sci.* 92:5620-5624 (1995), each of which is incorporated herein by reference in its entirety.

F. Altered antioxidant content or composition, such as alteration of tocopherol or tocotrienols. For example, see U.S. Pat. Nos. 6,787,683 and 7,154,029 and WO 00/68393 involving the manipulation of antioxidant levels through alteration of a phytyl prenyl transferase (ppt), WO 03/082899 through alteration of a homogentisate geranyl geranyl transferase (hgg), each of which is incorporated herein by reference in its entirety.

G. Altered essential seed amino acids. For example, see U.S. Pat. No. 6,127,600 (method of increasing accumulation of essential amino acids in seeds), U.S. Pat. No. 6,080,913 (binary methods of increasing accumulation of essential amino acids in seeds), U.S. Pat. No. 5,990,389 (high lysine), WO 99/40209 (alteration of amino acid compositions in seeds), WO 99/29882 (methods for altering amino acid content of proteins), U.S. Pat. No. 5,850,016 (alteration of amino acid compositions in seeds), WO 98/20133 (proteins with enhanced levels of essential amino acids), U.S. Pat. No. 5,885,802 (high methionine), U.S. Pat. No. 5,885,801 (high threonine), U.S. Pat. No. 6,664,445 (plant amino acid biosynthetic enzymes), U.S. Pat. No. 6,459,019 (increased lysine and threonine), U.S. Pat. No. 6,441,274 (plant tryptophan synthase beta subunit), U.S. Pat. No. 6,346,403 (methionine metabolic enzymes), U.S. Pat. No. 5,939,599 (high sulfur), U.S. Pat. No. 5,912,414 (increased methionine), WO 98/56935 (plant amino acid biosynthetic enzymes), WO 98/45458 (engineered seed protein having higher percentage of essential amino acids), WO 98/42831 (increased lysine), U.S. Pat. No. 5,633,436 (increasing sulfur amino acid content), U.S. Pat. No. 5,559,223 (synthetic storage proteins with defined structure containing programmable levels of essential amino acids for improvement of the nutritional value of plants), WO 96/01905 (increased threonine), WO 95/15392 (increased lysine), U.S. Pat. Nos. 6,930,225, 7,179,955, U.S. Patent Application No. 2004/0068767, U.S. Pat. No. 6,803,498, WO 01/79516, and WO 00/09706 (Ces A: cellulose synthase), U.S. Pat. No. 6,194,638 (hemicellulose), U.S. Pat. No. 6,399,859 and U.S. Pat. No. 7,098,381 (UDPGdH), U.S. Pat. No. 6,194,638 (RGP), each of which is incorporated herein by reference in its entirety.

4. Genes that Control Male Sterility:
There are several methods of conferring genetic male sterility available, such as multiple mutant genes at separate locations within the genome that confer male sterility, as disclosed in U.S. Pat. Nos. 4,654,465 and 4,727,219 to Brar, et al. and chromosomal translocations as described by Patterson in U.S. Pat. Nos. 3,861,709 and 3,710,511, each of which is incorporated herein by reference in its entirety. In addition to these methods, Albertsen, et al., U.S. Pat. No. 5,432,068, incorporated by reference in its entirety, describe a system of nuclear male sterility which includes: identifying a gene which is critical to male fertility; silencing this native gene which is critical to male fertility; removing the native

promoter from the essential male fertility gene and replacing it with an inducible promoter; inserting this genetically engineered gene back into the plant; and thus creating a plant that is male sterile because the inducible promoter is not "on" resulting in the male fertility gene not being transcribed. Fertility is restored by inducing, or turning "on", the promoter, which in turn allows the gene that confers male fertility to be transcribed.

A. Introduction of a deacetylase gene under the control of a tapetum-specific promoter and with the application of the chemical N-Ac-PPT. See PCT Application No. WO 01/29237, incorporated herein by reference in its entirety.

B. Introduction of various stamen-specific promoters. See PCT Application Nos. WO 92/13956 and WO 92/13957, each of which is incorporated herein by reference in its entirety.

C. Introduction of the barnase and the barstar genes. See Paul, et al., *Plant Mol. Biol.* 19:611-622, 1992).

For additional examples of nuclear male and female sterility systems and genes, see U.S. Pat. Nos. 5,859,341; 6,297,426; 5,478,369; 5,824,524; 5,850,014; and 6,265,640, each of which is incorporated herein by reference in its entirety.

5. Genes that Create a Site for Site-Specific DNA Integration:
This includes the introduction of FRT sites that may be used in the FLP/FRT system and/or Lox sites that may be used in the Cre/Loxp system. For example, see Lyznik, et al., Site-Specific Recombination for Genetic Engineering in Plants, *Plant Cell Rep* (2003) 21:925-932 and WO 99/25821, each of which is incorporated herein by reference in its entirety. Other systems that may be used include the Gin recombinase of phage Mu (Maeser, et al., 1991; Vicki Chandler, *The Maize Handbook* ch. 118, Springer-Verlag, 1994), the Pin recombinase of *E. coli* (Enomoto, et al., 1983), and the R/RS system of the pSRi plasmid (Araki, et al., 1992).

6. Genes that Affect Abiotic Stress Resistance:
Genes that affect abiotic stress resistance (including but not limited to flowering, pod and seed development, enhancement of nitrogen utilization efficiency, altered nitrogen responsiveness, drought resistance or tolerance, cold resistance or tolerance, and salt resistance or tolerance) and increased yield under stress. For example, see WO 00/73475 where water use efficiency is altered through alteration of malate; U.S. Pat. Nos. 5,892,009, 5,965,705, 5,929,305, 5,891,859, 6,417,428, 6,664,446, 6,706,866, 6,717,034, 6,801,104, international patent publications WO 2000/060089, WO 2001/026459, WO 2001/035725, WO 2001/034726, WO 2001/035727, WO 2001/036444, WO 2001/036597, WO 2001/036598, WO 2002/015675, WO 2002/017430, WO 2002/077185, WO 2002/079403, WO 2003/013227, WO 2003/013228, WO 2003/014327, WO 2004/031349, WO 2004/076638, WO 98/09521, and WO 99/38977, each of which is incorporated by reference in its entirety, describing genes, including CBF genes and transcription factors effective in mitigating the negative effects of freezing, high salinity, and drought on plants, as well as conferring other positive effects on plant phenotype; U.S. Patent Application No. 2004/0148654 and international publication WO 01/36596, each of which is incorporated herein by reference in its entirety, where abscisic acid is altered in plants resulting in improved plant phenotype such as increased yield and/or increased tolerance to abiotic

stress; WO 2000/006341, WO 04/090143, U.S. Pat. Nos. 7,531,723 and 6,992,237, each of which is incorporated herein by reference in its entirety, where cytokinin expression is modified resulting in plants with increased stress tolerance, such as drought tolerance, and/or increased yield. Also see WO 02/02776, WO 2003/052063, JP2002281975, U.S. Pat. No. 6,084,153, WO 01/64898, U.S. Pat. Nos. 6,177,275 and 6,107,547 (enhancement of nitrogen utilization and altered nitrogen responsiveness). For ethylene alteration, see U.S. Application Nos. 2004/0128719, 2003/0166197 and WO 2000/32761, each of which is incorporated herein by reference in its entirety. For plant transcription factors or transcriptional regulators of abiotic stress, see e.g. U.S. Application No. 2004/0098764 or U.S. Application No. 2004/0078852, each of which is incorporated herein by reference in its entirety.

Other genes and transcription factors that affect plant growth and agronomic traits such as yield, flowering, plant growth and/or plant structure, can be introduced or introgressed into plants, see e.g. WO 97/49811 (LHY), WO 98/56918 (ESD4), WO 97/10339 and U.S. Pat. Nos. 6,573,430 (TFL), 6,713,663 (FT), 6,794,560, 6,307,126 (GAI), WO 96/14414 (CON), WO 96/38560, WO 01/21822 (VRN1), WO 00/44918 (VRN2), WO 99/49064 (GI), WO 00/46358 (FR1), WO 97/29123, WO 99/09174 (D8 and Rht), and WO 2004/076638 and WO 2004/031349 (transcription factors), each of which is incorporated herein by reference in its entirety.

METHODS FOR CANOLA TRANSFORMATION

Numerous methods for plant transformation have been developed including biological and physical plant transformation protocols. See, for example, Miki, et al., "Procedures for Introducing Foreign DNA into Plants" in *Methods in Plant Molecular Biology and Biotechnology*, Glick B. R. and Thompson, J. E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 67-88. In addition, expression vectors and in vitro culture methods for plant cell or tissue transformation and regeneration of plants are available. See, for example, Gruber, et al., "Vectors for Plant Transformation" in *Methods in Plant Molecular Biology and Biotechnology*, Glick B. R. and Thompson, J. E. Eds. (CRC Press, Inc., Boca Raton, 1993) pages 89-119.

A. *Agrobacterium*-Mediated Transformation—One method for introducing an expression vector into plants is based on the natural transformation system of *Agrobacterium*. See, for example, Horsch, et al., *Science* 227:1229 (1985). *A. tumefaciens* and *A. rhizogenes* are plant pathogenic soil bacteria which genetically transform plant cells. The Ti and Ri plasmids of *A. tumefaciens* and *A. rhizogenes*, respectively, carry genes responsible for genetic transformation of the plant. See, for example, Kado, C. I., *Crit. Rev. Plant Sci.* 10:1 (1991). Descriptions of *Agrobacterium* vector systems and methods for *Agrobacterium*-mediated gene transfer are provided by Gruber, et al., supra, Miki, et al., supra, and Moloney, et al., *Plant Cell Reports* 8:238 (1989). See also, U.S. Pat. No. 5,563,055 (Townsend and Thomas), incorporated herein by reference in its entirety.

B. Direct Gene Transfer—Several methods of plant transformation, collectively referred to as direct gene transfer, have been developed as an alternative to *Agrobacterium*-mediated transformation. A generally applicable method of plant transformation is microprojectile-mediated transformation wherein DNA is carried on the

surface of microprojectiles measuring 1 to 4 μm . The expression vector is introduced into plant tissues with a biolistic device that accelerates the microprojectiles to speeds of 300 to 600 m/s which is sufficient to penetrate plant cell walls and membranes. Sanford, et al., *Part. Sci. Technol.* 5:27 (1987), Sanford, J. C., *Trends Biotech.* 6:299 (1988), Klein, et al., *Bio/Technology* 6:559-563 (1988), Sanford, J. C., *Physiol Plant* 7:206 (1990), Klein, et al., *Biotechnology* 10:268 (1992). See also U.S. Pat. No. 5,015,580 (Christou, et al.) and U.S. Pat. No. 5,322,783 (Tomes, et al.), each of which is incorporated herein by reference in its entirety.

Another method for physical delivery of DNA to plants is sonication of target cells. Zhang, et al., *Bio/Technology* 9:996 (1991).

Alternatively, liposome and spheroplast fusion have been used to introduce expression vectors into plants. Deshayes, et al., *EMBO J.* 4:2731 (1985), Christou, et al., *Proc Natl. Acad. Sci. U.S.A.* 84:3962 (1987). Direct uptake of DNA into protoplasts using CaCl_2 precipitation, polyvinyl alcohol or poly-L-ornithine has also been reported. Hain, et al., *Mol. Gen. Genet.* 199:161 (1985) and Draper, et al., *Plant Cell Physiol.* 23:451 (1982). Electroporation of protoplasts and whole cells and tissues have also been described. Donn, et al., In Abstracts of VIIth International Congress on Plant Cell and Tissue Culture IAPTC, A2-38, p 53 (1990); D'Halluin, et al., *Plant Cell* 4:1495-1505 (1992) and Spencer, et al., *Plant Mol. Biol.* 24:51-61 (1994).

Following transformation of canola target tissues, expression of the above-described selectable marker genes allows for preferential selection of transformed cells, tissues and/or plants, using regeneration and selection methods well-known in the art.

The foregoing methods for transformation would typically be used for producing a transgenic variety. The transgenic variety could then be crossed with another (non-transformed or transformed) variety, in order to produce a new transgenic variety. Alternatively, a genetic trait which has been engineered into a particular canola line using the foregoing transformation techniques could be moved into another line using traditional backcrossing techniques that are well-known in the plant breeding arts. For example, a backcrossing approach could be used to move an engineered trait from a public, non-elite variety into an elite variety, or from a variety containing a foreign gene in its genome into a variety or varieties which do not contain that gene. As used herein, "crossing" can refer to a simple X by Y cross, or the process of backcrossing, depending on the context.

Genetic Marker Profile Through SSR and First Generation Progeny

In addition to phenotypic observations, a plant can also be identified by its genotype. The genotype of a plant can be characterized through a genetic marker profile which can identify plants of the same variety or a related variety or be used to determine or validate a pedigree. Genetic marker profiles can be obtained by techniques such as RFLP, Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARs), Amplified Fragment Length Polymorphisms (AFLPs), SSRs which are also referred to as Microsatellites, and SNPs. For exemplary methodologies, see Glick, et al., 1993. *Methods in Plant Molecular Biology and Biotechnology*. CRC Press, Boca Raton.

Particular markers used for these purposes are not limited to any particular set of markers, but are envisioned to include any type of marker and marker profile which provides a means of distinguishing varieties. One method of comparison is to use only homozygous loci for SCV470336.

In addition to being used for identification of canola line SCV470336 and plant parts and plant cells of line SCV470336, the genetic profile may be used to identify a canola plant produced through the use of SCV470336 or to verify a pedigree for progeny plants produced through the use of SCV470336. The genetic marker profile is also useful in breeding and developing backcross conversions.

The present invention comprises a canola plant characterized by molecular and physiological data obtained from the representative sample of said variety deposited with the American Type Culture Collection (ATCC). Further provided by the invention is a canola plant formed by the combination of the disclosed canola plant or plant cell with another canola plant or cell and comprising the homozygous alleles of the variety.

Means of performing genetic marker profiles using SSR polymorphisms are well-known in the art. SSRs are genetic markers based on polymorphisms in repeated nucleotide sequences, such as microsatellites. A marker system based on SSRs can be highly informative in linkage analysis relative to other marker systems in that multiple alleles may be present. Another advantage of this type of marker is that, through use of flanking primers, detection of SSRs can be achieved, for example, by PCR, thereby eliminating the need for labor-intensive Southern hybridization. The PCR detection is done by use of two oligonucleotide primers flanking the polymorphic segment of repetitive DNA. Repeated cycles of heat denaturation of the DNA followed by annealing of the primers to their complementary sequences at low temperatures, and extension of the annealed primers with DNA polymerase, comprise the major part of the methodology. For example, see Batley, J., et al. 2007. *Mol. Ecol. Notes* (Online Early Articles) and Plieske, J., et al., 2001. *Theor. Appl. Genet.* 102:689-694.

Following amplification, markers can be scored by electrophoresis of the amplification products. Scoring of marker genotype is based on the size of the amplified fragment, which may be measured by the number of base pairs of the fragment. While variation in the primer used or in laboratory procedures can affect the reported fragment size, relative values should remain constant regardless of the specific primer or laboratory used. When comparing varieties it is preferable if all SSR profiles are performed in the same lab.

The SSR profile of canola plant SCV470336 can be used to identify plants comprising SCV470336 as a parent, since such plants will comprise the same homozygous alleles as SCV470336. Because the canola variety is essentially homozygous at all relevant loci, most loci should have only one type of allele present. In contrast, a genetic marker profile of an F_1 progeny should be the sum of those parents, e.g., if one parent was homozygous for allele x at a particular locus, and the other parent homozygous for allele y at that locus, then the F_1 progeny will be xy (heterozygous) at that locus. Subsequent generations of progeny produced by selection and breeding are expected to be of genotype x (homozygous), y (homozygous), or xy (heterozygous) for that locus position. When the F_1 plant is selfed or sibbed for successive filial generations, the locus should be either x or y for that position.

In addition, plants and plant parts substantially benefiting from the use of SCV470336 in their development, such as SCV470336 comprising a backcross conversion, transgene, or genetic sterility factor, may be identified by having a molecular marker profile with a high percent identity to

SCV470336. Such a percent identity might be 95%, 96%, 97%, 98%, 99%, 99.5% or 99.9% identical to SCV470336.

The SSR profile of SCV470336 also can be used to identify essentially derived varieties and other progeny varieties developed from the use of SCV470336, as well as cells and other plant parts thereof. Such plants may be developed using the markers identified in WO 00/31964, U.S. Pat. No. 6,162,967 and U.S. Pat. No. 7,288,386, each of which is incorporated herein by reference in its entirety. Progeny plants and plant parts produced using SCV470336 may be identified by having a molecular marker profile of at least 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 99.5% genetic contribution from canola variety, as measured by either percent identity or percent similarity. Such progeny may be further characterized as being within a pedigree distance of SCV470336, such as within 1, 2, 3, 4, or 5 or less cross-pollinations to a canola plant other than SCV470336 or a plant that has SCV470336 as a progenitor. Unique molecular profiles may be identified with other molecular tools such as SNPs and RFLPs.

While determining the SSR genetic marker profile of the plants described supra, several unique SSR profiles may also be identified which did not appear in either parent of such plant. Such unique SSR profiles may arise during the breeding process from recombination or mutation. A combination of several unique alleles provides a means of identifying a plant variety, an F_1 progeny produced from such variety, and progeny produced from such variety.

Single-Gene Conversions

When the term "canola plant" is used in the context of the present invention, this also includes any single gene conversions of that variety. The term "single gene converted plant," as used herein, refers to those canola plants which are developed by a plant breeding technique called backcrossing wherein essentially all of the desired morphological and physiological characteristics of a variety are recovered in addition to the single gene transferred into the variety via the backcrossing technique. Backcrossing methods can be used with the present invention to improve or introduce a characteristic into the variety. The term "backcrossing" as used herein refers to the repeated crossing of a hybrid progeny back to the recurrent parent, i.e., backcrossing 1, 2, 3, 4, 5, 6, 7, 8 or more times to the recurrent parent. The parental canola plant that contributes the gene for the desired characteristic is termed the nonrecurrent or donor parent. This terminology refers to the fact that the nonrecurrent parent is used one time in the backcross protocol and therefore does not recur. The parental canola plant to which the gene or genes from the nonrecurrent parent are transferred is known as the recurrent parent as it is used for several rounds in the backcrossing protocol (Poehlman & Sleper, 1994; Fehr, *Principles of Cultivar Development* (1987)). In a typical backcross protocol, the original variety of interest (recurrent parent) is crossed to a second variety (nonrecurrent parent) that carries the single gene of interest to be transferred. The resulting progeny from this cross are then crossed again to the recurrent parent and the process is repeated until a canola plant is obtained wherein essentially all of the desired morphological and physiological characteristics of the recurrent parent are recovered in the converted plant, in addition to the single transferred gene from the nonrecurrent parent.

The selection of a suitable recurrent parent is an important step for a successful backcrossing procedure. The goal of a

backcross protocol is to alter or substitute a single trait or characteristic in the original variety. To accomplish this, a single gene of the recurrent variety is modified or substituted with the desired gene from the nonrecurrent parent, while retaining essentially all of the rest of the desired genetic, and therefore the desired physiological and morphological, constitution of the original variety. The choice of the particular nonrecurrent parent will depend on the purpose of the backcross; one of the major purposes is to add some agronomically important trait to the plant. The exact backcrossing protocol will depend on the characteristic or trait being altered to determine an appropriate testing protocol. Although backcrossing methods are simplified when the characteristic being transferred is a dominant allele, a recessive allele may also be transferred. In this instance it may be necessary to introduce a test of the progeny to determine if the desired characteristic has been successfully transferred.

Many single gene traits have been identified that are not regularly selected for in the development of a new variety but that can be improved by backcrossing techniques. Single gene traits may or may not be transgenic; examples of these traits include but are not limited to, male sterility, waxy starch, herbicide tolerance, resistance for bacterial, fungal, or viral disease, insect resistance, male fertility, enhanced nutritional quality, industrial usage, yield stability and yield enhancement. These genes are generally inherited through the nucleus. Several of these single gene traits are described in U.S. Pat. Nos. 5,959,185; 5,973,234 and 5,977,445, each of which is incorporated herein by reference in its entirety.

Introduction of a New Trait or Locus into SCV470336

Line SCV470336 represents a new base genetic variety into which a new locus or trait may be introgressed. Direct transformation and backcrossing represent two important methods that can be used to accomplish such an introgression. The term backcross conversion and single locus conversion are used interchangeably to designate the product of a backcrossing program.

Backcross Conversions of SCV470336

A backcross conversion of SCV470336 occurs when DNA sequences are introduced through backcrossing with SCV470336 utilized as the recurrent parent. Both naturally occurring and transgenic DNA sequences may be introduced through backcrossing techniques. A backcross conversion may produce a plant with a trait or locus conversion in at least two or more backcrosses, including at least 2 crosses, at least 3 crosses, at least 4 crosses, or at least 5 crosses. Molecular marker assisted breeding or selection may be utilized to reduce the number of backcrosses necessary to achieve the backcross conversion. For example, see Openshaw, S. J., et al., Marker-assisted Selection in Backcross Breeding. In: *Proceedings Symposium of the Analysis of Molecular Data*, August 1994, Crop Science Society of America, Corvallis, Oreg., where it is demonstrated that a backcross conversion can be made in as few as two backcrosses.

The complexity of the backcross conversion method depends on the type of trait being transferred (single genes or closely linked genes as vs. unlinked genes), the level of expression of the trait, the type of inheritance (cytoplasmic or nuclear) and the types of parents included in the cross. It is understood by those of ordinary skill in the art that for single gene traits that are relatively easy to classify, the backcross method is effective and relatively easy to manage (see

Hanauer, et al. in *Corn and Corn Improvement*, Sprague and Dudley, Third Ed. 1998). Desired traits that may be transferred through backcross conversion include, but are not limited to, sterility (nuclear and cytoplasmic), fertility restoration, nutritional enhancements, drought tolerance, nitrogen utilization, altered fatty acid profile, altered seed amino acid levels, altered seed oil levels, low phytate, industrial enhancements, disease resistance (bacterial, fungal or viral), insect resistance and herbicide tolerance. In addition, an introgression site itself, such as an FRT site, Lox site or other site specific integration site, may be inserted by backcrossing and utilized for direct insertion of one or more genes of interest into a specific plant variety. In some embodiments of the invention, the number of loci that may be backcrossed into SCV470336 is at least 1, 2, 3, 4, or 5 and/or no more than 6, 5, 4, 3, or 2. A single locus may contain several transgenes, such as a transgene for disease resistance that, in the same expression vector, also contains a transgene for herbicide tolerance. The gene for herbicide tolerance may be used as a selectable marker and/or as a phenotypic trait. A single locus conversion of site specific integration system allows for the integration of multiple genes at the converted loci.

The backcross conversion may result from either the transfer of a dominant allele or a recessive allele. Selection of progeny containing the trait of interest is accomplished by direct selection for a trait associated with a dominant allele. Transgenes transferred via backcrossing typically function as a dominant single gene trait and are relatively easy to classify. Selection of progeny for a trait that is transferred via a recessive allele requires growing and selfing the first backcross generation to determine which plants carry the recessive alleles. Recessive traits may require additional progeny testing in successive backcross generations to determine the presence of the locus of interest. The last backcross generation is usually selfed to give pure breeding progeny for the gene(s) being transferred, although a backcross conversion with a stably introgressed trait may also be maintained by further backcrossing to the recurrent parent with selection for the converted trait.

Along with selection for the trait of interest, progeny are selected for the phenotype of the recurrent parent. The backcross is a form of inbreeding, and the features of the recurrent parent are automatically recovered after successive backcrosses. Poehlman, *Breeding Field Crops*, P. 204 (1987). Poehlman suggests from one to four or more backcrosses, but as noted above, the number of backcrosses necessary can be reduced with the use of molecular markers. Other factors, such as a genetically similar donor parent, may also reduce the number of backcrosses necessary. As noted by Poehlman, backcrossing is easiest for simply inherited, dominant and easily recognized traits.

One process for adding or modifying a trait or locus in canola line SCV470336 comprises crossing SCV470336 plants grown from SCV470336 seed with plants of another canola variety that comprise the desired trait or locus, selecting F₁ progeny plants that comprise the desired trait or locus to produce selected F₁ progeny plants, crossing the selected progeny plants with the SCV470336 plants to produce backcross progeny plants, selecting for backcross progeny plants that have the desired trait or locus and the morphological characteristics of canola line SCV470336 to produce selected backcross progeny plants; and backcrossing to SCV470336 three or more times in succession to produce selected fourth or higher backcross progeny plants that comprise said trait or locus. The modified SCV470336 may be further characterized as having essentially all of the physiological and morphological characteristics of canola line SCV470336 listed in

Table 1 and/or may be characterized by percent similarity or identity to SCV470336 as determined by SSR markers. The above method may be utilized with fewer backcrosses in appropriate situations, such as when the donor parent is highly related or markers are used in the selection step. Desired traits that may be used include those nucleic acids known in the art, some of which are listed herein, that will affect traits through nucleic acid expression or inhibition. Desired loci include the introgression of FRT, Lox and other sites for site specific integration, which may also affect a desired trait if a functional nucleic acid is inserted at the integration site.

In addition, the above process and other similar processes described herein may be used to produce first generation progeny canola seed by adding a step at the end of the process that comprises crossing SCV470336 with the introgressed trait or locus with a different canola plant and harvesting the resultant first generation progeny canola seed.

Tissue Culture of Canola

Further production of the SCV470336 line can occur by tissue culture and regeneration. Tissue culture of various tissues of canola and regeneration of plants therefrom is known and widely published. For example, reference may be had to Chuong, et al., "A Simple Culture Method for *Brassica* Hypocotyl Protoplasts", *Plant Cell Reports* 4:4-6 (1985); Barsby, T. L., et al., "A Rapid and Efficient Alternative Procedure for the Regeneration of Plants from Hypocotyl Protoplasts of *Brassica napus*", *Plant Cell Reports*, (Spring, 1996); Kartha, K., et al., "In vitro Plant Formation from Stem Explants of Rape", *Physiol. Plant*, 31:217-220 (1974); Narasimhulu, S., et al., "Species Specific Shoot Regeneration Response of Cotyledonary Explants of Brassicas", *Plant Cell Reports*, (Spring 1988); Swanson, E., "Microspore Culture in *Brassica*", *Methods in Molecular Biology*, Vol. 6, Chapter 17, p. 159 (1990). Thus, another aspect of this invention is to provide cells which upon growth and differentiation produce canola plants having the physiological and morphological characteristics of canola line SCV470336.

As used herein, the term "tissue culture" indicates a composition comprising isolated cells of the same or a different type or a collection of such cells organized into parts of a plant. Exemplary types of tissue cultures are protoplasts, calli, plant clumps, and plant cells that can generate tissue culture that are intact in plants or parts of plants, such as embryos, pollen, flowers, seeds, pods, leaves, stems, roots, root tips, anthers, and pistils. Means for preparing and maintaining plant tissue culture are well-known in the art. By way of example, a tissue culture comprising organs has been used to produce regenerated plants. U.S. Pat. Nos. 5,959,185, 5,973,234 and 5,977,445, describe certain techniques, each of which is incorporated herein by reference in its entirety.

Using SCV470336 to Develop Other Canola Varieties

Canola varieties such as SCV470336 are typically developed for use in seed and grain production. However, canola varieties such as SCV470336 also provide a source of breeding material that may be used to develop new canola varieties. Plant breeding techniques known in the art and used in a canola plant breeding program include, but are not limited to, recurrent selection, mass selection, bulk selection, mass selection, backcrossing, pedigree breeding, open pollination breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, mak-

ing double haploids, and transformation. Often combinations of these techniques are used. The development of canola varieties in a plant breeding program requires, in general, the development and evaluation of homozygous varieties. For a general description of rapeseed and canola breeding, see Downey, et al., 1987. Rapeseed and Mustard in Fehr, W. R. (ed.), *Principles of Cultivar Development* p. 437-486. N.Y., Macmillan and Co.; Thompson, K. F., 1983. Breeding winter oilseed rape *Brassica napus*. *Adv. Appl. Bio.* 7:1-104; and Ward, et al., 1985. *Oilseed Rape*. Farming Press Ltd., Wharfedale Road, Ipswich, Suffolk.

Additional Breeding Methods

This invention, in an embodiment, is directed to methods for producing a canola plant by crossing a first parent canola plant with a second parent canola plant wherein either the first or second parent canola plant is line SCV470336. The other parent may be any other canola plant, such as a canola plant that is part of a synthetic or natural population. Any such methods using canola line SCV470336 are part of this invention: selfing, sibbing, backcrosses, mass selection, pedigree breeding, bulk selection, hybrid production, or crosses to populations. These methods are well-known in the art and some of the more commonly used breeding methods are described below. Descriptions of breeding methods can be found in one of several reference books (e.g., Allard, *Principles of Plant Breeding*, 1960; Simmonds, *Principles of Crop Improvement*, 1979; Sneep, et al., 1979).

The following describes breeding methods that may be used with canola line SCV470336 in the development of further canola plants. One such embodiment is a method for developing a line SCV470336 progeny canola plant in a canola plant breeding program comprising: obtaining the canola plant, or a part thereof, of line SCV470336 utilizing said plant or plant part as a source of breeding material and selecting a canola line SCV470336 progeny plant with molecular markers in common with line SCV470336 and/or with morphological and/or physiological characteristics selected from the characteristics listed in Table 1. Breeding steps that may be used in the canola plant breeding program include pedigree breeding, backcrossing, mutation breeding, and recurrent selection. In conjunction with these steps, techniques such as RFLP-enhanced selection, genetic marker enhanced selection (for example SSR markers) and the making of double haploids may be utilized.

Another method involves producing a population of canola line SCV470336 progeny canola plants, comprising crossing line SCV470336 with another canola plant, thereby producing a population of canola plants, which, on average, derive 50% of their alleles from canola line SCV470336. A plant of this population may be selected and repeatedly selfed or sibbed with a canola line resulting from these successive filial generations. One embodiment of this invention is the canola line produced by this method and that has obtained at least 50% of its alleles from canola line SCV470336.

One of ordinary skill in the art of plant breeding would know how to evaluate the traits of two plant varieties to determine if there is no significant difference between the two traits expressed by those varieties. For example, see Fehr and Walt, *Principles of Cultivar Development*, p 261-286 (1987). Thus, the invention includes, in an embodiment, canola line SCV470336 progeny canola plants comprising a combination of at least two line SCV470336 traits selected from the group consisting of those listed in Table 1. In another embodiment, the line SCV470336 comprises any combination of traits described herein. In either embodiment, the progeny

canola plant may not be significantly different for said traits than canola line SCV470336 as determined at the 5% significance level when grown in the same environmental conditions. Using techniques described herein, molecular markers may be used to identify said progeny plant as a canola line SCV470336 progeny plant. Mean trait values may be used to determine whether trait differences are significant, and the traits may then be measured on plants grown under the same environmental conditions. Once such a variety is developed its value is substantial since it is important to advance the germplasm base as a whole in order to maintain or improve traits such as yield, disease resistance, pest resistance, and plant performance in extreme environmental conditions.

Progeny of canola line SCV470336 may also be characterized through their filial relationship with canola line SCV470336, as for example, being within a certain number of breeding crosses of canola line SCV470336. A breeding cross is a cross made to introduce new genetics into the progeny, and is distinguished from a cross, such as a self or a sib cross, made to select among existing genetic alleles. The lower the number of breeding crosses in the pedigree, the closer the relationship between canola line SCV470336 and its progeny. For example, progeny produced by the methods described herein may be within 1, 2, 3, 4 or 5 breeding crosses of canola line SCV470336.

Pedigree Breeding

Pedigree breeding starts with the crossing of two genotypes, such as SCV470336 and another canola variety having one or more desirable characteristics that is lacking or which complements SCV470336. If the two original parents do not provide all the desired characteristics, other sources can be included in the breeding population. In the pedigree method, superior plants are selfed and selected in successive filial generations. In the succeeding filial generations the heterozygous condition gives way to homogeneous varieties as a result of self-pollination and selection. Typically in the pedigree method of breeding, five or more successive filial generations of selfing and selection is practiced: F_1 to F_2 ; F_2 to F_3 ; F_3 to F_4 ; F_4 to F_5 , etc. After a sufficient amount of inbreeding, successive filial generations will serve to increase seed of the developed variety. The developed variety may comprise homozygous alleles at about 95% or more of its loci.

In addition to being used to create a backcross conversion, backcrossing can also be used in combination with pedigree breeding. As discussed previously, backcrossing can be used to transfer one or more specifically desirable traits from one variety, the donor parent, to a developed variety called the recurrent parent, which has overall good agronomic characteristics yet lacks that desirable trait or traits. However, the same procedure can be used to move the progeny toward the genotype of the recurrent parent but at the same time retain many components of the non-recurrent parent by stopping the backcrossing at an early stage and proceeding with selfing and selection. For example, a canola variety may be crossed with another variety to produce a first generation progeny plant. The first generation progeny plant may then be backcrossed to one of its parent varieties to create a BC1 or BC2. Progeny are selfed and selected so that the newly developed variety has many of the attributes of the recurrent parent and yet several of the desired attributes of the non-recurrent parent. This approach leverages the value and strengths of the recurrent parent for use in new canola varieties.

Therefore, an embodiment of this invention is a method of making a backcross conversion of canola line SCV470336, comprising the steps of crossing a plant of canola line

SCV470336 with a donor plant comprising a desired trait, selecting an F_1 progeny plant comprising the desired trait, and backcrossing the selected F_1 progeny plant to a plant of canola line SCV470336. This method may further comprise the step of obtaining a molecular marker profile of canola line SCV470336 and using the molecular marker profile to select for a progeny plant with the desired trait and the molecular marker profile of SCV470336. In one embodiment, the desired trait is a mutant gene or transgene present in the donor parent.

Recurrent Selection and Mass Selection

Recurrent selection is a method used in a plant breeding program to improve a population of plants. SCV470336 is suitable for use in a recurrent selection program. The method entails individual plants cross pollinating with each other to form progeny. The progeny are grown and the superior progeny selected by any number of selection methods, which include individual plant, half-sib progeny, full-sib progeny and selfed progeny. The selected progeny are cross pollinated with each other to form progeny for another population. This population is planted and again superior plants are selected to cross pollinate with each other. Recurrent selection is a cyclical process and therefore can be repeated as many times as desired. The objective of recurrent selection is to improve the traits of a population. The improved population can then be used as a source of breeding material to obtain new varieties for commercial or breeding use, including the production of a synthetic line. A synthetic line is the resultant progeny formed by the intercrossing of several selected varieties.

Mass selection is a useful technique when used in conjunction with molecular marker enhanced selection. In mass selection seeds from individuals are selected based on phenotype or genotype. These selected seeds are then bulked and used to grow the next generation. Bulk selection requires growing a population of plants in a bulk plot, allowing the plants to self-pollinate, harvesting the seed in bulk and then using a sample of the seed harvested in bulk to plant the next generation. Also, instead of self pollination, directed pollination could be used as part of the breeding program.

Mutation Breeding

Mutation breeding is another method of introducing new traits into canola line SCV470336. Mutations that occur spontaneously or are artificially induced can be useful sources of variability for a plant breeder. The goal of artificial mutagenesis is to increase the rate of mutation for a desired characteristic. Mutation rates can be increased by many different means including temperature, long-term seed storage, tissue culture conditions, radiation; such as X-rays, Gamma rays (e.g. cobalt 60 or cesium 137), neutrons, (product of nuclear fission by uranium 235 in an atomic reactor), Beta radiation (emitted from radioisotopes such as phosphorus 32 or carbon 14), or ultraviolet radiation (from 2500 to 2900 nm), or chemical mutagens (such as base analogues (5-bromo-uracil), related compounds (8-ethoxy caffeine), antibiotics (streptonigrin), alkylating agents (sulfur mustard, nitrogen mustards, epoxides, ethylenamines, sulfates, sulfonates, sulfones, lactones), azide, hydroxylamine, nitrous acid, or acridines. Once a desired trait is observed through mutagenesis the trait may then be incorporated into existing germplasm by traditional breeding techniques. Details of mutation breeding can be found in Fehr, 1993. *Principles of Cultivar Development*, Macmillan Publishing Company. In addition, mutations created in other canola plants may be

used to produce a backcross conversion of canola line SCV470336 that comprises such mutation.

Breeding with Molecular Markers

Molecular markers, which include markers identified through the use of techniques such as Isozyme Electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed Polymerase Chain Reaction (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions (SCARS), Amplified Fragment Length Polymorphisms (AFLPs), Simple Sequence Repeats (SSRs) and Single Nucleotide Polymorphisms (SNPs), may be used in plant breeding methods utilizing canola line SCV470336. One use of molecular markers is QTL mapping. QTL mapping is the use of markers, which are known to be closely linked to alleles that have measurable effects on a quantitative trait. Selection in the breeding process is based upon the accumulation of markers linked to the positive effecting alleles and/or the elimination of the markers linked to the negative effecting alleles from the plant's genome.

Molecular markers can also be used during the breeding process for the selection of qualitative traits. For example, markers closely linked to alleles or markers containing sequences within the actual alleles of interest can be used to select plants that contain the alleles of interest during a backcrossing breeding program. The markers can also be used to select for the genome of the recurrent parent and against the genome of the donor parent. Using this procedure can minimize the amount of genome from the donor parent that remains in the selected plants. It can also be used to reduce the number of crosses back to the recurrent parent needed in a backcrossing program. The use of molecular markers in the selection process is often called genetic marker enhanced selection. Molecular markers may also be used to identify and exclude certain sources of germplasm as parental varieties or ancestors of a plant by providing a means of tracking genetic profiles through crosses.

Production of Double Haploids

The production of double haploids can also be used for the development of plants with a homozygous phenotype in the breeding program. For example, a canola plant for which canola line SCV470336 is a parent can be used to produce double haploid plants. Double haploids are produced by the doubling of a set of chromosomes (1 N) from a heterozygous plant to produce a completely homozygous individual. For example, see Wan, et al., (1989) "Efficient Production of Doubled Haploid Plants Through Colchicine Treatment of Anther-Derived Maize Callus", *Theor. Appl. Genet.*, 77:889-892 and U.S. Pat. No. 7,135,615, incorporated herein by reference in its entirety. This can be advantageous because the process omits the generations of selfing needed to obtain a homozygous plant from a heterozygous source.

Haploid induction systems have been developed for various plants to produce haploid tissues, plants and seeds. Methods for obtaining haploid plants are also disclosed in Kobayashi, M., et al., *J. Heredity* 71(1):9-14, 1980; Pollacsek, M., *Agronomie (Paris)* 12(3):247-251, 1992; Cho-Un-Haing, et al., *J. Plant Biol.*, 1996, 39 (3):185-188; Verdoodt, L., et al., February 1998, 96(2):294-300; Genetic Manipulation in Plant Breeding, Proceedings International Symposium Organized by EUCARPIA, Sep. 8-13, 1985, Berlin, Germany; Chalyk, et al., 1994, *Maize Genet. Coop. Newsletter* 68:47; Chalyk, S.

Thus, an embodiment of this invention is a process for making a substantially homozygous SCV470336 progeny plant by producing or obtaining a seed from the cross of SCV470336 and another canola plant and applying double haploid methods to the F₁ seed or F₁ plant or to any successive filial generation. Based on studies in maize and currently being conducted in canola, such methods would decrease the number of generations required to produce a variety with similar genetics or characteristics to SCV470336. See Bernardo, R. and Kahler, A. L., *Theor. Appl. Genet.* 102:986-992, 2001.

In particular, a process of making seed retaining the molecular marker profile of canola line SCV470336 is contemplated, such process comprising obtaining or producing F₁ seed for which canola line SCV470336 is a parent, inducing doubled haploids to create progeny without the occurrence of meiotic segregation, obtaining the molecular marker profile of canola line SCV470336, and selecting progeny that retain the molecular marker profile of SCV470336.

A pollination control system and effective transfer of pollen from one parent to the other offers improved plant breeding and an effective method for producing hybrid canola seed and plants. For example, the ogura cytoplasmic male sterility (cms) system, developed via protoplast fusion between radish (*Raphanus sativus*) and rapeseed (*Brassica napus*) is one of the most frequently used methods of hybrid production. It provides stable expression of the male sterility trait (Ogura 1968), Pelletier, et al. (1983) and an effective nuclear restorer gene (Heyn 1976).

In developing improved new *Brassica* hybrid varieties, breeders use self-incompatible (SI), cytoplasmic male sterile (CMS) and nuclear male sterile (NMS) *Brassica* plants as the female parent. In using these plants, breeders are attempting to improve the efficiency of seed production and the quality of the F₁ hybrids and to reduce the breeding costs. When hybridization is conducted without using SI, CMS or NMS plants, it is more difficult to obtain and isolate the desired traits in the progeny (F₁ generation) because the parents are capable of undergoing both cross-pollination and self-pollination. If one of the parents is a SI, CMS or NMS plant that is incapable of producing pollen, only cross pollination will occur. By eliminating the pollen of one parental variety in a cross, a plant breeder is assured of obtaining hybrid seed of uniform quality, provided that the parents are of uniform quality and the breeder conducts a single cross.

In one instance, production of F₁ hybrids includes crossing a CMS *Brassica* female parent, with a pollen producing male *Brassica* parent. To reproduce effectively, however, the male parent of the F₁ hybrid must have a fertility restorer gene (Rf gene). The presence of an Rf gene means that the F₁ generation will not be completely or partially sterile, so that either self-pollination or cross pollination may occur. Self pollination of the F₁ generation to produce several subsequent generations is important to ensure that a desired trait is heritable and stable and that a new variety has been isolated.

An example of a *Brassica* plant which is cytoplasmic male sterile and used for breeding is ogura (OGU) cytoplasmic male sterile (R. Pellan-Delourme, et al., 1987). A fertility restorer for ogura cytoplasmic male sterile plants has been transferred from *Raphanus sativus* (radish) to *Brassica* by Institut. National de Recherche Agricole (INRA) in Rennes, France (Pelletier, et al., 1987). The restorer gene, Rf1 originating from radish, is described in WO 92/05251 and in Delourme, et al., (1991). Improved versions of this restorer have been developed. For example, see WO 98/27806 "Oilseed *brassica* containing an improved fertility restorer gene for ogura cytoplasmic male sterility".

Other sources and refinements of CMS sterility in canola include the Polima cytoplasmic male sterile plant, as well as those of U.S. Pat. No. 5,789,566, "DNA sequence imparting cytoplasmic male sterility, mitochondrial genome, nuclear genome, mitochondria and plant containing said sequence and process for the preparation of hybrids"; U.S. Pat. No. 5,973,233 "Cytoplasmic male sterility system production canola hybrids"; and WO 97/02737 "Cytoplasmic male sterility system producing canola hybrids"; EP patent application 0 599042A "Methods for introducing a fertility restorer gene and for producing F₁ hybrids of *Brassica* plants thereby"; U.S. Pat. No. 6,229,072 "Cytoplasmic male sterility system production canola hybrids"; U.S. Pat. No. 4,658,085 "Hybridization using cytoplasmic male sterility, cytoplasmic herbicide tolerance, and herbicide tolerance from nuclear genes"; each of which is incorporated herein by reference in its entirety.

Further, as a result of the advances in sterility systems, lines are developed that can be used as an open pollinated variety (i.e. a pureline line sold to the grower for planting) and/or as a sterile inbred (female) used in the production of F₁ hybrid seed. In the latter case, favorable combining ability with a restorer (male) would be desirable. The resulting hybrid seed would then be sold to the grower for planting.

The development of a canola hybrid in a canola plant breeding program involves three steps: (1) the selection of plants from various germplasm pools for initial breeding crosses; (2) the selfing of the selected plants from the breeding crosses for several generations to produce a series of inbred lines, which, although different from each other, breed true and are highly uniform; and (3) crossing the selected inbred lines with different inbred lines to produce the hybrids. During the inbreeding process in canola, the vigor of the lines decreases. Vigor is restored when two different inbred lines are crossed to produce the hybrid. An important consequence of the homozygosity and homogeneity of the inbred lines is that the hybrid between a defined pair of inbreds will always be the same. Once the inbreds that give a superior hybrid have been identified, the hybrid seed can be reproduced indefinitely as long as the homogeneity of the inbred parents is maintained.

Combining ability of a line, as well as the performance of the line, is a factor in the selection of improved canola lines that may be used as inbreds. Combining ability refers to a line's contribution as a parent when crossed with other lines to form hybrids. The hybrids formed for the purpose of selecting superior lines are designated test crosses. One way of measuring combining ability is by using breeding values. Breeding values are based on the overall mean of a number of test crosses. This mean is then adjusted to remove environmental effects and it is adjusted for known genetic relationships among the lines.

Hybrid seed production requires inactivation of pollen produced by the female parent. Incomplete inactivation of the pollen provides the potential for self-pollination. This inadvertently self-pollinated seed may be unintentionally harvested and packaged with hybrid seed. Similarly, because the male parent is grown next to the female parent in the field there is also the potential that the male selfed seed could be unintentionally harvested and packaged with the hybrid seed. Once the seed from the hybrid bag is planted, it is possible to identify and select these self-pollinated plants. These self-pollinated plants will be genetically equivalent to one of the inbred lines used to produce the hybrid. Though the possibility of inbreds being included in hybrid seed bags exists, the occurrence is rare because much care is taken to avoid such inclusions. These self-pollinated plants can be identified and

selected by one skilled in the art, either through visual or molecular methods. *Brassica napus* canola plants, absent the use of sterility systems, are recognized to commonly be self-fertile with approximately 70 to 90 percent of the seed normally forming as the result of self-pollination. The percentage of cross pollination may be further enhanced when populations of recognized insect pollinators at a given growing site are greater. Thus open pollination is often used in commercial canola production.

Descriptions of other breeding methods that are commonly used for different traits and crops can be found in one of several reference books (e.g., Allard, 1960; Simmonds, 1979; Sneep, et al., 1979; Fehr, 1987).

INDUSTRIAL USES

Currently *Brassica napus* canola is recognized as an increasingly important oilseed crop and a source of meal in many parts of the world. The oil as removed from the seeds commonly contains a lesser concentration of endogenously formed saturated fatty acids than other vegetable oils and is well suited for use in the production of salad oil or other food products or in cooking or frying applications. The oil also finds utility in industrial applications. Additionally, the meal component of the seeds can be used as a nutritious protein concentrate for livestock.

Canola oil has the lowest level of saturated fatty acids of all vegetable oils. "Canola" refers to rapeseed (*Brassica*) which has a erucic acid (C_{22:1}) content of at most 2 percent by weight based on the total fatty acid content of a seed, and which produces, after crushing, an air-dried meal containing less than 30 micromoles (μmol) per gram of defatted (oil-free) meal. These types of rapeseed are distinguished by their edibility in comparison to more traditional varieties of the species.

Canola line SCV470336 can be used in the production of an edible vegetable oil or other food products in accordance with known techniques. The solid meal component derived from seeds can be used as a nutritious livestock feed. Parts of the plant not used for human or animal food can be used for biofuel.

TABLES

In Table 2, selected oil quality characteristics of the seed of canola line SCV470336 are compared with oil quality characteristics of the proprietary canola lines referenced in Table 1, SCV378221 and SCV204738. The data in Table 2 includes results on seed samples collected from 20 testing locations and are presented as averages of the values observed. Column 1 shows the variety, column 2 shows the percent saturated fatty acid content, column 3 shows the percent oleic acid content, column 4 shows the percent linoleic acid content and column 5 shows the percent linolenic acid content for each variety.

Compared to the two canola lines SCV378221 and SCV204738, the averages presented in Table 2 indicate that seed of canola line SCV470336 of the present invention has a percent saturated fatty acid content within a normal range, an average percent oleic acid content, an average percent linoleic acid content, and an average percent linolenic acid content that is between that of SCV378221 and SCV204738.

TABLE 2

Oil Quality Characteristics of SCV470336 Compared to Two Proprietary Canola Lines				
1 Variety	2 Sat. Fatty Acids %	3 Oleic Acid %	4 Linoleic Acid %	5 Linolenic Acid %
SCV470336	7.84%	66.63%	17.46%	6.34%
SCV378221	7.61%	64.17%	19.28%	7.09%
SCV204738	7.49%	66.93%	17.41%	6.43%

In Table 3, selected characteristics of a single cross hybrid G88124 containing canola line SCV470336 are compared with characteristics of two commercial canola check varieties. Trial data values are shown for G88124 and the average of the two commercial canola lines, Q2 and 46A45, with the values shown being representative of data collected from a specified number of trial locations ("No. Locs"). Column 1 shows the variety, column 2 shows the yield as a percent of the average of the check varieties, column 3 shows the plant lodging ratings, column 4 shows the days to maturity, column 5 shows the percent saturated fatty acid content, column 6 shows the plant height, column 7 shows the percent glucosinolate content in micromoles, column 8 shows the percent oil content, column 9 shows the percent protein content, column 10 shows the resistance rating to blackleg disease, and column 11 shows the resistance rating to *Fusarium* wilt disease.

Compared to the average of the values recorded for Check Q2 and Check 46A45, the hybrid (G88124) containing SCV470336 of the present invention has higher yield, a lodging rating that is comparable, a days to maturity rating that is slightly earlier, a comparable percent saturated fat content, a comparable plant height, a lower glucosinolate content, a higher percent oil content, and a comparable percent protein content, with resistance to blackleg and *Fusarium* wilt diseases.

TABLE 3

Characteristics of a Hybrid Containing SCV470336 Compared to Two Commercial Varieties*										
1 Var.	2 Yield %	3 Lodging rating	4 DMat Days	5 Sats %	6 Height cm	7 Gluc µm/g	8 Oil %	9 Prot %	10 BL rating	11 FW rating
Q2	98.8	1.2	106.6	7.02	105.2	12.87	46.65	45.21	MR	R
46A65	101.2	1.4	106.1	6.71	101.9	13.95	47.43	46.30	MR	R
AVG of CHECKS	100	1.3	106.3	6.86	103.5	13.41	47.04	45.75	MR	R
G88124	125.0	1.3	105.3	6.87	103.2	9.45	50.31	45.69	R	R
No. Loc	34	34	34	34	34	34	34	34	3	3

*Hybrid G88124 compared to commercial varieties Q2 and 46A65.

Note:

Q2 and 46A65 are used as check varieties in the official Canadian variety registration trials conducted by the Western Canada Canola/Rapeseed Recommending Committee, Inc. Data shown for each variety and characteristic are the mean values over all zones and years tested. "NA" means not available.

DEPOSIT INFORMATION

A deposit of the Monsanto Canada Inc. proprietary canola line designated SCV470336 disclosed above and recited in the appended claims has been made with the American Type Culture Collection (ATCC), 10801 University Boulevard, Manassas, Va. 20110. The date of deposit was Jan. 28, 2010. The deposit of 2,500 seeds was taken from the same deposit maintained by Monsanto Canada Inc. since prior to the filing date of this application. All restrictions upon the deposit have been removed, and the deposit is intended to meet all of the requirements of 37 C.F.R. 1.801-1.809. The ATCC accession

number is PTA-10628. The deposit will be maintained in the depository for a period of 30 years, or 5 years after the last request, or for the effective life of the patent, whichever is longer, and will be replaced as necessary during that period.

All references cited in this specification, including without limitation, all papers, publications, patents, patent applications, presentations, texts, reports, manuscripts, brochures, books, interne postings, journal articles, periodicals, and the like, are hereby incorporated by reference into this specification in their entireties. The discussion of the references herein is intended merely to summarize the assertions made by their authors and no admission is made that any reference constitutes prior art. Applicants reserve the right to challenge the accuracy and pertinence of the cited references.

Although embodiments of the invention have been described using specific terms, devices, and methods, such description is for illustrative purposes only. The words used are words of description rather than of limitation. It is to be understood that changes and variations may be made by those of ordinary skill in the art without departing from the spirit or the scope of the present invention, which is set forth in the following claims. In addition, it should be understood that aspects of the various embodiments may be interchanged either in whole or in part. Therefore, the spirit and scope of the appended claims should not be limited to the description of the versions contained therein.

What is claimed is:

1. A seed of canola line SCV470336, a representative sample of seed of which was deposited under ATCC Accession No. PTA-10628.
2. A canola plant, or a part thereof, produced by growing the seed of claim 1.
3. A tissue culture produced from protoplasts or cells from the plant of claim 2, wherein said cells or protoplasts of the tissue culture are produced from a plant part selected from the

group consisting of leaf, pollen, embryo, cotyledon, hypocotyl, meristematic cell, root, root tip, anther, pistil, flower, shoot, stem, petiole and pod.

4. A canola plant regenerated from the tissue culture of claim 3, wherein the plant has essentially all of the morphological and physiological characteristics of line SCV470336, as shown in Table 1.

5. A method for producing a canola seed comprising crossing two canola plants and harvesting the resultant canola seed, wherein at least one of the two canola plants is the canola plant of claim 2.

6. A canola seed produced by the method of claim 5.

7. A canola plant, or a part thereof, produced by growing said seed of claim 6.

8. A method for producing a male sterile canola plant wherein the method comprises transforming the canola plant of claim 2 with a nucleic acid molecule that confers male sterility.

9. A male sterile canola plant produced by the method of claim 8.

10. A method of producing an herbicide tolerant canola plant wherein the method comprises transforming the canola plant of claim 2 with a transgene that confers herbicide tolerance to an herbicide selected from the group consisting of imidazolinone, sulfonylurea, glyphosate, glufosinate, 2,4-D, Dicamba, L-phosphinothricin, triazine, hydroxyphenylpyruvate dioxygenase inhibitor, protoporphyrinogen oxidase inhibitor, phenoxy proprionic acid, cyclohexone, and benzonitrile.

11. An herbicide tolerant canola plant produced by the method of claim 10.

12. A method of producing an insect- or pest-resistant canola plant wherein the method comprises transforming the canola plant of claim 2 with a transgene that confers insect or pest resistance.

13. An insect- or pest-resistant canola plant produced by the method of claim 12.

14. The canola plant of claim 13, wherein the transgene encodes a *Bacillus thuringiensis* endotoxin.

15. A method of producing a disease resistant canola plant wherein the method comprises transforming the canola plant of claim 2 with a transgene that confers disease resistance.

16. A disease resistant canola plant produced by the method of claim 15.

17. A method of producing a canola plant with modified fatty acid metabolism or modified carbohydrate metabolism wherein the method comprises transforming the canola plant of claim 2 with a transgene encoding a protein selected from the group consisting of fructosyltransferase, levansucrase, alpha-amylase, invertase, and starch branching enzyme or encoding an antisense of stearyl-ACP desaturase.

18. A canola plant having modified fatty acid metabolism or modified carbohydrate metabolism produced by the method of claim 17.

19. Meal prepared from the seed of the canola line of claim 1.

20. An unblended canola oil extracted from the seed of the canola line of claim 1.

21. A method of introducing a desired trait into canola line SCV470336 wherein the method comprises:

- (a) crossing a SCV470336 plant, wherein a representative sample of seed was deposited under ATCC Accession

No. PTA-10628, with a plant of another canola line that comprises a desired trait to produce progeny plants, wherein the desired trait is selected from the group consisting of male sterility, herbicide tolerance, insect resistance, pest resistance, modified fatty acid metabolism, modified carbohydrate metabolism, modified seed yield, modified oil percent, modified protein percent, modified lodging resistance, and resistance to bacterial disease, fungal disease, or viral disease;

(b) selecting one or more progeny plants that have the desired trait to produce selected progeny plants;

(c) crossing the selected progeny plants with the SCV470336 plants to produce backcross progeny plants;

(d) selecting for backcross progeny plants that have the desired trait and essentially all of the physiological and morphological characteristics of canola line SCV470336 listed in Table 1; and

(e) repeating the crossing the selected progeny step and selecting for backcross progeny step two or more times to produce selected third or higher backcross progeny plants that comprise the desired trait and essentially all of the physiological and morphological characteristics of canola line SCV470336 as shown in Table 1.

22. A canola plant produced by the method of claim 21, wherein the plant has the desired trait.

23. The canola plant of claim 22, wherein the desired trait is herbicide tolerance and the resistance is conferred to an herbicide selected from the group consisting of imidazolinone, sulfonylurea, glyphosate, glufosinate, 2,4-D, Dicamba, L-phosphinothricin, triazine, hydroxyphenylpyruvate dioxygenase inhibitor, protoporphyrinogen oxidase inhibitor, phenoxy proprionic acid, cyclohexone and benzonitrile.

24. The canola plant of claim 22, wherein the desired trait is insect resistance and the insect resistance is conferred by a transgene encoding a *Bacillus thuringiensis* endotoxin.

25. The canola plant of claim 22, wherein the desired trait is modified fatty acid metabolism or modified carbohydrate metabolism and said desired trait is conferred by a nucleic acid encoding a protein selected from the group consisting of phytase, fructosyltransferase, levansucrase, alpha-amylase, invertase and starch branching enzyme or encoding an antisense of stearyl-ACP desaturase.

26. A method of producing a male sterile canola seed wherein the method comprises crossing the canola plant of claim 2 with a male sterile canola plant and harvesting the resultant seed.

* * * * *

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,143,488 B2
APPLICATION NO. : 12/713594
DATED : March 27, 2012
INVENTOR(S) : Kevin T. Stone et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

Column 1,

Line 64, replace "FIG. 2A through 2C" with --FIGS. 2A through 2C--

Column 3,

Line 8, replace "layer 20A" with --layer 20a--

Column 6,

Line 40, after "end" delete "most"


Column 10,

Line 21, replace "lavers" with --layers--

Column 10,

Line 24, replace "laver" with --layer--

Signed and Sealed this
Twenty-ninth Day of May, 2012



David J. Kappos
Director of the United States Patent and Trademark Office

UNITED STATES PATENT AND TRADEMARK OFFICE
CERTIFICATE OF CORRECTION

PATENT NO. : 8,143,488 B2
APPLICATION NO. : 12/713594
DATED : March 27, 2012
INVENTOR(S) : Dale Burns et al.

Page 1 of 1

It is certified that error appears in the above-identified patent and that said Letters Patent is hereby corrected as shown below:

On the title page: Add omitted reference -- 2008/0256651 A1 10/2008 Chungu et al. -- to page 2 of the U.S. Patent Documents, right column, between listed reference 15 and reference 16 of the issued Patent.

Replace the numbers "12/713,637" with the numbers -- 12/721,637 -- on Page 7, of the OTHER PUBLICATIONS, right column, line 5 of the issued Patent.

Replace the term "Meichers" with the term -- Melchers -- in Column 16, line 20 of the issued Patent.

Replace the term "Hanauer" with the term -- Hallauer -- in Column 26, line 1 of the issued Patent.

Replace the term "interne" with the term -- internet -- in Column 36, line 8 of the issued Patent.

Signed and Sealed this
Twenty-eighth Day of August, 2012



David J. Kappos
Director of the United States Patent and Trademark Office