



US008158156B2

(12) **United States Patent**  
**Matthews et al.**

(10) **Patent No.:** **US 8,158,156 B2**  
(45) **Date of Patent:** **\*Apr. 17, 2012**

(54) **ABUSE-DETERRENT MULTI-LAYER  
PHARMACEUTICAL COMPOSITION  
COMPRISING AN OPIOID ANTAGONIST AND  
AN OPIOID AGONIST**

(75) Inventors: **Frank Matthews**, Edison, NJ (US);  
**Garth Boehm**, Westfield, NJ (US);  
**Lijuan Tang**, Flemington, NJ (US);  
**Alfred Liang**, Edison, NJ (US)

(73) Assignee: **Alpharma Pharmaceuticals, LLC**,  
Bridgewater, NJ (US)

(\*) Notice: Subject to any disclaimer, the term of this  
patent is extended or adjusted under 35  
U.S.C. 154(b) by 1164 days.

This patent is subject to a terminal dis-  
claimer.

(21) Appl. No.: **11/820,499**

(22) Filed: **Jun. 19, 2007**

(65) **Prior Publication Data**

US 2008/0233197 A1 Sep. 25, 2008

(51) **Int. Cl.**

**A61K 9/16** (2006.01)

**A61P 43/00** (2006.01)

(52) **U.S. Cl.** ..... **424/490**

(58) **Field of Classification Search** ..... None  
See application file for complete search history.

(56) **References Cited**

**U.S. PATENT DOCUMENTS**

2,770,569 A	11/1956	Fromherz et al.
2,945,049 A	7/1960	Chang et al.
2,981,641 A	4/1961	O'Neill
3,071,509 A	1/1963	O'Neill
3,332,950 A	7/1967	Blumberg et al.
3,493,657 A	2/1970	Lewenstein et al.
3,676,557 A	7/1972	Lachman et al.
3,689,574 A	9/1972	Engelhart
3,773,955 A	11/1973	Pachter et al.
3,845,770 A	11/1974	Theeuwes et al.
3,852,058 A	12/1974	Mezei et al.
3,879,555 A	4/1975	Pachter et al.
3,916,889 A	11/1975	Russell
3,965,256 A	6/1976	Leslie
3,966,940 A	6/1976	Pachter et al.
3,980,766 A	9/1976	Shaw et al.
3,993,073 A	11/1976	Zaffaroni
4,008,719 A	2/1977	Theeuwes et al.
4,036,228 A	7/1977	Theeuwes
4,063,064 A	12/1977	Saunders et al.
4,088,864 A	5/1978	Theeuwes et al.
4,111,201 A	9/1978	Theeuwes
4,111,202 A	9/1978	Theeuwes
4,176,186 A	11/1979	Goldberg
4,237,140 A	12/1980	Dudzinski
4,293,539 A	10/1981	Ludwig et al.
4,327,725 A	5/1982	Cortese et al.
4,366,310 A	12/1982	Leslie
4,401,672 A	8/1983	Portoghese et al.
4,443,428 A	4/1984	Oshlack et al.
4,451,470 A	5/1984	Ganti
4,457,933 A	7/1984	Gordon et al.

4,464,378 A	8/1984	Hussain et al.
4,519,801 A	5/1985	Edgren
4,573,995 A	3/1986	Chen et al.
4,576,604 A	3/1986	Guittard et al.
4,582,835 A	4/1986	Lewis et al.
4,588,580 A	5/1986	Gale et al.
4,597,961 A	7/1986	Etscorn
4,606,909 A	8/1986	Bechgaard
4,608,376 A	8/1986	Pasternak
4,612,008 A	9/1986	Wong et al.
4,626,539 A	12/1986	Aungst et al.
4,655,766 A	4/1987	Theeuwes et al.
4,661,492 A	4/1987	Lewis et al.
4,719,215 A	1/1988	Goldberg
4,730,048 A	3/1988	Portoghese et al.
4,760,069 A	7/1988	Rzeszotarski et al.
4,769,372 A	9/1988	Kreek et al.
4,783,456 A	11/1988	Glassman
4,785,000 A	11/1988	Kreek et al.
4,803,208 A	2/1989	Pasternak
4,806,341 A	2/1989	Chien et al.
4,806,543 A	2/1989	Choi
4,806,558 A	2/1989	Wuest et al.
4,828,836 A	5/1989	Elger et al.
4,834,965 A	5/1989	Martani et al.
4,834,984 A	5/1989	Goldie et al.
4,834,985 A	5/1989	Elger et al.

(Continued)

**FOREIGN PATENT DOCUMENTS**

AU 9211180 9/1992

(Continued)

**OTHER PUBLICATIONS**

Pakkanen, Jukka. S. , University of Helsinki. "Upregulation and  
Functionality of Neuronal Nicotinic Acetylcholine Receptors," Aca-  
demic Dissertation, 2006.

Bodmeier, Roland, et al. "The influence of Buffer Species and  
Strength of Diltiazem HCl Release from Beads Coated with the  
Aqueous Cationic Polymer Dispersions, Eudragit RS, RL 30 D,"  
Pharmaceutical Research, 13(1), 52-56, 1996.

Wagner, Karl G. et al "Influence of chloride ion exchange on the  
permeability and drug release of Eudragit RS 30 D films," Journal of  
Controlled Release, 82, (2002).

Felton, Linda A., et al. "Influence of Insoluble Excipients on Film  
Coating Systems," Drug Development and Industrial Pharmacy,  
28(3), 225-243 (2002).

Knop, K., et al. "Influence of Surfactants of Different Charge and  
Concentration on Drug Release from Pellets Coated with an Aqueous  
Dispersion of Quaternary Acrylic Polymers," S.T.P. Pharma Sci-  
ences, 7(6), 507-512 (1997).

(Continued)

*Primary Examiner* — Robert A Wax

*Assistant Examiner* — Danah Al-Awadi

(74) *Attorney, Agent, or Firm* — J. Michael Dixon; Gregg C.  
Benson

(57) **ABSTRACT**

Provided herein is a pharmaceutical composition comprising  
an antagonist, an agonist, a seal coat, and a sequestering  
polymer, wherein the antagonist, agonist, seal coat and at  
least one sequestering polymer are all components of a single  
unit, and wherein the seal coat forms a layer physically sepa-  
rating the antagonist from the agonist from one another.  
Methods for manufacturing such a pharmaceutical composi-  
tion are also provided.

**65 Claims, 14 Drawing Sheets**

U.S. PATENT DOCUMENTS		
4,844,907	A	7/1989 Elger et al.
4,844,909	A	7/1989 Goldie et al.
4,844,910	A	7/1989 Leslie et al.
4,846,199	A	7/1989 Rose
4,861,598	A	8/1989 Oshlack et al.
4,861,781	A	8/1989 Goldberg
4,867,985	A	9/1989 Heafield et al.
4,873,076	A	10/1989 Fishman et al.
4,882,335	A	11/1989 Sinclair
4,889,860	A	12/1989 Rzeszotarski et al.
4,935,428	A	6/1990 Lewis
4,935,429	A	6/1990 Dackis et al.
4,940,587	A	7/1990 Jenkins et al.
4,957,681	A	9/1990 Klimesch et al.
4,970,075	A	11/1990 Oshlack
4,987,136	A	1/1991 Kreek et al.
4,990,341	A	2/1991 Goldie et al.
5,021,053	A	6/1991 Barclay et al.
5,071,646	A	12/1991 Malkowska et al.
5,075,341	A	12/1991 Mendelson et al.
5,086,058	A	2/1992 Sinclair et al.
5,091,189	A	2/1992 Heafield et al.
5,096,715	A	3/1992 Sinclair
5,102,887	A	4/1992 Goldberg
5,130,311	A	7/1992 Guillaumet et al.
5,149,538	A	9/1992 Granger et al.
5,189,064	A	2/1993 Blum et al.
5,198,229	A	3/1993 Wong et al.
5,202,128	A	4/1993 Morella et al.
5,215,758	A	6/1993 Krishnamurthy
5,219,858	A	6/1993 Parnell
5,225,440	A	7/1993 London et al.
5,226,331	A	7/1993 Thompson et al.
5,236,714	A	8/1993 Lee et al.
5,256,669	A	10/1993 Askanazi et al.
5,266,331	A	11/1993 Oshlack et al.
5,266,574	A	11/1993 Zagon et al.
5,273,760	A	12/1993 Oshlack et al.
5,286,493	A	2/1994 Oshlack et al.
5,312,389	A	5/1994 Theeuwes et al.
5,316,759	A	5/1994 Rose et al.
5,317,022	A	5/1994 Borsodi et al.
5,321,012	A	6/1994 Mayer et al.
5,324,351	A	6/1994 Oshlack et al.
5,326,571	A	7/1994 Wright et al.
5,330,766	A	7/1994 Morella et al.
5,336,691	A	8/1994 Raffa et al.
5,352,680	A	10/1994 Portoghese et al.
5,352,683	A	10/1994 Mayer et al.
5,356,467	A	10/1994 Oshlack et al.
5,356,900	A	10/1994 Bihari et al.
5,362,496	A	11/1994 Baker et al.
5,376,384	A	12/1994 Eichel et al.
5,376,662	A	12/1994 Ockert
5,378,474	A	1/1995 Morella et al.
5,385,903	A	1/1995 Steppuhn et al.
5,403,595	A	4/1995 Kitchell et al.
5,409,944	A	4/1995 Black et al.
5,411,745	A	5/1995 Oshlack et al.
5,426,112	A	6/1995 Zagon et al.
5,436,265	A	7/1995 Black et al.
5,451,408	A	9/1995 Mezei et al.
5,457,208	A	10/1995 Portoghese et al.
5,460,826	A	10/1995 Merrill et al.
5,472,712	A	12/1995 Oshlack et al.
5,472,943	A	12/1995 Crain et al.
5,475,995	A	12/1995 Livingston
5,478,573	A	12/1995 Eichel et al.
5,478,577	A	12/1995 Sackler et al.
5,486,362	A	1/1996 Kitchell et al.
5,500,227	A	3/1996 Oshlack et al.
5,502,058	A	3/1996 Mayer et al.
5,507,277	A	4/1996 Rubsamen et al.
5,508,042	A	4/1996 Oshlack et al.
5,508,043	A	4/1996 Krishnamurthy
5,510,368	A	4/1996 Lau et al.
5,512,578	A	4/1996 Crain et al.
5,514,680	A	5/1996 Weber et al.
5,521,213	A	5/1996 Prasit et al.
5,529,787	A	6/1996 Merrill et al.
5,529,790	A	6/1996 Eichel
5,529,813	A	6/1996 Kobsa et al.
5,534,492	A	7/1996 Aston et al.
5,536,752	A	7/1996 Ducharme et al.
5,549,912	A	8/1996 Oshlack et al.
5,550,142	A	8/1996 Ducharme et al.
5,552,422	A	9/1996 Gauthier et al.
5,556,838	A	9/1996 Mayer et al.
5,565,455	A	10/1996 Bjork et al.
5,574,052	A	11/1996 Rose et al.
5,578,725	A	11/1996 Portoghese et al.
5,580,578	A	12/1996 Oshlack et al.
5,580,876	A	12/1996 Crain et al.
5,585,348	A	12/1996 Crain et al.
5,591,452	A	1/1997 Miller et al.
5,593,994	A	1/1997 Batt et al.
5,601,845	A	2/1997 Buxton et al.
5,604,253	A	2/1997 Lau et al.
5,604,260	A	2/1997 Guay et al.
5,616,601	A	4/1997 Khanna et al.
5,622,722	A	4/1997 Knott et al.
5,624,932	A	4/1997 Qin et al.
5,633,259	A	5/1997 Qin et al.
5,639,476	A	6/1997 Oshlack et al.
5,639,780	A	6/1997 Lau et al.
5,656,295	A	8/1997 Oshlack et al.
5,670,172	A	9/1997 Buxton et al.
5,672,360	A	9/1997 Sackler et al.
5,681,585	A	10/1997 Oshlack et al.
5,763,452	A	6/1998 Miller et al.
5,767,125	A	6/1998 Crain et al.
5,780,057	A	7/1998 Conte
5,780,479	A	7/1998 Kim
5,811,126	A	9/1998 Krishnamurthy
5,834,024	A	11/1998 Heinicke et al.
5,834,477	A	11/1998 Mioduszewski
5,843,480	A	12/1998 Miller et al.
5,849,240	A	12/1998 Miller et al.
5,858,017	A	1/1999 Demopulos et al.
5,860,950	A	1/1999 Demopulos et al.
5,866,164	A	2/1999 Kuczynski et al.
5,869,097	A	2/1999 Wong et al.
5,879,705	A	3/1999 Heafield et al.
5,880,132	A	3/1999 Hill
5,891,471	A	4/1999 Miller et al.
5,908,848	A	6/1999 Miller et al.
5,919,473	A	7/1999 Elkhoury
5,935,975	A	8/1999 Rose et al.
5,942,241	A	8/1999 Chasin et al.
5,958,452	A	9/1999 Oshlack et al.
5,958,459	A	9/1999 Chasin et al.
5,965,161	A	10/1999 Oshlack et al.
5,965,163	A	10/1999 Miller et al.
5,968,547	A	10/1999 Reder et al.
5,968,551	A	10/1999 Oshlack et al.
5,972,954	A	10/1999 Foss
5,998,434	A	12/1999 Mitch et al.
6,024,982	A	2/2000 Oshlack et al.
6,033,687	A	3/2000 Heinicke et al.
6,068,855	A	5/2000 Leslie et al.
6,077,532	A	6/2000 Malkowska et al.
6,077,533	A	6/2000 Oshlack et al.
6,103,258	A	8/2000 Simon
6,103,261	A	8/2000 Chasin et al.
6,103,734	A	8/2000 Legarda Ibanez
6,143,322	A	11/2000 Sackler et al.
6,143,328	A	11/2000 Heafield et al.
6,149,940	A	11/2000 Maggi et al.
6,162,467	A	12/2000 Miller et al.
6,210,714	B1	4/2001 Oshlack et al.
6,214,385	B1	4/2001 Heinicke et al.
6,228,863	B1	5/2001 Palermo et al.
6,254,887	B1	7/2001 Miller et al.
6,261,599	B1	7/2001 Oshlack et al.
6,277,384	B1	8/2001 Kaiko et al.
6,294,195	B1	9/2001 Oshlack et al.
6,306,438	B1	10/2001 Oshlack et al.

6,316,031	B1	11/2001	Oshlack				
6,326,027	B1	12/2001	Miller et al.				
6,335,033	B2	1/2002	Oshlack et al.	CA	2229621	9/1998	
6,375,957	B1	4/2002	Kaiko et al.	DE	2222039	11/1972	
6,387,404	B2	5/2002	Oshlack et al.	DE	4325465	A1	2/1995
6,399,096	B1	6/2002	Miller et al.	DE	29719704	U1	3/1998
6,475,494	B2	11/2002	Kaiko et al.	DE	19651551		6/1998
6,627,635	B2	9/2003	Palermo et al.	EP	0074105		3/1983
6,635,277	B2	10/2003	Sharma et al.	EP	0144243	A1	6/1985
6,696,066	B2	2/2004	Kaiko et al.	EP	0185472	A1	6/1986
6,696,088	B2	2/2004	Oshlack et al.	EP	0193355	A2	9/1986
6,716,449	B2	4/2004	Oshlack et al.	EP	0220805	A3	5/1987
6,765,010	B2	7/2004	Crain et al.	EP	0232877	A2	8/1987
6,865,444	B2	3/2005	Howard	EP	0205282		12/1987
6,905,709	B2	6/2005	Oshlack et al.	EP	0319243	A1	6/1989
7,067,151	B1	6/2006	Heinicke et al.	EP	0352361		1/1990
7,070,806	B2	7/2006	Oshlack et al.	EP	502642		9/1992
7,125,561	B2	10/2006	Sackler	EP	0502642	A1	9/1992
7,144,587	B2	12/2006	Oshlack et al.	EP	0647448	A1	4/1995
7,172,767	B2	2/2007	Kaiko et al.	EP	0913152	A1	5/1999
7,332,182	B2	2/2008	Sackler	EP	1041987	B1	4/2006
7,384,653	B2	6/2008	Wright, IV et al.	GB	751767		7/1956
7,419,686	B2	9/2008	Kaiko et al.	GB	769517		3/1957
2002/0010127	A1	1/2002	Oshlack et al.	GB	791644		3/1958
2002/0058673	A1	5/2002	Kaiko et al.	GB	808269		1/1959
2003/0004177	A1	1/2003	Kao et al.	GB	2196848		10/1987
2003/0026838	A1	2/2003	Farrell	WO	WO 8303197		9/1983
2003/0049317	A1	3/2003	Lindsay	WO	WO 8701282	A2	3/1987
2003/0064099	A1	4/2003	Oshlack et al.	WO	WO 9004965	A1	5/1990
2003/0065002	A1	4/2003	Caruso et al.	WO	WO 9406426	A1	3/1994
2003/0068392	A1	4/2003	Sackler	WO	WO 9503804	A1	2/1995
2003/0073714	A1	4/2003	Breder et al.	WO	WO 9602251	A1	2/1996
2003/0124061	A1	7/2003	Roberts	WO	WO 9732573		9/1997
2003/0124185	A1	7/2003	Oshlack et al.	WO	WO 9733566	A2	9/1997
2003/0143269	A1	7/2003	Oshlack et al.	WO	WO 9825613	A2	6/1998
2003/0157167	A1	8/2003	Kao et al.	WO	WO 9835679	A1	8/1998
2003/0157168	A1	8/2003	Breder et al.	WO	WO 9932119		7/1999
2004/0024004	A1	2/2004	Sherman et al.	WO	WO 9932120	A1	7/1999
2004/0024006	A1	2/2004	Simon	WO	WO 0001377	A2	1/2000
2004/0092542	A1	5/2004	Oshlack et al.	WO	WO 0038649	A1	7/2000
2004/0110781	A1	6/2004	Harmon et al.	WO	WO 0132180	A2	5/2001
2004/0131552	A1	7/2004	Boehm	WO	WO 0137785	A2	5/2001
2004/0176402	A1	9/2004	Oshlack et al.	WO	WO 0152851	A1	7/2001
2004/0186121	A1	9/2004	Oshlack et al.	WO	WO 0158447	A1	8/2001
2004/0202717	A1	10/2004	Mehta	WO	WO 0158451		8/2001
2004/0241218	A1	12/2004	Tavares et al.	WO	WO 0168080	A2	9/2001
2005/0095291	A1	5/2005	Oshlack et al.	WO	WO 0185257	A2	11/2001
2005/0163856	A1	7/2005	Maloney et al.	WO	WO 0193852	A2	12/2001
2005/0181046	A1	8/2005	Oshlack et al.	WO	WO 02092059		11/2002
2005/0191244	A1	9/2005	Bartholomaus et al.	WO	WO 2004026256		4/2004
2005/0192309	A1	9/2005	Palermo et al.	WO	WO 2004026283		4/2004
2006/0039970	A1	2/2006	Oshlack et al.	WO	WO 2004/091512		10/2004
2006/0104909	A1	5/2006	Vaghefi et al.	WO	WO 2004093801		11/2004
2006/0182801	A1	8/2006	Breder et al.	WO	WO 2005018616		3/2005
2006/0182891	A1	8/2006	Usoskin et al.	WO	WO 2005055981		6/2005
2006/0198881	A1	9/2006	Howard et al.	WO	WO 2005081825		9/2005
2006/0257460	A1	11/2006	Jansen et al.	WO	WO 2006130471		12/2006
2006/0269605	A1	11/2006	Lizio et al.	WO	WO 2008/011169		1/2008
2007/0014732	A1	1/2007	Sackler	WO	WO 2009/088673		7/2009
2007/0020339	A1	1/2007	Bear				
2007/0042045	A1	2/2007	Lizio et al.				
2007/0185145	A1	8/2007	Royds				
2007/0196396	A1	8/2007	Pilgaonkar et al.				
2008/0069871	A1	3/2008	Vaughn et al.				
2008/0069881	A1	3/2008	Caruso et al.				
2008/0075768	A1	3/2008	Vaughn et al.				
2008/0075770	A1	3/2008	Vaughn et al.				
2008/0075771	A1	3/2008	Vaughn et al.				
2008/0166405	A1	7/2008	Mehta				
2008/0233156	A1	9/2008	Matthews et al.				
2008/0233197	A1	9/2008	Matthews et al.				
2008/0280975	A1	11/2008	Badul				
2008/0306104	A2	12/2008	Oshlack				
2008/0318993	A1	12/2008	Ahdieh				
2008/0318994	A1	12/2008	Ahdieh				
2009/0028942	A1	1/2009	Kandi et al.				
2009/0131466	A1	5/2009	Liang et al.				
2009/0162450	A1	6/2009	Matthews et al.				

## FOREIGN PATENT DOCUMENTS

CA	2229621	9/1998
DE	2222039	11/1972
DE	4325465	A1 2/1995
DE	29719704	U1 3/1998
DE	19651551	6/1998
EP	0074105	3/1983
EP	0144243	A1 6/1985
EP	0185472	A1 6/1986
EP	0193355	A2 9/1986
EP	0220805	A3 5/1987
EP	0232877	A2 8/1987
EP	0205282	12/1987
EP	0319243	A1 6/1989
EP	0352361	1/1990
EP	502642	9/1992
EP	0502642	A1 9/1992
EP	0647448	A1 4/1995
EP	0913152	A1 5/1999
EP	1041987	B1 4/2006
GB	751767	7/1956
GB	769517	3/1957
GB	791644	3/1958
GB	808269	1/1959
GB	2196848	10/1987
WO	WO 8303197	9/1983
WO	WO 8701282	A2 3/1987
WO	WO 9004965	A1 5/1990
WO	WO 9406426	A1 3/1994
WO	WO 9503804	A1 2/1995
WO	WO 9602251	A1 2/1996
WO	WO 9732573	9/1997
WO	WO 9733566	A2 9/1997
WO	WO 9825613	A2 6/1998
WO	WO 9835679	A1 8/1998
WO	WO 9932119	7/1999
WO	WO 9932120	A1 7/1999
WO	WO 0001377	A2 1/2000
WO	WO 0038649	A1 7/2000
WO	WO 0132180	A2 5/2001
WO	WO 0137785	A2 5/2001
WO	WO 0152851	A1 7/2001
WO	WO 0158447	A1 8/2001
WO	WO 0158451	8/2001
WO	WO 0168080	A2 9/2001
WO	WO 0185257	A2 11/2001
WO	WO 0193852	A2 12/2001
WO	WO 02092059	11/2002
WO	WO 2004026256	4/2004
WO	WO 2004026283	4/2004
WO	WO 2004/091512	10/2004
WO	WO 2004093801	11/2004
WO	WO 2005018616	3/2005
WO	WO 2005055981	6/2005
WO	WO 2005081825	9/2005
WO	WO 2006130471	12/2006
WO	WO 2008/011169	1/2008
WO	WO 2009/088673	7/2009

## OTHER PUBLICATIONS

Rao, et al., "Effect of Sodium Lauryl Sulfate on the Release of Rifampicin from Guar Gum Matrix", *Indian J. of Pharm. Sci.* pp. 404-406 (Sep.-Oct. 2000).

Abdulla, et al., "Axotomy reduces the effect of analgesic opioids yet increases the effect of nociceptin on dorsal root . . .", *J. Neuro Sci* 1998, p. 9685-9694, vol. 18.

Alvarez-Fuentes, et al., "Effectiveness of Repeated Administration of a New Oral Naltrexone . . ."; *J. Pharm Pharmacol* (2001), p. 1201-1205, v. 53.

Alvarez-Fuentes, et al., "Preclinical Study of an Oral Controlled Release Naltrexone Complex in Mice"; *J. Pharm Pharmacol* (2000), p. 659-663, v. 52.

Barton et al., "Intranasal Administration of Naloxone by Paramedics"; *Prehospital Emergency Care* (2002), p. 54-58, vol. 6, No. 1.


Baum et al., "The Impact of the Addition of Naloxone on the Use and Abuse of Pentazocine"; *Public Health Reports*, (1987) p. 426-429. v. 102(4).

- Blachly, M.D., "Naloxone in Opiate Addiction"; *Current Psychiatric Therapies* (1976) p. 209-213.
- Bloom et al., *Clinical Studies with Naloxone/Methadone in a Ratio of 1:20*; 5. sup.th National Conference on Methadone Treatment (1973), p. 1342-1349, vol. 2.
- Brennscheidt et al., *Pharmacokinetics of Nortilidine and Naloxone . . .*, *Arzneim.-Forsch/Drug Res.* (2000), p. 1015-1022, vol. 50.
- Bigelow et al., "Abuse Liability Assessment . . ."; Dept. of Psychiatry and Behavioral Sciences—The Johns Hopkins University School of Medicine. p. 145-149.
- Bromm, et al., "A Sensitive Method to Evaluate Effects of Analgesics in Man"; *Meth and Find Exptl Clin Pharmacol*, (1983) p. 545-551, v. 5 No. 8.
- Bullingham, et al., "Clinical Pharmacokinetics of Narcotic Agonist-Antagonist Drugs"; *Clinical Pharm.* (1983) p. 332-343, v. 8.
- Cherny, N., "Opioid Analgesics"; *Drugs* (1996), p. 713-737, v. 51 (5).
- Fraser, Albert D., et al., "Clinical Toxicology of Drugs Used in the Treatment of Opiate Dependency"; *Clinical Toxicology I*, (1990), p. 375-386, vol. 10, No. 2.
- Caruso et al., "Methadone and Naloxone in Combination (Naldone. RTM.) for the Treatment of Heroin Addicts"; Bristol Laboratories, p. 1336-1341F.
- Crabtree et al., "Review of Naltrexone, a long-acting opiate antagonist"; *Clinical Pharmacy*, (1984) p. 273-280, vol. 3.
- Crain et al., "Ultra-low concentrations of naloxone selectively antagonize excitatory . . ." *Proc. Natl. Acad. Sci* (1995), p. 10540-10544, vol. 92.
- Czarnecki et al. "The Use of Photography in the Clinical Evaluation of Unequal Pupils." *Canad J Ophthal* 1979, p. 297-302, v. 14.
- Driscoll, P. "Comparative Toxicity for Mice . . ." *Praeventimedizin*, 1972, p. 211-213, v. 17 (4).
- Gal et al., "Prolonged Blockade of Opioid Effect with Oral Nalmefene"; *Clin Pharmacol Ther.* (1986) p. 537-542.
- Ghodse, et al., "Opioid analgesics and narcotic antagonists"; *Side Effects of Drugs* (2000), p. 96-113, Annual 23, Ch. 8.
- Gonzalez et al., "Naltrexone: A Review of its Pharmacodynamic and Pharmacokinetic Properties and Therapeutic . . ." *Drugs* (1988), p. 192-213., v. 35.
- Holmes et al., *Anesth. Analg.* (1993), p. 1166-73, v. 77.
- Goodman et al., *Pharmacological Basis of Therapeutics*, 3rd Edition, 1965, pp. 274-279.
- Gawin, Fh. "Chronic Neuropharmacology of Cocaine", *Jr. of Clin. Psychiatry*, Feb. 1988, p. 11-16. v. 49(2).
- Gan et al., "Opioid-Sparing Effects of a Low-Dose Infusion of Naloxone in Patient-Administered Morphine Sulfate," *Anesthesiology* (1997), 1075-1080, v. 87(5).
- Glick et al "Titration of Oral nicotine Intake . . ." *Nature*, 1971, p. 207-208, v. 233.
- Hanson, *Analgesic, Antipyretic and Anti-Inflammatory Drugs in Remington's Science and Practice of Pharmacy*, 1995, p. 1207, v. 2.
- Harris et al. "Protection Against Disopropyfluorophosphate Intox." *Arch. of Pharm.* 1984. p. 64-69, v.327.
- Harris, LS. *Central Neuropunoral Systems*, *Fed Proceed.*, Jan.-Feb. 1970, p. 28-32, v. 20:1.
- Hawkes et al., "Effect of an enteric-release formulation of naloxone on intestinal transit in volunteers taking codeine"; *Aliment Pharm Ther* (2001), p. 625-630, vol. 15.
- Levine, et al. "Potentiation of Pentazocine Analgesia by Low-dose Naloxone", *J. Clin. Invest.*, 1988, p. 1574-1577, vol. 82.
- Hogger et al., "Comparison of tilidine/naloxone, tramadol and bromfenac in experimental pain . . ." *International J. Clin Pharm and Ther* (1999), p. 377-385, vol. 37, No. 8.
- Hussain et al., "Buccal and oral bioavailability of naloxone and naltrexone in rats"; (1987), p. 127-130, vol. 36.
- King et al., "Naltrexone Biotransformation and Incidence of Subjective Side Effects: A Preliminary Study"; *Alcoholism: Clin and Exp Res* (1997), p. 906-909, vol. 21, No. 5.
- Kurland et al., "Naloxone and the Narcotic Abuser: A Controlled Study of Partial Blockade"; *Inter. J. of the Addictions* (1974), p. 663-672, vol. 9, No. 5.
- Martin et al., "Bioavailability Investigation of a New Tilidine/Naloxone Liquid Formulation Compared . . ." *Arzneim.-Forsch./Drug Res.* (1999), p. 599-607, vol. 49.
- Lehmann et al., "Influence of Naloxone on the Postoperative Analgesic and Respiratory Effects of Buprenorphine"; *Eur J. Clin. Pharm.* (1988), 343-352, vol. 34.
- Jasinski D.R., "Assessment of the Abuse Potentiality of Morphinelike Drugs (Methods Used in Man)"; *Drug Addiction* (1977) p. 197-258.
- Johnson et al., "Buprenorphine and Naloxone for Heroin Dependence"; *Substance Use Disorders*(2000) p. 519-526.
- Tai et al., "Naltrexone: An Antagonist Therapy for Heroin Addiction"; *NIDA* (1997) 5 pages.
- Stine et al., "Reduction of Opiate Withdrawal-like Symptoms by Cocaine Abuse during . . ." *Am. J. Drug Alcohol Abuse* (1994), p. 445-458, vol. 20, (4).
- Stine et al., "Use of Drug Combinations in Treatment of Opioid Withdrawal"; *J. of Clinical Psych.* (1992), p. 203-209, vol. 12, No. 3.
- Suzuki, et al., "Morphine conditioned place preference after chronic treatment with naloxone . . ." *Research Communications in Substances of Abuse.* (1991), vol. 12(3), p. 191-131.
- Tennant et al "Withdrawal From Nicotine Dependence . . .", *NIDA Research Monograph*, 1985, p. 291-297, v. 55.
- Walsh et al., "Effects of Naltrexone on Response to Intravenous Cocain, Hydromorphone and their Combination in Humans," (1996).
- Way et al., "Responsivity to Naloxone during Morphine Dependence"; *Annals New York Academy of Sciences*, p. 252-261.
- Wodak Alex, "Drug Treatment for Opioid Dependence"; *Australian Prescriber* (2001), p. 4-6, vol. 24, No. 1.
- Parwartikar et al., "Naloxone-Methadone Combination for the Treatment of Opiate Dependence"; *Missouri Institute of Psychiatry*, p. 1350-1354.
- Wikler et al., *N-Allylnormorphine: Effects of single doses and precipitation of acute . . . N-Allylnormorphine During Narcotic Addiction* (1953) p. 8-20.
- Vaccarino et al., *Pain*, 1989, p. 103-109, v. 36.
- Yuan et al., *Clinical Trials and Therapeutics*, (Apr. 1997) 467-475, v. 61.
- Zaks et al., "Naloxone Treatment of Opiate Dependence"; *JAMA* (1971), p. 2108-2110, vol. 215, No. 13.
- Alvarez-Fuentes, et al., "Study of a complexation process between naltrexone and Eudragit . . ." *Int'l Journal of Pharmaceutics* 148 (1997) 219-230.
- W.K. Asbeck, et al. "Critical Pigment Volume Relationships", *Ind. Eng. Chem.*, 1949, (41), 1470-1475.
- Bashaw, et al. "Relative bioavailability of controlled-release oral morphine sulfate . . ." *Int'l Journal of Clinical Pharmacology and Therapeutics*, vol. 33 No. 9 1995(524-529).
- Bierwagen, et al., "Studies of the Effects of Particle Size Distribution on the Packing Efficiency of Particles", *Powder Tech.*, 10(1974) 111-119.
- Bodmeier, et al., "Dry and wet strengths of polymeric films prepared from an aqueous colloidal polymer dispersion, Eudragit RS30D", *Int'l Journ. of Pharma.*, 96(1993) 129-138.
- Bodmeier, et al. "Mechanical Properties of Dry and Wet Cellulosic and Acrylic Films Prepared from Aqueous Colloidal Polymer . . .", *Pharmaceutical Research*, vol. 11, No. 6 (1994).
- Childs, et al. "Crystal Engineering Approach to Forming Cocrystals of Amine Hydrochlorides With Organic Acids . . ." *J. Am. Chem. SOc.* 2004, 126 p. 13335-13342.
- Fernandez-Arevalo, et al., "Water-Insoluble Complexes of Morphine and Carteolol-Eudragit L.A. . . .", *Supplied by the British Library*, p. 158-181.
- Gould, "Salt selection for basic drugs", *International Journal of Pharmaceutics*, 33 (1986) 201-207.
- Holgado, et al., "Physical characterization of carteolol: Eudragit L binding interaction", *International Journal of Pharmaceutics* 114 (1995) p. 13-21.
- Judson, et al., "Chapter 1. Uses of Naloxone in the Diagnosis and Treatment of Heroin Addiction", p. 1-13.
- Kelly, et al., "Prevalence and characteristics of opioid use in the US adult population", *PAIN* 138 (2008) p. 507-513.

- Lang, et al., "Cardiovascular responses to injections of cholinomimetic drugs into the cerebral ventricles . . .", *Br. J. Pharmac.* (1973), 47, p. 196-205.
- McGinity, et al., "In Vitro Adsorption of Various Pharmaceuticals to Montmorillonite", *Journal of Pharmaceutical Sciences*, vol. 65, Jun. 1976 p. 896-902.
- Mendelson, et al. "Buprenorphine and naloxone interactions in opiate-dependent volunteers", *Clinical Pharmacology & Therapeutics*, (1996) p. 105-114.
- Meyer, et al. "A Behavioral Paradigm for the Evaluation of Narcotic Antagonists", *Arch. Gen Psychiatry*, vol. 33 (1976), pp. 371-377.
- Pakkanen, "Upregulation and Functionality of Neuronal Nicotinic Acetylcholine Receptors", *Div. of Pharm. and Toxicology Faculty Pharm. Univ. Helsinki*, p. 1-89.
- Patel, et al. "Nanoclays for Polymer nanocomposites, paints, inks, greases and cosmetics formulations, . . .", *Bull. Mater. Sci.*, vol. 29, No. 2 (2006) pp. 133-145.
- Peachey, et al. "Assessment of Opioid Dependence with Naloxone", *British Journal of Addiction* (1988) 83, pp. 193-201.
- Sunshine, et al. "Analgesic Efficacy of Pentazocine Versus a Pentazocine-Naloxone Combination Following Oral Administration", *The Clinical Journal of Pain* (1988) pp. 35-40.
- Rogers, et al., *Glass Transitions of the Poly-(n-Alkyl Methacrylates)*, *J. Phys. Chem.*, 1957, 61 (7), pp. 985-991.
- Bierwagen et al., "Recent Studies of Particle Packing in Organic Coatings"; *Progress in Organic Coatings*, 35 (1999) pp. 1-9.
- Okhamafe, et al., "Interaction Phenomena in Pharmaceutical Film Coating and Testing Methods" *Int. J. Pharmaceutics*; vol. 39 (1987) pp. 1-21.
- Farris, R.J.; "Prediction of the Viscosity of Multimodal Suspensions from Unimodal Viscosity Data" *Trans. Soc. Rheology*, vol. 12, No. 2, (1968) pp. 281-301.
- Knop, "Influence of Buffer Solution Composition on Drug Release from Pellets Coated with Neutral and Quaternary . . .", *Euro. J. Pharm Sci.*, vol. 4, (1996) pp. 293-300.
- Ozturk, et al. "Mechanism of Release from Pellets Coated with an Ethylcellulose Based Film" *J. Cont. Release*, vol. 14, (1990) pp. 203-213.
- Briscoe, B. et al. "Rheology of Solvent-Cast Polymer Films", *Journal of Applied Polymer Science*, vol. 28 (1983) No. 3, pp. 3827-3848.
- Michelson, A.W, "Ion Exchange", *Chem. Engineering*, Mar. 18, 1963 pp. 163-182.
- Hercules Inc., "Physical and Chemical Properties", pp. 1-34.
- Dressman J. et al., "Dissolution Testing as a Prognostic Tool for Oral Drug Absorption: . . ." *Pharm. Research*, 15 (1) pp. 11-22 (1998).
- Galia, et al., "Evaluation of Various Dissolution Media for Predicting In Vivo Performance of Class I and II Drugs" *Pharm. Research*, 15 (5) pp. 698-705 (1998).
- Vertzoni, et al., "Dissolution media simulating the intraluminal composition of the small intestine: . . ." *J. of Pharmacy and Pharm.*, vol. 56, 2004; pp. 453-462.
- Pederson, et al., "Dissolution of Hydrocortisone in Human and Simulated Intestinal Fluids" *Pharm. Research*, vol. 17, No. 2, 2000; pp. 183-189.
- Powell, et al., "Application of the critical precipitation essay to complex samples: aluminum binding . . ." *Chem. Speciation and Bioavailability*; vol. 16, No. 3, 2004, pp. 97-104.
- Pharm. Coatings Bulletin* 102-3, "Influence of Plasticizers on the Dissolution and Physical Properties of Ethyl Cellulose Films and Coated Beads" pp. 1-8 (1995).
- Hjartstam, et al. "Effect of Hydroxyl Group Content in Ethyl Cellulose on Permeability in Free Films and Coated Membranes", *J. Appl. Polymer Sci.*, V. 72, pp. 529-535 (1999).
- Feller, et al. "Evaluation of Cellulose Ethers for Conservation", *The Getty Conserv. Inst.*, (1990) pp. 1-161.
- Rhom GmbH & Co. KG, Specifications and test methods for Eudragit RL 100 and Eudragit RL PO Eudragit RS 100 and Eudragit RS PO, (Sep. 2004), pp. 1-4.
- Lehmann, et al. "Practical Course in Film Coating of Pharmaceutical Dosage Forms with Eudragit", *Pharma Polymers* (2001) pp. 1-199.
- Blackie Academic and Prof. "Introduction to Surfactant Analysis", (1992) pp. 1-3.
- Kaufman, et al., "Narcotic and Narcotic Antagonist pKa's and Partition Coefficients and Their Significance in Clinical Practice", *Drug and Alcohol Depend.* 1 (1975-76) pp. 103-114.
- The Dow Chem. Co., Technical Information, "Dowex Ion Exchange Resins—Using Ion Exchange Resin Selectivity Coefficients", pp. 1-3.
- van der Wel, "Moisture in Organic Coatings—a review", *Progress in Organic Coatings* 37 (1999) pp. 1-14.


**FIGURE 1**

**NT release from Eudragit RS-coated pellets without SLS**



**FIGURE 2**


**Effect of SLS levels in Eudragit RS coat on Naltrexone (NT) release**



- RB 358-88 — □ —
- RB 358-73 — △ —
- RB 358-83 — × —

**FIGURE 3**

**Effect of SLS levels in Eudragit RS coat on NT release**



**FIGURE 4**

**Effect of talc levels in Eudragit RS coat on NT release**





FIGURE 5

Naltrexone Dissolution Profile vs. % RS Neutralization at 26% (v/v) Talc



34.5 – 39 micron 

51.5 – 55.5 micron 

63 – 66 micron 



FIGURE 6

Naltrexone Dissolution Profile vs. Talc Level at  
41% RS Neutralization




FIGURE 7




FIGURE 8




FIGURE 9

**Plasma Naltrexone levels for PI-1460 and PI-1461**




FIGURE 10

**Plasma 6-beta-naltrexol levels for PI-1460 and PI-1461**




FIGURE 11




FIGURE 12




FIGURE 13




FIGURE 14

**Plasma Naltrexone levels for PI-1465 and PI-1466**




FIGURE 15

**Plasma 6-beta-naltrexol levels for PI-1465 and PI-1466**




FIGURE 16




FIGURE 17




FIGURE 18




FIGURE 19






FIGURE 20

**Plasma Naltrexone levels for PI 1510**






FIGURE 21


**Plasma 6-beta-Naltrexol levels**



**Figure 22A. Manufacturing Process**



**Figure 22B. Manufacturing Process**



1

**ABUSE-DETERRENT MULTI-LAYER  
PHARMACEUTICAL COMPOSITION  
COMPRISING AN OPIOID ANTAGONIST AND  
AN OPIOID AGONIST**

RELATED APPLICATIONS

This application claims priority to U.S. Ser. No. 60/814,949 filed Jun. 19, 2006.

FIELD OF THE INVENTION

This invention pertains to a sequestering subunit comprising an antagonist and a blocking agent, and related compositions and methods of use, such as in the prevention of abuse of a therapeutic agent.

BACKGROUND OF THE INVENTION

Opioids, also called opioid agonists, are a class of drugs that exhibit opium-like or morphine-like properties. The opioids are employed primarily as moderate to strong analgesics, but have many other pharmacological effects as well, including drowsiness, respiratory depression, changes in mood, and mental clouding without a resulting loss of consciousness. Because of these other pharmacological effects, opioids have become the subject of dependence and abuse. Therefore, a major concern associated with the use of opioids is the diversion of these drugs from the illicit user, e.g., an addict.

Physical dependence may develop upon repeated administrations or extended use of opioids. Physical dependence is gradually manifested after stopping opioid use or is precipitously manifested (e.g., within a few minutes) after administration of a narcotic antagonist (referred to "precipitated withdrawal"). Depending upon the drug upon which dependence has been established and the duration of use and dose, symptoms of withdrawal vary in number and kind, duration and severity. The most common symptoms of the withdrawal syndrome include anorexia, weight loss, pupillary dilation, chills alternating with excessive sweating, abdominal cramps, nausea, vomiting, muscle spasms, hyperirritability, lacrimation, rinorrhea, goose flesh and increased heart rate. Natural abstinence syndromes typically begin to occur 24-48 hours after the last dose, reach maximum intensity about the third day and may not begin to decrease until the third week. Precipitated abstinence syndromes produced by administration of an opioid antagonist vary in intensity and duration with the dose and the specific antagonist, but generally vary from a few minutes to several hours in length.

Psychological dependence or addiction to opioids is characterized by drug-seeking behavior directed toward achieving euphoria and escape from, e.g., psychosocioeconomic pressures. An addict will continue to administer opioids for non-medicinal purposes and in the face of self-harm.

Although opioids, such as morphine, hydromorphone, hydrocodone and oxycodone, are effective in the management of pain, there has been an increase in their abuse by individuals who are psychologically dependent on opioids or who misuse opioids for non-therapeutic reasons. Previous experience with other opioids has demonstrated a decreased abuse potential when opioids are administered in combination with a narcotic antagonist, especially in patients who are ex-addicts (Weinhold et al., *Drug and Alcohol Dependence* 30:263-274 (1992); and Mendelson et al., *Clin. Pharm. Ther.* 60:105-114 (1996)). These combinations, however, do not contain the opioid antagonist that is in a sequestered form.

2

Rather, the opioid antagonist is released in the gastrointestinal system when orally administered and is made available for absorption, relying on the physiology of the host to metabolize differentially the agonist and antagonist and negate the agonist effects.

Previous attempts to control the abuse potential associated with opioid analgesics include, for example, the combination of pentazocine and naloxone in tablets, commercially available in the United States as Talwin®Nx from Sanofi-Winthrop, Canterbury, Australia. Talwin®Nx contains pentazocine hydrochloride equivalent to 50 mg base and naloxone hydrochloride equivalent to 0.5 mg base. Talwin®Nx is indicated for the relief of moderate to severe pain. The amount of naloxone present in this combination has low activity when taken orally, and minimally interferes with the pharmacologic action of pentazocine. However, this amount of naloxone given parenterally has profound antagonistic action to narcotic analgesics. Thus, the inclusion of naloxone is intended to curb a form of misuse of oral pentazocine, which occurs when the dosage form is solubilized and injected. Therefore, this dosage has lower potential for parenteral misuse than previous oral pentazocine formulations. However, it is still subject to patient misuse and abuse by the oral route, for example, by the patient taking multiple doses at once. A fixed combination therapy comprising tilidine (50 mg) and naloxone (4 mg) has been available in Germany for the management of severe pain since 1978 (Valoron®N, Goedecke). The rationale for the combination of these drugs is effective pain relief and the prevention of tilidine addiction through naloxone-induced antagonisms at the tilidine receptors. A fixed combination of buprenorphine and naloxone was introduced in 1991 in New Zealand (Terngesic®Nx, Reckitt & Colman) for the treatment of pain.

International Patent Application No. PCT/US01/04346 (WO 01/58451) to Euroceltique, S. A., describes the use of a pharmaceutical composition that contains a substantially non-releasing opioid antagonist and a releasing opioid agonist as separate subunits that are combined into a pharmaceutical dosage form, e.g., tablet or capsule. However, because the agonist and antagonist are in separate subunits, they can be readily separated. Further, providing the agonist and antagonist as separate subunits, tablets are more difficult to form due to the mechanical sensitivity of some subunits comprising a sequestering agent.

The benefits of the abuse-resistant dosage form are especially great in connection with oral dosage forms of strong opioid agonists (e.g., morphine, hydromorphone, oxycodone or hydrocodone), which provide valuable analgesics but are prone to being abused. This is particularly true for sustained-release opioid agonist products, which have a large dose of a desirable opioid agonist intended to be released over a period of time in each dosage unit. Drug abusers take such sustained release product and crush, grind, extract or otherwise damage the product so that the full contents of the dosage form become available for immediate absorption.

Such abuse-resistant, sustained-release dosage forms have been described in the art (see, for example, U.S. Application Nos. 2003/0124185 and 2003/0044458). However, it is believed that substantial amounts of the opioid antagonist or other antagonist found in these sequestered forms are released over time (usually less than 24 hours) due to the osmotic pressure that builds up in the core of the sequestered form, as water permeates through the sequestered form into the core. The high osmotic pressure inside the core of the sequestered form causes the opioid antagonist or antagonist

to be pushed out of the sequestered form, thereby causing the opioid antagonist or antagonist to be released from the sequestered form.

In view of the foregoing drawbacks of the sequestered forms of the prior art, there exists a need in the art for a sequestered form of an opioid antagonist or other antagonist that is not substantially released from the sequestered form due to osmotic pressure. The invention provides such a sequestering form of an opioid antagonist or antagonist. This and other objects and advantages of the invention, as well as additional inventive features, will be apparent from the description of the invention provided herein.

#### BRIEF SUMMARY OF THE INVENTION

Provided herein is a pharmaceutical composition comprising an antagonist, an agonist, a seal coat, and a sequestering polymer, wherein the antagonist, agonist, seal coat and at least one sequestering polymer are all components of a single unit, and wherein the seal coat forms a layer physically separating the antagonist from the agonist from one another. In one embodiment, a multi-layer pharmaceutical composition comprising an agonist and an antagonist thereof, wherein the agonist and antagonist are not in contact with one another in the intact form of the composition, wherein the agonist is substantially released and the antagonist is substantially sequestered upon administration to a human being is provided. Methods for manufacturing such a pharmaceutical composition are also provided. In another embodiment, a method for measuring the amount of antagonist or derivative thereof in a biological sample, the antagonist or derivative having been released from a pharmaceutical composition in vivo, the method comprising the USP paddle method at 37° C., 100 rpm, but further comprising incubation in a buffer containing a surfactant.

#### BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1. Naltrexone (NT) release from Eudragit® RS-coated pellets without SLS.

FIG. 2. Effect of SLS levels in Eudragit®RS coat on naltrexone (NT) release.

FIG. 3. Effect of SLS levels in Eudragit® RS coat on naltrexone (NT) release.

FIG. 4. Effect of talc levels in Eudragit® RS coat on naltrexone (NT) release.

FIG. 5. Naltrexone dissolution profile vs. Eudragit® RS neutralization at 26% (v/v) talc.

FIG. 6. Naltrexone dissolution profile vs. talc level at 41% Eudragit® RS neutralization.

FIG. 7. Plasma naltrexone levels for naltrexone (NTX) solution.

FIG. 8. Plasma 6-beta-naltrexol levels for naltrexone (NTX) solution.

FIG. 9. Plasma naltrexone levels for PI-1460 and PI-1461.

FIG. 10. Plasma 6-beta naltrexol levels for PI-1460 and PI-1461.

FIG. 11. Plasma naltrexone levels for PI-1462 and PI-1463.

FIG. 12. Plasma 6-beta-naltrexol levels for PI-1462 and PI-1463.

FIG. 13. Percent naltrexone (NT) release for PI-1465 and PI-1466.

FIG. 14. Plasma naltrexone levels for PI-1465 and PI-1466.

FIG. 15. Plasma 6-beta-naltrexol levels for PI-1465 and PI-1466.

FIG. 16. Plasma naltrexone levels for PI-1465 and PI-1466 (fast and fed).

FIG. 17. Plasma 6-beta-naltrexol levels for PI-1465 and PI-1466 (fast and fed).

FIG. 18. Plasma naltrexone levels for PI-1495 and PI-1496 (fast and fed).

FIG. 19. Plasma 6-beta-naltrexol levels for PI-1495 and PI-1496 (fast and fed).

FIG. 20. Plasma naltrexone levels for PI-1510 (fast and fed).

FIG. 21. Plasma 6-beta-naltrexol levels for PI-1510 (fast and fed).

FIGS. 22 A and B. Exemplary manufacturing process for multi-layer naltrexone-morphine pharmaceutical composition.

#### DETAILED DESCRIPTION OF THE INVENTION

Provided herein are compositions and methods for administering a multiple active agents to a mammal in a from and manner that minimizes the effects of either active agent upon the other in vivo. In certain embodiments, at least two active agents are formulated as part of a pharmaceutical composition. A first active agent may provide a therapeutic effect in vivo. The second active agent may be an antagonist of the first active agent, and may be useful in preventing misuse of the composition. For instance, where the first active agent is a narcotic, the second active agent may be an antagonist of the narcotic. The composition remains intact during normal usage by patients and the antagonist is not released. However, upon tampering with the composition, the antagonist may be released thereby preventing the narcotic from having its intended effect. In certain embodiments, the active agents are both contained within a single unit, such as a bead, in the form of layers. The active agents may be formulated with a substantially impermeable barrier as, for example, a controlled-release composition, such that release of the antagonist from the composition is minimized. In certain embodiments, the antagonist is released in in vitro assays but is substantially not released in vivo. In vitro and in vivo release of the active agent from the composition may be measured by any of several well-known techniques. For instance, in vivo release may be determined by measuring the plasma levels of the active agent or metabolites thereof (i.e., AUC, Cmax).

In one embodiment, the invention provides a sequestering subunit comprising an opioid antagonist and a blocking agent, wherein the blocking agent substantially prevents release of the opioid antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. This sequestering subunit is incorporated into a single pharmaceutical unit that also includes an opioid agonist. The pharmaceutical unit thus includes a core portion to which the opioid antagonist is applied. A seal coat is then optionally applied upon the antagonist. Upon the seal coat is then applied a composition comprising the pharmaceutically active agent. An additional layer containing the same or a different blocking agent may then be applied such that the opioid agonist is released in the digestive tract over time (i.e., controlled release). Thus, the opioid antagonist and the opioid agonist are both contained within a single pharmaceutical unit, which is typically in the form of a bead.

The term “sequestering subunit” as used herein refers to any means for containing an antagonist and preventing or substantially preventing the release thereof in the gastrointestinal tract when intact, i.e., when not tampered with. The term “blocking agent” as used herein refers to the means by which the sequestering subunit is able to prevent substantially the

antagonist from being released. The blocking agent may be a sequestering polymer, for instance, as described in greater detail below.

The terms “substantially prevents,” “prevents,” or any words stemming therefrom, as used herein, means that the antagonist is substantially not released from the sequestering subunit in the gastrointestinal tract. By “substantially not released” is meant that the antagonist may be released in a small amount, but the amount released does not affect or does not significantly affect the analgesic efficacy when the dosage form is orally administered to a host, e.g., a mammal (e.g., a human), as intended. The terms “substantially prevents,” “prevents,” or any words stemming therefrom, as used herein, does not necessarily imply a complete or 100% prevention. Rather, there are varying degrees of prevention of which one of ordinary skill in the art recognizes as having a potential benefit. In this regard, the blocking agent substantially prevents or prevents the release of the antagonist to the extent that at least about 80% of the antagonist is prevented from being released from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. Preferably, the blocking agent prevents release of at least about 90% of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. More preferably, the blocking agent prevents release of at least about 95% of the antagonist from the sequestering subunit. Most preferably, the blocking agent prevents release of at least about 99% of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours.

For purposes of this invention, the amount of the antagonist released after oral administration can be measured in-vitro by dissolution testing as described in the United States Pharmacopeia (USP26) in chapter <711> Dissolution. For example, using 900 mL of 0.1 N HCl, Apparatus 2 (Paddle), 75 rpm, at 37° C. to measure release at various times from the dosage unit. Other methods of measuring the release of an antagonist from a sequestering subunit over a given period of time are known in the art (see, e.g., USP26).

Without being bound to any particular theory, it is believed that the sequestering subunit of the invention overcomes the limitations of the sequestered forms of an antagonist known in the art in that the sequestering subunit of the invention reduces osmotically-driven release of the antagonist from the sequestering subunit. Furthermore, it is believed that the present inventive sequestering subunit reduces the release of the antagonist for a longer period of time (e.g., greater than 24 hours) in comparison to the sequestered forms of antagonists known in the art. The fact that the sequestered subunit of the invention provides a longer prevention of release of the antagonist is particularly relevant, since precipitated withdrawal could occur after the time for which the therapeutic agent is released and acts. It is well known that the gastrointestinal tract transit time for individuals varies greatly within the population. Hence, the residue of the dosage form may be retained in the tract for longer than 24 hours, and in some cases for longer than 48 hours. It is further well known that opioid analgesics cause decreased bowel motility, further prolonging gastrointestinal tract transit time. Currently, sustained-release forms having an effect over a 24 hour time period have been approved by the Food and Drug Administration. In this regard, the present inventive sequestering subunit provides prevention of release of, the antagonist for a time period that is greater than 24 hours when the sequestering subunit has not been tampered.

The sequestering subunit of the invention is designed to prevent substantially the release of the antagonist when

intact. By “intact” is meant that a dosage form has not undergone tampering. The term “tampering” is meant to include any manipulation by mechanical, thermal and/or chemical means, which changes the physical properties of the dosage form. The tampering can be, for example, crushing, shearing, grinding, chewing, dissolution in a solvent, heating (for example, greater than about 45° C.), or any combination thereof. When the sequestering subunit of the invention has been tampered with, the antagonist is immediately released from the sequestering subunit.

By “subunit” is meant to include a composition, mixture, particle; etc., that can provide a dosage form (e.g., an oral dosage form) when combined with another subunit.

The subunit can be in the form of a bead, pellet, granule, spheroid, or the like, and can be combined with additional same or different subunits, in the form of a capsule, tablet or the like, to provide a dosage form, e.g., an oral dosage form. The subunit may also be part of a larger, single unit, forming part of that unit, such as a layer. For instance, the subunit may be a core coated with an antagonist and a seal coat; this subunit may then be coated with additional compositions including a pharmaceutically active agent such as an opioid agonist.

By “antagonist of a therapeutic agent” is meant any drug or molecule, naturally-occurring or synthetic that binds to the same target molecule (e.g., a receptor) of the therapeutic agent, yet does not produce a therapeutic, intracellular, or in vivo response. In this regard, the antagonist of a therapeutic agent binds to the receptor of the therapeutic agent, thereby preventing the therapeutic agent from acting on the receptor. In the case of opioids, an antagonist may prevent the achievement of a “high” in the host.

The antagonist can be any agent that negates the effect of the therapeutic agent or produces an unpleasant or punishing stimulus or effect, which will deter or cause avoidance of tampering with the sequestering subunit or compositions comprising the same. Desirably, the antagonist does not harm a host by its administration or consumption but has properties that deter its administration or consumption, e.g., by chewing and swallowing or by crushing and snorting, for example. The antagonist can have a strong or foul taste or smell, provide a burning or tingling sensation, cause a lachrymation response, nausea, vomiting, or any other unpleasant or repugnant sensation, or color tissue, for example. Preferably, the antagonist is selected from the group consisting of an antagonist of a therapeutic agent, a bittering agent, a dye, a gelling agent, and an irritant. Exemplary antagonists include capsaicin, dye, bittering agents and emetics. The antagonist can comprise a single type of antagonist (e.g., a capsaicin), multiple forms of a single type of antagonist (e.g., a capasin and an analogue thereof), or a combination of different types of antagonists (e.g., one or more bittering agents and one or more gelling agents). Desirably, the amount of antagonist in the sequestering subunit of the invention is not toxic to the host.

In the instance when the therapeutic agent is an opioid agonist, the antagonist preferably is an opioid antagonist, such as naltrexone, naloxone, nalmefene, cyclazacine, levallorphan, derivatives or complexes thereof, pharmaceutically acceptable salts thereof, and combinations thereof. More preferably, the opioid antagonist is naloxone or naltrexone. By “opioid antagonist” is meant to include one or more opioid antagonists, either alone or in combination, and is further meant to include partial antagonists, pharmaceutically acceptable salts thereof, stereoisomers thereof, ethers thereof, esters thereof, and combinations thereof. The pharmaceutically acceptable salts include metal salts, such as sodium salt, potassium salt, cesium salt, and the like; alkaline

earth metals, such as calcium salt, magnesium salt, and the like; organic amine salts, such as triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N-dibenzylethylenediamine salt, and the like; inorganic acid salts, such as hydrochloride, hydrobromide, sulfate, phosphate, and the like; organic acid salts, such as formate, acetate, trifluoroacetate, maleate, tartrate, and the like; sulfonates, such as methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like; amino acid salts, such as arginate, asparaginate, glutamate, and the like. In certain embodiments, the amount of the opioid antagonist can be about 10 ng to about 275 mg. In a preferred embodiment, when the antagonist is naltrexone, it is preferable that the intact dosage form releases less than 0.125 mg or less within 24 hours, with 0.25 mg or greater of naltrexone released after 1 hour when the dosage form is crushed or chewed.

In a preferred embodiment, the opioid antagonist comprises naloxone. Naloxone is an opioid antagonist, which is almost void of agonist effects. Subcutaneous doses of up to 12 mg of naloxone produce no discernable subjective effects, and 24 mg naloxone causes only slight drowsiness. Small doses (0.4-0.8 mg) of naloxone given intramuscularly or intravenously in man prevent or promptly reverse the effects of morphine-like opioid agonist. One mg of naloxone intravenously has been reported to block completely the effect of 25 mg of heroin. The effects of naloxone are seen almost immediately after intravenous administration. The drug is absorbed after oral administration, but has been reported to be metabolized into an inactive form rapidly in its first passage through the liver, such that it has been reported to have significantly lower potency than when parenterally administered. Oral dosages of more than 1 g have been reported to be almost completely metabolized in less than 24 hours. It has been reported that 25% of naloxone administered sublingually is absorbed (Weinberg et al., *Clin. Pharmacol. Ther.* 44:335-340 (1988)).

In another preferred embodiment, the opioid antagonist comprises naltrexone. In the treatment of patients previously addicted to opioids, naltrexone has been used in large oral doses (over 100 mg) to prevent euphorogenic effects of opioid agonists. Naltrexone has been reported to exert strong preferential blocking action against mu over delta sites. Naltrexone is known as a synthetic congener of oxymorphone with no opioid agonist properties, and differs in structure from oxymorphone by the replacement of the methyl group located on the nitrogen atom of oxymorphone with a cyclopropylmethyl group. The hydrochloride salt of naltrexone is soluble in water up to about 100 mg/cc. The pharmacological and pharmacokinetic properties of naltrexone have been evaluated in multiple animal and clinical studies. See, e.g., Gonzalez et al. *Drugs* 35:192-213 (1988). Following oral administration, naltrexone is rapidly absorbed (within 1 hour) and has an oral bioavailability ranging from 5-40%. Naltrexone's protein binding is approximately 21% and the volume of distribution following single-dose administration is 16.1 L/kg.

Naltrexone is commercially available in tablet form (Revia®, DuPont (Wilmington, Del.)) for the treatment of alcohol dependence and for the blockade of exogenously administered opioids. See, e.g., Revia (naltrexone hydrochloride tablets), Physician's Desk Reference, 51<sup>st</sup> ed., Montvale, N.J.; and *Medical Economics* 51:957-959 (1997). A dosage of 50 mg Revia® blocks the pharmacological effects of 25 mg IV administered heroin for up to 24 hours. It is known that, when coadministered with morphine, heroin or other opioids on a chronic basis, naltrexone blocks the development of physical dependence to opioids. It is believed that the method

by which naltrexone blocks the effects of heroin is by competitively binding at the opioid receptors. Naltrexone has been used to treat narcotic addiction by complete blockade of the effects of opioids. It has been found that the most successful use of naltrexone for a narcotic addiction is with narcotic addicts having good prognosis, as part of a comprehensive occupational or rehabilitative program involving behavioral control or other compliance-enhancing methods. For treatment of narcotic dependence with naltrexone, it is desirable that the patient be opioid-free for at least 7-10 days. The initial dosage of naltrexone for such purposes has typically been about 25 mg, and if no withdrawal signs occur, the dosage may be increased to 50 mg per day. A daily dosage of 50 mg is considered to produce adequate clinical blockade of the actions of parenterally administered opioids. Naltrexone also has been used for the treatment of alcoholism as an adjunct with social and psychotherapeutic methods. Other preferred opioid antagonists include, for example, cyclazocine and naltrexone, both of which have cyclopropylmethyl substitutions on the nitrogen, retain much of their efficacy by the oral route, and last longer, with durations approaching 24 hours after oral administration.

The antagonist may also be a bittering agent. The term "bittering agent" as used herein refers to any agent that provides an unpleasant taste to the host upon inhalation and/or swallowing of a tampered dosage form comprising the sequestering subunit. With the inclusion of a bittering agent, the intake of the tampered dosage form produces a bitter taste upon inhalation or oral administration, which, in certain embodiments, spoils or hinders the pleasure of obtaining a high from the tampered dosage form, and preferably prevents the abuse of the dosage form.

Various bittering agents can be employed including, for example, and without limitation, natural, artificial and synthetic flavor oils and flavoring aromatics and/or oils, oleoresins and extracts derived from plants, leaves, flowers, fruits, and so forth, and combinations thereof. Non-limiting representative flavor oils include spearmint oil, peppermint oil, eucalyptus oil, oil of nutmeg, allspice, mace, oil of bitter almonds, menthol and the like. Also useful bittering agents are artificial, natural and synthetic fruit flavors such as citrus oils, including lemon, orange, lime, and grapefruit, fruit essences, and so forth. Additional bittering agents include sucrose derivatives (e.g., sucrose octaacetate), chlorosucrose derivatives, quinine sulphate, and the like. A preferred bittering agent for use in the invention is Denatonium Benzoate NF-Anhydrous, sold under the name Bitrex™ (Macfarlan Smith Limited, Edinburgh, UK). A bittering agent can be added to the formulation in an amount of less than about 50% by weight, preferably less than about 10% by weight, more preferably less than about 5% by weight of the dosage form, and most preferably in an amount ranging from about 0.1 to 1.0 percent by weight of the dosage form, depending on the particular bittering agent(s) used.

Alternatively, the antagonist may be a dye. The term "dye" as used herein refers to any agent that causes discoloration of the tissue in contact. In this regard, if the sequestering subunit is tampered with and the contents are snorted, the dye will discolor the nasal tissues and surrounding tissues thereof. Preferred dyes are those that can bind strongly with subcutaneous tissue proteins and are well-known in the art. Dyes useful in applications ranging from, for example, food coloring to tattooing, are exemplary dyes suitable for the invention. Food coloring dyes include, but are not limited to FD&C Green #3 and FD&C Blue #1, as well as any other FD&C or

D&C color. Such food dyes are commercially available through companies, such as Voigt Global Distribution (Kansas City, Mo.).

The antagonist may alternatively be an irritant. The term “irritant” as used herein includes a compound used to impart an irritating, e.g., burning or uncomfortable, sensation to an abuser administering a tampered dosage form of the invention. Use of an irritant will discourage an abuser from tampering with the dosage form and thereafter inhaling, injecting, or swallowing the tampered dosage form. Preferably, the irritant is released when the dosage form is tampered with and provides a burning or irritating effect to the abuser upon inhalation, injection, and/or swallowing the tampered dosage form. Various irritants can be employed including, for example, and without limitation, capsaicin, a capsaicin analog with similar type properties as capsaicin, and the like. Some capsaicin analogues or derivatives include, for example, and without limitation, resiniferatoxin, tinyatoxin, heptanoylisobutylamide, heptanoyl guaiacylamide, other isobutylamides or guaiacylamides, dihydrocapsaicin, homovanillyl octylester, nonanoyl vanillylamide, or other compounds of the class known as vanilloids. Resiniferatoxin is described, for example, in U.S. Pat. No. 5,290,816. 4,812,446 describes capsaicin analogs and methods for their preparation. Furthermore, U.S. Pat. No. 4,424,205 cites Newman, “Natural and Synthetic Pepper-Flavored Substances,” published in 1954 as listing pungency of capsaicin-like analogs. Ton et al., *British Journal of Pharmacology* 10:175-182 (1955), discusses pharmacological actions of capsaicin and its analogs. With the inclusion of an irritant (e.g., capsaicin) in the dosage form, the irritant imparts a burning or discomforting quality to the abuser to discourage the inhalation, injection, or oral administration of the tampered dosage form, and preferably to prevent the abuse of the dosage form. Suitable capsaicin compositions include capsaicin (trans 8-methyl-N-vanillyl-6-noneamide) or analogues thereof in a concentration between about 0.00125% and 50% by weight, preferably between about 1% and about 7.5% by weight, and most preferably, between about 1% and about 5% by weight.

The antagonist may also be a gelling agent. The term “gelling agent” as used herein refers to any agent that provides a gel-like quality to the tampered dosage form, which slows the absorption of the therapeutic agent, which is formulated with the sequestering subunit, such that a host is less likely to obtain a rapid “high.” In certain preferred embodiments, when the dosage form is tampered with and exposed to a small amount (e.g., less than about 10 ml) of an aqueous liquid (e.g., water), the dosage form will be unsuitable for injection and/or inhalation. Upon the addition of the aqueous liquid, the tampered dosage form preferably becomes thick and viscous, rendering it unsuitable for injection. The term “unsuitable for injection” is defined for purposes of the invention to mean that one would have substantial difficulty injecting the dosage form (e.g., due to pain upon administration or difficulty pushing the dosage form through a syringe) due to the viscosity imparted on the dosage form, thereby reducing the potential for abuse of the therapeutic agent in the dosage form. In certain embodiments, the gelling agent is present in such an amount in the dosage form that attempts at evaporation (by the application of heat) to an aqueous mixture of the dosage form in an effort to produce a higher concentration of the therapeutic agent, produces a highly viscous substance unsuitable for injection. When nasally inhaling the tampered dosage form, the gelling agent can become gel-like upon administration to the nasal passages, due to the moisture of the mucous membranes. This also makes such formulations aversive to nasal administration, as the gel will stick to the

nasal passage and minimize absorption of the abusable substance. Various gelling agents may be employed including, for example, and without limitation, sugars or sugar-derived alcohols, such as mannitol, sorbitol, and the like, starch and starch derivatives, cellulose derivatives, such as microcrystalline cellulose, sodium carboxymethyl cellulose, methylcellulose, ethyl cellulose, hydroxyethyl cellulose, hydroxypropyl cellulose, and hydroxypropyl methylcellulose, attapulgites, bentonites, dextrans, alginates, carrageenan, gum tragacant, gum acacia, guar gum, xanthan gum, pectin, gelatin, kaolin, lecithin, magnesium aluminum silicate, the carbomers and carbopols, polyvinylpyrrolidone, polyethylene glycol, polyethylene oxide, polyvinyl alcohol, silicon dioxide, surfactants, mixed surfactant/wetting agent systems, emulsifiers, other polymeric materials, and mixtures thereof; etc. In certain preferred embodiments, the gelling agent is xanthan gum. In other preferred embodiments, the gelling agent of the invention is pectin. The pectin or pectic substances useful for this invention include not only purified or isolated pectates but also crude natural pectin sources, such as apple, citrus or sugar beet residues, which have been subjected, when necessary, to esterification or de-esterification, e.g., by alkali or enzymes. Preferably, the pectins used in this invention are derived from citrus fruits, such as lime, lemon, grapefruit, and orange. With the inclusion of a gelling agent in the dosage form, the gelling agent preferably imparts a gel-like quality to the dosage form upon tampering that spoils or hinders the pleasure of obtaining a rapid high from due to the gel-like consistency of the tampered dosage form in contact with the mucous membrane, and in certain embodiments, prevents the abuse of the dosage form by minimizing absorption, e.g., in the nasal passages. A gelling agent can be added to the formulation in a ratio of gelling agent to opioid agonist of from about 1:40 to about 40:1 by weight, preferably from about 1:1 to about 30:1 by weight, and more preferably from about 2:1 to about 10:1 by weight of the opioid agonist. In certain other embodiments, the dosage form forms a viscous gel having a viscosity of at least about 10 cP after the dosage form is tampered with by dissolution in an aqueous liquid (from about 0.5 to about 10 ml and preferably from 1 to about 5 ml). Most preferably, the resulting mixture will have a viscosity of at least about 60 cP.

The “blocking agent” prevents or substantially prevents the release of the antagonist in the gastrointestinal tract for a time period that is greater than 24 hours, e.g., between 24 and 25 hours, 30 hours, 35 hours, 40 hours, 45 hours, 48 hours, 50 hours, 55 hours, 60 hours, 65 hours, 70 hours, 72 hours, 75 hours, 80 hours, 85 hours, 90 hours, 95 hours, or 100 hours; etc. Preferably, the time period for which the release of the antagonist is prevented or substantially prevented in the gastrointestinal tract is at least about 48 hours. More preferably, the blocking agent prevents or substantially prevents the release for a time period of at least about 72 hours.

The blocking agent of the present inventive sequestering subunit can be a system comprising a first antagonist-impermeable material and a core. By “antagonist-impermeable material” is meant any material that is substantially impermeable to the antagonist, such that the antagonist is substantially not released from the sequestering subunit. The term “substantially impermeable” as used herein does not necessarily imply complete or 100% impermeability. Rather, there are varying degrees of impermeability of which one of ordinary skill in the art recognizes as having a potential benefit. In this regard, the antagonist-impermeable material substantially prevents or prevents the release of the antagonist to an extent that at least about 80% of the antagonist is prevented from being released from the sequestering subunit in the



gastrointestinal tract for a time period that is greater than 24 hours. Preferably, the antagonist-impermeable material prevents release of at least about 90% of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. More preferably, the antagonist-impermeable material prevents release of at least about 95% of the antagonist from the sequestering subunit. Most preferably, the antagonist-impermeable material prevents release of at least about 99% of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. The antagonist-impermeable material prevents or substantially prevents the release of the antagonist in the gastrointestinal tract for a time period that is greater than 24 hours, and desirably, at least about 48 hours. More desirably, the antagonist-impermeable material prevents or substantially prevents the release of the adverse agent from the sequestering subunit for a time period of at least about 72 hours.

Preferably, the first antagonist-impermeable material comprises a hydrophobic material, such that the antagonist is not released or substantially not released during its transit through the gastrointestinal tract when administered orally as intended, without having been tampered with. Suitable hydrophobic materials for use in the invention are described herein and set forth below. The hydrophobic material is preferably a pharmaceutically acceptable hydrophobic material.

It is also preferred that the first antagonist-impermeable material comprises a polymer insoluble in the gastrointestinal tract. One of ordinary skill in the art appreciates that a polymer that is insoluble in the gastrointestinal tract will prevent the release of the antagonist upon ingestion of the sequestering subunit. The polymer may be a cellulose or an acrylic polymer. Desirably, the cellulose is selected from the group consisting of ethylcellulose, cellulose acetate, cellulose propionate, cellulose acetate propionate, cellulose acetate butyrate, cellulose acetate phthalate, cellulose triacetate, and combinations thereof. Ethylcellulose includes, for example, one that has an ethoxy content of about 44 to about 55%. Ethylcellulose can be used in the form of an aqueous dispersion, an alcoholic solution, or a solution in other suitable solvents. The cellulose can have a degree of substitution (D.S.) on the anhydroglucose unit, from greater than zero and up to 3 inclusive. By "degree of substitution" is meant the average number of hydroxyl groups on the anhydroglucose unit of the cellulose polymer that are replaced by a substituting group. Representative materials include a polymer selected from the group consisting of cellulose acylate, cellulose diacylate, cellulose triacylate, cellulose acetate, cellulose diacetate, cellulose triacetate, monocellulose alkanylate, dicellulose alkanylate, tricellulose alkanylate, monocellulose alkenylates, dicellulose alkenylates, tricellulose alkenylates, monocellulose aroylates, dicellulose aroylates, and tricellulose aroylates.

More specific celluloses include cellulose propionate having a D.S. of 1.8 and a propyl content of 39.2 to 45 and a hydroxy content of 2.8 to 5.4%; cellulose acetate butyrate having a D.S. of 1.8, an acetyl content of 13 to 15% and a butyryl content of 34 to 39%; cellulose acetate butyrate having an acetyl content of 2 to 29%, a butyryl content of 17 to 53% and a hydroxy content of 0.5 to 4.7%; cellulose triacylate having a D.S. of 2.9 to 3, such as cellulose triacetate, cellulose trivalerate, cellulose trilaurate, cellulose tripatmitate, cellulose trisuccinate, and cellulose trioctanoate; cellulose diacylates having a D.S. of 2.2 to 2.6, such as cellulose disuccinate, cellulose dipalmitate, cellulose dioctanoate, cellulose dipentanoate, and coesters of cellulose, such as cellulose acetate butyrate, cellulose acetate octanoate butyrate, and cellulose

acetate propionate. Additional cellulose polymers that may be used to prepare the sequestering subunit include acetaldehyde dimethyl cellulose acetate, cellulose acetate ethylcarbamate, cellulose acetate methycarbamate, and cellulose acetate dimethylaminocellulose acetate.

The acrylic polymer preferably is selected from the group consisting of methacrylic polymers, acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, poly(acrylic acid), poly(methacrylic acid), methacrylic acid alkylamide copolymer, poly(methyl methacrylate), polymethacrylate, poly(methyl methacrylate) copolymer, polyacrylamide, aminoalkyl methacrylate copolymer, poly(methacrylic acid anhydride), glycidyl methacrylate copolymers, and combinations thereof. An acrylic polymer useful for preparation of a sequestering subunit of the invention includes acrylic resins comprising copolymers synthesized from acrylic and methacrylic acid esters (e.g., the copolymer of acrylic acid lower alkyl ester and methacrylic acid lower alkyl ester) containing about 0.02 to about 0.03 mole of a tri(lower alkyl) ammonium group per mole of the acrylic and methacrylic monomer used. An example of a suitable acrylic resin is ammonio methacrylate copolymer NF21, a polymer manufactured by Rohm Pharma GmbH, Darmstadt, Germany, and sold under the Eudragit® trademark. Eudragit® is a water-insoluble copolymer of ethyl acrylate (EA), methyl methacrylate (MM) and trimethylammoniummethyl methacrylate chloride (TAM) in which the molar ratio of TAM to the remaining components (EA and MM) is 1:40. Acrylic resins, such as Eudragit®, can be used in the form of an aqueous dispersion or as a solution in suitable solvents. Preferred acrylic polymers include copolymers of acrylic and methacrylic acid esters with a low content in quaternary ammonium groups such as Eudragit® RL PO (Type A) and Eudragit® RS PO (Type B; as used herein, "Eudragit® RS") (as described the monographs Ammonio Methacrylate Copolymer Type A Ph. Eur., Ammonio Methacrylate Copolymer Type B Ph. Eur., Ammonio Methacrylate Copolymer, Type A and B USP/NF, and Aminoalkylmethacrylate Copolymer RS JPE).

In another preferred embodiment, the antagonist-impermeable material is selected from the group consisting of polylactic acid, polyglycolic acid, a co-polymer of polylactic acid and polyglycolic acid, and combinations thereof. In certain other embodiments, the hydrophobic material includes a biodegradable polymer comprising a poly(lactic/glycolic acid) ("PLGA"), a polylactide, a polyglycolide, a polyanhydride, a polyorthoester, polycaprolactones, polyphosphazenes, polysaccharides, proteinaceous polymers, polyesters, polydioxanone, polygluconate, polylactic-acid-polyethylene oxide copolymers, poly(hydroxybutyrate), polyphosphoester or combinations thereof. Preferably, the biodegradable polymer comprises a poly(lactic/glycolic acid), a copolymer of lactic and glycolic acid, having a molecular weight of about 2,000 to about 500,000 daltons. The ratio of lactic acid to glycolic acid is preferably from about 100:1 to about 25:75, with the ratio of lactic acid to glycolic acid of about 65:35 being more preferred.

Poly(lactic/glycolic acid) can be prepared by the procedures set forth in U.S. Pat. No. 4,293,539 (Ludwig et al.), which is incorporated herein by reference. In brief, Ludwig prepares the copolymer by condensation of lactic acid and glycolic acid in the presence of a readily removable polymerization catalyst (e.g., a strong ion-exchange resin such as Dowex HCR-W2-H). The amount of catalyst is not critical to the polymerization, but typically is from about 0.01 to about 20 parts by weight relative to the total weight of combined

lactic acid and glycolic acid. The polymerization reaction can be conducted without solvents at a temperature from about 100° C. to about 250° C. for about 48 to about 96 hours, preferably under a reduced pressure to facilitate removal of water and by-products. Poly(lactic/glycolic acid) is then recovered by filtering the molten reaction mixture in an organic solvent, such as dichloromethane or acetone, and then filtering to remove the catalyst.

Suitable plasticizers for use in the sequestering subunit include, for example, acetyl triethyl citrate, acetyl tributyl citrate, triethyl citrate, diethyl phthalate, dibutyl phthalate (DBP), acetyltri-N-butyl citrate (ATBC), or dibutyl sebacate, which can be admixed with the polymer. Other additives such as coloring agents may also be used in making the present inventive sequestering subunit.

In certain embodiments, additives may be included in the compositions that improve the sequestering characteristics of the sequestering subunit. As described below, the ratio of additives or components with respect to other additives or components may be modified to enhance or delay improve sequestration of the agent contained within the subunit. Various amounts of a functional additive (i.e., a charge-neutralizing additive) may be included to vary the release of an antagonist, particularly where a water-soluble core (i.e., a sugar sphere) is utilized. For instance, it has been determined that the inclusion of a low amount of charge-neutralizing additive relative to sequestering polymer on a weight-by-weight basis may cause decreased release of the antagonist.

In certain embodiments, a surfactant may serve as a charge-neutralizing additive. Such neutralization may in certain embodiments reduce the swelling of the sequestering polymer by hydration of positively charged groups contained therein. Surfactants (ionic or non-ionic) may also be used in preparing the sequestering subunit. It is preferred that the surfactant be ionic. Suitable exemplary agents include, for example, alkylaryl sulphonates, alcohol sulphates, sulphosuccinates, sulphosuccinamates, sarcosinates or taurates and others. Additional examples include but are not limited to ethoxylated castor oil, benzalkonium chloride, polyglycolized glycerides, acetylated monoglycerides, sorbitan fatty acid esters, poloxamers, polyoxyethylene fatty acid esters, polyoxyethylene derivatives, monoglycerides or ethoxylated derivatives thereof, diglycerides or polyoxyethylene derivatives thereof, sodium docusate, sodium lauryl sulfate, dioctyl sodium sulphosuccinate, sodium lauryl sarcosinate and sodium methyl cocoyl taurate, magnesium lauryl sulfate, triethanolamine, cetrimide, sucrose laurate and other sucrose esters, glucose (dextrose) esters, simethicone, ocoxynol, dioctyl sodiumsulfo succinate, polyglycolized glycerides, sodiumdodecylbenzene sulfonate, dialkyl sodiumsulfo succinate, fatty alcohols such as lauryl, cetyl, and steryl, glycerylestes, cholic acid or derivatives thereof, lecithins, and phospholipids. These agents are typically characterized as ionic (i.e., anionic or cationic) or nonionic. In certain embodiments described herein, an anionic surfactant such as sodium lauryl sulfate (SLS) is preferably used (U.S. Pat. Nos. 5,725,883; 7,201,920; EP 502642A1; Shokri, et al. Pharm. Sci. 2003. *The effect of sodium lauryl sulphate on the release of diazepam from solid dispersions prepared by cogrinding technique.* Wells, et al. *Effect of Anionic Surfactants on the Release of Chlorpheniramine Maleate From an Inert, Heterogeneous Matrix.* Drug Development and Industrial Pharmacy 18(2) (1992): 175-186. Rao, et al. "Effect of Sodium Lauryl Sulfate on the Release of Rifampicin from Guar Gum Matrix." Indian Journal of Pharmaceutical Science (2000): 404-406; Knop, et al. *Influence of surfactants of different charge and concentration on drug release from pellets coated*

*with an aqueous dispersion of quaternary acrylic polymers.* STP Pharma Sciences, Vol. 7, No. 6, (1997) 507-512). Other suitable agents are known in the art.

As shown herein, SLS is particularly useful in combination with Eudragit RS when the sequestering subunit is built upon a sugar sphere substrate. The inclusion of SLS at less than approximately 6.3% on a weight-to-weight basis relative to the sequestering polymer (i.e., Eudragit RS) may provide a charge neutralizing function (theoretically 20% and 41% neutralization, respectfully), and thereby significantly slow the release of the active agent encapsulated thereby (i.e., the antagonist naltrexone). Inclusion of more than approximately 6.3% SLS relative to the sequestering polymer appears to increase release of the antagonist from the sequestering subunit. With respect to SLS used in conjunction with Eudragit® RS, it is preferred that the SLS is present at approximately 1%, 2%, 3%, 4% or 5%, and typically less than 6% on a w/w basis relative to the sequestering polymer (i.e., Eudragit® RS). In preferred embodiments, SLS may be present at approximately 1.6% or approximately 3.3% relative to the sequestering polymer. As discussed above, many agents (i.e., surfactants) may substitute for SLS in the compositions disclosed herein.

Additionally useful agents include those that may physically block migration of the antagonist from the subunit and/or enhance the hydrophobicity of the barrier. One exemplary agent is talc, which is commonly used in pharmaceutical compositions (Pawar et al. *Agglomeration of Ibuprofen With Talc by Novel Crystallo-Co-Agglomeration Technique.* AAPS PharmSciTech. 2004; 5(4): article 55). As shown in the Examples, talc is especially useful where the sequestering subunit is built upon a sugar sphere core. Any form of talc may be used, so long as it does not detrimentally affect the function of the composition. Most talc results from the alteration of dolomite (CaMg(CO<sub>3</sub>)<sub>2</sub> or magnesite (MgO) in the presence of excess dissolved silica (SiO<sub>2</sub>) or by altering serpentine or quartzite. Talc may include minerals such as tremolite (CaMg<sub>3</sub>(SiO<sub>3</sub>)<sub>4</sub>), serpentine (3MgO.2SiO<sub>2</sub>.2H<sub>2</sub>O), anthophyllite (Mg<sub>7</sub>(OH)<sub>2</sub>(Si<sub>4</sub>O<sub>11</sub>)<sub>2</sub>), magnesite, mica, chlorite, dolomite, the calcite form of calcium carbonate (CaCO<sub>3</sub>), iron oxide, carbon, quartz, and/or manganese oxide. The presence of such impurities may be acceptable in the compositions described herein provided the function of the talc is maintained. It is preferred that that talc be USP grade. As mentioned above, the function of talc as described herein is to enhance the hydrophobicity and therefore the functionality of the sequestering polymer. Many substitutes for talc may be utilized in the compositions described herein as may be determined by one of skill in the art.

It has been determined that the ratio of talc to sequestering polymer may make a dramatic difference in the functionality of the compositions described herein. For instance, the Examples described below demonstrate that the talc to sequestering polymer ratio (w/w) is important with respect to compositions designed to prevent the release of naltrexone therefrom. It is shown therein that inclusion of an approximately equivalent amount (on a weight-by-weight basis) of talc and Eudragit® RS results in a very low naltrexone release profile. In contrast, significantly lower or higher both a lower (69% w/w) and a higher (151% w/w) talc:Eudragit® RS ratios result in increased release of naltrexone release. Thus, where talc and Eudragit® RS are utilized, it is preferred that talc is present at approximately 75%, 80%, 85%, 90%, 95%, 100%, 105%, 110%, 115%, 120% or 125% w/w relative to Eudragit® RS. As described above, the most beneficial ratio for other additives or components will vary and may be determined using standard experimental procedures.

In certain embodiments, such as where a water-soluble core is utilized, it is useful to include agents that may affect the osmotic pressure of the composition (i.e., an osmotic pressure regulating agent) (see, in general, WO 2005/046561 A2 and WO 2005/046649 A2 relating to Eudragit®). This agent is preferably applied to the Eudragit® RS/talc layer described above. In a pharmaceutical unit comprising a sequestering subunit overlaid by an active agent (i.e., a controlled-release agonist preparation), the osmotic pressure regulating agent is preferably positioned immediately beneath the active agent layer. Suitable osmotic pressure regulating agents may include, for instance, hydroxypropyl-methyl cellulose (HPMC) or chloride ions (i.e., from NaCl), or a combination of HPMC and chloride ions (i.e., from NaCl). Other ions that may be useful include bromide or iodide. The combination of sodium chloride and HPMC may be prepared in water or in a mixture of ethanol and water, for instance. HPMC is commonly utilized in pharmaceutical compositions (see, for example, U.S. Pat. Nos. 7,226,620 and 7,229,982). In certain embodiments, HPMC may have a molecular weight ranging from about 10,000 to about 1,500,000, and typically from about 5000 to about 10,000 (low molecular weight HPMC). The specific gravity of HPMC is typically from about 1.19 to about 1.31, with an average specific gravity of about 1.26 and a viscosity of about 3600 to 5600. HPMC may be a water-soluble synthetic polymer. Examples of suitable, commercially available hydroxypropyl methylcellulose polymers include Methocel K100 LV and Methocel K4M (Dow). Other HPMC additives are known in the art and may be suitable in preparing the compositions described herein. As shown in the Examples, the inclusion of NaCl (with HPMC) was found to have positively affect sequestration of naltrexone by Eudragit® RS. In certain embodiments, it is preferred that the charge-neutralizing additive (i.e., NaCl) is included at less than approximately 1, 2, 3, 4, 5, 6, 7, 8, 9, or 10% on a weight-by-weight basis with respect to the sequestering polymer. In other preferred embodiments, the charge-neutralizing additive is present at approximately 4% on a weight-by-weight basis with respect to the sequestering polymer.

Thus, in one embodiment, a sequestering subunit built upon a sugar sphere substrate is provided comprising a sequestering polymer (i.e., Eudragit® RS) in combination with several optimizing agents, including sodium lauryl sulfate (SLS) as a charge-neutralizing agent to reduce swelling of the film by hydration of the positively charged groups on the polymer; talc to create a solid impermeable obstacle to naltrexone transport through the film and as a hydrophobicity-enhancing agent; and a chloride ion (i.e., as NaCl) as an osmotic pressure reducing agent. The ratio of each of the additional ingredients relative to the sequestering polymer was surprisingly found to be important to the function of the sequestering subunit. For instance, the Examples provide a sequestering subunit including a sequestering polymer and the optimizing agents SLS at less than 6%, preferably 1-4%, and even more preferably 1.6% or 3.3% on a w/w basis relative to Eudragit RS; talc in an amount approximately equal to Eudragit® RS (on a w/w basis); and, NaCl present at approximately 4% on a w/w basis relative to Eudragit® RS.

Methods of making any of the sequestering subunits of the invention are known in the art. See, for example, *Remington. The Science and Practice of Pharmacy*, Alfonso R. Genaro (ed), 20<sup>th</sup> edition, and Example 2 set forth below. The sequestering subunits can be prepared by any suitable method to provide, for example, beads, pellets, granules, spheroids, and the like. Spheroids or beads, coated with an active ingredient can be prepared, for example, by dissolving the active ingre-

dient in water and then spraying the solution onto a substrate, for example, nu pariel 18/20 beads, using a Wurster insert. Optionally, additional ingredients are also added prior to coating the beads in order to assist the active ingredient in binding to the substrates, and/or to color the solution; etc. The resulting substrate-active material optionally can be over-coated with a barrier material to separate the therapeutically active agent from the next coat of material, e.g., release-retarding or sequestering material. Preferably, the barrier material is a material comprising hydroxypropyl methylcellulose. However, any film-former known in the art can be used. Preferably, the barrier material does not affect the dissolution rate of the final product.

Pellets comprising an active ingredient can be prepared, for example, by a melt pelletization technique. Typical of such techniques is when the active ingredient in finely divided form is combined with a binder (also in particulate form) and other optional inert ingredients, and thereafter the mixture is pelletized, e.g., by mechanically working the mixture in a high shear mixer to form the pellets (e.g., pellets, granules, spheres, beads; etc., collectively referred to herein as “pellets”). Thereafter, the pellets can be sieved in order to obtain pellets of the requisite size. The binder material is preferably in particulate form and has a melting point above about 40° C. Suitable binder substances include, for example, hydrogenated castor oil, hydrogenated vegetable oil, other hydrogenated fats, fatty alcohols, fatty acid esters, fatty acid glycerides, and the like.

The diameter of the extruder aperture or exit port also can be adjusted to vary the thickness of the extruded strands. Furthermore, the exit part of the extruder need not be round; it can be oblong, rectangular; etc. The exiting strands can be reduced to particles using a hot wire cutter, guillotine; etc.

The melt-extruded multiparticulate system can be, for example, in the form of granules, spheroids, pellets, or the like, depending upon the extruder exit orifice. The terms “melt-extruded multiparticulate(s)” and “melt-extruded multiparticulate system(s)” and “melt-extruded particles” are used interchangeably herein and include a plurality of sub-units, preferably within a range of similar size and/or shape. The melt-extruded multiparticulates are preferably in a range of from about 0.1 to about 12 mm in length and have a diameter of from about 0.1 to about 5 mm. In addition, the melt-extruded multiparticulates can be any geometrical shape within this size range. Alternatively, the extrudate can simply be cut into desired lengths and divided into unit doses of the therapeutically active agent without the need of a spheronization step.

The substrate also can be prepared via a granulation technique. Generally, melt-granulation techniques involve melting a normally solid hydrophobic material, e.g., a wax, and incorporating an active ingredient therein. To obtain a sustained-release dosage form, it can be necessary to incorporate an additional hydrophobic material.

A coating composition can be applied onto a substrate by spraying it onto the substrate using any suitable spray equipment. For example, a Wurster fluidized-bed system can be used in which an air flow from underneath, fluidizes the coated material and effects drying, while the insoluble polymer coating is sprayed on. The thickness of the coating will depend on the characteristics of the particular coating composition, and can be determined by using routine experimentation.

Any manner of preparing a subunit can be employed. By way of example, a subunit in the form of a pellet or the like can be prepared by co-extruding a material comprising the opioid agonist and a material comprising the opioid antagonist and/

or antagonist in sequestered form. Optionally, the opioid agonist composition can cover, e.g., overcoat, the material comprising the antagonist and/or antagonist in sequestered form. A bead, for example, can be prepared by coating a substrate comprising an opioid antagonist and/or an antagonist in sequestered form with a solution comprising an opioid agonist.

The sequestering subunits of the invention are particularly well-suited for use in compositions comprising the sequestering subunit and a therapeutic agent in releasable form. In this regard, the invention also provides a composition comprising any of the sequestering subunits of the invention and a therapeutic agent in releasable form. By "releasable form" is meant to include immediate release, intermediate release, and sustained-release forms. The therapeutic agent can be formulated to provide immediate release of the therapeutic agent. In preferred embodiments, the composition provides sustained-release of the therapeutic agent.

The therapeutic agent applied upon the sequestering subunit may be any medicament. The therapeutic agent of the present inventive compositions can be any medicinal agent used for the treatment of a condition or disease, a pharmaceutically acceptable salt thereof, or an analogue of either of the foregoing. The therapeutic agent can be, for example, an analgesic (e.g., an opioid agonist, aspirin, acetaminophen, non-steroidal anti-inflammatory drugs ("NSAIDS"), N-methyl-D-aspartate ("NMDA") receptor antagonists, cyclooxygenase-II inhibitors ("COX-II inhibitors"), and glycine receptor antagonists), an antibacterial agent, an anti-viral agent, an anti-microbial agent, anti-infective agent, a chemotherapeutic, an immunosuppressant agent, an antitussive, an expectorant, a decongestant, an antihistamine drugs, a decongestant, antihistamine drugs, and the like. Preferably, the therapeutic agent is one that is addictive (physically and/or psychologically) upon repeated use and typically leads to abuse of the therapeutic agent. In this regard, the therapeutic agent can be any opioid agonist as discussed herein.

The therapeutic agent can be an opioid agonist. By "opioid" is meant to include a drug, hormone, or other chemical or biological substance, natural or synthetic, having a sedative, narcotic, or otherwise similar effect(s) to those containing opium or its natural or synthetic derivatives. By "opioid agonist," sometimes used herein interchangeably with terms "opioid" and "opioid analgesic," is meant to include one or more opioid agonists, either alone or in combination, and is further meant to include the base of the opioid, mixed or combined agonist-antagonists, partial agonists, pharmaceutically acceptable salts thereof, stereoisomers thereof, ethers thereof, esters thereof, and combinations thereof.

Opioid agonists include, for example, alfentanil, allylpropidine, alphaprodine, anileridine, benzylmorphine, bezitramide, buprenorphine, butorphanol, clonitazene, codeine, cyclazocine, desomorphine, dextromoramide, dezocine, diampromide, dihydrocodeine, dihydroetorphine, dihydromorphine, dimenoxadol, dimephtanol, dimethylthiambutene, dioxaphetyl butyrate, dipipanone, eptazocine, ethoheptazine, ethylmethylthiambutene, ethylmorphine, etonitazene, etorphine, fentanyl, heroin, hydrocodone, hydromorphone, hydroxypethidine, isomethadone, ketobemidone, levallorphan, levorphanol, levophenacymorphan, lofentanil, meperidine, meptazinol, metazocine, methadone, metopon, morphine, myrophine, nalbuphine, narceine, nicomorphine, norlevorphanol, normethadone, nalorphine, normorphine, norpipanone, opium, oxycodone, oxymorphone, papavereturn, pentazocine, phenadoxone, phenazocine, phenomorphan, phenoperidine, piminodine, piritramide, proph-

eptazine, promedol, properidine, propiram, propoxyphene, sufentanil, tramadol, tilidine, derivatives or complexes thereof, pharmaceutically acceptable salts thereof, and combinations thereof. Preferably, the opioid agonist is selected from the group consisting of hydrocodone, hydromorphone, oxycodone, dihydrocodeine, codeine, dihydromorphine, morphine, buprenorphine, derivatives or complexes thereof, pharmaceutically acceptable salts thereof, and combinations thereof. Most preferably, the opioid agonist is morphine, hydromorphone, oxycodone or hydrocodone. In a preferred embodiment, the opioid agonist comprises oxycodone or hydrocodone and is present in the dosage form in an amount of about 15 to about 45 mg, and the opioid antagonist comprises naltrexone and is present in the dosage form in an amount of about 0.5 to about 5 mg.

Equianalgesic doses of these opioids, in comparison to a 15 mg dose of hydrocodone, are set forth in Table 1 below:

TABLE I

Equianalgesic Doses of Opioids	
Opioid	Calculated Dose (mg)
Oxycodone	13.5
Codeine	90.0
Hydrocodone	15.0
Hydromorphone	3.375
Levorphanol	1.8
Meperidine	135.0
Methadone	9.0
Morphine	27.0

Hydrocodone is a semisynthetic narcotic analgesic and antitussive with multiple nervous system and gastrointestinal actions. Chemically, hydrocodone is 4,5-epoxy-3-methoxy-17-methylmorphinan-6-one, and is also known as dihydrocodeinone. Like other opioids, hydrocodone can be habit-forming and can produce drug dependence of the morphine type. Like other opium derivatives, excess doses of hydrocodone will depress respiration.

Oral hydrocodone is also available in Europe (e.g., Belgium, Germany, Greece, Italy, Luxembourg, Norway and Switzerland) as an antitussive agent. A parenteral formulation is also available in Germany as an antitussive agent. For use as an analgesic, hydrocodone bitartrate is commonly available in the United States only as a fixed combination with non-opiate drugs (e.g., ibuprofen, acetaminophen, aspirin; etc.) for relief of moderate to moderately severe pain.

A common dosage form of hydrocodone is in combination with acetaminophen and is commercially available, for example, as Lortab® in the United States from UCB Pharma, Inc. (Brussels, Belgium), as 2.5/500 mg, 5/500 mg, 7.5/500 mg and 10/500 mg hydrocodone/acetaminophen tablets. Tablets are also available in the ratio of 7.5 mg hydrocodone bitartrate and 650 mg acetaminophen and a 7.5 mg hydrocodone bitartrate and 750 mg acetaminophen. Hydrocodone, in combination with aspirin, is given in an oral dosage form to adults generally in 1-2 tablets every 4-6 hours as needed to alleviate pain. The tablet form is 5 mg hydrocodone bitartrate and 224 mg aspirin with 32 mg caffeine; or 5 mg hydrocodone bitartrate and 500 mg aspirin. Another formulation comprises hydrocodone bitartrate and ibuprofen. Vicoprofen®, commercially available in the U.S. from Knoll Laboratories (Mount Olive, N.J.), is a tablet containing 7.5 mg hydrocodone bitartrate and 200 mg ibuprofen. The invention is contemplated to encompass all such formulations, with the inclusion of the opioid antagonist and/or antagonist in sequestered form as part of a subunit comprising an opioid agonist.

Oxycodone, chemically known as 4,5-epoxy-14-hydroxy-3-methoxy-17-methylmorphinan-6-one, is an opioid agonist whose principal therapeutic action is analgesia. Other therapeutic effects of oxycodone include anxiolysis, euphoria and feelings of relaxation. The precise mechanism of its analgesic action is not known, but specific CNS opioid receptors for endogenous compounds with opioid-like activity have been identified throughout the brain and spinal cord and play a role in the analgesic effects of this drug. Oxycodone is commercially available in the United States, e.g., as Oxycotin® from Purdue Pharma L.P. (Stamford, Conn.), as controlled-release tablets for oral administration containing 10 mg, 20 mg, 40 mg or 80 mg oxycodone hydrochloride, and as OxyIR™, also from Purdue Pharma L.P., as immediate-release capsules containing 5 mg oxycodone hydrochloride. The invention is contemplated to encompass all such formulations, with the inclusion of an opioid antagonist and/or antagonist in sequestered form as part of a subunit comprising an opioid agonist.

Oral hydromorphone is commercially available in the United States, e.g., as Dilaudid® from Abbott Laboratories (Chicago, Ill.). Oral morphine is commercially available in the United States, e.g., as Kadian® from Faulding Laboratories (Piscataway, N.J.).

Exemplary NSAIDS include ibuprofen, diclofenac, naproxen, benoxaprofen, flurbiprofen, fenoprofen, flubufen, ketoprofen, indoprofen, piroprofen, carprofen, oxaprozin, pramoprofen, muprofen, trioxaprofen, suprofen, aminoprofen, tiaprofenic acid, fluprofen, bucloxic acid, indomethacin, sulindac, tolmetin, zomepirac, tiopinac, zidometacin, acemetacin, fentiazac, clidanac, oxpinac, mefenamic acid, meclofenamic acid, flufenamic acid, niflumic acid, tolfenamic acid, diflurisal, flufenisal, piroxicam, sudoxicam or isoxicam, and the like. Useful dosages of these drugs are well-known.

Exemplary NMDA receptor medicaments include morphinans, such as dextromethorphan or dextrophan, ketamine, d-methadone, and pharmaceutically acceptable salts thereof, and encompass drugs that block a major intracellular consequence of NMDA-receptor activation, e.g., a ganglioside, such as (6-aminothexyl)-5-chloro-1-naphthalenesulfonamide. These drugs are stated to inhibit the development of tolerance to and/or dependence on addictive drugs, e.g., narcotic analgesics, such as morphine, codeine; etc., in U.S. Pat. Nos. 5,321,012 and 5,556,838 (both to Mayer et al.), both of which are incorporated herein by reference, and to treat chronic pain in U.S. Pat. No. 5,502,058 (Mayer et al.), incorporated herein by reference. The NMDA agonist can be included alone or in combination with a local anesthetic, such as lidocaine, as described in these patents by Mayer et al.

COX-2 inhibitors have been reported in the art, and many chemical compounds are known to produce inhibition of cyclooxygenase-2. COX-2 inhibitors are described, for example, in U.S. Pat. Nos. 5,616,601; 5,604,260; 5,593,994; 5,550,142; 5,536,752; 5,521,213; 5,475,995; 5,639,780; 5,604,253; 5,552,422; 5,510,368; 5,436,265; 5,409,944 and 5,130,311, all of which are incorporated herein by reference. Certain preferred COX-2 inhibitors include celecoxib (SC-58635), DUP-697, flosulide (CGP-28238), meloxicam, 6-methoxy-2-naphthylacetic acid (6-NMA), MK-966 (also known as Vioxx), nabumetone (prodrug for 6-MNA), nimesulide, NS-398, SC-5766, SC-58215, T-614, or combinations thereof. Dosage levels of COX-2 inhibitor on the order of from about 0.005 mg to about 140 mg per kilogram of body weight per day have been shown to be therapeutically effective in combination with an opioid analgesic. Alternatively,

about 0.25 mg to about 7 g per patient per day of a COX-2 inhibitor can be administered in combination with an opioid analgesic.

The treatment of chronic pain via the use of glycine receptor antagonists and the identification of such drugs is described in U.S. Pat. No. 5,514,680 (Weber et al.), which is incorporated herein by reference.

In embodiments in which the opioid agonist comprises hydrocodone, the sustained-release oral dosage forms can include analgesic doses from about 8 mg to about 50 mg of hydrocodone per dosage unit. In sustained-release oral dosage forms where hydromorphone is the therapeutically active opioid, it is included in an amount from about 2 mg to about 64 mg hydromorphone hydrochloride. In another embodiment, the opioid agonist comprises morphine, and the sustained-release oral dosage forms of the invention include from about 2.5 mg to about 800 mg morphine, by weight. In yet another embodiment, the opioid agonist comprises oxycodone and the sustained-release oral dosage forms include from about 2.5 mg to about 800 mg oxycodone. In certain preferred embodiments, the sustained-release oral dosage forms include from about 20 mg to about 30 mg oxycodone. Controlled release oxycodone formulations are known in the art. The following documents describe various controlled-release oxycodone formulations suitable for use in the invention described herein, and processes for their manufacture: U.S. Pat. Nos. 5,266,331; 5,549,912; 5,508,042; and 5,656,295, which are incorporated herein by reference. The opioid agonist can comprise tramadol and the sustained-release oral dosage forms can include from about 25 mg to 800 mg tramadol per dosage unit.

The therapeutic agent in sustained-release form is preferably a particle of therapeutic agent that is combined with a release-retarding or sequestering material. The release-retarding or sequestering material is preferably a material that permits release of the therapeutic agent at a sustained rate in an aqueous medium. The release-retarding or sequestering material can be selectively chosen so as to achieve, in combination, with the other stated properties, a desired in vitro release rate.

In a preferred embodiment, the oral dosage form of the invention can be formulated to provide for an increased duration of therapeutic action allowing once-daily dosing. In general, a release-retarding or sequestering material is used to provide the increased duration of therapeutic action. Preferably, the once-daily dosing is provided by the dosage forms and methods described in U.S. patent application Ser. No. 10/667,570 to Boehm, entitled "Sustained-Release Opioid Formulations and Method of Use," filed on Sep. 22, 2003, and incorporated herein by reference.

Preferred release-retarding or sequestering materials include acrylic polymers, alkylcelluloses, shellac, zein, hydrogenated vegetable oil, hydrogenated castor oil, and combinations thereof. In certain preferred embodiments, the release-retarding or sequestering material is a pharmaceutically acceptable acrylic polymer, including acrylic acid and methacrylic acid copolymers, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, aminoalkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), methacrylic acid alkylamide copolymer, poly(methyl methacrylate), poly(methacrylic acid anhydride), methyl methacrylate, polymethacrylate, poly(methyl methacrylate) copolymer, polyacrylamide, aminoalkyl methacrylate copolymer, and glycidyl methacrylate copolymers. In certain preferred embodiments, the acrylic polymer comprises one or more ammonio methacrylate copolymers. Ammonio methacrylate copolymers are well-known in the

art, and are described in NF21, the 21<sup>st</sup> edition of the National Formulary, published by the United States Pharmacopeial Convention Inc. (Rockville, Md.), as fully polymerized copolymers of acrylic and methacrylic acid esters with a low content of quaternary ammonium groups. In other preferred 5 embodiments, the release-retarding or sequestering material is an alkyl cellulosic material, such as ethylcellulose. Those skilled in the art will appreciate that other cellulosic polymers, including other alkyl cellulosic polymers, can be substituted for part or all of the ethylcellulose.

Release-modifying agents, which affect the release properties of the release-retarding or sequestering material, also can be used. In a preferred embodiment, the release-modifying agent functions as a pore-former. The pore-former can be organic or inorganic, and include materials that can be dissolved, extracted or leached from the coating in the environment of use. The pore-former can comprise one or more hydrophilic polymers, such as hydroxypropylmethylcellulose. In certain preferred embodiments, the release-modifying agent is selected from hydroxypropylmethylcellulose, 10 lactose, metal stearates, and combinations thereof.

The release-retarding or sequestering material can also include an erosion-promoting agent, such as starch and gums; a release-modifying agent useful for making microporous lamina in the environment of use, such as polycarbonates 15 comprised of linear polyesters of carbonic acid in which carbonate groups reoccur in the polymer chain; and/or a semi-permeable polymer.

The release-retarding or sequestering material can also include an exit means comprising at least one passageway, orifice, or the like. The passageway can be formed by such methods as those disclosed in U.S. Pat. Nos. 3,845,770; 3,916,889; 4,063,064; and 4,088,864, which are incorporated herein by reference. The passageway can have any shape, such as round, triangular, square, elliptical, irregular; etc. 20

In certain embodiments, the therapeutic agent in sustained-release form can include a plurality of substrates comprising the active ingredient, which substrates are coated with a sustained-release coating comprising a release-retarding or sequestering material. 25

The sustained-release preparations of the invention can be made in conjunction with any multiparticulate system, such as beads, ion-exchange resin beads, spheroids, microspheres, seeds, pellets, granules, and other multiparticulate systems in order to obtain a desired sustained-release of the therapeutic agent. The multiparticulate system can be presented in a capsule or in any other suitable unit dosage form.

In certain preferred embodiments, more than one multiparticulate system can be used, each exhibiting different characteristics, such as pH dependence of release, time for release in various media (e.g., acid, base, simulated intestinal fluid), release in vivo, size and composition. 30

To obtain a sustained-release of the therapeutic agent in a manner sufficient to provide a therapeutic effect for the sustained durations, the therapeutic agent can be coated with an amount of release-retarding or sequestering material sufficient to obtain a weight gain level from about 2 to about 30%, although the coat can be greater or lesser depending upon the physical properties of the particular therapeutic agent utilized and the desired release rate, among other things. Moreover, 35 there can be more than one release-retarding or sequestering material used in the coat, as well as various other pharmaceutical excipients.

Solvents typically used for the release-retarding or sequestering material include pharmaceutically acceptable solvents, such as water, methanol, ethanol, methylene chloride and combinations thereof. 40

In certain embodiments of the invention, the release-retarding or sequestering material is in the form of a coating comprising an aqueous dispersion of a hydrophobic polymer. The inclusion of an effective amount of a plasticizer in the aqueous dispersion of hydrophobic polymer will further 5 improve the physical properties of the film. For example, because ethylcellulose has a relatively high glass transition temperature and does not form flexible films under normal coating conditions, it is necessary to plasticize the ethylcellulose before using the same as a coating material. Generally, 10 the amount of plasticizer included in a coating solution is based on the concentration of the film-former, e.g., most often from about 1 to about 50 percent by weight of the film-former. Concentrations of the plasticizer, however, can be determined by routine experimentation. 15

Examples of plasticizers for ethylcellulose and other celluloses include dibutyl sebacate, diethyl phthalate, triethyl citrate, tributyl citrate, and triacetin, although it is possible that other plasticizers (such as acetylated monoglycerides, phthalate esters, castor oil; etc.) can be used. A plasticizer that is not leached into the aqueous phase such as DBS is preferred. 20

Examples of plasticizers for the acrylic polymers include citric acid esters, such as triethyl citrate NF21, tributyl citrate, dibutyl phthalate (DBP), acetyltri-N-butyl citrate (ATBC), and possibly 1,2-propylene glycol, polyethylene glycols, propylene glycol, diethyl phthalate, castor oil, and triacetin, although it is possible that other plasticizers (such as acetylated monoglycerides, phthalate esters, castor oil; etc.) can be used. 25

The sustained-release profile of drug release in the formulations of the invention (either in vivo or in vitro) can be altered, for example, by using more than one release-retarding or sequestering material, varying the thickness of the release-retarding or sequestering material, changing the particular release-retarding or sequestering material used, altering the relative amounts of release-retarding or sequestering material, altering the manner in which the plasticizer is added (e.g., when the sustained-release coating is derived from an aqueous dispersion of hydrophobic polymer), by varying the amount of plasticizer relative to retardant material, by the inclusion of additional ingredients or excipients, by altering the method of manufacture; etc. 30

In certain other embodiments, the oral dosage form can utilize a multiparticulate sustained-release matrix. In certain 35 embodiments, the sustained-release matrix comprises a hydrophilic and/or hydrophobic polymer, such as gums, cellulose ethers, acrylic resins and protein-derived materials. Of these polymers, the cellulose ethers, specifically hydroxyalkylcelluloses and carboxyalkylcelluloses, are preferred. The oral dosage form can contain between about 1% and about 80% (by weight) of at least one hydrophilic or hydrophobic polymer. 40

The hydrophobic material is preferably selected from the group consisting of alkylcelluloses, acrylic and methacrylic acid polymers and copolymers, shellac, zein, hydrogenated castor oil, hydrogenated vegetable oil, or mixtures thereof. Preferably, the hydrophobic material is a pharmaceutically acceptable acrylic polymer, including acrylic acid and methacrylic acid copolymers, methyl methacrylate, methyl methacrylate copolymers, ethoxyethyl methacrylates, cyanoethyl methacrylate, aminoalkyl methacrylate copolymer, poly(acrylic acid), poly(methacrylic acid), methacrylic acid alkylamine copolymer, poly(methyl methacrylate), poly(methacrylic acid) (anhydride), polymethacrylate, polyacrylamide, poly(methacrylic acid anhydride), and glycidyl methacrylate copolymers. In other embodiments, the hydrophobic material 45

can also include hydroxyalkylcelluloses such as hydroxypropylmethylcellulose and mixtures of the foregoing.

Preferred hydrophobic materials are water-insoluble with more or less pronounced hydrophobic trends. Preferably, the hydrophobic material has a melting point from about 30° C. to about 200° C., more preferably from about 45° C. to about 90° C. The hydrophobic material can include neutral or synthetic waxes, fatty alcohols (such as lauryl, myristyl, stearyl, cetyl or preferably cetostearyl alcohol), fatty acids, including fatty acid esters, fatty acid glycerides (mono-, di-, and tri-glycerides), hydrogenated fats, hydrocarbons, normal waxes, stearic acid, stearyl alcohol and hydrophobic and hydrophilic materials having hydrocarbon backbones. Suitable waxes include beeswax, glycowax, castor wax, carnauba wax and wax-like substances, e.g., material normally solid at room temperature and having a melting point of from about 30° C. to about 100° C.

Preferably, a combination of two or more hydrophobic materials are included in the matrix formulations. If an additional hydrophobic material is included, it is preferably a natural or synthetic wax, a fatty acid, a fatty alcohol, or mixtures thereof. Examples include beeswax, carnauba wax, stearic acid and stearyl alcohol.

In other embodiments, the sustained-release matrix comprises digestible, long-chain (e.g., C<sub>8</sub>-C<sub>50</sub>, preferably C<sub>12</sub>-C<sub>40</sub>), substituted or unsubstituted hydrocarbons, such as fatty acids, fatty alcohols, glyceryl esters of fatty acids, mineral and vegetable oils and waxes. Hydrocarbons having a melting point of between about 25° C. and about 90° C. are preferred. Of these long-chain hydrocarbon materials, fatty (aliphatic) alcohols are preferred. The oral dosage form can contain up to about 60% (by weight) of at least one digestible, long-chain hydrocarbon. Further, the sustained-release matrix can contain up to 60% (by weight) of at least one polyalkylene glycol.

In a preferred embodiment, the matrix comprises at least one water-soluble hydroxyalkyl cellulose, at least one C<sub>12</sub>-C<sub>36</sub>, preferably C<sub>14</sub>-C<sub>22</sub>, aliphatic alcohol and, optionally, at least one polyalkylene glycol. The at least one hydroxyalkyl cellulose is preferably a hydroxy (C<sub>1</sub>-C<sub>6</sub>) alkyl cellulose, such as hydroxypropylcellulose, hydroxypropylmethylcellulose and, preferably, hydroxyethyl cellulose. The amount of the at least one hydroxyalkyl cellulose in the oral dosage form will be determined, amongst other things, by the precise rate of opioid release required. The amount of the at least one aliphatic alcohol in the present oral dosage form will be determined by the precise rate of opioid release required. However, it will also depend on whether the at least one polyalkylene glycol is absent from the oral dosage form.

In certain embodiments, a spherizing agent, together with the active ingredient, can be spherized to form spheroids. Microcrystalline cellulose and hydrous lactose impalpable are examples of such agents. Additionally (or alternatively), the spheroids can contain a water-insoluble polymer, preferably an acrylic polymer, an acrylic copolymer, such as a methacrylic acid-ethyl acrylate copolymer, or ethyl cellulose. In such embodiments, the sustained-release coating will generally include a water-insoluble material such as (a) a wax, either alone or in admixture with a fatty alcohol, or (b) shellac or zein.

The sustained-release unit can be prepared by any suitable method. For example, a plasticized aqueous dispersion of the release-retarding or sequestering material can be applied onto the subunit comprising the opioid agonist. A sufficient amount of the aqueous dispersion of release-retarding or sequestering material to obtain a predetermined sustained-release of the opioid agonist when the coated substrate is exposed to aqueous solutions, e.g., gastric fluid, is preferably

applied, taking into account the physical characteristics of the opioid agonist, the manner of incorporation of the plasticizer; etc. Optionally, a further overcoat of a film-former, such as Opadry (Colorcon, West Point, Va.), can be applied after coating with the release-retarding or sequestering material.

The subunit can be cured in order to obtain a stabilized release rate of the therapeutic agent. In embodiments employing an acrylic coating, a stabilized product can be preferably obtained by subjecting the subunit to oven curing at a temperature above the glass transition temperature of the plasticized acrylic polymer for the required time period. The optimum temperature and time for the particular formulation can be determined by routine experimentation.

Once prepared, the subunit can be combined with at least one additional subunit and, optionally, other excipients or drugs to provide an oral dosage form. In addition to the above ingredients, a sustained-release matrix also can contain suitable quantities of other materials, e.g., diluents, lubricants, binders, granulating aids, colorants, flavorants and glidants that are conventional in the pharmaceutical art.

Optionally and preferably, the mechanical fragility of any of the sequestering subunits described herein is the same as the mechanical fragility of the therapeutic agent in releasable form. In this regard, tampering with the composition of the invention in a manner to obtain the therapeutic agent will result in the destruction of the sequestering subunit, such that the antagonist is released and mixed in with the therapeutic agent. Consequently, the antagonist cannot be separated from the therapeutic agent, and the therapeutic agent cannot be administered in the absence of the antagonist. Methods of assaying the mechanical fragility of the sequestering subunit and of a therapeutic agent are known in the art.

The composition of the invention can be in any suitable dosage form or formulation, (see, e.g., *Pharmaceutics and Pharmacy Practice*, J. B. Lippincott Company, Philadelphia, Pa., Banker and Chalmers, eds., pages 238-250 (1982)). Pharmaceutically acceptable salts of the antagonist or agonist agents discussed herein include metal salts, such as sodium salt, potassium salt, cesium salt, and the like; alkaline earth metals, such as calcium salt, magnesium salt, and the like; organic amine salts, such as triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, and the like; inorganic acid salts, such as hydrochloride, hydrobromide, sulfate, phosphate, and the like; organic acid salts, such as formate, acetate, trifluoroacetate, maleate, tartrate, and the like; sulfonates, such as methanesulfonate, benzenesulfonate, p-toluenesulfonate, and the like; amino acid salts, such as arginate, asparinate, glutamate, and the like. Formulations suitable for oral administration can consist of (a) liquid solutions, such as an effective amount of the inhibitor dissolved in diluents, such as water, saline, or orange juice; (b) capsules, sachets, tablets, lozenges, and troches, each containing a predetermined amount of the active ingredient, as solids or granules; (c) powders; (d) suspensions in an appropriate liquid; and (e) suitable emulsions. Liquid formulations may include diluents, such as water and alcohols, for example, ethanol, benzyl alcohol, and the polyethylene alcohols, either with or without the addition of a pharmaceutically acceptable surfactant. Capsule forms can be of the ordinary hard- or soft-shelled gelatin type containing, for example, surfactants, lubricants, and inert fillers, such as lactose, sucrose, calcium phosphate, and corn starch. Tablet forms can include one or more of lactose, sucrose, mannitol, corn starch, potato starch, alginic acid, microcrystalline cellulose, acacia, gelatin, guar gum, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, calcium stearate, zinc stearate, stearic

acid, and other excipients, colorants, diluents, buffering agents, disintegrating agents, moistening agents, preservatives, flavoring agents, and pharmacologically compatible excipients. Lozenge forms can comprise the active ingredient in a flavor, usually sucrose and acacia or tragacanth, as well as pastilles comprising the active ingredient in an inert base, such as gelatin and glycerin, or sucrose and acacia, emulsions, gels, and the like containing, in addition to the active ingredient, such excipients as are known in the art.

One of ordinary skill in the art will readily appreciate that the compositions of the invention can be modified in any number of ways, such that the therapeutic efficacy of the composition is increased through the modification. For instance, the therapeutic agent or sequestering subunit could be conjugated either directly or indirectly through a linker to a targeting moiety. The practice of conjugating therapeutic agents or sequestering subunits to targeting moieties is known in the art. See, for instance, Wadwa et al., *J. Drug Targeting* 3: 111 (1995), and U.S. Pat. No. 5,087,616. The term "targeting moiety" as used herein, refers to any molecule or agent that specifically recognizes and binds to a cell-surface receptor, such that the targeting moiety directs the delivery of the therapeutic agent or sequestering subunit to a population of cells on which the receptor is expressed. Targeting moieties include, but are not limited to, antibodies, or fragments thereof, peptides, hormones, growth factors, cytokines, and any other naturally- or non-naturally-existing ligands, which bind to cell-surface receptors. The term "linker" as used herein, refers to any agent or molecule that bridges the therapeutic agent or sequestering subunit to the targeting moiety. One of ordinary skill in the art recognizes that sites on the therapeutic agent or sequestering subunit, which are not necessary for the function of the agent or sequestering subunit, are ideal sites for attaching a linker and/or a targeting moiety, provided that the linker and/or targeting moiety, once attached to the agent or sequestering subunit, do(es) not interfere with the function of the therapeutic agent or sequestering subunit.

With respect to the present inventive compositions, the composition is preferably an oral dosage form. By "oral dosage form" is meant to include a unit dosage form prescribed or intended for oral administration comprising subunits. Desirably, the composition comprises the sequestering subunit coated with the therapeutic agent in releasable form, thereby forming a composite subunit comprising the sequestering subunit and the therapeutic agent. Accordingly, the invention further provides a capsule suitable for oral administration comprising a plurality of such composite subunits.

Alternatively, the oral dosage form can comprise any of the sequestering subunits of the invention in combination with a therapeutic agent subunit, wherein the therapeutic agent subunit comprises the therapeutic agent in releasable form. In this respect, the invention provides a capsule suitable for oral administration comprising a plurality of sequestering subunits of the invention and a plurality of therapeutic subunits, each of which comprises a therapeutic agent in releasable form.

The invention further provides tablets comprising a sequestering subunit of the invention and a therapeutic agent in releasable form. For instance, the invention provides a tablet suitable for oral administration comprising a first layer comprising any of the sequestering subunits of the invention and a second layer comprising therapeutic agent in releasable form, wherein the first layer is coated with the second layer. The first layer can comprise a plurality of sequestering subunits. Alternatively, the first layer can be or can consist of a single sequestering subunit. The therapeutic agent in releas-

able form can be in the form of a therapeutic agent subunit and the second layer can comprise a plurality of therapeutic subunits. Alternatively, the second layer can comprise a single substantially homogeneous layer comprising the therapeutic agent in releasable form.

When the blocking agent is a system comprising a first antagonist-impermeable material and a core, the sequestering subunit can be in one of several different forms. For example, the system can further comprise a second antagonist-impermeable material, in which case the sequestering unit comprises an antagonist, a first antagonist-impermeable material, a second antagonist-impermeable material, and a core. In this instance, the core is coated with the first antagonist-impermeable material, which, in turn, is coated with the antagonist, which, in turn, is coated with the second antagonist-impermeable material. The first antagonist-impermeable material and second antagonist-impermeable material substantially prevent release of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. In some instances, it is preferable that the first antagonist-impermeable material is the same as the second antagonist-impermeable material. In other instances, the first antagonist-impermeable material is different from the second antagonist-impermeable material. It is within the skill of the ordinary artisan to determine whether or not the first and second antagonist-impermeable materials should be the same or different. Factors that influence the decision as to whether the first and second antagonist-impermeable materials should be the same or different can include whether a layer to be placed over the antagonist-impermeable material requires certain properties to prevent dissolving part or all of the antagonist-impermeable layer when applying the next layer or properties to promote adhesion of a layer to be applied over the antagonist-impermeable layer.

Alternatively, the antagonist can be incorporated into the core, and the core is coated with the first antagonist-impermeable material. In this case, the invention provides a sequestering subunit comprising an antagonist, a core and a first antagonist-impermeable material, wherein the antagonist is incorporated into the core and the core is coated with the first antagonist-impermeable material, and wherein the first antagonist-impermeable material substantially prevents release of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. By "incorporate" and words stemming therefrom, as used herein is meant to include any means of incorporation, e.g., homogeneous dispersion of the antagonist throughout the core, a single layer of the antagonist coated on top of a core, or a multi-layer system of the antagonist, which comprises the core.

In another alternative embodiment, the core comprises a water-insoluble material, and the core is coated with the antagonist, which, in turn, is coated with the first antagonist-impermeable material. In this case, the invention further provides a sequestering subunit comprising an antagonist, a first antagonist-impermeable material, and a core, which comprises a water-insoluble material, wherein the core is coated with the antagonist, which, in turn, is coated with the first antagonist-impermeable material, and wherein the first antagonist-impermeable material substantially prevents release of the antagonist from the sequestering subunit in the gastrointestinal tract for a time period that is greater than 24 hours. The term "water-insoluble material" as used herein means any material that is substantially water-insoluble. The term "substantially water-insoluble" does not necessarily refer to complete or 100% water-insolubility. Rather, there are varying degrees of water insolubility of which one of



ordinary skill in the art recognizes as having a potential benefit. Preferred water-insoluble materials include, for example, microcrystalline cellulose, a calcium salt, and a wax. Calcium salts include, but are not limited to, a calcium phosphate (e.g., hydroxyapatite, apatite; etc.), calcium carbonate, calcium sulfate, calcium stearate, and the like. Waxes include, for example, carnuba wax, beeswax, petroleum wax, candelilla wax, and the like.

In one embodiment, the sequestering subunit includes an antagonist and a seal coat where the seal coat forms a layer physically separating the antagonist within the sequestering subunit from the agonist which is layered upon the sequestering subunit. In one embodiment, the seal coat comprises one or more of an osmotic pressure regulating agent, a charge-neutralizing additive, a sequestering polymer hydrophobicity-enhancing additive, and a first sequestering polymer (each having been described above). In such embodiments, it is preferred that the osmotic pressure regulating agent, charge-neutralizing additive, and/or sequestering polymer hydrophobicity-enhancing additive, respectively where present, are present in proportion to the first sequestering polymer such that no more than 10% of the antagonist is released from the intact dosage form. Where an opioid antagonist is used in the sequestering subunit and the intact dosage form includes an opioid agonist, it is preferred that ratio of the osmotic pressure regulating agent, charge-neutralizing additive, and/or sequestering polymer hydrophobicity-enhancing additive, respectively where present, in relation to the first sequestering polymer is such that the physiological effect of the opioid agonist is not diminished when the composition is in its intact dosage form or during the normal course digestion in the patient. Release may be determined as described above using the USP paddle method (optionally using a buffer containing a surfactant such as Triton X-100) or measured from plasma after administration to a patient in the fed or non-fed state. In one embodiment, plasma naltrexone levels are determined; in others, plasma 6-beta naltrexol levels are determined. Standard tests may be utilized to ascertain the antagonist's effect on agonist function (i.e., reduction of pain).

The sequestering subunit of the invention can have a blocking agent that is a tether to which the antagonist is attached. The term "tether" as used herein refers to any means by which the antagonist is tethered or attached to the interior of the sequestering subunit, such that the antagonist is not released, unless the sequestering subunit is tampered with. In this instance, a tether-antagonist complex is formed. The complex is coated with a tether-impermeable material, thereby substantially preventing release of the antagonist from the subunit. The term "tether-impermeable material" as used herein refers to any material that substantially prevents or prevents the tether from permeating through the material. The tether preferably is an ion exchange resin bead.

The invention further provides a tablet suitable for oral administration comprising a single layer comprising a therapeutic agent in releasable form and a plurality of any of the sequestering subunits of the invention dispersed throughout the layer of the therapeutic agent in releasable form. The invention also provides a tablet in which the therapeutic agent in releasable form is in the form of a therapeutic agent subunit and the tablet comprises an at least substantially homogeneous mixture of a plurality of sequestering subunits and a plurality of subunits comprising the therapeutic agent.

In preferred embodiments, oral dosage forms are prepared to include an effective amount of melt-extruded: subunits in the form of multiparticles within a capsule. For example, a plurality of the melt-extruded multiparticulates can be placed

in a gelatin capsule in an amount sufficient to provide an effective release dose when ingested and contacted by gastric fluid.

In another preferred embodiment, the subunits, e.g., in the form of multiparticulates, can be compressed into an oral tablet using conventional tableting equipment using standard techniques. Techniques and compositions for making tablets (compressed and molded), capsules (hard and soft gelatin) and pills are also described in *Remington's Pharmaceutical Sciences*, (Aurthur Osol., editor), 1553-1593 (1980), which is incorporated herein by reference. Excipients in tablet formulation can include, for example, an inert diluent such as lactose, granulating and disintegrating agents, such as cornstarch, binding agents, such as starch, and lubricating agents, such as magnesium stearate. In yet another preferred embodiment, the subunits are added during the extrusion process and the extrudate can be shaped into tablets as set forth in U.S. Pat. No. 4,957,681 (Klimesch et al.), which is incorporated herein by reference.

Optionally, the sustained-release, melt-extruded, multiparticulate systems or tablets can be coated, or the gelatin capsule can be further coated, with a sustained-release coating, such as the sustained-release coatings described herein. Such coatings are particularly useful when the subunit comprises an opioid agonist in releasable form, but not in sustained-release form. The coatings preferably include a sufficient amount of a hydrophobic material to obtain a weight gain level form about 2 to about 30 percent, although the overcoat can be greater, depending upon the physical properties of the particular opioid analgesic utilized and the desired release rate, among other things.

The melt-extruded dosage forms can further include combinations of melt-extruded multiparticulates containing one or more of the therapeutically active agents before being encapsulated. Furthermore, the dosage forms can also include an amount of an immediate release therapeutic agent for prompt therapeutic effect. The immediate release therapeutic agent can be incorporated or coated on the surface of the subunits after preparation of the dosage forms (e.g., controlled-release coating or matrix-based). The dosage forms can also contain a combination of controlled-release beads and matrix multiparticulates to achieve a desired effect.

The sustained-release formulations preferably slowly release the therapeutic agent, e.g., when ingested and exposed to gastric fluids, and then to intestinal fluids. The sustained-release profile of the melt-extruded formulations can be altered, for example, by varying the amount of retardant, e.g., hydrophobic material, by varying the amount of plasticizer relative to hydrophobic material, by the inclusion of additional ingredients or excipients, by altering the method of manufacture; etc.

In other embodiments, the melt-extruded material is prepared without the inclusion of the subunits, which are added thereafter to the extrudate. Such formulations can have the subunits and other drugs blended together with the extruded matrix material, and then the mixture is tableted in order to provide a slow release of the therapeutic agent or other drugs. Such formulations can be particularly advantageous, for example, when the therapeutically active agent included in the formulation is sensitive to temperatures needed for softening the hydrophobic material and/or the retardant material.

In certain embodiments, the release of the antagonist of the sequestering subunit or composition is expressed in terms of a ratio of the release achieved after tampering, e.g., by crushing or chewing, relative to the amount released from the intact formulation. The ratio is, therefore, expressed as Crushed: Whole, and it is desired that this ratio have a numerical range

of at least about 4:1 or greater (e.g., crushed release within 1 hour/intact release in 24 hours). In certain embodiments, the ratio of the therapeutic agent and the antagonist, present in the sequestering subunit, is about 1:1 to about 50:1 by weight, preferably about 1:1 to about 20:1 by weight or 15:1 to about 30:1 by weight. The weight ratio of the therapeutic agent to antagonist refers to the weight of the active ingredients. Thus, for example, the weight of the therapeutic agent excludes the weight of the coating, matrix, or other component that renders the antagonist sequestered, or other possible excipients associated with the antagonist particles. In certain preferred embodiments, the ratio is about 1:1 to about 10:1 by weight. Because in certain embodiments the antagonist is in a sequestered form, the amount of such antagonist within the dosage form can be varied more widely than the therapeutic agent/antagonist combination dosage forms, where both are available for release upon administration, as the formulation does not depend on differential metabolism or hepatic clearance for proper functioning. For safety reasons, the amount of the antagonist present in a substantially non-releasable form is selected as not to be harmful to humans, even if fully released under conditions of tampering.

Thus, in certain embodiments, a pharmaceutical composition comprising an antagonist in direct contact with a seal coat, an agonist in direct contact with the seal coat and a sequestering polymer but not the antagonist, wherein the antagonist and agonist are present within a single multilayer pharmaceutical unit, is provided. In others, pharmaceutical compositions comprising a pharmaceutical dosing unit consisting essentially of a multiple layer bead comprising an antagonist and an agonist that are not in direct contact with one another are provided. In yet others, pharmaceutical composition comprising a plurality of pharmaceutically active units wherein each unit comprises an antagonist, an agonist, a seal coat, and a sequestering polymer wherein the antagonist and the agonist are not in direct contact with one another. In still others, pharmaceutical compositions comprising a pharmaceutically inert support material such as a sugar sphere, an antagonist in direct contact with the support material, a seal coat in direct contact with the antagonist and an agonist, and a sequestering polymer in direct contact with the agonist are provided. In preferred embodiments, multiple layer pharmaceutical compositions comprising an agonist and an antagonist within distinct layers of the composition, wherein at least 90-95% of the antagonist is sequestered for at least 24 hours following administration to a human being are provided. In a particularly preferred embodiment, a pharmaceutical composition comprising naltrexone within a sequestering subunit and morphine in contact with the subunit but not the naltrexone, wherein administration of the composition to a human being results in the release of substantially all of the morphine from the composition but less than 5-10% of the naltrexone from the composition within 24 hours of administration, is provided. Also provided are methods for preparing pharmaceutical compositions by, for example, adhering an antagonist to a pharmaceutically inert support material, coating the antagonist with a seal coat that includes a sequestering polymer, coating the seal coat with an agonist, and coating the agonist with a release-retarding or sequestering material. In another embodiment, a method for measuring the amount of antagonist or derivative thereof in a biological sample, the antagonist or derivative having been released from a pharmaceutical composition in vivo, the method comprising the USP paddle method at 37° C., 100 rpm, but further comprising incubation in a buffer containing a surfactant such as Triton X-100, for example.

A particularly preferred embodiment comprises a multiple layer pharmaceutical is described in the Examples is multi-layer naltrexone/morphine dosing unit in an abuse-resistant dosage form. Naltrexone is contained in a sequestering subunit comprising a seal coat comprising Eudragit® RS and the optimization agents SLS, talc and chloride ions that together prevent release of naltrexone upon hydration. Overlaid onto the sequestering subunit is a layer comprising morphine that is released upon hydration in pH 7.5 buffer; the naltrexone, however, remains within the sequestering subunit under these conditions. If the unit is modified by, for example, crushing the unit, the sequestering subunit is crushed as well causing the release of both morphine and naltrexone therefrom.

Thus, the compositions are particularly well-suited for use in preventing abuse of a therapeutic agent. In this regard, the invention also provides a method of preventing abuse of a therapeutic agent by a human being. The method comprises incorporating the therapeutic agent into any of the compositions of the invention. Upon administration of the composition of the invention to the person, the antagonist is substantially prevented from being released in the gastrointestinal tract for a time period that is greater than 24 hours. However, if a person tampers with the compositions, the sequestering subunit, which is mechanically fragile, will break and thereby allow the antagonist to be released. Since the mechanical fragility of the sequestering subunit is the same as the therapeutic agent in releasable form, the antagonist will be mixed with the therapeutic agent, such that separation between the two components is virtually impossible.

A better understanding of the present invention and of its many advantages will be had from the following examples, given by way of illustration.

## EXAMPLES

The preparations and experiments described below were actually performed. In certain cases, however, the present tense is utilized.

### Example 1

#### Formulation Evaluation

##### A. Exclusion of Charge-neutralization Additive (SLS)

	RB 380-56	
	Gram per batch	Percent
<u>Seal-coated sugar spheres</u>		
Sugar spheres	577.9	51.8
Ethylcellulose N50	46.2	4.1
Talc	123.3	11.1
Dibutyl Sebacate	4.6	0.4
<u>Naltrexone cores</u>		
Seal-coated sugar spheres	(752.0)	(67.4)
Naltrexone HCl	27.2	2.4
Klucel LF	5.2	0.5
Talc	12.8	1.1
Ascorbic acid	2.8	0.3
<u>Naltrexone pellets</u>		
Naltrexone cores	(800.0)	(71.7)
Eudragit RS	150.0	13.5

31

-continued

	RB 380-56	
	Gram per batch	Percent
Sodium lauryl sulfate	0.0	0.0
Talc	150.0	13.5
Dibutyl Sebacate	15.0	1.3
Total	1115.0	100.0

Method of Preparation:

- Ethylcellulose and dibutyl sebacate were dissolved into ethanol and talc dispersed into the solution.
- The dispersion from 1 was sprayed onto sugar spheres in a Wurster to form seal-coated sugar spheres.
- Klucel L F and ascorbic acid were dissolved into a 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
- The naltrexone dispersion from 3 was sprayed onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
- Eudragit RS and dibutyl debacate were dissolved into ethanol and talc dispersed into the solution.
- The dispersion from 5 was sprayed onto the naltrexone cores from 4 in a Wurster to form naltrexone pellets.
- Pellets were dried at 50° C. for 48 hours.
- The resulting pellets had a Eudragit RS coat thickness of 47 µm.

Drug Release Results

Dissolution conditions: USP paddle method at 37° C. and 100 rpm, 1 hour in 500 mL of 0.1N HCl followed by 72 hours in 500 mL of 0.05M pH 7.5 phosphate buffer.

Conclusions: The results are shown in FIG. 1. The exclusion of SLS from the Naltrexone pellet (Eudragit RS) coat results in rapid release of Naltrexone, with more than 90% release in 24 hours.

B. Variable Amounts of SLS (Eudragit RS Coat Thickness of 53 µm)

	Batch Number					
	RB 358-88		RB 358-73		RB 358-83	
	Gm per batch	Percent	Gm per batch	Percent	Gm per batch	Percent
Seal-coated sugar spheres						
Sugar spheres	646.1	50.1	646.1	50.0	646.1	49.8
Ethylcellulose N50	48.5	3.8	48.5	3.7	48.5	3.7
Talc	126.0	9.8	126.0	9.7	126.0	9.7
Dibutyl Sebacate	4.9	0.4	4.9	0.4	4.9	0.4
Magnesium stearate	19.4	1.5	19.4	1.5	19.4	1.5
Sodium lauryl sulfate	1.9	0.2	1.9	0.1	1.9	0.1
Naltrexone cores						
Seal-coated sugar spheres	(846.7)	(65.6)	(846.7)	(65.5)	(846.7)	(65.2)
Naltrexone HCl	29.5	2.3	29.5	2.3	29.5	2.3
Klucel LF	5.9	0.5	5.9	0.5	5.9	0.5
Talc	17.8	1.4	17.8	1.4	17.8	1.4
Naltrexone						

32

-continued

	Batch Number					
	RB 358-88		RB 358-73		RB 358-83	
	Gm per batch	Percent	Gm per batch	Percent	Gm per batch	Percent
pellets						
Naltrexone cores	(900.0)	(69.7)	(900.0)	(69.6)	(900.0)	(69.3)
Eudragit RS	184.6	14.3	184.3	14.3	183.7	14.2
Sodium lauryl sulfate	3.0	0.23	6.1	0.47	12.3	0.95
Talc	184.6	14.3	184.3	14.3	183.7	14.2
Dibutyl Sebacate	18.5	1.4	18.4	1.4	18.4	1.4
Total	1290.7	100.0	1293.2	100.0	1298.1	100.0

Method of preparation:

- Ethylcellulose, sodium lauryl sulfate and dibutyl sebacate were dissolved into ethanol, and then talc and magnesium stearate were dispersed into the solution.
- The dispersion from 1 was sprayed onto sugar spheres in a Wurster to form seal-coated sugar spheres.
- Klucel LF was dissolved into a 20:80 mixture of water and ethanol. Naltrexone HCl and talc were then dispersed into the solution.
- The naltrexone dispersion from 3 was then sprayed onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
- Eudragit RS, sodium lauryl sulfate and dibutyl debacate were dissolved into ethanol, and talc dispersed into the solution.
- The dispersion from 5 was sprayed onto naltrexone cores from 4 in a Wurster to form naltrexone pellets.
- The pellets were dried at 50° C. for 13-16.5 hours.
- The resulting pellets had a Eudragit RS coat thickness of 51-53 µm.

Drug Release Results

Dissolution conditions: USP paddle method at 37° C. and 100 rpm, 72 hours in 500 mL of 0.05M pH 7.5 phosphate buffer

Conclusions: The results are shown in FIG. 2. Addition of a small amount of SLS (1.6% w/w of Eudragit RS) results in charge neutralization of Eudragit RS (theoretically 20% neutralization), and significantly slows down the release of naltrexone. Further addition of SLS (3.2% w/w of Eudragit RS) leads to further Eudragit RS charge neutralization (theoretically 41% neutralization), and dramatically slows down release of naltrexone. Still higher amount of SLS (6.3% w/w of Eudragit RS), however, results in higher naltrexone release, possibly due to plasticizing effect of SLS.

3. Different Levels of SLS (Eudragit RS Coat Thickness of 65 µm)

	Batch Number					
	RB 358-88A		RB 358-73A		RB 358-83A	
	Gm per batch	Percent	Gm per batch	Percent	Gm per batch	Percent
Seal-coated sugar spheres						
Sugar spheres	646.1	45.5	646.1	45.4	646.1	45.1
Ethylcellulose N50	48.5	3.4	48.5	3.4	48.5	3.4

-continued

	Batch Number					
	RB 358-88A		RB 358-73A		RB 358-83A	
	Gm per batch	Percent	Gm per batch	Percent	Gm per batch	Percent
Talc	126.0	8.9	126.0	8.8	126.0	8.8
Dibutyl Sebacate	4.9	0.3	4.9	0.3	4.9	0.3
Magnesium stearate	19.4	1.4	19.4	1.4	19.4	1.4
Sodium lauryl sulfate	1.9	0.1	1.9	0.1	1.9	0.1
Naltrexone cores						
Seal-coated sugar spheres	(846.7)	(59.6)	(846.7)	(59.4)	(846.7)	(59.1)
Naltrexone HCl	29.5	2.1	29.5	2.1	29.5	2.1
Klucel LF	5.9	0.4	5.9	0.4	5.9	0.4
Talc	17.8	1.3	17.8	1.2	17.8	1.2
Naltrexone pellets						
Naltrexone cores	(900.0)	(63.4)	(900.0)	(63.2)	(900.0)	(62.8)
Eudragit RS	245.8	17.3	245.8	17.3	245.8	17.2
Sodium lauryl sulfate	4.0	0.3	8.2	0.6	16.4	1.1
Talc	245.8	17.3	245.8	17.3	245.8	17.2
Dibutyl Sebacate	24.6	1.7	24.6	1.7	24.6	1.7
Total	1420.2	100.0	1424.4	100.0	1432.6	100.0

Method of Preparation

1. Ethylcellulose, sodium lauryl sulfate and dibutyl sebacate were dissolved into ethanol; talc and magnesium stearate were then dispersed into the solution.
2. The dispersion from 1 was sprayed onto sugar spheres in a Wurster to form seal-coated sugar spheres.
3. Klucel LF was dissolved into a 20:80 mixture of water and ethanol; naltrexone HCl and talc were then dispersed into the solution.
4. The naltrexone dispersion from 3 was then sprayed onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
5. Eudragit RS, sodium lauryl sulfate and dibutyl sebacate were dissolved into ethanol; talc was then dispersed into the solution.
6. The dispersion from 5 was sprayed onto naltrexone cores from 4 in a Wurster to form naltrexone pellets.
7. The pellets were dried at 50° C. for 13-16.5 hours.
8. The resulting pellets had a Eudragit RS coat thickness of 63-67 µm.

Drug Release Results

Dissolution conditions: USP paddle method at 37° C. and 100 rpm, 72 hours in 500 mL of 0.05M pH 7.5 phosphate buffer

Conclusions: The results are shown in FIG. 3. As described above, there is an optimal ratio of SLS to Eudragit RS.

B. Talc Content Relative to Eudragit RS Polymer

	Batch Number					
	RB 358-93		RB 358-73A		RB 358-78	
	Gm per batch	Percent	Gm per batch	Percent	Gm per batch	Percent
Seal-coated sugar spheres						
Sugar spheres	646.1	46.5	646.1	45.4	646.1	43.9
Ethylcellulose	48.5	3.5	48.5	3.4	48.5	3.3
N50						
Talc	126.0	9.1	126.0	8.8	126.0	8.6
Dibutyl Sebacate	4.9	0.4	4.9	0.3	4.9	0.3
Magnesium stearate	19.4	1.4	19.4	1.4	19.4	1.3
Sodium lauryl sulfate	1.9	0.1	1.9	0.1	1.9	0.1
Naltrexone cores						
Seal-coated sugar spheres	(846.7)	(61.0)	(846.7)	(59.4)	(846.7)	(57.5)
Naltrexone HCl	29.5	2.1	29.5	2.1	29.5	2.0
Klucel LF	5.9	0.4	5.9	0.4	5.9	0.4
Talc	17.8	1.3	17.8	1.2	17.8	1.2
Naltrexone pellets						
Naltrexone cores	(900.0)	(64.8)	(900.0)	(63.2)	(900.0)	(61.1)
Eudragit RS	266.5	19.2	245.8	17.3	216.7	14.7
Sodium lauryl sulfate	8.8	0.6	8.2	0.6	7.2	0.5
Talc	186.2	13.4	245.8	17.3	326.3	22.2
Dibutyl Sebacate	26.6	1.9	24.6	1.7	21.7	1.5
Total	1388.1	100.0	1424.4	100.0	1471.9	100.0

Method of Preparation

1. Dissolve Ethylcellulose, sodium lauryl sulfate and dibutyl sebacate into ethanol, then disperse talc and magnesium stearate into the solution.
2. Spray the dispersion from 1 onto sugar spheres in a Wurster to form seal-coated sugar spheres.
3. Dissolve Klucel LF into 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
4. Spray the naltrexone dispersion from 3 onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
5. Dissolve Eudragit RS, sodium lauryl sulfate and dibutyl sebacate into ethanol. Disperse talc into the solution.
6. Spray the dispersion from 5 onto naltrexone cores from 4 in a Wurster to form naltrexone pellets.
7. Pellets are dried at 50° C. for 13-16.5 hours.
8. Resulting pellets have Eudragit RS coat thickness of 63-67 µm.

Drug Release Results

Dissolution conditions: USP paddle method at 37° C. and 100 rpm, 72 hours in 500 mL of 0.05M pH 7.5 phosphate buffer

Conclusions. The results of this assay are shown in FIG. 4, and demonstrate that there is an optimal ratio of talc to Eudragit RS (approximately 1:1). Talc increases the hydrophobicity of the Eudragit RS coat, but also reduces film integrity at high amount. FIG. 5 demonstrates the transition point in the behavior of the film. FIG. 6 demonstrates that there is a

distinct optimum in the relationship between film permeability and talc content when using a sugar sphere core.  
 C. Effects of Osmotic Pressure Reducing Agents on Top of Eudragit RS Coat

	Percent Batch Number			
	RB 362-28	RB 362-48	RB 362-67	RB 362-65
Naltrexone cores				
Naltrexone HCl	1.10	0.93	0.89	1.00
Sugar (#20-25 mesh)	24.48	20.59	19.80	22.15
HPC (Klucel LF)	0.22	0.19		
HPMC, 3 cps			0.18	0.20
Citric acid			0.004	0.004
Ascorbic acid			0.004	0.004
BHA			0.004	0.004
Talc	0.66	0.56	0.54	0.60
Naltrexone pellets				
Naltrexone cores	(26.47)	(22.26)	(21.41)	(23.95)
Eudragit RS PO	10.64	8.95	8.62	9.64
SLS	0.36	0.30	0.29	0.33
DBS	1.06	0.89	0.85	0.95
Talc	10.89	9.16	8.62	9.64
Naltrexone-morphine cores				
Naltrexone pellets	(49.41)	(41.55)	(39.78)	(44.50)
Morphine sulfate	26.05	21.70	21.70	24.76
Confectioner's sugar		13.66	9.32	
Sodium chloride			6.43	7.01
HPMC, 3 cps	2.32	3.46	3.13	4.10
Naltrexone-morphine pellets				
Naltrexone-morphine cores	(77.78)	(80.37)	(80.37)	(80.37)
Ethylcellulose N50	7.48	7.07	7.07	7.07
PEG 6000	3.59	2.88	2.81	2.62
Eudragit L100-55	2.10	1.70	1.77	1.96
DEP	1.65	1.44	1.44	1.44
Talc	7.41	6.54	6.54	6.54
Total	100.00	100.00	100.00	100.00

Method of preparation:

1. Klucel LF or HPMC (with or without citric acid, ascorbic acid and butylated hydroxyanisole) was dissolved into 20:80 mixture of water and ethanol; naltrexone HCl and talc were dispersed into the solution.
2. The naltrexone dispersion from 1 was sprayed onto sugar spheres in a Wurster to form naltrexone cores.
3. Eudragit RS, sodium lauryl sulfate and dibutyl debacate were dissolved into ethanol; talc was then dispersed into the solution.
4. The dispersion from 3 was sprayed onto naltrexone cores from 2 in a Wurster to form naltrexone pellets.
5. The Naltrexone pellets were dried at 50° C. for either 12 hours (RB 362-28 and RB 362-48) or 65 hours (RB 362-67 and RB 362-65).
6. The resulting pellets had a Eudragit RS coat thickness of 85-90 μm.
7. Sodium chloride and hypromellose were then dissolved into water.
8. HPMC was dissolved into either water or mixture of ethanol and water.
9. Sodium chloride was dissolved into the HPMC solution from 8.
10. Confectioner's sugar was dispersed into the HPMC solution from 8.

11. Morphine sulfate was dispersed into the HPMC solution from 8.
12.
  - a. For RB 362-28, spray onto naltrexone pellets in 5 in a rotor the solution from 8, followed by the dispersion from 11, to form naltrexone-morphine cores.
  - b. For RB 362-48, spray onto naltrexone pellets in 5 in a rotor the solution from 8, followed by the dispersion from 10, followed by the solution from 8, and followed by the dispersion from 11, to form naltrexone-morphine cores.
  - c. For RB 362-67, spray onto naltrexone pellets in 5 in a rotor the solution from 9, followed by the dispersion from 10, followed by the solution from 8, and followed by the dispersion from 11, to form naltrexone-morphine cores.
  - d. For RB 362-65, spray onto naltrexone pellets in 5 in a rotor the solution from 9, followed by the solution from 8, and followed by the dispersion from 11, to form naltrexone-morphine cores.
13. Ethylcellulose, PEG 6000, Eudragit L100-55 and diethyl phthalate were dissolved into ethanol and talc was dispersed into the solution.
14. The dispersion from 13 was sprayed onto naltrexone-morphine cores in 12 to form naltrexone-morphine pellets.

Drug Release Results:

Dissolution conditions: USP paddle method at 37° C. and 100 rpm, 72 hours in 500 mL of 0.05M pH 7.5 phosphate buffer; or, USP paddle method at 37° C. and 100 rpm, 1 hour in 0.1N HCl, followed by 72 hours in 0.05M pH 7.5 phosphate buffer

Results:

Batch Number	% NT release at the end of dissolution	
RB 362-28	Naltrexone pellet	2
	Naltrexone-morphine pellet	7.9
RB 362-48	Naltrexone pellet	2
	Naltrexone-morphine pellet	68.5
RB 362-67	Naltrexone pellet	0
	Naltrexone-morphine pellet	25
RB 362-65	Naltrexone pellet	0.2
	Naltrexone-morphine pellet	1.4

Conclusions: Sugar has a detrimental effect on NT release. The use of NaCl/HPMC provides the desired NT release profile.

II. Proof of Concept Study, 16 mg Naltrexone HCl (20-727-1N)

	PI-1460		PI-1461	
	mg/unit	Percent	mg/unit	Percent
Naltrexone HCl	8	2.23	8	2.07
Sugar sphere (#20-25 mesh)	177.9	49.6		
Celllets (#20-25 mesh)			228.3	59.1
HPC (Klucel LF)	1.6	0.4	1.6	0.4
Talc	4.8	1.3	4.8	1.2
Eudragit RS PO	77.3	21.5	66.2	17.2
SLS	2.6	0.7	2.3	0.6
DBS	7.7	2.1	6.6	1.7
Talc	79.1	22.0	68.2	17.7
Total	359	100.0	386	100.0

A. Method of Preparation—

1. Dissolve Klucel LF into 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
2. Spray the naltrexone dispersion from 1 onto sugar spheres (for PI-1460) or Cellets (for PI-1461) in a Wurster to form naltrexone cores.
3. Dissolve Eudragit RS, sodium lauryl sulfate and dibutyl sebacate into ethanol. Disperse talc into the solution.
4. Spray the dispersion from 3 onto naltrexone cores from 2 in a Wurster to form naltrexone pellets.
5. The naltrexone pellets are dried in an oven at 50° C. for 12 hours.
6. Resulting pellets have Eudragit RS coat thickness of 90 μm (for PI-1460) and 60 μm (for PI-1461).
7. The pellets are filled into capsules.

B. In-vitro Drug Release—

Method	USP paddle method at 37° C. and 100 rpm; 1 hour in 0.1N HCl, then 72 hours in 0.05M pH 7.5 phosphate buffer
Results	Percent of NT released at 73 hours for PI-1460 = 2% Percent of NT released at 73 hours for PI-1461 = 0%

C. In-vivo Biostudy—

Single-dose, open-label, two-period pilot study in 26 healthy subjects under fasting conditions:

Period 1: Oral liquid containing 16 mg naltrexone (N=26)

Period 2: 2 capsules of PI-1460 (N=13) or PI-1461 (N=13)

Blood samples were withdrawn from prior to dosing and from 0.5 to 72 hours after dosing, and analyzed for plasma naltrexone and 6-beta-Naltrexol levels. Limit of quantitation was 20.0 pg/mL for naltrexone and 0.250 pg/mL for 6-beta-Naltrexol. The data is shown in FIGS. 7-10.

Summary of Pharmacokinetic Results—

	6-beta-Naltrexol			Naltrexone		
	NTX Solution	2 capsules of PI-1460	2 capsules of PI-1461	NTX Solution	2 capsules of PI-1460	2 capsules of PI-1461
Tmax (hr)	0.75	43.02	32.01	0.75	24.38 (N = 4)	23.21 (N = 10)
Cmax (pg/mL)	24600	298	834	2950	22.4 (N = 11)	60.7
AUC <sub>last</sub> (pg * h/mL)	205800	10460	32530	8925	200.2 (N = 11)	1258
AUC <sub>linf</sub> (pg * h/mL)	212700			9569		
				(N = 23)		
Relative Bioavailability to an oral solution:						
Cmax Ratio (Capsule/Solution)		1.21%	3.39%		0.76%	2.06%
AUC <sub>last</sub> Ratio (Capsule/Solution)		5.08%	15.80%		2.24%	14.08%

N = 26 for Solution, unless specified otherwise

N = 12 for PI-1460 or PI-1461, unless specified otherwise

D. Conclusion—

1. Plasma 6-beta-naltrexol levels provide a more accurate indicator of bioavailability than plasma NT levels, due to its higher plasma levels and higher analytical sensitivity.
2. Using 6-beta-naltrexol AUC<sub>last</sub> ratio of capsules to solution as indicator of cumulative in vivo NT release, significant sequestering of naltrexone is observed to 72 hours under fasting condition. Using Cellets as seed cores resulted in three times higher observed in vivo NT release than sugar. However, NT pellets using Cellet have lower RS coat thickness than Sugar (60 μm versus 90 μm), because at 60 μm, Cellet NT pellets have slightly better in vitro dissolution performance than Sugar NT pellets at 90 μm.

15 III. Optimization Study #1, Morphine Sulfate and Naltrexone 60 mg/2.4 mg (ALPH-KNT-002)

	PI-1462		PI-1463	
	mg/unit	Percent	mg/unit	Percent
Naltrexone cores				
Naltrexone HCl	2.4	0.96	2.4	0.94
Cellets (#20-25 mesh)	67.1	26.8	59.8	23.4
HPC (Klucel LF)	0.5	0.2	0.5	0.2
Citric acid	0.01	0.0040	0.01	0.004
Ascorbic acid	0.01	0.0040	0.01	0.004
BHA	0.01	0.0040	0.01	0.004
Talc	1.38	0.6	1.57	0.6
Subtotal	71.4	28.5	64.3	25.1
Naltrexone pellets				
Naltrexone cores	(71.4)	(28.5)	(64.3)	(25.1)
Eudragit RS PO	19.5	7.8	26	10.2
SLS	0.7	0.3	0.9	0.4
DBS	2	0.8	2.6	1.0
Talc	20	8.0	26.6	10.4
Subtotal	113.6	45.4	120.4	47.1

-continued

	PI-1462		PI-1463	
	mg/unit	Percent	mg/unit	Percent
Naltrexone-morphine cores				
Naltrexone pellets	(113.6)	(45.4)	(120.4)	(47.1)
Morphine sulfate	58.7	23.5	56.3	22.0
Sodium chloride	16.6	6.6	16.6	6.5
HPMC, 3 cps	13.6	5.4	13.5	5.3
Subtotal	202.5	80.9	206.8	80.8
Naltrexone-morphine pellets				
Naltrexone-morphine cores	(202.5)	(80.9)	(206.8)	(80.8)
Ethylcellulose N50	16	6.4	16.4	6.4
PEG 6000	7.4	3.0	7.6	3.0
Eudragit L100-55	3.5	1.4	3.6	1.4
DEP	3.3	1.3	3.4	1.3
Talc	17.5	7.0	18	7.0
Total	250.2	100.0	255.8	100.0

A. Method of Preparation—

1. Dissolve Klucel LF, citric acid, ascorbic acid and butylated hydroxyanisole into 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
2. Spray the naltrexone dispersion from 1 onto Cellets in a Wurster to form naltrexone cores.
3. Dissolve Eudragit RS, sodium lauryl sulfate and dibutyl debacate into ethanol. Disperse talc into the solution.
4. Spray the dispersion from 3 onto naltrexone cores from 2 in a Wurster to form naltrexone pellets.
5. The Naltrexone pellets are dried at 50° C. for 48 hours.
6. Resulting pellets have a Eudragit RS coat thickness of 60 µm for PI-1462 and 90 µm for PI-1463.
7. Dissolve sodium chloride and hypromellose into water.
8. Dissolve hypromellose into 10:90 mixture of water and ethanol. Disperse morphine sulfate into the solution.
9. Spray the solution from 7 followed by the dispersion from 8 onto naltrexone pellets in 5 in a rotor to form naltrexone-morphine cores.
10. Dissolve ethylcellulose, PEG 6000, Eudragit L100-55 and diethyl phthalate into ethanol. Disperse talc into the solution.
11. Spray the dispersion from 10 onto naltrexone-morphine cores in 9 to form naltrexone-morphine pellets.
12. The pellets are filled into capsules.

B. In-vitro Drug Release—

Method	USP paddle method at 37° C. and 100 rpm 1 hour in 0.1N HCl, then 72 hours in 0.05M pH 7.5 phosphate buffer
Results	Percent of NT released at 73 hours for PI-1462 = 0% Percent of NT released at 73 hours for PI-1463 = 0%

C. In-vivo Study

This is a single-dose, open-label, single-period study in which two groups of eight subjects received one dose of either PI-1462 or PI-1463 under fasting condition. Blood samples were drawn prior to dose administration and at 0.5 to 168 hours post-dose. Limits of quantitation are 4.00 pg/mL for naltrexone and 0.250 pg/mL for 6-beta-naltrexol. The data is shown in FIGS. 11-12.

2. Summary of Pharmacokinetics Parameters

	6-beta-Naltrexol		Naltrexone	
	PI-1462	PI-1463	PI-1462	PI-1463
Tmax (hr)	49.52	40.53	42.03	37.75 (N = 3)
Cmax (pg/mL)	349	285	25.3	35.5
AUC <sub>last</sub> (pg * h/mL)	16850	11130	705.1	835.0
AUC <sub>∞</sub> (pg * h/mL)	17040	11170	1057 (N = 4)	1711 (N = 3)
T <sub>1/2</sub> (hr)	18.18	14.49	14.15 (N = 4)	8.89 (N = 3)
Relative Bioavailability to an oral solution (Dose-adjusted)				
Cmax Ratio (Test/Solution)	9.46%	7.72%	5.71%	8.02%
AUC <sub>last</sub> Ratio (Test/Solution)	54.58%	36.05%	52.67%	62.37%
AUC <sub>∞</sub> Ratio (Test/Solution)	53.41%	35.01%	78.95%	119.2%

N = 8, unless specified otherwise

3. Conclusion

- a. Plasma 6-beta-naltrexol levels provide more consistent indication of bioavailability than Naltrexone.
- b. There is significant release in-vivo in both formulations, as indicated by relative bioavailability based on AUC<sub>∞</sub> ratios. 90 µm coat thickness results in less release than 60 µm. Comparing PI-1463 (Opt #1) with PI-1461 (POC), the coating of morphine/NaCl/Kadian ER coat on top of Naltrexone pellet causes more than three-fold increase in NT release.
- c. 7-day duration of study allows 6-beta-naltrexol to return to baseline.
- d. There is clearly no in vitro/in vivo correlation regarding NT release, using conventional buffer system. In vitro dissolution shows 0% NT release at the end of 72 hours, but in vivo data reveals significant NT release.

IV. Optimization Studies #2 and #3, Morphine sulfate and Naltrexone HCl 60 mg/2.4 mg (20-778-1N and 20-779-1N)

	PI-1465		PI-1466	
	mg/unit	Percent	mg/unit	Percent
Sealed-coated sugar spheres				
Sugar spheres (#20-25 mesh)	52.1	16.0	53.1	14.6
Ethylcellulose N50	3.9	1.2	3.98	1.1
Mag Stearate	1.6	0.5	1.6	0.4
Dibutyl Sebacate	0.4	0.1	0.4	0.1
Talc	10	3.1	10.27	2.8
Subtotal	68.0	20.9	69.4	19.0
Naltrexone cores				
Sealed sugar spheres	(68.0)	(20.9)	(69.4)	(19.0)
Naltrexone HCl	2.4	0.74	2.4	0.66
HPC (Klucel LF)	0.5	0.2	0.5	0.1
Citric acid	0.01	0.0031	0.01	0.0027
Ascorbic acid	0.01	0.0031	0.01	0.0027
Butylated Hydroxyanisole	0.01	0.0031	0.01	0.0027
Talc	1.4	0.4	1.43	0.4
Subtotal	72.3	22.3	73.7	20.2
Naltrexone pellets				
Naltrexone cores	(144.7)	(44.5)	(147.4)	(40.4)
Eudragit RS PO	25.4	7.8	38.7	10.6

-continued

	PI-1465		PI-1466	
	mg/unit	Percent	mg/unit	Percent
Sodium lauryl sulfate	0.9	0.3	1.31	0.4
Dibutyl Sebecate	2.53	0.8	3.87	1.1
Talc	26	8.0	38.7	10.6
Subtotal	199.5	61.4	230.0	63.1
Naltrexone-morphine cores				
Naltrexone pellets	(199.5)	(61.4)	(230.0)	(63.1)
Morphine sulfate	59.3	18.2	59.5	16.3
Sodium chloride	17.5	5.4	20.1	5.5
Hypromellose 2910,	14.2	4.4	15.1	4.1
3 cps				
Subtotal	290.5	89.4	324.7	89.0
Naltrexone-morphine pellets				
Naltrexone-morphine cores	(290.5)	(89.4)	(324.7)	(89.0)
Ethylcellulose N50	11.51	3.5	13.1	3.6
Polyethylene glycol 6000	5.3	1.6	6.1	1.7
Eudragit L100-55	2.1	0.6	2.85	0.8
Diethyl Phthalate	2.4	0.7	2.8	0.8
Talc	13.23	4.1	15.2	4.2
Total	325.0	100.0	364.8	100.0

## A. Method of Preparation—

- Dissolve Ethylcellulose and dibutyl sebacate into ethanol, then disperse talc and magnesium stearate into the solution.
- Spray the dispersion from 1 onto sugar spheres in a Wurster to form seal-coated sugar spheres (25  $\mu\text{m}$  seal coat thickness).
- Dissolve Klucel LF, citric acid, ascorbic acid and butylated hydroxyanisole into 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
- Spray the naltrexone dispersion from 3 onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
- Dissolve Eudragit RS, sodium lauryl sulfate and dibutyl sebacate into ethanol. Disperse talc into the solution.
- Spray the dispersion from 5 onto naltrexone cores from 4 in a Wurster to form naltrexone pellets.
- The Naltrexone pellets are dried at 50° C. for 48 hours.
- Resulting pellets have a Eudragit RS coat thickness of 90  $\mu\text{m}$  for PI-1465 and 120  $\mu\text{m}$  for PI-1466.
- Dissolve sodium chloride and hypromellose into water.
- Dissolve hypromellose into 10:90 mixture of water and ethanol. Disperse morphine sulfate into the solution.
- Spray the solution from 9 followed by the dispersion from 10 onto naltrexone pellets in 7 in a rotor to form naltrexone-morphine cores.
- Dissolve ethylcellulose, PEG 6000, Eudragit L100-55 and diethyl phthalate into ethanol. Disperse talc into the solution.
- Spray the dispersion from 12 onto naltrexone-morphine cores in 11 to form naltrexone-morphine pellets.
- The pellets are filled into capsules.

## B. In-vitro Drug Release—

- Method USP paddle method at 37° C. and 100 rpm  
1 hour in 0.1N HCl, then 72 hours in 0.05M pH 7.5 phosphate buffer  
Results Percent of NT released at 73 hours for PI-1465 = 1%  
Percent of NT released at 73 hours for PI-1466 = 0%
- Method USP paddle method at 37° C. and 100 rpm  
72 hrs in 0.2% Triton X-100/0.2% sodium acetate/  
0.002N HCl, pH 5.5  
The data is shown in FIG. 13.

## C. In-vivo Study #1

This is a single-dose, open-label, single-period study in which two groups of eight subjects received one dose of either PI-1465 or PI-1466 under fasting condition. Blood samples were drawn prior to dose administration and at 0.5 to 168 hours post-dose. Limits of quantitation are 4.00 pg/mL for naltrexone and 0.250 pg/mL for 6-beta-naltrexol. The data is shown in FIGS. 14-15.

## 2. Summary of Pharmacokinetics Parameters

	6-beta-Naltrexol		Naltrexone	
	PI-1465	PI-1466	PI-1465	PI-1466
T <sub>max</sub> (hr)	58.51	79.50	50.30 (N = 7)	45.17 (N = 3)
C <sub>max</sub> (pg/mL)	1060	72.6	139.3	46.2
AUC <sub>last</sub> (pg * h/mL)	54693	23473	3713	744
AUC <sub>∞</sub> (pg * h/mL)	56260	23940	7213 (N = 4)	5943 (N = 2)
T <sub>1/2</sub> (hr)	20.90	15.09	16.47 (N = 4)	34.10 (N = 2)
Relative Bioavailability to an oral solution (Dose-adjusted)				
C <sub>max</sub> Ratio (Test/Solution)	4.31%	1.97%	4.72%	1.57%
AUC <sub>last</sub> Ratio (Test/Solution)	26.58%	11.41%	41.60%	8.34%
AUC <sub>∞</sub> Ratio (Test/Solution)	26.45%	11.26%	75.38%	62.11%

N = 8, unless specified otherwise

## 3. Conclusions

- Presence of surfactant in the dissolution medium (second in-vitro drug release method) provides better in-vitro-in-vivo-correlation than buffer alone (first in-vitro drug release method).
- Kadian NT pellets (additional layering of NaCl/morphine/Kadian ER coat on top of naltrexone pellets) had a higher release of naltrexone in vivo than Naltrexone pellets alone. PI-1465 containing the seal coat and the same naltrexone pellet coat thickness as PI-1466 from POC without seal coat (90  $\mu\text{m}$ ), had more than 5 times more release of naltrexone. Even an increase in Naltrexone pellet coat thickness to 120  $\mu\text{m}$  (PI-1466) still gave twice the release of naltrexone.

## D. In-vivo Study #2

This is a single-dose, open-label, single-period study in which four groups of four healthy subjects received a single dose of either PI-1465 or PI-1466 under either fasting or fed conditions. Blood samples were drawn prior to dose administration and at 0.5 to 168 hours post-dose. Limits of quantitation are 4.00 pg/mL for naltrexone and 0.250 pg/mL for 6-beta-naltrexol. The data is shown in FIGS. 16-17.



## 43

1. Summary of Pharmacokinetic Parameters  
a. Naltrexone

	PI-1465		PI-1466	
	Fast	Fed	Fast	Fed
Tmax (hr)	72.00	26.67 (N = 3)	60.00 (N = 2)	32.00 (N = 3)
Cmax (pg/mL)	107.3	279.3	35.73	262
AUC <sub>last</sub> (pg * h/mL)	2825	4135	1319	4611
AUC <sub>∞</sub> (pg * h/mL)	3593 (N = 1)	6787 (N = 2)	3651 (N = 2)	—
T <sub>1/2</sub> (hr)	15.26 (N = 1)	20.98 (N = 2)	24.75 (N = 2)	—
Relative Bioavailability to an oral solution(Dose-adjusted)				
Cmax Ratio (Test/Solution)	3.64%	9.47%	1.21%	8.89%
AUC <sub>last</sub> Ratio (Test/Solution)	31.65%	46.33%	14.78%	51.66%
AUC <sub>∞</sub> Ratio (Test/Solution)	37.55%	70.93%	38.15%	—

N = 4, unless specified otherwise

## b. 6-beta-Naltrexol levels

	PI-1465		PI-1466	
	Fast	Fed	Fast	Fed
Tmax (hr)	69.00	29.00	69.00	36.00
Cmax (pg/mL)	1280	3787	873	2680
AUC <sub>last</sub> (pg * h/mL)	53307	120400	47140	78533
AUC <sub>∞</sub> (pg * h/mL)	53547	122533	47920	78867
T <sub>1/2</sub> (hr)	19.21	18.17	20.69	20.19
Relative Bioavailability to an oral solution				
Cmax Ratio (Test/Solution)	5.20%	15.39%	3.55%	10.89%
AUC <sub>last</sub> Ratio (Test/Solution)	25.90%	58.50%	22.91%	38.16%
AUC <sub>∞</sub> Ratio (Test/Solution)	25.17%	57.61%	22.53%	37.08%

N = 4, unless specified otherwise

## 2. Conclusion

a. There is significant food effect, where the lag time was reduced and NT release was increased in the presence of food. There is a two-fold increase in NT release for PI-1465 and 1.5-fold increase for PI-1466 in the presence of food.

b. There is some subject group variability. Comparing PI-1466 in both in-vivo study #1 and #2, although the same product was used, for fasting condition, there was a two-fold difference in AUC. For PI-1465, the AUC was similar between the two studies.

## V. Optimization Study #4, Morphine sulfate and Naltrexone HCl 60 mg/4.8 mg (20-780-1N)

	PI-1495		PI-1496	
	mg/unit	Percent	mg/unit	Percent
Sealed-coated sugar spheres				
Sugar spheres (#25-30 mesh)	37.2	11.7	37.1	11.9
Ethylcellulose N50	6.2	1.9	6.2	2.0

## 44

-continued

	PI-1495		PI-1496	
	mg/unit	Percent	mg/unit	Percent
Mag Stearate	2.5	0.8	2.5	0.8
DBS	0.6	0.2	0.6	0.2
Talc	15.5	4.9	15.5	5.0
Subtotal	62.0	19.4	61.9	19.9
Naltrexone cores				
Sealed sugar spheres	(62.0)	(19.4)	(61.9)	(19.9)
Naltrexone HCl	4.8	1.50	4.8	1.54
HPC (Klucel LF)	0.9	0.3	0.9	0.3
Ascorbic acid	0.5	0.2	0.5	0.2
Talc	2.27	0.7	2.24	0.7
Subtotal	70.5	22.1	70.3	22.6
Naltrexone pellets				
Naltrexone cores	(70.5)	(22.1)	(70.3)	(22.6)
Eudragit RS PO	53.3	16.7	53.3	17.1
SLS	1.8	0.6	1.8	0.6
DBS	5.36	1.7	5.36	1.7
Talc	52.1	16.3	52.1	16.8
Subtotal	183.0	57.4	182.9	58.8
Naltrexone-morphine cores				
Naltrexone pellets	(183.0)	(57.4)	(182.9)	(58.8)
Morphine sulfate	59.9	18.8	59.7	19.2
Sodium chloride	11.2	3.5		
HPC (Klucel LF)	7.3	2.3	4.76	1.5
HPMC, 3 cps			7.6	2.4
Subtotal	261.4	82.0	255.0	82.0
Naltrexone-morphine pellets				
Naltrexone-morphine cores	(261.4)	(82.0)	(255.0)	(82.0)
Ethylcellulose N50	19.81	6.2	19.31	6.2
PEG 6000	9.16	2.9	8.9	2.9
Eudragit L100-55	4.3	1.3	4.2	1.4
DEP	4.12	1.3	4	1.3
Talc	20.13	6.3	19.62	6.3
Total	319.0	100.0	311.0	100.0

## A. Method of Preparation—

1. Dissolve Ethylcellulose and dibutyl sebacate into ethanol, then disperse talc and magnesium stearate into the solution.
2. Spray the dispersion from 1 onto sugar spheres in a Wurster to form seal-coated sugar spheres (50 μm seal coat).
3. Dissolve Klucel LF and ascorbic acid into 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
4. Spray the naltrexone dispersion from 3 onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
5. Dissolve Eudragit RS, sodium lauryl sulfate and dibutyl sebacate into ethanol. Disperse talc into the solution.
6. Spray the dispersion from 5 onto naltrexone cores from 4 in a Wurster to form naltrexone pellets.
7. The Naltrexone pellets are dried at 50° C. for 48 hours.
8. Resulting pellets have a Eudragit RS coat thickness of 150 μm for both PI-1495 PI-1496.
9. (Only for PI-1495) Dissolve sodium chloride and hypromellose into water.
10. Dissolve hypromellose into 10:90 mixture of water and ethanol. Disperse morphine sulfate into the solution.
11. (Only for PI-1495) Spray the solution from 9 followed by the dispersion from 10 onto naltrexone pellets in 7 in a rotor to form naltrexone-morphine cores.

45

12. (Only for PI-1496) Spray the dispersion from 10 onto naltrexone pellets in 7 in a rotor to form naltrexone-morphine cores.
13. Dissolve ethylcellulose, PEG 6000, Eudragit L100-55 and diethyl phthalate into ethanol. Disperse talc into the solution.
14. Spray the dispersion from 12 onto naltrexone-morphine cores in 11 or 12 to form naltrexone-morphine pellets.
15. The pellets are filled into capsules.

B. In-vitro Drug Release—

1.	Method	USP paddle method at 37° C. and 100 rpm 1 hour in 0.1N HCl, then 72 hours in 0.05M pH 7.5 phosphate buffer
	Results	Percent of NT released at 73 hours for PI-1495 = 0% Percent of NT released at 73 hours for PI-1496 = 0%
2.	Method	USP paddle method at 37° C. and 100 rpm 72 hrs in 0.2% Triton X-100/0.2% sodium acetate/ 0.002N HCl, pH 5.5
	Results	Percent of NT released at 73 hours for PI-1495 = 0% Percent of NT released at 73 hours for PI-1496 = 0%

C. In-vivo Study

This is a single-dose, open-label, two period study in which two groups of eight subjects received one dose of either PI-1495 or PI-1496. Each subject received an assigned treatment sequence based on a randomization schedule under fasting and non-fasting conditions. Blood samples were drawn prior to dose administration and at 0.5 to 168 hours post-dose. Limits of quantitation are 4.00 pg/mL for naltrexone and 0.250 pg/mL for 6-beta-naltrexol. The data is shown in FIGS. 18-19.

2. Summary of Pharmacokinetic Parameters

a. Naltrexone

	PI-1495		PI-1496	
	Fast	Fed	Fast	Fed
Tmax (hr)	54.00 (N = 2)	14.34 (N = 3)	55.20 (N = 5)	41.60 (N = 5)
Cmax (pg/mL)	8.53	6.32 (N = 7)	24.23 (N = 7)	45.67 (N = 7)
AUC <sub>last</sub> (pg * h/mL)	100.8	75.9 (N = 7)	500.6 (N = 7)	1265 (N = 7)
AUC <sub>∞</sub> (pg * h/mL)	—	—	2105.3 (N = 2)	3737 (N = 2)
T <sub>1/2</sub> (hr)	—	—	44.56 (N = 2)	33.17 (N = 2)
Relative Bioavailability to an oral solution (Dose-adjusted)				
Cmax Ratio (Test/Solution)	0.29%	0.21%	0.82%	1.55%
AUC <sub>last</sub> Ratio (Test/Solution)	1.13%	0.85%	5.61%	14.17%
AUC <sub>∞</sub> Ratio (Test/Solution)	—	—	22.0%	39.1%

N = 8, unless specified otherwise

b. 6-beta-Naltrexol levels

	PI-1495		PI-1496	
	Fast	Fed	Fast	Fed
Tmax (hr)	69.00	41.44 (N = 7)	70.51	67.63
Cmax (pg/mL)	116.3	151.7 (N = 7)	303.3	656.7

46

-continued

	PI-1495		PI-1496	
	Fast	Fed	Fast	Fed
AUC <sub>last</sub> (pg * h/mL)	5043	7332 (N = 7)	14653	27503
AUC <sub>∞</sub> (pg * h/mL)	5607	8449 (N = 6)	14930	27827
T <sub>1/2</sub> (hr)	20.97	16.69 (N = 7)	16.29	22.59
Relative Bioavailability to an oral solution (Dose-adjusted)				
Cmax Ratio (Test/Solution)	0.47%	0.62%	1.23%	2.67%
AUC <sub>last</sub> Ratio (Test/Solution)	2.45%	3.45%	7.12%	13.36%
AUC <sub>∞</sub> Ratio (Test/Solution)	2.64%	3.97%	7.02%	13.08%

N = 8, unless specified otherwise

3. Conclusion

- a. Kadian NT pellets with naltrexone pellet coat thickness of 150 μm had comparable naltrexone release as NT pellets with 90 μm coat thickness. This comparable NT release may also be attributed from the presence of 50 μm seal coat on the sugar spheres used in Kadian NT pellets.
- b. Significant NT sequestering was observed, both at fasting (>97%) and fed states (>96%).
- c. Kadian NT pellets containing sodium chloride immediately above the naltrexone pellet coat (PI-1495) had half the release of naltrexone compared to Kadian NT pellet without sodium chloride (PI-1496), consistent with in vitro results.
- d. There is again food effect observed. Lag time was significantly reduced.

VI. Optimization Study #5, Morphine sulfate and Naltrexone HCl 60 mg/2.4 mg (20-903-AU)

	PI-1510	
	mg/unit	Percent
<u>Sealed sugar spheres</u>		
Sugar spheres (#25-30 mesh)	39.9	12.2
Ethylcellulose N50	6.5	2.0
Mag Stearate	2.6	0.8
DBS	0.7	0.2
Talc	16.7	5.1
Subtotal	66.4	20.3
<u>Naltrexone cores</u>		
Sealed sugar spheres	(66.4)	(20.3)
Naltrexone HCl	2.4	0.73
HPC (Klucel LF)	0.5	0.1
Ascorbic acid	0.2	0.1
Talc	1.1	0.4
Subtotal	70.6	21.6
<u>Naltrexone pellets</u>		
Naltrexone cores	(70.6)	(21.6)
Eudragit RS PO	53.0	16.2
SLS	1.8	0.6
DBS	5.3	1.6
Talc	53.0	16.2
Subtotal	183.7	56.2

-continued

	PI-1510	
	mg/unit	Percent
<u>Naltrexone-morphine cores</u>		
Naltrexone pellets	(183.7)	(56.2)
Morphine sulfate	60.1	18.4
Sodium chloride	12.5	3.8
HPC (Klucel LF)	6.2	1.9
Subtotal	262.4	80.2
<u>Naltrexone-morphine pellets</u>		
Naltrexone-morphine cores	(262.4)	(80.2)
Ethylcellulose N50	22.9	7.0
PEG 6000	10.6	3.2
Eudragit L100-55	5.0	1.5
DEP	4.7	1.5
Talc	21.5	6.6
Total	327.1	100.0

B. Method of Preparation—

1. Dissolve Ethylcellulose and dibutyl sebacate into ethanol, then disperse talc and magnesium stearate into the solution.
2. Spray the dispersion from 1 onto sugar spheres in a Wurster to form seal-coated sugar spheres (50 µm seal coat).
3. Dissolve Klucel LF and ascorbic acid into 20:80 mixture of water and ethanol. Disperse naltrexone HCl and talc into the solution.
4. Spray the naltrexone dispersion from 3 onto seal-coated sugar spheres from 2 in a Wurster to form naltrexone cores.
5. Dissolve Eudragit RS, sodium lauryl sulfate and dibutyl sebacate into ethanol. Disperse talc into the solution.
6. Spray the dispersion from 5 onto naltrexone cores from 4 in a Wurster to form naltrexone pellets.
7. The Naltrexone pellets are dried at 50° C. for 48 hours.
8. Resulting pellets have a Eudragit RS coat thickness of 150 µm.
9. Dissolve sodium chloride and hypromellose into water.
10. Dissolve hypromellose into 10:90 mixture of water and ethanol. Disperse morphine sulfate into the solution.
11. Spray the solution from 9 followed by the dispersion from 10 onto naltrexone pellets in 7 in a rotor to form naltrexone-morphine cores.
12. Dissolve ethylcellulose, PEG 6000, Eudragit L100-55 and diethyl phthalate into ethanol. Disperse talc into the solution.
13. Spray the dispersion from 12 onto naltrexone-morphine cores in 11 or 12 to form naltrexone-morphine pellets.
14. The pellets are filled into capsules.

B. In-vitro Drug Release—

1.	Method	USP paddle method at 37° C. and 100 rpm 1 hour in 0.1N HCl, then 72 hours in 0.05M pH 7.5 phosphate buffer
	Results	Percent of NT released at 73 hours for = 0%
2.	Method	USP paddle method at 37° C. and 100 rpm 72 hrs in 0.2% Triton X-100/0.2% sodium acetate/ 0.002N HCl, pH 5.5
	Results	Percent of NT released at 73 hours = 0%

C. In-vivo Study

This is a single-dose, open-label, two period study in which eight subjects were randomized to receive one dose of PI-1510 under either fasted or fed state during Study Period 1 and alternate fasted or fed state for Study Period 2. Blood samples were drawn prior to dose administration and at 0.5 to 168 hours post-dose. Limits of quantitation are 4.00 pg/mL for naltrexone and 0.250 pg/mL for 6-beta-naltrexol. The data is shown in FIGS. 20 and 21.

2. Summary of Pharmacokinetic Parameters

a. 6-beta-Naltrexol Levels

	PI-1510	
	Fast	Fed
Tmax (hr)	45.00 (N = 6)	57.29 (N = 7)
Cmax (pg/mL)	16.1	25.0
AUC <sub>last</sub> (pg * h/mL)	609.2	1057
AUC <sub>∞</sub> (pg * h/mL)	1233	1431 (N = 6)
T <sub>1/2</sub> (hr)	17.36	17.48 (N = 6)
Relative Bioavailability to an oral solution (Dose-adjusted)		
Cmax Ratio (Test/Solution)	0.44%	0.68%
AUC <sub>last</sub> Ratio (Test/Solution)	1.97%	3.42%
AUC <sub>∞</sub> Ratio (Test/Solution)	3.86%	4.49%

N = 8, unless specified otherwise

3. Conclusion

- a. PI-1510 and PI-1495 are comparable. The reduction in naltrexone loading in the pellets (from 1.5% in PI-1495 to 0.7% in PI-1510) does not seem to affect NT release.
- b. Significant NT sequestering was observed, both at fasting (>96%) and fed states (>95%).
- c. The food effect observed was modest in terms of total NT release. However, the lag time was significantly reduced in the presence of food. There were subjects with multiple peaks of release.

VII. Summary of NT Release From All in-vivo Studies

BA (Cmax)=Relative bioavailability based on Cmax=Dose-adjusted ratio of Cmax (NT/KNT pellet) to (Cmax)(NT soln)

BA (AUC last)=Relative bioavailability based on AUC last=Dose-adjusted ratio of AUC last (NT/KNT pellet) to AU

BA (AUC inf)=Relative bioavailability based on AUC inf=Dose-adjusted ratio of AUC inf (NT/KNT pellet)

Total in-vivo cumulative NT release can be extrapolated from BA (AUC inf) calculations from 6-beta-Naltrexol plasma levels

	BA (Cmax) (%)	BA (AUC last) (%)	BA (AUC inf) (%)
POC			
PI-1460 Fast			
Avg ± SD	1.2 ± 0.9	5.1 ± 3.1	
Range	0.32-2.99	1.92-10.65	
PI-1461 Fast			
Avg ± SD	3.1 ± 2.4	15.8 ± 11.9	
Range	0.7-10.3	2.8-49.2	

-continued

	BA (Cmax) (%)	BA (AUC last) (%)	BA (AUC inf) (%)
<b>OPTIM. #1</b>			
<b>PI-1462 Fast</b>			
Avg ± SD	9.5 ± 2.8	54.6 ± 21.0	53.4 ± 20.6
Range	5.7-13.0	26.3-86.3	25.6-84.4
<b>PI-1463 Fast</b>			
Avg ± SD	7.7 ± 3.7	36.1 ± 18.2	35.0 ± 17.7
Range	0.8-12.4	3.9-59.2	3.8-57.3
<b>OPTIM. #2 and #3</b>			
<b>PI-1465</b>			
<b>Fast 1</b>			
Avg ± SD	4.3 ± 6.2	26.6 ± 35.4	26.4 ± 35.0
Range	0.1-18.6	0.1-111.6	0.1-110.5
<b>Fast 2</b>			
Avg ± SD	5.2 ± 3.9	25.9 ± 15.7	25.2 ± 15.2
Range	1.8-10.5	9.6-41.5	9.4-40.2
<b>Fed</b>			
Avg ± SD	15.4 ± 12.5	58.5 ± 34.6	57.6 ± 34.4
Range	1.4-31.2	11.9-90.6	11.5-90.6
<b>OPTIM. #2 and #3</b>			
<b>PI-1466</b>			
<b>Fast 1</b>			
Avg ± SD	2.0 ± 2.3	11.4 ± 11.8	11.3 ± 11.4
Range	0.2-5.9	1.1-30.0	11.1-29.1
<b>Fast 2</b>			
Avg ± SD	3.6 ± 3.9	22.9-25.6	22.5 ± 24.9
Range	0.5-8.6	1.8-57.4	1.8-56.1
<b>Fed</b>			
Avg ± SD	10.9 ± 12.7	38.2 ± 40.0	37.1 ± 38.9
Range	0.3-28.5	1.7-90.3	1.6-87.7
<b>OPTIM. #4</b>			
<b>PI-1495</b>			
<b>Fast</b>			
Avg ± SD	0.5 ± 0.5	2.5 ± 2.3	2.6 ± 2.4
Range	0.1-1.4	5.9-0.3	0.3-5.7
<b>Fed</b>			
Avg ± SD	3.0 ± 6.7	10.2 ± 19.4	11.3 ± 20.0
Range	0.1-19.4	0.2-57.0	0.2-55.4
<b>Fed (-Subject 1)</b>			
Avg ± SD	0.6 ± 0.9	3.6 ± 4.9	4.0 ± 5.0
Range	0.1-2.5	0.2-13.8	0.2-13.4
<b>PI-1496</b>			
<b>Fast</b>			
Avg ± SD	1.2 ± 0.9	7.1 ± 4.6	7.0 ± 4.6
Range	0.1-2.7	0.6-14.2	0.6-14.5
<b>Fed</b>			
Avg ± SD	2.7 ± 2.9	13.4 ± 12.6	13.1 ± 12.3
Range	0.1-7.6	0.1-31.6	0.4-30.7
<b>OPTIM. #5</b>			
<b>PI-1510</b>			
<b>Fast</b>			
Avg	0.4	2.0	3.9
<b>Fed</b>			
Avg	0.7	3.4	4.5

While the present invention has been described in terms of the preferred embodiments, it is understood that variations and modifications will occur to those skilled in the art. Therefore, it is intended that the appended claims cover all such equivalent variations that come within the scope of the invention as claimed.

What is claimed is:

1. A method of treating moderate to severe pain in a human, the method comprising: orally administering to said human an intact composition in an amount sufficient to treat the moderate to severe pain;
  - a. a water-soluble core;
  - b. an opioid antagonist comprising layer coating the core, wherein the opioid antagonist is selected from the group consisting of naltrexone, naloxone, nalmefene, cyclazacine, levallorphan, salts of these molecules, and combinations thereof;
  - c. a sequestering polymer layer coating the opioid antagonist comprising layer;
  - d. an osmotic pressure regulating agent comprising layer coating the sequestering polymer layer, wherein the osmotic pressure regulating agent is selected from the group consisting of sodium chloride, sodium bromide, sodium iodide, hydroxypropyl methyl cellulose, and combinations thereof;
  - e. an opioid agonist comprising layer coating the osmotic pressure regulating agent comprising layer, wherein the opioid agonist is selected from the group consisting of morphine, oxycodone, hydrocodone, hydromorphone, dihydrocodeine, codeine, dihydromorphone, buprenorphine, salts of these molecules, and combinations thereof; and
  - f. a controlled release layer coating the opioid agonist comprising layer;
 wherein the sequestering polymer layer comprises copolymers of acrylic and methacrylic acid esters with quaternary ammonium groups, a surfactant in an amount from 1.6% to 6.3% of the copolymers of acrylic and methacrylic acid esters with quaternary ammonium groups on a weight-to-weight basis, and talc in an amount of from 75% to 125% of the copolymers of acrylic and methacrylic acid esters with quaternary ammonium groups on a weight-to-weight basis;
  - wherein the surfactant is selected from the group consisting of sodium lauryl sulfate, sodium docusate, dioctyl sodium sulphosuccinate, sodium lauryl sarcosinate, sodium methyl cocyl taurate, magnesium lauryl sulfate, dioctyl sodiumsulfosuccinate, sodiumdodecylbenzene sulfonate, and combinations thereof; and
  - wherein when tested by a testing method, the composition sequesters at least 80% of the opioid antagonist as determined at 73 hours from the start of the testing method,
  - wherein the testing method comprises first placing the composition in 500 mL of a 0.1 N HCl solution for 1 hour at 37° C. using USP paddle method, 100 rotations per minute, and then placing the composition in 500 mL of a pH 7.5, 0.05 M phosphate buffer, for 72 hours at 37° C. using USP paddle method, 100 rotations per minute, and then determining the amount of the opioid antagonist sequestered.
2. The method of claim 1, wherein in the multi-layer pellets the surfactant is not comprised in the following:
  - the water-soluble core,
  - the opioid antagonist comprising layer,
  - the osmotic pressure regulating agent comprising layer,
  - the opioid agonist comprising layer, and
  - the controlled release layer.
3. The method of claim 1, wherein in the multi-layer pellets the opioid agonist comprising layer further comprises hydroxypropyl cellulose.

## 51

4. The method of claim 1, wherein in the multi-layer pellets the surfactant is not comprised in the following:

the water-soluble core,  
the opioid antagonist comprising layer,  
the osmotic pressure regulating agent comprising layer,  
the opioid agonist comprising layer, and  
the controlled release layer; and  
wherein the opioid agonist comprising layer further comprises hydroxypropyl cellulose.

5. The method of claim 1, wherein in the multi-layer pellets the osmotic pressure regulating agent comprises sodium chloride;

the surfactant is sodium lauryl sulfate; and  
the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

6. The method of claim 1, wherein in the multi-layer pellets the opioid antagonist is the salt of naltrexone;  
the osmotic pressure regulating agent comprises sodium chloride;

the surfactant is sodium lauryl sulfate; and  
the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

7. The method of claim 1, wherein in the multi-layer pellets the opioid antagonist is naltrexone HCl;  
the osmotic pressure regulating agent comprises sodium chloride;

the surfactant is sodium lauryl sulfate; and  
the opioid agonist comprising layer further comprises hydroxypropyl cellulose.

8. The method of claim 1, wherein in the multi-layer pellets,

the opioid antagonist is naltrexone HCl;  
the osmotic pressure regulating agent comprises sodium chloride;

the surfactant is sodium lauryl sulfate; and  
the opioid agonist comprising layer comprises a salt of morphine and further comprises hydroxypropyl cellulose.

9. The method of claim 1, wherein in the multi-layer pellets the opioid antagonist is naltrexone HCl;  
the osmotic pressure regulating agent comprises sodium chloride; and

the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

10. The method of claim 1, wherein in the multi-layer pellets

the opioid antagonist is naltrexone HCl;  
the surfactant is sodium lauryl sulfate; and  
the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

11. The method of claim 1, wherein in the multi-layer pellets

the opioid antagonist is naltrexone HCl;  
the surfactant is sodium lauryl sulfate;  
the osmotic pressure regulating agent comprises sodium bromide; and  
the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

12. The method of claim 1, wherein in the multi-layer pellets

the opioid antagonist is naltrexone HCl;  
the surfactant is sodium lauryl sulfate;  
the osmotic pressure regulating agent comprises sodium iodide; and  
the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

## 52

13. The method of claim 1, wherein in the multi-layer pellets,

the opioid antagonist is naltrexone HCl;  
the surfactant is sodium lauryl sulfate;  
the osmotic pressure regulating agent comprises hydroxypropyl methyl cellulose; and  
the opioid agonist comprising layer comprises morphine sulfate and further comprises hydroxypropyl cellulose.

14. The method of claim 1, wherein in the multi-layer pellets the surfactant is not comprised in the following:

the water-soluble core,  
the opioid antagonist comprising layer,  
the osmotic pressure regulating agent comprising layer,  
the opioid agonist comprising layer, and  
the controlled release layer.

15. The method of claim 1, wherein in the multi-layer pellets the controlled release layer comprises a cellulose; a polyalkylene glycol; an anionic copolymer based on methacrylic acid and ethylacrylate; and a plasticizer.

16. The method of claim 1, wherein in the multi-layer pellets the controlled release layer comprises a cellulose; a polyalkylene glycol; an anionic copolymer based on methacrylic acid and ethylacrylate; and a plasticizer.

17. The method of claim 1, wherein the composition does not comprise a bittering agent, a gelling agent, an irritant, or an emetic.

18. The method of claim 1, wherein in the multi-layer pellets opioid antagonist is not comprised in the following:

the water-soluble core,  
the sequestering polymer layer,  
the osmotic pressure regulating agent comprising layer,  
the opioid agonist comprising layer, and  
the controlled release layer.

19. The method of claim 1, wherein in the multi-layer pellets opioid agonist is not comprised in the following:

the water-soluble core,  
the opioid antagonist comprising layer,  
the sequestering polymer layer,  
the osmotic pressure regulating agent comprising layer, and  
the controlled release layer.

20. A method of treating moderate to severe pain in a human, the method comprising: orally administering to said human an intact composition in an amount sufficient to treat the moderate to severe pain;

wherein the composition comprises a plurality of multi-layer pellets comprising:

- a water-soluble core;
- an opioid antagonist comprising layer coating the core, wherein the opioid antagonist is selected from the group consisting of naltrexone, naloxone, nalmeferene, cyclazacine, levallorphan, salts of these molecules, and combinations thereof;
- a sequestering polymer layer coating the opioid antagonist comprising layer;
- an osmotic pressure regulating agent comprising layer coating the sequestering polymer layer, wherein the osmotic pressure regulating agent is selected from the group consisting of sodium chloride, sodium bromide, sodium iodide, hydroxypropyl methyl cellulose, and combinations thereof;
- an opioid agonist comprising layer coating the osmotic pressure regulating agent comprising layer, wherein the opioid agonist is selected from the group consisting of oxycodone, a salt of oxycodone, and combinations thereof; and

f. a controlled release layer coating the opioid agonist comprising layer;

wherein the sequestering polymer layer comprises

copolymers of acrylic and methacrylic acid esters with quaternary ammonium groups, a surfactant in an amount from 1.6% to 6.3% of the copolymers of acrylic and methacrylic acid esters with quaternary ammonium groups on a weight-to-weight basis, and talc in an amount of from 75% to 125% of the copolymers of acrylic and methacrylic acid esters with quaternary ammonium groups on a weight-to-weight basis;

wherein the surfactant is selected from the group consisting of sodium lauryl sulfate, sodium docusate, dioctyl sodium sulphosuccinate, sodium lauryl sarcosinate, sodium methyl cocyl taurate, magnesium lauryl sulfate, dioctyl sodiumsulfo succinate, sodiumdodecylbenzene sulfonate, and combinations thereof; and

wherein when tested by a testing method, the composition sequesters at least 80% of the opioid antagonist as determined at 73 hours from the start of the testing method,

wherein the testing method comprises first placing the composition in 500 mL of a 0.1 N HCl solution for 1 hour at 37° C. using USP paddle method, 100 rotations per minute, and then placing the composition in 500 mL of a pH 7.5, 0.05 M phosphate buffer, for 72 hours at 37° C. using USP paddle method, 100 rotations per minute, and then determining the amount of the opioid antagonist sequestered.

21. The method of claim 1, wherein the opioid antagonist is naltrexone or a salt thereof.

22. The method of claim 21, wherein the opioid agonist is morphine or a salt thereof.

23. The method of claim 21, wherein the opioid agonist is oxycodone or a salt thereof.

24. The method of claim 21, wherein the opioid agonist is hydrocodone or a salt thereof.

25. The method of claim 21, wherein the opioid agonist is hydromorphone or a salt thereof.

26. The method of claim 21, wherein the opioid agonist is dihydrocodeine or a salt thereof.

27. The method of claim 21, wherein the opioid agonist is codeine or a salt thereof.

28. The method of claim 21, wherein the opioid agonist is dihydromorphine or a salt thereof.

29. The method of claim 21, wherein the opioid agonist is buprenorphine or a salt thereof.

30. The method of claim 1, wherein the opioid antagonist is naloxone or a salt thereof.

31. The method of claim 30, wherein the opioid agonist is morphine or a salt thereof.

32. The method of claim 30, wherein the opioid agonist is oxycodone or a salt thereof.

33. The method of claim 30, wherein the opioid agonist is hydrocodone or a salt thereof.

34. The method of claim 30, wherein the opioid agonist is hydromorphone or a salt thereof.

35. The method of claim 30, wherein the opioid agonist is dihydrocodeine or a salt thereof.

36. The method of claim 30, wherein the opioid agonist is codeine or a salt thereof.

37. The method of claim 30, wherein the opioid agonist is dihydromorphine or a salt thereof.

38. The method of claim 30, wherein the opioid agonist is buprenorphine or a salt thereof.

39. The method of claim 1, wherein the opioid antagonist is nalmefene or a salt thereof.

40. The method of claim 39, wherein the opioid agonist is morphine or a salt thereof.

41. The method of claim 39, wherein the opioid agonist is oxycodone or a salt thereof.

42. The method of claim 39, wherein the opioid agonist is hydrocodone or a salt thereof.

43. The method of claim 39, wherein the opioid agonist is hydromorphone or a salt thereof.

44. The method of claim 39, wherein the opioid agonist is dihydrocodeine or a salt thereof.

45. The method of claim 39, wherein the opioid agonist is codeine or a salt thereof.

46. The method of claim 39, wherein the opioid agonist is dihydromorphine or a salt thereof.

47. The method of claim 39, wherein the opioid agonist is buprenorphine or a salt thereof.

48. The method of claim 1, wherein the opioid antagonist is cyclazacine or a salt thereof.

49. The method of claim 48, wherein the opioid agonist is morphine or a salt thereof.

50. The method of claim 48, wherein the opioid agonist is oxycodone or a salt thereof.

51. The method of claim 48, wherein the opioid agonist is hydrocodone or a salt thereof.

52. The method of claim 48, wherein the opioid agonist is hydromorphone or a salt thereof.

53. The method of claim 48, wherein the opioid agonist is dihydrocodeine or a salt thereof.

54. The method of claim 48, wherein the opioid agonist is codeine or a salt thereof.

55. The method of claim 48, wherein the opioid agonist is dihydromorphine or a salt thereof.

56. The method of claim 48, wherein the opioid agonist is buprenorphine or a salt thereof.

57. The method of claim 1, wherein the opioid antagonist is levallorphan or a salt thereof.

58. The method of claim 57, wherein the opioid agonist is morphine or a salt thereof.

59. The method of claim 57, wherein the opioid agonist is oxycodone or a salt thereof.

60. The method of claim 57, wherein the opioid agonist is hydrocodone or a salt thereof.

61. The method of claim 57, wherein the opioid agonist is hydromorphone or a salt thereof.

62. The method of claim 57, wherein the opioid agonist is dihydrocodeine or a salt thereof.

63. The method of claim 57, wherein the opioid agonist is codeine or a salt thereof.

64. The method of claim 57, wherein the opioid agonist is dihydromorphine or a salt thereof.

65. The method of claim 57, wherein the opioid agonist is buprenorphine or a salt thereof.