



US008188147B2

(12) **United States Patent**
Summar et al.

(10) **Patent No.:** **US 8,188,147 B2**
(45) **Date of Patent:** **May 29, 2012**

(54) **THERAPEUTIC METHODS EMPLOYING NITRIC OXIDE PRECURSORS**

(75) Inventors: **Marshall L. Summar**, Brentwood, TN (US); **Brian W. Christman**, Nashville, TN (US); **Frederick E. Barr**, Nashville, TN (US)

(73) Assignee: **Vanderbilt University**, Nashville, TN (US)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 201 days.

(21) Appl. No.: **12/122,117**

(22) Filed: **May 16, 2008**

(65) **Prior Publication Data**

US 2008/0234379 A1 Sep. 25, 2008

Related U.S. Application Data

(60) Division of application No. 10/785,374, filed on Feb. 24, 2004, which is a continuation-in-part of application No. 09/585,077, filed on Jun. 1, 2000, now Pat. No. 6,743,823, which is a continuation-in-part of application No. 09/323,472, filed on Jun. 1, 1999, now Pat. No. 6,346,382.

(51) **Int. Cl.**
A61K 31/195 (2006.01)

(52) **U.S. Cl.** **514/565**

(58) **Field of Classification Search** 514/565
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

4,196,265 A 4/1980 Koprowski et al.
4,458,066 A 7/1984 Caruthers et al.
4,554,101 A 11/1985 Hopp

4,683,195 A 7/1987 Mullis et al.
4,683,202 A 7/1987 Mullis
4,736,866 A 4/1988 Leder et al.
4,769,331 A 9/1988 Roizman et al.
4,965,188 A 10/1990 Mullis et al.
4,996,236 A * 2/1991 Nakamura et al. 514/659
5,162,215 A 11/1992 Bosselman et al.
5,217,997 A 6/1993 Levere
5,279,833 A 1/1994 Rose
5,286,634 A 2/1994 Stadler et al.
5,286,739 A 2/1994 Kilbourn et al.

(Continued)

FOREIGN PATENT DOCUMENTS

AU 3460571 A 4/1973

(Continued)

OTHER PUBLICATIONS

Akashi et al., "Citrulline, a Novel Compatible Solute in Drought-Tolerant Wild Watermelon Leaves, Is an Efficient Hydroxyl Radical Scavenger," FEBS Lett., vol. 508, No. 3, pp. 438-442 (Nov. 23, 2001). (Abstract).

(Continued)


Primary Examiner — San-Ming Hui

(74) *Attorney, Agent, or Firm* — Jenkins, Wilson, Taylor & Hunt, P.A.

(57) **ABSTRACT**

Isolated polynucleotide molecules and peptides encoded by these molecules are used in the analysis of human carbamyl phosphate synthetase I phenotypes, as well as in diagnostic and therapeutic applications, relating to a human carbamyl phosphate synthetase I polymorphism. By analyzing genomic DNA or amplified genomic DNA, or amplified cDNA derived from mRNA, it is possible to type a human carbamyl phosphate synthetase I with regard to the human carbamyl phosphate synthetase I polymorphism, for example, in the context of diagnosing and treating hepatic veno-occlusive disease (HVOD) associated with bone marrow transplants.

18 Claims, 11 Drawing Sheets



U.S. PATENT DOCUMENTS

5,334,380	A	8/1994	Kilbourn et al.	
5,385,940	A	1/1995	Moskowitz	
5,399,346	A	3/1995	Anderson et al.	
5,489,742	A	2/1996	Hammer et al.	
5,550,024	A	8/1996	Rothschild	
5,550,316	A	8/1996	Mintz	
5,573,933	A	11/1996	Seamark et al.	
5,614,396	A	3/1997	Bradley et al.	
5,625,125	A	4/1997	Bennett et al.	
5,641,484	A	6/1997	Hung et al.	
5,643,567	A	7/1997	Hung et al.	
5,646,008	A	7/1997	Thompson et al.	
5,648,061	A	7/1997	Bernstein et al.	
5,651,964	A	7/1997	Hung et al.	
5,741,957	A	4/1998	Deboer et al.	
5,767,160	A	6/1998	Kaesemeyer	
5,873,359	A	2/1999	Zapol et al.	
5,874,471	A *	2/1999	Waugh	514/563
5,891,459	A	4/1999	Cooke et al.	
6,337,321	B1	1/2002	Cooke et al.	
6,343,382	B2	2/2002	Sciglia	
6,346,382	B1	2/2002	Summar et al.	
6,358,536	B1	3/2002	Thomas	
6,642,208	B2	11/2003	Cooke et al.	
6,646,006	B2	11/2003	Cooke et al.	
6,689,810	B2	2/2004	Martin	
6,743,823	B1	6/2004	Summar et al.	
2001/0056068	A1	12/2001	Chwalisz et al.	
2002/0013288	A1	1/2002	Cooke et al.	
2003/0134332	A1	7/2003	Boykin	
2004/0235953	A1	11/2004	Summar et al.	
2006/0194728	A1	8/2006	Killian et al.	
2007/0026448	A1	2/2007	Ramanathan et al.	
2007/0184554	A1	8/2007	Teuscher et al.	
2009/0197964	A1	8/2009	Summar et al.	
2009/0312423	A1	12/2009	Summar et al.	

FOREIGN PATENT DOCUMENTS

EP	044119	A2	8/1991	
WO	WO 94/16740	A	8/1994	
WO	WO 00/06151		2/2000	
WO	WO 00/73322	A	12/2000	
WO	WO 01/56068	A1	12/2001	
WO	WO 2005/082042		9/2005	

OTHER PUBLICATIONS

Akisu et al., "Protective Effect of Dietary Supplementation with L-Arginine and L-Carnitine on Hypoxia/Reoxygenation-Induced Necrotizing Enterocolitis in Young Mice," *Biology of the Neonate*, vol. 81, pp. 260-265 (2002). (Abstract).

Alonso et al., "Affinity Cleavage of Carbomoyl-Phosphate Synthetase I Localizes Regions of the Enzyme Interacting with the Molecule of ATP that Phosphorylates Carbamate," *Eur. J. Biochem*, pp. 377-384 (1995).

Alves et al., "The SGOT/SGPT Ratio in Alcoholic Liver Disease," *Acta Med Port*, vol. 3, pp. 255-260 (1981).

Amin et al., "Arginine Supplementation Prevents Necrotizing Enterocolitis in the Premature Infant," *Journal of Pediatrics*, vol. 140, No. 4, pp. 425-431 (Apr. 2002).

Australian Patent Office Examination Report corresponding to Australian Patent Application No. 2005216270 dated Nov. 2, 2007.

Barr et al., "Effect of Cardiopulmonary Bypass on Urea Cycle Intermediates and Nitric Oxide Levels After Congenital Heart Surgery," *The Journal of Pediatrics*, vol. 142, No. 1, pp. 26-30 (2003).

Baumgartner et al., "Hyperammonemia with Reduced Ornithine, Arginine and Praline: a New Inborn Error Caused by a Mutation in the Gene Encoding Delta(1)-Pyrroline-5-Carboxylate Synthase," *Hum. Mol. Genet.*, vol. 9, No. 19, pp. 2853-2858 (Nov. 22, 2000). (Abstract).

Benedetto et al., "Increased L-citrulline/L-arginine Plasma Ratio in severe Preclampsia," *Obstet. Gynecol.*, vol. 96, No. 3, pp. 395-399 (Sep. 2000). (Abstract).

BIOSIS Database Accession No. PREV199699075415 [on-line] Biosciences Information service, Philadelphia, PA, US; 1996.

BIOSIS Database Accession No. PREV199799686392 [on-line] Biosciences Information service, Philadelphia, PA, US; 1997.

BIOSIS Database Accession No. PREV199800177848 [on-line] Biosciences Information service, Philadelphia, PA, US; Feb. 1998.

BIOSIS Database Accession No. PREV200200378590 [on-line] Biosciences Information service, Philadelphia, PA, US; May 2002.

BIOSIS Database Accession No. PREV200400055833 [on-line] Biosciences Information service, Philadelphia, PA, US; Dec. 2003.

Boger et al., "An Endogenous Inhibitor of Nitric Oxide Synthase Regulates Endothelial Adhesiveness for Monocytes," *J. Am. Cardiol.*, vol. 36, No. 7, pp. 2287-2295 (Dec. 2000). (Abstract).

Boger et al., "Is Asymmetric Dimethylarginine a Novel Mark of Atherosclerosis?" *Circulation*, vol. 101, No. 14, pp. 160-161 (Apr. 11, 2000). (Abstract).

Bredt, "Endogenous Nitric Oxide Synthesis: Biological Functions and Pathophysiology," *Free Radic. Res.*, vol. 31, No. 6, pp. 577-596 (Dec. 1999). (Abstract).

Bruno et al., "Population Pharmacokinetics and Pharmacokinetic-Pharmacodynamic Relationships for Docetaxel," *Invest. New Drugs*, vol. 19, No. 2, pp. 163-169 (May 2001). (Abstract).

Chan et al., "Asymmetric Dimethylarginine Increases Mononuclear Cell Adhesiveness in Hypercholesterolemic Humans," *Arterioscler. Thromb. Vasc. Biol.*, vol. 20, No. 4, pp. 1040-1046 (Apr. 2000). (Abstract).

Chwalisz et al., "Role of Nitric Oxide in Implantation and Menstruation," *Hum. Reprod.*, vol. 15, Suppl. 3, pp. 96-111 (Aug. 2000). (Abstract).

Cohen et al., "The SGOT/SGPT Ratio—an Indicator of Alcoholic Liver Disease," *Dig. Dis. Sci.*, vol. 24, No. 11, pp. 835-838 (Nov. 1979). (Abstract).

Communication pursuant to Article 94(3) EPC corresponding to European Patent Application No. 05 723 789.3-2123 dated Aug. 4, 2008.

Communication pursuant to Article 96(2) EPC corresponding to European Application No. 05 723 789.3-2123 dated Sep. 14, 2007.

Cooke et al., "Atherogenesis and the Arginine Hypothesis," *Curr. Atheroscler. Rep.*, vol. 3, No. 3, pp. 252-259 (May 2001). (Abstract)

Cooke, "Does ADMA Cause Endothelial Dysfunction?" *Arterioscler. Thromb. Vasc. Biol.*, vol. 20, No. 9, pp. 2032-2037 (Sep. 2000). (Abstract).

Cooke, "The Endothelium: a New Target for Therapy," *Vasc. Med.*, vol. 5, No. 1, pp. 49-53 (2000). (Abstract).

Davies et al., "Idiopathic Hyperammonemia: a Frequently Lethal Complication of Bone Marrow Transplantation," *Bone Marrow Transplantation*, vol. 17, pp. 1119-1125 (Jun. 1996).

Dulak et al., "Nitric Oxide Induces the Synthesis of Vascular Endothelial Growth Factor by Rat Vascular Smooth Muscle Cells," *Arterioscler. Thromb. Vasc.*, vol. 20, No. 3, pp. 659-666 (Mar. 2000). (Abstract).

Durand et al., "Acute Liver Failure in Infancy: a 14-Year Experience of a Pediatric Liver Transplantation Center," *J. Pediatr.*, vol. 139, No. 6, pp. 871-876 (Dec. 2000). (Abstract).

EMBASE Database Accession No. EMB-1990075119 [on-line] Elsevier Science Publishers, Amsterdam, NL; 1990.

EMBASE Database Accession No. EMB-2003218186 [on-line] Elsevier Science Publishers, Amsterdam, NL; May 27, 2003.

Erlich et al., "Recent advances in the polymerase chain reaction," *Science*, vol. 252 pp. 1643-1651 (Jun. 1991).

Fearon et al., "Genetic Analysis of Carbamyl Phosphate Synthetase I Deficiency," *Human Genetics*, vol. 70, No. 3, pp. 207-210 (1985).

Finckh et al., "Prenatal Diagnosis of Carbamoyl Phosphate Synthetase I Deficiency by Identification of a Missense Mutation in CPS1," *Human Mutation*, Wiley-Liss, NY, vol. 12, No. 3, pp. 206-211 (1998).

Gebhardt et al., "Treatment of Cirrhotic Rats with L-Ornithine-L-Aspartate Enhances Urea Synthesis and Lowers Serum Ammonia Levels," *The Journal of Pharmacology and Experimental Therapeutics*, vol. 283, No. 1, pp. 1-6 (1997).

Ghishan et al., "Polymerase Chain Reaction (PCR) Detectable Polymorphisms in the Prenatal Diagnosis of Carbamyl Phosphate Synthetase I Deficiency," *Gastroenterology*, vol. 106, No. 4 Suppl., p. A1028 (1994).

- Guillou et al., "Escherichia coli Carbamoyl-Phosphate Synthetase: Domains of Glutaminase and Synthetase Subunit Interaction," Proc. Natl. Acad. Sci. USA, pp. 8304-8308 (Nov. 14, 1989).
- Gur et al., "Determination of Hepatic Zinc Content in Chronic Liver Disease Due to Hepatitis B Virus," Hepatogastroenterology, vol. 45, No. 20, pp. 472-476 (Mar.-Apr. 1998). (Abstract).
- Guy et al., "Substructure of the Amidotransferase Domain of Mammalian Carbamyl Phosphate Synthetase," J. Biol. Chem., vol. 270, No. 5, pp. 2190-2197 (Feb. 3, 1995).
- Haraguchi et al., "Cloning and Sequence of a cDNA Encoding Human Carbamyl Phosphate Synthetase I Molecular Analysis of Hyperammonemia," Gene, vol. 107, No. 2, pp. 335-340 (1991).
- Hoshida et al., "Carbamyl Phosphate Synthetase I Deficiency. One Base Substitution in an Exon of the CPS I Gene Causes a 9-Basepair Deletion Due to Aberrant Splicing," J. Clin. Invest. vol. 91, No. 5, pp. 1884-1887 (May 1993).
- Ikeda et al., "Cardiovascular Effects of Citrullin in Ischemia/Reperfusion Injury Via a Non-Nitric Oxide-Mediated Mechanism," Methods Find Exp. Clin. Pharmacol., vol. 22, No. 7, pp. 563-571 (Sep. 2000). (Abstract).
- International Search Report and Written Opinion corresponding to PCT International Application No. PCT/US05/06081 dated Dec. 23, 2005.
- Jang et al., "Angiogenesis Is Impaired by Hypercholesterolemia: Role of Asymmetric Dimethylarginine," Circulation, vol. 102, No. 12, pp. 1414-1419 (Sep. 19, 2000). (Abstract).
- Kurowka, "Nitric Oxide Therapies in Vascular Diseases," Current Pharmaceutical Design, vol. 8: pp. 155-166 (2002).
- Lagace et al., "Rat Carbamyl-Phosphate Synthetase I Gene," J. Biol. Chem., vol. 262, No. 22, pp. 10415-10418 (Aug. 5, 1987).
- Lundman et al. "Mild-to-moderate Hypertriglyceridemia in Young Men Is Associated with Endothelial Dysfunction and Increased Plasma Concentrations of Asymmetric Dimethylarginine," J. Am. Coll. Cardiol., vol. 38, No. 1, pp. 111-116 (Jul. 2001). (Abstract).
- MacCallum, "Detection of PCR Amplified Products" PCR Essential Data, pp. 99-127 (1995).
- Maier et al., "Activities of Urea-Cycle Enzymes in Chronic Liver Disease," Klin. Wochenschr., vol. 57, No. 13, pp. 661-665 (Jul. 3, 1979). (Abstract).
- Maxwell et al., "Endothelial Dysfunction in Hypercholesterolemia Is Reserved by a Nutritional Product Designed to Enhance Nitric Oxide Activity," Cardiovasc. Drugs Ther., vol. 14, No. 3, pp. 309-316 (Jun. 2000). (Abstract).
- Maxwell et al., "L-arginine Enhances Aerobic Exercise Capacity in Association with Augmented Nitric Oxide Production," J. Appl. Physiol., vol. 90, No. 3, pp. 933-938 (Mar. 2001). (Abstract).
- Maxwell et al., "Nutritional Therapy for Peripheral Arterial Disease: a Double-Blind, Placebo-Controlled, Randomized Trial of HeartBar," Vasc. Med., vol. 5, No. 1, pp. 11-19 (2000). (Abstract). MEDLINE Database Accession No. NLM7600831 [on-line] US National Library of Medicine (NLM), Bethesda, MD, US; Jul. 1995.
- Mitani et al., "Prolonged Administration of L-Arginine Ameliorates Chronic Pulmonary Hypertension and Pulmonary Vascular Remodeling in Rats," Circulation, vol. 96, No. 2, pp. 689-697 (1997). (Abstract).
- Mize et al., "Urea Cycle Disorders," Mol. Genet. Basis. Neurol. Dis., 2nd ed., pp. 1151-1174 (1997).
- Mizutani et al., "Oral Administration of Arginine and Citrulline in the Treatment of Lysinuric Protein Intolerance," Tohoku J. of Exp. Med., vol. 142, pp. 15-24 (1998).
- Mori et al., "Regulation of Nitric Oxide Production by Arginine Metabolic Enzymes," Biochem. Biophys. Res. Commun., vol. 275, No. 3, pp. 715-719 (Sep. 7, 2000). (Abstract).
- Morrow, "The Isoprostanes: Their Quantification as an Index of Oxidant Stress Status in Vivo," Drug Metabolism Reviews, vol. 32, Nos. 3 & 4, pp. 377-385 (2000).
- Muriel, "Regulation of Nitric Oxide Synthesis in the Liver," J. Appl. Toxicol., vol. 20, No. 3, pp. 189-195 (May-Jun. 2000). (Abstract).
- Niebauer et al., "Gene Transfer of Nitric Oxide Synthase: Effects on Endothelial Biology," J. Am. Coll. Cardiol., vol. 34, No. 4, pp. 1201-1207 (Oct. 26, 1999). (Abstract).
- Niebauer et al., "Local L-arginine Delivery after Balloon Angioplasty Reduces Monocyte Binding and Induces Apoptosis," Circulation, vol. 100, No. 17, pp. 1830-1835 (Oct. 26, 1999). (Abstract).
- Notification Concerning Transmittal of International Preliminary Report on Patentability corresponding to PCT international application No. PCT/US05/06081 dated Sep. 8, 2006.
- Nyunoya et al., "Characterization and Derivation of the Gene Coding for Mitochondrial Carbamyl Phosphate Synthetase I of Rat," J. Biol. Chem., vol. 260, No. 16, pp. 9346-9356 (Aug. 5, 1985).
- O'Connor et al., "Nonalcoholic Fatty Liver (NASH Syndrome)," Gastroenterologist, vol. 5, No. 4, pp. 316-329 (Dec. 1997). (Abstract).
- Official Action corresponding to Chinese Patent Application No. 2005800112693.5 dated Aug. 22, 2008.
- Official Action corresponding to U.S. Appl. No. 10/785,374 dated Jul. 25, 2008.
- Official Action corresponding to U.S. Appl. No. 10/785,374 dated Sep. 14, 2007.
- Pearson et al., "Neonatal Pulmonary Hypertension," N. Engl. J. Med., vol. 344, No. 24, pp. 1832-1838 (Jun. 12, 2001).
- Pernow et al., "L-Arginine Protects from Ischemia-Reperfusion-Induced Endothelial Dysfunction in Humans in Vivo," J. Appl. Physiol., vol. 95, pp. 2218-2222 (2003). (Abstract).
- Saijyo et al., "Autonomic Nervous System Activity During Infusion of L-Arginine in Patients with Liver Cirrhosis," Liver, vol. 18, No. 1, pp. 27-31 (1998). (Abstract).
- Schwarzacher et al., "Local Intramural Delivery of L-arginine Enhances Nitric Oxide Generation and Inhibits Lesion Formation after Balloon Angioplasty," Circulation, vol. 95, No. 7, pp. 1863-1869 (Apr. 1, 1997). (Abstract).
- Schwarzacher et al., "Effects of Stenting on Adjacent Vascular Distensibility and Neointima Formation: Role of Nitric Oxide," Vasc. Med., vol. 6, No. 3, pp. 139-144 (2001). (Abstract).
- Stuhlinger et al., "Homocysteine Impairs the Nitric Oxide Synthase Pathway: Role of Asymmetric Dimethylarginine," Circulation, vol. 104, No. 21, pp. 2569-2575 (Nov. 20, 2001). (Abstract).
- Suzuki et al., "Effect of Local Delivery of L-Arginine on In-stent Restenosis in Humans," The American Journal of Cardiology, vol. 89, pp. 363-367 (Feb. 15, 2002).
- Stuhlinger et al., "Relationship between Insulin Resistance and an Endogenous Nitric Oxide Synthase Inhibitor," JAMA, vol. 287, No. 11, pp. 1420-1426 (Mar. 20, 2002). (Abstract).
- Treem et al., "Disorders of the Mitochondria," Semin. Liver Dis., vol. 18, No. 3, pp. 237-253 (1998). (Abstract).
- Uemura et al., "Rapid and Efficient Vascular Transport of Arginine Polymers Inhibits Myointimal Hyperplasia," vol. 102, No. 21, pp. 2629-2635 (Nov. 21, 2000). (Abstract).
- Vassal et al., "Busulfan Disposition and Hepatic Venous Occlusive Disease in Children Undergoing Bone Marrow Transplantation," Cancer Chemo. Phar., vol. 37, pp. 247-253 (1996).
- Vnencak-Jones et al., "Efficacy of Prenatal DNA Testing for Carbamyl Phosphate Synthetase I Deficiency," Am. Jour. Human Genetics, vol. 51, No. 4 Suppl., p. A357 (1992).
- Vosatka, "Persistent Pulmonary Hypertension of the Newborn," N. Engl. J. Med., vol. 346, No. 11, p. 864 (Mar. 14, 2002).
- Wagh et al., "Oral Citrulline as Arginine Precursor May Be Beneficial in Sickle Cell Disease: Early Phase Two Results," J. Natl. Med. Assoc., vol. 93, No. 10, pp. 363-371 (Oct. 2001). (Abstract).
- Interview Summary corresponding to U.S. Appl. No. 10/785,374 dated Jun. 1, 2009.
- Interview Summary corresponding to U.S. Appl. No. 09/585,077 dated Nov. 4, 2004.
- Notice of Allowance corresponding to U.S. Appl. No. 09/323,472 (Patent No. 6,346,382) dated Aug. 9, 2001.
- Notice of Allowance corresponding to U.S. Appl. No. 09/585,077 dated Oct. 8, 2003.
- Notification of Transmittal of International Search Report or the Declaration corresponding to International Application No. PCT/US00/15079 dated Aug. 22, 2000.
- Office Communication corresponding to U.S. Appl. No. 09/989,956 dated Oct. 18, 2006.

- Office Communication corresponding to U.S. Appl. No. 09/989,956 dated Jan. 30, 2006.
- Office Communication corresponding to U.S. Appl. No. 09/989,956 dated Jul. 1, 2004.
- Office Communication corresponding to U.S. Appl. No. 09/323,472 (Patent No. 6,346,382) dated Dec. 20, 2000.
- Office Communication corresponding to U.S. Appl. No. 09/585,077 dated Jul. 3, 2002.
- Olsvik, O. et al., "A nested PCR followed by magnetic separation of amplified fragments for detection of *Escherichia coli* Shigalike toxin genes," *Molecular and Cellular Probes*, vol. 5, pp. 429-435 (Dec. 1991).
- Saiki, R.K. "Amplification of Genomic DNA," Chapter 2 in *PCR Protocols: A Guide to Methods and Applications*, Innis, M.A. et al, eds., Academic Press, Inc., San Diego, pp. 13-21 (1990).
- Office Communication corresponding to Chinese Patent Application No. 20050012693.5 dated Oct. 30, 2009.
- Ahrens et al., "Consensus statement from a Conference for the Management of Patients With Urea Cycle Disorders," Supplement to the *Journal of Pediatrics*, vol. 138, No. 1, pp. S1-S5 (Jan. 2001).
- Barsotti, "Measurement of ammonia in blood," *The Journal of Pediatrics*, vol. 138, pp. S11-S20 (Jan. 2001).
- Batshaw et al., "Alternative pathway therapy for urea cycle disorders: Twenty years later," *J. Pediatr.*, vol. 138, pp. S46-S55 (2001).
- Leonard, J., "The nutritional management of urea cycle disorders," *The Journal of Pediatrics*, vol. 138, No. 1, pp. S40-S45 (2001).
- Singh et al., "Nutritional Management of Urea Cycle Disorders," *Crit Care Clin*, vol. 21, pp. S27-S35, (2005).
- Smith et al., "Urea Cycle Disorders: Clinical Presentation Outside the Newborn Period," *Crit Care Clin*, vol. 21, pp. S9-S17 (2005).
- Summar and Tuchman, "Proceedings of a Consensus Conference for the Management of Patients with Urea Cycle Disorders," *J. Pediatr.*, vol. 138, pp. S6-S10 (2001).
- Summar, M., "Current strategies for the management of neonatal urea cycle disorders," *J. Pediatr.*, vol. 138, pp. S30-S39 (2001).
- Office Action corresponding to European Patent Application No. 06 005 642.1-2403 dated Nov. 12, 2010.
- "Persistierende fetale Zirkulation (PFC-Syndrom)," In: Ania Carolina Muntau: "Intensivkurs Padiatrie", Elsevier Urban & Fischer, Munchen, pp. 17-18 (2009).
- Abman et al., "Role of endothelium-derived relaxing factor during transition of the pulmonary circulation at birth," *Am. J. Physiol.*, vol. 259, pp. H1921-H1927 (1990).
- Abman, S. H.; *Arch Dis Child Fetal Neonatal Ed* (2002) 87: F15-F18.
- Adelman et al., "In Vitro Deletional Mutagenesis for Bacterial Production of the 20,000-Dalton Form of Human Pituitary Growth Hormone," *DNA*, vol. 2, p. 183-193 (1983).
- Advisory Action corresponding to U.S. Appl. No. 12/364,078 dated May 23, 2011.
- Allen, J. and ATS subcommittee AoP, *Am J Respir Crit Care Med* (2003) 168: 356-396.
- Ananthkrishnan et al., "L-Citrulline ameliorates chronic hypoxia-induced pulmonary hypertension in newborn piglets," *Am. J. Physiol. Lung Cell. Mol. Physiol.*, vol. 297, pp. L506-L511 (2009).
- Artymiuk et al., "Biotin carboxylase comes into the fold," *Nature Struct. Biol.*, vol. 3, pp. 128-132 (1996).
- Aschner, J. L., "New Therapies for Pulmonary Hypertension in Neonates and Children," *Pediatr. Pulmonol. Suppl.*, vol. 26, pp. 132-135 (2004).
- Bachmann et al., *New England Journal of Medicine*, vol. 304, p. 543 (1981).
- Ballard et al., "Inhaled nitric oxide in preterm infants undergoing mechanical ventilation," *N. Engl. J. Med.*, vol. 355, No. 4, pp. 343-353 (2006).
- Barr et al., "Pharmacokinetics and safety of intravenously administered citrulline in children undergoing congenital heart surgery: Potential therapy for postoperative pulmonary hypertension," *The Journal of Thoracic and Cardiovascular Surgery*, vol. 134, No. 2, pp. 319-326 (Aug. 2007).
- Batista da Costa, Jr. et al., "Surgical treatment of intracranial aneurysms: six-year experience in Belo Horizonte, MG, Brazil," *Arq Neuro-Psiquiatr* (Sao Paulo), vol. 62, pp. 245-249 (2004).
- Batshaw and Brusilow, "Valproate-induced hyperammonemia," *Annals of Neurology*, vol. 11, No. 3, pp. 319-321 (1982).
- Bearman, S.I., "Venooclusive disease of the liver: Development of a model for predicting fatal outcome after marrow transplantation," *Journal of Clinical Oncology*, vol. 11, pp. 1729-1736 (1993).
- Beaucage et al., "Deoxynucleoside phosphoramidites—a new class of key intermediates for deoxypolynucleotide synthesis," *Tetrahedron Letters*, vol. 22, pp. 1859-1862 (1981).
- Beaumier, L., "Arginine: New and exciting developments for an 'old' amino acid," *Biomedical & Environmental Sciences*, vol. 9, pp. 296-315 (1996).
- Becker et al., *Archives of Biochemistry & Biophysics*, vol. 223, pp. 381-392 (1983).
- Berkenbosch et al., "Decreased synthesis and vasodilation to nitric oxide in piglets with hypoxia-induced pulmonary hypertension," *Am. J. Physiol. Lung Cell. Mol. Physiol.*, vol. 278, pp. L276-L283 (2000).
- Bernard et al., *Intensive Care Medicine*, vol. 20, pp. 225-232 (1994).
- Blum et al., "Oral L-arginine in Patients with Coronary Artery Disease on Medical Management," *Circulation*, vol. 101, pp. 2160-2164 (May 9, 2000).
- Boger, R. H., *Curr Opin Clin Nutr and Met Care* (2008) 11:55-61.
- Bourrier et al., *Prese Medicale*, vol. 17, pp. 2063-2066 (1988).
- Caroline et al., "Acute L-arginine supplement and cardiac surgery," *Canadian J. Anesth.*, vol. 43, p. A16 (1996).
- Castillo et al., "Whole body arginine metabolism and nitric oxide synthesis in newborns with persistent pulmonary hypertension," *Pediatr. Res.*, vol. 38, pp. 17-24 (1995).
- Castro-Gago et al., *Child Neuro Systems*, vol. 6, pp. 434-436 (1990).
- Cervera et al., "Photoaffinity labeling with UMP of lysine 992 of carbamyl phosphate synthetase from *Escherichia coli* allows identification of the binding site for the pyrimidine inhibitor," *Biochemistry*, vol. 35, pp. 7247-7255 (1996).
- Cheung et al., "Channeling of Urea Cycle Intermediates in Situ in Permeabilized Hepatocytes," *J. Biol. Chem.*, vol. 264, pp. 4038-4044 (1989).
- Chin-Dusting et al., "Dietary supplementation with L-arginine fails to restore endothelial function in forearm resistance arteries of patients with severe heart failure," *J. Am. Coll. Cardiol.*, vol. 27, pp. 1207-1213 (1996).
- Cohen P. P., *Current Topics in Cellular Regulation*, vol. 18, pp. 1-19 (1981).
- Conner et al., *Proc. Natl. Acad. Sci. U.S.A.* 80:278(1983).
- Coude et al., "Inhibition of ureagenesis by valproate in rat hepatocytes. Role of N-acetylglutamate and acetyl-CoA," *Biochem. J.*, vol. 216, pp. 233-236 (1983).
- Coude et al., *J. Clin. Invest.*, vol. 64, pp. 1544-1551 (1979).
- Coulter et al., *Lancet*, vol. 1, No. 8181, pp. 1310-1311 (1980).
- Crea et al., *Proc. Natl. Acad. Sci. USA*, vol. 75, pp. 5765-5769 (1978).
- de Groot et al., *Biochemical & Biophysical Research Communications*, vol. 124, pp. 882-888 (1984).
- Dimopoulou et al., "High Incidence of Neuroendocrine Dysfunction in Long-Term Survivors of Aneurysmal Subarachnoid Hemorrhage," *Stroke*, vol. 35, pp. 2884-2889 (2004).
- Eadie et al., "Valproate-associated hepatotoxicity and its biochemical mechanisms," *Med. Toxicol.*, vol. 3, pp. 85-106 (1998).
- Eichenlaub et al., "Mutants of the mini-F plasmid pML31 thermosensitive in replication," *J. Bacteriol.*, vol. 138, pp. 559-566 (1979).
- European Search Report corresponding to European Patent Application No. 10184493 dated Mar. 30, 2011.
- European Search Report corresponding to European Patent Application No. 09707788.7-2123/2247297 dated Apr. 28, 2011.
- Faber-Langendoen et al., *Bone Marrow Transplantation*, vol. 12, pp. 501-507 (1993).
- Fagan et al., "L-arginine reduces 1-10 right heart hypertrophy in hypoxia-induced pulmonary hypertension," *Biochemical and Biophysical Research Communications*, vol. 254, No. 1, pp. 100-103, (Jan. 8, 1999).
- Feng et al., "Effects of L-arginine on endothelial and cardiac function in rats with heart failure," *Eur. J. Pharmacol.*, vol. 376, pp. 37-44 (Jul. 2, 1999).

- Fike, C. D. et al., *J Appl Physiol* (2000) 88:1797-1803.
- Fike, C. D., et al., *American Journal of Physiology (Lung, Cellular and Molecular Physiology 18)* (1998) 274:L517-L526.
- Flett et al., "Aneurysmal Subarachnoid Hemorrhage: Management Strategies and Clinical Outcomes in a Regional Neuroscience Center," *AJNR Am. J. Neuroradiol.*, vol. 26, pp. 367-372 (2005).
- Genbank Accession No. AB005063, 1998.
- Genbank Accession No. AH005315, 1998.
- Genbank Accession No. BAA21088, 1998.
- Genbank Accession No. M11710, 1985.
- Genbank Accession No. M12318, 1985.
- Genbank Accession No. M12319, 1985.
- Genbank Accession No. M12320, 1985.
- Genbank Accession No. M12321, 1985.
- Genbank Accession No. M12322, 1985.
- Genbank Accession No. M12323, 1985.
- Genbank Accession No. M12324, 1985.
- Genbank Accession No. M12325, 1985.
- Genbank Accession No. M12326, 1985.
- Genbank Accession No. M12327, 1985.
- Genbank Accession No. M12328, 1985.
- Genbank Accession No. M12335, 1985.
- Genbank Accession No. M27174, 1987.
- Genbank Accession No. P03965, 1986.
- Genbank Accession No. P07258, 1988.
- Genbank Accession No. X67573, 1992.
- Gribskov et al., "Sigma factors from *E. coli*, *B. subtilis*, phage SP01, and phage T4 are homologous proteins," *Nucl. Acids. Res.*, vol. 14, pp. 6745-6763 (1986).
- Hauser et al., "Allopurinol-induced orotidinuria," *New England Journal of Medicine*, vol. 322, pp. 1641-1645 (1990).
- Hebert, P.C., "A simple multiple system organ failure scoring system predicts mortality of patients who have sepsis syndrome," *Chest*, vol. 104, pp. 230-235 (1993).
- Hladunewich et al., "Effect of L-arginine therapy on the glomerular injury of preeclampsia: a randomized controlled trial," *Obstet. Gynecol.*, vol. 107, pp. 886-895 (2006).
- Horiuchi et al., "Sex-Related Differences in Patients Treated Surgically for Aneurysmal Subarachnoid Hemorrhage," *Neurol. Med. Chir. (Tokyo)*, vol. 46, pp. 328-332 (2006).
- Jackson M. J., *Annual Review of Genetics*, vol. 20, pp. 431-464 (1986).
- Javid-Majd et al., "Comparison of the Functional Differences for the Homologous Residues within the Carboxy Phosphate and Carbamate Domains of Carbamoyl Phosphate Synthetase," *Biochemistry*, vol. 35, pp. 14362-14369 (1996).
- Jobe et al., "Bronchopulmonary Dysplasia," *Am. J. Respir. Crit. Care Med.*, vol. 163, pp. 1723-1729 (2001).
- Jones et al., "Venoocclusive disease of the liver following bone marrow transplantation," *Transplantation*, vol. 44, pp. 778-783 (1987).
- Kamoun et al., "Valproate-induced inhibition of urea synthesis," *Lancet*, vol. 1, p. 48 (1987).
- Kinouchi et al., "Prevention of Symptomatic Vasospasm After Aneurysmal Subarachnoid Hemorrhage by Intraoperative Cisternal Fibrinolysis Using Tissue-Type Plasminogen Activator Combined with Continuous Cisternal Drainage," *Neurol. Med. Chir. (Toykyo)*, vol. 44, pp. 569-577 (2004).
- Kinsella et al., "Low-dose inhalational nitric oxide in persistent pulmonary hypertension of the newborn," *Lancet*, vol. 340, pp. 819-820 (1992).
- Kinsella et al., "Bronchopulmonary dysplasia," *Lancet*, vol. 367, No. 9520, pp. 1421-1431 (Apr. 29, 2006).
- Kivisaari et al., "MR Imaging After Aneurysmal Subarachnoid Hemorrhage and Surgery: A Long-term Follow-up Study," *AJNR Am. J. Neuroradiol.*, vol. 22, pp. 1143-1148 (Jun./Jul. 2001).
- Kyte & Doolittle, "A simple method for displaying the hydrophobic character of a protein," *J. Mol. Biol.*, vol. 157, pp. 105-132 (1982).
- Landgren et al., *Science*, 241:1007, (1988).
- Landgren et al., *Science*, 242:229-237, (1988).
- Laursen et al., "Hypoxia-induced pulmonary vascular remodeling and right ventricular hypertrophy is unaltered by long-term oral L-arginine administration," *Vascul. Pharmacol.*, vol. 49, pp. 71-76 (2008).
- Lipsitz et al., "Endogenous nitric oxide and pulmonary vascular tone in the neonate," *J. Pediatr. Surg.*, vol. 31, pp. 137-140 (1996).
- Liu and Sommer, "Restriction endonuclease fingerprinting (REF): a sensitive method for screening mutations in long, contiguous segments of DNA," *Biotechniques*, vol. 18, No. 3, pp. 470-477 (1995).
- Liu, et al., *Am J Physiol Lung Cell Mol Physiol* (2006) 290:L2-L10.
- Lorente et al., "Modulation of systemic hemodynamics by exogenous L-arginine in normal and bacteremic sheep," *Crit. Care Med.*, vol. 27, pp. 2474-2479 (1999).
- Maniatis et. al., in *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y., pp. 280-281 (1982).
- Marrini et al., "Hepatic and renal contributions to valproic acid-induced hyperammonemia," *Neurology*, vol. 38, pp. 365-371(1988).
- Marshall et al., "Multiple organ dysfunction score: a reliable descriptor of a complex clinical outcome," *Critical Care Medicine*, vol. 23, pp. 1638-1652 (1995).
- Matuschak and Rinaldo, "Organ interactions in the adult respiratory distress syndrome during sepsis. Role of the liver in host defense," *Chest*, vol. 94, pp. 400-406 (1988).
- Matuschak et al., *American Review of Respiratory Disease*, vol. 141, pp. 1296-1306 (1990).
- Matuschak, G. M., "Lung-liver interactions in sepsis and multiple organ failure syndrome," *Clinics in Chest Medicine*, vol. 17, pp. 83-98 (1996).
- McCaffrey et al., "Effect of L-Arginine Infusion on Infants with Persistent Pulmonary Hypertension of the Newborn," *Biol. Neonate*, vol. 67, No. 4, pp. 240-243 (1995).
- McDonald et al., "Veno-occlusive disease of the liver and multiorgan failure after bone marrow transplantation: a cohort study of 355 patients," *Annals of Internal Medicine*, vol. 118, pp. 255-267 (1993).
- Meier, U., *Pharm Stat* (2006) 5:253-263.
- Meister, A., "Mechanism and regulation of the glutamine-dependent carbamyl phosphate synthetase of *Escherichia coli*," *Adv. Enzymol. Relat. Areas Mol. Biol.*, vol. 62, pp. 315-374 (1989).
- Messing et al., *Third Cleveland Symposium on Macro Molecular and Recombinant DNA*, Ed. Walton, A., (Elsevier, Amsterdam) (1981).
- Miller et al., "Effects of an acute dose of L-arginine during coronary angiography in patients with chronic renal failure: a randomized, parallel, double-blind clinical trial," *Am. J. Nephrol.*, vol. 23, pp. 91-95 (2003).
- Mitchell et al., "Syndrome of Idiopathic Hyperammonemia after High-Dose Chemotherapy: Review of Nine Cases," *American Journal of Medicine*, vol. 85, No. 5, pp. 662-667 (1988).
- Moncada et al., "The L-Arginine-Nitric Oxide Pathway," *New England Journal of Medicine*, vol. 329, pp. 2002-2012 (1993).
- Moorman et al., "Expression patterns of mRNAs for ammonia-metabolizing enzymes in the developing rat: the ontogenesis of hepatocyte heterogeneity," *Histochemical Journal*, vol. 22, pp. 457-468 (1990).
- Morris et al., "Hydroxyurea and arginine therapy: impact on nitric oxide production in sickle cell disease," *J. Pediatr. Hematol. Oncol.*, vol. 25, pp. 629-634 (2003).
- Mupanemunda, R. H., *Early Human Development* (1997) 47: 247-262.
- Needleman et al., "A general method applicable to the search for similarities in the amino acid sequence of two proteins," *J. Mol. Biol.*, vol. 48, pp. 443-453 (1970).
- Nelin et al., "L-arginine, but not D-arginine, increases nitric oxide production and vasodilates hypoxic neonatal pig lungs," *FASEB Journal*, vol. 18, No. 4-5, p. A327 (2004).
- Notification of Transmittal of the International Search Report and the Written Opinion of the International Searching Authority, or the Declaration corresponding to International Application No. PCT/US 09/32824 dated Apr. 21, 2009.
- Notification of Transmittal of the International Search Report and the Written Opinion of the International Searching Authority, or the Declaration corresponding to International Application No. PCT/US 09/32826 dated Mar. 25, 2009.
- Nuzum, "Urea Cycle Enzyme Adaptation to Dietary Protein in Primates," *Science*, vol. 172, pp. 1042-1043 (1971).
- Office Action corresponding to European Patent Application Serial No. 06 005 642.1-2403 dated Jul. 25, 2011.

- Office Action corresponding to EP Patent Application Serial No. 05723789 dated Mar. 31, 2011.
- Office Action corresponding to Israeli Patent Application No. 177224 dated Apr. 26, 2011.—Translation.
- Office Action corresponding to Mexican Patent Application No. PA/a/2006/009468 dated May 12, 2011.
- Office Action corresponding to U.S. Appl. No. 12/322,434 dated Apr. 5, 2011.
- Office Action corresponding to U.S. Appl. No. 12/364,078 dated Oct. 21, 2010.
- Office Action corresponding to U.S. Appl. No. 12/364,078 dated Mar. 4, 2010.
- Office Action corresponding to U.S. Appl. No. 10/785,374 dated Mar. 2, 2010.
- Office Action corresponding to U.S. Appl. No. 10/785,374 dated Sep. 27, 2011.
- Palmer et al., "L-Arginine is the Physiological Precursor for the Formation of Nitric Oxide in Endothelium-Dependent Relaxation," *Biochem. Biophys. Res. Commun.*, vol. 153, pp. 1251-1256 (1988).
- Paulus et al., "Myocardial contractile effects of L-arginine in the human allograft," *J. Am. Coll. Cardiol.*, vol. 29, pp. 1332-1338 (1997).
- Pierson, D.L., "A rapid colorimetric assay for carbamyl phosphate synthetase I," *J. Biochem. Biophys. Methods*, vol. 3, pp. 31-37 (1980).
- Price et al., "Prognostic Indicators for Blood and Marrow Transplant Patients Admitted to an Intensive Care Unit," *American Journal of Respiratory & Critical Care Medicine*, vol. 158, pp. 876-884 (1998).
- Rabier et al., "Effects of organic acids on the synthesis of citrulline by intact rat liver mitochondria," *Biochimie*, vol. 68, pp. 639-647 (1986).
- Rabier et al., "Propionate and succinate effects on acetyl glutamate biosynthesis by rat liver mitochondria," *Biochem. & Biophys. Research Comm.*, vol. 91, pp. 456-460 (1979).
- Rabinstein et al., "Patterns of Cerebral Infarction in Aneurysmal Subarachnoid Hemorrhage," *Stroke*, vol. 36, pp. 992-997 (2005).
- Raiha and Suihkonen, *J. Acta Paediatrica Scand.*, vol. 57, pp. 121-127 (1968).
- Ravnik et al., "Long-term Cognitive Deficits in Patients with Good Outcomes after Aneurysmal Subarachnoid Hemorrhage from Anterior Communicating Artery," *Croat. Med. J.*, vol. 47, pp. 253-263 (2006).
- Richardson et al., "Prevention and treatment of hepatic venoocclusive disease after high-dose cytoreductive therapy," *Leukemia & Lymphoma*, vol. 31, pp. 267-277 (1998).
- Rinaldo et al., "Nitric oxide inactivates xanthine dehydrogenase and xanthine oxidase in interferon-gamma-stimulated macrophages," *American Journal of Respiratory Cell & Molecular Biology*, vol. 11, pp. 625-630 (1994).
- Roberts, J. D., et al., "Inhaled nitric oxide in persistent pulmonary hypertension of the newborn," *Lancet*, vol. 340, pp. 818-819 (1992).
- Rodriguez-Aparicio, L. B. et al., *Biochemistry*, vol. 28, pp. 3070-3074 (1989).
- Romero et al., "Therapeutic use of citrulline in cardiovascular disease," *Cardiovasc. Drug Rev.*, vol. 24, pp. 275-290 (2006).
- Rubinfeld et al., "Withdrawing life support from mechanically ventilated recipients of bone marrow transplants: a case for evidence-based guidelines," *Annals of Internal Medicine*, vol. 125, pp. 625-633 (1996).
- Rubio and Grisolia, "Human Carbamoylphosphate Synthetase I," *Enzyme*, vol. 26, pp. 233-239 (1981).
- Rubio, V., (Review) *Biochemical Society Transactions*, vol. 21, pp. 198-202 (1993).
- Ryan, R. M., "A new look at bronchopulmonary dysplasia classification," *J. Perinatology*, vol. 26, pp. 207-209 (2006).
- Saiki et al., "A Novel Method for the Detection of Polymorphic Restriction Sites by Cleavage of Oligonucleotide Probes: Application to Sickle-Cell Anemia," *Bio/Technology*, vol. 3, pp. 1008-1012 (1985).
- Sakowitz et al., "Relation of Cerebral Energy Metabolism and Extracellular Nitrite and Nitrate Concentrations in Patients after Aneurysmal Subarachnoid Hemorrhage," *Journal of Cerebral Blood Flow and Metabolism*, vol. 21, pp. 1067-1076 (2001).
- Schmid, R. D., *Clin. Chim. Acta.*, vol. 74, pp. 39-42 (1977).
- Schreiber et al., "Inhaled nitric oxide in premature infants with respiratory distress syndrome," *New England J. Med.*, vol. 349, pp. 2099-2107 (2003).
- Schulman et al., "L-arginine therapy in acute myocardial infarction: the Vascular Interaction With Age in Myocardial Infarction (Vintage MI) randomized clinical trial," *JAMA*, vol. 295, pp. 58-64 (Jan. 4, 2006).
- Sercombe et al., "Cerebrovascular Inflammation Following Subarachnoid Hemorrhage," *Jpn. J. Pharmacol.*, vol. 88, pp. 227-249 (2002).
- Shigesada et al., *Journal of Biological Chemistry*, vol. 246, pp. 5588-5595 (1971).
- Shimoda L., et al., *Physiol Res* (2000) 49:549-560.
- Shulman et al., "Veno-occlusive disease of the liver after marrow transplantation: histological correlates of clinical signs and symptoms," *Hepatology*, vol. 19, pp. 1171-1181 (1994).
- Simko et al., "L-arginine fails to protect against myocardial remodeling in L-NAME-induced hypertension," *Eur. J. Clin. Invest.*, vol. 35, pp. 362-368 (2005).
- Smith et al., "Comparisons of biosequences," *Adv. Appl. Math.*, vol. 2, pp. 482-489 (1981).
- Smith et al., "Nitric oxide precursors and congenital heart surgery: A randomized controlled trial of oral citrulline," *The Journal of Thoracic and Cardiovascular Surgery*, vol. 132, No. 1, pp. 58-65 (Jul. 2006).
- Stapleton et al., "Comparison of the Functional Differences for the Homologous Residues Within the Carboxy Phosphate and Carbamate Domains of Carbamoyl Phosphate Synthetase," *Biochemistry*, vol. 35, pp. 14352-14361 (1996).
- Stier et al., "Dietary arginine fails to protect against cerebrovascular damage in stroke-prone hypertensive rats," *Brain Res.*, vol. 549, pp. 354-356 (May 24, 1991).
- Suarez et al., "Aneurysmal Subarachnoid Hemorrhage," *N. Engl. J. Med.*, vol. 354, pp. 387-396 (2006).
- Subhedar, N. V., *Acta Paediatr suppl* (2004) 444:29-32.
- Summar ML, et al., "Physical and linkage mapping of human carbamyl phosphate synthetase I (CPS1) and reassignment from 2p to 2q35," *Cytogenetics & Cell Genetics*, vol. 71, pp. 266-267 (1995).
- Summar, M. L., "Molecular genetic research into carbamoyl-phosphate synthase I: molecular defects and linkage markers," *Journal of Inherited Metabolic Disease*, vol. 21, Suppl. 1, pp. 30-39 (1998).
- Surdacki et al., "Lack of beneficial effects of L-arginine infusion in primary pulmonary hypertension," *Wien. Klin. Wochenschr.*, vol. 106, pp. 521-526 (1994).
- Takiguchi, "Transcriptional regulation of genes for ornithine cycle enzymes," *Biochem J.*, vol. 312, pp. 649-659 (1995).
- Thorell et al., "Optical Coherence Tomography: A New Method to Assess Aneurysm Healing," *J. Neurosurg.*, vol. 102, No. 2, pp. 348-354 (2005).
- Toh et al., *European Journal of Biochemistry*, vol. 215, pp. 687-696 (1993).
- Tse et al., "Hyperammonemia following allogeneic bone marrow transplantation," *American Journal of Hematology*, No. 38, pp. 140-141 (1991).
- Turley, J. E. et al., *Am J Physiol Lung Cell Mol Physiol* (2003) 284:L489-L500.
- van den Hoff et al., "Evolutionary relationships of the carbamoylphosphate synthetase genes," *Journal of Molecular Evolution*, vol. 41, pp. 813-832 (1995).
- van der Schaaf et al., "New Detected Aneurysms on Follow-Up Screening in Patients with Previously Clipped Intracranial Aneurysms: Comparison with DSA or CTA at the Time of SAH," *Stroke*, vol. 36, pp. 1753-1758 (2005).
- Vosatka et al., "Arginine Deficiency Accompanies Persistent Pulmonary Hypertension of the Newborn," *Biol. Neonate*, vol. 66, pp. 65-70 (1994).
- Warter et al., *Revue Neurologique*, vol. 139, pp. 753-757 (1983).
- Wetmur & Davidson, *J. Mol. Biol.*, vol. 31, pp. 349-370 (1968).

Wilson et al., "L-arginine supplementation in peripheral arterial disease: no benefit and possible harm," *Circulation*, vol. 116, pp. 188-195 (Jul. 10, 2007).

Wingard et al., *Bone Marrow Transplantation*, vol. 4, pp. 685-9 (1989).

Wong et al., "Use of Phenytoin and Other Anticonvulsant Prophylaxis in Patients with Aneurysmal Subarachnoid Hemorrhage Response," *Stroke*, vol. 36, p. 2532 (2005).

Zamora et al., "Plasma L-arginine concentration, oxygenation index, and systemic blood pressure in premature infants," *Crit. Care Med.*, vol. 26, pp. 1271-1276 (1998).

Office Communication corresponding to Chinese Patent Application No. 20050012693.5 dated May 4, 2010.

Examiner's Report corresponding to Australian Patent Application No. 2009203177 dated Jun. 8, 2010.

* cited by examiner




FIG. 1




FIG. 2




FIG. 3




FIG. 4




FIG. 5




FIG. 6




FIG. 7




FIG. 8




FIG. 9

1 ctacttctca tgttcagcaa tttcttcttc tttatgtttt aaattacatg ttccataaaa ataagaaat
 71 cactgtgata cggtaattga ttttttcatt ttaaattgag/(intron exon boundary)
 111 (U4295)
 CTGTTTGCCA CGGRAGCCAC ATCAGACTGG CTCACAGCCA ACAATGTCCC TGCCACCCCA GTGGCATGGC
 181
 CGTCTCAAGA AGGACAGAAT CCCAGCCTCT CTTCCATCAG AAA/ (intron exon boundary)
 224 GTCGGAGA GAAGGTAGTC TT L(135a)
 gtaagaacta ggcatactgt tttctgaaat aatttagagg attaaatttg agaaccagta tatgaatatt
 294
 caccttgctt gattgcaagt cttttaaaac aaatttaaaa atgaaacacat ttgtggatga ttgtcaagtt
 364 (L135b)
 tcaacttcca tcaactatgga atacataaag tcatgtgtac atggtgtatg gaaacgtggt tcaaaataact
 434
 tcttagtaag gatactttcc ttgacggaaa caagtggag tatgaagaat gtaatgcagc ac

Primer	Begins	Size	SEQ ID NO:
U4295	119	20	8
L135a	220	21	9
L135b	370	24	10
Spanner 1	agctgttgccacggaagcc		6
Spanner 2	cccagcctctcttccatcagaaagtaag		7
Pairs			
U4295 - L135a	101 base fragment		
U4295 - L135b	251 base fragment		
Spanner1 - Spanner2	119 base fragment		

FIG. 10

CPSI T1405 SEQUENCE (SEQ ID NO:4)

MTRILTAFKV VRTLKTGFGF TNVTAHQKWK FSRPGIRLLS VKAQTAHIVL EDGTKMKGYS
FGHPSSVAGE VVFNGLGGY PEAITDPAYK GQILTMANPI IGNGGAPDTT ALDELGLSKY
LENGIKVSG LLVLDYSKDY NHWLATKSLG QWLQEEKVPA IYGVDRMLT KIIRDKGTML
GKIEFEGQPV DFVDPNKQNL IAEVSTKDVK VYGKGNPTKV VAVDCGIKNN VIRLLVKRGA
EVHLPWNHD FTKMEYDGIL IAGGPGNPAL AEPLIQNVRK ILES DRKEPL FGISTGNLIT
GLAAGAKTYK MSMANRGQNG PVLNITNKQA FITAQNHGYA LDNTLPAGWK PLFVNVNDQT
NEGIMHESKP FFAVQFHPEV TPGPIDTEYL FDSFFSLIKK GKATTITSVL PKPALVASRV
EVSKVLILGS GGLSIGQAGE FDYSGSQAVK AMKEENVKTV LMNPNIASVQ TNEVGLKQAD
TVYFLPITPQ FVTEVIKAEQ PDGLILGMGG QTALNCGVEL FKRGLVKEYG VKVLGTSVES
IMATEDRQLF SDKLNEINEK IAPSFAVESI EDALKAADTI GYPVMIRSAY ALGGLGSGIC
PNRETLMDLS TKAFAMTNQI LVEKSVTGWK EIEYEVV RDA DDNCVTV CNM ENVDAMGVHT
GDSVVAPAQ TLSNAEFQML RRTSINVVRH LGIVGECNIQ FALHPTSM EY CII EVNARLS
RSSALASKAT GYPLAFIAAK IALGIPLPEI KNV VSGK TSA CFEPSLDY MV TKIPRWDLDR
FHGTSSRIGS SMKSVGEVMA IGRTFEESFQ KALRMCHPSI EGFTPRLPMN KEWPSNLDLR
KELSEPSSTR IYAIKAIDD NMSLDEIEKL TYIDKWFLYK MRDILNMEKT LKGLNSESMT
EETLKRAKEI GFSDKQISKC LGLTEAQTRE LRLKKNHPW VKQIDTLAAE YPSVTNYLYV
TYNGQEHVDN FDDHGMMVLG CGPYHIGSSV EFDWCAVSSI RTLRQLGKKT VVNCNPETV
STDFDECDKL YFEELSLERI LDIYHQEACG GCIISVGGQI PNNLAVPLYK NGVKIMGTSP
LQIDRAEDRS IFSAVLDELK VAQAPWKAVN TLNEALEFAK SVDYPCLLRP SYVLSGSAMN
VVFSEDEMCK FLEEATRV SQ EHPVLT K FV EGAREVEMDA VGKDGRVISH AISEHVEDAG
VHSGDATLML PTQTISQGAI EKVKDATRKI AKAF AISGPF NVQFLVKGND VLVI ECNLRA
SRSFPFVSKT LGVDFIDVAT KVMIGENVDE KHLPTLDHPI IPADYVAIKA PMFSWPRLRD
ADPILRCEMA STGEVACFGE GIHTAFLKAM LSTGFKIPQK GILIGIQSF RPRFLGVAEQ
LHNEGFKLFA TEATSDWLNA NNPATPVAV PSQEGQNPSL SSIRKLIRDG SIDLVINLPN
NNTK FVHDNY VIRRTAVDSG IPLL TNFQVT KLFAEAVQKS RKVDSKSLFH YRQYSAGKAA
X


FIG. 11

CPSI N1405 SEQUENCE (SEQ ID NO:2)

MTRILTAFKV VRTLKTGFGF TNVTAHQKWK FSRPGIRLLS VKAQTAHIVL EDGTKMKGYS
FGHPSSVAGE VVFNTGLGGY PEAITDPAYK GQILTMANPI IGNGGAPDTT ALDELGLSKY
LENGIKVSG LLVLDYSKDY NHWLATKSLG QWLQEEKVPA IYGVDRMLT KIIRDKGTML
GKIEFEGQPV DFVDPNKQNL IAEVSTKDVK VYGKGNPTKV VAVDCGIKNN VIRLLVKRGA
EVHLVPWNHD FTKMEYDGIL IAGGPGNPAL AEPLIQNVRK ILES DRKEPL FGISTGNLIT
GLAAGAKTYK MSMANRGQNQ PVLNITNKQA FITAQNHGYA LDNTLPAGWK PLFVNVNDQT
NEGIMHESKP FFAVQFHPEV TPGPIDTEYL FDSFFSLIKK GKATTITSVL PKPALVASRV
EVSKVLILGS GGLSIGQAGE FDYSGSQAVK AMKEENVKTV LMNPNIASVQ TNEVGLKQAD
TVYFLPITPQ FVTEVIKAEQ PDGLILGMGG QTALNCGVEL FKRGV LKEYG VKVLGTSVES
IMATEDRQLF SDKLNEINEK IAPSF AVESI EDALKAADTI GYPVMIRSAY ALGGLGSGIC
PNRETLMDLS TKAFAMTNQI LVEKSVTGWK EIEYEVVRDA DDNCVTVCNM ENVDAMGVHT
GDSVVVAPAQ TLSNAEFQML RRTSINVVRH LGIVGECNIQ FALHPTSMEY CIIEVNARLS
RSSALASKAT GYPLAFIAAK IALGIPLPEI KNVVSGK TSA CFEPSLDY MV TKIPRWDLDR
FHGTSSRIGS SMKSVGEVMA IGRTFEESFQ KALRMCHPSI EGFTPRLPMN KEWPSNLDLR
KELSEPSSTR IYAIKAIDD NMSLDEIEKL TYIDKWFLYK MRDILNMEKT LKGLNSESMT
EETLKRAKEI GFSDKQISKC LGLTEAQTRE LRLKKNHPW VKQIDTLAAE YPSVTNYLYV
TYNGQEHVDN FDDHGMMVLG CGPYHIGSSV EFDWCAVSSI RTLRQLGKKT VVNCNPETV
STDFDECDKL YFEELSLERI LDIYHQEACG GCIISVGGQI PNNLAVPLYK NGVKIMGTSP
LQIDRAEDRS IFSAVLDELK VAQAPWKAVN TLNEALEFAK SVDYPCLLRP SYVLSGSAMN
VFSEDEMKK FLEEATRV SQ EHPVVLTKFV EGAREVEMDA VGKDGRVISH AISEHVEDAG
VHSGDATLML PTQTISQGAI EKVKDATRKI AKAF AISGPF NVQFLVKGND VLIECNLRA
SRSFPFVSKT LGVDFIDVAT KVMIGENVDE KHLPTLDHPI IPADYVAIKA PMFSWPRLRD
ADPILRCEMA STGEVACFGE GIHTAFLKAM LSTGFKIPQK GILIGIQSF RPRFLGVAEQ
LHNEGFKLFA TEATSDWLNA NNVPANPVAW PSQEGQNPSL SSIRKLIRDG SIDLVINLPN
NNTK FVHDNY VIRRTAVDSG IPLLTNFQVT KLFAEAVQKS RKVDSKSLFH YRQYSAGKAA
X

FIG. 12

FIGURE 13



1

THERAPEUTIC METHODS EMPLOYING
NITRIC OXIDE PRECURSORS

RELATED APPLICATION INFORMATION

This is a divisional of U.S. patent application Ser. No. 10/785,374, filed Feb. 24, 2004, which is a continuation-in-part of U.S. patent application Ser. No. 09/585,077, filed Jun. 1, 2000, now U.S. Pat. No. 6,743,823 which is a continuation-in-part of U.S. patent application Ser. No. 09/323,472, filed Jun. 1, 1999, now U.S. Pat. No. 6,346,382, the entire contents of which are herein incorporated by reference.

GRANT STATEMENT

This work was supported by NIH grants R29-DK46965, NIH HL55198, NIH ES 09915 and NIH 1 P30 CA 68485. Thus, the U.S. Government has certain rights in the invention.

TECHNICAL FIELD

The present invention relates to isolated polynucleotide molecules useful for analyzing carbamyl phosphate synthetase I phenotypes, to peptides encoded by these molecules, and to the diagnostic and therapeutic uses thereof relating to a newly identified carbamyl phosphate synthetase I polymorphism. Among such uses are methods for determining the susceptibility of a subject to hyperammonemia, decreased production of arginine and to bone marrow transplant toxicity based on an analysis of a nucleic acid sample isolated from tissue biopsies from the subject.

TABLE OF ABBREVIATIONS

ABG—arterial blood gas(es)
ALI—acute lung injury
ASO—allele-specific oligonucleotide
ATP—adenosine triphosphate
BCAA—branched chain amino acid(s)
BMT—bone marrow transplant
BSA—bovine serum albumin
BuCy—busulfan, cyclophosphamide
BUN—blood urea nitrogen
CBVP16—cyclophosphamide, bis-chloroethylnitrosourea, etoposide
cc—cubic centimeters
CPSI—carbamyl phosphate synthetase I
CTC—cyclophosphamide, thiotepa, carboplatin
CVP16TBI—cyclophosphamide, etoposide, total body irradiation
ECMO—extracorporeal membrane oxygenation
fl—full length
GSHosc—glutathione synthetase
HAT—hypoxanthine, aminopterin, thymidine
HVOD—hepatic veno-occlusive disease
iNO—inhaled nitric oxide
KDa—kilodalton
KLH—keyhole limpet hemocyanin
l—liter
LAT—ligation activated translation
LCR—ligase chain reaction
MAS—meconium aspiration syndrome
NAG—n-acetyl glutamate
NASDA™—nucleic acid sequence-based amplification
NO or NO _x —nitric oxide
NOS—nitric oxide synthetase
O/C—ornithine/citrulline

2

PBSCT—peripheral blood stem-cell transplantation
PPHN—persistent pulmonary hypertension in newborns
PCR—polymerase chain reaction
RCR—repair chain reaction
5 RDS—respiratory distress syndrome
REF—restriction endonuclease finger-printing
RT—reverse transcriptase
SSCP—single strand conformation polymorphism
SDA—strand displacement activation
10 SNP—single nucleotide polymorphism
TC—thiotepa, cyclophosphamide
TEAA—total essential amino acids
uc—urea cycle
UCF—urea cycle function
15 VPA—valproic acid

BACKGROUND ART

The in vivo synthetic pathway for arginine commences with ornithine. Ornithine is combined with carbamyl phosphate to produce citrulline, which in turn is combined with aspartate, in the presence of adenosine triphosphate (ATP), to produce argininosuccinate. In the final step, fumarate is split from argininosuccinate, to produce arginine. The degradative pathway for arginine is by the hydrolytic action of arginase, to produce ornithine and urea. These reactions form the urea cycle. The urea cycle serves as the primary pathway for removing waste nitrogen produced by the metabolism of endogenous and exogenous proteins, and is shown schematically in FIG. 1.

Disruption of metabolic processes is a frequent side effect of chemotherapy. Indeed, the agents used in high-dose chemotherapy affect a number of cellular processes. Metabolic processes localized in chemo-sensitive tissues, such as the liver and gastrointestinal tract, face a particularly great risk to disruption.

The constant turn-over and processing of nitrogen involves all the tissues in the body, but the first critical steps of the urea cycle are limited to the liver and gut. The high-dose chemotherapy associated with bone marrow transplant (BMT) interferes with liver function and is toxic to the intestine. Idiopathic hyperammonemia, which is suggestive of urea cycle dysfunction, has been reported to be associated with high mortality in patients undergoing bone marrow transplant. Davies et al., *Bone Marrow Transplantation*, 17:1119-1125 (1996); Tse et al., *American Journal of Hematology*, 38:140-141 (1991); and Mitchell et al., *American Journal of Medicine*, 85:662-667 (1988).

A common complication of BMT is hepatic veno-occlusive disease (HVOD).

HVOD is associated with jaundice, increased liver size and disruption of normal hepatic blood flow. HVOD occurs in approximately 20 to 40% of patients and is associated with severe morbidity and mortality.

Nitric oxide (NO) plays a role in regulating vascular tone and in maintaining patency of hepatic and pulmonary venules following high-dose chemotherapy. Intact urea cycle function is important not only for excretion of ammonia but in maintaining adequate tissue levels of arginine, the precursor of NO.

Carbamyl phosphate synthetase I (CPSI) is the rate limiting enzyme catalyzing the first committed step of urea genesis via the urea cycle. CPSI is highly tissue specific, with function and production substantially limited to liver and intestines. Genomically encoded, CPSI is produced in the cytoplasm and transported into the mitochondria where it is cleaved into its mature 160 kDa monomeric form. The

enzyme combines ammonia and bicarbonate to form carbamyl with the expenditure of two ATP molecules and using the co-factor N-acetyl-glutamate (NAG).

Any genetic predisposition to decreased urea cycle function would lead to hyperammonemia and would likely contribute to the severity of disorders associated with sub-optimal urea cycle function, including BMT-related toxicity. Thus, there is a need in the art for characterization of alleles present in populations suffering from disorders associated with suboptimal urea cycle function, undergoing BMT or otherwise facing exposure to environmental or pharmacological hepatotoxins. In view of the role of CPSI in the urea cycle, there is a particular need for characterization of CPSI alleles present in such populations.

SUMMARY OF THE INVENTION

A method of screening for susceptibility to sub-optimal urea cycle function in a subject is disclosed. The method comprising the steps of: (a) obtaining a nucleic acid sample from the subject; and (b) detecting a polymorphism of a carbamyl phosphate synthase I (CPSI) gene in the nucleic acid sample from the subject, the presence of the polymorphism indicating that the susceptibility of the subject to sub-optimal urea cycle function. In accordance with the present invention, detection of the polymorphism is particularly provided with respect to determining the susceptibility of a subject to bone marrow transplant toxicity.

Preferably, the polymorphism of the carbamyl phosphate synthetase polypeptide comprises a C to A transversion in exon 36 of the CPSI gene, more preferably at nucleotide 4340 of a cDNA that corresponds to the CPSI gene. More preferably, the C to A transversion at nucleotide 4340 of the cDNA that corresponds to the CPSI gene further comprises a change in the triplet code from AAC to ACC, which encodes a CPSI polypeptide having an threonine moiety at amino acid 1405.

The present invention also provides an isolated and purified biologically active CPSI polypeptide. Preferably, a polypeptide of the invention is a recombinant polypeptide. More preferably, a polypeptide of the present invention comprises human CPSI having an asparagine moiety at amino acid 1405.

The present invention also provides an isolated and purified polynucleotide that encodes a biologically active CPSI polypeptide. In a preferred embodiment, a polynucleotide of the present invention comprises a DNA molecule from a human. More preferably, a polynucleotide of the present invention comprises a cDNA that corresponds to the CPSI gene and which includes a C to A transversion at nucleotide 4340. Even more preferably, a polynucleotide of the present invention further comprises a cDNA that corresponds to the CPSI gene that includes a change in the triplet code from ACC to AAC at nucleotide 4340, and encodes a CPSI polypeptide having an asparagine moiety at amino acid 1405.

Kits and reagents, including oligonucleotides, nucleic acid probes and antibodies suitable for use in carrying out the methods of the present invention and for use in detecting the polypeptides and polynucleotides of the present invention are also disclosed herein. Methods for preparing the polynucleotides and polypeptides of the present invention are also disclosed herein.

In a further embodiment, this invention pertains to therapeutic methods based upon a polymorphism of a carbamyl phosphate synthase I (CPSI) gene as described herein. Such therapeutic methods include administration of nitric oxide precursors in the treatment and prophylaxis of disorders mediated or modulated by sub-optimal urea cycle function

(e.g. bone marrow transplant toxicity) and gene therapy approaches using an isolated and purified polynucleotide of the present invention.

It is therefore an object of the present invention to provide polynucleotide molecules that can be used in analyzing carbamyl phosphate synthetase I (CPSI) in vertebrate subjects.

It is also an object of the present invention to provide for the determination of CPSI phenotype in vertebrate subjects and particularly human subjects, based on information obtained through the analysis of nucleic acids, including genomic DNA and cDNA, derived from tissues from the subject.

It is yet another object of the present invention to provide a ready technique for determining CPSI phenotype.

It is still a further object of the present invention to provide polypeptide and polynucleotide molecules for use in generating antibodies that distinguish between the different forms of CPSI which constitute the CPSI polymorphism.

It is yet a further object of the present invention is to provide methods for diagnosing and treating clinical syndromes related to and associated with the CPSI polymorphism.

Some of the objects of the invention having been stated hereinabove, other objects will become evident as the description proceeds, when taken in connection with the accompanying drawings and examples as best described hereinbelow.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a schematic of the urea cycle;

FIG. 2 is a schematic of the consensus CPSI protein which does not reflect recognized mutations;

FIG. 3 is a schematic of the consensus CPSI protein depicting several known mutations in the protein and depicting the T1405N polymorphism of the present invention;

FIG. 4 is a schematic of recognized post-transcriptional modification of CPSI;

FIG. 5 is a schematic of the human genomic locus for CPSI;

FIG. 6 is a schematic of a cloning strategy for a full length CPSI cDNA;

FIG. 7 is a schematic of an alternative cloning strategy for a full length CPSI cDNA;

FIG. 8 is a graphical depiction of the metabolic activity of the CPSI protein expressed in COS-7 cells;

FIG. 9 is a graphical presentation of the size and position of introns in CPSI cDNA;

FIG. 10 is a diagram of exon 36 (SEQ ID NO:5) showing the locations of preferred oligonucleotide primers of the present invention;

FIG. 11 presents the amino acid sequence of T1405 CPSI (SEQ ID NO:4) (stop codon translated as "X", 165049 MW, 1.163602e+07 CN), with the initial amino acid methionine considered to be at a -1 position;

FIG. 12 presents the amino acid sequence of N1405 CPSI (SEQ ID NO:2) (stop codon translated as "X", 165062 MW, 1.161634E+07 CN), with the initial amino acid methionine considered to be at a -1 position;

FIG. 13 is a graph of a concentration curve of plasma arginine levels.

DETAILED DESCRIPTION OF THE INVENTION

Disclosed herein is the surprising discovery of a polymorphism of carbamyl phosphate synthetase I (CPSI), the enzyme that catalyzes the rate limiting first step of the urea cycle. Particularly, the polymorphism is characterized by an

5

amino acid substitution, threonine/asparagine at amino acid 1405 (heterozygosity=0.44) in CPSI.

Also disclosed herein is the surprising observation that a single nucleotide change in the CPSI gene is responsible for the polymorphism of CPSI. Particularly, a C to A transversion with exon 36 of the CPSI gene changes the triplet code from ACC to AAC and leads to the T1405N change in the encoded CPSI polypeptide.

In light of these discoveries, manipulation of nucleic acid molecules derived from the tissues of vertebrate subjects can be effected to provide for the analysis of CPSI phenotypes, for the generation of peptides encoded by such nucleic acid molecules, and for diagnostic and therapeutic methods relating to the CPSI polymorphism. Nucleic acid molecules utilized in these contexts may be amplified, as described below, and generally include RNA, genomic DNA and cDNA derived from RNA.

A. General Considerations

Most of the currently available structural information on CPSI is derived from studies of the rat CPSI enzyme. The rat CPSI enzyme and the human CPSI enzyme each comprise a single polypeptide of 1,500 residues and exhibit about 95% sequence identity. Rat CPSI polypeptide and nucleic acid sequence information is disclosed by Nyunoya, H., et al., *Journal of Biological Chemistry* 260:9346-9356 (1985) and at GenBank accession numbers AH005315, M12335, M12328, M12327, M12326, M12325, M12324, M12323, M12322, M12321, M12320, M12319, M12318 and M11710, herein incorporated by reference. The structural information about rat CPSI is derived from sequence homology and substrate and co-factor binding studies; however, no crystallographic data is available.

Mature CPSI is modular in nature, containing 2 main regions. The first region, residues 39-406, is homologous to the small subunit of the heterodimeric CPS of *Escherichia coli*. Bacterial and yeast CPSI polypeptide and nucleic acid sequence information is disclosed at GenBank accession numbers AB005063, X67573, M27174, P07258, P03965, BAA21088, SYBYCP, SYBYCS, and SYECCS, herein incorporated by reference.


The other region, residues 417-1500 (referred to hereinafter as the "CPS domain"), is homologous to the large subunit of *E. coli* CPS. Meister, A., *Adv. Enzymol. Relat. Areas Mol. Biol.* 62:315-374 (1989). This subunit is responsible for carbamyl phosphate synthesis from ammonia and for the binding of the substrates and cofactors. Meister, A., *Adv. Enzymol. Relat. Areas Mol. Biol.* 62:315-374 (1989). The CPS domain arose by gene duplication and tandem fusion in the pro-genome, and, as depicted schematically in FIG. 2, is itself composed of two phosphorylation domains and a C-terminal regulatory domain involved in the binding of n-acetylglutamate (NAG). Nyunoya, H., et al., *Journal of Biological Chemistry* 260:9346-9356 (1985).

As depicted schematically in FIG. 2, residues 407-416 act as a bridge between the two major subunits, and residues 1-38 constitute the leader peptide that directs immature CPSI to the mitochondria prior to being removed. Continuing with FIG. 2, the small subunit-like region is composed of two approximately equal subdomains. The interaction subdomain, residues 39-212, corresponds to the region which, in the small subunit of the CPS from *E. coli*, is necessary for association with the large subunit. The glutaminase subdomain, residues 213-406, is homologous to several glutamine amidotransferases and to the region of CPSI that when generated free from other components exhibited considerable glutaminase activity, as described by Guillou, F., et al. *Proc Natl Acad Sci* 86:8304-8308 (1989); Nyunoya, H., et al., *Journal of Biologi-*

6

cal Chemistry 260:9346-9356 (1985); and Guy, H. I. et al., *Journal of Biological Chemistry* 270:2190-2197 (1995). Since CPSI has lost the cysteine residue necessary to split glutamine, the function of the glutaminase subdomain is uncertain in this enzyme.

The CPS domain (corresponding to the large subunit in *E. coli*) is believed to catalyze the synthesis of carbamyl phosphate from ammonia, according to the reaction:



As shown schematically in FIGS. 1 and 2, this reaction comprises three steps: bicarbonate phosphorylation by an ATP molecule that is designated herein as ATP_A, giving carboxyphosphate; carbamate synthesis from carboxyphosphate and ammonia; and carbamate phosphorylation by another ATP molecule (ATP_B), giving carbamyl phosphate, as described by Rubio, V. and Grisolia, S., *Enzyme* 26:233-239 (1981).

As shown schematically in FIG. 4, the CPS domain appears to have arisen by duplication and tandem fusion of the duplicated component; therefore, its amino and COOH-terminal halves are homologous, as described by Nyunoya, H., et al., *Journal of Biological Chemistry* 260:9346-9356 (1985)). Each homologous half comprises an amino- and a COOH-terminal domain of about 40 and 20 kDa, respectively, of which the domain of 40 kDa of the amino-half is believed to be involved in bicarbonate phosphorylation (bicarbonate phosphorylation domain, residues 417-788) (FIG. 2). The corresponding domain in the COOH-half is involved in carbamate phosphorylation via the carbamate phosphorylation domain, residues 969-1329 (FIG. 2), as described by Alonso, E. and Rubio, V., *European Journal of Biochemistry* 229:377-384 (1995)).

These phosphorylation domains are homologous to biotin carboxylase (Toh, H. et al., *European Journal of Biochemistry* 215:687-696 (1993)), an enzyme of known tri-dimensional structure that phosphorylates bicarbonate as well as DD-ligase and glutathione synthetase (GSHase), two enzymes that catalyze analogous reactions (Artymiuk, P. J. et al., *Nature Struct. Biol.* 3:128-132 (1996)). Thus, information on these enzymes is helpful in interpreting the mutations found in homologous domains in the patients with CPSI deficiency.

Referring again to FIG. 2, of the 20-kDa domains of the large subunit-like region, the function of the domain of the amino-terminal half, residues 789-968, remains to be established. In contrast, the corresponding COOH-terminal domain, residues 1330-1500, is called the allosteric domain, because the activator, n-acetylglutamate (NAG) of CPSI and the nucleotide effectors of the *E. coli* enzyme, UMP and IMP, bind in this domain, as described by Rodriguez-Aparicio, L. B. et al., *Biochemistry* 28:3070-3074 (1989) and Cervera, J. et al., *Biochemistry* 35:7247-7255 (1996).

A.1. Enzyme Processing.

Human CPSI mRNA encodes a 165 kDa, 1500 amino acid pre-protein. The amino terminus of this precursor contains 38 residues, including 8 basic residues, and 1 acidic residue with a Pro-Gly sequence 4 residues before the start of the mature enzyme (Nyunoya, H. et al., *Journal of Biological Chemistry* 260:9346-9356 (1985); Lagace, M. et al., *Journal of Biological Chemistry* 262:10415-10418 (1987). This highly con-

served signal sequence promotes enzyme entry into the mitochondrial matrix, where it is then removed to produce the 160 kDA mature enzyme.

A.2. Normal Expression of CPSI.

CPSI enzymatic activity is first detected in human fetal liver by 5-10 weeks gestation (Moorman, A. F. et al. *Histochemical Journal* 22:457-468 (1990)). By 20 weeks gestation, the level of CPSI reaches approximately 50% of the normal adult level, where it remains until birth, after which it gradually increases to adult levels by 20 years of age (Raiha, N. C. R. and Suihkonen, *J. Acta Paediatrica Scand* 57:121-127 (1968)). Tissue expression of CPSI is essentially limited to the liver, with trace amounts of activity in the intestine and kidney. When the liver develops its mature acinar structure in adulthood, CPSI is compartmentalized in parenchymal cells around the terminal portal venules (Moorman, A. F. et al. *Histochemical Journal* 22:457-468 (1990)).

In addition to its compartmentalization, several factors are known to be important in the regulation of CPSI activity and expression. For example, low or absent levels of ornithine decrease CPSI activity, presumably due to an inhibitory effect from accumulated carbamyl phosphate (CP) as described by Jackson, M. J. et al., *Annual Review of Genetics* 20:431-464 (1986); and Rubio, V., *Biochemical Society Transactions* 21:198-202 (1993)). Levels of both CPSI mRNA and enzyme increase with a high protein diet, and in response to glucagon and glucocorticoids (Jackson, M. J. et al., *Annual Review of Genetics* 20:431-464 (1986); de Groot, C. J., et al., *Biochemical & Biophysical Research Communications* 124:882-888 (1984)). In normal unstimulated hepatic tissue that has been examined, an abundance of CPSI mRNA has been observed.

B. Screening Techniques

In accordance with the present invention, a method of screening for susceptibility to sub-optimal urea cycle function resulting in decreased ammonia clearance and decreased arginine production in a subject is provided. The method comprises: (a) obtaining a nucleic acid sample from the subject; and (b) detecting a polymorphism of a carbamyl phosphate synthase I (CPSI) gene in the nucleic acid sample from the subject, the presence of the polymorphism indicating that the susceptibility of the subject to sub-optimal urea cycle function resulting in decreased ammonia clearance and decreased arginine production. In accordance with the present invention, detection of the polymorphism is particularly provided with respect to determining the susceptibility of a subject to bone marrow transplant toxicity.

It is further noted that the polymorphism of the present invention may be used to predict toxicity in a number of conditions beyond BMT or valproic acid administration as disclosed herein and in the Examples. The polymorphism is also implicated in the mediation or modulation of disrupted ammonia clearance and arginine production in situations such as adult hepatic cirrhosis, other medication toxicities, newborns with impaired hepatic function, and the like.

As used herein and in the claims, the term “polymorphism” refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A polymorphic marker is the locus at which divergence occurs. Preferred markers have at least two alleles, each occurring at frequency of greater than 1%. A polymorphic locus may be as small as one base pair.

Useful nucleic acid molecules according to the present invention include those which will specifically hybridize to CPSI sequences in the region of the C to A transversion at base 4340 and within exon 36 changing the triplet code from ACC to AAC. This transversion leads to the T1405N change in the encoded CPSI polypeptide. Typically these are at least

about 20 nucleotides in length and have the nucleotide sequence corresponding to the region of the C to A transversion at base 4340 of the consensus CPSI cDNA sequence (EC6.3.4.16), which changes the triplet code from ACC to AAC. The term “consensus sequence”, as used herein, is meant to refer to a nucleic acid or protein sequence for CSPI, the nucleic or amino acids of which are known to occur with high frequency in a population of individuals who carry the gene which codes for a normally functioning protein, or which nucleic acid itself has normal function.

Provided nucleic acid molecules can be labeled according to any technique known in the art, such as with radiolabels, fluorescent labels, enzymatic labels, sequence tags, etc. According to another aspect of the invention, the nucleic acid molecules contain the C to A transversion at base 4340. Such molecules can be used as allele-specific oligonucleotide probes to track a particular mutation, for example, through a family of subjects.

Body samples can be tested to determine whether the CPSI gene contains the C to A transversion at base 4340. Suitable body samples for testing include those comprising DNA, RNA or protein obtained from biopsies, including liver and intestinal tissue biopsies; or from blood, prenatal; or embryonic tissues, for example.

In one embodiment of the invention a pair of isolated oligonucleotide primers are provided: 5'-AGCTGTTTGC-CACGGAAGCC-3'(SEQ ID NO:6) and 5'-CCCAGC-CTCTCTTCCATCAGAAAGTAAG-3'(SEQ ID NO:7). These primers are derived from CPSI exon 36 (the location of the polymorphism of the present invention) and related intronic sequences (SEQ ID NO:5) and produce a 119 base pair fragment. Other primers derived from CPSI exon 36 (the location of the polymorphism of the present invention) and related intronic sequences (SEQ ID NO:5) are provided in SEQ ID NOs:8-10, in FIG. 10, and in Example 2 (SEQ ID NOs:15 and 16).

The oligonucleotide primers are useful in diagnosis of a subject at risk for hyperammonemia such as can result as a BMT complication or toxicity. The primers direct amplification of a target polynucleotide prior to sequencing. These unique CPSI exon 36 oligonucleotide primers were designed and produced based upon identification of the C to A transversion in exon 36.

In another embodiment of the invention isolated allele specific oligonucleotides are provided. Sequences substantially similar thereto are also provided in accordance with the present invention. The allele specific oligonucleotides are useful in diagnosis of a subject at risk for hyperammonemia, such as can result as a BMT complication or toxicity. These unique CPSI exon 36 oligonucleotide primers were designed and produced based upon identification of the C to A transversion in exon 36.

The terms “substantially complementary to” or “substantially the sequence of” refer to sequences which hybridize to the sequences provided (e.g. SEQ ID NOs: 5-10) under stringent conditions and/or sequences having sufficient homology with any of SEQ ID NOs: 5-10, such that the allele specific oligonucleotides of the invention hybridize to the sequence. The term “isolated” as used herein includes oligonucleotides substantially free of other nucleic acids, proteins, lipids, carbohydrates or other materials with which they may be associated, such association being either in cellular material or in a synthesis medium. A “target polynucleotide” or “target nucleic acid” refers to the nucleic acid sequence of interest e.g., a CPSI-encoding polynucleotide. Other primers which

can be used for primer hybridization are readily ascertainable to those of skill in the art based upon the disclosure herein of the CPSI polymorphism.

The primers of the invention embrace oligonucleotides of sufficient length and appropriate sequence so as to provide initiation of polymerization on a significant number of nucleic acids in the polymorphic locus. The CPSI locus is depicted schematically in FIG. 5. Specifically, the term “primer” as used herein refers to a sequence comprising two or more deoxyribonucleotides or ribonucleotides, preferably more than three, and more preferably more than eight and most preferably at least about 20 nucleotides of the CPSI gene wherein the DNA sequence contains the C to A transversion at base 4340 relative to CPSI contained in SEQ ID NO’s:1 and 3. The allele including cytosine (C) at base 4340 relative to CPSI is referred to herein as the “CPSIa allele”, the “T1405 allele”, or the “threonine-encoding allele”. The allele including adenosine (A) at base 4340 relative to CPSI is referred to herein as the “CPSIb allele”, the “N1405 allele”, or the “arginine-encoding allele”.

An oligonucleotide that distinguishes between the CPSIa and the CPSIb alleles of the CPSI gene, wherein said oligonucleotide hybridizes to a portion of said CPSI gene that includes nucleotide 4340 of the cDNA that corresponds to said CPSI gene when said nucleotide 4340 is adenosine, but does not hybridize with said portion of said CPSI gene when said nucleotide 4340 is cytosine is also provided in accordance with the present invention. An oligonucleotide that distinguishes between the CPSIa and the CPSIb alleles of the CPSI gene, wherein said oligonucleotide hybridizes to a portion of said CPSI gene that includes nucleotide 4340 of the cDNA that corresponds to said CPSI gene when said nucleotide 4340 is cytosine, but does not hybridize with said portion of said CPSI gene when said nucleotide 4340 is adenosine is also provided in accordance with the present invention. Such oligonucleotides are preferably between ten and thirty bases in length. Such oligonucleotides may optionally further comprises a detectable label.

Environmental conditions conducive to synthesis include the presence of nucleoside triphosphates and an agent for polymerization, such as DNA polymerase, and a suitable temperature and pH. The primer is preferably single stranded for maximum efficiency in amplification, but may be double stranded. If double stranded, the primer is first treated to separate its strands before being used to prepare extension products. The primer must be sufficiently long to prime the synthesis of extension products in the presence of the inducing agent for polymerization. The exact length of primer will depend on many factors, including temperature, buffer, and nucleotide composition. The oligonucleotide primer typically contains 12-20 or more nucleotides, although it may contain fewer nucleotides.

Primers of the invention are designed to be “substantially” complementary to each strand of the genomic locus to be amplified. This means that the primers must be sufficiently complementary to hybridize with their respective strands under conditions which allow the agent for polymerization to perform. In other words, the primers should have sufficient complementarity with the 5' and 3' sequences flanking the transversion to hybridize therewith and permit amplification of the genomic locus.

Oligonucleotide primers of the invention are employed in the amplification method which is an enzymatic chain reaction that produces exponential quantities of polymorphic locus relative to the number of reaction steps involved. Typically, one primer is complementary to the negative (-) strand of the polymorphic locus and the other is complementary to

the positive (+) strand. Annealing the primers to denatured nucleic acid followed by extension with an enzyme, such as the large fragment of DNA polymerase I (Klenow) and nucleotides, results in newly synthesized + and - strands containing the target polymorphic locus sequence. Because these newly synthesized sequences are also templates, repeated cycles of denaturing, primer annealing, and extension results in exponential production of the region (i.e., the target polymorphic locus sequence) defined by the primers. The product of the chain reaction is a discrete nucleic acid duplex with termini corresponding to the ends of the specific primers employed.

The oligonucleotide primers of the invention may be prepared using any suitable method, such as conventional phosphotriester and phosphodiester methods or automated embodiments thereof. In one such automated embodiment, diethylphosphoramidites are used as starting materials and may be synthesized as described by Beaucage et al., *Tetrahedron Letters* 22:1859-1862 (1981). One method for synthesizing oligonucleotides on a modified solid support is described in U.S. Pat. No. 4,458,066.

Any nucleic acid specimen, in purified or non-purified form, can be utilized as the starting nucleic acid or acids, providing it contains, or is suspected of containing, a nucleic acid sequence containing the polymorphic locus. Thus, the method may amplify, for example, DNA or RNA, including messenger RNA, wherein DNA or RNA may be single stranded or double stranded. In the event that RNA is to be used as a template, enzymes, and/or conditions optimal for reverse transcribing the template to DNA would be utilized. In addition, a DNA-RNA hybrid which contains one strand of each may be utilized. A mixture of nucleic acids may also be employed, or the nucleic acids produced in a previous amplification reaction herein, using the same or different primers may be so utilized. The specific nucleic acid sequence to be amplified, i.e., the polymorphic locus, may be a fraction of a larger molecule or can be present initially as a discrete molecule, so that the specific sequence constitutes the entire nucleic acid. It is not necessary that the sequence to be amplified be present initially in a pure form; it may be a minor fraction of a complex mixture, such as contained in whole human DNA.

DNA utilized herein may be extracted from a body sample, such as blood, tissue material, preferably liver tissue, and the like by a variety of techniques such as that described by Maniatis et. al. in *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y., p 280-281 (1982). If the extracted sample is impure, it may be treated before amplification with an amount of a reagent effective to open the cells, or animal cell membranes of the sample, and to expose and/or separate the strand(s) of the nucleic acid(s). This lysing and nucleic acid denaturing step to expose and separate the strands will allow amplification to occur much more readily.

The deoxyribonucleotide triphosphates dATP, dCTP, dGTP, and dTTP are added to the synthesis mixture, either separately or together with the primers, in adequate amounts and the resulting solution is heated to about 90-100° C. from about 1 to 10 minutes, preferably from 1 to 4 minutes. After this heating period, the solution is allowed to cool, which is preferable for the primer hybridization. To the cooled mixture is added an appropriate agent for effecting the primer extension reaction (called herein “agent for polymerization”), and the reaction is allowed to occur under conditions known in the art. The agent for polymerization may also be added together with the other reagents if it is heat stable. This synthesis (or amplification) reaction may occur at room temperature up to a temperature above which the agent for polymerization no

longer functions. Thus, for example, if DNA polymerase is used as the agent, the temperature is generally no greater than about 40° C. Most conveniently the reaction occurs at room temperature.

The agent for polymerization may be any compound or system which will function to accomplish the synthesis of primer extension products, including enzymes. Suitable enzymes for this purpose include, for example, *E. coli* DNA polymerase I, Klenow fragment of *E. coli* DNA polymerase, polymerase mutants, reverse transcriptase, other enzymes, including heat-stable enzymes (i.e., those enzymes which perform primer extension after being subjected to temperatures sufficiently elevated to cause denaturation), such as Taq polymerase. Suitable enzyme will facilitate combination of the nucleotides in the proper manner to form the primer extension products which are complementary to each polymorphic locus nucleic acid strand. Generally, the synthesis will be initiated at the 3' end of each primer and proceed in the 5' direction along the template strand, until synthesis terminates, producing molecules of different lengths.

The newly synthesized strand and its complementary nucleic acid strand will form a double-stranded molecule under hybridizing conditions described above and this hybrid is used in subsequent steps of the method. In the next step, the newly synthesized double-stranded molecule is subjected to denaturing conditions using any of the procedures described above to provide single-stranded molecules.

The steps of denaturing, annealing, and extension product synthesis can be repeated as often as needed to amplify the target polymorphic locus nucleic acid sequence to the extent necessary for detection. The amount of the specific nucleic acid sequence produced will accumulate in an exponential fashion. *PCR. A Practical Approach*, ILR Press, Eds. McPherson et al. (1992).

The amplification products may be detected by Southern blot analysis with or without using radioactive probes. In one such method, for example, a small sample of DNA containing a very low level of the nucleic acid sequence of the polymorphic locus is amplified, and analyzed via a Southern blotting technique or similarly, using dot blot analysis. The use of non-radioactive probes or labels is facilitated by the high level of the amplified signal. Alternatively, probes used to detect the amplified products can be directly or indirectly detectably labeled, for example, with a radioisotope, a fluorescent compound, a bioluminescent compound, a chemiluminescent compound, a metal chelator or an enzyme. Those of ordinary skill in the art will know of other suitable labels for binding to the probe, or will be able to ascertain such, using routine experimentation.

Sequences amplified by the methods of the invention can be further evaluated, detected, cloned, sequenced, and the like, either in solution or after binding to a solid support, by any method usually applied to the detection of a specific DNA sequence such as dideoxy sequencing, PCR, oligomer restriction (Saiki et al., *Bio/Technology* 3:1008-1012 (1985), allele-specific oligonucleotide (ASO) probe analysis (Conner et al., *Proc. Natl. Acad. Sci. U.S.A.* 80:278 (1983), oligonucleotide ligation assays (OLAs) (Landgren et. al., *Science* 241:1007, 1988), and the like. Molecular techniques for DNA analysis have been reviewed (Landgren et. al., *Science* 242:229-237, 1988).

Preferably, the method of amplifying is by PCR, as described herein and in U.S. Pat. Nos. 4,683,195; 4,683,202; and 4,965,188 each of which is hereby incorporated by reference; and as is commonly used by those of ordinary skill in the art. Alternative methods of amplification have been described and can also be employed as long as the CPSI locus

amplified by PCR using primers of the invention is similarly amplified by the alternative means. Such alternative amplification systems include but are not limited to self-sustained sequence replication, which begins with a short sequence of RNA of interest and a T7 promoter. Reverse transcriptase copies the RNA into cDNA and degrades the RNA, followed by reverse transcriptase polymerizing a second strand of DNA.

Another nucleic acid amplification technique is nucleic acid sequence-based amplification (NASBA™) which uses reverse transcription and T7 RNA polymerase and incorporates two primers to target its cycling scheme. NASBA™ amplification can begin with either DNA or RNA and finish with either, and amplifies to about 10⁸ copies within 60 to 90 minutes.

Alternatively, nucleic acid can be amplified by ligation activated transcription (LAT). LAT works from a single-stranded template with a single primer that is partially single-stranded and partially double-stranded. Amplification is initiated by ligating a cDNA to the promoter oligonucleotide and within a few hours, amplification is about 10⁸ to about 10⁹ fold. The QB replicase system can be utilized by attaching an RNA sequence called MDV-1 to RNA complementary to a DNA sequence of interest. Upon mixing with a sample, the hybrid RNA finds its complement among the specimen's mRNAs and binds, activating the replicase to copy the tag-along sequence of interest.

Another nucleic acid amplification technique, ligase chain reaction (LCR), works by using two differently labeled halves of a sequence of interest which are covalently bonded by ligase in the presence of the contiguous sequence in a sample, forming a new target. The repair chain reaction (RCR) nucleic acid amplification technique uses two complementary and target-specific oligonucleotide probe pairs, thermostable polymerase and ligase, and DNA nucleotides to geometrically amplify targeted sequences. A 2-base gap separates the oligo probe pairs, and the RCR fills and joins the gap, mimicking normal DNA repair.

Nucleic acid amplification by strand displacement activation (SDA) utilizes a short primer containing a recognition site for HincII with short overhang on the 5' end which binds to target DNA. A DNA polymerase fills in the part of the primer opposite the overhang with sulfur-containing adenine analogs. HincII is added but only cuts the unmodified DNA strand. A DNA polymerase that lacks 5' exonuclease activity enters at the site of the nick and begins to polymerize, displacing the initial primer strand downstream and building a new one which serves as more primer.

SDA produces greater than about a 10⁷-fold amplification in 2 hours at 37° C. Unlike PCR and LCR, SDA does not require instrumented temperature cycling. Another amplification system useful in the method of the invention is the QB Replicase System. Although PCR is the preferred method of amplification if the invention, these other methods can also be used to amplify the CPSI locus as described in the method of the invention. Thus, the term "amplification technique" as used herein and in the claims is meant to encompass all the foregoing methods.

In another embodiment of the invention a method is provided for diagnosing or identifying a subject having a predisposition or higher susceptibility to (at risk of) hyperammonemia, comprising sequencing a target nucleic acid of a sample from a subject by dideoxy sequencing, preferably following amplification of the target nucleic acid.

In another embodiment of the invention a method is provided for diagnosing a subject having a predisposition or higher susceptibility to (at risk of) hyperammonemia, com-

prising contacting a target nucleic acid of a sample from a subject with a reagent that detects the presence of the CPSI polymorphism and detecting the reagent.

Another method comprises contacting a target nucleic acid of a sample from a subject with a reagent that detects the presence of the C to A transversion at base 4340, i.e. within exon 36, and detecting the transversion. A number of hybridization methods are well known to those skilled in the art. Many of them are useful in carrying out the invention.

Hepatic veno-occlusive disease (HVOD) is a common toxicity in bone marrow transplant (BMT). It occurs in approximately 20 to 40% of patients and is associated with severe morbidity and mortality. In accordance with the present invention, the frequency of both CPSI alleles was tested in an HVOD and a non-HVOD group undergoing BMT in an effort to identify evidence of disequilibrium. The results indicated the CPSI polymorphism disclosed herein effects susceptibility to a BMT toxicity. Thus, a method of screening subjects for susceptibility to BMT toxicity, and particularly to HVOD, via detection of the CPSI polymorphism is provided in accordance with the present invention.

The materials for use in the method of the invention are ideally suited for the preparation of a diagnostic kit. Such a kit may comprise a carrier means being compartmentalized to receive in close confinement one or more container means such as vials, tubes, and the like, each of the container means comprising one of the separate elements to be used in the method. For example, one of the container means may comprise means for amplifying CPSI DNA, the means comprising the necessary enzyme(s) and oligonucleotide primers for amplifying said target DNA from the subject.

The oligonucleotide primers include primers having a sequence selected from the group including, but not limited to: SEQ ID NOs:6-10, or primer sequences substantially complementary or substantially homologous thereto. The target flanking 5' and 3' polynucleotide sequence has substantially the sequence set forth in SEQ ID NO:5, and sequences substantially complementary or homologous thereto. Other oligonucleotide primers for amplifying CPSI will be known or readily ascertainable to those of skill in the art given the disclosure of the present invention presented herein.

A kit in accordance with the present invention can further comprise a reagent or reagents for extracting a nucleic acid sample from a biological sample obtained from a subject. Any such reagents as would be readily apparent to one of ordinary skill in the art are contemplated to fall within the scope of the present invention. By way of particular example, a suitable lysis buffer for the tissue along with a suspension of glass beads for capturing the nucleic acid sample and an elution buffer for eluting the nucleic acid sample off of the glass beads comprise reagents for extracting a nucleic acid sample from a biological sample obtained from a subject.

Other examples include commercially available, such as the GENOMIC ISOLATION KIT A.S.A.P.TM (Boehringer Mannheim, Indianapolis, Ind.), Genomic DNA Isolation System (GIBCO BRL, Gaithersburg, Md.), ELU-QUIKTM DNA Purification Kit (Schleicher & Schuell, Keene, N.H.), DNA Extraction Kit (Stratagene, La Jolla, Calif.), TURBOGENTM Isolation Kit (Invitrogen, San Diego, Calif.), and the like. Use of these kits according to the manufacturer's instructions is generally acceptable for purification of DNA prior to practicing the methods of the present invention.

C. Definitions Affecting CPSI-Encoding Polynucleotide and CPSI Polypeptides Encoded by Same

In accordance with the present invention, purified and isolated CPSI-encoding polynucleotides and CPSI polypeptides encoded by same are provided. A particularly provided CPSI-

encoding polynucleotide comprises a CPSI encoding polynucleotide which includes a C to A transversion at base 4340, i.e. within exon 36, of the CPSI gene which changes the triplet code from ACC to AAC and leads to the T1405N change in the encoded CPSI polypeptide. The encoded CPSI polypeptide comprising the T1405N change is also particularly provided. Thus, allelic variant polynucleotides and polypeptides encoded by same are provided in accordance with the present invention. Further, a biologically active CPSI polypeptide is also provided in accordance with the present invention, as is a CPSI-encoding polynucleotide encoding such a CPSI polypeptide. Exemplary biological activities include the biological activity of mediating the first step of the urea cycle and the biological activity of cross-reacting with an anti-CPSI antibody.

The provided CPSI-encoding polynucleotides and polypeptides have broad utility given the biological significance of the urea cycle, as is known in the art. By way of example, the CPSI-encoding polynucleotides and polypeptides are useful in the preparation of screening assays and assay kits that are used to detect the presence of the proteins and nucleic acids of this invention in biological samples. Additionally, it is well known that isolated and purified polypeptides have utility as feed additives for livestock and polynucleotides encoding the polypeptides are thus useful in producing the polypeptides.

Preferably, the provided CPSI polynucleotides and polypeptides are isolated from vertebrate and invertebrate sources. Thus, homologs of CPSI, including, but not limited to, mammalian, yeast and bacterial homologs are provided in accordance with the present invention. Preferred mammalian homologs of CPSI members include, but are not limited to, rat and human homologs.

The terms "CPSI gene product", "CPSI protein" and "CPSI polypeptide" refer to proteins having amino acid sequences which are substantially identical to the native amino acid sequences in CPSI and which are biologically active in that they are capable of mediating the synthesis of carbamyl phosphate in the urea cycle, or cross-reacting with anti-CPSI antibodies raised against a CPSI polypeptide.

The terms "CPSI gene product", "CPSI protein" and "CPSI polypeptide" also include analogs of CPSI molecules which exhibit at least some biological activity in common with native CPSI gene products. Furthermore, those skilled in the art of mutagenesis will appreciate that other analogs, as yet undisclosed or undiscovered, may be used to construct CPSI analogs. There is no need for an "CPSI gene product", "CPSI protein" or "CPSI polypeptide" to comprise all, or substantially all of the amino acid sequence of a native CPSI gene product. Shorter or longer sequences are anticipated to be of use in the invention. Thus, the term "CPSI gene product" also includes fusion or recombinant CPSI polypeptides and proteins. Methods of preparing such proteins are described herein.

The terms "CPSI-encoding polynucleotide", "CPSI gene", "CPSI gene sequence" and "CPSI gene segment" refer to any DNA sequence that is substantially identical to a polynucleotide sequence encoding a CPSI gene product, CPSI protein or CPSI polypeptide as defined above. The terms also refer to RNA, or antisense sequences, compatible with such DNA sequences. A "CPSI-encoding polynucleotide", "CPSI gene", "CPSI gene sequence" and "CPSI gene segment" may also comprise any combination of associated control sequences.

The term "substantially identical", when used to define either a CPSI gene product or CPSI amino acid sequence, or a CPSI gene or CPSI nucleic acid sequence, means that a

15

particular sequence, for example, a mutant sequence, varies from the sequence of a natural CPSI by one or more deletions, substitutions, or additions, the net effect of which is to retain at least some of biological activity of CPSI. Alternatively, DNA analog sequences are “substantially identical” to specific DNA sequences disclosed herein if: (a) the DNA analog sequence is derived from coding regions of the natural CPSI gene; or (b) the DNA analog sequence is capable of hybridization of DNA sequences of (a) under moderately stringent conditions and which encode biologically active CPSI gene product; or (c) the DNA sequences are degenerative as a result of the genetic code to the DNA analog sequences defined in (a) and/or (b). Substantially identical analog proteins will be greater than about 60% identical to the corresponding sequence of the native protein. Sequences having lesser degrees of similarity but comparable biological activity are considered to be equivalents. In determining nucleic acid sequences, all subject nucleic acid sequences capable of encoding substantially similar amino acid sequences are considered to be substantially similar to a reference nucleic acid sequence, regardless of differences in codon sequences.

C.1. Percent Similarity

Percent similarity may be determined, for example, by comparing sequence information using the GAP computer program, available from the University of Wisconsin Geneticist Computer Group. The GAP program utilizes the alignment method of Needleman et al., *J. Mol. Biol.* 48:443 (1970), as revised by Smith et al., *Adv. Appl. Math.* 2:482 (1981). Briefly, the GAP program defines similarity as the number of aligned symbols (i.e. nucleotides or amino acids) which are similar, divided by the total number of symbols in the shorter of the two sequences. The preferred default parameters for the GAP program include: (1) a unitary comparison matrix (containing a value of 1 for identities and 0 for non-identities) of nucleotides and the weighted comparison matrix of Gribskov et al., *Nucl. Acids. Res.* 14:6745 (1986), as described by Schwartz et al., eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 357-358 (1979); (2) a penalty of 3.0 for each gap and an additional 0.01 penalty for each symbol and each gap; and (3) no penalty for end gaps. Other comparison techniques are described in the Examples.

The term “homology” describes a mathematically based comparison of sequence similarities which is used to identify genes or proteins with similar functions or motifs. Accordingly, the term “homology” is synonymous with the term “similarity” and “percent similarity” as defined above. Thus, the phrases “substantial homology” or “substantial similarity” have similar meanings.

C.2. Nucleic Acid Sequences

In certain embodiments, the invention concerns the use of CPSI genes and gene products that include within their respective sequences a sequence which is essentially that of a CPSI gene, or the corresponding protein. The term “a sequence essentially as that of a CPSI gene”, means that the sequence substantially corresponds to a portion of a CPSI polypeptide or CPSI encoding polynucleotide and has relatively few bases or amino acids (whether DNA or protein) which are not identical to those of a CPSI protein or CPSI gene, (or a biologically functional equivalent of, when referring to proteins). The term “biologically functional equivalent” is well understood in the art and is further defined in detail herein. Accordingly, sequences which have between about 70% and about 80%; or more preferably, between about 81% and about 90%; or even more preferably, between about 91% and about 99%; of amino acids which are identical or

16

functionally equivalent to the amino acids of a CPSI protein or CPSI gene, will be sequences which are “essentially the same”.

CPSI gene products and CPSI genes which have functionally equivalent codons are also covered by the invention. The term “functionally equivalent codon” is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine, and also to refer to codons that encode biologically equivalent amino acids (see Table 1).

TABLE 1

Table of the Genetic Code						
Amino Acids			Codons			
Alanine	Ala	A	GCA	GCC	GCG	GCU
Cysteine	Cys	C	UGC	UGU		
Aspartic Acid	Asp	D	GAC	GAU		
Glumatic acid	Glu	E	GAA	GAG		
Phenylalanine	Phe	F	UUC	UUU		
Glycine	Gly	G	GGA	GGC	GGG	GGU
Histidine	His	H	CAC	CAU		
Isoleucine	Ile	I	AUA	AUC	AUU	
Lysine	Lys	K	AAA	AAG		
Leucine	Leu	L	UUA	UUG	CUA	CUC
			CUG	CUU		
Methionine	Met	M	AUG			
Asparagine	Asn	N	AAC	AAU		
Proline	Pro	P	CCA	CCC	CCG	CCU
Glutamine	Gln	Q	CAA	CAG		
Arginine	Arg	R	AGA	AGG	CGA	CGC
			CGG	CGU		
Serine	Ser	S	ACG	AGU	UCA	UCC
			UCG	UCU		
Threonine	Thr	T	ACA	ACC	ACG	ACU
Valine	Val	V	GUA	GUC	GUG	GUU
Tryptophan	Trp	W	UGG			
Tyrosine	Tyr	Y	UAC	UAU		

It will also be understood that amino acid and nucleic acid sequences may include additional residues, such as additional N- or C-terminal amino acids or 5' or 3' sequences, and yet still be essentially as set forth in one of the sequences disclosed herein, so long as the sequence meets the criteria set forth above, including the maintenance of biological protein activity where protein expression is concerned. The addition of terminal sequences particularly applies to nucleic acid sequences which may, for example, include various non-coding sequences flanking either of the 5' or 3' portions of the coding region or may include various internal sequences, i.e., introns, which are known to occur within genes.

The present invention also encompasses the use of DNA segments which are complementary, or essentially complementary, to the sequences set forth in the specification. Nucleic acid sequences which are “complementary” are those which are base-pairing according to the standard Watson-Crick complementarity rules. As used herein, the term “complementary sequences” means nucleic acid sequences

which are substantially complementary, as may be assessed by the same nucleotide comparison set forth above, or as defined as being capable of hybridizing to the nucleic acid segment in question under relatively stringent conditions such as those described herein. A particular example of a contemplated complementary nucleic acid segment is an anti-sense oligonucleotide.

Nucleic acid hybridization will be affected by such conditions as salt concentration, temperature, or organic solvents, in addition to the base composition, length of the complementary strands, and the number of nucleotide base mismatches between the hybridizing nucleic acids, as will be readily appreciated by those skilled in the art. Stringent temperature conditions will generally include temperatures in excess of 30° C., typically in excess of 37° C., and preferably in excess of 45° C. Stringent salt conditions will ordinarily be less than 1,000 mM, typically less than 500 mM, and preferably less than 200 mM. However, the combination of parameters is much more important than the measure of any single parameter. (See e.g., Wetmur & Davidson, *J. Mol. Biol.* 31:349-370 (1968)).

Probe sequences may also hybridize specifically to duplex DNA under certain conditions to form triplex or other higher order DNA complexes. The preparation of such probes and suitable hybridization conditions are well known in the art.

As used herein, the term "DNA segment" refers to a DNA molecule which has been isolated free of total genomic DNA of a particular species. Furthermore, a DNA segment encoding a CPSI polypeptide refers to a DNA segment which contains CPSI coding sequences, yet is isolated away from, or purified free from, total genomic DNA of a source species, such as *Homo sapiens*. Included within the term "DNA segment" are DNA segments and smaller fragments of such segments, and also recombinant vectors, including, for example, plasmids, cosmids, phages, viruses, and the like.

Similarly, a DNA segment comprising an isolated or purified CPSI gene refers to a DNA segment including CPSI coding sequences isolated substantially away from other naturally occurring genes or protein encoding sequences. In this respect, the term "gene" is used for simplicity to refer to a functional protein, polypeptide or peptide encoding unit. As will be understood by those in the art, this functional term includes both genomic sequences and cDNA sequences. "Isolated substantially away from other coding sequences" means that the gene of interest, in this case, the CPSI gene, forms the significant part of the coding region of the DNA segment, and that the DNA segment does not contain large portions of naturally-occurring coding DNA, such as large chromosomal fragments or other functional genes or cDNA coding regions. Of course, this refers to the DNA segment as originally isolated, and does not exclude genes or coding regions later added to the segment by the hand of man.

In particular embodiments, the invention concerns isolated DNA segments and recombinant vectors incorporating DNA sequences which encode a CPSI polypeptide that includes within its amino acid sequence an amino acid sequence of any of SEQ ID NOs:2, 4, 12 and 14. In other particular embodiments, the invention concerns isolated DNA segments and recombinant vectors incorporating DNA sequences which encode a protein that includes within its amino acid sequence the amino acid sequence of a CPSI polypeptide corresponding to human tissues.

It will also be understood that this invention is not limited to the particular nucleic acid and amino acid sequences of SEQ ID NOs:1-4 and 11-14. Recombinant vectors and isolated DNA segments may therefore variously include the CPSI polypeptide-encoding region itself, include coding

regions bearing selected alterations or modifications in the basic coding region, or include encoded larger polypeptides which nevertheless include CPSI polypeptide-encoding regions or may encode biologically functional equivalent proteins or peptides which have variant amino acid sequences.

In certain embodiments, the invention concerns isolated DNA segments and recombinant vectors which encode a protein or peptide that includes within its amino acid sequence an amino acid sequence essentially as set forth in any of SEQ ID NOs:2, 4, 12 and 14. Naturally, where the DNA segment or vector encodes a full length CPSI gene product, the most preferred nucleic acid sequence is that which is essentially as set forth in any of SEQ ID NOs: 1, 3, 11 and 13 and which encode a protein that exhibits activity in the urea cycle, as may be determined by, for example, calorimetric assays to detect production of carbonyl phosphate from ammonia, as disclosed herein in Example 3.

The term "a sequence essentially as set forth in any of SEQ ID NO:2, 4, 12 and 14" means that the sequence substantially corresponds to a portion an amino acid sequence either of SEQ ID NOs:2, 4, 12 and 14 and has relatively few amino acids which are not identical to, or a biologically functional equivalent of, the amino acids of an amino acid sequence of any of SEQ ID NOs:2, 4, 12 and 14. The term "biologically functional equivalent" is well understood in the art and is further defined in detail herein. Accordingly, sequences, which have between about 70% and about 80%; or more preferably, between about 81% and about 90%; or even more preferably, between about 91% and about 99%; of amino acids which are identical or functionally equivalent to the amino acids in any of SEQ ID NOs: 2, 4, 12 and 14, will be sequences which "a sequence essentially as set forth in SEQ ID NOs:2, 4, 12 and 14".

In particular embodiments, the invention concerns gene therapy methods that use isolated DNA segments and recombinant vectors incorporating DNA sequences which encode a protein that includes within its amino acid sequence an amino acid sequence of any of SEQ ID NOs:2, 4, 12 and 14, SEQ ID NOs:2, 4, 12 and 14 including sequences which are derived from human tissue. In other particular embodiments, the invention concerns isolated DNA sequences and recombinant DNA vectors incorporating DNA sequences which encode a protein that includes within its amino acid sequence the amino acid sequence of the CPSI protein from human hepatic tissue.

In certain other embodiments, the invention concerns isolated DNA segments and recombinant vectors that include within their sequence a nucleic acid sequence essentially as set forth in any of SEQ ID NO:1, 3, 11 and 13. The term "a sequence essentially as set forth in any of SEQ ID NO:1, 3, 11 and 13" is used in the same sense as described above and means that the nucleic acid sequence substantially corresponds to a portion of any of SEQ ID NOs:1, 3, 11 and 13, respectively, and has relatively few codons which are not identical, or functionally equivalent, to the codons of any of SEQ ID NOs:1, 3, 11 and 13, respectively. Again, DNA segments which encode gene products exhibiting activity in the urea cycle, cross-reactivity with an anti-CPSI antibody, or other biological activity of the CPSI gene product will be most preferred. The term "functionally equivalent codon" is used herein to refer to codons that encode the same amino acid, such as the six codons for arginine or serine, and also to refer to codons that encode biologically equivalent amino acids (see Table 1).

The nucleic acid segments of the present invention, regardless of the length of the coding sequence itself, may be combined with other DNA sequences, such as promoters, enhanc-

ers, polyadenylation signals, additional restriction enzyme sites, multiple cloning sites, other coding segments, and the like, such that their overall length may vary considerably. It is therefore contemplated that a nucleic acid fragment of almost any length may be employed, with the total length preferably being limited by the ease of preparation and use in the intended recombinant DNA protocol. For example, nucleic acid fragments may be prepared which include a short stretch complementary to a nucleic acid sequence set for in any of SEQ ID NOs:1, 3, 11 and 13 respectively, such as about 10 nucleotides, and which are up to 10,000 or 5,000 base pairs in length, with segments of 3,000 being preferred in certain cases. DNA segments with total lengths of about 1,000, 500, 200, 100 and about 50 base pairs in length are also contemplated to be useful.

The DNA segments of the present invention encompass biologically functional equivalent CPSI proteins and peptides. Such sequences may arise as a consequence of codon redundancy and functional equivalency which are known to occur naturally within nucleic acid sequences and the proteins thus encoded. Alternatively, functionally equivalent proteins or peptides may be created via the application of recombinant DNA technology, in which changes in the protein structure may be engineered, based on considerations of the properties of the amino acids being exchanged, e.g. substitution of Ile and Leu at amino acids 4 and 5 is SEQ ID NOs:11-14. Changes designed by man may be introduced through the application of site-directed mutagenesis techniques, e.g., to introduce improvements to the antigenicity of the protein or to test CPSI mutants in order to examine activity in the urea cycle, or other activity at the molecular level.

If desired, one may also prepare fusion proteins and peptides, e.g., where the CPSI coding region is aligned within the same expression unit with other proteins or peptides having desired functions, such as for purification or immunodetection purposes (e.g., proteins which may be purified by affinity chromatography and enzyme label coding regions, respectively).

Recombinant vectors form important further aspects of the present invention. Particularly useful vectors are contemplated to be those vectors in which the coding portion of the DNA segment is positioned under the control of a promoter. The promoter may be in the form of the promoter which is naturally associated with the CPSI gene, e.g., in mammalian tissues, as may be obtained by isolating the 5' non-coding sequences located upstream of the coding segment or exon, for example, using recombinant cloning and/or PCR technology, in connection with the compositions disclosed herein.

In other embodiments, it is contemplated that certain advantages will be gained by positioning the coding DNA segment under the control of a recombinant, or heterologous, promoter. As used herein, a recombinant or heterologous promoter is intended to refer to a promoter that is not normally associated with a CPSI gene in its natural environment. Such promoters may include promoters isolated from bacterial, viral, eukaryotic, or mammalian cells. Naturally, it will be important to employ a promoter that effectively directs the expression of the DNA segment in the cell type chosen for expression. The use of promoter and cell type combinations for protein expression is generally known to those of skill in the art of molecular biology, for example, see Sambrook et al., 1989, incorporated herein by reference. The promoters employed may be constitutive, or inducible, and can be used under the appropriate conditions to direct high level expression of the introduced DNA segment, such as is advantageous in the large-scale production of recombinant proteins or peptides. Appropriate promoter systems provided for use in high-

level expression include, but are not limited to, the vaccinia virus promoter and the baculovirus promoter.

In an alternative embodiment, the present invention provides an expression vector comprising a polynucleotide that encodes a CPSI polypeptide having activity in the urea cycle, cross-reacting with an anti-CPSI antibody, or other biological activity in accordance with the present invention. Also preferably, an expression vector of the present invention comprises a polynucleotide that encodes a human CPSI gene product. More preferably, an expression vector of the present invention comprises a polynucleotide that encodes a polypeptide comprising an amino acid residue sequence of any of SEQ ID NOs:2, 4, 12 and 14. More preferably, an expression vector of the present invention comprises a polynucleotide comprising the nucleotide base sequence of any of SEQ ID NO:1, 3, 11 and 13.

Even more preferably, an expression vector of the invention comprises a polynucleotide operatively linked to an enhancer-promoter. More preferably still, an expression vector of the invention comprises a polynucleotide operatively linked to a prokaryotic promoter. Alternatively, an expression vector of the present invention comprises a polynucleotide operatively linked to an enhancer-promoter that is a eukaryotic promoter, and the expression vector further comprises a polyadenylation signal that is positioned 3' of the carboxy-terminal amino acid and within a transcriptional unit of the encoded polypeptide.

In yet another embodiment, the present invention provides a recombinant host cell transfected with a polynucleotide that encodes a CPSI polypeptide having activity in the modulation of the urea cycle, cross-reactivity with an anti-CPSI antibody, or other biological activity in accordance with the present invention. SEQ ID NOs: 1-4 and 11-14 set forth nucleotide and amino acid sequences from an exemplary vertebrate, human. Also provided by the present invention are homologous or biologically equivalent polynucleotides and CPSI polypeptides found in other vertebrates, including rat. Also provided by the present invention are homologous or biologically equivalent polynucleotides and CPSI polypeptides found in invertebrates, including bacteria and yeast.

Preferably, a recombinant host cell of the present invention is transfected with the polynucleotide that encodes human CPSI polypeptide. More preferably, a recombinant host cell of the present invention is transfected with the polynucleotide sequence of any of SEQ ID NOs:1, 3, 11 and 13. Even more preferably, a host cell of the invention is a eukaryotic host cell. Still more preferably, a recombinant host cell of the present invention is a vertebrate cell. Preferably, a recombinant host cell of the invention is a mammalian cell.

In another aspect, a recombinant host cell of the present invention is a prokaryotic host cell. Preferably, a recombinant host cell of the invention is a bacterial cell, preferably a strain of *Escherichia coli*. More preferably, a recombinant host cell comprises a polynucleotide under the transcriptional control of regulatory signals functional in the recombinant host cell, wherein the regulatory signals appropriately control expression of the CPSI polypeptide in a manner to enable all necessary transcriptional and post-transcriptional modification.

In yet another embodiment, the present invention provides a method of preparing a CPSI polypeptide comprising transfecting a cell with polynucleotide that encodes a CPSI polypeptide having activity in the urea cycle, cross-reacting with an anti-CPSI antibody, or other biological activity in accordance with the present invention, to produce a transformed host cell; and maintaining the transformed host cell under biological conditions sufficient for expression of the polypeptide. More preferably, the transformed host cell is a

eukaryotic cell. More preferably still, the eukaryotic cell is a vertebrate cell. Alternatively, the host cell is a prokaryotic cell. More preferably, the prokaryotic cell is a bacterial cell of *Escherichia coli*. Even more preferably, a polynucleotide transfected into the transformed cell comprises a nucleotide base sequence of any of SEQ ID NOs: 1, 3, 11 and 13. SEQ ID NOs: 1-4 and 11-14 set forth nucleotide and amino acid sequences for an exemplary vertebrate, human. Also provided by the present invention are homologues or biologically equivalent CPSI polynucleotides and polypeptides found in other vertebrates, particularly warm blooded vertebrates, and more particularly rat. Also provided by the present invention are homologous or biologically equivalent polynucleotides and CPSI polypeptides found in invertebrates, including bacteria and yeast.

As mentioned above, in connection with expression embodiments to prepare recombinant CPSI proteins and peptides, it is contemplated that longer DNA segments will most often be used, with DNA segments encoding the entire CPSI protein, functional domains or cleavage products thereof, being most preferred. However, it will be appreciated that the use of shorter DNA segments to direct the expression of CPSI peptides or epitopic core regions, such as may be used to generate anti-CPSI antibodies, also falls within the scope of the invention.

DNA segments which encode peptide antigens from about 15 to about 50 amino acids in length, or more preferably, from about 15 to about 30 amino acids in length are contemplated to be particularly useful. DNA segments encoding peptides will generally have a minimum coding length in the order of about 45 to about 150, or to about 90 nucleotides. DNA segments encoding full length proteins may have a minimum coding length on the order of about 4,500 to about 4,600 nucleotides for a protein in accordance with any of SEQ ID NOs: 2, 4, 12 and 14.

Naturally, the present invention also encompasses DNA segments which are complementary, or essentially complementary, to the sequences set forth in any of SEQ ID NOs: 1, 3, 11 and 13. The terms "complementary" and "essentially complementary" are defined above. Excepting intronic or flanking regions, details of which are disclosed graphically in FIG. 9, and allowing for the degeneracy of the genetic code, sequences which have between about 70% and about 80%; or more preferably, between about 81% and about 90%; or even more preferably, between about 91% and about 99%; of nucleotides which are identical or functionally equivalent (i.e. encoding the same amino acid) of nucleotides in any of SEQ ID NOs: 1, 3, 11 and 13 will be sequences which are "a sequence essentially as set forth in any of SEQ ID NOs: 1, 3, 11 and 13". Sequences which are essentially the same as those set forth in any of SEQ ID NOs: 1, 3, 11 and 13 may also be functionally defined as sequences which are capable of hybridizing to a nucleic acid segment containing the complement in any of SEQ ID NOs: 1, 3, 11 and 13 under relatively stringent conditions. Suitable relatively stringent hybridization conditions are described herein and will be well known to those of skill in the art.

C.2. Biologically Functional Equivalents

As mentioned above, modification and changes may be made in the structure of the CPSI proteins and peptides described herein and still obtain a molecule having like or otherwise desirable characteristics. For example, certain amino acids may be substituted for other amino acids in a protein structure without appreciable loss of interactive capacity with structures such as, for example, in the nucleus of a cell. Since it is the interactive capacity and nature of a protein that defines that protein's biological functional activ-

ity, certain amino acid sequence substitutions can be made in a protein sequence (or, of course, its underlying DNA coding sequence) and nevertheless obtain a protein with like or even countervailing properties (e.g., antagonistic v. agonistic). It is thus contemplated by applicants that various changes may be made in the sequence of the CPSI proteins and peptides (or underlying DNA) without appreciable loss of their biological utility or activity.

It is also well understood by the skilled artisan that, inherent in the definition of a biologically functional equivalent protein or peptide, is the concept that there is a limit to the number of changes that may be made within a defined portion of the molecule and still result in a molecule with an acceptable level of equivalent biological activity. Biologically functional equivalent peptides are thus defined herein as those peptides in which certain, not most or all, of the amino acids may be substituted. Of course, a plurality of distinct proteins/peptides with different substitutions may easily be made and used in accordance with the invention.

It is also well understood that where certain residues are shown to be particularly important to the biological or structural properties of a protein or peptide, e.g., residues in active sites, such residues may not generally be exchanged. This is the case in the present invention, where if any changes, for example, in the phosphorylation domains of a CPSI polypeptide, could result in a loss of an aspect of the utility of the resulting peptide for the present invention.

Amino acid substitutions, such as those which might be employed in modifying the CPSI proteins and peptides described herein, are generally based on the relative similarity of the amino acid side-chain substituents, for example, their hydrophobicity, hydrophilicity, charge, size, and the like. An analysis of the size, shape and type of the amino acid side-chain substituents reveals that arginine, lysine and histidine are all positively charged residues; that alanine, glycine and serine are all a similar size; and that phenylalanine, tryptophan and tyrosine all have a generally similar shape. Therefore, based upon these considerations, arginine, lysine and histidine; alanine, glycine and serine; and phenylalanine, tryptophan and tyrosine; are defined herein as biologically functional equivalents.

In making such changes, the hydropathic index of amino acids may be considered. Each amino acid has been assigned a hydropathic index on the basis of their hydrophobicity and charge characteristics, these are: isoleucine (+4.5); valine (+4.2); leucine (+3.8); phenylalanine (+2.8); cysteine/cystine (+2.5); methionine (+1.9); alanine (+1.8); glycine (-0.4); threonine (-0.7); serine (-0.8); tryptophan (-0.9); tyrosine (-1.3); proline (-1.6); histidine (-3.2); glutamate (-3.5); glutamine (-3.5); aspartate (-3.5); asparagine (-3.5); lysine (-3.9); and arginine (-4.5).

The importance of the hydropathic amino acid index in conferring interactive biological function on a protein is generally understood in the art (Kyte & Doolittle, *J. Mol. Biol.* 157:105-132 (1982), incorporated herein by reference). It is known that certain amino acids may be substituted for other amino acids having a similar hydropathic index or score and still retain a similar biological activity. In making changes based upon the hydropathic index, the substitution of amino acids whose hydropathic indices are within ± 2 of the original value is preferred, those which are within ± 1 of the original value are particularly preferred, and those within ± 0.5 of the original value are even more particularly preferred.

It is also understood in the art that the substitution of like amino acids can be made effectively on the basis of hydrophilicity. U.S. Pat. No. 4,554,101, incorporated herein by reference, states that the greatest local average hydrophilicity

of a protein, as governed by the hydrophilicity of its adjacent amino acids, correlates with its immunogenicity and antigenicity, i.e. with a biological property of the protein. It is understood that an amino acid can be substituted for another having a similar hydrophilicity value and still obtain a biologically equivalent protein.

As detailed in U.S. Pat. No. 4,554,101, the following hydrophilicity values have been assigned to amino acid residues: arginine (+3.0); lysine (+3.0); aspartate (+3.0±1); glutamate (+3.0±1); serine (+0.3); asparagine (+0.2); glutamine (+0.2); glycine (0); threonine (-0.4); proline (-0.5±1); alanine (-0.5); histidine (-0.5); cysteine (-1.0); methionine (-1.3); valine (-1.5); leucine (-1.8); isoleucine (-1.8); tyrosine (-2.3); phenylalanine (-2.5); tryptophan (-3.4).

In making changes based upon similar hydrophilicity values, the substitution of amino acids whose hydrophilicity values are within ±2 of the original value is preferred, those which are within ±1 of the original value are particularly preferred, and those within ±0.5 of the original value are even more particularly preferred.

While discussion has focused on functionally equivalent polypeptides arising from amino acid changes, it will be appreciated that these changes may be effected by alteration of the encoding DNA, taking into consideration also that the genetic code is degenerate and that two or more codons may code for the same amino acid.

C.3. Sequence Modification Techniques

Modifications to the CPSI proteins and peptides described herein may be carried out using techniques such as site directed mutagenesis. Site-specific mutagenesis is a technique useful in the preparation of individual peptides, or biologically functional equivalent proteins or peptides, through specific mutagenesis of the underlying DNA. The technique further provides a ready ability to prepare and test sequence variants, for example, incorporating one or more of the foregoing considerations, by introducing one or more nucleotide sequence changes into the DNA. Site-specific mutagenesis allows the production of mutants through the use of specific oligonucleotide sequences which encode the DNA sequence of the desired mutation, as well as a sufficient number of adjacent nucleotides, to provide a primer sequence of sufficient size and sequence complexity to form a stable duplex on both sides of the deletion junction being traversed. Typically, a primer of about 17 to 30 nucleotides in length is preferred, with about 5 to 10 residues on both sides of the junction of the sequence being altered.

In general, the technique of site-specific mutagenesis is well known in the art as exemplified by publications (e.g., Adelman et al., 1983). As will be appreciated, the technique typically employs a phage vector which exists in both a single stranded and double stranded form. Typical vectors useful in site-directed mutagenesis include vectors such as the M13 phage (Messing et al., 1981). These phage are readily commercially available and their use is generally well known to those skilled in the art. Double stranded plasmids are also routinely employed in site directed mutagenesis which eliminates the step of transferring the gene of interest from a plasmid to a phage.

In general, site-directed mutagenesis in accordance herewith is performed by first obtaining a single-stranded vector or melting apart the two strands of a double stranded vector which includes within its sequence a DNA sequence which encodes, for example, a human CPSI polypeptide. An oligonucleotide primer bearing the desired mutated sequence is prepared, generally synthetically, for example by the method of Crea et al. (1978). This primer is then annealed with the

single-stranded vector, and subjected to DNA polymerizing enzymes such as *E. coli* polymerase I Klenow fragment, in order to complete the synthesis of the mutation-bearing strand. Thus, a heteroduplex is formed wherein one strand encodes the original non-mutated sequence and the second strand bears the desired mutation. This heteroduplex vector is then used to transform appropriate cells, such as *E. coli* cells, and clones are selected which include recombinant vectors bearing the mutated sequence arrangement.

The preparation of sequence variants of the selected gene using site-directed mutagenesis is provided as a means of producing potentially useful CPSI polypeptide or other species having activity in the urea cycle and is not meant to be limiting as there are other ways in which sequence variants of these peptides may be obtained. For example, recombinant vectors encoding the desired genes may be treated with mutagenic agents to obtain sequence variants (see, e.g., a method described by Eichenlaub, 1979) for the mutagenesis of plasmid DNA using hydroxylamine.

C.4. Other Structural Equivalents

In addition to the CPSI peptidyl compounds described herein, the inventors also contemplate that other sterically similar compounds may be formulated to mimic the key portions of the peptide structure. Such compounds may be used in the same manner as the peptides of the invention and hence are also functional equivalents. The generation of a structural functional equivalent may be achieved by the techniques of modeling and chemical design known to those of skill in the art. It will be understood that all such sterically similar constructs fall within the scope of the present invention.

D. Introduction of Gene Products

Where the gene itself is employed to introduce the gene products, a convenient method of introduction will be through the use of a recombinant vector which incorporates the desired gene, together with its associated control sequences. The preparation of recombinant vectors is well known to those of skill in the art and described in many references, such as, for example, Sambrook et al. (1989), specifically incorporated herein by reference.

In vectors, it is understood that the DNA coding sequences to be expressed, in this case those encoding the CPSI gene products, are positioned adjacent to and under the control of a promoter. It is understood in the art that to bring a coding sequence under the control of such a promoter, one generally positions the 5' end of the transcription initiation site of the transcriptional reading frame of the gene product to be expressed between about 1 and about 50 nucleotides "downstream" of (i.e., 3' of) the chosen promoter. One may also desire to incorporate into the transcriptional unit of the vector an appropriate polyadenylation site (e.g., 5'-AATAAA-3'), if one was not contained within the original inserted DNA. Typically, these poly A addition sites are placed about 30 to 2000 nucleotides "downstream" of the coding sequence at a position prior to transcription termination.

While use of the control sequences of the specific gene (i.e., a CPSI promoter for a CPSI gene) will be preferred, there is no reason why other control sequences could not be employed, so long as they are compatible with the genotype of the cell being treated. Thus, one may mention other useful promoters by way of example, including, e.g., an SV40 early promoter, a long terminal repeat promoter from retrovirus, an actin promoter, a heat shock promoter, a metallothionein promoter, and the like.

As is known in the art, a promoter is a region of a DNA molecule typically within about 100 nucleotide pairs in front of (upstream of) the point at which transcription begins (i.e.,

a transcription start site). That region typically contains several types of DNA sequence elements that are located in similar relative positions in different genes. As used herein, the term "promoter" includes what is referred to in the art as an upstream promoter region, a promoter region or a promoter of a generalized eukaryotic RNA Polymerase II transcription unit.

Another type of discrete transcription regulatory sequence element is an enhancer. An enhancer provides specificity of time, location and expression level for a particular encoding region (e.g., gene). A major function of an enhancer is to increase the level of transcription of a coding sequence in a cell that contains one or more transcription factors that bind to that enhancer. Unlike a promoter, an enhancer can function when located at variable distances from transcription start sites so long as a promoter is present.

As used herein, the phrase "enhancer-promoter" means a composite unit that contains both enhancer and promoter elements. An enhancer-promoter is operatively linked to a coding sequence that encodes at least one gene product. As used herein, the phrase "operatively linked" means that an enhancer-promoter is connected to a coding sequence in such a way that the transcription of that coding sequence is controlled and regulated by that enhancer-promoter. Means for operatively linking an enhancer-promoter to a coding sequence are well known in the art. As is also well known in the art, the precise orientation and location relative to a coding sequence whose transcription is controlled, is dependent inter alia upon the specific nature of the enhancer-promoter. Thus, a TATA box minimal promoter is typically located from about 25 to about 30 base pairs upstream of a transcription initiation site and an upstream promoter element is typically located from about 100 to about 200 base pairs upstream of a transcription initiation site. In contrast, an enhancer can be located downstream from the initiation site and can be at a considerable distance from that site.

An enhancer-promoter used in a vector construct of the present invention can be any enhancer-promoter that drives expression in a cell to be transfected. By employing an enhancer-promoter with well-known properties, the level and pattern of gene product expression can be optimized.

For introduction of, for example, the human CPSI gene including allelic variations thereof, it is proposed that one will desire to preferably employ a vector construct that will deliver the desired gene to the affected cells. This will, of course, generally require that the construct be delivered to the targeted cells, for example, mammalian hepatic cells. It is proposed that this may be achieved most preferably by introduction of the desired gene through the use of a viral vector to carry the CPSI sequence to efficiently infect the cells. These vectors will preferably be an adenoviral, a retroviral, a vaccinia viral vector or adeno-associated virus. These vectors are preferred because they have been successfully used to deliver desired sequences to cells and tend to have a high infection efficiency. Suitable vector-CPSI gene constructs are adapted for administration as pharmaceutical compositions, as described herein below.

Commonly used viral promoters for expression vectors are derived from polyoma, cytomegalovirus, Adenovirus 2, and Simian Virus 40 (SV40). The early and late promoters of SV40 virus are particularly useful because both are obtained easily from the virus as a fragment which also contains the SV40 viral origin of replication. Smaller or larger SV40 fragments may also be used, provided there is included the approximately 250 bp sequence extending from the HindIII site toward the BglI site located in the viral origin of replication. Further, it is also possible, and often desirable, to utilize

promoter or control sequences normally associated with the desired gene sequence, provided such control sequences are compatible with the host cell systems.

The origin of replication may be provided either by construction of the vector to include an exogenous origin, such as may be derived from SV40 or other viral (e.g., Polyoma, Adeno, VSV, BPV) source, or may be provided by the host cell chromosomal replication mechanism. If the vector is integrated into the host cell chromosome, the latter is often sufficient.

Where a CPSI gene itself is employed it will be most convenient to simply use a wild type CPSI gene directly. The CPSI gene can thus comprise the threonine encoding allele such that amino acid 1405 of the encoded polypeptide comprises threonine. Alternatively, the CPSI gene comprises the arginine encoding allele such that amino acid 1405 of the encoded polypeptide comprises arginine. Additionally, it is envisioned that certain regions of a CPSI gene can be employed exclusively without employing an entire wild type CPSI gene or an entire allelic variant thereof. It is proposed that it will ultimately be preferable to employ the smallest region needed to modulate the urea cycle so that one is not introducing unnecessary DNA into cells which receive a CPSI gene construct. Techniques well known to those of skill in the art, such as the use of restriction enzymes, will allow for the generation of small regions of an exemplary CPSI gene. The ability of these regions to modulate the urea cycle can easily be determined by the assays reported in the Examples. In general, techniques for assessing the modulation of the urea cycle are known in the art.

D.1. Transgenic Animals

It is also provided within the scope of the present invention to prepare a transgenic non-human animal which expresses a CPSI gene of the present invention or in which expression of a CPSI gene is "knocked-out". Provided transgenic non-human animals express either the T1405 form of CPSI or the N1405 form of CPSI. A preferred transgenic animal is a mouse.

Techniques for the preparation of transgenic animals are known in the art. Exemplary techniques are described in U.S. Pat. No. 5,489,742 (transgenic rats); U.S. Pat. Nos. 4,736,866, 5,550,316, 5,614,396, 5,625,125 and 5,648,061 (transgenic mice); U.S. Pat. No. 5,573,933 (transgenic pigs); U.S. Pat. No. 5,162,215 (transgenic avian species) and U.S. Pat. No. 5,741,957 (transgenic bovine species), the entire contents of each of which are herein incorporated by reference.

With respect to an exemplary method for the preparation of a transgenic mouse, cloned recombinant or synthetic DNA sequences or DNA segments encoding a CPSI gene product are injected into fertilized mouse eggs. The injected eggs are implanted in pseudo pregnant females and are grown to term to provide transgenic mice whose cells express a CPSI gene product. Preferably, the injected sequences are constructed having promoter sequences connected so as to express the desired protein in hepatic cells of the transgenic mouse.

D.2. Gene Therapy

CPSI genes can be used for gene therapy in accordance with the present invention. Exemplary gene therapy methods, including liposomal transfection of nucleic acids into host cells, are described in U.S. Pat. Nos. 5,279,833; 5,286,634; 5,399,346; 5,646,008; 5,651,964; 5,641,484; and 5,643,567, the contents of each of which are herein incorporated by reference.

Briefly, CPSI gene therapy directed toward modulation of the urea cycle in a target cell is described. Target cells include but are not limited to hepatic cells and intestinal cells. In one embodiment, a therapeutic method of the present invention

provides a method for modulating of the urea cycle in a cell comprising the steps of: (a) delivering to the cell an effective amount of a DNA molecule comprising a polynucleotide that encodes a CPSI polypeptide that modulates the urea cycle; and (b) maintaining the cell under conditions sufficient for expression of said polypeptide.

Delivery is preferably accomplished by injecting the DNA molecule into the cell. Where the cell is in a subject delivering is preferably administering the DNA molecule into the circulatory system of the subject. In a preferred embodiment, administering comprises the steps of: (a) providing a vehicle that contains the DNA molecule; and (b) administering the vehicle to the subject.

A vehicle is preferably a cell transformed or transfected with the DNA molecule or a transfected cell derived from such a transformed or transfected cell. An exemplary and preferred transformed or transfected cell is a hepatic cell. Means for transforming or transfecting a cell with a DNA molecule of the present invention are set forth above.

Alternatively, the vehicle is a virus or an antibody that specifically infects or immunoreacts with an antigen of the tumor. Retroviruses used to deliver the constructs to the host target tissues generally are viruses in which the 3'-LTR (linear transfer region) has been inactivated. That is, these are enhancerless 3'-LTR's, often referred to as SIN (self-inactivating viruses) because after productive infection into the host cell, the 3'-LTR is transferred to the 5'-end and both viral LTR's are inactive with respect to transcriptional activity. A use of these viruses well known to those skilled in the art is to clone genes for which the regulatory elements of the cloned gene are inserted in the space between the two LTR's. An advantage of a viral infection system is that it allows for a very high level of infection into the appropriate recipient cell.

Antibodies have been used to target and deliver DNA molecules. An N-terminal modified poly-L-lysine (NPLL)-antibody conjugate readily forms a complex with plasmid DNA. A complex of monoclonal antibodies against a cell surface thrombomodulin conjugated with NPLL was used to target a foreign plasmid DNA to an antigen-expressing mouse lung endothelial cell line and mouse lung. Those targeted endothelial cells expressed the product encoded by that foreign DNA.

It is also envisioned that this embodiment of the present invention can be practiced using alternative viral or phage vectors, including retroviral vectors and vaccinia viruses whose genome has been manipulated in alternative ways so as to render the virus non-pathogenic. Methods for creating such a viral mutation are set forth in detail in U.S. Pat. No. 4,769,331, incorporated herein by reference.

By way of specific example, a human CPSI-encoding polynucleotide or a CPSI-encoding polynucleotide homolog from another warm-blooded vertebrate or a CPSI-encoding homolog from an invertebrate source, such as bacteria or yeast is introduced into isolated hepatic cells or other relevant cells. The re-injection of the transgene-carrying cells into the liver or other relevant tissues provides a treatment for susceptibility to hyperammonemia or other relevant diseases in human and animals.

E. Supplementation Therapy

In addition to its role in nitrogen clearance, the urea cycle is the body's intrinsic source of arginine which acts as a precursor of nitric oxide (NO), a potent vasodilator. Methods of treating suboptimal urea cycle function are provided in accordance with the present invention, including treatment by administration of nitric oxide precursors such as citrulline. Typically, the suboptimal urea cycle function is associated with the polymorphism disclosed herein. The sub-optimal

urea cycle function can further comprise hyperammonemia or decreased citrulline and/or arginine production.

The subject to be treated can be suffering from a disorder associated with sub-optimal urea cycle function, such as but not limited to a disorder associated with impaired production of nitric oxide precursors. Such disorders include but are not limited to disorders that involve impaired or damaged liver and/or gut tissue.

Representative disorders include but are not limited to hepatitis (including hepatitis A, B and C), sclerosis, asthma, pulmonary hypertension (including primary and secondary), bone marrow transplant toxicity in a subject undergoing bone marrow transplant, and combinations thereof.

The subject to be treated can also be exposed or about to be exposed to an environmental stimulus associated with sub-optimal urea cycle function. Such environmental stimuli include but are not limited to stimuli that involve impairment or damage to liver and/or gut tissue. Representative environmental stimuli include but are not limited to chemotherapy or other pharmaceutical therapy, cardiac surgery (represented in some situations as increased postoperative pulmonary vascular tone), increased oxidative stress, bone marrow transplant, sepsis, acute asthma attack, hypoxia, hepatotoxin exposure, and combinations thereof. Representative cardiac surgeries include repair of congenital heart defects, and further includes cardiopulmonary bypass used for correction of congenital heart defects. Cardiac defects associated with excess pulmonary blood flow, such as an atrioventricular septal defect (AVSD) or large unrestrictive ventricular septal defect (VSD) are representative cardiac defects. Sustained pulmonary overcirculation can cause hypertrophy and hyperreactivity of pulmonary vascular smooth muscle. Preoperatively, these patients often have congestive heart failure and poor weight gain. Surgical repair is scheduled as early as possible in order to reduce this postoperative complication.

Additional cardiac defect correct procedures are bidirectional Glenn and modified Fontan procedures. In such procedures patients with single ventricle lesions require surgical procedures where success depends on maintenance of low postoperative pulmonary vascular tone. Staged correction of a single ventricle lesion requires a series of 3 surgical procedures aimed at separating the pulmonary and systemic circulations. The first of these procedures, often performed in the neonatal period, is a Blalock-Taussig shunt for those patients with a hypoplastic right ventricle or a Norwood I procedure for those patients with hypoplastic left heart syndrome. The second surgery is a bidirectional Glenn shunt where superior vena cava flow is diverted directly into the pulmonary artery. The third and final stage is a modified Fontan procedure where inferior vena cava flow is diverted into the pulmonary artery, thereby completing separation of the pulmonary and systemic circulations. With the Glenn and Fontan procedures, pulmonary blood flow is entirely passive and relies on an adequate pressure gradient between the venous system (SVC and IVC pressure) and the PA pressure. Any elevation in the pulmonary vascular tone in the immediate postoperative period can lead to decreased pulmonary blood flow and a subsequent fall in cardiac output. On a longer term, elevated pulmonary vascular tone after these procedures can lead to persistent pleural effusions, prolonged requirement for pleural or mediastinal drainage tubes, prolonged ventilation, and prolonged ICU stays.

Additional cardiac defect correct procedures are Norwood I procedures. Postoperative care of infants with hypoplastic left heart syndrome (HLHS) undergoing a Norwood I procedure relies heavily on balancing pulmonary and systemic flow. Abrupt elevations in pulmonary vascular resistance can

cause significant hypoxemia and desaturation. Rarely, low pulmonary vascular resistance can be detrimental if blood flow is shunted to the lungs at the expense of systemic and coronary circulation. With refined surgical techniques and optimal sizing of the central shunt, this complication is much less common than problems with inadequate pulmonary blood flow.

Additional cardiac defect correct procedures are arterial Switch Procedures.

Transposition of the great arteries (TGA) is a complex cardiac lesion that requires surgical correction in the immediate neonatal period. Timing of the arterial switch procedure for correction of TGA specifically takes into account pulmonary vascular tone issues. Frequently, surgery is not performed until 5-7 days of age when perinatal pulmonary vascular tone has partially decreased. Because the right ventricle is the systemic ventricle before surgical correction, postoperative elevations in pulmonary vascular resistance are usually well tolerated and pulmonary artery pressure is usually not measured. However, if postoperative pulmonary vascular tone is increased, it may partially explain why some infants with favorable anatomy and short bypass times still have a complicated postoperative course.

A method of treating or preventing a disorder related to sub-optimal urea cycle function in a subject is provided in accordance with the present invention. The method comprises administering to the subject a therapeutically effective amount of a nitric oxide precursor, whereby treatment or prevention of the disorder is accomplished. The nitric oxide precursor can include but is not limited to citrulline, arginine and combinations thereof. In some embodiments, sub-optimal nitric oxide formation resulting from sub-optimal urea cycle function can be treated.

A method of treating or preventing a disorder selected from the group consisting hepatitis, cirrhosis, pulmonary hypertension (both primary and secondary), necrotizing enterocolitis (NEC), Acute Respiratory Distress Syndrome, ethnic specific endothelial dysfunction, erectile dysfunction, asthma, and combinations thereof, in a subject is also disclosed. In some embodiments the method comprises administering to a subject in need thereof a therapeutically effective amount of a nitric oxide precursor. The administering can be intravenous or oral administration. The nitric oxide precursor can be selected from the group consisting of citrulline, arginine and combinations thereof. In some embodiments the disorder is necrotizing enterocolitis (NEC) and the subject is a premature infant.

A method of raising a level of a nitric acid precursor in a subject in need thereof is also disclosed. In some embodiments the method comprises administering to the subject a therapeutically effective amount of a nitric oxide precursor, whereby a level of a nitric oxide precursor in the subject is raised. The administering can be intravenous or oral administration. The nitric oxide precursor can be selected from the group consisting of citrulline, arginine and combinations thereof.

Optionally, a supplementation therapy method of the present invention further comprises the step of initially detecting a polymorphism of a carbamyl phosphate synthase I (CPSI) gene in the subject. The polymorphism of the carbamyl phosphate synthetase polypeptide preferably comprises a C to A transversion within CPSI exon 36, more preferably comprises a C to A transversion at nucleotide 4340 of a cDNA that corresponds to the CPSI gene, and ever more preferably, the C to A transversion at nucleotide 4340 of the cDNA that corresponds to the CPSI gene further comprises a

change in the triplet code from AAC to ACC, which encodes a CPSI polypeptide having an threonine moiety at amino acid 1405.

A significant decrease in urea cycle intermediates (citrulline, arginine) was observed in subjects undergoing BMT associated with the T1405N CPSI polymorphism disclosed herein. In accordance with the present invention, a method for the treatment or prophylaxis of BMT toxicity, such as HVOD and/or acute lung injury, comprising administering a therapeutically effective amount of a NO precursor, such as citrulline and/or arginine, to a subject in need thereof is also provided in accordance with the present invention. Preferably, the T1405N CPSI polymorphism disclosed herein is present in the subject. More preferably, a therapeutically effective amount of citrulline is administered to the subject.

In accordance with the present invention, a method of reducing toxicity and/or the occurrence of HVOD in a subject undergoing BMT is thus provided. This method comprises administering the BMT subject an effective amount of arginine and/or citrulline, with citrulline being preferred, to bolster arginine and NO synthesis in the subject. The bolstering of arginine and NO synthesis in the subject will reduce and/or substantially prevent the occurrence of HVOD associated with BMT. Citrulline is a preferred supplementation agent given that it is more readily converted to NO. Additionally and preferably, subjects having the CPSI polymorphism of the present invention are contemplated to be preferred candidates for supplementation in accordance with this method.

The subject treated in the present invention in its many embodiments is desirably a human subject, although it is to be understood that the principles of the invention indicate that the invention is effective with respect to all vertebrate species, including warm-blooded vertebrates such as mammals and birds, which are intended to be included in the term "subject". In this context, a mammal is understood to include any mammalian species in which treatment of hyperammonemia, BMT toxicity and other diseases associated with impaired urea cycle function is desirable, particularly agricultural and domestic mammalian species.

Thus, contemplated is the treatment of mammals such as humans, as well as those mammals of importance due to being endangered (such as Siberian tigers), of economical importance (animals raised on farms for consumption by humans) and/or social importance (animals kept as pets or in zoos) to humans, for instance, carnivores other than humans (such as cats and dogs), swine (pigs, hogs, and wild boars), ruminants (such as cattle, oxen, sheep, giraffes, deer, goats, bison, and camels), and horses. Also contemplated is the treatment of birds, including the treatment of those kinds of birds that are endangered, kept in zoos, as well as fowl, and more particularly domesticated fowl, i.e., poultry, such as turkeys, chickens, ducks, geese, guinea fowl, and the like, as they are also of economical importance to humans. Thus, contemplated is the treatment of livestock, including, but not limited to, domesticated swine (pigs and hogs), ruminants, horses, poultry, and the like.

The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. For example, a formulation intended for administration to humans may contain from 0.5 mg to 5 g of active agent compounded with an appropriate and convenient amount of carrier material which may vary from about 5 to about 95 percent of the total composition. For example, in a human adult, the doses per person per administration are generally between 1 mg and 500 mg up to several times per day. Thus, dosage unit forms will generally contain between

from about 1 mg to about 500 mg of an active ingredient, typically 25 mg, 50 mg, 100 mg, 200 mg, 300 mg, 400 mg, 500 mg, 600 mg, 800 mg, or 1000 mg.

The nitric oxide precursor is administered in a dose ranging from about 0.01 mg to about 1,000 mg, preferably in a dose ranging from about 0.5 mg to about 500 mg, and more preferably in a dose ranging from about 1.0 mg to about 250 mg. The nitric oxide precursor can also be administered in a dose ranging from about 100 mg to about 30,000 mg, in some embodiments in a dose ranging from about 250 mg to about 1,000 mg. A representative dose is 3.8 g/m²/day of arginine or citrulline (molar equivalents, MW L-citrulline 175.2, MW L-arginine 174.2).

Representative intravenous citrulline solutions can comprise a 100 mg/ml (10%) solution. Representative intravenous citrulline dosages can comprise 200 mg/kg, 400 mg/kg, 600 mg/kg, and 800 mg/kg. In some embodiments, for example but not limited to a 600 or 800 mg/kg dosage, the dose can be decreased by an amount ranging from 50 mg/kg and 100 mg/kg to mitigate observed undesired effects on systemic blood pressure.

In a representative embodiment doses can be administered to a subject, prior to exposure to an environmental stimulus (e.g. one dose 30 minutes before initiation of a cardiac surgery such as cardiopulmonary bypass and/or up to 1, 2, 3, 4, 5, 6 or more dosages over a perioperative period, such as every 12 hours over a period of time prior to surgery) after exposure to an environmental stimulus (e.g. upon arrival to a postoperative care setting, and/or up to 1, 2, 3, 4, 5, 6 or more dosages over a postoperative period, such as every 12 hours over a period of time after surgery).

It will be understood, however, that the specific dose level for any particular subject will depend upon a variety of factors including the age, body weight, general health, sex, diet, time of administration, route of administration, rate of excretion, drug combination and the severity of the particular disease undergoing therapy.

F. Pharmaceutical Compositions

In a preferred embodiment, the present invention provides pharmaceutical compositions comprising a polypeptide or polynucleotide of the present invention and a physiologically acceptable carrier. More preferably, a pharmaceutical composition comprises a polynucleotide that encodes a biologically active CPSI polypeptide. Alternatively, provided pharmaceutical compositions comprise citrulline or arginine in dosages as described above.

A composition of the present invention is typically administered orally or parenterally in dosage unit formulations containing standard, well-known nontoxic physiologically acceptable carriers, adjuvants, and vehicles as desired. The term "parenteral" as used herein includes intravenous, intramuscular, intra-arterial injection, or infusion techniques.

Injectable preparations, for example sterile injectable aqueous or oleaginous suspensions, are formulated according to the known art using suitable dispersing or wetting agents and suspending agents. The sterile injectable preparation can also be a sterile injectable solution or suspension in a nontoxic parenterally acceptable diluent or solvent, for example, as a solution in 1,3-butanediol.

Among the acceptable vehicles and solvents that may be employed are water, Ringer's solution, and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose any bland fixed oil can be employed including synthetic mono- or di-glycerides. In addition, fatty acids such as oleic acid find use in the preparation of injectables.

Preferred carriers include neutral saline solutions buffered with phosphate, lactate, Tris, and the like. Of course, in the case of a pharmaceutical composition provided for use in gene therapy, one purifies the vector sufficiently to render it essentially free of undesirable contaminants, such as defective interfering adenovirus particles or endotoxins and other pyrogens such that it does not cause any untoward reactions in the individual receiving the vector construct. A preferred means of purifying the vector involves the use of buoyant density gradients, such as cesium chloride gradient centrifugation.

A transfected cell can also serve as a carrier. By way of example, a liver cell can be removed from an organism, transfected with a polynucleotide of the present invention using methods set forth above and then the transfected cell returned to the organism (e.g. injected intra-vascularly).

G. Generation of Antibodies

In still another embodiment, the present invention provides an antibody immunoreactive with a polypeptide or polynucleotide of the present invention. Preferably, an antibody of the invention is a monoclonal antibody. Means for preparing and characterizing antibodies are well known in the art (See, e.g., *Antibodies A Laboratory Manual*, E. Howell and D. Lane, Cold Spring Harbor Laboratory, 1988). More preferred antibodies distinguish between the different forms of CPSI which comprise the CPSI polymorphism.

Briefly, a polyclonal antibody is prepared by immunizing an animal with an immunogen comprising a polypeptide or polynucleotide of the present invention, and collecting antisera from that immunized animal. A wide range of animal species can be used for the production of antisera. Typically an animal used for production of anti-antisera is a rabbit, a mouse, a rat, a hamster or a guinea pig. Because of the relatively large blood volume of rabbits, a rabbit is a preferred choice for production of polyclonal antibodies.

As is well known in the art, a given polypeptide or polynucleotide may vary in its immunogenicity. It is often necessary therefore to couple the immunogen (e.g., a polypeptide or polynucleotide) of the present invention) with a carrier. Exemplary and preferred carriers are keyhole limpet hemocyanin (KLH) and bovine serum albumin (BSA). Other albumins such as ovalbumin, mouse serum albumin or rabbit serum albumin can also be used as carriers.

Means for conjugating a polypeptide or a polynucleotide to a carrier protein are well known in the art and include glutaraldehyde, m-maleimidobenzoyl-N-hydroxysuccinimide ester, carbodiimide and bis-biazotized benzidine.

As is also well known in the art, immunogenicity to a particular immunogen can be enhanced by the use of non-specific stimulators of the immune response known as adjuvants. Exemplary and preferred adjuvants include complete Freund's adjuvant, incomplete Freund's adjuvants and aluminum hydroxide adjuvant.

The amount of immunogen used of the production of polyclonal antibodies varies, inter alia, upon the nature of the immunogen as well as the animal used for immunization. A variety of routes can be used to administer the immunogen, e.g. subcutaneous, intramuscular, intradermal, intravenous and intraperitoneal. The production of polyclonal antibodies is monitored by sampling blood of the immunized animal at various points following immunization. When a desired level of immunogenicity is obtained, the immunized animal can be bled and the serum isolated and stored.

In another aspect, the present invention provides a method of producing an antibody immunoreactive with a CPSI polypeptide, the method comprising the steps of (a) transfecting recombinant host cells with a polynucleotide that encodes

that polypeptide; (b) culturing the host cells under conditions sufficient for expression of the polypeptide; (c) recovering the polypeptide; and (d) preparing antibodies to the polypeptide. Preferably, the CPSI polypeptide is capable of mediating the first step of the urea cycle, cross-reacting with anti-CPSI antibody, or other biological activity in accordance with the present invention. Even more preferably, the present invention provides antibodies prepared according to the method described above.

A monoclonal antibody of the present invention can be readily prepared through use of well-known techniques such as those exemplified in U.S. Pat. No 4,196,265, herein incorporated by reference. Typically, a technique involves first immunizing a suitable animal with a selected antigen (e.g., a polypeptide or polynucleotide of the present invention) in a manner sufficient to provide an immune response. Rodents such as mice and rats are preferred animals. Spleen cells from the immunized animal are then fused with cells of an immortal myeloma cell. Where the immunized animal is a mouse, a preferred myeloma cell is a murine NS-1 myeloma cell.

The fused spleen/myeloma cells are cultured in a selective medium to select fused spleen/myeloma cells from the parental cells. Fused cells are separated from the mixture of non-fused parental cells, for example, by the addition of agents that block the de novo synthesis of nucleotides in the tissue culture media. Exemplary and preferred agents are aminopterin, methotrexate, and azaserine. Aminopterin and methotrexate block de novo synthesis of both purines and pyrimidines, whereas azaserine blocks only purine synthesis. Where aminopterin or methotrexate is used, the media is supplemented with hypoxanthine and thymidine as a source of nucleotides. Where azaserine is used, the media is supplemented with hypoxanthine.

This culturing provides a population of hybridomas from which specific hybridomas are selected. Typically, selection of hybridomas is performed by culturing the cells by single-clone dilution in microtiter plates, followed by testing the individual clonal supernatants for reactivity with an antigen-polypeptides. The selected clones can then be propagated indefinitely to provide the monoclonal antibody.

By way of specific example, to produce an antibody of the present invention, mice are injected intraperitoneally with between about 1-200 μg of an antigen comprising a polypeptide of the present invention. B lymphocyte cells are stimulated to grow by injecting the antigen in association with an adjuvant such as complete Freund's adjuvant (a non-specific stimulator of the immune response containing killed *Mycobacterium tuberculosis*). At some time (e.g., at least two weeks) after the first injection, mice are boosted by injection with a second dose of the antigen mixed with incomplete Freund's adjuvant.

A few weeks after the second injection, mice are tail bled and the sera titered by immunoprecipitation against radiolabeled antigen. Preferably, the process of boosting and titering is repeated until a suitable titer is achieved. The spleen of the mouse with the highest titer is removed and the spleen lymphocytes are obtained by homogenizing the spleen with a syringe. Typically, a spleen from an immunized mouse contains approximately 5×10^7 to 2×10^8 lymphocytes.

Mutant lymphocyte cells known as myeloma cells are obtained from laboratory animals in which such cells have been induced to grow by a variety of well-known methods. Myeloma cells lack the salvage pathway of nucleotide biosynthesis. Because myeloma cells are tumor cells, they can be propagated indefinitely in tissue culture, and are thus denomi-

nated immortal. Numerous cultured cell lines of myeloma cells from mice and rats, such as murine NS-1 myeloma cells, have been established.

Myeloma cells are combined under conditions appropriate to foster fusion with the normal antibody-producing cells from the spleen of the mouse or rat injected with the antigen/polypeptide of the present invention. Fusion conditions include, for example, the presence of polyethylene glycol. The resulting fused cells are hybridoma cells. Like myeloma cells, hybridoma cells grow indefinitely in culture.

Hybridoma cells are separated from unfused myeloma cells by culturing in a selection medium such as HAT media (hypoxanthine, aminopterin, thymidine). Unfused myeloma cells lack the enzymes necessary to synthesize nucleotides from the salvage pathway because they are killed in the presence of aminopterin, methotrexate, or azaserine. Unfused lymphocytes also do not continue to grow in tissue culture. Thus, only cells that have successfully fused (hybridoma cells) can grow in the selection media.

Each of the surviving hybridoma cells produces a single antibody. These cells are then screened for the production of the specific antibody immunoreactive with an antigen/polypeptide of the present invention. Single cell hybridomas are isolated by limiting dilutions of the hybridomas. The hybridomas are serially diluted many times and, after the dilutions are allowed to grow, the supernatant is tested for the presence of the monoclonal antibody. The clones producing that antibody are then cultured in large amounts to produce an antibody of the present invention in convenient quantity.

By use of a monoclonal antibody of the present invention, specific polypeptides and polynucleotide of the invention can be recognized as antigens, and thus identified. Once identified, those polypeptides and polynucleotide can be isolated and purified by techniques such as antibody-affinity chromatography. In antibody-affinity chromatography, a monoclonal antibody is bound to a solid substrate and exposed to a solution containing the desired antigen. The antigen is removed from the solution through an immunospecific reaction with the bound antibody. The polypeptide or polynucleotide is then easily removed from the substrate and purified.

H. Detecting a Polynucleotide or a Polypeptide of the Present Invention

Alternatively, the present invention provides a method of detecting a polypeptide of the present invention, wherein the method comprises immunoreacting the polypeptides with antibodies prepared according to the methods described above to form antibody-polypeptide conjugates, and detecting the conjugates.

In yet another embodiment, the present invention provides a method of detecting messenger RNA transcripts that encode a polypeptide of the present invention, wherein the method comprises hybridizing the messenger RNA transcripts with polynucleotide sequences that encode the polypeptide to form duplexes; and detecting the duplex. Alternatively, the present invention provides a method of detecting DNA molecules that encode a polypeptide of the present invention, wherein the method comprises hybridizing DNA molecules with a polynucleotide that encodes that polypeptide to form duplexes; and detecting the duplexes.

The detection and screening assays disclosed herein can be used as a prognosis tool. Human CPSI-encoding polynucleotides as well as their protein products can be readily used in clinical setting as a prognostic indicator for screening for susceptibility to hyperammonemia and to other heritable CPSI-related diseases in humans.

The detection and screening assays disclosed herein can be also used as a part of a diagnostic method. Human CPSI-

encoding polynucleotides as well as their protein products can be readily used in clinical setting to diagnose susceptibility to hyperammonemia and to other heritable CPSI-related diseases in humans.

H.1. Screening Assays for a Polypeptide of the Present Invention

The present invention provides a method of screening a biological sample for the presence of a CPSI polypeptide. Preferably, the CPSI polypeptide possesses activity in the urea cycle, cross-reactivity with an anti-CPSI antibody, or other biological activity in accordance with the present invention. A biological sample to be screened can be a biological fluid such as extracellular or intracellular fluid or a cell or tissue extract or homogenate. A biological sample can also be an isolated cell (e.g., in culture) or a collection of cells such as in a tissue sample or histology sample. A tissue sample can be suspended in a liquid medium or fixed onto a solid support such as a microscope slide. Hepatic tissues comprise particularly contemplated tissues.

Preferably, antibodies which distinguish between the N1405 CPSI polypeptide and the T1405 CPSI polypeptide are provided. Such antibodies may compare polyclonal antibodies but are preferably monoclonal antibodies prepared as described hereinabove.

In accordance with a screening assay method, a biological sample is exposed to an antibody immunoreactive with the polypeptide whose presence is being assayed. Typically, exposure is accomplished by forming an admixture in a liquid medium that contains both the antibody and the candidate polypeptide. Either the antibody or the sample with the polypeptide can be affixed to a solid support (e.g., a column or a microtiter plate).

The biological sample is exposed to the antibody under biological reaction conditions and for a period of time sufficient for antibody-polypeptide conjugate formation. Biological reaction conditions include ionic composition and concentration, temperature, pH and the like.

Ionic composition and concentration can range from that of distilled water to a 2 molal solution of NaCl. Preferably, osmolality is from about 100 mosmols/l to about 400 mosmols/l and, more preferably from about 200 mosmols/l to about 300 mosmols/l. Temperature preferably is from about 4° C. to about 100° C., more preferably from about 15° C. to about 50° C. and, even more preferably from about 25° C. to about 40° C. pH is preferably from about a value of 4.0 to a value of about 9.0, more preferably from about a value of 6.5 to a value of about 8.5 and, even more preferably from about a value of 7.0 to a value of about 7.5. The only limit on biological reaction conditions is that the conditions selected allow for antibody-polypeptide conjugate formation and that the conditions do not adversely affect either the antibody or the polypeptide.

Exposure time will vary inter alia with the biological conditions used, the concentration of antibody and polypeptide and the nature of the sample (e.g., fluid or tissue sample). Means for determining exposure time are well known to one of ordinary skill in the art. Typically, where the sample is fluid and the concentration of polypeptide in that sample is about 10^{-10} M, exposure time is from about 10 minutes to about 200 minutes.

The presence of polypeptide in the sample is detected by detecting the formation and presence of antibody-polypeptide conjugates. Means for detecting such antibody-antigen (e.g., receptor polypeptide) conjugates or complexes are well known in the art and include such procedures as centrifugation, affinity chromatography and the like, binding of a secondary antibody to the antibody-candidate receptor complex.

In one embodiment, detection is accomplished by detecting an indicator affixed to the antibody. Exemplary and well known such indicators include radioactive labels (e.g., ^{32}P , ^{125}I , ^{14}C), a second antibody or an enzyme such as horse radish peroxidase. Means for affixing indicators to antibodies are well known in the art. Commercial kits are available.

H.2. Screening Assay for Anti-Polypeptide Antibody

In another aspect, the present invention provides a method of screening a biological sample for the presence of antibodies immunoreactive with a CPSI polypeptide. Preferably the CPSI polypeptide has activity in the urea cycle, cross-reactivity with an anti-CPSI antibody, or other biological activity in accordance with the present invention. In accordance with such a method, a biological sample is exposed to a CPSI polypeptide under biological conditions and for a period of time sufficient for antibody-polypeptide conjugate formation and the formed conjugates are detected.

H.3. Screening Assay for Polynucleotide that Encodes a CPSI Polypeptide of the Present Invention

A nucleic acid molecule and, particularly a probe molecule, can be used for hybridizing as an oligonucleotide probe to a nucleic acid source suspected of encoding a CPSI polypeptide of the present invention. Optimally, the CPSI polypeptide has activity in the urea cycle, cross-reactivity with an anti-CPSI antibody, or other biological activity in accordance with the present invention. The probing is usually accomplished by hybridizing the oligonucleotide to a DNA source suspected of possessing a CPSI gene. In some cases, the probes constitute only a single probe, and in others, the probes constitute a collection of probes based on a certain amino acid sequence or sequences of the polypeptide and account in their diversity for the redundancy inherent in the genetic code.

A suitable source of DNA for probing in this manner is capable of expressing a polypeptide of the present invention and can be a genomic library of a cell line of interest. Alternatively, a source of DNA can include total DNA from the cell line of interest. Once the hybridization method of the invention has identified a candidate DNA segment, one confirms that a positive clone has been obtained by further hybridization, restriction enzyme mapping, sequencing and/or expression and testing.

Alternatively, such DNA molecules can be used in a number of techniques including their use as: (1) diagnostic tools to detect normal and abnormal DNA sequences in DNA derived from subject's cells, such as a CPSI polymorphism described herein; (2) means for detecting and isolating other members of the polypeptide family and related polypeptides from a DNA library potentially containing such sequences; (3) primers for hybridizing to related sequences for the purpose of amplifying those sequences; (4) primers for altering native CPSI DNA sequences; as well as other techniques which rely on the similarity of the DNA sequences to those of the DNA segments herein disclosed.

As set forth above, in certain aspects, DNA sequence information provided by the invention allows for the preparation of relatively short DNA (or RNA) sequences (e.g., probes) that specifically hybridize to encoding sequences of a selected CPSI gene. In these aspects, nucleic acid probes of an appropriate length are prepared based on a consideration of the encoding sequence for a polypeptide of this invention. The ability of such nucleic acid probes to specifically hybridize to other encoding sequences lend them particular utility in a variety of embodiments. Most importantly, the probes can be used in a variety of assays for detecting the presence of complementary sequences in a given sample. However, other uses are envisioned, including the use of the sequence infor-

mation for the preparation of mutant species primers, or primers for use in preparing other genetic constructions.

To provide certain of the advantages in accordance with the invention, a preferred nucleic acid sequence employed for hybridization studies or assays includes probe sequences that are complementary to at least a 14 to 40 or so long nucleotide stretch of a nucleic acid sequence of the present invention, such as a sequence shown in any of SEQ ID NOs:1, 3, 11 and 13. A size of at least 14 nucleotides in length helps to ensure that the fragment is of sufficient length to form a duplex molecule that is both stable and selective. Molecules having complementary sequences over stretches greater than 14 bases in length are generally preferred, though, to increase stability and selectivity of the hybrid, and thereby improve the quality and degree of specific hybrid molecules obtained. One will generally prefer to design nucleic acid molecules having gene-complementary stretches of 14 to 20 nucleotides, or even longer where desired. Such fragments can be readily prepared by, for example, directly synthesizing the fragment by chemical means, by application of nucleic acid reproduction technology, such as the PCR technology of U.S. Pat. No. 4,683,202, herein incorporated by reference, or by introducing selected sequences into recombinant vectors for recombinant production.

Accordingly, a nucleotide sequence of the present invention can be used for its ability to selectively form duplex molecules with complementary stretches of the gene. Depending on the application envisioned, one employs varying conditions of hybridization to achieve varying degrees of selectivity of the probe toward the target sequence. For applications requiring a high degree of selectivity, one typically employs relatively stringent conditions to form the hybrids. For example, one selects relatively low salt and/or high temperature conditions, such as provided by 0.02M-0.15M salt at temperatures of about 50° C. to about 70° C. including particularly temperatures of about 55° C., about 60° C. and about 65° C. Such conditions are particularly selective, and tolerate little, if any, mismatch between the probe and the template or target strand.

Of course, for some applications, for example, where one desires to prepare mutants employing a mutant primer strand hybridized to an underlying template or where one seeks to isolate polypeptide coding sequences from related species, functional equivalents, or the like, less stringent hybridization conditions are typically needed to allow formation of the heteroduplex. Under such circumstances, one employs conditions such as 0.15M-0.9M salt, at temperatures ranging from about 20° C. to about 55° C., including particularly temperatures of about 25° C., about 37° C., about 45° C., and about 50° C. Cross-hybridizing species can thereby be readily identified as positively hybridizing signals with respect to control hybridizations. In any case, it is generally appreciated that conditions can be rendered more stringent by the addition of increasing amounts of formamide, which serves to destabilize the hybrid duplex in the same manner as increased temperature. Thus, hybridization conditions can be readily manipulated, and thus will generally be a method of choice depending on the desired results.

In certain embodiments, it is advantageous to employ a nucleic acid sequence of the present invention in combination with an appropriate means, such as a label, for determining hybridization. A wide variety of appropriate indicator means are known in the art, including radioactive, enzymatic or other ligands, such as avidin/biotin, which are capable of giving a detectable signal. In preferred embodiments, one likely employs an enzyme tag such as urease, alkaline phosphatase or peroxidase, instead of radioactive or other environmentally

undesirable reagents. In the case of enzyme tags, calorimetric indicator substrates are known which can be employed to provide a means visible to the human eye or spectrophotometrically, to identify specific hybridization with complementary nucleic acid-containing samples.

In general, it is envisioned that the hybridization probes described herein are useful both as reagents in solution hybridization as well as in embodiments employing a solid phase. In embodiments involving a solid phase, the sample containing test DNA (or RNA) is adsorbed or otherwise affixed to a selected matrix or surface. This fixed, single-stranded nucleic acid is then subjected to specific hybridization with selected probes under desired conditions. The selected conditions depend inter alia on the particular circumstances based on the particular criteria required (depending, for example, on the G+C contents, type of target nucleic acid, source of nucleic acid, size of hybridization probe, etc.). Following washing of the hybridized surface so as to remove nonspecifically bound probe molecules, specific hybridization is detected, or even quantified, by means of the label.

H.4. Assay Kits

In another aspect, the present invention provides a diagnostic assay kit for detecting the presence of a polypeptide of the present invention in biological samples, where the kit comprises a first container containing a first antibody capable of immunoreacting with the polypeptide, with the first antibody present in an amount sufficient to perform at least one assay. Preferably, the assay kits of the invention further comprise a second container containing a second antibody that immunoreacts with the first antibody. More preferably, the antibodies used in the assay kits of the present invention are monoclonal antibodies. Even more preferably, the first antibody is affixed to a solid support. More preferably still, the first and second antibodies comprise an indicator, and, preferably, the indicator is a radioactive label or an enzyme.

The present invention also provides a diagnostic kit for screening agents. Such a kit can contain a polypeptide of the present invention. The kit can contain reagents for detecting an interaction between an agent and a receptor of the present invention. The provided reagent can be radiolabeled. The kit can contain a known radiolabeled agent capable of binding or interacting with a receptor of the present invention.

In an alternative aspect, the present invention provides diagnostic assay kits for detecting the presence, in biological samples, of a polynucleotide that encodes a polypeptide of the present invention, the kits comprising a first container that contains a second polynucleotide identical or complementary to a segment of at least 10 contiguous nucleotide bases of, as a preferred example, in any of SEQ ID NOs:1, 3, 11 and 13.

In another embodiment, the present invention provides diagnostic assay kits for detecting the presence, in a biological sample, of antibodies immunoreactive with a polypeptide of the present invention, the kits comprising a first container containing a CPSI polypeptide, that immunoreacts with the antibodies, with the polypeptide present in an amount sufficient to perform at least one assay. Preferably, the CPSI polypeptide has activity in the urea cycle, cross-reactivity on an anti-CPSI antibody, or other biological activity in accordance with the present invention. The reagents of the kit can be provided as a liquid solution, attached to a solid support or as a dried powder. Preferably, when the reagent is provided in a liquid solution, the liquid solution is an aqueous solution. Preferably, when the reagent provided is attached to a solid support, the solid support can be chromatograph media or a microscope slide. When the reagent provided is a dry powder, the powder can be reconstituted by the addition of a suitable solvent. The solvent can be provided.

The following Examples have been included to illustrate preferred modes of the invention. Certain aspects of the following Examples are described in terms of techniques or procedures found or contemplated by the present inventors to work well in the practice of the invention. These Examples are exemplified through the use of standard laboratory practices of the inventors. In light of the present disclosure and the general level of skill in the art, those of skill will appreciate that the following Examples are intended to be exemplary only in that numerous changes, modification, and alterations can be employed without departing from the spirit and scope of the invention.

Materials and Methods Used in Examples 1-3

The following materials and methods are employed in each of Examples 1-3. Additional materials and methods are also described in each Example.

Clinical/Patient Recruitment: More than 200 patients undergoing BMT at Vanderbilt University Medical Center, Nashville, Tenn., have been enrolled in the BMT-Lung Injury Following Engraftment (LIFE) Study aimed at understanding mechanisms of acute lung injury and multiple organ failure after transplant. Consent was sought from consecutive patients undergoing BMT or PBSCT for treatment of malignancy. Definitions of organ failure (including HVOD) and reversal were prospectively defined and data was collected concurrently during hospitalization. Plasma, cell pellets, and urine were collected at study enrollment (before receiving chemotherapy) and on the day of transplantation (before marrow infusion) after completing ablative chemo-radiotherapy.

Amino Acid Analysis—Blood and urine were immediately centrifuged after collection. All samples were kept on ice, then stored at -70° C. until analyzed. Under these storage conditions, glutamine, cysteine and homocysteine are known to decrease, so these were not used in the analysis. Plasma amino acids were measured in the Vanderbilt Diagnostic Laboratories, Vanderbilt University, Nashville, Tenn. Briefly, a protein free extract of plasma was prepared by protein precipitation with sulfosalicylic acid and filtration through a $0.45\ \mu\text{m}$ ACRODISC™ 4 filter (Gelman Sciences, Ann Arbor, Mich.). Amino acids were separated by cation exchange chromatography using a four-component pH- and ionic strength-graded lithium citrate buffer system on a Beckmann 7300 amino acid analyzer (Beckmann, Palo Alto, Calif.). Post column derivatization of amino acids with ninhydrin allowed detection of primary amine amino acids at 570 nm, and secondary amines at 440 nm. Quantification was achieved by instrument calibration with standard reference materials (Sigma, St. Louis, Mo.).

Statistics. Plasma amino acid values were expressed as mean \pm SEM. Comparisons between baseline and post-chemotherapy amino acid values were made using Student's t-Test. Allelic frequency was compared between patients with and without HVOD using Chi square analysis.

Patients. Patients were identified from those enrolled in the BMT Lift Study at Vanderbilt University. DNA was isolated from pre-transplant blood or spun urine samples. HVOD status was determined using the Baltimore criteria:

- Bilirubin >2.0 mg/dl
- Hepatomegaly
- 2% sudden weight gain

Genotyping. DNA was isolated using a QIAmpJ blood kit (Qiagen). The T1405N polymorphism changes the DNA sequence as follows:

CCT-GCC-ACC-CCA-GTG	Normal
CCT-GCC-AAC-CCA-GTG	Change

The C to A transversion replaces the pyrimidine C with the purine A which destroys a MspI site. The use of a primer from within the 35th intron of CPSI and an exotic primer from exon 36 of the CPSI gene reliably PCR amplifies a 387 bp fragment encompassing the region containing the change. This combination gives a robust amplification. PCR Ready-to-Go™ beads are also used in amplification (Pharmacia).

The polymorphism was detected using a non-denaturing gel to take advantage of the secondary structures created by the C to A transversion. This change creates enough secondary structure to prevent reliable digestion by restriction enzymes (MspI) to detect the polymorphism. This change also interferes with direct sequence analysis unless ITP is substituted for GTP in the reaction. Non-denaturing gels take advantage of the secondary structures created by this change. Fifteen (15) individuals were compared by this method and sequence analysis.

To detect the DNA fragments in the gel, a silver staining technique was adapted. This inexpensive rapid method allowed visualization of bands shortly after electrophoresis.

Statistical Analysis. A sufficient sample size was obtained to perform Chi Square analysis on the results. The Hardy-Weinberg equation was used to calculate the expected frequencies for the genotypes ($p^2+2pq+q^2$). P values were obtained from a standard Chi Square table using 2 degrees of freedom.

Example 1

Alleles of CPSI Exonic Polymorphism (T1405N) are not in Hardy-Weinberg Equilibrium with the Presence or Absence of HVOD

In accordance with the present invention, a common polymorphism near the 3' end of the CPSI mRNA (about 0.44 heterozygosity) has been identified. Sequence analysis of this change revealed a C to A transversion at base 4340 changing the triplet code from ACC to AAC. This results in a substitution of asparagine for threonine at amino acid 1405 (referred to herein as "T1405N"). The threonine is within the allosteric domain, preceding the signature sequence PV(A/S)WP(T/S)(A/Q)E, a sequence that is important in the binding of the cofactor n-acetyl-glutamate (NAG).

In all known CPSIs activated by NAG, a threonine residue is among the two residues that precede the signature sequence. (Rubio, *Biochemical Society Transactions* 21:198-202 (1998)). On the basis of structure-function studies, hydrogen bond formation with the carbonyl oxygen of the acetamido group of NAG is felt to play a role in the binding of this activator. (Stapleton et al., *Biochemistry* 35:14352-14361 (1996); Javid-Majd et al., *Biochemistry* 35:14362-14369 (1996)). The substitution of the threonine side chain by asparagine is envisioned to alter the hydrogen bond formation with NAG and results in a qualitative change in CPSI enzymatic function and in sensitivity to the available pool of NAG. Although applicants do not wish to be bound by any particular theory of operation, it is speculated that based on the precedent of the effects of other xenobiotics, that limited availabil-

ity of NAG after escalated dose chemotherapy is one of the mechanisms promoting urea cycle dysfunction.

126 individuals were genotyped from the BMT Life Study group. 30 individuals manifested evidence of HVOD in this group (24%). 70 patients were genotyped from blood samples and 56 from urine cell pellets. Samples from 15 patients were reamplified via PCR and sequenced to confirm the consistency of the results.

Tables 2 and 3 show the results of genotype analysis for the T1405N polymorphism between HVOD+ and HVOD- patients. The C allele, also referred to herein as the CPSIa allele or the threonine encoding allele, has a frequency of 0.62 in the examined population and the A allele, also referred to herein as the CPSIb allele or the asparagine encoding allele, has a frequency of 0.38. The Chi Square value for the table is 4.3 (P=0.1) indicating that the polymorphism is probably not in Hardy-Weinburg equilibrium with the presence of HVOD. Thus, these results provide evidence for disequilibrium in the distribution of the T1405N alleles in BMT patients with HVOD, indicating that the polymorphism can be used to identify subjects who are susceptible to BMT toxicity.

TABLE 2

Genotype	HVOD+	HVOD-
CC	13 (expected 11.4)	32 (expected 36.5)
AC	16 (expected 14.1)	50 (expected 45.1)
AA	1 (expected 4.5)	14 (expected 14.4)

TABLE 3

Total alleles:	Expected Frequencies:
A: 96	AA: 0.15
C: 62	AC: 0.47
	CC: 0.38

Additional data gathered from a study of approximately 200 patients provided additional statistical evidence supporting the use of the polymorphism in detection of susceptibility to sub-optimal urea cycle function. This data was subjected to the statistical methods described above.

Bone marrow transplant toxicity results in significant morbidity and mortality. HVOD is associated with a poor prognosis in BMT patients. This study was undertaken to assess an association between the CPSI enzyme and the occurrence of HVOD. The T1405N polymorphism affects CPSI function. Its wide distribution in the population suggests that both forms provide adequate urea cycle function under normal conditions. The addition of metabolic stressors (such as high-dose chemotherapy) serves to lower CPSI efficiency below an effective threshold. Analysis of the data thus suggests that HVOD is more likely to occur in patients with the threonine encoding allele than those with the asparagine. The threonine encoding allele is shared by the rodent form of CPSI.

Example 2

Biochemical and Genetic Alterations in Carbamyl Phosphate Synthetase I in Patients with Post-Bone Marrow Transplant Complications

Bone marrow transplantation (BMT) and peripheral blood stem cell transplants (PBSCT) are increasingly being used as primary therapy for selected malignancies. Use of stem cell support for hematopoietic reconstitution allows for substan-

tial escalation in the dose of chemotherapy in an attempt to eradicate potentially lethal cancers. With improvements in prophylaxis for infection and prevention of disabling graft-versus-host disease, chemotherapy-induced organ dysfunction remains a significant barrier to more widespread use of this treatment.

Hepatic venoocclusive disease (HVOD), a clinical syndrome of hyperbilirubinemia (serum bilirubin >2.0 mg/dL), hepatomegaly, and fluid retention early after BMT, is a major dose-limiting toxicity after BMT, afflicting up to 54% of patients. Many patients developing HVOD after BMT will also meet the criteria for acute lung injury (ALI). Nearly half of patients with severe HVOD require mechanical ventilation, with an attendant mortality in excess of 90%. Such data underscore the large impact on mortality of sequential organ dysfunction, even in a young patient population, and reinforce the clinically important association of poor prognosis after acute lung injury in patients with hepatic dysfunction. The mechanisms responsible for this organ interaction remain incompletely understood.

In this Example, whether conditioning chemotherapy administered prior to BMT might affect early enzymes in the UC and secondarily predispose patients for hepatic dysfunction and multiple organ failure was analyzed. The plasma amino acid analyses supported the notions of both impaired UC function and decreased production of nitric oxide (NO_x). In light of these findings, patients were screened for the exonic single nucleotide polymorphism (SNP) in CPS-I disclosed herein. It was found that homozygosity for the SNP was associated with a decreased incidence of HVOD and enhanced early survival after BMT, consistent with a significant pharmacogenetic interaction.

Methods

Clinical/Patient Recruitment: Over the last three years 200 patients undergoing BMT at Vanderbilt University Medical Center have been sequentially enrolled in the Bone Marrow Transplant-Lung Injury Following Engraftment (BMT-LIFE) Study, a coordinated clinical-biochemical exploratory investigation aimed at understanding mechanisms of acute lung injury and multiple organ failure after transplant. Definitions of organ failure and reversal were prospectively defined and data was collected concurrently during hospitalization and until 60 days after BMT. Exclusion criteria included active viral and prior escalated dose therapy with hematopoietic stem cell support (either PBSCT or BMT).

Hepatic venoocclusive disease (HVOD) was identified in patients with bilirubin >2 mg/dL before 21 days after transplant with either weight gain >5% of baseline or new onset of tender hepatomegaly. Acute lung injury (ALI) was defined as bilateral infiltrates on chest roentgenogram for three consecutive dates with a ratio of partial pressure of oxygen in arterial blood to the fraction of inspired oxygen concentration (PaO₂/FiO₂) of less than 300 in the absence of clinical cardiac dysfunction. Patients alive 60 days after transplant were defined as survivors. Plasma, circulating cell pellets, and urine were collected at study enrollment (before receiving chemotherapy) and on the day of BMT, several days after completing high dose chemotherapy but before marrow infusion. Samples were aliquotted, and immediately placed on ice prior to storage at -80° C. before analysis.

Amino Acid Analysis. Amino acid analysis was performed on cryopreserved plasma samples from days -8 and 0 (pre-treatment and day of transplantation) in 60 patients. Patient samples were initially randomly selected for pilot studies; subsequently analyzed samples were specifically enriched to

include extra patients with the SNP AA genotype of CPS-I (see below) and additional patients with the post-BMT complications of HVOD and ALI. A protein free extract of plasma was prepared by protein precipitation with sulfosalicylic acid and filtration through a 0.45 μm Acrodisc 4 (Gelman Sciences, Ann Arbor, Mich.).

Amino acids were separated by cation exchange chromatography using a four-component pH- and ionic strength-graded lithium citrate buffer system on a Beckmann 7300 amino acid analyzer (Beckmann, Palo Alto, Calif.). Post column derivatization of amino acids with ninhydrin allowed detection of primary amine amino acids at 570 nm, and secondary amines at 440 nm. Quantitation was achieved by instrument calibration with standard reference materials (Sigma, St. Louis, Mo.). Citrulline, arginine, and ornithine were examined as measurable indices of flux of intermediates through the urea cycle.

Measurement of plasma nitric oxide metabolites (NO_x) Plasma NO_x was measured in a subgroup of patients using modified Griess reagents after samples were deproteinated and incubated with cadmium beads to convert nitrate to nitrite.

Detection of T1405N polymorphism. Oligonucleotide primers from within the 36th exon (CGGAAGCCACATCA-GACTGG (SEQ ID NO:15) and intron (GGAGAGT-GAACTTGACAATCATC (SEQ ID NO:16)) of CPS1 and the polymerase chain reaction (PCR) to reliably amplify a 251 bp fragment encompassing the region containing the change from genomic DNA obtained from buffy coat preparations or urinary sediment. This combination of primers gave reproducible amplification using PCR Ready-to-Go beads (Pharmacia) and PCR cycle conditions as follows: 35 cycles of 1 minute anneal at 55°C., 1 minute extension at 72°C., and 1 minute denaturation at 94°C.

After formamide treatment, samples were subjected to electrophoresis for 4 hours at 4°C. in a non-denaturing MDE™ gel (FMC, Rockland, Me.), then stained with silver nitrate to detect DNA fragments. Confirmatory genotyping of 17 individuals using both non-denaturing gel electrophoresis and direct sequence analysis yielded identical results. Patients were classified as having homozygous SNP genotypes of CC or AA, or as being heterozygous (AC). For comparison, using identical methods, a cohort of 100 patients with Alzheimer's disease was analyzed to assess the distribution of CPS1 SNP genotypes.

Statistical Analysis. Plasma amino acid levels before and after chemotherapy, and levels between groups of patients, were compared using Student's T-test or Wilcoxon's Rank Sum Test (if the data were not normally distributed). Distribution of genotypes of CPS1 was compared across groups by calculating allelic frequency for the entire group and searching for evidence of Hardy-Weinberg disequilibrium in specifically selected subgroups using P^2 analysis. Sensitivity, specificity, predictive values, and relative risk assessments were generated from two-by-two contingency tables constructed using specific amino acid values in groups of patients divided by presence and absence of specific clinical outcomes (e.g. HVOD, ALI, and death).

Results

Two hundred patients were enrolled in the BMT-LIFE Study. 52% underwent autologous transplant (mean age 46±1 years); 48% received allogeneic grafts (mean age 40±1 years). Of the patients undergoing allogeneic transplants, 24% received grafts from HLA-matched unrelated donors. Nearly two-thirds of the patients in the autologous group were

women, reflecting the increased prevalence of breast cancer in this population. The indications for transplant were diverse, but 79% of the patients were transplanted for breast cancer, leukemia, or non-Hodgkin's lymphoma. The different preparative regimens used prior to BMT included CTC (cyclophosphamide, thiotepa, carboplatin), BuCy (busulfan, cyclophosphamide), CVP16TBI (cyclophosphamide, etoposide, total body irradiation), CBVP16 (cyclophosphamide, bis-chloroethylnitrosourea, etoposide) and TC (thiotepa, cyclophosphamide).

Both morbidity and mortality are not uncommon after BMT. While the overall 60 day mortality in the study was 14%, it was 20% in patients receiving allografts. Complications of acute lung injury (ALI) and hepatic venoocclusive disease (HVOD) each occurred in 19% of the patients. These complications were more than twice as common in patients receiving allografts. In the group of patients developing HVOD, 62% (24/38) also met criteria for ALI during hospitalization. Only 38% (14/38) of the cases of ALI occurred in patients who never met criteria for HVOD.

A subset (60/200) of the patients, specifically enriched during sample selection with extra patients with CPS-I AA SNP genotype and additional patients with post-transplant complications, had plasma amino acid determinations before administration of chemotherapy and on the day of transplant. Comparison of levels of selected amino acids that participate in the UC (citrulline, ornithine, and arginine) before and after chemotherapy revealed significant differences. Citrulline levels fell in virtually all patients with a mean group decrease from 23.4±1.3 μM to 9.1±0.7 μM ($P<0.05$). Arginine levels rose by approximately 35% ($P<0.05$), and ornithine levels rose by 21% ($P<0.05$).

The ratio of ornithine/citrulline (O/C ratio), an index of flux through the early steps of the UC (i.e. lower values indicate better cycle flow), increased from 3.9±0.7 at study enrollment to 11.8±1.8 after induction chemotherapy ($P<0.05$). Shifts also occurred in amino acids that are not part of the UC. Levels of glycine and alanine, two aliphatic amino acids, fell significantly by 11% and 19%, respectively, in a pattern not consistent with decreased flux of intermediates through the cycle simply due to decreased protein intake (acute or chronic). Phenylalanine and methionine levels rose by 43% and 23%, respectively, suggesting subclinical hepatic dysfunction.

Baseline plasma levels of citrulline and the O/C ratios had prognostic importance. Sixty day survivors of BMT had higher baseline levels of citrulline than did nonsurvivors (24.4±1.3 vs 17.7±2.9 μM , respectively; $P<0.05$). The relative risk for death before 60 days after BMT was 2.92 for patients with an enrollment citrulline level less than 20. The negative predictive value for death of a plasma citrulline level greater than 20 μM was 90%. O/C ratios at enrollment were significantly lower in patients never developing either HVOD (2.8±0.2) or ALI (2.9±0.2) when compared to patients who subsequently developed these complications (5.8±1.9 and 6.5±2.7, respectively; $P<0.05$). Comparison of O/C ratios between 60 day survivors and nonsurvivors of BMT at study enrollment showed a trend toward lower values in survivors (3.3±0.2 vs. 6.9±3.9; $P=0.06$). The negative predictive value for death within 60 days after BMT associated with a baseline O/C ratio less than 2.5 was 92%.

Several urea cycle amino acid intermediate levels after preparative therapy, on the day of BMT, also had significance. Plasma arginine levels were higher in survivors (114.5±5.9 μM) when compared to nonsurvivors (92.3±10.4 μM) ($P<0.05$). O/C ratios were significantly higher, suggesting more impaired UCF, in patients who later developed ALI

when compared to those never developing severe lung dysfunction (18.4 ± 5.9 vs 9.5 ± 0.7 ; $P < 0.05$). Although the negative predictive value for development of ALI of a post-chemotherapy O/C ratio less than ten was high (86%), the relative risk for mortality associated with this threshold was only 1.44. There was a trend toward higher O/C ratios in patients on the day of BMT in patients who subsequently developed HVOD ($P = 0.09$).

Levels of nitric oxide metabolites (NO_x) in plasma were measured in 62 patients. Plasma NO_x levels fell 20% after induction therapy, from $40 \pm 2 \mu\text{M}$ at study enrollment to $32 \pm 2 \mu\text{M}$ on the day of BMT ($P < 0.05$). The median NO_x value on the day of BMT in 20 patients developing either HVOD or ALI was $28 \mu\text{M}$; for patients without such complications the plasma NO_x was $35 \mu\text{M}$. No clear differences between plasma NO_x was observed when patients with different CPSI SNP genotypes were compared.

To assess whether certain patients might have a genetic predisposition to develop morbid complications following induction therapy and BMT, all patients in the study were genotyped for a CPSI SNP. Of 200 patients, data was analyzed from 196 patients (i.e. 2 clinical exclusions; 2 unsuccessful PCR amplifications) to determine if the CPS-I C4340A SNP was in Hardy-Weinberg equilibrium with the development of HVOD. The distribution of CPSI SNP genotypes in patients undergoing BMT was identical to that of the control group (100 patients with Alzheimer's disease): 44% CC (wild type), 45% AC (heterozygous), and 11% AA (homozygous for the transversion). The attack rate of HVOD in those with the CC or AC genotype were 18% and 24%, respectively. There were no cases of HVOD in patients with the AA genotype.

Finding that this allelic distribution was not in Hardy-Weinberg equilibrium with the development of HVOD ($P^2 = 5.06$, $P < 0.05$) suggests that the SNP AA genotype alters susceptibility to hepatic toxicity following induction chemotherapy. There were also trends toward differences in mortality 60 days after BMT between the SNP genotypes. Nonsurvivors constituted 15% and 20% of the AC and CC genotype groups, respectively. Interestingly, all of the patients with the AA genotype survived 60 days after BMT ($P^2 = 3.36$; $P = 0.06$). Of note, almost all of the P^2 score came from the AA/survivor cell. There were no significant differences between patients with different SNP C4340A genotypes in the attack rate of ALI (16%, 15%, and 25% in the AA, AC, and CC groups, respectively). While ALI was associated with significant mortality in patients with either the AC or CC genotypes (71% and 66%, respectively), all patients with the AA genotype who developed ALI eventually had resolution of both bilateral pulmonary infiltrates on CXR and impaired gas exchange and survived 60 days after BMT.

Discussion

The data presented in this Example reflect a close association between HVOD and ALI in patients after BMT, with nearly two-thirds of patients with HVOD meeting criteria for ALI. In this study, 68% (26/38) of patients developing ALI required mechanical ventilation. Rubenfelt and Crawford have reported a meaningful survival, defined as extubation followed by discharge from the hospital with thirty day survival, of only 6% in patients requiring mechanical ventilation after BMT. See Rubenfeld, G. D. and Crawford, S. W., *Annals of Internal Medicine* (1996) 125:625-33.

HVOD remains the major dose limiting toxicity of escalated dose chemotherapy. It is clinically characterized by fluid retention, jaundice, ascites, and painful hepatic enlargement

occurring within 3 weeks of BMT. Autopsy studies of those non-surviving patients fulfilling these clinical criteria provide histological confirmation in >80% of cases and are consistent with the idea that enhanced local thrombosis might be an initiating event in the pathogenesis of HVOD.

The significant fall in citrulline levels and rise in plasma ornithine levels from patients undergoing BMT suggests a significant disturbance in flux of carbon intermediates through the hepatic UC in patients after induction chemotherapy. Analysis of the patterns of other amino acids argues that this effect is not simply due to decreased protein intake. In contrast to the patterns seen in patients with starvation, where levels of glycine and branched chain amino acids (BCAA) are usually significantly elevated, we observed a fall in glycine and no significant change in the BCAAs. Furthermore, starvation tends to increase activity of CPSI in liver and should not lead to increases in plasma ornithine.

The pretreatment ability of patients undergoing BMT to maintain flow of intermediates through the UC had particular prognostic importance. Sixty day nonsurvivors after BMT and those patients developing HVOD or ALI had significantly lower levels of citrulline and higher O/C ratios compared to patients who did not develop these complications. Of interest was the observation that nonsurvivors of BMT had lower plasma arginine values after induction therapy when compared to surviving patients. In light of the clustering of cells containing early UC enzymes about the terminal hepatic venules, local concentrations of both arginine and nitric oxide (NO) might be much higher and might play an important role in maintaining patency of these vessels and regulating regional hepatic blood flow. The studies showing a significant reduction in plasma NO_x levels after induction chemotherapy support the idea that NO production is altered during BMT.

The apparent discrepancy between apparently normal plasma levels of arginine on the day of transplant and markedly reduced plasma NO_x underscores the complex in vivo kinetics of arginine and citrulline flux across different organ beds. Stable isotope studies of whole body arginine homeostasis have indicated that only about 15% of plasma arginine turnover is associated with urea formation, and that only 1.2% of plasma arginine turnover is associated with NO formation. Furthermore, in vitro studies have documented substantial channeling of urea cycle intermediates, from citrulline to arginine, that is not influenced by exogenous provision of substrate. The ability of an individual patient to maintain urea cycle function and hepatic NO production during the stresses of induction chemotherapy can, in part, influence their resistance to complications after BMT.

Since there is no gender disparity in the occurrence of HVOD, we concentrated on potential pharmacogenetic issues related to CPSI, an autosomally encoded gene, rather than on the X-linked ornithine transcarbamylase gene. While characterizing the molecular changes underlying the causes of neonatal and late-onset CPSI deficiency, a common SNP near the 3' end of the CPSI mRNA (0.44 heterozygosity) was identified. This C4340A transversion encodes a predicted substitution of asparagine (AAC) for threonine (ACC) at amino acid 1405 (T1405N). This threonine is within the allosteric domain, preceding the sequence PV(A/S)WP(T/S)(A/Q)E important in the binding of a cofactor, n-acetyl-glutamate (NAG), that increases enzyme activity. Although applicants do not wish to be bound by any particular theory of operation, it is speculated that based on the precedent of the effects of other xenobiotics, that limited availability of NAG after escalated dose chemotherapy is one of the mechanisms promoting urea cycle dysfunction. Nonetheless, it appears that the presence of the CPS-I SNP AA genotype is associated with pro-

tection against the development of HVOD, resolution of ALI if it occurs, and improved 60 day survival after BMT. Thus, the data suggest that alteration in UC function plays a role in modifying liver-lung interaction during sepsis and acute lung injury.

In summary, this Example documents significant impairment in hepatic UC function in patients who receive escalated dose chemotherapy prior to BMT. Patients with more severe derangement in cycle function are more likely to develop morbid complications after BMT. Additionally, a significant association between a CPS-I C4340A SNP and both post-BMT complications and short-term survival has been found. Such data are useful in assessment of risk for patients undergoing BMT and provide a rationale for therapeutic attempts to support UC function during high-dose chemotherapy.

Example 3

Arcinine/Citrulline Supplementation Therapy

The added decrease in urea cycle products (arginine and citrulline) and increase in precursors (ammonia, glutamine, etc.) resulting from the polymorphism contribute to BMT associated toxicity. As part of the BMT Life Study, citrulline and arginine levels were measured in 10 patients undergoing BMT.

High-dose chemotherapy used in BMT disrupts normal functions of urea cycle enzymes and contributes to either the occurrence of or toxicity associated with HVOD. To further evaluate this information, an analysis of stored plasma from ten patients undergoing BMT before treatment and after completion of induction chemotherapy was performed. Amino acid profiles were determined from all samples. Particular attention was paid to the urea cycle intermediates citrulline, arginine, and ornithine. As shown in Table 4, a marked decrease in citrulline levels of all patients from a pre-treatment baseline mean of $24 \pm 3 \mu\text{mol/L}$ to a post-treatment mean of $8 \pm 1 \mu\text{mol/L}$ ($P < 0.001$). Plasma arginine levels fell from a mean of $91 \pm 6 \mu\text{mol/L}$ to $70 \pm 6 \mu\text{mol/L}$ ($P < 0.05$), despite the use of arginine-containing parenteral nutrition in several patients:

TABLE 4

Amino Acid	Pre Chemo.	Post Chemo.	P Value
citrulline	$24 \pm 3 \mu\text{M}$	$8 \pm 1 \mu\text{M}$	<0.001
arginine	$91 \pm 6 \mu\text{M}$	$70 \pm 6 \mu\text{M}$	0.03

The fall in citrulline and arginine was similar in patients who did and did not receive total parenteral nutrition and was the same in males and females. The decreases in citrulline suggest that there is a decrease in flow through the first steps of the urea cycle (FIG. 1).

Thus, in accordance with the present invention, a method of reducing toxicity and/or the occurrence of HVOD in a patient undergoing BMT is provided. This method comprises administering the BMT patient arginine and/or citrulline, with citrulline being preferred, in an amount effective to bolster arginine and NO synthesis in the patient. The bolstering of arginine and NO synthesis in the patient reduces and/or substantially prevents the occurrence of HVOD associated with BMT. Citrulline is a preferred supplementation agent given that it is more readily converted to NO.

Example 4

Construction of a Functional Full-Length CPSI Expression Clone

After attempting a number of strategies, a human CPSI cDNA expression clone containing the entire coding region was constructed. FIGS. 6 and 7 present schematic diagrams illustrating the method used to construct the expression clone. This clone has been completely sequenced and does not contain any changes from the consensus CPSI sequence which has been characterized in the art.

The ability of the clone to make CPSI protein was tested in COS-7 cells. COS-7 cells were chosen for their lack of native CPSI activity or production. A western blot analysis of the COS-7 cells transfected with the flCPSI-PCDNA3.1 construct was prepared. HepG2 cell extracts were used as a control as these liver-derived cells have retained CPSI activity. Untransfected COS-7 cells were used as a negative control. Unlike the untransfected COS-7 cells, the HepG2 and COS-7-flCPSI cells demonstrated the expected 160 kDa band using a rabbit anti-rat CPSI antibody. Additionally, a calorimetric assay was performed to detect the production of carbamyl phosphate from ammonia. As shown graphically in FIG. 8, the transfected cells demonstrated activity similar to HepG2 cells while untransfected COS-7 cells did not.

Site-directed mutagenesis has been performed on the T1405 containing CPSI insert and a copy with the N1405 polymorphic codon has been created. The N1405 polymorphic codon was sequenced for its entire length and no other changes were detected. The QuikChange™ (Stratagene) system, which takes advantage of the methylation introduced into DNA by host bacteria, was used to prepare this construct.

These constructs are used to provide a steady supply of recombinant CPSI protein as encoded by both alleles, (T1405, N1405) using COS cells and the respective CPSI/PCDNA 3.1 constructs as an expression system. Enzymatically active CPSI has been produced using this system, as shown by the graph in FIG. 8.

A component of these experiments is to determine the in vitro effect of the T1405N polymorphism on CPSI function. As discussed in Examples 1 and 2, this change affects the sensitivity of the enzyme to NAG concentrations. Screening of 20 individuals for the C to A change showed a heterozygosity rate of 50% with 25% of the group homozygous AA. This suggests that a significant portion of the general population has a potential qualitative abnormality in CPSI function. This abnormality, while silent under normal conditions, is unmasked by stressful conditions and toxins such as high-dose chemotherapy or valproic acid administration.

Comparison of the protein products is then done in stages. The first stage examines the physical characteristics of the expressed mRNA and protein. Using the flCPSI insert as a probe, Northern blots of message prepared from the expressing COS-7 cell lines are probed. Positive controls include HepG2 and human liver message. Negative controls were COS-7 cells transfected with empty cassette pCDNA3.1. The expressed flCPSI derived message is somewhat smaller than the native CPSI (4.9 kb vs. 5.7 kb) since the clone does not contain the 1 kb 3' untranslated region.

Using the same controls, Western blot analysis of cell lysates by SDS-PAGE are performed. Comassie blue staining is used to examine total protein production. For specific CPSI detection, a polyclonal rabbit anti-rat CPSI antibody is used. This antibody detects the expressed CPSI from COS-7 cells as well as the control samples. Finally, changes in the protein's structure are determined by examining the mobility

pattern by 2-D electrophoresis, a useful tool to detect conformational changes. Any large changes in confirmation likely explain the alteration in CPSI function for that mutation.

The next stage involves measuring the functional characteristics of the expressed enzymes. A sensitive calorimetric assay has been modified for this purpose (Pierson, D. L., *J. Biochem. Biophys. Methods*, 3:31-37 (1980)). The modified assay allows 4-5 analyses from 20-50 mg of tissue or cells. The tissue is first homogenized in 0.75M KCl. Small molecules, including ATP and NAG, are removed through a SEPHADEX™ G25 column (Boehringer). The reaction mix contains ammonium bicarbonate, ATP, magnesium DTT, n-acetylglutamate (NAG), and triethanolamine. The concentration of any reagent can be varied, and experiments on HepG2 cells show decreased activity with both low and high concentrations of NAG (0.50 mM). Absence of NAG in preliminary COS-7 cell expression experiments yields no measurable enzyme activity.

Since CPSI is an allosteric enzyme, it does not follow Michaelis-Menton kinetics under varying NAG concentrations; however, when the amount of NAG is fixed, the production of carbamyl phosphate is steady. As shown in FIG. 8, carbamyl phosphate production is measured by the addition of hydroxylamine to the solution after incubation at 37° C. for varying time periods (0, 5, 10, 20, 25, 30 minutes). This step, carried out at 95° C., also serves to inactivate the enzyme and prevent further production of carbamyl phosphate. The hydroxylamine converts the carbamyl phosphate to hydroxyurea which is subsequently treated with a sulfuric/acetic acid solution with butanedione to derive a compound with peak absorption at 458 nm. The reaction is then spun at 12,000×g for 15 minutes to remove precipitated protein. Next, the 458 nm absorbance is measured for each reaction. Activity typically begins to decrease after 20-30 minutes of reaction.

A number of expressing cell pellets are pooled for analysis. To ensure that activity measurements are based on consistent amounts of enzyme, expressed CPSI is quantified by Western blot analysis of the pooled sample using a CPSI antibody such as the rabbit anti-rat CPSI described hereinabove. Basal activity is first determined using fixed amounts of substrate and cofactor and a time course analysis. Varying amounts of ammonia bicarbonate, ATP, and NAG are then used to determine the binding efficiency for these elements. These elements are varied from 0 to 10-fold the normal amount. Enzyme activity is also measured after heat treatment of the homogenate. Protein labeling (pulse-chase) experiments are performed to determine the stability of the protein over time.

Stable CPSI protein expression is obtained using the methods described above. The establishment of stable transfected cell lines allows the production of sufficient quantities of both varieties of CPSI to carry out these studies. In activity studies, changes in activity for the N1405 as compared to the T1405 type of CPSI are noted. A change in the enzyme activity under varying concentrations of NAG is also noted. These results support the role of this polymorphism of the present invention in predicting susceptibility to sub-optimal urea cycle function and hyperammonemia and decreased arginine production associated therewith.

Example 5

Relationship of the T1405N Polymorphism and Urea Cycle Intermediates to the Ammonia Elevation Seen in Patients on Valproic Acid Therapy

Valproic acid (VPA) is a commonly used seizure medication, particularly for the treatment of absence seizures or as an

adjunct therapy of other seizure disorders. Toxicity from VPA treatment is a complex and multi-variant process and probably reflects several metabolic disruptions. Hyperammonemia and hepatic micro-vesicular steatosis and necrosis are the most commonly reported serious medical complications.

Although the development of toxic hyperammonemia involves only a small number of patients, it carries a significant morbidity and mortality, and several deaths have been attributed to this complication. The development of asymptomatic hyperammonemia (plasma ammonia level greater than 60 μmol/L) occurs within one hour of VPA administration, and is, however, relatively common.

Mechanisms of VPA-induced Hyperammonemia. The mechanisms by which VPA causes hyperammonemia has been the subject of some debate, and a number of different theories currently have support in the art. A renal model proposed that the changed in glutamine metabolism resulted in an increased ammonia load to the liver, while most other theories concentrate on different aspects of urea cycle function. See, for example, Warter et al., *Revue Neurologique*, 139:753-757 (1983). Since the urea cycle is the major mechanism for the removal of ammonia in humans, it is thought that hyperammonemia arises in some way from the inhibitory interactions of VPA and/or its metabolites with urea cycle function and capacity.

Evidence for urea cycle dysfunction in VPA therapy comes from a number of experimental and clinical observations aside from elevations in plasma ammonia described above. For example, Marrini et al. measured a reduction in both baseline and stimulated CPSI activity in non-nephrectomized animals following an amino acid and VPA load (Marrini et al., *Neurology* 38:365-371 (1988)). Marrini et al. also observed that nephrectomized rats injected with an amino acid load and VPA also developed hyperammonemia. Another group, Castro-Gago et al., measured serum amino acids in 22 epileptic children treated with VPA, and found reduction in aspartic acid and ornithine, implicating a decrease in urea cycle efficiency rather than an increase in precursors (Castro-Gago et al., *Childs Neurons System* 6:434-436 (1990)).

Significance of Carbamyl Phosphate Synthase I. Mechanisms of VPA-induced urea cycle deficits typically revolve around mitochondrial carbamyl phosphate synthetase I (CPSI). A patient with severe toxicity following VPA overdose was found to have 50% normal CPSI activity (Bourrier et al., *Prese Medicale* 17:2063-2066 (1988)). Applicants have observed several mild CPSI deficient patients who deteriorated when given valproic acid with ready reversal after discontinuation.

Role of NAG. N-acetylglutamate (NAG) is a required allosteric cofactor for CPSI. NAGA is synthesized from glutamate and acetyl CoA in mitochondria, with a cellular distribution that mirrors that of CPSI (Shigesada et al., *Journal of Biological Chemistry* 246: 5588-5595 (1971)). It is synthesized from glutamate (from amino acid catabolism) and acetyl CoA. There are several ways in which an alteration of NAG availability is envisaged to reduce the activity of CPSI. Genetic deficiencies in NAG synthetase have been observed, and this enzyme is known to be inhibited competitively by alternate substrates such as propionyl CoA or succinate (Bachmann et al., *New England Journal of Medicine* 304:543 (1981); Kamoun et al., *Lancet* 48 (1987); Coude et al., *J. Clin. Invest.* 64:1544-1551 (1979); Rabier et al., *Biochem. And Biophys. Research Comm.* 91:456-460 (1979); Rabier et al., *Biochimie* 68:639-647 (1986)). It has been shown experimentally that CPSI is inhibited in a competitive manner by the presence of increased amounts of propionyl CoA, and that VPA therapy causes an increase in blood pro-

pionate concentration (Coulter et al., *Lancet* 1 (8181): 1310-1311 (1980); Gruskay et al., *Ped. Res.* 15:475 (1981); Schmidt, R. D., *Clin. Chim. Acta.* 74:39-42 (1977)). VPA exposure has also been shown to decrease NAG concentrations in intact hepatocytes, by decreasing concentrations of both acetyl CoA and glutamine (Coude et al., *Biochem. J.* 216:233-236 (1983)). The decrease in glutamine concentration is attributed to inhibition of both pyruvate dehydrogenase and pyruvate carboxylase.

Alternatively, it has been suggested that depletion of mitochondrial acetyl CoA occurs because CoA is diverted on VPA therapy for the manufacture of valproyl CoA (Becker et al., *Archives of Biochemistry & Biophysics* 223:381-392 (1983)). It is well known that VPA also disrupts fatty acid β -oxidation, with resultant diminution of acetyl CoA (Eadie et al., *Med. Toxicol.* 3:85-106 (1998)). All these mechanisms could lead to a shortage in NAG since it is synthesized from acetyl CoA. Given the effects of VPA on NAG availability it follows that any change in the binding properties of CPSI for NAG would affect its activity.

Thus, this Example sets forth experimentation for determining correlation between the presence or absence of the polymorphism of the present invention in the CPSI gene with susceptibility to hyperammonemia using VPA as a model agent for the production of hyperammonemia. Initially, genomic DNA is isolated from patients who are beginning valproic acid therapy for genotyping for the T1405N polymorphism in accordance with the methods described herein, such as PCR amplification and use of non-denaturing gels. After genotyping these patients, pre- and post-treatment amino acid and ammonia determination is performed for these patients. Particularly, DNA is isolated from whole blood using the QIAmp™ (Qiagen) kit described in Example I.

Next, plasma total VPA concentration is determined by an enzyme-mediated immunoassay technique (EMIT™ Syva-Behring, San Jose, Calif. on a Syva 30R™ analyzer). This technique utilizes competitive binding for VPA antibody binding sites between VPA in the patient plasma and that complexed with the enzyme G6PDH. Release of the VPA enzyme complex from the antibody reactivates the enzyme, and its activity is assessed by the rate of formation of NADH upon addition of the substrate. NADH production is monitored via spectroscopy at 340 nanometers (nm). Free (non-protein bound) VPA is isolated from plasma using a centrifugal micro partition filter device with a 3000 Dalton cut-off (CENTRI FREE, Aimcon, Beverley, Mass.). The VPA concentration in the plasma ultra filtrate is measured as described for total VPA.

Data collected from VPA patients is analyzed for correlations between genotype and phenotype. Additionally, free and conjugated VPA fractionation are compared to evaluate effects on NAG production and availability. The latter comparison is prepared given that there are known effects of VPA on NAG availability. For example, VPA exposure has been shown to decrease NAG concentrations in intact hepatocytes by decreasing concentrations of both acetyl CoA and glutamine. See Coude et al., *Biochem. J.* 216:233-236 (1983). Thus, this comparison reflects that changes in the binding properties of CPSI for NAG affect the activity of CPSI.

Example 6

Detection of Additional Polymorphisms in CPSI

Using the techniques developed for mutation analysis of CPSI message, 10 non-CPSI deficient, unrelated patients are

screened for additional polymorphisms in the coding region. This is done using "illegitimate" transcripts from lymphoblastoid and fibroblast cell lines. Polymorphisms with a widespread effect on the population should be evident in this size sample. As used herein and in the claims, the term "polymorphism" refers to the occurrence of two or more genetically determined alternative sequences or alleles in a population. A polymorphic marker is the locus at which divergence occurs. Preferred markers have at least two alleles, each occurring at frequency of greater than 1%. A polymorphic locus may be as small as one base pair. Provided polymorphic markers thus include restriction fragment length polymorphisms, variable number of tandem repeats (VNTR's), hypervariable regions, minisatellites, dinucleotide repeats and tetranucleotide repeats.

A number of "mutation" detection techniques have been carried out, all of which are based on detectable changes in the mobility of non-denatured single-stranded DNA, as described by Summar, M., *J. Inherited Metabolic Disease* 21:30-39 (1998). Examples of CPSI mutations identified by these techniques are disclosed in FIG. 3. Due to the large size of the CPSI message (about 5,700 bases) a method to screen a large amount of DNA in a few reactions is preferred. Restriction endonuclease fingerprinting (REF) provides for the screening large DNA fragments, up to about 2,000 bp, with excellent sensitivity.

Reverse transcriptase reactions (RT) are carried out using 1 μ g of total RNA and either an oligo-dt primer or an antisense primer from the midpoint of the CPSI message. Using the RT product as template, PCR reactions are performed with 4 different primer sets creating 4 overlapping fragments spanning the 4,600 base coding region. Control PCR reactions are run with each set of experiments, to ensure that contaminating template is not amplified. Genomic DNA is not preferred for this study due to the size of the gene (80,000+bp), the number of introns (36), and that sequencing of the intron/exon boundaries for CPSI has not been completed. However, intronic locations are characterized graphically in FIG. 9.

The 4 overlapping RT/PCR products described above are used for mutation screening. Careful analysis of the restriction maps leads to the selection of three restriction enzymes for each fragment which cleave them into pieces ranging from 100-250 bp. Fragments of this size are ideal for single strand conformation polymorphism (SSCP) analysis. The enzymes are selected such that each fragment can be evenly evaluated across its length. Prior to digestion, the PCR products are purified by gel electrophoresis and isolation from the agarose slices. After 3 hours, the digested fragments are ethanol precipitated. These fragments are separated in a 6% non-denaturing polyacrylamide gel at 4° C. running at a constant 35 watts. These conditions maximize the detection of conformational changes in the single stranded fragments, as described by Liu, Q. and Sommer, S. S., *Biotechniques* 18(3):470-477 (1995). DNA detection is done by silver staining and the gels are scored for mobility shifts. Based on the location of any shifted fragment, direct sequence analysis of the RT/PCR product is performed using a cycle-sequencing protocol. To eliminate the possibility of a mutation resulting from Taq polymerase errors, a fresh RT product is amplified and sequenced in each case. The entire 4,600 bases of coding message is rapidly screened in this fashion. Any regions containing unclear areas are sequenced, looking for changes in the expected sequence.

The restriction digestion products of each RT/PCR fragment are isolated. These individual fragments are then run against the combined digestion in a non-denaturing gel as described above. By characterizing the fragment pattern in

this way, the portions of the CPSI message involved in any observed mobility shifts are readily identified.

Polymorphisms detected in these experiments are genotyped against the Centre d'Etude Polymorphsim Humanise (CEPH) parents panel to establish frequency. All changes are examined for their effect on codon use and those resulting in mis-sense mutations are examined using the CPSI characterization data disclosed herein.

The techniques described in Example 3 are used to express site-directed mutants containing these changes. Using this system the in vitro effects of the changes on CPSI production and activity are observed.

A T344A polymorphism was detected in CPSI. Oligonucleotide primers were used from the 10th exon (U1119: tactgctcagaatcatggc—SEQ ID NO:17) and intron (L110+37: tcataccaactgaacagg—SEQ ID NO:18) to amplify a 91 bp fragment containing the change. PCR cycle conditions were: 35 cycles of 1 minute anneal at 59° C., 1 minute extension at 72° C., and 1 minute denaturation at 94° C. Patients were classified as having either homozygous SNP genotypes of AA or TT, or as being heterozygous (AT). The adult population distribution of this polymorphism is 35% AA, 44% AT, and 21% TT.

A 118-CTT polymorphism was also detected in CPSI. Oligonucleotide primers were used from the 5' untranslated region (U5'-74: ggtaagagaaggaggagctg—SEQ ID NO:19) and intron (L175: aaccagtcttcagtgtcctca—SEQ ID NO:20) to amplify a 249 bp fragment containing the change. PCR cycle conditions were: 35 cycles of 1 minute anneal at 59EC, 1 minute extension at 72° C., and 1 minute denaturation at 94° C. Patients were classified as having either a homozygous genotype with the 118 trinucleotide insertion or deletion, or as being heterozygous. The adult population distribution of this polymorphism is 34% CTT-, 43% heterozygous, and 23% CTT+.

Example 7

Biochemical and Genetic Alterations in Carbamyl Phosphate Synthetase I in Neonatal Patients with Persistent Pulmonary Hypertension

This Example investigates the role of the limitation of endogenous NO production in the pathogenesis of persistent pulmonary hypertension (PPHN) in the sick term neonate. Endogenous NO is the product of the urea cycle intermediate arginine. Production of arginine depends on the rate-determining enzyme of the urea cycle, carbamyl phosphate synthetase (CPSI). Newborns possess less than half the normal urea cycle function making them particularly susceptible to minor changes in enzyme form and function. A common exonic polymorphism (T1405N) in CPSI has been observed which affects flow through the first step of the urea cycle.

In this Example, it was tested whether newborns who developed PPHN would have lower NO precursors (arginine and citrulline) than matched controls. Whether PPHN patients have predominantly the CC (threonine/threonine) or AC (asparagine/threonine) CPSI genotypes which are associated with lower function than AA (asparagine/asparagine) CPSI genotype was also analyzed.

Methods. Forty-seven neonates >2 kg, >35 weeks, and <72 hours old who were admitted to the Vanderbilt Neonatal Intensive Care Unit with (n=22) and without (n=25) echocardiographically-documented pulmonary hypertension were enrolled. Clinically important measures of the severity of respiratory distress were recorded. Ammonia levels and

plasma amino acid profiles were obtained. Genotypes were determined by running PCR-amplified DNA on nondenaturing MDE™ gels.

Results. Patients who developed PPHN had an average arginine of 21.5 $\mu\text{mol/l}$ while those who did not averaged 38.3 $\mu\text{mol/l}$ ($p=0.0004$). The citrulline averages were 6.1 $\mu\text{mol/l}$ and 10.3 $\mu\text{mol/l}$ respectively ($p=0.02$). The levels of arginine and citrulline were inversely correlated with the severity of hypoxemia as measured by oxygenation index, days of mechanical ventilation, and days requiring supplemental O_2 . Genotype analysis of PPHN patients for T1405N showed 5CCs, 17ACs, and 0AAs, whereas the controls had 7CCs, 16ACs, and 2AAs (Chi-square $p=0.005$ using the expected population allele frequency). Infants with the CC genotype had lower arginine and citrulline means (21.5 $\mu\text{mol/l}$ and 5.8 $\mu\text{mol/l}$) than infants with the AA genotype (31.5 $\mu\text{mol/l}$ and 13.5 $\mu\text{mol/l}$) consistent with a functional difference between the two forms of the enzyme.

Conclusions. This Example shows that the development of PPHN in sick newborns is associated with inadequate availability of the urea cycle intermediates arginine and citrulline. The T1405N polymorphism in the CPSI DNA leads to diminished enzyme function and subsequent lower levels of NO precursors.

Discussion. Carbamyl phosphate synthetase (CPS I) catalyzes the rate-determining step in the urea cycle thereby determining tissue levels of the urea cycle intermediates including arginine and citrulline. As disclosed herein, a widely distributed C to A exonic polymorphism in the CPS I gene changes a conserved threonine to an asparagine at position 1405 near the critical N-acetyl glutamate binding domain. Data has shown that the asparagine-containing version of CPSI displays more efficient kinetics in enzyme function studies.

The T1405N allele exhibits 50% heterozygosity and appears to be a silent variant in normal healthy adults. However, consequences of the qualitative change can be unmasked by stressful conditions. As disclosed in Examples 1-3, adults exposed to high-dose chemotherapy in preparation for bone marrow transplantation that the threonine-containing enzyme produces inadequate levels of arginine and citrulline and is associated with an increased incidence of hepatic veno-occlusive disease, acute lung injury, and death. As nitric oxide (NO) is generated in endothelial cells from L-arginine by nitric oxide synthetase (NOS), decreased levels of urea cycle intermediates could predispose to disturbances in vascular tone by limiting endogenous NO production.

In the prospective cohort study of this Example, the possibility that a similar process could be involved in the pathogenesis of persistent pulmonary hypertension of the newborn (PPHN) was investigated. Endogenously produced NO functions in regulation of pulmonary vascular resistance and in the transition from fetal to neonatal circulation. Lipsitz, E. C., et al. *J Pediatr Surg* (1996) 31:137-140; Abman, S. H., et al. *Am J Physiol* (1990) 259:H1921-H1927. Between 20 weeks gestation and term birth, CPSI production and function are less than 50% of adult levels. This physiologic deficiency could unmask the effect of the T1405N gene mutation particularly if coupled with other neonatal stresses affecting hepatic function; for instance, asphyxia or sepsis.

Patients eligible for this study included appropriately grown neonates ≥ 35 weeks gestation and ≥ 2 kg birthweight who were admitted to the Vanderbilt University Medical Center neo-natal intensive care unit (NICU) between Jul. 1, 1999 and Feb. 29, 2000 for symptoms of respiratory distress. Infants with multiple congenital anomalies, known genetic syndromes, and anatomic causes of pulmonary hypertension

(congenital diaphragmatic hernia, Potter=s syndrome, asphyxiating thoracic dystrophy, etc.) were excluded. Parental consent was obtained for all enrollees. Fifty-one neonates had 3 cc of blood drawn in the first 72 hours of life for plasma amino acid profiles, ammonia and BUN levels, nitric oxide metabolite determination, and CPS1 genotyping. Blood was drawn prior to blood transfusion, enteral or parenteral protein intake, inhaled nitric oxide administration, or ECMO cannulation.

Data collected on the enrollees included (1) baseline characteristics (birthweight, gestational age, sex, race, Apgar scores, primary diagnosis, any pulmonary complications, and the postnatal age at the time blood was drawn) and (2) measures of respiratory support (FiO₂, MAP, iNO, ECMO) and clinical response (ABGs, duration of mechanical ventilation and supplemental O₂, survival.) Maximum oxygenation index [OI=FiO₂×MAP/PaO₂] was used as a measure of the severity of respiratory distress. Predominant primary diagnoses included (1) birth asphyxia: 5-minute Apgar score <5 with a mixed acidosis on first ABG or cord blood gas plus evidence or neurologic dysfunction and other end-organ injury, (2) respiratory distress syndrome (RDS): clinical symptoms of respiratory distress with ground-glass lung fields and air bronchograms on chest X-ray plus combined hypercarbia/hypoxia on ABG (Note: given the gestational age of these neonates, infants with this picture could have had either surfactant-deficiency or congenital pneumonia; however, in no case was a positive tracheal aspirate culture obtained), and (3) meconium aspiration syndrome (MAS): history of meconium-staining at delivery plus clinical symptoms of respiratory distress, hypoxemia, and coarse infiltrates chest X-ray.

Infants were defined as having pulmonary hypertension (PPHN) if they developed significant hypoxemia (PaO₂<100 on 100% O₂>6 hours) with normal intracardiac anatomy and echocardiographic evidence of elevated pulmonary artery pressure. The latter was defined as (1) right-to-left or bidirectional ductal or foramen ovale flow or (2) elevated (>35 mmHg) pulmonary artery pressure based on Doppler estimate of the tricuspid regurgitation jet as read by a blinded third party.

Amino acid analysis was performed on fresh plasma samples in 47 patients. A protein free extract of plasma was prepared by protein precipitation with sulfosalicylic acid and filtration through a 0.45 μm Acrodisc 4 (Gelman Sciences, Ann Arbor, Mich.). Amino acids were separated by cation exchange chromatography using a four-component pH- and ionic strength-graded lithium citrate buffer system on a Beckmann 7300 amino acid analyzer (Beckmann, Palo Alto, Calif.). Post column derivatization of amino acids with ninhydrin allowed detection of primary amine amino acids at 570 nm, and secondary amines at 440 nm. Quantitation was achieved by instrument calibration with standard reference materials (Sigma, St. Louis, Mo.). Citrulline and arginine were detected as measurable indices of flux of intermediates through the urea cycle.

Measurement of plasma nitric oxide metabolites (NO_x). Plasma NO_x was measured in a subgroup of patients using modified Griess reagents after samples were deproteinated and incubated with cadmium beads to convert nitrate to nitrite.

SNP Detection. Oligonucleotide primers from within the 36th exon (U4295—SEQ ID NO:15) and intron (LI36—SEQ ID NO:16) of CPS1 and the polymerase chain reaction (PCR) to reliably amplify a 251 bp fragment encompassing the region containing the change from genomic DNA obtained from whole blood preparations. This combination of primers

gave reproducible amplification using Taq polymerase (Promega) and PCR cycle conditions as follows: 35 cycles of 1 minute anneal at 67° C., 1 minute extension at 72° C., and 1 minute denaturation at 94° C. After formamide treatment, samples were subjected to electrophoresis for 5 hours at 4° C. in a non-denaturing MDE™ gel (FMC, Rockland, Me.), then stained with silver nitrate to detect DNA fragments. Patients were classified as having homozygous SNP genotypes of CC or AA, or as being heterozygous (AC). Genotyping using nondenaturing gel electrophoresis and direct sequence analysis yielded identical results as those disclosed above. Thus, the adult population distribution of the T1405N polymorphism was determined to be: 45% CC, 44% AC, and 11% AA.

An identical technique to that described above was used to detect the T344A polymorphism. Oligonucleotide primers were used from the 10th exon (U1119: tactgctcagaatcatggc—SEQ ID NO:17) and intron (LI10+37: tcatcaccactgaacagg—SEQ ID NO:18) to amplify a 91 bp fragment containing the change. PCR cycle conditions were: 35 cycles of 1 minute anneal at 59, 1 minute extension at 72° C., and 1 minute denaturation at 94° C. Patients were classified as having either homozygous SNP genotypes of AA or TT, or as being heterozygous (AT). The adult population distribution of this polymorphism is 35% AA, 44% AT, and 21% TT.

An identical technique to that described above was used to detect the 118-CTT polymorphism. Oligonucleotide primers were used from the 5' untranslated region (U5'-74: gggttaagagaaggaggagctg—SEQ ID NO:19) and intron (L175: aacagctctcagtgctctca—SEQ ID NO:20) to amplify a 249 bp fragment containing the change. PCR cycle conditions were: 35 cycles of 1 minute anneal at 59° C., 1 minute extension at 72° C., and 1 minute denaturation at 94° C. Patients were classified as having either a homozygous genotype with the 118 trinucleotide insertion or deletion, or as being heterozygous. The adult population distribution of this polymorphism is 34% CTT-, 43% heterozygous, and 23% CTT+.

Ammonia and plasma amino acid levels were compared between groups of patients using Student's T-test. Distributions of genotypes of CPSI were compared across groups by calculating allelic frequency for the entire group and searching for evidence of Hardy-Weinberg disequilibrium in specifically selected subgroups using Chi-square analysis. Of the 51 neonates originally enrolled, 25 developed PPHN while 26 did not. There were no statistically significant differences in the baseline characteristics of the two groups including birthweight, gestational age, race, or the postnatal age in hours of the infants at enrollment. There was, however, a slight predominance of males in the control group.

The distribution of primary diagnoses was evenly distributed. In the PPHN group, 5 infants had birth asphyxia, 9 infants had RDS, 5 infants had meconium aspiration syndrome, and 6 infants had other diagnoses, including 4 infants with primary PPHN. In the control group, 4 infants had birth asphyxia, 8 infants had RDS, 3 infants had MAS, and 11 infants had other diagnoses. The other diagnoses included supraventricular tachycardia, anemia, birth trauma, and viral sepsis. No infant in the study had a positive bacterial blood culture.

As expected, infants who had PPHN complicate their primary pathology did develop more severe illness than the controls by some clinical criteria. Eight of the infants with PPHN required treatment with inhaled NO (iNO), 2 required ECMO, and 2 died (one infant with asphyxia and multiorgan-system failure on iNO; another infant with alveolar capillary dysplasia was withdrawn from ECMO.) Obviously, none of the controls were treated with iNO or ECMO; and there was no mortality in the control group.

Three infants in the PPHN group were excluded from analysis. The infant found to have alveolar capillary dysplasia on lung biopsy was considered to have an anatomical etiology for pulmonary hypertension. Another infant was mistakenly enrolled with a congenital diaphragmatic hernia, and the third was enrolled at 119 hours of age after TPN had been initiated. One infant in the control group was excluded from analysis after karyotype analysis revealed the etiology of his hypotonia to be Prader-Willi syndrome.

The infants who developed PPHN had significantly lower serum arginine and citrulline levels on amino acid analysis. The mean arginine level in PPHN cases was 21.5 ± 9.2 $\mu\text{mol/l}$ whereas the mean arginine of the control group was 38.3 ± 18.4 $\mu\text{mol/l}$ ($p=0.0004$). The mean citrulline in PPHN cases was 6.1 ± 3.6 $\mu\text{mol/l}$ compared to 10.3 ± 7 $\mu\text{mol/l}$ in the control group ($p=0.02$). There were no significant differences in the levels of other amino acids between the two groups, including glutamine, glycine, alanine, lysine, valine, ornithine, and leucine. The level of total essential amino acids (TEAA) was slightly lower in the PPHN cases, about 537 $\mu\text{mol/l}$ versus about 654 $\mu\text{mol/l}$, but this difference was not statistically significant ($p=0.08$). by birthweight, gestational age, or number of hours of postnatal life. The level of TEAA was found to be significantly higher in the four infants whose blood was drawn prior to six hours of age (about 1021.5 $\mu\text{mol/l}$ vs. about 542 $\mu\text{mol/l}$, $p=0.0026$). This difference is presumed to reflect the recent cessation of parenteral protein influx in these infants from the placental circulation.

No differences in arginine and citrulline levels were found when the primary diagnosis categories of asphyxia, RDS, MAS, and "other" were separately analyzed. In each group, infants with pulmonary hypertension tended to have lower values, but the results were not statistically significant given the small numbers of infants in each group. For example, asphyxiated infants with PPHN had a mean arginine of about 18.5 $\mu\text{mol/l}$ compared to about 52.7 $\mu\text{mol/l}$ in asphyxiated controls ($p=0.06$) and a mean citrulline of about 6.8 $\mu\text{mol/l}$ compared to about 14.3 $\mu\text{mol/l}$ ($p=0.04$).

There was an inverse relationship between the levels of serum arginine and citrulline and the severity of hypoxemia. Arginine and citrulline values fell progressively as oxygenation index increased, days of mechanical ventilation increased, and days requiring supplemental oxygen increased birthweight, gestational age, or number of hours of postnatal life. The NH_3 levels in infants with PPHN tended to be slightly higher than in controls (54 ± 18.1 $\mu\text{mol/l}$ vs. 45.6 ± 12 $\mu\text{mol/l}$) but these values were not statistically significant ($p=0.08$).

On CPS1 T1405N genotype analysis, of the 22 infants who developed PPHN, 5 were CC and 17 were AC. There were no AAs in the PPHN cases. In the 25 controls, there were 7 CCs, 16 ACs, and 2 AAs. These distributions of genotypes were then compared by calculating the expected allelic frequency for the entire group revealing evidence of Hardy-Weinberg disequilibrium in the PPHN group. On Chi-square analysis these two groups are significantly different from each other with a p -value=0.005. Of the two infants with the AA genotype, one infant had RDS while the other suffered from birth asphyxia. Neither infant ever achieved an $\text{OI} \geq 15$; both spent <1 week on the ventilator and <10 days on oxygen.

Infants with the CC genotype had mean arginine levels of 21.9 ± 7 $\mu\text{mol/l}$ and citrulline levels of 5.8 ± 1.8 $\mu\text{mol/l}$ while infants with the AA genotype had a mean arginine level of 31.5 ± 3.5 $\mu\text{mol/l}$ and a mean citrulline level of 13.5 ± 6.4 $\mu\text{mol/l}$. Again, given the small number of AAs, this data has difficulty reaching statistical significance with p -values of 0.1 and 0.006, respectively.

Intravenous Citrulline Supplementation Increases Plasma Arginine Levels in Piglets

Intravenous citrulline has not been previously used in a clinical model. This Example assessed the safety of IV citrulline and its effect on serum arginine levels in piglets. A total of 9 Duroc swine, aged 5-21 days, with a target minimum weight of 4 kg were utilized. All piglets underwent anesthetic induction and tracheostomy. Central lines were placed in the femoral artery and femoral vein and hemodynamics monitored continuously. Citrulline (600 mg/kg IV) was administered to 5 piglets. Saline was given to control animals. Serum amino acids were drawn before and each hour after citrulline administration.

Serum arginine levels peaked at 1-2 hours following IV citrulline administration and remained sustained above baseline three hours following, reaching significance at all time points compared to controls ($p < 0.001$). No hemodynamic instability was observed.

Arginine Levels ($\mu\text{mol/L}$) Following IV Citrulline

Treatment Group (n = 5)	Baseline	1 hour post	2 hours post	3 hours post
Citrulline (600 mg/kg)	131.5	535.0	559.8	498.4
Control (saline)	89.6	103.0	118.1	136.7
p-value	0.1582	<0.001	<0.001	<0.001

Mean Arterial Blood Pressures (mmHg) Following IV Citrulline

Treatment Group (n = 4)	Pre-dose	1 hour post	2 hours post	3 hours post
Citrulline (600 mg/kg)	67.0	67.4	64.8	62.2
Control (saline)	53.2	58.7	55.7	54.7

$p > .05$ at all time points

Pharmacokinetics: Based on the above data, the pharmacokinetics were calculated for both plasma citrulline and arginine levels after the single dose of IV citrulline. Pharmacokinetic data included plasma half-life ($t_{1/2}$), elimination constant (K_{el}), volume of distribution (V_d), and plasma clearance (CL_p).

Plasma citrulline levels rapidly increased and demonstrated a $t_{1/2}=1.5$ hrs, $K_{el}=0.462$ hr^{-1} , $V_d=2.25$ L, and $CL_p=1.05$ L/hr. However, the effect of citrulline on plasma arginine was of interest because it is the substrate for NO synthase. The concentration curve of plasma arginine levels is represented in FIG. 13. Based on this curve, the pharmacokinetics of plasma arginine are as follows: $t_{1/2}=18$ hrs; $K_{el}=0.039$ hr^{-1} ; $V_d=2.85$ L; $CL_p=0.11$ L/hr. The long half-life and slow clearance indicates that a single dose of IV citrulline is effective at maintaining increased plasma arginine levels over a fairly long interval without detrimental effects on hemodynamics.

REFERENCES

The references listed below as well as all references cited in the specification are incorporated herein by reference to the

extent that they supplement, explain, provide a background for or teach methodology, techniques and/or compositions employed herein.

- Abman, S. H., et al., *Am J Physiol* (1990) 259:H1921-H1927.
 Adelman et al., *DNA* 2:183 (1983).
 Alonso, E. and Rubio, V., *European Journal of Biochemistry* 229:377-384 (1995).
 Artymiuk, P. J. et al., *Nature Struct. Biol.* 3:128-132 (1996).
 Bachmann et al., *New England Journal of Medicine* 304:543 (1981).
 Batshaw M L, Brusilow S W. *Annals of Neurology* 1982; 11:319-21.
 Bearman S I, *Journal of Clinical Oncology* 1993; 11:1729-36.
 Beaucage et al., *Tetrahedron Letters* 22:1859-1862 (1981).
 Beaumier L, *Biomedical & Environmental Sciences* 1996; 9:296-315.
 Becker et al., *Archives of Biochemistry & Biophysics* 223: 381-392 (1983).
 Bernard G R, Artigas A, Brigham K L, et al. *Intensive Care Medicine* 1994; 20:225-32.
 Blau N, Duran, M., and Blaskovics, M. E. *Physician's Guide to the Laboratory Diagnosis of Metabolic Diseases* London: Chapman&Hall Medical, 1996.
 Bourrier et al., *Prese Medicale* 17:2063-2066 (1988).
 Castillo, L., et al., *Pediatr Res* (1995) 38:17-24.
 Castillo L, et al., *Journal of Biological Chemistry* 1989; 264: 4038-44.
 Castro-Gago et al., *Child Neuro Systems* 6:434-436 (1990).
 Cervera, J. et al., *Biochemistry* 35:7247-7255 (1996).
 Cohen P P. *Current Topics in Cellular Regulation* 1981; 18:1-19.
 Coude et al., *Biochem. J.* 216:233-236 (1983).
 Coude et al., *J. Clin. Invest.* 64: 1544-1551 (1979).
 Coulter et al., *Lancet* 1 (8181): 1310-1311 (1980).
 Crea et al., *Proc. Natl. Acad. Sci. USA* 75:5765 (1978).
 Davies et al., *Bone Marrow Transplantation* 17:1119-1125 (1996).
 de Groot, C. J., et al., *Biochemical & Biophysical Research Communications* 124:882-888 (1984).
 Eadie et al., *Med. Toxicol.* 3:85-106 (1998).
 Eichenlaub et al., *J. Bacteriol.* 138:559-566 (1979).
 Faber-Langendoen K, et al. *Bone Marrow Transplantation* (1993) 12:12501-7.
 Gribskov et al., *Nuc. Acids. Res.* 14:6745 (1986).
 Gruskay et al., *Ped. Res.* 15:475 (1981).
 Guillou, F., et al. *Proc Natl Acad Sci* 86:8304-8308 (1989).
 Guy, H. I. et al., *Journal of Biological Chemistry* 270:2190-2197 (1995).
 Hauser E R, et al., *New England Journal of Medicine* 1990; 322:1641-5.
 Hebert P C, *Chest* 1993; 104:230-5.
 Howell et al., *Antibodies A Laboratory Manual* Cold Spring Harbor Laboratory, (1988).
 Jackson, M. J. et al., *Annual Review of Genetics* 20:431-464 (1986).
 Jackson M J, *Annual Review of Genetics* 1986; 20:431-64.
 Javid-Majd et al., *Biochemistry* 35:14362-14369 (1996).
 Jones R J, et al. *Transplantation* 1987; 44:778-83.
 Kamoun et al., *Lancet* 48 (1987).
 Kinsella, J. P., et al., *Lancet* (1992) 340:819-820.
 Kyte & Doolittle, *J. Mol. Biol.* 157:105-132 (1982).
 Lagace, M. et al., *Journal of Biological Chemistry* 262: 10415-10418 (1987).
 Lipsitz, E. C., et al., *J Pediatr Surg* (1996) 31:137-140.
 Liu, Q. and Sommer, S. S., *Biotechniques* 18(3):470-477 (1995).

- Maniatis et. al. in *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor, N.Y., p 280-281 (1982).
 Marrini et al., *Neurology* 38:365-371 (1988).
 Marshall J C, et al., *Critical Care Medicine* (1995) 23:1638-52.
 Matuschak, G. M., *Clinics in Chest Medicine* (1996) 17:83-98.
 Matuschak, G. M. and Rinaldo, J. E., *Chest* (1988) 94:400-6.
 Matuschak et al., *American Review of Respiratory Disease* 1990; 141:1296-306.
 McCaffrey, M. J., et al., *Biol Neonate* (1995) 67:240-243.
 McDonald, G. B., et al., *Annals of Internal Medicine* 1993; 118:255-67.
 Meister, A., *Adv. Enzymol. Relat. Areas Mol. Biol.* 62:315-374 (1989).
 Messing et al., *Third Cleveland Symposium on Macro Molecular and Recombinant DNA* Ed. Walton, A., (Elsevier, Amsterdam) (1981).
 Mitchell R B, Wagner J E, Karp J E, et al. *American Journal of Medicine* 1988; 85:662-7.
 Mitchell et al., *Amer. J. Med.* 85:662-667 (1988).
 Moncada S, Higgs A. *New England Journal of Medicine* 1993; 329:2002-12.
 Moorman, A. F. et al. *Histochemical Journal* 22:457-468 (1990).
 Needleman et al., *J. Mol. Biol.* 48:443 (1970).
 Nuzum, C. T. and Snodgrass, P. J., *Science* 1971; 172:1042-3.
 Nyunoya, H., et al., *Journal of Biological Chemistry* 260: 9346-9356 (1985).
 Palmer R M J, et al., *Biochem Biophys Res Commun* (1988) 153:1251-1256.
PCR. A Practical Approach, ILR Press, Eds. McPherson, et al. (1992).
 Pierson, D. L., *J. Biochem. Biophys. Methods* 3:31-37 (1980).
 Price, K. J., et al., *American Journal of Respiratory & Critical Care Medicine* 1998; 158:876-84.
 Rabier et al., *Biochem. & Biophys. Research Comm.* 91:456-460 (1979).
 Rabier et al., *Biochimie* 68:639-647 (1986).
 Raiha, N. C. R. and Suihkonen, J. *Acta Paediatrica Scand* 57:121-127 (1968).
 Richardson, P. and Bearman, S. I., *Leukemia & Lymphoma* (1998) 31:267-77.
 Rinaldo J E, et al., *American Journal of Respiratory Cell & Molecular Biology* 1994; 11:625-30.
 Roberts, J. D., et al., *Lancet* (1992) 340:818-819.
 Rodriguez-Aparicio, L. B. et al., *Biochemistry* 28:3070-3074 (1989).
 Rubenfeld G D, Crawford S W. *Annals of Internal Medicine* 1996; 125:625-33.
 Rubio, V., (Review) *Biochemical Society Transactions* 21:198-202 (1993).
 Rubio, V. and Grisolia, S., *Enzyme* 26:233-239 (1981).
 Saiki et al., *Bio/Technology* 3:1008-1012 (1985).
 Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Press, Cold Spring Harbor, N.Y.) (1989).
 Schmidt, R. D., *Clin. Chim. Acta.* 74:39-42 (1977).
 Schwartz et al., eds., *Atlas of Protein Sequence and Structure*, National Biomedical Research Foundation, pp. 357-358 (1979).
 Shulman H M, et al., *Hepatology* 1994; 19:1171-81.
 Smith et al., *Adv. Appl. Math.* 2:482 (1981).
 Stapleton et al., *Biochemistry* 35:14352-14361 (1996).
 Summar M L. *Journal of Inherited Metabolic Disease* 1998; 21 Suppl 1:30-9.

Summar, M., J. *Inherited Metabolic Disease* 21:30-39 (1998).
 Summar M L, et al., *Cytogenetics & Cell Genetics* 1995; 71:266-7.
 Takiguchi MaM, M., *Biochem J.* (1995) 312:649-659.
 Toh, H. et al., *European Journal of Biochemistry* 215:687-696 (1993).
 Tse et al., *American Journal of Hematology* 38:140-141 (1991).
 U.S. Pat. No. 5,399,346
 U.S. Pat. No. 5,162,215
 U.S. Pat. No. 5,279,833
 U.S. Pat. No. 5,643,567
 U.S. Pat. No. 4,683,202
 U.S. Pat. No. 4,683,195
 U.S. Pat. No. 5,286,634
 U.S. Pat. No. 5,646,008
 U.S. Pat. No. 4,196,265
 U.S. Pat. No. 5,489,742
 U.S. Pat. No. 5,550,316
 U.S. Pat. No. 5,573,933
 U.S. Pat. No. 5,614,396

U.S. Pat. No. 5,625,125
 U.S. Pat. No. 5,641,484
 U.S. Pat. No. 5,648,061
 U.S. Pat. No. 5,651,964
 5 U.S. Pat. No. 4,965,188
 U.S. Pat. No. 4,769,331
 U.S. Pat. No. 5,741,957
 U.S. Pat. No. 4,458,066
 U.S. Pat. No. 4,554,101
 10 U.S. Pat. No. 4,736,866
 van den Hoff, M. J. et al, *Journal of Molecular Evolution* 41:813-832 (1995).
 Vosatka R J, et al., *Biol Neonate* (1994) 66:65-70.
 Warter et al., *Revue Neurologique* 139:753-757 (1983).
 15 Wingard J R, et al. *Bone Marrow Transplantation* 1989; 4:685-9.
 Zamora, S. A., et al., *Crit Care Med* (1998) 26: 1271-1276.
 It will be understood that various details of the invention may be changed without departing from the scope of the invention. Furthermore, the foregoing description is for the purpose of illustration only, and not for the purpose of limitation—the invention being defined by the claims.
 20

SEQUENCE LISTING

<160> NUMBER OF SEQ ID NOS: 23

<210> SEQ ID NO 1

<211> LENGTH: 5761

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: CDS

<222> LOCATION: (124)..(4626)

<400> SEQUENCE: 1

```

gtcagcctta aacactgact gcaccctcc cagatttctt ttacattaac taaaaagtct      60
tatcacacaa tctcataaaa tttatgtaat ttcatttaat tttagccaca aatcatcttc      120
aaa atg acg agg att ttg aca gct ttc aaa gtg gtg agg aca ctg aag      168
  Met Thr Arg Ile Leu Thr Ala Phe Lys Val Val Arg Thr Leu Lys
    1             5             10             15
act ggt ttt ggc ttt acc aat gtg act gca cac caa aaa tgg aaa ttt      216
  Thr Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe
    20             25             30
tca aga cct ggc atc agg ctc ctt tct gtc aag gca cag aca gca cac      264
  Ser Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His
    35             40             45
att gtc ctg gaa gat gga act aag atg aaa ggt tac tcc ttt ggc cat      312
  Ile Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His
    50             55             60
cca tcc tct gtt gct ggt gaa gtg gtt ttt aat act ggc ctg gga ggg      360
  Pro Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly
    65             70             75
tac cca gaa gct att act gac cct gcc tac aaa gga cag att ctc aca      408
  Tyr Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr
    80             85             90             95
atg gcc aac cct att att ggg aat ggt gga gct cct gat act act gct      456
  Met Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala
    100            105            110
ctg gat gaa ctg gga ctt agc aaa tat ttg gag tct aat gga atc aag      504
  Leu Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys
    115            120            125
ggt tca ggt ttg ctg gtg ctg gat tat agt aaa gac tac aac cac tgg      552

```

-continued

Val	Ser	Gly	Leu	Leu	Val	Leu	Asp	Tyr	Ser	Lys	Asp	Tyr	Asn	His	Trp	
		130					135					140				
ctg	gct	acc	aag	agt	tta	ggg	caa	tgg	cta	cag	gaa	gaa	aag	ggt	cct	600
Leu	Ala	Thr	Lys	Ser	Leu	Gly	Gln	Trp	Leu	Gln	Glu	Glu	Lys	Val	Pro	
	145					150					155					
gca	att	tat	gga	gtg	gac	aca	aga	atg	ctg	act	aaa	ata	att	cgg	gat	648
Ala	Ile	Tyr	Gly	Val	Asp	Thr	Arg	Met	Leu	Thr	Lys	Ile	Ile	Arg	Asp	
160					165					170					175	
aag	ggt	acc	atg	ctt	ggg	aag	att	gaa	ttt	gaa	ggt	cag	cct	gtg	gat	696
Lys	Gly	Thr	Met	Leu	Gly	Lys	Ile	Glu	Phe	Glu	Gly	Gln	Pro	Val	Asp	
				180					185					190		
ttt	gtg	gat	cca	aat	aaa	cag	aat	ttg	att	gct	gag	ggt	tca	acc	aag	744
Phe	Val	Asp	Pro	Asn	Lys	Gln	Asn	Leu	Ile	Ala	Glu	Val	Ser	Thr	Lys	
			195					200						205		
gat	gtc	aaa	gtg	tac	ggc	aaa	gga	aac	ccc	aca	aaa	gtg	gta	gct	gta	792
Asp	Val	Lys	Val	Tyr	Gly	Lys	Gly	Asn	Pro	Thr	Lys	Val	Val	Ala	Val	
		210					215					220				
gac	tgt	ggg	att	aaa	aac	aat	gta	atc	cgc	ctg	cta	gta	aag	cga	gga	840
Asp	Cys	Gly	Ile	Lys	Asn	Asn	Val	Ile	Arg	Leu	Leu	Val	Lys	Arg	Gly	
	225					230					235					
gct	gaa	gtg	cac	tta	ggt	ccc	tgg	aac	cat	gat	ttc	acc	aag	atg	gag	888
Ala	Glu	Val	His	Leu	Val	Pro	Trp	Asn	His	Asp	Phe	Thr	Lys	Met	Glu	
240					245					250					255	
tat	gat	ggg	att	ttg	atc	gcg	gga	gga	ccg	ggg	aac	cca	gct	ctt	gca	936
Tyr	Asp	Gly	Ile	Leu	Ile	Ala	Gly	Gly	Pro	Gly	Asn	Pro	Ala	Leu	Ala	
				260					265					270		
gaa	cca	cta	att	cag	aat	gtc	aga	aag	att	ttg	gag	agt	gat	cgc	aag	984
Glu	Pro	Leu	Ile	Gln	Asn	Val	Arg	Lys	Ile	Leu	Glu	Ser	Asp	Arg	Lys	
			275					280						285		
gag	cca	ttg	ttt	gga	atc	agt	aca	gga	aac	tta	ata	aca	gga	ttg	gct	1032
Glu	Pro	Leu	Phe	Gly	Ile	Ser	Thr	Gly	Asn	Leu	Ile	Thr	Gly	Leu	Ala	
		290					295					300				
gct	ggt	gcc	aaa	acc	tac	aag	atg	tcc	atg	gcc	aac	aga	ggg	cag	aat	1080
Ala	Gly	Ala	Lys	Thr	Tyr	Lys	Met	Ser	Met	Ala	Asn	Arg	Gly	Gln	Asn	
	305					310						315				
cag	cct	ggt	ttg	aat	atc	aca	aac	aaa	cag	gct	ttc	att	act	gct	cag	1128
Gln	Pro	Val	Leu	Asn	Ile	Thr	Asn	Lys	Gln	Ala	Phe	Ile	Thr	Ala	Gln	
320					325					330					335	
aat	cat	ggc	tat	gcc	ttg	gac	aac	acc	ctc	cct	gct	ggc	tgg	aaa	cca	1176
Asn	His	Gly	Tyr	Ala	Leu	Asp	Asn	Thr	Leu	Pro	Ala	Gly	Trp	Lys	Pro	
				340					345					350		
ctt	ttt	gtg	aat	gtc	aac	gat	caa	aca	aat	gag	ggg	att	atg	cat	gag	1224
Leu	Phe	Val	Asn	Val	Asn	Asp	Gln	Thr	Asn	Glu	Gly	Ile	Met	His	Glu	
			355					360						365		
agc	aaa	ccc	ttc	ttc	gct	gtg	cag	ttc	cac	cca	gag	gtc	acc	ccg	ggg	1272
Ser	Lys	Pro	Phe	Phe	Ala	Val	Gln	Phe	His	Pro	Glu	Val	Thr	Pro	Gly	
	370					375						380				
cca	ata	gac	act	gag	tac	ctg	ttt	gat	tcc	ttt	ttc	tca	ctg	ata	aag	1320
Pro	Ile	Asp	Thr	Glu	Tyr	Leu	Phe	Asp	Ser	Phe	Phe	Ser	Leu	Ile	Lys	
	385					390						395				
aaa	gga	aaa	gct	acc	acc	att	aca	tca	gtc	tta	ccg	aag	cca	gca	cta	1368
Lys	Gly	Lys	Ala	Thr	Thr	Ile	Thr	Ser	Val	Leu	Pro	Lys	Pro	Ala	Leu	
400					405					410					415	
ggt	gca	tct	cgg	ggt	gag	ggt	tcc	aaa	gtc	ctt	att	cta	gga	tca	gga	1416
Val	Ala	Ser	Arg	Val	Glu	Val	Ser	Lys	Val	Leu	Ile	Leu	Gly	Ser	Gly	
				420					425					430		
ggt	ctg	tcc	att	ggt	cag	gct	gga	gaa	ttt	gat	tac	tca	gga	tct	caa	1464
Gly	Leu	Ser	Ile	Gly	Gln	Ala	Gly	Glu	Phe	Asp	Tyr	Ser	Gly	Ser	Gln	
			435					440					445			
gct	gta	aaa	gcc	atg	aag	gaa	gaa	aat	gtc	aaa	act	ggt	ctg	atg	aac	1512

-continued

Ala	Val	Lys	Ala	Met	Lys	Glu	Glu	Asn	Val	Lys	Thr	Val	Leu	Met	Asn		
		450					455					460					
cca	aac	att	gca	tca	gtc	cag	acc	aat	gag	gtg	ggc	tta	aag	caa	gcg	1560	
Pro	Asn	Ile	Ala	Ser	Val	Gln	Thr	Asn	Glu	Val	Gly	Leu	Lys	Gln	Ala		
		465				470					475						
gat	act	gtc	tac	ttt	ctt	ccc	atc	acc	cct	cag	ttt	gtc	aca	gag	gtc	1608	
Asp	Thr	Val	Tyr	Phe	Leu	Pro	Ile	Thr	Pro	Gln	Phe	Val	Thr	Glu	Val		
		480			485					490					495		
atc	aag	gca	gaa	cag	cca	gat	ggg	tta	att	ctg	ggc	atg	ggt	ggc	cag	1656	
Ile	Lys	Ala	Glu	Gln	Pro	Asp	Gly	Leu	Ile	Leu	Gly	Met	Gly	Gly	Gln		
				500				505							510		
aca	gct	ctg	aac	tgt	gga	gtg	gaa	cta	ttc	aag	aga	ggt	gtg	ctc	aag	1704	
Thr	Ala	Leu	Asn	Cys	Gly	Val	Glu	Leu	Phe	Lys	Arg	Gly	Val	Leu	Lys		
			515					520						525			
gaa	tat	ggt	gtg	aaa	gtc	ctg	gga	act	tca	ggt	gag	tcc	att	atg	gct	1752	
Glu	Tyr	Gly	Val	Lys	Val	Leu	Gly	Thr	Ser	Val	Glu	Ser	Ile	Met	Ala		
		530					535					540					
acg	gaa	gac	agg	cag	ctg	ttt	tca	gat	aaa	cta	aat	gag	atc	aat	gaa	1800	
Thr	Glu	Asp	Arg	Gln	Leu	Phe	Ser	Asp	Lys	Leu	Asn	Glu	Ile	Asn	Glu		
		545				550					555						
aag	att	gct	cca	agt	ttt	gca	gtg	gaa	tcg	att	gag	gat	gca	ctg	aag	1848	
Lys	Ile	Ala	Pro	Ser	Phe	Ala	Val	Glu	Ser	Ile	Glu	Asp	Ala	Leu	Lys		
		560			565					570					575		
gca	gca	gac	acc	att	ggc	tac	cca	gtg	atg	atc	cgt	tcc	gcc	tat	gca	1896	
Ala	Ala	Asp	Thr	Ile	Gly	Tyr	Pro	Val	Met	Ile	Arg	Ser	Ala	Tyr	Ala		
				580					585						590		
ctg	ggt	ggg	tta	ggc	tca	ggc	atc	tgt	ccc	aac	aga	gag	act	ttg	atg	1944	
Leu	Gly	Gly	Leu	Gly	Ser	Gly	Ile	Cys	Pro	Asn	Arg	Glu	Thr	Leu	Met		
			595					600						605			
gac	ctc	agc	aca	aag	gcc	ttt	gct	atg	acc	aac	caa	att	ctg	gtg	gag	1992	
Asp	Leu	Ser	Thr	Lys	Ala	Phe	Ala	Met	Thr	Asn	Gln	Ile	Leu	Val	Glu		
		610					615						620				
aag	tca	gtg	aca	ggt	tgg	aaa	gaa	ata	gaa	tat	gaa	gtg	ggt	cga	gat	2040	
Lys	Ser	Val	Thr	Gly	Trp	Lys	Glu	Ile	Glu	Tyr	Glu	Val	Val	Arg	Asp		
		625				630					635						
gct	gat	gac	aat	tgt	gtc	act	gtc	tgt	aac	atg	gaa	aat	ggt	gat	gcc	2088	
Ala	Asp	Asp	Asn	Cys	Val	Thr	Val	Cys	Asn	Met	Glu	Asn	Val	Asp	Ala		
				645						650					655		
atg	ggt	ggt	cac	aca	ggt	gac	tca	ggt	ggt	gtg	gct	cct	gcc	cag	aca	2136	
Met	Gly	Val	His	Thr	Gly	Asp	Ser	Val	Val	Val	Ala	Pro	Ala	Gln	Thr		
				660						665					670		
ctc	tcc	aat	gcc	gag	ttt	cag	atg	ttg	aga	cgt	act	tca	atc	aat	ggt	2184	
Leu	Ser	Asn	Ala	Glu	Phe	Gln	Met	Leu	Arg	Arg	Thr	Ser	Ile	Asn	Val		
			675					680							685		
ggt	cgc	cac	ttg	ggc	att	gtg	ggt	gaa	tgc	aac	att	cag	ttt	gcc	ctt	2232	
Val	Arg	His	Leu	Gly	Ile	Val	Gly	Glu	Cys	Asn	Ile	Gln	Phe	Ala	Leu		
		690					695								700		
cat	cct	acc	tca	atg	gaa	tac	tgc	atc	att	gaa	gtg	aat	gcc	aga	ctg	2280	
His	Pro	Thr	Ser	Met	Glu	Tyr	Cys	Ile	Ile	Glu	Val	Asn	Ala	Arg	Leu		
		705				710						715					
tcc	cga	agc	tct	gct	ctg	gcc	tca	aaa	gcc	act	ggc	tac	cca	ttg	gca	2328	
Ser	Arg	Ser	Ser	Ala	Leu	Ala	Ser	Lys	Ala	Thr	Gly	Tyr	Pro	Leu	Ala		
				725						730					735		
ttc	att	gct	gca	aag	att	gcc	cta	gga	atc	cca	ctt	cca	gaa	att	aag	2376	
Phe	Ile	Ala	Ala	Lys	Ile	Ala	Leu	Gly	Ile	Pro	Leu	Pro	Glu	Ile	Lys		
				740					745						750		
aac	gtc	gta	tcc	ggg	aag	aca	tca	gcc	tgt	ttt	gaa	cct	agc	ctg	gat	2424	
Asn	Val	Val	Ser	Gly	Lys	Thr	Ser	Ala	Cys	Phe	Glu	Pro	Ser	Leu	Asp		
			755					760							765		
tac	atg	gtc	acc	aag	att	ccc	cgc	tgg	gat	ctt	gac	cgt	ttt	cat	gga	2472	

-continued

Tyr	Met	Val	Thr	Lys	Ile	Pro	Arg	Trp	Asp	Leu	Asp	Arg	Phe	His	Gly		
		770					775					780					
aca	tct	agc	cga	att	ggt	agc	tct	atg	aaa	agt	gta	gga	gag	gtc	atg	2520	
Thr	Ser	Ser	Arg	Ile	Gly	Ser	Ser	Met	Lys	Ser	Val	Gly	Glu	Val	Met		
	785					790					795						
gct	att	ggt	cgt	acc	ttt	gag	gag	agt	ttc	cag	aaa	gct	tta	cgg	atg	2568	
Ala	Ile	Gly	Arg	Thr	Phe	Glu	Glu	Ser	Phe	Gln	Lys	Ala	Leu	Arg	Met		
800					805					810					815		
tgc	cac	cca	tct	ata	gaa	ggt	ttc	act	ccc	cgt	ctc	cca	atg	aac	aaa	2616	
Cys	His	Pro	Ser	Ile	Glu	Gly	Phe	Thr	Pro	Arg	Leu	Pro	Met	Asn	Lys		
				820					825						830		
gaa	tgg	cca	tct	aat	tta	gat	ctt	aga	aaa	gag	ttg	tct	gaa	cca	agc	2664	
Glu	Trp	Pro	Ser	Asn	Leu	Asp	Leu	Arg	Lys	Glu	Leu	Ser	Glu	Pro	Ser		
			835					840						845			
agc	acg	cgt	atc	tat	gcc	att	gcc	aag	gcc	att	gat	gac	aac	atg	tcc	2712	
Ser	Thr	Arg	Ile	Tyr	Ala	Ile	Ala	Lys	Ala	Ile	Asp	Asp	Asn	Met	Ser		
		850					855						860				
ctt	gat	gag	att	gag	aag	ctc	aca	tac	att	gac	aag	tgg	ttt	ttg	tat	2760	
Leu	Asp	Glu	Ile	Glu	Lys	Leu	Thr	Tyr	Ile	Asp	Lys	Trp	Phe	Leu	Tyr		
	865					870					875						
aag	atg	cgt	gat	att	tta	aac	atg	gaa	aag	aca	ctg	aaa	ggg	ctc	aac	2808	
Lys	Met	Arg	Asp	Ile	Leu	Asn	Met	Glu	Lys	Thr	Leu	Lys	Gly	Leu	Asn		
880					885					890					895		
agt	gag	tcc	atg	aca	gaa	gaa	acc	ctg	aaa	agg	gca	aag	gag	att	ggg	2856	
Ser	Glu	Ser	Met	Thr	Glu	Glu	Thr	Leu	Lys	Arg	Ala	Lys	Glu	Ile	Gly		
				900					905						910		
ttc	tca	gat	aag	cag	att	tca	aaa	tgc	ctt	ggg	ctc	act	gag	gcc	cag	2904	
Phe	Ser	Asp	Lys	Gln	Ile	Ser	Lys	Cys	Leu	Gly	Leu	Thr	Glu	Ala	Gln		
			915					920						925			
aca	agg	gag	ctg	agg	tta	aag	aaa	aac	atc	cac	cct	tgg	ggt	aaa	cag	2952	
Thr	Arg	Glu	Leu	Arg	Leu	Lys	Lys	Asn	Ile	His	Pro	Trp	Val	Lys	Gln		
		930					935							940			
att	gat	aca	ctg	gct	gca	gaa	tac	cca	tca	gta	aca	aac	tat	ctc	tat	3000	
Ile	Asp	Thr	Leu	Ala	Ala	Glu	Tyr	Pro	Ser	Val	Thr	Asn	Tyr	Leu	Tyr		
	945					950						955					
gtt	acc	tac	aat	ggt	cag	gag	cat	gat	gtc	aat	ttt	gat	gac	cat	gga	3048	
Val	Thr	Tyr	Asn	Gly	Gln	Glu	His	Asp	Val	Asn	Phe	Asp	Asp	His	Gly		
960					965				970						975		
atg	atg	gtg	cta	ggc	tgt	ggt	cca	tat	cac	att	ggc	agc	agt	gtg	gaa	3096	
Met	Met	Val	Leu	Gly	Cys	Gly	Pro	Tyr	His	Ile	Gly	Ser	Ser	Val	Glu		
				980					985						990		
ttt	gat	tgg	tgt	gct	gtc	tct	agt	atc	cgc	aca	ctg	cgt	caa	ctt	ggc	3144	
Phe	Asp	Trp	Cys	Ala	Val	Ser	Ser	Ile	Arg	Thr	Leu	Arg	Gln	Leu	Gly		
			995					1000						1005			
aag	aag	acg	gtg	gtg	gtg	aat	tgc	aat	cct	gag	act	gtg	agc	aca		3189	
Lys	Lys	Thr	Val	Val	Val	Asn	Cys	Asn	Pro	Glu	Thr	Val	Ser	Thr			
			1010				1015						1020				
gac	ttt	gat	gag	tgt	gac	aaa	ctg	tac	ttt	gaa	gag	ttg	tcc	ttg		3234	
Asp	Phe	Asp	Glu	Cys	Asp	Lys	Leu	Tyr	Phe	Glu	Glu	Leu	Ser	Leu			
		1025					1030						1035				
gag	aga	atc	cta	gac	atc	tac	cat	cag	gag	gca	tgt	ggt	ggc	tgc		3279	
Glu	Arg	Ile	Leu	Asp	Ile	Tyr	His	Gln	Glu	Ala	Cys	Gly	Gly	Cys			
		1040					1045						1050				
atc	ata	tca	ggt	gga	ggc	cag	att	cca	aac	aac	ctg	gca	ggt	cct		3324	
Ile	Ile	Ser	Val	Gly	Gly	Gln	Ile	Pro	Asn	Asn	Leu	Ala	Val	Pro			
		1055					1060						1065				
cta	tac	aag	aat	ggt	gtc	aag	atc	atg	ggc	aca	agc	ccc	ctg	cag		3369	
Leu	Tyr	Lys	Asn	Gly	Val	Lys	Ile	Met	Gly	Thr	Ser	Pro	Leu	Gln			
		1070					1075						1080				
atc	gac	agg	gct	gag	gat	cgc	tcc	atc	ttc	tca	gct	gtc	ttg	gat		3414	

-continued

Ile Asp Arg	Ala Glu Asp Arg	Ser	Ile Phe Ser	Ala Val	Leu Asp	
1085		1090		1095		
gag ctg aag	gtg gct cag gca cct	tgg aaa gct gtt aat	act ttg	3459		
Glu Leu Lys	Val Ala Gln Ala Pro	Trp Lys Ala Val Asn	Thr Leu			
1100		1105	1110			
aat gaa gca	ctg gaa ttt gca aag	tct gtg gac tac ccc	tgc ttg	3504		
Asn Glu Ala	Leu Glu Phe Ala Lys	Ser Val Asp Tyr Pro	Cys Leu			
1115		1120	1125			
ttg agg cct	tcc tat gtt ttg agt	ggg tct gct atg aat	gtg gta	3549		
Leu Arg Pro	Ser Tyr Val Leu Ser	Gly Ser Ala Met Asn	Val Val			
1130		1135	1140			
ttc tct gag	gat gag atg aaa aaa	ttc cta gaa gag gcg	act aga	3594		
Phe Ser Glu	Asp Glu Met Lys Lys	Phe Leu Glu Glu Ala	Thr Arg			
1145		1150	1155			
gtt tct cag	gag cac cca gtg gtc	ctg aca aaa ttt gtt	gaa ggg	3639		
Val Ser Gln	Glu His Pro Val Val	Leu Thr Lys Phe Val	Glu Gly			
1160		1165	1170			
gcc cga gaa	gta gaa atg gac gct	gtt ggc aaa gat gga	agg gtt	3684		
Ala Arg Glu	Val Glu Met Asp Ala	Val Gly Lys Asp Gly	Arg Val			
1175		1180	1185			
atc tct cat	gcc atc tct gaa cat	gtt gaa gat gca ggt	gtc cac	3729		
Ile Ser His	Ala Ile Ser Glu His	Val Glu Asp Ala Gly	Val His			
1190		1195	1200			
tcg gga gat	gcc act ctg atg ctg	ccc aca caa acc atc	agc caa	3774		
Ser Gly Asp	Ala Thr Leu Met Leu	Pro Thr Gln Thr Ile	Ser Gln			
1205		1210	1215			
ggg gcc att	gaa aag gtg aag gat	gct acc cgg aag att	gca aag	3819		
Gly Ala Ile	Glu Lys Val Lys Asp	Ala Thr Arg Lys Ile	Ala Lys			
1220		1225	1230			
gct ttt gcc	atc tct ggt cca ttc	aac gtc caa ttt ctt	gtc aaa	3864		
Ala Phe Ala	Ile Ser Gly Pro Phe	Asn Val Gln Phe Leu	Val Lys			
1235		1240	1245			
gga aat gat	gtc ttg gtg att gag	tgt aac ttg aga gct	tct cga	3909		
Gly Asn Asp	Val Leu Val Ile Glu	Cys Asn Leu Arg Ala	Ser Arg			
1250		1255	1260			
tcc ttc ccc	ttt gtt tcc aag act	ctt ggg gtt gac ttc	att gat	3954		
Ser Phe Pro	Phe Val Ser Lys Thr	Leu Gly Val Asp Phe	Ile Asp			
1265		1270	1275			
gtg gcc acc	aag gtg atg att gga	gag aat gtt gat gag	aaa cat	3999		
Val Ala Thr	Lys Val Met Ile Gly	Glu Asn Val Asp Glu	Lys His			
1280		1285	1290			
ctt cca aca	ttg gac cat ccc ata	att cct gct gac tat	gtt gca	4044		
Leu Pro Thr	Leu Asp His Pro Ile	Ile Pro Ala Asp Tyr	Val Ala			
1295		1300	1305			
att aag gct	ccc atg ttt tcc tgg	ccc cgg ttg agg gat	gct gac	4089		
Ile Lys Ala	Pro Met Phe Ser Trp	Pro Arg Leu Arg Asp	Ala Asp			
1310		1315	1320			
ccc att ctg	aga tgt gag atg gct	tcc act gga gag gtg	gct tgc	4134		
Pro Ile Leu	Arg Cys Glu Met Ala	Ser Thr Gly Glu Val	Ala Cys			
1325		1330	1335			
ttt ggt gaa	ggt att cat aca gcc	ttc cta aag gca atg	ctt tcc	4179		
Phe Gly Glu	Gly Ile His Thr Ala	Phe Leu Lys Ala Met	Leu Ser			
1340		1345	1350			
aca gga ttt	aag ata ccc cag aaa	ggc atc ctg ata ggc	atc cag	4224		
Thr Gly Phe	Lys Ile Pro Gln Lys	Gly Ile Leu Ile Gly	Ile Gln			
1355		1360	1365			
caa tca ttc	cgg cca aga ttc ctt	ggt gtg gct gaa caa	tta cac	4269		
Gln Ser Phe	Arg Pro Arg Phe Leu	Gly Val Ala Glu Gln	Leu His			
1370		1375	1380			
aat gaa ggt	ttc aag ctg ttt gcc	acg gaa gcc aca tca	gac tgg	4314		

-continued

Asn	Glu	Gly	Phe	Lys	Leu	Phe	Ala	Thr	Glu	Ala	Thr	Ser	Asp	Trp	
		1385					1390					1395			
ctc	aac	gcc	aac	aat	gtc	cct	gcc	aac	cca	gtg	gca	tgg	ccg	tct	4359
Leu	Asn	Ala	Asn	Asn	Val	Pro	Ala	Asn	Pro	Val	Ala	Trp	Pro	Ser	
		1400					1405					1410			
caa	gaa	gga	cag	aat	ccc	agc	ctc	tct	tcc	atc	aga	aaa	ttg	att	4404
Gln	Glu	Gly	Gln	Asn	Pro	Ser	Leu	Ser	Ser	Ile	Arg	Lys	Leu	Ile	
		1415					1420					1425			
aga	gat	ggc	agc	att	gac	cta	gtg	att	aac	ctt	ccc	aac	aac	aac	4449
Arg	Asp	Gly	Ser	Ile	Asp	Leu	Val	Ile	Asn	Leu	Pro	Asn	Asn	Asn	
		1430					1435					1440			
act	aaa	ttt	gtc	cat	gat	aat	tat	gtg	att	cgg	agg	aca	gct	gtt	4494
Thr	Lys	Phe	Val	His	Asp	Asn	Tyr	Val	Ile	Arg	Arg	Thr	Ala	Val	
		1445					1450					1455			
gat	agt	gga	atc	cct	ctc	ctc	act	aat	ttt	cag	gtg	acc	aaa	ctt	4539
Asp	Ser	Gly	Ile	Pro	Leu	Leu	Thr	Asn	Phe	Gln	Val	Thr	Lys	Leu	
		1460					1465					1470			
ttt	gct	gaa	gct	gtg	cag	aaa	tct	cgc	aag	gtg	gac	tcc	aag	agt	4584
Phe	Ala	Glu	Ala	Val	Gln	Lys	Ser	Arg	Lys	Val	Asp	Ser	Lys	Ser	
		1475					1480					1485			
ctt	ttc	cac	tac	agg	cag	tac	agt	gct	gga	aaa	gca	gca	tag		4626
Leu	Phe	His	Tyr	Arg	Gln	Tyr	Ser	Ala	Gly	Lys	Ala	Ala			
		1490					1495					1500			
agatgcagac	accccagccc	cattattaa	tcaacctgag	ccacatgtta	tctaaaggaa										4686
ctgattcaca	actttctcag	agatgaatat	tgataactaa	acttcatttc	agtttacttt										4746
gttatgcctt	aatattctgt	gtcttttgca	attaaattgt	cagtcacttc	ttcaaacct										4806
tacagtcctt	cctaagttac	tcttcatgag	atttcatcca	tttactaata	ctgtattttt										4866
ggtaggactag	gcttgccctat	gtgcttatgt	gtagcttttt	actttttatg	gtgctgatta										4926
atgggtgatca	aggtaggaaa	agttgctggt	ctattttctg	aactctttct	atactttaag										4986
atactctatt	tttaaacac	tatctgcaaa	ctcaggacac	tttaacaggg	cagaatactc										5046
taaaaacttg	ataaatgaa	atatagattt	aatttatgaa	ccttccatca	tgatgtttgt										5106
gtattgcttc	tttttgatc	ctcattctca	cccatttggc	taatccagga	atattgttat										5166
cccttcccat	tatattgaag	ttgagaaatg	tgacagaggc	atttagagta	tggacttttc										5226
ttttcttttt	cttttcttt	ttttcttttt	gagatggagt	cacactctcc	aggetggagt										5286
gcagtggcac	aatctcggct	cactgcaatt	tgctctccc	aagttcaagc	gattctcctg										5346
ctttagacta	tggatttctt	taaggaatac	tggtttgcag	ttttgttttc	tggactatat										5406
cagcagatgg	tagacagtgt	ttatgtagat	gtgttgttgt	ttttatcatt	ggattttaac										5466
ttggcccagag	tgaataatc	agatttttgt	cattcacact	ctccccagt	tttgaataa										5526
cttgaagta	aggttcattc	ccttaagacg	atggattctg	ttgaactatg	gggtcccaca										5586
ctgcactatt	aattccacc	actgtaagg	caaggacacc	attcctteta	catataagaa										5646
aaaagtctct	ccccagggc	agcctttgtt	acttttaaat	atcttctgtt	attacaagtg										5706
ctctaattgt	gaacttttaa	ataaaatact	attaagaggt	aaaaaaaaa	aaaaa										5761

<210> SEQ ID NO 2
 <211> LENGTH: 1500
 <212> TYPE: PRT
 <213> ORGANISM: Homo sapiens

<400> SEQUENCE: 2

Met Thr Arg Ile Leu Thr Ala Phe Lys Val Val Arg Thr Leu Lys Thr
 1 5 10 15

-continued

Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe Ser
 20 25 30
 Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His Ile
 35 40 45
 Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His Pro
 50 55 60
 Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly Tyr
 65 70 75 80
 Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr Met
 85 90 95
 Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala Leu
 100 105 110
 Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys Val
 115 120 125
 Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp Leu
 130 135 140
 Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro Ala
 145 150 155 160
 Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp Lys
 165 170 175
 Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp Phe
 180 185 190
 Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys Asp
 195 200 205
 Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val Asp
 210 215 220
 Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly Ala
 225 230 235 240
 Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu Tyr
 245 250 255
 Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala Glu
 260 265 270
 Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys Glu
 275 280 285
 Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala Ala
 290 295 300
 Gly Ala Lys Thr Tyr Lys Met Ser Met Ala Asn Arg Gly Gln Asn Gln
 305 310 315 320
 Pro Val Leu Asn Ile Thr Asn Lys Gln Ala Phe Ile Thr Ala Gln Asn
 325 330 335
 His Gly Tyr Ala Leu Asp Asn Thr Leu Pro Ala Gly Trp Lys Pro Leu
 340 345 350
 Phe Val Asn Val Asn Asp Gln Thr Asn Glu Gly Ile Met His Glu Ser
 355 360 365
 Lys Pro Phe Phe Ala Val Gln Phe His Pro Glu Val Thr Pro Gly Pro
 370 375 380
 Ile Asp Thr Glu Tyr Leu Phe Asp Ser Phe Phe Ser Leu Ile Lys Lys
 385 390 395 400
 Gly Lys Ala Thr Thr Ile Thr Ser Val Leu Pro Lys Pro Ala Leu Val
 405 410 415
 Ala Ser Arg Val Glu Val Ser Lys Val Leu Ile Leu Gly Ser Gly Gly
 420 425 430
 Leu Ser Ile Gly Gln Ala Gly Glu Phe Asp Tyr Ser Gly Ser Gln Ala
 435 440 445

-continued

Val Lys Ala Met Lys Glu Glu Asn Val Lys Thr Val Leu Met Asn Pro
 450 455 460

Asn Ile Ala Ser Val Gln Thr Asn Glu Val Gly Leu Lys Gln Ala Asp
 465 470 475 480

Thr Val Tyr Phe Leu Pro Ile Thr Pro Gln Phe Val Thr Glu Val Ile
 485 490 495

Lys Ala Glu Gln Pro Asp Gly Leu Ile Leu Gly Met Gly Gly Gln Thr
 500 505 510

Ala Leu Asn Cys Gly Val Glu Leu Phe Lys Arg Gly Val Leu Lys Glu
 515 520 525

Tyr Gly Val Lys Val Leu Gly Thr Ser Val Glu Ser Ile Met Ala Thr
 530 535 540

Glu Asp Arg Gln Leu Phe Ser Asp Lys Leu Asn Glu Ile Asn Glu Lys
 545 550 555 560

Ile Ala Pro Ser Phe Ala Val Glu Ser Ile Glu Asp Ala Leu Lys Ala
 565 570 575

Ala Asp Thr Ile Gly Tyr Pro Val Met Ile Arg Ser Ala Tyr Ala Leu
 580 585 590

Gly Gly Leu Gly Ser Gly Ile Cys Pro Asn Arg Glu Thr Leu Met Asp
 595 600 605

Leu Ser Thr Lys Ala Phe Ala Met Thr Asn Gln Ile Leu Val Glu Lys
 610 615 620

Ser Val Thr Gly Trp Lys Glu Ile Glu Tyr Glu Val Val Arg Asp Ala
 625 630 635 640

Asp Asp Asn Cys Val Thr Val Cys Asn Met Glu Asn Val Asp Ala Met
 645 650 655

Gly Val His Thr Gly Asp Ser Val Val Val Ala Pro Ala Gln Thr Leu
 660 665 670

Ser Asn Ala Glu Phe Gln Met Leu Arg Arg Thr Ser Ile Asn Val Val
 675 680 685

Arg His Leu Gly Ile Val Gly Glu Cys Asn Ile Gln Phe Ala Leu His
 690 695 700

Pro Thr Ser Met Glu Tyr Cys Ile Ile Glu Val Asn Ala Arg Leu Ser
 705 710 715 720

Arg Ser Ser Ala Leu Ala Ser Lys Ala Thr Gly Tyr Pro Leu Ala Phe
 725 730 735

Ile Ala Ala Lys Ile Ala Leu Gly Ile Pro Leu Pro Glu Ile Lys Asn
 740 745 750

Val Val Ser Gly Lys Thr Ser Ala Cys Phe Glu Pro Ser Leu Asp Tyr
 755 760 765

Met Val Thr Lys Ile Pro Arg Trp Asp Leu Asp Arg Phe His Gly Thr
 770 775 780

Ser Ser Arg Ile Gly Ser Ser Met Lys Ser Val Gly Glu Val Met Ala
 785 790 795 800

Ile Gly Arg Thr Phe Glu Glu Ser Phe Gln Lys Ala Leu Arg Met Cys
 805 810 815

His Pro Ser Ile Glu Gly Phe Thr Pro Arg Leu Pro Met Asn Lys Glu
 820 825 830

Trp Pro Ser Asn Leu Asp Leu Arg Lys Glu Leu Ser Glu Pro Ser Ser
 835 840 845

Thr Arg Ile Tyr Ala Ile Ala Lys Ala Ile Asp Asp Asn Met Ser Leu
 850 855 860

Asp Glu Ile Glu Lys Leu Thr Tyr Ile Asp Lys Trp Phe Leu Tyr Lys

-continued

865	870	875	880
Met Arg Asp Ile Leu Asn Met Glu Lys Thr Leu Lys Gly Leu Asn Ser	885	890	895
Glu Ser Met Thr Glu Glu Thr Leu Lys Arg Ala Lys Glu Ile Gly Phe	900	905	910
Ser Asp Lys Gln Ile Ser Lys Cys Leu Gly Leu Thr Glu Ala Gln Thr	915	920	925
Arg Glu Leu Arg Leu Lys Lys Asn Ile His Pro Trp Val Lys Gln Ile	930	935	940
Asp Thr Leu Ala Ala Glu Tyr Pro Ser Val Thr Asn Tyr Leu Tyr Val	945	950	955
Thr Tyr Asn Gly Gln Glu His Asp Val Asn Phe Asp Asp His Gly Met	965	970	975
Met Val Leu Gly Cys Gly Pro Tyr His Ile Gly Ser Ser Val Glu Phe	980	985	990
Asp Trp Cys Ala Val Ser Ser Ile Arg Thr Leu Arg Gln Leu Gly Lys	995	1000	1005
Lys Thr Val Val Val Asn Cys Asn Pro Glu Thr Val Ser Thr Asp	1010	1015	1020
Phe Asp Glu Cys Asp Lys Leu Tyr Phe Glu Glu Leu Ser Leu Glu	1025	1030	1035
Arg Ile Leu Asp Ile Tyr His Gln Glu Ala Cys Gly Gly Cys Ile	1040	1045	1050
Ile Ser Val Gly Gly Gln Ile Pro Asn Asn Leu Ala Val Pro Leu	1055	1060	1065
Tyr Lys Asn Gly Val Lys Ile Met Gly Thr Ser Pro Leu Gln Ile	1070	1075	1080
Asp Arg Ala Glu Asp Arg Ser Ile Phe Ser Ala Val Leu Asp Glu	1085	1090	1095
Leu Lys Val Ala Gln Ala Pro Trp Lys Ala Val Asn Thr Leu Asn	1100	1105	1110
Glu Ala Leu Glu Phe Ala Lys Ser Val Asp Tyr Pro Cys Leu Leu	1115	1120	1125
Arg Pro Ser Tyr Val Leu Ser Gly Ser Ala Met Asn Val Val Phe	1130	1135	1140
Ser Glu Asp Glu Met Lys Lys Phe Leu Glu Glu Ala Thr Arg Val	1145	1150	1155
Ser Gln Glu His Pro Val Val Leu Thr Lys Phe Val Glu Gly Ala	1160	1165	1170
Arg Glu Val Glu Met Asp Ala Val Gly Lys Asp Gly Arg Val Ile	1175	1180	1185
Ser His Ala Ile Ser Glu His Val Glu Asp Ala Gly Val His Ser	1190	1195	1200
Gly Asp Ala Thr Leu Met Leu Pro Thr Gln Thr Ile Ser Gln Gly	1205	1210	1215
Ala Ile Glu Lys Val Lys Asp Ala Thr Arg Lys Ile Ala Lys Ala	1220	1225	1230
Phe Ala Ile Ser Gly Pro Phe Asn Val Gln Phe Leu Val Lys Gly	1235	1240	1245
Asn Asp Val Leu Val Ile Glu Cys Asn Leu Arg Ala Ser Arg Ser	1250	1255	1260
Phe Pro Phe Val Ser Lys Thr Leu Gly Val Asp Phe Ile Asp Val	1265	1270	1275

-continued

Ala Thr	Lys Val Met Ile Gly	Glu Asn Val Asp	Glu Lys His Leu
1280	1285		1290
Pro Thr	Leu Asp His Pro Ile	Ile Pro Ala Asp Tyr	Val Ala Ile
1295	1300		1305
Lys Ala	Pro Met Phe Ser Trp	Pro Arg Leu Arg Asp	Ala Asp Pro
1310	1315		1320
Ile Leu	Arg Cys Glu Met Ala	Ser Thr Gly Glu Val	Ala Cys Phe
1325	1330		1335
Gly Glu	Gly Ile His Thr Ala	Phe Leu Lys Ala Met	Leu Ser Thr
1340	1345		1350
Gly Phe	Lys Ile Pro Gln Lys	Gly Ile Leu Ile Gly	Ile Gln Gln
1355	1360		1365
Ser Phe	Arg Pro Arg Phe Leu	Gly Val Ala Glu Gln	Leu His Asn
1370	1375		1380
Glu Gly	Phe Lys Leu Phe Ala	Thr Glu Ala Thr Ser	Asp Trp Leu
1385	1390		1395
Asn Ala	Asn Asn Val Pro Ala	Asn Pro Val Ala Trp	Pro Ser Gln
1400	1405		1410
Glu Gly	Gln Asn Pro Ser Leu	Ser Ser Ile Arg Lys	Leu Ile Arg
1415	1420		1425
Asp Gly	Ser Ile Asp Leu Val	Ile Asn Leu Pro Asn	Asn Asn Thr
1430	1435		1440
Lys Phe	Val His Asp Asn Tyr	Val Ile Arg Arg Thr	Ala Val Asp
1445	1450		1455
Ser Gly	Ile Pro Leu Leu Thr	Asn Phe Gln Val Thr	Lys Leu Phe
1460	1465		1470
Ala Glu	Ala Val Gln Lys Ser	Arg Lys Val Asp Ser	Lys Ser Leu
1475	1480		1485
Phe His	Tyr Arg Gln Tyr Ser	Ala Gly Lys Ala Ala	
1490	1495		1500

<210> SEQ ID NO 3
 <211> LENGTH: 5761
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (124)..(4626)

<400> SEQUENCE: 3

gtcagcctta aacctgact gcaccctcc cagatttctt ttacattaac taaaagtct	60
tatcacacaa tctcataaaa tttatgtaat ttcatttaat tttagccaca aatcatcttc	120
aaa atg acg agg att ttg aca gct ttc aaa gtg gtg agg aca ctg aag	168
Met Thr Arg Ile Leu Thr Ala Phe Lys Val Val Arg Thr Leu Lys	
1 5 10 15	
act ggt ttt ggc ttt acc aat gtg act gca cac caa aaa tgg aaa ttt	216
Thr Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe	
20 25 30	
tca aga cct ggc atc agg ctc ctt tct gtc aag gca cag aca gca cac	264
Ser Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His	
35 40 45	
att gtc ctg gaa gat gga act aag atg aaa ggt tac tcc ttt ggc cat	312
Ile Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His	
50 55 60	
cca tcc tct gtt gct ggt gaa gtg gtt ttt aat act ggc ctg gga ggg	360
Pro Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly	
65 70 75	

-continued

tac cca gaa gct att act gac cct gcc tac aaa gga cag att ctc aca	408
Tyr Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr	
80 85 90 95	
atg gcc aac cct att att ggg aat ggt gga gct cct gat act act gct	456
Met Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala	
100 105 110	
ctg gat gaa ctg gga ctt agc aaa tat ttg gag tct aat gga atc aag	504
Leu Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys	
115 120 125	
gtt tca ggt ttg ctg gtg ctg gat tat agt aaa gac tac aac cac tgg	552
Val Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp	
130 135 140	
ctg gct acc aag agt tta ggg caa tgg cta cag gaa gaa aag gtt cct	600
Leu Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro	
145 150 155	
gca att tat gga gtg gac aca aga atg ctg act aaa ata att cgg gat	648
Ala Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp	
160 165 170 175	
aag ggt acc atg ctt ggg aag att gaa ttt gaa ggt cag cct gtg gat	696
Lys Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp	
180 185 190	
ttt gtg gat cca aat aaa cag aat ttg att gct gag gtt tca acc aag	744
Phe Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys	
195 200 205	
gat gtc aaa gtg tac ggc aaa gga aac ccc aca aaa gtg gta gct gta	792
Asp Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val	
210 215 220	
gac tgt ggg att aaa aac aat gta atc cgc ctg cta gta aag cga gga	840
Asp Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly	
225 230 235	
gct gaa gtg cac tta gtt ccc tgg aac cat gat ttc acc aag atg gag	888
Ala Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu	
240 245 250 255	
tat gat ggg att ttg atc gcg gga gga ccg ggg aac cca gct ctt gca	936
Tyr Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala	
260 265 270	
gaa cca cta att cag aat gtc aga aag att ttg gag agt gat cgc aag	984
Glu Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys	
275 280 285	
gag cca ttg ttt gga atc agt aca gga aac tta ata aca gga ttg gct	1032
Glu Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala	
290 295 300	
gct ggt gcc aaa acc tac aag atg tcc atg gcc aac aga ggg cag aat	1080
Ala Gly Ala Lys Thr Tyr Lys Met Ser Met Ala Asn Arg Gly Gln Asn	
305 310 315	
cag cct gtt ttg aat atc aca aac aaa cag gct ttc att act gct cag	1128
Gln Pro Val Leu Asn Ile Thr Asn Lys Gln Ala Phe Ile Thr Ala Gln	
320 325 330 335	
aat cat ggc tat gcc ttg gac aac acc ctc cct gct ggc tgg aaa cca	1176
Asn His Gly Tyr Ala Leu Asp Asn Thr Leu Pro Ala Gly Trp Lys Pro	
340 345 350	
ctt ttt gtg aat gtc aac gat caa aca aat gag ggg att atg cat gag	1224
Leu Phe Val Asn Val Asn Asp Gln Thr Asn Glu Gly Ile Met His Glu	
355 360 365	
agc aaa ccc ttc ttc gct gtg cag ttc cac cca gag gtc acc ccg ggg	1272
Ser Lys Pro Phe Phe Ala Val Gln Phe His Pro Glu Val Thr Pro Gly	
370 375 380	
cca ata gac act gag tac ctg ttt gat tcc ttt ttc tca ctg ata aag	1320
Pro Ile Asp Thr Glu Tyr Leu Phe Asp Ser Phe Phe Ser Leu Ile Lys	
385 390 395	

-continued

aaa gga aaa gct acc acc att aca tca gtc tta ccg aag cca gca cta	1368
Lys Gly Lys Ala Thr Thr Ile Thr Ser Val Leu Pro Lys Pro Ala Leu	
400 405 410 415	
gtt gca tct cgg gtt gag gtt tcc aaa gtc ctt att cta gga tca gga	1416
Val Ala Ser Arg Val Glu Val Ser Lys Val Leu Ile Leu Gly Ser Gly	
420 425 430	
ggt ctg tcc att ggt cag gct gga gaa ttt gat tac tca gga tct caa	1464
Gly Leu Ser Ile Gly Gln Ala Gly Glu Phe Asp Tyr Ser Gly Ser Gln	
435 440 445	
gct gta aaa gcc atg aag gaa gaa aat gtc aaa act gtt ctg atg aac	1512
Ala Val Lys Ala Met Lys Glu Glu Asn Val Lys Thr Val Leu Met Asn	
450 455 460	
cca aac att gca tca gtc cag acc aat gag gtg ggc tta aag caa gcg	1560
Pro Asn Ile Ala Ser Val Gln Thr Asn Glu Val Gly Leu Lys Gln Ala	
465 470 475	
gat act gtc tac ttt ctt ccc atc acc cct cag ttt gtc aca gag gtc	1608
Asp Thr Val Tyr Phe Leu Pro Ile Thr Pro Gln Phe Val Thr Glu Val	
480 485 490 495	
atc aag gca gaa cag cca gat ggg tta att ctg ggc atg ggt ggc cag	1656
Ile Lys Ala Glu Gln Pro Asp Gly Leu Ile Leu Gly Met Gly Gly Gln	
500 505 510	
aca gct ctg aac tgt gga gtg gaa cta ttc aag aga ggt gtg ctc aag	1704
Thr Ala Leu Asn Cys Gly Val Glu Leu Phe Lys Arg Gly Val Leu Lys	
515 520 525	
gaa tat ggt gtg aaa gtc ctg gga act tca gtt gag tcc att atg gct	1752
Glu Tyr Gly Val Lys Val Leu Gly Thr Ser Val Glu Ser Ile Met Ala	
530 535 540	
acg gaa gac agg cag ctg ttt tca gat aaa cta aat gag atc aat gaa	1800
Thr Glu Asp Arg Gln Leu Phe Ser Asp Lys Leu Asn Glu Ile Asn Glu	
545 550 555	
aag att gct cca agt ttt gca gtg gaa tcg att gag gat gca ctg aag	1848
Lys Ile Ala Pro Ser Phe Ala Val Glu Ser Ile Glu Asp Ala Leu Lys	
560 565 570 575	
gca gca gac acc att ggc tac cca gtg atg atc cgt tcc gcc tat gca	1896
Ala Ala Asp Thr Ile Gly Tyr Pro Val Met Ile Arg Ser Ala Tyr Ala	
580 585 590	
ctg ggt ggg tta ggc tca ggc atc tgt ccc aac aga gag act ttg atg	1944
Leu Gly Gly Leu Gly Ser Gly Ile Cys Pro Asn Arg Glu Thr Leu Met	
595 600 605	
gac ctc agc aca aag gcc ttt gct atg acc aac caa att ctg gtg gag	1992
Asp Leu Ser Thr Lys Ala Phe Ala Met Thr Asn Gln Ile Leu Val Glu	
610 615 620	
aag tca gtg aca ggt tgg aaa gaa ata gaa tat gaa gtg gtt cga gat	2040
Lys Ser Val Thr Gly Trp Lys Glu Ile Glu Tyr Glu Val Val Arg Asp	
625 630 635	
gct gat gac aat tgt gtc act gtc tgt aac atg gaa aat gtt gat gcc	2088
Ala Asp Asp Asn Cys Val Thr Val Cys Asn Met Glu Asn Val Asp Ala	
640 645 650 655	
atg ggt gtt cac aca ggt gac tca gtt gtt gtg gct cct gcc cag aca	2136
Met Gly Val His Thr Gly Asp Ser Val Val Val Ala Pro Ala Gln Thr	
660 665 670	
ctc tcc aat gcc gag ttt cag atg ttg aga cgt act tca atc aat gtt	2184
Leu Ser Asn Ala Glu Phe Gln Met Leu Arg Arg Thr Ser Ile Asn Val	
675 680 685	
gtt cgc cac ttg ggc att gtg ggt gaa tgc aac att cag ttt gcc ctt	2232
Val Arg His Leu Gly Ile Val Gly Glu Cys Asn Ile Gln Phe Ala Leu	
690 695 700	
cat cct acc tca atg gaa tac tgc atc att gaa gtg aat gcc aga ctg	2280
His Pro Thr Ser Met Glu Tyr Cys Ile Ile Glu Val Asn Ala Arg Leu	
705 710 715	

-continued

tcc cga agc tct gct ctg gcc tca aaa gcc act ggc tac cca ttg gca	2328
Ser Arg Ser Ser Ala Leu Ala Ser Lys Ala Thr Gly Tyr Pro Leu Ala	
720 725 730 735	
ttc att gct gca aag att gcc cta gga atc cca ctt cca gaa att aag	2376
Phe Ile Ala Ala Lys Ile Ala Leu Gly Ile Pro Leu Pro Glu Ile Lys	
740 745 750	
aac gtc gta tcc ggg aag aca tca gcc tgt ttt gaa cct agc ctg gat	2424
Asn Val Val Ser Gly Lys Thr Ser Ala Cys Phe Glu Pro Ser Leu Asp	
755 760 765	
tac atg gtc acc aag att ccc cgc tgg gat ctt gac cgt ttt cat gga	2472
Tyr Met Val Thr Lys Ile Pro Arg Trp Asp Leu Asp Arg Phe His Gly	
770 775 780	
aca tct agc cga att ggt agc tct atg aaa agt gta gga gag gtc atg	2520
Thr Ser Ser Arg Ile Gly Ser Ser Met Lys Ser Val Gly Glu Val Met	
785 790 795	
gct att ggt cgt acc ttt gag gag agt ttc cag aaa gct tta cgg atg	2568
Ala Ile Gly Arg Thr Phe Glu Glu Ser Phe Gln Lys Ala Leu Arg Met	
800 805 810 815	
tgc cac cca tct ata gaa ggt ttc act ccc cgt ctc cca atg aac aaa	2616
Cys His Pro Ser Ile Glu Gly Phe Thr Pro Arg Leu Pro Met Asn Lys	
820 825 830	
gaa tgg cca tct aat tta gat ctt aga aaa gag ttg tct gaa cca agc	2664
Glu Trp Pro Ser Asn Leu Asp Leu Arg Lys Glu Leu Ser Glu Pro Ser	
835 840 845	
agc acg cgt atc tat gcc att gcc aag gcc att gat gac aac atg tcc	2712
Ser Thr Arg Ile Tyr Ala Ile Ala Lys Ala Ile Asp Asp Asn Met Ser	
850 855 860	
ctt gat gag att gag aag ctc aca tac att gac aag tgg ttt ttg tat	2760
Leu Asp Glu Ile Glu Lys Leu Thr Tyr Ile Asp Lys Trp Phe Leu Tyr	
865 870 875	
aag atg cgt gat att tta aac atg gaa aag aca ctg aaa ggg ctc aac	2808
Lys Met Arg Asp Ile Leu Asn Met Glu Lys Thr Leu Lys Gly Leu Asn	
880 885 890 895	
agt gag tcc atg aca gaa gaa acc ctg aaa agg gca aag gag att ggg	2856
Ser Glu Ser Met Thr Glu Glu Thr Leu Lys Arg Ala Lys Glu Ile Gly	
900 905 910	
ttc tca gat aag cag att tca aaa tgc ctt ggg ctc act gag gcc cag	2904
Phe Ser Asp Lys Gln Ile Ser Lys Cys Leu Gly Leu Thr Glu Ala Gln	
915 920 925	
aca agg gag ctg agg tta aag aaa aac atc cac cct tgg gtt aaa cag	2952
Thr Arg Glu Leu Arg Leu Lys Lys Asn Ile His Pro Trp Val Lys Gln	
930 935 940	
att gat aca ctg gct gca gaa tac cca tca gta aca aac tat ctc tat	3000
Ile Asp Thr Leu Ala Ala Glu Tyr Pro Ser Val Thr Asn Tyr Leu Tyr	
945 950 955	
ggt acc tac aat ggt cag gag cat gat gtc aat ttt gat gac cat gga	3048
Val Thr Tyr Asn Gly Gln Glu His Asp Val Asn Phe Asp Asp His Gly	
960 965 970 975	
atg atg gtg cta ggc tgt ggt cca tat cac att ggc agc agt gtg gaa	3096
Met Met Val Leu Gly Cys Gly Pro Tyr His Ile Gly Ser Ser Val Glu	
980 985 990	
ttt gat tgg tgt gct gtc tct agt atc cgc aca ctg cgt caa ctt ggc	3144
Phe Asp Trp Cys Ala Val Ser Ser Ile Arg Thr Leu Arg Gln Leu Gly	
995 1000 1005	
aag aag acg gtg gtg gtg aat tgc aat cct gag act gtg agc aca	3189
Lys Lys Thr Val Val Val Asn Cys Asn Pro Glu Thr Val Ser Thr	
1010 1015 1020	
gac ttt gat gag tgt gac aaa ctg tac ttt gaa gag ttg tcc ttg	3234
Asp Phe Asp Glu Cys Asp Lys Leu Tyr Phe Glu Glu Leu Ser Leu	
1025 1030 1035	

-continued

gag aga atc cta gac atc tac cat cag gag gca tgt ggt ggc tgc	3279
Glu Arg Ile Leu Asp Ile Tyr His Gln Glu Ala Cys Gly Gly Cys	
1040 1045 1050	
atc ata tca gtt gga ggc cag att cca aac aac ctg gca gtt cct	3324
Ile Ile Ser Val Gly Gly Gln Ile Pro Asn Asn Leu Ala Val Pro	
1055 1060 1065	
cta tac aag aat ggt gtc aag atc atg ggc aca agc ccc ctg cag	3369
Leu Tyr Lys Asn Gly Val Lys Ile Met Gly Thr Ser Pro Leu Gln	
1070 1075 1080	
atc gac agg gct gag gat cgc tcc atc ttc tca gct gtc ttg gat	3414
Ile Asp Arg Ala Glu Asp Arg Ser Ile Phe Ser Ala Val Leu Asp	
1085 1090 1095	
gag ctg aag gtg gct cag gca cct tgg aaa gct gtt aat act ttg	3459
Glu Leu Lys Val Ala Gln Ala Pro Trp Lys Ala Val Asn Thr Leu	
1100 1105 1110	
aat gaa gca ctg gaa ttt gca aag tct gtg gac tac ccc tgc ttg	3504
Asn Glu Ala Leu Glu Phe Ala Lys Ser Val Asp Tyr Pro Cys Leu	
1115 1120 1125	
ttg agg cct tcc tat gtt ttg agt ggg tct gct atg aat gtg gta	3549
Leu Arg Pro Ser Tyr Val Leu Ser Gly Ser Ala Met Asn Val Val	
1130 1135 1140	
ttc tct gag gat gag atg aaa aaa ttc cta gaa gag gcg act aga	3594
Phe Ser Glu Asp Glu Met Lys Lys Phe Leu Glu Glu Ala Thr Arg	
1145 1150 1155	
gtt tct cag gag cac cca gtg gtc ctg aca aaa ttt gtt gaa ggg	3639
Val Ser Gln Glu His Pro Val Val Leu Thr Lys Phe Val Glu Gly	
1160 1165 1170	
gcc cga gaa gta gaa atg gac gct gtt ggc aaa gat gga agg gtt	3684
Ala Arg Glu Val Glu Met Asp Ala Val Gly Lys Asp Gly Arg Val	
1175 1180 1185	
atc tct cat gcc atc tct gaa cat gtt gaa gat gca ggt gtc cac	3729
Ile Ser His Ala Ile Ser Glu His Val Glu Asp Ala Gly Val His	
1190 1195 1200	
tcg gga gat gcc act ctg atg ctg ccc aca caa acc atc agc caa	3774
Ser Gly Asp Ala Thr Leu Met Leu Pro Thr Gln Thr Ile Ser Gln	
1205 1210 1215	
ggg gcc att gaa aag gtg aag gat gct acc cgg aag att gca aag	3819
Gly Ala Ile Glu Lys Val Lys Asp Ala Thr Arg Lys Ile Ala Lys	
1220 1225 1230	
gct ttt gcc atc tct ggt cca ttc aac gtc caa ttt ctt gtc aaa	3864
Ala Phe Ala Ile Ser Gly Pro Phe Asn Val Gln Phe Leu Val Lys	
1235 1240 1245	
gga aat gat gtc ttg gtg att gag tgt aac ttg aga gct tct cga	3909
Gly Asn Asp Val Leu Val Ile Glu Cys Asn Leu Arg Ala Ser Arg	
1250 1255 1260	
tcc ttc ccc ttt gtt tcc aag act ctt ggg gtt gac ttc att gat	3954
Ser Phe Pro Phe Val Ser Lys Thr Leu Gly Val Asp Phe Ile Asp	
1265 1270 1275	
gtg gcc acc aag gtg atg att gga gag aat gtt gat gag aaa cat	3999
Val Ala Thr Lys Val Met Ile Gly Glu Asn Val Asp Glu Lys His	
1280 1285 1290	
ctt cca aca ttg gac cat ccc ata att cct gct gac tat gtt gca	4044
Leu Pro Thr Leu Asp His Pro Ile Ile Pro Ala Asp Tyr Val Ala	
1295 1300 1305	
att aag gct ccc atg ttt tcc tgg ccc cgg ttg agg gat gct gac	4089
Ile Lys Ala Pro Met Phe Ser Trp Pro Arg Leu Arg Asp Ala Asp	
1310 1315 1320	
ccc att ctg aga tgt gag atg gct tcc act gga gag gtg gct tgc	4134
Pro Ile Leu Arg Cys Glu Met Ala Ser Thr Gly Glu Val Ala Cys	
1325 1330 1335	

-continued

ttt ggt gaa ggt att cat aca gcc ttc cta aag gca atg ctt tcc	4179
Phe Gly Glu Gly Ile His Thr Ala Phe Leu Lys Ala Met Leu Ser	
1340 1345 1350	
aca gga ttt aag ata ccc cag aaa ggc atc ctg ata ggc atc cag	4224
Thr Gly Phe Lys Ile Pro Gln Lys Gly Ile Leu Ile Gly Ile Gln	
1355 1360 1365	
caa tca ttc cgg cca aga ttc ctt ggt gtg gct gaa caa tta cac	4269
Gln Ser Phe Arg Pro Arg Phe Leu Gly Val Ala Glu Gln Leu His	
1370 1375 1380	
aat gaa ggt ttc aag ctg ttt gcc acg gaa gcc aca tca gac tgg	4314
Asn Glu Gly Phe Lys Leu Phe Ala Thr Glu Ala Thr Ser Asp Trp	
1385 1390 1395	
ctc aac gcc aac aat gtc cct gcc acc cca gtg gca tgg ccg tct	4359
Leu Asn Ala Asn Asn Val Pro Ala Thr Pro Val Ala Trp Pro Ser	
1400 1405 1410	
caa gaa gga cag aat ccc agc ctc tct tcc atc aga aaa ttg att	4404
Gln Glu Gly Gln Asn Pro Ser Leu Ser Ser Ile Arg Lys Leu Ile	
1415 1420 1425	
aga gat ggc agc att gac cta gtg att aac ctt ccc aac aac aac	4449
Arg Asp Gly Ser Ile Asp Leu Val Ile Asn Leu Pro Asn Asn Asn	
1430 1435 1440	
act aaa ttt gtc cat gat aat tat gtg att cgg agg aca gct gtt	4494
Thr Lys Phe Val His Asp Asn Tyr Val Ile Arg Arg Thr Ala Val	
1445 1450 1455	
gat agt gga atc cct ctc ctc act aat ttt cag gtg acc aaa ctt	4539
Asp Ser Gly Ile Pro Leu Leu Thr Asn Phe Gln Val Thr Lys Leu	
1460 1465 1470	
ttt gct gaa gct gtg cag aaa tct cgc aag gtg gac tcc aag agt	4584
Phe Ala Glu Ala Val Gln Lys Ser Arg Lys Val Asp Ser Lys Ser	
1475 1480 1485	
ctt ttc cac tac agg cag tac agt gct gga aaa gca gca tag	4626
Leu Phe His Tyr Arg Gln Tyr Ser Ala Gly Lys Ala Ala	
1490 1495 1500	
agatgcagac accccagccc cattattaata tcaacctgag ccacatgtta tctaaaggaa	4686
ctgattcaca actttctcag agatgaatat tgataactaa acttcatttc agtttacttt	4746
gttatgcctt aatattctgt gtcttttgca attaaattgt cagtcacttc ttcaaacct	4806
tacagtcctt cctaagttac tcttcatgag atttcatcca tttactaata ctgtattttt	4866
ggtggactag gcttgctat gtgcttatgt gtagcttttt actttttatg gtgctgatta	4926
atggtgatca aggtaggaaa agttgctgtt ctattttctg aactctttct atactttaag	4986
atactctatt tttaaaacac tatctgcaaa ctgaggacac tttacaggg cagaatactc	5046
taaaaacttg ataaaatgaa atatagattt aatttatgaa ccttccatca tgatgtttgt	5106
gtattgcttc tttttggatc ctcatctca cccatttggc taatccagga atattgttat	5166
cccttcccat tatattgaag ttgagaaatg tgacagaggc atttagagta tggacttttc	5226
ttttcttttt cttttctttt ttttcttttt gagatggagt cacactctcc aggetggagt	5286
gcagtggcac aatctcggct cactgcaatt tgcgtctccc aagttcaagc gattctcctg	5346
ctttagacta tggatttctt taaggaatac tggtttgag ttttgttttc tggactatat	5406
cagcagatgg tagacagtgt ttatgtagat gtgtgttgt ttttatcatt ggattttaac	5466
ttggcccagag tgaataatc agatttttgt cattcacact ctccccagt tttggaataa	5526
cttgaagta aggttcattc ccttaagacg atggattctg ttgaactatg gggcccaca	5586
ctgcactatt aattccacc actgtaagg caaggacacc attcctteta catataagaa	5646
aaaagtctct cccaagggc agcctttgtt acttttaaat attttctgtt attacaagtg	5706

-continued

 ctctaattgt gaacttttaa ataaaatact attaagaggt aaaaaaaaaa aaaaa 5761

<210> SEQ ID NO 4

<211> LENGTH: 1500

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 4

Met Thr Arg Ile Leu Thr Ala Phe Lys Val Val Arg Thr Leu Lys Thr
 1 5 10 15
 Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe Ser
 20 25 30
 Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His Ile
 35 40 45
 Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His Pro
 50 55 60
 Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly Tyr
 65 70 75 80
 Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr Met
 85 90 95
 Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala Leu
 100 105 110
 Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys Val
 115 120 125
 Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp Leu
 130 135 140
 Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro Ala
 145 150 155 160
 Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp Lys
 165 170 175
 Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp Phe
 180 185 190
 Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys Asp
 195 200 205
 Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val Asp
 210 215 220
 Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly Ala
 225 230 235 240
 Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu Tyr
 245 250 255
 Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala Glu
 260 265 270
 Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys Glu
 275 280 285
 Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala Ala
 290 295 300
 Gly Ala Lys Thr Tyr Lys Met Ser Met Ala Asn Arg Gly Gln Asn Gln
 305 310 315 320
 Pro Val Leu Asn Ile Thr Asn Lys Gln Ala Phe Ile Thr Ala Gln Asn
 325 330 335
 His Gly Tyr Ala Leu Asp Asn Thr Leu Pro Ala Gly Trp Lys Pro Leu
 340 345 350
 Phe Val Asn Val Asn Asp Gln Thr Asn Glu Gly Ile Met His Glu Ser
 355 360 365
 Lys Pro Phe Phe Ala Val Gln Phe His Pro Glu Val Thr Pro Gly Pro

-continued

370			375			380									
Ile	Asp	Thr	Glu	Tyr	Leu	Phe	Asp	Ser	Phe	Phe	Ser	Leu	Ile	Lys	Lys
385					390				395					400	
Gly	Lys	Ala	Thr	Thr	Ile	Thr	Ser	Val	Leu	Pro	Lys	Pro	Ala	Leu	Val
				405					410					415	
Ala	Ser	Arg	Val	Glu	Val	Ser	Lys	Val	Leu	Ile	Leu	Gly	Ser	Gly	Gly
			420					425					430		
Leu	Ser	Ile	Gly	Gln	Ala	Gly	Glu	Phe	Asp	Tyr	Ser	Gly	Ser	Gln	Ala
		435						440					445		
Val	Lys	Ala	Met	Lys	Glu	Glu	Asn	Val	Lys	Thr	Val	Leu	Met	Asn	Pro
	450						455				460				
Asn	Ile	Ala	Ser	Val	Gln	Thr	Asn	Glu	Val	Gly	Leu	Lys	Gln	Ala	Asp
465					470					475					480
Thr	Val	Tyr	Phe	Leu	Pro	Ile	Thr	Pro	Gln	Phe	Val	Thr	Glu	Val	Ile
				485						490				495	
Lys	Ala	Glu	Gln	Pro	Asp	Gly	Leu	Ile	Leu	Gly	Met	Gly	Gly	Gln	Thr
				500				505					510		
Ala	Leu	Asn	Cys	Gly	Val	Glu	Leu	Phe	Lys	Arg	Gly	Val	Leu	Lys	Glu
		515						520					525		
Tyr	Gly	Val	Lys	Val	Leu	Gly	Thr	Ser	Val	Glu	Ser	Ile	Met	Ala	Thr
	530						535				540				
Glu	Asp	Arg	Gln	Leu	Phe	Ser	Asp	Lys	Leu	Asn	Glu	Ile	Asn	Glu	Lys
545					550					555					560
Ile	Ala	Pro	Ser	Phe	Ala	Val	Glu	Ser	Ile	Glu	Asp	Ala	Leu	Lys	Ala
				565						570				575	
Ala	Asp	Thr	Ile	Gly	Tyr	Pro	Val	Met	Ile	Arg	Ser	Ala	Tyr	Ala	Leu
				580				585					590		
Gly	Gly	Leu	Gly	Ser	Gly	Ile	Cys	Pro	Asn	Arg	Glu	Thr	Leu	Met	Asp
		595					600						605		
Leu	Ser	Thr	Lys	Ala	Phe	Ala	Met	Thr	Asn	Gln	Ile	Leu	Val	Glu	Lys
	610						615				620				
Ser	Val	Thr	Gly	Trp	Lys	Glu	Ile	Glu	Tyr	Glu	Val	Val	Arg	Asp	Ala
625					630					635					640
Asp	Asp	Asn	Cys	Val	Thr	Val	Cys	Asn	Met	Glu	Asn	Val	Asp	Ala	Met
				645						650				655	
Gly	Val	His	Thr	Gly	Asp	Ser	Val	Val	Val	Ala	Pro	Ala	Gln	Thr	Leu
				660				665					670		
Ser	Asn	Ala	Glu	Phe	Gln	Met	Leu	Arg	Arg	Thr	Ser	Ile	Asn	Val	Val
		675					680						685		
Arg	His	Leu	Gly	Ile	Val	Gly	Glu	Cys	Asn	Ile	Gln	Phe	Ala	Leu	His
	690						695				700				
Pro	Thr	Ser	Met	Glu	Tyr	Cys	Ile	Ile	Glu	Val	Asn	Ala	Arg	Leu	Ser
705					710					715					720
Arg	Ser	Ser	Ala	Leu	Ala	Ser	Lys	Ala	Thr	Gly	Tyr	Pro	Leu	Ala	Phe
				725						730				735	
Ile	Ala	Ala	Lys	Ile	Ala	Leu	Gly	Ile	Pro	Leu	Pro	Glu	Ile	Lys	Asn
				740				745					750		
Val	Val	Ser	Gly	Lys	Thr	Ser	Ala	Cys	Phe	Glu	Pro	Ser	Leu	Asp	Tyr
		755					760						765		
Met	Val	Thr	Lys	Ile	Pro	Arg	Trp	Asp	Leu	Asp	Arg	Phe	His	Gly	Thr
	770						775				780				
Ser	Ser	Arg	Ile	Gly	Ser	Ser	Met	Lys	Ser	Val	Gly	Glu	Val	Met	Ala
					790					795					800

-continued

Ile	Gly	Arg	Thr	Phe	Glu	Glu	Ser	Phe	Gln	Lys	Ala	Leu	Arg	Met	Cys
				805					810					815	
His	Pro	Ser	Ile	Glu	Gly	Phe	Thr	Pro	Arg	Leu	Pro	Met	Asn	Lys	Glu
			820					825					830		
Trp	Pro	Ser	Asn	Leu	Asp	Leu	Arg	Lys	Glu	Leu	Ser	Glu	Pro	Ser	Ser
		835					840					845			
Thr	Arg	Ile	Tyr	Ala	Ile	Ala	Lys	Ala	Ile	Asp	Asp	Asn	Met	Ser	Leu
	850					855					860				
Asp	Glu	Ile	Glu	Lys	Leu	Thr	Tyr	Ile	Asp	Lys	Trp	Phe	Leu	Tyr	Lys
865					870				875						880
Met	Arg	Asp	Ile	Leu	Asn	Met	Glu	Lys	Thr	Leu	Lys	Gly	Leu	Asn	Ser
				885					890					895	
Glu	Ser	Met	Thr	Glu	Glu	Thr	Leu	Lys	Arg	Ala	Lys	Glu	Ile	Gly	Phe
			900					905					910		
Ser	Asp	Lys	Gln	Ile	Ser	Lys	Cys	Leu	Gly	Leu	Thr	Glu	Ala	Gln	Thr
		915					920					925			
Arg	Glu	Leu	Arg	Leu	Lys	Lys	Asn	Ile	His	Pro	Trp	Val	Lys	Gln	Ile
	930					935					940				
Asp	Thr	Leu	Ala	Ala	Glu	Tyr	Pro	Ser	Val	Thr	Asn	Tyr	Leu	Tyr	Val
945					950					955					960
Thr	Tyr	Asn	Gly	Gln	Glu	His	Asp	Val	Asn	Phe	Asp	Asp	His	Gly	Met
				965				970						975	
Met	Val	Leu	Gly	Cys	Gly	Pro	Tyr	His	Ile	Gly	Ser	Ser	Val	Glu	Phe
		980						985					990		
Asp	Trp	Cys	Ala	Val	Ser	Ser	Ile	Arg	Thr	Leu	Arg	Gln	Leu	Gly	Lys
		995					1000					1005			
Lys	Thr	Val	Val	Val	Asn	Cys	Asn	Pro	Glu	Thr	Val	Ser	Thr	Asp	
	1010					1015					1020				
Phe	Asp	Glu	Cys	Asp	Lys	Leu	Tyr	Phe	Glu	Glu	Leu	Ser	Leu	Glu	
	1025					1030					1035				
Arg	Ile	Leu	Asp	Ile	Tyr	His	Gln	Glu	Ala	Cys	Gly	Gly	Cys	Ile	
	1040					1045					1050				
Ile	Ser	Val	Gly	Gly	Gln	Ile	Pro	Asn	Asn	Leu	Ala	Val	Pro	Leu	
	1055					1060					1065				
Tyr	Lys	Asn	Gly	Val	Lys	Ile	Met	Gly	Thr	Ser	Pro	Leu	Gln	Ile	
	1070					1075					1080				
Asp	Arg	Ala	Glu	Asp	Arg	Ser	Ile	Phe	Ser	Ala	Val	Leu	Asp	Glu	
	1085					1090					1095				
Leu	Lys	Val	Ala	Gln	Ala	Pro	Trp	Lys	Ala	Val	Asn	Thr	Leu	Asn	
	1100					1105					1110				
Glu	Ala	Leu	Glu	Phe	Ala	Lys	Ser	Val	Asp	Tyr	Pro	Cys	Leu	Leu	
	1115					1120					1125				
Arg	Pro	Ser	Tyr	Val	Leu	Ser	Gly	Ser	Ala	Met	Asn	Val	Val	Phe	
	1130					1135					1140				
Ser	Glu	Asp	Glu	Met	Lys	Lys	Phe	Leu	Glu	Glu	Ala	Thr	Arg	Val	
	1145					1150					1155				
Ser	Gln	Glu	His	Pro	Val	Val	Leu	Thr	Lys	Phe	Val	Glu	Gly	Ala	
	1160					1165					1170				
Arg	Glu	Val	Glu	Met	Asp	Ala	Val	Gly	Lys	Asp	Gly	Arg	Val	Ile	
	1175					1180					1185				
Ser	His	Ala	Ile	Ser	Glu	His	Val	Glu	Asp	Ala	Gly	Val	His	Ser	
	1190					1195					1200				
Gly	Asp	Ala	Thr	Leu	Met	Leu	Pro	Thr	Gln	Thr	Ile	Ser	Gln	Gly	
	1205					1210					1215				

-continued

Ala	Ile	Glu	Lys	Val	Lys	Asp	Ala	Thr	Arg	Lys	Ile	Ala	Lys	Ala
1220						1225					1230			
Phe	Ala	Ile	Ser	Gly	Pro	Phe	Asn	Val	Gln	Phe	Leu	Val	Lys	Gly
1235						1240					1245			
Asn	Asp	Val	Leu	Val	Ile	Glu	Cys	Asn	Leu	Arg	Ala	Ser	Arg	Ser
1250						1255					1260			
Phe	Pro	Phe	Val	Ser	Lys	Thr	Leu	Gly	Val	Asp	Phe	Ile	Asp	Val
1265						1270					1275			
Ala	Thr	Lys	Val	Met	Ile	Gly	Glu	Asn	Val	Asp	Glu	Lys	His	Leu
1280						1285					1290			
Pro	Thr	Leu	Asp	His	Pro	Ile	Ile	Pro	Ala	Asp	Tyr	Val	Ala	Ile
1295						1300					1305			
Lys	Ala	Pro	Met	Phe	Ser	Trp	Pro	Arg	Leu	Arg	Asp	Ala	Asp	Pro
1310						1315					1320			
Ile	Leu	Arg	Cys	Glu	Met	Ala	Ser	Thr	Gly	Glu	Val	Ala	Cys	Phe
1325						1330					1335			
Gly	Glu	Gly	Ile	His	Thr	Ala	Phe	Leu	Lys	Ala	Met	Leu	Ser	Thr
1340						1345					1350			
Gly	Phe	Lys	Ile	Pro	Gln	Lys	Gly	Ile	Leu	Ile	Gly	Ile	Gln	Gln
1355						1360					1365			
Ser	Phe	Arg	Pro	Arg	Phe	Leu	Gly	Val	Ala	Glu	Gln	Leu	His	Asn
1370						1375					1380			
Glu	Gly	Phe	Lys	Leu	Phe	Ala	Thr	Glu	Ala	Thr	Ser	Asp	Trp	Leu
1385						1390					1395			
Asn	Ala	Asn	Asn	Val	Pro	Ala	Thr	Pro	Val	Ala	Trp	Pro	Ser	Gln
1400						1405					1410			
Glu	Gly	Gln	Asn	Pro	Ser	Leu	Ser	Ser	Ile	Arg	Lys	Leu	Ile	Arg
1415						1420					1425			
Asp	Gly	Ser	Ile	Asp	Leu	Val	Ile	Asn	Leu	Pro	Asn	Asn	Asn	Thr
1430						1435					1440			
Lys	Phe	Val	His	Asp	Asn	Tyr	Val	Ile	Arg	Arg	Thr	Ala	Val	Asp
1445						1450					1455			
Ser	Gly	Ile	Pro	Leu	Leu	Thr	Asn	Phe	Gln	Val	Thr	Lys	Leu	Phe
1460						1465					1470			
Ala	Glu	Ala	Val	Gln	Lys	Ser	Arg	Lys	Val	Asp	Ser	Lys	Ser	Leu
1475						1480					1485			
Phe	His	Tyr	Arg	Gln	Tyr	Ser	Ala	Gly	Lys	Ala	Ala			
1490						1495					1500			

<210> SEQ ID NO 5
 <211> LENGTH: 495
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: misc_feature
 <222> LOCATION: (1)..(495)
 <223> OTHER INFORMATION: n is a, c, g, or t

 <400> SEQUENCE: 5

ctacttctca	tgttcagcaa	tttcttcttc	tttatgtttt	aaattacatg	ttccataaaa	60
ataagaaatn	cactgtgata	cggttaattga	ttttttcatt	ttaaattgcag	ctgtttgcca	120
cggaagccac	atcagactgg	ctcaacgccca	acaatgtccc	tgccacccca	gtggcatggc	180
cgtctcaaga	aggacagaat	cccagcctct	cttccatcag	aaagtaagaa	ctaggcatac	240
tgttttctga	aataatttag	aggattaact	ttgagaacca	gtatatgaat	attcaccttg	300

-continued

```

cttgattgca agtcttttaa aacaaattta aaaatgaata catttggtga tgattgtcaa 360
gtttcactct ccatcactat ggaatacata acgtcatgtg tacatggtga tatgaaacgt 420
gtttcaaaat acttcttagt aaggatactt tccttgacgg aaacaagtga gagtatgaag 480
aatgtaatgc agcac 495

```

```

<210> SEQ ID NO 6
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 6

```

```

agctgtttgc cacggaagcc 20

```

```

<210> SEQ ID NO 7
<211> LENGTH: 28
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 7

```

```

cccagcctct cttccatcag aaagtaag 28

```

```

<210> SEQ ID NO 8
<211> LENGTH: 20
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 8

```

```

cacggaagcc acatcagact 20

```

```

<210> SEQ ID NO 9
<211> LENGTH: 21
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 9

```

```

ttctgatgga agagaggctt g 21

```

```

<210> SEQ ID NO 10
<211> LENGTH: 24
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens

```

```

<400> SEQUENCE: 10

```

```

agagtgaaac ttgacaatca tcca 24

```

```

<210> SEQ ID NO 11
<211> LENGTH: 5761
<212> TYPE: DNA
<213> ORGANISM: Homo sapiens
<220> FEATURE:
<221> NAME/KEY: CDS
<222> LOCATION: (124)..(4626)

```

```

<400> SEQUENCE: 11

```

```

gtcagcctta aacactgact gcaccctcc cagatttctt ttacattaac taaaaagtct 60

```

```

tatcacacaa tctcataaaa tttatgtaat ttcatttaat tttageccaca aatcatcttc 120

```

```

aaa atg acg agg tta ttg aca gct ttc aaa gtg gtg agg aca ctg aag 168
Met Thr Arg Leu Leu Thr Ala Phe Lys Val Val Arg Thr Leu Lys
1 5 10 15

```

```

act ggt ttt ggc ttt acc aat gtg act gca cac caa aaa tgg aaa ttt 216
Thr Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe
20 25 30

```

-continued

tca aga cct ggc atc agg ctc ctt tct gtc aag gca cag aca gca cac	264
Ser Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His	
35 40 45	
att gtc ctg gaa gat gga act aag atg aaa ggt tac tcc ttt ggc cat	312
Ile Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His	
50 55 60	
cca tcc tct gtt gct ggt gaa gtg gtt ttt aat act ggc ctg gga ggg	360
Pro Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly	
65 70 75	
tac cca gaa gct att act gac cct gcc tac aaa gga cag att ctc aca	408
Tyr Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr	
80 85 90 95	
atg gcc aac cct att att ggg aat ggt gga gct cct gat act act gct	456
Met Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala	
100 105 110	
ctg gat gaa ctg gga ctt agc aaa tat ttg gag tct aat gga atc aag	504
Leu Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys	
115 120 125	
gtt tca ggt ttg ctg gtg ctg gat tat agt aaa gac tac aac cac tgg	552
Val Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp	
130 135 140	
ctg gct acc aag agt tta ggg caa tgg cta cag gaa gaa aag gtt cct	600
Leu Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro	
145 150 155	
gca att tat gga gtg gac aca aga atg ctg act aaa ata att cgg gat	648
Ala Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp	
160 165 170 175	
aag ggt acc atg ctt ggg aag att gaa ttt gaa ggt cag cct gtg gat	696
Lys Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp	
180 185 190	
ttt gtg gat cca aat aaa cag aat ttg att gct gag gtt tca acc aag	744
Phe Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys	
195 200 205	
gat gtc aaa gtg tac ggc aaa gga aac ccc aca aaa gtg gta gct gta	792
Asp Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val	
210 215 220	
gac tgt ggg att aaa aac aat gta atc cgc ctg cta gta aag cga gga	840
Asp Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly	
225 230 235	
gct gaa gtg cac tta gtt ccc tgg aac cat gat ttc acc aag atg gag	888
Ala Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu	
240 245 250 255	
tat gat ggg att ttg atc gcg gga gga ccg ggg aac cca gct ctt gca	936
Tyr Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala	
260 265 270	
gaa cca cta att cag aat gtc aga aag att ttg gag agt gat cgc aag	984
Glu Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys	
275 280 285	
gag cca ttg ttt gga atc agt aca gga aac tta ata aca gga ttg gct	1032
Glu Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala	
290 295 300	
gct ggt gcc aaa acc tac aag atg tcc atg gcc aac aga ggg cag aat	1080
Ala Gly Ala Lys Thr Tyr Lys Met Ser Met Ala Asn Arg Gly Gln Asn	
305 310 315	
cag cct gtt ttg aat atc aca aac aaa cag gct ttc att act gct cag	1128
Gln Pro Val Leu Asn Ile Thr Asn Lys Gln Ala Phe Ile Thr Ala Gln	
320 325 330 335	
aat cat ggc tat gcc ttg gac aac acc ctc cct gct ggc tgg aaa cca	1176
Asn His Gly Tyr Ala Leu Asp Asn Thr Leu Pro Ala Gly Trp Lys Pro	
340 345 350	

-continued

ctt ttt gtg aat gtc aac gat caa aca aat gag ggg att atg cat gag	1224
Leu Phe Val Asn Val Asn Asp Gln Thr Asn Glu Gly Ile Met His Glu	
355 360 365	
agc aaa ccc ttc ttc gct gtg cag ttc cac cca gag gtc acc ccg ggg	1272
Ser Lys Pro Phe Phe Ala Val Gln Phe His Pro Glu Val Thr Pro Gly	
370 375 380	
cca ata gac act gag tac ctg ttt gat tcc ttt ttc tca ctg ata aag	1320
Pro Ile Asp Thr Glu Tyr Leu Phe Asp Ser Phe Phe Ser Leu Ile Lys	
385 390 395	
aaa gga aaa gct acc acc att aca tca gtc tta ccg aag cca gca cta	1368
Lys Gly Lys Ala Thr Thr Ile Thr Ser Val Leu Pro Lys Pro Ala Leu	
400 405 410 415	
ggt gca tct cgg gtt gag gtt tcc aaa gtc ctt att cta gga tca gga	1416
Val Ala Ser Arg Val Glu Val Ser Lys Val Leu Ile Leu Gly Ser Gly	
420 425 430	
ggt ctg tcc att ggt cag gct gga gaa ttt gat tac tca gga tct caa	1464
Gly Leu Ser Ile Gly Gln Ala Gly Glu Phe Asp Tyr Ser Gly Ser Gln	
435 440 445	
gct gta aaa gcc atg aag gaa gaa aat gtc aaa act gtt ctg atg aac	1512
Ala Val Lys Ala Met Lys Glu Glu Asn Val Lys Thr Val Leu Met Asn	
450 455 460	
cca aac att gca tca gtc cag acc aat gag gtg ggc tta aag caa gcg	1560
Pro Asn Ile Ala Ser Val Gln Thr Asn Glu Val Gly Leu Lys Gln Ala	
465 470 475	
gat act gtc tac ttt ctt ccc atc acc cct cag ttt gtc aca gag gtc	1608
Asp Thr Val Tyr Phe Leu Pro Ile Thr Pro Gln Phe Val Thr Glu Val	
480 485 490 495	
atc aag gca gaa cag cca gat ggg tta att ctg ggc atg ggt ggc cag	1656
Ile Lys Ala Glu Gln Pro Asp Gly Leu Ile Leu Gly Met Gly Gly Gln	
500 505 510	
aca gct ctg aac tgt gga gtg gaa cta ttc aag aga ggt gtg ctc aag	1704
Thr Ala Leu Asn Cys Gly Val Glu Leu Phe Lys Arg Gly Val Leu Lys	
515 520 525	
gaa tat ggt gtg aaa gtc ctg gga act tca gtt gag tcc att atg gct	1752
Glu Tyr Gly Val Lys Val Leu Gly Thr Ser Val Glu Ser Ile Met Ala	
530 535 540	
acg gaa gac agg cag ctg ttt tca gat aaa cta aat gag atc aat gaa	1800
Thr Glu Asp Arg Gln Leu Phe Ser Asp Lys Leu Asn Glu Ile Asn Glu	
545 550 555	
aag att gct cca agt ttt gca gtg gaa tcg att gag gat gca ctg aag	1848
Lys Ile Ala Pro Ser Phe Ala Val Glu Ser Ile Glu Asp Ala Leu Lys	
560 565 570 575	
gca gca gac acc att ggc tac cca gtg atg atc cgt tcc gcc tat gca	1896
Ala Ala Asp Thr Ile Gly Tyr Pro Val Met Ile Arg Ser Ala Tyr Ala	
580 585 590	
ctg ggt ggg tta ggc tca ggc atc tgt ccc aac aga gag act ttg atg	1944
Leu Gly Gly Leu Gly Ser Gly Ile Cys Pro Asn Arg Glu Thr Leu Met	
595 600 605	
gac ctc agc aca aag gcc ttt gct atg acc aac caa att ctg gtg gag	1992
Asp Leu Ser Thr Lys Ala Phe Ala Met Thr Asn Gln Ile Leu Val Glu	
610 615 620	
aag tca gtg aca ggt tgg aaa gaa ata gaa tat gaa gtg gtt cga gat	2040
Lys Ser Val Thr Gly Trp Lys Glu Ile Glu Tyr Glu Val Val Arg Asp	
625 630 635	
gct gat gac aat tgt gtc act gtc tgt aac atg gaa aat gtt gat gcc	2088
Ala Asp Asp Asn Cys Val Thr Val Cys Asn Met Glu Asn Val Asp Ala	
640 645 650 655	
atg ggt gtt cac aca ggt gac tca gtt gtt gtg gct cct gcc cag aca	2136
Met Gly Val His Thr Gly Asp Ser Val Val Val Ala Pro Ala Gln Thr	
660 665 670	

-continued

ctc tcc aat gcc gag ttt cag atg ttg aga cgt act tca atc aat gtt	2184
Leu Ser Asn Ala Glu Phe Gln Met Leu Arg Arg Thr Ser Ile Asn Val	
675 680 685	
gtt cgc cac ttg ggc att gtg ggt gaa tgc aac att cag ttt gcc ctt	2232
Val Arg His Leu Gly Ile Val Gly Glu Cys Asn Ile Gln Phe Ala Leu	
690 695 700	
cat cct acc tca atg gaa tac tgc atc att gaa gtg aat gcc aga ctg	2280
His Pro Thr Ser Met Glu Tyr Cys Ile Ile Glu Val Asn Ala Arg Leu	
705 710 715	
tcc cga agc tct gct ctg gcc tca aaa gcc act ggc tac cca ttg gca	2328
Ser Arg Ser Ser Ala Leu Ala Ser Lys Ala Thr Gly Tyr Pro Leu Ala	
720 725 730 735	
ttc att gct gca aag att gcc cta gga atc cca ctt cca gaa att aag	2376
Phe Ile Ala Ala Lys Ile Ala Leu Gly Ile Pro Leu Pro Glu Ile Lys	
740 745 750	
aac gtc gta tcc ggg aag aca tca gcc tgt ttt gaa cct agc ctg gat	2424
Asn Val Val Ser Gly Lys Thr Ser Ala Cys Phe Glu Pro Ser Leu Asp	
755 760 765	
tac atg gtc acc aag att ccc cgc tgg gat ctt gac cgt ttt cat gga	2472
Tyr Met Val Thr Lys Ile Pro Arg Trp Asp Leu Asp Arg Phe His Gly	
770 775 780	
aca tct agc cga att ggt agc tct atg aaa agt gta gga gag gtc atg	2520
Thr Ser Ser Arg Ile Gly Ser Ser Met Lys Ser Val Gly Glu Val Met	
785 790 795	
gct att ggt cgt acc ttt gag gag agt ttc cag aaa gct tta cgg atg	2568
Ala Ile Gly Arg Thr Phe Glu Glu Ser Phe Gln Lys Ala Leu Arg Met	
800 805 810 815	
tgc cac cca tct ata gaa ggt ttc act ccc cgt ctc cca atg aac aaa	2616
Cys His Pro Ser Ile Glu Gly Phe Thr Pro Arg Leu Pro Met Asn Lys	
820 825 830	
gaa tgg cca tct aat tta gat ctt aga aaa gag ttg tct gaa cca agc	2664
Glu Trp Pro Ser Asn Leu Asp Leu Arg Lys Glu Leu Ser Glu Pro Ser	
835 840 845	
agc acg cgt atc tat gcc att gcc aag gcc att gat gac aac atg tcc	2712
Ser Thr Arg Ile Tyr Ala Ile Ala Lys Ala Ile Asp Asp Asn Met Ser	
850 855 860	
ctt gat gag att gag aag ctc aca tac att gac aag tgg ttt ttg tat	2760
Leu Asp Glu Ile Glu Lys Leu Thr Tyr Ile Asp Lys Trp Phe Leu Tyr	
865 870 875	
aag atg cgt gat att tta aac atg gaa aag aca ctg aaa ggg ctc aac	2808
Lys Met Arg Asp Ile Leu Asn Met Glu Lys Thr Leu Lys Gly Leu Asn	
880 885 890 895	
agt gag tcc atg aca gaa gaa acc ctg aaa agg gca aag gag att ggg	2856
Ser Glu Ser Met Thr Glu Glu Thr Leu Lys Arg Ala Lys Glu Ile Gly	
900 905 910	
ttc tca gat aag cag att tca aaa tgc ctt ggg ctc act gag gcc cag	2904
Phe Ser Asp Lys Gln Ile Ser Lys Cys Leu Gly Leu Thr Glu Ala Gln	
915 920 925	
aca agg gag ctg agg tta aag aaa aac atc cac cct tgg gtt aaa cag	2952
Thr Arg Glu Leu Arg Leu Lys Lys Asn Ile His Pro Trp Val Lys Gln	
930 935 940	
att gat aca ctg gct gca gaa tac cca tca gta aca aac tat ctc tat	3000
Ile Asp Thr Leu Ala Ala Glu Tyr Pro Ser Val Thr Asn Tyr Leu Tyr	
945 950 955	
gtt acc tac aat ggt cag gag cat gat gtc aat ttt gat gac cat gga	3048
Val Thr Tyr Asn Gly Gln Glu His Asp Val Asn Phe Asp Asp His Gly	
960 965 970 975	
atg atg gtg cta ggc tgt ggt cca tat cac att ggc agc agt gtg gaa	3096
Met Met Val Leu Gly Cys Gly Pro Tyr His Ile Gly Ser Ser Val Glu	
980 985 990	

-continued

ttt gat tgg tgt gct gtc tct agt atc cgc aca ctg cgt caa ctt ggc	3144
Phe Asp Trp Cys Ala Val Ser Ser Ile Arg Thr Leu Arg Gln Leu Gly	
995 1000 1005	
aag aag acg gtg gtg gtg aat tgc aat cct gag act gtg agc aca	3189
Lys Lys Thr Val Val Val Asn Cys Asn Pro Glu Thr Val Ser Thr	
1010 1015 1020	
gac ttt gat gag tgt gac aaa ctg tac ttt gaa gag ttg tcc ttg	3234
Asp Phe Asp Glu Cys Asp Lys Leu Tyr Phe Glu Glu Leu Ser Leu	
1025 1030 1035	
gag aga atc cta gac atc tac cat cag gag gca tgt ggt ggc tgc	3279
Glu Arg Ile Leu Asp Ile Tyr His Gln Glu Ala Cys Gly Gly Cys	
1040 1045 1050	
atc ata tca gtt gga ggc cag att cca aac aac ctg gca gtt cct	3324
Ile Ile Ser Val Gly Gly Gln Ile Pro Asn Asn Leu Ala Val Pro	
1055 1060 1065	
cta tac aag aat ggt gtc aag atc atg ggc aca agc ccc ctg cag	3369
Leu Tyr Lys Asn Gly Val Lys Ile Met Gly Thr Ser Pro Leu Gln	
1070 1075 1080	
atc gac agg gct gag gat cgc tcc atc ttc tca gct gtc ttg gat	3414
Ile Asp Arg Ala Glu Asp Arg Ser Ile Phe Ser Ala Val Leu Asp	
1085 1090 1095	
gag ctg aag gtg gct cag gca cct tgg aaa gct gtt aat act ttg	3459
Glu Leu Lys Val Ala Gln Ala Pro Trp Lys Ala Val Asn Thr Leu	
1100 1105 1110	
aat gaa gca ctg gaa ttt gca aag tct gtg gac tac ccc tgc ttg	3504
Asn Glu Ala Leu Glu Phe Ala Lys Ser Val Asp Tyr Pro Cys Leu	
1115 1120 1125	
ttg agg cct tcc tat gtt ttg agt ggg tct gct atg aat gtg gta	3549
Leu Arg Pro Ser Tyr Val Leu Ser Gly Ser Ala Met Asn Val Val	
1130 1135 1140	
ttc tct gag gat gag atg aaa aaa ttc cta gaa gag gcg act aga	3594
Phe Ser Glu Asp Glu Met Lys Lys Phe Leu Glu Glu Ala Thr Arg	
1145 1150 1155	
gtt tct cag gag cac cca gtg gtc ctg aca aaa ttt gtt gaa ggg	3639
Val Ser Gln Glu His Pro Val Val Leu Thr Lys Phe Val Glu Gly	
1160 1165 1170	
gcc cga gaa gta gaa atg gac gct gtt ggc aaa gat gga agg gtt	3684
Ala Arg Glu Val Glu Met Asp Ala Val Gly Lys Asp Gly Arg Val	
1175 1180 1185	
atc tct cat gcc atc tct gaa cat gtt gaa gat gca ggt gtc cac	3729
Ile Ser His Ala Ile Ser Glu His Val Glu Asp Ala Gly Val His	
1190 1195 1200	
tcg gga gat gcc act ctg atg ctg ccc aca caa acc atc agc caa	3774
Ser Gly Asp Ala Thr Leu Met Leu Pro Thr Gln Thr Ile Ser Gln	
1205 1210 1215	
ggg gcc att gaa aag gtg aag gat gct acc cgg aag att gca aag	3819
Gly Ala Ile Glu Lys Val Lys Asp Ala Thr Arg Lys Ile Ala Lys	
1220 1225 1230	
gct ttt gcc atc tct ggt cca ttc aac gtc caa ttt ctt gtc aaa	3864
Ala Phe Ala Ile Ser Gly Pro Phe Asn Val Gln Phe Leu Val Lys	
1235 1240 1245	
gga aat gat gtc ttg gtg att gag tgt aac ttg aga gct tct cga	3909
Gly Asn Asp Val Leu Val Ile Glu Cys Asn Leu Arg Ala Ser Arg	
1250 1255 1260	
tcc ttc ccc ttt gtt tcc aag act ctt ggg gtt gac ttc att gat	3954
Ser Phe Pro Phe Val Ser Lys Thr Leu Gly Val Asp Phe Ile Asp	
1265 1270 1275	
gtg gcc acc aag gtg atg att gga gag aat gtt gat gag aaa cat	3999
Val Ala Thr Lys Val Met Ile Gly Glu Asn Val Asp Glu Lys His	
1280 1285 1290	

-continued

ctt cca aca	ttg gac cat	ccc ata	att cct gct	gac tat	gtt gca	4044
Leu Pro Thr	Leu Asp His	Pro Ile	Ile Pro Ala	Asp Tyr	Val Ala	
1295		1300		1305		
att aag gct	ccc atg ttt	tcc tgg	ccc cgg ttg	agg gat	gct gac	4089
Ile Lys Ala	Pro Met Phe	Ser Trp	Pro Arg Leu	Arg Asp	Ala Asp	
1310		1315		1320		
ccc att ctg	aga tgt gag	atg gct	tcc act gga	gag gtg	gct tgc	4134
Pro Ile Leu	Arg Cys Glu	Met Ala	Ser Thr Gly	Glu Val	Ala Cys	
1325		1330		1335		
ttt ggt gaa	ggt att cat	aca gcc	ttc cta aag	gca atg	ctt tcc	4179
Phe Gly Glu	Gly Ile His	Thr Ala	Phe Leu Lys	Ala Met	Leu Ser	
1340		1345		1350		
aca gga ttt	aag ata ccc	cag aaa	ggc atc ctg	ata ggc	atc cag	4224
Thr Gly Phe	Lys Ile Pro	Gln Lys	Gly Ile Leu	Ile Gly	Ile Gln	
1355		1360		1365		
caa tca ttc	cgg cca aga	ttc ctt	ggt gtg gct	gaa caa	tta cac	4269
Gln Ser Phe	Arg Pro Arg	Phe Leu	Gly Val Ala	Glu Gln	Leu His	
1370		1375		1380		
aat gaa ggt	ttc aag ctg	ttt gcc	acg gaa gcc	aca tca	gac tgg	4314
Asn Glu Gly	Phe Lys Leu	Phe Ala	Thr Glu Ala	Thr Ser	Asp Trp	
1385		1390		1395		
ctc aac gcc	aac aat gtc	cct gcc	aac cca gtg	gca tgg	cag tct	4359
Leu Asn Ala	Asn Asn Val	Pro Ala	Asn Pro Val	Ala Trp	Pro Ser	
1400		1405		1410		
caa gaa gga	cag aat ccc	agc ctc	tct tcc atc	aga aaa	ttg att	4404
Gln Glu Gly	Gln Asn Pro	Ser Leu	Ser Ser Ile	Arg Lys	Leu Ile	
1415		1420		1425		
aga gat ggc	agc att gac	cta gtg	att aac ctt	ccc aac	aac aac	4449
Arg Asp Gly	Ser Ile Asp	Leu Val	Ile Asn Leu	Pro Asn	Asn Asn	
1430		1435		1440		
act aaa ttt	gtc cat gat	aat tat	gtg att cgg	agg aca	gct gtt	4494
Thr Lys Phe	Val His Asp	Asn Tyr	Val Ile Arg	Arg Thr	Ala Val	
1445		1450		1455		
gat agt gga	atc cct ctc	ctc act	aat ttt cag	gtg acc	aaa ctt	4539
Asp Ser Gly	Ile Pro Leu	Leu Thr	Asn Phe Gln	Val Thr	Lys Leu	
1460		1465		1470		
ttt gct gaa	gct gtg cag	aaa tct	cgc aag gtg	gac tcc	aag agt	4584
Phe Ala Glu	Ala Val Gln	Lys Ser	Arg Lys Val	Asp Ser	Lys Ser	
1475		1480		1485		
ctt ttc cac	tac agg cag	tac agt	gct gga aaa	gca gca	tag	4626
Leu Phe His	Tyr Arg Gln	Tyr Ser	Ala Gly Lys	Ala Ala		
1490		1495		1500		
agatgcagac	acccagccc	cattattaa	tcaacctgag	ccacatgtta	tctaaaggaa	4686
ctgattcaca	actttctcag	agatgaatat	tgataactaa	acttcatttc	agtttacttt	4746
gttatgcctt	aatattctgt	gtcttttgca	attaaattgt	cagtcacttc	ttcaaacct	4806
tacagtcctt	cctaagttac	tcttcatgag	atttcatcca	tttactaata	ctgtattttt	4866
ggtggactag	gcttgccctat	gtgcttatgt	gtagcttttt	actttttatg	gtgctgatta	4926
atggtgatca	aggtaggaaa	agttgctggt	ctattttctg	aactctttct	atactttaag	4986
atactctatt	tttaaacac	tatctgcaaa	ctcaggacac	tttaacaggg	cagaatactc	5046
taaaaacttg	ataaatgaa	atatagattt	aatttatgaa	ccttccatca	tgatgtttgt	5106
gtattgcttc	tttttgatc	ctcattctca	cccatttggc	taatccagga	atattgttat	5166
cccttccat	tatattgaag	ttgagaaatg	tgacagaggc	atttagagta	tgacttttc	5226
ttttcttttt	ctttttcttt	ttttcttttt	gagatggagt	cacactctcc	aggetggagt	5286
gcagtggcac	aatctcggct	cactgcaatt	tgctctccc	aagttcaagc	gattctctg	5346

-continued

```

ctttagacta tggatttctt taaggaatac tggtttgcag ttttgtttctc tggactatat 5406
cagcagatgg tagacagtgt ttatgtagat gtgttgttgt ttttatcatt ggattttaac 5466
ttggccccgag tgaataatc agatttttgt cattcacact ctccccagt tttggaataa 5526
cttgaagta aggttcattc ccttaagacg atggattctg ttgaactatg ggggccaca 5586
ctgcactatt aattccacc actgtaaggg caaggacacc attccttcta catataagaa 5646
aaaagtctct cccaagggc agcctttgtt acttttaaat attttctgtt attacaagtg 5706
ctctaattgt gaacttttaa ataaaatact attaagaggt aaaaaaaaaa aaaaa 5761

```

<210> SEQ ID NO 12

<211> LENGTH: 1500

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 12

```

Met Thr Arg Leu Leu Thr Ala Phe Lys Val Val Arg Thr Leu Lys Thr
1           5           10           15
Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe Ser
20           25           30
Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His Ile
35           40           45
Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His Pro
50           55           60
Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly Tyr
65           70           75           80
Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr Met
85           90           95
Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala Leu
100          105          110
Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys Val
115          120          125
Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp Leu
130          135          140
Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro Ala
145          150          155          160
Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp Lys
165          170          175
Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp Phe
180          185          190
Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys Asp
195          200          205
Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val Asp
210          215          220
Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly Ala
225          230          235          240
Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu Tyr
245          250          255
Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala Glu
260          265          270
Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys Glu
275          280          285
Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala Ala
290          295          300

```

-continued

Gly	Ala	Lys	Thr	Tyr	Lys	Met	Ser	Met	Ala	Asn	Arg	Gly	Gln	Asn	Gln
305					310					315					320
Pro	Val	Leu	Asn	Ile	Thr	Asn	Lys	Gln	Ala	Phe	Ile	Thr	Ala	Gln	Asn
			325						330					335	
His	Gly	Tyr	Ala	Leu	Asp	Asn	Thr	Leu	Pro	Ala	Gly	Trp	Lys	Pro	Leu
			340					345					350		
Phe	Val	Asn	Val	Asn	Asp	Gln	Thr	Asn	Glu	Gly	Ile	Met	His	Glu	Ser
		355					360					365			
Lys	Pro	Phe	Phe	Ala	Val	Gln	Phe	His	Pro	Glu	Val	Thr	Pro	Gly	Pro
		370				375						380			
Ile	Asp	Thr	Glu	Tyr	Leu	Phe	Asp	Ser	Phe	Phe	Ser	Leu	Ile	Lys	Lys
385					390					395					400
Gly	Lys	Ala	Thr	Thr	Ile	Thr	Ser	Val	Leu	Pro	Lys	Pro	Ala	Leu	Val
				405					410					415	
Ala	Ser	Arg	Val	Glu	Val	Ser	Lys	Val	Leu	Ile	Leu	Gly	Ser	Gly	Gly
			420					425						430	
Leu	Ser	Ile	Gly	Gln	Ala	Gly	Glu	Phe	Asp	Tyr	Ser	Gly	Ser	Gln	Ala
		435					440					445			
Val	Lys	Ala	Met	Lys	Glu	Glu	Asn	Val	Lys	Thr	Val	Leu	Met	Asn	Pro
		450				455					460				
Asn	Ile	Ala	Ser	Val	Gln	Thr	Asn	Glu	Val	Gly	Leu	Lys	Gln	Ala	Asp
465					470					475					480
Thr	Val	Tyr	Phe	Leu	Pro	Ile	Thr	Pro	Gln	Phe	Val	Thr	Glu	Val	Ile
				485					490					495	
Lys	Ala	Glu	Gln	Pro	Asp	Gly	Leu	Ile	Leu	Gly	Met	Gly	Gly	Gln	Thr
			500					505						510	
Ala	Leu	Asn	Cys	Gly	Val	Glu	Leu	Phe	Lys	Arg	Gly	Val	Leu	Lys	Glu
			515				520						525		
Tyr	Gly	Val	Lys	Val	Leu	Gly	Thr	Ser	Val	Glu	Ser	Ile	Met	Ala	Thr
						535					540				
Glu	Asp	Arg	Gln	Leu	Phe	Ser	Asp	Lys	Leu	Asn	Glu	Ile	Asn	Glu	Lys
545					550					555					560
Ile	Ala	Pro	Ser	Phe	Ala	Val	Glu	Ser	Ile	Glu	Asp	Ala	Leu	Lys	Ala
				565						570				575	
Ala	Asp	Thr	Ile	Gly	Tyr	Pro	Val	Met	Ile	Arg	Ser	Ala	Tyr	Ala	Leu
			580					585					590		
Gly	Gly	Leu	Gly	Ser	Gly	Ile	Cys	Pro	Asn	Arg	Glu	Thr	Leu	Met	Asp
		595					600					605			
Leu	Ser	Thr	Lys	Ala	Phe	Ala	Met	Thr	Asn	Gln	Ile	Leu	Val	Glu	Lys
						615					620				
Ser	Val	Thr	Gly	Trp	Lys	Glu	Ile	Glu	Tyr	Glu	Val	Val	Arg	Asp	Ala
625					630					635					640
Asp	Asp	Asn	Cys	Val	Thr	Val	Cys	Asn	Met	Glu	Asn	Val	Asp	Ala	Met
				645					650					655	
Gly	Val	His	Thr	Gly	Asp	Ser	Val	Val	Val	Ala	Pro	Ala	Gln	Thr	Leu
			660					665						670	
Ser	Asn	Ala	Glu	Phe	Gln	Met	Leu	Arg	Arg	Thr	Ser	Ile	Asn	Val	Val
			675				680						685		
Arg	His	Leu	Gly	Ile	Val	Gly	Glu	Cys	Asn	Ile	Gln	Phe	Ala	Leu	His
						695					700				
Pro	Thr	Ser	Met	Glu	Tyr	Cys	Ile	Ile	Glu	Val	Asn	Ala	Arg	Leu	Ser
705					710					715					720
Arg	Ser	Ser	Ala	Leu	Ala	Ser	Lys	Ala	Thr	Gly	Tyr	Pro	Leu	Ala	Phe
				725					730						735

-continued

Ile Ala Ala Lys Ile Ala Leu Gly Ile Pro Leu Pro Glu Ile Lys Asn
 740 745 750
 Val Val Ser Gly Lys Thr Ser Ala Cys Phe Glu Pro Ser Leu Asp Tyr
 755 760 765
 Met Val Thr Lys Ile Pro Arg Trp Asp Leu Asp Arg Phe His Gly Thr
 770 775 780
 Ser Ser Arg Ile Gly Ser Ser Met Lys Ser Val Gly Glu Val Met Ala
 785 790 795 800
 Ile Gly Arg Thr Phe Glu Glu Ser Phe Gln Lys Ala Leu Arg Met Cys
 805 810 815
 His Pro Ser Ile Glu Gly Phe Thr Pro Arg Leu Pro Met Asn Lys Glu
 820 825 830
 Trp Pro Ser Asn Leu Asp Leu Arg Lys Glu Leu Ser Glu Pro Ser Ser
 835 840 845
 Thr Arg Ile Tyr Ala Ile Ala Lys Ala Ile Asp Asp Asn Met Ser Leu
 850 855 860
 Asp Glu Ile Glu Lys Leu Thr Tyr Ile Asp Lys Trp Phe Leu Tyr Lys
 865 870 875 880
 Met Arg Asp Ile Leu Asn Met Glu Lys Thr Leu Lys Gly Leu Asn Ser
 885 890 895
 Glu Ser Met Thr Glu Glu Thr Leu Lys Arg Ala Lys Glu Ile Gly Phe
 900 905 910
 Ser Asp Lys Gln Ile Ser Lys Cys Leu Gly Leu Thr Glu Ala Gln Thr
 915 920 925
 Arg Glu Leu Arg Leu Lys Lys Asn Ile His Pro Trp Val Lys Gln Ile
 930 935 940
 Asp Thr Leu Ala Ala Glu Tyr Pro Ser Val Thr Asn Tyr Leu Tyr Val
 945 950 955 960
 Thr Tyr Asn Gly Gln Glu His Asp Val Asn Phe Asp Asp His Gly Met
 965 970 975
 Met Val Leu Gly Cys Gly Pro Tyr His Ile Gly Ser Ser Val Glu Phe
 980 985 990
 Asp Trp Cys Ala Val Ser Ser Ile Arg Thr Leu Arg Gln Leu Gly Lys
 995 1000 1005
 Lys Thr Val Val Val Asn Cys Asn Pro Glu Thr Val Ser Thr Asp
 1010 1015 1020
 Phe Asp Glu Cys Asp Lys Leu Tyr Phe Glu Glu Leu Ser Leu Glu
 1025 1030 1035
 Arg Ile Leu Asp Ile Tyr His Gln Glu Ala Cys Gly Gly Cys Ile
 1040 1045 1050
 Ile Ser Val Gly Gly Gln Ile Pro Asn Asn Leu Ala Val Pro Leu
 1055 1060 1065
 Tyr Lys Asn Gly Val Lys Ile Met Gly Thr Ser Pro Leu Gln Ile
 1070 1075 1080
 Asp Arg Ala Glu Asp Arg Ser Ile Phe Ser Ala Val Leu Asp Glu
 1085 1090 1095
 Leu Lys Val Ala Gln Ala Pro Trp Lys Ala Val Asn Thr Leu Asn
 1100 1105 1110
 Glu Ala Leu Glu Phe Ala Lys Ser Val Asp Tyr Pro Cys Leu Leu
 1115 1120 1125
 Arg Pro Ser Tyr Val Leu Ser Gly Ser Ala Met Asn Val Val Phe
 1130 1135 1140
 Ser Glu Asp Glu Met Lys Lys Phe Leu Glu Glu Ala Thr Arg Val

-continued

1145	1150	1155
Ser Gln Glu His Pro Val Val Leu Thr Lys Phe Val Glu Gly Ala		
1160	1165	1170
Arg Glu Val Glu Met Asp Ala Val Gly Lys Asp Gly Arg Val Ile		
1175	1180	1185
Ser His Ala Ile Ser Glu His Val Glu Asp Ala Gly Val His Ser		
1190	1195	1200
Gly Asp Ala Thr Leu Met Leu Pro Thr Gln Thr Ile Ser Gln Gly		
1205	1210	1215
Ala Ile Glu Lys Val Lys Asp Ala Thr Arg Lys Ile Ala Lys Ala		
1220	1225	1230
Phe Ala Ile Ser Gly Pro Phe Asn Val Gln Phe Leu Val Lys Gly		
1235	1240	1245
Asn Asp Val Leu Val Ile Glu Cys Asn Leu Arg Ala Ser Arg Ser		
1250	1255	1260
Phe Pro Phe Val Ser Lys Thr Leu Gly Val Asp Phe Ile Asp Val		
1265	1270	1275
Ala Thr Lys Val Met Ile Gly Glu Asn Val Asp Glu Lys His Leu		
1280	1285	1290
Pro Thr Leu Asp His Pro Ile Ile Pro Ala Asp Tyr Val Ala Ile		
1295	1300	1305
Lys Ala Pro Met Phe Ser Trp Pro Arg Leu Arg Asp Ala Asp Pro		
1310	1315	1320
Ile Leu Arg Cys Glu Met Ala Ser Thr Gly Glu Val Ala Cys Phe		
1325	1330	1335
Gly Glu Gly Ile His Thr Ala Phe Leu Lys Ala Met Leu Ser Thr		
1340	1345	1350
Gly Phe Lys Ile Pro Gln Lys Gly Ile Leu Ile Gly Ile Gln Gln		
1355	1360	1365
Ser Phe Arg Pro Arg Phe Leu Gly Val Ala Glu Gln Leu His Asn		
1370	1375	1380
Glu Gly Phe Lys Leu Phe Ala Thr Glu Ala Thr Ser Asp Trp Leu		
1385	1390	1395
Asn Ala Asn Asn Val Pro Ala Asn Pro Val Ala Trp Pro Ser Gln		
1400	1405	1410
Glu Gly Gln Asn Pro Ser Leu Ser Ser Ile Arg Lys Leu Ile Arg		
1415	1420	1425
Asp Gly Ser Ile Asp Leu Val Ile Asn Leu Pro Asn Asn Asn Thr		
1430	1435	1440
Lys Phe Val His Asp Asn Tyr Val Ile Arg Arg Thr Ala Val Asp		
1445	1450	1455
Ser Gly Ile Pro Leu Leu Thr Asn Phe Gln Val Thr Lys Leu Phe		
1460	1465	1470
Ala Glu Ala Val Gln Lys Ser Arg Lys Val Asp Ser Lys Ser Leu		
1475	1480	1485
Phe His Tyr Arg Gln Tyr Ser Ala Gly Lys Ala Ala		
1490	1495	1500

<210> SEQ ID NO 13
 <211> LENGTH: 5761
 <212> TYPE: DNA
 <213> ORGANISM: Homo sapiens
 <220> FEATURE:
 <221> NAME/KEY: CDS
 <222> LOCATION: (124)..(4626)

-continued

<400> SEQUENCE: 13

gtcagcctta aacctgact gcaccctcc cagatttctt ttacattaac taaaaagtct	60
tatcacacaa tctcataaaa tttatgtaat ttcatttaat tttagccaca aatcatcttc	120
aaa atg acg agg att att aca gct ttc aaa gtg gtg agg aca ctg aag	168
Met Thr Arg Ile Ile Thr Ala Phe Lys Val Val Arg Thr Leu Lys	
1 5 10 15	
act ggt ttt ggc ttt acc aat gtg act gca cac caa aaa tgg aaa ttt	216
Thr Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe	
20 25 30	
tca aga cct ggc atc agg ctc ctt tct gtc aag gca cag aca gca cac	264
Ser Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His	
35 40 45	
att gtc ctg gaa gat gga act aag atg aaa ggt tac tcc ttt ggc cat	312
Ile Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His	
50 55 60	
cca tcc tct gtt gct ggt gaa gtg gtt ttt aat act ggc ctg gga ggg	360
Pro Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly	
65 70 75	
tac cca gaa gct att act gac cct gcc tac aaa gga cag att ctc aca	408
Tyr Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr	
80 85 90 95	
atg gcc aac cct att att ggg aat ggt gga gct cct gat act act gct	456
Met Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala	
100 105 110	
ctg gat gaa ctg gga ctt agc aaa tat ttg gag tct aat gga atc aag	504
Leu Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys	
115 120 125	
gtt tca ggt ttg ctg gtg ctg gat tat agt aaa gac tac aac cac tgg	552
Val Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp	
130 135 140	
ctg gct acc aag agt tta ggg caa tgg cta cag gaa gaa aag gtt cct	600
Leu Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro	
145 150 155	
gca att tat gga gtg gac aca aga atg ctg act aaa ata att cgg gat	648
Ala Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp	
160 165 170 175	
aag ggt acc atg ctt ggg aag att gaa ttt gaa ggt cag cct gtg gat	696
Lys Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp	
180 185 190	
ttt gtg gat cca aat aaa cag aat ttg att gct gag gtt tca acc aag	744
Phe Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys	
195 200 205	
gat gtc aaa gtg tac ggc aaa gga aac ccc aca aaa gtg gta gct gta	792
Asp Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val	
210 215 220	
gac tgt ggg att aaa aac aat gta atc cgc ctg cta gta aag cga gga	840
Asp Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly	
225 230 235	
gct gaa gtg cac tta gtt ccc tgg aac cat gat ttc acc aag atg gag	888
Ala Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu	
240 245 250 255	
tat gat ggg att ttg atc gcg gga gga ccg ggg aac cca gct ctt gca	936
Tyr Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala	
260 265 270	
gaa cca cta att cag aat gtc aga aag att ttg gag agt gat cgc aag	984
Glu Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys	
275 280 285	
gag cca ttg ttt gga atc agt aca gga aac tta ata aca gga ttg gct	1032
Glu Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala	

-continued

290	295	300	
gct ggt gcc aaa acc tac aag atg tcc atg gcc aac aga ggg cag aat Ala Gly Ala Lys Thr Tyr Lys Met Ser Met Ala Asn Arg Gly Gln Asn 305 310 315			1080
cag cct gtt ttg aat atc aca aac aaa cag gct ttc att act gct cag Gln Pro Val Leu Asn Ile Thr Asn Lys Gln Ala Phe Ile Thr Ala Gln 320 325 330 335			1128
aat cat ggc tat gcc ttg gac aac acc ctc cct gct ggc tgg aaa cca Asn His Gly Tyr Ala Leu Asp Asn Thr Leu Pro Ala Gly Trp Lys Pro 340 345 350			1176
ctt ttt gtg aat gtc aac gat caa aca aat gag ggg att atg cat gag Leu Phe Val Asn Val Asn Asp Gln Thr Asn Glu Gly Ile Met His Glu 355 360 365			1224
agc aaa ccc ttc ttc gct gtg cag ttc cac cca gag gtc acc ccg ggg Ser Lys Pro Phe Phe Ala Val Gln Phe His Pro Glu Val Thr Pro Gly 370 375 380			1272
cca ata gac act gag tac ctg ttt gat tcc ttt ttc tca ctg ata aag Pro Ile Asp Thr Glu Tyr Leu Phe Asp Ser Phe Phe Ser Leu Ile Lys 385 390 395			1320
aaa gga aaa gct acc acc att aca tca gtc tta ccg aag cca gca cta Lys Gly Lys Ala Thr Thr Ile Thr Ser Val Leu Pro Lys Pro Ala Leu 400 405 410 415			1368
gtt gca tct cgg gtt gag gtt tcc aaa gtc ctt att cta gga tca gga Val Ala Ser Arg Val Glu Val Ser Lys Val Leu Ile Leu Gly Ser Gly 420 425 430			1416
ggt ctg tcc att ggt cag gct gga gaa ttt gat tac tca gga tct caa Gly Leu Ser Ile Gly Gln Ala Gly Glu Phe Asp Tyr Ser Gly Ser Gln 435 440 445			1464
gct gta aaa gcc atg aag gaa gaa aat gtc aaa act gtt ctg atg aac Ala Val Lys Ala Met Lys Glu Glu Asn Val Lys Thr Val Leu Met Asn 450 455 460			1512
cca aac att gca tca gtc cag acc aat gag gtg ggc tta aag caa gcg Pro Asn Ile Ala Ser Val Gln Thr Asn Glu Val Gly Leu Lys Gln Ala 465 470 475			1560
gat act gtc tac ttt ctt ccc atc acc cct cag ttt gtc aca gag gtc Asp Thr Val Tyr Phe Leu Pro Ile Thr Pro Gln Phe Val Thr Glu Val 480 485 490 495			1608
atc aag gca gaa cag cca gat ggg tta att ctg ggc atg ggt ggc cag Ile Lys Ala Glu Gln Pro Asp Gly Leu Ile Leu Gly Met Gly Gly Gln 500 505 510			1656
aca gct ctg aac tgt gga gtg gaa cta ttc aag aga ggt gtg ctc aag Thr Ala Leu Asn Cys Gly Val Glu Leu Phe Lys Arg Gly Val Leu Lys 515 520 525			1704
gaa tat ggt gtg aaa gtc ctg gga act tca gtt gag tcc att atg gct Glu Tyr Gly Val Lys Val Leu Gly Thr Ser Val Glu Ser Ile Met Ala 530 535 540			1752
acg gaa gac agg cag ctg ttt tca gat aaa cta aat gag atc aat gaa Thr Glu Asp Arg Gln Leu Phe Ser Asp Lys Leu Asn Glu Ile Asn Glu 545 550 555			1800
aag att gct cca agt ttt gca gtg gaa tcg att gag gat gca ctg aag Lys Ile Ala Pro Ser Phe Ala Val Glu Ser Ile Glu Asp Ala Leu Lys 560 565 570 575			1848
gca gca gac acc att ggc tac cca gtg atg atc cgt tcc gcc tat gca Ala Ala Asp Thr Ile Gly Tyr Pro Val Met Ile Arg Ser Ala Tyr Ala 580 585 590			1896
ctg ggt ggg tta ggc tca ggc atc tgt ccc aac aga gag act ttg atg Leu Gly Gly Leu Gly Ser Gly Ile Cys Pro Asn Arg Glu Thr Leu Met 595 600 605			1944
gac ctc agc aca aag gcc ttt gct atg acc aac caa att ctg gtg gag Asp Leu Ser Thr Lys Ala Phe Ala Met Thr Asn Gln Ile Leu Val Glu			1992

-continued

610	615	620	
aag tca gtg aca ggt tgg Lys Ser Val Thr Gly Trp 625	aaa gaa ata gaa tat Lys Glu Ile Glu Tyr 630	gaa gtg gtt cga gat Glu Val Val Arg Asp 635	2040
gct gat gac aat tgt gtc Ala Asp Asp Asn Cys Val 640	act gtc tgt aac atg Thr Val Cys Asn Met 645	gaa aat gtt gat gcc Glu Asn Val Asp Ala 650 655	2088
atg ggt gtt cac aca ggt Met Gly Val His Thr Gly 660	gac tca gtt gtt gtg Asp Ser Val Val Val 665	gct cct gcc cag aca Ala Pro Ala Gln Thr 670	2136
ctc tcc aat gcc gag ttt Leu Ser Asn Ala Glu Phe 675	cag atg ttg aga cgt act Gln Met Leu Arg Arg Thr 680	tca atc aat gtt Ser Ile Asn Val 685	2184
gtt cgc cac ttg ggc att Val Arg His Leu Gly Ile 690	gtg ggt gaa tgc aac Val Gly Glu Cys Asn Ile 695	att cag ttt gcc ctt Ile Gln Phe Ala Leu 700	2232
cat cct acc tca atg gaa His Pro Thr Ser Met Glu 705	tac tgc atc att gaa Tyr Cys Ile Ile Glu Val 710 715	gtg aat gcc aga ctg Val Asn Ala Arg Leu	2280
tcc cga agc tct gct ctg Ser Arg Ser Ser Ala Leu 720 725	gcc tca aaa gcc act Ala Ser Lys Ala Thr Gly 730	ggc tac cca ttg gca Tyr Pro Leu Ala 735	2328
ttc att gct gca aag att Phe Ile Ala Ala Lys Ile 740	gcc cta gga atc cca Ala Leu Gly Ile Pro Leu 745	ctt cca gaa att aag Pro Leu Pro Glu Ile Lys 750	2376
aac gtc gta tcc ggg aag Asn Val Val Ser Gly Lys 755	aca tca gcc tgt ttt Thr Ser Ala Cys Phe Glu 760	cct agc ctg gat Pro Ser Leu Asp 765	2424
tac atg gtc acc aag att Tyr Met Val Thr Lys Ile 770	ccc cgc tgg gat ctt Pro Arg Trp Asp Leu Asp 775 780	gac cgt ttt cat gga Arg Phe His Gly	2472
aca tct agc cga att ggt Thr Ser Ser Arg Ile Gly 785	agc tct atg aaa agt Ser Ser Met Lys Ser Val 790 795	gta gga gag gtc atg Val Gly Glu Val Met	2520
gct att ggt cgt acc ttt Ala Ile Gly Arg Thr Phe 800 805	gag gag agt ttc cag Glu Glu Ser Phe Gln Lys 810	aaa gct tta cgg atg Ala Leu Arg Met 815	2568
tgc cac cca tct ata gaa Cys His Pro Ser Ile Glu 820	ggt ttc act ccc cgt Gly Phe Thr Pro Arg Leu 825	ctc cca atg aac aaa Pro Met Asn Lys 830	2616
gaa tgg cca tct aat tta Glu Trp Pro Ser Asn Leu 835	gat ctt aga aaa gag Asp Leu Arg Lys Glu Leu 840	ttg tct gaa cca agc Leu Ser Glu Pro Ser 845	2664
agc acg cgt atc tat gcc Ser Thr Arg Ile Tyr Ala 850	att gcc aag gcc att Ala Lys Ala Ile Asp Asp 855 860	gat gac aac atg tcc Asp Asn Met Ser	2712
ctt gat gag att gag aag Leu Asp Glu Ile Glu Lys 865 870	ctc aca tac att gac Leu Thr Tyr Ile Asp Lys 875	ttg ttt ttg tat Trp Phe Leu Tyr	2760
aag atg cgt gat att tta Lys Met Arg Asp Ile Leu 880 885	aac atg gaa aag aca Asn Met Glu Lys Thr Leu 890	ctg aaa ggg ctc aac Leu Lys Gly Leu Asn 895	2808
agt gag tcc atg aca gaa Ser Glu Ser Met Thr Glu 900	gaa gaa acc ctg aaa Glu Glu Thr Leu Lys Arg 905	gca aag gag att ggg Ala Lys Glu Ile Gly 910	2856
ttc tca gat aag cag att Phe Ser Asp Lys Gln Ile 915	tca aaa tgc ctt ggg Ser Lys Cys Leu Gly Leu 920	ctc act gag gcc cag Leu Thr Glu Ala Gln 925	2904
aca agg gag ctg agg tta Thr Arg Glu Leu Arg Leu 930	aag aaa aac atc cac Lys Lys Asn Ile His Pro 935	cct tgg gtt aaa cag Pro Trp Val Lys Gln 940	2952

-continued

930	935	940	
att gat aca ctg gct gca gaa tac cca tca gta aca aac tat ctc tat Ile Asp Thr Leu Ala Ala Glu Tyr Pro Ser Val Thr Asn Tyr Leu Tyr 945 950 955			3000
gtt acc tac aat ggt cag gag cat gat gtc aat ttt gat gac cat gga Val Thr Tyr Asn Gly Gln Glu His Asp Val Asn Phe Asp Asp His Gly 960 965 970 975			3048
atg atg gtg cta ggc tgt ggt cca tat cac att ggc agc agt gtg gaa Met Met Val Leu Gly Cys Gly Pro Tyr His Ile Gly Ser Ser Val Glu 980 985 990			3096
ttt gat tgg tgt gct gtc tct agt atc cgc aca ctg cgt caa ctt ggc Phe Asp Trp Cys Ala Val Ser Ser Ile Arg Thr Leu Arg Gln Leu Gly 995 1000 1005			3144
aag aag acg gtg gtg gtg aat tgc aat cct gag act gtg agc aca Lys Lys Thr Val Val Val Asn Cys Asn Pro Glu Thr Val Ser Thr 1010 1015 1020			3189
gac ttt gat gag tgt gac aaa ctg tac ttt gaa gag ttg tcc ttg Asp Phe Asp Glu Cys Asp Lys Leu Tyr Phe Glu Glu Leu Ser Leu 1025 1030 1035			3234
gag aga atc cta gac atc tac cat cag gag gca tgt ggt ggc tgc Glu Arg Ile Leu Asp Ile Tyr His Gln Glu Ala Cys Gly Gly Cys 1040 1045 1050			3279
atc ata tca gtt gga ggc cag att cca aac aac ctg gca gtt cct Ile Ile Ser Val Gly Gly Gln Ile Pro Asn Asn Leu Ala Val Pro 1055 1060 1065			3324
cta tac aag aat ggt gtc aag atc atg ggc aca agc ccc ctg cag Leu Tyr Lys Asn Gly Val Lys Ile Met Gly Thr Ser Pro Leu Gln 1070 1075 1080			3369
atc gac agg gct gag gat cgc tcc atc ttc tca gct gtc ttg gat Ile Asp Arg Ala Glu Asp Arg Ser Ile Phe Ser Ala Val Leu Asp 1085 1090 1095			3414
gag ctg aag gtg gct cag gca cct tgg aaa gct gtt aat act ttg Glu Leu Lys Val Ala Gln Ala Pro Trp Lys Ala Val Asn Thr Leu 1100 1105 1110			3459
aat gaa gca ctg gaa ttt gca aag tct gtg gac tac ccc tgc ttg Asn Glu Ala Leu Glu Phe Ala Lys Ser Val Asp Tyr Pro Cys Leu 1115 1120 1125			3504
ttg agg cct tcc tat gtt ttg agt ggg tct gct atg aat gtg gta Leu Arg Pro Ser Tyr Val Leu Ser Gly Ser Ala Met Asn Val Val 1130 1135 1140			3549
ttc tct gag gat gag atg aaa aaa ttc cta gaa gag gcg act aga Phe Ser Glu Asp Glu Met Lys Lys Phe Leu Glu Glu Ala Thr Arg 1145 1150 1155			3594
gtt tct cag gag cac cca gtg gtc ctg aca aaa ttt gtt gaa ggg Val Ser Gln Glu His Pro Val Val Leu Thr Lys Phe Val Glu Gly 1160 1165 1170			3639
gcc cga gaa gta gaa atg gac gct gtt ggc aaa gat gga agg gtt Ala Arg Glu Val Glu Met Asp Ala Val Gly Lys Asp Gly Arg Val 1175 1180 1185			3684
atc tct cat gcc atc tct gaa cat gtt gaa gat gca ggt gtc cac Ile Ser His Ala Ile Ser Glu His Val Glu Asp Ala Gly Val His 1190 1195 1200			3729
tcg gga gat gcc act ctg atg ctg ccc aca caa acc atc agc caa Ser Gly Asp Ala Thr Leu Met Leu Pro Thr Gln Thr Ile Ser Gln 1205 1210 1215			3774
ggg gcc att gaa aag gtg aag gat gct acc cgg aag att gca aag Gly Ala Ile Glu Lys Val Lys Asp Ala Thr Arg Lys Ile Ala Lys 1220 1225 1230			3819
gct ttt gcc atc tct ggt cca ttc aac gtc caa ttt ctt gtc aaa Ala Phe Ala Ile Ser Gly Pro Phe Asn Val Gln Phe Leu Val Lys 1235			3864

-continued

1235	1240	1245	
gga aat gat gtc ttg gtg att gag tgt aac ttg aga gct tct cga Gly Asn Asp Val Leu Val Ile Glu Cys Asn Leu Arg Ala Ser Arg 1250	1255	1260	3909
tcc ttc ccc ttt gtt tcc aag act ctt ggg gtt gac ttc att gat Ser Phe Pro Phe Val Ser Lys Thr Leu Gly Val Asp Phe Ile Asp 1265	1270	1275	3954
gtg gcc acc aag gtg atg att gga gag aat gtt gat gag aaa cat Val Ala Thr Lys Val Met Ile Gly Glu Asn Val Asp Glu Lys His 1280	1285	1290	3999
ctt cca aca ttg gac cat ccc ata att cct gct gac tat gtt gca Leu Pro Thr Leu Asp His Pro Ile Ile Pro Ala Asp Tyr Val Ala 1295	1300	1305	4044
att aag gct ccc atg ttt tcc tgg ccc cgg ttg agg gat gct gac Ile Lys Ala Pro Met Phe Ser Trp Pro Arg Leu Arg Asp Ala Asp 1310	1315	1320	4089
ccc att ctg aga tgt gag atg gct tcc act gga gag gtg gct tgc Pro Ile Leu Arg Cys Glu Met Ala Ser Thr Gly Glu Val Ala Cys 1325	1330	1335	4134
ttt ggt gaa ggt att cat aca gcc ttc cta aag gca atg ctt tcc Phe Gly Glu Gly Ile His Thr Ala Phe Leu Lys Ala Met Leu Ser 1340	1345	1350	4179
aca gga ttt aag ata ccc cag aaa ggc atc ctg ata ggc atc cag Thr Gly Phe Lys Ile Pro Gln Lys Gly Ile Leu Ile Gly Ile Gln 1355	1360	1365	4224
caa tca ttc cgg cca aga ttc ctt ggt gtg gct gaa caa tta cac Gln Ser Phe Arg Pro Arg Phe Leu Gly Val Ala Glu Gln Leu His 1370	1375	1380	4269
aat gaa ggt ttc aag ctg ttt gcc acg gaa gcc aca tca gac tgg Asn Glu Gly Phe Lys Leu Phe Ala Thr Glu Ala Thr Ser Asp Trp 1385	1390	1395	4314
ctc aac gcc aac aat gtc cct gcc aac cca gtg gca tgg ccg tct Leu Asn Ala Asn Asn Val Pro Ala Asn Pro Val Ala Trp Pro Ser 1400	1405	1410	4359
caa gaa gga cag aat ccc agc ctc tct tcc atc aga aaa ttg att Gln Glu Gly Gln Asn Pro Ser Leu Ser Ser Ile Arg Lys Leu Ile 1415	1420	1425	4404
aga gat ggc agc att gac cta gtg att aac ctt ccc aac aac aac Arg Asp Gly Ser Ile Asp Leu Val Ile Asn Leu Pro Asn Asn Asn 1430	1435	1440	4449
act aaa ttt gtc cat gat aat tat gtg att cgg agg aca gct gtt Thr Lys Phe Val His Asp Asn Tyr Val Ile Arg Arg Thr Ala Val 1445	1450	1455	4494
gat agt gga atc cct ctc ctc act aat ttt cag gtg acc aaa ctt Asp Ser Gly Ile Pro Leu Leu Thr Asn Phe Gln Val Thr Lys Leu 1460	1465	1470	4539
ttt gct gaa gct gtg cag aaa tct cgc aag gtg gac tcc aag agt Phe Ala Glu Ala Val Gln Lys Ser Arg Lys Val Asp Ser Lys Ser 1475	1480	1485	4584
ctt ttc cac tac agg cag tac agt gct gga aaa gca gca tag Leu Phe His Tyr Arg Gln Tyr Ser Ala Gly Lys Ala Ala 1490	1495	1500	4626
agatgcagac accccagccc cattattaa tcaacctgag ccacatgta tctaaaggaa			4686
ctgattcaca actttctcag agatgaatat tgataactaa acttcatttc agtttacttt			4746
gttatgcctt aatattctgt gtcttttgca attaaattgt cagtcacttc ttcaaacct			4806
tacagtcctt cctaagttac tcttcatgag atttcatcca tttactaata ctgtattttt			4866
ggtggactag gcttgcctat gtgcttatgt gtagcttttt actttttatg gtgctgatta			4926

-continued

```

atggtgatca aggtaggaaa agttgctggt ctattttctg aactctttct atactttaag 4986
atactctatt tttaaaacac tatctgcaaa ctcaggacac tttaacaggg cagaatactc 5046
taaaaacttg ataaaatgaa atatagattt aatttatgaa ccttccatca tgatgtttgt 5106
gtattgcttc tttttggatc ctcattctca cccatttggc taatccagga atattgttat 5166
cccttcccat tatattgaag ttgagaaatg tgacagaggg atttagagta tggacttttc 5226
ttttcttttt cttttctttt tttttctttt gagatggagt cacactctcc aggetggagt 5286
gcagtggcac aatctcggct cactgcaatt tgcgtctccc aagttcaagc gattctcctg 5346
ctttagacta tggatttctt taaggaatac tggtttgcag ttttgtttc tggactatat 5406
cagcagatgg tagacagtgt ttatgtagat gtgtgtttgt ttttatcatt ggattttaac 5466
ttggcccgag tgaataatc agatttttgt cattcacact ctccccagt tttggaataa 5526
cttgaagta aggttcattc ccttaagacg atggattctg ttgaactatg ggggtcccaca 5586
ctgcactatt aattccacc actgtaaggg caaggacacc attccttcta catataagaa 5646
aaaagtctct cccaagggc agcctttggt acttttaaat attttctgtt attacaagtg 5706
ctctaattgt gaacttttaa ataaaatact attaagaggt aaaaaaaaa aaaaa 5761

```

<210> SEQ ID NO 14

<211> LENGTH: 1500

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 14

```

Met Thr Arg Ile Ile Thr Ala Phe Lys Val Val Arg Thr Leu Lys Thr
1           5           10           15
Gly Phe Gly Phe Thr Asn Val Thr Ala His Gln Lys Trp Lys Phe Ser
20           25           30
Arg Pro Gly Ile Arg Leu Leu Ser Val Lys Ala Gln Thr Ala His Ile
35           40           45
Val Leu Glu Asp Gly Thr Lys Met Lys Gly Tyr Ser Phe Gly His Pro
50           55           60
Ser Ser Val Ala Gly Glu Val Val Phe Asn Thr Gly Leu Gly Gly Tyr
65           70           75           80
Pro Glu Ala Ile Thr Asp Pro Ala Tyr Lys Gly Gln Ile Leu Thr Met
85           90           95
Ala Asn Pro Ile Ile Gly Asn Gly Gly Ala Pro Asp Thr Thr Ala Leu
100          105          110
Asp Glu Leu Gly Leu Ser Lys Tyr Leu Glu Ser Asn Gly Ile Lys Val
115          120          125
Ser Gly Leu Leu Val Leu Asp Tyr Ser Lys Asp Tyr Asn His Trp Leu
130          135          140
Ala Thr Lys Ser Leu Gly Gln Trp Leu Gln Glu Glu Lys Val Pro Ala
145          150          155          160
Ile Tyr Gly Val Asp Thr Arg Met Leu Thr Lys Ile Ile Arg Asp Lys
165          170          175
Gly Thr Met Leu Gly Lys Ile Glu Phe Glu Gly Gln Pro Val Asp Phe
180          185          190
Val Asp Pro Asn Lys Gln Asn Leu Ile Ala Glu Val Ser Thr Lys Asp
195          200          205
Val Lys Val Tyr Gly Lys Gly Asn Pro Thr Lys Val Val Ala Val Asp
210          215          220
Cys Gly Ile Lys Asn Asn Val Ile Arg Leu Leu Val Lys Arg Gly Ala
225          230          235          240

```

-continued

Glu Val His Leu Val Pro Trp Asn His Asp Phe Thr Lys Met Glu Tyr
 245 250 255
 Asp Gly Ile Leu Ile Ala Gly Gly Pro Gly Asn Pro Ala Leu Ala Glu
 260 265 270
 Pro Leu Ile Gln Asn Val Arg Lys Ile Leu Glu Ser Asp Arg Lys Glu
 275 280 285
 Pro Leu Phe Gly Ile Ser Thr Gly Asn Leu Ile Thr Gly Leu Ala Ala
 290 295 300
 Gly Ala Lys Thr Tyr Lys Met Ser Met Ala Asn Arg Gly Gln Asn Gln
 305 310 315 320
 Pro Val Leu Asn Ile Thr Asn Lys Gln Ala Phe Ile Thr Ala Gln Asn
 325 330 335
 His Gly Tyr Ala Leu Asp Asn Thr Leu Pro Ala Gly Trp Lys Pro Leu
 340 345 350
 Phe Val Asn Val Asn Asp Gln Thr Asn Glu Gly Ile Met His Glu Ser
 355 360 365
 Lys Pro Phe Phe Ala Val Gln Phe His Pro Glu Val Thr Pro Gly Pro
 370 375 380
 Ile Asp Thr Glu Tyr Leu Phe Asp Ser Phe Phe Ser Leu Ile Lys Lys
 385 390 395 400
 Gly Lys Ala Thr Thr Ile Thr Ser Val Leu Pro Lys Pro Ala Leu Val
 405 410 415
 Ala Ser Arg Val Glu Val Ser Lys Val Leu Ile Leu Gly Ser Gly Gly
 420 425 430
 Leu Ser Ile Gly Gln Ala Gly Glu Phe Asp Tyr Ser Gly Ser Gln Ala
 435 440 445
 Val Lys Ala Met Lys Glu Glu Asn Val Lys Thr Val Leu Met Asn Pro
 450 455 460
 Asn Ile Ala Ser Val Gln Thr Asn Glu Val Gly Leu Lys Gln Ala Asp
 465 470 475 480
 Thr Val Tyr Phe Leu Pro Ile Thr Pro Gln Phe Val Thr Glu Val Ile
 485 490 495
 Lys Ala Glu Gln Pro Asp Gly Leu Ile Leu Gly Met Gly Gly Gln Thr
 500 505 510
 Ala Leu Asn Cys Gly Val Glu Leu Phe Lys Arg Gly Val Leu Lys Glu
 515 520 525
 Tyr Gly Val Lys Val Leu Gly Thr Ser Val Glu Ser Ile Met Ala Thr
 530 535 540
 Glu Asp Arg Gln Leu Phe Ser Asp Lys Leu Asn Glu Ile Asn Glu Lys
 545 550 555 560
 Ile Ala Pro Ser Phe Ala Val Glu Ser Ile Glu Asp Ala Leu Lys Ala
 565 570 575
 Ala Asp Thr Ile Gly Tyr Pro Val Met Ile Arg Ser Ala Tyr Ala Leu
 580 585 590
 Gly Gly Leu Gly Ser Gly Ile Cys Pro Asn Arg Glu Thr Leu Met Asp
 595 600 605
 Leu Ser Thr Lys Ala Phe Ala Met Thr Asn Gln Ile Leu Val Glu Lys
 610 615 620
 Ser Val Thr Gly Trp Lys Glu Ile Glu Tyr Glu Val Val Arg Asp Ala
 625 630 635 640
 Asp Asp Asn Cys Val Thr Val Cys Asn Met Glu Asn Val Asp Ala Met
 645 650 655
 Gly Val His Thr Gly Asp Ser Val Val Val Ala Pro Ala Gln Thr Leu

-continued

660					665					670					
Ser	Asn	Ala	Glu	Phe	Gln	Met	Leu	Arg	Arg	Thr	Ser	Ile	Asn	Val	Val
	675						680					685			
Arg	His	Leu	Gly	Ile	Val	Gly	Glu	Cys	Asn	Ile	Gln	Phe	Ala	Leu	His
	690					695					700				
Pro	Thr	Ser	Met	Glu	Tyr	Cys	Ile	Ile	Glu	Val	Asn	Ala	Arg	Leu	Ser
	705					710					715				720
Arg	Ser	Ser	Ala	Leu	Ala	Ser	Lys	Ala	Thr	Gly	Tyr	Pro	Leu	Ala	Phe
			725						730					735	
Ile	Ala	Ala	Lys	Ile	Ala	Leu	Gly	Ile	Pro	Leu	Pro	Glu	Ile	Lys	Asn
			740					745					750		
Val	Val	Ser	Gly	Lys	Thr	Ser	Ala	Cys	Phe	Glu	Pro	Ser	Leu	Asp	Tyr
			755				760					765			
Met	Val	Thr	Lys	Ile	Pro	Arg	Trp	Asp	Leu	Asp	Arg	Phe	His	Gly	Thr
	770					775					780				
Ser	Ser	Arg	Ile	Gly	Ser	Ser	Met	Lys	Ser	Val	Gly	Glu	Val	Met	Ala
	785					790					795				800
Ile	Gly	Arg	Thr	Phe	Glu	Glu	Ser	Phe	Gln	Lys	Ala	Leu	Arg	Met	Cys
				805					810					815	
His	Pro	Ser	Ile	Glu	Gly	Phe	Thr	Pro	Arg	Leu	Pro	Met	Asn	Lys	Glu
			820						825				830		
Trp	Pro	Ser	Asn	Leu	Asp	Leu	Arg	Lys	Glu	Leu	Ser	Glu	Pro	Ser	Ser
			835				840					845			
Thr	Arg	Ile	Tyr	Ala	Ile	Ala	Lys	Ala	Ile	Asp	Asp	Asn	Met	Ser	Leu
	850					855					860				
Asp	Glu	Ile	Glu	Lys	Leu	Thr	Tyr	Ile	Asp	Lys	Trp	Phe	Leu	Tyr	Lys
	865					870					875				880
Met	Arg	Asp	Ile	Leu	Asn	Met	Glu	Lys	Thr	Leu	Lys	Gly	Leu	Asn	Ser
			885						890					895	
Glu	Ser	Met	Thr	Glu	Glu	Thr	Leu	Lys	Arg	Ala	Lys	Glu	Ile	Gly	Phe
			900					905					910		
Ser	Asp	Lys	Gln	Ile	Ser	Lys	Cys	Leu	Gly	Leu	Thr	Glu	Ala	Gln	Thr
		915					920					925			
Arg	Glu	Leu	Arg	Leu	Lys	Lys	Asn	Ile	His	Pro	Trp	Val	Lys	Gln	Ile
	930						935				940				
Asp	Thr	Leu	Ala	Ala	Glu	Tyr	Pro	Ser	Val	Thr	Asn	Tyr	Leu	Tyr	Val
	945					950					955				960
Thr	Tyr	Asn	Gly	Gln	Glu	His	Asp	Val	Asn	Phe	Asp	Asp	His	Gly	Met
				965					970					975	
Met	Val	Leu	Gly	Cys	Gly	Pro	Tyr	His	Ile	Gly	Ser	Ser	Val	Glu	Phe
		980						985					990		
Asp	Trp	Cys	Ala	Val	Ser	Ser	Ile	Arg	Thr	Leu	Arg	Gln	Leu	Gly	Lys
		995					1000					1005			
Lys	Thr	Val	Val	Val	Asn	Cys	Asn	Pro	Glu	Thr	Val	Ser	Thr	Asp	
	1010						1015					1020			
Phe	Asp	Glu	Cys	Asp	Lys	Leu	Tyr	Phe	Glu	Glu	Leu	Ser	Leu	Glu	
	1025						1030				1035				
Arg	Ile	Leu	Asp	Ile	Tyr	His	Gln	Glu	Ala	Cys	Gly	Gly	Cys	Ile	
	1040					1045					1050				
Ile	Ser	Val	Gly	Gly	Gln	Ile	Pro	Asn	Asn	Leu	Ala	Val	Pro	Leu	
	1055					1060					1065				
Tyr	Lys	Asn	Gly	Val	Lys	Ile	Met	Gly	Thr	Ser	Pro	Leu	Gln	Ile	
	1070					1075					1080				

-continued

Asp	Arg	Ala	Glu	Asp	Arg	Ser	Ile	Phe	Ser	Ala	Val	Leu	Asp	Glu
1085						1090					1095			
Leu	Lys	Val	Ala	Gln	Ala	Pro	Trp	Lys	Ala	Val	Asn	Thr	Leu	Asn
1100						1105					1110			
Glu	Ala	Leu	Glu	Phe	Ala	Lys	Ser	Val	Asp	Tyr	Pro	Cys	Leu	Leu
1115						1120					1125			
Arg	Pro	Ser	Tyr	Val	Leu	Ser	Gly	Ser	Ala	Met	Asn	Val	Val	Phe
1130						1135					1140			
Ser	Glu	Asp	Glu	Met	Lys	Lys	Phe	Leu	Glu	Glu	Ala	Thr	Arg	Val
1145						1150					1155			
Ser	Gln	Glu	His	Pro	Val	Val	Leu	Thr	Lys	Phe	Val	Glu	Gly	Ala
1160						1165					1170			
Arg	Glu	Val	Glu	Met	Asp	Ala	Val	Gly	Lys	Asp	Gly	Arg	Val	Ile
1175						1180					1185			
Ser	His	Ala	Ile	Ser	Glu	His	Val	Glu	Asp	Ala	Gly	Val	His	Ser
1190						1195					1200			
Gly	Asp	Ala	Thr	Leu	Met	Leu	Pro	Thr	Gln	Thr	Ile	Ser	Gln	Gly
1205						1210					1215			
Ala	Ile	Glu	Lys	Val	Lys	Asp	Ala	Thr	Arg	Lys	Ile	Ala	Lys	Ala
1220						1225					1230			
Phe	Ala	Ile	Ser	Gly	Pro	Phe	Asn	Val	Gln	Phe	Leu	Val	Lys	Gly
1235						1240					1245			
Asn	Asp	Val	Leu	Val	Ile	Glu	Cys	Asn	Leu	Arg	Ala	Ser	Arg	Ser
1250						1255					1260			
Phe	Pro	Phe	Val	Ser	Lys	Thr	Leu	Gly	Val	Asp	Phe	Ile	Asp	Val
1265						1270					1275			
Ala	Thr	Lys	Val	Met	Ile	Gly	Glu	Asn	Val	Asp	Glu	Lys	His	Leu
1280						1285					1290			
Pro	Thr	Leu	Asp	His	Pro	Ile	Ile	Pro	Ala	Asp	Tyr	Val	Ala	Ile
1295						1300					1305			
Lys	Ala	Pro	Met	Phe	Ser	Trp	Pro	Arg	Leu	Arg	Asp	Ala	Asp	Pro
1310						1315					1320			
Ile	Leu	Arg	Cys	Glu	Met	Ala	Ser	Thr	Gly	Glu	Val	Ala	Cys	Phe
1325						1330					1335			
Gly	Glu	Gly	Ile	His	Thr	Ala	Phe	Leu	Lys	Ala	Met	Leu	Ser	Thr
1340						1345					1350			
Gly	Phe	Lys	Ile	Pro	Gln	Lys	Gly	Ile	Leu	Ile	Gly	Ile	Gln	Gln
1355						1360					1365			
Ser	Phe	Arg	Pro	Arg	Phe	Leu	Gly	Val	Ala	Glu	Gln	Leu	His	Asn
1370						1375					1380			
Glu	Gly	Phe	Lys	Leu	Phe	Ala	Thr	Glu	Ala	Thr	Ser	Asp	Trp	Leu
1385						1390					1395			
Asn	Ala	Asn	Asn	Val	Pro	Ala	Asn	Pro	Val	Ala	Trp	Pro	Ser	Gln
1400						1405					1410			
Glu	Gly	Gln	Asn	Pro	Ser	Leu	Ser	Ser	Ile	Arg	Lys	Leu	Ile	Arg
1415						1420					1425			
Asp	Gly	Ser	Ile	Asp	Leu	Val	Ile	Asn	Leu	Pro	Asn	Asn	Asn	Thr
1430						1435					1440			
Lys	Phe	Val	His	Asp	Asn	Tyr	Val	Ile	Arg	Arg	Thr	Ala	Val	Asp
1445						1450					1455			
Ser	Gly	Ile	Pro	Leu	Leu	Thr	Asn	Phe	Gln	Val	Thr	Lys	Leu	Phe
1460						1465					1470			
Ala	Glu	Ala	Val	Gln	Lys	Ser	Arg	Lys	Val	Asp	Ser	Lys	Ser	Leu
1475						1480					1485			

-continued

Phe His Tyr Arg Gln Tyr Ser Ala Gly Lys Ala Ala
 1490 1495 1500

<210> SEQ ID NO 15

<211> LENGTH: 20

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 15

cggaagccac atcagactgg

20

<210> SEQ ID NO 16

<211> LENGTH: 24

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 16

ggagagtgaa acttgacaat catc

24

<210> SEQ ID NO 17

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 17

tactgctcag aatcatggc

19

<210> SEQ ID NO 18

<211> LENGTH: 19

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 18

tcatcaccaa ctgaacagg

19

<210> SEQ ID NO 19

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 19

ggttaagaga aggaggagct g

21

<210> SEQ ID NO 20

<211> LENGTH: 21

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 20

aaccagtctt cagtgtcctc a

21

<210> SEQ ID NO 21

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

<400> SEQUENCE: 21

cctgccaccc cagtg

15

<210> SEQ ID NO 22

<211> LENGTH: 15

<212> TYPE: DNA

<213> ORGANISM: Homo sapiens

-continued

<400> SEQUENCE: 22

cctgccaacc cagtg

15

<210> SEQ ID NO 23

<211> LENGTH: 8

<212> TYPE: PRT

<213> ORGANISM: Homo sapiens

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (3)..(3)

<223> OTHER INFORMATION: X is A or S

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (6)..(6)

<223> OTHER INFORMATION: X is T or S

<220> FEATURE:

<221> NAME/KEY: MISC_FEATURE

<222> LOCATION: (7)..(7)

<223> OTHER INFORMATION: X is A or Q

<400> SEQUENCE: 23

Pro Val Xaa Trp Pro Xaa Xaa Glu
1 5

What is claimed is:

1. A method for treating or preventing hyperammonemia in a subject with a suboptimal urea cycle function, the method comprising administering to a neonatal or infant human subject in need thereof a therapeutically effective amount of citrulline, wherein the hyperammonemia is a result of urea cycle enzyme dysfunction, and whereby treatment or prevention of hyperammonemia is accomplished.

2. The method of claim 1, wherein the sub-optimal urea cycle function comprises decreased urea cycle intermediate production.

3. The method of claim 1, wherein the sub-optimal urea cycle function results in decreased ammonia clearance.

4. The method of claim 1, wherein the sub-optimal urea cycle function is associated with impaired liver function resulting from an impaired or damaged liver.

5. The method of claim 4, wherein the impaired liver function resulting from an impaired or damaged liver is associated with hepatic cirrhosis.

6. The method of claim 1, wherein the citrulline is administered in a dose ranging from about 100 mg to about 30,000 mg.

7. The method of claim 6, wherein the citrulline is administered in a dose ranging from about 250 mg to about 1,000 mg.

8. A method for increasing ammonia clearance in a subject suffering from suboptimal urea cycle function, the method comprising administering to the subject a therapeutically effective amount of citrulline, whereby a level of citrulline in the subject is raised and ammonia clearance is increased, wherein the subject is a neonate or infant and wherein the suboptimal urea cycle function is a result of urea cycle enzyme dysfunction.

25

9. The method of claim 8, wherein the sub-optimal urea cycle function comprises decreased urea cycle intermediate production.

10. The method of claim 8, wherein the sub-optimal urea cycle function is associated with impaired liver function resulting from an impaired or damaged liver.

11. The method of claim 10, wherein the impaired liver function resulting from an impaired or damaged liver is associated with hepatic cirrhosis.

12. The method of claim 8, wherein the citrulline is administered in a dose ranging from about 100 mg to about 30,000 mg.

13. The method of claim 12, wherein the citrulline is administered in a dose ranging from about 250 mg to about 1,000 mg.

14. The method of claim 1, wherein the administering is intravenously.

15. The method of claim 8, wherein the administering is intravenously.

16. The method of claim 1, wherein the urea cycle enzyme comprises a CPSI enzyme.

17. The method of claim 8, wherein the urea cycle enzyme comprises a CPSI enzyme.

18. A method of treating N-acetyl-glutamate (NAG) deficiency related hyperammonemia, said method comprising administering to a human subject in need thereof an amount of citrulline effective for therapy of NAG deficiency related hyperammonemia.

55

* * * * *